

Computer Algebra Independent Integration Tests

Summer 2023 edition with Rubi V 4.17.3

4-Trig-functions/4.3-Tangent/104-4.3.3.1-a+b-tan-^m-c+d-tan-ⁿ-
A+B-tan-

Nasser M. Abbasi

December 9, 2023

Compiled on December 9, 2023 at 12:17am

Contents

1	Introduction	2
2	detailed summary tables of results	20
3	Listing of integrals	272
4	Appendix	7652

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	3
1.2	Results	4
1.3	Time and leaf size Performance	7
1.4	Performance based on number of rules Rubi used	9
1.5	Performance based on number of steps Rubi used	10
1.6	Solved integrals histogram based on leaf size of result	11
1.7	Solved integrals histogram based on CPU time used	12
1.8	Leaf size vs. CPU time used	13
1.9	list of integrals with no known antiderivative	14
1.10	List of integrals solved by CAS but has no known antiderivative	14
1.11	list of integrals solved by CAS but failed verification	14
1.12	Timing	15
1.13	Verification	15
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [855]. This is test number [104].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 13.3.1 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (855)	0.00 (0)
Mathematica	95.79 (819)	4.21 (36)
Maple	91.58 (783)	8.42 (72)
Fricas	91.58 (783)	8.42 (72)
Mupad	61.75 (528)	38.25 (327)
Maxima	50.06 (428)	49.94 (427)
Giac	31.70 (271)	68.30 (584)
Sympy	24.44 (209)	75.56 (646)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	72.515	3.275	20.000	4.211
Maple	56.491	34.269	0.819	8.421
Maxima	42.222	7.719	0.117	49.942
Fricas	27.836	61.637	2.105	8.421
Giac	15.673	15.906	0.117	68.304
Sympy	12.749	8.421	3.275	75.556
Mupad	0.000	61.754	0.000	38.246

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima

and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	36	100.00	0.00	0.00
Fricas	72	100.00	0.00	0.00
Maple	72	100.00	0.00	0.00
Mupad	327	0.00	100.00	0.00
Maxima	427	42.86	12.41	44.73
Giac	584	55.99	33.90	10.10
Sympy	646	72.45	24.15	3.41

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.47
Maple	0.52
Sympy	0.81
Giac	0.91
Rubi	0.96
Mathematica	3.46
Fricas	4.07
Mupad	12.32

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mathematica	195.87	0.99	158.00	0.91
Rubi	199.12	0.99	189.00	1.00
Maxima	324.83	1.92	185.50	1.04
Giac	331.60	2.42	184.00	1.77
Sympy	509.15	3.98	258.00	2.06
Mupad	3189.57	13.63	221.00	1.39
Fricas	3680.38	13.86	489.00	2.57
Maple	224520.77	878.53	227.00	1.14

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following gives the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {196, 197, 198, 199, 200, 201, 202, 203, 226, 227, 228, 229, 230, 231, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 473, 474, 475, 476, 477, 478, 568, 569, 570, 571, 572, 573, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 657, 664}

Mathematica {823, 824, 829}

Maple {427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 634, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

1.14.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the Timofeev test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.14.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

1.14.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```


1.14.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer, 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer, Leaf size of result.
4. integer, Leaf size of the optimal antiderivative.
5. number, CPU time used to solve this integral. 0 if failed.
6. string, The integral in Latex format
7. string, The input used in CAS own syntax.
8. string, The result (antiderivative) produced by CAS in Latex format
9. string, The optimal antiderivative in Latex format.
10. integer, 0 or 1. Indicates if problem has known antiderivative or not
11. String, The result (antiderivative) in CAS own syntax.
12. String, The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String, Small string description of why the grade was given.
14. integer, 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer, Number of steps used.
16. integer, Number of rules used.
17. integer, Integrand leaf size.
18. real number, Ratio, Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String, The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2013
Design v0.01

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	21
2.2	Detailed conclusion table per each integral for all CAS systems	32
2.3	Detailed conclusion table specific for Rubi results	246

2.1 List of integrals sorted by grade for each CAS

2.1.1	Rubi	21
2.1.2	Mma	22
2.1.3	Maple	24
2.1.4	Fricas	25
2.1.5	Maxima	26
2.1.6	Giac	27
2.1.7	Mupad	29
2.1.8	Sympy	30

2.1.1 Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544,

545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830, 831, 832, 833, 834, 835, 836, 837, 838, 839, 840, 841, 842, 843, 844, 845, 846, 847, 848, 849, 850, 851, 852, 853, 854, 855 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.2 Mma

A grade { 1, 2, 3, 4, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 44, 45, 46, 47, 48, 51, 52, 53, 54, 55, 56, 59, 60, 61, 62, 63, 64, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 101, 102, 103, 105, 106, 107, 109, 110, 111, 112, 113, 114, 115, 116, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 140, 141, 142, 143, 146, 147, 148, 149, 150, 151, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 166, 167, 168, 169, 170, 177, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 219, 220, 221, 222, 223, 224, 225, 232, 233, 234, 235, 270, 298, 299, 300, 301, 304, 315, 317, 318, 319, 320, 321, 322, 323, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 338, 339, 340, 343, 344, 345, 346, 347, 348, 349, 352, 354, 355, 356, 360, 362, 363, 364, 367, 369, 370, 373, 374, 382, 383, 384, 400, 401, 416, 417, 418, 419, 420, 421, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481,

482, 483, 484, 485, 494, 495, 496, 497, 498, 499, 500, 501, 504, 505, 506, 507, 509, 510, 511, 512, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 771, 772, 773, 774, 775, 776, 779, 780, 781, 782, 783, 784, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 816, 817, 818, 819, 820, 821, 822, 823, 825, 830, 831, 832, 833, 834, 835, 838, 839, 840, 841, 842, 846, 847, 848, 849, 850, 851, 852, 853, 854, 855 }
}

B grade { 164, 165, 171, 172, 173, 174, 175, 176, 178, 302, 314, 324, 336, 337, 410, 463, 486, 548, 549, 550, 551, 677, 815, 824, 826, 827, 828, 829 }

C grade { 5, 6, 7, 8, 41, 42, 43, 49, 50, 57, 58, 65, 66, 100, 104, 108, 117, 118, 138, 139, 144, 145, 152, 153, 200, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 303, 305, 306, 307, 308, 309, 310, 311, 312, 313, 316, 341, 342, 350, 351, 353, 357, 358, 359, 361, 365, 366, 368, 371, 372, 375, 376, 377, 378, 379, 380, 381, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 402, 403, 404, 405, 406, 407, 408, 409, 411, 412, 413, 414, 415, 422, 423, 424, 425, 426, 446, 502, 503, 508, 513, 514, 515, 516, 631, 769, 770, 777, 778, 785, 786, 836, 837, 843, 844, 845 }

F normal fail { 212, 213, 214, 215, 216, 217, 218, 226, 227, 228, 229, 230, 231, 487, 488, 489, 490, 491, 492, 493, 567, 568, 569, 570, 571, 572, 573, 656, 657, 658, 659, 660, 661, 662, 663, 664 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.1.3 Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 119, 126, 127, 128, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 196, 197, 198, 199, 200, 201, 202, 203, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 513, 516, 517, 518, 521, 522, 523, 524, 527, 528, 529, 530, 531, 532, 533, 534, 535, 577, 578, 579, 580, 581, 582, 588, 592, 593, 594, 597, 598, 599, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 706, 707, 708, 709, 710, 711, 712, 713, 714, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 792, 793, 794, 795, 796, 797, 798, 801, 802, 803, 804, 806, 807, 808, 812, 813, 814, 815, 817, 819, 820, 825, 826, 827, 828, 829, 833, 834, 835, 839, 840, 841, 842, 846, 847, 848, 849, 850, 854, 855 }

B grade { 112, 113, 114, 115, 116, 117, 118, 120, 121, 122, 123, 124, 125, 129, 130, 131, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 514, 515, 519, 520, 525, 526, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 574, 575, 576, 583, 584, 585, 586, 587, 589, 590, 591, 595, 596, 600, 601, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 791, 799, 800, 805, 809, 810, 811, 816, 818, 821, 822, 823, 824, 830, 831, 832, 836, 837, 838, 843, 844, 845, 853 }

C grade { 462, 473, 474, 475, 476, 477, 478 }

F normal fail { 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 567, 568, 569, 570, 571, 572, 573, 656, 657, 658, 659, 660, 661, 662, 663, 664, 705, 715, 727, 851, 852 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.4 Fricas

A grade { 2, 3, 4, 5, 6, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 277, 278, 298, 299, 300, 301, 302, 304, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 365, 375, 665, 667, 668, 670, 671, 672, 673, 674, 677, 678, 679, 681, 682, 683, 684, 685, 686, 687, 688, 690, 691, 692, 694, 696, 697, 698, 699, 700, 701, 702, 703, 704, 708, 709, 710, 712, 713, 714, 716, 717, 718, 719, 720, 721, 722, 724, 725, 726, 728, 729, 730, 731, 732, 733, 734, 735, 737, 738, 739, 741, 742, 743, 744, 745, 746, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 785, 786, 792, 793, 794, 801, 802, 803, 804, 812, 813, 814, 815, 825, 826, 827, 828, 829, 833, 834, 835, 839, 840, 841, 842, 846, 847, 848, 849, 850, 853, 854 }

B grade { 1, 7, 8, 29, 30, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 275, 276, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 303, 305, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 366, 367, 368, 369, 370, 371, 372, 373, 374, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530,

531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 666, 675, 676, 689, 695, 706, 707, 740, 747, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 787, 788, 789, 790, 791, 795, 796, 797, 798, 799, 800, 805, 806, 807, 808, 809, 810, 811, 816, 817, 818, 819, 820, 821, 822, 823, 824, 830, 831, 832, 836, 837, 838, 843, 844, 845, 855 }

C grade { 416, 417, 418, 419, 420, 421, 606, 607, 608, 609, 610, 611, 669, 680, 693, 711, 723, 736 }

F normal fail { 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 567, 568, 569, 570, 571, 572, 573, 656, 657, 658, 659, 660, 661, 662, 663, 664, 705, 715, 727, 851, 852 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.1.5 Maxima

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 119, 120, 121, 125, 126, 127, 128, 129, 130, 131, 132, 133, 196, 197, 198, 199, 200, 201, 202, 203, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 295, 296, 297, 298, 299, 300, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 508, 512, 513, 514, 515, 516, 517, 518, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 666, 667, 668, 669, 670, 677, 678, 679, 680, 681, 682, 690, 691, 692, 693, 694, 695, 696, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 800, 801, 802, 803, 804, 811, 813, 814, 815, 824, 826, 827, 828, 829, 832, 833, 838, 841, 845, 846, 855 }

B grade { 112, 113, 114, 115, 116, 117, 118, 122, 123, 124, 292, 293, 294, 302, 373, 502, 503, 504, 505, 506, 507, 509, 510, 511, 536, 537, 538, 541, 542, 543, 547, 548, 549, 665, 676, 689, 787, 788, 789, 790, 795, 796, 797, 798, 799, 805, 806, 807, 808, 809, 810, 812, 816, 817, 818, 819, 820, 821, 822, 823, 825, 830, 831, 836, 844, 850 }

C grade { 301 }

F normal fail { 154, 155, 156, 161, 162, 163, 165, 169, 170, 173, 178, 204, 205, 206, 207, 212, 213, 214, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 231, 317, 318, 319, 321, 322, 323, 324, 325, 326, 328, 329, 330, 333, 335, 336, 343, 344, 345, 347, 348, 349, 352, 354, 355, 362, 375, 376, 377, 427, 428, 429, 431, 433, 434, 435, 436, 438, 440, 442, 443, 444, 446, 448, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 463, 464, 465, 466, 469, 470, 471, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 539, 540, 545, 546, 550, 553, 567, 568, 569, 570, 571, 572, 573, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 624, 625, 626, 627, 628, 629, 630, 631, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 656, 657, 658, 659, 660, 661, 662, 851, 852 }

F(-1) timeout fail { 157, 158, 159, 160, 164, 166, 167, 168, 171, 172, 174, 175, 176, 177, 230, 331, 332, 337, 338, 339, 350, 351, 356, 357, 358, 359, 360, 363, 364, 367, 369, 430, 432, 437, 439, 441, 445, 447, 449, 450, 461, 462, 467, 468, 472, 544, 551, 552, 623, 632, 655, 663, 664 }

F(-2) exception fail { 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 208, 209, 210, 211, 215, 216, 217, 320, 327, 334, 340, 341, 342, 346, 353, 361, 365, 366, 368, 370, 371, 372, 374, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 671, 672, 673, 674, 675, 683, 684, 685, 686, 687, 688, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 791, 792, 793, 794, 834, 835, 837, 839, 840, 842, 843, 847, 848, 849, 853, 854 }

2.1.6 Giac

A grade { 32, 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 235, 243, 244, 251, 252, 253, 260, 261, 262, 263, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 280, 281, 282, 283, 288, 297, 299, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 475, 668, 672, 679, 692, 711, 712, 713, 714, 720, 721, 722, 723, 724, 725, 726, 733, 734, 735, 736, 737, 738, 739 }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 33, 34, 35, 39, 232, 233, 234, 236, 237, 238, 239, 240, 241, 242, 245, 246, 247, 248, 249, 250, 254, 255, 256, 257, 258, 259, 264, 265, 266, 277, 278, 279, 284, 285, 286, 287, 289, 290, 291, 292, 293, 294, 295, 296, 298, 300, 302, 303, 304, 305, 373, 374, 375, 376, 477, 666, 667, 669, 670, 671, 673, 674, 675, 677, 678, 680, 681, 682, 683, 684, 685, 686, 687, 688, 690, 691, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 706, 707, 708, 709, 710, 716, 717, 718, 719, 728, 729, 730, 731, 732, 789, 798, 854 }

C grade { 301 }

F normal fail { 69, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 428, 473, 474, 476, 478, 479, 480, 481, 482, 483, 484, 485, 486, 489, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 522, 523, 524, 525, 526, 527, 528, 529, 530, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 600, 601, 602, 606, 607, 608, 609, 610, 611, 638, 643, 648, 656, 657, 658, 659, 660, 661, 662, 665, 676, 689, 705, 715, 727, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 788, 790, 791, 792, 793, 794, 796, 797, 799, 800, 801, 802, 803, 804, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830, 831, 832, 833, 834, 835, 836, 837, 838, 839, 840, 841, 842, 843, 844, 845, 846, 847, 848, 849, 850, 851, 852, 853, 855 }

F(-1) timeout fail { 67, 68, 70, 83, 84, 187, 188, 189, 193, 194, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 487, 488, 490, 491, 492, 599, 603, 604, 605, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 645, 650, 652, 654, 663, 664, 787, 795, 805, 806, 820 }

F(-2) exception fail { 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 190, 191, 192, 195, 521, 531, 612, 613, 614, 615, 616, 617, 618, 636, 637, 639, 640, 641, 642, 644, 646, 647,

649, 651, 653, 655 }

2.1.7 Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 155, 156, 180, 181, 196, 197, 198, 199, 200, 201, 202, 203, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 428, 429, 453, 454, 473, 474, 475, 476, 477, 478, 606, 607, 608, 609, 610, 611, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 706, 707, 708, 709, 710, 711, 712, 713, 714, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 790, 791, 792, 793, 794, 801, 802, 803, 804, 812, 813, 814, 815, 825, 826, 827, 828, 829, 832, 833, 834, 835, 839, 840, 841, 842, 846, 847, 848, 849, 850, 853, 854, 855 }

C grade { }

F normal fail { }

F(-1) timeout fail { 154, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 332, 427, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 479, 480, 481, 482, 483, 484,

485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 705, 715, 727, 787, 788, 789, 795, 796, 797, 798, 799, 800, 805, 806, 807, 808, 809, 810, 811, 816, 817, 818, 819, 820, 821, 822, 823, 824, 830, 831, 836, 837, 838, 843, 844, 845, 851, 852 }

F(-2) exception fail { }

2.1.8 Sympy

A grade { 2, 3, 5, 6, 9, 11, 12, 13, 14, 15, 16, 17, 19, 23, 24, 25, 26, 31, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 232, 233, 240, 241, 242, 243, 250, 251, 252, 259, 260, 261, 262, 263, 265, 266, 301, 305, 670, 671, 682, 683, 684, 696, 697, 698, 699, 706, 707, 708, 709, 710, 711, 712, 713, 714, 716, 717, 718, 719, 721, 722, 723, 724, 725, 726, 728, 729, 730, 733, 734, 735, 736, 737, 738, 739, 854 }

B grade { 1, 4, 7, 8, 10, 18, 20, 21, 22, 27, 28, 29, 30, 32, 234, 235, 236, 237, 238, 239, 244, 245, 246, 247, 248, 249, 253, 254, 255, 256, 257, 258, 264, 298, 299, 300, 302, 303, 304, 314, 665, 666, 667, 668, 672, 673, 674, 675, 676, 677, 678, 679, 681, 685, 686, 687, 688, 689, 690, 691, 692, 694, 695, 700, 701, 702, 703, 704, 720, 731, 732, 853 }

C grade { 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 306, 307, 308, 309, 310, 311, 312, 313, 315, 316, 669, 680, 693 }

F normal fail { 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 139, 140, 141, 142, 149, 150, 154, 155, 156, 157, 158, 159, 162, 163, 164, 165, 166, 171, 172, 173, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 192, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 228, 229, 230, 231, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 399, 400, 401, 402, 404, 405, 406, 407, 412, 413, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 435, 436, 437, 438, 439, 444, 445, 446, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 464, 465, 469, 470, 471, 472, 473, 474, 475,

476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 504, 505, 506, 507, 510, 511, 512, 516, 517, 518, 520, 521, 522, 523, 524, 525, 526, 527, 528, 530, 531, 532, 539, 540, 545, 546, 555, 556, 557, 558, 559, 560, 561, 563, 564, 567, 570, 571, 572, 575, 576, 577, 580, 581, 582, 586, 587, 588, 590, 591, 592, 593, 594, 595, 596, 597, 598, 600, 601, 602, 606, 607, 608, 609, 610, 611, 615, 616, 617, 618, 624, 625, 626, 638, 639, 640, 641, 643, 644, 645, 646, 649, 650, 653, 654, 655, 658, 659, 660, 662, 663, 664, 705, 715, 727, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 772, 773, 774, 775, 776, 777, 778, 780, 781, 782, 783, 784, 785, 786, 788, 789, 790, 791, 792, 793, 794, 797, 798, 799, 800, 801, 802, 808, 809, 810, 811, 812, 830, 831, 832, 833, 834, 835, 837, 838, 839, 840, 841, 842, 845, 846, 847, 848, 849, 850, 851, 852, 855 }

F(-1) timeout fail { 86, 87, 88, 89, 145, 146, 147, 148, 153, 160, 161, 167, 168, 169, 170, 174, 175, 176, 177, 189, 190, 191, 193, 194, 195, 212, 226, 227, 336, 337, 338, 339, 398, 403, 408, 409, 410, 411, 414, 415, 433, 434, 440, 441, 442, 443, 447, 448, 449, 450, 463, 466, 467, 468, 487, 502, 503, 508, 509, 513, 514, 515, 519, 529, 533, 534, 535, 536, 537, 538, 541, 542, 543, 544, 547, 548, 549, 550, 551, 552, 553, 554, 562, 565, 566, 568, 569, 573, 574, 578, 579, 583, 584, 585, 589, 599, 603, 604, 605, 612, 613, 614, 619, 620, 621, 622, 623, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 642, 647, 648, 651, 652, 656, 657, 661, 771, 779, 787, 795, 796, 803, 804, 805, 806, 807, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 836, 843, 844 }

F(-2) exception fail { 137, 138, 143, 144, 151, 152, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	91	91	86	95	82	170	167	284	82
N.S.	1	1.00	0.95	1.04	0.90	1.87	1.84	3.12	0.90
time (sec)	N/A	0.495	1.074	0.197	0.390	0.250	0.349	0.482	7.343

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	70	72	68	109	109	194	59
N.S.	1	1.00	1.01	1.04	0.99	1.58	1.58	2.81	0.86
time (sec)	N/A	0.365	0.331	0.050	0.337	0.239	0.282	0.337	7.891

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	66	50	50	64	53	103	38
N.S.	1	1.00	1.43	1.09	1.09	1.39	1.15	2.24	0.83
time (sec)	N/A	0.251	0.033	0.038	0.327	0.252	0.234	0.299	7.802

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	40	42	54	43	49	36	94	74	36
N.S.	1	1.05	1.35	1.08	1.22	0.90	2.35	1.85	0.90
time (sec)	N/A	0.397	0.052	0.559	0.305	0.251	1.031	0.400	7.701

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	46	97	59	64	62	53	104	39
N.S.	1	1.05	2.20	1.34	1.45	1.41	1.20	2.36	0.89
time (sec)	N/A	0.374	0.041	0.172	0.282	0.244	0.257	0.495	7.231

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	68	70	76	73	84	111	109	162	60
N.S.	1	1.03	1.12	1.07	1.24	1.63	1.60	2.38	0.88
time (sec)	N/A	0.491	0.419	0.168	0.286	0.246	0.314	0.609	7.329

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	91	102	101	103	166	168	221	80
N.S.	1	1.02	1.15	1.13	1.16	1.87	1.89	2.48	0.90
time (sec)	N/A	0.619	0.795	0.175	0.288	0.251	0.369	0.706	7.568

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	111	113	96	106	116	206	218	282	100
N.S.	1	1.02	0.86	0.95	1.05	1.86	1.96	2.54	0.90
time (sec)	N/A	0.764	0.909	0.198	0.294	0.250	0.722	0.853	8.460

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	141	147	101	121	112	236	236	408	153
N.S.	1	1.04	0.72	0.86	0.79	1.67	1.67	2.89	1.09
time (sec)	N/A	0.774	1.357	0.091	0.293	0.265	0.464	0.590	7.248

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	99	82	99	92	175	178	312	111
N.S.	1	0.93	0.77	0.93	0.86	1.64	1.66	2.92	1.04
time (sec)	N/A	0.492	0.824	0.077	0.299	0.253	0.346	0.435	7.653

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	70	58	76	71	121	122	214	76
N.S.	1	0.88	0.72	0.95	0.89	1.51	1.52	2.68	0.95
time (sec)	N/A	0.355	0.490	0.055	0.293	0.252	0.327	0.384	7.257

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	75	77	46	63	67	97	109	174	70
N.S.	1	1.03	0.61	0.84	0.89	1.29	1.45	2.32	0.93
time (sec)	N/A	0.600	0.502	0.174	0.293	0.248	1.218	0.636	8.131

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	79	81	54	74	74	102	109	155	87
N.S.	1	1.03	0.68	0.94	0.94	1.29	1.38	1.96	1.10
time (sec)	N/A	0.633	0.753	0.231	0.293	0.256	1.325	0.857	7.685

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	94	100	111	77	94	123	119	186	67
N.S.	1	1.06	1.18	0.82	1.00	1.31	1.27	1.98	0.71
time (sec)	N/A	0.628	0.500	0.245	0.301	0.245	0.331	1.084	7.399

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	126	143	104	111	181	182	255	93
N.S.	1	1.08	1.22	0.89	0.95	1.55	1.56	2.18	0.79
time (sec)	N/A	0.781	0.655	0.247	0.325	0.238	0.570	1.343	8.035

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	139	152	174	109	132	227	235	322	113
N.S.	1	1.09	1.25	0.78	0.95	1.63	1.69	2.32	0.81
time (sec)	N/A	0.940	1.134	0.268	0.300	0.256	0.536	1.102	7.918

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	182	192	123	146	132	291	292	504	230
N.S.	1	1.05	0.68	0.80	0.73	1.60	1.60	2.77	1.26
time (sec)	N/A	1.058	1.589	0.115	0.296	0.249	0.494	0.683	7.270

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	128	105	123	114	227	235	408	176
N.S.	1	0.93	0.76	0.89	0.83	1.64	1.70	2.96	1.28
time (sec)	N/A	0.596	1.017	0.097	0.302	0.252	0.468	0.535	7.564

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	99	73	100	96	175	184	312	125
N.S.	1	0.90	0.66	0.91	0.87	1.59	1.67	2.84	1.14
time (sec)	N/A	0.450	0.869	0.090	0.279	0.249	0.390	0.463	7.862

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	109	70	87	89	172	224	264	87
N.S.	1	1.02	0.65	0.81	0.83	1.61	2.09	2.47	0.81
time (sec)	N/A	0.811	0.876	0.211	0.294	0.259	1.597	0.859	7.336

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	118	94	84	84	141	219	257	76
N.S.	1	1.02	0.81	0.72	0.72	1.22	1.89	2.22	0.66
time (sec)	N/A	0.878	1.016	0.240	0.278	0.262	1.013	1.193	7.925

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	123	125	75	92	96	179	226	223	88
N.S.	1	1.02	0.61	0.75	0.78	1.46	1.84	1.81	0.72
time (sec)	N/A	0.904	1.097	0.207	0.294	0.265	1.029	1.143	8.115

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	134	145	89	104	115	181	182	254	93
N.S.	1	1.08	0.66	0.78	0.86	1.35	1.36	1.90	0.69
time (sec)	N/A	0.919	1.045	0.196	0.289	0.250	0.471	0.757	7.336

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	157	172	111	109	134	228	235	322	114
N.S.	1	1.10	0.71	0.69	0.85	1.45	1.50	2.05	0.73
time (sec)	N/A	1.087	1.448	0.245	0.296	0.265	1.186	0.827	7.967

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	180	195	207	130	151	287	296	391	140
N.S.	1	1.08	1.15	0.72	0.84	1.59	1.64	2.17	0.78
time (sec)	N/A	1.291	1.212	0.267	0.295	0.244	0.726	0.912	8.236

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	225	240	138	168	150	345	348	600	308
N.S.	1	1.07	0.61	0.75	0.67	1.53	1.55	2.67	1.37
time (sec)	N/A	1.422	1.827	0.193	0.282	0.251	0.818	0.821	7.544

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	168	157	119	145	132	279	291	504	240
N.S.	1	0.93	0.71	0.86	0.79	1.66	1.73	3.00	1.43
time (sec)	N/A	0.721	1.164	0.162	0.296	0.254	0.498	0.633	7.503

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	128	86	122	114	227	241	408	181
N.S.	1	0.91	0.61	0.87	0.81	1.62	1.72	2.91	1.29
time (sec)	N/A	0.559	1.019	0.111	0.279	0.249	0.442	0.543	7.205

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	142	144	89	109	107	246	289	332	133
N.S.	1	1.01	0.63	0.77	0.75	1.73	2.04	2.34	0.94
time (sec)	N/A	1.071	1.206	0.231	0.281	0.247	1.854	1.234	7.237

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	144	146	121	91	102	254	260	336	110
N.S.	1	1.01	0.84	0.63	0.71	1.76	1.81	2.33	0.76
time (sec)	N/A	1.133	1.373	0.244	0.282	0.267	1.302	0.792	8.384

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	162	122	105	108	255	252	317	102
N.S.	1	1.04	0.78	0.67	0.69	1.63	1.62	2.03	0.65
time (sec)	N/A	1.190	4.851	0.237	0.295	0.272	1.175	0.904	7.671

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	163	165	143	115	117	249	292	291	113
N.S.	1	1.01	0.88	0.71	0.72	1.53	1.79	1.79	0.69
time (sec)	N/A	1.198	1.148	0.234	0.292	0.250	2.472	0.975	7.625

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	177	193	91	111	136	228	235	322	114
N.S.	1	1.09	0.51	0.63	0.77	1.29	1.33	1.82	0.64
time (sec)	N/A	1.246	1.107	0.267	0.291	0.246	0.654	1.123	7.899

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	200	219	120	130	153	287	296	391	140
N.S.	1	1.10	0.60	0.65	0.76	1.44	1.48	1.96	0.70
time (sec)	N/A	1.457	1.650	0.266	0.290	0.243	2.522	1.216	7.938

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	223	245	150	143	172	332	347	459	162
N.S.	1	1.10	0.67	0.64	0.77	1.49	1.56	2.06	0.73
time (sec)	N/A	1.655	3.744	0.296	0.277	0.247	1.087	1.292	9.452

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	129	122	137	146	0	186	196	122	141
N.S.	1	0.95	1.06	1.13	0.00	1.44	1.52	0.95	1.09
time (sec)	N/A	0.599	2.048	0.098	0.000	0.249	0.411	0.606	7.479

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	99	103	125	0	127	151	99	95
N.S.	1	0.98	1.02	1.24	0.00	1.26	1.50	0.98	0.94
time (sec)	N/A	0.469	1.393	0.070	0.000	0.254	0.310	0.471	7.770

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	64	75	86	0	66	119	79	81
N.S.	1	0.96	1.12	1.28	0.00	0.99	1.78	1.18	1.21
time (sec)	N/A	0.379	0.503	0.058	0.000	0.242	0.215	0.357	7.833

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	56	54	0	42	87	83	45
N.S.	1	1.00	1.19	1.15	0.00	0.89	1.85	1.77	0.96
time (sec)	N/A	0.220	0.297	0.056	0.000	0.234	0.135	0.340	7.149

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	67	86	85	0	68	116	97	98
N.S.	1	1.08	1.39	1.37	0.00	1.10	1.87	1.56	1.58
time (sec)	N/A	0.419	0.824	0.141	0.000	0.253	0.196	0.428	7.976

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	102	98	140	0	129	158	133	126
N.S.	1	1.00	0.96	1.37	0.00	1.26	1.55	1.30	1.24
time (sec)	N/A	0.580	1.257	0.142	0.000	0.259	0.325	0.550	7.720

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	131	127	111	159	0	188	199	160	153
N.S.	1	0.97	0.85	1.21	0.00	1.44	1.52	1.22	1.17
time (sec)	N/A	0.737	1.816	0.149	0.000	0.259	0.415	0.691	7.937

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	155	150	114	185	0	249	253	184	174
N.S.	1	0.97	0.74	1.19	0.00	1.61	1.63	1.19	1.12
time (sec)	N/A	0.907	1.898	0.155	0.000	0.255	0.403	0.890	7.950

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	142	147	197	173	0	150	262	120	141
N.S.	1	1.04	1.39	1.22	0.00	1.06	1.85	0.85	0.99
time (sec)	N/A	0.730	1.506	0.104	0.000	0.259	0.437	0.684	7.780

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	106	158	122	0	84	223	107	114
N.S.	1	1.03	1.53	1.18	0.00	0.82	2.17	1.04	1.11
time (sec)	N/A	0.602	0.850	0.078	0.000	0.251	0.309	0.549	7.154

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	82	92	73	0	52	167	105	106
N.S.	1	1.08	1.21	0.96	0.00	0.68	2.20	1.38	1.39
time (sec)	N/A	0.374	0.670	0.068	0.000	0.242	0.195	0.427	8.132

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	79	89	73	0	54	162	110	70
N.S.	1	0.99	1.11	0.91	0.00	0.68	2.02	1.38	0.88
time (sec)	N/A	0.303	0.851	0.060	0.000	0.235	0.191	0.405	7.442

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	109	105	122	0	86	219	121	129
N.S.	1	1.15	1.11	1.28	0.00	0.91	2.31	1.27	1.36
time (sec)	N/A	0.649	1.755	0.161	0.000	0.244	0.274	0.604	7.371

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	141	147	130	177	0	152	267	161	164
N.S.	1	1.04	0.92	1.26	0.00	1.08	1.89	1.14	1.16
time (sec)	N/A	0.924	2.196	0.175	0.000	0.273	0.474	0.825	8.023

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	170	181	143	209	0	215	323	176	188
N.S.	1	1.06	0.84	1.23	0.00	1.26	1.90	1.04	1.11
time (sec)	N/A	1.096	2.711	0.174	0.000	0.254	0.481	1.132	7.914

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	191	203	270	214	0	174	337	142	184
N.S.	1	1.06	1.41	1.12	0.00	0.91	1.76	0.74	0.96
time (sec)	N/A	1.058	1.779	0.124	0.000	0.258	0.601	1.080	8.027

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	148	159	166	160	0	104	296	127	146
N.S.	1	1.07	1.12	1.08	0.00	0.70	2.00	0.86	0.99
time (sec)	N/A	0.860	2.135	0.099	0.000	0.251	0.562	0.817	7.688

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	124	135	147	128	0	78	258	129	111
N.S.	1	1.09	1.19	1.03	0.00	0.63	2.08	1.04	0.90
time (sec)	N/A	0.630	1.130	0.085	0.000	0.246	0.288	0.694	7.723

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	116	148	128	0	74	260	127	147
N.S.	1	1.05	1.35	1.16	0.00	0.67	2.36	1.15	1.34
time (sec)	N/A	0.471	1.530	0.083	0.000	0.246	0.294	0.559	7.124

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	112	111	109	128	0	76	258	129	111
N.S.	1	0.99	0.97	1.14	0.00	0.68	2.30	1.15	0.99
time (sec)	N/A	0.393	1.114	0.080	0.000	0.244	0.270	0.541	7.608

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	131	153	132	159	0	104	292	143	164
N.S.	1	1.17	1.01	1.21	0.00	0.79	2.23	1.09	1.25
time (sec)	N/A	0.898	1.601	0.193	0.000	0.264	0.392	0.947	8.297

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	183	197	164	214	0	173	340	182	197
N.S.	1	1.08	0.90	1.17	0.00	0.95	1.86	0.99	1.08
time (sec)	N/A	1.251	2.579	0.175	0.000	0.269	0.532	0.907	7.324

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	216	233	173	246	0	235	398	207	221
N.S.	1	1.08	0.80	1.14	0.00	1.09	1.84	0.96	1.02
time (sec)	N/A	1.480	3.572	0.188	0.000	0.262	0.920	1.298	8.252

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	185	203	264	197	0	120	359	154	178
N.S.	1	1.10	1.43	1.06	0.00	0.65	1.94	0.83	0.96
time (sec)	N/A	1.169	1.349	0.116	0.000	0.247	1.151	1.201	8.087

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	159	181	158	147	0	88	301	149	178
N.S.	1	1.14	0.99	0.92	0.00	0.55	1.89	0.94	1.12
time (sec)	N/A	0.911	1.693	0.096	0.000	0.239	0.403	0.924	7.392

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	145	153	144	109	0	78	241	151	135
N.S.	1	1.06	0.99	0.75	0.00	0.54	1.66	1.04	0.93
time (sec)	N/A	0.713	1.635	0.109	0.000	0.233	0.490	0.767	7.612

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	143	148	141	109	0	78	246	149	172
N.S.	1	1.03	0.99	0.76	0.00	0.55	1.72	1.04	1.20
time (sec)	N/A	0.584	0.788	0.095	0.000	0.241	0.309	0.636	8.424

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	145	143	132	147	0	88	299	154	143
N.S.	1	0.99	0.91	1.01	0.00	0.61	2.06	1.06	0.99
time (sec)	N/A	0.495	1.341	0.086	0.000	0.232	0.351	0.607	7.672

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	162	191	152	196	0	122	359	165	196
N.S.	1	1.18	0.94	1.21	0.00	0.75	2.22	1.02	1.21
time (sec)	N/A	1.171	1.721	0.203	0.000	0.254	0.450	0.913	7.298

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	220	241	191	251	0	190	406	205	226
N.S.	1	1.10	0.87	1.14	0.00	0.86	1.85	0.93	1.03
time (sec)	N/A	1.565	3.578	0.206	0.000	0.246	0.906	1.287	8.154

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	255	279	204	283	0	253	466	228	251
N.S.	1	1.09	0.80	1.11	0.00	0.99	1.83	0.89	0.98
time (sec)	N/A	1.877	4.603	0.247	0.000	0.257	0.734	1.076	9.057

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	194	213	130	163	153	440	0	0	216
N.S.	1	1.10	0.67	0.84	0.79	2.27	0.00	0.00	1.11
time (sec)	N/A	1.098	2.110	0.305	0.323	0.277	0.000	0.000	1.978

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	143	150	111	124	130	383	0	0	168
N.S.	1	1.05	0.78	0.87	0.91	2.68	0.00	0.00	1.17
time (sec)	N/A	0.705	1.275	0.179	0.315	0.257	0.000	0.000	8.276

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	105	107	92	82	107	332	0	0	120
N.S.	1	1.02	0.88	0.78	1.02	3.16	0.00	0.00	1.14
time (sec)	N/A	0.445	0.704	0.130	0.304	0.263	0.000	0.000	7.997

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	75	77	75	63	87	272	0	0	96
N.S.	1	1.03	1.00	0.84	1.16	3.63	0.00	0.00	1.28
time (sec)	N/A	0.310	0.300	0.135	0.314	0.261	0.000	0.000	7.491

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	89	79	72	113	447	0	0	493
N.S.	1	1.03	0.92	0.84	1.31	5.20	0.00	0.00	5.73
time (sec)	N/A	0.547	1.106	0.371	0.305	0.257	0.000	0.000	8.165

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	123	134	118	114	145	649	0	0	168
N.S.	1	1.09	0.96	0.93	1.18	5.28	0.00	0.00	1.37
time (sec)	N/A	0.806	2.058	0.289	0.331	0.269	0.000	0.000	8.364

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	184	138	146	202	730	0	0	702
N.S.	1	1.09	0.82	0.86	1.20	4.32	0.00	0.00	4.15
time (sec)	N/A	1.103	2.937	0.275	0.309	0.266	0.000	0.000	7.849

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	210	236	157	171	249	823	0	0	735
N.S.	1	1.12	0.75	0.81	1.19	3.92	0.00	0.00	3.50
time (sec)	N/A	1.441	4.705	0.289	0.311	0.282	0.000	0.000	8.445

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	197	213	133	164	153	467	0	0	211
N.S.	1	1.08	0.68	0.83	0.78	2.37	0.00	0.00	1.07
time (sec)	N/A	1.077	1.946	0.125	0.328	0.269	0.000	0.000	9.199

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	137	137	114	123	130	415	0	0	163
N.S.	1	1.00	0.83	0.90	0.95	3.03	0.00	0.00	1.19
time (sec)	N/A	0.565	1.045	0.130	0.297	0.265	0.000	0.000	7.950

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	106	98	99	111	359	0	0	139
N.S.	1	0.99	0.92	0.93	1.04	3.36	0.00	0.00	1.30
time (sec)	N/A	0.394	0.556	0.103	0.312	0.254	0.000	0.000	7.871

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	113	115	108	87	130	514	0	0	553
N.S.	1	1.02	0.96	0.77	1.15	4.55	0.00	0.00	4.89
time (sec)	N/A	0.769	0.638	0.306	0.309	0.258	0.000	0.000	8.009

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	132	122	111	145	685	0	0	2338
N.S.	1	1.06	0.98	0.89	1.16	5.48	0.00	0.00	18.70
time (sec)	N/A	0.836	3.436	0.296	0.309	0.282	0.000	0.000	9.996

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	182	139	148	203	762	0	0	3027
N.S.	1	1.06	0.81	0.87	1.19	4.46	0.00	0.00	17.70
time (sec)	N/A	1.127	3.154	0.288	0.322	0.266	0.000	0.000	9.401

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	234	158	179	253	856	0	0	3084
N.S.	1	1.10	0.74	0.84	1.19	4.02	0.00	0.00	14.48
time (sec)	N/A	1.489	5.411	0.300	0.308	0.272	0.000	0.000	9.303

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	246	266	151	206	176	531	0	0	258
N.S.	1	1.08	0.61	0.84	0.72	2.16	0.00	0.00	1.05
time (sec)	N/A	1.431	3.619	0.184	0.313	0.277	0.000	0.000	8.714

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	168	132	165	153	477	0	0	212
N.S.	1	0.98	0.77	0.96	0.89	2.79	0.00	0.00	1.24
time (sec)	N/A	0.664	1.538	0.125	0.313	0.261	0.000	0.000	1.673

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	141	137	116	141	134	421	0	0	188
N.S.	1	0.97	0.82	1.00	0.95	2.99	0.00	0.00	1.33
time (sec)	N/A	0.504	0.874	0.126	0.301	0.252	0.000	0.000	1.156

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	147	149	126	125	154	610	0	0	597
N.S.	1	1.01	0.86	0.85	1.05	4.15	0.00	0.00	4.06
time (sec)	N/A	1.012	1.172	0.299	0.296	0.270	0.000	0.000	8.144

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	158	166	129	131	163	705	0	0	2947
N.S.	1	1.05	0.82	0.83	1.03	4.46	0.00	0.00	18.65
time (sec)	N/A	1.103	1.436	0.304	0.301	0.264	0.000	0.000	8.994

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	173	181	141	143	206	774	0	0	2991
N.S.	1	1.05	0.82	0.83	1.19	4.47	0.00	0.00	17.29
time (sec)	N/A	1.157	5.859	0.282	0.313	0.274	0.000	0.000	9.474

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	217	232	160	179	249	868	0	0	3048
N.S.	1	1.07	0.74	0.82	1.15	4.00	0.00	0.00	14.05
time (sec)	N/A	1.468	5.542	0.326	0.304	0.275	0.000	0.000	9.409

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	261	285	177	206	293	944	0	0	3094
N.S.	1	1.09	0.68	0.79	1.12	3.62	0.00	0.00	11.85
time (sec)	N/A	1.892	4.771	0.339	0.307	0.284	0.000	0.000	9.720

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	205	218	130	167	157	447	0	0	236
N.S.	1	1.06	0.63	0.81	0.77	2.18	0.00	0.00	1.15
time (sec)	N/A	1.099	2.430	0.172	0.311	0.282	0.000	0.000	8.696

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	159	165	111	128	134	391	0	0	188
N.S.	1	1.04	0.70	0.81	0.84	2.46	0.00	0.00	1.18
time (sec)	N/A	0.757	1.867	0.141	0.325	0.266	0.000	0.000	8.182

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	109	111	109	88	110	327	0	0	141
N.S.	1	1.02	1.00	0.81	1.01	3.00	0.00	0.00	1.29
time (sec)	N/A	0.460	1.356	0.135	0.302	0.255	0.000	0.000	1.071

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	82	84	82	71	91	328	0	0	117
N.S.	1	1.02	1.00	0.87	1.11	4.00	0.00	0.00	1.43
time (sec)	N/A	0.314	0.504	0.138	0.305	0.257	0.000	0.000	0.890

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	114	125	120	99	133	575	0	0	515
N.S.	1	1.10	1.05	0.87	1.17	5.04	0.00	0.00	4.52
time (sec)	N/A	0.782	2.059	0.303	0.289	0.265	0.000	0.000	7.797

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	167	183	131	139	184	742	0	0	2961
N.S.	1	1.10	0.78	0.83	1.10	4.44	0.00	0.00	17.73
time (sec)	N/A	1.134	3.644	0.316	0.305	0.274	0.000	0.000	9.456

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	219	237	150	171	232	835	0	0	3037
N.S.	1	1.08	0.68	0.78	1.06	3.81	0.00	0.00	13.87
time (sec)	N/A	1.517	4.417	0.319	0.306	0.281	0.000	0.000	9.578

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	209	219	149	154	160	440	0	0	233
N.S.	1	1.05	0.71	0.74	0.77	2.11	0.00	0.00	1.11
time (sec)	N/A	1.110	2.937	0.156	0.310	0.276	0.000	0.000	8.807

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	167	166	148	116	137	375	0	0	186
N.S.	1	0.99	0.89	0.69	0.82	2.25	0.00	0.00	1.11
time (sec)	N/A	0.778	1.724	0.125	0.316	0.254	0.000	0.000	0.735

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	119	126	118	96	116	369	0	0	163
N.S.	1	1.06	0.99	0.81	0.97	3.10	0.00	0.00	1.37
time (sec)	N/A	0.525	1.369	0.117	0.366	0.252	0.000	0.000	8.110

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	121	119	76	96	113	372	0	0	162
N.S.	1	0.98	0.63	0.79	0.93	3.07	0.00	0.00	1.34
time (sec)	N/A	0.408	0.623	0.126	0.310	0.255	0.000	0.000	7.558

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	167	156	126	161	624	0	0	563
N.S.	1	1.07	1.00	0.81	1.03	4.00	0.00	0.00	3.61
time (sec)	N/A	1.073	2.789	0.256	0.305	0.260	0.000	0.000	7.954

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	217	233	168	168	215	814	0	0	3051
N.S.	1	1.07	0.77	0.77	0.99	3.75	0.00	0.00	14.06
time (sec)	N/A	1.516	3.581	0.247	0.332	0.278	0.000	0.000	9.810

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	268	287	182	198	259	903	0	0	3106
N.S.	1	1.07	0.68	0.74	0.97	3.37	0.00	0.00	11.59
time (sec)	N/A	1.918	5.455	0.270	0.310	0.277	0.000	0.000	9.929

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	255	272	162	181	185	457	0	0	279
N.S.	1	1.07	0.64	0.71	0.73	1.79	0.00	0.00	1.09
time (sec)	N/A	1.522	3.108	0.147	0.335	0.262	0.000	0.000	8.545

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	211	219	204	142	162	392	0	0	230
N.S.	1	1.04	0.97	0.67	0.77	1.86	0.00	0.00	1.09
time (sec)	N/A	1.146	3.884	0.136	0.289	0.264	0.000	0.000	7.682

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	167	179	177	124	141	393	0	0	187
N.S.	1	1.07	1.06	0.74	0.84	2.35	0.00	0.00	1.12
time (sec)	N/A	0.849	2.216	0.132	0.287	0.259	0.000	0.000	7.945

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	153	158	147	121	136	391	0	0	186
N.S.	1	1.03	0.96	0.79	0.89	2.56	0.00	0.00	1.22
time (sec)	N/A	0.650	2.082	0.133	0.313	0.269	0.000	0.000	8.353

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	155	153	76	123	139	392	0	0	205
N.S.	1	0.99	0.49	0.79	0.90	2.53	0.00	0.00	1.32
time (sec)	N/A	0.537	0.889	0.131	0.294	0.248	0.000	0.000	7.608

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	192	211	217	153	186	644	0	0	528
N.S.	1	1.10	1.13	0.80	0.97	3.35	0.00	0.00	2.75
time (sec)	N/A	1.390	3.278	0.311	0.304	0.270	0.000	0.000	7.832

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	259	283	269	195	240	834	0	0	3002
N.S.	1	1.09	1.04	0.75	0.93	3.22	0.00	0.00	11.59
time (sec)	N/A	1.917	6.375	0.302	0.309	0.289	0.000	0.000	10.088

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	312	336	298	225	283	923	0	0	3048
N.S.	1	1.08	0.96	0.72	0.91	2.96	0.00	0.00	9.77
time (sec)	N/A	2.315	8.701	0.299	0.316	0.290	0.000	0.000	10.015

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	130	130	108	272	202	480	0	142	161
N.S.	1	1.00	0.83	2.09	1.55	3.69	0.00	1.09	1.24
time (sec)	N/A	0.694	1.457	0.087	0.318	0.329	0.000	0.638	12.583

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	105	105	89	251	186	432	0	112	130
N.S.	1	1.00	0.85	2.39	1.77	4.11	0.00	1.07	1.24
time (sec)	N/A	0.560	0.928	0.035	0.307	0.257	0.000	0.559	10.240

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	80	71	226	170	372	0	82	99
N.S.	1	1.00	0.89	2.82	2.12	4.65	0.00	1.02	1.24
time (sec)	N/A	0.445	0.542	0.033	0.332	0.254	0.000	0.460	9.755

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	55	55	201	151	314	0	47	68
N.S.	1	1.00	1.00	3.65	2.75	5.71	0.00	0.85	1.24
time (sec)	N/A	0.327	0.588	0.032	0.285	0.248	0.000	0.493	9.178

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	53	54	202	151	367	0	46	67
N.S.	1	1.00	1.02	3.81	2.85	6.92	0.00	0.87	1.26
time (sec)	N/A	0.354	0.536	0.037	0.306	0.262	0.000	0.619	8.139

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	78	58	220	171	427	0	69	99
N.S.	1	1.00	0.74	2.82	2.19	5.47	0.00	0.88	1.27
time (sec)	N/A	0.487	0.594	0.034	0.303	0.280	0.000	0.720	9.543

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	103	57	236	187	486	0	93	123
N.S.	1	1.00	0.55	2.29	1.82	4.72	0.00	0.90	1.19
time (sec)	N/A	0.611	0.808	0.037	0.316	0.277	0.000	0.802	9.973

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	183	186	130	298	230	555	0	194	326
N.S.	1	1.02	0.71	1.63	1.26	3.03	0.00	1.06	1.78
time (sec)	N/A	1.042	4.676	0.105	0.291	0.337	0.000	1.139	13.162

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	161	111	275	212	501	0	160	291
N.S.	1	1.03	0.71	1.76	1.36	3.21	0.00	1.03	1.87
time (sec)	N/A	0.856	2.134	0.037	0.288	0.289	0.000	0.964	10.807

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	129	134	93	252	194	441	0	126	256
N.S.	1	1.04	0.72	1.95	1.50	3.42	0.00	0.98	1.98
time (sec)	N/A	0.707	1.364	0.033	0.289	0.275	0.000	0.754	9.109

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	104	107	75	229	174	389	0	92	221
N.S.	1	1.03	0.72	2.20	1.67	3.74	0.00	0.88	2.12
time (sec)	N/A	0.583	1.973	0.031	0.283	0.251	0.000	0.756	8.849

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	74	217	170	386	0	69	203
N.S.	1	1.00	0.76	2.21	1.73	3.94	0.00	0.70	2.07
time (sec)	N/A	0.554	1.365	0.035	0.425	0.260	0.000	0.836	7.710

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	105	86	222	177	441	0	79	222
N.S.	1	1.03	0.84	2.18	1.74	4.32	0.00	0.77	2.18
time (sec)	N/A	0.574	1.582	0.041	0.394	0.263	0.000	0.895	8.384

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	127	135	102	240	195	502	0	107	258
N.S.	1	1.06	0.80	1.89	1.54	3.95	0.00	0.84	2.03
time (sec)	N/A	0.765	2.650	0.032	0.383	0.267	0.000	1.089	9.500

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	162	120	258	213	561	0	136	293
N.S.	1	1.05	0.78	1.68	1.38	3.64	0.00	0.88	1.90
time (sec)	N/A	0.921	3.474	0.037	0.398	0.282	0.000	1.142	10.918

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	198	206	129	299	234	561	0	194	327
N.S.	1	1.04	0.65	1.51	1.18	2.83	0.00	0.98	1.65
time (sec)	N/A	1.179	3.182	0.032	0.410	0.316	0.000	1.111	13.580

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	181	112	276	216	498	0	160	292
N.S.	1	1.06	0.65	1.61	1.26	2.91	0.00	0.94	1.71
time (sec)	N/A	0.995	2.046	0.042	0.407	0.274	0.000	0.910	10.439

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	146	154	92	253	196	447	0	127	257
N.S.	1	1.05	0.63	1.73	1.34	3.06	0.00	0.87	1.76
time (sec)	N/A	0.870	2.989	0.033	0.447	0.266	0.000	0.874	8.593

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	134	137	94	241	190	405	0	109	239
N.S.	1	1.02	0.70	1.80	1.42	3.02	0.00	0.81	1.78
time (sec)	N/A	0.819	1.888	0.040	0.439	0.265	0.000	1.057	8.630

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	136	140	94	234	191	438	0	97	240
N.S.	1	1.03	0.69	1.72	1.40	3.22	0.00	0.71	1.76
time (sec)	N/A	0.828	2.003	0.036	0.397	0.262	0.000	1.060	8.656

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	144	152	100	240	197	504	0	107	258
N.S.	1	1.06	0.69	1.67	1.37	3.50	0.00	0.74	1.79
time (sec)	N/A	0.858	2.872	0.045	0.411	0.254	0.000	1.320	9.096

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	182	118	258	215	561	0	136	293
N.S.	1	1.08	0.70	1.53	1.27	3.32	0.00	0.80	1.73
time (sec)	N/A	1.047	3.658	0.037	0.412	0.292	0.000	1.362	11.297

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	306	270	138	166	0	707	0	163	305
N.S.	1	0.88	0.45	0.54	0.00	2.31	0.00	0.53	1.00
time (sec)	N/A	0.923	2.465	0.075	0.000	0.281	0.000	0.533	13.767

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	275	237	114	143	0	621	0	122	270
N.S.	1	0.86	0.41	0.52	0.00	2.26	0.00	0.44	0.98
time (sec)	N/A	0.714	1.562	0.054	0.000	0.283	0.000	0.484	12.586

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	236	206	96	124	0	570	0	97	184
N.S.	1	0.87	0.41	0.53	0.00	2.42	0.00	0.41	0.78
time (sec)	N/A	0.589	1.121	0.128	0.000	0.267	0.000	0.423	10.226

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-2)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	234	204	97	121	0	571	0	98	184
N.S.	1	0.87	0.41	0.52	0.00	2.44	0.00	0.42	0.79
time (sec)	N/A	0.577	1.599	0.081	0.000	0.266	0.000	0.592	9.107

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F(-2)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	267	237	115	140	0	703	0	113	266
N.S.	1	0.89	0.43	0.52	0.00	2.63	0.00	0.42	1.00
time (sec)	N/A	0.729	1.383	0.054	0.000	0.270	0.000	0.771	9.159

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	296	260	118	160	0	795	0	141	303
N.S.	1	0.88	0.40	0.54	0.00	2.69	0.00	0.48	1.02
time (sec)	N/A	0.868	1.560	0.066	0.000	0.289	0.000	0.898	11.365

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	316	289	191	159	0	664	0	143	334
N.S.	1	0.91	0.60	0.50	0.00	2.10	0.00	0.45	1.06
time (sec)	N/A	1.046	3.466	0.072	0.000	0.266	0.000	0.769	12.037

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	277	244	178	147	0	664	0	125	318
N.S.	1	0.88	0.64	0.53	0.00	2.40	0.00	0.45	1.15
time (sec)	N/A	0.860	3.231	0.098	0.000	0.262	0.000	0.652	11.367

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	279	246	176	144	0	658	0	123	318
N.S.	1	0.88	0.63	0.52	0.00	2.36	0.00	0.44	1.14
time (sec)	N/A	0.835	2.805	0.101	0.000	0.280	0.000	0.560	11.295

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-2)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	285	243	178	144	0	664	0	126	318
N.S.	1	0.85	0.62	0.51	0.00	2.33	0.00	0.44	1.12
time (sec)	N/A	0.866	2.419	0.090	0.000	0.270	0.000	0.868	11.105

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F(-2)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	318	286	180	157	0	763	0	143	338
N.S.	1	0.90	0.57	0.49	0.00	2.40	0.00	0.45	1.06
time (sec)	N/A	1.056	2.037	0.063	0.000	0.278	0.000	1.223	11.928

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	347	315	180	176	0	861	0	162	373
N.S.	1	0.91	0.52	0.51	0.00	2.48	0.00	0.47	1.07
time (sec)	N/A	1.246	1.967	0.059	0.000	0.293	0.000	1.495	12.218

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	393	372	235	204	0	779	0	208	431
N.S.	1	0.95	0.60	0.52	0.00	1.98	0.00	0.53	1.10
time (sec)	N/A	1.666	6.173	0.071	0.000	0.296	0.000	1.020	11.597

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	364	339	222	181	0	685	0	167	395
N.S.	1	0.93	0.61	0.50	0.00	1.88	0.00	0.46	1.09
time (sec)	N/A	1.376	4.600	0.063	0.000	0.276	0.000	0.950	11.532

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	307	285	197	163	0	681	0	135	308
N.S.	1	0.93	0.64	0.53	0.00	2.22	0.00	0.44	1.00
time (sec)	N/A	1.146	3.764	0.078	0.000	0.265	0.000	0.853	9.994

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	309	287	183	152	0	635	0	131	239
N.S.	1	0.93	0.59	0.49	0.00	2.06	0.00	0.42	0.77
time (sec)	N/A	1.116	4.444	0.076	0.000	0.269	0.000	0.779	8.364

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	317	291	179	152	0	631	0	131	239
N.S.	1	0.92	0.56	0.48	0.00	1.99	0.00	0.41	0.75
time (sec)	N/A	1.165	5.014	0.091	0.000	0.260	0.000	0.680	7.867

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-2)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	315	288	168	157	0	682	0	137	308
N.S.	1	0.91	0.53	0.50	0.00	2.17	0.00	0.43	0.98
time (sec)	N/A	1.133	4.487	0.080	0.000	0.262	0.000	1.249	7.868

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F(-2)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	364	338	194	177	0	784	0	167	389
N.S.	1	0.93	0.53	0.49	0.00	2.15	0.00	0.46	1.07
time (sec)	N/A	1.372	3.030	0.059	0.000	0.283	0.000	1.683	8.400

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	393	367	194	196	0	875	0	182	425
N.S.	1	0.93	0.49	0.50	0.00	2.23	0.00	0.46	1.08
time (sec)	N/A	1.593	3.105	0.069	0.000	0.294	0.000	0.723	9.985

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	200	213	278	700	0	771	0	0	0
N.S.	1	1.06	1.39	3.50	0.00	3.86	0.00	0.00	0.00
time (sec)	N/A	1.215	7.010	0.263	0.000	0.262	0.000	0.000	0.000

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	152	158	189	577	0	673	0	0	2225
N.S.	1	1.04	1.24	3.80	0.00	4.43	0.00	0.00	14.64
time (sec)	N/A	0.881	5.549	0.158	0.000	0.270	0.000	0.000	28.492

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	112	112	141	469	0	539	0	0	372
N.S.	1	1.00	1.26	4.19	0.00	4.81	0.00	0.00	3.32
time (sec)	N/A	0.627	2.110	0.141	0.000	0.265	0.000	0.000	13.420

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	90	90	102	278	0	435	0	0	0
N.S.	1	1.00	1.13	3.09	0.00	4.83	0.00	0.00	0.00
time (sec)	N/A	0.415	0.952	0.157	0.000	0.252	0.000	0.000	0.000

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	135	142	131	354	0	483	0	0	0
N.S.	1	1.05	0.97	2.62	0.00	3.58	0.00	0.00	0.00
time (sec)	N/A	0.655	2.067	0.140	0.000	0.250	0.000	0.000	0.000

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	178	196	154	555	0	543	0	0	0
N.S.	1	1.10	0.87	3.12	0.00	3.05	0.00	0.00	0.00
time (sec)	N/A	0.967	6.372	0.138	0.000	0.268	0.000	0.000	0.000

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	221	250	282	630	0	593	0	0	0
N.S.	1	1.13	1.28	2.85	0.00	2.68	0.00	0.00	0.00
time (sec)	N/A	1.288	7.496	0.143	0.000	0.259	0.000	0.000	0.000

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	248	267	415	650	0	911	0	0	0
N.S.	1	1.08	1.67	2.62	0.00	3.67	0.00	0.00	0.00
time (sec)	N/A	1.632	7.169	0.170	0.000	0.289	0.000	0.000	0.000

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	204	212	283	563	0	829	0	0	0
N.S.	1	1.04	1.39	2.76	0.00	4.06	0.00	0.00	0.00
time (sec)	N/A	1.237	4.836	0.146	0.000	0.274	0.000	0.000	0.000

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	161	257	484	0	743	0	0	0
N.S.	1	1.03	1.65	3.10	0.00	4.76	0.00	0.00	0.00
time (sec)	N/A	0.901	3.560	0.151	0.000	0.281	0.000	0.000	0.000

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F(-1)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	146	146	336	519	0	748	0	0	0
N.S.	1	1.00	2.30	3.55	0.00	5.12	0.00	0.00	0.00
time (sec)	N/A	0.876	6.886	0.175	0.000	0.263	0.000	0.000	0.000

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	137	140	304	616	0	524	0	0	0
N.S.	1	1.02	2.22	4.50	0.00	3.82	0.00	0.00	0.00
time (sec)	N/A	0.694	7.008	0.130	0.000	0.258	0.000	0.000	0.000

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	181	194	360	705	0	593	0	0	0
N.S.	1	1.07	1.99	3.90	0.00	3.28	0.00	0.00	0.00
time (sec)	N/A	0.979	8.087	0.134	0.000	0.248	0.000	0.000	0.000

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	225	248	416	794	0	638	0	0	0
N.S.	1	1.10	1.85	3.53	0.00	2.84	0.00	0.00	0.00
time (sec)	N/A	1.341	7.984	0.139	0.000	0.275	0.000	0.000	0.000

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	269	302	470	883	0	701	0	0	0
N.S.	1	1.12	1.75	3.28	0.00	2.61	0.00	0.00	0.00
time (sec)	N/A	1.705	8.393	0.140	0.000	0.264	0.000	0.000	0.000

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	298	320	563	740	0	1017	0	0	0
N.S.	1	1.07	1.89	2.48	0.00	3.41	0.00	0.00	0.00
time (sec)	N/A	2.036	7.263	0.166	0.000	0.320	0.000	0.000	0.000

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	252	265	503	651	0	932	0	0	0
N.S.	1	1.05	2.00	2.58	0.00	3.70	0.00	0.00	0.00
time (sec)	N/A	1.572	6.784	0.153	0.000	0.273	0.000	0.000	0.000

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F(-1)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	206	212	433	564	0	849	0	0	0
N.S.	1	1.03	2.10	2.74	0.00	4.12	0.00	0.00	0.00
time (sec)	N/A	1.246	10.687	0.141	0.000	0.280	0.000	0.000	0.000

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F(-1)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	196	202	485	563	0	857	0	0	0
N.S.	1	1.03	2.47	2.87	0.00	4.37	0.00	0.00	0.00
time (sec)	N/A	1.237	7.221	0.156	0.000	0.282	0.000	0.000	0.000

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	190	190	554	618	0	846	0	0	0
N.S.	1	1.00	2.92	3.25	0.00	4.45	0.00	0.00	0.00
time (sec)	N/A	1.157	7.316	0.164	0.000	0.283	0.000	0.000	0.000

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F(-1)	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	185	193	447	707	0	608	0	0	0
N.S.	1	1.04	2.42	3.82	0.00	3.29	0.00	0.00	0.00
time (sec)	N/A	1.019	7.691	0.145	0.000	0.267	0.000	0.000	0.000

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F(-1)	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	231	247	503	796	0	657	0	0	0
N.S.	1	1.07	2.18	3.45	0.00	2.84	0.00	0.00	0.00
time (sec)	N/A	1.361	8.201	0.136	0.000	0.272	0.000	0.000	0.000

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F(-1)	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	277	301	559	885	0	722	0	0	0
N.S.	1	1.09	2.02	3.19	0.00	2.61	0.00	0.00	0.00
time (sec)	N/A	1.677	8.816	0.156	0.000	0.269	0.000	0.000	0.000

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	323	355	613	974	0	771	0	0	0
N.S.	1	1.10	1.90	3.02	0.00	2.39	0.00	0.00	0.00
time (sec)	N/A	2.079	9.712	0.152	0.000	0.270	0.000	0.000	0.000

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	190	190	566	618	0	900	0	0	0
N.S.	1	1.00	2.98	3.25	0.00	4.74	0.00	0.00	0.00
time (sec)	N/A	1.239	11.383	0.142	0.000	0.279	0.000	0.000	0.000

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	205	214	178	1135	0	807	0	0	0
N.S.	1	1.04	0.87	5.54	0.00	3.94	0.00	0.00	0.00
time (sec)	N/A	1.234	2.755	0.197	0.000	0.341	0.000	0.000	0.000

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	160	157	894	0	718	0	0	4040
N.S.	1	1.03	1.01	5.73	0.00	4.60	0.00	0.00	25.90
time (sec)	N/A	0.879	2.100	0.152	0.000	0.331	0.000	0.000	28.331

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	99	111	633	0	416	0	0	426
N.S.	1	1.00	1.12	6.39	0.00	4.20	0.00	0.00	4.30
time (sec)	N/A	0.423	1.368	0.174	0.000	0.268	0.000	0.000	14.095

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	143	145	128	695	0	473	0	0	0
N.S.	1	1.01	0.90	4.86	0.00	3.31	0.00	0.00	0.00
time (sec)	N/A	0.685	1.568	0.177	0.000	0.277	0.000	0.000	0.000

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	191	199	151	740	0	534	0	0	0
N.S.	1	1.04	0.79	3.87	0.00	2.80	0.00	0.00	0.00
time (sec)	N/A	0.983	3.312	0.177	0.000	0.278	0.000	0.000	0.000

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	237	253	172	815	0	596	0	0	0
N.S.	1	1.07	0.73	3.44	0.00	2.51	0.00	0.00	0.00
time (sec)	N/A	1.354	3.816	0.181	0.000	0.283	0.000	0.000	0.000

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	203	211	179	1215	0	758	0	0	0
N.S.	1	1.04	0.88	5.99	0.00	3.73	0.00	0.00	0.00
time (sec)	N/A	1.215	4.580	0.187	0.000	0.349	0.000	0.000	0.000

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	150	154	148	860	0	443	0	0	0
N.S.	1	1.03	0.99	5.73	0.00	2.95	0.00	0.00	0.00
time (sec)	N/A	0.738	2.907	0.159	0.000	0.280	0.000	0.000	0.000

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	148	151	152	860	0	441	0	0	0
N.S.	1	1.02	1.03	5.81	0.00	2.98	0.00	0.00	0.00
time (sec)	N/A	0.734	2.001	0.170	0.000	0.272	0.000	0.000	0.000

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	194	199	166	923	0	511	0	0	0
N.S.	1	1.03	0.86	4.76	0.00	2.63	0.00	0.00	0.00
time (sec)	N/A	1.049	2.668	0.142	0.000	0.275	0.000	0.000	0.000

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	240	253	185	1004	0	569	0	0	0
N.S.	1	1.05	0.77	4.18	0.00	2.37	0.00	0.00	0.00
time (sec)	N/A	1.385	4.314	0.168	0.000	0.293	0.000	0.000	0.000

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	249	264	296	1530	0	777	0	0	0
N.S.	1	1.06	1.19	6.14	0.00	3.12	0.00	0.00	0.00
time (sec)	N/A	1.592	6.083	0.172	0.000	0.339	0.000	0.000	0.000

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	194	206	224	1086	0	459	0	0	0
N.S.	1	1.06	1.15	5.60	0.00	2.37	0.00	0.00	0.00
time (sec)	N/A	1.058	3.698	0.138	0.000	0.275	0.000	0.000	0.000

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	196	207	222	1086	0	461	0	0	0
N.S.	1	1.06	1.13	5.54	0.00	2.35	0.00	0.00	0.00
time (sec)	N/A	1.071	2.996	0.144	0.000	0.289	0.000	0.000	0.000

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	194	205	242	1086	0	460	0	0	0
N.S.	1	1.06	1.25	5.60	0.00	2.37	0.00	0.00	0.00
time (sec)	N/A	1.052	4.700	0.154	0.000	0.288	0.000	0.000	0.000

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	240	253	216	1148	0	529	0	0	0
N.S.	1	1.05	0.90	4.78	0.00	2.20	0.00	0.00	0.00
time (sec)	N/A	1.391	4.301	0.174	0.000	0.289	0.000	0.000	0.000

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	286	307	259	1229	0	588	0	0	0
N.S.	1	1.07	0.91	4.30	0.00	2.06	0.00	0.00	0.00
time (sec)	N/A	1.748	7.774	0.170	0.000	0.286	0.000	0.000	0.000

Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	201	139	186	159	167	405	0	0	365
N.S.	1	0.69	0.93	0.79	0.83	2.01	0.00	0.00	1.82
time (sec)	N/A	0.395	0.814	0.066	0.387	0.241	0.000	0.000	1.253

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	270	216	196	223	210	660	0	0	436
N.S.	1	0.80	0.73	0.83	0.78	2.44	0.00	0.00	1.61
time (sec)	N/A	0.841	5.251	0.093	0.390	0.256	0.000	0.000	10.294

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	232	171	165	181	187	571	0	0	390
N.S.	1	0.74	0.71	0.78	0.81	2.46	0.00	0.00	1.68
time (sec)	N/A	0.544	1.252	0.055	0.362	0.255	0.000	0.000	9.572

Problem 199	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	202	141	138	159	168	489	0	0	367
N.S.	1	0.70	0.68	0.79	0.83	2.42	0.00	0.00	1.82
time (sec)	N/A	0.400	0.662	0.046	0.379	0.265	0.000	0.000	9.384

Problem 200	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	289	218	127	248	252	711	0	0	1761
N.S.	1	0.75	0.44	0.86	0.87	2.46	0.00	0.00	6.09
time (sec)	N/A	0.762	4.287	0.224	0.413	0.274	0.000	0.000	8.448

Problem 201	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	342	267	251	294	298	1096	0	0	5825
N.S.	1	0.78	0.73	0.86	0.87	3.20	0.00	0.00	17.03
time (sec)	N/A	1.042	3.477	0.209	0.365	0.301	0.000	0.000	9.987

Problem 202	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	150	150	166	172	547	0	0	383
N.S.	1	0.70	0.70	0.78	0.81	2.57	0.00	0.00	1.80
time (sec)	N/A	0.399	0.895	0.076	0.350	0.261	0.000	0.000	8.518

Problem 203	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	150	198	166	171	493	0	0	390
N.S.	1	0.70	0.93	0.78	0.80	2.31	0.00	0.00	1.83
time (sec)	N/A	0.417	0.788	0.075	0.352	0.248	0.000	0.000	8.424

Problem 204	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	290	315	173	0	0	0	0	0	0
N.S.	1	1.09	0.60	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.820	3.095	0.000	0.000	0.000	0.000	0.000	0.000

Problem 205	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	205	266	119	0	0	0	0	0	0
N.S.	1	1.30	0.58	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.262	2.554	0.000	0.000	0.000	0.000	0.000	0.000

Problem 206	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	132	164	81	0	0	0	0	0	0
N.S.	1	1.24	0.61	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.809	1.573	0.000	0.000	0.000	0.000	0.000	0.000

Problem 207	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	70	97	52	0	0	0	0	0	0
N.S.	1	1.39	0.74	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.404	0.524	0.000	0.000	0.000	0.000	0.000	0.000

Problem 208	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	168	164	137	0	0	0	0	0	0
N.S.	1	0.98	0.82	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.574	1.890	0.000	0.000	0.000	0.000	0.000	0.000

Problem 209	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	226	227	175	0	0	0	0	0	0
N.S.	1	1.00	0.77	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.965	3.218	0.000	0.000	0.000	0.000	0.000	0.000

Problem 210	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	308	319	227	0	0	0	0	0	0
N.S.	1	1.04	0.74	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.505	4.328	0.000	0.000	0.000	0.000	0.000	0.000

Problem 211	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	386	405	277	0	0	0	0	0	0
N.S.	1	1.05	0.72	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.092	3.557	0.000	0.000	0.000	0.000	0.000	0.000

Problem 212	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	316	326	0	0	0	0	0	0	0
N.S.	1	1.03	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.677	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 213	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	227	235	0	0	0	0	0	0	0
N.S.	1	1.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.143	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 214	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	159	159	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.686	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 215	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	214	222	0	0	0	0	0	0	0
N.S.	1	1.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.057	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 216	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	285	296	0	0	0	0	0	0	0
N.S.	1	1.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.549	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 217	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	363	381	0	0	0	0	0	0	0
N.S.	1	1.05	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.155	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 218	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	167	167	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.621	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 219	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	245	260	189	0	0	0	0	0	0
N.S.	1	1.06	0.77	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.276	4.142	0.000	0.000	0.000	0.000	0.000	0.000

Problem 220	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	164	172	137	0	0	0	0	0	0
N.S.	1	1.05	0.84	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.789	1.480	0.000	0.000	0.000	0.000	0.000	0.000

Problem 221	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	111	117	87	0	0	0	0	0	0
N.S.	1	1.05	0.78	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.455	0.809	0.000	0.000	0.000	0.000	0.000	0.000

Problem 222	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	78	84	60	0	0	0	0	0	0
N.S.	1	1.08	0.77	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.310	0.280	0.000	0.000	0.000	0.000	0.000	0.000

Problem 223	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	97	103	124	0	0	0	0	0	0
N.S.	1	1.06	1.28	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.512	1.224	0.000	0.000	0.000	0.000	0.000	0.000

Problem 224	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	131	144	156	0	0	0	0	0	0
N.S.	1	1.10	1.19	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.760	2.923	0.000	0.000	0.000	0.000	0.000	0.000

Problem 225	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	185	205	221	0	0	0	0	0	0
N.S.	1	1.11	1.19	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.144	3.167	0.000	0.000	0.000	0.000	0.000	0.000

Problem 226	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	383	421	0	0	0	0	0	0	0
N.S.	1	1.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.064	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 227	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	291	330	0	0	0	0	0	0	0
N.S.	1	1.13	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.459	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 228	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	215	247	0	0	0	0	0	0	0
N.S.	1	1.15	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.992	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 229	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	158	176	0	0	0	0	0	0	0
N.S.	1	1.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.670	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 230	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F(-1)	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	194	220	0	0	0	0	0	0	0
N.S.	1	1.13	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.945	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 231	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	247	280	0	0	0	0	0	0	0
N.S.	1	1.13	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.400	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 232	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	87	87	86	88	86	85	136	937	84
N.S.	1	1.00	0.99	1.01	0.99	0.98	1.56	10.77	0.97
time (sec)	N/A	0.507	0.623	0.053	0.373	0.245	0.119	0.846	7.281

Problem 233	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	65	67	69	66	66	104	556	63
N.S.	1	1.00	1.03	1.06	1.02	1.02	1.60	8.55	0.97
time (sec)	N/A	0.365	0.308	0.041	0.404	0.256	0.120	0.510	8.431

Problem 234	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	59	47	50	50	73	289	55
N.S.	1	1.00	1.40	1.12	1.19	1.19	1.74	6.88	1.31
time (sec)	N/A	0.250	0.035	0.025	0.374	0.252	0.083	0.394	7.178

Problem 235	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	39	49	47	52	59	78	53	69
N.S.	1	1.05	1.32	1.27	1.41	1.59	2.11	1.43	1.86
time (sec)	N/A	0.399	0.065	0.114	0.401	0.275	0.211	0.421	7.384

Problem 236	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	45	88	65	68	73	121	119	87
N.S.	1	1.05	2.05	1.51	1.58	1.70	2.81	2.77	2.02
time (sec)	N/A	0.374	0.046	0.123	0.375	0.252	0.322	0.538	8.479

Problem 237	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	66	68	77	89	86	95	148	179	108
N.S.	1	1.03	1.17	1.35	1.30	1.44	2.24	2.71	1.64
time (sec)	N/A	0.510	0.502	0.152	0.388	0.249	0.559	0.628	7.277

Problem 238	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	87	89	101	106	104	121	178	237	127
N.S.	1	1.02	1.16	1.22	1.20	1.39	2.05	2.72	1.46
time (sec)	N/A	0.642	1.095	0.184	0.349	0.260	0.848	0.756	7.537

Problem 239	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	108	110	100	124	122	138	209	299	145
N.S.	1	1.02	0.93	1.15	1.13	1.28	1.94	2.77	1.34
time (sec)	N/A	0.778	1.250	0.201	0.322	0.262	1.119	0.905	7.873

Problem 240	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	148	160	221	148	147	146	246	2078	151
N.S.	1	1.08	1.49	1.00	0.99	0.99	1.66	14.04	1.02
time (sec)	N/A	0.830	6.223	0.075	0.383	0.250	0.149	1.843	7.331

Problem 241	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	112	112	172	120	120	119	192	1389	121
N.S.	1	1.00	1.54	1.07	1.07	1.06	1.71	12.40	1.08
time (sec)	N/A	0.545	1.909	0.056	0.325	0.257	0.129	1.046	7.322

Problem 242	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	87	87	96	90	91	91	143	811	91
N.S.	1	1.00	1.10	1.03	1.05	1.05	1.64	9.32	1.05
time (sec)	N/A	0.406	0.477	0.040	0.380	0.244	0.104	0.696	8.347

Problem 243	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	72	93	80	85	92	129	86	90
N.S.	1	1.03	1.33	1.14	1.21	1.31	1.84	1.23	1.29
time (sec)	N/A	0.458	0.319	0.132	0.337	0.261	0.305	0.697	7.512

Problem 244	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	74	100	84	93	112	167	118	100
N.S.	1	1.03	1.39	1.17	1.29	1.56	2.32	1.64	1.39
time (sec)	N/A	0.466	0.297	0.201	0.324	0.286	0.458	0.874	7.532

Problem 245	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	91	123	107	120	122	214	237	127
N.S.	1	1.03	1.40	1.22	1.36	1.39	2.43	2.69	1.44
time (sec)	N/A	0.630	0.367	0.265	0.318	0.256	0.822	1.104	8.041

Problem 246	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	118	121	152	136	149	157	260	334	156
N.S.	1	1.03	1.29	1.15	1.26	1.33	2.20	2.83	1.32
time (sec)	N/A	0.800	1.253	0.250	0.344	0.261	1.128	1.431	7.955

Problem 247	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	151	152	180	162	175	191	313	435	182
N.S.	1	1.01	1.19	1.07	1.16	1.26	2.07	2.88	1.21
time (sec)	N/A	0.975	3.099	0.285	0.335	0.267	1.846	1.180	8.040

Problem 248	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	201	216	241	207	214	213	384	3757	217
N.S.	1	1.07	1.20	1.03	1.06	1.06	1.91	18.69	1.08
time (sec)	N/A	1.060	2.373	0.096	0.324	0.260	0.183	3.792	7.309

Problem 249	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	165	165	209	180	179	178	311	2670	181
N.S.	1	1.00	1.27	1.09	1.08	1.08	1.88	16.18	1.10
time (sec)	N/A	0.741	1.580	0.088	0.347	0.252	0.159	2.182	8.244

Problem 250	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	140	130	141	143	142	240	1751	142
N.S.	1	1.00	0.93	1.01	1.02	1.01	1.71	12.51	1.01
time (sec)	N/A	0.593	1.036	0.069	0.329	0.255	0.137	1.396	8.272

Problem 251	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	119	115	121	124	133	204	129	118
N.S.	1	1.02	0.98	1.03	1.06	1.14	1.74	1.10	1.01
time (sec)	N/A	0.733	0.626	0.214	0.365	0.281	0.458	1.133	8.484

Problem 252	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	119	121	113	118	125	145	223	152	114
N.S.	1	1.02	0.95	0.99	1.05	1.22	1.87	1.28	0.96
time (sec)	N/A	0.763	0.589	0.225	0.349	0.281	0.769	1.435	7.700

Problem 253	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	127	129	126	136	142	162	262	193	135
N.S.	1	1.02	0.99	1.07	1.12	1.28	2.06	1.52	1.06
time (sec)	N/A	0.767	0.454	0.214	0.295	0.275	1.081	1.403	8.152

Problem 254	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	161	164	172	180	181	332	390	169
N.S.	1	1.05	1.06	1.12	1.17	1.18	2.16	2.53	1.10
time (sec)	N/A	1.021	1.284	0.204	0.286	0.288	1.810	0.971	8.168

Problem 255	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	191	204	199	209	215	225	400	528	204
N.S.	1	1.07	1.04	1.09	1.13	1.18	2.09	2.76	1.07
time (sec)	N/A	1.291	0.783	0.245	0.298	0.273	2.315	1.093	7.831

Problem 256	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	233	242	237	243	250	266	471	670	238
N.S.	1	1.04	1.02	1.04	1.07	1.14	2.02	2.88	1.02
time (sec)	N/A	1.572	1.240	0.283	0.299	0.270	4.175	1.121	8.430

Problem 257	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	263	275	290	274	290	289	536	6042	300
N.S.	1	1.05	1.10	1.04	1.10	1.10	2.04	22.97	1.14
time (sec)	N/A	1.375	6.104	0.152	0.298	0.266	0.249	7.744	7.985

Problem 258	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	226	226	257	242	246	245	437	4489	251
N.S.	1	1.00	1.14	1.07	1.09	1.08	1.93	19.86	1.11
time (sec)	N/A	1.013	3.536	0.134	0.298	0.274	0.193	4.841	8.516

Problem 259	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	202	202	240	201	202	201	347	3133	205
N.S.	1	1.00	1.19	1.00	1.00	1.00	1.72	15.51	1.01
time (sec)	N/A	0.891	3.995	0.098	0.285	0.252	0.165	2.814	7.078

Problem 260	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	172	174	149	172	175	185	291	191	151
N.S.	1	1.01	0.87	1.00	1.02	1.08	1.69	1.11	0.88
time (sec)	N/A	1.076	1.521	0.230	0.294	0.270	0.755	1.743	7.466

Problem 261	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	175	177	134	159	164	193	289	195	142
N.S.	1	1.01	0.77	0.91	0.94	1.10	1.65	1.11	0.81
time (sec)	N/A	1.141	1.106	0.236	0.307	0.277	1.001	1.282	7.975

Problem 262	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	186	192	140	168	173	199	309	224	149
N.S.	1	1.03	0.75	0.90	0.93	1.07	1.66	1.20	0.80
time (sec)	N/A	1.199	0.704	0.234	0.321	0.270	1.791	1.388	7.400

Problem 263	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	187	189	167	195	202	222	369	281	177
N.S.	1	1.01	0.89	1.04	1.08	1.19	1.97	1.50	0.95
time (sec)	N/A	1.168	1.175	0.243	0.272	0.284	2.290	1.511	8.016

Problem 264	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	225	238	211	240	246	249	459	584	218
N.S.	1	1.06	0.94	1.07	1.09	1.11	2.04	2.60	0.97
time (sec)	N/A	1.551	0.988	0.294	0.277	0.266	4.169	1.680	8.512

Problem 265	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	273	288	257	284	289	300	546	763	263
N.S.	1	1.05	0.94	1.04	1.06	1.10	2.00	2.79	0.96
time (sec)	N/A	1.912	1.724	0.308	0.274	0.276	5.550	1.724	8.617

Problem 266	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	323	338	299	330	333	350	643	943	307
N.S.	1	1.05	0.93	1.02	1.03	1.08	1.99	2.92	0.95
time (sec)	N/A	2.295	1.376	0.370	0.272	0.266	12.579	1.855	8.105

Problem 267	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	127	144	138	127	130	190	1297	135	144
N.S.	1	1.13	1.09	1.00	1.02	1.50	10.21	1.06	1.13
time (sec)	N/A	0.930	1.562	0.135	0.276	0.287	0.953	0.646	7.613

Problem 268	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	108	118	101	109	149	1015	110	117
N.S.	1	1.07	1.17	1.00	1.08	1.48	10.05	1.09	1.16
time (sec)	N/A	0.634	0.660	0.089	0.292	0.284	0.688	0.495	7.844

Problem 269	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	75	98	87	94	110	700	95	100
N.S.	1	0.94	1.22	1.09	1.18	1.38	8.75	1.19	1.25
time (sec)	N/A	0.505	0.180	0.079	0.268	0.288	0.548	0.366	7.706

Problem 270	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	58	58	66	82	88	76	524	94	93
N.S.	1	1.00	1.14	1.41	1.52	1.31	9.03	1.62	1.60
time (sec)	N/A	0.332	0.110	0.047	0.267	0.275	0.468	0.348	7.797

Problem 271	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	82	113	95	107	118	952	113	115
N.S.	1	1.02	1.41	1.19	1.34	1.48	11.90	1.41	1.44
time (sec)	N/A	0.454	0.426	0.192	0.269	0.270	1.050	0.441	9.399

Problem 272	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	111	138	122	131	177	2071	157	140
N.S.	1	1.08	1.34	1.18	1.27	1.72	20.11	1.52	1.36
time (sec)	N/A	0.685	0.947	0.214	0.278	0.288	1.997	0.568	9.865

Problem 273	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	137	153	163	152	158	234	2599	214	175
N.S.	1	1.12	1.19	1.11	1.15	1.71	18.97	1.56	1.28
time (sec)	N/A	1.071	1.488	0.297	0.273	0.294	2.853	0.702	10.898

Problem 274	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	192	194	189	200	293	3016	285	208
N.S.	1	1.14	1.15	1.12	1.18	1.73	17.85	1.69	1.23
time (sec)	N/A	1.451	2.955	0.304	0.281	0.295	4.199	0.866	10.065

Problem 275	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	208	225	193	172	220	434	4534	290	210
N.S.	1	1.08	0.93	0.83	1.06	2.09	21.80	1.39	1.01
time (sec)	N/A	1.189	6.004	0.161	0.284	0.336	1.399	0.708	8.326

Problem 276	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	157	171	146	155	197	311	3485	244	165
N.S.	1	1.09	0.93	0.99	1.25	1.98	22.20	1.55	1.05
time (sec)	N/A	0.787	2.892	0.119	0.318	0.308	1.049	0.578	7.843

Problem 277	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	115	126	140	145	185	221	2987	241	163
N.S.	1	1.10	1.22	1.26	1.61	1.92	25.97	2.10	1.42
time (sec)	N/A	0.570	2.269	0.070	0.301	0.265	0.843	0.430	8.187

Problem 278	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	111	122	190	141	177	222	2878	234	153
N.S.	1	1.10	1.71	1.27	1.59	2.00	25.93	2.11	1.38
time (sec)	N/A	0.532	2.333	0.062	0.285	0.268	0.827	0.404	8.467

Problem 279	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	137	161	183	163	208	323	4488	279	180
N.S.	1	1.18	1.34	1.19	1.52	2.36	32.76	2.04	1.31
time (sec)	N/A	0.819	0.961	0.279	0.288	0.298	1.798	0.657	9.157

Problem 280	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	192	222	193	196	262	465	8145	362	230
N.S.	1	1.16	1.01	1.02	1.36	2.42	42.42	1.89	1.20
time (sec)	N/A	1.189	3.770	0.417	0.287	0.321	2.921	0.873	10.847

Problem 281	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	250	288	220	227	325	590	9896	402	284
N.S.	1	1.15	0.88	0.91	1.30	2.36	39.58	1.61	1.14
time (sec)	N/A	1.696	4.961	0.486	0.289	0.349	4.008	0.907	11.861

Problem 282	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-2)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	331	363	275	263	389	890	0	505	335
N.S.	1	1.10	0.83	0.79	1.18	2.69	0.00	1.53	1.01
time (sec)	N/A	1.835	5.220	0.196	0.299	0.380	0.000	1.116	9.001

Problem 283	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-2)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	250	285	235	242	366	666	0	458	307
N.S.	1	1.14	0.94	0.97	1.46	2.66	0.00	1.83	1.23
time (sec)	N/A	1.276	6.374	0.166	0.308	0.343	0.000	0.854	8.271

Problem 284	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	189	216	288	223	333	478	0	410	280
N.S.	1	1.14	1.52	1.18	1.76	2.53	0.00	2.17	1.48
time (sec)	N/A	0.984	5.746	0.110	0.297	0.275	0.000	0.704	7.825

Problem 285	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	179	201	188	213	330	488	0	410	282
N.S.	1	1.12	1.05	1.19	1.84	2.73	0.00	2.29	1.58
time (sec)	N/A	0.832	3.970	0.118	0.307	0.276	0.000	0.577	7.878

Problem 286	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	175	197	243	208	321	482	0	409	279
N.S.	1	1.13	1.39	1.19	1.83	2.75	0.00	2.34	1.59
time (sec)	N/A	0.807	4.042	0.110	0.305	0.272	0.000	0.538	8.251

Problem 287	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	215	257	254	243	372	683	0	479	315
N.S.	1	1.20	1.18	1.13	1.73	3.18	0.00	2.23	1.47
time (sec)	N/A	1.339	3.877	0.442	0.305	0.331	0.000	1.000	10.319

Problem 288	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-2)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	287	333	288	289	454	917	0	560	380
N.S.	1	1.16	1.00	1.01	1.58	3.20	0.00	1.95	1.32
time (sec)	N/A	1.776	6.459	0.724	0.386	0.371	0.000	0.920	13.087

Problem 289	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	352	406	320	320	541	1065	0	812	434
N.S.	1	1.15	0.91	0.91	1.54	3.03	0.00	2.31	1.23
time (sec)	N/A	2.377	6.687	0.706	0.398	0.398	0.000	1.239	14.255

Problem 290	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	351	402	307	343	583	1113	0	719	486
N.S.	1	1.15	0.87	0.98	1.66	3.17	0.00	2.05	1.38
time (sec)	N/A	1.865	3.560	0.275	0.387	0.357	0.000	1.260	8.805

Problem 291	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	298	346	465	318	550	813	0	670	470
N.S.	1	1.16	1.56	1.07	1.85	2.73	0.00	2.25	1.58
time (sec)	N/A	1.554	6.367	0.210	0.418	0.293	0.000	0.983	9.101

Problem 292	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	B	F(-2)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	261	300	411	301	526	836	0	632	446
N.S.	1	1.15	1.57	1.15	2.02	3.20	0.00	2.42	1.71
time (sec)	N/A	1.319	6.322	0.209	0.398	0.325	0.000	0.828	7.871

Problem 293	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	B	F(-2)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	250	283	248	287	523	838	0	638	447
N.S.	1	1.13	0.99	1.15	2.09	3.35	0.00	2.55	1.79
time (sec)	N/A	1.168	1.323	0.220	0.368	0.303	0.000	0.694	8.454

Problem 294	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	B	B	F(-2)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	247	280	327	284	514	815	0	630	442
N.S.	1	1.13	1.32	1.15	2.08	3.30	0.00	2.55	1.79
time (sec)	N/A	1.154	6.281	0.209	0.530	0.303	0.000	0.675	8.430

Problem 295	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	302	358	308	335	580	1126	0	722	484
N.S.	1	1.19	1.02	1.11	1.92	3.73	0.00	2.39	1.60
time (sec)	N/A	1.893	3.542	0.681	0.304	0.379	0.000	0.991	11.613

Problem 296	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-2)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	399	459	357	400	698	1510	0	846	576
N.S.	1	1.15	0.89	1.00	1.75	3.78	0.00	2.12	1.44
time (sec)	N/A	2.471	6.576	1.340	0.307	0.443	0.000	1.308	12.846

Problem 297	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-2)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	477	545	417	429	815	1732	0	903	664
N.S.	1	1.14	0.87	0.90	1.71	3.63	0.00	1.89	1.39
time (sec)	N/A	3.103	6.730	1.081	0.314	0.533	0.000	1.112	14.986

Problem 298	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	26	30	30	31	53	187	28
N.S.	1	1.00	0.90	1.03	1.03	1.07	1.83	6.45	0.97
time (sec)	N/A	0.256	0.024	0.043	0.390	0.253	0.574	0.569	7.506

Problem 299	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	25	17	22	19	36	22	16
N.S.	1	1.00	1.56	1.06	1.38	1.19	2.25	1.38	1.00
time (sec)	N/A	0.201	0.008	0.028	0.377	0.246	0.400	0.440	7.978

Problem 300	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	18	17	19	37	99	17
N.S.	1	1.00	1.00	1.38	1.31	1.46	2.85	7.62	1.31
time (sec)	N/A	0.183	0.005	0.025	0.339	0.261	0.333	0.329	7.228

Problem 301	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	A	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	10	3	2	10	3
N.S.	1	1.00	1.00	1.33	3.33	1.00	0.67	3.33	1.00
time (sec)	N/A	0.138	0.000	0.016	0.355	0.237	0.068	0.319	7.313

Problem 302	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	14	25	13	29	20	49	59	27
N.S.	1	1.17	2.08	1.08	2.42	1.67	4.08	4.92	2.25
time (sec)	N/A	0.185	0.016	0.170	0.349	0.269	0.500	0.333	7.006

Problem 303	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	30	22	23	42	37	39	16
N.S.	1	1.00	1.76	1.29	1.35	2.47	2.18	2.29	0.94
time (sec)	N/A	0.198	0.023	0.159	0.295	0.253	0.416	0.356	7.124

Problem 304	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	32	35	26	40	53	76	124	37
N.S.	1	1.07	1.17	0.87	1.33	1.77	2.53	4.13	1.23
time (sec)	N/A	0.259	0.092	0.169	0.296	0.250	0.915	0.385	7.272

Problem 305	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	29	34	27	38	90	49	69	32
N.S.	1	0.94	1.10	0.87	1.23	2.90	1.58	2.23	1.03
time (sec)	N/A	0.254	0.020	0.171	0.313	0.259	0.711	0.410	7.142

Problem 306	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	115	108	93	104	144	782	105	114
N.S.	1	1.13	1.06	0.91	1.02	1.41	7.67	1.03	1.12
time (sec)	N/A	0.752	0.394	0.096	0.285	0.265	1.473	0.894	8.162

Problem 307	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	83	89	92	80	89	119	660	90	98
N.S.	1	1.07	1.11	0.96	1.07	1.43	7.95	1.08	1.18
time (sec)	N/A	0.564	0.385	0.085	0.278	0.265	1.044	0.669	7.209

Problem 308	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	81	77	79	59	75	95	442	76	81
N.S.	1	0.95	0.98	0.73	0.93	1.17	5.46	0.94	1.00
time (sec)	N/A	0.483	0.044	0.065	0.279	0.274	0.741	0.506	7.363

Problem 309	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	48	48	67	51	71	65	282	76	79
N.S.	1	1.00	1.40	1.06	1.48	1.35	5.88	1.58	1.65
time (sec)	N/A	0.323	0.110	0.051	0.291	0.253	0.578	0.385	8.753

Problem 310	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	77	51	72	64	272	77	76
N.S.	1	1.00	1.64	1.09	1.53	1.36	5.79	1.64	1.62
time (sec)	N/A	0.302	0.025	0.049	0.291	0.254	0.574	0.377	8.077

Problem 311	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	70	79	80	88	104	672	92	99
N.S.	1	1.01	1.14	1.16	1.28	1.51	9.74	1.33	1.43
time (sec)	N/A	0.402	0.147	0.241	0.290	0.263	1.431	0.461	7.401

Problem 312	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	92	97	95	105	147	1137	122	113
N.S.	1	1.08	1.14	1.12	1.24	1.73	13.38	1.44	1.33
time (sec)	N/A	0.595	0.399	0.242	0.292	0.271	2.654	0.525	7.563

Problem 313	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	112	125	107	115	130	192	1401	165	143
N.S.	1	1.12	0.96	1.03	1.16	1.71	12.51	1.47	1.28
time (sec)	N/A	0.846	0.652	0.293	0.291	0.272	3.825	0.598	7.298

Problem 314	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	62	33	35	44	39	36	49
N.S.	1	1.00	2.48	1.32	1.40	1.76	1.56	1.44	1.96
time (sec)	N/A	0.273	0.058	0.042	0.286	0.249	0.089	0.345	7.683

Problem 315	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	58	58	65	85	95	78	235	99	112
N.S.	1	1.00	1.12	1.47	1.64	1.34	4.05	1.71	1.93
time (sec)	N/A	0.334	0.129	0.053	0.278	0.286	0.455	0.361	8.682

Problem 316	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	112	187	125	161	191	1348	199	152
N.S.	1	1.11	1.85	1.24	1.59	1.89	13.35	1.97	1.50
time (sec)	N/A	0.514	2.580	0.082	0.300	0.257	0.750	0.422	7.811

Problem 317	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	233	260	212	937	0	1312	0	0	1093
N.S.	1	1.12	0.91	4.02	0.00	5.63	0.00	0.00	4.69
time (sec)	N/A	1.462	2.966	0.398	0.000	0.275	0.000	0.000	66.519

Problem 318	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	186	180	169	873	0	1265	0	0	938
N.S.	1	0.97	0.91	4.69	0.00	6.80	0.00	0.00	5.04
time (sec)	N/A	1.006	2.089	0.128	0.000	0.272	0.000	0.000	25.434

Problem 319	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	146	145	140	829	0	1228	0	0	864
N.S.	1	0.99	0.96	5.68	0.00	8.41	0.00	0.00	5.92
time (sec)	N/A	0.733	0.551	0.109	0.000	0.292	0.000	0.000	14.084

Problem 320	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	122	103	120	812	0	1199	0	0	845
N.S.	1	0.84	0.98	6.66	0.00	9.83	0.00	0.00	6.93
time (sec)	N/A	0.560	0.141	0.094	0.000	0.328	0.000	0.000	10.813

Problem 321	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	131	118	219	976	0	2452	0	0	9785
N.S.	1	0.90	1.67	7.45	0.00	18.72	0.00	0.00	74.69
time (sec)	N/A	0.988	0.655	0.237	0.000	0.530	0.000	0.000	10.088

Problem 322	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	167	156	235	1029	0	2579	0	0	10987
N.S.	1	0.93	1.41	6.16	0.00	15.44	0.00	0.00	65.79
time (sec)	N/A	1.229	2.651	0.224	0.000	0.878	0.000	0.000	8.826

Problem 323	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	219	218	271	1145	0	2691	0	0	14195
N.S.	1	1.00	1.24	5.23	0.00	12.29	0.00	0.00	64.82
time (sec)	N/A	1.747	4.917	0.246	0.000	3.173	0.000	0.000	9.457

Problem 324	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	279	286	566	1295	0	2769	0	0	16796
N.S.	1	1.03	2.03	4.64	0.00	9.92	0.00	0.00	60.20
time (sec)	N/A	2.309	6.494	0.226	0.000	9.350	0.000	0.000	9.611

Problem 325	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	214	209	252	1697	0	3148	0	0	2993
N.S.	1	0.98	1.18	7.93	0.00	14.71	0.00	0.00	13.99
time (sec)	N/A	1.284	2.796	0.151	0.000	0.475	0.000	0.000	82.098

Problem 326	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	175	160	192	1668	0	3102	0	0	2868
N.S.	1	0.91	1.10	9.53	0.00	17.73	0.00	0.00	16.39
time (sec)	N/A	0.919	1.392	0.106	0.000	0.457	0.000	0.000	34.894

Problem 327	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	150	131	140	1657	0	3058	0	0	2823
N.S.	1	0.87	0.93	11.05	0.00	20.39	0.00	0.00	18.82
time (sec)	N/A	0.760	0.545	0.090	0.000	0.473	0.000	0.000	20.844

Problem 328	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	152	139	144	1654	0	6165	0	0	20255
N.S.	1	0.91	0.95	10.88	0.00	40.56	0.00	0.00	133.26
time (sec)	N/A	1.298	0.408	0.244	0.000	2.671	0.000	0.000	11.797

Problem 329	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	158	282	1689	0	6248	0	0	21319
N.S.	1	0.93	1.67	9.99	0.00	36.97	0.00	0.00	126.15
time (sec)	N/A	1.398	0.641	0.235	0.000	3.339	0.000	0.000	9.838

Problem 330	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	219	218	195	1805	0	6357	0	0	23016
N.S.	1	1.00	0.89	8.24	0.00	29.03	0.00	0.00	105.10
time (sec)	N/A	1.848	2.704	0.260	0.000	6.199	0.000	0.000	9.952

Problem 331	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	278	282	241	1985	0	6452	0	0	25789
N.S.	1	1.01	0.87	7.14	0.00	23.21	0.00	0.00	92.77
time (sec)	N/A	2.407	5.974	0.222	0.000	15.332	0.000	0.000	10.787

Problem 332	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	252	246	296	2438	0	4963	0	0	0
N.S.	1	0.98	1.17	9.67	0.00	19.69	0.00	0.00	0.00
time (sec)	N/A	1.612	4.966	0.177	0.000	0.864	0.000	0.000	0.000

Problem 333	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	216	258	2407	0	4916	0	0	3932
N.S.	1	1.01	1.21	11.30	0.00	23.08	0.00	0.00	18.46
time (sec)	N/A	1.158	1.735	0.117	0.000	0.838	0.000	0.000	94.320

Problem 334	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	188	169	233	2387	0	4855	0	0	3863
N.S.	1	0.90	1.24	12.70	0.00	25.82	0.00	0.00	20.55
time (sec)	N/A	0.989	1.166	0.108	0.000	0.820	0.000	0.000	41.525

Problem 335	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	182	169	177	2373	0	9757	0	0	29441
N.S.	1	0.93	0.97	13.04	0.00	53.61	0.00	0.00	161.76
time (sec)	N/A	1.643	1.263	0.253	0.000	10.913	0.000	0.000	14.543

Problem 336	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F(-1)	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	196	186	400	2390	0	9815	0	0	31186
N.S.	1	0.95	2.04	12.19	0.00	50.08	0.00	0.00	159.11
time (sec)	N/A	1.779	1.129	0.236	0.000	13.312	0.000	0.000	11.722

Problem 337	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F(-1)	B	F(-1)	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	220	214	448	2488	0	9888	0	0	32561
N.S.	1	0.97	2.04	11.31	0.00	44.95	0.00	0.00	148.00
time (sec)	N/A	1.839	2.552	0.266	0.000	16.382	0.000	0.000	12.572

Problem 338	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	277	276	548	2668	0	9965	0	0	33949
N.S.	1	1.00	1.98	9.63	0.00	35.97	0.00	0.00	122.56
time (sec)	N/A	2.457	6.488	0.249	0.000	25.113	0.000	0.000	11.684

Problem 339	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	342	357	622	2914	0	10107	0	0	36736
N.S.	1	1.04	1.82	8.52	0.00	29.55	0.00	0.00	107.42
time (sec)	N/A	3.042	6.578	0.296	0.000	55.784	0.000	0.000	12.432

Problem 340	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	151	121	193	1375	0	1472	0	0	3441
N.S.	1	0.80	1.28	9.11	0.00	9.75	0.00	0.00	22.79
time (sec)	N/A	0.686	1.421	0.079	0.000	0.290	0.000	0.000	32.674

Problem 341	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	408	432	183	986	0	956	0	0	2529
N.S.	1	1.06	0.45	2.42	0.00	2.34	0.00	0.00	6.20
time (sec)	N/A	0.718	0.506	0.086	0.000	0.285	0.000	0.000	19.901

Problem 342	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	422	453	157	814	0	719	0	0	581
N.S.	1	1.07	0.37	1.93	0.00	1.70	0.00	0.00	1.38
time (sec)	N/A	0.731	0.260	0.073	0.000	0.269	0.000	0.000	10.209

Problem 343	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	221	170	2015	0	1767	0	0	3054
N.S.	1	1.04	0.80	9.46	0.00	8.30	0.00	0.00	14.34
time (sec)	N/A	1.193	4.341	0.181	0.000	0.297	0.000	0.000	16.395

Problem 344	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	166	157	139	1949	0	1725	0	0	2981
N.S.	1	0.95	0.84	11.74	0.00	10.39	0.00	0.00	17.96
time (sec)	N/A	0.828	1.687	0.125	0.000	0.306	0.000	0.000	11.913

Problem 345	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	124	109	118	1910	0	1706	0	0	2930
N.S.	1	0.88	0.95	15.40	0.00	13.76	0.00	0.00	23.63
time (sec)	N/A	0.567	0.558	0.120	0.000	0.304	0.000	0.000	11.279

Problem 346	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	83	101	1892	0	1669	0	0	2909
N.S.	1	0.81	0.99	18.55	0.00	16.36	0.00	0.00	28.52
time (sec)	N/A	0.438	0.121	0.504	0.000	0.279	0.000	0.000	10.167

Problem 347	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	131	118	170	4002	0	3458	0	0	7099
N.S.	1	0.90	1.30	30.55	0.00	26.40	0.00	0.00	54.19
time (sec)	N/A	0.940	0.812	0.279	0.000	0.815	0.000	0.000	12.117

Problem 348	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	164	201	4057	0	3582	0	0	9790
N.S.	1	0.97	1.19	24.01	0.00	21.20	0.00	0.00	57.93
time (sec)	N/A	1.272	3.144	0.225	0.000	1.994	0.000	0.000	9.957

Problem 349	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	224	230	360	4175	0	3688	0	0	13182
N.S.	1	1.03	1.61	18.64	0.00	16.46	0.00	0.00	58.85
time (sec)	N/A	1.701	6.319	0.236	0.000	7.510	0.000	0.000	10.907

Problem 350	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	264	273	300	3754	0	4421	0	0	5811
N.S.	1	1.03	1.14	14.22	0.00	16.75	0.00	0.00	22.01
time (sec)	N/A	1.380	3.795	0.183	0.000	0.709	0.000	0.000	23.823

Problem 351	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	167	186	248	3713	0	4379	0	0	5768
N.S.	1	1.11	1.49	22.23	0.00	26.22	0.00	0.00	34.54
time (sec)	N/A	0.938	1.408	0.176	0.000	0.697	0.000	0.000	16.124

Problem 352	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	141	151	229	3688	0	4329	0	0	5742
N.S.	1	1.07	1.62	26.16	0.00	30.70	0.00	0.00	40.72
time (sec)	N/A	0.640	1.524	0.103	0.000	0.698	0.000	0.000	15.338

Problem 353	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	144	113	3684	0	4318	0	0	5737
N.S.	1	1.04	0.82	26.70	0.00	31.29	0.00	0.00	41.57
time (sec)	N/A	0.605	0.230	0.092	0.000	0.687	0.000	0.000	14.920

Problem 354	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	195	186	7982	0	8820	0	0	26139
N.S.	1	1.14	1.09	46.68	0.00	51.58	0.00	0.00	152.86
time (sec)	N/A	1.358	1.336	0.233	0.000	4.109	0.000	0.000	15.353

Problem 355	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	219	253	208	8043	0	9075	0	0	38368
N.S.	1	1.16	0.95	36.73	0.00	41.44	0.00	0.00	175.20
time (sec)	N/A	1.815	4.168	0.240	0.000	17.367	0.000	0.000	12.654

Problem 356	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	285	324	409	8164	0	9219	0	0	42371
N.S.	1	1.14	1.44	28.65	0.00	32.35	0.00	0.00	148.67
time (sec)	N/A	2.352	6.279	0.245	0.000	42.423	0.000	0.000	12.504

Problem 357	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	371	395	450	4592	0	7358	0	0	9547
N.S.	1	1.06	1.21	12.38	0.00	19.83	0.00	0.00	25.73
time (sec)	N/A	2.169	6.389	0.198	0.000	2.204	0.000	0.000	47.632

Problem 358	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	261	298	309	4548	0	7284	0	0	9498
N.S.	1	1.14	1.18	17.43	0.00	27.91	0.00	0.00	36.39
time (sec)	N/A	1.446	3.638	0.143	0.000	2.204	0.000	0.000	37.094

Problem 359	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	198	225	260	4492	0	7218	0	0	9468
N.S.	1	1.14	1.31	22.69	0.00	36.45	0.00	0.00	47.82
time (sec)	N/A	1.075	1.155	0.117	0.000	2.235	0.000	0.000	27.682

Problem 360	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	188	213	325	4484	0	7209	0	0	9464
N.S.	1	1.13	1.73	23.85	0.00	38.35	0.00	0.00	50.34
time (sec)	N/A	0.925	3.868	0.127	0.000	2.258	0.000	0.000	26.558

Problem 361	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	185	206	115	4473	0	7181	0	0	9457
N.S.	1	1.11	0.62	24.18	0.00	38.82	0.00	0.00	51.12
time (sec)	N/A	0.928	0.267	0.110	0.000	2.214	0.000	0.000	27.143

Problem 362	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	224	272	242	12870	0	14700	0	0	45681
N.S.	1	1.21	1.08	57.46	0.00	65.62	0.00	0.00	203.93
time (sec)	N/A	1.866	5.470	0.228	0.000	19.726	0.000	0.000	18.988

Problem 363	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	289	343	306	12968	0	15082	0	0	67465
N.S.	1	1.19	1.06	44.87	0.00	52.19	0.00	0.00	233.44
time (sec)	N/A	2.375	5.430	0.259	0.000	57.597	0.000	0.000	15.511

Problem 364	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	364	423	593	13092	0	15325	0	0	71314
N.S.	1	1.16	1.63	35.97	0.00	42.10	0.00	0.00	195.92
time (sec)	N/A	3.081	6.363	0.274	0.000	111.451	0.000	0.000	14.666

Problem 365	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	A	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	362	403	88	662	0	399	0	0	3033
N.S.	1	1.11	0.24	1.83	0.00	1.10	0.00	0.00	8.38
time (sec)	N/A	0.613	0.082	0.120	0.000	0.257	0.000	0.000	9.979

Problem 366	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	406	426	88	1575	0	805	0	0	6453
N.S.	1	1.05	0.22	3.88	0.00	1.98	0.00	0.00	15.89
time (sec)	N/A	0.658	0.035	0.118	0.000	0.263	0.000	0.000	12.469

Problem 367	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	119	108	112	374	0	1674	0	0	2142
N.S.	1	0.91	0.94	3.14	0.00	14.07	0.00	0.00	18.00
time (sec)	N/A	0.879	0.091	0.343	0.000	0.291	0.000	0.000	10.113

Problem 368	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	123	125	106	1955	0	2174	0	0	9618
N.S.	1	1.02	0.86	15.89	0.00	17.67	0.00	0.00	78.20
time (sec)	N/A	0.595	0.139	0.088	0.000	0.305	0.000	0.000	21.213

Problem 369	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	176	166	1779	0	4578	0	0	7172
N.S.	1	1.14	1.08	11.55	0.00	29.73	0.00	0.00	46.57
time (sec)	N/A	1.244	1.248	0.330	0.000	0.390	0.000	0.000	14.086

Problem 370	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	85	109	1893	0	1115	0	0	2731
N.S.	1	0.83	1.07	18.56	0.00	10.93	0.00	0.00	26.77
time (sec)	N/A	0.413	0.203	0.090	0.000	0.263	0.000	0.000	10.321

Problem 371	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	132	129	154	2291	0	2320	0	0	5475
N.S.	1	0.98	1.17	17.36	0.00	17.58	0.00	0.00	41.48
time (sec)	N/A	0.608	0.432	0.117	0.000	0.295	0.000	0.000	13.948

Problem 372	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	B	F	F(-1)	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	174	182	156	3055	0	3695	0	0	8437
N.S.	1	1.05	0.90	17.56	0.00	21.24	0.00	0.00	48.49
time (sec)	N/A	0.858	0.313	0.100	0.000	0.344	0.000	0.000	24.858

Problem 373	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	35	45	729	6480	249	0	155	1410
N.S.	1	0.78	1.00	16.20	144.00	5.53	0.00	3.44	31.33
time (sec)	N/A	0.236	0.438	0.105	0.636	0.244	0.000	0.407	10.392

Problem 374	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	35	45	741	0	267	0	155	1410
N.S.	1	0.78	1.00	16.47	0.00	5.93	0.00	3.44	31.33
time (sec)	N/A	0.243	0.044	0.092	0.000	0.242	0.000	0.402	9.215

Problem 375	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	61	54	0	28	0	65	31
N.S.	1	1.00	2.03	1.80	0.00	0.93	0.00	2.17	1.03
time (sec)	N/A	0.203	0.767	0.225	0.000	0.248	0.000	0.329	0.860

Problem 376	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	61	52	0	47	0	57	35
N.S.	1	1.00	2.26	1.93	0.00	1.74	0.00	2.11	1.30
time (sec)	N/A	0.205	0.926	0.048	0.000	0.250	0.000	0.301	0.691

Problem 377	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	85	75	134	0	309	0	0	147
N.S.	1	1.00	0.88	1.58	0.00	3.64	0.00	0.00	1.73
time (sec)	N/A	0.381	0.106	0.231	0.000	0.254	0.000	0.000	8.935

Problem 378	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	278	255	151	275	227	2266	0	0	1522
N.S.	1	0.92	0.54	0.99	0.82	8.15	0.00	0.00	5.47
time (sec)	N/A	0.909	1.742	0.065	0.368	0.389	0.000	0.000	18.211

Problem 379	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	254	231	134	253	210	2252	0	0	1492
N.S.	1	0.91	0.53	1.00	0.83	8.87	0.00	0.00	5.87
time (sec)	N/A	0.748	1.104	0.037	0.388	0.381	0.000	0.000	14.410

Problem 380	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	229	206	114	229	192	2230	0	0	1456
N.S.	1	0.90	0.50	1.00	0.84	9.74	0.00	0.00	6.36
time (sec)	N/A	0.605	0.447	0.032	0.347	0.384	0.000	0.000	12.018

Problem 381	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	205	182	94	203	174	2216	0	0	1420
N.S.	1	0.89	0.46	0.99	0.85	10.81	0.00	0.00	6.93
time (sec)	N/A	0.494	0.178	0.037	0.373	0.380	0.000	0.000	10.383

Problem 382	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	205	182	158	203	174	2244	0	0	1420
N.S.	1	0.89	0.77	0.99	0.85	10.95	0.00	0.00	6.93
time (sec)	N/A	0.529	0.683	0.041	0.314	0.387	0.000	0.000	10.790

Problem 383	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	229	206	178	222	192	2275	0	0	1448
N.S.	1	0.90	0.78	0.97	0.84	9.93	0.00	0.00	6.32
time (sec)	N/A	0.652	0.830	0.039	0.399	0.414	0.000	0.000	11.751

Problem 384	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	254	231	198	240	211	2290	0	0	1473
N.S.	1	0.91	0.78	0.94	0.83	9.02	0.00	0.00	5.80
time (sec)	N/A	0.807	1.325	0.040	0.472	0.399	0.000	0.000	14.246

Problem 385	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	394	349	304	374	329	4337	0	0	3914
N.S.	1	0.89	0.77	0.95	0.84	11.01	0.00	0.00	9.93
time (sec)	N/A	1.405	6.145	0.102	0.466	0.749	0.000	0.000	29.534

Problem 386	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	360	317	178	333	302	4326	0	0	3869
N.S.	1	0.88	0.49	0.92	0.84	12.02	0.00	0.00	10.75
time (sec)	N/A	1.186	2.367	0.034	0.715	0.712	0.000	0.000	21.128

Problem 387	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	326	283	151	292	275	4283	0	0	3825
N.S.	1	0.87	0.46	0.90	0.84	13.14	0.00	0.00	11.73
time (sec)	N/A	0.967	1.314	0.035	0.372	0.731	0.000	0.000	15.104

Problem 388	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	294	249	119	251	248	4271	0	0	3773
N.S.	1	0.85	0.40	0.85	0.84	14.53	0.00	0.00	12.83
time (sec)	N/A	0.803	0.577	0.035	0.398	0.716	0.000	0.000	11.186

Problem 389	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	276	228	211	238	240	4280	0	0	3749
N.S.	1	0.83	0.76	0.86	0.87	15.51	0.00	0.00	13.58
time (sec)	N/A	0.738	0.988	0.035	0.370	0.739	0.000	0.000	10.687

Problem 390	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	283	235	119	243	248	4313	0	0	3745
N.S.	1	0.83	0.42	0.86	0.88	15.24	0.00	0.00	13.23
time (sec)	N/A	0.754	0.740	0.033	0.354	0.722	0.000	0.000	11.813

Problem 391	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	317	269	120	270	276	4325	0	0	3782
N.S.	1	0.85	0.38	0.85	0.87	13.64	0.00	0.00	11.93
time (sec)	N/A	0.920	0.661	0.033	0.366	0.729	0.000	0.000	15.107

Problem 392	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	463	395	221	432	398	6209	0	0	6774
N.S.	1	0.85	0.48	0.93	0.86	13.41	0.00	0.00	14.63
time (sec)	N/A	1.648	3.721	0.039	0.356	1.396	0.000	0.000	38.607

Problem 393	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	421	355	197	372	363	6157	0	0	6716
N.S.	1	0.84	0.47	0.88	0.86	14.62	0.00	0.00	15.95
time (sec)	N/A	1.379	2.178	0.035	0.331	1.437	0.000	0.000	25.335

Problem 394	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	380	312	153	314	327	6136	0	0	6657
N.S.	1	0.82	0.40	0.83	0.86	16.15	0.00	0.00	17.52
time (sec)	N/A	1.149	1.585	0.034	0.341	1.491	0.000	0.000	14.527

Problem 395	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	374	303	264	286	310	6137	0	0	7108
N.S.	1	0.81	0.71	0.76	0.83	16.41	0.00	0.00	19.01
time (sec)	N/A	1.135	2.860	0.038	0.329	1.466	0.000	0.000	12.194

Problem 396	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	372	302	165	278	310	6162	0	0	7578
N.S.	1	0.81	0.44	0.75	0.83	16.56	0.00	0.00	20.37
time (sec)	N/A	1.136	1.326	0.038	0.312	1.438	0.000	0.000	12.366

Problem 397	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	380	312	166	290	327	6162	0	0	7591
N.S.	1	0.82	0.44	0.76	0.86	16.22	0.00	0.00	19.98
time (sec)	N/A	1.209	1.436	0.040	0.327	1.450	0.000	0.000	14.602

Problem 398	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	325	289	187	292	258	6160	0	0	16441
N.S.	1	0.89	0.58	0.90	0.79	18.95	0.00	0.00	50.59
time (sec)	N/A	1.446	1.233	0.099	0.332	21.871	0.000	0.000	14.940

Problem 399	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	297	254	165	265	234	6092	0	0	15701
N.S.	1	0.86	0.56	0.89	0.79	20.51	0.00	0.00	52.87
time (sec)	N/A	1.092	0.372	0.067	0.309	9.278	0.000	0.000	15.087

Problem 400	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	278	228	195	244	216	5992	0	0	15090
N.S.	1	0.82	0.70	0.88	0.78	21.55	0.00	0.00	54.28
time (sec)	N/A	0.796	0.400	0.063	0.322	3.211	0.000	0.000	13.347

Problem 401	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	278	228	194	244	217	6024	0	0	14816
N.S.	1	0.82	0.70	0.88	0.78	21.67	0.00	0.00	53.29
time (sec)	N/A	0.814	0.367	0.055	0.326	4.612	0.000	0.000	13.411

Problem 402	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	297	254	153	265	235	6136	0	0	15318
N.S.	1	0.86	0.52	0.89	0.79	20.66	0.00	0.00	51.58
time (sec)	N/A	1.079	0.631	0.049	0.324	12.056	0.000	0.000	13.622

Problem 403	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	325	289	174	286	256	6284	0	0	16111
N.S.	1	0.89	0.54	0.88	0.79	19.34	0.00	0.00	49.57
time (sec)	N/A	1.450	3.628	0.050	0.322	27.010	0.000	0.000	15.152

Problem 404	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	436	379	275	351	377	11910	0	0	18313
N.S.	1	0.87	0.63	0.81	0.86	27.32	0.00	0.00	42.00
time (sec)	N/A	1.827	2.486	0.080	0.317	36.196	0.000	0.000	41.994

Problem 405	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	391	335	230	336	354	11824	0	0	17579
N.S.	1	0.86	0.59	0.86	0.91	30.24	0.00	0.00	44.96
time (sec)	N/A	1.382	2.009	0.053	0.352	21.054	0.000	0.000	38.269

Problem 406	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	391	334	220	332	341	11827	0	0	17089
N.S.	1	0.85	0.56	0.85	0.87	30.25	0.00	0.00	43.71
time (sec)	N/A	1.331	1.519	0.047	0.366	15.760	0.000	0.000	38.279

Problem 407	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	391	335	204	336	356	11831	0	0	17494
N.S.	1	0.86	0.52	0.86	0.91	30.26	0.00	0.00	44.74
time (sec)	N/A	1.402	1.220	0.049	0.334	25.292	0.000	0.000	39.001

Problem 408	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	439	381	239	352	396	12008	0	0	22667
N.S.	1	0.87	0.54	0.80	0.90	27.35	0.00	0.00	51.63
time (sec)	N/A	1.912	2.509	0.051	0.365	39.862	0.000	0.000	29.801

Problem 409	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	493	433	287	373	449	12216	0	0	24620
N.S.	1	0.88	0.58	0.76	0.91	24.78	0.00	0.00	49.94
time (sec)	N/A	2.502	4.194	0.058	0.345	75.099	0.000	0.000	27.533

Problem 410	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	600	528	1351	466	571	17991	0	0	27429
N.S.	1	0.88	2.25	0.78	0.95	29.98	0.00	0.00	45.72
time (sec)	N/A	2.745	6.450	0.156	0.366	166.932	0.000	0.000	71.273

Problem 411	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	534	466	690	451	551	17852	0	0	26614
N.S.	1	0.87	1.29	0.84	1.03	33.43	0.00	0.00	49.84
time (sec)	N/A	2.061	6.419	0.136	0.442	108.208	0.000	0.000	61.366

Problem 412	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	533	462	333	450	538	17891	0	0	25944
N.S.	1	0.87	0.62	0.84	1.01	33.57	0.00	0.00	48.68
time (sec)	N/A	2.038	6.306	0.046	0.415	75.164	0.000	0.000	62.636

Problem 413	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	531	464	552	451	537	17893	0	0	26133
N.S.	1	0.87	1.04	0.85	1.01	33.70	0.00	0.00	49.21
time (sec)	N/A	2.081	6.333	0.051	0.402	81.209	0.000	0.000	61.165

Problem 414	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	534	466	288	452	551	17859	0	0	26707
N.S.	1	0.87	0.54	0.85	1.03	33.44	0.00	0.00	50.01
time (sec)	N/A	2.205	4.940	0.058	0.401	118.654	0.000	0.000	63.373

Problem 415	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	601	528	585	466	607	18091	0	0	35300
N.S.	1	0.88	0.97	0.78	1.01	30.10	0.00	0.00	58.74
time (sec)	N/A	3.013	6.346	0.060	0.360	189.650	0.000	0.000	57.044

Problem 416	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	157	89	102	123	184	0	0	16727
N.S.	1	1.01	0.57	0.65	0.79	1.18	0.00	0.00	107.22
time (sec)	N/A	0.411	0.174	0.033	0.351	0.254	0.000	0.000	13.956

Problem 417	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	155	139	102	123	167	0	0	16060
N.S.	1	1.01	0.90	0.66	0.80	1.08	0.00	0.00	104.29
time (sec)	N/A	0.408	0.161	0.035	0.374	0.249	0.000	0.000	13.829

Problem 418	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	138	60	90	109	164	0	0	15753
N.S.	1	1.00	0.43	0.65	0.79	1.19	0.00	0.00	114.15
time (sec)	N/A	0.338	0.056	0.060	0.356	0.249	0.000	0.000	13.378

Problem 419	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	138	110	90	112	148	0	0	15437
N.S.	1	1.00	0.80	0.65	0.81	1.07	0.00	0.00	111.86
time (sec)	N/A	0.339	0.038	0.057	0.339	0.249	0.000	0.000	13.812

Problem 420	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	155	78	102	122	215	0	0	15569
N.S.	1	1.01	0.51	0.66	0.79	1.40	0.00	0.00	101.10
time (sec)	N/A	0.408	0.145	0.038	0.332	0.252	0.000	0.000	13.340

Problem 421	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	157	82	102	124	208	0	0	16545
N.S.	1	1.01	0.53	0.65	0.79	1.33	0.00	0.00	106.06
time (sec)	N/A	0.412	0.148	0.032	0.342	0.257	0.000	0.000	13.566

Problem 422	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	256	228	156	242	188	3238	0	0	18514
N.S.	1	0.89	0.61	0.95	0.73	12.65	0.00	0.00	72.32
time (sec)	N/A	1.015	0.171	0.051	0.337	0.340	0.000	0.000	42.437

Problem 423	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	237	203	228	226	173	3156	0	0	16878
N.S.	1	0.86	0.96	0.95	0.73	13.32	0.00	0.00	71.22
time (sec)	N/A	0.759	0.181	0.049	0.326	0.324	0.000	0.000	39.454

Problem 424	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	237	203	205	225	172	3160	0	0	17323
N.S.	1	0.86	0.86	0.95	0.73	13.33	0.00	0.00	73.09
time (sec)	N/A	0.734	0.095	0.048	0.326	0.322	0.000	0.000	37.529

Problem 425	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	237	203	229	226	172	3156	0	0	16598
N.S.	1	0.86	0.97	0.95	0.73	13.32	0.00	0.00	70.03
time (sec)	N/A	0.745	0.122	0.044	0.394	0.335	0.000	0.000	37.134

Problem 426	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	256	228	132	242	188	3314	0	0	22906
N.S.	1	0.89	0.52	0.95	0.73	12.95	0.00	0.00	89.48
time (sec)	N/A	0.977	0.516	0.040	0.333	0.343	0.000	0.000	28.978

Problem 427	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	264	265	304	2183234	0	16118	0	0	0
N.S.	1	1.00	1.15	8269.83	0.00	61.05	0.00	0.00	0.00
time (sec)	N/A	1.415	3.732	1.761	0.000	3.479	0.000	0.000	0.000

Problem 428	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	201	199	239	2180698	0	15912	0	0	61200
N.S.	1	0.99	1.19	10849.24	0.00	79.16	0.00	0.00	304.48
time (sec)	N/A	0.952	2.975	1.429	0.000	3.371	0.000	0.000	132.153

Problem 429	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	169	179	204	2178123	0	15940	0	0	1141
N.S.	1	1.06	1.21	12888.30	0.00	94.32	0.00	0.00	6.75
time (sec)	N/A	1.183	1.103	2.285	0.000	2.376	0.000	0.000	20.288

Problem 430	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	154	170	169	2176213	0	7865	0	0	0
N.S.	1	1.10	1.10	14131.25	0.00	51.07	0.00	0.00	0.00
time (sec)	N/A	0.851	1.118	1.061	0.000	1.410	0.000	0.000	0.000

Problem 431	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	199	222	194	2181119	0	7946	0	0	0
N.S.	1	1.12	0.97	10960.40	0.00	39.93	0.00	0.00	0.00
time (sec)	N/A	1.193	1.745	0.827	0.000	1.453	0.000	0.000	0.000

Problem 432	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	250	289	226	2183172	0	7964	0	0	0
N.S.	1	1.16	0.90	8732.69	0.00	31.86	0.00	0.00	0.00
time (sec)	N/A	1.760	3.432	1.404	0.000	1.420	0.000	0.000	0.000

Problem 433	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	314	355	265	2185304	0	8039	0	0	0
N.S.	1	1.13	0.84	6959.57	0.00	25.60	0.00	0.00	0.00
time (sec)	N/A	2.272	4.425	1.618	0.000	1.418	0.000	0.000	0.000

Problem 434	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	323	326	347	2403184	0	24426	0	0	0
N.S.	1	1.01	1.07	7440.20	0.00	75.62	0.00	0.00	0.00
time (sec)	N/A	1.965	4.650	0.898	0.000	5.198	0.000	0.000	0.000

Problem 435	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	268	263	290	2400957	0	24274	0	0	0
N.S.	1	0.98	1.08	8958.79	0.00	90.57	0.00	0.00	0.00
time (sec)	N/A	1.548	2.699	0.896	0.000	4.736	0.000	0.000	0.000

Problem 436	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	204	200	243	2396039	0	24220	0	0	0
N.S.	1	0.98	1.19	11745.29	0.00	118.73	0.00	0.00	0.00
time (sec)	N/A	1.100	1.111	0.954	0.000	4.766	0.000	0.000	0.000

Problem 437	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	209	200	416	2394883	0	24297	0	0	0
N.S.	1	0.96	1.99	11458.77	0.00	116.25	0.00	0.00	0.00
time (sec)	N/A	1.046	4.457	0.908	0.000	4.515	0.000	0.000	0.000

Problem 438	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	196	225	238	2398858	0	12024	0	0	0
N.S.	1	1.15	1.21	12239.07	0.00	61.35	0.00	0.00	0.00
time (sec)	N/A	1.337	0.998	0.911	0.000	2.464	0.000	0.000	0.000

Problem 439	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	259	289	286	2400946	0	12093	0	0	0
N.S.	1	1.12	1.10	9270.06	0.00	46.69	0.00	0.00	0.00
time (sec)	N/A	1.837	3.341	0.879	0.000	2.318	0.000	0.000	0.000

Problem 440	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	311	352	346	2403086	0	12128	0	0	0
N.S.	1	1.13	1.11	7726.96	0.00	39.00	0.00	0.00	0.00
time (sec)	N/A	2.371	5.664	0.977	0.000	2.464	0.000	0.000	0.000

Problem 441	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	382	430	474	2403057	0	12202	0	0	0
N.S.	1	1.13	1.24	6290.73	0.00	31.94	0.00	0.00	0.00
time (sec)	N/A	2.981	6.785	0.900	0.000	2.431	0.000	0.000	0.000

Problem 442	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	397	400	411	2659561	0	35458	0	0	0
N.S.	1	1.01	1.04	6699.15	0.00	89.31	0.00	0.00	0.00
time (sec)	N/A	2.564	4.984	1.062	0.000	9.414	0.000	0.000	0.000

Problem 443	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	316	325	345	2657119	0	35244	0	0	0
N.S.	1	1.03	1.09	8408.60	0.00	111.53	0.00	0.00	0.00
time (sec)	N/A	2.041	4.832	1.038	0.000	7.575	0.000	0.000	0.000

Problem 444	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	260	257	291	2654895	0	35212	0	0	0
N.S.	1	0.99	1.12	10211.13	0.00	135.43	0.00	0.00	0.00
time (sec)	N/A	1.542	2.863	1.033	0.000	7.660	0.000	0.000	0.000

Problem 445	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	241	237	358	2653772	0	35230	0	0	0
N.S.	1	0.98	1.49	11011.50	0.00	146.18	0.00	0.00	0.00
time (sec)	N/A	1.507	2.707	1.078	0.000	7.595	0.000	0.000	0.000

Problem 446	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	240	241	417	2654078	0	35349	0	0	0
N.S.	1	1.00	1.74	11058.66	0.00	147.29	0.00	0.00	0.00
time (sec)	N/A	1.427	4.473	1.023	0.000	7.332	0.000	0.000	0.000

Problem 447	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	247	283	321	2652302	0	18802	0	0	0
N.S.	1	1.15	1.30	10738.06	0.00	76.12	0.00	0.00	0.00
time (sec)	N/A	1.885	2.288	1.021	0.000	5.056	0.000	0.000	0.000

Problem 448	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	309	350	381	2654465	0	18856	0	0	0
N.S.	1	1.13	1.23	8590.50	0.00	61.02	0.00	0.00	0.00
time (sec)	N/A	2.387	6.805	1.055	0.000	4.852	0.000	0.000	0.000

Problem 449	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	378	428	543	2659448	0	18922	0	0	0
N.S.	1	1.13	1.44	7035.58	0.00	50.06	0.00	0.00	0.00
time (sec)	N/A	3.004	7.155	1.047	0.000	4.896	0.000	0.000	0.000

Problem 450	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	460	509	632	2660696	0	18981	0	0	0
N.S.	1	1.11	1.37	5784.12	0.00	41.26	0.00	0.00	0.00
time (sec)	N/A	3.683	7.313	5.880	0.000	4.906	0.000	0.000	0.000

Problem 451	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	253	255	356	1491744	0	16341	0	0	0
N.S.	1	1.01	1.41	5896.22	0.00	64.59	0.00	0.00	0.00
time (sec)	N/A	1.569	4.856	5.865	0.000	2.542	0.000	0.000	0.000

Problem 452	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	206	207	245	1889462	0	20328	0	0	0
N.S.	1	1.00	1.19	9172.15	0.00	98.68	0.00	0.00	0.00
time (sec)	N/A	0.950	2.010	0.951	0.000	4.848	0.000	0.000	0.000

Problem 453	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	168	168	205	1885950	0	20234	0	0	30600
N.S.	1	1.00	1.22	11225.89	0.00	120.44	0.00	0.00	182.14
time (sec)	N/A	1.094	1.404	0.951	0.000	4.151	0.000	0.000	103.958

Problem 454	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	123	123	137	1878820	0	9913	0	0	8223
N.S.	1	1.00	1.11	15274.96	0.00	80.59	0.00	0.00	66.85
time (sec)	N/A	0.601	0.261	0.921	0.000	2.976	0.000	0.000	59.965

Problem 455	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	159	166	172	1886236	0	10036	0	0	0
N.S.	1	1.04	1.08	11863.12	0.00	63.12	0.00	0.00	0.00
time (sec)	N/A	0.832	0.553	0.920	0.000	2.982	0.000	0.000	0.000

Problem 456	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	203	219	195	1888895	0	10067	0	0	0
N.S.	1	1.08	0.96	9304.90	0.00	49.59	0.00	0.00	0.00
time (sec)	N/A	1.148	2.318	0.893	0.000	3.625	0.000	0.000	0.000

Problem 457	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	256	282	227	1890924	0	10125	0	0	0
N.S.	1	1.10	0.89	7386.42	0.00	39.55	0.00	0.00	0.00
time (sec)	N/A	1.674	5.785	0.941	0.000	3.061	0.000	0.000	0.000

Problem 458	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	219	253	341	1560668	0	37564	0	0	0
N.S.	1	1.16	1.56	7126.34	0.00	171.53	0.00	0.00	0.00
time (sec)	N/A	1.141	2.034	1.549	0.000	10.950	0.000	0.000	0.000

Problem 459	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	170	202	239	1559497	0	18532	0	0	0
N.S.	1	1.19	1.41	9173.51	0.00	109.01	0.00	0.00	0.00
time (sec)	N/A	0.970	1.698	1.434	0.000	7.014	0.000	0.000	0.000

Problem 460	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	175	203	202	4927	0	18563	0	0	0
N.S.	1	1.16	1.15	28.15	0.00	106.07	0.00	0.00	0.00
time (sec)	N/A	0.954	0.885	29.704	0.000	6.947	0.000	0.000	0.000

Problem 461	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	216	258	248	5198	0	18689	0	0	0
N.S.	1	1.19	1.15	24.06	0.00	86.52	0.00	0.00	0.00
time (sec)	N/A	1.316	2.327	3.927	0.000	7.038	0.000	0.000	0.000

Problem 462	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F(-1)	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	276	316	299	9712	0	18754	0	0	0
N.S.	1	1.14	1.08	35.19	0.00	67.95	0.00	0.00	0.00
time (sec)	N/A	1.798	3.264	3.992	0.000	7.046	0.000	0.000	0.000

Problem 463	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	282	344	596	2978162	0	55768	0	0	0
N.S.	1	1.22	2.11	10560.86	0.00	197.76	0.00	0.00	0.00
time (sec)	N/A	1.732	6.463	2.183	0.000	22.134	0.000	0.000	0.000

Problem 464	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	244	293	308	2978130	0	27576	0	0	0
N.S.	1	1.20	1.26	12205.45	0.00	113.02	0.00	0.00	0.00
time (sec)	N/A	1.512	3.350	5.394	0.000	14.842	0.000	0.000	0.000

Problem 465	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	244	297	320	2978176	0	27593	0	0	0
N.S.	1	1.22	1.31	12205.64	0.00	113.09	0.00	0.00	0.00
time (sec)	N/A	1.483	3.738	2.542	0.000	14.523	0.000	0.000	0.000

Problem 466	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	247	299	273	2975233	0	27639	0	0	0
N.S.	1	1.21	1.11	12045.48	0.00	111.90	0.00	0.00	0.00
time (sec)	N/A	1.572	2.753	3.069	0.000	14.523	0.000	0.000	0.000

Problem 467	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	301	361	326	2976756	0	27749	0	0	0
N.S.	1	1.20	1.08	9889.55	0.00	92.19	0.00	0.00	0.00
time (sec)	N/A	1.986	4.033	2.102	0.000	14.454	0.000	0.000	0.000

Problem 468	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	359	424	383	2982515	0	27865	0	0	0
N.S.	1	1.18	1.07	8307.84	0.00	77.62	0.00	0.00	0.00
time (sec)	N/A	2.599	3.891	5.471	0.000	14.524	0.000	0.000	0.000

Problem 469	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	155	148	190	943434	0	9186	0	0	0
N.S.	1	0.95	1.23	6086.67	0.00	59.26	0.00	0.00	0.00
time (sec)	N/A	0.431	0.837	1.307	0.000	1.046	0.000	0.000	0.000

Problem 470	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	117	114	124	940499	0	4509	0	0	0
N.S.	1	0.97	1.06	8038.45	0.00	38.54	0.00	0.00	0.00
time (sec)	N/A	0.334	0.057	0.573	0.000	0.750	0.000	0.000	0.000

Problem 471	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	111	108	125	940264	0	4469	0	0	0
N.S.	1	0.97	1.13	8470.85	0.00	40.26	0.00	0.00	0.00
time (sec)	N/A	0.356	0.048	0.542	0.000	0.771	0.000	0.000	0.000

Problem 472	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	150	149	158	943929	0	4608	0	0	0
N.S.	1	0.99	1.05	6292.86	0.00	30.72	0.00	0.00	0.00
time (sec)	N/A	0.455	0.421	0.576	0.000	0.761	0.000	0.000	0.000

Problem 473	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	B	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	379	269	263	99	0	5405	0	0	3945
N.S.	1	0.71	0.69	0.26	0.00	14.26	0.00	0.00	10.41
time (sec)	N/A	0.702	1.074	0.560	0.000	1.890	0.000	0.000	25.427

Problem 474	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	B	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	377	267	347	97	0	2785	0	0	2537
N.S.	1	0.71	0.92	0.26	0.00	7.39	0.00	0.00	6.73
time (sec)	N/A	0.725	0.987	0.614	0.000	0.373	0.000	0.000	20.451

Problem 475	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	B	F	A	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	357	233	227	72	0	4121	0	147	3228
N.S.	1	0.65	0.64	0.20	0.00	11.54	0.00	0.41	9.04
time (sec)	N/A	0.544	0.459	0.426	0.000	0.441	0.000	3.432	20.960

Problem 476	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	B	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	357	233	305	69	0	6073	0	0	4562
N.S.	1	0.65	0.85	0.19	0.00	17.01	0.00	0.00	12.78
time (sec)	N/A	0.567	0.273	0.401	0.000	1.345	0.000	0.000	23.039

Problem 477	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	B	F	B	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	148	105	104	42	0	278	0	795	2982
N.S.	1	0.71	0.70	0.28	0.00	1.88	0.00	5.37	20.15
time (sec)	N/A	0.298	0.671	0.358	0.000	0.253	0.000	0.662	22.944

Problem 478	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	B	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	299	235	330	72	0	443	0	0	4308
N.S.	1	0.79	1.10	0.24	0.00	1.48	0.00	0.00	14.41
time (sec)	N/A	0.585	0.401	0.457	0.000	0.249	0.000	0.000	23.058

Problem 479	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	403	416	355	0	0	0	0	0	0
N.S.	1	1.03	0.88	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.435	4.833	0.000	0.000	0.000	0.000	0.000	0.000

Problem 480	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	267	283	232	0	0	0	0	0	0
N.S.	1	1.06	0.87	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.431	2.516	0.000	0.000	0.000	0.000	0.000	0.000

Problem 481	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	194	198	155	0	0	0	0	0	0
N.S.	1	1.02	0.80	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.850	0.744	0.000	0.000	0.000	0.000	0.000	0.000

Problem 482	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	127	127	108	0	0	0	0	0	0
N.S.	1	1.00	0.85	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.459	0.500	0.000	0.000	0.000	0.000	0.000	0.000

Problem 483	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	185	178	144	0	0	0	0	0	0
N.S.	1	0.96	0.78	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.834	1.012	0.000	0.000	0.000	0.000	0.000	0.000

Problem 484	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	282	292	239	0	0	0	0	0	0
N.S.	1	1.04	0.85	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.420	3.328	0.000	0.000	0.000	0.000	0.000	0.000

Problem 485	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	438	464	530	0	0	0	0	0	0
N.S.	1	1.06	1.21	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.456	6.408	0.000	0.000	0.000	0.000	0.000	0.000

Problem 486	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	659	704	1425	0	0	0	0	0	0
N.S.	1	1.07	2.16	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.055	6.475	0.000	0.000	0.000	0.000	0.000	0.000

Problem 487	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	193	193	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.666	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 488	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F(-1)	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	189	189	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.626	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 489	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	187	187	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.605	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 490	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F(-1)	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	187	187	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.619	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 491	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F(-1)	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	193	193	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.640	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 492	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F(-1)	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	193	193	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.640	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 493	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	183	183	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.531	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 494	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	387	412	384	0	0	0	0	0	0
N.S.	1	1.06	0.99	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.972	6.100	0.000	0.000	0.000	0.000	0.000	0.000

Problem 495	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	291	319	281	0	0	0	0	0	0
N.S.	1	1.10	0.97	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.285	2.612	0.000	0.000	0.000	0.000	0.000	0.000

Problem 496	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	219	234	169	0	0	0	0	0	0
N.S.	1	1.07	0.77	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.832	1.496	0.000	0.000	0.000	0.000	0.000	0.000

Problem 497	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	168	174	125	0	0	0	0	0	0
N.S.	1	1.04	0.74	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.531	0.286	0.000	0.000	0.000	0.000	0.000	0.000

Problem 498	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	143	145	120	0	0	0	0	0	0
N.S.	1	1.01	0.84	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.385	0.204	0.000	0.000	0.000	0.000	0.000	0.000

Problem 499	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	190	196	169	0	0	0	0	0	0
N.S.	1	1.03	0.89	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.784	0.432	0.000	0.000	0.000	0.000	0.000	0.000

Problem 500	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	228	237	202	0	0	0	0	0	0
N.S.	1	1.04	0.89	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.071	0.463	0.000	0.000	0.000	0.000	0.000	0.000

Problem 501	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	292	317	230	0	0	0	0	0	0
N.S.	1	1.09	0.79	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.652	0.593	0.000	0.000	0.000	0.000	0.000	0.000

Problem 502	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	103	55	551	190	434	0	0	0
N.S.	1	1.00	0.53	5.35	1.84	4.21	0.00	0.00	0.00
time (sec)	N/A	0.705	0.764	0.611	0.375	0.265	0.000	0.000	0.000

Problem 503	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	78	54	528	174	382	0	0	0
N.S.	1	1.00	0.69	6.77	2.23	4.90	0.00	0.00	0.00
time (sec)	N/A	0.543	0.531	0.451	0.401	0.248	0.000	0.000	0.000

Problem 504	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	53	64	508	155	316	0	0	0
N.S.	1	1.00	1.21	9.58	2.92	5.96	0.00	0.00	0.00
time (sec)	N/A	0.465	0.772	0.362	0.371	0.243	0.000	0.000	0.000

Problem 505	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	55	63	422	155	364	0	0	0
N.S.	1	1.00	1.15	7.67	2.82	6.62	0.00	0.00	0.00
time (sec)	N/A	0.480	0.727	0.364	0.760	0.249	0.000	0.000	0.000

Problem 506	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	80	86	220	177	429	0	0	0
N.S.	1	1.00	1.08	2.75	2.21	5.36	0.00	0.00	0.00
time (sec)	N/A	0.603	1.041	0.357	0.295	0.250	0.000	0.000	0.000

Problem 507	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	105	105	99	470	191	482	0	0	0
N.S.	1	1.00	0.94	4.48	1.82	4.59	0.00	0.00	0.00
time (sec)	N/A	0.699	1.988	0.403	0.309	0.268	0.000	0.000	0.000

Problem 508	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	A	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	128	133	74	253	198	448	0	0	0
N.S.	1	1.04	0.58	1.98	1.55	3.50	0.00	0.00	0.00
time (sec)	N/A	0.844	1.581	0.716	0.305	0.264	0.000	0.000	0.000

Problem 509	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	106	78	230	180	394	0	0	0
N.S.	1	1.03	0.76	2.23	1.75	3.83	0.00	0.00	0.00
time (sec)	N/A	0.713	3.075	0.704	0.300	0.251	0.000	0.000	0.000

Problem 510	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	99	69	218	174	385	0	0	0
N.S.	1	1.00	0.70	2.20	1.76	3.89	0.00	0.00	0.00
time (sec)	N/A	0.700	2.568	0.391	0.324	0.248	0.000	0.000	0.000

Problem 511	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	105	108	90	225	180	439	0	0	0
N.S.	1	1.03	0.86	2.14	1.71	4.18	0.00	0.00	0.00
time (sec)	N/A	0.723	4.508	0.373	0.298	0.252	0.000	0.000	0.000

Problem 512	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	130	138	97	243	199	505	0	0	0
N.S.	1	1.06	0.75	1.87	1.53	3.88	0.00	0.00	0.00
time (sec)	N/A	0.898	4.873	0.376	0.308	0.260	0.000	0.000	0.000

Problem 513	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	181	113	275	218	508	0	0	0
N.S.	1	1.06	0.66	1.61	1.27	2.97	0.00	0.00	0.00
time (sec)	N/A	1.149	1.926	1.313	0.293	0.284	0.000	0.000	0.000

Problem 514	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	A	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	146	154	113	252	200	449	0	0	0
N.S.	1	1.05	0.77	1.73	1.37	3.08	0.00	0.00	0.00
time (sec)	N/A	0.977	3.585	1.237	0.305	0.268	0.000	0.000	0.000

Problem 515	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	A	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	141	113	241	194	409	0	0	0
N.S.	1	1.02	0.82	1.75	1.41	2.96	0.00	0.00	0.00
time (sec)	N/A	0.945	2.999	1.210	0.310	0.265	0.000	0.000	0.000

Problem 516	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	142	146	113	236	197	442	0	0	0
N.S.	1	1.03	0.80	1.66	1.39	3.11	0.00	0.00	0.00
time (sec)	N/A	0.963	2.954	0.460	0.299	0.254	0.000	0.000	0.000

Problem 517	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	148	156	110	243	203	502	0	0	0
N.S.	1	1.05	0.74	1.64	1.37	3.39	0.00	0.00	0.00
time (sec)	N/A	1.067	5.937	0.422	0.303	0.269	0.000	0.000	0.000

Problem 518	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	173	186	132	261	221	563	0	0	0
N.S.	1	1.08	0.76	1.51	1.28	3.25	0.00	0.00	0.00
time (sec)	N/A	1.246	6.343	0.425	0.303	0.279	0.000	0.000	0.000

Problem 519	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	297	265	186	543	0	716	0	0	0
N.S.	1	0.89	0.63	1.83	0.00	2.41	0.00	0.00	0.00
time (sec)	N/A	1.100	3.469	0.407	0.000	0.271	0.000	0.000	0.000

Problem 520	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	268	234	159	513	0	623	0	0	0
N.S.	1	0.87	0.59	1.91	0.00	2.32	0.00	0.00	0.00
time (sec)	N/A	0.868	2.379	0.360	0.000	0.265	0.000	0.000	0.000

Problem 521	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	235	205	137	342	0	570	0	0	0
N.S.	1	0.87	0.58	1.46	0.00	2.43	0.00	0.00	0.00
time (sec)	N/A	0.707	3.151	0.433	0.000	0.256	0.000	0.000	0.000

Problem 522	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	237	207	136	127	0	571	0	0	0
N.S.	1	0.87	0.57	0.54	0.00	2.41	0.00	0.00	0.00
time (sec)	N/A	0.705	3.382	0.434	0.000	0.267	0.000	0.000	0.000

Problem 523	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	276	242	154	143	0	700	0	0	0
N.S.	1	0.88	0.56	0.52	0.00	2.54	0.00	0.00	0.00
time (sec)	N/A	0.872	4.020	0.375	0.000	0.266	0.000	0.000	0.000

Problem 524	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	307	267	178	161	0	794	0	0	0
N.S.	1	0.87	0.58	0.52	0.00	2.59	0.00	0.00	0.00
time (sec)	N/A	1.043	5.571	0.410	0.000	0.266	0.000	0.000	0.000

Problem 525	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	317	286	221	762	0	669	0	0	0
N.S.	1	0.90	0.70	2.40	0.00	2.11	0.00	0.00	0.00
time (sec)	N/A	1.241	5.415	0.407	0.000	0.266	0.000	0.000	0.000

Problem 526	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	284	248	198	681	0	666	0	0	0
N.S.	1	0.87	0.70	2.40	0.00	2.35	0.00	0.00	0.00
time (sec)	N/A	1.002	4.195	0.425	0.000	0.260	0.000	0.000	0.000

Problem 527	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	274	241	196	146	0	663	0	0	0
N.S.	1	0.88	0.72	0.53	0.00	2.42	0.00	0.00	0.00
time (sec)	N/A	0.985	3.162	0.505	0.000	0.274	0.000	0.000	0.000

Problem 528	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	284	246	197	144	0	666	0	0	0
N.S.	1	0.87	0.69	0.51	0.00	2.35	0.00	0.00	0.00
time (sec)	N/A	0.998	4.384	0.524	0.000	0.260	0.000	0.000	0.000

Problem 529	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	319	288	215	157	0	761	0	0	0
N.S.	1	0.90	0.67	0.49	0.00	2.39	0.00	0.00	0.00
time (sec)	N/A	1.265	4.939	0.448	0.000	0.273	0.000	0.000	0.000

Problem 530	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	367	342	248	180	0	688	0	0	0
N.S.	1	0.93	0.68	0.49	0.00	1.87	0.00	0.00	0.00
time (sec)	N/A	1.640	5.997	0.398	0.000	0.280	0.000	0.000	0.000

Problem 531	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	318	293	223	162	0	683	0	0	0
N.S.	1	0.92	0.70	0.51	0.00	2.15	0.00	0.00	0.00
time (sec)	N/A	1.330	4.588	0.450	0.000	0.266	0.000	0.000	0.000

Problem 532	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	316	291	196	155	0	635	0	0	0
N.S.	1	0.92	0.62	0.49	0.00	2.01	0.00	0.00	0.00
time (sec)	N/A	1.289	5.790	0.546	0.000	0.275	0.000	0.000	0.000

Problem 533	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	308	287	203	150	0	637	0	0	0
N.S.	1	0.93	0.66	0.49	0.00	2.07	0.00	0.00	0.00
time (sec)	N/A	1.284	4.332	0.520	0.000	0.262	0.000	0.000	0.000

Problem 534	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	310	288	217	158	0	685	0	0	0
N.S.	1	0.93	0.70	0.51	0.00	2.21	0.00	0.00	0.00
time (sec)	N/A	1.329	5.761	0.549	0.000	0.265	0.000	0.000	0.000

Problem 535	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	367	346	242	178	0	783	0	0	0
N.S.	1	0.94	0.66	0.49	0.00	2.13	0.00	0.00	0.00
time (sec)	N/A	1.624	7.085	0.599	0.000	0.279	0.000	0.000	0.000

Problem 536	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	198	217	154	638	1409	481	0	0	0
N.S.	1	1.10	0.78	3.22	7.12	2.43	0.00	0.00	0.00
time (sec)	N/A	1.288	5.878	0.678	0.743	0.256	0.000	0.000	0.000

Problem 537	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	155	163	122	561	1157	430	0	0	0
N.S.	1	1.05	0.79	3.62	7.46	2.77	0.00	0.00	0.00
time (sec)	N/A	0.926	4.150	0.561	0.502	0.254	0.000	0.000	0.000

Problem 538	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	111	102	442	558	370	0	0	0
N.S.	1	1.01	0.93	4.02	5.07	3.36	0.00	0.00	0.00
time (sec)	N/A	0.597	3.042	0.561	0.404	0.252	0.000	0.000	0.000

Problem 539	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	152	133	161	506	0	576	0	0	0
N.S.	1	0.88	1.06	3.33	0.00	3.79	0.00	0.00	0.00
time (sec)	N/A	0.815	3.436	0.556	0.000	0.258	0.000	0.000	0.000

Problem 540	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	192	179	233	316	0	793	0	0	0
N.S.	1	0.93	1.21	1.65	0.00	4.13	0.00	0.00	0.00
time (sec)	N/A	1.112	3.754	0.678	0.000	0.256	0.000	0.000	0.000

Problem 541	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	245	269	359	804	3787	568	0	0	0
N.S.	1	1.10	1.47	3.28	15.46	2.32	0.00	0.00	0.00
time (sec)	N/A	1.587	7.450	0.577	3.818	0.260	0.000	0.000	0.000

Problem 542	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	201	215	325	715	1457	510	0	0	0
N.S.	1	1.07	1.62	3.56	7.25	2.54	0.00	0.00	0.00
time (sec)	N/A	1.220	6.638	0.532	0.740	0.266	0.000	0.000	0.000

Problem 543	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	157	161	291	626	1113	460	0	0	0
N.S.	1	1.03	1.85	3.99	7.09	2.93	0.00	0.00	0.00
time (sec)	N/A	0.892	5.675	0.534	0.511	0.262	0.000	0.000	0.000

Problem 544	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	186	167	357	529	0	674	0	0	0
N.S.	1	0.90	1.92	2.84	0.00	3.62	0.00	0.00	0.00
time (sec)	N/A	1.102	7.775	0.553	0.000	0.260	0.000	0.000	0.000

Problem 545	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	196	182	280	492	0	827	0	0	0
N.S.	1	0.93	1.43	2.51	0.00	4.22	0.00	0.00	0.00
time (sec)	N/A	1.117	8.710	0.527	0.000	0.283	0.000	0.000	0.000

Problem 546	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	244	233	376	455	0	913	0	0	0
N.S.	1	0.95	1.54	1.86	0.00	3.74	0.00	0.00	0.00
time (sec)	N/A	1.476	10.717	0.659	0.000	0.273	0.000	0.000	0.000

Problem 547	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	297	322	580	895	4543	629	0	0	0
N.S.	1	1.08	1.95	3.01	15.30	2.12	0.00	0.00	0.00
time (sec)	N/A	1.952	11.115	1.599	4.828	0.272	0.000	0.000	0.000

Problem 548	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	251	268	524	806	4087	577	0	0	0
N.S.	1	1.07	2.09	3.21	16.28	2.30	0.00	0.00	0.00
time (sec)	N/A	1.553	9.422	1.667	1.689	0.266	0.000	0.000	0.000

Problem 549	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	205	214	468	717	1545	515	0	0	0
N.S.	1	1.04	2.28	3.50	7.54	2.51	0.00	0.00	0.00
time (sec)	N/A	1.218	8.458	1.586	0.691	0.264	0.000	0.000	0.000

Problem 550	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	230	211	575	628	0	780	0	0	0
N.S.	1	0.92	2.50	2.73	0.00	3.39	0.00	0.00	0.00
time (sec)	N/A	1.391	8.300	1.664	0.000	0.279	0.000	0.000	0.000

Problem 551	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F(-1)	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	236	223	506	573	0	849	0	0	0
N.S.	1	0.94	2.14	2.43	0.00	3.60	0.00	0.00	0.00
time (sec)	N/A	1.448	8.250	0.539	0.000	0.275	0.000	0.000	0.000

Problem 552	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	246	233	454	572	0	923	0	0	0
N.S.	1	0.95	1.85	2.33	0.00	3.75	0.00	0.00	0.00
time (sec)	N/A	1.471	11.226	0.527	0.000	0.270	0.000	0.000	0.000

Problem 553	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	292	286	524	542	0	1013	0	0	0
N.S.	1	0.98	1.79	1.86	0.00	3.47	0.00	0.00	0.00
time (sec)	N/A	1.851	11.274	0.578	0.000	0.288	0.000	0.000	0.000

Problem 554	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	211	220	141	748	0	487	0	0	0
N.S.	1	1.04	0.67	3.55	0.00	2.31	0.00	0.00	0.00
time (sec)	N/A	1.201	5.221	0.620	0.000	0.255	0.000	0.000	0.000

Problem 555	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	163	166	121	703	0	426	0	0	0
N.S.	1	1.02	0.74	4.31	0.00	2.61	0.00	0.00	0.00
time (sec)	N/A	0.900	4.095	0.601	0.000	0.250	0.000	0.000	0.000

Problem 556	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	119	120	111	641	0	423	0	0	0
N.S.	1	1.01	0.93	5.39	0.00	3.55	0.00	0.00	0.00
time (sec)	N/A	0.627	2.844	0.693	0.000	0.255	0.000	0.000	0.000

Problem 557	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	196	181	191	607	0	736	0	0	0
N.S.	1	0.92	0.97	3.10	0.00	3.76	0.00	0.00	0.00
time (sec)	N/A	1.112	2.833	0.569	0.000	0.259	0.000	0.000	0.000

Problem 558	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	214	220	196	939	0	465	0	0	0
N.S.	1	1.03	0.92	4.39	0.00	2.17	0.00	0.00	0.00
time (sec)	N/A	1.255	7.145	0.610	0.000	0.260	0.000	0.000	0.000

Problem 559	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	168	172	146	876	0	463	0	0	0
N.S.	1	1.02	0.87	5.21	0.00	2.76	0.00	0.00	0.00
time (sec)	N/A	0.892	5.064	0.569	0.000	0.260	0.000	0.000	0.000

Problem 560	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	170	175	148	663	0	461	0	0	0
N.S.	1	1.03	0.87	3.90	0.00	2.71	0.00	0.00	0.00
time (sec)	N/A	0.920	5.077	0.533	0.000	0.254	0.000	0.000	0.000

Problem 561	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	243	232	232	1233	0	782	0	0	0
N.S.	1	0.95	0.95	5.07	0.00	3.22	0.00	0.00	0.00
time (sec)	N/A	1.483	5.866	0.535	0.000	0.277	0.000	0.000	0.000

Problem 562	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	260	274	223	1166	0	482	0	0	0
N.S.	1	1.05	0.86	4.48	0.00	1.85	0.00	0.00	0.00
time (sec)	N/A	1.596	9.206	0.572	0.000	0.267	0.000	0.000	0.000

Problem 563	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	214	226	221	1094	0	481	0	0	0
N.S.	1	1.06	1.03	5.11	0.00	2.25	0.00	0.00	0.00
time (sec)	N/A	1.263	5.721	0.585	0.000	0.259	0.000	0.000	0.000

Problem 564	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	216	228	222	856	0	482	0	0	0
N.S.	1	1.06	1.03	3.96	0.00	2.23	0.00	0.00	0.00
time (sec)	N/A	1.260	5.633	0.543	0.000	0.253	0.000	0.000	0.000

Problem 565	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	214	227	224	1106	0	482	0	0	0
N.S.	1	1.06	1.05	5.17	0.00	2.25	0.00	0.00	0.00
time (sec)	N/A	1.284	7.131	0.543	0.000	0.250	0.000	0.000	0.000

Problem 566	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	289	285	320	1552	0	799	0	0	0
N.S.	1	0.99	1.11	5.37	0.00	2.76	0.00	0.00	0.00
time (sec)	N/A	1.820	9.973	0.546	0.000	0.266	0.000	0.000	0.000

Problem 567	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	179	200	0	0	0	0	0	0	0
N.S.	1	1.12	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.852	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 568	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	247	301	0	0	0	0	0	0	0
N.S.	1	1.22	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.519	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 569	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	194	241	0	0	0	0	0	0	0
N.S.	1	1.24	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.114	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 570	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	158	197	0	0	0	0	0	0	0
N.S.	1	1.25	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.841	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 571	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	215	268	0	0	0	0	0	0	0
N.S.	1	1.25	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.171	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 572	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	291	351	0	0	0	0	0	0	0
N.S.	1	1.21	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.671	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 573	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	383	442	0	0	0	0	0	0	0
N.S.	1	1.15	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.247	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 574	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	229	206	198	535	196	2267	0	0	0
N.S.	1	0.90	0.86	2.34	0.86	9.90	0.00	0.00	0.00
time (sec)	N/A	0.746	1.026	0.472	0.334	0.701	0.000	0.000	0.000

Problem 575	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	205	182	179	512	178	2212	0	0	0
N.S.	1	0.89	0.87	2.50	0.87	10.79	0.00	0.00	0.00
time (sec)	N/A	0.610	0.527	0.370	0.322	0.701	0.000	0.000	0.000

Problem 576	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	205	182	178	426	178	2216	0	0	0
N.S.	1	0.89	0.87	2.08	0.87	10.81	0.00	0.00	0.00
time (sec)	N/A	0.644	0.228	0.362	0.302	0.709	0.000	0.000	0.000

Problem 577	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	229	206	198	223	198	2239	0	0	0
N.S.	1	0.90	0.86	0.97	0.86	9.78	0.00	0.00	0.00
time (sec)	N/A	0.804	0.564	0.383	0.303	0.698	0.000	0.000	0.000

Problem 578	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	326	283	255	293	279	4325	0	0	0
N.S.	1	0.87	0.78	0.90	0.86	13.27	0.00	0.00	0.00
time (sec)	N/A	1.119	2.023	0.687	0.301	2.267	0.000	0.000	0.000

Problem 579	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	294	249	226	252	252	4305	0	0	0
N.S.	1	0.85	0.77	0.86	0.86	14.64	0.00	0.00	0.00
time (sec)	N/A	0.958	1.291	0.704	0.312	2.306	0.000	0.000	0.000

Problem 580	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	276	228	221	239	244	4248	0	0	0
N.S.	1	0.83	0.80	0.87	0.88	15.39	0.00	0.00	0.00
time (sec)	N/A	0.897	0.903	0.414	0.306	2.278	0.000	0.000	0.000

Problem 581	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	283	235	226	244	254	4281	0	0	0
N.S.	1	0.83	0.80	0.86	0.90	15.13	0.00	0.00	0.00
time (sec)	N/A	0.962	0.612	0.399	0.312	2.333	0.000	0.000	0.000

Problem 582	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	317	269	255	271	282	4293	0	0	0
N.S.	1	0.85	0.80	0.85	0.89	13.54	0.00	0.00	0.00
time (sec)	N/A	1.109	1.226	0.407	0.406	2.365	0.000	0.000	0.000

Problem 583	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	421	355	326	1192	366	6214	0	0	0
N.S.	1	0.84	0.77	2.83	0.87	14.76	0.00	0.00	0.00
time (sec)	N/A	1.587	3.912	1.182	0.452	6.606	0.000	0.000	0.000

Problem 584	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	380	312	286	1134	330	6162	0	0	0
N.S.	1	0.82	0.75	2.98	0.87	16.22	0.00	0.00	0.00
time (sec)	N/A	1.317	2.468	1.220	0.410	6.464	0.000	0.000	0.000

Problem 585	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	374	303	270	1104	314	6154	0	0	0
N.S.	1	0.81	0.72	2.95	0.84	16.45	0.00	0.00	0.00
time (sec)	N/A	1.292	2.193	1.220	0.652	6.379	0.000	0.000	0.000

Problem 586	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	372	302	270	1104	316	6105	0	0	0
N.S.	1	0.81	0.73	2.97	0.85	16.41	0.00	0.00	0.00
time (sec)	N/A	1.318	2.241	0.453	0.315	6.202	0.000	0.000	0.000

Problem 587	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	380	312	287	956	334	6147	0	0	0
N.S.	1	0.82	0.76	2.52	0.88	16.18	0.00	0.00	0.00
time (sec)	N/A	1.393	1.322	0.450	0.323	6.194	0.000	0.000	0.000

Problem 588	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	421	358	327	328	370	6167	0	0	0
N.S.	1	0.85	0.78	0.78	0.88	14.65	0.00	0.00	0.00
time (sec)	N/A	1.642	2.686	0.479	0.321	6.199	0.000	0.000	0.000

Problem 589	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F(-1)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	325	286	272	772	262	6260	0	0	0
N.S.	1	0.88	0.84	2.38	0.81	19.26	0.00	0.00	0.00
time (sec)	N/A	1.733	1.756	0.497	0.308	26.271	0.000	0.000	0.000

Problem 590	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	297	250	249	686	239	6056	0	0	0
N.S.	1	0.84	0.84	2.31	0.80	20.39	0.00	0.00	0.00
time (sec)	N/A	1.406	0.954	0.385	0.326	11.525	0.000	0.000	0.000

Problem 591	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	278	224	215	531	221	6020	0	0	0
N.S.	1	0.81	0.77	1.91	0.79	21.65	0.00	0.00	0.00
time (sec)	N/A	1.057	0.435	0.395	0.318	4.333	0.000	0.000	0.000

Problem 592	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	278	224	215	244	220	5992	0	0	0
N.S.	1	0.81	0.77	0.88	0.79	21.55	0.00	0.00	0.00
time (sec)	N/A	1.091	0.441	0.400	0.316	3.091	0.000	0.000	0.000

Problem 593	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	297	249	251	265	238	6092	0	0	0
N.S.	1	0.84	0.85	0.89	0.80	20.51	0.00	0.00	0.00
time (sec)	N/A	1.452	0.596	0.369	0.312	8.778	0.000	0.000	0.000

Problem 594	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	325	285	272	286	262	6176	0	0	0
N.S.	1	0.88	0.84	0.88	0.81	19.00	0.00	0.00	0.00
time (sec)	N/A	1.846	1.111	0.391	0.317	21.242	0.000	0.000	0.000

Problem 595	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	438	377	383	2217	383	11904	0	0	0
N.S.	1	0.86	0.87	5.06	0.87	27.18	0.00	0.00	0.00
time (sec)	N/A	2.110	5.877	0.391	0.317	37.586	0.000	0.000	0.000

Problem 596	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	392	331	341	1936	362	11827	0	0	0
N.S.	1	0.84	0.87	4.94	0.92	30.17	0.00	0.00	0.00
time (sec)	N/A	1.605	2.790	0.415	0.418	23.694	0.000	0.000	0.000

Problem 597	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	390	330	336	332	347	11827	0	0	0
N.S.	1	0.85	0.86	0.85	0.89	30.33	0.00	0.00	0.00
time (sec)	N/A	1.643	3.022	0.423	0.509	14.678	0.000	0.000	0.000

Problem 598	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	392	331	342	336	360	11824	0	0	0
N.S.	1	0.84	0.87	0.86	0.92	30.16	0.00	0.00	0.00
time (sec)	N/A	1.687	2.683	0.444	0.733	19.641	0.000	0.000	0.000

Problem 599	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F(-1)	F(-1)	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	437	375	390	351	402	11930	0	0	0
N.S.	1	0.86	0.89	0.80	0.92	27.30	0.00	0.00	0.00
time (sec)	N/A	2.260	3.628	0.458	0.332	35.180	0.000	0.000	0.000

Problem 600	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	601	524	617	4581	577	17988	0	0	0
N.S.	1	0.87	1.03	7.62	0.96	29.93	0.00	0.00	0.00
time (sec)	N/A	3.151	6.565	0.417	0.322	187.830	0.000	0.000	0.000

Problem 601	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	534	462	581	4191	558	17874	0	0	0
N.S.	1	0.87	1.09	7.85	1.04	33.47	0.00	0.00	0.00
time (sec)	N/A	2.377	6.474	0.401	0.342	124.594	0.000	0.000	0.000

Problem 602	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	534	459	576	451	544	17912	0	0	0
N.S.	1	0.86	1.08	0.84	1.02	33.54	0.00	0.00	0.00
time (sec)	N/A	2.359	6.436	0.514	0.327	85.897	0.000	0.000	0.000

Problem 603	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F(-1)	F(-1)	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	530	459	586	450	545	17911	0	0	0
N.S.	1	0.87	1.11	0.85	1.03	33.79	0.00	0.00	0.00
time (sec)	N/A	2.378	6.458	0.574	0.316	73.397	0.000	0.000	0.000

Problem 604	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F(-1)	F(-1)	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	534	463	610	451	556	17872	0	0	0
N.S.	1	0.87	1.14	0.84	1.04	33.47	0.00	0.00	0.00
time (sec)	N/A	2.399	6.533	0.503	0.309	106.832	0.000	0.000	0.000

Problem 605	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F(-1)	F(-1)	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	600	524	639	466	612	18011	0	0	0
N.S.	1	0.87	1.06	0.78	1.02	30.02	0.00	0.00	0.00
time (sec)	N/A	3.312	6.581	0.490	0.298	163.937	0.000	0.000	0.000

Problem 606	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	157	89	102	127	458	0	0	64
N.S.	1	1.01	0.57	0.65	0.81	2.94	0.00	0.00	0.41
time (sec)	N/A	0.456	0.252	0.283	0.299	0.259	0.000	0.000	9.929

Problem 607	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	155	140	102	127	422	0	0	65
N.S.	1	1.01	0.91	0.66	0.82	2.74	0.00	0.00	0.42
time (sec)	N/A	0.408	0.183	0.273	0.289	0.261	0.000	0.000	9.823

Problem 608	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	138	60	90	113	362	0	0	42
N.S.	1	1.00	0.43	0.65	0.82	2.62	0.00	0.00	0.30
time (sec)	N/A	0.351	0.069	0.280	0.292	0.250	0.000	0.000	9.508

Problem 609	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	138	110	90	116	386	0	0	47
N.S.	1	1.00	0.80	0.65	0.84	2.80	0.00	0.00	0.34
time (sec)	N/A	0.356	0.060	0.309	0.386	0.267	0.000	0.000	9.409

Problem 610	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	155	78	102	126	471	0	0	64
N.S.	1	1.01	0.51	0.66	0.82	3.06	0.00	0.00	0.42
time (sec)	N/A	0.429	0.146	0.267	0.421	0.258	0.000	0.000	9.024

Problem 611	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	157	82	102	128	503	0	0	65
N.S.	1	1.01	0.53	0.65	0.82	3.22	0.00	0.00	0.42
time (sec)	N/A	0.425	0.240	0.326	0.355	0.265	0.000	0.000	9.892

Problem 612	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	354	376	291	2185615	0	8179	0	0	0
N.S.	1	1.06	0.82	6174.05	0.00	23.10	0.00	0.00	0.00
time (sec)	N/A	2.638	4.488	1.378	0.000	1.396	0.000	0.000	0.000

Problem 613	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	290	310	252	2183483	0	8104	0	0	0
N.S.	1	1.07	0.87	7529.25	0.00	27.94	0.00	0.00	0.00
time (sec)	N/A	1.938	2.481	1.184	0.000	1.397	0.000	0.000	0.000

Problem 614	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	239	243	216	2181430	0	8070	0	0	0
N.S.	1	1.02	0.90	9127.32	0.00	33.77	0.00	0.00	0.00
time (sec)	N/A	1.395	2.023	1.372	0.000	1.393	0.000	0.000	0.000

Problem 615	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	194	191	189	2178676	0	7949	0	0	0
N.S.	1	0.98	0.97	11230.29	0.00	40.97	0.00	0.00	0.00
time (sec)	N/A	1.041	0.524	1.286	0.000	1.428	0.000	0.000	0.000

Problem 616	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	229	200	238	2178428	0	16183	0	0	0
N.S.	1	0.87	1.04	9512.79	0.00	70.67	0.00	0.00	0.00
time (sec)	N/A	1.342	0.855	1.406	0.000	2.444	0.000	0.000	0.000

Problem 617	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	261	220	293	2180708	0	16171	0	0	0
N.S.	1	0.84	1.12	8355.20	0.00	61.96	0.00	0.00	0.00
time (sec)	N/A	1.162	3.769	1.208	0.000	3.381	0.000	0.000	0.000

Problem 618	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	324	286	356	2181027	0	16383	0	0	0
N.S.	1	0.88	1.10	6731.56	0.00	50.56	0.00	0.00	0.00
time (sec)	N/A	1.588	5.202	1.355	0.000	3.669	0.000	0.000	0.000

Problem 619	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	422	451	495	2403002	0	12342	0	0	0
N.S.	1	1.07	1.17	5694.32	0.00	29.25	0.00	0.00	0.00
time (sec)	N/A	3.196	6.730	1.384	0.000	2.356	0.000	0.000	0.000

Problem 620	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	351	373	346	2403427	0	12268	0	0	0
N.S.	1	1.06	0.99	6847.37	0.00	34.95	0.00	0.00	0.00
time (sec)	N/A	2.547	5.526	1.447	0.000	2.420	0.000	0.000	0.000

Problem 621	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	299	310	286	2401287	0	12233	0	0	0
N.S.	1	1.04	0.96	8031.06	0.00	40.91	0.00	0.00	0.00
time (sec)	N/A	2.009	3.213	1.488	0.000	2.366	0.000	0.000	0.000

Problem 622	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	236	246	244	2399199	0	12148	0	0	0
N.S.	1	1.04	1.03	10166.10	0.00	51.47	0.00	0.00	0.00
time (sec)	N/A	1.522	1.111	1.487	0.000	2.426	0.000	0.000	0.000

Problem 623	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	269	221	266	2397600	0	24440	0	0	0
N.S.	1	0.82	0.99	8913.01	0.00	90.86	0.00	0.00	0.00
time (sec)	N/A	1.224	2.635	1.414	0.000	4.700	0.000	0.000	0.000

Problem 624	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	264	221	263	2398750	0	24471	0	0	0
N.S.	1	0.84	1.00	9086.17	0.00	92.69	0.00	0.00	0.00
time (sec)	N/A	1.290	1.407	1.499	0.000	4.890	0.000	0.000	0.000

Problem 625	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	328	284	310	2398591	0	24563	0	0	0
N.S.	1	0.87	0.95	7312.78	0.00	74.89	0.00	0.00	0.00
time (sec)	N/A	1.744	3.538	1.394	0.000	5.024	0.000	0.000	0.000

Problem 626	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	383	347	367	2403527	0	24697	0	0	0
N.S.	1	0.91	0.96	6275.53	0.00	64.48	0.00	0.00	0.00
time (sec)	N/A	2.148	6.167	1.636	0.000	5.509	0.000	0.000	0.000

Problem 627	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	500	530	653	2660853	0	19121	0	0	0
N.S.	1	1.06	1.31	5321.71	0.00	38.24	0.00	0.00	0.00
time (sec)	N/A	3.987	7.217	1.447	0.000	4.955	0.000	0.000	0.000

Problem 628	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	418	449	564	2658291	0	19062	0	0	0
N.S.	1	1.07	1.35	6359.55	0.00	45.60	0.00	0.00	0.00
time (sec)	N/A	3.215	7.012	1.642	0.000	4.981	0.000	0.000	0.000

Problem 629	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	349	371	382	2682	0	18996	0	0	0
N.S.	1	1.06	1.09	7.68	0.00	54.43	0.00	0.00	0.00
time (sec)	N/A	2.607	5.723	9.664	0.000	4.973	0.000	0.000	0.000

Problem 630	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	287	304	321	2518	0	18942	0	0	0
N.S.	1	1.06	1.12	8.77	0.00	66.00	0.00	0.00	0.00
time (sec)	N/A	2.049	3.099	1.622	0.000	4.948	0.000	0.000	0.000

Problem 631	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	300	262	417	2438	0	35564	0	0	0
N.S.	1	0.87	1.39	8.13	0.00	118.55	0.00	0.00	0.00
time (sec)	N/A	1.686	5.720	1.411	0.000	7.263	0.000	0.000	0.000

Problem 632	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	301	258	363	2668	0	35373	0	0	0
N.S.	1	0.86	1.21	8.86	0.00	117.52	0.00	0.00	0.00
time (sec)	N/A	1.725	3.400	0.443	0.000	7.576	0.000	0.000	0.000

Problem 633	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	320	278	311	2819	0	35483	0	0	0
N.S.	1	0.87	0.97	8.81	0.00	110.88	0.00	0.00	0.00
time (sec)	N/A	1.756	3.766	0.513	0.000	7.669	0.000	0.000	0.000

Problem 634	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	376	346	365	2657129	0	35533	0	0	0
N.S.	1	0.92	0.97	7066.83	0.00	94.50	0.00	0.00	0.00
time (sec)	N/A	2.264	6.843	1.418	0.000	7.919	0.000	0.000	0.000

Problem 635	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	457	421	431	3240	0	35731	0	0	0
N.S.	1	0.92	0.94	7.09	0.00	78.19	0.00	0.00	0.00
time (sec)	N/A	2.763	6.246	1.049	0.000	9.419	0.000	0.000	0.000

Problem 636	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	296	303	444	1890883	0	10265	0	0	0
N.S.	1	1.02	1.50	6388.12	0.00	34.68	0.00	0.00	0.00
time (sec)	N/A	1.855	6.308	2.067	0.000	2.929	0.000	0.000	0.000

Problem 637	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	243	240	213	1890882	0	10191	0	0	0
N.S.	1	0.99	0.88	7781.41	0.00	41.94	0.00	0.00	0.00
time (sec)	N/A	1.378	2.924	1.311	0.000	2.937	0.000	0.000	0.000

Problem 638	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	199	187	193	1886187	0	10120	0	0	0
N.S.	1	0.94	0.97	9478.33	0.00	50.85	0.00	0.00	0.00
time (sec)	N/A	1.044	1.766	1.145	0.000	3.021	0.000	0.000	0.000

Problem 639	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	163	144	157	1880801	0	10053	0	0	0
N.S.	1	0.88	0.96	11538.66	0.00	61.67	0.00	0.00	0.00
time (sec)	N/A	0.792	0.440	1.230	0.000	2.888	0.000	0.000	0.000

Problem 640	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	228	189	225	1887832	0	20493	0	0	0
N.S.	1	0.83	0.99	8279.96	0.00	89.88	0.00	0.00	0.00
time (sec)	N/A	1.340	1.601	0.978	0.000	4.345	0.000	0.000	0.000

Problem 641	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	266	228	354	1890515	0	20571	0	0	0
N.S.	1	0.86	1.33	7107.20	0.00	77.33	0.00	0.00	0.00
time (sec)	N/A	1.162	2.694	1.249	0.000	5.036	0.000	0.000	0.000

Problem 642	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	316	337	301	1564143	0	18876	0	0	0
N.S.	1	1.07	0.95	4949.82	0.00	59.73	0.00	0.00	0.00
time (sec)	N/A	2.075	4.471	2.002	0.000	7.082	0.000	0.000	0.000

Problem 643	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	256	279	255	1560397	0	18757	0	0	0
N.S.	1	1.09	1.00	6095.30	0.00	73.27	0.00	0.00	0.00
time (sec)	N/A	1.566	2.530	1.972	0.000	7.136	0.000	0.000	0.000

Problem 644	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	215	224	222	1561166	0	18703	0	0	0
N.S.	1	1.04	1.03	7261.24	0.00	86.99	0.00	0.00	0.00
time (sec)	N/A	1.179	1.305	1.959	0.000	7.112	0.000	0.000	0.000

Problem 645	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	210	223	259	1559507	0	18672	0	0	0
N.S.	1	1.06	1.23	7426.22	0.00	88.91	0.00	0.00	0.00
time (sec)	N/A	1.190	2.350	1.321	0.000	7.005	0.000	0.000	0.000

Problem 646	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	279	274	361	1560634	0	37815	0	0	0
N.S.	1	0.98	1.29	5593.67	0.00	135.54	0.00	0.00	0.00
time (sec)	N/A	1.354	2.448	2.033	0.000	11.540	0.000	0.000	0.000

Problem 647	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	399	445	385	2981214	0	27987	0	0	0
N.S.	1	1.12	0.96	7471.71	0.00	70.14	0.00	0.00	0.00
time (sec)	N/A	2.864	4.436	6.016	0.000	14.368	0.000	0.000	0.000

Problem 648	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	341	382	334	2976685	0	27817	0	0	0
N.S.	1	1.12	0.98	8729.28	0.00	81.57	0.00	0.00	0.00
time (sec)	N/A	2.222	4.430	2.835	0.000	14.411	0.000	0.000	0.000

Problem 649	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	287	320	293	2978354	0	27790	0	0	0
N.S.	1	1.11	1.02	10377.54	0.00	96.83	0.00	0.00	0.00
time (sec)	N/A	1.768	3.900	2.742	0.000	14.620	0.000	0.000	0.000

Problem 650	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	284	318	340	2978186	0	27743	0	0	0
N.S.	1	1.12	1.20	10486.57	0.00	97.69	0.00	0.00	0.00
time (sec)	N/A	1.723	4.908	2.011	0.000	14.813	0.000	0.000	0.000

Problem 651	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	284	314	328	2975109	0	27726	0	0	0
N.S.	1	1.11	1.15	10475.74	0.00	97.63	0.00	0.00	0.00
time (sec)	N/A	1.753	4.014	2.698	0.000	14.564	0.000	0.000	0.000

Problem 652	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	342	365	617	2981291	0	56055	0	0	0
N.S.	1	1.07	1.80	8717.23	0.00	163.90	0.00	0.00	0.00
time (sec)	N/A	1.997	6.437	2.862	0.000	24.495	0.000	0.000	0.000

Problem 653	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	151	128	145	762	0	4609	0	0	0
N.S.	1	0.85	0.96	5.05	0.00	30.52	0.00	0.00	0.00
time (sec)	N/A	0.468	0.125	16.237	0.000	0.786	0.000	0.000	0.000

Problem 654	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	157	134	144	1302	0	4649	0	0	0
N.S.	1	0.85	0.92	8.29	0.00	29.61	0.00	0.00	0.00
time (sec)	N/A	0.473	0.112	13.342	0.000	0.756	0.000	0.000	0.000

Problem 655	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-1)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	215	168	210	1148	0	9437	0	0	0
N.S.	1	0.78	0.98	5.34	0.00	43.89	0.00	0.00	0.00
time (sec)	N/A	0.568	1.062	13.457	0.000	1.119	0.000	0.000	0.000

Problem 656	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	195	216	0	0	0	0	0	0	0
N.S.	1	1.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.799	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 657	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	169	182	0	0	0	0	0	0	0
N.S.	1	1.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.765	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 658	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	167	188	0	0	0	0	0	0	0
N.S.	1	1.13	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.795	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 659	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	173	194	0	0	0	0	0	0	0
N.S.	1	1.12	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.788	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 660	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	173	194	0	0	0	0	0	0	0
N.S.	1	1.12	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.791	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 661	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	173	173	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.587	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 662	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	173	173	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.590	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 663	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F(-1)	F	F	F(-1)	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	167	167	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.571	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 664	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F(-1)	F	F	F(-1)	F(-1)
verified	N/A	No	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	169	161	0	0	0	0	0	0	0
N.S.	1	0.95	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.610	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 665	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	64	50	80	311	93	394	0	128
N.S.	1	1.02	0.79	1.27	4.94	1.48	6.25	0.00	2.03
time (sec)	N/A	0.308	0.997	1.219	0.453	0.261	0.571	0.000	1.204

Problem 666	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	53	77	56	95	100	155	112	100
N.S.	1	0.90	1.31	0.95	1.61	1.69	2.63	1.90	1.69
time (sec)	N/A	0.287	1.445	0.180	0.320	0.243	0.306	0.676	9.157

Problem 667	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	53	70	56	72	86	136	99	76
N.S.	1	0.90	1.19	0.95	1.22	1.46	2.31	1.68	1.29
time (sec)	N/A	0.285	3.125	0.139	0.351	0.228	0.278	0.521	8.930

Problem 668	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	66	53	54	49	55	74	117	87	50
N.S.	1	0.80	0.82	0.74	0.83	1.12	1.77	1.32	0.76
time (sec)	N/A	0.275	1.945	0.105	0.304	0.238	0.206	0.429	8.926

Problem 669	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	24	32	27	29	54	82	106	25
N.S.	1	0.75	1.00	0.84	0.91	1.69	2.56	3.31	0.78
time (sec)	N/A	0.215	0.041	0.063	0.309	0.242	0.150	0.347	8.980

Problem 670	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	66	50	50	64	53	103	38
N.S.	1	1.00	1.43	1.09	1.09	1.39	1.15	2.24	0.83
time (sec)	N/A	0.258	0.045	0.061	0.309	0.247	0.212	0.339	8.885

Problem 671	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	54	43	44	74	0	43	88	123	51
N.S.	1	0.80	0.81	1.37	0.00	0.80	1.63	2.28	0.94
time (sec)	N/A	0.291	2.236	0.114	0.000	0.249	0.188	0.391	8.856

Problem 672	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	44	46	0	46	153	79	51
N.S.	1	1.00	0.96	1.00	0.00	1.00	3.33	1.72	1.11
time (sec)	N/A	0.269	2.557	0.143	0.000	0.238	0.201	0.451	9.034

Problem 673	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	49	41	43	0	59	201	140	63
N.S.	1	0.89	0.75	0.78	0.00	1.07	3.65	2.55	1.15
time (sec)	N/A	0.283	0.488	0.148	0.000	0.240	0.246	0.640	8.917

Problem 674	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	57	51	41	44	0	81	304	200	73
N.S.	1	0.89	0.72	0.77	0.00	1.42	5.33	3.51	1.28
time (sec)	N/A	0.282	0.529	0.170	0.000	0.254	0.308	0.693	8.594

Problem 675	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	49	41	45	0	94	348	260	82
N.S.	1	0.89	0.75	0.82	0.00	1.71	6.33	4.73	1.49
time (sec)	N/A	0.281	1.676	0.187	0.000	0.244	0.394	0.975	8.661

Problem 676	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	109	102	89	167	660	206	1482	0	193
N.S.	1	0.94	0.82	1.53	6.06	1.89	13.60	0.00	1.77
time (sec)	N/A	0.364	2.178	1.045	0.451	0.259	1.252	0.000	11.539

Problem 677	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	82	270	83	151	152	243	180	158
N.S.	1	0.83	2.73	0.84	1.53	1.54	2.45	1.82	1.60
time (sec)	N/A	0.354	1.989	0.359	0.412	0.248	0.613	1.019	9.225

Problem 678	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	82	103	83	116	136	224	167	120
N.S.	1	0.83	1.04	0.84	1.17	1.37	2.26	1.69	1.21
time (sec)	N/A	0.326	2.307	0.273	0.373	0.236	0.493	0.888	9.283

Problem 679	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	82	95	83	101	124	206	155	108
N.S.	1	0.83	0.96	0.84	1.02	1.25	2.08	1.57	1.09
time (sec)	N/A	0.331	5.625	0.187	0.380	0.237	0.390	0.660	9.432

Problem 680	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	49	53	53	72	105	160	391	82
N.S.	1	0.79	0.85	0.85	1.16	1.69	2.58	6.31	1.32
time (sec)	N/A	0.290	0.170	0.110	0.354	0.225	0.281	0.541	8.736

Problem 681	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	64	51	54	51	53	98	158	120	50
N.S.	1	0.80	0.84	0.80	0.83	1.53	2.47	1.88	0.78
time (sec)	N/A	0.267	1.076	0.110	0.435	0.248	0.214	0.441	8.325

Problem 682	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	70	58	76	71	121	122	214	76
N.S.	1	0.88	0.72	0.95	0.89	1.51	1.52	2.68	0.95
time (sec)	N/A	0.361	0.371	0.095	0.373	0.245	0.317	0.411	8.431

Problem 683	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	93	65	84	146	0	111	134	271	105
N.S.	1	0.70	0.90	1.57	0.00	1.19	1.44	2.91	1.13
time (sec)	N/A	0.322	5.526	0.188	0.000	0.254	0.362	0.469	8.399

Problem 684	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	91	65	56	103	0	62	160	191	104
N.S.	1	0.71	0.62	1.13	0.00	0.68	1.76	2.10	1.14
time (sec)	N/A	0.322	4.527	0.126	0.000	0.250	0.288	0.542	8.437

Problem 685	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	93	76	62	69	0	49	167	156	87
N.S.	1	0.82	0.67	0.74	0.00	0.53	1.80	1.68	0.94
time (sec)	N/A	0.332	3.941	0.229	0.000	0.245	0.276	0.709	8.323

Problem 686	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	91	74	51	68	0	64	218	190	78
N.S.	1	0.81	0.56	0.75	0.00	0.70	2.40	2.09	0.86
time (sec)	N/A	0.332	5.301	0.149	0.000	0.246	0.328	0.886	8.455

Problem 687	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	78	62	69	0	89	332	292	108
N.S.	1	0.82	0.65	0.73	0.00	0.94	3.49	3.07	1.14
time (sec)	N/A	0.335	5.363	0.372	0.000	0.257	0.414	1.128	8.646

Problem 688	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	91	74	60	66	0	104	379	360	118
N.S.	1	0.81	0.66	0.73	0.00	1.14	4.16	3.96	1.30
time (sec)	N/A	0.327	5.406	0.187	0.000	0.252	0.523	1.248	8.476

Problem 689	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	151	138	154	271	1056	335	3665	0	323
N.S.	1	0.91	1.02	1.79	6.99	2.22	24.27	0.00	2.14
time (sec)	N/A	0.394	5.190	0.955	0.581	0.256	3.355	0.000	13.211

Problem 690	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	135	109	150	108	193	194	325	239	208
N.S.	1	0.81	1.11	0.80	1.43	1.44	2.41	1.77	1.54
time (sec)	N/A	0.382	5.892	0.637	0.299	0.246	0.934	1.289	8.717

Problem 691	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	135	109	98	108	166	182	306	227	174
N.S.	1	0.81	0.73	0.80	1.23	1.35	2.27	1.68	1.29
time (sec)	N/A	0.361	5.690	0.520	0.320	0.239	0.828	1.200	8.603

Problem 692	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	132	106	131	108	151	170	287	215	156
N.S.	1	0.80	0.99	0.82	1.14	1.29	2.17	1.63	1.18
time (sec)	N/A	0.354	2.090	0.352	0.324	0.242	0.641	0.965	8.476

Problem 693	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	84	61	65	75	106	147	224	754	120
N.S.	1	0.73	0.77	0.89	1.26	1.75	2.67	8.98	1.43
time (sec)	N/A	0.303	0.278	0.123	0.303	0.248	0.487	0.904	8.328

Problem 694	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	84	91	94	105	152	250	192	108
N.S.	1	0.83	0.90	0.93	1.04	1.50	2.48	1.90	1.07
time (sec)	N/A	0.333	5.539	0.234	0.308	0.237	0.412	0.642	8.307

Problem 695	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	55	70	63	72	132	218	164	72
N.S.	1	0.90	1.15	1.03	1.18	2.16	3.57	2.69	1.18
time (sec)	N/A	0.287	3.283	0.146	0.308	0.232	0.278	0.542	8.342

Problem 696	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	99	73	100	96	175	184	312	125
N.S.	1	0.90	0.66	0.91	0.87	1.59	1.67	2.84	1.14
time (sec)	N/A	0.462	0.918	0.134	0.313	0.264	0.369	0.489	8.479

Problem 697	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	119	81	120	178	0	164	206	306	139
N.S.	1	0.68	1.01	1.50	0.00	1.38	1.73	2.57	1.17
time (sec)	N/A	0.352	5.681	0.118	0.000	0.258	0.447	0.583	8.633

Problem 698	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	123	85	90	189	0	138	236	341	182
N.S.	1	0.69	0.73	1.54	0.00	1.12	1.92	2.77	1.48
time (sec)	N/A	0.339	5.603	0.188	0.000	0.250	0.509	0.644	9.096

Problem 699	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	129	101	74	124	0	77	212	242	141
N.S.	1	0.78	0.57	0.96	0.00	0.60	1.64	1.88	1.09
time (sec)	N/A	0.339	4.964	0.220	0.000	0.249	0.395	0.870	8.694

Problem 700	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	91	79	88	0	49	167	224	118
N.S.	1	0.92	0.80	0.89	0.00	0.49	1.69	2.26	1.19
time (sec)	N/A	0.311	3.528	0.241	0.000	0.250	0.374	1.026	8.698

Problem 701	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	122	96	78	87	0	64	218	292	128
N.S.	1	0.79	0.64	0.71	0.00	0.52	1.79	2.39	1.05
time (sec)	N/A	0.349	5.640	0.306	0.000	0.245	0.455	1.301	8.700

Problem 702	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	127	101	80	90	0	89	332	326	140
N.S.	1	0.80	0.63	0.71	0.00	0.70	2.61	2.57	1.10
time (sec)	N/A	0.345	5.668	0.188	0.000	0.246	0.531	1.582	8.709

Problem 703	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	99	80	89	0	104	379	428	151
N.S.	1	0.79	0.64	0.71	0.00	0.83	3.03	3.42	1.21
time (sec)	N/A	0.344	5.526	0.500	0.000	0.269	0.662	1.065	8.937

Problem 704	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	127	101	84	90	0	131	496	496	160
N.S.	1	0.80	0.66	0.71	0.00	1.03	3.91	3.91	1.26
time (sec)	N/A	0.352	5.554	0.275	0.000	0.258	0.718	1.057	9.222

Problem 705	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	115	119	98	0	0	0	0	0	0
N.S.	1	1.03	0.85	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.355	1.421	0.000	0.000	0.000	0.000	0.000	0.000

Problem 706	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	157	110	150	224	0	298	328	424	205
N.S.	1	0.70	0.96	1.43	0.00	1.90	2.09	2.70	1.31
time (sec)	N/A	0.380	1.508	0.172	0.000	0.252	0.507	0.660	8.427

Problem 707	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	121	85	124	191	0	219	265	307	136
N.S.	1	0.70	1.02	1.58	0.00	1.81	2.19	2.54	1.12
time (sec)	N/A	0.337	5.107	0.136	0.000	0.267	0.453	0.600	8.479

Problem 708	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	96	68	83	154	0	153	194	271	110
N.S.	1	0.71	0.86	1.60	0.00	1.59	2.02	2.82	1.15
time (sec)	N/A	0.336	4.419	0.165	0.000	0.270	0.366	0.507	8.812

Problem 709	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	57	48	47	83	0	67	112	123	54
N.S.	1	0.84	0.82	1.46	0.00	1.18	1.96	2.16	0.95
time (sec)	N/A	0.292	0.743	0.120	0.000	0.253	0.192	0.388	8.229

Problem 710	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	56	54	0	42	87	83	45
N.S.	1	1.00	1.19	1.15	0.00	0.89	1.85	1.77	0.96
time (sec)	N/A	0.220	0.276	0.095	0.000	0.254	0.132	0.381	8.370

Problem 711	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	C	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	50	43	54	0	58	165	50	40
N.S.	1	1.11	0.96	1.20	0.00	1.29	3.67	1.11	0.89
time (sec)	N/A	0.312	0.110	0.131	0.000	0.252	0.183	0.425	8.878

Problem 712	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	113	96	123	130	0	80	284	159	129
N.S.	1	0.85	1.09	1.15	0.00	0.71	2.51	1.41	1.14
time (sec)	N/A	0.368	5.629	0.187	0.000	0.251	0.265	0.547	9.063

Problem 713	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	149	123	149	195	0	92	328	180	161
N.S.	1	0.83	1.00	1.31	0.00	0.62	2.20	1.21	1.08
time (sec)	N/A	0.388	5.814	0.206	0.000	0.245	0.340	0.612	9.237

Problem 714	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	181	146	141	238	0	115	439	207	204
N.S.	1	0.81	0.78	1.31	0.00	0.64	2.43	1.14	1.13
time (sec)	N/A	0.405	3.779	0.213	0.000	0.250	0.412	0.822	9.826

Problem 715	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	115	119	152	0	0	0	0	0	0
N.S.	1	1.03	1.32	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.350	2.476	0.000	0.000	0.000	0.000	0.000	0.000

Problem 716	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	194	138	197	267	0	329	445	492	282
N.S.	1	0.71	1.02	1.38	0.00	1.70	2.29	2.54	1.45
time (sec)	N/A	0.404	6.122	0.324	0.000	0.271	0.691	1.028	8.527

Problem 717	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	158	113	129	239	0	249	377	423	207
N.S.	1	0.72	0.82	1.51	0.00	1.58	2.39	2.68	1.31
time (sec)	N/A	0.370	6.031	0.252	0.000	0.256	0.582	0.782	8.477

Problem 718	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	128	94	92	200	0	177	309	341	194
N.S.	1	0.73	0.72	1.56	0.00	1.38	2.41	2.66	1.52
time (sec)	N/A	0.355	5.643	0.200	0.000	0.260	0.495	0.662	9.069

Problem 719	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	73	58	114	0	88	206	191	104
N.S.	1	0.75	0.60	1.18	0.00	0.91	2.12	1.97	1.07
time (sec)	N/A	0.340	5.642	0.128	0.000	0.263	0.293	0.543	8.836

Problem 720	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	48	48	60	46	0	47	158	79	50
N.S.	1	1.00	1.25	0.96	0.00	0.98	3.29	1.65	1.04
time (sec)	N/A	0.259	1.207	0.152	0.000	0.247	0.205	0.473	8.313

Problem 721	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	80	79	89	73	0	54	162	110	70
N.S.	1	0.99	1.11	0.91	0.00	0.68	2.02	1.38	0.88
time (sec)	N/A	0.319	0.800	0.105	0.000	0.236	0.183	0.435	8.292

Problem 722	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	100	123	130	0	79	296	159	129
N.S.	1	0.85	1.05	1.11	0.00	0.68	2.53	1.36	1.10
time (sec)	N/A	0.371	5.695	0.210	0.000	0.237	0.264	0.554	8.587

Problem 723	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	C	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	71	77	53	96	0	90	360	63	53
N.S.	1	1.08	0.75	1.35	0.00	1.27	5.07	0.89	0.75
time (sec)	N/A	0.324	0.165	0.146	0.000	0.255	0.343	0.558	8.625

Problem 724	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	183	148	173	209	0	115	454	205	208
N.S.	1	0.81	0.95	1.14	0.00	0.63	2.48	1.12	1.14
time (sec)	N/A	0.415	5.997	0.277	0.000	0.252	0.421	0.785	9.094

Problem 725	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	221	177	213	281	0	127	498	225	247
N.S.	1	0.80	0.96	1.27	0.00	0.57	2.25	1.02	1.12
time (sec)	N/A	0.440	6.080	0.283	0.000	0.248	0.512	0.943	10.033

Problem 726	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	251	198	233	303	0	151	605	251	291
N.S.	1	0.79	0.93	1.21	0.00	0.60	2.41	1.00	1.16
time (sec)	N/A	0.460	6.529	0.351	0.000	0.247	0.582	0.793	10.573

Problem 727	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	115	119	210	0	0	0	0	0	0
N.S.	1	1.03	1.83	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.355	2.576	0.000	0.000	0.000	0.000	0.000	0.000

Problem 728	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	191	137	152	285	0	267	473	491	233
N.S.	1	0.72	0.80	1.49	0.00	1.40	2.48	2.57	1.22
time (sec)	N/A	0.403	5.012	0.257	0.000	0.260	0.774	1.231	8.787

Problem 729	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	164	118	107	248	0	197	403	409	266
N.S.	1	0.72	0.65	1.51	0.00	1.20	2.46	2.49	1.62
time (sec)	N/A	0.367	5.705	0.223	0.000	0.264	0.641	1.039	10.605

Problem 730	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	135	109	76	135	0	103	258	242	149
N.S.	1	0.81	0.56	1.00	0.00	0.76	1.91	1.79	1.10
time (sec)	N/A	0.348	5.908	0.221	0.000	0.256	0.390	0.881	8.678

Problem 731	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	82	62	69	0	49	172	156	87
N.S.	1	0.83	0.63	0.70	0.00	0.49	1.74	1.58	0.88
time (sec)	N/A	0.335	5.333	0.196	0.000	0.258	0.279	0.727	8.517

Problem 732	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	53	41	43	0	60	206	140	62
N.S.	1	0.90	0.69	0.73	0.00	1.02	3.49	2.37	1.05
time (sec)	N/A	0.282	5.140	0.138	0.000	0.244	0.270	0.635	8.469

Problem 733	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	112	111	109	128	0	76	258	129	111
N.S.	1	0.99	0.97	1.14	0.00	0.68	2.30	1.15	0.99
time (sec)	N/A	0.408	1.006	0.110	0.000	0.238	0.259	0.568	8.560

Problem 734	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	153	127	148	195	0	91	340	180	161
N.S.	1	0.83	0.97	1.27	0.00	0.59	2.22	1.18	1.05
time (sec)	N/A	0.389	5.832	0.278	0.000	0.239	0.348	0.632	8.733

Problem 735	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	185	150	172	209	0	115	452	205	208
N.S.	1	0.81	0.93	1.13	0.00	0.62	2.44	1.11	1.12
time (sec)	N/A	0.408	6.061	0.275	0.000	0.254	0.456	0.800	9.364

Problem 736	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	C	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	104	63	138	0	126	508	74	64
N.S.	1	1.05	0.64	1.39	0.00	1.27	5.13	0.75	0.65
time (sec)	N/A	0.324	0.181	0.136	0.000	0.248	0.549	0.767	8.730

Problem 737	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	251	198	221	261	0	151	605	251	286
N.S.	1	0.79	0.88	1.04	0.00	0.60	2.41	1.00	1.14
time (sec)	N/A	0.444	6.402	0.257	0.000	0.256	0.632	0.848	10.531

Problem 738	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	287	225	261	335	0	159	646	269	319
N.S.	1	0.78	0.91	1.17	0.00	0.55	2.25	0.94	1.11
time (sec)	N/A	0.477	6.795	0.385	0.000	0.249	0.797	0.888	10.407

Problem 739	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	319	248	278	392	0	185	753	295	352
N.S.	1	0.78	0.87	1.23	0.00	0.58	2.36	0.92	1.10
time (sec)	N/A	0.499	7.199	0.475	0.000	0.259	0.844	0.946	10.578

Problem 740	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	63	58	55	48	108	0	0	101
N.S.	1	1.02	0.94	0.89	0.77	1.74	0.00	0.00	1.63
time (sec)	N/A	0.313	3.569	0.389	0.302	0.353	0.000	0.000	12.347

Problem 741	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	63	60	55	48	96	0	0	101
N.S.	1	1.02	0.97	0.89	0.77	1.55	0.00	0.00	1.63
time (sec)	N/A	0.293	2.227	0.382	0.308	0.306	0.000	0.000	13.274

Problem 742	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	63	54	55	48	78	0	0	99
N.S.	1	1.02	0.87	0.89	0.77	1.26	0.00	0.00	1.60
time (sec)	N/A	0.300	1.546	0.286	0.346	0.260	0.000	0.000	10.868

Problem 743	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	60	61	45	66	48	66	0	0	102
N.S.	1	1.02	0.75	1.10	0.80	1.10	0.00	0.00	1.70
time (sec)	N/A	0.289	1.176	0.347	0.299	0.247	0.000	0.000	0.567

Problem 744	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	58	59	42	53	48	55	0	0	164
N.S.	1	1.02	0.72	0.91	0.83	0.95	0.00	0.00	2.83
time (sec)	N/A	0.294	1.587	0.252	0.308	0.256	0.000	0.000	8.429

Problem 745	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	60	61	57	53	45	75	0	0	170
N.S.	1	1.02	0.95	0.88	0.75	1.25	0.00	0.00	2.83
time (sec)	N/A	0.301	2.554	0.274	0.227	0.249	0.000	0.000	9.062

Problem 746	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	63	58	53	46	92	0	0	232
N.S.	1	1.02	0.94	0.85	0.74	1.48	0.00	0.00	3.74
time (sec)	N/A	0.308	3.721	0.219	0.220	0.255	0.000	0.000	9.963

Problem 747	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	63	58	53	46	113	0	0	157
N.S.	1	1.02	0.94	0.85	0.74	1.82	0.00	0.00	2.53
time (sec)	N/A	0.305	5.644	0.272	0.223	0.277	0.000	0.000	10.376

Problem 748	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	105	98	104	84	78	149	0	0	132
N.S.	1	0.93	0.99	0.80	0.74	1.42	0.00	0.00	1.26
time (sec)	N/A	0.392	5.759	0.406	0.214	0.421	0.000	0.000	11.859

Problem 749	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	105	98	104	84	78	133	0	0	132
N.S.	1	0.93	0.99	0.80	0.74	1.27	0.00	0.00	1.26
time (sec)	N/A	0.378	5.304	0.444	0.224	0.325	0.000	0.000	12.744

Problem 750	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	105	98	105	84	78	119	0	0	130
N.S.	1	0.93	1.00	0.80	0.74	1.13	0.00	0.00	1.24
time (sec)	N/A	0.381	4.245	0.318	0.213	0.276	0.000	0.000	11.704

Problem 751	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	96	76	84	78	104	0	0	241
N.S.	1	0.93	0.74	0.82	0.76	1.01	0.00	0.00	2.34
time (sec)	N/A	0.360	3.190	0.439	0.213	0.258	0.000	0.000	10.353

Problem 752	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	94	62	94	81	93	0	0	176
N.S.	1	0.93	0.61	0.93	0.80	0.92	0.00	0.00	1.74
time (sec)	N/A	0.371	4.093	0.284	0.214	0.259	0.000	0.000	9.187

Problem 753	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	94	81	79	79	81	0	0	158
N.S.	1	0.93	0.80	0.78	0.78	0.80	0.00	0.00	1.56
time (sec)	N/A	0.380	5.537	0.280	0.228	0.246	0.000	0.000	9.006

Problem 754	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	96	77	80	76	102	0	0	208
N.S.	1	0.93	0.75	0.78	0.74	0.99	0.00	0.00	2.02
time (sec)	N/A	0.376	5.740	0.293	0.227	0.254	0.000	0.000	9.771

Problem 755	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	105	98	79	80	76	123	0	0	167
N.S.	1	0.93	0.75	0.76	0.72	1.17	0.00	0.00	1.59
time (sec)	N/A	0.382	6.018	0.244	0.214	0.273	0.000	0.000	10.116

Problem 756	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	144	131	123	121	104	179	0	0	349
N.S.	1	0.91	0.85	0.84	0.72	1.24	0.00	0.00	2.42
time (sec)	N/A	0.397	6.802	0.688	0.220	0.531	0.000	0.000	12.901

Problem 757	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	144	131	128	121	104	173	0	0	349
N.S.	1	0.91	0.89	0.84	0.72	1.20	0.00	0.00	2.42
time (sec)	N/A	0.384	5.970	0.597	0.210	0.384	0.000	0.000	12.108

Problem 758	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	144	131	128	121	104	153	0	0	335
N.S.	1	0.91	0.89	0.84	0.72	1.06	0.00	0.00	2.33
time (sec)	N/A	0.391	4.884	0.391	0.221	0.315	0.000	0.000	11.700

Problem 759	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	142	129	93	121	104	137	0	0	313
N.S.	1	0.91	0.65	0.85	0.73	0.96	0.00	0.00	2.20
time (sec)	N/A	0.371	3.957	0.374	0.305	0.280	0.000	0.000	12.550

Problem 760	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	127	83	135	108	127	0	0	351
N.S.	1	0.91	0.59	0.96	0.77	0.91	0.00	0.00	2.51
time (sec)	N/A	0.386	5.194	0.261	0.340	0.260	0.000	0.000	11.890

Problem 761	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	127	95	118	105	117	0	0	221
N.S.	1	0.91	0.68	0.84	0.75	0.84	0.00	0.00	1.58
time (sec)	N/A	0.400	5.825	0.230	0.306	0.264	0.000	0.000	10.457

Problem 762	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	127	100	105	107	102	0	0	208
N.S.	1	0.91	0.71	0.75	0.76	0.73	0.00	0.00	1.49
time (sec)	N/A	0.391	6.249	0.236	0.214	0.255	0.000	0.000	9.893

Problem 763	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	142	129	95	105	102	123	0	0	161
N.S.	1	0.91	0.67	0.74	0.72	0.87	0.00	0.00	1.13
time (sec)	N/A	0.393	6.375	0.283	0.228	0.263	0.000	0.000	10.108

Problem 764	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	220	188	148	192	190	466	0	0	298
N.S.	1	0.85	0.67	0.87	0.86	2.12	0.00	0.00	1.35
time (sec)	N/A	0.412	7.748	0.227	0.295	0.302	0.000	0.000	1.368

Problem 765	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	180	161	134	150	167	406	0	0	245
N.S.	1	0.89	0.74	0.83	0.93	2.26	0.00	0.00	1.36
time (sec)	N/A	0.377	6.182	0.266	0.302	0.273	0.000	0.000	9.386

Problem 766	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	144	132	108	111	138	330	0	0	189
N.S.	1	0.92	0.75	0.77	0.96	2.29	0.00	0.00	1.31
time (sec)	N/A	0.374	3.905	0.284	0.315	0.253	0.000	0.000	9.199

Problem 767	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	109	108	98	90	116	326	0	0	159
N.S.	1	0.99	0.90	0.83	1.06	2.99	0.00	0.00	1.46
time (sec)	N/A	0.349	2.889	0.309	0.303	0.260	0.000	0.000	8.936

Problem 768	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	141	137	113	121	143	355	0	0	212
N.S.	1	0.97	0.80	0.86	1.01	2.52	0.00	0.00	1.50
time (sec)	N/A	0.375	3.906	0.204	0.299	0.257	0.000	0.000	9.292

Problem 769	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	184	171	103	141	167	388	0	0	261
N.S.	1	0.93	0.56	0.77	0.91	2.11	0.00	0.00	1.42
time (sec)	N/A	0.410	4.712	0.270	0.295	0.258	0.000	0.000	9.700

Problem 770	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	223	205	127	168	195	418	0	0	308
N.S.	1	0.92	0.57	0.75	0.87	1.87	0.00	0.00	1.38
time (sec)	N/A	0.421	5.847	0.203	0.301	0.286	0.000	0.000	10.098

Problem 771	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	275	229	196	220	244	513	0	0	402
N.S.	1	0.83	0.71	0.80	0.89	1.87	0.00	0.00	1.46
time (sec)	N/A	0.424	7.917	0.283	0.312	0.282	0.000	0.000	8.954

Problem 772	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	238	202	182	179	221	458	0	0	349
N.S.	1	0.85	0.76	0.75	0.93	1.92	0.00	0.00	1.47
time (sec)	N/A	0.407	6.894	0.292	0.304	0.267	0.000	0.000	8.915

Problem 773	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	199	173	167	139	193	381	0	0	294
N.S.	1	0.87	0.84	0.70	0.97	1.91	0.00	0.00	1.48
time (sec)	N/A	0.385	6.202	0.279	0.319	0.267	0.000	0.000	8.723

Problem 774	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	160	146	144	118	166	372	0	0	267
N.S.	1	0.91	0.90	0.74	1.04	2.32	0.00	0.00	1.67
time (sec)	N/A	0.370	6.221	0.286	0.337	0.273	0.000	0.000	8.723

Problem 775	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	159	155	145	120	172	362	0	0	264
N.S.	1	0.97	0.91	0.75	1.08	2.28	0.00	0.00	1.66
time (sec)	N/A	0.373	5.535	0.271	0.308	0.278	0.000	0.000	8.834

Problem 776	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	195	182	153	149	195	390	0	0	305
N.S.	1	0.93	0.78	0.76	1.00	2.00	0.00	0.00	1.56
time (sec)	N/A	0.414	6.238	0.285	0.294	0.261	0.000	0.000	9.051

Problem 777	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	226	216	155	176	217	420	0	0	353
N.S.	1	0.96	0.69	0.78	0.96	1.86	0.00	0.00	1.56
time (sec)	N/A	0.427	5.945	0.248	0.311	0.271	0.000	0.000	9.298

Problem 778	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	273	250	142	196	245	446	0	0	399
N.S.	1	0.92	0.52	0.72	0.90	1.63	0.00	0.00	1.46
time (sec)	N/A	0.434	5.841	0.181	0.304	0.285	0.000	0.000	9.880

Problem 779	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	291	243	201	207	266	471	0	0	441
N.S.	1	0.84	0.69	0.71	0.91	1.62	0.00	0.00	1.52
time (sec)	N/A	0.420	7.271	0.257	0.314	0.278	0.000	0.000	9.172

Problem 780	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	252	216	193	168	240	402	0	0	386
N.S.	1	0.86	0.77	0.67	0.95	1.60	0.00	0.00	1.53
time (sec)	N/A	0.387	6.712	0.245	0.423	0.270	0.000	0.000	9.281

Problem 781	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	189	174	147	212	403	0	0	360
N.S.	1	0.89	0.82	0.69	1.00	1.89	0.00	0.00	1.69
time (sec)	N/A	0.392	6.504	0.216	0.394	0.269	0.000	0.000	9.351

Problem 782	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	211	196	174	139	212	395	0	0	360
N.S.	1	0.93	0.82	0.66	1.00	1.87	0.00	0.00	1.71
time (sec)	N/A	0.392	6.437	0.286	0.446	0.259	0.000	0.000	9.101

Problem 783	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	209	200	173	147	220	386	0	0	355
N.S.	1	0.96	0.83	0.70	1.05	1.85	0.00	0.00	1.70
time (sec)	N/A	0.387	5.758	0.211	0.421	0.259	0.000	0.000	9.054

Problem 784	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	245	227	211	178	241	408	0	0	394
N.S.	1	0.93	0.86	0.73	0.98	1.67	0.00	0.00	1.61
time (sec)	N/A	0.412	5.066	0.291	0.487	0.272	0.000	0.000	9.201

Problem 785	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	274	261	191	205	263	438	0	0	443
N.S.	1	0.95	0.70	0.75	0.96	1.60	0.00	0.00	1.62
time (sec)	N/A	0.437	6.144	0.230	0.392	0.272	0.000	0.000	9.491

Problem 786	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	311	295	187	232	291	464	0	0	490
N.S.	1	0.95	0.60	0.75	0.94	1.49	0.00	0.00	1.58
time (sec)	N/A	0.456	6.410	0.210	0.401	0.300	0.000	0.000	9.985

Problem 787	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	272	265	171	349	1343	604	0	0	0
N.S.	1	0.97	0.63	1.28	4.94	2.22	0.00	0.00	0.00
time (sec)	N/A	0.445	6.621	0.461	1.294	0.273	0.000	0.000	0.000

Problem 788	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	217	216	149	285	1085	547	0	0	0
N.S.	1	1.00	0.69	1.31	5.00	2.52	0.00	0.00	0.00
time (sec)	N/A	0.424	5.309	0.314	0.848	0.273	0.000	0.000	0.000

Problem 789	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	164	167	129	223	771	459	0	1747	0
N.S.	1	1.02	0.79	1.36	4.70	2.80	0.00	10.65	0.00
time (sec)	N/A	0.395	3.709	0.317	0.632	0.267	0.000	14.170	0.000

Problem 790	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	104	110	104	121	447	293	0	0	133
N.S.	1	1.06	1.00	1.16	4.30	2.82	0.00	0.00	1.28
time (sec)	N/A	0.355	2.415	0.322	0.546	0.263	0.000	0.000	10.464

Problem 791	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	109	115	110	321	0	332	0	0	266
N.S.	1	1.06	1.01	2.94	0.00	3.05	0.00	0.00	2.44
time (sec)	N/A	0.366	3.719	0.391	0.000	0.266	0.000	0.000	11.888

Problem 792	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	111	81	84	0	96	0	0	145
N.S.	1	1.09	0.79	0.82	0.00	0.94	0.00	0.00	1.42
time (sec)	N/A	0.375	4.939	0.424	0.000	0.243	0.000	0.000	1.232

Problem 793	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	155	167	110	98	0	113	0	0	171
N.S.	1	1.08	0.71	0.63	0.00	0.73	0.00	0.00	1.10
time (sec)	N/A	0.399	6.203	0.391	0.000	0.247	0.000	0.000	9.363

Problem 794	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	208	221	132	135	0	131	0	0	246
N.S.	1	1.06	0.63	0.65	0.00	0.63	0.00	0.00	1.18
time (sec)	N/A	0.417	6.609	0.424	0.000	0.247	0.000	0.000	10.130

Problem 795	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	279	265	456	412	1655	682	0	0	0
N.S.	1	0.95	1.63	1.48	5.93	2.44	0.00	0.00	0.00
time (sec)	N/A	0.432	11.541	0.347	2.239	0.284	0.000	0.000	0.000

Problem 796	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	226	216	206	350	1375	614	0	0	0
N.S.	1	0.96	0.91	1.55	6.08	2.72	0.00	0.00	0.00
time (sec)	N/A	0.410	7.766	0.367	1.129	0.280	0.000	0.000	0.000

Problem 797	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	157	159	181	179	853	433	0	0	0
N.S.	1	1.01	1.15	1.14	5.43	2.76	0.00	0.00	0.00
time (sec)	N/A	0.399	5.459	0.404	0.452	0.278	0.000	0.000	0.000

Problem 798	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	160	167	182	223	771	459	0	533	0
N.S.	1	1.04	1.14	1.39	4.82	2.87	0.00	3.33	0.00
time (sec)	N/A	0.389	4.374	0.383	0.473	0.281	0.000	1.528	0.000

Problem 799	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	170	149	497	599	438	0	0	0
N.S.	1	1.01	0.88	2.94	3.54	2.59	0.00	0.00	0.00
time (sec)	N/A	0.387	5.587	0.352	0.470	0.276	0.000	0.000	0.000

Problem 800	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	155	168	218	406	170	380	0	0	0
N.S.	1	1.08	1.41	2.62	1.10	2.45	0.00	0.00	0.00
time (sec)	N/A	0.406	7.251	0.393	0.416	0.286	0.000	0.000	0.000

Problem 801	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	111	111	90	154	98	0	0	190
N.S.	1	1.09	1.09	0.88	1.51	0.96	0.00	0.00	1.86
time (sec)	N/A	0.370	7.024	0.343	0.406	0.253	0.000	0.000	9.817

Problem 802	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	155	167	130	113	186	117	0	0	215
N.S.	1	1.08	0.84	0.73	1.20	0.75	0.00	0.00	1.39
time (sec)	N/A	0.417	6.887	0.405	0.415	0.245	0.000	0.000	10.302

Problem 803	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	208	219	131	136	260	136	0	0	290
N.S.	1	1.05	0.63	0.65	1.25	0.65	0.00	0.00	1.39
time (sec)	N/A	0.412	7.063	0.392	0.429	0.250	0.000	0.000	11.046

Problem 804	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	261	275	153	158	312	155	0	0	315
N.S.	1	1.05	0.59	0.61	1.20	0.59	0.00	0.00	1.21
time (sec)	N/A	0.456	7.699	0.362	0.432	0.251	0.000	0.000	12.508

Problem 805	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	288	267	231	478	2033	756	0	0	0
N.S.	1	0.93	0.80	1.66	7.06	2.62	0.00	0.00	0.00
time (sec)	N/A	0.452	11.105	0.475	4.577	0.278	0.000	0.000	0.000

Problem 806	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	210	148	221	1445	585	0	0	0
N.S.	1	0.99	0.69	1.04	6.78	2.75	0.00	0.00	0.00
time (sec)	N/A	0.413	8.533	0.395	0.942	0.276	0.000	0.000	0.000

Problem 807	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	222	216	222	350	1375	614	0	0	0
N.S.	1	0.97	1.00	1.58	6.19	2.77	0.00	0.00	0.00
time (sec)	N/A	0.414	7.934	0.375	1.098	0.289	0.000	0.000	0.000

Problem 808	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	217	216	198	285	1087	547	0	0	0
N.S.	1	1.00	0.91	1.31	5.01	2.52	0.00	0.00	0.00
time (sec)	N/A	0.411	5.744	0.343	0.583	0.279	0.000	0.000	0.000

Problem 809	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	227	221	170	565	993	530	0	0	0
N.S.	1	0.97	0.75	2.49	4.37	2.33	0.00	0.00	0.00
time (sec)	N/A	0.415	7.328	0.423	0.506	0.278	0.000	0.000	0.000

Problem 810	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	226	222	179	667	795	493	0	0	0
N.S.	1	0.98	0.79	2.95	3.52	2.18	0.00	0.00	0.00
time (sec)	N/A	0.430	7.633	0.351	0.485	0.265	0.000	0.000	0.000

Problem 811	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	203	218	203	507	214	418	0	0	0
N.S.	1	1.07	1.00	2.50	1.05	2.06	0.00	0.00	0.00
time (sec)	N/A	0.422	17.953	0.351	0.429	0.270	0.000	0.000	0.000

Problem 812	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	111	113	106	166	104	0	0	192
N.S.	1	1.09	1.11	1.04	1.63	1.02	0.00	0.00	1.88
time (sec)	N/A	0.371	6.898	0.384	0.430	0.245	0.000	0.000	9.961

Problem 813	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	155	167	130	119	198	125	0	0	217
N.S.	1	1.08	0.84	0.77	1.28	0.81	0.00	0.00	1.40
time (sec)	N/A	0.405	7.025	0.353	0.432	0.240	0.000	0.000	10.562

Problem 814	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	208	221	156	156	276	146	0	0	292
N.S.	1	1.06	0.75	0.75	1.33	0.70	0.00	0.00	1.40
time (sec)	N/A	0.422	16.384	0.417	0.610	0.247	0.000	0.000	12.413

Problem 815	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	261	275	577	169	332	167	0	0	191
N.S.	1	1.05	2.21	0.65	1.27	0.64	0.00	0.00	0.73
time (sec)	N/A	0.445	19.166	0.378	0.604	0.245	0.000	0.000	13.404

Problem 816	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	350	318	241	604	2603	882	0	0	0
N.S.	1	0.91	0.69	1.73	7.44	2.52	0.00	0.00	0.00
time (sec)	N/A	0.449	9.429	0.495	19.778	0.288	0.000	0.000	0.000

Problem 817	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	267	261	159	269	1901	693	0	0	0
N.S.	1	0.98	0.60	1.01	7.12	2.60	0.00	0.00	0.00
time (sec)	N/A	0.420	8.561	0.405	2.745	0.275	0.000	0.000	0.000

Problem 818	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	284	267	510	478	2033	756	0	0	0
N.S.	1	0.94	1.80	1.68	7.16	2.66	0.00	0.00	0.00
time (sec)	N/A	0.440	11.712	0.492	5.957	0.286	0.000	0.000	0.000

Problem 819	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	279	265	458	412	1657	682	0	0	0
N.S.	1	0.95	1.64	1.48	5.94	2.44	0.00	0.00	0.00
time (sec)	N/A	0.439	11.533	0.319	2.784	0.287	0.000	0.000	0.000

Problem 820	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F(-1)	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	272	265	221	349	1345	604	0	0	0
N.S.	1	0.97	0.81	1.28	4.94	2.22	0.00	0.00	0.00
time (sec)	N/A	0.431	7.961	0.354	1.058	0.274	0.000	0.000	0.000

Problem 821	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	283	270	183	627	1329	590	0	0	0
N.S.	1	0.95	0.65	2.22	4.70	2.08	0.00	0.00	0.00
time (sec)	N/A	0.441	8.512	0.403	0.710	0.270	0.000	0.000	0.000

Problem 822	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	285	273	196	731	1176	571	0	0	0
N.S.	1	0.96	0.69	2.56	4.13	2.00	0.00	0.00	0.00
time (sec)	N/A	0.459	9.274	0.378	0.555	0.284	0.000	0.000	0.000

Problem 823	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	283	274	528	833	936	515	0	0	0
N.S.	1	0.97	1.87	2.94	3.31	1.82	0.00	0.00	0.00
time (sec)	N/A	0.442	19.930	0.368	0.496	0.274	0.000	0.000	0.000

Problem 824	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	A	B	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	251	268	570	598	246	434	0	0	0
N.S.	1	1.07	2.27	2.38	0.98	1.73	0.00	0.00	0.00
time (sec)	N/A	0.441	19.877	0.343	0.442	0.278	0.000	0.000	0.000

Problem 825	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	111	111	106	166	104	0	0	192
N.S.	1	1.09	1.09	1.04	1.63	1.02	0.00	0.00	1.88
time (sec)	N/A	0.371	6.957	0.359	0.444	0.260	0.000	0.000	10.600

Problem 826	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	155	167	417	119	198	125	0	0	217
N.S.	1	1.08	2.69	0.77	1.28	0.81	0.00	0.00	1.40
time (sec)	N/A	0.405	18.824	0.369	0.635	0.248	0.000	0.000	10.820

Problem 827	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	208	221	495	156	276	146	0	0	167
N.S.	1	1.06	2.38	0.75	1.33	0.70	0.00	0.00	0.80
time (sec)	N/A	0.442	19.123	0.341	0.666	0.250	0.000	0.000	13.008

Problem 828	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	261	275	577	169	332	167	0	0	191
N.S.	1	1.05	2.21	0.65	1.27	0.64	0.00	0.00	0.73
time (sec)	N/A	0.451	19.377	0.377	1.087	0.254	0.000	0.000	12.602

Problem 829	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	F(-1)	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	314	329	655	206	412	188	0	0	229
N.S.	1	1.05	2.09	0.66	1.31	0.60	0.00	0.00	0.73
time (sec)	N/A	0.479	19.832	0.374	0.556	0.245	0.000	0.000	13.227

Problem 830	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	228	222	162	566	1319	548	0	0	0
N.S.	1	0.97	0.71	2.48	5.79	2.40	0.00	0.00	0.00
time (sec)	N/A	0.430	7.738	0.393	0.656	0.284	0.000	0.000	0.000

Problem 831	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	171	136	499	898	456	0	0	0
N.S.	1	1.01	0.80	2.95	5.31	2.70	0.00	0.00	0.00
time (sec)	N/A	0.402	4.597	0.382	0.473	0.272	0.000	0.000	0.000

Problem 832	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	116	119	323	138	348	0	0	250
N.S.	1	1.05	1.08	2.94	1.25	3.16	0.00	0.00	2.27
time (sec)	N/A	0.356	3.018	0.413	0.395	0.266	0.000	0.000	11.562

Problem 833	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	92	101	50	92	124	114	0	0	143
N.S.	1	1.10	0.54	1.00	1.35	1.24	0.00	0.00	1.55
time (sec)	N/A	0.345	2.489	0.474	0.391	0.248	0.000	0.000	0.804

Problem 834	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	157	163	97	110	0	147	0	0	146
N.S.	1	1.04	0.62	0.70	0.00	0.94	0.00	0.00	0.93
time (sec)	N/A	0.396	4.824	0.407	0.000	0.254	0.000	0.000	0.728

Problem 835	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	217	116	148	0	161	0	0	186
N.S.	1	1.02	0.54	0.69	0.00	0.76	0.00	0.00	0.87
time (sec)	N/A	0.447	6.241	0.467	0.000	0.257	0.000	0.000	9.339

Problem 836	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	287	275	160	733	1373	589	0	0	0
N.S.	1	0.96	0.56	2.55	4.78	2.05	0.00	0.00	0.00
time (sec)	N/A	0.461	8.382	0.382	0.828	0.273	0.000	0.000	0.000

Problem 837	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	229	224	141	669	0	511	0	0	0
N.S.	1	0.98	0.62	2.92	0.00	2.23	0.00	0.00	0.00
time (sec)	N/A	0.425	6.781	0.407	0.000	0.268	0.000	0.000	0.000

Problem 838	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	157	170	137	408	168	400	0	0	0
N.S.	1	1.08	0.87	2.60	1.07	2.55	0.00	0.00	0.00
time (sec)	N/A	0.416	5.233	0.340	0.414	0.275	0.000	0.000	0.000

Problem 839	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	104	113	81	103	0	92	0	0	195
N.S.	1	1.09	0.78	0.99	0.00	0.88	0.00	0.00	1.88
time (sec)	N/A	0.369	3.021	0.438	0.000	0.257	0.000	0.000	9.691

Problem 840	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	152	162	97	152	0	145	0	0	170
N.S.	1	1.07	0.64	1.00	0.00	0.95	0.00	0.00	1.12
time (sec)	N/A	0.409	3.167	0.398	0.000	0.284	0.000	0.000	9.356

Problem 841	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	152	160	79	113	200	150	0	0	198
N.S.	1	1.05	0.52	0.74	1.32	0.99	0.00	0.00	1.30
time (sec)	N/A	0.408	5.922	0.399	0.420	0.244	0.000	0.000	9.377

Problem 842	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	269	273	152	199	0	181	0	0	196
N.S.	1	1.01	0.57	0.74	0.00	0.67	0.00	0.00	0.73
time (sec)	N/A	0.458	6.749	0.386	0.000	0.264	0.000	0.000	9.536

Problem 843	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	343	327	228	899	0	610	0	0	0
N.S.	1	0.95	0.66	2.62	0.00	1.78	0.00	0.00	0.00
time (sec)	N/A	0.503	9.772	0.444	0.000	0.276	0.000	0.000	0.000

Problem 844	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	284	276	191	835	1025	533	0	0	0
N.S.	1	0.97	0.67	2.94	3.61	1.88	0.00	0.00	0.00
time (sec)	N/A	0.451	7.674	0.383	0.571	0.274	0.000	0.000	0.000

Problem 845	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	205	220	167	509	216	436	0	0	0
N.S.	1	1.07	0.81	2.48	1.05	2.13	0.00	0.00	0.00
time (sec)	N/A	0.414	7.654	0.410	0.422	0.285	0.000	0.000	0.000

Problem 846	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	104	113	92	92	153	98	0	0	240
N.S.	1	1.09	0.88	0.88	1.47	0.94	0.00	0.00	2.31
time (sec)	N/A	0.379	5.365	0.408	0.427	0.250	0.000	0.000	10.101

Problem 847	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	157	169	99	127	0	112	0	0	246
N.S.	1	1.08	0.63	0.81	0.00	0.71	0.00	0.00	1.57
time (sec)	N/A	0.404	3.526	0.427	0.000	0.249	0.000	0.000	10.323

Problem 848	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	212	216	115	186	0	161	0	0	246
N.S.	1	1.02	0.54	0.88	0.00	0.76	0.00	0.00	1.16
time (sec)	N/A	0.411	4.514	0.388	0.000	0.279	0.000	0.000	9.877

Problem 849	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	218	225	152	199	0	181	0	0	249
N.S.	1	1.03	0.70	0.91	0.00	0.83	0.00	0.00	1.14
time (sec)	N/A	0.426	6.656	0.391	0.000	0.251	0.000	0.000	9.971

Problem 850	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	206	218	90	124	334	185	0	0	249
N.S.	1	1.06	0.44	0.60	1.62	0.90	0.00	0.00	1.21
time (sec)	N/A	0.414	7.038	0.396	0.591	0.255	0.000	0.000	10.478

Problem 851	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	150	165	122	0	0	0	0	0	0
N.S.	1	1.10	0.81	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.391	1.133	0.000	0.000	0.000	0.000	0.000	0.000

Problem 852	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	147	152	133	0	0	0	0	0	0
N.S.	1	1.03	0.90	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.412	5.836	0.000	0.000	0.000	0.000	0.000	0.000

Problem 853	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	A	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	31	92	0	54	66	0	90
N.S.	1	1.00	0.94	2.79	0.00	1.64	2.00	0.00	2.73
time (sec)	N/A	0.299	5.453	0.738	0.000	0.241	0.638	0.000	8.459

Problem 854	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	104	109	93	154	0	84	296	185	159
N.S.	1	1.05	0.89	1.48	0.00	0.81	2.85	1.78	1.53
time (sec)	N/A	0.481	2.060	0.235	0.000	0.244	0.261	0.491	8.727

Problem 855	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	147	154	141	131	142	619	0	0	245
N.S.	1	1.05	0.96	0.89	0.97	4.21	0.00	0.00	1.67
time (sec)	N/A	0.637	2.517	0.314	0.330	0.250	0.000	0.000	1.588

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. The column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [600] had the largest ratio of [.878788000000000014]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	8	8	1.00	32	0.250
2	A	6	6	1.00	30	0.200
3	A	4	4	1.00	24	0.167
4	A	8	8	1.05	30	0.267
5	A	7	7	1.05	32	0.219
6	A	10	10	1.03	32	0.312
7	A	12	12	1.02	32	0.375
8	A	15	15	1.02	32	0.469
9	A	10	10	1.04	34	0.294
10	A	8	8	0.93	32	0.250
11	A	6	6	0.88	26	0.231
12	A	10	10	1.03	32	0.312
13	A	10	10	1.03	34	0.294
14	A	10	10	1.06	34	0.294
15	A	12	12	1.08	34	0.353
16	A	15	15	1.09	34	0.441
17	A	12	12	1.05	34	0.353
18	A	10	10	0.93	32	0.312
19	A	8	8	0.90	26	0.308
20	A	13	13	1.02	32	0.406
21	A	12	12	1.02	34	0.353
22	A	14	14	1.02	34	0.412

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
23	A	13	13	1.08	34	0.382
24	A	17	17	1.10	34	0.500
25	A	18	18	1.08	34	0.529
26	A	16	16	1.07	34	0.471
27	A	12	12	0.93	32	0.375
28	A	10	10	0.91	26	0.385
29	A	16	16	1.01	32	0.500
30	A	15	15	1.01	34	0.441
31	A	15	15	1.04	34	0.441
32	A	16	16	1.01	34	0.471
33	A	15	15	1.09	34	0.441
34	A	17	17	1.10	34	0.500
35	A	21	21	1.10	34	0.618
36	A	8	8	0.95	34	0.235
37	A	6	6	0.98	34	0.176
38	A	7	7	0.96	32	0.219
39	A	3	3	1.00	26	0.115
40	A	7	7	1.08	32	0.219
41	A	10	10	1.00	34	0.294
42	A	12	12	0.97	34	0.353
43	A	15	15	0.97	34	0.441
44	A	9	9	1.04	34	0.265
45	A	11	11	1.03	34	0.324
46	A	5	5	1.08	32	0.156
47	A	5	5	0.99	26	0.192
48	A	10	10	1.15	32	0.312
49	A	13	13	1.04	34	0.382
50	A	15	15	1.06	34	0.441
51	A	12	12	1.06	34	0.353
52	A	13	13	1.07	34	0.382
53	A	7	7	1.09	34	0.206
54	A	7	7	1.05	32	0.219
55	A	7	7	0.99	26	0.269
56	A	13	13	1.17	32	0.406

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
57	A	16	16	1.08	34	0.471
58	A	18	18	1.08	34	0.529
59	A	17	17	1.10	34	0.500
60	A	11	11	1.14	34	0.324
61	A	11	11	1.06	34	0.324
62	A	9	9	1.03	32	0.281
63	A	9	9	0.99	26	0.346
64	A	16	16	1.18	32	0.500
65	A	19	19	1.10	34	0.559
66	A	22	22	1.09	34	0.647
67	A	14	13	1.10	36	0.361
68	A	11	10	1.05	36	0.278
69	A	8	7	1.02	34	0.206
70	A	6	5	1.03	28	0.179
71	A	9	8	1.03	34	0.235
72	A	12	11	1.09	36	0.306
73	A	15	14	1.09	36	0.389
74	A	18	17	1.12	36	0.472
75	A	14	13	1.08	36	0.361
76	A	10	9	1.00	34	0.265
77	A	8	7	0.99	28	0.250
78	A	12	11	1.02	34	0.324
79	A	12	11	1.06	36	0.306
80	A	15	14	1.06	36	0.389
81	A	18	17	1.10	36	0.472
82	A	17	16	1.08	36	0.444
83	A	12	11	0.98	34	0.324
84	A	10	9	0.97	28	0.321
85	A	15	14	1.01	34	0.412
86	A	15	14	1.05	36	0.389
87	A	15	14	1.05	36	0.389
88	A	18	17	1.07	36	0.472
89	A	21	20	1.09	36	0.556
90	A	14	13	1.06	36	0.361

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
91	A	11	10	1.04	36	0.278
92	A	8	7	1.02	34	0.206
93	A	6	5	1.02	28	0.179
94	A	12	11	1.10	34	0.324
95	A	15	14	1.10	36	0.389
96	A	18	17	1.08	36	0.472
97	A	14	13	1.05	36	0.361
98	A	11	10	0.99	36	0.278
99	A	8	7	1.06	34	0.206
100	A	8	7	0.98	28	0.250
101	A	15	14	1.07	34	0.412
102	A	18	17	1.07	36	0.472
103	A	21	20	1.07	36	0.556
104	A	17	16	1.07	36	0.444
105	A	14	13	1.04	36	0.361
106	A	11	10	1.07	36	0.278
107	A	10	9	1.03	34	0.265
108	A	10	9	0.99	28	0.321
109	A	18	17	1.10	34	0.500
110	A	21	20	1.09	36	0.556
111	A	23	22	1.08	36	0.611
112	A	12	11	1.00	34	0.324
113	A	10	9	1.00	34	0.265
114	A	8	7	1.00	34	0.206
115	A	6	5	1.00	34	0.147
116	A	6	5	1.00	34	0.147
117	A	9	8	1.00	34	0.235
118	A	11	10	1.00	34	0.294
119	A	15	14	1.02	36	0.389
120	A	13	12	1.03	36	0.333
121	A	11	10	1.04	36	0.278
122	A	9	8	1.03	36	0.222
123	A	9	8	1.00	36	0.222
124	A	10	9	1.03	36	0.250

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
125	A	12	11	1.06	36	0.306
126	A	15	14	1.05	36	0.389
127	A	16	15	1.04	36	0.417
128	A	14	13	1.06	36	0.361
129	A	12	11	1.05	36	0.306
130	A	12	11	1.02	36	0.306
131	A	12	11	1.03	36	0.306
132	A	13	12	1.06	36	0.333
133	A	16	15	1.08	36	0.417
134	A	20	19	0.88	36	0.528
135	A	18	17	0.86	36	0.472
136	A	15	14	0.87	36	0.389
137	A	15	14	0.87	36	0.389
138	A	18	17	0.89	36	0.472
139	A	20	19	0.88	36	0.528
140	A	20	19	0.91	36	0.528
141	A	18	17	0.88	36	0.472
142	A	17	16	0.88	36	0.444
143	A	17	16	0.85	36	0.444
144	A	20	19	0.90	36	0.528
145	A	22	21	0.91	36	0.583
146	A	25	24	0.95	36	0.667
147	A	24	23	0.93	36	0.639
148	A	21	20	0.93	36	0.556
149	A	20	19	0.93	36	0.528
150	A	20	19	0.92	36	0.528
151	A	20	19	0.91	36	0.528
152	A	24	23	0.93	36	0.639
153	A	25	24	0.93	36	0.667
154	A	15	14	1.06	38	0.368
155	A	12	11	1.04	38	0.289
156	A	9	8	1.00	38	0.211
157	A	7	6	1.00	38	0.158
158	A	10	9	1.05	38	0.237

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
159	A	13	12	1.10	38	0.316
160	A	16	15	1.13	38	0.395
161	A	18	17	1.08	38	0.447
162	A	15	14	1.04	38	0.368
163	A	12	11	1.03	38	0.289
164	A	12	11	1.00	38	0.289
165	A	10	9	1.02	38	0.237
166	A	13	12	1.07	38	0.316
167	A	16	15	1.10	38	0.395
168	A	19	18	1.12	38	0.474
169	A	21	20	1.07	38	0.526
170	A	18	17	1.05	38	0.447
171	A	15	14	1.03	38	0.368
172	A	15	14	1.03	38	0.368
173	A	15	14	1.00	38	0.368
174	A	13	12	1.04	38	0.316
175	A	15	14	1.07	38	0.368
176	A	19	18	1.09	38	0.474
177	A	22	21	1.10	38	0.553
178	A	15	14	1.00	46	0.304
179	A	15	14	1.04	38	0.368
180	A	12	11	1.03	38	0.289
181	A	7	6	1.00	38	0.158
182	A	10	9	1.01	38	0.237
183	A	13	12	1.04	38	0.316
184	A	16	15	1.07	38	0.395
185	A	15	14	1.04	38	0.368
186	A	10	9	1.03	38	0.237
187	A	10	9	1.02	38	0.237
188	A	13	12	1.03	38	0.316
189	A	16	15	1.05	38	0.395
190	A	18	17	1.06	38	0.447
191	A	13	12	1.06	38	0.316
192	A	13	12	1.06	38	0.316

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
193	A	13	12	1.06	38	0.316
194	A	16	15	1.05	38	0.395
195	A	19	18	1.07	38	0.474
196	A	9	8	0.69	28	0.286
197	A	14	13	0.80	36	0.361
198	A	11	10	0.74	34	0.294
199	A	9	8	0.70	28	0.286
200	A	14	13	0.75	34	0.382
201	A	17	16	0.78	36	0.444
202	A	9	8	0.70	28	0.286
203	A	9	8	0.70	28	0.286
204	A	15	14	1.09	34	0.412
205	A	12	11	1.30	34	0.324
206	A	10	9	1.24	34	0.265
207	A	7	6	1.39	32	0.188
208	A	8	7	0.98	34	0.206
209	A	11	10	1.00	34	0.294
210	A	14	13	1.04	34	0.382
211	A	17	16	1.05	34	0.471
212	A	18	17	1.03	36	0.472
213	A	15	14	1.04	36	0.389
214	A	12	11	1.00	36	0.306
215	A	15	14	1.04	36	0.389
216	A	18	17	1.04	36	0.472
217	A	21	20	1.05	36	0.556
218	A	12	11	1.00	34	0.324
219	A	14	13	1.06	34	0.382
220	A	11	10	1.05	34	0.294
221	A	8	7	1.05	32	0.219
222	A	6	5	1.08	26	0.192
223	A	8	7	1.06	32	0.219
224	A	10	9	1.10	34	0.265
225	A	12	11	1.11	34	0.324
226	A	23	22	1.10	36	0.611

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
227	A	20	19	1.13	36	0.528
228	A	17	16	1.15	36	0.444
229	A	14	13	1.11	36	0.361
230	A	17	16	1.13	36	0.444
231	A	20	19	1.13	36	0.528
232	A	8	8	1.00	29	0.276
233	A	6	6	1.00	27	0.222
234	A	4	4	1.00	21	0.190
235	A	8	8	1.05	27	0.296
236	A	7	7	1.05	29	0.241
237	A	10	10	1.03	29	0.345
238	A	12	12	1.02	29	0.414
239	A	15	15	1.02	29	0.517
240	A	11	11	1.08	31	0.355
241	A	8	8	1.00	29	0.276
242	A	6	6	1.00	23	0.261
243	A	7	7	1.03	29	0.241
244	A	7	7	1.03	31	0.226
245	A	10	10	1.03	31	0.323
246	A	12	12	1.03	31	0.387
247	A	15	15	1.01	31	0.484
248	A	13	13	1.07	31	0.419
249	A	10	10	1.00	29	0.345
250	A	8	8	1.00	23	0.348
251	A	11	11	1.02	29	0.379
252	A	10	10	1.02	31	0.323
253	A	11	11	1.02	31	0.355
254	A	13	13	1.05	31	0.419
255	A	17	17	1.07	31	0.548
256	A	18	18	1.04	31	0.581
257	A	15	15	1.05	31	0.484
258	A	12	12	1.00	29	0.414
259	A	10	10	1.00	23	0.435
260	A	14	14	1.01	29	0.483

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
261	A	13	13	1.01	31	0.419
262	A	13	13	1.03	31	0.419
263	A	13	13	1.01	31	0.419
264	A	15	15	1.06	31	0.484
265	A	17	17	1.05	31	0.548
266	A	21	21	1.05	31	0.677
267	A	12	11	1.13	31	0.355
268	A	10	9	1.07	31	0.290
269	A	8	8	0.94	29	0.276
270	A	4	4	1.00	23	0.174
271	A	6	6	1.02	29	0.207
272	A	8	8	1.08	31	0.258
273	A	12	12	1.12	31	0.387
274	A	14	14	1.14	31	0.452
275	A	13	12	1.08	31	0.387
276	A	10	9	1.09	31	0.290
277	A	6	6	1.10	29	0.207
278	A	6	6	1.10	23	0.261
279	A	8	8	1.18	29	0.276
280	A	10	10	1.16	31	0.323
281	A	13	13	1.15	31	0.419
282	A	16	15	1.10	31	0.484
283	A	12	11	1.14	31	0.355
284	A	9	9	1.14	31	0.290
285	A	8	8	1.12	29	0.276
286	A	8	8	1.13	23	0.348
287	A	11	11	1.20	29	0.379
288	A	13	13	1.16	31	0.419
289	A	17	17	1.15	31	0.548
290	A	16	15	1.15	31	0.484
291	A	13	13	1.16	31	0.419
292	A	11	11	1.15	31	0.355
293	A	10	10	1.13	29	0.345
294	A	10	10	1.13	23	0.435

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
295	A	14	14	1.19	29	0.483
296	A	16	16	1.15	31	0.516
297	A	19	19	1.14	31	0.613
298	A	5	5	1.00	34	0.147
299	A	4	4	1.00	34	0.118
300	A	3	3	1.00	32	0.094
301	A	2	2	1.00	26	0.077
302	A	4	4	1.17	32	0.125
303	A	4	4	1.00	34	0.118
304	A	8	8	1.07	34	0.235
305	A	6	6	0.94	34	0.176
306	A	14	13	1.13	34	0.382
307	A	11	10	1.07	34	0.294
308	A	8	8	0.95	34	0.235
309	A	5	5	1.00	32	0.156
310	A	5	5	1.00	26	0.192
311	A	7	7	1.01	32	0.219
312	A	9	9	1.08	34	0.265
313	A	13	13	1.12	34	0.382
314	A	5	5	1.00	21	0.238
315	A	4	4	1.00	28	0.143
316	A	6	6	1.11	23	0.261
317	A	18	17	1.12	33	0.515
318	A	15	14	0.97	33	0.424
319	A	12	11	0.99	31	0.355
320	A	10	9	0.84	25	0.360
321	A	13	12	0.90	31	0.387
322	A	17	16	0.93	33	0.485
323	A	20	19	1.00	33	0.576
324	A	23	22	1.03	33	0.667
325	A	17	16	0.98	33	0.485
326	A	14	13	0.91	31	0.419
327	A	12	11	0.87	25	0.440
328	A	16	15	0.91	31	0.484

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
329	A	17	16	0.93	33	0.485
330	A	20	19	1.00	33	0.576
331	A	23	22	1.01	33	0.667
332	A	19	18	0.98	33	0.545
333	A	16	15	1.01	31	0.484
334	A	14	13	0.90	25	0.520
335	A	19	18	0.93	31	0.581
336	A	20	19	0.95	33	0.576
337	A	20	19	0.97	33	0.576
338	A	23	22	1.00	33	0.667
339	A	26	25	1.04	33	0.758
340	A	13	12	0.80	27	0.444
341	A	14	13	1.06	27	0.481
342	A	14	13	1.07	27	0.481
343	A	16	15	1.04	33	0.455
344	A	13	12	0.95	33	0.364
345	A	10	9	0.88	31	0.290
346	A	8	7	0.81	25	0.280
347	A	13	12	0.90	31	0.387
348	A	17	16	0.97	33	0.485
349	A	20	19	1.03	33	0.576
350	A	16	15	1.03	33	0.455
351	A	13	12	1.11	33	0.364
352	A	10	9	1.07	31	0.290
353	A	10	9	1.04	25	0.360
354	A	17	16	1.14	31	0.516
355	A	20	19	1.16	33	0.576
356	A	23	22	1.14	33	0.667
357	A	19	18	1.06	33	0.545
358	A	15	14	1.14	33	0.424
359	A	13	12	1.14	33	0.364
360	A	12	11	1.13	31	0.355
361	A	12	11	1.11	25	0.440
362	A	20	19	1.21	31	0.613

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
363	A	23	22	1.19	33	0.667
364	A	26	25	1.16	33	0.758
365	A	12	11	1.11	28	0.393
366	A	12	11	1.05	28	0.393
367	A	15	14	0.91	34	0.412
368	A	11	10	1.02	28	0.357
369	A	18	17	1.14	34	0.500
370	A	8	7	0.83	27	0.259
371	A	11	10	0.98	27	0.370
372	A	13	12	1.05	27	0.444
373	A	6	5	0.78	27	0.185
374	A	6	5	0.78	27	0.185
375	A	4	3	1.00	15	0.200
376	A	4	3	1.00	17	0.176
377	A	8	7	1.00	25	0.280
378	A	19	18	0.92	31	0.581
379	A	18	17	0.91	31	0.548
380	A	16	15	0.90	31	0.484
381	A	13	12	0.89	31	0.387
382	A	13	12	0.89	31	0.387
383	A	16	15	0.90	31	0.484
384	A	18	17	0.91	31	0.548
385	A	23	22	0.89	33	0.667
386	A	21	20	0.88	33	0.606
387	A	19	18	0.87	33	0.545
388	A	17	16	0.85	33	0.485
389	A	15	14	0.83	33	0.424
390	A	16	15	0.83	33	0.455
391	A	18	17	0.85	33	0.515
392	A	24	23	0.85	33	0.697
393	A	22	21	0.84	33	0.636
394	A	20	19	0.82	33	0.576
395	A	19	18	0.81	33	0.545
396	A	20	19	0.81	33	0.576

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
397	A	20	19	0.82	33	0.576
398	A	23	22	0.89	33	0.667
399	A	20	19	0.86	33	0.576
400	A	16	15	0.82	33	0.455
401	A	16	15	0.82	33	0.455
402	A	20	19	0.86	33	0.576
403	A	23	22	0.89	33	0.667
404	A	24	23	0.87	33	0.697
405	A	21	20	0.86	33	0.606
406	A	21	20	0.85	33	0.606
407	A	21	20	0.86	33	0.606
408	A	24	23	0.87	33	0.697
409	A	27	26	0.88	33	0.788
410	A	27	26	0.88	33	0.788
411	A	24	23	0.87	33	0.697
412	A	24	23	0.87	33	0.697
413	A	24	23	0.87	33	0.697
414	A	24	23	0.87	33	0.697
415	A	27	26	0.88	33	0.788
416	A	15	14	1.01	36	0.389
417	A	15	14	1.01	36	0.389
418	A	13	12	1.00	36	0.333
419	A	13	12	1.00	36	0.333
420	A	15	14	1.01	36	0.389
421	A	15	14	1.01	36	0.389
422	A	21	20	0.89	36	0.556
423	A	18	17	0.86	36	0.472
424	A	17	16	0.86	36	0.444
425	A	17	16	0.86	36	0.444
426	A	21	20	0.89	36	0.556
427	A	12	11	1.00	35	0.314
428	A	9	8	0.99	35	0.229
429	A	13	12	1.06	35	0.343
430	A	11	10	1.10	35	0.286

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
431	A	14	13	1.12	35	0.371
432	A	17	16	1.16	35	0.457
433	A	20	19	1.13	35	0.543
434	A	15	14	1.01	35	0.400
435	A	12	11	0.98	35	0.314
436	A	9	8	0.98	35	0.229
437	A	9	8	0.96	35	0.229
438	A	14	13	1.15	35	0.371
439	A	17	16	1.12	35	0.457
440	A	20	19	1.13	35	0.543
441	A	23	22	1.13	35	0.629
442	A	18	17	1.01	35	0.486
443	A	15	14	1.03	35	0.400
444	A	12	11	0.99	35	0.314
445	A	12	11	0.98	35	0.314
446	A	12	11	1.00	35	0.314
447	A	17	16	1.15	35	0.457
448	A	20	19	1.13	35	0.543
449	A	23	22	1.13	35	0.629
450	A	26	25	1.11	35	0.714
451	A	12	11	1.01	43	0.256
452	A	9	8	1.00	35	0.229
453	A	13	12	1.00	35	0.343
454	A	8	7	1.00	35	0.200
455	A	11	10	1.04	35	0.286
456	A	14	13	1.08	35	0.371
457	A	17	16	1.10	35	0.457
458	A	9	8	1.16	35	0.229
459	A	11	10	1.19	35	0.286
460	A	11	10	1.16	35	0.286
461	A	14	13	1.19	35	0.371
462	A	17	16	1.14	35	0.457
463	A	12	11	1.22	35	0.314
464	A	14	13	1.20	35	0.371

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
465	A	14	13	1.22	35	0.371
466	A	14	13	1.21	35	0.371
467	A	17	16	1.20	35	0.457
468	A	20	19	1.18	35	0.543
469	A	6	5	0.95	38	0.132
470	A	8	7	0.97	38	0.184
471	A	6	5	0.97	38	0.132
472	A	11	10	0.99	38	0.263
473	A	12	11	0.71	25	0.440
474	A	12	11	0.71	25	0.440
475	A	10	9	0.65	25	0.360
476	A	10	9	0.65	25	0.360
477	A	7	6	0.71	27	0.222
478	A	10	9	0.79	26	0.346
479	A	16	15	1.03	31	0.484
480	A	14	13	1.06	31	0.419
481	A	11	10	1.02	31	0.323
482	A	8	7	1.00	29	0.241
483	A	10	9	0.96	31	0.290
484	A	12	11	1.04	31	0.355
485	A	16	15	1.06	31	0.484
486	A	19	18	1.07	31	0.581
487	A	7	6	1.00	33	0.182
488	A	7	6	1.00	33	0.182
489	A	7	6	1.00	33	0.182
490	A	7	6	1.00	33	0.182
491	A	7	6	1.00	33	0.182
492	A	7	6	1.00	33	0.182
493	A	7	6	1.00	31	0.194
494	A	18	17	1.06	31	0.548
495	A	15	14	1.10	31	0.452
496	A	12	11	1.07	31	0.355
497	A	9	8	1.04	29	0.276
498	A	7	6	1.01	23	0.261

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
499	A	11	10	1.03	29	0.345
500	A	15	14	1.04	31	0.452
501	A	17	16	1.09	31	0.516
502	A	12	11	1.00	34	0.324
503	A	10	9	1.00	34	0.265
504	A	8	7	1.00	34	0.206
505	A	8	7	1.00	34	0.206
506	A	10	9	1.00	34	0.265
507	A	13	12	1.00	34	0.353
508	A	13	12	1.04	36	0.333
509	A	11	10	1.03	36	0.278
510	A	11	10	1.00	36	0.278
511	A	12	11	1.03	36	0.306
512	A	14	13	1.06	36	0.361
513	A	16	15	1.06	36	0.417
514	A	14	13	1.05	36	0.361
515	A	14	13	1.02	36	0.361
516	A	14	13	1.03	36	0.361
517	A	15	14	1.05	36	0.389
518	A	17	16	1.08	36	0.444
519	A	21	20	0.89	36	0.556
520	A	20	19	0.87	36	0.528
521	A	18	17	0.87	36	0.472
522	A	18	17	0.87	36	0.472
523	A	20	19	0.88	36	0.528
524	A	23	22	0.87	36	0.611
525	A	22	21	0.90	36	0.583
526	A	20	19	0.87	36	0.528
527	A	21	20	0.88	36	0.556
528	A	21	20	0.87	36	0.556
529	A	24	23	0.90	36	0.639
530	A	26	25	0.93	36	0.694
531	A	23	22	0.92	36	0.611
532	A	23	22	0.92	36	0.611

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
533	A	23	22	0.93	36	0.611
534	A	23	22	0.93	36	0.611
535	A	26	25	0.94	36	0.694
536	A	15	14	1.10	38	0.368
537	A	12	11	1.05	38	0.289
538	A	9	8	1.01	38	0.211
539	A	11	10	0.88	38	0.263
540	A	14	13	0.93	38	0.342
541	A	18	17	1.10	38	0.447
542	A	15	14	1.07	38	0.368
543	A	12	11	1.03	38	0.289
544	A	14	13	0.90	38	0.342
545	A	14	13	0.93	38	0.342
546	A	17	16	0.95	38	0.421
547	A	21	20	1.08	38	0.526
548	A	17	16	1.07	38	0.421
549	A	15	14	1.04	38	0.368
550	A	17	16	0.92	38	0.421
551	A	17	16	0.94	38	0.421
552	A	17	16	0.95	38	0.421
553	A	20	19	0.98	38	0.500
554	A	15	14	1.04	38	0.368
555	A	12	11	1.02	38	0.289
556	A	9	8	1.01	38	0.211
557	A	14	13	0.92	38	0.342
558	A	15	14	1.03	38	0.368
559	A	12	11	1.02	38	0.289
560	A	12	11	1.03	38	0.289
561	A	17	16	0.95	38	0.421
562	A	18	17	1.05	38	0.447
563	A	15	14	1.06	38	0.368
564	A	15	14	1.06	38	0.368
565	A	15	14	1.06	38	0.368
566	A	20	19	0.99	38	0.500

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
567	A	14	13	1.12	34	0.382
568	A	22	21	1.22	36	0.583
569	A	19	18	1.24	36	0.500
570	A	16	15	1.25	36	0.417
571	A	19	18	1.25	36	0.500
572	A	22	21	1.21	36	0.583
573	A	25	24	1.15	36	0.667
574	A	17	16	0.90	31	0.516
575	A	15	14	0.89	31	0.452
576	A	16	15	0.89	31	0.484
577	A	18	17	0.90	31	0.548
578	A	21	20	0.87	33	0.606
579	A	19	18	0.85	33	0.545
580	A	18	17	0.83	33	0.515
581	A	19	18	0.83	33	0.545
582	A	21	20	0.85	33	0.606
583	A	24	23	0.84	33	0.697
584	A	22	21	0.82	33	0.636
585	A	21	20	0.81	33	0.606
586	A	22	21	0.81	33	0.636
587	A	23	22	0.82	33	0.667
588	A	25	24	0.85	33	0.727
589	A	26	25	0.88	33	0.758
590	A	23	22	0.84	33	0.667
591	A	20	19	0.81	33	0.576
592	A	19	18	0.81	33	0.545
593	A	23	22	0.84	33	0.667
594	A	26	25	0.88	33	0.758
595	A	27	26	0.86	33	0.788
596	A	24	23	0.84	33	0.697
597	A	24	23	0.85	33	0.697
598	A	24	23	0.84	33	0.697
599	A	27	26	0.86	33	0.788
600	A	30	29	0.87	33	0.879

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
601	A	27	26	0.87	33	0.788
602	A	27	26	0.86	33	0.788
603	A	27	26	0.87	33	0.788
604	A	27	26	0.87	33	0.788
605	A	30	29	0.87	33	0.879
606	A	15	14	1.01	36	0.389
607	A	15	14	1.01	36	0.389
608	A	13	12	1.00	36	0.333
609	A	13	12	1.00	36	0.333
610	A	15	14	1.01	36	0.389
611	A	15	14	1.01	36	0.389
612	A	22	21	1.06	35	0.600
613	A	19	18	1.07	35	0.514
614	A	16	15	1.02	35	0.429
615	A	13	12	0.98	35	0.343
616	A	15	14	0.87	35	0.400
617	A	11	10	0.84	35	0.286
618	A	14	13	0.88	35	0.371
619	A	25	24	1.07	35	0.686
620	A	22	21	1.06	35	0.600
621	A	19	18	1.04	35	0.514
622	A	16	15	1.04	35	0.429
623	A	11	10	0.82	35	0.286
624	A	11	10	0.84	35	0.286
625	A	14	13	0.87	35	0.371
626	A	17	16	0.91	35	0.457
627	A	28	27	1.06	35	0.771
628	A	25	24	1.07	35	0.686
629	A	22	21	1.06	35	0.600
630	A	19	18	1.06	35	0.514
631	A	14	13	0.87	35	0.371
632	A	14	13	0.86	35	0.371
633	A	14	13	0.87	35	0.371
634	A	17	16	0.92	35	0.457

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
635	A	20	19	0.92	35	0.543
636	A	19	18	1.02	35	0.514
637	A	16	15	0.99	35	0.429
638	A	13	12	0.94	35	0.343
639	A	10	9	0.88	35	0.257
640	A	15	14	0.83	35	0.400
641	A	11	10	0.86	35	0.286
642	A	19	18	1.07	35	0.514
643	A	16	15	1.09	35	0.429
644	A	13	12	1.04	35	0.343
645	A	13	12	1.06	35	0.343
646	A	11	10	0.98	35	0.286
647	A	22	21	1.12	35	0.600
648	A	19	18	1.12	35	0.514
649	A	16	15	1.11	35	0.429
650	A	16	15	1.12	35	0.429
651	A	16	15	1.11	35	0.429
652	A	14	13	1.07	35	0.371
653	A	8	7	0.85	38	0.184
654	A	10	9	0.85	38	0.237
655	A	8	7	0.78	38	0.184
656	A	9	8	1.11	31	0.258
657	A	10	9	1.08	33	0.273
658	A	10	9	1.13	33	0.273
659	A	10	9	1.12	33	0.273
660	A	10	9	1.12	33	0.273
661	A	8	7	1.00	33	0.212
662	A	8	7	1.00	33	0.212
663	A	8	7	1.00	33	0.212
664	A	8	7	0.95	33	0.212
665	A	5	4	1.02	39	0.103
666	A	6	5	0.90	39	0.128
667	A	6	5	0.90	39	0.128
668	A	6	5	0.80	39	0.128

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
669	A	4	3	0.75	37	0.081
670	A	4	4	1.00	24	0.167
671	A	6	5	0.80	39	0.128
672	A	5	4	1.00	39	0.103
673	A	6	5	0.89	39	0.128
674	A	6	5	0.89	39	0.128
675	A	6	5	0.89	39	0.128
676	A	6	5	0.94	41	0.122
677	A	6	5	0.83	41	0.122
678	A	6	5	0.83	41	0.122
679	A	6	5	0.83	41	0.122
680	A	7	6	0.79	41	0.146
681	A	6	5	0.80	39	0.128
682	A	6	6	0.88	26	0.231
683	A	6	5	0.70	41	0.122
684	A	6	5	0.71	41	0.122
685	A	6	5	0.82	41	0.122
686	A	6	5	0.81	41	0.122
687	A	6	5	0.82	41	0.122
688	A	6	5	0.81	41	0.122
689	A	6	5	0.91	41	0.122
690	A	6	5	0.81	41	0.122
691	A	6	5	0.81	41	0.122
692	A	6	5	0.80	41	0.122
693	A	8	7	0.73	41	0.171
694	A	6	5	0.83	41	0.122
695	A	6	5	0.90	39	0.128
696	A	8	8	0.90	26	0.308
697	A	6	5	0.68	41	0.122
698	A	6	5	0.69	41	0.122
699	A	7	6	0.78	41	0.146
700	A	6	5	0.92	41	0.122
701	A	6	5	0.79	41	0.122
702	A	6	5	0.80	41	0.122

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
703	A	6	5	0.79	41	0.122
704	A	6	5	0.80	41	0.122
705	A	6	5	1.03	41	0.122
706	A	6	5	0.70	41	0.122
707	A	6	5	0.70	41	0.122
708	A	6	5	0.71	41	0.122
709	A	6	5	0.84	39	0.128
710	A	3	3	1.00	26	0.115
711	A	7	6	1.11	41	0.146
712	A	6	5	0.85	41	0.122
713	A	6	5	0.83	41	0.122
714	A	6	5	0.81	41	0.122
715	A	6	5	1.03	41	0.122
716	A	6	5	0.71	41	0.122
717	A	6	5	0.72	41	0.122
718	A	6	5	0.73	41	0.122
719	A	6	5	0.75	41	0.122
720	A	5	4	1.00	39	0.103
721	A	5	5	0.99	26	0.192
722	A	6	5	0.85	41	0.122
723	A	8	7	1.08	41	0.171
724	A	6	5	0.81	41	0.122
725	A	6	5	0.80	41	0.122
726	A	6	5	0.79	41	0.122
727	A	6	5	1.03	41	0.122
728	A	6	5	0.72	41	0.122
729	A	6	5	0.72	41	0.122
730	A	7	6	0.81	41	0.146
731	A	6	5	0.83	41	0.122
732	A	6	5	0.90	39	0.128
733	A	7	7	0.99	26	0.269
734	A	6	5	0.83	41	0.122
735	A	6	5	0.81	41	0.122
736	A	9	8	1.05	41	0.195

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
737	A	6	5	0.79	41	0.122
738	A	6	5	0.78	41	0.122
739	A	6	5	0.78	41	0.122
740	A	5	4	1.02	41	0.098
741	A	5	4	1.02	41	0.098
742	A	5	4	1.02	41	0.098
743	A	5	4	1.02	41	0.098
744	A	5	4	1.02	41	0.098
745	A	5	4	1.02	41	0.098
746	A	5	4	1.02	41	0.098
747	A	5	4	1.02	41	0.098
748	A	6	5	0.93	43	0.116
749	A	6	5	0.93	43	0.116
750	A	6	5	0.93	43	0.116
751	A	6	5	0.93	43	0.116
752	A	6	5	0.93	43	0.116
753	A	6	5	0.93	43	0.116
754	A	6	5	0.93	43	0.116
755	A	6	5	0.93	43	0.116
756	A	6	5	0.91	43	0.116
757	A	6	5	0.91	43	0.116
758	A	6	5	0.91	43	0.116
759	A	6	5	0.91	43	0.116
760	A	6	5	0.91	43	0.116
761	A	6	5	0.91	43	0.116
762	A	6	5	0.91	43	0.116
763	A	6	5	0.91	43	0.116
764	A	10	9	0.85	43	0.209
765	A	9	8	0.89	43	0.186
766	A	8	7	0.92	43	0.163
767	A	7	6	0.99	43	0.140
768	A	8	7	0.97	43	0.163
769	A	9	8	0.93	43	0.186
770	A	10	9	0.92	43	0.209

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
771	A	11	10	0.83	43	0.233
772	A	10	9	0.85	43	0.209
773	A	9	8	0.87	43	0.186
774	A	8	7	0.91	43	0.163
775	A	8	7	0.97	43	0.163
776	A	9	8	0.93	43	0.186
777	A	10	9	0.96	43	0.209
778	A	11	10	0.92	43	0.233
779	A	11	10	0.84	43	0.233
780	A	10	9	0.86	43	0.209
781	A	9	8	0.89	43	0.186
782	A	9	8	0.93	43	0.186
783	A	9	8	0.96	43	0.186
784	A	10	9	0.93	43	0.209
785	A	11	10	0.95	43	0.233
786	A	12	11	0.95	43	0.256
787	A	9	8	0.97	45	0.178
788	A	8	7	1.00	45	0.156
789	A	7	6	1.02	45	0.133
790	A	6	5	1.06	45	0.111
791	A	6	5	1.06	45	0.111
792	A	5	4	1.09	45	0.089
793	A	6	5	1.08	45	0.111
794	A	7	6	1.06	45	0.133
795	A	9	8	0.95	45	0.178
796	A	8	7	0.96	45	0.156
797	A	7	6	1.01	45	0.133
798	A	7	6	1.04	45	0.133
799	A	7	6	1.01	45	0.133
800	A	7	6	1.08	45	0.133
801	A	5	4	1.09	45	0.089
802	A	6	5	1.08	45	0.111
803	A	7	6	1.05	45	0.133
804	A	8	7	1.05	45	0.156

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
805	A	9	8	0.93	45	0.178
806	A	8	7	0.99	45	0.156
807	A	8	7	0.97	45	0.156
808	A	8	7	1.00	45	0.156
809	A	8	7	0.97	45	0.156
810	A	8	7	0.98	45	0.156
811	A	8	7	1.07	45	0.156
812	A	5	4	1.09	45	0.089
813	A	6	5	1.08	45	0.111
814	A	7	6	1.06	45	0.133
815	A	8	7	1.05	45	0.156
816	A	10	9	0.91	45	0.200
817	A	9	8	0.98	45	0.178
818	A	9	8	0.94	45	0.178
819	A	9	8	0.95	45	0.178
820	A	9	8	0.97	45	0.178
821	A	9	8	0.95	45	0.178
822	A	9	8	0.96	45	0.178
823	A	9	8	0.97	45	0.178
824	A	9	8	1.07	45	0.178
825	A	5	4	1.09	45	0.089
826	A	6	5	1.08	45	0.111
827	A	7	6	1.06	45	0.133
828	A	8	7	1.05	45	0.156
829	A	9	8	1.05	45	0.178
830	A	8	7	0.97	45	0.156
831	A	7	6	1.01	45	0.133
832	A	6	5	1.05	45	0.111
833	A	5	4	1.10	45	0.089
834	A	6	5	1.04	45	0.111
835	A	7	6	1.02	45	0.133
836	A	9	8	0.96	45	0.178
837	A	8	7	0.98	45	0.156
838	A	7	6	1.08	45	0.133

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
839	A	5	4	1.09	45	0.089
840	A	6	5	1.07	45	0.111
841	A	6	5	1.05	45	0.111
842	A	8	7	1.01	45	0.156
843	A	10	9	0.95	45	0.200
844	A	9	8	0.97	45	0.178
845	A	8	7	1.07	45	0.156
846	A	5	4	1.09	45	0.089
847	A	6	5	1.08	45	0.111
848	A	7	6	1.02	45	0.133
849	A	7	6	1.03	45	0.133
850	A	7	6	1.06	45	0.133
851	A	6	5	1.10	41	0.122
852	A	6	5	1.03	47	0.106
853	A	4	3	1.00	46	0.065
854	A	5	5	1.05	36	0.139
855	A	8	7	1.05	38	0.184

LISTING OF INTEGRALS

3.1	$\int \tan^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$	298
3.2	$\int \tan(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$	305
3.3	$\int (a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$	311
3.4	$\int \cot(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$	316
3.5	$\int \cot^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$	322
3.6	$\int \cot^3(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$	328
3.7	$\int \cot^4(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$	335
3.8	$\int \cot^5(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$	343
3.9	$\int \tan^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$	351
3.10	$\int \tan(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$	359
3.11	$\int (a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$	366
3.12	$\int \cot(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$	372
3.13	$\int \cot^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$	379
3.14	$\int \cot^3(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$	386
3.15	$\int \cot^4(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$	393
3.16	$\int \cot^5(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$	401
3.17	$\int \tan^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$	410
3.18	$\int \tan(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$	419
3.19	$\int (a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$	427
3.20	$\int \cot(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$	433
3.21	$\int \cot^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$	441
3.22	$\int \cot^3(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$	449
3.23	$\int \cot^4(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$	457
3.24	$\int \cot^5(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$	466
3.25	$\int \cot^6(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$	476
3.26	$\int \tan^2(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$	486
3.27	$\int \tan(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$	496
3.28	$\int (a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$	505

3.29	$\int \cot(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$	513
3.30	$\int \cot^2(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$	522
3.31	$\int \cot^3(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$	532
3.32	$\int \cot^4(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$	542
3.33	$\int \cot^5(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$	551
3.34	$\int \cot^6(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$	560
3.35	$\int \cot^7(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$	570
3.36	$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$	580
3.37	$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$	587
3.38	$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$	593
3.39	$\int \frac{A+B \tan(c+dx)}{a+ia \tan(c+dx)} dx$	599
3.40	$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$	604
3.41	$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$	610
3.42	$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$	617
3.43	$\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$	625
3.44	$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$	634
3.45	$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$	641
3.46	$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$	648
3.47	$\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^2} dx$	654
3.48	$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$	659
3.49	$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$	666
3.50	$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$	674
3.51	$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$	683
3.52	$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$	691
3.53	$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$	698
3.54	$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$	704
3.55	$\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^3} dx$	710
3.56	$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$	716
3.57	$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$	723
3.58	$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$	732
3.59	$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$	742
3.60	$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$	751
3.61	$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$	759
3.62	$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$	766

3.63	$\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^4} dx$	773
3.64	$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$	779
3.65	$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$	787
3.66	$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$	798
3.67	$\int \tan^3(c+dx) \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	809
3.68	$\int \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	818
3.69	$\int \tan(c+dx) \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	825
3.70	$\int \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	831
3.71	$\int \cot(c+dx) \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	837
3.72	$\int \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	844
3.73	$\int \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	852
3.74	$\int \cot^4(c+dx) \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	862
3.75	$\int \tan^2(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	873
3.76	$\int \tan(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	882
3.77	$\int (a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	889
3.78	$\int \cot(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	895
3.79	$\int \cot^2(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	903
3.80	$\int \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	912
3.81	$\int \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	922
3.82	$\int \tan^2(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	933
3.83	$\int \tan(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	943
3.84	$\int (a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	951
3.85	$\int \cot(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	958
3.86	$\int \cot^2(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	967
3.87	$\int \cot^3(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	977
3.88	$\int \cot^4(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	987
3.89	$\int \cot^5(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	998
3.90	$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$	1010
3.91	$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$	1019
3.92	$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$	1027
3.93	$\int \frac{A+B \tan(c+dx)}{\sqrt{a+ia \tan(c+dx)}} dx$	1033
3.94	$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$	1039
3.95	$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$	1047
3.96	$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$	1057
3.97	$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$	1068
3.98	$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$	1076
3.99	$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$	1083

3.100	$\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^{3/2}} dx$	1089
3.101	$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$	1095
3.102	$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$	1104
3.103	$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$	1115
3.104	$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$	1127
3.105	$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$	1136
3.106	$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$	1144
3.107	$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$	1151
3.108	$\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^{5/2}} dx$	1158
3.109	$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$	1165
3.110	$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$	1175
3.111	$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$	1187
3.112	$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$	1200
3.113	$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$	1208
3.114	$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$	1216
3.115	$\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$	1223
3.116	$\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$	1229
3.117	$\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$	1235
3.118	$\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$	1242
3.119	$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$	1250
3.120	$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$	1260
3.121	$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$	1270
3.122	$\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$	1278
3.123	$\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$	1286
3.124	$\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$	1294
3.125	$\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$	1302
3.126	$\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$	1311
3.127	$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$	1321
3.128	$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$	1332
3.129	$\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$	1342
3.130	$\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$	1351
3.131	$\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$	1359

3.132	$\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$	1368
3.133	$\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$	1377
3.134	$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$	1387
3.135	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$	1398
3.136	$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$	1408
3.137	$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))} dx$	1417
3.138	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} dx$	1426
3.139	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$	1437
3.140	$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$	1448
3.141	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$	1459
3.142	$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$	1470
3.143	$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^2} dx$	1481
3.144	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$	1491
3.145	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$	1503
3.146	$\int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$	1515
3.147	$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$	1528
3.148	$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$	1541
3.149	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$	1552
3.150	$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$	1563
3.151	$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3} dx$	1574
3.152	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$	1585
3.153	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$	1598
3.154	$\int \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	1612
3.155	$\int \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	1622
3.156	$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$	1631
3.157	$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$	1639
3.158	$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$	1645
3.159	$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$	1653
3.160	$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$	1662

3.161	$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	1672
3.162	$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	1684
3.163	$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$	1694
3.164	$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$	1703
3.165	$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$	1712
3.166	$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$	1720
3.167	$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$	1729
3.168	$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$	1739
3.169	$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	1751
3.170	$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	1764
3.171	$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$	1775
3.172	$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$	1785
3.173	$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$	1795
3.174	$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$	1805
3.175	$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$	1814
3.176	$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$	1824
3.177	$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$	1835
3.178	$\int \frac{(a+ia \tan(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c+dx) \right)}{\tan^{\frac{5}{2}}(c+dx)} dx$	1849
3.179	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$	1859
3.180	$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$	1869
3.181	$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)}} dx$	1878
3.182	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} dx$	1885
3.183	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} dx$	1893
3.184	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{7}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} dx$	1902
3.185	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$	1912
3.186	$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$	1922
3.187	$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}} dx$	1930
3.188	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$	1937
3.189	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$	1945

3.190	$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx$	1954
3.191	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx$	1965
3.192	$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx$	1973
3.193	$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{\frac{5}{2}}} dx$	1981
3.194	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{5}{2}}} dx$	1988
3.195	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{5}{2}}} dx$	1997
3.196	$\int \sqrt[3]{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	2007
3.197	$\int \tan^2(c+dx)(a+ia \tan(c+dx))^{\frac{2}{3}}(A+B \tan(c+dx)) dx$	2016
3.198	$\int \tan(c+dx)(a+ia \tan(c+dx))^{\frac{2}{3}}(A+B \tan(c+dx)) dx$	2027
3.199	$\int (a+ia \tan(c+dx))^{\frac{2}{3}}(A+B \tan(c+dx)) dx$	2037
3.200	$\int \cot(c+dx)(a+ia \tan(c+dx))^{\frac{2}{3}}(A+B \tan(c+dx)) dx$	2045
3.201	$\int \cot^2(c+dx)(a+ia \tan(c+dx))^{\frac{2}{3}}(A+B \tan(c+dx)) dx$	2055
3.202	$\int \frac{A+B \tan(c+dx)}{\sqrt[3]{a+ia \tan(c+dx)}} dx$	2067
3.203	$\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^{\frac{2}{3}}} dx$	2075
3.204	$\int \tan^m(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$	2083
3.205	$\int \tan^m(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$	2091
3.206	$\int \tan^m(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$	2098
3.207	$\int \tan^m(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$	2105
3.208	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$	2110
3.209	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$	2116
3.210	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$	2123
3.211	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$	2131
3.212	$\int \tan^m(c+dx)(a+ia \tan(c+dx))^{\frac{5}{2}}(A+B \tan(c+dx)) dx$	2140
3.213	$\int \tan^m(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}(A+B \tan(c+dx)) dx$	2149
3.214	$\int \tan^m(c+dx)\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	2157
3.215	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$	2164
3.216	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{3}{2}}} dx$	2172
3.217	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx$	2181
3.218	$\int \tan^m(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	2191
3.219	$\int \tan^3(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	2198
3.220	$\int \tan^2(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	2205
3.221	$\int \tan(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	2212
3.222	$\int (a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	2218
3.223	$\int \cot(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	2223
3.224	$\int \cot^2(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	2229
3.225	$\int \cot^3(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	2236

3.226	$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	2243
3.227	$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	2253
3.228	$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	2262
3.229	$\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$	2271
3.230	$\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$	2278
3.231	$\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$	2286
3.232	$\int \tan^2(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	2296
3.233	$\int \tan(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	2303
3.234	$\int (a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	2309
3.235	$\int \cot(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	2314
3.236	$\int \cot^2(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	2320
3.237	$\int \cot^3(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	2326
3.238	$\int \cot^4(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	2333
3.239	$\int \cot^5(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	2340
3.240	$\int \tan^2(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	2348
3.241	$\int \tan(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	2357
3.242	$\int (a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	2364
3.243	$\int \cot(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	2371
3.244	$\int \cot^2(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	2377
3.245	$\int \cot^3(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	2383
3.246	$\int \cot^4(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	2391
3.247	$\int \cot^5(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	2399
3.248	$\int \tan^2(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	2409
3.249	$\int \tan(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	2419
3.250	$\int (a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	2428
3.251	$\int \cot(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	2435
3.252	$\int \cot^2(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	2443
3.253	$\int \cot^3(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	2450
3.254	$\int \cot^4(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	2458
3.255	$\int \cot^5(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	2468
3.256	$\int \cot^6(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	2479
3.257	$\int \tan^2(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$	2490
3.258	$\int \tan(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$	2500
3.259	$\int (a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$	2510
3.260	$\int \cot(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$	2518
3.261	$\int \cot^2(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$	2527
3.262	$\int \cot^3(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$	2536
3.263	$\int \cot^4(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$	2545
3.264	$\int \cot^5(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$	2554

3.265	$\int \cot^6(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$	2565
3.266	$\int \cot^7(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$	2577
3.267	$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$	2590
3.268	$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$	2598
3.269	$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$	2605
3.270	$\int \frac{A+B \tan(c+dx)}{a+b \tan(c+dx)} dx$	2612
3.271	$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$	2618
3.272	$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$	2625
3.273	$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$	2632
3.274	$\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$	2640
3.275	$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	2650
3.276	$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	2659
3.277	$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	2667
3.278	$\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^2} dx$	2674
3.279	$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	2681
3.280	$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	2689
3.281	$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	2699
3.282	$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$	2710
3.283	$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$	2721
3.284	$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$	2730
3.285	$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$	2739
3.286	$\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^3} dx$	2747
3.287	$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$	2755
3.288	$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$	2764
3.289	$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$	2775
3.290	$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$	2787
3.291	$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$	2799
3.292	$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$	2810
3.293	$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$	2820
3.294	$\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^4} dx$	2829
3.295	$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$	2839
3.296	$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$	2850
3.297	$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$	2862
3.298	$\int \frac{\tan^3(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$	2876

3.299	$\int \frac{\tan^2(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$	2881
3.300	$\int \frac{\tan(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$	2886
3.301	$\int \frac{aB+bB \tan(c+dx)}{a+b \tan(c+dx)} dx$	2891
3.302	$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$	2895
3.303	$\int \frac{\cot^2(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$	2900
3.304	$\int \frac{\cot^3(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$	2905
3.305	$\int \frac{\cot^4(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$	2911
3.306	$\int \frac{\tan^4(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	2916
3.307	$\int \frac{\tan^3(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	2925
3.308	$\int \frac{\tan^2(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	2933
3.309	$\int \frac{\tan(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	2940
3.310	$\int \frac{aB+bB \tan(c+dx)}{(a+b \tan(c+dx))^2} dx$	2946
3.311	$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	2952
3.312	$\int \frac{\cot^2(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	2959
3.313	$\int \frac{\cot^3(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	2967
3.314	$\int \frac{3+\tan(c+dx)}{2-\tan(c+dx)} dx$	2976
3.315	$\int \frac{\frac{bE}{a}+B \tan(c+dx)}{a+b \tan(c+dx)} dx$	2981
3.316	$\int \frac{a+b \tan(c+dx)}{(b+a \tan(c+dx))^2} dx$	2987
3.317	$\int \tan^3(c+dx) \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	2994
3.318	$\int \tan^2(c+dx) \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	3005
3.319	$\int \tan(c+dx) \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	3016
3.320	$\int \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	3026
3.321	$\int \cot(c+dx) \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	3035
3.322	$\int \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	3044
3.323	$\int \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	3055
3.324	$\int \cot^4(c+dx) \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	3067
3.325	$\int \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	3081
3.326	$\int \tan(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	3091
3.327	$\int (a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	3100
3.328	$\int \cot(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	3108
3.329	$\int \cot^2(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	3117
3.330	$\int \cot^3(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	3127
3.331	$\int \cot^4(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	3138
3.332	$\int \tan^2(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	3151
3.333	$\int \tan(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	3161
3.334	$\int (a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	3171

3.335	$\int \cot(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	3180
3.336	$\int \cot^2(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	3191
3.337	$\int \cot^3(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	3202
3.338	$\int \cot^4(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	3212
3.339	$\int \cot^5(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	3224
3.340	$\int (-a+b \tan(c+dx))(a+b \tan(c+dx))^{5/2} dx$	3239
3.341	$\int (-a+b \tan(c+dx))(a+b \tan(c+dx))^{3/2} dx$	3248
3.342	$\int (-a+b \tan(c+dx))\sqrt{a+b \tan(c+dx)} dx$	3258
3.343	$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	3269
3.344	$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	3279
3.345	$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	3288
3.346	$\int \frac{A+B \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$	3296
3.347	$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	3303
3.348	$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	3312
3.349	$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	3322
3.350	$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	3333
3.351	$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	3344
3.352	$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	3353
3.353	$\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx$	3361
3.354	$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	3369
3.355	$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	3377
3.356	$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	3387
3.357	$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	3398
3.358	$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	3409
3.359	$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	3418
3.360	$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	3426
3.361	$\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^{5/2}} dx$	3434
3.362	$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	3442
3.363	$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	3452
3.364	$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	3463
3.365	$\int \frac{aB+bB \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$	3475
3.366	$\int \frac{aB+bB \tan(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx$	3484
3.367	$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	3494
3.368	$\int \frac{aB+bB \tan(c+dx)}{(a+b \tan(c+dx))^{5/2}} dx$	3503

3.369	$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	3511
3.370	$\int \frac{-a+b \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$	3521
3.371	$\int \frac{-a+b \tan(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx$	3528
3.372	$\int \frac{-a+b \tan(c+dx)}{(a+b \tan(c+dx))^{5/2}} dx$	3536
3.373	$\int \frac{1+i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$	3545
3.374	$\int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$	3553
3.375	$\int \frac{3+\tan(x)}{\sqrt{4+3 \tan(x)}} dx$	3560
3.376	$\int \frac{1-3 \tan(x)}{\sqrt{4+3 \tan(x)}} dx$	3565
3.377	$\int \frac{4-3 \tan(a+bx)}{\sqrt{4+3 \tan(a+bx)}} dx$	3570
3.378	$\int \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	3577
3.379	$\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	3588
3.380	$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	3598
3.381	$\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$	3608
3.382	$\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$	3618
3.383	$\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$	3629
3.384	$\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$	3640
3.385	$\int \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	3650
3.386	$\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	3663
3.387	$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	3675
3.388	$\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$	3686
3.389	$\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$	3697
3.390	$\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$	3707
3.391	$\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$	3717
3.392	$\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	3728
3.393	$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	3740
3.394	$\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$	3752
3.395	$\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$	3763
3.396	$\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$	3774
3.397	$\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$	3785
3.398	$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$	3796
3.399	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$	3808
3.400	$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$	3819

3.401	$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx$	3829
3.402	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$	3839
3.403	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$	3850
3.404	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	3862
3.405	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	3875
3.406	$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	3886
3.407	$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))^2}} dx$	3898
3.408	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$	3910
3.409	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$	3923
3.410	$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$	3937
3.411	$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$	3952
3.412	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$	3966
3.413	$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$	3979
3.414	$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))^3}} dx$	3993
3.415	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$	4007
3.416	$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$	4022
3.417	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$	4031
3.418	$\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$	4040
3.419	$\int \frac{aB+bB \tan(c+dx)}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx$	4048
3.420	$\int \frac{aB+bB \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$	4056
3.421	$\int \frac{aB+bB \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$	4065
3.422	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	4074
3.423	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	4086
3.424	$\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	4097
3.425	$\int \frac{aB+bB \tan(c+dx)}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))^2}} dx$	4108
3.426	$\int \frac{aB+bB \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$	4119
3.427	$\int \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	4131
3.428	$\int \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	4139
3.429	$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$	4146
3.430	$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$	4154

3.431	$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$	4161
3.432	$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$	4169
3.433	$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$	4178
3.434	$\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	4189
3.435	$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	4198
3.436	$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$	4206
3.437	$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$	4213
3.438	$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$	4220
3.439	$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$	4229
3.440	$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$	4239
3.441	$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$	4250
3.442	$\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	4264
3.443	$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	4273
3.444	$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$	4282
3.445	$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$	4290
3.446	$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$	4298
3.447	$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$	4306
3.448	$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$	4316
3.449	$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$	4327
3.450	$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$	4340
3.451	$\int \frac{(a+b \tan(c+dx))^{5/2}\left(\frac{3bB}{2a}+B \tan(c+dx)\right)}{\tan^{\frac{5}{2}}(c+dx)} dx$	4356
3.452	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	4364
3.453	$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	4370
3.454	$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx$	4378
3.455	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx$	4385
3.456	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx$	4392
3.457	$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{7}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx$	4400
3.458	$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	4409
3.459	$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	4416

3.460	$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx$	4423
3.461	$\int \frac{A+B \tan(c+dx)}{\tan^{3/2}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$	4431
3.462	$\int \frac{A+B \tan(c+dx)}{\tan^{5/2}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$	4439
3.463	$\int \frac{\tan^{5/2}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	4448
3.464	$\int \frac{\tan^{3/2}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	4456
3.465	$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	4465
3.466	$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx$	4473
3.467	$\int \frac{A+B \tan(c+dx)}{\tan^{3/2}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$	4481
3.468	$\int \frac{A+B \tan(c+dx)}{\tan^{5/2}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$	4490
3.469	$\int \frac{\tan^{3/2}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	4500
3.470	$\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	4505
3.471	$\int \frac{aB+bB \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx$	4511
3.472	$\int \frac{aB+bB \tan(c+dx)}{\tan^{3/2}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$	4516
3.473	$\int (a+b \tan(c+dx))^{2/3}(A+B \tan(c+dx)) dx$	4523
3.474	$\int \sqrt[3]{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	4532
3.475	$\int \frac{A+B \tan(c+dx)}{\sqrt[3]{a+b \tan(c+dx)}} dx$	4542
3.476	$\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^{2/3}} dx$	4551
3.477	$\int \frac{i-\tan(e+fx)}{\sqrt[3]{c+d \tan(e+fx)}} dx$	4559
3.478	$\int \frac{d-c \tan(e+fx)}{(c+d \tan(e+fx))^{2/3}} dx$	4567
3.479	$\int \tan^m(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$	4576
3.480	$\int \tan^m(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	4585
3.481	$\int \tan^m(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	4593
3.482	$\int \tan^m(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	4600
3.483	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$	4606
3.484	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	4613
3.485	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$	4621
3.486	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$	4631
3.487	$\int \tan^m(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	4642
3.488	$\int \tan^m(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	4648
3.489	$\int \tan^m(c+dx)\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	4654
3.490	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	4660
3.491	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	4666
3.492	$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	4672

3.493	$\int \tan^m(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$	4678
3.494	$\int \tan^4(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$	4684
3.495	$\int \tan^3(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$	4693
3.496	$\int \tan^2(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$	4701
3.497	$\int \tan(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$	4708
3.498	$\int (a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$	4714
3.499	$\int \cot(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$	4719
3.500	$\int \cot^2(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$	4726
3.501	$\int \cot^3(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$	4734
3.502	$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$	4743
3.503	$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$	4751
3.504	$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$	4758
3.505	$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$	4765
3.506	$\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$	4772
3.507	$\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$	4779
3.508	$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$	4787
3.509	$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$	4795
3.510	$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$	4803
3.511	$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$	4811
3.512	$\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$	4819
3.513	$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$	4828
3.514	$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$	4837
3.515	$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$	4845
3.516	$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$	4853
3.517	$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$	4861
3.518	$\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$	4870
3.519	$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$	4880
3.520	$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$	4890
3.521	$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$	4900
3.522	$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))} dx$	4910
3.523	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} dx$	4919
3.524	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$	4929
3.525	$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$	4940
3.526	$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$	4951
3.527	$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^2} dx$	4961

3.528	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$	4971
3.529	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$	4981
3.530	$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$	4992
3.531	$\int \frac{\sqrt{\cot(c+dx)(A+B \tan(c+dx))}}{(a+ia \tan(c+dx))^3} dx$	5004
3.532	$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+ia \tan(c+dx))^3}} dx$	5015
3.533	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$	5026
3.534	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$	5037
3.535	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$	5048
3.536	$\int \cot^{\frac{7}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	5060
3.537	$\int \cot^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	5069
3.538	$\int \cot^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	5077
3.539	$\int \sqrt{\cot(c+dx)} \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$	5085
3.540	$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$	5094
3.541	$\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	5104
3.542	$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	5115
3.543	$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	5125
3.544	$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	5134
3.545	$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	5143
3.546	$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$	5153
3.547	$\int \cot^{\frac{11}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	5163
3.548	$\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	5175
3.549	$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	5186
3.550	$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	5196
3.551	$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	5207
3.552	$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	5217
3.553	$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$	5227
3.554	$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$	5237
3.555	$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$	5246
3.556	$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$	5254
3.557	$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)} \sqrt{a+ia \tan(c+dx)}} dx$	5261
3.558	$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$	5270
3.559	$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$	5279
3.560	$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{3/2}} dx$	5286

3.561	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$	5294
3.562	$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$	5304
3.563	$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$	5313
3.564	$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{5/2}} dx$	5321
3.565	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$	5330
3.566	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$	5339
3.567	$\int \cot^m(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	5349
3.568	$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	5356
3.569	$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	5366
3.570	$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$	5375
3.571	$\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$	5383
3.572	$\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$	5392
3.573	$\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\cot^{\frac{5}{2}}(c+dx)} dx$	5402
3.574	$\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	5413
3.575	$\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	5423
3.576	$\int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$	5433
3.577	$\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$	5443
3.578	$\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	5453
3.579	$\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	5465
3.580	$\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	5476
3.581	$\int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$	5486
3.582	$\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$	5497
3.583	$\int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	5508
3.584	$\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	5521
3.585	$\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	5533
3.586	$\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	5544
3.587	$\int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$	5556
3.588	$\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$	5569
3.589	$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$	5582
3.590	$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$	5595
3.591	$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$	5607
3.592	$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))} dx$	5617
3.593	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$	5627
3.594	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$	5638

3.595	$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	5651
3.596	$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$	5665
3.597	$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^2} dx$	5677
3.598	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$	5689
3.599	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$	5701
3.600	$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$	5714
3.601	$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$	5730
3.602	$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^3} dx$	5744
3.603	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$	5758
3.604	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$	5772
3.605	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$	5786
3.606	$\int \frac{\cot^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$	5801
3.607	$\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$	5810
3.608	$\int \frac{\sqrt{\cot(c+dx)}(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$	5819
3.609	$\int \frac{aB+bB \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))} dx$	5827
3.610	$\int \frac{aB+bB \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$	5835
3.611	$\int \frac{aB+bB \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$	5844
3.612	$\int \cot^{\frac{9}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	5853
3.613	$\int \cot^{\frac{7}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	5864
3.614	$\int \cot^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	5874
3.615	$\int \cot^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	5883
3.616	$\int \sqrt{\cot(c+dx)} \sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$	5891
3.617	$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$	5899
3.618	$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$	5906
3.619	$\int \cot^{\frac{11}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	5914
3.620	$\int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	5929
3.621	$\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	5940
3.622	$\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	5949
3.623	$\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	5958
3.624	$\int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$	5965
3.625	$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$	5972
3.626	$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$	5980

3.627	$\int \cot^{\frac{13}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	5989
3.628	$\int \cot^{\frac{11}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	6005
3.629	$\int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	6018
3.630	$\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	6029
3.631	$\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	6039
3.632	$\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	6048
3.633	$\int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$	6057
3.634	$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$	6065
3.635	$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$	6073
3.636	$\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	6083
3.637	$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	6093
3.638	$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	6102
3.639	$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$	6109
3.640	$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}\sqrt{a+b \tan(c+dx)}} dx$	6115
3.641	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx$	6123
3.642	$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	6130
3.643	$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	6140
3.644	$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	6149
3.645	$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^{3/2}} dx$	6156
3.646	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$	6163
3.647	$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	6170
3.648	$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	6181
3.649	$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$	6191
3.650	$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^{5/2}} dx$	6200
3.651	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$	6209
3.652	$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$	6218
3.653	$\int \frac{\sqrt{\cot(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$	6227
3.654	$\int \frac{aB+bB \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^{3/2}} dx$	6234
3.655	$\int \frac{aB+bB \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$	6241
3.656	$\int \cot^m(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$	6248
3.657	$\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$	6254

3.658	$\int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$	6260
3.659	$\int \frac{(a+b \tan(c+dx))^n(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$	6267
3.660	$\int \frac{(a+b \tan(c+dx))^n(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$	6274
3.661	$\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$	6281
3.662	$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$	6287
3.663	$\int \frac{(a+b \tan(c+dx))^n(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$	6293
3.664	$\int \frac{(a+b \tan(c+dx))^n(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$	6299
3.665	$\int (a+ia \tan(e+fx))(A+B \tan(e+fx))(c-ictan(e+fx))^n dx$	6305
3.666	$\int (a+ia \tan(e+fx))(A+B \tan(e+fx))(c-ictan(e+fx))^4 dx$	6311
3.667	$\int (a+ia \tan(e+fx))(A+B \tan(e+fx))(c-ictan(e+fx))^3 dx$	6317
3.668	$\int (a+ia \tan(e+fx))(A+B \tan(e+fx))(c-ictan(e+fx))^2 dx$	6323
3.669	$\int (a+ia \tan(e+fx))(A+B \tan(e+fx))(c-ictan(e+fx)) dx$	6329
3.670	$\int (a+ia \tan(e+fx))(A+B \tan(e+fx)) dx$	6334
3.671	$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{c-ictan(e+fx)} dx$	6339
3.672	$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ictan(e+fx))^2} dx$	6345
3.673	$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ictan(e+fx))^3} dx$	6351
3.674	$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ictan(e+fx))^4} dx$	6357
3.675	$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ictan(e+fx))^5} dx$	6363
3.676	$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ictan(e+fx))^n dx$	6369
3.677	$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ictan(e+fx))^5 dx$	6376
3.678	$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ictan(e+fx))^4 dx$	6383
3.679	$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ictan(e+fx))^3 dx$	6389
3.680	$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ictan(e+fx))^2 dx$	6395
3.681	$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ictan(e+fx)) dx$	6401
3.682	$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx)) dx$	6407
3.683	$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{c-ictan(e+fx)} dx$	6413
3.684	$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ictan(e+fx))^2} dx$	6419
3.685	$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ictan(e+fx))^3} dx$	6425
3.686	$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ictan(e+fx))^4} dx$	6431
3.687	$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ictan(e+fx))^5} dx$	6437
3.688	$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ictan(e+fx))^6} dx$	6443
3.689	$\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx))^n dx$	6449
3.690	$\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx))^6 dx$	6456
3.691	$\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx))^5 dx$	6463
3.692	$\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx))^4 dx$	6470
3.693	$\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx))^3 dx$	6477
3.694	$\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx))^2 dx$	6484

3.695	$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ic \tan(e + fx)) dx$	6491
3.696	$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) dx$	6497
3.697	$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx$	6503
3.698	$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx$	6510
3.699	$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx$	6517
3.700	$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx$	6523
3.701	$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx$	6529
3.702	$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx$	6536
3.703	$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^7} dx$	6543
3.704	$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^8} dx$	6550
3.705	$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{a + ia \tan(e + fx)} dx$	6557
3.706	$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{a + ia \tan(e + fx)} dx$	6563
3.707	$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{a + ia \tan(e + fx)} dx$	6570
3.708	$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{a + ia \tan(e + fx)} dx$	6577
3.709	$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))}{a + ia \tan(e + fx)} dx$	6584
3.710	$\int \frac{A + B \tan(e + fx)}{a + ia \tan(e + fx)} dx$	6590
3.711	$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))} dx$	6595
3.712	$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^2} dx$	6601
3.713	$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^3} dx$	6607
3.714	$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^4} dx$	6613
3.715	$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^2} dx$	6619
3.716	$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^5}{(a + ia \tan(e + fx))^2} dx$	6625
3.717	$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^2} dx$	6633
3.718	$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{(a + ia \tan(e + fx))^2} dx$	6640
3.719	$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{(a + ia \tan(e + fx))^2} dx$	6647
3.720	$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx$	6653
3.721	$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2} dx$	6659
3.722	$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))} dx$	6664
3.723	$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^2} dx$	6670
3.724	$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^3} dx$	6676
3.725	$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^4} dx$	6682
3.726	$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^5} dx$	6689
3.727	$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^3} dx$	6697
3.728	$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^5}{(a + ia \tan(e + fx))^3} dx$	6703

3.729	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^4}{(a+ia \tan(e+fx))^3} dx$	6711
3.730	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^3}{(a+ia \tan(e+fx))^3} dx$	6718
3.731	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^2}{(a+ia \tan(e+fx))^3} dx$	6724
3.732	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))}{(a+ia \tan(e+fx))^3} dx$	6730
3.733	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3} dx$	6736
3.734	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))} dx$	6742
3.735	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^2} dx$	6748
3.736	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^3} dx$	6754
3.737	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^4} dx$	6761
3.738	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^5} dx$	6769
3.739	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^6} dx$	6777
3.740	$\int (a+ia \tan(e+fx))(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2} dx$	6785
3.741	$\int (a+ia \tan(e+fx))(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2} dx$	6791
3.742	$\int (a+ia \tan(e+fx))(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2} dx$	6797
3.743	$\int (a+ia \tan(e+fx))(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)} dx$	6803
3.744	$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$	6809
3.745	$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$	6815
3.746	$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$	6821
3.747	$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$	6827
3.748	$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2} dx$	6833
3.749	$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2} dx$	6840
3.750	$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2} dx$	6846
3.751	$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)} dx$	6853
3.752	$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$	6859
3.753	$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$	6865
3.754	$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$	6871
3.755	$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$	6878
3.756	$\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2} dx$	6885
3.757	$\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2} dx$	6892
3.758	$\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2} dx$	6899
3.759	$\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)} dx$	6906
3.760	$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$	6913
3.761	$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$	6920
3.762	$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$	6927
3.763	$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$	6934

3.764	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{a+ia \tan(e+fx)} dx$	6941
3.765	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{a+ia \tan(e+fx)} dx$	6949
3.766	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{a+ia \tan(e+fx)} dx$	6956
3.767	$\int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{a+ia \tan(e+fx)} dx$	6963
3.768	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))\sqrt{c-ic \tan(e+fx)}} dx$	6970
3.769	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} dx$	6977
3.770	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^{5/2}} dx$	6985
3.771	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^2} dx$	6993
3.772	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^2} dx$	7002
3.773	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^2} dx$	7010
3.774	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^2} dx$	7018
3.775	$\int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{(a+ia \tan(e+fx))^2} dx$	7025
3.776	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 \sqrt{c-ic \tan(e+fx)}} dx$	7032
3.777	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 (c-ic \tan(e+fx))^{3/2}} dx$	7040
3.778	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 (c-ic \tan(e+fx))^{5/2}} dx$	7048
3.779	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^3} dx$	7057
3.780	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^3} dx$	7065
3.781	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^3} dx$	7073
3.782	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^3} dx$	7081
3.783	$\int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{(a+ia \tan(e+fx))^3} dx$	7089
3.784	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3 \sqrt{c-ic \tan(e+fx)}} dx$	7097
3.785	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3 (c-ic \tan(e+fx))^{3/2}} dx$	7105
3.786	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3 (c-ic \tan(e+fx))^{5/2}} dx$	7114
3.787	$\int \sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2} dx$	7123
3.788	$\int \sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2} dx$	7132
3.789	$\int \sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2} dx$	7140
3.790	$\int \sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)} dx$	7148
3.791	$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$	7155
3.792	$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$	7162
3.793	$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$	7168
3.794	$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$	7174
3.795	$\int (a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2} dx$	7181
3.796	$\int (a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2} dx$	7190

3.797	$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ict \tan(e + fx))^{3/2} dx$	7199
3.798	$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) \sqrt{c - ict \tan(e + fx)} dx$	7207
3.799	$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{\sqrt{c - ict \tan(e + fx)}} dx$	7215
3.800	$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{3/2}} dx$	7223
3.801	$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{5/2}} dx$	7230
3.802	$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{7/2}} dx$	7236
3.803	$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{9/2}} dx$	7242
3.804	$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{11/2}} dx$	7249
3.805	$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ict \tan(e + fx))^{7/2} dx$	7257
3.806	$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ict \tan(e + fx))^{5/2} dx$	7266
3.807	$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ict \tan(e + fx))^{3/2} dx$	7274
3.808	$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) \sqrt{c - ict \tan(e + fx)} dx$	7283
3.809	$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{\sqrt{c - ict \tan(e + fx)}} dx$	7291
3.810	$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{3/2}} dx$	7300
3.811	$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{5/2}} dx$	7309
3.812	$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{7/2}} dx$	7317
3.813	$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{9/2}} dx$	7323
3.814	$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{11/2}} dx$	7329
3.815	$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{13/2}} dx$	7336
3.816	$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ict \tan(e + fx))^{9/2} dx$	7344
3.817	$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ict \tan(e + fx))^{7/2} dx$	7353
3.818	$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ict \tan(e + fx))^{5/2} dx$	7361
3.819	$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ict \tan(e + fx))^{3/2} dx$	7370
3.820	$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) \sqrt{c - ict \tan(e + fx)} dx$	7379
3.821	$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{\sqrt{c - ict \tan(e + fx)}} dx$	7388
3.822	$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{3/2}} dx$	7397
3.823	$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{5/2}} dx$	7407
3.824	$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{7/2}} dx$	7418
3.825	$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{9/2}} dx$	7429
3.826	$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{11/2}} dx$	7435
3.827	$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{13/2}} dx$	7442
3.828	$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{15/2}} dx$	7449
3.829	$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{17/2}} dx$	7457

3.830	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{\sqrt{a+ia \tan(e+fx)}} dx$	7468
3.831	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{\sqrt{a+ia \tan(e+fx)}} dx$	7477
3.832	$\int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{\sqrt{a+ia \tan(e+fx)}} dx$	7485
3.833	$\int \frac{A+B \tan(e+fx)}{\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}} dx$	7492
3.834	$\int \frac{A+B \tan(e+fx)}{\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{3/2}} dx$	7498
3.835	$\int \frac{A+B \tan(e+fx)}{\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{5/2}} dx$	7504
3.836	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^{3/2}} dx$	7511
3.837	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^{3/2}} dx$	7520
3.838	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^{3/2}} dx$	7528
3.839	$\int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{(a+ia \tan(e+fx))^{3/2}} dx$	7535
3.840	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{3/2}\sqrt{c-ic \tan(e+fx)}} dx$	7541
3.841	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{3/2}(c-ic \tan(e+fx))^{3/2}} dx$	7547
3.842	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{3/2}(c-ic \tan(e+fx))^{5/2}} dx$	7553
3.843	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^{5/2}} dx$	7560
3.844	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^{5/2}} dx$	7570
3.845	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^{5/2}} dx$	7580
3.846	$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^{5/2}} dx$	7588
3.847	$\int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{(a+ia \tan(e+fx))^{5/2}} dx$	7594
3.848	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2}\sqrt{c-ic \tan(e+fx)}} dx$	7600
3.849	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2}(c-ic \tan(e+fx))^{3/2}} dx$	7607
3.850	$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2}(c-ic \tan(e+fx))^{5/2}} dx$	7614
3.851	$\int (a+ia \tan(e+fx))^m (A+B \tan(e+fx))(c-ic \tan(e+fx))^n dx$	7621
3.852	$\int (a+ia \tan(e+fx))^{1+m} (A+B \tan(e+fx))(c-ic \tan(e+fx))^{-1-m} dx$	7627
3.853	$\int \frac{(c-ic \tan(e+fx))^n (-i(2+n)+(-2+n) \tan(e+fx))}{(-i+\tan(e+fx))^2} dx$	7634
3.854	$\int \frac{(A+B \tan(e+fx))(c+d \tan(e+fx))}{(a+ia \tan(e+fx))^2} dx$	7639
3.855	$\int \frac{(A+B \tan(e+fx))(c+d \tan(e+fx))}{(a+ia \tan(e+fx))^{3/2}} dx$	7645

3.1 $\int \tan^2(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.1.1	Optimal result	298
3.1.2	Mathematica [A] (verified)	298
3.1.3	Rubi [A] (verified)	299
3.1.4	Maple [A] (verified)	301
3.1.5	Fricas [B] (verification not implemented)	302
3.1.6	Sympy [B] (verification not implemented)	302
3.1.7	Maxima [A] (verification not implemented)	303
3.1.8	Giac [B] (verification not implemented)	303
3.1.9	Mupad [B] (verification not implemented)	304

3.1.1 Optimal result

Integrand size = 32, antiderivative size = 91

$$\int \tan^2(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= -a(A-iB)x + \frac{a(iA+B) \log(\cos(c+dx))}{d} + \frac{a(A-iB) \tan(c+dx)}{d}$$

$$+ \frac{a(iA+B) \tan^2(c+dx)}{2d} + \frac{iaB \tan^3(c+dx)}{3d}$$

```
output -a*(A-I*B)*x+a*(I*A+B)*ln(cos(d*x+c))/d+a*(A-I*B)*tan(d*x+c)/d+1/2*a*(I*A+B)*tan(d*x+c)^2/d+1/3*I*a*B*tan(d*x+c)^3/d
```

3.1.2 Mathematica [A] (verified)

Time = 1.07 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.95

$$\int \tan^2(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{a(-6(A-iB) \arctan(\tan(c+dx)) + 6(iA+B) \log(\cos(c+dx)) + 6(A-iB) \tan(c+dx) + 3(iA+B) \tan^3(c+dx))}{6d}$$

```
input Integrate[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]
```

output $(a*(-6*(A - I*B)*ArcTan[Tan[c + d*x]] + 6*(I*A + B)*Log[Cos[c + d*x]] + 6*(A - I*B)*Tan[c + d*x] + 3*(I*A + B)*Tan[c + d*x]^2 + (2*I)*B*Tan[c + d*x]^3)/(6*d)$

3.1.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4075, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow 3042 \\
 & \int \tan(c + dx)^2(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow 4075 \\
 & \int \tan^2(c + dx)(a(A - iB) + a(iA + B) \tan(c + dx)) dx + \frac{iaB \tan^3(c + dx)}{3d} \\
 & \quad \downarrow 3042 \\
 & \int \tan(c + dx)^2(a(A - iB) + a(iA + B) \tan(c + dx)) dx + \frac{iaB \tan^3(c + dx)}{3d} \\
 & \quad \downarrow 4011 \\
 & \int \tan(c + dx)(a(A - iB) \tan(c + dx) - a(iA + B)) dx + \frac{a(B + iA) \tan^2(c + dx)}{2d} + \frac{iaB \tan^3(c + dx)}{3d} \\
 & \quad \downarrow 3042 \\
 & \int \tan(c + dx)(a(A - iB) \tan(c + dx) - a(iA + B)) dx + \frac{a(B + iA) \tan^2(c + dx)}{2d} + \frac{iaB \tan^3(c + dx)}{3d} \\
 & \quad \downarrow 4008 \\
 & -a(B + iA) \int \tan(c + dx) dx + \frac{a(B + iA) \tan^2(c + dx)}{2d} + \frac{a(A - iB) \tan(c + dx)}{d} - ax(A - iB) + \frac{iaB \tan^3(c + dx)}{3d} \\
 & \quad \downarrow 3042
 \end{aligned}$$

3.1. $\int \tan^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\begin{aligned}
 & -a(B + iA) \int \tan(c + dx) dx + \frac{a(B + iA) \tan^2(c + dx)}{2d} + \frac{a(A - iB) \tan(c + dx)}{d} - ax(A - iB) + \\
 & \quad \frac{iaB \tan^3(c + dx)}{3d} \\
 & \quad \downarrow \text{3956} \\
 & \frac{a(B + iA) \tan^2(c + dx)}{2d} + \frac{a(A - iB) \tan(c + dx)}{d} + \frac{a(B + iA) \log(\cos(c + dx))}{d} - ax(A - iB) + \\
 & \quad \frac{iaB \tan^3(c + dx)}{3d}
 \end{aligned}$$

input `Int[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `-(a*(A - I*B)*x) + (a*(I*A + B)*Log[Cos[c + d*x]])/d + (a*(A - I*B)*Tan[c + d*x])/d + (a*(I*A + B)*Tan[c + d*x]^2)/(2*d) + ((I/3)*a*B*Tan[c + d*x]^3)/d`

3.1.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

3.1.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.04

method	result
parts	$\frac{(iaA+Ba)\left(\frac{\tan^2(dx+c)}{2} - \frac{\ln(1+\tan^2(dx+c))}{2}\right)}{d} + \frac{aA(\tan(dx+c)-\arctan(\tan(dx+c)))}{d} + \frac{iaB\left(\frac{\tan^3(dx+c)}{3} - \tan(dx+c)\right)}{d}$
norman	$(iaB - aA)x + \frac{(-iaB+aA)\tan(dx+c)}{d} + \frac{(iaA+Ba)(\tan^2(dx+c))}{2d} + \frac{iaB(\tan^3(dx+c))}{3d} - \frac{(iaA+Ba)\ln(1+\tan^2(dx+c))}{2d}$
derivativedivides	$a\left(\frac{iB(\tan^3(dx+c))}{3} + \frac{iA(\tan^2(dx+c))}{2} - iB\tan(dx+c) + \frac{B(\tan^2(dx+c))}{2} + A\tan(dx+c) + \frac{(-iA-B)\ln(1+\tan^2(dx+c))}{2}\right) + (iB - aA)x$
default	$a\left(\frac{iB(\tan^3(dx+c))}{3} + \frac{iA(\tan^2(dx+c))}{2} - iB\tan(dx+c) + \frac{B(\tan^2(dx+c))}{2} + A\tan(dx+c) + \frac{(-iA-B)\ln(1+\tan^2(dx+c))}{2}\right) + (iB - aA)x$
parallelrisc	$-\frac{-2iaB(\tan^3(dx+c))-3iA(\tan^2(dx+c))a-6iBxad+3iA\ln(1+\tan^2(dx+c))a+6Axad+6iB\tan(dx+c)a-3B(\tan^2(dx+c))}{6d}$
risc	$-\frac{2iaBc}{d} + \frac{2aAc}{d} + \frac{2a(6iAe^{4i(dx+c)}+9Be^{4i(dx+c)}+9iAe^{2i(dx+c)}+9Be^{2i(dx+c)}+3iA+4B)}{3d(e^{2i(dx+c)}+1)^3} + \frac{a\ln(e^{2i(dx+c)}+1)}{d}$

```
input int(tan(d*x+c)^2*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output ((I*a*A+B*a)/d*(1/2*tan(d*x+c)^2-1/2*ln(1+tan(d*x+c)^2))+a*A/d*(tan(d*x+c)-arctan(tan(d*x+c)))+I*a*B/d*(1/3*tan(d*x+c)^3-tan(d*x+c)+arctan(tan(d*x+c))))
```

3.1. $\int \tan^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

3.1.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 170 vs. $2(77) = 154$.

Time = 0.25 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.87

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{6(-2iA - 3B)ae^{(4i dx + 4i c)} + 18(-iA - B)ae^{(2i dx + 2i c)} + 2(-3iA - 4B)a + 3((-iA - B)ae^{(6i dx + 6i c)} + 3de^{(6i dx + 6i c)} + 3de^{(4i dx + 4i c)})}{3(de^{(6i dx + 6i c)} + 3de^{(4i dx + 4i c)})}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-1/3*(6*(-2*I*A - 3*B)*a*e^(4*I*d*x + 4*I*c) + 18*(-I*A - B)*a*e^(2*I*d*x + 2*I*c) + 2*(-3*I*A - 4*B)*a + 3*((-I*A - B)*a*e^(6*I*d*x + 6*I*c) + 3*(-I*A - B)*a*e^(4*I*d*x + 4*I*c) + 3*(-I*A - B)*a*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a)*log(e^(2*I*d*x + 2*I*c) + 1)/(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)`

3.1.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 167 vs. $2(75) = 150$.

Time = 0.35 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.84

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{ia(A - iB) \log(e^{2idx} + e^{-2ic})}{d} + \frac{6iAa + 8Ba + (18iAae^{2ic} + 18Bae^{2ic})e^{2idx} + (12iAae^{4ic} + 18Bae^{4ic})e^{4idx}}{3de^{6ic}e^{6idx} + 9de^{4ic}e^{4idx} + 9de^{2ic}e^{2idx} + 3d}$$

input `integrate(tan(d*x+c)**2*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `I*a*(A - I*B)*log(exp(2*I*d*x) + exp(-2*I*c))/d + (6*I*A*a + 8*B*a + (18*I*A*a*exp(2*I*c) + 18*B*a*exp(2*I*c))*exp(2*I*d*x) + (12*I*A*a*exp(4*I*c) + 18*B*a*exp(4*I*c))*exp(4*I*d*x))/(3*d*exp(6*I*c)*exp(6*I*d*x) + 9*d*exp(4*I*c)*exp(4*I*d*x) + 9*d*exp(2*I*c)*exp(2*I*d*x) + 3*d)`

3.1. $\int \tan^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

3.1.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.90

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{-2i Ba \tan(dx + c)^3 + 3(-iA - B)a \tan(dx + c)^2 + 6(dx + c)(A - iB)a + 3(iA + B)a \log(\tan(dx + c))}{6d}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/6*(-2*I*B*a*tan(d*x + c)^3 + 3*(-I*A - B)*a*tan(d*x + c)^2 + 6*(d*x + c)*(A - I*B)*a + 3*(I*A + B)*a*log(tan(d*x + c)^2 + 1) - 6*(A - I*B)*a*tan(d*x + c))/d`

3.1.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 284 vs. $2(77) = 154$.

Time = 0.48 (sec) , antiderivative size = 284, normalized size of antiderivative = 3.12

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{3i Aae^{(6i dx+6i c)} \log(e^{(2i dx+2i c)} + 1) + 3 Bae^{(6i dx+6i c)} \log(e^{(2i dx+2i c)} + 1) + 9i Aae^{(4i dx+4i c)} \log(e^{(2i dx+2i c)} + 1)}{6d}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `1/3*(3*I*A*a*e^(6*I*d*x + 6*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 3*B*a*e^(6*I*d*x + 6*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 9*I*A*a*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 9*B*a*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 9*I*A*a*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 9*B*a*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 12*I*A*a*e^(4*I*d*x + 4*I*c) + 18*B*a*e^(4*I*d*x + 4*I*c) + 18*I*A*a*e^(2*I*d*x + 2*I*c) + 18*B*a*e^(2*I*d*x + 2*I*c) + 3*I*A*a*log(e^(2*I*d*x + 2*I*c) + 1) + 3*B*a*log(e^(2*I*d*x + 2*I*c) + 1) + 6*I*A*a + 8*B*a)/(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)`

3.1. $\int \tan^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

3.1.9 Mupad [B] (verification not implemented)

Time = 7.34 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.90

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{\tan(c + dx)^2 \left(\frac{Ba}{2} + \frac{Aa1i}{2}\right)}{d} - \frac{\ln(\tan(c + dx) + 1i) (Ba + Aa1i)}{d}$$

$$+ \frac{\tan(c + dx) (Aa - Ba1i)}{d} + \frac{Ba \tan(c + dx)^3 1i}{3d}$$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`output `(tan(c + d*x)^2*((A*a*1i)/2 + (B*a)/2))/d - (log(tan(c + d*x) + 1i)*(A*a*1i + B*a))/d + (tan(c + d*x)*(A*a - B*a*1i))/d + (B*a*tan(c + d*x)^3*1i)/(3*d)`

3.2 $\int \tan(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.2.1	Optimal result	305
3.2.2	Mathematica [A] (verified)	305
3.2.3	Rubi [A] (verified)	306
3.2.4	Maple [A] (verified)	307
3.2.5	Fricas [A] (verification not implemented)	308
3.2.6	Sympy [A] (verification not implemented)	308
3.2.7	Maxima [A] (verification not implemented)	309
3.2.8	Giac [B] (verification not implemented)	309
3.2.9	Mupad [B] (verification not implemented)	310

3.2.1 Optimal result

Integrand size = 30, antiderivative size = 69

$$\int \tan(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= -a(iA+B)x - \frac{a(A-iB) \log(\cos(c+dx))}{d} + \frac{a(iA+B) \tan(c+dx)}{d} + \frac{iaB \tan^2(c+dx)}{2d}$$

```
output -a*(I*A+B)*x-a*(A-I*B)*ln(cos(d*x+c))/d+a*(I*A+B)*tan(d*x+c)/d+1/2*I*a*B*tan(d*x+c)^2/d
```

3.2.2 Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.01

$$\int \tan(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{a((-2iA-2B) \arctan(\tan(c+dx)) - 2(A-iB) \log(\cos(c+dx)) + 2(iA+B) \tan(c+dx) + iB \tan^2(c+dx))}{2d}$$

```
input Integrate[Tan[c + d*x]*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]
```

```
output (a*((( -2*I)*A - 2*B)*ArcTan[Tan[c + d*x]] - 2*(A - I*B)*Log[Cos[c + d*x]] + 2*(I*A + B)*Tan[c + d*x] + I*B*Tan[c + d*x]^2))/(2*d)
```

3.2. $\int \tan(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.2.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 4075, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4075} \\
 & \int \tan(c+dx)(a(A-iB)+a(iA+B) \tan(c+dx)) dx + \frac{iaB \tan^2(c+dx)}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c+dx)(a(A-iB)+a(iA+B) \tan(c+dx)) dx + \frac{iaB \tan^2(c+dx)}{2d} \\
 & \quad \downarrow \text{4008} \\
 & a(A-iB) \int \tan(c+dx) dx + \frac{a(B+iA) \tan(c+dx)}{d} - ax(B+iA) + \frac{iaB \tan^2(c+dx)}{2d} \\
 & \quad \downarrow \text{3042} \\
 & a(A-iB) \int \tan(c+dx) dx + \frac{a(B+iA) \tan(c+dx)}{d} - ax(B+iA) + \frac{iaB \tan^2(c+dx)}{2d} \\
 & \quad \downarrow \text{3956} \\
 & \frac{a(B+iA) \tan(c+dx)}{d} - \frac{a(A-iB) \log(\cos(c+dx))}{d} - ax(B+iA) + \frac{iaB \tan^2(c+dx)}{2d}
 \end{aligned}$$

input `Int[Tan[c + d*x]*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `-(a*(I*A + B)*x) - (a*(A - I*B)*Log[Cos[c + d*x]])/d + (a*(I*A + B)*Tan[c + d*x])/d + ((I/2)*a*B*Tan[c + d*x]^2)/d`

3.2. $\int \tan(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.2.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_. + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.2.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.04

method	result
derivativedivides	$\frac{a \left(\frac{iB \tan^2(dx+c)}{2} + iA \tan(dx+c) + B \tan(dx+c) + \frac{(-iB+A) \ln(1+\tan^2(dx+c))}{2} + (-iA-B) \arctan(\tan(dx+c)) \right)}{d}$
default	$a \left(\frac{iB \tan^2(dx+c)}{2} + iA \tan(dx+c) + B \tan(dx+c) + \frac{(-iB+A) \ln(1+\tan^2(dx+c))}{2} + (-iA-B) \arctan(\tan(dx+c)) \right) / d$
norman	$(-iaA - Ba) x + \frac{(iaA+Ba) \tan(dx+c)}{d} + \frac{iaB \tan^2(dx+c)}{2d} + \frac{(-iaB+aA) \ln(1+\tan^2(dx+c))}{2d}$
parts	$\frac{(iaA+Ba)(\tan(dx+c) - \arctan(\tan(dx+c)))}{d} + \frac{aA \ln(1+\tan^2(dx+c))}{2d} + \frac{iaB \left(\frac{\tan^2(dx+c)}{2} - \frac{\ln(1+\tan^2(dx+c))}{2} \right)}{d}$
parallelrisch	$-\frac{2iAxad - iaB \tan^2(dx+c) - 2iA \tan(dx+c)a + iB \ln(1+\tan^2(dx+c))a + 2Bxad - aA \ln(1+\tan^2(dx+c)) - 2B \tan(dx+c)}{2d}$
risch	$\frac{2aBc}{d} + \frac{2iaAc}{d} + \frac{2ia(iA e^{2i(dx+c)} + 2B e^{2i(dx+c)} + iA + B)}{d(e^{2i(dx+c)} + 1)^2} + \frac{ia \ln(e^{2i(dx+c)} + 1)B}{d} - \frac{a \ln(e^{2i(dx+c)} + 1)A}{d}$

3.2. $\int \tan(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

```
input int(tan(d*x+c)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/d*a*(1/2*I*B*tan(d*x+c)^2+I*A*tan(d*x+c)+B*tan(d*x+c)+1/2*(A-I*B)*ln(1+tan(d*x+c)^2)+(-I*A-B)*arctan(tan(d*x+c)))
```

3.2.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.58

$$\int \tan(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{2(A - 2iB)ae^{(2i dx + 2i c)} + 2(A - iB)a + ((A - iB)ae^{(4i dx + 4i c)} + 2(A - iB)ae^{(2i dx + 2i c)} + (A - iB)a)}{de^{(4i dx + 4i c)} + 2de^{(2i dx + 2i c)} + d}$$

```
input integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output -(2*(A - 2*I*B)*a*e^(2*I*d*x + 2*I*c) + 2*(A - I*B)*a + ((A - I*B)*a*e^(4*I*d*x + 4*I*c) + 2*(A - I*B)*a*e^(2*I*d*x + 2*I*c) + (A - I*B)*a)*log(e^(2*I*d*x + 2*I*c) + 1)/(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)
```

3.2.6 Sympy [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.58

$$\int \tan(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx = -\frac{a(A - iB) \log(e^{2idx} + e^{-2ic})}{d} + \frac{-2Aa + 2iBa + (-2Aae^{2ic} + 4iBae^{2ic})e^{2idx}}{de^{4ic}e^{4idx} + 2de^{2ic}e^{2idx} + d}$$

```
input integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)
```

```
output -a*(A - I*B)*log(exp(2*I*d*x) + exp(-2*I*c))/d + (-2*A*a + 2*I*B*a + (-2*A*a*exp(2*I*c) + 4*I*B*a*exp(2*I*c))*exp(2*I*d*x))/(d*exp(4*I*c)*exp(4*I*d*x) + 2*d*exp(2*I*c)*exp(2*I*d*x) + d)
```

3.2. $\int \tan(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

3.2.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.99

$$\int \tan(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{-i Ba \tan(dx + c)^2 - 2(dx + c)(-iA - B)a - (A - iB)a \log(\tan(dx + c)^2 + 1) + 2(-iA - B)a \tan(dx + c)}{2d}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(-I*B*a*tan(d*x + c)^2 - 2*(d*x + c)*(-I*A - B)*a - (A - I*B)*a*log(tan(d*x + c)^2 + 1) + 2*(-I*A - B)*a*tan(d*x + c))/d`

3.2.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 194 vs. 2(59) = 118.

Time = 0.34 (sec) , antiderivative size = 194, normalized size of antiderivative = 2.81

$$\int \tan(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{Aae^{(4i dx+4i c)} \log(e^{(2i dx+2i c)} + 1) - i Ba e^{(4i dx+4i c)} \log(e^{(2i dx+2i c)} + 1) + 2 Aae^{(2i dx+2i c)} \log(e^{(2i dx+2i c)})}{2d}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `-(A*a*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - I*B*a*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 2*A*a*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 2*I*B*a*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 2*A*a*e^(2*I*d*x + 2*I*c) - 4*I*B*a*e^(2*I*d*x + 2*I*c) + A*a*log(e^(2*I*d*x + 2*I*c) + 1) - I*B*a*log(e^(2*I*d*x + 2*I*c) + 1) + 2*A*a - 2*I*B*a)/(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)`

3.2.9 Mupad [B] (verification not implemented)

Time = 7.89 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.86

$$\int \tan(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx) + 1i)(Aa - Ba1i)}{d} + \frac{\tan(c + dx)(Ba + Aa1i)}{d} + \frac{Ba \tan(c + dx)^2 1i}{2d}$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`

output `(log(tan(c + d*x) + 1i)*(A*a - B*a*1i))/d + (tan(c + d*x)*(A*a*1i + B*a))/d + (B*a*tan(c + d*x)^2*1i)/(2*d)`

3.3 $\int (a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

3.3.1	Optimal result	311
3.3.2	Mathematica [A] (verified)	311
3.3.3	Rubi [A] (verified)	312
3.3.4	Maple [A] (verified)	313
3.3.5	Fricas [A] (verification not implemented)	313
3.3.6	Sympy [A] (verification not implemented)	314
3.3.7	Maxima [A] (verification not implemented)	314
3.3.8	Giac [B] (verification not implemented)	315
3.3.9	Mupad [B] (verification not implemented)	315

3.3.1 Optimal result

Integrand size = 24, antiderivative size = 46

$$\int (a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= a(A - iB)x - \frac{a(iA + B) \log(\cos(c + dx))}{d} + \frac{iaB \tan(c + dx)}{d}$$

output `a*(A-I*B)*x-a*(I*A+B)*ln(cos(d*x+c))/d+I*a*B*tan(d*x+c)/d`

3.3.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.43

$$\int (a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= aAx - \frac{iaB \arctan(\tan(c + dx))}{d} - \frac{iaA \log(\cos(c + dx))}{d}$$

$$- \frac{aB \log(\cos(c + dx))}{d} + \frac{iaB \tan(c + dx)}{d}$$

input `Integrate[(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `a*A*x - (I*a*B*ArcTan[Tan[c + d*x]])/d - (I*a*A*Log[Cos[c + d*x]])/d - (a*B*Log[Cos[c + d*x]])/d + (I*a*B*Tan[c + d*x])/d`

3.3.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4008} \\
 & a(B + iA) \int \tan(c + dx) dx + ax(A - iB) + \frac{iaB \tan(c + dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & a(B + iA) \int \tan(c + dx) dx + ax(A - iB) + \frac{iaB \tan(c + dx)}{d} \\
 & \quad \downarrow \text{3956} \\
 & -\frac{a(B + iA) \log(\cos(c + dx))}{d} + ax(A - iB) + \frac{iaB \tan(c + dx)}{d}
 \end{aligned}$$

input `Int[(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `a*(A - I*B)*x - (a*(I*A + B)*Log[Cos[c + d*x]])/d + (I*a*B*Tan[c + d*x])/d`

3.3.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

```
rule 4008 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)]), x_Symbol] :> Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f),
x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e,
f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]
```

3.3.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.09

method	result	size
derivativedivides	$\frac{a \left(iB \tan(dx+c) + \frac{(iA+B) \ln(1+\tan^2(dx+c))}{2} + (-iB+A) \arctan(\tan(dx+c)) \right)}{d}$	50
default	$\frac{a \left(iB \tan(dx+c) + \frac{(iA+B) \ln(1+\tan^2(dx+c))}{2} + (-iB+A) \arctan(\tan(dx+c)) \right)}{d}$	50
norman	$(-iaB + aA)x + \frac{iaB \tan(dx+c)}{d} + \frac{(iaA+Ba) \ln(1+\tan^2(dx+c))}{2d}$	52
parts	$Aax + \frac{(iaA+Ba) \ln(1+\tan^2(dx+c))}{2d} + \frac{iaB(\tan(dx+c) - \arctan(\tan(dx+c)))}{d}$	55
parallelrisc	$\frac{-2iBxad + ia \ln(1+\tan^2(dx+c))a + 2Axad + 2iB \tan(dx+c)a + B \ln(1+\tan^2(dx+c))a}{2d}$	61
risc	$\frac{2iaBc}{d} - \frac{2aAc}{d} - \frac{2aB}{d(e^{2i(dx+c)}+1)} - \frac{a \ln(e^{2i(dx+c)}+1)B}{d} - \frac{ia \ln(e^{2i(dx+c)}+1)A}{d}$	78

```
input int((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/d*a*(I*B*tan(d*x+c)+1/2*(I*A+B)*ln(1+tan(d*x+c)^2)+(A-I*B)*arctan(tan(d*
x+c)))
```

3.3.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.39

$$\int (a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= -\frac{2Ba - ((-iA - B)ae^{(2i dx+2i c)} + (-iA - B)a) \log(e^{(2i dx+2i c)} + 1)}{de^{(2i dx+2i c)} + d}$$

```
input integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")
```


output $-(2*B*a - ((-I*A - B)*a*e^{(2*I*d*x + 2*I*c)} + (-I*A - B)*a)*\log(e^{(2*I*d*x + 2*I*c)} + 1))/(d*e^{(2*I*d*x + 2*I*c)} + d)$

3.3.6 Sympy [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.15

$$\int (a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= -\frac{2Ba}{de^{2ic}e^{2idx} + d} - \frac{ia(A - iB) \log(e^{2idx} + e^{-2ic})}{d}$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output $-2*B*a/(d*\exp(2*I*c)*\exp(2*I*d*x) + d) - I*a*(A - I*B)*\log(\exp(2*I*d*x) + \exp(-2*I*c))/d$

3.3.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.09

$$\int (a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2(dx + c)(A - iB)a - (-iA - B)a \log(\tan(dx + c)^2 + 1) + 2iBa \tan(dx + c)}{2d}$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output $1/2*(2*(d*x + c)*(A - I*B)*a - (-I*A - B)*a*\log(\tan(d*x + c)^2 + 1) + 2*I*B*a*\tan(d*x + c))/d$

3.3.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(40) = 80$.

Time = 0.30 (sec) , antiderivative size = 103, normalized size of antiderivative = 2.24

$$\int (a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{-i A a e^{(2i dx + 2i c)} \log(e^{(2i dx + 2i c)} + 1) - B a e^{(2i dx + 2i c)} \log(e^{(2i dx + 2i c)} + 1) - i A a \log(e^{(2i dx + 2i c)} + 1) - B a}{d e^{(2i dx + 2i c)} + d}$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `(-I*A*a*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - B*a*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - I*A*a*log(e^(2*I*d*x + 2*I*c) + 1) - B*a*log(e^(2*I*d*x + 2*I*c) + 1) - 2*B*a)/(d*e^(2*I*d*x + 2*I*c) + d)`

3.3.9 Mupad [B] (verification not implemented)

Time = 7.80 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.83

$$\int (a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx) + 1i) (B a + A a 1i)}{d} + \frac{B a \tan(c + dx) 1i}{d}$$

input `int((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`

output `(log(tan(c + d*x) + 1i)*(A*a*1i + B*a))/d + (B*a*tan(c + d*x)*1i)/d`

3.4 $\int \cot(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

3.4.1	Optimal result	316
3.4.2	Mathematica [A] (verified)	316
3.4.3	Rubi [A] (verified)	317
3.4.4	Maple [A] (verified)	319
3.4.5	Fricas [A] (verification not implemented)	319
3.4.6	Sympy [B] (verification not implemented)	320
3.4.7	Maxima [A] (verification not implemented)	320
3.4.8	Giac [B] (verification not implemented)	321
3.4.9	Mupad [B] (verification not implemented)	321

3.4.1 Optimal result

Integrand size = 30, antiderivative size = 40

$$\int \cot(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= a(iA + B)x - \frac{iaB \log(\cos(c + dx))}{d} + \frac{aA \log(\sin(c + dx))}{d}$$

output `a*(I*A+B)*x-I*a*B*ln(cos(d*x+c))/d+a*A*ln(sin(d*x+c))/d`

3.4.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.35

$$\int \cot(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= iaAx + aBx + \frac{aA \log(\cos(c + dx))}{d} - \frac{iaB \log(\cos(c + dx))}{d} + \frac{aA \log(\tan(c + dx))}{d}$$

input `Integrate[Cot[c + d*x]*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `I*a*A*x + a*B*x + (a*A*Log[Cos[c + d*x]])/d - (I*a*B*Log[Cos[c + d*x]])/d + (a*A*Log[Tan[c + d*x]])/d`

3.4. $\int \cot(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

3.4.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3042, 4072, 3042, 3956, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan(c+dx)} dx \\
 & \quad \downarrow \text{4072} \\
 & \int \cot(c+dx)(aA+a(iA+B) \tan(c+dx)) dx + iaB \int \tan(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{aA+a(iA+B) \tan(c+dx)}{\tan(c+dx)} dx + iaB \int \tan(c+dx) dx \\
 & \quad \downarrow \text{3956} \\
 & \int \frac{aA+a(iA+B) \tan(c+dx)}{\tan(c+dx)} dx - \frac{iaB \log(\cos(c+dx))}{d} \\
 & \quad \downarrow \text{4014} \\
 & aA \int \cot(c+dx) dx + ax(B+iA) - \frac{iaB \log(\cos(c+dx))}{d} \\
 & \quad \downarrow \text{3042} \\
 & aA \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx + ax(B+iA) - \frac{iaB \log(\cos(c+dx))}{d} \\
 & \quad \downarrow \text{25} \\
 & -aA \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx + ax(B+iA) - \frac{iaB \log(\cos(c+dx))}{d} \\
 & \quad \downarrow \text{3956} \\
 & ax(B+iA) + \frac{aA \log(-\sin(c+dx))}{d} - \frac{iaB \log(\cos(c+dx))}{d}
 \end{aligned}$$

input `Int[Cot[c + d*x]*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `a*(I*A + B)*x - (I*a*B*Log[Cos[c + d*x]])/d + (a*A*Log[-Sin[c + d*x]])/d`

3.4.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4072 `Int[((((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*(d/b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c - a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0]`

3.4.4 Maple [A] (verified)

Time = 0.56 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.08

method	result	size
parallelrisc	$\frac{a\left(\left(-\frac{A}{2} + \frac{iB}{2}\right) \ln(\sec^2(dx+c)) + A \ln(\tan(dx+c)) + (iA+B)xd\right)}{d}$	43
derivativedivides	$\frac{a\left(\frac{(iB-A) \ln(1+\tan^2(dx+c))}{2} + (iA+B) \arctan(\tan(dx+c)) + A \ln(\tan(dx+c))\right)}{d}$	51
default	$\frac{a\left(\frac{(iB-A) \ln(1+\tan^2(dx+c))}{2} + (iA+B) \arctan(\tan(dx+c)) + A \ln(\tan(dx+c))\right)}{d}$	51
norman	$(iaA + Ba)x + \frac{aA \ln(\tan(dx+c))}{d} - \frac{(-iaB+aA) \ln(1+\tan^2(dx+c))}{2d}$	51
risc	$-\frac{2iaAc}{d} - \frac{2aBc}{d} + \frac{aA \ln(e^{2i(dx+c)}-1)}{d} - \frac{ia \ln(e^{2i(dx+c)}+1)B}{d}$	57

input `int(cot(d*x+c)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `a*((-1/2*A+1/2*I*B)*ln(sec(d*x+c)^2)+A*ln(tan(d*x+c))+(I*A+B)*x*d)/d`

3.4.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.90

$$\int \cot(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{-iBa \log(e^{(2i dx+2i c)}+1) + Aa \log(e^{(2i dx+2i c)}-1)}{d}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `(-I*B*a*log(e^(2*I*d*x + 2*I*c) + 1) + A*a*log(e^(2*I*d*x + 2*I*c) - 1))/d`

3.4.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 94 vs. $2(36) = 72$.

Time = 1.03 (sec) , antiderivative size = 94, normalized size of antiderivative = 2.35

$$\int \cot(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{Aa \log\left(\frac{-Aa - iBa}{Aae^{2ic} + iBae^{2ic}} + e^{2idx}\right)}{d} - \frac{iBa \log\left(\frac{Aa + iBa}{Aae^{2ic} + iBae^{2ic}} + e^{2idx}\right)}{d}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `A*a*log((-A*a - I*B*a)/(A*a*exp(2*I*c) + I*B*a*exp(2*I*c)) + exp(2*I*d*x))/d - I*B*a*log((A*a + I*B*a)/(A*a*exp(2*I*c) + I*B*a*exp(2*I*c)) + exp(2*I*d*x))/d`

3.4.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.22

$$\int \cot(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2(dx + c)(iA + B)a - (A - iB)a \log(\tan(dx + c)^2 + 1) + 2Aa \log(\tan(dx + c))}{2d}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(2*(d*x + c)*(I*A + B)*a - (A - I*B)*a*log(tan(d*x + c)^2 + 1) + 2*A*a*log(tan(d*x + c)))/d`

3.4.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 74 vs. $2(36) = 72$.

Time = 0.40 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.85

$$\int \cot(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{i Ba \log(\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1) + i Ba \log(\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1) - Aa \log(\tan(\frac{1}{2} dx + \frac{1}{2} c)) + 2(Aa - I^2 B a)}{d}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `-(I*B*a*log(tan(1/2*d*x + 1/2*c) + 1) + I*B*a*log(tan(1/2*d*x + 1/2*c) - 1) - A*a*log(tan(1/2*d*x + 1/2*c)) + 2*(A*a - I*B*a)*log(tan(1/2*d*x + 1/2*c) + I))/d`

3.4.9 Mupad [B] (verification not implemented)

Time = 7.70 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.90

$$\int \cot(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{A a \ln(\tan(c + dx))}{d} - \frac{a \ln(\tan(c + dx) + 1i) (A - B 1i)}{d}$$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`

output `(A*a*log(tan(c + d*x)))/d - (a*log(tan(c + d*x) + 1i)*(A - B*1i))/d`

3.5 $\int \cot^2(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.5.1	Optimal result	322
3.5.2	Mathematica [C] (verified)	322
3.5.3	Rubi [A] (verified)	323
3.5.4	Maple [A] (verified)	325
3.5.5	Fricas [A] (verification not implemented)	325
3.5.6	Sympy [A] (verification not implemented)	326
3.5.7	Maxima [A] (verification not implemented)	326
3.5.8	Giac [B] (verification not implemented)	326
3.5.9	Mupad [B] (verification not implemented)	327

3.5.1 Optimal result

Integrand size = 32, antiderivative size = 44

$$\int \cot^2(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= -a(A-iB)x - \frac{aA \cot(c+dx)}{d} + \frac{a(iA+B) \log(\sin(c+dx))}{d}$$

output

```
-a*(A-I*B)*x-a*A*cot(d*x+c)/d+a*(I*A+B)*ln(sin(d*x+c))/d
```

3.5.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.04 (sec) , antiderivative size = 97, normalized size of antiderivative = 2.20

$$\int \cot^2(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= iaBx - \frac{aA \cot(c+dx) \text{Hypergeometric2F1}(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(c+dx))}{d}$$

$$+ \frac{iaA \log(\cos(c+dx))}{d} + \frac{aB \log(\cos(c+dx))}{d}$$

$$+ \frac{iaA \log(\tan(c+dx))}{d} + \frac{aB \log(\tan(c+dx))}{d}$$

input `Integrate[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `I*a*B*x - (a*A*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2])/d + (I*a*A*Log[Cos[c + d*x]])/d + (a*B*Log[Cos[c + d*x]])/d + (I*a*A*Log[Tan[c + d*x]])/d + (a*B*Log[Tan[c + d*x]])/d`

3.5.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.05, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.219$, Rules used = {3042, 4074, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan(c + dx)^2} dx \\
 & \quad \downarrow \text{4074} \\
 & -\frac{aA \cot(c + dx)}{d} + \int \cot(c + dx)(a(iA + B) - a(A - iB) \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & -\frac{aA \cot(c + dx)}{d} + \int \frac{a(iA + B) - a(A - iB) \tan(c + dx)}{\tan(c + dx)} dx \\
 & \quad \downarrow \text{4014} \\
 & a(B + iA) \int \cot(c + dx) dx - ax(A - iB) - \frac{aA \cot(c + dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & a(B + iA) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - ax(A - iB) - \frac{aA \cot(c + dx)}{d} \\
 & \quad \downarrow \text{25} \\
 & -a(B + iA) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - ax(A - iB) - \frac{aA \cot(c + dx)}{d} \\
 & \quad \downarrow \text{3956}
 \end{aligned}$$

3.5. $\int \cot^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\frac{a(B + iA) \log(-\sin(c + dx))}{d} - ax(A - iB) - \frac{aA \cot(c + dx)}{d}$$

input `Int[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `-(a*(A - I*B)*x) - (a*A*Cot[c + d*x])/d + (a*(I*A + B)*Log[-Sin[c + d*x]])/d`

3.5.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4074 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.5.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.34

method	result	size
parallelrisc	$\frac{\left(\left(-\frac{iA}{2}-\frac{B}{2}\right)\ln(\sec^2(dx+c))+iA+B\right)\ln(\tan(dx+c))-A\cot(dx+c)+xd(iB-A)a}{d}$	59
derivativedivides	$\frac{a\left(\frac{(-iA-B)\ln(1+\tan^2(dx+c))}{2}+(iB-A)\arctan(\tan(dx+c))-\frac{A}{\tan(dx+c)}+(iA+B)\ln(\tan(dx+c))\right)}{d}$	69
default	$\frac{a\left(\frac{(-iA-B)\ln(1+\tan^2(dx+c))}{2}+(iB-A)\arctan(\tan(dx+c))-\frac{A}{\tan(dx+c)}+(iA+B)\ln(\tan(dx+c))\right)}{d}$	69
risc	$-\frac{2iaBc}{d}+\frac{2aAc}{d}-\frac{2iaA}{d(e^{2i(dx+c)}-1)}+\frac{a\ln(e^{2i(dx+c)}-1)B}{d}+\frac{ia\ln(e^{2i(dx+c)}-1)A}{d}$	78
norman	$\frac{(iaB-aA)x\tan(dx+c)-\frac{aA}{d}}{\tan(dx+c)}+\frac{(iaA+Ba)\ln(\tan(dx+c))}{d}-\frac{(iaA+Ba)\ln(1+\tan^2(dx+c))}{2d}$	82

input `int(cot(d*x+c)^2*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `((-1/2*I*A-1/2*B)*ln(sec(d*x+c)^2)+(I*A+B)*ln(tan(d*x+c))-A*cot(d*x+c)+x*d*(-A+I*B))*a/d`

3.5.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.41

$$\int \cot^2(c+dx)(a+ia\tan(c+dx))(A+B\tan(c+dx))dx$$

$$= \frac{-2iAa + ((iA+B)ae^{(2i dx+2i c)} + (-iA-B)a)\log(e^{(2i dx+2i c)} - 1)}{de^{(2i dx+2i c)} - d}$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `(-2*I*A*a + ((I*A + B)*a*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a)*log(e^(2*I*d*x + 2*I*c) - 1))/(d*e^(2*I*d*x + 2*I*c) - d)`

3.5. $\int \cot^2(c+dx)(a+ia\tan(c+dx))(A+B\tan(c+dx))dx$

3.5.6 Sympy [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.20

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= -\frac{2iAa}{de^{2ic}e^{2idx} - d} + \frac{ia(A - iB) \log(e^{2idx} - e^{-2ic})}{d}$$

input `integrate(cot(d*x+c)**2*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)), x)`

output `-2*I*A*a/(d*exp(2*I*c)*exp(2*I*d*x) - d) + I*a*(A - I*B)*log(exp(2*I*d*x) - exp(-2*I*c))/d`

3.5.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.45

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx =$$

$$\frac{2(dx + c)(A - iB)a + (iA + B)a \log(\tan(dx + c)^2 + 1) - 2(iA + B)a \log(\tan(dx + c)) + \frac{2Aa}{\tan(dx + c)}}{2d}$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)), x, algorithm="maxima")`

output `-1/2*(2*(d*x + c)*(A - I*B)*a + (I*A + B)*a*log(tan(d*x + c)^2 + 1) - 2*(I*A + B)*a*log(tan(d*x + c)) + 2*A*a/tan(d*x + c))/d`

3.5.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 104 vs. $2(40) = 80$.

Time = 0.49 (sec) , antiderivative size = 104, normalized size of antiderivative = 2.36

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{Aa \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 4(-iAa - Ba) \log\left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + i\right) + 2(iAa + Ba) \log\left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right) + \dots}{2d}$$

3.5. $\int \cot^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `1/2*(A*a*tan(1/2*d*x + 1/2*c) + 4*(-I*A*a - B*a)*log(tan(1/2*d*x + 1/2*c) + I) + 2*(I*A*a + B*a)*log(tan(1/2*d*x + 1/2*c)) + (-2*I*A*a*tan(1/2*d*x + 1/2*c) - 2*B*a*tan(1/2*d*x + 1/2*c) - A*a)/tan(1/2*d*x + 1/2*c))/d`

3.5.9 Mupad [B] (verification not implemented)

Time = 7.23 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.89

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= -\frac{A a \cot(c + dx)}{d} + \frac{a \operatorname{atan}(2 \tan(c + dx) + 1i) (B + A 1i) 2i}{d}$$

input `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`

output `(a*atan(2*tan(c + d*x) + 1i)*(A*1i + B)*2i)/d - (A*a*cot(c + d*x))/d`

3.6 $\int \cot^3(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.6.1	Optimal result	328
3.6.2	Mathematica [C] (verified)	328
3.6.3	Rubi [A] (verified)	329
3.6.4	Maple [A] (verified)	331
3.6.5	Fricas [A] (verification not implemented)	332
3.6.6	Sympy [A] (verification not implemented)	332
3.6.7	Maxima [A] (verification not implemented)	333
3.6.8	Giac [B] (verification not implemented)	333
3.6.9	Mupad [B] (verification not implemented)	334

3.6.1 Optimal result

Integrand size = 32, antiderivative size = 68

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= -a(iA+B)x - \frac{a(iA+B) \cot(c+dx)}{d} - \frac{aA \cot^2(c+dx)}{2d} - \frac{a(A-iB) \log(\sin(c+dx))}{d}$$

```
output -a*(I*A+B)*x-a*(I*A+B)*cot(d*x+c)/d-1/2*a*A*cot(d*x+c)^2/d-a*(A-I*B)*ln(sin(d*x+c))/d
```

3.6.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.42 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.12

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx =$$

$$\frac{a(A \cot^2(c+dx) + 2(iA+B) \cot(c+dx) \text{Hypergeometric2F1}(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(c+dx)) + 2(A-iB))}{2d}$$

```
input Integrate[Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]
```

output
$$\frac{-1/2*(a*(A*\text{Cot}[c + d*x]^2 + 2*(I*A + B)*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[-1/2, 1, 1/2, -\text{Tan}[c + d*x]^2] + 2*(A - I*B)*(\text{Log}[\text{Cos}[c + d*x]] + \text{Log}[\text{Tan}[c + d*x]])))/d}{d}$$

3.6.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3042, 4074, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot^3(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\ & \quad \downarrow 3042 \\ & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan(c + dx)^3} dx \\ & \quad \downarrow 4074 \\ & -\frac{aA \cot^2(c + dx)}{2d} + \int \cot^2(c + dx)(a(iA + B) - a(A - iB) \tan(c + dx)) dx \\ & \quad \downarrow 3042 \\ & -\frac{aA \cot^2(c + dx)}{2d} + \int \frac{a(iA + B) - a(A - iB) \tan(c + dx)}{\tan(c + dx)^2} dx \\ & \quad \downarrow 4012 \\ & \int -\cot(c + dx)(a(A - iB) + a(iA + B) \tan(c + dx)) dx - \frac{a(B + iA) \cot(c + dx)}{d} - \frac{aA \cot^2(c + dx)}{2d} \\ & \quad \downarrow 25 \\ & -\int \cot(c + dx)(a(A - iB) + a(iA + B) \tan(c + dx)) dx - \frac{a(B + iA) \cot(c + dx)}{d} - \frac{aA \cot^2(c + dx)}{2d} \\ & \quad \downarrow 3042 \\ & -\int \frac{a(A - iB) + a(iA + B) \tan(c + dx)}{\tan(c + dx)} dx - \frac{a(B + iA) \cot(c + dx)}{d} - \frac{aA \cot^2(c + dx)}{2d} \\ & \quad \downarrow 4014 \\ & -a(A - iB) \int \cot(c + dx) dx - \frac{a(B + iA) \cot(c + dx)}{d} - ax(B + iA) - \frac{aA \cot^2(c + dx)}{2d} \end{aligned}$$

3.6. $\int \cot^3(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \downarrow 3042 \\
& -a(A - iB) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - \frac{a(B + iA) \cot(c + dx)}{d} - ax(B + iA) - \frac{aA \cot^2(c + dx)}{2d} \\
& \downarrow 25 \\
& a(A - iB) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - \frac{a(B + iA) \cot(c + dx)}{d} - ax(B + iA) - \frac{aA \cot^2(c + dx)}{2d} \\
& \downarrow 3956 \\
& -\frac{a(B + iA) \cot(c + dx)}{d} - \frac{a(A - iB) \log(-\sin(c + dx))}{d} - ax(B + iA) - \frac{aA \cot^2(c + dx)}{2d}
\end{aligned}$$

input `Int[Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `-(a*(I*A + B)*x) - (a*(I*A + B)*Cot[c + d*x])/d - (a*A*Cot[c + d*x]^2)/(2*d) - (a*(A - I*B)*Log[-Sin[c + d*x]])/d`

3.6.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)
)*(x_)], x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

3.6.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.07

method	result
parallelrisc	$-\frac{\left(\left(-\frac{A}{2} + \frac{iB}{2}\right) \ln(\sec^2(dx+c)) + (-iB+A) \ln(\tan(dx+c)) + \frac{A(\cot^2(dx+c))}{2} + \cot(dx+c)(iA+B) + (iA+B)xd\right)a}{d}$
derivativedivides	$a \frac{\left(\frac{(-iB+A) \ln(1+\tan^2(dx+c))}{2} + (-iA-B) \arctan(\tan(dx+c)) - \frac{A}{2 \tan(dx+c)^2} + (iB-A) \ln(\tan(dx+c)) - \frac{iA+B}{\tan(dx+c)}\right)}{d}$
default	$a \frac{\left(\frac{(-iB+A) \ln(1+\tan^2(dx+c))}{2} + (-iA-B) \arctan(\tan(dx+c)) - \frac{A}{2 \tan(dx+c)^2} + (iB-A) \ln(\tan(dx+c)) - \frac{iA+B}{\tan(dx+c)}\right)}{d}$
norman	$\frac{(-iaA - Ba)x(\tan^2(dx+c)) - \frac{aA}{2d} - \frac{(iaA + Ba)\tan(dx+c)}{d}}{\tan(dx+c)^2} - \frac{(-iaB + aA) \ln(\tan(dx+c))}{d} + \frac{(-iaB + aA) \ln(1 + \tan^2(dx+c))}{2d}$
risc	$\frac{2aBc}{d} + \frac{2iaAc}{d} - \frac{2ia(2iAe^{2i(dx+c)} + Be^{2i(dx+c)} - iA - B)}{d(e^{2i(dx+c)} - 1)^2} + \frac{ia \ln(e^{2i(dx+c)} - 1)B}{d} - \frac{aA \ln(e^{2i(dx+c)} - 1)}{d}$

```
input int(cot(d*x+c)^3*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

```
output -((-1/2*A+1/2*I*B)*ln(sec(d*x+c)^2)+(A-I*B)*ln(tan(d*x+c))+1/2*A*cot(d*x+c
)^2+cot(d*x+c)*(I*A+B)+(I*A+B)*x*d)*a/d
```

3.6. $\int \cot^3(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

3.6.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.63

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2(2A - iB)ae^{(2idx+2ic)} - 2(A - iB)a - ((A - iB)ae^{(4idx+4ic)} - 2(A - iB)ae^{(2idx+2ic)} + (A - iB)a)}{de^{(4idx+4ic)} - 2de^{(2idx+2ic)} + d}$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `(2*(2*A - I*B)*a*e^(2*I*d*x + 2*I*c) - 2*(A - I*B)*a - ((A - I*B)*a*e^(4*I*d*x + 4*I*c) - 2*(A - I*B)*a*e^(2*I*d*x + 2*I*c) + (A - I*B)*a)*log(e^(2*I*d*x + 2*I*c) - 1)/(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)`

3.6.6 Sympy [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.60

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= -\frac{a(A - iB) \log(e^{2idx} - e^{-2ic})}{d} + \frac{-2Aa + 2iBa + (4Aae^{2ic} - 2iBae^{2ic})e^{2idx}}{de^{4ic}e^{4idx} - 2de^{2ic}e^{2idx} + d}$$

input `integrate(cot(d*x+c)**3*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `-a*(A - I*B)*log(exp(2*I*d*x) - exp(-2*I*c))/d + (-2*A*a + 2*I*B*a + (4*A*a*exp(2*I*c) - 2*I*B*a*exp(2*I*c))*exp(2*I*d*x))/(d*exp(4*I*c)*exp(4*I*d*x) - 2*d*exp(2*I*c)*exp(2*I*d*x) + d)`

3.6.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.24

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2(dx + c)(-iA - B)a + (A - iB)a \log(\tan(dx + c)^2 + 1) - 2(A - iB)a \log(\tan(dx + c)) + \frac{2(-iA - B)}{\tan}}{2d}$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(2*(d*x + c)*(-I*A - B)*a + (A - I*B)*a*log(tan(d*x + c)^2 + 1) - 2*(A - I*B)*a*log(tan(d*x + c)) + (2*(-I*A - B)*a*tan(d*x + c) - A*a)/tan(d*x + c)^2)/d`

3.6.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 162 vs. 2(60) = 120.

Time = 0.61 (sec) , antiderivative size = 162, normalized size of antiderivative = 2.38

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx =$$

$$\frac{Aa \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 4i Aa \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 4Ba \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 16(Aa - iBa) \log\left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right)}{2d}$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `-1/8*(A*a*tan(1/2*d*x + 1/2*c)^2 - 4*I*A*a*tan(1/2*d*x + 1/2*c) - 4*B*a*tan(1/2*d*x + 1/2*c) - 16*(A*a - I*B*a)*log(tan(1/2*d*x + 1/2*c) + I) + 8*(A*a - I*B*a)*log(tan(1/2*d*x + 1/2*c)) - (12*A*a*tan(1/2*d*x + 1/2*c)^2 - 12*I*B*a*tan(1/2*d*x + 1/2*c)^2 - 4*I*A*a*tan(1/2*d*x + 1/2*c) - 4*B*a*tan(1/2*d*x + 1/2*c) - A*a)/tan(1/2*d*x + 1/2*c)^2)/d`

3.6.9 Mupad [B] (verification not implemented)

Time = 7.33 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.88

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= -\frac{\frac{Aa}{2} + \tan(c + dx) (Ba + Aa \text{li})}{d \tan(c + dx)^2} - \frac{a \operatorname{atan}(2 \tan(c + dx) + \text{li}) (A - B \text{li}) 2i}{d}$$

input `int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`

output `- ((A*a)/2 + tan(c + d*x)*(A*a*1i + B*a))/(d*tan(c + d*x)^2) - (a*atan(2*tan(c + d*x) + 1i)*(A - B*1i)*2i)/d`

3.7 $\int \cot^4(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.7.1	Optimal result	335
3.7.2	Mathematica [C] (verified)	335
3.7.3	Rubi [A] (verified)	336
3.7.4	Maple [A] (verified)	339
3.7.5	Fricas [B] (verification not implemented)	339
3.7.6	Sympy [B] (verification not implemented)	340
3.7.7	Maxima [A] (verification not implemented)	340
3.7.8	Giac [B] (verification not implemented)	341
3.7.9	Mupad [B] (verification not implemented)	341

3.7.1 Optimal result

Integrand size = 32, antiderivative size = 89

$$\int \cot^4(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= a(A-iB)x + \frac{a(A-iB) \cot(c+dx)}{d} - \frac{a(iA+B) \cot^2(c+dx)}{2d}$$

$$- \frac{aA \cot^3(c+dx)}{3d} - \frac{a(iA+B) \log(\sin(c+dx))}{d}$$

```
output a*(A-I*B)*x+a*(A-I*B)*cot(d*x+c)/d-1/2*a*(I*A+B)*cot(d*x+c)^2/d-1/3*a*A*co
t(d*x+c)^3/d-a*(I*A+B)*ln(sin(d*x+c))/d
```

3.7.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.79 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.15

$$\int \cot^4(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx =$$

$$\frac{a(2A \cot^3(c+dx) \text{Hypergeometric2F1}(-\frac{3}{2}, 1, -\frac{1}{2}, -\tan^2(c+dx)) + 6iB \cot(c+dx) \text{Hypergeometric2F1}(\frac{3}{2}, 1, \frac{5}{2}, -\tan^2(c+dx)))}{d}$$

input `Integrate[Cot[c + d*x]^4*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `-1/6*(a*(2*A*Cot[c + d*x]^3*Hypergeometric2F1[-3/2, 1, -1/2, -Tan[c + d*x]^2] + (6*I)*B*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2] + 3*(I*A + B)*(Cot[c + d*x]^2 + 2*(Log[Cos[c + d*x]] + Log[Tan[c + d*x]])))/d`

3.7.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3042, 4074, 3042, 4012, 25, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^4(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan(c + dx)^4} dx \\
 & \quad \downarrow \text{4074} \\
 & -\frac{aA \cot^3(c + dx)}{3d} + \int \cot^3(c + dx)(a(iA + B) - a(A - iB) \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & -\frac{aA \cot^3(c + dx)}{3d} + \int \frac{a(iA + B) - a(A - iB) \tan(c + dx)}{\tan(c + dx)^3} dx \\
 & \quad \downarrow \text{4012} \\
 & \int -\cot^2(c + dx)(a(A - iB) + a(iA + B) \tan(c + dx)) dx - \frac{a(B + iA) \cot^2(c + dx)}{2d} - \frac{aA \cot^3(c + dx)}{3d} \\
 & \quad \downarrow \text{25} \\
 & -\int \cot^2(c + dx)(a(A - iB) + a(iA + B) \tan(c + dx)) dx - \frac{a(B + iA) \cot^2(c + dx)}{2d} - \frac{aA \cot^3(c + dx)}{3d} \\
 & \quad \downarrow \text{3042} \\
 & -\int \frac{a(A - iB) + a(iA + B) \tan(c + dx)}{\tan(c + dx)^2} dx - \frac{a(B + iA) \cot^2(c + dx)}{2d} - \frac{aA \cot^3(c + dx)}{3d}
 \end{aligned}$$

3.7. $\int \cot^4(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \downarrow 4012 \\
& - \int \cot(c+dx)(a(iA+B) - a(A-iB)\tan(c+dx))dx - \frac{a(B+iA)\cot^2(c+dx)}{2d} + \\
& \quad \frac{a(A-iB)\cot(c+dx)}{d} - \frac{aA\cot^3(c+dx)}{3d} \\
& \downarrow 3042 \\
& - \int \frac{a(iA+B) - a(A-iB)\tan(c+dx)}{\tan(c+dx)} dx - \frac{a(B+iA)\cot^2(c+dx)}{2d} + \frac{a(A-iB)\cot(c+dx)}{d} - \\
& \quad \frac{aA\cot^3(c+dx)}{3d} \\
& \downarrow 4014 \\
& -a(B+iA) \int \cot(c+dx)dx - \frac{a(B+iA)\cot^2(c+dx)}{2d} + \frac{a(A-iB)\cot(c+dx)}{d} + ax(A-iB) - \\
& \quad \frac{aA\cot^3(c+dx)}{3d} \\
& \downarrow 3042 \\
& -a(B+iA) \int -\tan\left(c+dx + \frac{\pi}{2}\right) dx - \frac{a(B+iA)\cot^2(c+dx)}{2d} + \frac{a(A-iB)\cot(c+dx)}{d} + \\
& \quad ax(A-iB) - \frac{aA\cot^3(c+dx)}{3d} \\
& \downarrow 25 \\
& a(B+iA) \int \tan\left(\frac{1}{2}(2c+\pi) + dx\right) dx - \frac{a(B+iA)\cot^2(c+dx)}{2d} + \frac{a(A-iB)\cot(c+dx)}{d} + \\
& \quad ax(A-iB) - \frac{aA\cot^3(c+dx)}{3d} \\
& \downarrow 3956 \\
& -\frac{a(B+iA)\cot^2(c+dx)}{2d} + \frac{a(A-iB)\cot(c+dx)}{d} - \frac{a(B+iA)\log(-\sin(c+dx))}{d} + ax(A-iB) - \\
& \quad \frac{aA\cot^3(c+dx)}{3d}
\end{aligned}$$

input `Int[Cot[c + d*x]^4*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `a*(A - I*B)*x + (a*(A - I*B)*Cot[c + d*x])/d - (a*(I*A + B)*Cot[c + d*x]^2)/(2*d) - (a*A*Cot[c + d*x]^3)/(3*d) - (a*(I*A + B)*Log[-Sin[c + d*x]])/d`

3.7.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`
- rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`
- rule 4074 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.7.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.13

method	result
derivativedivides	$\frac{a \left(\frac{(iA+B) \ln(1+\tan^2(dx+c))}{2} + (-iB+A) \arctan(\tan(dx+c)) - \frac{A}{3 \tan(dx+c)^3} - \frac{iB-A}{\tan(dx+c)} + (-iA-B) \ln(\tan(dx+c)) - \frac{iA}{2 \tan(dx+c)} \right)}{d}$
default	$\frac{a \left(\frac{(iA+B) \ln(1+\tan^2(dx+c))}{2} + (-iB+A) \arctan(\tan(dx+c)) - \frac{A}{3 \tan(dx+c)^3} - \frac{iB-A}{\tan(dx+c)} + (-iA-B) \ln(\tan(dx+c)) - \frac{iA}{2 \tan(dx+c)} \right)}{d}$
norman	$\frac{(-iaB+aA) \frac{(\tan^2(dx+c))}{d} + (-iaB+aA)x \frac{(\tan^3(dx+c))}{\tan(dx+c)^3} - \frac{aA}{3d} - \frac{(iaA+Ba) \tan(dx+c)}{2d} - \frac{(iaA+Ba) \ln(\tan(dx+c))}{d} + \frac{(iaA+Ba) \ln(\tan(dx+c))}{d}}{\tan(dx+c)^3}$
parallelrisch	$-\frac{a(6iBdx+6iA \ln(\tan(dx+c))-3iA \ln(\sec^2(dx+c))-6Adx+3iA(\cot^2(dx+c))+6iB \cot(dx+c)+6B \ln(\tan(dx+c)))-3a}{6d}$
risch	$\frac{2iaBc}{d} - \frac{2aAc}{d} + \frac{2a(9iA e^{4i(dx+c)}+6B e^{4i(dx+c)}-9iA e^{2i(dx+c)}-9B e^{2i(dx+c)}+4iA+3B)}{3d(e^{2i(dx+c)}-1)^3} - \frac{a \ln(e^{2i(dx+c)}-1)B}{d}$

input `int(cot(d*x+c)^4*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `a/d*(1/2*(I*A+B)*ln(1+tan(d*x+c)^2)+(A-I*B)*arctan(tan(d*x+c))-1/3*A/tan(d*x+c)^3-(-A+I*B)/tan(d*x+c)+(-I*A-B)*ln(tan(d*x+c))-1/2*(I*A+B)/tan(d*x+c)^2)`

3.7.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 166 vs. 2(77) = 154.

Time = 0.25 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.87

$$\int \cot^4(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx = \frac{6(-3iA-2B)ae^{(4i dx+4i c)} + 18(iA+B)ae^{(2i dx+2i c)} + 2(-4iA-3B)a + 3((iA+B)ae^{(6i dx+6i c)} + 3(de^{(6i dx+6i c)} - 3de^{(4i dx+4i c)} + 3de^{(2i dx+2i c)} - 3de^{(0i dx+0i c)}))}{3(de^{(6i dx+6i c)} - 3de^{(4i dx+4i c)} + 3de^{(2i dx+2i c)} - 3de^{(0i dx+0i c)})}$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output
$$\begin{aligned} & -1/3*(6*(-3*I*A - 2*B))*a*e^{(4*I*d*x + 4*I*c)} + 18*(I*A + B)*a*e^{(2*I*d*x + 2*I*c)} \\ & + 2*(-4*I*A - 3*B)*a + 3*((I*A + B))*a*e^{(6*I*d*x + 6*I*c)} + 3*(-I*A - B)*a*e^{(4*I*d*x + 4*I*c)} \\ & + 3*(I*A + B)*a*e^{(2*I*d*x + 2*I*c)} + (-I*A - B)*a*\log(e^{(2*I*d*x + 2*I*c)} - 1)/(d*e^{(6*I*d*x + 6*I*c)} - 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} - d) \end{aligned}$$

3.7.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 168 vs. $2(73) = 146$.

Time = 0.37 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.89

$$\begin{aligned} & \int \cot^4(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\ & = -\frac{ia(A - iB) \log(e^{2idx} - e^{-2ic})}{d} \\ & \quad + \frac{8iAa + 6Ba + (-18iAae^{2ic} - 18Bae^{2ic})e^{2idx} + (18iAae^{4ic} + 12Bae^{4ic})e^{4idx}}{3de^{6ic}e^{6idx} - 9de^{4ic}e^{4idx} + 9de^{2ic}e^{2idx} - 3d} \end{aligned}$$

input `integrate(cot(d*x+c)**4*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output
$$\begin{aligned} & -I*a*(A - I*B)*\log(\exp(2*I*d*x) - \exp(-2*I*c))/d + (8*I*A*a + 6*B*a + (-18 \\ & *I*A*a*\exp(2*I*c) - 18*B*a*\exp(2*I*c))*\exp(2*I*d*x) + (18*I*A*a*\exp(4*I*c) \\ & + 12*B*a*\exp(4*I*c))*\exp(4*I*d*x))/(3*d*\exp(6*I*c)*\exp(6*I*d*x) - 9*d*\exp \\ & (4*I*c)*\exp(4*I*d*x) + 9*d*\exp(2*I*c)*\exp(2*I*d*x) - 3*d) \end{aligned}$$

3.7.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.16

$$\begin{aligned} & \int \cot^4(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\ & = \frac{6(dx + c)(A - iB)a - 3(-iA - B)a \log(\tan(dx + c)^2 + 1) + 6(-iA - B)a \log(\tan(dx + c)) + \frac{6(A-i)}{d}}{6d} \end{aligned}$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

3.7. $\int \cot^4(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

output $1/6*(6*(d*x + c)*(A - I*B)*a - 3*(-I*A - B)*a*\log(\tan(d*x + c)^2 + 1) + 6*(-I*A - B)*a*\log(\tan(d*x + c)) + (6*(A - I*B)*a*\tan(d*x + c)^2 + 3*(-I*A - B)*a*\tan(d*x + c) - 2*A*a)/\tan(d*x + c)^3)/d$

3.7.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 221 vs. $2(77) = 154$.

Time = 0.71 (sec) , antiderivative size = 221, normalized size of antiderivative = 2.48

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{Aa \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 3i Aa \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 3Ba \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 15Aa \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 12i Ba}{d}$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output $1/24*(A*a*\tan(1/2*d*x + 1/2*c)^3 - 3*I*A*a*\tan(1/2*d*x + 1/2*c)^2 - 3*B*a*\tan(1/2*d*x + 1/2*c)^2 - 15*A*a*\tan(1/2*d*x + 1/2*c) + 12*I*B*a*\tan(1/2*d*x + 1/2*c) + 48*(I*A*a + B*a)*\log(\tan(1/2*d*x + 1/2*c) + I) - 24*(I*A*a + B*a)*\log(\tan(1/2*d*x + 1/2*c)) - (-44*I*A*a*\tan(1/2*d*x + 1/2*c)^3 - 44*B*a*\tan(1/2*d*x + 1/2*c)^3 - 15*A*a*\tan(1/2*d*x + 1/2*c)^2 + 12*I*B*a*\tan(1/2*d*x + 1/2*c)^2 + 3*I*A*a*\tan(1/2*d*x + 1/2*c) + 3*B*a*\tan(1/2*d*x + 1/2*c) + A*a)/\tan(1/2*d*x + 1/2*c)^3)/d$

3.7.9 Mupad [B] (verification not implemented)

Time = 7.57 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.90

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= -\frac{(-Aa + Ba \operatorname{li}) \tan(c + dx)^2 + \left(\frac{Ba}{2} + \frac{Aa \operatorname{li}}{2}\right) \tan(c + dx) + \frac{Aa}{3}}{d \tan(c + dx)^3}$$

$$- \frac{a \operatorname{atan}(2 \tan(c + dx) + \operatorname{li}) (B + A \operatorname{li}) 2i}{d}$$

3.7. $\int \cot^4(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

input `int(cot(c + d*x)^4*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`

output `- ((A*a)/3 + tan(c + d*x)*((A*a*1i)/2 + (B*a)/2) - tan(c + d*x)^2*(A*a - B*a*1i))/(d*tan(c + d*x)^3) - (a*atan(2*tan(c + d*x) + 1i)*(A*1i + B)*2i)/d`

3.8 $\int \cot^5(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.8.1	Optimal result	343
3.8.2	Mathematica [C] (verified)	343
3.8.3	Rubi [A] (verified)	344
3.8.4	Maple [A] (verified)	347
3.8.5	Fricas [B] (verification not implemented)	348
3.8.6	Sympy [B] (verification not implemented)	348
3.8.7	Maxima [A] (verification not implemented)	349
3.8.8	Giac [B] (verification not implemented)	349
3.8.9	Mupad [B] (verification not implemented)	350

3.8.1 Optimal result

Integrand size = 32, antiderivative size = 111

$$\int \cot^5(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= a(iA+B)x + \frac{a(iA+B) \cot(c+dx)}{d} + \frac{a(A-iB) \cot^2(c+dx)}{2d}$$

$$- \frac{a(iA+B) \cot^3(c+dx)}{3d} - \frac{aA \cot^4(c+dx)}{4d} + \frac{a(A-iB) \log(\sin(c+dx))}{d}$$

```
output a*(I*A+B)*x+a*(I*A+B)*cot(d*x+c)/d+1/2*a*(A-I*B)*cot(d*x+c)^2/d-1/3*a*(I*A+B)*cot(d*x+c)^3/d-1/4*a*A*cot(d*x+c)^4/d+a*(A-I*B)*ln(sin(d*x+c))/d
```

3.8.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.91 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.86

$$\int \cot^5(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx =$$

$$\frac{a(-6(A-iB) \cot^2(c+dx) + 3A \cot^4(c+dx) + 4(iA+B) \cot^3(c+dx) \text{Hypergeometric2F1}(-\frac{3}{2}, 1, -12d$$

input `Integrate[Cot[c + d*x]^5*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `-1/12*(a*(-6*(A - I*B)*Cot[c + d*x]^2 + 3*A*Cot[c + d*x]^4 + 4*(I*A + B)*Cot[c + d*x]^3*Hypergeometric2F1[-3/2, 1, -1/2, -Tan[c + d*x]^2] - 12*(A - I*B)*(Log[Cos[c + d*x]] + Log[Tan[c + d*x]])))/d`

3.8.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.02, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.469$, Rules used = {3042, 4074, 3042, 4012, 25, 3042, 4012, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^5(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan(c + dx)^5} dx \\
 & \quad \downarrow 4074 \\
 & -\frac{aA \cot^4(c + dx)}{4d} + \int \cot^4(c + dx)(a(iA + B) - a(A - iB) \tan(c + dx)) dx \\
 & \quad \downarrow 3042 \\
 & -\frac{aA \cot^4(c + dx)}{4d} + \int \frac{a(iA + B) - a(A - iB) \tan(c + dx)}{\tan(c + dx)^4} dx \\
 & \quad \downarrow 4012 \\
 & \int -\cot^3(c + dx)(a(A - iB) + a(iA + B) \tan(c + dx)) dx - \frac{a(B + iA) \cot^3(c + dx)}{3d} - \frac{aA \cot^4(c + dx)}{4d} \\
 & \quad \downarrow 25 \\
 & -\int \cot^3(c + dx)(a(A - iB) + a(iA + B) \tan(c + dx)) dx - \frac{a(B + iA) \cot^3(c + dx)}{3d} - \frac{aA \cot^4(c + dx)}{4d} \\
 & \quad \downarrow 3042 \\
 & -\int \frac{a(A - iB) + a(iA + B) \tan(c + dx)}{\tan(c + dx)^3} dx - \frac{a(B + iA) \cot^3(c + dx)}{3d} - \frac{aA \cot^4(c + dx)}{4d}
 \end{aligned}$$

3.8. $\int \cot^5(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \downarrow 4012 \\
& - \int \cot^2(c+dx)(a(iA+B) - a(A-iB)\tan(c+dx))dx - \frac{a(B+iA)\cot^3(c+dx)}{3d} + \\
& \quad \frac{a(A-iB)\cot^2(c+dx)}{2d} - \frac{aA\cot^4(c+dx)}{4d} \\
& \downarrow 3042 \\
& - \int \frac{a(iA+B) - a(A-iB)\tan(c+dx)}{\tan(c+dx)^2} dx - \frac{a(B+iA)\cot^3(c+dx)}{3d} + \frac{a(A-iB)\cot^2(c+dx)}{2d} - \\
& \quad \frac{aA\cot^4(c+dx)}{4d} \\
& \downarrow 4012 \\
& - \int -\cot(c+dx)(a(A-iB) + a(iA+B)\tan(c+dx))dx - \frac{a(B+iA)\cot^3(c+dx)}{3d} + \\
& \quad \frac{a(A-iB)\cot^2(c+dx)}{2d} + \frac{a(B+iA)\cot(c+dx)}{d} - \frac{aA\cot^4(c+dx)}{4d} \\
& \downarrow 25 \\
& \int \cot(c+dx)(a(A-iB) + a(iA+B)\tan(c+dx))dx - \frac{a(B+iA)\cot^3(c+dx)}{3d} + \\
& \quad \frac{a(A-iB)\cot^2(c+dx)}{2d} + \frac{a(B+iA)\cot(c+dx)}{d} - \frac{aA\cot^4(c+dx)}{4d} \\
& \downarrow 3042 \\
& \int \frac{a(A-iB) + a(iA+B)\tan(c+dx)}{\tan(c+dx)} dx - \frac{a(B+iA)\cot^3(c+dx)}{3d} + \frac{a(A-iB)\cot^2(c+dx)}{2d} + \\
& \quad \frac{a(B+iA)\cot(c+dx)}{d} - \frac{aA\cot^4(c+dx)}{4d} \\
& \downarrow 4014 \\
& a(A-iB) \int \cot(c+dx)dx - \frac{a(B+iA)\cot^3(c+dx)}{3d} + \frac{a(A-iB)\cot^2(c+dx)}{2d} + \\
& \quad \frac{a(B+iA)\cot(c+dx)}{d} + ax(B+iA) - \frac{aA\cot^4(c+dx)}{4d} \\
& \downarrow 3042 \\
& a(A-iB) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx - \frac{a(B+iA)\cot^3(c+dx)}{3d} + \frac{a(A-iB)\cot^2(c+dx)}{2d} + \\
& \quad \frac{a(B+iA)\cot(c+dx)}{d} + ax(B+iA) - \frac{aA\cot^4(c+dx)}{4d} \\
& \downarrow 25
\end{aligned}$$

$$\begin{aligned}
 & -a(A - iB) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - \frac{a(B + iA) \cot^3(c + dx)}{3d} + \frac{a(A - iB) \cot^2(c + dx)}{2d} + \\
 & \quad \frac{a(B + iA) \cot(c + dx)}{d} + ax(B + iA) - \frac{aA \cot^4(c + dx)}{4d} \\
 & \quad \downarrow \text{3956} \\
 & -\frac{a(B + iA) \cot^3(c + dx)}{3d} + \frac{a(A - iB) \cot^2(c + dx)}{2d} + \frac{a(B + iA) \cot(c + dx)}{d} + \\
 & \quad \frac{a(A - iB) \log(-\sin(c + dx))}{d} + ax(B + iA) - \frac{aA \cot^4(c + dx)}{4d}
 \end{aligned}$$

input `Int[Cot[c + d*x]^5*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `a*(I*A + B)*x + (a*(I*A + B)*Cot[c + d*x])/d + (a*(A - I*B)*Cot[c + d*x]^2)/(2*d) - (a*(I*A + B)*Cot[c + d*x]^3)/(3*d) - (a*A*Cot[c + d*x]^4)/(4*d) + (a*(A - I*B)*Log[-Sin[c + d*x]])/d`

3.8.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

3.8.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.95

method	result
parallelrisch	$\frac{a \left(\left(-\frac{A}{2} + \frac{iB}{2} \right) \ln(\sec^2(dx+c)) + (-iB+A) \ln(\tan(dx+c)) - \frac{A(\cot^4(dx+c))}{4} + (\cot^3(dx+c)) \left(-\frac{iA}{3} - \frac{B}{3} \right) + (\cot^2(dx+c)) \left(\frac{A}{3} + \frac{iB}{3} \right) \right)}{d}$
derivativedivides	$\frac{a \left(\frac{(iB-A) \ln(1+\tan^2(dx+c))}{2} + (iA+B) \arctan(\tan(dx+c)) - \frac{A}{4 \tan(dx+c)^4} - \frac{iB-A}{2 \tan(dx+c)^2} + (-iB+A) \ln(\tan(dx+c)) - \frac{-iA}{\tan(dx+c)} \right)}{d}$
default	$\frac{a \left(\frac{(iB-A) \ln(1+\tan^2(dx+c))}{2} + (iA+B) \arctan(\tan(dx+c)) - \frac{A}{4 \tan(dx+c)^4} - \frac{iB-A}{2 \tan(dx+c)^2} + (-iB+A) \ln(\tan(dx+c)) - \frac{-iA}{\tan(dx+c)} \right)}{d}$
norman	$\frac{\frac{(iA+Ba) \tan^3(dx+c)}{d} + (iA+Ba)x \tan^4(dx+c) - \frac{aA}{4d} + \frac{(-iA+Ba) \tan^2(dx+c)}{2d} - \frac{(iA+Ba) \tan(dx+c)}{3d}}{\tan(dx+c)^4} + \frac{(-iA+Ba)}{3d}$
risch	$-\frac{2aBc}{d} - \frac{2iaAc}{d} + \frac{2ia(12iA e^{6i(dx+c)} + 9B e^{6i(dx+c)} - 18iA e^{4i(dx+c)} - 18B e^{4i(dx+c)} + 16iA e^{2i(dx+c)} + 13B e^{2i(dx+c)} + 13iA)}{3d(e^{2i(dx+c)} - 1)^4}$

```
input int(cot(d*x+c)^5*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

```
output a*((-1/2*A+1/2*I*B)*ln(sec(d*x+c)^2)+(A-I*B)*ln(tan(d*x+c))-1/4*A*cot(d*x+
c)^4+cot(d*x+c)^3*(-1/3*I*A-1/3*B)+cot(d*x+c)^2*(1/2*A-1/2*I*B)+cot(d*x+c)
*(I*A+B)+(I*A+B)*x*d)/d
```

3.8. $\int \cot^5(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

3.8.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 206 vs. 2(95) = 190.

Time = 0.25 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.86

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{6(4A - 3iB)ae^{(6idx+6ic)} - 36(A - iB)ae^{(4idx+4ic)} + 2(16A - 13iB)ae^{(2idx+2ic)} - 8(A - iB)a - 3}{3(de^{(8idx+8ic)} - 4)}$$

input `integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-1/3*(6*(4*A - 3*I*B)*a*e^(6*I*d*x + 6*I*c) - 36*(A - I*B)*a*e^(4*I*d*x + 4*I*c) + 2*(16*A - 13*I*B)*a*e^(2*I*d*x + 2*I*c) - 8*(A - I*B)*a - 3*((A - I*B)*a*e^(8*I*d*x + 8*I*c) - 4*(A - I*B)*a*e^(6*I*d*x + 6*I*c) + 6*(A - I*B)*a*e^(4*I*d*x + 4*I*c) - 4*(A - I*B)*a*e^(2*I*d*x + 2*I*c) + (A - I*B)*a)*log(e^(2*I*d*x + 2*I*c) - 1))/(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)`

3.8.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 218 vs. 2(92) = 184.

Time = 0.72 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.96

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{a(A - iB) \log(e^{2idx} - e^{-2ic})}{d} + \frac{8Aa - 8iBa + (-32Aae^{2ic} + 26iBae^{2ic})e^{2idx} + (36Aae^{4ic} - 36iBae^{4ic})e^{4idx} + (-24Aae^{6ic} + 18iBae^{6ic})e^{6idx}}{3de^{8ic}e^{8idx} - 12de^{6ic}e^{6idx} + 18de^{4ic}e^{4idx} - 12de^{2ic}e^{2idx} + 3d}$$

input `integrate(cot(d*x+c)**5*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `a*(A - I*B)*log(exp(2*I*d*x) - exp(-2*I*c))/d + (8*A*a - 8*I*B*a + (-32*A*a*exp(2*I*c) + 26*I*B*a*exp(2*I*c))*exp(2*I*d*x) + (36*A*a*exp(4*I*c) - 36*I*B*a*exp(4*I*c))*exp(4*I*d*x) + (-24*A*a*exp(6*I*c) + 18*I*B*a*exp(6*I*c))*exp(6*I*d*x))/(3*d*exp(8*I*c)*exp(8*I*d*x) - 12*d*exp(6*I*c)*exp(6*I*d*x) + 18*d*exp(4*I*c)*exp(4*I*d*x) - 12*d*exp(2*I*c)*exp(2*I*d*x) + 3*d)`

3.8. $\int \cot^5(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

3.8.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.05

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{12(dx + c)(iA + B)a - 6(A - iB)a \log(\tan(dx + c)^2 + 1) + 12(A - iB)a \log(\tan(dx + c)) - \frac{12(-iA - B)a \tan(dx + c)^3 - 6(A - iB)a \tan(dx + c)^2 + 4(IA + B)a \tan(dx + c) + 3Aa}{\tan(dx + c)^4}}{12d}$$

input `integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/12*(12*(d*x + c)*(I*A + B)*a - 6*(A - I*B)*a*log(tan(d*x + c)^2 + 1) + 12*(A - I*B)*a*log(tan(d*x + c)) - (12*(-I*A - B)*a*tan(d*x + c)^3 - 6*(A - I*B)*a*tan(d*x + c)^2 + 4*(I*A + B)*a*tan(d*x + c) + 3*A*a)/tan(d*x + c)^4)/d`

3.8.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 282 vs. 2(95) = 190.

Time = 0.85 (sec) , antiderivative size = 282, normalized size of antiderivative = 2.54

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx =$$

$$\frac{3Aa \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^4 - 8iAa \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 8Ba \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 36Aa \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + 24iAa \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 3Aa}{12d}$$

input `integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `-1/192*(3*A*a*tan(1/2*d*x + 1/2*c)^4 - 8*I*A*a*tan(1/2*d*x + 1/2*c)^3 - 8*B*a*tan(1/2*d*x + 1/2*c)^3 - 36*A*a*tan(1/2*d*x + 1/2*c)^2 + 24*I*B*a*tan(1/2*d*x + 1/2*c)^2 + 120*I*A*a*tan(1/2*d*x + 1/2*c) + 120*B*a*tan(1/2*d*x + 1/2*c) + 384*(A*a - I*B*a)*log(tan(1/2*d*x + 1/2*c) + I) - 192*(A*a - I*B*a)*log(tan(1/2*d*x + 1/2*c)) + (400*A*a*tan(1/2*d*x + 1/2*c)^4 - 400*I*B*a*tan(1/2*d*x + 1/2*c)^4 - 120*I*A*a*tan(1/2*d*x + 1/2*c)^3 - 120*B*a*tan(1/2*d*x + 1/2*c)^3 - 36*A*a*tan(1/2*d*x + 1/2*c)^2 + 24*I*B*a*tan(1/2*d*x + 1/2*c)^2 + 8*I*A*a*tan(1/2*d*x + 1/2*c) + 8*B*a*tan(1/2*d*x + 1/2*c) + 3*A*a)/tan(1/2*d*x + 1/2*c)^4)/d`

3.8. $\int \cot^5(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

3.8.9 Mupad [B] (verification not implemented)

Time = 8.46 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.90

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx =$$

$$\frac{(-Ba - Aa1i) \tan(c + dx)^3 + \left(-\frac{Aa}{2} + \frac{Ba1i}{2}\right) \tan(c + dx)^2 + \left(\frac{Ba}{3} + \frac{Aa1i}{3}\right) \tan(c + dx) + \frac{Aa}{4}}{d \tan(c + dx)^4}$$

$$+ \frac{a \operatorname{atan}(2 \tan(c + dx) + 1i) (A - B1i) 2i}{d}$$

input `int(cot(c + d*x)^5*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`

output `(a*atan(2*tan(c + d*x) + 1i)*(A - B*1i)*2i)/d - ((A*a)/4 + tan(c + d*x)*((A*a*1i)/3 + (B*a)/3) - tan(c + d*x)^3*(A*a*1i + B*a) - tan(c + d*x)^2*((A*a)/2 - (B*a*1i)/2))/(d*tan(c + d*x)^4)`

3.9 $\int \tan^2(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.9.1	Optimal result	351
3.9.2	Mathematica [A] (verified)	352
3.9.3	Rubi [A] (verified)	352
3.9.4	Maple [A] (verified)	355
3.9.5	Fricas [A] (verification not implemented)	356
3.9.6	Sympy [A] (verification not implemented)	356
3.9.7	Maxima [A] (verification not implemented)	357
3.9.8	Giac [B] (verification not implemented)	357
3.9.9	Mupad [B] (verification not implemented)	358

3.9.1 Optimal result

Integrand size = 34, antiderivative size = 141

$$\begin{aligned} & \int \tan^2(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx \\ &= -2a^2(A-iB)x + \frac{2a^2(iA+B) \log(\cos(c+dx))}{d} \\ & \quad + \frac{2a^2(A-iB) \tan(c+dx)}{d} + \frac{a^2(iA+B) \tan^2(c+dx)}{d} \\ & \quad - \frac{a^2(4A-5iB) \tan^3(c+dx)}{12d} + \frac{iB \tan^3(c+dx) (a^2+ia^2 \tan(c+dx))}{4d} \end{aligned}$$

output

```
-2*a^2*(A-I*B)*x+2*a^2*(I*A+B)*ln(cos(d*x+c))/d+2*a^2*(A-I*B)*tan(d*x+c)/d
+a^2*(I*A+B)*tan(d*x+c)^2/d-1/12*a^2*(4*A-5*I*B)*tan(d*x+c)^3/d+1/4*I*B*ta
n(d*x+c)^3*(a^2+I*a^2*tan(d*x+c))/d
```

3.9.2 Mathematica [A] (verified)

Time = 1.36 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.72

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{a^2(-4iA - 5B - 24i(A - iB) \log(i + \tan(c + dx)) + 24(A - iB) \tan(c + dx) + 12(iA + B) \tan^2(c + dx))}{12d}$$

input `Integrate[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(a^2*((-4*I)*A - 5*B - (24*I)*(A - I*B)*Log[I + Tan[c + d*x]] + 24*(A - I*B)*Tan[c + d*x] + 12*(I*A + B)*Tan[c + d*x]^2 - 4*(A - (2*I)*B)*Tan[c + d*x]^3 - 3*B*Tan[c + d*x]^4))/(12*d)`

3.9.3 Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.04, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3042, 4077, 3042, 4075, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \tan(c + dx)^2(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$\downarrow 4077$$

$$\frac{1}{4} \int \tan^2(c + dx)(i \tan(c + dx)a + a)(a(4A - 3iB) + a(4iA + 5B) \tan(c + dx)) dx + \frac{iB \tan^3(c + dx)(a^2 + ia^2 \tan(c + dx))}{4d}$$

$$\downarrow 3042$$

$$\frac{1}{4} \int \tan(c + dx)^2(i \tan(c + dx)a + a)(a(4A - 3iB) + a(4iA + 5B) \tan(c + dx)) dx + \frac{iB \tan^3(c + dx)(a^2 + ia^2 \tan(c + dx))}{4d}$$

3.9. $\int \tan^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\begin{array}{c} \downarrow 4075 \\ \frac{1}{4} \left(\int \tan^2(c+dx) (8(A-iB)a^2 + 8(iA+B)\tan(c+dx)a^2) dx - \frac{a^2(4A-5iB)\tan^3(c+dx)}{3d} \right) + \\ \frac{iB \tan^3(c+dx) (a^2 + ia^2 \tan(c+dx))}{4d} \end{array}$$

$$\begin{array}{c} \downarrow 3042 \\ \frac{1}{4} \left(\int \tan(c+dx)^2 (8(A-iB)a^2 + 8(iA+B)\tan(c+dx)a^2) dx - \frac{a^2(4A-5iB)\tan^3(c+dx)}{3d} \right) + \\ \frac{iB \tan^3(c+dx) (a^2 + ia^2 \tan(c+dx))}{4d} \end{array}$$

$$\begin{array}{c} \downarrow 4011 \\ \frac{1}{4} \left(\int \tan(c+dx) (8a^2(A-iB)\tan(c+dx) - 8a^2(iA+B)) dx - \frac{a^2(4A-5iB)\tan^3(c+dx)}{3d} + \frac{4a^2(B+iA)\tan^2(c+dx)}{d} \right) + \\ \frac{iB \tan^3(c+dx) (a^2 + ia^2 \tan(c+dx))}{4d} \end{array}$$

$$\begin{array}{c} \downarrow 3042 \\ \frac{1}{4} \left(\int \tan(c+dx) (8a^2(A-iB)\tan(c+dx) - 8a^2(iA+B)) dx - \frac{a^2(4A-5iB)\tan^3(c+dx)}{3d} + \frac{4a^2(B+iA)\tan^2(c+dx)}{d} \right) + \\ \frac{iB \tan^3(c+dx) (a^2 + ia^2 \tan(c+dx))}{4d} \end{array}$$

$$\begin{array}{c} \downarrow 4008 \\ \frac{1}{4} \left(-8a^2(B+iA) \int \tan(c+dx) dx - \frac{a^2(4A-5iB)\tan^3(c+dx)}{3d} + \frac{4a^2(B+iA)\tan^2(c+dx)}{d} + \frac{8a^2(A-iB)\tan^2(c+dx)}{d} \right) + \\ \frac{iB \tan^3(c+dx) (a^2 + ia^2 \tan(c+dx))}{4d} \end{array}$$

$$\begin{array}{c} \downarrow 3042 \\ \frac{1}{4} \left(-8a^2(B+iA) \int \tan(c+dx) dx - \frac{a^2(4A-5iB)\tan^3(c+dx)}{3d} + \frac{4a^2(B+iA)\tan^2(c+dx)}{d} + \frac{8a^2(A-iB)\tan^2(c+dx)}{d} \right) + \\ \frac{iB \tan^3(c+dx) (a^2 + ia^2 \tan(c+dx))}{4d} \end{array}$$

$$\begin{array}{c} \downarrow 3956 \\ \frac{1}{4} \left(-8a^2(B+iA) \int \tan(c+dx) dx - \frac{a^2(4A-5iB)\tan^3(c+dx)}{3d} + \frac{4a^2(B+iA)\tan^2(c+dx)}{d} + \frac{8a^2(A-iB)\tan^2(c+dx)}{d} \right) + \\ \frac{iB \tan^3(c+dx) (a^2 + ia^2 \tan(c+dx))}{4d} \end{array}$$

$$\frac{1}{4} \left(-\frac{a^2(4A - 5iB) \tan^3(c + dx)}{3d} + \frac{4a^2(B + iA) \tan^2(c + dx)}{d} + \frac{8a^2(A - iB) \tan(c + dx)}{d} + \frac{8a^2(B + iA) \log(\cos(c + dx))}{d} \right) + \frac{iB \tan^3(c + dx) (a^2 + ia^2 \tan(c + dx))}{4d}$$

input `Int[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `((I/4)*B*Tan[c + d*x]^3*(a^2 + I*a^2*Tan[c + d*x]))/d + (-8*a^2*(A - I*B)*x + (8*a^2*(I*A + B)*Log[Cos[c + d*x]])/d + (8*a^2*(A - I*B)*Tan[c + d*x])/d + (4*a^2*(I*A + B)*Tan[c + d*x]^2)/d - (a^2*(4*A - (5*I)*B)*Tan[c + d*x]^3)/(3*d))/4`

3.9.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.9. $\int \tan^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

```
rule 4077 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan
[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c -
a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] &&
GtQ[m, 1] && !LtQ[n, -1]
```

3.9.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.86

method	result
derivativedivides	$\frac{a^2 \left(\frac{2iB(\tan^3(dx+c))}{3} - \frac{B(\tan^4(dx+c))}{4} + iA(\tan^2(dx+c)) - \frac{A(\tan^3(dx+c))}{3} - 2iB \tan(dx+c) + B(\tan^2(dx+c)) + 2A \tan(dx+c) \right)}{d}$
default	$\frac{a^2 \left(\frac{2iB(\tan^3(dx+c))}{3} - \frac{B(\tan^4(dx+c))}{4} + iA(\tan^2(dx+c)) - \frac{A(\tan^3(dx+c))}{3} - 2iB \tan(dx+c) + B(\tan^2(dx+c)) + 2A \tan(dx+c) \right)}{d}$
norman	$(2iB a^2 - 2A a^2) x + \frac{(iA a^2 + B a^2)(\tan^2(dx+c))}{d} - \frac{(-2iB a^2 + A a^2)(\tan^3(dx+c))}{3d} + \frac{2(-iB a^2 + A a^2) \tan(dx+c)}{d}$
parts	$\frac{(2iA a^2 + B a^2) \left(\frac{(\tan^2(dx+c))}{2} - \frac{\ln(1+\tan^2(dx+c))}{2} \right)}{d} + \frac{(2iB a^2 - A a^2) \left(\frac{(\tan^3(dx+c))}{3} - \tan(dx+c) + \arctan(\tan(dx+c)) \right)}{d}$
parallelrisch	$-\frac{8iB(\tan^3(dx+c))a^2 + 3B a^2(\tan^4(dx+c)) - 12iA(\tan^2(dx+c))a^2 + 4A(\tan^3(dx+c))a^2 - 24iB x a^2 d + 12iA \ln(1+\tan^2(dx+c))a^2}{12d}$
risch	$-\frac{4ia^2Bc}{d} + \frac{4a^2Ac}{d} + \frac{2a^2(15iA e^{6i(dx+c)} + 21B e^{6i(dx+c)} + 33iA e^{4i(dx+c)} + 36B e^{4i(dx+c)} + 25iA e^{2i(dx+c)} + 29B e^{2i(dx+c)} + 1)}{3d(e^{2i(dx+c)} + 1)^4}$

```
input int(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)), x, method=_RETURNVER
BOSE)
```

```
output 1/d*a^2*(2/3*I*B*tan(d*x+c)^3-1/4*B*tan(d*x+c)^4+I*A*tan(d*x+c)^2-1/3*A*ta
n(d*x+c)^3-2*I*B*tan(d*x+c)+B*tan(d*x+c)^2+2*A*tan(d*x+c)+1/2*(-2*B-2*I*A)
*ln(1+tan(d*x+c)^2)+(-2*A+2*I*B)*arctan(tan(d*x+c)))
```

3.9. $\int \tan^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.9.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.67

$$\int \tan^2(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx = \frac{2(3(-5iA-7B)a^2e^{(6i dx+6i c)} + 3(-11iA-12B)a^2e^{(4i dx+4i c)} + (-25iA-29B)a^2e^{(2i dx+2i c)} + (-7iA-8B)a^2e^{(0i dx+0i c)})}{3}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-2/3*(3*(-5*I*A - 7*B)*a^2*e^(6*I*d*x + 6*I*c) + 3*(-11*I*A - 12*B)*a^2*e^(4*I*d*x + 4*I*c) + (-25*I*A - 29*B)*a^2*e^(2*I*d*x + 2*I*c) + (-7*I*A - 8*B)*a^2 + 3*((-I*A - B)*a^2*e^(8*I*d*x + 8*I*c) + 4*(-I*A - B)*a^2*e^(6*I*d*x + 6*I*c) + 6*(-I*A - B)*a^2*e^(4*I*d*x + 4*I*c) + 4*(-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*log(e^(2*I*d*x + 2*I*c) + 1)/(d*e^(8*I*d*x + 8*I*c) + 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*x + 2*I*c) + d)`

3.9.6 Sympy [A] (verification not implemented)

Time = 0.46 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.67

$$\int \tan^2(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx = \frac{2ia^2(A-iB) \log(e^{2idx} + e^{-2ic})}{d} + \frac{14iAa^2 + 16Ba^2 + (50iAa^2e^{2ic} + 58Ba^2e^{2ic})e^{2idx} + (66iAa^2e^{4ic} + 72Ba^2e^{4ic})e^{4idx} + (30iAa^2e^{6ic} + 42Ba^2e^{6ic})e^{6idx}}{3de^{8ic}e^{8idx} + 12de^{6ic}e^{6idx} + 18de^{4ic}e^{4idx} + 12de^{2ic}e^{2idx} + 3d}$$

input `integrate(tan(d*x+c)**2*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `2*I*a**2*(A - I*B)*log(exp(2*I*d*x) + exp(-2*I*c))/d + (14*I*A*a**2 + 16*B*a**2 + (50*I*A*a**2*exp(2*I*c) + 58*B*a**2*exp(2*I*c))*exp(2*I*d*x) + (66*I*A*a**2*exp(4*I*c) + 72*B*a**2*exp(4*I*c))*exp(4*I*d*x) + (30*I*A*a**2*exp(6*I*c) + 42*B*a**2*exp(6*I*c))*exp(6*I*d*x))/(3*d*exp(8*I*c)*exp(8*I*d*x) + 12*d*exp(6*I*c)*exp(6*I*d*x) + 18*d*exp(4*I*c)*exp(4*I*d*x) + 12*d*exp(2*I*c)*exp(2*I*d*x) + 3*d)`

3.9.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.79

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{3Ba^2 \tan(dx + c)^4 + 4(A - 2iB)a^2 \tan(dx + c)^3 + 12(-iA - B)a^2 \tan(dx + c)^2 + 24(dx + c)(A - B)a^2 \tan(dx + c) + 24(dx + c)^2(A - B)a^2}{12d}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/12*(3*B*a^2*tan(d*x + c)^4 + 4*(A - 2*I*B)*a^2*tan(d*x + c)^3 + 12*(-I*A - B)*a^2*tan(d*x + c)^2 + 24*(d*x + c)*(A - I*B)*a^2 - 12*(-I*A - B)*a^2*log(tan(d*x + c)^2 + 1) - 24*(A - I*B)*a^2*tan(d*x + c))/d`

3.9.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 408 vs. 2(123) = 246.

Time = 0.59 (sec) , antiderivative size = 408, normalized size of antiderivative = 2.89

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{2(-3iAa^2e^{(8i dx+8i c)} \log(e^{(2i dx+2i c)} + 1) - 3Ba^2e^{(8i dx+8i c)} \log(e^{(2i dx+2i c)} + 1) - 12iAa^2e^{(6i dx+6i c)} \log(e^{(2i dx+2i c)} + 1) + 24(dx + c)(A - B)a^2 \tan(dx + c) + 24(dx + c)^2(A - B)a^2}{12d}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

$$\begin{aligned}
& -2/3*(-3*I*A*a^2*e^{(8*I*d*x + 8*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 3*B*a^2*e^{(8*I*d*x + 8*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 12*I*A*a^2*e^{(6*I*d*x + 6*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 12*B*a^2*e^{(6*I*d*x + 6*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 18*I*A*a^2*e^{(4*I*d*x + 4*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 18*B*a^2*e^{(4*I*d*x + 4*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 12*I*A*a^2*e^{(2*I*d*x + 2*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 12*B*a^2*e^{(2*I*d*x + 2*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 15*I*A*a^2*e^{(6*I*d*x + 6*I*c)} - 21*B*a^2*e^{(6*I*d*x + 6*I*c)} - 33*I*A*a^2*e^{(4*I*d*x + 4*I*c)} - 36*B*a^2*e^{(4*I*d*x + 4*I*c)} - 25*I*A*a^2*e^{(2*I*d*x + 2*I*c)} - 29*B*a^2*e^{(2*I*d*x + 2*I*c)} - 3*I*A*a^2*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 3*B*a^2*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 7*I*A*a^2 - 8*B*a^2)/(d*e^{(8*I*d*x + 8*I*c)} + 4*d*e^{(6*I*d*x + 6*I*c)} + 6*d*e^{(4*I*d*x + 4*I*c)} + 4*d*e^{(2*I*d*x + 2*I*c)} + d)
\end{aligned}$$

3.9.9 Mupad [B] (verification not implemented)

Time = 7.25 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.09

$$\begin{aligned}
& \int \tan^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
& = \frac{\tan(c + dx)^3 \left(\frac{a^2(B+A \operatorname{li}) \operatorname{li}}{3} + \frac{B a^2 \operatorname{li}}{3} \right)}{d} - \frac{\tan(c + dx) (-A a^2 + a^2 (B + A \operatorname{li}) \operatorname{li} + B a^2 \operatorname{li})}{d} \\
& + \frac{\tan(c + dx)^2 \left(\frac{a^2(B+A \operatorname{li})}{2} + \frac{B a^2}{2} + \frac{A a^2 \operatorname{li}}{2} \right)}{d} \\
& - \frac{\ln(\tan(c + dx) + \operatorname{li}) (2 B a^2 + A a^2 2i)}{d} - \frac{B a^2 \tan(c + dx)^4}{4 d}
\end{aligned}$$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2,x)`

output

$$\begin{aligned}
& (\tan(c + d*x)^3*((a^2*(A*1i + B)*1i)/3 + (B*a^2*1i)/3))/d - (\tan(c + d*x)* \\
& (a^2*(A*1i + B)*1i - A*a^2 + B*a^2*1i))/d + (\tan(c + d*x)^2*((A*a^2*1i)/2 \\
& + (a^2*(A*1i + B))/2 + (B*a^2)/2))/d - (\log(\tan(c + d*x) + 1i)*(A*a^2*2i + \\
& 2*B*a^2))/d - (B*a^2*tan(c + d*x)^4)/(4*d)
\end{aligned}$$

3.10 $\int \tan(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.10.1	Optimal result	359
3.10.2	Mathematica [A] (verified)	359
3.10.3	Rubi [A] (verified)	360
3.10.4	Maple [A] (verified)	362
3.10.5	Fricas [A] (verification not implemented)	363
3.10.6	Sympy [B] (verification not implemented)	363
3.10.7	Maxima [A] (verification not implemented)	364
3.10.8	Giac [B] (verification not implemented)	364
3.10.9	Mupad [B] (verification not implemented)	365

3.10.1 Optimal result

Integrand size = 32, antiderivative size = 107

$$\int \tan(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= -2a^2(iA+B)x - \frac{2a^2(A-iB) \log(\cos(c+dx))}{d} + \frac{a^2(iA+B) \tan(c+dx)}{d}$$

$$+ \frac{A(a+ia \tan(c+dx))^2}{2d} - \frac{iB(a+ia \tan(c+dx))^3}{3ad}$$

```
output -2*a^2*(I*A+B)*x-2*a^2*(A-I*B)*ln(cos(d*x+c))/d+a^2*(I*A+B)*tan(d*x+c)/d+1
/2*A*(a+I*a*tan(d*x+c))^2/d-1/3*I*B*(a+I*a*tan(d*x+c))^3/a/d
```

3.10.2 Mathematica [A] (verified)

Time = 0.82 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.77

$$\int \tan(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{a^2(3A-2iB+12(A-iB) \log(i+\tan(c+dx))+12(iA+B) \tan(c+dx)-3(A-2iB) \tan^2(c+dx)-6d)}{6d}$$

```
input Integrate[Tan[c + d*x]*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]
```

output $(a^2(3A - (2I)B + 12(A - I)B)\text{Log}[I + \text{Tan}[c + d*x]] + 12(I*A + B)*\text{Tan}[c + d*x] - 3(A - (2I)B)*\text{Tan}[c + d*x]^2 - 2*B*\text{Tan}[c + d*x]^3)/(6*d)$

3.10.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.93, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4075, 3042, 4010, 3042, 3958, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4075} \\
 & \int (i \tan(c + dx)a + a)^2(A \tan(c + dx) - B) dx - \frac{iB(a + ia \tan(c + dx))^3}{3ad} \\
 & \quad \downarrow \text{3042} \\
 & \int (i \tan(c + dx)a + a)^2(A \tan(c + dx) - B) dx - \frac{iB(a + ia \tan(c + dx))^3}{3ad} \\
 & \quad \downarrow \text{4010} \\
 & -(B + iA) \int (i \tan(c + dx)a + a)^2 dx + \frac{A(a + ia \tan(c + dx))^2}{2d} - \frac{iB(a + ia \tan(c + dx))^3}{3ad} \\
 & \quad \downarrow \text{3042} \\
 & -(B + iA) \int (i \tan(c + dx)a + a)^2 dx + \frac{A(a + ia \tan(c + dx))^2}{2d} - \frac{iB(a + ia \tan(c + dx))^3}{3ad} \\
 & \quad \downarrow \text{3958} \\
 & -(B + iA) \left(2ia^2 \int \tan(c + dx) dx - \frac{a^2 \tan(c + dx)}{d} + 2a^2 x \right) + \frac{A(a + ia \tan(c + dx))^2}{2d} - \\
 & \quad \frac{iB(a + ia \tan(c + dx))^3}{3ad} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.10. $\int \tan(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& -(B + iA) \left(2ia^2 \int \tan(c + dx) dx - \frac{a^2 \tan(c + dx)}{d} + 2a^2 x \right) + \frac{A(a + ia \tan(c + dx))^2}{2d} - \\
& \quad \frac{iB(a + ia \tan(c + dx))^3}{3ad} \\
& \quad \downarrow \text{3956} \\
& -(B + iA) \left(-\frac{a^2 \tan(c + dx)}{d} - \frac{2ia^2 \log(\cos(c + dx))}{d} + 2a^2 x \right) + \frac{A(a + ia \tan(c + dx))^2}{2d} - \\
& \quad \frac{iB(a + ia \tan(c + dx))^3}{3ad}
\end{aligned}$$

input `Int[Tan[c + d*x]*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(A*(a + I*a*Tan[c + d*x])^2)/(2*d) - ((I/3)*B*(a + I*a*Tan[c + d*x])^3)/(a*d) - (I*A + B)*(2*a^2*x - ((2*I)*a^2*Log[Cos[c + d*x]])/d - (a^2*Tan[c + d*x])/d)`

3.10.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 3958 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^2, x_Symbol] := Simp[(a^2 - b^2)*x, x] + (Simp[b^2*(Tan[c + d*x]/d), x] + Simp[2*a*b Int[Tan[c + d*x], x], x]) /; FreeQ[{a, b, c, d}, x]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`


```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

3.10.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.93

method	result
derivativedivides	$\frac{a^2 \left(iB(\tan^2(dx+c)) - \frac{B(\tan^3(dx+c))}{3} + 2iA \tan(dx+c) - \frac{A(\tan^2(dx+c))}{2} + 2B \tan(dx+c) + \frac{(-2iB+2A) \ln(1+\tan^2(dx+c))}{2} \right)}{d}$
default	$\frac{a^2 \left(iB(\tan^2(dx+c)) - \frac{B(\tan^3(dx+c))}{3} + 2iA \tan(dx+c) - \frac{A(\tan^2(dx+c))}{2} + 2B \tan(dx+c) + \frac{(-2iB+2A) \ln(1+\tan^2(dx+c))}{2} \right)}{d}$
norman	$\frac{(-2iA a^2 - 2B a^2) x - \frac{(-2iB a^2 + A a^2)(\tan^2(dx+c))}{2d} + \frac{2(iA a^2 + B a^2) \tan(dx+c)}{d} - \frac{B a^2(\tan^3(dx+c))}{3d}}{6d}$
parallelrisch	$\frac{12iA x a^2 d - 6iB(\tan^2(dx+c)) a^2 + 2B a^2(\tan^3(dx+c)) - 12iA \tan(dx+c) a^2 + 3A(\tan^2(dx+c)) a^2 + 6iB \ln(1+\tan^2(dx+c))}{6d}$
parts	$\frac{(2iA a^2 + B a^2)(\tan(dx+c) - \arctan(\tan(dx+c)))}{d} + \frac{(2iB a^2 - A a^2) \left(\frac{\tan^2(dx+c)}{2} - \frac{\ln(1+\tan^2(dx+c))}{2} \right)}{d} + \frac{A \ln(1+\tan^2(dx+c))}{d}$
risch	$\frac{4a^2 Bc}{d} + \frac{4ia^2 Ac}{d} + \frac{2ia^2 (9iA e^{4i(dx+c)} + 15B e^{4i(dx+c)} + 15iA e^{2i(dx+c)} + 18B e^{2i(dx+c)} + 6iA + 7B)}{3d(e^{2i(dx+c)} + 1)^3} + \frac{2ia^2 \ln(e^{2i(dx+c)} + 1)}{d}$

```
input int(tan(d*x+c)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

```
output 1/d*a^2*(I*B*tan(d*x+c)^2-1/3*B*tan(d*x+c)^3+2*I*A*tan(d*x+c)-1/2*A*tan(d*
x+c)^2+2*B*tan(d*x+c)+1/2*(2*A-2*I*B)*ln(1+tan(d*x+c)^2)+(-2*B-2*I*A)*arct
an(tan(d*x+c)))
```

3.10. $\int \tan(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.10.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.64

$$\int \tan(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{2(3(3A - 5iB)a^2e^{(4i dx + 4i c)} + 3(5A - 6iB)a^2e^{(2i dx + 2i c)} + (6A - 7iB)a^2 + 3((A - iB)a^2e^{(6i dx + 6i c)} + 3(de^{(6i dx + 6i c)} + 3de^{(4i dx + 4i c)} +$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-2/3*(3*(3*A - 5*I*B)*a^2*e^(4*I*d*x + 4*I*c) + 3*(5*A - 6*I*B)*a^2*e^(2*I*d*x + 2*I*c) + (6*A - 7*I*B)*a^2 + 3*((A - I*B)*a^2*e^(6*I*d*x + 6*I*c) + 3*(A - I*B)*a^2*e^(4*I*d*x + 4*I*c) + 3*(A - I*B)*a^2*e^(2*I*d*x + 2*I*c) + (A - I*B)*a^2)*log(e^(2*I*d*x + 2*I*c) + 1)/(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)`

3.10.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 178 vs. 2(88) = 176.

Time = 0.35 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.66

$$\int \tan(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = -\frac{2a^2(A - iB) \log(e^{2idx} + e^{-2ic})}{d} + \frac{-12Aa^2 + 14iBa^2 + (-30Aa^2e^{2ic} + 36iBa^2e^{2ic})e^{2idx} + (-18Aa^2e^{4ic} + 30iBa^2e^{4ic})e^{4idx}}{3de^{6ic}e^{6idx} + 9de^{4ic}e^{4idx} + 9de^{2ic}e^{2idx} + 3d}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `-2*a**2*(A - I*B)*log(exp(2*I*d*x) + exp(-2*I*c))/d + (-12*A*a**2 + 14*I*B*a**2 + (-30*A*a**2*exp(2*I*c) + 36*I*B*a**2*exp(2*I*c))*exp(2*I*d*x) + (-18*A*a**2*exp(4*I*c) + 30*I*B*a**2*exp(4*I*c))*exp(4*I*d*x))/(3*d*exp(6*I*c)*exp(6*I*d*x) + 9*d*exp(4*I*c)*exp(4*I*d*x) + 9*d*exp(2*I*c)*exp(2*I*d*x) + 3*d)`

3.10.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.86

$$\int \tan(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{2Ba^2 \tan(dx + c)^3 + 3(A - 2iB)a^2 \tan(dx + c)^2 + 12(dx + c)(iA + B)a^2 - 6(A - iB)a^2 \log(\tan(dx + c) + 1)}{6d}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/6*(2*B*a^2*tan(d*x + c)^3 + 3*(A - 2*I*B)*a^2*tan(d*x + c)^2 + 12*(d*x + c)*(I*A + B)*a^2 - 6*(A - I*B)*a^2*log(tan(d*x + c)^2 + 1) + 12*(-I*A - B)*a^2*tan(d*x + c))/d`

3.10.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 312 vs. 2(91) = 182.

Time = 0.44 (sec) , antiderivative size = 312, normalized size of antiderivative = 2.92

$$\int \tan(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{2(3Aa^2e^{(6i dx+6i c)} \log(e^{(2i dx+2i c)} + 1) - 3iBa^2e^{(6i dx+6i c)} \log(e^{(2i dx+2i c)} + 1) + 9Aa^2e^{(4i dx+4i c)} \log(e^{(2i dx+2i c)} + 1) - 9iBa^2e^{(4i dx+4i c)} \log(e^{(2i dx+2i c)} + 1) + 9Aa^2e^{(2i dx+2i c)} \log(e^{(2i dx+2i c)} + 1) - 9iBa^2e^{(2i dx+2i c)} \log(e^{(2i dx+2i c)} + 1) + 9Aa^2e^{(4i dx+4i c)} - 15iBa^2e^{(4i dx+4i c)} + 15Aa^2e^{(2i dx+2i c)} - 18iBa^2e^{(2i dx+2i c)} + 3Aa^2 \log(e^{(2i dx+2i c)} + 1) - 3iBa^2 \log(e^{(2i dx+2i c)} + 1) + 6Aa^2 - 7iBa^2)/(d * e^{(6i dx+6i c)} + 3d * e^{(4i dx+4i c)} + 3d * e^{(2i dx+2i c)} + d)}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `-2/3*(3*A*a^2*e^(6*I*d*x + 6*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 3*I*B*a^2*e^(6*I*d*x + 6*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 9*A*a^2*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 9*I*B*a^2*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 9*A*a^2*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 9*I*B*a^2*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 9*A*a^2*e^(4*I*d*x + 4*I*c) - 15*I*B*a^2*e^(4*I*d*x + 4*I*c) + 15*A*a^2*e^(2*I*d*x + 2*I*c) - 18*I*B*a^2*e^(2*I*d*x + 2*I*c) + 3*A*a^2*log(e^(2*I*d*x + 2*I*c) + 1) - 3*I*B*a^2*log(e^(2*I*d*x + 2*I*c) + 1) + 6*A*a^2 - 7*I*B*a^2)/(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)`

3.10. $\int \tan(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.10.9 Mupad [B] (verification not implemented)

Time = 7.65 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.04

$$\int \tan(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{\tan(c + dx)^2 \left(\frac{a^2 (B + A i) i}{2} + \frac{B a^2 i}{2} \right)}{d} + \frac{\tan(c + dx) (a^2 (B + A i) + B a^2 + A a^2 i)}{d}$$

$$+ \frac{\ln(\tan(c + dx) + i) (2 A a^2 - B a^2 2i)}{d} - \frac{B a^2 \tan(c + dx)^3}{3 d}$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2,x)`

output `(tan(c + d*x)^2*((a^2*(A*1i + B)*1i)/2 + (B*a^2*1i)/2))/d + (tan(c + d*x)*(A*a^2*1i + a^2*(A*1i + B) + B*a^2))/d + (log(tan(c + d*x) + 1i)*(2*A*a^2 - B*a^2*2i))/d - (B*a^2*tan(c + d*x)^3)/(3*d)`

3.11 $\int (a + ia \tan(c + dx))^2 (A + B \tan(c + dx)) dx$

3.11.1	Optimal result	366
3.11.2	Mathematica [A] (verified)	366
3.11.3	Rubi [A] (verified)	367
3.11.4	Maple [A] (verified)	368
3.11.5	Fricas [A] (verification not implemented)	369
3.11.6	Sympy [A] (verification not implemented)	369
3.11.7	Maxima [A] (verification not implemented)	370
3.11.8	Giac [B] (verification not implemented)	370
3.11.9	Mupad [B] (verification not implemented)	371

3.11.1 Optimal result

Integrand size = 26, antiderivative size = 80

$$\begin{aligned} & \int (a + ia \tan(c + dx))^2 (A + B \tan(c + dx)) dx \\ &= 2a^2(A - iB)x - \frac{2a^2(iA + B) \log(\cos(c + dx))}{d} \\ & \quad - \frac{a^2(A - iB) \tan(c + dx)}{d} + \frac{B(a + ia \tan(c + dx))^2}{2d} \end{aligned}$$

output `2*a^2*(A-I*B)*x-2*a^2*(I*A+B)*ln(cos(d*x+c))/d-a^2*(A-I*B)*tan(d*x+c)/d+1/2*B*(a+I*a*tan(d*x+c))^2/d`

3.11.2 Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.72

$$\begin{aligned} & \int (a + ia \tan(c + dx))^2 (A + B \tan(c + dx)) dx \\ &= \frac{a^2(B + 4(iA + B) \log(i + \tan(c + dx))) - 2(A - 2iB) \tan(c + dx) - B \tan^2(c + dx)}{2d} \end{aligned}$$

input `Integrate[(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(a^2*(B + 4*(I*A + B)*Log[I + Tan[c + d*x]] - 2*(A - (2*I)*B)*Tan[c + d*x] - B*Tan[c + d*x]^2)/(2*d)`

3.11. $\int (a + ia \tan(c + dx))^2 (A + B \tan(c + dx)) dx$

3.11.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.88, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3042, 4010, 3042, 3958, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(c + dx))^2 (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(c + dx))^2 (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4010} \\
 & (A - iB) \int (i \tan(c + dx)a + a)^2 dx + \frac{B(a + ia \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \int (i \tan(c + dx)a + a)^2 dx + \frac{B(a + ia \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{3958} \\
 & (A - iB) \left(2ia^2 \int \tan(c + dx) dx - \frac{a^2 \tan(c + dx)}{d} + 2a^2 x \right) + \frac{B(a + ia \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \left(2ia^2 \int \tan(c + dx) dx - \frac{a^2 \tan(c + dx)}{d} + 2a^2 x \right) + \frac{B(a + ia \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{3956} \\
 & (A - iB) \left(-\frac{a^2 \tan(c + dx)}{d} - \frac{2ia^2 \log(\cos(c + dx))}{d} + 2a^2 x \right) + \frac{B(a + ia \tan(c + dx))^2}{2d}
 \end{aligned}$$

input `Int[(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(B*(a + I*a*Tan[c + d*x])^2)/(2*d) + (A - I*B)*(2*a^2*x - ((2*I)*a^2*Log[Cos[c + d*x]]))/d - (a^2*Tan[c + d*x])/d`

3.11.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 3958 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_.)]^2, x_Symbol] := Simp[(a^2 - b^2)*x, x] + (Simp[b^2*(Tan[c + d*x]/d), x] + Simp[2*a*b Int[Tan[c + d*x], x], x]) /; FreeQ[{a, b, c, d}, x]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)]^(m_))*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^(m/(f*m))), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

3.11.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.95

method	result
derivativedivides	$\frac{a^2 \left(-\frac{B(\tan^2(dx+c))}{2} - A \tan(dx+c) + 2iB \tan(dx+c) + \frac{(2iA+2B) \ln(1+\tan^2(dx+c))}{2} + (-2iB+2A) \arctan(\tan(dx+c)) \right)}{d}$
default	$\frac{a^2 \left(-\frac{B(\tan^2(dx+c))}{2} - A \tan(dx+c) + 2iB \tan(dx+c) + \frac{(2iA+2B) \ln(1+\tan^2(dx+c))}{2} + (-2iB+2A) \arctan(\tan(dx+c)) \right)}{d}$
norman	$(-2iB a^2 + 2A a^2) x - \frac{(-2iB a^2 + A a^2) \tan(dx+c)}{d} - \frac{B a^2 (\tan^2(dx+c))}{2d} + \frac{(iA a^2 + B a^2) \ln(1+\tan^2(dx+c))}{d}$
parallelrisch	$\frac{-4iBx a^2 d + 2iA \ln(1+\tan^2(dx+c)) a^2 + 4Ax a^2 d + 4iB \tan(dx+c) a^2 - B(\tan^2(dx+c)) a^2 - 2A \tan(dx+c) a^2 + 2B \ln(1+\tan^2(dx+c)) a^2}{2d}$
parts	$A a^2 x + \frac{(2iA a^2 + B a^2) \ln(1+\tan^2(dx+c))}{2d} + \frac{(2iB a^2 - A a^2) (\tan(dx+c) - \arctan(\tan(dx+c)))}{d} - \frac{B a^2 \left(\frac{\tan^2(dx+c)}{2} \right)}{d}$
risch	$\frac{4ia^2 Bc}{d} - \frac{4a^2 Ac}{d} - \frac{2a^2 (iA e^{2i(dx+c)} + 3B e^{2i(dx+c)} + iA + 2B)}{d(e^{2i(dx+c)} + 1)^2} - \frac{2a^2 \ln(e^{2i(dx+c)} + 1) B}{d} - \frac{2ia^2 \ln(e^{2i(dx+c)} + 1) A}{d}$

input `int((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

3.11. $\int (a + ia \tan(c + dx))^2 (A + B \tan(c + dx)) dx$

output $1/d*a^2*(-1/2*B*\tan(d*x+c)^2-A*\tan(d*x+c)+2*I*B*\tan(d*x+c)+1/2*(2*B+2*I*A)*\ln(1+\tan(d*x+c)^2)+(2*A-2*I*B)*\arctan(\tan(d*x+c)))$

3.11.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.51

$$\int (a + ia \tan(c + dx))^2 (A + B \tan(c + dx)) dx = \frac{2((iA + 3B)a^2 e^{(2i dx + 2i c)} + (iA + 2B)a^2 + ((iA + B)a^2 e^{(4i dx + 4i c)} + 2(iA + B)a^2 e^{(2i dx + 2i c)} + (iA + B)a^2 e^{(4i dx + 4i c)} + 2d e^{(2i dx + 2i c)} + d$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output $-2*((I*A + 3*B)*a^2*e^{(2*I*d*x + 2*I*c)} + (I*A + 2*B)*a^2 + ((I*A + B)*a^2*e^{(4*I*d*x + 4*I*c)} + 2*(I*A + B)*a^2*e^{(2*I*d*x + 2*I*c)} + (I*A + B)*a^2)*\log(e^{(2*I*d*x + 2*I*c)} + 1)/(d*e^{(4*I*d*x + 4*I*c)} + 2*d*e^{(2*I*d*x + 2*I*c)} + d)$

3.11.6 Sympy [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.52

$$\int (a + ia \tan(c + dx))^2 (A + B \tan(c + dx)) dx = -\frac{2ia^2(A - iB) \log(e^{2idx} + e^{-2ic})}{d} + \frac{-2iAa^2 - 4Ba^2 + (-2iAa^2 e^{2ic} - 6Ba^2 e^{2ic}) e^{2idx}}{de^{4ic}e^{4idx} + 2de^{2ic}e^{2idx} + d}$$

input `integrate((a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output $-2*I*a**2*(A - I*B)*\log(\exp(2*I*d*x) + \exp(-2*I*c))/d + (-2*I*A*a**2 - 4*B*a**2 + (-2*I*A*a**2*\exp(2*I*c) - 6*B*a**2*\exp(2*I*c))*\exp(2*I*d*x))/(d*\exp(4*I*c)*\exp(4*I*d*x) + 2*d*\exp(2*I*c)*\exp(2*I*d*x) + d)$

3.11.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.89

$$\int (a + ia \tan(c + dx))^2 (A + B \tan(c + dx)) dx = \frac{Ba^2 \tan(dx + c)^2 - 4(dx + c)(A - iB)a^2 - 2(iA + B)a^2 \log(\tan(dx + c)^2 + 1) + 2(A - 2iB)a^2 \tan(dx + c)}{2d}$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(B*a^2*tan(d*x + c)^2 - 4*(d*x + c)*(A - I*B)*a^2 - 2*(I*A + B)*a^2*log(tan(d*x + c)^2 + 1) + 2*(A - 2*I*B)*a^2*tan(d*x + c))/d`

3.11.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 214 vs. $2(70) = 140$.

Time = 0.38 (sec) , antiderivative size = 214, normalized size of antiderivative = 2.68

$$\int (a + ia \tan(c + dx))^2 (A + B \tan(c + dx)) dx = \frac{2(iAa^2 e^{(4i dx + 4i c)} \log(e^{(2i dx + 2i c)} + 1) + Ba^2 e^{(4i dx + 4i c)} \log(e^{(2i dx + 2i c)} + 1) + 2iAa^2 e^{(2i dx + 2i c)} \log(e^{(2i dx + 2i c)} + 1) + 2(A - 2iB)a^2 \tan(dx + c))}{2d}$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `-2*(I*A*a^2*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + B*a^2*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 2*I*A*a^2*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 2*B*a^2*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + I*A*a^2*e^(2*I*d*x + 2*I*c) + 3*B*a^2*e^(2*I*d*x + 2*I*c) + I*A*a^2*log(e^(2*I*d*x + 2*I*c) + 1) + B*a^2*log(e^(2*I*d*x + 2*I*c) + 1) + I*A*a^2 + 2*B*a^2)/(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)`

3.11.9 Mupad [B] (verification not implemented)

Time = 7.26 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.95

$$\int (a + ia \tan(c + dx))^2 (A + B \tan(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx) + 1i) (2B a^2 + A a^2 2i)}{d}$$

$$+ \frac{\tan(c + dx) (a^2 (B + A 1i) 1i + B a^2 1i)}{d} - \frac{B a^2 \tan(c + dx)^2}{2d}$$

input `int((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2,x)`output `(log(tan(c + d*x) + 1i)*(A*a^2*2i + 2*B*a^2))/d + (tan(c + d*x)*(a^2*(A*1i + B)*1i + B*a^2*1i))/d - (B*a^2*tan(c + d*x)^2)/(2*d)`

3.12 $\int \cot(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.12.1	Optimal result	372
3.12.2	Mathematica [A] (verified)	372
3.12.3	Rubi [A] (verified)	373
3.12.4	Maple [A] (verified)	375
3.12.5	Fricas [A] (verification not implemented)	376
3.12.6	Sympy [A] (verification not implemented)	376
3.12.7	Maxima [A] (verification not implemented)	377
3.12.8	Giac [B] (verification not implemented)	377
3.12.9	Mupad [B] (verification not implemented)	378

3.12.1 Optimal result

Integrand size = 32, antiderivative size = 75

$$\int \cot(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= 2a^2(iA+B)x + \frac{a^2(A-2iB) \log(\cos(c+dx))}{d}$$

$$+ \frac{a^2A \log(\sin(c+dx))}{d} + \frac{iB(a^2+ia^2 \tan(c+dx))}{d}$$

```
output 2*a^2*(I*A+B)*x+a^2*(A-2*I*B)*ln(cos(d*x+c))/d+a^2*A*ln(sin(d*x+c))/d+I*B*(a^2+I*a^2*tan(d*x+c))/d
```

3.12.2 Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.61

$$\int \cot(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{a^2(A \log(\tan(c+dx)) - 2(A-iB) \log(i+\tan(c+dx)) - B \tan(c+dx))}{d}$$

```
input Integrate[Cot[c + d*x]*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]
```

output $(a^2*(A*\text{Log}[\text{Tan}[c + d*x]] - 2*(A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]] - B*\text{Tan}[c + d*x]))/d$

3.12.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3042, 4077, 3042, 4072, 3042, 3956, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\tan(c + dx)} dx \\
 & \quad \downarrow \text{4077} \\
 & \int \cot(c + dx)(i \tan(c + dx)a + a)(aA + a(iA + 2B) \tan(c + dx)) dx + \frac{iB(a^2 + ia^2 \tan(c + dx))}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(i \tan(c + dx)a + a)(aA + a(iA + 2B) \tan(c + dx))}{\tan(c + dx)} dx + \frac{iB(a^2 + ia^2 \tan(c + dx))}{d} \\
 & \quad \downarrow \text{4072} \\
 & -\left(a^2(A - 2iB) \int \tan(c + dx) dx\right) + \int \cot(c + dx) \left(Aa^2 + 2(iA + B) \tan(c + dx)a^2\right) dx + \\
 & \quad \frac{iB(a^2 + ia^2 \tan(c + dx))}{d} \\
 & \quad \downarrow \text{3042} \\
 & -\left(a^2(A - 2iB) \int \tan(c + dx) dx\right) + \int \frac{Aa^2 + 2(iA + B) \tan(c + dx)a^2}{\tan(c + dx)} dx + \\
 & \quad \frac{iB(a^2 + ia^2 \tan(c + dx))}{d} \\
 & \quad \downarrow \text{3956} \\
 & \int \frac{Aa^2 + 2(iA + B) \tan(c + dx)a^2}{\tan(c + dx)} dx + \frac{a^2(A - 2iB) \log(\cos(c + dx))}{d} + \frac{iB(a^2 + ia^2 \tan(c + dx))}{d}
 \end{aligned}$$

3.12. $\int \cot(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \downarrow 4014 \\
& a^2 A \int \cot(c + dx) dx + \frac{a^2(A - 2iB) \log(\cos(c + dx))}{d} + 2a^2 x(B + iA) + \frac{iB(a^2 + ia^2 \tan(c + dx))}{d} \\
& \downarrow 3042 \\
& a^2 A \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + \frac{a^2(A - 2iB) \log(\cos(c + dx))}{d} + 2a^2 x(B + iA) + \\
& \quad \frac{iB(a^2 + ia^2 \tan(c + dx))}{d} \\
& \downarrow 25 \\
& -a^2 A \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + \frac{a^2(A - 2iB) \log(\cos(c + dx))}{d} + 2a^2 x(B + iA) + \\
& \quad \frac{iB(a^2 + ia^2 \tan(c + dx))}{d} \\
& \downarrow 3956 \\
& \frac{a^2(A - 2iB) \log(\cos(c + dx))}{d} + 2a^2 x(B + iA) + \frac{a^2 A \log(-\sin(c + dx))}{d} + \frac{iB(a^2 + ia^2 \tan(c + dx))}{d}
\end{aligned}$$

input `Int[Cot[c + d*x]*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `2*a^2*(I*A + B)*x + (a^2*(A - (2*I)*B)*Log[Cos[c + d*x]])/d + (a^2*A*Log[-Sin[c + d*x]])/d + (I*B*(a^2 + I*a^2*Tan[c + d*x]))/d`

3.12.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4072 Int((((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_
.)*(x_)]))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*(d/
b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c
- a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d
, e, f, A, B}, x] && NeQ[b*c - a*d, 0]
```

```
rule 4077 Int(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan
[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c -
a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a
, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] &&
GtQ[m, 1] && !LtQ[n, -1]
```

3.12.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.84

method	result
parallelrisch	$\frac{a^2(2iAxd+iB \ln(\sec^2(dx+c))+2Bdx-A \ln(\sec^2(dx+c))+A \ln(\tan(dx+c))-B \tan(dx+c))}{d}$
norman	$(2iAa^2 + 2Ba^2)x - \frac{Ba^2 \tan(dx+c)}{d} + \frac{Aa^2 \ln(\tan(dx+c))}{d} - \frac{(-iBa^2+Aa^2) \ln(1+\tan^2(dx+c))}{d}$
derivativedivides	$\frac{a^2 \left(\frac{(-2iB+2A) \ln(\cot^2(dx+c)+1)}{2} + (2iA+2B) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)) \right) + (2iB-A) \ln(\cot(dx+c)) + \frac{B}{\cot(dx+c)} \right)}{d}$
default	$\frac{a^2 \left(\frac{(-2iB+2A) \ln(\cot^2(dx+c)+1)}{2} + (2iA+2B) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)) \right) + (2iB-A) \ln(\cot(dx+c)) + \frac{B}{\cot(dx+c)} \right)}{d}$
risch	$-\frac{4ia^2Ac}{d} - \frac{4a^2Bc}{d} - \frac{2iBa^2}{d(e^{2i(dx+c)}+1)} + \frac{Aa^2 \ln(e^{2i(dx+c)}-1)}{d} - \frac{2ia^2 \ln(e^{2i(dx+c)}+1)B}{d} + \frac{a^2 \ln(e^{2i(dx+c)}+1)}{d}$

3.12. $\int \cot(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

```
input int(cot(d*x+c)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/d*a^2*(2*I*A*x*d+I*B*ln(sec(d*x+c)^2)+2*B*d*x-A*ln(sec(d*x+c)^2)+A*ln(tan(d*x+c))-B*tan(d*x+c))
```

3.12.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.29

$$\int \cot(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{-2iBa^2 + ((A-2iB)a^2 e^{(2i dx+2i c)} + (A-2iB)a^2) \log(e^{(2i dx+2i c)} + 1) + (Aa^2 e^{(2i dx+2i c)} + Aa^2) \log(e^{(2i dx+2i c)} + d)}{de^{(2i dx+2i c)} + d}$$

```
input integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output (-2*I*B*a^2 + ((A - 2*I*B)*a^2*e^(2*I*d*x + 2*I*c) + (A - 2*I*B)*a^2)*log(e^(2*I*d*x + 2*I*c) + 1) + (A*a^2*e^(2*I*d*x + 2*I*c) + A*a^2)*log(e^(2*I*d*x + 2*I*c) - 1))/(d*e^(2*I*d*x + 2*I*c) + d)
```

3.12.6 Sympy [A] (verification not implemented)

Time = 1.22 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.45

$$\int \cot(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{Aa^2 \log(e^{2idx} - e^{-2ic})}{d} - \frac{2iBa^2}{de^{2ic}e^{2idx} + d}$$

$$+ \frac{a^2(A-2iB) \log\left(e^{2idx} + \frac{(-iAa^2 - Ba^2 + ia^2(A-2iB))e^{-2ic}}{Ba^2}\right)}{d}$$

```
input integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)
```

```
output A*a**2*log(exp(2*I*d*x) - exp(-2*I*c))/d - 2*I*B*a**2/(d*exp(2*I*c)*exp(2*I*d*x) + d) + a**2*(A - 2*I*B)*log(exp(2*I*d*x) + (-I*A*a**2 - B*a**2 + I*a**2*(A - 2*I*B))*exp(-2*I*c)/(B*a**2))/d
```

3.12. $\int \cot(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.12.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.89

$$\int \cot(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{2(dx + c)(-iA - B)a^2 + (A - iB)a^2 \log(\tan(dx + c)^2 + 1) - Aa^2 \log(\tan(dx + c)) + Ba^2 \tan(dx + c)}{d}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-(2*(d*x + c)*(-I*A - B)*a^2 + (A - I*B)*a^2*log(tan(d*x + c)^2 + 1) - A*a^2*log(tan(d*x + c)) + B*a^2*tan(d*x + c))/d`

3.12.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 174 vs. 2(67) = 134.

Time = 0.64 (sec) , antiderivative size = 174, normalized size of antiderivative = 2.32

$$\int \cot(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{Aa^2 \log(\tan(\frac{1}{2} dx + \frac{1}{2} c)) + (Aa^2 - 2iBa^2) \log(\tan(\frac{1}{2} dx + \frac{1}{2} c) + 1) - 4(Aa^2 - iBa^2) \log(\tan(\frac{1}{2} dx + \frac{1}{2} c) - 1) + (Aa^2 - 2iB*a^2) \log(\tan(\frac{1}{2} dx + \frac{1}{2} c) + I) + (Aa^2 - 2iB*a^2) \log(\tan(\frac{1}{2} dx + \frac{1}{2} c) - I) - (Aa^2 \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 2iB*a^2 \tan(\frac{1}{2} dx + \frac{1}{2} c) - Aa^2 + 2iB*a^2) / (\tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 1)}{d}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `(A*a^2*log(tan(1/2*d*x + 1/2*c)) + (A*a^2 - 2*I*B*a^2)*log(tan(1/2*d*x + 1/2*c) + 1) - 4*(A*a^2 - I*B*a^2)*log(tan(1/2*d*x + 1/2*c) + I) + (A*a^2 - 2*I*B*a^2)*log(tan(1/2*d*x + 1/2*c) - 1) - (A*a^2*tan(1/2*d*x + 1/2*c)^2 - 2*I*B*a^2*tan(1/2*d*x + 1/2*c) - A*a^2 + 2*I*B*a^2)/(tan(1/2*d*x + 1/2*c)^2 - 1))/d`

3.12.9 Mupad [B] (verification not implemented)

Time = 8.13 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.93

$$\int \cot(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{A a^2 \ln(\tan(c + dx))}{d} - \frac{2 A a^2 \ln(\tan(c + dx) + 1i)}{d}$$

$$- \frac{B a^2 \tan(c + dx)}{d} + \frac{B a^2 \ln(\tan(c + dx) + 1i) 2i}{d}$$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2,x)`output `(A*a^2*log(tan(c + d*x)))/d - (2*A*a^2*log(tan(c + d*x) + 1i))/d + (B*a^2*log(tan(c + d*x) + 1i)*2i)/d - (B*a^2*tan(c + d*x))/d`

3.13 $\int \cot^2(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.13.1	Optimal result	379
3.13.2	Mathematica [A] (verified)	379
3.13.3	Rubi [A] (verified)	380
3.13.4	Maple [A] (verified)	382
3.13.5	Fricas [A] (verification not implemented)	383
3.13.6	Sympy [A] (verification not implemented)	383
3.13.7	Maxima [A] (verification not implemented)	384
3.13.8	Giac [B] (verification not implemented)	384
3.13.9	Mupad [B] (verification not implemented)	385

3.13.1 Optimal result

Integrand size = 34, antiderivative size = 79

$$\begin{aligned} & \int \cot^2(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx \\ &= -2a^2(A-iB)x + \frac{a^2B \log(\cos(c+dx))}{d} \\ & \quad + \frac{a^2(2iA+B) \log(\sin(c+dx))}{d} - \frac{A \cot(c+dx)(a^2+ia^2 \tan(c+dx))}{d} \end{aligned}$$

output `-2*a^2*(A-I*B)*x+a^2*B*ln(cos(d*x+c))/d+a^2*(2*I*A+B)*ln(sin(d*x+c))/d-A*cot(d*x+c)*(a^2+I*a^2*tan(d*x+c))/d`

3.13.2 Mathematica [A] (verified)

Time = 0.75 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.68

$$\begin{aligned} & \int \cot^2(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx \\ &= \frac{a^2(-A \cot(c+dx) + (2iA+B) \log(\tan(c+dx)) - 2i(A-iB) \log(i+\tan(c+dx)))}{d} \end{aligned}$$

input `Integrate[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output $(a^2*(-(A*\cot[c + d*x]) + ((2*I)*A + B)*\log[\tan[c + d*x]] - (2*I)*(A - I*B)*\log[I + \tan[c + d*x]]))/d$

3.13.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3042, 4076, 3042, 4072, 3042, 3956, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\tan(c + dx)^2} dx \\
 & \quad \downarrow \text{4076} \\
 & \int \cot(c + dx)(i \tan(c + dx)a + a)(a(2iA + B) + iaB \tan(c + dx)) dx - \\
 & \quad \frac{A \cot(c + dx)(a^2 + ia^2 \tan(c + dx))}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(i \tan(c + dx)a + a)(a(2iA + B) + iaB \tan(c + dx))}{\tan(c + dx)} dx - \frac{A \cot(c + dx)(a^2 + ia^2 \tan(c + dx))}{d} \\
 & \quad \downarrow \text{4072} \\
 & \int \cot(c + dx)(a^2(2iA + B) - 2a^2(A - iB) \tan(c + dx)) dx + a^2(-B) \int \tan(c + dx) dx - \\
 & \quad \frac{A \cot(c + dx)(a^2 + ia^2 \tan(c + dx))}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{a^2(2iA + B) - 2a^2(A - iB) \tan(c + dx)}{\tan(c + dx)} dx + a^2(-B) \int \tan(c + dx) dx - \\
 & \quad \frac{A \cot(c + dx)(a^2 + ia^2 \tan(c + dx))}{d} \\
 & \quad \downarrow \text{3956}
 \end{aligned}$$

$$\begin{aligned}
& \int \frac{a^2(2iA + B) - 2a^2(A - iB) \tan(c + dx)}{\tan(c + dx)} dx - \frac{A \cot(c + dx) (a^2 + ia^2 \tan(c + dx))}{d} + \\
& \quad \frac{a^2 B \log(\cos(c + dx))}{d} \\
& \quad \downarrow \text{4014} \\
& a^2(B + 2iA) \int \cot(c + dx) dx - 2a^2 x(A - iB) - \frac{A \cot(c + dx) (a^2 + ia^2 \tan(c + dx))}{d} + \\
& \quad \frac{a^2 B \log(\cos(c + dx))}{d} \\
& \quad \downarrow \text{3042} \\
& a^2(B + 2iA) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - 2a^2 x(A - iB) - \frac{A \cot(c + dx) (a^2 + ia^2 \tan(c + dx))}{d} + \\
& \quad \frac{a^2 B \log(\cos(c + dx))}{d} \\
& \quad \downarrow \text{25} \\
& -a^2(B + 2iA) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - 2a^2 x(A - iB) - \\
& \quad \frac{A \cot(c + dx) (a^2 + ia^2 \tan(c + dx))}{d} + \frac{a^2 B \log(\cos(c + dx))}{d} \\
& \quad \downarrow \text{3956} \\
& \frac{a^2(B + 2iA) \log(-\sin(c + dx))}{d} - 2a^2 x(A - iB) - \frac{A \cot(c + dx) (a^2 + ia^2 \tan(c + dx))}{d} + \\
& \quad \frac{a^2 B \log(\cos(c + dx))}{d}
\end{aligned}$$

input `Int[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `-2*a^2*(A - I*B)*x + (a^2*B*Log[Cos[c + d*x]])/d + (a^2*((2*I)*A + B)*Log[-Sin[c + d*x]])/d - (A*Cot[c + d*x]*(a^2 + I*a^2*Tan[c + d*x]))/d`

3.13.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

3.13. $\int \cot^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4072 `Int((((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*(d/b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c - a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0]`

rule 4076 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.13.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.94

method	result
parallelrisch	$\frac{a^2(2iBdx+2iA \ln(\tan(dx+c))-iA \ln(\sec^2(dx+c))-2Adx-A \cot(dx+c)+B \ln(\tan(dx+c))-B \ln(\sec^2(dx+c)))}{d}$
derivativedivides	$-\frac{A a^2(dx+c)+B a^2 \ln(\cos(dx+c))+2iA a^2 \ln(\sin(dx+c))+2iB a^2(dx+c)+A a^2(-\cot(dx+c)-dx-c)+B a^2 \ln(\sin(dx+c))}{d}$
default	$-\frac{A a^2(dx+c)+B a^2 \ln(\cos(dx+c))+2iA a^2 \ln(\sin(dx+c))+2iB a^2(dx+c)+A a^2(-\cot(dx+c)-dx-c)+B a^2 \ln(\sin(dx+c))}{d}$
norman	$\frac{(2iB a^2-2A a^2)x \tan(dx+c)-\frac{A a^2}{d}}{\tan(dx+c)} + \frac{(2iA a^2+B a^2) \ln(\tan(dx+c))}{d} - \frac{(iA a^2+B a^2) \ln(1+\tan^2(dx+c))}{d}$
risch	$-\frac{4ia^2Bc}{d} + \frac{4a^2Ac}{d} - \frac{2iAa^2}{d(e^{2i(dx+c)}-1)} + \frac{a^2 \ln(e^{2i(dx+c)}-1)B}{d} + \frac{2ia^2 \ln(e^{2i(dx+c)}-1)A}{d} + \frac{a^2 \ln(e^{2i(dx+c)}+1)}{d}$

3.13. $\int \cot^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

input `int(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output $a^2(2iBdx + 2IA \ln(\tan(dx+c)) - IA \ln(\sec(dx+c)^2) - 2Adx - A \cot(dx+c) + B \ln(\tan(dx+c)) - B \ln(\sec(dx+c)^2))/d$

3.13.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.29

$$\int \cot^2(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{-2iAa^2 + (Ba^2 e^{2i dx+2i c} - Ba^2) \log(e^{2i dx+2i c} + 1) + ((2iA+B)a^2 e^{2i dx+2i c} + (-2iA-B)a^2) \log(d e^{2i dx+2i c} - d)}{d e^{2i dx+2i c} - d}$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output $(-2IAa^2 + (Ba^2 e^{2I dx+2I c} - Ba^2) \log(e^{2I dx+2I c} + 1) + ((2IA+B)a^2 e^{2I dx+2I c} + (-2IA-B)a^2) \log(e^{2I dx+2I c} - 1))/(d e^{2I dx+2I c} - d)$

3.13.6 Sympy [A] (verification not implemented)

Time = 1.32 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.38

$$\int \cot^2(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= -\frac{2iAa^2}{d e^{2ic} e^{2idx} - d} + \frac{Ba^2 \log(e^{2idx} + e^{-2ic})}{d}$$

$$+ \frac{ia^2 \cdot (2A - iB) \log\left(e^{2idx} + \frac{(Aa^2 - iBa^2 - a^2 \cdot (2A - iB))e^{-2ic}}{Aa^2}\right)}{d}$$

input `integrate(cot(d*x+c)**2*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output $-2IAa^2/(d \exp(2Ic) \exp(2I dx) - d) + Ba^2 \log(\exp(2I dx) + \exp(-2Ic))/d + IA^2(2A - IB) \log(\exp(2I dx) + (Aa^2 - IBa^2 - a^2(2A - IB)) \exp(-2Ic)/(Aa^2))/d$

3.13. $\int \cot^2(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.13.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.94

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{2(dx + c)(A - iB)a^2 - (-iA - B)a^2 \log(\tan(dx + c)^2 + 1) - (2iA + B)a^2 \log(\tan(dx + c)) + \frac{A}{\tan(dx + c)}}{d}$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-(2*(d*x + c)*(A - I*B)*a^2 - (-I*A - B)*a^2*log(tan(d*x + c)^2 + 1) - (2*I*A + B)*a^2*log(tan(d*x + c)) + A*a^2/tan(d*x + c))/d`

3.13.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 155 vs. 2(73) = 146.

Time = 0.86 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.96

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{2Ba^2 \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) + 1) + 2Ba^2 \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1) + Aa^2 \tan(\frac{1}{2}dx + \frac{1}{2}c) - 8(iAa^2 + B)}{d}$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `1/2*(2*B*a^2*log(tan(1/2*d*x + 1/2*c) + 1) + 2*B*a^2*log(tan(1/2*d*x + 1/2*c) - 1) + A*a^2*tan(1/2*d*x + 1/2*c) - 8*(I*A*a^2 + B*a^2)*log(tan(1/2*d*x + 1/2*c) + I) + 2*(2*I*A*a^2 + B*a^2)*log(tan(1/2*d*x + 1/2*c)) + (-4*I*A*a^2*tan(1/2*d*x + 1/2*c) - 2*B*a^2*tan(1/2*d*x + 1/2*c) - A*a^2)/tan(1/2*d*x + 1/2*c))/d`

3.13.9 Mupad [B] (verification not implemented)

Time = 7.69 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.10

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{B a^2 \ln(\tan(c + dx))}{d} - \frac{2 B a^2 \ln(\tan(c + dx) + 1i)}{d} - \frac{A a^2 \cot(c + dx)}{d}$$

$$+ \frac{A a^2 \ln(\tan(c + dx)) 2i}{d} - \frac{A a^2 \ln(\tan(c + dx) + 1i) 2i}{d}$$

input `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2,x)`output `(A*a^2*log(tan(c + d*x))*2i)/d + (B*a^2*log(tan(c + d*x)))/d - (A*a^2*log(tan(c + d*x) + 1i)*2i)/d - (2*B*a^2*log(tan(c + d*x) + 1i))/d - (A*a^2*cot(c + d*x))/d`

3.14 $\int \cot^3(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.14.1	Optimal result	386
3.14.2	Mathematica [A] (verified)	386
3.14.3	Rubi [A] (verified)	387
3.14.4	Maple [A] (verified)	390
3.14.5	Fricas [A] (verification not implemented)	390
3.14.6	Sympy [A] (verification not implemented)	391
3.14.7	Maxima [A] (verification not implemented)	391
3.14.8	Giac [B] (verification not implemented)	392
3.14.9	Mupad [B] (verification not implemented)	392

3.14.1 Optimal result

Integrand size = 34, antiderivative size = 94

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= -2a^2(iA+B)x - \frac{a^2(3iA+2B) \cot(c+dx)}{2d}$$

$$- \frac{2a^2(A-iB) \log(\sin(c+dx))}{d} - \frac{A \cot^2(c+dx)(a^2+ia^2 \tan(c+dx))}{2d}$$

output

```
-2*a^2*(I*A+B)*x-1/2*a^2*(3*I*A+2*B)*cot(d*x+c)/d-2*a^2*(A-I*B)*ln(sin(d*x+c))/d-1/2*A*cot(d*x+c)^2*(a^2+I*a^2*tan(d*x+c))/d
```

3.14.2 Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.18

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= a^2 \left(-\frac{2iA \cot(c+dx)}{d} - \frac{B \cot(c+dx)}{d} - \frac{A \cot^2(c+dx)}{2d} - \frac{2A \log(\tan(c+dx))}{d} \right.$$

$$\left. + \frac{2iB \log(\tan(c+dx))}{d} + \frac{2A \log(i + \tan(c+dx))}{d} - \frac{2iB \log(i + \tan(c+dx))}{d} \right)$$

input `Integrate[Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `a^2*(((-2*I)*A*Cot[c + d*x])/d - (B*Cot[c + d*x])/d - (A*Cot[c + d*x]^2)/(2*d) - (2*A*Log[Tan[c + d*x]])/d + ((2*I)*B*Log[Tan[c + d*x]])/d + (2*A*Log[I + Tan[c + d*x]])/d - ((2*I)*B*Log[I + Tan[c + d*x]])/d)`

3.14.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.06, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3042, 4076, 3042, 4074, 27, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^3(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\tan(c + dx)^3} dx \\
 & \quad \downarrow \text{4076} \\
 & \frac{1}{2} \int \cot^2(c + dx)(i \tan(c + dx)a + a)(a(3iA + 2B) - a(A - 2iB) \tan(c + dx)) dx - \\
 & \quad \frac{A \cot^2(c + dx) (a^2 + ia^2 \tan(c + dx))}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \frac{(i \tan(c + dx)a + a)(a(3iA + 2B) - a(A - 2iB) \tan(c + dx))}{\tan(c + dx)^2} dx - \\
 & \quad \frac{A \cot^2(c + dx) (a^2 + ia^2 \tan(c + dx))}{2d} \\
 & \quad \downarrow \text{4074} \\
 & \frac{1}{2} \left(\int -4 \cot(c + dx) ((A - iB)a^2 + (iA + B) \tan(c + dx)a^2) dx - \frac{a^2(2B + 3iA) \cot(c + dx)}{d} \right) - \\
 & \quad \frac{A \cot^2(c + dx) (a^2 + ia^2 \tan(c + dx))}{2d} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.14. $\int \cot^3(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \frac{1}{2} \left(-4 \int \cot(c+dx) \left((A-iB)a^2 + (iA+B)\tan(c+dx)a^2 \right) dx - \frac{a^2(2B+3iA)\cot(c+dx)}{d} \right) - \\
& \quad \frac{A \cot^2(c+dx) (a^2 + ia^2 \tan(c+dx))}{2d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \left(-4 \int \frac{(A-iB)a^2 + (iA+B)\tan(c+dx)a^2}{\tan(c+dx)} dx - \frac{a^2(2B+3iA)\cot(c+dx)}{d} \right) - \\
& \quad \frac{A \cot^2(c+dx) (a^2 + ia^2 \tan(c+dx))}{2d} \\
& \quad \downarrow \text{4014} \\
& \frac{1}{2} \left(-4 \left(a^2(A-iB) \int \cot(c+dx) dx + a^2 x(B+iA) \right) - \frac{a^2(2B+3iA)\cot(c+dx)}{d} \right) - \\
& \quad \frac{A \cot^2(c+dx) (a^2 + ia^2 \tan(c+dx))}{2d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \left(-4 \left(a^2(A-iB) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx + a^2 x(B+iA) \right) - \frac{a^2(2B+3iA)\cot(c+dx)}{d} \right) - \\
& \quad \frac{A \cot^2(c+dx) (a^2 + ia^2 \tan(c+dx))}{2d} \\
& \quad \downarrow \text{25} \\
& \frac{1}{2} \left(-4 \left(a^2 x(B+iA) - a^2(A-iB) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx \right) - \frac{a^2(2B+3iA)\cot(c+dx)}{d} \right) - \\
& \quad \frac{A \cot^2(c+dx) (a^2 + ia^2 \tan(c+dx))}{2d} \\
& \quad \downarrow \text{3956} \\
& \frac{1}{2} \left(-\frac{a^2(2B+3iA)\cot(c+dx)}{d} - 4 \left(\frac{a^2(A-iB)\log(-\sin(c+dx))}{d} + a^2 x(B+iA) \right) \right) - \\
& \quad \frac{A \cot^2(c+dx) (a^2 + ia^2 \tan(c+dx))}{2d}
\end{aligned}$$

input `Int[Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `((-(a^2*((3*I)*A + 2*B)*Cot[c + d*x])/d) - 4*(a^2*(I*A + B)*x + (a^2*(A - I*B)*Log[-Sin[c + d*x]]/d))/2 - (A*Cot[c + d*x]^2*(a^2 + I*a^2*Tan[c + d*x]))/(2*d)`

3.14.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4014 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`
- rule 4074 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`
- rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.14.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.82

method	result
parallelrisc	$\frac{2a^2 \left(\left(-\frac{A}{2} + \frac{iB}{2} \right) \ln(\sec^2(dx+c)) + (-iB+A) \ln(\tan(dx+c)) + \frac{A(\cot^2(dx+c))}{4} + \cot(dx+c) \left(iA + \frac{B}{2} \right) + (iA+B)xd \right)}{d}$
derivativedivides	$\frac{-A a^2 \ln(\sin(dx+c)) - B a^2(dx+c) + 2iA a^2(-\cot(dx+c) - dx - c) + 2iB a^2 \ln(\sin(dx+c)) + A a^2 \left(-\frac{\cot^2(dx+c)}{2} - \ln(\sin(dx+c)) \right)}{d}$
default	$\frac{-A a^2 \ln(\sin(dx+c)) - B a^2(dx+c) + 2iA a^2(-\cot(dx+c) - dx - c) + 2iB a^2 \ln(\sin(dx+c)) + A a^2 \left(-\frac{\cot^2(dx+c)}{2} - \ln(\sin(dx+c)) \right)}{d}$
risc	$\frac{4a^2 Bc}{d} + \frac{4ia^2 Ac}{d} - \frac{2ia^2(3iA e^{2i(dx+c)} + B e^{2i(dx+c)} - 2iA - B)}{d(e^{2i(dx+c)} - 1)^2} + \frac{2ia^2 \ln(e^{2i(dx+c)} - 1)B}{d} - \frac{2A a^2 \ln(e^{2i(dx+c)} - 1)}{d}$
norman	$\frac{(-2iA a^2 - 2B a^2)x(\tan^2(dx+c)) - \frac{A a^2}{2d} - \frac{(2iA a^2 + B a^2) \tan(dx+c)}{d}}{\tan(dx+c)^2} + \frac{(-iB a^2 + A a^2) \ln(1 + \tan^2(dx+c))}{d} - \frac{2(-iB a^2 + A a^2)}{d}$

```
input int(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output -2*a^2*((-1/2*A+1/2*I*B)*ln(sec(d*x+c)^2)+(A-I*B)*ln(tan(d*x+c))+1/4*A*cot(d*x+c)^2+cot(d*x+c)*(I*A+1/2*B)+(I*A+B)*x*d)/d
```

3.14.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.31

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{2((3A - iB)a^2 e^{(2i dx + 2i c)} - (2A - iB)a^2 - ((A - iB)a^2 e^{(4i dx + 4i c)} - 2(A - iB)a^2 e^{(2i dx + 2i c)} + (A - iB)a^2))}{de^{(4i dx + 4i c)} - 2de^{(2i dx + 2i c)} + d}$$

```
input integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output 2*((3*A - I*B)*a^2*e^(2*I*d*x + 2*I*c) - (2*A - I*B)*a^2 - ((A - I*B)*a^2*e^(4*I*d*x + 4*I*c) - 2*(A - I*B)*a^2*e^(2*I*d*x + 2*I*c) + (A - I*B)*a^2)*log(e^(2*I*d*x + 2*I*c) - 1)/(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)
```

3.14. $\int \cot^3(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.14.6 Sympy [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.27

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= -\frac{2a^2(A - iB) \log(e^{2idx} - e^{-2ic})}{d} + \frac{-4Aa^2 + 2iBa^2 + (6Aa^2e^{2ic} - 2iBa^2e^{2ic})e^{2idx}}{de^{4ic}e^{4idx} - 2de^{2ic}e^{2idx} + d}$$

input `integrate(cot(d*x+c)**3*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`output `-2*a**2*(A - I*B)*log(exp(2*I*d*x) - exp(-2*I*c))/d + (-4*A*a**2 + 2*I*B*a**2 + (6*A*a**2*exp(2*I*c) - 2*I*B*a**2*exp(2*I*c))*exp(2*I*d*x))/(d*exp(4*I*c)*exp(4*I*d*x) - 2*d*exp(2*I*c)*exp(2*I*d*x) + d)`**3.14.7 Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.00

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$\frac{4(dx + c)(iA + B)a^2 - 2(A - iB)a^2 \log(\tan(dx + c)^2 + 1) + 4(A - iB)a^2 \log(\tan(dx + c)) - \frac{2(-2)}{2d}}{2d}$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`output `-1/2*(4*(d*x + c)*(I*A + B)*a^2 - 2*(A - I*B)*a^2*log(tan(d*x + c)^2 + 1) + 4*(A - I*B)*a^2*log(tan(d*x + c)) - (2*(-2*I*A - B)*a^2*tan(d*x + c) - A*a^2)/tan(d*x + c)^2)/d`

3.14.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 186 vs. $2(82) = 164$.

Time = 1.08 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.98

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$Aa^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 8i Aa^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 4Ba^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 32(Aa^2 - iBa^2) \log\left(\tan\left(\frac{1}{2}\right.\right.$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output
$$-1/8*(A*a^2*\tan(1/2*d*x + 1/2*c)^2 - 8*I*A*a^2*\tan(1/2*d*x + 1/2*c) - 4*B*a^2*\tan(1/2*d*x + 1/2*c) - 32*(A*a^2 - I*B*a^2)*\log(\tan(1/2*d*x + 1/2*c) + I) + 16*(A*a^2 - I*B*a^2)*\log(\tan(1/2*d*x + 1/2*c)) - (24*A*a^2*\tan(1/2*d*x + 1/2*c)^2 - 24*I*B*a^2*\tan(1/2*d*x + 1/2*c)^2 - 8*I*A*a^2*\tan(1/2*d*x + 1/2*c) - 4*B*a^2*\tan(1/2*d*x + 1/2*c) - A*a^2)/\tan(1/2*d*x + 1/2*c)^2)/d$$

3.14.9 Mupad [B] (verification not implemented)

Time = 7.40 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.71

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= -\frac{\frac{Aa^2}{2} + \tan(c + dx)(Ba^2 + Aa^2 2i)}{d \tan(c + dx)^2} - \frac{4a^2 \operatorname{atan}(2 \tan(c + dx) + 1i)(B + A 1i)}{d}$$

input `int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2,x)`

output
$$-((A*a^2)/2 + \tan(c + d*x)*(A*a^2*2i + B*a^2))/(d*\tan(c + d*x)^2) - (4*a^2*\operatorname{atan}(2*\tan(c + d*x) + 1i)*(A*1i + B))/d$$

3.15 $\int \cot^4(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.15.1	Optimal result	393
3.15.2	Mathematica [A] (verified)	393
3.15.3	Rubi [A] (verified)	394
3.15.4	Maple [A] (verified)	397
3.15.5	Fricas [A] (verification not implemented)	398
3.15.6	Sympy [A] (verification not implemented)	399
3.15.7	Maxima [A] (verification not implemented)	399
3.15.8	Giac [B] (verification not implemented)	400
3.15.9	Mupad [B] (verification not implemented)	400

3.15.1 Optimal result

Integrand size = 34, antiderivative size = 117

$$\int \cot^4(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= 2a^2(A-iB)x + \frac{2a^2(A-iB) \cot(c+dx)}{d} - \frac{a^2(4iA+3B) \cot^2(c+dx)}{6d}$$

$$- \frac{2a^2(iA+B) \log(\sin(c+dx))}{d} - \frac{A \cot^3(c+dx) (a^2+ia^2 \tan(c+dx))}{3d}$$

```
output 2*a^2*(A-I*B)*x+2*a^2*(A-I*B)*cot(d*x+c)/d-1/6*a^2*(4*I*A+3*B)*cot(d*x+c)^2/d-2*a^2*(I*A+B)*ln(sin(d*x+c))/d-1/3*A*cot(d*x+c)^3*(a^2+I*a^2*tan(d*x+c))/d
```

3.15.2 Mathematica [A] (verified)

Time = 0.66 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.22

$$\int \cot^4(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= a^2 \left(\frac{2A \cot(c+dx)}{d} - \frac{2iB \cot(c+dx)}{d} - \frac{iA \cot^2(c+dx)}{d} - \frac{B \cot^2(c+dx)}{2d} \right.$$

$$- \frac{A \cot^3(c+dx)}{3d} - \frac{2iA \log(\tan(c+dx))}{d} - \frac{2B \log(\tan(c+dx))}{d}$$

$$\left. + \frac{2iA \log(i + \tan(c+dx))}{d} + \frac{2B \log(i + \tan(c+dx))}{d} \right)$$

input `Integrate[Cot[c + d*x]^4*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `a^2*((2*A*Cot[c + d*x])/d - ((2*I)*B*Cot[c + d*x])/d - (I*A*Cot[c + d*x]^2)/d - (B*Cot[c + d*x]^2)/(2*d) - (A*Cot[c + d*x]^3)/(3*d) - ((2*I)*A*Log[Tan[c + d*x]])/d - (2*B*Log[Tan[c + d*x]])/d + ((2*I)*A*Log[I + Tan[c + d*x]])/d + (2*B*Log[I + Tan[c + d*x]])/d)`

3.15.3 Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.08, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3042, 4076, 3042, 4074, 27, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^4(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\tan(c + dx)^4} dx \\
 & \quad \downarrow \text{4076} \\
 & \frac{1}{3} \int \cot^3(c + dx)(i \tan(c + dx)a + a)(a(4iA + 3B) - a(2A - 3iB) \tan(c + dx)) dx - \\
 & \quad \frac{A \cot^3(c + dx) (a^2 + ia^2 \tan(c + dx))}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \frac{(i \tan(c + dx)a + a)(a(4iA + 3B) - a(2A - 3iB) \tan(c + dx))}{\tan(c + dx)^3} dx - \\
 & \quad \frac{A \cot^3(c + dx) (a^2 + ia^2 \tan(c + dx))}{3d} \\
 & \quad \downarrow \text{4074} \\
 & \frac{1}{3} \left(\int -6 \cot^2(c + dx) ((A - iB)a^2 + (iA + B) \tan(c + dx)a^2) dx - \frac{a^2(3B + 4iA) \cot^2(c + dx)}{2d} \right) - \\
 & \quad \frac{A \cot^3(c + dx) (a^2 + ia^2 \tan(c + dx))}{3d} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.15. $\int \cot^4(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\frac{1}{3} \left(-6 \int \cot^2(c+dx) \left((A-iB)a^2 + (iA+B)\tan(c+dx)a^2 \right) dx - \frac{a^2(3B+4iA)\cot^2(c+dx)}{2d} \right) - \frac{A \cot^3(c+dx) (a^2 + ia^2 \tan(c+dx))}{3d}$$

↓ 3042

$$\frac{1}{3} \left(-6 \int \frac{(A-iB)a^2 + (iA+B)\tan(c+dx)a^2}{\tan(c+dx)^2} dx - \frac{a^2(3B+4iA)\cot^2(c+dx)}{2d} \right) - \frac{A \cot^3(c+dx) (a^2 + ia^2 \tan(c+dx))}{3d}$$

↓ 4012

$$\frac{1}{3} \left(-6 \left(\int \cot(c+dx) (a^2(iA+B) - a^2(A-iB)\tan(c+dx)) dx - \frac{a^2(A-iB)\cot(c+dx)}{d} \right) - \frac{a^2(3B+4iA)\cot^2(c+dx)}{2d} \right) - \frac{A \cot^3(c+dx) (a^2 + ia^2 \tan(c+dx))}{3d}$$

↓ 3042

$$\frac{1}{3} \left(-6 \left(\int \frac{a^2(iA+B) - a^2(A-iB)\tan(c+dx)}{\tan(c+dx)} dx - \frac{a^2(A-iB)\cot(c+dx)}{d} \right) - \frac{a^2(3B+4iA)\cot^2(c+dx)}{2d} \right) - \frac{A \cot^3(c+dx) (a^2 + ia^2 \tan(c+dx))}{3d}$$

↓ 4014

$$\frac{1}{3} \left(-6 \left(a^2(B+iA) \int \cot(c+dx) dx - \frac{a^2(A-iB)\cot(c+dx)}{d} - (a^2x(A-iB)) \right) - \frac{a^2(3B+4iA)\cot^2(c+dx)}{2d} \right) - \frac{A \cot^3(c+dx) (a^2 + ia^2 \tan(c+dx))}{3d}$$

↓ 3042

$$\frac{1}{3} \left(-6 \left(a^2(B+iA) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx - \frac{a^2(A-iB)\cot(c+dx)}{d} - (a^2x(A-iB)) \right) - \frac{a^2(3B+4iA)\cot^2(c+dx)}{2d} \right) - \frac{A \cot^3(c+dx) (a^2 + ia^2 \tan(c+dx))}{3d}$$

↓ 25

$$\frac{1}{3} \left(-6 \left(-a^2(B+iA) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx - \frac{a^2(A-iB)\cot(c+dx)}{d} - (a^2x(A-iB)) \right) - \frac{a^2(3B+4iA)\cot^2(c+dx)}{2d} \right) - \frac{A \cot^3(c+dx) (a^2 + ia^2 \tan(c+dx))}{3d}$$

↓ 3956

$$\frac{1}{3} \left(-\frac{a^2(3B + 4iA) \cot^2(c + dx)}{2d} - 6 \left(-\frac{a^2(A - iB) \cot(c + dx)}{d} + \frac{a^2(B + iA) \log(-\sin(c + dx))}{d} - (a^2x(A - iB) \cot^3(c + dx)) \right) \right) \frac{A \cot^3(c + dx) (a^2 + ia^2 \tan(c + dx))}{3d}$$

input `Int[Cot[c + d*x]^4*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(-1/2*(a^2*((4*I)*A + 3*B)*Cot[c + d*x]^2)/d - 6*(-(a^2*(A - I*B)*x) - (a^2*(A - I*B)*Cot[c + d*x])/d + (a^2*(I*A + B)*Log[-Sin[c + d*x]])/d)/3 - (A*Cot[c + d*x]^3*(a^2 + I*a^2*Tan[c + d*x]))/(3*d)`

3.15.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)
)*(x_)], x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

```
rule 4076 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

3.15.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.89

$$3.15. \quad \int \cot^4(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

method	result
parallelrisch	$\frac{2a^2 \left(\left(-\frac{iA}{2} - \frac{B}{2} \right) \ln(\sec^2(dx+c)) + (iA+B) \ln(\tan(dx+c)) + \frac{7A(\cot^3(dx+c))}{6} + (\cot^2(dx+c)) \left(\frac{iA}{2} + \frac{B}{4} \right) + (-A(\csc^2(dx+c))) \right)}{d}$
derivativedivides	$\frac{-A a^2 (-\cot(dx+c) - dx - c) - B a^2 \ln(\sin(dx+c)) + 2iA a^2 \left(-\frac{(\cot^2(dx+c))}{2} - \ln(\sin(dx+c)) \right) + 2iB a^2 (-\cot(dx+c) - dx - c)}{d}$
default	$\frac{-A a^2 (-\cot(dx+c) - dx - c) - B a^2 \ln(\sin(dx+c)) + 2iA a^2 \left(-\frac{(\cot^2(dx+c))}{2} - \ln(\sin(dx+c)) \right) + 2iB a^2 (-\cot(dx+c) - dx - c)}{d}$
risch	$\frac{4ia^2 Bc}{d} - \frac{4a^2 Ac}{d} + \frac{2a^2 (15iA e^{4i(dx+c)} + 9B e^{4i(dx+c)} - 18iA e^{2i(dx+c)} - 15B e^{2i(dx+c)} + 7iA + 6B)}{3d(e^{2i(dx+c)} - 1)^3} - \frac{2a^2 \ln(e^{2i(dx+c)})}{d}$
norman	$\frac{(-2iB a^2 + 2A a^2)x(\tan^3(dx+c)) - \frac{A a^2}{3d} + \frac{2(-iB a^2 + A a^2)(\tan^2(dx+c))}{d} - \frac{(2iA a^2 + B a^2)\tan(dx+c)}{2d}}{\tan(dx+c)^3} + \frac{(iA a^2 + B a^2) \ln(\tan(dx+c))}{d}$

input `int(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `-2*a^2*((-1/2*I*A-1/2*B)*ln(sec(d*x+c)^2)+(I*A+B)*ln(tan(d*x+c))+7/6*A*cot(d*x+c)^3+cot(d*x+c)^2*(1/2*I*A+1/4*B)+(-A*csc(d*x+c)^2+I*B)*cot(d*x+c)+x*d*(-A+I*B))/d`

3.15.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.55

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{2(3(-5iA - 3B)a^2 e^{(4i dx + 4i c)} + 3(6iA + 5B)a^2 e^{(2i dx + 2i c)} + (-7iA - 6B)a^2 + 3((iA + B)a^2 e^{(6i dx + 6i c)} - 3de^{(6i dx + 6i c)} - 3de^{(4i dx + 4i c)}))}{3(de^{(6i dx + 6i c)} - 3de^{(4i dx + 4i c)})}$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-2/3*(3*(-5*I*A - 3*B)*a^2*e^(4*I*d*x + 4*I*c) + 3*(6*I*A + 5*B)*a^2*e^(2*I*d*x + 2*I*c) + (-7*I*A - 6*B)*a^2 + 3*((I*A + B)*a^2*e^(6*I*d*x + 6*I*c) + 3*(-I*A - B)*a^2*e^(4*I*d*x + 4*I*c) + 3*(I*A + B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*log(e^(2*I*d*x + 2*I*c) - 1))/(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)`

3.15. $\int \cot^4(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.15.6 Sympy [A] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.56

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= -\frac{2ia^2(A - iB) \log(e^{2idx} - e^{-2ic})}{d} + \frac{14iAa^2 + 12Ba^2 + (-36iAa^2e^{2ic} - 30Ba^2e^{2ic})e^{2idx} + (30iAa^2e^{4ic} + 18Ba^2e^{4ic})e^{4idx}}{3de^{6ic}e^{6idx} - 9de^{4ic}e^{4idx} + 9de^{2ic}e^{2idx} - 3d}$$

input `integrate(cot(d*x+c)**4*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)), x)`

output `-2*I*a**2*(A - I*B)*log(exp(2*I*d*x) - exp(-2*I*c))/d + (14*I*A*a**2 + 12*B*a**2 + (-36*I*A*a**2*exp(2*I*c) - 30*B*a**2*exp(2*I*c))*exp(2*I*d*x) + (30*I*A*a**2*exp(4*I*c) + 18*B*a**2*exp(4*I*c))*exp(4*I*d*x))/(3*d*exp(6*I*c)*exp(6*I*d*x) - 9*d*exp(4*I*c)*exp(4*I*d*x) + 9*d*exp(2*I*c)*exp(2*I*d*x) - 3*d)`

3.15.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.95

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{12(dx + c)(A - iB)a^2 + 6(iA + B)a^2 \log(\tan(dx + c)^2 + 1) - 12(iA + B)a^2 \log(\tan(dx + c)) + \frac{12(A + B)a^2 \tan(dx + c)}{d}}{6d}$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)), x, algorithm="maxima")`

output `1/6*(12*(d*x + c)*(A - I*B)*a^2 + 6*(I*A + B)*a^2*log(tan(d*x + c)^2 + 1) - 12*(I*A + B)*a^2*log(tan(d*x + c)) + (12*(A - I*B)*a^2*tan(d*x + c)^2 + 3*(-2*I*A - B)*a^2*tan(d*x + c) - 2*A*a^2)/tan(d*x + c)^3/d`

3.15.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 255 vs. $2(103) = 206$.

Time = 1.34 (sec) , antiderivative size = 255, normalized size of antiderivative = 2.18

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{Aa^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 6i Aa^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 3Ba^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 27Aa^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 24A^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 24A^2 B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 24A^2 B^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 24A^2 B^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 96i(-Aa^2 - Ba^2) \log(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + i) + 48(-Aa^2 - Ba^2) \log(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)) - (-88iAa^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 88Ba^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 27Aa^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 24iB \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 6iAa^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 3Ba^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + Aa^2)/\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3}{d}$$

```
input integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")
```

```
output 1/24*(A*a^2*tan(1/2*d*x + 1/2*c)^3 - 6*I*A*a^2*tan(1/2*d*x + 1/2*c)^2 - 3*B*a^2*tan(1/2*d*x + 1/2*c)^2 - 27*A*a^2*tan(1/2*d*x + 1/2*c) + 24*I*B*a^2*tan(1/2*d*x + 1/2*c) - 96*(-I*A*a^2 - B*a^2)*log(tan(1/2*d*x + 1/2*c) + I) + 48*(-I*A*a^2 - B*a^2)*log(tan(1/2*d*x + 1/2*c)) - (-88*I*A*a^2*tan(1/2*d*x + 1/2*c)^3 - 88*B*a^2*tan(1/2*d*x + 1/2*c)^3 - 27*A*a^2*tan(1/2*d*x + 1/2*c)^2 + 24*I*B*a^2*tan(1/2*d*x + 1/2*c)^2 + 6*I*A*a^2*tan(1/2*d*x + 1/2*c) + 3*B*a^2*tan(1/2*d*x + 1/2*c) + A*a^2)/tan(1/2*d*x + 1/2*c)^3/d
```

3.15.9 Mupad [B] (verification not implemented)

Time = 8.04 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.79

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{\frac{Aa^2}{3} - \tan(c + dx)^2(2Aa^2 - Ba^2 2i) + \tan(c + dx)\left(\frac{Ba^2}{2} + Aa^2 1i\right)}{d \tan(c + dx)^3} - \frac{a^2 \operatorname{atan}(2 \tan(c + dx) + 1i)(B + A 1i) 4i}{d}$$

```
input int(cot(c + d*x)^4*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2,x)
```

```
output - ((A*a^2)/3 - tan(c + d*x)^2*(2*A*a^2 - B*a^2*2i) + tan(c + d*x)*(A*a^2*1i + (B*a^2)/2))/(d*tan(c + d*x)^3) - (a^2*atan(2*tan(c + d*x) + 1i)*(A*1i + B)*4i)/d
```

3.15. $\int \cot^4(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.16 $\int \cot^5(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.16.1	Optimal result	401
3.16.2	Mathematica [A] (verified)	402
3.16.3	Rubi [A] (verified)	402
3.16.4	Maple [A] (verified)	406
3.16.5	Fricas [A] (verification not implemented)	407
3.16.6	Sympy [A] (verification not implemented)	407
3.16.7	Maxima [A] (verification not implemented)	408
3.16.8	Giac [B] (verification not implemented)	408
3.16.9	Mupad [B] (verification not implemented)	409

3.16.1 Optimal result

Integrand size = 34, antiderivative size = 139

$$\int \cot^5(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= 2a^2(iA+B)x + \frac{2a^2(iA+B) \cot(c+dx)}{d}$$

$$+ \frac{a^2(A-iB) \cot^2(c+dx)}{d} - \frac{a^2(5iA+4B) \cot^3(c+dx)}{12d}$$

$$+ \frac{2a^2(A-iB) \log(\sin(c+dx))}{d} - \frac{A \cot^4(c+dx) (a^2+ia^2 \tan(c+dx))}{4d}$$

output

```
2*a^2*(I*A+B)*x+2*a^2*(I*A+B)*cot(d*x+c)/d+a^2*(A-I*B)*cot(d*x+c)^2/d-1/12
*a^2*(5*I*A+4*B)*cot(d*x+c)^3/d+2*a^2*(A-I*B)*ln(sin(d*x+c))/d-1/4*A*cot(d
*x+c)^4*(a^2+I*a^2*tan(d*x+c))/d
```


3.16.2 Mathematica [A] (verified)

Time = 1.13 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.25

$$\int \cot^5(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= a^2 \left(\frac{2iA \cot(c+dx)}{d} + \frac{2B \cot(c+dx)}{d} + \frac{A \cot^2(c+dx)}{d} - \frac{iB \cot^2(c+dx)}{d} \right.$$

$$\left. - \frac{2iA \cot^3(c+dx)}{3d} - \frac{B \cot^3(c+dx)}{3d} - \frac{A \cot^4(c+dx)}{4d} + \frac{2A \log(\tan(c+dx))}{d} \right.$$

$$\left. - \frac{2iB \log(\tan(c+dx))}{d} - \frac{2A \log(i + \tan(c+dx))}{d} + \frac{2iB \log(i + \tan(c+dx))}{d} \right)$$

input `Integrate[Cot[c + d*x]^5*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `a^2*(((2*I)*A*Cot[c + d*x])/d + (2*B*Cot[c + d*x])/d + (A*Cot[c + d*x]^2)/d - (I*B*Cot[c + d*x]^2)/d - (((2*I)/3)*A*Cot[c + d*x]^3)/d - (B*Cot[c + d*x]^3)/(3*d) - (A*Cot[c + d*x]^4)/(4*d) + (2*A*Log[Tan[c + d*x]])/d - ((2*I)*B*Log[Tan[c + d*x]])/d - (2*A*Log[I + Tan[c + d*x]])/d + ((2*I)*B*Log[I + Tan[c + d*x]])/d)`

3.16.3 Rubi [A] (verified)

Time = 0.94 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.09, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.441$, Rules used = {3042, 4076, 3042, 4074, 27, 3042, 4012, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^5(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan(c+dx)^5} dx$$

$$\downarrow \text{4076}$$

$$\frac{1}{4} \int \cot^4(c+dx)(i \tan(c+dx)a+a)(a(5iA+4B)-a(3A-4iB) \tan(c+dx)) dx - \frac{A \cot^4(c+dx)(a^2+ia^2 \tan(c+dx))}{4d}$$

3.16. $\int \cot^5(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{1}{4} \int \frac{(i \tan(c+dx)a+a)(a(5iA+4B)-a(3A-4iB)\tan(c+dx))}{\tan(c+dx)^4} dx - \\
& \quad \frac{A \cot^4(c+dx)(a^2+ia^2 \tan(c+dx))}{4d} \\
& \downarrow 4074 \\
& \frac{1}{4} \left(\int -8 \cot^3(c+dx) ((A-iB)a^2+(iA+B)\tan(c+dx)a^2) dx - \frac{a^2(4B+5iA)\cot^3(c+dx)}{3d} \right) - \\
& \quad \frac{A \cot^4(c+dx)(a^2+ia^2 \tan(c+dx))}{4d} \\
& \downarrow 27 \\
& \frac{1}{4} \left(-8 \int \cot^3(c+dx) ((A-iB)a^2+(iA+B)\tan(c+dx)a^2) dx - \frac{a^2(4B+5iA)\cot^3(c+dx)}{3d} \right) - \\
& \quad \frac{A \cot^4(c+dx)(a^2+ia^2 \tan(c+dx))}{4d} \\
& \downarrow 3042 \\
& \frac{1}{4} \left(-8 \int \frac{(A-iB)a^2+(iA+B)\tan(c+dx)a^2}{\tan(c+dx)^3} dx - \frac{a^2(4B+5iA)\cot^3(c+dx)}{3d} \right) - \\
& \quad \frac{A \cot^4(c+dx)(a^2+ia^2 \tan(c+dx))}{4d} \\
& \downarrow 4012 \\
& \frac{1}{4} \left(-8 \left(\int \cot^2(c+dx) (a^2(iA+B)-a^2(A-iB)\tan(c+dx)) dx - \frac{a^2(A-iB)\cot^2(c+dx)}{2d} \right) - \frac{a^2(4B+5iA)\cot^3(c+dx)}{3d} \right) - \\
& \quad \frac{A \cot^4(c+dx)(a^2+ia^2 \tan(c+dx))}{4d} \\
& \downarrow 3042 \\
& \frac{1}{4} \left(-8 \left(\int \frac{a^2(iA+B)-a^2(A-iB)\tan(c+dx)}{\tan(c+dx)^2} dx - \frac{a^2(A-iB)\cot^2(c+dx)}{2d} \right) - \frac{a^2(4B+5iA)\cot^3(c+dx)}{3d} \right) - \\
& \quad \frac{A \cot^4(c+dx)(a^2+ia^2 \tan(c+dx))}{4d} \\
& \downarrow 4012 \\
& \frac{1}{4} \left(-8 \left(\int -\cot(c+dx) ((A-iB)a^2+(iA+B)\tan(c+dx)a^2) dx - \frac{a^2(A-iB)\cot^2(c+dx)}{2d} - \frac{a^2(B+iA)\cot}{d} \right) - \right. \\
& \quad \left. \frac{A \cot^4(c+dx)(a^2+ia^2 \tan(c+dx))}{4d} \right)
\end{aligned}$$

3.16. $\int \cot^5(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

↓ 25

$$\frac{1}{4} \left(-8 \left(- \int \cot(c+dx) ((A-iB)a^2 + (iA+B)\tan(c+dx)a^2) dx - \frac{a^2(A-iB)\cot^2(c+dx)}{2d} - \frac{a^2(B+iA)\cot(c+dx)}{d} \right) - \frac{A \cot^4(c+dx) (a^2 + ia^2 \tan(c+dx))}{4d} \right)$$

↓ 3042

$$\frac{1}{4} \left(-8 \left(- \int \frac{(A-iB)a^2 + (iA+B)\tan(c+dx)a^2}{\tan(c+dx)} dx - \frac{a^2(A-iB)\cot^2(c+dx)}{2d} - \frac{a^2(B+iA)\cot(c+dx)}{d} \right) - \frac{A \cot^4(c+dx) (a^2 + ia^2 \tan(c+dx))}{4d} \right)$$

↓ 4014

$$\frac{1}{4} \left(-8 \left(-a^2(A-iB) \int \cot(c+dx) dx - \frac{a^2(A-iB)\cot^2(c+dx)}{2d} - \frac{a^2(B+iA)\cot(c+dx)}{d} - a^2x(B+iA) \right) - \frac{A \cot^4(c+dx) (a^2 + ia^2 \tan(c+dx))}{4d} \right)$$

↓ 3042

$$\frac{1}{4} \left(-8 \left(-a^2(A-iB) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx - \frac{a^2(A-iB)\cot^2(c+dx)}{2d} - \frac{a^2(B+iA)\cot(c+dx)}{d} - a^2x(B+iA) \right) - \frac{A \cot^4(c+dx) (a^2 + ia^2 \tan(c+dx))}{4d} \right)$$

↓ 25

$$\frac{1}{4} \left(-8 \left(a^2(A-iB) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx - \frac{a^2(A-iB)\cot^2(c+dx)}{2d} - \frac{a^2(B+iA)\cot(c+dx)}{d} - a^2x(B+iA) \right) - \frac{A \cot^4(c+dx) (a^2 + ia^2 \tan(c+dx))}{4d} \right)$$

↓ 3956

$$\frac{1}{4} \left(-\frac{a^2(4B+5iA)\cot^3(c+dx)}{3d} - 8 \left(-\frac{a^2(A-iB)\cot^2(c+dx)}{2d} - \frac{a^2(B+iA)\cot(c+dx)}{d} - \frac{a^2(A-iB)\log(-s)}{d} \right) - \frac{A \cot^4(c+dx) (a^2 + ia^2 \tan(c+dx))}{4d} \right)$$

input `Int[Cot[c + d*x]^5*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

```
output (-1/3*(a^2*((5*I)*A + 4*B)*Cot[c + d*x]^3)/d - 8*(-(a^2*(I*A + B)*x) - (a^
2*(I*A + B)*Cot[c + d*x])/d - (a^2*(A - I*B)*Cot[c + d*x]^2)/(2*d) - (a^2*
(A - I*B)*Log[-Sin[c + d*x]])/d)/4 - (A*Cot[c + d*x]^4*(a^2 + I*a^2*Tan[c
+ d*x]))/(4*d)
```

3.16.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3956 Int[tan[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4012 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]
```

```
rule 4014 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

```
rule 4076 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

3.16.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.78

method	result
parallelrisch	$2a^2 \left(\left(-\frac{A}{2} + \frac{iB}{2} \right) \ln(\sec^2(dx+c)) + (-iB+A) \ln(\tan(dx+c)) - \frac{A(\cot^4(dx+c))}{8} + (\cot^3(dx+c)) \left(-\frac{iA}{3} - \frac{B}{6} \right) + (\cot^2(dx+c)) \right)$
derivativedivides	$-A a^2 \left(-\frac{(\cot^2(dx+c))}{2} - \ln(\sin(dx+c)) \right) - B a^2 (-\cot(dx+c) - dx - c) + 2iA a^2 \left(-\frac{(\cot^3(dx+c))}{3} + \cot(dx+c) + dx + c \right) + 2a^2$
default	$-A a^2 \left(-\frac{(\cot^2(dx+c))}{2} - \ln(\sin(dx+c)) \right) - B a^2 (-\cot(dx+c) - dx - c) + 2iA a^2 \left(-\frac{(\cot^3(dx+c))}{3} + \cot(dx+c) + dx + c \right) + 2a^2$
risch	$-\frac{4a^2 Bc}{d} - \frac{4ia^2 Ac}{d} + \frac{2ia^2 (21iA e^{6i(dx+c)} + 15B e^{6i(dx+c)} - 36iA e^{4i(dx+c)} - 33B e^{4i(dx+c)} + 29iA e^{2i(dx+c)} + 25B e^{2i(dx+c)} - 15B)}{3d(e^{2i(dx+c)} - 1)^4}$
norman	$\frac{(-iB a^2 + A a^2)(\tan^2(dx+c))}{d} + (2iA a^2 + 2B a^2)x(\tan^4(dx+c)) - \frac{A a^2}{4d} + \frac{2(iA a^2 + B a^2)(\tan^3(dx+c))}{d} - \frac{(2iA a^2 + B a^2)\tan(dx+c)}{3d}$

```
input int(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

3.16. $\int \cot^5(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

output $2*a^2*((-1/2*A+1/2*I*B)*\ln(\sec(d*x+c)^2)+(A-I*B)*\ln(\tan(d*x+c))-1/8*A*\cot(d*x+c)^4+\cot(d*x+c)^3*(-1/3*I*A-1/6*B)+\cot(d*x+c)^2*(1/2*A-1/2*I*B)+\cot(d*x+c)*(I*A+B)+(I*A+B)*x*d)/d$

3.16.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.63

$$\int \cot^5(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx = \frac{2(3(7A-5iB)a^2e^{(6i dx+6i c)} - 3(12A-11iB)a^2e^{(4i dx+4i c)} + (29A-25iB)a^2e^{(2i dx+2i c)} - (8A-7iB)a^2e^{(i dx+i c)} + 3A^2e^{(i dx+i c)})}{3(de^{(8i dx+8i c)} - 1)}$$

input `integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output $-2/3*(3*(7*A - 5*I*B)*a^2*e^{(6*I*d*x + 6*I*c)} - 3*(12*A - 11*I*B)*a^2*e^{(4*I*d*x + 4*I*c)} + (29*A - 25*I*B)*a^2*e^{(2*I*d*x + 2*I*c)} - (8*A - 7*I*B)*a^2 - 3*((A - I*B)*a^2*e^{(8*I*d*x + 8*I*c)} - 4*(A - I*B)*a^2*e^{(6*I*d*x + 6*I*c)} + 6*(A - I*B)*a^2*e^{(4*I*d*x + 4*I*c)} - 4*(A - I*B)*a^2*e^{(2*I*d*x + 2*I*c)} + (A - I*B)*a^2)*\log(e^{(2*I*d*x + 2*I*c)} - 1)/(d*e^{(8*I*d*x + 8*I*c)} - 4*d*e^{(6*I*d*x + 6*I*c)} + 6*d*e^{(4*I*d*x + 4*I*c)} - 4*d*e^{(2*I*d*x + 2*I*c)} + d)$

3.16.6 Sympy [A] (verification not implemented)

Time = 0.54 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.69

$$\int \cot^5(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx = \frac{2a^2(A-iB) \log(e^{2idx} - e^{-2ic})}{d} + \frac{16Aa^2 - 14iBa^2 + (-58Aa^2e^{2ic} + 50iBa^2e^{2ic})e^{2idx} + (72Aa^2e^{4ic} - 66iBa^2e^{4ic})e^{4idx} + (-42Aa^2e^{6ic} + 36iBa^2e^{6ic})e^{6idx} + 3A^2e^{8ic}}{3de^{8ic}e^{8idx} - 12de^{6ic}e^{6idx} + 18de^{4ic}e^{4idx} - 12de^{2ic}e^{2idx} + 3d}$$

input `integrate(cot(d*x+c)**5*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output $2*a**2*(A - I*B)*\log(\exp(2*I*d*x) - \exp(-2*I*c))/d + (16*A*a**2 - 14*I*B*a**2 + (-58*A*a**2*\exp(2*I*c) + 50*I*B*a**2*\exp(2*I*c))*\exp(2*I*d*x) + (72*A*a**2*\exp(4*I*c) - 66*I*B*a**2*\exp(4*I*c))*\exp(4*I*d*x) + (-42*A*a**2*\exp(6*I*c) + 30*I*B*a**2*\exp(6*I*c))*\exp(6*I*d*x))/(3*d*\exp(8*I*c)*\exp(8*I*d*x) - 12*d*\exp(6*I*c)*\exp(6*I*d*x) + 18*d*\exp(4*I*c)*\exp(4*I*d*x) - 12*d*\exp(2*I*c)*\exp(2*I*d*x) + 3*d)$

3.16.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.95

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{24(dx + c)(-iA - B)a^2 + 12(A - iB)a^2 \log(\tan(dx + c)^2 + 1) - 24(A - iB)a^2 \log(\tan(dx + c))}{12d}$$

input `integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output $-1/12*(24*(d*x + c)*(-I*A - B)*a^2 + 12*(A - I*B)*a^2*\log(\tan(d*x + c)^2 + 1) - 24*(A - I*B)*a^2*\log(\tan(d*x + c)) - (24*(I*A + B)*a^2*\tan(d*x + c)^3 + 12*(A - I*B)*a^2*\tan(d*x + c)^2 + 4*(-2*I*A - B)*a^2*\tan(d*x + c) - 3*A*a^2)/\tan(d*x + c)^4)/d$

3.16.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 322 vs. $2(123) = 246$.

Time = 1.10 (sec) , antiderivative size = 322, normalized size of antiderivative = 2.32

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{3Aa^2 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^4 - 16iAa^2 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 8Ba^2 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 60Aa^2 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)}{12d}$$

input `integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

3.16. $\int \cot^5(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

output
$$\begin{aligned} & -1/192*(3*A*a^2*\tan(1/2*d*x + 1/2*c)^4 - 16*I*A*a^2*\tan(1/2*d*x + 1/2*c)^3 \\ & - 8*B*a^2*\tan(1/2*d*x + 1/2*c)^3 - 60*A*a^2*\tan(1/2*d*x + 1/2*c)^2 + 48*I \\ & *B*a^2*\tan(1/2*d*x + 1/2*c)^2 + 240*I*A*a^2*\tan(1/2*d*x + 1/2*c) + 216*B*a \\ & ^2*\tan(1/2*d*x + 1/2*c) + 768*(A*a^2 - I*B*a^2)*\log(\tan(1/2*d*x + 1/2*c) + \\ & I) - 384*(A*a^2 - I*B*a^2)*\log(\tan(1/2*d*x + 1/2*c)) + (800*A*a^2*\tan(1/2 \\ & *d*x + 1/2*c)^4 - 800*I*B*a^2*\tan(1/2*d*x + 1/2*c)^4 - 240*I*A*a^2*\tan(1/2 \\ & *d*x + 1/2*c)^3 - 216*B*a^2*\tan(1/2*d*x + 1/2*c)^3 - 60*A*a^2*\tan(1/2*d*x \\ & + 1/2*c)^2 + 48*I*B*a^2*\tan(1/2*d*x + 1/2*c)^2 + 16*I*A*a^2*\tan(1/2*d*x + \\ & 1/2*c) + 8*B*a^2*\tan(1/2*d*x + 1/2*c) + 3*A*a^2)/\tan(1/2*d*x + 1/2*c)^4)/d \end{aligned}$$

3.16.9 Mupad [B] (verification not implemented)

Time = 7.92 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.81

$$\begin{aligned} & \int \cot^5(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\ & = \frac{\tan(c + dx)^2 (Aa^2 - Ba^2 li) + \tan(c + dx)^3 (2Ba^2 + Aa^2 2i) - \frac{Aa^2}{4} - \tan(c + dx) \left(\frac{Ba^2}{3} + \frac{Aa^2 2i}{3} \right)}{d \tan(c + dx)^4} \\ & + \frac{4a^2 \operatorname{atan}(2 \tan(c + dx) + li) (B + A li)}{d} \end{aligned}$$

input `int(cot(c + d*x)^5*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2,x)`

output
$$\begin{aligned} & (\tan(c + d*x)^2*(A*a^2 - B*a^2*1i) + \tan(c + d*x)^3*(A*a^2*2i + 2*B*a^2) - \\ & (A*a^2)/4 - \tan(c + d*x)*((A*a^2*2i)/3 + (B*a^2)/3))/d*\tan(c + d*x)^4 + \\ & (4*a^2*\operatorname{atan}(2*\tan(c + d*x) + 1i)*(A*1i + B))/d \end{aligned}$$

3.17 $\int \tan^2(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.17.1	Optimal result	410
3.17.2	Mathematica [A] (verified)	411
3.17.3	Rubi [A] (verified)	411
3.17.4	Maple [A] (verified)	414
3.17.5	Fricas [A] (verification not implemented)	415
3.17.6	Sympy [A] (verification not implemented)	416
3.17.7	Maxima [A] (verification not implemented)	416
3.17.8	Giac [B] (verification not implemented)	417
3.17.9	Mupad [B] (verification not implemented)	418

3.17.1 Optimal result

Integrand size = 34, antiderivative size = 182

$$\begin{aligned} & \int \tan^2(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx \\ &= -4a^3(A-iB)x + \frac{4a^3(iA+B) \log(\cos(c+dx))}{d} \\ & \quad + \frac{4a^3(A-iB) \tan(c+dx)}{d} + \frac{2a^3(iA+B) \tan^2(c+dx)}{d} \\ & \quad - \frac{a^3(45A-47iB) \tan^3(c+dx)}{60d} + \frac{iaB \tan^3(c+dx)(a+ia \tan(c+dx))^2}{5d} \\ & \quad - \frac{(5A-7iB) \tan^3(c+dx)(a^3+ia^3 \tan(c+dx))}{20d} \end{aligned}$$

output

```
-4*a^3*(A-I*B)*x+4*a^3*(I*A+B)*ln(cos(d*x+c))/d+4*a^3*(A-I*B)*tan(d*x+c)/d
+2*a^3*(I*A+B)*tan(d*x+c)^2/d-1/60*a^3*(45*A-47*I*B)*tan(d*x+c)^3/d+1/5*I*
a*B*tan(d*x+c)^3*(a+I*a*tan(d*x+c))^2/d-1/20*(5*A-7*I*B)*tan(d*x+c)^3*(a^3
+I*a^3*tan(d*x+c))/d
```

3.17.2 Mathematica [A] (verified)

Time = 1.59 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.68

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{a^3(-15iA - 17B - 240i(A - iB) \log(i + \tan(c + dx)) + 240(A - iB) \tan(c + dx) + 120(iA + B) \tan^2(c + dx))}{60d}$$

input `Integrate[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(a^3*((-15*I)*A - 17*B - (240*I)*(A - I*B)*Log[I + Tan[c + d*x]] + 240*(A - I*B)*Tan[c + d*x] + 120*(I*A + B)*Tan[c + d*x]^2 + (-60*A + (80*I)*B)*Tan[c + d*x]^3 - (15*I)*(A - (3*I)*B)*Tan[c + d*x]^4 - (12*I)*B*Tan[c + d*x]^5))/(60*d)`

3.17.3 Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.05, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3042, 4077, 3042, 4077, 3042, 4075, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^2(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4077}$$

$$\frac{1}{5} \int \tan^2(c + dx)(i \tan(c + dx)a + a)^2(a(5A - 3iB) + a(5iA + 7B) \tan(c + dx)) dx + \frac{iaB \tan^3(c + dx)(a + ia \tan(c + dx))^2}{5d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{5} \int \tan(c + dx)^2(i \tan(c + dx)a + a)^2(a(5A - 3iB) + a(5iA + 7B) \tan(c + dx)) dx + \frac{iaB \tan^3(c + dx)(a + ia \tan(c + dx))^2}{5d}$$

3.17. $\int \tan^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\downarrow 4077$$

$$\frac{1}{5} \left(\frac{1}{4} \int \tan^2(c+dx) (i \tan(c+dx)a + a) ((35A - 33iB)a^2 + (45iA + 47B) \tan(c+dx)a^2) dx - \frac{(5A - 7iB) \tan^3(c+dx)}{5d} \right) + \frac{iaB \tan^3(c+dx)(a + ia \tan(c+dx))^2}{5d}$$

$$\downarrow 3042$$

$$\frac{1}{5} \left(\frac{1}{4} \int \tan(c+dx)^2 (i \tan(c+dx)a + a) ((35A - 33iB)a^2 + (45iA + 47B) \tan(c+dx)a^2) dx - \frac{(5A - 7iB) \tan^3(c+dx)}{5d} \right) + \frac{iaB \tan^3(c+dx)(a + ia \tan(c+dx))^2}{5d}$$

$$\downarrow 4075$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\int \tan^2(c+dx) (80(A - iB)a^3 + 80(iA + B) \tan(c+dx)a^3) dx - \frac{a^3(45A - 47iB) \tan^3(c+dx)}{3d} \right) - \frac{(5A - 7iB) \tan^3(c+dx)}{5d} \right) + \frac{iaB \tan^3(c+dx)(a + ia \tan(c+dx))^2}{5d}$$

$$\downarrow 3042$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\int \tan(c+dx)^2 (80(A - iB)a^3 + 80(iA + B) \tan(c+dx)a^3) dx - \frac{a^3(45A - 47iB) \tan^3(c+dx)}{3d} \right) - \frac{(5A - 7iB) \tan^3(c+dx)}{5d} \right) + \frac{iaB \tan^3(c+dx)(a + ia \tan(c+dx))^2}{5d}$$

$$\downarrow 4011$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\int \tan(c+dx) (80a^3(A - iB) \tan(c+dx) - 80a^3(iA + B)) dx - \frac{a^3(45A - 47iB) \tan^3(c+dx)}{3d} + \frac{40a^3(B - iA) \tan^2(c+dx)}{3d} \right) - \frac{(5A - 7iB) \tan^3(c+dx)}{5d} \right) + \frac{iaB \tan^3(c+dx)(a + ia \tan(c+dx))^2}{5d}$$

$$\downarrow 3042$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\int \tan(c+dx) (80a^3(A - iB) \tan(c+dx) - 80a^3(iA + B)) dx - \frac{a^3(45A - 47iB) \tan^3(c+dx)}{3d} + \frac{40a^3(B - iA) \tan^2(c+dx)}{3d} \right) - \frac{(5A - 7iB) \tan^3(c+dx)}{5d} \right) + \frac{iaB \tan^3(c+dx)(a + ia \tan(c+dx))^2}{5d}$$

$$\downarrow 4008$$

$$3.17. \quad \int \tan^2(c+dx)(a + ia \tan(c+dx))^3(A + B \tan(c+dx)) dx$$

$$\frac{1}{5} \left(\frac{1}{4} \left(-80a^3(B + iA) \int \tan(c + dx) dx - \frac{a^3(45A - 47iB) \tan^3(c + dx)}{3d} + \frac{40a^3(B + iA) \tan^2(c + dx)}{d} + \frac{80a^3(A - iB) \tan(c + dx)}{d} + \frac{80a^3(B + iA)}{d} \right) + \frac{iaB \tan^3(c + dx)(a + ia \tan(c + dx))^2}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(-80a^3(B + iA) \int \tan(c + dx) dx - \frac{a^3(45A - 47iB) \tan^3(c + dx)}{3d} + \frac{40a^3(B + iA) \tan^2(c + dx)}{d} + \frac{80a^3(A - iB) \tan(c + dx)}{d} + \frac{80a^3(B + iA)}{d} \right) + \frac{iaB \tan^3(c + dx)(a + ia \tan(c + dx))^2}{5d} \right)$$

↓ 3956

$$\frac{1}{5} \left(\frac{1}{4} \left(-\frac{a^3(45A - 47iB) \tan^3(c + dx)}{3d} + \frac{40a^3(B + iA) \tan^2(c + dx)}{d} + \frac{80a^3(A - iB) \tan(c + dx)}{d} + \frac{80a^3(B + iA)}{d} \right) + \frac{iaB \tan^3(c + dx)(a + ia \tan(c + dx))^2}{5d} \right)$$

input `Int[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]), x]`

output `((I/5)*a*B*Tan[c + d*x]^3*(a + I*a*Tan[c + d*x])^2)/d + (-1/4*((5*A - (7*I)*B)*Tan[c + d*x]^3*(a^3 + I*a^3*Tan[c + d*x]))/d + (-80*a^3*(A - I*B)*x + (80*a^3*(I*A + B)*Log[Cos[c + d*x]])/d + (80*a^3*(A - I*B)*Tan[c + d*x])/d + (40*a^3*(I*A + B)*Tan[c + d*x]^2)/d - (a^3*(45*A - (47*I)*B)*Tan[c + d*x]^3)/(3*d))/4/5`

3.17.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4011 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4077 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.17.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.80

$$3.17. \quad \int \tan^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

method	result
derivativedivides	$\frac{a^3 \left(-\frac{iB(\tan^5(dx+c))}{5} - \frac{iA(\tan^4(dx+c))}{4} + \frac{4iB(\tan^3(dx+c))}{3} - \frac{3B(\tan^4(dx+c))}{4} + 2iA(\tan^2(dx+c)) - A(\tan^3(dx+c)) - 4iB \right)}{d}$
default	$\frac{a^3 \left(-\frac{iB(\tan^5(dx+c))}{5} - \frac{iA(\tan^4(dx+c))}{4} + \frac{4iB(\tan^3(dx+c))}{3} - \frac{3B(\tan^4(dx+c))}{4} + 2iA(\tan^2(dx+c)) - A(\tan^3(dx+c)) - 4iB \right)}{d}$
norman	$(4iB a^3 - 4A a^3) x - \frac{(iA a^3 + 3B a^3)(\tan^4(dx+c))}{4d} - \frac{(-4iB a^3 + 3A a^3)(\tan^3(dx+c))}{3d} + \frac{4(-iB a^3 + A a^3) \tan^2(dx+c)}{d}$
parallelrisch	$- \frac{12iB a^3(\tan^5(dx+c)) + 15iA(\tan^4(dx+c))a^3 - 80iB(\tan^3(dx+c))a^3 + 45B(\tan^4(dx+c))a^3 - 120iA(\tan^2(dx+c))a^3 + 69iA^2(\tan^2(dx+c))a^3}{15d(e^{2i(dx+c)}+1)^5}$
risch	$- \frac{8ia^3Bc}{d} + \frac{8a^3Ac}{d} + \frac{2a^3(180iA e^{8i(dx+c)} + 240B e^{8i(dx+c)} + 525iA e^{6i(dx+c)} + 585B e^{6i(dx+c)} + 615iA e^{4i(dx+c)} + 69iA^2 e^{4i(dx+c)})}{15d(e^{2i(dx+c)}+1)^5}$
parts	$\frac{(-iA a^3 - 3B a^3) \left(\frac{(\tan^4(dx+c))}{4} - \frac{(\tan^2(dx+c))}{2} + \frac{\ln(1+\tan^2(dx+c))}{2} \right)}{d} + \frac{(3iA a^3 + B a^3) \left(\frac{(\tan^2(dx+c))}{2} - \frac{\ln(1+\tan^2(dx+c))}{2} \right)}{d}$

input `int(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*a^3*(-1/5*I*B*tan(d*x+c)^5-1/4*I*A*tan(d*x+c)^4+4/3*I*B*tan(d*x+c)^3-3/4*B*tan(d*x+c)^4+2*I*A*tan(d*x+c)^2-A*tan(d*x+c)^3-4*I*B*tan(d*x+c)+2*B*tan(d*x+c)^2+4*A*tan(d*x+c)+1/2*(-4*I*A-4*B)*ln(1+tan(d*x+c)^2)+(4*I*B-4*A)*arctan(tan(d*x+c))`

3.17.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.60

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{2(60(-3iA - 4B)a^3 e^{(8i dx + 8i c)} + 15(-35iA - 39B)a^3 e^{(6i dx + 6i c)} + 5(-123iA - 139B)a^3 e^{(4i dx + 4i c)})}{15d(e^{2i(dx+c)}+1)^5}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output
$$\frac{-2/15*(60*(-3*I*A - 4*B))*a^3*e^{(8*I*d*x + 8*I*c)} + 15*(-35*I*A - 39*B)*a^3*e^{(6*I*d*x + 6*I*c)} + 5*(-123*I*A - 139*B)*a^3*e^{(4*I*d*x + 4*I*c)} + 5*(-69*I*A - 77*B)*a^3*e^{(2*I*d*x + 2*I*c)} + (-75*I*A - 83*B)*a^3 + 30*((-I*A - B)*a^3*e^{(10*I*d*x + 10*I*c)} + 5*(-I*A - B)*a^3*e^{(8*I*d*x + 8*I*c)} + 10*(-I*A - B)*a^3*e^{(6*I*d*x + 6*I*c)} + 10*(-I*A - B)*a^3*e^{(4*I*d*x + 4*I*c)} + 5*(-I*A - B)*a^3*e^{(2*I*d*x + 2*I*c)} + (-I*A - B)*a^3)*\log(e^{(2*I*d*x + 2*I*c)} + 1))/(d*e^{(10*I*d*x + 10*I*c)} + 5*d*e^{(8*I*d*x + 8*I*c)} + 10*d*e^{(6*I*d*x + 6*I*c)} + 10*d*e^{(4*I*d*x + 4*I*c)} + 5*d*e^{(2*I*d*x + 2*I*c)} + d)$$

3.17.6 Sympy [A] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 292, normalized size of antiderivative = 1.60

$$\int \tan^2(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx = \frac{4ia^3(A-iB) \log(e^{2idx} + e^{-2ic})}{d} + \frac{150iAa^3 + 166Ba^3 + (690iAa^3e^{2ic} + 770Ba^3e^{2ic})e^{2idx} + (1230iAa^3e^{4ic} + 1390Ba^3e^{4ic})e^{4idx} + (1050iAa^3e^{6ic} + 1170Ba^3e^{6ic})e^{6idx} + (360iAa^3e^{8ic} + 480Ba^3e^{8ic})e^{8idx} + 150d^2e^{10ic}}{15de^{10ic}e^{10idx} + 75de^{8ic}e^{8idx} + 150de^{6ic}e^{6idx} + 150de^{4ic}e^{4idx} + 150de^{2ic}e^{2idx} + 15d}$$

input `integrate(tan(d*x+c)**2*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)), x)`

output
$$4*I*a**3*(A - I*B)*\log(\exp(2*I*d*x) + \exp(-2*I*c))/d + (150*I*A*a**3 + 166*B*a**3 + (690*I*A*a**3*\exp(2*I*c) + 770*B*a**3*\exp(2*I*c))*\exp(2*I*d*x) + (1230*I*A*a**3*\exp(4*I*c) + 1390*B*a**3*\exp(4*I*c))*\exp(4*I*d*x) + (1050*I*A*a**3*\exp(6*I*c) + 1170*B*a**3*\exp(6*I*c))*\exp(6*I*d*x) + (360*I*A*a**3*\exp(8*I*c) + 480*B*a**3*\exp(8*I*c))*\exp(8*I*d*x))/(15*d*\exp(10*I*c)*\exp(10*I*d*x) + 75*d*\exp(8*I*c)*\exp(8*I*d*x) + 150*d*\exp(6*I*c)*\exp(6*I*d*x) + 150*d*\exp(4*I*c)*\exp(4*I*d*x) + 75*d*\exp(2*I*c)*\exp(2*I*d*x) + 15*d)$$

3.17.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.73

$$\int \tan^2(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx = \frac{12iBa^3 \tan(dx+c)^5 + 15(iA+3B)a^3 \tan(dx+c)^4 + 20(3A-4iB)a^3 \tan(dx+c)^3 + 120(-iA-3B)a^3 \tan(dx+c)^2 + 60(3A-4iB)a^3 \tan(dx+c) + 60Aa^3}{15d \tan^2(dx+c) + 15d}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output
$$\frac{-1/60*(12*I*B*a^3*\tan(d*x + c)^5 + 15*(I*A + 3*B)*a^3*\tan(d*x + c)^4 + 20*(3*A - 4*I*B)*a^3*\tan(d*x + c)^3 + 120*(-I*A - B)*a^3*\tan(d*x + c)^2 + 240*(d*x + c)*(A - I*B)*a^3 + 120*(I*A + B)*a^3*\log(\tan(d*x + c)^2 + 1) - 240*(A - I*B)*a^3*\tan(d*x + c))/d$$

3.17.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 504 vs. $2(158) = 316$.

Time = 0.68 (sec) , antiderivative size = 504, normalized size of antiderivative = 2.77

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{2(-30i Aa^3 e^{(10i dx + 10i c)} \log(e^{(2i dx + 2i c)} + 1) - 30 B a^3 e^{(10i dx + 10i c)} \log(e^{(2i dx + 2i c)} + 1) - 150i A a^3 e^{(8i dx + 8i c)} \log(e^{(2i dx + 2i c)} + 1) - 150i B a^3 e^{(8i dx + 8i c)} \log(e^{(2i dx + 2i c)} + 1) - 300i A a^3 e^{(6i dx + 6i c)} \log(e^{(2i dx + 2i c)} + 1) - 300i B a^3 e^{(6i dx + 6i c)} \log(e^{(2i dx + 2i c)} + 1) - 300i A a^3 e^{(4i dx + 4i c)} \log(e^{(2i dx + 2i c)} + 1) - 300i B a^3 e^{(4i dx + 4i c)} \log(e^{(2i dx + 2i c)} + 1) - 150i A a^3 e^{(2i dx + 2i c)} \log(e^{(2i dx + 2i c)} + 1) - 150i B a^3 e^{(2i dx + 2i c)} \log(e^{(2i dx + 2i c)} + 1) - 180i A a^3 e^{(8i dx + 8i c)} - 240i B a^3 e^{(8i dx + 8i c)} - 525i A a^3 e^{(6i dx + 6i c)} - 585i B a^3 e^{(6i dx + 6i c)} - 615i A a^3 e^{(4i dx + 4i c)} - 695i B a^3 e^{(4i dx + 4i c)} - 345i A a^3 e^{(2i dx + 2i c)} - 385i B a^3 e^{(2i dx + 2i c)} - 30i A a^3 \log(e^{(2i dx + 2i c)} + 1) - 30i B a^3 \log(e^{(2i dx + 2i c)} + 1) - 75i A a^3 - 83i B a^3)/(d e^{(10i dx + 10i c)} + 5d e^{(8i dx + 8i c)} + 10d e^{(6i dx + 6i c)} + 10d e^{(4i dx + 4i c)} + 5d e^{(2i dx + 2i c)} + d)$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output
$$\frac{-2/15*(-30*I*A*a^3*e^{(10*I*d*x + 10*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 30*B*a^3*e^{(10*I*d*x + 10*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 150*I*A*a^3*e^{(8*I*d*x + 8*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 150*B*a^3*e^{(8*I*d*x + 8*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 300*I*A*a^3*e^{(6*I*d*x + 6*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 300*B*a^3*e^{(6*I*d*x + 6*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 300*I*A*a^3*e^{(4*I*d*x + 4*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 300*B*a^3*e^{(4*I*d*x + 4*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 150*I*A*a^3*e^{(2*I*d*x + 2*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 150*B*a^3*e^{(2*I*d*x + 2*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 180*I*A*a^3*e^{(8*I*d*x + 8*I*c)} - 240*B*a^3*e^{(8*I*d*x + 8*I*c)} - 525*I*A*a^3*e^{(6*I*d*x + 6*I*c)} - 585*B*a^3*e^{(6*I*d*x + 6*I*c)} - 615*I*A*a^3*e^{(4*I*d*x + 4*I*c)} - 695*B*a^3*e^{(4*I*d*x + 4*I*c)} - 345*I*A*a^3*e^{(2*I*d*x + 2*I*c)} - 385*B*a^3*e^{(2*I*d*x + 2*I*c)} - 30*I*A*a^3*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 30*B*a^3*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 75*I*A*a^3 - 83*B*a^3)/(d*e^{(10*I*d*x + 10*I*c)} + 5*d*e^{(8*I*d*x + 8*I*c)} + 10*d*e^{(6*I*d*x + 6*I*c)} + 10*d*e^{(4*I*d*x + 4*I*c)} + 5*d*e^{(2*I*d*x + 2*I*c)} + d)$$

3.17. $\int \tan^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.17.9 Mupad [B] (verification not implemented)

Time = 7.27 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.26

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{\tan(c + dx)^3 \left(\frac{Ba^3 1i}{3} - \frac{a^3(2A - B 1i)}{3} + \frac{a^3(2B + A 1i) 1i}{3} \right)}{d} + \frac{\tan(c + dx) (Aa^3 - Ba^3 1i + a^3(2A - B 1i) - a^3(2B + A 1i) 1i)}{d}$$

$$- \frac{\tan(c + dx)^4 \left(\frac{Ba^3}{4} + \frac{a^3(2B + A 1i)}{4} \right) \ln(\tan(c + dx) + 1i) (4Ba^3 + Aa^3 4i)}{d} + \frac{\tan(c + dx)^2 \left(\frac{Aa^3 1i}{2} + \frac{Ba^3}{2} + \frac{a^3(2A - B 1i) 1i}{2} + \frac{a^3(2B + A 1i)}{2} \right)}{d} - \frac{Ba^3 \tan(c + dx)^5 1i}{5d}$$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3,x)`output `(tan(c + d*x)^3*((B*a^3*1i)/3 - (a^3*(2*A - B*1i))/3 + (a^3*(A*1i + 2*B)*1i)/3))/d + (tan(c + d*x)*(A*a^3 - B*a^3*1i + a^3*(2*A - B*1i) - a^3*(A*1i + 2*B)*1i))/d - (tan(c + d*x)^4*((B*a^3)/4 + (a^3*(A*1i + 2*B))/4))/d - (log(tan(c + d*x) + 1i)*(A*a^3*4i + 4*B*a^3))/d + (tan(c + d*x)^2*((A*a^3*1i)/2 + (B*a^3)/2 + (a^3*(2*A - B*1i)*1i)/2 + (a^3*(A*1i + 2*B))/2))/d - (B*a^3*tan(c + d*x)^5*1i)/(5*d)`

3.18 $\int \tan(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.18.1	Optimal result	419
3.18.2	Mathematica [A] (verified)	419
3.18.3	Rubi [A] (verified)	420
3.18.4	Maple [A] (verified)	423
3.18.5	Fricas [A] (verification not implemented)	423
3.18.6	Sympy [B] (verification not implemented)	424
3.18.7	Maxima [A] (verification not implemented)	424
3.18.8	Giac [B] (verification not implemented)	425
3.18.9	Mupad [B] (verification not implemented)	426

3.18.1 Optimal result

Integrand size = 32, antiderivative size = 138

$$\int \tan(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= -4a^3(iA+B)x - \frac{4a^3(A-iB) \log(\cos(c+dx))}{d} + \frac{2a^3(iA+B) \tan(c+dx)}{d}$$

$$+ \frac{a(A-iB)(a+ia \tan(c+dx))^2}{2d} + \frac{A(a+ia \tan(c+dx))^3}{3d} - \frac{iB(a+ia \tan(c+dx))^4}{4ad}$$

```
output -4*a^3*(I*A+B)*x-4*a^3*(A-I*B)*ln(cos(d*x+c))/d+2*a^3*(I*A+B)*tan(d*x+c)/d
+1/2*a*(A-I*B)*(a+I*a*tan(d*x+c))^2/d+1/3*A*(a+I*a*tan(d*x+c))^3/d-1/4*I*B
*(a+I*a*tan(d*x+c))^4/a/d
```

3.18.2 Mathematica [A] (verified)

Time = 1.02 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.76

$$\int \tan(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{a^3(4A - 3iB + 48(A - iB) \log(i + \tan(c+dx)) + 48(iA + B) \tan(c+dx) - 6(3A - 4iB) \tan^2(c+dx) - \dots}{12d}$$

input `Integrate[Tan[c + d*x]*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output $(a^3(4A - (3I)B + 48(A - I)B)\text{Log}[I + \text{Tan}[c + d*x]] + 48(I)A + B)\text{Tan}[c + d*x] - 6(3A - (4I)B)\text{Tan}[c + d*x]^2 - (4I)(A - (3I)B)\text{Tan}[c + d*x]^3 - (3I)B\text{Tan}[c + d*x]^4)/(12*d)$

3.18.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.93, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3042, 4075, 3042, 4010, 3042, 3959, 3042, 3958, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4075} \\
 & \int (i \tan(c + dx)a + a)^3(A \tan(c + dx) - B) dx - \frac{iB(a + ia \tan(c + dx))^4}{4ad} \\
 & \quad \downarrow \text{3042} \\
 & \int (i \tan(c + dx)a + a)^3(A \tan(c + dx) - B) dx - \frac{iB(a + ia \tan(c + dx))^4}{4ad} \\
 & \quad \downarrow \text{4010} \\
 & -(B + iA) \int (i \tan(c + dx)a + a)^3 dx + \frac{A(a + ia \tan(c + dx))^3}{3d} - \frac{iB(a + ia \tan(c + dx))^4}{4ad} \\
 & \quad \downarrow \text{3042} \\
 & -(B + iA) \int (i \tan(c + dx)a + a)^3 dx + \frac{A(a + ia \tan(c + dx))^3}{3d} - \frac{iB(a + ia \tan(c + dx))^4}{4ad} \\
 & \quad \downarrow \text{3959}
 \end{aligned}$$

$$\begin{aligned}
& -(B + iA) \left(2a \int (i \tan(c + dx)a + a)^2 dx + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \frac{A(a + ia \tan(c + dx))^3}{3d} - \\
& \quad \frac{iB(a + ia \tan(c + dx))^4}{4ad} \\
& \quad \downarrow \text{3042} \\
& -(B + iA) \left(2a \int (i \tan(c + dx)a + a)^2 dx + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \frac{A(a + ia \tan(c + dx))^3}{3d} - \\
& \quad \frac{iB(a + ia \tan(c + dx))^4}{4ad} \\
& \quad \downarrow \text{3958} \\
& -(B + iA) \left(2a \left(2ia^2 \int \tan(c + dx) dx - \frac{a^2 \tan(c + dx)}{d} + 2a^2 x \right) + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \\
& \quad \frac{A(a + ia \tan(c + dx))^3}{3d} - \frac{iB(a + ia \tan(c + dx))^4}{4ad} \\
& \quad \downarrow \text{3042} \\
& -(B + iA) \left(2a \left(2ia^2 \int \tan(c + dx) dx - \frac{a^2 \tan(c + dx)}{d} + 2a^2 x \right) + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \\
& \quad \frac{A(a + ia \tan(c + dx))^3}{3d} - \frac{iB(a + ia \tan(c + dx))^4}{4ad} \\
& \quad \downarrow \text{3956} \\
& -(B + iA) \left(2a \left(-\frac{a^2 \tan(c + dx)}{d} - \frac{2ia^2 \log(\cos(c + dx))}{d} + 2a^2 x \right) + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \\
& \quad \frac{A(a + ia \tan(c + dx))^3}{3d} - \frac{iB(a + ia \tan(c + dx))^4}{4ad}
\end{aligned}$$

input `Int[Tan[c + d*x]*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(A*(a + I*a*Tan[c + d*x])^3)/(3*d) - ((I/4)*B*(a + I*a*Tan[c + d*x])^4)/(a*d) - (I*A + B)*(((I/2)*a*(a + I*a*Tan[c + d*x])^2)/d + 2*a*(2*a^2*x - ((2*I)*a^2*Log[Cos[c + d*x]]))/d - (a^2*Tan[c + d*x])/d)`

3.18.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 3958 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^2, x_Symbol] := Simp[(a^2 - b^2)*x, x] + (Simp[b^2*(Tan[c + d*x]/d), x] + Simp[2*a*b Int[Tan[c + d*x], x], x]) /; FreeQ[{a, b, c, d}, x]`

rule 3959 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.18.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.89

method	result
derivativedivides	$a^3 \left(-\frac{iB(\tan^4(dx+c))}{4} - \frac{iA(\tan^3(dx+c))}{3} + 2iB(\tan^2(dx+c)) - B(\tan^3(dx+c)) + 4iA \tan(dx+c) - \frac{3A(\tan^2(dx+c))}{2} + 4B \tan(dx+c) \right) \frac{1}{d}$
default	$a^3 \left(-\frac{iB(\tan^4(dx+c))}{4} - \frac{iA(\tan^3(dx+c))}{3} + 2iB(\tan^2(dx+c)) - B(\tan^3(dx+c)) + 4iA \tan(dx+c) - \frac{3A(\tan^2(dx+c))}{2} + 4B \tan(dx+c) \right) \frac{1}{d}$
norman	$(-4iA a^3 - 4B a^3) x - \frac{(iA a^3 + 3B a^3)(\tan^3(dx+c))}{3d} - \frac{(-4iB a^3 + 3A a^3)(\tan^2(dx+c))}{2d} + \frac{4(iA a^3 + B a^3) \tan(dx+c)}{d}$
parallelrisch	$-\frac{3iB a^3(\tan^4(dx+c)) + 4iA(\tan^3(dx+c))a^3 + 48iA x a^3 d - 24iB(\tan^2(dx+c))a^3 + 12B(\tan^3(dx+c))a^3 - 48iA \tan(dx+c)}{12a^3 d}$
risch	$\frac{8a^3 Bc}{d} + \frac{8ia^3 Ac}{d} + \frac{2ia^3(24iA e^{6i(dx+c)} + 36B e^{6i(dx+c)} + 57iA e^{4i(dx+c)} + 69B e^{4i(dx+c)} + 46iA e^{2i(dx+c)} + 54B e^{2i(dx+c)})}{3d(e^{2i(dx+c)} + 1)^4}$
parts	$\frac{(-iA a^3 - 3B a^3) \left(\frac{(\tan^3(dx+c))}{3} - \tan(dx+c) + \arctan(\tan(dx+c)) \right)}{d} + \frac{(3iA a^3 + B a^3)(\tan(dx+c) - \arctan(\tan(dx+c)))}{d}$

input `int(tan(d*x+c)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*a^3*(-1/4*I*B*tan(d*x+c)^4-1/3*I*A*tan(d*x+c)^3+2*I*B*tan(d*x+c)^2-B*tan(d*x+c)^3+4*I*A*tan(d*x+c)-3/2*A*tan(d*x+c)^2+4*B*tan(d*x+c)+1/2*(-4*I*B+4*A)*ln(1+tan(d*x+c)^2)+(-4*I*A-4*B)*arctan(tan(d*x+c)))`

3.18.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.64

$$\int \tan(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx =$$

$$\frac{2(12(2A - 3iB)a^3 e^{6i dx + 6i c} + 3(19A - 23iB)a^3 e^{4i dx + 4i c} + 2(23A - 27iB)a^3 e^{2i dx + 2i c} + (13A - 3iB)a^3)}{3(d e^{8i c})}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

3.18. $\int \tan(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

output
$$\begin{aligned} & -2/3*(12*(2*A - 3*I*B)*a^3*e^{(6*I*d*x + 6*I*c)} + 3*(19*A - 23*I*B)*a^3*e^{(4*I*d*x + 4*I*c)} + 2*(23*A - 27*I*B)*a^3*e^{(2*I*d*x + 2*I*c)} + (13*A - 15*I*B)*a^3 + 6*((A - I*B)*a^3*e^{(8*I*d*x + 8*I*c)} + 4*(A - I*B)*a^3*e^{(6*I*d*x + 6*I*c)} + 6*(A - I*B)*a^3*e^{(4*I*d*x + 4*I*c)} + 4*(A - I*B)*a^3*e^{(2*I*d*x + 2*I*c)} + (A - I*B)*a^3)*\log(e^{(2*I*d*x + 2*I*c)} + 1))/(d*e^{(8*I*d*x + 8*I*c)} + 4*d*e^{(6*I*d*x + 6*I*c)} + 6*d*e^{(4*I*d*x + 4*I*c)} + 4*d*e^{(2*I*d*x + 2*I*c)} + d) \end{aligned}$$

3.18.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 235 vs. $2(114) = 228$.

Time = 0.47 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.70

$$\int \tan(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx = -\frac{4a^3(A-iB) \log(e^{2idx} + e^{-2ic})}{d} + \frac{-26Aa^3 + 30iBa^3 + (-92Aa^3e^{2ic} + 108iBa^3e^{2ic})e^{2idx} + (-114Aa^3e^{4ic} + 138iBa^3e^{4ic})e^{4idx} + (-48Aa^3e^{6ic} + 72iBa^3e^{6ic})e^{6idx}}{3de^{8ic}e^{8idx} + 12de^{6ic}e^{6idx} + 18de^{4ic}e^{4idx} + 12de^{2ic}e^{2idx} + 3d}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)), x)`

output
$$\begin{aligned} & -4*a**3*(A - I*B)*\log(\exp(2*I*d*x) + \exp(-2*I*c))/d + (-26*A*a**3 + 30*I*B*a**3 + (-92*A*a**3*\exp(2*I*c) + 108*I*B*a**3*\exp(2*I*c))*\exp(2*I*d*x) + (-114*A*a**3*\exp(4*I*c) + 138*I*B*a**3*\exp(4*I*c))*\exp(4*I*d*x) + (-48*A*a**3*\exp(6*I*c) + 72*I*B*a**3*\exp(6*I*c))*\exp(6*I*d*x))/(3*d*\exp(8*I*c)*\exp(8*I*d*x) + 12*d*\exp(6*I*c)*\exp(6*I*d*x) + 18*d*\exp(4*I*c)*\exp(4*I*d*x) + 12*d*\exp(2*I*c)*\exp(2*I*d*x) + 3*d) \end{aligned}$$

3.18.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.83

$$\int \tan(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx = \frac{3iBa^3 \tan(dx+c)^4 + 4(iA+3B)a^3 \tan(dx+c)^3 + 6(3A-4iB)a^3 \tan(dx+c)^2 + 48(dx+c)(iA+3B)a^3 \tan(dx+c) + 48Aa^3}{12d}$$

3.18. $\int \tan(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output
$$-1/12*(3*I*B*a^3*\tan(d*x + c)^4 + 4*(I*A + 3*B)*a^3*\tan(d*x + c)^3 + 6*(3*A - 4*I*B)*a^3*\tan(d*x + c)^2 + 48*(d*x + c)*(I*A + B)*a^3 - 24*(A - I*B)*a^3*\log(\tan(d*x + c)^2 + 1) + 48*(-I*A - B)*a^3*\tan(d*x + c))/d$$

3.18.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 408 vs. $2(116) = 232$.

Time = 0.53 (sec) , antiderivative size = 408, normalized size of antiderivative = 2.96

$$\int \tan(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{2(6Aa^3e^{(8i dx + 8i c)} \log(e^{(2i dx + 2i c)} + 1) - 6iBa^3e^{(8i dx + 8i c)} \log(e^{(2i dx + 2i c)} + 1) + 24Aa^3e^{(6i dx + 6i c)} \log(e^{(2i dx + 2i c)} + 1) - 24iBae^{(6i dx + 6i c)} \log(e^{(2i dx + 2i c)} + 1) + 36Aa^3e^{(4i dx + 4i c)} \log(e^{(2i dx + 2i c)} + 1) - 36iBae^{(4i dx + 4i c)} \log(e^{(2i dx + 2i c)} + 1) + 24Aa^3e^{(2i dx + 2i c)} \log(e^{(2i dx + 2i c)} + 1) - 24iBae^{(2i dx + 2i c)} \log(e^{(2i dx + 2i c)} + 1) + 24Aa^3e^{(6i dx + 6i c)} - 36iBae^{(6i dx + 6i c)} + 57Aa^3e^{(4i dx + 4i c)} - 69iBae^{(4i dx + 4i c)} + 46Aa^3e^{(2i dx + 2i c)} - 54iBae^{(2i dx + 2i c)} + 6Aa^3 \log(e^{(2i dx + 2i c)} + 1) - 6iBae^3 \log(e^{(2i dx + 2i c)} + 1) + 13Aa^3 - 15iBae^3)/(d * e^{(8i dx + 8i c)} + 4 * d * e^{(6i dx + 6i c)} + 6 * d * e^{(4i dx + 4i c)} + 4 * d * e^{(2i dx + 2i c)} + d)$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output
$$-2/3*(6*A*a^3*e^{(8*I*d*x + 8*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 6*I*B*a^3*e^{(8*I*d*x + 8*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) + 24*A*a^3*e^{(6*I*d*x + 6*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 24*I*B*a^3*e^{(6*I*d*x + 6*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) + 36*A*a^3*e^{(4*I*d*x + 4*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 36*I*B*a^3*e^{(4*I*d*x + 4*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) + 24*A*a^3*e^{(2*I*d*x + 2*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 24*I*B*a^3*e^{(2*I*d*x + 2*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) + 24*A*a^3*e^{(6*I*d*x + 6*I*c)} - 36*I*B*a^3*e^{(6*I*d*x + 6*I*c)} + 57*A*a^3*e^{(4*I*d*x + 4*I*c)} - 69*I*B*a^3*e^{(4*I*d*x + 4*I*c)} + 46*A*a^3*e^{(2*I*d*x + 2*I*c)} - 54*I*B*a^3*e^{(2*I*d*x + 2*I*c)} + 6*A*a^3*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 6*I*B*a^3*\log(e^{(2*I*d*x + 2*I*c)} + 1) + 13*A*a^3 - 15*I*B*a^3)/(d*e^{(8*I*d*x + 8*I*c)} + 4*d*e^{(6*I*d*x + 6*I*c)} + 6*d*e^{(4*I*d*x + 4*I*c)} + 4*d*e^{(2*I*d*x + 2*I*c)} + d)$$

3.18. $\int \tan(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.18.9 Mupad [B] (verification not implemented)

Time = 7.56 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.28

$$\int \tan(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{\tan(c + dx)^2 \left(\frac{B a^3 i i}{2} - \frac{a^3 (2A - B i i)}{2} + \frac{a^3 (2B + A i i) i i}{2} \right)}{d}$$

$$+ \frac{\tan(c + dx) (A a^3 i i + B a^3 + a^3 (2A - B i i) i i + a^3 (2B + A i i))}{d}$$

$$- \frac{\tan(c + dx)^3 \left(\frac{B a^3}{3} + \frac{a^3 (2B + A i i)}{3} \right)}{d}$$

$$+ \frac{\ln(\tan(c + dx) + i i) (4A a^3 - B a^3 4i)}{d} - \frac{B a^3 \tan(c + dx)^4 i i}{4d}$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3,x)`output `(tan(c + d*x)^2*((B*a^3*1i)/2 - (a^3*(2*A - B*1i))/2 + (a^3*(A*1i + 2*B)*1i)/2))/d + (tan(c + d*x)*(A*a^3*1i + B*a^3 + a^3*(2*A - B*1i)*1i + a^3*(A*1i + 2*B)))/d - (tan(c + d*x)^3*((B*a^3)/3 + (a^3*(A*1i + 2*B))/3))/d + (log(tan(c + d*x) + 1i)*(4*A*a^3 - B*a^3*4i))/d - (B*a^3*tan(c + d*x)^4*1i)/(4*d)`

3.19 $\int (a + ia \tan(c + dx))^3 (A + B \tan(c + dx)) dx$

3.19.1	Optimal result	427
3.19.2	Mathematica [A] (verified)	427
3.19.3	Rubi [A] (verified)	428
3.19.4	Maple [A] (verified)	430
3.19.5	Fricas [A] (verification not implemented)	430
3.19.6	Sympy [A] (verification not implemented)	431
3.19.7	Maxima [A] (verification not implemented)	431
3.19.8	Giac [B] (verification not implemented)	432
3.19.9	Mupad [B] (verification not implemented)	432

3.19.1 Optimal result

Integrand size = 26, antiderivative size = 110

$$\int (a + ia \tan(c + dx))^3 (A + B \tan(c + dx)) dx$$

$$= 4a^3(A - iB)x - \frac{4a^3(iA + B) \log(\cos(c + dx))}{d} - \frac{2a^3(A - iB) \tan(c + dx)}{d}$$

$$+ \frac{a(iA + B)(a + ia \tan(c + dx))^2}{2d} + \frac{B(a + ia \tan(c + dx))^3}{3d}$$

output `4*a^3*(A-I*B)*x-4*a^3*(I*A+B)*ln(cos(d*x+c))/d-2*a^3*(A-I*B)*tan(d*x+c)/d+1/2*a*(I*A+B)*(a+I*a*tan(d*x+c))^2/d+1/3*B*(a+I*a*tan(d*x+c))^3/d`

3.19.2 Mathematica [A] (verified)

Time = 0.87 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.66

$$\int (a + ia \tan(c + dx))^3 (A + B \tan(c + dx)) dx$$

$$= \frac{B(a + ia \tan(c + dx))^3 + \frac{3}{2}a^3(iA + B) (8 \log(i + \tan(c + dx)) + 6i \tan(c + dx) - \tan^2(c + dx))}{3d}$$

input `Integrate[(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(B*(a + I*a*Tan[c + d*x])^3 + (3*a^3*(I*A + B)*(8*Log[I + Tan[c + d*x]] + (6*I)*Tan[c + d*x] - Tan[c + d*x]^2))/2)/(3*d)`

3.19.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.90, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3042, 4010, 3042, 3959, 3042, 3958, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(c + dx))^3 (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(c + dx))^3 (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4010} \\
 & (A - iB) \int (i \tan(c + dx)a + a)^3 dx + \frac{B(a + ia \tan(c + dx))^3}{3d} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \int (i \tan(c + dx)a + a)^3 dx + \frac{B(a + ia \tan(c + dx))^3}{3d} \\
 & \quad \downarrow \text{3959} \\
 & (A - iB) \left(2a \int (i \tan(c + dx)a + a)^2 dx + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \frac{B(a + ia \tan(c + dx))^3}{3d} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \left(2a \int (i \tan(c + dx)a + a)^2 dx + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \frac{B(a + ia \tan(c + dx))^3}{3d} \\
 & \quad \downarrow \text{3958} \\
 & (A - iB) \left(2a \left(2ia^2 \int \tan(c + dx) dx - \frac{a^2 \tan(c + dx)}{d} + 2a^2 x \right) + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \\
 & \quad \frac{B(a + ia \tan(c + dx))^3}{3d} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \left(2a \left(2ia^2 \int \tan(c + dx) dx - \frac{a^2 \tan(c + dx)}{d} + 2a^2 x \right) + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \\
 & \quad \frac{B(a + ia \tan(c + dx))^3}{3d} \\
 & \quad \downarrow \text{3956}
 \end{aligned}$$

$$(A - iB) \left(2a \left(-\frac{a^2 \tan(c + dx)}{d} - \frac{2ia^2 \log(\cos(c + dx))}{d} + 2a^2 x \right) + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \frac{B(a + ia \tan(c + dx))^3}{3d}$$

input `Int[(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(B*(a + I*a*Tan[c + d*x])^3)/(3*d) + (A - I*B)*(((I/2)*a*(a + I*a*Tan[c + d*x])^2)/d + 2*a*(2*a^2*x - ((2*I)*a^2*Log[Cos[c + d*x]])/d - (a^2*Tan[c + d*x])/d)`

3.19.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 3958 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^2, x_Symbol] := Simp[(a^2 - b^2)*x, x] + (Simp[b^2*(Tan[c + d*x]/d), x] + Simp[2*a*b Int[Tan[c + d*x], x], x]) /; FreeQ[{a, b, c, d}, x]`

rule 3959 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

3.19.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.91

method	result
derivativedivides	$\frac{a^3 \left(-\frac{iB(\tan^3(dx+c))}{3} - \frac{iA(\tan^2(dx+c))}{2} + 4iB \tan(dx+c) - \frac{3B(\tan^2(dx+c))}{2} - 3A \tan(dx+c) + \frac{(4iA+4B) \ln(1+\tan^2(dx+c))}{2} \right)}{d}$
default	$\frac{a^3 \left(-\frac{iB(\tan^3(dx+c))}{3} - \frac{iA(\tan^2(dx+c))}{2} + 4iB \tan(dx+c) - \frac{3B(\tan^2(dx+c))}{2} - 3A \tan(dx+c) + \frac{(4iA+4B) \ln(1+\tan^2(dx+c))}{2} \right)}{d}$
norman	$\frac{(-4iB a^3 + 4A a^3) x - \frac{(iA a^3 + 3B a^3)(\tan^2(dx+c))}{2d} - \frac{(-4iB a^3 + 3A a^3) \tan(dx+c)}{d} - \frac{iB a^3 (\tan^3(dx+c))}{3d}}{-2iB(\tan^3(dx+c))a^3 - 3iA(\tan^2(dx+c))a^3 - 24iBx a^3 d + 12iA \ln(1+\tan^2(dx+c))a^3 + 24Ax a^3 d + 24iB \tan(dx+c)a^3 - 9}$
parallelrisch	$\frac{-2iB(\tan^3(dx+c))a^3 - 3iA(\tan^2(dx+c))a^3 - 24iBx a^3 d + 12iA \ln(1+\tan^2(dx+c))a^3 + 24Ax a^3 d + 24iB \tan(dx+c)a^3 - 9}{6d}$
risch	$\frac{\frac{8ia^3 Bc}{d} - \frac{8a^3 Ac}{d} - \frac{2a^3(12iA e^{4i(dx+c)} + 24B e^{4i(dx+c)} + 21iA e^{2i(dx+c)} + 33B e^{2i(dx+c)} + 9iA + 13B)}{3d(e^{2i(dx+c)} + 1)^3} - \frac{4a^3 \ln(e^{2i(dx+c)} + 1)}{d}}{d}$
parts	$A a^3 x + \frac{(-iA a^3 - 3B a^3) \left(\frac{\tan^2(dx+c)}{2} - \frac{\ln(1+\tan^2(dx+c))}{2} \right)}{d} + \frac{(3iA a^3 + B a^3) \ln(1+\tan^2(dx+c))}{2d} + \frac{(3iB a^3 - 3A a^3)}{2d}$

input `int((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*a^3*(-1/3*I*B*tan(d*x+c)^3-1/2*I*A*tan(d*x+c)^2+4*I*B*tan(d*x+c)-3/2*B*tan(d*x+c)^2-3*A*tan(d*x+c)+1/2*(4*I*A+4*B)*ln(1+tan(d*x+c)^2)+(-4*I*B+4*A)*arctan(tan(d*x+c)))`

3.19.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.59

$$\int (a + ia \tan(c + dx))^3 (A + B \tan(c + dx)) dx = \frac{2(12(iA + 2B)a^3 e^{(4i dx + 4i c)} + 3(7iA + 11B)a^3 e^{(2i dx + 2i c)} + (9iA + 13B)a^3 + 6((iA + B)a^3 e^{(6i dx + 6i c)} - 3(de^{(6i dx + 6i c)} + 3de^{(4i dx + 4i c)}))}{3(de^{(6i dx + 6i c)} + 3de^{(4i dx + 4i c)})}$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output
$$\begin{aligned} & -2/3*(12*(I*A + 2*B)*a^3*e^(4*I*d*x + 4*I*c) + 3*(7*I*A + 11*B)*a^3*e^(2*I \\ & *d*x + 2*I*c) + (9*I*A + 13*B)*a^3 + 6*((I*A + B)*a^3*e^(6*I*d*x + 6*I*c) \\ & + 3*(I*A + B)*a^3*e^(4*I*d*x + 4*I*c) + 3*(I*A + B)*a^3*e^(2*I*d*x + 2*I*c) \\ &) + (I*A + B)*a^3)*\log(e^(2*I*d*x + 2*I*c) + 1))/(d*e^(6*I*d*x + 6*I*c) + \\ & 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d) \end{aligned}$$

3.19.6 Sympy [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.67

$$\int (a + ia \tan(c + dx))^3 (A + B \tan(c + dx)) dx = -\frac{4ia^3(A - iB) \log(e^{2idx} + e^{-2ic})}{d} + \frac{-18iAa^3 - 26Ba^3 + (-42iAa^3e^{2ic} - 66Ba^3e^{2ic})e^{2idx} + (-24iAa^3e^{4ic} - 48Ba^3e^{4ic})e^{4idx}}{3de^{6ic}e^{6idx} + 9de^{4ic}e^{4idx} + 9de^{2ic}e^{2idx} + 3d}$$

input `integrate((a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output
$$\begin{aligned} & -4*I*a**3*(A - I*B)*\log(\exp(2*I*d*x) + \exp(-2*I*c))/d + (-18*I*A*a**3 - 26 \\ & *B*a**3 + (-42*I*A*a**3*\exp(2*I*c) - 66*B*a**3*\exp(2*I*c))*\exp(2*I*d*x) + \\ & (-24*I*A*a**3*\exp(4*I*c) - 48*B*a**3*\exp(4*I*c))*\exp(4*I*d*x))/(3*d*\exp(6* \\ & I*c)*\exp(6*I*d*x) + 9*d*\exp(4*I*c)*\exp(4*I*d*x) + 9*d*\exp(2*I*c)*\exp(2*I*d \\ & *x) + 3*d) \end{aligned}$$

3.19.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.87

$$\int (a + ia \tan(c + dx))^3 (A + B \tan(c + dx)) dx = \frac{2iBa^3 \tan(dx + c)^3 + 3(iA + 3B)a^3 \tan(dx + c)^2 - 24(dx + c)(A - iB)a^3 + 12(-iA - B)a^3 \log(\tan(dx + c)^2 + 1)}{6d}$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output
$$\begin{aligned} & -1/6*(2*I*B*a^3*\tan(d*x + c)^3 + 3*(I*A + 3*B)*a^3*\tan(d*x + c)^2 - 24*(d* \\ & x + c)*(A - I*B)*a^3 + 12*(-I*A - B)*a^3*\log(\tan(d*x + c)^2 + 1) + 6*(3*A \\ & - 4*I*B)*a^3*\tan(d*x + c))/d \end{aligned}$$

3.19.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 312 vs. $2(94) = 188$.

Time = 0.46 (sec) , antiderivative size = 312, normalized size of antiderivative = 2.84

$$\int (a + ia \tan(c + dx))^3 (A + B \tan(c + dx)) dx =$$

$$\frac{2(6i Aa^3 e^{(6i dx + 6i c)} \log(e^{(2i dx + 2i c)} + 1) + 6Ba^3 e^{(6i dx + 6i c)} \log(e^{(2i dx + 2i c)} + 1) + 18i Aa^3 e^{(4i dx + 4i c)} \log(e^{(2i dx + 2i c)} + 1) + 18i Ba^3 e^{(4i dx + 4i c)} \log(e^{(2i dx + 2i c)} + 1) + 12i Aa^3 e^{(4i dx + 4i c)} + 24i Ba^3 e^{(4i dx + 4i c)} + 21i Aa^3 e^{(2i dx + 2i c)} + 33i Ba^3 e^{(2i dx + 2i c)} + 6i Aa^3 \log(e^{(2i dx + 2i c)} + 1) + 6i Ba^3 \log(e^{(2i dx + 2i c)} + 1) + 9i Aa^3 + 13i Ba^3) / (d e^{(6i dx + 6i c)} + 3d e^{(4i dx + 4i c)} + 3d e^{(2i dx + 2i c)} + d)}$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `-2/3*(6*I*A*a^3*e^(6*I*d*x + 6*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 6*B*a^3*e^(6*I*d*x + 6*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 18*I*A*a^3*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 18*B*a^3*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 18*I*A*a^3*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 18*B*a^3*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 12*I*A*a^3*e^(4*I*d*x + 4*I*c) + 24*B*a^3*e^(4*I*d*x + 4*I*c) + 21*I*A*a^3*e^(2*I*d*x + 2*I*c) + 33*B*a^3*e^(2*I*d*x + 2*I*c) + 6*I*A*a^3*log(e^(2*I*d*x + 2*I*c) + 1) + 6*B*a^3*log(e^(2*I*d*x + 2*I*c) + 1) + 9*I*A*a^3 + 13*B*a^3)/(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)`

3.19.9 Mupad [B] (verification not implemented)

Time = 7.86 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.14

$$\int (a + ia \tan(c + dx))^3 (A + B \tan(c + dx)) dx$$

$$= -\frac{\tan(c + dx)^2 \left(\frac{Ba^3}{2} + \frac{a^3(2B + A1i)}{2} \right)}{d} + \frac{\ln(\tan(c + dx) + 1i) (4Ba^3 + Aa^3 4i)}{d}$$

$$+ \frac{\tan(c + dx) (Ba^3 1i - a^3(2A - B1i) + a^3(2B + A1i) 1i)}{d} - \frac{Ba^3 \tan(c + dx)^3 1i}{3d}$$

input `int((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3,x)`

output `(log(tan(c + d*x) + 1i)*(A*a^3*4i + 4*B*a^3))/d - (tan(c + d*x)^2*((B*a^3)/2 + (a^3*(A*1i + 2*B))/2))/d + (tan(c + d*x)*(B*a^3*1i - a^3*(2*A - B*1i) + a^3*(A*1i + 2*B)*1i))/d - (B*a^3*tan(c + d*x)^3*1i)/(3*d)`

3.19. $\int (a + ia \tan(c + dx))^3 (A + B \tan(c + dx)) dx$

3.20 $\int \cot(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.20.1	Optimal result	433
3.20.2	Mathematica [A] (verified)	433
3.20.3	Rubi [A] (verified)	434
3.20.4	Maple [A] (verified)	437
3.20.5	Fricas [A] (verification not implemented)	438
3.20.6	Sympy [B] (verification not implemented)	438
3.20.7	Maxima [A] (verification not implemented)	439
3.20.8	Giac [B] (verification not implemented)	439
3.20.9	Mupad [B] (verification not implemented)	440

3.20.1 Optimal result

Integrand size = 32, antiderivative size = 107

$$\int \cot(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= 4a^3(iA+B)x + \frac{a^3(3A-4iB) \log(\cos(c+dx))}{d} + \frac{a^3A \log(\sin(c+dx))}{d}$$

$$+ \frac{iaB(a+ia \tan(c+dx))^2}{2d} - \frac{(A-2iB)(a^3+ia^3 \tan(c+dx))}{d}$$

```
output 4*a^3*(I*A+B)*x+a^3*(3*A-4*I*B)*ln(cos(d*x+c))/d+a^3*A*ln(sin(d*x+c))/d+1/
2*I*a*B*(a+I*a*tan(d*x+c))^2/d-(A-2*I*B)*(a^3+I*a^3*tan(d*x+c))/d
```

3.20.2 Mathematica [A] (verified)

Time = 0.88 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.65

$$\int \cot(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{a^3(2A \log(\tan(c+dx)) - 8(A-iB) \log(i+\tan(c+dx)) + (-2iA-6B) \tan(c+dx) - iB \tan^2(c+dx))}{2d}$$

```
input Integrate[Cot[c + d*x]*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]
```


output $(a^3(2A \operatorname{Log}[\operatorname{Tan}[c + dx]] - 8(A - I*B) \operatorname{Log}[I + \operatorname{Tan}[c + dx]] + ((-2*I)*A - 6*B) \operatorname{Tan}[c + dx] - I*B \operatorname{Tan}[c + dx]^2))/(2*d)$

3.20.3 Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.02, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.406$, Rules used = {3042, 4077, 27, 3042, 4077, 3042, 4072, 3042, 3956, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\tan(c + dx)} dx \\
 & \quad \downarrow \text{4077} \\
 & \frac{1}{2} \int 2 \cot(c + dx)(i \tan(c + dx)a + a)^2(aA + a(iA + 2B) \tan(c + dx)) dx + \frac{iaB(a + ia \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{27} \\
 & \int \cot(c + dx)(i \tan(c + dx)a + a)^2(aA + a(iA + 2B) \tan(c + dx)) dx + \frac{iaB(a + ia \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(i \tan(c + dx)a + a)^2(aA + a(iA + 2B) \tan(c + dx))}{\tan(c + dx)} dx + \frac{iaB(a + ia \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{4077} \\
 & \int \cot(c + dx)(i \tan(c + dx)a + a) (Aa^2 + (3iA + 4B) \tan(c + dx)a^2) dx - \\
 & \quad \frac{(A - 2iB)(a^3 + ia^3 \tan(c + dx))}{d} + \frac{iaB(a + ia \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(i \tan(c + dx)a + a) (Aa^2 + (3iA + 4B) \tan(c + dx)a^2)}{\tan(c + dx)} dx - \\
 & \quad \frac{(A - 2iB)(a^3 + ia^3 \tan(c + dx))}{d} + \frac{iaB(a + ia \tan(c + dx))^2}{2d}
 \end{aligned}$$

3.20. $\int \cot(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \downarrow 4072 \\
& -\left(a^3(3A - 4iB) \int \tan(c + dx) dx\right) + \int \cot(c + dx) (Aa^3 + 4(iA + B) \tan(c + dx)a^3) dx - \\
& \quad \frac{(A - 2iB)(a^3 + ia^3 \tan(c + dx))}{d} + \frac{iaB(a + ia \tan(c + dx))^2}{2d} \\
& \downarrow 3042 \\
& -\left(a^3(3A - 4iB) \int \tan(c + dx) dx\right) + \int \frac{Aa^3 + 4(iA + B) \tan(c + dx)a^3}{\tan(c + dx)} dx - \\
& \quad \frac{(A - 2iB)(a^3 + ia^3 \tan(c + dx))}{d} + \frac{iaB(a + ia \tan(c + dx))^2}{2d} \\
& \downarrow 3956 \\
& \int \frac{Aa^3 + 4(iA + B) \tan(c + dx)a^3}{\tan(c + dx)} dx - \frac{(A - 2iB)(a^3 + ia^3 \tan(c + dx))}{d} + \\
& \quad \frac{a^3(3A - 4iB) \log(\cos(c + dx))}{d} + \frac{iaB(a + ia \tan(c + dx))^2}{2d} \\
& \downarrow 4014 \\
& a^3 A \int \cot(c + dx) dx - \frac{(A - 2iB)(a^3 + ia^3 \tan(c + dx))}{d} + \frac{a^3(3A - 4iB) \log(\cos(c + dx))}{d} + \\
& \quad 4a^3 x(B + iA) + \frac{iaB(a + ia \tan(c + dx))^2}{2d} \\
& \downarrow 3042 \\
& a^3 A \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - \frac{(A - 2iB)(a^3 + ia^3 \tan(c + dx))}{d} + \\
& \quad \frac{a^3(3A - 4iB) \log(\cos(c + dx))}{d} + 4a^3 x(B + iA) + \frac{iaB(a + ia \tan(c + dx))^2}{2d} \\
& \downarrow 25 \\
& -a^3 A \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - \frac{(A - 2iB)(a^3 + ia^3 \tan(c + dx))}{d} + \\
& \quad \frac{a^3(3A - 4iB) \log(\cos(c + dx))}{d} + 4a^3 x(B + iA) + \frac{iaB(a + ia \tan(c + dx))^2}{2d} \\
& \downarrow 3956 \\
& -\frac{(A - 2iB)(a^3 + ia^3 \tan(c + dx))}{d} + \frac{a^3(3A - 4iB) \log(\cos(c + dx))}{d} + 4a^3 x(B + iA) + \\
& \quad \frac{a^3 A \log(-\sin(c + dx))}{d} + \frac{iaB(a + ia \tan(c + dx))^2}{2d}
\end{aligned}$$

input `Int[Cot[c + d*x]*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output $4a^3(I A + B)x + (a^3(3A - (4I)B)\text{Log}[\text{Cos}[c + dx]])/d + (a^3A\text{Log}[-\text{Sin}[c + dx]])/d + ((I/2)a^3B(a + I a \text{Tan}[c + dx])^2)/d - ((A - (2I)B)(a^3 + I a^3 \text{Tan}[c + dx]))/d$

3.20.3.1 Defintions of rubi rules used

rule 25 $\text{Int}[-(F x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{Int}[F x, x], x]$

rule 27 $\text{Int}[(a)(F x), x_Symbol] \rightarrow \text{Simp}[a \text{Int}[F x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F x, (b)(G x)] /; \text{FreeQ}[b, x]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3956 $\text{Int}[\text{tan}[(c) + (d)(x)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + dx], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

rule 4014 $\text{Int}[((c) + (d)\text{tan}[(e) + (f)(x)])/((a) + (b)\text{tan}[(e) + (f)(x)]), x_Symbol] \rightarrow \text{Simp}[(a c + b d)(x/(a^2 + b^2)), x] + \text{Simp}[(b c - a d)/(a^2 + b^2) \text{Int}[(b - a \text{Tan}[e + f x])/(a + b \text{Tan}[e + f x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b c - a d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[a c + b d, 0]$

rule 4072 $\text{Int}[(((A) + (B)\text{tan}[(e) + (f)(x)])((c) + (d)\text{tan}[(e) + (f)(x)]))/((a) + (b)\text{tan}[(e) + (f)(x)]), x_Symbol] \rightarrow \text{Simp}[B(d/b) \text{Int}[\text{Tan}[e + f x], x], x] + \text{Simp}[1/b \text{Int}[\text{Simp}[A b c + (A b d + B(b c - a d))\text{Tan}[e + f x], x]/(a + b \text{Tan}[e + f x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[b c - a d, 0]$

```
rule 4077 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan
[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c -
a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] &&
GtQ[m, 1] && !LtQ[n, -1]
```

3.20.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.81

method	result
parallelrisch	$-\frac{a^3(-8iAxd+iB(\tan^2(dx+c))+2iA \tan(dx+c)-4iB \ln(\sec^2(dx+c))-8Bdx+4A \ln(\sec^2(dx+c))-2A \ln(\tan(dx+c)))}{2d}$
derivativedivides	$\frac{a^3 \left(\frac{(4iB-4A) \ln(\cot^2(dx+c)+1)}{2} + (-4iA-4B) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)) \right) + (-4iB+3A) \ln(\cot(dx+c)) - \frac{iA+3B}{\cot(dx+c)} - \frac{iB}{2 \cot(dx+c)} \right)}{d}$
default	$\frac{a^3 \left(\frac{(4iB-4A) \ln(\cot^2(dx+c)+1)}{2} + (-4iA-4B) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)) \right) + (-4iB+3A) \ln(\cot(dx+c)) - \frac{iA+3B}{\cot(dx+c)} - \frac{iB}{2 \cot(dx+c)} \right)}{d}$
norman	$(4iA a^3 + 4B a^3) x - \frac{(iA a^3 + 3B a^3) \tan(dx+c)}{d} - \frac{iB a^3 (\tan^2(dx+c))}{2d} + \frac{A a^3 \ln(\tan(dx+c))}{d} - \frac{2(-iB a^3)}{d}$
risch	$-\frac{8a^3 Bc}{d} - \frac{8ia^3 Ac}{d} - \frac{2ia^3 (iA e^{2i(dx+c)} + 4B e^{2i(dx+c)} + iA + 3B)}{d(e^{2i(dx+c)} + 1)^2} - \frac{4ia^3 \ln(e^{2i(dx+c)} + 1)B}{d} + \frac{3a^3 \ln(e^{2i(dx+c)} + 1)}{d}$

```
input int(cot(d*x+c)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

```
output -1/2*a^3*(-8*I*A*x*d+I*B*tan(d*x+c)^2+2*I*A*tan(d*x+c)-4*I*B*ln(sec(d*x+c)
^2)-8*B*d*x+4*A*ln(sec(d*x+c)^2)-2*A*ln(tan(d*x+c))+6*B*tan(d*x+c))/d
```

3.20. $\int \cot(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.20.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.61

$$\int \cot(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{2(A - 4iB)a^3 e^{(2i dx + 2i c)} + 2(A - 3iB)a^3 + ((3A - 4iB)a^3 e^{(4i dx + 4i c)} + 2(3A - 4iB)a^3 e^{(2i dx + 2i c)} + (3A - 4iB)a^3 e^{(4i dx + 4i c)} + 2de^{(2i dx + 2i c)})}{de^{(4i dx + 4i c)} + 2de^{(2i dx + 2i c)} + d}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `(2*(A - 4*I*B)*a^3*e^(2*I*d*x + 2*I*c) + 2*(A - 3*I*B)*a^3 + ((3*A - 4*I*B)*a^3*e^(4*I*d*x + 4*I*c) + 2*(3*A - 4*I*B)*a^3*e^(2*I*d*x + 2*I*c) + (3*A - 4*I*B)*a^3)*log(e^(2*I*d*x + 2*I*c) + 1) + (A*a^3*e^(4*I*d*x + 4*I*c) + 2*A*a^3*e^(2*I*d*x + 2*I*c) + A*a^3)*log(e^(2*I*d*x + 2*I*c) - 1)/(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)`

3.20.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 224 vs. 2(94) = 188.

Time = 1.60 (sec) , antiderivative size = 224, normalized size of antiderivative = 2.09

$$\int \cot(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{Aa^3 \log\left(\frac{-Aa^3 + 2iBa^3}{Aa^3 e^{2ic} - 2iBa^3 e^{2ic}} + e^{2idx}\right)}{d} + \frac{a^3 \cdot (3A - 4iB) \log\left(e^{2idx} + \frac{-2Aa^3 + 2iBa^3 + a^3 \cdot (3A - 4iB)}{Aa^3 e^{2ic} - 2iBa^3 e^{2ic}}\right)}{d}$$

$$+ \frac{2Aa^3 - 6iBa^3 + (2Aa^3 e^{2ic} - 8iBa^3 e^{2ic}) e^{2idx}}{de^{4ic} e^{4idx} + 2de^{2ic} e^{2idx} + d}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `A*a**3*log((-A*a**3 + 2*I*B*a**3)/(A*a**3*exp(2*I*c) - 2*I*B*a**3*exp(2*I*c)) + exp(2*I*d*x))/d + a**3*(3*A - 4*I*B)*log(exp(2*I*d*x) + (-2*A*a**3 + 2*I*B*a**3 + a**3*(3*A - 4*I*B))/(A*a**3*exp(2*I*c) - 2*I*B*a**3*exp(2*I*c)))/d + (2*A*a**3 - 6*I*B*a**3 + (2*A*a**3*exp(2*I*c) - 8*I*B*a**3*exp(2*I*c))*exp(2*I*d*x))/(d*exp(4*I*c)*exp(4*I*d*x) + 2*d*exp(2*I*c)*exp(2*I*d*x) + d)`

3.20. $\int \cot(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.20.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.83

$$\int \cot(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{i Ba^3 \tan(dx + c)^2 + 8(dx + c)(-iA - B)a^3 + 4(A - iB)a^3 \log(\tan(dx + c)^2 + 1) - 2Aa^3 \log(\tan(dx + c))}{2d}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(I*B*a^3*tan(d*x + c)^2 + 8*(d*x + c)*(-I*A - B)*a^3 + 4*(A - I*B)*a^3*log(tan(d*x + c)^2 + 1) - 2*A*a^3*log(tan(d*x + c)) + 2*(I*A + 3*B)*a^3*tan(d*x + c))/d`

3.20.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 264 vs. 2(93) = 186.

Time = 0.86 (sec) , antiderivative size = 264, normalized size of antiderivative = 2.47

$$\int \cot(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{2Aa^3 \log(\tan(\frac{1}{2}dx + \frac{1}{2}c)) + 2(3Aa^3 - 4iBa^3) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) + 1) - 16(Aa^3 - iBa^3) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1) + 2(3Aa^3 - 4iB*a^3) \log(\tan(\frac{1}{2}d*x + 1/2*c) + 1) - 16*(A*a^3 - I*B*a^3) \log(\tan(\frac{1}{2}d*x + 1/2*c) + I) + 2*(3*A*a^3 - 4*I*B*a^3) \log(\tan(\frac{1}{2}d*x + 1/2*c) - 1) - (9*A*a^3 \tan(\frac{1}{2}d*x + 1/2*c)^4 - 12*I*B*a^3 \tan(\frac{1}{2}d*x + 1/2*c)^4 - 4*I*A*a^3 \tan(\frac{1}{2}d*x + 1/2*c)^3 - 12*B*a^3 \tan(\frac{1}{2}d*x + 1/2*c)^3 - 18*A*a^3 \tan(\frac{1}{2}d*x + 1/2*c)^2 + 28*I*B*a^3 \tan(\frac{1}{2}d*x + 1/2*c)^2 + 4*I*A*a^3 \tan(\frac{1}{2}d*x + 1/2*c) + 12*B*a^3 \tan(\frac{1}{2}d*x + 1/2*c) + 9*A*a^3 - 12*I*B*a^3) / (\tan(\frac{1}{2}d*x + 1/2*c)^2 - 1)^2}{d}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `1/2*(2*A*a^3*log(tan(1/2*d*x + 1/2*c)) + 2*(3*A*a^3 - 4*I*B*a^3)*log(tan(1/2*d*x + 1/2*c) + 1) - 16*(A*a^3 - I*B*a^3)*log(tan(1/2*d*x + 1/2*c) + I) + 2*(3*A*a^3 - 4*I*B*a^3)*log(tan(1/2*d*x + 1/2*c) - 1) - (9*A*a^3*tan(1/2*d*x + 1/2*c)^4 - 12*I*B*a^3*tan(1/2*d*x + 1/2*c)^4 - 4*I*A*a^3*tan(1/2*d*x + 1/2*c)^3 - 12*B*a^3*tan(1/2*d*x + 1/2*c)^3 - 18*A*a^3*tan(1/2*d*x + 1/2*c)^2 + 28*I*B*a^3*tan(1/2*d*x + 1/2*c)^2 + 4*I*A*a^3*tan(1/2*d*x + 1/2*c) + 12*B*a^3*tan(1/2*d*x + 1/2*c) + 9*A*a^3 - 12*I*B*a^3)/(tan(1/2*d*x + 1/2*c)^2 - 1)^2/d`

3.20. $\int \cot(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.20.9 Mupad [B] (verification not implemented)

Time = 7.34 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.81

$$\int \cot(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{A a^3 \ln(\tan(c + dx))}{d} - \frac{\tan(c + dx) (B a^3 + a^3 (2B + A i))}{d}$$

$$- \frac{4 a^3 \ln(\tan(c + dx) + i) (A - B i)}{d} - \frac{B a^3 \tan(c + dx)^2 i}{2d}$$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*i)^3,x)`output `(A*a^3*log(tan(c + d*x)))/d - (tan(c + d*x)*(B*a^3 + a^3*(A*i + 2*B)))/d
- (4*a^3*log(tan(c + d*x) + i)*(A - B*i))/d - (B*a^3*tan(c + d*x)^2*i)/
(2*d)`

3.21 $\int \cot^2(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.21.1	Optimal result	441
3.21.2	Mathematica [A] (verified)	441
3.21.3	Rubi [A] (verified)	442
3.21.4	Maple [A] (verified)	445
3.21.5	Fricas [A] (verification not implemented)	446
3.21.6	Sympy [B] (verification not implemented)	446
3.21.7	Maxima [A] (verification not implemented)	447
3.21.8	Giac [B] (verification not implemented)	447
3.21.9	Mupad [B] (verification not implemented)	448

3.21.1 Optimal result

Integrand size = 34, antiderivative size = 116

$$\int \cot^2(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= -4a^3(A-iB)x + \frac{a^3(iA+3B) \log(\cos(c+dx))}{d} + \frac{a^3(3iA+B) \log(\sin(c+dx))}{d}$$

$$- \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^2}{d} + \frac{(iA-B)(a^3+ia^3 \tan(c+dx))}{d}$$

```
output -4*a^3*(A-I*B)*x+a^3*(I*A+3*B)*ln(cos(d*x+c))/d+a^3*(3*I*A+B)*ln(sin(d*x+c
)))/d-a*A*cot(d*x+c)*(a+I*a*tan(d*x+c))^2/d+(I*A-B)*(a^3+I*a^3*tan(d*x+c))/
d
```

3.21.2 Mathematica [A] (verified)

Time = 1.02 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.81

$$\int \cot^2(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= a^3 \left(-\frac{A \cot(c+dx)}{d} + \frac{3iA \log(\tan(c+dx))}{d} + \frac{B \log(\tan(c+dx))}{d} \right.$$

$$\left. - \frac{4iA \log(i+\tan(c+dx))}{d} - \frac{4B \log(i+\tan(c+dx))}{d} - \frac{iB \tan(c+dx)}{d} \right)$$

input `Integrate[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `a^3*(-((A*Cot[c + d*x])/d) + ((3*I)*A*Log[Tan[c + d*x]])/d + (B*Log[Tan[c + d*x]])/d - ((4*I)*A*Log[I + Tan[c + d*x]])/d - (4*B*Log[I + Tan[c + d*x]])/d - (I*B*Tan[c + d*x])/d)`

3.21.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3042, 4076, 3042, 4077, 3042, 4072, 3042, 3956, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\tan(c + dx)^2} dx \\
 & \quad \downarrow \text{4076} \\
 & \int \cot(c + dx)(i \tan(c + dx)a + a)^2(a(3iA + B) + a(A + iB) \tan(c + dx)) dx - \\
 & \quad \frac{aA \cot(c + dx)(a + ia \tan(c + dx))^2}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(i \tan(c + dx)a + a)^2(a(3iA + B) + a(A + iB) \tan(c + dx))}{\tan(c + dx)} dx - \\
 & \quad \frac{aA \cot(c + dx)(a + ia \tan(c + dx))^2}{d} \\
 & \quad \downarrow \text{4077} \\
 & \int \cot(c + dx)(i \tan(c + dx)a + a) (a^2(3iA + B) - a^2(A - 3iB) \tan(c + dx)) dx + \\
 & \quad \frac{(-B + iA)(a^3 + ia^3 \tan(c + dx))}{d} - \frac{aA \cot(c + dx)(a + ia \tan(c + dx))^2}{d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.21. $\int \cot^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \int \frac{(i \tan(c+dx)a + a)(a^2(3iA+B) - a^2(A-3iB)\tan(c+dx))}{\tan(c+dx)} dx + \\
& \frac{(-B+iA)(a^3+ia^3\tan(c+dx))}{d} - \frac{aA \cot(c+dx)(a+ia\tan(c+dx))^2}{d} \\
& \quad \downarrow \text{4072} \\
& - \left(a^3(3B+iA) \int \tan(c+dx) dx \right) + \int \cot(c+dx) (a^3(3iA+B) - 4a^3(A-iB)\tan(c+dx)) dx + \\
& \frac{(-B+iA)(a^3+ia^3\tan(c+dx))}{d} - \frac{aA \cot(c+dx)(a+ia\tan(c+dx))^2}{d} \\
& \quad \downarrow \text{3042} \\
& - \left(a^3(3B+iA) \int \tan(c+dx) dx \right) + \int \frac{a^3(3iA+B) - 4a^3(A-iB)\tan(c+dx)}{\tan(c+dx)} dx + \\
& \frac{(-B+iA)(a^3+ia^3\tan(c+dx))}{d} - \frac{aA \cot(c+dx)(a+ia\tan(c+dx))^2}{d} \\
& \quad \downarrow \text{3956} \\
& \int \frac{a^3(3iA+B) - 4a^3(A-iB)\tan(c+dx)}{\tan(c+dx)} dx + \frac{(-B+iA)(a^3+ia^3\tan(c+dx))}{d} + \\
& \frac{a^3(3B+iA)\log(\cos(c+dx))}{d} - \frac{aA \cot(c+dx)(a+ia\tan(c+dx))^2}{d} \\
& \quad \downarrow \text{4014} \\
& a^3(B+3iA) \int \cot(c+dx) dx + \frac{(-B+iA)(a^3+ia^3\tan(c+dx))}{d} + \\
& \frac{a^3(3B+iA)\log(\cos(c+dx))}{d} - 4a^3x(A-iB) - \frac{aA \cot(c+dx)(a+ia\tan(c+dx))^2}{d} \\
& \quad \downarrow \text{3042} \\
& a^3(B+3iA) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx + \frac{(-B+iA)(a^3+ia^3\tan(c+dx))}{d} + \\
& \frac{a^3(3B+iA)\log(\cos(c+dx))}{d} - 4a^3x(A-iB) - \frac{aA \cot(c+dx)(a+ia\tan(c+dx))^2}{d} \\
& \quad \downarrow \text{25} \\
& -a^3(B+3iA) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx + \frac{(-B+iA)(a^3+ia^3\tan(c+dx))}{d} + \\
& \frac{a^3(3B+iA)\log(\cos(c+dx))}{d} - 4a^3x(A-iB) - \frac{aA \cot(c+dx)(a+ia\tan(c+dx))^2}{d} \\
& \quad \downarrow \text{3956} \\
& \frac{(-B+iA)(a^3+ia^3\tan(c+dx))}{d} + \frac{a^3(B+3iA)\log(-\sin(c+dx))}{d} + \\
& \frac{a^3(3B+iA)\log(\cos(c+dx))}{d} - 4a^3x(A-iB) - \frac{aA \cot(c+dx)(a+ia\tan(c+dx))^2}{d}
\end{aligned}$$

3.21. $\int \cot^2(c+dx)(a+ia\tan(c+dx))^3(A+B\tan(c+dx)) dx$

input `Int[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `-4*a^3*(A - I*B)*x + (a^3*(I*A + 3*B)*Log[Cos[c + d*x]])/d + (a^3*((3*I)*A + B)*Log[-Sin[c + d*x]])/d - (a*A*Cot[c + d*x]*(a + I*a*Tan[c + d*x])^2)/d + ((I*A - B)*(a^3 + I*a^3*Tan[c + d*x]))/d`

3.21.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4072 `Int[(((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*(d/b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c - a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0]`

```
rule 4076 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

```
rule 4077 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan
[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c -
a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] &&
GtQ[m, 1] && !LtQ[n, -1]
```

3.21.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.72

method	result
parallelrisch	$\frac{a^3(4iBdx+3iA \ln(\tan(dx+c))-2iA \ln(\sec^2(dx+c))-4Adx-iB \tan(dx+c)-A \cot(dx+c)+B \ln(\tan(dx+c))-2B \ln(\sec(dx+c)))}{d}$
derivativedivides	$\frac{a^3 \left(-A \cot(dx+c) + \frac{(-4iA-4B) \ln(\cot^2(dx+c)+1)}{2} + (-4iB+4A) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)) \right) + (iA+3B) \ln(\cot(dx+c)) - \frac{1}{\cot(dx+c)} \right)}{d}$
default	$\frac{a^3 \left(-A \cot(dx+c) + \frac{(-4iA-4B) \ln(\cot^2(dx+c)+1)}{2} + (-4iB+4A) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)) \right) + (iA+3B) \ln(\cot(dx+c)) - \frac{1}{\cot(dx+c)} \right)}{d}$
norman	$\frac{(4iB a^3 - 4A a^3)x \tan(dx+c) - \frac{A a^3}{d} - \frac{iB a^3 (\tan^2(dx+c))}{d}}{\tan(dx+c)} + \frac{(3iA a^3 + B a^3) \ln(\tan(dx+c))}{d} - \frac{2(iA a^3 + B a^3) \ln(1+\tan(dx+c))}{d}$
risch	$-\frac{8ia^3Bc}{d} + \frac{8a^3Ac}{d} + \frac{2a^3(-iA e^{2i(dx+c)} + B e^{2i(dx+c)} - iA - B)}{d(e^{2i(dx+c)}+1)(e^{2i(dx+c)}-1)} + \frac{3a^3 \ln(e^{2i(dx+c)}+1)B}{d} + \frac{ia^3 \ln(e^{2i(dx+c)}+1)}{d}$

```
input int(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

$$3.21. \int \cot^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

output $a^3(4iBdx+3iA\ln(\tan(dx+c))-2iA\ln(\sec(dx+c)^2)-4A*dx-iB*\tan(dx+c)-A*\cot(dx+c)+B*\ln(\tan(dx+c))-2B*\ln(\sec(dx+c)^2))/d$

3.21.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.22

$$\int \cot^2(c+dx)(a+ia\tan(c+dx))^3(A+B\tan(c+dx))dx = \frac{2(iA-B)a^3e^{(2i dx+2i c)} + 2(iA+B)a^3 - ((iA+3B)a^3e^{(4i dx+4i c)} + (-iA-3B)a^3) \log(e^{(2i dx+2i c)} - d)}{de^{(4i dx+4i c)} - d}$$

input `integrate(cot(d*x+c)^2*(a+i*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output $-(2*(I*A - B)*a^3*e^{(2*I*d*x + 2*I*c)} + 2*(I*A + B)*a^3 - ((I*A + 3*B)*a^3*e^{(4*I*d*x + 4*I*c)} + (-I*A - 3*B)*a^3)*\log(e^{(2*I*d*x + 2*I*c)} + 1) - ((3*I*A + B)*a^3*e^{(4*I*d*x + 4*I*c)} + (-3*I*A - B)*a^3)*\log(e^{(2*I*d*x + 2*I*c)} - 1))/(d*e^{(4*I*d*x + 4*I*c)} - d)$

3.21.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 219 vs. $2(99) = 198$.

Time = 1.01 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.89

$$\begin{aligned} & \int \cot^2(c+dx)(a+ia\tan(c+dx))^3(A+B\tan(c+dx))dx \\ &= \frac{ia^3(A-3iB)\log\left(e^{2idx} + \frac{2Aa^3-2iBa^3-a^3(A-3iB)}{Aa^3e^{2ic}+iBa^3e^{2ic}}\right)}{d} \\ &+ \frac{ia^3 \cdot (3A-iB)\log\left(e^{2idx} + \frac{2Aa^3-2iBa^3-a^3(3A-iB)}{Aa^3e^{2ic}+iBa^3e^{2ic}}\right)}{d} \\ &+ \frac{-2iAa^3-2Ba^3+(-2iAa^3e^{2ic}+2Ba^3e^{2ic})e^{2idx}}{de^{4ic}e^{4idx}-d} \end{aligned}$$

input `integrate(cot(d*x+c)**2*(a+i*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

3.21. $\int \cot^2(c+dx)(a+ia\tan(c+dx))^3(A+B\tan(c+dx))dx$

output $I*a^{3*(A - 3*I*B)*\log(\exp(2*I*d*x) + (2*A*a^{3*} - 2*I*B*a^{3*} - a^{3*(A - 3*I*B)})/(A*a^{3*}\exp(2*I*c) + I*B*a^{3*}\exp(2*I*c)))/d + I*a^{3*(3*A - I*B)*\log(\exp(2*I*d*x) + (2*A*a^{3*} - 2*I*B*a^{3*} - a^{3*(3*A - I*B)})/(A*a^{3*}\exp(2*I*c) + I*B*a^{3*}\exp(2*I*c)))/d + (-2*I*A*a^{3*} - 2*B*a^{3*} + (-2*I*A*a^{3*}\exp(2*I*c) + 2*B*a^{3*}\exp(2*I*c))*\exp(2*I*d*x))/(d*\exp(4*I*c)*\exp(4*I*d*x) - d)$

3.21.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.72

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{4(dx + c)(A - iB)a^3 + 2(iA + B)a^3 \log(\tan(dx + c)^2 + 1) - (3iA + B)a^3 \log(\tan(dx + c)) + iBa^3}{d}$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output $-(4*(d*x + c)*(A - I*B)*a^3 + 2*(I*A + B)*a^3*\log(\tan(d*x + c)^2 + 1) - (3*I*A + B)*a^3*\log(\tan(d*x + c)) + I*B*a^3*\tan(d*x + c) + A*a^3/\tan(d*x + c))/d$

3.21.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 257 vs. $2(104) = 208$.

Time = 1.19 (sec) , antiderivative size = 257, normalized size of antiderivative = 2.22

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{3Aa^3 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 6(iAa^3 + 3Ba^3) \log\left(\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 1\right) - 48(iAa^3 + Ba^3) \log\left(\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)\right)}{d}$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

3.21. $\int \cot^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

output $\frac{1}{6}(3Aa^3 \tan(1/2dx + 1/2c) + 6(IAa^3 + 3Ba^3) \log(\tan(1/2dx + 1/2c) + 1) - 48(IAa^3 + Ba^3) \log(\tan(1/2dx + 1/2c) + I) + 6(IAa^3 + 3Ba^3) \log(\tan(1/2dx + 1/2c) - 1) - 6(-3IAa^3 - Ba^3) \log(\tan(1/2dx + 1/2c)) + (-10IAa^3 \tan(1/2dx + 1/2c)^3 - 14Ba^3 \tan(1/2dx + 1/2c)^3 - 3Aa^3 \tan(1/2dx + 1/2c)^2 + 12IBa^3 \tan(1/2dx + 1/2c)^2 + 10IAa^3 \tan(1/2dx + 1/2c) + 14Ba^3 \tan(1/2dx + 1/2c) + 3Aa^3) / (\tan(1/2dx + 1/2c)^3 - \tan(1/2dx + 1/2c)) / d$

3.21.9 Mupad [B] (verification not implemented)

Time = 7.93 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.66

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{a^3 \ln(\tan(c + dx)) (B + A3i)}{d} - \frac{4a^3 \ln(\tan(c + dx) + 1i) (B + A1i)}{d}$$

$$- \frac{Aa^3 \cot(c + dx)}{d} - \frac{Ba^3 \tan(c + dx) 1i}{d}$$

input `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3,x)`

output $(a^3 \log(\tan(c + d*x))(A3i + B))/d - (4a^3 \log(\tan(c + d*x) + 1i)(A1i + B))/d - (Aa^3 \cot(c + d*x))/d - (Ba^3 \tan(c + d*x)1i)/d$

3.22 $\int \cot^3(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.22.1	Optimal result	449
3.22.2	Mathematica [A] (verified)	449
3.22.3	Rubi [A] (verified)	450
3.22.4	Maple [A] (verified)	453
3.22.5	Fricas [A] (verification not implemented)	454
3.22.6	Sympy [B] (verification not implemented)	454
3.22.7	Maxima [A] (verification not implemented)	455
3.22.8	Giac [B] (verification not implemented)	455
3.22.9	Mupad [B] (verification not implemented)	456

3.22.1 Optimal result

Integrand size = 34, antiderivative size = 123

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= -4a^3(iA+B)x + \frac{ia^3B \log(\cos(c+dx))}{d} - \frac{a^3(4A-3iB) \log(\sin(c+dx))}{d}$$

$$- \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^2}{2d} - \frac{(2iA+B) \cot(c+dx)(a^3+ia^3 \tan(c+dx))}{d}$$

```
output -4*a^3*(I*A+B)*x+I*a^3*B*ln(cos(d*x+c))/d-a^3*(4*A-3*I*B)*ln(sin(d*x+c))/d
-1/2*a*A*cot(d*x+c)^2*(a+I*a*tan(d*x+c))^2/d-(2*I*A+B)*cot(d*x+c)*(a^3+I*a
^3*tan(d*x+c))/d
```

3.22.2 Mathematica [A] (verified)

Time = 1.10 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.61

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{a^3((-6iA-2B) \cot(c+dx) - A \cot^2(c+dx) + (-8A+6iB) \log(\tan(c+dx)) + 8(A-iB) \log(i+\tan(c+dx)))}{2d}$$

input `Integrate[Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(a^3*(((6*I)*A - 2*B)*Cot[c + d*x] - A*Cot[c + d*x]^2 + (-8*A + (6*I)*B)*Log[Tan[c + d*x]] + 8*(A - I*B)*Log[I + Tan[c + d*x]]))/(2*d)`

3.22.3 Rubi [A] (verified)

Time = 0.90 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.02, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {3042, 4076, 27, 3042, 4076, 25, 3042, 4072, 3042, 3956, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^3(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\tan(c + dx)^3} dx \\
 & \quad \downarrow \text{4076} \\
 & \frac{1}{2} \int 2 \cot^2(c + dx)(i \tan(c + dx)a + a)^2(a(2iA + B) + iaB \tan(c + dx)) dx - \\
 & \quad \frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{27} \\
 & \int \cot^2(c + dx)(i \tan(c + dx)a + a)^2(a(2iA + B) + iaB \tan(c + dx)) dx - \\
 & \quad \frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(i \tan(c + dx)a + a)^2(a(2iA + B) + iaB \tan(c + dx))}{\tan(c + dx)^2} dx - \frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{4076} \\
 & \int -\cot(c + dx)(i \tan(c + dx)a + a)((4A - 3iB)a^2 + B \tan(c + dx)a^2) dx - \\
 & \quad \frac{(B + 2iA) \cot(c + dx)(a^3 + ia^3 \tan(c + dx))}{d} - \frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^2}{2d}
 \end{aligned}$$

3.22. $\int \cot^3(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \downarrow 25 \\
& - \int \cot(c+dx)(i \tan(c+dx)a+a)((4A-3iB)a^2+B \tan(c+dx)a^2) dx - \\
& \frac{(B+2iA) \cot(c+dx)(a^3+ia^3 \tan(c+dx))}{d} - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^2}{2d} \\
& \downarrow 3042 \\
& - \int \frac{(i \tan(c+dx)a+a)((4A-3iB)a^2+B \tan(c+dx)a^2)}{\tan(c+dx)} dx - \\
& \frac{(B+2iA) \cot(c+dx)(a^3+ia^3 \tan(c+dx))}{d} - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^2}{2d} \\
& \downarrow 4072 \\
& - \int \cot(c+dx)((4A-3iB)a^3+4(iA+B) \tan(c+dx)a^3) dx - ia^3B \int \tan(c+dx) dx - \\
& \frac{(B+2iA) \cot(c+dx)(a^3+ia^3 \tan(c+dx))}{d} - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^2}{2d} \\
& \downarrow 3042 \\
& - \int \frac{(4A-3iB)a^3+4(iA+B) \tan(c+dx)a^3}{\tan(c+dx)} dx - ia^3B \int \tan(c+dx) dx - \\
& \frac{(B+2iA) \cot(c+dx)(a^3+ia^3 \tan(c+dx))}{d} - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^2}{2d} \\
& \downarrow 3956 \\
& - \int \frac{(4A-3iB)a^3+4(iA+B) \tan(c+dx)a^3}{\tan(c+dx)} dx - \frac{(B+2iA) \cot(c+dx)(a^3+ia^3 \tan(c+dx))}{d} + \\
& \frac{ia^3B \log(\cos(c+dx))}{d} - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^2}{2d} \\
& \downarrow 4014 \\
& -a^3(4A-3iB) \int \cot(c+dx) dx - \frac{(B+2iA) \cot(c+dx)(a^3+ia^3 \tan(c+dx))}{d} - 4a^3x(B+iA) + \\
& \frac{ia^3B \log(\cos(c+dx))}{d} - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^2}{2d} \\
& \downarrow 3042 \\
& -a^3(4A-3iB) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx - \frac{(B+2iA) \cot(c+dx)(a^3+ia^3 \tan(c+dx))}{d} - \\
& 4a^3x(B+iA) + \frac{ia^3B \log(\cos(c+dx))}{d} - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^2}{2d} \\
& \downarrow 25
\end{aligned}$$

$$\begin{aligned}
& a^3(4A - 3iB) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - \frac{(B + 2iA) \cot(c + dx) (a^3 + ia^3 \tan(c + dx))}{d} - \\
& 4a^3x(B + iA) + \frac{ia^3B \log(\cos(c + dx))}{d} - \frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^2}{2d} \\
& \qquad \qquad \qquad \downarrow \text{3956} \\
& -\frac{a^3(4A - 3iB) \log(-\sin(c + dx))}{d} - \frac{(B + 2iA) \cot(c + dx) (a^3 + ia^3 \tan(c + dx))}{d} - 4a^3x(B + \\
& iA) + \frac{ia^3B \log(\cos(c + dx))}{d} - \frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^2}{2d}
\end{aligned}$$

input `Int[Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `-4*a^3*(I*A + B)*x + (I*a^3*B*Log[Cos[c + d*x]])/d - (a^3*(4*A - (3*I)*B)*Log[-Sin[c + d*x]])/d - (a*A*Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^2)/(2*d) - (((2*I)*A + B)*Cot[c + d*x]*(a^3 + I*a^3*Tan[c + d*x]))/d`

3.22.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4072 `Int[(((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*(d/b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c - a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0]`

rule 4076 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.22.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.75

method	result
derivativedivides	$-\frac{a^3 \left(\frac{(4iB-4A) \ln(1+\tan^2(dx+c))}{2} + (4iA+4B) \arctan(\tan(dx+c)) + (-3iB+4A) \ln(\tan(dx+c)) - \frac{-3iA-B}{\tan(dx+c)} + \frac{A}{2 \tan(dx+c)} \right)}{d}$
default	$-\frac{a^3 \left(\frac{(4iB-4A) \ln(1+\tan^2(dx+c))}{2} + (4iA+4B) \arctan(\tan(dx+c)) + (-3iB+4A) \ln(\tan(dx+c)) - \frac{-3iA-B}{\tan(dx+c)} + \frac{A}{2 \tan(dx+c)} \right)}{d}$
parallelrisch	$-\frac{a^3 (8iAxd - 6iB \ln(\tan(dx+c)) + 4iB \ln(\sec^2(dx+c)) + 8Bdx + 6iA \cot(dx+c) + 8A \ln(\tan(dx+c)) - 4A \ln(\sec^2(dx+c)))}{2d}$
norman	$\frac{(-4iA a^3 - 4B a^3)x \tan^2(dx+c) - \frac{A a^3}{2d} - \frac{(3iA a^3 + B a^3) \tan(dx+c)}{d}}{\tan(dx+c)^2} - \frac{(-3iB a^3 + 4A a^3) \ln(\tan(dx+c))}{d} + \frac{2(-iB a^3)}{d}$
risch	$\frac{8a^3 Bc}{d} + \frac{8ia^3 Ac}{d} - \frac{2ia^3 (4iA e^{2i(dx+c)} + B e^{2i(dx+c)} - 3iA - B)}{d(e^{2i(dx+c)} - 1)^2} + \frac{3ia^3 \ln(e^{2i(dx+c)} - 1)B}{d} - \frac{4A a^3 \ln(e^{2i(dx+c)} - 1)}{d}$

input `int(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `-a^3/d*(1/2*(4*I*B-4*A)*ln(1+tan(d*x+c)^2)+(4*I*A+4*B)*arctan(tan(d*x+c))+ (4*A-3*I*B)*ln(tan(d*x+c))-(-3*I*A-B)/tan(d*x+c)+1/2*A/tan(d*x+c)^2)`

3.22. $\int \cot^3(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.22.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.46

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{2(4A - iB)a^3 e^{(2i dx + 2i c)} - 2(3A - iB)a^3 + (iBa^3 e^{(4i dx + 4i c)} - 2iBa^3 e^{(2i dx + 2i c)} + iBa^3) \log(e^{(2i dx + 2i c)})}{de^{(4i dx + 4i c)} - 2d}$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output $(2*(4*A - I*B)*a^3*e^{(2*I*d*x + 2*I*c)} - 2*(3*A - I*B)*a^3 + (I*B*a^3*e^{(4*I*d*x + 4*I*c)} - 2*I*B*a^3*e^{(2*I*d*x + 2*I*c)} + I*B*a^3)*\log(e^{(2*I*d*x + 2*I*c)} + 1) - ((4*A - 3*I*B)*a^3*e^{(4*I*d*x + 4*I*c)} - 2*(4*A - 3*I*B)*a^3*e^{(2*I*d*x + 2*I*c)} + (4*A - 3*I*B)*a^3)*\log(e^{(2*I*d*x + 2*I*c)} - 1))/(d*e^{(4*I*d*x + 4*I*c)} - 2*d*e^{(2*I*d*x + 2*I*c)} + d)$

3.22.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 226 vs. $2(109) = 218$.

Time = 1.03 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.84

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{iBa^3 \log\left(\frac{2Aa^3 - iBa^3}{2Aa^3 e^{2ic} - iBa^3 e^{2ic}} + e^{2idx}\right)}{d} - \frac{a^3 \cdot (4A - 3iB) \log\left(e^{2idx} + \frac{2Aa^3 - 2iBa^3 - a^3 \cdot (4A - 3iB)}{2Aa^3 e^{2ic} - iBa^3 e^{2ic}}\right)}{d}$$

$$+ \frac{-6Aa^3 + 2iBa^3 + (8Aa^3 e^{2ic} - 2iBa^3 e^{2ic}) e^{2idx}}{de^{4ic} e^{4idx} - 2de^{2ic} e^{2idx} + d}$$

input `integrate(cot(d*x+c)**3*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output $I*B*a**3*\log((2*A*a**3 - I*B*a**3)/(2*A*a**3*\exp(2*I*c) - I*B*a**3*\exp(2*I*c)) + \exp(2*I*d*x))/d - a**3*(4*A - 3*I*B)*\log(\exp(2*I*d*x) + (2*A*a**3 - 2*I*B*a**3 - a**3*(4*A - 3*I*B))/(2*A*a**3*\exp(2*I*c) - I*B*a**3*\exp(2*I*c)))/d + (-6*A*a**3 + 2*I*B*a**3 + (8*A*a**3*\exp(2*I*c) - 2*I*B*a**3*\exp(2*I*c))*\exp(2*I*d*x))/(d*\exp(4*I*c)*\exp(4*I*d*x) - 2*d*\exp(2*I*c)*\exp(2*I*d*x) + d)$

3.22. $\int \cot^3(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.22.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.78

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{8(dx + c)(iA + B)a^3 - 4(A - iB)a^3 \log(\tan(dx + c)^2 + 1) + 2(4A - 3iB)a^3 \log(\tan(dx + c)) - 2Aa^3}{2d}$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(8*(d*x + c)*(I*A + B)*a^3 - 4*(A - I*B)*a^3*log(tan(d*x + c)^2 + 1) + 2*(4*A - 3*I*B)*a^3*log(tan(d*x + c)) - (2*(-3*I*A - B)*a^3*tan(d*x + c) - A*a^3)/tan(d*x + c)^2)/d`

3.22.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 223 vs. 2(109) = 218.

Time = 1.14 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.81

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{Aa^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 8iBa^3 \log\left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right) - 8iBa^3 \log\left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 1\right) - 12iAa^3}{2d}$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `-1/8*(A*a^3*tan(1/2*d*x + 1/2*c)^2 - 8*I*B*a^3*log(tan(1/2*d*x + 1/2*c) + 1) - 8*I*B*a^3*log(tan(1/2*d*x + 1/2*c) - 1) - 12*I*A*a^3*tan(1/2*d*x + 1/2*c) - 4*B*a^3*tan(1/2*d*x + 1/2*c) - 64*(A*a^3 - I*B*a^3)*log(tan(1/2*d*x + 1/2*c) + I) + 8*(4*A*a^3 - 3*I*B*a^3)*log(tan(1/2*d*x + 1/2*c)) - (48*A*a^3*tan(1/2*d*x + 1/2*c)^2 - 36*I*B*a^3*tan(1/2*d*x + 1/2*c)^2 - 12*I*A*a^3*tan(1/2*d*x + 1/2*c) - 4*B*a^3*tan(1/2*d*x + 1/2*c) - A*a^3)/tan(1/2*d*x + 1/2*c)^2)/d`

3.22. $\int \cot^3(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.22.9 Mupad [B] (verification not implemented)

Time = 8.12 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.72

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= -\frac{\frac{Aa^3}{2} + \tan(c + dx)(Ba^3 + Aa^3 3i)}{d \tan(c + dx)^2} - \frac{a^3 \ln(\tan(c + dx))(4A - B 3i)}{d}$$

$$+ \frac{4a^3 \ln(\tan(c + dx) + 1i)(A - B 1i)}{d}$$

input `int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3,x)`output `(4*a^3*log(tan(c + d*x) + 1i)*(A - B*1i))/d - (a^3*log(tan(c + d*x))*(4*A - B*3i))/d - ((A*a^3)/2 + tan(c + d*x)*(A*a^3*3i + B*a^3))/(d*tan(c + d*x)^2)`

3.23 $\int \cot^4(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.23.1	Optimal result	457
3.23.2	Mathematica [A] (verified)	458
3.23.3	Rubi [A] (verified)	458
3.23.4	Maple [A] (verified)	462
3.23.5	Fricas [A] (verification not implemented)	462
3.23.6	Sympy [A] (verification not implemented)	463
3.23.7	Maxima [A] (verification not implemented)	463
3.23.8	Giac [B] (verification not implemented)	464
3.23.9	Mupad [B] (verification not implemented)	464

3.23.1 Optimal result

Integrand size = 34, antiderivative size = 134

$$\int \cot^4(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= 4a^3(A-iB)x + \frac{a^3(17A-15iB) \cot(c+dx)}{6d}$$

$$- \frac{4a^3(iA+B) \log(\sin(c+dx))}{d} - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^2}{3d}$$

$$- \frac{(5iA+3B) \cot^2(c+dx)(a^3+ia^3 \tan(c+dx))}{6d}$$

```
output 4*a^3*(A-I*B)*x+1/6*a^3*(17*A-15*I*B)*cot(d*x+c)/d-4*a^3*(I*A+B)*ln(sin(d*x+c))/d-1/3*a*A*cot(d*x+c)^3*(a+I*a*tan(d*x+c))^2/d-1/6*(5*I*A+3*B)*cot(d*x+c)^2*(a^3+I*a^3*tan(d*x+c))/d
```


3.23.2 Mathematica [A] (verified)

Time = 1.04 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.66

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{a^3(6(4A - 3iB) \cot(c + dx) + (-9iA - 3B) \cot^2(c + dx) - 2A \cot^3(c + dx) - 24i(A - iB)(\log(\tan(c + dx) + \cot(c + dx))))}{6d}$$

input `Integrate[Cot[c + d*x]^4*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(a^3*(6*(4*A - (3*I)*B)*Cot[c + d*x] + ((-9*I)*A - 3*B)*Cot[c + d*x]^2 - 2*A*Cot[c + d*x]^3 - (24*I)*(A - I*B)*(Log[Tan[c + d*x]] - Log[I + Tan[c + d*x]])))/(6*d)`

3.23.3 Rubi [A] (verified)

Time = 0.92 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.08, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.382$, Rules used = {3042, 4076, 3042, 4076, 25, 3042, 4074, 27, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\tan(c + dx)^4} dx$$

$$\downarrow 4076$$

$$\frac{1}{3} \int \cot^3(c + dx)(i \tan(c + dx)a + a)^2(a(5iA + 3B) - a(A - 3iB) \tan(c + dx)) dx - \frac{aA \cot^3(c + dx)(a + ia \tan(c + dx))^2}{3d}$$

$$\downarrow 3042$$

$$\frac{1}{3} \int \frac{(i \tan(c + dx)a + a)^2(a(5iA + 3B) - a(A - 3iB) \tan(c + dx))}{\tan(c + dx)^3} dx - \frac{aA \cot^3(c + dx)(a + ia \tan(c + dx))^2}{3d}$$

3.23. $\int \cot^4(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\downarrow 4076$$

$$\frac{1}{3} \left(\frac{1}{2} \int -\cot^2(c+dx)(i \tan(c+dx)a+a) ((17A-15iB)a^2+(7iA+9B)\tan(c+dx)a^2) dx - \frac{(3B+5iA)\cot^2(c+dx)}{2d} \right) - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^2}{3d}$$

$$\downarrow 25$$

$$\frac{1}{3} \left(-\frac{1}{2} \int \cot^2(c+dx)(i \tan(c+dx)a+a) ((17A-15iB)a^2+(7iA+9B)\tan(c+dx)a^2) dx - \frac{(3B+5iA)\cot^2(c+dx)}{2d} \right) - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^2}{3d}$$

$$\downarrow 3042$$

$$\frac{1}{3} \left(-\frac{1}{2} \int \frac{(i \tan(c+dx)a+a) ((17A-15iB)a^2+(7iA+9B)\tan(c+dx)a^2)}{\tan(c+dx)^2} dx - \frac{(3B+5iA)\cot^2(c+dx)(a^3+a)}{2d} \right) - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^2}{3d}$$

$$\downarrow 4074$$

$$\frac{1}{3} \left(\frac{1}{2} \left(\frac{a^3(17A-15iB)\cot(c+dx)}{d} - \int 24 \cot(c+dx) (a^3(iA+B) - a^3(A-iB)\tan(c+dx)) dx \right) - \frac{(3B+5iA)\cot^2(c+dx)}{2d} \right) - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^2}{3d}$$

$$\downarrow 27$$

$$\frac{1}{3} \left(\frac{1}{2} \left(\frac{a^3(17A-15iB)\cot(c+dx)}{d} - 24 \int \cot(c+dx) (a^3(iA+B) - a^3(A-iB)\tan(c+dx)) dx \right) - \frac{(3B+5iA)\cot^2(c+dx)}{2d} \right) - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^2}{3d}$$

$$\downarrow 3042$$

$$\frac{1}{3} \left(\frac{1}{2} \left(\frac{a^3(17A-15iB)\cot(c+dx)}{d} - 24 \int \frac{a^3(iA+B) - a^3(A-iB)\tan(c+dx)}{\tan(c+dx)} dx \right) - \frac{(3B+5iA)\cot^2(c+dx)}{2d} \right) - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^2}{3d}$$

$$\downarrow 4014$$

3.23. $\int \cot^4(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\frac{1}{3} \left(\frac{1}{2} \left(\frac{a^3(17A - 15iB) \cot(c + dx)}{d} - 24 \left(a^3(B + iA) \int \cot(c + dx) dx - a^3x(A - iB) \right) \right) - \frac{(3B + 5iA) \cot^2(c + dx)}{3d} \right) \frac{aA \cot^3(c + dx)(a + ia \tan(c + dx))^2}{3d}$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{2} \left(\frac{a^3(17A - 15iB) \cot(c + dx)}{d} - 24 \left(a^3(B + iA) \int -\tan \left(c + dx + \frac{\pi}{2} \right) dx - a^3x(A - iB) \right) \right) - \frac{(3B + 5iA) \cot^2(c + dx)}{3d} \right) \frac{aA \cot^3(c + dx)(a + ia \tan(c + dx))^2}{3d}$$

↓ 25

$$\frac{1}{3} \left(\frac{1}{2} \left(\frac{a^3(17A - 15iB) \cot(c + dx)}{d} - 24 \left(-a^3(B + iA) \int \tan \left(\frac{1}{2}(2c + \pi) + dx \right) dx - (a^3x(A - iB)) \right) \right) - \frac{(3B + 5iA) \cot^2(c + dx)}{3d} \right) \frac{aA \cot^3(c + dx)(a + ia \tan(c + dx))^2}{3d}$$

↓ 3956

$$\frac{1}{3} \left(\frac{1}{2} \left(\frac{a^3(17A - 15iB) \cot(c + dx)}{d} - 24 \left(\frac{a^3(B + iA) \log(-\sin(c + dx))}{d} - a^3x(A - iB) \right) \right) - \frac{(3B + 5iA) \cot^2(c + dx)}{3d} \right) \frac{aA \cot^3(c + dx)(a + ia \tan(c + dx))^2}{3d}$$

input `Int[Cot[c + d*x]^4*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `-1/3*(a*A*Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^2)/d + (((a^3*(17*A - (15*I)*B)*Cot[c + d*x])/d - 24*(-(a^3*(A - I*B)*x) + (a^3*(I*A + B)*Log[-Sin[c + d*x]]))/d))/2 - (((5*I)*A + 3*B)*Cot[c + d*x]^2*(a^3 + I*a^3*Tan[c + d*x]))/(2*d))/3`

3.23.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`
- rule 4074 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`
- rule 4076 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.23.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.78

method	result
parallelrisc	$\frac{4a^3 \left(\left(-\frac{iA}{2} - \frac{B}{2} \right) \ln(\sec^2(dx+c)) + (iA+B) \ln(\tan(dx+c)) + \frac{13A(\cot^3(dx+c))}{12} + \cot^2(dx+c) \left(\frac{3iA}{8} + \frac{B}{8} \right) + (-A \csc^2(dx+c)) \right)}{d}$
derivativedivides	$\frac{a^3 \left(-\frac{3iA(\cot^2(dx+c))}{2} - \frac{A(\cot^3(dx+c))}{3} - 3iB \cot(dx+c) - \frac{B(\cot^2(dx+c))}{2} + 4A \cot(dx+c) + \frac{(4iA+4B) \ln(\cot^2(dx+c)+1)}{2} \right)}{d}$
default	$\frac{a^3 \left(-\frac{3iA(\cot^2(dx+c))}{2} - \frac{A(\cot^3(dx+c))}{3} - 3iB \cot(dx+c) - \frac{B(\cot^2(dx+c))}{2} + 4A \cot(dx+c) + \frac{(4iA+4B) \ln(\cot^2(dx+c)+1)}{2} \right)}{d}$
risc	$\frac{8ia^3Bc}{d} - \frac{8a^3Ac}{d} + \frac{2a^3(24iAe^{4i(dx+c)} + 12Be^{4i(dx+c)} - 33iAe^{2i(dx+c)} - 21Be^{2i(dx+c)} + 13iA + 9B)}{3d(e^{2i(dx+c)} - 1)^3} - \frac{4a^3 \ln(e^{2i(dx+c)} + 1)}{d}$
norman	$\frac{\left(-3iB a^3 + 4A a^3 \right) \left(\tan^2(dx+c) \right)}{d} + \frac{\left(-4iB a^3 + 4A a^3 \right) x \left(\tan^3(dx+c) \right) - \frac{A a^3}{3d} - \frac{\left(3iA a^3 + B a^3 \right) \tan(dx+c)}{2d}}{\tan(dx+c)^3} - \frac{4(iA a^3 + B a^3) \ln(\tan(dx+c))}{d}$

input `int(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `-4*a^3*((-1/2*I*A-1/2*B)*ln(sec(d*x+c)^2)+(I*A+B)*ln(tan(d*x+c))+13/12*A*cot(d*x+c)^3+cot(d*x+c)^2*(3/8*I*A+1/8*B)+(-A*csc(d*x+c)^2+3/4*I*B)*cot(d*x+c)+x*d*(-A+I*B))/d`

3.23.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.35

$$\int \cot^4(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx = \frac{2(12(-2iA-B)a^3e^{4i dx+4i c} + 3(11iA+7B)a^3e^{2i dx+2i c} + (-13iA-9B)a^3 + 6((iA+B)a^3e^{6i dx+6i c} - 3de^{6i dx+6i c} - 3de^{4i dx+4i c}))}{3(de^{6i dx+6i c} - 3de^{4i dx+4i c})}$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output
$$\begin{aligned} & -2/3*(12*(-2*I*A - B)*a^3*e^(4*I*d*x + 4*I*c) + 3*(11*I*A + 7*B)*a^3*e^(2* \\ & I*d*x + 2*I*c) + (-13*I*A - 9*B)*a^3 + 6*((I*A + B)*a^3*e^(6*I*d*x + 6*I*c \\ &) + 3*(-I*A - B)*a^3*e^(4*I*d*x + 4*I*c) + 3*(I*A + B)*a^3*e^(2*I*d*x + 2* \\ & I*c) + (-I*A - B)*a^3)*\log(e^(2*I*d*x + 2*I*c) - 1))/(d*e^(6*I*d*x + 6*I*c \\ &) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d) \end{aligned}$$

3.23.6 Sympy [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.36

$$\begin{aligned} & \int \cot^4(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ & = -\frac{4ia^3(A - iB) \log(e^{2idx} - e^{-2ic})}{d} \\ & + \frac{26iAa^3 + 18Ba^3 + (-66iAa^3e^{2ic} - 42Ba^3e^{2ic})e^{2idx} + (48iAa^3e^{4ic} + 24Ba^3e^{4ic})e^{4idx}}{3de^{6ic}e^{6idx} - 9de^{4ic}e^{4idx} + 9de^{2ic}e^{2idx} - 3d} \end{aligned}$$

input `integrate(cot(d*x+c)**4*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output
$$\begin{aligned} & -4*I*a**3*(A - I*B)*\log(\exp(2*I*d*x) - \exp(-2*I*c))/d + (26*I*A*a**3 + 18* \\ & B*a**3 + (-66*I*A*a**3*\exp(2*I*c) - 42*B*a**3*\exp(2*I*c))*\exp(2*I*d*x) + (\\ & 48*I*A*a**3*\exp(4*I*c) + 24*B*a**3*\exp(4*I*c))*\exp(4*I*d*x))/(3*d*\exp(6*I* \\ & c)*\exp(6*I*d*x) - 9*d*\exp(4*I*c)*\exp(4*I*d*x) + 9*d*\exp(2*I*c)*\exp(2*I*d*x \\ &) - 3*d) \end{aligned}$$

3.23.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.86

$$\begin{aligned} & \int \cot^4(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ & = \frac{24(dx + c)(A - iB)a^3 - 12(-iA - B)a^3 \log(\tan(dx + c)^2 + 1) - 24(iA + B)a^3 \log(\tan(dx + c)) + 6}{6d} \end{aligned}$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output $1/6*(24*(d*x + c)*(A - I*B)*a^3 - 12*(-I*A - B)*a^3*\log(\tan(d*x + c))^2 + 1) - 24*(I*A + B)*a^3*\log(\tan(d*x + c)) + (6*(4*A - 3*I*B)*a^3*\tan(d*x + c)^2 + 3*(-3*I*A - B)*a^3*\tan(d*x + c) - 2*A*a^3)/\tan(d*x + c)^3/d$

3.23.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 254 vs. $2(116) = 232$.

Time = 0.76 (sec) , antiderivative size = 254, normalized size of antiderivative = 1.90

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{Aa^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 9i Aa^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 3Ba^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 51Aa^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 36Aa^3}{d}$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output $1/24*(A*a^3*\tan(1/2*d*x + 1/2*c)^3 - 9*I*A*a^3*\tan(1/2*d*x + 1/2*c)^2 - 3*B*a^3*\tan(1/2*d*x + 1/2*c)^2 - 51*A*a^3*\tan(1/2*d*x + 1/2*c) + 36*I*B*a^3*\tan(1/2*d*x + 1/2*c) - 192*(-I*A*a^3 - B*a^3)*\log(\tan(1/2*d*x + 1/2*c) + I) - 96*(I*A*a^3 + B*a^3)*\log(\tan(1/2*d*x + 1/2*c)) - (-176*I*A*a^3*\tan(1/2*d*x + 1/2*c)^3 - 176*B*a^3*\tan(1/2*d*x + 1/2*c)^3 - 51*A*a^3*\tan(1/2*d*x + 1/2*c)^2 + 36*I*B*a^3*\tan(1/2*d*x + 1/2*c)^2 + 9*I*A*a^3*\tan(1/2*d*x + 1/2*c) + 3*B*a^3*\tan(1/2*d*x + 1/2*c) + A*a^3)/\tan(1/2*d*x + 1/2*c)^3/d$

3.23.9 Mupad [B] (verification not implemented)

Time = 7.34 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.69

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= -\frac{\frac{Aa^3}{3} - \tan(c + dx)^2(4Aa^3 - Ba^3 3i) + \tan(c + dx)\left(\frac{Ba^3}{2} + \frac{Aa^3 3i}{2}\right)}{d \tan(c + dx)^3} - \frac{a^3 \operatorname{atan}(2 \tan(c + dx) + i) (B + A i) 8i}{d}$$

3.23. $\int \cot^4(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

input `int(cot(c + d*x)^4*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3,x)`

output `- ((A*a^3)/3 - tan(c + d*x)^2*(4*A*a^3 - B*a^3*3i) + tan(c + d*x)*((A*a^3*3i)/2 + (B*a^3)/2))/(d*tan(c + d*x)^3) - (a^3*atan(2*tan(c + d*x) + 1i)*(A*1i + B)*8i)/d`

3.24 $\int \cot^5(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.24.1	Optimal result	466
3.24.2	Mathematica [A] (verified)	467
3.24.3	Rubi [A] (verified)	467
3.24.4	Maple [A] (verified)	472
3.24.5	Fricas [A] (verification not implemented)	472
3.24.6	Sympy [A] (verification not implemented)	473
3.24.7	Maxima [A] (verification not implemented)	473
3.24.8	Giac [B] (verification not implemented)	474
3.24.9	Mupad [B] (verification not implemented)	475

3.24.1 Optimal result

Integrand size = 34, antiderivative size = 157

$$\int \cot^5(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= 4a^3(iA+B)x + \frac{4a^3(iA+B) \cot(c+dx)}{d} + \frac{a^3(15A-14iB) \cot^2(c+dx)}{12d}$$

$$+ \frac{4a^3(A-iB) \log(\sin(c+dx))}{d} - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^2}{4d}$$

$$- \frac{(3iA+2B) \cot^3(c+dx)(a^3+ia^3 \tan(c+dx))}{6d}$$

output

```
4*a^3*(I*A+B)*x+4*a^3*(I*A+B)*cot(d*x+c)/d+1/12*a^3*(15*A-14*I*B)*cot(d*x+c)^2/d+4*a^3*(A-I*B)*ln(sin(d*x+c))/d-1/4*a*A*cot(d*x+c)^4*(a+I*a*tan(d*x+c))^2/d-1/6*(3*I*A+2*B)*cot(d*x+c)^3*(a^3+I*a^3*tan(d*x+c))/d
```

3.24.2 Mathematica [A] (verified)

Time = 1.45 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.71

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{ia^3(-((3A - 4iB)(i + \cot(c + dx))^3) + 3iA \cot(c + dx)(i + \cot(c + dx))^3 - 6i(A - iB)(6i \cot(c + dx) + \cot^2(c + dx)))}{12d}$$

input `Integrate[Cot[c + d*x]^5*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `((I/12)*a^3*(-((3*A - (4*I)*B)*(I + Cot[c + d*x])^3) + (3*I)*A*Cot[c + d*x]*(I + Cot[c + d*x])^3 - (6*I)*(A - I*B)*((6*I)*Cot[c + d*x] + Cot[c + d*x])^2 + 8*Log[Tan[c + d*x]] - 8*Log[I + Tan[c + d*x]]))/d`

3.24.3 Rubi [A] (verified)

Time = 1.09 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.10, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4076, 27, 3042, 4076, 25, 3042, 4074, 27, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\tan(c + dx)^5} dx$$

$$\downarrow \text{4076}$$

$$\frac{1}{4} \int 2 \cot^4(c + dx)(i \tan(c + dx)a + a)^2(a(3iA + 2B) - a(A - 2iB) \tan(c + dx)) dx - \frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^2}{4d}$$

$$\downarrow \text{27}$$

$$\frac{1}{2} \int \cot^4(c + dx)(i \tan(c + dx)a + a)^2(a(3iA + 2B) - a(A - 2iB) \tan(c + dx)) dx - \frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^2}{4d}$$

3.24. $\int \cot^5(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{1}{2} \int \frac{(i \tan(c+dx)a+a)^2(a(3iA+2B)-a(A-2iB)\tan(c+dx))}{\tan(c+dx)^4} dx - \\ & \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^2}{4d} \\ & \downarrow 4076 \end{aligned}$$

$$\begin{aligned} & \frac{1}{2} \left(\frac{1}{3} \int -\cot^3(c+dx)(i \tan(c+dx)a+a) ((15A-14iB)a^2+(9iA+10B)\tan(c+dx)a^2) dx - \frac{(2B+3iA)\cot^3}{3d} \right) \\ & \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^2}{4d} \\ & \downarrow 25 \end{aligned}$$

$$\begin{aligned} & \frac{1}{2} \left(-\frac{1}{3} \int \cot^3(c+dx)(i \tan(c+dx)a+a) ((15A-14iB)a^2+(9iA+10B)\tan(c+dx)a^2) dx - \frac{(2B+3iA)\cot^3}{3d} \right) \\ & \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^2}{4d} \\ & \downarrow 3042 \end{aligned}$$

$$\begin{aligned} & \frac{1}{2} \left(-\frac{1}{3} \int \frac{(i \tan(c+dx)a+a) ((15A-14iB)a^2+(9iA+10B)\tan(c+dx)a^2)}{\tan(c+dx)^3} dx - \frac{(2B+3iA)\cot^3(c+dx)(a^3)}{3d} \right) \\ & \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^2}{4d} \\ & \downarrow 4074 \end{aligned}$$

$$\begin{aligned} & \frac{1}{2} \left(\frac{1}{3} \left(\frac{a^3(15A-14iB)\cot^2(c+dx)}{2d} - \int 24 \cot^2(c+dx) (a^3(iA+B)-a^3(A-iB)\tan(c+dx)) dx \right) - \frac{(2B+3iA)\cot^3}{3d} \right) \\ & \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^2}{4d} \\ & \downarrow 27 \end{aligned}$$

$$\begin{aligned} & \frac{1}{2} \left(\frac{1}{3} \left(\frac{a^3(15A-14iB)\cot^2(c+dx)}{2d} - 24 \int \cot^2(c+dx) (a^3(iA+B)-a^3(A-iB)\tan(c+dx)) dx \right) - \frac{(2B+3iA)\cot^3}{3d} \right) \\ & \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^2}{4d} \\ & \downarrow 3042 \end{aligned}$$

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{a^3(15A - 14iB) \cot^2(c + dx)}{2d} - 24 \int \frac{a^3(iA + B) - a^3(A - iB) \tan(c + dx)}{\tan(c + dx)^2} dx \right) - \frac{(2B + 3iA) \cot^3(c + dx)}{3d} \right) - \frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^2}{4d}$$

↓ 4012

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{a^3(15A - 14iB) \cot^2(c + dx)}{2d} - 24 \left(\int -\cot(c + dx) ((A - iB)a^3 + (iA + B) \tan(c + dx)a^3) dx - \frac{a^3(B + iA) \cot(c + dx)}{d} \right) \right) - \frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^2}{4d} \right)$$

↓ 25

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{a^3(15A - 14iB) \cot^2(c + dx)}{2d} - 24 \left(- \int \cot(c + dx) ((A - iB)a^3 + (iA + B) \tan(c + dx)a^3) dx - \frac{a^3(B + iA) \cot(c + dx)}{d} \right) \right) - \frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^2}{4d} \right)$$

↓ 3042

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{a^3(15A - 14iB) \cot^2(c + dx)}{2d} - 24 \left(- \int \frac{(A - iB)a^3 + (iA + B) \tan(c + dx)a^3}{\tan(c + dx)} dx - \frac{a^3(B + iA) \cot(c + dx)}{d} \right) \right) - \frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^2}{4d} \right)$$

↓ 4014

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{a^3(15A - 14iB) \cot^2(c + dx)}{2d} - 24 \left(-a^3(A - iB) \int \cot(c + dx) dx - \frac{a^3(B + iA) \cot(c + dx)}{d} - (a^3x(B + iA) \cot(c + dx))' \right) \right) - \frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^2}{4d} \right)$$

↓ 3042

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{a^3(15A - 14iB) \cot^2(c + dx)}{2d} - 24 \left(-a^3(A - iB) \int -\tan \left(c + dx + \frac{\pi}{2} \right) dx - \frac{a^3(B + iA) \cot(c + dx)}{d} - (a^3x(B + iA) \cot(c + dx))' \right) \right) - \frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^2}{4d} \right)$$

↓ 25

3.24. $\int \cot^5(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{a^3(15A - 14iB) \cot^2(c + dx)}{2d} - 24 \left(a^3(A - iB) \int \tan \left(\frac{1}{2}(2c + \pi) + dx \right) dx - \frac{a^3(B + iA) \cot(c + dx)}{d} - \frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^2}{4d} \right) \right) \right)$$

↓ 3956

$$\frac{1}{2} \left(\frac{1}{3} \left(\frac{a^3(15A - 14iB) \cot^2(c + dx)}{2d} - 24 \left(-\frac{a^3(B + iA) \cot(c + dx)}{d} - \frac{a^3(A - iB) \log(-\sin(c + dx))}{d} - (a^3 x(B + iA) \cot(c + dx) - \frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^2}{4d}) \right) \right) \right)$$

input `Int[Cot[c + d*x]^5*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `-1/4*(a*A*Cot[c + d*x]^4*(a + I*a*Tan[c + d*x])^2)/d + (((a^3*(15*A - (14*I)*B)*Cot[c + d*x]^2)/(2*d) - 24*(-(a^3*(I*A + B)*x) - (a^3*(I*A + B)*Cot[c + d*x])/d - (a^3*(A - I*B)*Log[-Sin[c + d*x]]/d))/3 - (((3*I)*A + 2*B)*Cot[c + d*x]^3*(a^3 + I*a^3*Tan[c + d*x]))/(3*d))/2`

3.24.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4074 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

rule 4076 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.24.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.69

method	result
parallelrisc	$\frac{4a^3 \left(\left(-\frac{A}{2} + \frac{iB}{2} \right) \ln(\sec^2(dx+c)) + (-iB+A) \ln(\tan(dx+c)) - \frac{A(\cot^4(dx+c))}{16} + (\cot^3(dx+c)) \left(-\frac{iA}{4} - \frac{B}{12} \right) + (\cot^2(dx+c)) \right)}{d}$
derivativedivides	$\frac{a^3 \left(-iA(\cot^3(dx+c)) - \frac{A(\cot^4(dx+c))}{4} - \frac{3iB(\cot^2(dx+c))}{2} - \frac{B(\cot^3(dx+c))}{3} + 4iA \cot(dx+c) + 2A(\cot^2(dx+c)) + 4 \cot(dx+c) \right)}{d}$
default	$\frac{a^3 \left(-iA(\cot^3(dx+c)) - \frac{A(\cot^4(dx+c))}{4} - \frac{3iB(\cot^2(dx+c))}{2} - \frac{B(\cot^3(dx+c))}{3} + 4iA \cot(dx+c) + 2A(\cot^2(dx+c)) + 4 \cot(dx+c) \right)}{d}$
risc	$-\frac{8a^3 Bc}{d} - \frac{8ia^3 Ac}{d} + \frac{2ia^3 (36iA e^{6i(dx+c)} + 24B e^{6i(dx+c)} - 69iA e^{4i(dx+c)} - 57B e^{4i(dx+c)} + 54iA e^{2i(dx+c)} + 46B e^{2i(dx+c)})}{3d(e^{2i(dx+c)} - 1)^4}$
norman	$\frac{(4iA a^3 + 4B a^3)x(\tan^4(dx+c)) - \frac{A a^3}{4d} + \frac{(-3iB a^3 + 4A a^3)(\tan^2(dx+c))}{2d} - \frac{(3iA a^3 + B a^3)\tan(dx+c)}{3d} + \frac{4(iA a^3 + B a^3)(\tan^3(dx+c))}{d}}{\tan(dx+c)^4}$

input `int(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `4*a^3*((-1/2*A+1/2*I*B)*ln(sec(d*x+c)^2)+(A-I*B)*ln(tan(d*x+c))-1/16*A*cot(d*x+c)^4+cot(d*x+c)^3*(-1/4*I*A-1/12*B)+cot(d*x+c)^2*(-3/8*I*B+1/2*A)+cot(d*x+c)*(I*A+B)+(I*A+B)*x*d)/d`

3.24.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.45

$$\int \cot^5(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx = \frac{2(12(3A-2iB)a^3 e^{6i dx+6i c} - 3(23A-19iB)a^3 e^{4i dx+4i c} + 2(27A-23iB)a^3 e^{2i dx+2i c} - (15A-15iB)a^3 e^{i dx+i c})}{3(d e^{8i c} - 3)}$$

input `integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output
$$-2/3*(12*(3*A - 2*I*B)*a^3*e^(6*I*d*x + 6*I*c) - 3*(23*A - 19*I*B)*a^3*e^(4*I*d*x + 4*I*c) + 2*(27*A - 23*I*B)*a^3*e^(2*I*d*x + 2*I*c) - (15*A - 13*I*B)*a^3 - 6*((A - I*B)*a^3*e^(8*I*d*x + 8*I*c) - 4*(A - I*B)*a^3*e^(6*I*d*x + 6*I*c) + 6*(A - I*B)*a^3*e^(4*I*d*x + 4*I*c) - 4*(A - I*B)*a^3*e^(2*I*d*x + 2*I*c) + (A - I*B)*a^3)*log(e^(2*I*d*x + 2*I*c) - 1)/(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)$$

3.24.6 Sympy [A] (verification not implemented)

Time = 1.19 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.50

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{4a^3(A - iB) \log(e^{2idx} - e^{-2ic})}{d} + \frac{30Aa^3 - 26iBa^3 + (-108Aa^3e^{2ic} + 92iBa^3e^{2ic})e^{2idx} + (138Aa^3e^{4ic} - 114iBa^3e^{4ic})e^{4idx} + (-72Aa^3e^{6ic} + 54iBa^3e^{6ic})e^{6idx} + 3d}{3de^{8ic}e^{8idx} - 12de^{6ic}e^{6idx} + 18de^{4ic}e^{4idx} - 12de^{2ic}e^{2idx} + 3d}$$

input `integrate(cot(d*x+c)**5*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output
$$4*a**3*(A - I*B)*log(exp(2*I*d*x) - exp(-2*I*c))/d + (30*A*a**3 - 26*I*B*a**3 + (-108*A*a**3*exp(2*I*c) + 92*I*B*a**3*exp(2*I*c))*exp(2*I*d*x) + (138*A*a**3*exp(4*I*c) - 114*I*B*a**3*exp(4*I*c))*exp(4*I*d*x) + (-72*A*a**3*exp(6*I*c) + 54*I*B*a**3*exp(6*I*c))*exp(6*I*d*x))/(3*d*exp(8*I*c)*exp(8*I*d*x) - 12*d*exp(6*I*c)*exp(6*I*d*x) + 18*d*exp(4*I*c)*exp(4*I*d*x) - 12*d*exp(2*I*c)*exp(2*I*d*x) + 3*d)$$

3.24.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.85

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{48(dx + c)(-iA - B)a^3 + 24(A - iB)a^3 \log(\tan(dx + c)^2 + 1) - 48(A - iB)a^3 \log(\tan(dx + c))}{12d}$$

input `integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output
$$\frac{-1/12*(48*(d*x + c)*(-I*A - B)*a^3 + 24*(A - I*B)*a^3*\log(\tan(d*x + c)^2 + 1) - 48*(A - I*B)*a^3*\log(\tan(d*x + c)) - (48*(I*A + B)*a^3*\tan(d*x + c)^3 + 6*(4*A - 3*I*B)*a^3*\tan(d*x + c)^2 + 4*(-3*I*A - B)*a^3*\tan(d*x + c) - 3*A*a^3)/\tan(d*x + c)^4}{d}$$

3.24.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 322 vs. $2(137) = 274$.

Time = 0.83 (sec) , antiderivative size = 322, normalized size of antiderivative = 2.05

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx =$$

$$3 A a^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 - 24i A a^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 8 B a^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 108 A a^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2$$

input `integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output
$$\frac{-1/192*(3*A*a^3*\tan(1/2*d*x + 1/2*c)^4 - 24*I*A*a^3*\tan(1/2*d*x + 1/2*c)^3 - 8*B*a^3*\tan(1/2*d*x + 1/2*c)^3 - 108*A*a^3*\tan(1/2*d*x + 1/2*c)^2 + 72*I*B*a^3*\tan(1/2*d*x + 1/2*c)^2 + 456*I*A*a^3*\tan(1/2*d*x + 1/2*c) + 408*B*a^3*\tan(1/2*d*x + 1/2*c) + 1536*(A*a^3 - I*B*a^3)*\log(\tan(1/2*d*x + 1/2*c) + I) - 768*(A*a^3 - I*B*a^3)*\log(\tan(1/2*d*x + 1/2*c)) + (1600*A*a^3*\tan(1/2*d*x + 1/2*c)^4 - 1600*I*B*a^3*\tan(1/2*d*x + 1/2*c)^4 - 456*I*A*a^3*\tan(1/2*d*x + 1/2*c)^3 - 408*B*a^3*\tan(1/2*d*x + 1/2*c)^3 - 108*A*a^3*\tan(1/2*d*x + 1/2*c)^2 + 72*I*B*a^3*\tan(1/2*d*x + 1/2*c)^2 + 24*I*A*a^3*\tan(1/2*d*x + 1/2*c) + 8*B*a^3*\tan(1/2*d*x + 1/2*c) + 3*A*a^3)/\tan(1/2*d*x + 1/2*c)^4}{d}$$

3.24.9 Mupad [B] (verification not implemented)

Time = 7.97 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.73

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{\tan(c + dx)^2 \left(2 A a^3 - \frac{B a^3 3i}{2}\right) + \tan(c + dx)^3 (4 B a^3 + A a^3 4i) - \frac{A a^3}{4} - \tan(c + dx) \left(\frac{B a^3}{3} + A a^3 1i\right)}{d \tan(c + dx)^4} + \frac{8 a^3 \operatorname{atan}(2 \tan(c + dx) + 1i) (B + A 1i)}{d}$$

input `int(cot(c + d*x)^5*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3,x)`

output `(tan(c + d*x)^2*(2*A*a^3 - (B*a^3*3i)/2) + tan(c + d*x)^3*(A*a^3*4i + 4*B*a^3) - (A*a^3)/4 - tan(c + d*x)*(A*a^3*1i + (B*a^3)/3))/(d*tan(c + d*x)^4) + (8*a^3*atan(2*tan(c + d*x) + 1i)*(A*1i + B))/d`

3.25 $\int \cot^6(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.25.1	Optimal result	476
3.25.2	Mathematica [A] (verified)	477
3.25.3	Rubi [A] (verified)	477
3.25.4	Maple [A] (verified)	482
3.25.5	Fricas [A] (verification not implemented)	483
3.25.6	Sympy [A] (verification not implemented)	483
3.25.7	Maxima [A] (verification not implemented)	484
3.25.8	Giac [B] (verification not implemented)	484
3.25.9	Mupad [B] (verification not implemented)	485

3.25.1 Optimal result

Integrand size = 34, antiderivative size = 180

$$\int \cot^6(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= -4a^3(A-iB)x - \frac{4a^3(A-iB) \cot(c+dx)}{d}$$

$$+ \frac{2a^3(iA+B) \cot^2(c+dx)}{d} + \frac{a^3(47A-45iB) \cot^3(c+dx)}{60d}$$

$$+ \frac{4a^3(iA+B) \log(\sin(c+dx))}{d} - \frac{aA \cot^5(c+dx)(a+ia \tan(c+dx))^2}{5d}$$

$$- \frac{(7iA+5B) \cot^4(c+dx)(a^3+ia^3 \tan(c+dx))}{20d}$$

output

```
-4*a^3*(A-I*B)*x-4*a^3*(A-I*B)*cot(d*x+c)/d+2*a^3*(I*A+B)*cot(d*x+c)^2/d+1/60*a^3*(47*A-45*I*B)*cot(d*x+c)^3/d+4*a^3*(I*A+B)*ln(sin(d*x+c))/d-1/5*a*A*cot(d*x+c)^5*(a+I*a*tan(d*x+c))^2/d-1/20*(7*I*A+5*B)*cot(d*x+c)^4*(a^3+I*a^3*tan(d*x+c))/d
```

3.25.2 Mathematica [A] (verified)

Time = 1.21 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.15

$$\int \cot^6(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= a^3 \left(-\frac{4A \cot(c+dx)}{d} + \frac{4iB \cot(c+dx)}{d} + \frac{2iA \cot^2(c+dx)}{d} + \frac{2B \cot^2(c+dx)}{d} \right. \\ \left. + \frac{4A \cot^3(c+dx)}{3d} - \frac{iB \cot^3(c+dx)}{d} - \frac{3iA \cot^4(c+dx)}{4d} - \frac{B \cot^4(c+dx)}{4d} \right. \\ \left. - \frac{A \cot^5(c+dx)}{5d} + \frac{4iA \log(\tan(c+dx))}{d} + \frac{4B \log(\tan(c+dx))}{d} \right. \\ \left. - \frac{4iA \log(i + \tan(c+dx))}{d} - \frac{4B \log(i + \tan(c+dx))}{d} \right)$$

input `Integrate[Cot[c + d*x]^6*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `a^3*((-4*A*Cot[c + d*x])/d + ((4*I)*B*Cot[c + d*x])/d + ((2*I)*A*Cot[c + d*x]^2)/d + (2*B*Cot[c + d*x]^2)/d + (4*A*Cot[c + d*x]^3)/(3*d) - (I*B*Cot[c + d*x]^3)/d - (((3*I)/4)*A*Cot[c + d*x]^4)/d - (B*Cot[c + d*x]^4)/(4*d) - (A*Cot[c + d*x]^5)/(5*d) + ((4*I)*A*Log[Tan[c + d*x]])/d + (4*B*Log[Tan[c + d*x]])/d - ((4*I)*A*Log[I + Tan[c + d*x]])/d - (4*B*Log[I + Tan[c + d*x]])/d)`

3.25.3 Rubi [A] (verified)

Time = 1.29 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.08, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.529$, Rules used = {3042, 4076, 3042, 4076, 25, 3042, 4074, 27, 3042, 4012, 25, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^6(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan(c+dx)^6} dx$$

$$\begin{aligned}
& \downarrow 4076 \\
& \frac{1}{5} \int \cot^5(c+dx)(i \tan(c+dx)a+a)^2(a(7iA+5B)-a(3A-5iB)\tan(c+dx))dx - \\
& \quad \frac{aA \cot^5(c+dx)(a+ia \tan(c+dx))^2}{5d} \\
& \downarrow 3042 \\
& \frac{1}{5} \int \frac{(i \tan(c+dx)a+a)^2(a(7iA+5B)-a(3A-5iB)\tan(c+dx))}{\tan(c+dx)^5} dx - \\
& \quad \frac{aA \cot^5(c+dx)(a+ia \tan(c+dx))^2}{5d} \\
& \downarrow 4076 \\
& \frac{1}{5} \left(\frac{1}{4} \int -\cot^4(c+dx)(i \tan(c+dx)a+a) ((47A-45iB)a^2+(33iA+35B)\tan(c+dx)a^2) dx - \frac{(5B+7iA)\cot^4(c+dx)(a+ia \tan(c+dx))^2}{4d} \right. \\
& \quad \left. \frac{aA \cot^5(c+dx)(a+ia \tan(c+dx))^2}{5d} \right) \\
& \downarrow 25 \\
& \frac{1}{5} \left(-\frac{1}{4} \int \cot^4(c+dx)(i \tan(c+dx)a+a) ((47A-45iB)a^2+(33iA+35B)\tan(c+dx)a^2) dx - \frac{(5B+7iA)\cot^4(c+dx)(a+ia \tan(c+dx))^2}{4d} \right. \\
& \quad \left. \frac{aA \cot^5(c+dx)(a+ia \tan(c+dx))^2}{5d} \right) \\
& \downarrow 3042 \\
& \frac{1}{5} \left(-\frac{1}{4} \int \frac{(i \tan(c+dx)a+a) ((47A-45iB)a^2+(33iA+35B)\tan(c+dx)a^2)}{\tan(c+dx)^4} dx - \frac{(5B+7iA)\cot^4(c+dx)(a+ia \tan(c+dx))^2}{4d} \right. \\
& \quad \left. \frac{aA \cot^5(c+dx)(a+ia \tan(c+dx))^2}{5d} \right) \\
& \downarrow 4074 \\
& \frac{1}{5} \left(\frac{1}{4} \left(\frac{a^3(47A-45iB)\cot^3(c+dx)}{3d} - \int 80 \cot^3(c+dx) (a^3(iA+B)-a^3(A-iB)\tan(c+dx)) dx \right) - \frac{(5B+7iA)\cot^4(c+dx)(a+ia \tan(c+dx))^2}{4d} \right. \\
& \quad \left. \frac{aA \cot^5(c+dx)(a+ia \tan(c+dx))^2}{5d} \right) \\
& \downarrow 27
\end{aligned}$$

3.25. $\int \cot^6(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{a^3(47A - 45iB) \cot^3(c + dx)}{3d} - 80 \int \cot^3(c + dx) (a^3(iA + B) - a^3(A - iB) \tan(c + dx)) dx \right) - \frac{(5B + 7iA) \cot^4(c + dx)}{4d} \right) - \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^2}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{a^3(47A - 45iB) \cot^3(c + dx)}{3d} - 80 \int \frac{a^3(iA + B) - a^3(A - iB) \tan(c + dx)}{\tan(c + dx)^3} dx \right) - \frac{(5B + 7iA) \cot^4(c + dx)}{4d} \right) - \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^2}{5d}$$

↓ 4012

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{a^3(47A - 45iB) \cot^3(c + dx)}{3d} - 80 \left(\int -\cot^2(c + dx) ((A - iB)a^3 + (iA + B) \tan(c + dx)a^3) dx - \frac{a^3(B - iA) \cot^3(c + dx)}{3d} \right) \right) - \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^2}{5d}$$

↓ 25

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{a^3(47A - 45iB) \cot^3(c + dx)}{3d} - 80 \left(- \int \cot^2(c + dx) ((A - iB)a^3 + (iA + B) \tan(c + dx)a^3) dx - \frac{a^3(B - iA) \cot^3(c + dx)}{3d} \right) \right) - \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^2}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{a^3(47A - 45iB) \cot^3(c + dx)}{3d} - 80 \left(- \int \frac{(A - iB)a^3 + (iA + B) \tan(c + dx)a^3}{\tan(c + dx)^2} dx - \frac{a^3(B + iA) \cot^2(c + dx)}{2d} \right) \right) - \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^2}{5d}$$

↓ 4012

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{a^3(47A - 45iB) \cot^3(c + dx)}{3d} - 80 \left(- \int \cot(c + dx) (a^3(iA + B) - a^3(A - iB) \tan(c + dx)) dx - \frac{a^3(B + iA) \cot^3(c + dx)}{3d} \right) \right) - \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^2}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{a^3(47A - 45iB) \cot^3(c + dx)}{3d} - 80 \left(- \int \frac{a^3(iA + B) - a^3(A - iB) \tan(c + dx)}{\tan(c + dx)} dx - \frac{a^3(B + iA) \cot^2(c + dx)}{2d} \right) \right) \right. \\ \left. \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^2}{5d} \right) \\ \downarrow 4014$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{a^3(47A - 45iB) \cot^3(c + dx)}{3d} - 80 \left(-a^3(B + iA) \int \cot(c + dx) dx - \frac{a^3(B + iA) \cot^2(c + dx)}{2d} + \frac{a^3(A - iB) \cot(c + dx)}{d} \right) \right) \right. \\ \left. \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^2}{5d} \right) \\ \downarrow 3042$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{a^3(47A - 45iB) \cot^3(c + dx)}{3d} - 80 \left(-a^3(B + iA) \int -\tan \left(c + dx + \frac{\pi}{2} \right) dx - \frac{a^3(B + iA) \cot^2(c + dx)}{2d} + \frac{a^3(A - iB) \cot(c + dx)}{d} \right) \right) \right. \\ \left. \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^2}{5d} \right) \\ \downarrow 25$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{a^3(47A - 45iB) \cot^3(c + dx)}{3d} - 80 \left(a^3(B + iA) \int \tan \left(\frac{1}{2}(2c + \pi) + dx \right) dx - \frac{a^3(B + iA) \cot^2(c + dx)}{2d} + \frac{a^3(A - iB) \cot(c + dx)}{d} \right) \right) \right. \\ \left. \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^2}{5d} \right) \\ \downarrow 3956$$

$$\frac{1}{5} \left(\frac{1}{4} \left(\frac{a^3(47A - 45iB) \cot^3(c + dx)}{3d} - 80 \left(-\frac{a^3(B + iA) \cot^2(c + dx)}{2d} + \frac{a^3(A - iB) \cot(c + dx)}{d} - \frac{a^3(B + iA) \log|\cot(c + dx)|}{d} \right) \right) \right. \\ \left. \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^2}{5d} \right)$$

input `Int[Cot[c + d*x]^6*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `-1/5*(a*A*Cot[c + d*x]^5*(a + I*a*Tan[c + d*x])^2)/d + (((a^3*(47*A - (45*I)*B)*Cot[c + d*x]^3)/(3*d) - 80*(a^3*(A - I*B)*x + (a^3*(A - I*B)*Cot[c + d*x])/d - (a^3*(I*A + B)*Cot[c + d*x]^2)/(2*d) - (a^3*(I*A + B)*Log[-Sin[c + d*x]])/d))/4 - (((7*I)*A + 5*B)*Cot[c + d*x]^4*(a^3 + I*a^3*Tan[c + d*x]))/(4*d))/5`

3.25.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`
- rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`
- rule 4074 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`


```
rule 4076 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

3.25.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.72

method	result
parallelrisch	$4 \left(\left(-\frac{iA}{2} - \frac{B}{2} \right) \ln(\sec^2(dx+c)) + (iA+B) \ln(\tan(dx+c)) - \frac{A(\cot^5(dx+c))}{20} + (\cot^4(dx+c)) \left(-\frac{3iA}{16} - \frac{B}{16} \right) + (\cot^3(dx+c)) \left(-\frac{3iA}{16} - \frac{B}{16} \right) \right) \frac{d}{d}$
derivativedivides	$a^3 \left(-\frac{3iA(\cot^4(dx+c))}{4} - \frac{A(\cot^5(dx+c))}{5} - iB(\cot^3(dx+c)) - \frac{B(\cot^4(dx+c))}{4} + 2iA(\cot^2(dx+c)) + \frac{4A(\cot^3(dx+c))}{3} + 4iB \cot(dx+c) \right) \frac{d}{d}$
default	$a^3 \left(-\frac{3iA(\cot^4(dx+c))}{4} - \frac{A(\cot^5(dx+c))}{5} - iB(\cot^3(dx+c)) - \frac{B(\cot^4(dx+c))}{4} + 2iA(\cot^2(dx+c)) + \frac{4A(\cot^3(dx+c))}{3} + 4iB \cot(dx+c) \right) \frac{d}{d}$
risch	$-\frac{8ia^3Bc}{d} + \frac{8a^3Ac}{d} - \frac{2a^3(240iAe^{8i(dx+c)} + 180Be^{8i(dx+c)} - 585iAe^{6i(dx+c)} - 525Be^{6i(dx+c)} + 695iAe^{4i(dx+c)} + 615Be^{4i(dx+c)} - 15d(e^{2i(dx+c)} - 1)^5)}{15d(e^{2i(dx+c)} - 1)^5}$
norman	$\frac{(4iBa^3 - 4Aa^3)x(\tan^5(dx+c)) - \frac{Aa^3}{5d} + \frac{(-3iBa^3 + 4Aa^3)(\tan^2(dx+c))}{3d} - \frac{(3iAa^3 + Ba^3)\tan(dx+c)}{4d} - \frac{4(-iBa^3 + Aa^3)(\tan^4(dx+c))}{d}}{\tan(dx+c)^5}$

```
input int(cot(d*x+c)^6*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

```
output 4*((-1/2*I*A-1/2*B)*ln(sec(d*x+c)^2)+(I*A+B)*ln(tan(d*x+c))-1/20*A*cot(d*x
+c)^5+cot(d*x+c)^4*(-3/16*I*A-1/16*B)+cot(d*x+c)^3*(-1/4*I*B+1/3*A)+cot(d*
x+c)^2*(1/2*I*A+1/2*B)+(-A+I*B)*cot(d*x+c)+x*d*(-A+I*B))*a^3/d
```

3.25. $\int \cot^6(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.25.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.59

$$\int \cot^6(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{2(60(4iA + 3B)a^3e^{(8i dx + 8i c)} + 15(-39iA - 35B)a^3e^{(6i dx + 6i c)} + 5(139iA + 123B)a^3e^{(4i dx + 4i c)} + 5(-77iA - 69B)a^3e^{(2i dx + 2i c)} + (83iA + 75B)a^3 + 30((-iA - B)a^3e^{(10i dx + 10i c)} + 5(iA + B)a^3e^{(8i dx + 8i c)} + 10(-iA - B)a^3e^{(6i dx + 6i c)} + 10(iA + B)a^3e^{(4i dx + 4i c)} + 5(-iA - B)a^3e^{(2i dx + 2i c)} + (iA + B)a^3)\log(e^{(2i dx + 2i c)} - 1))/(d e^{(10i dx + 10i c)} - 5d e^{(8i dx + 8i c)} + 10d e^{(6i dx + 6i c)} - 10d e^{(4i dx + 4i c)} + 5d e^{(2i dx + 2i c)} - d)}$$

input `integrate(cot(d*x+c)^6*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-2/15*(60*(4*I*A + 3*B))*a^3*e^(8*I*d*x + 8*I*c) + 15*(-39*I*A - 35*B))*a^3*e^(6*I*d*x + 6*I*c) + 5*(139*I*A + 123*B))*a^3*e^(4*I*d*x + 4*I*c) + 5*(-77*I*A - 69*B))*a^3*e^(2*I*d*x + 2*I*c) + (83*I*A + 75*B))*a^3 + 30*((-I*A - B))*a^3*e^(10*I*d*x + 10*I*c) + 5*(I*A + B))*a^3*e^(8*I*d*x + 8*I*c) + 10*(-I*A - B))*a^3*e^(6*I*d*x + 6*I*c) + 10*(I*A + B))*a^3*e^(4*I*d*x + 4*I*c) + 5*(-I*A - B))*a^3*e^(2*I*d*x + 2*I*c) + (I*A + B))*a^3*log(e^(2*I*d*x + 2*I*c) - 1)/(d*e^(10*I*d*x + 10*I*c) - 5*d*e^(8*I*d*x + 8*I*c) + 10*d*e^(6*I*d*x + 6*I*c) - 10*d*e^(4*I*d*x + 4*I*c) + 5*d*e^(2*I*d*x + 2*I*c) - d)`

3.25.6 Sympy [A] (verification not implemented)

Time = 0.73 (sec) , antiderivative size = 296, normalized size of antiderivative = 1.64

$$\int \cot^6(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{4ia^3(A - iB) \log(e^{2idx} - e^{-2ic})}{d} + \frac{-166iAa^3 - 150Ba^3 + (770iAa^3e^{2ic} + 690Ba^3e^{2ic})e^{2idx} + (-1390iAa^3e^{4ic} - 1230Ba^3e^{4ic})e^{4idx} + (1170iAa^3e^{6ic} + 1050Ba^3e^{6ic})e^{6idx} + (-480iAa^3e^{8ic} - 360Ba^3e^{8ic})e^{8idx}}{15de^{10ic}e^{10idx} - 75de^{8ic}e^{8idx} + 150de^{6ic}e^{6idx} - 150de^{4ic}e^{4idx} + 15d}$$

input `integrate(cot(d*x+c)**6*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `4*I*a**3*(A - I*B)*log(exp(2*I*d*x) - exp(-2*I*c))/d + (-166*I*A*a**3 - 150*B*a**3 + (770*I*A*a**3*exp(2*I*c) + 690*B*a**3*exp(2*I*c))*exp(2*I*d*x) + (-1390*I*A*a**3*exp(4*I*c) - 1230*B*a**3*exp(4*I*c))*exp(4*I*d*x) + (1170*I*A*a**3*exp(6*I*c) + 1050*B*a**3*exp(6*I*c))*exp(6*I*d*x) + (-480*I*A*a**3*exp(8*I*c) - 360*B*a**3*exp(8*I*c))*exp(8*I*d*x))/(15*d*exp(10*I*c)*exp(10*I*d*x) - 75*d*exp(8*I*c)*exp(8*I*d*x) + 150*d*exp(6*I*c)*exp(6*I*d*x) - 150*d*exp(4*I*c)*exp(4*I*d*x) + 15*d)`

3.25. $\int \cot^6(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.25.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.84

$$\int \cot^6(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{240(dx + c)(A - iB)a^3 + 120(iA + B)a^3 \log(\tan(dx + c)^2 + 1) + 240(-iA - B)a^3 \log(\tan(dx + c))}{60d}$$

input `integrate(cot(d*x+c)^6*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/60*(240*(d*x + c)*(A - I*B)*a^3 + 120*(I*A + B)*a^3*log(tan(d*x + c)^2 + 1) + 240*(-I*A - B)*a^3*log(tan(d*x + c)) + (240*(A - I*B)*a^3*tan(d*x + c)^4 - 120*(I*A + B)*a^3*tan(d*x + c)^3 - 20*(4*A - 3*I*B)*a^3*tan(d*x + c)^2 - 15*(-3*I*A - B)*a^3*tan(d*x + c) + 12*A*a^3)/tan(d*x + c)^5/d`

3.25.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 391 vs. 2(158) = 316.

Time = 0.91 (sec) , antiderivative size = 391, normalized size of antiderivative = 2.17

$$\int \cot^6(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{6Aa^3 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^5 - 45iAa^3 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^4 - 15Ba^3 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^4 - 190Aa^3 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)}{60d}$$

input `integrate(cot(d*x+c)^6*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output $\frac{1}{960}(6A^3 \tan^5(\frac{1}{2}dx + \frac{1}{2}c) - 45IA^3 \tan^4(\frac{1}{2}dx + \frac{1}{2}c) - 15B^3 \tan^4(\frac{1}{2}dx + \frac{1}{2}c) - 190A^3 \tan^3(\frac{1}{2}dx + \frac{1}{2}c) + 120IB^3 \tan^3(\frac{1}{2}dx + \frac{1}{2}c) + 660IA^3 \tan^2(\frac{1}{2}dx + \frac{1}{2}c) + 540B^3 \tan^2(\frac{1}{2}dx + \frac{1}{2}c) + 2460A^3 \tan(\frac{1}{2}dx + \frac{1}{2}c) - 2280IB^3 \tan(\frac{1}{2}dx + \frac{1}{2}c) - 7680(IA^3 + B^3) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) + I) - 3840(-IA^3 - B^3) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c)) + (-8768IA^3 \tan^5(\frac{1}{2}dx + \frac{1}{2}c) - 8768B^3 \tan^5(\frac{1}{2}dx + \frac{1}{2}c) - 2460A^3 \tan^4(\frac{1}{2}dx + \frac{1}{2}c) + 2280IB^3 \tan^4(\frac{1}{2}dx + \frac{1}{2}c) + 660IA^3 \tan^3(\frac{1}{2}dx + \frac{1}{2}c) + 540B^3 \tan^3(\frac{1}{2}dx + \frac{1}{2}c) + 190A^3 \tan^2(\frac{1}{2}dx + \frac{1}{2}c) - 120IB^3 \tan^2(\frac{1}{2}dx + \frac{1}{2}c) - 45IA^3 \tan(\frac{1}{2}dx + \frac{1}{2}c) - 15B^3 \tan(\frac{1}{2}dx + \frac{1}{2}c) - 6A^3) / \tan(\frac{1}{2}dx + \frac{1}{2}c)^5) / d$

3.25.9 Mupad [B] (verification not implemented)

Time = 8.24 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.78

$$\int \cot^6(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx =$$

$$\frac{\frac{Aa^3}{5} - \tan(c + dx)^2 \left(\frac{4Aa^3}{3} - Ba^3 1i \right) + \tan(c + dx)^4 (4Aa^3 - Ba^3 4i) - \tan(c + dx)^3 (2Ba^3 + Aa^3)}{d \tan(c + dx)^5} + \frac{a^3 \operatorname{atan}(2 \tan(c + dx) + 1i) (B + A 1i) 8i}{d}$$

input `int(cot(c + d*x)^6*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3,x)`

output $(a^3 \operatorname{atan}(2 \tan(c + d*x) + 1i) (A 1i + B) 8i) / d - (\tan(c + d*x)^4 (4A^3 - B^3 4i) - \tan(c + d*x)^2 ((4A^3) / 3 - B^3 1i) - \tan(c + d*x)^3 (A^3 2i + 2B^3)) + (A^3) / 5 + \tan(c + d*x) ((A^3 3i) / 4 + (B^3) / 4) / (d \tan(c + d*x)^5)$

3.26 $\int \tan^2(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.26.1	Optimal result	486
3.26.2	Mathematica [A] (verified)	487
3.26.3	Rubi [A] (verified)	487
3.26.4	Maple [A] (verified)	491
3.26.5	Fricas [A] (verification not implemented)	492
3.26.6	Sympy [A] (verification not implemented)	492
3.26.7	Maxima [A] (verification not implemented)	493
3.26.8	Giac [B] (verification not implemented)	493
3.26.9	Mupad [B] (verification not implemented)	494

3.26.1 Optimal result

Integrand size = 34, antiderivative size = 225

$$\begin{aligned} & \int \tan^2(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx \\ &= -8a^4(A-iB)x + \frac{8a^4(iA+B) \log(\cos(c+dx))}{d} \\ & \quad + \frac{8a^4(A-iB) \tan(c+dx)}{d} + \frac{4a^4(iA+B) \tan^2(c+dx)}{d} \\ & \quad - \frac{a^4(92A-93iB) \tan^3(c+dx)}{60d} + \frac{iaB \tan^3(c+dx)(a+ia \tan(c+dx))^3}{6d} \\ & \quad - \frac{(2A-3iB) \tan^3(c+dx) (a^2+ia^2 \tan(c+dx))^2}{10d} \\ & \quad - \frac{(12A-13iB) \tan^3(c+dx) (a^4+ia^4 \tan(c+dx))}{20d} \end{aligned}$$

output

```
-8*a^4*(A-I*B)*x+8*a^4*(I*A+B)*ln(cos(d*x+c))/d+8*a^4*(A-I*B)*tan(d*x+c)/d
+4*a^4*(I*A+B)*tan(d*x+c)^2/d-1/60*a^4*(92*A-93*I*B)*tan(d*x+c)^3/d+1/6*I*
a*B*tan(d*x+c)^3*(a+I*a*tan(d*x+c))^3/d-1/10*(2*A-3*I*B)*tan(d*x+c)^3*(a^2
+I*a^2*tan(d*x+c))^2/d-1/20*(12*A-13*I*B)*tan(d*x+c)^3*(a^4+I*a^4*tan(d*x+
c))/d
```

3.26.2 Mathematica [A] (verified)

Time = 1.83 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.61

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{a^4(-92iA - 93B - 480i(A - iB) \log(i + \tan(c + dx)) + 480(A - iB) \tan(c + dx) + 240(iA + B) \tan^2(c + dx) + 20(7A - (8i)B) \tan^3(c + dx) + ((-60i)A - 105B) \tan^4(c + dx) + 12(A - (4i)B) \tan^5(c + dx) + 10B \tan^6(c + dx))}{60d}$$

input `Integrate[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(a^4*((-92*I)*A - 93*B - (480*I)*(A - I*B)*Log[I + Tan[c + d*x]] + 480*(A - I*B)*Tan[c + d*x] + 240*(I*A + B)*Tan[c + d*x]^2 - 20*(7*A - (8*I)*B)*Tan[c + d*x]^3 + ((-60*I)*A - 105*B)*Tan[c + d*x]^4 + 12*(A - (4*I)*B)*Tan[c + d*x]^5 + 10*B*Tan[c + d*x]^6))/(60*d)`

3.26.3 Rubi [A] (verified)

Time = 1.42 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.07, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.471$, Rules used = {3042, 4077, 27, 3042, 4077, 27, 3042, 4077, 3042, 4075, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^2(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4077}$$

$$\frac{1}{6} \int 3 \tan^2(c + dx)(i \tan(c + dx)a + a)^3(a(2A - iB) + a(2iA + 3B) \tan(c + dx)) dx + \frac{iaB \tan^3(c + dx)(a + ia \tan(c + dx))^3}{6d}$$

$$\downarrow \text{27}$$

$$\frac{1}{2} \int \tan^2(c+dx)(i \tan(c+dx)a+a)^3(a(2A-iB)+a(2iA+3B)\tan(c+dx))dx + \frac{iaB \tan^3(c+dx)(a+ia \tan(c+dx))^3}{6d}$$

↓ 3042

$$\frac{1}{2} \int \tan(c+dx)^2(i \tan(c+dx)a+a)^3(a(2A-iB)+a(2iA+3B)\tan(c+dx))dx + \frac{iaB \tan^3(c+dx)(a+ia \tan(c+dx))^3}{6d}$$

↓ 4077

$$\frac{1}{2} \left(\frac{1}{5} \int 2 \tan^2(c+dx)(i \tan(c+dx)a+a)^2((8A-7iB)a^2+(12iA+13B)\tan(c+dx)a^2)dx - \frac{(2A-3iB)\tan^3(c+dx)(a+ia \tan(c+dx))^3}{6d} \right)$$

↓ 27

$$\frac{1}{2} \left(\frac{2}{5} \int \tan^2(c+dx)(i \tan(c+dx)a+a)^2((8A-7iB)a^2+(12iA+13B)\tan(c+dx)a^2)dx - \frac{(2A-3iB)\tan^3(c+dx)(a+ia \tan(c+dx))^3}{6d} \right)$$

↓ 3042

$$\frac{1}{2} \left(\frac{2}{5} \int \tan(c+dx)^2(i \tan(c+dx)a+a)^2((8A-7iB)a^2+(12iA+13B)\tan(c+dx)a^2)dx - \frac{(2A-3iB)\tan^3(c+dx)(a+ia \tan(c+dx))^3}{6d} \right)$$

↓ 4077

$$\frac{1}{2} \left(\frac{2}{5} \left(\frac{1}{4} \int \tan^2(c+dx)(i \tan(c+dx)a+a)((68A-67iB)a^3+(92iA+93B)\tan(c+dx)a^3)dx - \frac{(12A-13iB)\tan^3(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right)$$

↓ 3042

$$\frac{1}{2} \left(\frac{2}{5} \left(\frac{1}{4} \int \tan(c+dx)^2(i \tan(c+dx)a+a)((68A-67iB)a^3+(92iA+93B)\tan(c+dx)a^3)dx - \frac{(12A-13iB)\tan^3(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right)$$

↓ 4075

$$\frac{1}{2} \left(\frac{2}{5} \left(\frac{1}{4} \left(\int \tan^2(c+dx) (160(A-iB)a^4 + 160(iA+B)\tan(c+dx)a^4) dx - \frac{a^4(92A-93iB)\tan^3(c+dx)}{3d} \right) - \frac{iaB \tan^3(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right)$$

↓ 3042

$$\frac{1}{2} \left(\frac{2}{5} \left(\frac{1}{4} \left(\int \tan(c+dx)^2 (160(A-iB)a^4 + 160(iA+B)\tan(c+dx)a^4) dx - \frac{a^4(92A-93iB)\tan^3(c+dx)}{3d} \right) - \frac{iaB \tan^3(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right)$$

↓ 4011

$$\frac{1}{2} \left(\frac{2}{5} \left(\frac{1}{4} \left(\int \tan(c+dx) (160a^4(A-iB)\tan(c+dx) - 160a^4(iA+B)) dx - \frac{a^4(92A-93iB)\tan^3(c+dx)}{3d} + \frac{80a^4(B+iA)\tan^2(c+dx)}{d} \right) - \frac{iaB \tan^3(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right)$$

↓ 3042

$$\frac{1}{2} \left(\frac{2}{5} \left(\frac{1}{4} \left(\int \tan(c+dx) (160a^4(A-iB)\tan(c+dx) - 160a^4(iA+B)) dx - \frac{a^4(92A-93iB)\tan^3(c+dx)}{3d} + \frac{80a^4(B+iA)\tan^2(c+dx)}{d} + \frac{160a^4(B+iA)\tan(c+dx)}{d} \right) - \frac{iaB \tan^3(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right)$$

↓ 4008

$$\frac{1}{2} \left(\frac{2}{5} \left(\frac{1}{4} \left(-160a^4(B+iA) \int \tan(c+dx) dx - \frac{a^4(92A-93iB)\tan^3(c+dx)}{3d} + \frac{80a^4(B+iA)\tan^2(c+dx)}{d} + \frac{160a^4(B+iA)\tan(c+dx)}{d} \right) - \frac{iaB \tan^3(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right)$$

↓ 3042

$$\frac{1}{2} \left(\frac{2}{5} \left(\frac{1}{4} \left(-160a^4(B+iA) \int \tan(c+dx) dx - \frac{a^4(92A-93iB)\tan^3(c+dx)}{3d} + \frac{80a^4(B+iA)\tan^2(c+dx)}{d} + \frac{160a^4(B+iA)\tan(c+dx)}{d} + \frac{160a^4(B+iA)}{d} \right) - \frac{iaB \tan^3(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right)$$

↓ 3956

$$\frac{1}{2} \left(\frac{2}{5} \left(\frac{1}{4} \left(-\frac{a^4(92A - 93iB) \tan^3(c + dx)}{3d} + \frac{80a^4(B + iA) \tan^2(c + dx)}{d} + \frac{160a^4(A - iB) \tan(c + dx)}{d} + \frac{160a^4(I}{6d} \frac{iaB \tan^3(c + dx)(a + ia \tan(c + dx))^3}{6d} \right) \right) \right)$$

input `Int[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `((I/6)*a*B*Tan[c + d*x]^3*(a + I*a*Tan[c + d*x])^3)/d + (-1/5*((2*A - (3*I)*B)*Tan[c + d*x]^3*(a^2 + I*a^2*Tan[c + d*x])^2)/d + (2*(-1/4*((12*A - (13*I)*B)*Tan[c + d*x]^3*(a^4 + I*a^4*Tan[c + d*x])))/d + (-160*a^4*(A - I*B)*x + (160*a^4*(I*A + B)*Log[Cos[c + d*x]])/d + (160*a^4*(A - I*B)*Tan[c + d*x])/d + (80*a^4*(I*A + B)*Tan[c + d*x]^2)/d - (a^4*(92*A - (93*I)*B)*Tan[c + d*x]^3)/(3*d))/4)/5)/2`

3.26.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4077 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.26.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.75

method	result
derivativedivides	$a^4 \left(-\frac{4iB(\tan^5(dx+c))}{5} + \frac{B(\tan^6(dx+c))}{6} - iA(\tan^4(dx+c)) + \frac{A(\tan^5(dx+c))}{5} + \frac{8iB(\tan^3(dx+c))}{3} - \frac{7B(\tan^4(dx+c))}{4} + 4iA(\tan^2(dx+c)) \right)$
default	$a^4 \left(-\frac{4iB(\tan^5(dx+c))}{5} + \frac{B(\tan^6(dx+c))}{6} - iA(\tan^4(dx+c)) + \frac{A(\tan^5(dx+c))}{5} + \frac{8iB(\tan^3(dx+c))}{3} - \frac{7B(\tan^4(dx+c))}{4} + 4iA(\tan^2(dx+c)) \right)$
norman	$(8iB a^4 - 8A a^4) x - \frac{(4iA a^4 + 7B a^4)(\tan^4(dx+c))}{4d} - \frac{(-8iB a^4 + 7A a^4)(\tan^3(dx+c))}{3d} + \frac{8(-iB a^4 + A a^4)}{d}$
parallelrisc	$\frac{240iA \ln(1+\tan^2(dx+c))a^4 - 10B a^4(\tan^6(dx+c)) - 160iB(\tan^3(dx+c))a^4 - 12A(\tan^5(dx+c))a^4 - 240iA(\tan^2(dx+c))a^4}{d}$
risc	$-\frac{16ia^4Bc}{d} + \frac{16a^4Ac}{d} + \frac{4a^4(210iA e^{10i(dx+c)} + 270B e^{10i(dx+c)} + 765iA e^{8i(dx+c)} + 855B e^{8i(dx+c)} + 1210iA e^{6i(dx+c)} + 150B e^{6i(dx+c)} + 1210iA e^{4i(dx+c)} + 150B e^{4i(dx+c)} + 1210iA e^{2i(dx+c)} + 150B e^{2i(dx+c)} + 1210iA e^{0i(dx+c)} + 150B e^{0i(dx+c)})}{d}$
parts	$\frac{(-4iA a^4 - 6B a^4) \left(\frac{\tan^4(dx+c)}{4} - \frac{\tan^2(dx+c)}{2} + \frac{\ln(1+\tan^2(dx+c))}{2} \right)}{d} + \frac{(-4iB a^4 + A a^4) \left(\frac{\tan^5(dx+c)}{5} - \frac{\tan^3(dx+c)}{3} + \frac{\tan(dx+c)}{d} \right)}{d}$

input `int(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)), x, method=_RETURNVERBOSE)`

$$3.26. \quad \int \tan^2(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

output $1/d*a^4*(-4/5*I*B*\tan(d*x+c)^5+1/6*B*\tan(d*x+c)^6-I*A*\tan(d*x+c)^4+1/5*A*\tan(d*x+c)^5+8/3*I*B*\tan(d*x+c)^3-7/4*B*\tan(d*x+c)^4+4*I*A*\tan(d*x+c)^2-7/3*A*\tan(d*x+c)^3-8*I*B*\tan(d*x+c)+4*B*\tan(d*x+c)^2+8*A*\tan(d*x+c)+1/2*(-8*B-8*I*A)*\ln(1+\tan(d*x+c)^2)+(-8*A+8*I*B)*\arctan(\tan(d*x+c))$

3.26.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 345, normalized size of antiderivative = 1.53

$$\int \tan^2(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx = \frac{4(30(-7iA-9B)a^4e^{(10i dx+10i c)}+45(-17iA-19B)a^4e^{(8i dx+8i c)}+10(-121iA-135B)a^4e^{(6i dx+6i c)})}{d}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output $-4/15*(30*(-7*I*A-9*B)*a^4*e^{(10*I*d*x+10*I*c)}+45*(-17*I*A-19*B)*a^4*e^{(8*I*d*x+8*I*c)}+10*(-121*I*A-135*B)*a^4*e^{(6*I*d*x+6*I*c)}+15*(-68*I*A-75*B)*a^4*e^{(4*I*d*x+4*I*c)}+6*(-74*I*A-81*B)*a^4*e^{(2*I*d*x+2*I*c)}+(-79*I*A-86*B)*a^4+30*((-I*A-B)*a^4*e^{(12*I*d*x+12*I*c)}+6*(-I*A-B)*a^4*e^{(10*I*d*x+10*I*c)}+15*(-I*A-B)*a^4*e^{(8*I*d*x+8*I*c)}+20*(-I*A-B)*a^4*e^{(6*I*d*x+6*I*c)}+15*(-I*A-B)*a^4*e^{(4*I*d*x+4*I*c)}+6*(-I*A-B)*a^4*e^{(2*I*d*x+2*I*c)}+(-I*A-B)*a^4)*\log(e^{(2*I*d*x+2*I*c)}+1)/(d*e^{(12*I*d*x+12*I*c)}+6*d*e^{(10*I*d*x+10*I*c)}+15*d*e^{(8*I*d*x+8*I*c)}+20*d*e^{(6*I*d*x+6*I*c)}+15*d*e^{(4*I*d*x+4*I*c)}+6*d*e^{(2*I*d*x+2*I*c)}+d)$

3.26.6 Sympy [A] (verification not implemented)

Time = 0.82 (sec) , antiderivative size = 348, normalized size of antiderivative = 1.55

$$\int \tan^2(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx = \frac{8ia^4(A-iB) \log(e^{2idx}+e^{-2ic})}{d} + \frac{316iAa^4+344Ba^4+(1776iAa^4e^{2ic}+1944Ba^4e^{2ic})e^{2idx}+(4080iAa^4e^{4ic}+4500Ba^4e^{4ic})e^{4idx}+(4840iAa^4e^{6ic}+5400Ba^4e^{6ic})e^{6idx}+(2592iAa^4e^{8ic}+2916Ba^4e^{8ic})e^{8idx}+(1296iAa^4e^{10ic}+1458Ba^4e^{10ic})e^{10idx}+(648iAa^4e^{12ic}+729Ba^4e^{12ic})e^{12idx}}{15de^{12ic}e^{12idx}+90de^{10ic}e^{10idx}+225de^{8ic}e^{8idx}+45de^{6ic}e^{6idx}+9de^{4ic}e^{4idx}+de^{2ic}e^{2idx}+d}$$

input `integrate(tan(d*x+c)**2*(a+I*a*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)`

3.26. $\int \tan^2(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

output $8*I*a**4*(A - I*B)*\log(\exp(2*I*d*x) + \exp(-2*I*c))/d + (316*I*A*a**4 + 344*B*a**4 + (1776*I*A*a**4*\exp(2*I*c) + 1944*B*a**4*\exp(2*I*c))*\exp(2*I*d*x) + (4080*I*A*a**4*\exp(4*I*c) + 4500*B*a**4*\exp(4*I*c))*\exp(4*I*d*x) + (4840*I*A*a**4*\exp(6*I*c) + 5400*B*a**4*\exp(6*I*c))*\exp(6*I*d*x) + (3060*I*A*a**4*\exp(8*I*c) + 3420*B*a**4*\exp(8*I*c))*\exp(8*I*d*x) + (840*I*A*a**4*\exp(10*I*c) + 1080*B*a**4*\exp(10*I*c))*\exp(10*I*d*x))/(15*d*\exp(12*I*c)*\exp(12*I*d*x) + 90*d*\exp(10*I*c)*\exp(10*I*d*x) + 225*d*\exp(8*I*c)*\exp(8*I*d*x) + 300*d*\exp(6*I*c)*\exp(6*I*d*x) + 225*d*\exp(4*I*c)*\exp(4*I*d*x) + 90*d*\exp(2*I*c)*\exp(2*I*d*x) + 15*d)$

3.26.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.67

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{10 B a^4 \tan(dx + c)^6 + 12(A - 4i B) a^4 \tan(dx + c)^5 - 15(4i A + 7 B) a^4 \tan(dx + c)^4 - 20(7 A - 8i B) a^4 \tan(dx + c)^3 - 240(-i A - B) a^4 \tan(dx + c)^2 + 480(A - i B) a^4 \tan(dx + c)}{d}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output $1/60*(10*B*a^4*\tan(d*x + c)^6 + 12*(A - 4*I*B)*a^4*\tan(d*x + c)^5 - 15*(4*I*A + 7*B)*a^4*\tan(d*x + c)^4 - 20*(7*A - 8*I*B)*a^4*\tan(d*x + c)^3 - 240*(-I*A - B)*a^4*\tan(d*x + c)^2 - 480*(d*x + c)*(A - I*B)*a^4 - 240*(I*A + B)*a^4*\log(\tan(d*x + c)^2 + 1) + 480*(A - I*B)*a^4*\tan(d*x + c))/d$

3.26.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 600 vs. $2(195) = 390$.

Time = 0.82 (sec) , antiderivative size = 600, normalized size of antiderivative = 2.67

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$\frac{4(-30i A a^4 e^{(12i dx + 12i c)} \log(e^{(2i dx + 2i c)} + 1) - 30 B a^4 e^{(12i dx + 12i c)} \log(e^{(2i dx + 2i c)} + 1) - 180i A a^4 e^{(10i dx + 10i c)} \log(e^{(2i dx + 2i c)} + 1) - 180i B a^4 e^{(10i dx + 10i c)} \log(e^{(2i dx + 2i c)} + 1) - 180i A a^4 e^{(8i dx + 8i c)} \log(e^{(2i dx + 2i c)} + 1) - 180i B a^4 e^{(8i dx + 8i c)} \log(e^{(2i dx + 2i c)} + 1) - 180i A a^4 e^{(6i dx + 6i c)} \log(e^{(2i dx + 2i c)} + 1) - 180i B a^4 e^{(6i dx + 6i c)} \log(e^{(2i dx + 2i c)} + 1) - 180i A a^4 e^{(4i dx + 4i c)} \log(e^{(2i dx + 2i c)} + 1) - 180i B a^4 e^{(4i dx + 4i c)} \log(e^{(2i dx + 2i c)} + 1) - 180i A a^4 e^{(2i dx + 2i c)} \log(e^{(2i dx + 2i c)} + 1) - 180i B a^4 \log(e^{(2i dx + 2i c)} + 1))}{d}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `-4/15*(-30*I*A*a^4*e^(12*I*d*x + 12*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 30*B*a^4*e^(12*I*d*x + 12*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 180*I*A*a^4*e^(10*I*d*x + 10*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 180*B*a^4*e^(10*I*d*x + 10*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 450*I*A*a^4*e^(8*I*d*x + 8*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 450*B*a^4*e^(8*I*d*x + 8*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 600*I*A*a^4*e^(6*I*d*x + 6*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 600*B*a^4*e^(6*I*d*x + 6*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 450*I*A*a^4*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 450*B*a^4*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 180*I*A*a^4*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 180*B*a^4*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 210*I*A*a^4*e^(10*I*d*x + 10*I*c) - 270*B*a^4*e^(10*I*d*x + 10*I*c) - 765*I*A*a^4*e^(8*I*d*x + 8*I*c) - 855*B*a^4*e^(8*I*d*x + 8*I*c) - 1210*I*A*a^4*e^(6*I*d*x + 6*I*c) - 1350*B*a^4*e^(6*I*d*x + 6*I*c) - 1020*I*A*a^4*e^(4*I*d*x + 4*I*c) - 1125*B*a^4*e^(4*I*d*x + 4*I*c) - 444*I*A*a^4*e^(2*I*d*x + 2*I*c) - 486*B*a^4*e^(2*I*d*x + 2*I*c) - 30*I*A*a^4*log(e^(2*I*d*x + 2*I*c) + 1) - 30*B*a^4*log(e^(2*I*d*x + 2*I*c) + 1) - 79*I*A*a^4 - 86*B*a^4)/(d*e^(12*I*d*x + 12*I*c) + 6*d*e^(10*I*d*x + 10*I*c) + 15*d*e^(8*I*d*x + 8*I*c) + 20*d*e^(6*I*d*x + 6*I*c) + 15*d*e^(4*I*d*x + 4*I*c) + 6*d*e^(2*I*d*x + 2*I*c) + d)`

3.26.9 Mupad [B] (verification not implemented)

Time = 7.54 (sec) , antiderivative size = 308, normalized size of antiderivative = 1.37

$$\int \tan^2(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= \frac{\tan(c+dx)^3 \left(-a^4(A-B \operatorname{li}) + \frac{a^4(B+A 3i) \operatorname{li}}{3} + \frac{B a^4 \operatorname{li}}{3} + \frac{a^4(3B+A 1i) \operatorname{li}}{3} \right)}{d} - \frac{\tan(c+dx) (-A a^4 - 3 a^4(A-B \operatorname{li}) + a^4(B+A 3i) \operatorname{li} + B a^4 \operatorname{li} + a^4(3B+A 1i) \operatorname{li})}{d} - \frac{\tan(c+dx)^5 \left(\frac{B a^4 \operatorname{li}}{5} + \frac{a^4(3B+A 1i) \operatorname{li}}{5} \right)}{d} - \frac{\ln(\tan(c+dx) + \operatorname{li}) (8 B a^4 + A a^4 8i)}{d} + \frac{\tan(c+dx)^2 \left(\frac{A a^4 \operatorname{li}}{2} + \frac{a^4(A-B 1i) 3i}{2} + \frac{a^4(B+A 3i)}{2} + \frac{B a^4}{2} + \frac{a^4(3B+A 1i)}{2} \right)}{d} - \frac{\tan(c+dx)^4 \left(\frac{a^4(A-B 1i) 3i}{4} + \frac{B a^4}{4} + \frac{a^4(3B+A 1i)}{4} \right)}{d} + \frac{B a^4 \tan(c+dx)^6}{6 d}$$

3.26. $\int \tan^2(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^4,x)`

output `(tan(c + d*x)^3*((a^4*(A*3i + B)*1i)/3 - a^4*(A - B*1i) + (B*a^4*1i)/3 + (a^4*(A*1i + 3*B)*1i)/3))/d - (tan(c + d*x)*(a^4*(A*3i + B)*1i - 3*a^4*(A - B*1i) - A*a^4 + B*a^4*1i + a^4*(A*1i + 3*B)*1i))/d - (tan(c + d*x)^5*((B*a^4*1i)/5 + (a^4*(A*1i + 3*B)*1i)/5))/d - (log(tan(c + d*x) + 1i)*(A*a^4*8i + 8*B*a^4))/d + (tan(c + d*x)^2*((A*a^4*1i)/2 + (a^4*(A - B*1i)*3i)/2 + (a^4*(A*3i + B))/2 + (B*a^4)/2 + (a^4*(A*1i + 3*B))/2))/d - (tan(c + d*x)^4*((a^4*(A - B*1i)*3i)/4 + (B*a^4)/4 + (a^4*(A*1i + 3*B))/4))/d + (B*a^4*tan(c + d*x)^6)/(6*d)`

3.27 $\int \tan(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.27.1	Optimal result	496
3.27.2	Mathematica [A] (verified)	497
3.27.3	Rubi [A] (verified)	497
3.27.4	Maple [A] (verified)	500
3.27.5	Fricas [A] (verification not implemented)	501
3.27.6	Sympy [B] (verification not implemented)	501
3.27.7	Maxima [A] (verification not implemented)	502
3.27.8	Giac [B] (verification not implemented)	502
3.27.9	Mupad [B] (verification not implemented)	503

3.27.1 Optimal result

Integrand size = 32, antiderivative size = 168

$$\int \tan(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= -8a^4(iA+B)x - \frac{8a^4(A-iB) \log(\cos(c+dx))}{d} + \frac{4a^4(iA+B) \tan(c+dx)}{d}$$

$$+ \frac{a(A-iB)(a+ia \tan(c+dx))^3}{3d} + \frac{A(a+ia \tan(c+dx))^4}{4d}$$

$$- \frac{iB(a+ia \tan(c+dx))^5}{5ad} + \frac{(A-iB)(a^2+ia^2 \tan(c+dx))^2}{d}$$

output

```
-8*a^4*(I*A+B)*x-8*a^4*(A-I*B)*ln(cos(d*x+c))/d+4*a^4*(I*A+B)*tan(d*x+c)/d
+1/3*a*(A-I*B)*(a+I*a*tan(d*x+c))^3/d+1/4*A*(a+I*a*tan(d*x+c))^4/d-1/5*I*B
*(a+I*a*tan(d*x+c))^5/a/d+(A-I*B)*(a^2+I*a^2*tan(d*x+c))^2/d
```

3.27.2 Mathematica [A] (verified)

Time = 1.16 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.71

$$\int \tan(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{a^4(95A - 92iB + 480(A - iB) \log(i + \tan(c + dx)) + 480(iA + B) \tan(c + dx) - 30(7A - 8iB) \tan^2(c + dx) + 15(A - 4iB) \tan^3(c + dx) + 12B \tan^4(c + dx))}{60d}$$

input `Integrate[Tan[c + d*x]*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(a^4*(95*A - (92*I)*B + 480*(A - I*B)*Log[I + Tan[c + d*x]] + 480*(I*A + B)*Tan[c + d*x] - 30*(7*A - (8*I)*B)*Tan[c + d*x]^2 + ((-80*I)*A - 140*B)*Tan[c + d*x]^3 + 15*(A - (4*I)*B)*Tan[c + d*x]^4 + 12*B*Tan[c + d*x]^5)/(60*d)`

3.27.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.93, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3042, 4075, 3042, 4010, 3042, 3959, 3042, 3959, 3042, 3958, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4075}$$

$$\int (i \tan(c + dx)a + a)^4(A \tan(c + dx) - B) dx - \frac{iB(a + ia \tan(c + dx))^5}{5ad}$$

$$\downarrow \text{3042}$$

$$\int (i \tan(c + dx)a + a)^4(A \tan(c + dx) - B) dx - \frac{iB(a + ia \tan(c + dx))^5}{5ad}$$

$$\downarrow \text{4010}$$

3.27. $\int \tan(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& -(B + iA) \int (i \tan(c + dx)a + a)^4 dx + \frac{A(a + ia \tan(c + dx))^4}{4d} - \frac{iB(a + ia \tan(c + dx))^5}{5ad} \\
& \quad \downarrow \text{3042} \\
& -(B + iA) \int (i \tan(c + dx)a + a)^4 dx + \frac{A(a + ia \tan(c + dx))^4}{4d} - \frac{iB(a + ia \tan(c + dx))^5}{5ad} \\
& \quad \downarrow \text{3959} \\
& -(B + iA) \left(2a \int (i \tan(c + dx)a + a)^3 dx + \frac{ia(a + ia \tan(c + dx))^3}{3d} \right) + \frac{A(a + ia \tan(c + dx))^4}{4d} - \\
& \quad \frac{iB(a + ia \tan(c + dx))^5}{5ad} \\
& \quad \downarrow \text{3042} \\
& -(B + iA) \left(2a \int (i \tan(c + dx)a + a)^3 dx + \frac{ia(a + ia \tan(c + dx))^3}{3d} \right) + \frac{A(a + ia \tan(c + dx))^4}{4d} - \\
& \quad \frac{iB(a + ia \tan(c + dx))^5}{5ad} \\
& \quad \downarrow \text{3959} \\
& iA \left(2a \left(2a \int (i \tan(c + dx)a + a)^2 dx + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \frac{ia(a + ia \tan(c + dx))^3}{3d} \right) + \\
& \quad \frac{A(a + ia \tan(c + dx))^4}{4d} - \frac{iB(a + ia \tan(c + dx))^5}{5ad} \\
& \quad \downarrow \text{3042} \\
& iA \left(2a \left(2a \int (i \tan(c + dx)a + a)^2 dx + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \frac{ia(a + ia \tan(c + dx))^3}{3d} \right) + \\
& \quad \frac{A(a + ia \tan(c + dx))^4}{4d} - \frac{iB(a + ia \tan(c + dx))^5}{5ad} \\
& \quad \downarrow \text{3958} \\
& iA \left(2a \left(2a \left(2ia^2 \int \tan(c + dx) dx - \frac{a^2 \tan(c + dx)}{d} + 2a^2 x \right) + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \frac{ia(a + ia \tan(c + dx))^3}{3d} \right) + \\
& \quad \frac{A(a + ia \tan(c + dx))^4}{4d} - \frac{iB(a + ia \tan(c + dx))^5}{5ad} \\
& \quad \downarrow \text{3042} \\
& iA \left(2a \left(2a \left(2ia^2 \int \tan(c + dx) dx - \frac{a^2 \tan(c + dx)}{d} + 2a^2 x \right) + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \frac{ia(a + ia \tan(c + dx))^3}{3d} \right) + \\
& \quad \frac{A(a + ia \tan(c + dx))^4}{4d} - \frac{iB(a + ia \tan(c + dx))^5}{5ad}
\end{aligned}$$

3.27. $\int \tan(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$

$$\begin{array}{c}
 \downarrow \text{3956} \\
 iA \left(2a \left(2a \left(-\frac{a^2 \tan(c+dx)}{d} - \frac{2ia^2 \log(\cos(c+dx))}{d} + 2a^2 x \right) + \frac{ia(a+ia \tan(c+dx))^2}{2d} \right) + \frac{ia(a+ia \tan(c+dx))}{3d} \right. \\
 \left. \frac{A(a+ia \tan(c+dx))^4}{4d} - \frac{iB(a+ia \tan(c+dx))^5}{5ad} \right)
 \end{array}$$

input `Int[Tan[c + d*x]*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(A*(a + I*a*Tan[c + d*x])^4)/(4*d) - ((I/5)*B*(a + I*a*Tan[c + d*x])^5)/(a*d) - (I*A + B)*(((I/3)*a*(a + I*a*Tan[c + d*x])^3)/d + 2*a*(((I/2)*a*(a + I*a*Tan[c + d*x])^2)/d + 2*a*(2*a^2*x - ((2*I)*a^2*Log[Cos[c + d*x]])/d - (a^2*Tan[c + d*x])/d))`

3.27.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 3958 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_.)])^2, x_Symbol] := Simp[(a^2 - b^2)*x, x] + (Simp[b^2*(Tan[c + d*x]/d), x] + Simp[2*a*b Int[Tan[c + d*x], x], x]) /; FreeQ[{a, b, c, d}, x]`

rule 3959 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

$$3.27. \quad \int \tan(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

3.27.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.86

method	result
derivativedivides	$\frac{a^4 \left(-iB(\tan^4(dx+c)) + \frac{B(\tan^5(dx+c))}{5} - \frac{4iA(\tan^3(dx+c))}{3} + \frac{A(\tan^4(dx+c))}{4} + 4iB(\tan^2(dx+c)) - \frac{7B(\tan^3(dx+c))}{3} + 8iA \right)}{d}$
default	$\frac{a^4 \left(-iB(\tan^4(dx+c)) + \frac{B(\tan^5(dx+c))}{5} - \frac{4iA(\tan^3(dx+c))}{3} + \frac{A(\tan^4(dx+c))}{4} + 4iB(\tan^2(dx+c)) - \frac{7B(\tan^3(dx+c))}{3} + 8iA \right)}{d}$
norman	$\frac{(-8iA a^4 - 8B a^4) x - \frac{(4iA a^4 + 7B a^4)(\tan^3(dx+c))}{3d} - \frac{(-8iB a^4 + 7A a^4)(\tan^2(dx+c))}{2d} + \frac{8(iA a^4 + B a^4)}{d}}{d}$
parallelrisch	$\frac{60iB(\tan^4(dx+c))a^4 - 12B a^4(\tan^5(dx+c)) + 80iA(\tan^3(dx+c))a^4 - 15A(\tan^4(dx+c))a^4 + 480iA x a^4 d - 240iB(\tan^2(dx+c))a^4}{15d(e^{2i(dx+c)} + 1)^5}$
risch	$\frac{16a^4 Bc}{d} + \frac{16ia^4 Ac}{d} + \frac{4ia^4(150iA e^{8i(dx+c)} + 210B e^{8i(dx+c)} + 465iA e^{6i(dx+c)} + 555B e^{6i(dx+c)} + 565iA e^{4i(dx+c)} + 615B e^{4i(dx+c)} + 615iA e^{2i(dx+c)} + 615B e^{2i(dx+c)} + 615iA)}{15d(e^{2i(dx+c)} + 1)^5}$
parts	$\frac{(-4iA a^4 - 6B a^4) \left(\frac{(\tan^3(dx+c))}{3} - \tan(dx+c) + \arctan(\tan(dx+c)) \right)}{d} + \frac{(-4iB a^4 + A a^4) \left(\frac{(\tan^4(dx+c))}{4} - \frac{(\tan^2(dx+c))}{2} \right)}{d}$

```
input int(tan(d*x+c)*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

```
output 1/d*a^4*(-I*B*tan(d*x+c)^4+1/5*B*tan(d*x+c)^5-4/3*I*A*tan(d*x+c)^3+1/4*A*t
an(d*x+c)^4+4*I*B*tan(d*x+c)^2-7/3*B*tan(d*x+c)^3+8*I*A*tan(d*x+c)-7/2*A*t
an(d*x+c)^2+8*B*tan(d*x+c)+1/2*(8*A-8*I*B)*ln(1+tan(d*x+c)^2)+(-8*B-8*I*A)
*arctan(tan(d*x+c)))
```

3.27. $\int \tan(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$

3.27.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.66

$$\int \tan(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx =$$

$$4(30(5A-7iB)a^4e^{(8i dx+8i c)} + 15(31A-37iB)a^4e^{(6i dx+6i c)} + 5(113A-131iB)a^4e^{(4i dx+4i c)} + 5(6$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output `-4/15*(30*(5*A - 7*I*B)*a^4*e^(8*I*d*x + 8*I*c) + 15*(31*A - 37*I*B)*a^4*e^(6*I*d*x + 6*I*c) + 5*(113*A - 131*I*B)*a^4*e^(4*I*d*x + 4*I*c) + 5*(64*A - 73*I*B)*a^4*e^(2*I*d*x + 2*I*c) + (70*A - 79*I*B)*a^4 + 30*((A - I*B)*a^4*e^(10*I*d*x + 10*I*c) + 5*(A - I*B)*a^4*e^(8*I*d*x + 8*I*c) + 10*(A - I*B)*a^4*e^(6*I*d*x + 6*I*c) + 10*(A - I*B)*a^4*e^(4*I*d*x + 4*I*c) + 5*(A - I*B)*a^4*e^(2*I*d*x + 2*I*c) + (A - I*B)*a^4)*log(e^(2*I*d*x + 2*I*c) + 1))/(d*e^(10*I*d*x + 10*I*c) + 5*d*e^(8*I*d*x + 8*I*c) + 10*d*e^(6*I*d*x + 6*I*c) + 10*d*e^(4*I*d*x + 4*I*c) + 5*d*e^(2*I*d*x + 2*I*c) + d)`

3.27.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 291 vs. 2(138) = 276.

Time = 0.50 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.73

$$\int \tan(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx = -\frac{8a^4(A-iB) \log(e^{2idx} + e^{-2ic})}{d}$$

$$+ \frac{-280Aa^4 + 316iBa^4 + (-1280Aa^4e^{2ic} + 1460iBa^4e^{2ic})e^{2idx} + (-2260Aa^4e^{4ic} + 2620iBa^4e^{4ic})e^{4idx} + 15de^{10ic}e^{10idx} + 75de^{8ic}e^{8idx} + 150de^{6ic}e^{6idx} + 150de^{4ic}e^{4idx}}{15de^{10ic}e^{10idx} + 75de^{8ic}e^{8idx} + 150de^{6ic}e^{6idx} + 150de^{4ic}e^{4idx}}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)`

output
$$\begin{aligned} & -8a^{**4}(A - I*B)*\log(\exp(2*I*d*x) + \exp(-2*I*c))/d + (-280*A*a^{**4} + 316*I \\ & *B*a^{**4} + (-1280*A*a^{**4}*\exp(2*I*c) + 1460*I*B*a^{**4}*\exp(2*I*c))*\exp(2*I*d*x \\ &) + (-2260*A*a^{**4}*\exp(4*I*c) + 2620*I*B*a^{**4}*\exp(4*I*c))*\exp(4*I*d*x) + (- \\ & 1860*A*a^{**4}*\exp(6*I*c) + 2220*I*B*a^{**4}*\exp(6*I*c))*\exp(6*I*d*x) + (-600*A* \\ & a^{**4}*\exp(8*I*c) + 840*I*B*a^{**4}*\exp(8*I*c))*\exp(8*I*d*x)/(15*d*\exp(10*I*c) \\ & * \exp(10*I*d*x) + 75*d*\exp(8*I*c)*\exp(8*I*d*x) + 150*d*\exp(6*I*c)*\exp(6*I*d \\ & *x) + 150*d*\exp(4*I*c)*\exp(4*I*d*x) + 75*d*\exp(2*I*c)*\exp(2*I*d*x) + 15*d \end{aligned}$$

3.27.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.79

$$\int \tan(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{12 B a^4 \tan(dx + c)^5 + 15 (A - 4i B) a^4 \tan(dx + c)^4 - 20 (4i A + 7 B) a^4 \tan(dx + c)^3 - 30 (7 A - 8i B) a^4 \tan(dx + c)^2 + 15 (A - 4i B) a^4 \tan(dx + c) + 15 a^4 \log(\tan(dx + c)^2 + 1)}{d}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output
$$\begin{aligned} & 1/60*(12*B*a^4*\tan(d*x + c)^5 + 15*(A - 4*I*B)*a^4*\tan(d*x + c)^4 - 20*(4* \\ & I*A + 7*B)*a^4*\tan(d*x + c)^3 - 30*(7*A - 8*I*B)*a^4*\tan(d*x + c)^2 - 480* \\ & (d*x + c)*(I*A + B)*a^4 + 240*(A - I*B)*a^4*\log(\tan(d*x + c)^2 + 1) - 480* \\ & (-I*A - B)*a^4*\tan(d*x + c))/d \end{aligned}$$

3.27.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 504 vs. $2(142) = 284$.

Time = 0.63 (sec) , antiderivative size = 504, normalized size of antiderivative = 3.00

$$\int \tan(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$\frac{4(30 A a^4 e^{(10i dx + 10i c)} \log(e^{(2i dx + 2i c)} + 1) - 30i B a^4 e^{(10i dx + 10i c)} \log(e^{(2i dx + 2i c)} + 1) + 150 A a^4 e^{(8i dx + 8i c)} \log(e^{(2i dx + 2i c)} + 1) - 150i B a^4 e^{(8i dx + 8i c)} \log(e^{(2i dx + 2i c)} + 1) + 150 A a^4 \log(e^{(2i dx + 2i c)} + 1) - 150i B a^4 \log(e^{(2i dx + 2i c)} + 1))}{d}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

3.27. $\int \tan(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$

output

```

-4/15*(30*A*a^4*e^(10*I*d*x + 10*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 30*I*
B*a^4*e^(10*I*d*x + 10*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 150*A*a^4*e^(8*
I*d*x + 8*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 150*I*B*a^4*e^(8*I*d*x + 8*I
*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 300*A*a^4*e^(6*I*d*x + 6*I*c)*log(e^(2*
I*d*x + 2*I*c) + 1) - 300*I*B*a^4*e^(6*I*d*x + 6*I*c)*log(e^(2*I*d*x + 2*I
*c) + 1) + 300*A*a^4*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 30
0*I*B*a^4*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 150*A*a^4*e^(
2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 150*I*B*a^4*e^(2*I*d*x + 2
*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 150*A*a^4*e^(8*I*d*x + 8*I*c) - 210*I
*B*a^4*e^(8*I*d*x + 8*I*c) + 465*A*a^4*e^(6*I*d*x + 6*I*c) - 555*I*B*a^4*e
^(6*I*d*x + 6*I*c) + 565*A*a^4*e^(4*I*d*x + 4*I*c) - 655*I*B*a^4*e^(4*I*d*
x + 4*I*c) + 320*A*a^4*e^(2*I*d*x + 2*I*c) - 365*I*B*a^4*e^(2*I*d*x + 2*I*
c) + 30*A*a^4*log(e^(2*I*d*x + 2*I*c) + 1) - 30*I*B*a^4*log(e^(2*I*d*x + 2
*I*c) + 1) + 70*A*a^4 - 79*I*B*a^4)/(d*e^(10*I*d*x + 10*I*c) + 5*d*e^(8*I*
d*x + 8*I*c) + 10*d*e^(6*I*d*x + 6*I*c) + 10*d*e^(4*I*d*x + 4*I*c) + 5*d*e
^(2*I*d*x + 2*I*c) + d)

```

3.27.9 Mupad [B] (verification not implemented)

Time = 7.50 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.43

$$\begin{aligned}
 & \int \tan(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx \\
 &= \frac{\tan(c + dx)^2 \left(-\frac{3a^4(A - B \operatorname{li})}{2} + \frac{a^4(B + A 3i) \operatorname{li}}{2} + \frac{B a^4 \operatorname{li}}{2} + \frac{a^4(3B + A \operatorname{li}) \operatorname{li}}{2} \right)}{d} \\
 &+ \frac{\tan(c + dx) (A a^4 \operatorname{li} + a^4(A - B \operatorname{li}) 3i + a^4(B + A 3i) + B a^4 + a^4(3B + A \operatorname{li}))}{d} \\
 &- \frac{\tan(c + dx)^4 \left(\frac{B a^4 \operatorname{li}}{4} + \frac{a^4(3B + A \operatorname{li}) \operatorname{li}}{4} \right)}{d} + \frac{\ln(\tan(c + dx) + \operatorname{li}) (8A a^4 - B a^4 8i)}{d} \\
 &- \frac{\tan(c + dx)^3 \left(a^4(A - B \operatorname{li}) \operatorname{li} + \frac{B a^4}{3} + \frac{a^4(3B + A \operatorname{li})}{3} \right)}{d} + \frac{B a^4 \tan(c + dx)^5}{5d}
 \end{aligned}$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^4,x)`

output $(\tan(c + dx)^2((a^4(A^3i + B)1i)/2 - (3a^4(A - B1i))/2 + (Ba^41i)/2 + (a^4(A1i + 3B)1i)/2))/d + (\tan(c + dx)(Aa^41i + a^4(A - B1i)3i + a^4(A3i + B) + Ba^4 + a^4(A1i + 3B)))/d - (\tan(c + dx)^4((Ba^41i)/4 + (a^4(A1i + 3B)1i)/4))/d + (\log(\tan(c + dx) + 1)(8Aa^4 - Ba^48i))/d - (\tan(c + dx)^3(a^4(A - B1i)1i + (Ba^4)/3 + (a^4(A1i + 3B))/3))/d + (Ba^4\tan(c + dx)^5)/(5d)$

3.28 $\int (a + ia \tan(c + dx))^4 (A + B \tan(c + dx)) dx$

3.28.1	Optimal result	505
3.28.2	Mathematica [A] (verified)	505
3.28.3	Rubi [A] (verified)	506
3.28.4	Maple [A] (verified)	508
3.28.5	Fricas [A] (verification not implemented)	509
3.28.6	Sympy [B] (verification not implemented)	510
3.28.7	Maxima [A] (verification not implemented)	510
3.28.8	Giac [B] (verification not implemented)	511
3.28.9	Mupad [B] (verification not implemented)	511

3.28.1 Optimal result

Integrand size = 26, antiderivative size = 140

$$\begin{aligned} & \int (a + ia \tan(c + dx))^4 (A + B \tan(c + dx)) dx \\ &= 8a^4(A - iB)x - \frac{8a^4(iA + B) \log(\cos(c + dx))}{d} \\ & \quad - \frac{4a^4(A - iB) \tan(c + dx)}{d} + \frac{a(iA + B)(a + ia \tan(c + dx))^3}{3d} \\ & \quad + \frac{B(a + ia \tan(c + dx))^4}{4d} + \frac{(iA + B)(a^2 + ia^2 \tan(c + dx))^2}{d} \end{aligned}$$

```
output 8*a^4*(A-I*B)*x-8*a^4*(I*A+B)*ln(cos(d*x+c))/d-4*a^4*(A-I*B)*tan(d*x+c)/d+
1/3*a*(I*A+B)*(a+I*a*tan(d*x+c))^3/d+1/4*B*(a+I*a*tan(d*x+c))^4/d+(I*A+B)*
(a^2+I*a^2*tan(d*x+c))^2/d
```

3.28.2 Mathematica [A] (verified)

Time = 1.02 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.61

$$\begin{aligned} & \int (a + ia \tan(c + dx))^4 (A + B \tan(c + dx)) dx \\ &= \frac{B(a + ia \tan(c + dx))^4 + \frac{4}{3}a^4(iA + B)(4 + 24 \log(i + \tan(c + dx)) + 21i \tan(c + dx) - 6 \tan^2(c + dx)) - 6 \tan^2(c + dx)}{4d} \end{aligned}$$

input `Integrate[(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(B*(a + I*a*Tan[c + d*x])^4 + (4*a^4*(I*A + B)*(4 + 24*Log[I + Tan[c + d*x]]) + (21*I)*Tan[c + d*x] - 6*Tan[c + d*x]^2 - I*Tan[c + d*x]^3))/3)/(4*d)`

3.28.3 Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.91, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {3042, 4010, 3042, 3959, 3042, 3959, 3042, 3958, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(c + dx))^4 (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(c + dx))^4 (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4010} \\
 & (A - iB) \int (i \tan(c + dx)a + a)^4 dx + \frac{B(a + ia \tan(c + dx))^4}{4d} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \int (i \tan(c + dx)a + a)^4 dx + \frac{B(a + ia \tan(c + dx))^4}{4d} \\
 & \quad \downarrow \text{3959} \\
 & (A - iB) \left(2a \int (i \tan(c + dx)a + a)^3 dx + \frac{ia(a + ia \tan(c + dx))^3}{3d} \right) + \frac{B(a + ia \tan(c + dx))^4}{4d} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \left(2a \int (i \tan(c + dx)a + a)^3 dx + \frac{ia(a + ia \tan(c + dx))^3}{3d} \right) + \frac{B(a + ia \tan(c + dx))^4}{4d} \\
 & \quad \downarrow \text{3959}
 \end{aligned}$$

$$iB) \left(2a \left(2a \int (i \tan(c + dx)a + a)^2 dx + \frac{(A - ia(a + ia \tan(c + dx))^2)}{2d} \right) + \frac{ia(a + ia \tan(c + dx))^3}{3d} \right) + \frac{B(a + ia \tan(c + dx))^4}{4d}$$

↓ 3042

$$iB) \left(2a \left(2a \int (i \tan(c + dx)a + a)^2 dx + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \frac{ia(a + ia \tan(c + dx))^3}{3d} \right) + \frac{B(a + ia \tan(c + dx))^4}{4d}$$

↓ 3958

$$iB) \left(2a \left(2a \left(2ia^2 \int \tan(c + dx) dx - \frac{a^2 \tan(c + dx)}{d} + 2a^2 x \right) + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \frac{ia(a + ia \tan(c + dx))^3}{3d} \right) + \frac{B(a + ia \tan(c + dx))^4}{4d}$$

↓ 3042

$$iB) \left(2a \left(2a \left(2ia^2 \int \tan(c + dx) dx - \frac{a^2 \tan(c + dx)}{d} + 2a^2 x \right) + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \frac{ia(a + ia \tan(c + dx))^3}{3d} \right) + \frac{B(a + ia \tan(c + dx))^4}{4d}$$

↓ 3956

$$iB) \left(2a \left(2a \left(-\frac{a^2 \tan(c + dx)}{d} - \frac{2ia^2 \log(\cos(c + dx))}{d} + 2a^2 x \right) + \frac{ia(a + ia \tan(c + dx))^2}{2d} \right) + \frac{ia(a + ia \tan(c + dx))^3}{3d} \right) + \frac{B(a + ia \tan(c + dx))^4}{4d}$$

input `Int[(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(B*(a + I*a*Tan[c + d*x])^4)/(4*d) + (A - I*B)*(((I/3)*a*(a + I*a*Tan[c + d*x])^3)/d + 2*a*(((I/2)*a*(a + I*a*Tan[c + d*x])^2)/d + 2*a*(2*a^2*x - ((2*I)*a^2*Log[Cos[c + d*x]]))/d - (a^2*Tan[c + d*x])/d))`

3.28.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 3958 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^2, x_Symbol] := Simp[(a^2 - b^2)*x, x] + (Simp[b^2*(Tan[c + d*x]/d), x] + Simp[2*a*b Int[Tan[c + d*x], x], x]) /; FreeQ[{a, b, c, d}, x]`

rule 3959 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

3.28.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.87

method	result
derivativdivides	$a^4 \left(-\frac{4iB(\tan^3(dx+c))}{3} + \frac{B(\tan^4(dx+c))}{4} - 2iA(\tan^2(dx+c)) + \frac{A(\tan^3(dx+c))}{3} + 8iB \tan(dx+c) - \frac{7B(\tan^2(dx+c))}{2} - 7A \tan(dx+c) \right) \frac{1}{d}$
default	$a^4 \left(-\frac{4iB(\tan^3(dx+c))}{3} + \frac{B(\tan^4(dx+c))}{4} - 2iA(\tan^2(dx+c)) + \frac{A(\tan^3(dx+c))}{3} + 8iB \tan(dx+c) - \frac{7B(\tan^2(dx+c))}{2} - 7A \tan(dx+c) \right) \frac{1}{d}$
norman	$(-8iB a^4 + 8A a^4) x - \frac{(4iA a^4 + 7B a^4)(\tan^2(dx+c))}{2d} - \frac{(-8iB a^4 + 7A a^4) \tan(dx+c)}{d} + \frac{(-4iB a^4 + A a^4) \ln(1+\tan^2(dx+c))}{3d}$
parallelrisch	$\frac{-16iB(\tan^3(dx+c))a^4 + 3B(\tan^4(dx+c))a^4 - 24iA(\tan^2(dx+c))a^4 + 4A(\tan^3(dx+c))a^4 - 96iBx a^4 d + 48iA \ln(1+\tan^2(dx+c))}{12d}$
risch	$\frac{16ia^4Bc}{d} - \frac{16a^4Ac}{d} - \frac{4a^4(18iA e^{6i(dx+c)} + 30B e^{6i(dx+c)} + 45iA e^{4i(dx+c)} + 63B e^{4i(dx+c)} + 38iA e^{2i(dx+c)} + 50B e^{2i(dx+c)})}{3d(e^{2i(dx+c)}+1)^4}$
parts	$A a^4 x + \frac{(-4iA a^4 - 6B a^4) \left(\frac{\tan^2(dx+c)}{2} - \frac{\ln(1+\tan^2(dx+c))}{2} \right)}{d} + \frac{(-4iB a^4 + A a^4) \left(\frac{\tan^3(dx+c)}{3} - \tan(dx+c) \right)}{d}$

```
input int((a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/d*a^4*(-4/3*I*B*tan(d*x+c)^3+1/4*B*tan(d*x+c)^4-2*I*A*tan(d*x+c)^2+1/3*A
*tan(d*x+c)^3+8*I*B*tan(d*x+c)-7/2*B*tan(d*x+c)^2-7*A*tan(d*x+c)+1/2*(8*B+
8*I*A)*ln(1+tan(d*x+c)^2)+(8*A-8*I*B)*arctan(tan(d*x+c)))
```

3.28.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.62

$$\int (a + ia \tan(c + dx))^4 (A + B \tan(c + dx)) dx = \frac{4(6(3iA + 5B)a^4 e^{(6i dx + 6i c)} + 9(5iA + 7B)a^4 e^{(4i dx + 4i c)} + 2(19iA + 25B)a^4 e^{(2i dx + 2i c)} + (11iA + 11B)a^4 e^{(0i dx + 0i c)})}{3(d e^{(8i dx + 8i c)} + 1)}$$

```
input integrate((a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output -4/3*(6*(3*I*A + 5*B))*a^4*e^(6*I*d*x + 6*I*c) + 9*(5*I*A + 7*B))*a^4*e^(4*I
*d*x + 4*I*c) + 2*(19*I*A + 25*B))*a^4*e^(2*I*d*x + 2*I*c) + (11*I*A + 14*B
)*a^4 + 6*((I*A + B))*a^4*e^(8*I*d*x + 8*I*c) + 4*(I*A + B))*a^4*e^(6*I*d*x
+ 6*I*c) + 6*(I*A + B))*a^4*e^(4*I*d*x + 4*I*c) + 4*(I*A + B))*a^4*e^(2*I*d*
x + 2*I*c) + (I*A + B))*a^4*log(e^(2*I*d*x + 2*I*c) + 1)/(d*e^(8*I*d*x +
8*I*c) + 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*
x + 2*I*c) + d)
```

3.28. $\int (a + ia \tan(c + dx))^4 (A + B \tan(c + dx)) dx$

3.28.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 241 vs. $2(116) = 232$.

Time = 0.44 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.72

$$\int (a + ia \tan(c + dx))^4 (A + B \tan(c + dx)) dx = -\frac{8ia^4(A - iB) \log(e^{2idx} + e^{-2ic})}{d} + \frac{-44iAa^4 - 56Ba^4 + (-152iAa^4e^{2ic} - 200Ba^4e^{2ic})e^{2idx} + (-180iAa^4e^{4ic} - 252Ba^4e^{4ic})e^{4idx} + (-72iAa^4e^{6ic} - 120Ba^4e^{6ic})e^{6idx}}{3de^{8ic}e^{8idx} + 12de^{6ic}e^{6idx} + 18de^{4ic}e^{4idx} + 12de^{2ic}e^{2idx} + 3d}$$

input `integrate((a+I*a*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)`

output `-8*I*a**4*(A - I*B)*log(exp(2*I*d*x) + exp(-2*I*c))/d + (-44*I*A*a**4 - 56*B*a**4 + (-152*I*A*a**4*exp(2*I*c) - 200*B*a**4*exp(2*I*c))*exp(2*I*d*x) + (-180*I*A*a**4*exp(4*I*c) - 252*B*a**4*exp(4*I*c))*exp(4*I*d*x) + (-72*I*A*a**4*exp(6*I*c) - 120*B*a**4*exp(6*I*c))*exp(6*I*d*x))/(3*d*exp(8*I*c)*exp(8*I*d*x) + 12*d*exp(6*I*c)*exp(6*I*d*x) + 18*d*exp(4*I*c)*exp(4*I*d*x) + 12*d*exp(2*I*c)*exp(2*I*d*x) + 3*d)`

3.28.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.81

$$\int (a + ia \tan(c + dx))^4 (A + B \tan(c + dx)) dx = \frac{3Ba^4 \tan(dx + c)^4 + 4(A - 4iB)a^4 \tan(dx + c)^3 - 6(4iA + 7B)a^4 \tan(dx + c)^2 + 96(dx + c)(A - iB)a^4 \tan(dx + c) + 96(dx + c)^2(A - iB)a^4}{12d}$$

input `integrate((a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/12*(3*B*a^4*tan(d*x + c)^4 + 4*(A - 4*I*B)*a^4*tan(d*x + c)^3 - 6*(4*I*A + 7*B)*a^4*tan(d*x + c)^2 + 96*(d*x + c)*(A - I*B)*a^4 - 48*(-I*A - B)*a^4*log(tan(d*x + c)^2 + 1) - 12*(7*A - 8*I*B)*a^4*tan(d*x + c))/d`

3.28.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 408 vs. $2(120) = 240$.

Time = 0.54 (sec) , antiderivative size = 408, normalized size of antiderivative = 2.91

$$\int (a + ia \tan(c + dx))^4 (A + B \tan(c + dx)) dx =$$

$$4 (6i Aa^4 e^{(8i dx + 8i c)} \log(e^{(2i dx + 2i c)} + 1) + 6 Ba^4 e^{(8i dx + 8i c)} \log(e^{(2i dx + 2i c)} + 1) + 24i Aa^4 e^{(6i dx + 6i c)} \log(e^{(2i dx + 2i c)} + 1) + 24i Ba^4 e^{(6i dx + 6i c)} \log(e^{(2i dx + 2i c)} + 1) + 36i Aa^4 e^{(4i dx + 4i c)} \log(e^{(2i dx + 2i c)} + 1) + 36i Ba^4 e^{(4i dx + 4i c)} \log(e^{(2i dx + 2i c)} + 1) + 24i Aa^4 e^{(2i dx + 2i c)} \log(e^{(2i dx + 2i c)} + 1) + 24i Ba^4 e^{(2i dx + 2i c)} \log(e^{(2i dx + 2i c)} + 1) + 18i Aa^4 e^{(6i dx + 6i c)} + 30i Ba^4 e^{(6i dx + 6i c)} + 45i Aa^4 e^{(4i dx + 4i c)} + 63i Ba^4 e^{(4i dx + 4i c)} + 38i Aa^4 e^{(2i dx + 2i c)} + 50i Ba^4 e^{(2i dx + 2i c)} + 6i Aa^4 \log(e^{(2i dx + 2i c)} + 1) + 6i Ba^4 \log(e^{(2i dx + 2i c)} + 1) + 11i Aa^4 + 14i Ba^4) / (d e^{(8i dx + 8i c)} + 4d e^{(6i dx + 6i c)} + 6d e^{(4i dx + 4i c)} + 4d e^{(2i dx + 2i c)} + d)$$

input `integrate((a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

```
-4/3*(6*I*A*a^4*e^(8*I*d*x + 8*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 6*B*a^4
*e^(8*I*d*x + 8*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 24*I*A*a^4*e^(6*I*d*x
+ 6*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 24*B*a^4*e^(6*I*d*x + 6*I*c)*log(e
^(2*I*d*x + 2*I*c) + 1) + 36*I*A*a^4*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x +
2*I*c) + 1) + 36*B*a^4*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) +
24*I*A*a^4*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 24*B*a^4*e^(
2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) + 18*I*A*a^4*e^(6*I*d*x + 6*
I*c) + 30*B*a^4*e^(6*I*d*x + 6*I*c) + 45*I*A*a^4*e^(4*I*d*x + 4*I*c) + 63*
B*a^4*e^(4*I*d*x + 4*I*c) + 38*I*A*a^4*e^(2*I*d*x + 2*I*c) + 50*B*a^4*e^(2
*I*d*x + 2*I*c) + 6*I*A*a^4*log(e^(2*I*d*x + 2*I*c) + 1) + 6*B*a^4*log(e^(
2*I*d*x + 2*I*c) + 1) + 11*I*A*a^4 + 14*B*a^4)/(d*e^(8*I*d*x + 8*I*c) + 4*
d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*x + 2*I*c)
+ d)
```

3.28.9 Mupad [B] (verification not implemented)

Time = 7.21 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.29

$$\int (a + ia \tan(c + dx))^4 (A + B \tan(c + dx)) dx$$

$$= \frac{Ba^4 \tan(c + dx)^4}{4d} + \frac{\ln(\tan(c + dx) + i) (8Ba^4 + Aa^4 8i)}{d}$$

$$- \frac{\tan(c + dx)^2 \left(\frac{a^4(A-Bi)3i}{2} + \frac{Ba^4}{2} + \frac{a^4(3B+Ai)}{2} \right)}{d}$$

$$+ \frac{\tan(c + dx) (-3a^4(A - Bi) + a^4(B + A3i) li + Ba^4 li + a^4(3B + Ai) li)}{d}$$

$$- \frac{\tan(c + dx)^3 \left(\frac{Ba^4 li}{3} + \frac{a^4(3B+Ai) li}{3} \right)}{d}$$

3.28. $\int (a + ia \tan(c + dx))^4 (A + B \tan(c + dx)) dx$

input `int((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^4,x)`

output `(log(tan(c + d*x) + 1i)*(A*a^4*8i + 8*B*a^4))/d - (tan(c + d*x)^3*((B*a^4*1i)/3 + (a^4*(A*1i + 3*B)*1i)/3))/d - (tan(c + d*x)^2*((a^4*(A - B*1i)*3i)/2 + (B*a^4)/2 + (a^4*(A*1i + 3*B))/2))/d + (tan(c + d*x)*(a^4*(A*3i + B)*1i - 3*a^4*(A - B*1i) + B*a^4*1i + a^4*(A*1i + 3*B)*1i))/d + (B*a^4*tan(c + d*x)^4)/(4*d)`

3.29 $\int \cot(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.29.1	Optimal result	513
3.29.2	Mathematica [A] (verified)	514
3.29.3	Rubi [A] (verified)	514
3.29.4	Maple [A] (verified)	518
3.29.5	Fricas [B] (verification not implemented)	518
3.29.6	Sympy [B] (verification not implemented)	519
3.29.7	Maxima [A] (verification not implemented)	520
3.29.8	Giac [B] (verification not implemented)	520
3.29.9	Mupad [B] (verification not implemented)	521

3.29.1 Optimal result

Integrand size = 32, antiderivative size = 142

$$\int \cot(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= 8a^4(iA+B)x + \frac{a^4(7A-8iB) \log(\cos(c+dx))}{d}$$

$$+ \frac{a^4 A \log(\sin(c+dx))}{d} + \frac{iaB(a+ia \tan(c+dx))^3}{3d}$$

$$- \frac{(A-2iB)(a^2+ia^2 \tan(c+dx))^2}{2d} - \frac{(3A-4iB)(a^4+ia^4 \tan(c+dx))}{d}$$

```
output 8*a^4*(I*A+B)*x+a^4*(7*A-8*I*B)*ln(cos(d*x+c))/d+a^4*A*ln(sin(d*x+c))/d+1/
3*I*a*B*(a+I*a*tan(d*x+c))^3/d-1/2*(A-2*I*B)*(a^2+I*a^2*tan(d*x+c))^2/d-(3
*A-4*I*B)*(a^4+I*a^4*tan(d*x+c))/d
```


3.29.2 Mathematica [A] (verified)

Time = 1.21 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.63

$$\int \cot(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{a^4(6A \log(\tan(c + dx)) - 8(A - iB)(1 + 6 \log(i + \tan(c + dx))) + (-24iA - 42B) \tan(c + dx) + 3(A -$$

$$6d$$

input `Integrate[Cot[c + d*x]*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(a^4*(6*A*Log[Tan[c + d*x]] - 8*(A - I*B)*(1 + 6*Log[I + Tan[c + d*x]]) + ((-24*I)*A - 42*B)*Tan[c + d*x] + 3*(A - (4*I)*B)*Tan[c + d*x]^2 + 2*B*Tan[c + d*x]^3))/(6*d)`

3.29.3 Rubi [A] (verified)

Time = 1.07 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.01, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4077, 27, 3042, 4077, 27, 3042, 4077, 3042, 4072, 3042, 3956, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + ia \tan(c + dx))^4(A + B \tan(c + dx))}{\tan(c + dx)} dx$$

$$\downarrow \text{4077}$$

$$\frac{1}{3} \int 3 \cot(c + dx)(i \tan(c + dx)a + a)^3(aA + a(iA + 2B) \tan(c + dx)) dx + \frac{iaB(a + ia \tan(c + dx))^3}{3d}$$

$$\downarrow \text{27}$$

$$\int \cot(c + dx)(i \tan(c + dx)a + a)^3(aA + a(iA + 2B) \tan(c + dx)) dx + \frac{iaB(a + ia \tan(c + dx))^3}{3d}$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
& \int \frac{(i \tan(c+dx)a+a)^3(aA+a(iA+2B)\tan(c+dx))}{\tan(c+dx)} dx + \frac{iaB(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 4077 \\
& \frac{1}{2} \int 2 \cot(c+dx)(i \tan(c+dx)a+a)^2 (Aa^2+(3iA+4B)\tan(c+dx)a^2) dx - \\
& \quad \frac{(A-2iB)(a^2+ia^2 \tan(c+dx))^2}{2d} + \frac{iaB(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 27 \\
& \int \cot(c+dx)(i \tan(c+dx)a+a)^2 (Aa^2+(3iA+4B)\tan(c+dx)a^2) dx - \\
& \quad \frac{(A-2iB)(a^2+ia^2 \tan(c+dx))^2}{2d} + \frac{iaB(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 3042 \\
& \int \frac{(i \tan(c+dx)a+a)^2 (Aa^2+(3iA+4B)\tan(c+dx)a^2)}{\tan(c+dx)} dx - \\
& \quad \frac{(A-2iB)(a^2+ia^2 \tan(c+dx))^2}{2d} + \frac{iaB(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 4077 \\
& \int \cot(c+dx)(i \tan(c+dx)a+a) (Aa^3+(7iA+8B)\tan(c+dx)a^3) dx - \\
& \frac{(3A-4iB)(a^4+ia^4 \tan(c+dx))}{d} - \frac{(A-2iB)(a^2+ia^2 \tan(c+dx))^2}{2d} + \frac{iaB(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 3042 \\
& \int \frac{(i \tan(c+dx)a+a) (Aa^3+(7iA+8B)\tan(c+dx)a^3)}{\tan(c+dx)} dx - \\
& \frac{(3A-4iB)(a^4+ia^4 \tan(c+dx))}{d} - \frac{(A-2iB)(a^2+ia^2 \tan(c+dx))^2}{2d} + \frac{iaB(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 4072 \\
& - \left(a^4(7A-8iB) \int \tan(c+dx) dx \right) + \int \cot(c+dx) (Aa^4+8(iA+B)\tan(c+dx)a^4) dx - \\
& \frac{(3A-4iB)(a^4+ia^4 \tan(c+dx))}{d} - \frac{(A-2iB)(a^2+ia^2 \tan(c+dx))^2}{2d} + \frac{iaB(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 3042 \\
& - \left(a^4(7A-8iB) \int \tan(c+dx) dx \right) + \int \frac{Aa^4+8(iA+B)\tan(c+dx)a^4}{\tan(c+dx)} dx - \\
& \frac{(3A-4iB)(a^4+ia^4 \tan(c+dx))}{d} - \frac{(A-2iB)(a^2+ia^2 \tan(c+dx))^2}{2d} + \frac{iaB(a+ia \tan(c+dx))^3}{3d}
\end{aligned}$$

3.29. $\int \cot(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \int \frac{Aa^4 + 8(iA + B) \tan(c + dx)a^4}{\tan(c + dx)} dx - \frac{(3A - 4iB)(a^4 + ia^4 \tan(c + dx))}{d} + \\
& \frac{a^4(7A - 8iB) \log(\cos(c + dx))}{d} - \frac{(A - 2iB)(a^2 + ia^2 \tan(c + dx))^2}{2d} + \frac{iaB(a + ia \tan(c + dx))^3}{3d} \\
& \downarrow \text{3956} \\
& a^4 A \int \cot(c + dx) dx - \frac{(3A - 4iB)(a^4 + ia^4 \tan(c + dx))}{d} + \frac{a^4(7A - 8iB) \log(\cos(c + dx))}{d} + \\
& 8a^4 x(B + iA) - \frac{(A - 2iB)(a^2 + ia^2 \tan(c + dx))^2}{2d} + \frac{iaB(a + ia \tan(c + dx))^3}{3d} \\
& \downarrow \text{4014} \\
& a^4 A \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - \frac{(3A - 4iB)(a^4 + ia^4 \tan(c + dx))}{d} + \\
& \frac{a^4(7A - 8iB) \log(\cos(c + dx))}{d} + 8a^4 x(B + iA) - \frac{(A - 2iB)(a^2 + ia^2 \tan(c + dx))^2}{2d} + \\
& \frac{iaB(a + ia \tan(c + dx))^3}{3d} \\
& \downarrow \text{3042} \\
& -a^4 A \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - \frac{(3A - 4iB)(a^4 + ia^4 \tan(c + dx))}{d} + \\
& \frac{a^4(7A - 8iB) \log(\cos(c + dx))}{d} + 8a^4 x(B + iA) - \frac{(A - 2iB)(a^2 + ia^2 \tan(c + dx))^2}{2d} + \\
& \frac{iaB(a + ia \tan(c + dx))^3}{3d} \\
& \downarrow \text{25} \\
& -a^4 A \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - \frac{(3A - 4iB)(a^4 + ia^4 \tan(c + dx))}{d} + \\
& \frac{a^4(7A - 8iB) \log(\cos(c + dx))}{d} + 8a^4 x(B + iA) - \frac{(A - 2iB)(a^2 + ia^2 \tan(c + dx))^2}{2d} + \\
& \frac{iaB(a + ia \tan(c + dx))^3}{3d} \\
& \downarrow \text{3956} \\
& -\frac{(3A - 4iB)(a^4 + ia^4 \tan(c + dx))}{d} + \frac{a^4(7A - 8iB) \log(\cos(c + dx))}{d} + 8a^4 x(B + iA) + \\
& \frac{a^4 A \log(-\sin(c + dx))}{d} - \frac{(A - 2iB)(a^2 + ia^2 \tan(c + dx))^2}{2d} + \frac{iaB(a + ia \tan(c + dx))^3}{3d}
\end{aligned}$$

input `Int[Cot[c + d*x]*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `8*a^4*(I*A + B)*x + (a^4*(7*A - (8*I)*B)*Log[Cos[c + d*x]]/d + (a^4*A*Log[-Sin[c + d*x]]/d + ((I/3)*a*B*(a + I*a*Tan[c + d*x])^3)/d - ((A - (2*I)*B)*(a^2 + I*a^2*Tan[c + d*x])^2)/(2*d) - ((3*A - (4*I)*B)*(a^4 + I*a^4*Tan[c + d*x]))/d`

3.29. $\int \cot(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$

3.29.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4014 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`
- rule 4072 `Int((((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]))/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*(d/b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c - a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0]`
- rule 4077 `Int(((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.29.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.77

method	result
parallelrisch	$-\frac{a^4(-48iAxd+12iB(\tan^2(dx+c))-2B(\tan^3(dx+c))+24iA \tan(dx+c)-3A(\tan^2(dx+c))-24iB \ln(\sec^2(dx+c))-48iB \ln(\tan(dx+c)))}{6d}$
derivativedivides	$a^4 \left(\frac{(8iB-8A) \ln(\cot^2(dx+c)+1)}{2} + (-8iA-8B) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)) \right) + (-8iB+7A) \ln(\cot(dx+c)) - \frac{4iA+7B}{\cot(dx+c)} - \frac{4iB-8A}{2 \cot(dx+c)} \right) / d$
default	$a^4 \left(\frac{(8iB-8A) \ln(\cot^2(dx+c)+1)}{2} + (-8iA-8B) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)) \right) + (-8iB+7A) \ln(\cot(dx+c)) - \frac{4iA+7B}{\cot(dx+c)} - \frac{4iB-8A}{2 \cot(dx+c)} \right) / d$
norman	$(8iA a^4 + 8B a^4) x - \frac{(4iA a^4 + 7B a^4) \tan(dx+c)}{d} + \frac{(-4iB a^4 + A a^4) (\tan^2(dx+c))}{2d} + \frac{B a^4 (\tan^3(dx+c))}{3d} + \frac{(-4iA a^4 + 7B a^4) \ln(\sec^2(dx+c))}{2d} - \frac{4iB a^4 \ln(\tan(dx+c))}{d}$
risch	$-\frac{16a^4 Bc}{d} - \frac{16ia^4 Ac}{d} - \frac{2ia^4(15iA e^{4i(dx+c)} + 36B e^{4i(dx+c)} + 27iA e^{2i(dx+c)} + 54B e^{2i(dx+c)} + 12iA + 22B)}{3d(e^{2i(dx+c)} + 1)^3} - \frac{8ia^4}{d}$

input `int(cot(d*x+c)*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `-1/6*a^4*(-48*I*A*x*d+12*I*B*tan(d*x+c)^2-2*B*tan(d*x+c)^3+24*I*A*tan(d*x+c)-3*A*tan(d*x+c)^2-24*I*B*ln(sec(d*x+c)^2)-48*B*d*x+24*A*ln(sec(d*x+c)^2)-6*A*ln(tan(d*x+c))+42*B*tan(d*x+c))/d`

3.29.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 246 vs. 2(122) = 244.

Time = 0.25 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.73

$$\int \cot(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= \frac{6(5A-12iB)a^4 e^{(4i dx+4i c)} + 54(A-2iB)a^4 e^{(2i dx+2i c)} + 4(6A-11iB)a^4 + 3((7A-8iB)a^4 e^{(6i dx+6i c)} + 12iAa^4 e^{(4i dx+4i c)} + 12iBa^4 e^{(2i dx+2i c)} + 12iAa^4 + 12iBa^4)}{3d(e^{2i(dx+c)}+1)^3}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fricas")`

```
output 1/3*(6*(5*A - 12*I*B)*a^4*e^(4*I*d*x + 4*I*c) + 54*(A - 2*I*B)*a^4*e^(2*I*
d*x + 2*I*c) + 4*(6*A - 11*I*B)*a^4 + 3*((7*A - 8*I*B)*a^4*e^(6*I*d*x + 6*
I*c) + 3*(7*A - 8*I*B)*a^4*e^(4*I*d*x + 4*I*c) + 3*(7*A - 8*I*B)*a^4*e^(2*
I*d*x + 2*I*c) + (7*A - 8*I*B)*a^4)*log(e^(2*I*d*x + 2*I*c) + 1) + 3*(A*a^
4*e^(6*I*d*x + 6*I*c) + 3*A*a^4*e^(4*I*d*x + 4*I*c) + 3*A*a^4*e^(2*I*d*x +
2*I*c) + A*a^4)*log(e^(2*I*d*x + 2*I*c) - 1))/(d*e^(6*I*d*x + 6*I*c) + 3*
d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)
```

3.29.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 289 vs. $2(121) = 242$.

Time = 1.85 (sec) , antiderivative size = 289, normalized size of antiderivative = 2.04

$$\int \cot(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{Aa^4 \log\left(\frac{-3Aa^4 + 4iBa^4}{3Aa^4 e^{2ic} - 4iBa^4 e^{2ic}} + e^{2idx}\right)}{d}$$

$$+ \frac{a^4 \cdot (7A - 8iB) \log\left(e^{2idx} + \frac{-4Aa^4 + 4iBa^4 + a^4 \cdot (7A - 8iB)}{3Aa^4 e^{2ic} - 4iBa^4 e^{2ic}}\right)}{d}$$

$$+ \frac{24Aa^4 - 44iBa^4 + (54Aa^4 e^{2ic} - 108iBa^4 e^{2ic}) e^{2idx} + (30Aa^4 e^{4ic} - 72iBa^4 e^{4ic}) e^{4idx}}{3de^{6ic} e^{6idx} + 9de^{4ic} e^{4idx} + 9de^{2ic} e^{2idx} + 3d}$$

```
input integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)
```

```
output A***4*log((-3*A*a***4 + 4*I*B*a***4)/(3*A*a***4*exp(2*I*c) - 4*I*B*a***4*exp(
2*I*c)) + exp(2*I*d*x))/d + a***4*(7*A - 8*I*B)*log(exp(2*I*d*x) + (-4*A*a*
**4 + 4*I*B*a***4 + a***4*(7*A - 8*I*B))/(3*A*a***4*exp(2*I*c) - 4*I*B*a***4*ex
p(2*I*c)))/d + (24*A*a***4 - 44*I*B*a***4 + (54*A*a***4*exp(2*I*c) - 108*I*B*
a***4*exp(2*I*c))*exp(2*I*d*x) + (30*A*a***4*exp(4*I*c) - 72*I*B*a***4*exp(4*
I*c))*exp(4*I*d*x))/(3*d*exp(6*I*c)*exp(6*I*d*x) + 9*d*exp(4*I*c)*exp(4*I*
d*x) + 9*d*exp(2*I*c)*exp(2*I*d*x) + 3*d)
```

3.29.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.75

$$\int \cot(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{2Ba^4 \tan(dx + c)^3 + 3(A - 4iB)a^4 \tan(dx + c)^2 - 48(dx + c)(-iA - B)a^4 - 24(A - iB)a^4 \log(\tan(dx + c))}{6d}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/6*(2*B*a^4*tan(d*x + c)^3 + 3*(A - 4*I*B)*a^4*tan(d*x + c)^2 - 48*(d*x + c)*(-I*A - B)*a^4 - 24*(A - I*B)*a^4*log(tan(d*x + c)^2 + 1) + 6*A*a^4*log(tan(d*x + c)) - 6*(4*I*A + 7*B)*a^4*tan(d*x + c))/d`

3.29.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 332 vs. 2(122) = 244.

Time = 1.23 (sec) , antiderivative size = 332, normalized size of antiderivative = 2.34

$$\int \cot(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{6Aa^4 \log(\tan(\frac{1}{2}dx + \frac{1}{2}c)) + 6(7Aa^4 - 8iBa^4) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) + 1) - 96(Aa^4 - iBa^4) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1) + 6(7Aa^4 - 8iB^2a^4) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) + I) + 6(7Aa^4 - 8iB^2a^4) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1) - (77Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^6 - 88iB^2a^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^6 - 48iAa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 - 84B^2a^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 - 243Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^4 + 312iB^2a^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^4 + 96iAa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 + 184B^2a^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 + 243Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 312iB^2a^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 48iAa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) - 84B^2a^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) - 77Aa^4 + 88iB^2a^4)}{(\tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 1)^3}d$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `1/6*(6*A*a^4*log(tan(1/2*d*x + 1/2*c)) + 6*(7*A*a^4 - 8*I*B*a^4)*log(tan(1/2*d*x + 1/2*c) + 1) - 96*(A*a^4 - I*B*a^4)*log(tan(1/2*d*x + 1/2*c) + I) + 6*(7*A*a^4 - 8*I*B*a^4)*log(tan(1/2*d*x + 1/2*c) - 1) - (77*A*a^4*tan(1/2*d*x + 1/2*c)^6 - 88*I*B*a^4*tan(1/2*d*x + 1/2*c)^6 - 48*I*A*a^4*tan(1/2*d*x + 1/2*c)^5 - 84*B*a^4*tan(1/2*d*x + 1/2*c)^5 - 243*A*a^4*tan(1/2*d*x + 1/2*c)^4 + 312*I*B*a^4*tan(1/2*d*x + 1/2*c)^4 + 96*I*A*a^4*tan(1/2*d*x + 1/2*c)^3 + 184*B*a^4*tan(1/2*d*x + 1/2*c)^3 + 243*A*a^4*tan(1/2*d*x + 1/2*c)^2 - 312*I*B*a^4*tan(1/2*d*x + 1/2*c)^2 - 48*I*A*a^4*tan(1/2*d*x + 1/2*c) - 84*B*a^4*tan(1/2*d*x + 1/2*c) - 77*A*a^4 + 88*I*B*a^4)/(tan(1/2*d*x + 1/2*c)^2 - 1)^3/d`

3.29.9 Mupad [B] (verification not implemented)

Time = 7.24 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.94

$$\int \cot(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{A a^4 \ln(\tan(c + dx))}{d} - \frac{\tan(c + dx) (a^4 (A - B i) 3i + B a^4 + a^4 (3B + A i))}{d}$$

$$- \frac{\tan(c + dx)^2 \left(\frac{B a^4 i}{2} + \frac{a^4 (3B + A i) i}{2} \right)}{d}$$

$$- \frac{8 a^4 \ln(\tan(c + dx) + i) (A - B i)}{d} + \frac{B a^4 \tan(c + dx)^3}{3 d}$$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^4,x)`output `(A*a^4*log(tan(c + d*x)))/d - (tan(c + d*x)*(a^4*(A - B*1i)*3i + B*a^4 + a^4*(A*1i + 3*B)))/d - (tan(c + d*x)^2*((B*a^4*1i)/2 + (a^4*(A*1i + 3*B)*1i)/2))/d - (8*a^4*log(tan(c + d*x) + 1i)*(A - B*1i))/d + (B*a^4*tan(c + d*x)^3)/(3*d)`

3.30 $\int \cot^2(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.30.1	Optimal result	522
3.30.2	Mathematica [A] (verified)	523
3.30.3	Rubi [A] (verified)	523
3.30.4	Maple [A] (verified)	527
3.30.5	Fricas [B] (verification not implemented)	528
3.30.6	Sympy [B] (verification not implemented)	529
3.30.7	Maxima [A] (verification not implemented)	530
3.30.8	Giac [B] (verification not implemented)	530
3.30.9	Mupad [B] (verification not implemented)	531

3.30.1 Optimal result

Integrand size = 34, antiderivative size = 144

$$\begin{aligned} & \int \cot^2(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx \\ &= -8a^4(A-iB)x + \frac{a^4(4iA+7B) \log(\cos(c+dx))}{d} \\ & \quad + \frac{a^4(4iA+B) \log(\sin(c+dx))}{d} - \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^3}{d} \\ & \quad + \frac{(2iA-B)(a^2+ia^2 \tan(c+dx))^2}{2d} - \frac{3B(a^4+ia^4 \tan(c+dx))}{d} \end{aligned}$$

output

```
-8*a^4*(A-I*B)*x+a^4*(4*I*A+7*B)*ln(cos(d*x+c))/d+a^4*(4*I*A+B)*ln(sin(d*x+c))/d-a*A*cot(d*x+c)*(a+I*a*tan(d*x+c))^3/d+1/2*(2*I*A-B)*(a^2+I*a^2*tan(d*x+c))^2/d-3*B*(a^4+I*a^4*tan(d*x+c))/d
```

3.30.2 Mathematica [A] (verified)

Time = 1.37 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.84

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= a^4 \left(-\frac{A \cot(c + dx)}{d} + \frac{4iA \log(\tan(c + dx))}{d} + \frac{B \log(\tan(c + dx))}{d} \right. \\ \left. - \frac{8iA \log(i + \tan(c + dx))}{d} - \frac{8B \log(i + \tan(c + dx))}{d} + \frac{A \tan(c + dx)}{d} \right. \\ \left. - \frac{4iB \tan(c + dx)}{d} + \frac{B \tan^2(c + dx)}{2d} \right)$$

input `Integrate[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `a^4*(-((A*Cot[c + d*x])/d) + ((4*I)*A*Log[Tan[c + d*x]])/d + (B*Log[Tan[c + d*x]])/d - ((8*I)*A*Log[I + Tan[c + d*x]])/d - (8*B*Log[I + Tan[c + d*x]])/d + (A*Tan[c + d*x])/d - ((4*I)*B*Tan[c + d*x])/d + (B*Tan[c + d*x]^2)/(2*d))`

3.30.3 Rubi [A] (verified)

Time = 1.13 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.441$, Rules used = {3042, 4076, 3042, 4077, 27, 3042, 4077, 3042, 4072, 3042, 3956, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + ia \tan(c + dx))^4(A + B \tan(c + dx))}{\tan(c + dx)^2} dx$$

$$\downarrow \text{4076}$$

$$\int \cot(c + dx)(i \tan(c + dx)a + a)^3(a(4iA + B) + a(2A + iB) \tan(c + dx)) dx - \frac{aA \cot(c + dx)(a + ia \tan(c + dx))^3}{d}$$

$$\begin{aligned}
& \int \frac{(i \tan(c+dx)a+a)^3(a(4iA+B)+a(2A+iB)\tan(c+dx))}{\tan(c+dx)} dx - \\
& \quad \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^3}{d} \\
& \quad \downarrow 3042 \\
& \frac{1}{2} \int 2 \cot(c+dx)(i \tan(c+dx)a+a)^2((4iA+B)a^2+3iB \tan(c+dx)a^2) dx + \\
& \quad \frac{(-B+2iA)(a^2+ia^2 \tan(c+dx))^2}{2d} - \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^3}{d} \\
& \quad \downarrow 4077 \\
& \int \cot(c+dx)(i \tan(c+dx)a+a)^2((4iA+B)a^2+3iB \tan(c+dx)a^2) dx + \\
& \quad \frac{(-B+2iA)(a^2+ia^2 \tan(c+dx))^2}{2d} - \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^3}{d} \\
& \quad \downarrow 27 \\
& \int \cot(c+dx)(i \tan(c+dx)a+a)^2((4iA+B)a^2+3iB \tan(c+dx)a^2) dx + \\
& \quad \frac{(-B+2iA)(a^2+ia^2 \tan(c+dx))^2}{2d} - \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^3}{d} \\
& \quad \downarrow 3042 \\
& \int \frac{(i \tan(c+dx)a+a)^2((4iA+B)a^2+3iB \tan(c+dx)a^2)}{\tan(c+dx)} dx + \\
& \quad \frac{(-B+2iA)(a^2+ia^2 \tan(c+dx))^2}{2d} - \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^3}{d} \\
& \quad \downarrow 4077 \\
& \int \cot(c+dx)(i \tan(c+dx)a+a)(a^3(4iA+B)-a^3(4A-7iB)\tan(c+dx)) dx - \\
& \quad \frac{3B(a^4+ia^4 \tan(c+dx))}{d} + \frac{(-B+2iA)(a^2+ia^2 \tan(c+dx))^2}{2d} - \\
& \quad \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^3}{d} \\
& \quad \downarrow 3042 \\
& \int \frac{(i \tan(c+dx)a+a)(a^3(4iA+B)-a^3(4A-7iB)\tan(c+dx))}{\tan(c+dx)} dx - \frac{3B(a^4+ia^4 \tan(c+dx))}{d} + \\
& \quad \frac{(-B+2iA)(a^2+ia^2 \tan(c+dx))^2}{2d} - \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^3}{d} \\
& \quad \downarrow 4072 \\
& - \left(a^4(7B+4iA) \int \tan(c+dx) dx \right) + \int \cot(c+ \\
& dx)(a^4(4iA+B)-8a^4(A-iB)\tan(c+dx)) dx - \frac{3B(a^4+ia^4 \tan(c+dx))}{d} + \\
& \quad \frac{(-B+2iA)(a^2+ia^2 \tan(c+dx))^2}{2d} - \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^3}{d}
\end{aligned}$$

3.30. $\int \cot^2(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \downarrow 3042 \\
& -\left(a^4(7B + 4iA) \int \tan(c + dx) dx\right) + \int \frac{a^4(4iA + B) - 8a^4(A - iB) \tan(c + dx)}{\tan(c + dx)} dx - \\
& \quad \frac{3B(a^4 + ia^4 \tan(c + dx))}{d} + \frac{(-B + 2iA)(a^2 + ia^2 \tan(c + dx))^2}{\frac{2d}{aA \cot(c + dx)(a + ia \tan(c + dx))^3}} - \\
& \quad \downarrow 3956 \\
& \int \frac{a^4(4iA + B) - 8a^4(A - iB) \tan(c + dx)}{\tan(c + dx)} dx + \frac{a^4(7B + 4iA) \log(\cos(c + dx))}{d} - \\
& \quad \frac{3B(a^4 + ia^4 \tan(c + dx))}{d} + \frac{(-B + 2iA)(a^2 + ia^2 \tan(c + dx))^2}{\frac{2d}{aA \cot(c + dx)(a + ia \tan(c + dx))^3}} - \\
& \quad \downarrow 4014 \\
& a^4(B + 4iA) \int \cot(c + dx) dx + \frac{a^4(7B + 4iA) \log(\cos(c + dx))}{d} - 8a^4x(A - iB) - \\
& \quad \frac{3B(a^4 + ia^4 \tan(c + dx))}{d} + \frac{(-B + 2iA)(a^2 + ia^2 \tan(c + dx))^2}{\frac{2d}{aA \cot(c + dx)(a + ia \tan(c + dx))^3}} - \\
& \quad \downarrow 3042 \\
& a^4(B + 4iA) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + \frac{a^4(7B + 4iA) \log(\cos(c + dx))}{d} - 8a^4x(A - iB) - \\
& \quad \frac{3B(a^4 + ia^4 \tan(c + dx))}{d} + \frac{(-B + 2iA)(a^2 + ia^2 \tan(c + dx))^2}{\frac{2d}{aA \cot(c + dx)(a + ia \tan(c + dx))^3}} - \\
& \quad \downarrow 25 \\
& -a^4(B + 4iA) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + \frac{a^4(7B + 4iA) \log(\cos(c + dx))}{d} - 8a^4x(A - iB) - \\
& \quad \frac{3B(a^4 + ia^4 \tan(c + dx))}{d} + \frac{(-B + 2iA)(a^2 + ia^2 \tan(c + dx))^2}{\frac{2d}{aA \cot(c + dx)(a + ia \tan(c + dx))^3}} - \\
& \quad \downarrow 3956
\end{aligned}$$

$$\frac{a^4(B + 4iA) \log(-\sin(c + dx))}{d} + \frac{a^4(7B + 4iA) \log(\cos(c + dx))}{d} - 8a^4x(A - iB) - \frac{3B(a^4 + ia^4 \tan(c + dx))}{d} + \frac{(-B + 2iA)(a^2 + ia^2 \tan(c + dx))^2}{2d} - \frac{aA \cot(c + dx)(a + ia \tan(c + dx))^3}{d}$$

input `Int[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `-8*a^4*(A - I*B)*x + (a^4*((4*I)*A + 7*B)*Log[Cos[c + d*x]])/d + (a^4*((4*I)*A + B)*Log[-Sin[c + d*x]])/d - (a*A*Cot[c + d*x]*(a + I*a*Tan[c + d*x])^3)/d + (((2*I)*A - B)*(a^2 + I*a^2*Tan[c + d*x])^2)/(2*d) - (3*B*(a^4 + I*a^4*Tan[c + d*x]))/d`

3.30.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

```
rule 4072 Int[(((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*(d/b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c - a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0]
```

```
rule 4076 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

```
rule 4077 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]
```

3.30.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.63

$$3.30. \quad \int \cot^2(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

method	result
parallelrisch	$\frac{4a^4 \left((-iA-B) \ln(\sec^2(dx+c)) + \left(iA + \frac{B}{4} \right) \ln(\tan(dx+c)) + \frac{B(\tan^2(dx+c))}{8} + \left(-iB + \frac{A}{4} \right) \tan(dx+c) - \frac{A \cot(dx+c)}{4} + 2xd \right)}{d}$
derivativedivides	$\frac{a^4 \left(-A \cot(dx+c) + \frac{(-8iA-8B) \ln(\cot^2(dx+c)+1)}{2} + (-8iB+8A) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)) \right) + (4iA+7B) \ln(\cot(dx+c)) - \frac{4}{\cot} \right)}{d}$
default	$\frac{a^4 \left(-A \cot(dx+c) + \frac{(-8iA-8B) \ln(\cot^2(dx+c)+1)}{2} + (-8iB+8A) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)) \right) + (4iA+7B) \ln(\cot(dx+c)) - \frac{4}{\cot} \right)}{d}$
norman	$\frac{(8iB a^4 - 8A a^4) x \tan(dx+c) + \frac{(-4iB a^4 + A a^4) (\tan^2(dx+c))}{d} - \frac{A a^4}{d} + \frac{B a^4 (\tan^3(dx+c))}{2d}}{\tan(dx+c)} + \frac{(4iA a^4 + B a^4) \ln(\tan(dx+c))}{d}$
risch	$-\frac{16ia^4 Bc}{d} + \frac{16a^4 Ac}{d} + \frac{2a^4 (5B e^{4i(dx+c)} - 2iA e^{2i(dx+c)} - B e^{2i(dx+c)} - 2iA - 4B)}{d(e^{2i(dx+c)}+1)^2(e^{2i(dx+c)}-1)} + \frac{7a^4 \ln(e^{2i(dx+c)}+1)B}{d} + \frac{4i}{d}$

input `int(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `4*a^4*((-I*A-B)*ln(sec(d*x+c)^2)+(I*A+1/4*B)*ln(tan(d*x+c))+1/8*B*tan(d*x+c)^2+(-I*B+1/4*A)*tan(d*x+c)-1/4*A*cot(d*x+c)+2*x*d*(-A+I*B))/d`

3.30.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 254 vs. 2(126) = 252.

Time = 0.27 (sec) , antiderivative size = 254, normalized size of antiderivative = 1.76

$$\int \cot^2(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= \frac{10Ba^4e^{(4idx+4ic)} - 2(2iA+B)a^4e^{(2idx+2ic)} - 4(iA+2B)a^4 + ((4iA+7B)a^4e^{(6idx+6ic)} + (4iA+7B))}{d}$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fricas")`

```
output (10*B*a^4*e^(4*I*d*x + 4*I*c) - 2*(2*I*A + B)*a^4*e^(2*I*d*x + 2*I*c) - 4*
(I*A + 2*B)*a^4 + ((4*I*A + 7*B)*a^4*e^(6*I*d*x + 6*I*c) + (4*I*A + 7*B)*a
^4*e^(4*I*d*x + 4*I*c) + (-4*I*A - 7*B)*a^4*e^(2*I*d*x + 2*I*c) + (-4*I*A
- 7*B)*a^4)*log(e^(2*I*d*x + 2*I*c) + 1) + ((4*I*A + B)*a^4*e^(6*I*d*x + 6
*I*c) + (4*I*A + B)*a^4*e^(4*I*d*x + 4*I*c) + (-4*I*A - B)*a^4*e^(2*I*d*x
+ 2*I*c) + (-4*I*A - B)*a^4)*log(e^(2*I*d*x + 2*I*c) - 1))/(d*e^(6*I*d*x +
6*I*c) + d*e^(4*I*d*x + 4*I*c) - d*e^(2*I*d*x + 2*I*c) - d)
```

3.30.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 260 vs. $2(124) = 248$.

Time = 1.30 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.81

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{ia^4 \cdot (4A - 7iB) \log\left(e^{2idx} + \frac{(-4iAa^4 - 4Ba^4 + ia^4 \cdot (4A - 7iB))e^{-2ic}}{3Ba^4}\right)}{d}$$

$$+ \frac{ia^4 \cdot (4A - iB) \log\left(e^{2idx} + \frac{(-4iAa^4 - 4Ba^4 + ia^4 \cdot (4A - iB))e^{-2ic}}{3Ba^4}\right)}{d}$$

$$+ \frac{-4iAa^4 + 10Ba^4 e^{4ic} e^{4idx} - 8Ba^4 + (-4iAa^4 e^{2ic} - 2Ba^4 e^{2ic}) e^{2idx}}{de^{6ic} e^{6idx} + de^{4ic} e^{4idx} - de^{2ic} e^{2idx} - d}$$

```
input integrate(cot(d*x+c)**2*(a+I*a*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)
```

```
output I*a**4*(4*A - 7*I*B)*log(exp(2*I*d*x) + (-4*I*A*a**4 - 4*B*a**4 + I*a**4*(
4*A - 7*I*B))*exp(-2*I*c)/(3*B*a**4))/d + I*a**4*(4*A - I*B)*log(exp(2*I*d
*x) + (-4*I*A*a**4 - 4*B*a**4 + I*a**4*(4*A - I*B))*exp(-2*I*c)/(3*B*a**4
))/d + (-4*I*A*a**4 + 10*B*a**4*exp(4*I*c)*exp(4*I*d*x) - 8*B*a**4 + (-4*I*
A*a**4*exp(2*I*c) - 2*B*a**4*exp(2*I*c))*exp(2*I*d*x))/(d*exp(6*I*c)*exp(6
*I*d*x) + d*exp(4*I*c)*exp(4*I*d*x) - d*exp(2*I*c)*exp(2*I*d*x) - d)
```


3.30.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.71

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{Ba^4 \tan(dx + c)^2 - 16(dx + c)(A - iB)a^4 - 8(iA + B)a^4 \log(\tan(dx + c)^2 + 1) + 2(4iA + B)a^4 \log(\tan(dx + c))}{2d}$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(B*a^4*tan(d*x + c)^2 - 16*(d*x + c)*(A - I*B)*a^4 - 8*(I*A + B)*a^4*log(tan(d*x + c)^2 + 1) + 2*(4*I*A + B)*a^4*log(tan(d*x + c)) + 2*(A - 4*I*B)*a^4*tan(d*x + c) - 2*A*a^4/tan(d*x + c))/d`

3.30.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 336 vs. $2(126) = 252$.

Time = 0.79 (sec) , antiderivative size = 336, normalized size of antiderivative = 2.33

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{Aa^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 2(4iAa^4 + 7Ba^4) \log\left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1\right) - 32(iAa^4 + Ba^4) \log\left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right)}{2d}$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output $\frac{1}{2}(Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 2(4IAa^4 + 7Ba^4) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) + 1) - 32(IAa^4 + Ba^4) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) + I) - 2(-4IAa^4 - 7Ba^4) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1) - 2(-4IAa^4 - Ba^4) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c)) - (8IAa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 2Ba^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) + Aa^4) / \tan(\frac{1}{2}dx + \frac{1}{2}c) - (12IAa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^4 + 21Ba^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^4 + 4Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 16IBa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 24IAa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 46Ba^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 4Aa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 16IBa^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 12IAa^4 + 21Ba^4) / (\tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 1)^2) / d$

3.30.9 Mupad [B] (verification not implemented)

Time = 8.38 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.76

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{B a^4 \tan(c + dx)^2}{2d} + \frac{a^4 \ln(\tan(c + dx)) (B + A 4i)}{d}$$

$$- \frac{8 a^4 \ln(\tan(c + dx) + 1i) (B + A 1i)}{d} - \frac{A a^4 \cot(c + dx)}{d}$$

$$- \frac{\tan(c + dx) (B a^4 1i + a^4 (3 B + A 1i) 1i)}{d}$$

input `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^4,x)`

output $(a^4 \log(\tan(c + d*x))(A*4i + B))/d - (\tan(c + d*x)*(B*a^4*1i + a^4*(A*1i + 3*B)*1i))/d - (8*a^4 \log(\tan(c + d*x) + 1i)*(A*1i + B))/d - (A*a^4 \cot(c + d*x))/d + (B*a^4 \tan(c + d*x)^2)/(2*d)$

3.31 $\int \cot^3(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.31.1	Optimal result	532
3.31.2	Mathematica [A] (verified)	533
3.31.3	Rubi [A] (verified)	533
3.31.4	Maple [A] (verified)	537
3.31.5	Fricas [A] (verification not implemented)	538
3.31.6	Sympy [A] (verification not implemented)	539
3.31.7	Maxima [A] (verification not implemented)	539
3.31.8	Giac [B] (verification not implemented)	540
3.31.9	Mupad [B] (verification not implemented)	540

3.31.1 Optimal result

Integrand size = 34, antiderivative size = 156

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= -8a^4(iA+B)x - \frac{a^4(A-4iB) \log(\cos(c+dx))}{d}$$

$$- \frac{a^4(7A-4iB) \log(\sin(c+dx))}{d} - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^3}{d}$$

$$- \frac{(5iA+2B) \cot(c+dx)(a^2+ia^2 \tan(c+dx))^2}{2d} - \frac{3A(a^4+ia^4 \tan(c+dx))}{d}$$

```
output -8*a^4*(I*A+B)*x-a^4*(A-4*I*B)*ln(cos(d*x+c))/d-a^4*(7*A-4*I*B)*ln(sin(d*x+c))/d-1/2*a*A*cot(d*x+c)^2*(a+I*a*tan(d*x+c))^3/d-1/2*(5*I*A+2*B)*cot(d*x+c)*(a^2+I*a^2*tan(d*x+c))^2/d-3*A*(a^4+I*a^4*tan(d*x+c))/d
```

3.31.2 Mathematica [A] (verified)

Time = 4.85 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.78

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= a^4 \left(-\frac{4iA \cot(c+dx)}{d} - \frac{B \cot(c+dx)}{d} - \frac{A \cot^2(c+dx)}{2d} - \frac{7A \log(\tan(c+dx))}{d} \right. \\ \left. + \frac{4iB \log(\tan(c+dx))}{d} + \frac{8A \log(i + \tan(c+dx))}{d} - \frac{8iB \log(i + \tan(c+dx))}{d} + \frac{B \tan(c+dx)}{d} \right)$$

input `Integrate[Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `a^4*(((4*I)*A*Cot[c + d*x])/d - (B*Cot[c + d*x])/d - (A*Cot[c + d*x]^2)/(2*d) - (7*A*Log[Tan[c + d*x]])/d + ((4*I)*B*Log[Tan[c + d*x]])/d + (8*A*Log[I + Tan[c + d*x]])/d - ((8*I)*B*Log[I + Tan[c + d*x]])/d + (B*Tan[c + d*x])/d)`

3.31.3 Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.441$, Rules used = {3042, 4076, 3042, 4076, 27, 3042, 4077, 3042, 4072, 3042, 3956, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a+ia \tan(c+dx))^4(A+B \tan(c+dx))}{\tan(c+dx)^3} dx$$

$$\downarrow \text{4076}$$

$$\frac{1}{2} \int \cot^2(c+dx)(i \tan(c+dx)a+a)^3(a(5iA+2B)+a(A+2iB) \tan(c+dx)) dx - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^3}{2d}$$

$$\begin{array}{c} \downarrow 3042 \\ \frac{1}{2} \int \frac{(i \tan(c+dx)a+a)^3(a(5iA+2B)+a(A+2iB)\tan(c+dx))}{\tan(c+dx)^2} dx - \\ \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^3}{2d} \\ \downarrow 4076 \end{array}$$

$$\begin{array}{c} \frac{1}{2} \left(\int -2 \cot(c+dx)(i \tan(c+dx)a+a)^2 (a^2(7A-4iB)-3ia^2A \tan(c+dx)) dx - \frac{(2B+5iA) \cot(c+dx)(a^2}{d} \right. \\ \left. \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^3}{2d} \right) \\ \downarrow 27 \end{array}$$

$$\begin{array}{c} \frac{1}{2} \left(-2 \int \cot(c+dx)(i \tan(c+dx)a+a)^2 (a^2(7A-4iB)-3ia^2A \tan(c+dx)) dx - \frac{(2B+5iA) \cot(c+dx)(a^2}{d} \right. \\ \left. \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^3}{2d} \right) \\ \downarrow 3042 \end{array}$$

$$\begin{array}{c} \frac{1}{2} \left(-2 \int \frac{(i \tan(c+dx)a+a)^2 (a^2(7A-4iB)-3ia^2A \tan(c+dx))}{\tan(c+dx)} dx - \frac{(2B+5iA) \cot(c+dx)(a^2+ia^2 \tan(c+dx))}{d} \right. \\ \left. \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^3}{2d} \right) \\ \downarrow 4077 \end{array}$$

$$\begin{array}{c} \frac{1}{2} \left(-2 \left(\int \cot(c+dx)(i \tan(c+dx)a+a) ((7A-4iB)a^3+(iA+4B)\tan(c+dx)a^3) dx + \frac{3A(a^4+ia^4 \tan(c+dx))}{d} \right) \right. \\ \left. \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^3}{2d} \right) \\ \downarrow 3042 \end{array}$$

$$\begin{array}{c} \frac{1}{2} \left(-2 \left(\int \frac{(i \tan(c+dx)a+a) ((7A-4iB)a^3+(iA+4B)\tan(c+dx)a^3)}{\tan(c+dx)} dx + \frac{3A(a^4+ia^4 \tan(c+dx))}{d} \right) \right. \\ \left. \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^3}{2d} \right) \\ \downarrow 4072 \end{array}$$

$$\frac{1}{2} \left(-2 \left(- \left(a^4(A - 4iB) \int \tan(c + dx) dx \right) + \int \cot(c + dx) \left((7A - 4iB)a^4 + 8(iA + B) \tan(c + dx)a^4 \right) dx + \frac{3A}{d} \right) \right. \\ \left. \frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^3}{2d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{2} \left(-2 \left(- \left(a^4(A - 4iB) \int \tan(c + dx) dx \right) + \int \frac{(7A - 4iB)a^4 + 8(iA + B) \tan(c + dx)a^4}{\tan(c + dx)} dx + \frac{3A(a^4 + ia^4 \tan(c + dx))}{d} \right) \right. \\ \left. \frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^3}{2d} \right) \\ \downarrow \text{3956}$$

$$\frac{1}{2} \left(-2 \left(\int \frac{(7A - 4iB)a^4 + 8(iA + B) \tan(c + dx)a^4}{\tan(c + dx)} dx + \frac{a^4(A - 4iB) \log(\cos(c + dx))}{d} + \frac{3A(a^4 + ia^4 \tan(c + dx))}{d} \right) \right. \\ \left. \frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^3}{2d} \right) \\ \downarrow \text{4014}$$

$$\frac{1}{2} \left(-2 \left(a^4(7A - 4iB) \int \cot(c + dx) dx + \frac{a^4(A - 4iB) \log(\cos(c + dx))}{d} + 8a^4x(B + iA) + \frac{3A(a^4 + ia^4 \tan(c + dx))}{d} \right) \right. \\ \left. \frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^3}{2d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{2} \left(-2 \left(a^4(7A - 4iB) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + \frac{a^4(A - 4iB) \log(\cos(c + dx))}{d} + 8a^4x(B + iA) + \frac{3A(a^4 + ia^4 \tan(c + dx))}{d} \right) \right. \\ \left. \frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^3}{2d} \right) \\ \downarrow \text{25}$$

$$\frac{1}{2} \left(-2 \left(-a^4(7A - 4iB) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + \frac{a^4(A - 4iB) \log(\cos(c + dx))}{d} + 8a^4x(B + iA) + \frac{3A(a^4 + ia^4 \tan(c + dx))}{d} \right) \right. \\ \left. \frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^3}{2d} \right) \\ \downarrow \text{3956}$$

$$\frac{1}{2} \left(-2 \left(\frac{a^4(7A - 4iB) \log(-\sin(c + dx))}{d} + \frac{a^4(A - 4iB) \log(\cos(c + dx))}{d} + 8a^4x(B + iA) + \frac{3A(a^4 + ia^4 \tan(c + dx))}{d} \right) + \frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^3}{2d} \right)$$

input `Int[Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `-1/2*(a*A*Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^3)/d + (-((((5*I)*A + 2*B)*Cot[c + d*x]*(a^2 + I*a^2*Tan[c + d*x])^2)/d) - 2*(8*a^4*(I*A + B)*x + (a^4*(A - (4*I)*B)*Log[Cos[c + d*x]])/d + (a^4*(7*A - (4*I)*B)*Log[-Sin[c + d*x]])/d + (3*A*(a^4 + I*a^4*Tan[c + d*x]))/d))/2`

3.31.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4072 `Int[(((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*(d/b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c - a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0]`

rule 4076 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 4077 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.31.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.67

$$3.31. \quad \int \cot^3(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

method	result
parallelrisc	$-\frac{a^4(16iAxd+A(\cot^2(dx+c))+8iA\cot(dx+c)-8iB\ln(\tan(dx+c))+8iB\ln(\sec^2(dx+c))+16Bdx+14A\ln(\tan(dx+c)))}{2d}$
derivativdivides	$a^4\left(-\frac{A(\cot^2(dx+c))}{2}-4iA\cot(dx+c)-\cot(dx+c)B+\frac{(-8iB+8A)\ln(\cot^2(dx+c)+1)}{2}+(8iA+8B)\left(\frac{\pi}{2}-\operatorname{arccot}(\cot(dx+c))\right)\right)$
default	$a^4\left(-\frac{A(\cot^2(dx+c))}{2}-4iA\cot(dx+c)-\cot(dx+c)B+\frac{(-8iB+8A)\ln(\cot^2(dx+c)+1)}{2}+(8iA+8B)\left(\frac{\pi}{2}-\operatorname{arccot}(\cot(dx+c))\right)\right)$
norman	$\frac{(-8iAa^4-8Ba^4)x(\tan^2(dx+c))+\frac{Ba^4(\tan^3(dx+c))}{d}-\frac{Aa^4}{2d}-\frac{(4iAa^4+Ba^4)\tan(dx+c)}{d}}{\tan(dx+c)^2}-\frac{(-4iBa^4+7Aa^4)\ln(\tan(dx+c))}{d}$
risc	$\frac{16a^4Bc}{d}+\frac{16ia^4Ac}{d}-\frac{2ia^4(5iAe^{4i(dx+c)}+iAe^{2i(dx+c)}+2Be^{2i(dx+c)}-4iA-2B)}{d(e^{2i(dx+c)}+1)(e^{2i(dx+c)}-1)^2}+\frac{4ia^4\ln(e^{2i(dx+c)}-1)B}{d}-\frac{7Aa^4}{d}$

input `int(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `-1/2*a^4*(16*I*A*x*d+A*cot(d*x+c)^2+8*I*A*cot(d*x+c)-8*I*B*ln(tan(d*x+c))+8*I*B*ln(sec(d*x+c)^2)+16*B*d*x+14*A*ln(tan(d*x+c))-8*A*ln(sec(d*x+c)^2)+2*cot(d*x+c)*B-2*B*tan(d*x+c))/d`

3.31.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.63

$$\int \cot^3(c+dx)(a+ia\tan(c+dx))^4(A+B\tan(c+dx))dx = \frac{10Aa^4e^{(4i dx+4i c)}+2(A-2iB)a^4e^{(2i dx+2i c)}-4(2A-iB)a^4-((A-4iB)a^4e^{(6i dx+6i c)}-(A-4iB)a^4}{d}$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `(10*A*a^4*e^(4*I*d*x+4*I*c)+2*(A-2*I*B)*a^4*e^(2*I*d*x+2*I*c)-4*(2*A-I*B)*a^4-((A-4*I*B)*a^4*e^(6*I*d*x+6*I*c)-(A-4*I*B)*a^4*e^(4*I*d*x+4*I*c)-(A-4*I*B)*a^4*e^(2*I*d*x+2*I*c)+(A-4*I*B)*a^4)*log(e^(2*I*d*x+2*I*c)+1)-((7*A-4*I*B)*a^4*e^(6*I*d*x+6*I*c)-(7*A-4*I*B)*a^4*e^(4*I*d*x+4*I*c)-(7*A-4*I*B)*a^4*e^(2*I*d*x+2*I*c)+(7*A-4*I*B)*a^4)*log(e^(2*I*d*x+2*I*c)-1)/(d*e^(6*I*d*x+6*I*c)-d*e^(4*I*d*x+4*I*c)-d*e^(2*I*d*x+2*I*c)+d)`

3.31. $\int \cot^3(c+dx)(a+ia\tan(c+dx))^4(A+B\tan(c+dx))dx$

3.31.6 Sympy [A] (verification not implemented)

Time = 1.18 (sec) , antiderivative size = 252, normalized size of antiderivative = 1.62

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= -\frac{a^4(A - 4iB) \log\left(e^{2idx} + \frac{(4Aa^4 - 4iBa^4 - a^4(A - 4iB))e^{-2ic}}{3Aa^4}\right)}{d}$$

$$- \frac{a^4 \cdot (7A - 4iB) \log\left(e^{2idx} + \frac{(4Aa^4 - 4iBa^4 - a^4(7A - 4iB))e^{-2ic}}{3Aa^4}\right)}{d}$$

$$+ \frac{10Aa^4 e^{4ic} e^{4idx} - 8Aa^4 + 4iBa^4 + (2Aa^4 e^{2ic} - 4iBa^4 e^{2ic}) e^{2idx}}{de^{6ic} e^{6idx} - de^{4ic} e^{4idx} - de^{2ic} e^{2idx} + d}$$

input `integrate(cot(d*x+c)**3*(a+I*a*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)`

output `-a**4*(A - 4*I*B)*log(exp(2*I*d*x) + (4*A*a**4 - 4*I*B*a**4 - a**4*(A - 4*I*B))*exp(-2*I*c)/(3*A*a**4))/d - a**4*(7*A - 4*I*B)*log(exp(2*I*d*x) + (4*A*a**4 - 4*I*B*a**4 - a**4*(7*A - 4*I*B))*exp(-2*I*c)/(3*A*a**4))/d + (10*A*a**4*exp(4*I*c)*exp(4*I*d*x) - 8*A*a**4 + 4*I*B*a**4 + (2*A*a**4*exp(2*I*c) - 4*I*B*a**4*exp(2*I*c))*exp(2*I*d*x))/(d*exp(6*I*c)*exp(6*I*d*x) - d*exp(4*I*c)*exp(4*I*d*x) - d*exp(2*I*c)*exp(2*I*d*x) + d)`

3.31.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.69

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$\frac{16(dx + c)(iA + B)a^4 - 8(A - iB)a^4 \log(\tan(dx + c)^2 + 1) + 2(7A - 4iB)a^4 \log(\tan(dx + c)) -}{2d}$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(16*(d*x + c)*(I*A + B)*a^4 - 8*(A - I*B)*a^4*log(tan(d*x + c)^2 + 1) + 2*(7*A - 4*I*B)*a^4*log(tan(d*x + c)) - 2*B*a^4*tan(d*x + c) - (2*(-4*I*A - B)*a^4*tan(d*x + c) - A*a^4)/tan(d*x + c)^2)/d`

3.31.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 317 vs. $2(138) = 276$.

Time = 0.90 (sec) , antiderivative size = 317, normalized size of antiderivative = 2.03

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$\frac{Aa^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 16i Aa^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 4Ba^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 8(Aa^4 - 4i Ba^4) \log(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right))}{d}$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output
$$\frac{-1/8*(A*a^4*\tan(1/2*d*x + 1/2*c)^2 - 16*I*A*a^4*\tan(1/2*d*x + 1/2*c) - 4*B*a^4*\tan(1/2*d*x + 1/2*c) + 8*(A*a^4 - 4*I*B*a^4)*\log(\tan(1/2*d*x + 1/2*c) + 1) - 128*(A*a^4 - I*B*a^4)*\log(\tan(1/2*d*x + 1/2*c) + I) + 8*(A*a^4 - 4*I*B*a^4)*\log(\tan(1/2*d*x + 1/2*c) - 1) + 8*(7*A*a^4 - 4*I*B*a^4)*\log(\tan(1/2*d*x + 1/2*c)) - 8*(A*a^4*\tan(1/2*d*x + 1/2*c)^2 - 4*I*B*a^4*\tan(1/2*d*x + 1/2*c)^2 - 2*B*a^4*\tan(1/2*d*x + 1/2*c) - A*a^4 + 4*I*B*a^4)/(\tan(1/2*d*x + 1/2*c)^2 - 1) - (84*A*a^4*\tan(1/2*d*x + 1/2*c)^2 - 48*I*B*a^4*\tan(1/2*d*x + 1/2*c)^2 - 16*I*A*a^4*\tan(1/2*d*x + 1/2*c) - 4*B*a^4*\tan(1/2*d*x + 1/2*c) - A*a^4)/\tan(1/2*d*x + 1/2*c)^2}{d}$$

3.31.9 Mupad [B] (verification not implemented)

Time = 7.67 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.65

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{B a^4 \tan(c + dx)}{d} - \frac{a^4 \ln(\tan(c + dx)) (7 A - B 4i)}{d}$$

$$+ \frac{8 a^4 \ln(\tan(c + dx) + 1i) (A - B 1i)}{d}$$

$$- \frac{\cot(c + dx)^2 \left(\frac{A a^4}{2} + \tan(c + dx) (B a^4 + A a^4 4i) \right)}{d}$$

input `int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^4,x)`

3.31. $\int \cot^3(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$

output $(8a^4 \log(\tan(c + dx) + 1i)(A - B1i))/d - (a^4 \log(\tan(c + dx))(7A - B4i))/d - (\cot(c + dx)^2((Aa^4)/2 + \tan(c + dx)(Aa^44i + Ba^4)))/d + (Ba^4 \tan(c + dx))/d$

3.32 $\int \cot^4(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.32.1	Optimal result	542
3.32.2	Mathematica [A] (verified)	543
3.32.3	Rubi [A] (verified)	543
3.32.4	Maple [A] (verified)	547
3.32.5	Fricas [A] (verification not implemented)	548
3.32.6	Sympy [B] (verification not implemented)	548
3.32.7	Maxima [A] (verification not implemented)	549
3.32.8	Giac [A] (verification not implemented)	549
3.32.9	Mupad [B] (verification not implemented)	550

3.32.1 Optimal result

Integrand size = 34, antiderivative size = 163

$$\int \cot^4(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= 8a^4(A-iB)x - \frac{a^4B \log(\cos(c+dx))}{d} - \frac{a^4(8iA+7B) \log(\sin(c+dx))}{d}$$

$$- \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^3}{3d}$$

$$- \frac{(2iA+B) \cot^2(c+dx)(a^2+ia^2 \tan(c+dx))^2}{2d}$$

$$+ \frac{(4A-3iB) \cot(c+dx)(a^4+ia^4 \tan(c+dx))}{d}$$

output

```
8*a^4*(A-I*B)*x-a^4*B*ln(cos(d*x+c))/d-a^4*(8*I*A+7*B)*ln(sin(d*x+c))/d-1/3*a*A*cot(d*x+c)^3*(a+I*a*tan(d*x+c))^3/d-1/2*(2*I*A+B)*cot(d*x+c)^2*(a^2+I*a^2*tan(d*x+c))^2/d+(4*A-3*I*B)*cot(d*x+c)*(a^4+I*a^4*tan(d*x+c))/d
```

3.32.2 Mathematica [A] (verified)

Time = 1.15 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.88

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= a^4 \left(\frac{7A \cot(c + dx)}{d} - \frac{4iB \cot(c + dx)}{d} - \frac{2iA \cot^2(c + dx)}{d} - \frac{B \cot^2(c + dx)}{2d} \right. \\ \left. - \frac{A \cot^3(c + dx)}{3d} - \frac{8iA \log(\tan(c + dx))}{d} - \frac{7B \log(\tan(c + dx))}{d} \right. \\ \left. + \frac{8iA \log(i + \tan(c + dx))}{d} + \frac{8B \log(i + \tan(c + dx))}{d} \right)$$

input `Integrate[Cot[c + d*x]^4*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `a^4*((7*A*Cot[c + d*x])/d - ((4*I)*B*Cot[c + d*x])/d - ((2*I)*A*Cot[c + d*x]^2)/d - (B*Cot[c + d*x]^2)/(2*d) - (A*Cot[c + d*x]^3)/(3*d) - ((8*I)*A*Log[Tan[c + d*x]])/d - (7*B*Log[Tan[c + d*x]])/d + ((8*I)*A*Log[I + Tan[c + d*x]])/d + (8*B*Log[I + Tan[c + d*x]])/d)`

3.32.3 Rubi [A] (verified)

Time = 1.20 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.01, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.471$, Rules used = {3042, 4076, 27, 3042, 4076, 27, 3042, 4076, 3042, 4072, 3042, 3956, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + ia \tan(c + dx))^4(A + B \tan(c + dx))}{\tan(c + dx)^4} dx$$

$$\downarrow \text{4076}$$

$$\frac{1}{3} \int 3 \cot^3(c + dx)(i \tan(c + dx)a + a)^3(a(2iA + B) + iaB \tan(c + dx)) dx - \frac{aA \cot^3(c + dx)(a + ia \tan(c + dx))^3}{3d}$$

$$\begin{aligned}
& \int \cot^3(c+dx)(i \tan(c+dx)a+a)^3(a(2iA+B)+iaB \tan(c+dx))dx - \\
& \quad \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 27 \\
& \int \frac{(i \tan(c+dx)a+a)^3(a(2iA+B)+iaB \tan(c+dx))}{\tan(c+dx)^3} dx - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 3042 \\
& \frac{1}{2} \int -2 \cot^2(c+dx)(i \tan(c+dx)a+a)^2((4A-3iB)a^2+B \tan(c+dx)a^2) dx - \\
& \quad \frac{(B+2iA) \cot^2(c+dx)(a^2+ia^2 \tan(c+dx))^2}{2d} - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 4076 \\
& - \int \cot^2(c+dx)(i \tan(c+dx)a+a)^2((4A-3iB)a^2+B \tan(c+dx)a^2) dx - \\
& \quad \frac{(B+2iA) \cot^2(c+dx)(a^2+ia^2 \tan(c+dx))^2}{2d} - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 27 \\
& - \int \cot^2(c+dx)(i \tan(c+dx)a+a)^2((4A-3iB)a^2+B \tan(c+dx)a^2) dx - \\
& \quad \frac{(B+2iA) \cot^2(c+dx)(a^2+ia^2 \tan(c+dx))^2}{2d} - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 3042 \\
& - \int \frac{(i \tan(c+dx)a+a)^2((4A-3iB)a^2+B \tan(c+dx)a^2)}{\tan(c+dx)^2} dx - \\
& \quad \frac{(B+2iA) \cot^2(c+dx)(a^2+ia^2 \tan(c+dx))^2}{2d} - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 4076 \\
& - \int \cot(c+dx)(i \tan(c+dx)a+a)((8iA+7B)a^3+iB \tan(c+dx)a^3) dx + \\
& \quad \frac{(4A-3iB) \cot(c+dx)(a^4+ia^4 \tan(c+dx))}{d} - \frac{(B+2iA) \cot^2(c+dx)(a^2+ia^2 \tan(c+dx))^2}{2d} - \\
& \quad \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 3042 \\
& - \int \frac{(i \tan(c+dx)a+a)((8iA+7B)a^3+iB \tan(c+dx)a^3)}{\tan(c+dx)} dx + \\
& \quad \frac{(4A-3iB) \cot(c+dx)(a^4+ia^4 \tan(c+dx))}{d} - \frac{(B+2iA) \cot^2(c+dx)(a^2+ia^2 \tan(c+dx))^2}{2d} - \\
& \quad \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^3}{3d} \\
& \quad \downarrow 4072
\end{aligned}$$

3.32. $\int \cot^4(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& - \int \cot(c+dx) (a^4(8iA+7B) - 8a^4(A-iB)\tan(c+dx)) dx + a^4B \int \tan(c+dx) dx + \\
& \frac{(4A-3iB)\cot(c+dx)(a^4+ia^4\tan(c+dx))}{d} - \frac{(B+2iA)\cot^2(c+dx)(a^2+ia^2\tan(c+dx))^2}{2d} - \\
& \frac{aA\cot^3(c+dx)(a+ia\tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{3042} \\
& - \int \frac{a^4(8iA+7B) - 8a^4(A-iB)\tan(c+dx)}{\tan(c+dx)} dx + a^4B \int \tan(c+dx) dx + \\
& \frac{(4A-3iB)\cot(c+dx)(a^4+ia^4\tan(c+dx))}{d} - \frac{(B+2iA)\cot^2(c+dx)(a^2+ia^2\tan(c+dx))^2}{2d} - \\
& \frac{aA\cot^3(c+dx)(a+ia\tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{3956} \\
& - \int \frac{a^4(8iA+7B) - 8a^4(A-iB)\tan(c+dx)}{\tan(c+dx)} dx + \\
& \frac{(4A-3iB)\cot(c+dx)(a^4+ia^4\tan(c+dx))}{d} - \frac{a^4B\log(\cos(c+dx))}{d} - \\
& \frac{(B+2iA)\cot^2(c+dx)(a^2+ia^2\tan(c+dx))^2}{2d} - \frac{aA\cot^3(c+dx)(a+ia\tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{4014} \\
& -a^4(7B+8iA) \int \cot(c+dx) dx + \frac{(4A-3iB)\cot(c+dx)(a^4+ia^4\tan(c+dx))}{d} + 8a^4x(A-iB) - \\
& \frac{a^4B\log(\cos(c+dx))}{d} - \frac{(B+2iA)\cot^2(c+dx)(a^2+ia^2\tan(c+dx))^2}{2d} - \\
& \frac{aA\cot^3(c+dx)(a+ia\tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{3042} \\
& -a^4(7B+8iA) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx + \frac{(4A-3iB)\cot(c+dx)(a^4+ia^4\tan(c+dx))}{d} + \\
& 8a^4x(A-iB) - \frac{a^4B\log(\cos(c+dx))}{d} - \frac{(B+2iA)\cot^2(c+dx)(a^2+ia^2\tan(c+dx))^2}{2d} - \\
& \frac{aA\cot^3(c+dx)(a+ia\tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{25} \\
& a^4(7B+8iA) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx + \frac{(4A-3iB)\cot(c+dx)(a^4+ia^4\tan(c+dx))}{d} + \\
& 8a^4x(A-iB) - \frac{a^4B\log(\cos(c+dx))}{d} - \frac{(B+2iA)\cot^2(c+dx)(a^2+ia^2\tan(c+dx))^2}{2d} - \\
& \frac{aA\cot^3(c+dx)(a+ia\tan(c+dx))^3}{3d}
\end{aligned}$$

3.32. $\int \cot^4(c+dx)(a+ia\tan(c+dx))^4(A+B\tan(c+dx)) dx$

$$\begin{aligned} & \downarrow \text{3956} \\ & -\frac{a^4(7B + 8iA) \log(-\sin(c + dx))}{d} + \frac{(4A - 3iB) \cot(c + dx) (a^4 + ia^4 \tan(c + dx))}{d} + 8a^4x(A - \\ & iB) - \frac{a^4B \log(\cos(c + dx))}{d} - \frac{(B + 2iA) \cot^2(c + dx) (a^2 + ia^2 \tan(c + dx))^2}{3d} - \\ & \frac{aA \cot^3(c + dx)(a + ia \tan(c + dx))^3}{3d} \end{aligned}$$

input `Int[Cot[c + d*x]^4*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `8*a^4*(A - I*B)*x - (a^4*B*Log[Cos[c + d*x]])/d - (a^4*((8*I)*A + 7*B)*Log[-Sin[c + d*x]])/d - (a*A*Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^3)/(3*d) - (((2*I)*A + B)*Cot[c + d*x]^2*(a^2 + I*a^2*Tan[c + d*x])^2)/(2*d) + ((4*A - (3*I)*B)*Cot[c + d*x]*(a^4 + I*a^4*Tan[c + d*x]))/d`

3.32.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4072 `Int[(((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*(d/b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c - a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0]`

rule 4076 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.32.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.71

method	result
derivativedivides	$\frac{a^4 \left(-2iA(\cot^2(dx+c)) - \frac{A(\cot^3(dx+c))}{3} - 4iB \cot(dx+c) - \frac{B(\cot^2(dx+c))}{2} + 7A \cot(dx+c) + \frac{(8iA+8B) \ln(\cot^2(dx+c)+1)}{2} \right)}{d}$
default	$\frac{a^4 \left(-2iA(\cot^2(dx+c)) - \frac{A(\cot^3(dx+c))}{3} - 4iB \cot(dx+c) - \frac{B(\cot^2(dx+c))}{2} + 7A \cot(dx+c) + \frac{(8iA+8B) \ln(\cot^2(dx+c)+1)}{2} \right)}{d}$
parallelrisch	$-\frac{a^4(2A(\cot^3(dx+c)) + 12iA(\cot^2(dx+c)) + 48iBdx + 48iA \ln(\tan(dx+c)) - 24iA \ln(\sec^2(dx+c)) - 48Adx + 3B(\cot^2(dx+c)))}{6d}$
norman	$\frac{\left(\frac{-4iB a^4 + 7A a^4}{d} \right) (\tan^2(dx+c)) + (-8iB a^4 + 8A a^4) x (\tan^3(dx+c)) - \frac{A a^4}{3d} - \frac{(4iA a^4 + B a^4) \tan(dx+c)}{2d}}{\tan(dx+c)^3} + \frac{4(iA a^4 + B a^4) \ln(\tan(dx+c))}{d}$
risch	$\frac{16ia^4Bc}{d} - \frac{16a^4Ac}{d} + \frac{2a^4(36iA e^{4i(dx+c)} + 15B e^{4i(dx+c)} - 54iA e^{2i(dx+c)} - 27B e^{2i(dx+c)} + 22iA + 12B)}{3d(e^{2i(dx+c)} - 1)^3} - \frac{7a^4 \ln(\tan(dx+c))}{d}$

input `int(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `a^4/d*(-2*I*A*cot(d*x+c)^2-1/3*A*cot(d*x+c)^3-4*I*B*cot(d*x+c)-1/2*B*cot(d*x+c)^2+7*A*cot(d*x+c)+1/2*(8*B+8*I*A)*ln(cot(d*x+c)^2+1)+(-8*A+8*I*B)*(1/2*Pi-arccot(cot(d*x+c)))-B*ln(cot(d*x+c)))`

3.32. $\int \cot^4(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$

3.32.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.53

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$\frac{6(-12iA - 5B)a^4 e^{(4i dx + 4i c)} + 54(2iA + B)a^4 e^{(2i dx + 2i c)} + 4(-11iA - 6B)a^4 + 3(Ba^4 e^{(6i dx + 6i c)} -$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output
$$\frac{-1/3*(6*(-12*I*A - 5*B))*a^4*e^{(4*I*d*x + 4*I*c)} + 54*(2*I*A + B)*a^4*e^{(2*I*d*x + 2*I*c)} + 4*(-11*I*A - 6*B)*a^4 + 3*(B*a^4*e^{(6*I*d*x + 6*I*c)} - 3*B*a^4*e^{(4*I*d*x + 4*I*c)} + 3*B*a^4*e^{(2*I*d*x + 2*I*c)} - B*a^4)*\log(e^{(2*I*d*x + 2*I*c)} + 1) + 3*((8*I*A + 7*B)*a^4*e^{(6*I*d*x + 6*I*c)} + 3*(-8*I*A - 7*B)*a^4*e^{(4*I*d*x + 4*I*c)} + 3*(8*I*A + 7*B)*a^4*e^{(2*I*d*x + 2*I*c)} + (-8*I*A - 7*B)*a^4)*\log(e^{(2*I*d*x + 2*I*c)} - 1))/(d*e^{(6*I*d*x + 6*I*c)} - 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} - d)$$

3.32.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 292 vs. 2(143) = 286.

Time = 2.47 (sec) , antiderivative size = 292, normalized size of antiderivative = 1.79

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= -\frac{Ba^4 \log\left(\frac{4Aa^4 - 3iBa^4}{4Aa^4 e^{2ic} - 3iBa^4 e^{2ic}} + e^{2idx}\right)}{d}$$

$$- \frac{ia^4 \cdot (8A - 7iB) \log\left(e^{2idx} + \frac{4Aa^4 - 4iBa^4 - a^4 \cdot (8A - 7iB)}{4Aa^4 e^{2ic} - 3iBa^4 e^{2ic}}\right)}{d}$$

$$+ \frac{44iAa^4 + 24Ba^4 + (-108iAa^4 e^{2ic} - 54Ba^4 e^{2ic}) e^{2idx} + (72iAa^4 e^{4ic} + 30Ba^4 e^{4ic}) e^{4idx}}{3de^{6ic} e^{6idx} - 9de^{4ic} e^{4idx} + 9de^{2ic} e^{2idx} - 3d}$$

input `integrate(cot(d*x+c)**4*(a+I*a*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)`

output
$$\begin{aligned} & -B a^{4} \log \left(\frac{4 A a^{4} - 3 I B a^{4}}{4 A a^{4} \exp(2 I c) - 3 I B a^{4} \exp(2 I c)} + \exp(2 I d x) \right) / d - I a^{4} (8 A - 7 I B) \log \left(\exp(2 I d x) + \frac{4 A a^{4} - 4 I B a^{4} - a^{4} (8 A - 7 I B)}{4 A a^{4} \exp(2 I c) - 3 I B a^{4} \exp(2 I c)} \right) / d \\ & + \frac{(44 I A a^{4} + 24 B a^{4} + (-108 I A a^{4} \exp(2 I c) - 54 B a^{4} \exp(2 I c)) \exp(2 I d x) + (72 I A a^{4} \exp(4 I c) + 30 B a^{4} \exp(4 I c)) \exp(4 I d x))}{(3 d \exp(6 I c) \exp(6 I d x) - 9 d \exp(4 I c) \exp(4 I d x) + 9 d \exp(2 I c) \exp(2 I d x) - 3 d)} \end{aligned}$$

3.32.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.72

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{48(dx + c)(A - iB)a^4 - 24(-iA - B)a^4 \log(\tan(dx + c)^2 + 1) + 6(-8iA - 7B)a^4 \log(\tan(dx + c))}{6d}$$

input `integrate(cot(dx+c)^4*(a+I*a*tan(dx+c))^4*(A+B*tan(dx+c)),x, algorithm="maxima")`

output
$$\frac{1}{6} \frac{(48(dx + c)(A - I B) a^4 - 24(-I A - B) a^4 \log(\tan(dx + c)^2 + 1) + 6(-8 I A - 7 B) a^4 \log(\tan(dx + c)) + (6(7 A - 4 I B) a^4 \tan(dx + c)^2 + 3(-4 I A - B) a^4 \tan(dx + c) - 2 A a^4) / \tan(dx + c)^3) / d}{6d}$$

3.32.8 Giac [A] (verification not implemented)

Time = 0.98 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.79

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{A a^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 12 i A a^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 3 B a^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 24 B a^4 \log\left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right)}{6d}$$

input `integrate(cot(dx+c)^4*(a+I*a*tan(dx+c))^4*(A+B*tan(dx+c)),x, algorithm="giac")`

output $\frac{1}{24}(A^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 12IA^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 3B^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 24B^4 \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) + 1) - 24B^4 \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) - 1) - 87A^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 48IB^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) - 384(-IA^4 - B^4) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c) + I) - 24(8IA^4 + 7B^4) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c)) - (-352IA^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 308B^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 87A^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^2 + 48IB^4 \tan(\frac{1}{2}dx + \frac{1}{2}c)^2 + 12IA^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 3B^4 \tan(\frac{1}{2}dx + \frac{1}{2}c) + A^4) / \tan(\frac{1}{2}dx + \frac{1}{2}c)^3) / d$

3.32.9 Mupad [B] (verification not implemented)

Time = 7.62 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.69

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= -\frac{\frac{Aa^4}{3} - \tan(c + dx)^2(7Aa^4 - Ba^4 4i) + \tan(c + dx) \left(\frac{Ba^4}{2} + Aa^4 2i\right)}{d \tan(c + dx)^3} - \frac{a^4 \ln(\tan(c + dx))(7B + A8i)}{d} + \frac{8a^4 \ln(\tan(c + dx) + 1i)(B + A1i)}{d}$$

input `int(cot(c + d*x)^4*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^4,x)`

output $(8a^4 \log(\tan(c + d*x) + 1i)(A1i + B))/d - (a^4 \log(\tan(c + d*x))(A8i + 7B))/d - ((Aa^4)/3 - \tan(c + d*x)^2(7Aa^4 - Ba^4 4i) + \tan(c + d*x)(Aa^4 2i + (Ba^4)/2)) / (d \tan(c + d*x)^3)$

3.33 $\int \cot^5(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.33.1	Optimal result	551
3.33.2	Mathematica [A] (verified)	552
3.33.3	Rubi [A] (verified)	552
3.33.4	Maple [A] (verified)	556
3.33.5	Fricas [A] (verification not implemented)	557
3.33.6	Sympy [A] (verification not implemented)	557
3.33.7	Maxima [A] (verification not implemented)	558
3.33.8	Giac [B] (verification not implemented)	558
3.33.9	Mupad [B] (verification not implemented)	559

3.33.1 Optimal result

Integrand size = 34, antiderivative size = 177

$$\begin{aligned} & \int \cot^5(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx \\ &= 8a^4(iA+B)x + \frac{a^4(67iA+64B) \cot(c+dx)}{12d} \\ & \quad + \frac{8a^4(A-iB) \log(\sin(c+dx))}{d} - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^3}{4d} \\ & \quad - \frac{(7iA+4B) \cot^3(c+dx)(a^2+ia^2 \tan(c+dx))^2}{12d} \\ & \quad + \frac{(19A-16iB) \cot^2(c+dx)(a^4+ia^4 \tan(c+dx))}{12d} \end{aligned}$$

output

```
8*a^4*(I*A+B)*x+1/12*a^4*(67*I*A+64*B)*cot(d*x+c)/d+8*a^4*(A-I*B)*ln(sin(d*x+c))/d-1/4*a*A*cot(d*x+c)^4*(a+I*a*tan(d*x+c))^3/d-1/12*(7*I*A+4*B)*cot(d*x+c)^3*(a^2+I*a^2*tan(d*x+c))^2/d+1/12*(19*A-16*I*B)*cot(d*x+c)^2*(a^4+I*a^4*tan(d*x+c))/d
```

3.33.2 Mathematica [A] (verified)

Time = 1.11 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.51

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{a^4(-3A(i + \cot(c + dx))^4 + 4(A - iB)(21i \cot(c + dx) + 6 \cot^2(c + dx) - i \cot^3(c + dx) + 24(\log(\tan(c + dx))))}{12d}$$

input `Integrate[Cot[c + d*x]^5*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(a^4*(-3*A*(I + Cot[c + d*x])^4 + 4*(A - I*B)*((21*I)*Cot[c + d*x] + 6*Cot[c + d*x]^2 - I*Cot[c + d*x]^3 + 24*(Log[Tan[c + d*x]] - Log[I + Tan[c + d*x]])))/(12*d)`

3.33.3 Rubi [A] (verified)

Time = 1.25 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.09, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.441$, Rules used = {3042, 4076, 3042, 4076, 27, 3042, 4076, 3042, 4074, 27, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a + ia \tan(c + dx))^4(A + B \tan(c + dx))}{\tan(c + dx)^5} dx$$

$$\downarrow 4076$$

$$\frac{1}{4} \int \cot^4(c + dx)(i \tan(c + dx)a + a)^3(a(7iA + 4B) - a(A - 4iB) \tan(c + dx)) dx - \frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^3}{4d}$$

$$\downarrow 3042$$

$$\frac{1}{4} \int \frac{(i \tan(c + dx)a + a)^3(a(7iA + 4B) - a(A - 4iB) \tan(c + dx))}{\tan(c + dx)^4} dx - \frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^3}{4d}$$

3.33. $\int \cot^5(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$

↓ 4076

$$\frac{1}{4} \left(\frac{1}{3} \int -2 \cot^3(c+dx)(i \tan(c+dx)a+a)^2 ((19A-16iB)a^2+(5iA+8B)\tan(c+dx)a^2) dx - \frac{(4B+7iA)\cot^3(c+dx)}{3d} \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^3}{4d}$$

↓ 27

$$\frac{1}{4} \left(-\frac{2}{3} \int \cot^3(c+dx)(i \tan(c+dx)a+a)^2 ((19A-16iB)a^2+(5iA+8B)\tan(c+dx)a^2) dx - \frac{(4B+7iA)\cot^3(c+dx)}{3d} \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^3}{4d}$$

↓ 3042

$$\frac{1}{4} \left(-\frac{2}{3} \int \frac{(i \tan(c+dx)a+a)^2 ((19A-16iB)a^2+(5iA+8B)\tan(c+dx)a^2)}{\tan(c+dx)^3} dx - \frac{(4B+7iA)\cot^3(c+dx)(a^2)}{3d} \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^3}{4d}$$

↓ 4076

$$\frac{1}{4} \left(-\frac{2}{3} \left(\frac{1}{2} \int \cot^2(c+dx)(i \tan(c+dx)a+a)(a^3(67iA+64B)-a^3(29A-32iB)\tan(c+dx)) dx - \frac{(19A-16iB)\cot^2(c+dx)}{2d} \right) \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^3}{4d}$$

↓ 3042

$$\frac{1}{4} \left(-\frac{2}{3} \left(\frac{1}{2} \int \frac{(i \tan(c+dx)a+a)(a^3(67iA+64B)-a^3(29A-32iB)\tan(c+dx))}{\tan(c+dx)^2} dx - \frac{(19A-16iB)\cot^2(c+dx)}{2d} \right) \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^3}{4d}$$

↓ 4074

$$\frac{1}{4} \left(-\frac{2}{3} \left(\frac{1}{2} \left(\int -96 \cot(c+dx)((A-iB)a^4+(iA+B)\tan(c+dx)a^4) dx - \frac{a^4(64B+67iA)\cot(c+dx)}{d} \right) \right) \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^3}{4d}$$

↓ 27

3.33. $\int \cot^5(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

$$\frac{1}{4} \left(-\frac{2}{3} \left(\frac{1}{2} \left(-96 \int \cot(c+dx) ((A-iB)a^4 + (iA+B)\tan(c+dx)a^4) dx - \frac{a^4(64B+67iA)\cot(c+dx)}{d} \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^3}{4d} \right) \right)$$

↓ 3042

$$\frac{1}{4} \left(-\frac{2}{3} \left(\frac{1}{2} \left(-96 \int \frac{(A-iB)a^4 + (iA+B)\tan(c+dx)a^4}{\tan(c+dx)} dx - \frac{a^4(64B+67iA)\cot(c+dx)}{d} \right) - \frac{(19A-16iB)\cot(c+dx)}{d} \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^3}{4d} \right)$$

↓ 4014

$$\frac{1}{4} \left(-\frac{2}{3} \left(\frac{1}{2} \left(-96 \left(a^4(A-iB) \int \cot(c+dx) dx + a^4x(B+iA) \right) - \frac{a^4(64B+67iA)\cot(c+dx)}{d} \right) - \frac{(19A-16iB)\cot(c+dx)}{d} \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^3}{4d} \right)$$

↓ 3042

$$\frac{1}{4} \left(-\frac{2}{3} \left(\frac{1}{2} \left(-96 \left(a^4(A-iB) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx + a^4x(B+iA) \right) - \frac{a^4(64B+67iA)\cot(c+dx)}{d} \right) - \frac{(19A-16iB)\cot(c+dx)}{d} \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^3}{4d} \right)$$

↓ 25

$$\frac{1}{4} \left(-\frac{2}{3} \left(\frac{1}{2} \left(-96 \left(a^4x(B+iA) - a^4(A-iB) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx \right) - \frac{a^4(64B+67iA)\cot(c+dx)}{d} \right) - \frac{(19A-16iB)\cot(c+dx)}{d} \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^3}{4d} \right)$$

↓ 3956

$$\frac{1}{4} \left(-\frac{2}{3} \left(\frac{1}{2} \left(-\frac{a^4(64B+67iA)\cot(c+dx)}{d} - 96 \left(\frac{a^4(A-iB)\log(-\sin(c+dx))}{d} + a^4x(B+iA) \right) \right) - \frac{(19A-16iB)\cot(c+dx)}{d} \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^3}{4d} \right)$$

input `Int[Cot[c + d*x]^5*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output
$$-1/4*(a*A*\cot[c + d*x]^4*(a + I*a*\tan[c + d*x])^3)/d + (-1/3*(((7*I)*A + 4*B)*\cot[c + d*x]^3*(a^2 + I*a^2*\tan[c + d*x])^2)/d - (2*((-(a^4*((67*I)*A + 64*B)*\cot[c + d*x])/d) - 96*(a^4*(I*A + B)*x + (a^4*(A - I*B)*\log[-\sin[c + d*x]]))/d)/2 - ((19*A - (16*I)*B)*\cot[c + d*x]^2*(a^4 + I*a^4*\tan[c + d*x]))/(2*d))/3)/4$$

3.33.3.1 Defintions of rubi rules used

rule 25 $\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{Int}[F_x, x], x]$

rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x) /; \text{FreeQ}[b, x]]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3956 $\text{Int}[\tan[(c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\log[\text{RemoveContent}[\cos[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

rule 4014 $\text{Int}[((c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_)])/(a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(a*c + b*d)*(x/(a^2 + b^2)), x] + \text{Simp}[(b*c - a*d)/(a^2 + b^2) \text{Int}[(b - a*\tan[e + f*x])/(a + b*\tan[e + f*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[a*c + b*d, 0]$

rule 4074 $\text{Int}[(a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_)]^{(m_)}*((A_.) + (B_.)*\tan[(e_.) + (f_.)*(x_)]*(c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(b*c - a*d)*(A*b - a*B)*((a + b*\tan[e + f*x])^{(m + 1)})/(b*f*(m + 1)*(a^2 + b^2)), x] + \text{Simp}[1/(a^2 + b^2) \text{Int}[(a + b*\tan[e + f*x])^{(m + 1)}*\text{Simp}[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*\tan[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$

```
rule 4076 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m -
n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

3.33.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.63

method	result
parallelrisch	$\frac{8a^4 \left(\left(-\frac{A}{2} + \frac{iB}{2} \right) \ln(\sec^2(dx+c)) + (-iB+A) \ln(\tan(dx+c)) - \frac{A(\cot^4(dx+c))}{32} + (\cot^3(dx+c)) \left(-\frac{iA}{6} - \frac{B}{24} \right) + (\cot^2(dx+c)) \right)}{d}$
derivativedivides	$\frac{a^4 \left(-\frac{4iA(\cot^3(dx+c))}{3} - \frac{A(\cot^4(dx+c))}{4} - 2iB(\cot^2(dx+c)) - \frac{B(\cot^3(dx+c))}{3} + 8iA \cot(dx+c) + \frac{7A(\cot^2(dx+c))}{2} + 7 \cot(dx+c) \right)}{d}$
default	$\frac{a^4 \left(-\frac{4iA(\cot^3(dx+c))}{3} - \frac{A(\cot^4(dx+c))}{4} - 2iB(\cot^2(dx+c)) - \frac{B(\cot^3(dx+c))}{3} + 8iA \cot(dx+c) + \frac{7A(\cot^2(dx+c))}{2} + 7 \cot(dx+c) \right)}{d}$
risch	$-\frac{16a^4 Bc}{d} - \frac{16ia^4 Ac}{d} + \frac{4ia^4 (30iA e^{6i(dx+c)} + 18B e^{6i(dx+c)} - 63iA e^{4i(dx+c)} - 45B e^{4i(dx+c)} + 50iA e^{2i(dx+c)} + 38B)}{3d(e^{2i(dx+c)} - 1)^4}$
norman	$\frac{\left(\frac{8iA a^4 + 7B a^4}{d} (\tan^3(dx+c)) + (8iA a^4 + 8B a^4) x (\tan^4(dx+c)) - \frac{A a^4}{4d} + \frac{(-4iB a^4 + 7A a^4) (\tan^2(dx+c))}{2d} - \frac{(4iA a^4 + B a^4) \tan(dx+c)}{3d} \right)}{\tan(dx+c)^4}$

```
input int(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

```
output 8*a^4*((-1/2*A+1/2*I*B)*ln(sec(d*x+c)^2)+(A-I*B)*ln(tan(d*x+c))-1/32*A*cot
(d*x+c)^4+cot(d*x+c)^3*(-1/6*I*A-1/24*B)+cot(d*x+c)^2*(-1/4*I*B+7/16*A)+co
t(d*x+c)*(I*A+7/8*B)+(I*A+B)*x*d)/d
```

3.33. $\int \cot^5(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$

3.33.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.29

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx = \frac{4(6(5A - 3iB)a^4e^{(6idx+6ic)} - 9(7A - 5iB)a^4e^{(4idx+4ic)} + 2(25A - 19iB)a^4e^{(2idx+2ic)} - (14A - 11iB)a^4e^{(0idx+0ic)})}{3(de^{(8idx+8ic)} - 1)}$$

input `integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output `-4/3*(6*(5*A - 3*I*B)*a^4*e^(6*I*d*x + 6*I*c) - 9*(7*A - 5*I*B)*a^4*e^(4*I*d*x + 4*I*c) + 2*(25*A - 19*I*B)*a^4*e^(2*I*d*x + 2*I*c) - (14*A - 11*I*B)*a^4 - 6*((A - I*B)*a^4*e^(8*I*d*x + 8*I*c) - 4*(A - I*B)*a^4*e^(6*I*d*x + 6*I*c) + 6*(A - I*B)*a^4*e^(4*I*d*x + 4*I*c) - 4*(A - I*B)*a^4*e^(2*I*d*x + 2*I*c) + (A - I*B)*a^4)*log(e^(2*I*d*x + 2*I*c) - 1)/(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)`

3.33.6 Sympy [A] (verification not implemented)

Time = 0.65 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.33

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx = \frac{8a^4(A - iB) \log(e^{2idx} - e^{-2ic})}{d} + \frac{56Aa^4 - 44iBa^4 + (-200Aa^4e^{2ic} + 152iBa^4e^{2ic})e^{2idx} + (252Aa^4e^{4ic} - 180iBa^4e^{4ic})e^{4idx} + (-120Aa^4e^{6ic} + 72iBa^4e^{6ic})e^{6idx} + (-14Aa^4e^{8ic} + 11iBa^4e^{8ic})e^{8idx}}{3de^{8ic}e^{8idx} - 12de^{6ic}e^{6idx} + 18de^{4ic}e^{4idx} - 12de^{2ic}e^{2idx} + 3d}$$

input `integrate(cot(d*x+c)**5*(a+I*a*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)`

output `8*a**4*(A - I*B)*log(exp(2*I*d*x) - exp(-2*I*c))/d + (56*A*a**4 - 44*I*B*a**4 + (-200*A*a**4*exp(2*I*c) + 152*I*B*a**4*exp(2*I*c))*exp(2*I*d*x) + (252*A*a**4*exp(4*I*c) - 180*I*B*a**4*exp(4*I*c))*exp(4*I*d*x) + (-120*A*a**4*exp(6*I*c) + 72*I*B*a**4*exp(6*I*c))*exp(6*I*d*x))/(3*d*exp(8*I*c)*exp(8*I*d*x) - 12*d*exp(6*I*c)*exp(6*I*d*x) + 18*d*exp(4*I*c)*exp(4*I*d*x) - 12*d*exp(2*I*c)*exp(2*I*d*x) + 3*d)`

3.33.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.77

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$\frac{96(dx + c)(-iA - B)a^4 + 48(A - iB)a^4 \log(\tan(dx + c)^2 + 1) - 96(A - iB)a^4 \log(\tan(dx + c)) - 12d}{12d}$$

input `integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/12*(96*(d*x + c)*(-I*A - B)*a^4 + 48*(A - I*B)*a^4*log(tan(d*x + c)^2 + 1) - 96*(A - I*B)*a^4*log(tan(d*x + c)) - (12*(8*I*A + 7*B)*a^4*tan(d*x + c)^3 + 6*(7*A - 4*I*B)*a^4*tan(d*x + c)^2 + 4*(-4*I*A - B)*a^4*tan(d*x + c) - 3*A*a^4)/tan(d*x + c)^4/d`

3.33.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 322 vs. 2(153) = 306.

Time = 1.12 (sec) , antiderivative size = 322, normalized size of antiderivative = 1.82

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$\frac{3Aa^4 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^4 - 32iAa^4 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 8Ba^4 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 180Aa^4 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + 120Aa^4 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 60Aa^4}{12d}$$

input `integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output
$$\begin{aligned} & -1/192*(3*A*a^4*\tan(1/2*d*x + 1/2*c)^4 - 32*I*A*a^4*\tan(1/2*d*x + 1/2*c)^3 \\ & - 8*B*a^4*\tan(1/2*d*x + 1/2*c)^3 - 180*A*a^4*\tan(1/2*d*x + 1/2*c)^2 + 96* \\ & I*B*a^4*\tan(1/2*d*x + 1/2*c)^2 + 864*I*A*a^4*\tan(1/2*d*x + 1/2*c) + 696*B* \\ & a^4*\tan(1/2*d*x + 1/2*c) + 3072*(A*a^4 - I*B*a^4)*\log(\tan(1/2*d*x + 1/2*c) \\ & + I) - 1536*(A*a^4 - I*B*a^4)*\log(\tan(1/2*d*x + 1/2*c)) + (3200*A*a^4*\tan \\ & (1/2*d*x + 1/2*c)^4 - 3200*I*B*a^4*\tan(1/2*d*x + 1/2*c)^4 - 864*I*A*a^4*\tan \\ & (1/2*d*x + 1/2*c)^3 - 696*B*a^4*\tan(1/2*d*x + 1/2*c)^3 - 180*A*a^4*\tan(1/ \\ & 2*d*x + 1/2*c)^2 + 96*I*B*a^4*\tan(1/2*d*x + 1/2*c)^2 + 32*I*A*a^4*\tan(1/2* \\ & d*x + 1/2*c) + 8*B*a^4*\tan(1/2*d*x + 1/2*c) + 3*A*a^4)/\tan(1/2*d*x + 1/2*c \\ &)^4)/d \end{aligned}$$

3.33.9 Mupad [B] (verification not implemented)

Time = 7.90 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.64

$$\begin{aligned} & \int \cot^5(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx \\ & = \frac{\tan(c + dx)^2 \left(\frac{7Aa^4}{2} - Ba^4 2i \right) + \tan(c + dx)^3 (7Ba^4 + Aa^4 8i) - \frac{Aa^4}{4} - \tan(c + dx) \left(\frac{Ba^4}{3} + \frac{Aa^4 4i}{3} \right)}{d \tan(c + dx)^4} \\ & + \frac{16a^4 \operatorname{atan}(2 \tan(c + dx) + 1i) (B + A 1i)}{d} \end{aligned}$$

input `int(cot(c + d*x)^5*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^4,x)`

output
$$\begin{aligned} & (\tan(c + d*x)^2*((7*A*a^4)/2 - B*a^4*2i) + \tan(c + d*x)^3*(A*a^4*8i + 7*B* \\ & a^4) - (A*a^4)/4 - \tan(c + d*x)*((A*a^4*4i)/3 + (B*a^4)/3))/ (d*\tan(c + d*x \\ &)^4) + (16*a^4*\operatorname{atan}(2*\tan(c + d*x) + 1i)*(A*1i + B))/d \end{aligned}$$

3.34 $\int \cot^6(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.34.1	Optimal result	560
3.34.2	Mathematica [A] (verified)	561
3.34.3	Rubi [A] (verified)	561
3.34.4	Maple [A] (verified)	566
3.34.5	Fricas [A] (verification not implemented)	566
3.34.6	Sympy [A] (verification not implemented)	567
3.34.7	Maxima [A] (verification not implemented)	567
3.34.8	Giac [B] (verification not implemented)	568
3.34.9	Mupad [B] (verification not implemented)	569

3.34.1 Optimal result

Integrand size = 34, antiderivative size = 200

$$\int \cot^6(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= -8a^4(A-iB)x - \frac{8a^4(A-iB) \cot(c+dx)}{d} + \frac{a^4(148iA+145B) \cot^2(c+dx)}{60d}$$

$$+ \frac{8a^4(iA+B) \log(\sin(c+dx))}{d} - \frac{aA \cot^5(c+dx)(a+ia \tan(c+dx))^3}{5d}$$

$$- \frac{(8iA+5B) \cot^4(c+dx) (a^2+ia^2 \tan(c+dx))^2}{20d}$$

$$+ \frac{(28A-25iB) \cot^3(c+dx) (a^4+ia^4 \tan(c+dx))}{30d}$$

output

```
-8*a^4*(A-I*B)*x-8*a^4*(A-I*B)*cot(d*x+c)/d+1/60*a^4*(148*I*A+145*B)*cot(d*x+c)^2/d+8*a^4*(I*A+B)*ln(sin(d*x+c))/d-1/5*a*A*cot(d*x+c)^5*(a+I*a*tan(d*x+c))^3/d-1/20*(8*I*A+5*B)*cot(d*x+c)^4*(a^2+I*a^2*tan(d*x+c))^2/d+1/30*(28*A-25*I*B)*cot(d*x+c)^3*(a^4+I*a^4*tan(d*x+c))/d
```

3.34.2 Mathematica [A] (verified)

Time = 1.65 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.60

$$\int \cot^6(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{a^4(-3i(4A - 5iB)(i + \cot(c + dx))^4 - 12A \cot(c + dx)(i + \cot(c + dx))^4 + 20(A - iB)(-21 \cot(c + dx) + \cot^3(c + dx)))}{60d}$$

input `Integrate[Cot[c + d*x]^6*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(a^4*((-3*I)*(4*A - (5*I)*B)*(I + Cot[c + d*x])^4 - 12*A*Cot[c + d*x]*(I + Cot[c + d*x])^4 + 20*(A - I*B)*(-21*Cot[c + d*x] + (6*I)*Cot[c + d*x]^2 + Cot[c + d*x]^3 + (24*I)*(Log[Tan[c + d*x]] - Log[I + Tan[c + d*x]]))))/(60*d)`

3.34.3 Rubi [A] (verified)

Time = 1.46 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.10, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4076, 3042, 4076, 27, 3042, 4076, 3042, 4074, 27, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^6(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + ia \tan(c + dx))^4(A + B \tan(c + dx))}{\tan(c + dx)^6} dx$$

$$\downarrow \text{4076}$$

$$\frac{1}{5} \int \cot^5(c + dx)(i \tan(c + dx)a + a)^3(a(8iA + 5B) - a(2A - 5iB) \tan(c + dx)) dx - \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^3}{5d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{5} \int \frac{(i \tan(c + dx)a + a)^3 (a(8iA + 5B) - a(2A - 5iB) \tan(c + dx))}{\tan(c + dx)^5} dx - \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^3}{5d}$$

↓ 4076

$$\frac{1}{5} \left(\frac{1}{4} \int -2 \cot^4(c + dx)(i \tan(c + dx)a + a)^2 ((28A - 25iB)a^2 + 3(4iA + 5B) \tan(c + dx)a^2) dx - \frac{(5B + 8iA) \cot^4(c + dx)(a + ia \tan(c + dx))^3}{5d} \right)$$

↓ 27

$$\frac{1}{5} \left(-\frac{1}{2} \int \cot^4(c + dx)(i \tan(c + dx)a + a)^2 ((28A - 25iB)a^2 + 3(4iA + 5B) \tan(c + dx)a^2) dx - \frac{(5B + 8iA) \cot^4(c + dx)(a + ia \tan(c + dx))^3}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(-\frac{1}{2} \int \frac{(i \tan(c + dx)a + a)^2 ((28A - 25iB)a^2 + 3(4iA + 5B) \tan(c + dx)a^2)}{\tan(c + dx)^4} dx - \frac{(5B + 8iA) \cot^4(c + dx)(a + ia \tan(c + dx))^3}{4d} \right)$$

↓ 4076

$$\frac{1}{5} \left(\frac{1}{2} \left(\frac{(28A - 25iB) \cot^3(c + dx)(a^4 + ia^4 \tan(c + dx))}{3d} - \frac{1}{3} \int \cot^3(c + dx)(i \tan(c + dx)a + a)(a^3(148iA + 145B) - a^3(92iA + 145B) \tan(c + dx))}{\tan(c + dx)^3} dx - \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^3}{5d} \right) \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{2} \left(\frac{(28A - 25iB) \cot^3(c + dx)(a^4 + ia^4 \tan(c + dx))}{3d} - \frac{1}{3} \int \frac{(i \tan(c + dx)a + a)(a^3(148iA + 145B) - a^3(92iA + 145B) \tan(c + dx))}{\tan(c + dx)^3} dx - \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^3}{5d} \right) \right)$$

↓ 4074

3.34. $\int \cot^6(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$

$$\frac{1}{5} \left(\frac{1}{2} \left(\frac{1}{3} \left(\frac{a^4(145B + 148iA) \cot^2(c + dx)}{2d} - \int -240 \cot^2(c + dx) ((A - iB)a^4 + (iA + B) \tan(c + dx)a^4) dx \right) + \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^3}{5d} \right) \right) \downarrow 27$$

$$\frac{1}{5} \left(\frac{1}{2} \left(\frac{1}{3} \left(240 \int \cot^2(c + dx) ((A - iB)a^4 + (iA + B) \tan(c + dx)a^4) dx + \frac{a^4(145B + 148iA) \cot^2(c + dx)}{2d} \right) + \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^3}{5d} \right) \right) \downarrow 3042$$

$$\frac{1}{5} \left(\frac{1}{2} \left(\frac{1}{3} \left(240 \int \frac{(A - iB)a^4 + (iA + B) \tan(c + dx)a^4}{\tan(c + dx)^2} dx + \frac{a^4(145B + 148iA) \cot^2(c + dx)}{2d} \right) + \frac{(28A - 25iB) \cot^2(c + dx)}{2d} + \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^3}{5d} \right) \right) \downarrow 4012$$

$$\frac{1}{5} \left(\frac{1}{2} \left(\frac{1}{3} \left(240 \left(\int \cot(c + dx) (a^4(iA + B) - a^4(A - iB) \tan(c + dx)) dx - \frac{a^4(A - iB) \cot(c + dx)}{d} \right) + \frac{a^4(145B + 148iA) \cot^2(c + dx)}{2d} \right) + \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^3}{5d} \right) \right) \downarrow 3042$$

$$\frac{1}{5} \left(\frac{1}{2} \left(\frac{1}{3} \left(240 \left(\int \frac{a^4(iA + B) - a^4(A - iB) \tan(c + dx)}{\tan(c + dx)} dx - \frac{a^4(A - iB) \cot(c + dx)}{d} \right) + \frac{a^4(145B + 148iA) \cot^2(c + dx)}{2d} \right) + \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^3}{5d} \right) \right) \downarrow 4014$$

$$\frac{1}{5} \left(\frac{1}{2} \left(\frac{1}{3} \left(240 \left(a^4(B + iA) \int \cot(c + dx) dx - \frac{a^4(A - iB) \cot(c + dx)}{d} - (a^4 x(A - iB)) \right) + \frac{a^4(145B + 148iA) \cot^2(c + dx)}{2d} \right) + \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^3}{5d} \right) \right) \downarrow 3042$$

3.34. $\int \cot^6(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$

$$\frac{1}{5} \left(\frac{1}{2} \left(\frac{1}{3} \left(240 \left(a^4(B + iA) \int -\tan \left(c + dx + \frac{\pi}{2} \right) dx - \frac{a^4(A - iB) \cot(c + dx)}{d} - (a^4 x(A - iB)) \right) + \frac{a^4(145B + 148iA) \cot^2(c + dx)}{2d} + 240 \left(-\frac{a^4(A - iB)x}{d} + \frac{a^4(A - iB) \cot(c + dx)}{d} \right) \right) \right) \right) \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^3}{5d}$$

↓ 25

$$\frac{1}{5} \left(\frac{1}{2} \left(\frac{1}{3} \left(240 \left(-a^4(B + iA) \int \tan \left(\frac{1}{2}(2c + \pi) + dx \right) dx - \frac{a^4(A - iB) \cot(c + dx)}{d} - (a^4 x(A - iB)) \right) + \frac{a^4(145B + 148iA) \cot^2(c + dx)}{2d} + 240 \left(-\frac{a^4(A - iB)x}{d} + \frac{a^4(A - iB) \cot(c + dx)}{d} \right) \right) \right) \right) \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^3}{5d}$$

↓ 3956

$$\frac{1}{5} \left(\frac{1}{2} \left(\frac{(28A - 25iB) \cot^3(c + dx) (a^4 + ia^4 \tan(c + dx))}{3d} + \frac{1}{3} \left(\frac{a^4(145B + 148iA) \cot^2(c + dx)}{2d} + 240 \left(-\frac{a^4(A - iB)x}{d} + \frac{a^4(A - iB) \cot(c + dx)}{d} \right) \right) \right) \right) \frac{aA \cot^5(c + dx)(a + ia \tan(c + dx))^3}{5d}$$

input `Int[Cot[c + d*x]^6*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `-1/5*(a*A*Cot[c + d*x]^5*(a + I*a*Tan[c + d*x])^3)/d + (-1/4*(((8*I)*A + 5*B)*Cot[c + d*x]^4*(a^2 + I*a^2*Tan[c + d*x])^2)/d + (((a^4*(((148*I)*A + 145*B)*Cot[c + d*x]^2)/(2*d) + 240*(-(a^4*(A - I*B)*x) - (a^4*(A - I*B)*Cot[c + d*x])/d + (a^4*(I*A + B)*Log[-Sin[c + d*x]])/d))/3 + ((28*A - (25*I)*B)*Cot[c + d*x]^3*(a^4 + I*a^4*Tan[c + d*x]))/(3*d))/2)/5`

3.34.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4074 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

rule 4076 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.34.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.65

method	result
parallelrisc	$\frac{8a^4 \left(\left(-\frac{iA}{2} - \frac{B}{2} \right) \ln(\sec^2(dx+c)) + (iA+B) \ln(\tan(dx+c)) - \frac{A(\cot^5(dx+c))}{40} + (\cot^4(dx+c)) \left(-\frac{iA}{8} - \frac{B}{32} \right) + (\cot^3(dx+c)) \left(\frac{iA}{8} + \frac{B}{32} \right) \right)}{d}$
derivativedivides	$\frac{a^4 \left(-iA(\cot^4(dx+c)) - \frac{A(\cot^5(dx+c))}{5} - \frac{4iB(\cot^3(dx+c))}{3} - \frac{B(\cot^4(dx+c))}{4} + 4iA(\cot^2(dx+c)) + \frac{7A(\cot^3(dx+c))}{3} + 8iB \cot(dx+c) \right)}{d}$
default	$\frac{a^4 \left(-iA(\cot^4(dx+c)) - \frac{A(\cot^5(dx+c))}{5} - \frac{4iB(\cot^3(dx+c))}{3} - \frac{B(\cot^4(dx+c))}{4} + 4iA(\cot^2(dx+c)) + \frac{7A(\cot^3(dx+c))}{3} + 8iB \cot(dx+c) \right)}{d}$
risc	$-\frac{16ia^4Bc}{d} + \frac{16a^4Ac}{d} - \frac{4a^4(210iAe^{8i(dx+c)} + 150Be^{8i(dx+c)} - 555iAe^{6i(dx+c)} - 465Be^{6i(dx+c)} + 655iAe^{4i(dx+c)} + 15d(e^{2i(dx+c)} - 1)^5)}{15d(e^{2i(dx+c)} - 1)^5}$
norman	$\frac{(8iBa^4 - 8Aa^4)x(\tan^5(dx+c)) - \frac{Aa^4}{5d} + \frac{(-4iBa^4 + 7Aa^4)(\tan^2(dx+c))}{3d} - \frac{(4iAa^4 + Ba^4)\tan(dx+c)}{4d} - \frac{8(-iBa^4 + Aa^4)(\tan^4(dx+c))}{d}}{\tan(dx+c)^5}$

input `int(cot(d*x+c)^6*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `8*a^4*((-1/2*I*A-1/2*B)*ln(sec(d*x+c)^2)+(I*A+B)*ln(tan(d*x+c))-1/40*A*cot(d*x+c)^5+cot(d*x+c)^4*(-1/8*I*A-1/32*B)+cot(d*x+c)^3*(-1/6*I*B+7/24*A)+cot(d*x+c)^2*(1/2*I*A+7/16*B)+(-A+I*B)*cot(d*x+c)+x*d*(-A+I*B))/d`

3.34.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.44

$$\int \cot^6(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx = \frac{4(30(7iA+5B)a^4e^{(8i dx+8i c)} + 15(-37iA-31B)a^4e^{(6i dx+6i c)} + 5(131iA+113B)a^4e^{(4i dx+4i c)} + 5(-131iA-113B)a^4e^{(2i dx+2i c)} + 5(-131iA-113B)a^4e^{(0i dx+0i c)})}{d}$$

input `integrate(cot(d*x+c)^6*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output
$$-4/15*(30*(7*I*A + 5*B)*a^4*e^{(8*I*d*x + 8*I*c)} + 15*(-37*I*A - 31*B)*a^4*e^{(6*I*d*x + 6*I*c)} + 5*(131*I*A + 113*B)*a^4*e^{(4*I*d*x + 4*I*c)} + 5*(-73*I*A - 64*B)*a^4*e^{(2*I*d*x + 2*I*c)} + (79*I*A + 70*B)*a^4 + 30*((-I*A - B)*a^4*e^{(10*I*d*x + 10*I*c)} + 5*(I*A + B)*a^4*e^{(8*I*d*x + 8*I*c)} + 10*(-I*A - B)*a^4*e^{(6*I*d*x + 6*I*c)} + 10*(I*A + B)*a^4*e^{(4*I*d*x + 4*I*c)} + 5*(-I*A - B)*a^4*e^{(2*I*d*x + 2*I*c)} + (I*A + B)*a^4)*\log(e^{(2*I*d*x + 2*I*c)} - 1)/(d*e^{(10*I*d*x + 10*I*c)} - 5*d*e^{(8*I*d*x + 8*I*c)} + 10*d*e^{(6*I*d*x + 6*I*c)} - 10*d*e^{(4*I*d*x + 4*I*c)} + 5*d*e^{(2*I*d*x + 2*I*c)} - d)$$

3.34.6 Sympy [A] (verification not implemented)

Time = 2.52 (sec) , antiderivative size = 296, normalized size of antiderivative = 1.48

$$\int \cot^6(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx = \frac{8ia^4(A - iB) \log(e^{2idx} - e^{-2ic})}{d} + \frac{-316iAa^4 - 280Ba^4 + (1460iAa^4e^{2ic} + 1280Ba^4e^{2ic})e^{2idx} + (-2620iAa^4e^{4ic} - 2260Ba^4e^{4ic})e^{4idx} + (2220iAa^4e^{6ic} + 1860Ba^4e^{6ic})e^{6idx} + (-840iAa^4e^{8ic} - 600Ba^4e^{8ic})e^{8idx}}{15de^{10ic}e^{10idx} - 75de^{8ic}e^{8idx} + 150de^{6ic}e^{6idx} - 150de^{4ic}e^{4idx} - 15d}$$

input `integrate(cot(d*x+c)**6*(a+I*a*tan(d*x+c))**4*(A+B*tan(d*x+c)), x)`

output
$$8*I*a**4*(A - I*B)*\log(\exp(2*I*d*x) - \exp(-2*I*c))/d + (-316*I*A*a**4 - 280*B*a**4 + (1460*I*A*a**4*\exp(2*I*c) + 1280*B*a**4*\exp(2*I*c))*\exp(2*I*d*x) + (-2620*I*A*a**4*\exp(4*I*c) - 2260*B*a**4*\exp(4*I*c))*\exp(4*I*d*x) + (2220*I*A*a**4*\exp(6*I*c) + 1860*B*a**4*\exp(6*I*c))*\exp(6*I*d*x) + (-840*I*A*a**4*\exp(8*I*c) - 600*B*a**4*\exp(8*I*c))*\exp(8*I*d*x))/(15*d*\exp(10*I*c)*\exp(10*I*d*x) - 75*d*\exp(8*I*c)*\exp(8*I*d*x) + 150*d*\exp(6*I*c)*\exp(6*I*d*x) - 150*d*\exp(4*I*c)*\exp(4*I*d*x) + 75*d*\exp(2*I*c)*\exp(2*I*d*x) - 15*d)$$

3.34.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.76

$$\int \cot^6(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx = \frac{480(dx + c)(A - iB)a^4 + 240(iA + B)a^4 \log(\tan(dx + c)^2 + 1) + 480(-iA - B)a^4 \log(\tan(dx + c) + \cot(dx + c))}{60d}$$

input `integrate(cot(d*x+c)^6*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)), x, algorithm="maxima")`

output
$$\frac{-1/60*(480*(d*x + c)*(A - I*B)*a^4 + 240*(I*A + B)*a^4*\log(\tan(d*x + c)^2 + 1) + 480*(-I*A - B)*a^4*\log(\tan(d*x + c)) + (480*(A - I*B)*a^4*\tan(d*x + c)^4 - 30*(8*I*A + 7*B)*a^4*\tan(d*x + c)^3 - 20*(7*A - 4*I*B)*a^4*\tan(d*x + c)^2 - 15*(-4*I*A - B)*a^4*\tan(d*x + c) + 12*A*a^4)/\tan(d*x + c)^5}{d}$$

3.34.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 391 vs. $2(174) = 348$.

Time = 1.22 (sec) , antiderivative size = 391, normalized size of antiderivative = 1.96

$$\int \cot^6(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$6 Aa^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 60i Aa^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 - 15 Ba^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 - 310 Aa^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)$$

input `integrate(cot(d*x+c)^6*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output
$$\frac{1/960*(6*A*a^4*\tan(1/2*d*x + 1/2*c)^5 - 60*I*A*a^4*\tan(1/2*d*x + 1/2*c)^4 - 15*B*a^4*\tan(1/2*d*x + 1/2*c)^4 - 310*A*a^4*\tan(1/2*d*x + 1/2*c)^3 + 160*I*B*a^4*\tan(1/2*d*x + 1/2*c)^3 + 1200*I*A*a^4*\tan(1/2*d*x + 1/2*c)^2 + 900*B*a^4*\tan(1/2*d*x + 1/2*c)^2 + 4740*A*a^4*\tan(1/2*d*x + 1/2*c) - 4320*I*B*a^4*\tan(1/2*d*x + 1/2*c) - 15360*(I*A*a^4 + B*a^4)*\log(\tan(1/2*d*x + 1/2*c) + I) - 7680*(-I*A*a^4 - B*a^4)*\log(\tan(1/2*d*x + 1/2*c)) + (-17536*I*A*a^4*\tan(1/2*d*x + 1/2*c)^5 - 17536*B*a^4*\tan(1/2*d*x + 1/2*c)^5 - 4740*A*a^4*\tan(1/2*d*x + 1/2*c)^4 + 4320*I*B*a^4*\tan(1/2*d*x + 1/2*c)^4 + 1200*I*A*a^4*\tan(1/2*d*x + 1/2*c)^3 + 900*B*a^4*\tan(1/2*d*x + 1/2*c)^3 + 310*A*a^4*\tan(1/2*d*x + 1/2*c)^2 - 160*I*B*a^4*\tan(1/2*d*x + 1/2*c)^2 - 60*I*A*a^4*\tan(1/2*d*x + 1/2*c) - 15*B*a^4*\tan(1/2*d*x + 1/2*c) - 6*A*a^4)/\tan(1/2*d*x + 1/2*c)^5}{d}$$

3.34.9 Mupad [B] (verification not implemented)

Time = 7.94 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.70

$$\int \cot^6(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$\frac{\frac{Aa^4}{5} - \tan(c + dx)^2 \left(\frac{7Aa^4}{3} - \frac{Ba^4 4i}{3} \right) + \tan(c + dx)^4 (8Aa^4 - Ba^4 8i) - \tan(c + dx)^3 \left(\frac{7Ba^4}{2} + Aa^4 4i \right)}{d \tan(c + dx)^5}$$

$$+ \frac{a^4 \operatorname{atan}(2 \tan(c + dx) + 1i) (B + A 1i) 16i}{d}$$

input `int(cot(c + d*x)^6*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^4,x)`output `(a^4*atan(2*tan(c + d*x) + 1i)*(A*1i + B)*16i)/d - (tan(c + d*x)^4*(8*A*a^4 - B*a^4*8i) - tan(c + d*x)^2*((7*A*a^4)/3 - (B*a^4*4i)/3) - tan(c + d*x)^3*(A*a^4*4i + (7*B*a^4)/2) + (A*a^4)/5 + tan(c + d*x)*(A*a^4*1i + (B*a^4/4)))/(d*tan(c + d*x)^5)`

3.35 $\int \cot^7(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.35.1	Optimal result	570
3.35.2	Mathematica [A] (verified)	571
3.35.3	Rubi [A] (verified)	571
3.35.4	Maple [A] (verified)	576
3.35.5	Fricas [A] (verification not implemented)	577
3.35.6	Sympy [A] (verification not implemented)	577
3.35.7	Maxima [A] (verification not implemented)	578
3.35.8	Giac [B] (verification not implemented)	578
3.35.9	Mupad [B] (verification not implemented)	579

3.35.1 Optimal result

Integrand size = 34, antiderivative size = 223

$$\begin{aligned} & \int \cot^7(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx \\ &= -8a^4(iA+B)x - \frac{8a^4(iA+B) \cot(c+dx)}{d} \\ & \quad - \frac{4a^4(A-iB) \cot^2(c+dx)}{d} + \frac{a^4(93iA+92B) \cot^3(c+dx)}{60d} \\ & \quad - \frac{8a^4(A-iB) \log(\sin(c+dx))}{d} - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d} \\ & \quad - \frac{(3iA+2B) \cot^5(c+dx) (a^2+ia^2 \tan(c+dx))^2}{10d} \\ & \quad + \frac{(13A-12iB) \cot^4(c+dx) (a^4+ia^4 \tan(c+dx))}{20d} \end{aligned}$$

output

```
-8*a^4*(I*A+B)*x-8*a^4*(I*A+B)*cot(d*x+c)/d-4*a^4*(A-I*B)*cot(d*x+c)^2/d+1/60*a^4*(93*I*A+92*B)*cot(d*x+c)^3/d-8*a^4*(A-I*B)*ln(sin(d*x+c))/d-1/6*a*A*cot(d*x+c)^6*(a+I*a*tan(d*x+c))^3/d-1/10*(3*I*A+2*B)*cot(d*x+c)^5*(a^2+I*a^2*tan(d*x+c))^2/d+1/20*(13*A-12*I*B)*cot(d*x+c)^4*(a^4+I*a^4*tan(d*x+c))/d
```

3.35.2 Mathematica [A] (verified)

Time = 3.74 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.67

$$\int \cot^7(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{a^4((13A - 12iB)(i + \cot(c + dx))^4 - 4i(2A - 3iB) \cot(c + dx)(i + \cot(c + dx))^4 - 10A \cot^2(c + dx)(i + \cot(c + dx))^4 - 10A \cot^4(c + dx)(i + \cot(c + dx))^2 + 10A \cot^6(c + dx)(i + \cot(c + dx))^2 - 10A \cot^8(c + dx)(i + \cot(c + dx))^2)}{60d}$$

input `Integrate[Cot[c + d*x]^7*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(a^4*((13*A - (12*I)*B)*(I + Cot[c + d*x])^4 - (4*I)*(2*A - (3*I)*B)*Cot[c + d*x]*(I + Cot[c + d*x])^4 - 10*A*Cot[c + d*x]^2*(I + Cot[c + d*x])^4 + 20*(I*A + B)*(-21*Cot[c + d*x] + (6*I)*Cot[c + d*x]^2 + Cot[c + d*x]^3 + (24*I)*(Log[Tan[c + d*x]] - Log[I + Tan[c + d*x]]))))/(60*d)`

3.35.3 Rubi [A] (verified)

Time = 1.66 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.10, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.618$, Rules used = {3042, 4076, 27, 3042, 4076, 27, 3042, 4076, 3042, 4074, 27, 3042, 4012, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^7(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + ia \tan(c + dx))^4(A + B \tan(c + dx))}{\tan(c + dx)^7} dx$$

$$\downarrow \text{4076}$$

$$\frac{1}{6} \int 3 \cot^6(c + dx)(i \tan(c + dx)a + a)^3(a(3iA + 2B) - a(A - 2iB) \tan(c + dx)) dx - \frac{aA \cot^6(c + dx)(a + ia \tan(c + dx))^3}{6d}$$

$$\downarrow \text{27}$$

$$\frac{1}{2} \int \cot^6(c+dx)(i \tan(c+dx)a+a)^3(a(3iA+2B)-a(A-2iB)\tan(c+dx))dx - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d}$$

↓ 3042

$$\frac{1}{2} \int \frac{(i \tan(c+dx)a+a)^3(a(3iA+2B)-a(A-2iB)\tan(c+dx))}{\tan(c+dx)^6} dx - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d}$$

↓ 4076

$$\frac{1}{2} \left(\frac{1}{5} \int -2 \cot^5(c+dx)(i \tan(c+dx)a+a)^2((13A-12iB)a^2+(7iA+8B)\tan(c+dx)a^2) dx - \frac{(2B+3iA)\cot^5(c+dx)}{5d} \right) - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d}$$

↓ 27

$$\frac{1}{2} \left(-\frac{2}{5} \int \cot^5(c+dx)(i \tan(c+dx)a+a)^2((13A-12iB)a^2+(7iA+8B)\tan(c+dx)a^2) dx - \frac{(2B+3iA)\cot^5(c+dx)}{5d} \right) - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d}$$

↓ 3042

$$\frac{1}{2} \left(-\frac{2}{5} \int \frac{(i \tan(c+dx)a+a)^2((13A-12iB)a^2+(7iA+8B)\tan(c+dx)a^2)}{\tan(c+dx)^5} dx - \frac{(2B+3iA)\cot^5(c+dx)}{5d} \right) - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d}$$

↓ 4076

$$\frac{1}{2} \left(-\frac{2}{5} \left(\frac{1}{4} \int \cot^4(c+dx)(i \tan(c+dx)a+a)(a^3(93iA+92B)-a^3(67A-68iB)\tan(c+dx)) dx - \frac{(13A-12iB)\cot^4(c+dx)}{4d} \right) \right) - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d}$$

↓ 3042

$$\frac{1}{2} \left(-\frac{2}{5} \left(\frac{1}{4} \int \frac{(i \tan(c+dx)a+a)(a^3(93iA+92B)-a^3(67A-68iB)\tan(c+dx))}{\tan(c+dx)^4} dx - \frac{(13A-12iB)\cot^4(c+dx)}{4d} \right) \right) - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d}$$

3.35. $\int \cot^7(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

↓ 4074

$$\frac{1}{2} \left(-\frac{2}{5} \left(\frac{1}{4} \left(\int -160 \cot^3(c+dx) ((A-iB)a^4 + (iA+B)\tan(c+dx)a^4) dx - \frac{a^4(92B+93iA)\cot^3(c+dx)}{3d} \right) - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right)$$

↓ 27

$$\frac{1}{2} \left(-\frac{2}{5} \left(\frac{1}{4} \left(-160 \int \cot^3(c+dx) ((A-iB)a^4 + (iA+B)\tan(c+dx)a^4) dx - \frac{a^4(92B+93iA)\cot^3(c+dx)}{3d} \right) - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right)$$

↓ 3042

$$\frac{1}{2} \left(-\frac{2}{5} \left(\frac{1}{4} \left(-160 \int \frac{(A-iB)a^4 + (iA+B)\tan(c+dx)a^4}{\tan(c+dx)^3} dx - \frac{a^4(92B+93iA)\cot^3(c+dx)}{3d} \right) - \frac{(13A-12iB)}{3d} - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right)$$

↓ 4012

$$\frac{1}{2} \left(-\frac{2}{5} \left(\frac{1}{4} \left(-160 \left(\int \cot^2(c+dx) (a^4(iA+B) - a^4(A-iB)\tan(c+dx)) dx - \frac{a^4(A-iB)\cot^2(c+dx)}{2d} \right) - \frac{a^4}{3d} - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right)$$

↓ 3042

$$\frac{1}{2} \left(-\frac{2}{5} \left(\frac{1}{4} \left(-160 \left(\int \frac{a^4(iA+B) - a^4(A-iB)\tan(c+dx)}{\tan(c+dx)^2} dx - \frac{a^4(A-iB)\cot^2(c+dx)}{2d} \right) - \frac{a^4(92B+93iA)\cot^3(c+dx)}{3d} - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right)$$

↓ 4012

$$\frac{1}{2} \left(-\frac{2}{5} \left(\frac{1}{4} \left(-160 \left(\int -\cot(c+dx) ((A-iB)a^4 + (iA+B)\tan(c+dx)a^4) dx - \frac{a^4(A-iB)\cot^2(c+dx)}{2d} - \frac{a^4}{3d} - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right) \right)$$

↓ 25

3.35. $\int \cot^7(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

$$\frac{1}{2} \left(-\frac{2}{5} \left(\frac{1}{4} \left(-160 \left(- \int \cot(c+dx) ((A-iB)a^4 + (iA+B) \tan(c+dx)a^4) dx - \frac{a^4(A-iB) \cot^2(c+dx)}{2d} - \frac{a^4(B+iA) \cot(c+dx)}{d} - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right) \right)$$

↓ 3042

$$\frac{1}{2} \left(-\frac{2}{5} \left(\frac{1}{4} \left(-160 \left(- \int \frac{(A-iB)a^4 + (iA+B) \tan(c+dx)a^4}{\tan(c+dx)} dx - \frac{a^4(A-iB) \cot^2(c+dx)}{2d} - \frac{a^4(B+iA) \cot(c+dx)}{d} - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right) \right)$$

↓ 4014

$$\frac{1}{2} \left(-\frac{2}{5} \left(\frac{1}{4} \left(-160 \left(-a^4(A-iB) \int \cot(c+dx) dx - \frac{a^4(A-iB) \cot^2(c+dx)}{2d} - \frac{a^4(B+iA) \cot(c+dx)}{d} - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right) \right)$$

↓ 3042

$$\frac{1}{2} \left(-\frac{2}{5} \left(\frac{1}{4} \left(-160 \left(-a^4(A-iB) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx - \frac{a^4(A-iB) \cot^2(c+dx)}{2d} - \frac{a^4(B+iA) \cot(c+dx)}{d} - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right) \right)$$

↓ 25

$$\frac{1}{2} \left(-\frac{2}{5} \left(\frac{1}{4} \left(-160 \left(a^4(A-iB) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx - \frac{a^4(A-iB) \cot^2(c+dx)}{2d} - \frac{a^4(B+iA) \cot(c+dx)}{d} - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right) \right)$$

↓ 3956

$$\frac{1}{2} \left(-\frac{2}{5} \left(\frac{1}{4} \left(-\frac{a^4(92B+93iA) \cot^3(c+dx)}{3d} - 160 \left(-\frac{a^4(A-iB) \cot^2(c+dx)}{2d} - \frac{a^4(B+iA) \cot(c+dx)}{d} - \frac{aA \cot^6(c+dx)(a+ia \tan(c+dx))^3}{6d} \right) \right) \right)$$

input `Int[Cot[c + d*x]^7*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

```
output -1/6*(a*A*Cot[c + d*x]^6*(a + I*a*Tan[c + d*x])^3)/d + (-1/5*(((3*I)*A + 2
*B)*Cot[c + d*x]^5*(a^2 + I*a^2*Tan[c + d*x])^2)/d - (2*((-1/3*(a^4*((93*I
)*A + 92*B)*Cot[c + d*x]^3)/d - 160*(-(a^4*(I*A + B)*x) - (a^4*(I*A + B)*C
ot[c + d*x])/d - (a^4*(A - I*B)*Cot[c + d*x]^2)/(2*d) - (a^4*(A - I*B)*Log
[-Sin[c + d*x]]/d))/4 - ((13*A - (12*I)*B)*Cot[c + d*x]^4*(a^4 + I*a^4*Ta
n[c + d*x]))/(4*d))/5)/2
```

3.35.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4012 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]
```

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

```
rule 4076 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

3.35.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.64

method	result
parallelrisch	$\frac{8a^4 \left(\left(-\frac{A}{2} + \frac{iB}{2} \right) \ln(\sec^2(dx+c)) + (-iB+A) \ln(\tan(dx+c)) + \frac{A(\cot^6(dx+c))}{48} + (\cot^5(dx+c)) \left(\frac{iA}{10} + \frac{B}{40} \right) + (\cot^4(dx+c)) \right)}{d}$
derivativedivides	$a^4 \left(-\frac{4iA(\cot^5(dx+c))}{5} - \frac{A(\cot^6(dx+c))}{6} - iB(\cot^4(dx+c)) - \frac{B(\cot^5(dx+c))}{5} + \frac{8iA(\cot^3(dx+c))}{3} + \frac{7A(\cot^4(dx+c))}{4} + 4iB(\cot^4(dx+c)) \right)$
default	$a^4 \left(-\frac{4iA(\cot^5(dx+c))}{5} - \frac{A(\cot^6(dx+c))}{6} - iB(\cot^4(dx+c)) - \frac{B(\cot^5(dx+c))}{5} + \frac{8iA(\cot^3(dx+c))}{3} + \frac{7A(\cot^4(dx+c))}{4} + 4iB(\cot^4(dx+c)) \right)$
risch	$\frac{16a^4 Bc}{d} + \frac{16ia^4 Ac}{d} - \frac{4ia^4 (270iA e^{10i(dx+c)} + 210B e^{10i(dx+c)} - 855iA e^{8i(dx+c)} - 765B e^{8i(dx+c)} + 1350iA e^{6i(dx+c)} - 1350B e^{6i(dx+c)} - 1350iA e^{4i(dx+c)} - 1350B e^{4i(dx+c)} + 1350iA e^{2i(dx+c)} + 1350B e^{2i(dx+c)} - 1350iA e^{0i(dx+c)} - 1350B e^{0i(dx+c)})}{15d}$
norman	$\frac{(-8iA a^4 - 8B a^4)x(\tan^6(dx+c)) - \frac{A a^4}{6d} + \frac{(-4iB a^4 + 7A a^4)(\tan^2(dx+c))}{4d} - \frac{(4iA a^4 + B a^4) \tan(dx+c)}{5d} - \frac{4(-iB a^4 + A a^4) \tan(dx+c)}{d}}{\tan(dx+c)^6}$

```
input int(cot(d*x+c)^7*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

3.35. $\int \cot^7(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$

output $-8a^4((-1/2A+1/2I*B)*\ln(\sec(dx+c)^2)+(A-I*B)*\ln(\tan(dx+c))+1/48A*\cot(dx+c)^6+\cot(dx+c)^5*(1/10I*A+1/40*B)+\cot(dx+c)^4*(1/8I*B-7/32*A)+\cot(dx+c)^3*(-1/3I*A-7/24*B)+\cot(dx+c)^2*(1/2A-1/2I*B)+\cot(dx+c)*(I*A+B)+(I*A+B)*x*d)/d$

3.35.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 332, normalized size of antiderivative = 1.49

$$\int \cot^7(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= \frac{4(30(9A-7iB)a^4e^{10i dx+10ic} - 45(19A-17iB)a^4e^{8i dx+8ic} + 10(135A-121iB)a^4e^{6i dx+6ic} - 15$$

input `integrate(cot(dx+c)^7*(a+I*a*tan(dx+c))^4*(A+B*tan(dx+c)),x, algorithm="fricas")`

output $4/15*(30*(9A-7I*B)*a^4*e^{(10I*d*x+10I*c)} - 45*(19A-17I*B)*a^4*e^{(8I*d*x+8I*c)} + 10*(135A-121I*B)*a^4*e^{(6I*d*x+6I*c)} - 15*(75A-68I*B)*a^4*e^{(4I*d*x+4I*c)} + 6*(81A-74I*B)*a^4*e^{(2I*d*x+2I*c)} - (86A-79I*B)*a^4 - 30*((A-I*B)*a^4*e^{(12I*d*x+12I*c)} - 6*(A-I*B)*a^4*e^{(10I*d*x+10I*c)} + 15*(A-I*B)*a^4*e^{(8I*d*x+8I*c)} - 20*(A-I*B)*a^4*e^{(6I*d*x+6I*c)} + 15*(A-I*B)*a^4*e^{(4I*d*x+4I*c)} - 6*(A-I*B)*a^4*e^{(2I*d*x+2I*c)} + (A-I*B)*a^4)*\log(e^{(2I*d*x+2I*c)} - 1))/(d*e^{(12I*d*x+12I*c)} - 6*d*e^{(10I*d*x+10I*c)} + 15*d*e^{(8I*d*x+8I*c)} - 20*d*e^{(6I*d*x+6I*c)} + 15*d*e^{(4I*d*x+4I*c)} - 6*d*e^{(2I*d*x+2I*c)} + d)$

3.35.6 Sympy [A] (verification not implemented)

Time = 1.09 (sec) , antiderivative size = 347, normalized size of antiderivative = 1.56

$$\int \cot^7(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx = -\frac{8a^4(A-iB) \log(e^{2idx} - e^{-2ic})}{d}$$

$$+ \frac{-344Aa^4 + 316iBa^4 + (1944Aa^4e^{2ic} - 1776iBa^4e^{2ic})e^{2idx} + (-4500Aa^4e^{4ic} + 4080iBa^4e^{4ic})e^{4idx} + (5$$

$$15de^{12ic}e^{12idx} - 90de^{10ic}e^{10idx} + 225de^{8ic}e^{8idx} - 15de^{6ic}e^{6idx} + 15de^{4ic}e^{4idx} - 15de^{2ic}e^{2idx} + 15d)}{15de^{12ic}e^{12idx} - 90de^{10ic}e^{10idx} + 225de^{8ic}e^{8idx} - 15de^{6ic}e^{6idx} + 15de^{4ic}e^{4idx} - 15de^{2ic}e^{2idx} + 15d}$$

input `integrate(cot(dx+c)**7*(a+I*a*tan(dx+c))**4*(A+B*tan(dx+c)),x)`

3.35. $\int \cot^7(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

output
$$\begin{aligned} & -8a^4(A - IB)\log(\exp(2I dx) - \exp(-2Ic))/d + (-344Aa^4 + 316I \\ & *Ba^4 + (1944Aa^4\exp(2Ic) - 1776IBa^4\exp(2Ic))\exp(2I dx) \\ & + (-4500Aa^4\exp(4Ic) + 4080IBa^4\exp(4Ic))\exp(4I dx) + (54 \\ & 00Aa^4\exp(6Ic) - 4840IBa^4\exp(6Ic))\exp(6I dx) + (-3420Aa \\ & **4\exp(8Ic) + 3060IBa^4\exp(8Ic))\exp(8I dx) + (1080Aa^4\exp \\ & (10Ic) - 840IBa^4\exp(10Ic))\exp(10I dx)/(15d\exp(12Ic)\exp(\\ & 12I dx) - 90d\exp(10Ic)\exp(10I dx) + 225d\exp(8Ic)\exp(8I dx) \\ & - 300d\exp(6Ic)\exp(6I dx) + 225d\exp(4Ic)\exp(4I dx) - 90d\exp(\\ & 2Ic)\exp(2I dx) + 15d) \end{aligned}$$

3.35.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.77

$$\int \cot^7(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$480(dx + c)(iA + B)a^4 - 240(A - iB)a^4 \log(\tan(dx + c)^2 + 1) + 480(A - iB)a^4 \log(\tan(dx + c))$$

input `integrate(cot(d*x+c)^7*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output
$$\begin{aligned} & -1/60*(480*(d*x + c)*(I*A + B)*a^4 - 240*(A - I*B)*a^4*\log(\tan(d*x + c)^2 \\ & + 1) + 480*(A - I*B)*a^4*\log(\tan(d*x + c)) - (480*(-I*A - B)*a^4*\tan(d*x + \\ & c)^5 - 240*(A - I*B)*a^4*\tan(d*x + c)^4 + 20*(8*I*A + 7*B)*a^4*\tan(d*x + \\ & c)^3 + 15*(7*A - 4*I*B)*a^4*\tan(d*x + c)^2 + 12*(-4*I*A - B)*a^4*\tan(d*x + \\ & c) - 10*A*a^4)/\tan(d*x + c)^6)/d \end{aligned}$$

3.35.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 459 vs. $2(195) = 390$.

Time = 1.29 (sec) , antiderivative size = 459, normalized size of antiderivative = 2.06

$$\int \cot^7(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$5Aa^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^6 - 48iAa^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 12Ba^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 240Aa^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)$$

input `integrate(cot(d*x+c)^7*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output
$$\begin{aligned} & -1/1920*(5*A*a^4*\tan(1/2*d*x + 1/2*c)^6 - 48*I*A*a^4*\tan(1/2*d*x + 1/2*c)^5 \\ & - 12*B*a^4*\tan(1/2*d*x + 1/2*c)^5 - 240*A*a^4*\tan(1/2*d*x + 1/2*c)^4 + 1 \\ & 20*I*B*a^4*\tan(1/2*d*x + 1/2*c)^4 + 880*I*A*a^4*\tan(1/2*d*x + 1/2*c)^3 + 6 \\ & 20*B*a^4*\tan(1/2*d*x + 1/2*c)^3 + 2835*A*a^4*\tan(1/2*d*x + 1/2*c)^2 - 2400 \\ & *I*B*a^4*\tan(1/2*d*x + 1/2*c)^2 - 10080*I*A*a^4*\tan(1/2*d*x + 1/2*c) - 948 \\ & 0*B*a^4*\tan(1/2*d*x + 1/2*c) - 30720*(A*a^4 - I*B*a^4)*\log(\tan(1/2*d*x + 1 \\ & /2*c) + I) + 15360*(A*a^4 - I*B*a^4)*\log(\tan(1/2*d*x + 1/2*c)) - (37632*A* \\ & a^4*\tan(1/2*d*x + 1/2*c)^6 - 37632*I*B*a^4*\tan(1/2*d*x + 1/2*c)^6 - 10080* \\ & I*A*a^4*\tan(1/2*d*x + 1/2*c)^5 - 9480*B*a^4*\tan(1/2*d*x + 1/2*c)^5 - 2835* \\ & A*a^4*\tan(1/2*d*x + 1/2*c)^4 + 2400*I*B*a^4*\tan(1/2*d*x + 1/2*c)^4 + 880*I \\ & *A*a^4*\tan(1/2*d*x + 1/2*c)^3 + 620*B*a^4*\tan(1/2*d*x + 1/2*c)^3 + 240*A*a \\ & ^4*\tan(1/2*d*x + 1/2*c)^2 - 120*I*B*a^4*\tan(1/2*d*x + 1/2*c)^2 - 48*I*A*a^4 \\ & *\tan(1/2*d*x + 1/2*c) - 12*B*a^4*\tan(1/2*d*x + 1/2*c) - 5*A*a^4)/\tan(1/2* \\ & d*x + 1/2*c)^6)/d \end{aligned}$$

3.35.9 Mupad [B] (verification not implemented)

Time = 9.45 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.73

$$\int \cot^7(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx = \frac{\tan(c + dx)^4(4Aa^4 - Ba^4 4i) - \tan(c + dx)^2 \left(\frac{7Aa^4}{4} - Ba^4 1i \right) + \tan(c + dx)^5(8Ba^4 + Aa^4 8i) - \tan(c + dx)^6(4Aa^4 + Ba^4 4i)}{d \tan(c + dx)^6} - \frac{16a^4 \operatorname{atan}(2 \tan(c + dx) + 1i)(B + A 1i)}{d}$$

input `int(cot(c + d*x)^7*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^4,x)`

output
$$\begin{aligned} & -(\tan(c + d*x)^4*(4*A*a^4 - B*a^4*4i) - \tan(c + d*x)^2*((7*A*a^4)/4 - B*a \\ & ^4*1i) + \tan(c + d*x)^5*(A*a^4*8i + 8*B*a^4) - \tan(c + d*x)^3*((A*a^4*8i)/ \\ & 3 + (7*B*a^4)/3) + (A*a^4)/6 + \tan(c + d*x)*((A*a^4*4i)/5 + (B*a^4)/5))/(d \\ & * \tan(c + d*x)^6) - (16*a^4*\operatorname{atan}(2*\tan(c + d*x) + 1i)*(A*1i + B))/d \end{aligned}$$

3.36 $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

3.36.1	Optimal result	580
3.36.2	Mathematica [A] (verified)	580
3.36.3	Rubi [A] (verified)	581
3.36.4	Maple [A] (verified)	583
3.36.5	Fricas [A] (verification not implemented)	584
3.36.6	Sympy [A] (verification not implemented)	584
3.36.7	Maxima [F(-2)]	585
3.36.8	Giac [A] (verification not implemented)	585
3.36.9	Mupad [B] (verification not implemented)	586

3.36.1 Optimal result

Integrand size = 34, antiderivative size = 129

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{3(iA-B)x}{2a} - \frac{(A+2iB) \log(\cos(c+dx))}{ad} - \frac{3(iA-B) \tan(c+dx)}{2ad} - \frac{(A+2iB) \tan^2(c+dx)}{2ad} + \frac{(iA-B) \tan^3(c+dx)}{2d(a+ia \tan(c+dx))}$$

output $\frac{3}{2}*(I*A-B)*x/a - (A+2*I*B)*\ln(\cos(d*x+c))/a/d - 3/2*(I*A-B)*\tan(d*x+c)/a/d - 1/2*(A+2*I*B)*\tan(d*x+c)^2/a/d + 1/2*(I*A-B)*\tan(d*x+c)^3/d/(a+I*a*\tan(d*x+c))$

3.36.2 Mathematica [A] (verified)

Time = 2.05 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.06

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{\frac{2(2A+iB) \tan^2(c+dx)}{a+ia \tan(c+dx)} + \frac{2B \tan^3(c+dx)}{a+ia \tan(c+dx)} + \frac{(5A+7iB) \log(i-\tan(c+dx)) - (A-iB) \log(i+\tan(c+dx)) + \frac{6(-iA+B)}{-i+\tan(c+dx)}}{a}}{4d}$$

input `Integrate[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `((2*(2*A + I*B)*Tan[c + d*x]^2)/(a + I*a*Tan[c + d*x]) + (2*B*Tan[c + d*x]^3)/(a + I*a*Tan[c + d*x]) + ((5*A + (7*I)*B)*Log[I - Tan[c + d*x]] - (A - I*B)*Log[I + Tan[c + d*x]] + (6*((-I)*A + B))/(-I + Tan[c + d*x]))/a)/(4*d)`

3.36.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.95, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3042, 4078, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

↓ 3042

$$\int \frac{\tan(c+dx)^3(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

↓ 4078

$$\frac{(-B+iA) \tan^3(c+dx)}{2d(a+ia \tan(c+dx))} - \frac{\int \tan^2(c+dx)(3a(iA-B) + 2a(A+2iB) \tan(c+dx)) dx}{2a^2}$$

↓ 3042

$$\frac{(-B+iA) \tan^3(c+dx)}{2d(a+ia \tan(c+dx))} - \frac{\int \tan(c+dx)^2(3a(iA-B) + 2a(A+2iB) \tan(c+dx)) dx}{2a^2}$$

↓ 4011

$$\frac{(-B+iA) \tan^3(c+dx)}{2d(a+ia \tan(c+dx))} - \frac{\int \tan(c+dx)(3a(iA-B) \tan(c+dx) - 2a(A+2iB)) dx + \frac{a(A+2iB) \tan^2(c+dx)}{d}}{2a^2}$$

↓ 3042

$$\frac{(-B+iA) \tan^3(c+dx)}{2d(a+ia \tan(c+dx))} - \frac{\int \tan(c+dx)(3a(iA-B) \tan(c+dx) - 2a(A+2iB)) dx + \frac{a(A+2iB) \tan^2(c+dx)}{d}}{2a^2}$$

3.36. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

$$\begin{array}{c}
\downarrow 4008 \\
\frac{(-B + iA) \tan^3(c + dx)}{2d(a + ia \tan(c + dx))} - \\
\frac{-2a(A + 2iB) \int \tan(c + dx) dx + \frac{a(A+2iB) \tan^2(c+dx)}{d} + \frac{3a(-B+iA) \tan(c+dx)}{d} - 3ax(-B + iA)}{2a^2} \\
\downarrow 3042 \\
\frac{(-B + iA) \tan^3(c + dx)}{2d(a + ia \tan(c + dx))} - \\
\frac{-2a(A + 2iB) \int \tan(c + dx) dx + \frac{a(A+2iB) \tan^2(c+dx)}{d} + \frac{3a(-B+iA) \tan(c+dx)}{d} - 3ax(-B + iA)}{2a^2} \\
\downarrow 3956 \\
\frac{(-B + iA) \tan^3(c + dx)}{2d(a + ia \tan(c + dx))} - \\
\frac{\frac{a(A+2iB) \tan^2(c+dx)}{d} + \frac{3a(-B+iA) \tan(c+dx)}{d} + \frac{2a(A+2iB) \log(\cos(c+dx))}{d} - 3ax(-B + iA)}{2a^2}
\end{array}$$

input `Int[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `((I*A - B)*Tan[c + d*x]^3)/(2*d*(a + I*a*Tan[c + d*x])) - (-3*a*(I*A - B)*x + (2*a*(A + (2*I)*B)*Log[Cos[c + d*x]])/d + (3*a*(I*A - B)*Tan[c + d*x])/d + (a*(A + (2*I)*B)*Tan[c + d*x]^2)/d)/(2*a^2)`

3.36.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

```
rule 4011 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

```
rule 4078 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.36.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.13

method	result
norman	$\frac{(-iA+B)\left(\frac{\tan^3(dx+c)}{ad} - \frac{3(-iA+B)x}{2a} + \frac{2iB+A}{2ad} + \frac{3(-iA+B)\tan(dx+c)}{2ad} - \frac{3(-iA+B)x\tan^2(dx+c)}{2a} - \frac{iB\tan^4(dx+c)}{2ad}\right)}{1+\tan^2(dx+c)} + \frac{2iB}{ad}$
risch	$-\frac{7xB}{2a} + \frac{5ixA}{2a} + \frac{ie^{-2i(dx+c)}B}{4ad} + \frac{e^{-2i(dx+c)}A}{4ad} - \frac{4Bc}{ad} + \frac{2iAc}{ad} + \frac{2i(-iAe^{2i(dx+c)}-iA+B)}{da(e^{2i(dx+c)}+1)^2} - \frac{2i\ln(e^{2i(dx+c)}+1)}{ad}$
derivativedivides	$\frac{B\tan(dx+c)}{da} - \frac{iB(\tan^2(dx+c))}{2da} - \frac{iA\tan(dx+c)}{da} + \frac{A\ln(1+\tan^2(dx+c))}{2da} + \frac{3iA\arctan(\tan(dx+c))}{2da} + \frac{iB\ln(1+\tan^2(dx+c))}{2da}$
default	$\frac{B\tan(dx+c)}{da} - \frac{iB(\tan^2(dx+c))}{2da} - \frac{iA\tan(dx+c)}{da} + \frac{A\ln(1+\tan^2(dx+c))}{2da} + \frac{3iA\arctan(\tan(dx+c))}{2da} + \frac{iB\ln(1+\tan^2(dx+c))}{2da}$

```
input int(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

```
output (1/a/d*(-I*A+B)*tan(d*x+c)^3-3/2/a*(-I*A+B)*x+1/2*(A+2*I*B)/a/d+3/2/a/d*(-
I*A+B)*tan(d*x+c)-3/2/a*(-I*A+B)*x*tan(d*x+c)^2-1/2*I*B/a/d*tan(d*x+c)^4)/
(1+tan(d*x+c)^2)+1/2*(A+2*I*B)/a/d*ln(1+tan(d*x+c)^2)
```

3.36.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.44

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{2(-5iA+7B)dx e^{(6i dx+6i c)} + (4(-5iA+7B)dx - 9A - iB)e^{(4i dx+4i c)} + 2((-5iA+7B)dx - 5A - iB)e^{(2i dx+2i c)}}{4(ade^{(6i dx+6i c)} + 2ade^{(4i dx+4i c)} + ade^{(2i dx+2i c)})}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="fracas")`

output `-1/4*(2*(-5*I*A + 7*B)*d*x*e^(6*I*d*x + 6*I*c) + (4*(-5*I*A + 7*B)*d*x - 9*A - I*B)*e^(4*I*d*x + 4*I*c) + 2*((-5*I*A + 7*B)*d*x - 5*A - 5*I*B)*e^(2*I*d*x + 2*I*c) + 4*((A + 2*I*B)*e^(6*I*d*x + 6*I*c) + 2*(A + 2*I*B)*e^(4*I*d*x + 4*I*c) + (A + 2*I*B)*e^(2*I*d*x + 2*I*c))*log(e^(2*I*d*x + 2*I*c) + 1) - A - I*B)/(a*d*e^(6*I*d*x + 6*I*c) + 2*a*d*e^(4*I*d*x + 4*I*c) + a*d*e^(2*I*d*x + 2*I*c))`

3.36.6 Sympy [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.52

$$\begin{aligned} & \int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx \\ &= \frac{2Ae^{2ic}e^{2idx} + 2A + 2iB}{ade^{4ic}e^{4idx} + 2ade^{2ic}e^{2idx} + ad} \\ &+ \begin{cases} \frac{(A+iB)e^{-2ic}e^{-2idx}}{4ad} & \text{for } ade^{2ic} \neq 0 \\ x \left(-\frac{5iA-7B}{2a} + \frac{(5iAe^{2ic}-iA-7Be^{2ic}+B)e^{-2ic}}{2a} \right) & \text{otherwise} \end{cases} \\ &+ \frac{x(5iA-7B)}{2a} - \frac{(A+2iB) \log(e^{2idx} + e^{-2ic})}{ad} \end{aligned}$$

input `integrate(tan(d*x+c)**3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)`

output `(2*A*exp(2*I*c)*exp(2*I*d*x) + 2*A + 2*I*B)/(a*d*exp(4*I*c)*exp(4*I*d*x) + 2*a*d*exp(2*I*c)*exp(2*I*d*x) + a*d) + Piecewise(((A + I*B)*exp(-2*I*c)*exp(-2*I*d*x)/(4*a*d), Ne(a*d*exp(2*I*c), 0)), (x*(-(5*I*A - 7*B)/(2*a) + (5*I*A*exp(2*I*c) - I*A - 7*B*exp(2*I*c) + B)*exp(-2*I*c)/(2*a)), True)) + x*(5*I*A - 7*B)/(2*a) - (A + 2*I*B)*log(exp(2*I*d*x) + exp(-2*I*c))/(a*d)`

3.36.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.36.8 Giac [A] (verification not implemented)

Time = 0.61 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.95

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \frac{\frac{(A-iB)\log(\tan(dx+c)+i)}{a} - \frac{(5A+7iB)\log(\tan(dx+c)-i)}{a} + \frac{2(iBa\tan(dx+c)^2+2iAa\tan(dx+c)-2Ba\tan(dx+c))}{a^2} + \frac{5A\tan(dx+c)}{4d}}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")
```

```
output -1/4*((A - I*B)*log(tan(d*x + c) + I)/a - (5*A + 7*I*B)*log(tan(d*x + c) - I)/a + 2*(I*B*a*tan(d*x + c)^2 + 2*I*A*a*tan(d*x + c) - 2*B*a*tan(d*x + c))/a^2 + (5*A*tan(d*x + c) + 7*I*B*tan(d*x + c) - 3*I*A + 5*B)/(a*(tan(d*x + c) - I)))/d
```


3.36.9 Mupad [B] (verification not implemented)

Time = 7.48 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.09

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{\frac{A}{2a} - \frac{A+B1i}{2a} + \frac{A+B2i}{2a}}{d(1+\tan(c+dx)1i)} - \frac{\tan(c+dx)\left(-\frac{B}{a} + \frac{A1i}{a}\right)}{d} + \frac{\ln(\tan(c+dx)+1i)(B+A1i)1i}{4ad} + \frac{\ln(\tan(c+dx)-1i)(5A+B7i)}{4ad} - \frac{B \tan(c+dx)^2 1i}{2ad}$$

input `int((tan(c + d*x)^3*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i),x)`output `(A/(2*a) - (A + B*1i)/(2*a) + (A + B*2i)/(2*a))/(d*(tan(c + d*x)*1i + 1)) - (tan(c + d*x)*((A*1i)/a - B/a))/d + (log(tan(c + d*x) + 1i)*(A*1i + B)*1i)/(4*a*d) + (log(tan(c + d*x) - 1i)*(5*A + B*7i))/(4*a*d) - (B*tan(c + d*x)^2*1i)/(2*a*d)`

3.37 $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

3.37.1	Optimal result	587
3.37.2	Mathematica [A] (verified)	587
3.37.3	Rubi [A] (verified)	588
3.37.4	Maple [A] (verified)	590
3.37.5	Fricas [A] (verification not implemented)	590
3.37.6	Sympy [A] (verification not implemented)	591
3.37.7	Maxima [F(-2)]	591
3.37.8	Giac [A] (verification not implemented)	592
3.37.9	Mupad [B] (verification not implemented)	592

3.37.1 Optimal result

Integrand size = 34, antiderivative size = 101

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{(A+3iB)x}{2a} + \frac{(iA-B) \log(\cos(c+dx))}{ad} - \frac{(A+3iB) \tan(c+dx)}{2ad} + \frac{(iA-B) \tan^2(c+dx)}{2d(a+ia \tan(c+dx))}$$

```
output 1/2*(A+3*I*B)*x/a+(I*A-B)*ln(cos(d*x+c))/a/d-1/2*(A+3*I*B)*tan(d*x+c)/a/d+
1/2*(I*A-B)*tan(d*x+c)^2/d/(a+I*a*tan(d*x+c))
```

3.37.2 Mathematica [A] (verified)

Time = 1.39 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.02

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{\frac{4B \tan^2(c+dx)}{a+ia \tan(c+dx)} + \frac{(-3iA+5B) \log(i-\tan(c+dx))-(iA+B) \log(i+\tan(c+dx))-\frac{2(A+3iB)}{-i+\tan(c+dx)}}{a}}{4d}$$

```
input Integrate[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]
```

output $((4*B*\text{Tan}[c + d*x]^2)/(a + I*a*\text{Tan}[c + d*x]) + (((-3*I)*A + 5*B)*\text{Log}[I - \text{Tan}[c + d*x]] - (I*A + B)*\text{Log}[I + \text{Tan}[c + d*x]] - (2*(A + (3*I)*B)))/(-I + \text{Tan}[c + d*x]))/a)/(4*d)$

3.37.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.98, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {3042, 4078, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

↓ 3042

$$\int \frac{\tan(c+dx)^2(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

↓ 4078

$$\frac{(-B+ia) \tan^2(c+dx)}{2d(a+ia \tan(c+dx))} - \frac{\int \tan(c+dx)(2a(iA-B) + a(A+3iB) \tan(c+dx)) dx}{2a^2}$$

↓ 3042

$$\frac{(-B+ia) \tan^2(c+dx)}{2d(a+ia \tan(c+dx))} - \frac{\int \tan(c+dx)(2a(iA-B) + a(A+3iB) \tan(c+dx)) dx}{2a^2}$$

↓ 4008

$$\frac{(-B+ia) \tan^2(c+dx)}{2d(a+ia \tan(c+dx))} - \frac{2a(-B+ia) \int \tan(c+dx) dx + \frac{a(A+3iB) \tan(c+dx)}{d} - ax(A+3iB)}{2a^2}$$

↓ 3042

$$\frac{(-B+ia) \tan^2(c+dx)}{2d(a+ia \tan(c+dx))} - \frac{2a(-B+ia) \int \tan(c+dx) dx + \frac{a(A+3iB) \tan(c+dx)}{d} - ax(A+3iB)}{2a^2}$$

↓ 3956

$$\frac{(-B+ia) \tan^2(c+dx)}{2d(a+ia \tan(c+dx))} - \frac{\frac{a(A+3iB) \tan(c+dx)}{d} - \frac{2a(-B+ia) \log(\cos(c+dx))}{d} - ax(A+3iB)}{2a^2}$$

input $\text{Int}[(\text{Tan}[c + d*x]^2*(A + B*\text{Tan}[c + d*x]))/(a + I*a*\text{Tan}[c + d*x]), x]$

3.37. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

```
output ((I*A - B)*Tan[c + d*x]^2)/(2*d*(a + I*a*Tan[c + d*x])) - (-a*(A + (3*I)*
B)*x) - (2*a*(I*A - B)*Log[Cos[c + d*x]])/d + (a*(A + (3*I)*B)*Tan[c + d*x
])/d)/(2*a^2)
```

3.37.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4008 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_.)]), x_Symbol] :> Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f),
x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e,
f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]
```

```
rule 4078 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] :> Sim
p[(-(A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.37.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.24

method	result
norman	$\frac{(3iB+A)x}{2a} + \frac{-iA+B}{2ad} - \frac{(3iB+A)\tan(dx+c)}{2ad} + \frac{(3iB+A)x(\tan^2(dx+c))}{2a} - \frac{iB(\tan^3(dx+c))}{ad} + \frac{(-iA+B)\ln(1+\tan^2(dx+c))}{2ad}$
derivativedivides	$-\frac{iB\tan(dx+c)}{da} - \frac{A}{2da(\tan(dx+c)-i)} - \frac{iB}{2da(\tan(dx+c)-i)} - \frac{iA\ln(1+\tan^2(dx+c))}{2da} + \frac{A\arctan(\tan(dx+c))}{2da} +$
default	$-\frac{iB\tan(dx+c)}{da} - \frac{A}{2da(\tan(dx+c)-i)} - \frac{iB}{2da(\tan(dx+c)-i)} - \frac{iA\ln(1+\tan^2(dx+c))}{2da} + \frac{A\arctan(\tan(dx+c))}{2da} +$
risch	$\frac{5ixB}{2a} + \frac{3xA}{2a} + \frac{e^{-2i(dx+c)}B}{4ad} - \frac{ie^{-2i(dx+c)}A}{4ad} + \frac{2iBc}{ad} + \frac{2Ac}{ad} + \frac{2B}{da(e^{2i(dx+c)}+1)} - \frac{\ln(e^{2i(dx+c)}+1)B}{ad} + i$

input `int(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output $(1/2*(A+3*I*B)*x/a+1/2/a/d*(-I*A+B)-1/2*(A+3*I*B)*\tan(d*x+c)/a/d+1/2*(A+3*I*B)/a*x*\tan(d*x+c)^2-I*B/a/d*\tan(d*x+c)^3)/(1+\tan(d*x+c)^2)+1/2/a/d*(-I*A+B)*\ln(1+\tan(d*x+c)^2)$

3.37.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.26

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$$

$$= \frac{2(3A+5iB)dx e^{(4i dx+4i c)} + (2(3A+5iB)dx - iA+9B)e^{(2i dx+2i c)} - 4((-iA+B)e^{(4i dx+4i c)} + (-iA+9B)e^{(2i dx+2i c)})}{4(ade^{(4i dx+4i c)} + ade^{(2i dx+2i c)})}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="fracas")`

output $1/4*(2*(3*A+5*I*B)*d*x*e^{(4*I*d*x+4*I*c)} + (2*(3*A+5*I*B)*d*x - I*A+9*B)*e^{(2*I*d*x+2*I*c)} - 4*((-I*A+B)*e^{(4*I*d*x+4*I*c)} + (-I*A+9*B)*e^{(2*I*d*x+2*I*c)})*\log(e^{(2*I*d*x+2*I*c)}+1) - I*A+B)/(a*d*e^{(4*I*d*x+4*I*c)} + a*d*e^{(2*I*d*x+2*I*c)})$

3.37.6 Sympy [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.50

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= \frac{2B}{ade^{2ic}e^{2idx} + ad} + \begin{cases} \frac{(-iA+B)e^{-2ic}e^{-2idx}}{4ad} & \text{for } ade^{2ic} \neq 0 \\ x \left(-\frac{3A+5iB}{2a} + \frac{(3Ae^{2ic}-A+5iBe^{2ic}-iB)e^{-2ic}}{2a} \right) & \text{otherwise} \end{cases}$$

$$+ \frac{x(3A+5iB)}{2a} + \frac{i(A+iB) \log(e^{2idx} + e^{-2ic})}{ad}$$

```
input integrate(tan(d*x+c)**2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)
```

```
output 2*B/(a*d*exp(2*I*c)*exp(2*I*d*x) + a*d) + Piecewise((( -I*A + B)*exp(-2*I*c)
)*exp(-2*I*d*x)/(4*a*d), Ne(a*d*exp(2*I*c), 0)), (x*(-(3*A + 5*I*B)/(2*a)
+ (3*A*exp(2*I*c) - A + 5*I*B*exp(2*I*c) - I*B)*exp(-2*I*c)/(2*a)), True))
+ x*(3*A + 5*I*B)/(2*a) + I*(A + I*B)*log(exp(2*I*d*x) + exp(-2*I*c))/(a*
d)
```

3.37.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="m
axima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.37.8 Giac [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.98

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$$

$$= \frac{\frac{(-iA-B)\log(\tan(dx+c)+i)}{a} - \frac{(3iA-5B)\log(\tan(dx+c)-i)}{a} - \frac{4iB\tan(dx+c)}{a} - \frac{-3iA\tan(dx+c)+5B\tan(dx+c)-A-3iB}{a(\tan(dx+c)-i)}}{4d}$$

```
input integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")
```

```
output 1/4*((-I*A - B)*log(tan(d*x + c) + I)/a - (3*I*A - 5*B)*log(tan(d*x + c) - I)/a - 4*I*B*tan(d*x + c)/a - (-3*I*A*tan(d*x + c) + 5*B*tan(d*x + c) - A - 3*I*B)/(a*(tan(d*x + c) - I)))/d
```

3.37.9 Mupad [B] (verification not implemented)

Time = 7.77 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.94

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = -\frac{\ln(\tan(c+dx)+i)(B+A1i)}{4ad} - \frac{B\tan(c+dx)1i}{ad} - \frac{(A+B1i)1i}{2ad(1+\tan(c+dx)1i)} - \frac{\ln(\tan(c+dx)-i)(-5B+A3i)}{4ad}$$

```
input int((tan(c + d*x)^2*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i),x)
```

```
output - (log(tan(c + d*x) + 1i)*(A*1i + B))/(4*a*d) - (B*tan(c + d*x)*1i)/(a*d) - ((A + B*1i)*1i)/(2*a*d*(tan(c + d*x)*1i + 1)) - (log(tan(c + d*x) - 1i)*(A*3i - 5*B))/(4*a*d)
```

3.38 $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

3.38.1	Optimal result	593
3.38.2	Mathematica [A] (verified)	593
3.38.3	Rubi [A] (verified)	594
3.38.4	Maple [A] (verified)	596
3.38.5	Fricas [A] (verification not implemented)	596
3.38.6	Sympy [A] (verification not implemented)	597
3.38.7	Maxima [F(-2)]	597
3.38.8	Giac [A] (verification not implemented)	598
3.38.9	Mupad [B] (verification not implemented)	598

3.38.1 Optimal result

Integrand size = 32, antiderivative size = 67

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = -\frac{(iA-B)x}{2a} + \frac{iB \log(\cos(c+dx))}{ad} - \frac{A+iB}{2ad(1+i \tan(c+dx))}$$

output `-1/2*(I*A-B)*x/a+I*B*ln(cos(d*x+c))/a/d+1/2*(-A-I*B)/a/d/(1+I*tan(d*x+c))`

3.38.2 Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.12

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{-((A+3iB) \log(i-\tan(c+dx))) + (A-iB) \log(i+\tan(c+dx)) + \frac{2i(A+iB)}{-i+\tan(c+dx)}}{4ad}$$

input `Integrate[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `-((A + (3*I)*B)*Log[I - Tan[c + d*x]]) + (A - I*B)*Log[I + Tan[c + d*x]] + ((2*I)*(A + I*B))/(-I + Tan[c + d*x])/(4*a*d)`

3.38.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.96, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.219$, Rules used = {3042, 4072, 27, 3042, 3956, 4009, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx \\
 & \quad \downarrow \text{4072} \\
 & -\frac{i\int \frac{(iA-B)\tan(c+dx)}{i\tan(c+dx)+1} dx}{a} - \frac{iB\int \tan(c+dx) dx}{a} \\
 & \quad \downarrow \text{27} \\
 & -\frac{i(-B+iA)\int \frac{\tan(c+dx)}{i\tan(c+dx)+1} dx}{a} - \frac{iB\int \tan(c+dx) dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{i(-B+iA)\int \frac{\tan(c+dx)}{i\tan(c+dx)+1} dx}{a} - \frac{iB\int \tan(c+dx) dx}{a} \\
 & \quad \downarrow \text{3956} \\
 & \frac{iB\log(\cos(c+dx))}{ad} - \frac{i(-B+iA)\int \frac{\tan(c+dx)}{i\tan(c+dx)+1} dx}{a} \\
 & \quad \downarrow \text{4009} \\
 & \frac{iB\log(\cos(c+dx))}{ad} - \frac{i(-B+iA)\left(-\frac{i\int 1dx}{2} - \frac{1}{2d(1+i\tan(c+dx))}\right)}{a} \\
 & \quad \downarrow \text{24} \\
 & \frac{iB\log(\cos(c+dx))}{ad} - \frac{i(-B+iA)\left(-\frac{1}{2d(1+i\tan(c+dx))} - \frac{ix}{2}\right)}{a}
 \end{aligned}$$

input `Int[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

3.38. $\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$

```
output (I*B*Log[Cos[c + d*x]])/(a*d) - (I*(I*A - B)*((-1/2*I)*x - 1/(2*d*(1 + I*T
an[c + d*x]))))/a
```

3.38.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4009 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a
*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x
], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2
, 0] && LtQ[m, 0]
```

```
rule 4072 Int[(((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_
.)*(x_)]))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*(d/
b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c
- a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d
, e, f, A, B}, x] && NeQ[b*c - a*d, 0]
```

3.38.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.28

method	result
risch	$\frac{3xB}{2a} - \frac{ixA}{2a} - \frac{ie^{-2i(dx+c)}B}{4ad} - \frac{e^{-2i(dx+c)}A}{4ad} + \frac{2Bc}{ad} + \frac{i \ln(e^{2i(dx+c)}+1)B}{ad}$
derivativdivides	$-\frac{iA \arctan(\tan(dx+c))}{2da} - \frac{iB \ln(1+\tan^2(dx+c))}{2da} + \frac{B \arctan(\tan(dx+c))}{2da} + \frac{iA}{2da(\tan(dx+c)-i)} - \frac{B}{2da(\tan(dx+c)+i)}$
default	$-\frac{iA \arctan(\tan(dx+c))}{2da} - \frac{iB \ln(1+\tan^2(dx+c))}{2da} + \frac{B \arctan(\tan(dx+c))}{2da} + \frac{iA}{2da(\tan(dx+c)-i)} - \frac{B}{2da(\tan(dx+c)+i)}$
norman	$\frac{(-iA+B)x}{2a} - \frac{iB+A}{2ad} - \frac{(-iA+B) \tan(dx+c)}{2ad} + \frac{(-iA+B)x(\tan^2(dx+c))}{2a} - \frac{iB \ln(1+\tan^2(dx+c))}{2da}$

input `int(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `3/2*x/a*B-1/2*I*x/a*A-1/4*I/a/d*exp(-2*I*(d*x+c))*B-1/4/a/d*exp(-2*I*(d*x+c))*A+2/a/d*B*c+I*B/a/d*ln(exp(2*I*(d*x+c))+1)`

3.38.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.99

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= -\frac{(2(iA-3B)dx e^{(2i dx+2i c)} - 4i B e^{(2i dx+2i c)} \log(e^{(2i dx+2i c)}+1) + A+iB)e^{(-2i dx-2i c)}}{4ad}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="fricas")`

output `-1/4*(2*(I*A - 3*B)*d*x*e^(2*I*d*x + 2*I*c) - 4*I*B*e^(2*I*d*x + 2*I*c))*log(e^(2*I*d*x + 2*I*c) + 1) + A + I*B)*e^(-2*I*d*x - 2*I*c)/(a*d)`

3.38.6 Sympy [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.78

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$$

$$= \frac{iB \log(e^{2idx} + e^{-2ic})}{ad} + \begin{cases} \frac{(-A-iB)e^{-2ic}e^{-2idx}}{4ad} & \text{for } ade^{2ic} \neq 0 \\ x\left(-\frac{iA+3B}{2a} + \frac{(-iAe^{2ic}+iA+3Be^{2ic}-B)e^{-2ic}}{2a}\right) & \text{otherwise} \end{cases} + \frac{x(-iA+3B)}{2a}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)`output `I*B*log(exp(2*I*d*x) + exp(-2*I*c))/(a*d) + Piecewise(((-A - I*B)*exp(-2*I*c)*exp(-2*I*d*x)/(4*a*d), Ne(a*d*exp(2*I*c), 0)), (x*(-(-I*A + 3*B)/(2*a) + (-I*A*exp(2*I*c) + I*A + 3*B*exp(2*I*c) - B)*exp(-2*I*c)/(2*a)), True)) + x*(-I*A + 3*B)/(2*a)`**3.38.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="maxima")`output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.38.8 Giac [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.18

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$$

$$= \frac{\frac{(A-iB)\log(\tan(dx+c)+i)}{a} - \frac{(A+3iB)\log(\tan(dx+c)-i)}{a} + \frac{A\tan(dx+c)+3iB\tan(dx+c)+iA+B}{a(\tan(dx+c)-i)}}{4d}$$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")
```

```
output 1/4*((A - I*B)*log(tan(d*x + c) + I)/a - (A + 3*I*B)*log(tan(d*x + c) - I)/a + (A*tan(d*x + c) + 3*I*B*tan(d*x + c) + I*A + B)/(a*(tan(d*x + c) - I)))/d
```

3.38.9 Mupad [B] (verification not implemented)

Time = 7.83 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.21

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = -\frac{\frac{A}{2a} + \frac{B1i}{2a}}{d(1+\tan(c+dx)1i)} + \frac{\ln(\tan(c+dx)+1i)(A-B1i)}{4ad} - \frac{\ln(\tan(c+dx)-i)(A+B3i)}{4ad}$$

```
input int((tan(c + d*x)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i),x)
```

```
output (log(tan(c + d*x) + 1i)*(A - B*1i))/(4*a*d) - (A/(2*a) + (B*1i)/(2*a))/(d*(tan(c + d*x)*1i + 1)) - (log(tan(c + d*x) - 1i)*(A + B*3i))/(4*a*d)
```

3.39 $\int \frac{A+B \tan(c+dx)}{a+ia \tan(c+dx)} dx$

3.39.1	Optimal result	599
3.39.2	Mathematica [A] (verified)	599
3.39.3	Rubi [A] (verified)	600
3.39.4	Maple [A] (verified)	601
3.39.5	Fricas [A] (verification not implemented)	601
3.39.6	Sympy [A] (verification not implemented)	602
3.39.7	Maxima [F(-2)]	602
3.39.8	Giac [B] (verification not implemented)	602
3.39.9	Mupad [B] (verification not implemented)	603

3.39.1 Optimal result

Integrand size = 26, antiderivative size = 47

$$\int \frac{A + B \tan(c + dx)}{a + ia \tan(c + dx)} dx = \frac{(A - iB)x}{2a} + \frac{iA - B}{2d(a + ia \tan(c + dx))}$$

output `1/2*(A-I*B)*x/a+1/2*(I*A-B)/d/(a+I*a*tan(d*x+c))`

3.39.2 Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.19

$$\int \frac{A + B \tan(c + dx)}{a + ia \tan(c + dx)} dx = \frac{(A - iB) \arctan(\tan(c + dx))}{2ad} - \frac{A + iB}{2ad(i - \tan(c + dx))}$$

input `Integrate[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x]),x]`

output `((A - I*B)*ArcTan[Tan[c + d*x]])/(2*a*d) - (A + I*B)/(2*a*d*(I - Tan[c + d*x]))`

3.39.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {3042, 4009, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{A + B \tan(c + dx)}{a + ia \tan(c + dx)} dx \\
 \downarrow 3042 \\
 \int \frac{A + B \tan(c + dx)}{a + ia \tan(c + dx)} dx \\
 \downarrow 4009 \\
 \frac{(A - iB) \int 1 dx}{2a} + \frac{-B + iA}{2d(a + ia \tan(c + dx))} \\
 \downarrow 24 \\
 \frac{-B + iA}{2d(a + ia \tan(c + dx))} + \frac{x(A - iB)}{2a}
 \end{array}$$

input `Int[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x]),x]`

output `((A - I*B)*x)/(2*a) + (I*A - B)/(2*d*(a + I*a*Tan[c + d*x]))`

3.39.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4009 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a
*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x
], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2
, 0] && LtQ[m, 0]
```

3.39.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.15

method	result	size
risch	$-\frac{i x B}{2 a} + \frac{x A}{2 a} - \frac{e^{-2 i(d x+c)} B}{4 a d} + \frac{i e^{-2 i(d x+c)} A}{4 a d}$	54
derivativedivides	$\frac{A}{2 d a(\tan(d x+c)-i)} + \frac{i B}{2 d a(\tan(d x+c)-i)} + \frac{A \arctan(\tan(d x+c))}{2 d a} - \frac{i B \arctan(\tan(d x+c))}{2 d a}$	76
default	$\frac{A}{2 d a(\tan(d x+c)-i)} + \frac{i B}{2 d a(\tan(d x+c)-i)} + \frac{A \arctan(\tan(d x+c))}{2 d a} - \frac{i B \arctan(\tan(d x+c))}{2 d a}$	76
norman	$\frac{(-i B+A) x}{2 a} - \frac{-i A+B}{2 a d} + \frac{(-i B+A) x(\tan^2(d x+c))}{2 a} + \frac{(i B+A) \tan(d x+c)}{2 a d}$ $\frac{1}{1+\tan^2(d x+c)}$	81

```
input int((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output -1/2*I*x/a*B+1/2*x/a*A-1/4/a/d*exp(-2*I*(d*x+c))*B+1/4*I/a/d*exp(-2*I*(d*x
+c))*A
```

3.39.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.89

$$\int \frac{A + B \tan(c + dx)}{a + i a \tan(c + dx)} dx = \frac{(2(A - i B) dx e^{(2i dx + 2i c)} + i A - B) e^{(-2i dx - 2i c)}}{4 a d}$$

```
input integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="fricas")
```

```
output 1/4*(2*(A - I*B)*d*x*e^(2*I*d*x + 2*I*c) + I*A - B)*e^(-2*I*d*x - 2*I*c)/(
a*d)
```


3.39.6 Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.85

$$\int \frac{A + B \tan(c + dx)}{a + ia \tan(c + dx)} dx$$

$$= \begin{cases} \frac{(iA-B)e^{-2ic}e^{-2idx}}{4ad} & \text{for } ade^{2ic} \neq 0 \\ x \left(-\frac{A-iB}{2a} + \frac{(Ae^{2ic}+A-iBe^{2ic}+iB)e^{-2ic}}{2a} \right) & \text{otherwise} \end{cases} + \frac{x(A-iB)}{2a}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)`output `Piecewise(((I*A - B)*exp(-2*I*c)*exp(-2*I*d*x)/(4*a*d), Ne(a*d*exp(2*I*c), 0)), (x*(-(A - I*B)/(2*a) + (A*exp(2*I*c) + A - I*B*exp(2*I*c) + I*B)*exp(-2*I*c)/(2*a)), True)) + x*(A - I*B)/(2*a)`**3.39.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{a + ia \tan(c + dx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="maxima")`output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`**3.39.8 Giac [B] (verification not implemented)**Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 83 vs. $2(35) = 70$.

Time = 0.34 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.77

$$\int \frac{A + B \tan(c + dx)}{a + ia \tan(c + dx)} dx$$

$$= -\frac{\frac{(-iA-B)\log(\tan(dx+c)+i)}{a} + \frac{(iA+B)\log(\tan(dx+c)-i)}{a} + \frac{-iA\tan(dx+c)-B\tan(dx+c)-3A-iB}{a(\tan(dx+c)-i)}}{4d}$$

3.39. $\int \frac{A+B \tan(c+dx)}{a+ia \tan(c+dx)} dx$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")`

output `-1/4*((-I*A - B)*log(tan(d*x + c) + I)/a + (I*A + B)*log(tan(d*x + c) - I)/a + (-I*A*tan(d*x + c) - B*tan(d*x + c) - 3*A - I*B)/(a*(tan(d*x + c) - I)))/d`

3.39.9 Mupad [B] (verification not implemented)

Time = 7.15 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.96

$$\int \frac{A + B \tan(c + dx)}{a + ia \tan(c + dx)} dx = \frac{-\frac{B}{2a} + \frac{A i}{2a}}{d (1 + \tan(c + dx) i)} - \frac{x (B + A i) i}{2 a}$$

input `int((A + B*tan(c + d*x))/(a + a*tan(c + d*x)*1i),x)`

output `((A*1i)/(2*a) - B/(2*a))/(d*(tan(c + d*x)*1i + 1)) - (x*(A*1i + B)*1i)/(2*a)`

3.40 $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

3.40.1	Optimal result	604
3.40.2	Mathematica [A] (verified)	604
3.40.3	Rubi [A] (verified)	605
3.40.4	Maple [A] (verified)	607
3.40.5	Fricas [A] (verification not implemented)	607
3.40.6	Sympy [A] (verification not implemented)	608
3.40.7	Maxima [F(-2)]	608
3.40.8	Giac [A] (verification not implemented)	609
3.40.9	Mupad [B] (verification not implemented)	609

3.40.1 Optimal result

Integrand size = 32, antiderivative size = 62

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = -\frac{(iA-B)x}{2a} + \frac{A \log(\sin(c+dx))}{ad} + \frac{A+iB}{2d(a+ia \tan(c+dx))}$$

output `-1/2*(I*A-B)*x/a+A*ln(sin(d*x+c))/a/d+1/2*(A+I*B)/d/(a+I*a*tan(d*x+c))`

3.40.2 Mathematica [A] (verified)

Time = 0.82 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.39

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{(3A+iB) \log(i-\tan(c+dx)) - 4A \log(\tan(c+dx)) + (A-iB) \log(i+\tan(c+dx)) + \frac{2i(A+iB)}{-i+\tan(c+dx)}}{4ad}$$

input `Integrate[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `-1/4*((3*A + I*B)*Log[I - Tan[c + d*x]] - 4*A*Log[Tan[c + d*x]] + (A - I*B)*Log[I + Tan[c + d*x]] + ((2*I)*(A + I*B))/(-I + Tan[c + d*x]))/(a*d)`

3.40.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.08, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.219$, Rules used = {3042, 4079, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)(a+ia \tan(c+dx))} dx \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \cot(c+dx)(2aA-a(iA-B) \tan(c+dx)) dx}{2a^2} + \frac{A+iB}{2d(a+ia \tan(c+dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{2aA-a(iA-B) \tan(c+dx)}{\tan(c+dx)} dx}{2a^2} + \frac{A+iB}{2d(a+ia \tan(c+dx))} \\
 & \quad \downarrow \text{4014} \\
 & \frac{2aA \int \cot(c+dx) dx - ax(-B+iA)}{2a^2} + \frac{A+iB}{2d(a+ia \tan(c+dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2aA \int -\tan(c+dx+\frac{\pi}{2}) dx - ax(-B+iA)}{2a^2} + \frac{A+iB}{2d(a+ia \tan(c+dx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{-2aA \int \tan(\frac{1}{2}(2c+\pi)+dx) dx - ax(-B+iA)}{2a^2} + \frac{A+iB}{2d(a+ia \tan(c+dx))} \\
 & \quad \downarrow \text{3956} \\
 & \frac{\frac{2aA \log(-\sin(c+dx))}{d} - ax(-B+iA)}{2a^2} + \frac{A+iB}{2d(a+ia \tan(c+dx))}
 \end{aligned}$$

input `Int[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

3.40. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

output $(-(a*(I*A - B)*x) + (2*a*A*Log[-Sin[c + d*x]])/d)/(2*a^2) + (A + I*B)/(2*d*(a + I*a*Tan[c + d*x]))$

3.40.3.1 Defintions of rubi rules used

rule 25 $Int[-(Fx_), x_Symbol] \rightarrow Simp[Identity[-1] Int[Fx, x], x]$

rule 3042 $Int[u_, x_Symbol] \rightarrow Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]$

rule 3956 $Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]$

rule 4014 $Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] \rightarrow Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] \&\& NeQ[b*c - a*d, 0] \&\& NeQ[a^2 + b^2, 0] \&\& NeQ[a*c + b*d, 0]$

rule 4079 $Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^m*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)]*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])^n, x_Symbol] \rightarrow Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] \&\& NeQ[b*c - a*d, 0] \&\& EqQ[a^2 + b^2, 0] \&\& LtQ[m, 0] \&\& !GtQ[n, 0]$

3.40.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.37

method	result
risch	$\frac{xB}{2a} - \frac{3ixA}{2a} + \frac{ie^{-2i(dx+c)}B}{4ad} + \frac{e^{-2i(dx+c)}A}{4ad} - \frac{2iAc}{ad} + \frac{A \ln(e^{2i(dx+c)}-1)}{ad}$
derivativedivides	$-\frac{A \ln(1+\tan^2(dx+c))}{2da} - \frac{iA \arctan(\tan(dx+c))}{2da} + \frac{B \arctan(\tan(dx+c))}{2da} - \frac{iA}{2da(\tan(dx+c)-i)} + \frac{B}{2da(\tan(dx+c)-i)}$
default	$-\frac{A \ln(1+\tan^2(dx+c))}{2da} - \frac{iA \arctan(\tan(dx+c))}{2da} + \frac{B \arctan(\tan(dx+c))}{2da} - \frac{iA}{2da(\tan(dx+c)-i)} + \frac{B}{2da(\tan(dx+c)-i)}$
norman	$\frac{\frac{iB+A}{2ad} + \frac{(-iA+B)x}{2a} + \frac{(-iA+B)\tan(dx+c)}{2ad} + \frac{(-iA+B)x(\tan^2(dx+c))}{2a}}{1+\tan^2(dx+c)} + \frac{A \ln(\tan(dx+c))}{ad} - \frac{A \ln(1+\tan^2(dx+c))}{2da}$

input `int(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/2*x/a*B-3/2*I*x/a*A+1/4*I/a/d*exp(-2*I*(d*x+c))*B+1/4/a/d*exp(-2*I*(d*x+c))*A-2*I*A/a/d*c+A/a/d*ln(exp(2*I*(d*x+c))-1)`

3.40.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.10

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= -\frac{(2(3iA-B)dx e^{(2i dx+2i c)} - 4A e^{(2i dx+2i c)} \log(e^{(2i dx+2i c)} - 1) - A - iB) e^{(-2i dx-2i c)}}{4ad}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="fricas")`

output `-1/4*(2*(3*I*A - B)*d*x*e^(2*I*d*x + 2*I*c) - 4*A*e^(2*I*d*x + 2*I*c)*log(e^(2*I*d*x + 2*I*c) - 1) - A - I*B)*e^(-2*I*d*x - 2*I*c)/(a*d)`

3.40.6 Sympy [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.87

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$$

$$= \frac{A \log(e^{2idx} - e^{-2ic})}{ad} + \begin{cases} \frac{(A+IB)e^{-2ic}e^{-2idx}}{4ad} & \text{for } ade^{2ic} \neq 0 \\ x\left(-\frac{3iA+B}{2a} + \frac{(-3iAe^{2ic}-iA+Be^{2ic}+B)e^{-2ic}}{2a}\right) & \text{otherwise} \end{cases} + \frac{x(-3iA+B)}{2a}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)`

output `A*log(exp(2*I*d*x) - exp(-2*I*c))/(a*d) + Piecewise(((A + I*B)*exp(-2*I*c)*exp(-2*I*d*x)/(4*a*d), Ne(a*d*exp(2*I*c), 0)), (x*(-(-3*I*A + B)/(2*a) + (-3*I*A*exp(2*I*c) - I*A + B*exp(2*I*c) + B)*exp(-2*I*c)/(2*a)), True)) + x*(-3*I*A + B)/(2*a)`

3.40.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.40.8 Giac [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.56

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{\frac{(A-iB) \log(\tan(dx+c)+i)}{a} + \frac{(3A+iB) \log(\tan(dx+c)-i)}{a} - \frac{4A \log(\tan(dx+c))}{a} - \frac{3A \tan(dx+c)+iB \tan(dx+c)-5iA+3B}{a(\tan(dx+c)-i)}}{4d}$$

```
input integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")
```

```
output -1/4*((A - I*B)*log(tan(d*x + c) + I)/a + (3*A + I*B)*log(tan(d*x + c) - I)/a - 4*A*log(tan(d*x + c))/a - (3*A*tan(d*x + c) + I*B*tan(d*x + c) - 5*I*A + 3*B)/(a*(tan(d*x + c) - I)))/d
```

3.40.9 Mupad [B] (verification not implemented)

Time = 7.98 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.58

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{\frac{A}{2a} + \frac{B1i}{2a}}{d(1 + \tan(c+dx)1i)} + \frac{A \ln(\tan(c+dx))}{ad} + \frac{\ln(\tan(c+dx) + 1i)(B + A1i)1i}{4ad} - \frac{\ln(\tan(c+dx) - i)(3A + B1i)}{4ad}$$

```
input int((cot(c + d*x)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i),x)
```

```
output (A/(2*a) + (B*1i)/(2*a))/(d*(tan(c + d*x)*1i + 1)) + (A*log(tan(c + d*x)))/(a*d) + (log(tan(c + d*x) + 1i)*(A*1i + B)*1i)/(4*a*d) - (log(tan(c + d*x) - 1i)*(3*A + B*1i))/(4*a*d)
```


$$3.41 \quad \int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

3.41.1	Optimal result	610
3.41.2	Mathematica [C] (verified)	610
3.41.3	Rubi [A] (verified)	611
3.41.4	Maple [A] (verified)	613
3.41.5	Fricas [A] (verification not implemented)	614
3.41.6	Sympy [A] (verification not implemented)	614
3.41.7	Maxima [F(-2)]	615
3.41.8	Giac [A] (verification not implemented)	615
3.41.9	Mupad [B] (verification not implemented)	616

3.41.1 Optimal result

Integrand size = 34, antiderivative size = 102

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = -\frac{(3A+iB)x}{2a} - \frac{(3A+iB) \cot(c+dx)}{2ad} - \frac{(iA-B) \log(\sin(c+dx))}{ad} + \frac{(A+iB) \cot(c+dx)}{2d(a+ia \tan(c+dx))}$$

output `-1/2*(3*A+I*B)*x/a-1/2*(3*A+I*B)*cot(d*x+c)/a/d-(I*A-B)*ln(sin(d*x+c))/a/d+1/2*(A+I*B)*cot(d*x+c)/d/(a+I*a*tan(d*x+c))`

3.41.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.26 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.96

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{\frac{(A+iB) \cot^2(c+dx)}{i+\cot(c+dx)} - (3A+iB) \cot(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(c+dx)\right) + 2(-iA+B) \log(\sin(c+dx))}{2ad}$$

input `Integrate[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `((A + I*B)*Cot[c + d*x]^2)/(I + Cot[c + d*x]) - (3*A + I*B)*Cot[c + d*x]*
Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2] + 2*((-I)*A + B)*(Log[Cos
[c + d*x]] + Log[Tan[c + d*x]])/(2*a*d)`

3.41.3 Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3042, 4079, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{a + ia \tan(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^2(a + ia \tan(c + dx))} dx \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \cot^2(c + dx)(a(3A + iB) - 2a(iA - B) \tan(c + dx)) dx}{2a^2} + \frac{(A + iB) \cot(c + dx)}{2d(a + ia \tan(c + dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a(3A + iB) - 2a(iA - B) \tan(c + dx)}{\tan(c + dx)^2} dx}{2a^2} + \frac{(A + iB) \cot(c + dx)}{2d(a + ia \tan(c + dx))} \\
 & \quad \downarrow \text{4012} \\
 & \frac{\int -\cot(c + dx)(2a(iA - B) + a(3A + iB) \tan(c + dx)) dx - \frac{a(3A + iB) \cot(c + dx)}{d}}{2a^2} + \frac{(A + iB) \cot(c + dx)}{2d(a + ia \tan(c + dx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{-\int \cot(c + dx)(2a(iA - B) + a(3A + iB) \tan(c + dx)) dx - \frac{a(3A + iB) \cot(c + dx)}{d}}{2a^2} + \frac{(A + iB) \cot(c + dx)}{2d(a + ia \tan(c + dx))}
 \end{aligned}$$

3.41. $\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{a + ia \tan(c + dx)} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& - \int \frac{2a(iA-B)+a(3A+iB)\tan(c+dx)}{\tan(c+dx)} dx - \frac{a(3A+iB)\cot(c+dx)}{d} + \frac{(A+iB)\cot(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow \text{4014} \\
& \frac{-2a(-B+iA)\int\cot(c+dx)dx - \frac{a(3A+iB)\cot(c+dx)}{d} - ax(3A+iB)}{2a^2} + \frac{(A+iB)\cot(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow \text{3042} \\
& \frac{-2a(-B+iA)\int-\tan(c+dx+\frac{\pi}{2})dx - \frac{a(3A+iB)\cot(c+dx)}{d} - ax(3A+iB)}{2a^2} + \frac{(A+iB)\cot(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow \text{25} \\
& \frac{2a(-B+iA)\int\tan(\frac{1}{2}(2c+\pi)+dx)dx - \frac{a(3A+iB)\cot(c+dx)}{d} - ax(3A+iB)}{2a^2} + \frac{(A+iB)\cot(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow \text{3956} \\
& \frac{-\frac{a(3A+iB)\cot(c+dx)}{d} - \frac{2a(-B+iA)\log(-\sin(c+dx))}{d} - ax(3A+iB)}{2a^2} + \frac{(A+iB)\cot(c+dx)}{2d(a+ia\tan(c+dx))}
\end{aligned}$$

input `Int[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `(-(a*(3*A + I*B)*x) - (a*(3*A + I*B)*Cot[c + d*x])/d - (2*a*(I*A - B)*Log[-Sin[c + d*x]])/d)/(2*a^2) + ((A + I*B)*Cot[c + d*x])/(2*d*(a + I*a*Tan[c + d*x]))`

3.41.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.41.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.37

method	result
risch	$-\frac{3ixB}{2a} - \frac{5xA}{2a} + \frac{e^{-2i(dx+c)}B}{4ad} - \frac{ie^{-2i(dx+c)}A}{4ad} - \frac{2iBc}{ad} - \frac{2Ac}{ad} - \frac{2iA}{ad(e^{2i(dx+c)}-1)} + \frac{\ln(e^{2i(dx+c)}-1)B}{ad}$
norman	$-\frac{A}{ad} - \frac{(iB+3A)(\tan^2(dx+c))}{2ad} - \frac{(iB+3A)x \tan(dx+c)}{2a \tan(dx+c)(1+\tan^2(dx+c))} - \frac{(iB+3A)x(\tan^3(dx+c))}{2a} + \frac{(-iA+B)\tan(dx+c)}{2ad} + \frac{(-iA+B)\ln(\tan(dx+c))}{ad}$
derivativedivides	$-\frac{A}{2da(\tan(dx+c)-i)} - \frac{iB}{2da(\tan(dx+c)-i)} + \frac{iA \ln(1+\tan^2(dx+c))}{2da} - \frac{3A \arctan(\tan(dx+c))}{2da} - \frac{B \ln(1+\tan^2(dx+c))}{2da}$
default	$-\frac{A}{2da(\tan(dx+c)-i)} - \frac{iB}{2da(\tan(dx+c)-i)} + \frac{iA \ln(1+\tan^2(dx+c))}{2da} - \frac{3A \arctan(\tan(dx+c))}{2da} - \frac{B \ln(1+\tan^2(dx+c))}{2da}$

3.41.
$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

```
input int(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

```
output -3/2*I*x/a*B-5/2*x/a*A+1/4/a/d*exp(-2*I*(d*x+c))*B-1/4*I/a/d*exp(-2*I*(d*x
+c))*A-2*I/a/d*B*c-2/a/d*A*c-2*I*A/a/d/(exp(2*I*(d*x+c))-1)+1/a/d*ln(exp(2
*I*(d*x+c))-1)*B-I/a/d*ln(exp(2*I*(d*x+c))-1)*A
```

3.41.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.26

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{2(5A+3iB)dx e^{(4i dx+4i c)} - (2(5A+3iB)dx - 9iA+B)e^{(2i dx+2i c)} + 4((iA-B)e^{(4i dx+4i c)} + (-iA+B)e^{(2i dx+2i c)})}{4(ade^{(4i dx+4i c)} - ade^{(2i dx+2i c)})}$$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="f
ricas")
```

```
output -1/4*(2*(5*A + 3*I*B)*d*x*e^(4*I*d*x + 4*I*c) - (2*(5*A + 3*I*B)*d*x - 9*I
*A + B)*e^(2*I*d*x + 2*I*c) + 4*((I*A - B)*e^(4*I*d*x + 4*I*c) + (-I*A + B
)*e^(2*I*d*x + 2*I*c))*log(e^(2*I*d*x + 2*I*c) - 1) - I*A + B)/(a*d*e^(4*I
*d*x + 4*I*c) - a*d*e^(2*I*d*x + 2*I*c))
```

3.41.6 Sympy [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.55

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = -\frac{2iA}{ade^{2ic}e^{2idx} - ad} + \begin{cases} \frac{(-iA+B)e^{-2ic}e^{-2idx}}{4ad} & \text{for } ade^{2ic} \neq 0 \\ x \left(-\frac{5A-3iB}{2a} + \frac{(-5Ae^{2ic}-A-3iBe^{2ic}-iB)e^{-2ic}}{2a} \right) & \text{otherwise} \end{cases} + \frac{x(-5A-3iB)}{2a} - \frac{i(A+iB) \log(e^{2idx} - e^{-2ic})}{ad}$$

input `integrate(cot(d*x+c)**2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)`

output `-2*I*A/(a*d*exp(2*I*c)*exp(2*I*d*x) - a*d) + Piecewise(((-I*A + B)*exp(-2*I*c)*exp(-2*I*d*x)/(4*a*d), Ne(a*d*exp(2*I*c), 0)), (x*(-(-5*A - 3*I*B)/(2*a) + (-5*A*exp(2*I*c) - A - 3*I*B*exp(2*I*c) - I*B)*exp(-2*I*c)/(2*a)), True)) + x*(-5*A - 3*I*B)/(2*a) - I*(A + I*B)*log(exp(2*I*d*x) - exp(-2*I*c))/(a*d)`

3.41.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{a + ia \tan(c + dx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.41.8 Giac [A] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.30

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{a + ia \tan(c + dx)} dx = \frac{\frac{2(iA+B)\log(\tan(dx+c)+i)}{a} + \frac{2(-5iA+3B)\log(\tan(dx+c)-i)}{a} + \frac{8(iA-B)\log(\tan(dx+c))}{a} + \frac{A \tan(dx+c)^2 - iB \tan(dx+c)^2 - 13iA \tan(dx+c) - 8A}{(-i \tan(dx+c)^2 - \tan(dx+c))a}}{8d}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")`

output `-1/8*(2*(I*A + B)*log(tan(d*x + c) + I)/a + 2*(-5*I*A + 3*B)*log(tan(d*x + c) - I)/a + 8*(I*A - B)*log(tan(d*x + c))/a + (A*tan(d*x + c)^2 - I*B*tan(d*x + c)^2 - 13*I*A*tan(d*x + c) + 3*B*tan(d*x + c) - 8*A)/((-I*tan(d*x + c)^2 - tan(d*x + c))*a))/d`

3.41. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

3.41.9 Mupad [B] (verification not implemented)

Time = 7.72 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.24

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = -\frac{\frac{A}{a} + \tan(c+dx) \left(-\frac{B}{2a} + \frac{A3i}{2a}\right)}{d(\tan(c+dx)^2 1i + \tan(c+dx))} - \frac{\ln(\tan(c+dx))(-B+Ai)}{ad} - \frac{\ln(\tan(c+dx)+1i)(B+Ai)}{4ad} + \frac{\ln(\tan(c+dx)-i)(-3B+5i)}{4ad}$$

input `int((cot(c + d*x)^2*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i),x)`output `(log(tan(c + d*x) - 1i)*(A*5i - 3*B))/(4*a*d) - (log(tan(c + d*x))*(A*1i - B))/(a*d) - (log(tan(c + d*x) + 1i)*(A*1i + B))/(4*a*d) - (A/a + tan(c + d*x)*((A*3i)/(2*a) - B/(2*a)))/(d*(tan(c + d*x) + tan(c + d*x)^2*1i))`

$$3.42 \quad \int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

3.42.1 Optimal result 617
 3.42.2 Mathematica [C] (verified) 617
 3.42.3 Rubi [A] (verified) 618
 3.42.4 Maple [A] (verified) 621
 3.42.5 Fricas [A] (verification not implemented) 621
 3.42.6 Sympy [A] (verification not implemented) 622
 3.42.7 Maxima [F(-2)] 623
 3.42.8 Giac [A] (verification not implemented) 623
 3.42.9 Mupad [B] (verification not implemented) 624

3.42.1 Optimal result

Integrand size = 34, antiderivative size = 131

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{3(iA-B)x}{2a} + \frac{3(iA-B) \cot(c+dx)}{2ad} - \frac{(2A+iB) \cot^2(c+dx)}{2ad} - \frac{(2A+iB) \log(\sin(c+dx))}{ad} + \frac{(A+iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))}$$

output $\frac{3}{2}*(I*A-B)*x/a+3/2*(I*A-B)*\cot(d*x+c)/a/d-1/2*(2*A+I*B)*\cot(d*x+c)^2/a/d-(2*A+I*B)*\ln(\sin(d*x+c))/a/d+1/2*(A+I*B)*\cot(d*x+c)^2/d/(a+I*a*\tan(d*x+c))$

3.42.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.82 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.85

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{(A+iB) \cot^3(c+dx)}{i+\cot(c+dx)} + 3i(A+iB) \cot(c+dx) \text{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(c+dx)\right) - (2A+iB) \cot(c+dx)$$

$2ad$

input `Integrate[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `((A + I*B)*Cot[c + d*x]^3)/(I + Cot[c + d*x]) + (3*I)*(A + I*B)*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2] - (2*A + I*B)*(Cot[c + d*x]^2 + 2*(Log[Cos[c + d*x]] + Log[Tan[c + d*x]])))/(2*a*d)`

3.42.3 Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.97, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3042, 4079, 3042, 4012, 25, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{a + ia \tan(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^3(a + ia \tan(c + dx))} dx \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \cot^3(c + dx)(2a(2A + iB) - 3a(iA - B) \tan(c + dx)) dx}{2a^2} + \frac{(A + iB) \cot^2(c + dx)}{2d(a + ia \tan(c + dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{2a(2A + iB) - 3a(iA - B) \tan(c + dx)}{\tan(c + dx)^3} dx}{2a^2} + \frac{(A + iB) \cot^2(c + dx)}{2d(a + ia \tan(c + dx))} \\
 & \quad \downarrow \text{4012} \\
 & \frac{\int -\cot^2(c + dx)(3a(iA - B) + 2a(2A + iB) \tan(c + dx)) dx}{2a^2} - \frac{a(2A + iB) \cot^2(c + dx)}{d} + \\
 & \quad \frac{(A + iB) \cot^2(c + dx)}{2d(a + ia \tan(c + dx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{-\int \cot^2(c + dx)(3a(iA - B) + 2a(2A + iB) \tan(c + dx)) dx}{2a^2} - \frac{a(2A + iB) \cot^2(c + dx)}{d} + \\
 & \quad \frac{(A + iB) \cot^2(c + dx)}{2d(a + ia \tan(c + dx))}
 \end{aligned}$$

3.42. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{-\int \frac{3a(iA-B)+2a(2A+iB)\tan(c+dx)}{\tan(c+dx)^2} dx - \frac{a(2A+iB)\cot^2(c+dx)}{d}}{2a^2} + \frac{(A+iB)\cot^2(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow 4012 \\
& \frac{-\int \cot(c+dx)(2a(2A+iB)-3a(iA-B)\tan(c+dx))dx - \frac{a(2A+iB)\cot^2(c+dx)}{d} + \frac{3a(-B+iA)\cot(c+dx)}{d}}{2a^2} + \\
& \quad \frac{(A+iB)\cot^2(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow 3042 \\
& \frac{-\int \frac{2a(2A+iB)-3a(iA-B)\tan(c+dx)}{\tan(c+dx)} dx - \frac{a(2A+iB)\cot^2(c+dx)}{d} + \frac{3a(-B+iA)\cot(c+dx)}{d}}{2a^2} + \\
& \quad \frac{(A+iB)\cot^2(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow 4014 \\
& \frac{-2a(2A+iB)\int \cot(c+dx)dx - \frac{a(2A+iB)\cot^2(c+dx)}{d} + \frac{3a(-B+iA)\cot(c+dx)}{d} + 3ax(-B+iA)}{2a^2} + \\
& \quad \frac{(A+iB)\cot^2(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow 3042 \\
& \frac{-2a(2A+iB)\int -\tan(c+dx+\frac{\pi}{2})dx - \frac{a(2A+iB)\cot^2(c+dx)}{d} + \frac{3a(-B+iA)\cot(c+dx)}{d} + 3ax(-B+iA)}{2a^2} + \\
& \quad \frac{(A+iB)\cot^2(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow 25 \\
& \frac{2a(2A+iB)\int \tan(\frac{1}{2}(2c+\pi)+dx)dx - \frac{a(2A+iB)\cot^2(c+dx)}{d} + \frac{3a(-B+iA)\cot(c+dx)}{d} + 3ax(-B+iA)}{2a^2} + \\
& \quad \frac{(A+iB)\cot^2(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow 3956 \\
& \frac{-\frac{a(2A+iB)\cot^2(c+dx)}{d} + \frac{3a(-B+iA)\cot(c+dx)}{d} - \frac{2a(2A+iB)\log(-\sin(c+dx))}{d} + 3ax(-B+iA)}{2a^2} + \\
& \quad \frac{(A+iB)\cot^2(c+dx)}{2d(a+ia\tan(c+dx))}
\end{aligned}$$

3.42. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$

input `Int[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `(3*a*(I*A - B)*x + (3*a*(I*A - B)*Cot[c + d*x])/d - (a*(2*A + I*B)*Cot[c + d*x]^2)/d - (2*a*(2*A + I*B)*Log[-Sin[c + d*x]])/d)/(2*a^2) + ((A + I*B)*Cot[c + d*x]^2)/(2*d*(a + I*a*Tan[c + d*x]))`

3.42.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

```
rule 4079 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.42.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.21

method	result
risch	$-\frac{5xB}{2a} + \frac{7ixA}{2a} - \frac{ie^{-2i(dx+c)}B}{4ad} - \frac{e^{-2i(dx+c)}A}{4ad} - \frac{2Bc}{ad} + \frac{4iAc}{ad} - \frac{2i(Be^{2i(dx+c)}+iA-B)}{ad(e^{2i(dx+c)}-1)^2} - \frac{i \ln(e^{2i(dx+c)}-1)}{ad}$
norman	$-\frac{A}{2ad} - \frac{(-iA+B)\tan(dx+c)}{ad} - \frac{3(-iA+B)(\tan^3(dx+c))}{2ad} - \frac{3(-iA+B)x(\tan^2(dx+c))}{2a} - \frac{3(-iA+B)x(\tan^4(dx+c))}{2a} - \frac{(iB+2A)(\tan^2(dx+c))}{2ad}$
derivativedivides	$\frac{A \ln(1+\tan^2(dx+c))}{da} + \frac{3iA \arctan(\tan(dx+c))}{2da} + \frac{iB \ln(1+\tan^2(dx+c))}{2da} - \frac{3B \arctan(\tan(dx+c))}{2da} + \frac{iA}{2da \tan(dx+c)}$
default	$\frac{A \ln(1+\tan^2(dx+c))}{da} + \frac{3iA \arctan(\tan(dx+c))}{2da} + \frac{iB \ln(1+\tan^2(dx+c))}{2da} - \frac{3B \arctan(\tan(dx+c))}{2da} + \frac{iA}{2da \tan(dx+c)}$

```
input int(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

```
output -5/2*x/a*B+7/2*I*x/a*A-1/4*I/a/d*exp(-2*I*(d*x+c))*B-1/4/a/d*exp(-2*I*(d*x
+c))*A-2/a/d*B*c+4*I/a/d*A*c-2*I*(B*exp(2*I*(d*x+c))+I*A-B)/a/d/(exp(2*I*(
d*x+c))-1)^2-I/a/d*ln(exp(2*I*(d*x+c))-1)*B-2*A/a/d*ln(exp(2*I*(d*x+c))-1)
```

3.42.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.44

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{2(-7iA+5B)dx e^{6i dx+6i c} + (4(7iA-5B)dx + A + 9iB)e^{4i dx+4i c} + 2((-7iA+5B)dx - 5A - 4(ade^{6i dx+6i c})}{4(ade^{6i dx+6i c})}$$

3.42. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="fricas")`

output `-1/4*(2*(-7*I*A + 5*B)*d*x*e^(6*I*d*x + 6*I*c) + (4*(7*I*A - 5*B)*d*x + A + 9*I*B)*e^(4*I*d*x + 4*I*c) + 2*((-7*I*A + 5*B)*d*x - 5*A - 5*I*B)*e^(2*I*d*x + 2*I*c) + 4*((2*A + I*B)*e^(6*I*d*x + 6*I*c) - 2*(2*A + I*B)*e^(4*I*d*x + 4*I*c) + (2*A + I*B)*e^(2*I*d*x + 2*I*c))*log(e^(2*I*d*x + 2*I*c) - 1) + A + I*B)/(a*d*e^(6*I*d*x + 6*I*c) - 2*a*d*e^(4*I*d*x + 4*I*c) + a*d*e^(2*I*d*x + 2*I*c))`

3.42.6 Sympy [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.52

$$\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{a + ia \tan(c + dx)} dx$$

$$= \frac{2A - 2iBe^{2ic}e^{2idx} + 2iB}{ade^{4ic}e^{4idx} - 2ade^{2ic}e^{2idx} + ad}$$

$$+ \begin{cases} \frac{(-A-iB)e^{-2ic}e^{-2idx}}{4ad} & \text{for } ade^{2ic} \neq 0 \\ x \left(-\frac{7iA-5B}{2a} + \frac{(7iAe^{2ic}+iA-5Be^{2ic}-B)e^{-2ic}}{2a} \right) & \text{otherwise} \end{cases}$$

$$+ \frac{x(7iA - 5B)}{2a} - \frac{(2A + iB) \log(e^{2idx} - e^{-2ic})}{ad}$$

input `integrate(cot(d*x+c)**3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)`

output `(2*A - 2*I*B*exp(2*I*c)*exp(2*I*d*x) + 2*I*B)/(a*d*exp(4*I*c)*exp(4*I*d*x) - 2*a*d*exp(2*I*c)*exp(2*I*d*x) + a*d) + Piecewise(((-A - I*B)*exp(-2*I*c)*exp(-2*I*d*x)/(4*a*d), Ne(a*d*exp(2*I*c), 0)), (x*(-(7*I*A - 5*B)/(2*a) + (7*I*A*exp(2*I*c) + I*A - 5*B*exp(2*I*c) - B)*exp(-2*I*c)/(2*a)), True)) + x*(7*I*A - 5*B)/(2*a) - (2*A + I*B)*log(exp(2*I*d*x) - exp(-2*I*c))/(a*d)`

3.42.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.42.8 Giac [A] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.22

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$$

$$= \frac{(A-iB)\log(\tan(dx+c)+i)}{a} + \frac{(7A+5iB)\log(\tan(dx+c)-i)}{a} - \frac{4(2A+iB)\log(\tan(dx+c))}{a} - \frac{7A\tan(dx+c)+5iB\tan(dx+c)-9iA+7B}{a(\tan(dx+c)-i)} + \frac{7A-5iB}{a(\tan(dx+c)+i)}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")
```

```
output 1/4*((A - I*B)*log(tan(d*x + c) + I)/a + (7*A + 5*I*B)*log(tan(d*x + c) - I)/a - 4*(2*A + I*B)*log(tan(d*x + c))/a - (7*A*tan(d*x + c) + 5*I*B*tan(d*x + c) - 9*I*A + 7*B)/(a*(tan(d*x + c) - I)) + 2*(6*A*tan(d*x + c)^2 + 3*I*B*tan(d*x + c)^2 + 2*I*A*tan(d*x + c) - 2*B*tan(d*x + c) - A)/(a*tan(d*x + c)^2))/d
```

3.42.9 Mupad [B] (verification not implemented)

Time = 7.94 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.17

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$$

$$= -\frac{\tan(c+dx)^2\left(\frac{3A}{2a} + \frac{B3i}{2a}\right) + \frac{A}{2a} - \tan(c+dx)\left(-\frac{B}{a} + \frac{A1i}{2a}\right)}{d(\tan(c+dx)^3 1i + \tan(c+dx)^2)}$$

$$- \frac{\ln(\tan(c+dx))(2A+B1i)}{ad} + \frac{\ln(\tan(c+dx)+1i)(A-B1i)}{4ad}$$

$$+ \frac{\ln(\tan(c+dx)-i)(7A+B5i)}{4ad}$$

input `int((cot(c + d*x)^3*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i),x)`

output `(log(tan(c + d*x) + 1i)*(A - B*1i))/(4*a*d) - (log(tan(c + d*x))*(2*A + B*1i))/(a*d) - (tan(c + d*x)^2*((3*A)/(2*a) + (B*3i)/(2*a)) + A/(2*a) - tan(c + d*x)*((A*1i)/(2*a) - B/a))/(d*(tan(c + d*x)^2 + tan(c + d*x)^3*1i)) + (log(tan(c + d*x) - 1i)*(7*A + B*5i))/(4*a*d)`

3.43 $\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

3.43.1 Optimal result 625
 3.43.2 Mathematica [C] (verified) 626
 3.43.3 Rubi [A] (verified) 626
 3.43.4 Maple [A] (verified) 630
 3.43.5 Fricas [A] (verification not implemented) 630
 3.43.6 Sympy [A] (verification not implemented) 631
 3.43.7 Maxima [F(-2)] 632
 3.43.8 Giac [A] (verification not implemented) 632
 3.43.9 Mupad [B] (verification not implemented) 633

3.43.1 Optimal result

Integrand size = 34, antiderivative size = 155

$$\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \frac{(5A+3iB)x}{2a} + \frac{(5A+3iB) \cot(c+dx)}{2ad} + \frac{(iA-B) \cot^2(c+dx)}{ad} - \frac{(5A+3iB) \cot^3(c+dx)}{6ad} + \frac{2(iA-B) \log(\sin(c+dx))}{ad} + \frac{(A+iB) \cot^3(c+dx)}{2d(a+ia \tan(c+dx))}$$

```
output 1/2*(5*A+3*I*B)*x/a+1/2*(5*A+3*I*B)*cot(d*x+c)/a/d+(I*A-B)*cot(d*x+c)^2/a/d-1/6*(5*A+3*I*B)*cot(d*x+c)^3/a/d+2*(I*A-B)*ln(sin(d*x+c))/a/d+1/2*(A+I*B)*cot(d*x+c)^3/d/(a+I*a*tan(d*x+c))
```


3.43.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.90 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.74

$$\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= \frac{\frac{3(A+iB)\cot^4(c+dx)}{i+\cot(c+dx)} - (5A+3iB)\cot^3(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, -\tan^2(c+dx)\right) + 6i(A+iB)}{6ad}$$

input `Integrate[(Cot[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `((3*(A + I*B)*Cot[c + d*x]^4)/(I + Cot[c + d*x]) - (5*A + (3*I)*B)*Cot[c + d*x]^3*Hypergeometric2F1[-3/2, 1, -1/2, -Tan[c + d*x]^2] + (6*I)*(A + I*B))*(Cot[c + d*x]^2 + 2*(Log[Cos[c + d*x]] + Log[Tan[c + d*x]]))/(6*a*d)`

3.43.3 Rubi [A] (verified)

Time = 0.91 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.97, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.441$, Rules used = {3042, 4079, 3042, 4012, 25, 3042, 4012, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$\downarrow 3042$$

$$\int \frac{A+B \tan(c+dx)}{\tan(c+dx)^4(a+ia \tan(c+dx))} dx$$

$$\downarrow 4079$$

$$\int \frac{\cot^4(c+dx)(a(5A+3iB) - 4a(iA-B) \tan(c+dx))}{2a^2} dx + \frac{(A+iB)\cot^3(c+dx)}{2d(a+ia \tan(c+dx))}$$

$$\downarrow 3042$$

$$\int \frac{a(5A+3iB) - 4a(iA-B) \tan(c+dx)}{2a^2 \tan(c+dx)^4} dx + \frac{(A+iB)\cot^3(c+dx)}{2d(a+ia \tan(c+dx))}$$

3.43. $\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

$$\begin{aligned}
& \downarrow 4012 \\
& \frac{\int -\cot^3(c+dx)(4a(iA-B) + a(5A+3iB)\tan(c+dx))dx - \frac{a(5A+3iB)\cot^3(c+dx)}{3d}}{2a^2} + \\
& \quad \frac{(A+iB)\cot^3(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow 25 \\
& \frac{-\int \cot^3(c+dx)(4a(iA-B) + a(5A+3iB)\tan(c+dx))dx - \frac{a(5A+3iB)\cot^3(c+dx)}{3d}}{2a^2} + \\
& \quad \frac{(A+iB)\cot^3(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow 3042 \\
& \frac{-\int \frac{4a(iA-B)+a(5A+3iB)\tan(c+dx)}{\tan(c+dx)^3} dx - \frac{a(5A+3iB)\cot^3(c+dx)}{3d}}{2a^2} + \frac{(A+iB)\cot^3(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow 4012 \\
& \frac{-\int \cot^2(c+dx)(a(5A+3iB) - 4a(iA-B)\tan(c+dx))dx - \frac{a(5A+3iB)\cot^3(c+dx)}{3d} + \frac{2a(-B+iA)\cot^2(c+dx)}{d}}{2a^2} + \\
& \quad \frac{(A+iB)\cot^3(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow 3042 \\
& \frac{-\int \frac{a(5A+3iB)-4a(iA-B)\tan(c+dx)}{\tan(c+dx)^2} dx - \frac{a(5A+3iB)\cot^3(c+dx)}{3d} + \frac{2a(-B+iA)\cot^2(c+dx)}{d}}{2a^2} + \\
& \quad \frac{(A+iB)\cot^3(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow 4012 \\
& \frac{-\int -\cot(c+dx)(4a(iA-B) + a(5A+3iB)\tan(c+dx))dx - \frac{a(5A+3iB)\cot^3(c+dx)}{3d} + \frac{2a(-B+iA)\cot^2(c+dx)}{d} + \frac{a(5A+3iB)\cot^2(c+dx)}{d}}{2a^2} + \\
& \quad \frac{(A+iB)\cot^3(c+dx)}{2d(a+ia\tan(c+dx))} \\
& \downarrow 25 \\
& \frac{\int \cot(c+dx)(4a(iA-B) + a(5A+3iB)\tan(c+dx))dx - \frac{a(5A+3iB)\cot^3(c+dx)}{3d} + \frac{2a(-B+iA)\cot^2(c+dx)}{d} + \frac{a(5A+3iB)\cot^2(c+dx)}{d}}{2a^2} + \\
& \quad \frac{(A+iB)\cot^3(c+dx)}{2d(a+ia\tan(c+dx))}
\end{aligned}$$

3.43. $\int \frac{\cot^4(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$

$$\int \frac{4a(iA-B)+a(5A+3iB)\tan(c+dx)}{\tan(c+dx)} dx - \frac{a(5A+3iB)\cot^3(c+dx)}{3d} + \frac{2a(-B+iA)\cot^2(c+dx)}{d} + \frac{a(5A+3iB)\cot(c+dx)}{d} + \frac{2a^2}{2d(a+ia\tan(c+dx))} \frac{(A+iB)\cot^3(c+dx)}{2d(a+ia\tan(c+dx))}$$

$$4a(-B+iA)\int \cot(c+dx)dx - \frac{a(5A+3iB)\cot^3(c+dx)}{3d} + \frac{2a(-B+iA)\cot^2(c+dx)}{d} + \frac{a(5A+3iB)\cot(c+dx)}{d} + ax(5A+3iB) + \frac{2a^2}{2d(a+ia\tan(c+dx))} \frac{(A+iB)\cot^3(c+dx)}{2d(a+ia\tan(c+dx))}$$

$$4a(-B+iA)\int -\tan\left(c+dx+\frac{\pi}{2}\right)dx - \frac{a(5A+3iB)\cot^3(c+dx)}{3d} + \frac{2a(-B+iA)\cot^2(c+dx)}{d} + \frac{a(5A+3iB)\cot(c+dx)}{d} + ax(5A+3iB) + \frac{2a^2}{2d(a+ia\tan(c+dx))} \frac{(A+iB)\cot^3(c+dx)}{2d(a+ia\tan(c+dx))}$$

$$-4a(-B+iA)\int \tan\left(\frac{1}{2}(2c+\pi)+dx\right)dx - \frac{a(5A+3iB)\cot^3(c+dx)}{3d} + \frac{2a(-B+iA)\cot^2(c+dx)}{d} + \frac{a(5A+3iB)\cot(c+dx)}{d} + ax(5A+3iB) + \frac{2a^2}{2d(a+ia\tan(c+dx))} \frac{(A+iB)\cot^3(c+dx)}{2d(a+ia\tan(c+dx))}$$

$$-\frac{a(5A+3iB)\cot^3(c+dx)}{3d} + \frac{2a(-B+iA)\cot^2(c+dx)}{d} + \frac{a(5A+3iB)\cot(c+dx)}{d} + \frac{4a(-B+iA)\log(-\sin(c+dx))}{d} + ax(5A+3iB) + \frac{2a^2}{2d(a+ia\tan(c+dx))} \frac{(A+iB)\cot^3(c+dx)}{2d(a+ia\tan(c+dx))}$$

input `Int[(Cot[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `(a*(5*A + (3*I)*B)*x + (a*(5*A + (3*I)*B)*Cot[c + d*x])/d + (2*a*(I*A - B)*Cot[c + d*x]^2)/d - (a*(5*A + (3*I)*B)*Cot[c + d*x]^3)/(3*d) + (4*a*(I*A - B)*Log[-Sin[c + d*x]])/d)/(2*a^2) + ((A + I*B)*Cot[c + d*x]^3)/(2*d*(a + I*a*Tan[c + d*x]))`

$$3.43. \int \frac{\cot^4(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$$

3.43.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`
- rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`
- rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.43.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.19

method	result
risch	$\frac{7ixB}{2a} + \frac{9xA}{2a} - \frac{e^{-2i(dx+c)}B}{4ad} + \frac{ie^{-2i(dx+c)}A}{4ad} + \frac{4iBc}{ad} + \frac{4Ac}{ad} + \frac{4iAe^{4i(dx+c)} - 6iAe^{2i(dx+c)} + 2Be^{2i(dx+c)} + 1}{ad(e^{2i(dx+c)} - 1)^3}$
norman	$-\frac{A}{3ad} - \frac{(-iA+B)\tan(dx+c)}{2ad} + \frac{(3iB+5A)(\tan^2(dx+c))}{3ad} + \frac{(3iB+5A)(\tan^4(dx+c))}{2ad} + \frac{(3iB+5A)x(\tan^3(dx+c))}{2a} + \frac{(3iB+5A)x(\tan^5(dx+c))}{2a}$ $\frac{1}{\tan(dx+c)^3(1+\tan^2(dx+c))}$
derivativedivides	$\frac{iB}{2da(\tan(dx+c)-i)} + \frac{5A \arctan(\tan(dx+c))}{2da} + \frac{B \ln(1+\tan^2(dx+c))}{da} + \frac{iB}{ad \tan(dx+c)} + \frac{A}{2da(\tan(dx+c)-i)} + \frac{3}{2da}$
default	$\frac{iB}{2da(\tan(dx+c)-i)} + \frac{5A \arctan(\tan(dx+c))}{2da} + \frac{B \ln(1+\tan^2(dx+c))}{da} + \frac{iB}{ad \tan(dx+c)} + \frac{A}{2da(\tan(dx+c)-i)} + \frac{3}{2da}$

input `int(cot(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `7/2*I*x/a*B+9/2*x/a*A-1/4/a/d*exp(-2*I*(d*x+c))*B+1/4*I/a/d*exp(-2*I*(d*x+c))*A+4*I/a/d*B*c+4/a/d*A*c+2/3*(6*I*A*exp(4*I*(d*x+c))-9*I*A*exp(2*I*(d*x+c))+3*B*exp(2*I*(d*x+c))+7*I*A-3*B)/a/d/(exp(2*I*(d*x+c))-1)^3-2/a/d*ln(exp(2*I*(d*x+c))-1)*B+2*I/a/d*ln(exp(2*I*(d*x+c))-1)*A`

3.43.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.61

$$\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= \frac{6(9A+7iB)dx e^{(8i dx+8i c)} - 3(6(9A+7iB)dx - 17iA+B)e^{(6i dx+6i c)} + 3(6(9A+7iB)dx - 27iA+B)}{...}$$

input `integrate(cot(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="fracas")`

```
output 1/12*(6*(9*A + 7*I*B)*d*x*e^(8*I*d*x + 8*I*c) - 3*(6*(9*A + 7*I*B)*d*x - 1
7*I*A + B)*e^(6*I*d*x + 6*I*c) + 3*(6*(9*A + 7*I*B)*d*x - 27*I*A + 11*B)*e
^(4*I*d*x + 4*I*c) - (6*(9*A + 7*I*B)*d*x - 65*I*A + 33*B)*e^(2*I*d*x + 2*
I*c) - 24*((-I*A + B)*e^(8*I*d*x + 8*I*c) + 3*(I*A - B)*e^(6*I*d*x + 6*I*c
) + 3*(-I*A + B)*e^(4*I*d*x + 4*I*c) + (I*A - B)*e^(2*I*d*x + 2*I*c))*log(
e^(2*I*d*x + 2*I*c) - 1) - 3*I*A + 3*B)/(a*d*e^(8*I*d*x + 8*I*c) - 3*a*d*e
^(6*I*d*x + 6*I*c) + 3*a*d*e^(4*I*d*x + 4*I*c) - a*d*e^(2*I*d*x + 2*I*c))
```

3.43.6 Sympy [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.63

$$\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= \frac{12iAe^{4ic}e^{4idx} + 14iA - 6B + (-18iAe^{2ic} + 6Be^{2ic})e^{2idx}}{3ade^{6ic}e^{6idx} - 9ade^{4ic}e^{4idx} + 9ade^{2ic}e^{2idx} - 3ad}$$

$$+ \begin{cases} \frac{(iA-B)e^{-2ic}e^{-2idx}}{4ad} & \text{for } ade^{2ic} \neq 0 \\ x \left(-\frac{9A+7iB}{2a} + \frac{(9Ae^{2ic}+A+7iBe^{2ic}+iB)e^{-2ic}}{2a} \right) & \text{otherwise} \end{cases}$$

$$+ \frac{x(9A+7iB)}{2a} + \frac{2i(A+iB) \log(e^{2idx} - e^{-2ic})}{ad}$$

```
input integrate(cot(d*x+c)**4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)
```

```
output (12*I*A*exp(4*I*c)*exp(4*I*d*x) + 14*I*A - 6*B + (-18*I*A*exp(2*I*c) + 6*B
*exp(2*I*c))*exp(2*I*d*x))/(3*a*d*exp(6*I*c)*exp(6*I*d*x) - 9*a*d*exp(4*I*
c)*exp(4*I*d*x) + 9*a*d*exp(2*I*c)*exp(2*I*d*x) - 3*a*d) + Piecewise(((I*A
- B)*exp(-2*I*c)*exp(-2*I*d*x)/(4*a*d), Ne(a*d*exp(2*I*c), 0)), (x*(-(9*A
+ 7*I*B)/(2*a) + (9*A*exp(2*I*c) + A + 7*I*B*exp(2*I*c) + I*B)*exp(-2*I*c
))/(2*a)), True)) + x*(9*A + 7*I*B)/(2*a) + 2*I*(A + I*B)*log(exp(2*I*d*x)
- exp(-2*I*c))/(a*d)
```

3.43.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^4(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(cot(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.43.8 Giac [A] (verification not implemented)

Time = 0.89 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.19

$$\int \frac{\cot^4(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \frac{\frac{3(-iA-B)\log(\tan(dx+c)+i)}{a} + \frac{3(9iA-7B)\log(\tan(dx+c)-i)}{a} + \frac{24(-iA+B)\log(\tan(dx+c))}{a} + \frac{3(-9iA\tan(dx+c)+7B\tan(dx+c))}{a(\tan(dx+c)-i)}}{12d}$$

```
input integrate(cot(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")
```

```
output -1/12*(3*(-I*A - B)*log(tan(d*x + c) + I)/a + 3*(9*I*A - 7*B)*log(tan(d*x + c) - I)/a + 24*(-I*A + B)*log(tan(d*x + c))/a + 3*(-9*I*A*tan(d*x + c) + 7*B*tan(d*x + c) - 11*A - 9*I*B)/(a*(tan(d*x + c) - I)) + 2*I*(22*A*tan(d*x + c)^3 + 22*I*B*tan(d*x + c)^3 + 12*I*A*tan(d*x + c)^2 - 6*B*tan(d*x + c)^2 - 3*A*tan(d*x + c) - 3*I*B*tan(d*x + c) - 2*I*A)/(a*tan(d*x + c)^3))/d
```

3.43.9 Mupad [B] (verification not implemented)

Time = 7.95 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.12

$$\int \frac{\cot^4(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$$

$$= \frac{\tan(c+dx)^2 \left(\frac{3A}{2a} + \frac{B1i}{2a}\right) + \tan(c+dx)^3 \left(-\frac{3B}{2a} + \frac{A5i}{2a}\right) - \frac{A}{3a} + \tan(c+dx) \left(-\frac{B}{2a} + \frac{A1i}{6a}\right)}{d \left(\tan(c+dx)^4 1i + \tan(c+dx)^3\right)}$$

$$+ \frac{2 \ln(\tan(c+dx)) (-B + A 1i)}{a d} + \frac{\ln(\tan(c+dx) + 1i) (B + A 1i)}{4 a d}$$

$$- \frac{\ln(\tan(c+dx) - i) (-7 B + A 9i)}{4 a d}$$

```
input int((cot(c + d*x)^4*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i),x)
```

```
output (tan(c + d*x)^2*((3*A)/(2*a) + (B*1i)/(2*a)) + tan(c + d*x)^3*((A*5i)/(2*a)
) - (3*B)/(2*a)) - A/(3*a) + tan(c + d*x)*((A*1i)/(6*a) - B/(2*a)))/(d*(ta
n(c + d*x)^3 + tan(c + d*x)^4*1i)) + (2*log(tan(c + d*x))*(A*1i - B))/(a*d
) + (log(tan(c + d*x) + 1i)*(A*1i + B))/(4*a*d) - (log(tan(c + d*x) - 1i)*
(A*9i - 7*B))/(4*a*d)
```


3.44 $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

3.44.1 Optimal result 634
 3.44.2 Mathematica [A] (verified) 634
 3.44.3 Rubi [A] (verified) 635
 3.44.4 Maple [A] (verified) 637
 3.44.5 Fricas [A] (verification not implemented) 638
 3.44.6 Sympy [A] (verification not implemented) 638
 3.44.7 Maxima [F(-2)] 639
 3.44.8 Giac [A] (verification not implemented) 639
 3.44.9 Mupad [B] (verification not implemented) 640

3.44.1 Optimal result

Integrand size = 34, antiderivative size = 142

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = -\frac{3(iA-3B)x}{4a^2} + \frac{(A+2iB) \log(\cos(c+dx))}{a^2d} + \frac{3(iA-3B) \tan(c+dx)}{4a^2d} + \frac{(A+2iB) \tan^2(c+dx)}{2a^2d(1+i \tan(c+dx))} + \frac{(iA-B) \tan^3(c+dx)}{4d(a+ia \tan(c+dx))^2}$$

output

```
-3/4*(I*A-3*B)*x/a^2+(A+2*I*B)*ln(cos(d*x+c))/a^2/d+3/4*(I*A-3*B)*tan(d*x+c)/a^2/d+1/2*(A+2*I*B)*tan(d*x+c)^2/a^2/d/(1+I*tan(d*x+c))+1/4*(I*A-B)*tan(d*x+c)^3/d/(a+I*a*tan(d*x+c))^2
```

3.44.2 Mathematica [A] (verified)

Time = 1.51 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.39

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = \frac{(7A+17iB) \log(i-\tan(c+dx)) + (A-iB) \log(i+\tan(c+dx)) + 2(-3iA+9B+(7iA-17B) \log(i$$

input `Integrate[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `((7*A + (17*I)*B)*Log[I - Tan[c + d*x]] + (A - I*B)*Log[I + Tan[c + d*x]] + 2*((-3*I)*A + 9*B + ((7*I)*A - 17*B)*Log[I - Tan[c + d*x]] + (I*A + B)*Log[I + Tan[c + d*x]])*Tan[c + d*x] + (8*A + (28*I)*B - (7*A + (17*I)*B)*Log[I - Tan[c + d*x]] - (A - I*B)*Log[I + Tan[c + d*x]])*Tan[c + d*x]^2 - 8*B*Tan[c + d*x]^3)/(8*a^2*d*(-I + Tan[c + d*x])^2)`

3.44.3 Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.04, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4078, 3042, 4078, 27, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^3(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+iA) \tan^3(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{\int \frac{\tan^2(c+dx)(3a(iA-B)+a(A+5iB) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{4a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B+iA) \tan^3(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{\int \frac{\tan(c+dx)^2(3a(iA-B)+a(A+5iB) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{4a^2} \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+iA) \tan^3(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{\int -2 \tan(c+dx)(4a^2(A+2iB)-3a^2(iA-3B) \tan(c+dx)) dx}{2a^2} - \frac{2(A+2iB) \tan^2(c+dx)}{d(1+i \tan(c+dx))} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B+iA) \tan^3(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{\int \tan(c+dx)(4a^2(A+2iB)-3a^2(iA-3B) \tan(c+dx)) dx}{a^2} - \frac{2(A+2iB) \tan^2(c+dx)}{d(1+i \tan(c+dx))}
 \end{aligned}$$

3.44. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^3(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int \tan(c+dx)(4a^2(A+2iB) - 3a^2(iA-3B) \tan(c+dx)) dx}{a^2} - \frac{2(A+2iB) \tan^2(c+dx)}{d(1+i \tan(c+dx))} \\
& \downarrow 4008 \\
& \frac{(-B + iA) \tan^3(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{4a^2(A+2iB) \int \tan(c+dx) dx - \frac{3a^2(-3B+iA) \tan(c+dx)}{d} + 3a^2x(-3B+iA)}{a^2} - \frac{2(A+2iB) \tan^2(c+dx)}{d(1+i \tan(c+dx))} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^3(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{4a^2(A+2iB) \int \tan(c+dx) dx - \frac{3a^2(-3B+iA) \tan(c+dx)}{d} + 3a^2x(-3B+iA)}{a^2} - \frac{2(A+2iB) \tan^2(c+dx)}{d(1+i \tan(c+dx))} \\
& \downarrow 3956 \\
& \frac{(-B + iA) \tan^3(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{-\frac{3a^2(-3B+iA) \tan(c+dx)}{d} - \frac{4a^2(A+2iB) \log(\cos(c+dx))}{d} + 3a^2x(-3B+iA)}{a^2} - \frac{2(A+2iB) \tan^2(c+dx)}{d(1+i \tan(c+dx))} \\
& \downarrow 4a^2
\end{aligned}$$

input `Int[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `((I*A - B)*Tan[c + d*x]^3)/(4*d*(a + I*a*Tan[c + d*x])^2) - ((-2*(A + (2*I)*B)*Tan[c + d*x]^2)/(d*(1 + I*Tan[c + d*x])) + (3*a^2*(I*A - 3*B)*x - (4*a^2*(A + (2*I)*B)*Log[Cos[c + d*x]])/d - (3*a^2*(I*A - 3*B)*Tan[c + d*x])/d)/a^2)/(4*a^2)`

3.44.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

$$3.44. \int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4078 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.44.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.22

method	result
derivativedivides	$-\frac{B \tan(dx+c)}{d a^2} - \frac{A}{4d a^2 (\tan(dx+c)-i)^2} - \frac{iB}{4d a^2 (\tan(dx+c)-i)^2} - \frac{A \ln(1+\tan^2(dx+c))}{2d a^2} - \frac{3iA \arctan(\tan(dx+c))}{4d a^2}$
default	$-\frac{B \tan(dx+c)}{d a^2} - \frac{A}{4d a^2 (\tan(dx+c)-i)^2} - \frac{iB}{4d a^2 (\tan(dx+c)-i)^2} - \frac{A \ln(1+\tan^2(dx+c))}{2d a^2} - \frac{3iA \arctan(\tan(dx+c))}{4d a^2}$
risch	$\frac{17xB}{4a^2} - \frac{7ixA}{4a^2} - \frac{3ie^{-2i(dx+c)}B}{4a^2d} - \frac{e^{-2i(dx+c)}A}{2a^2d} + \frac{ie^{-4i(dx+c)}B}{16a^2d} + \frac{e^{-4i(dx+c)}A}{16a^2d} + \frac{4Bc}{a^2d} - \frac{2iAc}{a^2d} - \frac{2}{da^2(e^{2i(dx+c)} + 1)}$

input `int(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output
$$-1/d/a^2*B*\tan(d*x+c)-1/4/d/a^2/(\tan(d*x+c)-I)^2*A-1/4*I/d/a^2/(\tan(d*x+c)-I)^2*B-1/2/d/a^2*A*\ln(1+\tan(d*x+c)^2)-3/4*I/d/a^2*A*\arctan(\tan(d*x+c))-I/d/a^2*B*\ln(1+\tan(d*x+c)^2)+9/4/d/a^2*B*\arctan(\tan(d*x+c))+5/4*I/d/a^2/(\tan(d*x+c)-I)*A-7/4/d/a^2/(\tan(d*x+c)-I)*B$$

3.44.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.06

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx =$$

$$\frac{4(7iA-17B)dx e^{(6i dx+6i c)} + 4((7iA-17B)dx + 2A + 11iB)e^{(4i dx+4i c)} + (7A+11iB)e^{(2i dx+2i c)}}{16(a^2 d e^{(6i dx+6i c)} + a^2 d e^{(4i dx+4i c)})}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="fracas")
```

```
output -1/16*(4*(7*I*A - 17*B)*d*x*e^(6*I*d*x + 6*I*c) + 4*((7*I*A - 17*B)*d*x + 2*A + 11*I*B)*e^(4*I*d*x + 4*I*c) + (7*A + 11*I*B)*e^(2*I*d*x + 2*I*c) - 16*((A + 2*I*B)*e^(6*I*d*x + 6*I*c) + (A + 2*I*B)*e^(4*I*d*x + 4*I*c))*log(e^(2*I*d*x + 2*I*c) + 1) - A - I*B)/(a^2*d*e^(6*I*d*x + 6*I*c) + a^2*d*e^(4*I*d*x + 4*I*c))
```

3.44.6 Sympy [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.85

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= -\frac{2iB}{a^2 d e^{2ic} e^{2idx} + a^2 d}$$

$$+ \begin{cases} \frac{((4Aa^2 d e^{2ic} + 4iBa^2 d e^{2ic})e^{-4idx} + (-32Aa^2 d e^{4ic} - 48iBa^2 d e^{4ic})e^{-2idx})e^{-6ic}}{64a^4 d^2} & \text{for } a^4 d^2 e^{6ic} \neq 0 \\ x \left(-\frac{7iA+17B}{4a^2} + \frac{(-7iAe^{4ic} + 4iAe^{2ic} - iA + 17Be^{4ic} - 6Be^{2ic} + B)e^{-4ic}}{4a^2} \right) & \text{otherwise} \end{cases}$$

$$+ \frac{x(-7iA+17B)}{4a^2} + \frac{(A+2iB)\log(e^{2idx} + e^{-2ic})}{a^2 d}$$

```
input integrate(tan(d*x+c)**3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**2,x)
```

```
output -2*I*B/(a**2*d*exp(2*I*c)*exp(2*I*d*x) + a**2*d) + Piecewise((((4*A*a**2*d
*exp(2*I*c) + 4*I*B*a**2*d*exp(2*I*c))*exp(-4*I*d*x) + (-32*A*a**2*d*exp(4
*I*c) - 48*I*B*a**2*d*exp(4*I*c))*exp(-2*I*d*x))*exp(-6*I*c)/(64*a**4*d**2
), Ne(a**4*d**2*exp(6*I*c), 0)), (x*(-(-7*I*A + 17*B)/(4*a**2) + (-7*I*A*exp
(4*I*c) + 4*I*A*exp(2*I*c) - I*A + 17*B*exp(4*I*c) - 6*B*exp(2*I*c) + B)
*exp(-4*I*c)/(4*a**2)), True)) + x*(-7*I*A + 17*B)/(4*a**2) + (A + 2*I*B)*
log(exp(2*I*d*x) + exp(-2*I*c))/(a**2*d)
```

3.44.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^3(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm=
"maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.44.8 Giac [A] (verification not implemented)

Time = 0.68 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.85

$$\int \frac{\tan^3(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx =$$

$$\frac{\frac{2(A-iB) \log(\tan(dx+c)+i)}{a^2} + \frac{2(7A+17iB) \log(\tan(dx+c)-i)}{a^2} + \frac{16B \tan(dx+c)}{a^2} - \frac{21A \tan(dx+c)^2 + 51iB \tan(dx+c)^2 - 22iA \tan(dx+c)}{a^2(\tan(dx+c)-i)}}{16d}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm=
"giac")
```

```
output -1/16*(2*(A - I*B)*log(tan(d*x + c) + I)/a^2 + 2*(7*A + 17*I*B)*log(tan(d*
x + c) - I)/a^2 + 16*B*tan(d*x + c)/a^2 - (21*A*tan(d*x + c)^2 + 51*I*B*ta
n(d*x + c)^2 - 22*I*A*tan(d*x + c) + 74*B*tan(d*x + c) - 5*A - 27*I*B)/(a^
2*(tan(d*x + c) - I)^2))/d
```

3.44. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

3.44.9 Mupad [B] (verification not implemented)

Time = 7.78 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.99

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \frac{\frac{(A+B2i)1i}{a^2} + \frac{B}{2a^2} - \tan(c+dx) \left(\frac{5(A+B2i)}{4a^2} - \frac{B3i}{4a^2} \right)}{d(\tan(c+dx)^2 1i + 2\tan(c+dx) - i)} + \frac{\ln(\tan(c+dx) + 1i)(B+A1i) 1i}{8a^2 d} - \frac{B\tan(c+dx)}{a^2 d} - \frac{\ln(\tan(c+dx) - i)(7A+B17i)}{8a^2 d}$$

input `int((tan(c + d*x)^3*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^2,x)`output `(((A + B*2i)*1i)/a^2 + B/(2*a^2) - tan(c + d*x)*((5*(A + B*2i))/(4*a^2) - (B*3i)/(4*a^2)))/(d*(2*tan(c + d*x) + tan(c + d*x)^2*1i - 1i)) + (log(tan(c + d*x) + 1i)*(A*1i + B)*1i)/(8*a^2*d) - (B*tan(c + d*x))/(a^2*d) - (log(tan(c + d*x) - 1i)*(7*A + B*17i))/(8*a^2*d)`

3.45
$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.45.1 Optimal result 641
 3.45.2 Mathematica [A] (verified) 641
 3.45.3 Rubi [A] (verified) 642
 3.45.4 Maple [A] (verified) 644
 3.45.5 Fricas [A] (verification not implemented) 645
 3.45.6 Sympy [A] (verification not implemented) 645
 3.45.7 Maxima [F(-2)] 646
 3.45.8 Giac [A] (verification not implemented) 646
 3.45.9 Mupad [B] (verification not implemented) 647

3.45.1 Optimal result

Integrand size = 34, antiderivative size = 103

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = -\frac{(A+3iB)x}{4a^2} + \frac{B \log(\cos(c+dx))}{a^2d} + \frac{iA-3B}{4a^2d(1+i \tan(c+dx))} + \frac{(iA-B) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2}$$

output `-1/4*(A+3*I*B)*x/a^2+B*ln(cos(d*x+c))/a^2/d+1/4*(I*A-3*B)/a^2/d/(1+I*tan(d*x+c))+1/4*(I*A-B)*tan(d*x+c)^2/d/(a+I*a*tan(d*x+c))^2`

3.45.2 Mathematica [A] (verified)

Time = 0.85 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.53

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = \frac{\sec^2(c+dx)(-2iA+4B+\cos(2(c+dx))(2iA-4B+(-iA+7B) \log(i-\tan(c+dx)))+(iA+B) \log(i+\tan(c+dx)))}{8a^2d(-1-i \tan(c+dx))}$$

input `Integrate[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output $(\text{Sec}[c + d*x]^2*((-2*I)*A + 4*B + \text{Cos}[2*(c + d*x)]*((2*I)*A - 4*B + ((-I)*A + 7*B)*\text{Log}[I - \text{Tan}[c + d*x]] + (I*A + B)*\text{Log}[I + \text{Tan}[c + d*x]]) + (-A - (3*I)*B + (A + (7*I)*B)*\text{Log}[I - \text{Tan}[c + d*x]] - (A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]])*\text{Sin}[2*(c + d*x)])/(8*a^2*d*(-I + \text{Tan}[c + d*x])^2)$

3.45.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.324$, Rules used = {3042, 4078, 27, 3042, 4072, 25, 27, 3042, 3956, 4009, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(c+dx)^2(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx \\ & \quad \downarrow \text{4078} \\ & \frac{(-B+IA) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{\int \frac{2 \tan(c+dx)(a(iA-B)+2iaB \tan(c+dx))}{i \tan(c+dx)a+a} dx}{4a^2} \\ & \quad \downarrow \text{27} \\ & \frac{(-B+IA) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{\int \frac{\tan(c+dx)(a(iA-B)+2iaB \tan(c+dx))}{i \tan(c+dx)a+a} dx}{2a^2} \\ & \quad \downarrow \text{3042} \\ & \frac{(-B+IA) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{\int \frac{\tan(c+dx)(a(iA-B)+2iaB \tan(c+dx))}{i \tan(c+dx)a+a} dx}{2a^2} \\ & \quad \downarrow \text{4072} \\ & \frac{(-B+IA) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{2B \int \tan(c+dx) dx - \frac{i \int -\frac{a^2(A+3iB) \tan(c+dx)}{i \tan(c+dx)a+a} dx}{a}}{2a^2} \\ & \quad \downarrow \text{25} \\ & \frac{(-B+IA) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{2B \int \tan(c+dx) dx + \frac{i \int \frac{a^2(A+3iB) \tan(c+dx)}{i \tan(c+dx)a+a} dx}{a}}{2a^2} \end{aligned}$$

3.45. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{(-B + iA) \tan^2(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{2B \int \tan(c + dx) dx + ia(A + 3iB) \int \frac{\tan(c+dx)}{i \tan(c+dx)a+a} dx}{2a^2} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^2(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{2B \int \tan(c + dx) dx + ia(A + 3iB) \int \frac{\tan(c+dx)}{i \tan(c+dx)a+a} dx}{2a^2} \\
& \downarrow 3956 \\
& \frac{(-B + iA) \tan^2(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{-\frac{2B \log(\cos(c+dx))}{d} + ia(A + 3iB) \int \frac{\tan(c+dx)}{i \tan(c+dx)a+a} dx}{2a^2} \\
& \downarrow 4009 \\
& \frac{(-B + iA) \tan^2(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{-\frac{2B \log(\cos(c+dx))}{d} + ia(A + 3iB) \left(-\frac{i \int 1 dx}{2a} - \frac{1}{2d(a+ia \tan(c+dx))} \right)}{2a^2} \\
& \downarrow 24 \\
& \frac{(-B + iA) \tan^2(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{-\frac{2B \log(\cos(c+dx))}{d} + ia(A + 3iB) \left(-\frac{1}{2d(a+ia \tan(c+dx))} - \frac{ix}{2a} \right)}{2a^2}
\end{aligned}$$

input `Int[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `((I*A - B)*Tan[c + d*x]^2)/(4*d*(a + I*a*Tan[c + d*x])^2) - ((-2*B*Log[Cos[c + d*x]])/d + I*a*(A + (3*I)*B)*(((-1/2*I)*x)/a - 1/(2*d*(a + I*a*Tan[c + d*x]))))/ (2*a^2)`

3.45.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

rule 4072 `Int[(((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*(d/b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c - a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0]`

rule 4078 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.45.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.18

method	result
risch	$-\frac{7ixB}{4a^2} - \frac{xA}{4a^2} - \frac{e^{-2i(dx+c)}B}{2a^2d} + \frac{ie^{-2i(dx+c)}A}{4a^2d} + \frac{e^{-4i(dx+c)}B}{16a^2d} - \frac{ie^{-4i(dx+c)}A}{16a^2d} - \frac{2iBc}{a^2d} + \frac{B \ln(e^{2i(dx+c)}+1)}{a^2d}$
derivativedivides	$\frac{iA}{4da^2(\tan(dx+c)-i)^2} - \frac{B}{4da^2(\tan(dx+c)-i)^2} - \frac{A \arctan(\tan(dx+c))}{4da^2} - \frac{B \ln(1+\tan^2(dx+c))}{2da^2} - \frac{3iB \arctan(\tan(dx+c))}{4da^2}$
default	$\frac{iA}{4da^2(\tan(dx+c)-i)^2} - \frac{B}{4da^2(\tan(dx+c)-i)^2} - \frac{A \arctan(\tan(dx+c))}{4da^2} - \frac{B \ln(1+\tan^2(dx+c))}{2da^2} - \frac{3iB \arctan(\tan(dx+c))}{4da^2}$

3.45. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

```
input int(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x,method=_RETURNVER
BOSE)
```

```
output -7/4*I*x/a^2*B-1/4*x/a^2*A-1/2/a^2/d*exp(-2*I*(d*x+c))*B+1/4*I/a^2/d*exp(-
2*I*(d*x+c))*A+1/16/a^2/d*exp(-4*I*(d*x+c))*B-1/16*I/a^2/d*exp(-4*I*(d*x+c
))*A-2*I*B/a^2/d*c+B/a^2/d*ln(exp(2*I*(d*x+c))+1)
```

3.45.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.82

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = \frac{(4(A+7iB)dx e^{(4i dx+4i c)} - 16 B e^{(4i dx+4i c)} \log(e^{(2i dx+2i c)} + 1) + 4(-iA+2B)e^{(2i dx+2i c)} + iA - B)}{16 a^2 d}$$

```
input integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm=
"fricas")
```

```
output -1/16*(4*(A + 7*I*B)*d*x*e^(4*I*d*x + 4*I*c) - 16*B*e^(4*I*d*x + 4*I*c)*lo
g(e^(2*I*d*x + 2*I*c) + 1) + 4*(-I*A + 2*B)*e^(2*I*d*x + 2*I*c) + I*A - B)
*e^(-4*I*d*x - 4*I*c)/(a^2*d)
```

3.45.6 Sympy [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 223, normalized size of antiderivative = 2.17

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = \frac{B \log(e^{2idx} + e^{-2ic})}{a^2 d} + \begin{cases} \frac{((-4iAa^2de^{2ic}+4Ba^2de^{2ic})e^{-4idx}+(16iAa^2de^{4ic}-32Ba^2de^{4ic})e^{-2idx})e^{-6ic}}{64a^4d^2} & \text{for } a^4d^2e^{6ic} \neq 0 \\ x\left(-\frac{-A-7iB}{4a^2} + \frac{(-Ae^{4ic}+2Ae^{2ic}-A-7iBe^{4ic}+4iBe^{2ic}-iB)e^{-4ic}}{4a^2}\right) & \text{otherwise} \end{cases} + \frac{x(-A-7iB)}{4a^2}$$

```
input integrate(tan(d*x+c)**2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**2,x)
```

```
output B*log(exp(2*I*d*x) + exp(-2*I*c))/(a**2*d) + Piecewise(((((-4*I*A*a**2*d*exp(2*I*c) + 4*B*a**2*d*exp(2*I*c))*exp(-4*I*d*x) + (16*I*A*a**2*d*exp(4*I*c) - 32*B*a**2*d*exp(4*I*c))*exp(-2*I*d*x))*exp(-6*I*c)/(64*a**4*d**2), Ne(a**4*d**2*exp(6*I*c), 0)), (x*(-(-A - 7*I*B)/(4*a**2) + (-A*exp(4*I*c) + 2*A*exp(2*I*c) - A - 7*I*B*exp(4*I*c) + 4*I*B*exp(2*I*c) - I*B)*exp(-4*I*c)/(4*a**2)), True)) + x*(-A - 7*I*B)/(4*a**2)
```

3.45.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.45.8 Giac [A] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.04

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx = \frac{\frac{2(iA+B)\log(\tan(dx+c)+i)}{a^2} + \frac{2(-iA+7B)\log(\tan(dx+c)-i)}{a^2} + \frac{3iA\tan(dx+c)^2 - 21B\tan(dx+c)^2 - 6A\tan(dx+c) + 22iB\tan(dx+c) + 5iA + 5B}{a^2(\tan(dx+c)-i)^2}}{16d}$$

```
input integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")
```

```
output -1/16*(2*(I*A + B)*log(tan(d*x + c) + I)/a^2 + 2*(-I*A + 7*B)*log(tan(d*x + c) - I)/a^2 + (3*I*A*tan(d*x + c)^2 - 21*B*tan(d*x + c)^2 - 6*A*tan(d*x + c) + 22*I*B*tan(d*x + c) + 5*I*A + 5*B)/(a^2*(tan(d*x + c) - I)^2))/d
```

3.45. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

3.45.9 Mupad [B] (verification not implemented)

Time = 7.15 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.11

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = \frac{\frac{A}{2a^2} + \frac{B1i}{a^2} + \tan(c+dx) \left(-\frac{5B}{4a^2} + \frac{A3i}{4a^2}\right)}{d (\tan(c+dx)^2 1i + 2 \tan(c+dx) - i)} - \frac{\ln(\tan(c+dx) + 1i) (B + A 1i)}{8a^2 d} + \frac{\ln(\tan(c+dx) - i) (-7B + A 1i)}{8a^2 d}$$

input `int((tan(c + d*x)^2*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^2,x)`output `(A/(2*a^2) + (B*1i)/a^2 + tan(c + d*x)*((A*3i)/(4*a^2) - (5*B)/(4*a^2)))/(d*(2*tan(c + d*x) + tan(c + d*x)^2*1i - 1i)) - (log(tan(c + d*x) + 1i)*(A*1i + B))/(8*a^2*d) + (log(tan(c + d*x) - 1i)*(A*1i - 7*B))/(8*a^2*d)`

3.46 $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

3.46.1	Optimal result	648
3.46.2	Mathematica [A] (verified)	648
3.46.3	Rubi [A] (verified)	649
3.46.4	Maple [A] (verified)	650
3.46.5	Fricas [A] (verification not implemented)	651
3.46.6	Sympy [A] (verification not implemented)	651
3.46.7	Maxima [F(-2)]	652
3.46.8	Giac [A] (verification not implemented)	652
3.46.9	Mupad [B] (verification not implemented)	652

3.46.1 Optimal result

Integrand size = 32, antiderivative size = 76

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = -\frac{(iA+B)x}{4a^2} + \frac{A+3iB}{4a^2d(1+i \tan(c+dx))} - \frac{A+iB}{4d(a+ia \tan(c+dx))^2}$$

```
output -1/4*(I*A+B)*x/a^2+1/4*(A+3*I*B)/a^2/d/(1+I*tan(d*x+c))+1/4*(-A-I*B)/d/(a+I*a*tan(d*x+c))^2
```

3.46.2 Mathematica [A] (verified)

Time = 0.67 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.21

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = \frac{\sec^2(c+dx)(-4iB+(A+4iAdx+B(i+4dx)) \cos(2(c+dx)) + (-iA+B-4Adx+4iBdx) \sin(2(c+dx)))}{16a^2d(-i+\tan(c+dx))^2}$$

```
input Integrate[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]
```

```
output (Sec[c + d*x]^2*((-4*I)*B + (A + (4*I)*A*d*x + B*(I + 4*d*x))*Cos[2*(c + d*x)] + ((-I)*A + B - 4*A*d*x + (4*I)*B*d*x)*Sin[2*(c + d*x)])/(16*a^2*d*(-I + Tan[c + d*x])^2)
```

3.46.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.08, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4073, 3042, 4009, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx \\
 & \quad \downarrow \text{4073} \\
 & -\frac{i \int \frac{a(A+iB)+2aB\tan(c+dx)}{i\tan(c+dx)a+a} dx}{2a^2} - \frac{A+iB}{4d(a+ia\tan(c+dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{i \int \frac{a(A+iB)+2aB\tan(c+dx)}{i\tan(c+dx)a+a} dx}{2a^2} - \frac{A+iB}{4d(a+ia\tan(c+dx))^2} \\
 & \quad \downarrow \text{4009} \\
 & -\frac{i\left(\frac{1}{2}(A-iB) \int 1dx + \frac{-3B+iA}{2d(1+i\tan(c+dx))}\right)}{2a^2} - \frac{A+iB}{4d(a+ia\tan(c+dx))^2} \\
 & \quad \downarrow \text{24} \\
 & -\frac{i\left(\frac{-3B+iA}{2d(1+i\tan(c+dx))} + \frac{1}{2}x(A-iB)\right)}{2a^2} - \frac{A+iB}{4d(a+ia\tan(c+dx))^2}
 \end{aligned}$$

input `Int[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `((-1/2*I)*(((A - I*B)*x)/2 + (I*A - 3*B)/(2*d*(1 + I*Tan[c + d*x]))))/a^2 - (A + I*B)/(4*d*(a + I*a*Tan[c + d*x])^2)`

3.46.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

- rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

- rule 4073 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(A*b - a*B))*(a*c + b*d)*((a + b*Tan[e + f*x])^m/(2*a^2*f*m)), x] + Simp[1/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[A*b*c + a*B*c + a*A*d + b*B*d + 2*a*B*d*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && EqQ[a^2 + b^2, 0]`

3.46.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.96

method	result
risch	$-\frac{x B}{4 a^2} - \frac{i x A}{4 a^2} + \frac{i e^{-2 i(d x+c)} B}{4 a^2 d} - \frac{i e^{-4 i(d x+c)} B}{16 a^2 d} - \frac{e^{-4 i(d x+c)} A}{16 a^2 d}$
derivativedivides	$-\frac{i A \arctan(\tan(dx+c))}{4 d a^2} - \frac{B \arctan(\tan(dx+c))}{4 d a^2} + \frac{A}{4 d a^2(\tan(dx+c)-i)^2} + \frac{i B}{4 d a^2(\tan(dx+c)-i)^2} - \frac{i A}{4 d a^2(\tan(dx+c)-i)^2}$
default	$-\frac{i A \arctan(\tan(dx+c))}{4 d a^2} - \frac{B \arctan(\tan(dx+c))}{4 d a^2} + \frac{A}{4 d a^2(\tan(dx+c)-i)^2} + \frac{i B}{4 d a^2(\tan(dx+c)-i)^2} - \frac{i A}{4 d a^2(\tan(dx+c)-i)^2}$

```
input int(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x,method=_RETURNVERBOSE)
```

```
output -1/4*x/a^2*B-1/4*I*x/a^2*A+1/4*I*B/a^2/d*exp(-2*I*(d*x+c))-1/16*I/a^2/d*exp(-4*I*(d*x+c))*B-1/16/a^2/d*exp(-4*I*(d*x+c))*A
```

3.46. $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

3.46.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.68

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= -\frac{(4(iA+B)dx e^{(4i dx+4i c)} - 4i B e^{(2i dx+2i c)} + A+i B) e^{(-4i dx-4i c)}}{16 a^2 d}$$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="fracas")
```

```
output -1/16*(4*(I*A + B)*d*x*e^(4*I*d*x + 4*I*c) - 4*I*B*e^(2*I*d*x + 2*I*c) + A + I*B)*e^(-4*I*d*x - 4*I*c)/(a^2*d)
```

3.46.6 Sympy [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 167, normalized size of antiderivative = 2.20

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= \begin{cases} \frac{(16iBa^2de^{4ic}e^{-2idx} + (-4Aa^2de^{2ic} - 4iBa^2de^{2ic})e^{-4idx})e^{-6ic}}{64a^4d^2} & \text{for } a^4d^2e^{6ic} \neq 0 \\ x\left(-\frac{-iA-B}{4a^2} + \frac{(-iAe^{4ic} + iA - Be^{4ic} + 2Be^{2ic} - B)e^{-4ic}}{4a^2}\right) & \text{otherwise} \end{cases} + \frac{x(-iA - B)}{4a^2}$$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**2,x)
```

```
output Piecewise((((16*I*B*a**2*d*exp(4*I*c)*exp(-2*I*d*x) + (-4*A*a**2*d*exp(2*I*c) - 4*I*B*a**2*d*exp(2*I*c))*exp(-4*I*d*x))*exp(-6*I*c)/(64*a**4*d**2), Ne(a**4*d**2*exp(6*I*c), 0)), (x*(-(-I*A - B)/(4*a**2) + (-I*A*exp(4*I*c) + I*A - B*exp(4*I*c) + 2*B*exp(2*I*c) - B)*exp(-4*I*c)/(4*a**2)), True)) + x*(-I*A - B)/(4*a**2)
```

3.46.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.46.8 Giac [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.38

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx = \frac{\frac{2(A-iB)\log(\tan(dx+c)+i)}{a^2} - \frac{2(A-iB)\log(\tan(dx+c)-i)}{a^2} + \frac{3A\tan(dx+c)^2 - 3iB\tan(dx+c)^2 - 10iA\tan(dx+c) + 6B\tan(dx+c) - 3A - 5i}{a^2(\tan(dx+c)-i)^2}}{16d}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")`

output `1/16*(2*(A - I*B)*log(tan(d*x + c) + I)/a^2 - 2*(A - I*B)*log(tan(d*x + c) - I)/a^2 + (3*A*tan(d*x + c)^2 - 3*I*B*tan(d*x + c)^2 - 10*I*A*tan(d*x + c) + 6*B*tan(d*x + c) - 3*A - 5*I*B)/(a^2*(tan(d*x + c) - I)^2))/d`

3.46.9 Mupad [B] (verification not implemented)

Time = 8.13 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.39

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx = \frac{\frac{B}{2a^2} + \tan(c + dx) \left(\frac{A}{4a^2} + \frac{B3i}{4a^2} \right)}{d \left(\tan(c + dx)^2 li + 2 \tan(c + dx) - i \right)} + \frac{\ln(\tan(c + dx) - i) (B + A li) li}{8a^2 d} + \frac{\ln(\tan(c + dx) + li) (A - B li)}{8a^2 d}$$

3.46. $\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$

input `int((tan(c + d*x)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^2,x)`

output `(B/(2*a^2) + tan(c + d*x)*(A/(4*a^2) + (B*3i)/(4*a^2)))/(d*(2*tan(c + d*x) + tan(c + d*x)^2*1i - 1i)) + (log(tan(c + d*x) - 1i)*(A*1i + B)*1i)/(8*a^2*d) + (log(tan(c + d*x) + 1i)*(A - B*1i))/(8*a^2*d)`

$$3.47 \quad \int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^2} dx$$

3.47.1	Optimal result	654
3.47.2	Mathematica [A] (verified)	654
3.47.3	Rubi [A] (verified)	655
3.47.4	Maple [A] (verified)	656
3.47.5	Fricas [A] (verification not implemented)	657
3.47.6	Sympy [A] (verification not implemented)	657
3.47.7	Maxima [F(-2)]	658
3.47.8	Giac [A] (verification not implemented)	658
3.47.9	Mupad [B] (verification not implemented)	658

3.47.1 Optimal result

Integrand size = 26, antiderivative size = 80

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^2} dx = \frac{(A - iB)x}{4a^2} + \frac{iA - B}{4d(a + ia \tan(c + dx))^2} + \frac{iA + B}{4d(a^2 + ia^2 \tan(c + dx))}$$

output `1/4*(A-I*B)*x/a^2+1/4*(I*A-B)/d/(a+I*a*tan(d*x+c))^2+1/4*(I*A+B)/d/(a^2+I*a^2*tan(d*x+c))`

3.47.2 Mathematica [A] (verified)

Time = 0.85 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.11

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^2} dx = \frac{(A - iB) \arctan(\tan(c + dx))}{4a^2d} - \frac{iA - B}{4a^2d(i - \tan(c + dx))^2} - \frac{A - iB}{4a^2d(i - \tan(c + dx))}$$

input `Integrate[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x])^2,x]`

output `((A - I*B)*ArcTan[Tan[c + d*x]])/(4*a^2*d) - (I*A - B)/(4*a^2*d*(I - Tan[c + d*x])^2) - (A - I*B)/(4*a^2*d*(I - Tan[c + d*x]))`

3.47. $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^2} dx$

3.47.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.99, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$, Rules used = {3042, 4009, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{4009} \\
 & \frac{(A - iB) \int \frac{1}{i \tan(c+dx)a+a} dx}{2a} + \frac{-B + iA}{4d(a + ia \tan(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A - iB) \int \frac{1}{i \tan(c+dx)a+a} dx}{2a} + \frac{-B + iA}{4d(a + ia \tan(c + dx))^2} \\
 & \quad \downarrow \text{3960} \\
 & \frac{(A - iB) \left(\frac{\int 1 dx}{2a} + \frac{i}{2d(a+ia \tan(c+dx))} \right)}{2a} + \frac{-B + iA}{4d(a + ia \tan(c + dx))^2} \\
 & \quad \downarrow \text{24} \\
 & \frac{-B + iA}{4d(a + ia \tan(c + dx))^2} + \frac{(A - iB) \left(\frac{x}{2a} + \frac{i}{2d(a+ia \tan(c+dx))} \right)}{2a}
 \end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x])^2,x]`

output `(I*A - B)/(4*d*(a + I*a*Tan[c + d*x])^2) + ((A - I*B)*(x/(2*a) + (I/2)/(d*(a + I*a*Tan[c + d*x]))))/(2*a)`

3.47.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`
- rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

3.47.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.91

method	result
risch	$-\frac{ixB}{4a^2} + \frac{xA}{4a^2} + \frac{ie^{-2i(dx+c)}A}{4a^2d} - \frac{e^{-4i(dx+c)}B}{16a^2d} + \frac{ie^{-4i(dx+c)}A}{16a^2d}$
derivativedivides	$\frac{A \arctan(\tan(dx+c))}{4da^2} - \frac{iB \arctan(\tan(dx+c))}{4da^2} + \frac{A}{4da^2(\tan(dx+c)-i)} - \frac{iB}{4da^2(\tan(dx+c)-i)} - \frac{iA}{4da^2(\tan(dx+c))}$
default	$\frac{A \arctan(\tan(dx+c))}{4da^2} - \frac{iB \arctan(\tan(dx+c))}{4da^2} + \frac{A}{4da^2(\tan(dx+c)-i)} - \frac{iB}{4da^2(\tan(dx+c)-i)} - \frac{iA}{4da^2(\tan(dx+c))}$

input `int((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `-1/4*I*x/a^2*B+1/4*x/a^2*A+1/4*I/a^2/d*exp(-2*I*(d*x+c))*A-1/16/a^2/d*exp(-4*I*(d*x+c))*B+1/16*I/a^2/d*exp(-4*I*(d*x+c))*A`

3.47. $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^2} dx$

3.47.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.68

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^2} dx$$

$$= \frac{(4(A - iB)dx e^{(4i dx + 4i c)} + 4i A e^{(2i dx + 2i c)} + i A - B) e^{(-4i dx - 4i c)}}{16 a^2 d}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="fracas")`output `1/16*(4*(A - I*B)*d*x*e^(4*I*d*x + 4*I*c) + 4*I*A*e^(2*I*d*x + 2*I*c) + I*A - B)*e^(-4*I*d*x - 4*I*c)/(a^2*d)`**3.47.6 Sympy [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 162, normalized size of antiderivative = 2.02

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^2} dx$$

$$= \begin{cases} \frac{(16iAa^2 d e^{4ic} e^{-2idx} + (4iAa^2 d e^{2ic} - 4Ba^2 d e^{2ic}) e^{-4idx}) e^{-6ic}}{64a^4 d^2} & \text{for } a^4 d^2 e^{6ic} \neq 0 \\ x \left(-\frac{A-iB}{4a^2} + \frac{(Ae^{4ic} + 2Ae^{2ic} + A-iBe^{4ic} + iB)e^{-4ic}}{4a^2} \right) & \text{otherwise} \end{cases} + \frac{x(A - iB)}{4a^2}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**2,x)`output `Piecewise(((16*I*A*a**2*d*exp(4*I*c)*exp(-2*I*d*x) + (4*I*A*a**2*d*exp(2*I*c) - 4*B*a**2*d*exp(2*I*c))*exp(-4*I*d*x))*exp(-6*I*c)/(64*a**4*d**2), Ne(a**4*d**2*exp(6*I*c), 0)), (x*(-(A - I*B)/(4*a**2) + (A*exp(4*I*c) + 2*A*exp(2*I*c) + A - I*B*exp(4*I*c) + I*B)*exp(-4*I*c)/(4*a**2)), True)) + x*(A - I*B)/(4*a**2)`

3.47.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^2} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.47.8 Giac [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.38

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^2} dx = \frac{\frac{2(-iA-B)\log(\tan(dx+c)+i)}{a^2} - \frac{2(-iA-B)\log(\tan(dx+c)-i)}{a^2} - \frac{3iA \tan(dx+c)^2 + 3B \tan(dx+c)^2 + 10A \tan(dx+c) - 10iB \tan(dx+c)}{a^2(\tan(dx+c)-i)^2}}{16d}$$

```
input integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")
```

```
output -1/16*(2*(-I*A - B)*log(tan(d*x + c) + I)/a^2 - 2*(-I*A - B)*log(tan(d*x + c) - I)/a^2 - (3*I*A*tan(d*x + c)^2 + 3*B*tan(d*x + c)^2 + 10*A*tan(d*x + c) - 10*I*B*tan(d*x + c) - 11*I*A - 3*B)/(a^2*(tan(d*x + c) - I)^2)/d
```

3.47.9 Mupad [B] (verification not implemented)

Time = 7.44 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.88

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^2} dx = \frac{\frac{A}{2a^2} + \tan(c + dx) \left(\frac{B}{4a^2} + \frac{A1i}{4a^2} \right)}{d (\tan(c + dx)^2 1i + 2 \tan(c + dx) - i)} - \frac{x (B + A 1i) 1i}{4a^2}$$

```
input int((A + B*tan(c + d*x))/(a + a*tan(c + d*x)*1i)^2,x)
```

```
output (A/(2*a^2) + tan(c + d*x)*((A*1i)/(4*a^2) + B/(4*a^2)))/(d*(2*tan(c + d*x) + tan(c + d*x)^2*1i - 1i)) - (x*(A*1i + B)*1i)/(4*a^2)
```

3.47. $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^2} dx$

3.48 $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

3.48.1 Optimal result 659
 3.48.2 Mathematica [A] (verified) 659
 3.48.3 Rubi [A] (verified) 660
 3.48.4 Maple [A] (verified) 662
 3.48.5 Fricas [A] (verification not implemented) 663
 3.48.6 Sympy [A] (verification not implemented) 663
 3.48.7 Maxima [F(-2)] 664
 3.48.8 Giac [A] (verification not implemented) 664
 3.48.9 Mupad [B] (verification not implemented) 665

3.48.1 Optimal result

Integrand size = 32, antiderivative size = 95

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = -\frac{(3iA-B)x}{4a^2} + \frac{A \log(\sin(c+dx))}{a^2d} + \frac{3A+iB}{4a^2d(1+i \tan(c+dx))} + \frac{A+iB}{4d(a+ia \tan(c+dx))^2}$$

output `-1/4*(3*I*A-B)*x/a^2+A*ln(sin(d*x+c))/a^2/d+1/4*(3*A+I*B)/a^2/d/(1+I*tan(d*x+c))+1/4*(A+I*B)/d/(a+I*a*tan(d*x+c))^2`

3.48.2 Mathematica [A] (verified)

Time = 1.75 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.11

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = \frac{(7A+iB) \log(i-\tan(c+dx)) - 8A \log(\tan(c+dx)) + (A-iB) \log(i+\tan(c+dx)) + \frac{2(A+iB)}{(-i+\tan(c+dx))}}{8a^2d}$$

input `Integrate[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output
$$\frac{-1/8*((7*A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]] - 8*A*\text{Log}[\text{Tan}[c + d*x]] + (A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]] + (2*(A + I*B))/(-I + \text{Tan}[c + d*x])^2 - (2*((-3*I)*A + B))/(-I + \text{Tan}[c + d*x]))/(a^2*d)}$$

3.48.3 Rubi [A] (verified)

Time = 0.65 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.15, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3042, 4079, 27, 3042, 4079, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx \\ & \quad \downarrow 3042 \\ & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)(a+ia \tan(c+dx))^2} dx \\ & \quad \downarrow 4079 \\ & \frac{\int \frac{2 \cot(c+dx)(2aA-a(iA-B) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{4a^2} + \frac{A+iB}{4d(a+ia \tan(c+dx))^2} \\ & \quad \downarrow 27 \\ & \frac{\int \frac{\cot(c+dx)(2aA-a(iA-B) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{2a^2} + \frac{A+iB}{4d(a+ia \tan(c+dx))^2} \\ & \quad \downarrow 3042 \\ & \frac{\int \frac{2aA-a(iA-B) \tan(c+dx)}{\tan(c+dx)(i \tan(c+dx)a+a)} dx}{2a^2} + \frac{A+iB}{4d(a+ia \tan(c+dx))^2} \\ & \quad \downarrow 4079 \\ & \frac{\int \frac{\cot(c+dx)(4a^2A-a^2(3iA-B) \tan(c+dx))}{2a^2} dx}{2a^2} + \frac{3A+iB}{2d(1+i \tan(c+dx))} + \frac{A+iB}{4d(a+ia \tan(c+dx))^2} \\ & \quad \downarrow 3042 \\ & \frac{\int \frac{4a^2A-a^2(3iA-B) \tan(c+dx)}{\tan(c+dx)} dx}{2a^2} + \frac{3A+iB}{2d(1+i \tan(c+dx))} + \frac{A+iB}{4d(a+ia \tan(c+dx))^2} \end{aligned}$$

3.48.
$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$\begin{aligned}
& \downarrow 4014 \\
& \frac{\frac{4a^2 A \int \cot(c+dx) dx - a^2 x(-B+3iA)}{2a^2} + \frac{3A+iB}{2d(1+i \tan(c+dx))}}{2a^2} + \frac{A+iB}{4d(a+ia \tan(c+dx))^2} \\
& \downarrow 3042 \\
& \frac{\frac{4a^2 A \int -\tan(c+dx+\frac{\pi}{2}) dx - a^2 x(-B+3iA)}{2a^2} + \frac{3A+iB}{2d(1+i \tan(c+dx))}}{2a^2} + \frac{A+iB}{4d(a+ia \tan(c+dx))^2} \\
& \downarrow 25 \\
& \frac{\frac{-4a^2 A \int \tan(\frac{1}{2}(2c+\pi)+dx) dx - (a^2 x(-B+3iA))}{2a^2} + \frac{3A+iB}{2d(1+i \tan(c+dx))}}{2a^2} + \frac{A+iB}{4d(a+ia \tan(c+dx))^2} \\
& \downarrow 3956 \\
& \frac{\frac{4a^2 A \log\left(\frac{-\sin(c+dx)}{d}\right) - a^2 x(-B+3iA)}{2a^2} + \frac{3A+iB}{2d(1+i \tan(c+dx))}}{2a^2} + \frac{A+iB}{4d(a+ia \tan(c+dx))^2}
\end{aligned}$$

input `Int[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `((-(a^2*((3*I)*A - B)*x) + (4*a^2*A*Log[-Sin[c + d*x]])/d)/(2*a^2) + (3*A + I*B)/(2*d*(1 + I*Tan[c + d*x]))/(2*a^2) + (A + I*B)/(4*d*(a + I*a*Tan[c + d*x])^2)`

3.48.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

3.48. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4079 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.48.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.28

method	result
risch	$\frac{x B}{4 a^2} - \frac{7 i x A}{4 a^2} + \frac{i e^{-2 i(d x+c)} B}{4 a^2 d} + \frac{e^{-2 i(d x+c)} A}{2 a^2 d} + \frac{i e^{-4 i(d x+c)} B}{16 a^2 d} + \frac{e^{-4 i(d x+c)} A}{16 a^2 d} - \frac{2 i A c}{a^2 d} + \frac{A \ln(e^{2 i(d x+c)} - 1)}{a^2 d}$
derivativedivides	$-\frac{A}{4 d a^2(\tan(dx+c)-i)^2} - \frac{i B}{4 d a^2(\tan(dx+c)-i)^2} - \frac{A \ln(1+\tan^2(dx+c))}{2 d a^2} - \frac{3 i A \arctan(\tan(dx+c))}{4 d a^2} + \frac{B \arctan(\tan(dx+c))}{4 d a^2}$
default	$-\frac{A}{4 d a^2(\tan(dx+c)-i)^2} - \frac{i B}{4 d a^2(\tan(dx+c)-i)^2} - \frac{A \ln(1+\tan^2(dx+c))}{2 d a^2} - \frac{3 i A \arctan(\tan(dx+c))}{4 d a^2} + \frac{B \arctan(\tan(dx+c))}{4 d a^2}$

```
input int(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x,method=_RETURNVERBO
SE)
```

```
output 1/4*x/a^2*B-7/4*I*x/a^2*A+1/4*I*B/a^2/d*exp(-2*I*(d*x+c))+1/2/a^2/d*exp(-2
*I*(d*x+c))*A+1/16*I/a^2/d*exp(-4*I*(d*x+c))*B+1/16/a^2/d*exp(-4*I*(d*x+c)
)*A-2*I/a^2/d*A*c+1/a^2*A/d*ln(exp(2*I*(d*x+c))-1)
```

$$3.48. \int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.48.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.91

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = \frac{(4(7iA-B)dx e^{(4i dx+4i c)} - 16Ae^{(4i dx+4i c)} \log(e^{(2i dx+2i c)} - 1) - 4(2A+iB)e^{(2i dx+2i c)} - A-iB)e^{-4i dx-4i c}}{16a^2 d}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="fracas")`

output `-1/16*(4*(7*I*A - B)*d*x*e^(4*I*d*x + 4*I*c) - 16*A*e^(4*I*d*x + 4*I*c)*log(e^(2*I*d*x + 2*I*c) - 1) - 4*(2*A + I*B)*e^(2*I*d*x + 2*I*c) - A - I*B)*e^(-4*I*d*x - 4*I*c)/(a^2*d)`

3.48.6 Sympy [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 219, normalized size of antiderivative = 2.31

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = \frac{A \log(e^{2idx} - e^{-2ic})}{a^2 d} + \begin{cases} \frac{((4Aa^2 de^{2ic} + 4iBa^2 de^{2ic})e^{-4idx} + (32Aa^2 de^{4ic} + 16iBa^2 de^{4ic})e^{-2idx})e^{-6ic}}{64a^4 d^2} & \text{for } a^4 d^2 e^{6ic} \neq 0 \\ x \left(-\frac{7iA+B}{4a^2} + \frac{(-7iAe^{4ic} - 4iAe^{2ic} - iA + Be^{4ic} + 2Be^{2ic} + B)e^{-4ic}}{4a^2} \right) & \text{otherwise} \end{cases} + \frac{x(-7iA+B)}{4a^2}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**2,x)`

output `A*log(exp(2*I*d*x) - exp(-2*I*c))/(a**2*d) + Piecewise((((4*A*a**2*d*exp(2*I*c) + 4*I*B*a**2*d*exp(2*I*c))*exp(-4*I*d*x) + (32*A*a**2*d*exp(4*I*c) + 16*I*B*a**2*d*exp(4*I*c))*exp(-2*I*d*x))*exp(-6*I*c)/(64*a**4*d**2), Ne(a**4*d**2*exp(6*I*c), 0)), (x*(-(-7*I*A + B)/(4*a**2) + (-7*I*A*exp(4*I*c) - 4*I*A*exp(2*I*c) - I*A + B*exp(4*I*c) + 2*B*exp(2*I*c) + B)*exp(-4*I*c)/(4*a**2)), True)) + x*(-7*I*A + B)/(4*a**2)`

3.48.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.48.8 Giac [A] (verification not implemented)

Time = 0.60 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.27

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx =$$

$$\frac{\frac{2(A-iB)\log(\tan(dx+c)+i)}{a^2} + \frac{2(7A+iB)\log(\tan(dx+c)-i)}{a^2} - \frac{16A\log(\tan(dx+c))}{a^2} - \frac{21A\tan(dx+c)^2+3iB\tan(dx+c)^2-54iA\tan(dx+c)-16d}{a^2(\tan(dx+c)-i)^2}}{16d}$$

```
input integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")
```

```
output -1/16*(2*(A - I*B)*log(tan(d*x + c) + I)/a^2 + 2*(7*A + I*B)*log(tan(d*x + c) - I)/a^2 - 16*A*log(tan(d*x + c))/a^2 - (21*A*tan(d*x + c)^2 + 3*I*B*tan(d*x + c)^2 - 54*I*A*tan(d*x + c) + 10*B*tan(d*x + c) - 37*A - 11*I*B)/(a^2*(tan(d*x + c) - I)^2))/d
```

3.48.9 Mupad [B] (verification not implemented)

Time = 7.37 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.36

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \frac{\frac{B}{2a^2} - \frac{A1i}{a^2} + \tan(c+dx) \left(\frac{3A}{4a^2} + \frac{B1i}{4a^2}\right)}{d(\tan(c+dx)^2 1i + 2\tan(c+dx) - i)} + \frac{A \ln(\tan(c+dx))}{a^2 d} - \frac{\ln(\tan(c+dx) + 1i)(A - B 1i)}{8a^2 d} - \frac{\ln(\tan(c+dx) - i)(7A + B 1i)}{8a^2 d}$$

input `int((cot(c + d*x)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^2,x)`output `(B/(2*a^2) - (A*1i)/a^2 + tan(c + d*x)*((3*A)/(4*a^2) + (B*1i)/(4*a^2)))/(d*(2*tan(c + d*x) + tan(c + d*x)^2*1i - 1i)) + (A*log(tan(c + d*x)))/(a^2*d) - (log(tan(c + d*x) + 1i)*(A - B*1i))/(8*a^2*d) - (log(tan(c + d*x) - 1i)*(7*A + B*1i))/(8*a^2*d)`

$$3.49 \quad \int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.49.1	Optimal result	666
3.49.2	Mathematica [C] (verified)	666
3.49.3	Rubi [A] (verified)	667
3.49.4	Maple [A] (verified)	670
3.49.5	Fricas [A] (verification not implemented)	671
3.49.6	Sympy [A] (verification not implemented)	671
3.49.7	Maxima [F(-2)]	672
3.49.8	Giac [A] (verification not implemented)	672
3.49.9	Mupad [B] (verification not implemented)	673

3.49.1 Optimal result

Integrand size = 34, antiderivative size = 141

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = -\frac{3(3A+iB)x}{4a^2} - \frac{3(3A+iB) \cot(c+dx)}{4a^2d} - \frac{(2iA-B) \log(\sin(c+dx))}{a^2d} + \frac{(2A+iB) \cot(c+dx)}{2a^2d(1+i \tan(c+dx))} + \frac{(A+iB) \cot(c+dx)}{4d(a+ia \tan(c+dx))^2}$$

output
$$-3/4*(3*A+I*B)*x/a^2-3/4*(3*A+I*B)*\cot(d*x+c)/a^2/d-(2*I*A-B)*\ln(\sin(d*x+c))/a^2/d+1/2*(2*A+I*B)*\cot(d*x+c)/a^2/d/(1+I*\tan(d*x+c))+1/4*(A+I*B)*\cot(d*x+c)/d/(a+I*a*\tan(d*x+c))^2$$

3.49.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 2.20 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.92

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = \frac{2(A+iB) \cot^3(c+dx)}{(i+\cot(c+dx))^2} + \frac{4(2A+iB) \cot^2(c+dx)}{i+\cot(c+dx)} - \frac{6(3A+iB) \cot(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(c+dx)\right)}{8a^2d}$$

3.49.
$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

input `Integrate[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `((2*(A + I*B)*Cot[c + d*x]^3)/(I + Cot[c + d*x])^2 + (4*(2*A + I*B)*Cot[c + d*x]^2)/(I + Cot[c + d*x]) - 6*(3*A + I*B)*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2] + 8*((-2*I)*A + B)*(Log[Cos[c + d*x]] + Log[Tan[c + d*x]]))/(8*a^2*d)`

3.49.3 Rubi [A] (verified)

Time = 0.92 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.04, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.382$, Rules used = {3042, 4079, 3042, 4079, 27, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^2(a+ia \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{\cot^2(c+dx)(a(5A+iB)-3a(iA-B) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{4a^2} + \frac{(A+iB) \cot(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a(5A+iB)-3a(iA-B) \tan(c+dx)}{\tan(c+dx)^2(i \tan(c+dx)a+a)} dx}{4a^2} + \frac{(A+iB) \cot(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{2 \cot^2(c+dx)(3a^2(3A+iB)-4a^2(2iA-B) \tan(c+dx))}{2a^2} dx}{4a^2} + \frac{2(2A+iB) \cot(c+dx)}{d(1+i \tan(c+dx))} + \frac{(A+iB) \cot(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\cot^2(c+dx)(3a^2(3A+iB)-4a^2(2iA-B) \tan(c+dx))}{a^2} dx}{4a^2} + \frac{2(2A+iB) \cot(c+dx)}{d(1+i \tan(c+dx))} + \frac{(A+iB) \cot(c+dx)}{4d(a+ia \tan(c+dx))^2}
 \end{aligned}$$

3.49. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{\int \frac{3a^2(3A+iB)-4a^2(2iA-B)\tan(c+dx)}{\tan(c+dx)^2} dx + \frac{2(2A+iB)\cot(c+dx)}{d(1+i\tan(c+dx))}}{4a^2} + \frac{(A+iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} \\
& \downarrow 4012 \\
& \frac{\int -\cot(c+dx)(4(2iA-B)a^2+3(3A+iB)\tan(c+dx)a^2) dx - \frac{3a^2(3A+iB)\cot(c+dx)}{d} + \frac{2(2A+iB)\cot(c+dx)}{d(1+i\tan(c+dx))}}{4a^2} + \\
& \quad \frac{(A+iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} \\
& \downarrow 25 \\
& \frac{-\int \cot(c+dx)(4(2iA-B)a^2+3(3A+iB)\tan(c+dx)a^2) dx - \frac{3a^2(3A+iB)\cot(c+dx)}{d} + \frac{2(2A+iB)\cot(c+dx)}{d(1+i\tan(c+dx))}}{4a^2} + \\
& \quad \frac{(A+iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} \\
& \downarrow 3042 \\
& \frac{-\int \frac{4(2iA-B)a^2+3(3A+iB)\tan(c+dx)a^2}{\tan(c+dx)} dx - \frac{3a^2(3A+iB)\cot(c+dx)}{d} + \frac{2(2A+iB)\cot(c+dx)}{d(1+i\tan(c+dx))}}{4a^2} + \frac{(A+iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} \\
& \downarrow 4014 \\
& \frac{-4a^2(-B+2iA)\int \cot(c+dx) dx - \frac{3a^2(3A+iB)\cot(c+dx)}{d} - 3a^2x(3A+iB) + \frac{2(2A+iB)\cot(c+dx)}{d(1+i\tan(c+dx))}}{4a^2} + \\
& \quad \frac{(A+iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} \\
& \downarrow 3042 \\
& \frac{-4a^2(-B+2iA)\int -\tan(c+dx+\frac{\pi}{2}) dx - \frac{3a^2(3A+iB)\cot(c+dx)}{d} - 3a^2x(3A+iB) + \frac{2(2A+iB)\cot(c+dx)}{d(1+i\tan(c+dx))}}{4a^2} + \\
& \quad \frac{(A+iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} \\
& \downarrow 25 \\
& \frac{4a^2(-B+2iA)\int \tan(\frac{1}{2}(2c+\pi)+dx) dx - \frac{3a^2(3A+iB)\cot(c+dx)}{d} - 3a^2x(3A+iB) + \frac{2(2A+iB)\cot(c+dx)}{d(1+i\tan(c+dx))}}{4a^2} + \\
& \quad \frac{(A+iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} \\
& \downarrow 3956
\end{aligned}$$

3.49. $\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$

$$\frac{-\frac{3a^2(3A+iB)\cot(c+dx)}{d} - \frac{4a^2(-B+2iA)\log(-\sin(c+dx))}{d} - 3a^2x(3A+iB)}{a^2} + \frac{2(2A+iB)\cot(c+dx)}{d(1+i\tan(c+dx))} + \frac{4a^2(A+iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2}$$

input `Int[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `((-3*a^2*(3*A + I*B)*x - (3*a^2*(3*A + I*B)*Cot[c + d*x])/d - (4*a^2*((2*I)*A - B)*Log[-Sin[c + d*x]])/d)/a^2 + (2*(2*A + I*B)*Cot[c + d*x])/(d*(1 + I*Tan[c + d*x]))/(4*a^2) + ((A + I*B)*Cot[c + d*x])/(4*d*(a + I*a*Tan[c + d*x])^2)`

3.49.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)
)*(x_)], x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4079 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.49.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.26

method	result
risch	$-\frac{7ixB}{4a^2} - \frac{17xA}{4a^2} + \frac{e^{-2i(dx+c)}B}{2a^2d} - \frac{3ie^{-2i(dx+c)}A}{4a^2d} + \frac{e^{-4i(dx+c)}B}{16a^2d} - \frac{ie^{-4i(dx+c)}A}{16a^2d} - \frac{2iBc}{a^2d} - \frac{4Ac}{a^2d} - \frac{1}{a^2d} \ln(e^{2i(dx+c)} - 1)$
derivativedivides	$\frac{iA}{4da^2(\tan(dx+c)-i)^2} - \frac{B}{4da^2(\tan(dx+c)-i)^2} - \frac{B \ln(1+\tan^2(dx+c))}{2da^2} - \frac{3iB \arctan(\tan(dx+c))}{4da^2} + \frac{iA \ln(1+\tan^2(dx+c))}{a^2d}$
default	$\frac{iA}{4da^2(\tan(dx+c)-i)^2} - \frac{B}{4da^2(\tan(dx+c)-i)^2} - \frac{B \ln(1+\tan^2(dx+c))}{2da^2} - \frac{3iB \arctan(\tan(dx+c))}{4da^2} + \frac{iA \ln(1+\tan^2(dx+c))}{a^2d}$

```
input int(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x,method=_RETURNVER
BOSE)
```

```
output -7/4*I*x/a^2*B-17/4*x/a^2*A+1/2/a^2/d*exp(-2*I*(d*x+c))*B-3/4*I/a^2/d*exp(
-2*I*(d*x+c))*A+1/16/a^2/d*exp(-4*I*(d*x+c))*B-1/16*I/a^2/d*exp(-4*I*(d*x+
c))*A-2*I*B/a^2/d*c-4/a^2/d*A*c-2*I*A/a^2/d/(exp(2*I*(d*x+c))-1)+1/a^2/d*ln
(exp(2*I*(d*x+c))-1)*B-2*I/a^2/d*ln(exp(2*I*(d*x+c))-1)*A
```

$$3.49. \int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.49.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.08

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx =$$

$$\frac{4(17A+7iB)dx e^{(6i dx+6i c)} - 4((17A+7iB)dx - 11iA+2B)e^{(4i dx+4i c)} - (11iA-7B)e^{(2i dx+2i c)}}{16(a^2 d e^{(6i dx+6i c)} - a^2 d e^{(4i dx+4i c)})}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="fracas")`

output `-1/16*(4*(17*A + 7*I*B)*d*x*e^(6*I*d*x + 6*I*c) - 4*((17*A + 7*I*B)*d*x - 11*I*A + 2*B)*e^(4*I*d*x + 4*I*c) - (11*I*A - 7*B)*e^(2*I*d*x + 2*I*c) + 16*((2*I*A - B)*e^(6*I*d*x + 6*I*c) + (-2*I*A + B)*e^(4*I*d*x + 4*I*c))*log(e^(2*I*d*x + 2*I*c) - 1) - I*A + B)/(a^2*d*e^(6*I*d*x + 6*I*c) - a^2*d*e^(4*I*d*x + 4*I*c))`

3.49.6 Sympy [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.89

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$= -\frac{2iA}{a^2 d e^{2ic} e^{2idx} - a^2 d}$$

$$+ \begin{cases} \frac{((-4iAa^2 d e^{2ic} + 4Ba^2 d e^{2ic})e^{-4idx} + (-48iAa^2 d e^{4ic} + 32Ba^2 d e^{4ic})e^{-2idx})e^{-6ic}}{64a^4 d^2} & \text{for } a^4 d^2 e^{6ic} \neq 0 \\ x \left(-\frac{17A-7iB}{4a^2} + \frac{(-17Ae^{4ic} - 6Ae^{2ic} - A - 7iBe^{4ic} - 4iBe^{2ic} - iB)e^{-4ic}}{4a^2} \right) & \text{otherwise} \end{cases}$$

$$+ \frac{x(-17A-7iB)}{4a^2} - \frac{i(2A+iB) \log(e^{2idx} - e^{-2ic})}{a^2 d}$$

input `integrate(cot(d*x+c)**2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**2,x)`

```
output -2*I*A/(a**2*d*exp(2*I*c)*exp(2*I*d*x) - a**2*d) + Piecewise(((((-4*I*A*a**
2*d*exp(2*I*c) + 4*B*a**2*d*exp(2*I*c))*exp(-4*I*d*x) + (-48*I*A*a**2*d*ex
p(4*I*c) + 32*B*a**2*d*exp(4*I*c))*exp(-2*I*d*x))*exp(-6*I*c)/(64*a**4*d**
2), Ne(a**4*d**2*exp(6*I*c), 0)), (x*(-(-17*A - 7*I*B)/(4*a**2) + (-17*A*exp
(4*I*c) - 6*A*exp(2*I*c) - A - 7*I*B*exp(4*I*c) - 4*I*B*exp(2*I*c) - I*B
)*exp(-4*I*c)/(4*a**2)), True)) + x*(-17*A - 7*I*B)/(4*a**2) - I*(2*A + I*
B)*log(exp(2*I*d*x) - exp(-2*I*c))/(a**2*d)
```

3.49.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm=
"maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.49.8 Giac [A] (verification not implemented)

Time = 0.82 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.14

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx$$

$$= \frac{\frac{2(-iA-B)\log(\tan(dx+c)+i)}{a^2} - \frac{2(-17iA+7B)\log(\tan(dx+c)-i)}{a^2} - \frac{16(2iA-B)\log(\tan(dx+c))}{a^2} - \frac{16(-2iA\tan(dx+c)+B\tan(dx+c)+A)}{a^2\tan(dx+c)}}{16d}$$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm=
"giac")
```

```
output 1/16*(2*(-I*A - B)*log(tan(d*x + c) + I)/a^2 - 2*(-17*I*A + 7*B)*log(tan(d
*x + c) - I)/a^2 - 16*(2*I*A - B)*log(tan(d*x + c))/a^2 - 16*(-2*I*A*tan(d
*x + c) + B*tan(d*x + c) + A)/(a^2*tan(d*x + c)) - (51*I*A*tan(d*x + c)^2
- 21*B*tan(d*x + c)^2 + 122*A*tan(d*x + c) + 54*I*B*tan(d*x + c) - 75*I*A
+ 37*B)/(a^2*(tan(d*x + c) - I)^2))/d
```

3.49. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

3.49.9 Mupad [B] (verification not implemented)

Time = 8.02 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.16

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= -\frac{\tan(c+dx)^2\left(-\frac{3B}{4a^2} + \frac{A9i}{4a^2}\right) - \frac{A1i}{a^2} + \tan(c+dx)\left(\frac{7A}{2a^2} + \frac{B1i}{a^2}\right)}{d\left(\tan(c+dx)^3 1i + 2\tan(c+dx)^2 - \tan(c+dx) 1i\right)}$$

$$- \frac{\ln(\tan(c+dx))(-B+A2i)}{a^2 d} - \frac{\ln(\tan(c+dx)+1i)(B+A1i)}{8a^2 d}$$

$$+ \frac{\ln(\tan(c+dx)-i)(-7B+A17i)}{8a^2 d}$$

```
input int((cot(c + d*x)^2*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^2,x)
```

```
output (log(tan(c + d*x) - 1i)*(A*17i - 7*B))/(8*a^2*d) - (log(tan(c + d*x))*(A*2
i - B))/(a^2*d) - (log(tan(c + d*x) + 1i)*(A*1i + B))/(8*a^2*d) - (tan(c +
d*x)^2*((A*9i)/(4*a^2) - (3*B)/(4*a^2)) - (A*1i)/a^2 + tan(c + d*x)*((7*A
)/(2*a^2) + (B*1i)/a^2))/(d*(2*tan(c + d*x)^2 - tan(c + d*x)*1i + tan(c +
d*x)^3*1i))
```


3.50 $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

3.50.1	Optimal result	674
3.50.2	Mathematica [C] (verified)	675
3.50.3	Rubi [A] (verified)	675
3.50.4	Maple [A] (verified)	679
3.50.5	Fricas [A] (verification not implemented)	679
3.50.6	Sympy [A] (verification not implemented)	680
3.50.7	Maxima [F(-2)]	681
3.50.8	Giac [A] (verification not implemented)	681
3.50.9	Mupad [B] (verification not implemented)	682

3.50.1 Optimal result

Integrand size = 34, antiderivative size = 170

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = \frac{3(5iA-3B)x}{4a^2} + \frac{3(5iA-3B) \cot(c+dx)}{4a^2d} - \frac{(2A+iB) \cot^2(c+dx)}{a^2d} - \frac{2(2A+iB) \log(\sin(c+dx))}{a^2d} + \frac{(5A+3iB) \cot^2(c+dx)}{4a^2d(1+i \tan(c+dx))} + \frac{(A+iB) \cot^2(c+dx)}{4d(a+ia \tan(c+dx))^2}$$

```
output 3/4*(5*I*A-3*B)*x/a^2+3/4*(5*I*A-3*B)*cot(d*x+c)/a^2/d-(2*A+I*B)*cot(d*x+c)^2/a^2/d-2*(2*A+I*B)*ln(sin(d*x+c))/a^2/d+1/4*(5*A+3*I*B)*cot(d*x+c)^2/a^2/d/(1+I*tan(d*x+c))+1/4*(A+I*B)*cot(d*x+c)^2/d/(a+I*a*tan(d*x+c))^2
```

3.50.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 2.71 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.84

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$= \frac{\frac{2(A+iB) \cot^4(c+dx)}{(i+\cot(c+dx))^2} + \frac{2(5A+3iB) \cot^3(c+dx)}{i+\cot(c+dx)} + 6(5iA-3B) \cot(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(c+dx)\right)}{8a^2d}$$

input `Integrate[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `((2*(A + I*B)*Cot[c + d*x]^4)/(I + Cot[c + d*x])^2 + (2*(5*A + (3*I)*B)*Cot[c + d*x]^3)/(I + Cot[c + d*x]) + 6*((5*I)*A - 3*B)*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2] - 8*(2*A + I*B)*(Cot[c + d*x]^2 + 2*(Log[Cos[c + d*x]] + Log[Tan[c + d*x]])))/(8*a^2*d)`

3.50.3 Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.06, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.441$, Rules used = {3042, 4079, 27, 3042, 4079, 3042, 4012, 25, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A+B \tan(c+dx)}{\tan(c+dx)^3(a+ia \tan(c+dx))^2} dx$$

$$\downarrow \text{4079}$$

$$\frac{\int \frac{2 \cot^3(c+dx)(a(3A+iB)-2a(iA-B) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{4a^2} + \frac{(A+iB) \cot^2(c+dx)}{4d(a+ia \tan(c+dx))^2}$$

$$\downarrow \text{27}$$

3.50. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
& \frac{\int \frac{\cot^3(c+dx)(a(3A+iB)-2a(iA-B)\tan(c+dx))}{i \tan(c+dx)a+a} dx}{2a^2} + \frac{(A+iB)\cot^2(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{a(3A+iB)-2a(iA-B)\tan(c+dx)}{\tan(c+dx)^3(i \tan(c+dx)a+a)} dx}{2a^2} + \frac{(A+iB)\cot^2(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
& \quad \downarrow 4079 \\
& \frac{\int \cot^3(c+dx)(8a^2(2A+iB)-3a^2(5iA-3B)\tan(c+dx)) dx}{2a^2} + \frac{(5A+3iB)\cot^2(c+dx)}{2d(1+i \tan(c+dx))} + \frac{(A+iB)\cot^2(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{8a^2(2A+iB)-3a^2(5iA-3B)\tan(c+dx)}{\tan(c+dx)^3} dx}{2a^2} + \frac{(5A+3iB)\cot^2(c+dx)}{2d(1+i \tan(c+dx))} + \frac{(A+iB)\cot^2(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
& \quad \downarrow 4012 \\
& \frac{\int -\cot^2(c+dx)(3(5iA-3B)a^2+8(2A+iB)\tan(c+dx)a^2) dx - \frac{4a^2(2A+iB)\cot^2(c+dx)}{d}}{2a^2} + \frac{(5A+3iB)\cot^2(c+dx)}{2d(1+i \tan(c+dx))} + \\
& \quad \frac{2a^2}{4d(a+ia \tan(c+dx))^2} \\
& \quad \downarrow 25 \\
& \frac{-\int \cot^2(c+dx)(3(5iA-3B)a^2+8(2A+iB)\tan(c+dx)a^2) dx - \frac{4a^2(2A+iB)\cot^2(c+dx)}{d}}{2a^2} + \frac{(5A+3iB)\cot^2(c+dx)}{2d(1+i \tan(c+dx))} + \\
& \quad \frac{2a^2}{4d(a+ia \tan(c+dx))^2} \\
& \quad \downarrow 3042 \\
& \frac{-\int \frac{3(5iA-3B)a^2+8(2A+iB)\tan(c+dx)a^2}{\tan(c+dx)^2} dx - \frac{4a^2(2A+iB)\cot^2(c+dx)}{d}}{2a^2} + \frac{(5A+3iB)\cot^2(c+dx)}{2d(1+i \tan(c+dx))} + \\
& \quad \frac{2a^2}{4d(a+ia \tan(c+dx))^2} \\
& \quad \downarrow 4012 \\
& \frac{-\int \cot(c+dx)(8a^2(2A+iB)-3a^2(5iA-3B)\tan(c+dx)) dx - \frac{4a^2(2A+iB)\cot^2(c+dx)}{d} + \frac{3a^2(-3B+5iA)\cot(c+dx)}{d}}{2a^2} + \frac{(5A+3iB)\cot^2(c+dx)}{2d(1+i \tan(c+dx))} + \\
& \quad \frac{2a^2}{4d(a+ia \tan(c+dx))^2}
\end{aligned}$$

3.50. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{-\int \frac{8a^2(2A+iB)-3a^2(5iA-3B)\tan(c+dx)}{\tan(c+dx)} dx - \frac{4a^2(2A+iB)\cot^2(c+dx)}{d} + \frac{3a^2(-3B+5iA)\cot(c+dx)}{d}}{2a^2} + \frac{(5A+3iB)\cot^2(c+dx)}{2d(1+i\tan(c+dx))} + \\ & \frac{2a^2}{4d(a+ia\tan(c+dx))^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 4014 \\ & \frac{-8a^2(2A+iB)\int \cot(c+dx)dx - \frac{4a^2(2A+iB)\cot^2(c+dx)}{d} + \frac{3a^2(-3B+5iA)\cot(c+dx)}{d} + 3a^2x(-3B+5iA)}{2a^2} + \frac{(5A+3iB)\cot^2(c+dx)}{2d(1+i\tan(c+dx))} + \\ & \frac{2a^2}{4d(a+ia\tan(c+dx))^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{-8a^2(2A+iB)\int -\tan(c+dx+\frac{\pi}{2})dx - \frac{4a^2(2A+iB)\cot^2(c+dx)}{d} + \frac{3a^2(-3B+5iA)\cot(c+dx)}{d} + 3a^2x(-3B+5iA)}{2a^2} + \frac{(5A+3iB)\cot^2(c+dx)}{2d(1+i\tan(c+dx))} + \\ & \frac{2a^2}{4d(a+ia\tan(c+dx))^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 25 \\ & \frac{8a^2(2A+iB)\int \tan(\frac{1}{2}(2c+\pi)+dx)dx - \frac{4a^2(2A+iB)\cot^2(c+dx)}{d} + \frac{3a^2(-3B+5iA)\cot(c+dx)}{d} + 3a^2x(-3B+5iA)}{2a^2} + \frac{(5A+3iB)\cot^2(c+dx)}{2d(1+i\tan(c+dx))} + \\ & \frac{2a^2}{4d(a+ia\tan(c+dx))^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 3956 \\ & \frac{-\frac{4a^2(2A+iB)\cot^2(c+dx)}{d} + \frac{3a^2(-3B+5iA)\cot(c+dx)}{d} - \frac{8a^2(2A+iB)\log(-\sin(c+dx))}{d} + 3a^2x(-3B+5iA)}{2a^2} + \frac{(5A+3iB)\cot^2(c+dx)}{2d(1+i\tan(c+dx))} + \\ & \frac{2a^2}{4d(a+ia\tan(c+dx))^2} \end{aligned}$$

input `Int[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

$$3.50. \quad \int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

```
output ((3*a^2*((5*I)*A - 3*B)*x + (3*a^2*((5*I)*A - 3*B)*Cot[c + d*x])/d - (4*a^
2*(2*A + I*B)*Cot[c + d*x]^2)/d - (8*a^2*(2*A + I*B)*Log[-Sin[c + d*x]])/d
)/(2*a^2) + ((5*A + (3*I)*B)*Cot[c + d*x]^2)/(2*d*(1 + I*Tan[c + d*x]))/(
2*a^2) + ((A + I*B)*Cot[c + d*x]^2)/(4*d*(a + I*a*Tan[c + d*x])^2)
```

3.50.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4012 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]
```

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4079 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.50.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.23

method	result
risch	$-\frac{17xB}{4a^2} + \frac{31ixA}{4a^2} - \frac{3ie^{-2i(dx+c)}B}{4a^2d} - \frac{e^{-2i(dx+c)}A}{a^2d} - \frac{ie^{-4i(dx+c)}B}{16a^2d} - \frac{e^{-4i(dx+c)}A}{16a^2d} - \frac{4Bc}{a^2d} + \frac{8iAc}{a^2d} - \frac{2i(-}{da^2}$
derivativedivides	$\frac{A}{4da^2(\tan(dx+c)-i)^2} + \frac{iB}{4da^2(\tan(dx+c)-i)^2} - \frac{5B}{4da^2(\tan(dx+c)-i)} + \frac{15iA \arctan(\tan(dx+c))}{4da^2} + \frac{2A \ln(1+\tan}{da^2}$
default	$\frac{A}{4da^2(\tan(dx+c)-i)^2} + \frac{iB}{4da^2(\tan(dx+c)-i)^2} - \frac{5B}{4da^2(\tan(dx+c)-i)} + \frac{15iA \arctan(\tan(dx+c))}{4da^2} + \frac{2A \ln(1+\tan}{da^2}$

```
input int(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x,method=_RETURNVER
BOSE)
```

```
output -17/4*x/a^2*B+31/4*I*x/a^2*A-3/4*I/a^2/d*exp(-2*I*(d*x+c))*B-1/a^2/d*exp(-
2*I*(d*x+c))*A-1/16*I/a^2/d*exp(-4*I*(d*x+c))*B-1/16/a^2/d*exp(-4*I*(d*x+c
))*A-4/a^2/d*B*c+8*I/a^2/d*A*c-2*I*(-I*A*exp(2*I*(d*x+c))+B*exp(2*I*(d*x+c
)))+2*I*A-B)/a^2/d/(exp(2*I*(d*x+c))-1)^2-2*I/a^2/d*ln(exp(2*I*(d*x+c))-1)*
B-4/a^2*A/d*ln(exp(2*I*(d*x+c))-1)
```

3.50.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.26

$$\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx = \frac{4(-31iA + 17B)dx e^{8i dx + 8i c} + 4(2(31iA - 17B)dx + 12A + 11iB)e^{6i dx + 6i c} + 4(-31iA + 17B)}{...}$$

3.50. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="fricas")`

output `-1/16*(4*(-31*I*A + 17*B)*d*x*e^(8*I*d*x + 8*I*c) + 4*(2*(31*I*A - 17*B)*d*x + 12*A + 11*I*B)*e^(6*I*d*x + 6*I*c) + (4*(-31*I*A + 17*B)*d*x - 95*A - 55*I*B)*e^(4*I*d*x + 4*I*c) + 2*(7*A + 5*I*B)*e^(2*I*d*x + 2*I*c) + 32*((2*A + I*B)*e^(8*I*d*x + 8*I*c) - 2*(2*A + I*B)*e^(6*I*d*x + 6*I*c) + (2*A + I*B)*e^(4*I*d*x + 4*I*c))*log(e^(2*I*d*x + 2*I*c) - 1) + A + I*B)/(a^2*d*e^(8*I*d*x + 8*I*c) - 2*a^2*d*e^(6*I*d*x + 6*I*c) + a^2*d*e^(4*I*d*x + 4*I*c))`

3.50.6 Sympy [A] (verification not implemented)

Time = 0.48 (sec) , antiderivative size = 323, normalized size of antiderivative = 1.90

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$= \frac{4A + 2iB + (-2Ae^{2ic} - 2iBe^{2ic})e^{2idx}}{a^2de^{4ic}e^{4idx} - 2a^2de^{2ic}e^{2idx} + a^2d}$$

$$+ \begin{cases} \frac{((-4Aa^2de^{2ic} - 4iBa^2de^{2ic})e^{-4idx} + (-64Aa^2de^{4ic} - 48iBa^2de^{4ic})e^{-2idx})e^{-6ic}}{64a^4d^2} & \text{for } a^4d^2e^{6ic} \neq 0 \\ x \left(-\frac{31iA-17B}{4a^2} + \frac{(31iAe^{4ic} + 8iAe^{2ic} + iA - 17Be^{4ic} - 6Be^{2ic} - B)e^{-4ic}}{4a^2} \right) & \text{otherwise} \end{cases}$$

$$+ \frac{x(31iA - 17B)}{4a^2} - \frac{2 \cdot (2A + iB) \log(e^{2idx} - e^{-2ic})}{a^2d}$$

input `integrate(cot(d*x+c)**3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**2,x)`

output `(4*A + 2*I*B + (-2*A*exp(2*I*c) - 2*I*B*exp(2*I*c))*exp(2*I*d*x))/(a**2*d*exp(4*I*c)*exp(4*I*d*x) - 2*a**2*d*exp(2*I*c)*exp(2*I*d*x) + a**2*d) + Piecewise(((((-4*A*a**2*d*exp(2*I*c) - 4*I*B*a**2*d*exp(2*I*c))*exp(-4*I*d*x) + (-64*A*a**2*d*exp(4*I*c) - 48*I*B*a**2*d*exp(4*I*c))*exp(-2*I*d*x))*exp(-6*I*c)/(64*a**4*d**2), Ne(a**4*d**2*exp(6*I*c), 0)), (x*(-(31*I*A - 17*B)/(4*a**2) + (31*I*A*exp(4*I*c) + 8*I*A*exp(2*I*c) + I*A - 17*B*exp(4*I*c) - 6*B*exp(2*I*c) - B)*exp(-4*I*c)/(4*a**2)), True)) + x*(31*I*A - 17*B)/(4*a**2) - 2*(2*A + I*B)*log(exp(2*I*d*x) - exp(-2*I*c))/(a**2*d)`

3.50.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm=
"maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.50.8 Giac [A] (verification not implemented)

Time = 1.13 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.04

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$\frac{4(A-iB)\log(\tan(dx+c)+i)}{a^2} + \frac{4(31A+17iB)\log(\tan(dx+c)-i)}{a^2} - \frac{64(2A+iB)\log(\tan(dx+c))}{a^2} + \frac{3A\tan(dx+c)^4-3iB\tan(dx+c)^4+114A\tan(dx+c)^3-114iB\tan(dx+c)^3+173A\tan(dx+c)^2+115iB\tan(dx+c)^2-32iA\tan(dx+c)+32B\tan(dx+c)+16A}{32d}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm=
"giac")
```

```
output 1/32*(4*(A - I*B)*log(tan(d*x + c) + I)/a^2 + 4*(31*A + 17*I*B)*log(tan(d*
x + c) - I)/a^2 - 64*(2*A + I*B)*log(tan(d*x + c))/a^2 + (3*A*tan(d*x + c)
^4 - 3*I*B*tan(d*x + c)^4 + 114*I*A*tan(d*x + c)^3 - 78*B*tan(d*x + c)^3 +
173*A*tan(d*x + c)^2 + 115*I*B*tan(d*x + c)^2 - 32*I*A*tan(d*x + c) + 32*
B*tan(d*x + c) + 16*A)/((tan(d*x + c)^2 - I*tan(d*x + c))^2*a^2))/d
```

3.50. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$

3.50.9 Mupad [B] (verification not implemented)

Time = 7.91 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.11

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$= \frac{\tan(c+dx)^2 \left(-\frac{7B}{2a^2} + \frac{A11i}{2a^2}\right) - \tan(c+dx)^3 \left(\frac{15A}{4a^2} + \frac{B9i}{4a^2}\right) + \frac{A1i}{2a^2} + \tan(c+dx) \left(\frac{A}{a^2} + \frac{B1i}{a^2}\right)}{d \left(\tan(c+dx)^4 1i + 2 \tan(c+dx)^3 - \tan(c+dx)^2 1i\right)}$$

$$- \frac{2 \ln(\tan(c+dx)) (2A+B1i)}{a^2 d} + \frac{\ln(\tan(c+dx)+1i) (A-B1i)}{8a^2 d}$$

$$+ \frac{\ln(\tan(c+dx)-i) (31A+B17i)}{8a^2 d}$$

input `int((cot(c + d*x)^3*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^2,x)`output `(tan(c + d*x)^2*((A*11i)/(2*a^2) - (7*B)/(2*a^2)) - tan(c + d*x)^3*((15*A)/(4*a^2) + (B*9i)/(4*a^2)) + (A*1i)/(2*a^2) + tan(c + d*x)*(A/a^2 + (B*1i)/a^2))/(d*(2*tan(c + d*x)^3 - tan(c + d*x)^2*1i + tan(c + d*x)^4*1i)) - (2*log(tan(c + d*x))*(2*A + B*1i))/(a^2*d) + (log(tan(c + d*x) + 1i)*(A - B*1i))/(8*a^2*d) + (log(tan(c + d*x) - 1i)*(31*A + B*17i))/(8*a^2*d)`

3.51
$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.51.1 Optimal result 683
 3.51.2 Mathematica [A] (verified) 683
 3.51.3 Rubi [A] (verified) 684
 3.51.4 Maple [A] (verified) 687
 3.51.5 Fricas [A] (verification not implemented) 687
 3.51.6 Sympy [A] (verification not implemented) 688
 3.51.7 Maxima [F(-2)] 689
 3.51.8 Giac [A] (verification not implemented) 689
 3.51.9 Mupad [B] (verification not implemented) 690

3.51.1 Optimal result

Integrand size = 34, antiderivative size = 191

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= -\frac{(7A+25iB)x}{8a^3} - \frac{(iA-3B) \log(\cos(c+dx))}{a^3d} + \frac{(7A+25iB) \tan(c+dx)}{8a^3d}$$

$$+ \frac{(iA-B) \tan^4(c+dx)}{6d(a+ia \tan(c+dx))^3} + \frac{(5A+11iB) \tan^3(c+dx)}{24ad(a+ia \tan(c+dx))^2} - \frac{(iA-3B) \tan^2(c+dx)}{2d(a^3+ia^3 \tan(c+dx))}$$

output

```
-1/8*(7*A+25*I*B)*x/a^3-(I*A-3*B)*ln(cos(d*x+c))/a^3/d+1/8*(7*A+25*I*B)*tan(d*x+c)/a^3/d+1/6*(I*A-B)*tan(d*x+c)^4/d/(a+I*a*tan(d*x+c))^3+1/24*(5*A+11*I*B)*tan(d*x+c)^3/a/d/(a+I*a*tan(d*x+c))^2-1/2*(I*A-3*B)*tan(d*x+c)^2/d/(a^3+I*a^3*tan(d*x+c))
```

3.51.2 Mathematica [A] (verified)

Time = 1.78 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.41

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{-3((15A+49iB) \log(i-\tan(c+dx)) + (A-iB) \log(i+\tan(c+dx))) + 3(14iA-50B+3(-15iA+49iB) \tan(c+dx))}{(a+ia \tan(c+dx))^3}$$

input `Integrate[(Tan[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x
]`

output `(-3*((15*A + (49*I)*B)*Log[I - Tan[c + d*x]] + (A - I*B)*Log[I + Tan[c + d
x]]) + 3((14*I)*A - 50*B + 3*((-15*I)*A + 49*B)*Log[I - Tan[c + d*x]] +
((-3*I)*A - 3*B)*Log[I + Tan[c + d*x]])*Tan[c + d*x] + 3*(-2*(17*A + (63*I
)*B) + 3*(15*A + (49*I)*B)*Log[I - Tan[c + d*x]] + 3*(A - I*B)*Log[I + Tan
[c + d*x]])*Tan[c + d*x]^2 + ((-68*I)*A + 284*B + ((45*I)*A - 147*B)*Log[I
- Tan[c + d*x]] + 3*(I*A + B)*Log[I + Tan[c + d*x]])*Tan[c + d*x]^3 + (48
*I)*B*Tan[c + d*x]^4)/(48*a^3*d*(-I + Tan[c + d*x])^3)`

3.51.3 Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3042, 4078, 3042, 4078, 27, 3042, 4078, 27, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^4(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+ia) \tan^4(c+dx)}{6d(a+ia \tan(c+dx))^3} - \int \frac{\tan^3(c+dx)(4a(ia-B)+a(A+7iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B+ia) \tan^4(c+dx)}{6d(a+ia \tan(c+dx))^3} - \int \frac{\tan(c+dx)^3(4a(ia-B)+a(A+7iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+ia) \tan^4(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int -\frac{3 \tan^2(c+dx)(a^2(5A+11iB)-a^2(3iA-13B) \tan(c+dx))}{4a^2} dx}{6a^2} - \frac{a(5A+11iB) \tan^3(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.51. $\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{(-B + iA) \tan^4(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \int \frac{\tan^2(c+dx) (a^2(5A+11iB) - a^2(3iA-13B) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{4a^2} - \frac{a(5A+11iB) \tan^3(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{(-B + iA) \tan^4(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \int \frac{\tan(c+dx)^2 (a^2(5A+11iB) - a^2(3iA-13B) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{4a^2} - \frac{a(5A+11iB) \tan^3(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 4078 \\
 & \frac{(-B + iA) \tan^4(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{4a^2(-3B+iA) \tan^2(c+dx)}{d(a+ia \tan(c+dx))} - \frac{\int 2 \tan(c+dx) (8(iA-3B)a^3 + (7A+25iB) \tan(c+dx)a^3) dx}{2a^2} \right)}{4a^2} - \frac{a(5A+11iB) \tan^3(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & \frac{(-B + iA) \tan^4(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{4a^2(-3B+iA) \tan^2(c+dx)}{d(a+ia \tan(c+dx))} - \frac{\int \tan(c+dx) (8(iA-3B)a^3 + (7A+25iB) \tan(c+dx)a^3) dx}{a^2} \right)}{4a^2} - \frac{a(5A+11iB) \tan^3(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{(-B + iA) \tan^4(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{4a^2(-3B+iA) \tan^2(c+dx)}{d(a+ia \tan(c+dx))} - \frac{\int \tan(c+dx) (8(iA-3B)a^3 + (7A+25iB) \tan(c+dx)a^3) dx}{a^2} \right)}{4a^2} - \frac{a(5A+11iB) \tan^3(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 4008 \\
 & \frac{(-B + iA) \tan^4(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{4a^2(-3B+iA) \tan^2(c+dx)}{d(a+ia \tan(c+dx))} - \frac{8a^3(-3B+iA) \int \tan(c+dx) dx + \frac{a^3(7A+25iB) \tan(c+dx)}{d} - (a^3 x(7A+25iB))}{a^2} \right)}{4a^2} - \frac{a(5A+11iB) \tan^3(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 3042
 \end{aligned}$$

3.51. $\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\frac{\frac{(-B + iA) \tan^4(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{4a^2(-3B + iA) \tan^2(c + dx)}{d(a + ia \tan(c + dx))} - \frac{8a^3(-3B + iA) \int \tan(c + dx) dx + \frac{a^3(7A + 25iB) \tan(c + dx)}{d} - (a^3 x(7A + 25iB)) \right)}{4a^2}}{6a^2} - \frac{a(5A + 11iB) \tan^3(c + dx)}{4d(a + ia \tan(c + dx))^2}}$$

↓ 3956

$$\frac{\frac{(-B + iA) \tan^4(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{4a^2(-3B + iA) \tan^2(c + dx)}{d(a + ia \tan(c + dx))} - \frac{a^3(7A + 25iB) \tan(c + dx)}{d} - \frac{8a^3(-3B + iA) \log(\cos(c + dx))}{a^2} - (a^3 x(7A + 25iB)) \right)}{4a^2}}{6a^2} - \frac{a(5A + 11iB) \tan^3(c + dx)}{4d(a + ia \tan(c + dx))^2}}$$

input `Int[(Tan[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `((I*A - B)*Tan[c + d*x]^4)/(6*d*(a + I*a*Tan[c + d*x])^3) - (-1/4*(a*(5*A + (11*I)*B)*Tan[c + d*x]^3)/(d*(a + I*a*Tan[c + d*x])^2) + (3*((4*a^2*(I*A - 3*B)*Tan[c + d*x]^2)/(d*(a + I*a*Tan[c + d*x])) - (-a^3*(7*A + (25*I)*B)*x) - (8*a^3*(I*A - 3*B)*Log[Cos[c + d*x]])/d + (a^3*(7*A + (25*I)*B)*Tan[c + d*x])/d)/a^2))/(4*a^2))/(6*a^2)`

3.51.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

3.51. $\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

```
rule 4078 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Sim
p[(- (A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.51.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.12

method	result
risch	$-\frac{49ixB}{8a^3} - \frac{15xA}{8a^3} - \frac{23e^{-2i(dx+c)}B}{16a^3d} + \frac{11ie^{-2i(dx+c)}A}{16a^3d} + \frac{7e^{-4i(dx+c)}B}{32a^3d} - \frac{5ie^{-4i(dx+c)}A}{32a^3d} - \frac{e^{-6i(dx+c)}B}{48a^3d} + \dots$
derivativedivides	$\frac{iB \tan(dx+c)}{d a^3} - \frac{A}{6d a^3 (\tan(dx+c)-i)^3} - \frac{iB}{6d a^3 (\tan(dx+c)-i)^3} - \frac{3B \ln(1+\tan^2(dx+c))}{2d a^3} - \frac{25iB \arctan(\tan(dx+c))}{8d a^3}$
default	$\frac{iB \tan(dx+c)}{d a^3} - \frac{A}{6d a^3 (\tan(dx+c)-i)^3} - \frac{iB}{6d a^3 (\tan(dx+c)-i)^3} - \frac{3B \ln(1+\tan^2(dx+c))}{2d a^3} - \frac{25iB \arctan(\tan(dx+c))}{8d a^3}$

```
input int(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVER
BOSE)
```

```
output -49/8*I*x/a^3*B-15/8*x/a^3*A-23/16/a^3/d*exp(-2*I*(d*x+c))*B+11/16*I/a^3/d
*exp(-2*I*(d*x+c))*A+7/32/a^3/d*exp(-4*I*(d*x+c))*B-5/32*I/a^3/d*exp(-4*I*
(d*x+c))*A-1/48/a^3/d*exp(-6*I*(d*x+c))*B+1/48*I/a^3/d*exp(-6*I*(d*x+c))*A
-6*I/a^3/d*B*c-2/a^3/d*A*c-2/d/a^3*B/(exp(2*I*(d*x+c))+1)+3/a^3/d*ln(exp(2
*I*(d*x+c))+1)*B-I/a^3/d*ln(exp(2*I*(d*x+c))+1)*A
```

3.51.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.91

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \frac{12(15A+49iB)dx e^{(8i dx+8i c)} + 6(2(15A+49iB)dx - 11iA + 55B)e^{(6i dx+6i c)} + 3(-17iA + 39B)e^{(4i dx+4i c)}}{96(a^3)}$$

3.51. $\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

```
input integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm=
"fricas")
```

```
output -1/96*(12*(15*A + 49*I*B)*d*x*e^(8*I*d*x + 8*I*c) + 6*(2*(15*A + 49*I*B)*d
*x - 11*I*A + 55*B)*e^(6*I*d*x + 6*I*c) + 3*(-17*I*A + 39*B)*e^(4*I*d*x +
4*I*c) - (-13*I*A + 19*B)*e^(2*I*d*x + 2*I*c) + 96*((I*A - 3*B)*e^(8*I*d*x
+ 8*I*c) + (I*A - 3*B)*e^(6*I*d*x + 6*I*c))*log(e^(2*I*d*x + 2*I*c) + 1)
- 2*I*A + 2*B)/(a^3*d*e^(8*I*d*x + 8*I*c) + a^3*d*e^(6*I*d*x + 6*I*c))
```

3.51.6 Sympy [A] (verification not implemented)

Time = 0.60 (sec) , antiderivative size = 337, normalized size of antiderivative = 1.76

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = -\frac{2B}{a^3 d e^{2ic} e^{2idx} + a^3 d} + \left\{ \frac{((512iAa^6 d^2 e^{6ic} - 512Ba^6 d^2 e^{6ic})e^{-6idx} + (-3840iAa^6 d^2 e^{8ic} + 5376Ba^6 d^2 e^{8ic})e^{-4idx} + (16896iAa^6 d^2 e^{10ic} - 35328Ba^6 d^2 e^{10ic})e^{-2idx})e^{-2ic}}{24576a^9 d^3} \right. \\ \left. + x \left(-\frac{15A-49iB}{8a^3} + \frac{(-15Ae^{6ic} + 11Ae^{4ic} - 5Ae^{2ic} + A - 49iBe^{6ic} + 23iBe^{4ic} - 7iBe^{2ic} + iB)e^{-6ic}}{8a^3} \right) \right. \\ \left. + \frac{x(-15A - 49iB)}{8a^3} - \frac{i(A + 3iB) \log(e^{2idx} + e^{-2ic})}{a^3 d} \right.$$

```
input integrate(tan(d*x+c)**4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)
```

```
output -2*B/(a**3*d*exp(2*I*c)*exp(2*I*d*x) + a**3*d) + Piecewise((((512*I*A*a**6
*d**2*exp(6*I*c) - 512*B*a**6*d**2*exp(6*I*c))*exp(-6*I*d*x) + (-3840*I*A
*a**6*d**2*exp(8*I*c) + 5376*B*a**6*d**2*exp(8*I*c))*exp(-4*I*d*x) + (16896
*I*A*a**6*d**2*exp(10*I*c) - 35328*B*a**6*d**2*exp(10*I*c))*exp(-2*I*d*x))
*exp(-12*I*c)/(24576*a**9*d**3), Ne(a**9*d**3*exp(12*I*c), 0)), (x*(-(15*
A - 49*I*B)/(8*a**3) + (-15*A*exp(6*I*c) + 11*A*exp(4*I*c) - 5*A*exp(2*I*c)
) + A - 49*I*B*exp(6*I*c) + 23*I*B*exp(4*I*c) - 7*I*B*exp(2*I*c) + I*B)*ex
p(-6*I*c)/(8*a**3)), True)) + x*(-15*A - 49*I*B)/(8*a**3) - I*(A + 3*I*B)*
log(exp(2*I*d*x) + exp(-2*I*c))/(a**3*d)
```

3.51.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm=
"maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.51.8 Giac [A] (verification not implemented)

Time = 1.08 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.74

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$= \frac{\frac{6(iA+B)\log(\tan(dx+c)+i)}{a^3} - \frac{6(-15iA+49B)\log(\tan(dx+c)-i)}{a^3} + \frac{96iB\tan(dx+c)}{a^3} - \frac{165iA\tan(dx+c)^3 - 539B\tan(dx+c)^3 + 291A\tan(dx+c)^2 - 171iA\tan(dx+c) + 981B\tan(dx+c) - 29A - 259iB}{a^3(\tan(dx+c) - I)^3}}{96d}$$

```
input integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm=
"giac")
```

```
output 1/96*(6*(I*A + B)*log(tan(d*x + c) + I)/a^3 - 6*(-15*I*A + 49*B)*log(tan(d
*x + c) - I)/a^3 + 96*I*B*tan(d*x + c)/a^3 - (165*I*A*tan(d*x + c)^3 - 539
*B*tan(d*x + c)^3 + 291*A*tan(d*x + c)^2 + 1245*I*B*tan(d*x + c)^2 - 171*I
*A*tan(d*x + c) + 981*B*tan(d*x + c) - 29*A - 259*I*B)/(a^3*(tan(d*x + c)
- I)^3))/d
```

3.51. $\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

3.51.9 Mupad [B] (verification not implemented)

Time = 8.03 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.96

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$= \frac{\tan(c+dx) \left(\frac{B7i}{2a^3} + \frac{(-3B+A1i)27i}{8a^3} \right) + \frac{4B}{3a^3} - \tan(c+dx)^2 \left(\frac{5B}{2a^3} + \frac{17(-3B+A1i)}{8a^3} \right) + \frac{17(-3B+A1i)}{12a^3}}{d(-\tan(c+dx)^3 1i - 3\tan(c+dx)^2 + \tan(c+dx) 3i + 1)}$$

$$+ \frac{\ln(\tan(c+dx) + 1i)(B+A1i)}{16a^3d} + \frac{B\tan(c+dx) 1i}{a^3d}$$

$$+ \frac{\ln(\tan(c+dx) - i)(-49B+A15i)}{16a^3d}$$

input `int((tan(c + d*x)^4*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3,x)`output `(tan(c + d*x)*((B*7i)/(2*a^3) + ((A*1i - 3*B)*27i)/(8*a^3)) + (4*B)/(3*a^3) - tan(c + d*x)^2*((5*B)/(2*a^3) + (17*(A*1i - 3*B))/(8*a^3)) + (17*(A*1i - 3*B))/(12*a^3))/(d*(tan(c + d*x)*3i - 3*tan(c + d*x)^2 - tan(c + d*x)^3*1i + 1)) + (log(tan(c + d*x) + 1i)*(A*1i + B))/(16*a^3*d) + (B*tan(c + d*x)*1i)/(a^3*d) + (log(tan(c + d*x) - 1i)*(A*15i - 49*B))/(16*a^3*d)`

3.52
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.52.1 Optimal result 691
 3.52.2 Mathematica [A] (verified) 691
 3.52.3 Rubi [A] (verified) 692
 3.52.4 Maple [A] (verified) 695
 3.52.5 Fricas [A] (verification not implemented) 695
 3.52.6 Sympy [A] (verification not implemented) 696
 3.52.7 Maxima [F(-2)] 696
 3.52.8 Giac [A] (verification not implemented) 697
 3.52.9 Mupad [B] (verification not implemented) 697

3.52.1 Optimal result

Integrand size = 34, antiderivative size = 148

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \frac{(iA-7B)x}{8a^3} - \frac{iB \log(\cos(c+dx))}{a^3d} + \frac{(iA-B) \tan^3(c+dx)}{6d(a+ia \tan(c+dx))^3} + \frac{(A+3iB) \tan^2(c+dx)}{8ad(a+ia \tan(c+dx))^2} + \frac{A+7iB}{8d(a^3+ia^3 \tan(c+dx))}$$

```
output 1/8*(I*A-7*B)*x/a^3-I*B*ln(cos(d*x+c))/a^3/d+1/6*(I*A-B)*tan(d*x+c)^3/d/(a
+I*a*tan(d*x+c))^3+1/8*(A+3*I*B)*tan(d*x+c)^2/a/d/(a+I*a*tan(d*x+c))^2+1/8
*(A+7*I*B)/d/(a^3+I*a^3*tan(d*x+c))
```

3.52.2 Mathematica [A] (verified)

Time = 2.13 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.12

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \frac{i(3i((A+15iB) \log(i-\tan(c+dx)) - (A-iB) \log(i+\tan(c+dx))) + 6(A+7iB) \tan(c+dx) + 2}{48}$$

input `Integrate[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x
]`

output `((-1/48*I)*((3*I)*((A + (15*I)*B)*Log[I - Tan[c + d*x]] - (A - I*B)*Log[I + Tan[c + d*x]]) + 6*(A + (7*I)*B)*Tan[c + d*x] + 24*B*Tan[c + d*x]^2 - 2*(A + (7*I)*B)*Tan[c + d*x]^3 + (2*Tan[c + d*x]^4*(-6*(A + I*B) + ((-3*I)*A + 9*B)*Tan[c + d*x] + (A + (7*I)*B)*Tan[c + d*x]^2))/(-I + Tan[c + d*x])^3)/(a^3*d)`

3.52.3 Rubi [A] (verified)

Time = 0.86 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.07, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.382$, Rules used = {3042, 4078, 27, 3042, 4078, 27, 3042, 4072, 27, 3042, 3956, 4009, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^3(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+IA) \tan^3(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{3 \tan^2(c+dx)(a(iA-B)+2iaB \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{6a^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B+IA) \tan^3(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{\tan^2(c+dx)(a(iA-B)+2iaB \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{2a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B+IA) \tan^3(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{\tan(c+dx)^2(a(iA-B)+2iaB \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{2a^2} \\
 & \quad \downarrow \text{4078}
 \end{aligned}$$

3.52. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{(-B + iA) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int -\frac{2 \tan(c+dx)((A+3iB)a^2+4B \tan(c+dx)a^2)}{i \tan(c+dx)a+a} dx}{4a^2} - \frac{a(A+3iB) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & \frac{(-B + iA) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{\tan(c+dx)((A+3iB)a^2+4B \tan(c+dx)a^2)}{i \tan(c+dx)a+a} dx}{2a^2} - \frac{a(A+3iB) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{(-B + iA) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{\tan(c+dx)((A+3iB)a^2+4B \tan(c+dx)a^2)}{i \tan(c+dx)a+a} dx}{2a^2} - \frac{a(A+3iB) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 4072 \\
 & \frac{(-B + iA) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{-\int \frac{a^3(iA-7B) \tan(c+dx)}{i \tan(c+dx)a+a} dx}{a} - \frac{4iaB \int \tan(c+dx) dx}{2a^2} - \frac{a(A+3iB) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & \frac{(-B + iA) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{-ia^2(-7B+iA) \int \frac{\tan(c+dx)}{i \tan(c+dx)a+a} dx - 4iaB \int \tan(c+dx) dx}{2a^2} - \frac{a(A+3iB) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{(-B + iA) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{-ia^2(-7B+iA) \int \frac{\tan(c+dx)}{i \tan(c+dx)a+a} dx - 4iaB \int \tan(c+dx) dx}{2a^2} - \frac{a(A+3iB) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 3956 \\
 & \frac{(-B + iA) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\frac{4iaB \log(\cos(c+dx))}{d} - ia^2(-7B+iA) \int \frac{\tan(c+dx)}{i \tan(c+dx)a+a} dx}{2a^2} - \frac{a(A+3iB) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 4009 \\
 & \frac{(-B + iA) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\frac{4iaB \log(\cos(c+dx))}{d} - ia^2(-7B+iA) \left(-\frac{i \int 1 dx}{2a} - \frac{1}{2d(a+ia \tan(c+dx))} \right)}{2a^2} - \frac{a(A+3iB) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 24 \\
 & \frac{(-B + iA) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\frac{4iaB \log(\cos(c+dx))}{d} - ia^2(-7B+iA) \left(-\frac{1}{2d(a+ia \tan(c+dx))} - \frac{ix}{2a} \right)}{2a^2} - \frac{a(A+3iB) \tan^2(c+dx)}{4d(a+ia \tan(c+dx))^2}
 \end{aligned}$$

3.52. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

input `Int[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `((I*A - B)*Tan[c + d*x]^3)/(6*d*(a + I*a*Tan[c + d*x])^3) - (-1/4*(a*(A + (3*I)*B)*Tan[c + d*x]^2)/(d*(a + I*a*Tan[c + d*x])^2) + (((4*I)*a*B*Log[Cos[c + d*x]])/d - I*a^2*(I*A - 7*B)*((-1/2*I)*x)/a - 1/(2*d*(a + I*a*Tan[c + d*x]))) / (2*a^2) / (2*a^2)`

3.52.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

rule 4072 `Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]) / ((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*(d/b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c - a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0]`

```
rule 4078 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(- (A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.52.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.08

method	result
risch	$-\frac{15xB}{8a^3} + \frac{ixA}{8a^3} + \frac{11ie^{-2i(dx+c)}B}{16a^3d} + \frac{3e^{-2i(dx+c)}A}{16a^3d} - \frac{5ie^{-4i(dx+c)}B}{32a^3d} - \frac{3e^{-4i(dx+c)}A}{32a^3d} + \frac{ie^{-6i(dx+c)}B}{48a^3d} + e$
derivativedivides	$\frac{iA \arctan(\tan(dx+c))}{8da^3} + \frac{iB \ln(1+\tan^2(dx+c))}{2da^3} - \frac{7B \arctan(\tan(dx+c))}{8da^3} + \frac{17B}{8da^3(\tan(dx+c)-i)} - \frac{7iA}{8da^3(\tan(dx+c)-i)}$
default	$\frac{iA \arctan(\tan(dx+c))}{8da^3} + \frac{iB \ln(1+\tan^2(dx+c))}{2da^3} - \frac{7B \arctan(\tan(dx+c))}{8da^3} + \frac{17B}{8da^3(\tan(dx+c)-i)} - \frac{7iA}{8da^3(\tan(dx+c)-i)}$

```
input int(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVER
BOSE)
```

```
output -15/8*x/a^3*B+1/8*I*x/a^3*A+11/16*I/a^3/d*exp(-2*I*(d*x+c))*B+3/16/a^3/d*e
xp(-2*I*(d*x+c))*A-5/32*I/a^3/d*exp(-4*I*(d*x+c))*B-3/32/a^3/d*exp(-4*I*(d
*x+c))*A+1/48*I/a^3/d*exp(-6*I*(d*x+c))*B+1/48/a^3/d*exp(-6*I*(d*x+c))*A-2
*B/a^3/d*c-I*B/a^3/d*ln(exp(2*I*(d*x+c))+1)
```

3.52.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.70

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \frac{(12(-iA+15B)dx e^{(6i dx+6i c)} + 96i B e^{(6i dx+6i c)} \log(e^{(2i dx+2i c)} + 1) - 6(3A+11iB)e^{(4i dx+4i c)} + 3(3A+11iB)e^{(2i dx+2i c)} + 3(3A+11iB))}{96 a^3 d}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm=
"fracas")
```

3.52. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

```
output -1/96*(12*(-I*A + 15*B)*d*x*e^(6*I*d*x + 6*I*c) + 96*I*B*e^(6*I*d*x + 6*I*c)*log(e^(2*I*d*x + 2*I*c) + 1) - 6*(3*A + 11*I*B)*e^(4*I*d*x + 4*I*c) + 3*(3*A + 5*I*B)*e^(2*I*d*x + 2*I*c) - 2*A - 2*I*B)*e^(-6*I*d*x - 6*I*c)/(a^3*d)
```

3.52.6 Sympy [A] (verification not implemented)

Time = 0.56 (sec) , antiderivative size = 296, normalized size of antiderivative = 2.00

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = -\frac{iB \log(e^{2idx} + e^{-2ic})}{a^3 d} + \left\{ \frac{((512Aa^6 d^2 e^{6ic} + 512iBa^6 d^2 e^{6ic})e^{-6idx} + (-2304Aa^6 d^2 e^{8ic} - 3840iBa^6 d^2 e^{8ic})e^{-4idx} + (4608Aa^6 d^2 e^{10ic} + 16896iBa^6 d^2 e^{10ic})e^{-2idx})e^{-12ic}}{24576a^9 d^3} \right. \\ \left. + x \left(-\frac{iA-15B}{8a^3} + \frac{(iAe^{6ic} - 3iAe^{4ic} + 3iAe^{2ic} - iA - 15Be^{6ic} + 11Be^{4ic} - 5Be^{2ic} + B)e^{-6ic}}{8a^3} \right) \right. \\ \left. + \frac{x(iA - 15B)}{8a^3} \right.$$

```
input integrate(tan(d*x+c)**3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)
```

```
output -I*B*log(exp(2*I*d*x) + exp(-2*I*c))/(a**3*d) + Piecewise((((512*A*a**6*d**2*exp(6*I*c) + 512*I*B*a**6*d**2*exp(6*I*c))*exp(-6*I*d*x) + (-2304*A*a**6*d**2*exp(8*I*c) - 3840*I*B*a**6*d**2*exp(8*I*c))*exp(-4*I*d*x) + (4608*A*a**6*d**2*exp(10*I*c) + 16896*I*B*a**6*d**2*exp(10*I*c))*exp(-2*I*d*x))*exp(-12*I*c)/(24576*a**9*d**3), Ne(a**9*d**3*exp(12*I*c), 0)), (x*(-(I*A - 15*B)/(8*a**3) + (I*A*exp(6*I*c) - 3*I*A*exp(4*I*c) + 3*I*A*exp(2*I*c) - I*A - 15*B*exp(6*I*c) + 11*B*exp(4*I*c) - 5*B*exp(2*I*c) + B)*exp(-6*I*c)/(8*a**3)), True)) + x*(I*A - 15*B)/(8*a**3)
```

3.52.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")
```

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

3.52.8 Giac [A] (verification not implemented)

Time = 0.82 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.86

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \frac{\frac{6(A-iB)\log(\tan(dx+c)+i)}{a^3} - \frac{6(A+15iB)\log(\tan(dx+c)-i)}{a^3} + \frac{11A\tan(dx+c)^3+165iB\tan(dx+c)^3+51iA\tan(dx+c)^2+291B\tan(dx+c)}{a^3(\tan(dx+c)-i)^5}}{96d}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output
$$\frac{-1/96*(6*(A - I*B)*\log(\tan(d*x + c) + I)/a^3 - 6*(A + 15*I*B)*\log(\tan(d*x + c) - I)/a^3 + (11*A*\tan(d*x + c)^3 + 165*I*B*\tan(d*x + c)^3 + 51*I*A*\tan(d*x + c)^2 + 291*B*\tan(d*x + c)^2 + 75*A*\tan(d*x + c) - 171*I*B*\tan(d*x + c) - 29*I*A - 29*B)/(a^3*(\tan(d*x + c) - I)^3))/d}$$

3.52.9 Mupad [B] (verification not implemented)

Time = 7.69 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.99

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \frac{\frac{5A}{12a^3} - \tan(c+dx)^2 \left(\frac{7A}{8a^3} + \frac{B17i}{8a^3} \right) + \frac{B17i}{12a^3} + \tan(c+dx) \left(-\frac{27B}{8a^3} + \frac{A9i}{8a^3} \right)}{d \left(-\tan(c+dx)^3 \operatorname{li} - 3 \tan(c+dx)^2 + \tan(c+dx) 3i + 1 \right)} - \frac{\ln(\tan(c+dx) + 1i) (A - B 1i)}{16 a^3 d} + \frac{\ln(\tan(c+dx) - i) (A + B 15i)}{16 a^3 d}$$

input `int((tan(c + d*x)^3*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3,x)`

output
$$\left(\frac{5A}{12a^3} - \tan(c+d*x)^2 \left(\frac{7A}{8a^3} + \frac{B*17i}{8a^3} \right) + \frac{B*17i}{12a^3} + \tan(c+d*x) \left(-\frac{27B}{8a^3} + \frac{A*9i}{8a^3} \right) \right) / (d * (\tan(c+d*x)*3i - 3*\tan(c+d*x)^2 - \tan(c+d*x)^3*1i + 1)) - (\log(\tan(c+d*x) + 1i)*(A - B*1i))/(16*a^3*d) + (\log(\tan(c+d*x) - 1i)*(A + B*15i))/(16*a^3*d)$$

3.53 $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

3.53.1 Optimal result 698
 3.53.2 Mathematica [A] (verified) 698
 3.53.3 Rubi [A] (verified) 699
 3.53.4 Maple [A] (verified) 701
 3.53.5 Fricas [A] (verification not implemented) 701
 3.53.6 Sympy [A] (verification not implemented) 702
 3.53.7 Maxima [F(-2)] 702
 3.53.8 Giac [A] (verification not implemented) 703
 3.53.9 Mupad [B] (verification not implemented) 703

3.53.1 Optimal result

Integrand size = 34, antiderivative size = 124

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = -\frac{(A-iB)x}{8a^3} + \frac{(iA-B) \tan^2(c+dx)}{6d(a+ia \tan(c+dx))^3} + \frac{iA-7B}{24ad(a+ia \tan(c+dx))^2} + \frac{iA+17B}{24d(a^3+ia^3 \tan(c+dx))}$$

output `-1/8*(A-I*B)*x/a^3+1/6*(I*A-B)*tan(d*x+c)^2/d/(a+I*a*tan(d*x+c))^3+1/24*(I*A-7*B)/a/d/(a+I*a*tan(d*x+c))^2+1/24*(I*A+17*B)/d/(a^3+I*a^3*tan(d*x+c))`

3.53.2 Mathematica [A] (verified)

Time = 1.13 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.19

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \frac{\sec^3(c+dx)(-9(A-iB) \cos(c+dx) + 2(A+iB-6iAdx-6Bdx) \cos(3(c+dx)) - 3iA \sin(c+dx) - \dots)}{96a^3d(-i + \dots)}$$

input `Integrate[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output $(\text{Sec}[c + d*x]^3*(-9*(A - I*B)*\text{Cos}[c + d*x] + 2*(A + I*B - (6*I)*A*d*x - 6*B*d*x)*\text{Cos}[3*(c + d*x)] - (3*I)*A*\text{Sin}[c + d*x] - 27*B*\text{Sin}[c + d*x] - (2*I)*A*\text{Sin}[3*(c + d*x)] + 2*B*\text{Sin}[3*(c + d*x)] + 12*A*d*x*\text{Sin}[3*(c + d*x)] - (12*I)*B*d*x*\text{Sin}[3*(c + d*x)]))/(96*a^3*d*(-I + \text{Tan}[c + d*x])^3)$

3.53.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.09, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.206$, Rules used = {3042, 4078, 3042, 4073, 3042, 4009, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

↓ 3042

$$\int \frac{\tan(c+dx)^2(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

↓ 4078

$$\frac{(-B+iA) \tan^2(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{\tan(c+dx)(2a(iA-B)-a(A-5iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{6a^2}$$

↓ 3042

$$\frac{(-B+iA) \tan^2(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{\tan(c+dx)(2a(iA-B)-a(A-5iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{6a^2}$$

↓ 4073

$$\frac{(-B+iA) \tan^2(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{i \int \frac{a^2(iA-7B)-2a^2(A-5iB) \tan(c+dx)}{i \tan(c+dx)a+a} dx}{2a^2} - \frac{a(-7B+iA)}{4d(a+ia \tan(c+dx))^2}$$

↓ 3042

$$\frac{(-B+iA) \tan^2(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{i \int \frac{a^2(iA-7B)-2a^2(A-5iB) \tan(c+dx)}{i \tan(c+dx)a+a} dx}{2a^2} - \frac{a(-7B+iA)}{4d(a+ia \tan(c+dx))^2}$$

↓ 4009

3.53. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\frac{(-B + iA) \tan^2(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{i \left(\frac{3}{2} a(B + iA) \int 1 dx + \frac{a^2(A - 17iB)}{2d(a + ia \tan(c + dx))} \right)}{2a^2} - \frac{a(-7B + iA)}{4d(a + ia \tan(c + dx))^2}$$

↓ 24

$$\frac{(-B + iA) \tan^2(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{i \left(\frac{a^2(A - 17iB)}{2d(a + ia \tan(c + dx))} + \frac{3}{2} ax(B + iA) \right)}{2a^2} - \frac{a(-7B + iA)}{4d(a + ia \tan(c + dx))^2}$$

input `Int[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `((I*A - B)*Tan[c + d*x]^2)/(6*d*(a + I*a*Tan[c + d*x])^3) - (-1/4*(a*(I*A - 7*B))/(d*(a + I*a*Tan[c + d*x])^2) - ((I/2)*((3*a*(I*A + B)*x)/2 + (a^2*(A - (17*I)*B))/(2*d*(a + I*a*Tan[c + d*x]))))/a^2)/(6*a^2)`

3.53.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4009 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

rule 4073 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)], x_Symbol] := Simp[(-(A*b - a*B))*(a*c + b*d)*((a + b*Tan[e + f*x])^m/(2*a^2*f*m)), x] + Simp[1/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[A*b*c + a*B*c + a*A*d + b*B*d + 2*a*B*d*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && EqQ[a^2 + b^2, 0]`

```
rule 4078 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(- (A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.53.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.03

method	result
risch	$\frac{ixB}{8a^3} - \frac{xA}{8a^3} + \frac{3e^{-2i(dx+c)}B}{16a^3d} + \frac{ie^{-2i(dx+c)}A}{16a^3d} - \frac{3e^{-4i(dx+c)}B}{32a^3d} + \frac{ie^{-4i(dx+c)}A}{32a^3d} + \frac{e^{-6i(dx+c)}B}{48a^3d} - \frac{ie^{-6i(dx+c)}A}{48a^3d}$
derivativedivides	$\frac{A}{6da^3(\tan(dx+c)-i)^3} + \frac{iB}{6da^3(\tan(dx+c)-i)^3} - \frac{A}{8da^3(\tan(dx+c)-i)} - \frac{7iB}{8da^3(\tan(dx+c)-i)} - \frac{A \arctan(\tan(dx+c))}{8da^3}$
default	$\frac{A}{6da^3(\tan(dx+c)-i)^3} + \frac{iB}{6da^3(\tan(dx+c)-i)^3} - \frac{A}{8da^3(\tan(dx+c)-i)} - \frac{7iB}{8da^3(\tan(dx+c)-i)} - \frac{A \arctan(\tan(dx+c))}{8da^3}$

```
input int(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVER
BOSE)
```

```
output 1/8*I*x/a^3*B-1/8*x/a^3*A+3/16/a^3/d*exp(-2*I*(d*x+c))*B+1/16*I/a^3/d*exp(
-2*I*(d*x+c))*A-3/32/a^3/d*exp(-4*I*(d*x+c))*B+1/32*I/a^3/d*exp(-4*I*(d*x+
c))*A+1/48/a^3/d*exp(-6*I*(d*x+c))*B-1/48*I/a^3/d*exp(-6*I*(d*x+c))*A
```

3.53.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.63

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \frac{(12(A-iB)dx e^{(6i dx+6i c)} + 6(-iA-3B)e^{(4i dx+4i c)} + 3(-iA+3B)e^{(2i dx+2i c)} + 2iA-2B)e^{(-6i dx)}}{96a^3d}$$

```
input integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm=
"fracas")
```

3.53. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

output
$$\frac{-1/96*(12*(A - I*B)*d*x*e^{(6*I*d*x + 6*I*c)} + 6*(-I*A - 3*B)*e^{(4*I*d*x + 4*I*c)} + 3*(-I*A + 3*B)*e^{(2*I*d*x + 2*I*c)} + 2*I*A - 2*B)*e^{(-6*I*d*x - 6*I*c)}}{(a^3*d)}$$

3.53.6 Sympy [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 258, normalized size of antiderivative = 2.08

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$= \left\{ \begin{array}{l} \frac{((-512iAa^6d^2e^{6ic}+512Ba^6d^2e^{6ic})e^{-6idx}+(768iAa^6d^2e^{8ic}-2304Ba^6d^2e^{8ic})e^{-4idx}+(1536iAa^6d^2e^{10ic}+4608Ba^6d^2e^{10ic})e^{-2idx})e^{-12ic}}{24576a^9d^3} \\ x\left(-\frac{-A+iB}{8a^3} + \frac{(-Ae^{6ic}+Ae^{4ic}+Ae^{2ic}-A+iBe^{6ic}-3iBe^{4ic}+3iBe^{2ic}-iB)e^{-6ic}}{8a^3}\right) \\ + \frac{x(-A+iB)}{8a^3} \end{array} \right.$$

input `integrate(tan(d*x+c)**2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)`

output `Piecewise(((((-512*I*A*a**6*d**2*exp(6*I*c) + 512*B*a**6*d**2*exp(6*I*c))*exp(-6*I*d*x) + (768*I*A*a**6*d**2*exp(8*I*c) - 2304*B*a**6*d**2*exp(8*I*c))*exp(-4*I*d*x) + (1536*I*A*a**6*d**2*exp(10*I*c) + 4608*B*a**6*d**2*exp(10*I*c))*exp(-2*I*d*x))*exp(-12*I*c)/(24576*a**9*d**3), Ne(a**9*d**3*exp(12*I*c), 0)), (x*(-(-A + I*B)/(8*a**3) + (-A*exp(6*I*c) + A*exp(4*I*c) + A*exp(2*I*c) - A + I*B*exp(6*I*c) - 3*I*B*exp(4*I*c) + 3*I*B*exp(2*I*c) - I*B)*exp(-6*I*c)/(8*a**3)), True)) + x*(-A + I*B)/(8*a**3)`

3.53.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.53.8 Giac [A] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.04

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx =$$

$$\frac{6(iA+B)\log(\tan(dx+c)+i)}{a^3} + \frac{6(-iA-B)\log(\tan(dx+c)-i)}{a^3} + \frac{11iA\tan(dx+c)^3+11B\tan(dx+c)^3+45A\tan(dx+c)^2+51iB\tan(dx+c)}{a^3(\tan(dx+c)-i)^3}$$

$$96d$$

```
input integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm=
"giac")
```

```
output -1/96*(6*(I*A + B)*log(tan(d*x + c) + I)/a^3 + 6*(-I*A - B)*log(tan(d*x +
c) - I)/a^3 + (11*I*A*tan(d*x + c)^3 + 11*B*tan(d*x + c)^3 + 45*A*tan(d*x
+ c)^2 + 51*I*B*tan(d*x + c)^2 - 21*I*A*tan(d*x + c) + 75*B*tan(d*x + c) -
3*A - 29*I*B)/(a^3*(tan(d*x + c) - I)^3))/d
```

3.53.9 Mupad [B] (verification not implemented)

Time = 7.72 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.90

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$= \frac{\tan(c+dx)^2 \left(-\frac{7B}{8a^3} + \frac{A1i}{8a^3} \right) + \frac{A1i}{12a^3} + \frac{5B}{12a^3} - \tan(c+dx) \left(\frac{A}{8a^3} - \frac{B9i}{8a^3} \right) + \frac{x(B+A1i)1i}{8a^3}}{d \left(-\tan(c+dx)^3 1i - 3\tan(c+dx)^2 + \tan(c+dx) 3i + 1 \right)}$$

```
input int((tan(c + d*x)^2*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3,x)
```

```
output (tan(c + d*x)^2*((A*1i)/(8*a^3) - (7*B)/(8*a^3)) + (A*1i)/(12*a^3) + (5*B)
/(12*a^3) - tan(c + d*x)*(A/(8*a^3) - (B*9i)/(8*a^3)))/(d*(tan(c + d*x)*3i
- 3*tan(c + d*x)^2 - tan(c + d*x)^3*1i + 1)) + (x*(A*1i + B)*1i)/(8*a^3)
```

3.54 $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

3.54.1	Optimal result	704
3.54.2	Mathematica [A] (verified)	704
3.54.3	Rubi [A] (verified)	705
3.54.4	Maple [A] (verified)	707
3.54.5	Fricas [A] (verification not implemented)	707
3.54.6	Sympy [A] (verification not implemented)	708
3.54.7	Maxima [F(-2)]	708
3.54.8	Giac [A] (verification not implemented)	709
3.54.9	Mupad [B] (verification not implemented)	709

3.54.1 Optimal result

Integrand size = 32, antiderivative size = 110

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = -\frac{(iA+B)x}{8a^3} - \frac{A+iB}{6d(a+ia \tan(c+dx))^3} + \frac{A+3iB}{8ad(a+ia \tan(c+dx))^2} + \frac{A-iB}{8d(a^3+ia^3 \tan(c+dx))}$$

output `-1/8*(I*A+B)*x/a^3+1/6*(-A-I*B)/d/(a+I*a*tan(d*x+c))^3+1/8*(A+3*I*B)/a/d/(a+I*a*tan(d*x+c))^2+1/8*(A-I*B)/d/(a^3+I*a^3*tan(d*x+c))`

3.54.2 Mathematica [A] (verified)

Time = 1.53 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.35

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \frac{(\cos(3(c+dx)) - i \sin(3(c+dx)))(3(A+3iB) \cos(c+dx) - 2(A+6iAdx + B(i+6dx)) \cos(3(c+dx)))}{(a+ia \tan(c+dx))^3}$$

input `Integrate[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output $((\text{Cos}[3*(c + d*x)] - I*\text{Sin}[3*(c + d*x)])*(3*(A + (3*I)*B)*\text{Cos}[c + d*x] - 2*(A + (6*I)*A*d*x + B*(I + 6*d*x))*\text{Cos}[3*(c + d*x)] + (9*I)*A*\text{Sin}[c + d*x] - 3*B*\text{Sin}[c + d*x] + (2*I)*A*\text{Sin}[3*(c + d*x)] - 2*B*\text{Sin}[3*(c + d*x)] + 12*A*d*x*\text{Sin}[3*(c + d*x)] - (12*I)*B*d*x*\text{Sin}[3*(c + d*x)]))/(96*a^3*d)$

3.54.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.05, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.219$, Rules used = {3042, 4073, 3042, 4009, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx \\ & \quad \downarrow \text{4073} \\ & -\frac{i \int \frac{a(A+iB)+2aB\tan(c+dx)}{(i\tan(c+dx)a+a)^2} dx}{2a^2} - \frac{A+iB}{6d(a+ia\tan(c+dx))^3} \\ & \quad \downarrow \text{3042} \\ & -\frac{i \int \frac{a(A+iB)+2aB\tan(c+dx)}{(i\tan(c+dx)a+a)^2} dx}{2a^2} - \frac{A+iB}{6d(a+ia\tan(c+dx))^3} \\ & \quad \downarrow \text{4009} \\ & -\frac{i\left(\frac{1}{2}(A-iB) \int \frac{1}{i\tan(c+dx)a+a} dx + \frac{a(-3B+iA)}{4d(a+ia\tan(c+dx))^2}\right)}{2a^2} - \frac{A+iB}{6d(a+ia\tan(c+dx))^3} \\ & \quad \downarrow \text{3042} \\ & -\frac{i\left(\frac{1}{2}(A-iB) \int \frac{1}{i\tan(c+dx)a+a} dx + \frac{a(-3B+iA)}{4d(a+ia\tan(c+dx))^2}\right)}{2a^2} - \frac{A+iB}{6d(a+ia\tan(c+dx))^3} \\ & \quad \downarrow \text{3960} \\ & -\frac{i\left(\frac{1}{2}(A-iB)\left(\frac{\int 1dx}{2a} + \frac{i}{2d(a+ia\tan(c+dx))}\right) + \frac{a(-3B+iA)}{4d(a+ia\tan(c+dx))^2}\right)}{2a^2} - \frac{A+iB}{6d(a+ia\tan(c+dx))^3} \end{aligned}$$

3.54. $\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

$$\downarrow 24$$

$$-\frac{i\left(\frac{a(-3B+iA)}{4d(a+ia\tan(c+dx))^2} + \frac{1}{2}(A-iB)\left(\frac{x}{2a} + \frac{i}{2d(a+ia\tan(c+dx))}\right)\right)}{2a^2} - \frac{A+iB}{6d(a+ia\tan(c+dx))^3}$$

input `Int[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `-1/6*(A + I*B)/(d*(a + I*a*Tan[c + d*x])^3) - ((I/2)*((a*(I*A - 3*B))/(4*d*(a + I*a*Tan[c + d*x])^2) + ((A - I*B)*(x/(2*a) + (I/2)/(d*(a + I*a*Tan[c + d*x]))))/2))/a^2`

3.54.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

rule 4073 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(A*b - a*B))*(a*c + b*d)*((a + b*Tan[e + f*x])^m/(2*a^2*f*m)), x] + Simp[1/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[A*b*c + a*B*c + a*A*d + b*B*d + 2*a*B*d*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && EqQ[a^2 + b^2, 0]`

3.54.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.16

method	result
risch	$-\frac{x B}{8 a^3} - \frac{i x A}{8 a^3} + \frac{i e^{-2 i(dx+c)} B}{16 a^3 d} + \frac{e^{-2 i(dx+c)} A}{16 a^3 d} + \frac{i e^{-4 i(dx+c)} B}{32 a^3 d} - \frac{e^{-4 i(dx+c)} A}{32 a^3 d} - \frac{i e^{-6 i(dx+c)} B}{48 a^3 d} - \frac{e^{-6 i(dx+c)} A}{48 a^3 d}$
derivativedivides	$-\frac{A}{8 d a^3 (\tan(dx+c)-i)^2} - \frac{3 i B}{8 d a^3 (\tan(dx+c)-i)^2} - \frac{i A \arctan(\tan(dx+c))}{8 d a^3} - \frac{B \arctan(\tan(dx+c))}{8 d a^3} - \frac{i A}{6 d a^3 (\tan(dx+c)-i)}$
default	$-\frac{A}{8 d a^3 (\tan(dx+c)-i)^2} - \frac{3 i B}{8 d a^3 (\tan(dx+c)-i)^2} - \frac{i A \arctan(\tan(dx+c))}{8 d a^3} - \frac{B \arctan(\tan(dx+c))}{8 d a^3} - \frac{i A}{6 d a^3 (\tan(dx+c)-i)}$

input `int(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output
$$-1/8*x/a^3*B-1/8*I*x/a^3*A+1/16*I/a^3/d*\exp(-2*I*(d*x+c))*B+1/16/a^3/d*\exp(-2*I*(d*x+c))*A+1/32*I/a^3/d*\exp(-4*I*(d*x+c))*B-1/32/a^3/d*\exp(-4*I*(d*x+c))*A-1/48*I/a^3/d*\exp(-6*I*(d*x+c))*B-1/48/a^3/d*\exp(-6*I*(d*x+c))*A$$

3.54.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.67

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \frac{(12(iA+B)dx e^{(6i dx+6i c)} - 6(A+iB)e^{(4i dx+4i c)} + 3(A-iB)e^{(2i dx+2i c)} + 2A+2iB)e^{(-6i dx-6i c)}}{96 a^3 d}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="fracas")`

output
$$-1/96*(12*(I*A+B)*d*x*e^{(6*I*d*x+6*I*c)}-6*(A+I*B)*e^{(4*I*d*x+4*I*c)}+3*(A-I*B)*e^{(2*I*d*x+2*I*c)}+2*A+2*I*B)*e^{(-6*I*d*x-6*I*c)}/(a^3*d)$$

3.54.6 Sympy [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 260, normalized size of antiderivative = 2.36

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$= \begin{cases} \frac{((-512Aa^6d^2e^{6ic}-512iBa^6d^2e^{6ic})e^{-6idx}+(-768Aa^6d^2e^{8ic}+768iBa^6d^2e^{8ic})e^{-4idx}+(1536Aa^6d^2e^{10ic}+1536iBa^6d^2e^{10ic})e^{-2idx})e^{-12ic}}{24576a^9d^3} \\ x\left(-\frac{-iA-B}{8a^3} + \frac{(-iAe^{6ic}-iAe^{4ic}+iAe^{2ic}+iA-Be^{6ic}+Be^{4ic}+Be^{2ic}-B)e^{-6ic}}{8a^3}\right) \\ + \frac{x(-iA-B)}{8a^3} \end{cases}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)`

output `Piecewise(((((-512*A*a**6*d**2*exp(6*I*c) - 512*I*B*a**6*d**2*exp(6*I*c))*exp(-6*I*d*x) + (-768*A*a**6*d**2*exp(8*I*c) + 768*I*B*a**6*d**2*exp(8*I*c))*exp(-4*I*d*x) + (1536*A*a**6*d**2*exp(10*I*c) + 1536*I*B*a**6*d**2*exp(10*I*c))*exp(-2*I*d*x))*exp(-12*I*c)/(24576*a**9*d**3), Ne(a**9*d**3*exp(12*I*c), 0)), (x*(-(-I*A - B)/(8*a**3) + (-I*A*exp(6*I*c) - I*A*exp(4*I*c) + I*A*exp(2*I*c) + I*A - B*exp(6*I*c) + B*exp(4*I*c) + B*exp(2*I*c) - B)*exp(-6*I*c)/(8*a**3)), True)) + x*(-I*A - B)/(8*a**3)`

3.54.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.54.8 Giac [A] (verification not implemented)

Time = 0.56 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.15

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$= \frac{\frac{6(A-iB)\log(\tan(dx+c)+i)}{a^3} - \frac{6(A-iB)\log(\tan(dx+c)-i)}{a^3} + \frac{11A\tan(dx+c)^3 - 11iB\tan(dx+c)^3 - 45iA\tan(dx+c)^2 - 45B\tan(dx+c)^2 - 69iA - 69B}{a^3(\tan(dx+c)-i)^3}}{96d}$$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")
```

```
output 1/96*(6*(A - I*B)*log(tan(d*x + c) + I)/a^3 - 6*(A - I*B)*log(tan(d*x + c) - I)/a^3 + (11*A*tan(d*x + c)^3 - 11*I*B*tan(d*x + c)^3 - 45*I*A*tan(d*x + c)^2 - 45*B*tan(d*x + c)^2 - 69*A*tan(d*x + c) + 21*I*B*tan(d*x + c) + 19*I*A + 3*B)/(a^3*(tan(d*x + c) - I)^3)/d
```

3.54.9 Mupad [B] (verification not implemented)

Time = 7.12 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.34

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$= \frac{\frac{A}{12a^3} - \tan(c+dx)^2 \left(\frac{A}{8a^3} - \frac{B1i}{8a^3} \right) + \frac{B1i}{12a^3} + \tan(c+dx) \left(-\frac{B}{8a^3} + \frac{A3i}{8a^3} \right)}{d \left(-\tan(c+dx)^3 1i - 3\tan(c+dx)^2 + \tan(c+dx) 3i + 1 \right)} + \frac{\ln(\tan(c+dx) - i) (B + A 1i) 1i}{16a^3 d} + \frac{\ln(\tan(c+dx) + i) (A - B 1i)}{16a^3 d}$$

```
input int((tan(c + d*x)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3,x)
```

```
output (A/(12*a^3) - tan(c + d*x)^2*(A/(8*a^3) - (B*1i)/(8*a^3)) + (B*1i)/(12*a^3) + tan(c + d*x)*((A*3i)/(8*a^3) - B/(8*a^3)))/(d*(tan(c + d*x)*3i - 3*tan(c + d*x)^2 - tan(c + d*x)^3*1i + 1)) + (log(tan(c + d*x) - 1i)*(A*1i + B*1i))/(16*a^3*d) + (log(tan(c + d*x) + 1i)*(A - B*1i))/(16*a^3*d)
```

3.55 $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^3} dx$

3.55.1	Optimal result	710
3.55.2	Mathematica [A] (verified)	710
3.55.3	Rubi [A] (verified)	711
3.55.4	Maple [A] (verified)	712
3.55.5	Fricas [A] (verification not implemented)	713
3.55.6	Sympy [A] (verification not implemented)	713
3.55.7	Maxima [F(-2)]	714
3.55.8	Giac [A] (verification not implemented)	714
3.55.9	Mupad [B] (verification not implemented)	715

3.55.1 Optimal result

Integrand size = 26, antiderivative size = 112

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^3} dx = \frac{(A - iB)x}{8a^3} + \frac{iA - B}{6d(a + ia \tan(c + dx))^3} + \frac{iA + B}{8ad(a + ia \tan(c + dx))^2} + \frac{iA + B}{8d(a^3 + ia^3 \tan(c + dx))}$$

```
output 1/8*(A-I*B)*x/a^3+1/6*(I*A-B)/d/(a+I*a*tan(d*x+c))^3+1/8*(I*A+B)/a/d/(a+I*a*tan(d*x+c))^2+1/8*(I*A+B)/d/(a^3+I*a^3*tan(d*x+c))
```

3.55.2 Mathematica [A] (verified)

Time = 1.11 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.97

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^3} dx = \frac{-10A + 2iB + 3(iA + B) \arctan(\tan(c + dx)) \sec^3(c + dx)(\cos(3(c + dx)) + i \sin(3(c + dx))) + (-9iA - 9iB) \tan(c + dx)}{24a^3d(-i + \tan(c + dx))^3}$$

```
input Integrate[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x])^3,x]
```

```
output (-10*A + (2*I)*B + 3*(I*A + B)*ArcTan[Tan[c + d*x]]*Sec[c + d*x]^3*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)]) + ((-9*I)*A - 9*B)*Tan[c + d*x] + 3*(A - I*B)*Tan[c + d*x]^2)/(24*a^3*d*(-I + Tan[c + d*x])^3)
```

3.55.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.99, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.269$, Rules used = {3042, 4009, 3042, 3960, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^3} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^3} dx \\
 & \quad \downarrow 4009 \\
 & \frac{(A - iB) \int \frac{1}{(i \tan(c+dx)a+a)^2} dx}{2a} + \frac{-B + iA}{6d(a + ia \tan(c + dx))^3} \\
 & \quad \downarrow 3042 \\
 & \frac{(A - iB) \int \frac{1}{(i \tan(c+dx)a+a)^2} dx}{2a} + \frac{-B + iA}{6d(a + ia \tan(c + dx))^3} \\
 & \quad \downarrow 3960 \\
 & \frac{(A - iB) \left(\frac{\int \frac{1}{i \tan(c+dx)a+a} dx}{2a} + \frac{i}{4d(a+ia \tan(c+dx))^2} \right)}{2a} + \frac{-B + iA}{6d(a + ia \tan(c + dx))^3} \\
 & \quad \downarrow 3042 \\
 & \frac{(A - iB) \left(\frac{\int \frac{1}{i \tan(c+dx)a+a} dx}{2a} + \frac{i}{4d(a+ia \tan(c+dx))^2} \right)}{2a} + \frac{-B + iA}{6d(a + ia \tan(c + dx))^3} \\
 & \quad \downarrow 3960 \\
 & \frac{(A - iB) \left(\frac{\int \frac{1 dx}{2a} + \frac{i}{2d(a+ia \tan(c+dx))}}{2a} + \frac{i}{4d(a+ia \tan(c+dx))^2} \right)}{2a} + \frac{-B + iA}{6d(a + ia \tan(c + dx))^3} \\
 & \quad \downarrow 24 \\
 & \frac{-B + iA}{6d(a + ia \tan(c + dx))^3} + \frac{(A - iB) \left(\frac{\frac{x}{2a} + \frac{i}{2d(a+ia \tan(c+dx))}}{2a} + \frac{i}{4d(a+ia \tan(c+dx))^2} \right)}{2a}
 \end{aligned}$$

3.55. $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^3} dx$

input `Int[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x])^3,x]`

output
$$\frac{(I*A - B)/(6*d*(a + I*a*Tan[c + d*x])^3) + ((A - I*B)*((I/4)/(d*(a + I*a*Tan[c + d*x])^2) + (x/(2*a) + (I/2)/(d*(a + I*a*Tan[c + d*x]))/(2*a)))/(2*a)}{a}$$

3.55.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

rule 4009 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

3.55.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.14

method	result
risch	$-\frac{ixB}{8a^3} + \frac{xA}{8a^3} + \frac{e^{-2i(dx+c)}B}{16a^3d} + \frac{3ie^{-2i(dx+c)}A}{16a^3d} - \frac{e^{-4i(dx+c)}B}{32a^3d} + \frac{3ie^{-4i(dx+c)}A}{32a^3d} - \frac{e^{-6i(dx+c)}B}{48a^3d} + \frac{ie^{-6i(dx+c)}A}{48a^3d}$
derivativedivides	$\frac{A \arctan(\tan(dx+c))}{8da^3} - \frac{iB \arctan(\tan(dx+c))}{8da^3} - \frac{iA}{8da^3(\tan(dx+c)-i)^2} - \frac{B}{8da^3(\tan(dx+c)-i)^2} + \frac{A}{8da^3(\tan(dx+c)+i)^2}$
default	$\frac{A \arctan(\tan(dx+c))}{8da^3} - \frac{iB \arctan(\tan(dx+c))}{8da^3} - \frac{iA}{8da^3(\tan(dx+c)-i)^2} - \frac{B}{8da^3(\tan(dx+c)-i)^2} + \frac{A}{8da^3(\tan(dx+c)+i)^2}$

input `int((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

$$3.55. \quad \int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^3} dx$$

output $-1/8*I*x/a^3*B+1/8*x/a^3*A+1/16/a^3/d*\exp(-2*I*(d*x+c))*B+3/16*I/a^3/d*\exp(-2*I*(d*x+c))*A-1/32/a^3/d*\exp(-4*I*(d*x+c))*B+3/32*I/a^3/d*\exp(-4*I*(d*x+c))*A-1/48/a^3/d*\exp(-6*I*(d*x+c))*B+1/48*I/a^3/d*\exp(-6*I*(d*x+c))*A$

3.55.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.68

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^3} dx$$

$$= \frac{(12(A - iB)dx e^{(6i dx + 6i c)} - 6(-3iA - B)e^{(4i dx + 4i c)} - 3(-3iA + B)e^{(2i dx + 2i c)} + 2iA - 2B)e^{(-6i dx - 6i c)}}{96 a^3 d}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="fracas")`

output $1/96*(12*(A - I*B)*d*x*e^{(6*I*d*x + 6*I*c)} - 6*(-3*I*A - B)*e^{(4*I*d*x + 4*I*c)} - 3*(-3*I*A + B)*e^{(2*I*d*x + 2*I*c)} + 2*I*A - 2*B)*e^{(-6*I*d*x - 6*I*c)}/(a^3*d)$

3.55.6 Sympy [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 258, normalized size of antiderivative = 2.30

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^3} dx$$

$$= \left\{ \frac{((512iAa^6 d^2 e^{6ic} - 512Ba^6 d^2 e^{6ic})e^{-6idx} + (2304iAa^6 d^2 e^{8ic} - 768Ba^6 d^2 e^{8ic})e^{-4idx} + (4608iAa^6 d^2 e^{10ic} + 1536Ba^6 d^2 e^{10ic})e^{-2idx})e^{-12ic}}{24576a^9 d^3} \right.$$

$$\left. x \left(-\frac{A-iB}{8a^3} + \frac{(Ae^{6ic} + 3Ae^{4ic} + 3Ae^{2ic} + A - iBe^{6ic} - iBe^{4ic} + iBe^{2ic} + iB)e^{-6ic}}{8a^3} \right) \right.$$

$$\left. + \frac{x(A - iB)}{8a^3} \right.$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)`


```
output Piecewise((((512*I*A*a**6*d**2*exp(6*I*c) - 512*B*a**6*d**2*exp(6*I*c))*exp(-6*I*d*x) + (2304*I*A*a**6*d**2*exp(8*I*c) - 768*B*a**6*d**2*exp(8*I*c))*exp(-4*I*d*x) + (4608*I*A*a**6*d**2*exp(10*I*c) + 1536*B*a**6*d**2*exp(10*I*c))*exp(-2*I*d*x))*exp(-12*I*c)/(24576*a**9*d**3), Ne(a**9*d**3*exp(12*I*c), 0)), (x*(-(A - I*B)/(8*a**3) + (A*exp(6*I*c) + 3*A*exp(4*I*c) + 3*A*exp(2*I*c) + A - I*B*exp(6*I*c) - I*B*exp(4*I*c) + I*B*exp(2*I*c) + I*B)*exp(-6*I*c)/(8*a**3)), True)) + x*(A - I*B)/(8*a**3)
```

3.55.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^3} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.55.8 Giac [A] (verification not implemented)

Time = 0.54 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.15

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^3} dx = \frac{\frac{6(-iA-B)\log(\tan(dx+c)+i)}{a^3} + \frac{6(iA+B)\log(\tan(dx+c)-i)}{a^3} + \frac{-11iA \tan(dx+c)^3 - 11B \tan(dx+c)^3 - 45A \tan(dx+c)^2 + 45iB \tan(dx+c)}{a^3(\tan(dx+c)-i)^3}}{96d}$$

```
input integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")
```

```
output -1/96*(6*(-I*A - B)*log(tan(d*x + c) + I)/a^3 + 6*(I*A + B)*log(tan(d*x + c) - I)/a^3 + (-11*I*A*tan(d*x + c)^3 - 11*B*tan(d*x + c)^3 - 45*A*tan(d*x + c)^2 + 45*I*B*tan(d*x + c)^2 + 69*I*A*tan(d*x + c) + 69*B*tan(d*x + c) + 51*A - 19*I*B)/(a^3*(tan(d*x + c) - I)^3)/d
```

3.55.9 Mupad [B] (verification not implemented)

Time = 7.61 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.99

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^3} dx$$

$$= -\frac{\tan(c + dx)^2 \left(\frac{B}{8a^3} + \frac{A1i}{8a^3}\right) - \frac{A5i}{12a^3} - \frac{B}{12a^3} + \tan(c + dx) \left(\frac{3A}{8a^3} - \frac{B3i}{8a^3}\right)}{d \left(-\tan(c + dx)^3 1i - 3 \tan(c + dx)^2 + \tan(c + dx) 3i + 1\right)} - \frac{x(B + A 1i) 1i}{8a^3}$$

input `int((A + B*tan(c + d*x))/(a + a*tan(c + d*x)*1i)^3,x)`output `- (tan(c + d*x)^2*((A*1i)/(8*a^3) + B/(8*a^3)) - (A*5i)/(12*a^3) - B/(12*a^3) + tan(c + d*x)*((3*A)/(8*a^3) - (B*3i)/(8*a^3)))/(d*(tan(c + d*x)*3i - 3*tan(c + d*x)^2 - tan(c + d*x)^3*1i + 1)) - (x*(A*1i + B)*1i)/(8*a^3)`

3.56 $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

3.56.1	Optimal result	716
3.56.2	Mathematica [A] (verified)	716
3.56.3	Rubi [A] (verified)	717
3.56.4	Maple [A] (verified)	720
3.56.5	Fricas [A] (verification not implemented)	720
3.56.6	Sympy [A] (verification not implemented)	721
3.56.7	Maxima [F(-2)]	721
3.56.8	Giac [A] (verification not implemented)	722
3.56.9	Mupad [B] (verification not implemented)	722

3.56.1 Optimal result

Integrand size = 32, antiderivative size = 131

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = -\frac{(7iA-B)x}{8a^3} + \frac{A \log(\sin(c+dx))}{a^3d}$$

$$+ \frac{A+iB}{6d(a+ia \tan(c+dx))^3}$$

$$+ \frac{3A+iB}{8ad(a+ia \tan(c+dx))^2}$$

$$+ \frac{7A+iB}{8d(a^3+ia^3 \tan(c+dx))}$$

output `-1/8*(7*I*A-B)*x/a^3+A*ln(sin(d*x+c))/a^3/d+1/6*(A+I*B)/d/(a+I*a*tan(d*x+c))^3+1/8*(3*A+I*B)/a/d/(a+I*a*tan(d*x+c))^2+1/8*(7*A+I*B)/d/(a^3+I*a^3*tan(d*x+c))`

3.56.2 Mathematica [A] (verified)

Time = 1.60 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.01

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{-3(15A+iB) \log(i-\tan(c+dx)) + 48A \log(\tan(c+dx)) - 3(A-iB) \log(i+\tan(c+dx)) + \frac{8i(A+iB)}{-i+\tan(c+dx)}}{48a^3d}$$

input `Integrate[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `(-3*(15*A + I*B)*Log[I - Tan[c + d*x]] + 48*A*Log[Tan[c + d*x]] - 3*(A - I*B)*Log[I + Tan[c + d*x]] + ((8*I)*(A + I*B))/(-I + Tan[c + d*x])^3 - (6*(3*A + I*B))/(-I + Tan[c + d*x])^2 + (6*((-7*I)*A + B))/(-I + Tan[c + d*x])/(48*a^3*d)`

3.56.3 Rubi [A] (verified)

Time = 0.90 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.17, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.406$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)(a+ia \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{3 \cot(c+dx)(2aA-a(iA-B) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{6a^2} + \frac{A+iB}{6d(a+ia \tan(c+dx))^3} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\cot(c+dx)(2aA-a(iA-B) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{2a^2} + \frac{A+iB}{6d(a+ia \tan(c+dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{2aA-a(iA-B) \tan(c+dx)}{\tan(c+dx)(i \tan(c+dx)a+a)^2} dx}{2a^2} + \frac{A+iB}{6d(a+ia \tan(c+dx))^3} \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{2 \cot(c+dx)(4a^2A-a^2(3iA-B) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{4a^2} + \frac{a(3A+iB)}{4d(a+ia \tan(c+dx))^2} + \frac{A+iB}{6d(a+ia \tan(c+dx))^3} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.56. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
& \frac{\int \frac{\cot(c+dx)(4a^2A - a^2(3iA-B)\tan(c+dx))}{i \tan(c+dx)a+a} dx}{2a^2} + \frac{a(3A+iB)}{4d(a+ia \tan(c+dx))^2} + \frac{A+iB}{6d(a+ia \tan(c+dx))^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{4a^2A - a^2(3iA-B)\tan(c+dx)}{\tan(c+dx)(i \tan(c+dx)a+a)} dx}{2a^2} + \frac{a(3A+iB)}{4d(a+ia \tan(c+dx))^2} + \frac{A+iB}{6d(a+ia \tan(c+dx))^3} \\
& \quad \downarrow \text{4079} \\
& \frac{\int \frac{\cot(c+dx)(8a^3A - a^3(7iA-B)\tan(c+dx))}{2a^2} dx}{2a^2} + \frac{a^2(7A+iB)}{2d(a+ia \tan(c+dx))} + \frac{a(3A+iB)}{4d(a+ia \tan(c+dx))^2} + \frac{A+iB}{6d(a+ia \tan(c+dx))^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{8a^3A - a^3(7iA-B)\tan(c+dx)}{\tan(c+dx)} dx}{2a^2} + \frac{a^2(7A+iB)}{2d(a+ia \tan(c+dx))} + \frac{a(3A+iB)}{4d(a+ia \tan(c+dx))^2} + \frac{A+iB}{6d(a+ia \tan(c+dx))^3} \\
& \quad \downarrow \text{4014} \\
& \frac{8a^3A \int \cot(c+dx) dx - a^3x(-B+7iA)}{2a^2} + \frac{a^2(7A+iB)}{2d(a+ia \tan(c+dx))} + \frac{a(3A+iB)}{4d(a+ia \tan(c+dx))^2} + \frac{A+iB}{6d(a+ia \tan(c+dx))^3} \\
& \quad \downarrow \text{3042} \\
& \frac{8a^3A \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx - a^3x(-B+7iA)}{2a^2} + \frac{a^2(7A+iB)}{2d(a+ia \tan(c+dx))} + \frac{a(3A+iB)}{4d(a+ia \tan(c+dx))^2} + \frac{A+iB}{6d(a+ia \tan(c+dx))^3} \\
& \quad \downarrow \text{25} \\
& \frac{-8a^3A \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx - (a^3x(-B+7iA))}{2a^2} + \frac{a^2(7A+iB)}{2d(a+ia \tan(c+dx))} + \frac{a(3A+iB)}{4d(a+ia \tan(c+dx))^2} + \frac{A+iB}{6d(a+ia \tan(c+dx))^3} \\
& \quad \downarrow \text{3956} \\
& \frac{\frac{a^2(7A+iB)}{2d(a+ia \tan(c+dx))} + \frac{8a^3A \log\left(\frac{-\sin(c+dx)}{d}\right) - a^3x(-B+7iA)}{2a^2}}{2a^2} + \frac{a(3A+iB)}{4d(a+ia \tan(c+dx))^2} + \frac{A+iB}{6d(a+ia \tan(c+dx))^3}
\end{aligned}$$

input `Int[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

3.56. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

```
output (A + I*B)/(6*d*(a + I*a*Tan[c + d*x])^3) + ((a*(3*A + I*B))/(4*d*(a + I*a*
Tan[c + d*x])^2) + ((-(a^3*((7*I)*A - B)*x) + (8*a^3*A*Log[-Sin[c + d*x]])
/d)/(2*a^2) + (a^2*(7*A + I*B))/(2*d*(a + I*a*Tan[c + d*x])))/(2*a^2))/(2*
a^2)
```

3.56.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4079 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.56.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.21

method	result
risch	$\frac{x B}{8a^3} - \frac{15ixA}{8a^3} + \frac{3ie^{-2i(dx+c)}B}{16a^3d} + \frac{11e^{-2i(dx+c)}A}{16a^3d} + \frac{3ie^{-4i(dx+c)}B}{32a^3d} + \frac{5e^{-4i(dx+c)}A}{32a^3d} + \frac{ie^{-6i(dx+c)}B}{48a^3d} + \frac{e^{-6i(dx+c)}A}{48a^3d}$
derivativedivides	$-\frac{A \ln(1+\tan^2(dx+c))}{2a^3d} - \frac{7iA \arctan(\tan(dx+c))}{8da^3} + \frac{B \arctan(\tan(dx+c))}{8da^3} + \frac{iA}{6da^3(\tan(dx+c)-i)^3} - \frac{B}{6da^3(\tan(dx+c)-i)}$
default	$-\frac{A \ln(1+\tan^2(dx+c))}{2a^3d} - \frac{7iA \arctan(\tan(dx+c))}{8da^3} + \frac{B \arctan(\tan(dx+c))}{8da^3} + \frac{iA}{6da^3(\tan(dx+c)-i)^3} - \frac{B}{6da^3(\tan(dx+c)-i)}$

input `int(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/8*x/a^3*B-15/8*I*x/a^3*A+3/16*I/a^3/d*exp(-2*I*(d*x+c))*B+11/16/a^3/d*exp(-2*I*(d*x+c))*A+3/32*I/a^3/d*exp(-4*I*(d*x+c))*B+5/32/a^3/d*exp(-4*I*(d*x+c))*A+1/48*I/a^3/d*exp(-6*I*(d*x+c))*B+1/48/a^3/d*exp(-6*I*(d*x+c))*A-2*I*A/a^3/d*c+A/a^3/d*ln(exp(2*I*(d*x+c))-1)`

3.56.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.79

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \frac{(12(15iA-B)dx e^{(6i dx+6i c)} - 96A e^{(6i dx+6i c)} \log(e^{(2i dx+2i c)} - 1) - 6(11A+3iB)e^{(4i dx+4i c)} - 3(5A - B))}{96a^3d}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="fricas")`

output `-1/96*(12*(15*I*A - B)*d*x*e^(6*I*d*x + 6*I*c) - 96*A*e^(6*I*d*x + 6*I*c)*log(e^(2*I*d*x + 2*I*c) - 1) - 6*(11*A + 3*I*B)*e^(4*I*d*x + 4*I*c) - 3*(5*A + 3*I*B)*e^(2*I*d*x + 2*I*c) - 2*A - 2*I*B)*e^(-6*I*d*x - 6*I*c)/(a^3*d)`

3.56.6 Sympy [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 292, normalized size of antiderivative = 2.23

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \frac{A \log(e^{2idx} - e^{-2ic})}{a^3 d}$$

$$+ \left\{ \frac{((512Aa^6 d^2 e^{6ic} + 512iBa^6 d^2 e^{6ic})e^{-6idx} + (3840Aa^6 d^2 e^{8ic} + 2304iBa^6 d^2 e^{8ic})e^{-4idx} + (16896Aa^6 d^2 e^{10ic} + 4608iBa^6 d^2 e^{10ic})e^{-2idx})e^{-12ic}}{24576a^9 d^3} \right.$$

$$+ \left. x \left(-\frac{15iA+B}{8a^3} + \frac{(-15iAe^{6ic} - 11iAe^{4ic} - 5iAe^{2ic} - iA + Be^{6ic} + 3Be^{4ic} + 3Be^{2ic} + B)e^{-6ic}}{8a^3} \right) \right.$$

$$+ \frac{x(-15iA+B)}{8a^3}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)`

output `A*log(exp(2*I*d*x) - exp(-2*I*c))/(a**3*d) + Piecewise((((512*A*a**6*d**2*exp(6*I*c) + 512*I*B*a**6*d**2*exp(6*I*c))*exp(-6*I*d*x) + (3840*A*a**6*d**2*exp(8*I*c) + 2304*I*B*a**6*d**2*exp(8*I*c))*exp(-4*I*d*x) + (16896*A*a**6*d**2*exp(10*I*c) + 4608*I*B*a**6*d**2*exp(10*I*c))*exp(-2*I*d*x))*exp(-12*I*c)/(24576*a**9*d**3), Ne(a**9*d**3*exp(12*I*c), 0)), (x*(-(-15*I*A + B)/(8*a**3) + (-15*I*A*exp(6*I*c) - 11*I*A*exp(4*I*c) - 5*I*A*exp(2*I*c) - I*A + B*exp(6*I*c) + 3*B*exp(4*I*c) + 3*B*exp(2*I*c) + B)*exp(-6*I*c)/(8*a**3)), True)) + x*(-15*I*A + B)/(8*a**3)`

3.56.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.56.8 Giac [A] (verification not implemented)

Time = 0.95 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.09

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx =$$

$$\frac{6(A-iB) \log(\tan(dx+c)+i)}{a^3} + \frac{6(15A+iB) \log(\tan(dx+c)-i)}{a^3} - \frac{96A \log(\tan(dx+c))}{a^3} - \frac{165A \tan(dx+c)^3 + 11iB \tan(dx+c)^3 - 579iA \tan(dx+c)^2 - 699iB \tan(dx+c)^2 + 301iA - 51iB}{a^3(\tan(dx+c) - i)}$$

$$96d$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output `-1/96*(6*(A - I*B)*log(tan(d*x + c) + I)/a^3 + 6*(15*A + I*B)*log(tan(d*x + c) - I)/a^3 - 96*A*log(tan(d*x + c))/a^3 - (165*A*tan(d*x + c)^3 + 11*I*B*tan(d*x + c)^3 - 579*I*A*tan(d*x + c)^2 + 45*B*tan(d*x + c)^2 - 699*A*tan(d*x + c) - 69*I*B*tan(d*x + c) + 301*I*A - 51*B)/(a^3*(tan(d*x + c) - I)^3)/d`

3.56.9 Mupad [B] (verification not implemented)

Time = 8.30 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.25

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{\frac{17A}{12a^3} - \tan(c+dx)^2 \left(\frac{7A}{8a^3} + \frac{B1i}{8a^3} \right) + \frac{B5i}{12a^3} + \tan(c+dx) \left(-\frac{3B}{8a^3} + \frac{A17i}{8a^3} \right)}{d \left(-\tan(c+dx)^3 1i - 3 \tan(c+dx)^2 + \tan(c+dx) 3i + 1 \right)}$$

$$+ \frac{A \ln(\tan(c+dx))}{a^3 d} + \frac{\ln(\tan(c+dx) + 1i) (B + A 1i) 1i}{16 a^3 d}$$

$$- \frac{\ln(\tan(c+dx) - i) (15 A + B 1i)}{16 a^3 d}$$

input `int((cot(c + d*x)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3,x)`

output `((17*A)/(12*a^3) - tan(c + d*x)^2*((7*A)/(8*a^3) + (B*1i)/(8*a^3)) + (B*5i)/(12*a^3) + tan(c + d*x)*((A*17i)/(8*a^3) - (3*B)/(8*a^3)))/(d*(tan(c + d*x)*3i - 3*tan(c + d*x)^2 - tan(c + d*x)^3*1i + 1)) + (A*log(tan(c + d*x)))/(a^3*d) + (log(tan(c + d*x) + 1i)*(A*1i + B)*1i)/(16*a^3*d) - (log(tan(c + d*x) - 1i)*(15*A + B*1i))/(16*a^3*d)`

3.56. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

3.57 $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

3.57.1 Optimal result 723
 3.57.2 Mathematica [C] (verified) 724
 3.57.3 Rubi [A] (verified) 724
 3.57.4 Maple [A] (verified) 728
 3.57.5 Fricas [A] (verification not implemented) 729
 3.57.6 Sympy [A] (verification not implemented) 729
 3.57.7 Maxima [F(-2)] 730
 3.57.8 Giac [A] (verification not implemented) 730
 3.57.9 Mupad [B] (verification not implemented) 731

3.57.1 Optimal result

Integrand size = 34, antiderivative size = 183

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = -\frac{(25A+7iB)x}{8a^3} - \frac{(25A+7iB) \cot(c+dx)}{8a^3d} - \frac{(3iA-B) \log(\sin(c+dx))}{a^3d} + \frac{(A+iB) \cot(c+dx)}{6d(a+ia \tan(c+dx))^3} + \frac{(11A+5iB) \cot(c+dx)}{24ad(a+ia \tan(c+dx))^2} + \frac{(3A+iB) \cot(c+dx)}{2d(a^3+ia^3 \tan(c+dx))}$$

```
output -1/8*(25*A+7*I*B)*x/a^3-1/8*(25*A+7*I*B)*cot(d*x+c)/a^3/d-(3*I*A-B)*ln(sin
(d*x+c))/a^3/d+1/6*(A+I*B)*cot(d*x+c)/d/(a+I*a*tan(d*x+c))^3+1/24*(11*A+5*
I*B)*cot(d*x+c)/a/d/(a+I*a*tan(d*x+c))^2+1/2*(3*A+I*B)*cot(d*x+c)/d/(a^3+I
*a^3*tan(d*x+c))
```

3.57.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 2.58 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.90

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{\frac{4(A+iB) \cot^4(c+dx)}{(i+\cot(c+dx))^3} + \frac{(11A+5iB) \cot^3(c+dx)}{(i+\cot(c+dx))^2} + \frac{12(3A+iB) \cot^2(c+dx)}{i+\cot(c+dx)} - 3((25A+7iB) \cot(c+dx) \operatorname{Hypergeometric2F1}[-1/2, 1, 1/2, -\tan(c+dx)^2] + 8((3i)A-B)(\operatorname{Log}[\cos(c+dx)] + \operatorname{Log}[\tan(c+dx)]))}{24a^3d}$$

input `Integrate[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `((4*(A + I*B)*Cot[c + d*x]^4)/(I + Cot[c + d*x])^3 + ((11*A + (5*I)*B)*Cot[c + d*x]^3)/(I + Cot[c + d*x])^2 + (12*(3*A + I*B)*Cot[c + d*x]^2)/(I + Cot[c + d*x]) - 3*((25*A + (7*I)*B)*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2] + 8*((3*I)*A - B)*(Log[Cos[c + d*x]] + Log[Tan[c + d*x]])))/(24*a^3*d)`

3.57.3 Rubi [A] (verified)

Time = 1.25 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.08, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.471$, Rules used = {3042, 4079, 3042, 4079, 27, 3042, 4079, 27, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A+B \tan(c+dx)}{\tan(c+dx)^2(a+ia \tan(c+dx))^3} dx$$

$$\downarrow \text{4079}$$

$$\frac{\int \frac{\cot^2(c+dx)(a(7A+iB)-4a(iA-B) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{6a^2} + \frac{(A+iB) \cot(c+dx)}{6d(a+ia \tan(c+dx))^3}$$

3.57. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{a(7A+iB)-4a(iA-B)\tan(c+dx)}{\tan(c+dx)^2(i\tan(c+dx)a+a)^2} dx + \frac{(A+iB)\cot(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{3\cot^2(c+dx)(a^2(13A+3iB)-a^2(11iA-5B)\tan(c+dx))}{i\tan(c+dx)a+a} dx + \frac{a(11A+5iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2}}{6a^2} + \frac{(A+iB)\cot(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 4079 \\
& \frac{3\int \frac{\cot^2(c+dx)(a^2(13A+3iB)-a^2(11iA-5B)\tan(c+dx))}{i\tan(c+dx)a+a} dx + \frac{a(11A+5iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2}}{6a^2} + \frac{(A+iB)\cot(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 27 \\
& \frac{3\int \frac{\cot^2(c+dx)(a^2(13A+3iB)-a^2(11iA-5B)\tan(c+dx))}{i\tan(c+dx)a+a} dx + \frac{a(11A+5iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2}}{6a^2} + \frac{(A+iB)\cot(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 3042 \\
& \frac{3\int \frac{a^2(13A+3iB)-a^2(11iA-5B)\tan(c+dx)}{\tan(c+dx)^2(i\tan(c+dx)a+a)} dx + \frac{a(11A+5iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2}}{6a^2} + \frac{(A+iB)\cot(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 4079 \\
& \frac{3\left(\frac{\int 2\cot^2(c+dx)(a^3(25A+7iB)-8a^3(3iA-B)\tan(c+dx))}{2a^2} dx + \frac{4a^2(3A+iB)\cot(c+dx)}{d(a+ia\tan(c+dx))}\right)}{4a^2} + \frac{a(11A+5iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} + \\
& \quad \frac{6a^2}{(A+iB)\cot(c+dx)} \\
& \quad \frac{(A+iB)\cot(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 27 \\
& \frac{3\left(\frac{\int \cot^2(c+dx)(a^3(25A+7iB)-8a^3(3iA-B)\tan(c+dx))}{a^2} dx + \frac{4a^2(3A+iB)\cot(c+dx)}{d(a+ia\tan(c+dx))}\right)}{4a^2} + \frac{a(11A+5iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} + \\
& \quad \frac{6a^2}{(A+iB)\cot(c+dx)} \\
& \quad \frac{(A+iB)\cot(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 3042 \\
& \frac{3\left(\frac{\int \frac{a^3(25A+7iB)-8a^3(3iA-B)\tan(c+dx)}{\tan(c+dx)^2} dx + \frac{4a^2(3A+iB)\cot(c+dx)}{d(a+ia\tan(c+dx))}}{4a^2}\right)}{4a^2} + \frac{a(11A+5iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} + \\
& \quad \frac{6a^2}{(A+iB)\cot(c+dx)} \\
& \quad \frac{(A+iB)\cot(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 4012
\end{aligned}$$

3.57. $\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

$$3 \left(\frac{\int -\cot(c+dx) (8(3iA-B)a^3 + (25A+7iB)\tan(c+dx)a^3) dx - \frac{a^3(25A+7iB)\cot(c+dx)}{d}}{a^2} + \frac{4a^2(3A+iB)\cot(c+dx)}{d(a+ia\tan(c+dx))} \right) + \frac{a(11A+5iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} +$$

$$\frac{6a^2}{4a^2} \frac{(A+iB)\cot(c+dx)}{6d(a+ia\tan(c+dx))^3}$$

↓ 25

$$3 \left(\frac{-\int \cot(c+dx) (8(3iA-B)a^3 + (25A+7iB)\tan(c+dx)a^3) dx - \frac{a^3(25A+7iB)\cot(c+dx)}{d}}{a^2} + \frac{4a^2(3A+iB)\cot(c+dx)}{d(a+ia\tan(c+dx))} \right) + \frac{a(11A+5iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} +$$

$$\frac{6a^2}{4a^2} \frac{(A+iB)\cot(c+dx)}{6d(a+ia\tan(c+dx))^3}$$

↓ 3042

$$3 \left(\frac{-\int \frac{8(3iA-B)a^3 + (25A+7iB)\tan(c+dx)a^3}{\tan(c+dx)} dx - \frac{a^3(25A+7iB)\cot(c+dx)}{d}}{a^2} + \frac{4a^2(3A+iB)\cot(c+dx)}{d(a+ia\tan(c+dx))} \right) + \frac{a(11A+5iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} +$$

$$\frac{6a^2}{4a^2} \frac{(A+iB)\cot(c+dx)}{6d(a+ia\tan(c+dx))^3}$$

↓ 4014

$$3 \left(\frac{-8a^3(-B+3iA)\int \cot(c+dx) dx - \frac{a^3(25A+7iB)\cot(c+dx)}{d} - (a^3x(25A+7iB))}{a^2} + \frac{4a^2(3A+iB)\cot(c+dx)}{d(a+ia\tan(c+dx))} \right) + \frac{a(11A+5iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} +$$

$$\frac{6a^2}{4a^2} \frac{(A+iB)\cot(c+dx)}{6d(a+ia\tan(c+dx))^3}$$

↓ 3042

$$3 \left(\frac{-8a^3(-B+3iA)\int -\tan(c+dx+\frac{\pi}{2}) dx - \frac{a^3(25A+7iB)\cot(c+dx)}{d} - (a^3x(25A+7iB))}{a^2} + \frac{4a^2(3A+iB)\cot(c+dx)}{d(a+ia\tan(c+dx))} \right) + \frac{a(11A+5iB)\cot(c+dx)}{4d(a+ia\tan(c+dx))^2} +$$

$$\frac{6a^2}{4a^2} \frac{(A+iB)\cot(c+dx)}{6d(a+ia\tan(c+dx))^3}$$

↓ 25

3.57. $\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

$$\frac{3 \left(\frac{8a^3(-B+3iA) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx - \frac{a^3(25A+7iB) \cot(c+dx)}{d} - (a^3 x(25A+7iB)) + \frac{4a^2(3A+iB) \cot(c+dx)}{d(a+ia \tan(c+dx))} \right)}{4a^2} + \frac{a(11A+5iB) \cot(c+dx)}{4d(a+ia \tan(c+dx))^2} + \frac{6a^2 (A+iB) \cot(c+dx)}{6d(a+ia \tan(c+dx))^3}$$

↓ 3956

$$\frac{3 \left(\frac{4a^2(3A+iB) \cot(c+dx)}{d(a+ia \tan(c+dx))} + \frac{-\frac{a^3(25A+7iB) \cot(c+dx)}{d} - \frac{8a^3(-B+3iA) \log(-\sin(c+dx))}{a^2} - (a^3 x(25A+7iB)) \right)}{4a^2} + \frac{a(11A+5iB) \cot(c+dx)}{4d(a+ia \tan(c+dx))^2} + \frac{6a^2 (A+iB) \cot(c+dx)}{6d(a+ia \tan(c+dx))^3}$$

input `Int[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `((A + I*B)*Cot[c + d*x])/(6*d*(a + I*a*Tan[c + d*x])^3) + ((a*(11*A + (5*I)*B)*Cot[c + d*x])/(4*d*(a + I*a*Tan[c + d*x])^2) + (3*((-(a^3*(25*A + (7*I)*B)*x) - (a^3*(25*A + (7*I)*B)*Cot[c + d*x])/d - (8*a^3*((3*I)*A - B)*Log[-Sin[c + d*x]])/d)/a^2 + (4*a^2*(3*A + I*B)*Cot[c + d*x])/(d*(a + I*a*Tan[c + d*x])))/(4*a^2)/(6*a^2)`

3.57.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

3.57. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

```
rule 4012 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]
```

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4079 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d)), x] + Simp[1/(2*a*m*(b*c - a*d) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.57.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.17

method	result
risch	$-\frac{15xB}{8a^3} - \frac{49xA}{8a^3} + \frac{11e^{-2i(dx+c)}B}{16a^3d} - \frac{23ie^{-2i(dx+c)}A}{16a^3d} + \frac{5e^{-4i(dx+c)}B}{32a^3d} - \frac{7ie^{-4i(dx+c)}A}{32a^3d} + \frac{e^{-6i(dx+c)}B}{48a^3d}$
derivativedivides	$-\frac{A \cot(dx+c)}{a^3d} + \frac{3iA \ln(\cot^2(dx+c)+1)}{2a^3d} + \frac{25A(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)))}{8a^3d} - \frac{B \ln(\cot^2(dx+c)+1)}{2a^3d} + \frac{7iB(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)))}{8a^3d}$
default	$-\frac{A \cot(dx+c)}{a^3d} + \frac{3iA \ln(\cot^2(dx+c)+1)}{2a^3d} + \frac{25A(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)))}{8a^3d} - \frac{B \ln(\cot^2(dx+c)+1)}{2a^3d} + \frac{7iB(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)))}{8a^3d}$

```
input int(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVER
BOSE)
```

$$3.57. \quad \int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

output
$$-15/8*I*x/a^3*B-49/8*x/a^3*A+11/16/a^3/d*\exp(-2*I*(d*x+c))*B-23/16*I/a^3/d*\exp(-2*I*(d*x+c))*A+5/32/a^3/d*\exp(-4*I*(d*x+c))*B-7/32*I/a^3/d*\exp(-4*I*(d*x+c))*A+1/48/a^3/d*\exp(-6*I*(d*x+c))*B-1/48*I/a^3/d*\exp(-6*I*(d*x+c))*A-2*I/a^3/d*B*c-6/a^3/d*A*c-2*I*A/a^3/d/(\exp(2*I*(d*x+c))-1)+1/a^3/d*\ln(\exp(2*I*(d*x+c))-1)*B-3*I/a^3/d*\ln(\exp(2*I*(d*x+c))-1)*A$$

3.57.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.95

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \frac{12(49A+15iB)dx e^{(8i dx+8ic)} - 6(2(49A+15iB)dx - 55iA+11B)e^{(6i dx+6ic)} + 3(-39iA+17B)}{96(a^3}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="fricas")`

output
$$-1/96*(12*(49*A + 15*I*B)*d*x*e^{(8*I*d*x + 8*I*c)} - 6*(2*(49*A + 15*I*B)*d*x - 55*I*A + 11*B)*e^{(6*I*d*x + 6*I*c)} + 3*(-39*I*A + 17*B)*e^{(4*I*d*x + 4*I*c)} - (19*I*A - 13*B)*e^{(2*I*d*x + 2*I*c)} + 96*((3*I*A - B)*e^{(8*I*d*x + 8*I*c)} + (-3*I*A + B)*e^{(6*I*d*x + 6*I*c)})*\log(e^{(2*I*d*x + 2*I*c)} - 1) - 2*I*A + 2*B)/(a^3*d*e^{(8*I*d*x + 8*I*c)} - a^3*d*e^{(6*I*d*x + 6*I*c)})$$

3.57.6 Sympy [A] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 340, normalized size of antiderivative = 1.86

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = -\frac{2iA}{a^3 d e^{2ic} e^{2idx} - a^3 d} + \left\{ \frac{((-512iAa^6 d^2 e^{6ic} + 512Ba^6 d^2 e^{6ic})e^{-6idx} + (-5376iAa^6 d^2 e^{8ic} + 3840Ba^6 d^2 e^{8ic})e^{-4idx} + (-35328iAa^6 d^2 e^{10ic} + 16896Ba^6 d^2 e^{10ic})e^{-2idx})}{24576a^9 d^3} \right. \\ \left. + x \left(-\frac{49A-15iB}{8a^3} + \frac{(-49Ae^{6ic}-23Ae^{4ic}-7Ae^{2ic}-A-15iBe^{6ic}-11iBe^{4ic}-5iBe^{2ic}-iB)e^{-6ic}}{8a^3} \right) \right. \\ \left. + \frac{x(-49A-15iB)}{8a^3} - \frac{i(3A+iB)\log(e^{2idx} - e^{-2ic})}{a^3 d} \right.$$

input `integrate(cot(d*x+c)**2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)`

3.57.
$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$


```
output -2*I*A/(a**3*d*exp(2*I*c)*exp(2*I*d*x) - a**3*d) + Piecewise(((((-512*I*A*a
**6*d**2*exp(6*I*c) + 512*B*a**6*d**2*exp(6*I*c))*exp(-6*I*d*x) + (-5376*I
*A*a**6*d**2*exp(8*I*c) + 3840*B*a**6*d**2*exp(8*I*c))*exp(-4*I*d*x) + (-3
5328*I*A*a**6*d**2*exp(10*I*c) + 16896*B*a**6*d**2*exp(10*I*c))*exp(-2*I*d
*x))*exp(-12*I*c)/(24576*a**9*d**3), Ne(a**9*d**3*exp(12*I*c), 0)), (x*(-(
-49*A - 15*I*B)/(8*a**3) + (-49*A*exp(6*I*c) - 23*A*exp(4*I*c) - 7*A*exp(2
*I*c) - A - 15*I*B*exp(6*I*c) - 11*I*B*exp(4*I*c) - 5*I*B*exp(2*I*c) - I*B
)*exp(-6*I*c)/(8*a**3)), True)) + x*(-49*A - 15*I*B)/(8*a**3) - I*(3*A + I
*B)*log(exp(2*I*d*x) - exp(-2*I*c))/(a**3*d)
```

3.57.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^3} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm=
"maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.57.8 Giac [A] (verification not implemented)

Time = 0.91 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.99

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^3} dx =$$

$$\frac{6(iA+B)\log(\tan(dx+c)+i)}{a^3} + \frac{6(-49iA+15B)\log(\tan(dx+c)-i)}{a^3} + \frac{96(3iA-B)\log(\tan(dx+c))}{a^3} + \frac{96(-3iA\tan(dx+c)+B\tan(dx+c))}{a^3\tan(dx+c)}$$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm=
"giac")
```

3.57. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

output
$$\begin{aligned} & -1/96*(6*(I*A + B)*\log(\tan(d*x + c) + I)/a^3 + 6*(-49*I*A + 15*B)*\log(\tan(d*x + c) - I)/a^3 + 96*(3*I*A - B)*\log(\tan(d*x + c))/a^3 + 96*(-3*I*A*\tan(d*x + c) + B*\tan(d*x + c) + A)/(a^3*\tan(d*x + c)) + (539*A*\tan(d*x + c)^3 + 165*I*B*\tan(d*x + c)^3 - 1821*I*A*\tan(d*x + c)^2 + 579*B*\tan(d*x + c)^2 - 2085*A*\tan(d*x + c) - 699*I*B*\tan(d*x + c) + 819*I*A - 301*B)/(a^3*(I*\tan(d*x + c) + 1)^3))/d \end{aligned}$$

3.57.9 Mupad [B] (verification not implemented)

Time = 7.32 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.08

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^3} dx =$$

$$\frac{\tan(c + dx)^3 \left(\frac{25A}{8a^3} + \frac{B7i}{8a^3} \right) - \tan(c + dx)^2 \left(-\frac{17B}{8a^3} + \frac{A63i}{8a^3} \right) + \frac{A1i}{a^3} - \tan(c + dx) \left(\frac{71A}{12a^3} + \frac{B17i}{12a^3} \right)}{d \left(\tan(c + dx)^4 - \tan(c + dx)^3 3i - 3 \tan(c + dx)^2 + \tan(c + dx) 1i \right)}$$

$$- \frac{\ln(\tan(c + dx)) (-B + A3i)}{a^3 d} - \frac{\ln(\tan(c + dx) + 1i) (B + A1i)}{16 a^3 d}$$

$$+ \frac{\ln(\tan(c + dx) - i) (-15B + A49i)}{16 a^3 d}$$

input `int((cot(c + d*x)^2*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3,x)`

output
$$\begin{aligned} & (\log(\tan(c + d*x) - 1i)*(A*49i - 15*B))/(16*a^3*d) - (\log(\tan(c + d*x))*(A*3i - B))/(a^3*d) - (\log(\tan(c + d*x) + 1i)*(A*1i + B))/(16*a^3*d) - (\tan(c + d*x)^3*((25*A)/(8*a^3) + (B*7i)/(8*a^3)) - \tan(c + d*x)^2*((A*63i)/(8*a^3) - (17*B)/(8*a^3)) + (A*1i)/a^3 - \tan(c + d*x)*((71*A)/(12*a^3) + (B*17i)/(12*a^3)))/(d*(\tan(c + d*x)*1i - 3*\tan(c + d*x)^2 - \tan(c + d*x)^3*3i + \tan(c + d*x)^4)) \end{aligned}$$

3.58 $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

3.58.1	Optimal result	732
3.58.2	Mathematica [C] (verified)	733
3.58.3	Rubi [A] (verified)	733
3.58.4	Maple [A] (verified)	738
3.58.5	Fricas [A] (verification not implemented)	738
3.58.6	Sympy [A] (verification not implemented)	739
3.58.7	Maxima [F(-2)]	740
3.58.8	Giac [A] (verification not implemented)	740
3.58.9	Mupad [B] (verification not implemented)	741

3.58.1 Optimal result

Integrand size = 34, antiderivative size = 216

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{5(11iA-5B)x}{8a^3} + \frac{5(11iA-5B) \cot(c+dx)}{8a^3d} - \frac{(7A+3iB) \cot^2(c+dx)}{2a^3d}$$

$$- \frac{(7A+3iB) \log(\sin(c+dx))}{a^3d} + \frac{(A+iB) \cot^2(c+dx)}{6d(a+ia \tan(c+dx))^3}$$

$$+ \frac{(13A+7iB) \cot^2(c+dx)}{24ad(a+ia \tan(c+dx))^2} + \frac{5(11A+5iB) \cot^2(c+dx)}{24d(a^3+ia^3 \tan(c+dx))}$$

```
output 5/8*(11*I*A-5*B)*x/a^3+5/8*(11*I*A-5*B)*cot(d*x+c)/a^3/d-1/2*(7*A+3*I*B)*c
ot(d*x+c)^2/a^3/d-(7*A+3*I*B)*ln(sin(d*x+c))/a^3/d+1/6*(A+I*B)*cot(d*x+c)^
2/d/(a+I*a*tan(d*x+c))^3+1/24*(13*A+7*I*B)*cot(d*x+c)^2/a/d/(a+I*a*tan(d*x
+c))^2+5/24*(11*A+5*I*B)*cot(d*x+c)^2/d/(a^3+I*a^3*tan(d*x+c))
```

3.58.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 3.57 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.80

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{4(A+iB) \cot^5(c+dx)}{(i+\cot(c+dx))^3} + \frac{(13A+7iB) \cot^4(c+dx)}{(i+\cot(c+dx))^2} + \frac{5(11A+5iB) \cot^3(c+dx)}{i+\cot(c+dx)} + 15(11iA-5B) \cot(c+dx) \operatorname{Hypergeometric2F1}[-1/2, 1, 1/2, -\tan(c+dx)^2 - 12(7A+(3i)B)(\cot(c+dx)^2 + 2(\operatorname{Log}[\operatorname{Cos}[c+dx]] + \operatorname{Log}[\operatorname{Tan}[c+dx]]))]/(24a^3d)$$

input `Integrate[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `((4*(A + I*B)*Cot[c + d*x]^5)/(I + Cot[c + d*x])^3 + ((13*A + (7*I)*B)*Cot[c + d*x]^4)/(I + Cot[c + d*x])^2 + (5*(11*A + (5*I)*B)*Cot[c + d*x]^3)/(I + Cot[c + d*x]) + 15*((11*I)*A - 5*B)*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2 - 12*(7*A + (3*I)*B)*(Cot[c + d*x]^2 + 2*(Log[Cos[c + d*x]] + Log[Tan[c + d*x]]))))/(24*a^3*d)`

3.58.3 Rubi [A] (verified)

Time = 1.48 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.08, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.529$, Rules used = {3042, 4079, 3042, 4079, 27, 3042, 4079, 27, 3042, 4012, 25, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A+B \tan(c+dx)}{\tan(c+dx)^3(a+ia \tan(c+dx))^3} dx$$

$$\downarrow \text{4079}$$

$$\frac{\int \frac{\cot^3(c+dx)(2a(4A+iB)-5a(iA-B) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{6a^2} + \frac{(A+iB) \cot^2(c+dx)}{6d(a+ia \tan(c+dx))^3}$$

3.58. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{2a(4A+iB)-5a(iA-B)\tan(c+dx)}{\tan(c+dx)^3(i\tan(c+dx)a+a)^2} dx + \frac{(A+iB)\cot^2(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{2\cot^3(c+dx)(a^2(29A+11iB)-2a^2(13iA-7B)\tan(c+dx))}{i\tan(c+dx)a+a} dx}{4a^2} + \frac{a(13A+7iB)\cot^2(c+dx)}{4d(a+ia\tan(c+dx))^2} + \frac{(A+iB)\cot^2(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 4079 \\
& \frac{\int \frac{\cot^3(c+dx)(a^2(29A+11iB)-2a^2(13iA-7B)\tan(c+dx))}{i\tan(c+dx)a+a} dx}{2a^2} + \frac{a(13A+7iB)\cot^2(c+dx)}{4d(a+ia\tan(c+dx))^2} + \frac{(A+iB)\cot^2(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{a^2(29A+11iB)-2a^2(13iA-7B)\tan(c+dx)}{\tan(c+dx)^3(i\tan(c+dx)a+a)} dx}{2a^2} + \frac{a(13A+7iB)\cot^2(c+dx)}{4d(a+ia\tan(c+dx))^2} + \frac{(A+iB)\cot^2(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 3042 \\
& \frac{\int 3\cot^3(c+dx)(8a^3(7A+3iB)-5a^3(11iA-5B)\tan(c+dx))}{2a^2} dx + \frac{5a^2(11A+5iB)\cot^2(c+dx)}{2d(a+ia\tan(c+dx))} + \frac{a(13A+7iB)\cot^2(c+dx)}{4d(a+ia\tan(c+dx))^2} + \\
& \quad \frac{6a^2}{2a^2} \frac{(A+iB)\cot^2(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 27 \\
& \frac{3\int \cot^3(c+dx)(8a^3(7A+3iB)-5a^3(11iA-5B)\tan(c+dx))}{2a^2} dx + \frac{5a^2(11A+5iB)\cot^2(c+dx)}{2d(a+ia\tan(c+dx))} + \frac{a(13A+7iB)\cot^2(c+dx)}{4d(a+ia\tan(c+dx))^2} + \\
& \quad \frac{6a^2}{2a^2} \frac{(A+iB)\cot^2(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 3042 \\
& \frac{3\int \frac{8a^3(7A+3iB)-5a^3(11iA-5B)\tan(c+dx)}{\tan(c+dx)^3} dx}{2a^2} + \frac{5a^2(11A+5iB)\cot^2(c+dx)}{2d(a+ia\tan(c+dx))} + \frac{a(13A+7iB)\cot^2(c+dx)}{4d(a+ia\tan(c+dx))^2} + \\
& \quad \frac{6a^2}{2a^2} \frac{(A+iB)\cot^2(c+dx)}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 4012
\end{aligned}$$

3.58. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

$$\frac{3 \left(\int -\cot^2(c+dx) \left(5(11iA-5B)a^3 + 8(7A+3iB) \tan(c+dx)a^3 \right) dx - \frac{4a^3(7A+3iB) \cot^2(c+dx)}{d} \right)}{2a^2} + \frac{5a^2(11A+5iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{a(13A+7iB) \cot^2(c+dx)}{4d(a+ia \tan(c+dx))^2} +$$

$$\frac{6a^2}{6d(a+ia \tan(c+dx))^3} (A+iB) \cot^2(c+dx)$$

↓ 25

$$\frac{3 \left(-\int \cot^2(c+dx) \left(5(11iA-5B)a^3 + 8(7A+3iB) \tan(c+dx)a^3 \right) dx - \frac{4a^3(7A+3iB) \cot^2(c+dx)}{d} \right)}{2a^2} + \frac{5a^2(11A+5iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{a(13A+7iB) \cot^2(c+dx)}{4d(a+ia \tan(c+dx))^2} +$$

$$\frac{6a^2}{6d(a+ia \tan(c+dx))^3} (A+iB) \cot^2(c+dx)$$

↓ 3042

$$\frac{3 \left(-\int \frac{5(11iA-5B)a^3 + 8(7A+3iB) \tan(c+dx)a^3}{\tan(c+dx)^2} dx - \frac{4a^3(7A+3iB) \cot^2(c+dx)}{d} \right)}{2a^2} + \frac{5a^2(11A+5iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{a(13A+7iB) \cot^2(c+dx)}{4d(a+ia \tan(c+dx))^2} +$$

$$\frac{6a^2}{6d(a+ia \tan(c+dx))^3} (A+iB) \cot^2(c+dx)$$

↓ 4012

$$\frac{3 \left(-\int \cot(c+dx) \left(8a^3(7A+3iB) - 5a^3(11iA-5B) \tan(c+dx) \right) dx - \frac{4a^3(7A+3iB) \cot^2(c+dx)}{d} + \frac{5a^3(-5B+11iA) \cot(c+dx)}{d} \right)}{2a^2} + \frac{5a^2(11A+5iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{a(13A+7iB) \cot^2(c+dx)}{4d(a+ia \tan(c+dx))^2} +$$

$$\frac{6a^2}{6d(a+ia \tan(c+dx))^3} (A+iB) \cot^2(c+dx)$$

↓ 3042

$$\frac{3 \left(-\int \frac{8a^3(7A+3iB) - 5a^3(11iA-5B) \tan(c+dx)}{\tan(c+dx)} dx - \frac{4a^3(7A+3iB) \cot^2(c+dx)}{d} + \frac{5a^3(-5B+11iA) \cot(c+dx)}{d} \right)}{2a^2} + \frac{5a^2(11A+5iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{a(13A+7iB) \cot^2(c+dx)}{4d(a+ia \tan(c+dx))^2} +$$

$$\frac{6a^2}{6d(a+ia \tan(c+dx))^3} (A+iB) \cot^2(c+dx)$$

↓ 4014

3.58. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\frac{3 \left(-8a^3(7A+3iB) \int \cot(c+dx) dx - \frac{4a^3(7A+3iB)}{d} \cot^2(c+dx) + \frac{5a^3(-5B+11iA)}{d} \cot(c+dx) + 5a^3 x(-5B+11iA) \right)}{2a^2} + \frac{5a^2(11A+5iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{a(13A+7iB)}{4d(a+ia \tan(c+dx))}$$

$$\frac{(A+iB) \cot^2(c+dx)}{6d(a+ia \tan(c+dx))^3} \cdot 6a^2$$

↓ 3042

$$\frac{3 \left(-8a^3(7A+3iB) \int -\tan(c+dx+\frac{\pi}{2}) dx - \frac{4a^3(7A+3iB)}{d} \cot^2(c+dx) + \frac{5a^3(-5B+11iA)}{d} \cot(c+dx) + 5a^3 x(-5B+11iA) \right)}{2a^2} + \frac{5a^2(11A+5iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{a(13A+7iB)}{4d(a+ia \tan(c+dx))}$$

$$\frac{(A+iB) \cot^2(c+dx)}{6d(a+ia \tan(c+dx))^3} \cdot 6a^2$$

↓ 25

$$\frac{3 \left(8a^3(7A+3iB) \int \tan(\frac{1}{2}(2c+\pi)+dx) dx - \frac{4a^3(7A+3iB)}{d} \cot^2(c+dx) + \frac{5a^3(-5B+11iA)}{d} \cot(c+dx) + 5a^3 x(-5B+11iA) \right)}{2a^2} + \frac{5a^2(11A+5iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{a(13A+7iB)}{4d(a+ia \tan(c+dx))}$$

$$\frac{(A+iB) \cot^2(c+dx)}{6d(a+ia \tan(c+dx))^3} \cdot 6a^2$$

↓ 3956

$$\frac{5a^2(11A+5iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{3 \left(-\frac{4a^3(7A+3iB)}{d} \cot^2(c+dx) + \frac{5a^3(-5B+11iA)}{d} \cot(c+dx) - \frac{8a^3(7A+3iB) \log(-\sin(c+dx))}{d} + 5a^3 x(-5B+11iA) \right)}{2a^2} + \frac{a(13A+7iB)}{4d(a+ia \tan(c+dx))}$$

$$\frac{(A+iB) \cot^2(c+dx)}{6d(a+ia \tan(c+dx))^3} \cdot 6a^2$$

input `Int[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `((A + I*B)*Cot[c + d*x]^2)/(6*d*(a + I*a*Tan[c + d*x])^3) + ((a*(13*A + (7*I)*B)*Cot[c + d*x]^2)/(4*d*(a + I*a*Tan[c + d*x])^2) + ((3*(5*a^3*((11*I)*A - 5*B)*x + (5*a^3*((11*I)*A - 5*B)*Cot[c + d*x])/d - (4*a^3*(7*A + (3*I)*B)*Cot[c + d*x]^2)/d - (8*a^3*(7*A + (3*I)*B)*Log[-Sin[c + d*x]])/d))/(2*a^2) + (5*a^2*(11*A + (5*I)*B)*Cot[c + d*x]^2)/(2*d*(a + I*a*Tan[c + d*x])))/(2*a^2))/(6*a^2)`

3.58. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

3.58.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`
- rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`
- rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.58.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.14

method	result
risch	$-\frac{49xB}{8a^3} + \frac{111ixA}{8a^3} - \frac{23ie^{-2i(dx+c)}B}{16a^3d} - \frac{39e^{-2i(dx+c)}A}{16a^3d} - \frac{7ie^{-4i(dx+c)}B}{32a^3d} - \frac{9e^{-4i(dx+c)}A}{32a^3d} - \frac{ie^{-6i(dx+c)}B}{48a^3d}$
derivativedivides	$-\frac{A(\cot^2(dx+c))}{2a^3d} + \frac{9iB}{8a^3d(i+\cot(dx+c))^2} - \frac{B\cot(dx+c)}{a^3d} + \frac{7A\ln(\cot^2(dx+c)+1)}{2a^3d} - \frac{iA}{6a^3d(i+\cot(dx+c))^3} + \frac{3}{3}$
default	$-\frac{A(\cot^2(dx+c))}{2a^3d} + \frac{9iB}{8a^3d(i+\cot(dx+c))^2} - \frac{B\cot(dx+c)}{a^3d} + \frac{7A\ln(\cot^2(dx+c)+1)}{2a^3d} - \frac{iA}{6a^3d(i+\cot(dx+c))^3} + \frac{3}{3}$

input `int(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output
$$-\frac{49}{8} \frac{x}{a^3} B + \frac{111}{8} \frac{I x}{a^3} A - \frac{23}{16} \frac{I}{a^3} \frac{1}{d} \exp(-2I(d*x+c)) B - \frac{39}{16} \frac{1}{a^3} \frac{1}{d} \exp(-2I(d*x+c)) A - \frac{7}{32} \frac{I}{a^3} \frac{1}{d} \exp(-4I(d*x+c)) B - \frac{9}{32} \frac{1}{a^3} \frac{1}{d} \exp(-4I(d*x+c)) A - \frac{1}{48} \frac{I}{a^3} \frac{1}{d} \exp(-6I(d*x+c)) B - \frac{1}{48} \frac{1}{a^3} \frac{1}{d} \exp(-6I(d*x+c)) A - \frac{6B}{a^3} \frac{1}{d} c + \frac{14I}{a^3} \frac{1}{d} A c - 2I(-2IA \exp(2I(d*x+c)) + B \exp(2I(d*x+c))) + 3IA - B / a^3 d / (\exp(2I(d*x+c)) - 1)^2 - 3I/a^3 d \ln(\exp(2I(d*x+c)) - 1) * B - 7A/a^3 d \ln(\exp(2I(d*x+c)) - 1)$$

3.58.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.09

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \frac{12(-111iA+49B)dx e^{(10i dx+10i c)} + 6(4(111iA-49B)dx + 103A + 55iB)e^{(8i dx+8i c)} + 3(4(-111i$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="fracas")`

```
output -1/96*(12*(-111*I*A + 49*B)*d*x*e^(10*I*d*x + 10*I*c) + 6*(4*(111*I*A - 49
*B)*d*x + 103*A + 55*I*B)*e^(8*I*d*x + 8*I*c) + 3*(4*(-111*I*A + 49*B)*d*x
- 339*A - 149*I*B)*e^(6*I*d*x + 6*I*c) + 14*(13*A + 7*I*B)*e^(4*I*d*x + 4
*I*c) + (23*A + 17*I*B)*e^(2*I*d*x + 2*I*c) + 96*((7*A + 3*I*B)*e^(10*I*d*
x + 10*I*c) - 2*(7*A + 3*I*B)*e^(8*I*d*x + 8*I*c) + (7*A + 3*I*B)*e^(6*I*d
*x + 6*I*c))*log(e^(2*I*d*x + 2*I*c) - 1) + 2*A + 2*I*B)/(a^3*d*e^(10*I*d*
x + 10*I*c) - 2*a^3*d*e^(8*I*d*x + 8*I*c) + a^3*d*e^(6*I*d*x + 6*I*c))
```

3.58.6 Sympy [A] (verification not implemented)

Time = 0.92 (sec) , antiderivative size = 398, normalized size of antiderivative = 1.84

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \frac{6A+2iB+(-4Ae^{2ic}-2iBe^{2ic})e^{2idx}}{a^3de^{4ic}e^{4idx}-2a^3de^{2ic}e^{2idx}+a^3d} + \left\{ \frac{((-512Aa^6d^2e^{6ic}-512iBa^6d^2e^{6ic})e^{-6idx}+(-6912Aa^6d^2e^{8ic}-5376iBa^6d^2e^{8ic})e^{-4idx}+(-59904Aa^6d^2e^{10ic}-35328iBa^6d^2e^{10ic})e^{-2idx})}{24576a^9d^3} \right. \\ \left. + x \left(-\frac{111iA-49B}{8a^3} + \frac{(111iAe^{6ic}+39iAe^{4ic}+9iAe^{2ic}+iA-49Be^{6ic}-23Be^{4ic}-7Be^{2ic}-B)e^{-6ic}}{8a^3} \right) \right. \\ \left. + \frac{x(111iA-49B)}{8a^3} - \frac{(7A+3iB) \log(e^{2idx}-e^{-2ic})}{a^3d} \right.$$

```
input integrate(cot(d*x+c)**3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)
```

```
output (6*A + 2*I*B + (-4*A*exp(2*I*c) - 2*I*B*exp(2*I*c))*exp(2*I*d*x))/(a**3*d*
exp(4*I*c)*exp(4*I*d*x) - 2*a**3*d*exp(2*I*c)*exp(2*I*d*x) + a**3*d) + Pie
cewise(((((-512*A*a**6*d**2*exp(6*I*c) - 512*I*B*a**6*d**2*exp(6*I*c))*exp(
-6*I*d*x) + (-6912*A*a**6*d**2*exp(8*I*c) - 5376*I*B*a**6*d**2*exp(8*I*c))
*exp(-4*I*d*x) + (-59904*A*a**6*d**2*exp(10*I*c) - 35328*I*B*a**6*d**2*exp
(10*I*c))*exp(-2*I*d*x))*exp(-12*I*c)/(24576*a**9*d**3), Ne(a**9*d**3*exp(
12*I*c), 0)), (x*(-(111*I*A - 49*B)/(8*a**3) + (111*I*A*exp(6*I*c) + 39*I*
A*exp(4*I*c) + 9*I*A*exp(2*I*c) + I*A - 49*B*exp(6*I*c) - 23*B*exp(4*I*c)
- 7*B*exp(2*I*c) - B)*exp(-6*I*c)/(8*a**3)), True)) + x*(111*I*A - 49*B)/(
8*a**3) - (7*A + 3*I*B)*log(exp(2*I*d*x) - exp(-2*I*c))/(a**3*d)
```

3.58.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm=
"maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.58.8 Giac [A] (verification not implemented)

Time = 1.30 (sec) , antiderivative size = 207, normalized size of antiderivative = 0.96

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$= \frac{6(A-iB)\log(\tan(dx+c)+i)}{a^3} + \frac{6(111A+49iB)\log(\tan(dx+c)-i)}{a^3} - \frac{96(7A+3iB)\log(\tan(dx+c))}{a^3} + \frac{48(21A\tan(dx+c)^2+9iB\tan(dx+c))}{a^3 \tan(dx+c)}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm=
"giac")
```

```
output 1/96*(6*(A - I*B)*log(tan(d*x + c) + I)/a^3 + 6*(111*A + 49*I*B)*log(tan(d
*x + c) - I)/a^3 - 96*(7*A + 3*I*B)*log(tan(d*x + c))/a^3 + 48*(21*A*tan(d
*x + c)^2 + 9*I*B*tan(d*x + c)^2 + 6*I*A*tan(d*x + c) - 2*B*tan(d*x + c) -
A)/(a^3*tan(d*x + c)^2) + (1221*I*A*tan(d*x + c)^3 - 539*B*tan(d*x + c)^3
+ 4035*A*tan(d*x + c)^2 + 1821*I*B*tan(d*x + c)^2 - 4491*I*A*tan(d*x + c)
+ 2085*B*tan(d*x + c) - 1693*A - 819*I*B)/(a^3*(I*tan(d*x + c) + 1)^3))/d
```

3.58. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

3.58.9 Mupad [B] (verification not implemented)

Time = 8.25 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.02

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx =$$

$$-\frac{\frac{A}{2a^3} + \tan(c+dx)^3 \left(-\frac{63B}{8a^3} + \frac{A137i}{8a^3}\right) + \tan(c+dx)^2 \left(\frac{149A}{12a^3} + \frac{B71i}{12a^3}\right) - \tan(c+dx)^4 \left(\frac{55A}{8a^3} + \frac{B25i}{8a^3}\right) - \tan(c+dx)^5 \left(\frac{11A}{8a^3} + \frac{B7i}{8a^3}\right)}{d \left(-\tan(c+dx)^5 1i - 3 \tan(c+dx)^4 + \tan(c+dx)^3 3i + \tan(c+dx)^2\right)}$$

$$-\frac{\ln(\tan(c+dx)) (7A+B3i)}{a^3 d} + \frac{\ln(\tan(c+dx)+1i) (A-B1i)}{16a^3 d}$$

$$+\frac{\ln(\tan(c+dx)-i) (111A+B49i)}{16a^3 d}$$

```
input int((cot(c + d*x)^3*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3,x)
```

```
output (log(tan(c + d*x) + 1i)*(A - B*1i))/(16*a^3*d) - (log(tan(c + d*x))*(7*A +
B*3i))/(a^3*d) - (tan(c + d*x)^3*((A*137i)/(8*a^3) - (63*B)/(8*a^3)) - ta
n(c + d*x)^4*((55*A)/(8*a^3) + (B*25i)/(8*a^3)) + tan(c + d*x)^2*((149*A)/
(12*a^3) + (B*71i)/(12*a^3)) + A/(2*a^3) - tan(c + d*x)*((A*3i)/(2*a^3) -
B/a^3))/(d*(tan(c + d*x)^2 + tan(c + d*x)^3*3i - 3*tan(c + d*x)^4 - tan(c
+ d*x)^5*1i)) + (log(tan(c + d*x) - 1i)*(111*A + B*49i))/(16*a^3*d)
```

3.59 $\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

3.59.1	Optimal result	742
3.59.2	Mathematica [A] (verified)	743
3.59.3	Rubi [A] (verified)	743
3.59.4	Maple [A] (verified)	747
3.59.5	Fricas [A] (verification not implemented)	748
3.59.6	Sympy [A] (verification not implemented)	748
3.59.7	Maxima [F(-2)]	749
3.59.8	Giac [A] (verification not implemented)	749
3.59.9	Mupad [B] (verification not implemented)	750

3.59.1 Optimal result

Integrand size = 34, antiderivative size = 185

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \frac{(A+15iB)x}{16a^4} - \frac{B \log(\cos(c+dx))}{a^4d} - \frac{iA-15B}{16a^4d(1+i \tan(c+dx))} - \frac{(iA-7B) \tan^2(c+dx)}{16a^4d(1+i \tan(c+dx))^2} + \frac{(iA-B) \tan^4(c+dx)}{8d(a+ia \tan(c+dx))^4} + \frac{(A+3iB) \tan^3(c+dx)}{12ad(a+ia \tan(c+dx))^3}$$

output

```
1/16*(A+15*I*B)*x/a^4-B*ln(cos(d*x+c))/a^4/d+1/16*(-I*A+15*B)/a^4/d/(1+I*tan(d*x+c))-1/16*(I*A-7*B)*tan(d*x+c)^2/a^4/d/(1+I*tan(d*x+c))^2+1/8*(I*A-B)*tan(d*x+c)^4/d/(a+I*a*tan(d*x+c))^4+1/12*(A+3*I*B)*tan(d*x+c)^3/a/d/(a+I*a*tan(d*x+c))^3
```

3.59.2 Mathematica [A] (verified)

Time = 1.35 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.43

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$$

$$= \frac{\sec^4(c+dx)(18iA-48B+8(-4iA+21B)\cos(2(c+dx))+2\cos(4(c+dx))(7iA-60B+(-3iA+93B)\cos(2(c+dx))))}{(a+ia\tan(c+dx))^4}$$

input `Integrate[(Tan[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]`

output `(Sec[c + d*x]^4*((18*I)*A - 48*B + 8*((-4*I)*A + 21*B)*Cos[2*(c + d*x)] + 2*Cos[4*(c + d*x)]*((7*I)*A - 60*B + ((-3*I)*A + 93*B)*Log[I - Tan[c + d*x]]) + 3*(I*A + B)*Log[I + Tan[c + d*x]]) + 16*A*Sin[2*(c + d*x)] + (144*I)*B*Sin[2*(c + d*x)] - 11*A*Sin[4*(c + d*x)] - (117*I)*B*Sin[4*(c + d*x)] + 6*A*Log[I - Tan[c + d*x]]*Sin[4*(c + d*x)] + (186*I)*B*Log[I - Tan[c + d*x]]*Sin[4*(c + d*x)] - 6*A*Log[I + Tan[c + d*x]]*Sin[4*(c + d*x)] + (6*I)*B*Log[I + Tan[c + d*x]]*Sin[4*(c + d*x)])/(192*a^4*d*(-I + Tan[c + d*x])^4)`

3.59.3 Rubi [A] (verified)

Time = 1.17 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.10, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4078, 27, 3042, 4078, 27, 3042, 4078, 27, 3042, 4072, 25, 27, 3042, 3956, 4009, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)^4(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$$

$$\downarrow \text{4078}$$

$$\frac{(-B+iA)\tan^4(c+dx)}{8d(a+ia\tan(c+dx))^4} - \frac{\int \frac{4\tan^3(c+dx)(a(iA-B)+2iaB\tan(c+dx))}{(i\tan(c+dx)a+a)^3} dx}{8a^2}$$

3.59. $\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{\int \frac{\tan^3(c+dx)(a(iA-B)+2iaB \tan(c+dx))}{(i \tan(c+dx)a+a)^3} dx}{2a^2} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{\int \frac{\tan(c+dx)^3(a(iA-B)+2iaB \tan(c+dx))}{(i \tan(c+dx)a+a)^3} dx}{2a^2} \\
& \downarrow 4078 \\
& \frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{\int -\frac{3 \tan^2(c+dx)((A+3iB)a^2+4B \tan(c+dx)a^2)}{(i \tan(c+dx)a+a)^2} dx}{2a^2} - \frac{a(A+3iB) \tan^3(c+dx)}{6d(a+ia \tan(c+dx))^3} \\
& \downarrow 27 \\
& \frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{\int \frac{\tan^2(c+dx)((A+3iB)a^2+4B \tan(c+dx)a^2)}{(i \tan(c+dx)a+a)^2} dx}{2a^2} - \frac{a(A+3iB) \tan^3(c+dx)}{6d(a+ia \tan(c+dx))^3} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{\int \frac{\tan(c+dx)^2((A+3iB)a^2+4B \tan(c+dx)a^2)}{(i \tan(c+dx)a+a)^2} dx}{2a^2} - \frac{a(A+3iB) \tan^3(c+dx)}{6d(a+ia \tan(c+dx))^3} \\
& \downarrow 4078 \\
& \frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{\int \frac{2 \tan(c+dx)((iA-7B)a^3+8iB \tan(c+dx)a^3)}{i \tan(c+dx)a+a} dx}{2a^2} - \frac{a(A+3iB) \tan^3(c+dx)}{6d(a+ia \tan(c+dx))^3} \\
& \downarrow 27 \\
& \frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{\int \frac{\tan(c+dx)((iA-7B)a^3+8iB \tan(c+dx)a^3)}{i \tan(c+dx)a+a} dx}{2a^2} - \frac{a(A+3iB) \tan^3(c+dx)}{6d(a+ia \tan(c+dx))^3} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{\int \frac{\tan(c+dx)((iA-7B)a^3+8iB \tan(c+dx)a^3)}{i \tan(c+dx)a+a} dx}{2a^2} - \frac{a(A+3iB) \tan^3(c+dx)}{6d(a+ia \tan(c+dx))^3}
\end{aligned}$$

3.59. $\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

$$\begin{aligned}
& \frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{(-7B + iA) \tan^2(c + dx)}{4d(1 + i \tan(c + dx))^2} - \frac{\int \frac{\tan(c + dx) ((iA - 7B)a^3 + 8iB \tan(c + dx)a^3)}{i \tan(c + dx)a + a} dx}{2a^2} - \frac{a(A + 3iB) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} \\
& \frac{2a^2}{2a^2} \\
& \quad \downarrow 4072 \\
& \frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{(-7B + iA) \tan^2(c + dx)}{4d(1 + i \tan(c + dx))^2} - \frac{8a^2 B \int \tan(c + dx) dx - \frac{i \int \frac{a^4(A + 15iB) \tan(c + dx)}{i \tan(c + dx)a + a} dx}{a}}{2a^2} - \frac{a(A + 3iB) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} \\
& \frac{2a^2}{2a^2} \\
& \quad \downarrow 25 \\
& \frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{(-7B + iA) \tan^2(c + dx)}{4d(1 + i \tan(c + dx))^2} - \frac{8a^2 B \int \tan(c + dx) dx + \frac{i \int \frac{a^4(A + 15iB) \tan(c + dx)}{i \tan(c + dx)a + a} dx}{a}}{2a^2} - \frac{a(A + 3iB) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} \\
& \frac{2a^2}{2a^2} \\
& \quad \downarrow 27 \\
& \frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{(-7B + iA) \tan^2(c + dx)}{4d(1 + i \tan(c + dx))^2} - \frac{8a^2 B \int \tan(c + dx) dx + ia^3(A + 15iB) \int \frac{\tan(c + dx)}{i \tan(c + dx)a + a} dx}{2a^2} - \frac{a(A + 3iB) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} \\
& \frac{2a^2}{2a^2} \\
& \quad \downarrow 3042 \\
& \frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{(-7B + iA) \tan^2(c + dx)}{4d(1 + i \tan(c + dx))^2} - \frac{8a^2 B \int \tan(c + dx) dx + ia^3(A + 15iB) \int \frac{\tan(c + dx)}{i \tan(c + dx)a + a} dx}{2a^2} - \frac{a(A + 3iB) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} \\
& \frac{2a^2}{2a^2} \\
& \quad \downarrow 3956 \\
& \frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{(-7B + iA) \tan^2(c + dx)}{4d(1 + i \tan(c + dx))^2} - \frac{8a^2 B \log(\cos(c + dx))}{d} + ia^3(A + 15iB) \int \frac{\tan(c + dx)}{i \tan(c + dx)a + a} dx}{2a^2} - \frac{a(A + 3iB) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3} \\
& \frac{2a^2}{2a^2} \\
& \quad \downarrow 4009
\end{aligned}$$

3.59. $\int \frac{\tan^4(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^4} dx$

$$\frac{\frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{(-7B + iA) \tan^2(c + dx)}{4d(1 + i \tan(c + dx))^2} - \frac{8a^2 B \log(\cos(c + dx))}{d} + ia^3(A + 15iB) \left(-\frac{i \int 1 dx}{2a} - \frac{1}{2d(a + ia \tan(c + dx))} \right)}{2a^2} - \frac{a(A + 3iB) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3}}{2a^2}$$

↓ 24

$$\frac{\frac{(-B + iA) \tan^4(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{(-7B + iA) \tan^2(c + dx)}{4d(1 + i \tan(c + dx))^2} - \frac{8a^2 B \log(\cos(c + dx))}{d} + ia^3(A + 15iB) \left(-\frac{1}{2d(a + ia \tan(c + dx))} - \frac{ix}{2a} \right)}{2a^2} - \frac{a(A + 3iB) \tan^3(c + dx)}{6d(a + ia \tan(c + dx))^3}}{2a^2}$$

```
input Int[(Tan[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]
```

```
output ((I*A - B)*Tan[c + d*x]^4)/(8*d*(a + I*a*Tan[c + d*x])^4) - (-1/6*(a*(A + (3*I)*B)*Tan[c + d*x]^3)/(d*(a + I*a*Tan[c + d*x])^3) + (((I*A - 7*B)*Tan[c + d*x]^2)/(4*d*(1 + I*Tan[c + d*x])^2) - ((-8*a^2*B*Log[Cos[c + d*x]])/d + I*a^3*(A + (15*I)*B)*((( -1/2*I)*x)/a - 1/(2*d*(a + I*a*Tan[c + d*x])))))/(2*a^2))/(2*a^2))/(2*a^2)
```

3.59.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

3.59. $\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

```
rule 4009 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a
*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x
], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2
, 0] && LtQ[m, 0]
```

```
rule 4072 Int((((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_
.)*(x_)]))/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*(d/
b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c
- a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d
, e, f, A, B}, x] && NeQ[b*c - a*d, 0]
```

```
rule 4078 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-(A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.59.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.06

method	result
risch	$\frac{31ixB}{16a^4} + \frac{xA}{16a^4} + \frac{13e^{-2i(dx+c)}B}{16da^4} - \frac{ie^{-2i(dx+c)}A}{8da^4} - \frac{e^{-4i(dx+c)}B}{4da^4} + \frac{3ie^{-4i(dx+c)}A}{32da^4} + \frac{e^{-6i(dx+c)}B}{16da^4} - \frac{ie^{-6i(dx+c)}A}{24da^4}$
derivativedivides	$-\frac{17iA}{16da^4(\tan(dx+c)-i)^2} + \frac{A \arctan(\tan(dx+c))}{16da^4} + \frac{B \ln(1+\tan^2(dx+c))}{2da^4} + \frac{15iB \arctan(\tan(dx+c))}{16da^4} - \frac{15iA \arctan(\tan(dx+c))}{16da^4}$
default	$-\frac{17iA}{16da^4(\tan(dx+c)-i)^2} + \frac{A \arctan(\tan(dx+c))}{16da^4} + \frac{B \ln(1+\tan^2(dx+c))}{2da^4} + \frac{15iB \arctan(\tan(dx+c))}{16da^4} - \frac{15iA \arctan(\tan(dx+c))}{16da^4}$

```
input int(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x,method=_RETURNVER
BOSE)
```

$$3.59. \int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$$

output $31/16*I*x/a^4*B+1/16*x/a^4*A+13/16/d/a^4*\exp(-2*I*(d*x+c))*B-1/8*I/d/a^4*\exp(-2*I*(d*x+c))*A-1/4/d/a^4*\exp(-4*I*(d*x+c))*B+3/32*I/d/a^4*\exp(-4*I*(d*x+c))*A+1/16/d/a^4*\exp(-6*I*(d*x+c))*B-1/24*I/d/a^4*\exp(-6*I*(d*x+c))*A-1/128/d/a^4*\exp(-8*I*(d*x+c))*B+1/128*I/d/a^4*\exp(-8*I*(d*x+c))*A+2*I*B/d/a^4*c-B/d/a^4*\ln(\exp(2*I*(d*x+c))+1)$

3.59.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.65

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx = \frac{(24(A+31iB)dx e^{(8i dx+8i c)} - 384 B e^{(8i dx+8i c)} \log(e^{(2i dx+2i c)} + 1) - 24(2i A - 13 B) e^{(6i dx+6i c)} - 12(-3i A + 8 B) e^{(4i dx+4i c)} - 8(2i A - 3 B) e^{(2i dx+2i c)} + 3i A - 3 B) e^{(-8i dx-8i c)}}{384 a^4 d}$$

input `integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="fricas")`

output $1/384*(24*(A + 31*I*B)*d*x*e^{(8*I*d*x + 8*I*c)} - 384*B*e^{(8*I*d*x + 8*I*c)}*\log(e^{(2*I*d*x + 2*I*c)} + 1) - 24*(2*I*A - 13*B)*e^{(6*I*d*x + 6*I*c)} - 12*(-3*I*A + 8*B)*e^{(4*I*d*x + 4*I*c)} - 8*(2*I*A - 3*B)*e^{(2*I*d*x + 2*I*c)} + 3*I*A - 3*B)*e^{(-8*I*d*x - 8*I*c)}/(a^4*d)$

3.59.6 Sympy [A] (verification not implemented)

Time = 1.15 (sec) , antiderivative size = 359, normalized size of antiderivative = 1.94

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx = -\frac{B \log(e^{2idx} + e^{-2ic})}{a^4 d} + \left\{ \frac{((24576iAa^{12}d^3e^{12ic} - 24576Ba^{12}d^3e^{12ic})e^{-8idx} + (-131072iAa^{12}d^3e^{14ic} + 196608Ba^{12}d^3e^{14ic})e^{-6idx} + (294912iAa^{12}d^3e^{16ic} - 786432Ba^{12}d^3e^{16ic})e^{-4idx} + (131072iAa^{12}d^3e^{18ic} - 3145728a^{16}d^4)e^{-2idx} + 3145728a^{16}d^4)}{3145728a^{16}d^4} \right. \\ \left. + x \left(-\frac{A+31iB}{16a^4} + \frac{(Ae^{8ic} - 4Ae^{6ic} + 6Ae^{4ic} - 4Ae^{2ic} + A + 31iBe^{8ic} - 26iBe^{6ic} + 16iBe^{4ic} - 6iBe^{2ic} + iB)e^{-8ic}}{16a^4} \right) \right. \\ \left. + \frac{x(A+31iB)}{16a^4} \right.$$

input `integrate(tan(d*x+c)**4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**4,x)`

3.59. $\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$

```
output -B*log(exp(2*I*d*x) + exp(-2*I*c))/(a**4*d) + Piecewise((((24576*I*A*a**12
*d**3*exp(12*I*c) - 24576*B*a**12*d**3*exp(12*I*c))*exp(-8*I*d*x) + (-1310
72*I*A*a**12*d**3*exp(14*I*c) + 196608*B*a**12*d**3*exp(14*I*c))*exp(-6*I*
d*x) + (294912*I*A*a**12*d**3*exp(16*I*c) - 786432*B*a**12*d**3*exp(16*I*c
))*exp(-4*I*d*x) + (-393216*I*A*a**12*d**3*exp(18*I*c) + 2555904*B*a**12*d
**3*exp(18*I*c))*exp(-2*I*d*x))*exp(-20*I*c)/(3145728*a**16*d**4), Ne(a**1
6*d**4*exp(20*I*c), 0)), (x*(-(A + 31*I*B)/(16*a**4) + (A*exp(8*I*c) - 4*A
*exp(6*I*c) + 6*A*exp(4*I*c) - 4*A*exp(2*I*c) + A + 31*I*B*exp(8*I*c) - 26
*I*B*exp(6*I*c) + 16*I*B*exp(4*I*c) - 6*I*B*exp(2*I*c) + I*B)*exp(-8*I*c)/
(16*a**4)), True)) + x*(A + 31*I*B)/(16*a**4)
```

3.59.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^4(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^4} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm=
"maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.59.8 Giac [A] (verification not implemented)

Time = 1.20 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.83

$$\int \frac{\tan^4(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^4} dx =$$

$$\frac{12(-iA-B)\log(\tan(dx+c)+i)}{a^4} - \frac{12(-iA+31B)\log(\tan(dx+c)-i)}{a^4} - \frac{25iA\tan(dx+c)^4 - 775B\tan(dx+c)^4 - 260A\tan(dx+c)^3 + 1924iB\tan(dx+c)^3}{a^4}$$

384 d

```
input integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm=
"giac")
```

output
$$\frac{-1/384*(12*(-I*A - B)*\log(\tan(dx + c) + I)/a^4 - 12*(-I*A + 31*B)*\log(\tan(dx + c) - I)/a^4 - (25*I*A*\tan(dx + c)^4 - 775*B*\tan(dx + c)^4 - 260*A*\tan(dx + c)^3 + 1924*I*B*\tan(dx + c)^3 + 522*I*A*\tan(dx + c)^2 + 1866*B*\tan(dx + c)^2 + 388*A*\tan(dx + c) - 772*I*B*\tan(dx + c) - 103*I*A - 103*B)/(a^4*(\tan(dx + c) - I)^4))/d$$

3.59.9 Mupad [B] (verification not implemented)

Time = 8.09 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.96

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$$

$$= \frac{\tan(c+dx)^2 \left(-\frac{29B}{4a^4} + \frac{A7i}{4a^4}\right) - \tan(c+dx)^3 \left(\frac{15A}{16a^4} + \frac{B49i}{16a^4}\right) - \frac{A1i}{3a^4} + \frac{7B}{4a^4} + \tan(c+dx) \left(\frac{61A}{48a^4} + \frac{B97i}{16a^4}\right)}{d \left(\tan(c+dx)^4 - \tan(c+dx)^3 4i - 6 \tan(c+dx)^2 + \tan(c+dx) 4i + 1\right)}$$

$$+ \frac{\ln(\tan(c+dx) + 1i) (B + A 1i)}{32 a^4 d} - \frac{\ln(\tan(c+dx) - i) (A + B 31i) 1i}{32 a^4 d}$$

input `int((tan(c + d*x)^4*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^4,x)`

output
$$\frac{(\tan(c + d*x)^2*((A*7i)/(4*a^4) - (29*B)/(4*a^4)) - \tan(c + d*x)^3*((15*A)/(16*a^4) + (B*49i)/(16*a^4)) - (A*1i)/(3*a^4) + (7*B)/(4*a^4) + \tan(c + d*x)*((61*A)/(48*a^4) + (B*97i)/(16*a^4)))/(d*(\tan(c + d*x)*4i - 6*\tan(c + d*x)^2 - \tan(c + d*x)^3*4i + \tan(c + d*x)^4 + 1)) + (\log(\tan(c + d*x) + 1i)*(A*1i + B))/(32*a^4*d) - (\log(\tan(c + d*x) - 1i)*(A + B*31i)*1i)/(32*a^4*d)}$$

3.60 $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

3.60.1 Optimal result 751
 3.60.2 Mathematica [A] (verified) 751
 3.60.3 Rubi [A] (verified) 752
 3.60.4 Maple [A] (verified) 755
 3.60.5 Fricas [A] (verification not implemented) 756
 3.60.6 Sympy [A] (verification not implemented) 756
 3.60.7 Maxima [F(-2)] 757
 3.60.8 Giac [A] (verification not implemented) 757
 3.60.9 Mupad [B] (verification not implemented) 758

3.60.1 Optimal result

Integrand size = 34, antiderivative size = 159

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \frac{(iA+B)x}{16a^4} - \frac{A-13iB}{48a^4d(1+i \tan(c+dx))^2} + \frac{5A-29iB}{48a^4d(1+i \tan(c+dx))} + \frac{(iA-B) \tan^3(c+dx)}{8d(a+ia \tan(c+dx))^4} + \frac{(A+5iB) \tan^2(c+dx)}{24ad(a+ia \tan(c+dx))^3}$$

```
output 1/16*(I*A+B)*x/a^4+1/48*(-A+13*I*B)/a^4/d/(1+I*tan(d*x+c))^2+1/48*(5*A-29*I*B)/a^4/d/(1+I*tan(d*x+c))+1/8*(I*A-B)*tan(d*x+c)^3/d/(a+I*a*tan(d*x+c))^4+1/24*(A+5*I*B)*tan(d*x+c)^2/a/d/(a+I*a*tan(d*x+c))^3
```

3.60.2 Mathematica [A] (verified)

Time = 1.69 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.99

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \frac{\sec^4(c+dx)(36iB+16(A-4iB) \cos(2(c+dx))+3(A+iB+8iAdx+8Bdx) \cos(4(c+dx))+32iA \sin(4(c+dx)))}{384a^4d}$$

input `Integrate[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x
]`

output `(Sec[c + d*x]^4*((36*I)*B + 16*(A - (4*I)*B)*Cos[2*(c + d*x)] + 3*(A + I*B
+ (8*I)*A*d*x + 8*B*d*x)*Cos[4*(c + d*x)] + (32*I)*A*Sin[2*(c + d*x)] + 3
2*B*Sin[2*(c + d*x)] - (3*I)*A*Sin[4*(c + d*x)] + 3*B*Sin[4*(c + d*x)] - 2
4*A*d*x*Sin[4*(c + d*x)] + (24*I)*B*d*x*Sin[4*(c + d*x)]))/(384*a^4*d*(-I
+ Tan[c + d*x])^4)`

3.60.3 Rubi [A] (verified)

Time = 0.91 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.14, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.324$, Rules used = {3042, 4078, 3042, 4078, 27, 3042, 4073, 25, 3042, 4009, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^3(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+iA) \tan^3(c+dx)}{8d(a+ia \tan(c+dx))^4} - \frac{\int \frac{\tan^2(c+dx)(3a(iA-B)-a(A-7iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^3} dx}{8a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B+iA) \tan^3(c+dx)}{8d(a+ia \tan(c+dx))^4} - \frac{\int \frac{\tan(c+dx)^2(3a(iA-B)-a(A-7iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^3} dx}{8a^2} \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+iA) \tan^3(c+dx)}{8d(a+ia \tan(c+dx))^4} - \frac{\int -\frac{4 \tan(c+dx)((A+5iB)a^2+2(iA+4B) \tan(c+dx)a^2)}{(i \tan(c+dx)a+a)^2} dx}{6a^2} - \frac{a(A+5iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^3} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.60. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

$$\frac{(-B + iA) \tan^3(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{2 \int \frac{\tan(c+dx)((A+5iB)a^2+2(iA+4B) \tan(c+dx)a^2)}{(i \tan(c+dx)a+a)^2} dx}{3a^2} - \frac{a(A+5iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^3}$$

↓ 3042

$$\frac{(-B + iA) \tan^3(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{2 \int \frac{\tan(c+dx)((A+5iB)a^2+2(iA+4B) \tan(c+dx)a^2)}{(i \tan(c+dx)a+a)^2} dx}{3a^2} - \frac{a(A+5iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^3}$$

↓ 4073

$$\frac{(-B + iA) \tan^3(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{2 \left(\frac{A-13iB}{4d(1+i \tan(c+dx))^2} - \frac{i \int \frac{a^3(A-13iB)-4a^3(iA+4B) \tan(c+dx)}{i \tan(c+dx)a+a} dx}{2a^2} \right)}{3a^2} - \frac{a(A+5iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^3}$$

↓ 25

$$\frac{(-B + iA) \tan^3(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{2 \left(\frac{i \int \frac{a^3(A-13iB)-4a^3(iA+4B) \tan(c+dx)}{i \tan(c+dx)a+a} dx}{2a^2} + \frac{A-13iB}{4d(1+i \tan(c+dx))^2} \right)}{3a^2} - \frac{a(A+5iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^3}$$

↓ 3042

$$\frac{(-B + iA) \tan^3(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{2 \left(\frac{i \int \frac{a^3(A-13iB)-4a^3(iA+4B) \tan(c+dx)}{i \tan(c+dx)a+a} dx}{2a^2} + \frac{A-13iB}{4d(1+i \tan(c+dx))^2} \right)}{3a^2} - \frac{a(A+5iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^3}$$

↓ 4009

$$\frac{(-B + iA) \tan^3(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{2 \left(\frac{i \left(\frac{a^3(29B+5iA)}{2d(a+ia \tan(c+dx))} - \frac{3}{2}a^2(A-iB) \int 1 dx \right)}{2a^2} + \frac{A-13iB}{4d(1+i \tan(c+dx))^2} \right)}{3a^2} - \frac{a(A+5iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^3}$$

↓ 24

3.60. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

$$\frac{(-B + iA) \tan^3(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{2 \left(\frac{i \left(\frac{a^3(29B + 5iA)}{2d(a + ia \tan(c + dx))} - \frac{3}{2} a^2 x(A - iB) \right)}{2a^2} + \frac{A - 13iB}{4d(1 + i \tan(c + dx))^2} \right)}{3a^2} - \frac{a(A + 5iB) \tan^2(c + dx)}{3d(a + ia \tan(c + dx))^3}$$

input `Int[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]`

output `((I*A - B)*Tan[c + d*x]^3)/(8*d*(a + I*a*Tan[c + d*x])^4) - (-1/3*(a*(A + (5*I)*B)*Tan[c + d*x]^2)/(d*(a + I*a*Tan[c + d*x])^3) + (2*((A - (13*I)*B)/(4*d*(1 + I*Tan[c + d*x])^2) + ((I/2)*((-3*a^2*(A - I*B)*x)/2 + (a^3*((5*I)*A + 29*B))/(2*d*(a + I*a*Tan[c + d*x]))))/a^2)/(3*a^2)/(8*a^2)`

3.60.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

```
rule 4073 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-
A*b - a*B)*(a*c + b*d)*((a + b*Tan[e + f*x])^m/(2*a^2*f*m)), x] + Simp[1/(
2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[A*b*c + a*B*c + a*A*d + b*B*
d + 2*a*B*d*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && LtQ[m, -1] && EqQ[a^2 + b^2, 0]
```

```
rule 4078 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Sim
p[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.60.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.92

method	result
risch	$\frac{x B}{16 a^4} + \frac{i x A}{16 a^4} - \frac{i e^{-2 i(d x+c)} B}{8 d a^4} + \frac{e^{-2 i(d x+c)} A}{16 d a^4} + \frac{3 i B e^{-4 i(d x+c)}}{32 d a^4} - \frac{i e^{-6 i(d x+c)} B}{24 d a^4} - \frac{e^{-6 i(d x+c)} A}{48 d a^4} + \frac{i e^{-8 i(d x+c)}}{128 d a^4}$
derivativedivides	$-\frac{17 i B}{16 d a^4(\tan(dx+c)-i)^2} + \frac{i A}{16 d a^4(\tan(dx+c)-i)} + \frac{B \arctan(\tan(dx+c))}{16 d a^4} + \frac{A}{8 d a^4(\tan(dx+c)-i)^4} + \frac{i}{8 d a^4(\tan(dx+c)-i)}$
default	$-\frac{17 i B}{16 d a^4(\tan(dx+c)-i)^2} + \frac{i A}{16 d a^4(\tan(dx+c)-i)} + \frac{B \arctan(\tan(dx+c))}{16 d a^4} + \frac{A}{8 d a^4(\tan(dx+c)-i)^4} + \frac{i}{8 d a^4(\tan(dx+c)-i)}$

```
input int(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x,method=_RETURNVER
BOSE)
```

```
output 1/16*x/a^4*B+1/16*I*x/a^4*A-1/8*I/d/a^4*exp(-2*I*(d*x+c))*B+1/16/d/a^4*exp
(-2*I*(d*x+c))*A+3/32*I*B/d/a^4*exp(-4*I*(d*x+c))-1/24*I/d/a^4*exp(-6*I*(d
*x+c))*B-1/48/d/a^4*exp(-6*I*(d*x+c))*A+1/128*I/d/a^4*exp(-8*I*(d*x+c))*B+
1/128/d/a^4*exp(-8*I*(d*x+c))*A
```

$$3.60. \int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$$

3.60.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.55

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \frac{(24(-iA-B)dx e^{(8i dx+8i c)} - 24(A-2iB)e^{(6i dx+6i c)} - 36i B e^{(4i dx+4i c)} + 8(A+2iB)e^{(2i dx+2i c)} - 3A - 3iB)e^{(-8i dx-8i c)}}{384 a^4 d}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="fracas")`

output `-1/384*(24*(-I*A - B)*d*x*e^(8*I*d*x + 8*I*c) - 24*(A - 2*I*B)*e^(6*I*d*x + 6*I*c) - 36*I*B*e^(4*I*d*x + 4*I*c) + 8*(A + 2*I*B)*e^(2*I*d*x + 2*I*c) - 3*A - 3*I*B)*e^(-8*I*d*x - 8*I*c)/(a^4*d)`

3.60.6 Sympy [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 301, normalized size of antiderivative = 1.89

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \left\{ \begin{array}{l} \frac{(294912iBa^{12}d^3e^{16ic}e^{-4idx} + (24576Aa^{12}d^3e^{12ic} + 24576iBa^{12}d^3e^{12ic})e^{-8idx} + (-65536Aa^{12}d^3e^{14ic} - 131072iBa^{12}d^3e^{14ic})e^{-6idx} + (196608 - 3145728a^{16}d^4))e^{-8ic}}{16a^4} \\ x \left(-\frac{iA+B}{16a^4} + \frac{(iAe^{8ic} - 2iAe^{6ic} + 2iAe^{2ic} - iA + Be^{8ic} - 4Be^{6ic} + 6Be^{4ic} - 4Be^{2ic} + B)e^{-8ic}}{16a^4} \right) \\ + \frac{x(iA+B)}{16a^4} \end{array} \right.$$

input `integrate(tan(d*x+c)**3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**4,x)`

output `Piecewise(((294912*I*B*a**12*d**3*exp(16*I*c)*exp(-4*I*d*x) + (24576*A*a**12*d**3*exp(12*I*c) + 24576*I*B*a**12*d**3*exp(12*I*c))*exp(-8*I*d*x) + (-65536*A*a**12*d**3*exp(14*I*c) - 131072*I*B*a**12*d**3*exp(14*I*c))*exp(-6*I*d*x) + (196608*A*a**12*d**3*exp(18*I*c) - 393216*I*B*a**12*d**3*exp(18*I*c))*exp(-2*I*d*x))*exp(-20*I*c)/(3145728*a**16*d**4), Ne(a**16*d**4*exp(20*I*c), 0)), (x*(-(I*A + B)/(16*a**4) + (I*A*exp(8*I*c) - 2*I*A*exp(6*I*c) + 2*I*A*exp(2*I*c) - I*A + B*exp(8*I*c) - 4*B*exp(6*I*c) + 6*B*exp(4*I*c) - 4*B*exp(2*I*c) + B)*exp(-8*I*c)/(16*a**4)), True)) + x*(I*A + B)/(16*a**4)`

3.60. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

3.60.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm=
"maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.60.8 Giac [A] (verification not implemented)

Time = 0.92 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.94

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx =$$

$$\frac{12(A-iB)\log(\tan(dx+c)+i)}{a^4} - \frac{12(A-iB)\log(\tan(dx+c)-i)}{a^4} + \frac{25A\tan(dx+c)^4 - 25iB\tan(dx+c)^4 - 124iA\tan(dx+c)^3 + 260B\tan(dx+c)^3 - 124A\tan(dx+c)^2 + 260iB\tan(dx+c)^2 - 4iA\tan(dx+c) - 388B\tan(dx+c) - 7A + 103iB}{384d}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm=
"giac")
```

```
output -1/384*(12*(A - I*B)*log(tan(d*x + c) + I)/a^4 - 12*(A - I*B)*log(tan(d*x
+ c) - I)/a^4 + (25*A*tan(d*x + c)^4 - 25*I*B*tan(d*x + c)^4 - 124*I*A*tan
(d*x + c)^3 + 260*B*tan(d*x + c)^3 - 54*A*tan(d*x + c)^2 - 522*I*B*tan(d*x
+ c)^2 - 4*I*A*tan(d*x + c) - 388*B*tan(d*x + c) - 7*A + 103*I*B)/(a^4*(t
an(d*x + c) - I)^4))/d
```

3.60.9 Mupad [B] (verification not implemented)

Time = 7.39 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.12

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$$

$$= \frac{\frac{A}{12a^4} + \tan(c+dx)^3 \left(-\frac{15B}{16a^4} + \frac{A1i}{16a^4}\right) - \tan(c+dx)^2 \left(\frac{A}{4a^4} - \frac{B7i}{4a^4}\right) - \frac{B1i}{3a^4} + \tan(c+dx) \left(\frac{61B}{48a^4} + \frac{A13i}{48a^4}\right)}{d(\tan(c+dx)^4 - \tan(c+dx)^3 4i - 6\tan(c+dx)^2 + \tan(c+dx) 4i + 1)}$$

$$+ \frac{\ln(\tan(c+dx) - i)(A - B1i)}{32a^4 d} + \frac{\ln(\tan(c+dx) + i)(B + A1i) 1i}{32a^4 d}$$

input `int((tan(c + d*x)^3*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^4,x)`output `(tan(c + d*x)^3*((A*1i)/(16*a^4) - (15*B)/(16*a^4)) - tan(c + d*x)^2*(A/(4*a^4) - (B*7i)/(4*a^4)) + A/(12*a^4) - (B*1i)/(3*a^4) + tan(c + d*x)*((A*13i)/(48*a^4) + (61*B)/(48*a^4)))/(d*(tan(c + d*x)*4i - 6*tan(c + d*x)^2 - tan(c + d*x)^3*4i + tan(c + d*x)^4 + 1)) + (log(tan(c + d*x) - 1i)*(A - B*1i))/(32*a^4*d) + (log(tan(c + d*x) + 1i)*(A*1i + B)*1i)/(32*a^4*d)`

3.61 $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

3.61.1	Optimal result	759
3.61.2	Mathematica [A] (verified)	759
3.61.3	Rubi [A] (verified)	760
3.61.4	Maple [A] (verified)	763
3.61.5	Fricas [A] (verification not implemented)	763
3.61.6	Sympy [A] (verification not implemented)	764
3.61.7	Maxima [F(-2)]	764
3.61.8	Giac [A] (verification not implemented)	765
3.61.9	Mupad [B] (verification not implemented)	765

3.61.1 Optimal result

Integrand size = 34, antiderivative size = 145

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = -\frac{(A-iB)x}{16a^4} + \frac{iA+5B}{16a^4d(1+i \tan(c+dx))^2} - \frac{iA+B}{16a^4d(1+i \tan(c+dx))} + \frac{(iA-B) \tan^2(c+dx)}{8d(a+ia \tan(c+dx))^4} - \frac{B}{6ad(a+ia \tan(c+dx))^3}$$

output

```
-1/16*(A-I*B)*x/a^4+1/16*(I*A+5*B)/a^4/d/(1+I*tan(d*x+c))^2+1/16*(-I*A-B)/a^4/d/(1+I*tan(d*x+c))+1/8*(I*A-B)*tan(d*x+c)^2/d/(a+I*a*tan(d*x+c))^4-1/6*B/a/d/(a+I*a*tan(d*x+c))^3
```

3.61.2 Mathematica [A] (verified)

Time = 1.63 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.99

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \frac{(\cos(4(c+dx)) - i \sin(4(c+dx)))(-12iA - 16B \cos(2(c+dx)) + 3(iA - B + 8Adx - 8iBdx) \cos(4(c+dx)))}{(a+ia \tan(c+dx))^4}$$

input `Integrate[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]`

output `-1/384*((Cos[4*(c + d*x)] - I*Sin[4*(c + d*x)])*((-12*I)*A - 16*B*Cos[2*(c + d*x)] + 3*(I*A - B + 8*A*d*x - (8*I)*B*d*x)*Cos[4*(c + d*x)] - (32*I)*B*Sin[2*(c + d*x)] + 3*A*Sin[4*(c + d*x)] + (3*I)*B*Sin[4*(c + d*x)] + (24*I)*A*d*x*Sin[4*(c + d*x)] + 24*B*d*x*Sin[4*(c + d*x)])))/(a^4*d)`

3.61.3 Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.06, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.324$, Rules used = {3042, 4078, 27, 3042, 4073, 27, 3042, 4009, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^2(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+ia) \tan^2(c+dx)}{8d(a+ia \tan(c+dx))^4} - \frac{\int \frac{2 \tan(c+dx)(a(iA-B)-a(A-3iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^3} dx}{8a^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B+ia) \tan^2(c+dx)}{8d(a+ia \tan(c+dx))^4} - \frac{\int \frac{\tan(c+dx)(a(iA-B)-a(A-3iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^3} dx}{4a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B+ia) \tan^2(c+dx)}{8d(a+ia \tan(c+dx))^4} - \frac{\int \frac{\tan(c+dx)(a(iA-B)-a(A-3iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^3} dx}{4a^2} \\
 & \quad \downarrow \text{4073} \\
 & \frac{(-B+ia) \tan^2(c+dx)}{8d(a+ia \tan(c+dx))^4} - \frac{2aB}{3d(a+ia \tan(c+dx))^3} - \frac{i \int \frac{2(2Ba^2+(A-3iB) \tan(c+dx)a^2)}{(i \tan(c+dx)a+a)^2} dx}{2a^2}
 \end{aligned}$$

3.61. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{(-B + iA) \tan^2(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{i \int \frac{2Ba^2 + (A - 3iB) \tan(c + dx) a^2}{(i \tan(c + dx) a + a)^2} dx}{4a^2} + \frac{2aB}{3d(a + ia \tan(c + dx))^3} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^2(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{i \int \frac{2Ba^2 + (A - 3iB) \tan(c + dx) a^2}{(i \tan(c + dx) a + a)^2} dx}{4a^2} + \frac{2aB}{3d(a + ia \tan(c + dx))^3} \\
& \downarrow 4009 \\
& \frac{(-B + iA) \tan^2(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{i \left(-\frac{1}{2} a(B + iA) \int \frac{1}{i \tan(c + dx) a + a} dx - \frac{A - 5iB}{4d(1 + i \tan(c + dx))^2} \right)}{4a^2} + \frac{2aB}{3d(a + ia \tan(c + dx))^3} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^2(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{i \left(-\frac{1}{2} a(B + iA) \int \frac{1}{i \tan(c + dx) a + a} dx - \frac{A - 5iB}{4d(1 + i \tan(c + dx))^2} \right)}{4a^2} + \frac{2aB}{3d(a + ia \tan(c + dx))^3} \\
& \downarrow 3960 \\
& \frac{(-B + iA) \tan^2(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{i \left(-\frac{1}{2} a(B + iA) \left(\frac{\int 1 dx}{2a} + \frac{i}{2d(a + ia \tan(c + dx))} \right) - \frac{A - 5iB}{4d(1 + i \tan(c + dx))^2} \right)}{4a^2} + \frac{2aB}{3d(a + ia \tan(c + dx))^3} \\
& \downarrow 24 \\
& \frac{(-B + iA) \tan^2(c + dx)}{8d(a + ia \tan(c + dx))^4} - \frac{i \left(-\frac{1}{2} a(B + iA) \left(\frac{x}{2a} + \frac{i}{2d(a + ia \tan(c + dx))} \right) - \frac{A - 5iB}{4d(1 + i \tan(c + dx))^2} \right)}{4a^2} + \frac{2aB}{3d(a + ia \tan(c + dx))^3}
\end{aligned}$$

input `Int[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]`

output `((I*A - B)*Tan[c + d*x]^2)/(8*d*(a + I*a*Tan[c + d*x])^4) - ((2*a*B)/(3*d*(a + I*a*Tan[c + d*x])^3) + (I*(-1/4*(A - (5*I)*B))/(d*(1 + I*Tan[c + d*x])^2) - (a*(I*A + B)*(x/(2*a) + (I/2)/(d*(a + I*a*Tan[c + d*x]))))/2))/a^2)/(4*a^2)`

3.61.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3960 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`
- rule 4009 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`
- rule 4073 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-(A*b - a*B))*(a*c + b*d)*((a + b*Tan[e + f*x])^m/(2*a^2*f*m)), x] + Simp[1/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[A*b*c + a*B*c + a*A*d + b*B*d + 2*a*B*d*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && EqQ[a^2 + b^2, 0]`
- rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^n, x_Symbol] := Simp[(-(A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.61.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.75

method	result
risch	$\frac{ixB}{16a^4} - \frac{xA}{16a^4} + \frac{e^{-2i(dx+c)}B}{16da^4} + \frac{ie^{-4i(dx+c)}A}{32da^4} - \frac{e^{-6i(dx+c)}B}{48da^4} + \frac{e^{-8i(dx+c)}B}{128da^4} - \frac{ie^{-8i(dx+c)}A}{128da^4}$
derivativedivides	$-\frac{iA}{8da^4(\tan(dx+c)-i)^4} - \frac{A \arctan(\tan(dx+c))}{16da^4} + \frac{iB \arctan(\tan(dx+c))}{16da^4} - \frac{A}{4da^4(\tan(dx+c)-i)^3} + \frac{iB}{16da^4(\tan(dx+c)-i)^3}$
default	$-\frac{iA}{8da^4(\tan(dx+c)-i)^4} - \frac{A \arctan(\tan(dx+c))}{16da^4} + \frac{iB \arctan(\tan(dx+c))}{16da^4} - \frac{A}{4da^4(\tan(dx+c)-i)^3} + \frac{iB}{16da^4(\tan(dx+c)-i)^3}$

input `int(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x,method=_RETURNVERBOSE)`

output $\frac{1}{16}I*x/a^4*B - \frac{1}{16}*x/a^4*A + \frac{1}{16}/d/a^4*\exp(-2*I*(d*x+c))*B + \frac{1}{32}*I*A/d/a^4*\exp(-4*I*(d*x+c)) - \frac{1}{48}/d/a^4*\exp(-6*I*(d*x+c))*B + \frac{1}{128}/d/a^4*\exp(-8*I*(d*x+c))*B - \frac{1}{128}*I/d/a^4*\exp(-8*I*(d*x+c))*A$

3.61.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.54

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \frac{(24(A-iB)dx e^{(8i dx+8i c)} - 24B e^{(6i dx+6i c)} - 12i A e^{(4i dx+4i c)} + 8B e^{(2i dx+2i c)} + 3i A - 3B) e^{(-8i dx-8i c)}}{384a^4d}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="fracas")`

output $\frac{-1/384*(24*(A - I*B)*d*x*e^{(8*I*d*x + 8*I*c)} - 24*B*e^{(6*I*d*x + 6*I*c)} - 12*I*A*e^{(4*I*d*x + 4*I*c)} + 8*B*e^{(2*I*d*x + 2*I*c)} + 3*I*A - 3*B)*e^{(-8*I*d*x - 8*I*c)}}{(a^4*d)}$

3.61.6 Sympy [A] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.66

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$$

$$= \left\{ \frac{(98304iAa^{12}d^3e^{16ic}e^{-4idx}+196608Ba^{12}d^3e^{18ic}e^{-2idx}-65536Ba^{12}d^3e^{14ic}e^{-6idx}+(-24576iAa^{12}d^3e^{12ic}+24576Ba^{12}d^3e^{12ic})e^{-8idx})e^{-20ic}}{3145728a^{16}d^4} \right.$$

$$\left. x \left(-\frac{-A+iB}{16a^4} + \frac{(-Ae^{8ic}+2Ae^{4ic}-A+iBe^{8ic}-2iBe^{6ic}+2iBe^{2ic}-iB)e^{-8ic}}{16a^4} \right) \right.$$

$$\left. + \frac{x(-A+iB)}{16a^4} \right.$$

input `integrate(tan(d*x+c)**2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**4,x)`

output `Piecewise(((98304*I*A*a**12*d**3*exp(16*I*c)*exp(-4*I*d*x) + 196608*B*a**12*d**3*exp(18*I*c)*exp(-2*I*d*x) - 65536*B*a**12*d**3*exp(14*I*c)*exp(-6*I*d*x) + (-24576*I*A*a**12*d**3*exp(12*I*c) + 24576*B*a**12*d**3*exp(12*I*c))*exp(-8*I*d*x))*exp(-20*I*c)/(3145728*a**16*d**4), Ne(a**16*d**4*exp(20*I*c), 0)), (x*(-(-A + I*B)/(16*a**4) + (-A*exp(8*I*c) + 2*A*exp(4*I*c) - A + I*B*exp(8*I*c) - 2*I*B*exp(6*I*c) + 2*I*B*exp(2*I*c) - I*B)*exp(-8*I*c)/(16*a**4)), True)) + x*(-A + I*B)/(16*a**4)`

3.61.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.61.8 Giac [A] (verification not implemented)

Time = 0.77 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.04

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx =$$

$$\frac{12(iA+B)\log(\tan(dx+c)+i)}{a^4} + \frac{12(-iA-B)\log(\tan(dx+c)-i)}{a^4} + \frac{25iA\tan(dx+c)^4+25B\tan(dx+c)^4+124A\tan(dx+c)^3-124iB\tan(dx+c)^2-124iA\tan(dx+c)+25iA-7B}{a^4} + 384d$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="giac")`

output `-1/384*(12*(I*A + B)*log(tan(d*x + c) + I)/a^4 + 12*(-I*A - B)*log(tan(d*x + c) - I)/a^4 + (25*I*A*tan(d*x + c)^4 + 25*B*tan(d*x + c)^4 + 124*A*tan(d*x + c)^3 - 124*I*B*tan(d*x + c)^3 - 246*I*A*tan(d*x + c)^2 - 54*B*tan(d*x + c)^2 - 124*A*tan(d*x + c) - 4*I*B*tan(d*x + c) + 25*I*A - 7*B)/(a^4*(tan(d*x + c) - I)^4)/d`

3.61.9 Mupad [B] (verification not implemented)

Time = 7.61 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.93

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$$

$$= \frac{\tan(c+dx)^2\left(-\frac{B}{4a^4} + \frac{A1i}{4a^4}\right) - \tan(c+dx)^3\left(\frac{A}{16a^4} - \frac{B1i}{16a^4}\right) + \frac{B}{12a^4} + \tan(c+dx)\left(\frac{A}{16a^4} + \frac{B13i}{48a^4}\right)}{d\left(\tan(c+dx)^4 - \tan(c+dx)^34i - 6\tan(c+dx)^2 + \tan(c+dx)4i + 1\right)} + \frac{x(B+A1i)1i}{16a^4}$$

input `int((tan(c + d*x)^2*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^4,x)`

output `(tan(c + d*x)^2*((A*1i)/(4*a^4) - B/(4*a^4)) - tan(c + d*x)^3*(A/(16*a^4) - (B*1i)/(16*a^4)) + B/(12*a^4) + tan(c + d*x)*(A/(16*a^4) + (B*13i)/(48*a^4)))/(d*(tan(c + d*x)*4i - 6*tan(c + d*x)^2 - tan(c + d*x)^3*4i + tan(c + d*x)^4 + 1)) + (x*(A*1i + B)*1i)/(16*a^4)`

3.62 $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

3.62.1	Optimal result	766
3.62.2	Mathematica [A] (verified)	766
3.62.3	Rubi [A] (verified)	767
3.62.4	Maple [A] (verified)	769
3.62.5	Fricas [A] (verification not implemented)	770
3.62.6	Sympy [A] (verification not implemented)	770
3.62.7	Maxima [F(-2)]	771
3.62.8	Giac [A] (verification not implemented)	771
3.62.9	Mupad [B] (verification not implemented)	772

3.62.1 Optimal result

Integrand size = 32, antiderivative size = 143

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = -\frac{(iA+B)x}{16a^4} - \frac{A+iB}{8d(a+ia \tan(c+dx))^4} + \frac{A+3iB}{12ad(a+ia \tan(c+dx))^3} + \frac{A-iB}{16d(a^2+ia^2 \tan(c+dx))^2} + \frac{A-iB}{16d(a^4+ia^4 \tan(c+dx))}$$

output `-1/16*(I*A+B)*x/a^4+1/8*(-A-I*B)/d/(a+I*a*tan(d*x+c))^4+1/12*(A+3*I*B)/a/d/(a+I*a*tan(d*x+c))^3+1/16*(A-I*B)/d/(a^2+I*a^2*tan(d*x+c))^2+1/16*(A-I*B)/d/(a^4+I*a^4*tan(d*x+c))`

3.62.2 Mathematica [A] (verified)

Time = 0.79 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.99

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \frac{\sec^4(c+dx)(12iB+16A \cos(2(c+dx)) - 3(A+8iA dx + B(i+8dx)) \cos(4(c+dx))) + 32iA \sin(2(c+dx))}{384a^4d(-i + \tan(c+dx))}$$

input `Integrate[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]`

output `(Sec[c + d*x]^4*((12*I)*B + 16*A*Cos[2*(c + d*x)] - 3*(A + (8*I)*A*d*x + B*(I + 8*d*x))*Cos[4*(c + d*x)] + (32*I)*A*Sin[2*(c + d*x)] + (3*I)*A*Sin[4*(c + d*x)] - 3*B*Sin[4*(c + d*x)] + 24*A*d*x*Sin[4*(c + d*x)] - (24*I)*B*d*x*Sin[4*(c + d*x)])/(384*a^4*d*(-I + Tan[c + d*x])^4)`

3.62.3 Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.03, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.281$, Rules used = {3042, 4073, 3042, 4009, 3042, 3960, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx \\
 & \quad \downarrow \text{4073} \\
 & -\frac{i \int \frac{a(A+iB)+2aB \tan(c+dx)}{(i \tan(c+dx)a+a)^3} dx}{2a^2} - \frac{A+iB}{8d(a+ia \tan(c+dx))^4} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{i \int \frac{a(A+iB)+2aB \tan(c+dx)}{(i \tan(c+dx)a+a)^3} dx}{2a^2} - \frac{A+iB}{8d(a+ia \tan(c+dx))^4} \\
 & \quad \downarrow \text{4009} \\
 & -\frac{i \left(\frac{1}{2}(A-iB) \int \frac{1}{(i \tan(c+dx)a+a)^2} dx + \frac{a(-3B+iA)}{6d(a+ia \tan(c+dx))^3} \right)}{2a^2} - \frac{A+iB}{8d(a+ia \tan(c+dx))^4} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{i \left(\frac{1}{2}(A-iB) \int \frac{1}{(i \tan(c+dx)a+a)^2} dx + \frac{a(-3B+iA)}{6d(a+ia \tan(c+dx))^3} \right)}{2a^2} - \frac{A+iB}{8d(a+ia \tan(c+dx))^4} \\
 & \quad \downarrow \text{3960}
 \end{aligned}$$

3.62. $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

$$\begin{array}{c}
 \frac{i \left(\frac{1}{2}(A - iB) \left(\frac{\int \frac{1}{i \tan(c+dx)a+a} dx}{2a} + \frac{i}{4d(a+ia \tan(c+dx))^2} \right) + \frac{a(-3B+iA)}{6d(a+ia \tan(c+dx))^3} \right)}{\frac{2a^2}{A+iB}} \\
 \frac{A+iB}{8d(a+ia \tan(c+dx))^4} \\
 \downarrow \text{3042} \\
 \frac{i \left(\frac{1}{2}(A - iB) \left(\frac{\int \frac{1}{i \tan(c+dx)a+a} dx}{2a} + \frac{i}{4d(a+ia \tan(c+dx))^2} \right) + \frac{a(-3B+iA)}{6d(a+ia \tan(c+dx))^3} \right)}{\frac{2a^2}{A+iB}} \\
 \frac{A+iB}{8d(a+ia \tan(c+dx))^4} \\
 \downarrow \text{3960} \\
 \frac{i \left(\frac{1}{2}(A - iB) \left(\frac{\int \frac{1 dx}{2a} + \frac{i}{2d(a+ia \tan(c+dx))}}{2a} + \frac{i}{4d(a+ia \tan(c+dx))^2} \right) + \frac{a(-3B+iA)}{6d(a+ia \tan(c+dx))^3} \right)}{\frac{2a^2}{A+iB}} \\
 \frac{A+iB}{8d(a+ia \tan(c+dx))^4} \\
 \downarrow \text{24} \\
 \frac{i \left(\frac{a(-3B+iA)}{6d(a+ia \tan(c+dx))^3} + \frac{1}{2}(A - iB) \left(\frac{\frac{x}{2a} + \frac{i}{2d(a+ia \tan(c+dx))}}{2a} + \frac{i}{4d(a+ia \tan(c+dx))^2} \right) \right)}{\frac{2a^2}{A+iB}} \\
 \frac{A+iB}{8d(a+ia \tan(c+dx))^4}
 \end{array}$$

input `Int[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]`

output `-1/8*(A + I*B)/(d*(a + I*a*Tan[c + d*x])^4) - ((I/2)*((a*(I*A - 3*B))/(6*d*(a + I*a*Tan[c + d*x])^3) + ((A - I*B)*((I/4)/(d*(a + I*a*Tan[c + d*x])^2) + (x/(2*a) + (I/2)/(d*(a + I*a*Tan[c + d*x])))/(2*a)))/2))/a^2`

3.62.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

rule 4009 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

rule 4073 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-(A*b - a*B))*(a*c + b*d)*((a + b*Tan[e + f*x])^m/(2*a^2*f*m)), x] + Simp[1/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[A*b*c + a*B*c + a*A*d + b*B*d + 2*a*B*d*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && EqQ[a^2 + b^2, 0]`

3.62.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.76

method	result
risch	$-\frac{xB}{16a^4} - \frac{ixA}{16a^4} + \frac{e^{-2i(dx+c)}A}{16da^4} + \frac{iBe^{-4i(dx+c)}}{32da^4} - \frac{e^{-6i(dx+c)}A}{48da^4} - \frac{ie^{-8i(dx+c)}B}{128da^4} - \frac{e^{-8i(dx+c)}A}{128da^4}$
derivativedivides	$-\frac{iA \arctan(\tan(dx+c))}{16da^4} - \frac{B}{16da^4(\tan(dx+c)-i)} - \frac{iA}{16da^4(\tan(dx+c)-i)} - \frac{iB}{8da^4(\tan(dx+c)-i)^4} - \frac{B \arctan(\tan(dx+c))}{16da^4}$
default	$-\frac{iA \arctan(\tan(dx+c))}{16da^4} - \frac{B}{16da^4(\tan(dx+c)-i)} - \frac{iA}{16da^4(\tan(dx+c)-i)} - \frac{iB}{8da^4(\tan(dx+c)-i)^4} - \frac{B \arctan(\tan(dx+c))}{16da^4}$

input `int(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `-1/16*x/a^4*B-1/16*I*x/a^4*A+1/16/d/a^4*exp(-2*I*(d*x+c))*A+1/32*I*B/d/a^4*exp(-4*I*(d*x+c))-1/48/d/a^4*exp(-6*I*(d*x+c))*A-1/128*I/d/a^4*exp(-8*I*(d*x+c))*B-1/128/d/a^4*exp(-8*I*(d*x+c))*A`

3.62.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.55

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx = \frac{(24(iA+B)dx e^{(8i dx+8i c)} - 24Ae^{(6i dx+6i c)} - 12iBe^{(4i dx+4i c)} + 8Ae^{(2i dx+2i c)} + 3A + 3iB)e^{(-8i dx-8i c)}}{384a^4d}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="fracas")`

output `-1/384*(24*(I*A + B)*d*x*e^(8*I*d*x + 8*I*c) - 24*A*e^(6*I*d*x + 6*I*c) - 12*I*B*e^(4*I*d*x + 4*I*c) + 8*A*e^(2*I*d*x + 2*I*c) + 3*A + 3*I*B)*e^(-8*I*d*x - 8*I*c)/(a^4*d)`

3.62.6 Sympy [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.72

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx = \left\{ \frac{(196608Aa^{12}d^3e^{18ic}e^{-2idx} - 65536Aa^{12}d^3e^{14ic}e^{-6idx} + 98304iBa^{12}d^3e^{16ic}e^{-4idx} + (-24576Aa^{12}d^3e^{12ic} - 24576iBa^{12}d^3e^{12ic})e^{-8idx})e^{-20ic}}{3145728a^{16}d^4} \right. \\ \left. x \left(-\frac{iA-B}{16a^4} + \frac{(-iAe^{8ic} - 2iAe^{6ic} + 2iAe^{2ic} + iA - Be^{8ic} + 2Be^{4ic} - B)e^{-8ic}}{16a^4} \right) \right. \\ \left. + \frac{x(-iA - B)}{16a^4} \right.$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**4,x)`

output `Piecewise((((196608*A*a**12*d**3*exp(18*I*c)*exp(-2*I*d*x) - 65536*A*a**12*d**3*exp(14*I*c)*exp(-6*I*d*x) + 98304*I*B*a**12*d**3*exp(16*I*c)*exp(-4*I*d*x) + (-24576*A*a**12*d**3*exp(12*I*c) - 24576*I*B*a**12*d**3*exp(12*I*c))*exp(-8*I*d*x))*exp(-20*I*c)/(3145728*a**16*d**4), Ne(a**16*d**4*exp(20*I*c), 0)), (x*(-(-I*A - B)/(16*a**4) + (-I*A*exp(8*I*c) - 2*I*A*exp(6*I*c) + 2*I*A*exp(2*I*c) + I*A - B*exp(8*I*c) + 2*B*exp(4*I*c) - B)*exp(-8*I*c)/(16*a**4)), True)) + x*(-I*A - B)/(16*a**4)`

3.62.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.62.8 Giac [A] (verification not implemented)

Time = 0.64 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.04

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$$

$$= \frac{12(A-iB)\log(\tan(dx+c)+i) - 12(A-iB)\log(\tan(dx+c)-i) + 25A\tan(dx+c)^4 - 25iB\tan(dx+c)^4 - 124iA\tan(dx+c)^3 - 124B\tan(dx+c)^3}{384d}$$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="giac")
```

```
output 1/384*(12*(A - I*B)*log(tan(d*x + c) + I)/a^4 - 12*(A - I*B)*log(tan(d*x + c) - I)/a^4 + (25*A*tan(d*x + c)^4 - 25*I*B*tan(d*x + c)^4 - 124*I*A*tan(d*x + c)^3 - 124*B*tan(d*x + c)^3 - 246*A*tan(d*x + c)^2 + 246*I*B*tan(d*x + c)^2 + 252*I*A*tan(d*x + c) + 124*B*tan(d*x + c) + 57*A - 25*I*B)/(a^4*(tan(d*x + c) - I)^4)/d
```

3.62.9 Mupad [B] (verification not implemented)

Time = 8.42 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.20

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx =$$

$$-\frac{\tan(c+dx)^2\left(\frac{A}{4a^4}-\frac{B1i}{4a^4}\right)+\tan(c+dx)^3\left(\frac{B}{16a^4}+\frac{A1i}{16a^4}\right)-\frac{A}{12a^4}-\tan(c+dx)\left(\frac{B}{16a^4}+\frac{A19i}{48a^4}\right)}{d\left(\tan(c+dx)^4-\tan(c+dx)^34i-6\tan(c+dx)^2+\tan(c+dx)4i+1\right)}$$

$$+\frac{\ln(\tan(c+dx)-i)(B+A1i)1i}{32a^4d}+\frac{\ln(\tan(c+dx)+1i)(A-B1i)}{32a^4d}$$

input `int((tan(c + d*x)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^4,x)`output `(log(tan(c + d*x) - 1i)*(A*1i + B)*1i)/(32*a^4*d) - (tan(c + d*x)^2*(A/(4*a^4) - (B*1i)/(4*a^4)) + tan(c + d*x)^3*((A*1i)/(16*a^4) + B/(16*a^4)) - A/(12*a^4) - tan(c + d*x)*((A*19i)/(48*a^4) + B/(16*a^4)))/(d*(tan(c + d*x)^4i - 6*tan(c + d*x)^2 - tan(c + d*x)^3*4i + tan(c + d*x)^4 + 1)) + (log(tan(c + d*x) + 1i)*(A - B*1i))/(32*a^4*d)`

3.63 $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^4} dx$

3.63.1	Optimal result	773
3.63.2	Mathematica [A] (verified)	773
3.63.3	Rubi [A] (verified)	774
3.63.4	Maple [A] (verified)	776
3.63.5	Fricas [A] (verification not implemented)	776
3.63.6	Sympy [A] (verification not implemented)	777
3.63.7	Maxima [F(-2)]	777
3.63.8	Giac [A] (verification not implemented)	778
3.63.9	Mupad [B] (verification not implemented)	778

3.63.1 Optimal result

Integrand size = 26, antiderivative size = 145

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^4} dx = \frac{(A - iB)x}{16a^4} + \frac{iA - B}{8d(a + ia \tan(c + dx))^4} + \frac{iA + B}{12ad(a + ia \tan(c + dx))^3} + \frac{iA + B}{16d(a^2 + ia^2 \tan(c + dx))^2} + \frac{iA + B}{16d(a^4 + ia^4 \tan(c + dx))}$$

output `1/16*(A-I*B)*x/a^4+1/8*(I*A-B)/d/(a+I*a*tan(d*x+c))^4+1/12*(I*A+B)/a/d/(a+I*a*tan(d*x+c))^3+1/16*(I*A+B)/d/(a^2+I*a^2*tan(d*x+c))^2+1/16*(I*A+B)/d/(a^4+I*a^4*tan(d*x+c))`

3.63.2 Mathematica [A] (verified)

Time = 1.34 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.91

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^4} dx = \frac{\sec^3(c + dx)(18iA \cos(c + dx) + 2(7iA + 4B) \cos(3(c + dx)) - (A - iB)(5 \sin(c + dx) + 11 \sin(3(c + dx)))}{96a^4d(-i + \tan(c + dx))}$$

input `Integrate[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x])^4,x]`

output $(\text{Sec}[c + d*x]^3*((18*I)*A*\text{Cos}[c + d*x] + 2*((7*I)*A + 4*B)*\text{Cos}[3*(c + d*x)] - (A - I*B)*(5*\text{Sin}[c + d*x] + 11*\text{Sin}[3*(c + d*x)]) + 6*(A - I*B)*\text{ArcTan}[\text{Tan}[c + d*x]]*\text{Sec}[c + d*x]*(\text{Cos}[4*(c + d*x)] + I*\text{Sin}[4*(c + d*x)])))/(96*a^4*d*(-I + \text{Tan}[c + d*x])^4)$

3.63.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.99, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.346$, Rules used = {3042, 4009, 3042, 3960, 3042, 3960, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^4} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^4} dx \\ & \quad \downarrow \text{4009} \\ & \frac{(A - iB) \int \frac{1}{(i \tan(c+dx)a+a)^3} dx}{2a} + \frac{-B + iA}{8d(a + ia \tan(c + dx))^4} \\ & \quad \downarrow \text{3042} \\ & \frac{(A - iB) \int \frac{1}{(i \tan(c+dx)a+a)^3} dx}{2a} + \frac{-B + iA}{8d(a + ia \tan(c + dx))^4} \\ & \quad \downarrow \text{3960} \\ & \frac{(A - iB) \left(\int \frac{1}{(i \tan(c+dx)a+a)^2} dx + \frac{i}{6d(a+ia \tan(c+dx))^3} \right)}{2a} + \frac{-B + iA}{8d(a + ia \tan(c + dx))^4} \\ & \quad \downarrow \text{3042} \\ & \frac{(A - iB) \left(\int \frac{1}{(i \tan(c+dx)a+a)^2} dx + \frac{i}{6d(a+ia \tan(c+dx))^3} \right)}{2a} + \frac{-B + iA}{8d(a + ia \tan(c + dx))^4} \\ & \quad \downarrow \text{3960} \end{aligned}$$

$$\begin{aligned}
& \frac{(A - iB) \left(\frac{\int \frac{1}{i \tan(c+dx)a+a} dx}{2a} + \frac{i}{4d(a+ia \tan(c+dx))^2} + \frac{i}{6d(a+ia \tan(c+dx))^3} \right)}{2a} + \frac{-B + iA}{8d(a + ia \tan(c + dx))^4} \\
& \quad \downarrow \text{3042} \\
& \frac{(A - iB) \left(\frac{\int \frac{1}{i \tan(c+dx)a+a} dx}{2a} + \frac{i}{4d(a+ia \tan(c+dx))^2} + \frac{i}{6d(a+ia \tan(c+dx))^3} \right)}{2a} + \frac{-B + iA}{8d(a + ia \tan(c + dx))^4} \\
& \quad \downarrow \text{3960} \\
& \frac{(A - iB) \left(\frac{\int \frac{1 dx}{2a} + \frac{i}{2d(a+ia \tan(c+dx))}}{2a} + \frac{i}{4d(a+ia \tan(c+dx))^2} + \frac{i}{6d(a+ia \tan(c+dx))^3} \right)}{2a} + \frac{-B + iA}{8d(a + ia \tan(c + dx))^4} \\
& \quad \downarrow \text{24} \\
& \frac{-B + iA}{8d(a + ia \tan(c + dx))^4} + \frac{(A - iB) \left(\frac{\frac{x}{2a} + \frac{i}{2d(a+ia \tan(c+dx))}}{2a} + \frac{i}{4d(a+ia \tan(c+dx))^2} + \frac{i}{6d(a+ia \tan(c+dx))^3} \right)}{2a}
\end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x])^4, x]`

output `(I*A - B)/(8*d*(a + I*a*Tan[c + d*x])^4) + ((A - I*B)*((I/6)/(d*(a + I*a*Tan[c + d*x])^3) + ((I/4)/(d*(a + I*a*Tan[c + d*x])^2) + (x/(2*a) + (I/2)/(d*(a + I*a*Tan[c + d*x])))/(2*a)))/(2*a)))/(2*a)`

3.63.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

rule 4009 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

3.63.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.01

method	result
risch	$-\frac{i x B}{16 a^4} + \frac{x A}{16 a^4} + \frac{e^{-2 i(d x+c)} B}{16 d a^4} + \frac{i e^{-2 i(d x+c)} A}{8 d a^4} + \frac{3 i e^{-4 i(d x+c)} A}{32 d a^4} - \frac{e^{-6 i(d x+c)} B}{48 d a^4} + \frac{i e^{-6 i(d x+c)} A}{24 d a^4} - \frac{e^{-8 i(d x+c)} B}{128 d a^4}$
derivativedivides	$-\frac{i A}{16 d a^4(\tan(dx+c)-i)^2} + \frac{A \arctan(\tan(dx+c))}{16 d a^4} - \frac{i B}{16 d a^4(\tan(dx+c)-i)} - \frac{i B \arctan(\tan(dx+c))}{16 d a^4} - \frac{e^{-8 i(d x+c)} B}{16 d a^4(\tan(dx+c)-i)}$
default	$-\frac{i A}{16 d a^4(\tan(dx+c)-i)^2} + \frac{A \arctan(\tan(dx+c))}{16 d a^4} - \frac{i B}{16 d a^4(\tan(dx+c)-i)} - \frac{i B \arctan(\tan(dx+c))}{16 d a^4} - \frac{e^{-8 i(d x+c)} B}{16 d a^4(\tan(dx+c)-i)}$

input `int((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `-1/16*I*x/a^4*B+1/16*x/a^4*A+1/16/d/a^4*exp(-2*I*(d*x+c))*B+1/8*I/d/a^4*exp(-2*I*(d*x+c))*A+3/32*I/d/a^4*exp(-4*I*(d*x+c))*A-1/48/d/a^4*exp(-6*I*(d*x+c))*B+1/24*I/d/a^4*exp(-6*I*(d*x+c))*A-1/128/d/a^4*exp(-8*I*(d*x+c))*B+1/128*I/d/a^4*exp(-8*I*(d*x+c))*A`

3.63.5 Fracas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.61

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^4} dx$$

$$= \frac{(24(A - i B)dx e^{8i(dx+ci)} - 24(-2iA - B)e^{6i(dx+6ic)} + 36iAe^{4i(dx+4ic)} - 8(-2iA + B)e^{2i(dx+2ic)} + 3iA)}{384a^4d}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="fracas")`

3.63. $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^4} dx$

output $\frac{1}{384}*(24*(A - I*B)*d*x*e^{(8*I*d*x + 8*I*c)} - 24*(-2*I*A - B)*e^{(6*I*d*x + 6*I*c)} + 36*I*A*e^{(4*I*d*x + 4*I*c)} - 8*(-2*I*A + B)*e^{(2*I*d*x + 2*I*c)} + 3*I*A - 3*B)*e^{(-8*I*d*x - 8*I*c)}/(a^4*d)$

3.63.6 Sympy [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 299, normalized size of antiderivative = 2.06

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^4} dx$$

$$= \left\{ \frac{(294912iAa^{12}d^3e^{16ic}e^{-4idx} + (24576iAa^{12}d^3e^{12ic} - 24576Ba^{12}d^3e^{12ic})e^{-8idx} + (131072iAa^{12}d^3e^{14ic} - 65536Ba^{12}d^3e^{14ic})e^{-6idx} + (393216iAa^{12}d^3e^{16ic} - 3145728a^{16}d^4))e^{-8ic}}{3145728a^{16}d^4} \right.$$

$$\left. x \left(-\frac{A-iB}{16a^4} + \frac{(Ae^{8ic} + 4Ae^{6ic} + 6Ae^{4ic} + 4Ae^{2ic} + A - iBe^{8ic} - 2iBe^{6ic} + 2iBe^{2ic} + iB)e^{-8ic}}{16a^4} \right) \right.$$

$$\left. + \frac{x(A - iB)}{16a^4} \right.$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**4,x)`

output `Piecewise((((294912*I*A*a**12*d**3*exp(16*I*c)*exp(-4*I*d*x) + (24576*I*A*a**12*d**3*exp(12*I*c) - 24576*B*a**12*d**3*exp(12*I*c))*exp(-8*I*d*x) + (131072*I*A*a**12*d**3*exp(14*I*c) - 65536*B*a**12*d**3*exp(14*I*c))*exp(-6*I*d*x) + (393216*I*A*a**12*d**3*exp(18*I*c) + 196608*B*a**12*d**3*exp(18*I*c))*exp(-2*I*d*x))*exp(-20*I*c)/(3145728*a**16*d**4), Ne(a**16*d**4*exp(20*I*c), 0)), (x*(-(A - I*B)/(16*a**4) + (A*exp(8*I*c) + 4*A*exp(6*I*c) + 6*A*exp(4*I*c) + 4*A*exp(2*I*c) + A - I*B*exp(8*I*c) - 2*I*B*exp(6*I*c) + 2*I*B*exp(2*I*c) + I*B)*exp(-8*I*c)/(16*a**4)), True)) + x*(A - I*B)/(16*a**4)`

3.63.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^4} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.63. $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^4} dx$

3.63.8 Giac [A] (verification not implemented)

Time = 0.61 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.06

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^4} dx =$$

$$\frac{12(-iA-B)\log(\tan(dx+c)+i)}{a^4} - \frac{12(-iA-B)\log(\tan(dx+c)-i)}{a^4} - \frac{25iA \tan(dx+c)^4 + 25B \tan(dx+c)^4 + 124A \tan(dx+c)^3 - 124iB \tan(dx+c)^3 + 124A \tan(dx+c)^2 - 124iB \tan(dx+c)^2 + 124A \tan(dx+c) - 124iB \tan(dx+c) + 153iA + 57B}{384d}$$

```
input integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="giac")
```

```
output -1/384*(12*(-I*A - B)*log(tan(d*x + c) + I)/a^4 - 12*(-I*A - B)*log(tan(d*x + c) - I)/a^4 - (25*I*A*tan(d*x + c)^4 + 25*B*tan(d*x + c)^4 + 124*A*tan(d*x + c)^3 - 124*I*B*tan(d*x + c)^3 - 246*I*A*tan(d*x + c)^2 - 246*B*tan(d*x + c)^2 - 252*A*tan(d*x + c) + 252*I*B*tan(d*x + c) + 153*I*A + 57*B)/(a^4*(tan(d*x + c) - I)^4)/d
```

3.63.9 Mupad [B] (verification not implemented)

Time = 7.67 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.99

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^4} dx$$

$$= \frac{\frac{B}{12a^4} + \tan(c + dx)^3 \left(\frac{A}{16a^4} - \frac{B1i}{16a^4} \right) + \frac{A1i}{3a^4} - \tan(c + dx)^2 \left(\frac{B}{4a^4} + \frac{A1i}{4a^4} \right) - \tan(c + dx) \left(\frac{19A}{48a^4} - \frac{B19i}{48a^4} \right)}{d \left(\tan(c + dx)^4 - \tan(c + dx)^3 4i - 6 \tan(c + dx)^2 + \tan(c + dx) 4i + 1 \right)}$$

$$- \frac{x(B + A1i)1i}{16a^4}$$

```
input int((A + B*tan(c + d*x))/(a + a*tan(c + d*x)*1i)^4,x)
```

```
output (tan(c + d*x)^3*(A/(16*a^4) - (B*1i)/(16*a^4)) - tan(c + d*x)^2*((A*1i)/(4*a^4) + B/(4*a^4)) + (A*1i)/(3*a^4) + B/(12*a^4) - tan(c + d*x)*((19*A)/(48*a^4) - (B*19i)/(48*a^4)))/(d*(tan(c + d*x)*4i - 6*tan(c + d*x)^2 - tan(c + d*x)^3*4i + tan(c + d*x)^4 + 1)) - (x*(A*1i + B)*1i)/(16*a^4)
```

3.64 $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

3.64.1	Optimal result	779
3.64.2	Mathematica [A] (verified)	780
3.64.3	Rubi [A] (verified)	780
3.64.4	Maple [A] (verified)	784
3.64.5	Fricas [A] (verification not implemented)	784
3.64.6	Sympy [A] (verification not implemented)	785
3.64.7	Maxima [F(-2)]	785
3.64.8	Giac [A] (verification not implemented)	786
3.64.9	Mupad [B] (verification not implemented)	786

3.64.1 Optimal result

Integrand size = 32, antiderivative size = 162

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = -\frac{(15iA-B)x}{16a^4} + \frac{A \log(\sin(c+dx))}{a^4d} + \frac{7A+iB}{16a^4d(1+i \tan(c+dx))^2} + \frac{15A+iB}{16a^4d(1+i \tan(c+dx))} + \frac{A+iB}{8d(a+ia \tan(c+dx))^4} + \frac{3A+iB}{12ad(a+ia \tan(c+dx))^3}$$

output

```
-1/16*(15*I*A-B)*x/a^4+A*ln(sin(d*x+c))/a^4/d+1/16*(7*A+I*B)/a^4/d/(1+I*tan(d*x+c))^2+1/16*(15*A+I*B)/a^4/d/(1+I*tan(d*x+c))+1/8*(A+I*B)/d/(a+I*a*tan(d*x+c))^4+1/12*(3*A+I*B)/a/d/(a+I*a*tan(d*x+c))^3
```

3.64.2 Mathematica [A] (verified)

Time = 1.72 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.94

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$$

$$= \frac{-3(31A+iB) \log(i-\tan(c+dx)) + 96A \log(\tan(c+dx)) - 3(A-iB) \log(i+\tan(c+dx)) + \frac{12(A+iB)}{(-i+\tan(c+dx))}}{96a^4d}$$

input `Integrate[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]`

output `(-3*(31*A + I*B)*Log[I - Tan[c + d*x]] + 96*A*Log[Tan[c + d*x]] - 3*(A - I*B)*Log[I + Tan[c + d*x]] + (12*(A + I*B))/(-I + Tan[c + d*x])^4 + ((24*I)*A - 8*B)/(-I + Tan[c + d*x])^3 - (6*(7*A + I*B))/(-I + Tan[c + d*x])^2 + (6*((-15*I)*A + B))/(-I + Tan[c + d*x]))/(96*a^4*d)`

3.64.3 Rubi [A] (verified)

Time = 1.17 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.18, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A+B \tan(c+dx)}{\tan(c+dx)(a+ia \tan(c+dx))^4} dx$$

$$\downarrow \text{4079}$$

$$\frac{\int \frac{4 \cot(c+dx)(2aA-a(iA-B) \tan(c+dx))}{(i \tan(c+dx)a+a)^3} dx}{8a^2} + \frac{A+iB}{8d(a+ia \tan(c+dx))^4}$$

$$\downarrow \text{27}$$

$$\frac{\int \frac{\cot(c+dx)(2aA-a(iA-B) \tan(c+dx))}{(i \tan(c+dx)a+a)^3} dx}{2a^2} + \frac{A+iB}{8d(a+ia \tan(c+dx))^4}$$

3.64. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

$$\begin{aligned}
& \int \frac{2aA - a(iA - B) \tan(c+dx)}{\tan(c+dx)(i \tan(c+dx)a + a)^3} dx + \frac{A + iB}{8d(a + ia \tan(c + dx))^4} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{3 \cot(c+dx)(4a^2 A - a^2(3iA - B) \tan(c+dx))}{6a^2} dx}{2a^2} + \frac{a(3A + iB)}{6d(a + ia \tan(c+dx))^3} + \frac{A + iB}{8d(a + ia \tan(c + dx))^4} \\
& \quad \downarrow 4079 \\
& \frac{\int \frac{\cot(c+dx)(4a^2 A - a^2(3iA - B) \tan(c+dx))}{2a^2} dx}{2a^2} + \frac{a(3A + iB)}{6d(a + ia \tan(c+dx))^3} + \frac{A + iB}{8d(a + ia \tan(c + dx))^4} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{4a^2 A - a^2(3iA - B) \tan(c+dx)}{\tan(c+dx)(i \tan(c+dx)a + a)^2} dx}{2a^2} + \frac{a(3A + iB)}{6d(a + ia \tan(c+dx))^3} + \frac{A + iB}{8d(a + ia \tan(c + dx))^4} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{2 \cot(c+dx)(8a^3 A - a^3(7iA - B) \tan(c+dx))}{4a^2} dx}{2a^2} + \frac{7A + iB}{4d(1 + i \tan(c+dx))^2} + \frac{a(3A + iB)}{6d(a + ia \tan(c+dx))^3} + \frac{A + iB}{8d(a + ia \tan(c + dx))^4} \\
& \quad \downarrow 4079 \\
& \frac{\int \frac{\cot(c+dx)(8a^3 A - a^3(7iA - B) \tan(c+dx))}{2a^2} dx}{2a^2} + \frac{7A + iB}{4d(1 + i \tan(c+dx))^2} + \frac{a(3A + iB)}{6d(a + ia \tan(c+dx))^3} + \frac{A + iB}{8d(a + ia \tan(c + dx))^4} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{8a^3 A - a^3(7iA - B) \tan(c+dx)}{\tan(c+dx)(i \tan(c+dx)a + a)} dx}{2a^2} + \frac{7A + iB}{4d(1 + i \tan(c+dx))^2} + \frac{a(3A + iB)}{6d(a + ia \tan(c+dx))^3} + \frac{A + iB}{8d(a + ia \tan(c + dx))^4} \\
& \quad \downarrow 3042 \\
& \frac{\int \cot(c+dx)(16a^4 A - a^4(15iA - B) \tan(c+dx)) dx}{2a^2} + \frac{a^3(15A + iB)}{2d(a + ia \tan(c+dx))} + \frac{7A + iB}{4d(1 + i \tan(c+dx))^2} + \frac{a(3A + iB)}{6d(a + ia \tan(c+dx))^3} + \frac{A + iB}{8d(a + ia \tan(c + dx))^4} \\
& \quad \downarrow 4079 \\
& \frac{2a^2}{8d(a + ia \tan(c + dx))^4} + \frac{A + iB}{8d(a + ia \tan(c + dx))^4}
\end{aligned}$$

3.64. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{\int \frac{16a^4 A - a^4(15iA - B) \tan(c+dx)}{2a^2} dx + \frac{a^3(15A+iB)}{2d(a+ia \tan(c+dx))} + \frac{7A+iB}{4d(1+i \tan(c+dx))^2} + \frac{a(3A+iB)}{6d(a+ia \tan(c+dx))^3} + \frac{2a^2}{A+iB}}{8d(a+ia \tan(c+dx))^4} \\
& \downarrow 4014 \\
& \frac{\frac{16a^4 A \int \cot(c+dx) dx - a^4 x(-B+15iA)}{2a^2} + \frac{a^3(15A+iB)}{2d(a+ia \tan(c+dx))} + \frac{7A+iB}{4d(1+i \tan(c+dx))^2} + \frac{a(3A+iB)}{6d(a+ia \tan(c+dx))^3} + \frac{2a^2}{A+iB}}{8d(a+ia \tan(c+dx))^4} \\
& \downarrow 3042 \\
& \frac{\frac{16a^4 A \int -\tan(c+dx+\frac{\pi}{2}) dx - a^4 x(-B+15iA)}{2a^2} + \frac{a^3(15A+iB)}{2d(a+ia \tan(c+dx))} + \frac{7A+iB}{4d(1+i \tan(c+dx))^2} + \frac{a(3A+iB)}{6d(a+ia \tan(c+dx))^3} + \frac{2a^2}{A+iB}}{8d(a+ia \tan(c+dx))^4} \\
& \downarrow 25 \\
& \frac{\frac{-16a^4 A \int \tan(\frac{1}{2}(2c+\pi)+dx) dx - (a^4 x(-B+15iA))}{2a^2} + \frac{a^3(15A+iB)}{2d(a+ia \tan(c+dx))} + \frac{7A+iB}{4d(1+i \tan(c+dx))^2} + \frac{a(3A+iB)}{6d(a+ia \tan(c+dx))^3} + \frac{2a^2}{A+iB}}{8d(a+ia \tan(c+dx))^4} \\
& \downarrow 3956 \\
& \frac{\frac{a^3(15A+iB)}{2d(a+ia \tan(c+dx))} + \frac{16a^4 A \log(-\sin(c+dx)) - a^4 x(-B+15iA)}{2a^2} + \frac{7A+iB}{4d(1+i \tan(c+dx))^2} + \frac{a(3A+iB)}{6d(a+ia \tan(c+dx))^3} + \frac{2a^2}{A+iB}}{8d(a+ia \tan(c+dx))^4}
\end{aligned}$$

input `Int[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]`

```
output (A + I*B)/(8*d*(a + I*a*Tan[c + d*x])^4) + ((a*(3*A + I*B))/(6*d*(a + I*a*
Tan[c + d*x])^3) + ((7*A + I*B)/(4*d*(1 + I*Tan[c + d*x])^2) + ((-(a^4*((1
5*I)*A - B)*x) + (16*a^4*A*Log[-Sin[c + d*x]])/d)/(2*a^2) + (a^3*(15*A + I
*B))/(2*d*(a + I*a*Tan[c + d*x])))/(2*a^2))/(2*a^2))/(2*a^2)
```

3.64.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4079 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.64.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.21

method	result
risch	$\frac{x B}{16 a^4} - \frac{31 i x A}{16 a^4} + \frac{i e^{-2 i(dx+c)} B}{8 d a^4} + \frac{13 e^{-2 i(dx+c)} A}{16 d a^4} + \frac{3 i B e^{-4 i(dx+c)}}{32 d a^4} + \frac{e^{-4 i(dx+c)} A}{4 d a^4} + \frac{i e^{-6 i(dx+c)} B}{24 d a^4} + \frac{e^{-6 i(dx+c)} A}{24 d a^4}$
derivativedivides	$-\frac{i B}{16 d a^4 (\tan(dx+c)-i)^2} + \frac{B}{16 d a^4 (\tan(dx+c)-i)} + \frac{i B}{8 d a^4 (\tan(dx+c)-i)^4} + \frac{B \arctan(\tan(dx+c))}{16 d a^4} - \frac{A \ln(1+\tan(dx+c))}{2 d a^4}$
default	$-\frac{i B}{16 d a^4 (\tan(dx+c)-i)^2} + \frac{B}{16 d a^4 (\tan(dx+c)-i)} + \frac{i B}{8 d a^4 (\tan(dx+c)-i)^4} + \frac{B \arctan(\tan(dx+c))}{16 d a^4} - \frac{A \ln(1+\tan(dx+c))}{2 d a^4}$

input `int(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `1/16*x/a^4*B-31/16*I*x/a^4*A+1/8*I/d/a^4*exp(-2*I*(d*x+c))*B+13/16/d/a^4*exp(-2*I*(d*x+c))*A+3/32*I*B/d/a^4*exp(-4*I*(d*x+c))+1/4/d/a^4*exp(-4*I*(d*x+c))*A+1/24*I/d/a^4*exp(-6*I*(d*x+c))*B+1/16/d/a^4*exp(-6*I*(d*x+c))*A+1/128*I/d/a^4*exp(-8*I*(d*x+c))*B+1/128/d/a^4*exp(-8*I*(d*x+c))*A-2*I/a^4*A/d*c+1/a^4*A/d*ln(exp(2*I*(d*x+c))-1)`

3.64.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.75

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \frac{(24(31iA-B)dx e^{(8i dx+8i c)} - 384Ae^{(8i dx+8i c)} \log(e^{(2i dx+2i c)} - 1) - 24(13A+2iB)e^{(6i dx+6i c)} - 12(2iA-3iB)e^{(4i dx+4i c)} - 8(3A+2iB)e^{(2i dx+2i c)} - 3A - 3iB)e^{(-8i dx-8i c)}}{384 a^4 d}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="fricas")`

output `-1/384*(24*(31*I*A - B)*d*x*e^(8*I*d*x + 8*I*c) - 384*A*e^(8*I*d*x + 8*I*c)*log(e^(2*I*d*x + 2*I*c) - 1) - 24*(13*A + 2*I*B)*e^(6*I*d*x + 6*I*c) - 12*(8*A + 3*I*B)*e^(4*I*d*x + 4*I*c) - 8*(3*A + 2*I*B)*e^(2*I*d*x + 2*I*c) - 3*A - 3*I*B)*e^(-8*I*d*x - 8*I*c)/(a^4*d)`

3.64.6 Sympy [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 359, normalized size of antiderivative = 2.22

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx = \frac{A \log(e^{2idx} - e^{-2ic})}{a^4 d} + \left\{ \frac{((24576Aa^{12}d^3e^{12ic} + 24576iBa^{12}d^3e^{12ic})e^{-8idx} + (196608Aa^{12}d^3e^{14ic} + 131072iBa^{12}d^3e^{14ic})e^{-6idx} + (786432Aa^{12}d^3e^{16ic} + 294912iBa^{12}d^3e^{16ic})e^{-4idx} + (2555904Aa^{12}d^3e^{18ic} + 393216iBa^{12}d^3e^{18ic})e^{-2idx})e^{-20ic}}{3145728a^{16}d^4} \right. \\ \left. + x \left(-\frac{31iA+B}{16a^4} + \frac{(-31iAe^{8ic} - 26iAe^{6ic} - 16iAe^{4ic} - 6iAe^{2ic} - iA + Be^{8ic} + 4Be^{6ic} + 6Be^{4ic} + 4Be^{2ic} + B)e^{-8ic}}{16a^4} \right) \right. \\ \left. + \frac{x(-31iA+B)}{16a^4} \right.$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**4,x)`

output `A*log(exp(2*I*d*x) - exp(-2*I*c))/(a**4*d) + Piecewise((((24576*A*a**12*d**3*exp(12*I*c) + 24576*I*B*a**12*d**3*exp(12*I*c))*exp(-8*I*d*x) + (196608*A*a**12*d**3*exp(14*I*c) + 131072*I*B*a**12*d**3*exp(14*I*c))*exp(-6*I*d*x) + (786432*A*a**12*d**3*exp(16*I*c) + 294912*I*B*a**12*d**3*exp(16*I*c))*exp(-4*I*d*x) + (2555904*A*a**12*d**3*exp(18*I*c) + 393216*I*B*a**12*d**3*exp(18*I*c))*exp(-2*I*d*x))*exp(-20*I*c)/(3145728*a**16*d**4), Ne(a**16*d**4*exp(20*I*c), 0)), (x*(-(31*I*A + B)/(16*a**4) + (-31*I*A*exp(8*I*c) - 26*I*A*exp(6*I*c) - 16*I*A*exp(4*I*c) - 6*I*A*exp(2*I*c) - I*A + B*exp(8*I*c) + 4*B*exp(6*I*c) + 6*B*exp(4*I*c) + 4*B*exp(2*I*c) + B)*exp(-8*I*c)/(16*a**4)), True)) + x*(-31*I*A + B)/(16*a**4)`

3.64.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.64.8 Giac [A] (verification not implemented)

Time = 0.91 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.02

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \frac{12(A-iB) \log(\tan(dx+c)+i)}{a^4} + \frac{12(31A+iB) \log(\tan(dx+c)-i)}{a^4} - \frac{384A \log(\tan(dx+c))}{a^4} - \frac{775A \tan(dx+c)^4 + 25iB \tan(dx+c)^4 - 3460iB \tan(dx+c)^3 + 124B \tan(dx+c)^3 - 5898A \tan(dx+c)^2 - 246iB \tan(dx+c)^2 + 4612iA \tan(dx+c) - 252iB \tan(dx+c) + 1447A + 153iB}{a^4(\tan(dx+c) - i)^4} / d$$

384 d

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="giac")`

output `-1/384*(12*(A - I*B)*log(tan(d*x + c) + I)/a^4 + 12*(31*A + I*B)*log(tan(d*x + c) - I)/a^4 - 384*A*log(tan(d*x + c))/a^4 - (775*A*tan(d*x + c)^4 + 25*I*B*tan(d*x + c)^4 - 3460*I*A*tan(d*x + c)^3 + 124*B*tan(d*x + c)^3 - 5898*A*tan(d*x + c)^2 - 246*I*B*tan(d*x + c)^2 + 4612*I*A*tan(d*x + c) - 252*I*B*tan(d*x + c) + 1447*A + 153*I*B)/(a^4*(tan(d*x + c) - I)^4))/d`

3.64.9 Mupad [B] (verification not implemented)

Time = 7.30 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.21

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \frac{\frac{7A}{4a^4} - \tan(c+dx)^3 \left(-\frac{B}{16a^4} + \frac{A15i}{16a^4}\right) - \tan(c+dx)^2 \left(\frac{13A}{4a^4} + \frac{B1i}{4a^4}\right) + \frac{B1i}{3a^4} + \tan(c+dx) \left(-\frac{19B}{48a^4} + \frac{A63i}{16a^4}\right)}{d(\tan(c+dx)^4 - \tan(c+dx)^3 4i - 6 \tan(c+dx)^2 + \tan(c+dx) 4i + 1)} + \frac{A \ln(\tan(c+dx))}{a^4 d} + \frac{\ln(\tan(c+dx) + 1i)(B + A 1i) 1i}{32 a^4 d} - \frac{\ln(\tan(c+dx) - i)(31A + B 1i)}{32 a^4 d}$$

input `int((cot(c + d*x)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^4,x)`

output `((7*A)/(4*a^4) - tan(c + d*x)^3*((A*15i)/(16*a^4) - B/(16*a^4)) - tan(c + d*x)^2*((13*A)/(4*a^4) + (B*1i)/(4*a^4)) + (B*1i)/(3*a^4) + tan(c + d*x)*((A*63i)/(16*a^4) - (19*B)/(48*a^4)))/(d*(tan(c + d*x)*4i - 6*tan(c + d*x)^2 - tan(c + d*x)^3*4i + tan(c + d*x)^4 + 1)) + (A*log(tan(c + d*x)))/(a^4*d) + (log(tan(c + d*x) + 1i)*(A*1i + B)*1i)/(32*a^4*d) - (log(tan(c + d*x) - 1i)*(31*A + B*1i))/(32*a^4*d)`

3.64. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

3.65 $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

3.65.1 Optimal result 787
 3.65.2 Mathematica [C] (verified) 788
 3.65.3 Rubi [A] (verified) 788
 3.65.4 Maple [A] (verified) 793
 3.65.5 Fricas [A] (verification not implemented) 794
 3.65.6 Sympy [A] (verification not implemented) 795
 3.65.7 Maxima [F(-2)] 795
 3.65.8 Giac [A] (verification not implemented) 796
 3.65.9 Mupad [B] (verification not implemented) 796

3.65.1 Optimal result

Integrand size = 34, antiderivative size = 220

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$$

$$= -\frac{5(13A+3iB)x}{16a^4} - \frac{5(13A+3iB) \cot(c+dx)}{16a^4d} - \frac{(4iA-B) \log(\sin(c+dx))}{a^4d}$$

$$+ \frac{(31A+9iB) \cot(c+dx)}{48a^4d(1+i \tan(c+dx))^2} + \frac{(4A+iB) \cot(c+dx)}{2a^4d(1+i \tan(c+dx))}$$

$$+ \frac{(A+iB) \cot(c+dx)}{8d(a+ia \tan(c+dx))^4} + \frac{(7A+3iB) \cot(c+dx)}{24ad(a+ia \tan(c+dx))^3}$$

```
output -5/16*(13*A+3*I*B)*x/a^4-5/16*(13*A+3*I*B)*cot(d*x+c)/a^4/d-(4*I*A-B)*ln(sin(d*x+c))/a^4/d+1/48*(31*A+9*I*B)*cot(d*x+c)/a^4/d/(1+I*tan(d*x+c))^2+1/2*(4*A+I*B)*cot(d*x+c)/a^4/d/(1+I*tan(d*x+c))+1/8*(A+I*B)*cot(d*x+c)/d/(a+I*a*tan(d*x+c))^4+1/24*(7*A+3*I*B)*cot(d*x+c)/a/d/(a+I*a*tan(d*x+c))^3
```

3.65.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 3.58 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.87

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$$

$$= \frac{6(A+iB) \cot^5(c+dx)}{(i+\cot(c+dx))^4} + \frac{2(7A+3iB) \cot^4(c+dx)}{(i+\cot(c+dx))^3} + \frac{(31A+9iB) \cot^3(c+dx)}{(i+\cot(c+dx))^2} + \frac{24(4A+iB) \cot^2(c+dx)}{i+\cot(c+dx)} - 15(13A+3iB) \cot(c+dx) + 48 \dots$$

input `Integrate[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]`

output `((6*(A + I*B)*Cot[c + d*x]^5)/(I + Cot[c + d*x])^4 + (2*(7*A + (3*I)*B)*Cot[c + d*x]^4)/(I + Cot[c + d*x])^3 + ((31*A + (9*I)*B)*Cot[c + d*x]^3)/(I + Cot[c + d*x])^2 + (24*(4*A + I*B)*Cot[c + d*x]^2)/(I + Cot[c + d*x]) - 15*(13*A + (3*I)*B)*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2] + 48*((-4*I)*A + B)*(Log[Cos[c + d*x]] + Log[Tan[c + d*x]]))/(48*a^4*d)`

3.65.3 Rubi [A] (verified)

Time = 1.57 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.10, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.559$, Rules used = {3042, 4079, 3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A+B \tan(c+dx)}{\tan(c+dx)^2(a+ia \tan(c+dx))^4} dx$$

$$\downarrow \text{4079}$$

$$\frac{\int \frac{\cot^2(c+dx)(a(9A+iB)-5a(iA-B) \tan(c+dx))}{(i \tan(c+dx)a+a)^3} dx}{8a^2} + \frac{(A+iB) \cot(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

3.65. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

$$\begin{aligned}
& \int \frac{a(9A+iB)-5a(iA-B)\tan(c+dx)}{\tan(c+dx)^2(i\tan(c+dx)a+a)^3} dx + \frac{(A+iB)\cot(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{4\cot^2(c+dx)(a^2(17A+3iB)-2a^2(7iA-3B)\tan(c+dx))}{(i\tan(c+dx)a+a)^2} dx}{8a^2} + \frac{a(7A+3iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^3} + \frac{(A+iB)\cot(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 4079 \\
& \frac{2\int \frac{\cot^2(c+dx)(a^2(17A+3iB)-2a^2(7iA-3B)\tan(c+dx))}{(i\tan(c+dx)a+a)^2} dx}{8a^2} + \frac{a(7A+3iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^3} + \frac{(A+iB)\cot(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 27 \\
& \frac{2\int \frac{a^2(17A+3iB)-2a^2(7iA-3B)\tan(c+dx)}{\tan(c+dx)^2(i\tan(c+dx)a+a)^2} dx}{8a^2} + \frac{a(7A+3iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^3} + \frac{(A+iB)\cot(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 3042 \\
& \frac{2\int \frac{a^2(17A+3iB)-2a^2(7iA-3B)\tan(c+dx)}{\tan(c+dx)^2(i\tan(c+dx)a+a)^2} dx}{8a^2} + \frac{a(7A+3iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^3} + \frac{(A+iB)\cot(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 4079 \\
& \frac{2\left(\int \frac{3\cot^2(c+dx)(a^3(33A+7iB)-a^3(31iA-9B)\tan(c+dx))}{i\tan(c+dx)a+a} dx + \frac{(31A+9iB)\cot(c+dx)}{4d(1+i\tan(c+dx))^2}\right)}{3a^2} + \frac{a(7A+3iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^3} + \\
& \quad \frac{8a^2}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 27 \\
& \frac{2\left(\int \frac{3\cot^2(c+dx)(a^3(33A+7iB)-a^3(31iA-9B)\tan(c+dx))}{i\tan(c+dx)a+a} dx + \frac{(31A+9iB)\cot(c+dx)}{4d(1+i\tan(c+dx))^2}\right)}{3a^2} + \frac{a(7A+3iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^3} + \\
& \quad \frac{8a^2}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 3042 \\
& \frac{2\left(\int \frac{3\int \frac{a^3(33A+7iB)-a^3(31iA-9B)\tan(c+dx)}{\tan(c+dx)^2(i\tan(c+dx)a+a)} dx + \frac{(31A+9iB)\cot(c+dx)}{4d(1+i\tan(c+dx))^2}}{4a^2} dx + \frac{(31A+9iB)\cot(c+dx)}{4d(1+i\tan(c+dx))^2}\right)}{3a^2} + \frac{a(7A+3iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^3} + \\
& \quad \frac{8a^2}{8d(a+ia\tan(c+dx))^4}
\end{aligned}$$

3.65. $\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$

↓ 4079

$$2 \left(\frac{3 \left(\frac{\int 2 \cot^2(c+dx) (5a^4(13A+3iB) - 16a^4(4iA-B) \tan(c+dx)) dx}{2a^2} + \frac{8a^3(4A+iB) \cot(c+dx)}{d(a+ia \tan(c+dx))} \right)}{4a^2} + \frac{(31A+9iB) \cot(c+dx)}{4d(1+i \tan(c+dx))^2} \right)}{3a^2} + \frac{a(7A+3iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))^3} + \frac{8a^2 (A+iB) \cot(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 27

$$2 \left(\frac{3 \left(\frac{\int \cot^2(c+dx) (5a^4(13A+3iB) - 16a^4(4iA-B) \tan(c+dx)) dx}{a^2} + \frac{8a^3(4A+iB) \cot(c+dx)}{d(a+ia \tan(c+dx))} \right)}{4a^2} + \frac{(31A+9iB) \cot(c+dx)}{4d(1+i \tan(c+dx))^2} \right)}{3a^2} + \frac{a(7A+3iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))^3} + \frac{8a^2 (A+iB) \cot(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 3042

$$2 \left(\frac{3 \left(\frac{\int \frac{5a^4(13A+3iB) - 16a^4(4iA-B) \tan(c+dx)}{\tan(c+dx)^2} dx}{4a^2} + \frac{8a^3(4A+iB) \cot(c+dx)}{d(a+ia \tan(c+dx))} \right)}{3a^2} + \frac{(31A+9iB) \cot(c+dx)}{4d(1+i \tan(c+dx))^2} \right)}{3a^2} + \frac{a(7A+3iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))^3} + \frac{8a^2 (A+iB) \cot(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 4012

$$2 \left(\frac{3 \left(\frac{\int -\cot(c+dx) (16(4iA-B)a^4 + 5(13A+3iB) \tan(c+dx)a^4) dx - 5a^4(13A+3iB) \cot(c+dx)}{a^2} + \frac{8a^3(4A+iB) \cot(c+dx)}{d(a+ia \tan(c+dx))} \right)}{4a^2} + \frac{(31A+9iB) \cot(c+dx)}{4d(1+i \tan(c+dx))^2} \right)}{3a^2} + \frac{a(7A+3iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))^3} + \frac{8a^2 (A+iB) \cot(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 25

3.65. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

$$2 \left(\frac{3 \left(\frac{-\int \cot(c+dx) (16(4iA-B)a^4 + 5(13A+3iB) \tan(c+dx)a^4) dx - 5a^4(13A+3iB) \cot(c+dx)}{a^2} + \frac{8a^3(4A+iB) \cot(c+dx)}{d(a+ia \tan(c+dx))} \right)}{4a^2} + \frac{(31A+9iB) \cot(c+dx)}{4d(1+i \tan(c+dx))^2} \right) + \frac{a(7A+3iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))} + \frac{8a^2}{3a^2} \frac{(A+iB) \cot(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 3042

$$2 \left(\frac{3 \left(\frac{-\int \frac{16(4iA-B)a^4 + 5(13A+3iB) \tan(c+dx)a^4}{\tan(c+dx)} dx - 5a^4(13A+3iB) \cot(c+dx)}{a^2} + \frac{8a^3(4A+iB) \cot(c+dx)}{d(a+ia \tan(c+dx))} \right)}{4a^2} + \frac{(31A+9iB) \cot(c+dx)}{4d(1+i \tan(c+dx))^2} \right) + \frac{a(7A+3iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))} + \frac{8a^2}{3a^2} \frac{(A+iB) \cot(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 4014

$$2 \left(\frac{3 \left(\frac{-16a^4(-B+4iA) \int \cot(c+dx) dx - 5a^4(13A+3iB) \cot(c+dx) - 5a^4x(13A+3iB)}{a^2} + \frac{8a^3(4A+iB) \cot(c+dx)}{d(a+ia \tan(c+dx))} \right)}{4a^2} + \frac{(31A+9iB) \cot(c+dx)}{4d(1+i \tan(c+dx))^2} \right) + \frac{a(7A+3iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))} + \frac{8a^2}{3a^2} \frac{(A+iB) \cot(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 3042

$$2 \left(\frac{3 \left(\frac{-16a^4(-B+4iA) \int -\tan(c+dx + \frac{\pi}{2}) dx - 5a^4(13A+3iB) \cot(c+dx) - 5a^4x(13A+3iB)}{a^2} + \frac{8a^3(4A+iB) \cot(c+dx)}{d(a+ia \tan(c+dx))} \right)}{4a^2} + \frac{(31A+9iB) \cot(c+dx)}{4d(1+i \tan(c+dx))^2} \right) + \frac{a(7A+3iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))} + \frac{8a^2}{3a^2} \frac{(A+iB) \cot(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 25

3.65. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

$$\begin{aligned}
 & \left(\frac{3 \left(\frac{16a^4(-B+4iA) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx - \frac{5a^4(13A+3iB) \cot(c+dx) - 5a^4x(13A+3iB)}{a^2} + \frac{8a^3(4A+iB) \cot(c+dx)}{d(a+ia \tan(c+dx))} \right)}{4a^2} + \frac{(31A+9iB) \cot(c+dx)}{4d(1+i \tan(c+dx))^2} \right)}{3a^2} + \frac{a(7A+3iB)}{3d(a+ia \tan(c+dx))} \\
 & \frac{(A+iB) \cot(c+dx)}{8d(a+ia \tan(c+dx))^4} \cdot 8a^2 \\
 & \quad \downarrow \text{3956} \\
 & \left(\frac{3 \left(\frac{8a^3(4A+iB) \cot(c+dx)}{d(a+ia \tan(c+dx))} + \frac{-\frac{5a^4(13A+3iB) \cot(c+dx) - 16a^4(-B+4iA) \log(-\sin(c+dx)) - 5a^4x(13A+3iB)}{a^2}}{4a^2} \right)}{3a^2} + \frac{(31A+9iB) \cot(c+dx)}{4d(1+i \tan(c+dx))^2} \right)}{3a^2} + \frac{a(7A+3iB)}{3d(a+ia \tan(c+dx))} \\
 & \frac{(A+iB) \cot(c+dx)}{8d(a+ia \tan(c+dx))^4} \cdot 8a^2
 \end{aligned}$$

input `Int[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]`

output `((A + I*B)*Cot[c + d*x])/(8*d*(a + I*a*Tan[c + d*x])^4) + ((a*(7*A + (3*I)*B)*Cot[c + d*x])/(3*d*(a + I*a*Tan[c + d*x])^3) + (2*((((31*A + (9*I)*B)*Cot[c + d*x])/(4*d*(1 + I*Tan[c + d*x])^2) + (3*((-5*a^4*(13*A + (3*I)*B)*x - (5*a^4*(13*A + (3*I)*B)*Cot[c + d*x])/d - (16*a^4*((4*I)*A - B)*Log[-Sin[c + d*x]])/d)/a^2 + (8*a^3*(4*A + I*B)*Cot[c + d*x])/(d*(a + I*a*Tan[c + d*x])))))/(4*a^2)))/(3*a^2))/(8*a^2)`

3.65.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.65. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4079 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.65.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.14

method	result
risch	$-\frac{ie^{-6i(dx+c)}A}{12da^4} - \frac{129xA}{16a^4} + \frac{13e^{-2i(dx+c)}B}{16da^4} - \frac{15ie^{-4i(dx+c)}A}{32da^4} + \frac{e^{-4i(dx+c)}B}{4da^4} - \frac{2iBc}{da^4} + \frac{e^{-6i(dx+c)}B}{16da^4} - \dots$
derivativedivides	$-\frac{15iB}{16da^4(\tan(dx+c)-i)} + \frac{B}{8da^4(\tan(dx+c)-i)^4} - \frac{B \ln(1+\tan^2(dx+c))}{2da^4} + \frac{17iA}{16da^4(\tan(dx+c)-i)^2} + \frac{i}{4da^4(\tan(dx+c)-i)}$
default	$-\frac{15iB}{16da^4(\tan(dx+c)-i)} + \frac{B}{8da^4(\tan(dx+c)-i)^4} - \frac{B \ln(1+\tan^2(dx+c))}{2da^4} + \frac{17iA}{16da^4(\tan(dx+c)-i)^2} + \frac{i}{4da^4(\tan(dx+c)-i)}$

3.65. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

input `int(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/12*I/d/a^4*\exp(-6*I*(d*x+c))*A-129/16*x/a^4*A+13/16/d/a^4*\exp(-2*I*(d*x+c))*B-15/32*I/d/a^4*\exp(-4*I*(d*x+c))*A+1/4/d/a^4*\exp(-4*I*(d*x+c))*B-2*I/a^4/d*B*c+1/16/d/a^4*\exp(-6*I*(d*x+c))*B-1/128*I/d/a^4*\exp(-8*I*(d*x+c))*A+1/128/d/a^4*\exp(-8*I*(d*x+c))*B-2*I*A/a^4/d/(\exp(2*I*(d*x+c))-1)-9/4*I/d/a^4*\exp(-2*I*(d*x+c))*A-4*I/a^4/d*\ln(\exp(2*I*(d*x+c))-1)*A-8/a^4/d*A*c-31/16*I*x/a^4*B+1/a^4/d*\ln(\exp(2*I*(d*x+c))-1)*B \end{aligned}$$

3.65.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.86

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \frac{24(129A+31iB)dx e^{(10i dx+10i c)} - 24((129A+31iB)dx - 68iA + 13B)e^{(8i dx+8i c)} + 36(-19iA + 6iB)e^{(6i dx+6i c)} + 4*(-37iA + 18B)e^{(4i dx+4i c)} - (29iA - 21B)e^{(2i dx+2i c)} + 384((4iA - B)e^{(10i dx+10i c)} + (-4iA + B)e^{(8i dx+8i c)}) * \log(e^{(2i dx+2i c)} - 1) - 3iA + 3B}{a^4 d e^{(10i dx+10i c)} - a^4 d e^{(8i dx+8i c)}}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="fricas")`

output
$$\begin{aligned} & -1/384*(24*(129*A + 31*I*B)*d*x*e^{(10*I*d*x + 10*I*c)} - 24*((129*A + 31*I*B)*d*x - 68*I*A + 13*B)*e^{(8*I*d*x + 8*I*c)} + 36*(-19*I*A + 6*B)*e^{(6*I*d*x + 6*I*c)} + 4*(-37*I*A + 18*B)*e^{(4*I*d*x + 4*I*c)} - (29*I*A - 21*B)*e^{(2*I*d*x + 2*I*c)} + 384*((4*I*A - B)*e^{(10*I*d*x + 10*I*c)} + (-4*I*A + B)*e^{(8*I*d*x + 8*I*c)}) * \log(e^{(2*I*d*x + 2*I*c)} - 1) - 3*I*A + 3*B)/(a^4*d*e^{(10*I*d*x + 10*I*c)} - a^4*d*e^{(8*I*d*x + 8*I*c)}) \end{aligned}$$

3.65.6 Sympy [A] (verification not implemented)

Time = 0.91 (sec) , antiderivative size = 406, normalized size of antiderivative = 1.85

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = -\frac{2iA}{a^4 d e^{2ic} e^{2idx} - a^4 d} + \left\{ \frac{((-24576iAa^{12}d^3e^{12ic}+24576Ba^{12}d^3e^{12ic})e^{-8idx}+(-262144iAa^{12}d^3e^{14ic}+196608Ba^{12}d^3e^{14ic})e^{-6idx}+(-1474560iAa^{12}d^3e^{16ic}+786432Ba^{12}d^3e^{16ic})e^{-4idx}+(-7077888iAa^{12}d^3e^{18ic}+2555904Ba^{12}d^3e^{18ic})e^{-2idx})e^{-20ic}}{3145728a^{16}d^4} \right. \\ \left. + x \left(-\frac{129A-31iB}{16a^4} + \frac{(-129Ae^{8ic}-72Ae^{6ic}-30Ae^{4ic}-8Ae^{2ic}-A-31iBe^{8ic}-26iBe^{6ic}-16iBe^{4ic}-6iBe^{2ic}-iB)e^{-8ic}}{16a^4} \right) \right. \\ \left. + \frac{x(-129A-31iB)}{16a^4} - \frac{i(4A+iB) \log(e^{2idx} - e^{-2ic})}{a^4 d} \right.$$

input `integrate(cot(d*x+c)**2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**4,x)`

output `-2*I*A/(a**4*d*exp(2*I*c)*exp(2*I*d*x) - a**4*d) + Piecewise(((((-24576*I*A*a**12*d**3*exp(12*I*c) + 24576*B*a**12*d**3*exp(12*I*c))*exp(-8*I*d*x) + (-262144*I*A*a**12*d**3*exp(14*I*c) + 196608*B*a**12*d**3*exp(14*I*c))*exp(-6*I*d*x) + (-1474560*I*A*a**12*d**3*exp(16*I*c) + 786432*B*a**12*d**3*exp(16*I*c))*exp(-4*I*d*x) + (-7077888*I*A*a**12*d**3*exp(18*I*c) + 2555904*B*a**12*d**3*exp(18*I*c))*exp(-2*I*d*x))*exp(-20*I*c)/(3145728*a**16*d**4), Ne(a**16*d**4*exp(20*I*c), 0)), (x*(-(-129*A - 31*I*B)/(16*a**4) + (-129*A*exp(8*I*c) - 72*A*exp(6*I*c) - 30*A*exp(4*I*c) - 8*A*exp(2*I*c) - A - 31*I*B*exp(8*I*c) - 26*I*B*exp(6*I*c) - 16*I*B*exp(4*I*c) - 6*I*B*exp(2*I*c) - I*B)*exp(-8*I*c)/(16*a**4)), True)) + x*(-129*A - 31*I*B)/(16*a**4) - I*(4*A + I*B)*log(exp(2*I*d*x) - exp(-2*I*c))/(a**4*d)`

3.65.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.65.8 Giac [A] (verification not implemented)

Time = 1.29 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.93

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \frac{12(-iA-B) \log(\tan(dx+c)+i)}{a^4} - \frac{12(-129iA+31B) \log(\tan(dx+c)-i)}{a^4} - \frac{384(4iA-B) \log(\tan(dx+c))}{a^4} - \frac{384(-4iA \tan(dx+c)+B \tan(dx+c))}{a^4 \tan(dx+c)}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="giac")`

output `1/384*(12*(-I*A - B)*log(tan(d*x + c) + I)/a^4 - 12*(-129*I*A + 31*B)*log(tan(d*x + c) - I)/a^4 - 384*(4*I*A - B)*log(tan(d*x + c))/a^4 - 384*(-4*I*A*tan(d*x + c) + B*tan(d*x + c) + A)/(a^4*tan(d*x + c)) - (3225*I*A*tan(d*x + c)^4 - 775*B*tan(d*x + c)^4 + 14076*A*tan(d*x + c)^3 + 3460*I*B*tan(d*x + c)^3 - 23286*I*A*tan(d*x + c)^2 + 5898*B*tan(d*x + c)^2 - 17404*A*tan(d*x + c) - 4612*I*B*tan(d*x + c) + 5017*I*A - 1447*B)/(a^4*(tan(d*x + c) - I)^4))/d`

3.65.9 Mupad [B] (verification not implemented)

Time = 8.15 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.03

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \frac{\frac{A}{a^4} + \tan(c+dx)^4 \left(\frac{65A}{16a^4} + \frac{B15i}{16a^4} \right) - \tan(c+dx)^2 \left(\frac{851A}{48a^4} + \frac{B63i}{16a^4} \right) - \tan(c+dx)^3 \left(-\frac{13B}{4a^4} + \frac{A57i}{4a^4} \right) + \tan(c+dx)}{d \left(\tan(c+dx)^5 - \tan(c+dx)^4 4i - 6 \tan(c+dx)^3 + \tan(c+dx)^2 4i + \tan(c+dx) \right)} - \frac{\ln(\tan(c+dx))(-B+4Ai)}{a^4 d} - \frac{\ln(\tan(c+dx)+1i)(B+4Ai)}{32a^4 d} + \frac{\ln(\tan(c+dx)-i)(-31B+A129i)}{32a^4 d}$$

input `int((cot(c + d*x)^2*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^4,x)`

output $(\log(\tan(c + dx) - 1i)(A*129i - 31*B))/(32*a^4*d) - (\log(\tan(c + dx))*(A*4i - B))/(a^4*d) - (\log(\tan(c + dx) + 1i)(A*1i + B))/(32*a^4*d) - (\tan(c + dx)^4*((65*A)/(16*a^4) + (B*15i)/(16*a^4)) - \tan(c + dx)^3*((A*57i)/(4*a^4) - (13*B)/(4*a^4)) - \tan(c + dx)^2*((851*A)/(48*a^4) + (B*63i)/(16*a^4)) + A/a^4 + \tan(c + dx)*((A*26i)/(3*a^4) - (7*B)/(4*a^4)))/(d*(\tan(c + dx) + \tan(c + dx)^2*4i - 6*\tan(c + dx)^3 - \tan(c + dx)^4*4i + \tan(c + dx)^5))$

3.66 $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

3.66.1	Optimal result	798
3.66.2	Mathematica [C] (verified)	799
3.66.3	Rubi [A] (verified)	799
3.66.4	Maple [A] (verified)	805
3.66.5	Fricas [A] (verification not implemented)	805
3.66.6	Sympy [A] (verification not implemented)	806
3.66.7	Maxima [F(-2)]	807
3.66.8	Giac [A] (verification not implemented)	807
3.66.9	Mupad [B] (verification not implemented)	808

3.66.1 Optimal result

Integrand size = 34, antiderivative size = 255

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$$

$$= \frac{5(35iA-13B)x}{16a^4} + \frac{5(35iA-13B) \cot(c+dx)}{16a^4d} - \frac{(11A+4iB) \cot^2(c+dx)}{2a^4d}$$

$$- \frac{(11A+4iB) \log(\sin(c+dx))}{a^4d} + \frac{(43A+17iB) \cot^2(c+dx)}{48a^4d(1+i \tan(c+dx))^2}$$

$$+ \frac{5(35A+13iB) \cot^2(c+dx)}{48a^4d(1+i \tan(c+dx))} + \frac{(A+iB) \cot^2(c+dx)}{8d(a+ia \tan(c+dx))^4} + \frac{(2A+iB) \cot^2(c+dx)}{6ad(a+ia \tan(c+dx))^3}$$

```
output 5/16*(35*I*A-13*B)*x/a^4+5/16*(35*I*A-13*B)*cot(d*x+c)/a^4/d-1/2*(11*A+4*I
*B)*cot(d*x+c)^2/a^4/d-(11*A+4*I*B)*ln(sin(d*x+c))/a^4/d+1/48*(43*A+17*I*B
)*cot(d*x+c)^2/a^4/d/(1+I*tan(d*x+c))^2+5/48*(35*A+13*I*B)*cot(d*x+c)^2/a^
4/d/(1+I*tan(d*x+c))+1/8*(A+I*B)*cot(d*x+c)^2/d/(a+I*a*tan(d*x+c))^4+1/6*(
2*A+I*B)*cot(d*x+c)^2/a/d/(a+I*a*tan(d*x+c))^3
```

3.66.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 4.60 (sec) , antiderivative size = 204, normalized size of antiderivative = 0.80

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$$

$$= \frac{6(A+iB) \cot^6(c+dx)}{(i+\cot(c+dx))^4} + \frac{8(2A+iB) \cot^5(c+dx)}{(i+\cot(c+dx))^3} + \frac{(43A+17iB) \cot^4(c+dx)}{(i+\cot(c+dx))^2} + \frac{5(35A+13iB) \cot^3(c+dx)}{i+\cot(c+dx)} + 15(35iA-13B) \cot(c+dx)$$

input `Integrate[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]`

output `((6*(A + I*B)*Cot[c + d*x]^6)/(I + Cot[c + d*x])^4 + (8*(2*A + I*B)*Cot[c + d*x]^5)/(I + Cot[c + d*x])^3 + ((43*A + (17*I)*B)*Cot[c + d*x]^4)/(I + Cot[c + d*x])^2 + (5*(35*A + (13*I)*B)*Cot[c + d*x]^3)/(I + Cot[c + d*x]) + 15*((35*I)*A - 13*B)*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2 - 24*(11*A + (4*I)*B)*(Cot[c + d*x]^2 + 2*(Log[Cos[c + d*x]] + Log[Tan[c + d*x]])))/(48*a^4*d)`

3.66.3 Rubi [A] (verified)

Time = 1.88 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.09, number of steps used = 22, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.647$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4012, 25, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A+B \tan(c+dx)}{\tan(c+dx)^3(a+ia \tan(c+dx))^4} dx$$

$$\downarrow \text{4079}$$

$$\frac{\int \frac{2 \cot^3(c+dx)(a(5A+iB)-3a(iA-B) \tan(c+dx))}{(i \tan(c+dx)a+a)^3} dx}{8a^2} + \frac{(A+iB) \cot^2(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

3.66. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

$$\begin{aligned}
& \int \frac{\cot^3(c+dx)(a(5A+iB)-3a(iA-B)\tan(c+dx))}{(i\tan(c+dx)a+a)^3} dx + \frac{(A+iB)\cot^2(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 27 \\
& \int \frac{a(5A+iB)-3a(iA-B)\tan(c+dx)}{\tan(c+dx)^3(i\tan(c+dx)a+a)^3} dx + \frac{(A+iB)\cot^2(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 3042 \\
& \int \frac{2\cot^3(c+dx)(a^2(23A+7iB)-10a^2(2iA-B)\tan(c+dx))}{(i\tan(c+dx)a+a)^2} dx + \frac{2a(2A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^3} + \frac{(A+iB)\cot^2(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 4079 \\
& \int \frac{\cot^3(c+dx)(a^2(23A+7iB)-10a^2(2iA-B)\tan(c+dx))}{(i\tan(c+dx)a+a)^2} dx + \frac{2a(2A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^3} + \frac{(A+iB)\cot^2(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 27 \\
& \int \frac{a^2(23A+7iB)-10a^2(2iA-B)\tan(c+dx)}{\tan(c+dx)^3(i\tan(c+dx)a+a)^2} dx + \frac{2a(2A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^3} + \frac{(A+iB)\cot^2(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 3042 \\
& \int \frac{a^2(23A+7iB)-10a^2(2iA-B)\tan(c+dx)}{\tan(c+dx)^3(i\tan(c+dx)a+a)^2} dx + \frac{2a(2A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^3} + \frac{(A+iB)\cot^2(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 4079 \\
& \int \frac{2\cot^3(c+dx)(a^3(89A+31iB)-2a^3(43iA-17B)\tan(c+dx))}{i\tan(c+dx)a+a} dx + \frac{(43A+17iB)\cot^2(c+dx)}{4d(1+i\tan(c+dx))^2} + \frac{2a(2A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^3} + \\
& \quad \frac{4a^2}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 27 \\
& \int \frac{\cot^3(c+dx)(a^3(89A+31iB)-2a^3(43iA-17B)\tan(c+dx))}{i\tan(c+dx)a+a} dx + \frac{(43A+17iB)\cot^2(c+dx)}{4d(1+i\tan(c+dx))^2} + \frac{2a(2A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^3} + \\
& \quad \frac{4a^2}{8d(a+ia\tan(c+dx))^4} \\
& \quad \downarrow 3042
\end{aligned}$$

3.66. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$

$$\frac{\int \frac{a^3(89A+31iB)-2a^3(43iA-17B)\tan(c+dx)}{\tan(c+dx)^3(i\tan(c+dx)a+a)} dx + \frac{(43A+17iB)\cot^2(c+dx)}{4d(1+i\tan(c+dx))^2} + \frac{2a(2A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^3}}{3a^2} + \frac{4a^2(A+iB)\cot^2(c+dx)}{8d(a+ia\tan(c+dx))^4}$$

↓ 4079

$$\frac{\int 3\cot^3(c+dx)\left(\frac{16a^4(11A+4iB)-5a^4(35iA-13B)\tan(c+dx)}{2a^2}\right) dx + \frac{5a^3(35A+13iB)\cot^2(c+dx)}{2d(a+ia\tan(c+dx))} + \frac{(43A+17iB)\cot^2(c+dx)}{4d(1+i\tan(c+dx))^2} + \frac{2a(2A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^3}}{3a^2} + \frac{4a^2(A+iB)\cot^2(c+dx)}{8d(a+ia\tan(c+dx))^4}$$

↓ 27

$$\frac{3\int \frac{16a^4(11A+4iB)-5a^4(35iA-13B)\tan(c+dx)}{2a^2} dx + \frac{5a^3(35A+13iB)\cot^2(c+dx)}{2d(a+ia\tan(c+dx))} + \frac{(43A+17iB)\cot^2(c+dx)}{4d(1+i\tan(c+dx))^2} + \frac{2a(2A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^3}}{3a^2} + \frac{4a^2(A+iB)\cot^2(c+dx)}{8d(a+ia\tan(c+dx))^4}$$

↓ 3042

$$\frac{3\int \frac{16a^4(11A+4iB)-5a^4(35iA-13B)\tan(c+dx)}{2a^2} dx + \frac{5a^3(35A+13iB)\cot^2(c+dx)}{2d(a+ia\tan(c+dx))} + \frac{(43A+17iB)\cot^2(c+dx)}{4d(1+i\tan(c+dx))^2} + \frac{2a(2A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^3}}{3a^2} + \frac{4a^2(A+iB)\cot^2(c+dx)}{8d(a+ia\tan(c+dx))^4}$$

↓ 4012

$$\frac{3\left(\int -\cot^2(c+dx)\left(5(35iA-13B)a^4+16(11A+4iB)\tan(c+dx)a^4\right) dx - \frac{8a^4(11A+4iB)\cot^2(c+dx)}{d}\right) + \frac{5a^3(35A+13iB)\cot^2(c+dx)}{2d(a+ia\tan(c+dx))} + \frac{(43A+17iB)\cot^2(c+dx)}{4d(1+i\tan(c+dx))^2}}{3a^2} + \frac{4a^2(A+iB)\cot^2(c+dx)}{8d(a+ia\tan(c+dx))^4}$$

↓ 25

3.66. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$

$$\frac{3 \left(- \int \cot^2(c+dx) \left(5(35iA-13B)a^4 + 16(11A+4iB) \tan(c+dx)a^4 \right) dx - \frac{8a^4(11A+4iB) \cot^2(c+dx)}{d} \right)}{2a^2} + \frac{5a^3(35A+13iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{(43A+17iB) \cot^2(c+dx)}{4d(1+i \tan(c+dx))^2} + \dots$$

$$\frac{(A+iB) \cot^2(c+dx)}{8d(a+ia \tan(c+dx))^4} \quad 4a^2$$

↓ 3042

$$\frac{3 \left(- \int \frac{5(35iA-13B)a^4 + 16(11A+4iB) \tan(c+dx)a^4}{\tan(c+dx)^2} dx - \frac{8a^4(11A+4iB) \cot^2(c+dx)}{d} \right)}{2a^2} + \frac{5a^3(35A+13iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{(43A+17iB) \cot^2(c+dx)}{4d(1+i \tan(c+dx))^2} + \frac{2a(2A+iB)}{3d(a+ia \tan(c+dx))} + \dots$$

$$\frac{(A+iB) \cot^2(c+dx)}{8d(a+ia \tan(c+dx))^4} \quad 4a^2$$

↓ 4012

$$\frac{3 \left(- \int \cot(c+dx) \left(16a^4(11A+4iB) - 5a^4(35iA-13B) \tan(c+dx) \right) dx - \frac{8a^4(11A+4iB) \cot^2(c+dx)}{d} + \frac{5a^4(-13B+35iA) \cot(c+dx)}{d} \right)}{2a^2} + \frac{5a^3(35A+13iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \dots$$

$$\frac{(A+iB) \cot^2(c+dx)}{8d(a+ia \tan(c+dx))^4} \quad 4a^2$$

↓ 3042

$$\frac{3 \left(- \int \frac{16a^4(11A+4iB) - 5a^4(35iA-13B) \tan(c+dx)}{\tan(c+dx)} dx - \frac{8a^4(11A+4iB) \cot^2(c+dx)}{d} + \frac{5a^4(-13B+35iA) \cot(c+dx)}{d} \right)}{2a^2} + \frac{5a^3(35A+13iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{(43A+17iB) \cot^2(c+dx)}{4d(1+i \tan(c+dx))^2} + \dots$$

$$\frac{(A+iB) \cot^2(c+dx)}{8d(a+ia \tan(c+dx))^4} \quad 4a^2$$

↓ 4014

$$\frac{3 \left(-16a^4(11A+4iB) \int \cot(c+dx) dx - \frac{8a^4(11A+4iB) \cot^2(c+dx)}{d} + \frac{5a^4(-13B+35iA) \cot(c+dx)}{d} + 5a^4 x(-13B+35iA) \right)}{2a^2} + \frac{5a^3(35A+13iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{(43A+17iB) \cot^2(c+dx)}{4d(1+i \tan(c+dx))^2} + \dots$$

$$\frac{(A+iB) \cot^2(c+dx)}{8d(a+ia \tan(c+dx))^4} \quad 4a^2$$

↓ 3042

3.66. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

$$\begin{aligned}
 & \frac{3 \left(-16a^4(11A+4iB) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx - \frac{8a^4(11A+4iB) \cot^2(c+dx)}{d} + \frac{5a^4(-13B+35iA) \cot(c+dx)}{d} + 5a^4x(-13B+35iA) \right)}{2a^2} + \frac{5a^3(35A+13iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{(43A+17iB) \cot(c+dx)}{4d} \\
 & \frac{(A+iB) \cot^2(c+dx)}{8d(a+ia \tan(c+dx))^4} \\
 & \quad \downarrow 25 \\
 & \frac{3 \left(16a^4(11A+4iB) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx - \frac{8a^4(11A+4iB) \cot^2(c+dx)}{d} + \frac{5a^4(-13B+35iA) \cot(c+dx)}{d} + 5a^4x(-13B+35iA) \right)}{2a^2} + \frac{5a^3(35A+13iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{(43A+17iB) \cot(c+dx)}{4d} \\
 & \frac{(A+iB) \cot^2(c+dx)}{8d(a+ia \tan(c+dx))^4} \\
 & \quad \downarrow 3956 \\
 & \frac{5a^3(35A+13iB) \cot^2(c+dx)}{2d(a+ia \tan(c+dx))} + \frac{3 \left(-\frac{8a^4(11A+4iB) \cot^2(c+dx)}{d} + \frac{5a^4(-13B+35iA) \cot(c+dx)}{d} - \frac{16a^4(11A+4iB) \log(-\sin(c+dx))}{d} + 5a^4x(-13B+35iA) \right)}{2a^2} + \frac{(43A+17iB) \cot(c+dx)}{4d} \\
 & \frac{(A+iB) \cot^2(c+dx)}{8d(a+ia \tan(c+dx))^4}
 \end{aligned}$$

```
input Int[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]
```

```
output ((A + I*B)*Cot[c + d*x]^2)/(8*d*(a + I*a*Tan[c + d*x])^4) + ((2*a*(2*A + I*B)*Cot[c + d*x]^2)/(3*d*(a + I*a*Tan[c + d*x])^3) + (((43*A + (17*I)*B)*Cot[c + d*x]^2)/(4*d*(1 + I*Tan[c + d*x])^2) + ((3*(5*a^4*((35*I)*A - 13*B)*x + (5*a^4*((35*I)*A - 13*B)*Cot[c + d*x])/d - (8*a^4*(11*A + (4*I)*B)*Cot[c + d*x]^2)/d - (16*a^4*(11*A + (4*I)*B)*Log[-Sin[c + d*x]])/d))/(2*a^2) + (5*a^3*(35*A + (13*I)*B)*Cot[c + d*x]^2)/(2*d*(a + I*a*Tan[c + d*x])))/(2*a^2)/(3*a^2)/(4*a^2)
```

3.66. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

3.66.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`
- rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`
- rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.66.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.11

method	result
risch	$-\frac{129xB}{16a^4} - \frac{ie^{-8i(dx+c)}B}{128da^4} - \frac{2i(-3iAe^{2i(dx+c)}+Be^{2i(dx+c)}+4iA-B)}{a^4d(e^{2i(dx+c)}-1)^2} - \frac{75e^{-2i(dx+c)}A}{16da^4} + \frac{22iAc}{a^4d} - \frac{3e^{-4i(dx+c)}}{4da^4}$
derivativedivides	$-\frac{A}{8da^4(\tan(dx+c)-i)^4} + \frac{4iA}{da^4\tan(dx+c)} - \frac{49B}{16da^4(\tan(dx+c)-i)} - \frac{iB}{8da^4(\tan(dx+c)-i)^4} + \frac{31A}{16da^4(\tan(dx+c))}$
default	$-\frac{A}{8da^4(\tan(dx+c)-i)^4} + \frac{4iA}{da^4\tan(dx+c)} - \frac{49B}{16da^4(\tan(dx+c)-i)} - \frac{iB}{8da^4(\tan(dx+c)-i)^4} + \frac{31A}{16da^4(\tan(dx+c))}$

input `int(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x,method=_RETURNVERBOSE)`

output
$$-129/16*x/a^4*B-1/128*I/d/a^4*\exp(-8*I*(d*x+c))*B-2*I*(-3*I*A*\exp(2*I*(d*x+c))+B*\exp(2*I*(d*x+c))+4*I*A-B)/a^4/d/(\exp(2*I*(d*x+c))-1)^2-75/16/d/a^4*\exp(-2*I*(d*x+c))*A+22*I/a^4/d*A*c-3/4/d/a^4*\exp(-4*I*(d*x+c))*A-9/4*I/d/a^4*\exp(-2*I*(d*x+c))*B-5/48/d/a^4*\exp(-6*I*(d*x+c))*A+351/16*I*x/a^4*A-1/128/d/a^4*\exp(-8*I*(d*x+c))*A-15/32*I/d/a^4*\exp(-4*I*(d*x+c))*B-8/a^4/d*B*c-4*I/a^4/d*\ln(\exp(2*I*(d*x+c))-1)*B-1/12*I/d/a^4*\exp(-6*I*(d*x+c))*B-11/a^4*A/d*\ln(\exp(2*I*(d*x+c))-1)$$

3.66.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 253, normalized size of antiderivative = 0.99

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx = \frac{72(-117iA+43B)dx e^{(12i dx+12i c)} + 24(6(117iA-43B)dx + 171A + 68iB)e^{(10i dx+10i c)} + 12(6(-1$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="fricas")`

output
$$-1/384*(72*(-117*I*A + 43*B)*d*x*e^(12*I*d*x + 12*I*c) + 24*(6*(117*I*A - 43*B)*d*x + 171*A + 68*I*B)*e^(10*I*d*x + 10*I*c) + 12*(6*(-117*I*A + 43*B)*d*x - 532*A - 193*I*B)*e^(8*I*d*x + 8*I*c) + 8*(158*A + 67*I*B)*e^(6*I*d*x + 6*I*c) + (211*A + 119*I*B)*e^(4*I*d*x + 4*I*c) + 2*(17*A + 13*I*B)*e^(2*I*d*x + 2*I*c) + 384*((11*A + 4*I*B)*e^(12*I*d*x + 12*I*c) - 2*(11*A + 4*I*B)*e^(10*I*d*x + 10*I*c) + (11*A + 4*I*B)*e^(8*I*d*x + 8*I*c))*log(e^(2*I*d*x + 2*I*c) - 1) + 3*A + 3*I*B)/(a^4*d*e^(12*I*d*x + 12*I*c) - 2*a^4*d*e^(10*I*d*x + 10*I*c) + a^4*d*e^(8*I*d*x + 8*I*c))$$

3.66.6 Sympy [A] (verification not implemented)

Time = 0.73 (sec) , antiderivative size = 466, normalized size of antiderivative = 1.83

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \frac{8A+2iB+(-6Ae^{2ic}-2iBe^{2ic})e^{2idx}}{a^4de^{4ic}e^{4idx}-2a^4de^{2ic}e^{2idx}+a^4d} + \left\{ \frac{((-24576Aa^{12}d^3e^{12ic}-24576iBa^{12}d^3e^{12ic})e^{-8idx}+(-327680Aa^{12}d^3e^{14ic}-262144iBa^{12}d^3e^{14ic})e^{-6idx}+(-2359296Aa^{12}d^3e^{16ic}-1474560iBa^{12}d^3e^{16ic})e^{-4idx}+(-14745600Aa^{12}d^3e^{18ic}-7077888iBa^{12}d^3e^{18ic})e^{-2idx})}{3145728a^{16}d^4} \right. \\ \left. + x\left(-\frac{351iA-129B}{16a^4} + \frac{(351iAe^{8ic}+150iAe^{6ic}+48iAe^{4ic}+10iAe^{2ic}+iA-129Be^{8ic}-72Be^{6ic}-30Be^{4ic}-8Be^{2ic}-B)e^{-8ic}}{16a^4}\right) \right. \\ \left. + \frac{x(351iA-129B)}{16a^4} - \frac{(11A+4iB)\log(e^{2idx}-e^{-2ic})}{a^4d} \right.$$

input `integrate(cot(d*x+c)**3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**4,x)`

output
$$(8*A + 2*I*B + (-6*A*\exp(2*I*c) - 2*I*B*\exp(2*I*c))*\exp(2*I*d*x))/(a**4*d*\exp(4*I*c)*\exp(4*I*d*x) - 2*a**4*d*\exp(2*I*c)*\exp(2*I*d*x) + a**4*d) + \text{Piecewise}(\left(\left(\left(-24576*A*a**12*d**3*\exp(12*I*c) - 24576*I*B*a**12*d**3*\exp(12*I*c)\right)*\exp(-8*I*d*x) + \left(-327680*A*a**12*d**3*\exp(14*I*c) - 262144*I*B*a**12*d**3*\exp(14*I*c)\right)*\exp(-6*I*d*x) + \left(-2359296*A*a**12*d**3*\exp(16*I*c) - 1474560*I*B*a**12*d**3*\exp(16*I*c)\right)*\exp(-4*I*d*x) + \left(-14745600*A*a**12*d**3*\exp(18*I*c) - 7077888*I*B*a**12*d**3*\exp(18*I*c)\right)*\exp(-2*I*d*x)\right)*\exp(-20*I*c)/(3145728*a**16*d**4), \text{Ne}(a**16*d**4*\exp(20*I*c), 0)), (x*(-(351*I*A - 129*B)/(16*a**4) + (351*I*A*\exp(8*I*c) + 150*I*A*\exp(6*I*c) + 48*I*A*\exp(4*I*c) + 10*I*A*\exp(2*I*c) + I*A - 129*B*\exp(8*I*c) - 72*B*\exp(6*I*c) - 30*B*\exp(4*I*c) - 8*B*\exp(2*I*c) - B)*\exp(-8*I*c)/(16*a**4)), \text{True})) + x*(351*I*A - 129*B)/(16*a**4) - (11*A + 4*I*B)*\log(\exp(2*I*d*x) - \exp(-2*I*c))/(a**4*d)$$

3.66.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm=
"maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.66.8 Giac [A] (verification not implemented)

Time = 1.08 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.89

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$$

$$= \frac{12(A-iB)\log(\tan(dx+c)+i)}{a^4} + \frac{36(117A+43iB)\log(\tan(dx+c)-i)}{a^4} - \frac{384(11A+4iB)\log(\tan(dx+c))}{a^4} + \frac{192(33A\tan(dx+c)^2+12iB\tan(dx+c))}{a^4}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm=
"giac")
```

```
output 1/384*(12*(A - I*B)*log(tan(d*x + c) + I)/a^4 + 36*(117*A + 43*I*B)*log(ta
n(d*x + c) - I)/a^4 - 384*(11*A + 4*I*B)*log(tan(d*x + c))/a^4 + 192*(33*A
*tan(d*x + c)^2 + 12*I*B*tan(d*x + c)^2 + 8*I*A*tan(d*x + c) - 2*B*tan(d*x
+ c) - A)/(a^4*tan(d*x + c)^2) - (8775*A*tan(d*x + c)^4 + 3225*I*B*tan(d*
x + c)^4 - 37764*I*A*tan(d*x + c)^3 + 14076*B*tan(d*x + c)^3 - 61386*A*tan
(d*x + c)^2 - 23286*I*B*tan(d*x + c)^2 + 44804*I*A*tan(d*x + c) - 17404*B*
tan(d*x + c) + 12455*A + 5017*I*B)/(a^4*(tan(d*x + c) - I)^4))/d
```

3.66.9 Mupad [B] (verification not implemented)

Time = 9.06 (sec) , antiderivative size = 251, normalized size of antiderivative = 0.98

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$$

$$= \frac{\tan(c+dx)^4 \left(\frac{153A}{4a^4} + \frac{B57i}{4a^4}\right) + \tan(c+dx)^5 \left(-\frac{65B}{16a^4} + \frac{A175i}{16a^4}\right) - \tan(c+dx)^2 \left(\frac{271A}{12a^4} + \frac{B26i}{3a^4}\right) - \tan(c+dx)^3}{d(\tan(c+dx)^6 - \tan(c+dx)^5 4i - 6\tan(c+dx)^4 + \tan(c+dx)^3)}$$

$$- \frac{\ln(\tan(c+dx))(11A+B4i)}{a^4 d} + \frac{\ln(\tan(c+dx)+1i)(A-B1i)}{32a^4 d}$$

$$+ \frac{\ln(\tan(c+dx)-i)(351A+B129i)}{32a^4 d}$$

```
input int((cot(c + d*x)^3*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^4,x)
```

```
output (tan(c + d*x)^4*((153*A)/(4*a^4) + (B*57i)/(4*a^4)) + tan(c + d*x)^5*((A*175i)/(16*a^4) - (65*B)/(16*a^4)) - tan(c + d*x)^2*((271*A)/(12*a^4) + (B*26i)/(3*a^4)) - tan(c + d*x)^3*((A*2269i)/(48*a^4) - (851*B)/(48*a^4)) - A/(2*a^4) + tan(c + d*x)*((A*2i)/a^4 - B/a^4))/(d*(tan(c + d*x)^2 + tan(c + d*x)^3*4i - 6*tan(c + d*x)^4 - tan(c + d*x)^5*4i + tan(c + d*x)^6)) - (log(tan(c + d*x))*(11*A + B*4i))/(a^4*d) + (log(tan(c + d*x) + 1i)*(A - B*1i))/(32*a^4*d) + (log(tan(c + d*x) - 1i)*(351*A + B*129i))/(32*a^4*d)
```

3.67 $\int \tan^3(c+dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.67.1	Optimal result	809
3.67.2	Mathematica [A] (verified)	810
3.67.3	Rubi [A] (verified)	810
3.67.4	Maple [A] (verified)	814
3.67.5	Fricas [B] (verification not implemented)	814
3.67.6	Sympy [F]	815
3.67.7	Maxima [A] (verification not implemented)	815
3.67.8	Giac [F(-1)]	816
3.67.9	Mupad [B] (verification not implemented)	816

3.67.1 Optimal result

Integrand size = 36, antiderivative size = 194

$$\int \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\sqrt{2}\sqrt{a}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{8(7A - iB)\sqrt{a + ia \tan(c + dx)}}{35d}$$

$$+ \frac{2(7A - iB) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{35d}$$

$$+ \frac{2B \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d} - \frac{2(7A - 31iB)(a + ia \tan(c + dx))^{3/2}}{105ad}$$

output

```
(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)*a^(1/2)/d-8/35*(7*A-I*B)*(a+I*a*tan(d*x+c))^(1/2)/d+2/35*(7*A-I*B)*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^2/d+2/7*B*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^3/d-2/105*(7*A-31*I*B)*(a+I*a*tan(d*x+c))^(3/2)/a/d
```


3.67.2 Mathematica [A] (verified)

Time = 2.11 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.67

$$\int \tan^3(c+dx)\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$= \frac{\sqrt{2}\sqrt{a}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{2\sqrt{a+ia\tan(c+dx)}(-91A+43iB+(-7iA-31B)\tan(c+dx)+3(7A-iB)\tan^2(c+dx)+15B\tan^3(c+dx))}{105d}$$

input `Integrate[Tan[c + d*x]^3*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[2]*Sqrt[a]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d + (2*Sqrt[a + I*a*Tan[c + d*x]]*(-91*A + (43*I)*B + ((-7*I)*A - 31*B)*Tan[c + d*x] + 3*(7*A - I*B)*Tan[c + d*x]^2 + 15*B*Tan[c + d*x]^3))/(105*d)`

3.67.3 Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.10, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.361$, Rules used = {3042, 4080, 27, 3042, 4080, 27, 3042, 4075, 3042, 4010, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^3(c+dx)\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow 3042$$

$$\int \tan(c+dx)^3\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow 4080$$

$$\frac{2\int -\frac{1}{2}\tan^2(c+dx)\sqrt{i\tan(c+dx)a+a(6aB-a(7A-iB)\tan(c+dx))}dx}{7a} + \frac{2B\tan^3(c+dx)\sqrt{a+ia\tan(c+dx)}}{7d}$$

$$\downarrow 27$$

3.67. $\int \tan^3(c+dx)\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$

$$\begin{aligned}
 & \frac{2B \tan^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{7d} - \frac{\int \tan^2(c+dx) \sqrt{i \tan(c+dx)a+a(6aB-a(7A-iB) \tan(c+dx))} dx}{7a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2B \tan^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{7d} - \frac{\int \tan(c+dx)^2 \sqrt{i \tan(c+dx)a+a(6aB-a(7A-iB) \tan(c+dx))} dx}{7a} \\
 & \quad \downarrow \text{4080} \\
 & \frac{2B \tan^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{7d} - \frac{2 \int \frac{1}{2} \tan(c+dx) \sqrt{i \tan(c+dx)a+a(4(7A-iB)a^2+(7iA+31B) \tan(c+dx)a^2)} dx}{5a} - \frac{2a(7A-iB) \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d} \\
 & \quad \downarrow \text{27} \\
 & \frac{2B \tan^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{7d} - \frac{\int \tan(c+dx) \sqrt{i \tan(c+dx)a+a(4(7A-iB)a^2+(7iA+31B) \tan(c+dx)a^2)} dx}{5a} - \frac{2a(7A-iB) \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2B \tan^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{7d} - \frac{\int \tan(c+dx) \sqrt{i \tan(c+dx)a+a(4(7A-iB)a^2+(7iA+31B) \tan(c+dx)a^2)} dx}{5a} - \frac{2a(7A-iB) \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d} \\
 & \quad \downarrow \text{4075} \\
 & \frac{2B \tan^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{7d} - \frac{\int \sqrt{i \tan(c+dx)a+a(4a^2(7A-iB) \tan(c+dx)-a^2(7iA+31B))} dx + \frac{2a(7A-31iB)(a+ia \tan(c+dx))^{3/2}}{3d}}{5a} - \frac{2a(7A-iB) \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2B \tan^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{7d} - \frac{\int \sqrt{i \tan(c+dx)a+a(4a^2(7A-iB) \tan(c+dx)-a^2(7iA+31B))} dx + \frac{2a(7A-31iB)(a+ia \tan(c+dx))^{3/2}}{3d}}{5a} - \frac{2a(7A-iB) \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d} \\
 & \quad \downarrow \text{4010}
 \end{aligned}$$

3.67. $\int \tan^3(c+dx) \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx$

$$\frac{2B \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d} - \frac{-35a^2(B+IA) \int \sqrt{i \tan(c+dx)a+adx} + \frac{8a^2(7A-iB)\sqrt{a+ia \tan(c+dx)}}{d} + \frac{2a(7A-31iB)(a+ia \tan(c+dx))^{3/2}}{3d}}{5a} - \frac{2a(7A-iB) \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d}}{7a}$$

↓ 3042

$$\frac{2B \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d} - \frac{-35a^2(B+IA) \int \sqrt{i \tan(c+dx)a+adx} + \frac{8a^2(7A-iB)\sqrt{a+ia \tan(c+dx)}}{d} + \frac{2a(7A-31iB)(a+ia \tan(c+dx))^{3/2}}{3d}}{5a} - \frac{2a(7A-iB) \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d}}{7a}$$

↓ 3961

$$\frac{2B \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d} - \frac{70ia^3(B+IA) \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a} + \frac{8a^2(7A-iB)\sqrt{a+ia \tan(c+dx)}}{d} + \frac{2a(7A-31iB)(a+ia \tan(c+dx))^{3/2}}{3d}}{5a} - \frac{2a(7A-iB) \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d}}{7a}$$

↓ 219

$$\frac{2B \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d} - \frac{35i\sqrt{2}a^{5/2}(B+IA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + \frac{8a^2(7A-iB)\sqrt{a+ia \tan(c+dx)}}{d} + \frac{2a(7A-31iB)(a+ia \tan(c+dx))^{3/2}}{3d}}{5a} - \frac{2a(7A-iB) \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d}}{7a}$$

input `Int[Tan[c + d*x]^3*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(2*B*Tan[c + d*x]^3*Sqrt[a + I*a*Tan[c + d*x]])/(7*d) - ((-2*a*(7*A - I*B) *Tan[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]])/(5*d) + (((35*I)*Sqrt[2]*a^(5/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d + (8 *a^2*(7*A - I*B)*Sqrt[a + I*a*Tan[c + d*x]]/d + (2*a*(7*A - (31*I)*B)*(a + I*a*Tan[c + d*x])^(3/2))/(3*d))/(5*a))/(7*a)`

3.67.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3961 `Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`
- rule 4010 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`
- rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`
- rule 4080 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`

3.67.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.84

method	result
derivativedivides	$\frac{2iB(a+ia \tan(dx+c))^{\frac{7}{2}} - 4iBa(a+ia \tan(dx+c))^{\frac{5}{2}} - 2Aa(a+ia \tan(dx+c))^{\frac{5}{2}} + 4iB a^2(a+ia \tan(dx+c))^{\frac{3}{2}} + 2A a^2(a+ia \tan(dx+c))^{\frac{3}{2}}}{a^3 d}$
default	$\frac{2iB(a+ia \tan(dx+c))^{\frac{7}{2}} - 4iBa(a+ia \tan(dx+c))^{\frac{5}{2}} - 2Aa(a+ia \tan(dx+c))^{\frac{5}{2}} + 4iB a^2(a+ia \tan(dx+c))^{\frac{3}{2}} + 2A a^2(a+ia \tan(dx+c))^{\frac{3}{2}}}{a^3 d}$
parts	$\frac{2A \left(-\frac{(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} + \frac{a(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - a^2 \sqrt{a+ia \tan(dx+c)} + \frac{a^{\frac{5}{2}} \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}}\right)}{2} \right)}{d a^2} + \frac{2iB}{a^3 d}$

input `int((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output
$$\frac{2/d/a^3*(1/7*I*B*(a+I*a*\tan(d*x+c))^{(7/2)}-2/5*I*B*a*(a+I*a*\tan(d*x+c))^{(5/2)}-1/5*A*a*(a+I*a*\tan(d*x+c))^{(5/2)}+2/3*I*B*a^2*(a+I*a*\tan(d*x+c))^{(3/2)}+1/3*A*a^2*(a+I*a*\tan(d*x+c))^{(3/2)}-A*a^3*(a+I*a*\tan(d*x+c))^{(1/2)}+1/2*a^{(7/2)}*(A-I*B)*2^{(1/2)}*\operatorname{arctanh}(1/2*(a+I*a*\tan(d*x+c))^{(1/2)}*2^{(1/2)}/a^{(1/2)})}{a^3 d}$$

3.67.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 440 vs. 2(151) = 302.

Time = 0.28 (sec) , antiderivative size = 440, normalized size of antiderivative = 2.27

$$\int \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{105 \sqrt{2} (de^{(6i dx+6i c)} + 3 de^{(4i dx+4i c)} + 3 de^{(2i dx+2i c)} + d) \sqrt{\frac{(A^2-2i AB-B^2)a}{d^2}} \log \left(-\frac{4 \left((-i A-B) a e^{(i dx+i c)} - (i de^{(2i dx+2i c)} + d) \sqrt{\frac{(A^2-2i AB-B^2)a}{d^2}} \right)}{2 \sqrt{a+ia \tan(c+dx)}} \right)}{a^3 d}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^3*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output $1/210*(105*\sqrt{2}*(d*e^{(6*I*d*x + 6*I*c)} + 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} + d)*\sqrt{(A^2 - 2*I*A*B - B^2)*a/d^2}*\log(-4*((-I*A - B)*a*e^{(I*d*x + I*c)} - (I*d*e^{(2*I*d*x + 2*I*c)} + I*d)*\sqrt{(A^2 - 2*I*A*B - B^2)*a/d^2}*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}))e^{(-I*d*x - I*c)}/(I*A + B)) - 105*\sqrt{2}*(d*e^{(6*I*d*x + 6*I*c)} + 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} + d)*\sqrt{(A^2 - 2*I*A*B - B^2)*a/d^2}*\log(-4*((-I*A - B)*a*e^{(I*d*x + I*c)} - (-I*d*e^{(2*I*d*x + 2*I*c)} - I*d)*\sqrt{(A^2 - 2*I*A*B - B^2)*a/d^2}*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}))e^{(-I*d*x - I*c)}/(I*A + B)) - 4*\sqrt{2}*((119*A - 92*I*B)*e^{(7*I*d*x + 7*I*c)} + 7*(37*A - 16*I*B)*e^{(5*I*d*x + 5*I*c)} + 35*(7*A - 4*I*B)*e^{(3*I*d*x + 3*I*c)} + 105*A*e^{(I*d*x + I*c)})*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})/(d*e^{(6*I*d*x + 6*I*c)} + 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} + d)$

3.67.6 Sympy [F]

$$\int \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \sqrt{ia (\tan(c + dx) - i)} (A + B \tan(c + dx)) \tan^3(c + dx) dx$$

input `integrate((a+I*a*tan(d*x+c))**(1/2)*tan(d*x+c)**3*(A+B*tan(d*x+c)),x)`

output `Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))*tan(c + d*x)**3, x)`

3.67.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.79

$$\int \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx =$$

$$\frac{105 \sqrt{2} (A - i B) a^{\frac{9}{2}} \log\left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}}\right) - 60i (ia \tan(dx+c) + a)^{\frac{7}{2}} Ba + 84 (ia \tan(dx+c) + a)^{\frac{5}{2}} Ba}{210 a^4 d}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

3.67. $\int \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

output
$$-1/210*(105*\sqrt{2}*(A - I*B)*a^{(9/2)}*\log(-(\sqrt{2}*\sqrt{a} - \sqrt{I*a*\tan(dx + c) + a})/(\sqrt{2}*\sqrt{a} + \sqrt{I*a*\tan(dx + c) + a}))) - 60*I*(I*a*\tan(dx + c) + a)^{(7/2)}*B*a + 84*(I*a*\tan(dx + c) + a)^{(5/2)}*(A + 2*I*B)*a^2 - 140*(I*a*\tan(dx + c) + a)^{(3/2)}*(A + 2*I*B)*a^3 + 420*\sqrt{I*a*\tan(dx + c) + a}*A*a^4/(a^4*d)$$

3.67.8 Giac [F(-1)]

Timed out.

$$\int \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output Timed out

3.67.9 Mupad [B] (verification not implemented)

Time = 1.98 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.11

$$\begin{aligned} & \int \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= -\frac{2A \sqrt{a + a \tan(c + dx)} \operatorname{li}}{d} + \frac{2A (a + a \tan(c + dx) \operatorname{li})^{3/2}}{3ad} \\ & \quad - \frac{2A (a + a \tan(c + dx) \operatorname{li})^{5/2}}{5a^2 d} + \frac{B (a + a \tan(c + dx) \operatorname{li})^{3/2} 4i}{3ad} \\ & \quad - \frac{B (a + a \tan(c + dx) \operatorname{li})^{5/2} 4i}{5a^2 d} + \frac{B (a + a \tan(c + dx) \operatorname{li})^{7/2} 2i}{7a^3 d} \\ & \quad - \frac{\sqrt{2} B \sqrt{-a} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2\sqrt{-a}}\right) \operatorname{li}}{d} - \frac{\sqrt{2} A \sqrt{a} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}} \operatorname{li}}{2\sqrt{a}}\right) \operatorname{li}}{d} \end{aligned}$$

input `int(tan(c + d*x)^3*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2),x)`

output $(2A(a + a\tan(c + dx)i)^{3/2})/(3ad) - (2A(a + a\tan(c + dx)i)^{1/2})/d - (2A(a + a\tan(c + dx)i)^{5/2})/(5a^2d) + (B(a + a\tan(c + dx)i)^{3/2}i)/(3ad) - (B(a + a\tan(c + dx)i)^{5/2}i)/(5a^2d) + (B(a + a\tan(c + dx)i)^{7/2}i)/(7a^3d) - (2^{1/2}B(-a)^{1/2})\operatorname{atan}((2^{1/2}(a + a\tan(c + dx)i)^{1/2})/(2(-a)^{1/2}))i/d - (2^{1/2}Aa^{1/2})\operatorname{atan}((2^{1/2}(a + a\tan(c + dx)i)^{1/2}i)/(2a^{1/2}))i/d$

3.68 $\int \tan^2(c+dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.68.1	Optimal result	818
3.68.2	Mathematica [A] (verified)	818
3.68.3	Rubi [A] (verified)	819
3.68.4	Maple [A] (verified)	822
3.68.5	Fricas [B] (verification not implemented)	822
3.68.6	Sympy [F]	823
3.68.7	Maxima [A] (verification not implemented)	823
3.68.8	Giac [F(-1)]	824
3.68.9	Mupad [B] (verification not implemented)	824

3.68.1 Optimal result

Integrand size = 36, antiderivative size = 143

$$\int \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\sqrt{2}\sqrt{a}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{8B\sqrt{a + ia \tan(c + dx)}}{5d}$$

$$+ \frac{2B \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{2(5iA + B)(a + ia \tan(c + dx))^{3/2}}{15ad}$$

output `(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)*a^(1/2)/d-8/5*B*(a+I*a*tan(d*x+c))^(1/2)/d+2/5*B*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^2/d-2/15*(5*I*A+B)*(a+I*a*tan(d*x+c))^(3/2)/a/d`

3.68.2 Mathematica [A] (verified)

Time = 1.28 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.78

$$\int \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{15\sqrt{2}\sqrt{a}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + 2\sqrt{a + ia \tan(c + dx)}(-5iA - 13B + (5A - iB) \tan(c + dx))}{15d}$$

input `Integrate[Tan[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x
]`

output `(15*Sqrt[2]*Sqrt[a]*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*
Sqrt[a])] + 2*Sqrt[a + I*a*Tan[c + d*x]]*((-5*I)*A - 13*B + (5*A - I*B)*Ta
n[c + d*x] + 3*B*Tan[c + d*x]^2))/(15*d)`

3.68.3 Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {3042, 4080, 27, 3042, 4075, 3042, 4010, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c + dx)^2 \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4080} \\
 & \frac{2 \int -\frac{1}{2} \tan(c + dx) \sqrt{i \tan(c + dx) a + a(4aB - a(5A - iB) \tan(c + dx))} dx}{5a} + \\
 & \quad \frac{2B \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} \\
 & \quad \downarrow \text{27} \\
 & \frac{2B \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \\
 & \frac{\int \tan(c + dx) \sqrt{i \tan(c + dx) a + a(4aB - a(5A - iB) \tan(c + dx))} dx}{5a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2B \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \\
 & \frac{\int \tan(c + dx) \sqrt{i \tan(c + dx) a + a(4aB - a(5A - iB) \tan(c + dx))} dx}{5a} \\
 & \quad \downarrow \text{4075}
 \end{aligned}$$

3.68. $\int \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \frac{2B \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d} - \\
& \frac{\int \sqrt{i \tan(c+dx)a+a}(a(5A-iB)+4aB \tan(c+dx))dx + \frac{2(B+5iA)(a+ia \tan(c+dx))^{3/2}}{3d}}{5a} \\
& \quad \downarrow \text{3042} \\
& \frac{2B \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d} - \\
& \frac{\int \sqrt{i \tan(c+dx)a+a}(a(5A-iB)+4aB \tan(c+dx))dx + \frac{2(B+5iA)(a+ia \tan(c+dx))^{3/2}}{3d}}{5a} \\
& \quad \downarrow \text{4010} \\
& \frac{2B \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d} - \\
& \frac{5a(A-iB) \int \sqrt{i \tan(c+dx)a+adx} + \frac{2(B+5iA)(a+ia \tan(c+dx))^{3/2}}{3d} + \frac{8aB \sqrt{a+ia \tan(c+dx)}}{d}}{5a} \\
& \quad \downarrow \text{3042} \\
& \frac{2B \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d} - \\
& \frac{5a(A-iB) \int \sqrt{i \tan(c+dx)a+adx} + \frac{2(B+5iA)(a+ia \tan(c+dx))^{3/2}}{3d} + \frac{8aB \sqrt{a+ia \tan(c+dx)}}{d}}{5a} \\
& \quad \downarrow \text{3961} \\
& \frac{2B \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d} - \\
& \frac{10ia^2(A-iB) \int \frac{1}{a-ia \tan(c+dx)} d \sqrt{i \tan(c+dx)a+a} + \frac{2(B+5iA)(a+ia \tan(c+dx))^{3/2}}{3d} + \frac{8aB \sqrt{a+ia \tan(c+dx)}}{d}}{5a} \\
& \quad \downarrow \text{219} \\
& \frac{2B \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d} - \\
& \frac{5i\sqrt{2}a^{3/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + \frac{2(B+5iA)(a+ia \tan(c+dx))^{3/2}}{3d} + \frac{8aB \sqrt{a+ia \tan(c+dx)}}{d}}{5a}
\end{aligned}$$

input `Int[Tan[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(2*B*Tan[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]])/(5*d) - (((-5*I)*Sqrt[2]*a^(3/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d + (8*a*B*Sqrt[a + I*a*Tan[c + d*x]])/d + (2*((5*I)*A + B)*(a + I*a*Tan[c + d*x])^(3/2))/(3*d))/(5*a)`

3.68. $\int \tan^2(c+dx) \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx$

3.68.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3961 `Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`
- rule 4010 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`
- rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`
- rule 4080 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`

3.68.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.87

method	result
derivativedivides	$\frac{2i \left(\frac{iB(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} - \frac{iBa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - \frac{Aa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + ia^2 B \sqrt{a+ia \tan(dx+c)} + \frac{a^{\frac{5}{2}}(-iB+A)\sqrt{2} \arctan\left(\frac{\sqrt{a+ia \tan(dx+c)}}{2\sqrt{a}}\right)}{2} \right)}{da^2}$
default	$\frac{2i \left(\frac{iB(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} - \frac{iBa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - \frac{Aa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + ia^2 B \sqrt{a+ia \tan(dx+c)} + \frac{a^{\frac{5}{2}}(-iB+A)\sqrt{2} \arctan\left(\frac{\sqrt{a+ia \tan(dx+c)}}{2\sqrt{a}}\right)}{2} \right)}{da^2}$
parts	$\frac{2iA \left(-\frac{(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + \frac{a^{\frac{3}{2}}\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{2\sqrt{a}}\right)}{2} \right)}{da} + \frac{2B \left(-\frac{(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} + \frac{a(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} \right)}{da}$

```
input int((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 2*I/d/a^2*(1/5*I*B*(a+I*a*tan(d*x+c))^(5/2)-1/3*I*B*a*(a+I*a*tan(d*x+c))^(3/2)-1/3*A*a*(a+I*a*tan(d*x+c))^(3/2)+I*a^2*B*(a+I*a*tan(d*x+c))^(1/2)+1/2*a^(5/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2)))
```

3.68.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 383 vs. 2(112) = 224.

Time = 0.26 (sec) , antiderivative size = 383, normalized size of antiderivative = 2.68

$$\int \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx = \frac{15\sqrt{2}(de^{(4i dx+4i c)} + 2de^{(2i dx+2i c)} + d) \sqrt{-\frac{(A^2-2iAB-B^2)a}{d^2}} \log \left(-\frac{4 \left((-iA-B)ae^{(i dx+i c)} + (de^{(2i dx+2i c)}+d) \sqrt{-\frac{(A^2-2iAB-B^2)a}{d^2}} \right)}{iA+B} \right)}{da^2}$$

```
input integrate((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^2*(A+B*tan(d*x+c)),x, algorithm="fracas")
```

3.68. $\int \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

```
output -1/30*(15*sqrt(2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(A^2 - 2*I*A*B - B^2)*a/d^2)*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) + (d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(A^2 - 2*I*A*B - B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/(I*A + B)) - 15*sqrt(2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(A^2 - 2*I*A*B - B^2)*a/d^2)*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) - (d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(A^2 - 2*I*A*B - B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/(I*A + B)) + 4*sqrt(2)*((10*I*A + 17*B)*e^(5*I*d*x + 5*I*c) + 10*(I*A + 2*B)*e^(3*I*d*x + 3*I*c) + 15*B*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)
```

3.68.6 Sympy [F]

$$\begin{aligned} & \int \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= \int \sqrt{ia (\tan(c + dx) - i)} (A + B \tan(c + dx)) \tan^2(c + dx) dx \end{aligned}$$

```
input integrate((a+I*a*tan(d*x+c))**(1/2)*tan(d*x+c)**2*(A+B*tan(d*x+c)),x)
```

```
output Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))*tan(c + d*x)**2, x)
```

3.68.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.91

$$\begin{aligned} & \int \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx = \\ & \frac{i \left(15 \sqrt{2} (A - i B) a^{\frac{7}{2}} \log \left(-\frac{\sqrt{2} \sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2} \sqrt{a} + \sqrt{ia \tan(dx+c)+a}} \right) - 12i (ia \tan(dx+c) + a)^{\frac{5}{2}} Ba + 20 (ia \tan(dx+c) + a)^{\frac{3}{2}} \right)}{30 a^3 d} \end{aligned}$$

```
input integrate((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^2*(A+B*tan(d*x+c)),x, algorithm="maxima")
```

output
$$-1/30*I*(15*\sqrt{2}*(A - I*B)*a^{7/2}*\log(-(\sqrt{2}*\sqrt{a} - \sqrt{I*a*\tan(dx + c) + a})/(\sqrt{2}*\sqrt{a} + \sqrt{I*a*\tan(dx + c) + a})) - 12*I*(I*a*\tan(dx + c) + a)^{5/2}*B*a + 20*(I*a*\tan(dx + c) + a)^{3/2}*(A + I*B)*a^2 - 60*I*\sqrt{I*a*\tan(dx + c) + a}*B*a^3)/(a^3*d)$$

3.68.8 Giac [F(-1)]

Timed out.

$$\int \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output Timed out

3.68.9 Mupad [B] (verification not implemented)

Time = 8.28 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.17

$$\begin{aligned} & \int \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= -\frac{2B \sqrt{a + a \tan(c + dx)} \operatorname{li}}{d} - \frac{A(a + a \tan(c + dx) \operatorname{li})^{3/2} 2i}{3ad} \\ &+ \frac{2B(a + a \tan(c + dx) \operatorname{li})^{3/2}}{3ad} - \frac{2B(a + a \tan(c + dx) \operatorname{li})^{5/2}}{5a^2 d} \\ &+ \frac{\sqrt{2} A \sqrt{-a} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2\sqrt{-a}}\right) \operatorname{li}}{d} - \frac{\sqrt{2} B \sqrt{a} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2\sqrt{a}}\right) \operatorname{li}}{d} \end{aligned}$$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2),x)`

output
$$(2*B*(a + a*\tan(c + d*x)*1i)^{3/2})/(3*a*d) - (A*(a + a*\tan(c + d*x)*1i)^{3/2}*2i)/(3*a*d) - (2*B*(a + a*\tan(c + d*x)*1i)^{1/2})/d - (2*B*(a + a*\tan(c + d*x)*1i)^{5/2})/(5*a^2*d) + (2^{1/2}*A*(-a)^{1/2}*\operatorname{atan}((2^{1/2}*(a + a*\tan(c + d*x)*1i)^{1/2})/(2*(-a)^{1/2}))*1i)/d - (2^{1/2}*B*a^{1/2}*\operatorname{atan}((2^{1/2}*(a + a*\tan(c + d*x)*1i)^{1/2})*1i)/(2*a^{1/2}))*1i)/d$$

3.68.
$$\int \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

3.69 $\int \tan(c+dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.69.1	Optimal result	825
3.69.2	Mathematica [A] (verified)	825
3.69.3	Rubi [A] (verified)	826
3.69.4	Maple [A] (verified)	828
3.69.5	Fricas [B] (verification not implemented)	828
3.69.6	Sympy [F]	829
3.69.7	Maxima [A] (verification not implemented)	829
3.69.8	Giac [F]	830
3.69.9	Mupad [B] (verification not implemented)	830

3.69.1 Optimal result

Integrand size = 34, antiderivative size = 105

$$\int \tan(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= -\frac{\sqrt{2}\sqrt{a}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{2A\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2iB(a + ia \tan(c + dx))^{3/2}}{3ad}$$

```
output -(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)*a^(1/2)/d+2*A*(a+I*a*tan(d*x+c))^(1/2)/d-2/3*I*B*(a+I*a*tan(d*x+c))^(3/2)/a/d
```

3.69.2 Mathematica [A] (verified)

Time = 0.70 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.88

$$\int \tan(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{-3\sqrt{2}\sqrt{a}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + 2\sqrt{a + ia \tan(c + dx)}(3A - iB + B \tan(c + dx))}{3d}$$

input `Integrate[Tan[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(-3*Sqrt[2]*Sqrt[a]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] + 2*Sqrt[a + I*a*Tan[c + d*x]]*(3*A - I*B + B*Tan[c + d*x]))/(3*d)`

3.69.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.02, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.206$, Rules used = {3042, 4075, 3042, 4010, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4075} \\
 & \int \sqrt{i \tan(c + dx) a + a} (A \tan(c + dx) - B) dx - \frac{2iB(a + ia \tan(c + dx))^{3/2}}{3ad} \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{i \tan(c + dx) a + a} (A \tan(c + dx) - B) dx - \frac{2iB(a + ia \tan(c + dx))^{3/2}}{3ad} \\
 & \quad \downarrow \text{4010} \\
 & -(B + iA) \int \sqrt{i \tan(c + dx) a + a} dx + \frac{2A \sqrt{a + ia \tan(c + dx)}}{d} - \frac{2iB(a + ia \tan(c + dx))^{3/2}}{3ad} \\
 & \quad \downarrow \text{3042} \\
 & -(B + iA) \int \sqrt{i \tan(c + dx) a + a} dx + \frac{2A \sqrt{a + ia \tan(c + dx)}}{d} - \frac{2iB(a + ia \tan(c + dx))^{3/2}}{3ad} \\
 & \quad \downarrow \text{3961}
 \end{aligned}$$

$$\frac{2ia(B + iA) \int \frac{1}{a - ia \tan(c + dx)} d\sqrt{i \tan(c + dx)a + a}}{d} + \frac{2A\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2iB(a + ia \tan(c + dx))^{3/2}}{3ad}$$

↓ 219

$$\frac{i\sqrt{2}\sqrt{a}(B + iA)\operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{2A\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2iB(a + ia \tan(c + dx))^{3/2}}{3ad}$$

input `Int[Tan[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(I*Sqrt[2]*Sqrt[a]*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])]/d + (2*A*Sqrt[a + I*a*Tan[c + d*x]])/d - (((2*I)/3)*B*(a + I*a*Tan[c + d*x])^(3/2))/(a*d)`

3.69.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

3.69.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.78

method	result
derivativedivides	$\frac{-\frac{2iB(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + 2aA\sqrt{a+ia \tan(dx+c)} - a^{\frac{3}{2}}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{ad}$
default	$\frac{-\frac{2iB(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + 2aA\sqrt{a+ia \tan(dx+c)} - a^{\frac{3}{2}}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{ad}$
parts	$\frac{A\left(2\sqrt{a+ia \tan(dx+c)} - \sqrt{a}\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)\right)}{d} + \frac{2iB\left(-\frac{(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + \frac{a^{\frac{3}{2}}\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2}\right)}{da}$

```
input int((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 2/d/a*(-1/3*I*B*(a+I*a*tan(d*x+c))^(3/2)+a*A*(a+I*a*tan(d*x+c))^(1/2)-1/2*
a^(3/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/
2)))
```

3.69.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 332 vs. 2(80) = 160.

Time = 0.26 (sec) , antiderivative size = 332, normalized size of antiderivative = 3.16

$$\int \tan(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx =$$

$$3\sqrt{2} \left(de^{(2i dx + 2i c)} + d \right) \sqrt{\frac{(A^2 - 2i AB - B^2)a}{d^2}} \log \left(-\frac{4 \left((-iA - B) a e^{(i dx + i c)} - (i de^{(2i dx + 2i c)} + i d) \sqrt{\frac{(A^2 - 2i AB - B^2)a}{d^2}} \sqrt{e^{(2i dx + 2i c)}} \right)}{iA + B} \right)$$

3.69. $\int \tan(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

```
input integrate((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)*(A+B*tan(d*x+c)),x, algorithm
m="fricas")
```

```
output -1/6*(3*sqrt(2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt((A^2 - 2*I*A*B - B^2)*a/d
^2)*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) - (I*d*e^(2*I*d*x + 2*I*c) + I*d)
*sqrt((A^2 - 2*I*A*B - B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-
I*d*x - I*c)/(I*A + B)) - 3*sqrt(2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt((A^2
- 2*I*A*B - B^2)*a/d^2)*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) - (-I*d*e^(2*
I*d*x + 2*I*c) - I*d)*sqrt((A^2 - 2*I*A*B - B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x
+ 2*I*c) + 1)))*e^(-I*d*x - I*c)/(I*A + B)) - 4*sqrt(2)*((3*A - 2*I*B)*e^
(3*I*d*x + 3*I*c) + 3*A*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)
)/(d*e^(2*I*d*x + 2*I*c) + d)
```

3.69.6 Sympy [F]

$$\begin{aligned} & \int \tan(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= \int \sqrt{ia (\tan(c + dx) - i)} (A + B \tan(c + dx)) \tan(c + dx) dx \end{aligned}$$

```
input integrate((a+I*a*tan(d*x+c))**(1/2)*tan(d*x+c)*(A+B*tan(d*x+c)),x)
```

```
output Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))*tan(c + d*x), x
)
```

3.69.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.02

$$\begin{aligned} & \int \tan(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= \frac{3\sqrt{2}(A - iB)a^{\frac{5}{2}} \log\left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}}\right) - 4i(i a \tan(dx+c) + a)^{\frac{3}{2}}Ba + 12\sqrt{ia \tan(dx+c) + a}A}{6a^2d} \end{aligned}$$

```
input integrate((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)*(A+B*tan(d*x+c)),x, algorithm
m="maxima")
```

3.69. $\int \tan(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

output $1/6*(3*\sqrt{2}*(A - I*B)*a^{5/2}*\log(-(\sqrt{2}*\sqrt{a} - \sqrt{I*a*\tan(dx + c) + a})/(\sqrt{2}*\sqrt{a} + \sqrt{I*a*\tan(dx + c) + a})) - 4*I*(I*a*\tan(dx + c) + a)^{3/2}*B*a + 12*\sqrt{I*a*\tan(dx + c) + a}*A*a^2)/(a^{2*d})$

3.69.8 Giac [F]

$$\begin{aligned} & \int \tan(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A) \sqrt{ia \tan(dx + c) + a \tan(dx + c)} dx \end{aligned}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)*tan(d*x + c), x)`

3.69.9 Mupad [B] (verification not implemented)

Time = 8.00 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.14

$$\begin{aligned} & \int \tan(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= \frac{2A \sqrt{a + a \tan(c + dx)} \operatorname{li}}{d} - \frac{B(a + a \tan(c + dx) \operatorname{li})^{3/2} 2i}{3ad} \\ &+ \frac{\sqrt{2} B \sqrt{-a} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx)} \operatorname{li}}{2\sqrt{-a}}\right) \operatorname{li}}{d} - \frac{\sqrt{2} A \sqrt{a} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx)} \operatorname{li}}{2\sqrt{a}}\right)}{d} \end{aligned}$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2),x)`

output $(2*A*(a + a*\tan(c + d*x)*1i)^{1/2})/d - (B*(a + a*\tan(c + d*x)*1i)^{3/2}*2i)/(3*a*d) + (2^{1/2}*B*(-a)^{1/2}*\operatorname{atan}((2^{1/2}*(a + a*\tan(c + d*x)*1i)^{1/2}))/((2*(-a)^{1/2}))*1i)/d - (2^{1/2}*A*a^{1/2}*\operatorname{atanh}((2^{1/2}*(a + a*\tan(c + d*x)*1i)^{1/2}))/((2*a^{1/2}))/d$

3.70 $\int \sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$

3.70.1	Optimal result	831
3.70.2	Mathematica [A] (verified)	831
3.70.3	Rubi [A] (verified)	832
3.70.4	Maple [A] (verified)	833
3.70.5	Fricas [B] (verification not implemented)	834
3.70.6	Sympy [F]	834
3.70.7	Maxima [A] (verification not implemented)	835
3.70.8	Giac [F(-1)]	835
3.70.9	Mupad [B] (verification not implemented)	835

3.70.1 Optimal result

Integrand size = 28, antiderivative size = 75

$$\int \sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$= -\frac{\sqrt{2}\sqrt{a}(iA + B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{2B\sqrt{a + ia \tan(c + dx)}}{d}$$

output `-(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)*a^(1/2)/d+2*B*(a+I*a*tan(d*x+c))^(1/2)/d`

3.70.2 Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00

$$\int \sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$= \frac{-i\sqrt{2}\sqrt{a}(A - iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + 2B\sqrt{a + ia \tan(c + dx)}}{d}$$

input `Integrate[Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `((-I)*Sqrt[2]*Sqrt[a]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] + 2*B*Sqrt[a + I*a*Tan[c + d*x]])/d`

3.70.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$, Rules used = {3042, 4010, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4010} \\
 & (A - iB) \int \sqrt{i \tan(c + dx)a + adx} + \frac{2B\sqrt{a + ia \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \int \sqrt{i \tan(c + dx)a + adx} + \frac{2B\sqrt{a + ia \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{3961} \\
 & \frac{2B\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2ia(A - iB) \int \frac{1}{a - ia \tan(c + dx)} d\sqrt{i \tan(c + dx)a + a}}{d} \\
 & \quad \downarrow \text{219} \\
 & \frac{2B\sqrt{a + ia \tan(c + dx)}}{d} - \frac{i\sqrt{2}\sqrt{a}(A - iB)\operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}
 \end{aligned}$$

input `Int[Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `((-I)*Sqrt[2]*Sqrt[a]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d + (2*B*Sqrt[a + I*a*Tan[c + d*x]])/d`

3.70.3.1 Defintions of rubi rules used

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3961 Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[-2*(b/d)
Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a
, b, c, d}, x] && EqQ[a^2 + b^2, 0]
```

```
rule 4010 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] :> Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp
[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e
, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]
```

3.70.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.84

method	result	s
derivativedivides	$\frac{2i \left(-iB\sqrt{a+ia \tan(dx+c)} - \frac{\sqrt{a}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2} \right)}{d}$	6
default	$\frac{2i \left(-iB\sqrt{a+ia \tan(dx+c)} - \frac{\sqrt{a}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2} \right)}{d}$	6
parts	$-\frac{iA\sqrt{a}\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{d} + \frac{B \left(2\sqrt{a+ia \tan(dx+c)} - \sqrt{a}\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right) \right)}{d}$	9

```
input int((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 2*I/d*(-I*B*(a+I*a*tan(d*x+c))^(1/2)-1/2*a^(1/2)*(A-I*B)*2^(1/2)*arctanh(1
/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2)))
```

3.70. $\int \sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$

3.70.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 272 vs. 2(59) = 118.

Time = 0.26 (sec) , antiderivative size = 272, normalized size of antiderivative = 3.63

$$\int \sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$= \frac{4\sqrt{2}B\sqrt{\frac{a}{e^{2i dx+2i c}+1}}e^{i dx+i c} + \sqrt{2}d\sqrt{-\frac{(A^2-2i AB-B^2)a}{d^2}} \log\left(-\frac{4\left((-i A-B)ae^{i dx+i c}+(de^{2i dx+2i c}+d)\sqrt{-\frac{(A^2-2i AB-B^2)a}{d^2}}\right)}{i A+B}}\right)}{1}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/2*(4*sqrt(2)*B*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*e^(I*d*x + I*c) + sqrt(2)*d*sqrt(-(A^2 - 2*I*A*B - B^2)*a/d^2)*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) + (d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(A^2 - 2*I*A*B - B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/(I*A + B) - sqrt(2)*d*sqrt(-(A^2 - 2*I*A*B - B^2)*a/d^2)*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) - (d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(A^2 - 2*I*A*B - B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/(I*A + B))/d`

3.70.6 Sympy [F]

$$\int \sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$= \int \sqrt{ia (\tan(c + dx) - i)}(A + B \tan(c + dx)) dx$$

input `integrate((a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x)), x)`

3.70.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.16

$$\int \sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$= \frac{i \left(\sqrt{2}(A - iB)a^{\frac{3}{2}} \log \left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}} \right) - 4i \sqrt{ia \tan(dx+c)+a}Ba \right)}{2ad}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`output `1/2*I*(sqrt(2)*(A - I*B)*a^(3/2)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - 4*I*sqrt(I*a*tan(d*x + c) + a)*B*a)/(a*d)`**3.70.8 Giac [F(-1)]**

Timed out.

$$\int \sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`output `Timed out`**3.70.9 Mupad [B] (verification not implemented)**

Time = 7.49 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.28

$$\int \sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$= \frac{2B \sqrt{a + a \tan(c + dx)} \operatorname{li} \left(\frac{\sqrt{2}\sqrt{a+a \tan(c+dx)} \operatorname{li}}{2\sqrt{-a}} \right)}{d} - \frac{\sqrt{2}A \sqrt{-a} \operatorname{atan} \left(\frac{\sqrt{2}\sqrt{a+a \tan(c+dx)} \operatorname{li}}{2\sqrt{-a}} \right)}{d}$$

$$- \frac{\sqrt{2}B \sqrt{a} \operatorname{atanh} \left(\frac{\sqrt{2}\sqrt{a+a \tan(c+dx)} \operatorname{li}}{2\sqrt{a}} \right)}{d}$$

3.70. $\int \sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$

input `int((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2),x)`

output `(2*B*(a + a*tan(c + d*x)*1i)^(1/2))/d - (2^(1/2)*A*(-a)^(1/2)*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2)))*1i)/d - (2^(1/2)*B*a^(1/2)*atanh((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*a^(1/2))))/d`

3.71 $\int \cot(c+dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.71.1	Optimal result	837
3.71.2	Mathematica [A] (verified)	837
3.71.3	Rubi [A] (verified)	838
3.71.4	Maple [A] (verified)	840
3.71.5	Fricas [B] (verification not implemented)	840
3.71.6	Sympy [F]	842
3.71.7	Maxima [A] (verification not implemented)	842
3.71.8	Giac [F]	842
3.71.9	Mupad [B] (verification not implemented)	843

3.71.1 Optimal result

Integrand size = 34, antiderivative size = 86

$$\int \cot(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= -\frac{2\sqrt{a}A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{d} + \frac{\sqrt{2}\sqrt{a}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}$$

output `-2*A*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))*a^(1/2)/d+(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)*a^(1/2)/d`

3.71.2 Mathematica [A] (verified)

Time = 1.11 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.92

$$\int \cot(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\sqrt{a} \left(-2A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) + \sqrt{2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) \right)}{d}$$

input `Integrate[Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[a]*(-2*A*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] + Sqrt[2]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])]))/d`

3.71. $\int \cot(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.71.3 Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.03, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot(c+dx) \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx))}{\tan(c+dx)} dx \\
 & \quad \downarrow \text{4083} \\
 & (B+iA) \int \sqrt{i \tan(c+dx) a+adx} + \frac{A \int \cot(c+dx) (a-ia \tan(c+dx)) \sqrt{i \tan(c+dx) a+adx}}{a} dx \\
 & \quad \downarrow \text{3042} \\
 & (B+iA) \int \sqrt{i \tan(c+dx) a+adx} + \frac{A \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx) a+adx}}{\tan(c+dx)} dx}{a} \\
 & \quad \downarrow \text{3961} \\
 & \frac{A \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx) a+adx}}{\tan(c+dx)} dx}{a} - \frac{2ia(B+iA) \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx) a+adx}}{d} \\
 & \quad \downarrow \text{219} \\
 & \frac{A \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx) a+adx}}{\tan(c+dx)} dx}{a} - \frac{i\sqrt{2}\sqrt{a}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} \\
 & \quad \downarrow \text{4082} \\
 & \frac{aA \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx) a+adx}} d \tan(c+dx)}{d} - \frac{i\sqrt{2}\sqrt{a}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} \\
 & \quad \downarrow \text{73} \\
 & \frac{2iA \int \frac{1}{i - \frac{i \tan(c+dx) a+adx}{a}} d\sqrt{i \tan(c+dx) a+adx}}{d} - \frac{i\sqrt{2}\sqrt{a}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} \\
 & \quad \downarrow \text{221}
 \end{aligned}$$

3.71. $\int \cot(c+dx) \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx$

$$\frac{i\sqrt{2}\sqrt{a}(B + iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{2\sqrt{a}A\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{a}}\right)}{d}$$

input `Int[Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(-2*Sqrt[a]*A*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d - (I*Sqrt[2]*Sqrt[a]*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d`

3.71.3.1 Defintions of rubi rules used

rule 73 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

```
rule 4083 Int[(((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(
A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A
*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*T
an[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

3.71.4 Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$2a \left(\frac{(iB-A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right) - A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{\sqrt{a}}\right)}{d} \right)$	72
default	$2a \left(\frac{(iB-A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right) - A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{\sqrt{a}}\right)}{d} \right)$	72

```
input int(cot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 2/d*a*(-1/2*(-A+I*B)*2^(1/2)/a^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*
2^(1/2)/a^(1/2))-A/a^(1/2)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2)))
```

3.71.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 447 vs. 2(65) = 130.

3.71. $\int \cot(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

Time = 0.26 (sec) , antiderivative size = 447, normalized size of antiderivative = 5.20

$$\int \cot(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{1}{2} \sqrt{2} \sqrt{\frac{(A^2 - 2iAB - B^2)a}{d^2}} \log \left(-\frac{4 \left((-iA - B)ae^{(idx+ic)} - (ide^{(2idx+2ic)} + id) \sqrt{\frac{(A^2 - 2iAB - B^2)a}{d^2}} \sqrt{\frac{a}{e^{(2idx+2ic)} + 1}} \right)}{iA + B} \right)$$

$$- \frac{1}{2} \sqrt{2} \sqrt{\frac{(A^2 - 2iAB - B^2)a}{d^2}} \log \left(-\frac{4 \left((-iA - B)ae^{(idx+ic)} - (-ide^{(2idx+2ic)} - id) \sqrt{\frac{(A^2 - 2iAB - B^2)a}{d^2}} \sqrt{\frac{a}{e^{(2idx+2ic)} + 1}} \right)}{iA + B} \right)$$

$$- \frac{1}{2} \sqrt{\frac{A^2 a}{d^2}} \log \left(\frac{16 \left(3Aa^2 e^{(2idx+2ic)} + Aa^2 + 2\sqrt{2}(ade^{(3idx+3ic)} + ade^{(idx+ic)}) \sqrt{\frac{A^2 a}{d^2}} \sqrt{\frac{a}{e^{(2idx+2ic)} + 1}} \right)}{A} \right) e^{(-2idx - 2ic)}$$

$$+ \frac{1}{2} \sqrt{\frac{A^2 a}{d^2}} \log \left(\frac{16 \left(3Aa^2 e^{(2idx+2ic)} + Aa^2 - 2\sqrt{2}(ade^{(3idx+3ic)} + ade^{(idx+ic)}) \sqrt{\frac{A^2 a}{d^2}} \sqrt{\frac{a}{e^{(2idx+2ic)} + 1}} \right)}{A} \right) e^{(-2idx - 2ic)}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/2*sqrt(2)*sqrt((A^2 - 2*I*A*B - B^2)*a/d^2)*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) - (I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt((A^2 - 2*I*A*B - B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/(I*A + B) - 1/2*sqrt(2)*sqrt((A^2 - 2*I*A*B - B^2)*a/d^2)*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) - (-I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt((A^2 - 2*I*A*B - B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/(I*A + B) - 1/2*sqrt(A^2*a/d^2)*log(16*(3*A*a^2*e^(2*I*d*x + 2*I*c) + A*a^2 + 2*sqrt(2)*(a*d*e^(3*I*d*x + 3*I*c) + a*d*e^(I*d*x + I*c))*sqrt(A^2*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x - 2*I*c)/A + 1/2*sqrt(A^2*a/d^2)*log(16*(3*A*a^2*e^(2*I*d*x + 2*I*c) + A*a^2 - 2*sqrt(2)*(a*d*e^(3*I*d*x + 3*I*c) + a*d*e^(I*d*x + I*c))*sqrt(A^2*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x - 2*I*c)/A)`

3.71.6 Sympy [F]

$$\begin{aligned} & \int \cot(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= \int \sqrt{ia (\tan(c + dx) - i)} (A + B \tan(c + dx)) \cot(c + dx) dx \end{aligned}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))*cot(c + d*x), x)`

3.71.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.31

$$\begin{aligned} & \int \cot(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= -\frac{\sqrt{2}(A - iB)\sqrt{a} \log\left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}}\right) - 2A\sqrt{a} \log\left(\frac{\sqrt{ia \tan(dx+c)+a} - \sqrt{a}}{\sqrt{ia \tan(dx+c)+a} + \sqrt{a}}\right)}{2d} \end{aligned}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(sqrt(2)*(A - I*B)*sqrt(a)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - 2*A*sqrt(a)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a))))/d`

3.71.8 Giac [F]

$$\begin{aligned} & \int \cot(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A) \sqrt{ia \tan(dx + c) + a} \cot(dx + c) dx \end{aligned}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)*cot(d*x + c), x)`

3.71.9 Mupad [B] (verification not implemented)

Time = 8.16 (sec) , antiderivative size = 493, normalized size of antiderivative = 5.73

$$\int \cot(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx =$$

$$\frac{2A\sqrt{a} \operatorname{atanh}\left(\frac{16A^3a^{9/2}d\sqrt{a+a\tan(c+dx)}\operatorname{li}}{16dA^3a^5+32i dA^2Ba^5+16dAB^2a^5} + \frac{16AB^2a^{9/2}d\sqrt{a+a\tan(c+dx)}\operatorname{li}}{16dA^3a^5+32i dA^2Ba^5+16dAB^2a^5} + \frac{A^2Ba^{9/2}d\sqrt{a+a\tan(c+dx)}\operatorname{li}}{16dA^3a^5+32i dA^2Ba^5+16dAB^2a^5}\right)}{d} + \frac{\sqrt{2}\sqrt{-a} \operatorname{atan}\left(\frac{4\sqrt{2}A^3(-a)^{9/2}d\sqrt{a+a\tan(c+dx)}\operatorname{li}}{8dA^3a^5+8i dA^2Ba^5+24dAB^2a^5-8i dB^3a^5} - \frac{\sqrt{2}B^3(-a)^{9/2}d\sqrt{a+a\tan(c+dx)}\operatorname{li}4i}{8dA^3a^5+8i dA^2Ba^5+24dAB^2a^5-8i dB^3a^5} + \frac{12\sqrt{2}AB^2}{8dA^3a^5+8i d}\right)}{d}$$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2),x)`

output `(2^(1/2)*(-a)^(1/2)*atan((4*2^(1/2)*A^3*(-a)^(9/2)*d*(a + a*tan(c + d*x)*1i)^(1/2))/(8*A^3*a^5*d - B^3*a^5*d*8i + 24*A*B^2*a^5*d + A^2*B*a^5*d*8i) - (2^(1/2)*B^3*(-a)^(9/2)*d*(a + a*tan(c + d*x)*1i)^(1/2)*4i)/(8*A^3*a^5*d - B^3*a^5*d*8i + 24*A*B^2*a^5*d + A^2*B*a^5*d*8i) + (12*2^(1/2)*A*B^2*(-a)^(9/2)*d*(a + a*tan(c + d*x)*1i)^(1/2))/(8*A^3*a^5*d - B^3*a^5*d*8i + 24*A*B^2*a^5*d + A^2*B*a^5*d*8i) + (2^(1/2)*A^2*B*(-a)^(9/2)*d*(a + a*tan(c + d*x)*1i)^(1/2)*4i)/(8*A^3*a^5*d - B^3*a^5*d*8i + 24*A*B^2*a^5*d + A^2*B*a^5*d*8i))*(A*1i + B)*1i)/d - (2*A*a^(1/2)*atanh((16*A^3*a^(9/2)*d*(a + a*tan(c + d*x)*1i)^(1/2))/(16*A^3*a^5*d + 16*A*B^2*a^5*d + A^2*B*a^5*d*32i) + (16*A*B^2*a^(9/2)*d*(a + a*tan(c + d*x)*1i)^(1/2))/(16*A^3*a^5*d + 16*A*B^2*a^5*d + A^2*B*a^5*d*32i) + (A^2*B*a^(9/2)*d*(a + a*tan(c + d*x)*1i)^(1/2))*32i)/(16*A^3*a^5*d + 16*A*B^2*a^5*d + A^2*B*a^5*d*32i)))/d`

3.72 $\int \cot^2(c+dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.72.1	Optimal result	844
3.72.2	Mathematica [A] (verified)	844
3.72.3	Rubi [A] (verified)	845
3.72.4	Maple [A] (verified)	848
3.72.5	Fricas [B] (verification not implemented)	849
3.72.6	Sympy [F]	850
3.72.7	Maxima [A] (verification not implemented)	850
3.72.8	Giac [F]	851
3.72.9	Mupad [B] (verification not implemented)	851

3.72.1 Optimal result

Integrand size = 36, antiderivative size = 123

$$\int \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= -\frac{\sqrt{a}(iA + 2B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{d} + \frac{\sqrt{2}\sqrt{a}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{A \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

output $-(I*A+2*B)*\operatorname{arctanh}\left(\frac{(a+I*a*\tan(d*x+c))^{1/2}}{a^{1/2}}\right)*a^{1/2}/d+(I*A+B)*\operatorname{arctanh}\left(\frac{1/2*(a+I*a*\tan(d*x+c))^{1/2}*2^{1/2}}{a^{1/2}}\right)*2^{1/2}*a^{1/2}/d-A*\cot(d*x+c)*(a+I*a*\tan(d*x+c))^{1/2}/d$

3.72.2 Mathematica [A] (verified)

Time = 2.06 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.96

$$\int \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{-i\sqrt{a}(A - 2iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) + \sqrt{2}\sqrt{a}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) - A \cot(c + dx) \sqrt{a}}{d}$$

3.72. $\int \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

input `Integrate[Cot[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `((-I)*Sqrt[a]*(A - (2*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] + Sqrt[2]*Sqrt[a]*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] - A*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]])/d`

3.72.3 Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.09, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.306$, Rules used = {3042, 4081, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx))}{\tan(c + dx)^2} dx \\
 & \quad \downarrow \text{4081} \\
 & \frac{\int \frac{1}{2} \cot(c + dx) \sqrt{ia \tan(c + dx) a + a} (a(iA + 2B) - aA \tan(c + dx)) dx}{\frac{A \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \cot(c + dx) \sqrt{ia \tan(c + dx) a + a} (a(iA + 2B) - aA \tan(c + dx)) dx}{\frac{2a A \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\sqrt{ia \tan(c + dx) a + a} (a(iA + 2B) - aA \tan(c + dx))}{\tan(c + dx)} dx}{2a} - \frac{A \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{4083}
 \end{aligned}$$

3.72. $\int \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

$$\frac{(2B + iA) \int \cot(c + dx)(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx)a + adx} - 2a(A - iB) \int \sqrt{i \tan(c + dx)a + adx}}{d}$$

$$\frac{A \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

↓ 3042

$$\frac{(2B + iA) \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} dx - 2a(A - iB) \int \sqrt{i \tan(c + dx)a + adx}}{d}$$

$$\frac{A \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

↓ 3961

$$\frac{4ia^2(A - iB) \int \frac{1}{a - ia \tan(c + dx)} d \sqrt{i \tan(c + dx)a + a} + (2B + iA) \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} dx}{d}$$

$$\frac{A \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

↓ 219

$$\frac{(2B + iA) \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} dx + \frac{2i\sqrt{2}a^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{d}$$

$$\frac{A \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

↓ 4082

$$\frac{a^2(2B + iA) \int \frac{\cot(c + dx)}{\sqrt{i \tan(c + dx)a + a}} d \tan(c + dx) + \frac{2i\sqrt{2}a^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{d}$$

$$\frac{A \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

↓ 73

$$\frac{2i\sqrt{2}a^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right) - \frac{2ia(2B + iA) \int \frac{1}{i - i(i \tan(c + dx)a + a)} d \sqrt{i \tan(c + dx)a + a}}{d}}{d}$$

$$\frac{A \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

↓ 221

$$\frac{2i\sqrt{2}a^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right) - \frac{2a^{3/2}(2B + iA) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{a}}\right)}{d}}{d}$$

$$\frac{A \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

3.72. $\int \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

input `Int[Cot[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `((-2*a^(3/2)*(I*A + 2*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d + ((2*I)*Sqrt[2]*a^(3/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d)/(2*a) - (A*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]])/d`

3.72.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

```
rule 4082 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 + b^2, 0] && EqQ[A*b + a*B, 0]
```

```
rule 4083 Int((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(
A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A
*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*T
an[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

3.72.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.93

method	result	size
derivativedivides	$2ia^2 \left(-\frac{(iB-A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2a^{\frac{3}{2}}} + \frac{iA\sqrt{a+ia \tan(dx+c)}}{2a \tan(dx+c)} - \frac{(-2iB+A) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{\sqrt{a}}\right)}{a \cdot 2\sqrt{a}} \right)$	114
default	$2ia^2 \left(-\frac{(iB-A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2a^{\frac{3}{2}}} + \frac{iA\sqrt{a+ia \tan(dx+c)}}{2a \tan(dx+c)} - \frac{(-2iB+A) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{\sqrt{a}}\right)}{a \cdot 2\sqrt{a}} \right)$	114

```
input int(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_RETUR
NVERBOSE)
```

$$3.72. \int \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

output $2*I/d*a^2*(-1/2*(-A+I*B)/a^{(3/2)}*2^{(1/2)}*\operatorname{arctanh}(1/2*(a+I*a*\tan(dx+c))^{(1/2)}*2^{(1/2)}/a^{(1/2)}))+1/a*(1/2*I*A*(a+I*a*\tan(dx+c))^{(1/2)}/a/\tan(dx+c)-1/2*(A-2*I*B)/a^{(1/2)}*\operatorname{arctanh}((a+I*a*\tan(dx+c))^{(1/2)}/a^{(1/2)}))$

3.72.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 649 vs. $2(95) = 190$.

Time = 0.27 (sec) , antiderivative size = 649, normalized size of antiderivative = 5.28

$$\int \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx =$$

$$2\sqrt{2}(de^{(2i dx + 2i c)} - d) \sqrt{-\frac{(A^2 - 2i AB - B^2)a}{d^2}} \log \left(-\frac{4 \left((-i A - B) a e^{(i dx + i c)} + (d e^{(2i dx + 2i c)} + d) \sqrt{-\frac{(A^2 - 2i AB - B^2)a}{d^2}} \sqrt{e^{(2i c)}} \right)}{i A + B} \right)$$

input `integrate(cot(dx+c)^2*(a+I*a*tan(dx+c))^(1/2)*(A+B*tan(dx+c)),x, algorithm="fricas")`

output $-1/4*(2*\sqrt{2}*(d*e^{(2*I*d*x + 2*I*c)} - d)*\sqrt{-(A^2 - 2*I*A*B - B^2)*a/d^2}*\log(-4*((-I*A - B)*a*e^{(I*d*x + I*c)} + (d*e^{(2*I*d*x + 2*I*c)} + d)*\sqrt{-(A^2 - 2*I*A*B - B^2)*a/d^2}*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}))e^{(-I*d*x - I*c)/(I*A + B)} - 2*\sqrt{2}*(d*e^{(2*I*d*x + 2*I*c)} - d)*\sqrt{-(A^2 - 2*I*A*B - B^2)*a/d^2}*\log(-4*((-I*A - B)*a*e^{(I*d*x + I*c)} - (d*e^{(2*I*d*x + 2*I*c)} + d)*\sqrt{-(A^2 - 2*I*A*B - B^2)*a/d^2}*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}))e^{(-I*d*x - I*c)/(I*A + B)} - (d*e^{(2*I*d*x + 2*I*c)} - d)*\sqrt{-(A^2 - 4*I*A*B - 4*B^2)*a/d^2}*\log(-16*(3*(-I*A - 2*B)*a^2*e^{(2*I*d*x + 2*I*c)} + (-I*A - 2*B)*a^2 + 2*\sqrt{2}*(a*d*e^{(3*I*d*x + 3*I*c)} + a*d*e^{(I*d*x + I*c)}))*\sqrt{-(A^2 - 4*I*A*B - 4*B^2)*a/d^2}*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}))e^{(-2*I*d*x - 2*I*c)/(I*A + 2*B)} + (d*e^{(2*I*d*x + 2*I*c)} - d)*\sqrt{-(A^2 - 4*I*A*B - 4*B^2)*a/d^2}*\log(-16*(3*(-I*A - 2*B)*a^2*e^{(2*I*d*x + 2*I*c)} + (-I*A - 2*B)*a^2 - 2*\sqrt{2}*(a*d*e^{(3*I*d*x + 3*I*c)} + a*d*e^{(I*d*x + I*c)}))*\sqrt{-(A^2 - 4*I*A*B - 4*B^2)*a/d^2}*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}))e^{(-2*I*d*x - 2*I*c)/(I*A + 2*B)} + 4*\sqrt{2}*(I*A*e^{(3*I*d*x + 3*I*c)} + I*A*e^{(I*d*x + I*c)})*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})/(d*e^{(2*I*d*x + 2*I*c)} - d)$

3.72.6 Sympy [F]

$$\begin{aligned} & \int \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= \int \sqrt{ia (\tan(c + dx) - i)} (A + B \tan(c + dx)) \cot^2(c + dx) dx \end{aligned}$$

input `integrate(cot(d*x+c)**2*(a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))*cot(c + d*x)**2, x)`

3.72.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.18

$$\begin{aligned} & \int \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx = \\ & \frac{i \left(\frac{\sqrt{2}(A - iB) \log\left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c) + a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c) + a}}\right)}{\sqrt{a}} - \frac{(A - 2iB) \log\left(\frac{\sqrt{ia \tan(dx+c) + a} - \sqrt{a}}{\sqrt{ia \tan(dx+c) + a} + \sqrt{a}}\right)}{\sqrt{a}} - \frac{2i \sqrt{ia \tan(dx+c) + a} A}{a \tan(dx+c)} \right)}{2d} a \end{aligned}$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*I*(sqrt(2)*(A - I*B)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a)))/sqrt(a) - (A - 2*I*B)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a)))/sqrt(a) - 2*I*sqrt(I*a*tan(d*x + c) + a)*A/(a*tan(d*x + c))*a/d`

3.72.8 Giac [F]

$$\int \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{ia \tan(dx + c) + a} \cot(dx + c)^2 dx$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)*cot(d*x + c)^2, x)`

3.72.9 Mupad [B] (verification not implemented)

Time = 8.36 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.37

$$\int \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx =$$

$$\frac{\cot(c + dx) \left(A \sqrt{a + a \tan(c + dx)} \operatorname{li} + A \sqrt{a} \tan(c + dx) \operatorname{atanh} \left(\frac{\sqrt{a + a \tan(c + dx)} \operatorname{li}}{\sqrt{a}} \right) \operatorname{li} + 2 B \sqrt{a} \tan(c + dx) \right)}{d}$$

input `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*I)^(1/2),x)`

output `-(cot(c + d*x)*(A*(a + a*tan(c + d*x)*I)^(1/2) + A*a^(1/2)*tan(c + d*x)*a*tanh((a + a*tan(c + d*x)*I)^(1/2)/a^(1/2))*I + 2*B*a^(1/2)*tan(c + d*x)*atanh((a + a*tan(c + d*x)*I)^(1/2)/a^(1/2)) - 2^(1/2)*A*a^(1/2)*atanh((2^(1/2)*(a + a*tan(c + d*x)*I)^(1/2))/(2*a^(1/2)))*tan(c + d*x)*I - 2^(1/2)*B*a^(1/2)*atanh((2^(1/2)*(a + a*tan(c + d*x)*I)^(1/2))/(2*a^(1/2)))*tan(c + d*x)))/d`

3.73 $\int \cot^3(c+dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.73.1 Optimal result	852
3.73.2 Mathematica [A] (verified)	852
3.73.3 Rubi [A] (verified)	853
3.73.4 Maple [A] (verified)	857
3.73.5 Fricas [B] (verification not implemented)	858
3.73.6 Sympy [F]	859
3.73.7 Maxima [A] (verification not implemented)	859
3.73.8 Giac [F]	860
3.73.9 Mupad [B] (verification not implemented)	860

3.73.1 Optimal result

Integrand size = 36, antiderivative size = 169

$$\int \cot^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\sqrt{a}(7A - 4iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{4d} - \frac{\sqrt{2}\sqrt{a}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}$$

$$- \frac{(iA + 4B) \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{4d} - \frac{A \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d}$$

```
output 1/4*(7*A-4*I*B)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))*a^(1/2)/d-(A-I*B)
)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)*a^(1/2)/d-
1/4*(I*A+4*B)*cot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)/d-1/2*A*cot(d*x+c)^2*(a+
I*a*tan(d*x+c))^(1/2)/d
```

3.73.2 Mathematica [A] (verified)

Time = 2.94 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.82

$$\int \cot^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\sqrt{a}(7A - 4iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) - 4\sqrt{2}\sqrt{a}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) - \cot(c + dx)(iA + B \tan(c + dx)) \sqrt{a + ia \tan(c + dx)}}{4d}$$

input `Integrate[Cot[c + d*x]^3*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[a]*(7*A - (4*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] - 4*Sqrt[2]*Sqrt[a]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])]) - Cot[c + d*x]*(I*A + 4*B + 2*A*Cot[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]]/(4*d)`

3.73.3 Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.09, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3042, 4081, 27, 3042, 4081, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx))}{\tan(c + dx)^3} dx \\
 & \quad \downarrow \text{4081} \\
 & \frac{\int \frac{1}{2} \cot^2(c + dx) \sqrt{i \tan(c + dx) a + a(a(iA + 4B) - 3aA \tan(c + dx))} dx}{\frac{2a}{A \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \cot^2(c + dx) \sqrt{i \tan(c + dx) a + a(a(iA + 4B) - 3aA \tan(c + dx))} dx}{\frac{4a}{A \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\sqrt{i \tan(c + dx) a + a(a(iA + 4B) - 3aA \tan(c + dx))}}{\tan(c + dx)^2} dx}{4a} - \frac{A \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} \\
 & \quad \downarrow \text{4081}
 \end{aligned}$$

3.73. $\int \cot^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

$$\frac{\int -\frac{1}{2} \cot(c+dx) \sqrt{i \tan(c+dx)a+a} ((7A-4iB)a^2+(iA+4B) \tan(c+dx)a^2) dx}{a} - \frac{a(4B+iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{4a}{2d} \frac{A \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 27

$$\frac{\int \cot(c+dx) \sqrt{i \tan(c+dx)a+a} ((7A-4iB)a^2+(iA+4B) \tan(c+dx)a^2) dx}{2a} - \frac{a(4B+iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{4a}{2d} \frac{A \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 3042

$$\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} ((7A-4iB)a^2+(iA+4B) \tan(c+dx)a^2)}{\tan(c+dx)} dx}{2a} - \frac{a(4B+iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{4a}{2d} \frac{A \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 4083

$$\frac{8a^2(B+iA) \int \sqrt{i \tan(c+dx)a+adx+a} (7A-4iB) \int \cot(c+dx) (a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{2a} - \frac{a(4B+iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{4a}{2d} \frac{A \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 3042

$$\frac{8a^2(B+iA) \int \sqrt{i \tan(c+dx)a+adx+a} (7A-4iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{\tan(c+dx)} dx}{2a} - \frac{a(4B+iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{4a}{2d} \frac{A \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 3961

$$\frac{a(7A-4iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{\tan(c+dx)} dx - \frac{16ia^3(B+iA) \int \frac{1}{a-ia \tan(c+dx)} d \sqrt{i \tan(c+dx)a+adx}}{2a}}{2a} - \frac{a(4B+iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{4a}{2d} \frac{A \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 219

3.73. $\int \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx$

$$\frac{a(7A-4iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{8i\sqrt{2}a^{5/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{a(4B+iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}}{2a} = \frac{A \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{4a}{2d}$$

↓ 4082

$$\frac{a^3(7A-4iB) \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx) - \frac{8i\sqrt{2}a^{5/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{a(4B+iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}}{2a} = \frac{A \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{4a}{2d}$$

↓ 73

$$\frac{2ia^2(7A-4iB) \int \frac{1}{i - \frac{i \tan(c+dx)a+a}{a}} d \sqrt{i \tan(c+dx)a+a} - \frac{8i\sqrt{2}a^{5/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{a(4B+iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}}{2a} = \frac{A \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{4a}{2d}$$

↓ 221

$$\frac{-\frac{2a^{5/2}(7A-4iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{d} - \frac{8i\sqrt{2}a^{5/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{a(4B+iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}}{2a} = \frac{A \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

```
input Int[Cot[c + d*x]^3*sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]
```

```
output -1/2*(A*Cot[c + d*x]^2*sqrt[a + I*a*Tan[c + d*x]])/d + (-1/2*((-2*a^(5/2)*(7*A - (4*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d - ((8*I)*Sqrt[2]*a^(5/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d)/a - (a*(I*A + 4*B)*Cot[c + d*x]*sqrt[a + I*a*Tan[c + d*x]])/d)/(4*a)
```

3.73. $\int \cot^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.73.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`
- rule 4081 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.73.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.86

method	result
derivativedivides	$2a^3 \left(-\frac{(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}}\right)}{2a^{\frac{5}{2}}} + \frac{-\left(-\frac{iB}{2} + \frac{A}{8}\right)(a+ia \tan(dx+c))^{\frac{3}{2}} + \left(\frac{1}{2}iaB + \frac{1}{8}aA\right)\sqrt{a+ia \tan(dx+c)}}{a^2 \tan(dx+c)^2} + \frac{(-4d)}{a^2} \right)$
default	$2a^3 \left(-\frac{(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}}\right)}{2a^{\frac{5}{2}}} + \frac{-\left(-\frac{iB}{2} + \frac{A}{8}\right)(a+ia \tan(dx+c))^{\frac{3}{2}} + \left(\frac{1}{2}iaB + \frac{1}{8}aA\right)\sqrt{a+ia \tan(dx+c)}}{a^2 \tan(dx+c)^2} + \frac{(-4d)}{a^2} \right)$

input `int(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `2/d*a^3*(-1/2*(A-I*B)/a^(5/2)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))+1/a^2*(-((-1/2*I*B+1/8*A)*(a+I*a*tan(d*x+c))^(3/2)+(1/2*I*a*B+1/8*a*A)*(a+I*a*tan(d*x+c))^(1/2))/a^2/tan(d*x+c)^2+1/8*(7*A-4*I*B)/a^(1/2)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2)))`

$$3.73. \int \cot^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

3.73.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 730 vs. $2(130) = 260$.

Time = 0.27 (sec) , antiderivative size = 730, normalized size of antiderivative = 4.32

$$\int \cot^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx =$$

$$8\sqrt{2} \left(de^{(4i dx + 4i c)} - 2de^{(2i dx + 2i c)} + d \right) \sqrt{\frac{(A^2 - 2i AB - B^2)a}{d^2}} \log \left(-\frac{4 \left((-i A - B) a e^{(i dx + i c)} - (i de^{(2i dx + 2i c)} + i d) \sqrt{\frac{(A^2 - 2i AB - B^2)a}{d^2}} \right)}{i A + B} \right)$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output

```
-1/16*(8*sqrt(2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt((A^2 - 2*I*A*B - B^2)*a/d^2)*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) - (I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt((A^2 - 2*I*A*B - B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/(I*A + B) - 8*sqrt(2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt((A^2 - 2*I*A*B - B^2)*a/d^2)*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) - (-I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt((A^2 - 2*I*A*B - B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/(I*A + B) + (d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt((49*A^2 - 56*I*A*B - 16*B^2)*a/d^2)*log(-16*(3*(-7*I*A - 4*B)*a^2*e^(2*I*d*x + 2*I*c) + (-7*I*A - 4*B)*a^2 + 2*sqrt(2)*(I*a*d*e^(3*I*d*x + 3*I*c) + I*a*d*e^(I*d*x + I*c))*sqrt((49*A^2 - 56*I*A*B - 16*B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/(7*I*A + 4*B) - (d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt((49*A^2 - 56*I*A*B - 16*B^2)*a/d^2)*log(-16*(3*(-7*I*A - 4*B)*a^2*e^(2*I*d*x + 2*I*c) + (-7*I*A - 4*B)*a^2 + 2*sqrt(2)*(-I*a*d*e^(3*I*d*x + 3*I*c) - I*a*d*e^(I*d*x + I*c))*sqrt((49*A^2 - 56*I*A*B - 16*B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/(7*I*A + 4*B) - 4*sqrt(2)*((3*A - 4*I*B)*e^(5*I*d*x + 5*I*c) + 4*A*e^(3*I*d*x + 3*I*c) + (A + 4*I*B)*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)
```

3.73.6 Sympy [F]

$$\int \cot^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \sqrt{ia (\tan(c + dx) - i)} (A + B \tan(c + dx)) \cot^3(c + dx) dx$$

input `integrate(cot(d*x+c)**3*(a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))*cot(c + d*x)**3, x)`

3.73.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.20

$$\int \cot^3(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{a^2 \left(\frac{4\sqrt{2}(A-iB) \log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia \tan(dx+c)+a}}\right)}{a^{\frac{3}{2}}} - \frac{(7A-4iB) \log\left(\frac{\sqrt{ia \tan(dx+c)+a}-\sqrt{a}}{\sqrt{ia \tan(dx+c)+a}+\sqrt{a}}\right)}{a^{\frac{3}{2}}} + \frac{2\left((ia \tan(dx+c)+a)^{\frac{3}{2}}(A-4iB)+\sqrt{ia \tan(dx+c)+a}\right)}{(ia \tan(dx+c)+a)^2 a - 2(ia \tan(dx+c)+a)} \right)}{8d}$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/8*a^2*(4*sqrt(2)*(A - I*B)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a)))/a^(3/2) - (7*A - 4*I*B)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a)))/a^(3/2) + 2*((I*a*tan(d*x + c) + a)^(3/2)*(A - 4*I*B) + sqrt(I*a*tan(d*x + c) + a)*(A + 4*I*B)*a)/((I*a*tan(d*x + c) + a)^2*a - 2*(I*a*tan(d*x + c) + a)*a^2 + a^3))/d`

3.73.8 Giac [F]

$$\int \cot^3(c+dx)\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$= \int (B\tan(dx+c)+A)\sqrt{ia\tan(dx+c)+a}\cot(dx+c)^3dx$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x+c)+A)*sqrt(I*a*tan(d*x+c)+a)*cot(d*x+c)^3,x)`

3.73.9 Mupad [B] (verification not implemented)

Time = 7.85 (sec) , antiderivative size = 702, normalized size of antiderivative = 4.15

$$\int \cot^3(c+dx)\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$= \frac{\frac{(Aa^2+Ba^24i)\sqrt{a+a\tan(c+dx)}\operatorname{li}}{4d} + \frac{(Aa-Ba4i)(a+a\tan(c+dx)\operatorname{li})^{3/2}}{4d}}{(a+a\tan(c+dx)\operatorname{li})^2 - 2a(a+a\tan(c+dx)\operatorname{li}) + a^2}$$

$$- \frac{\operatorname{atan}\left(\frac{17A^3a^4d\sqrt{-\frac{a}{2}}\sqrt{a+a\tan(c+dx)}\operatorname{li}}{17dA^3a^5-9idA^2Ba^5+24dAB^2a^5-16idB^3a^5} - \frac{B^3a^4d\sqrt{-\frac{a}{2}}\sqrt{a+a\tan(c+dx)}\operatorname{li}16i}{17dA^3a^5-9idA^2Ba^5+24dAB^2a^5-16idB^3a^5} + \frac{24AB^2a^4d\sqrt{-\frac{a}{2}}\sqrt{a+a\tan(c+dx)}\operatorname{li}16i}{17dA^3a^5-9idA^2Ba^5+24dAB^2a^5-16idB^3a^5}\right)}{d}$$

$$+ \frac{\sqrt{-a}\operatorname{atan}\left(\frac{119A^3(-a)^{9/2}d\sqrt{a+a\tan(c+dx)}\operatorname{li}}{4\left(\frac{119dA^3a^5}{4}-3idA^2Ba^5+36dAB^2a^5-16idB^3a^5\right)} - \frac{B^3(-a)^{9/2}d\sqrt{a+a\tan(c+dx)}\operatorname{li}16i}{\frac{119dA^3a^5}{4}-3idA^2Ba^5+36dAB^2a^5-16idB^3a^5} + \frac{36dAB^2a^4d\sqrt{-\frac{a}{2}}\sqrt{a+a\tan(c+dx)}\operatorname{li}16i}{\frac{119dA^3a^5}{4}-3idA^2Ba^5+36dAB^2a^5-16idB^3a^5}\right)}{4d}$$

input `int(cot(c+d*x)^3*(A+B*tan(c+d*x))*(a+a*tan(c+d*x)*1i)^(1/2),x)`

output

$$\begin{aligned}
& \left(\frac{((Aa^2 + B^2a^4i)(a + a\tan(c + dx)1i)^{1/2})}{(4d)} + \frac{((Aa - B^2a^4i)(a + a\tan(c + dx)1i)^{3/2})}{(4d)} \right) / \left((a + a\tan(c + dx)1i)^2 - 2a(a + a\tan(c + dx)1i) + a^2 \right) - \frac{\operatorname{atan}\left(\frac{17A^3a^4d(-a/2)^{1/2}(a + a\tan(c + dx)1i)^{1/2}}{17A^3a^5d - B^3a^5d16i + 24AB^2a^5d - A^2B^2a^5d9i}\right) - (B^3a^4d(-a/2)^{1/2}(a + a\tan(c + dx)1i)^{1/2}16i)}{(17A^3a^5d - B^3a^5d16i + 24AB^2a^5d - A^2B^2a^5d9i)} + \frac{(24AB^2a^4d(-a/2)^{1/2}(a + a\tan(c + dx)1i)^{1/2})}{(17A^3a^5d - B^3a^5d16i + 24AB^2a^5d - A^2B^2a^5d9i)} - \frac{(A^2B^2a^4d(-a/2)^{1/2}(a + a\tan(c + dx)1i)^{1/2}9i)}{(17A^3a^5d - B^3a^5d16i + 24AB^2a^5d - A^2B^2a^5d9i)) * (A1i + B) * (-a/2)^{1/2} * 2i}{d} + \frac{(-a)^{1/2} * \operatorname{atan}\left(\frac{119A^3(-a)^{9/2}d(a + a\tan(c + dx)1i)^{1/2}}{4(119A^3a^5d/4 - B^3a^5d16i + 36AB^2a^5d - A^2B^2a^5d3i)}\right) - (B^3(-a)^{9/2}d(a + a\tan(c + dx)1i)^{1/2}16i)}{\left(\frac{119A^3a^5d}{4} - B^3a^5d16i + 36AB^2a^5d - A^2B^2a^5d3i\right) + (36AB^2(-a)^{9/2}d(a + a\tan(c + dx)1i)^{1/2})}{\left(\frac{119A^3a^5d}{4} - B^3a^5d16i + 36AB^2a^5d - A^2B^2a^5d3i\right) - (A^2B(-a)^{9/2}d(a + a\tan(c + dx)1i)^{1/2}3i)} \left(\frac{119A^3a^5d}{4} - B^3a^5d16i + 36AB^2a^5d - A^2B^2a^5d3i \right) * (A7i + 4B) * 1i}{(4d)}
\end{aligned}$$

3.74 $\int \cot^4(c+dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.74.1 Optimal result	862
3.74.2 Mathematica [A] (verified)	863
3.74.3 Rubi [A] (verified)	863
3.74.4 Maple [A] (verified)	868
3.74.5 Fricas [B] (verification not implemented)	869
3.74.6 Sympy [F]	870
3.74.7 Maxima [A] (verification not implemented)	870
3.74.8 Giac [F]	871
3.74.9 Mupad [B] (verification not implemented)	871

3.74.1 Optimal result

Integrand size = 36, antiderivative size = 210

$$\int \cot^4(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\sqrt{a}(9iA + 14B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{8d} - \frac{\sqrt{2}\sqrt{a}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}$$

$$+ \frac{(7A - 2iB) \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{8d}$$

$$- \frac{(iA + 6B) \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{12d} - \frac{A \cot^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{3d}$$

```
output 1/8*(9*I*A+14*B)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))*a^(1/2)/d-(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)*a^(1/2)/d+1/8*(7*A-2*I*B)*cot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)/d-1/12*(I*A+6*B)*cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(1/2)/d-1/3*A*cot(d*x+c)^3*(a+I*a*tan(d*x+c))^(1/2)/d
```

3.74.2 Mathematica [A] (verified)

Time = 4.70 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.75

$$\int \cot^4(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx = \frac{-3\sqrt{a}(9iA + 14B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) + 24\sqrt{2}\sqrt{a}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + \cot(c + dx)}{24d}$$

input `Integrate[Cot[c + d*x]^4*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `-1/24*(-3*Sqrt[a]*((9*I)*A + 14*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] + 24*Sqrt[2]*Sqrt[a]*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] + Cot[c + d*x]*(-21*A + (6*I)*B + 2*(I*A + 6*B)*Cot[c + d*x] + 8*A*Cot[c + d*x]^2)*Sqrt[a + I*a*Tan[c + d*x]])/d`

3.74.3 Rubi [A] (verified)

Time = 1.44 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.12, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.472$, Rules used = {3042, 4081, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot^4(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx))}{\tan(c + dx)^4} dx \\ & \quad \downarrow \text{4081} \\ & \int \frac{\frac{1}{2} \cot^3(c + dx) \sqrt{i \tan(c + dx) a + a} (a(iA + 6B) - 5aA \tan(c + dx)) dx}{\frac{3a}{3d} A \cot^3(c + dx) \sqrt{a + ia \tan(c + dx)}} dx \\ & \quad \downarrow \text{27} \end{aligned}$$

3.74. $\int \cot^4(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

$$\begin{aligned}
 & \frac{\int \cot^3(c+dx) \sqrt{i \tan(c+dx)a+a(a(iA+6B)-5aA \tan(c+dx))} dx}{\frac{6a}{A \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{\int \frac{\sqrt{i \tan(c+dx)a+a(a(iA+6B)-5aA \tan(c+dx))}}{\tan(c+dx)^3} dx}{6a} - \frac{A \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \\
 & \qquad \qquad \qquad \downarrow 4081 \\
 & \frac{\int -\frac{3}{2} \cot^2(c+dx) \sqrt{i \tan(c+dx)a+a((7A-2iB)a^2+(iA+6B) \tan(c+dx)a^2)} dx}{2a} - \frac{a(6B+iA) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & \frac{3 \int \cot^2(c+dx) \sqrt{i \tan(c+dx)a+a((7A-2iB)a^2+(iA+6B) \tan(c+dx)a^2)} dx}{4a} - \frac{a(6B+iA) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{3 \int \frac{\sqrt{i \tan(c+dx)a+a((7A-2iB)a^2+(iA+6B) \tan(c+dx)a^2)}}{\tan(c+dx)^2} dx}{4a} - \frac{a(6B+iA) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} \\
 & \qquad \qquad \qquad \downarrow 4081 \\
 & \frac{3 \left(\frac{\int \frac{1}{2} \cot(c+dx) \sqrt{i \tan(c+dx)a+a(a^3(9iA+14B)-a^3(7A-2iB) \tan(c+dx))} dx}{a} - \frac{a^2(7A-2iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} - \frac{a(6B+iA) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & \frac{3 \left(\frac{\int \cot(c+dx) \sqrt{i \tan(c+dx)a+a(a^3(9iA+14B)-a^3(7A-2iB) \tan(c+dx))} dx}{2a} - \frac{a^2(7A-2iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} - \frac{a(6B+iA) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} \\
 & \qquad \qquad \qquad \downarrow 3042
 \end{aligned}$$

3.74. $\int \cot^4(c+dx) \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx$

$$\frac{3 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(9iA+14B) - a^3(7A-2iB) \tan(c+dx))}{\tan(c+dx)} dx - \frac{a^2(7A-2iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} - \frac{a(6B+iA) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

$$\frac{A \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \quad 6a$$

↓ 4083

$$\frac{3 \left(\frac{a^2(14B+9iA) \int \cot(c+dx)(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{2a} - 16a^3(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+adx}}{\tan(c+dx)} dx - \frac{a^2(7A-2iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} - \frac{a(6B+iA) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

$$\frac{A \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \quad 6a$$

↓ 3042

$$\frac{3 \left(\frac{a^2(14B+9iA) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{\tan(c+dx)} dx - 16a^3(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+adx}}{\tan(c+dx)} dx - \frac{a^2(7A-2iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} - \frac{a(6B+iA) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

$$\frac{A \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \quad 6a$$

↓ 3961

$$\frac{3 \left(\frac{32ia^4(A-iB) \int \frac{1}{a-ia \tan(c+dx)} d \sqrt{i \tan(c+dx)a+adx}}{2a} + \frac{a^2(14B+9iA) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{\tan(c+dx)} dx}{2a} - \frac{a^2(7A-2iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} - \frac{a(6B+iA) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

$$\frac{A \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \quad 6a$$

↓ 219

$$\frac{3 \left(\frac{a^2(14B+9iA) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{\tan(c+dx)} dx + \frac{16i\sqrt{2}a^{7/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{2a} - \frac{a^2(7A-2iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} - \frac{a(6B+iA) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

$$\frac{A \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \quad 6a$$

↓ 4082

3.74. $\int \cot^4(c+dx) \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx$

$$\begin{aligned}
 & \frac{3 \left(\frac{a^4(14B+9iA) \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{2a} + \frac{16i\sqrt{2}a^{7/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{a^2(7A-2iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} \\
 & \frac{A \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \quad 6a \\
 & \quad \downarrow \quad 73 \\
 & \frac{3 \left(\frac{16i\sqrt{2}a^{7/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{2ia^3(14B+9iA) \int \frac{1}{i-i \tan(c+dx)a+a} d \sqrt{i \tan(c+dx)a+a}}{2a} - \frac{a^2(7A-2iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} \\
 & \frac{A \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \quad 6a \\
 & \quad \downarrow \quad 221 \\
 & \frac{3 \left(\frac{16i\sqrt{2}a^{7/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{2a^{7/2}(14B+9iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{d} - \frac{a^2(7A-2iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} \\
 & \frac{A \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \quad 6a
 \end{aligned}$$

input `Int[Cot[c + d*x]^4*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `-1/3*(A*Cot[c + d*x]^3*Sqrt[a + I*a*Tan[c + d*x]])/d + (-1/2*(a*(I*A + 6*B)*Cot[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]])/d - (3*(((-2*a^(7/2))*((9*I)*A + 14*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d + ((16*I)*Sqrt[2]*a^(7/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d)/(2*a) - (a^2*(7*A - (2*I)*B)*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]])/d)/(4*a))/(6*a)`

3.74.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`
- rule 4081 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.74.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.81

method	result
derivativedivides	$2ia^4 \left(-\frac{(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2a^{\frac{7}{2}}} - \frac{i \left(\left(-\frac{iB}{8} + \frac{7A}{16}\right)(a+ia \tan(dx+c))^{\frac{5}{2}} - \frac{5Aa(a+ia \tan(dx+c))^{\frac{3}{2}}}{6} + \left(\frac{9}{16}Aa^2\right) \right)}{a^3 \tan(dx+c)^3} \right)$
default	$2ia^4 \left(-\frac{(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2a^{\frac{7}{2}}} - \frac{i \left(\left(-\frac{iB}{8} + \frac{7A}{16}\right)(a+ia \tan(dx+c))^{\frac{5}{2}} - \frac{5Aa(a+ia \tan(dx+c))^{\frac{3}{2}}}{6} + \left(\frac{9}{16}Aa^2\right) \right)}{a^3 \tan(dx+c)^3} \right)$

input `int(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `2*I/d*a^4*(-1/2*(A-I*B)/a^(7/2)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))-1/a^3*(-I*((-1/8*I*B+7/16*A)*(a+I*a*tan(d*x+c))^(5/2)-5/6*A*a*(a+I*a*tan(d*x+c))^(3/2)+(9/16*A*a^2+1/8*I*B*a^2)*(a+I*a*tan(d*x+c))^(1/2)))/a^3/tan(d*x+c)^3-1/16*(-14*I*B+9*A)/a^(1/2)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))`

$$3.74. \int \cot^4(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

3.74.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 823 vs. $2(163) = 326$.

Time = 0.28 (sec) , antiderivative size = 823, normalized size of antiderivative = 3.92

$$\int \cot^4(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fracas")
```

```
output 1/96*(48*sqrt(2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(A^2 - 2*I*A*B - B^2)*a/d^2)*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) + (d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(A^2 - 2*I*A*B - B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/(I*A + B) - 48*sqrt(2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(A^2 - 2*I*A*B - B^2)*a/d^2)*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) - (d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(A^2 - 2*I*A*B - B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/(I*A + B) - 3*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(81*A^2 - 252*I*A*B - 196*B^2)*a/d^2)*log(-16*(3*(-9*I*A - 14*B)*a^2*e^(2*I*d*x + 2*I*c) + (-9*I*A - 14*B)*a^2 + 2*sqrt(2)*(a*d*e^(3*I*d*x + 3*I*c) + a*d*e^(I*d*x + I*c)))*sqrt(-(81*A^2 - 252*I*A*B - 196*B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x - 2*I*c)/(9*I*A + 14*B) + 3*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(81*A^2 - 252*I*A*B - 196*B^2)*a/d^2)*log(-16*(3*(-9*I*A - 14*B)*a^2*e^(2*I*d*x + 2*I*c) + (-9*I*A - 14*B)*a^2 - 2*sqrt(2)*(a*d*e^(3*I*d*x + 3*I*c) + a*d*e^(I*d*x + I*c)))*sqrt(-(81*A^2 - 252*I*A*B - 196*B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x - 2*I*c)/(9*I*A + 14*B) + 4*sqrt(2)*((31*I*A + 18*B)*e^(7*I*d*x + 7*I*c) + (5*I*A + 6*B)*e^(5*I*d*x + 5*I*c) + (I*A - 18*B)*e^(3*I*d*x + 3*I*c) - 3*(-9*I*A + ...
```

3.74.6 Sympy [F]

$$\int \cot^4(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \sqrt{ia (\tan(c + dx) - i)} (A + B \tan(c + dx)) \cot^4(c + dx) dx$$

input `integrate(cot(d*x+c)**4*(a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))*cot(c + d*x)**4, x)`

3.74.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.19

$$\int \cot^4(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{ia^3 \left(\frac{2 \left(3(ia \tan(dx+c)+a)^{\frac{5}{2}}(7A-2iB) - 40(ia \tan(dx+c)+a)^{\frac{3}{2}}Aa + 3\sqrt{ia \tan(dx+c)+a}(9A+2iB)a^2 \right)}{(ia \tan(dx+c)+a)^3 a^2 - 3(ia \tan(dx+c)+a)^2 a^3 + 3(ia \tan(dx+c)+a)a^4 - a^5} \right) + \frac{24\sqrt{2}(A-iB) \log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{i}}{\sqrt{2}\sqrt{a}+\sqrt{i}}\right)}{a^{\frac{5}{2}}}}{48d}$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/48*I*a^3*(2*(3*(I*a*tan(d*x + c) + a)^(5/2)*(7*A - 2*I*B) - 40*(I*a*tan(d*x + c) + a)^(3/2)*A*a + 3*sqrt(I*a*tan(d*x + c) + a)*(9*A + 2*I*B)*a^2)/((I*a*tan(d*x + c) + a)^3*a^2 - 3*(I*a*tan(d*x + c) + a)^2*a^3 + 3*(I*a*tan(d*x + c) + a)*a^4 - a^5) + 24*sqrt(2)*(A - I*B)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a)))/a^(5/2) - 3*(9*A - 14*I*B)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a)))/a^(5/2))/d`

3.74.8 Giac [F]

$$\int \cot^4(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{ia \tan(dx + c) + a} \cot(dx + c)^4 dx$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)*cot(d*x + c)^4, x)`

3.74.9 Mupad [B] (verification not implemented)

Time = 8.45 (sec) , antiderivative size = 735, normalized size of antiderivative = 3.50

$$\int \cot^4(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx =$$

$$\frac{\frac{(9 A a^3 + B a^3 2i) \sqrt{a + a \tan(c + dx)} \operatorname{li} \operatorname{li}}{8 d} + \frac{(7 A a - B a 2i) (a + a \tan(c + dx) \operatorname{li})^{5/2} \operatorname{li}}{8 d} - \frac{A a^2 (a + a \tan(c + dx) \operatorname{li})^{3/2} 5i}{3 d}}{3 a (a + a \tan(c + dx) \operatorname{li})^2 - 3 a^2 (a + a \tan(c + dx) \operatorname{li}) - (a + a \tan(c + dx) \operatorname{li})^3 + a^3}$$

$$+ \frac{\operatorname{atan}\left(\frac{47 \sqrt{32} A^3 a^{9/2} d \sqrt{a + a \tan(c + dx)} \operatorname{li}}{8 (47i d A^3 a^5 + 51 d A^2 B a^5 + 64i d A B^2 a^5 + 68 d B^3 a^5)} - \frac{\sqrt{32} B^3 a^{9/2} d \sqrt{a + a \tan(c + dx)} \operatorname{li} 17i}{2 (47i d A^3 a^5 + 51 d A^2 B a^5 + 64i d A B^2 a^5 + 68 d B^3 a^5)} + \frac{8 \sqrt{32}}{47i d A^3 a^5 + 51 d A^2 B a^5 + 64i d A B^2 a^5 + 68 d B^3 a^5}\right)}{d}$$

$$+ \frac{\sqrt{a} \operatorname{atan}\left(\frac{423 A^3 a^{9/2} d \sqrt{a + a \tan(c + dx)} \operatorname{li}}{8 \left(\frac{423i d A^3 a^5}{8} + \frac{347 d A^2 B a^5}{4} + \frac{139i d A B^2 a^5}{2} + 119 d B^3 a^5\right)} - \frac{B^3 a^{9/2} d \sqrt{a + a \tan(c + dx)} \operatorname{li} 119i}{423i d A^3 a^5 + 347 d A^2 B a^5 + 139i d A B^2 a^5 + 119 d B^3 a^5} + \frac{d}{2 \left(\frac{423i d A^3 a^5}{8} + \frac{347 d A^2 B a^5}{4} + \frac{139i d A B^2 a^5}{2} + 119 d B^3 a^5\right)}\right)}{8}$$

input `int(cot(c + d*x)^4*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*I)^(1/2),x)`

output

$$\begin{aligned}
& (a^{1/2} \operatorname{atan}((423A^3a^{9/2}d(a + a \tan(c + dx)1i)^{1/2}) / (8((A^3a^5d423i)/8 + 119B^3a^5d + (AB^2a^5d139i)/2 + (347A^2Ba^5d)/4))) \\
& - (B^3a^{9/2}d(a + a \tan(c + dx)1i)^{1/2}119i) / ((A^3a^5d423i)/8 + 119B^3a^5d + (AB^2a^5d139i)/2 + (347A^2Ba^5d)/4) + (139AB^2a^{9/2}d(a + a \tan(c + dx)1i)^{1/2}) / (2((A^3a^5d423i)/8 + 119B^3a^5d + (AB^2a^5d139i)/2 + (347A^2Ba^5d)/4)) - (A^2Ba^{9/2}d(a + a \tan(c + dx)1i)^{1/2}347i) / (4((A^3a^5d423i)/8 + 119B^3a^5d + (AB^2a^5d139i)/2 + (347A^2Ba^5d)/4)) * (A9i + 14B)1i / (8d) - \\
& (\operatorname{atan}((47 \cdot 32^{1/2})A^3a^{9/2}d(a + a \tan(c + dx)1i)^{1/2}) / (8(A^3a^5d47i + 68B^3a^5d + AB^2a^5d64i + 51A^2Ba^5d))) - (32^{1/2})B^3a^{9/2}d(a + a \tan(c + dx)1i)^{1/2}17i) / (2(A^3a^5d47i + 68B^3a^5d + AB^2a^5d64i + 51A^2Ba^5d)) + (8 \cdot 32^{1/2})AB^2a^{9/2}d(a + a \tan(c + dx)1i)^{1/2}) / (A^3a^5d47i + 68B^3a^5d + AB^2a^5d64i + 51A^2Ba^5d) - (32^{1/2})A^2Ba^{9/2}d(a + a \tan(c + dx)1i)^{1/2}51i) / (8(A^3a^5d47i + 68B^3a^5d + AB^2a^5d64i + 51A^2Ba^5d)) * (A1i + B)(a/32)^{1/2}8i) / d - (((9Aa^3 + Ba^32i) * (a + a \tan(c + dx)1i)^{1/2}1i) / (8d) + ((7Aa - Ba2i) * (a + a \tan(c + dx)1i)^{5/2}1i) / (8d) - (Aa^2(a + a \tan(c + dx)1i)^{3/2}5i) / (3d)) / (3a(a + a \tan(c + dx)1i)^2 - 3a^2(a + a \tan(c + dx)1i) - (a + a \tan(c + dx)1i)^3 + a^3)
\end{aligned}$$

3.75 $\int \tan^2(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.75.1	Optimal result	873
3.75.2	Mathematica [A] (verified)	874
3.75.3	Rubi [A] (verified)	874
3.75.4	Maple [A] (verified)	878
3.75.5	Fricas [B] (verification not implemented)	879
3.75.6	Sympy [F]	879
3.75.7	Maxima [A] (verification not implemented)	880
3.75.8	Giac [F]	880
3.75.9	Mupad [B] (verification not implemented)	881

3.75.1 Optimal result

Integrand size = 36, antiderivative size = 197

$$\int \tan^2(c+dx)(a + ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{2\sqrt{2}a^{3/2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{8a(7iA+8B)\sqrt{a+ia \tan(c+dx)}}{35d} + \frac{2a(7iA+8B)\tan^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{35d} + \frac{2iaB \tan^3(c+dx)\sqrt{a+ia \tan(c+dx)}}{7d} - \frac{4(21iA+19B)(a+ia \tan(c+dx))^{3/2}}{105d}$$

```
output 2*a^(3/2)*(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)/d-8/35*a*(7*I*A+8*B)*(a+I*a*tan(d*x+c))^(1/2)/d+2/35*a*(7*I*A+8*B)*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^2/d+2/7*I*a*B*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^3/d-4/105*(21*I*A+19*B)*(a+I*a*tan(d*x+c))^(3/2)/d
```


3.75.2 Mathematica [A] (verified)

Time = 1.95 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.68

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{210\sqrt{2}a^{3/2}(iA + B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + 2a\sqrt{a + ia \tan(c + dx)}(-126iA - 134B + (42A - (38I)*B)*\tan[c + d*x] + 3*((7*I)*A + 8*B)*\tan[c + d*x]^2 + (15*I)*B*\tan[c + d*x]^3)/(105*d)}{105d}$$

input `Integrate[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(210*Sqrt[2]*a^(3/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] + 2*a*Sqrt[a + I*a*Tan[c + d*x]]*((-126*I)*A - 134*B + (42*A - (38*I)*B)*Tan[c + d*x] + 3*((7*I)*A + 8*B)*Tan[c + d*x]^2 + (15*I)*B*Tan[c + d*x]^3)/(105*d)`

3.75.3 Rubi [A] (verified)

Time = 1.08 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.08, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.361$, Rules used = {3042, 4077, 27, 3042, 4080, 25, 3042, 4075, 3042, 4010, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(c + dx)^2(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4077} \\ & \frac{2}{7} \int \frac{1}{2} \tan^2(c + dx) \sqrt{i \tan(c + dx)a + a(a(7A - 6iB) + a(7iA + 8B) \tan(c + dx))} dx + \\ & \quad \frac{2iaB \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\frac{1}{7} \int \tan^2(c + dx) \sqrt{i \tan(c + dx)a + a(a(7A - 6iB) + a(7iA + 8B) \tan(c + dx))} dx + \frac{2iaB \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d}$$

↓ 3042

$$\frac{1}{7} \int \tan(c + dx)^2 \sqrt{i \tan(c + dx)a + a(a(7A - 6iB) + a(7iA + 8B) \tan(c + dx))} dx + \frac{2iaB \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d}$$

↓ 4080

$$\frac{1}{7} \left(\frac{2 \int -\tan(c + dx) \sqrt{i \tan(c + dx)a + a(2a^2(7iA + 8B) - a^2(21A - 19iB) \tan(c + dx))} dx}{5a} + \frac{2a(8B + 7iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} + \frac{2iaB \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d} \right)$$

↓ 25

$$\frac{1}{7} \left(\frac{2a(8B + 7iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{2 \int \tan(c + dx) \sqrt{i \tan(c + dx)a + a(2a^2(7iA + 8B) - a^2(21A - 19iB) \tan(c + dx))} dx}{5a} + \frac{2iaB \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d} \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{2a(8B + 7iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{2 \int \tan(c + dx) \sqrt{i \tan(c + dx)a + a(2a^2(7iA + 8B) - a^2(21A - 19iB) \tan(c + dx))} dx}{5a} + \frac{2iaB \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d} \right)$$

↓ 4075

$$\frac{1}{7} \left(\frac{2a(8B + 7iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{2 \left(\int \sqrt{i \tan(c + dx)a + a((21A - 19iB)a^2 + 2(7iA + 8B) \tan(c + dx))} dx \right)}{5a} + \frac{2iaB \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d} \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{2a(8B + 7iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{2 \left(\int \sqrt{i \tan(c + dx) a + a} ((21A - 19iB)a^2 + 2(7iA + 8B)) \right)}{5a} \right) - \frac{2iaB \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d}$$

↓ 4010

$$\frac{1}{7} \left(\frac{2a(8B + 7iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{2 \left(35a^2(A - iB) \int \sqrt{i \tan(c + dx) a + a} dx + \frac{4a^2(8B + 7iA) \sqrt{a}}{d} \right)}{5a} \right) - \frac{2iaB \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d}$$

↓ 3042

$$\frac{1}{7} \left(\frac{2a(8B + 7iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{2 \left(35a^2(A - iB) \int \sqrt{i \tan(c + dx) a + a} dx + \frac{4a^2(8B + 7iA) \sqrt{a}}{d} \right)}{5a} \right) - \frac{2iaB \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d}$$

↓ 3961

$$\frac{1}{7} \left(\frac{2a(8B + 7iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{2 \left(-\frac{70ia^3(A - iB) \int \frac{1}{a - ia \tan(c + dx)} d \sqrt{i \tan(c + dx) a + a} + \frac{4a^2(8B + 7iA)}{d} \right)}{5a} \right) - \frac{2iaB \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d}$$

↓ 219

$$\frac{1}{7} \left(\frac{2a(8B + 7iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{2 \left(-\frac{35i\sqrt{2}a^{5/2}(A - iB) \operatorname{arctanh} \left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}} \right)}{d} + \frac{4a^2(8B + 7iA)}{d} \right)}{5a} \right) - \frac{2iaB \tan^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{7d}$$

input `Int[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

```
output ((2*I)/7)*a*B*Tan[c + d*x]^3*Sqrt[a + I*a*Tan[c + d*x]]/d + ((2*a*((7*I)
*A + 8*B)*Tan[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]])/(5*d) - (2*((-35*I)*
Sqrt[2]*a^(5/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt
[a])))/d + (4*a^2*((7*I)*A + 8*B)*Sqrt[a + I*a*Tan[c + d*x]]/d + (2*a*((2
1*I)*A + 19*B)*(a + I*a*Tan[c + d*x])^(3/2))/(3*d)))/(5*a))/7
```

3.75.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3961 Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d)
Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a
, b, c, d}, x] && EqQ[a^2 + b^2, 0]
```

```
rule 4010 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp
[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e
, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]
```

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f
*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

```
rule 4077 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan
[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c -
a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] &&
GtQ[m, 1] && !LtQ[n, -1]
```

```
rule 4080 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[
1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Sim
p[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*T
an[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]
```

3.75.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.83

method	result
derivativedivides	$\frac{2i \left(\frac{iB(a+ia \tan(dx+c))^{\frac{7}{2}}}{7} - \frac{iBa(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} - \frac{Aa(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} + \frac{iB a^2(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + ia^3 B \sqrt{a+ia \tan(dx+c)} \right)}{da^2}$
default	$\frac{2i \left(\frac{iB(a+ia \tan(dx+c))^{\frac{7}{2}}}{7} - \frac{iBa(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} - \frac{Aa(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} + \frac{iB a^2(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + ia^3 B \sqrt{a+ia \tan(dx+c)} \right)}{da^2}$
parts	$\frac{2iA \left(-\frac{(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} - a^2 \sqrt{a+ia \tan(dx+c)} + a^{\frac{5}{2}} \sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}} \right) \right)}{da} + \frac{2B \left(-\frac{(a+ia \tan(dx+c))^{\frac{7}{2}}}{7} \right)}{da}$

```
input int(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_RETUR
NVERBOSE)
```

```
output 2*I/d/a^2*(1/7*I*B*(a+I*a*tan(d*x+c))^(7/2)-1/5*I*B*a*(a+I*a*tan(d*x+c))^(
5/2)-1/5*A*a*(a+I*a*tan(d*x+c))^(5/2)+1/3*I*a^2*B*(a+I*a*tan(d*x+c))^(3/2)
+I*a^3*B*(a+I*a*tan(d*x+c))^(1/2)-A*a^3*(a+I*a*tan(d*x+c))^(1/2)+a^(7/2)*(
A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2)))
```

3.75. $\int \tan^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.75.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 467 vs. $2(152) = 304$.

Time = 0.27 (sec) , antiderivative size = 467, normalized size of antiderivative = 2.37

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx =$$

$$105 \sqrt{2} \sqrt{-\frac{(A^2 - 2iAB - B^2)a^3}{d^2}} (de^{(6i dx + 6i c)} + 3de^{(4i dx + 4i c)} + 3de^{(2i dx + 2i c)} + d) \log \left(\frac{4 \left((-iA - B)a^2 e^{(i dx + i c)} + \sqrt{-\frac{(A^2 - 2iAB - B^2)a^3}{d^2}} \right)}{\dots} \right)$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-1/105*(105*sqrt(2)*sqrt(-(A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((-I*A - B)*a^2*e^(I*d*x + I*c) + sqrt(-(A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a)) - 105*sqrt(2)*sqrt(-(A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((-I*A - B)*a^2*e^(I*d*x + I*c) - sqrt(-(A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a)) + 2*sqrt(2)*((189*I*A + 211*B)*a*e^(7*I*d*x + 7*I*c) + 7*(57*I*A + 53*B)*a*e^(5*I*d*x + 5*I*c) + 35*(9*I*A + 11*B)*a*e^(3*I*d*x + 3*I*c) + 105*(I*A + B)*a*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)`

3.75.6 Sympy [F]

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{3/2} (A + B \tan(c + dx)) \tan^2(c + dx) dx$$

input `integrate(tan(d*x+c)**2*(a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

3.75. $\int \tan^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

output `Integral((I*a*(tan(c + d*x) - I))**(3/2)*(A + B*tan(c + d*x))*tan(c + d*x)**2, x)`

3.75.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.78

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx =$$

$$\frac{i \left(105 \sqrt{2}(A - iB)a^{9/2} \log \left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}} \right) - 30i (ia \tan(dx+c) + a)^{7/2} Ba + 42 (ia \tan(dx+c) + a)^{5/2} (A + iB)a^2 - 70i (ia \tan(dx+c) + a)^{3/2} B a^3 + 210 \sqrt{ia \tan(dx+c) + a} (A - iB)a^4 \right)}{105 a^3 d}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/105*I*(105*sqrt(2)*(A - I*B)*a^(9/2)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - 30*I*(I*a*tan(d*x + c) + a)^(7/2)*B*a + 42*(I*a*tan(d*x + c) + a)^(5/2)*(A + I*B)*a^2 - 70*I*(I*a*tan(d*x + c) + a)^(3/2)*B*a^3 + 210*sqrt(I*a*tan(d*x + c) + a)*(A - I*B)*a^4)/(a^3*d)`

3.75.8 Giac [F]

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{3/2} \tan(dx + c)^2 dx$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)*tan(d*x + c)^2, x)`

3.75.9 Mupad [B] (verification not implemented)

Time = 9.20 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.07

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx =$$

$$\frac{2 B (a + a \tan(c + dx) \operatorname{li})^{3/2}}{3 d} - \frac{A a \sqrt{a + a \tan(c + dx) \operatorname{li}} 2i}{d}$$

$$- \frac{2 B a \sqrt{a + a \tan(c + dx) \operatorname{li}}}{d} - \frac{A (a + a \tan(c + dx) \operatorname{li})^{5/2} 2i}{5 a d}$$

$$+ \frac{2 B (a + a \tan(c + dx) \operatorname{li})^{5/2}}{5 a d} - \frac{2 B (a + a \tan(c + dx) \operatorname{li})^{7/2}}{7 a^2 d}$$

$$- \frac{\sqrt{2} A (-a)^{3/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2 \sqrt{-a}}\right) 2i}{d}$$

$$- \frac{\sqrt{2} B a^{3/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2 \sqrt{a}}\right) 2i}{d}$$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2),x)`output `(2*B*(a + a*tan(c + d*x)*1i)^(5/2))/(5*a*d) - (A*a*(a + a*tan(c + d*x)*1i)^(1/2)*2i)/d - (2*B*a*(a + a*tan(c + d*x)*1i)^(1/2))/d - (A*(a + a*tan(c + d*x)*1i)^(5/2)*2i)/(5*a*d) - (2*B*(a + a*tan(c + d*x)*1i)^(3/2))/(3*d) - (2*B*(a + a*tan(c + d*x)*1i)^(7/2))/(7*a^2*d) - (2^(1/2)*A*(-a)^(3/2)*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2)))*2i)/d - (2^(1/2)*B*a^(3/2)*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)*1i)/(2*a^(1/2)))*2i)/d`

3.76 $\int \tan(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.76.1	Optimal result	882
3.76.2	Mathematica [A] (verified)	882
3.76.3	Rubi [A] (verified)	883
3.76.4	Maple [A] (verified)	885
3.76.5	Fricas [B] (verification not implemented)	886
3.76.6	Sympy [F]	887
3.76.7	Maxima [A] (verification not implemented)	887
3.76.8	Giac [F]	887
3.76.9	Mupad [B] (verification not implemented)	888

3.76.1 Optimal result

Integrand size = 34, antiderivative size = 137

$$\int \tan(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx =$$

$$\frac{2\sqrt{2}a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{2a(A-iB)\sqrt{a+ia \tan(c+dx)}}{d}$$

$$+ \frac{2A(a+ia \tan(c+dx))^{3/2}}{3d} - \frac{2iB(a+ia \tan(c+dx))^{5/2}}{5ad}$$

output

```
-2*a^(3/2)*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2
^(1/2)/d+2*a*(A-I*B)*(a+I*a*tan(d*x+c))^(1/2)/d+2/3*A*(a+I*a*tan(d*x+c))^(
3/2)/d-2/5*I*B*(a+I*a*tan(d*x+c))^(5/2)/a/d
```

3.76.2 Mathematica [A] (verified)

Time = 1.05 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.83

$$\int \tan(c+dx)(a+ia \tan(c+dx))^{3/2}(A$$

$$+ B \tan(c+dx)) dx = \frac{-30\sqrt{2}a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + 2a\sqrt{a+ia \tan(c+dx)}(20A-18iB +$$

15d

input `Integrate[Tan[c + d*x]*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(-30*sqrt[2]*a^(3/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(sqrt[2]*sqrt[a])] + 2*a*sqrt[a + I*a*Tan[c + d*x]]*(20*A - (18*I)*B + ((5*I)*A + 6*B)*Tan[c + d*x] + (3*I)*B*Tan[c + d*x]^2))/(15*d)`

3.76.3 Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4075, 3042, 4010, 3042, 3959, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4075} \\
 & \int (i \tan(c + dx)a + a)^{3/2}(A \tan(c + dx) - B) dx - \frac{2iB(a + ia \tan(c + dx))^{5/2}}{5ad} \\
 & \quad \downarrow \text{3042} \\
 & \int (i \tan(c + dx)a + a)^{3/2}(A \tan(c + dx) - B) dx - \frac{2iB(a + ia \tan(c + dx))^{5/2}}{5ad} \\
 & \quad \downarrow \text{4010} \\
 & -(B + iA) \int (i \tan(c + dx)a + a)^{3/2} dx + \frac{2A(a + ia \tan(c + dx))^{3/2}}{3d} - \frac{2iB(a + ia \tan(c + dx))^{5/2}}{5ad} \\
 & \quad \downarrow \text{3042} \\
 & -(B + iA) \int (i \tan(c + dx)a + a)^{3/2} dx + \frac{2A(a + ia \tan(c + dx))^{3/2}}{3d} - \frac{2iB(a + ia \tan(c + dx))^{5/2}}{5ad} \\
 & \quad \downarrow \text{3959}
 \end{aligned}$$

$$\begin{aligned}
& -(B + iA) \left(2a \int \sqrt{i \tan(c + dx)a + adx} + \frac{2ia\sqrt{a + ia \tan(c + dx)}}{d} \right) + \\
& \quad \frac{2A(a + ia \tan(c + dx))^{3/2}}{3d} - \frac{2iB(a + ia \tan(c + dx))^{5/2}}{5ad} \\
& \quad \downarrow \text{3042} \\
& -(B + iA) \left(2a \int \sqrt{i \tan(c + dx)a + adx} + \frac{2ia\sqrt{a + ia \tan(c + dx)}}{d} \right) + \\
& \quad \frac{2A(a + ia \tan(c + dx))^{3/2}}{3d} - \frac{2iB(a + ia \tan(c + dx))^{5/2}}{5ad} \\
& \quad \downarrow \text{3961} \\
& -(B + iA) \left(\frac{2ia\sqrt{a + ia \tan(c + dx)}}{d} - \frac{4ia^2 \int \frac{1}{a - ia \tan(c + dx)} d\sqrt{i \tan(c + dx)a + a}}{d} \right) + \\
& \quad \frac{2A(a + ia \tan(c + dx))^{3/2}}{3d} - \frac{2iB(a + ia \tan(c + dx))^{5/2}}{5ad} \\
& \quad \downarrow \text{219} \\
& -(B + iA) \left(\frac{2ia\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2i\sqrt{2}a^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} \right) + \\
& \quad \frac{2A(a + ia \tan(c + dx))^{3/2}}{3d} - \frac{2iB(a + ia \tan(c + dx))^{5/2}}{5ad}
\end{aligned}$$

input `Int[Tan[c + d*x]*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(2*A*(a + I*a*Tan[c + d*x])^(3/2))/(3*d) - (((2*I)/5)*B*(a + I*a*Tan[c + d*x])^(5/2))/(a*d) - (I*A + B)*((-2*I)*Sqrt[2]*a^(3/2)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d + ((2*I)*a*Sqrt[a + I*a*Tan[c + d*x]])/d)`

3.76.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3959 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

rule 3961 `Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4010 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.76.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.90

method	result
derivativedivides	$\frac{-\frac{2iB(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} + \frac{2Aa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - 2ia^2 B \sqrt{a+ia \tan(dx+c)} + 2Aa^2 \sqrt{a+ia \tan(dx+c)} - 2a^{\frac{5}{2}}(-iB+A)\sqrt{2}}{ad}$
default	$\frac{-\frac{2iB(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} + \frac{2Aa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - 2ia^2 B \sqrt{a+ia \tan(dx+c)} + 2Aa^2 \sqrt{a+ia \tan(dx+c)} - 2a^{\frac{5}{2}}(-iB+A)\sqrt{2}}{ad}$
parts	$\frac{A \left(\frac{2(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + 2a \sqrt{a+ia \tan(dx+c)} - 2a^{\frac{3}{2}} \sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}} \right) \right)}{d} + \frac{2iB \left(-\frac{(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} \right)}{d}$

3.76. $\int \tan(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

input `int(tan(d*x+c)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)`

output `2/d/a*(-1/5*I*B*(a+I*a*tan(d*x+c))^(5/2)+1/3*A*a*(a+I*a*tan(d*x+c))^(3/2)-
I*B*a^2*(a+I*a*tan(d*x+c))^(1/2)+A*a^2*(a+I*a*tan(d*x+c))^(1/2)-a^(5/2)*(A
-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2)))`

3.76.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 415 vs. $2(104) = 208$.

Time = 0.27 (sec) , antiderivative size = 415, normalized size of antiderivative = 3.03

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx =$$

$$15\sqrt{2}\sqrt{\frac{(A^2 - 2iAB - B^2)a^3}{d^2}}(de^{4i dx + 4i c} + 2de^{2i dx + 2i c} + d) \log \left(\frac{4 \left((-iA - B)a^2 e^{i dx + i c} - \sqrt{\frac{(A^2 - 2iAB - B^2)a^3}{d^2}} (i de^{2i dx} \right)}{(-iA - B)a} \right)$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm
m="fricas")`

output `-1/15*(15*sqrt(2)*sqrt((A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(4*I*d*x + 4*I*
c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((-I*A - B)*a^2*e^(I*d*x + I*c) -
sqrt((A^2 - 2*I*A*B - B^2)*a^3/d^2)*(I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(a
/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/((-I*A - B)*a) - 15*sqrt(2)
*sqrt((A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d
*x + 2*I*c) + d)*log(4*((-I*A - B)*a^2*e^(I*d*x + I*c) - sqrt((A^2 - 2*I*A
*B - B^2)*a^3/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt(a/(e^(2*I*d*x + 2
*I*c) + 1))))*e^(-I*d*x - I*c)/((-I*A - B)*a) - 2*sqrt(2)*((25*A - 27*I*B)
*a*e^(5*I*d*x + 5*I*c) + 10*(4*A - 3*I*B)*a*e^(3*I*d*x + 3*I*c) + 15*(A -
I*B)*a*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(4*I*d*x +
4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)`

3.76. $\int \tan(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.76.6 Sympy [F]

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{3/2} (A + B \tan(c + dx)) \tan(c + dx) dx$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(3/2)*(A + B*tan(c + d*x))*tan(c + d*x), x)`

3.76.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.95

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{15\sqrt{2}(A - iB)a^{7/2} \log\left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}}\right) - 6i(ia \tan(dx+c) + a)^{5/2}Ba + 10(ia \tan(dx+c) + a)^{3/2}Aa^2 + 30\sqrt{ia \tan(dx+c) + a}(A - iB)a^3}{15a^2d}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/15*(15*sqrt(2)*(A - I*B)*a^(7/2)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - 6*I*(I*a*tan(d*x + c) + a)^(5/2)*B*a + 10*(I*a*tan(d*x + c) + a)^(3/2)*A*a^2 + 30*sqrt(I*a*tan(d*x + c) + a)*(A - I*B)*a^3)/(a^2*d)`

3.76.8 Giac [F]

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{3/2} \tan(dx + c) dx$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)*tan(d*x + c), x)`

3.76.9 Mupad [B] (verification not implemented)

Time = 7.95 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.19

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{2A(a + a \tan(c + dx) \operatorname{li})^{3/2}}{3d} + \frac{2Aa \sqrt{a + a \tan(c + dx) \operatorname{li}}}{d} - \frac{Ba \sqrt{a + a \tan(c + dx) \operatorname{li}}}{d} - \frac{B(a + a \tan(c + dx) \operatorname{li})^{5/2} 2i}{5ad} - \frac{\sqrt{2} B (-a)^{3/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2\sqrt{-a}}\right) 2i}{d} - \frac{2\sqrt{2} A a^{3/2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2\sqrt{a}}\right)}{d}$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2),x)`

output `(2*A*(a + a*tan(c + d*x)*1i)^(3/2))/(3*d) + (2*A*a*(a + a*tan(c + d*x)*1i)^(1/2))/d - (B*a*(a + a*tan(c + d*x)*1i)^(1/2)*2i)/d - (B*(a + a*tan(c + d*x)*1i)^(5/2)*2i)/(5*a*d) - (2^(1/2)*B*(-a)^(3/2)*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2)))*2i)/d - (2*2^(1/2)*A*a^(3/2)*atanh((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*a^(1/2))))/d`

3.77 $\int (a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx$

3.77.1	Optimal result	889
3.77.2	Mathematica [A] (verified)	889
3.77.3	Rubi [A] (verified)	890
3.77.4	Maple [A] (verified)	892
3.77.5	Fricas [B] (verification not implemented)	892
3.77.6	Sympy [F]	893
3.77.7	Maxima [A] (verification not implemented)	893
3.77.8	Giac [F]	894
3.77.9	Mupad [B] (verification not implemented)	894

3.77.1 Optimal result

Integrand size = 28, antiderivative size = 107

$$\int (a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx =$$

$$-\frac{2\sqrt{2}a^{3/2}(iA + B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}$$

$$+ \frac{2a(iA + B)\sqrt{a + ia \tan(c + dx)}}{d} + \frac{2B(a + ia \tan(c + dx))^{3/2}}{3d}$$

```
output -2*a^(3/2)*(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2
^(1/2)/d+2*a*(I*A+B)*(a+I*a*tan(d*x+c))^(1/2)/d+2/3*B*(a+I*a*tan(d*x+c))^(
3/2)/d
```

3.77.2 Mathematica [A] (verified)

Time = 0.56 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.92

$$\int (a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx =$$

$$\frac{-6i\sqrt{2}a^{3/2}(A - iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + 2a\sqrt{a + ia \tan(c + dx)}(3iA + 4B + ia \tan(c + dx))}{3d}$$

```
input Integrate[(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]
```


output $((-6*I)*\text{Sqrt}[2]*a^{(3/2)}*(A - I*B)*\text{ArcTanh}[\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]]/(\text{Sqrt}[2]*\text{Sqrt}[a])) + 2*a*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]*((3*I)*A + 4*B + I*B*\text{Tan}[c + d*x]))/(3*d)$

3.77.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.99, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4010, 3042, 3959, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4010} \\ & (A - iB) \int (i \tan(c + dx)a + a)^{3/2} dx + \frac{2B(a + ia \tan(c + dx))^{3/2}}{3d} \\ & \quad \downarrow \text{3042} \\ & (A - iB) \int (i \tan(c + dx)a + a)^{3/2} dx + \frac{2B(a + ia \tan(c + dx))^{3/2}}{3d} \\ & \quad \downarrow \text{3959} \\ & (A - iB) \left(2a \int \sqrt{i \tan(c + dx)a + a} dx + \frac{2ia \sqrt{a + ia \tan(c + dx)}}{d} \right) + \frac{2B(a + ia \tan(c + dx))^{3/2}}{3d} \\ & \quad \downarrow \text{3042} \\ & (A - iB) \left(2a \int \sqrt{i \tan(c + dx)a + a} dx + \frac{2ia \sqrt{a + ia \tan(c + dx)}}{d} \right) + \frac{2B(a + ia \tan(c + dx))^{3/2}}{3d} \\ & \quad \downarrow \text{3961} \\ & (A - iB) \left(\frac{2ia \sqrt{a + ia \tan(c + dx)}}{d} - \frac{4ia^2 \int \frac{1}{a - ia \tan(c + dx)} d \sqrt{i \tan(c + dx)a + a}}{d} \right) + \\ & \quad \frac{2B(a + ia \tan(c + dx))^{3/2}}{3d} \end{aligned}$$

3.77. $\int (a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx$

$$(A - iB) \left(\frac{2ia\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2i\sqrt{2}a^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} \right) + \frac{2B(a + ia \tan(c + dx))^{3/2}}{3d}$$

input `Int[(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(2*B*(a + I*a*Tan[c + d*x])^(3/2))/(3*d) + (A - I*B)*((-2*I)*Sqrt[2]*a^(3/2)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])]/d + ((2*I)*a*Sqrt[a + I*a*Tan[c + d*x]])/d)`

3.77.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3959 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

3.77.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.93

method	result
derivativedivides	$\frac{2i \left(-\frac{iB(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - iaB\sqrt{a+ia \tan(dx+c)} + aA\sqrt{a+ia \tan(dx+c)} - a^{\frac{3}{2}}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{2\sqrt{a}}\right) \right)}{d}$
default	$\frac{2i \left(-\frac{iB(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - iaB\sqrt{a+ia \tan(dx+c)} + aA\sqrt{a+ia \tan(dx+c)} - a^{\frac{3}{2}}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{2\sqrt{a}}\right) \right)}{d}$
parts	$\frac{2iAa \left(\sqrt{a+ia \tan(dx+c)} - \sqrt{a} \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}}\right) \right)}{d} + \frac{B \left(\frac{2(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + 2a\sqrt{a+ia \tan(dx+c)} \right)}{d}$

input `int((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `2*I/d*(-1/3*I*B*(a+I*a*tan(d*x+c))^(3/2)-I*a*B*(a+I*a*tan(d*x+c))^(1/2)+A*(a+I*a*tan(d*x+c))^(1/2)-a^(3/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2)))`

3.77.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 359 vs. 2(82) = 164.

Time = 0.25 (sec) , antiderivative size = 359, normalized size of antiderivative = 3.36

$$\int (a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx = \frac{3\sqrt{2}\sqrt{-\frac{(A^2-2iAB-B^2)a^3}{d^2}} (de^{(2i dx+2i c)} + d) \log \left(\frac{4 \left((-iA-B)a^2 e^{(i dx+ic)} + \sqrt{-\frac{(A^2-2iAB-B^2)a^3}{d^2}} \right) a^3}{(-iA-B)} \right)}{(-iA-B)}$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

```
output 1/3*(3*sqrt(2)*sqrt(-(A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c)
+ d)*log(4*((-I*A - B)*a^2*e^(I*d*x + I*c) + sqrt(-(A^2 - 2*I*A*B - B^2)*
a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^
(-I*d*x - I*c)/((-I*A - B)*a)) - 3*sqrt(2)*sqrt(-(A^2 - 2*I*A*B - B^2)*a^3
/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log(4*((-I*A - B)*a^2*e^(I*d*x + I*c) -
sqrt(-(A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(a/(e
^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a)) - 2*sqrt(2)*((-
3*I*A - 5*B)*a*e^(3*I*d*x + 3*I*c) + 3*(-I*A - B)*a*e^(I*d*x + I*c))*sqrt(
a/(e^(2*I*d*x + 2*I*c) + 1))/(d*e^(2*I*d*x + 2*I*c) + d)
```

3.77.6 Sympy [F]

$$\int (a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{3/2} (A + B \tan(c + dx)) dx$$

```
input integrate((a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)
```

```
output Integral((I*a*(tan(c + d*x) - I))**(3/2)*(A + B*tan(c + d*x)), x)
```

3.77.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.04

$$\int (a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx = \frac{i \left(3\sqrt{2}(A - iB)a^{5/2} \log \left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}} \right) - 2i (ia \tan(dx+c) + a)^{3/2} Ba + 6\sqrt{a} \right)}{3ad}$$

```
input integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")
```

```
output 1/3*I*(3*sqrt(2)*(A - I*B)*a^(5/2)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*
x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - 2*I*(I*a*ta
n(d*x + c) + a)^(3/2)*B*a + 6*sqrt(I*a*tan(d*x + c) + a)*(A - I*B)*a^2)/(a
*d)
```

3.77. $\int (a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx$

3.77.8 Giac [F]

$$\int (a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A) (ia \tan(dx + c) + a)^{3/2} dx$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2), x)`

3.77.9 Mupad [B] (verification not implemented)

Time = 7.87 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.30

$$\int (a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx = \frac{2B(a + a \tan(c + dx) \operatorname{li})^{3/2}}{3d} + \frac{Aa \sqrt{a + a \tan(c + dx) \operatorname{li}} \operatorname{li} 2i}{d} + \frac{2Ba \sqrt{a + a \tan(c + dx) \operatorname{li}}}{d} + \frac{\sqrt{2} A (-a)^{3/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2\sqrt{-a}}\right) 2i}{d} - \frac{2\sqrt{2} B a^{3/2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2\sqrt{a}}\right)}{d}$$

input `int((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2),x)`

output `(2*B*(a + a*tan(c + d*x)*1i)^(3/2))/(3*d) + (A*a*(a + a*tan(c + d*x)*1i)^(1/2)*2i)/d + (2*B*a*(a + a*tan(c + d*x)*1i)^(1/2))/d + (2^(1/2)*A*(-a)^(3/2)*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2)))*2i)/d - (2*2^(1/2)*B*a^(3/2)*atanh((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*a^(1/2))))/d`

3.78 $\int \cot(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.78.1	Optimal result	895
3.78.2	Mathematica [A] (verified)	895
3.78.3	Rubi [A] (verified)	896
3.78.4	Maple [A] (verified)	899
3.78.5	Fricas [B] (verification not implemented)	900
3.78.6	Sympy [F]	900
3.78.7	Maxima [A] (verification not implemented)	901
3.78.8	Giac [F]	901
3.78.9	Mupad [B] (verification not implemented)	902

3.78.1 Optimal result

Integrand size = 34, antiderivative size = 113

$$\int \cot(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx =$$

$$-\frac{2a^{3/2} A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{d}$$

$$+\frac{2\sqrt{2}a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{2iaB\sqrt{a+ia \tan(c+dx)}}{d}$$

```
output -2*a^(3/2)*A*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/d+2*a^(3/2)*(A-I*B)
*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)/d+2*I*a*B*(
a+I*a*tan(d*x+c))^(1/2)/d
```

3.78.2 Mathematica [A] (verified)

Time = 0.64 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.96

$$\int \cot(c+dx)(a+ia \tan(c+dx))^{3/2}(A$$

$$+B \tan(c+dx)) dx = \frac{-2a^{3/2} A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) + 2\sqrt{2}a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + 2iaB\sqrt{a+ia \tan(c+dx)}}{d}$$

input `Integrate[Cot[c + d*x]*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(-2*a^(3/2)*A*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] + 2*Sqrt[2]*a^(3/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] + (2*I)*a*B*Sqrt[a + I*a*Tan[c + d*x]])/d`

3.78.3 Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.324$, Rules used = {3042, 4077, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan(c + dx)} dx \\
 & \quad \downarrow \text{4077} \\
 & 2 \int \frac{1}{2} \cot(c + dx) \sqrt{i \tan(c + dx)a + a}(aA + a(iA + 2B) \tan(c + dx)) dx + \frac{2iaB \sqrt{a + ia \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{27} \\
 & \int \cot(c + dx) \sqrt{i \tan(c + dx)a + a}(aA + a(iA + 2B) \tan(c + dx)) dx + \frac{2iaB \sqrt{a + ia \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{i \tan(c + dx)a + a}(aA + a(iA + 2B) \tan(c + dx))}{\tan(c + dx)} dx + \frac{2iaB \sqrt{a + ia \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{4083} \\
 & 2a(B + iA) \int \sqrt{i \tan(c + dx)a + a} dx + A \int \cot(c + dx)(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx)a + a} dx + \frac{2iaB \sqrt{a + ia \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.78. $\int \cot(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& 2a(B + iA) \int \frac{\sqrt{i \tan(c + dx)a + a} dx}{\tan(c + dx)} + A \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} dx + \\
& \quad \frac{2iaB\sqrt{a + ia \tan(c + dx)}}{d} \\
& \quad \downarrow \text{3961} \\
& \quad \frac{4ia^2(B + iA) \int \frac{1}{a - ia \tan(c + dx)} d\sqrt{i \tan(c + dx)a + a}}{d} + \\
& \quad A \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} dx + \frac{2iaB\sqrt{a + ia \tan(c + dx)}}{d} \\
& \quad \downarrow \text{219} \\
& \quad A \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} dx - \frac{2i\sqrt{2}a^{3/2}(B + iA)\operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \\
& \quad \frac{2iaB\sqrt{a + ia \tan(c + dx)}}{d} \\
& \quad \downarrow \text{4082} \\
& \quad \frac{a^2A \int \frac{\cot(c + dx)}{\sqrt{i \tan(c + dx)a + a}} d \tan(c + dx)}{d} - \frac{2i\sqrt{2}a^{3/2}(B + iA)\operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \\
& \quad \frac{2iaB\sqrt{a + ia \tan(c + dx)}}{d} \\
& \quad \downarrow \text{73} \\
& \quad \frac{2iaA \int \frac{1}{i - \frac{i(i \tan(c + dx)a + a)}{a}} d\sqrt{i \tan(c + dx)a + a}}{d} - \frac{2i\sqrt{2}a^{3/2}(B + iA)\operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \\
& \quad \frac{2iaB\sqrt{a + ia \tan(c + dx)}}{d} \\
& \quad \downarrow \text{221} \\
& \quad \frac{2i\sqrt{2}a^{3/2}(B + iA)\operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{2a^{3/2}A\operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{a}}\right)}{d} + \\
& \quad \frac{2iaB\sqrt{a + ia \tan(c + dx)}}{d}
\end{aligned}$$

input `Int[Cot[c + d*x]*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(-2*a^(3/2)*A*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d - ((2*I)*Sqrt[2]*a^(3/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d + ((2*I)*a*B*Sqrt[a + I*a*Tan[c + d*x]])/d`

3.78.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`
- rule 4077 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int((((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.78.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$\frac{2a \left(iB \sqrt{a+ia \tan(dx+c)} + \sqrt{a} (-iB+A) \sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}} \right) - \sqrt{a} A \operatorname{arctanh} \left(\frac{\sqrt{a+ia \tan(dx+c)}}{\sqrt{a}} \right) \right)}{d}$	8
default	$\frac{2a \left(iB \sqrt{a+ia \tan(dx+c)} + \sqrt{a} (-iB+A) \sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}} \right) - \sqrt{a} A \operatorname{arctanh} \left(\frac{\sqrt{a+ia \tan(dx+c)}}{\sqrt{a}} \right) \right)}{d}$	8

input `int(cot(d*x+c)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_RETURNV ERBOSE)`

output `2/d*a*(I*B*(a+I*a*tan(d*x+c))^(1/2)+a^(1/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a +I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))-a^(1/2)*A*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2)))`

3.78. $\int \cot(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.78.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 514 vs. $2(86) = 172$.

Time = 0.26 (sec) , antiderivative size = 514, normalized size of antiderivative = 4.55

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx =$$

$$-4i \sqrt{2} B a \sqrt{\frac{a}{e^{(2i dx + 2i c)} + 1}} e^{(i dx + i c)} - 2 \sqrt{2} \sqrt{\frac{(A^2 - 2i AB - B^2)a^3}{d^2}} d \log \left(\frac{4 \left((-i A - B) a^2 e^{(i dx + i c)} - \sqrt{\frac{(A^2 - 2i AB - B^2)a^3}{d^2}} (i d e^{(i dx + i c)} - (-i A - B) a) \right)}{(-i A - B) a} \right)$$

```
input integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm
m="fricas")
```

```
output -1/2*(-4*I*sqrt(2)*B*a*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*e^(I*d*x + I*c) -
2*sqrt(2)*sqrt((A^2 - 2*I*A*B - B^2)*a^3/d^2)*d*log(4*((-I*A - B)*a^2*e^(
I*d*x + I*c) - sqrt((A^2 - 2*I*A*B - B^2)*a^3/d^2)*(I*d*e^(2*I*d*x + 2*I*c)
) + I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a
)) + 2*sqrt(2)*sqrt((A^2 - 2*I*A*B - B^2)*a^3/d^2)*d*log(4*((-I*A - B)*a^2
*e^(I*d*x + I*c) - sqrt((A^2 - 2*I*A*B - B^2)*a^3/d^2)*(-I*d*e^(2*I*d*x +
2*I*c) - I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A -
B)*a)) + sqrt(A^2*a^3/d^2)*d*log(16*(3*A*a^2*e^(2*I*d*x + 2*I*c) + A*a^2
+ 2*sqrt(2)*sqrt(A^2*a^3/d^2)*(d*e^(3*I*d*x + 3*I*c) + d*e^(I*d*x + I*c))*
sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x - 2*I*c)/A) - sqrt(A^2*a^3/
d^2)*d*log(16*(3*A*a^2*e^(2*I*d*x + 2*I*c) + A*a^2 - 2*sqrt(2)*sqrt(A^2*a^
3/d^2)*(d*e^(3*I*d*x + 3*I*c) + d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*
I*c) + 1)))e^(-2*I*d*x - 2*I*c)/A))/d
```

3.78.6 SymPy [F]

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{3/2} (A + B \tan(c + dx)) \cot(c + dx) dx$$

```
input integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)
```

3.78. $\int \cot(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

output `Integral((I*a*(tan(c + d*x) - I)**(3/2)*(A + B*tan(c + d*x))*cot(c + d*x), x)`

3.78.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.15

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{\sqrt{2}(A - iB)a^{3/2} \log\left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}}\right) - Aa^{3/2} \log\left(\frac{\sqrt{ia \tan(dx+c)+a} - \sqrt{a}}{\sqrt{ia \tan(dx+c)+a} + \sqrt{a}}\right) - 2i \sqrt{ia \tan(dx+c) + a} B}{d}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-(sqrt(2)*(A - I*B)*a^(3/2)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - A*a^(3/2)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a))) - 2*I*sqrt(I*a*tan(d*x + c) + a)*B*a)/d`

3.78.8 Giac [F]

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{3/2} \cot(dx + c) dx$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)*cot(d*x + c), x)`

3.78.9 Mupad [B] (verification not implemented)

Time = 8.01 (sec) , antiderivative size = 553, normalized size of antiderivative = 4.89

$$\int \cot(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{B a \sqrt{a+a \tan(c+dx)} \operatorname{li} 2i}{d} \\ - \frac{2 A \operatorname{atanh}\left(-\frac{32 A^3 a^6 d \sqrt{a^3} \sqrt{a+a \tan(c+dx)} \operatorname{li}}{-32 d A^3 a^8+128 i d A^2 B a^8+64 d A B^2 a^8} + \frac{64 A B^2 a^6 d \sqrt{a^3} \sqrt{a+a \tan(c+dx)} \operatorname{li}}{-32 d A^3 a^8+128 i d A^2 B a^8+64 d A B^2 a^8} + \frac{A^2 B a^6 d \sqrt{a^3} \sqrt{a+a \tan(c+dx)} \operatorname{li}}{-32 d A^3 a^8+128 i d A^2 B a^8+64 d A B^2 a^8}\right)}{d} \\ + \frac{2 \sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} A^3 a^6 d \sqrt{-a^3} \sqrt{a+a \tan(c+dx)} \operatorname{li} 16i}{32 d A^3 a^8-160 i d A^2 B a^8-192 d A B^2 a^8+64 i d B^3 a^8} - \frac{32 \sqrt{2} B^3 a^6 d \sqrt{-a^3} \sqrt{a+a \tan(c+dx)} \operatorname{li}}{32 d A^3 a^8-160 i d A^2 B a^8-192 d A B^2 a^8+64 i d B^3 a^8} - \frac{\sqrt{2} A^2 B a^6 d \sqrt{-a^3} \sqrt{a+a \tan(c+dx)} \operatorname{li}}{32 d A^3 a^8-160 i d A^2 B a^8-192 d A B^2 a^8+64 i d B^3 a^8}\right)}{d}$$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2),x)`

output `(B*a*(a + a*tan(c + d*x)*1i)^(1/2)*2i)/d - (2*A*atanh((64*A*B^2*a^6*d*(a^3)^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(64*A*B^2*a^8*d - 32*A^3*a^8*d + A^2*B*a^8*d*128i) - (32*A^3*a^6*d*(a^3)^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(64*A*B^2*a^8*d - 32*A^3*a^8*d + A^2*B*a^8*d*128i) + (A^2*B*a^6*d*(a^3)^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)*128i)/(64*A*B^2*a^8*d - 32*A^3*a^8*d + A^2*B*a^8*d*128i))*(a^3)^(1/2))/d + (2*2^(1/2)*atanh((2^(1/2)*A^3*a^6*d*(-a^3)^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)*16i)/(32*A^3*a^8*d + B^3*a^8*d*64i - 192*A*B^2*a^8*d - A^2*B*a^8*d*160i) - (32*2^(1/2)*B^3*a^6*d*(-a^3)^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(32*A^3*a^8*d + B^3*a^8*d*64i - 192*A*B^2*a^8*d - A^2*B*a^8*d*160i) - (2^(1/2)*A*B^2*a^6*d*(-a^3)^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)*96i)/(32*A^3*a^8*d + B^3*a^8*d*64i - 192*A*B^2*a^8*d - A^2*B*a^8*d*160i) + (80*2^(1/2)*A^2*B*a^6*d*(-a^3)^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(32*A^3*a^8*d + B^3*a^8*d*64i - 192*A*B^2*a^8*d - A^2*B*a^8*d*160i))*(A*1i + B)*(-a^3)^(1/2))/d`

3.79 $\int \cot^2(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.79.1	Optimal result	903
3.79.2	Mathematica [A] (verified)	903
3.79.3	Rubi [A] (verified)	904
3.79.4	Maple [A] (verified)	907
3.79.5	Fricas [B] (verification not implemented)	908
3.79.6	Sympy [F]	909
3.79.7	Maxima [A] (verification not implemented)	910
3.79.8	Giac [F]	910
3.79.9	Mupad [B] (verification not implemented)	911

3.79.1 Optimal result

Integrand size = 36, antiderivative size = 125

$$\int \cot^2(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx =$$

$$\frac{a^{3/2}(3iA+2B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{d} + \frac{2\sqrt{2}a^{3/2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{aA \cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{d}$$

```
output -a^(3/2)*(3*I*A+2*B)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/d+2*a^(3/2)
*(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)/d-a
*A*cot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)/d
```

3.79.2 Mathematica [A] (verified)

Time = 3.44 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.98

$$\int \cot^2(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx =$$

$$\frac{-ia^{3/2}(3A-2iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) + 2\sqrt{2}a^{3/2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}$$

input `Integrate[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `((-I)*a^(3/2)*(3*A - (2*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] + 2*Sqrt[2]*a^(3/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] - a*A*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]])/d`

3.79.3 Rubi [A] (verified)

Time = 0.84 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.306$, Rules used = {3042, 4076, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan(c + dx)^2} dx \\
 & \quad \downarrow \text{4076} \\
 & \int \frac{1}{2} \cot(c + dx) \sqrt{i \tan(c + dx)a + a(3iA + 2B) - a(A - 2iB) \tan(c + dx)} dx - \\
 & \quad \frac{aA \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2} \int \cot(c + dx) \sqrt{i \tan(c + dx)a + a(3iA + 2B) - a(A - 2iB) \tan(c + dx)} dx - \\
 & \quad \frac{aA \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \frac{\sqrt{i \tan(c + dx)a + a(3iA + 2B) - a(A - 2iB) \tan(c + dx)}}{\tan(c + dx)} dx - \\
 & \quad \frac{aA \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{4083}
 \end{aligned}$$

3.79. $\int \cot^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

$$\frac{1}{2} \left((2B + 3iA) \int \cot(c + dx)(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx)a + adx} - 4a(A - iB) \int \sqrt{i \tan(c + dx)a + adx} \right) - \frac{aA \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

↓ 3042

$$\frac{1}{2} \left((2B + 3iA) \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} dx - 4a(A - iB) \int \sqrt{i \tan(c + dx)a + adx} \right) - \frac{aA \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

↓ 3961

$$\frac{1}{2} \left(\frac{8ia^2(A - iB) \int \frac{1}{a - ia \tan(c + dx)} d \sqrt{i \tan(c + dx)a + a}}{d} + (2B + 3iA) \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} \right) - \frac{aA \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

↓ 219

$$\frac{1}{2} \left((2B + 3iA) \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} dx + \frac{4i\sqrt{2}a^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} \right) - \frac{aA \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

↓ 4082

$$\frac{1}{2} \left(\frac{a^2(2B + 3iA) \int \frac{\cot(c + dx)}{\sqrt{i \tan(c + dx)a + a}} d \tan(c + dx)}{d} + \frac{4i\sqrt{2}a^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} \right) - \frac{aA \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

↓ 73

$$\frac{1}{2} \left(\frac{4i\sqrt{2}a^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{2ia(2B + 3iA) \int \frac{1}{i - \frac{i \tan(c + dx)}{a}} d \sqrt{i \tan(c + dx)a + a}}{d} \right) - \frac{aA \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

↓ 221

3.79. $\int \cot^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

$$\frac{1}{2} \left(\frac{4i\sqrt{2}a^{3/2}(A - iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{2a^{3/2}(2B + 3iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{a}}\right)}{d} \right) - \frac{aA \cot(c + dx)\sqrt{a + ia \tan(c + dx)}}{d}$$

input `Int[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `((-2*a^(3/2)*((3*I)*A + 2*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]]/d + ((4*I)*Sqrt[2]*a^(3/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d)/2 - (a*A*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]])/d`

3.79.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 73 `Int[((a_) + (b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.79.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.89

method	result	size
derivativedivides	$\frac{2ia^2 \left(-\frac{(2iB-2A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2\sqrt{a}} + \frac{iA\sqrt{a+ia \tan(dx+c)}}{2a \tan(dx+c)} - \frac{(-2iB+3A) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{\sqrt{a}}\right)}{2\sqrt{a}} \right)}{d}$	1
default	$\frac{2ia^2 \left(-\frac{(2iB-2A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2\sqrt{a}} + \frac{iA\sqrt{a+ia \tan(dx+c)}}{2a \tan(dx+c)} - \frac{(-2iB+3A) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{\sqrt{a}}\right)}{2\sqrt{a}} \right)}{d}$	1

3.79. $\int \cot^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

```
input int(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 2*I/d*a^2*(-1/2*(-2*A+2*I*B)*2^(1/2)/a^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))+1/2*I*A*(a+I*a*tan(d*x+c))^(1/2)/a/tan(d*x+c)-1/2*(3*A-2*I*B)/a^(1/2)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2)))
```

3.79.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 685 vs. $2(98) = 196$.

Time = 0.28 (sec) , antiderivative size = 685, normalized size of antiderivative = 5.48

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx =$$

$$4\sqrt{2}\sqrt{-\frac{(A^2 - 2iAB - B^2)a^3}{d^2}}(de^{(2i dx + 2i c)} - d) \log \left(\frac{4 \left((-iA - B)a^2 e^{(i dx + i c)} + \sqrt{-\frac{(A^2 - 2iAB - B^2)a^3}{d^2}}(de^{(2i dx + 2i c)} + d) \sqrt{\frac{a}{e^{(2i dx + 2i c)}}}} \right)}{(-iA - B)a} \right)$$

```
input integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fracas")
```

```

output -1/4*(4*sqrt(2)*sqrt(-(A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c)
) - d)*log(4*((-I*A - B)*a^2*e^(I*d*x + I*c) + sqrt(-(A^2 - 2*I*A*B - B^2)
*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e
^(-I*d*x - I*c)/((-I*A - B)*a) - 4*sqrt(2)*sqrt(-(A^2 - 2*I*A*B - B^2)*a^
3/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log(4*((-I*A - B)*a^2*e^(I*d*x + I*c) -
sqrt(-(A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(a/(
e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a) - sqrt(-(9*A^2
- 12*I*A*B - 4*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log(-16*(3*(-3*I
*A - 2*B)*a^2*e^(2*I*d*x + 2*I*c) + (-3*I*A - 2*B)*a^2 + 2*sqrt(2)*sqrt(-(
9*A^2 - 12*I*A*B - 4*B^2)*a^3/d^2)*(d*e^(3*I*d*x + 3*I*c) + d*e^(I*d*x + I
*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x - 2*I*c)/(3*I*A + 2*B)
) + sqrt(-(9*A^2 - 12*I*A*B - 4*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*
log(-16*(3*(-3*I*A - 2*B)*a^2*e^(2*I*d*x + 2*I*c) + (-3*I*A - 2*B)*a^2 - 2
*sqrt(2)*sqrt(-(9*A^2 - 12*I*A*B - 4*B^2)*a^3/d^2)*(d*e^(3*I*d*x + 3*I*c)
+ d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x - 2*I*
c)/(3*I*A + 2*B)) + 4*sqrt(2)*(I*A*a*e^(3*I*d*x + 3*I*c) + I*A*a*e^(I*d*x
+ I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(2*I*d*x + 2*I*c) - d)

```

3.79.6 Sympy [F]

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{3/2} (A + B \tan(c + dx)) \cot^2(c + dx) dx$$

```

input integrate(cot(d*x+c)**2*(a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)

```

```

output Integral((I*a*(tan(c + d*x) - I))**(3/2)*(A + B*tan(c + d*x))*cot(c + d*x)
**2, x)

```

3.79.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.16

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx =$$

$$\frac{i \left(2 \sqrt{2}(A - i B) \sqrt{a} \log \left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}} \right) - (3A - 2i B) \sqrt{a} \log \left(\frac{\sqrt{ia \tan(dx+c)+a} - \sqrt{a}}{\sqrt{ia \tan(dx+c)+a} + \sqrt{a}} \right) - \frac{2i \sqrt{ia \tan(dx+c)+a}}{\tan(dx+c)} \right)}{2d}$$

```
input integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")
```

```
output -1/2*I*(2*sqrt(2)*(A - I*B)*sqrt(a)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - (3*A - 2*I*B)*sqrt(a)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a))) - 2*I*sqrt(I*a*tan(d*x + c) + a)*A/tan(d*x + c))*a/d
```

3.79.8 Giac [F]

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{\frac{3}{2}} \cot(dx + c)^2 dx$$

```
input integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")
```

```
output integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^2, x)
```

3.79.9 Mupad [B] (verification not implemented)

Time = 10.00 (sec) , antiderivative size = 2338, normalized size of antiderivative = 18.70

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2),x)
```

```
output - 2*atanh((6*d^4*(a + a*tan(c + d*x)*1i)^(1/2)*((3*B^2*a^3)/(2*d^2) - (17*
A^2*a^3)/(8*d^2) - ((A^4*a^18)/d^4 + (16*B^4*a^18)/d^4 - (8*A^2*B^2*a^18)/
d^4 + (A*B^3*a^18*32i)/d^4 + (A^3*B*a^18*8i)/d^4)^(1/2)/(8*a^6) + (A*B*a^3
*7i)/(2*d^2))^(1/2)*((A^4*a^18)/d^4 + (16*B^4*a^18)/d^4 - (8*A^2*B^2*a^18)
/d^4 + (A*B^3*a^18*32i)/d^4 + (A^3*B*a^18*8i)/d^4)^(1/2))/(A^3*a^11*d*10i
+ 32*B^3*a^11*d + A*B^2*a^11*d*72i - 32*A^2*B*a^11*d + A*a^2*d^3*((A^4*a^1
8)/d^4 + (16*B^4*a^18)/d^4 - (8*A^2*B^2*a^18)/d^4 + (A*B^3*a^18*32i)/d^4 +
(A^3*B*a^18*8i)/d^4)^(1/2)*2i) + (2*A^2*a^6*d^2*(a + a*tan(c + d*x)*1i)^(
1/2)*((3*B^2*a^3)/(2*d^2) - (17*A^2*a^3)/(8*d^2) - ((A^4*a^18)/d^4 + (16*B
^4*a^18)/d^4 - (8*A^2*B^2*a^18)/d^4 + (A*B^3*a^18*32i)/d^4 + (A^3*B*a^18*8
i)/d^4)^(1/2)/(8*a^6) + (A*B*a^3*7i)/(2*d^2))^(1/2))/(A^3*a^8*d*10i + 32*B
^3*a^8*d + A*B^2*a^8*d*72i - 32*A^2*B*a^8*d + (A*d^3*((A^4*a^18)/d^4 + (16
*B^4*a^18)/d^4 - (8*A^2*B^2*a^18)/d^4 + (A*B^3*a^18*32i)/d^4 + (A^3*B*a^18
*8i)/d^4)^(1/2)*2i)/a) + (8*B^2*a^6*d^2*(a + a*tan(c + d*x)*1i)^(1/2)*((3*
B^2*a^3)/(2*d^2) - (17*A^2*a^3)/(8*d^2) - ((A^4*a^18)/d^4 + (16*B^4*a^18)/
d^4 - (8*A^2*B^2*a^18)/d^4 + (A*B^3*a^18*32i)/d^4 + (A^3*B*a^18*8i)/d^4)^(
1/2)/(8*a^6) + (A*B*a^3*7i)/(2*d^2))^(1/2))/(A^3*a^8*d*10i + 32*B^3*a^8*d
+ A*B^2*a^8*d*72i - 32*A^2*B*a^8*d + (A*d^3*((A^4*a^18)/d^4 + (16*B^4*a^18
)/d^4 - (8*A^2*B^2*a^18)/d^4 + (A*B^3*a^18*32i)/d^4 + (A^3*B*a^18*8i)/d^4)
^(1/2)*2i)/a) + (A*B*a^6*d^2*(a + a*tan(c + d*x)*1i)^(1/2)*((3*B^2*a^3)...
```

3.80 $\int \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.80.1	Optimal result	912
3.80.2	Mathematica [A] (verified)	913
3.80.3	Rubi [A] (verified)	913
3.80.4	Maple [A] (verified)	918
3.80.5	Fricas [B] (verification not implemented)	918
3.80.6	Sympy [F]	919
3.80.7	Maxima [A] (verification not implemented)	920
3.80.8	Giac [F]	920
3.80.9	Mupad [B] (verification not implemented)	921

3.80.1 Optimal result

Integrand size = 36, antiderivative size = 171

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{a^{3/2}(11A-12iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{4d} - \frac{2\sqrt{2}a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{a(5iA+4B)\cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{4d} - \frac{aA \cot^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{2d}$$

output

```
1/4*a^(3/2)*(11*A-12*I*B)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/d-2*a^(3/2)*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)/d-1/4*a*(5*I*A+4*B)*cot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)/d-1/2*a*A*cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(1/2)/d
```

3.80.2 Mathematica [A] (verified)

Time = 3.15 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.81

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{a^{3/2}(11A-12iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) - 8\sqrt{2}a^{3/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{4d}$$

input `Integrate[Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(a^(3/2)*(11*A - (12*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] - 8*sqrt[2]*a^(3/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(sqrt[2]*Sqrt[a])] - a*Cot[c + d*x]*((5*I)*A + 4*B + 2*A*Cot[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]])/(4*d)`

3.80.3 Rubi [A] (verified)

Time = 1.13 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.06, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3042, 4076, 27, 3042, 4081, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan(c+dx)^3} dx \\ & \quad \downarrow \text{4076} \\ & \frac{1}{2} \int \frac{1}{2} \cot^2(c+dx) \sqrt{i \tan(c+dx)a + a(a(5iA+4B) - a(3A-4iB) \tan(c+dx))} dx - \\ & \quad \frac{aA \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\begin{aligned}
& \frac{1}{4} \int \cot^2(c+dx) \sqrt{i \tan(c+dx)a + a(a(5iA+4B) - a(3A-4iB) \tan(c+dx))} dx - \\
& \quad \frac{aA \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{4} \int \frac{\sqrt{i \tan(c+dx)a + a(a(5iA+4B) - a(3A-4iB) \tan(c+dx))}}{\tan(c+dx)^2} dx - \\
& \quad \frac{aA \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} \\
& \quad \downarrow \text{4081} \\
& \frac{1}{4} \left(\frac{\int -\frac{1}{2} \cot(c+dx) \sqrt{i \tan(c+dx)a + a((11A-12iB)a^2 + (5iA+4B) \tan(c+dx)a^2)} dx}{a} - \frac{a(4B+5iA) \cot(c+dx)}{d} \right) - \\
& \quad \frac{aA \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{4} \left(-\frac{\int \cot(c+dx) \sqrt{i \tan(c+dx)a + a((11A-12iB)a^2 + (5iA+4B) \tan(c+dx)a^2)} dx}{2a} - \frac{a(4B+5iA) \cot(c+dx)}{d} \right) - \\
& \quad \frac{aA \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{4} \left(-\frac{\int \frac{\sqrt{i \tan(c+dx)a + a((11A-12iB)a^2 + (5iA+4B) \tan(c+dx)a^2)}}{\tan(c+dx)} dx}{2a} - \frac{a(4B+5iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right) - \\
& \quad \frac{aA \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} \\
& \quad \downarrow \text{4083} \\
& \frac{1}{4} \left(-\frac{16a^2(B+iA) \int \sqrt{i \tan(c+dx)a + a} dx + a(11A-12iB) \int \cot(c+dx)(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a + a}}{2a} \right) - \\
& \quad \frac{aA \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.80. $\int \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\frac{1}{4} \left(-\frac{16a^2(B+iA) \int \sqrt{i \tan(c+dx)a+adx} + a(11A-12iB) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx}{2a} - \frac{a(4B+5iA)}{2d} \right) \frac{aA \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 3961

$$\frac{1}{4} \left(-\frac{a(11A-12iB) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{32ia^3(B+iA) \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a}}{d}}{2a} - \frac{a(4B+5iA)}{2d} \right) \frac{aA \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 219

$$\frac{1}{4} \left(-\frac{a(11A-12iB) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{16i\sqrt{2}a^{5/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{2a} - \frac{a(4B+5iA)}{2d} \right) \frac{aA \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 4082

$$\frac{1}{4} \left(-\frac{a^3(11A-12iB) \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx) - \frac{16i\sqrt{2}a^{5/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{2a} - \frac{a(4B+5iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right) \frac{aA \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 73

$$\frac{1}{4} \left(-\frac{2ia^2(11A-12iB) \int \frac{1}{i - \frac{i(i \tan(c+dx)a+a)}{a}} d\sqrt{i \tan(c+dx)a+a} - \frac{16i\sqrt{2}a^{5/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{2a} - \frac{a(4B+5iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right) \frac{aA \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 221

3.80. $\int \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\frac{1}{4} \left(-\frac{2a^{5/2}(11A-12iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{a}}\right)}{d} - \frac{16i\sqrt{2}a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{a(4B+5iA)\cot(c+dx)}{2a} - \frac{aA\cot^2(c+dx)\sqrt{a+ia\tan(c+dx)}}{2d} \right)$$

input `Int[Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `-1/2*(a*A*Cot[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]])/d + (-1/2*((-2*a^(5/2)*(11*A - (12*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d - ((16*I)*Sqrt[2]*a^(5/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d)/a - (a*((5*I)*A + 4*B)*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]])/d)/4`

3.80.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 73 `Int[((a_) + (b_)*(x_)^m)*((c_) + (d_)*(x_)^n), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntegerQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.80.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.87

method	result
derivativedivides	$2a^3 \left(-\frac{(-2iB+2A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2a^{\frac{3}{2}}} + \frac{-\left(\frac{iB}{2} + \frac{5A}{8}\right)(a+ia \tan(dx+c))^{\frac{3}{2}} + \left(\frac{1}{2}iaB - \frac{3}{8}aA\right)\sqrt{a+ia \tan(dx+c)}}{a^2 \tan(dx+c)^2} \right) + \dots$
default	$2a^3 \left(-\frac{(-2iB+2A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2a^{\frac{3}{2}}} + \frac{-\left(\frac{iB}{2} + \frac{5A}{8}\right)(a+ia \tan(dx+c))^{\frac{3}{2}} + \left(\frac{1}{2}iaB - \frac{3}{8}aA\right)\sqrt{a+ia \tan(dx+c)}}{a^2 \tan(dx+c)^2} \right) + \dots$

```
input int(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 2/d*a^3*(-1/2*(2*A-2*I*B)/a^(3/2)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))+1/a*(-((-1/2*I*B+5/8*A)*(a+I*a*tan(d*x+c))^(3/2)+(1/2*I*a*B-3/8*a*A)*(a+I*a*tan(d*x+c))^(1/2))/a^2/tan(d*x+c)^2+1/8*(11*A-12*I*B)/a^(1/2)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2)))
```

3.80.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 762 vs. 2(132) = 264.

Time = 0.27 (sec) , antiderivative size = 762, normalized size of antiderivative = 4.46

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx =$$

$$16\sqrt{2}\sqrt{\frac{(A^2-2iAB-B^2)a^3}{d^2}}(de^{(4i dx+4i c)} - 2de^{(2i dx+2i c)} + d) \log \left(\frac{4 \left((-iA-B)a^2 e^{(i dx+i c)} - \sqrt{\frac{(A^2-2iAB-B^2)a^3}{d^2}} \right) (i de^{(2i dx+2i c)} + d)}{(-iA-B)a} \right)$$

```
input integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

output

```

-1/16*(16*sqrt(2)*sqrt((A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((-I*A - B)*a^2*e^(I*d*x + I*c) - sqrt((A^2 - 2*I*A*B - B^2)*a^3/d^2)*(I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/((-I*A - B)*a)) - 16*sqrt(2)*sqrt((A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((-I*A - B)*a^2*e^(I*d*x + I*c) - sqrt((A^2 - 2*I*A*B - B^2)*a^3/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/((-I*A - B)*a)) + sqrt((121*A^2 - 264*I*A*B - 144*B^2)*a^3/d^2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log(-16*(3*(-11*I*A - 12*B)*a^2*e^(2*I*d*x + 2*I*c) + (-11*I*A - 12*B)*a^2 + 2*sqrt(2)*sqrt((121*A^2 - 264*I*A*B - 144*B^2)*a^3/d^2)*(I*d*e^(3*I*d*x + 3*I*c) + I*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/(11*I*A + 12*B)) - sqrt((121*A^2 - 264*I*A*B - 144*B^2)*a^3/d^2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log(-16*(3*(-11*I*A - 12*B)*a^2*e^(2*I*d*x + 2*I*c) + (-11*I*A - 12*B)*a^2 + 2*sqrt(2)*sqrt((121*A^2 - 264*I*A*B - 144*B^2)*a^3/d^2))*(-I*d*e^(3*I*d*x + 3*I*c) - I*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/(11*I*A + 12*B)) - 4*sqrt(2)*((7*A - 4*I*B)*a*e^(5*I*d*x + 5*I*c) + 4*A*a*e^(3*I*d*x + 3*I*c) - (3*A - 4*I*B)*a*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)

```

3.80.6 Sympy [F]

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{3/2}(A + B \tan(c + dx)) \cot^3(c + dx) dx$$

input `integrate(cot(d*x+c)**3*(a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(3/2)*(A + B*tan(c + d*x))*cot(c + d*x)**3, x)`

3.80.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.19

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}(A + B \tan(c+dx)) dx = \frac{\left(\frac{8\sqrt{2}(A-iB) \log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia \tan(dx+c)+a}}\right)}{\sqrt{a}} - \frac{(11A-12iB) \log\left(\frac{\sqrt{ia \tan(dx+c)+a}-\sqrt{a}}{\sqrt{ia \tan(dx+c)+a}+\sqrt{a}}\right)}{\sqrt{a}} + \frac{2((ia \tan(dx+c)+a)}{(ia \tan(dx+c)+a)} \right)}{8d}$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/8*(8*sqrt(2)*(A - I*B)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a)))/sqrt(a) - (11*A - 12*I*B)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a)))/sqrt(a) + 2*((I*a*tan(d*x + c) + a)^(3/2)*(5*A - 4*I*B) - sqrt(I*a*tan(d*x + c) + a)*(3*A - 4*I*B)*a)/((I*a*tan(d*x + c) + a)^2 - 2*(I*a*tan(d*x + c) + a)*a + a^2))*a^2/d`

3.80.8 Giac [F]

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}(A + B \tan(c+dx)) dx = \int (B \tan(dx+c) + A)(ia \tan(dx+c) + a)^{3/2} \cot(dx+c)^3 dx$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^3, x)`

3.80.9 Mupad [B] (verification not implemented)

Time = 9.40 (sec) , antiderivative size = 3027, normalized size of antiderivative = 17.70

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2),x)
```

```
output 2*atanh((3*d^4*(a + a*tan(c + d*x)*1i)^(1/2)*((249*A^2*a^3)/(128*d^2) - ((
49*A^4*a^18)/(4*d^4) + (64*B^4*a^18)/d^4 + (40*A^2*B^2*a^18)/d^4 + (A*B^3*
a^18*64i)/d^4 + (A^3*B*a^18*28i)/d^4)^(1/2)/(64*a^6) - (17*B^2*a^3)/(8*d^2
) - (A*B*a^3*65i)/(16*d^2))^(1/2)*((49*A^4*a^18)/(4*d^4) + (64*B^4*a^18)/d
^4 + (40*A^2*B^2*a^18)/d^4 + (A*B^3*a^18*64i)/d^4 + (A^3*B*a^18*28i)/d^4)^(
1/2))/(2*((133*A^3*a^11*d)/16 - B^3*a^11*d*20i + 29*A*B^2*a^11*d + (A^2*B
*a^11*d*3i)/4 + (3*A*a^2*d^3*((49*A^4*a^18)/(4*d^4) + (64*B^4*a^18)/d^4 +
(40*A^2*B^2*a^18)/d^4 + (A*B^3*a^18*64i)/d^4 + (A^3*B*a^18*28i)/d^4)^(1/2)
)/8 - (B*a^2*d^3*((49*A^4*a^18)/(4*d^4) + (64*B^4*a^18)/d^4 + (40*A^2*B^2*
a^18)/d^4 + (A*B^3*a^18*64i)/d^4 + (A^3*B*a^18*28i)/d^4)^(1/2)*1i)/2)) + (
7*A^2*a^6*d^2*(a + a*tan(c + d*x)*1i)^(1/2)*((249*A^2*a^3)/(128*d^2) - ((4
9*A^4*a^18)/(4*d^4) + (64*B^4*a^18)/d^4 + (40*A^2*B^2*a^18)/d^4 + (A*B^3*a
^18*64i)/d^4 + (A^3*B*a^18*28i)/d^4)^(1/2)/(64*a^6) - (17*B^2*a^3)/(8*d^2)
- (A*B*a^3*65i)/(16*d^2))^(1/2))/(4*((133*A^3*a^8*d)/16 - B^3*a^8*d*20i +
29*A*B^2*a^8*d + (A^2*B*a^8*d*3i)/4 + (3*A*d^3*((49*A^4*a^18)/(4*d^4) + (
64*B^4*a^18)/d^4 + (40*A^2*B^2*a^18)/d^4 + (A*B^3*a^18*64i)/d^4 + (A^3*B*a
^18*28i)/d^4)^(1/2))/(8*a) - (B*d^3*((49*A^4*a^18)/(4*d^4) + (64*B^4*a^18)
/d^4 + (40*A^2*B^2*a^18)/d^4 + (A*B^3*a^18*64i)/d^4 + (A^3*B*a^18*28i)/d^4
)^(1/2)*1i)/(2*a))) + (4*B^2*a^6*d^2*(a + a*tan(c + d*x)*1i)^(1/2)*((249*A
^2*a^3)/(128*d^2) - ((49*A^4*a^18)/(4*d^4) + (64*B^4*a^18)/d^4 + (40*A^...
```


3.81 $\int \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.81.1	Optimal result	922
3.81.2	Mathematica [A] (verified)	923
3.81.3	Rubi [A] (verified)	923
3.81.4	Maple [A] (verified)	929
3.81.5	Fricas [B] (verification not implemented)	929
3.81.6	Sympy [F]	930
3.81.7	Maxima [A] (verification not implemented)	931
3.81.8	Giac [F]	931
3.81.9	Mupad [B] (verification not implemented)	932

3.81.1 Optimal result

Integrand size = 36, antiderivative size = 213

$$\int \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{a^{3/2}(23iA+22B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{8d} - \frac{2\sqrt{2}a^{3/2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{a(9A-10iB)\cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{8d} - \frac{a(7iA+6B)\cot^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{12d} - \frac{aA\cot^3(c+dx)\sqrt{a+ia \tan(c+dx)}}{3d}$$

```
output 1/8*a^(3/2)*(23*I*A+22*B)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/d-2*a^(3/2)*(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)/d+1/8*a*(9*A-10*I*B)*cot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)/d-1/12*a*(7*I*A+6*B)*cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(1/2)/d-1/3*a*A*cot(d*x+c)^3*(a+I*a*tan(d*x+c))^(1/2)/d
```

3.81.2 Mathematica [A] (verified)

Time = 5.41 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.74

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx =$$

$$\frac{-3a^{3/2}(23iA + 22B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) + 48\sqrt{2}a^{3/2}(iA + B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + a \cot(c + dx)}{24d}$$

input `Integrate[Cot[c + d*x]^4*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `-1/24*(-3*a^(3/2)*((23*I)*A + 22*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] + 48*Sqrt[2]*a^(3/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] + a*Cot[c + d*x]*(-27*A + (30*I)*B + 2*((7*I)*A + 6*B)*Cot[c + d*x] + 8*A*Cot[c + d*x]^2)*Sqrt[a + I*a*Tan[c + d*x]]/d`

3.81.3 Rubi [A] (verified)

Time = 1.49 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.10, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.472$, Rules used = {3042, 4076, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan(c + dx)^4} dx$$

$$\downarrow \text{4076}$$

$$\frac{1}{3} \int \frac{1}{2} \cot^3(c + dx) \sqrt{i \tan(c + dx)a + a(a(7iA + 6B) - a(5A - 6iB) \tan(c + dx))} dx -$$

$$\frac{aA \cot^3(c + dx) \sqrt{a + ia \tan(c + dx)}}{3d}$$

$$\downarrow \text{27}$$

3.81. $\int \cot^4(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

$$\begin{aligned}
 & \frac{1}{6} \int \cot^3(c+dx) \sqrt{i \tan(c+dx)a + a(7iA+6B) - a(5A-6iB) \tan(c+dx)} dx - \\
 & \quad \frac{aA \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{6} \int \frac{\sqrt{i \tan(c+dx)a + a(7iA+6B) - a(5A-6iB) \tan(c+dx)}}{\tan(c+dx)^3} dx - \\
 & \quad \frac{aA \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \\
 & \quad \downarrow \text{4081} \\
 & \frac{1}{6} \left(\frac{\int -\frac{3}{2} \cot^2(c+dx) \sqrt{i \tan(c+dx)a + a((9A-10iB)a^2 + (7iA+6B) \tan(c+dx)a^2)} dx}{2a} - \frac{a(6B+7iA) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \right) \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{6} \left(-\frac{3 \int \cot^2(c+dx) \sqrt{i \tan(c+dx)a + a((9A-10iB)a^2 + (7iA+6B) \tan(c+dx)a^2)} dx}{4a} - \frac{a(6B+7iA) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \right) \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{6} \left(-\frac{3 \int \frac{\sqrt{i \tan(c+dx)a + a((9A-10iB)a^2 + (7iA+6B) \tan(c+dx)a^2)}}{\tan(c+dx)^2} dx}{4a} - \frac{a(6B+7iA) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} \right) \\
 & \quad \frac{aA \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \\
 & \quad \downarrow \text{4081} \\
 & \frac{1}{6} \left(-\frac{3 \left(\frac{\int \frac{1}{2} \cot(c+dx) \sqrt{i \tan(c+dx)a + a(23iA+22B) - a^3(9A-10iB) \tan(c+dx)} dx}{a} - \frac{a^2(9A-10iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} - \frac{aA \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \right) \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.81. $\int \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\frac{1}{6} \left(\frac{3 \left(\frac{\int \cot(c+dx) \sqrt{i \tan(c+dx) a + a} (a^3(23iA+22B) - a^3(9A-10iB) \tan(c+dx)) dx}{2a} - \frac{a^2(9A-10iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} - \frac{aA \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \right) - a(6B+7iA) \cot(c+dx)$$

↓ 3042

$$\frac{1}{6} \left(\frac{3 \left(\frac{\int \frac{\sqrt{i \tan(c+dx) a + a} (a^3(23iA+22B) - a^3(9A-10iB) \tan(c+dx))}{\tan(c+dx)} dx}{2a} - \frac{a^2(9A-10iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} - \frac{aA \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \right) - a(6B+7iA) \cot(c+dx)$$

↓ 4083

$$\frac{1}{6} \left(\frac{3 \left(\frac{a^2(22B+23iA) \int \cot(c+dx) (a-ia \tan(c+dx)) \sqrt{i \tan(c+dx) a + a dx} - 32a^3(A-iB) \int \sqrt{i \tan(c+dx) a + a dx}}{2a} - \frac{a^2(9A-10iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} - \frac{aA \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \right) - a(6B+7iA) \cot(c+dx)$$

↓ 3042

$$\frac{1}{6} \left(\frac{3 \left(\frac{a^2(22B+23iA) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx) a + a}}{\tan(c+dx)} dx - 32a^3(A-iB) \int \sqrt{i \tan(c+dx) a + a dx}}{2a} - \frac{a^2(9A-10iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} - \frac{aA \cot^3(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \right) - a(6B+7iA) \cot(c+dx)$$

↓ 3961

$$\frac{1}{6} \left(\frac{3 \left(\frac{64ia^4(A-iB) \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a} + a^2(22B+23iA) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{a^2(9A-10iB) \cot(c+dx)\sqrt{a}}{d} \right)}{4a} \right)$$

$$\frac{aA \cot^3(c+dx)\sqrt{a+ia \tan(c+dx)}}{3d}$$

↓ 219

$$\frac{1}{6} \left(\frac{3 \left(\frac{a^2(22B+23iA) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx + \frac{32i\sqrt{2}a^{7/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{a^2(9A-10iB) \cot(c+dx)\sqrt{a}}{d} \right)}{4a} \right)$$

$$\frac{aA \cot^3(c+dx)\sqrt{a+ia \tan(c+dx)}}{3d}$$

↓ 4082

$$\frac{1}{6} \left(\frac{3 \left(\frac{a^4(22B+23iA) \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx) + \frac{32i\sqrt{2}a^{7/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{a^2(9A-10iB) \cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{d} \right)}{4a} \right)$$

$$\frac{aA \cot^3(c+dx)\sqrt{a+ia \tan(c+dx)}}{3d}$$

↓ 73

$$\frac{1}{6} \left(\frac{3 \left(\frac{32i\sqrt{2}a^{7/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{2ia^3(22B+23iA) \int \frac{1}{i-i \tan(c+dx)a+a} d\sqrt{i \tan(c+dx)a+a}}{2a} - \frac{a^2(9A-10iB) \cot(c+dx)\sqrt{a}}{d} \right)}{4a} \right)$$

$$\frac{aA \cot^3(c+dx)\sqrt{a+ia \tan(c+dx)}}{3d}$$

↓ 221

3.81. $\int \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\frac{1}{6} \left(\frac{3 \left(\frac{32i\sqrt{2}a^{7/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{2a^{7/2}(22B+23iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{a}}\right)}{d} - \frac{a^2(9A-10iB)\cot(c+dx)\sqrt{a+ia\tan(c+dx)}}{d} \right)}{4a} \right) - \frac{aA \cot^3(c+dx)\sqrt{a+ia\tan(c+dx)}}{3d}$$

input `Int[Cot[c + d*x]^4*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `-1/3*(a*A*Cot[c + d*x]^3*Sqrt[a + I*a*Tan[c + d*x]])/d + (-1/2*(a*((7*I)*A + 6*B)*Cot[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]])/d - (3*(((-2*a^(7/2)*((23*I)*A + 22*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d + ((32*I)*Sqrt[2]*a^(7/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])]))/d)/(2*a) - (a^2*(9*A - (10*I)*B)*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]])/d)/(4*a))/6`

3.81.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 73 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

3.81. $\int \cot^4(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int[(((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.81.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.84

method	result
derivativedivides	$2ia^4 \left(-\frac{(-2iB+2A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}}\right)}{2a^{\frac{5}{2}}} + \frac{i\left(\left(\frac{5iB}{8} - \frac{9A}{16}\right)(a+ia \tan(dx+c))^{\frac{5}{2}} + (-iaB + \frac{5}{8}aA)(a+ia \tan(dx+c))^{\frac{5}{2}}\right)}{a^3 \tan(dx+c)^3} \right)$
default	$2ia^4 \left(-\frac{(-2iB+2A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}}\right)}{2a^{\frac{5}{2}}} + \frac{i\left(\left(\frac{5iB}{8} - \frac{9A}{16}\right)(a+ia \tan(dx+c))^{\frac{5}{2}} + (-iaB + \frac{5}{8}aA)(a+ia \tan(dx+c))^{\frac{5}{2}}\right)}{a^3 \tan(dx+c)^3} \right)$

input `int(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `2*I/d*a^4*(-1/2*(2*A-2*I*B)/a^(5/2)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))+1/a^2*(-I*((5/8*I*B-9/16*A)*(a+I*a*tan(d*x+c))^(5/2))+(-I*a*B+5/6*a*A)*(a+I*a*tan(d*x+c))^(3/2)+(3/8*I*B*a^2-7/16*A*a^2)*(a+I*a*tan(d*x+c))^(1/2))/a^3/tan(d*x+c)^3+1/16*(-22*I*B+23*A)/a^(1/2)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2)))`

3.81.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 856 vs. 2(166) = 332.

Time = 0.27 (sec) , antiderivative size = 856, normalized size of antiderivative = 4.02

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fracas")`


```

output 1/96*(96*sqrt(2)*sqrt(-(A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*log(4*((-I*A - B)*a^2*e^(I*d*x + I*c) + sqrt(-(A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/((-I*A - B)*a) - 96*sqrt(2)*sqrt(-(A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*log(4*((-I*A - B)*a^2*e^(I*d*x + I*c) - sqrt(-(A^2 - 2*I*A*B - B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/((-I*A - B)*a) - 3*sqrt(-(529*A^2 - 1012*I*A*B - 484*B^2)*a^3/d^2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*log(-16*(3*(-23*I*A - 22*B)*a^2*e^(2*I*d*x + 2*I*c) + (-23*I*A - 22*B)*a^2 + 2*sqrt(2)*sqrt(-(529*A^2 - 1012*I*A*B - 484*B^2)*a^3/d^2)*(d*e^(3*I*d*x + 3*I*c) + d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/(23*I*A + 22*B) + 3*sqrt(-(529*A^2 - 1012*I*A*B - 484*B^2)*a^3/d^2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*log(-16*(3*(-23*I*A - 22*B)*a^2*e^(2*I*d*x + 2*I*c) + (-23*I*A - 22*B)*a^2 - 2*sqrt(2)*sqrt(-(529*A^2 - 1012*I*A*B - 484*B^2)*a^3/d^2)*(d*e^(3*I*d*x + 3*I*c) + d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/(23*I*A + 22*B) - 4*sqrt(2)*(7*(-7*I*A - 6*B)*a*e^(7*I*d*x + 7*I*c) - (11*I*A - 18*B)*a*e^(5*I*d*x + 5*I*c) - (-17*...

```

3.81.6 Sympy [F]

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{3/2}(A + B \tan(c + dx)) \cot^4(c + dx) dx$$

```
input integrate(cot(d*x+c)**4*(a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)
```

```
output Integral((I*a*(tan(c + d*x) - I))**(3/2)*(A + B*tan(c + d*x))*cot(c + d*x)**4, x)
```

3.81.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.19

$$\int \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}(A + B \tan(c+dx)) dx = \frac{ia^3 \left(\frac{48\sqrt{2}(A-iB) \log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia \tan(dx+c)+a}}\right)}{a^{3/2}} - \frac{3(23A-22iB) \log\left(\frac{\sqrt{ia \tan(dx+c)+a}-\sqrt{a}}{\sqrt{ia \tan(dx+c)+a}+\sqrt{a}}\right)}{a^{3/2}} + \frac{2(3ia \tan(dx+c)+a)}{48d} \right)}{48d}$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/48*I*a^3*(48*sqrt(2)*(A - I*B)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a)))/a^(3/2) - 3*(23*A - 22*I*B)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a)))/a^(3/2) + 2*(3*(I*a*tan(d*x + c) + a)^(5/2)*(9*A - 10*I*B) - 8*(I*a*tan(d*x + c) + a)^(3/2)*(5*A - 6*I*B)*a + 3*sqrt(I*a*tan(d*x + c) + a)*(7*A - 6*I*B)*a^2)/((I*a*tan(d*x + c) + a)^3*a - 3*(I*a*tan(d*x + c) + a)^2*a^2 + 3*(I*a*tan(d*x + c) + a)*a^3 - a^4))/d`

3.81.8 Giac [F]

$$\int \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}(A + B \tan(c+dx)) dx = \int (B \tan(dx+c) + A)(ia \tan(dx+c) + a)^{3/2} \cot(dx+c)^4 dx$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^4, x)`

3.81.9 Mupad [B] (verification not implemented)

Time = 9.30 (sec) , antiderivative size = 3084, normalized size of antiderivative = 14.48

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int(cot(c + d*x)^4*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2),x)`

output `2*atanh((6*d^4*(a + a*tan(c + d*x)*1i)^(1/2)*((249*B^2*a^3)/(128*d^2) - (1041*A^2*a^3)/(512*d^2) - ((289*A^4*a^18)/(64*d^4) + (49*B^4*a^18)/(4*d^4) + (101*A^2*B^2*a^18)/(8*d^4) + (A*B^3*a^18*21i)/(2*d^4) + (A^3*B*a^18*51i)/(8*d^4))^(1/2)/(64*a^6) + (A*B*a^3*509i)/(128*d^2))^(1/2)*((289*A^4*a^18)/(64*d^4) + (49*B^4*a^18)/(4*d^4) + (101*A^2*B^2*a^18)/(8*d^4) + (A*B^3*a^18*21i)/(2*d^4) + (A^3*B*a^18*51i)/(8*d^4))^(1/2))/((A^3*a^11*d*663i)/32 + (133*B^3*a^11*d)/4 + (A*B^2*a^11*d*387i)/8 + (89*A^2*B*a^11*d)/16 + (A*a^2*d^3*((289*A^4*a^18)/(64*d^4) + (49*B^4*a^18)/(4*d^4) + (101*A^2*B^2*a^18)/(8*d^4) + (A*B^3*a^18*21i)/(2*d^4) + (A^3*B*a^18*51i)/(8*d^4))^(1/2)*7i)/4 + (3*B*a^2*d^3*((289*A^4*a^18)/(64*d^4) + (49*B^4*a^18)/(4*d^4) + (101*A^2*B^2*a^18)/(8*d^4) + (A*B^3*a^18*21i)/(2*d^4) + (A^3*B*a^18*51i)/(8*d^4))^(1/2))/2) + (17*A^2*a^6*d^2*(a + a*tan(c + d*x)*1i)^(1/2)*((249*B^2*a^3)/(128*d^2) - (1041*A^2*a^3)/(512*d^2) - ((289*A^4*a^18)/(64*d^4) + (49*B^4*a^18)/(4*d^4) + (101*A^2*B^2*a^18)/(8*d^4) + (A*B^3*a^18*21i)/(2*d^4) + (A^3*B*a^18*51i)/(8*d^4))^(1/2)/(64*a^6) + (A*B*a^3*509i)/(128*d^2))^(1/2))/(4*((A^3*a^8*d*663i)/32 + (133*B^3*a^8*d)/4 + (A*B^2*a^8*d*387i)/8 + (89*A^2*B*a^8*d)/16 + (A*d^3*((289*A^4*a^18)/(64*d^4) + (49*B^4*a^18)/(4*d^4) + (101*A^2*B^2*a^18)/(8*d^4) + (A*B^3*a^18*21i)/(2*d^4) + (A^3*B*a^18*51i)/(8*d^4))^(1/2)*7i)/(4*a) + (3*B*d^3*((289*A^4*a^18)/(64*d^4) + (49*B^4*a^18)/(4*d^4) + (101*A^2*B^2*a^18)/(8*d^4) + (A*B^3*a^18*21i)/(2*d^4) + ...`

3.82 $\int \tan^2(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.82.1	Optimal result	933
3.82.2	Mathematica [A] (verified)	934
3.82.3	Rubi [A] (verified)	934
3.82.4	Maple [A] (verified)	939
3.82.5	Fricas [B] (verification not implemented)	939
3.82.6	Sympy [F]	940
3.82.7	Maxima [A] (verification not implemented)	940
3.82.8	Giac [F]	941
3.82.9	Mupad [B] (verification not implemented)	941

3.82.1 Optimal result

Integrand size = 36, antiderivative size = 246

$$\int \tan^2(c+dx)(a + ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \frac{4\sqrt{2}a^{5/2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) - \frac{8a^2(45iA+46B)\sqrt{a+ia \tan(c+dx)}}{105d} + \frac{2a^2(45iA+46B)\tan^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{105d} - \frac{2a^2(3A-4iB)\tan^3(c+dx)\sqrt{a+ia \tan(c+dx)}}{21d} - \frac{8a(60iA+59B)(a+ia \tan(c+dx))^{3/2}}{315d} + \frac{2iaB \tan^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{9d}$$

```
output 4*a^(5/2)*(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)/d-8/105*a^2*(45*I*A+46*B)*(a+I*a*tan(d*x+c))^(1/2)/d+2/105*a^2*(45*I*A+46*B)*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^2/d-2/21*a^2*(3*A-4*I*B)*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^3/d-8/315*a*(60*I*A+59*B)*(a+I*a*tan(d*x+c))^(3/2)/d+2/9*I*a*B*tan(d*x+c)^3*(a+I*a*tan(d*x+c))^(3/2)/d
```

3.82.2 Mathematica [A] (verified)

Time = 3.62 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.61

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{2a^2 \left(630\sqrt{2}\sqrt{a}(iA + B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + \sqrt{a + ia \tan(c + dx)}(-780iA - 780B) \right)}{315d}$$

input `Integrate[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(2*a^2*(630*Sqrt[2]*Sqrt[a]*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] + Sqrt[a + I*a*Tan[c + d*x]]*((-780*I)*A - 788*B + 4*(60*A - (59*I)*B)*Tan[c + d*x] + 3*((45*I)*A + 46*B)*Tan[c + d*x]^2 + (-45*A + (95*I)*B)*Tan[c + d*x]^3 - 35*B*Tan[c + d*x]^4))/(315*d)`

3.82.3 Rubi [A] (verified)

Time = 1.43 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.08, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 4077, 27, 3042, 4077, 27, 3042, 4080, 27, 3042, 4075, 3042, 4010, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(c + dx)^2(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4077} \\ & \frac{2}{9} \int \frac{3}{2} \tan^2(c + dx)(i \tan(c + dx)a + a)^{3/2}(a(3A - 2iB) + a(3iA + 4B) \tan(c + dx)) dx + \\ & \quad \frac{2iaB \tan^3(c + dx)(a + ia \tan(c + dx))^{3/2}}{9d} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.82. $\int \tan^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\frac{1}{3} \int \tan^2(c+dx)(i \tan(c+dx)a+a)^{3/2}(a(3A-2iB)+a(3iA+4B)\tan(c+dx))dx + \frac{2iaB \tan^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{9d}$$

↓ 3042

$$\frac{1}{3} \int \tan(c+dx)^2(i \tan(c+dx)a+a)^{3/2}(a(3A-2iB)+a(3iA+4B)\tan(c+dx))dx + \frac{2iaB \tan^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{9d}$$

↓ 4077

$$\frac{1}{3} \left(\frac{2}{7} \int \frac{1}{2} \tan^2(c+dx) \sqrt{i \tan(c+dx)a+a} ((39A-38iB)a^2+(45iA+46B)\tan(c+dx)a^2) dx - \frac{2a^2(3A-4iB)t}{9d} \right) + \frac{2iaB \tan^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{9d}$$

↓ 27

$$\frac{1}{3} \left(\frac{1}{7} \int \tan^2(c+dx) \sqrt{i \tan(c+dx)a+a} ((39A-38iB)a^2+(45iA+46B)\tan(c+dx)a^2) dx - \frac{2a^2(3A-4iB)t}{9d} \right) + \frac{2iaB \tan^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{9d}$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{7} \int \tan(c+dx)^2 \sqrt{i \tan(c+dx)a+a} ((39A-38iB)a^2+(45iA+46B)\tan(c+dx)a^2) dx - \frac{2a^2(3A-4iB)t}{9d} \right) + \frac{2iaB \tan^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{9d}$$

↓ 4080

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2 \int -2 \tan(c+dx) \sqrt{i \tan(c+dx)a+a} (a^3(45iA+46B)-a^3(60A-59iB)\tan(c+dx)) dx}{5a} + \frac{2a^2(46B+45iA)\tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d} \right) - \frac{4 \int \tan(c+dx) \sqrt{i \tan(c+dx)a+a} (a^3(45iA+46B)-a^3(60A-59iB)\tan(c+dx)) dx}{5a} \right) + \frac{2iaB \tan^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{9d}$$

↓ 27

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46B+45iA)\tan^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{5d} - \frac{4 \int \tan(c+dx) \sqrt{i \tan(c+dx)a+a} (a^3(45iA+46B)-a^3(60A-59iB)\tan(c+dx)) dx}{5a} \right) \right) + \frac{2iaB \tan^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{9d}$$

3.82. $\int \tan^2(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46B + 45iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{4 \int \tan(c + dx) \sqrt{i \tan(c + dx)a + a(a^3(45iA + 46B))}}{5a} \right) - \frac{2iaB \tan^3(c + dx)(a + ia \tan(c + dx))^{3/2}}{9d} \right)$$

↓ 4075

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46B + 45iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{4 \left(\int \sqrt{i \tan(c + dx)a + a((60A - 59iB)a^3 + (45iA))} \right)}{5a} \right) - \frac{2iaB \tan^3(c + dx)(a + ia \tan(c + dx))^{3/2}}{9d} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46B + 45iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{4 \left(\int \sqrt{i \tan(c + dx)a + a((60A - 59iB)a^3 + (45iA))} \right)}{5a} \right) - \frac{2iaB \tan^3(c + dx)(a + ia \tan(c + dx))^{3/2}}{9d} \right)$$

↓ 4010

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46B + 45iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{4 \left(105a^3(A - iB) \int \sqrt{i \tan(c + dx)a + adx} + \frac{2a^3(46B + 45iA)}{3} \right)}{5a} \right) - \frac{2iaB \tan^3(c + dx)(a + ia \tan(c + dx))^{3/2}}{9d} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46B + 45iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{4 \left(105a^3(A - iB) \int \sqrt{i \tan(c + dx)a + adx} + \frac{2a^3(46B + 45iA)}{3} \right)}{5a} \right) - \frac{2iaB \tan^3(c + dx)(a + ia \tan(c + dx))^{3/2}}{9d} \right)$$

↓ 3961

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46B + 45iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{4 \left(-\frac{210ia^4(A-iB) \int \frac{1}{a-ia \tan(c+dx)} dx \sqrt{i \tan(c+dx)a+a} + 2a^3}{d} \right)}{9d} \right) \right)$$

↓ 219

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46B + 45iA) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \frac{4 \left(-\frac{105i\sqrt{2}a^{7/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + 2a^3}{d} \right)}{9d} \right) \right)$$

```
input Int[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]
```

```
output (((2*I)/9)*a*B*Tan[c + d*x]^3*(a + I*a*Tan[c + d*x])^(3/2)/d + ((-2*a^2*(3*A - (4*I)*B)*Tan[c + d*x]^3*Sqrt[a + I*a*Tan[c + d*x]])/(7*d) + ((2*a^2*((45*I)*A + 46*B)*Tan[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]])/(5*d) - (4*((-105*I)*Sqrt[2]*a^(7/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d + (2*a^3*((45*I)*A + 46*B)*Sqrt[a + I*a*Tan[c + d*x]]/d + (2*a^2*((60*I)*A + 59*B)*(a + I*a*Tan[c + d*x])^(3/2))/(3*d)))/(5*a))/7)/3
```

3.82.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

3.82. $\int \tan^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^(m/(f*m))), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4077 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

rule 4080 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`

3.82.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 206, normalized size of antiderivative = 0.84

method	result
derivativedivides	$2i \left(\frac{iB(a+ia \tan(dx+c))^{\frac{9}{2}}}{9} - \frac{iBa(a+ia \tan(dx+c))^{\frac{7}{2}}}{7} - \frac{Aa(a+ia \tan(dx+c))^{\frac{7}{2}}}{7} + \frac{ia^2B(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} + \frac{iBa^3(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} \right)$
default	$2i \left(\frac{iB(a+ia \tan(dx+c))^{\frac{9}{2}}}{9} - \frac{iBa(a+ia \tan(dx+c))^{\frac{7}{2}}}{7} - \frac{Aa(a+ia \tan(dx+c))^{\frac{7}{2}}}{7} + \frac{ia^2B(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} + \frac{iBa^3(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} \right)$
parts	$2iA \left(-\frac{(a+ia \tan(dx+c))^{\frac{7}{2}}}{7} - \frac{(a+ia \tan(dx+c))^{\frac{3}{2}}a^2}{3} - 2a^3\sqrt{a+ia \tan(dx+c)} + 2a^{\frac{7}{2}}\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right) \right) + da$

input `int(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output
$$2*I/d/a^2*(1/9*I*B*(a+I*a*tan(d*x+c))^(9/2)-1/7*I*B*a*(a+I*a*tan(d*x+c))^(7/2)-1/7*A*a*(a+I*a*tan(d*x+c))^(7/2)+1/5*I*a^2*B*(a+I*a*tan(d*x+c))^(5/2)+1/3*I*B*a^3*(a+I*a*tan(d*x+c))^(3/2)-1/3*A*a^3*(a+I*a*tan(d*x+c))^(3/2)+2*I*B*a^4*(a+I*a*tan(d*x+c))^(1/2)-2*A*a^4*(a+I*a*tan(d*x+c))^(1/2)+2*a^(9/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2)))$$

3.82.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 531 vs. 2(193) = 386.

Time = 0.28 (sec) , antiderivative size = 531, normalized size of antiderivative = 2.16

$$\int \tan^2(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx =$$

$$2 \left(315\sqrt{2}\sqrt{-\frac{(A^2-2iAB-B^2)a^5}{d^2}}(de^{(8i dx+8i c)} + 4de^{(6i dx+6i c)} + 6de^{(4i dx+4i c)} + 4de^{(2i dx+2i c)} + d) \log \left(\frac{4(-i \dots)}{\dots} \right) \right)$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output

```
-2/315*(315*sqrt(2)*sqrt(-(A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(8*I*d*x + 8
*I*c) + 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*x
+ 2*I*c) + d)*log(4*((-I*A - B)*a^3*e^(I*d*x + I*c) + sqrt(-(A^2 - 2*I*A*
B - B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(a/(e^(2*I*d*x + 2*I*c)
+ 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 315*sqrt(2)*sqrt(-(A^2 - 2*I*A
*B - B^2)*a^5/d^2)*(d*e^(8*I*d*x + 8*I*c) + 4*d*e^(6*I*d*x + 6*I*c) + 6*d*
e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((-I*A - B)*a^3*e
^(I*d*x + I*c) - sqrt(-(A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*
c) + d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a^
2)) + 2*sqrt(2)*(2*(300*I*A + 323*B)*a^2*e^(9*I*d*x + 9*I*c) + 27*(65*I*A
+ 61*B)*a^2*e^(7*I*d*x + 7*I*c) + 63*(35*I*A + 37*B)*a^2*e^(5*I*d*x + 5*I*
c) + 1365*(I*A + B)*a^2*e^(3*I*d*x + 3*I*c) + 315*(I*A + B)*a^2*e^(I*d*x +
I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(8*I*d*x + 8*I*c) + 4*d*e^(
6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*x + 2*I*c) + d)
```

3.82.6 Sympy [F]

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{5/2}(A + B \tan(c + dx)) \tan^2(c + dx) dx$$

input `integrate(tan(d*x+c)**2*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(5/2)*(A + B*tan(c + d*x))*tan(c + d*x)**2, x)`

3.82.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.72

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{2i \left(315 \sqrt{2}(A - iB)a^{\frac{11}{2}} \log \left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}} \right) - 35i (ia \tan(dx+c) + a)^{\frac{9}{2}}Ba + 45 (ia \tan(dx+c) + a)^{\frac{7}{2}}Ba \right)}{d}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

3.82. $\int \tan^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

output
$$\begin{aligned} & -2/315*I*(315*\sqrt{2}*(A - I*B)*a^{(11/2)}*\log(-(\sqrt{2}*\sqrt{a} - \sqrt{I*a*\tan(dx + c) + a})/(\sqrt{2}*\sqrt{a} + \sqrt{I*a*\tan(dx + c) + a})) - 35*I* \\ & (I*a*\tan(dx + c) + a)^{(9/2)}*B*a + 45*(I*a*\tan(dx + c) + a)^{(7/2)}*(A + I* \\ & B)*a^2 - 63*I*(I*a*\tan(dx + c) + a)^{(5/2)}*B*a^3 + 105*(I*a*\tan(dx + c) + \\ & a)^{(3/2)}*(A - I*B)*a^4 + 630*\sqrt{I*a*\tan(dx + c) + a}*(A - I*B)*a^5)/(a \\ & ^3*d) \end{aligned}$$

3.82.8 Giac [F]

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{5/2} \tan(dx + c)^2 dx$$

input `integrate(tan(dx+c)^2*(a+I*a*tan(dx+c))^(5/2)*(A+B*tan(dx+c)),x, algorithm="giac")`

output `integrate((B*tan(dx + c) + A)*(I*a*tan(dx + c) + a)^(5/2)*tan(dx + c)^2, x)`

3.82.9 Mupad [B] (verification not implemented)

Time = 8.71 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.05

$$\begin{aligned} & \int \tan^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \\ & -\frac{2B(a + a \tan(c + dx) \operatorname{li})^{5/2}}{5d} - \frac{Aa(a + a \tan(c + dx) \operatorname{li})^{3/2} 2i}{3d} \\ & -\frac{2Ba(a + a \tan(c + dx) \operatorname{li})^{3/2}}{3d} - \frac{Aa^2 \sqrt{a + a \tan(c + dx) \operatorname{li}} 4i}{d} \\ & -\frac{A(a + a \tan(c + dx) \operatorname{li})^{7/2} 2i}{7ad} - \frac{4Ba^2 \sqrt{a + a \tan(c + dx) \operatorname{li}}}{d} \\ & +\frac{2B(a + a \tan(c + dx) \operatorname{li})^{7/2}}{7ad} - \frac{2B(a + a \tan(c + dx) \operatorname{li})^{9/2}}{9a^2 d} \\ & +\frac{\sqrt{2}A(-a)^{5/2} \operatorname{atan}\left(\frac{\sqrt{2}\sqrt{a+a \tan(c+dx) \operatorname{li}}}{2\sqrt{-a}}\right) 4i}{d} \\ & -\frac{\sqrt{2}Ba^{5/2} \operatorname{atan}\left(\frac{\sqrt{2}\sqrt{a+a \tan(c+dx) \operatorname{li}} \operatorname{li}}{2\sqrt{a}}\right) 4i}{d} \end{aligned}$$

3.82. $\int \tan^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),x)`

output $(2*B*(a + a*\tan(c + d*x)*1i)^{(7/2)})/(7*a*d) - (A*a*(a + a*\tan(c + d*x)*1i)^{(3/2)*2i)/(3*d) - (2*B*a*(a + a*\tan(c + d*x)*1i)^{(3/2)})/(3*d) - (A*a^2*(a + a*\tan(c + d*x)*1i)^{(1/2)*4i)/d - (A*(a + a*\tan(c + d*x)*1i)^{(7/2)*2i)/(7*a*d) - (4*B*a^2*(a + a*\tan(c + d*x)*1i)^{(1/2)})/d - (2*B*(a + a*\tan(c + d*x)*1i)^{(5/2)})/(5*d) - (2*B*(a + a*\tan(c + d*x)*1i)^{(9/2)})/(9*a^2*d) + (2^{(1/2)*A*(-a)^{(5/2)*\operatorname{atan}((2^{(1/2)*(a + a*\tan(c + d*x)*1i)^{(1/2)})/(2*(-a)^{(1/2)})})}*4i)/d - (2^{(1/2)*B*a^{(5/2)*\operatorname{atan}((2^{(1/2)*(a + a*\tan(c + d*x)*1i)^{(1/2)})/(2*a^{(1/2)})})}*4i)/d$

3.83 $\int \tan(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.83.1	Optimal result	943
3.83.2	Mathematica [A] (verified)	944
3.83.3	Rubi [A] (verified)	944
3.83.4	Maple [A] (verified)	947
3.83.5	Fricas [B] (verification not implemented)	947
3.83.6	Sympy [F]	948
3.83.7	Maxima [A] (verification not implemented)	948
3.83.8	Giac [F(-1)]	949
3.83.9	Mupad [B] (verification not implemented)	949

3.83.1 Optimal result

Integrand size = 34, antiderivative size = 171

$$\int \tan(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx =$$

$$\frac{4\sqrt{2}a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}$$

$$+ \frac{4a^2(A-iB)\sqrt{a+ia \tan(c+dx)}}{d} + \frac{2a(A-iB)(a+ia \tan(c+dx))^{3/2}}{3d}$$

$$+ \frac{2A(a+ia \tan(c+dx))^{5/2}}{5d} - \frac{2iB(a+ia \tan(c+dx))^{7/2}}{7ad}$$

output

```
-4*a^(5/2)*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2
^(1/2)/d+4*a^2*(A-I*B)*(a+I*a*tan(d*x+c))^(1/2)/d+2/3*a*(A-I*B)*(a+I*a*tan
(d*x+c))^(3/2)/d+2/5*A*(a+I*a*tan(d*x+c))^(5/2)/d-2/7*I*B*(a+I*a*tan(d*x+c
))^(7/2)/a/d
```

3.83.2 Mathematica [A] (verified)

Time = 1.54 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.77

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{-420\sqrt{2}a^{5/2}(A - iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + 2a^2\sqrt{a + ia \tan(c + dx)}(266A - 260B + B \tan(c + dx))}{105d}$$

input `Integrate[Tan[c + d*x]*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(-420*Sqrt[2]*a^(5/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] + 2*a^2*Sqrt[a + I*a*Tan[c + d*x]]*(266*A - (260*I)*B + ((77*I)*A + 80*B)*Tan[c + d*x] + (-21*A + (45*I)*B)*Tan[c + d*x]^2 - 15*B*Tan[c + d*x]^3)/(105*d)`

3.83.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.98, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.324$, Rules used = {3042, 4075, 3042, 4010, 3042, 3959, 3042, 3959, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4075} \\ & \int (i \tan(c + dx)a + a)^{5/2}(A \tan(c + dx) - B) dx - \frac{2iB(a + ia \tan(c + dx))^{7/2}}{7ad} \\ & \quad \downarrow \text{3042} \\ & \int (i \tan(c + dx)a + a)^{5/2}(A \tan(c + dx) - B) dx - \frac{2iB(a + ia \tan(c + dx))^{7/2}}{7ad} \end{aligned}$$

$$\begin{aligned}
& \downarrow 4010 \\
& -(B + iA) \int (i \tan(c + dx)a + a)^{5/2} dx + \frac{2A(a + ia \tan(c + dx))^{5/2}}{5d} - \frac{2iB(a + ia \tan(c + dx))^{7/2}}{7ad} \\
& \downarrow 3042 \\
& -(B + iA) \int (i \tan(c + dx)a + a)^{5/2} dx + \frac{2A(a + ia \tan(c + dx))^{5/2}}{5d} - \frac{2iB(a + ia \tan(c + dx))^{7/2}}{7ad} \\
& \downarrow 3959 \\
& -(B + iA) \left(2a \int (i \tan(c + dx)a + a)^{3/2} dx + \frac{2ia(a + ia \tan(c + dx))^{3/2}}{3d} \right) + \\
& \quad \frac{2A(a + ia \tan(c + dx))^{5/2}}{5d} - \frac{2iB(a + ia \tan(c + dx))^{7/2}}{7ad} \\
& \downarrow 3042 \\
& -(B + iA) \left(2a \int (i \tan(c + dx)a + a)^{3/2} dx + \frac{2ia(a + ia \tan(c + dx))^{3/2}}{3d} \right) + \\
& \quad \frac{2A(a + ia \tan(c + dx))^{5/2}}{5d} - \frac{2iB(a + ia \tan(c + dx))^{7/2}}{7ad} \\
& \downarrow 3959 \\
& iA \left(2a \left(2a \int \sqrt{i \tan(c + dx)a + adx} + \frac{2ia\sqrt{a + ia \tan(c + dx)}}{d} \right) + \frac{2ia(a + ia \tan(c + dx))^{3/2}}{3d} \right) + \\
& \quad \frac{2A(a + ia \tan(c + dx))^{5/2}}{5d} - \frac{2iB(a + ia \tan(c + dx))^{7/2}}{7ad} \\
& \downarrow 3042 \\
& iA \left(2a \left(2a \int \sqrt{i \tan(c + dx)a + adx} + \frac{2ia\sqrt{a + ia \tan(c + dx)}}{d} \right) + \frac{2ia(a + ia \tan(c + dx))^{3/2}}{3d} \right) + \\
& \quad \frac{2A(a + ia \tan(c + dx))^{5/2}}{5d} - \frac{2iB(a + ia \tan(c + dx))^{7/2}}{7ad} \\
& \downarrow 3961 \\
& iA \left(2a \left(\frac{2ia\sqrt{a + ia \tan(c + dx)}}{d} - \frac{4ia^2 \int \frac{1}{a - ia \tan(c + dx)} d\sqrt{i \tan(c + dx)a + a}}{d} \right) + \frac{2ia(a + ia \tan(c + dx))^{3/2}}{3d} \right) + \\
& \quad \frac{2A(a + ia \tan(c + dx))^{5/2}}{5d} - \frac{2iB(a + ia \tan(c + dx))^{7/2}}{7ad} \\
& \downarrow 219
\end{aligned}$$

$$iA \left(2a \left(\frac{2ia\sqrt{a+ia\tan(c+dx)}}{d} - \frac{2i\sqrt{2}a^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} \right) + \frac{2ia(a+ia\tan(c+dx))^{3/2}}{3d} \right) + \frac{2A(a+ia\tan(c+dx))^{5/2}}{5d} - \frac{2iB(a+ia\tan(c+dx))^{7/2}}{7ad}$$

input `Int[Tan[c + d*x]*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(2*A*(a + I*a*Tan[c + d*x])^(5/2))/(5*d) - (((2*I)/7)*B*(a + I*a*Tan[c + d*x])^(7/2))/(a*d) - (I*A + B)*(((2*I)/3)*a*(a + I*a*Tan[c + d*x])^(3/2))/d + 2*a*(((2*I)*Sqrt[2]*a^(3/2)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d + ((2*I)*a*Sqrt[a + I*a*Tan[c + d*x]])/d)`

3.83.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3959 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

3.83.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.96

method	result
derivativedivides	$\frac{-\frac{2iB(a+ia \tan(dx+c))^{\frac{7}{2}}}{7} + \frac{2Aa(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} - \frac{2iB a^2(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + \frac{2A a^2(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - 4ia^3 B \sqrt{a+ia \tan(dx+c)}}{ad}$
default	$\frac{-\frac{2iB(a+ia \tan(dx+c))^{\frac{7}{2}}}{7} + \frac{2Aa(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} - \frac{2iB a^2(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + \frac{2A a^2(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - 4ia^3 B \sqrt{a+ia \tan(dx+c)}}{ad}$
parts	$\frac{A \left(\frac{2(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} + \frac{2a(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + 4a^2 \sqrt{a+ia \tan(dx+c)} - 4a^{\frac{5}{2}} \sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}} \right) \right)}{d} + \dots$

```
input int(tan(d*x+c)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 2/d/a*(-1/7*I*B*(a+I*a*tan(d*x+c))^(7/2)+1/5*A*a*(a+I*a*tan(d*x+c))^(5/2)-
1/3*I*B*a^2*(a+I*a*tan(d*x+c))^(3/2)+1/3*A*a^2*(a+I*a*tan(d*x+c))^(3/2)-2*
I*B*a^3*(a+I*a*tan(d*x+c))^(1/2)+2*A*a^3*(a+I*a*tan(d*x+c))^(1/2)-2*a^(7/2)
)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2)))
```

3.83.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 477 vs. 2(130) = 260.

Time = 0.26 (sec) , antiderivative size = 477, normalized size of antiderivative = 2.79

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx =$$

$$\frac{2 \left(105 \sqrt{2} \sqrt{\frac{(A^2 - 2iAB - B^2)a^5}{d^2}} (de^{(6i dx + 6i c)} + 3de^{(4i dx + 4i c)} + 3de^{(2i dx + 2i c)} + d) \log \left(\frac{4 \left((-iA - B)a^3 e^{(i dx + i c)} - \sqrt{\dots} \right)}{\dots} \right) \right)}{\dots}$$

3.83. $\int \tan(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-2/105*(105*sqrt(2)*sqrt((A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((-I*A - B)*a^3*e^(I*d*x + I*c) - sqrt((A^2 - 2*I*A*B - B^2)*a^5/d^2)*(I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/((-I*A - B)*a^2) - 105*sqrt(2)*sqrt((A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((-I*A - B)*a^3*e^(I*d*x + I*c) - sqrt((A^2 - 2*I*A*B - B^2)*a^5/d^2)*(I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/((-I*A - B)*a^2) - 2*sqrt(2)*(2*(91*A - 100*I*B)*a^2*e^(7*I*d*x + 7*I*c) + 7*(61*A - 55*I*B)*a^2*e^(5*I*d*x + 5*I*c) + 350*(A - I*B)*a^2*e^(3*I*d*x + 3*I*c) + 105*(A - I*B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)`

3.83.6 Sympy [F]

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{5/2} (A + B \tan(c + dx)) \tan(c + dx) dx$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(5/2)*(A + B*tan(c + d*x))*tan(c + d*x), x)`

3.83.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.89

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{2 \left(105 \sqrt{2} (A - i B) a^{\frac{9}{2}} \log \left(-\frac{\sqrt{2} \sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2} \sqrt{a} + \sqrt{ia \tan(dx+c)+a}} \right) - 15i (ia \tan(dx+c) + a)^{\frac{7}{2}} Ba + \right.}{\left. \right)}$$

3.83. $\int \tan(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output
$$\frac{2}{105} \cdot (105 \sqrt{2}) \cdot (A - I \cdot B) \cdot a^{9/2} \cdot \log\left(\frac{-\sqrt{2} \sqrt{a} - \sqrt{I \cdot a \cdot \tan(dx + c) + a}}{\sqrt{2} \sqrt{a} + \sqrt{I \cdot a \cdot \tan(dx + c) + a}}\right) - 15 \cdot I \cdot (I \cdot a \cdot \tan(dx + c) + a)^{7/2} \cdot B \cdot a + 21 \cdot (I \cdot a \cdot \tan(dx + c) + a)^{5/2} \cdot A \cdot a^2 + 35 \cdot (I \cdot a \cdot \tan(dx + c) + a)^{3/2} \cdot (A - I \cdot B) \cdot a^3 + 210 \cdot \sqrt{I \cdot a \cdot \tan(dx + c) + a} \cdot (A - I \cdot B) \cdot a^4 / (a^2 \cdot d)$$

3.83.8 Giac [F(-1)]

Timed out.

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output Timed out

3.83.9 Mupad [B] (verification not implemented)

Time = 1.67 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.24

$$\begin{aligned} & \int \tan(c + dx)(a \\ & + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{2 A (a + a \tan(c + dx) \operatorname{li})^{5/2}}{5 d} \\ & + \frac{2 A a (a + a \tan(c + dx) \operatorname{li})^{3/2}}{3 d} - \frac{B a (a + a \tan(c + dx) \operatorname{li})^{3/2} 2i}{3 d} \\ & + \frac{4 A a^2 \sqrt{a + a \tan(c + dx) \operatorname{li}}}{d} - \frac{B a^2 \sqrt{a + a \tan(c + dx) \operatorname{li}} 4i}{d} \\ & - \frac{B (a + a \tan(c + dx) \operatorname{li})^{7/2} 2i}{7 a d} + \frac{\sqrt{2} B (-a)^{5/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2 \sqrt{-a}}\right) 4i}{d} \\ & + \frac{\sqrt{2} A a^{5/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2 \sqrt{a}}\right) 4i}{d} \end{aligned}$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),x)`

output `(2*A*(a + a*tan(c + d*x)*1i)^(5/2))/(5*d) + (2*A*a*(a + a*tan(c + d*x)*1i)^(3/2))/(3*d) - (B*a*(a + a*tan(c + d*x)*1i)^(3/2)*2i)/(3*d) + (4*A*a^2*(a + a*tan(c + d*x)*1i)^(1/2))/d - (B*a^2*(a + a*tan(c + d*x)*1i)^(1/2)*4i)/d - (B*(a + a*tan(c + d*x)*1i)^(7/2)*2i)/(7*a*d) + (2^(1/2)*B*(-a)^(5/2)*a*tan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2)))*4i)/d + (2^(1/2)*A*a^(5/2)*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)*1i)/(2*a^(1/2)))*4i)/d`

3.84 $\int (a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx$

3.84.1	Optimal result	951
3.84.2	Mathematica [A] (verified)	951
3.84.3	Rubi [A] (verified)	952
3.84.4	Maple [A] (verified)	954
3.84.5	Fricas [B] (verification not implemented)	955
3.84.6	Sympy [F]	956
3.84.7	Maxima [A] (verification not implemented)	956
3.84.8	Giac [F(-1)]	956
3.84.9	Mupad [B] (verification not implemented)	957

3.84.1 Optimal result

Integrand size = 28, antiderivative size = 141

$$\int (a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx =$$

$$-\frac{4\sqrt{2}a^{5/2}(iA + B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{4a^2(iA + B)\sqrt{a + ia \tan(c + dx)}}{d}$$

$$+ \frac{2a(iA + B)(a + ia \tan(c + dx))^{3/2}}{3d} + \frac{2B(a + ia \tan(c + dx))^{5/2}}{5d}$$

```
output -4*a^(5/2)*(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2
^(1/2)/d+4*a^2*(I*A+B)*(a+I*a*tan(d*x+c))^(1/2)/d+2/3*a*(I*A+B)*(a+I*a*tan
(d*x+c))^(3/2)/d+2/5*B*(a+I*a*tan(d*x+c))^(5/2)/d
```

3.84.2 Mathematica [A] (verified)

Time = 0.87 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.82

$$\int (a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx =$$

$$\frac{-60i\sqrt{2}a^{5/2}(A - iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + 2a^2\sqrt{a + ia \tan(c + dx)}(35iA + 38B)}{15d}$$

```
input Integrate[(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]
```

output $((-60*I)*\text{Sqrt}[2]*a^{(5/2)}*(A - I*B)*\text{ArcTanh}[\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]]/(\text{Sqrt}[2]*\text{Sqrt}[a])) + 2*a^2*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]*((35*I)*A + 38*B + (-5*A + (11*I)*B)*\text{Tan}[c + d*x] - 3*B*\text{Tan}[c + d*x]^2)/(15*d)$

3.84.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.97, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.321$, Rules used = {3042, 4010, 3042, 3959, 3042, 3959, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4010} \\ & (A - iB) \int (i \tan(c + dx)a + a)^{5/2} dx + \frac{2B(a + ia \tan(c + dx))^{5/2}}{5d} \\ & \quad \downarrow \text{3042} \\ & (A - iB) \int (i \tan(c + dx)a + a)^{5/2} dx + \frac{2B(a + ia \tan(c + dx))^{5/2}}{5d} \\ & \quad \downarrow \text{3959} \\ & (A - iB) \left(2a \int (i \tan(c + dx)a + a)^{3/2} dx + \frac{2ia(a + ia \tan(c + dx))^{3/2}}{3d} \right) + \\ & \quad \frac{2B(a + ia \tan(c + dx))^{5/2}}{5d} \\ & \quad \downarrow \text{3042} \\ & (A - iB) \left(2a \int (i \tan(c + dx)a + a)^{3/2} dx + \frac{2ia(a + ia \tan(c + dx))^{3/2}}{3d} \right) + \\ & \quad \frac{2B(a + ia \tan(c + dx))^{5/2}}{5d} \\ & \quad \downarrow \text{3959} \end{aligned}$$

$$iB) \left(2a \left(2a \int \sqrt{i \tan(c+dx)a+adx} + \frac{2ia\sqrt{a+ia \tan(c+dx)}}{d} \right) + \frac{2ia(a+ia \tan(c+dx))^{3/2}}{3d} \right) + \frac{2B(a+ia \tan(c+dx))^{5/2}}{5d}$$

↓ 3042

$$iB) \left(2a \left(2a \int \sqrt{i \tan(c+dx)a+adx} + \frac{2ia\sqrt{a+ia \tan(c+dx)}}{d} \right) + \frac{2ia(a+ia \tan(c+dx))^{3/2}}{3d} \right) + \frac{2B(a+ia \tan(c+dx))^{5/2}}{5d}$$

↓ 3961

$$iB) \left(2a \left(\frac{2ia\sqrt{a+ia \tan(c+dx)}}{d} - \frac{4ia^2 \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a}}{d} \right) + \frac{2ia(a+ia \tan(c+dx))^{3/2}}{3d} \right) + \frac{2B(a+ia \tan(c+dx))^{5/2}}{5d}$$

↓ 219

$$iB) \left(2a \left(\frac{2ia\sqrt{a+ia \tan(c+dx)}}{d} - \frac{2i\sqrt{2}a^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} \right) + \frac{2ia(a+ia \tan(c+dx))^{3/2}}{3d} \right) + \frac{2B(a+ia \tan(c+dx))^{5/2}}{5d}$$

input `Int[(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(2*B*(a + I*a*Tan[c + d*x])^(5/2))/(5*d) + (A - I*B)*(((2*I)/3)*a*(a + I*a*Tan[c + d*x])^(3/2))/d + 2*a*(((-2*I)*Sqrt[2]*a^(3/2)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d + ((2*I)*a*Sqrt[a + I*a*Tan[c + d*x]])/d)`

3.84.3.1 Defintions of rubi rules used

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3959 Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a +
b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*
x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n
, 1]
```

```
rule 3961 Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d)
Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a
, b, c, d}, x] && EqQ[a^2 + b^2, 0]
```

```
rule 4010 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp
[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e
, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]
```

3.84.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{2i \left(-\frac{iB(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} - \frac{iBa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + \frac{Aa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - 2ia^2B\sqrt{a+ia \tan(dx+c)} + 2Aa^2\sqrt{a+ia \tan(dx+c)} \right)}{d}$
default	$\frac{2i \left(-\frac{iB(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} - \frac{iBa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + \frac{Aa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - 2ia^2B\sqrt{a+ia \tan(dx+c)} + 2Aa^2\sqrt{a+ia \tan(dx+c)} \right)}{d}$
parts	$\frac{2iAa \left(\frac{(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + 2a\sqrt{a+ia \tan(dx+c)} - 2a^{\frac{3}{2}}\sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}} \right) \right)}{d} + \frac{B \left(\frac{2(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} \right)}{d}$

3.84. $\int (a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx$

```
input int((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 2*I/d*(-1/5*I*B*(a+I*a*tan(d*x+c))^(5/2)-1/3*I*B*a*(a+I*a*tan(d*x+c))^(3/2)
)+1/3*A*a*(a+I*a*tan(d*x+c))^(3/2)-2*I*B*a^2*(a+I*a*tan(d*x+c))^(1/2)+2*A*
a^2*(a+I*a*tan(d*x+c))^(1/2)-2*a^(5/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*
tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))
```

3.84.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 421 vs. $2(108) = 216$.

Time = 0.25 (sec) , antiderivative size = 421, normalized size of antiderivative = 2.99

$$\int (a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx = \frac{2 \left(15 \sqrt{2} \sqrt{-\frac{(A^2 - 2iAB - B^2)a^5}{d^2}} (de^{(4i dx + 4i c)} + 2 de^{(2i dx + 2i c)} + d) \log \left(\frac{4 \left((-iA - B)a^3 e^{(i dx + i c)} + \dots \right)}{\dots} \right) \right)}{\dots}$$

```
input integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output 2/15*(15*sqrt(2)*sqrt(-(A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(4*I*d*x + 4*I*
c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((-I*A - B)*a^3*e^(I*d*x + I*c) +
sqrt(-(A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(a/(e
^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 15*sqrt(2)*
sqrt(-(A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d
*x + 2*I*c) + d)*log(4*((-I*A - B)*a^3*e^(I*d*x + I*c) - sqrt(-(A^2 - 2*I*
A*B - B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(a/(e^(2*I*d*x + 2*I*c
) + 1))))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 2*sqrt(2)*(2*(-10*I*A - 13*B
)*a^2*e^(5*I*d*x + 5*I*c) + 35*(-I*A - B)*a^2*e^(3*I*d*x + 3*I*c) + 15*(-I
*A - B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(4*I*
d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)
```

3.84.6 Sympy [F]

$$\int (a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{5/2} (A + B \tan(c + dx)) dx$$

input `integrate((a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(5/2)*(A + B*tan(c + d*x)), x)`

3.84.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.95

$$\int (a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx = \frac{2i \left(15 \sqrt{2} (A - iB) a^{7/2} \log \left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}} \right) - 3i (ia \tan(dx+c) + a)^{5/2} Ba + 5 \right)}{15 ad}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `2/15*I*(15*sqrt(2)*(A - I*B)*a^(7/2)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - 3*I*(I*a*tan(d*x + c) + a)^(5/2)*B*a + 5*(I*a*tan(d*x + c) + a)^(3/2)*(A - I*B)*a^2 + 30*sqrt(I*a*tan(d*x + c) + a)*(A - I*B)*a^3)/(a*d)`

3.84.8 Giac [F(-1)]

Timed out.

$$\int (a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.84.9 Mupad [B] (verification not implemented)

Time = 1.16 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.33

$$\begin{aligned}
\int (a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx &= \frac{2 B (a + a \tan(c + dx) \operatorname{li})^{5/2}}{5 d} \\
&+ \frac{A a (a + a \tan(c + dx) \operatorname{li})^{3/2} 2i}{3 d} + \frac{2 B a (a + a \tan(c + dx) \operatorname{li})^{3/2}}{3 d} \\
&+ \frac{A a^2 \sqrt{a + a \tan(c + dx) \operatorname{li}} 4i}{d} + \frac{4 B a^2 \sqrt{a + a \tan(c + dx) \operatorname{li}}}{d} \\
&- \frac{\sqrt{2} A (-a)^{5/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2 \sqrt{-a}}\right) 4i}{d} \\
&+ \frac{\sqrt{2} B a^{5/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2 \sqrt{a}}\right) 4i}{d}
\end{aligned}$$

input `int((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),x)`output `(2*B*(a + a*tan(c + d*x)*1i)^(5/2))/(5*d) + (A*a*(a + a*tan(c + d*x)*1i)^(3/2)*2i)/(3*d) + (2*B*a*(a + a*tan(c + d*x)*1i)^(3/2))/(3*d) + (A*a^2*(a + a*tan(c + d*x)*1i)^(1/2)*4i)/d + (4*B*a^2*(a + a*tan(c + d*x)*1i)^(1/2))/d - (2^(1/2)*A*(-a)^(5/2)*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2)))*4i)/d + (2^(1/2)*B*a^(5/2)*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)*1i)/(2*a^(1/2)))*4i)/d`

3.85 $\int \cot(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.85.1	Optimal result	958
3.85.2	Mathematica [A] (verified)	958
3.85.3	Rubi [A] (verified)	959
3.85.4	Maple [A] (verified)	963
3.85.5	Fricas [B] (verification not implemented)	963
3.85.6	Sympy [F]	964
3.85.7	Maxima [A] (verification not implemented)	964
3.85.8	Giac [F]	965
3.85.9	Mupad [B] (verification not implemented)	965

3.85.1 Optimal result

Integrand size = 34, antiderivative size = 147

$$\int \cot(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx =$$

$$-\frac{2a^{5/2} A \operatorname{Arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{d} + \frac{4\sqrt{2}a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}$$

$$-\frac{2a^2(A-2iB)\sqrt{a+ia \tan(c+dx)}}{d} + \frac{2iaB(a+ia \tan(c+dx))^{3/2}}{3d}$$

output

```
-2*a^(5/2)*A*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/d+4*a^(5/2)*(A-I*B)
*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)/d-2*a^2*(A-
2*I*B)*(a+I*a*tan(d*x+c))^(1/2)/d+2/3*I*a*B*(a+I*a*tan(d*x+c))^(3/2)/d
```

3.85.2 Mathematica [A] (verified)

Time = 1.17 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.86

$$\int \cot(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx =$$

$$-\frac{2a^2\left(3\sqrt{a}A \operatorname{Arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) - 6\sqrt{2}\sqrt{a}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + \sqrt{a+ia \tan(c+dx)}\right)}{3d}$$

input `Integrate[Cot[c + d*x]*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(-2*a^2*(3*Sqrt[a]*A*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] - 6*Sqrt[2]*Sqrt[a]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] + Sqrt[a + I*a*Tan[c + d*x]]*(3*A - (7*I)*B + B*Tan[c + d*x]))/(3*d)`

3.85.3 Rubi [A] (verified)

Time = 1.01 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {3042, 4077, 27, 3042, 4077, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan(c + dx)} dx \\
 & \quad \downarrow \text{4077} \\
 & \frac{2}{3} \int \frac{3}{2} \cot(c + dx)(i \tan(c + dx)a + a)^{3/2}(aA + a(iA + 2B) \tan(c + dx)) dx + \frac{2iaB(a + ia \tan(c + dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{27} \\
 & \int \cot(c + dx)(i \tan(c + dx)a + a)^{3/2}(aA + a(iA + 2B) \tan(c + dx)) dx + \frac{2iaB(a + ia \tan(c + dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(i \tan(c + dx)a + a)^{3/2}(aA + a(iA + 2B) \tan(c + dx))}{\tan(c + dx)} dx + \frac{2iaB(a + ia \tan(c + dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{4077} \\
 & 2 \int \frac{1}{2} \cot(c + dx) \sqrt{i \tan(c + dx)a + a} (Aa^2 + (3iA + 4B) \tan(c + dx)a^2) dx - \\
 & \quad \frac{2a^2(A - 2iB) \sqrt{a + ia \tan(c + dx)}}{d} + \frac{2iaB(a + ia \tan(c + dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.85. $\int \cot(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \int \cot(c+dx) \sqrt{i \tan(c+dx)a+a} (Aa^2 + (3iA+4B) \tan(c+dx)a^2) dx - \\
& \quad \frac{2a^2(A-2iB)\sqrt{a+ia \tan(c+dx)}}{d} + \frac{2iaB(a+ia \tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{3042} \\
& \int \frac{\sqrt{i \tan(c+dx)a+a} (Aa^2 + (3iA+4B) \tan(c+dx)a^2)}{\tan(c+dx)} dx - \\
& \quad \frac{2a^2(A-2iB)\sqrt{a+ia \tan(c+dx)}}{d} + \frac{2iaB(a+ia \tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{4083} \\
& 4a^2(B+iA) \int \sqrt{i \tan(c+dx)a+adx} + aA \int \cot(c+dx)(a-ia \tan(c+ \\
& dx)) \sqrt{i \tan(c+dx)a+adx} - \frac{2a^2(A-2iB)\sqrt{a+ia \tan(c+dx)}}{d} + \frac{2iaB(a+ia \tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{3042} \\
& 4a^2(B+iA) \int \sqrt{i \tan(c+dx)a+adx} + aA \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \\
& \quad \frac{2a^2(A-2iB)\sqrt{a+ia \tan(c+dx)}}{d} + \frac{2iaB(a+ia \tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{3961} \\
& \quad \frac{8ia^3(B+iA) \int \frac{1}{a-ia \tan(c+dx)} d \sqrt{i \tan(c+dx)a+a}}{d} + \\
& aA \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{2a^2(A-2iB)\sqrt{a+ia \tan(c+dx)}}{d} + \\
& \quad \frac{2iaB(a+ia \tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{219} \\
& aA \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{4i\sqrt{2}a^{5/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \\
& \quad \frac{2a^2(A-2iB)\sqrt{a+ia \tan(c+dx)}}{d} + \frac{2iaB(a+ia \tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{4082} \\
& a^3A \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx) - \frac{4i\sqrt{2}a^{5/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \\
& \quad \frac{2a^2(A-2iB)\sqrt{a+ia \tan(c+dx)}}{d} + \frac{2iaB(a+ia \tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{73}
\end{aligned}$$

3.85. $\int \cot(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \frac{2ia^2 A \int \frac{1}{i - \frac{i(\tan(c+dx)a+a)}{a}} d\sqrt{i \tan(c+dx)a+a}}{\frac{d}{2a^2(A-2iB)\sqrt{a+ia \tan(c+dx)}} + \frac{d}{2iaB(a+ia \tan(c+dx))^{3/2}}} - \frac{4i\sqrt{2}a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{3d} \\
& \quad \downarrow \text{221} \\
& \frac{4i\sqrt{2}a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\frac{d}{2a^2(A-2iB)\sqrt{a+ia \tan(c+dx)}} + \frac{d}{2iaB(a+ia \tan(c+dx))^{3/2}}} - \frac{2a^{5/2}A\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{3d}
\end{aligned}$$

input `Int[Cot[c + d*x]*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(-2*a^(5/2)*A*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d - ((4*I)*Sqrt[2]*a^(5/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d - (2*a^2*(A - (2*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/d + (((2*I)/3)*a*B*(a + I*a*Tan[c + d*x])^(3/2))/d`

3.85.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m+1)-1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4077 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int[((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x]))], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.85.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.85

method	result
derivativedivides	$\frac{2a \left(\frac{iB(a+ia \tan(dx+c))^{3/2}}{3} + 2iaB\sqrt{a+ia \tan(dx+c)} - aA\sqrt{a+ia \tan(dx+c)} + 2a^{3/2}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{2\sqrt{a}}\right) \right)}{d}$
default	$\frac{2a \left(\frac{iB(a+ia \tan(dx+c))^{3/2}}{3} + 2iaB\sqrt{a+ia \tan(dx+c)} - aA\sqrt{a+ia \tan(dx+c)} + 2a^{3/2}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{2\sqrt{a}}\right) \right)}{d}$

input `int(cot(d*x+c)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)`

output `2/d*a*(1/3*I*B*(a+I*a*tan(d*x+c))^(3/2)+2*I*B*a*(a+I*a*tan(d*x+c))^(1/2)-a
A(a+I*a*tan(d*x+c))^(1/2)+2*a^(3/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*t
an(d*x+c))^(1/2)*2^(1/2)/a^(1/2))-A*a^(3/2)*arctanh((a+I*a*tan(d*x+c))^(1/
2)/a^(1/2)))`

3.85.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 610 vs. 2(112) = 224.

Time = 0.27 (sec) , antiderivative size = 610, normalized size of antiderivative = 4.15

$$\int \cot(c+dx)(a+ia \tan(c+dx))^{5/2}(A + B \tan(c+dx)) dx = \frac{12\sqrt{2}\sqrt{\frac{(A^2-2iAB-B^2)a^5}{d^2}}(de^{(2i dx+2i c)} + d) \log \left(\frac{4 \left((-iA-B)a^3 e^{(i dx+i c)} - \sqrt{\frac{(A^2-2iAB-B^2)a^5}{d^2}}(i de^{(2i dx+2i c)} - (-iA-B)) \right)}{(-iA-B)} \right)}{(-iA-B)}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm
m="fracas")`

```
output 1/6*(12*sqrt(2)*sqrt((A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c)
+ d)*log(4*((-I*A - B)*a^3*e^(I*d*x + I*c) - sqrt((A^2 - 2*I*A*B - B^2)*a
^5/d^2)*(I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))
*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 12*sqrt(2)*sqrt((A^2 - 2*I*A*B - B^2
)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log(4*((-I*A - B)*a^3*e^(I*d*x + I*
c) - sqrt((A^2 - 2*I*A*B - B^2)*a^5/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) - I*d)*
sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 3*
sqrt(A^2*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log(16*(3*A*a^3*e^(2*I*d*x +
2*I*c) + A*a^3 + 2*sqrt(2)*sqrt(A^2*a^5/d^2)*(d*e^(3*I*d*x + 3*I*c) + d*e
^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x - 2*I*c)/(A
*a)) + 3*sqrt(A^2*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log(16*(3*A*a^3*e^(
2*I*d*x + 2*I*c) + A*a^3 - 2*sqrt(2)*sqrt(A^2*a^5/d^2)*(d*e^(3*I*d*x + 3*I
*c) + d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x -
2*I*c)/(A*a)) - 4*sqrt(2)*((3*A - 8*I*B)*a^2*e^(3*I*d*x + 3*I*c) + 3*(A -
2*I*B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(2*I*d
*x + 2*I*c) + d)
```

3.85.6 Sympy [F]

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{5/2}(A + B \tan(c + dx)) \cot(c + dx) dx$$

```
input integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)
```

```
output Integral((I*a*(tan(c + d*x) - I))**(5/2)*(A + B*tan(c + d*x))*cot(c + d*x)
, x)
```

3.85.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.05

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{6\sqrt{2}(A - iB)a^{5/2} \log\left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}}\right) - 3Aa^{5/2} \log\left(\frac{\sqrt{ia \tan(dx+c)+a} - \sqrt{a}}{\sqrt{ia \tan(dx+c)+a} + \sqrt{a}}\right) - 2i(ia \tan(dx+c) + a)}{3d}$$

3.85. $\int \cot(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/3*(6*sqrt(2)*(A - I*B)*a^(5/2)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - 3*A*a^(5/2)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a))) - 2*I*(I*a*tan(d*x + c) + a)^(3/2)*B*a + 6*sqrt(I*a*tan(d*x + c) + a)*(A - 2*I*B)*a^2)/d`

3.85.8 Giac [F]

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{5/2} \cot(dx + c) dx$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)*cot(d*x + c), x)`

3.85.9 Mupad [B] (verification not implemented)

Time = 8.14 (sec) , antiderivative size = 597, normalized size of antiderivative = 4.06

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx =$$

$$-\left(\frac{2a^2(A + B \operatorname{li})}{d} - \frac{Ba^2 6i}{d}\right) \sqrt{a + a \tan(c + dx)} \operatorname{li} + \frac{Ba(a + a \tan(c + dx) \operatorname{li})^{3/2} 2i}{3d}$$

$$- \frac{A \operatorname{atan}\left(\frac{A^3 a^8 d \sqrt{a^5} \sqrt{a + a \tan(c + dx)} \operatorname{li} 224i}{-224 d A^3 a^{11} + 512i d A^2 B a^{11} + 256 d A B^2 a^{11}} - \frac{A B^2 a^8 d \sqrt{a^5} \sqrt{a + a \tan(c + dx)} \operatorname{li} 256i}{-224 d A^3 a^{11} + 512i d A^2 B a^{11} + 256 d A B^2 a^{11}} + \frac{512 A^2 B a^8 d \sqrt{a^5} \sqrt{a + a \tan(c + dx)} \operatorname{li} 256i}{-224 d A^3 a^{11} + 512i d A^2 B a^{11} + 256 d A B^2 a^{11}}\right)}{d}$$

$$+ \frac{\sqrt{2} \operatorname{atan}\left(\frac{224 \sqrt{2} A^3 a^8 d \sqrt{-a^5} \sqrt{a + a \tan(c + dx)} \operatorname{li}}{448 d A^3 a^{11} - 1472i d A^2 B a^{11} - 1536 d A B^2 a^{11} + 512i d B^3 a^{11}} + \frac{\sqrt{2} B^3 a^8 d \sqrt{-a^5} \sqrt{a + a \tan(c + dx)} \operatorname{li} 256i}{448 d A^3 a^{11} - 1472i d A^2 B a^{11} - 1536 d A B^2 a^{11} + 512i d B^3 a^{11}}\right)}{d}$$

3.85. $\int \cot(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),x)`

output
$$\begin{aligned} & (B*a*(a + a*\tan(c + d*x)*1i)^{(3/2)*2i}/(3*d) - ((2*a^2*(A + B*1i))/d - (B* \\ & a^2*6i)/d)*(a + a*\tan(c + d*x)*1i)^{(1/2)} - (A*\operatorname{atan}((A^3*a^8*d*(a^5)^{(1/2)}* \\ & (a + a*\tan(c + d*x)*1i)^{(1/2)*224i}/(256*A*B^2*a^{11*d} - 224*A^3*a^{11*d} + A \\ & ^2*B*a^{11*d*512i}) - (A*B^2*a^8*d*(a^5)^{(1/2)}*(a + a*\tan(c + d*x)*1i)^{(1/2)} \\ & *256i)/(256*A*B^2*a^{11*d} - 224*A^3*a^{11*d} + A^2*B*a^{11*d*512i}) + (512*A^2* \\ & B*a^8*d*(a^5)^{(1/2)}*(a + a*\tan(c + d*x)*1i)^{(1/2)})/(256*A*B^2*a^{11*d} - 224 \\ & *A^3*a^{11*d} + A^2*B*a^{11*d*512i}))* (a^5)^{(1/2)*2i}/d + (2^{(1/2)}*\operatorname{atan}((224*2 \\ & ^{(1/2)}*A^3*a^8*d*(-a^5)^{(1/2)}*(a + a*\tan(c + d*x)*1i)^{(1/2)})/(448*A^3*a^{11} \\ & *d + B^3*a^{11*d*512i} - 1536*A*B^2*a^{11*d} - A^2*B*a^{11*d*1472i}) + (2^{(1/2)}* \\ & B^3*a^8*d*(-a^5)^{(1/2)}*(a + a*\tan(c + d*x)*1i)^{(1/2)*256i}/(448*A^3*a^{11*d} \\ & + B^3*a^{11*d*512i} - 1536*A*B^2*a^{11*d} - A^2*B*a^{11*d*1472i}) - (768*2^{(1/2)} \\ &)*A*B^2*a^8*d*(-a^5)^{(1/2)}*(a + a*\tan(c + d*x)*1i)^{(1/2)})/(448*A^3*a^{11*d} \\ & + B^3*a^{11*d*512i} - 1536*A*B^2*a^{11*d} - A^2*B*a^{11*d*1472i}) - (2^{(1/2)}*A^2 \\ & *B*a^8*d*(-a^5)^{(1/2)}*(a + a*\tan(c + d*x)*1i)^{(1/2)*736i}/(448*A^3*a^{11*d} \\ & + B^3*a^{11*d*512i} - 1536*A*B^2*a^{11*d} - A^2*B*a^{11*d*1472i}))* (A*1i + B)*(- \\ & a^5)^{(1/2)*4i}/d \end{aligned}$$

3.86 $\int \cot^2(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.86.1 Optimal result	967
3.86.2 Mathematica [A] (verified)	968
3.86.3 Rubi [A] (verified)	968
3.86.4 Maple [A] (verified)	973
3.86.5 Fricas [B] (verification not implemented)	973
3.86.6 Sympy [F(-1)]	974
3.86.7 Maxima [A] (verification not implemented)	974
3.86.8 Giac [F]	975
3.86.9 Mupad [B] (verification not implemented)	975

3.86.1 Optimal result

Integrand size = 36, antiderivative size = 158

$$\int \cot^2(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx =$$

$$\frac{a^{5/2}(5iA+2B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{d}$$

$$+ \frac{4\sqrt{2}a^{5/2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}$$

$$+ \frac{a^2(iA-2B)\sqrt{a+ia \tan(c+dx)}}{d} - \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^{3/2}}{d}$$

output

```
-a^(5/2)*(5*I*A+2*B)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/d+4*a^(5/2)
*(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)/d+a
^2*(I*A-2*B)*(a+I*a*tan(d*x+c))^(1/2)/d-a*A*cot(d*x+c)*(a+I*a*tan(d*x+c))^(
(3/2)/d
```

3.86.2 Mathematica [A] (verified)

Time = 1.44 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.82

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{-ia^{5/2}(5A - 2iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) + 4\sqrt{2}a^{5/2}(iA + B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}$$

input `Integrate[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `((-I)*a^(5/2)*(5*A - (2*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] + 4*Sqrt[2]*a^(5/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] + a^2*(-2*B - A*Cot[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]])/d`

3.86.3 Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.05, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3042, 4076, 27, 3042, 4077, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan(c + dx)^2} dx \\ & \quad \downarrow \text{4076} \\ & \int \frac{1}{2} \cot(c + dx)(i \tan(c + dx)a + a)^{3/2}(a(5iA + 2B) + a(A + 2iB) \tan(c + dx)) dx - \\ & \quad \frac{aA \cot(c + dx)(a + ia \tan(c + dx))^{3/2}}{d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\frac{1}{2} \int \cot(c+dx)(i \tan(c+dx)a+a)^{3/2}(a(5iA+2B)+a(A+2iB)\tan(c+dx))dx - \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^{3/2}}{d}$$

↓ 3042

$$\frac{1}{2} \int \frac{(i \tan(c+dx)a+a)^{3/2}(a(5iA+2B)+a(A+2iB)\tan(c+dx))}{\tan(c+dx)} dx - \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^{3/2}}{d}$$

↓ 4077

$$\frac{1}{2} \left(2 \int \frac{1}{2} \cot(c+dx) \sqrt{i \tan(c+dx)a+a} (a^2(5iA+2B) - 3a^2(A-2iB)\tan(c+dx)) dx + \frac{2a^2(-2B+iA)\sqrt{a+ia \tan(c+dx)}}{d} \right) - \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^{3/2}}{d}$$

↓ 27

$$\frac{1}{2} \left(\int \cot(c+dx) \sqrt{i \tan(c+dx)a+a} (a^2(5iA+2B) - 3a^2(A-2iB)\tan(c+dx)) dx + \frac{2a^2(-2B+iA)\sqrt{a+ia \tan(c+dx)}}{d} \right) - \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^{3/2}}{d}$$

↓ 3042

$$\frac{1}{2} \left(\int \frac{\sqrt{i \tan(c+dx)a+a} (a^2(5iA+2B) - 3a^2(A-2iB)\tan(c+dx))}{\tan(c+dx)} dx + \frac{2a^2(-2B+iA)\sqrt{a+ia \tan(c+dx)}}{d} \right) - \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^{3/2}}{d}$$

↓ 4083

$$\frac{1}{2} \left(-8a^2(A-iB) \int \sqrt{i \tan(c+dx)a+adx} + a(2B+5iA) \int \cot(c+dx)(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+} \right) - \frac{aA \cot(c+dx)(a+ia \tan(c+dx))^{3/2}}{d}$$

↓ 3042

$$\frac{1}{2} \left(-8a^2(A - iB) \int \frac{\sqrt{i \tan(c + dx)a + a} dx + a(2B + 5iA) \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} dx + \frac{2a^2}{d} \frac{aA \cot(c + dx)(a + ia \tan(c + dx))^{3/2}}{d} \right)$$

↓ 3961

$$\frac{1}{2} \left(\frac{16ia^3(A - iB) \int \frac{1}{a - ia \tan(c + dx)} d\sqrt{i \tan(c + dx)a + a}}{d} + a(2B + 5iA) \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} dx + \frac{aA \cot(c + dx)(a + ia \tan(c + dx))^{3/2}}{d} \right)$$

↓ 219

$$\frac{1}{2} \left(a(2B + 5iA) \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} dx + \frac{8i\sqrt{2}a^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{aA \cot(c + dx)(a + ia \tan(c + dx))^{3/2}}{d} \right)$$

↓ 4082

$$\frac{1}{2} \left(\frac{a^3(2B + 5iA) \int \frac{\cot(c + dx)}{\sqrt{i \tan(c + dx)a + a}} d \tan(c + dx)}{d} + \frac{8i\sqrt{2}a^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{2a^2(-2B + iA)}{d} + \frac{aA \cot(c + dx)(a + ia \tan(c + dx))^{3/2}}{d} \right)$$

↓ 73

$$\frac{1}{2} \left(-\frac{2ia^2(2B + 5iA) \int \frac{1}{i - \frac{i \tan(c + dx)a + a}{a}} d\sqrt{i \tan(c + dx)a + a}}{d} + \frac{8i\sqrt{2}a^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{aA \cot(c + dx)(a + ia \tan(c + dx))^{3/2}}{d} \right)$$

↓ 221

$$\frac{1}{2} \left(-\frac{2a^{5/2}(2B + 5iA) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{a}}\right)}{d} + \frac{8i\sqrt{2}a^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{2a^2(-2B + iA)}{d} + \frac{aA \cot(c + dx)(a + ia \tan(c + dx))^{3/2}}{d} \right)$$

input `Int[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `-((a*A*Cot[c + d*x]*(a + I*a*Tan[c + d*x])^(3/2))/d) + ((-2*a^(5/2)*((5*I)*A + 2*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d + ((8*I)*Sqrt[2]*a^(5/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d + (2*a^2*(I*A - 2*B)*Sqrt[a + I*a*Tan[c + d*x]])/d)/2`

3.86.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 73 `Int[((a_) + (b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 4077 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int[(((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.86.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.83

method	result
derivativedivides	$\frac{2ia^2 \left(iB\sqrt{a+ia \tan(dx+c)}+2\sqrt{a}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right) - a \left(-\frac{iA\sqrt{a+ia \tan(dx+c)}}{2a \tan(dx+c)} + \frac{(-2iB+5A)a}{2a \tan(dx+c)} \right) \right)}{d}$
default	$\frac{2ia^2 \left(iB\sqrt{a+ia \tan(dx+c)}+2\sqrt{a}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right) - a \left(-\frac{iA\sqrt{a+ia \tan(dx+c)}}{2a \tan(dx+c)} + \frac{(-2iB+5A)a}{2a \tan(dx+c)} \right) \right)}{d}$

input `int(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `2*I/d*a^2*(I*B*(a+I*a*tan(d*x+c))^(1/2)+2*a^(1/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))-a*(-1/2*I*A*(a+I*a*tan(d*x+c))^(1/2)/a/tan(d*x+c)+1/2*(5*A-2*I*B)/a^(1/2)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2)))`

3.86.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 705 vs. 2(125) = 250.

Time = 0.26 (sec) , antiderivative size = 705, normalized size of antiderivative = 4.46

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx =$$

$$8\sqrt{2}\sqrt{-\frac{(A^2-2iAB-B^2)a^5}{d^2}}(de^{(2i dx+2i c)} - d) \log \left(\frac{4 \left((-iA-B)a^3e^{(i dx+i c)} + \sqrt{-\frac{(A^2-2iAB-B^2)a^5}{d^2}}(de^{(2i dx+2i c)}+d)\sqrt{\frac{a}{e^{(2i dx+2i c)}}}} \right)}{(-iA-B)a^2} \right)$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output

```

-1/4*(8*sqrt(2)*sqrt(-(A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c)
) - d)*log(4*((-I*A - B)*a^3*e^(I*d*x + I*c) + sqrt(-(A^2 - 2*I*A*B - B^2)
*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e
^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 8*sqrt(2)*sqrt(-(A^2 - 2*I*A*B - B^2)*
a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log(4*((-I*A - B)*a^3*e^(I*d*x + I*c)
- sqrt(-(A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(a
/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - sqrt(-(2
5*A^2 - 20*I*A*B - 4*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log(-16*(3*
(-5*I*A - 2*B)*a^3*e^(2*I*d*x + 2*I*c) + (-5*I*A - 2*B)*a^3 + 2*sqrt(2)*sq
rt(-(25*A^2 - 20*I*A*B - 4*B^2)*a^5/d^2)*(d*e^(3*I*d*x + 3*I*c) + d*e^(I*d
*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x - 2*I*c)/((5*I*A
+ 2*B)*a) + sqrt(-(25*A^2 - 20*I*A*B - 4*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2
*I*c) - d)*log(-16*(3*(-5*I*A - 2*B)*a^3*e^(2*I*d*x + 2*I*c) + (-5*I*A - 2
*B)*a^3 - 2*sqrt(2)*sqrt(-(25*A^2 - 20*I*A*B - 4*B^2)*a^5/d^2)*(d*e^(3*I*d
*x + 3*I*c) + d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*
I*d*x - 2*I*c)/((5*I*A + 2*B)*a) + 4*sqrt(2)*((I*A + 2*B)*a^2*e^(3*I*d*x
+ 3*I*c) + (I*A - 2*B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) +
1)))/(d*e^(2*I*d*x + 2*I*c) - d)

```

3.86.6 Sympy [F(-1)]

Timed out.

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**2*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output Timed out

3.86.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.03

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx =$$

$$\frac{i \left(4 \sqrt{2}(A - iB) a^{\frac{3}{2}} \log \left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}} \right) - (5A - 2iB) a^{\frac{3}{2}} \log \left(\frac{\sqrt{ia \tan(dx+c)+a} - \sqrt{a}}{\sqrt{ia \tan(dx+c)+a} + \sqrt{a}} \right) - 4i \sqrt{ia \tan(dx+c)+a} \right)}{2d}$$

3.86. $\int \cot^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*I*(4*sqrt(2)*(A - I*B)*a^(3/2)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - (5*A - 2*I*B)*a^(3/2)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a))) - 4*I*sqrt(I*a*tan(d*x + c) + a)*B*a - 2*I*sqrt(I*a*tan(d*x + c) + a)*A*a/tan(d*x + c))*a/d`

3.86.8 Giac [F]

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{5/2} \cot(dx + c)^2 dx$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^2, x)`

3.86.9 Mupad [B] (verification not implemented)

Time = 8.99 (sec) , antiderivative size = 2947, normalized size of antiderivative = 18.65

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),x)`

output $\operatorname{atan}((d^4(a + a\tan(c + dx)*i))^{1/2} * (((49A^4a^{22})/d^4 + (784B^4a^{22})/d^4 - (2328A^2B^2a^{22})/d^4 + (AB^3a^{22}*2464i)/d^4 - (A^3Ba^{22}*616i)/d^4)^{1/2} / (8a^6) - (57A^2a^5)/(8d^2) + (9B^2a^5)/(2d^2) + (ABa^5*21i)/(2d^2))^{1/2} * (((49A^4a^{22})/d^4 + (784B^4a^{22})/d^4 - (2328A^2B^2a^{22})/d^4 + (AB^3a^{22}*2464i)/d^4 - (A^3Ba^{22}*616i)/d^4)^{1/2} * 6i) / (A^3a^{14}*d*126i - 336B^3a^{14}*d - AB^2a^{14}*d*1032i + 876A^2B*a^{14}*d + Aa^3*d^3 * ((49A^4a^{22})/d^4 + (784B^4a^{22})/d^4 - (2328A^2B^2a^{22})/d^4 + (AB^3a^{22}*2464i)/d^4 - (A^3Ba^{22}*616i)/d^4)^{1/2} * 2i - 4B*a^3*d^3 * ((49A^4a^{22})/d^4 + (784B^4a^{22})/d^4 - (2328A^2B^2a^{22})/d^4 + (AB^3a^{22}*2464i)/d^4 - (A^3Ba^{22}*616i)/d^4)^{1/2})) + (A^2a^8*d^2*(a + a\tan(c + dx)*i))^{1/2} * (((49A^4a^{22})/d^4 + (784B^4a^{22})/d^4 - (2328A^2B^2a^{22})/d^4 + (AB^3a^{22}*2464i)/d^4 - (A^3Ba^{22}*616i)/d^4)^{1/2} / (8a^6) - (57A^2a^5)/(8d^2) + (9B^2a^5)/(2d^2) + (ABa^5*21i)/(2d^2))^{1/2} * 14i) / (A^3a^{11}*d*126i - 336B^3a^{11}*d + A*d^3 * ((49A^4a^{22})/d^4 + (784B^4a^{22})/d^4 - (2328A^2B^2a^{22})/d^4 + (AB^3a^{22}*2464i)/d^4 - (A^3Ba^{22}*616i)/d^4)^{1/2} * 2i - 4B*d^3 * ((49A^4a^{22})/d^4 + (784B^4a^{22})/d^4 - (2328A^2B^2a^{22})/d^4 + (AB^3a^{22}*2464i)/d^4 - (A^3Ba^{22}*616i)/d^4)^{1/2} - AB^2a^{11}*d*1032i + 876A^2B*a^{11}*d) - (B^2a^8*d^2 * (a + a\tan(c + dx)*i))^{1/2} * (((49A^4a^{22})/d^4 + (784B^4a^{22})/d^4 - (2328A^2B^2a^{22})/d^4 + (AB^3a^{22}*2464i)/d^4 - (A^3Ba^{22}*616i)/d^4...$

3.87 $\int \cot^3(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.87.1	Optimal result	977
3.87.2	Mathematica [A] (verified)	978
3.87.3	Rubi [A] (verified)	978
3.87.4	Maple [A] (verified)	982
3.87.5	Fricas [B] (verification not implemented)	983
3.87.6	Sympy [F(-1)]	984
3.87.7	Maxima [A] (verification not implemented)	984
3.87.8	Giac [F]	985
3.87.9	Mupad [B] (verification not implemented)	985

3.87.1 Optimal result

Integrand size = 36, antiderivative size = 173

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \frac{a^{5/2}(23A-20iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{4d} - \frac{4\sqrt{2}a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{a^2(7iA+4B)\cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{4d} - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^{3/2}}{2d}$$

```
output 1/4*a^(5/2)*(23*A-20*I*B)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/d-4*a^(5/2)*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(1/2)/d-1/4*a^2*(7*I*A+4*B)*cot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)/d-1/2*a*A*cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(3/2)/d
```


3.87.2 Mathematica [A] (verified)

Time = 5.86 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.82

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \frac{a^{5/2}(23A-20iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) - 16\sqrt{2}a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{4d}$$

input `Integrate[Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(a^(5/2)*(23*A - (20*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] - 16*Sqrt[2]*a^(5/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] - a^2*Cot[c + d*x]*((9*I)*A + 4*B + 2*A*Cot[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]])/(4*d)`

3.87.3 Rubi [A] (verified)

Time = 1.16 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.05, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3042, 4076, 27, 3042, 4076, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot^3(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan(c+dx)^3} dx \\ & \quad \downarrow \text{4076} \\ & \frac{1}{2} \int \frac{1}{2} \cot^2(c+dx)(i \tan(c+dx)a+a)^{3/2}(a(7iA+4B) - a(A-4iB) \tan(c+dx)) dx - \\ & \quad \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^{3/2}}{2d} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\frac{1}{4} \int \cot^2(c+dx)(i \tan(c+dx)a+a)^{3/2}(a(7iA+4B)-a(A-4iB)\tan(c+dx))dx - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^{3/2}}{2d}$$

↓ 3042

$$\frac{1}{4} \int \frac{(i \tan(c+dx)a+a)^{3/2}(a(7iA+4B)-a(A-4iB)\tan(c+dx))}{\tan(c+dx)^2} dx - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^{3/2}}{2d}$$

↓ 4076

$$\frac{1}{4} \left(\int -\frac{1}{2} \cot(c+dx) \sqrt{i \tan(c+dx)a+a} ((23A-20iB)a^2+3(3iA+4B)\tan(c+dx)a^2) dx - \frac{a^2(4B+7iA) \cot(c+dx)}{d} \right) - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^{3/2}}{2d}$$

↓ 27

$$\frac{1}{4} \left(-\frac{1}{2} \int \cot(c+dx) \sqrt{i \tan(c+dx)a+a} ((23A-20iB)a^2+3(3iA+4B)\tan(c+dx)a^2) dx - \frac{a^2(4B+7iA) \cot(c+dx)}{d} \right) - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^{3/2}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(-\frac{1}{2} \int \frac{\sqrt{i \tan(c+dx)a+a} ((23A-20iB)a^2+3(3iA+4B)\tan(c+dx)a^2)}{\tan(c+dx)} dx - \frac{a^2(4B+7iA) \cot(c+dx) \sqrt{i \tan(c+dx)a+a}}{d} \right) - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^{3/2}}{2d}$$

↓ 4083

$$\frac{1}{4} \left(\frac{1}{2} \left(-32a^2(B+iA) \int \sqrt{i \tan(c+dx)a+adx} - a(23A-20iB) \int \cot(c+dx)(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a} dx \right) \right) - \frac{aA \cot^2(c+dx)(a+ia \tan(c+dx))^{3/2}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{2} \left(-32a^2(B + iA) \int \frac{\sqrt{i \tan(c + dx)a + adx} - a(23A - 20iB) \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} dx}{\frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^{3/2}}{2d}} \right) \right)$$

↓ 3961

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{64ia^3(B + iA) \int \frac{1}{a - ia \tan(c + dx)} d\sqrt{i \tan(c + dx)a + a} - a(23A - 20iB) \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} dx}{\frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^{3/2}}{2d}} \right) \right)$$

↓ 219

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{32i\sqrt{2}a^{5/2}(B + iA) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right) - a(23A - 20iB) \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\tan(c + dx)} dx}{\frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^{3/2}}{2d}} \right) \right)$$

↓ 4082

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{32i\sqrt{2}a^{5/2}(B + iA) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right) - \frac{a^3(23A - 20iB) \int \frac{\cot(c + dx)}{\sqrt{i \tan(c + dx)a + a}} d \tan(c + dx)}{d} - \frac{a^2(4B - 3A)}{d}}{\frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^{3/2}}{2d}} \right) \right)$$

↓ 73

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{2ia^2(23A - 20iB) \int \frac{1}{i - i(i \tan(c + dx)a + a)} d\sqrt{i \tan(c + dx)a + a} + \frac{32i\sqrt{2}a^{5/2}(B + iA) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{\frac{aA \cot^2(c + dx)(a + ia \tan(c + dx))^{3/2}}{2d}} \right) \right)$$

↓ 221

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{2a^{5/2}(23A - 20iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{a}}\right) + \frac{32i\sqrt{2}a^{5/2}(B + iA) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{d} - \frac{a^2(4B - 3A)}{d} \right) \right)$$

↓

3.87. $\int \cot^3(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

input `Int[Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `-1/2*(a*A*Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^(3/2))/d + (((2*a^(5/2)*(2
3*A - (20*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d + ((32*I)*S
qrt[2]*a^(5/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[
a]]))/d)/2 - (a^2*((7*I)*A + 4*B)*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]]
/d)/4`

3.87.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d)
Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a
, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int[(((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.87.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.83

method	result
derivativedivides	$2a^3 \left(-\frac{(-4iB+4A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2\sqrt{a}} - \frac{\left(-\frac{iB}{2} + \frac{9A}{8}\right)(a+ia \tan(dx+c))^{\frac{3}{2}} + \left(\frac{1}{2}iaB - \frac{7}{8}aA\right)\sqrt{a+ia \tan(dx+c)}}{a^2 \tan(dx+c)^2} + \dots \right)$
default	$2a^3 \left(-\frac{(-4iB+4A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2\sqrt{a}} - \frac{\left(-\frac{iB}{2} + \frac{9A}{8}\right)(a+ia \tan(dx+c))^{\frac{3}{2}} + \left(\frac{1}{2}iaB - \frac{7}{8}aA\right)\sqrt{a+ia \tan(dx+c)}}{a^2 \tan(dx+c)^2} + \dots \right)$

input `int(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)), x, method=_RETURNVERBOSE)`

$$3.87. \int \cot^3(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

output $2/d*a^3*(-1/2*(-4*I*B+4*A)*2^(1/2)/a^(1/2)*\operatorname{arctanh}(1/2*(a+I*a*\tan(dx+c))^(1/2)*2^(1/2)/a^(1/2))-((-1/2*I*B+9/8*A)*(a+I*a*\tan(dx+c))^(3/2)+(1/2*I*a*B-7/8*a*A)*(a+I*a*\tan(dx+c))^(1/2))/a^2/\tan(dx+c)^2+1/8*(23*A-20*I*B)/a^(1/2)*\operatorname{arctanh}((a+I*a*\tan(dx+c))^(1/2)/a^(1/2))$

3.87.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 774 vs. $2(134) = 268$.

Time = 0.27 (sec) , antiderivative size = 774, normalized size of antiderivative = 4.47

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(dx+c)^3*(a+I*a*tan(dx+c))^(5/2)*(A+B*tan(dx+c)),x, algorithm="fracas")`

output $-1/16*(32*\sqrt{2}*\sqrt{(A^2 - 2*I*A*B - B^2)*a^5/d^2}*(d*e^{(4*I*d*x + 4*I*c)} - 2*d*e^{(2*I*d*x + 2*I*c)} + d)*\log(4*((-I*A - B)*a^3*e^{(I*d*x + I*c)} - \sqrt{(A^2 - 2*I*A*B - B^2)*a^5/d^2}*(I*d*e^{(2*I*d*x + 2*I*c)} + I*d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}))e^{-I*d*x - I*c}/((-I*A - B)*a^2)) - 32*\sqrt{2}*\sqrt{(A^2 - 2*I*A*B - B^2)*a^5/d^2}*(d*e^{(4*I*d*x + 4*I*c)} - 2*d*e^{(2*I*d*x + 2*I*c)} + d)*\log(4*((-I*A - B)*a^3*e^{(I*d*x + I*c)} - \sqrt{(A^2 - 2*I*A*B - B^2)*a^5/d^2}*(-I*d*e^{(2*I*d*x + 2*I*c)} - I*d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}))e^{-I*d*x - I*c}/((-I*A - B)*a^2)) + \sqrt{(529*A^2 - 920*I*A*B - 400*B^2)*a^5/d^2}*(d*e^{(4*I*d*x + 4*I*c)} - 2*d*e^{(2*I*d*x + 2*I*c)} + d)*\log(-16*(3*(-23*I*A - 20*B)*a^3*e^{(2*I*d*x + 2*I*c)} + (-23*I*A - 20*B)*a^3 + 2*\sqrt{2}*\sqrt{(529*A^2 - 920*I*A*B - 400*B^2)*a^5/d^2}*(I*d*e^{(3*I*d*x + 3*I*c)} + I*d*e^{(I*d*x + I*c)})*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}))e^{-2*I*d*x - 2*I*c}/((23*I*A + 20*B)*a) - \sqrt{(529*A^2 - 920*I*A*B - 400*B^2)*a^5/d^2}*(d*e^{(4*I*d*x + 4*I*c)} - 2*d*e^{(2*I*d*x + 2*I*c)} + d)*\log(-16*(3*(-23*I*A - 20*B)*a^3*e^{(2*I*d*x + 2*I*c)} + (-23*I*A - 20*B)*a^3 + 2*\sqrt{2}*\sqrt{(529*A^2 - 920*I*A*B - 400*B^2)*a^5/d^2}*(-I*d*e^{(3*I*d*x + 3*I*c)} - I*d*e^{(I*d*x + I*c)})*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}))e^{-2*I*d*x - 2*I*c}/((23*I*A + 20*B)*a) - 4*\sqrt{2}*((11*A - 4*I*B)*a^2*e^{(5*I*d*x + 5*I*c)} + 4*A*a^2*e^{(3*I*d*x + 3*I*c)} - (7*A - 4*I*B)*a^2*e^{(I*d*x + I*c)})*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})/(d*e^{(4*I*d*x + 4*I*c)} - 2*d*e^{(2*I*d*x + 2*I*c)} + d)$

3.87.6 Sympy [F(-1)]

Timed out.

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**3*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.87.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.19

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{\left(16 \sqrt{2}(A - iB)\sqrt{a} \log\left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}}\right) - (23A - 20iB)\sqrt{a} \log\left(\frac{\sqrt{ia \tan(dx+c)+a}}{\sqrt{ia \tan(dx+c)+a}}\right)\right)}{8d}$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/8*(16*sqrt(2)*(A - I*B)*sqrt(a)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - (23*A - 20*I*B)*sqrt(a)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a))) + 2*((I*a*tan(d*x + c) + a)^(3/2)*(9*A - 4*I*B)*a - sqrt(I*a*tan(d*x + c) + a)*(7*A - 4*I*B)*a^2)/((I*a*tan(d*x + c) + a)^2 - 2*(I*a*tan(d*x + c) + a)*a + a^2))*a^2/d`

3.87.8 Giac [F]

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{5/2} \cot(dx + c)^3 dx$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^3, x)`

3.87.9 Mupad [B] (verification not implemented)

Time = 9.47 (sec) , antiderivative size = 2991, normalized size of antiderivative = 17.29

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),x)`

output

$$\begin{aligned}
& 2*\operatorname{atanh}((17*A^2*a^8*d^2*(a + a*\tan(c + d*x)*i))^{1/2}*((1041*A^2*a^5)/(128 \\
& *d^2) - ((289*A^4*a^22)/(4*d^4) + (3136*B^4*a^22)/d^4 - (1752*A^2*B^2*a^22 \\
&)/d^4 + (A*B^3*a^22*5824i)/d^4 + (A^3*B*a^22*884i)/d^4)^{1/2})/(64*a^6) - (\\
& 57*B^2*a^5)/(8*d^2) - (A*B*a^5*243i)/(16*d^2))^{1/2})/(4*((663*A^3*a^11*d) \\
& /16 - B^3*a^11*d*252i - (7*A*d^3*((289*A^4*a^22)/(4*d^4) + (3136*B^4*a^22) \\
& /d^4 - (1752*A^2*B^2*a^22)/d^4 + (A*B^3*a^22*5824i)/d^4 + (A^3*B*a^22*884i \\
&)/d^4)^{1/2}))/8 + (B*d^3*((289*A^4*a^22)/(4*d^4) + (3136*B^4*a^22)/d^4 - (\\
& 1752*A^2*B^2*a^22)/d^4 + (A*B^3*a^22*5824i)/d^4 + (A^3*B*a^22*884i)/d^4)^{1/2} \\
& *i)/2 + 507*A*B^2*a^11*d + (A^2*B*a^11*d*861i)/4) - (3*d^4*(a + a*\tan \\
& (c + d*x)*i))^{1/2}*((1041*A^2*a^5)/(128*d^2) - ((289*A^4*a^22)/(4*d^4) + \\
& (3136*B^4*a^22)/d^4 - (1752*A^2*B^2*a^22)/d^4 + (A*B^3*a^22*5824i)/d^4 + \\
& (A^3*B*a^22*884i)/d^4)^{1/2})/(64*a^6) - (57*B^2*a^5)/(8*d^2) - (A*B*a^5*24 \\
& 3i)/(16*d^2))^{1/2}*((289*A^4*a^22)/(4*d^4) + (3136*B^4*a^22)/d^4 - (1752* \\
& A^2*B^2*a^22)/d^4 + (A*B^3*a^22*5824i)/d^4 + (A^3*B*a^22*884i)/d^4)^{1/2} \\
& / (2*((663*A^3*a^14*d)/16 - B^3*a^14*d*252i + 507*A*B^2*a^14*d + (A^2*B*a^1 \\
& 4*d*861i)/4 - (7*A*a^3*d^3*((289*A^4*a^22)/(4*d^4) + (3136*B^4*a^22)/d^4 - \\
& (1752*A^2*B^2*a^22)/d^4 + (A*B^3*a^22*5824i)/d^4 + (A^3*B*a^22*884i)/d^4) \\
& ^{1/2}))/8 + (B*a^3*d^3*((289*A^4*a^22)/(4*d^4) + (3136*B^4*a^22)/d^4 - (17 \\
& 52*A^2*B^2*a^22)/d^4 + (A*B^3*a^22*5824i)/d^4 + (A^3*B*a^22*884i)/d^4)^{1/2} \\
& *i)/2)) + (28*B^2*a^8*d^2*(a + a*\tan(c + d*x)*i))^{1/2}*((1041*A^2*a...
\end{aligned}$$

3.88 $\int \cot^4(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.88.1	Optimal result	987
3.88.2	Mathematica [A] (verified)	988
3.88.3	Rubi [A] (verified)	988
3.88.4	Maple [A] (verified)	993
3.88.5	Fricas [B] (verification not implemented)	994
3.88.6	Sympy [F(-1)]	995
3.88.7	Maxima [A] (verification not implemented)	996
3.88.8	Giac [F]	996
3.88.9	Mupad [B] (verification not implemented)	997

3.88.1 Optimal result

Integrand size = 36, antiderivative size = 217

$$\int \cot^4(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \frac{a^{5/2}(45iA+46B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{8d} - \frac{4\sqrt{2}a^{5/2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{a^2(19A-18iB)\cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{8d} - \frac{a^2(3iA+2B)\cot^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{4d} - \frac{aA\cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d}$$

output $\frac{1}{8}a^{5/2}(45iA+46B)\operatorname{arctanh}\left(\frac{(a+i a \tan(dx+c))^{1/2}}{a^{1/2}}\right)/d-4a^{5/2}(iA+B)\operatorname{arctanh}\left(\frac{1}{2}\frac{(a+i a \tan(dx+c))^{1/2}}{a^{1/2}}\right)^2/d+1/8a^2(19A-18iB)\cot(dx+c)(a+i a \tan(dx+c))^{1/2}/d-1/4a^2(3iA+2B)\cot^2(dx+c)(a+i a \tan(dx+c))^{1/2}/d-1/3aA\cot^3(dx+c)(a+i a \tan(dx+c))^{3/2}/d$

3.88.2 Mathematica [A] (verified)

Time = 5.54 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.74

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{-3a^{5/2}(45iA + 46B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) + 96\sqrt{2}a^{5/2}(iA + B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) + a^2 \cot(c + dx)}{24d}$$

input `Integrate[Cot[c + d*x]^4*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `-1/24*(-3*a^(5/2)*((45*I)*A + 46*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] + 96*Sqrt[2]*a^(5/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] + a^2*Cot[c + d*x]*(-57*A + (54*I)*B + 2*((13*I)*A + 6*B)*Cot[c + d*x] + 8*A*Cot[c + d*x]^2)*Sqrt[a + I*a*Tan[c + d*x]])/d`

3.88.3 Rubi [A] (verified)

Time = 1.47 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.07, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.472$, Rules used = {3042, 4076, 27, 3042, 4076, 27, 3042, 4081, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot^4(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan(c + dx)^4} dx \\ & \quad \downarrow \text{4076} \\ & \frac{1}{3} \int \frac{3}{2} \cot^3(c + dx)(i \tan(c + dx)a + a)^{3/2}(a(3iA + 2B) - a(A - 2iB) \tan(c + dx)) dx - \\ & \quad \frac{aA \cot^3(c + dx)(a + ia \tan(c + dx))^{3/2}}{3d} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.88. $\int \cot^4(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\frac{1}{2} \int \cot^3(c+dx)(i \tan(c+dx)a+a)^{3/2}(a(3iA+2B)-a(A-2iB)\tan(c+dx))dx - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d}$$

↓ 3042

$$\frac{1}{2} \int \frac{(i \tan(c+dx)a+a)^{3/2}(a(3iA+2B)-a(A-2iB)\tan(c+dx))}{\tan(c+dx)^3} dx - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d}$$

↓ 4076

$$\frac{1}{2} \left(\frac{1}{2} \int -\frac{1}{2} \cot^2(c+dx)\sqrt{i \tan(c+dx)a+a}((19A-18iB)a^2+(13iA+14B)\tan(c+dx)a^2) dx - \frac{a^2(2B+3iA)}{3d} \right) - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d}$$

↓ 27

$$\frac{1}{2} \left(-\frac{1}{4} \int \cot^2(c+dx)\sqrt{i \tan(c+dx)a+a}((19A-18iB)a^2+(13iA+14B)\tan(c+dx)a^2) dx - \frac{a^2(2B+3iA)}{2d} \right) - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d}$$

↓ 3042

$$\frac{1}{2} \left(-\frac{1}{4} \int \frac{\sqrt{i \tan(c+dx)a+a}((19A-18iB)a^2+(13iA+14B)\tan(c+dx)a^2)}{\tan(c+dx)^2} dx - \frac{a^2(2B+3iA)\cot^2(c+dx)}{2d} \right) - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d}$$

↓ 4081

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19A-18iB)\cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{d} - \frac{\int \frac{1}{2} \cot(c+dx)\sqrt{i \tan(c+dx)a+a}(a^3(45iA+46B)-a^2(2B+3iA)\cot^2(c+dx))}{a} \right) - \frac{aA \cot^3(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d} \right)$$

↓ 27

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19A - 18iB) \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d} - \frac{\int \cot(c + dx) \sqrt{i \tan(c + dx) a + a} (a^3(45iA + 46B) - \dots)}{2a} \right) \right. \\ \left. \frac{aA \cot^3(c + dx) (a + ia \tan(c + dx))^{3/2}}{3d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19A - 18iB) \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d} - \frac{\int \frac{\sqrt{i \tan(c + dx) a + a} (a^3(45iA + 46B) - a^3(19A - 18iB) \tan(c + dx))}{\tan(c + dx)} dx}{2a} \right) \right. \\ \left. \frac{aA \cot^3(c + dx) (a + ia \tan(c + dx))^{3/2}}{3d} \right) \\ \downarrow \text{4083}$$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19A - 18iB) \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d} - \frac{a^2(46B + 45iA) \int \cot(c + dx) (a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + a}}{2a} \right) \right. \\ \left. \frac{aA \cot^3(c + dx) (a + ia \tan(c + dx))^{3/2}}{3d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19A - 18iB) \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d} - \frac{a^2(46B + 45iA) \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + a}}{\tan(c + dx)} dx - 6}{2a} \right) \right. \\ \left. \frac{aA \cot^3(c + dx) (a + ia \tan(c + dx))^{3/2}}{3d} \right) \\ \downarrow \text{3961}$$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19A - 18iB) \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d} - \frac{128ia^4(A - iB) \int \frac{1}{a - ia \tan(c + dx)} d \sqrt{i \tan(c + dx) a + a}}{d} + a^2(46B + \dots)}{2a} \right) \right. \\ \left. \frac{aA \cot^3(c + dx) (a + ia \tan(c + dx))^{3/2}}{3d} \right) \\ \downarrow \text{219}$$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19A - 18iB) \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d} - \frac{a^2(46B + 45iA) \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + a}}{\tan(c + dx)} dx + \dots}{2a} \right) \right. \\ \left. \frac{aA \cot^3(c + dx) (a + ia \tan(c + dx))^{3/2}}{3d} \right)$$

3.88. $\int \cot^4(c + dx) (a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx$

↓ 4082

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19A - 18iB) \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d} - \frac{a^4(46B + 45iA) \int \frac{\cot(c + dx)}{\sqrt{i \tan(c + dx)a + a}} d \tan(c + dx)}{d} + \frac{64i\sqrt{2}a^{7/2}(A - iB)}{2a} \right) \right. \\ \left. - \frac{aA \cot^3(c + dx)(a + ia \tan(c + dx))^{3/2}}{3d} \right)$$

↓ 73

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19A - 18iB) \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d} - \frac{64i\sqrt{2}a^{7/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{2ia^3(46B + 45iA)}{2a} \right) \right. \\ \left. - \frac{aA \cot^3(c + dx)(a + ia \tan(c + dx))^{3/2}}{3d} \right)$$

↓ 221

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19A - 18iB) \cot(c + dx) \sqrt{a + ia \tan(c + dx)}}{d} - \frac{64i\sqrt{2}a^{7/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{2a^{7/2}(46B + 45iA)}{2a} \right) \right. \\ \left. - \frac{aA \cot^3(c + dx)(a + ia \tan(c + dx))^{3/2}}{3d} \right)$$

input `Int[Cot[c + d*x]^4*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `-1/3*(a*A*Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^(3/2))/d + (-1/2*(a^2*((3*I)*A + 2*B)*Cot[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]])/d + (-1/2*((-2*a^(7/2))*((45*I)*A + 46*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]]))/d + ((6*4*I)*Sqrt[2]*a^(7/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d)/a + (a^2*(19*A - (18*I)*B)*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]])/d)/4)/2`

3.88.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`
- rule 4076 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2)), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

```
rule 4082 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 + b^2, 0] && EqQ[A*b + a*B, 0]
```

```
rule 4083 Int((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(
A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A
*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*T
an[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

3.88.4 Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.82

method	result
derivativedivides	$2ia^4 \left(-\frac{(-4iB+4A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2a^{\frac{3}{2}}} + \frac{i\left(\frac{9iB}{8}-\frac{19A}{16}\right)(a+ia \tan(dx+c))^{\frac{5}{2}} + (-2iaB+\frac{11}{6}aA)(a+ia \tan(dx+c))^{\frac{5}{2}}}{a^3 \tan(dx+c)^3} \right)$
default	$2ia^4 \left(-\frac{(-4iB+4A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2a^{\frac{3}{2}}} + \frac{i\left(\frac{9iB}{8}-\frac{19A}{16}\right)(a+ia \tan(dx+c))^{\frac{5}{2}} + (-2iaB+\frac{11}{6}aA)(a+ia \tan(dx+c))^{\frac{5}{2}}}{a^3 \tan(dx+c)^3} \right)$

3.88. $\int \cot^4(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$


```
input int(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 2*I/d*a^4*(-1/2*(-4*I*B+4*A)/a^(3/2)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))+1/a*(-I*((9/8*I*B-19/16*A)*(a+I*a*tan(d*x+c))^(5/2))+(-2*I*a*B+11/6*a*A)*(a+I*a*tan(d*x+c))^(3/2)+(7/8*I*B*a^2-13/16*A*a^2)*(a+I*a*tan(d*x+c))^(1/2))/a^3/tan(d*x+c)^3+1/16*(-46*I*B+45*A)/a^(1/2)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2)))
```

3.88.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 868 vs. $2(170) = 340$.

Time = 0.28 (sec) , antiderivative size = 868, normalized size of antiderivative = 4.00

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fracas")
```

```

output 1/96*(192*sqrt(2)*sqrt(-(A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(6*I*d*x + 6*I
*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*log(4*((-I*A
- B)*a^3*e^(I*d*x + I*c) + sqrt(-(A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(2*I*
d*x + 2*I*c) + d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/((-I
*A - B)*a^2)) - 192*sqrt(2)*sqrt(-(A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(6*I
*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*log
(4*((-I*A - B)*a^3*e^(I*d*x + I*c) - sqrt(-(A^2 - 2*I*A*B - B^2)*a^5/d^2)*
(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x -
I*c)/((-I*A - B)*a^2)) - 3*sqrt(-(2025*A^2 - 4140*I*A*B - 2116*B^2)*a^5/d
^2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*
I*c) - d)*log(-16*(3*(-45*I*A - 46*B)*a^3*e^(2*I*d*x + 2*I*c) + (-45*I*A -
46*B)*a^3 + 2*sqrt(2)*sqrt(-(2025*A^2 - 4140*I*A*B - 2116*B^2)*a^5/d^2)*(
d*e^(3*I*d*x + 3*I*c) + d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1
)))e^(-2*I*d*x - 2*I*c)/((45*I*A + 46*B)*a)) + 3*sqrt(-(2025*A^2 - 4140*I
*A*B - 2116*B^2)*a^5/d^2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c)
+ 3*d*e^(2*I*d*x + 2*I*c) - d)*log(-16*(3*(-45*I*A - 46*B)*a^3*e^(2*I*d*x
+ 2*I*c) + (-45*I*A - 46*B)*a^3 - 2*sqrt(2)*sqrt(-(2025*A^2 - 4140*I*A*B
- 2116*B^2)*a^5/d^2)*(d*e^(3*I*d*x + 3*I*c) + d*e^(I*d*x + I*c))*sqrt(a/(e
^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((45*I*A + 46*B)*a)) + 4*sq
rt(2)*((91*I*A + 66*B)*a^2*e^(7*I*d*x + 7*I*c) - 7*(I*A + 6*B)*a^2*e^(5...

```

3.88.6 Sympy [F(-1)]

Timed out.

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

```
input integrate(cot(d*x+c)**4*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)
```

```
output Timed out
```

3.88.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.15

$$\int \cot^4(c+dx)(a+ia \tan(c+dx))^{5/2}(A + B \tan(c+dx)) dx = \frac{i \left(\frac{96\sqrt{2}(A-iB) \log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia \tan(dx+c)+a}}\right)}{\sqrt{a}} - \frac{3(45A-46iB) \log\left(\frac{\sqrt{ia \tan(dx+c)+a}-\sqrt{a}}{\sqrt{ia \tan(dx+c)+a}+\sqrt{a}}\right)}{\sqrt{a}} + \frac{2(3ia \tan(dx+c)+a)^{5/2}}{48d} \right)}{48d}$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/48*I*(96*sqrt(2)*(A - I*B)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a)))/sqrt(a) - 3*(45*A - 46*I*B)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a)))/sqrt(a) + 2*(3*(I*a*tan(d*x + c) + a)^(5/2)*(19*A - 18*I*B) - 8*(I*a*tan(d*x + c) + a)^(3/2)*(11*A - 12*I*B)*a + 3*sqrt(I*a*tan(d*x + c) + a)*(13*A - 14*I*B)*a^2)/((I*a*tan(d*x + c) + a)^3 - 3*(I*a*tan(d*x + c) + a)^2*a + 3*(I*a*tan(d*x + c) + a)*a^2 - a^3))*a^3/d`

3.88.8 Giac [F]

$$\int \cot^4(c+dx)(a+ia \tan(c+dx))^{5/2}(A + B \tan(c+dx)) dx = \int (B \tan(dx+c) + A)(ia \tan(dx+c) + a)^{5/2} \cot(dx+c)^4 dx$$

input `integrate(cot(d*x+c)^4*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^4, x)`

3.88.9 Mupad [B] (verification not implemented)

Time = 9.41 (sec) , antiderivative size = 3048, normalized size of antiderivative = 14.05

$$\int \cot^4(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(cot(c + d*x)^4*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),x)
```

```
output 2*atanh((23*A^2*a^8*d^2*(a + a*tan(c + d*x)*1i)^(1/2)*((1041*B^2*a^5)/(128
*d^2) - (4073*A^2*a^5)/(512*d^2) - ((529*A^4*a^22)/(64*d^4) + (289*B^4*a^2
2)/(4*d^4) + (149*A^2*B^2*a^22)/(8*d^4) + (A*B^3*a^22*187i)/(2*d^4) + (A^3
*B*a^22*253i)/(8*d^4))^(1/2)/(64*a^6) + (A*B*a^5*2059i)/(128*d^2))^(1/2))/
(4*((A^3*a^11*d*1771i)/32 + (663*B^3*a^11*d)/4 - (A*d^3*((529*A^4*a^22)/(6
4*d^4) + (289*B^4*a^22)/(4*d^4) + (149*A^2*B^2*a^22)/(8*d^4) + (A*B^3*a^22
*187i)/(2*d^4) + (A^3*B*a^22*253i)/(8*d^4))^(1/2)*13i)/4 - (7*B*d^3*((529*
A^4*a^22)/(64*d^4) + (289*B^4*a^22)/(4*d^4) + (149*A^2*B^2*a^22)/(8*d^4) +
(A*B^3*a^22*187i)/(2*d^4) + (A^3*B*a^22*253i)/(8*d^4))^(1/2))/2 + (A*B^2*
a^11*d*2167i)/8 - (797*A^2*B*a^11*d)/16)) - (6*d^4*(a + a*tan(c + d*x)*1i)
^(1/2)*((1041*B^2*a^5)/(128*d^2) - (4073*A^2*a^5)/(512*d^2) - ((529*A^4*a^
22)/(64*d^4) + (289*B^4*a^22)/(4*d^4) + (149*A^2*B^2*a^22)/(8*d^4) + (A*B^
3*a^22*187i)/(2*d^4) + (A^3*B*a^22*253i)/(8*d^4))^(1/2)/(64*a^6) + (A*B*a^
5*2059i)/(128*d^2))^(1/2)*((529*A^4*a^22)/(64*d^4) + (289*B^4*a^22)/(4*d^4
) + (149*A^2*B^2*a^22)/(8*d^4) + (A*B^3*a^22*187i)/(2*d^4) + (A^3*B*a^22*2
53i)/(8*d^4))^(1/2))/((A^3*a^14*d*1771i)/32 + (663*B^3*a^14*d)/4 + (A*B^2*
a^14*d*2167i)/8 - (797*A^2*B*a^14*d)/16 - (A*a^3*d^3*((529*A^4*a^22)/(64*d
^4) + (289*B^4*a^22)/(4*d^4) + (149*A^2*B^2*a^22)/(8*d^4) + (A*B^3*a^22*18
7i)/(2*d^4) + (A^3*B*a^22*253i)/(8*d^4))^(1/2)*13i)/4 - (7*B*a^3*d^3*((529
*A^4*a^22)/(64*d^4) + (289*B^4*a^22)/(4*d^4) + (149*A^2*B^2*a^22)/(8*d^...
```

3.89 $\int \cot^5(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.89.1	Optimal result	998
3.89.2	Mathematica [A] (verified)	999
3.89.3	Rubi [A] (verified)	999
3.89.4	Maple [A] (verified)	1005
3.89.5	Fricas [B] (verification not implemented)	1006
3.89.6	Sympy [F(-1)]	1007
3.89.7	Maxima [A] (verification not implemented)	1007
3.89.8	Giac [F]	1008
3.89.9	Mupad [B] (verification not implemented)	1008

3.89.1 Optimal result

Integrand size = 36, antiderivative size = 261

$$\int \cot^5(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx =$$

$$\frac{3a^{5/2}(121A - 120iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{64d}$$

$$+ \frac{4\sqrt{2}a^{5/2}(A - iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}$$

$$+ \frac{a^2(149iA + 152B) \cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{64d}$$

$$+ \frac{a^2(107A - 104iB) \cot^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{96d}$$

$$- \frac{a^2(11iA + 8B) \cot^3(c+dx)\sqrt{a+ia \tan(c+dx)}}{24d}$$

$$- \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d}$$

output

```
-3/64*a^(5/2)*(121*A-120*I*B)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/d+
4*a^(5/2)*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))*2^(
1/2)/d+1/64*a^2*(149*I*A+152*B)*cot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)/d+1/9
6*a^2*(107*A-104*I*B)*cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(1/2)/d-1/24*a^2*(11
*I*A+8*B)*cot(d*x+c)^3*(a+I*a*tan(d*x+c))^(1/2)/d-1/4*a*A*cot(d*x+c)^4*(a
+I*a*tan(d*x+c))^(3/2)/d
```

3.89.2 Mathematica [A] (verified)

Time = 4.77 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.68

$$\int \cot^5(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \frac{-9a^{5/2}(121A-120iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)+768\sqrt{2}a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{1}$$

input `Integrate[Cot[c + d*x]^5*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(-9*a^(5/2)*(121*A - (120*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] + 768*Sqrt[2]*a^(5/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] + a^2*Cot[c + d*x]*((447*I)*A + 456*B + (214*A - (208*I)*B)*Cot[c + d*x] + ((-136*I)*A - 64*B)*Cot[c + d*x]^2 - 48*A*Cot[c + d*x]^3)*Sqrt[a + I*a*Tan[c + d*x]])/(192*d)`

3.89.3 Rubi [A] (verified)

Time = 1.89 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.09, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 4076, 27, 3042, 4076, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot^5(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan(c+dx)^5} dx \\ & \quad \downarrow \text{4076} \\ & \frac{1}{4} \int \frac{1}{2} \cot^4(c+dx)(i \tan(c+dx)a+a)^{3/2}(a(11iA+8B)-a(5A-8iB) \tan(c+dx)) dx - \\ & \quad \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.89. $\int \cot^5(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\frac{1}{8} \int \cot^4(c+dx)(i \tan(c+dx)a+a)^{3/2}(a(11iA+8B)-a(5A-8iB)\tan(c+dx))dx - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \int \frac{(i \tan(c+dx)a+a)^{3/2}(a(11iA+8B)-a(5A-8iB)\tan(c+dx))}{\tan(c+dx)^4} dx - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d}$$

↓ 4076

$$\frac{1}{8} \left(\frac{1}{3} \int -\frac{1}{2} \cot^3(c+dx) \sqrt{i \tan(c+dx)a+a} ((107A-104iB)a^2 + (85iA+88B)\tan(c+dx)a^2) dx - \frac{a^2(8B+11iA)}{2a} \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d}$$

↓ 27

$$\frac{1}{8} \left(-\frac{1}{6} \int \cot^3(c+dx) \sqrt{i \tan(c+dx)a+a} ((107A-104iB)a^2 + (85iA+88B)\tan(c+dx)a^2) dx - \frac{a^2(8B+11iA)}{2a} \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(-\frac{1}{6} \int \frac{\sqrt{i \tan(c+dx)a+a} ((107A-104iB)a^2 + (85iA+88B)\tan(c+dx)a^2)}{\tan(c+dx)^3} dx - \frac{a^2(8B+11iA) \cot^3(c+dx)}{2a} \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d}$$

↓ 4081

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107A-104iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \int \frac{3}{2} \cot^2(c+dx) \sqrt{i \tan(c+dx)a+a} (a^3(149iA+111B)) dx - \frac{a^3(149iA+111B)}{2a} \right) \right) - \frac{aA \cot^4(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d}$$

↓ 27

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107A - 104iB) \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \int \cot^2(c + dx) \sqrt{i \tan(c + dx) a + a} (a^3(149iA + 152B) - a^3(107A - 104iB) \tan(c + dx))}{4a} \right) - \frac{aA \cot^4(c + dx) (a + ia \tan(c + dx))^{3/2}}{4d} \right)$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107A - 104iB) \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \int \frac{\sqrt{i \tan(c + dx) a + a} (a^3(149iA + 152B) - a^3(107A - 104iB) \tan(c + dx))}{\tan(c + dx)^2}}{4a} \right) - \frac{aA \cot^4(c + dx) (a + ia \tan(c + dx))^{3/2}}{4d} \right)$$

↓ 4081

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107A - 104iB) \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(\int -\frac{1}{2} \cot(c + dx) \sqrt{i \tan(c + dx) a + a} (3(121A - 120iB) a^4 + (149iA + 152B) \tan(c + dx))}{a} \right)}{4d} \right) - \frac{aA \cot^4(c + dx) (a + ia \tan(c + dx))^{3/2}}{4d} \right)$$

↓ 27

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107A - 104iB) \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(-\frac{\int \cot(c + dx) \sqrt{i \tan(c + dx) a + a} (3(121A - 120iB) a^4 + (149iA + 152B) \tan(c + dx))}{2a} \right)}{4d} \right) - \frac{aA \cot^4(c + dx) (a + ia \tan(c + dx))^{3/2}}{4d} \right)$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107A - 104iB) \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(-\frac{\int \frac{\sqrt{i \tan(c + dx) a + a} (3(121A - 120iB) a^4 + (149iA + 152B) \tan(c + dx))}{\tan(c + dx)}}{2a} \right)}{4d} \right) - \frac{aA \cot^4(c + dx) (a + ia \tan(c + dx))^{3/2}}{4d} \right)$$

↓ 4083

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107A - 104iB) \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(-\frac{512a^4(B+iA) \int \sqrt{i \tan(c+dx)a+adx} + 3a^3(121A-120iB)}{2a} \right)}{4d} \right) \right)$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107A - 104iB) \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(-\frac{512a^4(B+iA) \int \sqrt{i \tan(c+dx)a+adx} + 3a^3(121A-120iB)}{2a} \right)}{4d} \right) \right)$$

↓ 3961

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107A - 104iB) \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(-\frac{3a^3(121A-120iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx}{2a} \right)}{4d} \right) \right)$$

↓ 219

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107A - 104iB) \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(-\frac{3a^3(121A-120iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx}{2a} \right)}{4d} \right) \right)$$

↓ 4082

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107A - 104iB) \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - 3 \left(-\frac{3a^5(121A - 120iB) \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx) - \frac{512i\sqrt{2}a^9}{2a}}{2a} \right) \right) \right)$$

$$\frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^{3/2}}{4d}$$

↓ 73

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107A - 104iB) \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - 3 \left(-\frac{6ia^4(121A - 120iB) \int \frac{1}{i - \frac{i \tan(c+dx)}{a}} d \sqrt{i \tan(c+dx)a+a}}{2a} \right) \right) \right)$$

$$\frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^{3/2}}{4d}$$

↓ 221

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107A - 104iB) \cot^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - 3 \left(-\frac{6a^{9/2}(121A - 120iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) - \frac{512i\sqrt{2}}{2a}}{d} \right) \right) \right)$$

$$\frac{aA \cot^4(c + dx)(a + ia \tan(c + dx))^{3/2}}{4d}$$

```
input Int[Cot[c + d*x]^5*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]
```

```
output -1/4*(a*A*Cot[c + d*x]^4*(a + I*a*Tan[c + d*x])^(3/2))/d + (-1/3*(a^2*((11*I)*A + 8*B)*Cot[c + d*x]^3*Sqrt[a + I*a*Tan[c + d*x]])/d + ((a^2*(107*A - (104*I)*B)*Cot[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]])/(2*d) - (3*(-1/2*((-6*a^(9/2)*(121*A - (120*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d - ((512*I)*Sqrt[2]*a^(9/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d)/a - (a^3*((149*I)*A + 152*B)*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]]/d))/(4*a))/6)/8
```

3.89. $\int \cot^5(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.89.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`
- rule 4076 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2)), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.89.4 Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 206, normalized size of antiderivative = 0.79

method	result
derivativedivides	$2a^5 \left(-\frac{(4iB-4A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2a^{\frac{5}{2}}} - \frac{\left(-\frac{19iB}{16} + \frac{149A}{128}\right)(a+ia \tan(dx+c))^{\frac{7}{2}} + \left(\frac{145}{48}iaB - \frac{1127}{384}aA\right)(a+ia \tan(dx+c))^{\frac{5}{2}}}{2a^{\frac{5}{2}}}\right)$
default	$2a^5 \left(-\frac{(4iB-4A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2a^{\frac{5}{2}}} - \frac{\left(-\frac{19iB}{16} + \frac{149A}{128}\right)(a+ia \tan(dx+c))^{\frac{7}{2}} + \left(\frac{145}{48}iaB - \frac{1127}{384}aA\right)(a+ia \tan(dx+c))^{\frac{5}{2}}}{2a^{\frac{5}{2}}}\right)$

input `int(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

$$3.89. \int \cot^5(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

```
output 2/d*a^5*(-1/2*(4*I*B-4*A)/a^(5/2)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))-1/a^2*((-19/16*I*B+149/128*A)*(a+I*a*tan(d*x+c))^(7/2)+(145/48*I*a*B-1127/384*a*A)*(a+I*a*tan(d*x+c))^(5/2)+(-127/48*I*a^2*B+1049/384*A*a^2)*(a+I*a*tan(d*x+c))^(3/2)+(13/16*I*B*a^3-107/128*A*a^3)*(a+I*a*tan(d*x+c))^(1/2))/a^4/tan(d*x+c)^4+3/128*(121*A-120*I*B)/a^(1/2)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))
```

3.89.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 944 vs. $2(206) = 412$.

Time = 0.28 (sec) , antiderivative size = 944, normalized size of antiderivative = 3.62

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output 1/768*(1536*sqrt(2)*sqrt((A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((-I*A - B)*a^3*e^(I*d*x + I*c) - sqrt((A^2 - 2*I*A*B - B^2)*a^5/d^2)*(I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 1536*sqrt(2)*sqrt((A^2 - 2*I*A*B - B^2)*a^5/d^2)*(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((-I*A - B)*a^3*e^(I*d*x + I*c) - sqrt((A^2 - 2*I*A*B - B^2)*a^5/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) + 9*sqrt((14641*A^2 - 29040*I*A*B - 14400*B^2)*a^5/d^2)*(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)*log(16*(3*(-121*I*A - 120*B)*a^3*e^(2*I*d*x + 2*I*c) + (-121*I*A - 120*B)*a^3 + 2*sqrt(2)*sqrt((14641*A^2 - 29040*I*A*B - 14400*B^2)*a^5/d^2)*(I*d*e^(3*I*d*x + 3*I*c) + I*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/((-121*I*A - 120*B)*a) - 9*sqrt((14641*A^2 - 29040*I*A*B - 14400*B^2)*a^5/d^2)*(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)*log(16*(3*(-121*I*A - 120*B)*a^3*e^(2*I*d*x + 2*I*c) + (-121*I*A - 120*B)*a^3 + 2*sqrt(2)*sqrt((14641*A^2 - 29040*I*A*B - 14400*B^2)*a^5/d^2)*(-I*d*e^(3*I*d*x + 3*I*c) - I*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d...
```

3.89. $\int \cot^5(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.89.6 Sympy [F(-1)]

Timed out.

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**5*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.89.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 293, normalized size of antiderivative = 1.12

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx =$$

$$a^4 \left(\frac{768 \sqrt{2}(A - iB) \log\left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c) + a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c) + a}}\right)}{a^{\frac{3}{2}}} - \frac{9(121A - 120iB) \log\left(\frac{\sqrt{ia \tan(dx+c) + a} - \sqrt{a}}{\sqrt{ia \tan(dx+c) + a} + \sqrt{a}}\right)}{a^{\frac{3}{2}}} + \frac{2(3(ia \tan(dx+c) + a)^{\frac{7}{2}}(149A - 152iB) - (ia \tan(dx+c) + a)^{\frac{5}{2}}(1127A - 1160iB)a + (ia \tan(dx+c) + a)^{\frac{3}{2}}(1049A - 1016iB)a^2 - 3\sqrt{ia \tan(dx+c) + a}(107A - 104iB)a^3)/((ia \tan(dx+c) + a)^4 a - 4(ia \tan(dx+c) + a)^3 a^2 + 6(ia \tan(dx+c) + a)^2 a^3 - 4(ia \tan(dx+c) + a)a^4 + a^5)}{d} \right)$$

384 d

input `integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/384*a^4*(768*sqrt(2)*(A - I*B)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a)))/a^(3/2) - 9*(121*A - 120*I*B)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a)))/a^(3/2) + 2*(3*(I*a*tan(d*x + c) + a)^(7/2)*(149*A - 152*I*B) - (I*a*tan(d*x + c) + a)^(5/2)*(1127*A - 1160*I*B)*a + (I*a*tan(d*x + c) + a)^(3/2)*(1049*A - 1016*I*B)*a^2 - 3*sqrt(I*a*tan(d*x + c) + a)*(107*A - 104*I*B)*a^3)/((I*a*tan(d*x + c) + a)^4*a - 4*(I*a*tan(d*x + c) + a)^3*a^2 + 6*(I*a*tan(d*x + c) + a)^2*a^3 - 4*(I*a*tan(d*x + c) + a)*a^4 + a^5))/d`

3.89.8 Giac [F]

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{5/2} \cot(dx + c)^5 dx$$

input `integrate(cot(d*x+c)^5*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^5, x)`

3.89.9 Mupad [B] (verification not implemented)

Time = 9.72 (sec) , antiderivative size = 3094, normalized size of antiderivative = 11.85

$$\int \cot^5(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int(cot(c + d*x)^5*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),x)`

output
$$\begin{aligned} &(((107*A*a^6 - B*a^6*104i)*(a + a*\tan(c + d*x)*1i)^{(1/2)})/(64*d) - ((149*A \\ &*a^3 - B*a^3*152i)*(a + a*\tan(c + d*x)*1i)^{(7/2)})/(64*d) - ((1049*A*a^5 - \\ &B*a^5*1016i)*(a + a*\tan(c + d*x)*1i)^{(3/2)})/(192*d) + ((1127*A*a^4 - B*a^4 \\ &*1160i)*(a + a*\tan(c + d*x)*1i)^{(5/2)})/(192*d))/((a + a*\tan(c + d*x)*1i)^4 \\ &- 4*a^3*(a + a*\tan(c + d*x)*1i) - 4*a*(a + a*\tan(c + d*x)*1i)^3 + 6*a^2*(\\ &a + a*\tan(c + d*x)*1i)^2 + a^4) - 2*atanh((384*d^4*(a + a*\tan(c + d*x)*1i) \\ &^{(1/2)}*((485809*A^4*a^22)/(262144*d^4) + (529*B^4*a^22)/(64*d^4) + (11229 \\ &*A^2*B^2*a^22)/(2048*d^4) + (A*B^3*a^22*1127i)/(128*d^4) + (A^3*B*a^22*341 \\ &53i)/(8192*d^4))^{(1/2)})/(64*a^6) + (262841*A^2*a^5)/(32768*d^2) - (4073*B^2 \\ &*a^5)/(512*d^2) - (A*B*a^5*32719i)/(2048*d^2))^{(1/2)}*((485809*A^4*a^22)/(2 \\ &62144*d^4) + (529*B^4*a^22)/(64*d^4) + (11229*A^2*B^2*a^22)/(2048*d^4) + (\\ &A*B^3*a^22*1127i)/(128*d^4) + (A^3*B*a^22*34153i)/(8192*d^4))^{(1/2)})/((431 \\ &443*A^3*a^14*d)/256 - B^3*a^14*d*3542i + (21783*A*B^2*a^14*d)/4 + (A^2*B*a \\ &^14*d*6993i)/32 + 214*A*a^3*d^3*((485809*A^4*a^22)/(262144*d^4) + (529*B^4 \\ &*a^22)/(64*d^4) + (11229*A^2*B^2*a^22)/(2048*d^4) + (A*B^3*a^22*1127i)/(12 \\ &8*d^4) + (A^3*B*a^22*34153i)/(8192*d^4))^{(1/2)} - B*a^3*d^3*((485809*A^4*a^ \\ &22)/(262144*d^4) + (529*B^4*a^22)/(64*d^4) + (11229*A^2*B^2*a^22)/(2048*d^ \\ &4) + (A*B^3*a^22*1127i)/(128*d^4) + (A^3*B*a^22*34153i)/(8192*d^4))^{(1/2)}* \\ &208i) + (697*A^2*a^8*d^2*(a + a*\tan(c + d*x)*1i)^{(1/2)}*((485809*A^4*a^22) \\ &/ (262144*d^4) + (529*B^4*a^22)/(64*d^4) + (11229*A^2*B^2*a^22)/(2048*d^... \end{aligned}$$

3.90
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

3.90.1 Optimal result 1010
 3.90.2 Mathematica [A] (verified) 1011
 3.90.3 Rubi [A] (verified) 1011
 3.90.4 Maple [A] (verified) 1015
 3.90.5 Fricas [B] (verification not implemented) 1016
 3.90.6 Sympy [F] 1016
 3.90.7 Maxima [A] (verification not implemented) 1017
 3.90.8 Giac [F] 1017
 3.90.9 Mupad [B] (verification not implemented) 1018

3.90.1 Optimal result

Integrand size = 36, antiderivative size = 205

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}} + \frac{(iA-B)\tan^3(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} + \frac{4(5A+7iB)\sqrt{a+ia \tan(c+dx)}}{5ad} - \frac{(5A+7iB)\tan^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{5ad} - \frac{(25A+23iB)(a+ia \tan(c+dx))^{3/2}}{15a^2d}$$

```
output 1/2*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/d*2^(1/2)
)/a^(1/2)+4/5*(5*A+7*I*B)*(a+I*a*tan(d*x+c))^(1/2)/a/d-1/5*(5*A+7*I*B)*(a+
I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^2/a/d+(I*A-B)*tan(d*x+c)^3/d/(a+I*a*tan(d
*x+c))^(1/2)-1/15*(25*A+23*I*B)*(a+I*a*tan(d*x+c))^(3/2)/a^2/d
```

3.90.2 Mathematica [A] (verified)

Time = 2.43 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.63

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}} + \frac{35A+61iB+(10iA-38B) \tan(c+dx)+2(5A+iB) \tan^2(c+dx)+6B \tan^3(c+dx)}{15d\sqrt{a+ia \tan(c+dx)}}$$

input `Integrate[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `((A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/(Sqrt[2]*Sqrt[a]*d) + (35*A + (61*I)*B + ((10*I)*A - 38*B)*Tan[c + d*x] + 2*(5*A + I*B)*Tan[c + d*x]^2 + 6*B*Tan[c + d*x]^3)/(15*d*Sqrt[a + I*a*Tan[c + d*x]])`

3.90.3 Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.06, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.361$, Rules used = {3042, 4078, 27, 3042, 4080, 27, 3042, 4075, 3042, 4010, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(c+dx)^3(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx \\ & \quad \downarrow \text{4078} \\ & \frac{(-B+iA) \tan^3(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \\ & \frac{\int \frac{1}{2} \tan^2(c+dx) \sqrt{i \tan(c+dx)a + a(6a(iA-B) + a(5A+7iB) \tan(c+dx))} dx}{a^2} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.90. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

$$\begin{aligned}
& \frac{(-B + iA) \tan^3(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\int \tan^2(c + dx) \sqrt{i \tan(c + dx)a + a(6a(iA - B) + a(5A + 7iB) \tan(c + dx))} dx}{2a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(-B + iA) \tan^3(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\int \tan(c + dx)^2 \sqrt{i \tan(c + dx)a + a(6a(iA - B) + a(5A + 7iB) \tan(c + dx))} dx}{2a^2} \\
& \quad \downarrow \text{4080} \\
& \frac{(-B + iA) \tan^3(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{2 \int -\frac{1}{2} \tan(c + dx) \sqrt{i \tan(c + dx)a + a(4a^2(5A + 7iB) - a^2(25iA - 23B) \tan(c + dx))} dx}{5a} + \frac{2a(5A + 7iB) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d}}{2a^2} \\
& \quad \downarrow \text{27} \\
& \frac{(-B + iA) \tan^3(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\frac{2a(5A + 7iB) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \int \tan(c + dx) \sqrt{i \tan(c + dx)a + a(4a^2(5A + 7iB) - a^2(25iA - 23B) \tan(c + dx))} dx}{5a}}{2a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(-B + iA) \tan^3(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\frac{2a(5A + 7iB) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \int \tan(c + dx) \sqrt{i \tan(c + dx)a + a(4a^2(5A + 7iB) - a^2(25iA - 23B) \tan(c + dx))} dx}{5a}}{2a^2} \\
& \quad \downarrow \text{4075} \\
& \frac{(-B + iA) \tan^3(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\frac{2a(5A + 7iB) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \int \sqrt{i \tan(c + dx)a + a((25iA - 23B)a^2 + 4(5A + 7iB) \tan(c + dx)a^2)} dx - \frac{2a(25A + 23iB)(a + ia \tan(c + dx))}{3d}}{5a}}{2a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(-B + iA) \tan^3(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\frac{2a(5A + 7iB) \tan^2(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d} - \int \sqrt{i \tan(c + dx)a + a((25iA - 23B)a^2 + 4(5A + 7iB) \tan(c + dx)a^2)} dx - \frac{2a(25A + 23iB)(a + ia \tan(c + dx))}{3d}}{5a}}{2a^2} \\
& \quad \downarrow \text{4010}
\end{aligned}$$

3.90. $\int \frac{\tan^3(c + dx)(A + B \tan(c + dx))}{\sqrt{a + ia \tan(c + dx)}} dx$

$$\frac{\frac{(-B + iA) \tan^3(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2a(5A+7iB) \tan^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{5d} - \frac{5a^2(B+iA) \int \sqrt{i \tan(c+dx)a+adx}}{5a^2} + \frac{8a^2(5A+7iB)\sqrt{a+ia \tan(c+dx)}}{5a} - \frac{2a(25A+23iB)(a+ia \tan(c+dx))^3}{3d}}{2a^2}$$

↓ 3042

$$\frac{\frac{(-B + iA) \tan^3(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2a(5A+7iB) \tan^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{5d} - \frac{5a^2(B+iA) \int \sqrt{i \tan(c+dx)a+adx}}{5a^2} + \frac{8a^2(5A+7iB)\sqrt{a+ia \tan(c+dx)}}{5a} - \frac{2a(25A+23iB)(a+ia \tan(c+dx))^3}{3d}}{2a^2}$$

↓ 3961

$$\frac{\frac{(-B + iA) \tan^3(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2a(5A+7iB) \tan^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{5d} - \frac{10ia^3(B+iA) \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a}}{2a^2} + \frac{8a^2(5A+7iB)\sqrt{a+ia \tan(c+dx)}}{5a} - \frac{2a(25A+23iB)(a+ia \tan(c+dx))^3}{3d}}{2a^2}$$

↓ 219

$$\frac{\frac{(-B + iA) \tan^3(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2a(5A+7iB) \tan^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{5d} - \frac{5i\sqrt{2}a^{5/2}(B+iA)\arctanh\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{8a^2(5A+7iB)\sqrt{a+ia \tan(c+dx)}}{5a} - \frac{2a(25A+23iB)(a+ia \tan(c+dx))^3}{3d}}{2a^2}$$

input `Int[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `((I*A - B)*Tan[c + d*x]^3)/(d*Sqrt[a + I*a*Tan[c + d*x]]) - ((2*a*(5*A + (7*I)*B)*Tan[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]])/(5*d) - (((-5*I)*Sqrt[2]*a^(5/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d + (8*a^2*(5*A + (7*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/d - (2*a*(25*A + (23*I)*B)*(a + I*a*Tan[c + d*x])^(3/2))/(3*d))/(5*a))/(2*a^2)`

3.90. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

3.90.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_)] /; FreeQ[b, x]`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3961 `Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`
- rule 4010 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`
- rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`
- rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

```
rule 4080 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[
1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Sim
p[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*T
an[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c
- a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]
```

3.90.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.81

method	result
derivativedivides	$\frac{\frac{2iB(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} - \frac{4iBa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - \frac{2Aa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + 4ia^2 B \sqrt{a+ia \tan(dx+c)} + 2A a^2 \sqrt{a+ia \tan(dx+c)}}{a^3 d}$
default	$\frac{\frac{2iB(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} - \frac{4iBa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - \frac{2Aa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + 4ia^2 B \sqrt{a+ia \tan(dx+c)} + 2A a^2 \sqrt{a+ia \tan(dx+c)}}{a^3 d}$
parts	$\frac{2A \left(-\frac{(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + a \sqrt{a+ia \tan(dx+c)} + \frac{a^{\frac{3}{2}} \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}}\right)}{4} + \frac{a^2}{2\sqrt{a+ia \tan(dx+c)}} \right)}{d a^2} + \frac{2iB \left(\frac{(a+ia \tan(dx+c))^{\frac{5}{2}}}{5} - \frac{4iBa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - \frac{2Aa(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + 4ia^2 B \sqrt{a+ia \tan(dx+c)} + 2A a^2 \sqrt{a+ia \tan(dx+c)} \right)}{a^3 d}$

```
input int(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x,method=_RETUR
NVERBOSE)
```

```
output 2/d/a^3*(1/5*I*B*(a+I*a*tan(d*x+c))^(5/2)-2/3*I*B*a*(a+I*a*tan(d*x+c))^(3/
2)-1/3*A*a*(a+I*a*tan(d*x+c))^(3/2)+2*I*a^2*B*(a+I*a*tan(d*x+c))^(1/2)+A*a
^2*(a+I*a*tan(d*x+c))^(1/2)+1/4*a^(5/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a
*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))+1/2*a^3*(A+I*B)/(a+I*a*tan(d*x+c))^(1/
2))
```

3.90.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 447 vs. $2(163) = 326$.

Time = 0.28 (sec) , antiderivative size = 447, normalized size of antiderivative = 2.18

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx =$$

$$15\sqrt{2}(ade^{(5i dx+5i c)} + 2ade^{(3i dx+3i c)} + ade^{(i dx+i c)})\sqrt{\frac{A^2-2iAB-B^2}{ad^2}} \log\left(-\frac{4\left((-iA-B)ae^{(i dx+i c)}+(iade^{(2i dx+2i c)}\right)}{\dots}\right)$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="fricas")
```

```
output -1/60*(15*sqrt(2)*(a*d*e^(5*I*d*x + 5*I*c) + 2*a*d*e^(3*I*d*x + 3*I*c) + a*d*e^(I*d*x + I*c))*sqrt((A^2 - 2*I*A*B - B^2)/(a*d^2))*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) + (I*a*d*e^(2*I*d*x + 2*I*c) + I*a*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a*d^2)))*e^(-I*d*x - I*c)/(I*A + B)) - 15*sqrt(2)*(a*d*e^(5*I*d*x + 5*I*c) + 2*a*d*e^(3*I*d*x + 3*I*c) + a*d*e^(I*d*x + I*c))*sqrt((A^2 - 2*I*A*B - B^2)/(a*d^2))*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) + (-I*a*d*e^(2*I*d*x + 2*I*c) - I*a*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a*d^2)))*e^(-I*d*x - I*c)/(I*A + B)) - 2*sqrt(2)*((35*A + 103*I*B)*e^(6*I*d*x + 6*I*c) + 5*(25*A + 41*I*B)*e^(4*I*d*x + 4*I*c) + 15*(7*A + 11*I*B)*e^(2*I*d*x + 2*I*c) + 15*A + 15*I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(a*d*e^(5*I*d*x + 5*I*c) + 2*a*d*e^(3*I*d*x + 3*I*c) + a*d*e^(I*d*x + I*c))
```

3.90.6 SymPy [F]

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{(A+B\tan(c+dx))\tan^3(c+dx)}{\sqrt{ia(\tan(c+dx)-i)}} dx$$

```
input integrate(tan(d*x+c)**3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/2),x)
```

```
output Integral((A + B*tan(c + d*x))*tan(c + d*x)**3/sqrt(I*a*(tan(c + d*x) - I)), x)
```

3.90. $\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx$

3.90.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.77

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \frac{15\sqrt{2}(A-iB)a^{\frac{7}{2}}\log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia\tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia\tan(dx+c)+a}}\right)-24i(ia\tan(dx+c)+a)^{\frac{5}{2}}Ba+40(ia\tan(dx+c)+a)^{\frac{3}{2}}(A+2iB)a^2-120\sqrt{2}(ia\tan(dx+c)+a)(A+2iB)a^3-60(A+iB)a^4}{60a^4d}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="maxima")
```

```
output -1/60*(15*sqrt(2)*(A - I*B)*a^(7/2)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - 24*I*(I*a*tan(d*x + c) + a)^(5/2)*B*a + 40*(I*a*tan(d*x + c) + a)^(3/2)*(A + 2*I*B)*a^2 - 120*sqrt(I*a*tan(d*x + c) + a)*(A + 2*I*B)*a^3 - 60*(A + I*B)*a^4/sqrt(I*a*tan(d*x + c) + a))/(a^4*d)
```

3.90.8 Giac [F]

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^3}{\sqrt{ia\tan(dx+c)+a}} dx$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")
```

```
output integrate((B*tan(d*x + c) + A)*tan(d*x + c)^3/sqrt(I*a*tan(d*x + c) + a), x)
```


3.90.9 Mupad [B] (verification not implemented)

Time = 8.70 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.15

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \frac{A}{d\sqrt{a+a\tan(c+dx)} \operatorname{li}} + \frac{B \operatorname{li}}{d\sqrt{a+a\tan(c+dx)} \operatorname{li}} + \frac{2A\sqrt{a+a\tan(c+dx)} \operatorname{li}}{ad} - \frac{2A(a+a\tan(c+dx) \operatorname{li})^{3/2}}{3a^2d} + \frac{B\sqrt{a+a\tan(c+dx)} \operatorname{li} 4i}{ad} - \frac{B(a+a\tan(c+dx) \operatorname{li})^{3/2} 4i}{3a^2d} + \frac{B(a+a\tan(c+dx) \operatorname{li})^{5/2} 2i}{5a^3d} + \frac{\sqrt{2} B \operatorname{atan}\left(\frac{\sqrt{2}\sqrt{a+a\tan(c+dx)} \operatorname{li}}{2\sqrt{-a}}\right) \operatorname{li}}{2\sqrt{-a}d} + \frac{\sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2}\sqrt{a+a\tan(c+dx)} \operatorname{li} \operatorname{li}}{2\sqrt{a}}\right) \operatorname{li}}{2\sqrt{a}d}$$

input `int((tan(c + d*x)^3*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2),x)`output `A/(d*(a + a*tan(c + d*x)*1i)^(1/2)) + (B*1i)/(d*(a + a*tan(c + d*x)*1i)^(1/2)) + (2*A*(a + a*tan(c + d*x)*1i)^(1/2))/(a*d) - (2*A*(a + a*tan(c + d*x)*1i)^(3/2))/(3*a^2*d) + (B*(a + a*tan(c + d*x)*1i)^(1/2)*4i)/(a*d) - (B*(a + a*tan(c + d*x)*1i)^(3/2)*4i)/(3*a^2*d) + (B*(a + a*tan(c + d*x)*1i)^(5/2)*2i)/(5*a^3*d) + (2^(1/2)*B*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)))/(2*(-a)^(1/2)))*1i)/(2*(-a)^(1/2)*d) - (2^(1/2)*A*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)*1i)/(2*a^(1/2)))*1i)/(2*a^(1/2)*d)`

3.91 $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

3.91.1	Optimal result	1019
3.91.2	Mathematica [A] (verified)	1020
3.91.3	Rubi [A] (verified)	1020
3.91.4	Maple [A] (verified)	1023
3.91.5	Fricas [B] (verification not implemented)	1024
3.91.6	Sympy [F]	1024
3.91.7	Maxima [A] (verification not implemented)	1025
3.91.8	Giac [F]	1025
3.91.9	Mupad [B] (verification not implemented)	1026

3.91.1 Optimal result

Integrand size = 36, antiderivative size = 159

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \frac{(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}} + \frac{(iA-B)\tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{4(iA-B)\sqrt{a+ia \tan(c+dx)}}{ad} + \frac{(3iA-5B)(a+ia \tan(c+dx))^{3/2}}{3a^2d}$$

output $\frac{1}{2}(iA+B)\operatorname{arctanh}\left(\frac{1}{2}(a+i a \tan(dx+c))^{1/2} 2^{1/2}/a^{1/2}\right)/d 2^{1/2}/a^{1/2}-4(iA-B)(a+i a \tan(dx+c))^{1/2}/a d+(iA-B)\tan^2(dx+c)/d/(a+i a \tan(dx+c))^{1/2}+1/3(3iA-5B)(a+i a \tan(dx+c))^{3/2}/a^2/d$

3.91.2 Mathematica [A] (verified)

Time = 1.87 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.70

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx$$

$$= \frac{(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}} + \frac{-9iA+7B+(6A+2iB)\tan(c+dx)+2B\tan^2(c+dx)}{3d\sqrt{a+ia\tan(c+dx)}}$$

input `Integrate[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `((I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/(Sqrt[2]*Sqrt[a]*d) + ((-9*I)*A + 7*B + (6*A + (2*I)*B)*Tan[c + d*x] + 2*B*Tan[c + d*x]^2)/(3*d*Sqrt[a + I*a*Tan[c + d*x]])`

3.91.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.04, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {3042, 4078, 27, 3042, 4075, 3042, 4010, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)^2(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx$$

$$\downarrow \text{4078}$$

$$\frac{(-B+iA)\tan^2(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} - \frac{\int \frac{1}{2}\tan(c+dx)\sqrt{i\tan(c+dx)a+a(4a(iA-B)+a(3A+5iB)\tan(c+dx))}dx}{a^2}$$

3.91. $\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{(-B + iA) \tan^2(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\int \tan(c + dx) \sqrt{i \tan(c + dx) a + a(4a(iA - B) + a(3A + 5iB) \tan(c + dx))} dx}{2a^2} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^2(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\int \tan(c + dx) \sqrt{i \tan(c + dx) a + a(4a(iA - B) + a(3A + 5iB) \tan(c + dx))} dx}{2a^2} \\
& \downarrow 4075 \\
& \frac{(-B + iA) \tan^2(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\int \sqrt{i \tan(c + dx) a + a(4a(iA - B) \tan(c + dx) - a(3A + 5iB))} dx - \frac{2(-5B + 3iA)(a + ia \tan(c + dx))^{3/2}}{3d}}{2a^2} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^2(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\int \sqrt{i \tan(c + dx) a + a(4a(iA - B) \tan(c + dx) - a(3A + 5iB))} dx - \frac{2(-5B + 3iA)(a + ia \tan(c + dx))^{3/2}}{3d}}{2a^2} \\
& \downarrow 4010 \\
& \frac{(-B + iA) \tan^2(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{a(A - iB) \int \sqrt{i \tan(c + dx) a + a} dx - \frac{2(-5B + 3iA)(a + ia \tan(c + dx))^{3/2}}{3d} + \frac{8a(-B + iA) \sqrt{a + ia \tan(c + dx)}}{d}}{2a^2} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^2(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{a(A - iB) \int \sqrt{i \tan(c + dx) a + a} dx - \frac{2(-5B + 3iA)(a + ia \tan(c + dx))^{3/2}}{3d} + \frac{8a(-B + iA) \sqrt{a + ia \tan(c + dx)}}{d}}{2a^2} \\
& \downarrow 3961 \\
& \frac{(-B + iA) \tan^2(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{2ia^2(A - iB) \int \frac{1}{a - ia \tan(c + dx)} d\sqrt{i \tan(c + dx) a + a} + \frac{8a(-B + iA) \sqrt{a + ia \tan(c + dx)}}{d} - \frac{2(-5B + 3iA)(a + ia \tan(c + dx))^{3/2}}{3d}}{2a^2}
\end{aligned}$$

3.91. $\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{\sqrt{a + ia \tan(c + dx)}} dx$

$$\begin{aligned} & \downarrow 219 \\ & \frac{(-B + iA) \tan^2(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\ & \frac{i\sqrt{2}a^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{8a(-B + iA)\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2(-5B + 3iA)(a + ia \tan(c + dx))^{3/2}}{3d} \\ & \hline & 2a^2 \end{aligned}$$

input `Int[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `((I*A - B)*Tan[c + d*x]^2)/(d*Sqrt[a + I*a*Tan[c + d*x]]) - (((-I)*Sqrt[2]*a^(3/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d + (8*a*(I*A - B)*Sqrt[a + I*a*Tan[c + d*x]])/d - (2*((3*I)*A - 5*B)*(a + I*a*Tan[c + d*x])^(3/2))/(3*d))/(2*a^2)`

3.91.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

```
rule 4078 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.91.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.81

method	result
derivativedivides	$2i \left(\frac{iB(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - iaB\sqrt{a+ia \tan(dx+c)} - aA\sqrt{a+ia \tan(dx+c)} + \frac{a^{\frac{3}{2}}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{4} \right) \frac{1}{da^2}$
default	$2i \left(\frac{iB(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - iaB\sqrt{a+ia \tan(dx+c)} - aA\sqrt{a+ia \tan(dx+c)} + \frac{a^{\frac{3}{2}}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{4} \right) \frac{1}{da^2}$
parts	$2iA \left(-\sqrt{a+ia \tan(dx+c)} + \frac{\sqrt{a}\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{4} - \frac{a}{2\sqrt{a+ia \tan(dx+c)}} \right) \frac{1}{da} + 2B \left(-\frac{(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + a \right) \frac{1}{da}$

```
input int(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x,method=_RETUR
NVERBOSE)
```

```
output 2*I/d/a^2*(1/3*I*B*(a+I*a*tan(d*x+c))^(3/2)-I*a*B*(a+I*a*tan(d*x+c))^(1/2)
-a*A*(a+I*a*tan(d*x+c))^(1/2)+1/4*a^(3/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I
*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))-1/2*a^2*(A+I*B)/(a+I*a*tan(d*x+c))^(
1/2))
```

$$3.91. \int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

3.91.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 391 vs. $2(127) = 254$.

Time = 0.27 (sec) , antiderivative size = 391, normalized size of antiderivative = 2.46

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx =$$

$$3\sqrt{2}\left(ade^{(3i dx+3i c)} + ade^{(i dx+i c)}\right)\sqrt{-\frac{A^2-2iAB-B^2}{ad^2}} \log\left(-\frac{4\left((-iA-B)ae^{(i dx+i c)}+(ade^{(2i dx+2i c)}+ad)\sqrt{\frac{a}{e^{(2i dx+2i c)}+iA+B}}\right)}{iA+B}\right)$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="fricas")`

output `-1/12*(3*sqrt(2)*(a*d*e^(3*I*d*x + 3*I*c) + a*d*e^(I*d*x + I*c))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a*d^2))*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) + (a*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a*d^2)))*e^(-I*d*x - I*c)/(I*A + B)) - 3*sqrt(2)*(a*d*e^(3*I*d*x + 3*I*c) + a*d*e^(I*d*x + I*c))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a*d^2))*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) - (a*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a*d^2)))*e^(-I*d*x - I*c)/(I*A + B)) + 2*sqrt(2)*((15*I*A - 7*B)*e^(4*I*d*x + 4*I*c) + 18*(I*A - B)*e^(2*I*d*x + 2*I*c) + 3*I*A - 3*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(a*d*e^(3*I*d*x + 3*I*c) + a*d*e^(I*d*x + I*c))`

3.91.6 Sympy [F]

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{(A+B\tan(c+dx))\tan^2(c+dx)}{\sqrt{ia(\tan(c+dx)-i)}} dx$$

input `integrate(tan(d*x+c)**2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**2/sqrt(I*a*(tan(c + d*x) - I)), x)`

3.91. $\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx$

3.91.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.84

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \frac{i\left(3\sqrt{2}(A-iB)a^{\frac{5}{2}}\log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia\tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia\tan(dx+c)+a}}\right)-8i\left(ia\tan(dx+c)+a\right)^{\frac{3}{2}}Ba+24\sqrt{ia\tan(dx+c)}\right)}{12a^3d}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `-1/12*I*(3*sqrt(2)*(A - I*B)*a^(5/2)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - 8*I*(I*a*tan(d*x + c) + a)^(3/2)*B*a + 24*sqrt(I*a*tan(d*x + c) + a)*(A + I*B)*a^2 + 12*(A + I*B)*a^3/sqrt(I*a*tan(d*x + c) + a))/(a^3*d)`

3.91.8 Giac [F]

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^2}{\sqrt{ia\tan(dx+c)+a}} dx$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^2/sqrt(I*a*tan(d*x + c) + a), x)`

3.91.9 Mupad [B] (verification not implemented)

Time = 8.18 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.18

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = -\frac{A \operatorname{li}}{d \sqrt{a+a\tan(c+dx)} \operatorname{li}} + \frac{B}{d \sqrt{a+a\tan(c+dx)} \operatorname{li}} - \frac{A \sqrt{a+a\tan(c+dx)} \operatorname{li} 2i}{ad} + \frac{2B \sqrt{a+a\tan(c+dx)} \operatorname{li}}{ad} - \frac{2B(a+a\tan(c+dx) \operatorname{li})^{3/2}}{3a^2 d} - \frac{\sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2}\sqrt{a+a\tan(c+dx)} \operatorname{li}}{2\sqrt{-a}}\right) \operatorname{li}}{2\sqrt{-a} d} - \frac{\sqrt{2} B \operatorname{atan}\left(\frac{\sqrt{2}\sqrt{a+a\tan(c+dx)} \operatorname{li} \operatorname{li}}{2\sqrt{a}}\right) \operatorname{li}}{2\sqrt{a} d}$$

input `int((tan(c + d*x))^2*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2),x)`

output `B/(d*(a + a*tan(c + d*x)*1i)^(1/2)) - (A*1i)/(d*(a + a*tan(c + d*x)*1i)^(1/2)) - (A*(a + a*tan(c + d*x)*1i)^(1/2)*2i)/(a*d) + (2*B*(a + a*tan(c + d*x)*1i)^(1/2))/(a*d) - (2*B*(a + a*tan(c + d*x)*1i)^(3/2))/(3*a^2*d) - (2^(1/2)*A*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2))))*1i)/(2*(-a)^(1/2)*d) - (2^(1/2)*B*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)*1i)/(2*a^(1/2))))*1i)/(2*a^(1/2)*d)`

$$3.92 \quad \int \frac{\tan(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

3.92.1 Optimal result 1027
 3.92.2 Mathematica [A] (verified) 1027
 3.92.3 Rubi [A] (verified) 1028
 3.92.4 Maple [A] (verified) 1030
 3.92.5 Fricas [B] (verification not implemented) 1030
 3.92.6 Sympy [F] 1031
 3.92.7 Maxima [A] (verification not implemented) 1031
 3.92.8 Giac [F] 1032
 3.92.9 Mupad [B] (verification not implemented) 1032

3.92.1 Optimal result

Integrand size = 34, antiderivative size = 109

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = -\frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}} - \frac{A+iB}{d\sqrt{a+ia \tan(c+dx)}} - \frac{2iB\sqrt{a+ia \tan(c+dx)}}{ad}$$

```
output -1/2*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/d*2^(1/2)/a^(1/2)+(-A-I*B)/d/(a+I*a*tan(d*x+c))^(1/2)-2*I*B*(a+I*a*tan(d*x+c))^(1/2)/a/d
```

3.92.2 Mathematica [A] (verified)

Time = 1.36 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.00

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = -\frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}} - \frac{A+iB}{d\sqrt{a+ia \tan(c+dx)}} - \frac{2iB\sqrt{a+ia \tan(c+dx)}}{ad}$$

input `Integrate[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x
]`

output `-(((A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/(Sqrt[
2]*Sqrt[a]*d)) - (A + I*B)/(d*Sqrt[a + I*a*Tan[c + d*x]]) - ((2*I)*B*Sqrt[
a + I*a*Tan[c + d*x]])/(a*d)`

3.92.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.02, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.206$, Rules used = {3042, 4075, 3042, 4009, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx \\
 & \quad \downarrow \text{4075} \\
 & \int \frac{A\tan(c+dx)-B}{\sqrt{i\tan(c+dx)a+a}} dx - \frac{2iB\sqrt{a+ia\tan(c+dx)}}{ad} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A\tan(c+dx)-B}{\sqrt{i\tan(c+dx)a+a}} dx - \frac{2iB\sqrt{a+ia\tan(c+dx)}}{ad} \\
 & \quad \downarrow \text{4009} \\
 & -\frac{(B+iA)\int\sqrt{i\tan(c+dx)a+adx}}{2a} - \frac{A+iB}{d\sqrt{a+ia\tan(c+dx)}} - \frac{2iB\sqrt{a+ia\tan(c+dx)}}{ad} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{(B+iA)\int\sqrt{i\tan(c+dx)a+adx}}{2a} - \frac{A+iB}{d\sqrt{a+ia\tan(c+dx)}} - \frac{2iB\sqrt{a+ia\tan(c+dx)}}{ad} \\
 & \quad \downarrow \text{3961}
 \end{aligned}$$

3.92. $\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx$

$$\frac{i(B + iA) \int \frac{1}{a - ia \tan(c + dx)} d\sqrt{i \tan(c + dx)a + a}}{d} - \frac{A + iB}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2iB\sqrt{a + ia \tan(c + dx)}}{ad}$$

↓ 219

$$\frac{i(B + iA) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}} - \frac{A + iB}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2iB\sqrt{a + ia \tan(c + dx)}}{ad}$$

input `Int[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `(I*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/(Sqrt[2]*Sqrt[a]*d) - (A + I*B)/(d*Sqrt[a + I*a*Tan[c + d*x]]) - ((2*I)*B*Sqrt[a + I*a*Tan[c + d*x]])/(a*d)`

3.92.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

3.92.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.81

method	result
derivativedivides	$\frac{-2iB\sqrt{a+ia \tan(dx+c)} - \frac{a(iB+A)}{\sqrt{a+ia \tan(dx+c)}} - \frac{\sqrt{a}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2}}{ad}$
default	$\frac{-2iB\sqrt{a+ia \tan(dx+c)} - \frac{a(iB+A)}{\sqrt{a+ia \tan(dx+c)}} - \frac{\sqrt{a}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2}}{ad}$
parts	$\frac{A\left(-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2\sqrt{a}} - \frac{1}{\sqrt{a+ia \tan(dx+c)}}\right)}{d} + \frac{2iB\left(-\sqrt{a+ia \tan(dx+c)} + \frac{\sqrt{a}\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{4}\right)}{da}$

```
input int(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output 2/d/a*(-I*B*(a+I*a*tan(d*x+c))^(1/2)-1/2*a*(A+I*B)/(a+I*a*tan(d*x+c))^(1/2
)-1/4*a^(1/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)
/a^(1/2)))
```

3.92.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 327 vs. 2(84) = 168.

Time = 0.25 (sec) , antiderivative size = 327, normalized size of antiderivative = 3.00

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

$$= \left(\sqrt{2}ad\sqrt{\frac{A^2-2iAB-B^2}{ad^2}} e^{(i dx+i c)} \log \left(-\frac{4\left((-iA-B)ae^{(i dx+i c)}+(iade^{(2i dx+2i c)}+iad)\sqrt{\frac{a}{e^{(2i dx+2i c)}+1}}\sqrt{\frac{A^2-2iAB-B^2}{ad^2}}\right)}{iA+B} \right) e^{(-i dx-i c)} \right)$$

3.92. $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="fricas")`

output `1/4*(sqrt(2)*a*d*sqrt((A^2 - 2*I*A*B - B^2)/(a*d^2))*e^(I*d*x + I*c)*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) + (I*a*d*e^(2*I*d*x + 2*I*c) + I*a*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a*d^2)))*e^(-I*d*x - I*c)/(I*A + B) - sqrt(2)*a*d*sqrt((A^2 - 2*I*A*B - B^2)/(a*d^2))*e^(I*d*x + I*c)*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) + (-I*a*d*e^(2*I*d*x + 2*I*c) - I*a*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a*d^2)))*e^(-I*d*x - I*c)/(I*A + B) - 2*sqrt(2)*((A + 5*I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/(a*d)`

3.92.6 Sympy [F]

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{(A+B\tan(c+dx))\tan(c+dx)}{\sqrt{ia(\tan(c+dx)-i)}} dx$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)/sqrt(I*a*(tan(c + d*x) - I)), x)`

3.92.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.01

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \frac{\sqrt{2}(A-iB)a^{\frac{3}{2}} \log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia\tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia\tan(dx+c)+a}}\right) - 8i\sqrt{ia\tan(dx+c)+a}Ba - \frac{4(A+iB)a^2}{\sqrt{ia\tan(dx+c)+a}}}{4a^2d}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output $1/4*(\text{sqrt}(2)*(A - I*B)*a^{(3/2)}*\log(-(\text{sqrt}(2)*\text{sqrt}(a) - \text{sqrt}(I*a*\tan(dx + c) + a))/(\text{sqrt}(2)*\text{sqrt}(a) + \text{sqrt}(I*a*\tan(dx + c) + a))) - 8*I*\text{sqrt}(I*a*\tan(dx + c) + a)*B*a - 4*(A + I*B)*a^2/\text{sqrt}(I*a*\tan(dx + c) + a)/(a^2*d)$

3.92.8 Giac [F]

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{\sqrt{a + ia \tan(c + dx)}} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)}{\sqrt{ia \tan(dx + c) + a}} dx$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)/sqrt(I*a*tan(d*x + c) + a), x)`

3.92.9 Mupad [B] (verification not implemented)

Time = 1.07 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.29

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{\sqrt{a + ia \tan(c + dx)}} dx = -\frac{A}{d \sqrt{a + a \tan(c + dx)} \operatorname{li}} - \frac{B \operatorname{li}}{d \sqrt{a + a \tan(c + dx)} \operatorname{li}} - \frac{B \sqrt{a + a \tan(c + dx)} \operatorname{li}^2}{B \sqrt{a + a \tan(c + dx)} \operatorname{li}^2} - \frac{a d}{\sqrt{2} B \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx)} \operatorname{li}}{2 \sqrt{-a}}\right) \operatorname{li}} - \frac{2 \sqrt{-a} d}{\sqrt{2} A \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx)} \operatorname{li}}{2 \sqrt{a}}\right)} - \frac{\sqrt{2} A \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx)} \operatorname{li}}{2 \sqrt{a}}\right)}{2 \sqrt{a} d}$$

input `int((tan(c + d*x)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2),x)`

output $-A/(d*(a + a*\tan(c + d*x)*1i)^{(1/2)}) - (B*1i)/(d*(a + a*\tan(c + d*x)*1i)^{(1/2)}) - (B*(a + a*\tan(c + d*x)*1i)^{(1/2)*2i}/(a*d) - (2^{(1/2)}*B*\operatorname{atan}((2^{(1/2)}*(a + a*\tan(c + d*x)*1i)^{(1/2)})/(2*(-a)^{(1/2)})))*1i)/(2*(-a)^{(1/2)}*d) - (2^{(1/2)}*A*\operatorname{atanh}((2^{(1/2)}*(a + a*\tan(c + d*x)*1i)^{(1/2)})/(2*a^{(1/2)})))/(2*a^{(1/2)}*d)$

3.93 $\int \frac{A+B \tan(c+dx)}{\sqrt{a+ia \tan(c+dx)}} dx$

3.93.1	Optimal result	1033
3.93.2	Mathematica [A] (verified)	1033
3.93.3	Rubi [A] (verified)	1034
3.93.4	Maple [A] (verified)	1035
3.93.5	Fricas [B] (verification not implemented)	1036
3.93.6	Sympy [F]	1036
3.93.7	Maxima [A] (verification not implemented)	1037
3.93.8	Giac [F]	1037
3.93.9	Mupad [B] (verification not implemented)	1037

3.93.1 Optimal result

Integrand size = 28, antiderivative size = 82

$$\int \frac{A + B \tan(c + dx)}{\sqrt{a + ia \tan(c + dx)}} dx = -\frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}} + \frac{iA - B}{d\sqrt{a + ia \tan(c + dx)}}$$

```
output -1/2*(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/d*2^(1/2)/a^(1/2)+(I*A-B)/d/(a+I*a*tan(d*x+c))^(1/2)
```

3.93.2 Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00

$$\int \frac{A + B \tan(c + dx)}{\sqrt{a + ia \tan(c + dx)}} dx = -\frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}} + \frac{iA - B}{d\sqrt{a + ia \tan(c + dx)}}$$

```
input Integrate[(A + B*Tan[c + d*x])/Sqrt[a + I*a*Tan[c + d*x]],x]
```

```
output -(((I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/(Sqrt[2]*Sqrt[a]*d)) + (I*A - B)/(d*Sqrt[a + I*a*Tan[c + d*x]])
```


3.93.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.02, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$, Rules used = {3042, 4009, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\sqrt{a + ia \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\sqrt{a + ia \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4009} \\
 & \frac{(A - iB) \int \sqrt{i \tan(c + dx)a + adx}}{2a} + \frac{-B + iA}{d\sqrt{a + ia \tan(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A - iB) \int \sqrt{i \tan(c + dx)a + adx}}{2a} + \frac{-B + iA}{d\sqrt{a + ia \tan(c + dx)}} \\
 & \quad \downarrow \text{3961} \\
 & \frac{-B + iA}{d\sqrt{a + ia \tan(c + dx)}} - \frac{i(A - iB) \int \frac{1}{a - ia \tan(c + dx)} d\sqrt{i \tan(c + dx)a + a}}{d} \\
 & \quad \downarrow \text{219} \\
 & \frac{-B + iA}{d\sqrt{a + ia \tan(c + dx)}} - \frac{i(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}}
 \end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `((-I)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/(Sqrt[2]*Sqrt[a]*d) + (I*A - B)/(d*Sqrt[a + I*a*Tan[c + d*x]])`

3.93.3.1 Defintions of rubi rules used

- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

- rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

- rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

3.93.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.87

method	result
derivativedivides	$\frac{2i \left(-\frac{\left(\frac{A}{2} - \frac{iB}{2}\right)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right) - \frac{-\frac{A}{2} - \frac{iB}{2}}{\sqrt{a+ia \tan(dx+c)}}}{d} \right)}{d}$
default	$\frac{2i \left(-\frac{\left(\frac{A}{2} - \frac{iB}{2}\right)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right) - \frac{-\frac{A}{2} - \frac{iB}{2}}{\sqrt{a+ia \tan(dx+c)}}}{d} \right)}{d}$
parts	$\frac{2iAa \left(-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{4a^{\frac{3}{2}}} + \frac{1}{2a\sqrt{a+ia \tan(dx+c)}} \right)}{d} + \frac{B \left(-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right) - \frac{1}{\sqrt{a+ia \tan(dx+c)}}}{d} \right)}{d}$

```
input int((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x,method=_RETURNVERBOSE)
```

3.93. $\int \frac{A+B \tan(c+dx)}{\sqrt{a+ia \tan(c+dx)}} dx$

output $2*I/d*(-1/2*(1/2*A-1/2*I*B)*2^(1/2)/a^(1/2)*\operatorname{arctanh}(1/2*(a+I*a*\tan(dx+c))$
 $^(1/2)*2^(1/2)/a^(1/2))-(-1/2*A-1/2*I*B)/(a+I*a*\tan(dx+c))^(1/2)$

3.93.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 328 vs. $2(63) = 126$.

Time = 0.26 (sec) , antiderivative size = 328, normalized size of antiderivative = 4.00

$$\int \frac{A + B \tan(c + dx)}{\sqrt{a + ia \tan(c + dx)}} dx$$

$$= \left(\sqrt{2}ad\sqrt{-\frac{A^2-2iAB-B^2}{ad^2}} e^{(idx+ic)} \log \left(-\frac{4 \left((-iA-B)ae^{(idx+ic)} + (ade^{(2idx+2ic)} + ad) \sqrt{\frac{a}{e^{(2idx+2ic)}+1}} \sqrt{-\frac{A^2-2iAB-B^2}{ad^2}} \right)}{iA+B} \right) \right)$$

input `integrate((A+B*tan(dx+c))/(a+I*a*tan(dx+c))^(1/2),x, algorithm="fricas")`

output $1/4*(\sqrt{2})*a*d*\sqrt{-(A^2 - 2*I*A*B - B^2)/(a*d^2)}*e^{(I*d*x + I*c)}*\log(-4*((-I*A - B)*a*e^{(I*d*x + I*c)} + (a*d*e^{(2*I*d*x + 2*I*c)} + a*d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*\sqrt{-(A^2 - 2*I*A*B - B^2)/(a*d^2)})*e^{(-I*d*x - I*c)/(I*A + B)} - \sqrt{2}*a*d*\sqrt{-(A^2 - 2*I*A*B - B^2)/(a*d^2)}*e^{(I*d*x + I*c)}*\log(-4*((-I*A - B)*a*e^{(I*d*x + I*c)} - (a*d*e^{(2*I*d*x + 2*I*c)} + a*d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*\sqrt{-(A^2 - 2*I*A*B - B^2)/(a*d^2)})*e^{(-I*d*x - I*c)/(I*A + B)} - 2*\sqrt{2}*((-I*A + B)*e^{(2*I*d*x + 2*I*c)} - I*A + B)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*e^{(-I*d*x - I*c)/(a*d)}$

3.93.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{a + ia \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{ia (\tan(c + dx) - i)}} dx$$

input `integrate((A+B*tan(dx+c))/(a+I*a*tan(dx+c))**(1/2),x)`

output `Integral((A + B*tan(c + dx))/sqrt(I*a*(tan(c + dx) - I)), x)`

3.93.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.11

$$\int \frac{A + B \tan(c + dx)}{\sqrt{a + ia \tan(c + dx)}} dx$$

$$= \frac{i \left(\sqrt{2}(A - iB)\sqrt{a} \log \left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}} \right) + \frac{4(A+iB)a}{\sqrt{ia \tan(dx+c)+a}} \right)}{4ad}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="maxima")`output `1/4*I*(sqrt(2)*(A - I*B)*sqrt(a)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) + 4*(A + I*B)*a/sqrt(I*a*tan(d*x + c) + a))/(a*d)`**3.93.8 Giac [F]**

$$\int \frac{A + B \tan(c + dx)}{\sqrt{a + ia \tan(c + dx)}} dx = \int \frac{B \tan(dx + c) + A}{\sqrt{ia \tan(dx + c) + a}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")`output `integrate((B*tan(d*x + c) + A)/sqrt(I*a*tan(d*x + c) + a), x)`**3.93.9 Mupad [B] (verification not implemented)**

Time = 0.89 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.43

$$\int \frac{A + B \tan(c + dx)}{\sqrt{a + ia \tan(c + dx)}} dx = \frac{A \operatorname{li}}{d \sqrt{a + a \tan(c + dx)} \operatorname{li}} - \frac{B}{d \sqrt{a + a \tan(c + dx)} \operatorname{li}}$$

$$+ \frac{\sqrt{2} A \operatorname{atan} \left(\frac{\sqrt{2} \sqrt{a+a \tan(c+dx)} \operatorname{li}}{2 \sqrt{-a}} \right) \operatorname{li}}{2 \sqrt{-a} d}$$

$$- \frac{\sqrt{2} B \operatorname{atanh} \left(\frac{\sqrt{2} \sqrt{a+a \tan(c+dx)} \operatorname{li}}{2 \sqrt{a}} \right)}{2 \sqrt{a} d}$$

input `int((A + B*tan(c + d*x))/(a + a*tan(c + d*x)*1i)^(1/2),x)`

output `(A*1i)/(d*(a + a*tan(c + d*x)*1i)^(1/2)) - B/(d*(a + a*tan(c + d*x)*1i)^(1/2)) + (2^(1/2)*A*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2)))*1i)/(2*(-a)^(1/2)*d) - (2^(1/2)*B*atanh((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*a^(1/2))))/(2*a^(1/2)*d)`

3.94 $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

3.94.1	Optimal result	1039
3.94.2	Mathematica [A] (verified)	1039
3.94.3	Rubi [A] (verified)	1040
3.94.4	Maple [A] (verified)	1043
3.94.5	Fricas [B] (verification not implemented)	1044
3.94.6	Sympy [F]	1045
3.94.7	Maxima [A] (verification not implemented)	1045
3.94.8	Giac [F]	1045
3.94.9	Mupad [B] (verification not implemented)	1046

3.94.1 Optimal result

Integrand size = 34, antiderivative size = 114

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = -\frac{2A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ad}} + \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}} + \frac{A+iB}{d\sqrt{a+ia \tan(c+dx)}}$$

output

```
-2*A*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/d/a^(1/2)+1/2*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/d*2^(1/2)/a^(1/2)+(A+I*B)/d/(a+I*a*tan(d*x+c))^(1/2)
```

3.94.2 Mathematica [A] (verified)

Time = 2.06 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.05

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \frac{i \left(\frac{4iA \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{a}} - \frac{\sqrt{2}(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{a}} - \frac{2(iA-B)}{\sqrt{a+ia \tan(c+dx)}} \right)}{2d}$$

input `Integrate[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x
]`

output `((I/2)*(((4*I)*A*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]]/Sqrt[a] - (S
qrt[2]*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])))/Sq
rt[a] - (2*(I*A - B))/Sqrt[a + I*a*Tan[c + d*x]]))/d`

3.94.3 Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.10, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.324$, Rules used = {3042, 4079, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)\sqrt{a+ia \tan(c+dx)}} dx \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{1}{2} \cot(c+dx) \sqrt{i \tan(c+dx)a+a(2aA-a(iA-B) \tan(c+dx))} dx}{a^2} + \frac{A+iB}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \cot(c+dx) \sqrt{i \tan(c+dx)a+a(2aA-a(iA-B) \tan(c+dx))} dx}{2a^2} + \frac{A+iB}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\sqrt{i \tan(c+dx)a+a(2aA-a(iA-B) \tan(c+dx))}}{\tan(c+dx)} dx}{2a^2} + \frac{A+iB}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \quad \downarrow \text{4083} \\
 & \frac{a(B+iA) \int \sqrt{i \tan(c+dx)a+adx} + 2A \int \cot(c+dx)(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{2a^2} + \frac{A+iB}{d\sqrt{a+ia \tan(c+dx)}}
 \end{aligned}$$

3.94. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{a(B+iA) \int \sqrt{i \tan(c+dx)a+adx} + 2A \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx}{\frac{2a^2}{A+iB} d\sqrt{a+ia \tan(c+dx)}} + \\
& \downarrow 3961 \\
& \frac{2A \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{2ia^2(B+iA) \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a}}{\frac{2a^2}{A+iB} d\sqrt{a+ia \tan(c+dx)}}}{\frac{2a^2}{A+iB} d\sqrt{a+ia \tan(c+dx)}} + \\
& \downarrow 219 \\
& \frac{2A \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{i\sqrt{2}a^{3/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{\frac{2a^2}{A+iB} d\sqrt{a+ia \tan(c+dx)}} + \\
& \downarrow 4082 \\
& \frac{2a^2 A \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx) - \frac{i\sqrt{2}a^{3/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{2a^2} + \frac{A+iB}{d\sqrt{a+ia \tan(c+dx)}} \\
& \downarrow 73 \\
& \frac{-\frac{4iaA \int \frac{1}{i-i \tan(c+dx)a+a} d\sqrt{i \tan(c+dx)a+a}}{d} - \frac{i\sqrt{2}a^{3/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{\frac{2a^2}{A+iB} d\sqrt{a+ia \tan(c+dx)}} + \\
& \downarrow 221 \\
& \frac{-\frac{i\sqrt{2}a^{3/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{4a^{3/2}A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{d}}{2a^2} + \frac{A+iB}{d\sqrt{a+ia \tan(c+dx)}}
\end{aligned}$$

input `Int[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`


```
output ((-4*a^(3/2)*A*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d - (I*Sqrt[2]
*a^(3/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/
d)/(2*a^2) + (A + I*B)/(d*Sqrt[a + I*a*Tan[c + d*x]])
```

3.94.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3961 Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d)
Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a
, b, c, d}, x] && EqQ[a^2 + b^2, 0]
```

```
rule 4079 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

```
rule 4082 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], Tan[e + f*x]],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 + b^2, 0] && EqQ[A*b + a*B, 0]
```

```
rule 4083 Int((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(
A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A
*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*T
an[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

3.94.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.87

method	result	size
derivativedivides	$2a \left(-\frac{(iB-A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{4a^{\frac{3}{2}}} - \frac{-iB-A}{2a\sqrt{a+ia \tan(dx+c)}} - \frac{A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{\sqrt{a}}\right)}{a^{\frac{3}{2}}} \right) / d$	99
default	$2a \left(-\frac{(iB-A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{4a^{\frac{3}{2}}} - \frac{-iB-A}{2a\sqrt{a+ia \tan(dx+c)}} - \frac{A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{\sqrt{a}}\right)}{a^{\frac{3}{2}}} \right) / d$	99

```
input int(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2), x, method=_RETURNV
ERBOSE)
```

$$3.94. \int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

output $2/d*a*(-1/4*(-A+I*B)/a^{(3/2)}*2^{(1/2)}*\operatorname{arctanh}(1/2*(a+I*a*\tan(d*x+c))^{(1/2)}*2^{(1/2)}/a^{(1/2)})-1/2*(-A-I*B)/a/(a+I*a*\tan(d*x+c))^{(1/2)}-1/a^{(3/2)}*A*\operatorname{arctanh}((a+I*a*\tan(d*x+c))^{(1/2)}/a^{(1/2)}))$

3.94.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 575 vs. $2(88) = 176$.

Time = 0.27 (sec) , antiderivative size = 575, normalized size of antiderivative = 5.04

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \frac{\sqrt{2}ad\sqrt{\frac{A^2-2iAB-B^2}{ad^2}}e^{(idx+ic)}\log\left(-\frac{4\left((-iA-B)ae^{(idx+ic)}+(iade^{(2idx+2ic)}+iad)\sqrt{\frac{a}{e^{(2idx+2ic)}+1}}\sqrt{\frac{A^2-2iAB-B^2}{ad^2}}\right)}{iA+B}\right)}{1}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="fricas")`

output $-1/4*(\sqrt{2}*a*d*\sqrt{(A^2 - 2*I*A*B - B^2)/(a*d^2)})*e^{(I*d*x + I*c)}*\log(-4*((-I*A - B)*a*e^{(I*d*x + I*c)} + (I*a*d*e^{(2*I*d*x + 2*I*c)} + I*a*d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*\sqrt{(A^2 - 2*I*A*B - B^2)/(a*d^2)})*e^{(-I*d*x - I*c)/(I*A + B)} - \sqrt{2}*a*d*\sqrt{(A^2 - 2*I*A*B - B^2)/(a*d^2)})*e^{(I*d*x + I*c)}*\log(-4*((-I*A - B)*a*e^{(I*d*x + I*c)} + (-I*a*d*e^{(2*I*d*x + 2*I*c)} - I*a*d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*\sqrt{(A^2 - 2*I*A*B - B^2)/(a*d^2)})*e^{(-I*d*x - I*c)/(I*A + B)} + 2*a*d*\sqrt{A^2/(a*d^2)})*e^{(I*d*x + I*c)}*\log(16*(3*A*a^2*e^{(2*I*d*x + 2*I*c)} + A*a^2 + 2*\sqrt{2}*(a^2*d*e^{(3*I*d*x + 3*I*c)} + a^2*d*e^{(I*d*x + I*c)})*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*\sqrt{A^2/(a*d^2)})*e^{(-2*I*d*x - 2*I*c)/A} - 2*a*d*\sqrt{A^2/(a*d^2)})*e^{(I*d*x + I*c)}*\log(16*(3*A*a^2*e^{(2*I*d*x + 2*I*c)} + A*a^2 - 2*\sqrt{2}*(a^2*d*e^{(3*I*d*x + 3*I*c)} + a^2*d*e^{(I*d*x + I*c)})*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*\sqrt{A^2/(a*d^2)})*e^{(-2*I*d*x - 2*I*c)/A} - 2*\sqrt{2}*((A + I*B)*e^{(2*I*d*x + 2*I*c)} + A + I*B)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*e^{(-I*d*x - I*c)/(a*d)}$

3.94.6 Sympy [F]

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \int \frac{(A+B \tan(c+dx)) \cot(c+dx)}{\sqrt{ia(\tan(c+dx)-i)}} dx$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)/sqrt(I*a*(tan(c + d*x) - I)), x)`

3.94.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.17

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

$$= -\frac{\sqrt{2}(A-iB) \log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia \tan(dx+c)+a}}\right)}{\sqrt{a}} - \frac{4A \log\left(\frac{\sqrt{ia \tan(dx+c)+a}-\sqrt{a}}{\sqrt{ia \tan(dx+c)+a}+\sqrt{a}}\right)}{\sqrt{a}} - \frac{4(A+iB)}{\sqrt{ia \tan(dx+c)+a}}$$

$$4d$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `-1/4*(sqrt(2)*(A - I*B)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a)))/sqrt(a) - 4*A*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a)))/sqrt(a) - 4*(A + I*B)/sqrt(I*a*tan(d*x + c) + a))/d`

3.94.8 Giac [F]

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \int \frac{(B \tan(dx+c) + A) \cot(dx+c)}{\sqrt{ia \tan(dx+c) + a}} dx$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)/sqrt(I*a*tan(d*x + c) + a), x)`

3.94.9 Mupad [B] (verification not implemented)

Time = 7.80 (sec) , antiderivative size = 515, normalized size of antiderivative = 4.52

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \frac{A+B \operatorname{li}}{d \sqrt{a+a \tan(c+dx)} \operatorname{li}}$$

$$- \frac{2A \operatorname{atanh}\left(\frac{28A^3 a^{3/2} d \sqrt{a+a \tan(c+dx)} \operatorname{li}}{28dA^3 a^2+8idA^2 B a^2+4dA B^2 a^2} + \frac{4AB^2 a^{3/2} d \sqrt{a+a \tan(c+dx)} \operatorname{li}}{28dA^3 a^2+8idA^2 B a^2+4dA B^2 a^2} + \frac{A^2 B a^{3/2} d \sqrt{a+a \tan(c+dx)} \operatorname{li} 8i}{28dA^3 a^2+8idA^2 B a^2+4dA B^2 a^2}\right)}{\sqrt{a} d}$$

$$+ \frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} A^3 (-a)^{3/2} d \sqrt{a+a \tan(c+dx)} \operatorname{li} 7i}{2(7dA^3 a^2-5idA^2 B a^2+3dA B^2 a^2-1idB^3 a^2)} + \frac{\sqrt{2} B^3 (-a)^{3/2} d \sqrt{a+a \tan(c+dx)} \operatorname{li}}{2(7dA^3 a^2-5idA^2 B a^2+3dA B^2 a^2-1idB^3 a^2)} + \frac{\sqrt{2} A B^2 (-a)^{3/2} d \sqrt{a+a \tan(c+dx)} \operatorname{li} 8i}{2(7dA^3 a^2-5idA^2 B a^2+3dA B^2 a^2-1idB^3 a^2)}\right)}{2\sqrt{-a} d}$$

```
input int((cot(c + d*x)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2),x)
```

```
output (A + B*1i)/(d*(a + a*tan(c + d*x)*1i)^(1/2)) - (2*A*atanh((28*A^3*a^(3/2)*
d*(a + a*tan(c + d*x)*1i)^(1/2))/(28*A^3*a^2*d + 4*A*B^2*a^2*d + A^2*B*a^2
*d*8i) + (4*A*B^2*a^(3/2)*d*(a + a*tan(c + d*x)*1i)^(1/2))/(28*A^3*a^2*d +
4*A*B^2*a^2*d + A^2*B*a^2*d*8i) + (A^2*B*a^(3/2)*d*(a + a*tan(c + d*x)*1i
)^(1/2)*8i)/(28*A^3*a^2*d + 4*A*B^2*a^2*d + A^2*B*a^2*d*8i)))/(a^(1/2)*d)
+ (2^(1/2)*atanh((2^(1/2)*A^3*(-a)^(3/2)*d*(a + a*tan(c + d*x)*1i)^(1/2)*7
i)/(2*(7*A^3*a^2*d - B^3*a^2*d*1i + 3*A*B^2*a^2*d - A^2*B*a^2*d*5i)) + (2^(
1/2)*B^3*(-a)^(3/2)*d*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(7*A^3*a^2*d - B^
3*a^2*d*1i + 3*A*B^2*a^2*d - A^2*B*a^2*d*5i)) + (2^(1/2)*A*B^2*(-a)^(3/2)*
d*(a + a*tan(c + d*x)*1i)^(1/2)*3i)/(2*(7*A^3*a^2*d - B^3*a^2*d*1i + 3*A*B
^2*a^2*d - A^2*B*a^2*d*5i)) + (5*2^(1/2)*A^2*B*(-a)^(3/2)*d*(a + a*tan(c +
d*x)*1i)^(1/2))/(2*(7*A^3*a^2*d - B^3*a^2*d*1i + 3*A*B^2*a^2*d - A^2*B*a^
2*d*5i)))*(A*1i + B))/(2*(-a)^(1/2)*d)
```

3.95
$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

3.95.1 Optimal result 1047
 3.95.2 Mathematica [A] (verified) 1048
 3.95.3 Rubi [A] (verified) 1048
 3.95.4 Maple [A] (verified) 1053
 3.95.5 Fricas [B] (verification not implemented) 1053
 3.95.6 Sympy [F] 1054
 3.95.7 Maxima [A] (verification not implemented) 1055
 3.95.8 Giac [F] 1055
 3.95.9 Mupad [B] (verification not implemented) 1056

3.95.1 Optimal result

Integrand size = 36, antiderivative size = 167

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \frac{(iA-2B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ad}} + \frac{(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}} + \frac{(A+iB)\cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{(2A+iB)\cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{ad}$$

```
output (I*A-2*B)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/d/a^(1/2)+1/2*(I*A+B)*
arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/d*2^(1/2)/a^(1/2)+(A
+I*B)*cot(d*x+c)/d/(a+I*a*tan(d*x+c))^(1/2)-(2*A+I*B)*cot(d*x+c)*(a+I*a*ta
n(d*x+c))^(1/2)/a/d
```

3.95.2 Mathematica [A] (verified)

Time = 3.64 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.78

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \frac{i(A+2iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ad}} + \frac{(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}} + \frac{-2iA+B-A \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}}$$

input `Integrate[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `(I*(A + (2*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]]/(Sqrt[a]*d) + ((I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/(Sqrt[2]*Sqrt[a]*d) + ((-2*I)*A + B - A*Cot[c + d*x])/(d*Sqrt[a + I*a*Tan[c + d*x]])`

3.95.3 Rubi [A] (verified)

Time = 1.13 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.10, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3042, 4079, 27, 3042, 4081, 25, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

↓ 3042

$$\int \frac{A+B \tan(c+dx)}{\tan(c+dx)^2 \sqrt{a+ia \tan(c+dx)}} dx$$

↓ 4079

$$\begin{aligned}
& \frac{\int \frac{1}{2} \cot^2(c+dx) \sqrt{i \tan(c+dx)a+a(2a(2A+iB)-3a(iA-B)\tan(c+dx))} dx}{a^2} + \\
& \quad \frac{(A+iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 27 \\
& \frac{\int \cot^2(c+dx) \sqrt{i \tan(c+dx)a+a(2a(2A+iB)-3a(iA-B)\tan(c+dx))} dx}{2a^2} + \\
& \quad \frac{(A+iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\sqrt{i \tan(c+dx)a+a(2a(2A+iB)-3a(iA-B)\tan(c+dx))}}{\tan(c+dx)^2} dx}{2a^2} + \frac{(A+iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 4081 \\
& \frac{\int -\cot(c+dx) \sqrt{i \tan(c+dx)a+a((iA-2B)a^2+(2A+iB)\tan(c+dx)a^2)} dx}{a} - \frac{2a(2A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \\
& \quad \frac{(A+iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 25 \\
& \frac{\int \cot(c+dx) \sqrt{i \tan(c+dx)a+a((iA-2B)a^2+(2A+iB)\tan(c+dx)a^2)} dx}{a} - \frac{2a(2A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \\
& \quad \frac{(A+iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\sqrt{i \tan(c+dx)a+a((iA-2B)a^2+(2A+iB)\tan(c+dx)a^2)}}{\tan(c+dx)} dx}{a} - \frac{2a(2A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \\
& \quad \frac{(A+iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 4083 \\
& \frac{-a^2(A-iB) \int \sqrt{i \tan(c+dx)a+adx+a(-2B+iA) \int \cot(c+dx)(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}}{a} - \frac{2a(2A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \\
& \quad \frac{(A+iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 3042
\end{aligned}$$

3.95. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

$$\begin{aligned}
& \frac{a^2(A-iB) \int \sqrt{i \tan(c+dx)a+adx+a}(-2B+iA) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx}{a} - \frac{2a(2A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \\
& \frac{2a^2}{d\sqrt{a+ia \tan(c+dx)}} \frac{(A+iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow \text{3961} \\
& \frac{a(-2B+iA) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{2ia^3(A-iB) \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a}}{a}}{a} - \frac{2a(2A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \\
& \frac{2a^2}{d\sqrt{a+ia \tan(c+dx)}} \frac{(A+iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow \text{219} \\
& \frac{a(-2B+iA) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{i\sqrt{2}a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{a} - \frac{2a(2A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \\
& \frac{2a^2}{d\sqrt{a+ia \tan(c+dx)}} \frac{(A+iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow \text{4082} \\
& \frac{a^3(-2B+iA) \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx) - \frac{i\sqrt{2}a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{a} - \frac{2a(2A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \\
& \frac{2a^2}{d\sqrt{a+ia \tan(c+dx)}} \frac{(A+iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow \text{73} \\
& \frac{2ia^2(-2B+iA) \int \frac{1}{i-\frac{1}{i \tan(c+dx)a+a}} d\sqrt{i \tan(c+dx)a+a} - \frac{i\sqrt{2}a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{a} - \frac{2a(2A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \\
& \frac{2a^2}{d\sqrt{a+ia \tan(c+dx)}} \frac{(A+iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow \text{221} \\
& \frac{2a^{5/2}(-2B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) - \frac{i\sqrt{2}a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{a} - \frac{2a(2A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \\
& \frac{2a^2}{d\sqrt{a+ia \tan(c+dx)}} \frac{(A+iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}}
\end{aligned}$$

3.95. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

input `Int[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `((A + I*B)*Cot[c + d*x])/(d*Sqrt[a + I*a*Tan[c + d*x]]) + (-(((-2*a^(5/2)*
(I*A - 2*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d - (I*Sqrt[2]*a^(5/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d)/
a) - (2*a*(2*A + I*B)*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]])/d)/(2*a^2)`

3.95.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d)
Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a
, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4079 $\text{Int}[(a_+ + (b_-)\tan[(e_-) + (f_-)(x_+)]^{(m_-)}((A_-) + (B_-)\tan[(e_-) + (f_-)(x_+)])(c_-) + (d_-)\tan[(e_-) + (f_-)(x_+)]^{(n_-)}, x_Symbol] \rightarrow \text{Simp}[(aA + bB)(a + b\tan[e + fx])^m((c + d\tan[e + fx])^{(n+1)}(2fm*(b*c - a*d))), x] + \text{Simp}[1/(2a*m*(b*c - a*d)) \text{Int}[(a + b\tan[e + fx])^{(m+1)}(c + d\tan[e + fx])^n \text{Simp}[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*\tan[e + fx], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[a^2 + b^2, 0] \&\& \text{LtQ}[m, 0] \&\& !\text{GtQ}[n, 0]$

rule 4081 $\text{Int}[(a_+ + (b_-)\tan[(e_-) + (f_-)(x_+)]^{(m_-)}((A_-) + (B_-)\tan[(e_-) + (f_-)(x_+)])(c_-) + (d_-)\tan[(e_-) + (f_-)(x_+)]^{(n_-)}, x_Symbol] \rightarrow \text{Simp}[(A*d - B*c)(a + b\tan[e + fx])^m((c + d\tan[e + fx])^{(n+1)}(f*(n+1)*(c^2 + d^2))), x] - \text{Simp}[1/(a*(n+1)*(c^2 + d^2)) \text{Int}[(a + b\tan[e + fx])^m(c + d\tan[e + fx])^{(n+1)} \text{Simp}[A*(b*d*m - a*c*(n+1)) - B*(b*c*m + a*d*(n+1)) - a*(B*c - A*d)*(m + n + 1)*\tan[e + fx], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[a^2 + b^2, 0] \&\& \text{LtQ}[n, -1]$

rule 4082 $\text{Int}[(a_+ + (b_-)\tan[(e_-) + (f_-)(x_+)]^{(m_-)}((A_-) + (B_-)\tan[(e_-) + (f_-)(x_+)])(c_-) + (d_-)\tan[(e_-) + (f_-)(x_+)]^{(n_-)}, x_Symbol] \rightarrow \text{Simp}[b*(B/f) \text{Subst}[\text{Int}[(a + b*x)^{(m-1)}(c + d*x)^n, x], x, \tan[e + fx]], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, m, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[a^2 + b^2, 0] \&\& \text{EqQ}[A*b + a*B, 0]$

rule 4083 $\text{Int}[(a_+ + (b_-)\tan[(e_-) + (f_-)(x_+)]^{(m_-)}((A_-) + (B_-)\tan[(e_-) + (f_-)(x_+)])/((c_-) + (d_-)\tan[(e_-) + (f_-)(x_+)]), x_Symbol] \rightarrow \text{Simp}[(A*b + a*B)/(b*c + a*d) \text{Int}[(a + b\tan[e + fx])^m, x], x] - \text{Simp}[(B*c - A*d)/(b*c + a*d) \text{Int}[(a + b\tan[e + fx])^m((a - b\tan[e + fx])/(c + d\tan[e + fx])), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[a^2 + b^2, 0] \&\& \text{NeQ}[A*b + a*B, 0]$

3.95.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.83

method	result
derivativedivides	$2ia^2 \left(-\frac{iB+A}{2a^2\sqrt{a+ia\tan(dx+c)}} - \frac{(iB-A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{4a^{\frac{5}{2}}} + \frac{iA\sqrt{a+ia\tan(dx+c)}}{2a\tan(dx+c)} + \frac{(2iB+A) \operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(dx+c)}}{\sqrt{a}}\right)}{a^2} \right) \frac{1}{d}$
default	$2ia^2 \left(-\frac{iB+A}{2a^2\sqrt{a+ia\tan(dx+c)}} - \frac{(iB-A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{4a^{\frac{5}{2}}} + \frac{iA\sqrt{a+ia\tan(dx+c)}}{2a\tan(dx+c)} + \frac{(2iB+A) \operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(dx+c)}}{\sqrt{a}}\right)}{a^2} \right) \frac{1}{d}$

input `int(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output `2*I/d*a^2*(-1/2/a^2*(A+I*B)/(a+I*a*tan(d*x+c))^(1/2)-1/4*(-A+I*B)/a^(5/2)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))+1/a^2*(1/2*I*A*(a+I*a*tan(d*x+c))^(1/2)/a/tan(d*x+c)+1/2*(A+2*I*B)/a^(1/2)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2)))`

3.95.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 742 vs. 2(135) = 270.

Time = 0.27 (sec) , antiderivative size = 742, normalized size of antiderivative = 4.44

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \frac{\sqrt{2}(ade^{(3i dx+3i c)} - ade^{(i dx+i c)})\sqrt{-\frac{A^2-2iAB-B^2}{ad^2}} \log\left(-\frac{4\left((-iA-B)ae^{(i dx+i c)}+(ade^{(2i dx+2i c)}+ad)\sqrt{\frac{a}{e^{(2i dx+2i c)}+1}}\right)}{iA+B}\right)}{ad^2}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x,algorithm="fracas")`

output

```
-1/4*(sqrt(2)*(a*d*e^(3*I*d*x + 3*I*c) - a*d*e^(I*d*x + I*c))*sqrt(-(A^2 -
2*I*A*B - B^2)/(a*d^2))*log(-4*((-I*A - B)*a*e^(I*d*x + I*c) + (a*d*e^(2*
I*d*x + 2*I*c) + a*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-(A^2 - 2*I*A
*B - B^2)/(a*d^2)))*e^(-I*d*x - I*c)/(I*A + B)) - sqrt(2)*(a*d*e^(3*I*d*x
+ 3*I*c) - a*d*e^(I*d*x + I*c))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a*d^2))*log(-
4*((-I*A - B)*a*e^(I*d*x + I*c) - (a*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt(a/(
e^(2*I*d*x + 2*I*c) + 1))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a*d^2)))*e^(-I*d*x
- I*c)/(I*A + B)) - (a*d*e^(3*I*d*x + 3*I*c) - a*d*e^(I*d*x + I*c))*sqrt(-
(A^2 + 4*I*A*B - 4*B^2)/(a*d^2))*log(-16*(3*(I*A - 2*B)*a^2*e^(2*I*d*x + 2
*I*c) + (I*A - 2*B)*a^2 + 2*sqrt(2)*(a^2*d*e^(3*I*d*x + 3*I*c) + a^2*d*e^(
I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-(A^2 + 4*I*A*B - 4*B
^2)/(a*d^2)))*e^(-2*I*d*x - 2*I*c)/(-I*A + 2*B)) + (a*d*e^(3*I*d*x + 3*I*c
) - a*d*e^(I*d*x + I*c))*sqrt(-(A^2 + 4*I*A*B - 4*B^2)/(a*d^2))*log(-16*(3
*(I*A - 2*B)*a^2*e^(2*I*d*x + 2*I*c) + (I*A - 2*B)*a^2 - 2*sqrt(2)*(a^2*d*
e^(3*I*d*x + 3*I*c) + a^2*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) +
1))*sqrt(-(A^2 + 4*I*A*B - 4*B^2)/(a*d^2)))*e^(-2*I*d*x - 2*I*c)/(-I*A +
2*B)) + 2*sqrt(2)*((3*I*A - B)*e^(4*I*d*x + 4*I*c) + 2*I*A*e^(2*I*d*x + 2*
I*c) - I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(a*d*e^(3*I*d*x + 3*I*c
) - a*d*e^(I*d*x + I*c))
```

3.95.6 Sympy [F]

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \int \frac{(A+B \tan(c+dx)) \cot^2(c+dx)}{\sqrt{ia(\tan(c+dx)-i)}} dx$$

input `integrate(cot(d*x+c)**2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**2/sqrt(I*a*(tan(c + d*x) - I)), x)`

3.95.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.10

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx =$$

$$ia \left(\frac{\sqrt{2}(A-iB) \log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia \tan(dx+c)+a}}\right)}{a^{\frac{3}{2}}} + \frac{2(A+2iB) \log\left(\frac{\sqrt{ia \tan(dx+c)+a}-\sqrt{a}}{\sqrt{ia \tan(dx+c)+a}+\sqrt{a}}\right)}{a^{\frac{3}{2}}} + \frac{4((ia \tan(dx+c)+a)(2A+iB)-(A+iB)a)}{(ia \tan(dx+c)+a)^{\frac{3}{2}}a-\sqrt{ia \tan(dx+c)+a}} \right) + \frac{}{4d}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `-1/4*I*a*(sqrt(2)*(A - I*B)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a)))/a^(3/2) + 2*(A + 2*I*B)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a)))/a^(3/2) + 4*((I*a*tan(d*x + c) + a)*(2*A + I*B) - (A + I*B)*a)/((I*a*tan(d*x + c) + a)^(3/2)*a - sqrt(I*a*tan(d*x + c) + a)*a^2))/d`

3.95.8 Giac [F]

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \int \frac{(B \tan(dx+c) + A) \cot(dx+c)^2}{\sqrt{ia \tan(dx+c) + a}} dx$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^2/sqrt(I*a*tan(d*x + c) + a), x)`

3.95.9 Mupad [B] (verification not implemented)

Time = 9.46 (sec) , antiderivative size = 2961, normalized size of antiderivative = 17.73

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \text{Too large to display}$$

input `int((cot(c + d*x)^2*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2),x)`

output `2*atanh((3*d^4*(a + a*tan(c + d*x)*1i)^(1/2)*((9*B^2)/(16*a*d^2) - (3*A^2)/(16*a*d^2) - ((A^4*a^10)/d^4 + (49*B^4*a^10)/d^4 - (114*A^2*B^2*a^10)/d^4 - (A*B^3*a^10*140i)/d^4 + (A^3*B*a^10*20i)/d^4)^(1/2))/(16*a^6) - (A*B^3i)/(8*a*d^2))^(1/2)*((A^4*a^10)/d^4 + (49*B^4*a^10)/d^4 - (114*A^2*B^2*a^10)/d^4 - (A*B^3*a^10*140i)/d^4 + (A^3*B*a^10*20i)/d^4)^(1/2))/((A^3*a^5*d*1i)/2 + (35*B^3*a^5*d)/2 + (A*d^3*((A^4*a^10)/d^4 + (49*B^4*a^10)/d^4 - (114*A^2*B^2*a^10)/d^4 - (A*B^3*a^10*140i)/d^4 + (A^3*B*a^10*20i)/d^4)^(1/2))*3i)/2 - (3*B*d^3*((A^4*a^10)/d^4 + (49*B^4*a^10)/d^4 - (114*A^2*B^2*a^10)/d^4 - (A*B^3*a^10*140i)/d^4 + (A^3*B*a^10*20i)/d^4)^(1/2))/2 - (A*B^2*a^5*d*57i)/2 - (15*A^2*B*a^5*d)/2) + (A^2*a^2*d^2*(a + a*tan(c + d*x)*1i)^(1/2)*((9*B^2)/(16*a*d^2) - (3*A^2)/(16*a*d^2) - ((A^4*a^10)/d^4 + (49*B^4*a^10)/d^4 - (114*A^2*B^2*a^10)/d^4 - (A*B^3*a^10*140i)/d^4 + (A^3*B*a^10*20i)/d^4)^(1/2))/(16*a^6) - (A*B^3i)/(8*a*d^2))^(1/2))/((A^3*a^2*d*1i)/2 + (35*B^3*a^2*d)/2 - (A*B^2*a^2*d*57i)/2 - (15*A^2*B*a^2*d)/2 + (A*d^3*((A^4*a^10)/d^4 + (49*B^4*a^10)/d^4 - (114*A^2*B^2*a^10)/d^4 - (A*B^3*a^10*140i)/d^4 + (A^3*B*a^10*20i)/d^4)^(1/2))*3i)/(2*a^3) - (3*B*d^3*((A^4*a^10)/d^4 + (49*B^4*a^10)/d^4 - (114*A^2*B^2*a^10)/d^4 - (A*B^3*a^10*140i)/d^4 + (A^3*B*a^10*20i)/d^4)^(1/2))/(2*a^3)) - (7*B^2*a^2*d^2*(a + a*tan(c + d*x)*1i)^(1/2)*((9*B^2)/(16*a*d^2) - (3*A^2)/(16*a*d^2) - ((A^4*a^10)/d^4 + (49*B^4*a^10)/d^4 - (114*A^2*B^2*a^10)/d^4 - (A*B^3*a^10*140i)/d^4 + (A^3*B*a^10...)`

3.96 $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

3.96.1	Optimal result	1057
3.96.2	Mathematica [A] (verified)	1058
3.96.3	Rubi [A] (verified)	1058
3.96.4	Maple [A] (verified)	1063
3.96.5	Fricas [B] (verification not implemented)	1064
3.96.6	Sympy [F]	1065
3.96.7	Maxima [A] (verification not implemented)	1065
3.96.8	Giac [F]	1066
3.96.9	Mupad [B] (verification not implemented)	1066

3.96.1 Optimal result

Integrand size = 36, antiderivative size = 219

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \frac{(11A+4iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{4\sqrt{ad}} - \frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}} + \frac{(A+iB)\cot^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} + \frac{(7iA-8B)\cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{4ad} - \frac{(3A+2iB)\cot^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{2ad}$$

output $\frac{1}{4}(11A+4iB)\operatorname{arctanh}\left(\frac{(a+Ia*\tan(dx+c))^{1/2}/a^{1/2}}{d/a^{1/2}}\right)-\frac{1}{2}(A-I*B)\operatorname{arctanh}\left(\frac{1/2*(a+Ia*\tan(dx+c))^{1/2}*2^{1/2}/a^{1/2}}{d*2^{1/2}/a^{1/2}}\right)+(A+I*B)*\cot(dx+c)^2/d/(a+Ia*\tan(dx+c))^{1/2}+\frac{1}{4}(7iA-8B)*\cot(dx+c)*(a+Ia*\tan(dx+c))^{1/2}/a/d-\frac{1}{2}(3A+2iB)*\cot(dx+c)^2*(a+Ia*\tan(dx+c))^{1/2}/a/d$

3.96.2 Mathematica [A] (verified)

Time = 4.42 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.68

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

$$= \frac{(11A+4iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{a}} - \frac{2\sqrt{2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{a}} + \frac{-7A-8iB+i(A+4iB) \cot(c+dx)-2A \cot^2(c+dx)}{\sqrt{a+ia \tan(c+dx)}} + \frac{}{4d}$$

input `Integrate[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `((((11*A + (4*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/Sqrt[a] - (2*Sqrt[2]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/Sqrt[a] + (-7*A - (8*I)*B + I*(A + (4*I)*B)*Cot[c + d*x] - 2*A*Cot[c + d*x]^2)/Sqrt[a + I*a*Tan[c + d*x]])/(4*d)`

3.96.3 Rubi [A] (verified)

Time = 1.52 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.08, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.472$, Rules used = {3042, 4079, 27, 3042, 4081, 25, 3042, 4081, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A+B \tan(c+dx)}{\tan^3(c+dx) \sqrt{a+ia \tan(c+dx)}} dx$$

$$\downarrow \text{4079}$$

$$\int \frac{\frac{1}{2} \cot^3(c+dx) \sqrt{i \tan(c+dx) a + a(2a(3A+2iB) - 5a(iA-B) \tan(c+dx))} dx}{d \sqrt{a+ia \tan(c+dx)}} + \frac{(A+iB) \cot^2(c+dx)}{d \sqrt{a+ia \tan(c+dx)}}$$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{\int \cot^3(c+dx) \sqrt{i \tan(c+dx)a+a(2a(3A+2iB)-5a(iA-B)\tan(c+dx))} dx}{2a^2} + \\
 & \quad \frac{(A+iB) \cot^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \downarrow 3042 \\
 & \frac{\int \frac{\sqrt{i \tan(c+dx)a+a(2a(3A+2iB)-5a(iA-B)\tan(c+dx))}}{\tan(c+dx)^3} dx}{2a^2} + \frac{(A+iB) \cot^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \downarrow 4081 \\
 & \frac{\int -\cot^2(c+dx) \sqrt{i \tan(c+dx)a+a((7iA-8B)a^2+3(3A+2iB)\tan(c+dx)a^2)} dx}{2a} - \frac{a(3A+2iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \\
 & \quad \frac{2a^2}{(A+iB) \cot^2(c+dx)} \\
 & \quad \frac{d\sqrt{a+ia \tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \downarrow 25 \\
 & \frac{\int \cot^2(c+dx) \sqrt{i \tan(c+dx)a+a((7iA-8B)a^2+3(3A+2iB)\tan(c+dx)a^2)} dx}{2a} - \frac{a(3A+2iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \\
 & \quad \frac{2a^2}{(A+iB) \cot^2(c+dx)} \\
 & \quad \frac{d\sqrt{a+ia \tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \downarrow 3042 \\
 & - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a((7iA-8B)a^2+3(3A+2iB)\tan(c+dx)a^2)}}{\tan(c+dx)^2} dx}{2a} - \frac{a(3A+2iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \\
 & \quad \frac{2a^2}{(A+iB) \cot^2(c+dx)} \\
 & \quad \frac{d\sqrt{a+ia \tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \downarrow 4081 \\
 & - \frac{\int \frac{1}{2} \cot(c+dx) \sqrt{i \tan(c+dx)a+a(a^3(11A+4iB)-a^3(7iA-8B)\tan(c+dx))} dx}{a} - \frac{a^2(-8B+7iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} - \frac{a(3A+2iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \\
 & \quad \frac{2a^2}{(A+iB) \cot^2(c+dx)} \\
 & \quad \frac{d\sqrt{a+ia \tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \downarrow 27
 \end{aligned}$$

3.96. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

$$\frac{\int \cot(c+dx) \sqrt{i \tan(c+dx)a+a} \left(a^3(11A+4iB) - a^3(7iA-8B) \tan(c+dx) \right) dx}{2a} - \frac{a^2(-8B+7iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} - \frac{a(3A+2iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{(A+iB) \cot^2(c+dx)}{d \sqrt{a+ia \tan(c+dx)}} \quad 2a^2$$

↓ 3042

$$\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} \left(a^3(11A+4iB) - a^3(7iA-8B) \tan(c+dx) \right)}{\tan(c+dx)} dx}{2a} - \frac{a^2(-8B+7iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} - \frac{a(3A+2iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{(A+iB) \cot^2(c+dx)}{d \sqrt{a+ia \tan(c+dx)}} \quad 2a^2$$

↓ 4083

$$\frac{4a^3(B+iA) \int \sqrt{i \tan(c+dx)a+adx} + a^2(11A+4iB) \int \cot(c+dx)(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{2a} - \frac{a^2(-8B+7iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} - \frac{a(3A+2iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{(A+iB) \cot^2(c+dx)}{d \sqrt{a+ia \tan(c+dx)}} \quad 2a^2$$

↓ 3042

$$\frac{4a^3(B+iA) \int \sqrt{i \tan(c+dx)a+adx} + a^2(11A+4iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx}{2a} - \frac{a^2(-8B+7iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} - \frac{a(3A+2iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{(A+iB) \cot^2(c+dx)}{d \sqrt{a+ia \tan(c+dx)}} \quad 2a^2$$

↓ 3961

$$\frac{a^2(11A+4iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{8ia^4(B+iA) \int \frac{1}{a-ia \tan(c+dx)} d \sqrt{i \tan(c+dx)a+a}}{2a}}{2a} - \frac{a^2(-8B+7iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} - \frac{a(3A+2iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{(A+iB) \cot^2(c+dx)}{d \sqrt{a+ia \tan(c+dx)}} \quad 2a^2$$

↓ 219

3.96. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

$$\frac{a^2(11A+4iB) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{4i\sqrt{2}a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{2a} - \frac{a^2(-8B+7iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}}{2a^2} - a(3A+2B) \frac{(A+iB) \cot^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}}$$

↓ 4082

$$\frac{a^4(11A+4iB) \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx) - \frac{4i\sqrt{2}a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{2a} - \frac{a^2(-8B+7iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}}{2a^2} - a(3A+2B) \frac{(A+iB) \cot^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}}$$

↓ 73

$$\frac{2ia^3(11A+4iB) \int \frac{1}{i - \frac{i(i \tan(c+dx)a+a)}{d}} d\sqrt{i \tan(c+dx)a+a} - \frac{4i\sqrt{2}a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{2a} - \frac{a^2(-8B+7iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}}{2a^2} - a(3A+2B) \frac{(A+iB) \cot^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}}$$

↓ 221

$$\frac{2a^{7/2}(11A+4iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) - \frac{4i\sqrt{2}a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{2a} - \frac{a^2(-8B+7iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}}{2a^2} - a(3A+2B) \frac{(A+iB) \cot^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}}$$

input `Int[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `((A + I*B)*Cot[c + d*x]^2)/(d*Sqrt[a + I*a*Tan[c + d*x]]) + (-((a*(3*A + (2*I)*B)*Cot[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]])/d) - (((-2*a^(7/2)*(11*A + (4*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d) - ((4*I)*Sqrt[2]*a^(7/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d)/(2*a) - (a^2*((7*I)*A - 8*B)*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]])/d)/(2*a))/(2*a^2)`

3.96. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

3.96.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`
- rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d)), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2)), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

```
rule 4082 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 + b^2, 0] && EqQ[A*b + a*B, 0]
```

```
rule 4083 Int((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(
A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A
*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*T
an[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

3.96.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.78

method	result
derivativedivides	$2a^3 \left(-\frac{iB+A}{2a^3 \sqrt{a+ia \tan(dx+c)}} - \frac{(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{4a^{\frac{7}{2}}} \right) + \frac{\left(-\frac{iB}{2} - \frac{3A}{8}\right)(a+ia \tan(dx+c))^{\frac{3}{2}} + \left(\frac{1}{2}iaB + \frac{5}{8}aA\right)}{a^2 \tan(dx+c)^2}$
default	$2a^3 \left(-\frac{iB+A}{2a^3 \sqrt{a+ia \tan(dx+c)}} - \frac{(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{4a^{\frac{7}{2}}} \right) + \frac{\left(-\frac{iB}{2} - \frac{3A}{8}\right)(a+ia \tan(dx+c))^{\frac{3}{2}} + \left(\frac{1}{2}iaB + \frac{5}{8}aA\right)}{a^2 \tan(dx+c)^2}$

```
input int(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x,method=_RETUR
NVERBOSE)
```

$$3.96. \int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

```
output 2/d*a^3*(-1/2/a^3*(A+I*B)/(a+I*a*tan(d*x+c))^(1/2)-1/4*(A-I*B)/a^(7/2)*2^(
1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))+1/a^3*(-((-1/2*
I*B-3/8*A)*(a+I*a*tan(d*x+c))^(3/2)+(1/2*I*a*B+5/8*a*A)*(a+I*a*tan(d*x+c))
^(1/2))/a^2/tan(d*x+c)^2+1/8*(11*A+4*I*B)/a^(1/2)*arctanh((a+I*a*tan(d*x+c
))^(1/2)/a^(1/2))))
```

3.96.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 835 vs. $2(172) = 344$.

Time = 0.28 (sec) , antiderivative size = 835, normalized size of antiderivative = 3.81

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algori
thm="fricas")
```

```
output 1/16*(4*sqrt(2)*(a*d*e^(5*I*d*x + 5*I*c) - 2*a*d*e^(3*I*d*x + 3*I*c) + a*d
*e^(I*d*x + I*c))*sqrt((A^2 - 2*I*A*B - B^2)/(a*d^2))*log(-4*((-I*A - B)*a
*e^(I*d*x + I*c) + (I*a*d*e^(2*I*d*x + 2*I*c) + I*a*d)*sqrt(a/(e^(2*I*d*x
+ 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a*d^2)))*e^(-I*d*x - I*c)/(I*A
+ B)) - 4*sqrt(2)*(a*d*e^(5*I*d*x + 5*I*c) - 2*a*d*e^(3*I*d*x + 3*I*c) + a
*d*e^(I*d*x + I*c))*sqrt((A^2 - 2*I*A*B - B^2)/(a*d^2))*log(-4*((-I*A - B)
*a*e^(I*d*x + I*c) + (-I*a*d*e^(2*I*d*x + 2*I*c) - I*a*d)*sqrt(a/(e^(2*I*d
*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a*d^2)))*e^(-I*d*x - I*c)/(I
*A + B)) + (a*d*e^(5*I*d*x + 5*I*c) - 2*a*d*e^(3*I*d*x + 3*I*c) + a*d*e^(I
*d*x + I*c))*sqrt((121*A^2 + 88*I*A*B - 16*B^2)/(a*d^2))*log(-16*(3*(11*I*
A - 4*B)*a^2*e^(2*I*d*x + 2*I*c) + (11*I*A - 4*B)*a^2 + 2*sqrt(2)*(I*a^2*d
*e^(3*I*d*x + 3*I*c) + I*a^2*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c
) + 1))*sqrt((121*A^2 + 88*I*A*B - 16*B^2)/(a*d^2)))*e^(-2*I*d*x - 2*I*c)/
(-11*I*A + 4*B)) - (a*d*e^(5*I*d*x + 5*I*c) - 2*a*d*e^(3*I*d*x + 3*I*c) +
a*d*e^(I*d*x + I*c))*sqrt((121*A^2 + 88*I*A*B - 16*B^2)/(a*d^2))*log(-16*(
3*(11*I*A - 4*B)*a^2*e^(2*I*d*x + 2*I*c) + (11*I*A - 4*B)*a^2 + 2*sqrt(2)*
(-I*a^2*d*e^(3*I*d*x + 3*I*c) - I*a^2*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*
x + 2*I*c) + 1))*sqrt((121*A^2 + 88*I*A*B - 16*B^2)/(a*d^2)))*e^(-2*I*d*x
- 2*I*c)/(-11*I*A + 4*B)) - 4*sqrt(2)*(3*(A + 2*I*B)*e^(6*I*d*x + 6*I*c) -
2*(3*A + I*B)*e^(4*I*d*x + 4*I*c) - (7*A + 6*I*B)*e^(2*I*d*x + 2*I*c) ...
```

3.96.6 Sympy [F]

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \int \frac{(A+B \tan(c+dx)) \cot^3(c+dx)}{\sqrt{ia(\tan(c+dx)-i)}} dx$$

input `integrate(cot(d*x+c)**3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**3/sqrt(I*a*(tan(c + d*x) - I)), x)`

3.96.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.06

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx =$$

$$a^2 \left(\frac{2 \left((ia \tan(dx+c)+a)^2 (7A+8iB) - (ia \tan(dx+c)+a) (13A+12iB)a + 4(A+iB)a^2 \right)}{(ia \tan(dx+c)+a)^{\frac{5}{2}} a^2 - 2(ia \tan(dx+c)+a)^{\frac{3}{2}} a^3 + \sqrt{ia \tan(dx+c)+a} a a^4} - \frac{2\sqrt{2}(A-iB) \log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia \tan(dx+c)+a}}\right)}{a^{\frac{5}{2}}} \right) \frac{1}{8d}$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `-1/8*a^2*(2*((I*a*tan(d*x + c) + a)^2*(7*A + 8*I*B) - (I*a*tan(d*x + c) + a)*(13*A + 12*I*B)*a + 4*(A + I*B)*a^2)/((I*a*tan(d*x + c) + a)^(5/2)*a^2 - 2*(I*a*tan(d*x + c) + a)^(3/2)*a^3 + sqrt(I*a*tan(d*x + c) + a)*a^4) - 2*sqrt(2)*(A - I*B)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a)))/a^(5/2) + (11*A + 4*I*B)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a)))/a^(5/2))/d`

3.96.8 Giac [F]

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^3}{\sqrt{ia\tan(dx+c)+a}} dx$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x+c)+A)*cot(d*x+c)^3/sqrt(I*a*tan(d*x+c)+a),x)`

3.96.9 Mupad [B] (verification not implemented)

Time = 9.58 (sec) , antiderivative size = 3037, normalized size of antiderivative = 13.87

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \text{Too large to display}$$

input `int((cot(c+d*x)^3*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*1i)^(1/2),x)`

output

$$\begin{aligned}
& 2*\operatorname{atanh}((12*d^4*(a + a*\tan(c + d*x)*i)^{(1/2))*((129*A^2)/(128*a*d^2) - ((1 \\
& 2769*A^4*a^{10})/(4*d^4) + (16*B^4*a^{10})/d^4 - (3156*A^2*B^2*a^{10})/d^4 - (A* \\
& B^3*a^{10}*416i)/d^4 + (A^3*B*a^{10}*5876i)/d^4)^{(1/2)})/(64*a^6) - (3*B^2)/(16* \\
& a*d^2) + (A*B*9i)/(16*a*d^2))^{(1/2))*((12769*A^4*a^{10})/(4*d^4) + (16*B^4*a^{10} \\
& 10)/d^4 - (3156*A^2*B^2*a^{10})/d^4 - (A*B^3*a^{10}*416i)/d^4 + (A^3*B*a^{10}*58 \\
& 76i)/d^4)^{(1/2)})/(B^3*a^5*d*8i - (1469*A^3*a^5*d)/2 + 9*A*d^3*((12769*A^4* \\
& a^{10})/(4*d^4) + (16*B^4*a^{10})/d^4 - (3156*A^2*B^2*a^{10})/d^4 - (A*B^3*a^{10}* \\
& 416i)/d^4 + (A^3*B*a^{10}*5876i)/d^4)^{(1/2)} + B*d^3*((12769*A^4*a^{10})/(4*d^4 \\
&) + (16*B^4*a^{10})/d^4 - (3156*A^2*B^2*a^{10})/d^4 - (A*B^3*a^{10}*416i)/d^4 + \\
& (A^3*B*a^{10}*5876i)/d^4)^{(1/2)}*6i + 156*A*B^2*a^5*d - A^2*B*a^5*d*789i) - (\\
& 226*A^2*a^2*d^2*(a + a*\tan(c + d*x)*i)^{(1/2))*((129*A^2)/(128*a*d^2) - ((1 \\
& 2769*A^4*a^{10})/(4*d^4) + (16*B^4*a^{10})/d^4 - (3156*A^2*B^2*a^{10})/d^4 - (A* \\
& B^3*a^{10}*416i)/d^4 + (A^3*B*a^{10}*5876i)/d^4)^{(1/2)})/(64*a^6) - (3*B^2)/(16* \\
& a*d^2) + (A*B*9i)/(16*a*d^2))^{(1/2)})/(B^3*a^2*d*8i - (1469*A^3*a^2*d)/2 + \\
& 156*A*B^2*a^2*d - A^2*B*a^2*d*789i + (9*A*d^3*((12769*A^4*a^{10})/(4*d^4) + \\
& (16*B^4*a^{10})/d^4 - (3156*A^2*B^2*a^{10})/d^4 - (A*B^3*a^{10}*416i)/d^4 + (A^3 \\
& *B*a^{10}*5876i)/d^4)^{(1/2)})/a^3 + (B*d^3*((12769*A^4*a^{10})/(4*d^4) + (16*B^ \\
& 4*a^{10})/d^4 - (3156*A^2*B^2*a^{10})/d^4 - (A*B^3*a^{10}*416i)/d^4 + (A^3*B*a^1 \\
& 0*5876i)/d^4)^{(1/2)}*6i)/a^3) + (16*B^2*a^2*d^2*(a + a*\tan(c + d*x)*i)^{(1/ \\
& 2))*((129*A^2)/(128*a*d^2) - ((12769*A^4*a^{10})/(4*d^4) + (16*B^4*a^{10})/d...
\end{aligned}$$

3.97 $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

3.97.1	Optimal result	1068
3.97.2	Mathematica [A] (verified)	1068
3.97.3	Rubi [A] (verified)	1069
3.97.4	Maple [A] (verified)	1072
3.97.5	Fricas [B] (verification not implemented)	1073
3.97.6	Sympy [F]	1074
3.97.7	Maxima [A] (verification not implemented)	1074
3.97.8	Giac [F]	1074
3.97.9	Mupad [B] (verification not implemented)	1075

3.97.1 Optimal result

Integrand size = 36, antiderivative size = 209

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{(iA-B)\tan^3(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} + \frac{(3A+5iB)\tan^2(c+dx)}{2ad\sqrt{a+ia \tan(c+dx)}} - \frac{2(3A+5iB)\sqrt{a+ia \tan(c+dx)}}{a^2d} + \frac{(11A+21iB)(a+ia \tan(c+dx))^{3/2}}{6a^3d}$$

```
output 1/4*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(3/2)/
d*2^(1/2)-2*(3*A+5*I*B)*(a+I*a*tan(d*x+c))^(1/2)/a^2/d+1/2*(3*A+5*I*B)*tan
(d*x+c)^2/a/d/(a+I*a*tan(d*x+c))^(1/2)+1/3*(I*A-B)*tan(d*x+c)^3/d/(a+I*a*t
an(d*x+c))^(3/2)+1/6*(11*A+21*I*B)*(a+I*a*tan(d*x+c))^(3/2)/a^3/d
```

3.97.2 Mathematica [A] (verified)

Time = 2.94 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.71

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{25iA-39B-3(13A+19iB)\tan(c+dx)+12(-iA+B)\tan^2(c+dx)-4iB \tan^3(c+dx)}{6ad(-i+\tan(c+dx))\sqrt{a+ia \tan(c+dx)}}$$

input `Integrate[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `((A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/(2*Sqrt[2]*a^(3/2)*d) + ((25*I)*A - 39*B - 3*(13*A + (19*I)*B)*Tan[c + d*x] + 12*(-I)*A + B)*Tan[c + d*x]^2 - (4*I)*B*Tan[c + d*x]^3)/(6*a*d*(-I + Tan[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]])`

3.97.3 Rubi [A] (verified)

Time = 1.11 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.05, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.361$, Rules used = {3042, 4078, 27, 3042, 4078, 27, 3042, 4075, 3042, 4010, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^3(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+iA) \tan^3(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{3 \tan^2(c+dx)(2a(iA-B)+a(A+3iB) \tan(c+dx))}{2\sqrt{i \tan(c+dx)a+a}} dx}{3a^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B+iA) \tan^3(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{\tan^2(c+dx)(2a(iA-B)+a(A+3iB) \tan(c+dx))}{\sqrt{i \tan(c+dx)a+a}} dx}{2a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B+iA) \tan^3(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{\tan(c+dx)^2(2a(iA-B)+a(A+3iB) \tan(c+dx))}{\sqrt{i \tan(c+dx)a+a}} dx}{2a^2} \\
 & \quad \downarrow \text{4078}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{(-B + iA) \tan^3(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\int -\frac{1}{2} \tan(c+dx) \sqrt{i \tan(c+dx) a + a(4a^2(3A+5iB) - a^2(11iA-21B) \tan(c+dx)) dx}}{a^2} - \frac{a(3A+5iB) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \frac{2a^2}{} \downarrow 27 \\
 & \frac{(-B + iA) \tan^3(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\int \tan(c+dx) \sqrt{i \tan(c+dx) a + a(4a^2(3A+5iB) - a^2(11iA-21B) \tan(c+dx)) dx}}{2a^2} - \frac{a(3A+5iB) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \frac{2a^2}{} \downarrow 3042 \\
 & \frac{(-B + iA) \tan^3(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\int \tan(c+dx) \sqrt{i \tan(c+dx) a + a(4a^2(3A+5iB) - a^2(11iA-21B) \tan(c+dx)) dx}}{2a^2} - \frac{a(3A+5iB) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \frac{2a^2}{} \downarrow 4075 \\
 & \frac{(-B + iA) \tan^3(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\int \sqrt{i \tan(c+dx) a + a((11iA-21B)a^2 + 4(3A+5iB) \tan(c+dx)a^2) dx - \frac{2a(11A+21iB)(a+ia \tan(c+dx))^{3/2}}{3d}}{2a^2} - \frac{a(3A+5iB) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \frac{2a^2}{} \downarrow 3042 \\
 & \frac{(-B + iA) \tan^3(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\int \sqrt{i \tan(c+dx) a + a((11iA-21B)a^2 + 4(3A+5iB) \tan(c+dx)a^2) dx - \frac{2a(11A+21iB)(a+ia \tan(c+dx))^{3/2}}{3d}}{2a^2} - \frac{a(3A+5iB) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \frac{2a^2}{} \downarrow 4010 \\
 & \frac{(-B + iA) \tan^3(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{-\left(a^2(B+iA) \int \sqrt{i \tan(c+dx) a + a dx}\right) + \frac{8a^2(3A+5iB) \sqrt{a+ia \tan(c+dx)}}{d} - \frac{2a(11A+21iB)(a+ia \tan(c+dx))^{3/2}}{3d}}{2a^2} - \frac{a(3A+5iB) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \frac{2a^2}{} \downarrow 3042 \\
 & \frac{(-B + iA) \tan^3(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{-\left(a^2(B+iA) \int \sqrt{i \tan(c+dx) a + a dx}\right) + \frac{8a^2(3A+5iB) \sqrt{a+ia \tan(c+dx)}}{d} - \frac{2a(11A+21iB)(a+ia \tan(c+dx))^{3/2}}{3d}}{2a^2} - \frac{a(3A+5iB) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \frac{2a^2}{}
 \end{aligned}$$

3.97. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

$$\begin{array}{c}
 \downarrow \text{3961} \\
 \frac{(-B + iA) \tan^3(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \\
 \frac{2ia^3(B + ia) \int \frac{1}{a - ia \tan(c + dx)} d\sqrt{i \tan(c + dx)a + a} + \frac{8a^2(3A + 5iB)\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2a(11A + 21iB)(a + ia \tan(c + dx))^{3/2}}{3d}}{2a^2} - \frac{a(3A + 5iB) \tan^2(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} \\
 \hline
 2a^2 \\
 \downarrow \text{219} \\
 \frac{(-B + iA) \tan^3(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \\
 \frac{i\sqrt{2}a^{5/2}(B + ia) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right) + \frac{8a^2(3A + 5iB)\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2a(11A + 21iB)(a + ia \tan(c + dx))^{3/2}}{3d}}{2a^2} - \frac{a(3A + 5iB) \tan^2(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} \\
 \hline
 2a^2
 \end{array}$$

input `Int[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `((I*A - B)*Tan[c + d*x]^3)/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) - (-((a*(3*A + (5*I)*B)*Tan[c + d*x]^2)/(d*Sqrt[a + I*a*Tan[c + d*x]])) + ((I*Sqrt[2]*a^(5/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d + (8*a^2*(3*A + (5*I)*B)*Sqrt[a + I*a*Tan[c + d*x])/d - (2*a*(11*A + (21*I)*B)*(a + I*a*Tan[c + d*x])^(3/2))/(3*d))/(2*a^2))/(2*a^2)`

3.97.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F x_), x_Symbol] := Simp[a Int[F x, x], x] /; FreeQ[a, x] && !MatchQ[F x, (b_)*(G x_) /; FreeQ[b, x]]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 3961 Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d)
  Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a
  , b, c, d}, x] && EqQ[a^2 + b^2, 0]
```

```
rule 4010 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
  (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp
  [(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e
  , f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]
```

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
  + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
  *d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
  x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
  d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

```
rule 4078 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
  p[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
  x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
  x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
  *A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
  && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.97.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.74

method	result
derivativedivides	$\frac{2iB(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - 4iaB\sqrt{a+ia \tan(dx+c)} - 2aA\sqrt{a+ia \tan(dx+c)} - \frac{a^2(7iB+5A)}{2\sqrt{a+ia \tan(dx+c)}} + \frac{a^3(iB+A)}{3(a+ia \tan(dx+c))^{\frac{3}{2}}} + \frac{a^{\frac{3}{2}}(-)}{a^3d}$
default	$\frac{2iB(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} - 4iaB\sqrt{a+ia \tan(dx+c)} - 2aA\sqrt{a+ia \tan(dx+c)} - \frac{a^2(7iB+5A)}{2\sqrt{a+ia \tan(dx+c)}} + \frac{a^3(iB+A)}{3(a+ia \tan(dx+c))^{\frac{3}{2}}} + \frac{a^{\frac{3}{2}}(-)}{a^3d}$
parts	$\frac{2A \left(-\sqrt{a+ia \tan(dx+c)} + \frac{\sqrt{a} \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}}\right)}{8} - \frac{5a}{4\sqrt{a+ia \tan(dx+c)}} + \frac{a^2}{6(a+ia \tan(dx+c))^{\frac{3}{2}}} \right)}{da^2} + \frac{2iB \left(a \right)}{+}$

3.97. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

```
input int(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x,method=_RETURNVERBOSE)
```

```
output 2/d/a^3*(1/3*I*B*(a+I*a*tan(d*x+c))^(3/2)-2*I*a*B*(a+I*a*tan(d*x+c))^(1/2)-a*A*(a+I*a*tan(d*x+c))^(1/2)-1/4*a^2*(5*A+7*I*B)/(a+I*a*tan(d*x+c))^(1/2)+1/6*a^3*(A+I*B)/(a+I*a*tan(d*x+c))^(3/2)+1/8*a^(3/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2)))
```

3.97.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 440 vs. $2(162) = 324$.

Time = 0.28 (sec) , antiderivative size = 440, normalized size of antiderivative = 2.11

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx =$$

$$3 \sqrt{\frac{1}{2}} (a^2 de^{(5i dx+5i c)} + a^2 de^{(3i dx+3i c)}) \sqrt{\frac{A^2-2i AB-B^2}{a^3 d^2}} \log \left(-\frac{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (i a^2 de^{(2i dx+2i c)} + i a^2 d) \sqrt{\frac{a}{e^{(2i dx+2i c)}+1}} \sqrt{\frac{A^2-2i AB-B^2}{a^3 d^2}} \right)}{i A+B} \right)$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="fracas")
```

```
output -1/12*(3*sqrt(1/2)*(a^2*d*e^(5*I*d*x + 5*I*c) + a^2*d*e^(3*I*d*x + 3*I*c))*sqrt((A^2 - 2*I*A*B - B^2)/(a^3*d^2))*log(-4*(sqrt(2)*sqrt(1/2)*(I*a^2*d*e^(2*I*d*x + 2*I*c) + I*a^2*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a^3*d^2)) + (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B) - 3*sqrt(1/2)*(a^2*d*e^(5*I*d*x + 5*I*c) + a^2*d*e^(3*I*d*x + 3*I*c))*sqrt((A^2 - 2*I*A*B - B^2)/(a^3*d^2))*log(-4*(sqrt(2)*sqrt(1/2)*(-I*a^2*d*e^(2*I*d*x + 2*I*c) - I*a^2*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a^3*d^2)) + (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) + sqrt(2)*(2*(19*A + 26*I*B)*e^(6*I*d*x + 6*I*c) + 3*(17*A + 29*I*B)*e^(4*I*d*x + 4*I*c) + 6*(2*A + 3*I*B)*e^(2*I*d*x + 2*I*c) - A - I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(a^2*d*e^(5*I*d*x + 5*I*c) + a^2*d*e^(3*I*d*x + 3*I*c))
```


3.97.6 Sympy [F]

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \int \frac{(A+B\tan(c+dx))\tan^3(c+dx)}{(ia(\tan(c+dx)-i))^{\frac{3}{2}}} dx$$

input `integrate(tan(d*x+c)**3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**3/(I*a*(tan(c + d*x) - I))**(3/2), x)`

3.97.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.77

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \frac{3\sqrt{2}(A-iB)a^{\frac{5}{2}} \log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia\tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia\tan(dx+c)+a}}\right) - 16i(ia\tan(dx+c)+a)^{\frac{3}{2}}Ba + 48\sqrt{ia\tan(dx+c)+a}}{24a^4d}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `-1/24*(3*sqrt(2)*(A - I*B)*a^(5/2)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - 16*I*(I*a*tan(d*x + c) + a)^(3/2)*B*a + 48*sqrt(I*a*tan(d*x + c) + a)*(A + 2*I*B)*a^2 + 4*(3*(I*a*tan(d*x + c) + a)*(5*A + 7*I*B)*a^3 - 2*(A + I*B)*a^4)/(I*a*tan(d*x + c) + a)^(3/2))/(a^4*d)`

3.97.8 Giac [F]

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \int \frac{(B\tan(dx+c)+A)\tan^3(dx+c)}{(ia\tan(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^3/(I*a*tan(d*x + c) + a)^(3/2), x)`

3.97.9 Mupad [B] (verification not implemented)

Time = 8.81 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.11

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \frac{\frac{B \operatorname{li}}{3d} - \frac{B(a+a\tan(c+dx)\operatorname{li})7i}{2ad}}{(a+a\tan(c+dx)\operatorname{li})^{3/2}} + \frac{\frac{Aa}{3} - \frac{5A(a+a\tan(c+dx)\operatorname{li})}{2}}{ad(a+a\tan(c+dx)\operatorname{li})^{3/2}} - \frac{2A\sqrt{a+a\tan(c+dx)\operatorname{li}}}{a^2d} - \frac{B\sqrt{a+a\tan(c+dx)\operatorname{li}}4i}{a^2d} + \frac{B(a+a\tan(c+dx)\operatorname{li})^{3/2}2i}{3a^3d} - \frac{\sqrt{2}B\operatorname{atan}\left(\frac{\sqrt{2}\sqrt{a+a\tan(c+dx)\operatorname{li}}}{2\sqrt{-a}}\right)\operatorname{li}}{4(-a)^{3/2}d} + \frac{\sqrt{2}A\operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{a+a\tan(c+dx)\operatorname{li}}}{2\sqrt{a}}\right)}{4a^{3/2}d}$$

input `int((tan(c + d*x)^3*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(3/2),x)`

output `((B*1i)/(3*d) - (B*(a + a*tan(c + d*x)*1i)*7i)/(2*a*d))/(a + a*tan(c + d*x)*1i)^(3/2) + ((A*a)/3 - (5*A*(a + a*tan(c + d*x)*1i))/2)/(a*d*(a + a*tan(c + d*x)*1i)^(3/2)) - (2*A*(a + a*tan(c + d*x)*1i)^(1/2))/(a^2*d) - (B*(a + a*tan(c + d*x)*1i)^(1/2)*4i)/(a^2*d) + (B*(a + a*tan(c + d*x)*1i)^(3/2)*2i)/(3*a^3*d) - (2^(1/2)*B*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2))))*1i)/(4*(-a)^(3/2)*d) + (2^(1/2)*A*atanh((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*a^(1/2))))/(4*a^(3/2)*d)`

3.98
$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$$

3.98.1	Optimal result	1076
3.98.2	Mathematica [A] (verified)	1076
3.98.3	Rubi [A] (verified)	1077
3.98.4	Maple [A] (verified)	1080
3.98.5	Fricas [B] (verification not implemented)	1080
3.98.6	Sympy [F]	1081
3.98.7	Maxima [A] (verification not implemented)	1081
3.98.8	Giac [F]	1082
3.98.9	Mupad [B] (verification not implemented)	1082

3.98.1 Optimal result

Integrand size = 36, antiderivative size = 167

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{(iA-B) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} + \frac{5iA-11B}{6ad\sqrt{a+ia \tan(c+dx)}} + \frac{(iA-7B)\sqrt{a+ia \tan(c+dx)}}{3a^2d}$$

```
output 1/4*(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(3/2)/
d*2^(1/2)+1/6*(5*I*A-11*B)/a/d/(a+I*a*tan(d*x+c))^(1/2)+1/3*(I*A-7*B)*(a+I
*a*tan(d*x+c))^(1/2)/a^2/d+1/3*(I*A-B)*tan(d*x+c)^2/d/(a+I*a*tan(d*x+c))^(
3/2)
```

3.98.2 Mathematica [A] (verified)

Time = 1.72 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.89

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{2B \tan^2(c+dx)}{d(a+ia \tan(c+dx))^{3/2}} + \frac{i \left(\frac{3\sqrt{2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{a}} - \frac{4a(A+7iB)}{(a+ia \tan(c+dx))^{3/2}} + \frac{6(3A+13iB)}{\sqrt{a+ia \tan(c+dx)}} \right)}{12ad}$$

input `Integrate[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `(2*B*Tan[c + d*x]^2)/(d*(a + I*a*Tan[c + d*x])^(3/2)) + ((I/12)*((3*sqrt[2] *(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(sqrt[2]*sqrt[a])]))/sqrt[a] - (4*a*(A + (7*I)*B))/(a + I*a*Tan[c + d*x])^(3/2) + (6*(3*A + (13*I)*B) /sqrt[a + I*a*Tan[c + d*x]])))/(a*d)`

3.98.3 Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.99, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {3042, 4078, 27, 3042, 4075, 3042, 4009, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^2(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+ia) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{\tan(c+dx)(4a(iA-B)+a(A+7iB) \tan(c+dx))}{2\sqrt{i \tan(c+dx)a+a}} dx}{3a^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B+ia) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{\tan(c+dx)(4a(iA-B)+a(A+7iB) \tan(c+dx))}{\sqrt{i \tan(c+dx)a+a}} dx}{6a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B+ia) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{\tan(c+dx)(4a(iA-B)+a(A+7iB) \tan(c+dx))}{\sqrt{i \tan(c+dx)a+a}} dx}{6a^2} \\
 & \quad \downarrow \text{4075} \\
 & \frac{(-B+ia) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{4a(iA-B) \tan(c+dx)-a(A+7iB)}{\sqrt{i \tan(c+dx)a+a}} dx - \frac{2(-7B+ia)\sqrt{a+ia \tan(c+dx)}}{d}}{6a^2}
 \end{aligned}$$

3.98. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^2(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\int \frac{4a(iA - B) \tan(c + dx) - a(A + 7iB)}{\sqrt{i \tan(c + dx)a + a}} dx - \frac{2(-7B + iA) \sqrt{a + ia \tan(c + dx)}}{d}}{6a^2} \\
& \downarrow 4009 \\
& \frac{(-B + iA) \tan^2(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\frac{3}{2}(A - iB) \int \sqrt{i \tan(c + dx)a + a} dx - \frac{a(-11B + 5iA)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2(-7B + iA) \sqrt{a + ia \tan(c + dx)}}{d}}{6a^2} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^2(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\frac{3}{2}(A - iB) \int \sqrt{i \tan(c + dx)a + a} dx - \frac{a(-11B + 5iA)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2(-7B + iA) \sqrt{a + ia \tan(c + dx)}}{d}}{6a^2} \\
& \downarrow 3961 \\
& \frac{(-B + iA) \tan^2(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\frac{3ia(A - iB) \int \frac{1}{a - ia \tan(c + dx)} d\sqrt{i \tan(c + dx)a + a} - \frac{a(-11B + 5iA)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2(-7B + iA) \sqrt{a + ia \tan(c + dx)}}{d}}{6a^2} \\
& \downarrow 219 \\
& \frac{(-B + iA) \tan^2(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\frac{3i\sqrt{a}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}d} - \frac{a(-11B + 5iA)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2(-7B + iA) \sqrt{a + ia \tan(c + dx)}}{d}}{6a^2}
\end{aligned}$$

input `Int[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `((I*A - B)*Tan[c + d*x]^2)/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) - (((-3*I)*Sqrt[a]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/(Sqrt[2]*d) - (a*((5*I)*A - 11*B))/(d*Sqrt[a + I*a*Tan[c + d*x]]) - (2*(I*A - 7*B)*Sqrt[a + I*a*Tan[c + d*x]])/d)/(6*a^2)`

3.98.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3961 `Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`
- rule 4009 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`
- rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`
- rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.98.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.69

method	result
derivativedivides	$\frac{2i \left(iB \sqrt{a+ia \tan(dx+c)} + \frac{a(5iB+3A)}{4\sqrt{a+ia \tan(dx+c)}} - \frac{a^2(iB+A)}{6(a+ia \tan(dx+c))^{\frac{3}{2}}} + \frac{\sqrt{a}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{8} \right)}{da^2}$
default	$\frac{2i \left(iB \sqrt{a+ia \tan(dx+c)} + \frac{a(5iB+3A)}{4\sqrt{a+ia \tan(dx+c)}} - \frac{a^2(iB+A)}{6(a+ia \tan(dx+c))^{\frac{3}{2}}} + \frac{\sqrt{a}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{8} \right)}{da^2}$
parts	$\frac{2iA \left(\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{8\sqrt{a}} + \frac{3}{4\sqrt{a+ia \tan(dx+c)}} - \frac{a}{6(a+ia \tan(dx+c))^{\frac{3}{2}}} \right)}{da} + \frac{2B \left(-\sqrt{a+ia \tan(dx+c)} + \frac{\sqrt{a}\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2\sqrt{a}} \right)}{da}$

input `int(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `2*I/d/a^2*(I*B*(a+I*a*tan(d*x+c))^(1/2)+1/4*a*(3*A+5*I*B)/(a+I*a*tan(d*x+c))^(1/2)-1/6*a^2*(A+I*B)/(a+I*a*tan(d*x+c))^(3/2)+1/8*a^(1/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2)))`

3.98.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 375 vs. 2(128) = 256.

Time = 0.25 (sec) , antiderivative size = 375, normalized size of antiderivative = 2.25

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{\left(3 \sqrt{\frac{1}{2}} a^2 d \sqrt{-\frac{A^2-2iAB-B^2}{a^3 d^2}} e^{(3i dx+3i c)} \log \left(-\frac{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (a^2 d e^{(2i dx+2i c)} + a^2 d) \sqrt{\frac{a}{e^{(2i dx+2i c)}+1}} \sqrt{-\frac{A^2-2iAB-B^2}{a^3 d^2}} + (-iA-B) a e^{(2i dx+2i c)}}{iA+B} \right) \right)}{da^2}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="fracas")`

output
$$-1/12*(3*\sqrt{1/2}*a^2*d*\sqrt{-(A^2 - 2*I*A*B - B^2)/(a^3*d^2)})*e^{(3*I*d*x + 3*I*c)}*\log(-4*(\sqrt{2}*\sqrt{1/2}*(a^2*d*e^{(2*I*d*x + 2*I*c)} + a^2*d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{-(A^2 - 2*I*A*B - B^2)/(a^3*d^2)}) + (-I*A - B)*a*e^{(I*d*x + I*c)})*e^{(-I*d*x - I*c)/(I*A + B)} - 3*\sqrt{1/2}*a^2*d*\sqrt{-(A^2 - 2*I*A*B - B^2)/(a^3*d^2)})*e^{(3*I*d*x + 3*I*c)}*\log(4*(\sqrt{2}*\sqrt{1/2}*(a^2*d*e^{(2*I*d*x + 2*I*c)} + a^2*d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{-(A^2 - 2*I*A*B - B^2)/(a^3*d^2)}) - (-I*A - B)*a*e^{(I*d*x + I*c)})*e^{(-I*d*x - I*c)/(I*A + B)} + \sqrt{2}*(2*(-4*I*A + 19*B)*e^{(4*I*d*x + 4*I*c)} - (7*I*A - 13*B)*e^{(2*I*d*x + 2*I*c)} + I*A - B)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*e^{(-3*I*d*x - 3*I*c)/(a^2*d)}$$

3.98.6 Sympy [F]

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \int \frac{(A+B\tan(c+dx))\tan^2(c+dx)}{(ia(\tan(c+dx)-i))^{3/2}} dx$$

input `integrate(tan(d*x+c)**2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**2/(I*a*(tan(c + d*x) - I))**(3/2), x)`

3.98.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.82

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \frac{i \left(3\sqrt{2}(A-iB)a^{\frac{3}{2}} \log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia\tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia\tan(dx+c)+a}}\right) - 48i\sqrt{ia\tan(dx+c)+a}Ba - \frac{4(3(ia\tan(dx+c)+a)(3A+5iB))}{(ia\tan(dx+c)-i)} \right)}{24a^3d}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output
$$-1/24*I*(3*\sqrt{2}*(A - I*B)*a^{(3/2)}*\log(-(\sqrt{2}*\sqrt{a} - \sqrt{I*a*\tan(d*x + c) + a})/(\sqrt{2}*\sqrt{a} + \sqrt{I*a*\tan(d*x + c) + a}))) - 48*I*\sqrt{I*a*\tan(d*x + c) + a}*B*a - 4*(3*(I*a*\tan(d*x + c) + a)*(3*A + 5*I*B)*a^2 - 2*(A + I*B)*a^3)/(I*a*\tan(d*x + c) + a)^{(3/2)}/(a^3*d)$$

3.98.
$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx$$

3.98.8 Giac [F]

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \int \frac{(B \tan(dx+c)+A) \tan(dx+c)^2}{(ia \tan(dx+c)+a)^{3/2}} dx$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(d*x+c)+A)*tan(d*x+c)^2/(I*a*tan(d*x+c)+a)^(3/2),x)`

3.98.9 Mupad [B] (verification not implemented)

Time = 0.74 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.11

$$\begin{aligned} \int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx &= -\frac{\frac{A \operatorname{li}}{3d} - \frac{A(a+a \tan(c+dx) \operatorname{li}) 3i}{2ad}}{(a+a \tan(c+dx) \operatorname{li})^{3/2}} \\ &+ \frac{\frac{Ba}{3} - \frac{5B(a+a \tan(c+dx) \operatorname{li})}{2}}{ad(a+a \tan(c+dx) \operatorname{li})^{3/2}} - \frac{2B \sqrt{a+a \tan(c+dx) \operatorname{li}}}{a^2 d} \\ &+ \frac{\sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a+a \tan(c+dx) \operatorname{li}}}{2\sqrt{-a}}\right) \operatorname{li}}{4(-a)^{3/2} d} + \frac{\sqrt{2} B \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{a+a \tan(c+dx) \operatorname{li}}}{2\sqrt{-a}}\right)}{4a^{3/2} d} \end{aligned}$$

input `int((tan(c+d*x)^2*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*1i)^(3/2),x)`

output `((B*a)/3 - (5*B*(a+a*tan(c+d*x)*1i))/2)/(a*d*(a+a*tan(c+d*x)*1i)^(3/2)) - ((A*1i)/(3*d) - (A*(a+a*tan(c+d*x)*1i)*3i)/(2*a*d))/(a+a*tan(c+d*x)*1i)^(3/2) - (2*B*(a+a*tan(c+d*x)*1i)^(1/2))/(a^2*d) + (2^(1/2)*A*atan((2^(1/2)*(a+a*tan(c+d*x)*1i)^(1/2))/(2*(-a)^(1/2)))*1i)/(4*(-a)^(3/2)*d) + (2^(1/2)*B*atanh((2^(1/2)*(a+a*tan(c+d*x)*1i)^(1/2))/(2*a^(1/2))))/(4*a^(3/2)*d)`

3.99 $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

3.99.1 Optimal result 1083
 3.99.2 Mathematica [A] (verified) 1083
 3.99.3 Rubi [A] (verified) 1084
 3.99.4 Maple [A] (verified) 1086
 3.99.5 Fricas [B] (verification not implemented) 1086
 3.99.6 Sympy [F] 1087
 3.99.7 Maxima [A] (verification not implemented) 1087
 3.99.8 Giac [F] 1088
 3.99.9 Mupad [B] (verification not implemented) 1088

3.99.1 Optimal result

Integrand size = 34, antiderivative size = 119

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = -\frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{2\sqrt{2}a^{3/2}d} - \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}} + \frac{A+3iB}{2ad\sqrt{a+ia \tan(c+dx)}}$$

output

```
-1/4*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(3/2)
/d*2^(1/2)+1/2*(A+3*I*B)/a/d/(a+I*a*tan(d*x+c))^(1/2)+1/3*(-A-I*B)/d/(a+I*
a*tan(d*x+c))^(3/2)
```

3.99.2 Mathematica [A] (verified)

Time = 1.37 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.99

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{i\left(\frac{3\sqrt{2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{a}} + \frac{4a(iA-B)}{(a+ia \tan(c+dx))^{3/2}} - \frac{6(iA-3B)}{\sqrt{a+ia \tan(c+dx)}}\right)}{12ad}$$

input

```
Integrate[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2)
,x]
```

output $((I/12)*((3*sqrt[2]*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(sqrt[2]*sqrt[a])])/sqrt[a] + (4*a*(I*A - B))/(a + I*a*Tan[c + d*x])^(3/2) - (6*(I*A - 3*B))/sqrt[a + I*a*Tan[c + d*x]]))/(a*d)$

3.99.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.06, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.206$, Rules used = {3042, 4073, 3042, 4009, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$$

↓ 3042

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$$

↓ 4073

$$-\frac{i \int \frac{a(A+iB)+2aB \tan(c+dx)}{\sqrt{i \tan(c+dx)a+a}} dx}{2a^2} - \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 3042

$$-\frac{i \int \frac{a(A+iB)+2aB \tan(c+dx)}{\sqrt{i \tan(c+dx)a+a}} dx}{2a^2} - \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 4009

$$-\frac{i \left(\frac{1}{2}(A-iB) \int \sqrt{i \tan(c+dx)a+adx} + \frac{a(-3B+iA)}{d\sqrt{a+ia \tan(c+dx)}} \right)}{2a^2} - \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 3042

$$-\frac{i \left(\frac{1}{2}(A-iB) \int \sqrt{i \tan(c+dx)a+adx} + \frac{a(-3B+iA)}{d\sqrt{a+ia \tan(c+dx)}} \right)}{2a^2} - \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 3961

$$-\frac{i \left(\frac{a(-3B+iA)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{ia(A-iB) \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a}}{d} \right)}{2a^2} - \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}}$$

3.99. $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

$$i \left(\frac{a(-3B+iA)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{i\sqrt{a}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}d} \right) - \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}}$$

input `Int[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `-1/3*(A + I*B)/(d*(a + I*a*Tan[c + d*x])^(3/2)) - ((I/2)*((-I)*Sqrt[a]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/(Sqrt[2]*d + (a*(I*A - 3*B))/(d*Sqrt[a + I*a*Tan[c + d*x]])))/a^2`

3.99.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

```
rule 4073 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-
A*b - a*B)*(a*c + b*d)*((a + b*Tan[e + f*x])^m/(2*a^2*f*m)), x] + Simp[1/(
2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[A*b*c + a*B*c + a*A*d + b*B*
d + 2*a*B*d*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && LtQ[m, -1] && EqQ[a^2 + b^2, 0]
```

3.99.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.81

method	result
derivativedivides	$\frac{2\left(-\frac{A}{4} - \frac{3iB}{4}\right)}{\sqrt{a+ia \tan(dx+c)}} - \frac{a(iB+A)}{3(a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{\left(\frac{A}{4} - \frac{iB}{4}\right)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{\sqrt{a}}$
default	$\frac{2\left(-\frac{A}{4} - \frac{3iB}{4}\right)}{\sqrt{a+ia \tan(dx+c)}} - \frac{a(iB+A)}{3(a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{\left(\frac{A}{4} - \frac{iB}{4}\right)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{\sqrt{a}}$
parts	$\frac{A\left(-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{4a^{\frac{3}{2}}} - \frac{1}{3(a+ia \tan(dx+c))^{\frac{3}{2}}} + \frac{1}{2a\sqrt{a+ia \tan(dx+c)}}\right)}{d} + \frac{2iB\left(\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{8\sqrt{a}}\right)}{d}$

```
input int(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output 2/d/a*(-(-1/4*A-3/4*I*B)/(a+I*a*tan(d*x+c))^(1/2)-1/6*a*(A+I*B)/(a+I*a*tan
(d*x+c))^(3/2)-1/2*(1/4*A-1/4*I*B)*2^(1/2)/a^(1/2)*arctanh(1/2*(a+I*a*tan
(d*x+c))^(1/2)*2^(1/2)/a^(1/2)))
```

3.99.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 369 vs. 2(88) = 176.

Time = 0.25 (sec) , antiderivative size = 369, normalized size of antiderivative = 3.10

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{\left(3\sqrt{\frac{1}{2}}a^2d\sqrt{\frac{A^2-2iAB-B^2}{a^3d^2}}e^{(3i dx+3i c)} \log\left(-\frac{4\left(\sqrt{2}\sqrt{\frac{1}{2}}(ia^2de^{(2i dx+2i c)}+i\right)}{2\sqrt{a}}\right)}{d}\right)}{d}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="fricas")`

output `1/12*(3*sqrt(1/2)*a^2*d*sqrt((A^2 - 2*I*A*B - B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(I*a^2*d*e^(2*I*d*x + 2*I*c) + I*a^2*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a^3*d^2)) + (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 3*sqrt(1/2)*a^2*d*sqrt((A^2 - 2*I*A*B - B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(-I*a^2*d*e^(2*I*d*x + 2*I*c) - I*a^2*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a^3*d^2)) + (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) + sqrt(2)*(2*(A + 4*I*B)*e^(4*I*d*x + 4*I*c) + (A + 7*I*B)*e^(2*I*d*x + 2*I*c) - A - I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-3*I*d*x - 3*I*c)/(a^2*d)`

3.99.6 Sympy [F]

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \int \frac{(A+B\tan(c+dx))\tan(c+dx)}{(ia(\tan(c+dx)-i))^{3/2}} dx$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)/(I*a*(tan(c + d*x) - I))**(3/2), x)`

3.99.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.97

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \frac{3\sqrt{2}(A-iB)\sqrt{a}\log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia\tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia\tan(dx+c)+a}}\right) + \frac{4(3(ia\tan(dx+c)+a)}{(ia\tan(dx+c)+a)}}{24a^2d}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output $1/24*(3*\sqrt{2}*(A - I*B)*\sqrt{a}*\log(-(\sqrt{2}*\sqrt{a} - \sqrt{I*a*\tan(dx + c) + a))/(\sqrt{2}*\sqrt{a} + \sqrt{I*a*\tan(dx + c) + a})) + 4*(3*(I*a*\tan(dx + c) + a)*(A + 3*I*B)*a - 2*(A + I*B)*a^2)/(I*a*\tan(dx + c) + a)^{(3/2)}/(a^2*d)$

3.99.8 Giac [F]

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)}{(ia \tan(dx + c) + a)^{3/2}} dx$$

input `integrate(tan(dx+c)*(A+B*tan(dx+c))/(a+I*a*tan(dx+c))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(dx + c) + A)*tan(dx + c)/(I*a*tan(dx + c) + a)^(3/2), x)`

3.99.9 Mupad [B] (verification not implemented)

Time = 8.11 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.37

$$\begin{aligned} & \int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{3/2}} dx = \\ & -\frac{\frac{B \operatorname{li} - \frac{B(a+a \tan(c+dx) \operatorname{li}) 3i}{2ad}}{(a + a \tan(c + dx) \operatorname{li})^{3/2}} - \frac{\frac{A}{3} - \frac{A(a+a \tan(c+dx) \operatorname{li})}{2a}}{d(a + a \tan(c + dx) \operatorname{li})^{3/2}}}{4(-a)^{3/2}d} - \frac{\sqrt{2} A \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{a+a \tan(c+dx) \operatorname{li}}}{2\sqrt{a}}\right)}{4a^{3/2}d} \end{aligned}$$

input `int((tan(c + d*x)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(3/2),x)`

output $(2^{(1/2)}*B*\operatorname{atan}((2^{(1/2)}*(a + a*\tan(c + d*x)*1i)^{(1/2)})/(2*(-a)^{(1/2)}))*1i)/(4*(-a)^{(3/2)}*d) - (A/3 - (A*(a + a*\tan(c + d*x)*1i))/(2*a))/(d*(a + a*\tan(c + d*x)*1i)^{(3/2)}) - ((B*1i)/(3*d) - (B*(a + a*\tan(c + d*x)*1i)*3i)/(2*a*d))/(a + a*\tan(c + d*x)*1i)^{(3/2)} - (2^{(1/2)}*A*\operatorname{atanh}((2^{(1/2)}*(a + a*\tan(c + d*x)*1i)^{(1/2)})/(2*a^{(1/2)})))/(4*a^{(3/2)}*d)$

3.100 $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^{3/2}} dx$

3.100.1 Optimal result 1089
 3.100.2 Mathematica [C] (verified) 1089
 3.100.3 Rubi [A] (verified) 1090
 3.100.4 Maple [A] (verified) 1092
 3.100.5 Fricas [B] (verification not implemented) 1092
 3.100.6 Sympy [F] 1093
 3.100.7 Maxima [A] (verification not implemented) 1093
 3.100.8 Giac [F] 1094
 3.100.9 Mupad [B] (verification not implemented) 1094

3.100.1 Optimal result

Integrand size = 28, antiderivative size = 121

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{3/2}} dx = -\frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{iA - B}{3d(a + ia \tan(c + dx))^{3/2}} + \frac{iA + B}{2ad\sqrt{a + ia \tan(c + dx)}}$$

output `-1/4*(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(3/2)/d*2^(1/2)+1/2*(I*A+B)/a/d/(a+I*a*tan(d*x+c))^(1/2)+1/3*(I*A-B)/d/(a+I*a*tan(d*x+c))^(3/2)`

3.100.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.62 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.63

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{3/2}} dx = \frac{2iA - 2B - 3(A - iB) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{1}{2}(1 + i \tan(c + dx))\right)}{6d(a + ia \tan(c + dx))^{3/2}}$$

input `Integrate[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `((2*I)*A - 2*B - 3*(A - I*B)*Hypergeometric2F1[-1/2, 1, 1/2, (1 + I*Tan[c + d*x])/2]*(-I + Tan[c + d*x]))/(6*d*(a + I*a*Tan[c + d*x])^(3/2))`

3.100. $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^{3/2}} dx$

3.100.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4009, 3042, 3960, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{4009} \\
 & \frac{(A - iB) \int \frac{1}{\sqrt{i \tan(c+dx)a+a}} dx}{2a} + \frac{-B + iA}{3d(a + ia \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A - iB) \int \frac{1}{\sqrt{i \tan(c+dx)a+a}} dx}{2a} + \frac{-B + iA}{3d(a + ia \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3960} \\
 & \frac{(A - iB) \left(\frac{\int \sqrt{i \tan(c+dx)a+adx}}{2a} + \frac{i}{d\sqrt{a+ia \tan(c+dx)}} \right)}{2a} + \frac{-B + iA}{3d(a + ia \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A - iB) \left(\frac{\int \sqrt{i \tan(c+dx)a+adx}}{2a} + \frac{i}{d\sqrt{a+ia \tan(c+dx)}} \right)}{2a} + \frac{-B + iA}{3d(a + ia \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3961} \\
 & \frac{(A - iB) \left(\frac{i}{d\sqrt{a+ia \tan(c+dx)}} - \frac{i \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a}}{d} \right)}{2a} + \frac{-B + iA}{3d(a + ia \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{219} \\
 & \frac{(A - iB) \left(\frac{i}{d\sqrt{a+ia \tan(c+dx)}} - \frac{i \operatorname{arctanh} \left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}} \right)}{\sqrt{2}\sqrt{ad}} \right)}{2a} + \frac{-B + iA}{3d(a + ia \tan(c + dx))^{3/2}}
 \end{aligned}$$

3.100. $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^{3/2}} dx$

input `Int[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `(I*A - B)/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) + ((A - I*B)*((-I)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/(Sqrt[2]*Sqrt[a]*d) + I/(d*Sqrt[a + I*a*Tan[c + d*x]])))/(2*a)`

3.100.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

3.100.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.79

method	result
derivativedivides	$\frac{2i \left(-\frac{-\frac{A}{2} - \frac{iB}{2}}{3(a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{iB-A}{4a\sqrt{a+ia \tan(dx+c)}} - \frac{(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{8a^{\frac{3}{2}}} \right)}{d}$
default	$\frac{2i \left(-\frac{-\frac{A}{2} - \frac{iB}{2}}{3(a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{iB-A}{4a\sqrt{a+ia \tan(dx+c)}} - \frac{(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{8a^{\frac{3}{2}}} \right)}{d}$
parts	$\frac{2iAa \left(-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{8a^{\frac{5}{2}}} + \frac{1}{4a^2\sqrt{a+ia \tan(dx+c)}} + \frac{1}{6a(a+ia \tan(dx+c))^{\frac{3}{2}}} \right)}{d} + \frac{B \left(-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{4a^{\frac{3}{2}}} \right)}{d}$

input `int((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `2*I/d*(-1/3*(-1/2*A-1/2*I*B)/(a+I*a*tan(d*x+c))^(3/2)-1/4/a*(-A+I*B)/(a+I*a*tan(d*x+c))^(1/2)-1/8*(A-I*B)/a^(3/2)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2)))`

3.100.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 372 vs. 2(90) = 180.

Time = 0.26 (sec) , antiderivative size = 372, normalized size of antiderivative = 3.07

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{3/2}} dx = \frac{\left(3 \sqrt{\frac{1}{2}} a^2 d \sqrt{-\frac{A^2 - 2iAB - B^2}{a^3 d^2}} e^{(3i dx + 3i c)} \log \left(-\frac{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (a^2 d e^{(2i dx + 2i c)} + a^2 d) \sqrt{e^{(2i dx + 2i c)}} \right)}{4a^{\frac{3}{2}}} \right)}{d} \right)}{d}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="fricas")`

output $1/12*(3*\sqrt{1/2}*a^{2*d}*\sqrt{-(A^2 - 2*I*A*B - B^2)/(a^3*d^2)}*e^{(3*I*d*x + 3*I*c)}*\log(-4*(\sqrt{2}*\sqrt{1/2}*(a^{2*d}*e^{(2*I*d*x + 2*I*c)} + a^{2*d})*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{-(A^2 - 2*I*A*B - B^2)/(a^3*d^2)} + (-I*A - B)*a*e^{(I*d*x + I*c)})*e^{(-I*d*x - I*c)/(I*A + B)} - 3*\sqrt{1/2}*a^{2*d}*\sqrt{-(A^2 - 2*I*A*B - B^2)/(a^3*d^2)}*e^{(3*I*d*x + 3*I*c)}*\log(4*(\sqrt{2}*\sqrt{1/2}*(a^{2*d}*e^{(2*I*d*x + 2*I*c)} + a^{2*d})*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{-(A^2 - 2*I*A*B - B^2)/(a^3*d^2)} - (-I*A - B)*a*e^{(I*d*x + I*c)})*e^{(-I*d*x - I*c)/(I*A + B)} - \sqrt{2}*(2*(-2*I*A - B)*e^{(4*I*d*x + 4*I*c)} - (5*I*A + B)*e^{(2*I*d*x + 2*I*c)} - I*A + B)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*e^{(-3*I*d*x - 3*I*c)/(a^2*d)}$

3.100.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{(ia (\tan(c + dx) - i))^{3/2}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))/(I*a*(tan(c + d*x) - I))**(3/2), x)`

3.100.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.93

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{3/2}} dx = \frac{i \left(\frac{3\sqrt{2}(A-iB) \log\left(\frac{-\sqrt{2}\sqrt{a}-\sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia \tan(dx+c)+a}}\right) + \frac{4(3(ia \tan(dx+c)+a)(A-iB)+2(A+iB)a)}{(ia \tan(dx+c)+a)^{3/2}}}{24ad} \right)}{24ad}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output $1/24*I*(3*\sqrt{2}*(A - I*B)*\log(-(\sqrt{2}*\sqrt{a} - \sqrt{I*a*\tan(d*x + c) + a})/(\sqrt{2}*\sqrt{a} + \sqrt{I*a*\tan(d*x + c) + a}))/\sqrt{a} + 4*(3*(I*a*\tan(d*x + c) + a)*(A - I*B) + 2*(A + I*B)*a)/(I*a*\tan(d*x + c) + a)^{3/2})/(a*d)$

3.100.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^{3/2}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/(I*a*tan(d*x + c) + a)^(3/2), x)`

3.100.9 Mupad [B] (verification not implemented)

Time = 7.56 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.34

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{3/2}} dx = \frac{\frac{A \operatorname{li}}{3d} + \frac{A(a+a \tan(c+dx) \operatorname{li}) \operatorname{li}}{2ad}}{(a + a \tan(c + dx) \operatorname{li})^{3/2}} - \frac{\frac{B}{3} - \frac{B(a+a \tan(c+dx) \operatorname{li})}{2a}}{d(a + a \tan(c + dx) \operatorname{li})^{3/2}}$$

$$- \frac{\sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a+a \tan(c+dx) \operatorname{li}}}{2\sqrt{-a}}\right) \operatorname{li}}{4(-a)^{3/2} d} - \frac{\sqrt{2} B \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{a+a \tan(c+dx) \operatorname{li}}}{2\sqrt{a}}\right)}{4a^{3/2} d}$$

input `int((A + B*tan(c + d*x))/(a + a*tan(c + d*x)*1i)^(3/2),x)`

output `((A*1i)/(3*d) + (A*(a + a*tan(c + d*x)*1i)*1i)/(2*a*d))/(a + a*tan(c + d*x)*1i)^(3/2) - (B/3 - (B*(a + a*tan(c + d*x)*1i))/(2*a))/(d*(a + a*tan(c + d*x)*1i)^(3/2)) - (2^(1/2)*A*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2)))*1i)/(4*(-a)^(3/2)*d) - (2^(1/2)*B*atanh((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*a^(1/2)))/(4*a^(3/2)*d)`

3.101
$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$$

3.101.1 Optimal result 1095
 3.101.2 Mathematica [A] (verified) 1095
 3.101.3 Rubi [A] (verified) 1096
 3.101.4 Maple [A] (verified) 1100
 3.101.5 Fricas [B] (verification not implemented) 1100
 3.101.6 Sympy [F] 1101
 3.101.7 Maxima [A] (verification not implemented) 1101
 3.101.8 Giac [F] 1102
 3.101.9 Mupad [B] (verification not implemented) 1102

3.101.1 Optimal result

Integrand size = 34, antiderivative size = 156

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx =$$

$$-\frac{2A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{a^{3/2}d} + \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{2\sqrt{2}a^{3/2}d}$$

$$+ \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}} + \frac{3A+iB}{2ad\sqrt{a+ia \tan(c+dx)}}$$

output `-2*A*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/a^(3/2)/d+1/4*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(3/2)/d*2^(1/2)+1/2*(3*A+I*B)/a/d/(a+I*a*tan(d*x+c))^(1/2)+1/3*(A+I*B)/d/(a+I*a*tan(d*x+c))^(3/2)`

3.101.2 Mathematica [A] (verified)

Time = 2.79 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.00

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{-6a^3 A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) + \frac{3a^3(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{2\sqrt{2}}}{3a^{9/2}d}$$

input `Integrate[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`

output $(-6*a^3*A*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] + (3*a^3*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/(2*Sqrt[2]) + (a^(9/2)*(A + I*B))/(a + I*a*Tan[c + d*x])^(3/2) + (3*a^(7/2)*(3*A + I*B))/(2*Sqrt[a + I*a*Tan[c + d*x]])/(3*a^(9/2)*d)$

3.101.3 Rubi [A] (verified)

Time = 1.07 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.07, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)(a+ia \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{3 \cot(c+dx)(2aA-a(iA-B) \tan(c+dx))}{2\sqrt{i \tan(c+dx)a+a}} dx}{3a^2} + \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\cot(c+dx)(2aA-a(iA-B) \tan(c+dx))}{\sqrt{i \tan(c+dx)a+a}} dx}{2a^2} + \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{2aA-a(iA-B) \tan(c+dx)}{\tan(c+dx)\sqrt{i \tan(c+dx)a+a}} dx}{2a^2} + \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{\frac{1}{2} \cot(c+dx)\sqrt{i \tan(c+dx)a+a}(4a^2A-a^2(3iA-B) \tan(c+dx))}{a^2} dx}{2a^2} + \frac{a(3A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \\
 & \quad \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.101. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \frac{\int \cot(c+dx) \sqrt{i \tan(c+dx) a+a} (4a^2 A - a^2 (3iA-B) \tan(c+dx)) dx}{2a^2} + \frac{a(3A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sqrt{i \tan(c+dx) a+a} (4a^2 A - a^2 (3iA-B) \tan(c+dx))}{\tan(c+dx)} dx}{2a^2} + \frac{a(3A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{4083} \\
& \frac{a^2(B+iA) \int \sqrt{i \tan(c+dx) a+adx} + 4aA \int \cot(c+dx) (a-ia \tan(c+dx)) \sqrt{i \tan(c+dx) a+adx}}{2a^2} + \frac{a(3A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{a^2(B+iA) \int \sqrt{i \tan(c+dx) a+adx} + 4aA \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx) a+a}}{\tan(c+dx)} dx}{2a^2} + \frac{a(3A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{3961} \\
& \frac{4aA \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx) a+a}}{\tan(c+dx)} dx - \frac{2ia^3(B+iA) \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx) a+a}}{2a^2}}{2a^2} + \frac{a(3A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{219} \\
& \frac{4aA \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx) a+a}}{\tan(c+dx)} dx - \frac{i\sqrt{2}a^{5/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{2a^2} + \frac{a(3A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{4082} \\
& \frac{4a^3 A \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx) a+a}} d \tan(c+dx) - \frac{i\sqrt{2}a^{5/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{2a^2} + \frac{a(3A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{73}
\end{aligned}$$

3.101. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \frac{8ia^2 A \int \frac{1}{i(i \tan(c+dx)a+a)} d\sqrt{i \tan(c+dx)a+a} - i\sqrt{2}a^{5/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{2a^2} + \frac{a(3A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \\
& \frac{2a^2}{3d(a+ia \tan(c+dx))^{3/2}} \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{221} \\
& \frac{i\sqrt{2}a^{5/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right) - 8a^{5/2}A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{2a^2} + \frac{a(3A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \\
& \frac{2a^2}{3d(a+ia \tan(c+dx))^{3/2}} \frac{A+iB}{3d(a+ia \tan(c+dx))^{3/2}}
\end{aligned}$$

input `Int[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `(A + I*B)/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) + (((-8*a^(5/2)*A*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d - (I*Sqrt[2]*a^(5/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d)/(2*a^2) + (a*(3*A + I*B))/(d*Sqrt[a + I*a*Tan[c + d*x]])/(2*a^2)`

3.101.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

3.101. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int[((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x]))], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.101.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.81

method	result
derivativedivides	$2a \left(-\frac{-iB-3A}{4a^2 \sqrt{a+ia \tan(dx+c)}} - \frac{-iB-A}{6a(a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{(iB-A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{8a^{\frac{5}{2}}} - \frac{A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{\sqrt{a}}\right)}{a^{\frac{5}{2}}} \right) dx$
default	$2a \left(-\frac{-iB-3A}{4a^2 \sqrt{a+ia \tan(dx+c)}} - \frac{-iB-A}{6a(a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{(iB-A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{8a^{\frac{5}{2}}} - \frac{A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{\sqrt{a}}\right)}{a^{\frac{5}{2}}} \right) dx$

input `int(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x,method=_RETURNV
ERBOSE)`

output `2/d*a*(-1/4/a^2*(-3*A-I*B)/(a+I*a*tan(d*x+c))^(1/2)-1/6*(-A-I*B)/a/(a+I*a*
tan(d*x+c))^(3/2)-1/8*(-A+I*B)/a^(5/2)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+
c))^(1/2)*2^(1/2)/a^(1/2))-1/a^(5/2)*A*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(
1/2)))`

3.101.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 624 vs. 2(117) = 234.

Time = 0.26 (sec) , antiderivative size = 624, normalized size of antiderivative = 4.00

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx =$$

$$\left(3 \sqrt{\frac{1}{2}} a^2 d \sqrt{\frac{A^2-2iAB-B^2}{a^3 d^2}} e^{(3i dx+3i c)} \log \left(-\frac{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (i a^2 d e^{(2i dx+2i c)} + i a^2 d) \sqrt{\frac{a}{e^{(2i dx+2i c)}+1}} \sqrt{\frac{A^2-2iAB-B^2}{a^3 d^2}} + (-iA-B) a e^{i(c+dx)} \right)}{iA+B} \right) \right)$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm
m="fracas")`

output

```
-1/12*(3*sqrt(1/2)*a^2*d*sqrt((A^2 - 2*I*A*B - B^2)/(a^3*d^2))*e^(3*I*d*x
+ 3*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(I*a^2*d*e^(2*I*d*x + 2*I*c) + I*a^2*d)
*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a^3*d^2)) +
(-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B) - 3*sqrt(1/2)*a
^2*d*sqrt((A^2 - 2*I*A*B - B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(-4*(sqr
t(2)*sqrt(1/2)*(-I*a^2*d*e^(2*I*d*x + 2*I*c) - I*a^2*d)*sqrt(a/(e^(2*I*d*x
+ 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a^3*d^2)) + (-I*A - B)*a*e^(I*
d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B) + 6*a^2*d*sqrt(A^2/(a^3*d^2))*e^(3
*I*d*x + 3*I*c)*log(16*(3*A*a^2*e^(2*I*d*x + 2*I*c) + A*a^2 + 2*sqrt(2)*(a
^3*d*e^(3*I*d*x + 3*I*c) + a^3*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I
*c) + 1))*sqrt(A^2/(a^3*d^2)))*e^(-2*I*d*x - 2*I*c)/A - 6*a^2*d*sqrt(A^2/
(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(16*(3*A*a^2*e^(2*I*d*x + 2*I*c) + A*a^2
- 2*sqrt(2)*(a^3*d*e^(3*I*d*x + 3*I*c) + a^3*d*e^(I*d*x + I*c))*sqrt(a/(e
^(2*I*d*x + 2*I*c) + 1))*sqrt(A^2/(a^3*d^2)))*e^(-2*I*d*x - 2*I*c)/A - sq
rt(2)*(2*(5*A + 2*I*B)*e^(4*I*d*x + 4*I*c) + (11*A + 5*I*B)*e^(2*I*d*x + 2
*I*c) + A + I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-3*I*d*x - 3*I*c)/(
a^2*d)
```

3.101.6 Sympy [F]

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \int \frac{(A+B \tan(c+dx)) \cot(c+dx)}{(ia(\tan(c+dx)-i))^{3/2}} dx$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(3/2), x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)/(I*a*(tan(c + d*x) - I))**(3/2), x)`

3.101.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.03

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{3\sqrt{2}(A-iB) \log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia \tan(dx+c)+a}}\right)}{a^{3/2}} - \frac{24A \log\left(\frac{\sqrt{ia \tan(dx+c)+a}-\sqrt{a}}{\sqrt{ia \tan(dx+c)+a}+\sqrt{a}}\right)}{a^{3/2}} - \frac{4(3(ia \tan(dx+c)+a)(3A+iB)+2(A+iB)a)}{(ia \tan(dx+c)+a)^{3/2} a}$$

24d

3.101. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output
$$-1/24*(3*\sqrt{2}*(A - I*B)*\log(-(\sqrt{2}*\sqrt{a} - \sqrt{I*a*\tan(dx + c) + a})/(\sqrt{2}*\sqrt{a} + \sqrt{I*a*\tan(dx + c) + a}))/a^{3/2} - 24*A*\log((\sqrt{I*a*\tan(dx + c) + a} - \sqrt{a})/(\sqrt{I*a*\tan(dx + c) + a} + \sqrt{a}))/a^{3/2} - 4*(3*(I*a*\tan(dx + c) + a)*(3*A + I*B) + 2*(A + I*B)*a)/((I*a*\tan(dx + c) + a)^{3/2}*a)/d$$

3.101.8 Giac [F]

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{(B \tan(dx + c) + A) \cot(dx + c)}{(ia \tan(dx + c) + a)^{3/2}} dx$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)/(I*a*tan(d*x + c) + a)^(3/2), x)`

3.101.9 Mupad [B] (verification not implemented)

Time = 7.95 (sec) , antiderivative size = 563, normalized size of antiderivative = 3.61

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{3/2}} dx = \frac{\frac{A+B li}{3d} + \frac{(3A+B li)(a+a \tan(c+dx) li)}{2ad}}{(a + a \tan(c + dx) li)^{3/2}} + \frac{2A \operatorname{atanh}\left(\frac{31A^3 d \sqrt{a+a \tan(c+dx) li}}{\sqrt{a^3} \left(\frac{31A^3 d}{a} + \frac{AB^2 d}{a} + \frac{A^2 B d 2i}{a}\right)} + \frac{AB^2 d \sqrt{a+a \tan(c+dx) li}}{\sqrt{a^3} \left(\frac{31A^3 d}{a} + \frac{AB^2 d}{a} + \frac{A^2 B d 2i}{a}\right)} + \frac{A^2 B d \sqrt{a+a \tan(c+dx) li} 2i}{\sqrt{a^3} \left(\frac{31A^3 d}{a} + \frac{AB^2 d}{a} + \frac{A^2 B d 2i}{a}\right)}\right)}{d \sqrt{a^3}} + \frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} A^3 d \sqrt{-a^3} \sqrt{a+a \tan(c+dx) li} 3li}{16 \left(\frac{31dA^3 a^2}{8} - \frac{29i d A^2 B a^2}{8} + \frac{3dAB^2 a^2}{8} - \frac{li d B^3 a^2}{8}\right)} + \frac{\sqrt{2} B^3 d \sqrt{-a^3} \sqrt{a+a \tan(c+dx) li}}{16 \left(\frac{31dA^3 a^2}{8} - \frac{29i d A^2 B a^2}{8} + \frac{3dAB^2 a^2}{8} - \frac{li d B^3 a^2}{8}\right)} + \frac{\sqrt{2} A B}{16 \left(\frac{31dA^3 a^2}{8} - \frac{29i d A^2 B a^2}{8} + \frac{3dAB^2 a^2}{8} - \frac{li d B^3 a^2}{8}\right)}\right)}{4a^3 d}$$

input `int((cot(c + d*x)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*li)^(3/2),x)`

3.101. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

output $((A + B*i)/(3*d) + ((3*A + B*i)*(a + a*\tan(c + d*x)*i))/(2*a*d))/(a + a*\tan(c + d*x)*i)^{3/2} - (2*A*\operatorname{atanh}((31*A^3*d*(a + a*\tan(c + d*x)*i)^{1/2}))/((a^3)^{1/2}*((31*A^3*d)/a + (A*B^2*d)/a + (A^2*B*d*2i)/a)) + (A*B^2*d*(a + a*\tan(c + d*x)*i)^{1/2})/((a^3)^{1/2}*((31*A^3*d)/a + (A*B^2*d)/a + (A^2*B*d*2i)/a)) + (A^2*B*d*(a + a*\tan(c + d*x)*i)^{1/2}*2i)/((a^3)^{1/2}*((31*A^3*d)/a + (A*B^2*d)/a + (A^2*B*d*2i)/a)))/(d*(a^3)^{1/2}) + (2^{1/2})*\operatorname{atanh}(2^{1/2}*A^3*d*(-a^3)^{1/2}*(a + a*\tan(c + d*x)*i)^{1/2}*3i)/(16*((31*A^3*a^2*d)/8 - (B^3*a^2*d*1i)/8 + (3*A*B^2*a^2*d)/8 - (A^2*B*a^2*d*29i)/8)) + (2^{1/2})*B^3*d*(-a^3)^{1/2}*(a + a*\tan(c + d*x)*i)^{1/2})/(16*((31*A^3*a^2*d)/8 - (B^3*a^2*d*1i)/8 + (3*A*B^2*a^2*d)/8 - (A^2*B*a^2*d*29i)/8)) + (2^{1/2})*A*B^2*d*(-a^3)^{1/2}*(a + a*\tan(c + d*x)*i)^{1/2}*3i)/(16*((31*A^3*a^2*d)/8 - (B^3*a^2*d*1i)/8 + (3*A*B^2*a^2*d)/8 - (A^2*B*a^2*d*29i)/8)) + (29*2^{1/2})*A^2*B*d*(-a^3)^{1/2}*(a + a*\tan(c + d*x)*i)^{1/2})/(16*((31*A^3*a^2*d)/8 - (B^3*a^2*d*1i)/8 + (3*A*B^2*a^2*d)/8 - (A^2*B*a^2*d*29i)/8))* (A*i + B)*(-a^3)^{1/2})/(4*a^3*d)$

3.102
$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$$

3.102.1 Optimal result 1104
 3.102.2 Mathematica [A] (verified) 1105
 3.102.3 Rubi [A] (verified) 1105
 3.102.4 Maple [A] (verified) 1110
 3.102.5 Fracas [B] (verification not implemented) 1111
 3.102.6 Sympy [F] 1112
 3.102.7 Maxima [A] (verification not implemented) 1113
 3.102.8 Giac [F] 1113
 3.102.9 Mupad [B] (verification not implemented) 1114

3.102.1 Optimal result

Integrand size = 36, antiderivative size = 217

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{(3iA-2B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{a^{3/2}d} + \frac{(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{(A+iB)\cot(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} + \frac{(13A+7iB)\cot(c+dx)}{6ad\sqrt{a+ia \tan(c+dx)}} - \frac{(7A+3iB)\cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{2a^2d}$$

output

```
(3*I*A-2*B)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/a^(3/2)/d+1/4*(I*A+B)
)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(3/2)/d*2^(1/2)+
1/6*(13*A+7*I*B)*cot(d*x+c)/a/d/(a+I*a*tan(d*x+c))^(1/2)-1/2*(7*A+3*I*B)*c
ot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)/a^2/d+1/3*(A+I*B)*cot(d*x+c)/d/(a+I*a*t
an(d*x+c))^(3/2)
```

3.102.2 Mathematica [A] (verified)

Time = 3.58 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.77

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \frac{12i(3A+2iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{a}}\right) + 3\sqrt{2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{a}}\right)}{12a^3}$$

input `Integrate[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2), x]`

output `((12*I)*(3*A + (2*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] + 3*Sqrt[2]*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] - (2*Sqrt[a]*(-3*(7*A + (3*I)*B) + ((29*I)*A - 11*B)*Cot[c + d*x] + 6*A*Cot[c + d*x]^2))/((I + Cot[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]]))/(12*a^(3/2)*d)`

3.102.3 Rubi [A] (verified)

Time = 1.52 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.07, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.472$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4081, 25, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A+B\tan(c+dx)}{\tan(c+dx)^2(a+ia\tan(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{4079} \\ & \frac{\int \frac{\cot^2(c+dx)(2a(4A+iB)-5a(iA-B)\tan(c+dx))}{2\sqrt{i\tan(c+dx)a+a}} dx}{3a^2} + \frac{(A+iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{\cot^2(c+dx)(2a(4A+iB)-5a(iA-B)\tan(c+dx))}{\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} + \frac{(A+iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \end{aligned}$$

$$\begin{aligned}
 & \int \frac{2a(4A+iB)-5a(iA-B)\tan(c+dx)}{\tan(c+dx)^2\sqrt{i\tan(c+dx)a+a}} dx + \frac{(A+iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{3}{2}\cot^2(c+dx)\sqrt{i\tan(c+dx)a+a}(2a^2(7A+3iB)-a^2(13iA-7B)\tan(c+dx)) dx}{a^2} + \frac{a(13A+7iB)\cot(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} + \\
 & \quad \downarrow \text{4079} \\
 & \frac{6a^2}{3d(a+ia\tan(c+dx))^{3/2}} \frac{(A+iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{3\int \cot^2(c+dx)\sqrt{i\tan(c+dx)a+a}(2a^2(7A+3iB)-a^2(13iA-7B)\tan(c+dx)) dx}{2a^2} + \frac{a(13A+7iB)\cot(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} + \\
 & \quad \downarrow \text{3042} \\
 & \frac{3\int \frac{\sqrt{i\tan(c+dx)a+a}(2a^2(7A+3iB)-a^2(13iA-7B)\tan(c+dx))}{\tan(c+dx)^2} dx}{6a^2} + \frac{a(13A+7iB)\cot(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} + \frac{(A+iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \\
 & \quad \downarrow \text{4081} \\
 & \frac{3\left(\frac{\int -\cot(c+dx)\sqrt{i\tan(c+dx)a+a}(2(3iA-2B)a^3+(7A+3iB)\tan(c+dx)a^3) dx}{a} - \frac{2a^2(7A+3iB)\cot(c+dx)\sqrt{a+ia\tan(c+dx)}}{d}\right)}{2a^2} + \frac{a(13A+7iB)\cot(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} + \\
 & \quad \downarrow \text{25} \\
 & \frac{3\left(-\frac{\int \cot(c+dx)\sqrt{i\tan(c+dx)a+a}(2(3iA-2B)a^3+(7A+3iB)\tan(c+dx)a^3) dx}{a} - \frac{2a^2(7A+3iB)\cot(c+dx)\sqrt{a+ia\tan(c+dx)}}{d}\right)}{2a^2} + \frac{a(13A+7iB)\cot(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} + \\
 & \quad \downarrow \text{3042} \\
 & \frac{6a^2}{3d(a+ia\tan(c+dx))^{3/2}} \frac{(A+iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}}
 \end{aligned}$$

3.102. $\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx$

$$\frac{3 \left(-\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (2(3iA-2B)a^3+(7A+3iB) \tan(c+dx)a^3)}{\tan(c+dx)} dx}{a} - \frac{2a^2(7A+3iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} + \frac{a(13A+7iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} +$$

$$\frac{6a^2 (A+iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 4083

$$\frac{3 \left(-\frac{a^3(A-iB) \int \sqrt{i \tan(c+dx)a+adx} + 2a^2(-2B+3iA) \int \cot(c+dx)(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{a} - \frac{2a^2(7A+3iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} + \frac{a(13A+7iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} +$$

$$\frac{6a^2 (A+iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 3042

$$\frac{3 \left(-\frac{a^3(A-iB) \int \sqrt{i \tan(c+dx)a+adx} + 2a^2(-2B+3iA) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{\tan(c+dx)} dx}{a} - \frac{2a^2(7A+3iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} + \frac{a(13A+7iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} +$$

$$\frac{6a^2 (A+iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 3961

$$\frac{3 \left(-\frac{2a^2(-2B+3iA) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{\tan(c+dx)} dx}{a} - \frac{2ia^4(A-iB) \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+adx}}{d} - \frac{2a^2(7A+3iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} + \frac{a(13A+7iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} +$$

$$\frac{6a^2 (A+iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 219

$$\frac{3 \left(-\frac{2a^2(-2B+3iA) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{\tan(c+dx)} dx}{a} - \frac{i\sqrt{2}a^{7/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{2a^2(7A+3iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} + \frac{a(13A+7iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} +$$

$$\frac{6a^2 (A+iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 4082

3.102. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & 3 \left(-\frac{2a^4(-2B+3iA) \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{2a^2} - \frac{i\sqrt{2}a^{7/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{a} - \frac{2a^2(7A+3iB) \cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{d} \right) + \frac{a(3)}{c} \\
 & \frac{(A+iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \quad 6a^2 \\
 & \quad \downarrow \text{73} \\
 & 3 \left(-\frac{4ia^3(-2B+3iA) \int \frac{1}{i-i \tan(c+dx)a+a} d \sqrt{i \tan(c+dx)a+a}}{2a^2} - \frac{i\sqrt{2}a^{7/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{a} - \frac{2a^2(7A+3iB) \cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{d} \right) + \frac{a(3)}{c} \\
 & \frac{(A+iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \quad 6a^2 \\
 & \quad \downarrow \text{221} \\
 & 3 \left(-\frac{4a^{7/2}(-2B+3iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{2a^2} - \frac{i\sqrt{2}a^{7/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{a} - \frac{2a^2(7A+3iB) \cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{d} \right) + \frac{a(3)}{c} \\
 & \frac{(A+iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \quad 6a^2
 \end{aligned}$$

input `Int[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `((A + I*B)*Cot[c + d*x])/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) + ((a*(13*A + (7*I)*B)*Cot[c + d*x])/(d*Sqrt[a + I*a*Tan[c + d*x]]) + (3*(-(((4*a^(7/2) * ((3*I)*A - 2*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d - (I*Sqrt[2]*a^(7/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d)/a - (2*a^2*(7*A + (3*I)*B)*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]])/d))/(2*a^2))/(6*a^2)`

3.102.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`
- rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d)), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2)), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.102.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.77

method	result
derivativedivides	$2ia^2 \left(-\frac{3iB+5A}{4a^3\sqrt{a+ia \tan(dx+c)}} - \frac{iB+A}{6a^2(a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{(iB-A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{8a^{\frac{7}{2}}} + \frac{iA\sqrt{a+ia \tan(dx+c)}}{2a \tan(dx+c)} + \dots \right) / d$
default	$2ia^2 \left(-\frac{3iB+5A}{4a^3\sqrt{a+ia \tan(dx+c)}} - \frac{iB+A}{6a^2(a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{(iB-A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{8a^{\frac{7}{2}}} + \frac{iA\sqrt{a+ia \tan(dx+c)}}{2a \tan(dx+c)} + \dots \right) / d$

input `int(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

3.102. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

output $2*I/d*a^2*(-1/4/a^3*(5*A+3*I*B)/(a+I*a*\tan(d*x+c))^{1/2}-1/6/a^2*(A+I*B)/(a+I*a*\tan(d*x+c))^{3/2}-1/8*(-A+I*B)/a^{7/2}*2^{1/2}*\operatorname{arctanh}(1/2*(a+I*a*\tan(d*x+c))^{1/2}*2^{1/2}/a^{1/2}))+1/a^3*(1/2*I*A*(a+I*a*\tan(d*x+c))^{1/2}/a/\tan(d*x+c)+1/2*(3*A+2*I*B)/a^{1/2}*\operatorname{arctanh}((a+I*a*\tan(d*x+c))^{1/2}/a^{1/2}))$

3.102.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 814 vs. $2(170) = 340$.

Time = 0.28 (sec) , antiderivative size = 814, normalized size of antiderivative = 3.75

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx =$$

$$3\sqrt{\frac{1}{2}}(a^2de^{(5i dx+5i c)} - a^2de^{(3i dx+3i c)})\sqrt{-\frac{A^2-2iAB-B^2}{a^3d^2}}\log\left(-\frac{4\left(\sqrt{2}\sqrt{\frac{1}{2}}(a^2de^{(2i dx+2i c)}+a^2d)\sqrt{\frac{a}{e^{(2i dx+2i c)}+1}}\sqrt{-\frac{A^2-2iAB-B^2}{a^3d^2}}\right)}{iA+B}\right)$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="fracas")`

```

output -1/12*(3*sqrt(1/2)*(a^2*d*e^(5*I*d*x + 5*I*c) - a^2*d*e^(3*I*d*x + 3*I*c))
*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^3*d^2))*log(-4*(sqrt(2)*sqrt(1/2)*(a^2*d*e
^(2*I*d*x + 2*I*c) + a^2*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-(A^2 -
2*I*A*B - B^2)/(a^3*d^2)) + (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c
)/(I*A + B)) - 3*sqrt(1/2)*(a^2*d*e^(5*I*d*x + 5*I*c) - a^2*d*e^(3*I*d*x +
3*I*c))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^3*d^2))*log(4*(sqrt(2)*sqrt(1/2)*(
a^2*d*e^(2*I*d*x + 2*I*c) + a^2*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(
-(A^2 - 2*I*A*B - B^2)/(a^3*d^2)) - (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*
x - I*c)/(I*A + B)) - 3*(a^2*d*e^(5*I*d*x + 5*I*c) - a^2*d*e^(3*I*d*x + 3*
I*c))*sqrt(-(9*A^2 + 12*I*A*B - 4*B^2)/(a^3*d^2))*log(-16*(3*(3*I*A - 2*B)
*a^2*e^(2*I*d*x + 2*I*c) + (3*I*A - 2*B)*a^2 + 2*sqrt(2)*(a^3*d*e^(3*I*d*x
+ 3*I*c) + a^3*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(
-(9*A^2 + 12*I*A*B - 4*B^2)/(a^3*d^2)))*e^(-2*I*d*x - 2*I*c)/(-3*I*A + 2*B
)) + 3*(a^2*d*e^(5*I*d*x + 5*I*c) - a^2*d*e^(3*I*d*x + 3*I*c))*sqrt(-(9*A^
2 + 12*I*A*B - 4*B^2)/(a^3*d^2))*log(-16*(3*(3*I*A - 2*B)*a^2*e^(2*I*d*x +
2*I*c) + (3*I*A - 2*B)*a^2 - 2*sqrt(2)*(a^3*d*e^(3*I*d*x + 3*I*c) + a^3*d
*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-(9*A^2 + 12*I*A*
B - 4*B^2)/(a^3*d^2)))*e^(-2*I*d*x - 2*I*c)/(-3*I*A + 2*B)) + sqrt(2)*(2*(
14*I*A - 5*B)*e^(6*I*d*x + 6*I*c) - (-13*I*A + B)*e^(4*I*d*x + 4*I*c) + 2*
(-8*I*A + 5*B)*e^(2*I*d*x + 2*I*c) - I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*...

```

3.102.6 Sympy [F]

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \int \frac{(A+B \tan(c+dx)) \cot^2(c+dx)}{(ia(\tan(c+dx)-i))^{\frac{3}{2}}} dx$$

```

input integrate(cot(d*x+c)**2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(3/2),x)

```

```

output Integral((A + B*tan(c + d*x))*cot(c + d*x)**2/(I*a*(tan(c + d*x) - I))**(3
/2), x)

```

3.102.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.99

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx =$$

$$i a \left(\frac{4 \left(3 (i a \tan(dx+c)+a)^2 (7 A+3 i B) - (i a \tan(dx+c)+a) (13 A+7 i B) a - 2 (A+i B) a^2 \right)}{(i a \tan(dx+c)+a)^{\frac{5}{2}} a^2 - (i a \tan(dx+c)+a)^{\frac{3}{2}} a^3} + \frac{3 \sqrt{2} (A-i B) \log \left(-\frac{\sqrt{2} \sqrt{a}-\sqrt{i a \tan(dx+c)+a}}{\sqrt{2} \sqrt{a}+\sqrt{i a \tan(dx+c)+a}} \right)}{a^{\frac{5}{2}}} \right) +$$

$24 d$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")
```

```
output -1/24*I*a*(4*(3*(I*a*tan(d*x+c)+a)^2*(7*A+3*I*B)-(I*a*tan(d*x+c)+a)*(13*A+7*I*B)*a-2*(A+I*B)*a^2)/((I*a*tan(d*x+c)+a)^(5/2)*a^2-(I*a*tan(d*x+c)+a)^(3/2)*a^3)+3*sqrt(2)*(A-I*B)*log(-(sqrt(2)*sqrt(a)-sqrt(I*a*tan(d*x+c)+a))/(sqrt(2)*sqrt(a)+sqrt(I*a*tan(d*x+c)+a)))/a^(5/2)+12*(3*A+2*I*B)*log((sqrt(I*a*tan(d*x+c)+a)-sqrt(a))/(sqrt(I*a*tan(d*x+c)+a)+sqrt(a)))/a^(5/2))/d
```

3.102.8 Giac [F]

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \int \frac{(B \tan(dx+c)+A) \cot(dx+c)^2}{(i a \tan(dx+c)+a)^{\frac{3}{2}}} dx$$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")
```

```
output integrate((B*tan(d*x+c)+A)*cot(d*x+c)^2/(I*a*tan(d*x+c)+a)^(3/2),x)
```


3.102.9 Mupad [B] (verification not implemented)

Time = 9.81 (sec) , antiderivative size = 3051, normalized size of antiderivative = 14.06

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `int((cot(c + d*x)^2*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(3/2),x)`

output

```

2*atanh((3*d^4*(a + a*tan(c + d*x)*1i)^(1/2)*((33*B^2)/(64*a^3*d^2) - (73*
A^2)/(64*a^3*d^2) - ((5041*A^4*a^6)/d^4 + (961*B^4*a^6)/d^4 - (14006*A^2*B
^2*a^6)/d^4 - (A*B^3*a^6*6076i)/d^4 + (A^3*B*a^6*13916i)/d^4)^(1/2)/(64*a^
6) - (A*B*47i)/(32*a^3*d^2))^(1/2)*((5041*A^4*a^6)/d^4 + (961*B^4*a^6)/d^4
- (14006*A^2*B^2*a^6)/d^4 - (A*B^3*a^6*6076i)/d^4 + (A^3*B*a^6*13916i)/d^
4)^(1/2))/((A^3*a^2*d*781i)/4 + (279*B^3*a^2*d)/4 - (A*B^2*a^2*d*1223i)/4
- (1717*A^2*B*a^2*d)/4 + (A*d^3*((5041*A^4*a^6)/d^4 + (961*B^4*a^6)/d^4 -
(14006*A^2*B^2*a^6)/d^4 - (A*B^3*a^6*6076i)/d^4 + (A^3*B*a^6*13916i)/d^4)^(
1/2)*13i)/(4*a) - (7*B*d^3*((5041*A^4*a^6)/d^4 + (961*B^4*a^6)/d^4 - (140
06*A^2*B^2*a^6)/d^4 - (A*B^3*a^6*6076i)/d^4 + (A^3*B*a^6*13916i)/d^4)^(1/2
))/(4*a)) + (71*A^2*d^2*(a + a*tan(c + d*x)*1i)^(1/2)*((33*B^2)/(64*a^3*d^
2) - (73*A^2)/(64*a^3*d^2) - ((5041*A^4*a^6)/d^4 + (961*B^4*a^6)/d^4 - (14
006*A^2*B^2*a^6)/d^4 - (A*B^3*a^6*6076i)/d^4 + (A^3*B*a^6*13916i)/d^4)^(1/
2)/(64*a^6) - (A*B*47i)/(32*a^3*d^2))^(1/2))/((A^3*d*781i)/(4*a) + (279*B^
3*d)/(4*a) - (A*B^2*d*1223i)/(4*a) - (1717*A^2*B*d)/(4*a) + (A*d^3*((5041*
A^4*a^6)/d^4 + (961*B^4*a^6)/d^4 - (14006*A^2*B^2*a^6)/d^4 - (A*B^3*a^6*60
76i)/d^4 + (A^3*B*a^6*13916i)/d^4)^(1/2)*13i)/(4*a^4) - (7*B*d^3*((5041*A^
4*a^6)/d^4 + (961*B^4*a^6)/d^4 - (14006*A^2*B^2*a^6)/d^4 - (A*B^3*a^6*6076
i)/d^4 + (A^3*B*a^6*13916i)/d^4)^(1/2))/(4*a^4)) - (31*B^2*d^2*(a + a*tan(
c + d*x)*1i)^(1/2)*((33*B^2)/(64*a^3*d^2) - (73*A^2)/(64*a^3*d^2) - ((5...
```

3.103 $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

3.103.1 Optimal result	1115
3.103.2 Mathematica [A] (verified)	1116
3.103.3 Rubi [A] (verified)	1116
3.103.4 Maple [A] (verified)	1122
3.103.5 Fracas [B] (verification not implemented)	1123
3.103.6 Sympy [F]	1124
3.103.7 Maxima [A] (verification not implemented)	1124
3.103.8 Giac [F]	1125
3.103.9 Mupad [B] (verification not implemented)	1125

3.103.1 Optimal result

Integrand size = 36, antiderivative size = 268

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{(23A+12iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{4a^{3/2}d} - \frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{2\sqrt{2}a^{3/2}d} + \frac{(A+iB)\cot^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} + \frac{(17A+11iB)\cot^2(c+dx)}{6ad\sqrt{a+ia \tan(c+dx)}} + \frac{7(3iA-2B)\cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{4a^2d} - \frac{(22A+13iB)\cot^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{6a^2d}$$

output

```
1/4*(23*A+12*I*B)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/a^(3/2)/d-1/4*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(3/2)/d*2^(1/2)+1/6*(17*A+11*I*B)*cot(d*x+c)^2/a/d/(a+I*a*tan(d*x+c))^(1/2)+7/4*(3*I*A-2*B)*cot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)/a^2/d-1/6*(22*A+13*I*B)*cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(1/2)/a^2/d+1/3*(A+I*B)*cot(d*x+c)^2/d/(a+I*a*tan(d*x+c))^(3/2)
```

3.103.2 Mathematica [A] (verified)

Time = 5.45 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.68

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{3(23A+12iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right) - 3\sqrt{2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{(a+ia \tan(c+dx))^{3/2}}$$

input `Integrate[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `(3*(23*A + (12*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]] - 3*Sqrt[2]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])] - (Sqrt[a]*((63*I)*A - 42*B + (82*A + (58*I)*B)*Cot[c + d*x] + 3*((-3*I)*A + 4*B)*Cot[c + d*x]^2 + 6*A*Cot[c + d*x]^3))/((I + Cot[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]]))/(12*a^(3/2)*d)`

3.103.3 Rubi [A] (verified)

Time = 1.92 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.07, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^3(a+ia \tan(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{4079} \\ & \frac{\int \frac{\cot^3(c+dx)(2a(5A+2iB)-7a(iA-B) \tan(c+dx))}{2\sqrt{i \tan(c+dx)a+a}} dx}{3a^2} + \frac{(A+iB) \cot^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\begin{aligned}
& \frac{\int \frac{\cot^3(c+dx)(2a(5A+2iB)-7a(iA-B)\tan(c+dx))}{\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} + \frac{(A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{2a(5A+2iB)-7a(iA-B)\tan(c+dx)}{\tan(c+dx)^3\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} + \frac{(A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{4079} \\
& \frac{\int \frac{1}{2}\cot^3(c+dx)\sqrt{i\tan(c+dx)a+a}(4a^2(22A+13iB)-5a^2(17iA-11B)\tan(c+dx)) dx}{a^2} + \frac{a(17A+11iB)\cot^2(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} + \\
& \quad \frac{6a^2}{3d(a+ia\tan(c+dx))^{3/2}} \frac{(A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \cot^3(c+dx)\sqrt{i\tan(c+dx)a+a}(4a^2(22A+13iB)-5a^2(17iA-11B)\tan(c+dx)) dx}{2a^2} + \frac{a(17A+11iB)\cot^2(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} + \\
& \quad \frac{6a^2}{3d(a+ia\tan(c+dx))^{3/2}} \frac{(A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sqrt{i\tan(c+dx)a+a}(4a^2(22A+13iB)-5a^2(17iA-11B)\tan(c+dx))}{\tan(c+dx)^3} dx}{2a^2} + \frac{a(17A+11iB)\cot^2(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} + \\
& \quad \frac{6a^2}{3d(a+ia\tan(c+dx))^{3/2}} \frac{(A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{4081} \\
& \frac{\int -6\cot^2(c+dx)\sqrt{i\tan(c+dx)a+a}(7(3iA-2B)a^3+(22A+13iB)\tan(c+dx)a^3) dx}{2a} - \frac{2a^2(22A+13iB)\cot^2(c+dx)\sqrt{a+ia\tan(c+dx)}}{d} + \frac{a(17A+11iB)\cot^2(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} \\
& \quad \frac{6a^2}{3d(a+ia\tan(c+dx))^{3/2}} \frac{(A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{27} \\
& \frac{3\int \cot^2(c+dx)\sqrt{i\tan(c+dx)a+a}(7(3iA-2B)a^3+(22A+13iB)\tan(c+dx)a^3) dx}{a} - \frac{2a^2(22A+13iB)\cot^2(c+dx)\sqrt{a+ia\tan(c+dx)}}{d} + \frac{a(17A+11iB)\cot^2(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} \\
& \quad \frac{6a^2}{3d(a+ia\tan(c+dx))^{3/2}} \frac{(A+iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.103. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{3 \int \frac{\sqrt{i \tan(c+dx)a+a} (7(3iA-2B)a^3+(22A+13iB) \tan(c+dx)a^3)}{\tan(c+dx)^2} dx - \frac{2a^2(22A+13iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} + \frac{a(17A+11iB) \cot^2(c+dx)}{d \sqrt{a+ia \tan(c+dx)}}}{2a^2} + \\
 & \frac{6a^2}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 4081 \\
 & \frac{3 \left(\int \frac{\frac{1}{2} \cot(c+dx) \sqrt{i \tan(c+dx)a+a} (a^4(23A+12iB)-7a^4(3iA-2B) \tan(c+dx))}{a} dx - \frac{7a^3(-2B+3iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right) - \frac{2a^2(22A+13iB) \cot^2(c+dx)}{d}}{2a^2} \\
 & \frac{6a^2}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 27 \\
 & \frac{3 \left(\int \frac{\cot(c+dx) \sqrt{i \tan(c+dx)a+a} (a^4(23A+12iB)-7a^4(3iA-2B) \tan(c+dx))}{2a} dx - \frac{7a^3(-2B+3iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right) - \frac{2a^2(22A+13iB) \cot^2(c+dx)}{d}}{2a^2} \\
 & \frac{6a^2}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 3042 \\
 & \frac{3 \left(\int \frac{\frac{\sqrt{i \tan(c+dx)a+a} (a^4(23A+12iB)-7a^4(3iA-2B) \tan(c+dx))}{2a} dx - \frac{7a^3(-2B+3iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right) - \frac{2a^2(22A+13iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}}{2a^2} \\
 & \frac{6a^2}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 4083 \\
 & \frac{3 \left(\frac{2a^4(B+iA) \int \sqrt{i \tan(c+dx)a+adx} + a^3(23A+12iB) \int \cot(c+dx)(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{2a} - \frac{7a^3(-2B+3iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right) - 2a^2}{2a^2} \\
 & \frac{6a^2}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 3042
 \end{aligned}$$

3.103. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

$$\frac{3 \left(\frac{2a^4(B+iA) \int \sqrt{i \tan(c+dx)a+adx+a^3(23A+12iB) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx}{2a} - \frac{7a^3(-2B+3iA) \cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{d} \right)}{a} - \frac{2a^2(22A+13iB)}{2a^2}$$

$$\frac{(A+iB) \cot^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \quad 6a^2$$

↓ 3961

$$\frac{3 \left(\frac{a^3(23A+12iB) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx}{2a} - \frac{4ia^5(B+iA) \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a}}{d} - \frac{7a^3(-2B+3iA) \cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{d} \right)}{a} - \frac{2a^2}{2a^2}$$

$$\frac{(A+iB) \cot^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \quad 6a^2$$

↓ 219

$$\frac{3 \left(\frac{a^3(23A+12iB) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx}{2a} - \frac{2i\sqrt{2}a^{9/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{7a^3(-2B+3iA) \cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{d} \right)}{a} - \frac{2a^2}{2a^2}$$

$$\frac{(A+iB) \cot^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \quad 6a^2$$

↓ 4082

$$\frac{3 \left(\frac{a^5(23A+12iB) \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{d} - \frac{2i\sqrt{2}a^{9/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{7a^3(-2B+3iA) \cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{d} \right)}{a} - \frac{2a^2}{2a^2}$$

$$\frac{(A+iB) \cot^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \quad 6a^2$$

↓ 73

3.103. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{3 \left(\frac{2ia^4(23A+12iB) \int \frac{1}{i - i \tan(c+dx)a+a} d\sqrt{i \tan(c+dx)a+a} - \frac{2i\sqrt{2}a^{9/2}(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{7a^3(-2B+3iA) \cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} \\
 & \frac{6a^2}{(A+iB) \cot^2(c+dx)} \\
 & \frac{3d(a+ia \tan(c+dx))^{3/2}}{\downarrow 221} \\
 & \frac{3 \left(\frac{2a^{9/2}(23A+12iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{d} - \frac{2i\sqrt{2}a^{9/2}(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{2a^2(22A+13iB) \cot^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} \\
 & \frac{6a^2}{(A+iB) \cot^2(c+dx)} \\
 & \frac{3d(a+ia \tan(c+dx))^{3/2}}{
 \end{aligned}$$

input `Int[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `((A + I*B)*Cot[c + d*x]^2)/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) + ((a*(17*A + (11*I)*B)*Cot[c + d*x]^2)/(d*Sqrt[a + I*a*Tan[c + d*x]]) + (((-2*a^2*(22*A + (13*I)*B)*Cot[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]])/d - (3*((-2*a^(9/2)*(23*A + (12*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d - ((2*I)*Sqrt[2]*a^(9/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d)/(2*a) - (7*a^3*((3*I)*A - 2*B)*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]])/d)/a)/(2*a^2))/(6*a^2)`

3.103.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

3.103. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

rule 4081 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.103.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.74

method	result
derivativedivides	$2a^3 \left(-\frac{5iB+7A}{4a^4 \sqrt{a+ia \tan(dx+c)}} - \frac{iB+A}{6a^3 (a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}}\right)}{8a^{\frac{9}{2}}} \right) + \frac{\left(-\frac{iB}{2} - \frac{7A}{8}\right)(a+ia \tan(dx+c))}{d}$
default	$2a^3 \left(-\frac{5iB+7A}{4a^4 \sqrt{a+ia \tan(dx+c)}} - \frac{iB+A}{6a^3 (a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}}\right)}{8a^{\frac{9}{2}}} \right) + \frac{\left(-\frac{iB}{2} - \frac{7A}{8}\right)(a+ia \tan(dx+c))}{d}$

input `int(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output `2/d*a^3*(-1/4/a^4*(7*A+5*I*B)/(a+I*a*tan(d*x+c))^(1/2)-1/6/a^3*(A+I*B)/(a+I*a*tan(d*x+c))^(3/2)-1/8/a^(9/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))+1/a^4*(-((-1/2*I*B-7/8*A)*(a+I*a*tan(d*x+c))^(3/2)+(1/2*I*a*B+9/8*a*A)*(a+I*a*tan(d*x+c))^(1/2))/a^2/tan(d*x+c)^2+1/8*(23*A+12*I*B)/a^(1/2)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))))`

3.103. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

3.103.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 903 vs. $2(209) = 418$.

Time = 0.28 (sec) , antiderivative size = 903, normalized size of antiderivative = 3.37

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorith
thm="fracas")
```

```
output 1/48*(12*sqrt(1/2)*(a^2*d*e^(7*I*d*x + 7*I*c) - 2*a^2*d*e^(5*I*d*x + 5*I*c)
) + a^2*d*e^(3*I*d*x + 3*I*c))*sqrt((A^2 - 2*I*A*B - B^2)/(a^3*d^2))*log(-
4*(sqrt(2)*sqrt(1/2)*(I*a^2*d*e^(2*I*d*x + 2*I*c) + I*a^2*d)*sqrt(a/(e^(2*
I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a^3*d^2)) + (-I*A - B)*a*
e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 12*sqrt(1/2)*(a^2*d*e^(7*I*
d*x + 7*I*c) - 2*a^2*d*e^(5*I*d*x + 5*I*c) + a^2*d*e^(3*I*d*x + 3*I*c))*sq
rt((A^2 - 2*I*A*B - B^2)/(a^3*d^2))*log(-4*(sqrt(2)*sqrt(1/2)*(-I*a^2*d*e^
(2*I*d*x + 2*I*c) - I*a^2*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 -
2*I*A*B - B^2)/(a^3*d^2)) + (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c
)/(I*A + B)) + 3*(a^2*d*e^(7*I*d*x + 7*I*c) - 2*a^2*d*e^(5*I*d*x + 5*I*c)
+ a^2*d*e^(3*I*d*x + 3*I*c))*sqrt((529*A^2 + 552*I*A*B - 144*B^2)/(a^3*d^2
))*log(-16*(3*(23*I*A - 12*B)*a^2*e^(2*I*d*x + 2*I*c) + (23*I*A - 12*B)*a^
2 + 2*sqrt(2)*(I*a^3*d*e^(3*I*d*x + 3*I*c) + I*a^3*d*e^(I*d*x + I*c))*sqrt
(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((529*A^2 + 552*I*A*B - 144*B^2)/(a^3*d^
2)))*e^(-2*I*d*x - 2*I*c)/(-23*I*A + 12*B)) - 3*(a^2*d*e^(7*I*d*x + 7*I*c)
- 2*a^2*d*e^(5*I*d*x + 5*I*c) + a^2*d*e^(3*I*d*x + 3*I*c))*sqrt((529*A^2
+ 552*I*A*B - 144*B^2)/(a^3*d^2))*log(-16*(3*(23*I*A - 12*B)*a^2*e^(2*I*d*
x + 2*I*c) + (23*I*A - 12*B)*a^2 + 2*sqrt(2)*(-I*a^3*d*e^(3*I*d*x + 3*I*c)
- I*a^3*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((529*A^
2 + 552*I*A*B - 144*B^2)/(a^3*d^2)))*e^(-2*I*d*x - 2*I*c)/(-23*I*A + 12...
```

3.103.6 Sympy [F]

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \int \frac{(A+B \tan(c+dx)) \cot^3(c+dx)}{(ia(\tan(c+dx)-i))^{\frac{3}{2}}} dx$$

input `integrate(cot(d*x+c)**3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**3/(I*a*(tan(c + d*x) - I))**(3/2), x)`

3.103.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 259, normalized size of antiderivative = 0.97

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = a^2 \left(\frac{2(21(ia \tan(dx+c)+a)^3(3A+2iB)-(ia \tan(dx+c)+a)^2(107A+68iB)a+2(ia \tan(dx+c)+a)(17A+11iB)a^2+4(A+iB)a^3)}{(ia \tan(dx+c)+a)^{\frac{7}{2}}a^3-2(ia \tan(dx+c)+a)^{\frac{5}{2}}a^4+(ia \tan(dx+c)+a)^{\frac{3}{2}}a^5} - \frac{3\sqrt{2}(A-I*B)\log(-\sqrt{2}\sqrt{a}-\sqrt{ia \tan(dx+c)+a})}{\sqrt{2}\sqrt{a}+\sqrt{ia \tan(dx+c)+a}} \right) / d$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `-1/24*a^2*(2*(21*(I*a*tan(d*x + c) + a)^3*(3*A + 2*I*B) - (I*a*tan(d*x + c) + a)^2*(107*A + 68*I*B)*a + 2*(I*a*tan(d*x + c) + a)*(17*A + 11*I*B)*a^2 + 4*(A + I*B)*a^3)/((I*a*tan(d*x + c) + a)^(7/2)*a^3 - 2*(I*a*tan(d*x + c) + a)^(5/2)*a^4 + (I*a*tan(d*x + c) + a)^(3/2)*a^5) - 3*sqrt(2)*(A - I*B)*log(-sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))/a^(7/2) + 3*(23*A + 12*I*B)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a)))/a^(7/2))/d`

3.103.8 Giac [F]

$$\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{(B \tan(dx + c) + A) \cot(dx + c)^3}{(ia \tan(dx + c) + a)^{3/2}} dx$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^3/(I*a*tan(d*x + c) + a)^(3/2), x)`

3.103.9 Mupad [B] (verification not implemented)

Time = 9.93 (sec) , antiderivative size = 3106, normalized size of antiderivative = 11.59

$$\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `int((cot(c + d*x)^3*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(3/2),x)`

output

$$\begin{aligned}
& 2*\operatorname{atanh}((48*d^4*(a + a*\tan(c + d*x)*i)^{(1/2))*((531*A^2)/(128*a^3*d^2) - (\\
& (277729*A^4*a^6)/(4*d^4) + (5041*B^4*a^6)/d^4 - (114701*A^2*B^2*a^6)/d^4 - \\
& (A*B^3*a^6*39476i)/d^4 + (A^3*B*a^6*146506i)/d^4)^{(1/2)})/(64*a^6) - (73*B^ \\
& 2)/(64*a^3*d^2) + (A*B*137i)/(32*a^3*d^2))^{(1/2))*((277729*A^4*a^6)/(4*d^4) \\
& + (5041*B^4*a^6)/d^4 - (114701*A^2*B^2*a^6)/d^4 - (A*B^3*a^6*39476i)/d^4 \\
& + (A^3*B*a^6*146506i)/d^4)^{(1/2)})/(B^3*a^2*d*3124i - 25296*A^3*a^2*d + 190 \\
& 48*A*B^2*a^2*d - A^2*B*a^2*d*38282i + (88*A*d^3*((277729*A^4*a^6)/(4*d^4) \\
& + (5041*B^4*a^6)/d^4 - (114701*A^2*B^2*a^6)/d^4 - (A*B^3*a^6*39476i)/d^4 + \\
& (A^3*B*a^6*146506i)/d^4)^{(1/2)})/a + (B*d^3*((277729*A^4*a^6)/(4*d^4) + (5 \\
& 041*B^4*a^6)/d^4 - (114701*A^2*B^2*a^6)/d^4 - (A*B^3*a^6*39476i)/d^4 + (A^ \\
& 3*B*a^6*146506i)/d^4)^{(1/2)}*52i)/a) - (4216*A^2*d^2*(a + a*\tan(c + d*x)*i \\
&)^{(1/2))*((531*A^2)/(128*a^3*d^2) - ((277729*A^4*a^6)/(4*d^4) + (5041*B^4*a \\
& ^6)/d^4 - (114701*A^2*B^2*a^6)/d^4 - (A*B^3*a^6*39476i)/d^4 + (A^3*B*a^6*1 \\
& 46506i)/d^4)^{(1/2)})/(64*a^6) - (73*B^2)/(64*a^3*d^2) + (A*B*137i)/(32*a^3*d \\
& ^2))^{(1/2)})/(B^3*d*3124i)/a - (25296*A^3*d)/a + (19048*A*B^2*d)/a - (A^2* \\
& B*d*38282i)/a + (88*A*d^3*((277729*A^4*a^6)/(4*d^4) + (5041*B^4*a^6)/d^4 - \\
& (114701*A^2*B^2*a^6)/d^4 - (A*B^3*a^6*39476i)/d^4 + (A^3*B*a^6*146506i)/d \\
& ^4)^{(1/2)})/a^4 + (B*d^3*((277729*A^4*a^6)/(4*d^4) + (5041*B^4*a^6)/d^4 - (\\
& 114701*A^2*B^2*a^6)/d^4 - (A*B^3*a^6*39476i)/d^4 + (A^3*B*a^6*146506i)/d^4 \\
&)^{(1/2)}*52i)/a^4) + (1136*B^2*d^2*(a + a*\tan(c + d*x)*i)^{(1/2))*((531*A...
\end{aligned}$$

3.104
$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$$

3.104.1 Optimal result 1127
 3.104.2 Mathematica [C] (verified) 1127
 3.104.3 Rubi [A] (verified) 1128
 3.104.4 Maple [A] (verified) 1132
 3.104.5 Fricas [B] (verification not implemented) 1133
 3.104.6 Sympy [F] 1133
 3.104.7 Maxima [A] (verification not implemented) 1134
 3.104.8 Giac [F] 1134
 3.104.9 Mupad [B] (verification not implemented) 1135

3.104.1 Optimal result

Integrand size = 36, antiderivative size = 255

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = -\frac{(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{4\sqrt{2}a^{5/2}d} + \frac{(iA-B)\tan^4(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} + \frac{(11A+21iB)\tan^3(c+dx)}{30ad(a+ia \tan(c+dx))^{3/2}} - \frac{(39iA-89B)\tan^2(c+dx)}{20a^2d\sqrt{a+ia \tan(c+dx)}} + \frac{(39iA-89B)\sqrt{a+ia \tan(c+dx)}}{5a^3d} - \frac{(151iA-361B)(a+ia \tan(c+dx))^{3/2}}{60a^4d}$$

```
output -1/8*(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(5/2)
/d*2^(1/2)+1/5*(39*I*A-89*B)*(a+I*a*tan(d*x+c))^(1/2)/a^3/d-1/20*(39*I*A-8
9*B)*tan(d*x+c)^2/a^2/d/(a+I*a*tan(d*x+c))^(1/2)+1/5*(I*A-B)*tan(d*x+c)^4/
d/(a+I*a*tan(d*x+c))^(5/2)+1/30*(11*A+21*I*B)*tan(d*x+c)^3/a/d/(a+I*a*tan(
d*x+c))^(3/2)-1/60*(151*I*A-361*B)*(a+I*a*tan(d*x+c))^(3/2)/a^4/d
```

3.104.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 3.11 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.64

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{15(iA+B)\operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{1}{2}(1+i \tan(c+dx))\right)}{(a+ia \tan(c+dx))^{5/2}}$$

input `Integrate[(Tan[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `(15*(I*A + B)*Hypergeometric2F1[-1/2, 1, 1/2, (1 + I*Tan[c + d*x])/2]*(-I + Tan[c + d*x])^2 - 2*((151*I)*A - 361*B - 5*(77*A + (179*I)*B)*Tan[c + d*x] + 60*((-5*I)*A + 11*B)*Tan[c + d*x]^2 + 20*(3*A + (5*I)*B)*Tan[c + d*x]^3 + 20*B*Tan[c + d*x]^4)/(60*a^2*d*(-I + Tan[c + d*x])^2*Sqrt[a + I*a*Tan[c + d*x]])`

3.104.3 Rubi [A] (verified)

Time = 1.52 (sec) , antiderivative size = 272, normalized size of antiderivative = 1.07, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 4078, 27, 3042, 4078, 27, 3042, 4078, 27, 3042, 4075, 3042, 4010, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^4(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+iA) \tan^4(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\tan^3(c+dx)(8a(iA-B)+a(3A+13iB) \tan(c+dx))}{2(i \tan(c+dx)a+a)^{3/2}} dx}{5a^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B+iA) \tan^4(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\tan^3(c+dx)(8a(iA-B)+a(3A+13iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^{3/2}} dx}{10a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B+iA) \tan^4(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\tan(c+dx)^3(8a(iA-B)+a(3A+13iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^{3/2}} dx}{10a^2} \\
 & \quad \downarrow \text{4078}
 \end{aligned}$$

3.104. $\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{(-B + iA) \tan^4(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\int -\frac{3 \tan^2(c+dx)(2a^2(11A+21iB)-a^2(17iA-47B) \tan(c+dx))}{2\sqrt{i \tan(c+dx)a+a}} dx}{3a^2} - \frac{a(11A+21iB) \tan^3(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \frac{10a^2}{27} \\
 & \frac{(-B + iA) \tan^4(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\int \frac{\tan^2(c+dx)(2a^2(11A+21iB)-a^2(17iA-47B) \tan(c+dx))}{\sqrt{i \tan(c+dx)a+a}} dx}{2a^2} - \frac{a(11A+21iB) \tan^3(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \frac{10a^2}{3042} \\
 & \frac{(-B + iA) \tan^4(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\int \frac{\tan(c+dx)^2(2a^2(11A+21iB)-a^2(17iA-47B) \tan(c+dx))}{\sqrt{i \tan(c+dx)a+a}} dx}{2a^2} - \frac{a(11A+21iB) \tan^3(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \frac{10a^2}{4078} \\
 & \frac{(-B + iA) \tan^4(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\frac{a^2(-89B+39iA) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \int \frac{1}{2} \tan(c+dx) \sqrt{i \tan(c+dx)a+a} (4(39iA-89B)a^3 + (151A+361iB) \tan(c+dx)a^3) dx}{a^2}}{2a^2} - \frac{a(11A+21iB) \tan^3(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \frac{10a^2}{27} \\
 & \frac{(-B + iA) \tan^4(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\frac{a^2(-89B+39iA) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \int \tan(c+dx) \sqrt{i \tan(c+dx)a+a} (4(39iA-89B)a^3 + (151A+361iB) \tan(c+dx)a^3) dx}{2a^2}}{2a^2} - \frac{a(11A+21iB) \tan^3(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \frac{10a^2}{3042} \\
 & \frac{(-B + iA) \tan^4(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\frac{a^2(-89B+39iA) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \int \tan(c+dx) \sqrt{i \tan(c+dx)a+a} (4(39iA-89B)a^3 + (151A+361iB) \tan(c+dx)a^3) dx}{2a^2}}{2a^2} - \frac{a(11A+21iB) \tan^3(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \frac{10a^2}{4075} \\
 & \frac{(-B + iA) \tan^4(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\frac{a^2(-89B+39iA) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \int \sqrt{i \tan(c+dx)a+a} (4a^3(39iA-89B) \tan(c+dx) - a^3(151A+361iB)) dx}{2a^2} - \frac{2a^2(-361B+151iA)(a+ia \tan(c+dx))^{3/2}}{3d}}{2a^2} - \frac{a(11A+21iB) \tan^3(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \frac{10a^2}{10a^2}
 \end{aligned}$$

3.104. $\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{(-B + iA) \tan^4(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \\ & \frac{\frac{a^2(-89B+39iA) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{\int \sqrt{i \tan(c+dx)a+a} (4a^3(39iA-89B) \tan(c+dx) - a^3(151A+361iB)) dx - \frac{2a^2(-361B+151iA)(a+ia \tan(c+dx))^{3/2}}{3d}}{2a^2}}{10a^2} - \frac{a(11A+2iB)}{3d(a+ia)} \end{aligned}$$

$$\begin{aligned} & \downarrow 4010 \\ & \frac{(-B + iA) \tan^4(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \\ & \frac{\frac{a^2(-89B+39iA) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{5a^3(A-iB) \int \sqrt{i \tan(c+dx)a+adx} + \frac{8a^3(-89B+39iA)\sqrt{a+ia \tan(c+dx)}}{d} - \frac{2a^2(-361B+151iA)(a+ia \tan(c+dx))^{3/2}}{3d}}{2a^2}}{10a^2} - \frac{a(11A+2iB)}{3d(a+ia)} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{(-B + iA) \tan^4(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \\ & \frac{\frac{a^2(-89B+39iA) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{5a^3(A-iB) \int \sqrt{i \tan(c+dx)a+adx} + \frac{8a^3(-89B+39iA)\sqrt{a+ia \tan(c+dx)}}{d} - \frac{2a^2(-361B+151iA)(a+ia \tan(c+dx))^{3/2}}{3d}}{2a^2}}{10a^2} - \frac{a(11A+2iB)}{3d(a+ia)} \end{aligned}$$

$$\begin{aligned} & \downarrow 3961 \\ & \frac{(-B + iA) \tan^4(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \\ & \frac{\frac{a^2(-89B+39iA) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{10ia^4(A-iB) \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a} + \frac{8a^3(-89B+39iA)\sqrt{a+ia \tan(c+dx)}}{d} - \frac{2a^2(-361B+151iA)(a+ia \tan(c+dx))^{3/2}}{3d}}{2a^2}}{10a^2} - \frac{a(11A+2iB)}{3d(a+ia)} \end{aligned}$$

$$\begin{aligned} & \downarrow 219 \\ & \frac{(-B + iA) \tan^4(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \\ & \frac{\frac{a^2(-89B+39iA) \tan^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{5i\sqrt{2}a^{7/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} + \frac{8a^3(-89B+39iA)\sqrt{a+ia \tan(c+dx)}}{d} - \frac{2a^2(-361B+151iA)(a+ia \tan(c+dx))^{3/2}}{3d}}{2a^2}}{10a^2} - \frac{a(11A+2iB)}{3d(a+ia)} \end{aligned}$$

input `Int[(Tan[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2), x]`

```
output ((I*A - B)*Tan[c + d*x]^4)/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) - (-1/3*(a*(
11*A + (21*I)*B)*Tan[c + d*x]^3)/(d*(a + I*a*Tan[c + d*x])^(3/2)) + ((a^2*
((39*I)*A - 89*B)*Tan[c + d*x]^2)/(d*Sqrt[a + I*a*Tan[c + d*x]]) - (((-5*I
)*Sqrt[2]*a^(7/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sq
rt[a])))/d + (8*a^3*((39*I)*A - 89*B)*Sqrt[a + I*a*Tan[c + d*x]])/d - (2*a
^2*((151*I)*A - 361*B)*(a + I*a*Tan[c + d*x])^(3/2))/(3*d)/(2*a^2))/(2*a
^2))/(10*a^2)
```

3.104.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 219 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3961 Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d)
Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a
, b, c, d}, x] && EqQ[a^2 + b^2, 0]
```

```
rule 4010 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp
[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e
, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]
```

```
rule 4075 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_)
+ (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f
*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

```
rule 4078 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-(A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.104.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 181, normalized size of antiderivative = 0.71

method	result
derivativedivides	$2i \left(-\frac{iB(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + 3iaB \sqrt{a+ia \tan(dx+c)} + aA \sqrt{a+ia \tan(dx+c)} + \frac{a^2(31iB+17A)}{8\sqrt{a+ia \tan(dx+c)}} - \frac{a^3(9iB+7A)}{12(a+ia \tan(dx+c))^{\frac{3}{2}}} \right) + \frac{\dots}{da^4}$
default	$2i \left(-\frac{iB(a+ia \tan(dx+c))^{\frac{3}{2}}}{3} + 3iaB \sqrt{a+ia \tan(dx+c)} + aA \sqrt{a+ia \tan(dx+c)} + \frac{a^2(31iB+17A)}{8\sqrt{a+ia \tan(dx+c)}} - \frac{a^3(9iB+7A)}{12(a+ia \tan(dx+c))^{\frac{3}{2}}} \right) + \frac{\dots}{da^4}$
parts	$2iA \left(\sqrt{a+ia \tan(dx+c)} - \frac{\sqrt{a} \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)} \sqrt{2}}{2\sqrt{a}}\right)}{16} + \frac{17a}{8\sqrt{a+ia \tan(dx+c)}} - \frac{7a^2}{12(a+ia \tan(dx+c))^{\frac{3}{2}}} + \frac{a^3}{10(a+ia \tan(dx+c))^{\frac{5}{2}}} \right) + \frac{\dots}{da^3}$

```
input int(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x,method=_RETUR
NVERBOSE)
```

```
output 2*I/d/a^4*(-1/3*I*B*(a+I*a*tan(d*x+c))^(3/2)+3*I*a*B*(a+I*a*tan(d*x+c))^(1
/2)+a*A*(a+I*a*tan(d*x+c))^(1/2)+1/8*a^2*(31*I*B+17*A)/(a+I*a*tan(d*x+c))^(
1/2)-1/12*a^3*(9*I*B+7*A)/(a+I*a*tan(d*x+c))^(3/2)+1/10*a^4*(A+I*B)/(a+I*
a*tan(d*x+c))^(5/2)-1/16*a^(3/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*
x+c))^(1/2)*2^(1/2)/a^(1/2)))
```

3.104.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 457 vs. $2(198) = 396$.

Time = 0.26 (sec) , antiderivative size = 457, normalized size of antiderivative = 1.79

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \frac{15\sqrt{\frac{1}{2}}(a^3de^{(7i dx+7i c)} + a^3de^{(5i dx+5i c)})\sqrt{-\frac{A^2-2iAB-B^2}{a^5d^2}}\log\left(-\frac{4}{\dots}\right)}{\dots}$$

input `integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="fracas")`

output `1/120*(15*sqrt(1/2)*(a^3*d*e^(7*I*d*x + 7*I*c) + a^3*d*e^(5*I*d*x + 5*I*c))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2))*log(-4*(sqrt(2)*sqrt(1/2)*(a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2)) + (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 15*sqrt(1/2)*(a^3*d*e^(7*I*d*x + 7*I*c) + a^3*d*e^(5*I*d*x + 5*I*c))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2))*log(4*(sqrt(2)*sqrt(1/2)*(a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2)) - (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) + sqrt(2)*((463*I*A - 983*B)*e^(8*I*d*x + 8*I*c) - 3*(-219*I*A + 509*B)*e^(6*I*d*x + 6*I*c) - 12*(-14*I*A + 29*B)*e^(4*I*d*x + 4*I*c) + (-23*I*A + 33*B)*e^(2*I*d*x + 2*I*c) + 3*I*A - 3*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))/(a^3*d*e^(7*I*d*x + 7*I*c) + a^3*d*e^(5*I*d*x + 5*I*c))`

3.104.6 Sympy [F]

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\tan^4(c+dx)}{(ia(\tan(c+dx)-i))^{5/2}} dx$$

input `integrate(tan(d*x+c)**4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**4/(I*a*(tan(c + d*x) - I))**(5/2), x)`

3.104.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.73

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \frac{i \left(15\sqrt{2}(A-iB)a^{5/2} \log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia\tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia\tan(dx+c)+a}}\right) - 160i(ia\tan(c+dx)) \right)}{(a+ia\tan(c+dx))^{5/2}}$$

```
input integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")
```

```
output 1/240*I*(15*sqrt(2)*(A - I*B)*a^(5/2)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - 160*I*(I*a*tan(d*x + c) + a)^(3/2)*B*a + 480*sqrt(I*a*tan(d*x + c) + a)*(A + 3*I*B)*a^2 + 4*(15*(I*a*tan(d*x + c) + a)^2*(17*A + 31*I*B)*a^3 - 10*(I*a*tan(d*x + c) + a)*(7*A + 9*I*B)*a^4 + 12*(A + I*B)*a^5)/(I*a*tan(d*x + c) + a)^(5/2)/(a^5*d)
```

3.104.8 Giac [F]

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^4}{(ia\tan(dx+c)+a)^{5/2}} dx$$

```
input integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")
```

```
output integrate((B*tan(d*x + c) + A)*tan(d*x + c)^4/(I*a*tan(d*x + c) + a)^(5/2), x)
```

3.104.9 Mupad [B] (verification not implemented)

Time = 8.55 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.09

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \frac{\frac{A \operatorname{li}}{5d} - \frac{A(a+a\tan(c+dx)\operatorname{li})7i}{6ad} + \frac{A(a+a\tan(c+dx)\operatorname{li})^2 17i}{4a^2 d}}{(a+a\tan(c+dx)\operatorname{li})^{5/2}}$$

$$+ \frac{A\sqrt{a+a\tan(c+dx)\operatorname{li}}2i}{a^3 d} - \frac{6B\sqrt{a+a\tan(c+dx)\operatorname{li}}}{a^3 d}$$

$$+ \frac{2B(a+a\tan(c+dx)\operatorname{li})^{3/2}}{3a^4 d} - \frac{\frac{Ba^2}{5} + \frac{31B(a+a\tan(c+dx)\operatorname{li})^2}{4} - \frac{3Ba(a+a\tan(c+dx)\operatorname{li})}{2}}{a^2 d(a+a\tan(c+dx)\operatorname{li})^{5/2}}$$

$$+ \frac{\sqrt{2}A\operatorname{atan}\left(\frac{\sqrt{2}\sqrt{a+a\tan(c+dx)\operatorname{li}}}{2\sqrt{-a}}\right)\operatorname{li}}{8(-a)^{5/2}d} + \frac{\sqrt{2}B\operatorname{atan}\left(\frac{\sqrt{2}\sqrt{a+a\tan(c+dx)\operatorname{li}}}{2\sqrt{-a}}\right)\operatorname{li}}{8a^{5/2}d}$$

input `int((tan(c + d*x)^4*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2),x)`output `((A*1i)/(5*d) - (A*(a + a*tan(c + d*x)*1i)*7i)/(6*a*d) + (A*(a + a*tan(c + d*x)*1i)^2*17i)/(4*a^2*d))/(a + a*tan(c + d*x)*1i)^(5/2) + (A*(a + a*tan(c + d*x)*1i)^(1/2)*2i)/(a^3*d) - (6*B*(a + a*tan(c + d*x)*1i)^(1/2))/(a^3*d) + (2*B*(a + a*tan(c + d*x)*1i)^(3/2))/(3*a^4*d) - ((B*a^2)/5 + (31*B*(a + a*tan(c + d*x)*1i)^2)/4 - (3*B*a*(a + a*tan(c + d*x)*1i))/2)/(a^2*d*(a + a*tan(c + d*x)*1i)^(5/2)) + (2^(1/2)*A*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2)))*1i)/(8*(-a)^(5/2)*d) + (2^(1/2)*B*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)*1i)/(2*a^(1/2)))*1i)/(8*a^(5/2)*d)`

3.105
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$$

3.105.1 Optimal result 1136
 3.105.2 Mathematica [A] (verified) 1136
 3.105.3 Rubi [A] (verified) 1137
 3.105.4 Maple [A] (verified) 1140
 3.105.5 Fricas [B] (verification not implemented) 1141
 3.105.6 Sympy [F] 1141
 3.105.7 Maxima [A] (verification not implemented) 1142
 3.105.8 Giac [F] 1142
 3.105.9 Mupad [B] (verification not implemented) 1143

3.105.1 Optimal result

Integrand size = 36, antiderivative size = 211

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{4\sqrt{2}a^{5/2}d} + \frac{(iA-B) \tan^3(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} + \frac{(7A+17iB) \tan^2(c+dx)}{30ad(a+ia \tan(c+dx))^{3/2}} + \frac{41A+151iB}{60a^2d\sqrt{a+ia \tan(c+dx)}} + \frac{(13A+83iB)\sqrt{a+ia \tan(c+dx)}}{30a^3d}$$

output `1/8*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(5/2)/d*2^(1/2)+1/60*(41*A+151*I*B)/a^2/d/(a+I*a*tan(d*x+c))^(1/2)+1/30*(13*A+83*I*B)*(a+I*a*tan(d*x+c))^(1/2)/a^3/d+1/5*(I*A-B)*tan(d*x+c)^3/d/(a+I*a*tan(d*x+c))^(5/2)+1/30*(7*A+17*I*B)*tan(d*x+c)^2/a/d/(a+I*a*tan(d*x+c))^(3/2)`

3.105.2 Mathematica [A] (verified)

Time = 3.88 (sec) , antiderivative size = 204, normalized size of antiderivative = 0.97

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{-240a^{5/2}(A+5iB) \tan^2(c+dx) + 240a^{5/2}B \tan^3(c+dx) + i(7A+17iB) \tan^2(c+dx)}{(a+ia \tan(c+dx))^{5/2}}$$

input `Integrate[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output $(-240*a^{(5/2)}*(A + (5*I)*B)*\text{Tan}[c + d*x]^2 + 240*a^{(5/2)}*B*\text{Tan}[c + d*x]^3 + I*(72*a^{(5/2)}*((3*I)*A - 13*B) + 20*a^{(5/2)}*(19*A + (77*I)*B)*(-I + \text{Tan}[c + d*x]) + 30*\text{Sqrt}[a]*(I*A + B)*(a + I*a*\text{Tan}[c + d*x])^2 - 15*\text{Sqrt}[2]*(I*A + B)*\text{ArcTanh}[\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]/(\text{Sqrt}[2]*\text{Sqrt}[a])]*(a + I*a*\text{Tan}[c + d*x])^{(5/2)})/(120*a^{(5/2)}*d*(a + I*a*\text{Tan}[c + d*x])^{(5/2)})$

3.105.3 Rubi [A] (verified)

Time = 1.15 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.04, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.361$, Rules used = {3042, 4078, 27, 3042, 4078, 27, 3042, 4075, 3042, 4009, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(c+dx)^3(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{4078} \\ & \frac{(-B+IA) \tan^3(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\tan^2(c+dx)(6a(iA-B)+a(A+11iB) \tan(c+dx))}{2(i \tan(c+dx)a+a)^{3/2}} dx}{5a^2} \\ & \quad \downarrow \text{27} \\ & \frac{(-B+IA) \tan^3(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\tan^2(c+dx)(6a(iA-B)+a(A+11iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^{3/2}} dx}{10a^2} \\ & \quad \downarrow \text{3042} \\ & \frac{(-B+IA) \tan^3(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\tan(c+dx)^2(6a(iA-B)+a(A+11iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^{3/2}} dx}{10a^2} \\ & \quad \downarrow \text{4078} \\ & \frac{(-B+IA) \tan^3(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int -\frac{\tan(c+dx)(4a^2(7A+17iB)-a^2(13iA-83B) \tan(c+dx))}{2\sqrt{i \tan(c+dx)a+a}} dx}{3a^2} - \frac{a(7A+17iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}}{10a^2} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.105. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\frac{(-B + iA) \tan^3(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\int \frac{\tan(c+dx)(4a^2(7A+17iB) - a^2(13iA-83B) \tan(c+dx))}{\sqrt{i \tan(c+dx)a+a}} dx}{6a^2} - \frac{a(7A+17iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 3042

$$\frac{(-B + iA) \tan^3(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\int \frac{\tan(c+dx)(4a^2(7A+17iB) - a^2(13iA-83B) \tan(c+dx))}{\sqrt{i \tan(c+dx)a+a}} dx}{6a^2} - \frac{a(7A+17iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 4075

$$\frac{(-B + iA) \tan^3(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\int \frac{(13iA-83B)a^2+4(7A+17iB) \tan(c+dx)a^2}{\sqrt{i \tan(c+dx)a+a}} dx - \frac{2a(13A+83iB)\sqrt{a+ia \tan(c+dx)}}{d}}{6a^2} - \frac{a(7A+17iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

10a²

↓ 3042

$$\frac{(-B + iA) \tan^3(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\int \frac{(13iA-83B)a^2+4(7A+17iB) \tan(c+dx)a^2}{\sqrt{i \tan(c+dx)a+a}} dx - \frac{2a(13A+83iB)\sqrt{a+ia \tan(c+dx)}}{d}}{6a^2} - \frac{a(7A+17iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

10a²

↓ 4009

$$-\frac{15}{2} a(B+iA) \int \sqrt{i \tan(c+dx)a+adx} - \frac{a^2(41A+15iB)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{2a(13A+83iB)\sqrt{a+ia \tan(c+dx)}}{d}}{6a^2} - \frac{a(7A+17iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

10a²

↓ 3042

$$-\frac{15}{2} a(B+iA) \int \sqrt{i \tan(c+dx)a+adx} - \frac{a^2(41A+15iB)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{2a(13A+83iB)\sqrt{a+ia \tan(c+dx)}}{d}}{6a^2} - \frac{a(7A+17iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

10a²

↓ 3961

$$\frac{15ia^2(B+iA) \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a} - \frac{a^2(41A+15iB)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{2a(13A+83iB)\sqrt{a+ia \tan(c+dx)}}{d}}{6a^2} - \frac{a(7A+17iB) \tan^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

10a²

↓ 219

3.105. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\frac{(-B + iA) \tan^3(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{15ia^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}d} - \frac{a^2(41A+151iB)}{6a^2} - \frac{2a(13A+83iB)\sqrt{a+ia \tan(c+dx)}}{d} - \frac{a(7A+17iB)\tan^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

$10a^2$

input `Int[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `((I*A - B)*Tan[c + d*x]^3)/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) - (-1/3*(a*(7*A + (17*I)*B)*Tan[c + d*x]^2)/(d*(a + I*a*Tan[c + d*x])^(3/2)) + (((15*I)*a^(3/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a]])/(Sqrt[2]*d) - (a^2*(41*A + (151*I)*B))/(d*Sqrt[a + I*a*Tan[c + d*x]]) - (2*a*(13*A + (83*I)*B)*Sqrt[a + I*a*Tan[c + d*x]]/d)/(6*a^2))/(10*a^2)`

3.105.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4078 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.105.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.67

method	result
derivativedivides	$\frac{2iB\sqrt{a+ia \tan(dx+c)} + \frac{a(17iB+7A)}{4\sqrt{a+ia \tan(dx+c)}} - \frac{a^2(7iB+5A)}{6(a+ia \tan(dx+c))^{\frac{3}{2}}} + \frac{a^3(iB+A)}{5(a+ia \tan(dx+c))^{\frac{5}{2}}} + \frac{\sqrt{a}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{2\sqrt{a}}\right)}{8}}{a^3d}$
default	$\frac{2iB\sqrt{a+ia \tan(dx+c)} + \frac{a(17iB+7A)}{4\sqrt{a+ia \tan(dx+c)}} - \frac{a^2(7iB+5A)}{6(a+ia \tan(dx+c))^{\frac{3}{2}}} + \frac{a^3(iB+A)}{5(a+ia \tan(dx+c))^{\frac{5}{2}}} + \frac{\sqrt{a}(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{2\sqrt{a}}\right)}{8}}{a^3d}$
parts	$\frac{2A \left(\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{16\sqrt{a}} + \frac{7}{8\sqrt{a+ia \tan(dx+c)}} - \frac{5a}{12(a+ia \tan(dx+c))^{\frac{3}{2}}} + \frac{a^2}{10(a+ia \tan(dx+c))^{\frac{5}{2}}} \right) + 2iB \left(\sqrt{a+ia \tan(dx+c)} \right)}{da^2}$

input `int(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2), x, method=_RETURNVERBOSE)`

output `2/d/a^3*(I*B*(a+I*a*tan(d*x+c))^(1/2)+1/8*a*(7*A+17*I*B)/(a+I*a*tan(d*x+c))^(1/2)-1/12*a^2*(5*A+7*I*B)/(a+I*a*tan(d*x+c))^(3/2)+1/10*a^3*(A+I*B)/(a+I*a*tan(d*x+c))^(5/2)+1/16*a^(1/2)*(A-I*B)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2)))`

3.105.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 392 vs. $2(162) = 324$.

Time = 0.26 (sec) , antiderivative size = 392, normalized size of antiderivative = 1.86

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx =$$

$$\left(15\sqrt{\frac{1}{2}}a^3d\sqrt{\frac{A^2-2iAB-B^2}{a^5d^2}}e^{(5i dx+5i c)} \log\left(-\frac{4\left(\sqrt{2}\sqrt{\frac{1}{2}}(ia^3de^{(2i dx+2i c)}+ia^3d)\sqrt{\frac{a}{e^{(2i dx+2i c)}+1}}\sqrt{\frac{A^2-2iAB-B^2}{a^5d^2}}+(-iA-B)ae^{(2i dx+2i c)}\right)}{iA+B}\right) \right)$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="fracas")`

output `-1/120*(15*sqrt(1/2)*a^3*d*sqrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(I*a^3*d*e^(2*I*d*x + 2*I*c) + I*a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2)) + (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 15*sqrt(1/2)*a^3*d*sqrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(-I*a^3*d*e^(2*I*d*x + 2*I*c) - I*a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2)) + (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - sqrt(2)*((83*A + 463*I*B)*e^(6*I*d*x + 6*I*c) + 2*(32*A + 97*I*B)*e^(4*I*d*x + 4*I*c) - 2*(8*A + 13*I*B)*e^(2*I*d*x + 2*I*c) + 3*A + 3*I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*e^(-5*I*d*x - 5*I*c)/(a^3*d)`

3.105.6 Sympy [F]

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\tan^3(c+dx)}{(ia(\tan(c+dx)-i))^{5/2}} dx$$

input `integrate(tan(d*x+c)**3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**3/(I*a*(tan(c + d*x) - I))**(5/2), x)`

3.105. $\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx$

3.105.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.77

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx =$$

$$\frac{15\sqrt{2}(A-iB)a^{\frac{3}{2}}\log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia\tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia\tan(dx+c)+a}}\right)-480i\sqrt{ia\tan(dx+c)+a}Ba-\frac{4(15(ia\tan(dx+c)+a)^2(7A+)}{240a^4d}}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `-1/240*(15*sqrt(2)*(A - I*B)*a^(3/2)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a))) - 480*I*sqrt(I*a*tan(d*x + c) + a)*B*a - 4*(15*(I*a*tan(d*x + c) + a)^2*(7*A + 17*I*B))*a^2 - 10*(I*a*tan(d*x + c) + a)*(5*A + 7*I*B)*a^3 + 12*(A + I*B)*a^4)/(I*a*tan(d*x + c) + a)^(5/2))/(a^4*d)`

3.105.8 Giac [F]

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^3}{(ia\tan(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^3/(I*a*tan(d*x + c) + a)^(5/2), x)`

3.105.9 Mupad [B] (verification not implemented)

Time = 7.68 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.09

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \frac{\frac{B \operatorname{li}}{5d} - \frac{B(a+a\tan(c+dx)\operatorname{li})7i}{6ad} + \frac{B(a+a\tan(c+dx)\operatorname{li})^2 17i}{4a^2 d}}{(a+a\tan(c+dx)\operatorname{li})^{5/2}} + \frac{B\sqrt{a+a\tan(c+dx)\operatorname{li}}2i}{a^3 d} + \frac{\frac{Aa^2}{5} + \frac{7A(a+a\tan(c+dx)\operatorname{li})^2}{4} - \frac{5Aa(a+a\tan(c+dx)\operatorname{li})}{6}}{a^2 d(a+a\tan(c+dx)\operatorname{li})^{5/2}} + \frac{\sqrt{2}B\operatorname{atan}\left(\frac{\sqrt{2}\sqrt{a+a\tan(c+dx)\operatorname{li}}}{2\sqrt{-a}}\right)\operatorname{li}}{8(-a)^{5/2}d} + \frac{\sqrt{2}A\operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{a+a\tan(c+dx)\operatorname{li}}}{2\sqrt{a}}\right)}{8a^{5/2}d}$$

input `int((tan(c + d*x))^3*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2),x)`output `((B*1i)/(5*d) - (B*(a + a*tan(c + d*x)*1i)*7i)/(6*a*d) + (B*(a + a*tan(c + d*x)*1i)^2*17i)/(4*a^2*d))/(a + a*tan(c + d*x)*1i)^(5/2) + (B*(a + a*tan(c + d*x)*1i)^(1/2)*2i)/(a^3*d) + ((A*a^2)/5 + (7*A*(a + a*tan(c + d*x)*1i)^2)/4 - (5*A*a*(a + a*tan(c + d*x)*1i))/6)/(a^2*d*(a + a*tan(c + d*x)*1i)^(5/2)) + (2^(1/2)*B*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2)))*1i)/(8*(-a)^(5/2)*d) + (2^(1/2)*A*atanh((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*a^(1/2))))/(8*a^(5/2)*d)`

3.106
$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$$

3.106.1 Optimal result 1144
 3.106.2 Mathematica [A] (verified) 1144
 3.106.3 Rubi [A] (verified) 1145
 3.106.4 Maple [A] (verified) 1148
 3.106.5 Fracas [B] (verification not implemented) 1148
 3.106.6 Sympy [F] 1149
 3.106.7 Maxima [A] (verification not implemented) 1149
 3.106.8 Giac [F] 1150
 3.106.9 Mupad [B] (verification not implemented) 1150

3.106.1 Optimal result

Integrand size = 36, antiderivative size = 167

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{4\sqrt{2}a^{5/2}d} + \frac{(iA-B) \tan^2(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} + \frac{3iA-13B}{30ad(a+ia \tan(c+dx))^{3/2}} - \frac{iA-31B}{20a^2d\sqrt{a+ia \tan(c+dx)}}$$

```
output 1/8*(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(5/2)/
d*2^(1/2)+1/20*(-I*A+31*B)/a^2/d/(a+I*a*tan(d*x+c))^(1/2)+1/5*(I*A-B)*tan(
d*x+c)^2/d/(a+I*a*tan(d*x+c))^(5/2)+1/30*(3*I*A-13*B)/a/d/(a+I*a*tan(d*x+c
))^3/2
```

3.106.2 Mathematica [A] (verified)

Time = 2.22 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.06

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = -\frac{2B \tan^2(c+dx)}{d(a+ia \tan(c+dx))^{5/2}} + i \left(\frac{15\sqrt{2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{a^{3/2}} - \frac{24a(A-9iB)}{(a+ia \tan(c+dx))^{5/2}} + \frac{20(3A-19iB)}{(a+ia \tan(c+dx))^{3/2}} - \frac{30(A-iB)}{a\sqrt{a+ia \tan(c+dx)}} \right) + \frac{}{120ad}$$

input `Integrate[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `(-2*B*Tan[c + d*x]^2)/(d*(a + I*a*Tan[c + d*x])^(5/2)) + ((I/120)*((15*sqrt[2]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(sqrt[2]*sqrt[a])])/a^(3/2) - (24*a*(A - (9*I)*B))/(a + I*a*Tan[c + d*x])^(5/2) + (20*(3*A - (19*I)*B))/(a + I*a*Tan[c + d*x])^(3/2) - (30*(A - I*B))/(a*sqrt[a + I*a*Tan[c + d*x]])))/(a*d)`

3.106.3 Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.07, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {3042, 4078, 27, 3042, 4073, 3042, 4009, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^2(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+ia) \tan^2(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\tan(c+dx)(4a(iA-B)-a(A-9iB) \tan(c+dx))}{2(i \tan(c+dx)a+a)^{3/2}} dx}{5a^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B+ia) \tan^2(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\tan(c+dx)(4a(iA-B)-a(A-9iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^{3/2}} dx}{10a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B+ia) \tan^2(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\tan(c+dx)(4a(iA-B)-a(A-9iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^{3/2}} dx}{10a^2} \\
 & \quad \downarrow \text{4073}
 \end{aligned}$$

$$\begin{aligned}
& \frac{(-B + iA) \tan^2(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{i \int \frac{a^2(3iA - 13B) - 2a^2(A - 9iB) \tan(c + dx)}{\sqrt{i \tan(c + dx)a + a}} dx}{2a^2} - \frac{a(-13B + 3iA)}{3d(a + ia \tan(c + dx))^{3/2}} \\
& \qquad \qquad \qquad \downarrow 3042 \\
& \frac{(-B + iA) \tan^2(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{i \int \frac{a^2(3iA - 13B) - 2a^2(A - 9iB) \tan(c + dx)}{\sqrt{i \tan(c + dx)a + a}} dx}{2a^2} - \frac{a(-13B + 3iA)}{3d(a + ia \tan(c + dx))^{3/2}} \\
& \qquad \qquad \qquad \downarrow 4009 \\
& \frac{(-B + iA) \tan^2(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{i \left(\frac{5}{2} a(B + iA) \int \sqrt{i \tan(c + dx)a + a} dx - \frac{a^2(A + 31iB)}{d\sqrt{a + ia \tan(c + dx)}} \right)}{2a^2} - \frac{a(-13B + 3iA)}{3d(a + ia \tan(c + dx))^{3/2}} \\
& \qquad \qquad \qquad \downarrow 3042 \\
& \frac{(-B + iA) \tan^2(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{i \left(\frac{5}{2} a(B + iA) \int \sqrt{i \tan(c + dx)a + a} dx - \frac{a^2(A + 31iB)}{d\sqrt{a + ia \tan(c + dx)}} \right)}{2a^2} - \frac{a(-13B + 3iA)}{3d(a + ia \tan(c + dx))^{3/2}} \\
& \qquad \qquad \qquad \downarrow 3961 \\
& \frac{(-B + iA) \tan^2(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{i \left(-\frac{5ia^2(B + iA) \int \frac{1}{a - ia \tan(c + dx)} d\sqrt{i \tan(c + dx)a + a}}{d} - \frac{a^2(A + 31iB)}{d\sqrt{a + ia \tan(c + dx)}} \right)}{2a^2} - \frac{a(-13B + 3iA)}{3d(a + ia \tan(c + dx))^{3/2}} \\
& \qquad \qquad \qquad \downarrow 219 \\
& \frac{(-B + iA) \tan^2(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{i \left(-\frac{5ia^{3/2}(B + iA) \operatorname{arctanh}\left(\frac{\sqrt{a + ia \tan(c + dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}d} - \frac{a^2(A + 31iB)}{d\sqrt{a + ia \tan(c + dx)}} \right)}{2a^2} - \frac{a(-13B + 3iA)}{3d(a + ia \tan(c + dx))^{3/2}}
\end{aligned}$$

input `Int[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `((I*A - B)*Tan[c + d*x]^2)/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) - (-1/3*(a*(3*I)*A - 13*B))/(d*(a + I*a*Tan[c + d*x])^(3/2)) - ((I/2)*((-5*I)*a^(3/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a]])/(Sqrt[2]*d) - (a^2*(A + (31*I)*B))/(d*Sqrt[a + I*a*Tan[c + d*x]]))/a^2/(10*a^2)`

3.106. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

3.106.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3961 `Int[Sqrt[(a_) + (b_)*tan[(c_) + (d_)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`
- rule 4009 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`
- rule 4073 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-(A*b - a*B))*((a + b*Tan[e + f*x])^m/(2*a^2*f*m)), x] + Simp[1/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[A*b*c + a*B*c + a*A*d + b*B*d + 2*a*B*d*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && EqQ[a^2 + b^2, 0]`
- rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^n_, x_Symbol] := Simp[(-(A*b - a*B))*((a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.106.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.74

method	result
derivativedivides	$2i \left(-\frac{\frac{A}{8} + \frac{7iB}{8}}{\sqrt{a+ia \tan(dx+c)}} + \frac{a(5iB+3A)}{12(a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{a^2(iB+A)}{10(a+ia \tan(dx+c))^{\frac{5}{2}}} - \frac{(-\frac{A}{8} + \frac{iB}{8})\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2\sqrt{a}} \right) \frac{1}{da^2}$
default	$2i \left(-\frac{\frac{A}{8} + \frac{7iB}{8}}{\sqrt{a+ia \tan(dx+c)}} + \frac{a(5iB+3A)}{12(a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{a^2(iB+A)}{10(a+ia \tan(dx+c))^{\frac{5}{2}}} - \frac{(-\frac{A}{8} + \frac{iB}{8})\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{2\sqrt{a}} \right) \frac{1}{da^2}$
parts	$2iA \left(\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{16a^{\frac{3}{2}}} + \frac{1}{4(a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{1}{8a\sqrt{a+ia \tan(dx+c)}} - \frac{a}{10(a+ia \tan(dx+c))^{\frac{5}{2}}} \right) + 2B \left(\frac{\sqrt{2}}{2\sqrt{a}} \right)$

```
input int(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x,method=_RETURNVERBOSE)
```

```
output 2*I/d/a^2*(-(1/8*A+7/8*I*B)/(a+I*a*tan(d*x+c))^(1/2)+1/12*a*(3*A+5*I*B)/(a+I*a*tan(d*x+c))^(3/2)-1/10*a^2*(A+I*B)/(a+I*a*tan(d*x+c))^(5/2)-1/2*(-1/8*A+1/8*I*B)*2^(1/2)/a^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2)))
```

3.106.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 393 vs. 2(128) = 256.

Time = 0.26 (sec) , antiderivative size = 393, normalized size of antiderivative = 2.35

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx =$$

$$\left(15 \sqrt{\frac{1}{2}} a^3 d \sqrt{-\frac{A^2-2iAB-B^2}{a^5 d^2}} e^{(5i dx+5i c)} \log \left(-\frac{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (a^3 d e^{(2i dx+2i c)} + a^3 d) \sqrt{\frac{a}{e^{(2i dx+2i c)}+1}} \sqrt{-\frac{A^2-2iAB-B^2}{a^5 d^2}} + (-iA-B) \right)}{iA+B} \right) \right)$$

```
input integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x,algorithm="fracas")
```

```
output -1/120*(15*sqrt(1/2)*a^3*d*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2))*e^(5*I*d
*x + 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*
sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2)) +
(-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B) - 15*sqrt(1/2)*
a^3*d*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(4*(sq
rt(2)*sqrt(1/2)*(a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt(a/(e^(2*I*d*x + 2
*I*c) + 1))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2)) - (-I*A - B)*a*e^(I*d*x
+ I*c))*e^(-I*d*x - I*c)/(I*A + B) - sqrt(2)*((-3*I*A + 83*B)*e^(6*I*d*x
+ 6*I*c) - 2*(-3*I*A - 32*B)*e^(4*I*d*x + 4*I*c) - 2*(-3*I*A + 8*B)*e^(2*
I*d*x + 2*I*c) - 3*I*A + 3*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))e^(-5*I*d
*x - 5*I*c)/(a^3*d)
```

3.106.6 Sympy [F]

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \int \frac{(A+B \tan(c+dx)) \tan^2(c+dx)}{(ia(\tan(c+dx)-i))^{5/2}} dx$$

```
input integrate(tan(d*x+c)**2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(5/2),x)
```

```
output Integral((A + B*tan(c + d*x))*tan(c + d*x)**2/(I*a*(tan(c + d*x) - I))**(5
/2), x)
```

3.106.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.84

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx =$$

$$i \left(15 \sqrt{2}(A-iB) \sqrt{a} \log \left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia \tan(dx+c)+a}} \right) + \frac{4 \left(15(ia \tan(dx+c)+a)^2(A+7iB)a-10(ia \tan(dx+c)+a)(3A+5iB)a \right)}{(ia \tan(dx+c)+a)^{5/2}} \right)$$

$$240 a^3 d$$

```
input integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algori
thm="maxima")
```

output
$$-1/240*I*(15*\sqrt{2}*(A - I*B)*\sqrt{a}*\log(-(\sqrt{2}*\sqrt{a} - \sqrt{I*a*\tan(dx + c) + a})/(\sqrt{2}*\sqrt{a} + \sqrt{I*a*\tan(dx + c) + a}))) + 4*(15*(I*a*\tan(dx + c) + a)^2*(A + 7*I*B)*a - 10*(I*a*\tan(dx + c) + a)*(3*A + 5*I*B)*a^2 + 12*(A + I*B)*a^3)/(I*a*\tan(dx + c) + a)^{(5/2)}/(a^3*d)$$

3.106.8 Giac [F]

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)^2}{(ia \tan(dx + c) + a)^{5/2}} dx$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^2/(I*a*tan(d*x + c) + a)^(5/2), x)`

3.106.9 Mupad [B] (verification not implemented)

Time = 7.95 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.12

$$\begin{aligned} \int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{5/2}} dx &= \frac{A \operatorname{li}}{20 d (a + a \tan(c + dx) \operatorname{li})^{5/2}} \\ &+ \frac{A \tan(c + dx)^2 \operatorname{li}}{4 d (a + a \tan(c + dx) \operatorname{li})^{5/2}} + \frac{\frac{B a^2}{5} + \frac{7 B (a + a \tan(c + dx) \operatorname{li})^2}{4} - \frac{5 B a (a + a \tan(c + dx) \operatorname{li})}{6}}{a^2 d (a + a \tan(c + dx) \operatorname{li})^{5/2}} \\ &- \frac{\sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2 \sqrt{-a}}\right) \operatorname{li}}{8 (-a)^{5/2} d} + \frac{\sqrt{2} B \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{a + a \tan(c + dx) \operatorname{li}}}{2 \sqrt{a}}\right)}{8 a^{5/2} d} \end{aligned}$$

input `int((tan(c + d*x)^2*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2),x)`

output
$$(A*1i)/(20*d*(a + a*tan(c + d*x)*1i)^(5/2)) + (A*tan(c + d*x)^2*1i)/(4*d*(a + a*tan(c + d*x)*1i)^(5/2)) + ((B*a^2)/5 + (7*B*(a + a*tan(c + d*x)*1i)^2)/4 - (5*B*a*(a + a*tan(c + d*x)*1i))/6)/(a^2*d*(a + a*tan(c + d*x)*1i)^(5/2)) - (2^(1/2)*A*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2)))*1i)/(8*(-a)^(5/2)*d) + (2^(1/2)*B*atanh((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*a^(1/2)))/(8*a^(5/2)*d)$$

3.107
$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$$

3.107.1 Optimal result 1151
 3.107.2 Mathematica [A] (verified) 1151
 3.107.3 Rubi [A] (verified) 1152
 3.107.4 Maple [A] (verified) 1155
 3.107.5 Fricas [B] (verification not implemented) 1155
 3.107.6 Sympy [F] 1156
 3.107.7 Maxima [A] (verification not implemented) 1156
 3.107.8 Giac [F] 1157
 3.107.9 Mupad [B] (verification not implemented) 1157

3.107.1 Optimal result

Integrand size = 34, antiderivative size = 153

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = -\frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{4\sqrt{2}a^{5/2}d} - \frac{A+iB}{5d(a+ia \tan(c+dx))^{5/2}} + \frac{A+3iB}{6ad(a+ia \tan(c+dx))^{3/2}} + \frac{A-iB}{4a^2d\sqrt{a+ia \tan(c+dx)}}$$

output

```
-1/8*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(5/2)
/d*2^(1/2)+1/4*(A-I*B)/a^2/d/(a+I*a*tan(d*x+c))^(1/2)+1/5*(-A-I*B)/d/(a+I*
a*tan(d*x+c))^(5/2)+1/6*(A+3*I*B)/a/d/(a+I*a*tan(d*x+c))^(3/2)
```

3.107.2 Mathematica [A] (verified)

Time = 2.08 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.96

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{i \left(\frac{15\sqrt{2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{a^{3/2}} + \frac{24a(iA-B)}{(a+ia \tan(c+dx))^{5/2}} - \frac{20(iA-B)}{(a+ia \tan(c+dx))^{3/2}} \right)}{120ad}$$

input

```
Integrate[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2)
,x]
```

output $((I/120)*((15*\text{Sqrt}[2]*(I*A + B)*\text{ArcTanh}[\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]]/(\text{Sqrt}[2]*\text{Sqrt}[a]))/a^{(3/2)} + (24*a*(I*A - B))/(a + I*a*\text{Tan}[c + d*x])^{(5/2)} - (20*(I*A - 3*B))/(a + I*a*\text{Tan}[c + d*x])^{(3/2)} - (30*(I*A + B))/(a*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])))/(a*d)$

3.107.3 Rubi [A] (verified)

Time = 0.65 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4073, 3042, 4009, 3042, 3960, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{4073} \\ & -\frac{i \int \frac{a(A+iB)+2aB\tan(c+dx)}{(i\tan(c+dx)a+a)^{3/2}} dx}{2a^2} - \frac{A+iB}{5d(a+ia\tan(c+dx))^{5/2}} \\ & \quad \downarrow \text{3042} \\ & -\frac{i \int \frac{a(A+iB)+2aB\tan(c+dx)}{(i\tan(c+dx)a+a)^{3/2}} dx}{2a^2} - \frac{A+iB}{5d(a+ia\tan(c+dx))^{5/2}} \\ & \quad \downarrow \text{4009} \\ & -\frac{i\left(\frac{1}{2}(A-iB) \int \frac{1}{\sqrt{i\tan(c+dx)a+a}} dx + \frac{a(-3B+iA)}{3d(a+ia\tan(c+dx))^{3/2}}\right)}{2a^2} - \frac{A+iB}{5d(a+ia\tan(c+dx))^{5/2}} \\ & \quad \downarrow \text{3042} \\ & -\frac{i\left(\frac{1}{2}(A-iB) \int \frac{1}{\sqrt{i\tan(c+dx)a+a}} dx + \frac{a(-3B+iA)}{3d(a+ia\tan(c+dx))^{3/2}}\right)}{2a^2} - \frac{A+iB}{5d(a+ia\tan(c+dx))^{5/2}} \\ & \quad \downarrow \text{3960} \end{aligned}$$

$$\begin{aligned}
& \frac{i\left(\frac{1}{2}(A-iB)\left(\frac{\int \sqrt{i \tan(c+dx)a+adx}}{2a} + \frac{i}{d\sqrt{a+ia \tan(c+dx)}}\right) + \frac{a(-3B+iA)}{3d(a+ia \tan(c+dx))^{3/2}}\right)}{\frac{2a^2}{A+iB}} \\
& \qquad \qquad \qquad \frac{2a^2}{5d(a+ia \tan(c+dx))^{5/2}} \\
& \qquad \qquad \qquad \downarrow \text{3042} \\
& \frac{i\left(\frac{1}{2}(A-iB)\left(\frac{\int \sqrt{i \tan(c+dx)a+adx}}{2a} + \frac{i}{d\sqrt{a+ia \tan(c+dx)}}\right) + \frac{a(-3B+iA)}{3d(a+ia \tan(c+dx))^{3/2}}\right)}{\frac{2a^2}{A+iB}} \\
& \qquad \qquad \qquad \frac{2a^2}{5d(a+ia \tan(c+dx))^{5/2}} \\
& \qquad \qquad \qquad \downarrow \text{3961} \\
& \frac{i\left(\frac{1}{2}(A-iB)\left(\frac{i}{d\sqrt{a+ia \tan(c+dx)}} - \frac{i \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a}}{d}\right) + \frac{a(-3B+iA)}{3d(a+ia \tan(c+dx))^{3/2}}\right)}{\frac{2a^2}{A+iB}} \\
& \qquad \qquad \qquad \frac{2a^2}{5d(a+ia \tan(c+dx))^{5/2}} \\
& \qquad \qquad \qquad \downarrow \text{219} \\
& \frac{i\left(\frac{1}{2}(A-iB)\left(\frac{i}{d\sqrt{a+ia \tan(c+dx)}} - \frac{i \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{2}\sqrt{ad}}\right) + \frac{a(-3B+iA)}{3d(a+ia \tan(c+dx))^{3/2}}\right)}{\frac{2a^2}{A+iB}} \\
& \qquad \qquad \qquad \frac{2a^2}{5d(a+ia \tan(c+dx))^{5/2}}
\end{aligned}$$

input `Int[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `-1/5*(A + I*B)/(d*(a + I*a*Tan[c + d*x])^(5/2)) - ((I/2)*((a*(I*A - 3*B))/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) + ((A - I*B)*((-I)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a]])/(Sqrt[2]*Sqrt[a]*d) + I/(d*Sqrt[a + I*a*Tan[c + d*x]])))/2)/a^2`

3.107.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

rule 4073 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(A*b - a*B))*((a + b*Tan[e + f*x])^m/(2*a^2*f*m)), x] + Simp[1/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[A*b*c + a*B*c + a*A*d + b*B*d + 2*a*B*d*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && EqQ[a^2 + b^2, 0]`

3.107.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.79

method	result
derivativedivides	$\frac{2\left(-\frac{A}{4}-\frac{3iB}{4}\right)}{3(a+ia \tan(dx+c))^{\frac{3}{2}}}-\frac{a(iB+A)}{5(a+ia \tan(dx+c))^{\frac{5}{2}}}-\frac{iB-A}{4a\sqrt{a+ia \tan(dx+c)}}-\frac{(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{8a^{\frac{3}{2}}}}$
default	$\frac{2\left(-\frac{A}{4}-\frac{3iB}{4}\right)}{3(a+ia \tan(dx+c))^{\frac{3}{2}}}-\frac{a(iB+A)}{5(a+ia \tan(dx+c))^{\frac{5}{2}}}-\frac{iB-A}{4a\sqrt{a+ia \tan(dx+c)}}-\frac{(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{8a^{\frac{3}{2}}}}$
parts	$A\left(\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{8a^{\frac{5}{2}}}-\frac{1}{5(a+ia \tan(dx+c))^{\frac{5}{2}}}+\frac{1}{4a^2\sqrt{a+ia \tan(dx+c)}}+\frac{1}{6a(a+ia \tan(dx+c))^{\frac{3}{2}}}\right)+\frac{2iB}{d}\left(\frac{\sqrt{a+ia \tan(dx+c)}}{a}\right)$

input `int(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x,method=_RETURNV
ERBOSE)`

output `2/d/a*(-1/3*(-1/4*A-3/4*I*B)/(a+I*a*tan(d*x+c))^(3/2)-1/10*a*(A+I*B)/(a+I*
a*tan(d*x+c))^(5/2)-1/8/a*(-A+I*B)/(a+I*a*tan(d*x+c))^(1/2)-1/16*(A-I*B)/a
^(3/2)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2)))`

3.107.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 391 vs. 2(114) = 228.

Time = 0.27 (sec) , antiderivative size = 391, normalized size of antiderivative = 2.56

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{\left(15 \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{A^2-2iAB-B^2}{a^5 d^2}} e^{(5i dx+5i c)} \log\left(-\frac{4\left(\sqrt{2}\sqrt{\frac{1}{2}}(i a^3 d e^{(2i dx+2i c)}+\dots\right)}{a^5 d^2}\right)\right)}{a^5 d^2}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm
m="fracas")`

```
output 1/120*(15*sqrt(1/2)*a^3*d*sqrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2))*e^(5*I*d*x
+ 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(I*a^3*d*e^(2*I*d*x + 2*I*c) + I*a^3*d
)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2))
+ (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 15*sqrt(1/2)
*a^3*d*sqrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-4*(s
qrt(2)*sqrt(1/2)*(-I*a^3*d*e^(2*I*d*x + 2*I*c) - I*a^3*d)*sqrt(a/(e^(2*I*d
*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2)) + (-I*A - B)*a*e^(
I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) + sqrt(2)*((17*A - 3*I*B)*e^(6*I
*d*x + 6*I*c) + 2*(8*A + 3*I*B)*e^(4*I*d*x + 4*I*c) - 2*(2*A - 3*I*B)*e^(2
*I*d*x + 2*I*c) - 3*A - 3*I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-5*I*
d*x - 5*I*c)/(a^3*d)
```

3.107.6 Sympy [F]

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \int \frac{(A+B \tan(c+dx)) \tan(c+dx)}{(ia(\tan(c+dx)-i))^{5/2}} dx$$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(5/2),x)
```

```
output Integral((A + B*tan(c + d*x))*tan(c + d*x)/(I*a*(tan(c + d*x) - I))**(5/2)
, x)
```

3.107.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.89

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{15\sqrt{2}(A-iB) \log\left(\frac{-\sqrt{2}\sqrt{a}-\sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia \tan(dx+c)+a}}\right)}{\sqrt{a}} + \frac{4\left(15(ia \tan(dx+c)+a)^2(A-iB)+10\right)}{240 a^2 d} (ia \tan(dx+c)+a)$$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm
m="maxima")
```

```
output 1/240*(15*sqrt(2)*(A - I*B)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c)
+ a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a)))/sqrt(a) + 4*(15*(I*a
*tan(d*x + c) + a)^2*(A - I*B) + 10*(I*a*tan(d*x + c) + a)*(A + 3*I*B)*a -
12*(A + I*B)*a^2)/(I*a*tan(d*x + c) + a)^(5/2))/(a^2*d)
```

3.107. $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

3.107.8 Giac [F]

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)}{(ia\tan(dx+c)+a)^{5/2}} dx$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)/(I*a*tan(d*x + c) + a)^(5/2), x)`

3.107.9 Mupad [B] (verification not implemented)

Time = 8.35 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.22

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \frac{-\frac{A}{5} + \frac{A(a+a\tan(c+dx)\text{li})}{6a} + \frac{A(a+a\tan(c+dx)\text{li})^2}{4a^2}}{d(a+a\tan(c+dx)\text{li})^{5/2}} + \frac{B\text{li}}{20d(a+a\tan(c+dx)\text{li})^{5/2}} + \frac{B\tan(c+dx)^2\text{li}}{4d(a+a\tan(c+dx)\text{li})^{5/2}} - \frac{\sqrt{2}B\text{atan}\left(\frac{\sqrt{2}\sqrt{a+a\tan(c+dx)\text{li}}}{2\sqrt{-a}}\right)\text{li}}{8(-a)^{5/2}d} - \frac{\sqrt{2}A\text{atanh}\left(\frac{\sqrt{2}\sqrt{a+a\tan(c+dx)\text{li}}}{2\sqrt{-a}}\right)}{8a^{5/2}d}$$

input `int((tan(c + d*x)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2),x)`

output `((A*(a + a*tan(c + d*x)*1i))/(6*a) - A/5 + (A*(a + a*tan(c + d*x)*1i)^2)/(4*a^2))/(d*(a + a*tan(c + d*x)*1i)^(5/2)) + (B*1i)/(20*d*(a + a*tan(c + d*x)*1i)^(5/2)) + (B*tan(c + d*x)^2*1i)/(4*d*(a + a*tan(c + d*x)*1i)^(5/2)) - (2^(1/2)*B*atan((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2)))*1i)/(8*(-a)^(5/2)*d) - (2^(1/2)*A*atanh((2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*a^(1/2)))/(8*a^(5/2)*d)`

3.108 $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^{5/2}} dx$

3.108.1 Optimal result 1158
 3.108.2 Mathematica [C] (verified) 1158
 3.108.3 Rubi [A] (verified) 1159
 3.108.4 Maple [A] (verified) 1161
 3.108.5 Fracas [B] (verification not implemented) 1162
 3.108.6 Sympy [F] 1162
 3.108.7 Maxima [A] (verification not implemented) 1163
 3.108.8 Giac [F] 1163
 3.108.9 Mupad [B] (verification not implemented) 1163

3.108.1 Optimal result

Integrand size = 28, antiderivative size = 155

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{5/2}} dx = -\frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{4\sqrt{2}a^{5/2}d} + \frac{iA - B}{5d(a + ia \tan(c + dx))^{5/2}} + \frac{iA + B}{6ad(a + ia \tan(c + dx))^{3/2}} + \frac{iA + B}{4a^2d\sqrt{a + ia \tan(c + dx)}}$$

output `-1/8*(I*A+B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(5/2)/d*2^(1/2)+1/4*(I*A+B)/a^2/d/(a+I*a*tan(d*x+c))^(1/2)+1/5*(I*A-B)/d/(a+I*a*tan(d*x+c))^(5/2)+1/6*(I*A+B)/a/d/(a+I*a*tan(d*x+c))^(3/2)`

3.108.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.89 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.49

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{5/2}} dx = \frac{6iA - 6B - 5(A - iB) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{1}{2}(1 + i \tan(c + dx))\right)}{30d(a + ia \tan(c + dx))^{5/2}}$$

input `Integrate[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `((6*I)*A - 6*B - 5*(A - I*B)*Hypergeometric2F1[-3/2, 1, -1/2, (1 + I*Tan[c + d*x])/2]*(-I + Tan[c + d*x]))/(30*d*(a + I*a*Tan[c + d*x])^(5/2))`

3.108. $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^{5/2}} dx$

3.108.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.99, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.321$, Rules used = {3042, 4009, 3042, 3960, 3042, 3960, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{4009} \\
 & \frac{(A - iB) \int \frac{1}{(i \tan(c+dx)a+a)^{3/2}} dx}{2a} + \frac{-B + iA}{5d(a + ia \tan(c + dx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A - iB) \int \frac{1}{(i \tan(c+dx)a+a)^{3/2}} dx}{2a} + \frac{-B + iA}{5d(a + ia \tan(c + dx))^{5/2}} \\
 & \quad \downarrow \text{3960} \\
 & \frac{(A - iB) \left(\frac{\int \frac{1}{\sqrt{i \tan(c+dx)a+a}} dx}{2a} + \frac{i}{3d(a+ia \tan(c+dx))^{3/2}} \right)}{2a} + \frac{-B + iA}{5d(a + ia \tan(c + dx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A - iB) \left(\frac{\int \frac{1}{\sqrt{i \tan(c+dx)a+a}} dx}{2a} + \frac{i}{3d(a+ia \tan(c+dx))^{3/2}} \right)}{2a} + \frac{-B + iA}{5d(a + ia \tan(c + dx))^{5/2}} \\
 & \quad \downarrow \text{3960} \\
 & \frac{(A - iB) \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+adx}}{2a} + \frac{i}{d\sqrt{a+ia \tan(c+dx)}}}{2a} + \frac{i}{3d(a+ia \tan(c+dx))^{3/2}} \right)}{2a} + \frac{-B + iA}{5d(a + ia \tan(c + dx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A - iB) \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+adx}}{2a} + \frac{i}{d\sqrt{a+ia \tan(c+dx)}}}{2a} + \frac{i}{3d(a+ia \tan(c+dx))^{3/2}} \right)}{2a} + \frac{-B + iA}{5d(a + ia \tan(c + dx))^{5/2}}
 \end{aligned}$$

3.108. $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \downarrow \text{3961} \\
 & \frac{(A - iB) \left(\frac{\frac{i}{d\sqrt{a+ia \tan(c+dx)}} - \frac{i \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a}}{2a}}{2a} + \frac{i}{3d(a+ia \tan(c+dx))^{3/2}} \right)}{-B + iA} + \\
 & \frac{2a}{5d(a + ia \tan(c + dx))^{5/2}} \\
 & \downarrow \text{219} \\
 & \frac{(A - iB) \left(\frac{\frac{i}{d\sqrt{a+ia \tan(c+dx)}} - \frac{i \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2\sqrt{a}}}\right)}{2a}}{2a} + \frac{i}{3d(a+ia \tan(c+dx))^{3/2}} \right)}{-B + iA} + \\
 & \frac{2a}{5d(a + ia \tan(c + dx))^{5/2}}
 \end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `(I*A - B)/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) + ((A - I*B)*((I/3)/(d*(a + I*a*Tan[c + d*x])^(3/2)) + (((-I)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a]])/(Sqrt[2]*Sqrt[a]*d) + I/(d*Sqrt[a + I*a*Tan[c + d*x]])))/(2*a)))/(2*a)`

3.108.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

3.108.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.79

method	result
derivativedivides	$2i \left(-\frac{-\frac{A}{2} - \frac{iB}{2}}{5(a+ia \tan(dx+c))^{\frac{5}{2}}} - \frac{iB-A}{12a(a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{iB-A}{8a^2 \sqrt{a+ia \tan(dx+c)}} - \frac{(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{16a^{\frac{5}{2}}} \right) \frac{1}{d}$
default	$2i \left(-\frac{-\frac{A}{2} - \frac{iB}{2}}{5(a+ia \tan(dx+c))^{\frac{5}{2}}} - \frac{iB-A}{12a(a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{iB-A}{8a^2 \sqrt{a+ia \tan(dx+c)}} - \frac{(-iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{16a^{\frac{5}{2}}} \right) \frac{1}{d}$
parts	$2iAa \left(-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{16a^{\frac{7}{2}}} + \frac{1}{8a^3 \sqrt{a+ia \tan(dx+c)}} + \frac{1}{12a^2(a+ia \tan(dx+c))^{\frac{3}{2}}} + \frac{1}{10a(a+ia \tan(dx+c))^{\frac{5}{2}}} \right) \frac{1}{d} + \dots$

input `int((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `2*I/d*(-1/5*(-1/2*A-1/2*I*B)/(a+I*a*tan(d*x+c))^(5/2)-1/12/a*(-A+I*B)/(a+I*a*tan(d*x+c))^(3/2)-1/8/a^2*(-A+I*B)/(a+I*a*tan(d*x+c))^(1/2)-1/16*(A-I*B)/a^(5/2)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))`

3.108.
$$\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^{5/2}} dx$$

3.108.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 392 vs. $2(118) = 236$.

Time = 0.25 (sec) , antiderivative size = 392, normalized size of antiderivative = 2.53

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{5/2}} dx = \frac{\left(15 \sqrt{\frac{1}{2}} a^3 d \sqrt{-\frac{A^2 - 2iAB - B^2}{a^5 d^2}} e^{(5i dx + 5i c)} \log \left(-\frac{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (a^3 d e^{(2i dx + 2i c)} + a^3 d) \sqrt{\frac{1}{e^{(2i dx + 2i c)}}}} \right)}{\right)} \right)}{\left(a + ia \tan(c + dx) \right)^{5/2}}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="fricas")`

output `1/120*(15*sqrt(1/2)*a^3*d*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2)) + (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 15*sqrt(1/2)*a^3*d*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(4*(sqrt(2)*sqrt(1/2)*(a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2)) - (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) + sqrt(2)*((23*I*A + 17*B)*e^(6*I*d*x + 6*I*c) - 2*(-17*I*A - 8*B)*e^(4*I*d*x + 4*I*c) - 2*(-7*I*A + 2*B)*e^(2*I*d*x + 2*I*c) + 3*I*A - 3*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*e^(-5*I*d*x - 5*I*c)/(a^3*d)`

3.108.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{(ia (\tan(c + dx) - i))^{5/2}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))/(I*a*(tan(c + d*x) - I))**(5/2), x)`

3.108.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.90

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{5/2}} dx = \frac{i \left(\frac{15 \sqrt{2}(A - iB) \log\left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c)+a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c)+a}}\right)}{a^{3/2}} + \frac{4 \left(15 (ia \tan(dx+c)+a)^2 (A - iB) + 10 (ia \tan(dx+c)+a)\right)}{(ia \tan(dx+c)+a)^2} \right)}{240 ad}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")`output `1/240*I*(15*sqrt(2)*(A - I*B)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a)))/a^(3/2) + 4*(15*(I*a*tan(d*x + c) + a)^2*(A - I*B) + 10*(I*a*tan(d*x + c) + a)*(A - I*B)*a + 12*(A + I*B)*a^2)/((I*a*tan(d*x + c) + a)^(5/2)*a)/(a*d)`**3.108.8 Giac [F]**

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^{5/2}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`output `integrate((B*tan(d*x + c) + A)/(I*a*tan(d*x + c) + a)^(5/2), x)`**3.108.9 Mupad [B] (verification not implemented)**

Time = 7.61 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.32

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{5/2}} dx = \frac{\frac{A \operatorname{li}}{5d} + \frac{A(a+a \tan(c+dx) \operatorname{li}) \operatorname{li}}{6ad} + \frac{A(a+a \tan(c+dx) \operatorname{li})^2 \operatorname{li}}{4a^2 d}}{(a + a \tan(c + dx) \operatorname{li})^{5/2}} + \frac{-\frac{B}{5} + \frac{B(a+a \tan(c+dx) \operatorname{li})}{6a} + \frac{B(a+a \tan(c+dx) \operatorname{li})^2}{4a^2}}{d(a + a \tan(c + dx) \operatorname{li})^{5/2}} + \frac{\sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2}\sqrt{a+a \tan(c+dx) \operatorname{li}}}{2\sqrt{-a}}\right) \operatorname{li}}{8(-a)^{5/2} d} - \frac{\sqrt{2} B \operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{a+a \tan(c+dx) \operatorname{li}}}{2\sqrt{a}}\right)}{8a^{5/2} d}$$

input `int((A + B*tan(c + d*x))/(a + a*tan(c + d*x)*1i)^(5/2),x)`

output `((A*1i)/(5*d) + (A*(a + a*tan(c + d*x)*1i)*1i)/(6*a*d) + (A*(a + a*tan(c + d*x)*1i)^2*1i)/(4*a^2*d))/(a + a*tan(c + d*x)*1i)^(5/2) + ((B*(a + a*tan(c + d*x)*1i))/(6*a) - B/5 + (B*(a + a*tan(c + d*x)*1i)^2)/(4*a^2))/(d*(a + a*tan(c + d*x)*1i)^(5/2)) + (2^(1/2)*A*atan(2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*(-a)^(1/2)))*1i)/(8*(-a)^(5/2)*d) - (2^(1/2)*B*atanh(2^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2))/(2*a^(1/2)))/(8*a^(5/2)*d)`

3.109
$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$$

3.109.1 Optimal result 1165
 3.109.2 Mathematica [A] (verified) 1165
 3.109.3 Rubi [A] (verified) 1166
 3.109.4 Maple [A] (verified) 1170
 3.109.5 Fricas [B] (verification not implemented) 1171
 3.109.6 Sympy [F] 1172
 3.109.7 Maxima [A] (verification not implemented) 1172
 3.109.8 Giac [F] 1173
 3.109.9 Mupad [B] (verification not implemented) 1173

3.109.1 Optimal result

Integrand size = 34, antiderivative size = 192

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = -\frac{2A \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{a^{5/2}d} + \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{4\sqrt{2}a^{5/2}d} + \frac{A+iB}{5d(a+ia \tan(c+dx))^{5/2}} + \frac{3A+iB}{6ad(a+ia \tan(c+dx))^{3/2}} + \frac{7A+iB}{4a^2d\sqrt{a+ia \tan(c+dx)}}$$

output `-2*A*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/a^(5/2)/d+1/8*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(5/2)/d*2^(1/2)+1/4*(7*A+I*B)/a^2/d/(a+I*a*tan(d*x+c))^(1/2)+1/5*(A+I*B)/d/(a+I*a*tan(d*x+c))^(5/2)+1/6*(3*A+I*B)/a/d/(a+I*a*tan(d*x+c))^(3/2)`

3.109.2 Mathematica [A] (verified)

Time = 3.28 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.13

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{3a^{13/2}(A+iB) + (a+ia \tan(c+dx)) \left(\frac{5}{2}a^{11/2}(3A+iB) + \frac{15}{8}a^4(a \right)}{...}$$

input `Integrate[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `(3*a^(13/2)*(A + I*B) + (a + I*a*Tan[c + d*x])*((5*a^(11/2)*(3*A + I*B))/2 + (15*a^4*(a + I*a*Tan[c + d*x])*(2*Sqrt[a]*(7*A + I*B) - 16*A*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[a + I*a*Tan[c + d*x]] + Sqrt[2]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])]*Sqrt[a + I*a*Tan[c + d*x]]))/8))/(15*a^(13/2)*d*(a + I*a*Tan[c + d*x])^(5/2))`

3.109.3 Rubi [A] (verified)

Time = 1.39 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.10, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)(a+ia \tan(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{5 \cot(c+dx)(2aA-a(iA-B) \tan(c+dx))}{2(i \tan(c+dx)a+a)^{3/2}} dx}{5a^2} + \frac{A+iB}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\cot(c+dx)(2aA-a(iA-B) \tan(c+dx))}{(i \tan(c+dx)a+a)^{3/2}} dx}{2a^2} + \frac{A+iB}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{2aA-a(iA-B) \tan(c+dx)}{\tan(c+dx)(i \tan(c+dx)a+a)^{3/2}} dx}{2a^2} + \frac{A+iB}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow \text{4079}
 \end{aligned}$$

3.109. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{\int \frac{3 \cot(c+dx) (4a^2 A - a^2 (3iA - B) \tan(c+dx))}{2\sqrt{i \tan(c+dx)a+a} \cdot 3a^2} dx + \frac{a(3A+iB)}{3d(a+ia \tan(c+dx))^{3/2}}}{2a^2} + \frac{A+iB}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{\cot(c+dx) (4a^2 A - a^2 (3iA - B) \tan(c+dx))}{\sqrt{i \tan(c+dx)a+a} \cdot 2a^2} dx + \frac{a(3A+iB)}{3d(a+ia \tan(c+dx))^{3/2}}}{2a^2} + \frac{A+iB}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{4a^2 A - a^2 (3iA - B) \tan(c+dx)}{\tan(c+dx) \sqrt{i \tan(c+dx)a+a}} dx + \frac{a(3A+iB)}{3d(a+ia \tan(c+dx))^{3/2}}}{2a^2} + \frac{A+iB}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 4079 \\
 & \frac{\int \frac{\frac{1}{2} \cot(c+dx) \sqrt{i \tan(c+dx)a+a} (8a^3 A - a^3 (7iA - B) \tan(c+dx))}{a^2} dx + \frac{a^2 (7A+iB)}{d\sqrt{a+ia \tan(c+dx)}}}{2a^2} + \frac{a(3A+iB)}{3d(a+ia \tan(c+dx))^{3/2}} + \\
 & \quad \frac{2a^2}{A+iB} \\
 & \quad \frac{A+iB}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{\cot(c+dx) \sqrt{i \tan(c+dx)a+a} (8a^3 A - a^3 (7iA - B) \tan(c+dx))}{2a^2} dx + \frac{a^2 (7A+iB)}{d\sqrt{a+ia \tan(c+dx)}}}{2a^2} + \frac{a(3A+iB)}{3d(a+ia \tan(c+dx))^{3/2}} + \\
 & \quad \frac{2a^2}{A+iB} \\
 & \quad \frac{A+iB}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (8a^3 A - a^3 (7iA - B) \tan(c+dx))}{\tan(c+dx) \cdot 2a^2} dx + \frac{a^2 (7A+iB)}{d\sqrt{a+ia \tan(c+dx)}}}{2a^2} + \frac{a(3A+iB)}{3d(a+ia \tan(c+dx))^{3/2}} + \\
 & \quad \frac{2a^2}{A+iB} \\
 & \quad \frac{A+iB}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 4083 \\
 & \frac{\frac{a^3 (B+iA) \int \sqrt{i \tan(c+dx)a+adx} + 8a^2 A \int \cot(c+dx)(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{2a^2} + \frac{a^2 (7A+iB)}{d\sqrt{a+ia \tan(c+dx)}}}{2a^2} + \frac{a(3A+iB)}{3d(a+ia \tan(c+dx))^{3/2}} + \\
 & \quad \frac{2a^2}{A+iB} \\
 & \quad \frac{A+iB}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 3042
 \end{aligned}$$

3.109. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{a^3(B+iA) \int \sqrt{i \tan(c+dx)a+adx} + 8a^2 A \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a} dx}{\tan(c+dx)}}{2a^2} + \frac{a^2(7A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \frac{a(3A+iB)}{3d(a+ia \tan(c+dx))^{3/2}} + \\
 & \frac{2a^2}{A+iB} \\
 & \frac{2a^2}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \downarrow 3961 \\
 & \frac{8a^2 A \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a} dx}{\tan(c+dx)} - \frac{2ia^4(B+iA) \int \frac{1}{a-ia \tan(c+dx)} d\sqrt{i \tan(c+dx)a+a}}{2a^2}}{2a^2} + \frac{a^2(7A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \frac{a(3A+iB)}{3d(a+ia \tan(c+dx))^{3/2}} + \\
 & \frac{2a^2}{A+iB} \\
 & \frac{2a^2}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \downarrow 219 \\
 & \frac{8a^2 A \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a} dx}{\tan(c+dx)} - \frac{i\sqrt{2}a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{2a^2}}{2a^2} + \frac{a^2(7A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \frac{a(3A+iB)}{3d(a+ia \tan(c+dx))^{3/2}} + \\
 & \frac{2a^2}{A+iB} \\
 & \frac{2a^2}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \downarrow 4082 \\
 & \frac{8a^4 A \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx) - \frac{i\sqrt{2}a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{2a^2}}{2a^2} + \frac{a^2(7A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \frac{a(3A+iB)}{3d(a+ia \tan(c+dx))^{3/2}} + \\
 & \frac{2a^2}{A+iB} \\
 & \frac{2a^2}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \downarrow 73 \\
 & \frac{16ia^3 A \int \frac{1}{i-i \tan(c+dx)a+a} d\sqrt{i \tan(c+dx)a+a} - \frac{i\sqrt{2}a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d}}{2a^2}}{2a^2} + \frac{a^2(7A+iB)}{d\sqrt{a+ia \tan(c+dx)}} + \frac{a(3A+iB)}{3d(a+ia \tan(c+dx))^{3/2}} + \\
 & \frac{2a^2}{A+iB} \\
 & \frac{2a^2}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \downarrow 221
 \end{aligned}$$

3.109. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\frac{\frac{a^2(7A+iB)}{d\sqrt{a+ia\tan(c+dx)}} + \frac{i\sqrt{2}a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{16a^{7/2}A\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{a}}\right)}{d}}{2a^2} + \frac{a(3A+iB)}{3d(a+ia\tan(c+dx))^{3/2}} + \frac{A+iB}{5d(a+ia\tan(c+dx))^{5/2}}$$

input `Int[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `(A + I*B)/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) + ((a*(3*A + I*B))/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) + (((-16*a^(7/2)*A*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d - (I*Sqrt[2]*a^(7/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d)/(2*a^2) + (a^2*(7*A + I*B))/(d*Sqrt[a + I*a*Tan[c + d*x]]))/(2*a^2))/(2*a^2)`

3.109.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 73 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]] , x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

rule 4082 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int((((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.109.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.80

method	result
derivativedivides	$2a \left(-\frac{-iB-7A}{8a^3 \sqrt{a+ia \tan(dx+c)}} - \frac{-iB-3A}{12a^2 (a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{-iB-A}{10a (a+ia \tan(dx+c))^{\frac{5}{2}}} - \frac{(iB-A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{16a^{\frac{7}{2}}} \right) \frac{d}{dx}$
default	$2a \left(-\frac{-iB-7A}{8a^3 \sqrt{a+ia \tan(dx+c)}} - \frac{-iB-3A}{12a^2 (a+ia \tan(dx+c))^{\frac{3}{2}}} - \frac{-iB-A}{10a (a+ia \tan(dx+c))^{\frac{5}{2}}} - \frac{(iB-A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{16a^{\frac{7}{2}}} \right) \frac{d}{dx}$

3.109. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

input `int(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x,method=_RETURNV
ERBOSE)`

output `2/d*a*(-1/8/a^3*(-7*A-I*B)/(a+I*a*tan(d*x+c))^(1/2)-1/12*(-3*A-I*B)/a^2/(a
+I*a*tan(d*x+c))^(3/2)-1/10*(-A-I*B)/a/(a+I*a*tan(d*x+c))^(5/2)-1/16*(-A+I
*B)/a^(7/2)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))-
1/a^(7/2)*A*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2)))`

3.109.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 644 vs. $2(145) = 290$.

Time = 0.27 (sec) , antiderivative size = 644, normalized size of antiderivative = 3.35

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx =$$

$$\left(15 \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{A^2-2iAB-B^2}{a^5 d^2}} e^{(5i dx+5i c)} \log \left(-\frac{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (i a^3 d e^{(2i dx+2i c)} + i a^3 d) \sqrt{\frac{a}{e^{(2i dx+2i c)}+1}} \sqrt{\frac{A^2-2iAB-B^2}{a^5 d^2}} + (-i A-B) a e^{(2i dx+2i c)} \right)}{i A+B} \right) \right)$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm
m="fricas")`

output

```
-1/120*(15*sqrt(1/2)*a^3*d*sqrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(I*a^3*d*e^(2*I*d*x + 2*I*c) + I*a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2)) + (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B) - 15*sqrt(1/2)*a^3*d*sqrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(-I*a^3*d*e^(2*I*d*x + 2*I*c) - I*a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2)) + (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B) + 60*a^3*d*sqrt(A^2/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(16*(3*A*a^2*e^(2*I*d*x + 2*I*c) + A*a^2 + 2*sqrt(2))*(a^4*d*e^(3*I*d*x + 3*I*c) + a^4*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(A^2/(a^5*d^2)))e^(-2*I*d*x - 2*I*c)/A - 60*a^3*d*sqrt(A^2/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(16*(3*A*a^2*e^(2*I*d*x + 2*I*c) + A*a^2 - 2*sqrt(2)*(a^4*d*e^(3*I*d*x + 3*I*c) + a^4*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(A^2/(a^5*d^2)))e^(-2*I*d*x - 2*I*c)/A - sqrt(2)*((123*A + 23*I*B)*e^(6*I*d*x + 6*I*c) + 2*(72*A + 17*I*B)*e^(4*I*d*x + 4*I*c) + 2*(12*A + 7*I*B)*e^(2*I*d*x + 2*I*c) + 3*A + 3*I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*e^(-5*I*d*x - 5*I*c)/(a^3*d)
```

3.109.6 Sympy [F]

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{(A + B \tan(c + dx)) \cot(c + dx)}{(ia (\tan(c + dx) - i))^{5/2}} dx$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)/(I*a*(tan(c + d*x) - I))**(5/2), x)`

3.109.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.97

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{5/2}} dx = \frac{15\sqrt{2}(A - iB) \log\left(-\frac{\sqrt{2}\sqrt{a} - \sqrt{ia \tan(dx+c) + a}}{\sqrt{2}\sqrt{a} + \sqrt{ia \tan(dx+c) + a}}\right)}{a^{5/2}} - \frac{240 A \log\left(\frac{\sqrt{ia \tan(dx+c) + a} - \sqrt{a}}{\sqrt{ia \tan(dx+c) + a} + \sqrt{a}}\right)}{a^{5/2}} - \frac{4 \left(15 (ia \tan(dx+c) + a)^2 (7A + iB) + 10 (ia \tan(dx+c) + a)\right)}{240 d (ia \tan(dx+c) + a)^{5/2} a^2}$$

3.109. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output
$$-1/240*(15*\sqrt{2}*(A - I*B)*\log(-(\sqrt{2}*\sqrt{a} - \sqrt{I*a*\tan(d*x + c) + a))/(\sqrt{2}*\sqrt{a} + \sqrt{I*a*\tan(d*x + c) + a}))/a^{5/2} - 240*A*\log((\sqrt{I*a*\tan(d*x + c) + a} - \sqrt{a})/(\sqrt{I*a*\tan(d*x + c) + a} + \sqrt{a}))/a^{5/2} - 4*(15*(I*a*\tan(d*x + c) + a)^2*(7*A + I*B) + 10*(I*a*\tan(d*x + c) + a)*(3*A + I*B)*a + 12*(A + I*B)*a^2)/((I*a*\tan(d*x + c) + a)^{5/2}*a^2))/d$$

3.109.8 Giac [F]

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)}{(ia\tan(dx+c)+a)^{5/2}} dx$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)/(I*a*tan(d*x + c) + a)^(5/2), x)`

3.109.9 Mupad [B] (verification not implemented)

Time = 7.83 (sec) , antiderivative size = 528, normalized size of antiderivative = 2.75

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \frac{A+B\operatorname{li}}{5d} + \frac{(3A+B\operatorname{li})(a+a\tan(c+dx)\operatorname{li})}{6ad} + \frac{(7A+B\operatorname{li})(a+a\tan(c+dx)\operatorname{li})^2}{4a^2d}$$

$$- \frac{2A\operatorname{atanh}\left(\frac{127A^3ad\sqrt{\frac{1}{a^3}}\sqrt{a+a\tan(c+dx)\operatorname{li}}}{127dA^3+2idA^2B+dAB^2} + \frac{AB^2ad\sqrt{\frac{1}{a^3}}\sqrt{a+a\tan(c+dx)\operatorname{li}}}{127dA^3+2idA^2B+dAB^2} + \frac{A^2Bad\sqrt{\frac{1}{a^3}}\sqrt{a+a\tan(c+dx)\operatorname{li}}}{127dA^3+2idA^2B+dAB^2}\right)\sqrt{\frac{1}{a^3}}}{8ad}$$

$$+ \frac{\sqrt{2}\operatorname{atanh}\left(\frac{\sqrt{2}A^3ad\sqrt{-\frac{1}{a^3}}\sqrt{a+a\tan(c+dx)\operatorname{li}}}{32\left(\frac{127dA^3}{16}-\frac{125idA^2B}{16}+\frac{3dAB^2}{16}-\frac{1idB^3}{16}\right)} + \frac{\sqrt{2}B^3ad\sqrt{-\frac{1}{a^3}}\sqrt{a+a\tan(c+dx)\operatorname{li}}}{32\left(\frac{127dA^3}{16}-\frac{125idA^2B}{16}+\frac{3dAB^2}{16}-\frac{1idB^3}{16}\right)} + \frac{\sqrt{2}AB^2ad\sqrt{-\frac{1}{a^3}}\sqrt{a+a\tan(c+dx)\operatorname{li}}}{32\left(\frac{127dA^3}{16}-\frac{125idA^2B}{16}+\frac{3dAB^2}{16}-\frac{1idB^3}{16}\right)}\right)}{8ad}$$

input `int((cot(c + d*x)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*I)^(5/2),x)`

3.109.
$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx$$

output

$$\begin{aligned}
& ((A + B*1i)/(5*d) + ((3*A + B*1i)*(a + a*\tan(c + d*x)*1i))/(6*a*d) + ((7*A \\
& + B*1i)*(a + a*\tan(c + d*x)*1i)^2)/(4*a^2*d))/(a + a*\tan(c + d*x)*1i)^{5/2} \\
& - (2*A*\operatorname{atanh}((127*A^3*a*d*(1/a^3)^{1/2}*(a + a*\tan(c + d*x)*1i)^{1/2}))/ \\
& (127*A^3*d + A*B^2*d + A^2*B*d*2i) + (A*B^2*a*d*(1/a^3)^{1/2}*(a + a*\tan(c \\
& + d*x)*1i)^{1/2}))/((127*A^3*d + A*B^2*d + A^2*B*d*2i) + (A^2*B*a*d*(1/a^3) \\
& ^{1/2}*(a + a*\tan(c + d*x)*1i)^{1/2}*2i))/((127*A^3*d + A*B^2*d + A^2*B*d*2i \\
&))*(1/a^3)^{1/2}))/ (a*d) + (2^{1/2}*\operatorname{atanh}((2^{1/2}*A^3*a*d*(-1/a^3)^{1/2}*(\\
& a + a*\tan(c + d*x)*1i)^{1/2}*127i)/(32*((127*A^3*d)/16 - (B^3*d*1i)/16 + (\\
& 3*A*B^2*d)/16 - (A^2*B*d*125i)/16))) + (2^{1/2}*B^3*a*d*(-1/a^3)^{1/2}*(a + \\
& a*\tan(c + d*x)*1i)^{1/2}))/((32*((127*A^3*d)/16 - (B^3*d*1i)/16 + (3*A*B^2* \\
& d)/16 - (A^2*B*d*125i)/16))) + (2^{1/2}*A*B^2*a*d*(-1/a^3)^{1/2}*(a + a*\tan \\
& (c + d*x)*1i)^{1/2}*3i)/(32*((127*A^3*d)/16 - (B^3*d*1i)/16 + (3*A*B^2*d)/ \\
& 16 - (A^2*B*d*125i)/16))) + (125*2^{1/2}*A^2*B*a*d*(-1/a^3)^{1/2}*(a + a*\tan \\
& (c + d*x)*1i)^{1/2}))/((32*((127*A^3*d)/16 - (B^3*d*1i)/16 + (3*A*B^2*d)/16 \\
& - (A^2*B*d*125i)/16)))*(A*1i + B)*(-1/a^3)^{1/2}))/ (8*a*d)
\end{aligned}$$

3.110 $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

3.110.1 Optimal result 1175
 3.110.2 Mathematica [A] (verified) 1176
 3.110.3 Rubi [A] (verified) 1176
 3.110.4 Maple [A] (verified) 1183
 3.110.5 Fracas [B] (verification not implemented) 1183
 3.110.6 Sympy [F] 1184
 3.110.7 Maxima [A] (verification not implemented) 1185
 3.110.8 Giac [F] 1185
 3.110.9 Mupad [B] (verification not implemented) 1186

3.110.1 Optimal result

Integrand size = 36, antiderivative size = 259

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{(5iA-2B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{a^{5/2}d} + \frac{(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{4\sqrt{2}a^{5/2}d} + \frac{(A+iB)\cot(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} + \frac{(19A+9iB)\cot(c+dx)}{30ad(a+ia \tan(c+dx))^{3/2}} + \frac{(41A+15iB)\cot(c+dx)}{12a^2d\sqrt{a+ia \tan(c+dx)}} - \frac{7(3A+iB)\cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{4a^3d}$$

output

```
(5*I*A-2*B)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/a^(5/2)/d+1/8*(I*A+B)
)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(5/2)/d*2^(1/2)+
1/12*(41*A+15*I*B)*cot(d*x+c)/a^2/d/(a+I*a*tan(d*x+c))^(1/2)-7/4*(3*A+I*B)
)*cot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)/a^3/d+1/5*(A+I*B)*cot(d*x+c)/d/(a+I*a
*tan(d*x+c))^(5/2)+1/30*(19*A+9*I*B)*cot(d*x+c)/a/d/(a+I*a*tan(d*x+c))^(3/
2)
```

3.110.2 Mathematica [A] (verified)

Time = 6.37 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.04

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \frac{12a^{15/2}(A+iB)\cot(c+dx) - \frac{1}{2}a^7(i+\cot(c+dx))\tan^2(c+dx)}{\dots}$$

input `Integrate[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `(12*a^(15/2)*(A + I*B)*Cot[c + d*x] - (a^7*(I + Cot[c + d*x])*Tan[c + d*x]^2*(2*Sqrt[a]*(-105*(3*A + I*B) + (5*I)*(85*A + (27*I)*B)*Cot[c + d*x] + 12*(6*A + I*B)*Cot[c + d*x]^2) + 120*(5*A + (2*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]]*(1 - I*Cot[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]] + 15*Sqrt[2]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])]*(1 - I*Cot[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]))/2)/(60*a^(15/2)*d*(a + I*a*Tan[c + d*x])^(5/2))`

3.110.3 Rubi [A] (verified)

Time = 1.92 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.09, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4081, 25, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A+B\tan(c+dx)}{\tan^2(c+dx)(a+ia\tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{4079} \\ & \frac{\int \frac{\cot^2(c+dx)(2a(6A+iB)-7a(iA-B)\tan(c+dx))}{2(i\tan(c+dx)a+a)^{3/2}} dx}{5a^2} + \frac{(A+iB)\cot(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.110. $\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \frac{\int \frac{\cot^2(c+dx)(2a(6A+iB)-7a(iA-B)\tan(c+dx))}{(i\tan(c+dx)a+a)^{3/2}} dx}{10a^2} + \frac{(A+iB)\cot(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{2a(6A+iB)-7a(iA-B)\tan(c+dx)}{\tan(c+dx)^2(i\tan(c+dx)a+a)^{3/2}} dx}{10a^2} + \frac{(A+iB)\cot(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow \text{4079} \\
& \frac{\int \frac{5\cot^2(c+dx)(2a^2(11A+3iB)-a^2(19iA-9B)\tan(c+dx))}{2\sqrt{i\tan(c+dx)a+a} \cdot 3a^2} dx}{10a^2} + \frac{a(19A+9iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} + \frac{(A+iB)\cot(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow \text{27} \\
& \frac{5\int \frac{\cot^2(c+dx)(2a^2(11A+3iB)-a^2(19iA-9B)\tan(c+dx))}{\sqrt{i\tan(c+dx)a+a} \cdot 6a^2} dx}{10a^2} + \frac{a(19A+9iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} + \frac{(A+iB)\cot(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{5\int \frac{2a^2(11A+3iB)-a^2(19iA-9B)\tan(c+dx)}{\tan(c+dx)^2\sqrt{i\tan(c+dx)a+a}} dx}{10a^2} + \frac{a(19A+9iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} + \frac{(A+iB)\cot(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow \text{4079} \\
& \frac{5\left(\frac{\int \frac{3}{2}\cot^2(c+dx)\sqrt{i\tan(c+dx)a+a}(14a^3(3A+iB)-a^3(41iA-15B)\tan(c+dx))}{a^2} dx + \frac{a^2(41A+15iB)\cot(c+dx)}{d\sqrt{a+ia\tan(c+dx)}}\right)}{6a^2} + \frac{a(19A+9iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} + \\
& \quad \frac{10a^2}{5d(a+ia\tan(c+dx))^{5/2}} \frac{(A+iB)\cot(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow \text{27} \\
& \frac{5\left(\frac{3\int \cot^2(c+dx)\sqrt{i\tan(c+dx)a+a}(14a^3(3A+iB)-a^3(41iA-15B)\tan(c+dx))}{2a^2} dx + \frac{a^2(41A+15iB)\cot(c+dx)}{d\sqrt{a+ia\tan(c+dx)}}\right)}{6a^2} + \frac{a(19A+9iB)\cot(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} + \\
& \quad \frac{10a^2}{5d(a+ia\tan(c+dx))^{5/2}} \frac{(A+iB)\cot(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.110. $\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{5 \left(\frac{3 \int \frac{\sqrt{i \tan(c+dx)a+a} (14a^3(3A+iB) - a^3(41iA-15B) \tan(c+dx))}{\tan(c+dx)^2} dx + \frac{a^2(41A+15iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \right)}{6a^2} + \frac{a(19A+9iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} + \\
 & \frac{10a^2}{5d(a+ia \tan(c+dx))^{5/2}} \frac{(A+iB) \cot(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 4081 \\
 & \frac{5 \left(\frac{3 \left(\frac{\int -\cot(c+dx) \sqrt{i \tan(c+dx)a+a} (4(5iA-2B)a^4 + 7(3A+iB) \tan(c+dx)a^4) dx}{a} - \frac{14a^3(3A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} + \frac{a^2(41A+15iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \right)}{6a^2} \\
 & \frac{10a^2}{5d(a+ia \tan(c+dx))^{5/2}} \frac{(A+iB) \cot(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 25 \\
 & \frac{5 \left(\frac{3 \left(-\frac{\int \cot(c+dx) \sqrt{i \tan(c+dx)a+a} (4(5iA-2B)a^4 + 7(3A+iB) \tan(c+dx)a^4) dx}{a} - \frac{14a^3(3A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} + \frac{a^2(41A+15iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \right)}{6a^2} \\
 & \frac{10a^2}{5d(a+ia \tan(c+dx))^{5/2}} \frac{(A+iB) \cot(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 3042 \\
 & \frac{5 \left(\frac{3 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (4(5iA-2B)a^4 + 7(3A+iB) \tan(c+dx)a^4)}{\tan(c+dx)} dx - \frac{14a^3(3A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} + \frac{a^2(41A+15iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} \right)}{6a^2} + \frac{a(19A+9iB) \cot(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \\
 & \frac{10a^2}{5d(a+ia \tan(c+dx))^{5/2}} \frac{(A+iB) \cot(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 4083
 \end{aligned}$$

3.110. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$5 \left(\frac{3 \left(-\frac{a^4(A-iB) \int \sqrt{i \tan(c+dx)a+adx} + 4a^3(-2B+5iA) \int \cot(c+dx)(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx} - 14a^3(3A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{2a^2} \right)}{6a^2} + \frac{a^2}{d \sqrt{a+ia \tan(c+dx)}} \right)$$

$$\frac{(A+iB) \cot(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} \quad 10a^2$$

↓ 3042

$$5 \left(\frac{3 \left(-\frac{a^4(A-iB) \int \sqrt{i \tan(c+dx)a+adx} + 4a^3(-2B+5iA) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{\tan(c+dx)} dx - 14a^3(3A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{2a^2} \right)}{6a^2} + \frac{a^2(41A+15i)}{d \sqrt{a+ia \tan(c+dx)}} \right)$$

$$\frac{(A+iB) \cot(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} \quad 10a^2$$

↓ 3961

$$5 \left(\frac{3 \left(-\frac{4a^3(-2B+5iA) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{\tan(c+dx)} dx - \frac{2ia^5(A-iB) \int \frac{1}{a-ia \tan(c+dx)} d \sqrt{i \tan(c+dx)a+adx}}{a} - 14a^3(3A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{2a^2} \right)}{6a^2} \right)$$

$$\frac{(A+iB) \cot(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} \quad 10a^2$$

↓ 219

$$5 \left(\frac{3 \left(-\frac{4a^3(-2B+5iA) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{\tan(c+dx)} dx - \frac{i\sqrt{2}a^{9/2}(A-iB) \operatorname{arctanh} \left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}} \right)}{d} - 14a^3(3A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{2a^2} \right)}{6a^2} \right)$$

$$\frac{(A+iB) \cot(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} \quad 10a^2$$

↓ 4082

3.110. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$5 \left(\frac{3 \left(-\frac{4a^5(-2B+5iA) \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{a} - \frac{i\sqrt{2}a^{9/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{14a^3(3A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} + \frac{a^2}{2a^2} \right)$$

$$\frac{(A+iB) \cot(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

73

$$5 \left(\frac{3 \left(-\frac{8ia^4(-2B+5iA) \int \frac{1}{i-i \tan(c+dx)a+a} d \sqrt{i \tan(c+dx)a+a}}{a} - \frac{i\sqrt{2}a^{9/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{14a^3(3A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} + \frac{a^2}{2a^2} \right)$$

$$\frac{(A+iB) \cot(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

221

$$5 \left(\frac{a^2(41A+15iB) \cot(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} + \frac{3 \left(-\frac{8a^{9/2}(-2B+5iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{d} - \frac{i\sqrt{2}a^{9/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{14a^3(3A+iB) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} \right)$$

$$\frac{(A+iB) \cot(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

input `Int[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

```
output ((A + I*B)*Cot[c + d*x])/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) + ((a*(19*A +
(9*I)*B)*Cot[c + d*x])/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) + (5*((a^2*(41*A
+ (15*I)*B)*Cot[c + d*x])/(d*Sqrt[a + I*a*Tan[c + d*x]]) + (3*(-((( -8*a^(
9/2))*((5*I)*A - 2*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d - (I*S
qrt[2]*a^(9/2)*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[
a])))/d)/a - (14*a^3*(3*A + I*B)*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x]])
/d)/(2*a^2)))/(6*a^2))/(10*a^2)
```

3.110.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3961 Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d)
Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a
, b, c, d}, x] && EqQ[a^2 + b^2, 0]
```

rule 4079 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]^m)((A_.) + (B_.)\tan[(e_.) + (f_.)x])((c_.) + (d_.)\tan[(e_.) + (f_.)x])^n, x_Symbol] \rightarrow \text{Simp}[(aA + bB)(a + b\tan[e + fx])^m((c + d\tan[e + fx])^{n+1}/(2f^m(b^2c - a^2d))), x] + \text{Simp}[1/(2a^m(b^2c - a^2d)) \text{Int}[(a + b\tan[e + fx])^{m+1}(c + d\tan[e + fx])^n \text{Simp}[A(b^2cm - a^2d(2m + n + 1)) + B(a^2cm - b^2d(n + 1)) + d(Ab - aB)(m + n + 1)\tan[e + fx], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, n\}, x] \&\& \text{NeQ}[b^2c - a^2d, 0] \&\& \text{EqQ}[a^2 + b^2, 0] \&\& \text{LtQ}[m, 0] \&\& \text{!GtQ}[n, 0]$

rule 4081 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]^m)((A_.) + (B_.)\tan[(e_.) + (f_.)x])((c_.) + (d_.)\tan[(e_.) + (f_.)x])^n, x_Symbol] \rightarrow \text{Simp}[(A^2d - B^2c)(a + b\tan[e + fx])^m((c + d\tan[e + fx])^{n+1}/(f^2(n+1)(c^2 + d^2))), x] - \text{Simp}[1/(a^2(n+1)(c^2 + d^2)) \text{Int}[(a + b\tan[e + fx])^{m+1}(c + d\tan[e + fx])^{n+1} \text{Simp}[A(b^2dm - a^2c(n+1)) - B(b^2cm + a^2d(n+1)) - a(B^2c - A^2d)(m + n + 1)\tan[e + fx], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, m\}, x] \&\& \text{NeQ}[b^2c - a^2d, 0] \&\& \text{EqQ}[a^2 + b^2, 0] \&\& \text{LtQ}[n, -1]$

rule 4082 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]^m)((A_.) + (B_.)\tan[(e_.) + (f_.)x])((c_.) + (d_.)\tan[(e_.) + (f_.)x])^n, x_Symbol] \rightarrow \text{Simp}[b(B/f) \text{Subst}[\text{Int}[(a + bx)^{m-1}(c + dx)^n, x], x, \tan[e + fx]], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, m, n\}, x] \&\& \text{NeQ}[b^2c - a^2d, 0] \&\& \text{EqQ}[a^2 + b^2, 0] \&\& \text{EqQ}[A^2b + a^2B, 0]$

rule 4083 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]^m)((A_.) + (B_.)\tan[(e_.) + (f_.)x])((c_.) + (d_.)\tan[(e_.) + (f_.)x])^n, x_Symbol] \rightarrow \text{Simp}[(A^2b + a^2B)/(b^2c + a^2d) \text{Int}[(a + b\tan[e + fx])^m, x], x] - \text{Simp}[(B^2c - A^2d)/(b^2c + a^2d) \text{Int}[(a + b\tan[e + fx])^m((a - b\tan[e + fx])/(c + d\tan[e + fx])), x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, m\}, x] \&\& \text{NeQ}[b^2c - a^2d, 0] \&\& \text{EqQ}[a^2 + b^2, 0] \&\& \text{NeQ}[A^2b + a^2B, 0]$

3.110.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 195, normalized size of antiderivative = 0.75

method	result
derivativedivides	$2ia^2 \left(-\frac{7iB+17A}{8a^4\sqrt{a+ia\tan(dx+c)}} - \frac{3iB+5A}{12a^3(a+ia\tan(dx+c))^{\frac{3}{2}}} - \frac{iB+A}{10a^2(a+ia\tan(dx+c))^{\frac{5}{2}}} - \frac{(iB-A)\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{16a^{\frac{9}{2}}} \right) dx$
default	$2ia^2 \left(-\frac{7iB+17A}{8a^4\sqrt{a+ia\tan(dx+c)}} - \frac{3iB+5A}{12a^3(a+ia\tan(dx+c))^{\frac{3}{2}}} - \frac{iB+A}{10a^2(a+ia\tan(dx+c))^{\frac{5}{2}}} - \frac{(iB-A)\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{16a^{\frac{9}{2}}} \right) dx$

input `int(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `2*I/d*a^2*(-1/8/a^4*(17*A+7*I*B)/(a+I*a*tan(d*x+c))^(1/2)-1/12/a^3*(5*A+3*I*B)/(a+I*a*tan(d*x+c))^(3/2)-1/10/a^2*(A+I*B)/(a+I*a*tan(d*x+c))^(5/2)-1/16*(-A+I*B)/a^(9/2)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))+1/a^4*(1/2*I*A*(a+I*a*tan(d*x+c))^(1/2)/a/tan(d*x+c)+1/2*(5*A+2*I*B)/a^(1/2)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2)))`

3.110.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 834 vs. 2(204) = 408.

Time = 0.29 (sec) , antiderivative size = 834, normalized size of antiderivative = 3.22

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x,algorithm="fracas")`

output

```

-1/120*(15*sqrt(1/2)*(a^3*d*e^(7*I*d*x + 7*I*c) - a^3*d*e^(5*I*d*x + 5*I*c
))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2))*log(-4*(sqrt(2)*sqrt(1/2)*(a^3*d
*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-(A^2
- 2*I*A*B - B^2)/(a^5*d^2)) + (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I
*c)/(I*A + B)) - 15*sqrt(1/2)*(a^3*d*e^(7*I*d*x + 7*I*c) - a^3*d*e^(5*I*d*
x + 5*I*c))*sqrt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2))*log(4*(sqrt(2)*sqrt(1/2
)*(a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sq
rt(-(A^2 - 2*I*A*B - B^2)/(a^5*d^2)) - (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I
*d*x - I*c)/(I*A + B)) - 30*(a^3*d*e^(7*I*d*x + 7*I*c) - a^3*d*e^(5*I*d*x
+ 5*I*c))*sqrt(-(25*A^2 + 20*I*A*B - 4*B^2)/(a^5*d^2))*log(-16*(3*(5*I*A -
2*B)*a^2*e^(2*I*d*x + 2*I*c) + (5*I*A - 2*B)*a^2 + 2*sqrt(2)*(a^4*d*e^(3*
I*d*x + 3*I*c) + a^4*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*
sqrt(-(25*A^2 + 20*I*A*B - 4*B^2)/(a^5*d^2)))*e^(-2*I*d*x - 2*I*c)/(-5*I*A
+ 2*B)) + 30*(a^3*d*e^(7*I*d*x + 7*I*c) - a^3*d*e^(5*I*d*x + 5*I*c))*sqrt
(-(25*A^2 + 20*I*A*B - 4*B^2)/(a^5*d^2))*log(-16*(3*(5*I*A - 2*B)*a^2*e^(2
*I*d*x + 2*I*c) + (5*I*A - 2*B)*a^2 - 2*sqrt(2)*(a^4*d*e^(3*I*d*x + 3*I*c)
+ a^4*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-(25*A^2
+ 20*I*A*B - 4*B^2)/(a^5*d^2)))*e^(-2*I*d*x - 2*I*c)/(-5*I*A + 2*B)) - sqr
t(2)*((-403*I*A + 123*B)*e^(8*I*d*x + 8*I*c) + (-151*I*A + 21*B)*e^(6*I*d*
x + 6*I*c) - 40*(-7*I*A + 3*B)*e^(4*I*d*x + 4*I*c) + (31*I*A - 21*B)*e^...

```

3.110.6 Sympy [F]

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \int \frac{(A+B \tan(c+dx)) \cot^2(c+dx)}{(ia(\tan(c+dx)-i))^{5/2}} dx$$

input `integrate(cot(d*x+c)**2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**2/(I*a*(tan(c + d*x) - I))**(5/2), x)`

3.110.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 240, normalized size of antiderivative = 0.93

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx =$$

$$ia \left(\frac{4(105(ia \tan(dx+c)+a)^3(3A+iB)-5(ia \tan(dx+c)+a)^2(41A+15iB)a-2(ia \tan(dx+c)+a)(19A+9iB)a^2-12(A+iB)a^3)}{(ia \tan(dx+c)+a)^{7/2}a^3-(ia \tan(dx+c)+a)^{5/2}a^4} \right) + \frac{15\sqrt{2}}{240d}$$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")
```

```
output -1/240*I*a*(4*(105*(I*a*tan(d*x + c) + a)^3*(3*A + I*B) - 5*(I*a*tan(d*x + c) + a)^2*(41*A + 15*I*B)*a - 2*(I*a*tan(d*x + c) + a)*(19*A + 9*I*B)*a^2 - 12*(A + I*B)*a^3)/((I*a*tan(d*x + c) + a)^(7/2)*a^3 - (I*a*tan(d*x + c) + a)^(5/2)*a^4) + 15*sqrt(2)*(A - I*B)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a)))/a^(7/2) + 120*(5*A + 2*I*B)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a)))/a^(7/2))/d
```

3.110.8 Giac [F]

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \int \frac{(B \tan(dx+c) + A) \cot(dx+c)^2}{(ia \tan(dx+c) + a)^{5/2}} dx$$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")
```

```
output integrate((B*tan(d*x + c) + A)*cot(d*x + c)^2/(I*a*tan(d*x + c) + a)^(5/2), x)
```


3.110.9 Mupad [B] (verification not implemented)

Time = 10.09 (sec) , antiderivative size = 3002, normalized size of antiderivative = 11.59

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `int((cot(c + d*x)^2*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2),x)`

output

```

2*atanh((12*a*d^4*(a + a*tan(c + d*x)*1i)^(1/2)*((129*B^2)/(256*a^5*d^2) -
(801*A^2)/(256*a^5*d^2) - ((638401*A^4*a^2)/(16*d^4) + (16129*B^4*a^2)/(1
6*d^4) - (307555*A^2*B^2*a^2)/(8*d^4) - (A*B^3*a^2*40767i)/(4*d^4) + (A^3*
B*a^2*256479i)/(4*d^4))^(1/2)/(64*a^6) - (A*B*319i)/(128*a^5*d^2))^(1/2)*
(638401*A^4*a^2)/(16*d^4) + (16129*B^4*a^2)/(16*d^4) - (307555*A^2*B^2*a^2
)/(8*d^4) - (A*B^3*a^2*40767i)/(4*d^4) + (A^3*B*a^2*256479i)/(4*d^4))^(1/2
))/((A^3*d*31161i)/8 + (2159*B^3*d)/8 - (A*B^2*d*15867i)/8 - (38621*A^2*B*
d)/8 + (A*d^3*((638401*A^4*a^2)/(16*d^4) + (16129*B^4*a^2)/(16*d^4) - (307
555*A^2*B^2*a^2)/(8*d^4) - (A*B^3*a^2*40767i)/(4*d^4) + (A^3*B*a^2*256479i
)/(4*d^4))^(1/2)*41i)/(2*a) - (15*B*d^3*((638401*A^4*a^2)/(16*d^4) + (1612
9*B^4*a^2)/(16*d^4) - (307555*A^2*B^2*a^2)/(8*d^4) - (A*B^3*a^2*40767i)/(4
*d^4) + (A^3*B*a^2*256479i)/(4*d^4))^(1/2))/(2*a)) + (799*A^2*a^2*d^2*(a +
a*tan(c + d*x)*1i)^(1/2)*((129*B^2)/(256*a^5*d^2) - (801*A^2)/(256*a^5*d^
2) - ((638401*A^4*a^2)/(16*d^4) + (16129*B^4*a^2)/(16*d^4) - (307555*A^2*B
^2*a^2)/(8*d^4) - (A*B^3*a^2*40767i)/(4*d^4) + (A^3*B*a^2*256479i)/(4*d^4)
)^(1/2)/(64*a^6) - (A*B*319i)/(128*a^5*d^2))^(1/2))/((A^3*d*31161i)/8 + (2
159*B^3*d)/8 - (A*B^2*d*15867i)/8 - (38621*A^2*B*d)/8 + (A*d^3*((638401*A^
4*a^2)/(16*d^4) + (16129*B^4*a^2)/(16*d^4) - (307555*A^2*B^2*a^2)/(8*d^4)
- (A*B^3*a^2*40767i)/(4*d^4) + (A^3*B*a^2*256479i)/(4*d^4))^(1/2)*41i)/(2*
a) - (15*B*d^3*((638401*A^4*a^2)/(16*d^4) + (16129*B^4*a^2)/(16*d^4) - ...

```

3.111
$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$$

3.111.1 Optimal result 1187
 3.111.2 Mathematica [A] (verified) 1188
 3.111.3 Rubi [A] (verified) 1188
 3.111.4 Maple [A] (verified) 1195
 3.111.5 Fracas [B] (verification not implemented) 1196
 3.111.6 Sympy [F] 1197
 3.111.7 Maxima [A] (verification not implemented) 1197
 3.111.8 Giac [F] 1198
 3.111.9 Mupad [B] (verification not implemented) 1198

3.111.1 Optimal result

Integrand size = 36, antiderivative size = 312

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{(43A+20iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{a}}\right)}{4a^{5/2}d} - \frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{4\sqrt{2}a^{5/2}d} + \frac{(A+iB)\cot^2(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} + \frac{(23A+13iB)\cot^2(c+dx)}{30ad(a+ia \tan(c+dx))^{3/2}} + \frac{(337A+167iB)\cot^2(c+dx)}{60a^2d\sqrt{a+ia \tan(c+dx)}} + \frac{21(2iA-B)\cot(c+dx)\sqrt{a+ia \tan(c+dx)}}{4a^3d} - \frac{(85A+41iB)\cot^2(c+dx)\sqrt{a+ia \tan(c+dx)}}{12a^3d}$$

```
output 1/4*(43*A+20*I*B)*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))/a^(5/2)/d-1/8*(A-I*B)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(1/2))/a^(5/2)/d*2^(1/2)+1/60*(337*A+167*I*B)*cot(d*x+c)^2/a^2/d/(a+I*a*tan(d*x+c))^(1/2)+21/4*(2*I*A-B)*cot(d*x+c)*(a+I*a*tan(d*x+c))^(1/2)/a^3/d-1/12*(85*A+41*I*B)*cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(1/2)/a^3/d+1/5*(A+I*B)*cot(d*x+c)^2/d/(a+I*a*tan(d*x+c))^(5/2)+1/30*(23*A+13*I*B)*cot(d*x+c)^2/a/d/(a+I*a*tan(d*x+c))^(3/2)
```

3.111.2 Mathematica [A] (verified)

Time = 8.70 (sec) , antiderivative size = 298, normalized size of antiderivative = 0.96

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \frac{12a^{17/2}(A+iB)\cot^2(c+dx) - \frac{1}{2}a^8(i+\cot(c+dx))\tan^2(c+dx)}{\dots}$$

input `Integrate[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `(12*a^(17/2)*(A + I*B)*Cot[c + d*x]^2 - (a^8*(I + Cot[c + d*x])*Tan[c + d*x]^2*(-1/2*(Sqrt[a]*Csc[c + d*x]^3*(-((961*A + (461*I)*B)*Cos[c + d*x]) + (793*A + (413*I)*B)*Cos[3*(c + d*x)] + 18*((-57*I)*A + 27*B + ((83*I)*A - 43*B)*Cos[2*(c + d*x)])*Sin[c + d*x])) - 30*(43*A + (20*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]]*(I + Cot[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]] + 15*Sqrt[2]*(A - I*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])]*(I + Cot[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]))/2)/(60*a^(17/2)*d*(a + I*a*Tan[c + d*x])^(5/2))`

3.111.3 Rubi [A] (verified)

Time = 2.31 (sec) , antiderivative size = 336, normalized size of antiderivative = 1.08, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.611$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4081, 27, 3042, 4081, 3042, 4083, 3042, 3961, 219, 4082, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A+B\tan(c+dx)}{\tan(c+dx)^3(a+ia\tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{4079} \\ & \frac{\int \frac{\cot^3(c+dx)(2a(7A+2iB)-9a(iA-B)\tan(c+dx))}{2(i\tan(c+dx)a+a)^{3/2}} dx}{5a^2} + \frac{(A+iB)\cot^2(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \end{aligned}$$

3.111. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \int \frac{\cot^3(c+dx)(2a(7A+2iB)-9a(iA-B)\tan(c+dx))}{(i\tan(c+dx)a+a)^{3/2}} dx + \frac{(A+iB)\cot^2(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 27 \\
& \int \frac{2a(7A+2iB)-9a(iA-B)\tan(c+dx)}{\tan(c+dx)^3(i\tan(c+dx)a+a)^{3/2}} dx + \frac{(A+iB)\cot^2(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 3042 \\
& \int \frac{\cot^3(c+dx)(4a^2(44A+19iB)-7a^2(23iA-13B)\tan(c+dx))}{2\sqrt{i\tan(c+dx)a+a}} dx + \frac{a(23A+13iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} + \frac{(A+iB)\cot^2(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 4079 \\
& \int \frac{\cot^3(c+dx)(4a^2(44A+19iB)-7a^2(23iA-13B)\tan(c+dx))}{\sqrt{i\tan(c+dx)a+a}} dx + \frac{a(23A+13iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} + \frac{(A+iB)\cot^2(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 27 \\
& \int \frac{\cot^3(c+dx)(4a^2(44A+19iB)-7a^2(23iA-13B)\tan(c+dx))}{6a^2} dx + \frac{a(23A+13iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} + \frac{(A+iB)\cot^2(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 3042 \\
& \int \frac{4a^2(44A+19iB)-7a^2(23iA-13B)\tan(c+dx)}{\tan(c+dx)^3\sqrt{i\tan(c+dx)a+a}} dx + \frac{a(23A+13iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} + \frac{(A+iB)\cot^2(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 4079 \\
& \frac{\int \frac{5}{2}\cot^3(c+dx)\sqrt{i\tan(c+dx)a+a}(4a^3(85A+41iB)-a^3(337iA-167B)\tan(c+dx)) dx + \frac{a^2(337A+167iB)\cot^2(c+dx)}{d\sqrt{a+ia\tan(c+dx)}}}{6a^2} + \frac{a(23A+13iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} + \\
& \quad \frac{10a^2}{5d(a+ia\tan(c+dx))^{5/2}} \frac{(A+iB)\cot^2(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 27 \\
& \frac{5\int \cot^3(c+dx)\sqrt{i\tan(c+dx)a+a}(4a^3(85A+41iB)-a^3(337iA-167B)\tan(c+dx)) dx + \frac{a^2(337A+167iB)\cot^2(c+dx)}{d\sqrt{a+ia\tan(c+dx)}}}{2a^2}}{6a^2} + \frac{a(23A+13iB)\cot^2(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} + \\
& \quad \frac{10a^2}{5d(a+ia\tan(c+dx))^{5/2}} \frac{(A+iB)\cot^2(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 3042
\end{aligned}$$

3.111. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{5 \int \frac{\sqrt{i \tan(c+dx)a+a} (4a^3(85A+41iB) - a^3(337iA-167B) \tan(c+dx))}{\tan(c+dx)^3} dx + \frac{a^2(337A+167iB) \cot^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} + \frac{a(23A+13iB) \cot^2(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}}{6a^2} + \\
 & \frac{10a^2}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 4081 \\
 & \frac{5 \left(\frac{\int -6 \cot^2(c+dx) \sqrt{i \tan(c+dx)a+a} (42(2iA-B)a^4 + (85A+41iB) \tan(c+dx)a^4) dx}{2a} - \frac{2a^3(85A+41iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} + \frac{a^2(337A+167iB) \cot^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} \\
 & \frac{10a^2}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 27 \\
 & \frac{5 \left(-\frac{3 \int \cot^2(c+dx) \sqrt{i \tan(c+dx)a+a} (42(2iA-B)a^4 + (85A+41iB) \tan(c+dx)a^4) dx}{a} - \frac{2a^3(85A+41iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} + \frac{a^2(337A+167iB) \cot^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} \\
 & \frac{10a^2}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 3042 \\
 & \frac{5 \left(\frac{3 \int \frac{\sqrt{i \tan(c+dx)a+a} (42(2iA-B)a^4 + (85A+41iB) \tan(c+dx)a^4)}{\tan(c+dx)^2} dx - \frac{2a^3(85A+41iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{2a^2} + \frac{a^2(337A+167iB) \cot^2(c+dx)}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} + a \\
 & \frac{10a^2}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 4081 \\
 & \frac{5 \left(\frac{3 \left(\frac{\int \cot(c+dx) \sqrt{i \tan(c+dx)a+a} (a^5(43A+20iB) - 21a^5(2iA-B) \tan(c+dx)) dx}{a} - \frac{42a^4(-B+2iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{a} - \frac{2a^3(85A+41iB) \cot^2(c+dx)}{d} \right)}{2a^2}}{6a^2} \\
 & \frac{10a^2}{5d(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow 3042
 \end{aligned}$$

3.111. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\left(\frac{3 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^5(43A+20iB)-21a^5(2iA-B) \tan(c+dx))}{\tan(c+dx)} dx - \frac{42a^4(-B+2iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{a} \right) - \frac{2a^3(85A+41iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{2a^2}{6a^2} \quad \frac{10a^2}{10a^2}$$

$$\frac{(A+iB) \cot^2(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

↓ 4083

$$\left(\frac{3 \left(\frac{a^5(B+iA) \int \sqrt{i \tan(c+dx)a+adx+a^4(43A+20iB) \int \cot(c+dx)(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{a} - \frac{42a^4(-B+2iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{a} \right) - \frac{2a^3(85A+41iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{2a^2}{6a^2} \quad \frac{10a^2}{10a^2}$$

$$\frac{(A+iB) \cot^2(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

↓ 3042

$$\left(\frac{3 \left(\frac{a^5(B+iA) \int \sqrt{i \tan(c+dx)a+adx+a^4(43A+20iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{42a^4(-B+2iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{a} \right) - \frac{2a^3(85A+41iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{2a^2}{6a^2} \quad \frac{10a^2}{10a^2}$$

$$\frac{(A+iB) \cot^2(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

↓ 3961

$$\left(\frac{3 \left(\frac{a^4(43A+20iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{2ia^6(B+iA) \int \frac{1}{a-ia \tan(c+dx)} d \sqrt{i \tan(c+dx)a+a}}{a} - \frac{42a^4(-B+2iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d} \right)}{a} \right) - \frac{2a^3(85A+41iB) \cot^2(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}$$

$$\frac{2a^2}{6a^2} \quad \frac{10a^2}{10a^2}$$

$$\frac{(A+iB) \cot^2(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

↓ 219

3.111. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\left(\frac{3}{5} \left(\frac{a^4(43A+20iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}}{\tan(c+dx)} dx - \frac{i\sqrt{2}a^{11/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{42a^4(-B+2iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}}{a} \right) \right)$$

$$\frac{(A+iB) \cot^2(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

$$\frac{4082}{5}$$

$$\frac{3}{5} \left(\frac{a^6(43A+20iB) \int \frac{\cot(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx) - \frac{i\sqrt{2}a^{11/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{42a^4(-B+2iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}}{a} \right)$$

$$\frac{(A+iB) \cot^2(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

$$\frac{73}{5}$$

$$\frac{3}{5} \left(\frac{2ia^5(43A+20iB) \int \frac{1}{i - \frac{i \tan(c+dx)a+a}{d}} d \sqrt{i \tan(c+dx)a+a} - \frac{i\sqrt{2}a^{11/2}(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(c+dx)}}{\sqrt{2}\sqrt{a}}\right)}{d} - \frac{42a^4(-B+2iA) \cot(c+dx) \sqrt{a+ia \tan(c+dx)}}{d}}{a} \right)$$

$$\frac{(A+iB) \cot^2(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

$$\frac{221}{5}$$

3.111. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\frac{\frac{a^2(337A+167iB)\cot^2(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} + \frac{2a^3(85A+41iB)\cot^2(c+dx)\sqrt{a+ia\tan(c+dx)}}{d} - \frac{2a^{11/2}(43A+20iB)\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(c+dx)}}{\sqrt{a}}\right)}{d} - \frac{i\sqrt{2}a^{11/2}(B+iA)}{a}}{6a^2} - \frac{2a^2}{10a^2} = \frac{(A+iB)\cot^2(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}}$$

input `Int[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `((A + I*B)*Cot[c + d*x]^2)/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) + ((a*(23*A + (13*I)*B)*Cot[c + d*x]^2)/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) + ((a^2*(337*A + (167*I)*B)*Cot[c + d*x]^2)/(d*Sqrt[a + I*a*Tan[c + d*x]])) + (5*((-2*a^3*(85*A + (41*I)*B)*Cot[c + d*x]^2*Sqrt[a + I*a*Tan[c + d*x]])/d - (3*((-2*a^(11/2)*(43*A + (20*I)*B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/Sqrt[a]])/d - (I*Sqrt[2]*a^(11/2)*(I*A + B)*ArcTanh[Sqrt[a + I*a*Tan[c + d*x]]/(Sqrt[2]*Sqrt[a])])/d)/a - (42*a^4*((2*I)*A - B)*Cot[c + d*x]*Sqrt[a + I*a*Tan[c + d*x])/d)/a))/(2*a^2)/(6*a^2)/(10*a^2)`

3.111.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

- rule 221 $\text{Int}[(a_ + (b_ \cdot)(x_)^2)^{-1}, x_ \text{Symbol}] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) \cdot \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] \text{ ; FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b]$
- rule 3042 $\text{Int}[u_ , x_ \text{Symbol}] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$
- rule 3961 $\text{Int}[\text{Sqrt}[a_ + (b_ \cdot) \cdot \tan[(c_ \cdot) + (d_ \cdot)(x_)]], x_ \text{Symbol}] \rightarrow \text{Simp}[-2 \cdot (b/d) \text{ Subst}[\text{Int}[1/(2 \cdot a - x^2), x], x, \text{Sqrt}[a + b \cdot \tan[c + d \cdot x]], x] \text{ ; FreeQ}\{a, b, c, d, x\} \ \&\& \ \text{EqQ}[a^2 + b^2, 0]$
- rule 4079 $\text{Int}[(a_ + (b_ \cdot) \cdot \tan[(e_ \cdot) + (f_ \cdot)(x_)])^{m_} \cdot ((A_ \cdot) + (B_ \cdot) \cdot \tan[(e_ \cdot) + (f_ \cdot)(x_)]) \cdot ((c_ \cdot) + (d_ \cdot) \cdot \tan[(e_ \cdot) + (f_ \cdot)(x_)])^{n_}, x_ \text{Symbol}] \rightarrow \text{Simp}[(a \cdot A + b \cdot B) \cdot (a + b \cdot \tan[e + f \cdot x])^m \cdot ((c + d \cdot \tan[e + f \cdot x])^{n+1} / (2 \cdot f \cdot m \cdot (b \cdot c - a \cdot d))), x] + \text{Simp}[1 / (2 \cdot a \cdot m \cdot (b \cdot c - a \cdot d)) \text{ Int}[(a + b \cdot \tan[e + f \cdot x])^{m+1} \cdot (c + d \cdot \tan[e + f \cdot x])^n \cdot \text{Simp}[A \cdot (b \cdot c \cdot m - a \cdot d \cdot (2 \cdot m + n + 1)) + B \cdot (a \cdot c \cdot m - b \cdot d \cdot (n + 1)) + d \cdot (A \cdot b - a \cdot B) \cdot (m + n + 1) \cdot \tan[e + f \cdot x], x], x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, A, B, n, x\} \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{LtQ}[m, 0] \ \&\& \ \text{!GtQ}[n, 0]$
- rule 4081 $\text{Int}[(a_ + (b_ \cdot) \cdot \tan[(e_ \cdot) + (f_ \cdot)(x_)])^{m_} \cdot ((A_ \cdot) + (B_ \cdot) \cdot \tan[(e_ \cdot) + (f_ \cdot)(x_)]) \cdot ((c_ \cdot) + (d_ \cdot) \cdot \tan[(e_ \cdot) + (f_ \cdot)(x_)])^{n_}, x_ \text{Symbol}] \rightarrow \text{Simp}[(A \cdot d - B \cdot c) \cdot (a + b \cdot \tan[e + f \cdot x])^m \cdot ((c + d \cdot \tan[e + f \cdot x])^{n+1} / (f \cdot (n + 1) \cdot (c^2 + d^2))), x] - \text{Simp}[1 / (a \cdot (n + 1) \cdot (c^2 + d^2)) \text{ Int}[(a + b \cdot \tan[e + f \cdot x])^m \cdot (c + d \cdot \tan[e + f \cdot x])^{n+1} \cdot \text{Simp}[A \cdot (b \cdot d \cdot m - a \cdot c \cdot (n + 1)) - B \cdot (b \cdot c \cdot m + a \cdot d \cdot (n + 1)) - a \cdot (B \cdot c - A \cdot d) \cdot (m + n + 1) \cdot \tan[e + f \cdot x], x], x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, A, B, m, x\} \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{LtQ}[n, -1]$
- rule 4082 $\text{Int}[(a_ + (b_ \cdot) \cdot \tan[(e_ \cdot) + (f_ \cdot)(x_)])^{m_} \cdot ((A_ \cdot) + (B_ \cdot) \cdot \tan[(e_ \cdot) + (f_ \cdot)(x_)]) \cdot ((c_ \cdot) + (d_ \cdot) \cdot \tan[(e_ \cdot) + (f_ \cdot)(x_)])^{n_}, x_ \text{Symbol}] \rightarrow \text{Simp}[b \cdot (B/f) \text{ Subst}[\text{Int}[(a + b \cdot x)^{m-1} \cdot (c + d \cdot x)^n, x], x, \tan[e + f \cdot x]], x] \text{ ; FreeQ}\{a, b, c, d, e, f, A, B, m, n, x\} \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[A \cdot b + a \cdot B, 0]$

```
rule 4083 Int[(((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
  (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(
  A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A
  *d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*T
  an[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c -
  a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

3.111.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 225, normalized size of antiderivative = 0.72

method	result
derivativedivides	$2a^3 \left(-\frac{17iB+31A}{8a^5\sqrt{a+ia\tan(dx+c)}} - \frac{5iB+7A}{12a^4(a+ia\tan(dx+c))^{\frac{3}{2}}} - \frac{iB+A}{10a^3(a+ia\tan(dx+c))^{\frac{5}{2}}} - \frac{(-iB+A)\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{16a^{\frac{11}{2}}} \right) d$
default	$2a^3 \left(-\frac{17iB+31A}{8a^5\sqrt{a+ia\tan(dx+c)}} - \frac{5iB+7A}{12a^4(a+ia\tan(dx+c))^{\frac{3}{2}}} - \frac{iB+A}{10a^3(a+ia\tan(dx+c))^{\frac{5}{2}}} - \frac{(-iB+A)\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{a+ia\tan(dx+c)}\sqrt{2}}{2\sqrt{a}}\right)}{16a^{\frac{11}{2}}} \right) d$

```
input int(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x,method=_RETUR
NVERBOSE)
```

```
output 2/d*a^3*(-1/8/a^5*(17*I*B+31*A)/(a+I*a*tan(d*x+c))^(1/2)-1/12/a^4*(7*A+5*I
*B)/(a+I*a*tan(d*x+c))^(3/2)-1/10/a^3*(A+I*B)/(a+I*a*tan(d*x+c))^(5/2)-1/1
6*(A-I*B)/a^(11/2)*2^(1/2)*arctanh(1/2*(a+I*a*tan(d*x+c))^(1/2)*2^(1/2)/a^(
1/2))+1/a^5*(-((-1/2*I*B-11/8*A)*(a+I*a*tan(d*x+c))^(3/2)+(1/2*I*a*B+13/8
*a*A)*(a+I*a*tan(d*x+c))^(1/2))/a^2/tan(d*x+c)^2+1/8*(43*A+20*I*B)/a^(1/2)
*arctanh((a+I*a*tan(d*x+c))^(1/2)/a^(1/2))))
```

3.111.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 923 vs. $2(243) = 486$.

Time = 0.29 (sec) , antiderivative size = 923, normalized size of antiderivative = 2.96

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorith
m="fracas")
```

```
output 1/240*(30*sqrt(1/2)*(a^3*d*e^(9*I*d*x + 9*I*c) - 2*a^3*d*e^(7*I*d*x + 7*I*
c) + a^3*d*e^(5*I*d*x + 5*I*c))*sqrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2))*log(
-4*(sqrt(2)*sqrt(1/2)*(I*a^3*d*e^(2*I*d*x + 2*I*c) + I*a^3*d)*sqrt(a/(e^(2
*I*d*x + 2*I*c) + 1))*sqrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2)) + (-I*A - B)*a
*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 30*sqrt(1/2)*(a^3*d*e^(9*I
*d*x + 9*I*c) - 2*a^3*d*e^(7*I*d*x + 7*I*c) + a^3*d*e^(5*I*d*x + 5*I*c))*s
qrt((A^2 - 2*I*A*B - B^2)/(a^5*d^2))*log(-4*(sqrt(2)*sqrt(1/2)*(-I*a^3*d*e
^(2*I*d*x + 2*I*c) - I*a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((A^2
- 2*I*A*B - B^2)/(a^5*d^2)) + (-I*A - B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*
c)/(I*A + B)) + 15*(a^3*d*e^(9*I*d*x + 9*I*c) - 2*a^3*d*e^(7*I*d*x + 7*I*c
) + a^3*d*e^(5*I*d*x + 5*I*c))*sqrt((1849*A^2 + 1720*I*A*B - 400*B^2)/(a^5
*d^2))*log(-16*(3*(43*I*A - 20*B)*a^2*e^(2*I*d*x + 2*I*c) + (43*I*A - 20*B
)*a^2 + 2*sqrt(2)*(I*a^4*d*e^(3*I*d*x + 3*I*c) + I*a^4*d*e^(I*d*x + I*c))*
sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((1849*A^2 + 1720*I*A*B - 400*B^2)/(
a^5*d^2)))*e^(-2*I*d*x - 2*I*c)/(-43*I*A + 20*B)) - 15*(a^3*d*e^(9*I*d*x +
9*I*c) - 2*a^3*d*e^(7*I*d*x + 7*I*c) + a^3*d*e^(5*I*d*x + 5*I*c))*sqrt((1
849*A^2 + 1720*I*A*B - 400*B^2)/(a^5*d^2))*log(-16*(3*(43*I*A - 20*B)*a^2*
e^(2*I*d*x + 2*I*c) + (43*I*A - 20*B)*a^2 + 2*sqrt(2)*(-I*a^4*d*e^(3*I*d*x
+ 3*I*c) - I*a^4*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqr
t((1849*A^2 + 1720*I*A*B - 400*B^2)/(a^5*d^2)))*e^(-2*I*d*x - 2*I*c)/(-...
```

3.111.6 Sympy [F]

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\cot^3(c+dx)}{(ia(\tan(c+dx)-i))^{\frac{5}{2}}} dx$$

input `integrate(cot(d*x+c)**3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**3/(I*a*(tan(c + d*x) - I))**(5/2), x)`

3.111.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 283, normalized size of antiderivative = 0.91

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = a^2 \left(\frac{4 \left(315 (ia \tan(dx+c)+a)^4 (2A+iB) - 5 (ia \tan(dx+c)+a)^3 (211A+104iB)a + (ia \tan(dx+c)+a)^2 (337A+167iB)a^2 + 2 (ia \tan(dx+c)+a) \right)}{(ia \tan(dx+c)+a)^{\frac{9}{2}} a^4 - 2 (ia \tan(dx+c)+a)^{\frac{7}{2}} a^5 + (ia \tan(dx+c)+a)^{\frac{5}{2}} a^6} \right)$$

240 d

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `-1/240*a^2*(4*(315*(I*a*tan(d*x + c) + a)^4*(2*A + I*B) - 5*(I*a*tan(d*x + c) + a)^3*(211*A + 104*I*B)*a + (I*a*tan(d*x + c) + a)^2*(337*A + 167*I*B)*a^2 + 2*(I*a*tan(d*x + c) + a)*(23*A + 13*I*B)*a^3 + 12*(A + I*B)*a^4)/((I*a*tan(d*x + c) + a)^(9/2)*a^4 - 2*(I*a*tan(d*x + c) + a)^(7/2)*a^5 + (I*a*tan(d*x + c) + a)^(5/2)*a^6) - 15*sqrt(2)*(A - I*B)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(d*x + c) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(d*x + c) + a)))/a^(9/2) + 30*(43*A + 20*I*B)*log((sqrt(I*a*tan(d*x + c) + a) - sqrt(a))/(sqrt(I*a*tan(d*x + c) + a) + sqrt(a)))/a^(9/2))/d`

3.111.8 Giac [F]

$$\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{(B \tan(dx + c) + A) \cot(dx + c)^3}{(ia \tan(dx + c) + a)^{5/2}} dx$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^3/(I*a*tan(d*x + c) + a)^(5/2), x)`

3.111.9 Mupad [B] (verification not implemented)

Time = 10.01 (sec) , antiderivative size = 3048, normalized size of antiderivative = 9.77

$$\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `int((cot(c + d*x)^3*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2),x)`

output

```

2*atanh((192*a*d^4*(a + a*tan(c + d*x)*1i)^(1/2)*((3699*A^2)/(256*a^5*d^2)
- ((13667809*A^4*a^2)/(16*d^4) + (638401*B^4*a^2)/(16*d^4) - (8877585*A^2
*B^2*a^2)/(8*d^4) - (A*B^3*a^2*1375079i)/(4*d^4) + (A^3*B*a^2*6362537i)/(4
*d^4))^(1/2)/(64*a^6) - (801*B^2)/(256*a^5*d^2) + (A*B*1719i)/(128*a^5*d^2
))^1/2)*((13667809*A^4*a^2)/(16*d^4) + (638401*B^4*a^2)/(16*d^4) - (88775
85*A^2*B^2*a^2)/(8*d^4) - (A*B^3*a^2*1375079i)/(4*d^4) + (A^3*B*a^2*636253
7i)/(4*d^4))^(1/2))/(B^3*d*62322i - 643278*A^3*d + 407502*A*B^2*d - A^2*B*
d*887274i + (680*A*d^3*((13667809*A^4*a^2)/(16*d^4) + (638401*B^4*a^2)/(16
*d^4) - (8877585*A^2*B^2*a^2)/(8*d^4) - (A*B^3*a^2*1375079i)/(4*d^4) + (A^
3*B*a^2*6362537i)/(4*d^4))^(1/2))/a + (B*d^3*((13667809*A^4*a^2)/(16*d^4)
+ (638401*B^4*a^2)/(16*d^4) - (8877585*A^2*B^2*a^2)/(8*d^4) - (A*B^3*a^2*1
375079i)/(4*d^4) + (A^3*B*a^2*6362537i)/(4*d^4))^(1/2)*328i)/a) - (59152*A
^2*a^2*d^2*(a + a*tan(c + d*x)*1i)^(1/2)*((3699*A^2)/(256*a^5*d^2) - ((136
67809*A^4*a^2)/(16*d^4) + (638401*B^4*a^2)/(16*d^4) - (8877585*A^2*B^2*a^2
)/(8*d^4) - (A*B^3*a^2*1375079i)/(4*d^4) + (A^3*B*a^2*6362537i)/(4*d^4))^(
1/2)/(64*a^6) - (801*B^2)/(256*a^5*d^2) + (A*B*1719i)/(128*a^5*d^2))^1/2
)/(B^3*d*62322i - 643278*A^3*d + 407502*A*B^2*d - A^2*B*d*887274i + (680*A
*d^3*((13667809*A^4*a^2)/(16*d^4) + (638401*B^4*a^2)/(16*d^4) - (8877585*A
^2*B^2*a^2)/(8*d^4) - (A*B^3*a^2*1375079i)/(4*d^4) + (A^3*B*a^2*6362537i)/
(4*d^4))^(1/2))/a + (B*d^3*((13667809*A^4*a^2)/(16*d^4) + (638401*B^4*a...

```

3.112 $\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.112.1 Optimal result	1200
3.112.2 Mathematica [A] (verified)	1200
3.112.3 Rubi [A] (verified)	1201
3.112.4 Maple [B] (verified)	1203
3.112.5 Fricas [B] (verification not implemented)	1204
3.112.6 Sympy [F]	1205
3.112.7 Maxima [B] (verification not implemented)	1205
3.112.8 Giac [A] (verification not implemented)	1206
3.112.9 Mupad [B] (verification not implemented)	1207

3.112.1 Optimal result

Integrand size = 34, antiderivative size = 130

$$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= -\frac{2\sqrt[4]{-1}a(iA+B) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{d} - \frac{2a(iA+B)\sqrt{\tan(c+dx)}}{d}$$

$$+ \frac{2a(A-iB)\tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2a(iA+B)\tan^{\frac{5}{2}}(c+dx)}{5d} + \frac{2iaB\tan^{\frac{7}{2}}(c+dx)}{7d}$$

```
output -2*(-1)^(1/4)*a*(I*A+B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d-2*a*(I*A+B)*
tan(d*x+c)^(1/2)/d+2/3*a*(A-I*B)*tan(d*x+c)^(3/2)/d+2/5*a*(I*A+B)*tan(d*x+
c)^(5/2)/d+2/7*I*a*B*tan(d*x+c)^(7/2)/d
```

3.112.2 Mathematica [A] (verified)

Time = 1.46 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.83

$$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{2a\left(-105\sqrt[4]{-1}(iA+B) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right) + \sqrt{\tan(c+dx)}(-105i(A-iB) + 35(A-iB)t\right)}{105d}$$

input `Integrate[Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(2*a*(-105*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + Sqrt[Tan[c + d*x]]*((-105*I)*(A - I*B) + 35*(A - I*B)*Tan[c + d*x] + 21*(I*A + B)*Tan[c + d*x]^2 + (15*I)*B*Tan[c + d*x]^3))/(105*d)`

3.112.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.324$, Rules used = {3042, 4075, 3042, 4011, 3042, 4011, 3042, 4011, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c+dx)^{5/2}(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4075} \\
 & \int \tan^{\frac{5}{2}}(c+dx)(a(A-iB)+a(iA+B) \tan(c+dx)) dx + \frac{2iaB \tan^{\frac{7}{2}}(c+dx)}{7d} \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c+dx)^{5/2}(a(A-iB)+a(iA+B) \tan(c+dx)) dx + \frac{2iaB \tan^{\frac{7}{2}}(c+dx)}{7d} \\
 & \quad \downarrow \text{4011} \\
 & \int \tan^{\frac{3}{2}}(c+dx)(a(A-iB) \tan(c+dx)-a(iA+B)) dx + \frac{2a(B+iA) \tan^{\frac{5}{2}}(c+dx)}{5d} + \frac{2iaB \tan^{\frac{7}{2}}(c+dx)}{7d} \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c+dx)^{3/2}(a(A-iB) \tan(c+dx)-a(iA+B)) dx + \frac{2a(B+iA) \tan^{\frac{5}{2}}(c+dx)}{5d} + \\
 & \quad \frac{2iaB \tan^{\frac{7}{2}}(c+dx)}{7d} \\
 & \quad \downarrow \text{4011}
 \end{aligned}$$

3.112. $\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \int \sqrt{\tan(c+dx)}(-a(A-iB) - a(iA+B)\tan(c+dx))dx + \frac{2a(B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d} + \\
& \quad \frac{2a(A-iB)\tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2iaB\tan^{\frac{7}{2}}(c+dx)}{7d} \\
& \quad \downarrow \text{3042} \\
& \int \sqrt{\tan(c+dx)}(-a(A-iB) - a(iA+B)\tan(c+dx))dx + \frac{2a(B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d} + \\
& \quad \frac{2a(A-iB)\tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2iaB\tan^{\frac{7}{2}}(c+dx)}{7d} \\
& \quad \downarrow \text{4011} \\
& \int \frac{a(iA+B) - a(A-iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}}dx + \frac{2a(B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d} + \\
& \quad \frac{2a(A-iB)\tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2a(B+iA)\sqrt{\tan(c+dx)}}{d} + \frac{2iaB\tan^{\frac{7}{2}}(c+dx)}{7d} \\
& \quad \downarrow \text{3042} \\
& \int \frac{a(iA+B) - a(A-iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}}dx + \frac{2a(B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d} + \\
& \quad \frac{2a(A-iB)\tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2a(B+iA)\sqrt{\tan(c+dx)}}{d} + \frac{2iaB\tan^{\frac{7}{2}}(c+dx)}{7d} \\
& \quad \downarrow \text{4016} \\
& \frac{2a^2(B+iA)^2 \int \frac{1}{a(iA+B)+a(A-iB)\tan(c+dx)}d\sqrt{\tan(c+dx)}}{d} + \frac{2a(B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d} + \\
& \quad \frac{2a(A-iB)\tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2a(B+iA)\sqrt{\tan(c+dx)}}{d} + \frac{2iaB\tan^{\frac{7}{2}}(c+dx)}{7d} \\
& \quad \downarrow \text{218} \\
& -\frac{2\sqrt[4]{-1}a(B+iA)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{d} + \frac{2a(B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d} + \\
& \quad \frac{2a(A-iB)\tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2a(B+iA)\sqrt{\tan(c+dx)}}{d} + \frac{2iaB\tan^{\frac{7}{2}}(c+dx)}{7d}
\end{aligned}$$

input `Int[Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(-2*(-1)^(1/4)*a*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]])/d - (2*a*(I*A + B)*Sqrt[Tan[c + d*x]])/d + (2*a*(A - I*B)*Tan[c + d*x]^(3/2))/(3*d) + (2*a*(I*A + B)*Tan[c + d*x]^(5/2))/(5*d) + (((2*I)/7)*a*B*Tan[c + d*x]^(7/2))/d`

3.112. $\int \tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))(A+B\tan(c+dx))dx$

3.112.3.1 Defintions of rubi rules used

- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`
- rule 4016 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`
- rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.112.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 271 vs. $2(105) = 210$.

Time = 0.09 (sec) , antiderivative size = 272, normalized size of antiderivative = 2.09

method	result
derivativedivides	$a \left(\frac{2iB \left(\tan^{\frac{7}{2}}(dx+c) \right)}{7} + \frac{2iA \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} + \frac{2B \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} - \frac{2iB \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + \frac{2A \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 2iA \sqrt{\tan(dx+c)} \right)$
default	$a \left(\frac{2iB \left(\tan^{\frac{7}{2}}(dx+c) \right)}{7} + \frac{2iA \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} + \frac{2B \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} - \frac{2iB \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + \frac{2A \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 2iA \sqrt{\tan(dx+c)} \right)$
parts	$\frac{(iaA+Ba) \left(\frac{2 \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} - 2 \sqrt{\tan(dx+c)} + \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)}{1-\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)} \right) + 2 \arctan \left(\frac{1+\sqrt{2} \left(\sqrt{\tan(dx+c)} \right)}{4} \right)}{d} \right)}{d}$

```
input int(tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 1/d*a*(2/7*I*B*tan(d*x+c)^(7/2)+2/5*I*A*tan(d*x+c)^(5/2)+2/5*B*tan(d*x+c)^(
5/2)-2/3*I*B*tan(d*x+c)^(3/2)+2/3*A*tan(d*x+c)^(3/2)-2*I*A*tan(d*x+c)^(1/
2)-2*B*tan(d*x+c)^(1/2)+1/4*(I*A+B)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2
))+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*
tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(-A+I*B)*2^(1
/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2
))+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*ta
n(d*x+c)^(1/2))))
```

3.112.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 480 vs. 2(100) = 200.

Time = 0.33 (sec) , antiderivative size = 480, normalized size of antiderivative = 3.69

$$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{105 \left(de^{(6i dx+6i c)} + 3 de^{(4i dx+4i c)} + 3 de^{(2i dx+2i c)} + d \right) \sqrt{-\frac{(-i A^2-2 AB+i B^2)a^2}{d^2}} \log \left(\frac{2 \left((A-i B) a e^{(2i dx+2i c)} + (de^{(2i dx+2i c)} + \dots \right)}{\dots} \right)}{\dots}$$

```
input integrate(tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorith
m="fricas")
```

3.112. $\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

output $\frac{1}{210}(105(d e^{6 I d x+6 I c}+3 d e^{4 I d x+4 I c}+3 d e^{2 I d x+2 I c}+d) \sqrt{-\left(-I A^2-2 A B+I B^2\right) a^2 / d^2} \log \left(2\left((A-I B) a e^{2 I d x+2 I c}+\left(d e^{2 I d x+2 I c}+d\right) \sqrt{-\left(-I A^2-2 A B+I B^2\right) a^2 / d^2}\right) \sqrt{\left(-I e^{2 I d x+2 I c}+I\right) / \left(e^{2 I d x+2 I c}+1\right)} e^{-2 I d x-2 I c} / \left((I A+B) a\right)\right)-105\left(d e^{6 I d x+6 I c}+3 d e^{4 I d x+4 I c}+3 d e^{2 I d x+2 I c}+d\right) \sqrt{-\left(-I A^2-2 A B+I B^2\right) a^2 / d^2} \log \left(2\left((A-I B) a e^{2 I d x+2 I c}-\left(d e^{2 I d x+2 I c}+d\right) \sqrt{-\left(-I A^2-2 A B+I B^2\right) a^2 / d^2}\right) \sqrt{\left(-I e^{2 I d x+2 I c}+I\right) / \left(e^{2 I d x+2 I c}+1\right)} e^{-2 I d x-2 I c} / \left((I A+B) a\right)\right)-4\left(\left(161 I A+176 B\right) a e^{6 I d x+6 I c}+\left(329 I A+284 B\right) a e^{4 I d x+4 I c}+\left(259 I A+304 B\right) a e^{2 I d x+2 I c}+\left(91 I A+76 B\right) a\right) \sqrt{\left(-I e^{2 I d x+2 I c}+I\right) / \left(e^{2 I d x+2 I c}+1\right)} / \left(d e^{6 I d x+6 I c}+3 d e^{4 I d x+4 I c}+3 d e^{2 I d x+2 I c}+d\right)$

3.112.6 Sympy [F]

$$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= ia \left(\int A \tan^{\frac{7}{2}}(c+dx) dx + \int B \tan^{\frac{9}{2}}(c+dx) dx + \int \left(-iA \tan^{\frac{5}{2}}(c+dx)\right) dx + \int \left(-iB \tan^{\frac{7}{2}}(c+dx)\right) dx \right)$$

input `integrate(tan(d*x+c)**(5/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `I*a*(Integral(A*tan(c+d*x)**(7/2),x)+Integral(B*tan(c+d*x)**(9/2),x)+Integral(-I*A*tan(c+d*x)**(5/2),x)+Integral(-I*B*tan(c+d*x)**(7/2),x))`

3.112.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 202 vs. $2(100) = 200$.

Time = 0.32 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.55

$$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx =$$

$$-120iBa \tan(dx+c)^{\frac{7}{2}} + 168(-iA-B)a \tan(dx+c)^{\frac{5}{2}} - 280(A-iB)a \tan(dx+c)^{\frac{3}{2}} + 840(iA +$$

3.112. $\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

input `integrate(tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/420*(-120*I*B*a*tan(d*x + c)^(7/2) + 168*(-I*A - B)*a*tan(d*x + c)^(5/2) - 280*(A - I*B)*a*tan(d*x + c)^(3/2) + 840*(I*A + B)*a*sqrt(tan(d*x + c)) - 105*(2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a)/d`

3.112.8 Giac [A] (verification not implemented)

Time = 0.64 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.09

$$\int \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{(i - 1) \sqrt{2}(Aa - iBa) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{d} - \frac{2\left(-15iBad^6 \tan(dx + c)^{\frac{7}{2}} - 21iAad^6 \tan(dx + c)^{\frac{5}{2}} - 21Bad^6 \tan(dx + c)^{\frac{5}{2}} - 35Aad^6 \tan(dx + c)^{\frac{3}{2}}\right)}{105d^7}$$

input `integrate(tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `(I - 1)*sqrt(2)*(A*a - I*B*a)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d - 2/105*(-15*I*B*a*d^6*tan(d*x + c)^(7/2) - 21*I*A*a*d^6*tan(d*x + c)^(5/2) - 21*B*a*d^6*tan(d*x + c)^(5/2) - 35*A*a*d^6*tan(d*x + c)^(3/2) + 35*I*B*a*d^6*tan(d*x + c)^(3/2) + 105*I*A*a*d^6*sqrt(tan(d*x + c)) + 105*B*a*d^6*sqrt(tan(d*x + c)))/d^7`

3.112.9 Mupad [B] (verification not implemented)

Time = 12.58 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.24

$$\begin{aligned}
& \int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx \\
&= \frac{2 A a \tan(c+dx)^{3/2}}{3 d} - \frac{A a \sqrt{\tan(c+dx)} 2i}{d} + \frac{A a \tan(c+dx)^{5/2} 2i}{5 d} \\
&\quad - \frac{2 B a \sqrt{\tan(c+dx)}}{d} - \frac{B a \tan(c+dx)^{3/2} 2i}{3 d} + \frac{2 B a \tan(c+dx)^{5/2}}{5 d} \\
&\quad + \frac{B a \tan(c+dx)^{7/2} 2i}{7 d} - \frac{(-1)^{1/4} A a \operatorname{atan}\left((-1)^{1/4} \sqrt{\tan(c+dx)} 1i\right) 2i}{d} \\
&\quad + \frac{\sqrt{2} B a \operatorname{atan}\left(\sqrt{2} \sqrt{\tan(c+dx)} \left(\frac{1}{2} - \frac{1}{2}i\right)\right) (1+1i)}{d}
\end{aligned}$$

input `int(tan(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`

output `(2*A*a*tan(c + d*x)^(3/2))/(3*d) - (A*a*tan(c + d*x)^(1/2)*2i)/d + (A*a*tan(c + d*x)^(5/2)*2i)/(5*d) - (2*B*a*tan(c + d*x)^(1/2))/d - (B*a*tan(c + d*x)^(3/2)*2i)/(3*d) + (2*B*a*tan(c + d*x)^(5/2))/(5*d) + (B*a*tan(c + d*x)^(7/2)*2i)/(7*d) - ((-1)^(1/4)*A*a*atan((-1)^(1/4)*tan(c + d*x)^(1/2)*1i)*2i)/d + (2^(1/2)*B*a*atan(2^(1/2)*tan(c + d*x)^(1/2)*(1/2 - 1i/2))*(1 + 1i))/d`

3.113 $\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.113.1 Optimal result	1208
3.113.2 Mathematica [A] (verified)	1208
3.113.3 Rubi [A] (verified)	1209
3.113.4 Maple [B] (verified)	1211
3.113.5 Fricas [B] (verification not implemented)	1212
3.113.6 Sympy [F]	1213
3.113.7 Maxima [B] (verification not implemented)	1213
3.113.8 Giac [A] (verification not implemented)	1214
3.113.9 Mupad [B] (verification not implemented)	1214

3.113.1 Optimal result

Integrand size = 34, antiderivative size = 105

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{2\sqrt[4]{-1}a(A-iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c+dx)}\right)}{d} + \frac{2a(A-iB) \sqrt{\tan(c+dx)}}{d}$$

$$+ \frac{2a(iA+B) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2iaB \tan^{\frac{5}{2}}(c+dx)}{5d}$$

```
output 2*(-1)^(1/4)*a*(A-I*B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d+2*a*(A-I*B)*tan(d*x+c)^(1/2)/d+2/3*a*(I*A+B)*tan(d*x+c)^(3/2)/d+2/5*I*a*B*tan(d*x+c)^(5/2)/d
```

3.113.2 Mathematica [A] (verified)

Time = 0.93 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.85

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{2a\left(15\sqrt[4]{-1}(A-iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c+dx)}\right) + \sqrt{\tan(c+dx)}(15(A-iB) + 5(iA+B) \tan(c+dx))\right)}{15d}$$

input `Integrate[Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x
]`

output `(2*a*(15*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + Sqrt
[Tan[c + d*x]]*(15*(A - I*B) + 5*(I*A + B)*Tan[c + d*x] + (3*I)*B*Tan[c +
d*x]^2)))/(15*d)`

3.113.3 Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4075, 3042, 4011, 3042, 4011, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(c + dx)^{3/2}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4075} \\ & \int \tan^{\frac{3}{2}}(c + dx)(a(A - iB) + a(iA + B) \tan(c + dx)) dx + \frac{2iaB \tan^{\frac{5}{2}}(c + dx)}{5d} \\ & \quad \downarrow \text{3042} \\ & \int \tan(c + dx)^{3/2}(a(A - iB) + a(iA + B) \tan(c + dx)) dx + \frac{2iaB \tan^{\frac{5}{2}}(c + dx)}{5d} \\ & \quad \downarrow \text{4011} \\ & \int \sqrt{\tan(c + dx)}(a(A - iB) \tan(c + dx) - a(iA + B)) dx + \frac{2a(B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d} + \\ & \quad \frac{2iaB \tan^{\frac{5}{2}}(c + dx)}{5d} \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{\tan(c + dx)}(a(A - iB) \tan(c + dx) - a(iA + B)) dx + \frac{2a(B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d} + \\ & \quad \frac{2iaB \tan^{\frac{5}{2}}(c + dx)}{5d} \end{aligned}$$

3.113. $\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \downarrow 4011 \\
& \int \frac{-a(A - iB) - a(iA + B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2a(B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d} + \\
& \quad \frac{2a(A - iB) \sqrt{\tan(c + dx)}}{d} + \frac{2iaB \tan^{\frac{5}{2}}(c + dx)}{5d} \\
& \downarrow 3042 \\
& \int \frac{-a(A - iB) - a(iA + B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2a(B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d} + \\
& \quad \frac{2a(A - iB) \sqrt{\tan(c + dx)}}{d} + \frac{2iaB \tan^{\frac{5}{2}}(c + dx)}{5d} \\
& \downarrow 4016 \\
& \frac{2a^2(A - iB)^2 \int \frac{1}{a(iA+B) \tan(c+dx) - a(A-iB)} d \sqrt{\tan(c + dx)}}{d} + \frac{2a(B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d} + \\
& \quad \frac{2a(A - iB) \sqrt{\tan(c + dx)}}{d} + \frac{2iaB \tan^{\frac{5}{2}}(c + dx)}{5d} \\
& \downarrow 218 \\
& \frac{2\sqrt[4]{-1}a(A - iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right)}{d} + \frac{2a(B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d} + \\
& \quad \frac{2a(A - iB) \sqrt{\tan(c + dx)}}{d} + \frac{2iaB \tan^{\frac{5}{2}}(c + dx)}{5d}
\end{aligned}$$

input `Int[Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(2*(-1)^(1/4)*a*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]])/d + (2*a*(A - I*B)*Sqrt[Tan[c + d*x]])/d + (2*a*(I*A + B)*Tan[c + d*x]^(3/2))/(3*d) + (((2*I)/5)*a*B*Tan[c + d*x]^(5/2))/d`

3.113.3.1 Defintions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4011 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

```
rule 4016 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_
)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b
*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]
```

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f
*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

3.113.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 250 vs. 2(85) = 170.

Time = 0.04 (sec) , antiderivative size = 251, normalized size of antiderivative = 2.39

method	result
derivativedivides	$a \left(\frac{2iB \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} + \frac{2iA \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + \frac{2B \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 2iB(\sqrt{\tan}(dx+c)) + 2A(\sqrt{\tan}(dx+c)) + \frac{(iB-A)\sqrt{2} \left(\ln\left(\frac{1}{1}\right) \right)}{1} \right)$
default	$a \left(\frac{2iB \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} + \frac{2iA \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + \frac{2B \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 2iB(\sqrt{\tan}(dx+c)) + 2A(\sqrt{\tan}(dx+c)) + \frac{(iB-A)\sqrt{2} \left(\ln\left(\frac{1}{1}\right) \right)}{1} \right)$
parts	$\frac{(iaA+Ba) \left(\frac{2 \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - \frac{\sqrt{2} \left(\ln\left(\frac{1-\sqrt{2}(\sqrt{\tan}(dx+c))+\tan(dx+c)}{1+\sqrt{2}(\sqrt{\tan}(dx+c))+\tan(dx+c)}\right) + 2 \arctan\left(\frac{1+\sqrt{2}(\sqrt{\tan}(dx+c))}{1}\right) + 2 \arctan\left(\frac{-1+\sqrt{2}(\sqrt{\tan}(dx+c))}{1}\right)}{4} \right)}{d}$

```
input int(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

$$3.113. \quad \int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

output $1/d*a*(2/5*I*B*\tan(d*x+c)^{(5/2)}+2/3*I*A*\tan(d*x+c)^{(3/2)}+2/3*B*\tan(d*x+c)^{(3/2)}-2*I*B*\tan(d*x+c)^{(1/2)}+2*A*\tan(d*x+c)^{(1/2)}+1/4*(-A+I*B)*2^{(1/2)}*(\ln((1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}))+1/4*(-I*A-B)*2^{(1/2)}*(\ln((1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}))$

3.113.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 432 vs. $2(81) = 162$.

Time = 0.26 (sec) , antiderivative size = 432, normalized size of antiderivative = 4.11

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx =$$

$$15 \left(d e^{(4i dx+4i c)} + 2 d e^{(2i dx+2i c)} + d \right) \sqrt{-\frac{(i A^2+2 A B-i B^2)a^2}{d^2}} \log \left(\frac{2 \left((A-i B) a e^{(2i dx+2i c)} + (i d e^{(2i dx+2i c)} + i d) \sqrt{-\frac{(i A^2+2 A B-i B^2)a^2}{d^2}} \right)}{(i A+B) \sqrt{-\frac{(i A^2+2 A B-i B^2)a^2}{d^2}}} \right)$$

input `integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output $-1/30*(15*(d*e^{(4*I*d*x + 4*I*c)} + 2*d*e^{(2*I*d*x + 2*I*c)} + d)*\sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2}*\log(2*((A - I*B)*a*e^{(2*I*d*x + 2*I*c)} + (I*d*e^{(2*I*d*x + 2*I*c)} + I*d)*\sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2}*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))}*e^{(-2*I*d*x - 2*I*c)})/((I*A + B)*a)) - 15*(d*e^{(4*I*d*x + 4*I*c)} + 2*d*e^{(2*I*d*x + 2*I*c)} + d)*\sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2}*\log(2*((A - I*B)*a*e^{(2*I*d*x + 2*I*c)} + (-I*d*e^{(2*I*d*x + 2*I*c)} - I*d)*\sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2}*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))}*e^{(-2*I*d*x - 2*I*c)})/((I*A + B)*a)) - 4*((20*A - 23*I*B)*a*e^{(4*I*d*x + 4*I*c)} + 6*(5*A - 4*I*B)*a*e^{(2*I*d*x + 2*I*c)} + (10*A - 13*I*B)*a)*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))}/(d*e^{(4*I*d*x + 4*I*c)} + 2*d*e^{(2*I*d*x + 2*I*c)} + d)$

3.113.6 Sympy [F]

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= ia \left(\int A \tan^{\frac{5}{2}}(c + dx) dx + \int B \tan^{\frac{7}{2}}(c + dx) dx + \int \left(-iA \tan^{\frac{3}{2}}(c + dx) \right) dx \right. \\ \left. + \int \left(-iB \tan^{\frac{5}{2}}(c + dx) \right) dx \right)$$

input `integrate(tan(d*x+c)**(3/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `I*a*(Integral(A*tan(c + d*x)**(5/2), x) + Integral(B*tan(c + d*x)**(7/2), x) + Integral(-I*A*tan(c + d*x)**(3/2), x) + Integral(-I*B*tan(c + d*x)**(5/2), x))`

3.113.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 186 vs. $2(81) = 162$.

Time = 0.31 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.77

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx =$$

$$\frac{-24iBa \tan(dx + c)^{\frac{5}{2}} + 40(-iA - B)a \tan(dx + c)^{\frac{3}{2}} - 120(A - iB)a \sqrt{\tan(dx + c)} - 15 \left(2\sqrt{2}(-\right)}{d}$$

input `integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/60*(-24*I*B*a*tan(d*x + c)^(5/2) + 40*(-I*A - B)*a*tan(d*x + c)^(3/2) - 120*(A - I*B)*a*sqrt(tan(d*x + c)) - 15*(2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a)/d`

3.113.8 Giac [A] (verification not implemented)

Time = 0.56 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.07

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{(i-1) \sqrt{2}(iAa+Ba) \arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx+c)}\right)}{d} - \frac{2\left(-3iBad^4 \tan(dx+c)^{\frac{5}{2}} - 5iAad^4 \tan(dx+c)^{\frac{3}{2}} - 5Bad^4 \tan(dx+c)^{\frac{3}{2}} - 15Aad^4 \sqrt{\tan(dx+c)} + \dots\right)}{15d^5}$$

input `integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `(I - 1)*sqrt(2)*(I*A*a + B*a)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d - 2/15*(-3*I*B*a*d^4*tan(d*x + c)^(5/2) - 5*I*A*a*d^4*tan(d*x + c)^(3/2) - 5*B*a*d^4*tan(d*x + c)^(3/2) - 15*A*a*d^4*sqrt(tan(d*x + c)) + 15*I*B*a*d^4*sqrt(tan(d*x + c)))/d^5`

3.113.9 Mupad [B] (verification not implemented)

Time = 10.24 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.24

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{2Aa \sqrt{\tan(c+dx)}}{d} + \frac{Aa \tan(c+dx)^{3/2} 2i}{3d} - \frac{Ba \sqrt{\tan(c+dx)} 2i}{d}$$

$$+ \frac{2Ba \tan(c+dx)^{3/2}}{3d} + \frac{Ba \tan(c+dx)^{5/2} 2i}{5d}$$

$$+ \frac{\sqrt{2}Aa \operatorname{atan}\left(\sqrt{2} \sqrt{\tan(c+dx)} \left(\frac{1}{2} - \frac{1}{2}i\right)\right) (-1-i)}{d}$$

$$- \frac{(-1)^{1/4} Ba \operatorname{atan}\left((-1)^{1/4} \sqrt{\tan(c+dx)} li\right) 2i}{d}$$

input `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`

output $(2Aa \tan(c + dx)^{1/2})/d + (Aa \tan(c + dx)^{3/2} 2i)/(3d) - (Ba \tan(c + dx)^{1/2} 2i)/d + (2Ba \tan(c + dx)^{3/2})/(3d) + (Ba \tan(c + dx)^{5/2} 2i)/(5d) - (2^{1/2} Aa \operatorname{atan}(2^{1/2} \tan(c + dx)^{1/2} (1/2 - 1i/2)) (1 + 1i))/d - ((-1)^{1/4} Ba \operatorname{atan}((-1)^{1/4} \tan(c + dx)^{1/2} 1i) 2i)/d$

3.114 $\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

3.114.1 Optimal result	1216
3.114.2 Mathematica [A] (verified)	1216
3.114.3 Rubi [A] (verified)	1217
3.114.4 Maple [B] (verified)	1219
3.114.5 Fricas [B] (verification not implemented)	1220
3.114.6 Sympy [F]	1220
3.114.7 Maxima [B] (verification not implemented)	1221
3.114.8 Giac [A] (verification not implemented)	1221
3.114.9 Mupad [B] (verification not implemented)	1222

3.114.1 Optimal result

Integrand size = 34, antiderivative size = 80

$$\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2\sqrt[4]{-1}a(iA + B) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right)}{d} + \frac{2a(iA + B)\sqrt{\tan(c + dx)}}{d} + \frac{2iaB \tan^{3/2}(c + dx)}{3d}$$

```
output 2*(-1)^(1/4)*a*(I*A+B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d+2*a*(I*A+B)*tan(d*x+c)^(1/2)/d+2/3*I*a*B*tan(d*x+c)^(3/2)/d
```

3.114.2 Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.89

$$\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2a\left(3\sqrt[4]{-1}(iA + B) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right) + \sqrt{\tan(c + dx)}(3iA + 3B + iB \tan(c + dx))\right)}{3d}$$

input `Integrate[Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x
]`

output `(2*a*(3*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + Sqrt[Tan[c + d*x]]*((3*I)*A + 3*B + I*B*Tan[c + d*x]))/(3*d)`

3.114.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.206$, Rules used = {3042, 4075, 3042, 4011, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4075} \\
 & \int \sqrt{\tan(c + dx)}(a(A - iB) + a(iA + B) \tan(c + dx)) dx + \frac{2iaB \tan^{\frac{3}{2}}(c + dx)}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\tan(c + dx)}(a(A - iB) + a(iA + B) \tan(c + dx)) dx + \frac{2iaB \tan^{\frac{3}{2}}(c + dx)}{3d} \\
 & \quad \downarrow \text{4011} \\
 & \int \frac{a(A - iB) \tan(c + dx) - a(iA + B)}{\sqrt{\tan(c + dx)}} dx + \frac{2a(B + iA) \sqrt{\tan(c + dx)}}{d} + \frac{2iaB \tan^{\frac{3}{2}}(c + dx)}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{a(A - iB) \tan(c + dx) - a(iA + B)}{\sqrt{\tan(c + dx)}} dx + \frac{2a(B + iA) \sqrt{\tan(c + dx)}}{d} + \frac{2iaB \tan^{\frac{3}{2}}(c + dx)}{3d} \\
 & \quad \downarrow \text{4016}
 \end{aligned}$$

3.114. $\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\frac{2a^2(B + iA)^2 \int \frac{1}{-a(iA+B) - a(A-iB)\tan(c+dx)} d\sqrt{\tan(c+dx)} + \frac{2a(B + iA)\sqrt{\tan(c+dx)}}{d} + \frac{2iaB \tan^{\frac{3}{2}}(c+dx)}{3d}$$

↓ 218

$$\frac{2\sqrt[4]{-1}a(B + iA) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{d} + \frac{2a(B + iA)\sqrt{\tan(c+dx)}}{d} + \frac{2iaB \tan^{\frac{3}{2}}(c+dx)}{3d}$$

input `Int[Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(2*(-1)^(1/4)*a*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]])/d + (2*a*(I*A + B)*Sqrt[Tan[c + d*x]])/d + (((2*I)/3)*a*B*Tan[c + d*x]^(3/2))/d`

3.114.3.1 Defintions of rubi rules used

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

3.114.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 225 vs. 2(65) = 130.

Time = 0.03 (sec) , antiderivative size = 226, normalized size of antiderivative = 2.82

method	result
derivativedivides	$a \left(\frac{2iB \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + 2iA(\sqrt{\tan(dx+c)}) + 2B(\sqrt{\tan(dx+c)}) + \frac{(-iA-B)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)} \right) + 2 \arctan \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)} \right)}{4} \right)$
default	$a \left(\frac{2iB \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + 2iA(\sqrt{\tan(dx+c)}) + 2B(\sqrt{\tan(dx+c)}) + \frac{(-iA-B)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)} \right) + 2 \arctan \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)} \right)}{4} \right)$
parts	$(iaA+Ba) \left(2(\sqrt{\tan(dx+c)}) - \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)} \right) + 2 \arctan \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)} \right) + 2 \arctan \left(-1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right)}{4} \right) \right) / d$

```
input int(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)), x, method=_RETURNV
ERBOSE)
```

```
output 1/d*a*(2/3*I*B*tan(d*x+c)^(3/2)+2*I*A*tan(d*x+c)^(1/2)+2*B*tan(d*x+c)^(1/2
)+1/4*(-I*A-B)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1
/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*a
rctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(A-I*B)*2^(1/2)*(ln((1-2^(1/2)*tan
(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arcta
n(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))
```

$$3.114. \int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

3.114.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 372 vs. $2(62) = 124$.

Time = 0.25 (sec) , antiderivative size = 372, normalized size of antiderivative = 4.65

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx =$$

$$3 \left(de^{(2i dx+2i c)} + d \right) \sqrt{-\frac{(-i A^2-2 AB+i B^2)a^2}{d^2}} \log \left(\frac{2 \left((A-i B) a e^{(2i dx+2i c)} + (d e^{(2i dx+2i c)} + d) \sqrt{-\frac{(-i A^2-2 AB+i B^2)a^2}{d^2}} \sqrt{\frac{-i}{e}} \right)}{(i A+B) a} \right)$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-1/6*(3*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*log(2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) + (d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a) - 3*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*log(2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) - (d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a) + 4*((-3*I*A - 4*B)*a*e^(2*I*d*x + 2*I*c) + (-3*I*A - 2*B)*a)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(2*I*d*x + 2*I*c) + d)`

3.114.6 Sympy [F]

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= ia \left(\int A \tan^{\frac{3}{2}}(c+dx) dx + \int B \tan^{\frac{5}{2}}(c+dx) dx + \int \left(-iA \sqrt{\tan(c+dx)} \right) dx \right. \\ \left. + \int \left(-iB \tan^{\frac{3}{2}}(c+dx) \right) dx \right)$$

input `integrate(tan(d*x+c)**(1/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `I*a*(Integral(A*tan(c + d*x)**(3/2), x) + Integral(B*tan(c + d*x)**(5/2), x) + Integral(-I*A*sqrt(tan(c + d*x)), x) + Integral(-I*B*tan(c + d*x)**(3/2), x))`

3.114.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 170 vs. $2(62) = 124$.

Time = 0.33 (sec) , antiderivative size = 170, normalized size of antiderivative = 2.12

$$\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx =$$

$$\frac{-8i Ba \tan(dx + c)^{\frac{3}{2}} + 24(-iA - B)a\sqrt{\tan(dx + c)} + 3\left(2\sqrt{2}((i - 1)A + (i + 1)B) \arctan\left(\frac{1}{2}\sqrt{2}\right)\right)}{d}$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/12*(-8*I*B*a*tan(d*x + c)^(3/2) + 24*(-I*A - B)*a*sqrt(tan(d*x + c)) + 3*(2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a)/d`

3.114.8 Giac [A] (verification not implemented)

Time = 0.46 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.02

$$\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{(i - 1) \sqrt{2}(Aa - i Ba) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2}\sqrt{\tan(dx + c)}\right)}{d}$$

$$- \frac{2\left(-i Bad^2 \tan(dx + c)^{\frac{3}{2}} - 3i Aad^2 \sqrt{\tan(dx + c)} - 3 Bad^2 \sqrt{\tan(dx + c)}\right)}{3d^3}$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output $-(I - 1)\sqrt{2}*(A*a - I*B*a)*\arctan(-\frac{1}{2}I - \frac{1}{2})\sqrt{2}*\sqrt{\tan(d*x + c))/d - 2/3*(-I*B*a*d^2*\tan(d*x + c)^{(3/2)} - 3*I*A*a*d^2*\sqrt{\tan(d*x + c)} - 3*B*a*d^2*\sqrt{\tan(d*x + c)})/d^3$

3.114.9 Mupad [B] (verification not implemented)

Time = 9.75 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.24

$$\begin{aligned} & \int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\ &= \frac{A a \sqrt{\tan(c + dx)} 2i}{d} + \frac{2 B a \sqrt{\tan(c + dx)}}{d} + \frac{B a \tan(c + dx)^{3/2} 2i}{3d} \\ & \quad - \frac{2(-1)^{1/4} A a \operatorname{atanh}\left((-1)^{1/4} \sqrt{\tan(c + dx)}\right)}{d} \\ & \quad + \frac{\sqrt{2} B a \operatorname{atan}\left(\sqrt{2} \sqrt{\tan(c + dx)} \left(\frac{1}{2} - \frac{1}{2}i\right)\right) (-1 - i)}{d} \end{aligned}$$

input `int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`

output $(A*a*\tan(c + d*x)^{(1/2)}*2i)/d + (2*B*a*\tan(c + d*x)^{(1/2)})/d + (B*a*\tan(c + d*x)^{(3/2)}*2i)/(3*d) - (2*(-1)^{(1/4)}*A*a*\operatorname{atanh}((-1)^{(1/4)}*\tan(c + d*x)^{(1/2)}))/d - (2^{(1/2)}*B*a*\operatorname{atan}(2^{(1/2)}*\tan(c + d*x)^{(1/2)}*(1/2 - 1i/2))*(1 + 1i))/d$

3.115
$$\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.115.1 Optimal result 1223
 3.115.2 Mathematica [A] (verified) 1223
 3.115.3 Rubi [A] (verified) 1224
 3.115.4 Maple [B] (verified) 1225
 3.115.5 Fricas [B] (verification not implemented) 1226
 3.115.6 Sympy [F] 1227
 3.115.7 Maxima [B] (verification not implemented) 1227
 3.115.8 Giac [A] (verification not implemented) 1228
 3.115.9 Mupad [B] (verification not implemented) 1228

3.115.1 Optimal result

Integrand size = 34, antiderivative size = 55

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= -\frac{2\sqrt[4]{-1}a(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d} + \frac{2iaB\sqrt{\tan(c + dx)}}{d}$$

output `-2*(-1)^(1/4)*a*(A-I*B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d+2*I*a*B*tan(d*x+c)^(1/2)/d`

3.115.2 Mathematica [A] (verified)

Time = 0.59 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= -\frac{2\sqrt[4]{-1}a(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d} + \frac{2iaB\sqrt{\tan(c + dx)}}{d}$$

input `Integrate[((a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output $(-2*(-1)^{(1/4)}*a*(A - I*B)*ArcTan[(-1)^{(3/4)}*Sqrt[Tan[c + d*x]]])/d + ((2*I)*a*B*Sqrt[Tan[c + d*x]])/d$

3.115.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.147$, Rules used = {3042, 4075, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\ & \quad \downarrow \text{4075} \\ & \int \frac{a(A - iB) + a(iA + B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2iaB \sqrt{\tan(c + dx)}}{d} \\ & \quad \downarrow \text{3042} \\ & \int \frac{a(A - iB) + a(iA + B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2iaB \sqrt{\tan(c + dx)}}{d} \\ & \quad \downarrow \text{4016} \\ & \frac{2a^2(A - iB)^2 \int \frac{1}{a(A - iB) - a(iA + B) \tan(c + dx)} d \sqrt{\tan(c + dx)}}{d} + \frac{2iaB \sqrt{\tan(c + dx)}}{d} \\ & \quad \downarrow \text{218} \\ & \frac{2iaB \sqrt{\tan(c + dx)}}{d} - \frac{2\sqrt[4]{-1}a(A - iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right)}{d} \end{aligned}$$

input $\text{Int}[(a + I*a*\text{Tan}[c + d*x])*(A + B*\text{Tan}[c + d*x])/Sqrt[\text{Tan}[c + d*x]],x]$

output $(-2*(-1)^{(1/4)}*a*(A - I*B)*ArcTan[(-1)^{(3/4)}*Sqrt[Tan[c + d*x]]])/d + ((2*I)*a*B*Sqrt[Tan[c + d*x]])/d$

3.115. $\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$

3.115.3.1 Defintions of rubi rules used

- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

- rule 4016 `Int[((c_) + (d_.)*tan[(e_) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_) + (f_.)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

- rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.115.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 200 vs. 2(45) = 90.

Time = 0.03 (sec) , antiderivative size = 201, normalized size of antiderivative = 3.65

method	result
derivativedivides	$a \left(2iB(\sqrt{\tan(dx+c)}) + \frac{(-iB+A)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4} \right)$
default	$a \left(2iB(\sqrt{\tan(dx+c)}) + \frac{(-iB+A)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4} \right)$
parts	$\frac{(iA+Ba)\sqrt{2} \left(\ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4d}$

3.115. $\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

input `int((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)`

output `1/d*a*(2*I*B*tan(d*x+c)^(1/2)+1/4*(A-I*B)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)
)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(
1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(I*A+B)
2^(1/2)(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)
^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/
2)*tan(d*x+c)^(1/2)))`

3.115.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 314 vs. $2(43) = 86$.

Time = 0.25 (sec) , antiderivative size = 314, normalized size of antiderivative = 5.71

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx =$$

$$-4i Ba \sqrt{\frac{-i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} + 1}} - \sqrt{\frac{(i A^2 + 2 AB - i B^2) a^2}{d^2}} d \log \left(\frac{2 \left((A - i B) a e^{(2i dx + 2i c)} + (i d e^{(2i dx + 2i c)} + i d) \sqrt{\frac{(i A^2 + 2 AB - i B^2) a^2}{d^2}} \right)}{(i A + B) a} \right)$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorith
m="fricas")`

output `-1/2*(-4*I*B*a*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)
) - sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*d*log(2*((A - I*B)*a*e^(2*I*d*x
+ 2*I*c) + (I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*
a^2/d^2)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(
-2*I*d*x - 2*I*c)/((I*A + B)*a)) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*
d*log(2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) + (-I*d*e^(2*I*d*x + 2*I*c) - I*d
) * sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2) * sqrt((-I*e^(2*I*d*x + 2*I*c) + I)
/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x - 2*I*c)/((I*A + B)*a))/d`

3.115.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= ia \left(\int A \sqrt{\tan(c + dx)} dx + \int B \tan^{\frac{3}{2}}(c + dx) dx + \int \left(-\frac{iA}{\sqrt{\tan(c + dx)}} \right) dx \right. \\ \left. + \int \left(-iB \sqrt{\tan(c + dx)} \right) dx \right)$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)**(1/2),x)`

output `I*a*(Integral(A*sqrt(tan(c + d*x)), x) + Integral(B*tan(c + d*x)**(3/2), x) + Integral(-I*A/sqrt(tan(c + d*x)), x) + Integral(-I*B*sqrt(tan(c + d*x)), x))`

3.115.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 151 vs. $2(43) = 86$.

Time = 0.28 (sec) , antiderivative size = 151, normalized size of antiderivative = 2.75

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx =$$

$$\frac{-8iBa\sqrt{\tan(dx+c)} + \left(2\sqrt{2}(-(i+1)A + (i-1)B)\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(dx+c)}\right)\right) + 2\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - 2\sqrt{\tan(dx+c)}\right)\right)\right)}{d}$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="maxima")`

output `-1/4*(-8*I*B*a*sqrt(tan(d*x + c)) + (2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a/d`

3.115.8 Giac [A] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.85

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \frac{(i - 1) \sqrt{2}(-i Aa - Ba) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{d} + \frac{2i Ba \sqrt{\tan(dx + c)}}{d}$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="giac")`

output `(I - 1)*sqrt(2)*(-I*A*a - B*a)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d + 2*I*B*a*sqrt(tan(d*x + c))/d`

3.115.9 Mupad [B] (verification not implemented)

Time = 9.18 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.24

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= -\frac{2(-1)^{1/4} B a \operatorname{atanh}\left((-1)^{1/4} \sqrt{\tan(c + dx)}\right)}{d} + \frac{B a \sqrt{\tan(c + dx)} 2i}{d}$$

$$+ \frac{\sqrt{2} A a \operatorname{atan}\left(\sqrt{2} \sqrt{\tan(c + dx)} \left(\frac{1}{2} - \frac{1}{2}i\right)\right) (1 + i)}{d}$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i))/tan(c + d*x)^(1/2),x)`

output `(B*a*tan(c + d*x)^(1/2)*2i)/d + (2^(1/2)*A*a*atan(2^(1/2)*tan(c + d*x)^(1/2)*(1/2 - 1i/2))*(1 + 1i))/d - (2*(-1)^(1/4)*B*a*atanh((-1)^(1/4)*tan(c + d*x)^(1/2)))/d`

3.116
$$\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.116.1 Optimal result 1229
 3.116.2 Mathematica [A] (verified) 1229
 3.116.3 Rubi [A] (verified) 1230
 3.116.4 Maple [B] (verified) 1231
 3.116.5 Fracas [B] (verification not implemented) 1232
 3.116.6 Sympy [F] 1233
 3.116.7 Maxima [B] (verification not implemented) 1233
 3.116.8 Giac [A] (verification not implemented) 1234
 3.116.9 Mupad [B] (verification not implemented) 1234

3.116.1 Optimal result

Integrand size = 34, antiderivative size = 53

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= -\frac{2\sqrt[4]{-1}a(iA + B) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right)}{d} - \frac{2aA}{d\sqrt{\tan(c + dx)}}$$

output `-2*(-1)^(1/4)*a*(I*A+B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d-2*a*A/d/tan(d*x+c)^(1/2)`

3.116.2 Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.02

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= -\frac{2ia\left(\sqrt[4]{-1}(A - iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right) - \frac{iA}{\sqrt{\tan(c + dx)}}\right)}{d}$$

input `Integrate[((a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output $((-2*I)*a*(-1)^{(1/4)}*(A - I*B)*ArcTan[(-1)^{(3/4)}*Sqrt[Tan[c + d*x]]] - (I*A)/Sqrt[Tan[c + d*x]])/d$

3.116.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.147$, Rules used = {3042, 4074, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx \\ & \quad \downarrow \text{4074} \\ & -\frac{2aA}{d\sqrt{\tan(c + dx)}} + \int \frac{a(iA + B) - a(A - iB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & -\frac{2aA}{d\sqrt{\tan(c + dx)}} + \int \frac{a(iA + B) - a(A - iB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx \\ & \quad \downarrow \text{4016} \\ & -\frac{2aA}{d\sqrt{\tan(c + dx)}} + \frac{2a^2(B + iA)^2 \int \frac{1}{a(iA+B)+a(A-iB)\tan(c+dx)} d\sqrt{\tan(c + dx)}}{d} \\ & \quad \downarrow \text{218} \\ & -\frac{2aA}{d\sqrt{\tan(c + dx)}} - \frac{2\sqrt{-1}a(B + iA) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d} \end{aligned}$$

input $\text{Int}[(a + I*a*\text{Tan}[c + d*x])*(A + B*\text{Tan}[c + d*x])/ \text{Tan}[c + d*x]^{(3/2)}, x]$

output $(-2*(-1)^{(1/4)}*a*(I*A + B)*ArcTan[(-1)^{(3/4)}*Sqrt[Tan[c + d*x]]])/d - (2*a*A)/(d*Sqrt[Tan[c + d*x]])$

3.116. $\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

3.116.3.1 Defintions of rubi rules used

- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

- rule 4016 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

- rule 4074 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.116.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 201 vs. 2(44) = 88.

Time = 0.04 (sec) , antiderivative size = 202, normalized size of antiderivative = 3.81

method	result
derivativedivides	$a \left(-\frac{2A}{\sqrt{\tan(dx+c)}} + \frac{(iA+B)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right)}{4} \right)$
default	$a \left(-\frac{2A}{\sqrt{\tan(dx+c)}} + \frac{(iA+B)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right)}{4} \right)$
parts	$\frac{(iaA+Ba)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right)}{4d}$

3.116. $\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

```
input int((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output 1/d*a*(-2*A/tan(d*x+c)^(1/2)+1/4*(I*A+B)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)
^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))) +2*arctan(1+2^(
1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(-A+I*B)
*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)
^(1/2)+tan(d*x+c))) +2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/
2)*tan(d*x+c)^(1/2))))
```

3.116.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 367 vs. $2(43) = 86$.

Time = 0.26 (sec) , antiderivative size = 367, normalized size of antiderivative = 6.92

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{(de^{(2i dx + 2i c)} - d) \sqrt{-\frac{(-i A^2 - 2AB + i B^2)a^2}{d^2}} \log \left(\frac{2 \left((A - i B) a e^{(2i dx + 2i c)} + (d e^{(2i dx + 2i c)} + d) \sqrt{-\frac{(-i A^2 - 2AB + i B^2)a^2}{d^2}} \right) \sqrt{\frac{-i e^{(2i dx + 2i c)}}{e^{(2i dx + 2i c)}}}}{(i A + B) a}} \right)}{1}$$

```
input integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorith
m="fricas")
```

```
output 1/2*((d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*1
og(2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) + (d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-
(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*
I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a)) - (d*e^(2*I*d*x
+ 2*I*c) - d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*log(2*((A - I*B)*a*e
^(2*I*d*x + 2*I*c) - (d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2*A*B + I
*B^2)*a^2/d^2)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)
))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a)) - 4*(I*A*a*e^(2*I*d*x + 2*I*c) + I*
A*a)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(2
*I*d*x + 2*I*c) - d)
```

3.116. $\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$

3.116.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= ia \left(\int \frac{A}{\sqrt{\tan(c + dx)}} dx + \int B \sqrt{\tan(c + dx)} dx + \int \left(-\frac{iA}{\tan^{\frac{3}{2}}(c + dx)} \right) dx \right. \\ \left. + \int \left(-\frac{iB}{\sqrt{\tan(c + dx)}} \right) dx \right)$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)**(3/2),x)`

output `I*a*(Integral(A/sqrt(tan(c + d*x)), x) + Integral(B*sqrt(tan(c + d*x)), x) + Integral(-I*A/tan(c + d*x)**(3/2), x) + Integral(-I*B/sqrt(tan(c + d*x)), x))`

3.116.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 151 vs. $2(43) = 86$.

Time = 0.31 (sec) , antiderivative size = 151, normalized size of antiderivative = 2.85

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{\left(2\sqrt{2}((i-1)A + (i+1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(dx+c)}\right)\right) + 2\sqrt{2}((i-1)A + (i+1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - 2\sqrt{\tan(dx+c)}\right)\right) - \sqrt{2}(-(i+1)A + (i-1)B) \log(\sqrt{2}\sqrt{\tan(dx+c)} + \tan(dx+c) + 1) + \sqrt{2}(-(i+1)A + (i-1)B) \log(-\sqrt{2}\sqrt{\tan(dx+c)} + \tan(dx+c) + 1) \right) a - 8Aa/\sqrt{\tan(dx+c)}}{d}$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="maxima")`

output `1/4*((2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a - 8*A*a/sqrt(tan(d*x + c))/d`

3.116. $\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

3.116.8 Giac [A] (verification not implemented)

Time = 0.62 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.87

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{(i - 1) \sqrt{2}(Aa - i Ba) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{d} - \frac{2Aa}{d \sqrt{\tan(dx + c)}}$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="giac")`

output `((I - 1)*sqrt(2)*(A*a - I*B*a)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d - 2*A*a/(d*sqrt(tan(d*x + c))))`

3.116.9 Mupad [B] (verification not implemented)

Time = 8.14 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.26

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{2(-1)^{1/4} Aa \operatorname{atanh}\left((-1)^{1/4} \sqrt{\tan(c + dx)}\right)}{d} - \frac{2Aa}{d \sqrt{\tan(c + dx)}}$$

$$+ \frac{\sqrt{2} B a \operatorname{atan}\left(\sqrt{2} \sqrt{\tan(c + dx)} \left(\frac{1}{2} - \frac{1}{2}i\right)\right) (1 + i)}{d}$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i))/tan(c + d*x)^(3/2),x)`

output `(2*(-1)^(1/4)*A*a*atanh((-1)^(1/4)*tan(c + d*x)^(1/2))/d - (2*A*a)/(d*tan(c + d*x)^(1/2)) + (2^(1/2)*B*a*atan(2^(1/2)*tan(c + d*x)^(1/2)*(1/2 - 1i/2))*(1 + 1i))/d`

$$3.117 \quad \int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.117.1 Optimal result	1235
3.117.2 Mathematica [C] (verified)	1235
3.117.3 Rubi [A] (verified)	1236
3.117.4 Maple [B] (verified)	1238
3.117.5 Fricas [B] (verification not implemented)	1239
3.117.6 Sympy [F]	1239
3.117.7 Maxima [B] (verification not implemented)	1240
3.117.8 Giac [A] (verification not implemented)	1240
3.117.9 Mupad [B] (verification not implemented)	1241

3.117.1 Optimal result

Integrand size = 34, antiderivative size = 78

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2\sqrt[4]{-1}a(A - iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right)}{d} - \frac{2aA}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2a(iA + B)}{d \sqrt{\tan(c + dx)}}$$

```
output 2*(-1)^(1/4)*a*(A-I*B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d-2*a*(I*A+B)/d
/tan(d*x+c)^(1/2)-2/3*a*A/d/tan(d*x+c)^(3/2)
```

3.117.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.59 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.74

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{-2aA - 6ia(A - iB) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, i \tan(c + dx)\right) \tan(c + dx)}{3d \tan^{\frac{3}{2}}(c + dx)}$$

```
input Integrate[((a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2)
,x]
```

output $(-2*a*A - (6*I)*a*(A - I*B)*\text{Hypergeometric2F1}[-1/2, 1, 1/2, I*\text{Tan}[c + d*x]]*\text{Tan}[c + d*x])/(3*d*\text{Tan}[c + d*x]^{(3/2)})$

3.117.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3042, 4074, 3042, 4012, 25, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx \\
 & \quad \downarrow \text{4074} \\
 & -\frac{2aA}{3d \tan^{\frac{3}{2}}(c + dx)} + \int \frac{a(iA + B) - a(A - iB) \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2aA}{3d \tan^{\frac{3}{2}}(c + dx)} + \int \frac{a(iA + B) - a(A - iB) \tan(c + dx)}{\tan(c + dx)^{3/2}} dx \\
 & \quad \downarrow \text{4012} \\
 & \int -\frac{a(A - iB) + a(iA + B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2a(B + iA)}{d\sqrt{\tan(c + dx)}} - \frac{2aA}{3d \tan^{\frac{3}{2}}(c + dx)} \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{a(A - iB) + a(iA + B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2a(B + iA)}{d\sqrt{\tan(c + dx)}} - \frac{2aA}{3d \tan^{\frac{3}{2}}(c + dx)} \\
 & \quad \downarrow \text{3042} \\
 & -\int \frac{a(A - iB) + a(iA + B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2a(B + iA)}{d\sqrt{\tan(c + dx)}} - \frac{2aA}{3d \tan^{\frac{3}{2}}(c + dx)} \\
 & \quad \downarrow \text{4016}
 \end{aligned}$$

3.117. $\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$

$$\frac{2a^2(A - iB)^2 \int \frac{1}{a(A - iB) - a(iA + B)\tan(c + dx)} d\sqrt{\tan(c + dx)}}{d} - \frac{2a(B + iA)}{d\sqrt{\tan(c + dx)}} - \frac{2aA}{3d\tan^{\frac{3}{2}}(c + dx)}$$

↓ 218

$$\frac{2\sqrt[4]{-1}a(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d} - \frac{2a(B + iA)}{d\sqrt{\tan(c + dx)}} - \frac{2aA}{3d\tan^{\frac{3}{2}}(c + dx)}$$

input `Int[((a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(2*(-1)^(1/4)*a*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]/d - (2*a*A)/(3*d*Tan[c + d*x]^(3/2)) - (2*a*(I*A + B))/(d*Sqrt[Tan[c + d*x]])`

3.117.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

3.117.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 219 vs. 2(64) = 128.

Time = 0.03 (sec) , antiderivative size = 220, normalized size of antiderivative = 2.82

method	result
derivativedivides	$a \left(-\frac{2A}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(iA+B)}{\sqrt{\tan(dx+c)}} + \frac{(iB-A)\sqrt{2}}{4} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(1-\sqrt{2}(\sqrt{\tan(dx+c)})) \right) \right)$
default	$a \left(-\frac{2A}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(iA+B)}{\sqrt{\tan(dx+c)}} + \frac{(iB-A)\sqrt{2}}{4} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(1-\sqrt{2}(\sqrt{\tan(dx+c)})) \right) \right)$
parts	$(iaA+Ba) \left(-\frac{\sqrt{2} \left(\ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4} - \dots \right)$

```
input int((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2), x, method=_RETURNV
ERBOSE)
```

```
output 1/d*a*(-2/3*A/tan(d*x+c)^(3/2)-2*(I*A+B)/tan(d*x+c)^(1/2)+1/4*(-A+I*B)*2^(
1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/
2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*t
an(d*x+c)^(1/2)))+1/4*(-I*A-B)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan
(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d
*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))
```

$$3.117. \int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.117.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 427 vs. $2(62) = 124$.

Time = 0.28 (sec) , antiderivative size = 427, normalized size of antiderivative = 5.47

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx =$$

$$3 \left(de^{(4i dx + 4i c)} - 2 de^{(2i dx + 2i c)} + d \right) \sqrt{-\frac{(iA^2 + 2AB - iB^2)a^2}{d^2}} \log \left(\frac{2 \left((A - iB)ae^{(2i dx + 2i c)} + (i de^{(2i dx + 2i c)} + i d) \sqrt{-\frac{(iA^2}{(iA+B)}}} \right)}{(iA+B)} \right)$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="fricas")`

output `-1/6*(3*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*log(2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) + (I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a) - 3*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*log(2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) + (-I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a) - 4*((4*A - 3*I*B)*a*e^(4*I*d*x + 4*I*c) + 2*A*a*e^(2*I*d*x + 2*I*c) - (2*A - 3*I*B)*a)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))/(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)`

3.117.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= ia \left(\int \frac{A}{\tan^{\frac{3}{2}}(c + dx)} dx + \int \frac{B}{\sqrt{\tan(c + dx)}} dx + \int \left(-\frac{iA}{\tan^{\frac{5}{2}}(c + dx)} \right) dx \right. \\ \left. + \int \left(-\frac{iB}{\tan^{\frac{3}{2}}(c + dx)} \right) dx \right)$$

3.117. $\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)**(5/2),x)`

output `I*a*(Integral(A/tan(c + d*x)**(3/2), x) + Integral(B/sqrt(tan(c + d*x)), x) + Integral(-I*A/tan(c + d*x)**(5/2), x) + Integral(-I*B/tan(c + d*x)**(3/2), x))`

3.117.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 171 vs. $2(62) = 124$.

Time = 0.30 (sec) , antiderivative size = 171, normalized size of antiderivative = 2.19

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{3 \left(2\sqrt{2}(-i+1)A + (i-1)B \right) \arctan \left(\frac{1}{2}\sqrt{2} \left(\sqrt{2} + 2\sqrt{\tan(dx+c)} \right) \right) + 2\sqrt{2}(-i+1)A + (i-1)B}{d}$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="maxima")`

output `1/12*(3*(2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a + 8*(3*(-I*A - B)*a*tan(d*x + c) - A*a)/tan(d*x + c)^(3/2))/d`

3.117.8 Giac [A] (verification not implemented)

Time = 0.72 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.88

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= -\frac{(i-1)\sqrt{2}(-iAa - Ba) \arctan \left(-\left(\frac{1}{2}i - \frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)} \right)}{d}$$

$$- \frac{2(3iAa \tan(dx+c) + 3Ba \tan(dx+c) + Aa)}{3d \tan(dx+c)^{\frac{3}{2}}}$$

3.117. $\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm m="giac")`

output `-(I - 1)*sqrt(2)*(-I*A*a - B*a)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d - 2/3*(3*I*A*a*tan(d*x + c) + 3*B*a*tan(d*x + c) + A*a)/(d*tan(d*x + c)^(3/2))`

3.117.9 Mupad [B] (verification not implemented)

Time = 9.54 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.27

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2(-1)^{1/4} B a \operatorname{atanh}\left((-1)^{1/4} \sqrt{\tan(c + dx)}\right)}{d} - \frac{2 B a}{d \sqrt{\tan(c + dx)}}$$

$$- \frac{\frac{2 A a}{3d} + \frac{A a \tan(c+dx) 2i}{d}}{\tan(c + dx)^{3/2}} + \frac{\sqrt{2} A a \operatorname{atan}\left(\sqrt{2} \sqrt{\tan(c + dx)} \left(\frac{1}{2} - \frac{1}{2}i\right)\right) (-1 - i)}{d}$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i))/tan(c + d*x)^(5/2),x)`

output `(2*(-1)^(1/4)*B*a*atanh((-1)^(1/4)*tan(c + d*x)^(1/2)))/d - (2*B*a)/(d*tan(c + d*x)^(1/2)) - (2^(1/2)*A*a*atan(2^(1/2)*tan(c + d*x)^(1/2)*(1/2 - 1i/2))*(1 + 1i))/d - ((2*A*a)/(3*d) + (A*a*tan(c + d*x)*2i)/d)/tan(c + d*x)^(3/2)`

3.118
$$\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

3.118.1 Optimal result 1242
 3.118.2 Mathematica [C] (verified) 1242
 3.118.3 Rubi [A] (verified) 1243
 3.118.4 Maple [B] (verified) 1245
 3.118.5 Fricas [B] (verification not implemented) 1246
 3.118.6 Sympy [F] 1247
 3.118.7 Maxima [B] (verification not implemented) 1248
 3.118.8 Giac [A] (verification not implemented) 1248
 3.118.9 Mupad [B] (verification not implemented) 1249

3.118.1 Optimal result

Integrand size = 34, antiderivative size = 103

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{2^{\frac{4}{5}} \sqrt{-1} a (iA + B) \arctan \left((-1)^{\frac{3}{4}} \sqrt{\tan(c + dx)} \right)}{d} - \frac{2aA}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{2a(iA + B)}{3d \tan^{\frac{3}{2}}(c + dx)} + \frac{2a(A - iB)}{d \sqrt{\tan(c + dx)}}$$

```
output 2*(-1)^(1/4)*a*(I*A+B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d+2*a*(A-I*B)/d
/tan(d*x+c)^(1/2)-2/5*a*A/d/tan(d*x+c)^(5/2)-2/3*a*(I*A+B)/d/tan(d*x+c)^(3
/2)
```

3.118.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.81 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.55

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{2a(-3A - 5i(A - iB) \text{Hypergeometric2F1}(-\frac{3}{2}, 1, -\frac{1}{2}, i \tan(c + dx)) \tan(c + dx))}{15d \tan^{\frac{5}{2}}(c + dx)}$$

input `Integrate[((a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(2*a*(-3*A - (5*I)*(A - I*B)*Hypergeometric2F1[-3/2, 1, -1/2, I*Tan[c + d*x]]*Tan[c + d*x]))/(15*d*Tan[c + d*x]^(5/2))`

3.118.3 Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3042, 4074, 3042, 4012, 25, 3042, 4012, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan(c + dx)^{7/2}} dx \\
 & \quad \downarrow \text{4074} \\
 & -\frac{2aA}{5d \tan^{\frac{5}{2}}(c + dx)} + \int \frac{a(iA + B) - a(A - iB) \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2aA}{5d \tan^{\frac{5}{2}}(c + dx)} + \int \frac{a(iA + B) - a(A - iB) \tan(c + dx)}{\tan(c + dx)^{5/2}} dx \\
 & \quad \downarrow \text{4012} \\
 & \int -\frac{a(A - iB) + a(iA + B) \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)} dx - \frac{2a(B + iA)}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2aA}{5d \tan^{\frac{5}{2}}(c + dx)} \\
 & \quad \downarrow \text{25} \\
 & -\int \frac{a(A - iB) + a(iA + B) \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)} dx - \frac{2a(B + iA)}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2aA}{5d \tan^{\frac{5}{2}}(c + dx)} \\
 & \quad \downarrow \text{3042} \\
 & -\int \frac{a(A - iB) + a(iA + B) \tan(c + dx)}{\tan(c + dx)^{3/2}} dx - \frac{2a(B + iA)}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2aA}{5d \tan^{\frac{5}{2}}(c + dx)}
 \end{aligned}$$

3.118. $\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
& - \int \frac{a(iA + B) - a(A - iB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2a(B + iA)}{3d \tan^{\frac{3}{2}}(c + dx)} + \frac{2a(A - iB)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{5d \tan^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{4012} \\
& - \int \frac{a(iA + B) - a(A - iB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2a(B + iA)}{3d \tan^{\frac{3}{2}}(c + dx)} + \frac{2a(A - iB)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{5d \tan^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& - \frac{2a^2(B + iA)^2 \int \frac{1}{a(iA + B) + a(A - iB) \tan(c + dx)} d \sqrt{\tan(c + dx)}}{d} - \frac{2a(B + iA)}{3d \tan^{\frac{3}{2}}(c + dx)} + \frac{2a(A - iB)}{d \sqrt{\tan(c + dx)}} - \\
& \quad \frac{2aA}{5d \tan^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{4016} \\
& \frac{2\sqrt[4]{-1}a(B + iA) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right)}{d} - \frac{2a(B + iA)}{3d \tan^{\frac{3}{2}}(c + dx)} + \frac{2a(A - iB)}{d \sqrt{\tan(c + dx)}} - \\
& \quad \frac{2aA}{5d \tan^{\frac{5}{2}}(c + dx)}
\end{aligned}$$

input `Int[((a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(2*(-1)^(1/4)*a*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]/d - (2*a*A)/(5*d*Tan[c + d*x]^(5/2)) - (2*a*(I*A + B))/(3*d*Tan[c + d*x]^(3/2)) + (2*a*(A - I*B))/(d*Sqrt[Tan[c + d*x]])`

3.118.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

3.118. $\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$

```
rule 4012 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]
```

```
rule 4016 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_
)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b
*Tan[e + f*x]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]
```

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

3.118.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 235 vs. 2(84) = 168.

Time = 0.04 (sec) , antiderivative size = 236, normalized size of antiderivative = 2.29

method	result
derivativedivides	$a \left(-\frac{2A}{5 \tan(dx+c)^{\frac{5}{2}}} - \frac{2(iB-A)}{\sqrt{\tan(dx+c)}} - \frac{2(iA+B)}{3 \tan(dx+c)^{\frac{3}{2}}} + \frac{(-iA-B)\sqrt{2}}{4} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) \right) \right)$
default	$a \left(-\frac{2A}{5 \tan(dx+c)^{\frac{5}{2}}} - \frac{2(iB-A)}{\sqrt{\tan(dx+c)}} - \frac{2(iA+B)}{3 \tan(dx+c)^{\frac{3}{2}}} + \frac{(-iA-B)\sqrt{2}}{4} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) \right) \right)$
parts	$(iA+Ba) \left(-\frac{2}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) \right)}{4} \right) \frac{1}{d}$

$$3.118. \int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

input `int((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x,method=_RETURNV
ERBOSE)`

output `1/d*a*(-2/5*A/tan(d*x+c)^(5/2)-2*(-A+I*B)/tan(d*x+c)^(1/2)-2/3*(I*A+B)/tan
(d*x+c)^(3/2)+1/4*(-I*A-B)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x
+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c
)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(A-I*B)*2^(1/2)*(ln((1
-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+
c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(
1/2))))`

3.118.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 486 vs. $2(81) = 162$.

Time = 0.28 (sec) , antiderivative size = 486, normalized size of antiderivative = 4.72

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx =$$

$$15 (de^{(6i dx+6i c)} - 3 de^{(4i dx+4i c)} + 3 de^{(2i dx+2i c)} - d) \sqrt{-\frac{(-i A^2 - 2 AB + i B^2) a^2}{d^2}} \log \left(\frac{2 \left((A - i B) a e^{(2i dx+2i c)} + (de^{(2i dx+2i c)} - d) \right)}{\dots} \right)$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm
m="fricas")`

output

```
-1/30*(15*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*log(2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) + (d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a) - 15*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*log(2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) - (d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a) + 4*((-23*I*A - 20*B)*a*e^(6*I*d*x + 6*I*c) + (I*A + 10*B)*a*e^(4*I*d*x + 4*I*c) + (11*I*A + 20*B)*a*e^(2*I*d*x + 2*I*c) + (-13*I*A - 10*B)*a)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))/(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)
```

3.118.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= ia \left(\int \frac{A}{\tan^{\frac{5}{2}}(c + dx)} dx + \int \frac{B}{\tan^{\frac{3}{2}}(c + dx)} dx + \int \left(-\frac{iA}{\tan^{\frac{7}{2}}(c + dx)} \right) dx + \int \left(-\frac{iB}{\tan^{\frac{5}{2}}(c + dx)} \right) dx \right)$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)**(7/2),x)`

output `I*a*(Integral(A/tan(c + d*x)**(5/2), x) + Integral(B/tan(c + d*x)**(3/2), x) + Integral(-I*A/tan(c + d*x)**(7/2), x) + Integral(-I*B/tan(c + d*x)**(5/2), x))`

3.118.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 187 vs. $2(81) = 162$.

Time = 0.32 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.82

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx =$$

$$\frac{15 \left(2\sqrt{2}((i-1)A + (i+1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(dx+c)}\right)\right) + 2\sqrt{2}((i-1)A + (i+1)B) \right)}{d}$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="maxima")`

output `-1/60*(15*(2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a - 8*(15*(A - I*B)*a*tan(d*x + c)^2 + 5*(-I*A - B)*a*tan(d*x + c) - 3*A*a)/tan(d*x + c)^(5/2))/d`

3.118.8 Giac [A] (verification not implemented)

Time = 0.80 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.90

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{(i-1)\sqrt{2}(Aa - iBa) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{d}$$

$$+ \frac{2(15Aa \tan(dx+c)^2 - 15iBa \tan(dx+c)^2 - 5iAa \tan(dx+c) - 5Ba \tan(dx+c) - 3Aa)}{15d \tan(dx+c)^{\frac{5}{2}}}$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="giac")`

output $-(I - 1)*\text{sqrt}(2)*(A*a - I*B*a)*\text{arctan}(-(1/2*I - 1/2)*\text{sqrt}(2)*\text{sqrt}(\tan(d*x + c)))/d + 2/15*(15*A*a*\tan(d*x + c)^2 - 15*I*B*a*\tan(d*x + c)^2 - 5*I*A*a*\tan(d*x + c) - 5*B*a*\tan(d*x + c) - 3*A*a)/(d*\tan(d*x + c)^{(5/2)})$

3.118.9 Mupad [B] (verification not implemented)

Time = 9.97 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.19

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= -\frac{\frac{2Ba}{3d} + \frac{Ba \tan(c+dx)2i}{d}}{\tan(c+dx)^{3/2}} + \frac{\sqrt{2} Ba \operatorname{atan}\left(\sqrt{2} \sqrt{\tan(c+dx)}\left(\frac{1}{2} - \frac{1}{2}i\right)\right) (-1 - i)}{d}$$

$$-\frac{2Aa\left(15(-1)^{1/4}\tan(c+dx)^{5/2}\operatorname{atanh}\left((-1)^{1/4}\sqrt{\tan(c+dx)}\right) - 15\tan(c+dx)^2 + 3 + \tan(c+dx)\right)}{15d\tan(c+dx)^{5/2}}$$

input $\text{int}(((A + B*\tan(c + d*x))*(a + a*\tan(c + d*x)*1i))/\tan(c + d*x)^{(7/2}),x)$

output $-((2*B*a)/(3*d) + (B*a*\tan(c + d*x)*2i)/d)/\tan(c + d*x)^{(3/2)} - (2^{(1/2)}*B*a*\operatorname{atan}(2^{(1/2)}*\tan(c + d*x)^{(1/2)}*(1/2 - 1i/2))*(1 + 1i))/d - (2*A*a*(\tan(c + d*x)*5i - 15*\tan(c + d*x)^2 + 15*(-1)^{(1/4)}*\tan(c + d*x)^{(5/2)}*\operatorname{atanh}((-1)^{(1/4)}*\tan(c + d*x)^{(1/2)} + 3))/(15*d*\tan(c + d*x)^{(5/2)})$

3.119 $\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.119.1 Optimal result	1250
3.119.2 Mathematica [A] (verified)	1251
3.119.3 Rubi [A] (verified)	1251
3.119.4 Maple [A] (verified)	1255
3.119.5 Fricas [B] (verification not implemented)	1256
3.119.6 Sympy [F]	1257
3.119.7 Maxima [A] (verification not implemented)	1257
3.119.8 Giac [A] (verification not implemented)	1258
3.119.9 Mupad [B] (verification not implemented)	1259

3.119.1 Optimal result

Integrand size = 36, antiderivative size = 183

$$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= -\frac{4\sqrt[4]{-1}a^2(iA+B) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{d} - \frac{4a^2(iA+B)\sqrt{\tan(c+dx)}}{d}$$

$$+ \frac{4a^2(A-iB) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{4a^2(iA+B) \tan^{\frac{5}{2}}(c+dx)}{5d}$$

$$- \frac{2a^2(9A-11iB) \tan^{\frac{7}{2}}(c+dx)}{63d} + \frac{2iB \tan^{\frac{7}{2}}(c+dx)(a^2+ia^2 \tan(c+dx))}{9d}$$

output

```
-4*(-1)^(1/4)*a^2*(I*A+B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d-4*a^2*(I*A+B)*tan(d*x+c)^(1/2)/d+4/3*a^2*(A-I*B)*tan(d*x+c)^(3/2)/d+4/5*a^2*(I*A+B)*tan(d*x+c)^(5/2)/d-2/63*a^2*(9*A-11*I*B)*tan(d*x+c)^(7/2)/d+2/9*I*B*tan(d*x+c)^(7/2)*(a^2+I*a^2*tan(d*x+c))/d
```

3.119.2 Mathematica [A] (verified)

Time = 4.68 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.71

$$\int \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{2a^2 \left((315 + 315i)\sqrt{2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{\tan(c+dx)}}{\sqrt{2}}\right) + \sqrt{\tan(c + dx)}(-630i(A - iB) + 210(A - iB)\tan(c + dx)) \right)}{315d}$$

input `Integrate[Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(2*a^2*((315 + 315*I)*Sqrt[2]*(A - I*B)*ArcTanh[((1 + I)*Sqrt[Tan[c + d*x]])/Sqrt[2]] + Sqrt[Tan[c + d*x]]*((-630*I)*(A - I*B) + 210*(A - I*B)*Tan[c + d*x] + 126*(I*A + B)*Tan[c + d*x]^2 - 45*(A - (2*I)*B)*Tan[c + d*x]^3 - 35*B*Tan[c + d*x]^4))/ (315*d)`

3.119.3 Rubi [A] (verified)

Time = 1.04 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.02, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3042, 4077, 27, 3042, 4075, 3042, 4011, 3042, 4011, 3042, 4011, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^{5/2}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4077}$$

$$\frac{2}{9} \int \frac{1}{2} \tan^{\frac{5}{2}}(c + dx)(i \tan(c + dx)a + a)(a(9A - 7iB) + a(9iA + 11B) \tan(c + dx)) dx + \frac{2iB \tan^{\frac{7}{2}}(c + dx)(a^2 + ia^2 \tan(c + dx))}{9d}$$

$$\downarrow \text{27}$$

3.119. $\int \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\frac{1}{9} \int \tan^{\frac{5}{2}}(c+dx) (i \tan(c+dx)a + a)(a(9A - 7iB) + a(9iA + 11B) \tan(c+dx)) dx + \frac{2iB \tan^{\frac{7}{2}}(c+dx) (a^2 + ia^2 \tan(c+dx))}{9d}$$

↓ 3042

$$\frac{1}{9} \int \tan(c+dx)^{5/2} (i \tan(c+dx)a + a)(a(9A - 7iB) + a(9iA + 11B) \tan(c+dx)) dx + \frac{2iB \tan^{\frac{7}{2}}(c+dx) (a^2 + ia^2 \tan(c+dx))}{9d}$$

↓ 4075

$$\frac{1}{9} \left(\int \tan^{\frac{5}{2}}(c+dx) (18(A - iB)a^2 + 18(iA + B) \tan(c+dx)a^2) dx - \frac{2a^2(9A - 11iB) \tan^{\frac{7}{2}}(c+dx)}{7d} \right) + \frac{2iB \tan^{\frac{7}{2}}(c+dx) (a^2 + ia^2 \tan(c+dx))}{9d}$$

↓ 3042

$$\frac{1}{9} \left(\int \tan(c+dx)^{5/2} (18(A - iB)a^2 + 18(iA + B) \tan(c+dx)a^2) dx - \frac{2a^2(9A - 11iB) \tan^{\frac{7}{2}}(c+dx)}{7d} \right) + \frac{2iB \tan^{\frac{7}{2}}(c+dx) (a^2 + ia^2 \tan(c+dx))}{9d}$$

↓ 4011

$$\frac{1}{9} \left(\int \tan^{\frac{3}{2}}(c+dx) (18a^2(A - iB) \tan(c+dx) - 18a^2(iA + B)) dx - \frac{2a^2(9A - 11iB) \tan^{\frac{7}{2}}(c+dx)}{7d} + \frac{36a^2(B + iA)}{7d} \right) + \frac{2iB \tan^{\frac{7}{2}}(c+dx) (a^2 + ia^2 \tan(c+dx))}{9d}$$

↓ 3042

$$\frac{1}{9} \left(\int \tan(c+dx)^{3/2} (18a^2(A - iB) \tan(c+dx) - 18a^2(iA + B)) dx - \frac{2a^2(9A - 11iB) \tan^{\frac{7}{2}}(c+dx)}{7d} + \frac{36a^2(B + iA)}{7d} \right) + \frac{2iB \tan^{\frac{7}{2}}(c+dx) (a^2 + ia^2 \tan(c+dx))}{9d}$$

↓ 4011

3.119. $\int \tan^{\frac{5}{2}}(c+dx) (a + ia \tan(c+dx))^2 (A + B \tan(c+dx)) dx$

$$\frac{1}{9} \left(\int \sqrt{\tan(c+dx)} (-18(A-iB)a^2 - 18(iA+B)\tan(c+dx)a^2) dx - \frac{2a^2(9A-11iB)\tan^{\frac{7}{2}}(c+dx)}{7d} + \frac{36a^2(B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d} \right. \\ \left. \frac{2iB\tan^{\frac{7}{2}}(c+dx)(a^2+ia^2\tan(c+dx))}{9d} \right) \\ \downarrow 3042$$

$$\frac{1}{9} \left(\int \sqrt{\tan(c+dx)} (-18(A-iB)a^2 - 18(iA+B)\tan(c+dx)a^2) dx - \frac{2a^2(9A-11iB)\tan^{\frac{7}{2}}(c+dx)}{7d} + \frac{36a^2(B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d} \right. \\ \left. \frac{2iB\tan^{\frac{7}{2}}(c+dx)(a^2+ia^2\tan(c+dx))}{9d} \right) \\ \downarrow 4011$$

$$\frac{1}{9} \left(\int \frac{18a^2(iA+B) - 18a^2(A-iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(9A-11iB)\tan^{\frac{7}{2}}(c+dx)}{7d} + \frac{36a^2(B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d} \right. \\ \left. \frac{2iB\tan^{\frac{7}{2}}(c+dx)(a^2+ia^2\tan(c+dx))}{9d} \right) \\ \downarrow 3042$$

$$\frac{1}{9} \left(\int \frac{18a^2(iA+B) - 18a^2(A-iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(9A-11iB)\tan^{\frac{7}{2}}(c+dx)}{7d} + \frac{36a^2(B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d} \right. \\ \left. \frac{2iB\tan^{\frac{7}{2}}(c+dx)(a^2+ia^2\tan(c+dx))}{9d} \right) \\ \downarrow 4016$$

$$\frac{1}{9} \left(\frac{648a^4(B+iA)^2 \int \frac{1}{18(iA+B)a^2+18(A-iB)\tan(c+dx)a^2} d\sqrt{\tan(c+dx)}}{d} - \frac{2a^2(9A-11iB)\tan^{\frac{7}{2}}(c+dx)}{7d} + \frac{36a^2(B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d} \right. \\ \left. \frac{2iB\tan^{\frac{7}{2}}(c+dx)(a^2+ia^2\tan(c+dx))}{9d} \right) \\ \downarrow 218$$

$$\frac{1}{9} \left(-\frac{36\sqrt[4]{-1}a^2(B+iA)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{d} - \frac{2a^2(9A-11iB)\tan^{\frac{7}{2}}(c+dx)}{7d} + \frac{36a^2(B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d} \right. \\ \left. \frac{2iB\tan^{\frac{7}{2}}(c+dx)(a^2+ia^2\tan(c+dx))}{9d} \right)$$

input `Int[Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `((2*I/9)*B*Tan[c + d*x]^(7/2)*(a^2 + I*a^2*Tan[c + d*x])/d + ((-36*(-1)^(1/4)*a^2*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]])/d - (36*a^2*(I*A + B)*Sqrt[Tan[c + d*x])/d + (12*a^2*(A - I*B)*Tan[c + d*x]^(3/2))/d + (36*a^2*(I*A + B)*Tan[c + d*x]^(5/2))/(5*d) - (2*a^2*(9*A - (11*I)*B)*Tan[c + d*x]^(7/2))/(7*d))/9`

3.119.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4016 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

```
rule 4077 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan
[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c -
a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] &&
GtQ[m, 1] && !LtQ[n, -1]
```

3.119.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 298, normalized size of antiderivative = 1.63

method	result
derivativedivides	$a^2 \left(-\frac{2B \left(\tan \frac{9}{2}(dx+c) \right)}{9} + \frac{4iB \left(\tan \frac{7}{2}(dx+c) \right)}{7} - \frac{2A \left(\tan \frac{7}{2}(dx+c) \right)}{7} + \frac{4iA \left(\tan \frac{5}{2}(dx+c) \right)}{5} + \frac{4B \left(\tan \frac{5}{2}(dx+c) \right)}{5} - \frac{4iB \left(\tan \frac{3}{2}(dx+c) \right)}{3} \right)$
default	$a^2 \left(-\frac{2B \left(\tan \frac{9}{2}(dx+c) \right)}{9} + \frac{4iB \left(\tan \frac{7}{2}(dx+c) \right)}{7} - \frac{2A \left(\tan \frac{7}{2}(dx+c) \right)}{7} + \frac{4iA \left(\tan \frac{5}{2}(dx+c) \right)}{5} + \frac{4B \left(\tan \frac{5}{2}(dx+c) \right)}{5} - \frac{4iB \left(\tan \frac{3}{2}(dx+c) \right)}{3} \right)$
parts	$(2iAa^2 + Ba^2) \left(\frac{2 \left(\tan \frac{5}{2}(dx+c) \right)}{5} - 2(\sqrt{\tan(dx+c)}) + \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})} \right)}{4} \right)$

```
input int(tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/d*a^2*(-2/9*B*tan(d*x+c)^(9/2)+4/7*I*B*tan(d*x+c)^(7/2)-2/7*A*tan(d*x+c)^(7/2)+4/5*I*A*tan(d*x+c)^(5/2)+4/5*B*tan(d*x+c)^(5/2)-4/3*I*B*tan(d*x+c)^(3/2)+4/3*A*tan(d*x+c)^(3/2)-4*I*A*tan(d*x+c)^(1/2)-4*B*tan(d*x+c)^(1/2)+1/4*(2*B+2*I*A)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(-2*A+2*I*B)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))
```

$$3.119. \int \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

3.119.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 555 vs. $2(145) = 290$.

Time = 0.34 (sec) , antiderivative size = 555, normalized size of antiderivative = 3.03

$$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{315 \sqrt{-\frac{(-iA^2-2AB+iB^2)a^4}{d^2}} (de^{(8i dx+8i c)} + 4de^{(6i dx+6i c)} + 6de^{(4i dx+4i c)} + 4de^{(2i dx+2i c)} + d) \log \left(-\frac{2 \left((A-iB) \right)}{\dots} \right)}{\dots}$$

```
input integrate(tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorith
m="fricas")
```

```
output 1/315*(315*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2)*(d*e^(8*I*d*x + 8*I*c)
+ 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*x + 2*I
*c) + d)*log(-2*((A - I*B)*a^2*e^(2*I*d*x + 2*I*c) + sqrt(-(-I*A^2 - 2*A*B
+ I*B^2)*a^4/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt((-I*e^(2*I*d*x + 2*I*c)
) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^2))
- 315*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2)*(d*e^(8*I*d*x + 8*I*c) + 4*d
*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*x + 2*I*c) +
d)*log(-2*((A - I*B)*a^2*e^(2*I*d*x + 2*I*c) - sqrt(-(-I*A^2 - 2*A*B + I*
B^2)*a^4/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I
))/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^2)) - 2*(
(1011*I*A + 1091*B)*a^2*e^(8*I*d*x + 8*I*c) + 10*(285*I*A + 262*B)*a^2*e^(
6*I*d*x + 6*I*c) + 42*(84*I*A + 89*B)*a^2*e^(4*I*d*x + 4*I*c) + 10*(219*I*
A + 214*B)*a^2*e^(2*I*d*x + 2*I*c) + (501*I*A + 491*B)*a^2)*sqrt((-I*e^(2*
I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(8*I*d*x + 8*I*c) + 4
*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*x + 2*I*c)
+ d)
```

3.119.6 Sympy [F]

$$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= -a^2 \left(\int (-A \tan^{\frac{5}{2}}(c+dx)) dx + \int A \tan^{\frac{9}{2}}(c+dx) dx + \int (-B \tan^{\frac{7}{2}}(c+dx)) dx \right. \\ \left. + \int B \tan^{\frac{11}{2}}(c+dx) dx + \int (-2iA \tan^{\frac{7}{2}}(c+dx)) dx + \int (-2iB \tan^{\frac{9}{2}}(c+dx)) dx \right)$$

input `integrate(tan(d*x+c)**(5/2)*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `-a**2*(Integral(-A*tan(c + d*x)**(5/2), x) + Integral(A*tan(c + d*x)**(9/2), x) + Integral(-B*tan(c + d*x)**(7/2), x) + Integral(B*tan(c + d*x)**(11/2), x) + Integral(-2*I*A*tan(c + d*x)**(7/2), x) + Integral(-2*I*B*tan(c + d*x)**(9/2), x))`

3.119.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.26

$$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx =$$

$$\frac{140 B a^2 \tan(dx+c)^{\frac{9}{2}} + 180 (A - 2i B) a^2 \tan(dx+c)^{\frac{7}{2}} + 504 (-i A - B) a^2 \tan(dx+c)^{\frac{5}{2}} - 840 (A -$$

input `integrate(tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/630*(140*B*a^2*tan(d*x + c)^(9/2) + 180*(A - 2*I*B)*a^2*tan(d*x + c)^(7/2) + 504*(-I*A - B)*a^2*tan(d*x + c)^(5/2) - 840*(A - I*B)*a^2*tan(d*x + c)^(3/2) + 2520*(I*A + B)*a^2*sqrt(tan(d*x + c)) - 315*(2*sqrt(2))*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a^2/d`

3.119. $\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.119.8 Giac [A] (verification not implemented)

Time = 1.14 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.06

$$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{(2i-2) \sqrt{2}(Aa^2-iBa^2) \arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx+c)}\right)}{d}$$

$$- \frac{2\left(35Ba^2d^8 \tan(dx+c)^{\frac{9}{2}}+45Aa^2d^8 \tan(dx+c)^{\frac{7}{2}}-90iBa^2d^8 \tan(dx+c)^{\frac{7}{2}}-126iAa^2d^8 \tan(dx+c)^{\frac{5}{2}}-126B^2a^2d^8 \tan(dx+c)^{\frac{5}{2}}-210A^2a^2d^8 \tan(dx+c)^{\frac{3}{2}}+210iB^2a^2d^8 \tan(dx+c)^{\frac{3}{2}}+630iA^2a^2d^8 \sqrt{\tan(dx+c)}+630B^2a^2d^8 \sqrt{\tan(dx+c)}\right)}{d^9}$$

input `integrate(tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `(2*I - 2)*sqrt(2)*(A*a^2 - I*B*a^2)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d - 2/315*(35*B*a^2*d^8*tan(d*x + c)^(9/2) + 45*A*a^2*d^8*tan(d*x + c)^(7/2) - 90*I*B*a^2*d^8*tan(d*x + c)^(7/2) - 126*I*A*a^2*d^8*tan(d*x + c)^(5/2) - 126*B*a^2*d^8*tan(d*x + c)^(5/2) - 210*A*a^2*d^8*tan(d*x + c)^(3/2) + 210*I*B*a^2*d^8*tan(d*x + c)^(3/2) + 630*I*A*a^2*d^8*sqrt(tan(d*x + c)) + 630*B*a^2*d^8*sqrt(tan(d*x + c)))/d^9`

3.119.9 Mupad [B] (verification not implemented)

Time = 13.16 (sec) , antiderivative size = 326, normalized size of antiderivative = 1.78

$$\begin{aligned}
& \int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx \\
&= \frac{4 A a^2 \tan(c+dx)^{3/2}}{3 d} - \frac{A a^2 \sqrt{\tan(c+dx)} 4i}{d} + \frac{A a^2 \tan(c+dx)^{5/2} 4i}{5 d} \\
&\quad - \frac{2 A a^2 \tan(c+dx)^{7/2}}{7 d} - \frac{4 B a^2 \sqrt{\tan(c+dx)}}{d} - \frac{B a^2 \tan(c+dx)^{3/2} 4i}{3 d} \\
&\quad + \frac{4 B a^2 \tan(c+dx)^{5/2}}{5 d} + \frac{B a^2 \tan(c+dx)^{7/2} 4i}{7 d} - \frac{2 B a^2 \tan(c+dx)^{9/2}}{9 d} \\
&\quad + \frac{\sqrt{2} A a^2 \ln\left(-4 A a^2 d + \sqrt{2} A a^2 d \sqrt{\tan(c+dx)}(-2-2i)\right)(1+i)}{d} \\
&\quad - \frac{\sqrt{4i} A a^2 \ln\left(-4 A a^2 d + 2 \sqrt{4i} A a^2 d \sqrt{\tan(c+dx)}\right)}{d} \\
&\quad + \frac{\sqrt{2} B a^2 \ln\left(B a^2 d 4i + \sqrt{2} B a^2 d \sqrt{\tan(c+dx)}(-2+2i)\right)(1-i)}{d} \\
&\quad - \frac{\sqrt{-4i} B a^2 \ln\left(B a^2 d 4i + 2 \sqrt{-4i} B a^2 d \sqrt{\tan(c+dx)}\right)}{d}
\end{aligned}$$

```
input int(tan(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2,x)
```

```
output (4*A*a^2*tan(c + d*x)^(3/2))/(3*d) - (A*a^2*tan(c + d*x)^(1/2)*4i)/d + (A*
a^2*tan(c + d*x)^(5/2)*4i)/(5*d) - (2*A*a^2*tan(c + d*x)^(7/2))/(7*d) - (4
*B*a^2*tan(c + d*x)^(1/2))/d - (B*a^2*tan(c + d*x)^(3/2)*4i)/(3*d) + (4*B*
a^2*tan(c + d*x)^(5/2))/(5*d) + (B*a^2*tan(c + d*x)^(7/2)*4i)/(7*d) - (2*B
*a^2*tan(c + d*x)^(9/2))/(9*d) + (2^(1/2)*A*a^2*log(- 4*A*a^2*d - 2^(1/2)*
A*a^2*d*tan(c + d*x)^(1/2)*(2 + 2i))*(1 + 1i))/d - (4i^(1/2)*A*a^2*log(2*4
i^(1/2)*A*a^2*d*tan(c + d*x)^(1/2) - 4*A*a^2*d))/d + (2^(1/2)*B*a^2*log(B*
a^2*d*4i - 2^(1/2)*B*a^2*d*tan(c + d*x)^(1/2)*(2 - 2i))*(1 - 1i))/d - ((-4
i)^(1/2)*B*a^2*log(B*a^2*d*4i + 2*(-4i)^(1/2)*B*a^2*d*tan(c + d*x)^(1/2))
/d
```

3.120 $\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.120.1 Optimal result	1260
3.120.2 Mathematica [A] (verified)	1261
3.120.3 Rubi [A] (verified)	1261
3.120.4 Maple [B] (verified)	1265
3.120.5 Fricas [B] (verification not implemented)	1266
3.120.6 Sympy [F]	1266
3.120.7 Maxima [A] (verification not implemented)	1267
3.120.8 Giac [A] (verification not implemented)	1267
3.120.9 Mupad [B] (verification not implemented)	1268

3.120.1 Optimal result

Integrand size = 36, antiderivative size = 156

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{4\sqrt{-1}a^2(A-iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{d} + \frac{4a^2(A-iB)\sqrt{\tan(c+dx)}}{d} + \frac{4a^2(iA+B)\tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2a^2(7A-9iB)\tan^{\frac{5}{2}}(c+dx)}{35d} + \frac{2iB \tan^{\frac{5}{2}}(c+dx)(a^2+ia^2 \tan(c+dx))}{7d}$$

output

```
4*(-1)^(1/4)*a^2*(A-I*B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d+4*a^2*(A-I*B)*tan(d*x+c)^(1/2)/d+4/3*a^2*(I*A+B)*tan(d*x+c)^(3/2)/d-2/35*a^2*(7*A-9*I*B)*tan(d*x+c)^(5/2)/d+2/7*I*B*tan(d*x+c)^(5/2)*(a^2+I*a^2*tan(d*x+c))/d
```

3.120.2 Mathematica [A] (verified)

Time = 2.13 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.71

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{2a^2 \left((105+105i)\sqrt{2}(iA+B) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{\tan(c+dx)}}{\sqrt{2}}\right) + \sqrt{\tan(c+dx)}(210(A-iB) + 70(iA+B) \tan(c+dx)) \right)}{105d}$$

input `Integrate[Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(2*a^2*((105 + 105*I)*Sqrt[2]*(I*A + B)*ArcTanh[((1 + I)*Sqrt[Tan[c + d*x]])/Sqrt[2]] + Sqrt[Tan[c + d*x]]*(210*(A - I*B) + 70*(I*A + B)*Tan[c + d*x] - 21*(A - (2*I)*B)*Tan[c + d*x]^2 - 15*B*Tan[c + d*x]^3))/(105*d)`

3.120.3 Rubi [A] (verified)

Time = 0.86 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.03, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 4077, 27, 3042, 4075, 3042, 4011, 3042, 4011, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c+dx)^{3/2}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4077}$$

$$\frac{2}{7} \int \frac{1}{2} \tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a)(a(7A-5iB)+a(7iA+9B) \tan(c+dx)) dx +$$

$$\frac{2iB \tan^{\frac{5}{2}}(c+dx)(a^2+ia^2 \tan(c+dx))}{7d}$$

$$\downarrow \text{27}$$

$$\frac{1}{7} \int \tan^{\frac{3}{2}}(c+dx) (i \tan(c+dx)a + a)(a(7A - 5iB) + a(7iA + 9B) \tan(c+dx)) dx + \frac{2iB \tan^{\frac{5}{2}}(c+dx) (a^2 + ia^2 \tan(c+dx))}{7d}$$

↓ 3042

$$\frac{1}{7} \int \tan(c+dx)^{3/2} (i \tan(c+dx)a + a)(a(7A - 5iB) + a(7iA + 9B) \tan(c+dx)) dx + \frac{2iB \tan^{\frac{5}{2}}(c+dx) (a^2 + ia^2 \tan(c+dx))}{7d}$$

↓ 4075

$$\frac{1}{7} \left(\int \tan^{\frac{3}{2}}(c+dx) (14(A - iB)a^2 + 14(iA + B) \tan(c+dx)a^2) dx - \frac{2a^2(7A - 9iB) \tan^{\frac{5}{2}}(c+dx)}{5d} \right) + \frac{2iB \tan^{\frac{5}{2}}(c+dx) (a^2 + ia^2 \tan(c+dx))}{7d}$$

↓ 3042

$$\frac{1}{7} \left(\int \tan(c+dx)^{3/2} (14(A - iB)a^2 + 14(iA + B) \tan(c+dx)a^2) dx - \frac{2a^2(7A - 9iB) \tan^{\frac{5}{2}}(c+dx)}{5d} \right) + \frac{2iB \tan^{\frac{5}{2}}(c+dx) (a^2 + ia^2 \tan(c+dx))}{7d}$$

↓ 4011

$$\frac{1}{7} \left(\int \sqrt{\tan(c+dx)} (14a^2(A - iB) \tan(c+dx) - 14a^2(iA + B)) dx - \frac{2a^2(7A - 9iB) \tan^{\frac{5}{2}}(c+dx)}{5d} + \frac{28a^2(B + iA)}{5d} \right) + \frac{2iB \tan^{\frac{5}{2}}(c+dx) (a^2 + ia^2 \tan(c+dx))}{7d}$$

↓ 3042

$$\frac{1}{7} \left(\int \sqrt{\tan(c+dx)} (14a^2(A - iB) \tan(c+dx) - 14a^2(iA + B)) dx - \frac{2a^2(7A - 9iB) \tan^{\frac{5}{2}}(c+dx)}{5d} + \frac{28a^2(B + iA)}{5d} \right) + \frac{2iB \tan^{\frac{5}{2}}(c+dx) (a^2 + ia^2 \tan(c+dx))}{7d}$$

↓ 4011

3.120. $\int \tan^{\frac{3}{2}}(c+dx) (a + ia \tan(c+dx))^2 (A + B \tan(c+dx)) dx$

$$\frac{1}{7} \left(\int \frac{-14(A - iB)a^2 - 14(iA + B) \tan(c + dx)a^2}{\sqrt{\tan(c + dx)}} dx - \frac{2a^2(7A - 9iB) \tan^{\frac{5}{2}}(c + dx)}{5d} + \frac{28a^2(B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d} \right. \\ \left. \frac{2iB \tan^{\frac{5}{2}}(c + dx) (a^2 + ia^2 \tan(c + dx))}{7d} \right)$$

↓ 3042

$$\frac{1}{7} \left(\int \frac{-14(A - iB)a^2 - 14(iA + B) \tan(c + dx)a^2}{\sqrt{\tan(c + dx)}} dx - \frac{2a^2(7A - 9iB) \tan^{\frac{5}{2}}(c + dx)}{5d} + \frac{28a^2(B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d} \right. \\ \left. \frac{2iB \tan^{\frac{5}{2}}(c + dx) (a^2 + ia^2 \tan(c + dx))}{7d} \right)$$

↓ 4016

$$\frac{1}{7} \left(\frac{392a^4(A - iB)^2 \int \frac{1}{14a^2(iA + B) \tan(c + dx) - 14a^2(A - iB)} d\sqrt{\tan(c + dx)}}{d} - \frac{2a^2(7A - 9iB) \tan^{\frac{5}{2}}(c + dx)}{5d} + \frac{28a^2(B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d} \right. \\ \left. \frac{2iB \tan^{\frac{5}{2}}(c + dx) (a^2 + ia^2 \tan(c + dx))}{7d} \right)$$

↓ 218

$$\frac{1}{7} \left(\frac{28\sqrt[4]{-1}a^2(A - iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right)}{d} - \frac{2a^2(7A - 9iB) \tan^{\frac{5}{2}}(c + dx)}{5d} + \frac{28a^2(B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d} \right. \\ \left. \frac{2iB \tan^{\frac{5}{2}}(c + dx) (a^2 + ia^2 \tan(c + dx))}{7d} \right)$$

input `Int[Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `((2*I)/7)*B*Tan[c + d*x]^(5/2)*(a^2 + I*a^2*Tan[c + d*x])/d + ((28*(-1)^(1/4)*a^2*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]])/d + (28*a^2*(A - I*B)*Sqrt[Tan[c + d*x]])/d + (28*a^2*(I*A + B)*Tan[c + d*x]^(3/2))/(3*d) - (2*a^2*(7*A - (9*I)*B)*Tan[c + d*x]^(5/2))/(5*d))/7`

3.120.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4011 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`
- rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`
- rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`
- rule 4077 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.120.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 274 vs. $2(130) = 260$.

Time = 0.04 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.76

method	result
derivativedivides	$a^2 \left(-\frac{2B \left(\tan^{\frac{7}{2}}(dx+c) \right)}{7} + \frac{4iB \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} - \frac{2A \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} + \frac{4iA \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + \frac{4B \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 4iB \left(\sqrt{\tan(dx+c)} \right) \right)$
default	$a^2 \left(-\frac{2B \left(\tan^{\frac{7}{2}}(dx+c) \right)}{7} + \frac{4iB \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} - \frac{2A \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} + \frac{4iA \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + \frac{4B \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 4iB \left(\sqrt{\tan(dx+c)} \right) \right)$
parts	$\frac{(2iAa^2 + Ba^2) \left(\frac{2 \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - \frac{\sqrt{2} \left(\ln \left(\frac{1 - \sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)}{1 + \sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)} \right) + 2 \arctan \left(\frac{1 + \sqrt{2} \left(\sqrt{\tan(dx+c)} \right)}{1} \right) + 2 \arctan \left(\frac{-1 + \sqrt{2} \left(\sqrt{\tan(dx+c)} \right)}{1} \right)}{4} \right)}{d}$

input `int(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*a^2*(-2/7*B*tan(d*x+c)^(7/2)+4/5*I*B*tan(d*x+c)^(5/2)-2/5*A*tan(d*x+c)^(5/2)+4/3*I*A*tan(d*x+c)^(3/2)+4/3*B*tan(d*x+c)^(3/2)-4*I*B*tan(d*x+c)^(1/2)+4*A*tan(d*x+c)^(1/2)+1/4*(-2*A+2*I*B)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(-2*B-2*I*A)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))`

3.120.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 501 vs. $2(124) = 248$.

Time = 0.29 (sec) , antiderivative size = 501, normalized size of antiderivative = 3.21

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx =$$

$$105 \sqrt{-\frac{(iA^2+2AB-iB^2)a^4}{d^2}} (de^{(6i dx+6i c)} + 3de^{(4i dx+4i c)} + 3de^{(2i dx+2i c)} + d) \log \left(-\frac{2 \left((A-iB)a^2 e^{(2i dx+2i c)} + \sqrt{\dots} \right)}{\dots} \right)$$

input `integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-1/105*(105*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*log(-2*((A - I*B)*a^2*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^2)) - 105*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*log(-2*((A - I*B)*a^2*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^2)) - 2*((301*A - 337*I*B)*a^2*e^(6*I*d*x + 6*I*c) + (679*A - 613*I*B)*a^2*e^(4*I*d*x + 4*I*c) + (539*A - 563*I*B)*a^2*e^(2*I*d*x + 2*I*c) + (161*A - 167*I*B)*a^2)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)`

3.120.6 Sympy [F]

$$\begin{aligned} & \int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx \\ &= -a^2 \left(\int (-A \tan^{\frac{3}{2}}(c+dx)) dx + \int A \tan^{\frac{7}{2}}(c+dx) dx + \int (-B \tan^{\frac{5}{2}}(c+dx)) dx \right. \\ & \quad \left. + \int B \tan^{\frac{9}{2}}(c+dx) dx + \int (-2iA \tan^{\frac{5}{2}}(c+dx)) dx + \int (-2iB \tan^{\frac{7}{2}}(c+dx)) dx \right) \end{aligned}$$

3.120. $\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

input `integrate(tan(d*x+c)**(3/2)*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `-a**2*(Integral(-A*tan(c + d*x)**(3/2), x) + Integral(A*tan(c + d*x)**(7/2), x) + Integral(-B*tan(c + d*x)**(5/2), x) + Integral(B*tan(c + d*x)**(9/2), x) + Integral(-2*I*A*tan(c + d*x)**(5/2), x) + Integral(-2*I*B*tan(c + d*x)**(7/2), x))`

3.120.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.36

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$\frac{60 Ba^2 \tan(dx + c)^{\frac{7}{2}} + 84(A - 2iB)a^2 \tan(dx + c)^{\frac{5}{2}} + 280(-iA - B)a^2 \tan(dx + c)^{\frac{3}{2}} - 840(A - iB)a^2 \tan(dx + c)^{\frac{1}{2}}}{d}$$

input `integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/210*(60*B*a^2*tan(d*x + c)^(7/2) + 84*(A - 2*I*B)*a^2*tan(d*x + c)^(5/2) + 280*(-I*A - B)*a^2*tan(d*x + c)^(3/2) - 840*(A - I*B)*a^2*sqrt(tan(d*x + c)) - 105*(2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a^2)/d`

3.120.8 Giac [A] (verification not implemented)

Time = 0.96 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.03

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{(2i - 2) \sqrt{2}(iAa^2 + Ba^2) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{d}$$

$$\frac{2 \left(15 Ba^2 d^6 \tan(dx + c)^{\frac{7}{2}} + 21 Aa^2 d^6 \tan(dx + c)^{\frac{5}{2}} - 42i Ba^2 d^6 \tan(dx + c)^{\frac{3}{2}} - 70i Aa^2 d^6 \tan(dx + c)^{\frac{1}{2}}\right)}{d}$$

105 d⁷

3.120. $\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

input `integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output $(2I - 2)\sqrt{2}(IAa^2 + Ba^2)\arctan\left(\frac{-1/2I - 1/2}{\sqrt{2}}\sqrt{\tan(dx + c)}\right)/d - 2/105(15Ba^2d^6\tan(dx + c)^{7/2} + 21Aa^2d^6\tan(dx + c)^{5/2} - 42IBa^2d^6\tan(dx + c)^{5/2} - 70IAa^2d^6\tan(dx + c)^{3/2} - 70Ba^2d^6\tan(dx + c)^{3/2} - 210Aa^2d^6\sqrt{\tan(dx + c)} + 210IBa^2d^6\sqrt{\tan(dx + c)})/d^7$

3.120.9 Mupad [B] (verification not implemented)

Time = 10.81 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.87

$$\begin{aligned} & \int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\ &= \frac{4Aa^2\sqrt{\tan(c + dx)}}{d} + \frac{Aa^2\tan(c + dx)^{3/2}4i}{3d} - \frac{2Aa^2\tan(c + dx)^{5/2}}{5d} \\ & - \frac{Ba^2\sqrt{\tan(c + dx)}4i}{d} + \frac{4Ba^2\tan(c + dx)^{3/2}}{3d} \\ & + \frac{Ba^2\tan(c + dx)^{5/2}4i}{5d} - \frac{2Ba^2\tan(c + dx)^{7/2}}{7d} \\ & + \frac{\sqrt{2}Aa^2\ln\left(-Aa^2d4i + \sqrt{2}Aa^2d\sqrt{\tan(c + dx)}(-2 + 2i)\right)(1 - i)}{d} \\ & - \frac{\sqrt{-4i}Aa^2\ln\left(-Aa^2d4i + 2\sqrt{-4i}Aa^2d\sqrt{\tan(c + dx)}\right)}{d} \\ & + \frac{\sqrt{2}Ba^2\ln\left(-4Ba^2d + \sqrt{2}Ba^2d\sqrt{\tan(c + dx)}(-2 - 2i)\right)(1 + i)}{d} \\ & - \frac{\sqrt{4i}Ba^2\ln\left(-4Ba^2d + 2\sqrt{4i}Ba^2d\sqrt{\tan(c + dx)}\right)}{d} \end{aligned}$$

input `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2,x)`

output $(4Aa^2 \tan(c + dx)^{1/2})/d + (Aa^2 \tan(c + dx)^{3/2} 4i)/(3d) - (2Aa^2 \tan(c + dx)^{5/2})/(5d) - (Ba^2 \tan(c + dx)^{1/2} 4i)/d + (4Baa^2 \tan(c + dx)^{3/2})/(3d) + (Ba^2 \tan(c + dx)^{5/2} 4i)/(5d) - (2Baa^2 \tan(c + dx)^{7/2})/(7d) + (2^{1/2} Aa^2 \log(-Aa^2 d 4i - 2^{1/2} Aa^2 d \tan(c + dx)^{1/2} (2 - 2i))(1 - i))/d - ((-4i)^{1/2} Aa^2 \log(2(-4i)^{1/2} Aa^2 d \tan(c + dx)^{1/2} - Aa^2 d 4i))/d + (2^{1/2} Baa^2 \log(-4Baa^2 d - 2^{1/2} Baa^2 d \tan(c + dx)^{1/2} (2 + 2i))(1 + i))/d - (4i^{1/2} Baa^2 \log(2 4i^{1/2} Baa^2 d \tan(c + dx)^{1/2} - 4Baa^2 d))/d$

3.121 $\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.121.1 Optimal result	1270
3.121.2 Mathematica [A] (verified)	1270
3.121.3 Rubi [A] (verified)	1271
3.121.4 Maple [B] (verified)	1274
3.121.5 Fricas [B] (verification not implemented)	1275
3.121.6 Sympy [F]	1275
3.121.7 Maxima [A] (verification not implemented)	1276
3.121.8 Giac [A] (verification not implemented)	1276
3.121.9 Mupad [B] (verification not implemented)	1277

3.121.1 Optimal result

Integrand size = 36, antiderivative size = 129

$$\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{4\sqrt[4]{-1}a^2(iA + B) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right)}{d} + \frac{4a^2(iA + B)\sqrt{\tan(c + dx)}}{d}$$

$$- \frac{2a^2(5A - 7iB) \tan^{3/2}(c + dx)}{15d} + \frac{2iB \tan^{3/2}(c + dx)(a^2 + ia^2 \tan(c + dx))}{5d}$$

```
output 4*(-1)^(1/4)*a^2*(I*A+B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d+4*a^2*(I*A+B)*tan(d*x+c)^(1/2)/d-2/15*a^2*(5*A-7*I*B)*tan(d*x+c)^(3/2)/d+2/5*I*B*tan(d*x+c)^(3/2)*(a^2+I*a^2*tan(d*x+c))/d
```

3.121.2 Mathematica [A] (verified)

Time = 1.36 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.72

$$\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$\frac{2a^2\left((15 + 15i)\sqrt{2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{\tan(c+dx)}}{\sqrt{2}}\right) + \sqrt{\tan(c + dx)}(-30iA - 30B + 5(A - 2iB) \tan(c + dx))\right)}{15d}$$

input `Integrate[Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(-2*a^2*((15 + 15*I)*Sqrt[2]*(A - I*B)*ArcTanh[((1 + I)*Sqrt[Tan[c + d*x]])/Sqrt[2]] + Sqrt[Tan[c + d*x]]*((-30*I)*A - 30*B + 5*(A - (2*I)*B)*Tan[c + d*x] + 3*B*Tan[c + d*x]^2)))/(15*d)`

3.121.3 Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.04, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {3042, 4077, 27, 3042, 4075, 3042, 4011, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4077} \\
 & \frac{2}{5} \int \frac{1}{2} \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)(a(5A-3iB)+a(5iA+7B) \tan(c+dx)) dx + \\
 & \quad \frac{2iB \tan^{\frac{3}{2}}(c+dx)(a^2+ia^2 \tan(c+dx))}{5d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{5} \int \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)(a(5A-3iB)+a(5iA+7B) \tan(c+dx)) dx + \\
 & \quad \frac{2iB \tan^{\frac{3}{2}}(c+dx)(a^2+ia^2 \tan(c+dx))}{5d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{5} \int \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)(a(5A-3iB)+a(5iA+7B) \tan(c+dx)) dx + \\
 & \quad \frac{2iB \tan^{\frac{3}{2}}(c+dx)(a^2+ia^2 \tan(c+dx))}{5d} \\
 & \quad \downarrow \text{4075}
 \end{aligned}$$

3.121. $\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

$$\frac{1}{5} \left(\int \sqrt{\tan(c+dx)} (10(A-iB)a^2 + 10(iA+B)\tan(c+dx)a^2) dx - \frac{2a^2(5A-7iB)\tan^{\frac{3}{2}}(c+dx)}{3d} \right) + \frac{2iB\tan^{\frac{3}{2}}(c+dx)(a^2+ia^2\tan(c+dx))}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\int \sqrt{\tan(c+dx)} (10(A-iB)a^2 + 10(iA+B)\tan(c+dx)a^2) dx - \frac{2a^2(5A-7iB)\tan^{\frac{3}{2}}(c+dx)}{3d} \right) + \frac{2iB\tan^{\frac{3}{2}}(c+dx)(a^2+ia^2\tan(c+dx))}{5d}$$

↓ 4011

$$\frac{1}{5} \left(\int \frac{10a^2(A-iB)\tan(c+dx) - 10a^2(iA+B)}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(5A-7iB)\tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{20a^2(B+iA)\sqrt{\tan(c+dx)}}{d} \right) + \frac{2iB\tan^{\frac{3}{2}}(c+dx)(a^2+ia^2\tan(c+dx))}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\int \frac{10a^2(A-iB)\tan(c+dx) - 10a^2(iA+B)}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(5A-7iB)\tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{20a^2(B+iA)\sqrt{\tan(c+dx)}}{d} \right) + \frac{2iB\tan^{\frac{3}{2}}(c+dx)(a^2+ia^2\tan(c+dx))}{5d}$$

↓ 4016

$$\frac{1}{5} \left(\frac{200a^4(B+iA)^2 \int \frac{1}{-10(iA+B)a^2 - 10(A-iB)\tan(c+dx)a^2} d\sqrt{\tan(c+dx)}}{d} - \frac{2a^2(5A-7iB)\tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{20a^2(B+iA)\sqrt{\tan(c+dx)}}{d} \right) + \frac{2iB\tan^{\frac{3}{2}}(c+dx)(a^2+ia^2\tan(c+dx))}{5d}$$

↓ 218

$$\frac{1}{5} \left(\frac{20\sqrt[4]{-1}a^2(B+iA)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{d} - \frac{2a^2(5A-7iB)\tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{20a^2(B+iA)\sqrt{\tan(c+dx)}}{d} \right) + \frac{2iB\tan^{\frac{3}{2}}(c+dx)(a^2+ia^2\tan(c+dx))}{5d}$$

input `Int[Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `((2*I)/5)*B*Tan[c + d*x]^(3/2)*(a^2 + I*a^2*Tan[c + d*x])/d + ((20*(-1)^(1/4)*a^2*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]])/d + (20*a^2*(I*A + B)*Sqrt[Tan[c + d*x]])/d - (2*a^2*(5*A - (7*I)*B)*Tan[c + d*x]^(3/2))/(3*d)/5`

3.121.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`


```
rule 4077 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan
[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c -
a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] &&
GtQ[m, 1] && !LtQ[n, -1]
```

3.121.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 251 vs. 2(108) = 216.

Time = 0.03 (sec) , antiderivative size = 252, normalized size of antiderivative = 1.95

method	result
derivativedivides	$a^2 \left(-\frac{2B \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} + \frac{4iB \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - \frac{2A \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + 4iA \left(\sqrt{\tan(dx+c)} \right) + 4B \left(\sqrt{\tan(dx+c)} \right) + \frac{(-2iA-2B)\sqrt{2}}{\dots} \right)$
default	$a^2 \left(-\frac{2B \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} + \frac{4iB \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - \frac{2A \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + 4iA \left(\sqrt{\tan(dx+c)} \right) + 4B \left(\sqrt{\tan(dx+c)} \right) + \frac{(-2iA-2B)\sqrt{2}}{\dots} \right)$
parts	$\frac{(2iAa^2 + Ba^2) \left(2\sqrt{\tan(dx+c)} - \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))} \right) + 2 \arctan \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})}{1-\sqrt{2}(\sqrt{\tan(dx+c)})} \right) + 2 \arctan(-1+\sqrt{\tan(dx+c)}) \right)}{4} \right)}{d}$

```
input int(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETUR
NVERBOSE)
```

```
output 1/d*a^2*(-2/5*B*tan(d*x+c)^(5/2)+4/3*I*B*tan(d*x+c)^(3/2)-2/3*A*tan(d*x+c)
^(3/2)+4*I*A*tan(d*x+c)^(1/2)+4*B*tan(d*x+c)^(1/2)+1/4*(-2*B-2*I*A)*2^(1/2)
*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+
tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(
d*x+c)^(1/2)))+1/4*(2*A-2*I*B)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan
(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d
*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))
```

3.121. $\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.121.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 441 vs. $2(103) = 206$.

Time = 0.27 (sec) , antiderivative size = 441, normalized size of antiderivative = 3.42

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx =$$

$$15 \sqrt{-\frac{(-iA^2-2AB+iB^2)a^4}{d^2}} \left(de^{(4i dx+4i c)} + 2 de^{(2i dx+2i c)} + d \right) \log \left(-\frac{2 \left((A-iB)a^2 e^{(2i dx+2i c)} + \sqrt{-\frac{(-iA^2-2AB+iB^2)a^4}{d^2}} \right)}{(-i \dots)} \right)$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output

```
-1/15*(15*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2)*(d*e^(4*I*d*x + 4*I*c) +
2*d*e^(2*I*d*x + 2*I*c) + d)*log(-2*((A - I*B)*a^2*e^(2*I*d*x + 2*I*c) +
sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt((
-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*
c)/((-I*A - B)*a^2)) - 15*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2)*(d*e^(4*
I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log(-2*((A - I*B)*a^2*e^(2*I
*d*x + 2*I*c) - sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2)*(d*e^(2*I*d*x + 2*
I*c) + d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^
(-2*I*d*x - 2*I*c)/((-I*A - B)*a^2)) + 2*((-35*I*A - 43*B)*a^2*e^(4*I*d*x
+ 4*I*c) + 6*(-10*I*A - 9*B)*a^2*e^(2*I*d*x + 2*I*c) + (-25*I*A - 23*B)*a^
2)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(4*I
*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)
```

3.121.6 Sympy [F]

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= -a^2 \left(\int \left(-A \sqrt{\tan(c+dx)} \right) dx + \int A \tan^{\frac{5}{2}}(c+dx) dx + \int \left(-B \tan^{\frac{3}{2}}(c+dx) \right) dx \right.$$

$$\left. + \int B \tan^{\frac{7}{2}}(c+dx) dx + \int \left(-2iA \tan^{\frac{3}{2}}(c+dx) \right) dx + \int \left(-2iB \tan^{\frac{5}{2}}(c+dx) \right) dx \right)$$

input `integrate(tan(d*x+c)**(1/2)*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `-a**2*(Integral(-A*sqrt(tan(c + d*x)), x) + Integral(A*tan(c + d*x)**(5/2), x) + Integral(-B*tan(c + d*x)**(3/2), x) + Integral(B*tan(c + d*x)**(7/2), x) + Integral(-2*I*A*tan(c + d*x)**(3/2), x) + Integral(-2*I*B*tan(c + d*x)**(5/2), x))`

3.121.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.50

$$\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$\frac{12 Ba^2 \tan(dx + c)^{\frac{5}{2}} + 20(A - 2iB)a^2 \tan(dx + c)^{\frac{3}{2}} + 120(-iA - B)a^2 \sqrt{\tan(dx + c)} + 15(2\sqrt{2}(($$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/30*(12*B*a^2*tan(d*x + c)^(5/2) + 20*(A - 2*I*B)*a^2*tan(d*x + c)^(3/2) + 120*(-I*A - B)*a^2*sqrt(tan(d*x + c)) + 15*(2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a^2)/d`

3.121.8 Giac [A] (verification not implemented)

Time = 0.75 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.98

$$\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{(2i - 2) \sqrt{2}(Aa^2 - iBa^2) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{d}$$

$$- \frac{2\left(3Ba^2d^4 \tan(dx + c)^{\frac{5}{2}} + 5Aa^2d^4 \tan(dx + c)^{\frac{3}{2}} - 10iBa^2d^4 \tan(dx + c)^{\frac{3}{2}} - 30iAa^2d^4 \sqrt{\tan(dx + c)}\right)}{15d^5}$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output
$$\frac{-(2I - 2)\sqrt{2}(Aa^2 - IBa^2)\arctan\left(\frac{1}{2}I - \frac{1}{2}\sqrt{2}\sqrt{\tan(dx+c)}\right)}{d} - \frac{2}{15}(3Ba^2d^4\tan(dx+c)^{5/2} + 5Aa^2d^4\tan(dx+c)^{3/2} - 10IBa^2d^4\tan(dx+c)^{3/2} - 30IAa^2d^4\sqrt{\tan(dx+c)} - 30Ba^2d^4\sqrt{\tan(dx+c)})/d^5$$

3.121.9 Mupad [B] (verification not implemented)

Time = 9.11 (sec) , antiderivative size = 256, normalized size of antiderivative = 1.98

$$\begin{aligned} & \int \sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2(A+B\tan(c+dx))dx \\ &= \frac{Aa^2\sqrt{\tan(c+dx)}4i}{d} - \frac{2Aa^2\tan(c+dx)^{3/2}}{3d} + \frac{4Ba^2\sqrt{\tan(c+dx)}}{d} \\ &+ \frac{Ba^2\tan(c+dx)^{3/2}4i}{3d} - \frac{2Ba^2\tan(c+dx)^{5/2}}{5d} \\ &+ \frac{\sqrt{2}Aa^2\ln\left(4Aa^2d+\sqrt{2}Aa^2d\sqrt{\tan(c+dx)}(-2-2i)\right)(1+i)}{d} \\ &- \frac{\sqrt{4i}Aa^2\ln\left(4Aa^2d+2\sqrt{4i}Aa^2d\sqrt{\tan(c+dx)}\right)}{d} \\ &+ \frac{\sqrt{2}Ba^2\ln\left(-Ba^2d4i+\sqrt{2}Ba^2d\sqrt{\tan(c+dx)}(-2+2i)\right)(1-i)}{d} \\ &- \frac{\sqrt{-4i}Ba^2\ln\left(-Ba^2d4i+2\sqrt{-4i}Ba^2d\sqrt{\tan(c+dx)}\right)}{d} \end{aligned}$$

input `int(tan(c+d*x)^(1/2)*(A+B*tan(c+d*x))*(a+a*tan(c+d*x)*1i)^2,x)`

output
$$\begin{aligned} & (Aa^2\tan(c+d*x)^{1/2}4i)/d - (2Aa^2\tan(c+d*x)^{3/2})/(3d) + (4Ba^2\tan(c+d*x)^{1/2})/d + (Ba^2\tan(c+d*x)^{3/2}4i)/(3d) - (2Ba^2\tan(c+d*x)^{5/2})/(5d) \\ &+ (2^{1/2}Aa^2\log(4Aa^2d-2^{1/2}Aa^2\tan(c+d*x)^{1/2}*(2+2i))*(1+i))/d - (4i^{1/2}Aa^2\log(4Aa^2d+2*4i^{1/2}Aa^2d\tan(c+d*x)^{1/2}))/d \\ &+ (2^{1/2}Ba^2\log(-Ba^2d*4i-2^{1/2}Ba^2d\tan(c+d*x)^{1/2}*(2-2i))*(1-i))/d - ((-4i)^{1/2}Ba^2\log(2*(-4i)^{1/2}Ba^2d\tan(c+d*x)^{1/2}-Ba^2d*4i))/d \end{aligned}$$

3.122
$$\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.122.1 Optimal result 1278
 3.122.2 Mathematica [A] (verified) 1278
 3.122.3 Rubi [A] (verified) 1279
 3.122.4 Maple [B] (verified) 1281
 3.122.5 Fricas [B] (verification not implemented) 1282
 3.122.6 Sympy [F] 1283
 3.122.7 Maxima [B] (verification not implemented) 1283
 3.122.8 Giac [A] (verification not implemented) 1284
 3.122.9 Mupad [B] (verification not implemented) 1285

3.122.1 Optimal result

Integrand size = 36, antiderivative size = 104

$$\int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= -\frac{4\sqrt{-1}a^2(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d}$$

$$- \frac{2a^2(3A - 5iB)\sqrt{\tan(c + dx)}}{3d} + \frac{2iB\sqrt{\tan(c + dx)}(a^2 + ia^2 \tan(c + dx))}{3d}$$

output `-4*(-1)^(1/4)*a^2*(A-I*B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d-2/3*a^2*(3*A-5*I*B)*tan(d*x+c)^(1/2)/d+2/3*I*B*tan(d*x+c)^(1/2)*(a^2+I*a^2*tan(d*x+c))/d`

3.122.2 Mathematica [A] (verified)

Time = 1.97 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.72

$$\int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx =$$

$$\frac{2a^2\left((3 + 3i)\sqrt{2}(iA + B)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{\tan(c+dx)}}{\sqrt{2}}\right) + \sqrt{\tan(c + dx)}(3A - 6iB + B \tan(c + dx))\right)}{3d}$$

input `Integrate[((a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]] ,x]`

output `(-2*a^2*((3 + 3*I)*Sqrt[2]*(I*A + B)*ArcTanh[((1 + I)*Sqrt[Tan[c + d*x]])/Sqrt[2]] + Sqrt[Tan[c + d*x]]*(3*A - (6*I)*B + B*Tan[c + d*x]))/(3*d)`

3.122.3 Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.03, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4077, 27, 3042, 4075, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
 & \quad \downarrow \text{4077} \\
 & \frac{2}{3} \int \frac{(i \tan(c + dx)a + a)(a(3A - iB) + a(3iA + 5B) \tan(c + dx))}{2\sqrt{\tan(c + dx)}} dx + \\
 & \quad \frac{2iB\sqrt{\tan(c + dx)}(a^2 + ia^2 \tan(c + dx))}{3d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{3} \int \frac{(i \tan(c + dx)a + a)(a(3A - iB) + a(3iA + 5B) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx + \\
 & \quad \frac{2iB\sqrt{\tan(c + dx)}(a^2 + ia^2 \tan(c + dx))}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \frac{(i \tan(c + dx)a + a)(a(3A - iB) + a(3iA + 5B) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx + \\
 & \quad \frac{2iB\sqrt{\tan(c + dx)}(a^2 + ia^2 \tan(c + dx))}{3d} \\
 & \quad \downarrow \text{4075}
 \end{aligned}$$

3.122. $\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

$$\begin{aligned}
& \frac{1}{3} \left(\int \frac{6(A-iB)a^2 + 6(iA+B)\tan(c+dx)a^2}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(3A-5iB)\sqrt{\tan(c+dx)}}{d} \right) + \\
& \quad \frac{2iB\sqrt{\tan(c+dx)}(a^2 + ia^2 \tan(c+dx))}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(\int \frac{6(A-iB)a^2 + 6(iA+B)\tan(c+dx)a^2}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(3A-5iB)\sqrt{\tan(c+dx)}}{d} \right) + \\
& \quad \frac{2iB\sqrt{\tan(c+dx)}(a^2 + ia^2 \tan(c+dx))}{3d} \\
& \quad \downarrow \text{4016} \\
& \frac{1}{3} \left(\frac{72a^4(A-iB)^2 \int \frac{1}{6a^2(A-iB)-6a^2(iA+B)\tan(c+dx)} d\sqrt{\tan(c+dx)}}{d} - \frac{2a^2(3A-5iB)\sqrt{\tan(c+dx)}}{d} \right) + \\
& \quad \frac{2iB\sqrt{\tan(c+dx)}(a^2 + ia^2 \tan(c+dx))}{3d} \\
& \quad \downarrow \text{218} \\
& \frac{1}{3} \left(-\frac{12\sqrt[4]{-1}a^2(A-iB)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{d} - \frac{2a^2(3A-5iB)\sqrt{\tan(c+dx)}}{d} \right) + \\
& \quad \frac{2iB\sqrt{\tan(c+dx)}(a^2 + ia^2 \tan(c+dx))}{3d}
\end{aligned}$$

input `Int[((a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `((-12*(-1)^(1/4)*a^2*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]])/d - (2*a^2*(3*A - (5*I)*B)*Sqrt[Tan[c + d*x]])/d)/3 + (((2*I)/3)*B*Sqrt[Tan[c + d*x]]*(a^2 + I*a^2*Tan[c + d*x]))/d`

3.122.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4077 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.122.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 228 vs. $2(86) = 172$.

Time = 0.03 (sec) , antiderivative size = 229, normalized size of antiderivative = 2.20

method	result
derivativedivides	$a^2 \left(-\frac{2B \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 2A \sqrt{\tan(dx+c)} + 4iB \sqrt{\tan(dx+c)} + \frac{(-2iB+2A)\sqrt{2}}{1-\sqrt{2}} \ln \left(\frac{1+\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)}{1-\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)} \right) + 2 \arctan \left(\frac{1+\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)}{1-\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)} \right) \right)$
default	$a^2 \left(-\frac{2B \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 2A \sqrt{\tan(dx+c)} + 4iB \sqrt{\tan(dx+c)} + \frac{(-2iB+2A)\sqrt{2}}{1-\sqrt{2}} \ln \left(\frac{1+\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)}{1-\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)} \right) + 2 \arctan \left(\frac{1+\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)}{1-\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)} \right) \right)$
parts	$\frac{(2iA a^2 + B a^2) \sqrt{2} \left(\ln \left(\frac{1-\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)}{1+\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)} \right) + 2 \arctan \left(1+\sqrt{2} \left(\sqrt{\tan(dx+c)} \right) \right) + 2 \arctan \left(-1+\sqrt{2} \left(\sqrt{\tan(dx+c)} \right) \right) \right)}{4d}$

```
input int((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/d*a^2*(-2/3*B*tan(d*x+c)^(3/2)-2*A*tan(d*x+c)^(1/2)+4*I*B*tan(d*x+c)^(1/2)+1/4*(2*A-2*I*B)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))+1/4*(2*B+2*I*A)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))
```

3.122.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 389 vs. 2(82) = 164.

Time = 0.25 (sec) , antiderivative size = 389, normalized size of antiderivative = 3.74

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= 3 \sqrt{-\frac{(i A^2 + 2 AB - i B^2) a^4}{d^2}} (de^{(2i dx + 2i c)} + d) \log \left(-\frac{2 \left((A - i B) a^2 e^{(2i dx + 2i c)} + \sqrt{-\frac{(i A^2 + 2 AB - i B^2) a^4}{d^2}} (i de^{(2i dx + 2i c)} + i d) \sqrt{\frac{-i e^{(2i dx + 2i c)}}{e^{(2i dx + 2i c)}}}} \right)}{(-i A - B) a^2} \right)$$

```
input integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="fracas")
```

3.122. $\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$

output $\frac{1}{3}*(3*\sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2}*(d*e^{(2*I*d*x + 2*I*c)} + d)*\log(-2*((A - I*B)*a^2*e^{(2*I*d*x + 2*I*c)} + \sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2}*(I*d*e^{(2*I*d*x + 2*I*c)} + I*d)*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))))*e^{(-2*I*d*x - 2*I*c)}/((-I*A - B)*a^2)) - 3*\sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2}*(d*e^{(2*I*d*x + 2*I*c)} + d)*\log(-2*((A - I*B)*a^2*e^{(2*I*d*x + 2*I*c)} + \sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2}*(-I*d*e^{(2*I*d*x + 2*I*c)} - I*d)*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))))*e^{(-2*I*d*x - 2*I*c)}/((-I*A - B)*a^2)) - 2*((3*A - 7*I*B)*a^2*e^{(2*I*d*x + 2*I*c)} + (3*A - 5*I*B)*a^2)*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))}/(d*e^{(2*I*d*x + 2*I*c)} + d)$

3.122.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= -a^2 \left(\int \left(-\frac{A}{\sqrt{\tan(c + dx)}} \right) dx + \int A \tan^{\frac{3}{2}}(c + dx) dx + \int \left(-B \sqrt{\tan(c + dx)} \right) dx \right. \\ \left. + \int B \tan^{\frac{5}{2}}(c + dx) dx + \int \left(-2iA \sqrt{\tan(c + dx)} \right) dx + \int \left(-2iB \tan^{\frac{3}{2}}(c + dx) \right) dx \right)$$

input `integrate((a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c))/tan(d*x+c)**(1/2),x)`

output `-a**2*(Integral(-A/sqrt(tan(c + d*x)), x) + Integral(A*tan(c + d*x)**(3/2), x) + Integral(-B*sqrt(tan(c + d*x)), x) + Integral(B*tan(c + d*x)**(5/2), x) + Integral(-2*I*A*sqrt(tan(c + d*x)), x) + Integral(-2*I*B*tan(c + d*x)**(3/2), x))`

3.122.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 174 vs. $2(82) = 164$.

Time = 0.28 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.67

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx =$$

$$\frac{4Ba^2 \tan(dx + c)^{\frac{3}{2}} + 12(A - 2iB)a^2 \sqrt{\tan(dx + c)} + 3 \left(2\sqrt{2}(-i + 1)A + (i - 1)B \right) \arctan \left(\frac{1}{2} \sqrt{2} \right)}{1}$$

3.122. $\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="maxima")`

output `-1/6*(4*B*a^2*tan(d*x + c)^(3/2) + 12*(A - 2*I*B)*a^2*sqrt(tan(d*x + c)) + 3*(2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a^2)/d`

3.122.8 Giac [A] (verification not implemented)

Time = 0.76 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.88

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \frac{(2i - 2) \sqrt{2} (-i A a^2 - B a^2) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{d}$$

$$- \frac{2 \left(B a^2 d^2 \tan(dx + c)^{\frac{3}{2}} + 3 A a^2 d^2 \sqrt{\tan(dx + c)} - 6i B a^2 d^2 \sqrt{\tan(dx + c)} \right)}{3 d^3}$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="giac")`

output `(2*I - 2)*sqrt(2)*(-I*A*a^2 - B*a^2)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d - 2/3*(B*a^2*d^2*tan(d*x + c)^(3/2) + 3*A*a^2*d^2*sqrt(tan(d*x + c)) - 6*I*B*a^2*d^2*sqrt(tan(d*x + c)))/d^3`

3.122.9 Mupad [B] (verification not implemented)

Time = 8.85 (sec) , antiderivative size = 221, normalized size of antiderivative = 2.12

$$\begin{aligned}
& \int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
&= -\frac{2 A a^2 \sqrt{\tan(c + dx)}}{d} + \frac{B a^2 \sqrt{\tan(c + dx)} 4i}{d} - \frac{2 B a^2 \tan(c + dx)^{3/2}}{3 d} \\
&+ \frac{\sqrt{2} A a^2 \ln \left(A a^2 d 4i + \sqrt{2} A a^2 d \sqrt{\tan(c + dx)} (-2 + 2i) \right) (1 - i)}{d} \\
&- \frac{\sqrt{-4i} A a^2 \ln \left(A a^2 d 4i + 2 \sqrt{-4i} A a^2 d \sqrt{\tan(c + dx)} \right)}{d} \\
&+ \frac{\sqrt{2} B a^2 \ln \left(4 B a^2 d + \sqrt{2} B a^2 d \sqrt{\tan(c + dx)} (-2 - 2i) \right) (1 + i)}{d} \\
&- \frac{\sqrt{4i} B a^2 \ln \left(4 B a^2 d + 2 \sqrt{4i} B a^2 d \sqrt{\tan(c + dx)} \right)}{d}
\end{aligned}$$

```
input int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2)/tan(c + d*x)^(1/2),x)
```

```
output (B*a^2*tan(c + d*x)^(1/2)*4i)/d - (2*A*a^2*tan(c + d*x)^(1/2))/d - (2*B*a^2*tan(c + d*x)^(3/2))/(3*d) + (2^(1/2)*A*a^2*log(A*a^2*d*4i - 2^(1/2)*A*a^2*d*tan(c + d*x)^(1/2)*(2 - 2i))*(1 - 1i))/d - ((-4i)^(1/2)*A*a^2*log(A*a^2*d*4i + 2*(-4i)^(1/2)*A*a^2*d*tan(c + d*x)^(1/2)))/d + (2^(1/2)*B*a^2*log(4*B*a^2*d - 2^(1/2)*B*a^2*d*tan(c + d*x)^(1/2)*(2 + 2i))*(1 + 1i))/d - (4i^(1/2)*B*a^2*log(4*B*a^2*d + 2*4i^(1/2)*B*a^2*d*tan(c + d*x)^(1/2)))/d
```

3.123
$$\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.123.1 Optimal result 1286
 3.123.2 Mathematica [A] (verified) 1286
 3.123.3 Rubi [A] (verified) 1287
 3.123.4 Maple [B] (verified) 1289
 3.123.5 Fricas [B] (verification not implemented) 1290
 3.123.6 Sympy [F] 1291
 3.123.7 Maxima [B] (verification not implemented) 1291
 3.123.8 Giac [A] (verification not implemented) 1292
 3.123.9 Mupad [B] (verification not implemented) 1292

3.123.1 Optimal result

Integrand size = 36, antiderivative size = 98

$$\int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= -\frac{4\sqrt{-1}a^2(iA + B) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d}$$

$$+ \frac{2a^2(iA - B)\sqrt{\tan(c + dx)}}{d} - \frac{2A(a^2 + ia^2 \tan(c + dx))}{d\sqrt{\tan(c + dx)}}$$

output `-4*(-1)^(1/4)*a^2*(I*A+B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d+2*a^2*(I*A-B)*tan(d*x+c)^(1/2)/d-2*A*(a^2+I*a^2*tan(d*x+c))/d/tan(d*x+c)^(1/2)`

3.123.2 Mathematica [A] (verified)

Time = 1.37 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.76

$$\int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= -\frac{2a^2\left(A - (1 + i)\sqrt{2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)\sqrt{\tan(c + dx)} + B \tan(c + dx)\right)}{d\sqrt{\tan(c + dx)}}$$

input `Integrate[((a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `(-2*a^2*(A - (1 + I)*Sqrt[2]*(A - I*B)*ArcTanh[((1 + I)*Sqrt[Tan[c + d*x]])/Sqrt[2]]*Sqrt[Tan[c + d*x]] + B*Tan[c + d*x]))/(d*Sqrt[Tan[c + d*x]])`

3.123.3 Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4076, 27, 3042, 4075, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx \\
 & \quad \downarrow \text{4076} \\
 & 2 \int \frac{(i \tan(c + dx)a + a)(a(3iA + B) + a(A + iB) \tan(c + dx))}{2\sqrt{\tan(c + dx)}} dx - \frac{2A(a^2 + ia^2 \tan(c + dx))}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(i \tan(c + dx)a + a)(a(3iA + B) + a(A + iB) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx - \frac{2A(a^2 + ia^2 \tan(c + dx))}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(i \tan(c + dx)a + a)(a(3iA + B) + a(A + iB) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx - \frac{2A(a^2 + ia^2 \tan(c + dx))}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{4075} \\
 & \int \frac{2a^2(iA + B) - 2a^2(A - iB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2a^2(-B + iA)\sqrt{\tan(c + dx)}}{d} - \\
 & \quad \frac{2A(a^2 + ia^2 \tan(c + dx))}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.123. $\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{2a^2(iA + B) - 2a^2(A - iB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2a^2(-B + iA) \sqrt{\tan(c + dx)}}{d} - \\
& \quad \frac{2A(a^2 + ia^2 \tan(c + dx))}{d \sqrt{\tan(c + dx)}} \\
& \quad \downarrow \text{4016} \\
& \frac{8a^4(B + iA)^2 \int \frac{1}{2(iA+B)a^2+2(A-iB)\tan(c+dx)a^2} d\sqrt{\tan(c+dx)}}{d} + \frac{2a^2(-B + iA) \sqrt{\tan(c + dx)}}{d} - \\
& \quad \frac{2A(a^2 + ia^2 \tan(c + dx))}{d \sqrt{\tan(c + dx)}} \\
& \quad \downarrow \text{218} \\
& - \frac{4\sqrt[4]{-1}a^2(B + iA) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right)}{d} + \frac{2a^2(-B + iA) \sqrt{\tan(c + dx)}}{d} - \\
& \quad \frac{2A(a^2 + ia^2 \tan(c + dx))}{d \sqrt{\tan(c + dx)}}
\end{aligned}$$

input `Int[((a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `(-4*(-1)^(1/4)*a^2*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]])/d + (2*a^2*(I*A - B)*Sqrt[Tan[c + d*x]])/d - (2*A*(a^2 + I*a^2*Tan[c + d*x]))/(d*Sqrt[Tan[c + d*x]])`

3.123.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.123.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 216 vs. 2(85) = 170.

Time = 0.04 (sec) , antiderivative size = 217, normalized size of antiderivative = 2.21

method	result
derivativedivides	$a^2 \left(-2B(\sqrt{\tan(dx+c)}) - \frac{2A}{\sqrt{\tan(dx+c)}} + \frac{(2iA+2B)\sqrt{2}}{4} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right) \right)$
default	$a^2 \left(-2B(\sqrt{\tan(dx+c)}) - \frac{2A}{\sqrt{\tan(dx+c)}} + \frac{(2iA+2B)\sqrt{2}}{4} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right) \right)$
parts	$\frac{(2iAa^2+B a^2)\sqrt{2}}{4d} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)$

3.123.
$$\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

input `int((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`

output `1/d*a^2*(-2*B*tan(d*x+c)^(1/2)-2*A/tan(d*x+c)^(1/2)+1/4*(2*B+2*I*A)*2^(1/2))*ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))+1/4*(-2*A+2*I*B)*2^(1/2)*ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))`

3.123.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 386 vs. $2(82) = 164$.

Time = 0.26 (sec) , antiderivative size = 386, normalized size of antiderivative = 3.94

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{\sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^4}{d^2}} (de^{(2i dx + 2i c)} - d) \log \left(-\frac{2 \left((A - iB)a^2 e^{(2i dx + 2i c)} + \sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^4}{d^2}} (de^{(2i dx + 2i c)} + d) \sqrt{\frac{-ie^{(2i dx + 2i c)}}{e^{(2i dx + 2i c)}}}}{(-iA - B)a^2} \right)}{\dots} \right)}{\dots}$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="fracas")`

output `(sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log(-2*((A - I*B)*a^2*e^(2*I*d*x + 2*I*c) + sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^2)) - sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log(-2*((A - I*B)*a^2*e^(2*I*d*x + 2*I*c) - sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^2)) - 2*((I*A + B)*a^2*e^(2*I*d*x + 2*I*c) + (I*A - B)*a^2)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(2*I*d*x + 2*I*c) - d)`

3.123.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= -a^2 \left(\int \left(-\frac{A}{\tan^{\frac{3}{2}}(c + dx)} \right) dx + \int A \sqrt{\tan(c + dx)} dx + \int \left(-\frac{B}{\sqrt{\tan(c + dx)}} \right) dx \right.$$

$$\left. + \int B \tan^{\frac{3}{2}}(c + dx) dx + \int \left(-\frac{2iA}{\sqrt{\tan(c + dx)}} \right) dx + \int \left(-2iB \sqrt{\tan(c + dx)} \right) dx \right)$$

input `integrate((a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c))/tan(d*x+c)**(3/2),x)`

output `-a**2*(Integral(-A/tan(c + d*x)**(3/2), x) + Integral(A*sqrt(tan(c + d*x)), x) + Integral(-B/sqrt(tan(c + d*x)), x) + Integral(B*tan(c + d*x)**(3/2), x) + Integral(-2*I*A/sqrt(tan(c + d*x)), x) + Integral(-2*I*B*sqrt(tan(c + d*x)), x))`

3.123.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 170 vs. 2(82) = 164.

Time = 0.43 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.73

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{4Ba^2 \sqrt{\tan(dx + c)} - \left(2\sqrt{2}((i-1)A + (i+1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(dx + c)}\right)\right) + 2\sqrt{2}((i+1)A - (i-1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - 2\sqrt{\tan(dx + c)}\right)\right) \right)}{\tan^{\frac{3}{2}}(c + dx)}$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="maxima")`

output `-1/2*(4*B*a^2*sqrt(tan(d*x + c)) - (2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a^2 + 4*A*a^2/sqrt(tan(d*x + c)))/d`

3.123. $\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

3.123.8 Giac [A] (verification not implemented)

Time = 0.84 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.70

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= -\frac{2 B a^2 \sqrt{\tan(dx + c)}}{d} + \frac{(2i - 2) \sqrt{2}(A a^2 - i B a^2) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{d} - \frac{2 A a^2}{d \sqrt{\tan(dx + c)}}$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="giac")`

output `-2*B*a^2*sqrt(tan(d*x + c))/d + (2*I - 2)*sqrt(2)*(A*a^2 - I*B*a^2)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d - 2*A*a^2/(d*sqrt(tan(d*x + c)))`

3.123.9 Mupad [B] (verification not implemented)

Time = 7.71 (sec) , antiderivative size = 203, normalized size of antiderivative = 2.07

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= -\frac{2 A a^2}{d \sqrt{\tan(c + dx)}} - \frac{2 B a^2 \sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{\sqrt{2} A a^2 \ln\left(-4 A a^2 d + \sqrt{2} A a^2 d \sqrt{\tan(c + dx)}(-2 - 2i)\right) (1 + 1i)}{d}$$

$$- \frac{\sqrt{4i} A a^2 \ln\left(-4 A a^2 d + 2 \sqrt{4i} A a^2 d \sqrt{\tan(c + dx)}\right)}{d}$$

$$+ \frac{\sqrt{2} B a^2 \ln\left(B a^2 d 4i + \sqrt{2} B a^2 d \sqrt{\tan(c + dx)}(-2 + 2i)\right) (1 - i)}{d}$$

$$- \frac{\sqrt{-4i} B a^2 \ln\left(B a^2 d 4i + 2 \sqrt{-4i} B a^2 d \sqrt{\tan(c + dx)}\right)}{d}$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2)/tan(c + d*x)^(3/2),x)`

output $(2^{(1/2)}*A*a^2*\log(-4*A*a^2*d - 2^{(1/2)}*A*a^2*d*\tan(c + d*x)^{(1/2)}*(2 + 2i))*(1 + i))/d - (2*B*a^2*\tan(c + d*x)^{(1/2)})/d - (2*A*a^2)/(d*\tan(c + d*x)^{(1/2)}) - (4i^{(1/2)}*A*a^2*\log(2*4i^{(1/2)}*A*a^2*d*\tan(c + d*x)^{(1/2)} - 4*A*a^2*d))/d + (2^{(1/2)}*B*a^2*\log(B*a^2*d*4i - 2^{(1/2)}*B*a^2*d*\tan(c + d*x)^{(1/2)}*(2 - 2i))*(1 - i))/d - ((-4i)^{(1/2)}*B*a^2*\log(B*a^2*d*4i + 2*(-4i)^{(1/2)}*B*a^2*d*\tan(c + d*x)^{(1/2)}))/d$

$$3.124 \quad \int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.124.1 Optimal result 1294
 3.124.2 Mathematica [A] (verified) 1294
 3.124.3 Rubi [A] (verified) 1295
 3.124.4 Maple [B] (verified) 1297
 3.124.5 Fricas [B] (verification not implemented) 1298
 3.124.6 Sympy [F] 1299
 3.124.7 Maxima [B] (verification not implemented) 1300
 3.124.8 Giac [A] (verification not implemented) 1300
 3.124.9 Mupad [B] (verification not implemented) 1301

3.124.1 Optimal result

Integrand size = 36, antiderivative size = 102

$$\int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{4\sqrt[4]{-1}a^2(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{3d\sqrt{\tan(c + dx)}} - \frac{2A(a^2 + ia^2 \tan(c + dx))}{3d \tan^{\frac{3}{2}}(c + dx)}$$

```
output 4*(-1)^(1/4)*a^2*(A-I*B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d-2/3*a^2*(5*I*A+3*B)/d/tan(d*x+c)^(1/2)-2/3*A*(a^2+I*a^2*tan(d*x+c))/d/tan(d*x+c)^(3/2)
```

3.124.2 Mathematica [A] (verified)

Time = 1.58 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.84

$$\int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2a^2\left(-A + (-6iA - 3B) \tan(c + dx) + (3 + 3i)\sqrt{2}(iA + B) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)\right) \tan^{\frac{3}{2}}(c + dx)}{3d \tan^{\frac{3}{2}}(c + dx)}$$

input `Integrate[((a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(2*a^2*(-A + ((-6*I)*A - 3*B)*Tan[c + d*x] + (3 + 3*I)*Sqrt[2]*(I*A + B)*ArcTanh[((1 + I)*Sqrt[Tan[c + d*x]])/Sqrt[2]]*Tan[c + d*x]^(3/2))/(3*d*Tan[c + d*x]^(3/2))`

3.124.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3042, 4076, 27, 3042, 4074, 27, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx$$

↓ 4076

$$\frac{2}{3} \int \frac{(i \tan(c + dx)a + a)(a(5iA + 3B) - a(A - 3iB) \tan(c + dx))}{2 \tan^{\frac{3}{2}}(c + dx)} dx - \frac{2A(a^2 + ia^2 \tan(c + dx))}{3d \tan^{\frac{3}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{3} \int \frac{(i \tan(c + dx)a + a)(a(5iA + 3B) - a(A - 3iB) \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx - \frac{2A(a^2 + ia^2 \tan(c + dx))}{3d \tan^{\frac{3}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \int \frac{(i \tan(c + dx)a + a)(a(5iA + 3B) - a(A - 3iB) \tan(c + dx))}{\tan(c + dx)^{3/2}} dx - \frac{2A(a^2 + ia^2 \tan(c + dx))}{3d \tan^{\frac{3}{2}}(c + dx)}$$

↓ 4074

$$\frac{1}{3} \left(\int -\frac{6((A - iB)a^2 + (iA + B) \tan(c + dx)a^2)}{\sqrt{\tan(c + dx)}} dx - \frac{2a^2(3B + 5iA)}{d\sqrt{\tan(c + dx)}} \right) - \frac{2A(a^2 + ia^2 \tan(c + dx))}{3d \tan^{\frac{3}{2}}(c + dx)}$$

3.124. $\int \frac{(a+ia \tan(c+dx))^2 (A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{1}{3} \left(-6 \int \frac{(A - iB)a^2 + (iA + B) \tan(c + dx)a^2}{\sqrt{\tan(c + dx)}} dx - \frac{2a^2(3B + 5iA)}{d\sqrt{\tan(c + dx)}} \right) - \frac{2A(a^2 + ia^2 \tan(c + dx))}{3d \tan^{\frac{3}{2}}(c + dx)} \\
& \downarrow 3042 \\
& \frac{1}{3} \left(-6 \int \frac{(A - iB)a^2 + (iA + B) \tan(c + dx)a^2}{\sqrt{\tan(c + dx)}} dx - \frac{2a^2(3B + 5iA)}{d\sqrt{\tan(c + dx)}} \right) - \frac{2A(a^2 + ia^2 \tan(c + dx))}{3d \tan^{\frac{3}{2}}(c + dx)} \\
& \downarrow 4016 \\
& \frac{1}{3} \left(-\frac{12a^4(A - iB)^2 \int \frac{1}{a^2(A - iB) - a^2(iA + B) \tan(c + dx)} d\sqrt{\tan(c + dx)}}{d} - \frac{2a^2(3B + 5iA)}{d\sqrt{\tan(c + dx)}} \right) - \\
& \quad \frac{2A(a^2 + ia^2 \tan(c + dx))}{3d \tan^{\frac{3}{2}}(c + dx)} \\
& \downarrow 218 \\
& \frac{1}{3} \left(\frac{12\sqrt[4]{-1}a^2(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d} - \frac{2a^2(3B + 5iA)}{d\sqrt{\tan(c + dx)}} \right) - \\
& \quad \frac{2A(a^2 + ia^2 \tan(c + dx))}{3d \tan^{\frac{3}{2}}(c + dx)}
\end{aligned}$$

input `Int[((a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `((12*(-1)^(1/4)*a^2*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]])/d - (2*a^2*((5*I)*A + 3*B))/(d*Sqrt[Tan[c + d*x]])/3 - (2*A*(a^2 + I*a^2*Tan[c + d*x]))/(3*d*Tan[c + d*x]^(3/2))`

3.124.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4074 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.124.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 221 vs. $2(85) = 170$.

Time = 0.04 (sec) , antiderivative size = 222, normalized size of antiderivative = 2.18

$$3.124. \quad \int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

method	result
derivativedivides	$a^2 \left(-\frac{2A}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(2iA+B)}{\sqrt{\tan(dx+c)}} + \frac{(2iB-2A)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4} \right)$
default	$a^2 \left(-\frac{2A}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(2iA+B)}{\sqrt{\tan(dx+c)}} + \frac{(2iB-2A)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4} \right)$
parts	$(2iA a^2 + B a^2) \left(-\frac{\sqrt{2} \left(\ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4} \right)$ d

input `int((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x,method=_RETURNVERBOSE)`

output `1/d*a^2*(-2/3*A/tan(d*x+c)^(3/2)-2*(2*I*A+B)/tan(d*x+c)^(1/2)+1/4*(-2*A+2*I*B)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(-2*B-2*I*A)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))`

3.124.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 441 vs. 2(82) = 164.

Time = 0.26 (sec) , antiderivative size = 441, normalized size of antiderivative = 4.32

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx =$$

$$3 \sqrt{-\frac{(i A^2 + 2 AB - i B^2) a^4}{d^2}} (de^{(4i dx + 4i c)} - 2 de^{(2i dx + 2i c)} + d) \log \left(-\frac{2 \left((A - i B) a^2 e^{(2i dx + 2i c)} + \sqrt{-\frac{(i A^2 + 2 AB - i B^2) a^4}{d^2}} \right)}{(-i A - \dots)} \right)$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="fracas")`

3.124.
$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

output

```
-1/3*(3*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(d*e^(4*I*d*x + 4*I*c) - 2*
d*e^(2*I*d*x + 2*I*c) + d)*log(-2*((A - I*B)*a^2*e^(2*I*d*x + 2*I*c) + sqrt
(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt((
-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x - 2*I*
c)/((-I*A - B)*a^2)) - 3*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(d*e^(4*I*
d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log(-2*((A - I*B)*a^2*e^(2*I*d
*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(-I*d*e^(2*I*d*x + 2*
I*c) - I*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e
^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^2)) - 2*((7*A - 3*I*B)*a^2*e^(4*I*d*x +
4*I*c) + 2*A*a^2*e^(2*I*d*x + 2*I*c) - (5*A - 3*I*B)*a^2)*sqrt((-I*e^(2*I
*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(4*I*d*x + 4*I*c) - 2*
d*e^(2*I*d*x + 2*I*c) + d)
```

3.124.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= -a^2 \left(\int \left(-\frac{A}{\tan^{\frac{5}{2}}(c + dx)} \right) dx + \int \frac{A}{\sqrt{\tan(c + dx)}} dx + \int \left(-\frac{B}{\tan^{\frac{3}{2}}(c + dx)} \right) dx \right.$$

$$\left. + \int B \sqrt{\tan(c + dx)} dx + \int \left(-\frac{2iA}{\tan^{\frac{3}{2}}(c + dx)} \right) dx + \int \left(-\frac{2iB}{\sqrt{\tan(c + dx)}} \right) dx \right)$$

input `integrate((a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c))/tan(d*x+c)**(5/2),x)`

output

```
-a**2*(Integral(-A/tan(c + d*x)**(5/2), x) + Integral(A/sqrt(tan(c + d*x))
, x) + Integral(-B/tan(c + d*x)**(3/2), x) + Integral(B*sqrt(tan(c + d*x))
, x) + Integral(-2*I*A/tan(c + d*x)**(3/2), x) + Integral(-2*I*B/sqrt(tan(
c + d*x)), x))
```

3.124.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 177 vs. $2(82) = 164$.

Time = 0.39 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.74

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{3 \left(2\sqrt{2}(-i + 1) A + (i - 1) B \right) \arctan \left(\frac{1}{2} \sqrt{2} \left(\sqrt{2} + 2 \sqrt{\tan(dx + c)} \right) \right) + 2\sqrt{2}(-i + 1) A + (i - 1) B}{1}$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="maxima")`

output `1/6*(3*(2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a^2 + 4*(3*(-2*I*A - B)*a^2*tan(d*x + c) - A*a^2)/tan(d*x + c)^(3/2))/d`

3.124.8 Giac [A] (verification not implemented)

Time = 0.89 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.77

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= - \frac{(2i - 2) \sqrt{2}(-i A a^2 - B a^2) \arctan \left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)} \right)}{d}$$

$$- \frac{2(6i A a^2 \tan(dx + c) + 3 B a^2 \tan(dx + c) + A a^2)}{3 d \tan(dx + c)^{\frac{3}{2}}}$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="giac")`

output `-(2*I - 2)*sqrt(2)*(-I*A*a^2 - B*a^2)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d - 2/3*(6*I*A*a^2*tan(d*x + c) + 3*B*a^2*tan(d*x + c) + A*a^2)/(d*tan(d*x + c)^(3/2))`

3.124. $\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

3.124.9 Mupad [B] (verification not implemented)

Time = 8.38 (sec) , antiderivative size = 222, normalized size of antiderivative = 2.18

$$\begin{aligned}
& \int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \\
&= -\frac{\frac{2Aa^2}{3d} + \frac{Aa^2 \tan(c+dx)4i}{d}}{\tan(c + dx)^{3/2}} - \frac{2Ba^2}{d\sqrt{\tan(c + dx)}} \\
&+ \frac{\sqrt{2}Aa^2 \ln\left(-Aa^2 d4i + \sqrt{2}Aa^2 d\sqrt{\tan(c + dx)}(-2 + 2i)\right)(1 - i)}{d} \\
&- \frac{\sqrt{-4i}Aa^2 \ln\left(-Aa^2 d4i + 2\sqrt{-4i}Aa^2 d\sqrt{\tan(c + dx)}\right)}{d} \\
&+ \frac{\sqrt{2}Ba^2 \ln\left(-4Ba^2 d + \sqrt{2}Ba^2 d\sqrt{\tan(c + dx)}(-2 - 2i)\right)(1 + i)}{d} \\
&- \frac{\sqrt{4i}Ba^2 \ln\left(-4Ba^2 d + 2\sqrt{4i}Ba^2 d\sqrt{\tan(c + dx)}\right)}{d}
\end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2)/tan(c + d*x)^(5/2),x)`

output `(2^(1/2)*A*a^2*log(-A*a^2*d*4i - 2^(1/2)*A*a^2*d*tan(c + d*x)^(1/2)*(2 - 2i))*(1 - 1i))/d - (2*B*a^2)/(d*tan(c + d*x)^(1/2)) - ((2*A*a^2)/(3*d) + (A*a^2*tan(c + d*x)*4i)/d)/tan(c + d*x)^(3/2) - ((-4i)^(1/2)*A*a^2*log(2*(-4i)^(1/2)*A*a^2*d*tan(c + d*x)^(1/2) - A*a^2*d*4i))/d + (2^(1/2)*B*a^2*log(-4*B*a^2*d - 2^(1/2)*B*a^2*d*tan(c + d*x)^(1/2)*(2 + 2i))*(1 + 1i))/d - (4i^(1/2)*B*a^2*log(2*4i^(1/2)*B*a^2*d*tan(c + d*x)^(1/2) - 4*B*a^2*d))/d`

3.125
$$\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

3.125.1 Optimal result 1302
 3.125.2 Mathematica [A] (verified) 1302
 3.125.3 Rubi [A] (verified) 1303
 3.125.4 Maple [B] (verified) 1306
 3.125.5 Fricas [B] (verification not implemented) 1307
 3.125.6 Sympy [F] 1308
 3.125.7 Maxima [A] (verification not implemented) 1308
 3.125.8 Giac [A] (verification not implemented) 1309
 3.125.9 Mupad [B] (verification not implemented) 1309

3.125.1 Optimal result

Integrand size = 36, antiderivative size = 127

$$\int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{4\sqrt{-1}a^2(iA + B) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{15d \tan^{\frac{3}{2}}(c + dx)} + \frac{d}{4a^2(A - iB)\sqrt{\tan(c + dx)}} - \frac{2A(a^2 + ia^2 \tan(c + dx))}{5d \tan^{\frac{5}{2}}(c + dx)}$$

output `4*(-1)^(1/4)*a^2*(I*A+B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d+4*a^2*(A-I*B)/d/tan(d*x+c)^(1/2)-2/15*a^2*(7*I*A+5*B)/d/tan(d*x+c)^(3/2)-2/5*A*(a^2+I*a^2*tan(d*x+c))/d/tan(d*x+c)^(5/2)`

3.125.2 Mathematica [A] (verified)

Time = 2.65 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.80

$$\int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx =$$

$$\frac{2a^2\left(3A + 5(2iA + B) \tan(c + dx) - 30(A - iB) \tan^2(c + dx) + (15 + 15i)\sqrt{2}(A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{\tan(c + dx)}}{1+i}\right)\right)}{15d \tan^{\frac{5}{2}}(c + dx)}$$

input `Integrate[((a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(-2*a^2*(3*A + 5*((2*I)*A + B)*Tan[c + d*x] - 30*(A - I*B)*Tan[c + d*x]^2 + (15 + 15*I)*Sqrt[2]*(A - I*B)*ArcTanh[((1 + I)*Sqrt[Tan[c + d*x]])/Sqrt[2]])*Tan[c + d*x]^(5/2))/(15*d*Tan[c + d*x]^(5/2))`

3.125.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.306$, Rules used = {3042, 4076, 27, 3042, 4074, 27, 3042, 4012, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan(c + dx)^{7/2}} dx$$

↓ 4076

$$\frac{2}{5} \int \frac{(i \tan(c + dx)a + a)(a(7iA + 5B) - a(3A - 5iB) \tan(c + dx))}{2 \tan^{\frac{5}{2}}(c + dx)} dx - \frac{2A(a^2 + ia^2 \tan(c + dx))}{5d \tan^{\frac{5}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{5} \int \frac{(i \tan(c + dx)a + a)(a(7iA + 5B) - a(3A - 5iB) \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx - \frac{2A(a^2 + ia^2 \tan(c + dx))}{5d \tan^{\frac{5}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{5} \int \frac{(i \tan(c + dx)a + a)(a(7iA + 5B) - a(3A - 5iB) \tan(c + dx))}{\tan(c + dx)^{5/2}} dx - \frac{2A(a^2 + ia^2 \tan(c + dx))}{5d \tan^{\frac{5}{2}}(c + dx)}$$

↓ 4074

$$\frac{1}{5} \left(\int -\frac{10((A - iB)a^2 + (iA + B) \tan(c + dx)a^2)}{\tan^{\frac{3}{2}}(c + dx)} dx - \frac{2a^2(5B + 7iA)}{3d \tan^{\frac{3}{2}}(c + dx)} \right) - \frac{2A(a^2 + ia^2 \tan(c + dx))}{5d \tan^{\frac{5}{2}}(c + dx)}$$

3.125. $\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{1}{5} \left(-10 \int \frac{(A - iB)a^2 + (iA + B) \tan(c + dx)a^2}{\tan^{\frac{3}{2}}(c + dx)} dx - \frac{2a^2(5B + 7iA)}{3d \tan^{\frac{3}{2}}(c + dx)} \right) - \\
& \quad \frac{2A(a^2 + ia^2 \tan(c + dx))}{5d \tan^{\frac{5}{2}}(c + dx)} \\
& \downarrow 3042 \\
& \frac{1}{5} \left(-10 \int \frac{(A - iB)a^2 + (iA + B) \tan(c + dx)a^2}{\tan(c + dx)^{3/2}} dx - \frac{2a^2(5B + 7iA)}{3d \tan^{\frac{3}{2}}(c + dx)} \right) - \\
& \quad \frac{2A(a^2 + ia^2 \tan(c + dx))}{5d \tan^{\frac{5}{2}}(c + dx)} \\
& \downarrow 4012 \\
& \frac{1}{5} \left(-10 \left(\int \frac{a^2(iA + B) - a^2(A - iB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2a^2(A - iB)}{d\sqrt{\tan(c + dx)}} \right) - \frac{2a^2(5B + 7iA)}{3d \tan^{\frac{3}{2}}(c + dx)} \right) - \\
& \quad \frac{2A(a^2 + ia^2 \tan(c + dx))}{5d \tan^{\frac{5}{2}}(c + dx)} \\
& \downarrow 3042 \\
& \frac{1}{5} \left(-10 \left(\int \frac{a^2(iA + B) - a^2(A - iB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2a^2(A - iB)}{d\sqrt{\tan(c + dx)}} \right) - \frac{2a^2(5B + 7iA)}{3d \tan^{\frac{3}{2}}(c + dx)} \right) - \\
& \quad \frac{2A(a^2 + ia^2 \tan(c + dx))}{5d \tan^{\frac{5}{2}}(c + dx)} \\
& \downarrow 4016 \\
& \frac{1}{5} \left(-10 \left(\frac{2a^4(B + iA)^2 \int \frac{1}{(iA+B)a^2+(A-iB)\tan(c+dx)a^2} d\sqrt{\tan(c+dx)}}{d} - \frac{2a^2(A - iB)}{d\sqrt{\tan(c + dx)}} \right) - \frac{2a^2(5B + 7iA)}{3d \tan^{\frac{3}{2}}(c + dx)} \right) - \\
& \quad \frac{2A(a^2 + ia^2 \tan(c + dx))}{5d \tan^{\frac{5}{2}}(c + dx)} \\
& \downarrow 218 \\
& \frac{1}{5} \left(-10 \left(-\frac{2\sqrt[4]{-1}a^2(B + iA) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right)}{d} - \frac{2a^2(A - iB)}{d\sqrt{\tan(c + dx)}} \right) - \frac{2a^2(5B + 7iA)}{3d \tan^{\frac{3}{2}}(c + dx)} \right) - \\
& \quad \frac{2A(a^2 + ia^2 \tan(c + dx))}{5d \tan^{\frac{5}{2}}(c + dx)}
\end{aligned}$$

3.125. $\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

input `Int[((a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(-10*((-2*(-1)^(1/4)*a^2*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]])/d - (2*a^2*(A - I*B))/(d*Sqrt[Tan[c + d*x]]) - (2*a^2*((7*I)*A + 5*B))/(3*d*Tan[c + d*x]^(3/2)))/5 - (2*A*(a^2 + I*a^2*Tan[c + d*x]))/(5*d*Tan[c + d*x]^(5/2))`

3.125.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`


```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

```
rule 4076 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

3.125.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 239 vs. 2(107) = 214.

Time = 0.03 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.89

method	result
derivativedivides	$a^2 \left(-\frac{2A}{5 \tan(dx+c)^{\frac{5}{2}}} - \frac{2(2iB-2A)}{\sqrt{\tan(dx+c)}} - \frac{2(2iA+B)}{3 \tan(dx+c)^{\frac{3}{2}}} + \frac{(-2iA-2B)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right)}{4} \right)$
default	$a^2 \left(-\frac{2A}{5 \tan(dx+c)^{\frac{5}{2}}} - \frac{2(2iB-2A)}{\sqrt{\tan(dx+c)}} - \frac{2(2iA+B)}{3 \tan(dx+c)^{\frac{3}{2}}} + \frac{(-2iA-2B)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right)}{4} \right)$
parts	$\frac{(2iA a^2 + B a^2) \left(-\frac{2}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right)}{4} \right)}{d}$

```
input int((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x,method=_RETURNVERBOSE)
```

$$3.125. \int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

output $1/d*a^2*(-2/5*A/\tan(d*x+c)^{(5/2)}-2*(-2*A+2*I*B)/\tan(d*x+c)^{(1/2)}-2/3*(2*I*A+B)/\tan(d*x+c)^{(3/2)}+1/4*(-2*B-2*I*A)*2^{(1/2)}*(\ln((1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+1/4*(2*A-2*I*B)*2^{(1/2)}*(\ln((1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}))$

3.125.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 502 vs. $2(103) = 206$.

Time = 0.27 (sec) , antiderivative size = 502, normalized size of antiderivative = 3.95

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx =$$

$$15 \sqrt{-\frac{(-i A^2 - 2AB + i B^2)a^4}{d^2}} (de^{(6i dx + 6i c)} - 3de^{(4i dx + 4i c)} + 3de^{(2i dx + 2i c)} - d) \log \left(-\frac{2 \left((A - i B)a^2 e^{(2i dx + 2i c)} + \sqrt{\dots} \right)}{\dots} \right)$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="fracas")`

output $-1/15*(15*\sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2}*(d*e^{(6*I*d*x + 6*I*c)} - 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} - d)*\log(-2*((A - I*B)*a^2*e^{(2*I*d*x + 2*I*c)} + \sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2}*(d*e^{(2*I*d*x + 2*I*c)} + d)*\sqrt{(-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1)}))e^{(-2*I*d*x - 2*I*c)}/((-I*A - B)*a^2)) - 15*\sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2}*(d*e^{(6*I*d*x + 6*I*c)} - 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} - d)*\log(-2*((A - I*B)*a^2*e^{(2*I*d*x + 2*I*c)} - \sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2}*(d*e^{(2*I*d*x + 2*I*c)} + d)*\sqrt{(-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1)}))e^{(-2*I*d*x - 2*I*c)}/((-I*A - B)*a^2)) + 2*((-43*I*A - 35*B)*a^2*e^{(6*I*d*x + 6*I*c)} + (11*I*A + 25*B)*a^2*e^{(4*I*d*x + 4*I*c)} + (31*I*A + 35*B)*a^2*e^{(2*I*d*x + 2*I*c)} + (-23*I*A - 25*B)*a^2)*\sqrt{(-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1)})/(d*e^{(6*I*d*x + 6*I*c)} - 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} - d)$

$$3.125. \int \frac{(a+ia \tan(c+dx))^2 (A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

3.125.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= -a^2 \left(\int \left(-\frac{A}{\tan^{\frac{7}{2}}(c + dx)} \right) dx + \int \frac{A}{\tan^{\frac{3}{2}}(c + dx)} dx + \int \left(-\frac{B}{\tan^{\frac{5}{2}}(c + dx)} \right) dx \right.$$

$$\left. + \int \frac{B}{\sqrt{\tan(c + dx)}} dx + \int \left(-\frac{2iA}{\tan^{\frac{5}{2}}(c + dx)} \right) dx + \int \left(-\frac{2iB}{\tan^{\frac{3}{2}}(c + dx)} \right) dx \right)$$

input `integrate((a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c))/tan(d*x+c)**(7/2),x)`

output `-a**2*(Integral(-A/tan(c + d*x)**(7/2), x) + Integral(A/tan(c + d*x)**(3/2), x) + Integral(-B/tan(c + d*x)**(5/2), x) + Integral(B/sqrt(tan(c + d*x)), x) + Integral(-2*I*A/tan(c + d*x)**(5/2), x) + Integral(-2*I*B/tan(c + d*x)**(3/2), x))`

3.125.7 Maxima [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.54

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx =$$

$$\frac{15 \left(2 \sqrt{2} ((i - 1) A + (i + 1) B) \arctan \left(\frac{1}{2} \sqrt{2} \left(\sqrt{2} + 2 \sqrt{\tan(dx + c)} \right) \right) + 2 \sqrt{2} ((i - 1) A + (i + 1) B) \right)}{\dots}$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="maxima")`

output `-1/30*(15*(2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a^2 - 4*(30*(A - I*B)*a^2*tan(d*x + c)^2 + 5*(-2*I*A - B)*a^2*tan(d*x + c) - 3*A*a^2)/tan(d*x + c)^(5/2)/d`

3.125. $\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

3.125.8 Giac [A] (verification not implemented)

Time = 1.09 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.84

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= - \frac{(2i - 2) \sqrt{2} (Aa^2 - iBa^2) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{d} + \frac{2(30Aa^2 \tan(dx + c)^2 - 30iBa^2 \tan(dx + c)^2 - 10iAa^2 \tan(dx + c) - 5Ba^2 \tan(dx + c) - 3Aa^2)}{15d \tan(dx + c)^{\frac{5}{2}}}$$

input `integrate((a+i*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="giac")`

output `-(2*I - 2)*sqrt(2)*(A*a^2 - I*B*a^2)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d + 2/15*(30*A*a^2*tan(d*x + c)^2 - 30*I*B*a^2*tan(d*x + c)^2 - 10*I*A*a^2*tan(d*x + c) - 5*B*a^2*tan(d*x + c) - 3*A*a^2)/(d*tan(d*x + c)^(5/2))`

3.125.9 Mupad [B] (verification not implemented)

Time = 9.50 (sec) , antiderivative size = 258, normalized size of antiderivative = 2.03

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= - \frac{\frac{2Aa^2}{5d} - \frac{4Aa^2 \tan(c+dx)^2}{d} + \frac{Aa^2 \tan(c+dx) 4i}{3d}}{\tan(c + dx)^{5/2}} - \frac{\frac{2Ba^2}{3d} + \frac{Ba^2 \tan(c+dx) 4i}{d}}{\tan(c + dx)^{3/2}}$$

$$+ \frac{\sqrt{2} A a^2 \ln\left(4 A a^2 d + \sqrt{2} A a^2 d \sqrt{\tan(c + dx)} (-2 - 2i)\right) (1 + 1i)}{d}$$

$$- \frac{\sqrt{4i} A a^2 \ln\left(4 A a^2 d + 2 \sqrt{4i} A a^2 d \sqrt{\tan(c + dx)}\right)}{d}$$

$$+ \frac{\sqrt{2} B a^2 \ln\left(-B a^2 d 4i + \sqrt{2} B a^2 d \sqrt{\tan(c + dx)} (-2 + 2i)\right) (1 - i)}{d}$$

$$- \frac{\sqrt{-4i} B a^2 \ln\left(-B a^2 d 4i + 2 \sqrt{-4i} B a^2 d \sqrt{\tan(c + dx)}\right)}{d}$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2)/tan(c + d*x)^(7/2),x)`

output $(2^{1/2}Aa^2 \log(4Aa^2d - 2^{1/2}Aa^2d \tan(c + dx)^{1/2}(2 + 2i))(1 + i)/d - ((2Ba^2)/(3d) + (Ba^2 \tan(c + dx)4i)/d)/\tan(c + dx)^{3/2} - ((2Aa^2)/(5d) + (Aa^2 \tan(c + dx)4i)/(3d) - (4Aa^2 \tan(c + dx)^2)/d)/\tan(c + dx)^{5/2} - (4i^{1/2}Aa^2 \log(4Aa^2d + 24i^{1/2}Aa^2d \tan(c + dx)^{1/2}))/d + (2^{1/2}Ba^2 \log(-Ba^2d4i - 2^{1/2}Ba^2d \tan(c + dx)^{1/2}(2 - 2i))(1 - i))/d - ((-4i)^{1/2}Ba^2 \log(2*(-4i)^{1/2}Ba^2d \tan(c + dx)^{1/2} - Ba^2d4i))/d$

3.125. $\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

3.126
$$\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

3.126.1 Optimal result 1311
 3.126.2 Mathematica [A] (verified) 1311
 3.126.3 Rubi [A] (verified) 1312
 3.126.4 Maple [A] (verified) 1316
 3.126.5 Fracas [B] (verification not implemented) 1317
 3.126.6 Sympy [F] 1318
 3.126.7 Maxima [A] (verification not implemented) 1318
 3.126.8 Giac [A] (verification not implemented) 1319
 3.126.9 Mupad [B] (verification not implemented) 1319

3.126.1 Optimal result

Integrand size = 36, antiderivative size = 154

$$\int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$= -\frac{4\sqrt[4]{-1}a^2(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d} - \frac{2a^2(9iA + 7B)}{35d \tan^{\frac{5}{2}}(c + dx)}$$

$$+ \frac{4a^2(A - iB)}{3d \tan^{\frac{3}{2}}(c + dx)} + \frac{4a^2(iA + B)}{d\sqrt{\tan(c + dx)}} - \frac{2A(a^2 + ia^2 \tan(c + dx))}{7d \tan^{\frac{7}{2}}(c + dx)}$$

output

```
-4*(-1)^(1/4)*a^2*(A-I*B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d+4*a^2*(I*A+B)/d/tan(d*x+c)^(1/2)-2/35*a^2*(9*I*A+7*B)/d/tan(d*x+c)^(5/2)+4/3*a^2*(A-I*B)/d/tan(d*x+c)^(3/2)-2/7*A*(a^2+I*a^2*tan(d*x+c))/d/tan(d*x+c)^(7/2)
```

3.126.2 Mathematica [A] (verified)

Time = 3.47 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.78

$$\int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$= \frac{2a^2\left(-15A + (-42iA - 21B) \tan(c + dx) + 70(A - iB) \tan^2(c + dx) + 210(iA + B) \tan^3(c + dx) + (105d \tan^{\frac{7}{2}}(c + dx))\right)}{105d \tan^{\frac{7}{2}}(c + dx)}$$

3.126.
$$\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

input `Integrate[((a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2),x]`

output `(2*a^2*(-15*A + ((-42*I)*A - 21*B)*Tan[c + d*x] + 70*(A - I*B)*Tan[c + d*x]^2 + 210*(I*A + B)*Tan[c + d*x]^3 + (105 - 105*I)*Sqrt[2]*(A - I*B)*ArcTanh[((1 + I)*Sqrt[Tan[c + d*x]])/Sqrt[2]]*Tan[c + d*x]^(7/2))/(105*d*Tan[c + d*x]^(7/2))`

3.126.3 Rubi [A] (verified)

Time = 0.92 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.05, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3042, 4076, 27, 3042, 4074, 27, 3042, 4012, 3042, 4012, 25, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan(c + dx)^{9/2}} dx$$

↓ 4076

$$\frac{2}{7} \int \frac{(i \tan(c + dx)a + a)(a(9iA + 7B) - a(5A - 7iB) \tan(c + dx))}{2 \tan^{\frac{7}{2}}(c + dx)} dx - \frac{2A(a^2 + ia^2 \tan(c + dx))}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{7} \int \frac{(i \tan(c + dx)a + a)(a(9iA + 7B) - a(5A - 7iB) \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx - \frac{2A(a^2 + ia^2 \tan(c + dx))}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \int \frac{(i \tan(c + dx)a + a)(a(9iA + 7B) - a(5A - 7iB) \tan(c + dx))}{\tan(c + dx)^{7/2}} dx - \frac{2A(a^2 + ia^2 \tan(c + dx))}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 4074

$$\begin{aligned}
& \frac{1}{7} \left(\int -\frac{14((A-iB)a^2 + (iA+B)\tan(c+dx)a^2)}{\tan^{\frac{5}{2}}(c+dx)} dx - \frac{2a^2(7B+9iA)}{5d\tan^{\frac{5}{2}}(c+dx)} \right) - \\
& \quad \frac{2A(a^2 + ia^2\tan(c+dx))}{7d\tan^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{7} \left(-14 \int \frac{(A-iB)a^2 + (iA+B)\tan(c+dx)a^2}{\tan^{\frac{5}{2}}(c+dx)} dx - \frac{2a^2(7B+9iA)}{5d\tan^{\frac{5}{2}}(c+dx)} \right) - \\
& \quad \frac{2A(a^2 + ia^2\tan(c+dx))}{7d\tan^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \left(-14 \int \frac{(A-iB)a^2 + (iA+B)\tan(c+dx)a^2}{\tan(c+dx)^{5/2}} dx - \frac{2a^2(7B+9iA)}{5d\tan^{\frac{5}{2}}(c+dx)} \right) - \\
& \quad \frac{2A(a^2 + ia^2\tan(c+dx))}{7d\tan^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow \text{4012} \\
& \frac{1}{7} \left(-14 \left(\int \frac{a^2(iA+B) - a^2(A-iB)\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)} dx - \frac{2a^2(A-iB)}{3d\tan^{\frac{3}{2}}(c+dx)} \right) - \frac{2a^2(7B+9iA)}{5d\tan^{\frac{5}{2}}(c+dx)} \right) - \\
& \quad \frac{2A(a^2 + ia^2\tan(c+dx))}{7d\tan^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \left(-14 \left(\int \frac{a^2(iA+B) - a^2(A-iB)\tan(c+dx)}{\tan(c+dx)^{3/2}} dx - \frac{2a^2(A-iB)}{3d\tan^{\frac{3}{2}}(c+dx)} \right) - \frac{2a^2(7B+9iA)}{5d\tan^{\frac{5}{2}}(c+dx)} \right) - \\
& \quad \frac{2A(a^2 + ia^2\tan(c+dx))}{7d\tan^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow \text{4012} \\
& \frac{1}{7} \left(-14 \left(\int -\frac{(A-iB)a^2 + (iA+B)\tan(c+dx)a^2}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(A-iB)}{3d\tan^{\frac{3}{2}}(c+dx)} - \frac{2a^2(B+iA)}{d\sqrt{\tan(c+dx)}} \right) - \frac{2a^2(7B+9iA)}{5d\tan^{\frac{5}{2}}(c+dx)} \right) - \\
& \quad \frac{2A(a^2 + ia^2\tan(c+dx))}{7d\tan^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow \text{25}
\end{aligned}$$

$$\frac{1}{7} \left(-14 \left(- \int \frac{(A - iB)a^2 + (iA + B) \tan(c + dx)a^2}{\sqrt{\tan(c + dx)}} dx - \frac{2a^2(A - iB)}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2a^2(B + iA)}{d\sqrt{\tan(c + dx)}} \right) - \frac{2a^2(7B + 9iA)}{5d \tan^{\frac{5}{2}}(c + dx)} \right) - \frac{2A(a^2 + ia^2 \tan(c + dx))}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \left(-14 \left(- \int \frac{(A - iB)a^2 + (iA + B) \tan(c + dx)a^2}{\sqrt{\tan(c + dx)}} dx - \frac{2a^2(A - iB)}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2a^2(B + iA)}{d\sqrt{\tan(c + dx)}} \right) - \frac{2a^2(7B + 9iA)}{5d \tan^{\frac{5}{2}}(c + dx)} \right) - \frac{2A(a^2 + ia^2 \tan(c + dx))}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 4016

$$\frac{1}{7} \left(-14 \left(- \frac{2a^4(A - iB)^2 \int \frac{1}{a^2(A - iB) - a^2(iA + B) \tan(c + dx)} d\sqrt{\tan(c + dx)}}{d} - \frac{2a^2(A - iB)}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2a^2(B + iA)}{d\sqrt{\tan(c + dx)}} \right) - \frac{2a^2(7B + 9iA)}{5d \tan^{\frac{5}{2}}(c + dx)} \right) - \frac{2A(a^2 + ia^2 \tan(c + dx))}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 218

$$\frac{1}{7} \left(-14 \left(\frac{2\sqrt[4]{-1}a^2(A - iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right)}{d} - \frac{2a^2(A - iB)}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2a^2(B + iA)}{d\sqrt{\tan(c + dx)}} \right) - \frac{2a^2(7B + 9iA)}{5d \tan^{\frac{5}{2}}(c + dx)} \right) - \frac{2A(a^2 + ia^2 \tan(c + dx))}{7d \tan^{\frac{7}{2}}(c + dx)}$$

input `Int[((a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2),x]`

output `(-14*((2*(-1)^(1/4)*a^2*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]])/d - (2*a^2*(A - I*B))/(3*d*Tan[c + d*x]^(3/2)) - (2*a^2*(I*A + B))/(d*Sqrt[Tan[c + d*x]])) - (2*a^2*((9*I)*A + 7*B))/(5*d*Tan[c + d*x]^(5/2))/7 - (2*A*(a^2 + I*a^2*Tan[c + d*x]))/(7*d*Tan[c + d*x]^(7/2))`

3.126.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`
- rule 4016 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`
- rule 4074 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

```
rule 4076 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m -
n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

3.126.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.68

method	result
derivativedivides	$a^2 \left(-\frac{2A}{7 \tan(dx+c)^{\frac{7}{2}}} - \frac{2(2iB-2A)}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(-2iA-2B)}{\sqrt{\tan(dx+c)}} - \frac{2(2iA+B)}{5 \tan(dx+c)^{\frac{5}{2}}} + \frac{(-2iB+2A)\sqrt{2}}{\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right)} + 2 \arctan\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) \right)$
default	$a^2 \left(-\frac{2A}{7 \tan(dx+c)^{\frac{7}{2}}} - \frac{2(2iB-2A)}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(-2iA-2B)}{\sqrt{\tan(dx+c)}} - \frac{2(2iA+B)}{5 \tan(dx+c)^{\frac{5}{2}}} + \frac{(-2iB+2A)\sqrt{2}}{\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right)} + 2 \arctan\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) \right)$
parts	$\frac{(2iAa^2 + Ba^2) \left(\frac{\sqrt{2} \left(\ln\left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})\right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})\right))}{4} \right)}{d}$

```
input int((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x,method=_RETUR
NVERBOSE)
```

```
output 1/d*a^2*(-2/7*A/tan(d*x+c)^(7/2)-2/3*(-2*A+2*I*B)/tan(d*x+c)^(3/2)-2*(-2*B
-2*I*A)/tan(d*x+c)^(1/2)-2/5*(2*I*A+B)/tan(d*x+c)^(5/2)+1/4*(2*A-2*I*B)*2^
(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1
/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*
tan(d*x+c)^(1/2)))+1/4*(2*B+2*I*A)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)
+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*t
an(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))
```

3.126.
$$\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

3.126.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 561 vs. $2(124) = 248$.

Time = 0.28 (sec) , antiderivative size = 561, normalized size of antiderivative = 3.64

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$= 105 \sqrt{-\frac{(iA^2 + 2AB - iB^2)a^4}{d^2}} (de^{(8i dx + 8i c)} - 4de^{(6i dx + 6i c)} + 6de^{(4i dx + 4i c)} - 4de^{(2i dx + 2i c)} + d) \log \left(-\frac{2((A - iB)^2}{\dots} \right)$$

```
input integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="fricas")
```

```
output 1/105*(105*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)*log(-2*((A - I*B)*a^2*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^2) - 105*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)*log(-2*((A - I*B)*a^2*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^2) - 2*((337*A - 301*I*B)*a^2*e^(8*I*d*x + 8*I*c) - 6*(46*A - 63*I*B)*a^2*e^(6*I*d*x + 6*I*c) - 10*(5*A - 14*I*B)*a^2*e^(4*I*d*x + 4*I*c) + 18*(22*A - 21*I*B)*a^2*e^(2*I*d*x + 2*I*c) - (167*A - 161*I*B)*a^2)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)
```

3.126.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$= -a^2 \left(\int \left(-\frac{A}{\tan^{\frac{9}{2}}(c + dx)} \right) dx + \int \frac{A}{\tan^{\frac{5}{2}}(c + dx)} dx + \int \left(-\frac{B}{\tan^{\frac{7}{2}}(c + dx)} \right) dx \right.$$

$$\left. + \int \frac{B}{\tan^{\frac{3}{2}}(c + dx)} dx + \int \left(-\frac{2iA}{\tan^{\frac{7}{2}}(c + dx)} \right) dx + \int \left(-\frac{2iB}{\tan^{\frac{5}{2}}(c + dx)} \right) dx \right)$$

input `integrate((a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c))/tan(d*x+c)**(9/2),x)`

output `-a**2*(Integral(-A/tan(c + d*x)**(9/2), x) + Integral(A/tan(c + d*x)**(5/2), x) + Integral(-B/tan(c + d*x)**(7/2), x) + Integral(B/tan(c + d*x)**(3/2), x) + Integral(-2*I*A/tan(c + d*x)**(7/2), x) + Integral(-2*I*B/tan(c + d*x)**(5/2), x))`

3.126.7 Maxima [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.38

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx =$$

$$\frac{105 \left(2\sqrt{2}(-i+1)A + (i-1)B \right) \arctan \left(\frac{1}{2}\sqrt{2} \left(\sqrt{2} + 2\sqrt{\tan(dx+c)} \right) \right) + 2\sqrt{2}(-i+1)A + (i-1)B}{\dots}$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="maxima")`

output `-1/210*(105*(2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a^2 - 4*(210*(I*A + B)*a^2*tan(d*x + c)^3 + 70*(A - I*B)*a^2*tan(d*x + c)^2 + 21*(-2*I*A - B)*a^2*tan(d*x + c) - 15*A*a^2)/tan(d*x + c)^(7/2)/d`

3.126. $\int \frac{(a+ia \tan(c+dx))^2 (A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

3.126.8 Giac [A] (verification not implemented)

Time = 1.14 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.88

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$= \frac{(2i - 2) \sqrt{2} (-i A a^2 - B a^2) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{d} - \frac{2(-210i A a^2 \tan(dx + c)^3 - 210 B a^2 \tan(dx + c)^3 - 70 A a^2 \tan(dx + c)^2 + 70i B a^2 \tan(dx + c)^2 + 42i A a^2 \tan(dx + c) + 21 B a^2 \tan(dx + c) + 15 A a^2)}{105 d \tan(dx + c)^{\frac{7}{2}}}$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="giac")`

output `(2*I - 2)*sqrt(2)*(-I*A*a^2 - B*a^2)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d - 2/105*(-210*I*A*a^2*tan(d*x + c)^3 - 210*B*a^2*tan(d*x + c)^3 - 70*A*a^2*tan(d*x + c)^2 + 70*I*B*a^2*tan(d*x + c)^2 + 42*I*A*a^2*tan(d*x + c) + 21*B*a^2*tan(d*x + c) + 15*A*a^2)/(d*tan(d*x + c)^(7/2))`

3.126.9 Mupad [B] (verification not implemented)

Time = 10.92 (sec) , antiderivative size = 293, normalized size of antiderivative = 1.90

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$= -\frac{\frac{2 A a^2}{7 d} + \frac{A a^2 \tan(c+dx) 4i}{5 d} - \frac{4 A a^2 \tan(c+dx)^2}{3 d} - \frac{A a^2 \tan(c+dx)^3 4i}{d}}{\tan(c + dx)^{7/2}} - \frac{\frac{2 B a^2}{5 d} - \frac{4 B a^2 \tan(c+dx)^2}{d} + \frac{B a^2 \tan(c+dx) 4i}{3 d}}{\tan(c + dx)^{5/2}}$$

$$+ \frac{\sqrt{2} A a^2 \ln\left(A a^2 d 4i + \sqrt{2} A a^2 d \sqrt{\tan(c + dx)} (-2 + 2i)\right) (1 - i)}{d}$$

$$- \frac{\sqrt{-4i} A a^2 \ln\left(A a^2 d 4i + 2 \sqrt{-4i} A a^2 d \sqrt{\tan(c + dx)}\right)}{d}$$

$$+ \frac{\sqrt{2} B a^2 \ln\left(4 B a^2 d + \sqrt{2} B a^2 d \sqrt{\tan(c + dx)} (-2 - 2i)\right) (1 + i)}{d}$$

$$- \frac{\sqrt{4i} B a^2 \ln\left(4 B a^2 d + 2 \sqrt{4i} B a^2 d \sqrt{\tan(c + dx)}\right)}{d}$$

3.126. $\int \frac{(a+ia \tan(c+dx))^2 (A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2)/tan(c + d*x)^(9/2),x)`

output $(2^{(1/2)}*A*a^2*\log(A*a^2*d*4i - 2^{(1/2)}*A*a^2*d*\tan(c + d*x)^{(1/2)}*(2 - 2i))*(1 - 1i))/d - ((2*B*a^2)/(5*d) + (B*a^2*\tan(c + d*x)*4i)/(3*d) - (4*B*a^2*\tan(c + d*x)^2)/d)/\tan(c + d*x)^{(5/2)} - ((2*A*a^2)/(7*d) + (A*a^2*\tan(c + d*x)*4i)/(5*d) - (4*A*a^2*\tan(c + d*x)^2)/(3*d) - (A*a^2*\tan(c + d*x)^3*4i)/d)/\tan(c + d*x)^{(7/2)} - ((-4i)^{(1/2)}*A*a^2*\log(A*a^2*d*4i + 2*(-4i)^{(1/2)}*A*a^2*d*\tan(c + d*x)^{(1/2)}))/d + (2^{(1/2)}*B*a^2*\log(4*B*a^2*d - 2^{(1/2)}*B*a^2*d*\tan(c + d*x)^{(1/2)}*(2 + 2i))*(1 + 1i))/d - (4i^{(1/2)}*B*a^2*\log(4*B*a^2*d + 2*4i^{(1/2)}*B*a^2*d*\tan(c + d*x)^{(1/2)}))/d$

3.127 $\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.127.1 Optimal result	1321
3.127.2 Mathematica [A] (verified)	1322
3.127.3 Rubi [A] (verified)	1322
3.127.4 Maple [A] (verified)	1326
3.127.5 Fricas [B] (verification not implemented)	1327
3.127.6 Sympy [F]	1328
3.127.7 Maxima [A] (verification not implemented)	1329
3.127.8 Giac [A] (verification not implemented)	1329
3.127.9 Mupad [B] (verification not implemented)	1330

3.127.1 Optimal result

Integrand size = 36, antiderivative size = 198

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{8\sqrt{-1}a^3(A-iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c+dx)}\right)}{d} + \frac{8a^3(A-iB)\sqrt{\tan(c+dx)}}{d} + \frac{8a^3(iA+B) \tan^{\frac{3}{2}}(c+dx)}{3d}$$

$$- \frac{16a^3(18A-19iB) \tan^{\frac{5}{2}}(c+dx)}{315d} + \frac{2iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2}{9d}$$

$$- \frac{2(9A-13iB) \tan^{\frac{5}{2}}(c+dx)(a^3+ia^3 \tan(c+dx))}{63d}$$

output

```
8*(-1)^(1/4)*a^3*(A-I*B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d+8*a^3*(A-I*
B)*tan(d*x+c)^(1/2)/d+8/3*a^3*(I*A+B)*tan(d*x+c)^(3/2)/d-16/315*a^3*(18*A-
19*I*B)*tan(d*x+c)^(5/2)/d+2/9*I*a*B*tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^2
/d-2/63*(9*A-13*I*B)*tan(d*x+c)^(5/2)*(a^3+I*a^3*tan(d*x+c))/d
```


3.127.2 Mathematica [A] (verified)

Time = 3.18 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.65

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{2a^3 \left(1260 \sqrt[4]{-1} (A-iB) \arctan \left((-1)^{3/4} \sqrt{\tan(c+dx)} \right) + \sqrt{\tan(c+dx)} (1260(A-iB) + 420(iA+B) \tan(c+dx)) \right)}{315d}$$

input `Integrate[Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(2*a^3*(1260*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + Sqrt[Tan[c + d*x]]*(1260*(A - I*B) + 420*(I*A + B)*Tan[c + d*x] - 63*(3*A - (4*I)*B)*Tan[c + d*x]^2 - (45*I)*(A - (3*I)*B)*Tan[c + d*x]^3 - (35*I)*B*Tan[c + d*x]^4))/(315*d)`

3.127.3 Rubi [A] (verified)

Time = 1.18 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.04, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3042, 4077, 27, 3042, 4077, 27, 3042, 4075, 3042, 4011, 3042, 4011, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c+dx)^{3/2}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4077}$$

$$\frac{2}{9} \int \frac{1}{2} \tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a)^2(a(9A-5iB)+a(9iA+13B) \tan(c+dx)) dx +$$

$$\frac{2iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2}{9d}$$

$$\downarrow \text{27}$$

3.127. $\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\frac{1}{9} \int \tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a)^2(a(9A-5iB)+a(9iA+13B)\tan(c+dx))dx + \frac{2iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2}{9d}$$

↓ 3042

$$\frac{1}{9} \int \tan(c+dx)^{3/2}(i \tan(c+dx)a+a)^2(a(9A-5iB)+a(9iA+13B)\tan(c+dx))dx + \frac{2iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2}{9d}$$

↓ 4077

$$\frac{1}{9} \left(\frac{2}{7} \int 2 \tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a) ((27A-25iB)a^2+2(18iA+19B)\tan(c+dx)a^2) dx - \frac{2(9A-13iB)}{9d} \right) + \frac{2iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2}{9d}$$

↓ 27

$$\frac{1}{9} \left(\frac{4}{7} \int \tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a) ((27A-25iB)a^2+2(18iA+19B)\tan(c+dx)a^2) dx - \frac{2(9A-13iB)}{9d} \right) + \frac{2iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2}{9d}$$

↓ 3042

$$\frac{1}{9} \left(\frac{4}{7} \int \tan(c+dx)^{3/2}(i \tan(c+dx)a+a) ((27A-25iB)a^2+2(18iA+19B)\tan(c+dx)a^2) dx - \frac{2(9A-13iB)}{9d} \right) + \frac{2iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2}{9d}$$

↓ 4075

$$\frac{1}{9} \left(\frac{4}{7} \left(\int \tan^{\frac{3}{2}}(c+dx) (63(A-iB)a^3+63(iA+B)\tan(c+dx)a^3) dx - \frac{4a^3(18A-19iB)\tan^{\frac{5}{2}}(c+dx)}{5d} \right) - \frac{2(9A-13iB)}{9d} \right) + \frac{2iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2}{9d}$$

↓ 3042

3.127. $\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\frac{1}{9} \left(\frac{4}{7} \left(\int \tan(c+dx)^{3/2} (63(A-iB)a^3 + 63(iA+B)\tan(c+dx)a^3) dx - \frac{4a^3(18A-19iB)\tan^{5/2}(c+dx)}{5d} \right) - \frac{2iaB \tan^{5/2}(c+dx)(a+ia \tan(c+dx))^2}{9d} \right) - 2$$

↓ 4011

$$\frac{1}{9} \left(\frac{4}{7} \left(\int \sqrt{\tan(c+dx)} (63a^3(A-iB)\tan(c+dx) - 63a^3(iA+B)) dx - \frac{4a^3(18A-19iB)\tan^{5/2}(c+dx)}{5d} + \frac{42a^3}{d} \right) - \frac{2iaB \tan^{5/2}(c+dx)(a+ia \tan(c+dx))^2}{9d} \right) - 2$$

↓ 3042

$$\frac{1}{9} \left(\frac{4}{7} \left(\int \sqrt{\tan(c+dx)} (63a^3(A-iB)\tan(c+dx) - 63a^3(iA+B)) dx - \frac{4a^3(18A-19iB)\tan^{5/2}(c+dx)}{5d} + \frac{42a^3}{d} \right) - \frac{2iaB \tan^{5/2}(c+dx)(a+ia \tan(c+dx))^2}{9d} \right) - 2$$

↓ 4011

$$\frac{1}{9} \left(\frac{4}{7} \left(\int \frac{-63(A-iB)a^3 - 63(iA+B)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx - \frac{4a^3(18A-19iB)\tan^{5/2}(c+dx)}{5d} + \frac{42a^3(B+iA)\tan^{3/2}(c+dx)}{d} \right) - \frac{2iaB \tan^{5/2}(c+dx)(a+ia \tan(c+dx))^2}{9d} \right) - 2$$

↓ 3042

$$\frac{1}{9} \left(\frac{4}{7} \left(\int \frac{-63(A-iB)a^3 - 63(iA+B)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx - \frac{4a^3(18A-19iB)\tan^{5/2}(c+dx)}{5d} + \frac{42a^3(B+iA)\tan^{3/2}(c+dx)}{d} \right) - \frac{2iaB \tan^{5/2}(c+dx)(a+ia \tan(c+dx))^2}{9d} \right) - 2$$

↓ 4016

$$\frac{1}{9} \left(\frac{4}{7} \left(\frac{7938a^6(A-iB)^2 \int \frac{1}{63a^3(iA+B)\tan(c+dx) - 63a^3(A-iB)} d\sqrt{\tan(c+dx)}}{d} - \frac{4a^3(18A-19iB)\tan^{5/2}(c+dx)}{5d} + \frac{42a^3}{d} \right) - \frac{2iaB \tan^{5/2}(c+dx)(a+ia \tan(c+dx))^2}{9d} \right) - 2$$

↓ 218

3.127. $\int \tan^{3/2}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\frac{1}{9} \left(\frac{4}{7} \left(\frac{126 \sqrt[4]{-1} a^3 (A - iB) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right)}{d} - \frac{4a^3 (18A - 19iB) \tan^{5/2}(c + dx)}{5d} + \frac{42a^3 (B + iA) \tan^{5/2}(c + dx)}{d} \right) + \frac{2iaB \tan^{5/2}(c + dx) (a + ia \tan(c + dx))^2}{9d} \right)$$

input `Int[Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `((2*I)/9)*a*B*Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^2/d + ((-2*(9*A - (13*I)*B)*Tan[c + d*x]^(5/2)*(a^3 + I*a^3*Tan[c + d*x]))/(7*d) + (4*((126*(-1)^(1/4)*a^3*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]])/d + (126*a^3*(A - I*B)*Sqrt[Tan[c + d*x]])/d + (42*a^3*(I*A + B)*Tan[c + d*x]^(3/2))/d - (4*a^3*(18*A - (19*I)*B)*Tan[c + d*x]^(5/2))/(5*d))/7)/9`

3.127.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x_, x], x] /; FreeQ[a, x] && !MatchQ[F_x_, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4077 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.127.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.51

method	result
derivativedivides	$a^3 \left(-\frac{2iB \left(\tan^{\frac{9}{2}}(dx+c) \right)}{9} - \frac{2iA \left(\tan^{\frac{7}{2}}(dx+c) \right)}{7} - \frac{6B \left(\tan^{\frac{7}{2}}(dx+c) \right)}{7} + \frac{8iB \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} - \frac{6A \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} + \frac{8iA \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} \right)$
default	$a^3 \left(-\frac{2iB \left(\tan^{\frac{9}{2}}(dx+c) \right)}{9} - \frac{2iA \left(\tan^{\frac{7}{2}}(dx+c) \right)}{7} - \frac{6B \left(\tan^{\frac{7}{2}}(dx+c) \right)}{7} + \frac{8iB \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} - \frac{6A \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} + \frac{8iA \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} \right)$
parts	$\frac{(-iA a^3 - 3B a^3) \left(\frac{2 \left(\tan^{\frac{7}{2}}(dx+c) \right)}{7} - \frac{2 \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + \frac{\sqrt{2} \left(\ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))) \right)}{4} \right)}{d}$

input `int(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

$$3.127. \quad \int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

```
output 1/d*a^3*(-2/9*I*B*tan(d*x+c)^(9/2)-2/7*I*A*tan(d*x+c)^(7/2)-6/7*B*tan(d*x+c)^(7/2)+8/5*I*B*tan(d*x+c)^(5/2)-6/5*A*tan(d*x+c)^(5/2)+8/3*I*A*tan(d*x+c)^(3/2)+8/3*B*tan(d*x+c)^(3/2)-8*I*B*tan(d*x+c)^(1/2)+8*A*tan(d*x+c)^(1/2)+1/4*(4*I*B-4*A)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))+1/4*(-4*I*A-4*B)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))
```

3.127.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 561 vs. $2(158) = 316$.

Time = 0.32 (sec) , antiderivative size = 561, normalized size of antiderivative = 2.83

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx =$$

$$2 \left(315 \sqrt{-\frac{(iA^2+2AB-iB^2)a^6}{d^2}} (de^{(8i dx+8i c)} + 4 de^{(6i dx+6i c)} + 6 de^{(4i dx+4i c)} + 4 de^{(2i dx+2i c)} + d) \log \left(-\frac{2}{\dots} \right) \right)$$

```
input integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fracas")
```

```
output -2/315*(315*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(d*e^(8*I*d*x + 8*I*c)
+ 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*x + 2*I
*c) + d)*log(-2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B
- I*B^2)*a^6/d^2)*(I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt((-I*e^(2*I*d*x + 2*
I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3
)) - 315*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(d*e^(8*I*d*x + 8*I*c) + 4
*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*x + 2*I*c)
+ d)*log(-2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I
*B^2)*a^6/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt((-I*e^(2*I*d*x + 2*I*
c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3))
- 2*((957*A - 1051*I*B)*a^3*e^(8*I*d*x + 8*I*c) + 5*(579*A - 547*I*B)*a^3
*e^(6*I*d*x + 6*I*c) + 21*(171*A - 173*I*B)*a^3*e^(4*I*d*x + 4*I*c) + 5*(4
29*A - 433*I*B)*a^3*e^(2*I*d*x + 2*I*c) + 4*(123*A - 124*I*B)*a^3)*sqrt((-
I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(8*I*d*x + 8*I
*c) + 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*x +
2*I*c) + d)
```

3.127.6 Sympy [F]

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= -ia^3 \left(\int (-3A \tan^{\frac{5}{2}}(c + dx)) dx + \int A \tan^{\frac{9}{2}}(c + dx) dx + \int (-3B \tan^{\frac{7}{2}}(c + dx)) dx \right.$$

$$+ \int B \tan^{\frac{11}{2}}(c + dx) dx + \int iA \tan^{\frac{3}{2}}(c + dx) dx + \int (-3iA \tan^{\frac{7}{2}}(c + dx)) dx$$

$$\left. + \int iB \tan^{\frac{5}{2}}(c + dx) dx + \int (-3iB \tan^{\frac{9}{2}}(c + dx)) dx \right)$$

```
input integrate(tan(d*x+c)**(3/2)*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)
```

```
output -I*a**3*(Integral(-3*A*tan(c + d*x)**(5/2), x) + Integral(A*tan(c + d*x)**
(9/2), x) + Integral(-3*B*tan(c + d*x)**(7/2), x) + Integral(B*tan(c + d*x
)**(11/2), x) + Integral(I*A*tan(c + d*x)**(3/2), x) + Integral(-3*I*A*tan
(c + d*x)**(7/2), x) + Integral(I*B*tan(c + d*x)**(5/2), x) + Integral(-3*
I*B*tan(c + d*x)**(9/2), x))
```

3.127.7 Maxima [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.18

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx =$$

$$70i Ba^3 \tan(dx+c)^{\frac{9}{2}} + 90(iA+3B)a^3 \tan(dx+c)^{\frac{7}{2}} + 126(3A-4iB)a^3 \tan(dx+c)^{\frac{5}{2}} + 840(-iA$$

```
input integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")
```

```
output -1/315*(70*I*B*a^3*tan(d*x+c)^(9/2)+90*(I*A+3*B)*a^3*tan(d*x+c)^(7/2)+126*(3*A-4*I*B)*a^3*tan(d*x+c)^(5/2)+840*(-I*A-B)*a^3*tan(d*x+c)^(3/2)-2520*(A-I*B)*a^3*sqrt(tan(d*x+c))+315*(2*sqrt(2)*((I+1)*A-(I-1)*B)*arctan(1/2*sqrt(2)*(sqrt(2)+2*sqrt(tan(d*x+c))))+2*sqrt(2)*((I+1)*A-(I-1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2)-2*sqrt(tan(d*x+c))))-sqrt(2)*((I-1)*A+(I+1)*B)*log(sqrt(2)*sqrt(tan(d*x+c))+tan(d*x+c)+1)+sqrt(2)*((I-1)*A+(I+1)*B)*log(-sqrt(2)*sqrt(tan(d*x+c))+tan(d*x+c)+1))*a^3)/d
```

3.127.8 Giac [A] (verification not implemented)

Time = 1.11 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.98

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{(4i-4)\sqrt{2}(iAa^3+Ba^3)\arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{d}$$

$$2\left(35iBa^3d^8 \tan(dx+c)^{\frac{9}{2}}+45iAa^3d^8 \tan(dx+c)^{\frac{7}{2}}+135Ba^3d^8 \tan(dx+c)^{\frac{5}{2}}+189Aa^3d^8 \tan(dx+c)^{\frac{3}{2}}\right)$$

```
input integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")
```


output $(4I - 4)\sqrt{2}(IA^3 + B^3)\arctan(-(1/2I - 1/2)\sqrt{2}\sqrt{\tan(dx + c)})/d - 2/315(35IB^3d^8\tan(dx + c)^{9/2} + 45IA^3d^8\tan(dx + c)^{7/2} + 135B^3d^8\tan(dx + c)^{7/2} + 189A^3d^8\tan(dx + c)^{5/2} - 252IB^3d^8\tan(dx + c)^{5/2} - 420IA^3d^8\tan(dx + c)^{3/2} - 420B^3d^8\tan(dx + c)^{3/2} - 1260A^3d^8\sqrt{\tan(dx + c)} + 1260IB^3d^8\sqrt{\tan(dx + c)})/d^9$

3.127.9 Mupad [B] (verification not implemented)

Time = 13.58 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.65

$$\begin{aligned} & \int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ &= \frac{8Aa^3\sqrt{\tan(c + dx)}}{d} + \frac{Aa^3\tan(c + dx)^{3/2}8i}{3d} - \frac{6Aa^3\tan(c + dx)^{5/2}}{5d} \\ & - \frac{Aa^3\tan(c + dx)^{7/2}2i}{7d} - \frac{Ba^3\sqrt{\tan(c + dx)}8i}{d} + \frac{8Ba^3\tan(c + dx)^{3/2}}{3d} \\ & + \frac{Ba^3\tan(c + dx)^{5/2}8i}{5d} - \frac{6Ba^3\tan(c + dx)^{7/2}}{7d} - \frac{Ba^3\tan(c + dx)^{9/2}2i}{9d} \\ & + \frac{\sqrt{2}Aa^3\ln\left(-Aa^3d8i + \sqrt{2}Aa^3d\sqrt{\tan(c + dx)}(-4 + 4i)\right)(2 - 2i)}{d} \\ & - \frac{\sqrt{-16i}Aa^3\ln\left(-Aa^3d8i + 2\sqrt{-16i}Aa^3d\sqrt{\tan(c + dx)}\right)}{d} \\ & + \frac{\sqrt{2}Ba^3\ln\left(-8Ba^3d + \sqrt{2}Ba^3d\sqrt{\tan(c + dx)}(-4 - 4i)\right)(2 + 2i)}{d} \\ & - \frac{\sqrt{16i}Ba^3\ln\left(-8Ba^3d + 2\sqrt{16i}Ba^3d\sqrt{\tan(c + dx)}\right)}{d} \end{aligned}$$

input `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3,x)`

output $(8Aa^3 \tan(c + dx)^{1/2})/d + (Aa^3 \tan(c + dx)^{3/2} 8i)/(3d) - (6Aa^3 \tan(c + dx)^{5/2})/(5d) - (Aa^3 \tan(c + dx)^{7/2} 2i)/(7d) - (Ba^3 \tan(c + dx)^{1/2} 8i)/d + (8Ba^3 \tan(c + dx)^{3/2})/(3d) + (Ba^3 \tan(c + dx)^{5/2} 8i)/(5d) - (6Ba^3 \tan(c + dx)^{7/2})/(7d) - (Ba^3 \tan(c + dx)^{9/2} 2i)/(9d) + (2^{1/2} Aa^3 \log(-Aa^3 d 8i - 2^{1/2}) Aa^3 d \tan(c + dx)^{1/2} (4 - 4i)) (2 - 2i)/d - ((-16i)^{1/2} Aa^3 \log(2(-16i)^{1/2} Aa^3 d \tan(c + dx)^{1/2} - Aa^3 d 8i))/d + (2^{1/2} Ba^3 \log(-8Ba^3 d - 2^{1/2} Ba^3 d \tan(c + dx)^{1/2} (4 + 4i)) (2 + 2i))/d - (16i)^{1/2} Ba^3 \log(2 \cdot 16i^{1/2} Ba^3 d \tan(c + dx)^{1/2} - 8Ba^3 d))/d$

3.128 $\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.128.1 Optimal result	1332
3.128.2 Mathematica [A] (verified)	1333
3.128.3 Rubi [A] (verified)	1333
3.128.4 Maple [A] (verified)	1337
3.128.5 Fricas [B] (verification not implemented)	1338
3.128.6 Sympy [F]	1339
3.128.7 Maxima [A] (verification not implemented)	1339
3.128.8 Giac [A] (verification not implemented)	1340
3.128.9 Mupad [B] (verification not implemented)	1340

3.128.1 Optimal result

Integrand size = 36, antiderivative size = 171

$$\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{8\sqrt{-1}a^3(iA + B) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right)}{d} + \frac{8a^3(iA + B)\sqrt{\tan(c + dx)}}{d}$$

$$- \frac{8a^3(21A - 23iB) \tan^{3/2}(c + dx)}{105d} + \frac{2iaB \tan^{3/2}(c + dx)(a + ia \tan(c + dx))^2}{7d}$$

$$- \frac{2(7A - 11iB) \tan^{3/2}(c + dx)(a^3 + ia^3 \tan(c + dx))}{35d}$$

output

```
8*(-1)^(1/4)*a^3*(I*A+B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d+8*a^3*(I*A+B)*tan(d*x+c)^(1/2)/d-8/105*a^3*(21*A-23*I*B)*tan(d*x+c)^(3/2)/d+2/7*I*a*B*tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^2/d-2/35*(7*A-11*I*B)*tan(d*x+c)^(3/2)*(a^3+I*a^3*tan(d*x+c))/d
```

3.128.2 Mathematica [A] (verified)

Time = 2.05 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.65

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{2a^3 \left(420\sqrt[4]{-1}(iA+B) \arctan \left((-1)^{3/4} \sqrt{\tan(c+dx)} \right) + \sqrt{\tan(c+dx)}(420(iA+B) - 35(3A-4iB) \tan(c+dx)) \right)}{105d}$$

input `Integrate[Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(2*a^3*(420*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + Sqrt[Tan[c + d*x]]*(420*(I*A + B) - 35*(3*A - (4*I)*B)*Tan[c + d*x] - (21*I)*(A - (3*I)*B)*Tan[c + d*x]^2 - (15*I)*B*Tan[c + d*x]^3))/(105*d)`

3.128.3 Rubi [A] (verified)

Time = 0.99 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.06, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.361$, Rules used = {3042, 4077, 27, 3042, 4077, 27, 3042, 4075, 3042, 4011, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

↓ 3042

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

↓ 4077

$$\frac{2}{7} \int \frac{1}{2} \sqrt{\tan(c+dx)}(i \tan(c+dx)a + a)^2(a(7A-3iB) + a(7iA+11B) \tan(c+dx)) dx + \frac{2iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2}{7d}$$

↓ 27

$$\frac{1}{7} \int \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^2(a(7A-3iB)+a(7iA+11B)\tan(c+dx))dx + \frac{2iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2}{7d}$$

↓ 3042

$$\frac{1}{7} \int \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^2(a(7A-3iB)+a(7iA+11B)\tan(c+dx))dx + \frac{2iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2}{7d}$$

↓ 4077

$$\frac{1}{7} \left(\frac{2}{5} \int 2\sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)(2(7A-6iB)a^2+(21iA+23B)\tan(c+dx)a^2)dx - \frac{2(7A-11iB)t}{7d} \right) + \frac{2iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2}{7d}$$

↓ 27

$$\frac{1}{7} \left(\frac{4}{5} \int \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)(2(7A-6iB)a^2+(21iA+23B)\tan(c+dx)a^2)dx - \frac{2(7A-11iB)t}{7d} \right) + \frac{2iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2}{7d}$$

↓ 3042

$$\frac{1}{7} \left(\frac{4}{5} \int \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)(2(7A-6iB)a^2+(21iA+23B)\tan(c+dx)a^2)dx - \frac{2(7A-11iB)t}{7d} \right) + \frac{2iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2}{7d}$$

↓ 4075

$$\frac{1}{7} \left(\frac{4}{5} \left(\int \sqrt{\tan(c+dx)}(35(A-iB)a^3+35(iA+B)\tan(c+dx)a^3)dx - \frac{2a^3(21A-23iB)\tan^{\frac{3}{2}}(c+dx)}{3d} \right) - \frac{2(7A-11iB)t}{7d} \right) + \frac{2iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2}{7d}$$

↓ 3042

$$\frac{1}{7} \left(\frac{4}{5} \left(\int \sqrt{\tan(c+dx)} (35(A-iB)a^3 + 35(iA+B)\tan(c+dx)a^3) dx - \frac{2a^3(21A-23iB)\tan^{\frac{3}{2}}(c+dx)}{3d} \right) - \frac{2iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2}{7d} \right) \downarrow 4011$$

$$\frac{1}{7} \left(\frac{4}{5} \left(\int \frac{35a^3(A-iB)\tan(c+dx) - 35a^3(iA+B)}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(21A-23iB)\tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{70a^3(B+iA)\sqrt{\tan(c+dx)}}{d} \right) - \frac{2iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2}{7d} \right) \downarrow 3042$$

$$\frac{1}{7} \left(\frac{4}{5} \left(\int \frac{35a^3(A-iB)\tan(c+dx) - 35a^3(iA+B)}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(21A-23iB)\tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{70a^3(B+iA)\sqrt{\tan(c+dx)}}{d} \right) - \frac{2iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2}{7d} \right) \downarrow 4016$$

$$\frac{1}{7} \left(\frac{4}{5} \left(\frac{2450a^6(B+iA)^2 \int \frac{1}{-35(iA+B)a^3 - 35(A-iB)\tan(c+dx)a^3} d\sqrt{\tan(c+dx)}}{d} - \frac{2a^3(21A-23iB)\tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{70a^3(B+iA)\sqrt{\tan(c+dx)}}{d} \right) - \frac{2iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2}{7d} \right) \downarrow 218$$

$$\frac{1}{7} \left(\frac{4}{5} \left(\frac{70\sqrt[4]{-1}a^3(B+iA) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{d} - \frac{2a^3(21A-23iB)\tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{70a^3(B+iA)\sqrt{\tan(c+dx)}}{d} \right) - \frac{2iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2}{7d} \right)$$

input `Int[Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

```
output ((2*I)/7)*a*B*Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^2/d + ((-2*(7*A
- (11*I)*B)*Tan[c + d*x]^(3/2)*(a^3 + I*a^3*Tan[c + d*x]))/(5*d) + (4*((70
*(-1)^(1/4)*a^3*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]])/d + (70*a
^3*(I*A + B)*Sqrt[Tan[c + d*x]]/d - (2*a^3*(21*A - (23*I)*B)*Tan[c + d*x]
^(3/2))/(3*d))/5)/7
```

3.128.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4011 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

```
rule 4016 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_
)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b
*Tan[e + f*x]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]
```

```
rule 4075 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_)
+ (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

rule 4077 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.128.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 276, normalized size of antiderivative = 1.61

method	result
derivativedivides	$a^3 \left(-\frac{2iB \left(\tan^{\frac{7}{2}}(dx+c) \right)}{7} - \frac{2iA \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} - \frac{6B \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} + \frac{8iB \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 2A \left(\tan^{\frac{3}{2}}(dx+c) \right) + 8iA \left(\sqrt{\tan(dx+c)} \right) \right)$
default	$a^3 \left(-\frac{2iB \left(\tan^{\frac{7}{2}}(dx+c) \right)}{7} - \frac{2iA \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} - \frac{6B \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} + \frac{8iB \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 2A \left(\tan^{\frac{3}{2}}(dx+c) \right) + 8iA \left(\sqrt{\tan(dx+c)} \right) \right)$
parts	$\frac{(-iA a^3 - 3B a^3) \left(\frac{2 \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} - 2 \left(\sqrt{\tan(dx+c)} \right) + \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)}{1-\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)} \right) + 2 \arctan \left(1+\sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right) \right)}{4} \right)}{d}$

input `int(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*a^3*(-2/7*I*B*tan(d*x+c)^(7/2)-2/5*I*A*tan(d*x+c)^(5/2)-6/5*B*tan(d*x+c)^(5/2)+8/3*I*B*tan(d*x+c)^(3/2)-2*A*tan(d*x+c)^(3/2)+8*I*A*tan(d*x+c)^(1/2)+8*B*tan(d*x+c)^(1/2)+1/4*(-4*I*A-4*B)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(-4*I*B+4*A)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))))`

3.128.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 498 vs. $2(137) = 274$.

Time = 0.27 (sec) , antiderivative size = 498, normalized size of antiderivative = 2.91

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx =$$

$$\frac{2 \left(105 \sqrt{-\frac{(-iA^2-2AB+iB^2)a^6}{d^2}} (de^{(6i dx+6i c)} + 3de^{(4i dx+4i c)} + 3de^{(2i dx+2i c)} + d) \log \left(-\frac{2 \left((A-iB)a^3e^{(2i dx+2i c)} \right)}{\dots} \right) \right)}{\dots}$$

```
input integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fracas")
```

```
output -2/105*(105*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2)*(d*e^(6*I*d*x + 6*I*c)
+ 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*log(-2*((A - I*B)
)*a^3*e^(2*I*d*x + 2*I*c) + sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2)*(d*e^(
2*I*d*x + 2*I*c) + d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*
c) + 1)))e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3)) - 105*sqrt(-(-I*A^2 - 2*A
*B + I*B^2)*a^6/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*
d*e^(2*I*d*x + 2*I*c) + d)*log(-2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) - sqr
t(-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt((-I*
e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-2*I*d*x - 2*I*c)/
((-I*A - B)*a^3)) + 2*((-273*I*A - 319*B)*a^3*e^(6*I*d*x + 6*I*c) + 2*(-33
6*I*A - 323*B)*a^3*e^(4*I*d*x + 4*I*c) + (-567*I*A - 551*B)*a^3*e^(2*I*d*x
+ 2*I*c) + 4*(-42*I*A - 41*B)*a^3)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(
2*I*d*x + 2*I*c) + 1)))/(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) +
3*d*e^(2*I*d*x + 2*I*c) + d)
```

3.128.6 Sympy [F]

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= -ia^3 \left(\int (-3A \tan^{\frac{3}{2}}(c+dx)) dx + \int A \tan^{\frac{7}{2}}(c+dx) dx + \int (-3B \tan^{\frac{5}{2}}(c+dx)) dx \right.$$

$$+ \int B \tan^{\frac{9}{2}}(c+dx) dx + \int iA \sqrt{\tan(c+dx)} dx + \int (-3iA \tan^{\frac{5}{2}}(c+dx)) dx$$

$$\left. + \int iB \tan^{\frac{3}{2}}(c+dx) dx + \int (-3iB \tan^{\frac{7}{2}}(c+dx)) dx \right)$$

input `integrate(tan(d*x+c)**(1/2)*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `-I*a**3*(Integral(-3*A*tan(c + d*x)**(3/2), x) + Integral(A*tan(c + d*x)**(7/2), x) + Integral(-3*B*tan(c + d*x)**(5/2), x) + Integral(B*tan(c + d*x)**(9/2), x) + Integral(I*A*sqrt(tan(c + d*x)), x) + Integral(-3*I*A*tan(c + d*x)**(5/2), x) + Integral(I*B*tan(c + d*x)**(3/2), x) + Integral(-3*I*B*tan(c + d*x)**(7/2), x))`

3.128.7 Maxima [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.26

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx =$$

$$\frac{30iBa^3 \tan(dx+c)^{\frac{7}{2}} + 42(iA+3B)a^3 \tan(dx+c)^{\frac{5}{2}} + 70(3A-4iB)a^3 \tan(dx+c)^{\frac{3}{2}} + 840(-iA -$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/105*(30*I*B*a^3*tan(d*x + c)^(7/2) + 42*(I*A + 3*B)*a^3*tan(d*x + c)^(5/2) + 70*(3*A - 4*I*B)*a^3*tan(d*x + c)^(3/2) + 840*(-I*A - B)*a^3*sqrt(tan(d*x + c)) + 105*(2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a^3)/d`

3.128. $\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.128.8 Giac [A] (verification not implemented)

Time = 0.91 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.94

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= -\frac{(4i-4)\sqrt{2}(Aa^3-iBa^3)\arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{d}$$

$$-\frac{2\left(15iBa^3d^6 \tan(dx+c)^{\frac{7}{2}}+21iAa^3d^6 \tan(dx+c)^{\frac{5}{2}}+63Ba^3d^6 \tan(dx+c)^{\frac{5}{2}}+105Aa^3d^6 \tan(dx+c)^{\frac{3}{2}}-140iBa^3d^6 \tan(dx+c)^{\frac{3}{2}}-420iAa^3d^6 \sqrt{\tan(dx+c)}-420Ba^3d^6 \sqrt{\tan(dx+c)}\right)}{105d^7}$$

input `integrate(tan(d*x+c)^(1/2)*(a+i*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `-(4*I - 4)*sqrt(2)*(A*a^3 - I*B*a^3)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d - 2/105*(15*I*B*a^3*d^6*tan(d*x + c)^(7/2) + 21*I*A*a^3*d^6*tan(d*x + c)^(5/2) + 63*B*a^3*d^6*tan(d*x + c)^(5/2) + 105*A*a^3*d^6*tan(d*x + c)^(3/2) - 140*I*B*a^3*d^6*tan(d*x + c)^(3/2) - 420*I*A*a^3*d^6*sqrt(tan(d*x + c)) - 420*B*a^3*d^6*sqrt(tan(d*x + c)))/d^7`

3.128.9 Mupad [B] (verification not implemented)

Time = 10.44 (sec) , antiderivative size = 292, normalized size of antiderivative = 1.71

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{Aa^3\sqrt{\tan(c+dx)}8i}{d} - \frac{2Aa^3 \tan(c+dx)^{3/2}}{d} - \frac{Aa^3 \tan(c+dx)^{5/2} 2i}{5d}$$

$$+ \frac{8Ba^3\sqrt{\tan(c+dx)}}{d} + \frac{Ba^3 \tan(c+dx)^{3/2} 8i}{3d}$$

$$- \frac{6Ba^3 \tan(c+dx)^{5/2}}{5d} - \frac{Ba^3 \tan(c+dx)^{7/2} 2i}{7d}$$

$$+ \frac{\sqrt{2}Aa^3 \ln\left(8Aa^3d + \sqrt{2}Aa^3d\sqrt{\tan(c+dx)}(-4-4i)\right)(2+2i)}{d}$$

$$- \frac{\sqrt{16i}Aa^3 \ln\left(8Aa^3d + 2\sqrt{16i}Aa^3d\sqrt{\tan(c+dx)}\right)}{d}$$

$$+ \frac{\sqrt{2}Ba^3 \ln\left(-Ba^3d8i + \sqrt{2}Ba^3d\sqrt{\tan(c+dx)}(-4+4i)\right)(2-2i)}{d}$$

$$- \frac{\sqrt{-16i}Ba^3 \ln\left(-Ba^3d8i + 2\sqrt{-16i}Ba^3d\sqrt{\tan(c+dx)}\right)}{d}$$

3.128. $\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

input `int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3,x)`

output
$$\begin{aligned} & (A*a^3*\tan(c + d*x)^{(1/2)}*8i)/d - (2*A*a^3*\tan(c + d*x)^{(3/2)})/d - (A*a^3* \\ & \tan(c + d*x)^{(5/2)}*2i)/(5*d) + (8*B*a^3*\tan(c + d*x)^{(1/2)})/d + (B*a^3*\tan \\ & (c + d*x)^{(3/2)}*8i)/(3*d) - (6*B*a^3*\tan(c + d*x)^{(5/2)})/(5*d) - (B*a^3*\tan \\ & (c + d*x)^{(7/2)}*2i)/(7*d) + (2^{(1/2)}*A*a^3*\log(8*A*a^3*d - 2^{(1/2)}*A*a^3* \\ & d*\tan(c + d*x)^{(1/2)}*(4 + 4i))*(2 + 2i))/d - (16i^{(1/2)}*A*a^3*\log(8*A*a^3* \\ & d + 2*16i^{(1/2)}*A*a^3*d*\tan(c + d*x)^{(1/2)}))/d + (2^{(1/2)}*B*a^3*\log(- B*a^ \\ & 3*d*8i - 2^{(1/2)}*B*a^3*d*\tan(c + d*x)^{(1/2)}*(4 - 4i))*(2 - 2i))/d - ((-16i \\ &)^{(1/2)}*B*a^3*\log(2*(-16i)^{(1/2)}*B*a^3*d*\tan(c + d*x)^{(1/2)} - B*a^3*d*8i)) \\ & /d \end{aligned}$$

3.129
$$\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.129.1 Optimal result 1342
 3.129.2 Mathematica [A] (verified) 1343
 3.129.3 Rubi [A] (verified) 1343
 3.129.4 Maple [B] (verified) 1346
 3.129.5 Fracas [B] (verification not implemented) 1347
 3.129.6 Sympy [F] 1348
 3.129.7 Maxima [A] (verification not implemented) 1349
 3.129.8 Giac [A] (verification not implemented) 1349
 3.129.9 Mupad [B] (verification not implemented) 1350

3.129.1 Optimal result

Integrand size = 36, antiderivative size = 146

$$\int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= -\frac{8\sqrt[4]{-1}a^3(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d}$$

$$- \frac{16a^3(5A - 6iB)\sqrt{\tan(c + dx)}}{15d} + \frac{2iaB\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2}{5d}$$

$$- \frac{2(5A - 9iB)\sqrt{\tan(c + dx)}(a^3 + ia^3 \tan(c + dx))}{15d}$$

output

```
-8*(-1)^(1/4)*a^3*(A-I*B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d-16/15*a^3*(5*A-6*I*B)*tan(d*x+c)^(1/2)/d+2/5*I*a*B*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^2/d-2/15*(5*A-9*I*B)*tan(d*x+c)^(1/2)*(a^3+I*a^3*tan(d*x+c))/d
```

3.129.2 Mathematica [A] (verified)

Time = 2.99 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.63

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \frac{2a^3 \left(60\sqrt[4]{-1}(A - iB) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right) + \sqrt{\tan(c + dx)}(45A - 60iB + 5(iA + 3B) \tan(c + dx)) \right)}{15d}$$

input `Integrate[((a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `(-2*a^3*(60*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + Sqrt[Tan[c + d*x]]*(45*A - (60*I)*B + 5*(I*A + 3*B)*Tan[c + d*x] + (3*I)*B*Tan[c + d*x]^2))/(15*d)`

3.129.3 Rubi [A] (verified)Time = 0.87 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.05, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.306$, Rules used = {3042, 4077, 27, 3042, 4077, 27, 3042, 4075, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\ & \quad \downarrow \text{4077} \\ & \frac{2}{5} \int \frac{(i \tan(c + dx)a + a)^2 (a(5A - iB) + a(5iA + 9B) \tan(c + dx))}{2\sqrt{\tan(c + dx)}} dx + \\ & \quad \frac{2iaB\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2}{5d} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.129. $\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

$$\frac{1}{5} \int \frac{(i \tan(c+dx)a + a)^2 (a(5A - iB) + a(5iA + 9B) \tan(c+dx))}{\sqrt{\tan(c+dx)} \frac{2iaB \sqrt{\tan(c+dx)} (a + ia \tan(c+dx))^2}{5d}} dx +$$

↓ 3042

$$\frac{1}{5} \int \frac{(i \tan(c+dx)a + a)^2 (a(5A - iB) + a(5iA + 9B) \tan(c+dx))}{\sqrt{\tan(c+dx)} \frac{2iaB \sqrt{\tan(c+dx)} (a + ia \tan(c+dx))^2}{5d}} dx +$$

↓ 4077

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{2(i \tan(c+dx)a + a) ((5A - 3iB)a^2 + 2(5iA + 6B) \tan(c+dx)a^2)}{\sqrt{\tan(c+dx)} \frac{2iaB \sqrt{\tan(c+dx)} (a + ia \tan(c+dx))^2}{5d}} dx - \frac{2(5A - 9iB) \sqrt{\tan(c+dx)} (a^3 + a)}{3d} \right)$$

↓ 27

$$\frac{1}{5} \left(\frac{4}{3} \int \frac{(i \tan(c+dx)a + a) ((5A - 3iB)a^2 + 2(5iA + 6B) \tan(c+dx)a^2)}{\sqrt{\tan(c+dx)} \frac{2iaB \sqrt{\tan(c+dx)} (a + ia \tan(c+dx))^2}{5d}} dx - \frac{2(5A - 9iB) \sqrt{\tan(c+dx)} (a^3 + a)}{3d} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{4}{3} \int \frac{(i \tan(c+dx)a + a) ((5A - 3iB)a^2 + 2(5iA + 6B) \tan(c+dx)a^2)}{\sqrt{\tan(c+dx)} \frac{2iaB \sqrt{\tan(c+dx)} (a + ia \tan(c+dx))^2}{5d}} dx - \frac{2(5A - 9iB) \sqrt{\tan(c+dx)} (a^3 + a)}{3d} \right)$$

↓ 4075

$$\frac{1}{5} \left(\frac{4}{3} \left(\int \frac{15(A - iB)a^3 + 15(iA + B) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)} \frac{2iaB \sqrt{\tan(c+dx)} (a + ia \tan(c+dx))^2}{5d}} dx - \frac{4a^3(5A - 6iB) \sqrt{\tan(c+dx)}}{d} \right) - \frac{2(5A - 9iB) \sqrt{\tan(c+dx)} (a^3 + a)}{3d} \right)$$

↓ 3042

3.129. $\int \frac{(a+ia \tan(c+dx))^3 (A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

$$\frac{1}{5} \left(\frac{4}{3} \left(\int \frac{15(A - iB)a^3 + 15(iA + B) \tan(c + dx)a^3}{\sqrt{\tan(c + dx)}} dx - \frac{4a^3(5A - 6iB)\sqrt{\tan(c + dx)}}{d} \right) - \frac{2(5A - 9iB)\sqrt{\tan(c + dx)}}{2iaB\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} \right)$$

↓ 4016

$$\frac{1}{5} \left(\frac{4}{3} \left(\frac{450a^6(A - iB)^2 \int \frac{1}{15a^3(A - iB) - 15a^3(iA + B) \tan(c + dx)} d\sqrt{\tan(c + dx)}}{d} - \frac{4a^3(5A - 6iB)\sqrt{\tan(c + dx)}}{d} \right) - \frac{2(5A - 9iB)\sqrt{\tan(c + dx)}}{2iaB\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} \right)$$

↓ 218

$$\frac{1}{5} \left(\frac{4}{3} \left(-\frac{30\sqrt[4]{-1}a^3(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d} - \frac{4a^3(5A - 6iB)\sqrt{\tan(c + dx)}}{d} \right) - \frac{2(5A - 9iB)\sqrt{\tan(c + dx)}}{2iaB\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} \right)$$

input `Int[((a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `((2*I)/5)*a*B*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^2/d + (4*((-30*(-1)^(1/4)*a^3*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]])/d - (4*a^3*(5*A - (6*I)*B)*Sqrt[Tan[c + d*x]])/d)/3 - (2*(5*A - (9*I)*B)*Sqrt[Tan[c + d*x]]*(a^3 + I*a^3*Tan[c + d*x]))/(3*d))/5`

3.129.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.129. $\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$

rule 4016 $\text{Int}[\frac{(c + (d \cdot \tan(e + f \cdot x)))}{\sqrt{(b \cdot \tan(e + f \cdot x) + (f \cdot x))}}, x_Symbol] \rightarrow \text{Simp}[2 \cdot (c^2/f) \text{ Subst}[\text{Int}[1/(b \cdot c - d \cdot x^2), x], x, \sqrt{b \cdot \tan[e + f \cdot x]}], x] /; \text{FreeQ}\{b, c, d, e, f\}, x\} \&\& \text{EqQ}[c^2 + d^2, 0]$

rule 4075 $\text{Int}[\frac{(a + (b \cdot \tan(e + f \cdot x)))^{m_1} \cdot ((A + (B \cdot \tan(e + f \cdot x) + (f \cdot x))) \cdot ((c + (d \cdot \tan(e + f \cdot x) + (f \cdot x))))}{x_Symbol} \rightarrow \text{Simp}[B \cdot d \cdot ((a + b \cdot \tan[e + f \cdot x])^{m+1} / (b \cdot f \cdot (m+1))), x] + \text{Int}[(a + b \cdot \tan[e + f \cdot x])^m \cdot \text{Simp}[A \cdot c - B \cdot d + (B \cdot c + A \cdot d) \cdot \tan[e + f \cdot x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, m\}, x\} \&\& \text{NeQ}[b \cdot c - a \cdot d, 0] \&\& \text{!LeQ}[m, -1]$

rule 4077 $\text{Int}[\frac{(a + (b \cdot \tan(e + f \cdot x)))^{m_1} \cdot ((A + (B \cdot \tan(e + f \cdot x) + (f \cdot x))) \cdot ((c + (d \cdot \tan(e + f \cdot x) + (f \cdot x))))^{n_1}}{x_Symbol} \rightarrow \text{Simp}[b \cdot B \cdot (a + b \cdot \tan[e + f \cdot x])^{m-1} \cdot ((c + d \cdot \tan[e + f \cdot x])^{n+1} / (d \cdot f \cdot (m+n))), x] + \text{Simp}[1/(d \cdot (m+n)) \text{ Int}[(a + b \cdot \tan[e + f \cdot x])^{m-1} \cdot (c + d \cdot \tan[e + f \cdot x])^n \cdot \text{Simp}[a \cdot A \cdot d \cdot (m+n) + B \cdot (a \cdot c \cdot (m-1) - b \cdot d \cdot (n+1)) - (B \cdot (b \cdot c - a \cdot d) \cdot (m-1) - d \cdot (A \cdot b + a \cdot B) \cdot (m+n)) \cdot \tan[e + f \cdot x], x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, n\}, x\} \&\& \text{NeQ}[b \cdot c - a \cdot d, 0] \&\& \text{EqQ}[a^2 + b^2, 0] \&\& \text{GtQ}[m, 1] \&\& \text{!LtQ}[n, -1]$

3.129.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 252 vs. 2(122) = 244.

Time = 0.03 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.73

method	result
derivativedivides	$a^3 \left(-\frac{2iB \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} - \frac{2iA \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 2B \left(\tan^{\frac{3}{2}}(dx+c) \right) + 8iB \left(\sqrt{\tan(dx+c)} \right) - 6A \left(\sqrt{\tan(dx+c)} \right) + \frac{(-4iB+4A)\sqrt{\tan(dx+c)}}{4} \right)$
default	$a^3 \left(-\frac{2iB \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} - \frac{2iA \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 2B \left(\tan^{\frac{3}{2}}(dx+c) \right) + 8iB \left(\sqrt{\tan(dx+c)} \right) - 6A \left(\sqrt{\tan(dx+c)} \right) + \frac{(-4iB+4A)\sqrt{\tan(dx+c)}}{4} \right)$
parts	$\frac{(-iA a^3 - 3B a^3) \left(\frac{2 \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - \frac{\sqrt{2} \left(\ln \left(\frac{1 - \sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)}{1 + \sqrt{2} \left(\sqrt{\tan(dx+c)} + \tan(dx+c) \right)} \right) + 2 \arctan \left(\frac{1 + \sqrt{2} \left(\sqrt{\tan(dx+c)} \right)}{1} \right) + 2 \arctan(-1)}{4} \right)}{d}$

3.129. $\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

input `int((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x,method=_RETURNVERBOSE)`

output `1/d*a^3*(-2/5*I*B*tan(d*x+c)^(5/2)-2/3*I*A*tan(d*x+c)^(3/2)-2*B*tan(d*x+c)^(3/2)+8*I*B*tan(d*x+c)^(1/2)-6*A*tan(d*x+c)^(1/2)+1/4*(-4*I*B+4*A)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(4*I*A+4*B)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))`

3.129.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 447 vs. $2(116) = 232$.

Time = 0.27 (sec) , antiderivative size = 447, normalized size of antiderivative = 3.06

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= 2 \left(15 \sqrt{-\frac{(iA^2 + 2AB - iB^2)a^6}{d^2}} (de^{(4i dx + 4i c)} + 2de^{(2i dx + 2i c)} + d) \log \left(-\frac{2 \left((A - iB)a^3 e^{(2i dx + 2i c)} + \sqrt{-\frac{(iA^2 + 2AB - iB^2)a^6}{d^2}} \right)}{(-i} \right. \right.$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="fracas")`

output `2/15*(15*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log(-2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3)) - 15*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log(-2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3)) - 2*((25*A - 39*I*B)*a^3*e^(4*I*d*x + 4*I*c) + 3*(15*A - 19*I*B)*a^3*e^(2*I*d*x + 2*I*c) + 4*(5*A - 6*I*B)*a^3)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)`

3.129.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= -ia^3 \left(\int (-3A \sqrt{\tan(c + dx)}) dx + \int A \tan^{\frac{5}{2}}(c + dx) dx \right. \\ \left. + \int (-3B \tan^{\frac{3}{2}}(c + dx)) dx + \int B \tan^{\frac{7}{2}}(c + dx) dx + \int \frac{iA}{\sqrt{\tan(c + dx)}} dx \right. \\ \left. + \int (-3iA \tan^{\frac{3}{2}}(c + dx)) dx + \int iB \sqrt{\tan(c + dx)} dx + \int (-3iB \tan^{\frac{5}{2}}(c + dx)) dx \right)$$

input `integrate((a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c))/tan(d*x+c)**(1/2),x)`

output `-I*a**3*(Integral(-3*A*sqrt(tan(c + d*x)), x) + Integral(A*tan(c + d*x)**(5/2), x) + Integral(-3*B*tan(c + d*x)**(3/2), x) + Integral(B*tan(c + d*x)**(7/2), x) + Integral(I*A/sqrt(tan(c + d*x)), x) + Integral(-3*I*A*tan(c + d*x)**(3/2), x) + Integral(I*B*sqrt(tan(c + d*x)), x) + Integral(-3*I*B*tan(c + d*x)**(5/2), x))`

3.129.7 Maxima [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.34

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx =$$

$$6i Ba^3 \tan(dx + c)^{\frac{5}{2}} + 10(iA + 3B)a^3 \tan(dx + c)^{\frac{3}{2}} + 30(3A - 4iB)a^3 \sqrt{\tan(dx + c)} - 15 \left(2\sqrt{2} \left((iA - 1)B \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2} + 2\sqrt{\tan(dx + c)})\right) + 2\sqrt{2} \left((iA + 1)A - (iA - 1)B \right) \arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2} - 2\sqrt{\tan(dx + c)})\right) \right) - \sqrt{2} \left((iA - 1)A + (iA + 1)B \right) \log(\sqrt{2}\sqrt{\tan(dx + c)}) + \tan(dx + c) + 1 + \sqrt{2} \left((iA - 1)A + (iA + 1)B \right) \log(-\sqrt{2}\sqrt{\tan(dx + c)}) + \tan(dx + c) + 1 \right) a^3 \right) / d$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="maxima")`

output `-1/15*(6*I*B*a^3*tan(d*x + c)^(5/2) + 10*(I*A + 3*B)*a^3*tan(d*x + c)^(3/2) + 30*(3*A - 4*I*B)*a^3*sqrt(tan(d*x + c)) - 15*(2*sqrt(2)*((I + 1)*A - (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I + 1)*A - (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c))) + tan(d*x + c) + 1) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a^3)/d`

3.129.8 Giac [A] (verification not implemented)

Time = 0.87 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.87

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \frac{(4i - 4) \sqrt{2}(-i Aa^3 - Ba^3) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{d}$$

$$+ \frac{2 \left(3i Ba^3 d^4 \tan(dx + c)^{\frac{5}{2}} + 5i Aa^3 d^4 \tan(dx + c)^{\frac{3}{2}} + 15 Ba^3 d^4 \tan(dx + c)^{\frac{3}{2}} + 45 Aa^3 d^4 \sqrt{\tan(dx + c)} \right)}{15 d^5}$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="giac")`

output `(4*I - 4)*sqrt(2)*(-I*A*a^3 - B*a^3)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d - 2/15*(3*I*B*a^3*d^4*tan(d*x + c)^(5/2) + 5*I*A*a^3*d^4*tan(d*x + c)^(3/2) + 15*B*a^3*d^4*tan(d*x + c)^(3/2) + 45*A*a^3*d^4*sqrt(tan(d*x + c)) - 60*I*B*a^3*d^4*sqrt(tan(d*x + c)))/d^5`

3.129.9 Mupad [B] (verification not implemented)

Time = 8.59 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.76

$$\begin{aligned}
& \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
&= -\frac{6 A a^3 \sqrt{\tan(c + dx)}}{d} - \frac{A a^3 \tan(c + dx)^{3/2} 2i}{3d} + \frac{B a^3 \sqrt{\tan(c + dx)} 8i}{d} \\
&\quad - \frac{2 B a^3 \tan(c + dx)^{3/2}}{d} - \frac{B a^3 \tan(c + dx)^{5/2} 2i}{5d} \\
&\quad + \frac{\sqrt{2} A a^3 \ln \left(A a^3 d 8i + \sqrt{2} A a^3 d \sqrt{\tan(c + dx)} (-4 + 4i) \right) (2 - 2i)}{d} \\
&\quad - \frac{\sqrt{-16i} A a^3 \ln \left(A a^3 d 8i + 2 \sqrt{-16i} A a^3 d \sqrt{\tan(c + dx)} \right)}{d} \\
&\quad + \frac{\sqrt{2} B a^3 \ln \left(8 B a^3 d + \sqrt{2} B a^3 d \sqrt{\tan(c + dx)} (-4 - 4i) \right) (2 + 2i)}{d} \\
&\quad - \frac{\sqrt{16i} B a^3 \ln \left(8 B a^3 d + 2 \sqrt{16i} B a^3 d \sqrt{\tan(c + dx)} \right)}{d}
\end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3)/tan(c + d*x)^(1/2),x)`output `(B*a^3*tan(c + d*x)^(1/2)*8i)/d - (A*a^3*tan(c + d*x)^(3/2)*2i)/(3*d) - (6*A*a^3*tan(c + d*x)^(1/2))/d - (2*B*a^3*tan(c + d*x)^(3/2))/d - (B*a^3*tan(c + d*x)^(5/2)*2i)/(5*d) + (2^(1/2)*A*a^3*log(A*a^3*d*8i - 2^(1/2)*A*a^3*d*tan(c + d*x)^(1/2)*(4 - 4i))*(2 - 2i))/d - ((-16i)^(1/2)*A*a^3*log(A*a^3*d*8i + 2*(-16i)^(1/2)*A*a^3*d*tan(c + d*x)^(1/2)))/d + (2^(1/2)*B*a^3*log(8*B*a^3*d - 2^(1/2)*B*a^3*d*tan(c + d*x)^(1/2)*(4 + 4i))*(2 + 2i))/d - (16i)^(1/2)*B*a^3*log(8*B*a^3*d + 2*16i^(1/2)*B*a^3*d*tan(c + d*x)^(1/2))/d`

3.130
$$\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.130.1 Optimal result 1351
 3.130.2 Mathematica [A] (verified) 1351
 3.130.3 Rubi [A] (verified) 1352
 3.130.4 Maple [B] (verified) 1355
 3.130.5 Fricas [B] (verification not implemented) 1356
 3.130.6 Sympy [F] 1356
 3.130.7 Maxima [A] (verification not implemented) 1357
 3.130.8 Giac [A] (verification not implemented) 1357
 3.130.9 Mupad [B] (verification not implemented) 1358

3.130.1 Optimal result

Integrand size = 36, antiderivative size = 134

$$\int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= -\frac{8\sqrt[4]{-1}a^3(iA + B) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d} - \frac{16a^3B\sqrt{\tan(c + dx)}}{3d}$$

$$- \frac{2aA(a + ia \tan(c + dx))^2}{d\sqrt{\tan(c + dx)}} + \frac{2(3iA - B)\sqrt{\tan(c + dx)}(a^3 + ia^3 \tan(c + dx))}{3d}$$

output `-8*(-1)^(1/4)*a^3*(I*A+B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d-16/3*a^3*B
 *tan(d*x+c)^(1/2)/d-2*a*A*(a+I*a*tan(d*x+c))^2/d/tan(d*x+c)^(1/2)+2/3*(3*I
 *A-B)*tan(d*x+c)^(1/2)*(a^3+I*a^3*tan(d*x+c))/d`

3.130.2 Mathematica [A] (verified)

Time = 1.89 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.70

$$\int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx =$$

$$-\frac{2ia^3\left(-3iA + 12\sqrt[4]{-1}(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right) \sqrt{\tan(c + dx)} + 3(A - 3iB) \tan(c + dx)\right)}{3d\sqrt{\tan(c + dx)}}$$

input `Integrate[((a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `(((-2*I)/3)*a^3*((-3*I)*A + 12*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sqrt[Tan[c + d*x]] + 3*(A - (3*I)*B)*Tan[c + d*x] + B*Tan[c + d*x]^2)/(d*Sqrt[Tan[c + d*x]])`

3.130.3 Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.306$, Rules used = {3042, 4076, 27, 3042, 4077, 27, 3042, 4075, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx \\
 & \quad \downarrow \text{4076} \\
 & 2 \int \frac{(i \tan(c + dx)a + a)^2 (a(5iA + B) + a(3A + iB) \tan(c + dx))}{2\sqrt{\tan(c + dx)}} dx - \frac{2aA(a + ia \tan(c + dx))^2}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(i \tan(c + dx)a + a)^2 (a(5iA + B) + a(3A + iB) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx - \frac{2aA(a + ia \tan(c + dx))^2}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(i \tan(c + dx)a + a)^2 (a(5iA + B) + a(3A + iB) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx - \frac{2aA(a + ia \tan(c + dx))^2}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{4077} \\
 & \frac{2}{3} \int \frac{2(i \tan(c + dx)a + a) ((3iA + B)a^2 + 2iB \tan(c + dx)a^2)}{\sqrt{\tan(c + dx)}} dx + \\
 & \frac{2(-B + 3iA)\sqrt{\tan(c + dx)}(a^3 + ia^3 \tan(c + dx))}{3d} - \frac{2aA(a + ia \tan(c + dx))^2}{d\sqrt{\tan(c + dx)}}
 \end{aligned}$$

3.130. $\int \frac{(a+ia \tan(c+dx))^3 (A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

↓ 27

$$\frac{4}{3} \int \frac{(i \tan(c+dx)a + a) ((3iA+B)a^2 + 2iB \tan(c+dx)a^2)}{\sqrt{\tan(c+dx)}} dx + \frac{2(-B+3iA)\sqrt{\tan(c+dx)}(a^3 + ia^3 \tan(c+dx))}{3d} - \frac{2aA(a + ia \tan(c+dx))^2}{d\sqrt{\tan(c+dx)}}$$

↓ 3042

$$\frac{4}{3} \int \frac{(i \tan(c+dx)a + a) ((3iA+B)a^2 + 2iB \tan(c+dx)a^2)}{\sqrt{\tan(c+dx)}} dx + \frac{2(-B+3iA)\sqrt{\tan(c+dx)}(a^3 + ia^3 \tan(c+dx))}{3d} - \frac{2aA(a + ia \tan(c+dx))^2}{d\sqrt{\tan(c+dx)}}$$

↓ 4075

$$\frac{4}{3} \left(-\frac{4a^3 B \sqrt{\tan(c+dx)}}{d} + \int \frac{3a^3(iA+B) - 3a^3(A-iB) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx \right) + \frac{2(-B+3iA)\sqrt{\tan(c+dx)}(a^3 + ia^3 \tan(c+dx))}{3d} - \frac{2aA(a + ia \tan(c+dx))^2}{d\sqrt{\tan(c+dx)}}$$

↓ 3042

$$\frac{4}{3} \left(-\frac{4a^3 B \sqrt{\tan(c+dx)}}{d} + \int \frac{3a^3(iA+B) - 3a^3(A-iB) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx \right) + \frac{2(-B+3iA)\sqrt{\tan(c+dx)}(a^3 + ia^3 \tan(c+dx))}{3d} - \frac{2aA(a + ia \tan(c+dx))^2}{d\sqrt{\tan(c+dx)}}$$

↓ 4016

$$\frac{4}{3} \left(-\frac{4a^3 B \sqrt{\tan(c+dx)}}{d} + \frac{18a^6(B+iA)^2 \int \frac{1}{3(iA+B)a^3 + 3(A-iB) \tan(c+dx)a^3} d\sqrt{\tan(c+dx)}}{d} \right) + \frac{2(-B+3iA)\sqrt{\tan(c+dx)}(a^3 + ia^3 \tan(c+dx))}{3d} - \frac{2aA(a + ia \tan(c+dx))^2}{d\sqrt{\tan(c+dx)}}$$

↓ 218

$$\frac{4}{3} \left(-\frac{4a^3 B \sqrt{\tan(c+dx)}}{d} - \frac{6\sqrt[4]{-1}a^3(B+iA) \arctan\left((-1)^{3/4} \sqrt{\tan(c+dx)}\right)}{d} \right) + \frac{2(-B+3iA)\sqrt{\tan(c+dx)}(a^3 + ia^3 \tan(c+dx))}{3d} - \frac{2aA(a + ia \tan(c+dx))^2}{d\sqrt{\tan(c+dx)}}$$

input `Int[((a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`


```
output (4*((-6*(-1)^(1/4)*a^3*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]])/d
- (4*a^3*B*Sqrt[Tan[c + d*x]])/d))/3 - (2*a*A*(a + I*a*Tan[c + d*x])^2)/(d
*Sqrt[Tan[c + d*x]]) + (2*((3*I)*A - B)*Sqrt[Tan[c + d*x]]*(a^3 + I*a^3*Ta
n[c + d*x]))/(3*d)
```

3.130.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4016 Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_
)]]], x_Symbol] := Simp[2*(c^2/f Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b
*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]
```

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

```
rule 4076 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m -
n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) +
b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

```
rule 4077 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan
[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c -
a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] &&
GtQ[m, 1] && !LtQ[n, -1]
```

3.130.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 240 vs. 2(114) = 228.

Time = 0.04 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.80

method	result
derivativedivides	$a^3 \left(-\frac{2iB \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 2iA \sqrt{\tan(dx+c)} - 6B \sqrt{\tan(dx+c)} - \frac{2A}{\sqrt{\tan(dx+c)}} + \frac{(4iA+4B)\sqrt{2}}{\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right)} \right)$
default	$a^3 \left(-\frac{2iB \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 2iA \sqrt{\tan(dx+c)} - 6B \sqrt{\tan(dx+c)} - \frac{2A}{\sqrt{\tan(dx+c)}} + \frac{(4iA+4B)\sqrt{2}}{\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right)} \right)$
parts	$\frac{(-iA a^3 - 3B a^3) \left(2\sqrt{\tan(dx+c)} - \frac{\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2\arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) + 2\arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right)}{4} \right)}{d}$

```
input int((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x,method=_RETURN
VERBOSE)
```

```
output 1/d*a^3*(-2/3*I*B*tan(d*x+c)^(3/2)-2*I*A*tan(d*x+c)^(1/2)-6*B*tan(d*x+c)^(
1/2)-2*A/tan(d*x+c)^(1/2)+1/4*(4*I*A+4*B)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)
)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(
1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(4*I*B-
4*A)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*
x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2
^(1/2)*tan(d*x+c)^(1/2))))
```

$$3.130. \int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.130.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 405 vs. $2(108) = 216$.

Time = 0.26 (sec) , antiderivative size = 405, normalized size of antiderivative = 3.02

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= 2 \left(3 \sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^6}{d^2}} (de^{(4i dx + 4i c)} - d) \log \left(-\frac{2 \left((A - iB)a^3 e^{(2i dx + 2i c)} + \sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^6}{d^2}} (de^{(2i dx + 2i c)} + d) \right)}{(-iA - B)a^3} \right) \right)$$

```
input integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="fricas")
```

```
output 2/3*(3*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2)*(d*e^(4*I*d*x + 4*I*c) - d)
*log(-2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) + sqrt(-(-I*A^2 - 2*A*B + I*B^2)
)*a^6/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(
e^(2*I*d*x + 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3)) - 3*sqrt
(-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2)*(d*e^(4*I*d*x + 4*I*c) - d)*log(-2*((A
- I*B)*a^3*e^(2*I*d*x + 2*I*c) - sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2)*
(d*e^(2*I*d*x + 2*I*c) + d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x
+ 2*I*c) + 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3)) - 2*((3*I*A + 5*B)*
a^3*e^(4*I*d*x + 4*I*c) + (3*I*A - B)*a^3*e^(2*I*d*x + 2*I*c) - 4*B*a^3)*s
qrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(4*I*d*x
+ 4*I*c) - d)
```

3.130.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= -ia^3 \left(\int \left(-\frac{3A}{\sqrt{\tan(c + dx)}} \right) dx + \int A \tan^{\frac{3}{2}}(c + dx) dx + \int \left(-3B \sqrt{\tan(c + dx)} \right) dx \right.$$

$$+ \int B \tan^{\frac{5}{2}}(c + dx) dx + \int \frac{iA}{\tan^{\frac{3}{2}}(c + dx)} dx + \int \left(-3iA \sqrt{\tan(c + dx)} \right) dx$$

$$\left. + \int \frac{iB}{\sqrt{\tan(c + dx)}} dx + \int \left(-3iB \tan^{\frac{3}{2}}(c + dx) \right) dx \right)$$

3.130. $\int \frac{(a+ia \tan(c+dx))^3 (A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

input `integrate((a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c))/tan(d*x+c)**(3/2),x)`

output `-I*a**3*(Integral(-3*A/sqrt(tan(c + d*x)), x) + Integral(A*tan(c + d*x)**(3/2), x) + Integral(-3*B*sqrt(tan(c + d*x)), x) + Integral(B*tan(c + d*x)**(5/2), x) + Integral(I*A/tan(c + d*x)**(3/2), x) + Integral(-3*I*A*sqrt(tan(c + d*x)), x) + Integral(I*B/sqrt(tan(c + d*x)), x) + Integral(-3*I*B*tan(c + d*x)**(3/2), x))`

3.130.7 Maxima [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.42

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{2i Ba^3 \tan(dx + c)^{\frac{3}{2}} + 6(iA + 3B)a^3 \sqrt{\tan(dx + c)} - 3 \left(2\sqrt{2}((i-1)A + (i+1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{\tan(dx + c)} + \tan(dx + c)\right)\right) \right)}{d}$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="maxima")`

output `-1/3*(2*I*B*a^3*tan(d*x + c)^(3/2) + 6*(I*A + 3*B)*a^3*sqrt(tan(d*x + c)) - 3*(2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a^3 + 6*A*a^3/sqrt(tan(d*x + c)))/d`

3.130.8 Giac [A] (verification not implemented)

Time = 1.06 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.81

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{2Aa^3}{d\sqrt{\tan(dx + c)}} + \frac{(4i - 4)\sqrt{2}(Aa^3 - iBa^3) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx + c)}\right)}{d}$$

$$- \frac{2\left(iBa^3d^2 \tan(dx + c)^{\frac{3}{2}} + 3iAa^3d^2 \sqrt{\tan(dx + c)} + 9Ba^3d^2 \sqrt{\tan(dx + c)}\right)}{3d^3}$$

3.130. $\int \frac{(a+ia \tan(c+dx))^3 (A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="giac")`

output `-2*A*a^3/(d*sqrt(tan(d*x + c))) + (4*I - 4)*sqrt(2)*(A*a^3 - I*B*a^3)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d - 2/3*(I*B*a^3*d^2*tan(d*x + c)^(3/2) + 3*I*A*a^3*d^2*sqrt(tan(d*x + c)) + 9*B*a^3*d^2*sqrt(tan(d*x + c)))/d^3`

3.130.9 Mupad [B] (verification not implemented)

Time = 8.63 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.78

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\ &= -\frac{2 A a^3}{d \sqrt{\tan(c + dx)}} - \frac{A a^3 \sqrt{\tan(c + dx)} 2i}{d} \\ & \quad - \frac{6 B a^3 \sqrt{\tan(c + dx)}}{d} - \frac{B a^3 \tan(c + dx)^{3/2} 2i}{3 d} \\ & \quad + \frac{\sqrt{2} A a^3 \ln\left(-8 A a^3 d + \sqrt{2} A a^3 d \sqrt{\tan(c + dx)}(-4 - 4i)\right) (2 + 2i)}{d} \\ & \quad - \frac{\sqrt{16i} A a^3 \ln\left(-8 A a^3 d + 2 \sqrt{16i} A a^3 d \sqrt{\tan(c + dx)}\right)}{d} \\ & \quad + \frac{\sqrt{2} B a^3 \ln\left(B a^3 d 8i + \sqrt{2} B a^3 d \sqrt{\tan(c + dx)}(-4 + 4i)\right) (2 - 2i)}{d} \\ & \quad - \frac{\sqrt{-16i} B a^3 \ln\left(B a^3 d 8i + 2 \sqrt{-16i} B a^3 d \sqrt{\tan(c + dx)}\right)}{d} \end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3)/tan(c + d*x)^(3/2),x)`

output `(2^(1/2)*A*a^3*log(-8*A*a^3*d - 2^(1/2)*A*a^3*d*tan(c + d*x)^(1/2)*(4 + 4i))*(2 + 2i))/d - (A*a^3*tan(c + d*x)^(1/2)*2i)/d - (6*B*a^3*tan(c + d*x)^(1/2))/d - (B*a^3*tan(c + d*x)^(3/2)*2i)/(3*d) - (2*A*a^3)/(d*tan(c + d*x)^(1/2)) - (16i^(1/2)*A*a^3*log(2*16i^(1/2)*A*a^3*d*tan(c + d*x)^(1/2) - 8*A*a^3*d))/d + (2^(1/2)*B*a^3*log(B*a^3*d*8i - 2^(1/2)*B*a^3*d*tan(c + d*x)^(1/2)*(4 - 4i))*(2 - 2i))/d - ((-16i)^(1/2)*B*a^3*log(B*a^3*d*8i + 2*(-16i)^(1/2)*B*a^3*d*tan(c + d*x)^(1/2)))/d`

3.131
$$\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.131.1 Optimal result 1359
 3.131.2 Mathematica [A] (verified) 1359
 3.131.3 Rubi [A] (verified) 1360
 3.131.4 Maple [B] (verified) 1363
 3.131.5 Fricas [B] (verification not implemented) 1364
 3.131.6 Sympy [F] 1365
 3.131.7 Maxima [A] (verification not implemented) 1365
 3.131.8 Giac [A] (verification not implemented) 1366
 3.131.9 Mupad [B] (verification not implemented) 1367

3.131.1 Optimal result

Integrand size = 36, antiderivative size = 136

$$\int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{8\sqrt[4]{-1}a^3(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d} - \frac{16a^3A\sqrt{\tan(c + dx)}}{3d}$$

$$- \frac{2aA(a + ia \tan(c + dx))^2}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2(7iA + 3B)(a^3 + ia^3 \tan(c + dx))}{3d\sqrt{\tan(c + dx)}}$$

output `8*(-1)^(1/4)*a^3*(A-I*B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d-16/3*a^3*A*tan(d*x+c)^(1/2)/d-2/3*a*A*(a+I*a*tan(d*x+c))^2/d/tan(d*x+c)^(3/2)-2/3*(7*I*A+3*B)*(a^3+I*a^3*tan(d*x+c))/d/tan(d*x+c)^(1/2)`

3.131.2 Mathematica [A] (verified)

Time = 2.00 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.69

$$\int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2a^3\left(-A + (-9iA - 3B) \tan(c + dx) + 12\sqrt[4]{-1}(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)\right) \tan^{\frac{3}{2}}(c + dx) -}{3d \tan^{\frac{3}{2}}(c + dx)}$$

input `Integrate[((a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(2*a^3*(-A + ((-9*I)*A - 3*B)*Tan[c + d*x] + 12*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Tan[c + d*x]^(3/2) - (3*I)*B*Tan[c + d*x]^2)/(3*d*Tan[c + d*x]^(3/2))`

3.131.3 Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.306$, Rules used = {3042, 4076, 27, 3042, 4076, 27, 3042, 4075, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx$$

↓ 4076

$$\frac{2}{3} \int \frac{(i \tan(c + dx)a + a)^2 (a(7iA + 3B) + a(A + 3iB) \tan(c + dx))}{2 \tan^{\frac{3}{2}}(c + dx)} dx - \frac{2aA(a + ia \tan(c + dx))^2}{3d \tan^{\frac{3}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{3} \int \frac{(i \tan(c + dx)a + a)^2 (a(7iA + 3B) + a(A + 3iB) \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx - \frac{2aA(a + ia \tan(c + dx))^2}{3d \tan^{\frac{3}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \int \frac{(i \tan(c + dx)a + a)^2 (a(7iA + 3B) + a(A + 3iB) \tan(c + dx))}{\tan(c + dx)^{3/2}} dx - \frac{2aA(a + ia \tan(c + dx))^2}{3d \tan^{\frac{3}{2}}(c + dx)}$$

↓ 4076

$$\frac{1}{3} \left(2 \int -\frac{2(i \tan(c + dx)a + a) (a^2(5A - 3iB) - 2ia^2A \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx - \frac{2(3B + 7iA) (a^3 + ia^3 \tan(c + dx))}{d\sqrt{\tan(c + dx)}} \right) - \frac{2aA(a + ia \tan(c + dx))^2}{3d \tan^{\frac{3}{2}}(c + dx)}$$

3.131. $\int \frac{(a+ia \tan(c+dx))^3 (A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

↓ 27

$$\frac{1}{3} \left(-4 \int \frac{(i \tan(c+dx)a + a) (a^2(5A - 3iB) - 2ia^2A \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx - \frac{2(3B + 7iA) (a^3 + ia^3 \tan(c+dx))}{d\sqrt{\tan(c+dx)}} \right) - \frac{2aA(a + ia \tan(c+dx))^2}{3d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{3} \left(-4 \int \frac{(i \tan(c+dx)a + a) (a^2(5A - 3iB) - 2ia^2A \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx - \frac{2(3B + 7iA) (a^3 + ia^3 \tan(c+dx))}{d\sqrt{\tan(c+dx)}} \right) - \frac{2aA(a + ia \tan(c+dx))^2}{3d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 4075

$$\frac{1}{3} \left(-4 \left(\frac{4a^3A\sqrt{\tan(c+dx)}}{d} + \int \frac{3(A - iB)a^3 + 3(iA + B) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx \right) - \frac{2(3B + 7iA) (a^3 + ia^3 \tan(c+dx))}{d\sqrt{\tan(c+dx)}} \right) - \frac{2aA(a + ia \tan(c+dx))^2}{3d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{3} \left(-4 \left(\frac{4a^3A\sqrt{\tan(c+dx)}}{d} + \int \frac{3(A - iB)a^3 + 3(iA + B) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx \right) - \frac{2(3B + 7iA) (a^3 + ia^3 \tan(c+dx))}{d\sqrt{\tan(c+dx)}} \right) - \frac{2aA(a + ia \tan(c+dx))^2}{3d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 4016

$$\frac{1}{3} \left(-4 \left(\frac{4a^3A\sqrt{\tan(c+dx)}}{d} + \frac{18a^6(A - iB)^2 \int \frac{1}{3a^3(A - iB) - 3a^3(iA + B) \tan(c+dx)} d\sqrt{\tan(c+dx)}}{d} \right) - \frac{2(3B + 7iA) (a^3 + ia^3 \tan(c+dx))}{d\sqrt{\tan(c+dx)}} \right) - \frac{2aA(a + ia \tan(c+dx))^2}{3d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 218

$$\frac{1}{3} \left(-4 \left(\frac{4a^3A\sqrt{\tan(c+dx)}}{d} - \frac{6\sqrt[4]{-1}a^3(A - iB) \arctan \left((-1)^{3/4} \sqrt{\tan(c+dx)} \right)}{d} \right) - \frac{2(3B + 7iA) (a^3 + ia^3 \tan(c+dx))}{d\sqrt{\tan(c+dx)}} \right) - \frac{2aA(a + ia \tan(c+dx))^2}{3d \tan^{\frac{3}{2}}(c+dx)}$$

3.131. $\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

input `Int[((a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(-2*a*A*(a + I*a*Tan[c + d*x])^2)/(3*d*Tan[c + d*x]^(3/2)) + (-4*((-6*(-1)^(1/4)*a^3*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]])/d + (4*a^3*A*Sqrt[Tan[c + d*x]]/d) - (2*((7*I)*A + 3*B)*(a^3 + I*a^3*Tan[c + d*x]))/(d*Sqrt[Tan[c + d*x]]))/3`

3.131.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

```
rule 4076 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

3.131.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 233 vs. 2(114) = 228.

Time = 0.04 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.72

method	result
derivativedivides	$a^3 \left(-2iB(\sqrt{\tan(dx+c)}) - \frac{2A}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(3iA+B)}{\sqrt{\tan(dx+c)}} + \frac{(4iB-4A)\sqrt{2}}{4} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right) \right)$
default	$a^3 \left(-2iB(\sqrt{\tan(dx+c)}) - \frac{2A}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(3iA+B)}{\sqrt{\tan(dx+c)}} + \frac{(4iB-4A)\sqrt{2}}{4} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right) \right)$
parts	$\frac{(-iA a^3 - 3B a^3)\sqrt{2}}{4d} \left(\ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right)$

```
input int((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2), x, method=_RETUR
NVERBOSE)
```

```
output 1/d*a^3*(-2*I*B*tan(d*x+c)^(1/2)-2/3*A/tan(d*x+c)^(3/2)-2*(3*I*A+B)/tan(d*
x+c)^(1/2)+1/4*(4*I*B-4*A)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x
+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c
)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(-4*I*A-4*B)*2^(1/2)*(
ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan
(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x
+c)^(1/2))))
```

$$3.131. \int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.131.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 438 vs. $2(110) = 220$.

Time = 0.26 (sec) , antiderivative size = 438, normalized size of antiderivative = 3.22

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx =$$

$$2 \left(3 \sqrt{-\frac{(iA^2 + 2AB - iB^2)a^6}{d^2}} (de^{4i dx + 4i c} - 2de^{2i dx + 2i c} + d) \log \left(-\frac{2 \left((A - iB)a^3 e^{2i dx + 2i c} + \sqrt{-\frac{(iA^2 + 2AB - iB^2)a^6}{d^2}} \right)}{(-} \right. \right.$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="fricas")`

output `-2/3*(3*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log(-2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3)) - 3*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log(-2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3)) - 2*((5*A - 3*I*B)*a^3*e^(4*I*d*x + 4*I*c) + (A + 3*I*B)*a^3*e^(2*I*d*x + 2*I*c) - 4*A*a^3)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)`

3.131.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= -ia^3 \left(\int \left(-\frac{3A}{\tan^{\frac{3}{2}}(c + dx)} \right) dx + \int A \sqrt{\tan(c + dx)} dx + \int \left(-\frac{3B}{\sqrt{\tan(c + dx)}} \right) dx \right.$$

$$+ \int B \tan^{\frac{3}{2}}(c + dx) dx + \int \frac{iA}{\tan^{\frac{5}{2}}(c + dx)} dx + \int \left(-\frac{3iA}{\sqrt{\tan(c + dx)}} \right) dx$$

$$\left. + \int \frac{iB}{\tan^{\frac{3}{2}}(c + dx)} dx + \int \left(-3iB \sqrt{\tan(c + dx)} \right) dx \right)$$

input `integrate((a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c))/tan(d*x+c)**(5/2),x)`

output `-I*a**3*(Integral(-3*A/tan(c + d*x)**(3/2), x) + Integral(A*sqrt(tan(c + d*x)), x) + Integral(-3*B/sqrt(tan(c + d*x)), x) + Integral(B*tan(c + d*x)**(3/2), x) + Integral(I*A/tan(c + d*x)**(5/2), x) + Integral(-3*I*A/sqrt(tan(c + d*x)), x) + Integral(I*B/tan(c + d*x)**(3/2), x) + Integral(-3*I*B*sqrt(tan(c + d*x)), x))`

3.131.7 Maxima [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.40

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx =$$

$$6i Ba^3 \sqrt{\tan(dx + c)} + 3 \left(2 \sqrt{2} ((i + 1) A - (i - 1) B) \arctan \left(\frac{1}{2} \sqrt{2} \left(\sqrt{2} + 2 \sqrt{\tan(dx + c)} \right) \right) \right) + 2 \sqrt{2}$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="maxima")`

output
$$\begin{aligned} & -1/3*(6*I*B*a^3*\sqrt{\tan(dx + c)} + 3*(2*\sqrt{2}*((I + 1)*A - (I - 1)*B)* \\ & \arctan(1/2*\sqrt{2}*(\sqrt{2} + 2*\sqrt{\tan(dx + c)}))) + 2*\sqrt{2}*((I + 1)* \\ & A - (I - 1)*B)*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2*\sqrt{\tan(dx + c)})) - \sqrt{2} \\ & *((I - 1)*A + (I + 1)*B)*\log(\sqrt{2}*\sqrt{\tan(dx + c)} + \tan(dx + c) \\ & + 1) + \sqrt{2}*((I - 1)*A + (I + 1)*B)*\log(-\sqrt{2}*\sqrt{\tan(dx + c)} + \\ & \tan(dx + c) + 1))*a^3 - 2*(3*(-3*I*A - B)*a^3*\tan(dx + c) - A*a^3)/\tan(d \\ & *x + c)^{(3/2)}/d \end{aligned}$$

3.131.8 Giac [A] (verification not implemented)

Time = 1.06 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.71

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \\ & = -\frac{2i Ba^3 \sqrt{\tan(dx + c)}}{d} \\ & \quad - \frac{(4i - 4) \sqrt{2}(-i Aa^3 - Ba^3) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{d} \\ & \quad + \frac{2(-9i Aa^3 \tan(dx + c) - 3Ba^3 \tan(dx + c) - Aa^3)}{3d \tan(dx + c)^{\frac{3}{2}}} \end{aligned}$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="giac")`

output
$$\begin{aligned} & -2*I*B*a^3*\sqrt{\tan(dx + c)}/d - (4*I - 4)*\sqrt{2}*(-I*A*a^3 - B*a^3)*\arctan \\ & \tan(-1/2*I - 1/2)*\sqrt{2}*\sqrt{\tan(dx + c)}/d + 2/3*(-9*I*A*a^3*\tan(dx \\ & + c) - 3*B*a^3*\tan(dx + c) - A*a^3)/(d*\tan(dx + c)^{(3/2)}) \end{aligned}$$

3.131.9 Mupad [B] (verification not implemented)

Time = 8.66 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.76

$$\begin{aligned}
& \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \\
&= -\frac{\frac{2Aa^3}{3d} + \frac{Aa^3 \tan(c+dx) 6i}{d}}{\tan(c + dx)^{3/2}} - \frac{2Ba^3}{d \sqrt{\tan(c + dx)}} - \frac{Ba^3 \sqrt{\tan(c + dx)} 2i}{d} \\
&+ \frac{\sqrt{2} A a^3 \ln\left(-A a^3 d 8i + \sqrt{2} A a^3 d \sqrt{\tan(c + dx)} (-4 + 4i)\right) (2 - 2i)}{d} \\
&- \frac{\sqrt{-16i} A a^3 \ln\left(-A a^3 d 8i + 2 \sqrt{-16i} A a^3 d \sqrt{\tan(c + dx)}\right)}{d} \\
&+ \frac{\sqrt{2} B a^3 \ln\left(-8 B a^3 d + \sqrt{2} B a^3 d \sqrt{\tan(c + dx)} (-4 - 4i)\right) (2 + 2i)}{d} \\
&- \frac{\sqrt{16i} B a^3 \ln\left(-8 B a^3 d + 2 \sqrt{16i} B a^3 d \sqrt{\tan(c + dx)}\right)}{d}
\end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3)/tan(c + d*x)^(5/2),x)`

output `(2^(1/2)*A*a^3*log(-A*a^3*d*8i - 2^(1/2)*A*a^3*d*tan(c + d*x)^(1/2)*(4 - 4i))*(2 - 2i))/d - (2*B*a^3)/(d*tan(c + d*x)^(1/2)) - (B*a^3*tan(c + d*x)^(1/2)*2i)/d - ((2*A*a^3)/(3*d) + (A*a^3*tan(c + d*x)*6i)/d)/tan(c + d*x)^(3/2) - ((-16i)^(1/2)*A*a^3*log(2*(-16i)^(1/2)*A*a^3*d*tan(c + d*x)^(1/2) - A*a^3*d*8i))/d + (2^(1/2)*B*a^3*log(-8*B*a^3*d - 2^(1/2)*B*a^3*d*tan(c + d*x)^(1/2)*(4 + 4i))*(2 + 2i))/d - (16i)^(1/2)*B*a^3*log(2*16i^(1/2)*B*a^3*d*tan(c + d*x)^(1/2) - 8*B*a^3*d))/d`

3.132
$$\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

3.132.1 Optimal result 1368
 3.132.2 Mathematica [A] (verified) 1368
 3.132.3 Rubi [A] (verified) 1369
 3.132.4 Maple [A] (verified) 1372
 3.132.5 Fricas [B] (verification not implemented) 1373
 3.132.6 Sympy [F] 1374
 3.132.7 Maxima [A] (verification not implemented) 1374
 3.132.8 Giac [A] (verification not implemented) 1375
 3.132.9 Mupad [B] (verification not implemented) 1376

3.132.1 Optimal result

Integrand size = 36, antiderivative size = 144

$$\int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{8\sqrt[4]{-1}a^3(iA + B) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d} + \frac{16a^3(6A - 5iB)}{15d\sqrt{\tan(c + dx)}}$$

$$- \frac{2aA(a + ia \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{2(9iA + 5B)(a^3 + ia^3 \tan(c + dx))}{15d \tan^{\frac{3}{2}}(c + dx)}$$

```
output 8*(-1)^(1/4)*a^3*(I*A+B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d+16/15*a^3*(
6*A-5*I*B)/d/tan(d*x+c)^(1/2)-2/5*a*A*(a+I*a*tan(d*x+c))^2/d/tan(d*x+c)^(5
/2)-2/15*(9*I*A+5*B)*(a^3+I*a^3*tan(d*x+c))/d/tan(d*x+c)^(3/2)
```

3.132.2 Mathematica [A] (verified)

Time = 2.87 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.69

$$\int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{2a^3\left(-3A + (-15iA - 5B) \tan(c + dx) + 15(4A - 3iB) \tan^2(c + dx) + 60\sqrt[4]{-1}(iA + B) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)\right)}{15d \tan^{\frac{5}{2}}(c + dx)}$$

input `Integrate[((a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(2*a^3*(-3*A + ((-15*I)*A - 5*B)*Tan[c + d*x] + 15*(4*A - (3*I)*B)*Tan[c + d*x]^2 + 60*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Tan[c + d*x]^(5/2))/(15*d*Tan[c + d*x]^(5/2))`

3.132.3 Rubi [A] (verified)

Time = 0.86 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.06, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 4076, 27, 3042, 4076, 27, 3042, 4074, 27, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan(c + dx)^{7/2}} dx$$

↓ 4076

$$\frac{2}{5} \int \frac{(i \tan(c + dx)a + a)^2 (a(9iA + 5B) - a(A - 5iB) \tan(c + dx))}{2 \tan^{\frac{5}{2}}(c + dx)} dx - \frac{2aA(a + ia \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{5} \int \frac{(i \tan(c + dx)a + a)^2 (a(9iA + 5B) - a(A - 5iB) \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx - \frac{2aA(a + ia \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{5} \int \frac{(i \tan(c + dx)a + a)^2 (a(9iA + 5B) - a(A - 5iB) \tan(c + dx))}{\tan(c + dx)^{5/2}} dx - \frac{2aA(a + ia \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)}$$

↓ 4076

$$\frac{1}{5} \left(\frac{2}{3} \int - \frac{2(i \tan(c + dx)a + a) (2(6A - 5iB)a^2 + (3iA + 5B) \tan(c + dx)a^2)}{\tan^{\frac{3}{2}}(c + dx)} dx - \frac{2(5B + 9iA) (a^3 + ia^3 \tan(c + dx))}{3d \tan^{\frac{3}{2}}(c + dx)} \right) - \frac{2aA(a + ia \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)}$$

3.132. $\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$

↓ 27

$$\frac{1}{5} \left(-\frac{4}{3} \int \frac{(i \tan(c+dx)a + a) (2(6A - 5iB)a^2 + (3iA + 5B) \tan(c+dx)a^2)}{\frac{\tan^{\frac{3}{2}}(c+dx)}{2aA(a + ia \tan(c+dx))^2}} dx - \frac{2(5B + 9iA) (a^3 + ia^3 \tan(c+dx))}{3d \tan^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(-\frac{4}{3} \int \frac{(i \tan(c+dx)a + a) (2(6A - 5iB)a^2 + (3iA + 5B) \tan(c+dx)a^2)}{\frac{\tan(c+dx)^{3/2}}{2aA(a + ia \tan(c+dx))^2}} dx - \frac{2(5B + 9iA) (a^3 + ia^3 \tan(c+dx))}{3d \tan^{\frac{3}{2}}(c+dx)} \right)$$

↓ 4074

$$\frac{1}{5} \left(-\frac{4}{3} \left(\int \frac{15(a^3(iA + B) - a^3(A - iB) \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx - \frac{4a^3(6A - 5iB)}{d\sqrt{\tan(c+dx)}} \right) - \frac{2(5B + 9iA) (a^3 + ia^3 \tan(c+dx))}{3d \tan^{\frac{3}{2}}(c+dx)} \right)$$

↓ 27

$$\frac{1}{5} \left(-\frac{4}{3} \left(15 \int \frac{a^3(iA + B) - a^3(A - iB) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{4a^3(6A - 5iB)}{d\sqrt{\tan(c+dx)}} \right) - \frac{2(5B + 9iA) (a^3 + ia^3 \tan(c+dx))}{3d \tan^{\frac{3}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(-\frac{4}{3} \left(15 \int \frac{a^3(iA + B) - a^3(A - iB) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{4a^3(6A - 5iB)}{d\sqrt{\tan(c+dx)}} \right) - \frac{2(5B + 9iA) (a^3 + ia^3 \tan(c+dx))}{3d \tan^{\frac{3}{2}}(c+dx)} \right)$$

↓ 4016

$$\frac{1}{5} \left(-\frac{4}{3} \left(\frac{30a^6(B + iA)^2 \int \frac{1}{(iA+B)a^3+(A-iB)\tan(c+dx)a^3} d\sqrt{\tan(c+dx)}}{d} - \frac{4a^3(6A - 5iB)}{d\sqrt{\tan(c+dx)}} \right) - \frac{2(5B + 9iA) (a^3 + ia^3 \tan(c+dx))}{3d \tan^{\frac{3}{2}}(c+dx)} \right)$$

3.132. $\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

↓ 218

$$\frac{1}{5} \left(-\frac{4}{3} \left(-\frac{30\sqrt[4]{-1}a^3(B+iA) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{d} - \frac{4a^3(6A-5iB)}{d\sqrt{\tan(c+dx)}} \right) - \frac{2(5B+9iA)(a^3+ia^3 \tan(c+dx))}{3d \tan^{3/2}(c+dx)} \right) - \frac{2aA(a+ia \tan(c+dx))^2}{5d \tan^{5/2}(c+dx)}$$

input `Int[((a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(-2*a*A*(a + I*a*Tan[c + d*x])^2)/(5*d*Tan[c + d*x]^(5/2)) + ((-4*((-30*(-1)^(1/4)*a^3*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]])/d - (4*a^3*(6*A - (5*I)*B))/(d*Sqrt[Tan[c + d*x]])))/3 - (2*((9*I)*A + 5*B)*(a^3 + I*a^3*Tan[c + d*x]))/(3*d*Tan[c + d*x]^(3/2))/5`

3.132.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4016 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

```
rule 4076 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

3.132.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.67

method	result
derivativedivides	$a^3 \left(-\frac{2A}{5 \tan(dx+c)^{\frac{5}{2}}} - \frac{2(3iA+B)}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(3iB-4A)}{\sqrt{\tan(dx+c)}} + \frac{(-4iA-4B)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right)}{4} \right)$
default	$a^3 \left(-\frac{2A}{5 \tan(dx+c)^{\frac{5}{2}}} - \frac{2(3iA+B)}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(3iB-4A)}{\sqrt{\tan(dx+c)}} + \frac{(-4iA-4B)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right)}{4} \right)$
parts	$\frac{(-iA a^3 - 3B a^3)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right)}{4d}$

```
input int((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2), x, method=_RETUR
NVERBOSE)
```

$$3.132. \int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

output $1/d*a^3*(-2/5*A/\tan(d*x+c)^{(5/2)}-2/3*(3*I*A+B)/\tan(d*x+c)^{(3/2)}-2*(3*I*B-4*A)/\tan(d*x+c)^{(1/2)}+1/4*(-4*I*A-4*B)*2^{(1/2)}*(\ln((1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}))+1/4*(-4*I*B+4*A)*2^{(1/2)}*(\ln((1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}))$

3.132.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 504 vs. $2(116) = 232$.

Time = 0.25 (sec) , antiderivative size = 504, normalized size of antiderivative = 3.50

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx =$$

$$2 \left(15 \sqrt{-\frac{(-i A^2 - 2AB + i B^2)a^6}{d^2}} (de^{(6i dx + 6i c)} - 3de^{(4i dx + 4i c)} + 3de^{(2i dx + 2i c)} - d) \log \left(-\frac{2 \left((A - i B)a^3 e^{(2i dx + 2i c)} \right)}{\dots} \right) \right)$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorith="fracas")`

output $-2/15*(15*\sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2}*(d*e^{(6*I*d*x + 6*I*c)} - 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} - d)*\log(-2*((A - I*B)*a^3*e^{(2*I*d*x + 2*I*c)} + \sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2}*(d*e^{(2*I*d*x + 2*I*c)} + d)*\sqrt{(-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1)}))e^{(-2*I*d*x - 2*I*c)}/((-I*A - B)*a^3)) - 15*\sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2}*(d*e^{(6*I*d*x + 6*I*c)} - 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} - d)*\log(-2*((A - I*B)*a^3*e^{(2*I*d*x + 2*I*c)} - \sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2}*(d*e^{(2*I*d*x + 2*I*c)} + d)*\sqrt{(-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1)}))e^{(-2*I*d*x - 2*I*c)}/((-I*A - B)*a^3)) + 2*((-39*I*A - 25*B)*a^3*e^{(6*I*d*x + 6*I*c)} + 2*(9*I*A + 10*B)*a^3*e^{(4*I*d*x + 4*I*c)} + (33*I*A + 25*B)*a^3*e^{(2*I*d*x + 2*I*c)} + 4*(-6*I*A - 5*B)*a^3)*\sqrt{(-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1)})/(d*e^{(6*I*d*x + 6*I*c)} - 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} - d)$

3.132. $\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

3.132.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= -ia^3 \left(\int \left(-\frac{3A}{\tan^{\frac{5}{2}}(c + dx)} \right) dx + \int \frac{A}{\sqrt{\tan(c + dx)}} dx + \int \left(-\frac{3B}{\tan^{\frac{3}{2}}(c + dx)} \right) dx \right.$$

$$+ \int B \sqrt{\tan(c + dx)} dx + \int \frac{iA}{\tan^{\frac{7}{2}}(c + dx)} dx + \int \left(-\frac{3iA}{\tan^{\frac{3}{2}}(c + dx)} \right) dx$$

$$\left. + \int \frac{iB}{\tan^{\frac{5}{2}}(c + dx)} dx + \int \left(-\frac{3iB}{\sqrt{\tan(c + dx)}} \right) dx \right)$$

input `integrate((a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c))/tan(d*x+c)**(7/2),x)`

output `-I*a**3*(Integral(-3*A/tan(c + d*x)**(5/2), x) + Integral(A/sqrt(tan(c + d*x)), x) + Integral(-3*B/tan(c + d*x)**(3/2), x) + Integral(B*sqrt(tan(c + d*x)), x) + Integral(I*A/tan(c + d*x)**(7/2), x) + Integral(-3*I*A/tan(c + d*x)**(3/2), x) + Integral(I*B/tan(c + d*x)**(5/2), x) + Integral(-3*I*B/sqrt(tan(c + d*x)), x))`

3.132.7 Maxima [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.37

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx =$$

$$\frac{15 \left(2 \sqrt{2} ((i - 1) A + (i + 1) B) \arctan \left(\frac{1}{2} \sqrt{2} \left(\sqrt{2} + 2 \sqrt{\tan(dx + c)} \right) \right) + 2 \sqrt{2} ((i - 1) A + (i + 1) B) \right)}{\dots}$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="maxima")`

output
$$\frac{-1/15*(15*(2*\sqrt{2})*((I - 1)*A + (I + 1)*B)*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2*\sqrt{\tan(dx + c)})) + 2*\sqrt{2}*((I - 1)*A + (I + 1)*B)*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2*\sqrt{\tan(dx + c)})) - \sqrt{2}*(-(I + 1)*A + (I - 1)*B) * \log(\sqrt{2}*\sqrt{\tan(dx + c)} + \tan(dx + c) + 1) + \sqrt{2}*(-(I + 1)*A + (I - 1)*B) * \log(-\sqrt{2}*\sqrt{\tan(dx + c)} + \tan(dx + c) + 1))*a^3 - 2*(15*(4*A - 3*I*B)*a^3*\tan(dx + c)^2 + 5*(-3*I*A - B)*a^3*\tan(dx + c) - 3*A*a^3)/\tan(dx + c)^{(5/2)}/d$$

3.132.8 Giac [A] (verification not implemented)

Time = 1.32 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.74

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= - \frac{(4i - 4) \sqrt{2} (Aa^3 - i Ba^3) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{d} + \frac{2(60 Aa^3 \tan(dx + c)^2 - 45i Ba^3 \tan(dx + c)^2 - 15i Aa^3 \tan(dx + c) - 5 Ba^3 \tan(dx + c) - 3 Aa^3)}{15 d \tan(dx + c)^{\frac{5}{2}}}$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="giac")`

output
$$-(4*I - 4)*\sqrt{2}*(A*a^3 - I*B*a^3)*\arctan(-1/2*I - 1/2)*\sqrt{2}*\sqrt{\tan(dx + c)}/d + 2/15*(60*A*a^3*\tan(dx + c)^2 - 45*I*B*a^3*\tan(dx + c)^2 - 15*I*A*a^3*\tan(dx + c) - 5*B*a^3*\tan(dx + c) - 3*A*a^3)/(d*\tan(dx + c)^{(5/2)})$$

3.132.9 Mupad [B] (verification not implemented)

Time = 9.10 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.79

$$\begin{aligned}
& \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx \\
&= -\frac{\frac{2Aa^3}{5d} - \frac{8Aa^3 \tan(c+dx)^2}{d} + \frac{Aa^3 \tan(c+dx)2i}{d}}{\tan(c + dx)^{5/2}} - \frac{\frac{2Ba^3}{3d} + \frac{Ba^3 \tan(c+dx)6i}{d}}{\tan(c + dx)^{3/2}} \\
&+ \frac{\sqrt{2} A a^3 \ln \left(8 A a^3 d + \sqrt{2} A a^3 d \sqrt{\tan(c + dx)} (-4 - 4i) \right) (2 + 2i)}{d} \\
&- \frac{\sqrt{16i} A a^3 \ln \left(8 A a^3 d + 2 \sqrt{16i} A a^3 d \sqrt{\tan(c + dx)} \right)}{d} \\
&+ \frac{\sqrt{2} B a^3 \ln \left(-B a^3 d 8i + \sqrt{2} B a^3 d \sqrt{\tan(c + dx)} (-4 + 4i) \right) (2 - 2i)}{d} \\
&- \frac{\sqrt{-16i} B a^3 \ln \left(-B a^3 d 8i + 2 \sqrt{-16i} B a^3 d \sqrt{\tan(c + dx)} \right)}{d}
\end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3)/tan(c + d*x)^(7/2),x)`

output `(2^(1/2)*A*a^3*log(8*A*a^3*d - 2^(1/2)*A*a^3*d*tan(c + d*x)^(1/2)*(4 + 4i))*(2 + 2i))/d - ((2*B*a^3)/(3*d) + (B*a^3*tan(c + d*x)*6i)/d)/tan(c + d*x)^(3/2) - ((2*A*a^3)/(5*d) + (A*a^3*tan(c + d*x)*2i)/d - (8*A*a^3*tan(c + d*x)^2)/d)/tan(c + d*x)^(5/2) - (16i^(1/2)*A*a^3*log(8*A*a^3*d + 2*16i^(1/2)*A*a^3*d*tan(c + d*x)^(1/2)))/d + (2^(1/2)*B*a^3*log(-B*a^3*d*8i - 2^(1/2)*B*a^3*d*tan(c + d*x)^(1/2)*(4 - 4i))*(2 - 2i))/d - ((-16i)^(1/2)*B*a^3*log(2*(-16i)^(1/2)*B*a^3*d*tan(c + d*x)^(1/2) - B*a^3*d*8i))/d`

3.133
$$\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

3.133.1 Optimal result 1377
 3.133.2 Mathematica [A] (verified) 1377
 3.133.3 Rubi [A] (verified) 1378
 3.133.4 Maple [A] (verified) 1382
 3.133.5 Fricas [B] (verification not implemented) 1383
 3.133.6 Sympy [F] 1384
 3.133.7 Maxima [A] (verification not implemented) 1385
 3.133.8 Giac [A] (verification not implemented) 1385
 3.133.9 Mupad [B] (verification not implemented) 1386

3.133.1 Optimal result

Integrand size = 36, antiderivative size = 169

$$\int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$= -\frac{8\sqrt[4]{-1}a^3(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d} + \frac{8a^3(23A - 21iB)}{105d \tan^{\frac{3}{2}}(c + dx)}$$

$$+ \frac{8a^3(iA + B)}{d\sqrt{\tan(c + dx)}} - \frac{2aA(a + ia \tan(c + dx))^2}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{2(11iA + 7B)(a^3 + ia^3 \tan(c + dx))}{35d \tan^{\frac{5}{2}}(c + dx)}$$

output

```
-8*(-1)^(1/4)*a^3*(A-I*B)*arctan((-1)^(3/4)*tan(d*x+c)^(1/2))/d+8*a^3*(I*A+B)/d/tan(d*x+c)^(1/2)+8/105*a^3*(23*A-21*I*B)/d/tan(d*x+c)^(3/2)-2/7*a*A*(a+I*a*tan(d*x+c))^2/d/tan(d*x+c)^(7/2)-2/35*(11*I*A+7*B)*(a^3+I*a^3*tan(d*x+c))/d/tan(d*x+c)^(5/2)
```

3.133.2 Mathematica [A] (verified)

Time = 3.66 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.70

$$\int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx =$$

$$\frac{2a^3\left(15A + 21(3iA + B) \tan(c + dx) - 35(4A - 3iB) \tan^2(c + dx) - 420i(A - iB) \tan^3(c + dx) + 420\right)}{105d \tan^{\frac{7}{2}}(c + dx)}$$

3.133.
$$\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

input `Integrate[((a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2),x]`

output `(-2*a^3*(15*A + 21*((3*I)*A + B)*Tan[c + d*x] - 35*(4*A - (3*I)*B)*Tan[c + d*x]^2 - (420*I)*(A - I*B)*Tan[c + d*x]^3 + 420*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Tan[c + d*x]^(7/2))/(105*d*Tan[c + d*x]^(7/2))`

3.133.3 Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.08, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3042, 4076, 27, 3042, 4076, 27, 3042, 4074, 27, 3042, 4012, 25, 3042, 4016, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan(c + dx)^{9/2}} dx \\ & \quad \downarrow \text{4076} \\ & \frac{2}{7} \int \frac{(i \tan(c + dx)a + a)^2 (a(11iA + 7B) - a(3A - 7iB) \tan(c + dx))}{2 \tan^{\frac{7}{2}}(c + dx) \frac{2aA(a + ia \tan(c + dx))^2}{7d \tan^{\frac{7}{2}}(c + dx)}} dx - \\ & \quad \downarrow \text{27} \\ & \frac{1}{7} \int \frac{(i \tan(c + dx)a + a)^2 (a(11iA + 7B) - a(3A - 7iB) \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx) \frac{2aA(a + ia \tan(c + dx))^2}{7d \tan^{\frac{7}{2}}(c + dx)}} dx - \\ & \quad \downarrow \text{3042} \\ & \frac{1}{7} \int \frac{(i \tan(c + dx)a + a)^2 (a(11iA + 7B) - a(3A - 7iB) \tan(c + dx))}{\tan(c + dx)^{7/2} \frac{2aA(a + ia \tan(c + dx))^2}{7d \tan^{\frac{7}{2}}(c + dx)}} dx - \end{aligned}$$

3.133. $\int \frac{(a+ia \tan(c+dx))^3 (A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

↓ 4076

$$\frac{1}{7} \left(\frac{2}{5} \int - \frac{2(i \tan(c+dx)a + a) ((23A - 21iB)a^2 + 2(6iA + 7B) \tan(c+dx)a^2)}{\frac{\tan^{\frac{5}{2}}(c+dx)}{2aA(a + ia \tan(c+dx))^2}} dx - \frac{2(7B + 11iA) (a^3 + ia^3 \tan(c+dx))}{5d \tan^{\frac{5}{2}}(c+dx)} \right)$$

↓ 27

$$\frac{1}{7} \left(-\frac{4}{5} \int \frac{(i \tan(c+dx)a + a) ((23A - 21iB)a^2 + 2(6iA + 7B) \tan(c+dx)a^2)}{\frac{\tan^{\frac{5}{2}}(c+dx)}{2aA(a + ia \tan(c+dx))^2}} dx - \frac{2(7B + 11iA) (a^3 + ia^3 \tan(c+dx))}{5d \tan^{\frac{5}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{7} \left(-\frac{4}{5} \int \frac{(i \tan(c+dx)a + a) ((23A - 21iB)a^2 + 2(6iA + 7B) \tan(c+dx)a^2)}{\frac{\tan(c+dx)^{5/2}}{2aA(a + ia \tan(c+dx))^2}} dx - \frac{2(7B + 11iA) (a^3 + ia^3 \tan(c+dx))}{5d \tan^{\frac{5}{2}}(c+dx)} \right)$$

↓ 4074

$$\frac{1}{7} \left(-\frac{4}{5} \left(\int \frac{35(a^3(iA + B) - a^3(A - iB) \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx - \frac{2a^3(23A - 21iB)}{3d \tan^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7B + 11iA) (a^3 + ia^3 \tan(c+dx))}{5d \tan^{\frac{5}{2}}(c+dx)} \right)$$

↓ 27

$$\frac{1}{7} \left(-\frac{4}{5} \left(35 \int \frac{a^3(iA + B) - a^3(A - iB) \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)} dx - \frac{2a^3(23A - 21iB)}{3d \tan^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7B + 11iA) (a^3 + ia^3 \tan(c+dx))}{5d \tan^{\frac{5}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{7} \left(-\frac{4}{5} \left(35 \int \frac{a^3(iA + B) - a^3(A - iB) \tan(c+dx)}{\tan(c+dx)^{3/2}} dx - \frac{2a^3(23A - 21iB)}{3d \tan^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7B + 11iA) (a^3 + ia^3 \tan(c+dx))}{5d \tan^{\frac{5}{2}}(c+dx)} \right)$$

3.133. $\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

↓ 4012

$$\frac{1}{7} \left(-\frac{4}{5} \left(35 \left(\int -\frac{(A-iB)a^3 + (iA+B)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(B+iA)}{d\sqrt{\tan(c+dx)}} \right) - \frac{2a^3(23A-21iB)}{3d\tan^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7B+1)}{7d\tan^{\frac{7}{2}}(c+dx)} \right)$$

↓ 25

$$\frac{1}{7} \left(-\frac{4}{5} \left(35 \left(-\int \frac{(A-iB)a^3 + (iA+B)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(B+iA)}{d\sqrt{\tan(c+dx)}} \right) - \frac{2a^3(23A-21iB)}{3d\tan^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7B+1)}{7d\tan^{\frac{7}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{7} \left(-\frac{4}{5} \left(35 \left(-\int \frac{(A-iB)a^3 + (iA+B)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(B+iA)}{d\sqrt{\tan(c+dx)}} \right) - \frac{2a^3(23A-21iB)}{3d\tan^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7B+1)}{7d\tan^{\frac{7}{2}}(c+dx)} \right)$$

↓ 4016

$$\frac{1}{7} \left(-\frac{4}{5} \left(35 \left(-\frac{2a^6(A-iB)^2 \int \frac{1}{a^3(A-iB)-a^3(iA+B)\tan(c+dx)} d\sqrt{\tan(c+dx)}}{d} - \frac{2a^3(B+iA)}{d\sqrt{\tan(c+dx)}} \right) - \frac{2a^3(23A-21iB)}{3d\tan^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7B+1)}{7d\tan^{\frac{7}{2}}(c+dx)} \right)$$

↓ 218

$$\frac{1}{7} \left(-\frac{4}{5} \left(35 \left(\frac{2\sqrt{-1}a^3(A-iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{d} - \frac{2a^3(B+iA)}{d\sqrt{\tan(c+dx)}} \right) - \frac{2a^3(23A-21iB)}{3d\tan^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7B+1)}{7d\tan^{\frac{7}{2}}(c+dx)} \right)$$

input `Int[((a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2),x]`

```
output (-2*a*A*(a + I*a*Tan[c + d*x])^2)/(7*d*Tan[c + d*x]^(7/2)) + ((-4*(35*((2*
(-1)^(1/4)*a^3*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]])/d - (2*a^3
*(I*A + B))/(d*Sqrt[Tan[c + d*x]])) - (2*a^3*(23*A - (21*I)*B))/(3*d*Tan[c
+ d*x]^(3/2))))/5 - (2*((11*I)*A + 7*B)*(a^3 + I*a^3*Tan[c + d*x]))/(5*d*
Tan[c + d*x]^(5/2))/7
```

3.133.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4012 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]
```

```
rule 4016 Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x
_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b
*Tan[e + f*x]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]
```

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

```
rule 4076 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

3.133.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.53

method	result
derivativedivides	$a^3 \left(-\frac{2A}{7 \tan(dx+c)^{\frac{7}{2}}} - \frac{2(-4iA-4B)}{\sqrt{\tan(dx+c)}} - \frac{2(3iA+B)}{5 \tan(dx+c)^{\frac{5}{2}}} - \frac{2(3iB-4A)}{3 \tan(dx+c)^{\frac{3}{2}}} + \frac{(-4iB+4A)\sqrt{2}}{\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right)} + 2 \arctan\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) \right)$
default	$a^3 \left(-\frac{2A}{7 \tan(dx+c)^{\frac{7}{2}}} - \frac{2(-4iA-4B)}{\sqrt{\tan(dx+c)}} - \frac{2(3iA+B)}{5 \tan(dx+c)^{\frac{5}{2}}} - \frac{2(3iB-4A)}{3 \tan(dx+c)^{\frac{3}{2}}} + \frac{(-4iB+4A)\sqrt{2}}{\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right)} + 2 \arctan\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) \right)$
parts	$\frac{(-iA a^3 - 3B a^3) \left(-\frac{\sqrt{2} \left(\ln\left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})\right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})\right)}{4} \right)}{d}$

```
input int((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2), x, method=_RETUR
NVERBOSE)
```

$$3.133. \int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

output $1/d*a^3*(-2/7*A/\tan(d*x+c)^{(7/2)}-2*(-4*I*A-4*B)/\tan(d*x+c)^{(1/2)}-2/5*(3*I*A+B)/\tan(d*x+c)^{(5/2)}-2/3*(3*I*B-4*A)/\tan(d*x+c)^{(3/2)}+1/4*(-4*I*B+4*A)*2^{(1/2)}*(\ln((1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}))+1/4*(4*I*A+4*B)*2^{(1/2)}*(\ln((1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}))$

3.133.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 561 vs. $2(137) = 274$.

Time = 0.29 (sec) , antiderivative size = 561, normalized size of antiderivative = 3.32

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$= 2 \left(105 \sqrt{-\frac{(iA^2 + 2AB - iB^2)a^6}{d^2}} (de^{(8i dx + 8i c)} - 4de^{(6i dx + 6i c)} + 6de^{(4i dx + 4i c)} - 4de^{(2i dx + 2i c)} + d) \log \left(-\frac{2(A + B \tan(c + dx))}{(A + B \tan(c + dx))^2} \right) \right)$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="fracas")`

```

output 2/105*(105*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(d*e^(8*I*d*x + 8*I*c) -
4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*
c) + d)*log(-2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B -
I*B^2)*a^6/d^2)*(I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt((-I*e^(2*I*d*x + 2*I
*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3)
) - 105*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(d*e^(8*I*d*x + 8*I*c) - 4*
d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c)
+ d)*log(-2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I*
B^2)*a^6/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt((-I*e^(2*I*d*x + 2*I*c
) + I)/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3))
- 2*((319*A - 273*I*B)*a^3*e^(8*I*d*x + 8*I*c) - 3*(109*A - 133*I*B)*a^3*e
^(6*I*d*x + 6*I*c) - 5*(19*A - 21*I*B)*a^3*e^(4*I*d*x + 4*I*c) + 3*(129*A
- 133*I*B)*a^3*e^(2*I*d*x + 2*I*c) - 4*(41*A - 42*I*B)*a^3)*sqrt((-I*e^(2*
I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(8*I*d*x + 8*I*c) - 4
*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c)
+ d)

```

3.133.6 Sympy [F]

$$\begin{aligned}
& \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx \\
&= -ia^3 \left(\int \left(-\frac{3A}{\tan^{\frac{7}{2}}(c + dx)} \right) dx + \int \frac{A}{\tan^{\frac{3}{2}}(c + dx)} dx + \int \left(-\frac{3B}{\tan^{\frac{5}{2}}(c + dx)} \right) dx \right. \\
&\quad \left. + \int \frac{B}{\sqrt{\tan(c + dx)}} dx + \int \frac{iA}{\tan^{\frac{9}{2}}(c + dx)} dx + \int \left(-\frac{3iA}{\tan^{\frac{5}{2}}(c + dx)} \right) dx \right. \\
&\quad \left. + \int \frac{iB}{\tan^{\frac{7}{2}}(c + dx)} dx + \int \left(-\frac{3iB}{\tan^{\frac{3}{2}}(c + dx)} \right) dx \right)
\end{aligned}$$

```

input integrate((a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c))/tan(d*x+c)**(9/2),x)

```

```

output -I*a**3*(Integral(-3*A/tan(c + d*x)**(7/2), x) + Integral(A/tan(c + d*x)**
(3/2), x) + Integral(-3*B/tan(c + d*x)**(5/2), x) + Integral(B/sqrt(tan(c
+ d*x)), x) + Integral(I*A/tan(c + d*x)**(9/2), x) + Integral(-3*I*A/tan(c
+ d*x)**(5/2), x) + Integral(I*B/tan(c + d*x)**(7/2), x) + Integral(-3*I*
B/tan(c + d*x)**(3/2), x))

```

3.133.7 Maxima [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.27

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$= \frac{105 \left(2\sqrt{2}((i+1)A - (i-1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(dx+c)}\right)\right) + 2\sqrt{2}((i+1)A - (i-1)B) \right)}{d}$$

```
input integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="maxima")
```

```
output 1/105*(105*(2*sqrt(2)*((I + 1)*A - (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I + 1)*A - (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*a^3 + 2*(420*(I*A + B)*a^3*tan(d*x + c)^3 + 35*(4*A - 3*I*B)*a^3*tan(d*x + c)^2 + 21*(-3*I*A - B)*a^3*tan(d*x + c) - 15*A*a^3)/tan(d*x + c)^(7/2)/d
```

3.133.8 Giac [A] (verification not implemented)

Time = 1.36 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.80

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$= \frac{(4i - 4) \sqrt{2}(-i A a^3 - B a^3) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{d}$$

$$= \frac{2(-420i A a^3 \tan(dx + c)^3 - 420 B a^3 \tan(dx + c)^3 - 140 A a^3 \tan(dx + c)^2 + 105i B a^3 \tan(dx + c)^2 + 63i A a^3 \tan(dx + c) + 21 B a^3 \tan(dx + c) + 15 A a^3)}{105 d \tan(dx + c)^{\frac{7}{2}}}$$

```
input integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="giac")
```

```
output (4*I - 4)*sqrt(2)*(-I*A*a^3 - B*a^3)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/d - 2/105*(-420*I*A*a^3*tan(d*x + c)^3 - 420*B*a^3*tan(d*x + c)^3 - 140*A*a^3*tan(d*x + c)^2 + 105*I*B*a^3*tan(d*x + c)^2 + 63*I*A*a^3*tan(d*x + c) + 21*B*a^3*tan(d*x + c) + 15*A*a^3)/(d*tan(d*x + c)^(7/2))
```

3.133. $\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

3.133.9 Mupad [B] (verification not implemented)

Time = 11.30 (sec) , antiderivative size = 293, normalized size of antiderivative = 1.73

$$\begin{aligned}
& \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx \\
&= -\frac{\frac{2Aa^3}{7d} + \frac{Aa^3 \tan(c+dx)6i}{5d} - \frac{8Aa^3 \tan(c+dx)^2}{3d} - \frac{Aa^3 \tan(c+dx)^3 8i}{d}}{\tan(c + dx)^{7/2}} \\
&\quad - \frac{\frac{2Ba^3}{5d} - \frac{8Ba^3 \tan(c+dx)^2}{d} + \frac{Ba^3 \tan(c+dx)2i}{d}}{\tan(c + dx)^{5/2}} \\
&\quad + \frac{\sqrt{2}Aa^3 \ln\left(Aa^3 d 8i + \sqrt{2}Aa^3 d \sqrt{\tan(c + dx)}(-4 + 4i)\right)(2 - 2i)}{d} \\
&\quad - \frac{\sqrt{-16i}Aa^3 \ln\left(Aa^3 d 8i + 2\sqrt{-16i}Aa^3 d \sqrt{\tan(c + dx)}\right)}{d} \\
&\quad + \frac{\sqrt{2}Ba^3 \ln\left(8Ba^3 d + \sqrt{2}Ba^3 d \sqrt{\tan(c + dx)}(-4 - 4i)\right)(2 + 2i)}{d} \\
&\quad - \frac{\sqrt{16i}Ba^3 \ln\left(8Ba^3 d + 2\sqrt{16i}Ba^3 d \sqrt{\tan(c + dx)}\right)}{d}
\end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3)/tan(c + d*x)^(9/2),x)`

output `(2^(1/2)*A*a^3*log(A*a^3*d*8i - 2^(1/2)*A*a^3*d*tan(c + d*x)^(1/2)*(4 - 4i))*(2 - 2i))/d - ((2*B*a^3)/(5*d) + (B*a^3*tan(c + d*x)*2i)/d - (8*B*a^3*tan(c + d*x)^2)/d)/tan(c + d*x)^(5/2) - ((2*A*a^3)/(7*d) + (A*a^3*tan(c + d*x)*6i)/(5*d) - (8*A*a^3*tan(c + d*x)^2)/(3*d) - (A*a^3*tan(c + d*x)^3*8i)/d)/tan(c + d*x)^(7/2) - ((-16i)^(1/2)*A*a^3*log(A*a^3*d*8i + 2*(-16i)^(1/2)*A*a^3*d*tan(c + d*x)^(1/2)))/d + (2^(1/2)*B*a^3*log(8*B*a^3*d - 2^(1/2)*B*a^3*d*tan(c + d*x)^(1/2)*(4 + 4i))*(2 + 2i))/d - (16i^(1/2)*B*a^3*log(8*B*a^3*d + 2*16i^(1/2)*B*a^3*d*tan(c + d*x)^(1/2)))/d`

3.134
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

3.134.1 Optimal result 1387
 3.134.2 Mathematica [A] (verified) 1388
 3.134.3 Rubi [A] (verified) 1388
 3.134.4 Maple [A] (verified) 1393
 3.134.5 Fricas [B] (verification not implemented) 1394
 3.134.6 Sympy [F] 1395
 3.134.7 Maxima [F(-2)] 1395
 3.134.8 Giac [A] (verification not implemented) 1395
 3.134.9 Mupad [B] (verification not implemented) 1396

3.134.1 Optimal result

Integrand size = 36, antiderivative size = 306

$$\begin{aligned} & \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx \\ &= -\frac{\left(\frac{1}{4} + \frac{i}{4}\right) \left((4+i)A + (1+6i)B\right) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}ad} \\ & \quad + \frac{\left(\frac{1}{4} + \frac{i}{4}\right) \left((4+i)A + (1+6i)B\right) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}ad} \\ & \quad - \frac{\left(\frac{1}{8} + \frac{i}{8}\right) \left((1+4i)A - (6+i)B\right) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{\sqrt{2}ad} \\ & \quad - \frac{\left((3-5i)A + (5+7i)B\right) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{8\sqrt{2}ad} \\ & \quad - \frac{5(iA-B)\sqrt{\tan(c+dx)}}{2ad} - \frac{(3A+7iB)\tan^{\frac{3}{2}}(c+dx)}{6ad} + \frac{(iA-B)\tan^{\frac{5}{2}}(c+dx)}{2d(a+ia \tan(c+dx))} \end{aligned}$$

output

```
(1/8+1/8*I)*((4+I)*A+(1+6*I)*B)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/a/d*2^(1/2)+(1/8+1/8*I)*((4+I)*A+(1+6*I)*B)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/a/d*2^(1/2)-(1/16+1/16*I)*((1+4*I)*A-(6+I)*B)*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a/d*2^(1/2)-1/16*((3-5*I)*A+(5+7*I)*B)*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a/d*2^(1/2)-5/2*(I*A-B)*tan(d*x+c)^(1/2)/a/d-1/6*(3*A+7*I*B)*tan(d*x+c)^(3/2)/a/d+1/2*(I*A-B)*tan(d*x+c)^(5/2)/d/(a+I*a*tan(d*x+c))
```

3.134.
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

3.134.2 Mathematica [A] (verified)

Time = 2.47 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.45

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= \frac{-3(-1)^{3/4}(A-iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c+dx)}\right) - 6(-1)^{3/4}(2A+3iB) \operatorname{arctanh}\left((-1)^{3/4} \sqrt{\tan(c+dx)}\right)}{6ad}$$

input `Integrate[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `(-3*(-1)^(3/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] - 6*(-1)^(3/4)*(2*A + (3*I)*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + (Sqrt[Tan[c + d*x]]*(-15*(A + I*B) + 4*((-3*I)*A + 2*B)*Tan[c + d*x] - (4*I)*B*Tan[c + d*x]^2))/(-I + Tan[c + d*x])/(6*a*d)`

3.134.3 Rubi [A] (verified)

Time = 0.92 (sec) , antiderivative size = 270, normalized size of antiderivative = 0.88, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.528$, Rules used = {3042, 4078, 27, 3042, 4011, 3042, 4011, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)^{5/2}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$\downarrow \text{4078}$$

$$\frac{(-B+ia) \tan^{\frac{5}{2}}(c+dx)}{2d(a+ia \tan(c+dx))} - \frac{\int \frac{1}{2} \tan^{\frac{3}{2}}(c+dx)(5a(iA-B) + a(3A+7iB) \tan(c+dx)) dx}{2a^2}$$

$$\downarrow \text{27}$$

3.134. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

$$\begin{aligned}
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \frac{\int \tan^{\frac{3}{2}}(c + dx)(5a(iA - B) + a(3A + 7iB) \tan(c + dx)) dx}{4a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \frac{\int \tan(c + dx)^{3/2}(5a(iA - B) + a(3A + 7iB) \tan(c + dx)) dx}{4a^2} \\
& \quad \downarrow \text{4011} \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \\
& \frac{\int \sqrt{\tan(c + dx)}(5a(iA - B) \tan(c + dx) - a(3A + 7iB)) dx + \frac{2a(3A + 7iB) \tan^{\frac{3}{2}}(c + dx)}{3d}}{4a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \\
& \frac{\int \sqrt{\tan(c + dx)}(5a(iA - B) \tan(c + dx) - a(3A + 7iB)) dx + \frac{2a(3A + 7iB) \tan^{\frac{3}{2}}(c + dx)}{3d}}{4a^2} \\
& \quad \downarrow \text{4011} \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \\
& \frac{\int \frac{-5a(iA - B) - a(3A + 7iB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2a(3A + 7iB) \tan^{\frac{3}{2}}(c + dx)}{3d} + \frac{10a(-B + iA) \sqrt{\tan(c + dx)}}{d}}{4a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \\
& \frac{\int \frac{-5a(iA - B) - a(3A + 7iB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2a(3A + 7iB) \tan^{\frac{3}{2}}(c + dx)}{3d} + \frac{10a(-B + iA) \sqrt{\tan(c + dx)}}{d}}{4a^2} \\
& \quad \downarrow \text{4017} \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \\
& \frac{2 \int \frac{-a(5(iA - B) + (3A + 7iB) \tan(c + dx))}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} + \frac{2a(3A + 7iB) \tan^{\frac{3}{2}}(c + dx)}{3d} + \frac{10a(-B + iA) \sqrt{\tan(c + dx)}}{d}}{4a^2} \\
& \quad \downarrow \text{25}
\end{aligned}$$

3.134. $\int \frac{\tan^{\frac{5}{2}}(c + dx)(A + B \tan(c + dx))}{a + ia \tan(c + dx)} dx$

$$\begin{aligned}
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \\
 & \frac{2 \int \frac{a(5(iA-B) + (3A+7iB) \tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{2a(3A+7iB) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{10a(-B+iA)\sqrt{\tan(c+dx)}}{d}}{4a^2} \\
 & \quad \downarrow 27 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \\
 & \frac{2a \int \frac{5(iA-B) + (3A+7iB) \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{2a(3A+7iB) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{10a(-B+iA)\sqrt{\tan(c+dx)}}{d}}{4a^2} \\
 & \quad \downarrow 1482 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \\
 & \frac{2a \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((1+4i)A - (6+i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \left(\frac{1}{2} + \frac{i}{2} \right) ((4+i)A + (1+6i)B) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} + \frac{2a(3A+7iB) \tan^{\frac{3}{2}}(c+dx)}{3d}}{4a^2} \\
 & \quad \downarrow 1476 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \\
 & \frac{2a \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((1+4i)A - (6+i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \left(\frac{1}{2} + \frac{i}{2} \right) ((4+i)A + (1+6i)B) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d}}{4a^2} \\
 & \quad \downarrow 1082 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \\
 & \frac{2a \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((1+4i)A - (6+i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \left(\frac{1}{2} + \frac{i}{2} \right) ((4+i)A + (1+6i)B) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)+1} d(1+\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d}}{4a^2} \\
 & \quad \downarrow 217 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \\
 & \frac{2a \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((1+4i)A - (6+i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \left(\frac{1}{2} + \frac{i}{2} \right) ((4+i)A + (1+6i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d}}{4a^2} \\
 & \quad \downarrow 1479
 \end{aligned}$$

3.134. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

$$\begin{aligned}
& \frac{2a \left(\left(\frac{1}{2} + \frac{i}{2}\right) ((1+4i)A - (6+i)B) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \left(\frac{1}{2} + \frac{i}{2}\right) ((4+i)A + (1+6i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d} \\
& \quad \downarrow 25 \\
& \frac{2a \left(\left(\frac{1}{2} + \frac{i}{2}\right) ((1+4i)A - (6+i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \left(\frac{1}{2} + \frac{i}{2}\right) ((4+i)A + (1+6i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d} \\
& \quad \downarrow 27 \\
& \frac{2a \left(\left(\frac{1}{2} + \frac{i}{2}\right) ((1+4i)A - (6+i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \left(\frac{1}{2} + \frac{i}{2}\right) ((4+i)A + (1+6i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d} \\
& \quad \downarrow 1103 \\
& \frac{2a \left(\left(\frac{1}{2} + \frac{i}{2}\right) ((4+i)A + (1+6i)B) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) + \left(\frac{1}{2} + \frac{i}{2}\right) ((1+4i)A - (6+i)B) \left(\frac{\log\left(\frac{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}{2\sqrt{2}}\right)}{\sqrt{2}} \right) \right)}{d}
\end{aligned}$$

input `Int[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `((I*A - B)*Tan[c + d*x]^(5/2))/(2*d*(a + I*a*Tan[c + d*x])) - ((-2*a*((1/2 + I/2)*((4 + I)*A + (1 + 6*I)*B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + (1/2 + I/2)*((1 + 4*I)*A - (6 + I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/d + (10*a*(I*A - B)*Sqrt[Tan[c + d*x]])/d + (2*a*(3*A + (7*I)*B)*Tan[c + d*x]^(3/2))/(3*d))/(4*a^2)`

$$3.134. \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

3.134.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*(a + b*Tan[e + f*x])^m/(f*m), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4078 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.134.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.54

method	result
derivativedivides	$-\frac{2iB \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + 2B(\sqrt{\tan(dx+c)}) - 2iA(\sqrt{\tan(dx+c)}) - \frac{i \left(-\frac{i(iB+A)(\sqrt{\tan(dx+c)})}{\tan(dx+c)-i} + \frac{4(2iA-3B) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} \right)}{2}$
default	$-\frac{2iB \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + 2B(\sqrt{\tan(dx+c)}) - 2iA(\sqrt{\tan(dx+c)}) - \frac{i \left(-\frac{i(iB+A)(\sqrt{\tan(dx+c)})}{\tan(dx+c)-i} + \frac{4(2iA-3B) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} \right)}{2}$

input `int(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,method=_RETURNV ERBOSE)`

$$3.134. \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

output $\frac{1}{d/a*(-2/3*I*B*\tan(dx+c)^{(3/2)}+2*B*\tan(dx+c)^{(1/2)}-2*I*A*\tan(dx+c)^{(1/2)}-1/2*I*(-I*(A+I*B)*\tan(dx+c)^{(1/2)}/(\tan(dx+c)-I)+4*(2*I*A-3*B)/(2^{(1/2)}-I*2^{(1/2)}))*\arctan(2*\tan(dx+c)^{(1/2)}/(2^{(1/2)}-I*2^{(1/2)})))+4*(-1/4*A+1/4*I*B)/(2^{(1/2)}+I*2^{(1/2)}))*\arctan(2*\tan(dx+c)^{(1/2)}/(2^{(1/2)}+I*2^{(1/2)}))}$

3.134.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 707 vs. $2(221) = 442$.

Time = 0.28 (sec) , antiderivative size = 707, normalized size of antiderivative = 2.31

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$$

$$3 \left(ade^{(4i dx+4i c)} + ade^{(2i dx+2i c)} \right) \sqrt{\frac{iA^2+2AB-iB^2}{a^2d^2}} \log \left(\frac{2 \left((ade^{(2i dx+2i c)}+ad) \sqrt{\frac{-ie^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}+1}} \sqrt{\frac{iA^2+2AB-iB^2}{a^2d^2}} + (A-iB) \right)}{iA+B} \right)$$

input `integrate(tan(dx+c)^(5/2)*(A+B*tan(dx+c))/(a+I*a*tan(dx+c)),x, algorithm="fracas")`

output $\frac{1}{24}*(3*(a*d*e^{(4*I*d*x + 4*I*c)} + a*d*e^{(2*I*d*x + 2*I*c)})*\sqrt{((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2))*\log(2*((a*d*e^{(2*I*d*x + 2*I*c)} + a*d)*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))*\sqrt{((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2)) + (A - I*B)*e^{(2*I*d*x + 2*I*c)}*e^{(-2*I*d*x - 2*I*c)/(I*A + B)} - 3*(a*d*e^{(4*I*d*x + 4*I*c)} + a*d*e^{(2*I*d*x + 2*I*c)})*\sqrt{((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2))*\log(-2*((a*d*e^{(2*I*d*x + 2*I*c)} + a*d)*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))*\sqrt{((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2)) - (A - I*B)*e^{(2*I*d*x + 2*I*c)}*e^{(-2*I*d*x - 2*I*c)/(I*A + B)} - 6*(a*d*e^{(4*I*d*x + 4*I*c)} + a*d*e^{(2*I*d*x + 2*I*c)})*\sqrt{((-4*I*A^2 + 12*A*B + 9*I*B^2)/(a^2*d^2))*\log(-((a*d*e^{(2*I*d*x + 2*I*c)} + a*d)*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))*\sqrt{((-4*I*A^2 + 12*A*B + 9*I*B^2)/(a^2*d^2)) + 2*A + 3*I*B}*e^{(-2*I*d*x - 2*I*c)/(a*d)} + 6*(a*d*e^{(4*I*d*x + 4*I*c)} + a*d*e^{(2*I*d*x + 2*I*c)})*\sqrt{((-4*I*A^2 + 12*A*B + 9*I*B^2)/(a^2*d^2))*\log(((a*d*e^{(2*I*d*x + 2*I*c)} + a*d)*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))*\sqrt{((-4*I*A^2 + 12*A*B + 9*I*B^2)/(a^2*d^2)) - 2*A - 3*I*B}*e^{(-2*I*d*x - 2*I*c)/(a*d)} + 2*((-27*I*A + 19*B)*e^{(4*I*d*x + 4*I*c)} - 2*(15*I*A - 19*B)*e^{(2*I*d*x + 2*I*c)} - 3*I*A + 3*B)*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))})/(a*d*e^{(4*I*d*x + 4*I*c)} + a*d*e^{(2*I*d*x + 2*I*c)})}$

3.134. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$

3.134.6 Sympy [F]

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = -\frac{i\left(\int \frac{A\tan^{\frac{5}{2}}(c+dx)}{\tan(c+dx)-i} dx + \int \frac{B\tan^{\frac{7}{2}}(c+dx)}{\tan(c+dx)-i} dx\right)}{a}$$

input `integrate(tan(d*x+c)**(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)`

output `-I*(Integral(A*tan(c+d*x)**(5/2)/(tan(c+d*x)-I),x)+Integral(B*tan(c+d*x)**(7/2)/(tan(c+d*x)-I),x))/a`

3.134.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.134.8 Giac [A] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.53

$$\begin{aligned} & \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx \\ &= \frac{(i+1)\sqrt{2}(2A+3iB)\arctan\left(\left(\frac{1}{2}i+\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{2ad} \\ & \quad + \frac{(i-1)\sqrt{2}(A-iB)\arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{4ad} \\ & \quad - \frac{A\sqrt{\tan(dx+c)}+iB\sqrt{\tan(dx+c)}}{2ad(\tan(dx+c)-i)} \\ & \quad - \frac{2\left(iBa^2d^2\tan(dx+c)^{\frac{3}{2}}+3iAa^2d^2\sqrt{\tan(dx+c)}-3Ba^2d^2\sqrt{\tan(dx+c)}\right)}{3a^3d^3} \end{aligned}$$

3.134. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")`

output `(1/2*I + 1/2)*sqrt(2)*(2*A + 3*I*B)*arctan((1/2*I + 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a*d) + (1/4*I - 1/4)*sqrt(2)*(A - I*B)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a*d) - 1/2*(A*sqrt(tan(d*x + c)) + I*B*sqrt(tan(d*x + c)))/(a*d*(tan(d*x + c) - I)) - 2/3*(I*B*a^2*d^2*tan(d*x + c)^(3/2) + 3*I*A*a^2*d^2*sqrt(tan(d*x + c)) - 3*B*a^2*d^2*sqrt(tan(d*x + c)))/(a^3*d^3)`

3.134.9 Mupad [B] (verification not implemented)

Time = 13.77 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.00

$$\begin{aligned} & \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx \\ &= \operatorname{atan}\left(\frac{ad\sqrt{\tan(c+dx)}\sqrt{-\frac{A^2 1i}{a^2 d^2}}}{A}\right) \sqrt{-\frac{A^2 1i}{a^2 d^2}} 2i \\ &\quad - \operatorname{atan}\left(\frac{ad\sqrt{\tan(c+dx)}\sqrt{\frac{A^2 1i}{16 a^2 d^2}}}{A}\right) \sqrt{\frac{A^2 1i}{16 a^2 d^2}} 2i \\ &\quad + \operatorname{atan}\left(\frac{2ad\sqrt{\tan(c+dx)}\sqrt{\frac{B^2 9i}{4 a^2 d^2}}}{3B}\right) \sqrt{\frac{B^2 9i}{4 a^2 d^2}} 2i \\ &\quad + \operatorname{atan}\left(\frac{4ad\sqrt{\tan(c+dx)}\sqrt{-\frac{B^2 1i}{16 a^2 d^2}}}{B}\right) \sqrt{-\frac{B^2 1i}{16 a^2 d^2}} 2i \\ &\quad - \frac{A\sqrt{\tan(c+dx)} 2i}{ad} + \frac{2B\sqrt{\tan(c+dx)}}{ad} - \frac{B\tan(c+dx)^{3/2} 2i}{3ad} \\ &\quad - \frac{A\sqrt{\tan(c+dx)} 1i}{2ad(1+\tan(c+dx) 1i)} + \frac{B\sqrt{\tan(c+dx)}}{2ad(1+\tan(c+dx) 1i)} \end{aligned}$$

input `int((tan(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i),x)`

output $\text{atan}((a*d*\tan(c + d*x))^{1/2}*(-(A^2*1i)/(a^2*d^2))^{1/2}*1i)/A*(-(A^2*1i)/(a^2*d^2))^{1/2}*2i - \text{atan}((a*d*\tan(c + d*x))^{1/2}*((A^2*1i)/(16*a^2*d^2))^{1/2}*4i)/A*((A^2*1i)/(16*a^2*d^2))^{1/2}*2i + \text{atan}((2*a*d*\tan(c + d*x))^{1/2}*((B^2*9i)/(4*a^2*d^2))^{1/2})/(3*B))*((B^2*9i)/(4*a^2*d^2))^{1/2}*2i + \text{atan}((4*a*d*\tan(c + d*x))^{1/2}*(-(B^2*1i)/(16*a^2*d^2))^{1/2})/B*(-(B^2*1i)/(16*a^2*d^2))^{1/2}*2i - (A*\tan(c + d*x))^{1/2}*2i)/(a*d) + (2*B*\tan(c + d*x))^{1/2})/(a*d) - (B*\tan(c + d*x))^{3/2}*2i)/(3*a*d) - (A*\tan(c + d*x))^{1/2}*1i)/(2*a*d*(\tan(c + d*x)*1i + 1)) + (B*\tan(c + d*x))^{1/2})/(2*a*d*(\tan(c + d*x)*1i + 1))$

3.134. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

3.135
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

3.135.1 Optimal result 1398
 3.135.2 Mathematica [A] (verified) 1399
 3.135.3 Rubi [A] (verified) 1399
 3.135.4 Maple [A] (verified) 1404
 3.135.5 Fricas [B] (verification not implemented) 1404
 3.135.6 Sympy [F] 1405
 3.135.7 Maxima [F(-2)] 1405
 3.135.8 Giac [A] (verification not implemented) 1406
 3.135.9 Mupad [B] (verification not implemented) 1407

3.135.1 Optimal result

Integrand size = 36, antiderivative size = 275

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= -\frac{((1-3i)A+(3+5i)B) \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{4\sqrt{2}ad}$$

$$- \frac{\left(\frac{1}{4}+\frac{i}{4}\right) \left((1+2i)A-(4+i)B\right) \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}ad}$$

$$- \frac{\left(\frac{1}{8}+\frac{i}{8}\right) \left((2+i)A+(1+4i)B\right) \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{\sqrt{2}ad}$$

$$+ \frac{\left(\frac{1}{8}+\frac{i}{8}\right) \left((2+i)A+(1+4i)B\right) \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{\sqrt{2}ad}$$

$$- \frac{(A+5iB)\sqrt{\tan(c+dx)}}{2ad} + \frac{(iA-B)\tan^{\frac{3}{2}}(c+dx)}{2d(a+ia \tan(c+dx))}$$

output

```
1/8*((1-3*I)*A+(3+5*I)*B)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/a/d*2^(1/2)-
(1/8+1/8*I)*((1+2*I)*A-(4+I)*B)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/a/d*2^(
1/2)-(1/16+1/16*I)*((2+I)*A+(1+4*I)*B)*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d
*x+c))/a/d*2^(1/2)+(1/16+1/16*I)*((2+I)*A+(1+4*I)*B)*ln(1+2^(1/2)*tan(d*x+
c)^(1/2)+tan(d*x+c))/a/d*2^(1/2)-1/2*(A+5*I*B)*tan(d*x+c)^(1/2)/a/d+1/2*(I
*A-B)*tan(d*x+c)^(3/2)/d/(a+I*a*tan(d*x+c))
```

3.135.
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

3.135.2 Mathematica [A] (verified)

Time = 1.56 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.41

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= \frac{\sqrt[4]{-1}(A-iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c+dx)}\right) - 2\sqrt[4]{-1}(A+2iB) \operatorname{arctanh}\left((-1)^{3/4} \sqrt{\tan(c+dx)}\right) + i\sqrt{\tan(c+dx)}}{2ad}$$

input `Integrate[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `((-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] - 2*(-1)^(1/4)*(A + (2*I)*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + (I*Sqrt[Tan[c + d*x]]*(A + (5*I)*B - 4*B*Tan[c + d*x]))/(-I + Tan[c + d*x]))/(2*a*d)`

3.135.3 Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 237, normalized size of antiderivative = 0.86, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.472$, Rules used = {3042, 4078, 27, 3042, 4011, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$\downarrow 3042$$

$$\int \frac{\tan(c+dx)^{3/2}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$\downarrow 4078$$

$$\frac{(-B+iA) \tan^{\frac{3}{2}}(c+dx)}{2d(a+ia \tan(c+dx))} - \frac{\int \frac{1}{2} \sqrt{\tan(c+dx)} (3a(iA-B) + a(A+5iB) \tan(c+dx)) dx}{2a^2}$$

$$\downarrow 27$$

$$\frac{(-B+iA) \tan^{\frac{3}{2}}(c+dx)}{2d(a+ia \tan(c+dx))} - \frac{\int \sqrt{\tan(c+dx)} (3a(iA-B) + a(A+5iB) \tan(c+dx)) dx}{4a^2}$$

3.135. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \frac{\int \sqrt{\tan(c + dx)}(3a(iA - B) + a(A + 5iB) \tan(c + dx)) dx}{4a^2} \\
 & \downarrow 4011 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \frac{\int \frac{3a(iA - B) \tan(c + dx) - a(A + 5iB)}{\sqrt{\tan(c + dx)}} dx + \frac{2a(A + 5iB) \sqrt{\tan(c + dx)}}{d}}{4a^2} \\
 & \downarrow 3042 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \frac{\int \frac{3a(iA - B) \tan(c + dx) - a(A + 5iB)}{\sqrt{\tan(c + dx)}} dx + \frac{2a(A + 5iB) \sqrt{\tan(c + dx)}}{d}}{4a^2} \\
 & \downarrow 4017 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \frac{2 \int -\frac{a(A + 5iB - 3(iA - B) \tan(c + dx))}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \frac{2a(A + 5iB) \sqrt{\tan(c + dx)}}{d}}{4a^2} \\
 & \downarrow 25 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \frac{\frac{2a(A + 5iB) \sqrt{\tan(c + dx)}}{d} - \frac{2 \int \frac{a(A + 5iB - 3(iA - B) \tan(c + dx))}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{4a^2}}{4a^2} \\
 & \downarrow 27 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \frac{\frac{2a(A + 5iB) \sqrt{\tan(c + dx)}}{d} - \frac{2a \int \frac{A + 5iB - 3(iA - B) \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{4a^2}}{4a^2} \\
 & \downarrow 1482 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \frac{\frac{2a(A + 5iB) \sqrt{\tan(c + dx)}}{d} - \frac{2a \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((2+i)A + (1+4i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \frac{1}{2} ((1-3i)A + (3+5i)B) \int \frac{\tan(c + dx) + 1}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} \right)}{4a^2}}{4a^2} \\
 & \downarrow 1476 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{2d(a + ia \tan(c + dx))} - \frac{\frac{2a(A + 5iB) \sqrt{\tan(c + dx)}}{d} - \frac{2a \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((2+i)A + (1+4i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \frac{1}{2} ((1-3i)A + (3+5i)B) \left(\frac{1}{2} \int \frac{1}{\tan(c + dx) - \sqrt{2}\sqrt{\tan(c + dx)}} \right)}{4a^2}}{4a^2} \\
 & \downarrow 1082
 \end{aligned}$$

3.135. $\int \frac{\tan^{\frac{3}{2}}(c + dx)(A + B \tan(c + dx))}{a + ia \tan(c + dx)} dx$

$$\frac{\frac{2a(A+5iB)\sqrt{\tan(c+dx)}}{d} - \frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{2d(a+ia\tan(c+dx))}}{4a^2} = \frac{2a\left(\left(\frac{1}{2}+\frac{i}{2}\right)((2+i)A+(1+4i)B)\int\frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}+\frac{1}{2}((1-3i)A+(3+5i)B)\left(\frac{\int\frac{1}{-\tan(c+dx)-1}d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}}-\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right)\right)\right)}{d}$$

↓ 217

$$\frac{\frac{2a(A+5iB)\sqrt{\tan(c+dx)}}{d} - \frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{2d(a+ia\tan(c+dx))}}{4a^2} = \frac{2a\left(\left(\frac{1}{2}+\frac{i}{2}\right)((2+i)A+(1+4i)B)\int\frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}+\frac{1}{2}((1-3i)A+(3+5i)B)\left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}}-\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right)\right)\right)}{d}$$

↓ 1479

$$\frac{\frac{2a(A+5iB)\sqrt{\tan(c+dx)}}{d} - \frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{2d(a+ia\tan(c+dx))}}{4a^2} = \frac{2a\left(\left(\frac{1}{2}+\frac{i}{2}\right)((2+i)A+(1+4i)B)\left(-\frac{\int\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}}{2\sqrt{2}}-\frac{\int\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}}{2\sqrt{2}}\right)\right)}{d}$$

↓ 25

$$\frac{\frac{2a(A+5iB)\sqrt{\tan(c+dx)}}{d} - \frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{2d(a+ia\tan(c+dx))}}{4a^2} = \frac{2a\left(\left(\frac{1}{2}+\frac{i}{2}\right)((2+i)A+(1+4i)B)\left(\frac{\int\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}}{2\sqrt{2}}+\frac{\int\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}}{2\sqrt{2}}\right)\right)}{d}$$

↓ 27

$$\frac{\frac{2a(A+5iB)\sqrt{\tan(c+dx)}}{d} - \frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{2d(a+ia\tan(c+dx))}}{4a^2} = \frac{2a\left(\left(\frac{1}{2}+\frac{i}{2}\right)((2+i)A+(1+4i)B)\left(\frac{\int\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}}{2\sqrt{2}}+\frac{1}{2}\int\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}\right)\right)}{d}$$

↓ 1103

$$\frac{\frac{2a(A+5iB)\sqrt{\tan(c+dx)}}{d} - \frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{2d(a+ia\tan(c+dx))}}{4a^2} = \frac{2a\left(\frac{1}{2}((1-3i)A+(3+5i)B)\left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}}-\frac{\arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}}\right)\right)+\left(\frac{1}{2}+\frac{i}{2}\right)((2+i)A+(1+4i)B)\left(\log\left|\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right|\right)}{d}$$

3.135. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$

input `Int[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `-1/4*((-2*a*(((1 - 3*I)*A + (3 + 5*I)*B)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 + (1/2 + I/2)*((2 + I)*A + (1 + 4*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/d + (2*a*(A + (5*I)*B)*Sqrt[Tan[c + d*x]]/d)/a^2 + ((I*A - B)*Tan[c + d*x]^(3/2))/(2*d*(a + I*a*Tan[c + d*x]))`

3.135.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4078 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.135.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.52

method	result
derivativedivides	$\frac{-2iB(\sqrt{\tan(dx+c)}) + \frac{i \left(-\frac{i(A-B)(\sqrt{\tan(dx+c)})}{\tan(dx+c)-i} - \frac{4(2iB+A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} \right)}{2}}{da} + \frac{4\left(-\frac{iA}{4} - \frac{B}{4}\right) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}$
default	$\frac{-2iB(\sqrt{\tan(dx+c)}) + \frac{i \left(-\frac{i(A-B)(\sqrt{\tan(dx+c)})}{\tan(dx+c)-i} - \frac{4(2iB+A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} \right)}{2}}{da} + \frac{4\left(-\frac{iA}{4} - \frac{B}{4}\right) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}$

input `int(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,method=_RETURNV
ERBOSE)`

output `1/d/a*(-2*I*B*tan(d*x+c)^(1/2)+1/2*I*(-I*(I*A-B)*tan(d*x+c)^(1/2)/(tan(d*x
+c)-I)-4*(A+2*I*B)/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-
I*2^(1/2))))+4*(-1/4*I*A-1/4*B)/(2^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1
/2)/(2^(1/2)+I*2^(1/2)))`

3.135.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 621 vs. 2(198) = 396.

Time = 0.28 (sec) , antiderivative size = 621, normalized size of antiderivative = 2.26

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= \left(ad \sqrt{\frac{-iA^2-2AB+iB^2}{a^2d^2}} e^{(2i dx+2i c)} \log \left(-\frac{2 \left((i ad e^{(2i dx+2i c)}+i ad) \sqrt{\frac{-i e^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}+1}} \sqrt{\frac{-iA^2-2AB+iB^2}{a^2d^2}} - (A-iB) e^{(2i dx+2i c)} \right)}{iA+B} \right) \right)$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorith
m="fracas")`

3.135. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

```
output 1/8*(a*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(-2*((I*a*d*e^(2*I*d*x + 2*I*c) + I*a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B) - a*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(-2*((-I*a*d*e^(2*I*d*x + 2*I*c) - I*a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) + 2*a*d*sqrt((I*A^2 - 4*A*B - 4*I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(((a*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*A^2 - 4*A*B - 4*I*B^2)/(a^2*d^2)) + I*A - 2*B)*e^(-2*I*d*x - 2*I*c)/(a*d)) - 2*a*d*sqrt((I*A^2 - 4*A*B - 4*I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(-((a*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*A^2 - 4*A*B - 4*I*B^2)/(a^2*d^2)) - I*A + 2*B)*e^(-2*I*d*x - 2*I*c)/(a*d)) - 2*((A + 9*I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*e^(-2*I*d*x - 2*I*c)/(a*d)
```

3.135.6 Sympy [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = -\frac{i \left(\int \frac{A \tan^{\frac{3}{2}}(c+dx)}{\tan(c+dx)-i} dx + \int \frac{B \tan^{\frac{5}{2}}(c+dx)}{\tan(c+dx)-i} dx \right)}{a}$$

```
input integrate(tan(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)
```

```
output -I*(Integral(A*tan(c + d*x)**(3/2)/(tan(c + d*x) - I), x) + Integral(B*tan(c + d*x)**(5/2)/(tan(c + d*x) - I), x))/a
```

3.135.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="maxima")
```

3.135. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

3.135.8 Giac [A] (verification not implemented)

Time = 0.48 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.44

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$$

$$= -\frac{(i+1)\sqrt{2}(iA-2B)\arctan\left(\left(\frac{1}{2}i+\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{2ad}$$

$$-\frac{(i-1)\sqrt{2}(-iA-B)\arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{4ad}$$

$$-\frac{2iB\sqrt{\tan(dx+c)}}{ad} - \frac{-iA\sqrt{\tan(dx+c)}+B\sqrt{\tan(dx+c)}}{2ad(\tan(dx+c)-i)}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")`

output `-(1/2*I + 1/2)*sqrt(2)*(I*A - 2*B)*arctan((1/2*I + 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a*d) - (1/4*I - 1/4)*sqrt(2)*(-I*A - B)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a*d) - 2*I*B*sqrt(tan(d*x + c))/(a*d) - 1/2*(-I*A*sqrt(tan(d*x + c)) + B*sqrt(tan(d*x + c)))/(a*d*(tan(d*x + c) - I))`

3.135.9 Mupad [B] (verification not implemented)

Time = 12.59 (sec) , antiderivative size = 270, normalized size of antiderivative = 0.98

$$\begin{aligned}
& \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx \\
&= -\operatorname{atan}\left(\frac{2ad \sqrt{\tan(c+dx)} \sqrt{\frac{A^2 li}{4a^2 d^2}}}{A}\right) \sqrt{\frac{A^2 li}{4a^2 d^2}} 2i \\
&\quad - \operatorname{atan}\left(\frac{4ad \sqrt{\tan(c+dx)} \sqrt{-\frac{A^2 li}{16a^2 d^2}}}{A}\right) \sqrt{-\frac{A^2 li}{16a^2 d^2}} 2i \\
&\quad + \operatorname{atan}\left(\frac{ad \sqrt{\tan(c+dx)} \sqrt{-\frac{B^2 li}{a^2 d^2}} li}{B}\right) \sqrt{-\frac{B^2 li}{a^2 d^2}} 2i \\
&\quad - \operatorname{atan}\left(\frac{ad \sqrt{\tan(c+dx)} \sqrt{\frac{B^2 li}{16a^2 d^2}} 4i}{B}\right) \sqrt{\frac{B^2 li}{16a^2 d^2}} 2i - \frac{B \sqrt{\tan(c+dx)} 2i}{ad} \\
&\quad - \frac{A \sqrt{\tan(c+dx)}}{2ad(1+\tan(c+dx) li)} - \frac{B \sqrt{\tan(c+dx)} li}{2ad(1+\tan(c+dx) li)}
\end{aligned}$$

```
input int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i),x)
```

```
output atan((a*d*tan(c + d*x)^(1/2)*(-(B^2*1i)/(a^2*d^2))^(1/2)*1i)/B)*(-(B^2*1i)/(a^2*d^2))^(1/2)*2i - atan((4*a*d*tan(c + d*x)^(1/2)*(-(A^2*1i)/(16*a^2*d^2))^(1/2))/A)*(-(A^2*1i)/(16*a^2*d^2))^(1/2)*2i - atan((2*a*d*tan(c + d*x)^(1/2)*((A^2*1i)/(4*a^2*d^2))^(1/2))/A)*((A^2*1i)/(4*a^2*d^2))^(1/2)*2i - atan((a*d*tan(c + d*x)^(1/2)*((B^2*1i)/(16*a^2*d^2))^(1/2)*4i)/B)*((B^2*1i)/(16*a^2*d^2))^(1/2)*2i - (B*tan(c + d*x)^(1/2)*2i)/(a*d) - (A*tan(c + d*x)^(1/2))/(2*a*d*(tan(c + d*x)*1i + 1)) - (B*tan(c + d*x)^(1/2)*1i)/(2*a*d*(tan(c + d*x)*1i + 1))
```

3.136
$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

3.136.1 Optimal result 1408
 3.136.2 Mathematica [A] (verified) 1409
 3.136.3 Rubi [A] (verified) 1409
 3.136.4 Maple [A] (verified) 1413
 3.136.5 Fricas [B] (verification not implemented) 1414
 3.136.6 Sympy [F] 1414
 3.136.7 Maxima [F(-2)] 1415
 3.136.8 Giac [A] (verification not implemented) 1415
 3.136.9 Mupad [B] (verification not implemented) 1416

3.136.1 Optimal result

Integrand size = 36, antiderivative size = 236

$$\begin{aligned} & \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx \\ &= -\frac{\left(\frac{1}{4}-\frac{i}{4}\right)(A+(2-i)B) \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}ad} \\ & \quad + \frac{\left(\frac{1}{4}-\frac{i}{4}\right)(A+(2-i)B) \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}ad} \\ & \quad + \frac{\left(\frac{1}{8}+\frac{i}{8}\right)(A-(2+i)B) \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{\sqrt{2}ad} \\ & \quad - \frac{\left(\frac{1}{8}+\frac{i}{8}\right)(A-(2+i)B) \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{\sqrt{2}ad} \\ & \quad + \frac{(iA-B)\sqrt{\tan(c+dx)}}{2d(a+ia \tan(c+dx))} \end{aligned}$$

```
output (1/8-1/8*I)*(A+(2-I)*B)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/a/d*2^(1/2)+(1/8-1/8*I)*(A+(2-I)*B)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/a/d*2^(1/2)+(1/16+1/16*I)*(A-(2+I)*B)*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a/d*2^(1/2)-(1/16+1/16*I)*(A-(2+I)*B)*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a/d*2^(1/2)+1/2*(I*A-B)*tan(d*x+c)^(1/2)/d/(a+I*a*tan(d*x+c))
```

3.136.2 Mathematica [A] (verified)

Time = 1.12 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.41

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$$

$$= \frac{\sqrt[4]{-1}(iA+B)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right) - 2\sqrt[4]{-1}B\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right) + \frac{(A+iB)\sqrt{\tan(c+dx)}}{-i+\tan(c+dx)}}{2ad}$$

input `Integrate[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `((-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] - 2*(-1)^(1/4)*B*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + ((A + I*B)*Sqrt[Tan[c + d*x]])/(-I + Tan[c + d*x]))/(2*a*d)`

3.136.3 Rubi [A] (verified)Time = 0.59 (sec) , antiderivative size = 206, normalized size of antiderivative = 0.87, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3042, 4078, 27, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$$

$$\downarrow 3042$$

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$$

$$\downarrow 4078$$

$$\frac{(-B+iA)\sqrt{\tan(c+dx)}}{2d(a+ia\tan(c+dx))} - \frac{\int \frac{a(iA-B)-a(A-3iB)\tan(c+dx)}{2\sqrt{\tan(c+dx)}} dx}{2a^2}$$

$$\downarrow 27$$

$$\frac{(-B+iA)\sqrt{\tan(c+dx)}}{2d(a+ia\tan(c+dx))} - \frac{\int \frac{a(iA-B)-a(A-3iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{4a^2}$$

3.136. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{2d(a + ia \tan(c + dx))} - \frac{\int \frac{a(iA - B) - a(A - 3iB)\tan(c + dx)}{\sqrt{\tan(c + dx)}} dx}{4a^2} \\
& \downarrow \text{4017} \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{2d(a + ia \tan(c + dx))} - \frac{\int \frac{a(iA - B - (A - 3iB)\tan(c + dx))}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{2a^2d} \\
& \downarrow \text{27} \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{2d(a + ia \tan(c + dx))} - \frac{\int \frac{iA - B - (A - 3iB)\tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{2ad} \\
& \downarrow \text{1482} \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{2d(a + ia \tan(c + dx))} - \frac{(\frac{1}{2} + \frac{i}{2})(A - (2 + i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - (\frac{1}{2} - \frac{i}{2})(A + (2 - i)B) \int \frac{\tan(c + dx) + 1}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{2ad} \\
& \downarrow \text{1476} \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{2d(a + ia \tan(c + dx))} - \frac{(\frac{1}{2} + \frac{i}{2})(A - (2 + i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - (\frac{1}{2} - \frac{i}{2})(A + (2 - i)B) \left(\frac{1}{2} \int \frac{1}{\tan(c + dx) - \sqrt{2}\sqrt{\tan(c + dx) + 1}} d\sqrt{\tan(c + dx)} \right)}{2ad} \\
& \downarrow \text{1082} \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{2d(a + ia \tan(c + dx))} - \frac{(\frac{1}{2} + \frac{i}{2})(A - (2 + i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - (\frac{1}{2} - \frac{i}{2})(A + (2 - i)B) \left(\frac{\int \frac{1}{-\tan(c + dx) - 1} d(1 - \sqrt{2}\sqrt{\tan(c + dx)})}{\sqrt{2}} \right)}{2ad} \\
& \downarrow \text{217} \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{2d(a + ia \tan(c + dx))} - \frac{(\frac{1}{2} + \frac{i}{2})(A - (2 + i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - (\frac{1}{2} - \frac{i}{2})(A + (2 - i)B) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\tan(c + dx) + 1}}{\sqrt{2}})}{\sqrt{2}} - \frac{\arctan(\dots)}{\dots} \right)}{2ad} \\
& \downarrow \text{1479}
\end{aligned}$$

3.136. $\int \frac{\sqrt{\tan(c + dx)}(A + B \tan(c + dx))}{a + ia \tan(c + dx)} dx$

$$\frac{\frac{(-B + iA)\sqrt{\tan(c + dx)}}{2d(a + ia \tan(c + dx))} - \left(\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)A}{2ad}$$

↓ 25

$$\frac{\frac{(-B + iA)\sqrt{\tan(c + dx)}}{2d(a + ia \tan(c + dx))} - \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)A}{2ad}$$

↓ 27

$$\frac{\frac{(-B + iA)\sqrt{\tan(c + dx)}}{2d(a + ia \tan(c + dx))} - \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c + dx)} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)A}{2ad}$$

↓ 1103

$$\frac{\frac{(-B + iA)\sqrt{\tan(c + dx)}}{2d(a + ia \tan(c + dx))} - \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)(A + (2 - i)B)}{2ad}$$

input `Int[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `-1/2*((-1/2 + I/2)*(A + (2 - I)*B)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + (1/2 + I/2)*(A - (2 + I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/(a*d) + ((I*A - B)*Sqrt[Tan[c + d*x]])/(2*d*(a + I*a*Tan[c + d*x]))`

3.136.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] & & NeQ[c^2 + d^2, 0]`

rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.136.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.53

method	result	size
derivativedivides	$i \frac{\left(-\frac{i(iB+A)(\sqrt{\tan(dx+c)})}{\tan(dx+c)-i} - \frac{4B \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2}-i\sqrt{2}}\right)}{\sqrt{2}-i\sqrt{2}} \right)}{2} + \frac{4\left(\frac{A}{4} - \frac{iB}{4}\right) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2}+i\sqrt{2}}\right)}{\sqrt{2}+i\sqrt{2}}$	124
default	$i \frac{\left(-\frac{i(iB+A)(\sqrt{\tan(dx+c)})}{\tan(dx+c)-i} - \frac{4B \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2}-i\sqrt{2}}\right)}{\sqrt{2}-i\sqrt{2}} \right)}{2} + \frac{4\left(\frac{A}{4} - \frac{iB}{4}\right) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2}+i\sqrt{2}}\right)}{\sqrt{2}+i\sqrt{2}}$	124

input `int(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,method=_RETURNV ERBOSE)`

output `1/d/a*(1/2*I*(-I*(A+I*B)*tan(d*x+c)^(1/2)/(tan(d*x+c)-I)-4*B/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2))))+4*(1/4*A-1/4*I*B)/(2^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))`

3.136.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 570 vs. $2(177) = 354$.

Time = 0.27 (sec) , antiderivative size = 570, normalized size of antiderivative = 2.42

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx =$$

$$\left(ad \sqrt{\frac{iA^2+2AB-iB^2}{a^2d^2}} e^{(2i dx+2i c)} \log \left(\frac{2 \left((ade^{(2i dx+2i c)}+ad) \sqrt{\frac{-ie^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}+1}} \sqrt{\frac{iA^2+2AB-iB^2}{a^2d^2}} + (A-iB)e^{(2i dx+2i c)} \right) e^{(-2i dx-2i c)}}{iA+B} \right) \right)$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="fricas")`

output

```
-1/8*(a*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(
2*((a*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2
*I*d*x + 2*I*c) + 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2)) + (A - I*B)*
e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - a*d*sqrt((I*A^2 + 2
*A*B - I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(-2*((a*d*e^(2*I*d*x + 2*I
*c) + a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sq
rt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(
-2*I*d*x - 2*I*c)/(I*A + B)) - 2*a*d*sqrt(I*B^2/(a^2*d^2))*e^(2*I*d*x + 2*
I*c)*log(((a*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I
)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(I*B^2/(a^2*d^2)) + I*B)*e^(-2*I*d*x - 2*
I*c)/(a*d)) + 2*a*d*sqrt(I*B^2/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(-((a*d*e
^(2*I*d*x + 2*I*c) + a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x +
2*I*c) + 1))*sqrt(I*B^2/(a^2*d^2)) - I*B)*e^(-2*I*d*x - 2*I*c)/(a*d)) - 2*
((I*A - B)*e^(2*I*d*x + 2*I*c) + I*A - B)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I
)/(e^(2*I*d*x + 2*I*c) + 1))*e^(-2*I*d*x - 2*I*c)/(a*d)
```

3.136.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = -\frac{i \left(\int \frac{A\sqrt{\tan(c+dx)}}{\tan(c+dx)-i} dx + \int \frac{B \tan^{\frac{3}{2}}(c+dx)}{\tan(c+dx)-i} dx \right)}{a}$$

input `integrate(tan(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)`

output `-I*(Integral(A*sqrt(tan(c + d*x))/(tan(c + d*x) - I), x) + Integral(B*tan(c + d*x)**(3/2)/(tan(c + d*x) - I), x))/a`

3.136.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.136.8 Giac [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.41

$$\begin{aligned} & \int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx \\ &= -\frac{(i-1)\sqrt{2}B\arctan\left(\left(\frac{1}{2}i+\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{2ad} \\ & \quad -\frac{(i-1)\sqrt{2}(A-iB)\arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{4ad} \\ & \quad +\frac{A\sqrt{\tan(dx+c)}+iB\sqrt{\tan(dx+c)}}{2ad(\tan(dx+c)-i)} \end{aligned}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")`

output $-(1/2*I - 1/2)*\sqrt{2}*B*\arctan((1/2*I + 1/2)*\sqrt{2}*\sqrt{\tan(d*x + c)})/(a*d) - (1/4*I - 1/4)*\sqrt{2}*(A - I*B)*\arctan(-(1/2*I - 1/2)*\sqrt{2}*\sqrt{\tan(d*x + c)})/(a*d) + 1/2*(A*\sqrt{\tan(d*x + c)} + I*B*\sqrt{\tan(d*x + c)})/(a*d*(\tan(d*x + c) - I))$

3.136.9 Mupad [B] (verification not implemented)

Time = 10.23 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.78

$$\begin{aligned} & \int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx \\ &= -\operatorname{atan}\left(\frac{2ad\sqrt{\tan(c+dx)}\sqrt{\frac{B^2 li}{4a^2 d^2}}}{B}\right)\sqrt{\frac{B^2 li}{4a^2 d^2}} 2i \\ &\quad - \operatorname{atan}\left(\frac{4ad\sqrt{\tan(c+dx)}\sqrt{-\frac{B^2 li}{16a^2 d^2}}}{B}\right)\sqrt{-\frac{B^2 li}{16a^2 d^2}} 2i \\ &\quad - \frac{2\sqrt{\frac{1}{16}i}A\operatorname{atanh}\left(4\sqrt{\frac{1}{16}i}\sqrt{\tan(c+dx)}\right)}{ad} \\ &\quad + \frac{A\sqrt{\tan(c+dx)}li}{2ad(1+\tan(c+dx)li)} - \frac{B\sqrt{\tan(c+dx)}}{2ad(1+\tan(c+dx)li)} \end{aligned}$$

input `int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i),x)`

output $(A*\tan(c + d*x)^{(1/2)*1i)/(2*a*d*(\tan(c + d*x)*1i + 1)) - \operatorname{atan}((4*a*d*\tan(c + d*x)^{(1/2)*(-B^2*1i)/(16*a^2*d^2)}^{(1/2)})/B)*(-B^2*1i)/(16*a^2*d^2)^{(1/2)*2i} - (2*(1i/16)^{(1/2)*A*\operatorname{atanh}(4*(1i/16)^{(1/2)*\tan(c + d*x)^{(1/2)})})/(a*d) - \operatorname{atan}((2*a*d*\tan(c + d*x)^{(1/2)*((B^2*1i)/(4*a^2*d^2)}^{(1/2)})/B)*((B^2*1i)/(4*a^2*d^2)^{(1/2)*2i} - (B*\tan(c + d*x)^{(1/2)})/(2*a*d*(\tan(c + d*x)*1i + 1))$

3.137
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))} dx$$

3.137.1 Optimal result 1417
 3.137.2 Mathematica [A] (verified) 1418
 3.137.3 Rubi [A] (verified) 1418
 3.137.4 Maple [A] (verified) 1422
 3.137.5 Fricas [B] (verification not implemented) 1423
 3.137.6 Sympy [F(-2)] 1423
 3.137.7 Maxima [F(-2)] 1424
 3.137.8 Giac [A] (verification not implemented) 1424
 3.137.9 Mupad [B] (verification not implemented) 1425

3.137.1 Optimal result

Integrand size = 36, antiderivative size = 234

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))} dx \\ &= -\frac{\left(\frac{1}{4} - \frac{i}{4}\right) ((2 + i)A + B) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}ad} \\ & \quad + \frac{\left(\frac{1}{4} - \frac{i}{4}\right) ((2 + i)A + B) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}ad} \\ & \quad - \frac{((3 + i)A - (1 + i)B) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{8\sqrt{2}ad} \\ & \quad + \frac{((3 + i)A - (1 + i)B) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{8\sqrt{2}ad} \\ & \quad + \frac{(A + iB)\sqrt{\tan(c + dx)}}{2d(a + ia \tan(c + dx))} \end{aligned}$$

output

```
(1/8-1/8*I)*((2+I)*A+B)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/a/d*2^(1/2)+(1/8-1/8*I)*((2+I)*A+B)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/a/d*2^(1/2)-1/16*((3+I)*A-(1+I)*B)*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a/d*2^(1/2)+1/16*((3+I)*A-(1+I)*B)*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a/d*2^(1/2)+1/2*(A+I*B)*tan(d*x+c)^(1/2)/d/(a+I*a*tan(d*x+c))
```


3.137.2 Mathematica [A] (verified)

Time = 1.60 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.41

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))} dx$$

$$= \frac{-\sqrt[4]{-1}(A - iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right) - 2\sqrt[4]{-1}A \operatorname{arctanh}\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right) + \frac{(-iA+B)\sqrt{\tan(c + dx)}}{-i + \tan(c + dx)}}{2ad}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])),x]`

output `((-((-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]) - 2*(-1)^(1/4)*A*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + (((-I)*A + B)*Sqrt[Tan[c + d*x]])/(-I + Tan[c + d*x]))/(2*a*d)`

3.137.3 Rubi [A] (verified)Time = 0.58 (sec) , antiderivative size = 204, normalized size of antiderivative = 0.87, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3042, 4079, 27, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))} dx$$

$$\downarrow 3042$$

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))} dx$$

$$\downarrow 4079$$

$$\frac{\int \frac{a(3A-iB)-a(iA-B) \tan(c+dx)}{2\sqrt{\tan(c+dx)}} dx}{2a^2} + \frac{(A + iB) \sqrt{\tan(c + dx)}}{2d(a + ia \tan(c + dx))}$$

$$\downarrow 27$$

$$\frac{\int \frac{a(3A-iB)-a(iA-B) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{4a^2} + \frac{(A + iB) \sqrt{\tan(c + dx)}}{2d(a + ia \tan(c + dx))}$$

3.137. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))} dx$

$$\begin{aligned}
 & \int \frac{a(3A-iB)-a(iA-B)\tan(c+dx)}{\sqrt{\tan(c+dx)}4a^2} dx + \frac{(A+iB)\sqrt{\tan(c+dx)}}{2d(a+ia\tan(c+dx))} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{a(3A-iB-(iA-B)\tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{2d(a+ia\tan(c+dx))} \\
 & \quad \downarrow \text{4017} \\
 & \int \frac{3A-iB-(iA-B)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{2d(a+ia\tan(c+dx))} \\
 & \quad \downarrow \text{27} \\
 & \quad \downarrow \text{1482} \\
 & \frac{\frac{1}{2}((3+i)A-(1+i)B)\int\frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}+(\frac{1}{2}-\frac{i}{2})(B+(2+i)A)\int\frac{\tan(c+dx)+1}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}}{2ad} + \\
 & \quad \frac{(A+iB)\sqrt{\tan(c+dx)}}{2d(a+ia\tan(c+dx))} \\
 & \quad \downarrow \text{1476} \\
 & \frac{\frac{1}{2}((3+i)A-(1+i)B)\int\frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}+(\frac{1}{2}-\frac{i}{2})(B+(2+i)A)\left(\frac{1}{2}\int\frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}\right)}{2ad} \\
 & \quad \frac{(A+iB)\sqrt{\tan(c+dx)}}{2d(a+ia\tan(c+dx))} \\
 & \quad \downarrow \text{1082} \\
 & \frac{\frac{1}{2}((3+i)A-(1+i)B)\int\frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}+(\frac{1}{2}-\frac{i}{2})(B+(2+i)A)\left(\frac{\int\frac{1}{-\tan(c+dx)-1}d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}}\right)}{2ad} \\
 & \quad \frac{(A+iB)\sqrt{\tan(c+dx)}}{2d(a+ia\tan(c+dx))} \\
 & \quad \downarrow \text{217} \\
 & \frac{\frac{1}{2}((3+i)A-(1+i)B)\int\frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}+(\frac{1}{2}-\frac{i}{2})(B+(2+i)A)\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}}-\arctan(\dots)\right)}{2ad} \\
 & \quad \frac{(A+iB)\sqrt{\tan(c+dx)}}{2d(a+ia\tan(c+dx))}
 \end{aligned}$$

3.137. $\int \frac{A+B\tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} dx$

$$\begin{aligned} & \downarrow 1479 \\ & \frac{\frac{1}{2}((3+i)A - (1+i)B) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \left(\frac{1}{2} - \frac{i}{2}\right) (B + (2+i)A)}{2ad}}{\frac{(A+iB)\sqrt{\tan(c+dx)}}{2d(a+ia\tan(c+dx))}} \end{aligned}$$

$$\begin{aligned} & \downarrow 25 \\ & \frac{\frac{1}{2}((3+i)A - (1+i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \left(\frac{1}{2} - \frac{i}{2}\right) (B + (2+i)A)}{2ad}}{\frac{(A+iB)\sqrt{\tan(c+dx)}}{2d(a+ia\tan(c+dx))}} \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{\frac{1}{2}((3+i)A - (1+i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \left(\frac{1}{2} - \frac{i}{2}\right) (B + (2+i)A)}{2ad}}{\frac{(A+iB)\sqrt{\tan(c+dx)}}{2d(a+ia\tan(c+dx))}} \end{aligned}$$

$$\begin{aligned} & \downarrow 1103 \\ & \frac{\left(\frac{1}{2} - \frac{i}{2}\right) (B + (2+i)A) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) + \frac{1}{2}((3+i)A - (1+i)B) \left(\frac{\log(\tan(c+dx))}{\sqrt{2}} \right)}{2ad}}{\frac{(A+iB)\sqrt{\tan(c+dx)}}{2d(a+ia\tan(c+dx))}} \end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])),x]`

output `((1/2 - I/2)*((2 + I)*A + B)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + (((3 + I)*A - (1 + I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/(2*a*d) + ((A + I*B)*Sqrt[Tan[c + d*x]])/(2*d*(a + I*a*Tan[c + d*x]))`

$$3.137. \quad \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))} dx$$

3.137.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.137.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.52

method	result	size
derivativedivides	$\frac{-\frac{i(iB+A)(\sqrt{\tan(dx+c)})}{2(\tan(dx+c)-i)} - \frac{2iA \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} + \frac{4\left(\frac{iA}{4} + \frac{B}{4}\right) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}}{da}$	121
default	$\frac{-\frac{i(iB+A)(\sqrt{\tan(dx+c)})}{2(\tan(dx+c)-i)} - \frac{2iA \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} + \frac{4\left(\frac{iA}{4} + \frac{B}{4}\right) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}}{da}$	121

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)), x, method=_RETURNV ERBOSE)`

output `1/d/a*(-1/2*I*(A+I*B)*tan(d*x+c)^(1/2)/(tan(d*x+c)-I)-2*I*A/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))+4*(1/4*I*A+1/4*B)/(2^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))`

$$3.137. \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))} dx$$

3.137.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 571 vs. $2(177) = 354$.

Time = 0.27 (sec) , antiderivative size = 571, normalized size of antiderivative = 2.44

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)(a + ia \tan(c + dx))}} dx =$$

$$\left(ad \sqrt{\frac{-iA^2 - 2AB + iB^2}{a^2 d^2}} e^{(2i dx + 2i c)} \log \left(-\frac{2 \left((i a d e^{(2i dx + 2i c)} + i a d) \sqrt{\frac{-i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} + 1}} \sqrt{\frac{-i A^2 - 2AB + i B^2}{a^2 d^2}} - (A - i B) e^{(2i dx + 2i c)} \right)}{i A + B} \right) \right)$$

```
input integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)),x, algorithm
m="fricas")
```

```
output -1/8*(a*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log
(-2*((I*a*d*e^(2*I*d*x + 2*I*c) + I*a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)
/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2)) - (A
- I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - a*d*sqrt((-I
*A^2 - 2*A*B + I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(-2*((-I*a*d*e^(2*
I*d*x + 2*I*c) - I*a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*
I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2)) - (A - I*B)*e^(2*I*d*x
+ 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 2*a*d*sqrt(I*A^2/(a^2*d^2))*e
^(2*I*d*x + 2*I*c)*log(((a*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt((-I*e^(2*I*d*
x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(I*A^2/(a^2*d^2)) + I*A)*e^
(-2*I*d*x - 2*I*c)/(a*d)) + 2*a*d*sqrt(I*A^2/(a^2*d^2))*e^(2*I*d*x + 2*I*c
)*log(-((a*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/
(e^(2*I*d*x + 2*I*c) + 1))*sqrt(I*A^2/(a^2*d^2)) - I*A)*e^(-2*I*d*x - 2*I*
c)/(a*d)) - 2*((A + I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*sqrt((-I*e^(2*I*d*
x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*e^(-2*I*d*x - 2*I*c)/(a*d)
```

3.137.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)(a + ia \tan(c + dx))}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+I*a*tan(d*x+c)),x)`

output Exception raised: TypeError >> Invalid comparison of non-real -I

3.137.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)),x, algorithm="maxima")`

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

3.137.8 Giac [A] (verification not implemented)

Time = 0.59 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.42

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))} dx \\ &= -\frac{(i - 1) \sqrt{2} A \arctan\left(\left(\frac{1}{2}i + \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{2 ad} \\ & \quad - \frac{(i - 1) \sqrt{2} (i A + B) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{4 ad} \\ & \quad - \frac{i A \sqrt{\tan(dx + c)} - B \sqrt{\tan(dx + c)}}{2 ad(\tan(dx + c) - i)} \end{aligned}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)),x, algorithm="giac")`

output $-(1/2*I - 1/2)*\sqrt{2}*A*\arctan((1/2*I + 1/2)*\sqrt{2}*\sqrt{\tan(d*x + c)})/(a*d) - (1/4*I - 1/4)*\sqrt{2}*(I*A + B)*\arctan(-(1/2*I - 1/2)*\sqrt{2}*\sqrt{\tan(d*x + c)})/(a*d) - 1/2*(I*A*\sqrt{\tan(d*x + c)} - B*\sqrt{\tan(d*x + c)})/(a*d*(\tan(d*x + c) - I))$

3.137. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))} dx$

3.137.9 Mupad [B] (verification not implemented)

Time = 9.11 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.79

$$\begin{aligned}
& \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))} dx \\
&= -\operatorname{atan}\left(\frac{2ad\sqrt{\tan(c + dx)}\sqrt{\frac{A^2 1i}{4a^2 d^2}}}{A}\right)\sqrt{\frac{A^2 1i}{4a^2 d^2}} 2i \\
&+ \operatorname{atan}\left(\frac{4ad\sqrt{\tan(c + dx)}\sqrt{-\frac{A^2 1i}{16a^2 d^2}}}{A}\right)\sqrt{-\frac{A^2 1i}{16a^2 d^2}} 2i \\
&- \frac{2\sqrt{\frac{1}{16}i}B \operatorname{atanh}\left(4\sqrt{\frac{1}{16}i}\sqrt{\tan(c + dx)}\right)}{ad} \\
&+ \frac{A\sqrt{\tan(c + dx)}}{2ad(1 + \tan(c + dx) 1i)} + \frac{B\sqrt{\tan(c + dx) 1i}}{2ad(1 + \tan(c + dx) 1i)}
\end{aligned}$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)),x)`

output `atan((4*a*d*tan(c + d*x)^(1/2)*(-(A^2*1i)/(16*a^2*d^2))^(1/2))/A)*(-(A^2*1i)/(16*a^2*d^2))^(1/2)*2i - atan((2*a*d*tan(c + d*x)^(1/2)*((A^2*1i)/(4*a^2*d^2))^(1/2))/A)*((A^2*1i)/(4*a^2*d^2))^(1/2)*2i - (2*(1i/16)^(1/2)*B*atanh(4*(1i/16)^(1/2)*tan(c + d*x)^(1/2)))/(a*d) + (A*tan(c + d*x)^(1/2))/(2*a*d*(tan(c + d*x)*1i + 1)) + (B*tan(c + d*x)^(1/2)*1i)/(2*a*d*(tan(c + d*x)*1i + 1))`

3.138
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} dx$$

3.138.1 Optimal result 1426
 3.138.2 Mathematica [C] (verified) 1427
 3.138.3 Rubi [A] (verified) 1427
 3.138.4 Maple [A] (verified) 1432
 3.138.5 Fracas [B] (verification not implemented) 1433
 3.138.6 Sympy [F(-2)] 1434
 3.138.7 Maxima [F(-2)] 1434
 3.138.8 Giac [A] (verification not implemented) 1434
 3.138.9 Mupad [B] (verification not implemented) 1435

3.138.1 Optimal result

Integrand size = 36, antiderivative size = 267

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$= \frac{((5 + 3i)A - (3 - i)B) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{4\sqrt{2}ad}$$

$$+ \frac{((-5 - 3i)A + (3 - i)B) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{4\sqrt{2}ad}$$

$$- \frac{\left(\frac{1}{8} - \frac{i}{8}\right) \left((4 + i)A + (1 + 2i)B\right) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{\sqrt{2}ad}$$

$$+ \frac{\left((5 - 3i)A + (3 + i)B\right) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{8\sqrt{2}ad}$$

$$- \frac{5A + iB}{2ad\sqrt{\tan(c + dx)}} + \frac{A + iB}{2d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))}$$

```
output -1/8*((5+3*I)*A+(-3+I)*B)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/a/d*2^(1/2)+
1/8*((-5-3*I)*A+(3-I)*B)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/a/d*2^(1/2)+(-
1/16+1/16*I)*((4+I)*A+(1+2*I)*B)*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))
/a/d*2^(1/2)+1/16*((5-3*I)*A+(3+I)*B)*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*
x+c))/a/d*2^(1/2)+1/2*(-5*A-I*B)/a/d/tan(d*x+c)^(1/2)+1/2*(A+I*B)/d/tan(d*
x+c)^(1/2)/(a+I*a*tan(d*x+c))
```

3.138.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.38 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.43

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$= \frac{-iA + B - 2(2A + iB) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -i \tan(c + dx)\right) (-i + \tan(c + dx)) - (A - iB) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, i \tan(c + dx)\right) (-i + \tan(c + dx))}{2ad\sqrt{\tan(c + dx)}(-i + \tan(c + dx))}$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])),x]`

output `((-I)*A + B - 2*(2*A + I*B)*Hypergeometric2F1[-1/2, 1, 1/2, (-I)*Tan[c + d*x]]*(-I + Tan[c + d*x]) - (A - I*B)*Hypergeometric2F1[-1/2, 1, 1/2, I*Tan[c + d*x]]*(-I + Tan[c + d*x]))/(2*a*d*Sqrt[Tan[c + d*x]]*(-I + Tan[c + d*x]))`

3.138.3 Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 237, normalized size of antiderivative = 0.89, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.472$, Rules used = {3042, 4079, 27, 3042, 4012, 25, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2}(a + ia \tan(c + dx))} dx$$

$$\downarrow \text{4079}$$

$$\frac{\int \frac{a(5A + iB) - 3a(iA - B) \tan(c + dx)}{2 \tan^{\frac{3}{2}}(c + dx)} dx}{2a^2} + \frac{A + iB}{2d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))}$$

$$\downarrow \text{27}$$

3.138. $\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx$

$$\begin{aligned}
& \frac{\int \frac{a(5A+iB)-3a(iA-B)\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)} dx}{4a^2} + \frac{A+iB}{2d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(5A+iB)-3a(iA-B)\tan(c+dx)}{\tan(c+dx)^{3/2}} dx}{4a^2} + \frac{A+iB}{2d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} \\
& \quad \downarrow \text{4012} \\
& \frac{\int -\frac{3a(iA-B)+a(5A+iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{2a(5A+iB)}{d\sqrt{\tan(c+dx)}}}{4a^2} + \frac{A+iB}{2d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} \\
& \quad \downarrow \text{25} \\
& \frac{-\int \frac{3a(iA-B)+a(5A+iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{2a(5A+iB)}{d\sqrt{\tan(c+dx)}}}{4a^2} + \frac{A+iB}{2d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} \\
& \quad \downarrow \text{3042} \\
& \frac{-\int \frac{3a(iA-B)+a(5A+iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{2a(5A+iB)}{d\sqrt{\tan(c+dx)}}}{4a^2} + \frac{A+iB}{2d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} \\
& \quad \downarrow \text{4017} \\
& \frac{-\frac{2\int \frac{a(3(iA-B)+(5A+iB)\tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} - \frac{2a(5A+iB)}{d\sqrt{\tan(c+dx)}}}{4a^2} + \frac{A+iB}{2d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} \\
& \quad \downarrow \text{27} \\
& \frac{-\frac{2a\int \frac{3(iA-B)+(5A+iB)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} - \frac{2a(5A+iB)}{d\sqrt{\tan(c+dx)}}}{4a^2} + \frac{A+iB}{2d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} \\
& \quad \downarrow \text{1482} \\
& \frac{2a\left(\frac{1}{2}((5+3i)A-(3-i)B)\int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \left(\frac{1}{2}-\frac{i}{2}\right)((4+i)A+(1+2i)B)\int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right) - \frac{2a(5A+iB)}{d\sqrt{\tan(c+dx)}}}{4a^2} + \\
& \quad \frac{A+iB}{2d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} \\
& \quad \downarrow \text{1476}
\end{aligned}$$

3.138. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))} dx$

$$\begin{aligned}
 & \frac{2a \left(\frac{1}{2}((5+3i)A - (3-i)B) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \left(\frac{1}{2} - \frac{i}{2} \right) ((4+i)A + (1+2i)B) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} \\
 & \frac{A + iB}{2d\sqrt{\tan(c+dx)}(a + ia \tan(c+dx))} \quad 4a^2 \\
 & \quad \downarrow \text{1082} \\
 & \frac{2a \left(\frac{1}{2}((5+3i)A - (3-i)B) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2} \right) ((4+i)A + (1+2i)B) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} \\
 & \frac{A + iB}{2d\sqrt{\tan(c+dx)}(a + ia \tan(c+dx))} \quad 4a^2 \\
 & \quad \downarrow \text{217} \\
 & \frac{2a \left(\frac{1}{2}((5+3i)A - (3-i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2} \right) ((4+i)A + (1+2i)B) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} \\
 & \frac{A + iB}{2d\sqrt{\tan(c+dx)}(a + ia \tan(c+dx))} \quad 4a^2 \\
 & \quad \downarrow \text{1479} \\
 & \frac{2a \left(\frac{1}{2}((5+3i)A - (3-i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2} \right) ((4+i)A + (1+2i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d} \\
 & \frac{A + iB}{2d\sqrt{\tan(c+dx)}(a + ia \tan(c+dx))} \quad 4a^2 \\
 & \quad \downarrow \text{25} \\
 & \frac{2a \left(\frac{1}{2}((5+3i)A - (3-i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2} \right) ((4+i)A + (1+2i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d} \\
 & \frac{A + iB}{2d\sqrt{\tan(c+dx)}(a + ia \tan(c+dx))} \quad 4a^2 \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.138. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

$$\frac{2a \left(\frac{1}{2}((5+3i)A-(3-i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2}-\frac{i}{2}\right)((4+i)A+(1+2i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} dx}{2\sqrt{2}} \right) \right)}{d} \frac{A+iB}{4a^2}$$

$$\frac{A+iB}{2d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))}$$

↓ 1103

$$\frac{2a \left(\frac{1}{2}((5+3i)A-(3-i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2}-\frac{i}{2}\right)((4+i)A+(1+2i)B) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d} \frac{A+iB}{4a^2}$$

$$\frac{A+iB}{2d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))}$$

```
input Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])),x]
```

```
output ((-2*a*(((5 + 3*I)*A - (3 - I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]/Sqrt[2]] + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]/Sqrt[2]]))/2 - (1/2 - I/2)*((4 + I)*A + (1 + 2*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/d - (2*a*(5*A + I*B))/(d*Sqrt[Tan[c + d*x]])/(4*a^2) + (A + I*B)/(2*d*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x]))
```

3.138.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

```
rule 4017 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_
)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sq
rt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] &
& NeQ[c^2 + d^2, 0]
```

```
rule 4079 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.138.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.52

method	result	size
derivativedivides	$\frac{\frac{i(iA-B)(\sqrt{\tan(dx+c)})}{2 \tan(dx+c)-2i} - \frac{2(iB+2A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}}}{da} - \frac{2A}{\sqrt{\tan(dx+c)}} + \frac{4\left(-\frac{A}{4} + \frac{iB}{4}\right) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}$	140
default	$\frac{\frac{i(iA-B)(\sqrt{\tan(dx+c)})}{2 \tan(dx+c)-2i} - \frac{2(iB+2A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}}}{da} - \frac{2A}{\sqrt{\tan(dx+c)}} + \frac{4\left(-\frac{A}{4} + \frac{iB}{4}\right) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}$	140

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 1/d/a*(1/2*I*(I*A-B)*tan(d*x+c)^(1/2)/(tan(d*x+c)-I)-2*(2*A+I*B)/(2^(1/2)-
I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))-2*A/tan(d*x+c)^(
1/2)+4*(-1/4*A+1/4*I*B)/(2^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(
1/2)+I*2^(1/2))))
```

$$3.138. \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} dx$$

3.138.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 703 vs. $2(200) = 400$.

Time = 0.27 (sec) , antiderivative size = 703, normalized size of antiderivative = 2.63

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$= \frac{(ade^{(4i dx + 4i c)} - ade^{(2i dx + 2i c)}) \sqrt{\frac{i A^2 + 2 AB - i B^2}{a^2 d^2}} \log \left(\frac{2 \left((ade^{(2i dx + 2i c)} + ad) \sqrt{\frac{-i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} + 1}} \sqrt{\frac{i A^2 + 2 AB - i B^2}{a^2 d^2}} + (A - i B) e^{(2i dx + 2i c)} \right)}{i A + B} \right)}{1}$$

```
input integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c)),x, algorithm="fricas")
```

```
output 1/8*((a*d*e^(4*I*d*x + 4*I*c) - a*d*e^(2*I*d*x + 2*I*c))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2))*log(2*((a*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - (a*d*e^(4*I*d*x + 4*I*c) - a*d*e^(2*I*d*x + 2*I*c))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2))*log(-2*((a*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) + 2*(a*d*e^(4*I*d*x + 4*I*c) - a*d*e^(2*I*d*x + 2*I*c))*sqrt((-4*I*A^2 + 4*A*B + I*B^2)/(a^2*d^2))*log(((a*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-4*I*A^2 + 4*A*B + I*B^2)/(a^2*d^2)) + 2*A + I*B)*e^(-2*I*d*x - 2*I*c)/(a*d)) - 2*(a*d*e^(4*I*d*x + 4*I*c) - a*d*e^(2*I*d*x + 2*I*c))*sqrt((-4*I*A^2 + 4*A*B + I*B^2)/(a^2*d^2))*log(-((a*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-4*I*A^2 + 4*A*B + I*B^2)/(a^2*d^2)) - 2*A - I*B)*e^(-2*I*d*x - 2*I*c)/(a*d)) + 2*((-9*I*A + B)*e^(4*I*d*x + 4*I*c) - 8*I*A*e^(2*I*d*x + 2*I*c) + I*A - B)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(a*d*e^(4*I*d*x + 4*I*c) - a*d*e^(2*I*d*x + 2*I*c))
```


3.138.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(3/2)/(a+I*a*tan(d*x+c)),x)`

output `Exception raised: TypeError >> Invalid comparison of non-real -I`

3.138.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.138.8 Giac [A] (verification not implemented)

Time = 0.77 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.42

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx \\ &= -\frac{(i + 1) \sqrt{2}(2A + iB) \arctan\left(\left(\frac{1}{2}i + \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{2ad} \\ &+ \frac{(i - 1) \sqrt{2}(A - iB) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{4ad} \\ &+ \frac{-5iA \tan(dx + c) + B \tan(dx + c) - 4A}{2\left(i \tan(dx + c)^{\frac{3}{2}} + \sqrt{\tan(dx + c)}\right)ad} \end{aligned}$$

3.138. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c)),x, algorithm="giac")`

output $-(1/2*I + 1/2)*\sqrt{2}*(2*A + I*B)*\arctan((1/2*I + 1/2)*\sqrt{2}*\sqrt{\tan(d*x + c)})/(a*d) + (1/4*I - 1/4)*\sqrt{2}*(A - I*B)*\arctan(-(1/2*I - 1/2)*\sqrt{2}*\sqrt{\tan(d*x + c)})/(a*d) + 1/2*(-5*I*A*\tan(d*x + c) + B*\tan(d*x + c) - 4*A)/((I*\tan(d*x + c))^(3/2) + \sqrt{\tan(d*x + c)})*a*d$

3.138.9 Mupad [B] (verification not implemented)

Time = 9.16 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.00

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx \\ &= 2 \operatorname{atanh} \left(\frac{ad \sqrt{\tan(c + dx)} \sqrt{-\frac{A^2 1i}{a^2 d^2}}}{A} \right) \sqrt{-\frac{A^2 1i}{a^2 d^2}} \\ &+ 2 \operatorname{atanh} \left(\frac{4ad \sqrt{\tan(c + dx)} \sqrt{\frac{A^2 1i}{16 a^2 d^2}}}{A} \right) \sqrt{\frac{A^2 1i}{16 a^2 d^2}} \\ &- \operatorname{atan} \left(\frac{2ad \sqrt{\tan(c + dx)} \sqrt{\frac{B^2 1i}{4 a^2 d^2}}}{B} \right) \sqrt{\frac{B^2 1i}{4 a^2 d^2}} 2i \\ &+ \operatorname{atan} \left(\frac{4ad \sqrt{\tan(c + dx)} \sqrt{-\frac{B^2 1i}{16 a^2 d^2}}}{B} \right) \sqrt{-\frac{B^2 1i}{16 a^2 d^2}} 2i \\ &- \frac{\frac{2A}{ad} + \frac{A \tan(c+dx) 5i}{2ad}}{\sqrt{\tan(c + dx) + \tan(c + dx)^{3/2} 1i}} + \frac{B \sqrt{\tan(c + dx)}}{2ad (1 + \tan(c + dx) 1i)} \end{aligned}$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + a*tan(c + d*x)*1i)),x)`

output

$$\begin{aligned}
& 2*\operatorname{atanh}((a*d*\tan(c + d*x)^{(1/2)}*(-(A^2*1i)/(a^2*d^2))^{(1/2)})/A)*(-(A^2*1i) \\
& / (a^2*d^2))^{(1/2)} + 2*\operatorname{atanh}((4*a*d*\tan(c + d*x)^{(1/2)}*((A^2*1i)/(16*a^2*d^2))^{(1/2)})/A)*((A^2*1i)/(16*a^2*d^2))^{(1/2)} - \operatorname{atan}((2*a*d*\tan(c + d*x)^{(1/2)}*((B^2*1i)/(4*a^2*d^2))^{(1/2)})/B)*((B^2*1i)/(4*a^2*d^2))^{(1/2)}*2i + \operatorname{atan} \\
& ((4*a*d*\tan(c + d*x)^{(1/2)}*(-(B^2*1i)/(16*a^2*d^2))^{(1/2)})/B)*(-(B^2*1i)/(16*a^2*d^2))^{(1/2)}*2i - ((2*A)/(a*d) + (A*\tan(c + d*x)*5i)/(2*a*d))/(\tan(c \\
& + d*x)^{(1/2)} + \tan(c + d*x)^{(3/2)}*1i) + (B*\tan(c + d*x)^{(1/2)})/(2*a*d*(\tan(c + d*x)*1i + 1))
\end{aligned}$$

3.138. $\int \frac{A+B \tan(c+dx)}{\tan^3(c+dx)(a+ia \tan(c+dx))} dx$

$$3.139 \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$$

3.139.1 Optimal result 1437
 3.139.2 Mathematica [C] (verified) 1438
 3.139.3 Rubi [A] (verified) 1438
 3.139.4 Maple [A] (verified) 1444
 3.139.5 Fricas [B] (verification not implemented) 1444
 3.139.6 Sympy [F] 1445
 3.139.7 Maxima [F(-2)] 1446
 3.139.8 Giac [A] (verification not implemented) 1446
 3.139.9 Mupad [B] (verification not implemented) 1447

3.139.1 Optimal result

Integrand size = 36, antiderivative size = 296

$$\begin{aligned} & \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx \\ &= \frac{((7-5i)A+(5+3i)B) \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{4\sqrt{2}ad} \\ & \quad - \frac{\left(\frac{1}{4}-\frac{i}{4}\right) ((6+i)A+(1+4i)B) \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}ad} \\ & \quad + \frac{((7+5i)A-(5-3i)B) \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{8\sqrt{2}ad} \\ & \quad + \frac{((-7-5i)A+(5-3i)B) \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{8\sqrt{2}ad} \\ & \quad - \frac{7A+3iB}{6ad \tan^{\frac{3}{2}}(c+dx)} + \frac{5(iA-B)}{2ad\sqrt{\tan(c+dx)}} + \frac{A+iB}{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} \end{aligned}$$

output

```
-1/8*((7-5*I)*A+(5+3*I)*B)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/a/d*2^(1/2)
+(-1/8+1/8*I)*((6+I)*A+(1+4*I)*B)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/a/d*2
^(1/2)+1/16*((7+5*I)*A+(-5+3*I)*B)*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c
))/a/d*2^(1/2)+1/16*((-7-5*I)*A+(5-3*I)*B)*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+t
an(d*x+c))/a/d*2^(1/2)+5/2*(I*A-B)/a/d/tan(d*x+c)^(1/2)+1/6*(-7*A-3*I*B)/a
/d/tan(d*x+c)^(3/2)+1/2*(A+I*B)/d/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))
```

3.139. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

3.139.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.56 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.40

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$= \frac{3(-iA + B) - 2(3A + 2iB) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, -i \tan(c + dx)\right) (-i + \tan(c + dx)) - (A - iB)}{6ad \tan^{\frac{3}{2}}(c + dx)(-i + \tan(c + dx))}$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])),x]`

output `(3*((-I)*A + B) - 2*(3*A + (2*I)*B)*Hypergeometric2F1[-3/2, 1, -1/2, (-I)*Tan[c + d*x]]*(-I + Tan[c + d*x]) - (A - I*B)*Hypergeometric2F1[-3/2, 1, -1/2, I*Tan[c + d*x]]*(-I + Tan[c + d*x]))/(6*a*d*Tan[c + d*x]^(3/2)*(-I + Tan[c + d*x]))`

3.139.3 Rubi [A] (verified)

Time = 0.87 (sec) , antiderivative size = 260, normalized size of antiderivative = 0.88, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.528$, Rules used = {3042, 4079, 27, 3042, 4012, 25, 3042, 4012, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$\downarrow 3042$$

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2}(a + ia \tan(c + dx))} dx$$

$$\downarrow 4079$$

$$\frac{\int \frac{a(7A+3iB)-5a(iA-B) \tan(c+dx)}{2 \tan^{\frac{5}{2}}(c+dx)} dx}{2a^2} + \frac{A + iB}{2d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))}$$

$$\downarrow 27$$

3.139. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

$$\begin{aligned}
& \frac{\int \frac{a(7A+3iB)-5a(iA-B)\tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)} dx}{4a^2} + \frac{A+iB}{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{a(7A+3iB)-5a(iA-B)\tan(c+dx)}{\tan(c+dx)^{5/2}} dx}{4a^2} + \frac{A+iB}{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} \\
& \quad \downarrow 4012 \\
& \frac{\int -\frac{5a(iA-B)+a(7A+3iB)\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)} dx - \frac{2a(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)}}{4a^2} + \frac{A+iB}{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} \\
& \quad \downarrow 25 \\
& \frac{-\int \frac{5a(iA-B)+a(7A+3iB)\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)} dx - \frac{2a(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)}}{4a^2} + \frac{A+iB}{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} \\
& \quad \downarrow 3042 \\
& \frac{-\int \frac{5a(iA-B)+a(7A+3iB)\tan(c+dx)}{\tan(c+dx)^{3/2}} dx - \frac{2a(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)}}{4a^2} + \frac{A+iB}{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} \\
& \quad \downarrow 4012 \\
& \frac{-\int \frac{a(7A+3iB)-5a(iA-B)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{2a(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\tan(c+dx)}}}{4a^2} + \frac{A+iB}{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} \\
& \quad \downarrow 3042 \\
& \frac{-\int \frac{a(7A+3iB)-5a(iA-B)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{2a(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\tan(c+dx)}}}{4a^2} + \frac{A+iB}{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} \\
& \quad \downarrow 4017 \\
& \frac{2 \int \frac{a(7A+3iB-5(iA-B)\tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{2a(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\tan(c+dx)}}}{4a^2} + \frac{A+iB}{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} \\
& \quad \downarrow 27
\end{aligned}$$

3.139. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

$$\begin{aligned}
 & \frac{-\frac{2a \int \frac{7A+3iB-5(iA-B)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} - \frac{2a(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\tan(c+dx)}}}{\frac{4a^2}{A+iB}} + \\
 & \frac{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))}{} \\
 & \quad \downarrow \text{1482} \\
 & \frac{2a \left(\frac{1}{2}((7+5i)A-(5-3i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((7-5i)A+(5+3i)B) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) - \frac{2a(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\tan(c+dx)}}}{\frac{4a^2}{A+iB}} \\
 & \frac{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))}{} \\
 & \quad \downarrow \text{1476} \\
 & \frac{2a \left(\frac{1}{2}((7+5i)A-(5-3i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((7-5i)A+(5+3i)B) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right) - \frac{2a(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\tan(c+dx)}}}{\frac{4a^2}{A+iB}} \\
 & \frac{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))}{} \\
 & \quad \downarrow \text{1082} \\
 & \frac{2a \left(\frac{1}{2}((7+5i)A-(5-3i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((7-5i)A+(5+3i)B) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) \right) - \frac{2a(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\tan(c+dx)}}}{\frac{4a^2}{A+iB}} \\
 & \frac{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))}{} \\
 & \quad \downarrow \text{217} \\
 & \frac{2a \left(\frac{1}{2}((7+5i)A-(5-3i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((7-5i)A+(5+3i)B) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) \right) - \frac{2a(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\tan(c+dx)}}}{\frac{4a^2}{A+iB}} \\
 & \frac{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))}{} \\
 & \quad \downarrow \text{1479}
 \end{aligned}$$

3.139. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

$$\begin{aligned}
& \frac{2a \left(\frac{1}{2}((7+5i)A - (5-3i)B) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}((7-5i)A + (5+3i)B) \right)}{d} \\
& \frac{A+iB}{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} \quad 4a^2 \\
& \quad \downarrow \quad 25 \\
& \frac{2a \left(\frac{1}{2}((7+5i)A - (5-3i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}((7-5i)A + (5+3i)B) \right)}{d} \\
& \frac{A+iB}{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} \quad 4a^2 \\
& \quad \downarrow \quad 27 \\
& \frac{2a \left(\frac{1}{2}((7+5i)A - (5-3i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2}((7-5i)A + (5+3i)B) \right)}{d} \\
& \frac{A+iB}{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} \quad 4a^2 \\
& \quad \downarrow \quad 1103 \\
& \frac{2a \left(\frac{1}{2}((7-5i)A + (5+3i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}((7+5i)A - (5-3i)B) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d} \\
& \frac{A+iB}{2d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} \quad 4a^2
\end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])),x]`

$$3.139. \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$$


```
output ((-2*a*(((7 - 5*I)*A + (5 + 3*I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*
x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 + (((7
+ 5*I)*A - (5 - 3*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c +
d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqr
t[2])))/2)/d - (2*a*(7*A + (3*I)*B))/(3*d*Tan[c + d*x]^(3/2)) + (10*a*(I*
A - B))/(d*Sqrt[Tan[c + d*x]])/(4*a^2) + (A + I*B)/(2*d*Tan[c + d*x]^(3/2
))*(a + I*a*Tan[c + d*x]))
```

3.139.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

```
rule 1103 Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

```
rule 1476 Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.139.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.54

method	result
derivativedivides	$\frac{i \left(\frac{i(A-B)(\sqrt{\tan(dx+c)})}{\tan(dx+c)-i} - \frac{4(2iB+3A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} \right)}{2} - \frac{2A}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(-iA+B)}{\sqrt{\tan(dx+c)}} + \frac{4\left(-\frac{iA}{4} - \frac{B}{4}\right) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}$
default	$\frac{i \left(\frac{i(iA-B)(\sqrt{\tan(dx+c)})}{\tan(dx+c)-i} - \frac{4(2iB+3A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} \right)}{2} - \frac{2A}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(-iA+B)}{\sqrt{\tan(dx+c)}} + \frac{4\left(-\frac{iA}{4} - \frac{B}{4}\right) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}$

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c)),x,method=_RETURNV
ERBOSE)`

output `1/d/a*(-1/2*I*(I*(I*A-B)*tan(d*x+c)^(1/2)/(tan(d*x+c)-I)-4*(3*A+2*I*B)/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2))))-2/3*A/tan(d*x+c)^(3/2)-2*(-I*A+B)/tan(d*x+c)^(1/2)+4*(-1/4*I*A-1/4*B)/(2^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))`

3.139.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 795 vs. 2(221) = 442.

Time = 0.29 (sec) , antiderivative size = 795, normalized size of antiderivative = 2.69

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$= \frac{3 \left(ade^{(6i dx+6i c)} - 2 ade^{(4i dx+4i c)} + ade^{(2i dx+2i c)} \right) \sqrt{\frac{-i A^2 - 2 AB + i B^2}{a^2 d^2}} \log \left(- \frac{2 \left((i ade^{(2i dx+2i c)} + i ad) \sqrt{\frac{-i e^{(2i dx+2i c)}}{e^{(2i dx+2i c)}} + \dots}} \right)}{\dots} \right)}{\dots}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c)),x, algorithm
m="fricas")`

3.139. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

output

```

1/24*(3*(a*d*e^(6*I*d*x + 6*I*c) - 2*a*d*e^(4*I*d*x + 4*I*c) + a*d*e^(2*I*
d*x + 2*I*c))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2))*log(-2*((I*a*d*e^(2
*I*d*x + 2*I*c) + I*a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2
*I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2)) - (A - I*B)*e^(2*I*d*
x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*(a*d*e^(6*I*d*x + 6*I*c) -
2*a*d*e^(4*I*d*x + 4*I*c) + a*d*e^(2*I*d*x + 2*I*c))*sqrt((-I*A^2 - 2*A*B
+ I*B^2)/(a^2*d^2))*log(-2*((-I*a*d*e^(2*I*d*x + 2*I*c) - I*a*d)*sqrt((-I
*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 - 2*A*B
+ I*B^2)/(a^2*d^2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/
(I*A + B)) - 6*(a*d*e^(6*I*d*x + 6*I*c) - 2*a*d*e^(4*I*d*x + 4*I*c) + a*d*
e^(2*I*d*x + 2*I*c))*sqrt((9*I*A^2 - 12*A*B - 4*I*B^2)/(a^2*d^2))*log(-(a
*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*
x + 2*I*c) + 1))*sqrt((9*I*A^2 - 12*A*B - 4*I*B^2)/(a^2*d^2)) + 3*I*A - 2*
B)*e^(-2*I*d*x - 2*I*c)/(a*d)) + 6*(a*d*e^(6*I*d*x + 6*I*c) - 2*a*d*e^(4*I
*d*x + 4*I*c) + a*d*e^(2*I*d*x + 2*I*c))*sqrt((9*I*A^2 - 12*A*B - 4*I*B^2)
/(a^2*d^2))*log(((a*d*e^(2*I*d*x + 2*I*c) + a*d)*sqrt((-I*e^(2*I*d*x + 2*I
*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((9*I*A^2 - 12*A*B - 4*I*B^2)/(a^2
*d^2)) - 3*I*A + 2*B)*e^(-2*I*d*x - 2*I*c)/(a*d)) - 2*((19*A + 27*I*B)*e^(
6*I*d*x + 6*I*c) - (19*A + 3*I*B)*e^(4*I*d*x + 4*I*c) - (35*A + 27*I*B)*e^(
2*I*d*x + 2*I*c) + 3*A + 3*I*B)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(...

```

3.139.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$= - \frac{i \left(\int \frac{A}{\tan^{\frac{7}{2}}(c + dx) - i \tan^{\frac{5}{2}}(c + dx)} dx + \int \frac{B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) - i \tan^{\frac{5}{2}}(c + dx)} dx \right)}{a}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(5/2)/(a+I*a*tan(d*x+c)),x)`

output `-I*(Integral(A/(tan(c + d*x)**(7/2) - I*tan(c + d*x)**(5/2)), x) + Integra
l(B*tan(c + d*x)/(tan(c + d*x)**(7/2) - I*tan(c + d*x)**(5/2)), x))/a`

3.139.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.139.8 Giac [A] (verification not implemented)

Time = 0.90 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.48

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$= - \frac{(i + 1) \sqrt{2}(-3i A + 2 B) \arctan\left(\left(\frac{1}{2}i + \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{2 ad}$$

$$+ \frac{(i - 1) \sqrt{2}(i A + B) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{4 ad}$$

$$- \frac{-i A \sqrt{\tan(dx + c)} + B \sqrt{\tan(dx + c)}}{2 ad(\tan(dx + c) - i)}$$

$$+ \frac{2i(3 A \tan(dx + c) + 3i B \tan(dx + c) + i A)}{3 ad \tan(dx + c)^{\frac{3}{2}}}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c)),x, algorithm="giac")`

output `-(1/2*I + 1/2)*sqrt(2)*(-3*I*A + 2*B)*arctan((1/2*I + 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a*d) + (1/4*I - 1/4)*sqrt(2)*(I*A + B)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a*d) - 1/2*(-I*A*sqrt(tan(d*x + c)) + B*sqrt(tan(d*x + c)))/(a*d*(tan(d*x + c) - I)) + 2/3*I*(3*A*tan(d*x + c) + 3*I*B*tan(d*x + c) + I*A)/(a*d*tan(d*x + c)^(3/2))`

3.139.9 Mupad [B] (verification not implemented)

Time = 11.37 (sec) , antiderivative size = 303, normalized size of antiderivative = 1.02

$$\begin{aligned}
& \int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx \\
&= \operatorname{atan} \left(\frac{2 a d \sqrt{\tan(c + dx)} \sqrt{\frac{A^2 9i}{4 a^2 d^2}}}{3 A} \right) \sqrt{\frac{A^2 9i}{4 a^2 d^2}} 2i \\
&\quad - \operatorname{atan} \left(\frac{4 a d \sqrt{\tan(c + dx)} \sqrt{-\frac{A^2 1i}{16 a^2 d^2}}}{A} \right) \sqrt{-\frac{A^2 1i}{16 a^2 d^2}} 2i \\
&\quad + 2 \operatorname{atanh} \left(\frac{a d \sqrt{\tan(c + dx)} \sqrt{-\frac{B^2 1i}{a^2 d^2}}}{B} \right) \sqrt{-\frac{B^2 1i}{a^2 d^2}} \\
&\quad + 2 \operatorname{atanh} \left(\frac{4 a d \sqrt{\tan(c + dx)} \sqrt{\frac{B^2 1i}{16 a^2 d^2}}}{B} \right) \sqrt{\frac{B^2 1i}{16 a^2 d^2}} \\
&\quad - \frac{\frac{2 A}{3 a d} + \frac{5 A \tan(c + dx)^2}{2 a d} - \frac{A \tan(c + dx) 4i}{3 a d}}{\tan(c + dx)^{3/2} + \tan(c + dx)^{5/2} 1i} - \frac{\frac{2 B}{a d} + \frac{B \tan(c + dx) 5i}{2 a d}}{\sqrt{\tan(c + dx)} + \tan(c + dx)^{3/2} 1i}
\end{aligned}$$

```
input int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + a*tan(c + d*x)*1i)),x)
```

```
output atan((2*a*d*tan(c + d*x)^(1/2)*((A^2*9i)/(4*a^2*d^2))^(1/2))/(3*A))*((A^2*
9i)/(4*a^2*d^2))^(1/2)*2i - atan((4*a*d*tan(c + d*x)^(1/2)*(-(A^2*1i)/(16*
a^2*d^2))^(1/2))/A)*(-(A^2*1i)/(16*a^2*d^2))^(1/2)*2i + 2*atanh((a*d*tan(c
+ d*x)^(1/2)*(-(B^2*1i)/(a^2*d^2))^(1/2))/B)*(-(B^2*1i)/(a^2*d^2))^(1/2)
+ 2*atanh((4*a*d*tan(c + d*x)^(1/2)*((B^2*1i)/(16*a^2*d^2))^(1/2))/B)*((B^
2*1i)/(16*a^2*d^2))^(1/2) - ((2*A)/(3*a*d) - (A*tan(c + d*x)*4i)/(3*a*d) +
(5*A*tan(c + d*x)^2)/(2*a*d))/(tan(c + d*x)^(3/2) + tan(c + d*x)^(5/2)*1i
) - ((2*B)/(a*d) + (B*tan(c + d*x)*5i)/(2*a*d))/(tan(c + d*x)^(1/2) + tan(
c + d*x)^(3/2)*1i)
```

3.140
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.140.1 Optimal result 1448
 3.140.2 Mathematica [A] (verified) 1449
 3.140.3 Rubi [A] (verified) 1449
 3.140.4 Maple [A] (verified) 1454
 3.140.5 Fricas [B] (verification not implemented) 1455
 3.140.6 Sympy [F] 1456
 3.140.7 Maxima [F(-2)] 1456
 3.140.8 Giac [A] (verification not implemented) 1456
 3.140.9 Mupad [B] (verification not implemented) 1457

3.140.1 Optimal result

Integrand size = 36, antiderivative size = 316

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$= \frac{((9+5i)A - (25-21i)B) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{16\sqrt{2}a^2d}$$

$$- \frac{((9+5i)A - (25-21i)B) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{16\sqrt{2}a^2d}$$

$$- \frac{\left(\frac{1}{32} - \frac{i}{32}\right) ((7+2i)A + (2+23i)B) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{\sqrt{2}a^2d}$$

$$+ \frac{\left(\frac{1}{32} - \frac{i}{32}\right) ((7+2i)A + (2+23i)B) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{\sqrt{2}a^2d}$$

$$+ \frac{5(iA - 5B)\sqrt{\tan(c+dx)}}{8a^2d} + \frac{(3A + 7iB) \tan^{\frac{3}{2}}(c+dx)}{8a^2d(1+i \tan(c+dx))} + \frac{(iA - B) \tan^{\frac{5}{2}}(c+dx)}{4d(a+ia \tan(c+dx))^2}$$

output

```
-1/32*((9+5*I)*A+(-25+21*I)*B)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/a^2/d*2
^(1/2)-1/32*((9+5*I)*A+(-25+21*I)*B)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/a^
2/d*2^(1/2)+(-1/64+1/64*I)*((7+2*I)*A+(2+23*I)*B)*ln(1-2^(1/2)*tan(d*x+c)^(
1/2)+tan(d*x+c))/a^2/d*2^(1/2)+(1/64-1/64*I)*((7+2*I)*A+(2+23*I)*B)*ln(1+
2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a^2/d*2^(1/2)+5/8*(I*A-5*B)*tan(d*x+c
)^(1/2)/a^2/d+1/8*(3*A+7*I*B)*tan(d*x+c)^(3/2)/a^2/d/(1+I*tan(d*x+c))+1/4*
(I*A-B)*tan(d*x+c)^(5/2)/d/(a+I*a*tan(d*x+c))^2
```

3.140.
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.140.2 Mathematica [A] (verified)

Time = 3.47 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.60

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$= \frac{2\sqrt[4]{-1}(iA+B) \arctan\left((-1)^{3/4} \sqrt{\tan(c+dx)}\right) \sec^2(c+dx)(\cos(2(c+dx)) + i \sin(2(c+dx))) + \sqrt[4]{-1}(-$$

input `Integrate[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])
^2,x]`

output `(2*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]
^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)]) + (-1)^(1/4)*((-7*I)*A + 23*B)*
ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] +
I*Sin[2*(c + d*x)]) + Sqrt[Tan[c + d*x]]*((-5*I)*A + 25*B + (7*A + (43*I)*
B)*Tan[c + d*x] - 16*B*Tan[c + d*x]^2))/(8*a^2*d*(-I + Tan[c + d*x])^2)`

3.140.3 Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 289, normalized size of antiderivative = 0.91,
number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.528$, Rules
used = {3042, 4078, 27, 3042, 4078, 25, 3042, 4011, 3042, 4017, 27, 1482, 1476, 1082, 217,
1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the
transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$\downarrow 3042$$

$$\int \frac{\tan(c+dx)^{5/2}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$\downarrow 4078$$

$$\frac{(-B+iA) \tan^{\frac{5}{2}}(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{\int \frac{\tan^{\frac{3}{2}}(c+dx)(5a(iA-B)+a(A+9iB) \tan(c+dx))}{2(i \tan(c+dx)a+a)} dx}{4a^2}$$

$$\downarrow 27$$

3.140. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{\tan^{\frac{3}{2}}(c+dx)(5a(iA-B)+a(A+9iB) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{8a^2} \\
& \quad \downarrow 3042 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{\tan(c+dx)^{3/2}(5a(iA-B)+a(A+9iB) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{8a^2} \\
& \quad \downarrow 4078 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int -\sqrt{\tan(c+dx)}(3a^2(3A+7iB)-5a^2(iA-5B) \tan(c+dx)) dx}{2a^2} - \frac{(3A+7iB) \tan^{\frac{3}{2}}(c+dx)}{d(1+i \tan(c+dx))} \\
& \quad \downarrow 25 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int \sqrt{\tan(c+dx)}(3a^2(3A+7iB)-5a^2(iA-5B) \tan(c+dx)) dx}{2a^2} - \frac{(3A+7iB) \tan^{\frac{3}{2}}(c+dx)}{d(1+i \tan(c+dx))} \\
& \quad \downarrow 3042 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int \sqrt{\tan(c+dx)}(3a^2(3A+7iB)-5a^2(iA-5B) \tan(c+dx)) dx}{2a^2} - \frac{(3A+7iB) \tan^{\frac{3}{2}}(c+dx)}{d(1+i \tan(c+dx))} \\
& \quad \downarrow 4011 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{5(iA-5B)a^2+3(3A+7iB) \tan(c+dx)a^2}{\sqrt{\tan(c+dx)}} dx - \frac{10a^2(-5B+iA)\sqrt{\tan(c+dx)}}{d}}{2a^2} - \frac{(3A+7iB) \tan^{\frac{3}{2}}(c+dx)}{d(1+i \tan(c+dx))} \\
& \quad \downarrow 3042 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{5(iA-5B)a^2+3(3A+7iB) \tan(c+dx)a^2}{\sqrt{\tan(c+dx)}} dx - \frac{10a^2(-5B+iA)\sqrt{\tan(c+dx)}}{d}}{2a^2} - \frac{(3A+7iB) \tan^{\frac{3}{2}}(c+dx)}{d(1+i \tan(c+dx))} \\
& \quad \downarrow 4017 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{2 \int \frac{a^2(5(iA-5B)+3(3A+7iB) \tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{10a^2(-5B+iA)\sqrt{\tan(c+dx)}}{d}}{2a^2} - \frac{(3A+7iB) \tan^{\frac{3}{2}}(c+dx)}{d(1+i \tan(c+dx))} \\
& \quad \downarrow 27
\end{aligned}$$

3.140. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\frac{\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{2a^2 \int \frac{5(iA - 5B) + 3(3A + 7iB) \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{10a^2(-5B + iA)\sqrt{\tan(c + dx)}}{d}}{2a^2}}{8a^2} - \frac{(3A + 7iB) \tan^{\frac{3}{2}}(c + dx)}{d(1 + i \tan(c + dx))}$$

↓ 1482

$$\frac{\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{2a^2 \left(\frac{1}{2}((9 + 5i)A - (25 - 21i)B) \int \frac{\tan(c + dx) + 1}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \left(\frac{1}{2} - \frac{i}{2}\right)((7 + 2i)A + (2 + 23i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} \right) - \frac{10a^2(-5B + iA)\sqrt{\tan(c + dx)}}{d}}{2a^2}}{8a^2} - \frac{(3A + 7iB) \tan^{\frac{3}{2}}(c + dx)}{d(1 + i \tan(c + dx))}$$

↓ 1476

$$\frac{\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{2a^2 \left(\frac{1}{2}((9 + 5i)A - (25 - 21i)B) \left(\frac{1}{2} \int \frac{1}{\tan(c + dx) - \sqrt{2}\sqrt{\tan(c + dx)} + 1} d\sqrt{\tan(c + dx)} + \frac{1}{2} \int \frac{1}{\tan(c + dx) + \sqrt{2}\sqrt{\tan(c + dx)} + 1} d\sqrt{\tan(c + dx)} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((7 + 2i)A + (2 + 23i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} \right)}{2a^2}}{8a^2} - \frac{(3A + 7iB) \tan^{\frac{3}{2}}(c + dx)}{d(1 + i \tan(c + dx))}$$

↓ 1082

$$\frac{\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{2a^2 \left(\frac{1}{2}((9 + 5i)A - (25 - 21i)B) \left(\frac{\int \frac{1}{-\tan(c + dx) - 1} d(1 - \sqrt{2}\sqrt{\tan(c + dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c + dx) - 1} d(\sqrt{2}\sqrt{\tan(c + dx)} + 1)}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((7 + 2i)A + (2 + 23i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} \right)}{2a^2}}{8a^2} - \frac{(3A + 7iB) \tan^{\frac{3}{2}}(c + dx)}{d(1 + i \tan(c + dx))}$$

↓ 217

$$\frac{\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{2a^2 \left(\frac{1}{2}((9 + 5i)A - (25 - 21i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c + dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c + dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((7 + 2i)A + (2 + 23i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} \right)}{2a^2}}{8a^2} - \frac{(3A + 7iB) \tan^{\frac{3}{2}}(c + dx)}{d(1 + i \tan(c + dx))}$$

↓ 1479

$$\frac{\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{2a^2 \left(\frac{1}{2}((9 + 5i)A - (25 - 21i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c + dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c + dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((7 + 2i)A + (2 + 23i)B) \left(-\frac{\int \frac{\sqrt{2} - 2\sqrt{\tan(c + dx)}}{\tan(c + dx) - \sqrt{2}\sqrt{\tan(c + dx)} + 1} d\sqrt{\tan(c + dx)}}{2\sqrt{2}} \right) \right)}{2a^2}}{8a^2} - \frac{(3A + 7iB) \tan^{\frac{3}{2}}(c + dx)}{d(1 + i \tan(c + dx))}$$

3.140. $\int \frac{\tan^{\frac{5}{2}}(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx$

$$\begin{aligned} & \downarrow 25 \\ & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \\ & \frac{2a^2 \left(\frac{1}{2}((9+5i)A - (25-21i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((7+2i)A + (2+23i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d \cdot 2a^2} \\ & \hline & 8a^2 \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \\ & \frac{2a^2 \left(\frac{1}{2}((9+5i)A - (25-21i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((7+2i)A + (2+23i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d \cdot 2a^2} \\ & \hline & 8a^2 \end{aligned}$$

$$\begin{aligned} & \downarrow 1103 \\ & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \\ & \frac{2a^2 \left(\frac{1}{2}((9+5i)A - (25-21i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((7+2i)A + (2+23i)B) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d \cdot 2a^2} \\ & \hline & 8a^2 \end{aligned}$$

input `Int[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `((I*A - B)*Tan[c + d*x]^(5/2))/(4*d*(a + I*a*Tan[c + d*x])^2) - (((2*a^2*((9 + 5*I)*A - (25 - 21*I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 - (1/2 - I/2)*((7 + 2*I)*A + (2 + 23*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/d - (10*a^2*(I*A - 5*B)*Sqrt[Tan[c + d*x]]/d)/(2*a^2) - ((3*A + (7*I)*B)*Tan[c + d*x]^(3/2))/(d*(1 + I*Tan[c + d*x]))/(8*a^2)`

3.140. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

3.140.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

$$3.140. \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4078 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.140.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.50

method	result
derivativedivides	$\frac{-2B(\sqrt{\tan(dx+c)}) + \frac{i \left(\frac{(-7iA + 11B)}{2} \left(\tan^{\frac{3}{2}}(dx+c) \right) + \left(-\frac{5A}{2} - \frac{9iB}{2} \right) (\sqrt{\tan(dx+c)}) \right)}{(\tan(dx+c)-i)^2} + \frac{(7iA-23B) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}}}{4} da^2$
default	$\frac{-2B(\sqrt{\tan(dx+c)}) + \frac{i \left(\frac{(-7iA + 11B)}{2} \left(\tan^{\frac{3}{2}}(dx+c) \right) + \left(-\frac{5A}{2} - \frac{9iB}{2} \right) (\sqrt{\tan(dx+c)}) \right)}{(\tan(dx+c)-i)^2} + \frac{(7iA-23B) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}}}{4} da^2$

input `int(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

$$3.140. \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

output $\frac{1}{d/a^2}(-2B\tan(dx+c)^{1/2}+1/4I(((-7/2IA+11/2B)\tan(dx+c)^{3/2}+(-5/2A-9/2IB)\tan(dx+c)^{1/2}))/(\tan(dx+c)-I)^2+(7IA-23B)/(2^{1/2}-I2^{1/2})*\arctan(2\tan(dx+c)^{1/2}/(2^{1/2}-I2^{1/2}))) + 1/2I*(IA+B)/(2^{1/2}+I2^{1/2})*\arctan(2\tan(dx+c)^{1/2}/(2^{1/2}+I2^{1/2})))$

3.140.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 664 vs. $2(235) = 470$.

Time = 0.27 (sec) , antiderivative size = 664, normalized size of antiderivative = 2.10

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= \left(2a^2d\sqrt{\frac{iA^2+2AB-iB^2}{a^4d^2}}e^{(4idx+4ic)} \log \left(\frac{2\left((a^2de^{(2idx+2ic)}+a^2d)\sqrt{\frac{-ie^{(2idx+2ic)}+i}{e^{(2idx+2ic)}+1}}\sqrt{\frac{iA^2+2AB-iB^2}{a^4d^2}}+(A-iB)e^{(2idx+2ic)} \right)}{iA+B} \right) \right) e^{(-$$

input `integrate(tan(dx+c)^(5/2)*(A+B*tan(dx+c))/(a+I*a*tan(dx+c))^2,x, algorithm="fracas")`

output $\frac{1}{32}*(2*a^2*d*\sqrt{((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))}*e^{(4*I*d*x + 4*I*c)}*\log(2*((a^2*d*e^{(2*I*d*x + 2*I*c)} + a^2*d)*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))*\sqrt{((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))} + (A - I*B)*e^{(2*I*d*x + 2*I*c)})*e^{(-2*I*d*x - 2*I*c)/(I*A + B)} - 2*a^2*d*\sqrt{((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))}*e^{(4*I*d*x + 4*I*c)}*\log(-2*((a^2*d*e^{(2*I*d*x + 2*I*c)} + a^2*d)*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))*\sqrt{((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))} - (A - I*B)*e^{(2*I*d*x + 2*I*c)})*e^{(-2*I*d*x - 2*I*c)/(I*A + B)} + a^2*d*\sqrt{((-49*I*A^2 + 322*A*B + 529*I*B^2)/(a^4*d^2))}*e^{(4*I*d*x + 4*I*c)}*\log(1/8*((a^2*d*e^{(2*I*d*x + 2*I*c)} + a^2*d)*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))*\sqrt{((-49*I*A^2 + 322*A*B + 529*I*B^2)/(a^4*d^2))} + 7*A + 23*I*B)*e^{(-2*I*d*x - 2*I*c)/(a^2*d)} - a^2*d*\sqrt{((-49*I*A^2 + 322*A*B + 529*I*B^2)/(a^4*d^2))}*e^{(4*I*d*x + 4*I*c)}*\log(-1/8*((a^2*d*e^{(2*I*d*x + 2*I*c)} + a^2*d)*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1))*\sqrt{((-49*I*A^2 + 322*A*B + 529*I*B^2)/(a^4*d^2))} - 7*A - 23*I*B)*e^{(-2*I*d*x - 2*I*c)/(a^2*d)} - 2*(6*(-I*A + 7*B)*e^{(4*I*d*x + 4*I*c)} - (5*I*A - 9*B)*e^{(2*I*d*x + 2*I*c)} + I*A - B)*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1)))*e^{(-4*I*d*x - 4*I*c)/(a^2*d)}$

$$3.140. \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

3.140.6 Sympy [F]

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= -\frac{\int \frac{A \tan^{\frac{5}{2}}(c+dx)}{\tan^2(c+dx)-2i\tan(c+dx)-1} dx + \int \frac{B \tan^{\frac{7}{2}}(c+dx)}{\tan^2(c+dx)-2i\tan(c+dx)-1} dx}{a^2}$$

input `integrate(tan(d*x+c)**(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**2,x)`

output `-(Integral(A*tan(c + d*x)**(5/2)/(tan(c + d*x)**2 - 2*I*tan(c + d*x) - 1), x) + Integral(B*tan(c + d*x)**(7/2)/(tan(c + d*x)**2 - 2*I*tan(c + d*x) - 1), x))/a**2`

3.140.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.140.8 Giac [A] (verification not implemented)

Time = 0.77 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.45

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= -\frac{(i+1)\sqrt{2}(7A+23iB)\arctan\left(\left(\frac{1}{2}i+\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{16a^2d}$$

$$+ \frac{(i-1)\sqrt{2}(A-iB)\arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{8a^2d} - \frac{2B\sqrt{\tan(dx+c)}}{a^2d}$$

$$+ \frac{7A\tan(dx+c)^{\frac{3}{2}} + 11iB\tan(dx+c)^{\frac{3}{2}} - 5iA\sqrt{\tan(dx+c)} + 9B\sqrt{\tan(dx+c)}}{8a^2d(\tan(dx+c)-i)^2}$$

3.140. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")`

output
$$-(1/16*I + 1/16)*\sqrt{2}*(7*A + 23*I*B)*\arctan((1/2*I + 1/2)*\sqrt{2}*\sqrt{\tan(d*x + c)})/(a^2*d) + (1/8*I - 1/8)*\sqrt{2}*(A - I*B)*\arctan(-(1/2*I - 1/2)*\sqrt{2}*\sqrt{\tan(d*x + c)})/(a^2*d) - 2*B*\sqrt{\tan(d*x + c)}/(a^2*d) + 1/8*(7*A*\tan(d*x + c)^{(3/2)} + 11*I*B*\tan(d*x + c)^{(3/2)} - 5*I*A*\sqrt{\tan(d*x + c)} + 9*B*\sqrt{\tan(d*x + c)})/(a^2*d*(\tan(d*x + c) - I)^2)$$

3.140.9 Mupad [B] (verification not implemented)

Time = 12.04 (sec) , antiderivative size = 334, normalized size of antiderivative = 1.06

$$\begin{aligned} & \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx \\ &= \frac{\frac{5A\sqrt{\tan(c+dx)}}{8a^2d} + \frac{A\tan(c+dx)^{3/2}7i}{8a^2d}}{\tan(c+dx)^2 li + 2\tan(c+dx) - i} + \frac{-\frac{11B\tan(c+dx)^{3/2}}{8a^2d} + \frac{B\sqrt{\tan(c+dx)}9i}{8a^2d}}{\tan(c+dx)^2 li + 2\tan(c+dx) - i} \\ &+ 2\operatorname{atanh}\left(\frac{8a^2d\sqrt{\tan(c+dx)}\sqrt{\frac{A^2 li}{64a^4d^2}}}{A}\right)\sqrt{\frac{A^2 li}{64a^4d^2}} \\ &+ 2\operatorname{atanh}\left(\frac{16a^2d\sqrt{\tan(c+dx)}\sqrt{-\frac{A^2 49i}{256a^4d^2}}}{7A}\right)\sqrt{-\frac{A^2 49i}{256a^4d^2}} \\ &- \frac{2B\sqrt{\tan(c+dx)}}{a^2d} + \operatorname{atan}\left(\frac{8a^2d\sqrt{\tan(c+dx)}\sqrt{-\frac{B^2 li}{64a^4d^2}}}{B}\right)\sqrt{-\frac{B^2 li}{64a^4d^2}} 2i \\ &- \operatorname{atan}\left(\frac{16a^2d\sqrt{\tan(c+dx)}\sqrt{\frac{B^2 529i}{256a^4d^2}}}{23B}\right)\sqrt{\frac{B^2 529i}{256a^4d^2}} 2i \end{aligned}$$

input `int((tan(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^2,x)`

output $((5A \tan(c + dx)^{1/2}) / (8a^2d) + (A \tan(c + dx)^{3/2} * 7i) / (8a^2d)) / (2 \tan(c + dx) + \tan(c + dx)^{2*1i - 1i}) + ((B \tan(c + dx)^{1/2} * 9i) / (8a^2d) - (11B \tan(c + dx)^{3/2}) / (8a^2d)) / (2 \tan(c + dx) + \tan(c + dx)^{2*1i - 1i}) + 2 \operatorname{atanh}((8a^2d \tan(c + dx)^{1/2} * ((A^2 * 1i) / (64a^4d^2))^{1/2}) / A) * ((A^2 * 1i) / (64a^4d^2))^{1/2} + 2 \operatorname{atanh}((16a^2d \tan(c + dx)^{1/2} * (-A^2 * 49i) / (256a^4d^2))^{1/2}) / (7A) * (-A^2 * 49i) / (256a^4d^2))^{1/2} + \operatorname{atan}((8a^2d \tan(c + dx)^{1/2} * (-B^2 * 1i) / (64a^4d^2))^{1/2}) / B * (-B^2 * 1i) / (64a^4d^2))^{1/2} * 2i - \operatorname{atan}((16a^2d \tan(c + dx)^{1/2} * (B^2 * 529i) / (256a^4d^2))^{1/2}) / (23B) * (B^2 * 529i) / (256a^4d^2))^{1/2} * 2i - (2B \tan(c + dx)^{1/2}) / (a^2d)$

3.140. $\int \frac{\tan^5(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

3.141
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.141.1 Optimal result 1459
 3.141.2 Mathematica [A] (verified) 1460
 3.141.3 Rubi [A] (verified) 1460
 3.141.4 Maple [A] (verified) 1465
 3.141.5 Fricas [B] (verification not implemented) 1466
 3.141.6 Sympy [F] 1467
 3.141.7 Maxima [F(-2)] 1467
 3.141.8 Giac [A] (verification not implemented) 1467
 3.141.9 Mupad [B] (verification not implemented) 1468

3.141.1 Optimal result

Integrand size = 36, antiderivative size = 277

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$= \frac{((1+3i)A+(9+5i)B) \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{16\sqrt{2}a^2d}$$

$$- \frac{((1+3i)A+(9+5i)B) \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{16\sqrt{2}a^2d}$$

$$+ \frac{((1-3i)A-(9-5i)B) \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{32\sqrt{2}a^2d}$$

$$- \frac{((1-3i)A-(9-5i)B) \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{32\sqrt{2}a^2d}$$

$$+ \frac{(A+5iB)\sqrt{\tan(c+dx)}}{8a^2d(1+i \tan(c+dx))} + \frac{(iA-B) \tan^{\frac{3}{2}}(c+dx)}{4d(a+ia \tan(c+dx))^2}$$

output

```
-1/32*((1+3*I)*A+(9+5*I)*B)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/a^2/d*2^(1/2)-1/32*((1+3*I)*A+(9+5*I)*B)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/a^2/d*2^(1/2)+1/64*((1-3*I)*A+(-9+5*I)*B)*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a^2/d*2^(1/2)-1/64*((1-3*I)*A+(-9+5*I)*B)*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a^2/d*2^(1/2)+1/8*(A+5*I*B)*tan(d*x+c)^(1/2)/a^2/d/(1+I*tan(d*x+c))+1/4*(I*A-B)*tan(d*x+c)^(3/2)/d/(a+I*a*tan(d*x+c))^2
```

3.141.
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.141.2 Mathematica [A] (verified)

Time = 3.23 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.64

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$= \frac{-2\sqrt[4]{-1}(A-iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right) \sec^2(c+dx)(\cos(2(c+dx)) + i \sin(2(c+dx))) + \sqrt[4]{-1}}$$

input `Integrate[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `(-2*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)]) + (-1)^(1/4)*(A - (7*I)*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)]) + Sqrt[Tan[c + d*x]]*(-A - (5*I)*B + ((-3*I)*A + 7*B)*Tan[c + d*x]))/(8*a^2*d*(-I + Tan[c + d*x])^2)`

3.141.3 Rubi [A] (verified)Time = 0.86 (sec) , antiderivative size = 244, normalized size of antiderivative = 0.88, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.472$, Rules used = {3042, 4078, 27, 3042, 4078, 25, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)^{3/2}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$\downarrow \text{4078}$$

$$\frac{(-B+iA) \tan^{\frac{3}{2}}(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{\int \frac{\sqrt{\tan(c+dx)}(3a(iA-B)-a(A-7iB) \tan(c+dx))}{2(i \tan(c+dx)a+a)} dx}{4a^2}$$

$$\downarrow \text{27}$$

3.141. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{\sqrt{\tan(c+dx)}(3a(iA-B)-a(A-7iB) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{8a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{\sqrt{\tan(c+dx)}(3a(iA-B)-a(A-7iB) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{8a^2} \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int -\frac{(A+5iB)a^2+3(iA+3B) \tan(c+dx)a^2}{\sqrt{\tan(c+dx)}} dx}{8a^2} - \frac{(A+5iB)\sqrt{\tan(c+dx)}}{d(1+i \tan(c+dx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{(A+5iB)a^2+3(iA+3B) \tan(c+dx)a^2}{\sqrt{\tan(c+dx)}} dx}{8a^2} - \frac{(A+5iB)\sqrt{\tan(c+dx)}}{d(1+i \tan(c+dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{(A+5iB)a^2+3(iA+3B) \tan(c+dx)a^2}{\sqrt{\tan(c+dx)}} dx}{8a^2} - \frac{(A+5iB)\sqrt{\tan(c+dx)}}{d(1+i \tan(c+dx))} \\
 & \quad \downarrow \text{4017} \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{a^2(A+5iB+3(iA+3B) \tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{8a^2} - \frac{(A+5iB)\sqrt{\tan(c+dx)}}{d(1+i \tan(c+dx))} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{A+5iB+3(iA+3B) \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{8a^2} - \frac{(A+5iB)\sqrt{\tan(c+dx)}}{d(1+i \tan(c+dx))} \\
 & \quad \downarrow \text{1482} \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \\
 & \frac{\frac{1}{2}((1-3i)A-(9-5i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((1+3i)A+(9+5i)B) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} - \frac{(A+5iB)\sqrt{\tan(c+dx)}}{d(1+i \tan(c+dx))} \\
 & \quad \downarrow \text{1476} \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1-3i)A-(9-5i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((1+3i)A+(9+5i)B) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} - \frac{(A+5iB)\sqrt{\tan(c+dx)}}{d(1+i \tan(c+dx))}
 \end{aligned}$$

3.141. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1-3i)A - (9-5i)B) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((1+3i)A + (9+5i)B) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right)}{8a^2}$$

↓ 1082

$$\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1-3i)A - (9-5i)B) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((1+3i)A + (9+5i)B) \left(\frac{\int \frac{1}{\tan(c+dx)-1} d(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right)}{8a^2}$$

↓ 217

$$\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1-3i)A - (9-5i)B) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((1+3i)A + (9+5i)B) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1 - \sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right)}{8a^2} - \frac{(A+5i)B}{d(1+i)}$$

↓ 1479

$$\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1-3i)A - (9-5i)B) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}((1+3i)A + (9+5i)B) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1 - \sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right)}{8a^2}$$

↓ 25

$$\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1-3i)A - (9-5i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}((1+3i)A + (9+5i)B) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1 - \sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right)}{8a^2}$$

↓ 27

3.141. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\frac{\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1-3i)A - (9-5i)B) \left(\int \frac{\sqrt{2-2\sqrt{\tan(c+dx)}}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2}((1+3i)A + (9+5i)B) \left(\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right) \right) + \frac{1}{2}((1-3i)A - (9-5i)B) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right)}{d}}{8a^2}$$

↓ 1103

$$\frac{\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1+3i)A + (9+5i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}) - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}} \right) + \frac{1}{2}((1-3i)A - (9-5i)B) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right)}{d}}{8a^2}$$

input `Int[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `-1/8*(((1 + 3*I)*A + (9 + 5*I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 + (((1 - 3*I)*A - (9 - 5*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d - ((A + (5*I)*B)*Sqrt[Tan[c + d*x]]/(d*(1 + I*Tan[c + d*x])))/a^2 + ((I*A - B)*Tan[c + d*x]^(3/2))/(4*d*(a + I*a*Tan[c + d*x])^2)`

3.141.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

3.141. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

```
rule 4078 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.141.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.53

method	result
derivativedivides	$\frac{i \left(\frac{\left(-\frac{7iB}{2} - \frac{3A}{2} \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) + \left(-\frac{5B}{2} + \frac{iA}{2} \right) \sqrt{\tan(dx+c)} - (-7iB+A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}} \right)}{(\tan(dx+c)-i)^2} \right)}{4} - \frac{i(-iB+A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}} \right)}{2(\sqrt{2+i\sqrt{2}})}$
default	$\frac{i \left(\frac{\left(-\frac{7iB}{2} - \frac{3A}{2} \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) + \left(-\frac{5B}{2} + \frac{iA}{2} \right) \sqrt{\tan(dx+c)} - (-7iB+A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}} \right)}{4} \right)}{da^2} - \frac{i(-iB+A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}} \right)}{2(\sqrt{2+i\sqrt{2}})}$

```
input int(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x,method=_RETUR
NVERBOSE)
```

```
output 1/d/a^2*(1/4*I*((( -7/2*I*B-3/2*A)*tan(d*x+c)^(3/2)+(-5/2*B+1/2*I*A)*tan(d*
x+c)^(1/2))/(tan(d*x+c)-I)^2-(-7*I*B+A)/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d
*x+c)^(1/2)/(2^(1/2)-I*2^(1/2))))-1/2*I*(A-I*B)/(2^(1/2)+I*2^(1/2))*arctan
(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2))))
```

$$3.141. \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.141.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 664 vs. $2(210) = 420$.

Time = 0.26 (sec) , antiderivative size = 664, normalized size of antiderivative = 2.40

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$= \left(2 a^2 d \sqrt{\frac{-i A^2 - 2 A B + i B^2}{a^4 d^2}} e^{(4i dx + 4i c)} \log \left(-\frac{2 \left((i a^2 d e^{(2i dx + 2i c)} + i a^2 d) \sqrt{\frac{-i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} + 1}} \sqrt{\frac{-i A^2 - 2 A B + i B^2}{a^4 d^2}} - (A - i B) e^{(2i dx + 2i c)} \right)}{i A + B} \right) \right)$$

```
input integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorith
thm="fricas")
```

```
output 1/32*(2*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)
*log(-2*((I*a^2*d*e^(2*I*d*x + 2*I*c) + I*a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I
*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2
)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 2*a^
2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(-2*((
-I*a^2*d*e^(2*I*d*x + 2*I*c) - I*a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/
(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2)) - (A -
I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) + a^2*d*sqrt((I
*A^2 + 14*A*B - 49*I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(1/8*((a^2*d*e
^(2*I*d*x + 2*I*c) + a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x
+ 2*I*c) + 1))*sqrt((I*A^2 + 14*A*B - 49*I*B^2)/(a^4*d^2)) + I*A + 7*B)*e^
(-2*I*d*x - 2*I*c)/(a^2*d)) - a^2*d*sqrt((I*A^2 + 14*A*B - 49*I*B^2)/(a^4*
d^2))*e^(4*I*d*x + 4*I*c)*log(-1/8*((a^2*d*e^(2*I*d*x + 2*I*c) + a^2*d)*sq
rt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*A^2 + 1
4*A*B - 49*I*B^2)/(a^4*d^2)) - I*A - 7*B)*e^(-2*I*d*x - 2*I*c)/(a^2*d)) +
2*(2*(A + 3*I*B)*e^(4*I*d*x + 4*I*c) + (A + 5*I*B)*e^(2*I*d*x + 2*I*c) - A
- I*B)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*e^(-
4*I*d*x - 4*I*c)/(a^2*d)
```

$$3.141. \quad \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.141.6 Sympy [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= -\frac{\int \frac{A\tan^{\frac{3}{2}}(c+dx)}{\tan^2(c+dx)-2i\tan(c+dx)-1} dx + \int \frac{B\tan^{\frac{5}{2}}(c+dx)}{\tan^2(c+dx)-2i\tan(c+dx)-1} dx}{a^2}$$

input `integrate(tan(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**2,x)`

output `-(Integral(A*tan(c + d*x)**(3/2)/(tan(c + d*x)**2 - 2*I*tan(c + d*x) - 1), x) + Integral(B*tan(c + d*x)**(5/2)/(tan(c + d*x)**2 - 2*I*tan(c + d*x) - 1), x))/a**2`

3.141.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.141.8 Giac [A] (verification not implemented)

Time = 0.65 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.45

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= -\frac{(i+1)\sqrt{2}(iA+7B)\arctan\left(\left(\frac{1}{2}i+\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{16a^2d}$$

$$+ \frac{(i-1)\sqrt{2}(iA+B)\arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{8a^2d}$$

$$- \frac{3iA\tan(dx+c)^{\frac{3}{2}}-7B\tan(dx+c)^{\frac{3}{2}}+A\sqrt{\tan(dx+c)}+5iB\sqrt{\tan(dx+c)}}{8a^2d(\tan(dx+c)-i)^2}$$

3.141. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")`

output
$$-(1/16*I + 1/16)*\sqrt{2}*(I*A + 7*B)*\arctan((1/2*I + 1/2)*\sqrt{2}*\sqrt{\tan(d*x + c)})/(a^2*d) + (1/8*I - 1/8)*\sqrt{2}*(I*A + B)*\arctan(-(1/2*I - 1/2)*\sqrt{2}*\sqrt{\tan(d*x + c)})/(a^2*d) - 1/8*(3*I*A*\tan(d*x + c)^{(3/2)} - 7*B*\tan(d*x + c)^{(3/2)} + A*\sqrt{\tan(d*x + c)} + 5*I*B*\sqrt{\tan(d*x + c)})/(a^2*d*(\tan(d*x + c) - I)^2)$$

3.141.9 Mupad [B] (verification not implemented)

Time = 11.37 (sec) , antiderivative size = 318, normalized size of antiderivative = 1.15

$$\begin{aligned} & \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx \\ &= -\frac{-\frac{3A\tan(c+dx)^{3/2}}{8a^2d} + \frac{A\sqrt{\tan(c+dx)}\operatorname{li}}{8a^2d}}{\tan(c+dx)^2\operatorname{li} + 2\tan(c+dx) - i} + \frac{\frac{5B\sqrt{\tan(c+dx)}}{8a^2d} + \frac{B\tan(c+dx)^{3/2}\operatorname{li}}{8a^2d}}{\tan(c+dx)^2\operatorname{li} + 2\tan(c+dx) - i} \\ & - \operatorname{atan}\left(\frac{8a^2d\sqrt{\tan(c+dx)}\sqrt{-\frac{A^2\operatorname{li}}{64a^4d^2}}}{A}\right)\sqrt{-\frac{A^2\operatorname{li}}{64a^4d^2}}2i \\ & - \operatorname{atan}\left(\frac{16a^2d\sqrt{\tan(c+dx)}\sqrt{\frac{A^2\operatorname{li}}{256a^4d^2}}}{A}\right)\sqrt{\frac{A^2\operatorname{li}}{256a^4d^2}}2i \\ & + 2\operatorname{atanh}\left(\frac{8a^2d\sqrt{\tan(c+dx)}\sqrt{\frac{B^2\operatorname{li}}{64a^4d^2}}}{B}\right)\sqrt{\frac{B^2\operatorname{li}}{64a^4d^2}} \\ & + 2\operatorname{atanh}\left(\frac{16a^2d\sqrt{\tan(c+dx)}\sqrt{-\frac{B^2\operatorname{li}}{256a^4d^2}}}{7B}\right)\sqrt{-\frac{B^2\operatorname{li}}{256a^4d^2}} \end{aligned}$$

input `int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^2,x)`

output $((5*B*\tan(c + d*x)^{(1/2)})/(8*a^2*d) + (B*\tan(c + d*x)^{(3/2)}*7i)/(8*a^2*d)) / (2*\tan(c + d*x) + \tan(c + d*x)^{2*1i} - 1i) - ((A*\tan(c + d*x)^{(1/2)}*1i)/(8*a^2*d) - (3*A*\tan(c + d*x)^{(3/2)})/(8*a^2*d)) / (2*\tan(c + d*x) + \tan(c + d*x)^{2*1i} - 1i) - \operatorname{atan}((8*a^2*d*\tan(c + d*x)^{(1/2)}*(-(A^2*1i)/(64*a^4*d^2))^{(1/2)})/A)*(-(A^2*1i)/(64*a^4*d^2))^{(1/2)}*2i - \operatorname{atan}((16*a^2*d*\tan(c + d*x)^{(1/2)}*((A^2*1i)/(256*a^4*d^2))^{(1/2)})/A)*((A^2*1i)/(256*a^4*d^2))^{(1/2)}*2i + 2*\operatorname{atanh}((8*a^2*d*\tan(c + d*x)^{(1/2)}*((B^2*1i)/(64*a^4*d^2))^{(1/2)})/B)*((B^2*1i)/(64*a^4*d^2))^{(1/2)} + 2*\operatorname{atanh}((16*a^2*d*\tan(c + d*x)^{(1/2)}*(-(B^2*49i)/(256*a^4*d^2))^{(1/2)})/(7*B))*(-(B^2*49i)/(256*a^4*d^2))^{(1/2)}$

3.141. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$3.142 \quad \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.142.1 Optimal result 1470
 3.142.2 Mathematica [A] (verified) 1471
 3.142.3 Rubi [A] (verified) 1471
 3.142.4 Maple [A] (verified) 1476
 3.142.5 Fricas [B] (verification not implemented) 1477
 3.142.6 Sympy [F] 1478
 3.142.7 Maxima [F(-2)] 1478
 3.142.8 Giac [A] (verification not implemented) 1478
 3.142.9 Mupad [B] (verification not implemented) 1479

3.142.1 Optimal result

Integrand size = 36, antiderivative size = 279

$$\begin{aligned} & \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx \\ &= \frac{((-1+3i)A+(1+3i)B) \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{16\sqrt{2}a^2d} \\ & \quad - \frac{((-1+3i)A+(1+3i)B) \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{16\sqrt{2}a^2d} \\ & \quad + \frac{((1+3i)A+(1-3i)B) \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{32\sqrt{2}a^2d} \\ & \quad - \frac{((1+3i)A+(1-3i)B) \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{32\sqrt{2}a^2d} \\ & \quad + \frac{(iA+3B)\sqrt{\tan(c+dx)}}{8a^2d(1+i \tan(c+dx))} + \frac{(iA-B)\sqrt{\tan(c+dx)}}{4d(a+ia \tan(c+dx))^2} \end{aligned}$$

output $-1/32*((-1+3*I)*A+(1+3*I)*B)*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/a^2/d*2^{(1/2)}-1/32*((-1+3*I)*A+(1+3*I)*B)*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/a^2/d*2^{(1/2)}+1/64*((1+3*I)*A+(1-3*I)*B)*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/a^2/d*2^{(1/2)}-1/64*((1+3*I)*A+(1-3*I)*B)*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/a^2/d*2^{(1/2)}+1/8*(I*A+3*B)*\tan(d*x+c)^{(1/2)}/a^2/d/(1+I*\tan(d*x+c))+1/4*(I*A-B)*\tan(d*x+c)^{(1/2)}/d/(a+I*a*\tan(d*x+c))^2$

3.142. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

3.142.2 Mathematica [A] (verified)

Time = 2.81 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.63

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= \frac{-2\sqrt[4]{-1}(iA+B)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)\sec^2(c+dx)(\cos(2(c+dx))+i\sin(2(c+dx))) + \sqrt[4]{-1}}$$

input `Integrate[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])
^2,x]`

output `(-2*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]
]^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)]) + (-1)^(1/4)*((-I)*A + B)*ArcT
anh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] + I*Si
n[2*(c + d*x)]) + Sqrt[Tan[c + d*x]]*((-3*I)*A - B + (A - (3*I)*B)*Tan[c +
d*x]))/(8*a^2*d*(-I + Tan[c + d*x])^2)`

3.142.3 Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 246, normalized size of antiderivative = 0.88, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 4078, 27, 3042, 4079, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$\downarrow 3042$$

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$\downarrow 4078$$

$$\frac{(-B+iA)\sqrt{\tan(c+dx)}}{4d(a+ia\tan(c+dx))^2} - \frac{\int \frac{a(iA-B)-a(3A-5iB)\tan(c+dx)}{2\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)} dx}{4a^2}$$

$$\downarrow 27$$

3.142. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$

$$\begin{aligned}
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{a(iA - B) - a(3A - 5iB)\tan(c + dx)}{\sqrt{\tan(c + dx)}(i \tan(c + dx)a + a)} dx}{8a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{a(iA - B) - a(3A - 5iB)\tan(c + dx)}{\sqrt{\tan(c + dx)}(i \tan(c + dx)a + a)} dx}{8a^2} \\
 & \quad \downarrow \text{4079} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{a^2(3iA + B) - a^2(A - 3iB)\tan(c + dx)}{\sqrt{\tan(c + dx)}} dx}{2a^2} - \frac{(3B + iA)\sqrt{\tan(c + dx)}}{d(1 + i \tan(c + dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{a^2(3iA + B) - a^2(A - 3iB)\tan(c + dx)}{\sqrt{\tan(c + dx)}} dx}{2a^2} - \frac{(3B + iA)\sqrt{\tan(c + dx)}}{d(1 + i \tan(c + dx))} \\
 & \quad \downarrow \text{4017} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{a^2(3iA + B) - a^2(A - 3iB)\tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{a^2 d} - \frac{(3B + iA)\sqrt{\tan(c + dx)}}{d(1 + i \tan(c + dx))} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\int \frac{3iA + B - (A - 3iB)\tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{d} - \frac{(3B + iA)\sqrt{\tan(c + dx)}}{d(1 + i \tan(c + dx))} \\
 & \quad \downarrow \text{1482} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1 + 3i)A + (1 - 3i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \frac{1}{2}((1 + 3i)B - (1 - 3i)A) \int \frac{\tan(c + dx) + 1}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{d} - \frac{(3B + iA)\sqrt{\tan(c + dx)}}{d(1 + i \tan(c + dx))} \\
 & \quad \downarrow \text{1476} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1 + 3i)A + (1 - 3i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \frac{1}{2}((1 + 3i)B - (1 - 3i)A) \left(\frac{1}{2} \int \frac{1}{\tan(c + dx) - \sqrt{2}\sqrt{\tan(c + dx) + 1}} d\sqrt{\tan(c + dx)} + \frac{1}{2} \int \frac{1}{\tan(c + dx) + \sqrt{2}\sqrt{\tan(c + dx) + 1}} d\sqrt{\tan(c + dx)} \right)}{d} \\
 & \quad \downarrow \text{1082} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1 + 3i)A + (1 - 3i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \frac{1}{2}((1 + 3i)B - (1 - 3i)A) \left(\frac{1}{2} \int \frac{1}{\tan(c + dx) - \sqrt{2}\sqrt{\tan(c + dx) + 1}} d\sqrt{\tan(c + dx)} + \frac{1}{2} \int \frac{1}{\tan(c + dx) + \sqrt{2}\sqrt{\tan(c + dx) + 1}} d\sqrt{\tan(c + dx)} \right)}{d}
 \end{aligned}$$

3.142. $\int \frac{\sqrt{\tan(c + dx)}(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx$

$$\frac{\frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1+3i)A+(1-3i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((1+3i)B-(1-3i)A) \left(\frac{\int \frac{1}{\tan(c+dx)-1} d\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right) - \int \frac{1}{\tan(c+dx)-1} d\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{d} \right)}{8a^2}}$$

217

$$\frac{\frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1+3i)A+(1-3i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((1+3i)B-(1-3i)A) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right)}{d} - \frac{(3B+iA)}{d(1+i)}}{8a^2}}$$

1479

$$\frac{\frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1+3i)A+(1-3i)B) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}((1+3i)B-(1-3i)A) \left(\frac{a}{d} \right)}{d}}{8a^2}}$$

25

$$\frac{\frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1+3i)A+(1-3i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}((1+3i)B-(1-3i)A) \left(\frac{\arctan(a)}{d} \right)}{d}}{8a^2}}$$

27

$$\frac{\frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1+3i)A+(1-3i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}} \right) + \frac{1}{2}((1+3i)B-(1-3i)A) \left(\frac{\arctan(a)}{d} \right)}{d}}{8a^2}}$$

1103

3.142. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\frac{\frac{(-B + iA)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2} - \frac{\frac{1}{2}((1+3i)B - (1-3i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}) - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}((1+3i)A + (1-3i)B) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right)}{d}}{8a^2}$$

input `Int[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `-1/8*((((-1 + 3*I)*A + (1 + 3*I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 + (((1 + 3*I)*A + (1 - 3*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d - ((I*A + 3*B)*Sqrt[Tan[c + d*x]]/(d*(1 + I*Tan[c + d*x])))/a^2 + ((I*A - B)*Sqrt[Tan[c + d*x]]/(4*d*(a + I*a*Tan[c + d*x])^2)`

3.142.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4078 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

```
rule 4079 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.142.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.52

method	result
derivativedivides	$\frac{\left(-\frac{3iB}{2} + \frac{A}{2}\right) \left(\tan^{\frac{3}{2}}(dx+c)\right) + \left(-\frac{B}{2} - \frac{3iA}{2}\right) (\sqrt{\tan}(dx+c)}{4(\tan(dx+c)-i)^2} - \frac{(iB+A) \arctan\left(\frac{2(\sqrt{\tan}(dx+c))}{\sqrt{2-i\sqrt{2}}}\right)}{4(\sqrt{2-i\sqrt{2}})} - \frac{i(iA+B) \arctan\left(\frac{2(\sqrt{\tan}(dx+c))}{\sqrt{2+i\sqrt{2}}}\right)}{2(\sqrt{2+i\sqrt{2}})}\right)}{da^2}$
default	$\frac{\left(-\frac{3iB}{2} + \frac{A}{2}\right) \left(\tan^{\frac{3}{2}}(dx+c)\right) + \left(-\frac{B}{2} - \frac{3iA}{2}\right) (\sqrt{\tan}(dx+c)}{4(\tan(dx+c)-i)^2} - \frac{(iB+A) \arctan\left(\frac{2(\sqrt{\tan}(dx+c))}{\sqrt{2-i\sqrt{2}}}\right)}{4(\sqrt{2-i\sqrt{2}})} - \frac{i(iA+B) \arctan\left(\frac{2(\sqrt{\tan}(dx+c))}{\sqrt{2+i\sqrt{2}}}\right)}{2(\sqrt{2+i\sqrt{2}})}\right)}{da^2}$

```
input int(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x,method=_RETUR
NVERBOSE)
```

```
output 1/d/a^2*(1/4*((-3/2*I*B+1/2*A)*tan(d*x+c)^(3/2)+(-1/2*B-3/2*I*A)*tan(d*x+c)
)^(1/2))/(tan(d*x+c)-I)^2-1/4*(A+I*B)/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x
+c)^(1/2)/(2^(1/2)-I*2^(1/2)))-1/2*I*(I*A+B)/(2^(1/2)+I*2^(1/2))*arctan(2*
tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))
```

3.142.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 658 vs. $2(212) = 424$.

Time = 0.28 (sec) , antiderivative size = 658, normalized size of antiderivative = 2.36

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx =$$

$$\left(2a^2d\sqrt{\frac{iA^2+2AB-iB^2}{a^4d^2}}e^{(4idx+4ic)} \log\left(\frac{2\left((a^2de^{(2idx+2ic)}+a^2d)\sqrt{\frac{-ie^{(2idx+2ic)}+i}{e^{(2idx+2ic)}+1}}\sqrt{\frac{iA^2+2AB-iB^2}{a^4d^2}}+(A-iB)e^{(2idx+2ic)}}{iA+B} \right)} \right)$$

```
input integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorith
thm="fracas")
```

```
output -1/32*(2*a^2*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)
*log(2*((a^2*d*e^(2*I*d*x + 2*I*c) + a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) +
I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2)) + (
A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 2*a^2*d*sq
rt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(-2*((a^2*d*e
^(2*I*d*x + 2*I*c) + a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x
+ 2*I*c) + 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2)) - (A - I*B)*e^(2*I*
d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - a^2*d*sqrt((-I*A^2 + 2*A*B
+ I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(1/8*((a^2*d*e^(2*I*d*x + 2*I*
c) + a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*s
qrt((-I*A^2 + 2*A*B + I*B^2)/(a^4*d^2)) + A + I*B)*e^(-2*I*d*x - 2*I*c)/(a
^2*d)) + a^2*d*sqrt((-I*A^2 + 2*A*B + I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c
)*log(-1/8*((a^2*d*e^(2*I*d*x + 2*I*c) + a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*
c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 + 2*A*B + I*B^2)/(a^4*d^2)
) - A - I*B)*e^(-2*I*d*x - 2*I*c)/(a^2*d)) + 2*(2*(-I*A - B)*e^(4*I*d*x +
4*I*c) - (3*I*A + B)*e^(2*I*d*x + 2*I*c) - I*A + B)*sqrt((-I*e^(2*I*d*x +
2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*e^(-4*I*d*x - 4*I*c)/(a^2*d)
```

3.142.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= -\frac{\int \frac{A\sqrt{\tan(c+dx)}}{\tan^2(c+dx)-2i\tan(c+dx)-1} dx + \int \frac{B\tan^{\frac{3}{2}}(c+dx)}{\tan^2(c+dx)-2i\tan(c+dx)-1} dx}{a^2}$$

input `integrate(tan(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**2,x)`

output `-(Integral(A*sqrt(tan(c + d*x))/(tan(c + d*x)**2 - 2*I*tan(c + d*x) - 1), x) + Integral(B*tan(c + d*x)**(3/2)/(tan(c + d*x)**2 - 2*I*tan(c + d*x) - 1), x))/a**2`

3.142.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.142.8 Giac [A] (verification not implemented)

Time = 0.56 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.44

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= -\frac{(i+1)\sqrt{2}(A+iB)\arctan\left(\left(\frac{1}{2}i+\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{16a^2d}$$

$$- \frac{(i-1)\sqrt{2}(A-iB)\arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{8a^2d}$$

$$+ \frac{A\tan(dx+c)^{\frac{3}{2}}-3iB\tan(dx+c)^{\frac{3}{2}}-3iA\sqrt{\tan(dx+c)}-B\sqrt{\tan(dx+c)}}{8a^2d(\tan(dx+c)-i)^2}$$

3.142. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")`

output
$$-(1/16*I + 1/16)*\sqrt{2}*(A + I*B)*\arctan((1/2*I + 1/2)*\sqrt{2}*\sqrt{\tan(d*x + c)})/(a^2*d) - (1/8*I - 1/8)*\sqrt{2}*(A - I*B)*\arctan(-(1/2*I - 1/2)*\sqrt{2}*\sqrt{\tan(d*x + c)})/(a^2*d) + 1/8*(A*\tan(d*x + c)^{(3/2)} - 3*I*B*\tan(d*x + c)^{(3/2)} - 3*I*A*\sqrt{\tan(d*x + c)} - B*\sqrt{\tan(d*x + c)})/(a^2*d*(\tan(d*x + c) - I)^2)$$

3.142.9 Mupad [B] (verification not implemented)

Time = 11.29 (sec) , antiderivative size = 318, normalized size of antiderivative = 1.14

$$\begin{aligned} & \int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx \\ &= \frac{\frac{3A\sqrt{\tan(c+dx)}}{8a^2d} + \frac{A\tan(c+dx)^{3/2}i}{8a^2d}}{\tan(c+dx)^2i + 2\tan(c+dx) - i} - \frac{-\frac{3B\tan(c+dx)^{3/2}}{8a^2d} + \frac{B\sqrt{\tan(c+dx)}i}{8a^2d}}{\tan(c+dx)^2i + 2\tan(c+dx) - i} \\ & - 2\operatorname{atanh}\left(\frac{8a^2d\sqrt{\tan(c+dx)}\sqrt{\frac{A^2i}{64a^4d^2}}}{A}\right)\sqrt{\frac{A^2i}{64a^4d^2}} \\ & + 2\operatorname{atanh}\left(\frac{16a^2d\sqrt{\tan(c+dx)}\sqrt{-\frac{A^2i}{256a^4d^2}}}{A}\right)\sqrt{-\frac{A^2i}{256a^4d^2}} \\ & - \operatorname{atan}\left(\frac{8a^2d\sqrt{\tan(c+dx)}\sqrt{-\frac{B^2i}{64a^4d^2}}}{B}\right)\sqrt{-\frac{B^2i}{64a^4d^2}}2i \\ & - \operatorname{atan}\left(\frac{16a^2d\sqrt{\tan(c+dx)}\sqrt{\frac{B^2i}{256a^4d^2}}}{B}\right)\sqrt{\frac{B^2i}{256a^4d^2}}2i \end{aligned}$$

input `int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^2,x)`

output $((3A \tan(c + dx)^{1/2}) / (8a^2d) + (A \tan(c + dx)^{3/2} i) / (8a^2d)) / (2 \tan(c + dx) + \tan(c + dx)^2 i - 1i) - ((B \tan(c + dx)^{1/2} i) / (8a^2d) - (3B \tan(c + dx)^{3/2})) / (8a^2d) / (2 \tan(c + dx) + \tan(c + dx)^2 i - 1i) - 2 \operatorname{atanh}((8a^2d \tan(c + dx)^{1/2} * ((A^2 i) / (64a^4d^2))^{1/2}) / A * ((A^2 i) / (64a^4d^2))^{1/2} + 2 \operatorname{atanh}((16a^2d \tan(c + dx)^{1/2} * (-A^2 i) / (256a^4d^2))^{1/2}) / A * (-A^2 i) / (256a^4d^2))^{1/2} - \operatorname{atan}((8a^2d \tan(c + dx)^{1/2} * (-B^2 i) / (64a^4d^2))^{1/2}) / B * (-B^2 i) / (64a^4d^2))^{1/2} * 2i - \operatorname{atan}((16a^2d \tan(c + dx)^{1/2} * (B^2 i) / (256a^4d^2))^{1/2}) / B * (B^2 i) / (256a^4d^2))^{1/2} * 2i$

3.143
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)(a+ia \tan(c+dx))^2}} dx$$

3.143.1 Optimal result 1481
 3.143.2 Mathematica [A] (verified) 1482
 3.143.3 Rubi [A] (verified) 1482
 3.143.4 Maple [A] (verified) 1487
 3.143.5 Fricas [B] (verification not implemented) 1487
 3.143.6 Sympy [F(-2)] 1488
 3.143.7 Maxima [F(-2)] 1488
 3.143.8 Giac [A] (verification not implemented) 1489
 3.143.9 Mupad [B] (verification not implemented) 1490

3.143.1 Optimal result

Integrand size = 36, antiderivative size = 285

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)(a + ia \tan(c + dx))^2}} dx$$

$$= \frac{\left(\frac{1}{16} + \frac{i}{16}\right) ((-2 + 7i)A + (1 + 2i)B) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}a^2d}$$

$$+ \frac{((9 - 5i)A + (1 - 3i)B) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{16\sqrt{2}a^2d}$$

$$+ \frac{\left(\frac{1}{32} + \frac{i}{32}\right) ((-7 + 2i)A + (2 + i)B) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{\sqrt{2}a^2d}$$

$$+ \frac{((9 + 5i)A - (1 + 3i)B) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{32\sqrt{2}a^2d}$$

$$+ \frac{(5A + iB)\sqrt{\tan(c + dx)}}{8a^2d(1 + i \tan(c + dx))} + \frac{(A + iB)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2}$$

output

```
(-1/32-1/32*I)*((-2+7*I)*A+(1+2*I)*B)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/
a^2/d*2^(1/2)+1/32*((9-5*I)*A+(1-3*I)*B)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2)
)/a^2/d*2^(1/2)+(1/64+1/64*I)*((-7+2*I)*A+(2+I)*B)*ln(1-2^(1/2)*tan(d*x+c)
^(1/2)+tan(d*x+c))/a^2/d*2^(1/2)+1/64*((9+5*I)*A-(1+3*I)*B)*ln(1+2^(1/2)*t
an(d*x+c)^(1/2)+tan(d*x+c))/a^2/d*2^(1/2)+1/8*(5*A+I*B)*tan(d*x+c)^(1/2)/a
^2/d/(1+I*tan(d*x+c))+1/4*(A+I*B)*tan(d*x+c)^(1/2)/d/(a+I*a*tan(d*x+c))^2
```


3.143.2 Mathematica [A] (verified)

Time = 2.42 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.62

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} dx$$

$$= \frac{2\sqrt[4]{-1}(A - iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right) \sec^2(c + dx)(\cos(2(c + dx)) + i \sin(2(c + dx))) + \sqrt[4]{-1}(7A - iB) \operatorname{ArcTan}\left[(-1)^{3/4} \sqrt{\tan(c + dx)}\right] \sec^2(c + dx)(\cos(2(c + dx)) + i \sin(2(c + dx)))}{8a^2 d (-I + \tan(c + dx))^2}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^2),x]`

output `(2*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)]) + (-1)^(1/4)*(7*A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)]) + Sqrt[Tan[c + d*x]]*(-7*A - (3*I)*B + ((-5*I)*A + B)*Tan[c + d*x])/(8*a^2*d*(-I + Tan[c + d*x])^2)`

3.143.3 Rubi [A] (verified)

Time = 0.87 (sec) , antiderivative size = 243, normalized size of antiderivative = 0.85, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 4079, 27, 3042, 4079, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} dx$$

↓ 4079

$$\frac{\int \frac{a(7A - iB) - 3a(iA - B) \tan(c + dx)}{2\sqrt{\tan(c + dx)}(i \tan(c + dx)a + a)} dx}{4a^2} + \frac{(A + iB)\sqrt{\tan(c + dx)}}{4d(a + ia \tan(c + dx))^2}$$

↓ 27

3.143. $\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} dx$

$$\begin{aligned}
 & \frac{\int \frac{a(7A-iB)-3a(iA-B)\tan(c+dx)}{\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)} dx}{8a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{4d(a+ia\tan(c+dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a(7A-iB)-3a(iA-B)\tan(c+dx)}{\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)} dx}{8a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{4d(a+ia\tan(c+dx))^2} \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{3a^2(3A-iB)-a^2(5iA-B)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{8a^2} + \frac{(5A+iB)\sqrt{\tan(c+dx)}}{d(1+i\tan(c+dx))} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{4d(a+ia\tan(c+dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{3a^2(3A-iB)-a^2(5iA-B)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{8a^2} + \frac{(5A+iB)\sqrt{\tan(c+dx)}}{d(1+i\tan(c+dx))} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{4d(a+ia\tan(c+dx))^2} \\
 & \quad \downarrow \text{4017} \\
 & \frac{\int \frac{a^2(3(3A-iB)-(5iA-B)\tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{8a^2} + \frac{(5A+iB)\sqrt{\tan(c+dx)}}{d(1+i\tan(c+dx))} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{4d(a+ia\tan(c+dx))^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{3(3A-iB)-(5iA-B)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{8a^2} + \frac{(5A+iB)\sqrt{\tan(c+dx)}}{d(1+i\tan(c+dx))} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{4d(a+ia\tan(c+dx))^2} \\
 & \quad \downarrow \text{1482} \\
 & \frac{\frac{1}{2}((9+5i)A-(1+3i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((9-5i)A+(1-3i)B) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} + \frac{(5A+iB)\sqrt{\tan(c+dx)}}{d(1+i\tan(c+dx))} + \\
 & \quad \frac{8a^2}{4d(a+ia\tan(c+dx))^2} \\
 & \quad \downarrow \text{1476} \\
 & \frac{\frac{1}{2}((9+5i)A-(1+3i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((9-5i)A+(1-3i)B) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right)}{8a^2} \\
 & \quad \frac{(A+iB)\sqrt{\tan(c+dx)}}{4d(a+ia\tan(c+dx))^2} \\
 & \quad \downarrow \text{1082}
 \end{aligned}$$

3.143. $\int \frac{A+B\tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} dx$

$$\frac{\frac{1}{2}((9+5i)A-(1+3i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((9-5i)A+(1-3i)B) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} \frac{d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} \frac{d(\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}}}{d}}{8a^2}$$

$$\frac{(A+iB)\sqrt{\tan(c+dx)}}{4d(a+ia\tan(c+dx))^2}$$

↓ 217

$$\frac{\frac{1}{2}((9+5i)A-(1+3i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}((9-5i)A+(1-3i)B) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)}+1}{\sqrt{2}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{d} \right) + \frac{(5A+iB)}{d(1+i}}$$

$$\frac{(A+iB)\sqrt{\tan(c+dx)}}{4d(a+ia\tan(c+dx))^2} \frac{8a^2}{8a^2}$$

↓ 1479

$$\frac{\frac{1}{2}((9+5i)A-(1+3i)B) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}((9-5i)A+(1-3i)B) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)}+1}{\sqrt{2}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{d} \right)}{8a^2}$$

$$\frac{(A+iB)\sqrt{\tan(c+dx)}}{4d(a+ia\tan(c+dx))^2}$$

↓ 25

$$\frac{\frac{1}{2}((9+5i)A-(1+3i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}((9-5i)A+(1-3i)B) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)}+1}{\sqrt{2}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{d} \right)}{8a^2}$$

$$\frac{(A+iB)\sqrt{\tan(c+dx)}}{4d(a+ia\tan(c+dx))^2}$$

↓ 27

$$\frac{\frac{1}{2}((9+5i)A-(1+3i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)}+1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2}((9-5i)A+(1-3i)B) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)}+1}{\sqrt{2}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{d} \right)}{8a^2}$$

$$\frac{(A+iB)\sqrt{\tan(c+dx)}}{4d(a+ia\tan(c+dx))^2}$$

↓ 1103

3.143. $\int \frac{A+B\tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} dx$

$$\frac{\frac{1}{2}((9-5i)A+(1-3i)B)\left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}}\right) + \frac{1}{2}((9+5i)A-(1+3i)B)\left(\frac{\log\left(\frac{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}{2\sqrt{2}}\right) - \log\left(\frac{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}}{2\sqrt{2}}\right)}{\sqrt{2}}\right)}{d} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{4d(a+ia\tan(c+dx))^2}$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^2),x]`

output `(((((9 - 5*I)*A + (1 - 3*I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 + (((9 + 5*I)*A - (1 + 3*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d + ((5*A + I*B)*Sqrt[Tan[c + d*x]]/(d*(1 + I*Tan[c + d*x])))/(8*a^2) + ((A + I*B)*Sqrt[Tan[c + d*x]])/(4*d*(a + I*a*Tan[c + d*x])^2)`

3.143.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

3.143. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)(a+ia \tan(c+dx))^2}} dx$

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.143.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.51

method	result
derivativedivides	$\frac{\left(\frac{5iA}{2} - \frac{B}{2}\right) \left(\tan^{\frac{3}{2}}(dx+c)\right) + \left(\frac{7A}{2} + \frac{3iB}{2}\right) (\sqrt{\tan(dx+c)})}{4(\tan(dx+c)-i)^2} - \frac{(7iA+B) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{4(\sqrt{2-i\sqrt{2}})} + \frac{i(-iB+A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2+i\sqrt{2}}}\right)}{2\sqrt{2}+2i\sqrt{2}}$
default	$\frac{\left(\frac{5iA}{2} - \frac{B}{2}\right) \left(\tan^{\frac{3}{2}}(dx+c)\right) + \left(\frac{7A}{2} + \frac{3iB}{2}\right) (\sqrt{\tan(dx+c)})}{4(\tan(dx+c)-i)^2} - \frac{(7iA+B) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{4(\sqrt{2-i\sqrt{2}})} + \frac{i(-iB+A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2+i\sqrt{2}}}\right)}{2\sqrt{2}+2i\sqrt{2}}$

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d/a^2*(-1/4*((5/2*I*A-1/2*B)*tan(d*x+c)^(3/2)+(7/2*A+3/2*I*B)*tan(d*x+c)^(1/2))/(tan(d*x+c)-I)^2-1/4*(7*I*A+B)/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))+1/2*I*(A-I*B)/(2^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))`

3.143.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 664 vs. 2(212) = 424.

Time = 0.27 (sec) , antiderivative size = 664, normalized size of antiderivative = 2.33

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} dx =$$

$$\left(2 a^2 d \sqrt{\frac{-i A^2 - 2 AB + i B^2}{a^4 d^2}} e^{(4i dx + 4i c)} \log \left(-\frac{2 \left((i a^2 d e^{(2i dx + 2i c)} + i a^2 d) \sqrt{\frac{-i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} + 1}} \sqrt{\frac{-i A^2 - 2 AB + i B^2}{a^4 d^2}} - (A - i B) e^{(2i dx + 2i c)} \right)}{i A + B} \right) \right)$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="fracas")`

output

```
-1/32*(2*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)
)*log(-2*((I*a^2*d*e^(2*I*d*x + 2*I*c) + I*a^2*d)*sqrt((-I*e^(2*I*d*x + 2*
I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^
2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 2*a
^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(-2*(
(-I*a^2*d*e^(2*I*d*x + 2*I*c) - I*a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)
/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2)) - (A
- I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - a^2*d*sqrt((
49*I*A^2 + 14*A*B - I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(1/8*((a^2*d*
e^(2*I*d*x + 2*I*c) + a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x
+ 2*I*c) + 1))*sqrt((49*I*A^2 + 14*A*B - I*B^2)/(a^4*d^2)) + 7*I*A + B)*e
^(-2*I*d*x - 2*I*c)/(a^2*d)) + a^2*d*sqrt((49*I*A^2 + 14*A*B - I*B^2)/(a^4
*d^2))*e^(4*I*d*x + 4*I*c)*log(-1/8*((a^2*d*e^(2*I*d*x + 2*I*c) + a^2*d)*s
qrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((49*I*A^2
+ 14*A*B - I*B^2)/(a^4*d^2)) - 7*I*A - B)*e^(-2*I*d*x - 2*I*c)/(a^2*d)) -
2*(2*(3*A + I*B)*e^(4*I*d*x + 4*I*c) + (7*A + 3*I*B)*e^(2*I*d*x + 2*I*c)
+ A + I*B)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*e
^(-4*I*d*x - 4*I*c)/(a^2*d)
```

3.143.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)(a + ia \tan(c + dx))^2}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+I*a*tan(d*x+c))**2,x)`

output `Exception raised: TypeError >> Invalid comparison of non-real -I`

3.143.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)(a + ia \tan(c + dx))^2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")`

3.143. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)(a+ia \tan(c+dx))^2}} dx$

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

3.143.8 Giac [A] (verification not implemented)

Time = 0.87 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.44

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} dx$$

$$= -\frac{(i + 1) \sqrt{2}(7iA + B) \arctan\left(\left(\frac{1}{2}i + \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{16a^2d}$$

$$+ \frac{(i - 1) \sqrt{2}(-iA - B) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{8a^2d}$$

$$- \frac{5iA \tan(dx + c)^{\frac{3}{2}} - B \tan(dx + c)^{\frac{3}{2}} + 7A \sqrt{\tan(dx + c)} + 3iB \sqrt{\tan(dx + c)}}{8a^2d(\tan(dx + c) - i)^2}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")`

output `-(1/16*I + 1/16)*sqrt(2)*(7*I*A + B)*arctan((1/2*I + 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^2*d) + (1/8*I - 1/8)*sqrt(2)*(-I*A - B)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^2*d) - 1/8*(5*I*A*tan(d*x + c)^(3/2) - B*tan(d*x + c)^(3/2) + 7*A*sqrt(tan(d*x + c)) + 3*I*B*sqrt(tan(d*x + c)))/(a^2*d*(tan(d*x + c) - I)^2)`

3.143.9 Mupad [B] (verification not implemented)

Time = 11.11 (sec) , antiderivative size = 318, normalized size of antiderivative = 1.12

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)(a + ia \tan(c + dx))^2}} dx$$

$$= -\frac{-\frac{5A \tan(c+dx)^{3/2}}{8a^2 d} + \frac{A \sqrt{\tan(c+dx)} 7i}{8a^2 d}}{\tan(c+dx)^2 li + 2 \tan(c+dx) - i} + \frac{\frac{3B \sqrt{\tan(c+dx)}}{8a^2 d} + \frac{B \tan(c+dx)^{3/2} li}{8a^2 d}}{\tan(c+dx)^2 li + 2 \tan(c+dx) - i}$$

$$+ \operatorname{atan}\left(\frac{8a^2 d \sqrt{\tan(c+dx)} \sqrt{-\frac{A^2 li}{64a^4 d^2}}}{A}\right) \sqrt{-\frac{A^2 li}{64a^4 d^2}} 2i$$

$$- \operatorname{atan}\left(\frac{16a^2 d \sqrt{\tan(c+dx)} \sqrt{\frac{A^2 49i}{256a^4 d^2}}}{7A}\right) \sqrt{\frac{A^2 49i}{256a^4 d^2}} 2i$$

$$- 2 \operatorname{atanh}\left(\frac{8a^2 d \sqrt{\tan(c+dx)} \sqrt{\frac{B^2 li}{64a^4 d^2}}}{B}\right) \sqrt{\frac{B^2 li}{64a^4 d^2}}$$

$$+ 2 \operatorname{atanh}\left(\frac{16a^2 d \sqrt{\tan(c+dx)} \sqrt{-\frac{B^2 li}{256a^4 d^2}}}{B}\right) \sqrt{-\frac{B^2 li}{256a^4 d^2}}$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*li)^2),x)`

output `((3*B*tan(c + d*x)^(1/2))/(8*a^2*d) + (B*tan(c + d*x)^(3/2)*li)/(8*a^2*d)) / (2*tan(c + d*x) + tan(c + d*x)^2*li - li) - ((A*tan(c + d*x)^(1/2)*7i)/(8*a^2*d) - (5*A*tan(c + d*x)^(3/2))/(8*a^2*d)) / (2*tan(c + d*x) + tan(c + d*x)^2*li - li) + atan((8*a^2*d*tan(c + d*x)^(1/2)*(-(A^2*li)/(64*a^4*d^2))^(1/2))/A)*(-(A^2*li)/(64*a^4*d^2))^(1/2)*2i - atan((16*a^2*d*tan(c + d*x)^(1/2)*((A^2*49i)/(256*a^4*d^2))^(1/2))/(7*A))*((A^2*49i)/(256*a^4*d^2))^(1/2)*2i - 2*atanh((8*a^2*d*tan(c + d*x)^(1/2)*((B^2*li)/(64*a^4*d^2))^(1/2))/B)*((B^2*li)/(64*a^4*d^2))^(1/2) + 2*atanh((16*a^2*d*tan(c + d*x)^(1/2)*(-(B^2*li)/(256*a^4*d^2))^(1/2))/B)*(-(B^2*li)/(256*a^4*d^2))^(1/2)`

$$3.144 \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$$

3.144.1 Optimal result	1491
3.144.2 Mathematica [C] (verified)	1492
3.144.3 Rubi [A] (verified)	1492
3.144.4 Maple [A] (verified)	1498
3.144.5 Fricas [B] (verification not implemented)	1499
3.144.6 Sympy [F(-2)]	1500
3.144.7 Maxima [F(-2)]	1500
3.144.8 Giac [A] (verification not implemented)	1500
3.144.9 Mupad [B] (verification not implemented)	1501

3.144.1 Optimal result

Integrand size = 36, antiderivative size = 318

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx \\ &= \frac{((25 + 21i)A - (9 - 5i)B) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{16\sqrt{2}a^2d} \\ & \quad - \frac{\left(\frac{1}{16} - \frac{i}{16}\right) ((2 + 23i)A - (7 + 2i)B) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}a^2d} \\ & \quad - \frac{\left(\frac{1}{32} - \frac{i}{32}\right) ((23 + 2i)A + (2 + 7i)B) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{\sqrt{2}a^2d} \\ & \quad + \frac{\left(\frac{1}{32} - \frac{i}{32}\right) ((23 + 2i)A + (2 + 7i)B) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{\sqrt{2}a^2d} \\ & \quad - \frac{5(5A + iB)}{8a^2d\sqrt{\tan(c + dx)}} + \frac{7A + 3iB}{8a^2d(1 + i \tan(c + dx))\sqrt{\tan(c + dx)}} \\ & \quad + \frac{A + iB}{4d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} \end{aligned}$$

3.144. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

output
$$-1/32*((25+21*I)*A+(-9+5*I)*B)*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/a^2/d*2^{(1/2)}+(-1/32+1/32*I)*((2+23*I)*A-(7+2*I)*B)*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/a^2/d*2^{(1/2)}+(-1/64+1/64*I)*((23+2*I)*A+(2+7*I)*B)*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/a^2/d*2^{(1/2)}+(1/64-1/64*I)*((23+2*I)*A+(2+7*I)*B)*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/a^2/d*2^{(1/2)}-5/8*(5*A+I*B)/a^2/d/\tan(d*x+c)^{(1/2)}+1/8*(7*A+3*I*B)/a^2/d/\tan(d*x+c)^{(1/2)}/(1+I*\tan(d*x+c))+1/4*(A+I*B)/d/\tan(d*x+c)^{(1/2)}/(a+I*a*\tan(d*x+c))^2$$

3.144.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 2.04 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.57

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx$$

$$= \frac{\sec^2(c + dx) (-2 \cos(c + dx)((9A + 5iB) \cos(c + dx) + (7iA - 3B) \sin(c + dx)) + 2(23A + 7iB) \operatorname{Hypergeometric2F1}[-1/2, 1, 1/2, I \tan(c + dx)]}{(16a^2 d \sqrt{\tan(c + dx)} (-I + \tan(c + dx))^2)}$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^2),x]`

output
$$(\operatorname{Sec}[c + d*x]^2*(-2*\operatorname{Cos}[c + d*x]*((9*A + (5*I)*B)*\operatorname{Cos}[c + d*x] + ((7*I)*A - 3*B)*\operatorname{Sin}[c + d*x]) + 2*(23*A + (7*I)*B)*\operatorname{Hypergeometric2F1}[-1/2, 1, 1/2, (-I)*\operatorname{Tan}[c + d*x]]*(\operatorname{Cos}[2*(c + d*x)] + I*\operatorname{Sin}[2*(c + d*x)]) + 4*(A - I*B)*\operatorname{Hypergeometric2F1}[-1/2, 1, 1/2, I*\operatorname{Tan}[c + d*x]]*(\operatorname{Cos}[2*(c + d*x)] + I*\operatorname{Sin}[2*(c + d*x)])))/(16*a^2*d*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]*(-I + \operatorname{Tan}[c + d*x])^2)$$

3.144.3 Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 286, normalized size of antiderivative = 0.90, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.528$, Rules used = {3042, 4079, 27, 3042, 4079, 3042, 4012, 25, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.144.
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$$

$$\begin{aligned}
& \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2}(a + ia \tan(c + dx))^2} dx \\
& \quad \downarrow \text{4079} \\
& \frac{\int \frac{a(9A+iB)-5a(iA-B) \tan(c+dx)}{2 \tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a)} dx}{4a^2} + \frac{A + iB}{4d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{a(9A+iB)-5a(iA-B) \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a)} dx}{8a^2} + \frac{A + iB}{4d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(9A+iB)-5a(iA-B) \tan(c+dx)}{\tan(c+dx)^{3/2}(i \tan(c+dx)a+a)} dx}{8a^2} + \frac{A + iB}{4d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} \\
& \quad \downarrow \text{4079} \\
& \frac{\int \frac{5a^2(5A+iB)-3a^2(7iA-3B) \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)} dx}{2a^2} + \frac{7A+3iB}{d(1+i \tan(c+dx))\sqrt{\tan(c+dx)}} + \\
& \quad \frac{8a^2}{A + iB} \\
& \quad \frac{A + iB}{4d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{5a^2(5A+iB)-3a^2(7iA-3B) \tan(c+dx)}{\tan(c+dx)^{3/2}} dx}{2a^2} + \frac{7A+3iB}{d(1+i \tan(c+dx))\sqrt{\tan(c+dx)}} + \\
& \quad \frac{8a^2}{A + iB} \\
& \quad \frac{A + iB}{4d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} \\
& \quad \downarrow \text{4012} \\
& \frac{\int -\frac{3(7iA-3B)a^2+5(5A+iB) \tan(c+dx)a^2}{\sqrt{\tan(c+dx)}} dx - \frac{10a^2(5A+iB)}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{7A+3iB}{d(1+i \tan(c+dx))\sqrt{\tan(c+dx)}} + \\
& \quad \frac{8a^2}{A + iB} \\
& \quad \frac{A + iB}{4d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} \\
& \quad \downarrow \text{25}
\end{aligned}$$

3.144. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
 & \frac{-\int \frac{3(7iA-3B)a^2+5(5A+iB)\tan(c+dx)a^2}{\sqrt{\tan(c+dx)}} dx - \frac{10a^2(5A+iB)}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{7A+3iB}{d(1+i\tan(c+dx))\sqrt{\tan(c+dx)}} + \\
 & \frac{8a^2}{A+iB} \\
 & \frac{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2}{\phantom{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\int \frac{3(7iA-3B)a^2+5(5A+iB)\tan(c+dx)a^2}{\sqrt{\tan(c+dx)}} dx - \frac{10a^2(5A+iB)}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{7A+3iB}{d(1+i\tan(c+dx))\sqrt{\tan(c+dx)}} + \\
 & \frac{8a^2}{A+iB} \\
 & \frac{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2}{\phantom{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2}} \\
 & \quad \downarrow \text{4017} \\
 & \frac{2\int \frac{a^2(3(7iA-3B)+5(5A+iB)\tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{10a^2(5A+iB)}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{7A+3iB}{d(1+i\tan(c+dx))\sqrt{\tan(c+dx)}} + \\
 & \frac{8a^2}{A+iB} \\
 & \frac{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2}{\phantom{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{2a^2\int \frac{3(7iA-3B)+5(5A+iB)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{10a^2(5A+iB)}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{7A+3iB}{d(1+i\tan(c+dx))\sqrt{\tan(c+dx)}} + \\
 & \frac{8a^2}{A+iB} \\
 & \frac{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2}{\phantom{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2}} \\
 & \quad \downarrow \text{1482} \\
 & \frac{2a^2\left(\frac{1}{2}((25+21i)A-(9-5i)B)\int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \left(\frac{1}{2}-\frac{i}{2}\right)((23+2i)A+(2+7i)B)\int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right) - \frac{10a^2(5A+iB)}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{7}{d(1+i\tan(c+dx))\sqrt{\tan(c+dx)}} + \\
 & \frac{8a^2}{A+iB} \\
 & \frac{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2}{\phantom{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2}} \\
 & \quad \downarrow \text{1476} \\
 & \frac{2a^2\left(\frac{1}{2}((25+21i)A-(9-5i)B)\left(\frac{1}{2}\int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2}\int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}\right) - \left(\frac{1}{2}-\frac{i}{2}\right)((23+2i)A+(2+7i)B)\int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right) - \frac{10a^2(5A+iB)}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{7}{d(1+i\tan(c+dx))\sqrt{\tan(c+dx)}} + \\
 & \frac{8a^2}{A+iB} \\
 & \frac{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2}{\phantom{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2}} \\
 & \quad \downarrow \text{1082}
 \end{aligned}$$

3.144. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^2} dx$

$$\begin{aligned}
 & \frac{2a^2 \left(\frac{1}{2}((25+21i)A - (9-5i)B) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} \frac{d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} \frac{d(\sqrt{2}\sqrt{\tan(c+dx)+1)}}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((23+2i)A + (2+7i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{2a^2} \\
 & \frac{A+iB}{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} \qquad 8a^2 \\
 & \quad \downarrow \text{217} \\
 & \frac{2a^2 \left(\frac{1}{2}((25+21i)A - (9-5i)B) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}) - \arctan(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}})}{d} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((23+2i)A + (2+7i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{2a^2} \\
 & \frac{A+iB}{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} \qquad 8a^2 \\
 & \quad \downarrow \text{1479} \\
 & \frac{2a^2 \left(\frac{1}{2}((25+21i)A - (9-5i)B) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}) - \arctan(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}})}{d} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((23+2i)A + (2+7i)B) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} \frac{d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{2a^2} \\
 & \frac{A+iB}{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} \qquad 8a^2 \\
 & \quad \downarrow \text{25} \\
 & \frac{2a^2 \left(\frac{1}{2}((25+21i)A - (9-5i)B) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}) - \arctan(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}})}{d} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((23+2i)A + (2+7i)B) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} \frac{d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{2a^2} \\
 & \frac{A+iB}{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} \qquad 8a^2 \\
 & \quad \downarrow \text{27} \\
 & \frac{2a^2 \left(\frac{1}{2}((25+21i)A - (9-5i)B) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}) - \arctan(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}})}{d} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((23+2i)A + (2+7i)B) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} \frac{d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{2a^2} \\
 & \frac{A+iB}{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} \qquad 8a^2
 \end{aligned}$$

3.144. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

↓ 1103

$$\frac{2a^2 \left(\frac{1}{2}((25+21i)A - (9-5i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2} \right) ((23+2i)A + (2+7i)B) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d} \frac{A+iB}{8a^2} \frac{1}{4d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^2), x]`

output `(((-2*a^2*(((25 + 21*I)*A - (9 - 5*I)*B)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])))/2 - (1/2 - I/2)*(((23 + 2*I)*A + (2 + 7*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]))))/d - (10*a^2*(5*A + I*B))/(d*Sqrt[Tan[c + d*x]])/(2*a^2) + (7*A + (3*I)*B)/(d*(1 + I*Tan[c + d*x])*Sqrt[Tan[c + d*x]])/(8*a^2) + (A + I*B)/(4*d*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^2)`

3.144.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`


```
rule 4079 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.144.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.49

method	result
derivativedivides	$\frac{\left(-\frac{9A}{2} - \frac{5iB}{2}\right)\left(\tan^{\frac{3}{2}}(dx+c)\right) + \left(\frac{11iA}{2} - \frac{7B}{2}\right)(\sqrt{\tan(dx+c)}) - (7iB+23A)\arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{4(\tan(dx+c)-i)^2} - \frac{(7iB+23A)\arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{4(\sqrt{2-i\sqrt{2}})} - \frac{2A}{\sqrt{\tan(dx+c)}} + \frac{i(iA+B)\arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{2\sqrt{\tan(dx+c)}}$
default	$\frac{\left(-\frac{9A}{2} - \frac{5iB}{2}\right)\left(\tan^{\frac{3}{2}}(dx+c)\right) + \left(\frac{11iA}{2} - \frac{7B}{2}\right)(\sqrt{\tan(dx+c)}) - (7iB+23A)\arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{4(\tan(dx+c)-i)^2} - \frac{(7iB+23A)\arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{4(\sqrt{2-i\sqrt{2}})} - \frac{2A}{\sqrt{\tan(dx+c)}} + \frac{i(iA+B)\arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{2\sqrt{\tan(dx+c)}}$

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^2,x,method=_RETUR
NVERBOSE)
```

```
output 1/d/a^2*(1/4*((-9/2*A-5/2*I*B)*tan(d*x+c)^(3/2)+(11/2*I*A-7/2*B)*tan(d*x+c
)^(1/2))/(tan(d*x+c)-I)^2-1/4*(23*A+7*I*B)/(2^(1/2)-I*2^(1/2))*arctan(2*ta
n(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))-2*A/tan(d*x+c)^(1/2)+1/2*I*(I*A+B)/(2^
(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2))))
```

$$3.144. \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$$

3.144.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 763 vs. $2(235) = 470$.

Time = 0.28 (sec) , antiderivative size = 763, normalized size of antiderivative = 2.40

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx$$

$$= \frac{2 \left(a^2 de^{(6i dx + 6i c)} - a^2 de^{(4i dx + 4i c)} \right) \sqrt{\frac{i A^2 + 2 AB - i B^2}{a^4 d^2}} \log \left(\frac{2 \left((a^2 de^{(2i dx + 2i c)} + a^2 d) \sqrt{\frac{-i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} + 1}} \sqrt{\frac{i A^2 + 2 AB - i B^2}{a^4 d^2}} + (A - I B) e^{(2i dx + 2i c)} \right)}{i A + B} \right)}{1}$$

```
input integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="fricas")
```

```
output 1/32*(2*(a^2*d*e^(6*I*d*x + 6*I*c) - a^2*d*e^(4*I*d*x + 4*I*c))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))*log(2*((a^2*d*e^(2*I*d*x + 2*I*c) + a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 2*(a^2*d*e^(6*I*d*x + 6*I*c) - a^2*d*e^(4*I*d*x + 4*I*c))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))*log(-2*((a^2*d*e^(2*I*d*x + 2*I*c) + a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) + (a^2*d*e^(6*I*d*x + 6*I*c) - a^2*d*e^(4*I*d*x + 4*I*c))*sqrt((-529*I*A^2 + 322*A*B + 49*I*B^2)/(a^4*d^2))*log(1/8*((a^2*d*e^(2*I*d*x + 2*I*c) + a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-529*I*A^2 + 322*A*B + 49*I*B^2)/(a^4*d^2)) + 23*A + 7*I*B)*e^(-2*I*d*x - 2*I*c)/(a^2*d)) - (a^2*d*e^(6*I*d*x + 6*I*c) - a^2*d*e^(4*I*d*x + 4*I*c))*sqrt((-529*I*A^2 + 322*A*B + 49*I*B^2)/(a^4*d^2))*log(-1/8*((a^2*d*e^(2*I*d*x + 2*I*c) + a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-529*I*A^2 + 322*A*B + 49*I*B^2)/(a^4*d^2)) - 23*A - 7*I*B)*e^(-2*I*d*x - 2*I*c)/(a^2*d)) - 2*(6*(7*I*A - B)*e^(6*I*d*x + 6*I*c) - (-33*I*A + B)*e^(4*I*d*x + 4*I*c) + 2*(-5*I*A + 3*B)*e^(2*I*d*x + 2*I*c) - I*A + B)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(a^2*d*e^(6*I*d*x + 6*I*c) - a^2*d*e^(4*I*d*x ...
```

3.144.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(3/2)/(a+I*a*tan(d*x+c))**2,x)`

output `Exception raised: TypeError >> Invalid comparison of non-real -I`

3.144.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.144.8 Giac [A] (verification not implemented)

Time = 1.22 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.45

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx \\ &= -\frac{(i + 1) \sqrt{2}(23 A + 7i B) \arctan\left(\left(\frac{1}{2}i + \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{16 a^2 d} \\ &+ \frac{(i - 1) \sqrt{2}(A - i B) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{8 a^2 d} - \frac{2 A}{a^2 d \sqrt{\tan(dx + c)}} \\ &- \frac{9 A \tan(dx + c)^{\frac{3}{2}} + 5i B \tan(dx + c)^{\frac{3}{2}} - 11i A \sqrt{\tan(dx + c)} + 7 B \sqrt{\tan(dx + c)}}{8 a^2 d (\tan(dx + c) - i)^2} \end{aligned}$$

3.144. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")`

output
$$-(1/16*I + 1/16)*\sqrt{2}*(23*A + 7*I*B)*\arctan((1/2*I + 1/2)*\sqrt{2}*\sqrt{\tan(d*x + c)})/(a^2*d) + (1/8*I - 1/8)*\sqrt{2}*(A - I*B)*\arctan(-(1/2*I - 1/2)*\sqrt{2}*\sqrt{\tan(d*x + c)})/(a^2*d) - 2*A/(a^2*d*\sqrt{\tan(d*x + c)}) - 1/8*(9*A*\tan(d*x + c)^{(3/2)} + 5*I*B*\tan(d*x + c)^{(3/2)} - 11*I*A*\sqrt{\tan(d*x + c)} + 7*B*\sqrt{\tan(d*x + c)})/(a^2*d*(\tan(d*x + c) - I)^2)$$

3.144.9 Mupad [B] (verification not implemented)

Time = 11.93 (sec) , antiderivative size = 338, normalized size of antiderivative = 1.06

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx \\ &= -\frac{-\frac{5B \tan(c+dx)^{3/2}}{8a^2d} + \frac{B \sqrt{\tan(c+dx)} 7i}{8a^2d}}{\tan(c+dx)^2 li + 2 \tan(c+dx) - i} \\ &+ 2 \operatorname{atanh}\left(\frac{8a^2d \sqrt{\tan(c+dx)} \sqrt{\frac{A^2 li}{64a^4d^2}}}{A}\right) \sqrt{\frac{A^2 li}{64a^4d^2}} \\ &+ 2 \operatorname{atanh}\left(\frac{16a^2d \sqrt{\tan(c+dx)} \sqrt{-\frac{A^2 529i}{256a^4d^2}}}{23A}\right) \sqrt{-\frac{A^2 529i}{256a^4d^2}} \\ &+ \operatorname{atan}\left(\frac{8a^2d \sqrt{\tan(c+dx)} \sqrt{-\frac{B^2 li}{64a^4d^2}}}{B}\right) \sqrt{-\frac{B^2 li}{64a^4d^2}} 2i \\ &- \operatorname{atan}\left(\frac{16a^2d \sqrt{\tan(c+dx)} \sqrt{\frac{B^2 49i}{256a^4d^2}}}{7B}\right) \sqrt{\frac{B^2 49i}{256a^4d^2}} 2i \\ &- \frac{\frac{43A \tan(c+dx)}{8a^2d} - \frac{A 2i}{a^2d} + \frac{A \tan(c+dx)^2 25i}{8a^2d}}{2 \tan(c+dx)^{3/2} - \sqrt{\tan(c+dx)} li + \tan(c+dx)^{5/2} li} \end{aligned}$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + a*tan(c + d*x)*I)^2),x)`

output

```

2*atanh((8*a^2*d*tan(c + d*x)^(1/2)*((A^2*1i)/(64*a^4*d^2))^(1/2))/A)*((A^
2*1i)/(64*a^4*d^2))^(1/2) - ((B*tan(c + d*x)^(1/2)*7i)/(8*a^2*d) - (5*B*ta
n(c + d*x)^(3/2))/(8*a^2*d))/(2*tan(c + d*x) + tan(c + d*x)^2*1i - 1i) + 2
*atanh((16*a^2*d*tan(c + d*x)^(1/2)*(-(A^2*529i)/(256*a^4*d^2))^(1/2))/(23
*A))*(-(A^2*529i)/(256*a^4*d^2))^(1/2) + atan((8*a^2*d*tan(c + d*x)^(1/2)*
(-(B^2*1i)/(64*a^4*d^2))^(1/2))/B)*(-(B^2*1i)/(64*a^4*d^2))^(1/2)*2i - ata
n((16*a^2*d*tan(c + d*x)^(1/2)*((B^2*49i)/(256*a^4*d^2))^(1/2))/(7*B))*((B
^2*49i)/(256*a^4*d^2))^(1/2)*2i - ((43*A*tan(c + d*x))/(8*a^2*d) - (A*2i)/
(a^2*d) + (A*tan(c + d*x)^2*25i)/(8*a^2*d))/(2*tan(c + d*x)^(3/2) - tan(c
+ d*x)^(1/2)*1i + tan(c + d*x)^(5/2)*1i)

```

3.144.
$$\int \frac{A+B \tan(c+dx)}{\tan^3(c+dx)(a+ia \tan(c+dx))^2} dx$$

$$3.145 \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$$

3.145.1 Optimal result	1503
3.145.2 Mathematica [C] (verified)	1504
3.145.3 Rubi [A] (verified)	1504
3.145.4 Maple [A] (verified)	1510
3.145.5 Fricas [B] (verification not implemented)	1511
3.145.6 Sympy [F(-1)]	1512
3.145.7 Maxima [F(-2)]	1513
3.145.8 Giac [A] (verification not implemented)	1513
3.145.9 Mupad [B] (verification not implemented)	1514

3.145.1 Optimal result

Integrand size = 36, antiderivative size = 347

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx \\ &= \frac{\left(\frac{1}{16} - \frac{i}{16}\right) \left((47 + 2i)A + (2 + 23i)B\right) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}a^2d} \\ & \quad - \frac{\left(\frac{1}{16} - \frac{i}{16}\right) \left((47 + 2i)A + (2 + 23i)B\right) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}a^2d} \\ & \quad + \frac{\left((49 + 45i)A - (25 - 21i)B\right) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{32\sqrt{2}a^2d} \\ & \quad - \frac{\left(\frac{1}{32} - \frac{i}{32}\right) \left((2 + 47i)A - (23 + 2i)B\right) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{\sqrt{2}a^2d} \\ & \quad - \frac{7(7A + 3iB)}{24a^2d \tan^{\frac{3}{2}}(c + dx)} + \frac{\sqrt{2}a^2d}{9A + 5iB} \\ & \quad + \frac{5(9iA - 5B)}{8a^2d \sqrt{\tan(c + dx)}} + \frac{8a^2d(1 + i \tan(c + dx)) \tan^{\frac{3}{2}}(c + dx)}{4d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} \end{aligned}$$

$$3.145. \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$$

output $(-1/32+1/32*I)*((47+2*I)*A+(2+23*I)*B)*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/a^2/d*2^{(1/2)}+(-1/32+1/32*I)*((47+2*I)*A+(2+23*I)*B)*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/a^2/d*2^{(1/2)}+1/64*((49+45*I)*A+(-25+21*I)*B)*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/a^2/d*2^{(1/2)}+(-1/64+1/64*I)*((2+47*I)*A-(23+2*I)*B)*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/a^2/d*2^{(1/2)}+5/8*(9*I*A-5*B)/a^2/d/\tan(d*x+c)^{(1/2)}-7/24*(7*A+3*I*B)/a^2/d/\tan(d*x+c)^{(3/2)}+1/8*(9*A+5*I*B)/a^2/d/(1+I*\tan(d*x+c))/\tan(d*x+c)^{(3/2)}+1/4*(A+I*B)/d/\tan(d*x+c)^{(3/2)}/(a+I*a*\tan(d*x+c))^2$

3.145.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.97 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.52

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx$$

$$= \frac{\sec^2(c + dx) (-6 \cos(c + dx)((11A + 7iB) \cos(c + dx) + (9iA - 5B) \sin(c + dx)) + 2(47A + 23iB) \operatorname{Hypergeometric2F1}[-3/2, 1, -1/2, (-I) \tan(c + dx)] * (\cos[2*(c + dx)] + I \sin[2*(c + dx)]) + 4*(A - I*B) * \operatorname{Hypergeometric2F1}[-3/2, 1, -1/2, I \tan(c + dx)] * (\cos[2*(c + dx)] + I \sin[2*(c + dx)])}}{(48*a^2*d*\tan(c + dx)^{(3/2)}*(-I + \tan(c + dx))^2)}$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^2),x]`

output $(\operatorname{Sec}[c + d*x]^2*(-6*\operatorname{Cos}[c + d*x]*((11*A + (7*I)*B)*\operatorname{Cos}[c + d*x] + ((9*I)*A - 5*B)*\operatorname{Sin}[c + d*x])) + 2*(47*A + (23*I)*B)*\operatorname{Hypergeometric2F1}[-3/2, 1, -1/2, (-I)*\operatorname{Tan}[c + d*x]]*(\operatorname{Cos}[2*(c + d*x)] + I*\operatorname{Sin}[2*(c + d*x)]) + 4*(A - I*B)*\operatorname{Hypergeometric2F1}[-3/2, 1, -1/2, I*\operatorname{Tan}[c + d*x]]*(\operatorname{Cos}[2*(c + d*x)] + I*\operatorname{Sin}[2*(c + d*x)])))/(48*a^2*d*\operatorname{Tan}[c + d*x]^(3/2)*(-I + \operatorname{Tan}[c + d*x])^2)$

3.145.3 Rubi [A] (verified)

Time = 1.25 (sec) , antiderivative size = 315, normalized size of antiderivative = 0.91, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3042, 4079, 27, 3042, 4079, 3042, 4012, 25, 3042, 4012, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.145. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
& \int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2}(a + ia \tan(c + dx))^2} dx \\
& \quad \downarrow \text{4079} \\
& \frac{\int \frac{a(11A+3iB)-7a(iA-B) \tan(c+dx)}{2 \tan^{\frac{5}{2}}(c+dx)(i \tan(c+dx)a+a)} dx}{4a^2} + \frac{A + iB}{4d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{a(11A+3iB)-7a(iA-B) \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(i \tan(c+dx)a+a)} dx}{8a^2} + \frac{A + iB}{4d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(11A+3iB)-7a(iA-B) \tan(c+dx)}{\tan(c+dx)^{5/2}(i \tan(c+dx)a+a)} dx}{8a^2} + \frac{A + iB}{4d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} \\
& \quad \downarrow \text{4079} \\
& \frac{\int \frac{7a^2(7A+3iB)-5a^2(9iA-5B) \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)} dx}{2a^2} + \frac{9A+5iB}{d(1+i \tan(c+dx)) \tan^{\frac{3}{2}}(c+dx)} + \\
& \quad \frac{8a^2}{A + iB} \\
& \quad \frac{A + iB}{4d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{7a^2(7A+3iB)-5a^2(9iA-5B) \tan(c+dx)}{\tan(c+dx)^{5/2}} dx}{2a^2} + \frac{9A+5iB}{d(1+i \tan(c+dx)) \tan^{\frac{3}{2}}(c+dx)} + \\
& \quad \frac{8a^2}{A + iB} \\
& \quad \frac{A + iB}{4d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} \\
& \quad \downarrow \text{4012} \\
& \frac{\int -\frac{5(9iA-5B)a^2+7(7A+3iB) \tan(c+dx)a^2}{\tan^{\frac{3}{2}}(c+dx)} dx - \frac{14a^2(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)}}{2a^2} + \frac{9A+5iB}{d(1+i \tan(c+dx)) \tan^{\frac{3}{2}}(c+dx)} + \\
& \quad \frac{8a^2}{A + iB} \\
& \quad \frac{A + iB}{4d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} \\
& \quad \downarrow \text{25}
\end{aligned}$$

3.145. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
 & \frac{-\int \frac{5(9iA-5B)a^2+7(7A+3iB)\tan(c+dx)a^2}{\tan^{\frac{3}{2}}(c+dx)} dx - \frac{14a^2(7A+3iB)}{3d\tan^{\frac{3}{2}}(c+dx)}}{2a^2} + \frac{9A+5iB}{d(1+i\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)} + \\
 & \frac{8a^2}{A+iB} \\
 & \frac{4d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^2}{} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\int \frac{5(9iA-5B)a^2+7(7A+3iB)\tan(c+dx)a^2}{\tan(c+dx)^{3/2}} dx - \frac{14a^2(7A+3iB)}{3d\tan^{\frac{3}{2}}(c+dx)}}{2a^2} + \frac{9A+5iB}{d(1+i\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)} + \\
 & \frac{8a^2}{A+iB} \\
 & \frac{4d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^2}{} \\
 & \quad \downarrow \text{4012} \\
 & \frac{-\int \frac{7a^2(7A+3iB)-5a^2(9iA-5B)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{14a^2(7A+3iB)}{3d\tan^{\frac{3}{2}}(c+dx)} + \frac{10a^2(-5B+9iA)}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{9A+5iB}{d(1+i\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)} + \\
 & \frac{8a^2}{A+iB} \\
 & \frac{4d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^2}{} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\int \frac{7a^2(7A+3iB)-5a^2(9iA-5B)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{14a^2(7A+3iB)}{3d\tan^{\frac{3}{2}}(c+dx)} + \frac{10a^2(-5B+9iA)}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{9A+5iB}{d(1+i\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)} + \\
 & \frac{8a^2}{A+iB} \\
 & \frac{4d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^2}{} \\
 & \quad \downarrow \text{4017} \\
 & \frac{2\int \frac{a^2(7(7A+3iB)-5(9iA-5B)\tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{14a^2(7A+3iB)}{3d\tan^{\frac{3}{2}}(c+dx)} + \frac{10a^2(-5B+9iA)}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{9A+5iB}{d(1+i\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)} + \\
 & \frac{8a^2}{A+iB} \\
 & \frac{4d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^2}{} \\
 & \quad \downarrow \text{27} \\
 & \frac{2a^2\int \frac{7(7A+3iB)-5(9iA-5B)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{14a^2(7A+3iB)}{3d\tan^{\frac{3}{2}}(c+dx)} + \frac{10a^2(-5B+9iA)}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{9A+5iB}{d(1+i\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)} + \\
 & \frac{8a^2}{A+iB} \\
 & \frac{4d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^2}{} \\
 & \quad \downarrow \text{1482}
 \end{aligned}$$

3.145. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^2} dx$

$$\frac{2a^2 \left(\frac{1}{2} ((49+45i)A - (25-21i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \left(\frac{1}{2} - \frac{i}{2}\right) ((47+2i)A + (2+23i)B) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} - \frac{14a^2(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{10a^2(-5B)}{d\sqrt{\tan(c+dx)}}$$

$$\frac{A+iB}{4d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} \quad 8a^2$$

↓ 1476

$$\frac{2a^2 \left(\frac{1}{2} ((49+45i)A - (25-21i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \left(\frac{1}{2} - \frac{i}{2}\right) ((47+2i)A + (2+23i)B) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d} - \frac{14a^2(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{10a^2(-5B)}{d\sqrt{\tan(c+dx)}}$$

$$\frac{A+iB}{4d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} \quad 8a^2$$

↓ 1082

$$\frac{2a^2 \left(\frac{1}{2} ((49+45i)A - (25-21i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \left(\frac{1}{2} - \frac{i}{2}\right) ((47+2i)A + (2+23i)B) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(1+\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d} - \frac{14a^2(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{10a^2(-5B)}{d\sqrt{\tan(c+dx)}}$$

$$\frac{A+iB}{4d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} \quad 8a^2$$

↓ 217

$$\frac{2a^2 \left(\frac{1}{2} ((49+45i)A - (25-21i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \left(\frac{1}{2} - \frac{i}{2}\right) ((47+2i)A + (2+23i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d} - \frac{14a^2(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{10a^2(-5B)}{d\sqrt{\tan(c+dx)}}$$

$$\frac{A+iB}{4d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} \quad 8a^2$$

↓ 1479

$$\frac{2a^2 \left(\frac{1}{2} ((49+45i)A - (25-21i)B) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \left(\frac{1}{2} - \frac{i}{2}\right) ((47+2i)A + (2+23i)B) \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d} - \frac{14a^2(7A+3iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{10a^2(-5B)}{d\sqrt{\tan(c+dx)}}$$

$$\frac{A+iB}{4d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} \quad 8a^2$$

3.145. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

↓ 25

$$\frac{2a^2 \left(\frac{1}{2}((49+45i)A - (25-21i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \left(\frac{1}{2} - \frac{i}{2}\right)((47+2i)A + (2+23i)B) \right)}{d}}{2a^2}$$

8a²

$$\frac{A + iB}{4d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2}$$

↓ 27

$$\frac{2a^2 \left(\frac{1}{2}((49+45i)A - (25-21i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \left(\frac{1}{2} - \frac{i}{2}\right)((47+2i)A + (2+23i)B) \right)}{d}}{2a^2}$$

8a²

$$\frac{A + iB}{4d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2}$$

↓ 1103

$$\frac{2a^2 \left(\left(\frac{1}{2} - \frac{i}{2}\right)((47+2i)A + (2+23i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}((49+45i)A - (25-21i)B) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)})}{2\sqrt{2}} \right) \right)}{d}}{2a^2}$$

8a²

$$\frac{A + iB}{4d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^2),x]`

output `(((-2*a^2*((1/2 - I/2)*((47 + 2*I)*A + (2 + 23*I)*B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])) + (((49 + 45*I)*A - (25 - 21*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2))/d - (14*a^2*(7*A + (3*I)*B))/(3*d*Tan[c + d*x]^(3/2)) + (10*a^2*((9*I)*A - 5*B))/(d*Sqrt[Tan[c + d*x]])/(2*a^2) + (9*A + (5*I)*B)/(d*(1 + I*Tan[c + d*x])*Tan[c + d*x]^(3/2))/(8*a^2) + (A + I*B)/(4*d*Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^2)`

3.145. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

3.145.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4017 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.145.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.51

method	result
derivativedivides	$i \frac{\left(\frac{(-13A - 9iB)}{2} \left(\tan^{\frac{3}{2}}(dx+c) \right) + \frac{(15iA - 11B)}{2} (\sqrt{\tan(dx+c)}) - (23iB+47A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right) \right)}{(\tan(dx+c)-i)^2} - \frac{2A}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}$
default	$i \frac{\left(\frac{(-13A - 9iB)}{2} \left(\tan^{\frac{3}{2}}(dx+c) \right) + \frac{(15iA - 11B)}{2} (\sqrt{\tan(dx+c)}) - (23iB+47A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right) \right)}{(\tan(dx+c)-i)^2} - \frac{2A}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}$

3.145. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d/a^2*(-1/4*I*((-13/2*A-9/2*I*B)*tan(d*x+c)^(3/2)+(15/2*I*A-11/2*B)*tan(d*x+c)^(1/2))/(tan(d*x+c)-I)^2-(47*A+23*I*B)/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))-2/3*A/tan(d*x+c)^(3/2)-2*(-2*I*A+B)/tan(d*x+c)^(1/2)-1/2*I*(A-I*B)/(2^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))`

3.145.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 861 vs. $2(258) = 516$.

Time = 0.29 (sec) , antiderivative size = 861, normalized size of antiderivative = 2.48

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="fracas")`

output

```

1/96*(6*(a^2*d*e^(8*I*d*x + 8*I*c) - 2*a^2*d*e^(6*I*d*x + 6*I*c) + a^2*d*e
^(4*I*d*x + 4*I*c))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2))*log(-2*((I*a^
2*d*e^(2*I*d*x + 2*I*c) + I*a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2
*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2)) - (A - I*B)
*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 6*(a^2*d*e^(8*I*d*
x + 8*I*c) - 2*a^2*d*e^(6*I*d*x + 6*I*c) + a^2*d*e^(4*I*d*x + 4*I*c))*sqrt
((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2))*log(-2*((-I*a^2*d*e^(2*I*d*x + 2*I*c)
- I*a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*s
qrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e
^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*(a^2*d*e^(8*I*d*x + 8*I*c) - 2*a^2*d*e^
(6*I*d*x + 6*I*c) + a^2*d*e^(4*I*d*x + 4*I*c))*sqrt((2209*I*A^2 - 2162*A*B
- 529*I*B^2)/(a^4*d^2))*log(-1/8*((a^2*d*e^(2*I*d*x + 2*I*c) + a^2*d)*sqr
t((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((2209*I*A^2
- 2162*A*B - 529*I*B^2)/(a^4*d^2)) + 47*I*A - 23*B)*e^(-2*I*d*x - 2*I*c)/
(a^2*d)) + 3*(a^2*d*e^(8*I*d*x + 8*I*c) - 2*a^2*d*e^(6*I*d*x + 6*I*c) + a^
2*d*e^(4*I*d*x + 4*I*c))*sqrt((2209*I*A^2 - 2162*A*B - 529*I*B^2)/(a^4*d^2
))*log(1/8*((a^2*d*e^(2*I*d*x + 2*I*c) + a^2*d)*sqrt((-I*e^(2*I*d*x + 2*I*
c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((2209*I*A^2 - 2162*A*B - 529*I*B^2
)/(a^4*d^2)) - 47*I*A + 23*B)*e^(-2*I*d*x - 2*I*c)/(a^2*d)) - 2*(2*(101*A
+ 63*I*B)*e^(8*I*d*x + 8*I*c) - (103*A + 27*I*B)*e^(6*I*d*x + 6*I*c) - ...

```

3.145.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(5/2)/(a+I*a*tan(d*x+c))**2,x)`

output `Timed out`

3.145.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.145.8 Giac [A] (verification not implemented)

Time = 1.49 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.47

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx \\ &= -\frac{(i + 1) \sqrt{2}(-47i A + 23 B) \arctan\left(\left(\frac{1}{2}i + \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{16 a^2 d} \\ &+ \frac{(i - 1) \sqrt{2}(i A + B) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{8 a^2 d} \\ &- \frac{2(-6i A \tan(dx + c) + 3 B \tan(dx + c) + A)}{3 a^2 d \tan(dx + c)^{\frac{3}{2}}} \\ &- \frac{-13i A \tan(dx + c)^{\frac{3}{2}} + 9 B \tan(dx + c)^{\frac{3}{2}} - 15 A \sqrt{\tan(dx + c)} - 11i B \sqrt{\tan(dx + c)}}{8 a^2 d (\tan(dx + c) - i)^2} \end{aligned}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")`

output `-(1/16*I + 1/16)*sqrt(2)*(-47*I*A + 23*B)*arctan((1/2*I + 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^2*d) + (1/8*I - 1/8)*sqrt(2)*(I*A + B)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^2*d) - 2/3*(-6*I*A*tan(d*x + c) + 3*B*tan(d*x + c) + A)/(a^2*d*tan(d*x + c)^(3/2)) - 1/8*(-13*I*A*tan(d*x + c)^(3/2) + 9*B*tan(d*x + c)^(3/2) - 15*A*sqrt(tan(d*x + c)) - 11*I*B*sqrt(tan(d*x + c)))/(a^2*d*(tan(d*x + c) - I)^2)`

3.145. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

3.145.9 Mupad [B] (verification not implemented)

Time = 12.22 (sec) , antiderivative size = 373, normalized size of antiderivative = 1.07

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx \\
 &= -\operatorname{atan}\left(\frac{8a^2 d \sqrt{\tan(c + dx)} \sqrt{-\frac{A^2 1i}{64a^4 d^2}}}{A}\right) \sqrt{-\frac{A^2 1i}{64a^4 d^2}} 2i \\
 &+ \operatorname{atan}\left(\frac{16a^2 d \sqrt{\tan(c + dx)} \sqrt{\frac{A^2 2209i}{256a^4 d^2}}}{47A}\right) \sqrt{\frac{A^2 2209i}{256a^4 d^2}} 2i \\
 &+ 2 \operatorname{atanh}\left(\frac{8a^2 d \sqrt{\tan(c + dx)} \sqrt{\frac{B^2 1i}{64a^4 d^2}}}{B}\right) \sqrt{\frac{B^2 1i}{64a^4 d^2}} \\
 &+ 2 \operatorname{atanh}\left(\frac{16a^2 d \sqrt{\tan(c + dx)} \sqrt{-\frac{B^2 529i}{256a^4 d^2}}}{23B}\right) \sqrt{-\frac{B^2 529i}{256a^4 d^2}} \\
 &+ \frac{\frac{8A \tan(c+dx)}{3a^2 d} - \frac{45A \tan(c+dx)^3}{8a^2 d} + \frac{A 2i}{3a^2 d} + \frac{A \tan(c+dx)^2 221i}{24a^2 d}}{2 \tan(c + dx)^{5/2} - \tan(c + dx)^{3/2} 1i + \tan(c + dx)^{7/2} 1i} \\
 &- \frac{\frac{43B \tan(c+dx)}{8a^2 d} - \frac{B 2i}{a^2 d} + \frac{B \tan(c+dx)^2 25i}{8a^2 d}}{2 \tan(c + dx)^{3/2} - \sqrt{\tan(c + dx)} 1i + \tan(c + dx)^{5/2} 1i}
 \end{aligned}$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + a*tan(c + d*x)*1i)^2),x)`

output `atan((16*a^2*d*tan(c + d*x)^(1/2)*((A^2*2209i)/(256*a^4*d^2))^(1/2))/(47*A))*((A^2*2209i)/(256*a^4*d^2))^(1/2)*2i - atan((8*a^2*d*tan(c + d*x)^(1/2))*((-A^2*1i)/(64*a^4*d^2))^(1/2))/A*(-(A^2*1i)/(64*a^4*d^2))^(1/2)*2i + 2*atanh((8*a^2*d*tan(c + d*x)^(1/2)*((B^2*1i)/(64*a^4*d^2))^(1/2))/B)*((B^2*1i)/(64*a^4*d^2))^(1/2) + 2*atanh((16*a^2*d*tan(c + d*x)^(1/2)*((-B^2*529i)/(256*a^4*d^2))^(1/2))/(23*B))*((-B^2*529i)/(256*a^4*d^2))^(1/2) + ((A*2i)/(3*a^2*d) + (8*A*tan(c + d*x))/(3*a^2*d) + (A*tan(c + d*x)^2*221i)/(24*a^2*d) - (45*A*tan(c + d*x)^3)/(8*a^2*d))/(2*tan(c + d*x)^(5/2) - tan(c + d*x)^(3/2)*1i + tan(c + d*x)^(7/2)*1i) - ((43*B*tan(c + d*x))/(8*a^2*d) - (B*2i)/(a^2*d) + (B*tan(c + d*x)^2*25i)/(8*a^2*d))/(2*tan(c + d*x)^(3/2) - tan(c + d*x)^(1/2)*1i + tan(c + d*x)^(5/2)*1i)`

3.146
$$\int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.146.1 Optimal result 1515
 3.146.2 Mathematica [A] (verified) 1516
 3.146.3 Rubi [A] (verified) 1517
 3.146.4 Maple [A] (verified) 1523
 3.146.5 Fricas [B] (verification not implemented) 1524
 3.146.6 Sympy [F(-1)] 1525
 3.146.7 Maxima [F(-2)] 1525
 3.146.8 Giac [A] (verification not implemented) 1525
 3.146.9 Mupad [B] (verification not implemented) 1526

3.146.1 Optimal result

Integrand size = 36, antiderivative size = 393

$$\begin{aligned} & \int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\ &= \frac{\left(\frac{1}{16} + \frac{i}{16}\right) \left((29+i)A + (1+76i)B\right) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}a^3d} \\ & \quad - \frac{\left(\frac{1}{16} + \frac{i}{16}\right) \left((29+i)A + (1+76i)B\right) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}a^3d} \\ & \quad - \frac{\left((28-30i)A + (75+77i)B\right) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{32\sqrt{2}a^3d} \\ & \quad - \frac{\left(\frac{1}{32} + \frac{i}{32}\right) \left((1+29i)A - (76+i)B\right) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{\sqrt{2}a^3d} \\ & \quad + \frac{15(2iA - 5B)\sqrt{\tan(c+dx)}}{8a^3d} + \frac{7(4A + 11iB)\tan^{\frac{3}{2}}(c+dx)}{24a^3d} \\ & \quad + \frac{(iA - B)\tan^{\frac{9}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} + \frac{(A + 2iB)\tan^{\frac{7}{2}}(c+dx)}{4ad(a+ia \tan(c+dx))^2} - \frac{3(2iA - 5B)\tan^{\frac{5}{2}}(c+dx)}{8d(a^3 + ia^3 \tan(c+dx))} \end{aligned}$$

output $(-1/32-1/32*I)*((29+I)*A+(1+76*I)*B)*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/a^{3/d*2^{(1/2)}}-(1/32+1/32*I)*((29+I)*A+(1+76*I)*B)*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/a^{3/d*2^{(1/2)}}-1/64*((28-30*I)*A+(75+77*I)*B)*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/a^{3/d*2^{(1/2)}}-(1/64+1/64*I)*((1+29*I)*A-(76+I)*B)*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/a^{3/d*2^{(1/2)}}+15/8*(2*I*A-5*B)*\tan(d*x+c)^{(1/2)}/a^{3/d+7/24*(4*A+11*I*B)*\tan(d*x+c)^{(3/2)}/a^{3/d+1/6*(I*A-B)*\tan(d*x+c)^{(9/2)}/d/(a+I*a*\tan(d*x+c))^{3+1/4*(A+2*I*B)*\tan(d*x+c)^{(7/2)}/a/d/(a+I*a*\tan(d*x+c))^{2-3/8*(2*I*A-5*B)*\tan(d*x+c)^{(5/2)}/d/(a^{3+I*a^{3*\tan(d*x+c))}}$

3.146.2 Mathematica [A] (verified)

Time = 6.17 (sec) , antiderivative size = 235, normalized size of antiderivative = 0.60

$$\int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{-3\sqrt[4]{-1}(A-iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right) \sec^3(c+dx)(\cos(3(c+dx))+i \sin(3(c+dx))) - 3\sqrt[4]{-1}}$$

input `Integrate[(Tan[c + d*x]^(9/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output $(-3*(-1)^{(1/4)}*(A - I*B)*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]]*\text{Sec}[c + d*x]^3*(\text{Cos}[3*(c + d*x)] + I*\text{Sin}[3*(c + d*x)]) - 3*(-1)^{(1/4)}*(29*A + (76*I)*B)*\text{ArcTanh}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]]*\text{Sec}[c + d*x]^3*(\text{Cos}[3*(c + d*x)] + I*\text{Sin}[3*(c + d*x)]) + \text{Sqrt}[\text{Tan}[c + d*x]]*(-45*(2*A + (5*I)*B) + ((-242*I)*A + 598*B)*\text{Tan}[c + d*x] + 3*(68*A + (163*I)*B)*\text{Tan}[c + d*x]^2 + (48*I)*(A + (2*I)*B)*\text{Tan}[c + d*x]^3 + (16*I)*B*\text{Tan}[c + d*x]^4)/(24*a^3*d*(-I + \text{Tan}[c + d*x])^3)$

3.146.3 Rubi [A] (verified)

Time = 1.67 (sec) , antiderivative size = 372, normalized size of antiderivative = 0.95, number of steps used = 25, number of rules used = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4078, 27, 3042, 4078, 27, 3042, 4078, 3042, 4011, 3042, 4011, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^{9/2}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+iA) \tan^{\frac{9}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{3 \tan^{\frac{7}{2}}(c+dx)(3a(iA-B)+a(A+5iB) \tan(c+dx))}{2(i \tan(c+dx)a+a)^2} dx}{6a^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B+iA) \tan^{\frac{9}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{\tan^{\frac{7}{2}}(c+dx)(3a(iA-B)+a(A+5iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{4a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B+iA) \tan^{\frac{9}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{\tan(c+dx)^{7/2}(3a(iA-B)+a(A+5iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{4a^2} \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+iA) \tan^{\frac{9}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int -\frac{2 \tan^{\frac{5}{2}}(c+dx)(7a^2(A+2iB)-a^2(5iA-16B) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{4a^2} - \frac{a(A+2iB) \tan^{\frac{7}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B+iA) \tan^{\frac{9}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{\tan^{\frac{5}{2}}(c+dx)(7a^2(A+2iB)-a^2(5iA-16B) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{2a^2} - \frac{a(A+2iB) \tan^{\frac{7}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.146. $\int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{(-B + iA) \tan^{\frac{9}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{\tan(c+dx)^{5/2} (7a^2(A+2iB) - a^2(5iA-16B) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{2a^2} - \frac{a(A+2iB) \tan^{\frac{7}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 4078 \\
 & \frac{(-B + iA) \tan^{\frac{9}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3a^2(-5B+2iA) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{\int \tan^{\frac{3}{2}}(c+dx) (15(2iA-5B)a^3+7(4A+11iB) \tan(c+dx)a^3) dx}{2a^2} - \frac{a(A+2iB) \tan^{\frac{7}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{(-B + iA) \tan^{\frac{9}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3a^2(-5B+2iA) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{\int \tan(c+dx)^{3/2} (15(2iA-5B)a^3+7(4A+11iB) \tan(c+dx)a^3) dx}{2a^2} - \frac{a(A+2iB) \tan^{\frac{7}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 4011 \\
 & \frac{(-B + iA) \tan^{\frac{9}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3a^2(-5B+2iA) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{\int \sqrt{\tan(c+dx)} (15a^3(2iA-5B) \tan(c+dx) - 7a^3(4A+11iB)) dx + \frac{14a^3(4A+11iB) \tan^{\frac{3}{2}}(c+dx)}{3d}}{2a^2} - \frac{a(A+2iB) \tan^{\frac{7}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \frac{(-B + iA) \tan^{\frac{9}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3a^2(-5B+2iA) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{\int \sqrt{\tan(c+dx)} (15a^3(2iA-5B) \tan(c+dx) - 7a^3(4A+11iB)) dx + \frac{14a^3(4A+11iB) \tan^{\frac{3}{2}}(c+dx)}{3d}}{2a^2} - \frac{a(A+2iB) \tan^{\frac{7}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 4011 \\
 & \frac{(-B + iA) \tan^{\frac{9}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3a^2(-5B+2iA) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{\int \frac{-15(2iA-5B)a^3-7(4A+11iB) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx + \frac{14a^3(4A+11iB) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{30a^3(-5B+2iA)\sqrt{\tan(c+dx)}}{d}}{2a^2} - \frac{a(A+2iB) \tan^{\frac{7}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \qquad \qquad \qquad \downarrow 3042
 \end{aligned}$$

3.146. $\int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\frac{(-B + iA) \tan^{\frac{9}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3a^2(-5B+2iA) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{\int \frac{-15(2iA-5B)a^3-7(4A+11iB) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx + \frac{14a^3(4A+11iB) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{30a^3(-5B+2iA)\sqrt{\tan(c+dx)}}{d}}{2a^2} - \frac{a(A+2iB)}{d(a+ia \tan(c+dx))}$$

↓ 4017

$$\frac{(-B + iA) \tan^{\frac{9}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3a^2(-5B+2iA) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{2 \int \frac{a^3(15(2iA-5B)+7(4A+11iB) \tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{14a^3(4A+11iB) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{30a^3(-5B+2iA)\sqrt{\tan(c+dx)}}{d}}{2a^2} - \frac{a(A+2iB)}{d(a+ia \tan(c+dx))}$$

↓ 25

$$\frac{(-B + iA) \tan^{\frac{9}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3a^2(-5B+2iA) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{2 \int \frac{a^3(15(2iA-5B)+7(4A+11iB) \tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{14a^3(4A+11iB) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{30a^3(-5B+2iA)\sqrt{\tan(c+dx)}}{d}}{2a^2} - \frac{a(A+2iB)}{d(a+ia \tan(c+dx))}$$

↓ 27

$$\frac{(-B + iA) \tan^{\frac{9}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3a^2(-5B+2iA) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{2a^3 \int \frac{15(2iA-5B)+7(4A+11iB) \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{14a^3(4A+11iB) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{30a^3(-5B+2iA)\sqrt{\tan(c+dx)}}{d}}{2a^2} - \frac{a(A+2iB)}{d(a+ia \tan(c+dx))}$$

↓ 1482

$$\frac{(-B + iA) \tan^{\frac{9}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3a^2(-5B+2iA) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((1+29i)A - (76+i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \left(\frac{1}{2} + \frac{i}{2} \right) ((29+i)A + (1+76i)B) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{2a^2} - \frac{a(A+2iB)}{d(a+ia \tan(c+dx))}$$

↓ 1476

$$\frac{(-B + iA) \tan^{\frac{9}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3a^2(-5B+2iA) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((1+29i)A - (76+i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \left(\frac{1}{2} + \frac{i}{2} \right) ((29+i)A + (1+76i)B) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}} \right) \right)}{2a^2} - \frac{a(A+2iB)}{d(a+ia \tan(c+dx))}$$

3.146. $\int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned} & \downarrow 1103 \\ & \frac{(-B + iA) \tan^{\frac{9}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\ & \frac{3a^2(-5B+2iA) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((29+i)A + (1+76i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \left(\frac{1}{2} + \frac{i}{2} \right) ((1+29i)A - (76+i)B) \right)}{d} \\ & \hline & 2a^2 \end{aligned}$$

input `Int[(Tan[c + d*x]^(9/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `((I*A - B)*Tan[c + d*x]^(9/2))/(6*d*(a + I*a*Tan[c + d*x])^3) - (-((a*(A + (2*I)*B)*Tan[c + d*x]^(7/2))/(d*(a + I*a*Tan[c + d*x])^2)) + ((3*a^2*((2*I)*A - 5*B)*Tan[c + d*x]^(5/2))/(d*(a + I*a*Tan[c + d*x])) - ((-2*a^3*((1/2 + I/2)*((29 + I)*A + (1 + 76*I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + (1/2 + I/2)*((1 + 29*I)*A - (76 + I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]))))/d + (30*a^3*((2*I)*A - 5*B)*Sqrt[Tan[c + d*x]]/d + (14*a^3*(4*A + (11*I)*B)*Tan[c + d*x]^(3/2))/(3*d)/(2*a^2))/(2*a^2)/(4*a^2)`

3.146.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

$$3.146. \int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

```
rule 4078 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.146.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 204, normalized size of antiderivative = 0.52

method	result
derivativedivides	$\frac{2iB \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 6B(\sqrt{\tan(dx+c)}) + 2iA(\sqrt{\tan(dx+c)}) + \frac{\left(\frac{-5i(7iB+4A) \left(\tan^{\frac{5}{2}}(dx+c) \right) + \left(-\frac{182iB}{3} - \frac{98A}{3} \right) \left(\tan^{\frac{3}{2}}(dx+c) \right)}{(\tan(dx+c)-i)^3} \right)}{da^3}$
default	$\frac{2iB \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - 6B(\sqrt{\tan(dx+c)}) + 2iA(\sqrt{\tan(dx+c)}) + \frac{\left(\frac{-5i(7iB+4A) \left(\tan^{\frac{5}{2}}(dx+c) \right) + \left(-\frac{182iB}{3} - \frac{98A}{3} \right) \left(\tan^{\frac{3}{2}}(dx+c) \right)}{(\tan(dx+c)-i)^3} \right)}{da^3}$

```
input int(tan(d*x+c)^(9/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETUR
NVERBOSE)
```

```
output 1/d/a^3*(2/3*I*B*tan(d*x+c)^(3/2)-6*B*tan(d*x+c)^(1/2)+2*I*A*tan(d*x+c)^(1
/2)+1/8*I*((-5*I*(7*I*B+4*A)*tan(d*x+c)^(5/2)+(-182/3*I*B-98/3*A)*tan(d*x+
c)^(3/2)+(-27*B+14*I*A)*tan(d*x+c)^(1/2))/(tan(d*x+c)-I)^3+2*(-76*B+29*I*A
)/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2))))+4*(1
/16*A-1/16*I*B)/(2^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2
^(1/2))))
```

$$3.146. \int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.146.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 779 vs. $2(294) = 588$.

Time = 0.30 (sec) , antiderivative size = 779, normalized size of antiderivative = 1.98

$$\int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx =$$

$$3 \left(a^3 de^{(8i dx+8i c)} + a^3 de^{(6i dx+6i c)} \right) \sqrt{\frac{i A^2+2 AB-i B^2}{a^6 d^2}} \log \left(\frac{2 \left((a^3 de^{(2i dx+2i c)}+a^3 d) \sqrt{\frac{-i e^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}+1}} \sqrt{\frac{i A^2+2 AB-i B^2}{a^6 d^2}} + \dots \right)}{i A+B} \right)$$

```
input integrate(tan(d*x+c)^(9/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="fricas")
```

```
output -1/96*(3*(a^3*d*e^(8*I*d*x + 8*I*c) + a^3*d*e^(6*I*d*x + 6*I*c))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2))*log(2*((a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*(a^3*d*e^(8*I*d*x + 8*I*c) + a^3*d*e^(6*I*d*x + 6*I*c))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2))*log(-2*((a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*(a^3*d*e^(8*I*d*x + 8*I*c) + a^3*d*e^(6*I*d*x + 6*I*c))*sqrt((-841*I*A^2 + 4408*A*B + 5776*I*B^2)/(a^6*d^2))*log(1/8*((a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-841*I*A^2 + 4408*A*B + 5776*I*B^2)/(a^6*d^2)) + 29*A + 76*I*B)*e^(-2*I*d*x - 2*I*c)/(a^3*d)) + 3*(a^3*d*e^(8*I*d*x + 8*I*c) + a^3*d*e^(6*I*d*x + 6*I*c))*sqrt((-841*I*A^2 + 4408*A*B + 5776*I*B^2)/(a^6*d^2))*log(-1/8*((a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-841*I*A^2 + 4408*A*B + 5776*I*B^2)/(a^6*d^2)) - 29*A - 76*I*B)*e^(-2*I*d*x - 2*I*c)/(a^3*d)) + 2*(2*(-73*I*A + 174*B)*e^(8*I*d*x + 8*I*c) - (187*I*A - 492*B)*e^(6*I*d*x + 6*I*c) + 3*(-11*I*A + 23*B)*e^(4*I*d*x + 4*I*c) - (-7*I*A + 10*B)*e^(2*I*d*x + 2*I*c) - I*A + B)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2...
```

$$3.146. \quad \int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.146.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(9/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)`

output `Timed out`

3.146.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(9/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.146.8 Giac [A] (verification not implemented)

Time = 1.02 (sec) , antiderivative size = 208, normalized size of antiderivative = 0.53

$$\begin{aligned} & \int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx \\ &= -\frac{(i+1)\sqrt{2}(29A+76iB)\arctan\left(\left(\frac{1}{2}i+\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{16a^3d} \\ & \quad -\frac{(i-1)\sqrt{2}(A-iB)\arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{16a^3d} \\ & \quad +\frac{60A\tan(dx+c)^{\frac{5}{2}}+105iB\tan(dx+c)^{\frac{5}{2}}-98iA\tan(dx+c)^{\frac{3}{2}}+182B\tan(dx+c)^{\frac{3}{2}}-42A\sqrt{\tan(dx+c)}}{24a^3d(\tan(dx+c)-i)^3} \\ & \quad -\frac{2\left(-iBa^6d^2\tan(dx+c)^{\frac{3}{2}}-3iAa^6d^2\sqrt{\tan(dx+c)}+9Ba^6d^2\sqrt{\tan(dx+c)}\right)}{3a^9d^3} \end{aligned}$$

3.146. $\int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

input `integrate(tan(d*x+c)^(9/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output
$$\begin{aligned} & -(1/16*I + 1/16)*\sqrt{2}*(29*A + 76*I*B)*\arctan((1/2*I + 1/2)*\sqrt{2}*\sqrt{\tan(dx + c)})/(a^3*d) - (1/16*I - 1/16)*\sqrt{2}*(A - I*B)*\arctan(-(1/2*I - 1/2)*\sqrt{2}*\sqrt{\tan(dx + c)})/(a^3*d) + 1/24*(60*A*\tan(dx + c)^{(5/2)} + 105*I*B*\tan(dx + c)^{(5/2)} - 98*I*A*\tan(dx + c)^{(3/2)} + 182*B*\tan(dx + c)^{(3/2)} - 42*A*\sqrt{\tan(dx + c)} - 81*I*B*\sqrt{\tan(dx + c)})/(a^3*d*(\tan(dx + c) - I)^3) - 2/3*(-I*B*a^6*d^2*\tan(dx + c)^{(3/2)} - 3*I*A*a^6*d^2*\sqrt{\tan(dx + c)} + 9*B*a^6*d^2*\sqrt{\tan(dx + c)})/(a^9*d^3) \end{aligned}$$

3.146.9 Mupad [B] (verification not implemented)

Time = 11.60 (sec) , antiderivative size = 431, normalized size of antiderivative = 1.10

$$\begin{aligned} & \int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx \\ & = \operatorname{atan}\left(\frac{a^3 d \sqrt{\tan(c+dx)} \sqrt{\frac{A^2 1i}{256 a^6 d^2}} 16i}{A}\right) \sqrt{\frac{A^2 1i}{256 a^6 d^2}} 2i \\ & \quad - \operatorname{atan}\left(\frac{a^3 d \sqrt{\tan(c+dx)} \sqrt{-\frac{A^2 841i}{256 a^6 d^2}} 16i}{29 A}\right) \sqrt{-\frac{A^2 841i}{256 a^6 d^2}} 2i \\ & \quad - \operatorname{atan}\left(\frac{16 a^3 d \sqrt{\tan(c+dx)} \sqrt{-\frac{B^2 1i}{256 a^6 d^2}}}{B}\right) \sqrt{-\frac{B^2 1i}{256 a^6 d^2}} 2i \\ & \quad - \operatorname{atan}\left(\frac{4 a^3 d \sqrt{\tan(c+dx)} \sqrt{\frac{B^2 361i}{16 a^6 d^2}}}{19 B}\right) \sqrt{\frac{B^2 361i}{16 a^6 d^2}} 2i \\ & \quad - \frac{\frac{49 A \tan(c+dx)^{3/2}}{12 a^3 d} - \frac{A \sqrt{\tan(c+dx)} 7i}{4 a^3 d} + \frac{A \tan(c+dx)^{5/2} 5i}{2 a^3 d}}{-\tan(c+dx)^3 1i - 3 \tan(c+dx)^2 + \tan(c+dx) 3i + 1} \\ & \quad - \frac{\frac{27 B \sqrt{\tan(c+dx)}}{8 a^3 d} - \frac{35 B \tan(c+dx)^{5/2}}{8 a^3 d} + \frac{B \tan(c+dx)^{3/2} 9i}{12 a^3 d}}{-\tan(c+dx)^3 1i - 3 \tan(c+dx)^2 + \tan(c+dx) 3i + 1} \\ & \quad + \frac{A \sqrt{\tan(c+dx)} 2i}{a^3 d} - \frac{6 B \sqrt{\tan(c+dx)}}{a^3 d} + \frac{B \tan(c+dx)^{3/2} 2i}{3 a^3 d} \end{aligned}$$

input `int((tan(c + d*x)^(9/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3,x)`

3.146.
$$\int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

output

$$\begin{aligned} & \operatorname{atan}\left(\frac{a^3 d \tan(c + dx)^{1/2} \left(\frac{A^2 i}{256 a^6 d^2}\right)^{1/2} 16i}{A}\right) \left(\frac{A^2 i}{256 a^6 d^2}\right)^{1/2} 2i - \operatorname{atan}\left(\frac{a^3 d \tan(c + dx)^{1/2} \left(-\frac{A^2 841i}{256 a^6 d^2}\right)^{1/2} 16i}{29A}\right) \left(-\frac{A^2 841i}{256 a^6 d^2}\right)^{1/2} 2i - \\ & \operatorname{atan}\left(\frac{16 a^3 d \tan(c + dx)^{1/2} \left(-\frac{B^2 i}{256 a^6 d^2}\right)^{1/2}}{B}\right) \left(-\frac{B^2 i}{256 a^6 d^2}\right)^{1/2} 2i - \operatorname{atan}\left(\frac{4 a^3 d \tan(c + dx)^{1/2} \left(\frac{B^2 361i}{16 a^6 d^2}\right)^{1/2}}{19B}\right) \left(\frac{B^2 361i}{16 a^6 d^2}\right)^{1/2} 2i - \\ & \left(\frac{49 A \tan(c + dx)^{3/2}}{12 a^3 d} - \frac{A \tan(c + dx)^{1/2} 7i}{4 a^3 d} + \frac{A \tan(c + dx)^{5/2} 5i}{2 a^3 d}\right) \left(\frac{\tan(c + dx) 3i - 3 \tan(c + dx)^2 - \tan(c + dx) 3i + 1}{\tan(c + dx) 3i - 3 \tan(c + dx)^2 - \tan(c + dx) 3i + 1}\right) - \\ & \left(\frac{27 B \tan(c + dx)^{1/2}}{8 a^3 d} + \frac{B \tan(c + dx)^{3/2} 9i}{12 a^3 d} - \frac{35 B \tan(c + dx)^{5/2}}{8 a^3 d}\right) \left(\frac{\tan(c + dx) 3i - 3 \tan(c + dx)^2 - \tan(c + dx) 3i + 1}{\tan(c + dx) 3i - 3 \tan(c + dx)^2 - \tan(c + dx) 3i + 1}\right) + \\ & \frac{A \tan(c + dx)^{1/2} 2i}{a^3 d} - \frac{6 B \tan(c + dx)^{1/2}}{a^3 d} + \frac{B \tan(c + dx)^{3/2} 2i}{3 a^3 d} \end{aligned}$$

3.146. $\int \frac{\tan^{\frac{9}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$3.147 \quad \int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.147.1 Optimal result	1528
3.147.2 Mathematica [A] (verified)	1529
3.147.3 Rubi [A] (verified)	1529
3.147.4 Maple [A] (verified)	1536
3.147.5 Fricas [B] (verification not implemented)	1537
3.147.6 Sympy [F(-1)]	1538
3.147.7 Maxima [F(-2)]	1538
3.147.8 Giac [A] (verification not implemented)	1538
3.147.9 Mupad [B] (verification not implemented)	1539

3.147.1 Optimal result

Integrand size = 36, antiderivative size = 364

$$\begin{aligned} & \int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\ &= -\frac{\left(\frac{1}{16} + \frac{i}{16}\right) \left((1+6i)A - (29+i)B\right) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}a^3d} \\ & \quad - \frac{\left((5-7i)A + (28+30i)B\right) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{16\sqrt{2}a^3d} \\ & \quad + \frac{\left(\frac{1}{32} + \frac{i}{32}\right) \left((6+i)A + (1+29i)B\right) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{\sqrt{2}a^3d} \\ & \quad - \frac{\left(\frac{1}{32} + \frac{i}{32}\right) \left((6+i)A + (1+29i)B\right) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{\sqrt{2}a^3d} \\ & \quad + \frac{5(A+6iB)\sqrt{\tan(c+dx)}}{8a^3d} + \frac{(iA-B)\tan^{\frac{7}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} \\ & \quad + \frac{(2A+5iB)\tan^{\frac{5}{2}}(c+dx)}{12ad(a+ia \tan(c+dx))^2} - \frac{7(iA-4B)\tan^{\frac{3}{2}}(c+dx)}{24d(a^3+ia^3 \tan(c+dx))} \end{aligned}$$

$$3.147. \quad \int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

output $(1/32+1/32*I)*((1+6*I)*A-(29+I)*B)*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/a^3/d*2^{(1/2)}-1/32*((5-7*I)*A+(28+30*I)*B)*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/a^3/d*2^{(1/2)}+(1/64+1/64*I)*((6+I)*A+(1+29*I)*B)*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/a^3/d*2^{(1/2)}-(1/64+1/64*I)*((6+I)*A+(1+29*I)*B)*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/a^3/d*2^{(1/2)}+5/8*(A+6*I*B)*\tan(d*x+c)^{(1/2)}/a^3/d+1/6*(I*A-B)*\tan(d*x+c)^{(7/2)}/d/(a+I*a*\tan(d*x+c))^3+1/12*(2*A+5*I*B)*\tan(d*x+c)^{(5/2)}/a/d/(a+I*a*\tan(d*x+c))^2-7/24*(I*A-4*B)*\tan(d*x+c)^{(3/2)}/d/(a^3+I*a^3*\tan(d*x+c))$

3.147.2 Mathematica [A] (verified)

Time = 4.60 (sec) , antiderivative size = 222, normalized size of antiderivative = 0.61

$$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{\sec^3(c+dx) \left(-12\sqrt[4]{-1}(iA+B) \arctan \left((-1)^{3/4} \sqrt{\tan(c+dx)} \right) (\cos(3(c+dx)) + i \sin(3(c+dx))) + 12 \right)}{a^3 + i a^3 \tan(c+dx)}$$

input `Integrate[(Tan[c + d*x]^(7/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output $(\text{Sec}[c + d*x]^3*(-12*(-1)^{(1/4)}*(I*A + B)*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]]*(\text{Cos}[3*(c + d*x)] + I*\text{Sin}[3*(c + d*x)]) + 12*(-1)^{(3/4)}*(6*A + (29*I)*B)*\text{ArcTanh}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]*(\text{Cos}[3*(c + d*x)] + I*\text{Sin}[3*(c + d*x)])] + (2*I)*((9*A + (33*I)*B)*\text{Cos}[c + d*x] + 21*(A + (7*I)*B)*\text{Cos}[3*(c + d*x)] + (2*I)*(19*A + (97*I)*B + (19*A + (145*I)*B)*\text{Cos}[2*(c + d*x)])*\text{Sin}[c + d*x])*\text{Sqrt}[\text{Tan}[c + d*x]]))/(96*a^3*d*(-I + \text{Tan}[c + d*x])^3)$

3.147.3 Rubi [A] (verified)

Time = 1.38 (sec) , antiderivative size = 339, normalized size of antiderivative = 0.93, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.639$, Rules used = {3042, 4078, 27, 3042, 4078, 27, 3042, 4078, 27, 3042, 4011, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.147. $\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
 & \int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{\tan(c+dx)^{7/2}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\
 & \quad \downarrow 4078 \\
 & \frac{(-B+iA) \tan^{\frac{7}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{\tan^{\frac{5}{2}}(c+dx)(7a(iA-B)+a(A+13iB) \tan(c+dx))}{2(i \tan(c+dx)a+a)^2} dx}{6a^2} \\
 & \quad \downarrow 27 \\
 & \frac{(-B+iA) \tan^{\frac{7}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{\tan^{\frac{5}{2}}(c+dx)(7a(iA-B)+a(A+13iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{12a^2} \\
 & \quad \downarrow 3042 \\
 & \frac{(-B+iA) \tan^{\frac{7}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{\tan(c+dx)^{5/2}(7a(iA-B)+a(A+13iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{12a^2} \\
 & \quad \downarrow 4078 \\
 & \frac{(-B+iA) \tan^{\frac{7}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int -\frac{2 \tan^{\frac{3}{2}}(c+dx)(5a^2(2A+5iB)-a^2(4iA-31B) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{4a^2} - \frac{a(2A+5iB) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \quad \downarrow 27 \\
 & \frac{(-B+iA) \tan^{\frac{7}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{\tan^{\frac{3}{2}}(c+dx)(5a^2(2A+5iB)-a^2(4iA-31B) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{2a^2} - \frac{a(2A+5iB) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \quad \downarrow 3042 \\
 & \frac{(-B+iA) \tan^{\frac{7}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{\tan(c+dx)^{3/2}(5a^2(2A+5iB)-a^2(4iA-31B) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{2a^2} - \frac{a(2A+5iB) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \quad \downarrow 4078 \\
 & \frac{(-B+iA) \tan^{\frac{7}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\frac{7a^2(-4B+iA) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{\int 3\sqrt{\tan(c+dx)}(7(iA-4B)a^3+5(A+6iB) \tan(c+dx)a^3) dx}{2a^2}}{2a^2} - \frac{a(2A+5iB) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \quad \downarrow 27
 \end{aligned}$$

3.147. $\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
& \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
& \frac{\frac{7a^2(-4B + iA) \tan^{\frac{3}{2}}(c + dx)}{d(a + ia \tan(c + dx))} - \frac{3 \int \sqrt{\tan(c + dx)} (7(iA - 4B)a^3 + 5(A + 6iB) \tan(c + dx)a^3) dx}{2a^2}}{2a^2} - \frac{a(2A + 5iB) \tan^{\frac{5}{2}}(c + dx)}{d(a + ia \tan(c + dx))^2} \\
& \frac{12a^2}{12a^2} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
& \frac{\frac{7a^2(-4B + iA) \tan^{\frac{3}{2}}(c + dx)}{d(a + ia \tan(c + dx))} - \frac{3 \int \sqrt{\tan(c + dx)} (7(iA - 4B)a^3 + 5(A + 6iB) \tan(c + dx)a^3) dx}{2a^2}}{2a^2} - \frac{a(2A + 5iB) \tan^{\frac{5}{2}}(c + dx)}{d(a + ia \tan(c + dx))^2} \\
& \frac{12a^2}{12a^2} \\
& \downarrow 4011 \\
& \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
& \frac{\frac{7a^2(-4B + iA) \tan^{\frac{3}{2}}(c + dx)}{d(a + ia \tan(c + dx))} - 3 \left(\int \frac{7a^3(iA - 4B) \tan(c + dx) - 5a^3(A + 6iB)}{\sqrt{\tan(c + dx)}} dx + \frac{10a^3(A + 6iB) \sqrt{\tan(c + dx)}}{d} \right)}{2a^2} - \frac{a(2A + 5iB) \tan^{\frac{5}{2}}(c + dx)}{d(a + ia \tan(c + dx))^2} \\
& \frac{12a^2}{12a^2} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
& \frac{\frac{7a^2(-4B + iA) \tan^{\frac{3}{2}}(c + dx)}{d(a + ia \tan(c + dx))} - 3 \left(\int \frac{7a^3(iA - 4B) \tan(c + dx) - 5a^3(A + 6iB)}{\sqrt{\tan(c + dx)}} dx + \frac{10a^3(A + 6iB) \sqrt{\tan(c + dx)}}{d} \right)}{2a^2} - \frac{a(2A + 5iB) \tan^{\frac{5}{2}}(c + dx)}{d(a + ia \tan(c + dx))^2} \\
& \frac{12a^2}{12a^2} \\
& \downarrow 4017 \\
& \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
& \frac{\frac{7a^2(-4B + iA) \tan^{\frac{3}{2}}(c + dx)}{d(a + ia \tan(c + dx))} - 3 \left(\frac{2 \int -\frac{a^3(5(A + 6iB) - 7(iA - 4B) \tan(c + dx))}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{d} + \frac{10a^3(A + 6iB) \sqrt{\tan(c + dx)}}{d} \right)}{2a^2} - \frac{a(2A + 5iB) \tan^{\frac{5}{2}}(c + dx)}{d(a + ia \tan(c + dx))^2} \\
& \frac{12a^2}{12a^2} \\
& \downarrow 25 \\
& \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
& \frac{\frac{7a^2(-4B + iA) \tan^{\frac{3}{2}}(c + dx)}{d(a + ia \tan(c + dx))} - 3 \left(\frac{10a^3(A + 6iB) \sqrt{\tan(c + dx)}}{d} - \frac{2 \int \frac{a^3(5(A + 6iB) - 7(iA - 4B) \tan(c + dx))}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{d} \right)}{2a^2} - \frac{a(2A + 5iB) \tan^{\frac{5}{2}}(c + dx)}{d(a + ia \tan(c + dx))^2} \\
& \frac{12a^2}{12a^2}
\end{aligned}$$

3.147. $\int \frac{\tan^{\frac{7}{2}}(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^3} dx$

$$\begin{aligned} & \downarrow 27 \\ & \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\ & \frac{7a^2(-4B+iA) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{3 \left(\frac{10a^3(A+6iB)\sqrt{\tan(c+dx)}}{d} - \frac{2a^3 \int \frac{5(A+6iB)-7(iA-4B) \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} \right)}{2a^2} \\ & \frac{a(2A+5iB) \tan^{\frac{5}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \end{aligned}$$

$$\begin{aligned} & \frac{12a^2}{2a^2} \\ & \downarrow 1482 \\ & \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\ & \frac{7a^2(-4B+iA) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{3 \left(\frac{10a^3(A+6iB)\sqrt{\tan(c+dx)}}{d} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((6+i)A + (1+29i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} ((5-7i)A + (28+30i)B) \int \frac{\tan(c+dx)}{\tan^2(c+dx)} \right)}{d} \right)}{2a^2} \end{aligned}$$

$$\begin{aligned} & \frac{12a^2}{2a^2} \\ & \downarrow 1476 \\ & \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\ & \frac{7a^2(-4B+iA) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{3 \left(\frac{10a^3(A+6iB)\sqrt{\tan(c+dx)}}{d} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((6+i)A + (1+29i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} ((5-7i)A + (28+30i)B) \left(\frac{1}{2} \int \frac{\tan(c+dx)}{\tan^2(c+dx)} \right) \right)}{d} \right)}{2a^2} \end{aligned}$$

$$\begin{aligned} & \frac{12a^2}{2a^2} \\ & \downarrow 1082 \\ & \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\ & \frac{7a^2(-4B+iA) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{3 \left(\frac{10a^3(A+6iB)\sqrt{\tan(c+dx)}}{d} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((6+i)A + (1+29i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} ((5-7i)A + (28+30i)B) \left(\frac{1}{2} \int \frac{\tan(c+dx)}{\tan^2(c+dx)} \right) \right)}{d} \right)}{2a^2} \end{aligned}$$

$$\begin{aligned} & \frac{12a^2}{2a^2} \\ & \downarrow 217 \\ & \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\ & \frac{7a^2(-4B+iA) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \frac{3 \left(\frac{10a^3(A+6iB)\sqrt{\tan(c+dx)}}{d} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((6+i)A + (1+29i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} ((5-7i)A + (28+30i)B) \left(\frac{\arctan(\tan(c+dx))}{d} \right) \right)}{d} \right)}{2a^2} \end{aligned}$$

3.147. $\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned} & \downarrow 1479 \\ & \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\ & \frac{7a^2(-4B+iA) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \left(\frac{10a^3(A+6iB)\sqrt{\tan(c+dx)}}{d} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((6+i)A + (1+29i)B) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}}{\tan(c+dx)} \right)}{3} \right)}{2a^2} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 25 \\ & \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\ & \frac{7a^2(-4B+iA) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \left(\frac{10a^3(A+6iB)\sqrt{\tan(c+dx)}}{d} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((6+i)A + (1+29i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}}}{\tan(c+dx)+\sqrt{2}} \right)}{3} \right)}{2a^2} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\ & \frac{7a^2(-4B+iA) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \left(\frac{10a^3(A+6iB)\sqrt{\tan(c+dx)}}{d} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((6+i)A + (1+29i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}} \right)}{3} \right)}{2a^2} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 1103 \\ & \frac{(-B + iA) \tan^{\frac{7}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\ & \frac{7a^2(-4B+iA) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))} - \left(\frac{10a^3(A+6iB)\sqrt{\tan(c+dx)}}{d} - \frac{2a^3 \left(\frac{1}{2} ((5-7i)A + (28+30i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \left(\frac{1}{2} + \frac{i}{2} \right) \right)}{3} \right)}{2a^2} \end{aligned}$$

3.147. $\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

input `Int[(Tan[c + d*x]^(7/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `((I*A - B)*Tan[c + d*x]^(7/2))/(6*d*(a + I*a*Tan[c + d*x])^3) - (-((a*(2*A + (5*I)*B)*Tan[c + d*x]^(5/2))/(d*(a + I*a*Tan[c + d*x])^2)) + ((-3*((-2*a^3*(((5 - 7*I)*A + (28 + 30*I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]])/Sqrt[2]))/2 + (1/2 + I/2)*((6 + I)*A + (1 + 29*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/d + (10*a^3*(A + (6*I)*B)*Sqrt[Tan[c + d*x]]/d)/(2*a^2) + (7*a^2*(I*A - 4*B)*Tan[c + d*x]^(3/2))/(d*(a + I*a*Tan[c + d*x]))/(2*a^2))/(12*a^2)`

3.147.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

3.147.
$$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x], Sqrt[b*Tan[e + f*x]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

```
rule 4078 Int[((a_) + (b_) * tan[(e_) + (f_) * (x_)])^(m_) * ((A_) + (B_) * tan[(e_) + (f_) * (x_)]) * ((c_) + (d_) * tan[(e_) + (f_) * (x_)])^(n_), x_Symbol] := Simp[(- (A * b - a * B)) * (a + b * Tan[e + f * x])^m * ((c + d * Tan[e + f * x])^n / (2 * a * f * m)), x] + Simp[1 / (2 * a^2 * m) Int[(a + b * Tan[e + f * x])^(m + 1) * (c + d * Tan[e + f * x])^(n - 1) * Simp[A * (a * c * m + b * d * n) - B * (b * c * m + a * d * n) - d * (b * B * (m - n) - a * A * (m + n)) * Tan[e + f * x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b * c - a * d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.147.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 181, normalized size of antiderivative = 0.50

method	result
derivativedivides	$\frac{2iB(\sqrt{\tan(dx+c)}) - \frac{i \left(\frac{-i(9iA-20B)}{8} \left(\tan^{\frac{5}{2}}(dx+c) \right) + \left(-\frac{38iA}{3} + \frac{98B}{3} \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) + (-14iB-5A)(\sqrt{\tan(dx+c)}) \right)}{d a^3}}{d a^3}$
default	$\frac{2iB(\sqrt{\tan(dx+c)}) - \frac{i \left(\frac{-i(9iA-20B)}{8} \left(\tan^{\frac{5}{2}}(dx+c) \right) + \left(-\frac{38iA}{3} + \frac{98B}{3} \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) + (-14iB-5A)(\sqrt{\tan(dx+c)}) \right)}{d a^3}}{d a^3}$

```
input int(tan(d*x+c)^(7/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)
```

```
output 1/d/a^3*(2*I*B*tan(d*x+c)^(1/2)-1/8*I*((-I*(9*I*A-20*B))*tan(d*x+c)^(5/2)+(-38/3*I*A+98/3*B)*tan(d*x+c)^(3/2)+(-5*A-14*I*B)*tan(d*x+c)^(1/2))/(tan(d*x+c)-I)^3-2*(29*I*B+6*A)/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))+4*(1/16*I*A+1/16*B)/(2^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))
```

3.147. $\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

3.147.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 685 vs. $2(271) = 542$.

Time = 0.28 (sec) , antiderivative size = 685, normalized size of antiderivative = 1.88

$$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx =$$

$$\left(3a^3d\sqrt{\frac{-iA^2-2AB+iB^2}{a^6d^2}}e^{(6idx+6ic)} \log\left(-\frac{2\left((ia^3de^{(2idx+2ic)}+ia^3d)\sqrt{\frac{-ie^{(2idx+2ic)}+i}{e^{(2idx+2ic)}+1}}\sqrt{\frac{-iA^2-2AB+iB^2}{a^6d^2}}-(A-iB)e^{(2i}}}{iA+B} \right. \right.$$

```
input integrate(tan(d*x+c)^(7/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorith
thm="fricas")
```

```
output -1/96*(3*a^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c
)*log(-2*((I*a^3*d*e^(2*I*d*x + 2*I*c) + I*a^3*d)*sqrt((-I*e^(2*I*d*x + 2*
I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^
2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*a
^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(-2*(
(-I*a^3*d*e^(2*I*d*x + 2*I*c) - I*a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)
/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2)) - (A
- I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) + 3*a^3*d*sqrt
((36*I*A^2 - 348*A*B - 841*I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(-1/8*
((a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^
(2*I*d*x + 2*I*c) + 1))*sqrt((36*I*A^2 - 348*A*B - 841*I*B^2)/(a^6*d^2)) +
6*I*A - 29*B)*e^(-2*I*d*x - 2*I*c)/(a^3*d)) - 3*a^3*d*sqrt((36*I*A^2 - 34
8*A*B - 841*I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(1/8*((a^3*d*e^(2*I*d
*x + 2*I*c) + a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c
) + 1))*sqrt((36*I*A^2 - 348*A*B - 841*I*B^2)/(a^6*d^2)) - 6*I*A + 29*B)*e
^(-2*I*d*x - 2*I*c)/(a^3*d)) - 2*(2*(10*A + 73*I*B)*e^(6*I*d*x + 6*I*c) +
(14*A + 41*I*B)*e^(4*I*d*x + 4*I*c) - (5*A + 8*I*B)*e^(2*I*d*x + 2*I*c) +
A + I*B)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*e^(
-6*I*d*x - 6*I*c)/(a^3*d)
```

$$3.147. \quad \int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

3.147.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(7/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)`

output `Timed out`

3.147.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(7/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.147.8 Giac [A] (verification not implemented)

Time = 0.95 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.46

$$\begin{aligned} & \int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx \\ &= -\frac{(i+1)\sqrt{2}(-6iA+29B)\arctan\left(\left(\frac{1}{2}i+\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{16a^3d} \\ &+ \frac{(i-1)\sqrt{2}(-iA-B)\arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{16a^3d} + \frac{2iB\sqrt{\tan(dx+c)}}{a^3d} \\ &+ \frac{-27iA\tan(dx+c)^{\frac{5}{2}}+60B\tan(dx+c)^{\frac{5}{2}}-38A\tan(dx+c)^{\frac{3}{2}}-98iB\tan(dx+c)^{\frac{3}{2}}+15iA\sqrt{\tan(dx+c)}}{24a^3d(\tan(dx+c)-i)^3} \end{aligned}$$

3.147. $\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

input `integrate(tan(d*x+c)^(7/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output `-(1/16*I + 1/16)*sqrt(2)*(-6*I*A + 29*B)*arctan((1/2*I + 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^3*d) + (1/16*I - 1/16)*sqrt(2)*(-I*A - B)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^3*d) + 2*I*B*sqrt(tan(d*x + c))/(a^3*d) + 1/24*(-27*I*A*tan(d*x + c)^(5/2) + 60*B*tan(d*x + c)^(5/2) - 38*A*tan(d*x + c)^(3/2) - 98*I*B*tan(d*x + c)^(3/2) + 15*I*A*sqrt(tan(d*x + c)) - 42*B*sqrt(tan(d*x + c)))/(a^3*d*(tan(d*x + c) - I)^3)`

3.147.9 Mupad [B] (verification not implemented)

Time = 11.53 (sec) , antiderivative size = 395, normalized size of antiderivative = 1.09

$$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \operatorname{atan}\left(\frac{8a^3 d \sqrt{\tan(c+dx)} \sqrt{\frac{A^2 9i}{64a^6 d^2}}}{3A}\right) \sqrt{\frac{A^2 9i}{64a^6 d^2}} 2i$$

$$+ \operatorname{atan}\left(\frac{16a^3 d \sqrt{\tan(c+dx)} \sqrt{-\frac{A^2 1i}{256a^6 d^2}}}{A}\right) \sqrt{-\frac{A^2 1i}{256a^6 d^2}} 2i$$

$$+ \operatorname{atan}\left(\frac{a^3 d \sqrt{\tan(c+dx)} \sqrt{\frac{B^2 1i}{256a^6 d^2}} 16i}{B}\right) \sqrt{\frac{B^2 1i}{256a^6 d^2}} 2i$$

$$- \operatorname{atan}\left(\frac{a^3 d \sqrt{\tan(c+dx)} \sqrt{-\frac{B^2 841i}{256a^6 d^2}} 16i}{29B}\right) \sqrt{-\frac{B^2 841i}{256a^6 d^2}} 2i$$

$$+ \frac{\frac{5A \sqrt{\tan(c+dx)}}{8a^3 d} - \frac{9A \tan(c+dx)^{5/2}}{8a^3 d} + \frac{A \tan(c+dx)^{3/2} 19i}{12a^3 d}}{-\tan(c+dx)^3 1i - 3 \tan(c+dx)^2 + \tan(c+dx) 3i + 1}$$

$$- \frac{\frac{49B \tan(c+dx)^{3/2}}{12a^3 d} - \frac{B \sqrt{\tan(c+dx)} 7i}{4a^3 d} + \frac{B \tan(c+dx)^{5/2} 5i}{2a^3 d}}{-\tan(c+dx)^3 1i - 3 \tan(c+dx)^2 + \tan(c+dx) 3i + 1} + \frac{B \sqrt{\tan(c+dx)} 2i}{a^3 d}$$

input `int((tan(c + d*x)^(7/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3,x)`

output $\operatorname{atan}\left(\frac{8a^3d\tan(c+dx)^{1/2}\left(\frac{A^2\cdot 9i}{64a^6d^2}\right)^{1/2}}{3A}\right)\left(\frac{A^2\cdot 9i}{64a^6d^2}\right)^{1/2}\cdot 2i + \operatorname{atan}\left(\frac{16a^3d\tan(c+dx)^{1/2}\left(-\frac{A^2\cdot 1i}{256a^6d^2}\right)^{1/2}}{A}\right)\left(-\frac{A^2\cdot 1i}{256a^6d^2}\right)^{1/2}\cdot 2i + \operatorname{atan}\left(\frac{a^3d\tan(c+dx)^{1/2}\left(\frac{B^2\cdot 1i}{256a^6d^2}\right)^{1/2}\cdot 16i}{B}\right)\left(\frac{B^2\cdot 1i}{256a^6d^2}\right)^{1/2}\cdot 2i - \operatorname{atan}\left(\frac{a^3d\tan(c+dx)^{1/2}\left(-\frac{B^2\cdot 841i}{256a^6d^2}\right)^{1/2}\cdot 16i}{29B}\right)\left(-\frac{B^2\cdot 841i}{256a^6d^2}\right)^{1/2}\cdot 2i + \left(\frac{5A\tan(c+dx)^{1/2}}{8a^3d} + \frac{A\tan(c+dx)^{3/2}\cdot 19i}{12a^3d} - \frac{9A\tan(c+dx)^{5/2}}{8a^3d}\right)\left(\frac{1}{\tan(c+dx)\cdot 3i - 3\tan(c+dx)^2 - \tan(c+dx)^3\cdot 1i + 1} - \left(\frac{49B\tan(c+dx)^{3/2}}{12a^3d} - \frac{B\tan(c+dx)^{1/2}\cdot 7i}{4a^3d} + \frac{B\tan(c+dx)^{5/2}\cdot 5i}{2a^3d}\right)\left(\frac{1}{\tan(c+dx)\cdot 3i - 3\tan(c+dx)^2 - \tan(c+dx)^3\cdot 1i + 1} + \frac{B\tan(c+dx)^{1/2}\cdot 2i}{a^3d}\right)\right)$

3.147. $\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

3.148 $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

3.148.1 Optimal result 1541
 3.148.2 Mathematica [A] (verified) 1542
 3.148.3 Rubi [A] (verified) 1542
 3.148.4 Maple [A] (verified) 1548
 3.148.5 Fricas [B] (verification not implemented) 1548
 3.148.6 Sympy [F(-1)] 1549
 3.148.7 Maxima [F(-2)] 1549
 3.148.8 Giac [A] (verification not implemented) 1550
 3.148.9 Mupad [B] (verification not implemented) 1551

3.148.1 Optimal result

Integrand size = 36, antiderivative size = 307

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{(2A+(5-7i)B) \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{16\sqrt{2}a^3d}$$

$$- \frac{(2A+(5-7i)B) \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{16\sqrt{2}a^3d}$$

$$- \frac{(2A-(5+7i)B) \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{32\sqrt{2}a^3d}$$

$$+ \frac{(2A-(5+7i)B) \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{32\sqrt{2}a^3d}$$

$$+ \frac{(iA-B) \tan^{\frac{5}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} + \frac{(A+4iB) \tan^{\frac{3}{2}}(c+dx)}{12ad(a+ia \tan(c+dx))^2} + \frac{5B\sqrt{\tan(c+dx)}}{8d(a^3+ia^3 \tan(c+dx))}$$

output

```
-1/32*(2*A+(5-7*I)*B)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/a^3/d*2^(1/2)-1/
32*(2*A+(5-7*I)*B)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/a^3/d*2^(1/2)-1/64*(
2*A-(5+7*I)*B)*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a^3/d*2^(1/2)+1/6
4*(2*A-(5+7*I)*B)*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a^3/d*2^(1/2)+
1/6*(I*A-B)*tan(d*x+c)^(5/2)/d/(a+I*a*tan(d*x+c))^3+1/12*(A+4*I*B)*tan(d*x
+c)^(3/2)/a/d/(a+I*a*tan(d*x+c))^2+5/8*B*tan(d*x+c)^(1/2)/d/(a^3+I*a^3*tan
(d*x+c))
```

3.148. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

3.148.2 Mathematica [A] (verified)

Time = 3.76 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.64

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \frac{\sec^3(c+dx) \left(-3\sqrt[4]{-1}(A-iB) \arctan \left((-1)^{3/4} \sqrt{\tan(c+dx)} \right) (\cos(3(c+dx)) + i \sin(3(c+dx))) + 3 \right)}{a^3 d (-I + \tan(c+dx))^3}$$

input `Integrate[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `-1/24*(Sec[c + d*x]^3*(-3*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)]) + 3*(-1)^(1/4)*(A - (6*I)*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)]) + Cos[c + d*x]*(3*(A + (2*I)*B) - 3*(A + (7*I)*B)*Cos[2*(c + d*x)] + ((-I)*A + 19*B)*Sin[2*(c + d*x)])*Sqrt[Tan[c + d*x]])/(a^3*d*(-I + Tan[c + d*x])^3)`

3.148.3 Rubi [A] (verified)

Time = 1.15 (sec) , antiderivative size = 285, normalized size of antiderivative = 0.93, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 4078, 27, 3042, 4078, 27, 3042, 4078, 25, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(c+dx)^{5/2}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\ & \quad \downarrow \text{4078} \\ & \frac{(-B+iA) \tan^{\frac{5}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{\tan^{\frac{3}{2}}(c+dx)(5a(iA-B)-a(A-11iB) \tan(c+dx))}{2(i \tan(c+dx)a+a)^2} dx}{6a^2} \end{aligned}$$

3.148. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{\tan^{\frac{3}{2}}(c+dx)(5a(iA-B) - a(A-11iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{12a^2} \\
 & \downarrow 3042 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{\tan(c+dx)^{3/2}(5a(iA-B) - a(A-11iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{12a^2} \\
 & \downarrow 4078 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int -\frac{6\sqrt{\tan(c+dx)}((A+4iB)a^2 + (iA+6B) \tan(c+dx)a^2)}{i \tan(c+dx)a+a} dx}{4a^2} - \frac{a(A+4iB) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \downarrow 27 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \int \frac{\sqrt{\tan(c+dx)}((A+4iB)a^2 + (iA+6B) \tan(c+dx)a^2)}{i \tan(c+dx)a+a} dx}{2a^2} - \frac{a(A+4iB) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \downarrow 3042 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \int \frac{\sqrt{\tan(c+dx)}((A+4iB)a^2 + (iA+6B) \tan(c+dx)a^2)}{i \tan(c+dx)a+a} dx}{2a^2} - \frac{a(A+4iB) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \downarrow 4078 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\int -\frac{5Ba^3 + (2A-7iB) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx - \frac{5a^2B \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))} \right)}{2a^2} - \frac{a(A+4iB) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \downarrow 25 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{\int \frac{5Ba^3 + (2A-7iB) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx}{2a^2} - \frac{5a^2B \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))} \right)}{2a^2} - \frac{a(A+4iB) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \downarrow 3042 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{\int \frac{5Ba^3 + (2A-7iB) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx}{2a^2} - \frac{5a^2B \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))} \right)}{2a^2} - \frac{a(A+4iB) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \downarrow 4017
 \end{aligned}$$

3.148. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
 & \frac{3 \left(\frac{\int \frac{a^3(5B+(2A-7iB)\tan(c+dx)) d\sqrt{\tan(c+dx)}}{\tan^2(c+dx)+1} - \frac{5a^2 B \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))} \right)}{2a^2} - \frac{a(A+4iB) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \frac{12a^2}{12a^2} \\
 & \downarrow 27 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{a \int \frac{5B+(2A-7iB)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{5a^2 B \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))} \right)}{2a^2} - \frac{a(A+4iB) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \frac{12a^2}{12a^2} \\
 & \downarrow 1482 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
 & \frac{3 \left(\frac{a \left(\frac{1}{2}(2A+(5-7i)B) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(2A-(5+7i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) - \frac{5a^2 B \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))} \right)}{2a^2} - \frac{a(A+4iB) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \frac{12a^2}{12a^2} \\
 & \downarrow 1476 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
 & \frac{3 \left(\frac{a \left(\frac{1}{2}(2A+(5-7i)B) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2}(2A-(5+7i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) - \frac{5a^2 B \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))} \right)}{2a^2} - \frac{a(A+4iB) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \frac{12a^2}{12a^2} \\
 & \downarrow 1082 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
 & \frac{3 \left(\frac{a \left(\frac{1}{2}(2A+(5-7i)B) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) - \frac{1}{2}(2A-(5+7i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) - \frac{5a^2 B \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))} \right)}{2a^2} - \frac{a(A+4iB) \tan^{\frac{3}{2}}(c+dx)}{d(a+ia \tan(c+dx))^2} \\
 & \frac{12a^2}{12a^2} \\
 & \downarrow 217
 \end{aligned}$$

3.148. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(a \left(\frac{1}{2}(2A + (5 - 7i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c + dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c + dx)})}{\sqrt{2}} \right) - \frac{1}{2}(2A - (5 + 7i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} \right) - \frac{5a^2 B \sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))}}{2a^2}$$

12a²

↓ 1479

$$\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(a \left(\frac{1}{2}(2A + (5 - 7i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c + dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c + dx)})}{\sqrt{2}} \right) - \frac{1}{2}(2A - (5 + 7i)B) \left(\int \frac{\sqrt{2} - 2\sqrt{\tan(c + dx)}}{\tan(c + dx) - \sqrt{2}\sqrt{\tan(c + dx)} + 1} d\sqrt{\tan(c + dx)} - \int \frac{\sqrt{2}}{\tan(c + dx) - \sqrt{2}\sqrt{\tan(c + dx)} + 1} d\sqrt{\tan(c + dx)} \right) \right)}{2a^2}$$

2a²

12a²

↓ 25

$$\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(a \left(\frac{1}{2}(2A + (5 - 7i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c + dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c + dx)})}{\sqrt{2}} \right) - \frac{1}{2}(2A - (5 + 7i)B) \left(\int \frac{\sqrt{2} - 2\sqrt{\tan(c + dx)}}{\tan(c + dx) - \sqrt{2}\sqrt{\tan(c + dx)} + 1} d\sqrt{\tan(c + dx)} + \int \frac{\sqrt{2}}{\tan(c + dx) - \sqrt{2}\sqrt{\tan(c + dx)} + 1} d\sqrt{\tan(c + dx)} \right) \right)}{2a^2}$$

2a²

12a²

↓ 27

$$\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(a \left(\frac{1}{2}(2A + (5 - 7i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c + dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c + dx)})}{\sqrt{2}} \right) - \frac{1}{2}(2A - (5 + 7i)B) \left(\int \frac{\sqrt{2} - 2\sqrt{\tan(c + dx)}}{\tan(c + dx) - \sqrt{2}\sqrt{\tan(c + dx)} + 1} d\sqrt{\tan(c + dx)} + \frac{1}{2} \int \frac{\sqrt{2}}{\tan(c + dx) - \sqrt{2}\sqrt{\tan(c + dx)} + 1} d\sqrt{\tan(c + dx)} \right) \right)}{2a^2}$$

2a²

12a²

↓ 1103

3.148. $\int \frac{\tan^{\frac{5}{2}}(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^3} dx$

$$\frac{\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{a \left(\frac{1}{2}(2A + (5 - 7i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(2A - (5 + 7i)B) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right)}{d}}{2a^2}}{12a^2}$$

input `Int[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `((I*A - B)*Tan[c + d*x]^(5/2))/(6*d*(a + I*a*Tan[c + d*x])^3) - (-((a*(A + (4*I)*B)*Tan[c + d*x]^(3/2))/(d*(a + I*a*Tan[c + d*x])^2)) + (3*((a*((2*A + (5 - 7*I)*B)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 - ((2*A - (5 + 7*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2))/d - (5*a^2*B*Sqrt[Tan[c + d*x]]/(d*(a + I*a*Tan[c + d*x]))))/(2*a^2))/(12*a^2)`

3.148.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

3.148. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4078 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.148.
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.148.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.53

method	result
derivativedivides	$\frac{i \left(\frac{-i(9iB+2A) \tan^{\frac{5}{2}}(dx+c) + \left(-\frac{38iB}{3} - \frac{2A}{3}\right) \tan^{\frac{3}{2}}(dx+c) - 5B(\sqrt{\tan}(dx+c)) - \frac{2(iA+6B) \arctan\left(\frac{2(\sqrt{\tan}(dx+c))}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}}}{\left(\tan(dx+c)-i\right)^3} \right)}{8} + \dots}{da^3}$
default	$\frac{i \left(\frac{-i(9iB+2A) \tan^{\frac{5}{2}}(dx+c) + \left(-\frac{38iB}{3} - \frac{2A}{3}\right) \tan^{\frac{3}{2}}(dx+c) - 5B(\sqrt{\tan}(dx+c)) - \frac{2(iA+6B) \arctan\left(\frac{2(\sqrt{\tan}(dx+c))}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}}}{\left(\tan(dx+c)-i\right)^3} \right)}{8} + \dots}{da^3}$

```
input int(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)
```

```
output 1/d/a^3*(-1/8*I*((-I*(9*I*B+2*A))*tan(d*x+c)^(5/2)+(-38/3*I*B-2/3*A)*tan(d*x+c)^(3/2)-5*B*tan(d*x+c)^(1/2)))/(tan(d*x+c)-I)^3-2*(I*A+6*B)/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))+4*(-1/16*A+1/16*I*B)/(2^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))
```

3.148.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 681 vs. 2(242) = 484.

Time = 0.27 (sec) , antiderivative size = 681, normalized size of antiderivative = 2.22

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{\left(3a^3 d \sqrt{\frac{iA^2+2AB-iB^2}{a^6 d^2}} e^{(6i dx+6i c)} \log \left(\frac{2 \left((a^3 d e^{(2i dx+2i c)} + a^3 d) \sqrt{\frac{-i e^{(2i dx+2i c)} + i}{e^{(2i dx+2i c)} + 1}} \sqrt{\frac{iA^2+2AB-iB^2}{a^6 d^2}} + (A-iB) e^{(2i dx+2i c)} \right)}{iA+B} \right) \right)}{iA+B}$$

```
input integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="fracas")
```

3.148. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

output $\frac{1}{96}(3a^3d\sqrt{(IA^2 + 2AB - IB^2)/(a^6d^2)})e^{(6Id*x + 6I*c)} \log(2((a^3d e^{(2Id*x + 2I*c)} + a^3d)\sqrt{(-Ie^{(2Id*x + 2I*c)} + I)/(e^{(2Id*x + 2I*c)} + 1)})\sqrt{(IA^2 + 2AB - IB^2)/(a^6d^2)} + (A - IB)e^{(2Id*x + 2I*c)})e^{(-2Id*x - 2I*c)/(IA + B)} - 3a^3d\sqrt{(IA^2 + 2AB - IB^2)/(a^6d^2)})e^{(6Id*x + 6I*c)} \log(-2((a^3d e^{(2Id*x + 2I*c)} + a^3d)\sqrt{(-Ie^{(2Id*x + 2I*c)} + I)/(e^{(2Id*x + 2I*c)} + 1)})\sqrt{(IA^2 + 2AB - IB^2)/(a^6d^2)} - (A - IB)e^{(2Id*x + 2I*c)})e^{(-2Id*x - 2I*c)/(IA + B)} + 3a^3d\sqrt{(-IA^2 - 12AB + 36IB^2)/(a^6d^2)})e^{(6Id*x + 6I*c)} \log(1/8((a^3d e^{(2Id*x + 2I*c)} + a^3d)\sqrt{(-Ie^{(2Id*x + 2I*c)} + I)/(e^{(2Id*x + 2I*c)} + 1)})\sqrt{(-IA^2 - 12AB + 36IB^2)/(a^6d^2)} + A - 6IB)e^{(-2Id*x - 2I*c)/(a^3d)} - 3a^3d\sqrt{(-IA^2 - 12AB + 36IB^2)/(a^6d^2)})e^{(6Id*x + 6I*c)} \log(-1/8((a^3d e^{(2Id*x + 2I*c)} + a^3d)\sqrt{(-Ie^{(2Id*x + 2I*c)} + I)/(e^{(2Id*x + 2I*c)} + 1)})\sqrt{(-IA^2 - 12AB + 36IB^2)/(a^6d^2)} - A + 6IB)e^{(-2Id*x - 2I*c)/(a^3d)} - 2*(2*(IA - 10B)e^{(6Id*x + 6I*c)} - (IA + 14B)e^{(4Id*x + 4I*c)} - (2IA - 5B)e^{(2Id*x + 2I*c)} + IA - B)\sqrt{(-Ie^{(2Id*x + 2I*c)} + I)/(e^{(2Id*x + 2I*c)} + 1)}))e^{(-6Id*x - 6I*c)/(a^3d)}$

3.148.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{5}{2}}(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)`

output `Timed out`

3.148.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{5}{2}}(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

3.148. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

3.148.8 Giac [A] (verification not implemented)

Time = 0.85 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.44

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= -\frac{(i+1) \sqrt{2}(A-6iB) \arctan\left(\left(\frac{1}{2}i+\frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx+c)}\right)}{16a^3d}$$

$$+ \frac{(i-1) \sqrt{2}(A-iB) \arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx+c)}\right)}{16a^3d}$$

$$- \frac{6A \tan(dx+c)^{\frac{5}{2}} + 27iB \tan(dx+c)^{\frac{5}{2}} - 2iA \tan(dx+c)^{\frac{3}{2}} + 38B \tan(dx+c)^{\frac{3}{2}} - 15iB \sqrt{\tan(dx+c)}}{24a^3d(\tan(dx+c)-i)^3}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output `-(1/16*I + 1/16)*sqrt(2)*(A - 6*I*B)*arctan((1/2*I + 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^3*d) + (1/16*I - 1/16)*sqrt(2)*(A - I*B)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^3*d) - 1/24*(6*A*tan(d*x + c)^(5/2) + 27*I*B*tan(d*x + c)^(5/2) - 2*I*A*tan(d*x + c)^(3/2) + 38*B*tan(d*x + c)^(3/2) - 15*I*B*sqrt(tan(d*x + c)))/(a^3*d*(tan(d*x + c) - I)^3)`

3.148.9 Mupad [B] (verification not implemented)

Time = 9.99 (sec) , antiderivative size = 308, normalized size of antiderivative = 1.00

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$= \frac{\frac{5B\sqrt{\tan(c+dx)}}{8a^3d} - \frac{9B\tan(c+dx)^{5/2}}{8a^3d} + \frac{B\tan(c+dx)^{3/2}19i}{12a^3d}}{-\tan(c+dx)^3 i - 3\tan(c+dx)^2 + \tan(c+dx) 3i + 1}$$

$$+ \frac{\frac{A\tan(c+dx)^{3/2}}{12a^3d} + \frac{A\tan(c+dx)^{5/2}1i}{4a^3d}}{-\tan(c+dx)^3 i - 3\tan(c+dx)^2 + \tan(c+dx) 3i + 1}$$

$$- \frac{(-1)^{1/4} A \operatorname{atan}\left((-1)^{1/4} \sqrt{\tan(c+dx)}\right)}{8a^3d} + \frac{(-1)^{1/4} A \operatorname{atanh}\left((-1)^{1/4} \sqrt{\tan(c+dx)}\right)}{8a^3d}$$

$$+ \operatorname{atan}\left(\frac{8a^3d\sqrt{\tan(c+dx)}\sqrt{\frac{B^2 9i}{64a^6d^2}}}{3B}\right) \sqrt{\frac{B^2 9i}{64a^6d^2}} 2i + \operatorname{atan}\left(\frac{16a^3d\sqrt{\tan(c+dx)}\sqrt{-\frac{B^2 1i}{256a^6d^2}}}{B}\right) \sqrt{-\frac{B^2 1i}{256a^6d^2}}$$

input `int((tan(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3,x)`output `atan((8*a^3*d*tan(c + d*x)^(1/2)*((B^2*9i)/(64*a^6*d^2))^(1/2))/(3*B))*((B^2*9i)/(64*a^6*d^2))^(1/2)*2i + atan((16*a^3*d*tan(c + d*x)^(1/2)*(-(B^2*1i)/(256*a^6*d^2))^(1/2))/B)*(-(B^2*1i)/(256*a^6*d^2))^(1/2)*2i + ((5*B*tan(c + d*x)^(1/2))/(8*a^3*d) + (B*tan(c + d*x)^(3/2)*19i)/(12*a^3*d) - (9*B*tan(c + d*x)^(5/2))/(8*a^3*d))/(tan(c + d*x)*3i - 3*tan(c + d*x)^2 - tan(c + d*x)^3*1i + 1) + ((A*tan(c + d*x)^(3/2))/(12*a^3*d) + (A*tan(c + d*x)^(5/2)*1i)/(4*a^3*d))/(tan(c + d*x)*3i - 3*tan(c + d*x)^2 - tan(c + d*x)^3*1i + 1) - ((-1)^(1/4)*A*atan((-1)^(1/4)*tan(c + d*x)^(1/2))/(8*a^3*d) + ((-1)^(1/4)*A*atanh((-1)^(1/4)*tan(c + d*x)^(1/2)))/(8*a^3*d)`

3.149
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.149.1 Optimal result 1552
 3.149.2 Mathematica [A] (verified) 1553
 3.149.3 Rubi [A] (verified) 1553
 3.149.4 Maple [A] (verified) 1558
 3.149.5 Fricas [B] (verification not implemented) 1559
 3.149.6 Sympy [F] 1560
 3.149.7 Maxima [F(-2)] 1561
 3.149.8 Giac [A] (verification not implemented) 1561
 3.149.9 Mupad [B] (verification not implemented) 1562

3.149.1 Optimal result

Integrand size = 36, antiderivative size = 309

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{((1+i)A+2B) \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{16\sqrt{2}a^3d}$$

$$- \frac{((1+i)A+2B) \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{16\sqrt{2}a^3d}$$

$$- \frac{((-1+i)A+2B) \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{32\sqrt{2}a^3d}$$

$$+ \frac{((-1+i)A+2B) \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{32\sqrt{2}a^3d}$$

$$+ \frac{(iA-B) \tan^{\frac{3}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} + \frac{iB\sqrt{\tan(c+dx)}}{4ad(a+ia \tan(c+dx))^2} + \frac{(A-2iB)\sqrt{\tan(c+dx)}}{8d(a^3+ia^3 \tan(c+dx))}$$

```
output -1/32*((1+I)*A+2*B)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/a^3/d*2^(1/2)-1/32
*((1+I)*A+2*B)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/a^3/d*2^(1/2)-1/64*((-1+
I)*A+2*B)*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a^3/d*2^(1/2)+1/64*((-
1+I)*A+2*B)*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a^3/d*2^(1/2)+1/6*(I
*A-B)*tan(d*x+c)^(3/2)/d/(a+I*a*tan(d*x+c))^3+1/4*I*B*tan(d*x+c)^(1/2)/a/d
/(a+I*a*tan(d*x+c))^2+1/8*(A-2*I*B)*tan(d*x+c)^(1/2)/d/(a^3+I*a^3*tan(d*x+
c))
```

3.149.
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.149.2 Mathematica [A] (verified)

Time = 4.44 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.59

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{3\sqrt[4]{-1}(iA+B) \arctan\left((-1)^{3/4} \sqrt{\tan(c+dx)}\right) \sec^3(c+dx)(\cos(3(c+dx)) + i \sin(3(c+dx))) - 3\sqrt[4]{-1}A \tan^{\frac{3}{2}}(c+dx)}{6a^2}$$

input `Integrate[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `(3*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^3*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)]) - 3*(-1)^(1/4)*B*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^3*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)]) - I*Sqrt[Tan[c + d*x]]*(-I + 3*Tan[c + d*x])*((-3*I)*A + (A - (2*I)*B)*Tan[c + d*x])/(24*a^3*d*(-I + Tan[c + d*x])^3)`

3.149.3 Rubi [A] (verified)Time = 1.12 (sec) , antiderivative size = 287, normalized size of antiderivative = 0.93, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.528$, Rules used = {3042, 4078, 27, 3042, 4078, 27, 3042, 4079, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$\downarrow 3042$$

$$\int \frac{\tan(c+dx)^{3/2}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$\downarrow 4078$$

$$\frac{(-B+iA) \tan^{\frac{3}{2}}(c+dx)}{6d(a+ia \tan(c+dx))^3} - \frac{\int \frac{3\sqrt{\tan(c+dx)}(a(iA-B)-a(A-3iB) \tan(c+dx))}{2(i \tan(c+dx)a+a)^2} dx}{6a^2}$$

$$\downarrow 27$$

3.149. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{\sqrt{\tan(c+dx)}(a(iA-B) - a(A-3iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{4a^2} \\
& \quad \downarrow 3042 \\
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{\sqrt{\tan(c+dx)}(a(iA-B) - a(A-3iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^2} dx}{4a^2} \\
& \quad \downarrow 4078 \\
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int -\frac{2(iBa^2 + (2iA+3B) \tan(c+dx)a^2)}{\sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)} dx}{4a^2} - \frac{iaB \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))^2} \\
& \quad \downarrow 27 \\
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{iBa^2 + (2iA+3B) \tan(c+dx)a^2}{\sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)} dx}{2a^2} - \frac{iaB \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))^2} \\
& \quad \downarrow 3042 \\
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{iBa^2 + (2iA+3B) \tan(c+dx)a^2}{\sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)} dx}{2a^2} - \frac{iaB \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))^2} \\
& \quad \downarrow 4079 \\
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{Aa^3 + (iA+2B) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx}{2a^2} - \frac{a^2(A-2iB) \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))} - \frac{iaB \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))^2} \\
& \quad \downarrow 3042 \\
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{Aa^3 + (iA+2B) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx}{2a^2} - \frac{a^2(A-2iB) \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))} - \frac{iaB \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))^2} \\
& \quad \downarrow 4017 \\
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{a^3(A+(iA+2B) \tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{a^2d} - \frac{a^2(A-2iB) \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))} - \frac{iaB \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))^2} \\
& \quad \downarrow 27 \\
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{a \int \frac{A+(iA+2B) \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} - \frac{a^2(A-2iB) \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))} - \frac{iaB \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))^2}
\end{aligned}$$

3.149. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{array}{c}
\downarrow 1482 \\
\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
\frac{a \left(\frac{1}{2}(-2B + (1-i)A) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(2B + (1+i)A) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} - \frac{a^2(A-2iB)\sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))} - \frac{iaB\sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))^2} \\
\hline
4a^2 \\
\downarrow 1476 \\
\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
\frac{a \left(\frac{1}{2}(-2B + (1-i)A) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(2B + (1+i)A) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d} \\
\hline
2a^2 \\
\hline
4a^2 \\
\downarrow 1082 \\
\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
\frac{a \left(\frac{1}{2}(-2B + (1-i)A) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(2B + (1+i)A) \left(\int \frac{1}{-\tan(c+dx)-1} \frac{d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \int \frac{1}{-\tan(c+dx)-1} \frac{d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \\
\hline
2a^2 \\
\hline
4a^2 \\
\downarrow 217 \\
\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
\frac{a \left(\frac{1}{2}(-2B + (1-i)A) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(2B + (1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d} - \frac{a^2(A-2iB)\sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))} \\
\hline
2a^2 \\
\hline
4a^2 \\
\downarrow 1479 \\
\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \\
\frac{a \left(\frac{1}{2}(-2B + (1-i)A) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(2B + (1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \\
\hline
2a^2 \\
\hline
4a^2 \\
\downarrow 25
\end{array}$$

$$3.149. \quad \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{a \left(\frac{1}{2}(-2B + (1-i)A) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2}(2B+(1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right)}{d} \right)}{2a^2} \frac{4a^2}{4a^2}$$

↓ 27

$$\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{a \left(\frac{1}{2}(-2B + (1-i)A) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2}(2B+(1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right)}{d} \right)}{2a^2} \frac{4a^2}{4a^2}$$

↓ 1103

$$\frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{6d(a + ia \tan(c + dx))^3} - \frac{a \left(\frac{1}{2}(2B+(1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(-2B+(1-i)A) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d} \right)}{2a^2} \frac{4a^2}{4a^2}$$

input `Int[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `((I*A - B)*Tan[c + d*x]^(3/2))/(6*d*(a + I*a*Tan[c + d*x])^3) - (((-I)*a*B*Sqrt[Tan[c + d*x]])/(d*(a + I*a*Tan[c + d*x])^2) + ((a((((1 + I)*A + 2*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]] + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])))/2 + (((1 - I)*A - 2*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d - (a^2*(A - (2*I)*B)*Sqrt[Tan[c + d*x]])/(d*(a + I*a*Tan[c + d*x]))/(2*a^2))/(4*a^2)`

3.149.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

3.149.
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] & NeQ[c^2 + d^2, 0]`

rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.149.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.49

method	result
derivativedivides	$\frac{-i(-2iB+A)\left(\tan^{\frac{5}{2}}(dx+c)\right)+\left(-\frac{10A}{3}+\frac{2iB}{3}\right)\left(\tan^{\frac{3}{2}}(dx+c)\right)+iA(\sqrt{\tan(dx+c)})-B\arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{8(\tan(dx+c)-i)^3} - \frac{4\left(-\frac{iA}{16}-\frac{B}{16}\right)\arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{4(\sqrt{2-i\sqrt{2}})} + \frac{4\left(-\frac{iA}{16}-\frac{B}{16}\right)\arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2}}$
default	$\frac{-i(-2iB+A)\left(\tan^{\frac{5}{2}}(dx+c)\right)+\left(-\frac{10A}{3}+\frac{2iB}{3}\right)\left(\tan^{\frac{3}{2}}(dx+c)\right)+iA(\sqrt{\tan(dx+c)})-B\arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{8(\tan(dx+c)-i)^3} - \frac{4\left(-\frac{iA}{16}-\frac{B}{16}\right)\arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{4(\sqrt{2-i\sqrt{2}})} + \frac{4\left(-\frac{iA}{16}-\frac{B}{16}\right)\arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2}}$

$$3.149. \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

```
input int(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)
```

```
output 1/d/a^3*(1/8*(-I*(A-2*I*B)*tan(d*x+c)^(5/2)+(-10/3*A+2/3*I*B)*tan(d*x+c)^(3/2)+I*A*tan(d*x+c)^(1/2))/(tan(d*x+c)-I)^3-1/4*B/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))+4*(-1/16*I*A-1/16*B)/(2^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))
```

3.149.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 635 vs. $2(242) = 484$.

Time = 0.27 (sec) , antiderivative size = 635, normalized size of antiderivative = 2.06

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \left(3a^3 d \sqrt{\frac{-iA^2-2AB+iB^2}{a^6 d^2}} e^{(6i dx+6i c)} \log \left(-\frac{2 \left((i a^3 d e^{(2i dx+2i c)} + i a^3 d) \sqrt{\frac{-i e^{(2i dx+2i c)} + i}{e^{(2i dx+2i c)} + 1}} \sqrt{\frac{-iA^2-2AB+iB^2}{a^6 d^2}} - (A-iB) e^{(2i dx+2i c)} \right)}{iA+B} \right) \right)$$

```
input integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="fracas")
```

output $\frac{1}{96}(3a^3d\sqrt{(-IA^2 - 2AB + IB^2)/(a^6d^2)})e^{(6Id*x + 6I*c)} \cdot \log(-2((Ia^3de^{(2Id*x + 2I*c)} + Ia^3d)\sqrt{(-Ie^{(2Id*x + 2I*c)} + I)/(e^{(2Id*x + 2I*c)} + 1)})\sqrt{(-IA^2 - 2AB + IB^2)/(a^6d^2)}) - (A - IB)e^{(2Id*x + 2I*c)})e^{(-2Id*x - 2I*c)/(IA + B)} - 3a^3d\sqrt{(-IA^2 - 2AB + IB^2)/(a^6d^2)})e^{(6Id*x + 6I*c)} \cdot \log(-2((-Ia^3de^{(2Id*x + 2I*c)} - Ia^3d)\sqrt{(-Ie^{(2Id*x + 2I*c)} + I)/(e^{(2Id*x + 2I*c)} + 1)})\sqrt{(-IA^2 - 2AB + IB^2)/(a^6d^2)}) - (A - IB)e^{(2Id*x + 2I*c)})e^{(-2Id*x - 2I*c)/(IA + B)} + 24a^3d\sqrt{(-1/64IB^2/(a^6d^2)})e^{(6Id*x + 6I*c)} \cdot \log(1/8(8(a^3de^{(2Id*x + 2I*c)} + a^3d)\sqrt{(-Ie^{(2Id*x + 2I*c)} + I)/(e^{(2Id*x + 2I*c)} + 1)})\sqrt{(-1/64IB^2/(a^6d^2)) + B})e^{(-2Id*x - 2I*c)/(a^3d)}) - 24a^3d\sqrt{(-1/64IB^2/(a^6d^2)})e^{(6Id*x + 6I*c)} \cdot \log(-1/8(8(a^3de^{(2Id*x + 2I*c)} + a^3d)\sqrt{(-Ie^{(2Id*x + 2I*c)} + I)/(e^{(2Id*x + 2I*c)} + 1)})\sqrt{(-1/64IB^2/(a^6d^2)) - B})e^{(-2Id*x - 2I*c)/(a^3d)}) + 2*(2*(2A - IB)e^{(6Id*x + 6I*c)} + (4A + IB)e^{(4Id*x + 4I*c)} - (A - 2IB)e^{(2Id*x + 2I*c)} - A - IB)\sqrt{(-Ie^{(2Id*x + 2I*c)} + I)/(e^{(2Id*x + 2I*c)} + 1)})e^{(-6Id*x - 6I*c)/(a^3d)}$

3.149.6 Sympy [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{i \left(\int \frac{A \tan^{\frac{3}{2}}(c+dx)}{\tan^3(c+dx) - 3i \tan^2(c+dx) - 3 \tan(c+dx) + i} dx + \int \frac{B \tan^{\frac{5}{2}}(c+dx)}{\tan^3(c+dx) - 3i \tan^2(c+dx) - 3 \tan(c+dx) + i} dx \right)}{a^3}$$

input `integrate(tan(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)`

output $I*(\text{Integral}(A*\tan(c + d*x)**(3/2)/(\tan(c + d*x)**3 - 3*I*\tan(c + d*x)**2 - 3*\tan(c + d*x) + I), x) + \text{Integral}(B*\tan(c + d*x)**(5/2)/(\tan(c + d*x)**3 - 3*I*\tan(c + d*x)**2 - 3*\tan(c + d*x) + I), x))/a**3$

3.149.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

3.149.8 Giac [A] (verification not implemented)

Time = 0.78 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.42

$$\begin{aligned} & \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx \\ &= -\frac{(i+1)\sqrt{2}B\arctan\left(\left(\frac{1}{2}i+\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{16a^3d} \\ & \quad + \frac{(i-1)\sqrt{2}(iA+B)\arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{16a^3d} \\ & \quad - \frac{3iA\tan(dx+c)^{\frac{5}{2}}+6B\tan(dx+c)^{\frac{5}{2}}+10A\tan(dx+c)^{\frac{3}{2}}-2iB\tan(dx+c)^{\frac{3}{2}}-3iA\sqrt{\tan(dx+c)}}{24a^3d(\tan(dx+c)-i)^3} \end{aligned}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output `-(1/16*I + 1/16)*sqrt(2)*B*arctan((1/2*I + 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^3*d) + (1/16*I - 1/16)*sqrt(2)*(I*A + B)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^3*d) - 1/24*(3*I*A*tan(d*x + c)^(5/2) + 6*B*tan(d*x + c)^(5/2) + 10*A*tan(d*x + c)^(3/2) - 2*I*B*tan(d*x + c)^(3/2) - 3*I*A*sqrt(tan(d*x + c)))/(a^3*d*(tan(d*x + c) - I)^3)`

3.149.9 Mupad [B] (verification not implemented)

Time = 8.36 (sec) , antiderivative size = 239, normalized size of antiderivative = 0.77

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$= \frac{\frac{A\sqrt{\tan(c+dx)}}{8a^3d} - \frac{A\tan(c+dx)^{5/2}}{8a^3d} + \frac{A\tan(c+dx)^{3/2}5i}{12a^3d}}{-\tan(c+dx)^3 i - 3\tan(c+dx)^2 + \tan(c+dx) 3i + 1}$$

$$+ \frac{\frac{B\tan(c+dx)^{3/2}}{12a^3d} + \frac{B\tan(c+dx)^{5/2} i}{4a^3d}}{-\tan(c+dx)^3 i - 3\tan(c+dx)^2 + \tan(c+dx) 3i + 1}$$

$$- \frac{(-1)^{1/4} B \operatorname{atan}\left((-1)^{1/4} \sqrt{\tan(c+dx)}\right)}{8a^3d} + \frac{(-1)^{1/4} B \operatorname{atanh}\left((-1)^{1/4} \sqrt{\tan(c+dx)}\right)}{8a^3d}$$

$$- \frac{\sqrt{-\frac{1}{256}i} A \operatorname{atan}\left(16 \sqrt{-\frac{1}{256}i} \sqrt{\tan(c+dx)}\right) 2i}{a^3d}$$

input `int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3,x)`output `((A*tan(c + d*x)^(1/2))/(8*a^3*d) + (A*tan(c + d*x)^(3/2)*5i)/(12*a^3*d) - (A*tan(c + d*x)^(5/2))/(8*a^3*d))/(tan(c + d*x)*3i - 3*tan(c + d*x)^2 - tan(c + d*x)^3*1i + 1) + ((B*tan(c + d*x)^(3/2))/(12*a^3*d) + (B*tan(c + d*x)^(5/2)*i)/(4*a^3*d))/(tan(c + d*x)*3i - 3*tan(c + d*x)^2 - tan(c + d*x)^3*1i + 1) - ((-1i/256)^(1/2)*A*atan(16*(-1i/256)^(1/2)*tan(c + d*x)^(1/2))*2i)/(a^3*d) - ((-1)^(1/4)*B*atan((-1)^(1/4)*tan(c + d*x)^(1/2)))/(8*a^3*d) + ((-1)^(1/4)*B*atanh((-1)^(1/4)*tan(c + d*x)^(1/2)))/(8*a^3*d)`

3.150
$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.150.1 Optimal result 1563
 3.150.2 Mathematica [A] (verified) 1564
 3.150.3 Rubi [A] (verified) 1564
 3.150.4 Maple [A] (verified) 1570
 3.150.5 Fricas [B] (verification not implemented) 1570
 3.150.6 Sympy [F] 1571
 3.150.7 Maxima [F(-2)] 1572
 3.150.8 Giac [A] (verification not implemented) 1572
 3.150.9 Mupad [B] (verification not implemented) 1573

3.150.1 Optimal result

Integrand size = 36, antiderivative size = 317

$$\begin{aligned} & \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\ &= \frac{\left(\frac{1}{16} + \frac{i}{16}\right) ((1+i)A+B) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}a^3d} \\ & \quad - \frac{\left(\frac{1}{16} + \frac{i}{16}\right) ((1+i)A+B) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}a^3d} \\ & \quad + \frac{(2iA+(1-i)B) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{32\sqrt{2}a^3d} \\ & \quad - \frac{(2iA+(1-i)B) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{32\sqrt{2}a^3d} \\ & \quad + \frac{(iA-B)\sqrt{\tan(c+dx)}}{6d(a+ia \tan(c+dx))^3} + \frac{(iA+2B)\sqrt{\tan(c+dx)}}{12ad(a+ia \tan(c+dx))^2} + \frac{B\sqrt{\tan(c+dx)}}{8d(a^3+ia^3 \tan(c+dx))} \end{aligned}$$

```
output (-1/32-1/32*I)*((1+I)*A+B)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/a^3/d*2^(1/2)
- (1/32+1/32*I)*((1+I)*A+B)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/a^3/d*2^(1/2)
+ 1/64*(2*I*A+(1-I)*B)*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a^3/d*2^(1/2)
- 1/64*(2*I*A+(1-I)*B)*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a^3/d*2^(1/2)
+ 1/6*(I*A-B)*tan(d*x+c)^(1/2)/d/(a+I*a*tan(d*x+c))^3+1/12*(I*A+2*B)*tan(d*x+c)^(1/2)/a/d/(a+I*a*tan(d*x+c))^2+1/8*B*tan(d*x+c)^(1/2)/d/(a^3+I*a^3*tan(d*x+c))
```

3.150.
$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.150.2 Mathematica [A] (verified)

Time = 5.01 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.56

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$= \frac{3(-1)^{3/4}(iA+B)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)\sec^3(c+dx)(\cos(3(c+dx))+i\sin(3(c+dx))) - 3\sqrt[4]{-1}}$$

input `Integrate[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])
^3,x]`

output `(3*(-1)^(3/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]
^3*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)]) - 3*(-1)^(1/4)*A*ArcTanh[(-1)^(
3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^3*(Cos[3*(c + d*x)] + I*Sin[3*(c + d
*x)]) - I*Sqrt[Tan[c + d*x]]*(-3*I + Tan[c + d*x])*(2*A - I*B + 3*B*Tan[c
+ d*x]))/(24*a^3*d*(-I + Tan[c + d*x])^3)`

3.150.3 Rubi [A] (verified)

Time = 1.16 (sec) , antiderivative size = 291, normalized size of antiderivative = 0.92, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.528$, Rules used = {3042, 4078, 27, 3042, 4079, 27, 3042, 4079, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$\downarrow 3042$$

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$\downarrow 4078$$

$$\frac{(-B+iA)\sqrt{\tan(c+dx)}}{6d(a+ia\tan(c+dx))^3} - \frac{\int \frac{a(iA-B)-a(5A-7iB)\tan(c+dx)}{2\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)^2} dx}{6a^2}$$

$$\downarrow 27$$

3.150. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{a(iA - B) - a(5A - 7iB)\tan(c + dx)}{\sqrt{\tan(c + dx)}(i \tan(c + dx)a + a)^2} dx}{12a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{a(iA - B) - a(5A - 7iB)\tan(c + dx)}{\sqrt{\tan(c + dx)}(i \tan(c + dx)a + a)^2} dx}{12a^2} \\
 & \quad \downarrow \text{4079} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{\int \frac{6(ia^2A - a^2(A - 2iB)\tan(c + dx))}{\sqrt{\tan(c + dx)}(i \tan(c + dx)a + a)} dx}{4a^2} - \frac{a(2B + iA)\sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{3 \int \frac{ia^2A - a^2(A - 2iB)\tan(c + dx)}{\sqrt{\tan(c + dx)}(i \tan(c + dx)a + a)} dx}{2a^2} - \frac{a(2B + iA)\sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{3 \int \frac{ia^2A - a^2(A - 2iB)\tan(c + dx)}{\sqrt{\tan(c + dx)}(i \tan(c + dx)a + a)} dx}{2a^2} - \frac{a(2B + iA)\sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))^2} \\
 & \quad \downarrow \text{4079} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{\int \frac{(2iA + B)a^3 + iB \tan(c + dx)a^3}{\sqrt{\tan(c + dx)}} dx}{2a^2} - \frac{a^2B\sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))} \right)}{2a^2} - \frac{a(2B + iA)\sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{\int \frac{(2iA + B)a^3 + iB \tan(c + dx)a^3}{\sqrt{\tan(c + dx)}} dx}{2a^2} - \frac{a^2B\sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))} \right)}{2a^2} - \frac{a(2B + iA)\sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))^2} \\
 & \quad \downarrow \text{4017} \\
 & \frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{\int \frac{a^3(2iA + B + iB \tan(c + dx))}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{a^2d} - \frac{a^2B\sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))} \right)}{2a^2} - \frac{a(2B + iA)\sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))^2} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.150. $\int \frac{\sqrt{\tan(c + dx)}(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^3} dx$

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{a \int \frac{2iA + B + iB \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{d} - \frac{a^2 B \sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))} \right)}{2a^2} - \frac{a(2B + iA)\sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))^2}}{12a^2}$$

↓ 1482

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{a \left(\frac{1}{2}(2iA + (1-i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \left(\frac{1}{2} + \frac{i}{2}\right)(B + (1+i)A) \int \frac{\tan(c + dx) + 1}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} \right)}{d} - \frac{a^2 B \sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))} \right)}{2a^2} - \frac{a(2B + iA)\sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))^2}}{12a^2}$$

↓ 1476

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{a \left(\frac{1}{2}(2iA + (1-i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \left(\frac{1}{2} + \frac{i}{2}\right)(B + (1+i)A) \left(\frac{1}{2} \int \frac{1}{\tan(c + dx) - \sqrt{2}\sqrt{\tan(c + dx)} + 1} d\sqrt{\tan(c + dx)} + \frac{1}{2} \int \frac{1}{\tan(c + dx) + \sqrt{2}\sqrt{\tan(c + dx)} + 1} d\sqrt{\tan(c + dx)} \right) \right)}{d} - \frac{a^2 B \sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))} \right)}{2a^2} - \frac{a(2B + iA)\sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))^2}}{12a^2}$$

↓ 1082

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{a \left(\frac{1}{2}(2iA + (1-i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \left(\frac{1}{2} + \frac{i}{2}\right)(B + (1+i)A) \left(\frac{\int \frac{1}{-\tan(c + dx) - 1} d(1 - \sqrt{2}\sqrt{\tan(c + dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c + dx) - 1} d(\sqrt{2}\sqrt{\tan(c + dx)} + 1)}{\sqrt{2}} \right) \right)}{d} - \frac{a^2 B \sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))} \right)}{2a^2} - \frac{a(2B + iA)\sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))^2}}{12a^2}$$

↓ 217

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{3 \left(\frac{a \left(\frac{1}{2}(2iA + (1-i)B) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \left(\frac{1}{2} + \frac{i}{2}\right)(B + (1+i)A) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c + dx)} + 1}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1 - \sqrt{2}\sqrt{\tan(c + dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) \right)}{d} - \frac{a^2 B \sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))} \right)}{2a^2} - \frac{a(2B + iA)\sqrt{\tan(c + dx)}}{d(a + ia \tan(c + dx))^2}}{12a^2}$$

↓ 1479

3.150. $\int \frac{\sqrt{\tan(c + dx)}(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^3} dx$

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{a \left(\frac{1}{2}(2iA + (1-i)B) \left(\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \left(\frac{1}{2} + \frac{i}{2}\right)(B+(1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{3d} \frac{2a^2}{12a^2}$$

25

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{a \left(\frac{1}{2}(2iA + (1-i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \left(\frac{1}{2} + \frac{i}{2}\right)(B+(1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{3d} \frac{2a^2}{12a^2}$$

27

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{a \left(\frac{1}{2}(2iA + (1-i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \left(\frac{1}{2} + \frac{i}{2}\right)(B+(1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{3d} \frac{2a^2}{12a^2}$$

1103

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} - \frac{a \left(\left(\frac{1}{2} + \frac{i}{2}\right)(B+(1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(2iA + (1-i)B) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx))}{2\sqrt{2}} \right) \right)}{3d} \frac{2a^2}{12a^2}$$

input `Int[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

3.150. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

```
output ((I*A - B)*Sqrt[Tan[c + d*x]]/(6*d*(a + I*a*Tan[c + d*x])^3) - ((a*(I*A
+ 2*B)*Sqrt[Tan[c + d*x]]/(d*(a + I*a*Tan[c + d*x])^2)) + (3*((a*((1/2 +
I/2)*((1 + I)*A + B)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) +
ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + (((2*I)*A + (1 - I)*B)*
(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 +
Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2))/d - (a^2*B*S
qrt[Tan[c + d*x]]/(d*(a + I*a*Tan[c + d*x])))/(2*a^2))/(12*a^2)
```

3.150.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

```
rule 1103 Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

```
rule 1476 Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.150.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.48

method	result
derivativedivides	$\frac{-iB \left(\tan^{\frac{5}{2}}(dx+c) \right) + \left(-\frac{10B}{3} - \frac{2iA}{3} \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) + (iB-2A)(\sqrt{\tan(dx+c)})}{8(\tan(dx+c)-i)^3} - \frac{A \arctan \left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}} \right)}{4(\sqrt{2-i\sqrt{2}})} + \frac{4 \left(\frac{A}{16} - \frac{iB}{16} \right) \arctan \left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}} \right)}{\sqrt{2+i\sqrt{2}}}$
default	$\frac{-iB \left(\tan^{\frac{5}{2}}(dx+c) \right) + \left(-\frac{10B}{3} - \frac{2iA}{3} \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) + (iB-2A)(\sqrt{\tan(dx+c)})}{8(\tan(dx+c)-i)^3} - \frac{A \arctan \left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}} \right)}{4(\sqrt{2-i\sqrt{2}})} + \frac{4 \left(\frac{A}{16} - \frac{iB}{16} \right) \arctan \left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}} \right)}{\sqrt{2+i\sqrt{2}}}$

input `int(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/d/a^3*(1/8*(-I*B*tan(d*x+c)^(5/2)+(-10/3*B-2/3*I*A)*tan(d*x+c)^(3/2)+(-2*A+I*B)*tan(d*x+c)^(1/2))/(tan(d*x+c)-I)^3-1/4*A/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))+4*(1/16*A-1/16*I*B)/(2^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))`

3.150.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 631 vs. 2(244) = 488.

Time = 0.26 (sec) , antiderivative size = 631, normalized size of antiderivative = 1.99

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx =$$

$$\left(3 a^3 d \sqrt{\frac{i A^2+2 A B-i B^2}{a^6 d^2}} e^{(6i dx+6i c)} \log \left(\frac{2 \left((a^3 d e^{(2i dx+2i c)}+a^3 d) \sqrt{\frac{-i e^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}+1}} \sqrt{\frac{i A^2+2 A B-i B^2}{a^6 d^2}}+(A-i B) e^{(2i dx+2i c)} \right)}{i A+B} \right) \right)$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="fracas")`

output

```
-1/96*(3*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)
*log(2*((a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) +
I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2)) + (
A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*a^3*d*sq
rt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(-2*((a^3*d*e
^(2*I*d*x + 2*I*c) + a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x
+ 2*I*c) + 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2)) - (A - I*B)*e^(2*I*
d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 24*a^3*d*sqrt(-1/64*I*A^2/
(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(1/8*(8*(a^3*d*e^(2*I*d*x + 2*I*c) + a^3
*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-1/6
4*I*A^2/(a^6*d^2)) + A)*e^(-2*I*d*x - 2*I*c)/(a^3*d)) + 24*a^3*d*sqrt(-1/6
4*I*A^2/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(-1/8*(8*(a^3*d*e^(2*I*d*x + 2*I
*c) + a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*
sqrt(-1/64*I*A^2/(a^6*d^2)) - A)*e^(-2*I*d*x - 2*I*c)/(a^3*d)) + 2*(2*(-I*
A - 2*B)*e^(6*I*d*x + 6*I*c) - (5*I*A + 4*B)*e^(4*I*d*x + 4*I*c) - (4*I*A
- B)*e^(2*I*d*x + 2*I*c) - I*A + B)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(
2*I*d*x + 2*I*c) + 1))*e^(-6*I*d*x - 6*I*c)/(a^3*d)
```

3.150.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{i \left(\int \frac{A\sqrt{\tan(c+dx)}}{\tan^3(c+dx)-3i \tan^2(c+dx)-3 \tan(c+dx)+i} dx + \int \frac{B \tan^{\frac{3}{2}}(c+dx)}{\tan^3(c+dx)-3i \tan^2(c+dx)-3 \tan(c+dx)+i} dx \right)}{a^3}$$

input `integrate(tan(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)`

output `I*(Integral(A*sqrt(tan(c + d*x))/(tan(c + d*x)**3 - 3*I*tan(c + d*x)**2 - 3*tan(c + d*x) + I), x) + Integral(B*tan(c + d*x)**(3/2)/(tan(c + d*x)**3 - 3*I*tan(c + d*x)**2 - 3*tan(c + d*x) + I), x))/a**3`

3.150.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.150.8 Giac [A] (verification not implemented)

Time = 0.68 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.41

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$= -\frac{(i+1)\sqrt{2}A\arctan\left(\left(\frac{1}{2}i+\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{16a^3d}$$

$$-\frac{(i-1)\sqrt{2}(A-iB)\arctan\left(-\left(\frac{1}{2}i-\frac{1}{2}\right)\sqrt{2}\sqrt{\tan(dx+c)}\right)}{16a^3d}$$

$$-\frac{3iB\tan(dx+c)^{\frac{5}{2}}+2iA\tan(dx+c)^{\frac{3}{2}}+10B\tan(dx+c)^{\frac{3}{2}}+6A\sqrt{\tan(dx+c)}-3iB\sqrt{\tan(dx+c)}}{24a^3d(\tan(dx+c)-i)^3}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output `-(1/16*I + 1/16)*sqrt(2)*A*arctan((1/2*I + 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^3*d) - (1/16*I - 1/16)*sqrt(2)*(A - I*B)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^3*d) - 1/24*(3*I*B*tan(d*x + c)^(5/2) + 2*I*A*tan(d*x + c)^(3/2) + 10*B*tan(d*x + c)^(3/2) + 6*A*sqrt(tan(d*x + c)) - 3*I*B*sqrt(tan(d*x + c)))/(a^3*d*(tan(d*x + c) - I)^3)`

3.150.9 Mupad [B] (verification not implemented)

Time = 7.87 (sec) , antiderivative size = 239, normalized size of antiderivative = 0.75

$$\begin{aligned}
& \int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx \\
&= \frac{\frac{B\sqrt{\tan(c+dx)}}{8a^3d} - \frac{B\tan(c+dx)^{5/2}}{8a^3d} + \frac{B\tan(c+dx)^{3/2}5i}{12a^3d}}{-\tan(c+dx)^3i - 3\tan(c+dx)^2 + \tan(c+dx)3i + 1} \\
&+ \frac{-\frac{A\tan(c+dx)^{3/2}}{12a^3d} + \frac{A\sqrt{\tan(c+dx)}i}{4a^3d}}{-\tan(c+dx)^3i - 3\tan(c+dx)^2 + \tan(c+dx)3i + 1} \\
&- \frac{(-1)^{1/4}A\operatorname{atan}\left((-1)^{1/4}\sqrt{\tan(c+dx)}\right)}{8a^3d} - \frac{(-1)^{1/4}A\operatorname{atanh}\left((-1)^{1/4}\sqrt{\tan(c+dx)}\right)}{8a^3d} \\
&- \frac{\sqrt{-\frac{1}{256}i}B\operatorname{atan}\left(16\sqrt{-\frac{1}{256}i}\sqrt{\tan(c+dx)}\right)2i}{a^3d}
\end{aligned}$$

input `int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3,x)`

output `((B*tan(c + d*x)^(1/2))/(8*a^3*d) + (B*tan(c + d*x)^(3/2)*5i)/(12*a^3*d) - (B*tan(c + d*x)^(5/2))/(8*a^3*d))/(tan(c + d*x)*3i - 3*tan(c + d*x)^2 - tan(c + d*x)^3*1i + 1) + ((A*tan(c + d*x)^(1/2)*1i)/(4*a^3*d) - (A*tan(c + d*x)^(3/2))/(12*a^3*d))/(tan(c + d*x)*3i - 3*tan(c + d*x)^2 - tan(c + d*x)^3*1i + 1) - ((-1)^(1/4)*A*atan((-1)^(1/4)*tan(c + d*x)^(1/2))/(8*a^3*d) - ((-1)^(1/4)*A*atanh((-1)^(1/4)*tan(c + d*x)^(1/2)))/(8*a^3*d) - ((-1i/256)^(1/2)*B*atan(16*(-1i/256)^(1/2)*tan(c + d*x)^(1/2))*2i)/(a^3*d)`

3.151
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3} dx$$

3.151.1 Optimal result 1574
 3.151.2 Mathematica [A] (verified) 1575
 3.151.3 Rubi [A] (verified) 1575
 3.151.4 Maple [A] (verified) 1581
 3.151.5 Fricas [B] (verification not implemented) 1581
 3.151.6 Sympy [F(-2)] 1582
 3.151.7 Maxima [F(-2)] 1582
 3.151.8 Giac [A] (verification not implemented) 1583
 3.151.9 Mupad [B] (verification not implemented) 1584

3.151.1 Optimal result

Integrand size = 36, antiderivative size = 315

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} dx \\ &= -\frac{((7 - 5i)A - 2iB) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{16\sqrt{2}a^3d} \\ &+ \frac{((7 - 5i)A - 2iB) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{16\sqrt{2}a^3d} \\ &- \frac{((7 + 5i)A - 2iB) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{32\sqrt{2}a^3d} \\ &+ \frac{((7 + 5i)A - 2iB) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{32\sqrt{2}a^3d} \\ &+ \frac{(A + iB)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3} + \frac{(4A + iB)\sqrt{\tan(c + dx)}}{12ad(a + ia \tan(c + dx))^2} + \frac{5A\sqrt{\tan(c + dx)}}{8d(a^3 + ia^3 \tan(c + dx))} \end{aligned}$$

```
output 1/32*((7-5*I)*A-2*I*B)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/a^3/d*2^(1/2)+1
/32*((7-5*I)*A-2*I*B)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/a^3/d*2^(1/2)-1/6
4*((7+5*I)*A-2*I*B)*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a^3/d*2^(1/2
)+1/64*((7+5*I)*A-2*I*B)*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/a^3/d*2
^(1/2)+1/6*(A+I*B)*tan(d*x+c)^(1/2)/d/(a+I*a*tan(d*x+c))^3+1/12*(4*A+I*B)*
tan(d*x+c)^(1/2)/a/d/(a+I*a*tan(d*x+c))^2+5/8*A*tan(d*x+c)^(1/2)/d/(a^3+I*
a^3*tan(d*x+c))
```

3.151.
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3} dx$$

3.151.2 Mathematica [A] (verified)

Time = 4.49 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.53

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} dx$$

$$= \frac{3\sqrt[4]{-1}(-6A + iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right) + \frac{-3\sqrt[4]{-1}(iA+B)\operatorname{arctan}\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)\sec^3(c+dx)(\cos(3(c+dx)))}{24a^3d}}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^3),x]`

output `(3*(-1)^(1/4)*(-6*A + I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + (-3*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^3*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)]) + Sqrt[Tan[c + d*x]]*((27*I)*A - 6*B - 2*(19*A + I*B)*Tan[c + d*x] - (15*I)*A*Tan[c + d*x]^2))/(-I + Tan[c + d*x])^3)/(24*a^3*d)`

3.151.3 Rubi [A] (verified)

Time = 1.13 (sec) , antiderivative size = 288, normalized size of antiderivative = 0.91, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.528$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} dx$$

↓ 4079

$$\frac{\int \frac{\alpha(11A - iB) - 5\alpha(iA - B)\tan(c + dx)}{2\sqrt{\tan(c + dx)}(i \tan(c + dx)a + a)^2} dx}{6a^2} + \frac{(A + iB)\sqrt{\tan(c + dx)}}{6d(a + ia \tan(c + dx))^3}$$

↓ 27

3.151. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
& \frac{\int \frac{a(11A-iB)-5a(iA-B)\tan(c+dx)}{\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)^2} dx}{12a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{a(11A-iB)-5a(iA-B)\tan(c+dx)}{\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)^2} dx}{12a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 4079 \\
& \frac{\int \frac{6(a^2(6A-iB)-a^2(4iA-B)\tan(c+dx))}{\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)} dx}{4a^2} + \frac{a(4A+iB)\sqrt{\tan(c+dx)}}{d(a+ia\tan(c+dx))^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 27 \\
& \frac{3 \int \frac{a^2(6A-iB)-a^2(4iA-B)\tan(c+dx)}{\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)} dx}{2a^2} + \frac{a(4A+iB)\sqrt{\tan(c+dx)}}{d(a+ia\tan(c+dx))^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 3042 \\
& \frac{3 \int \frac{a^2(6A-iB)-a^2(4iA-B)\tan(c+dx)}{\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)} dx}{2a^2} + \frac{a(4A+iB)\sqrt{\tan(c+dx)}}{d(a+ia\tan(c+dx))^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 4079 \\
& \frac{3 \left(\frac{\int \frac{a^3(7A-2iB)-5ia^3A\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{2a^2} + \frac{5a^2A\sqrt{\tan(c+dx)}}{d(a+ia\tan(c+dx))} \right)}{2a^2} + \frac{a(4A+iB)\sqrt{\tan(c+dx)}}{d(a+ia\tan(c+dx))^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 3042 \\
& \frac{3 \left(\frac{\int \frac{a^3(7A-2iB)-5ia^3A\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{2a^2} + \frac{5a^2A\sqrt{\tan(c+dx)}}{d(a+ia\tan(c+dx))} \right)}{2a^2} + \frac{a(4A+iB)\sqrt{\tan(c+dx)}}{d(a+ia\tan(c+dx))^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 4017 \\
& \frac{3 \left(\frac{\int \frac{a^3(-5i\tan(c+dx)A+7A-2iB)d\sqrt{\tan(c+dx)}}{\tan^2(c+dx)+1} dx}{a^2d} + \frac{5a^2A\sqrt{\tan(c+dx)}}{d(a+ia\tan(c+dx))} \right)}{2a^2} + \frac{a(4A+iB)\sqrt{\tan(c+dx)}}{d(a+ia\tan(c+dx))^2} + \\
& \quad \frac{12a^2}{6d(a+ia\tan(c+dx))^3} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{6d(a+ia\tan(c+dx))^3} \\
& \quad \downarrow 27
\end{aligned}$$

3.151. $\int \frac{A+B\tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3} dx$

$$3 \left(\frac{a \int \frac{-5i \tan(c+dx)A+7A-2iB}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{5a^2 A \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))}}{2a^2} \right) + \frac{a(4A+iB)\sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{6d(a+ia \tan(c+dx))^3}$$

↓ 1482

$$3 \left(\frac{a \left(\frac{1}{2} ((7+5i)A-2iB) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} ((7-5i)A-2iB) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) + \frac{5a^2 A \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))}}{2a^2} \right) + \frac{a(4A+iB)\sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))^2}$$

$$\frac{12a^2}{(A+iB)\sqrt{\tan(c+dx)}} \frac{1}{6d(a+ia \tan(c+dx))^3}$$

↓ 1476

$$3 \left(\frac{a \left(\frac{1}{2} ((7+5i)A-2iB) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} ((7-5i)A-2iB) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{2a^2} \right)$$

$$\frac{12a^2}{(A+iB)\sqrt{\tan(c+dx)}} \frac{1}{6d(a+ia \tan(c+dx))^3}$$

↓ 1082

$$3 \left(\frac{a \left(\frac{1}{2} ((7+5i)A-2iB) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} ((7-5i)A-2iB) \left(\int \frac{1}{-\tan(c+dx)-1} d \frac{(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \int \frac{1}{-\tan(c+dx)-1} d \frac{(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{2a^2} \right)$$

$$\frac{12a^2}{(A+iB)\sqrt{\tan(c+dx)}} \frac{1}{6d(a+ia \tan(c+dx))^3}$$

↓ 217

$$3 \left(\frac{a \left(\frac{1}{2} ((7+5i)A-2iB) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} ((7-5i)A-2iB) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) + \frac{5a^2 A \sqrt{\tan(c+dx)}}{d(a+ia \tan(c+dx))}}{2a^2} \right)$$

$$\frac{12a^2}{(A+iB)\sqrt{\tan(c+dx)}} \frac{1}{6d(a+ia \tan(c+dx))^3}$$

↓ 1479

3.151. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3} dx$

$$3 \left(\frac{a \left(\frac{1}{2}((7+5i)A-2iB) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}((7-5i)A-2iB) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right)}{d} \right)}{2a^2}$$

$$\frac{(A+iB)\sqrt{\tan(c+dx)}}{6d(a+ia\tan(c+dx))^3}$$

↓ 25

$$3 \left(\frac{a \left(\frac{1}{2}((7+5i)A-2iB) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}((7-5i)A-2iB) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right)}{d} \right)}{2a^2}$$

$$\frac{(A+iB)\sqrt{\tan(c+dx)}}{6d(a+ia\tan(c+dx))^3}$$

↓ 27

$$3 \left(\frac{a \left(\frac{1}{2}((7+5i)A-2iB) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2}((7-5i)A-2iB) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right)}{d} \right)}{2a^2}$$

$$\frac{(A+iB)\sqrt{\tan(c+dx)}}{6d(a+ia\tan(c+dx))^3}$$

↓ 1103

$$3 \left(\frac{\frac{5a^2 A \sqrt{\tan(c+dx)}}{d(a+ia\tan(c+dx))} + a \left(\frac{1}{2}((7-5i)A-2iB) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}((7+5i)A-2iB) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right)}{d} \right)}{2a^2}$$

$$\frac{(A+iB)\sqrt{\tan(c+dx)}}{6d(a+ia\tan(c+dx))^3}$$

3.151. $\int \frac{A+B\tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3} dx$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^3),x]`

output `((A + I*B)*Sqrt[Tan[c + d*x]])/(6*d*(a + I*a*Tan[c + d*x])^3) + ((a*(4*A + I*B)*Sqrt[Tan[c + d*x]])/(d*(a + I*a*Tan[c + d*x])^2) + (3*((a*(((7 - 5*I)*A - (2*I)*B)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]))/2 + (((7 + 5*I)*A - (2*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2))/d + (5*a^2*A*Sqrt[Tan[c + d*x]])/(d*(a + I*a*Tan[c + d*x]))/(2*a^2))/(12*a^2)`

3.151.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.151.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.50

method	result
derivativedivides	$-\frac{5iA \left(\tan^{\frac{5}{2}}(dx+c) \right) + \left(\frac{38A}{3} + \frac{2iB}{3} \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) + (-9iA+2B)(\sqrt{\tan(dx+c)})}{8(\tan(dx+c)-i)^3} - \frac{(6iA+B) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{4(\sqrt{2-i\sqrt{2}})} + \frac{4\left(\frac{iA}{16} + \frac{B}{16}\right)}{da^3}$
default	$-\frac{5iA \left(\tan^{\frac{5}{2}}(dx+c) \right) + \left(\frac{38A}{3} + \frac{2iB}{3} \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) + (-9iA+2B)(\sqrt{\tan(dx+c)})}{8(\tan(dx+c)-i)^3} - \frac{(6iA+B) \arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{4(\sqrt{2-i\sqrt{2}})} + \frac{4\left(\frac{iA}{16} + \frac{B}{16}\right)}{da^3}$

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/d/a^3*(-1/8*(5*I*A*tan(d*x+c)^(5/2)+(38/3*A+2/3*I*B)*tan(d*x+c)^(3/2)+(-9*I*A+2*B)*tan(d*x+c)^(1/2))/(tan(d*x+c)-I)^3-1/4*(6*I*A+B)/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))+4*(1/16*I*A+1/16*B)/(2^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))`

3.151.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 682 vs. 2(244) = 488.

Time = 0.26 (sec) , antiderivative size = 682, normalized size of antiderivative = 2.17

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} dx =$$

$$\left(3 a^3 d \sqrt{\frac{-i A^2 - 2 A B + i B^2}{a^6 d^2}} e^{(6i dx + 6i c)} \log \left(-\frac{2 \left((i a^3 d e^{(2i dx + 2i c)} + i a^3 d) \sqrt{\frac{-i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} + 1}} \sqrt{\frac{-i A^2 - 2 A B + i B^2}{a^6 d^2}} - (A - i B) e^{(2i dx + 2i c)} \right)}{i A + B} \right) \right)$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="fracas")`

output

```
-1/96*(3*a^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)
)*log(-2*((I*a^3*d*e^(2*I*d*x + 2*I*c) + I*a^3*d)*sqrt((-I*e^(2*I*d*x + 2*
I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^
2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*a
^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(-2*(
(-I*a^3*d*e^(2*I*d*x + 2*I*c) - I*a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)
/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2)) - (A
- I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*a^3*d*sqrt
((36*I*A^2 + 12*A*B - I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(1/8*((a^3*
d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d
*x + 2*I*c) + 1))*sqrt((36*I*A^2 + 12*A*B - I*B^2)/(a^6*d^2)) + 6*I*A + B)
*e^(-2*I*d*x - 2*I*c)/(a^3*d)) + 3*a^3*d*sqrt((36*I*A^2 + 12*A*B - I*B^2)/
(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(-1/8*((a^3*d*e^(2*I*d*x + 2*I*c) + a^3*
d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((36*I
*A^2 + 12*A*B - I*B^2)/(a^6*d^2)) - 6*I*A - B)*e^(-2*I*d*x - 2*I*c)/(a^3*d
)) - 2*(2*(10*A + I*B)*e^(6*I*d*x + 6*I*c) + (26*A + 5*I*B)*e^(4*I*d*x + 4
*I*c) + (7*A + 4*I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*sqrt((-I*e^(2*I*d*x +
2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*e^(-6*I*d*x - 6*I*c)/(a^3*d)
```

3.151.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+I*a*tan(d*x+c))**3,x)`

output `Exception raised: TypeError >> Invalid comparison of non-real -I`

3.151.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^3,x, algorith="maxima")`

3.151. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^3} dx$

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

3.151.8 Giac [A] (verification not implemented)

Time = 1.25 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.43

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} dx$$

$$= -\frac{(i + 1) \sqrt{2}(6iA + B) \arctan\left(\left(\frac{1}{2}i + \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{16 a^3 d}$$

$$+ \frac{(i - 1) \sqrt{2}(-iA - B) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{16 a^3 d}$$

$$- \frac{15iA \tan(dx + c)^{\frac{5}{2}} + 38A \tan(dx + c)^{\frac{3}{2}} + 2iB \tan(dx + c)^{\frac{3}{2}} - 27iA \sqrt{\tan(dx + c)} + 6B \sqrt{\tan(dx + c)}}{24 a^3 d (\tan(dx + c) - i)^3}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output `-(1/16*I + 1/16)*sqrt(2)*(6*I*A + B)*arctan((1/2*I + 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^3*d) + (1/16*I - 1/16)*sqrt(2)*(-I*A - B)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^3*d) - 1/24*(15*I*A*tan(d*x + c)^(5/2) + 38*A*tan(d*x + c)^(3/2) + 2*I*B*tan(d*x + c)^(3/2) - 27*I*A*sqrt(tan(d*x + c)) + 6*B*sqrt(tan(d*x + c)))/(a^3*d*(tan(d*x + c) - I)^3)`

3.151.9 Mupad [B] (verification not implemented)

Time = 7.87 (sec) , antiderivative size = 308, normalized size of antiderivative = 0.98

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} dx$$

$$= \frac{\frac{9A \sqrt{\tan(c+dx)}}{8a^3 d} - \frac{5A \tan(c+dx)^{5/2}}{8a^3 d} + \frac{A \tan(c+dx)^{3/2} 19i}{12a^3 d}}{-\tan(c + dx)^3 \operatorname{li} - 3 \tan(c + dx)^2 + \tan(c + dx) 3i + 1}$$

$$+ \frac{-\frac{B \tan(c+dx)^{3/2}}{12a^3 d} + \frac{B \sqrt{\tan(c+dx)} \operatorname{li}}{4a^3 d}}{-\tan(c + dx)^3 \operatorname{li} - 3 \tan(c + dx)^2 + \tan(c + dx) 3i + 1}$$

$$- \frac{(-1)^{1/4} B \operatorname{atan}\left((-1)^{1/4} \sqrt{\tan(c + dx)}\right)}{8a^3 d} - \frac{(-1)^{1/4} B \operatorname{atanh}\left((-1)^{1/4} \sqrt{\tan(c + dx)}\right)}{8a^3 d}$$

$$- \operatorname{atan}\left(\frac{8a^3 d \sqrt{\tan(c + dx)} \sqrt{\frac{A^2 9i}{64a^6 d^2}}}{3A}\right) \sqrt{\frac{A^2 9i}{64a^6 d^2}} 2i + \operatorname{atan}\left(\frac{16a^3 d \sqrt{\tan(c + dx)} \sqrt{-\frac{A^2 1i}{256a^6 d^2}}}{A}\right) \sqrt{-\frac{A^2 1i}{256a^6 d^2}}$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^3),x)`output `atan((16*a^3*d*tan(c + d*x)^(1/2)*(-(A^2*1i)/(256*a^6*d^2))^(1/2))/A)*(-(A^2*1i)/(256*a^6*d^2))^(1/2)*2i - atan((8*a^3*d*tan(c + d*x)^(1/2)*((A^2*9i)/(64*a^6*d^2))^(1/2))/(3*A))*((A^2*9i)/(64*a^6*d^2))^(1/2)*2i + ((9*A*tan(c + d*x)^(1/2))/(8*a^3*d) + (A*tan(c + d*x)^(3/2)*19i)/(12*a^3*d) - (5*A*tan(c + d*x)^(5/2))/(8*a^3*d))/(tan(c + d*x)*3i - 3*tan(c + d*x)^2 - tan(c + d*x)^3*1i + 1) + ((B*tan(c + d*x)^(1/2)*1i)/(4*a^3*d) - (B*tan(c + d*x)^(3/2))/(12*a^3*d))/(tan(c + d*x)*3i - 3*tan(c + d*x)^2 - tan(c + d*x)^3*1i + 1) - ((-1)^(1/4)*B*atan((-1)^(1/4)*tan(c + d*x)^(1/2))/(8*a^3*d) - ((-1)^(1/4)*B*atanh((-1)^(1/4)*tan(c + d*x)^(1/2)))/(8*a^3*d)`

$$3.152 \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$$

3.152.1 Optimal result	1585
3.152.2 Mathematica [C] (verified)	1586
3.152.3 Rubi [A] (verified)	1586
3.152.4 Maple [A] (verified)	1594
3.152.5 Fricas [B] (verification not implemented)	1594
3.152.6 Sympy [F(-2)]	1595
3.152.7 Maxima [F(-2)]	1596
3.152.8 Giac [A] (verification not implemented)	1596
3.152.9 Mupad [B] (verification not implemented)	1597

3.152.1 Optimal result

Integrand size = 36, antiderivative size = 364

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx \\ &= \frac{((30 + 28i)A - (7 - 5i)B) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{16\sqrt{2}a^3d} \\ & \quad - \frac{\left(\frac{1}{16} - \frac{i}{16}\right) ((1 + 29i)A - (6 + i)B) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}a^3d} \\ & \quad - \frac{\left(\frac{1}{32} - \frac{i}{32}\right) ((29 + i)A + (1 + 6i)B) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{\sqrt{2}a^3d} \\ & \quad + \frac{\left(\frac{1}{32} - \frac{i}{32}\right) ((29 + i)A + (1 + 6i)B) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{\sqrt{2}a^3d} \\ & \quad - \frac{5(6A + iB)}{8a^3d\sqrt{\tan(c + dx)}} + \frac{A + iB}{6d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} \\ & \quad + \frac{5A + 2iB}{12ad\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^2} + \frac{7(4A + iB)}{24d\sqrt{\tan(c + dx)}(a^3 + ia^3 \tan(c + dx))} \end{aligned}$$

3.152. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

output
$$\begin{aligned} & -1/32*((30+28*I)*A+(-7+5*I)*B)*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/a^3/d*2 \\ & ^{(1/2)}+(-1/32+1/32*I)*((1+29*I)*A-(6+I)*B)*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/ \\ & a^3/d*2^{(1/2)}+(-1/64+1/64*I)*((29+I)*A+(1+6*I)*B)*\ln(1-2^{(1/2)}*\tan(d*x \\ & +c)^{(1/2)}+\tan(d*x+c))/a^3/d*2^{(1/2)}+(1/64-1/64*I)*((29+I)*A+(1+6*I)*B)*\ln(\\ & 1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/a^3/d*2^{(1/2)}-5/8*(6*A+I*B)/a^3/d/t \\ & \tan(d*x+c)^{(1/2)}+1/6*(A+I*B)/d/\tan(d*x+c)^{(1/2)}/(a+I*a*\tan(d*x+c))^3+1/12*(\\ & 5*A+2*I*B)/a/d/\tan(d*x+c)^{(1/2)}/(a+I*a*\tan(d*x+c))^2+7/24*(4*A+I*B)/d/\tan(\\ & d*x+c)^{(1/2)}/(a^3+I*a^3*\tan(d*x+c)) \end{aligned}$$

3.152.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 3.03 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.53

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx$$

$$= \frac{\sec^3(c + dx) (2i \cos(c + dx)(7A + 4iB + (35A + 11iB) \cos(2(c + dx)) + (33iA - 9B) \sin(2(c + dx))) + 6$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^3),x]`

output
$$\begin{aligned} & (\text{Sec}[c + d*x]^3*((2*I)*\text{Cos}[c + d*x]*(7*A + (4*I)*B + (35*A + (11*I)*B)*\text{Cos} \\ & [2*(c + d*x)] + ((33*I)*A - 9*B)*\text{Sin}[2*(c + d*x)]) + 6*((-29*I)*A + 6*B)*\text{H} \\ & \text{ypergeometric2F1}[-1/2, 1, 1/2, (-I)*\text{Tan}[c + d*x]]*(\text{Cos}[3*(c + d*x)] + I*\text{Si} \\ & \text{n}[3*(c + d*x)]) + 6*(A - I*B)*\text{Hypergeometric2F1}[-1/2, 1, 1/2, I*\text{Tan}[c + d* \\ & x]]*((-I)*\text{Cos}[3*(c + d*x)] + \text{Sin}[3*(c + d*x)])))/(48*a^3*d*\text{Sqrt}[\text{Tan}[c + d* \\ & x]]*(-I + \text{Tan}[c + d*x])^3) \end{aligned}$$

3.152.3 Rubi [A] (verified)

Time = 1.37 (sec) , antiderivative size = 338, normalized size of antiderivative = 0.93, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.639$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4012, 25, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.152.
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$$

$$\begin{aligned}
& \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2}(a + ia \tan(c + dx))^3} dx \\
& \quad \downarrow \text{4079} \\
& \frac{\int \frac{a(13A+iB)-7a(iA-B) \tan(c+dx)}{2 \tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a)^2} dx}{6a^2} + \frac{A + iB}{6d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{a(13A+iB)-7a(iA-B) \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a)^2} dx}{12a^2} + \frac{A + iB}{6d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(13A+iB)-7a(iA-B) \tan(c+dx)}{\tan(c+dx)^{3/2}(i \tan(c+dx)a+a)^2} dx}{12a^2} + \frac{A + iB}{6d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} \\
& \quad \downarrow \text{4079} \\
& \frac{\int \frac{2(a^2(31A+4iB)-5a^2(5iA-2B) \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a)} dx}{4a^2} + \frac{a(5A+2iB)}{d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^2} + \\
& \quad \frac{12a^2}{A + iB} \\
& \quad \frac{A + iB}{6d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{a^2(31A+4iB)-5a^2(5iA-2B) \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a)} dx}{2a^2} + \frac{a(5A+2iB)}{d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^2} + \\
& \quad \frac{12a^2}{A + iB} \\
& \quad \frac{A + iB}{6d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a^2(31A+4iB)-5a^2(5iA-2B) \tan(c+dx)}{\tan(c+dx)^{3/2}(i \tan(c+dx)a+a)} dx}{2a^2} + \frac{a(5A+2iB)}{d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^2} + \\
& \quad \frac{12a^2}{A + iB} \\
& \quad \frac{A + iB}{6d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^3} \\
& \quad \downarrow \text{4079}
\end{aligned}$$

3.152. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
& \frac{\int \frac{3(5a^3(6A+iB)-7a^3(4iA-B)\tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx}{2a^2} + \frac{7a^2(4A+iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} + \frac{a(5A+2iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} + \\
& \frac{12a^2}{A+iB} \\
& \frac{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3}{27} \\
& \frac{3 \int \frac{5a^3(6A+iB)-7a^3(4iA-B)\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)} dx}{2a^2} + \frac{7a^2(4A+iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} + \frac{a(5A+2iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} + \\
& \frac{12a^2}{A+iB} \\
& \frac{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3}{3042} \\
& \frac{3 \int \frac{5a^3(6A+iB)-7a^3(4iA-B)\tan(c+dx)}{\tan(c+dx)^{3/2}} dx}{2a^2} + \frac{7a^2(4A+iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} + \frac{a(5A+2iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} + \\
& \frac{12a^2}{A+iB} \\
& \frac{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3}{4012} \\
& \frac{3 \left(\int -\frac{7(4iA-B)a^3+5(6A+iB)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx - \frac{10a^3(6A+iB)}{d\sqrt{\tan(c+dx)}} \right)}{2a^2} + \frac{7a^2(4A+iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} + \frac{a(5A+2iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} + \\
& \frac{12a^2}{A+iB} \\
& \frac{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3}{25} \\
& \frac{3 \left(-\int \frac{7(4iA-B)a^3+5(6A+iB)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx - \frac{10a^3(6A+iB)}{d\sqrt{\tan(c+dx)}} \right)}{2a^2} + \frac{7a^2(4A+iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} + \frac{a(5A+2iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} + \\
& \frac{12a^2}{A+iB} \\
& \frac{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3}{3042}
\end{aligned}$$

3.152. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^3} dx$

$$\frac{\frac{3 \left(- \int \frac{7(4iA-B)a^3 + 5(6A+iB)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx - \frac{10a^3(6A+iB)}{d\sqrt{\tan(c+dx)}} \right)}{2a^2} + \frac{7a^2(4A+iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} + \frac{a(5A+2iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} + \frac{12a^2}{A+iB}}{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3}$$

↓ 4017

$$\frac{\frac{3 \left(- \frac{2 \int \frac{a^3(7(4iA-B)+5(6A+iB)\tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{2a^2} - \frac{10a^3(6A+iB)}{d\sqrt{\tan(c+dx)}} \right)}{2a^2} + \frac{7a^2(4A+iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} + \frac{a(5A+2iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} + \frac{12a^2}{A+iB}}{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3}$$

↓ 27

$$\frac{\frac{3 \left(- \frac{2a^3 \int \frac{7(4iA-B)+5(6A+iB)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{2a^2} - \frac{10a^3(6A+iB)}{d\sqrt{\tan(c+dx)}} \right)}{2a^2} + \frac{7a^2(4A+iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} + \frac{a(5A+2iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} + \frac{12a^2}{A+iB}}{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3}$$

↓ 1482

$$\frac{\frac{3 \left(- \frac{2a^3 \left(\frac{1}{2}((30+28i)A-(7-5i)B) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \left(\frac{1}{2}-\frac{i}{2}\right)((29+i)A+(1+6i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) - \frac{10a^3(6A+iB)}{d\sqrt{\tan(c+dx)}} \right)}{2a^2} + \frac{7a^2(4A+iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} + \frac{a(5A+2iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} + \frac{12a^2}{A+iB}}{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3}$$

↓ 1476

$$\frac{\frac{3 \left(- \frac{2a^3 \left(\frac{1}{2}((30+28i)A-(7-5i)B) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \left(\frac{1}{2}-\frac{i}{2}\right)((29+i)A+(1+6i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) - \frac{10a^3(6A+iB)}{d\sqrt{\tan(c+dx)}} \right)}{2a^2} + \frac{7a^2(4A+iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} + \frac{a(5A+2iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^2} + \frac{12a^2}{A+iB}}{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3}$$

3.152. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

↓ 1082

$$3 \left(\frac{2a^3 \left(\frac{1}{2}((30+28i)A - (7-5i)B) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) - \left(\frac{1}{2}-\frac{i}{2}\right)((29+i)A+(1+6i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} \right)}{2a^2} \right)$$

$$\frac{A+iB}{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3} \qquad 12a^2$$

↓ 217

$$3 \left(\frac{2a^3 \left(\frac{1}{2}((30+28i)A - (7-5i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2}-\frac{i}{2}\right)((29+i)A+(1+6i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} \right)}{2a^2} \right)$$

$$\frac{A+iB}{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3} \qquad 12a^2$$

↓ 1479

$$3 \left(\frac{2a^3 \left(\frac{1}{2}((30+28i)A - (7-5i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2}-\frac{i}{2}\right)((29+i)A+(1+6i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}}}{2\sqrt{2}} \right) \right)}{d} \right)}{2a^2}$$

$$\frac{A+iB}{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3}$$

↓ 25

3.152. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\frac{2a^3 \left(\frac{1}{2}((30+28i)A - (7-5i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((29+i)A + (1+6i)B) \int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}+1} \frac{dx}{2\sqrt{2}} \right)}{3d} \frac{A+iB}{2a^2}$$

$$\frac{A+iB}{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3} \quad \downarrow \quad 27$$

$$\frac{2a^3 \left(\frac{1}{2}((30+28i)A - (7-5i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((29+i)A + (1+6i)B) \int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}+1} \frac{dx}{2\sqrt{2}} \right)}{3d} \frac{A+iB}{2a^2}$$

$$\frac{A+iB}{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3} \quad \downarrow \quad 1103$$

$$\frac{7a^2(4A+iB)}{d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))} + \frac{2a^3 \left(\frac{1}{2}((30+28i)A - (7-5i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)((29+i)A + (1+6i)B) \int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}+1} \frac{dx}{2\sqrt{2}} \right)}{3d} \frac{A+iB}{2a^2}$$

$$\frac{A+iB}{6d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^3}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^3),x]`

```
output (A + I*B)/(6*d*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^3) + ((a*(5*A + (
2*I)*B))/(d*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^2) + ((3*((-2*a^3*((
((30 + 28*I)*A - (7 - 5*I)*B)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sq
rt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])))/2 - (1/2 - I/2)*
((29 + I)*A + (1 + 6*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[
c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*
Sqrt[2])))/d - (10*a^3*(6*A + I*B))/(d*Sqrt[Tan[c + d*x]])))/(2*a^2) + (7
*a^2*(4*A + I*B))/(d*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x]))/(2*a^2))/
(12*a^2)
```

3.152.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; Fre
eQ[{a, b, c}, x]
```

```
rule 1103 Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

```
rule 1476 Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.152.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.49

method	result
derivativedivides	$\frac{i(14iA-5B)\left(\tan\frac{5}{2}(dx+c)\right)+\left(\frac{98iA}{3}-\frac{38B}{3}\right)\left(\tan\frac{3}{2}(dx+c)\right)+(9iB+20A)(\sqrt{\tan(dx+c)})}{8(\tan(dx+c)-i)^3} - \frac{(6iB+29A)\arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{4(\sqrt{2-i\sqrt{2}})}$
default	$\frac{i(14iA-5B)\left(\tan\frac{5}{2}(dx+c)\right)+\left(\frac{98iA}{3}-\frac{38B}{3}\right)\left(\tan\frac{3}{2}(dx+c)\right)+(9iB+20A)(\sqrt{\tan(dx+c)})}{8(\tan(dx+c)-i)^3} - \frac{(6iB+29A)\arctan\left(\frac{2(\sqrt{\tan(dx+c)})}{\sqrt{2-i\sqrt{2}}}\right)}{4(\sqrt{2-i\sqrt{2}})}$

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/d/a^3*(1/8*(I*(14*I*A-5*B)*tan(d*x+c)^(5/2)+(98/3*I*A-38/3*B)*tan(d*x+c)^(3/2)+(20*A+9*I*B)*tan(d*x+c)^(1/2))/(tan(d*x+c)-I)^3-1/4*(6*I*B+29*A)/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))-2*A/tan(d*x+c)^(1/2)+4*(-1/16*A+1/16*I*B)/(2^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))`

3.152.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 784 vs. 2(273) = 546.

Time = 0.28 (sec) , antiderivative size = 784, normalized size of antiderivative = 2.15

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx$$

$$= \frac{3 \left(a^3 de^{(8i dx + 8i c)} - a^3 de^{(6i dx + 6i c)} \right) \sqrt{\frac{i A^2 + 2 AB - i B^2}{a^6 d^2}} \log \left(\frac{2 \left((a^3 de^{(2i dx + 2i c)} + a^3 d) \sqrt{\frac{-i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} + 1}} \sqrt{\frac{i A^2 + 2 AB - i B^2}{a^6 d^2}} + (A + B) \right)}{i A + B} \right)}{1}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="fracas")`

```

output 1/96*(3*(a^3*d*e^(8*I*d*x + 8*I*c) - a^3*d*e^(6*I*d*x + 6*I*c))*sqrt((I*A^
2 + 2*A*B - I*B^2)/(a^6*d^2))*log(2*((a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*s
qrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*A^2 +
2*A*B - I*B^2)/(a^6*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2
*I*c)/(I*A + B)) - 3*(a^3*d*e^(8*I*d*x + 8*I*c) - a^3*d*e^(6*I*d*x + 6*I*c
))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2))*log(-2*((a^3*d*e^(2*I*d*x + 2*I
*c) + a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*
sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e
^(-2*I*d*x - 2*I*c)/(I*A + B)) + 3*(a^3*d*e^(8*I*d*x + 8*I*c) - a^3*d*e^(6
*I*d*x + 6*I*c))*sqrt((-841*I*A^2 + 348*A*B + 36*I*B^2)/(a^6*d^2))*log(1/8
*((a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e
^(2*I*d*x + 2*I*c) + 1))*sqrt((-841*I*A^2 + 348*A*B + 36*I*B^2)/(a^6*d^2))
+ 29*A + 6*I*B)*e^(-2*I*d*x - 2*I*c)/(a^3*d)) - 3*(a^3*d*e^(8*I*d*x + 8*I
*c) - a^3*d*e^(6*I*d*x + 6*I*c))*sqrt((-841*I*A^2 + 348*A*B + 36*I*B^2)/(a
^6*d^2))*log(-1/8*((a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt((-I*e^(2*I*d*x
+ 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-841*I*A^2 + 348*A*B + 36*
I*B^2)/(a^6*d^2)) - 29*A - 6*I*B)*e^(-2*I*d*x - 2*I*c)/(a^3*d)) - 2*(2*(73
*I*A - 10*B)*e^(8*I*d*x + 8*I*c) + 3*(35*I*A - 2*B)*e^(6*I*d*x + 6*I*c) -
(49*I*A - 19*B)*e^(4*I*d*x + 4*I*c) + 3*(-3*I*A + 2*B)*e^(2*I*d*x + 2*I*c)
- I*A + B)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)...

```

3.152.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \text{Exception raised: TypeError}$$

```

input integrate((A+B*tan(d*x+c))/tan(d*x+c)**(3/2)/(a+I*a*tan(d*x+c))**3,x)

```

```

output Exception raised: TypeError >> Invalid comparison of non-real -I

```

3.152.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.152.8 Giac [A] (verification not implemented)

Time = 1.68 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.46

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx$$

$$= -\frac{(i + 1) \sqrt{2}(29A + 6iB) \arctan\left(\left(\frac{1}{2}i + \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{16a^3d}$$

$$+ \frac{(i - 1) \sqrt{2}(A - iB) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{16a^3d} - \frac{2A}{a^3d \sqrt{\tan(dx + c)}}$$

$$- \frac{42iA \tan(dx + c)^{\frac{5}{2}} - 15B \tan(dx + c)^{\frac{5}{2}} + 98A \tan(dx + c)^{\frac{3}{2}} + 38iB \tan(dx + c)^{\frac{3}{2}} - 60iA \sqrt{\tan(dx + c)}}{24a^3d(-i \tan(dx + c) - 1)^3}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output `-(1/16*I + 1/16)*sqrt(2)*(29*A + 6*I*B)*arctan((1/2*I + 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^3*d) + (1/16*I - 1/16)*sqrt(2)*(A - I*B)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^3*d) - 2*A/(a^3*d*sqrt(tan(d*x + c))) - 1/24*(42*I*A*tan(d*x + c)^(5/2) - 15*B*tan(d*x + c)^(5/2) + 98*A*tan(d*x + c)^(3/2) + 38*I*B*tan(d*x + c)^(3/2) - 60*I*A*sqrt(tan(d*x + c)) + 27*B*sqrt(tan(d*x + c)))/(a^3*d*(-I*tan(d*x + c) - 1)^3)`

3.152.9 Mupad [B] (verification not implemented)

Time = 8.40 (sec) , antiderivative size = 389, normalized size of antiderivative = 1.07

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx \\
 &= 2 \operatorname{atanh} \left(\frac{16 a^3 d \sqrt{\tan(c + dx)} \sqrt{\frac{A^2 1i}{256 a^6 d^2}}}{A} \right) \sqrt{\frac{A^2 1i}{256 a^6 d^2}} \\
 &+ 2 \operatorname{atanh} \left(\frac{16 a^3 d \sqrt{\tan(c + dx)} \sqrt{-\frac{A^2 841i}{256 a^6 d^2}}}{29 A} \right) \sqrt{-\frac{A^2 841i}{256 a^6 d^2}} \\
 &- \operatorname{atan} \left(\frac{8 a^3 d \sqrt{\tan(c + dx)} \sqrt{\frac{B^2 9i}{64 a^6 d^2}}}{3 B} \right) \sqrt{\frac{B^2 9i}{64 a^6 d^2}} 2i \\
 &+ \operatorname{atan} \left(\frac{16 a^3 d \sqrt{\tan(c + dx)} \sqrt{-\frac{B^2 1i}{256 a^6 d^2}}}{B} \right) \sqrt{-\frac{B^2 1i}{256 a^6 d^2}} 2i \\
 &- \frac{\frac{2 A}{a^3 d} + \frac{A \tan(c+dx) 17i}{2 a^3 d} - \frac{121 A \tan(c+dx)^2}{12 a^3 d} - \frac{A \tan(c+dx)^3 15i}{4 a^3 d}}{\sqrt{\tan(c + dx)} + \tan(c + dx)^{3/2} 3i - 3 \tan(c + dx)^{5/2} - \tan(c + dx)^{7/2} 1i} \\
 &+ \frac{\frac{9 B \sqrt{\tan(c+dx)}}{8 a^3 d} - \frac{5 B \tan(c+dx)^{5/2}}{8 a^3 d} + \frac{B \tan(c+dx)^{3/2} 19i}{12 a^3 d}}{-\tan(c + dx)^3 1i - 3 \tan(c + dx)^2 + \tan(c + dx) 3i + 1}
 \end{aligned}$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + a*tan(c + d*x)*1i)^3),x)`

output `2*atanh((16*a^3*d*tan(c + d*x)^(1/2)*((A^2*1i)/(256*a^6*d^2))^(1/2))/A)*((A^2*1i)/(256*a^6*d^2))^(1/2) + 2*atanh((16*a^3*d*tan(c + d*x)^(1/2)*(-(A^2*841i)/(256*a^6*d^2))^(1/2))/(29*A))*(-(A^2*841i)/(256*a^6*d^2))^(1/2) - a*tan((8*a^3*d*tan(c + d*x)^(1/2)*((B^2*9i)/(64*a^6*d^2))^(1/2))/(3*B))*((B^2*9i)/(64*a^6*d^2))^(1/2)*2i + atan((16*a^3*d*tan(c + d*x)^(1/2)*(-(B^2*1i)/(256*a^6*d^2))^(1/2))/B)*(-(B^2*1i)/(256*a^6*d^2))^(1/2)*2i - ((2*A)/(a^3*d) + (A*tan(c + d*x)*17i)/(2*a^3*d) - (121*A*tan(c + d*x)^2)/(12*a^3*d) - (A*tan(c + d*x)^3*15i)/(4*a^3*d))/(tan(c + d*x)^(1/2) + tan(c + d*x)^(3/2)*3i - 3*tan(c + d*x)^(5/2) - tan(c + d*x)^(7/2)*1i) + ((9*B*tan(c + d*x)^(1/2))/(8*a^3*d) + (B*tan(c + d*x)^(3/2)*19i)/(12*a^3*d) - (5*B*tan(c + d*x)^(5/2))/(8*a^3*d))/(tan(c + d*x)*3i - 3*tan(c + d*x)^2 - tan(c + d*x)^3*1i + 1)`

3.153
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$$

3.153.1 Optimal result 1598
 3.153.2 Mathematica [C] (verified) 1599
 3.153.3 Rubi [A] (verified) 1600
 3.153.4 Maple [A] (verified) 1607
 3.153.5 Fricas [B] (verification not implemented) 1608
 3.153.6 Sympy [F(-1)] 1609
 3.153.7 Maxima [F(-2)] 1609
 3.153.8 Giac [A] (verification not implemented) 1609
 3.153.9 Mupad [B] (verification not implemented) 1610

3.153.1 Optimal result

Integrand size = 36, antiderivative size = 393

$$\begin{aligned} & \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx \\ &= \frac{\left(\frac{1}{16} - \frac{i}{16}\right) \left((76+i)A + (1+29i)B\right) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}a^3d} \\ & \quad - \frac{\left(\frac{1}{16} - \frac{i}{16}\right) \left((76+i)A + (1+29i)B\right) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}a^3d} \\ & \quad + \frac{\left((77+75i)A - (30-28i)B\right) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{32\sqrt{2}a^3d} \\ & \quad - \frac{\left(\frac{1}{32} - \frac{i}{32}\right) \left((1+76i)A - (29+i)B\right) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{\sqrt{2}a^3d} \\ & \quad - \frac{7(11A+4iB)}{24a^3d \tan^{\frac{3}{2}}(c+dx)} + \frac{15(5iA-2B)}{8a^3d \sqrt{\tan(c+dx)}} + \frac{A+iB}{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} \\ & \quad + \frac{2A+iB}{4ad \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} + \frac{3(5A+2iB)}{8d \tan^{\frac{3}{2}}(c+dx)(a^3+ia^3 \tan(c+dx))} \end{aligned}$$

output $(-1/32+1/32*I)*((76+I)*A+(1+29*I)*B)*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/a^{3/d*2^{(1/2)}}+(-1/32+1/32*I)*((76+I)*A+(1+29*I)*B)*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/a^{3/d*2^{(1/2)}}+1/64*((77+75*I)*A+(-30+28*I)*B)*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/a^{3/d*2^{(1/2)}}+(-1/64+1/64*I)*((1+76*I)*A-(29+I)*B)*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/a^{3/d*2^{(1/2)}}+15/8*(5*I*A-2*B)/a^{3/d/\tan(d*x+c)^{(1/2)}}-7/24*(11*A+4*I*B)/a^{3/d/\tan(d*x+c)^{(3/2)}}+1/6*(A+I*B)/d/\tan(d*x+c)^{(3/2)}/(a+I*a*\tan(d*x+c))^{3+1/4}*(2*A+I*B)/a/d/\tan(d*x+c)^{(3/2)}/(a+I*a*\tan(d*x+c))^{2+3/8}*(5*A+2*I*B)/d/\tan(d*x+c)^{(3/2)}/(a^{3+I*a^3*\tan(d*x+c)})$

3.153.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 3.10 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.49

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx$$

$$= \frac{\sec^3(c + dx) (2i \cos(c + dx)(8A + 5iB + (53A + 23iB) \cos(2(c + dx))) + (51iA - 21B) \sin(2(c + dx)))}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3}$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^3),x]`

output $(\text{Sec}[c + d*x]^{3*((2*I)*\text{Cos}[c + d*x]*(8*A + (5*I)*B + (53*A + (23*I)*B)*\text{Cos}[2*(c + d*x)] + ((51*I)*A - 21*B)*\text{Sin}[2*(c + d*x)])) + 2*((-76*I)*A + 29*B)*\text{Hypergeometric2F1}[-3/2, 1, -1/2, (-I)*\text{Tan}[c + d*x]]*(\text{Cos}[3*(c + d*x)] + I*\text{Sin}[3*(c + d*x)]) + 2*(A - I*B)*\text{Hypergeometric2F1}[-3/2, 1, -1/2, I*\text{Tan}[c + d*x]]*((-I)*\text{Cos}[3*(c + d*x)] + \text{Sin}[3*(c + d*x)])))/(48*a^3*d*\text{Tan}[c + d*x]^{(3/2)}*(-I + \text{Tan}[c + d*x])^3)$

3.153.3 Rubi [A] (verified)

Time = 1.59 (sec) , antiderivative size = 367, normalized size of antiderivative = 0.93, number of steps used = 25, number of rules used = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 3042, 4012, 25, 3042, 4012, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2}(a + ia \tan(c + dx))^3} dx \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{3(a(5A+iB)-3a(iA-B)\tan(c+dx))}{2 \tan^{\frac{5}{2}}(c+dx)(i \tan(c+dx)a+a)^2} dx}{6a^2} + \frac{A + iB}{6d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{a(5A+iB)-3a(iA-B)\tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(i \tan(c+dx)a+a)^2} dx}{4a^2} + \frac{A + iB}{6d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a(5A+iB)-3a(iA-B)\tan(c+dx)}{\tan(c+dx)^{5/2}(i \tan(c+dx)a+a)^2} dx}{4a^2} + \frac{A + iB}{6d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{2(a^2(16A+5iB)-7a^2(2iA-B)\tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)(i \tan(c+dx)a+a)} dx}{4a^2} + \frac{a(2A+iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} + \\
 & \quad \frac{4a^2}{A + iB} \\
 & \quad \frac{A + iB}{6d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.153. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{\int \frac{a^2(16A+5iB)-7a^2(2iA-B)\tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(i\tan(c+dx)a+a)} dx}{2a^2} + \frac{a(2A+iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} + \\
 & \frac{4a^2}{A+iB} \\
 & \frac{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3}{\phantom{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3}} \\
 & \quad \downarrow \quad 3042 \\
 & \frac{\int \frac{a^2(16A+5iB)-7a^2(2iA-B)\tan(c+dx)}{\tan(c+dx)^{5/2}(i \tan(c+dx)a+a)} dx}{2a^2} + \frac{a(2A+iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} + \\
 & \frac{4a^2}{A+iB} \\
 & \frac{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3}{\phantom{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3}} \\
 & \quad \downarrow \quad 4079 \\
 & \frac{\int \frac{7a^3(11A+4iB)-15a^3(5iA-2B)\tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)} dx}{2a^2} + \frac{3a^2(5A+2iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} + \frac{a(2A+iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} + \\
 & \frac{4a^2}{A+iB} \\
 & \frac{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3}{\phantom{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3}} \\
 & \quad \downarrow \quad 3042 \\
 & \frac{\int \frac{7a^3(11A+4iB)-15a^3(5iA-2B)\tan(c+dx)}{\tan(c+dx)^{5/2}} dx}{2a^2} + \frac{3a^2(5A+2iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} + \frac{a(2A+iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} + \\
 & \frac{4a^2}{A+iB} \\
 & \frac{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3}{\phantom{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3}} \\
 & \quad \downarrow \quad 4012 \\
 & \frac{\int -\frac{15(5iA-2B)a^3+7(11A+4iB)\tan(c+dx)a^3}{\tan^{\frac{3}{2}}(c+dx)} dx - \frac{14a^3(11A+4iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{3a^2(5A+2iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} + \frac{a(2A+iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} + \\
 & \frac{4a^2}{A+iB} \\
 & \frac{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3}{\phantom{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3}} \\
 & \quad \downarrow \quad 25
 \end{aligned}$$

3.153. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\frac{-\int \frac{15(5iA-2B)a^3+7(11A+4iB)\tan(c+dx)a^3}{\tan^{\frac{3}{2}}(c+dx)} dx - \frac{14a^3(11A+4iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{3a^2(5A+2iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} + \frac{a(2A+iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2}}{2a^2}}{2a^2} + \frac{4a^2}{A+iB} \frac{1}{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} +$$

↓ 3042

$$\frac{-\int \frac{15(5iA-2B)a^3+7(11A+4iB)\tan(c+dx)a^3}{\tan(c+dx)^{3/2}} dx - \frac{14a^3(11A+4iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{3a^2(5A+2iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} + \frac{a(2A+iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2}}{2a^2}}{2a^2} + \frac{4a^2}{A+iB} \frac{1}{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} +$$

↓ 4012

$$\frac{-\int \frac{7a^3(11A+4iB)-15a^3(5iA-2B)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{14a^3(11A+4iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{30a^3(-2B+5iA)}{d\sqrt{\tan(c+dx)}} + \frac{3a^2(5A+2iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} + \frac{a(2A+iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))}}{2a^2}}{2a^2} + \frac{4a^2}{A+iB} \frac{1}{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} +$$

↓ 3042

$$\frac{-\int \frac{7a^3(11A+4iB)-15a^3(5iA-2B)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{14a^3(11A+4iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{30a^3(-2B+5iA)}{d\sqrt{\tan(c+dx)}} + \frac{3a^2(5A+2iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} + \frac{a(2A+iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))}}{2a^2}}{2a^2} + \frac{4a^2}{A+iB} \frac{1}{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} +$$

↓ 4017

$$\frac{-2\int \frac{a^3(7(11A+4iB)-15(5iA-2B)\tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{14a^3(11A+4iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{30a^3(-2B+5iA)}{d\sqrt{\tan(c+dx)}} + \frac{3a^2(5A+2iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} + \frac{a(2A+iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))}}{2a^2}}{2a^2} + \frac{4a^2}{A+iB} \frac{1}{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} +$$

3.153. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

↓ 27

$$\frac{2a^3 \int \frac{7(11A+4iB)-15(5iA-2B)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{14a^3(11A+4iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{30a^3(-2B+5iA)}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{3a^2(5A+2iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} + \frac{a(2A+iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))}$$

$$\frac{A+iB}{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} \quad 4a^2$$

↓ 1482

$$\frac{2a^3 \left(\frac{1}{2}((77+75i)A-(30-28i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \left(\frac{1}{2}-\frac{i}{2}\right)((76+i)A+(1+29i)B) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) - \frac{14a^3(11A+4iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{30a^3(-2B+5iA)}{d\sqrt{\tan(c+dx)}}}{2a^2}$$

$$\frac{A+iB}{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} \quad 4a^2$$

↓ 1476

$$\frac{2a^3 \left(\frac{1}{2}((77+75i)A-(30-28i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \left(\frac{1}{2}-\frac{i}{2}\right)((76+i)A+(1+29i)B) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right) - \frac{14a^3(11A+4iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{30a^3(-2B+5iA)}{d\sqrt{\tan(c+dx)}}}{2a^2}$$

$$\frac{A+iB}{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} \quad 4a^2$$

↓ 1082

$$\frac{2a^3 \left(\frac{1}{2}((77+75i)A-(30-28i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \left(\frac{1}{2}-\frac{i}{2}\right)((76+i)A+(1+29i)B) \left(\int \frac{1}{-\tan(c+dx)-1} \frac{d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \int \frac{1}{-\tan(c+dx)-1} \frac{d(1+\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) - \frac{14a^3(11A+4iB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{30a^3(-2B+5iA)}{d\sqrt{\tan(c+dx)}}}{2a^2}$$

$$\frac{A+iB}{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} \quad 4a^2$$

↓ 217

3.153. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\frac{2a^3 \left(\frac{1}{2} ((77+75i)A - (30-28i)B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \left(\frac{1}{2} - \frac{i}{2} \right) ((76+i)A + (1+29i)B) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)}+1}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}} \right) \right)}{d}$$

$$\frac{2a^2}{2a^2}$$

$$\frac{4a^2}{2a^2}$$

$$\frac{A + iB}{6d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3}$$

↓ 1479

$$2a^3 \left(\frac{1}{2} ((77+75i)A - (30-28i)B) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \left(\frac{1}{2} - \frac{i}{2} \right) ((76+i)A + (1+29i)B) \right)$$

$$\frac{2a^2}{d}$$

$$\frac{2a^2}{2a^2}$$

$$\frac{2a^2}{2a^2}$$

$$\frac{A + iB}{6d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3}$$

↓ 25

$$2a^3 \left(\frac{1}{2} ((77+75i)A - (30-28i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \left(\frac{1}{2} - \frac{i}{2} \right) ((76+i)A + (1+29i)B) \right)$$

$$\frac{2a^2}{d}$$

$$\frac{2a^2}{2a^2}$$

$$\frac{2a^2}{2a^2}$$

$$\frac{A + iB}{6d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3}$$

↓ 27

$$2a^3 \left(\frac{1}{2} ((77+75i)A - (30-28i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)}+1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)} \right) + \left(\frac{1}{2} - \frac{i}{2} \right) ((76+i)A + (1+29i)B) \right)$$

$$\frac{2a^2}{d}$$

$$\frac{2a^2}{2a^2}$$

$$\frac{2a^2}{2a^2}$$

$$\frac{A + iB}{6d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3}$$

↓ 1103

3.153. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\frac{\frac{3a^2(5A+2iB)}{d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} + \frac{2a^3 \left(\left(\frac{1}{2} - \frac{i}{2} \right) ((76+i)A + (1+29i)B) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) + \frac{1}{2}((77+75i)A - (30-28i)B) \right)}{d}}{2a^2}}{A+iB}{6d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^3),x]`

output `(A + I*B)/(6*d*Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^3) + ((a*(2*A + I*B))/(d*Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^2) + (((-2*a^3*((1/2 - I/2)*((76 + I)*A + (1 + 29*I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + (((77 + 75*I)*A - (30 - 28*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2))/d - (14*a^3*(11*A + (4*I)*B))/(3*d*Tan[c + d*x]^(3/2)) + (30*a^3*((5*I)*A - 2*B))/(d*Sqrt[Tan[c + d*x]]))/(2*a^2) + (3*a^2*(5*A + (2*I)*B))/(d*Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])))/(2*a^2))/(4*a^2)`

3.153.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

$$3.153. \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$$

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] +
Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a
, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-
a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_
)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sq
rt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] &
& NeQ[c^2 + d^2, 0]`

```
rule 4079 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.153.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.50

method	result
derivativedivides	$\frac{i \left(\frac{27iA-14B}{3} \tan^{\frac{5}{2}}(dx+c) + \left(\frac{182iA}{3} - \frac{98B}{3} \right) \tan^{\frac{3}{2}}(dx+c) + (20iB+35A) \sqrt{\tan(dx+c)} \right)}{(\tan(dx+c)-i)^3} - \frac{2(29iB+76A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}}}{8 da^3}$
default	$\frac{i \left(\frac{27iA-14B}{3} \tan^{\frac{5}{2}}(dx+c) + \left(\frac{182iA}{3} - \frac{98B}{3} \right) \tan^{\frac{3}{2}}(dx+c) + (20iB+35A) \sqrt{\tan(dx+c)} \right)}{(\tan(dx+c)-i)^3} - \frac{2(29iB+76A) \arctan\left(\frac{2(\sqrt{\tan(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}}}{8 da^3}$

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^3,x,method=_RETUR
NVERBOSE)
```

```
output 1/d/a^3*(-1/8*I*((I*(27*I*A-14*B)*tan(d*x+c)^(5/2)+(182/3*I*A-98/3*B)*tan(
d*x+c)^(3/2)+(35*A+20*I*B)*tan(d*x+c)^(1/2))/(tan(d*x+c)-I)^3-2*(29*I*B+76
*A)/(2^(1/2)-I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2))))-2/
3*A/tan(d*x+c)^(3/2)-2*(-3*I*A+B)/tan(d*x+c)^(1/2)+4*(-1/16*I*A-1/16*B)/(2
^(1/2)+I*2^(1/2))*arctan(2*tan(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2))))
```

$$3.153. \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$$

3.153.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 875 vs. $2(296) = 592$.

Time = 0.29 (sec) , antiderivative size = 875, normalized size of antiderivative = 2.23

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \text{Too large to display}$$

```
input integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^3,x, algorith
thm="fricas")
```

```
output 1/96*(3*(a^3*d*e^(10*I*d*x + 10*I*c) - 2*a^3*d*e^(8*I*d*x + 8*I*c) + a^3*d
*e^(6*I*d*x + 6*I*c))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2))*log(-2*((I*
a^3*d*e^(2*I*d*x + 2*I*c) + I*a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^
(2*I*d*x + 2*I*c) + 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2)) - (A - I*
B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*(a^3*d*e^(10*I
*d*x + 10*I*c) - 2*a^3*d*e^(8*I*d*x + 8*I*c) + a^3*d*e^(6*I*d*x + 6*I*c))*
sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2))*log(-2*((-I*a^3*d*e^(2*I*d*x + 2*
I*c) - I*a^3*d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1
))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2)) - (A - I*B)*e^(2*I*d*x + 2*I*c
))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*(a^3*d*e^(10*I*d*x + 10*I*c) - 2*a^
3*d*e^(8*I*d*x + 8*I*c) + a^3*d*e^(6*I*d*x + 6*I*c))*sqrt((5776*I*A^2 - 44
08*A*B - 841*I*B^2)/(a^6*d^2))*log(-1/8*((a^3*d*e^(2*I*d*x + 2*I*c) + a^3*
d)*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((5776
*I*A^2 - 4408*A*B - 841*I*B^2)/(a^6*d^2)) + 76*I*A - 29*B)*e^(-2*I*d*x - 2
*I*c)/(a^3*d)) + 3*(a^3*d*e^(10*I*d*x + 10*I*c) - 2*a^3*d*e^(8*I*d*x + 8*I
*c) + a^3*d*e^(6*I*d*x + 6*I*c))*sqrt((5776*I*A^2 - 4408*A*B - 841*I*B^2)/
(a^6*d^2))*log(1/8*((a^3*d*e^(2*I*d*x + 2*I*c) + a^3*d)*sqrt((-I*e^(2*I*d*
x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((5776*I*A^2 - 4408*A*B - 8
41*I*B^2)/(a^6*d^2)) - 76*I*A + 29*B)*e^(-2*I*d*x - 2*I*c)/(a^3*d)) - 2*(2
*(174*A + 73*I*B)*e^(10*I*d*x + 10*I*c) - (144*A + 41*I*B)*e^(8*I*d*x + ...
```

3.153.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(5/2)/(a+I*a*tan(d*x+c))**3,x)`

output `Timed out`

3.153.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.153.8 Giac [A] (verification not implemented)

Time = 0.72 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.46

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx \\ &= -\frac{(i + 1) \sqrt{2}(-76i A + 29 B) \arctan\left(\left(\frac{1}{2}i + \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{16 a^3 d} \\ &+ \frac{(i - 1) \sqrt{2}(i A + B) \arctan\left(-\left(\frac{1}{2}i - \frac{1}{2}\right) \sqrt{2} \sqrt{\tan(dx + c)}\right)}{16 a^3 d} \\ &- \frac{225 A \tan(dx + c)^4 + 90i B \tan(dx + c)^4 - 598i A \tan(dx + c)^3 + 242 B \tan(dx + c)^3 - 489 A \tan(dx + c)^2 - 242 B \tan(dx + c)^2 - 24 \left(-i \tan(dx + c)\right)^{\frac{3}{2}} - \sqrt{\tan(dx + c)}}{24 \left(-i \tan(dx + c)\right)^{\frac{3}{2}} - \sqrt{\tan(dx + c)}} \end{aligned}$$

3.153. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output `-(1/16*I + 1/16)*sqrt(2)*(-76*I*A + 29*B)*arctan((1/2*I + 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^3*d) + (1/16*I - 1/16)*sqrt(2)*(I*A + B)*arctan(-(1/2*I - 1/2)*sqrt(2)*sqrt(tan(d*x + c)))/(a^3*d) - 1/24*(225*A*tan(d*x + c)^4 + 90*I*B*tan(d*x + c)^4 - 598*I*A*tan(d*x + c)^3 + 242*B*tan(d*x + c)^3 - 489*A*tan(d*x + c)^2 - 204*I*B*tan(d*x + c)^2 + 96*I*A*tan(d*x + c) - 48*B*tan(d*x + c) - 16*A)/((-I*tan(d*x + c)^(3/2) - sqrt(tan(d*x + c)))^3*a^3*d)`

3.153.9 Mupad [B] (verification not implemented)

Time = 9.98 (sec) , antiderivative size = 425, normalized size of antiderivative = 1.08

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx$$

$$= -\operatorname{atan}\left(\frac{16 a^3 d \sqrt{\tan(c + dx)} \sqrt{-\frac{A^2 1i}{256 a^6 d^2}}}{A}\right) \sqrt{-\frac{A^2 1i}{256 a^6 d^2}} 2i$$

$$+ \operatorname{atan}\left(\frac{4 a^3 d \sqrt{\tan(c + dx)} \sqrt{\frac{A^2 361i}{16 a^6 d^2}}}{19 A}\right) \sqrt{\frac{A^2 361i}{16 a^6 d^2}} 2i$$

$$- \frac{\frac{A 2i}{3 a^3 d} + \frac{4 A \tan(c+dx)}{a^3 d} + \frac{A \tan(c+dx)^2 163i}{8 a^3 d} - \frac{299 A \tan(c+dx)^3}{12 a^3 d} - \frac{A \tan(c+dx)^4 75i}{8 a^3 d}}{\tan(c + dx)^{9/2} - 3 \tan(c + dx)^{5/2} + \tan(c + dx)^{3/2} 1i - \tan(c + dx)^{7/2} 3i}$$

$$+ 2 \operatorname{atanh}\left(\frac{16 a^3 d \sqrt{\tan(c + dx)} \sqrt{\frac{B^2 1i}{256 a^6 d^2}}}{B}\right) \sqrt{\frac{B^2 1i}{256 a^6 d^2}} + 2 \operatorname{atanh}\left(\frac{16 a^3 d \sqrt{\tan(c + dx)} \sqrt{-\frac{B^2 841i}{256 a^6 d^2}}}{29 B}\right)$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + a*tan(c + d*x)*1i)^3),x)`

output

```
atan((4*a^3*d*tan(c + d*x)^(1/2)*((A^2*361i)/(16*a^6*d^2))^(1/2))/(19*A))*
((A^2*361i)/(16*a^6*d^2))^(1/2)*2i - atan((16*a^3*d*tan(c + d*x)^(1/2)*(-
A^2*1i)/(256*a^6*d^2))^(1/2))/A)*(-(A^2*1i)/(256*a^6*d^2))^(1/2)*2i - ((A*
2i)/(3*a^3*d) + (4*A*tan(c + d*x))/(a^3*d) + (A*tan(c + d*x)^2*163i)/(8*a^
3*d) - (299*A*tan(c + d*x)^3)/(12*a^3*d) - (A*tan(c + d*x)^4*75i)/(8*a^3*d
))/((tan(c + d*x)^(3/2)*1i - 3*tan(c + d*x)^(5/2) - tan(c + d*x)^(7/2)*3i +
tan(c + d*x)^(9/2)) + 2*atanh((16*a^3*d*tan(c + d*x)^(1/2)*((B^2*1i)/(256
*a^6*d^2))^(1/2))/B)*((B^2*1i)/(256*a^6*d^2))^(1/2) + 2*atanh((16*a^3*d*ta
n(c + d*x)^(1/2)*(-(B^2*841i)/(256*a^6*d^2))^(1/2))/(29*B))*(-(B^2*841i)/(
256*a^6*d^2))^(1/2) - ((2*B)/(a^3*d) + (B*tan(c + d*x)*17i)/(2*a^3*d) - (1
21*B*tan(c + d*x)^2)/(12*a^3*d) - (B*tan(c + d*x)^3*15i)/(4*a^3*d))/(tan(c
+ d*x)^(1/2) + tan(c + d*x)^(3/2)*3i - 3*tan(c + d*x)^(5/2) - tan(c + d*x
)^(7/2)*1i)
```

3.153.
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$$

3.154 $\int \tan^{\frac{3}{2}}(c+dx) \sqrt{a + ia \tan(c + dx)} (A+B \tan(c+dx)) dx$

3.154.1 Optimal result	1612
3.154.2 Mathematica [A] (verified)	1613
3.154.3 Rubi [A] (verified)	1613
3.154.4 Maple [B] (verified)	1617
3.154.5 Fricas [B] (verification not implemented)	1618
3.154.6 Sympy [F]	1619
3.154.7 Maxima [F]	1620
3.154.8 Giac [F(-2)]	1620
3.154.9 Mupad [F(-1)]	1620

3.154.1 Optimal result

Integrand size = 38, antiderivative size = 200

$$\int \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{(-1)^{3/4} \sqrt{a} (4iA + 7B) \arctan\left(\frac{(-1)^{3/4} \sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{4d}$$

$$+ \frac{(1+i)\sqrt{a}(iA+B) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$+ \frac{(4A-iB)\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}{4d} + \frac{B \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}}{2d}$$

output

```
1/4*(-1)^(3/4)*(4*I*A+7*B)*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I
*a*tan(d*x+c))^(1/2))*a^(1/2)/d+(1+I)*(I*A+B)*arctanh((1+I)*a^(1/2)*tan(d*
x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*a^(1/2)/d+1/4*(4*A-I*B)*tan(d*x+c)^(1
/2)*(a+I*a*tan(d*x+c))^(1/2)/d+1/2*B*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^(
3/2)/d
```

3.154.2 Mathematica [A] (verified)

Time = 7.01 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.39

$$\int \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{B \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d}$$

$$+ \frac{\frac{a(4A - iB) \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}}{2d} + \frac{i\sqrt{2}(-\frac{1}{4}ia^3(4A - iB) - \frac{1}{4}a^3(4iA + 7B)) \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right) \sqrt{\tan(c + dx)}}{d\sqrt{ia \tan(c + dx)}} - \frac{\sqrt[4]{-1}a^3(4iA + 7B)}{a}}{2a}$$

```
input Integrate[Tan[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]
```

```
output (B*Tan[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]])/(2*d) + ((a*(4*A - I*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(2*d) + (((-I)*Sqrt[2]*((-1/4*I)*a^3*(4*A - I*B) - (a^3*((4*I)*A + 7*B))/4)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]]*Sqrt[Tan[c + d*x]])/(d*Sqrt[I*a*Tan[c + d*x]]) - ((-1)^(1/4)*a^3*((4*I)*A + 7*B)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]]*Sqrt[1 + I*Tan[c + d*x]])/(2*d*Sqrt[a + I*a*Tan[c + d*x]]))/a)/(2*a)
```

3.154.3 Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.06, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4080, 27, 3042, 4080, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^{3/2} \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$\downarrow \text{4080}$$

$$\begin{aligned}
 & \frac{\int -\frac{1}{2}\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a(3aB-a(4A-iB)\tan(c+dx))}dx}{2a} + \\
 & \quad \frac{B\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}}{2d} \\
 & \quad \downarrow 27 \\
 & \quad \frac{B\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}}{2d} - \\
 & \quad \frac{\int \sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a(3aB-a(4A-iB)\tan(c+dx))}dx}{4a} \\
 & \quad \downarrow 3042 \\
 & \quad \frac{B\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}}{2d} - \\
 & \quad \frac{\int \sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a(3aB-a(4A-iB)\tan(c+dx))}dx}{4a} \\
 & \quad \downarrow 4080 \\
 & \quad \frac{B\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}}{2d} - \\
 & \quad \frac{\int \frac{\sqrt{i\tan(c+dx)a+a((4A-iB)a^2+(4iA+7B)\tan(c+dx)a^2)}}{2\sqrt{\tan(c+dx)}}dx}{a} - \frac{a(4A-iB)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} \\
 & \quad \downarrow 4a \\
 & \quad \downarrow 27 \\
 & \quad \frac{B\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}}{2d} - \\
 & \quad \frac{\int \frac{\sqrt{i\tan(c+dx)a+a((4A-iB)a^2+(4iA+7B)\tan(c+dx)a^2)}}{\sqrt{\tan(c+dx)}}dx}{2a} - \frac{a(4A-iB)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} \\
 & \quad \downarrow 4a \\
 & \quad \downarrow 3042 \\
 & \quad \frac{B\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}}{2d} - \\
 & \quad \frac{\int \frac{\sqrt{i\tan(c+dx)a+a((4A-iB)a^2+(4iA+7B)\tan(c+dx)a^2)}}{\sqrt{\tan(c+dx)}}dx}{2a} - \frac{a(4A-iB)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} \\
 & \quad \downarrow 4a \\
 & \quad \downarrow 4084 \\
 & \quad \frac{B\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}}{2d} - \\
 & \quad \frac{8a^2(A-iB)\int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx - a(4A-7iB)\int \frac{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx}{2a} - \frac{a(4A-iB)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} \\
 & \quad \downarrow 4a \\
 & \quad \downarrow 3042
 \end{aligned}$$

3.154. $\int \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$

$$\begin{aligned}
 & \frac{B \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{8a^2(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - a(4A-7iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} - \frac{a(4A-iB) \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)}}{d} \\
 & \qquad \qquad \qquad \downarrow 4027 \\
 & \frac{B \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{16ia^4(A-iB) \int \frac{1}{-2 \frac{\tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} - a(4A-7iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} - \frac{a(4A-iB) \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)}}{d} \\
 & \qquad \qquad \qquad \downarrow 218 \\
 & \frac{B \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{(8-8i)a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - a(4A-7iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} - \frac{a(4A-iB) \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)}}{d} \\
 & \qquad \qquad \qquad \downarrow 4082 \\
 & \frac{B \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{(8-8i)a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{a^3(4A-7iB) \int \frac{1}{\sqrt{\tan(c+dx)} \sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{2a} - \frac{a(4A-iB) \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)}}{d} \\
 & \qquad \qquad \qquad \downarrow 65 \\
 & \frac{B \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{(8-8i)a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^3(4A-7iB) \int \frac{1}{1 - \frac{ia \tan(c+dx)}{i \tan(c+dx)a+a}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{2a} - \frac{a(4A-iB) \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)}}{d} \\
 & \qquad \qquad \qquad \downarrow 216 \\
 & \frac{B \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{2\sqrt{-1}a^{5/2}(4A-7iB) \operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{(8-8i)a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{a(4A-iB) \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)}}{d}
 \end{aligned}$$

```
input Int[Tan[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]
```

3.154. $\int \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx$

output $(B \tan[c + dx]^{3/2} \sqrt{a + I a \tan[c + dx]}) / (2d) - (((2(-1)^{1/4} a^{5/2} (4A - (7I)B) \operatorname{ArcTan}[(1)^{3/4} \sqrt{a} \sqrt{\tan[c + dx]}) / \sqrt{a + I a \tan[c + dx]}]) / d + ((8 - 8I) a^{5/2} (A - I B) \operatorname{ArcTanh}[(1 + I) \sqrt{a} \sqrt{\tan[c + dx]}) / \sqrt{a + I a \tan[c + dx]}]) / d) / (2a) - (a(4A - I B) \sqrt{\tan[c + dx]} \sqrt{a + I a \tan[c + dx]}) / d) / (4a)$

3.154.3.1 Defintions of rubi rules used

rule 27 $\operatorname{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[Fx, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{MatchQ}[Fx, (b_*)(Gx_)] /; \operatorname{FreeQ}[b, x]$

rule 65 $\operatorname{Int}[1/(\sqrt{(b_*)(x_)} \sqrt{(c_*) + (d_*)(x_)}), x_Symbol] \rightarrow \operatorname{Simp}[2 \operatorname{Subst}[\operatorname{Int}[1/(b - dx^2), x], x, \sqrt{bx}/\sqrt{c + dx}], x] /; \operatorname{FreeQ}\{b, c, d\}, x] \ \&\& \ !\operatorname{GtQ}[c, 0]$

rule 216 $\operatorname{Int}[(a_*) + (b_*)(x_)^2]^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2] \operatorname{Rt}[b, 2])) \operatorname{ArcTan}[\operatorname{Rt}[b, 2](x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \ \operatorname{PosQ}[a/b] \ \&\& \ (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

rule 218 $\operatorname{Int}[(a_*) + (b_*)(x_)^2]^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[a/b, 2]/a) \operatorname{ArcTan}[x/\operatorname{Rt}[a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \ \operatorname{PosQ}[a/b]$

rule 3042 $\operatorname{Int}[u_, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$

rule 4027 $\operatorname{Int}[\sqrt{(a_*) + (b_*) \tan[(e_*) + (f_*)(x_)]}] / \sqrt{(c_*) + (d_*) \tan[(e_*) + (f_*)(x_)]}, x_Symbol] \rightarrow \operatorname{Simp}[-2a*(b/f) \operatorname{Subst}[\operatorname{Int}[1/(a*c - b*d - 2*a^2*x^2), x], x, \sqrt{c + d \tan[e + f*x]} / \sqrt{a + b \tan[e + f*x]}], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \operatorname{NeQ}[b*c - a*d, 0] \ \&\& \ \operatorname{EqQ}[a^2 + b^2, 0] \ \&\& \ \operatorname{NeQ}[c^2 + d^2, 0]$

rule 4080 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.154.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 699 vs. $2(159) = 318$.

Time = 0.26 (sec) , antiderivative size = 700, normalized size of antiderivative = 3.50

method	result
parts	$A \left(i\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \sqrt{ia} a+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia} \sqrt{-ia} \right)$
derivativedivides	$-\frac{\sqrt{a(1+i \tan(dx+c))} (\sqrt{\tan(dx+c)}) \left(4A\sqrt{ia} \ln \left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \sqrt{2} a-8iA\sqrt{a} \right)}{2d\sqrt{a \tan(dx+c)(1+i \tan(dx+c))}}$
default	$-\frac{\sqrt{a(1+i \tan(dx+c))} (\sqrt{\tan(dx+c)}) \left(4A\sqrt{ia} \ln \left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \sqrt{2} a-8iA\sqrt{a} \right)}{2d\sqrt{a \tan(dx+c)(1+i \tan(dx+c))}}$

3.154. $\int \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$


```
input int((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^(3/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

```
output 1/2*A/d*(I*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)
)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*(I*a)^(1/2)*a+2*(a*tan(d*x+c)
)*(1+I*tan(d*x+c))^(1/2)*(I*a)^(1/2)*(-I*a)^(1/2)+ln(1/2*(2*I*a*tan(d*x+c)
)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*
a)^(1/2)*a)*tan(d*x+c)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)/(a*tan(d*x+c)*(1+I
*tan(d*x+c)))^(1/2)/(I*a)^(1/2)/(-I*a)^(1/2)+1/8*B/d*(a*(1+I*tan(d*x+c)))^
(1/2)*tan(d*x+c)^(1/2)*(4*I*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+
c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*(I*a)^(1/2)
*a-4*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*ta
n(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*tan(d*x+c)*a+6*I*(a*ta
n(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)-4*(I*
a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2+7
*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(
1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d
*x+c)))^(1/2)*(I*a)^(1/2)*(-I*a)^(1/2)+7*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*ta
n(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a
)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)/(I*a)^(1/2)/(-I*a)^(1/2)/(-tan(d*x
+c)+I)
```

3.154.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 771 vs. $2(149) = 298$.

Time = 0.26 (sec) , antiderivative size = 771, normalized size of antiderivative = 3.86

$$\int \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx =$$

$$4\sqrt{2} \left(de^{(2i dx+2i c)} + d \right) \sqrt{-\frac{(iA^2+2AB-iB^2)a}{d^2}} \log \left(\frac{\left(i\sqrt{2}d\sqrt{-\frac{(iA^2+2AB-iB^2)a}{d^2}} e^{(i dx+i c)} + \sqrt{2}((iA+B)e^{(2i dx+2i c)} + iA+B) \right)}{iA+B} \right)$$

```
input integrate((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^(3/2)*(A+B*tan(d*x+c)),x, al
gorithm="fracas")
```

3.154. $\int \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx$

```

output -1/8*(4*sqrt(2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*
a/d^2)*log((I*sqrt(2)*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*e^(I*d*x + I*
c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x
+ 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)
))*e^(-I*d*x - I*c)/(I*A + B)) - 4*sqrt(2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqr
t(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*log((-I*sqrt(2)*d*sqrt(-(I*A^2 + 2*A*B -
I*B^2)*a/d^2)*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) +
I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) +
I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/(I*A + B)) - 2*sqrt(2)*((4
*A - 3*I*B)*e^(3*I*d*x + 3*I*c) + (4*A + I*B)*e^(I*d*x + I*c))*sqrt(a/(e^(
2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I
*c) + 1)) - (d*e^(2*I*d*x + 2*I*c) + d)*sqrt((-16*I*A^2 - 56*A*B + 49*I*B^
2)*a/d^2)*log((sqrt(2)*((4*I*A + 7*B)*e^(2*I*d*x + 2*I*c) + 4*I*A + 7*B)*s
qrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I
*d*x + 2*I*c) + 1)) + 2*I*d*sqrt((-16*I*A^2 - 56*A*B + 49*I*B^2)*a/d^2)*e^
(I*d*x + I*c))*e^(-I*d*x - I*c)/(4*I*A + 7*B)) + (d*e^(2*I*d*x + 2*I*c) +
d)*sqrt((-16*I*A^2 - 56*A*B + 49*I*B^2)*a/d^2)*log((sqrt(2)*((4*I*A + 7*B)
*e^(2*I*d*x + 2*I*c) + 4*I*A + 7*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt
((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) - 2*I*d*sqrt((-16
*I*A^2 - 56*A*B + 49*I*B^2)*a/d^2)*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(4...

```

3.154.6 Sympy [F]

$$\begin{aligned}
 & \int \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 &= \int \sqrt{ia (\tan(c + dx) - i)} (A + B \tan(c + dx)) \tan^{\frac{3}{2}}(c + dx) dx
 \end{aligned}$$

```
input integrate((a+I*a*tan(d*x+c))**(1/2)*tan(d*x+c)**(3/2)*(A+B*tan(d*x+c)),x)
```

```
output Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))*tan(c + d*x)**(
3/2), x)
```

3.154.7 Maxima [F]

$$\int \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{ia \tan(dx + c) + a} \tan(dx + c)^{\frac{3}{2}} dx$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)*tan(d*x + c)^(3/2), x)`

3.154.8 Giac [F(-2)]

Exception generated.

$$\int \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:The choice was done assuming 0=[0]Warning, replacing 0 by 15, a substitution variable should perhaps be purged.Warning`

3.154.9 Mupad [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^{3/2} (A + B \tan(c + dx)) \sqrt{a + a \tan(c + dx)} \operatorname{li} dx$$

input `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2),
x)`

output `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2),
x)`

3.155 $\int \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.155.1 Optimal result	1622
3.155.2 Mathematica [A] (verified)	1623
3.155.3 Rubi [A] (verified)	1623
3.155.4 Maple [B] (verified)	1627
3.155.5 Fricas [B] (verification not implemented)	1628
3.155.6 Sympy [F]	1629
3.155.7 Maxima [F]	1629
3.155.8 Giac [F(-2)]	1629
3.155.9 Mupad [B] (verification not implemented)	1630

3.155.1 Optimal result

Integrand size = 38, antiderivative size = 152

$$\int \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= -\frac{(-1)^{3/4} \sqrt{a} (2A - iB) \arctan\left(\frac{(-1)^{3/4} \sqrt{a} \sqrt{\tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right)}{d} - \frac{(1 + i) \sqrt{a} (A - iB) \operatorname{arctanh}\left(\frac{(1 + i) \sqrt{a} \sqrt{\tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right)}{d} + \frac{B \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}}{d}$$

output

```

-(-1)^(3/4)*(2*A-I*B)*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*a^(1/2)/d-(1+I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*a^(1/2)/d+B*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/d
    
```

3.155.2 Mathematica [A] (verified)

Time = 5.55 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.24

$$\int \sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$= \frac{\sqrt[4]{-1}a(2A-iB)\operatorname{arcsinh}\left(\sqrt[4]{-1}\sqrt{\tan(c+dx)}\right)\sqrt{1+i\tan(c+dx)} + \frac{aB(1+i\tan(c+dx))\tan(c+dx)-\sqrt{2}(A-iB)\operatorname{arctan}\left(\sqrt{\tan(c+dx)}\right)}{d\sqrt{a+ia\tan(c+dx)}}}{d\sqrt{a+ia\tan(c+dx)}}$$

input `Integrate[Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `((-1)^(1/4)*a*(2*A - I*B)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]]*Sqrt[1 + I*Tan[c + d*x]] + (a*B*(1 + I*Tan[c + d*x])*Tan[c + d*x] - Sqrt[2]*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/Sqrt[Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]])`

3.155.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.04, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4080, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow 3042$$

$$\int \sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow 4080$$

$$\frac{\int -\frac{\sqrt{i\tan(c+dx)a+a(aB-a(2A-iB)\tan(c+dx))}}{2\sqrt{\tan(c+dx)}}dx}{a} + \frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}$$

$$\downarrow 27$$

$$\begin{aligned}
 & \frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{\int \frac{\sqrt{i\tan(c+dx)a+a}(aB-a(2A-iB)\tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{2a} \\
 & \quad \downarrow 3042 \\
 & \frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{\int \frac{\sqrt{i\tan(c+dx)a+a}(aB-a(2A-iB)\tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{2a} \\
 & \quad \downarrow 4084 \\
 & \frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{2a(B+iA)\int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - (B+2iA)\int \frac{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} \\
 & \quad \downarrow 3042 \\
 & \frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{2a(B+iA)\int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - (B+2iA)\int \frac{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} \\
 & \quad \downarrow 4027 \\
 & \frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{4ia^3(B+iA)\int \frac{1}{\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}} - (B+2iA)\int \frac{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} \\
 & \quad \downarrow 218 \\
 & \frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{(2-2i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right) - (B+2iA)\int \frac{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} \\
 & \quad \downarrow 4082 \\
 & \frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{(2-2i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right) - a^2(B+2iA)\int \frac{1}{\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}} d \tan(c+dx)}{2a} \\
 & \quad \downarrow 65
 \end{aligned}$$

3.155. $\int \sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx)) dx$

$$\frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{(2-2i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} - \frac{2a^2(B+2iA)\int\frac{1}{1-\frac{ia\tan(c+dx)}{i\tan(c+dx)a+a}}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{d}$$

2a
↓ 216

$$\frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{2^4\sqrt{-1}a^{3/2}(B+2iA)\operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} + \frac{(2-2i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}$$

input `Int[Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `-1/2*((2*(-1)^(1/4)*a^(3/2)*((2*I)*A + B)*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d + ((2 - 2*I)*a^(3/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d)/a + (B*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/d`

3.155.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_)+(d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 216 `Int[((a_)+(b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 218 `Int[((a_)+(b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4080 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.155.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 576 vs. $2(123) = 246$.

Time = 0.16 (sec) , antiderivative size = 577, normalized size of antiderivative = 3.80

method	result
parts	$-\frac{A(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})a\left(i\sqrt{ia}\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)-\sqrt{ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))}\right)}{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})\left(iB\sqrt{ia}\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)a\tan(dx+c)\right)}$
derivativedivides	$-\frac{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})\left(iB\sqrt{ia}\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)a\tan(dx+c)\right)}{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})\left(iB\sqrt{ia}\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)a\tan(dx+c)\right)}$
default	$-\frac{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})\left(iB\sqrt{ia}\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)a\tan(dx+c)\right)}{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})\left(iB\sqrt{ia}\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)a\tan(dx+c)\right)}$

input `int(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_RETURVERBOSE)`

output
$$-\frac{1}{2}A/d\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))}^{1/2}a^{1/2}(I\sqrt{a})^{1/2}2^{1/2}\ln\left(\frac{2^{1/2}(-I\sqrt{a})^{1/2}(a\tan(dx+c)(1+i\tan(dx+c)))^{1/2}-I\sqrt{a+3a\tan(dx+c)}}{\tan(dx+c)+I}\right)-I\sqrt{a+3a\tan(dx+c)}}{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})\left(iB\sqrt{ia}\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)a\tan(dx+c)\right)}$$

3.155.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 673 vs. $2(116) = 232$.

Time = 0.27 (sec) , antiderivative size = 673, normalized size of antiderivative = 4.43

$$\int \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$= \frac{2\sqrt{2}B \sqrt{\frac{a}{e^{2i dx+2i c}+1}} \sqrt{\frac{-i e^{(2i dx+2i c)+i}}{e^{2i dx+2i c}+1}} e^{i dx+i c} - \sqrt{2}d \sqrt{\frac{(-i A^2-2 AB+i B^2)a}{d^2}} \log \left(\frac{\left(\sqrt{2}d \sqrt{\frac{(-i A^2-2 AB+i B^2)a}{d^2}} e^{i dx} \right)}{\dots} \right)}{\dots}$$

```
input integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output 1/2*(2*sqrt(2)*B*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*e^(I*d*x + I*c) - sqrt(2)*d*sqrt((-I*A^2 - 2*A*B + I*B^2)*a/d^2)*log((sqrt(2)*d*sqrt((-I*A^2 - 2*A*B + I*B^2)*a/d^2)*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/(I*A + B) + sqrt(2)*d*sqrt((-I*A^2 - 2*A*B + I*B^2)*a/d^2)*log(-(sqrt(2)*d*sqrt((-I*A^2 - 2*A*B + I*B^2)*a/d^2)*e^(I*d*x + I*c) - sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/(I*A + B) + d*sqrt((4*I*A^2 + 4*A*B - I*B^2)*a/d^2)*log((sqrt(2)*((2*I*A + B)*e^(2*I*d*x + 2*I*c) + 2*I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))) + 2*d*sqrt((4*I*A^2 + 4*A*B - I*B^2)*a/d^2)*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(2*I*A + B) - d*sqrt((4*I*A^2 + 4*A*B - I*B^2)*a/d^2)*log((sqrt(2)*((2*I*A + B)*e^(2*I*d*x + 2*I*c) + 2*I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))) - 2*d*sqrt((4*I*A^2 + 4*A*B - I*B^2)*a/d^2)*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(2*I*A + B))/d
```

3.155.6 Sympy [F]

$$\int \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$= \int \sqrt{ia(\tan(c+dx)-i)} (A+B \tan(c+dx)) \sqrt{\tan(c+dx)} dx$$

input `integrate(tan(d*x+c)**(1/2)*(a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))*sqrt(tan(c + d*x)), x)`

3.155.7 Maxima [F]

$$\int \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$= \int (B \tan(dx+c) + A) \sqrt{ia \tan(dx+c) + a} \sqrt{\tan(dx+c)} dx$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)*sqrt(tan(d*x + c)), x)`

3.155.8 Giac [F(-2)]

Exception generated.

$$\int \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx = \text{Exception raised: TypeError}$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:The choice was done assuming 0=[0]W
arning, replacing 0 by 32, a substitution variable should perhaps be purge
d.Warning

3.155.9 Mupad [B] (verification not implemented)

Time = 28.49 (sec) , antiderivative size = 2225, normalized size of antiderivative = 14.64

$$\int \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx = \text{Too large to display}$$

input `int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2),
x)`

output

$$\begin{aligned} & - ((B*\tan(c + d*x)^(3/2)*2i)/(d*((a + a*\tan(c + d*x)*1i)^(1/2) - a^(1/2))^(3 \\ & 3) + (2*B*\tan(c + d*x)^(1/2))/(a*d*((a + a*\tan(c + d*x)*1i)^(1/2) - a^(1/2) \\ &))) / (\tan(c + d*x)^2 / ((a + a*\tan(c + d*x)*1i)^(1/2) - a^(1/2))^4 - 1/a^2 + \\ & (\tan(c + d*x)*2i)/(a*((a + a*\tan(c + d*x)*1i)^(1/2) - a^(1/2))^2)) - ((-a \\ &)^(1/2)*\operatorname{atan}((A^4*(-a)^(21/2)*\tan(c + d*x)^(1/2)*(7168 - 7168i))/((a + a* \\ & \tan(c + d*x)*1i)^(1/2) - a^(1/2))*(A^4*a^10*3584i + B^4*a^10*512i - 4096*A \\ & *B^3*a^10 + 10240*A^3*B*a^10 - A^2*B^2*a^10*10240i - (3584*A^4*a^11*\tan(c \\ & + d*x))/((a + a*\tan(c + d*x)*1i)^(1/2) - a^(1/2))^2 - (512*B^4*a^11*\tan(c \\ & + d*x))/((a + a*\tan(c + d*x)*1i)^(1/2) - a^(1/2))^2 + (10240*A^2*B^2*a^11* \\ & \tan(c + d*x))/((a + a*\tan(c + d*x)*1i)^(1/2) - a^(1/2))^2 - (A*B^3*a^11*\tan(c \\ & + d*x)*4096i)/((a + a*\tan(c + d*x)*1i)^(1/2) - a^(1/2))^2 + (A^3*B*a^11* \\ & \tan(c + d*x)*10240i)/((a + a*\tan(c + d*x)*1i)^(1/2) - a^(1/2))^2)) + (B^ \\ & 4*(-a)^(21/2)*\tan(c + d*x)^(1/2)*(1024 - 1024i))/((a + a*\tan(c + d*x)*1i) \\ & ^{(1/2) - a^(1/2)}*(A^4*a^10*3584i + B^4*a^10*512i - 4096*A*B^3*a^10 + 1024 \\ & 0*A^3*B*a^10 - A^2*B^2*a^10*10240i - (3584*A^4*a^11*\tan(c + d*x))/((a + a* \\ & \tan(c + d*x)*1i)^(1/2) - a^(1/2))^2 - (512*B^4*a^11*\tan(c + d*x))/((a + a* \\ & \tan(c + d*x)*1i)^(1/2) - a^(1/2))^2 + (10240*A^2*B^2*a^11*\tan(c + d*x))/((\\ & a + a*\tan(c + d*x)*1i)^(1/2) - a^(1/2))^2 - (A*B^3*a^11*\tan(c + d*x)*4096i \\ &)/((a + a*\tan(c + d*x)*1i)^(1/2) - a^(1/2))^2 + (A^3*B*a^11*\tan(c + d*x)*1 \\ & 0240i)/((a + a*\tan(c + d*x)*1i)^(1/2) - a^(1/2))^2)) + (A*B^3*(-a)^(21/2)... \end{aligned}$$

$$3.156 \quad \int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.156.1 Optimal result	1631
3.156.2 Mathematica [A] (verified)	1631
3.156.3 Rubi [A] (verified)	1632
3.156.4 Maple [B] (verified)	1634
3.156.5 Fricas [B] (verification not implemented)	1635
3.156.6 Sympy [F]	1636
3.156.7 Maxima [F]	1637
3.156.8 Giac [F(-2)]	1637
3.156.9 Mupad [B] (verification not implemented)	1638

3.156.1 Optimal result

Integrand size = 38, antiderivative size = 112

$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

$$= -\frac{2(-1)^{3/4}\sqrt{a}B \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$- \frac{(1+i)\sqrt{a}(iA+B)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

```
output -2*(-1)^(3/4)*B*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*a^(1/2)/d-(1+I)*(I*A+B)*arctanh(((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*a^(1/2)/d
```

3.156.2 Mathematica [A] (verified)

Time = 2.11 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.26

$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

$$= \frac{a \left(\frac{\sqrt{2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)\sqrt{\tan(c+dx)}}{\sqrt{ia \tan(c+dx)}} + \frac{2^4\sqrt{-1}\operatorname{Barcsinh}\left(\frac{\sqrt[4]{-1}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)\sqrt{1+i \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}} \right)}{d}$$

3.156. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

input `Integrate[(Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `(a*((Sqrt[2]*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[Tan[c + d*x]])/Sqrt[I*a*Tan[c + d*x]] + (2*(-1)^(1/4)*B*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]])*Sqrt[1 + I*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d`

3.156.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
 & \quad \downarrow \text{4084} \\
 & (A - iB) \int \frac{\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx + \frac{iB \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \int \frac{\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx + \frac{iB \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx}{a} \\
 & \quad \downarrow \text{4027} \\
 & \frac{iB \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx}{a} - \frac{2ia^2(A - iB) \int \frac{1}{\frac{2 \tan(c + dx)a^2}{i \tan(c + dx)a + a} - ia} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{i \tan(c + dx)a + a}}}{d} \\
 & \quad \downarrow \text{218}
 \end{aligned}$$

3.156. $\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$

$$\begin{aligned}
& \frac{iB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{a} + \frac{(1-i)\sqrt{a}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} \\
& \quad \downarrow 4082 \\
& \frac{iaB \int \frac{1}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{d} + \frac{(1-i)\sqrt{a}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} \\
& \quad \downarrow 65 \\
& \frac{2iaB \int \frac{1}{1-\frac{ia \tan(c+dx)}{i \tan(c+dx)a+a}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} + \frac{(1-i)\sqrt{a}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} \\
& \quad \downarrow 216 \\
& \frac{(1-i)\sqrt{a}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2(-1)^{3/4}\sqrt{a}B \operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}
\end{aligned}$$

input `Int[(Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `(-2*(-1)^(3/4)*Sqrt[a]*B*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d + ((1 - I)*Sqrt[a]*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d`

3.156.3.1 Defintions of rubi rules used

rule 65 `Int[1/(Sqrt[(b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

3.156. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.156.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 468 vs. $2(89) = 178$.

Time = 0.14 (sec) , antiderivative size = 469, normalized size of antiderivative = 4.19

method	result
parts	$-\frac{iA\sqrt{2} \ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i}\right) a(\sqrt{\tan(dx+c)})\sqrt{a(1+i \tan(dx+c))}}{2d\sqrt{a \tan(dx+c)(1+i \tan(dx+c))}\sqrt{-ia}} - \frac{B(\sqrt{\tan(dx+c)})}{\sqrt{a(1+i \tan(dx+c))}}$
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))}(\sqrt{\tan(dx+c)})\left(iA\sqrt{ia}\sqrt{2} \ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i}\right)\tan(dx+c)}{\sqrt{a(1+i \tan(dx+c))}(\sqrt{\tan(dx+c)})\left(iA\sqrt{ia}\sqrt{2} \ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i}\right)\tan(dx+c)}\right)}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))}(\sqrt{\tan(dx+c)})\left(iA\sqrt{ia}\sqrt{2} \ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i}\right)\tan(dx+c)}{\sqrt{a(1+i \tan(dx+c))}(\sqrt{\tan(dx+c)})\left(iA\sqrt{ia}\sqrt{2} \ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i}\right)\tan(dx+c)}\right)}$

3.156.
$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

input `int((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x,method=_R
ETURNVERBOSE)`

output `-1/2*I*A/d*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)/(-I*a)^(1/2)-1/2*B/d*tan(d*x+c)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)*a*(I*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))-I*a)^(1/2)*2^(1/2)*tan(d*x+c)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))+2*I*(-I*a)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2)+a)/(I*a)^(1/2))/a*tan(d*x+c)^(1/2)/(I*a)^(1/2)/(-I*a)^(1/2)/(-tan(d*x+c)+I)`

3.156.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 539 vs. $2(84) = 168$.

Time = 0.27 (sec) , antiderivative size = 539, normalized size of antiderivative = 4.81

$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

$$= \frac{1}{2} \sqrt{2} \sqrt{-\frac{(iA^2+2AB-iB^2)a}{d^2}} \log \left(\frac{\left(i\sqrt{2}d\sqrt{-\frac{(iA^2+2AB-iB^2)a}{d^2}} e^{(idx+ic)} + \sqrt{2}((iA+B)e^{(2idx+2ic)} + iA+B) \right)}{iA+B} \right)$$

$$- \frac{1}{2} \sqrt{2} \sqrt{-\frac{(iA^2+2AB-iB^2)a}{d^2}} \log \left(\frac{\left(-i\sqrt{2}d\sqrt{-\frac{(iA^2+2AB-iB^2)a}{d^2}} e^{(idx+ic)} + \sqrt{2}((iA+B)e^{(2idx+2ic)} + iA+B) \right)}{iA+B} \right)$$

$$- \frac{1}{2} \sqrt{\frac{4iB^2a}{d^2}} \log \left(\frac{\left(\sqrt{2}(Be^{(2idx+2ic)} + B) \sqrt{\frac{a}{e^{(2idx+2ic)}+1}} \sqrt{\frac{-ie^{(2idx+2ic)}+i}{e^{(2idx+2ic)}+1}} + i\sqrt{\frac{4iB^2a}{d^2}} de^{(idx+ic)} \right) e^{(-idx-ic)}}{B} \right)$$

$$+ \frac{1}{2} \sqrt{\frac{4iB^2a}{d^2}} \log \left(\frac{\left(\sqrt{2}(Be^{(2idx+2ic)} + B) \sqrt{\frac{a}{e^{(2idx+2ic)}+1}} \sqrt{\frac{-ie^{(2idx+2ic)}+i}{e^{(2idx+2ic)}+1}} - i\sqrt{\frac{4iB^2a}{d^2}} de^{(idx+ic)} \right) e^{(-idx-ic)}}{B} \right)$$

3.156. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="fricas")`

output `1/2*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*log((I*sqrt(2)*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/(I*A + B)) - 1/2*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*log((-I*sqrt(2)*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/(I*A + B)) - 1/2*sqrt(4*I*B^2*a/d^2)*log((sqrt(2)*(B*e^(2*I*d*x + 2*I*c) + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + I*sqrt(4*I*B^2*a/d^2)*d*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/B) + 1/2*sqrt(4*I*B^2*a/d^2)*log((sqrt(2)*(B*e^(2*I*d*x + 2*I*c) + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) - I*sqrt(4*I*B^2*a/d^2)*d*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/B)`

3.156.6 Sympy [F]

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \int \frac{\sqrt{ia (\tan(c + dx) - i)}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

input `integrate((a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(1/2),x)`

output `Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))/sqrt(tan(c + d*x)), x)`

3.156.7 Maxima [F]

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)\sqrt{ia \tan(dx + c) + a}}{\sqrt{\tan(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)/sqrt(tan(d*x + c)), x)`

3.156.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:The choice was done assuming 0=[0]Warning, replacing 0 by -86, a substitution variable should perhaps be purged.Warnin`

3.156.9 Mupad [B] (verification not implemented)

Time = 13.42 (sec) , antiderivative size = 372, normalized size of antiderivative = 3.32

$$\begin{aligned}
& \int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx \\
&= \frac{B \sqrt{a} \ln \left(\frac{\sqrt{a} \sqrt{\tan(c+dx)}(2-2i)}{\sqrt{a+a \tan(c+dx)} li-\sqrt{a}} - \frac{a \tan(c+dx)}{(\sqrt{a+a \tan(c+dx)} li-\sqrt{a})^2} + 1i \right) \left(\frac{1}{2} + \frac{1}{2}i \right)}{d} \\
&+ \frac{\sqrt{2} B \sqrt{a} \ln \left(\sqrt{2}(1-i) + \frac{2 \sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+a \tan(c+dx)} li-\sqrt{a}} \right) (1+1i)}{d} \\
&- \frac{\sqrt{\frac{1}{2}i} B \sqrt{a} \ln \left(-\frac{a \tan(c+dx)}{(\sqrt{a+a \tan(c+dx)} li-\sqrt{a})^2} + \frac{2(-1)^{3/4} \sqrt{2} \sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+a \tan(c+dx)} li-\sqrt{a}} + 1i \right)}{d} \\
&- \frac{\sqrt{4i} B \sqrt{a} \ln \left((-1)^{3/4} + \frac{\sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+a \tan(c+dx)} li-\sqrt{a}} \right)}{d} \\
&+ \frac{2 \sqrt{\frac{1}{2}i} A \sqrt{-a} \operatorname{atanh} \left(\frac{2 \sqrt{\frac{1}{2}i} \sqrt{-a} \sqrt{\tan(c+dx)} (\sqrt{a+a \tan(c+dx)} li-\sqrt{a})}{a \tan(c+dx)-a li+\sqrt{a} \sqrt{a+a \tan(c+dx)} li li} \right)}{d}
\end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2))/tan(c + d*x)^(1/2),x)`

output `(B*a^(1/2)*log((a^(1/2)*tan(c + d*x)^(1/2)*(2 - 2i))/((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2)) - a^(1/2)) - (a*tan(c + d*x))/((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2))^2 + 1i)*(1/2 + 1i/2))/d + (2^(1/2)*B*a^(1/2)*log(2^(1/2)*(1 - 1i) + (2*a^(1/2)*tan(c + d*x)^(1/2))/((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2)))*(1 + 1i))/d - ((1i/2)^(1/2)*B*a^(1/2)*log((2*(-1)^(3/4)*2^(1/2)*a^(1/2)*tan(c + d*x)^(1/2))/((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2)) - (a*tan(c + d*x))/((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2))^2 + 1i))/d - (4i^(1/2)*B*a^(1/2)*log((-1)^(3/4) + (a^(1/2)*tan(c + d*x)^(1/2))/((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2))))/d + (2*(1i/2)^(1/2)*A*(-a)^(1/2)*atanh((2*(1i/2)^(1/2)*(-a)^(1/2)*tan(c + d*x)^(1/2)*((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2))))/(a*tan(c + d*x) - a*1i + a^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)*1i))/d`

$$3.157 \quad \int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.157.1 Optimal result	1639
3.157.2 Mathematica [A] (verified)	1639
3.157.3 Rubi [A] (verified)	1640
3.157.4 Maple [B] (verified)	1642
3.157.5 Fricas [B] (verification not implemented)	1642
3.157.6 Sympy [F]	1643
3.157.7 Maxima [F(-1)]	1643
3.157.8 Giac [F(-2)]	1644
3.157.9 Mupad [F(-1)]	1644

3.157.1 Optimal result

Integrand size = 38, antiderivative size = 90

$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

$$= \frac{(1+i)\sqrt{a}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}$$

```
output (1+I)*(A-I*B)*arctanh(((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*a^(1/2)/d-2*A*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(1/2)
```

3.157.2 Mathematica [A] (verified)

Time = 0.95 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.13

$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

$$= \frac{\sqrt{2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)\sqrt{ia \tan(c+dx)}-2A\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}$$

```
input Integrate[(Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]
```

3.157. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

output $(\text{Sqrt}[2]*(A - I*B)*\text{ArcTanh}[(\text{Sqrt}[2]*\text{Sqrt}[I*a*\text{Tan}[c + d*x]])/\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])*\text{Sqrt}[I*a*\text{Tan}[c + d*x]] - 2*A*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])/(d*\text{Sqrt}[\text{Tan}[c + d*x]])$

3.157.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx \\
 & \quad \downarrow \text{4081} \\
 & \frac{2 \int \frac{a(iA+B)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx}{a} - \frac{2A\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & (B + iA) \int \frac{\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx - \frac{2A\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & (B + iA) \int \frac{\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx - \frac{2A\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{4027} \\
 & \frac{2ia^2(B + iA) \int \frac{1}{\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} - \frac{2A\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{218} \\
 & \frac{(1 - i)\sqrt{a}(B + iA)\text{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2A\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}}
 \end{aligned}$$

3.157. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

input `Int[(Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `((1 - I)*Sqrt[a]*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d - (2*A*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]])`

3.157.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

3.157.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 277 vs. $2(75) = 150$.

Time = 0.16 (sec) , antiderivative size = 278, normalized size of antiderivative = 3.09

method	result
parts	$\frac{A \left(\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \tan(dx+c) a - 4 \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{-ia}}{2d\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{\tan(dx+c)}} \right)$
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} \left(iB\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}}{\tan(dx+c)+i} \right) a (\tan^2(dx+c)) + iA\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \right)}{\sqrt{a(1+i \tan(dx+c))} \left(iB\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}}{\tan(dx+c)+i} \right) a (\tan^2(dx+c)) + iA\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \right)}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))} \left(iB\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}}{\tan(dx+c)+i} \right) a (\tan^2(dx+c)) + iA\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \right)}{\sqrt{a(1+i \tan(dx+c))} \left(iB\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}}{\tan(dx+c)+i} \right) a (\tan^2(dx+c)) + iA\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \right)}$

input `int((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x,method=_R
ETURNVERBOSE)`

output
$$\frac{1/2*A/d*(2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*\tan(d*x+c)*a-4*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)*(-I*a)^{(1/2)}*(a*(1+I*\tan(d*x+c)))^{(1/2)/(-I*a)^{(1/2)}/(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)/\tan(d*x+c)^{(1/2)}-1/2*I*B/d*2^{(1/2)})*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*a*\tan(d*x+c)^{(1/2)}*(a*(1+I*\tan(d*x+c)))^{(1/2)}/(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)/(-I*a)^{(1/2)}}}$$

3.157.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 435 vs. $2(69) = 138$.

Time = 0.25 (sec) , antiderivative size = 435, normalized size of antiderivative = 4.83

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \sqrt{2}(de^{(2i dx + 2i c)} - d) \sqrt{-\frac{(-i A^2 - 2 AB + i B^2)a}{d^2}} \log \left(\frac{\left(\sqrt{2}d \sqrt{-\frac{(-i A^2 - 2 AB + i B^2)a}{d^2}} e^{(i dx + i c)} + \sqrt{2}((i A + B)e^{(2i dx + 2i c)} + i A + B) \right)}{i A + B} \right)$$

3.157.
$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="fricas")`

output `1/2*(sqrt(2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*log((sqrt(2)*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/(I*A + B) - sqrt(2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*log(-(sqrt(2)*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*e^(I*d*x + I*c) - sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/(I*A + B) - 4*sqrt(2)*(I*A*e^(3*I*d*x + 3*I*c) + I*A*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(2*I*d*x + 2*I*c) - d)`

3.157.6 Sympy [F]

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{\sqrt{ia (\tan(c + dx) - i)}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(3/2),x)`

output `Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))/tan(c + d*x)**(3/2), x)`

3.157.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="maxima")`

3.157. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

output Timed out

3.157.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[-74]Warning, replacing -74 by -81, a substitution vari

3.157.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(A + B \tan(c + dx)) \sqrt{a + a \tan(c + dx)} \operatorname{li}}{\tan(c + dx)^{3/2}} dx \end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2))/tan(c + d*x)^(3/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2))/tan(c + d*x)^(3/2), x)`

$$3.158 \quad \int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.158.1 Optimal result	1645
3.158.2 Mathematica [A] (verified)	1645
3.158.3 Rubi [A] (verified)	1646
3.158.4 Maple [B] (verified)	1649
3.158.5 Fricas [B] (verification not implemented)	1649
3.158.6 Sympy [F]	1650
3.158.7 Maxima [F(-1)]	1651
3.158.8 Giac [F(-2)]	1651
3.158.9 Mupad [F(-1)]	1651

3.158.1 Optimal result

Integrand size = 38, antiderivative size = 135

$$\begin{aligned} & \int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx \\ &= \frac{(1+i)\sqrt{a}(iA+B)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} \\ & \quad - \frac{2A\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2(iA+3B)\sqrt{a+ia \tan(c+dx)}}{3d\sqrt{\tan(c+dx)}} \end{aligned}$$

output

```
(1+I)*(I*A+B)*arctanh(((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*a^(1/2)/d-2/3*(I*A+3*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(1/2)-2/3*A*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(3/2)
```

3.158.2 Mathematica [A] (verified)

Time = 2.07 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.97

$$\begin{aligned} & \int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx \\ &= \frac{3\sqrt{2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \tan(c+dx)\sqrt{ia \tan(c+dx)} + 2\sqrt{a+ia \tan(c+dx)}(-A+(-iA}}{3d \tan^{\frac{3}{2}}(c+dx)} \end{aligned}$$

3.158. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

input `Integrate[(Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(3*Sqrt[2]*(I*A + B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Tan[c + d*x]*Sqrt[I*a*Tan[c + d*x]] + 2*Sqrt[a + I*a*Tan[c + d*x]]*(-A + ((-I)*A - 3*B)*Tan[c + d*x])/(3*d*Tan[c + d*x]^(3/2))`

3.158.3 Rubi [A] (verified)

Time = 0.65 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.05, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.237$, Rules used = {3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx \\
 & \quad \downarrow 4081 \\
 & \frac{2 \int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA+3B)-2aA \tan(c+dx))}{2 \tan^{\frac{3}{2}}(c+dx)} dx}{3a} - \frac{2A \sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA+3B)-2aA \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx}{3a} - \frac{2A \sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA+3B)-2aA \tan(c+dx))}{\tan(c+dx)^{3/2}} dx}{3a} - \frac{2A \sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \\
 & \quad \downarrow 4081
 \end{aligned}$$

3.158. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \frac{2 \int -\frac{3a^2(A-iB)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx - \frac{2a(3B+iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow 27 \\
 & \frac{-3a(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a(3B+iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow 3042 \\
 & \frac{-3a(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a(3B+iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow 4027 \\
 & \frac{6ia^3(A-iB) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} - \frac{2a(3B+iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow 218 \\
 & \frac{(3-3i)a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) - \frac{2a(3B+iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)}
 \end{aligned}$$

input `Int[(Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(-2*A*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) + (((-3 + 3*I)*a^(3/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a*(I*A + 3*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a)`

3.158. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

3.158.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2)), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

3.158.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 353 vs. 2(110) = 220.

Time = 0.14 (sec) , antiderivative size = 354, normalized size of antiderivative = 2.62

method	result
parts	$A \left(3i\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i} \right) (\tan^2(dx+c)) a - 4i \tan(dx+c) \sqrt{-ia} \sqrt{a \tan(dx+c)}$
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} \left(3iA\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i} \right) a (\tan^3(dx+c)) - 12B\sqrt{-ia}}{6d \tan(dx+c)^{\frac{3}{2}} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))}}$
default	$-\frac{\sqrt{a(1+i \tan(dx+c))} \left(3iA\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i} \right) a (\tan^3(dx+c)) - 12B\sqrt{-ia}}{\dots}$

```
input int((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x,method=_R
ETURNVERBOSE)
```

```
output 1/6*A/d/tan(d*x+c)^(3/2)*(3*I*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*
x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*tan(d*x+c
)^2*a-4*I*tan(d*x+c)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-4*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2))*(a*(1+I*tan(d*x+c)))^(
1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)/(-I*a)^(1/2)+1/2*B/d*(2^(1/2)*1
n((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*ta
n(d*x+c))/(tan(d*x+c)+I))*tan(d*x+c)*a-4*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(
1/2)*(-I*a)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)/(-I*a)^(1/2)/(a*tan(d*x+c)*(
1+I*tan(d*x+c)))^(1/2)/tan(d*x+c)^(1/2)
```

3.158.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 483 vs. 2(102) = 204.

Time = 0.25 (sec) , antiderivative size = 483, normalized size of antiderivative = 3.58

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx =$$

$$3\sqrt{2}(de^{(4i dx+4i c)} - 2de^{(2i dx+2i c)} + d)\sqrt{-\frac{(i A^2+2 AB-i B^2)a}{d^2}} \log \left(\frac{\left(i\sqrt{2}d\sqrt{-\frac{(i A^2+2 AB-i B^2)a}{d^2}}e^{(i dx+i c)} + \sqrt{2}((i A+B \right)}{\dots} \right)$$

3.158. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="fricas")`

output `-1/6*(3*sqrt(2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*log((I*sqrt(2)*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/(I*A + B)) - 3*sqrt(2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*log((-I*sqrt(2)*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/(I*A + B)) - 4*sqrt(2)*((2*A - 3*I*B)*e^(5*I*d*x + 5*I*c) + 2*A*e^(3*I*d*x + 3*I*c) + 3*I*B*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)`

3.158.6 Sympy [F]

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{\sqrt{ia (\tan(c + dx) - i)}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

input `integrate((a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(5/2),x)`

output `Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))/tan(c + d*x)**(5/2), x)`

3.158.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="maxima")`

output `Timed out`

3.158.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[-75]Warning, replacing -75 by -19, a substitution vari`

3.158.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(A + B \tan(c + dx)) \sqrt{a + a \tan(c + dx)} \operatorname{li}}{\tan(c + dx)^{5/2}} dx \end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2))/tan(c + d*x)^(5/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2))/tan(c + d*x)^(5/2), x)`

3.158.
$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.159
$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

3.159.1 Optimal result 1653
 3.159.2 Mathematica [A] (verified) 1654
 3.159.3 Rubi [A] (verified) 1654
 3.159.4 Maple [B] (verified) 1658
 3.159.5 Fricas [B] (verification not implemented) 1659
 3.159.6 Sympy [F] 1660
 3.159.7 Maxima [F(-1)] 1660
 3.159.8 Giac [F(-2)] 1660
 3.159.9 Mupad [F(-1)] 1661

3.159.1 Optimal result

Integrand size = 38, antiderivative size = 178

$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

$$= -\frac{(1+i)\sqrt{a}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)}$$

$$- \frac{2(iA+5B)\sqrt{a+ia \tan(c+dx)}}{15d \tan^{\frac{3}{2}}(c+dx)} + \frac{2(13A-5iB)\sqrt{a+ia \tan(c+dx)}}{15d \sqrt{\tan(c+dx)}}$$

output `(-1-I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*a^(1/2)/d+2/15*(13*A-5*I*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(1/2)-2/5*A*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(5/2)-2/15*(I*A+5*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(3/2)`

3.159.2 Mathematica [A] (verified)

Time = 6.37 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.87

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx =$$

$$i \left(\frac{15\sqrt{2}a(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \tan^3(c+dx)}{\sqrt{ia \tan(c+dx)}} + 2\sqrt{a + ia \tan(c + dx)}(-3iA + (A - 5iB) \tan(c + dx)) \right)$$

$$15d \tan^{\frac{5}{2}}(c + dx)$$

input `Integrate[(Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `((-1/15*I)*((15*Sqrt[2]*a*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])]/Sqrt[a + I*a*Tan[c + d*x]])*Tan[c + d*x]^3)/Sqrt[I*a*Tan[c + d*x]] + 2*Sqrt[a + I*a*Tan[c + d*x]]*((-3*I)*A + (A - (5*I)*B)*Tan[c + d*x] + ((13*I)*A + 5*B)*Tan[c + d*x]^2))/(d*Tan[c + d*x]^(5/2))`

3.159.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.10, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {3042, 4081, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan(c + dx)^{7/2}} dx$$

↓ 4081

$$\frac{2 \int \frac{\sqrt{ia \tan(c+dx)a+a(a(iA+5B)-4aA \tan(c+dx))}}{2 \tan^{\frac{5}{2}}(c+dx)} dx}{5a} - \frac{2A\sqrt{a + ia \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)}$$

↓ 27

3.159. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$\begin{array}{c}
\frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA+5B)-4aA \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx}{5a} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
\downarrow 3042 \\
\frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA+5B)-4aA \tan(c+dx))}{\tan(c+dx)^{5/2}} dx}{5a} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
\downarrow 4081 \\
\frac{2 \int -\frac{\sqrt{i \tan(c+dx)a+a}((13A-5iB)a^2+2(iA+5B) \tan(c+dx)a^2)}{2 \tan^{\frac{3}{2}}(c+dx)} dx}{3a} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
\hline
\frac{5a}{2A\sqrt{a+ia \tan(c+dx)}} \\
\frac{2A\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
\downarrow 27 \\
\frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((13A-5iB)a^2+2(iA+5B) \tan(c+dx)a^2)}{\tan^{\frac{3}{2}}(c+dx)} dx}{3a} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
\hline
\frac{5a}{2A\sqrt{a+ia \tan(c+dx)}} \\
\frac{2A\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
\downarrow 3042 \\
\frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((13A-5iB)a^2+2(iA+5B) \tan(c+dx)a^2)}{\tan(c+dx)^{3/2}} dx}{3a} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
\hline
\frac{5a}{2A\sqrt{a+ia \tan(c+dx)}} \\
\frac{2A\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
\downarrow 4081 \\
\frac{2 \int \frac{15a^3(iA+B)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx}{3a} - \frac{2a^2(13A-5iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
\hline
\frac{5a}{2A\sqrt{a+ia \tan(c+dx)}} \\
\frac{2A\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
\downarrow 27
\end{array}$$

3.159. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \frac{15a^2(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(13A-5iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
 & \frac{5a}{2A\sqrt{a+ia \tan(c+dx)}} \\
 & \frac{5d \tan^{\frac{5}{2}}(c+dx)}{5d \tan^{\frac{5}{2}}(c+dx)} \\
 & \downarrow \text{3042} \\
 & \frac{15a^2(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(13A-5iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
 & \frac{5a}{2A\sqrt{a+ia \tan(c+dx)}} \\
 & \frac{5d \tan^{\frac{5}{2}}(c+dx)}{5d \tan^{\frac{5}{2}}(c+dx)} \\
 & \downarrow \text{4027} \\
 & \frac{30ia^4(B+iA) \int \frac{1}{\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} - \frac{2a^2(13A-5iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
 & \frac{5a}{2A\sqrt{a+ia \tan(c+dx)}} \\
 & \frac{5d \tan^{\frac{5}{2}}(c+dx)}{5d \tan^{\frac{5}{2}}(c+dx)} \\
 & \downarrow \text{218} \\
 & \frac{(15-15i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) - \frac{2a^2(13A-5iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
 & \frac{5a}{2A\sqrt{a+ia \tan(c+dx)}} \\
 & \frac{5d \tan^{\frac{5}{2}}(c+dx)}{5d \tan^{\frac{5}{2}}(c+dx)}
 \end{aligned}$$

input `Int[(Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(-2*A*Sqrt[a + I*a*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) + ((-2*a*(I*A + 5*B)*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (((15 - 15*I)*a^(5/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a^2*(13*A - (5*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a))/(5*a)`

3.159. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

3.159.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2)), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

3.159.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 554 vs. $2(145) = 290$.

Time = 0.14 (sec) , antiderivative size = 555, normalized size of antiderivative = 3.12

method	result
parts	$-\frac{A\sqrt{a(1+i\tan(dx+c))}\left(15i\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)\right)}{(\tan^3(dx+c))a-15\sqrt{2}\ln}$
derivativedivides	$-\frac{\sqrt{a(1+i\tan(dx+c))}\left(15iB\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)\right)}{a(\tan^4(dx+c))+15iA\sqrt{2}}$
default	$-\frac{\sqrt{a(1+i\tan(dx+c))}\left(15iB\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)\right)}{a(\tan^4(dx+c))+15iA\sqrt{2}}$

```
input int((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x,method=_R
RETURNVERBOSE)
```

```
output -1/30*A/d*(a*(1+I*tan(d*x+c)))^(1/2)/tan(d*x+c)^(5/2)*(15*I*2^(1/2)*ln((2*
2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x
+c))/(tan(d*x+c)+I))*tan(d*x+c)^3*a-15*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*
tan(d*x+c)^4*a+52*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d
*x+c)^3-16*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)-5
6*I*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)+12*I*(
a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)/(a*tan(d*x+c)*(1+I*tan(
d*x+c)))^(1/2)/(-I*a)^(1/2)/(-tan(d*x+c)+I)+1/6*B/d/tan(d*x+c)^(3/2)*(3*I*
2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I
*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*tan(d*x+c)^2*a-4*I*tan(d*x+c)*(-I*a)^(1
/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-4*(a*tan(d*x+c)*(1+I*tan(d*x+c)
))^(1/2)*(-I*a)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*
x+c)))^(1/2)/(-I*a)^(1/2)
```

$$3.159. \int \frac{\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

3.159.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 543 vs. $2(135) = 270$.

Time = 0.27 (sec) , antiderivative size = 543, normalized size of antiderivative = 3.05

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx =$$

$$15\sqrt{2}(de^{(6i dx + 6i c)} - 3de^{(4i dx + 4i c)} + 3de^{(2i dx + 2i c)} - d)\sqrt{-\frac{(-iA^2 - 2AB + iB^2)a}{d^2}} \log\left(\frac{\left(\sqrt{2}d\sqrt{-\frac{(-iA^2 - 2AB + iB^2)a}{d^2}}\right)}{\dots}\right)$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="fracas")`

output `-1/30*(15*sqrt(2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*log((sqrt(2)*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/(I*A + B) - 15*sqrt(2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*log(-sqrt(2)*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*e^(I*d*x + I*c) - sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/(I*A + B) + 4*sqrt(2)*((-17*I*A - 10*B)*e^(7*I*d*x + 7*I*c) + 3*I*A*e^(5*I*d*x + 5*I*c) + 5*(I*A + 2*B)*e^(3*I*d*x + 3*I*c) - 15*I*A*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)`

3.159.6 Sympy [F]

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{\sqrt{ia (\tan(c + dx) - i)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

input `integrate((a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(7/2),x)`

output `Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))/tan(c + d*x)**(7/2), x)`

3.159.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="maxima")`

output `Timed out`

3.159.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx);;OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[66]Warning, replacing 66 by -57, a substituti on variab`

3.159.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx)) \sqrt{a + a \tan(c + dx)} \operatorname{li}}{\tan(c + dx)^{7/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2))/tan(c + d*x)^(7/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2))/tan(c + d*x)^(7/2), x)`

3.160
$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

3.160.1 Optimal result 1662
 3.160.2 Mathematica [A] (verified) 1663
 3.160.3 Rubi [A] (verified) 1663
 3.160.4 Maple [B] (verified) 1668
 3.160.5 Fricas [B] (verification not implemented) 1669
 3.160.6 Sympy [F(-1)] 1670
 3.160.7 Maxima [F(-1)] 1670
 3.160.8 Giac [F(-2)] 1670
 3.160.9 Mupad [F(-1)] 1671

3.160.1 Optimal result

Integrand size = 38, antiderivative size = 221

$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

$$= \frac{(1-i)\sqrt{a}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$- \frac{2A\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} - \frac{2(iA+7B)\sqrt{a+ia \tan(c+dx)}}{35d \tan^{\frac{5}{2}}(c+dx)}$$

$$+ \frac{2(31A-7iB)\sqrt{a+ia \tan(c+dx)}}{105d \tan^{\frac{3}{2}}(c+dx)} + \frac{2(43iA+91B)\sqrt{a+ia \tan(c+dx)}}{105d \sqrt{\tan(c+dx)}}$$

```
output (1-I)*(A-I*B)*arctanh(((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*a^(1/2)/d+2/105*(43*I*A+91*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(1/2)-2/7*A*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(7/2)-2/35*(I*A+7*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(5/2)+2/105*(31*A-7*I*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(3/2)
```

3.160.2 Mathematica [A] (verified)

Time = 7.50 (sec) , antiderivative size = 282, normalized size of antiderivative = 1.28

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx = -\frac{2A\sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} + \frac{2 \left(-\frac{a(iA+7B)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} + \frac{a^2(31A-7iB)\sqrt{a+ia \tan(c+dx)}}{6d \tan^{\frac{3}{2}}(c+dx)} + \frac{2 \left(\frac{105a^4(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)\sqrt{\tan(c+dx)}}{4\sqrt{2}d\sqrt{ia \tan(c+dx)}} + \frac{a^3(43iA+91B)\sqrt{a+ia \tan(c+dx)}}{4d\sqrt{\tan(c+dx)}} \right)}{3a} \right)}{5a} + \frac{a^3(43iA+91B)\sqrt{a+ia \tan(c+dx)}}{7a}$$

```
input Integrate[(Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2), x]
```

```
output (-2*A*Sqrt[a + I*a*Tan[c + d*x]])/(7*d*Tan[c + d*x]^(7/2)) + (2*(-1/5*(a*(I*A + 7*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Tan[c + d*x]^(5/2)) + (2*((a^2*(31*A - (7*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(6*d*Tan[c + d*x]^(3/2)) + (2*((105*a^4*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[Tan[c + d*x]])/(4*Sqrt[2]*d*Sqrt[I*a*Tan[c + d*x]]) + (a^3*((43*I)*A + 91*B)*Sqrt[a + I*a*Tan[c + d*x]])/(4*d*Sqrt[Tan[c + d*x]])))/(3*a)))/(5*a)))/(7*a)
```

3.160.3 Rubi [A] (verified)

Time = 1.29 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.13, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.395$, Rules used = {3042, 4081, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx \quad \downarrow \quad 3042$$

3.160. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan(c + dx)^{9/2}} dx \\
& \quad \downarrow 4081 \\
& \frac{2 \int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA+7B)-6aA \tan(c+dx))}{2 \tan^{\frac{7}{2}}(c+dx)} dx}{7a} - \frac{2A \sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA+7B)-6aA \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx}{7a} - \frac{2A \sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA+7B)-6aA \tan(c+dx))}{\tan(c+dx)^{7/2}} dx}{7a} - \frac{2A \sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} \\
& \quad \downarrow 4081 \\
& \frac{2 \int -\frac{\sqrt{i \tan(c+dx)a+a}((31A-7iB)a^2+4(iA+7B) \tan(c+dx)a^2)}{2 \tan^{\frac{5}{2}}(c+dx)} dx}{5a} - \frac{2a(7B+iA) \sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \quad \frac{7a}{2A \sqrt{a + ia \tan(c + dx)}} \\
& \quad \frac{7d \tan^{\frac{7}{2}}(c + dx)}{7d \tan^{\frac{7}{2}}(c + dx)} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((31A-7iB)a^2+4(iA+7B) \tan(c+dx)a^2)}{\tan^{\frac{5}{2}}(c+dx)} dx}{5a} - \frac{2a(7B+iA) \sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \quad \frac{7a}{2A \sqrt{a + ia \tan(c + dx)}} \\
& \quad \frac{7d \tan^{\frac{7}{2}}(c + dx)}{7d \tan^{\frac{7}{2}}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((31A-7iB)a^2+4(iA+7B) \tan(c+dx)a^2)}{\tan(c+dx)^{5/2}} dx}{5a} - \frac{2a(7B+iA) \sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \quad \frac{7a}{2A \sqrt{a + ia \tan(c + dx)}} \\
& \quad \frac{7d \tan^{\frac{7}{2}}(c + dx)}{7d \tan^{\frac{7}{2}}(c + dx)} \\
& \quad \downarrow 4081
\end{aligned}$$

3.160. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

$$\frac{2 \int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(43iA+91B)-2a^3(31A-7iB) \tan(c+dx)) dx}{2 \tan^{\frac{3}{2}}(c+dx)} - \frac{2a^2(31A-7iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7B+iA)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)}}{5a}$$

$$\frac{2A\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 27

$$\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(43iA+91B)-2a^3(31A-7iB) \tan(c+dx)) dx}{\tan^{\frac{3}{2}}(c+dx)} - \frac{2a^2(31A-7iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7B+iA)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)}}{5a}$$

$$\frac{2A\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(43iA+91B)-2a^3(31A-7iB) \tan(c+dx)) dx}{\tan(c+dx)^{3/2}} - \frac{2a^2(31A-7iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7B+iA)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)}}{5a}$$

$$\frac{2A\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 4081

$$\frac{2 \int -\frac{105a^4(A-iB)\sqrt{i \tan(c+dx)a+a} dx}{2\sqrt{\tan(c+dx)}} - \frac{2a^3(91B+43iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(31A-7iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7B+iA)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)}}{3a}$$

$$\frac{2A\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 27

$$\frac{-105a^3(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(91B+43iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(31A-7iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7B+iA)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)}}{5a}$$

$$\frac{2A\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

3.160. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

↓ 3042

$$\frac{-105a^3(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(91B+43iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(31A-7iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7B+iA)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)}}{5a} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 4027

$$\frac{210ia^5(A-iB) \int \frac{1}{-2 \tan(c+dx)a^2} dx - \frac{d\sqrt{\tan(c+dx)}}{i \tan(c+dx)a+a} - ia - \frac{2a^3(91B+43iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(31A-7iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7B+iA)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)}}{5a} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 218

$$\frac{(105-105i)a^{7/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^3(91B+43iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(31A-7iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7B+iA)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)}}{5a} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

input `Int[(Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2),x]`

output `(-2*A*Sqrt[a + I*a*Tan[c + d*x]])/(7*d*Tan[c + d*x]^(7/2)) + ((-2*a*(I*A + 7*B)*Sqrt[a + I*a*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) - ((-2*a^2*(31*A - (7*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) + (((-105 + 105*I)*a^(7/2)*(A - I*B)*ArcTanh[(((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/d - (2*a^3*((43*I)*A + 91*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a))/(5*a))/(7*a)`

3.160. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

3.160.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2)), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

3.160.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 629 vs. 2(180) = 360.

Time = 0.14 (sec) , antiderivative size = 630, normalized size of antiderivative = 2.85

method	result
parts	$A \left(105i\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}}{\tan(dx+c)+i} \right) (\tan^4(dx+c)a - 124\sqrt{-ia} (\tan^2(dx+c)) \sqrt{a} \right)$
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} \left(105iA\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}}{\tan(dx+c)+i} \right) a (\tan^5(dx+c)) - 364B\sqrt{-ia} \right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))} \left(105iA\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}}{\tan(dx+c)+i} \right) a (\tan^5(dx+c)) - 364B\sqrt{-ia} \right)}{\dots}$

```
input int((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x,method=_R
RETURNVERBOSE)
```

```
output -1/210*A/d/tan(d*x+c)^(7/2)*(105*I*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*tan(d*x+c)^4*a-124*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-172*I*tan(d*x+c)^3*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+12*I*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+60*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)/(-I*a)^(1/2)-1/30*B/d*(a*(1+I*tan(d*x+c)))^(1/2)/tan(d*x+c)^(5/2)*(15*I*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*tan(d*x+c)^3*a-15*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*tan(d*x+c)^4*a+52*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3-16*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)-56*I*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)+12*I*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)/(-I*a)^(1/2)/(-tan(d*x+c)+I)
```

3.160.
$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

3.160.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 593 vs. $2(168) = 336$.

Time = 0.26 (sec) , antiderivative size = 593, normalized size of antiderivative = 2.68

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$= \frac{105 \sqrt{2} (de^{(8i dx + 8i c)} - 4 de^{(6i dx + 6i c)} + 6 de^{(4i dx + 4i c)} - 4 de^{(2i dx + 2i c)} + d) \sqrt{-\frac{(i A^2 + 2 AB - i B^2)a}{d^2}} \log \left(\frac{i \sqrt{2} d \sqrt{-\frac{(i A^2 + 2 AB - i B^2)a}{d^2}}}{\dots} \right)}{\dots}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="fricas")`

output `1/210*(105*sqrt(2)*(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*log((I*sqrt(2)*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/(I*A + B) - 105*sqrt(2)*(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*log((-I*sqrt(2)*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/(I*A + B) - 4*sqrt(2)*((92*A - 119*I*B)*e^(9*I*d*x + 9*I*c) - 20*(A - 7*I*B)*e^(7*I*d*x + 7*I*c) + 14*(2*A + I*B)*e^(5*I*d*x + 5*I*c) + 140*(A - I*B)*e^(3*I*d*x + 3*I*c) + 105*I*B*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)`

3.160.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(9/2),x)`

output `Timed out`

3.160.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="maxima")`

output `Timed out`

3.160.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[15]Warning, replacing 15 by 99, a substitution variable`

3.160. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

3.160.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx)) \sqrt{a + a \tan(c + dx)} \operatorname{li}}{\tan(c + dx)^{9/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2))/tan(c + d*x)^(9/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2))/tan(c + d*x)^(9/2), x)`

3.161 $\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.161.1 Optimal result	1672
3.161.2 Mathematica [A] (verified)	1673
3.161.3 Rubi [A] (verified)	1673
3.161.4 Maple [B] (verified)	1679
3.161.5 Fricas [B] (verification not implemented)	1680
3.161.6 Sympy [F(-1)]	1681
3.161.7 Maxima [F]	1682
3.161.8 Giac [F(-2)]	1682
3.161.9 Mupad [F(-1)]	1682

3.161.1 Optimal result

Integrand size = 38, antiderivative size = 248

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{(-1)^{3/4} a^{3/2} (22iA + 23B) \arctan\left(\frac{(-1)^{3/4} \sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) + \frac{(2 + 2i)a^{3/2}(iA + B) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{a(10A - 9iB) \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}}{8d} + \frac{a(6iA + 7B) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{12d} + \frac{iaB \tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{3d}$$

```
output 1/8*(-1)^(3/4)*a^(3/2)*(22*I*A+23*B)*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d+(2+2*I)*a^(3/2)*(I*A+B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d+1/8*a*(10*A-9*I*B)*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/d+1/12*a*(6*I*A+7*B)*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^(3/2)/d+1/3*I*a*B*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^(5/2)/d
```

3.161.2 Mathematica [A] (verified)

Time = 7.17 (sec) , antiderivative size = 415, normalized size of antiderivative = 1.67

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{B \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{3d} + \frac{3a(2A - iB)\sqrt{\tan(c+dx)}(a + ia \tan(c+dx))^{3/2}}{4d} + \frac{i(-\frac{3}{4}ia^3(2A - iB) - \frac{3}{4}a^3(6iA + 7B))}{d\sqrt{\tan(c+dx)}} \left(-\frac{2i\sqrt{2}a \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a + ia \tan(c+dx)}}\right)\sqrt{ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{2ia^{3/2}}{a} \right)$$

input `Integrate[Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]), x]`

output `(B*Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(3/2))/(3*d) + ((3*a*(2*A - I*B)*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(3/2))/(4*d) + (((-I)*(((3*I)/4)*a^3*(2*A - I*B) - (3*a^3*((6*I)*A + 7*B))/4)*((-2*I)*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) + ((2*I)*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])))/a - (((3*I)/4)*a^3*((6*I)*A + 7*B)*(-((-1)^(3/4)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]]*Sqrt[a + I*a*Tan[c + d*x]])/Sqrt[1 + I*Tan[c + d*x]]) + Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]))/d)/(2*a))/(3*a)`

3.161.3 Rubi [A] (verified)

Time = 1.63 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.08, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.447$, Rules used = {3042, 4077, 27, 3042, 4080, 27, 3042, 4080, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$$

↓ 3042

3.161. $\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \int \tan(c+dx)^{3/2}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))dx \\
& \quad \downarrow 4077 \\
& \frac{1}{3} \int \frac{1}{2} \tan^{\frac{3}{2}}(c+dx) \sqrt{i \tan(c+dx)a+a}(a(6A-5iB)+a(6iA+7B) \tan(c+dx))dx + \\
& \quad \frac{iaB \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \\
& \quad \downarrow 27 \\
& \frac{1}{6} \int \tan^{\frac{3}{2}}(c+dx) \sqrt{i \tan(c+dx)a+a}(a(6A-5iB)+a(6iA+7B) \tan(c+dx))dx + \\
& \quad \frac{iaB \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \\
& \quad \downarrow 3042 \\
& \frac{1}{6} \int \tan(c+dx)^{3/2} \sqrt{i \tan(c+dx)a+a}(a(6A-5iB)+a(6iA+7B) \tan(c+dx))dx + \\
& \quad \frac{iaB \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \\
& \quad \downarrow 4080 \\
& \frac{1}{6} \left(\frac{\int -\frac{3}{2} \sqrt{\tan(c+dx)} \sqrt{i \tan(c+dx)a+a}(a^2(6iA+7B)-a^2(10A-9iB) \tan(c+dx)) dx}{2a} + \frac{a(7B+6iA) \tan^{\frac{3}{2}}(c+dx)}{2d} \right) \\
& \quad \frac{iaB \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \\
& \quad \downarrow 27 \\
& \frac{1}{6} \left(\frac{a(7B+6iA) \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{3 \int \sqrt{\tan(c+dx)} \sqrt{i \tan(c+dx)a+a}(a^2(6iA+7B)-a^2(10A-9iB) \tan(c+dx)) dx}{4a} \right) \\
& \quad \frac{iaB \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \\
& \quad \downarrow 3042 \\
& \frac{1}{6} \left(\frac{a(7B+6iA) \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{3 \int \sqrt{\tan(c+dx)} \sqrt{i \tan(c+dx)a+a}(a^2(6iA+7B)-a^2(10A-9iB) \tan(c+dx)) dx}{4a} \right) \\
& \quad \frac{iaB \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{3d} \\
& \quad \downarrow 4080
\end{aligned}$$

3.161. $\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\frac{1}{6} \left(\frac{a(7B + 6iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(\int \frac{\sqrt{i \tan(c+dx)a+a} \left((10A-9iB)a^3 + (22iA+23B) \tan(c+dx)a^3 \right) dx}{2\sqrt{\tan(c+dx)} a} - \frac{a^2(10A-9iB)}{4a} \right)}{4a} \right)$$

$$\frac{iaB \tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{3d}$$

↓ 27

$$\frac{1}{6} \left(\frac{a(7B + 6iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(\int \frac{\sqrt{i \tan(c+dx)a+a} \left((10A-9iB)a^3 + (22iA+23B) \tan(c+dx)a^3 \right) dx}{\sqrt{\tan(c+dx)} 2a} - \frac{a^2(10A-9iB)}{4a} \right)}{4a} \right)$$

$$\frac{iaB \tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{a(7B + 6iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(\int \frac{\sqrt{i \tan(c+dx)a+a} \left((10A-9iB)a^3 + (22iA+23B) \tan(c+dx)a^3 \right) dx}{\sqrt{\tan(c+dx)} 2a} - \frac{a^2(10A-9iB)}{4a} \right)}{4a} \right)$$

$$\frac{iaB \tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{3d}$$

↓ 4084

$$\frac{1}{6} \left(\frac{a(7B + 6iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(\frac{32a^3(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - a^2(22A-23iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{a+ia \tan(c+dx)}}{\sqrt{\tan(c+dx)}} dx}{2a} \right)}{4a} \right)$$

$$\frac{iaB \tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{3d}$$

↓ 3042

3.161. $\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

$$\frac{1}{6} \left(\frac{a(7B + 6iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(\frac{32a^3(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - a^2(22A-23iB) \int \frac{(a-ia \tan(c+dx))\sqrt{a+ia \tan(c+dx)}}{\sqrt{\tan(c+dx)}} dx}{2a} \right)}{4a} \right) - \frac{iaB \tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{3d} \downarrow 4027$$

$$\frac{1}{6} \left(\frac{a(7B + 6iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(\frac{-a^2(22A-23iB) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{64ia^5(A-iB) \int \dots}{2a}}{2a} \right)}{4a} \right) - \frac{iaB \tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{3d} \downarrow 218$$

$$\frac{1}{6} \left(\frac{a(7B + 6iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(\frac{(32-32i)a^{7/2}(A-iB) \operatorname{arctanh} \left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}} \right) - a^2(22A-23iB) \int \dots}{d} \right)}{2a} \right) - \frac{iaB \tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{3d} \downarrow 4082$$

$$\frac{1}{6} \left(\frac{a(7B + 6iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(\frac{(32-32i)a^{7/2}(A-iB) \operatorname{arctanh} \left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}} \right) - a^4(22A-23iB) \int \dots}{d} \right)}{2a} \right) - \frac{iaB \tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{3d}$$

3.161. $\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

$$\begin{aligned}
 & \downarrow 65 \\
 & \left(\frac{1}{6} \frac{a(7B + 6iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(\frac{(32 - 32i)a^{7/2}(A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^4(22A - 23iB)}{2a} \right)}{2a} \right. \\
 & \qquad \qquad \qquad \left. \frac{iaB \tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{3d} \right) \\
 & \qquad \qquad \qquad \downarrow 216 \\
 & \left(\frac{1}{6} \frac{a(7B + 6iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(\frac{2\sqrt[4]{-1}a^{7/2}(22A - 23iB) \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{(32 - 32i)a^{7/2}(A - iB)}{2a} \right)}{2a} \right. \\
 & \qquad \qquad \qquad \left. \frac{iaB \tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{3d} \right)
 \end{aligned}$$

input `Int[Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `((I/3)*a*B*Tan[c + d*x]^(5/2)*Sqrt[a + I*a*Tan[c + d*x]]/d + ((a*((6*I)*A + 7*B)*Tan[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]])/(2*d) - (3*(((-1)^(1/4)*a^(7/2)*(22*A - (23*I)*B)*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d + ((32 - 32*I)*a^(7/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d)/(2*a) - (a^2*(10*A - (9*I)*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]/d)/(4*a))/6`

3.161. $\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.161.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4077 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

```
rule 4080 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[
1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Sim
p[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*T
an[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c
- a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]
```

```
rule 4082 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 + b^2, 0] && EqQ[A*b + a*B, 0]
```

```
rule 4084 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x]
- Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[
e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

3.161.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 649 vs. $2(198) = 396$.

Time = 0.17 (sec) , antiderivative size = 650, normalized size of antiderivative = 2.62

method	result
derivativedivides	$\frac{(\sqrt{\tan(dx+c)} \sqrt{a(1+i \tan(dx+c))} a \left(16iB\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} (\tan^2(dx+c)) + 24iA\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} (\tan^2(dx+c)) + 24iA\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} (\tan^2(dx+c)) + 24iA\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} (\tan^2(dx+c)) \right)}{a^2}$
default	$\frac{(\sqrt{\tan(dx+c)} \sqrt{a(1+i \tan(dx+c))} a \left(16iB\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} (\tan^2(dx+c)) + 24iA\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} (\tan^2(dx+c)) + 24iA\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} (\tan^2(dx+c)) + 24iA\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} (\tan^2(dx+c)) \right)}{a^2}$
parts	$\frac{A(\sqrt{\tan(dx+c)} \sqrt{a(1+i \tan(dx+c))} a \left(-4i \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia} \sqrt{-ia} \tan(dx+c) - 10 \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia} \sqrt{-ia} \tan(dx+c) \right)}{a^2}$

3.161. $\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

```
input int(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

```
output 1/48/d*tan(d*x+c)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)*a*(16*I*B*(I*a)^(1/2)*
-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2+24*I*A*(I*a
)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+27*I
*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(
1/2)+a)/(I*a)^(1/2)*(-I*a)^(1/2)*a-54*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*(a*ta
n(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+28*B*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+
c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)-24*I*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/
2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)-I*a+3*a*tan(d*x+c))/
(tan(d*x+c)+I)*a-30*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d
*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2)*(-I*a)^(1/2)*a+60*A*(I*a)^(1/2)*
(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-48*I*ln(1/2*(2*I*a*tan(
d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2)
)*(-I*a)^(1/2)*a+24*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d
*x+c)*(1+I*tan(d*x+c))))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I)*a-48*ln(
1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+
a)/(I*a)^(1/2)*(-I*a)^(1/2)*a)/(I*a)^(1/2)/(-I*a)^(1/2)/(a*tan(d*x+c)*(1+
I*tan(d*x+c)))^(1/2)
```

3.161.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 911 vs. $2(184) = 368$.

Time = 0.29 (sec) , antiderivative size = 911, normalized size of antiderivative = 3.67

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, al
gorithm="fricas")
```

output

```

1/48*(48*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*(d*e^(4*I*d*x + 4*
I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log((I*sqrt(2)*sqrt(-(I*A^2 + 2*A*B -
I*B^2)*a^3/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a*e^(2*I*d*x + 2*I
*c) + (-I*A - B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x
+ 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a))
- 48*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*(d*e^(4*I*d*x + 4*I*c
) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log((-I*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*
B^2)*a^3/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a*e^(2*I*d*x + 2*I*c
) + (-I*A - B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x +
2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a)) +
2*sqrt(2)*(7*(6*A - 7*I*B)*a*e^(5*I*d*x + 5*I*c) + 2*(30*A - 19*I*B)*a*e^(
3*I*d*x + 3*I*c) + 3*(6*A - 7*I*B)*a*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x
+ 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1
) + 3*sqrt((-484*I*A^2 - 1012*A*B + 529*I*B^2)*a^3/d^2)*(d*e^(4*I*d*x + 4*
I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log((sqrt(2)*((22*I*A + 23*B)*a*e^(2*I
*d*x + 2*I*c) + (22*I*A + 23*B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(
(-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + 2*I*sqrt((-484*I
*A^2 - 1012*A*B + 529*I*B^2)*a^3/d^2)*d*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/
((22*I*A + 23*B)*a)) - 3*sqrt((-484*I*A^2 - 1012*A*B + 529*I*B^2)*a^3/d^2
)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log((sqrt(2)*((2...

```

3.161.6 Sympy [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(3/2)*(a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output Timed out

3.161.7 Maxima [F]

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \int (B \tan(dx+c) + A)(ia \tan(dx+c) + a)^{\frac{3}{2}} \tan(dx+c)^{\frac{3}{2}} dx$$

input `integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)*tan(d*x + c)^(3/2), x)`

3.161.8 Giac [F(-2)]

Exception generated.

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \text{Exception raised: TypeError}$$

input `integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:The choice was done assuming 0=[0]Warning, replacing 0 by -48, a substitution variable should perhaps be purged.Warnin`

3.161.9 Mupad [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \int \tan(c+dx)^{3/2} (A+B \tan(c+dx)) (a+a \tan(c+dx) li)^{3/2} dx$$

input `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2),
x)`

output `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2),
x)`

3.162 $\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.162.1 Optimal result	1684
3.162.2 Mathematica [A] (verified)	1685
3.162.3 Rubi [A] (verified)	1685
3.162.4 Maple [B] (verified)	1690
3.162.5 Fricas [B] (verification not implemented)	1691
3.162.6 Sympy [F]	1692
3.162.7 Maxima [F]	1692
3.162.8 Giac [F(-2)]	1692
3.162.9 Mupad [F(-1)]	1693

3.162.1 Optimal result

Integrand size = 38, antiderivative size = 204

$$\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx =$$

$$\frac{(-1)^{3/4} a^{3/2} (12A - 11iB) \arctan\left(\frac{(-1)^{3/4} \sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{4d}$$

$$- \frac{(2 + 2i)a^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$+ \frac{a(4iA + 5B) \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}}{4d}$$

$$+ \frac{iaB \tan^{3/2}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d}$$

```
output -1/4*(-1)^(3/4)*a^(3/2)*(12*A-11*I*B)*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)
^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d-(2+2*I)*a^(3/2)*(A-I*B)*arctanh((1+I)*a
^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d+1/4*a*(4*I*A+5*B)*tan(
d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/d+1/2*I*a*B*(a+I*a*tan(d*x+c))^(1/2)
*tan(d*x+c)^(3/2)/d
```

3.162.2 Mathematica [A] (verified)

Time = 4.84 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.39

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{a \left(\frac{a(4A-3iB) \left((-1)^{3/4} \operatorname{arcsinh} \left(\sqrt[4]{-1} \sqrt{\tan(c+dx)} \right) - \sqrt{1+i \tan(c+dx)} \sqrt{\tan(c+dx)} \right) (-i+\tan(c+dx))}{\sqrt{1+i \tan(c+dx)}} \right) - 2a}{\dots}$$

input `Integrate[Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(a*((a*(4*A - (3*I)*B))*((-1)^(3/4)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]] - Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]]*(-I + Tan[c + d*x]))/Sqrt[1 + I*Tan[c + d*x]] - 2*a*B*Sqrt[Tan[c + d*x]]*(-I + Tan[c + d*x])^2 + (8*(A - I*B)*Sqrt[I*a*Tan[c + d*x]]*(Sqrt[a]*ArcSinh[Sqrt[I*a*Tan[c + d*x]]]/Sqrt[a])*Sqrt[1 + I*Tan[c + d*x]] - Sqrt[2]*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[a + I*a*Tan[c + d*x]])/Sqrt[Tan[c + d*x]])/(4*d*Sqrt[a + I*a*Tan[c + d*x]])`

3.162.3 Rubi [A] (verified)Time = 1.24 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4077, 27, 3042, 4080, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4077}$$

$$\frac{1}{2} \int \frac{1}{2} \sqrt{\tan(c+dx)} \sqrt{i \tan(c+dx)a + a(a(4A-3iB) + a(4iA+5B) \tan(c+dx))} dx + \frac{iaB \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 27

$$\frac{1}{4} \int \sqrt{\tan(c+dx)} \sqrt{i \tan(c+dx)a + a(a(4A-3iB) + a(4iA+5B) \tan(c+dx))} dx + \frac{iaB \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \int \sqrt{\tan(c+dx)} \sqrt{i \tan(c+dx)a + a(a(4A-3iB) + a(4iA+5B) \tan(c+dx))} dx + \frac{iaB \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 4080

$$\frac{1}{4} \left(\frac{\int -\frac{\sqrt{i \tan(c+dx)a + a(a^2(4iA+5B) - a^2(12A-11iB) \tan(c+dx))}}{2\sqrt{\tan(c+dx)}} dx}{a} + \frac{a(5B+4iA) \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)}}{d} \right) + \frac{iaB \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 27

$$\frac{1}{4} \left(\frac{a(5B+4iA) \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)}}{d} - \frac{\int \frac{\sqrt{i \tan(c+dx)a + a(a^2(4iA+5B) - a^2(12A-11iB) \tan(c+dx))}}{\sqrt{\tan(c+dx)}} dx}{2a} \right) + \frac{iaB \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(\frac{a(5B+4iA) \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)}}{d} - \frac{\int \frac{\sqrt{i \tan(c+dx)a + a(a^2(4iA+5B) - a^2(12A-11iB) \tan(c+dx))}}{\sqrt{\tan(c+dx)}} dx}{2a} \right) + \frac{iaB \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d}$$

↓ 4084

3.162. $\int \sqrt{\tan(c+dx)} (a+ia \tan(c+dx))^{3/2} (A+B \tan(c+dx)) dx$

$$\frac{1}{4} \left(\frac{a(5B + 4iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{16a^2(B + iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - a(11B + 12iA) \int \frac{(a-ia)}{\sqrt{\tan(c+dx)}} dx}{2a} \right) - \frac{iaB \tan^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{2d} \downarrow 3042$$

$$\frac{1}{4} \left(\frac{a(5B + 4iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{16a^2(B + iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - a(11B + 12iA) \int \frac{(a-ia)}{\sqrt{\tan(c+dx)}} dx}{2a} \right) - \frac{iaB \tan^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{2d} \downarrow 4027$$

$$\frac{1}{4} \left(\frac{a(5B + 4iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{32ia^4(B+iA) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} - a(11B + 12iA) \int \frac{(a-ia)}{\sqrt{\tan(c+dx)}} dx}{2a} \right) - \frac{iaB \tan^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{2d} \downarrow 218$$

$$\frac{1}{4} \left(\frac{a(5B + 4iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{(16-16i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - a(11B + 12iA) \int \frac{(a-ia)}{\sqrt{\tan(c+dx)}} dx}{2a} \right) - \frac{iaB \tan^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{2d} \downarrow 4082$$

$$\frac{1}{4} \left(\frac{a(5B + 4iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{(16-16i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{a^3(11B+12iA)}{2a} \right) - \frac{iaB \tan^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{2d} \downarrow 65$$

$$\frac{1}{4} \left(\frac{a(5B + 4iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{(16-16i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^3(11B+12iA)}{2a} \right) - \frac{iaB \tan^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{2d}$$

↓ 216

$$\frac{1}{4} \left(\frac{a(5B + 4iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2\sqrt[4]{-1}a^{5/2}(11B+12iA) \operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{(16-16i)a^{5/2}}{2a} \right) - \frac{iaB \tan^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{2d}$$

input `Int[Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `((I/2)*a*B*Tan[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]]/d + (-1/2*((2*(-1)^(1/4)*a^(5/2)*((12*I)*A + 11*B)*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d + ((16 - 16*I)*a^(5/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d)/a + (a*((4*I)*A + 5*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]/d)/4`

3.162.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 65 `Int[1/(Sqrt[(b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

3.162. $\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

rule 218 $\text{Int}[(a_+) + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a) * \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

rule 3042 $\text{Int}[u_+, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4027 $\text{Int}[\text{Sqrt}[(a_+) + (b_+)\tan[(e_+) + (f_+)(x_+)]]/\text{Sqrt}[(c_+) + (d_+)\tan[(e_+) + (f_+)(x_+)]], x_Symbol] \rightarrow \text{Simp}[-2*a*(b/f) \ \text{Subst}[\text{Int}[1/(a*c - b*d - 2*a^2*x^2), x], x, \text{Sqrt}[c + d*\text{Tan}[e + f*x]]/\text{Sqrt}[a + b*\text{Tan}[e + f*x]]], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0]$

rule 4077 $\text{Int}[(a_+) + (b_+)\tan[(e_+) + (f_+)(x_+)])^{(m_+)} * ((A_+) + (B_+)\tan[(e_+) + (f_+)(x_+)])^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[b*B*(a + b*\text{Tan}[e + f*x])^{(m-1)} * ((c + d*\text{Tan}[e + f*x])^{(n+1)}) / (d*f*(m+n)), x] + \text{Simp}[1/(d*(m+n)) \ \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m-1)} * (c + d*\text{Tan}[e + f*x])^n * \text{Simp}[a*A*d*(m+n) + B*(a*c*(m-1) - b*d*(n+1)) - (B*(b*c - a*d)*(m-1) - d*(A*b + a*B)*(m+n))*\text{Tan}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{!LtQ}[n, -1]$

rule 4080 $\text{Int}[(a_+) + (b_+)\tan[(e_+) + (f_+)(x_+)])^{(m_+)} * ((A_+) + (B_+)\tan[(e_+) + (f_+)(x_+)])^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[B*(a + b*\text{Tan}[e + f*x])^m * ((c + d*\text{Tan}[e + f*x])^n / (f*(m+n))), x] + \text{Simp}[1/(a*(m+n)) \ \text{Int}[(a + b*\text{Tan}[e + f*x])^m * (c + d*\text{Tan}[e + f*x])^{(n-1)} * \text{Simp}[a*A*c*(m+n) - B*(b*c*m + a*d*n) + (a*A*d*(m+n) - B*(b*d*m - a*c*n))*\text{Tan}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, m\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[n, 0]$

rule 4082 $\text{Int}[(a_+) + (b_+)\tan[(e_+) + (f_+)(x_+)])^{(m_+)} * ((A_+) + (B_+)\tan[(e_+) + (f_+)(x_+)])^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[b*(B/f) \ \text{Subst}[\text{Int}[(a + b*x)^{(m-1)} * (c + d*x)^n, x], x, \text{Tan}[e + f*x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, m, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[A*b + a*B, 0]$


```
rule 4084 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x]
- Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[
e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

3.162.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 562 vs. $2(162) = 324$.

Time = 0.15 (sec) , antiderivative size = 563, normalized size of antiderivative = 2.76

method	result
derivativedivides	$\frac{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})a\left(-4iB\sqrt{ia}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))}\tan(dx+c)+4iA\ln\left(\frac{2ia\tan(dx+c)+2\sqrt{a\tan(dx+c)(1+i\tan(dx+c))}}{2\sqrt{ia}}\right)\right)}{...}$
default	$\frac{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})a\left(-4iB\sqrt{ia}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))}\tan(dx+c)+4iA\ln\left(\frac{2ia\tan(dx+c)+2\sqrt{a\tan(dx+c)(1+i\tan(dx+c))}}{2\sqrt{ia}}\right)\right)}{...}$
parts	$\frac{A(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})a\left(2i\sqrt{ia}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))}+3i\ln\left(\frac{2ia\tan(dx+c)+2\sqrt{a\tan(dx+c)(1+i\tan(dx+c))}}{2\sqrt{ia}}\right)\right)}{...}$

```
input int(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

```
output -1/8/d*tan(d*x+c)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)*a*(-4*I*B*(I*a)^(1/2)*(-
-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+4*I*A*ln(1/2*
(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(
I*a)^(1/2))*(-I*a)^(1/2)*a-8*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1
+I*tan(d*x+c)))^(1/2)-4*I*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a
+5*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a
)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a-10*B*(I*a)^(1/2)*(-I*a)^(1/2)*(a*ta
n(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+8*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x
+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*(-I*a)^(1/2)-4*(
I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c
))))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a-8*ln(1/2*(2*I*a*tan(d*x+c
)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a
)^(1/2)*a)/(I*a)^(1/2)/(-I*a)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
```

$$3.162. \int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$$

3.162.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 829 vs. $2(150) = 300$.

Time = 0.27 (sec) , antiderivative size = 829, normalized size of antiderivative = 4.06

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output 1/8*(8*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log((sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a)) - 8*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log(-(sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*d*e^(I*d*x + I*c) - sqrt(2)*((-I*A - B)*a*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a)) + 2*sqrt(2)*((4*I*A + 7*B)*a*e^(3*I*d*x + 3*I*c) + (4*I*A + 3*B)*a*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + sqrt((144*I*A^2 + 264*A*B - 121*I*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log((sqrt(2)*((12*I*A + 11*B)*a*e^(2*I*d*x + 2*I*c) + (12*I*A + 11*B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + 2*sqrt((144*I*A^2 + 264*A*B - 121*I*B^2)*a^3/d^2)*d*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/((12*I*A + 11*B)*a)) - sqrt((144*I*A^2 + 264*A*B - 121*I*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log((sqrt(2)*((12*I*A + 11*B)*a*e^(2*I*d*x + 2*I*c) + (12*I*A + 11*B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) - 2*sqrt((144...
```

3.162.6 Sympy [F]

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \int (ia(\tan(c+dx)-i))^{3/2}(A+B \tan(c+dx)) \sqrt{\tan(c+dx)} dx$$

input `integrate(tan(d*x+c)**(1/2)*(a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(3/2)*(A + B*tan(c + d*x))*sqrt(tan(c + d*x)), x)`

3.162.7 Maxima [F]

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \int (B \tan(dx+c) + A)(ia \tan(dx+c) + a)^{3/2} \sqrt{\tan(dx+c)} dx$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)*sqrt(tan(d*x + c)), x)`

3.162.8 Giac [F(-2)]

Exception generated.

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \text{Exception raised: TypeError}$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:The choice was done assuming 0=[0]Warning, replacing 0 by -28, a substitution variable should perhaps be purged.Warnin

3.162.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \int \sqrt{\tan(c+dx)}(A+B \tan(c+dx))(a+a \tan(c+dx) 1i)^{3/2} dx$$

input `int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2), x)`

output `int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2), x)`

3.163
$$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.163.1 Optimal result 1694
 3.163.2 Mathematica [A] (verified) 1695
 3.163.3 Rubi [A] (verified) 1695
 3.163.4 Maple [B] (verified) 1699
 3.163.5 Fricas [B] (verification not implemented) 1700
 3.163.6 Sympy [F] 1701
 3.163.7 Maxima [F] 1702
 3.163.8 Giac [F(-2)] 1702
 3.163.9 Mupad [F(-1)] 1702

3.163.1 Optimal result

Integrand size = 38, antiderivative size = 156

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx =$$

$$\frac{(-1)^{3/4}a^{3/2}(2iA + 3B) \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$- \frac{(2 + 2i)a^{3/2}(iA + B)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$+ \frac{iaB\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d}$$

output

```

-(-1)^(3/4)*a^(3/2)*(2*I*A+3*B)*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)
/(a+I*a*tan(d*x+c))^(1/2))/d-(2+2*I)*a^(3/2)*(I*A+B)*arctanh((1+I)*a^(1/2)
*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d+I*a*B*tan(d*x+c)^(1/2)*(a+I*
a*tan(d*x+c))^(1/2)/d
    
```

3.163.2 Mathematica [A] (verified)

Time = 3.56 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.65

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \frac{aB \left(\sqrt[4]{-1} \operatorname{arcsinh} \left(\sqrt[4]{-1} \sqrt{\tan(c + dx)} \right) + i \sqrt{1 + i \tan(c + dx)} \right)}{d \sqrt{1 + i \tan(c + dx)}} + \frac{2a(iA + B) \sqrt{ia \tan(c + dx)} \left(\sqrt{a} \operatorname{arcsinh} \left(\frac{\sqrt{ia \tan(c + dx)}}{\sqrt{a}} \right) \sqrt{1 + i \tan(c + dx)} - \sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{2} \sqrt{ia \tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}} \right) \right)}{d \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}}$$

input `Integrate[((a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `(a*B*((-1)^(1/4)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]] + I*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]) + (2*a*(I*A + B)*Sqrt[I*a*Tan[c + d*x]]*(Sqrt[a]*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]] - Sqrt[2]*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]]*Sqrt[a + I*a*Tan[c + d*x]]))/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])`

3.163.3 Rubi [A] (verified)Time = 0.90 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4077, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

↓ 4077

3.163. $\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$

$$\begin{aligned}
& \int \frac{\sqrt{i \tan(c+dx)a + a(a(2A-iB) + a(2iA+3B) \tan(c+dx))}}{2\sqrt{\tan(c+dx)}} dx + \\
& \quad \frac{iaB \sqrt{\tan(c+dx)} \sqrt{a + ia \tan(c+dx)}}{d} \\
& \quad \downarrow 27 \\
& \frac{1}{2} \int \frac{\sqrt{i \tan(c+dx)a + a(a(2A-iB) + a(2iA+3B) \tan(c+dx))}}{\sqrt{\tan(c+dx)}} dx + \\
& \quad \frac{iaB \sqrt{\tan(c+dx)} \sqrt{a + ia \tan(c+dx)}}{d} \\
& \quad \downarrow 3042 \\
& \frac{1}{2} \int \frac{\sqrt{i \tan(c+dx)a + a(a(2A-iB) + a(2iA+3B) \tan(c+dx))}}{\sqrt{\tan(c+dx)}} dx + \\
& \quad \frac{iaB \sqrt{\tan(c+dx)} \sqrt{a + ia \tan(c+dx)}}{d} \\
& \quad \downarrow 4084 \\
& \frac{1}{2} \left(4a(A-iB) \int \frac{\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx - (2A-3iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx \right) + \\
& \quad \frac{iaB \sqrt{\tan(c+dx)} \sqrt{a + ia \tan(c+dx)}}{d} \\
& \quad \downarrow 3042 \\
& \frac{1}{2} \left(4a(A-iB) \int \frac{\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx - (2A-3iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx \right) + \\
& \quad \frac{iaB \sqrt{\tan(c+dx)} \sqrt{a + ia \tan(c+dx)}}{d} \\
& \quad \downarrow 4027 \\
& \frac{1}{2} \left(\frac{8ia^3(A-iB) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} - (2A-3iB) \int \frac{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx \right) + \\
& \quad \frac{iaB \sqrt{\tan(c+dx)} \sqrt{a + ia \tan(c+dx)}}{d} \\
& \quad \downarrow 218
\end{aligned}$$

3.163. $\int \frac{(a+ia \tan(c+dx))^{3/2} (A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

$$\frac{1}{2} \left(\frac{(4-4i)a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} - (2A-3iB) \int \frac{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx \right. \\ \left. \frac{iaB\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} \right)$$

↓ 4082

$$\frac{1}{2} \left(\frac{(4-4i)a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} - \frac{a^2(2A-3iB) \int \frac{1}{\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}} d\tan(c+dx)}{d} \right. \\ \left. \frac{iaB\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} \right)$$

↓ 65

$$\frac{1}{2} \left(\frac{(4-4i)a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} - \frac{2a^2(2A-3iB) \int \frac{1}{1-\frac{ia\tan(c+dx)}{i\tan(c+dx)a+a}} d\frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{d} \right) + \\ \frac{iaB\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}$$

↓ 216

$$\frac{1}{2} \left(\frac{2\sqrt[4]{-1}a^{3/2}(2A-3iB)\operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} + \frac{(4-4i)a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} \right) + \\ \frac{iaB\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}$$

input `Int[((a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `((2*(-1)^(1/4)*a^(3/2)*(2*A - (3*I)*B)*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]]]/Sqrt[a + I*a*Tan[c + d*x]])/d + ((4 - 4*I)*a^(3/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]]]/Sqrt[a + I*a*Tan[c + d*x]]])/d)/2 + (I*a*B*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/d`

3.163.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4077 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B))*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.163.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 483 vs. $2(126) = 252$.

Time = 0.15 (sec) , antiderivative size = 484, normalized size of antiderivative = 3.10

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} (\sqrt{\tan(dx+c)}) a \left(-iB \ln \left(\frac{2ia \tan(dx+c) + 2\sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a + 2iB\sqrt{ia} \sqrt{-ia} \right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))} (\sqrt{\tan(dx+c)}) a \left(-iB \ln \left(\frac{2ia \tan(dx+c) + 2\sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a + 2iB\sqrt{ia} \sqrt{-ia} \right)}{\dots}$
parts	$\frac{A\sqrt{a(1+i \tan(dx+c))} (\sqrt{\tan(dx+c)}) a^2 \left(i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}{\tan(dx+c) + i} \right) + 4i\sqrt{-ia} \right)}{\dots}$

input `int((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x,method=_R
ETURNVERBOSE)`

output

```

1/2/d*(a*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^(1/2)*a*(-I*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2)))*(-I*a)^(1/2)*a+2*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*(I*a)^(1/2)*a+2*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a+2*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*(-I*a)^(1/2)-(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a+2*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a)/(I*a)^(1/2)/(-I*a)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)

```

3.163.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 743 vs. $2(116) = 232$.

Time = 0.28 (sec) , antiderivative size = 743, normalized size of antiderivative = 4.76

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \frac{2i \sqrt{2} B a \sqrt{\frac{a}{e^{(2i dx + 2i c)} + 1}} \sqrt{\frac{-i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} + 1}} e^{(i dx + i c)} - 2 \sqrt{2} \sqrt{a} \sqrt{\frac{a}{e^{(2i dx + 2i c)} + 1}}}{\sqrt{\tan(c + dx)}}$$

input

```

integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="fracas")

```

output

```

1/2*(2*I*sqrt(2)*B*a*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x
+ 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*e^(I*d*x + I*c) - 2*sqrt(2)*sqrt
(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*d*log((I*sqrt(2)*sqrt(-(I*A^2 + 2*A*B -
I*B^2)*a^3/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a*e^(2*I*d*x + 2*
I*c) + (-I*A - B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x
+ 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a)
) + 2*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*d*log((-I*sqrt(2)*sqr
t(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B
)*a*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*
sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x -
I*c)/((-I*A - B)*a) - sqrt((-4*I*A^2 - 12*A*B + 9*I*B^2)*a^3/d^2)*d*log((
sqrt(2)*((2*I*A + 3*B)*a*e^(2*I*d*x + 2*I*c) + (2*I*A + 3*B)*a)*sqrt(a/(e^
(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*
I*c) + 1)) + 2*I*sqrt((-4*I*A^2 - 12*A*B + 9*I*B^2)*a^3/d^2)*d*e^(I*d*x +
I*c))*e^(-I*d*x - I*c)/((2*I*A + 3*B)*a) + sqrt((-4*I*A^2 - 12*A*B + 9*I*
B^2)*a^3/d^2)*d*log((sqrt(2)*((2*I*A + 3*B)*a*e^(2*I*d*x + 2*I*c) + (2*I*A
+ 3*B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c)
+ I)/(e^(2*I*d*x + 2*I*c) + 1)) - 2*I*sqrt((-4*I*A^2 - 12*A*B + 9*I*B^2)*a
^3/d^2)*d*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/((2*I*A + 3*B)*a))/d

```

3.163.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \int \frac{(ia(\tan(c + dx) - i))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

input

```
integrate((a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(1/2),x)
```

output

```
Integral((I*a*(tan(c + d*x) - I))**(3/2)*(A + B*tan(c + d*x))/sqrt(tan(c +
d*x)), x)
```

3.163.7 Maxima [F]

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{3/2}}{\sqrt{\tan(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)/sqrt(tan(d*x + c)), x)`

3.163.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[80]Warning, replacing 80 by -82, a substitution on variab`

3.163.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) i)^{3/2}}{\sqrt{\tan(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2))/tan(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2))/tan(c + d*x)^(1/2), x)`

3.163. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

3.164
$$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.164.1 Optimal result 1703
 3.164.2 Mathematica [B] (verified) 1703
 3.164.3 Rubi [A] (verified) 1704
 3.164.4 Maple [B] (verified) 1708
 3.164.5 Fricas [B] (verification not implemented) 1709
 3.164.6 Sympy [F] 1710
 3.164.7 Maxima [F(-1)] 1710
 3.164.8 Giac [F(-2)] 1710
 3.164.9 Mupad [F(-1)] 1711

3.164.1 Optimal result

Integrand size = 38, antiderivative size = 146

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \frac{2\sqrt{-1}a^{3/2}B \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{(2 + 2i)a^{3/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2aA\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}}$$

output

```
2*(-1)^(1/4)*a^(3/2)*B*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)^(1/2))/d+(2+2*I)*a^(3/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)^(1/2))/d-2*a*A*(a+I*a*tan(d*x+c)^(1/2))/d/tan(d*x+c)^(1/2)
```

3.164.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 336 vs. 2(146) = 292.

Time = 6.89 (sec) , antiderivative size = 336, normalized size of antiderivative = 2.30

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = -\frac{2A(a + ia \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}} + \frac{i(a^2A + \frac{1}{2}ia^2(3iA+B))}{a} \left(-\frac{2i\sqrt{2}a \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{2ia^{3/2} \operatorname{arcsinh}\left(\frac{\sqrt{ia \tan(c+dx)}}{\sqrt{a}}\right) \sqrt{1+i \tan(c+dx)} \sqrt{ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)}} \right)$$

input `Integrate[((a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2), x]`

output `(-2*A*(a + I*a*Tan[c + d*x])^(3/2))/(d*Sqrt[Tan[c + d*x]]) + (2*(((-I)*(a^2*A + (I/2)*a^2*((3*I)*A + B))*((-2*I)*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) + ((2*I)*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])))/a + (I*a^2*A*(-(((-1)^(3/4)*ArcSinh[(-1)^(1/4)]*Sqrt[Tan[c + d*x]])*Sqrt[a + I*a*Tan[c + d*x]])/Sqrt[1 + I*Tan[c + d*x]]) + Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]))/d)/a`

3.164.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4076, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx \xrightarrow{3042} \int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx \xrightarrow{4076}$$

3.164. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{3/2}(c+dx)} dx$

$$\begin{aligned}
& 2 \int \frac{\sqrt{i \tan(c+dx)a + a(a(2iA+B) + iaB \tan(c+dx))}}{2\sqrt{\tan(c+dx)}} dx - \frac{2aA\sqrt{a + ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 27 \\
& \int \frac{\sqrt{i \tan(c+dx)a + a(a(2iA+B) + iaB \tan(c+dx))}}{\sqrt{\tan(c+dx)}} dx - \frac{2aA\sqrt{a + ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 3042 \\
& \int \frac{\sqrt{i \tan(c+dx)a + a(a(2iA+B) + iaB \tan(c+dx))}}{\sqrt{\tan(c+dx)}} dx - \frac{2aA\sqrt{a + ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 4084 \\
& 2a(B+iA) \int \frac{\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx - B \int \frac{(a - ia \tan(c+dx))\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx - \\
& \quad \frac{2aA\sqrt{a + ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 3042 \\
& 2a(B+iA) \int \frac{\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx - B \int \frac{(a - ia \tan(c+dx))\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx - \\
& \quad \frac{2aA\sqrt{a + ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 4027 \\
& \frac{4ia^3(B+iA) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} - \\
& B \int \frac{(a - ia \tan(c+dx))\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx - \frac{2aA\sqrt{a + ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 218 \\
& -B \int \frac{(a - ia \tan(c+dx))\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx + \\
& \frac{(2-2i)a^{3/2}(B+iA) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2aA\sqrt{a + ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 4082 \\
& -\frac{a^2B \int \frac{1}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{d} + \\
& \frac{(2-2i)a^{3/2}(B+iA) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2aA\sqrt{a + ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}
\end{aligned}$$

3.164. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \downarrow 65 \\
& -\frac{2a^2 B \int \frac{1}{1 - \frac{ia \tan(c+dx)}{a+a}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} + \frac{(2-2i)a^{3/2}(B+iA) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} \\
& \quad \frac{2aA\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \downarrow 216 \\
& \frac{(2-2i)a^{3/2}(B+iA) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{2\sqrt[4]{-1}a^{3/2}B \operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} \\
& \quad \frac{2aA\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}
\end{aligned}$$

input `Int[((a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `(2*(-1)^(1/4)*a^(3/2)*B*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]]]/Sqrt[a + I*a*Tan[c + d*x]])/d + ((2 - 2*I)*a^(3/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]]]/Sqrt[a + I*a*Tan[c + d*x]])/d - (2*a*A*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]])`

3.164.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

3.164. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.164.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 518 vs. $2(118) = 236$.

Time = 0.18 (sec) , antiderivative size = 519, normalized size of antiderivative = 3.55

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} a \left(4iA \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a \tan(dx+c) - i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{a(1+i \tan(dx+c))}}{\sqrt{-ia}} \right) \right)}{\sqrt{a(1+i \tan(dx+c))} a \left(4iA \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a \tan(dx+c) - i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{a(1+i \tan(dx+c))}}{\sqrt{-ia}} \right) \right)}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))} a \left(4iA \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a \tan(dx+c) - i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{a(1+i \tan(dx+c))}}{\sqrt{-ia}} \right) \right)}{\sqrt{a(1+i \tan(dx+c))} a \left(4iA \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a \tan(dx+c) - i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{a(1+i \tan(dx+c))}}{\sqrt{-ia}} \right) \right)}$
parts	$-\frac{A\sqrt{a(1+i \tan(dx+c))} a \left(i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - ia + 3a \tan(dx+c)}{\tan(dx+c)+i} \right) a \tan(dx+c) + \sqrt{ia} \sqrt{2} \right)}{\sqrt{a(1+i \tan(dx+c))} a \left(4iA \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a \tan(dx+c) - i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{a(1+i \tan(dx+c))}}{\sqrt{-ia}} \right) \right)}$

input `int((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{2} \frac{d \left(a(1+i \tan(dx+c)) \right)^{3/2} (A+B \tan(dx+c)) / \tan(dx+c)^{3/2}}{a(1+i \tan(dx+c))^{1/2} \left(4iA \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a \tan(dx+c) - i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{a(1+i \tan(dx+c))}}{\sqrt{-ia}} \right) \right) + \frac{A \sqrt{a(1+i \tan(dx+c))} a \left(i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - ia + 3a \tan(dx+c)}{\tan(dx+c)+i} \right) a \tan(dx+c) + \sqrt{ia} \sqrt{2} \right)}{\sqrt{a(1+i \tan(dx+c))} a \left(4iA \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a \tan(dx+c) - i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{a(1+i \tan(dx+c))}}{\sqrt{-ia}} \right) \right)}$$

3.164.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 748 vs. $2(110) = 220$.

Time = 0.26 (sec) , antiderivative size = 748, normalized size of antiderivative = 5.12

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx =$$

$$2\sqrt{2}\sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^3}{d^2}} (de^{(2i dx + 2i c)} - d) \log \left(\frac{\left(\sqrt{2}\sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^3}{d^2}} de^{(i dx + i c)} + \sqrt{2}((-iA - B)ae^{(2i dx + 2i c)} + (-iA - B)a \right)}{(-iA - B)a} \right)$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="fricas")`

output

```
-1/2*(2*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log((sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/((-I*A - B)*a) - 2*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log(-(sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*d*e^(I*d*x + I*c) - sqrt(2)*((-I*A - B)*a*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))))*e^(-I*d*x - I*c)/((-I*A - B)*a) + 4*sqrt(2)*(I*A*a*e^(3*I*d*x + 3*I*c) + I*A*a*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + sqrt(-4*I*B^2*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log((sqrt(2)*(B*a*e^(2*I*d*x + 2*I*c) + B*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + sqrt(-4*I*B^2*a^3/d^2)*d*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(B*a) - sqrt(-4*I*B^2*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log((sqrt(2)*(B*a*e^(2*I*d*x + 2*I*c) + B*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) - sqrt(-4*I*B^2*a^3/d^2)*d*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(B*a)))/(d*e^(2*I*d*x + 2*I*c) - d)
```

3.164.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \int \frac{(ia(\tan(c + dx) - i))^{3/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx$$

input `integrate((a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(3/2),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(3/2)*(A + B*tan(c + d*x))/tan(c + d*x)**(3/2), x)`

3.164.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="maxima")`

output `Timed out`

3.164.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx);OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[94]Warning, replacing 94 by -29, a substituti on variab`

3.164.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) 1i)^{3/2}}{\tan(c + dx)^{3/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2))/tan(c + d*x)^(3/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2))/tan(c + d*x)^(3/2), x)`

$$3.165 \quad \int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.165.1 Optimal result 1712
 3.165.2 Mathematica [B] (verified) 1712
 3.165.3 Rubi [A] (verified) 1713
 3.165.4 Maple [B] (verified) 1716
 3.165.5 Fricas [B] (verification not implemented) 1717
 3.165.6 Sympy [F] 1718
 3.165.7 Maxima [F] 1718
 3.165.8 Giac [F(-2)] 1719
 3.165.9 Mupad [F(-1)] 1719

3.165.1 Optimal result

Integrand size = 38, antiderivative size = 137

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = \frac{(2 + 2i)a^{3/2}(iA + B)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2aA\sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2a(4iA + 3B)\sqrt{a + ia \tan(c + dx)}}{3d\sqrt{\tan(c + dx)}}$$

```
output (2+2*I)*a^(3/2)*(I*A+B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d-2/3*a*(4*I*A+3*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(1/2)-2/3*a*A*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(3/2)
```

3.165.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 304 vs. 2(137) = 274.

Time = 7.01 (sec) , antiderivative size = 304, normalized size of antiderivative = 2.22

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = -\frac{2A(a + ia \tan(c + dx))^{3/2}}{3d \tan^{\frac{3}{2}}(c + dx)} + (iA + B) \left(\frac{2\sqrt{2}a \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} - \frac{2a^{3/2} \operatorname{arcsinh}\left(\frac{\sqrt{ia \tan(c+dx)}}{\sqrt{a}}\right) \sqrt{1 + i \tan(c + dx)}}{d\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}} \right)$$

3.165. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

input `Integrate[((a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(-2*A*(a + I*a*Tan[c + d*x])^(3/2))/(3*d*Tan[c + d*x]^(3/2)) + (I*A + B)*((2*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[I*a*Tan[c + d*x]]/(d*Sqrt[Tan[c + d*x]]) - (2*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + (2*(-1)^(1/4)*a*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]) - (2*a*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]])`

3.165.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.237$, Rules used = {3042, 4076, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx$$

↓ 4076

$$\frac{2}{3} \int \frac{\sqrt{ia \tan(c + dx)a + a(a(4iA + 3B) - a(2A - 3iB) \tan(c + dx))}}{2 \tan^{3/2}(c + dx)} dx - \frac{2aA \sqrt{a + ia \tan(c + dx)}}{3d \tan^{3/2}(c + dx)}$$

↓ 27

$$\frac{1}{3} \int \frac{\sqrt{ia \tan(c + dx)a + a(a(4iA + 3B) - a(2A - 3iB) \tan(c + dx))}}{\tan^{3/2}(c + dx)} dx - \frac{2aA \sqrt{a + ia \tan(c + dx)}}{3d \tan^{3/2}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \int \frac{\sqrt{ia \tan(c + dx)a + a(a(4iA + 3B) - a(2A - 3iB) \tan(c + dx))}}{\tan(c + dx)^{3/2}} dx - \frac{2aA \sqrt{a + ia \tan(c + dx)}}{3d \tan^{3/2}(c + dx)}$$

↓ 4081

3.165. $\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx$

$$\begin{aligned}
& \frac{1}{3} \left(\frac{2 \int -\frac{3a^2(A-iB)\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{a} - \frac{2a(3B+4iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) - \\
& \quad \frac{2aA\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow 27 \\
& \frac{1}{3} \left(-6a(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a(3B+4iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) - \\
& \quad \frac{2aA\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{3} \left(-6a(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a(3B+4iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) - \\
& \quad \frac{2aA\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow 4027 \\
& \frac{1}{3} \left(\frac{12ia^3(A-iB) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} - \frac{2a(3B+4iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) - \\
& \quad \frac{2aA\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow 218 \\
& \frac{1}{3} \left(-\frac{(6-6i)a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a(3B+4iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) - \\
& \quad \frac{2aA\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)}
\end{aligned}$$

input `Int[((a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

```
output (-2*a*A*Sqrt[a + I*a*Tan[c + d*x]]/(3*d*Tan[c + d*x]^(3/2)) + (((-6 + 6*I)
)*a^(3/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]]]/Sqrt[a +
I*a*Tan[c + d*x]]])/d - (2*a*((4*I)*A + 3*B)*Sqrt[a + I*a*Tan[c + d*x]]/(
d*Sqrt[Tan[c + d*x]]))/3
```

3.165.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4027 Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4076 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

3.165.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 615 vs. 2(112) = 224.

Time = 0.13 (sec) , antiderivative size = 616, normalized size of antiderivative = 4.50

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} a \left(-12iB \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a(\tan^2(dx+c))+3i\sqrt{ia} \right)}{}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))} a \left(-12iB \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a(\tan^2(dx+c))+3i\sqrt{ia} \right)}{}$
parts	$\frac{A\sqrt{a(1+i \tan(dx+c))} a \left(3i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - ia+3a \tan(dx+c)}{\tan(dx+c)+i} \right) a(\tan^2(dx+c))+12i \right)}{}$

```
input int((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x,method=_R
ETURNVERBOSE)
```

3.165.
$$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

output
$$-1/6/d*(a*(1+I*\tan(d*x+c)))^{1/2}*a/\tan(d*x+c)^{3/2}*(-12*I*B*\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}*(I*a)^{1/2}+a)/(I*a)^{1/2})*(-I*a)^{1/2}*a*\tan(d*x+c)^2+3*I*(I*a)^{1/2}*2^{1/2}*\ln((2*2^{1/2}*(-I*a)^{1/2}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}-I*a+3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*a*\tan(d*x+c)^2+16*I*A*(I*a)^{1/2}*(-I*a)^{1/2}*\tan(d*x+c)*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2}+12*A*\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}*(I*a)^{1/2}+a)/(I*a)^{1/2})*(-I*a)^{1/2}*a*\tan(d*x+c)^2+6*I*\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}*(I*a)^{1/2}+a)/(I*a)^{1/2})*(-I*a)^{1/2}*a*\tan(d*x+c)^2-3*(I*a)^{1/2}*2^{1/2}*\ln((2*2^{1/2}*(-I*a)^{1/2}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}-I*a+3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*a*\tan(d*x+c)^2+12*B*(I*a)^{1/2}*(-I*a)^{1/2}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2}*\tan(d*x+c)+6*\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}*(I*a)^{1/2}+a)/(I*a)^{1/2})*(-I*a)^{1/2}*a*\tan(d*x+c)^2+4*A*(I*a)^{1/2}*(-I*a)^{1/2}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2})/(I*a)^{1/2}/(-I*a)^{1/2}/(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2}$$

3.165.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 524 vs. 2(103) = 206.

Time = 0.26 (sec) , antiderivative size = 524, normalized size of antiderivative = 3.82

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \frac{3 \sqrt{2} \sqrt{-\frac{(i A^2 + 2 AB - i B^2) a^3}{d^2}} (de^{(4i dx + 4i c)} - 2 de^{(2i dx + 2i c)} + \dots)$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="fricas")`

output $\frac{1}{3}(3\sqrt{2})\sqrt{-(I^2A^2 + 2AB - I^2B^2)a^3/d^2}(d e^{4Ix + 4Ic} - 2d e^{2Ix + 2Ic} + d)\log((I\sqrt{2})\sqrt{-(I^2A^2 + 2AB - I^2B^2)a^3/d^2}d e^{Ix + Ic} + \sqrt{2}((-IA - B)a e^{2Ix + 2Ic} + (-IA - B)a)\sqrt{a/(e^{2Ix + 2Ic} + 1)})\sqrt{(-Ie^{2Ix + 2Ic} + I)/(e^{2Ix + 2Ic} + 1))}e^{-Ix - Ic}/((-IA - B)a) - 3\sqrt{2})\sqrt{-(I^2A^2 + 2AB - I^2B^2)a^3/d^2}(d e^{4Ix + 4Ic} - 2d e^{2Ix + 2Ic} + d)\log((-I\sqrt{2})\sqrt{-(I^2A^2 + 2AB - I^2B^2)a^3/d^2}d e^{Ix + Ic} + \sqrt{2}((-IA - B)a e^{2Ix + 2Ic} + (-IA - B)a)\sqrt{a/(e^{2Ix + 2Ic} + 1)})\sqrt{(-Ie^{2Ix + 2Ic} + I)/(e^{2Ix + 2Ic} + 1))}e^{-Ix - Ic}/((-IA - B)a) + 2\sqrt{2})((5A - 3IB)a e^{5Ix + 5Ic} + 2Aa e^{3Ix + 3Ic} - 3(A - IB)a e^{Ix + Ic})\sqrt{a/(e^{2Ix + 2Ic} + 1)})\sqrt{(-Ie^{2Ix + 2Ic} + I)/(e^{2Ix + 2Ic} + 1))}/(d e^{4Ix + 4Ic} - 2d e^{2Ix + 2Ic} + d)$

3.165.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \int \frac{(ia(\tan(c + dx) - i))^{3/2}(A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx$$

input `integrate((a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(5/2),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(3/2)*(A + B*tan(c + d*x))/tan(c + d*x)**(5/2), x)`

3.165.7 Maxima [F]

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{3/2}}{\tan(dx + c)^{5/2}} dx$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)/tan(d*x + c)^(5/2), x)`

3.165. $\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx$

3.165.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[78]Warning, replacing 78 by 64, a substitution variable`

3.165.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) i)^{3/2}}{\tan(c + dx)^{5/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*i)^(3/2))/tan(c + d*x)^(5/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*i)^(3/2))/tan(c + d*x)^(5/2), x)`

3.166
$$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{7/2}(c+dx)} dx$$

3.166.1 Optimal result 1720
 3.166.2 Mathematica [A] (verified) 1720
 3.166.3 Rubi [A] (verified) 1721
 3.166.4 Maple [B] (verified) 1725
 3.166.5 Fricas [B] (verification not implemented) 1726
 3.166.6 Sympy [F] 1726
 3.166.7 Maxima [F(-1)] 1727
 3.166.8 Giac [F(-2)] 1727
 3.166.9 Mupad [F(-1)] 1728

3.166.1 Optimal result

Integrand size = 38, antiderivative size = 181

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx =$$

$$-\frac{(2 + 2i)a^{3/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2aA\sqrt{a + ia \tan(c + dx)}}{5d \tan^{5/2}(c + dx)}$$

$$- \frac{2a(6iA + 5B)\sqrt{a + ia \tan(c + dx)}}{15d \tan^{3/2}(c + dx)} + \frac{4a(9A - 10iB)\sqrt{a + ia \tan(c + dx)}}{15d\sqrt{\tan(c + dx)}}$$

output $(-2-2*I)*a^{(3/2)}*(A-I*B)*\operatorname{arctanh}\left(\frac{(1+I)*a^{(1/2)}*\tan(d*x+c)^{(1/2)}}{(a+I*a*\tan(d*x+c))^{(1/2)}}\right)/d+4/15*a*(9*A-10*I*B)*(a+I*a*\tan(d*x+c))^{(1/2)}/d/\tan(d*x+c)^{(1/2)}-2/5*a*A*(a+I*a*\tan(d*x+c))^{(1/2)}/d/\tan(d*x+c)^{(5/2)}-2/15*a*(6*I*A+5*B)*(a+I*a*\tan(d*x+c))^{(1/2)}/d/\tan(d*x+c)^{(3/2)}$

3.166.2 Mathematica [A] (verified)

Time = 8.09 (sec) , antiderivative size = 360, normalized size of antiderivative = 1.99

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = -\frac{2A(a + ia \tan(c + dx))^{3/2}}{5d \tan^{5/2}(c + dx)}$$

$$+ 2\left(-\frac{a(3iA+5B)(a+ia \tan(c+dx))^{3/2}}{3d \tan^{3/2}(c+dx)} - \frac{5}{2}a(A - iB)\left(\frac{2\sqrt{2}a\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)\sqrt{ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^{3/2}\operatorname{arcsinh}\left(\frac{\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d\sqrt{\tan(c+dx)}}\right)\right)$$

3.166.
$$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{7/2}(c+dx)} dx$$

input `Integrate[((a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(-2*A*(a + I*a*Tan[c + d*x])^(3/2))/(5*d*Tan[c + d*x]^(5/2)) + (2*(-1/3*(a*((3*I)*A + 5*B)*(a + I*a*Tan[c + d*x])^(3/2))/(d*Tan[c + d*x]^(3/2)) - (5*a*(A - I*B)*((2*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - (2*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x])) + (2*(-1)^(1/4)*a*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]) - (2*a*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/2)/(5*a)`

3.166.3 Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.07, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {3042, 4076, 27, 3042, 4081, 25, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan(c + dx)^{7/2}} dx$$

↓ 4076

$$\frac{2}{5} \int \frac{\sqrt{ia \tan(c + dx)a + a(6iA + 5B) - a(4A - 5iB) \tan(c + dx)}}{2 \tan^{5/2}(c + dx)} dx - \frac{2aA \sqrt{a + ia \tan(c + dx)}}{5d \tan^{5/2}(c + dx)}$$

↓ 27

$$\frac{1}{5} \int \frac{\sqrt{ia \tan(c + dx)a + a(6iA + 5B) - a(4A - 5iB) \tan(c + dx)}}{\tan^{5/2}(c + dx)} dx - \frac{2aA \sqrt{a + ia \tan(c + dx)}}{5d \tan^{5/2}(c + dx)}$$

↓ 3042

$$\frac{1}{5} \int \frac{\sqrt{ia \tan(c + dx)a + a(6iA + 5B) - a(4A - 5iB) \tan(c + dx)}}{\tan(c + dx)^{5/2}} dx - \frac{2aA \sqrt{a + ia \tan(c + dx)}}{5d \tan^{5/2}(c + dx)}$$

3.166. $\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx$

$$\begin{aligned}
& \downarrow 4081 \\
& \frac{1}{5} \left(\frac{2 \int -\frac{\sqrt{i \tan(c+dx)a+a((9A-10iB)a^2+(6iA+5B) \tan(c+dx)a^2)} dx}{3a} - \frac{2a(5B+6iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right) - \\
& \quad \frac{2aA\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \downarrow 25 \\
& \frac{1}{5} \left(-\frac{2 \int \frac{\sqrt{i \tan(c+dx)a+a((9A-10iB)a^2+(6iA+5B) \tan(c+dx)a^2)} dx}{3a} - \frac{2a(5B+6iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right) - \\
& \quad \frac{2aA\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \downarrow 3042 \\
& \frac{1}{5} \left(-\frac{2 \int \frac{\sqrt{i \tan(c+dx)a+a((9A-10iB)a^2+(6iA+5B) \tan(c+dx)a^2)} dx}{3a \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(5B+6iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right) - \\
& \quad \frac{2aA\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \downarrow 4081 \\
& \frac{1}{5} \left(\frac{2 \left(\frac{2 \int \frac{15a^3(iA+B)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx}{a} - \frac{2a^2(9A-10iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)}{3a} - \frac{2a(5B+6iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right) - \\
& \quad \frac{2aA\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \downarrow 27 \\
& \frac{1}{5} \left(-\frac{2 \left(15a^2(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(9A-10iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)}{3a} - \frac{2a(5B+6iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right) - \\
& \quad \frac{2aA\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)}
\end{aligned}$$

3.166. $\int \frac{(a+ia \tan(c+dx))^{\frac{3}{2}}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$\begin{array}{c}
 \downarrow 3042 \\
 \frac{1}{5} \left(\frac{2 \left(15a^2(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(9A-10iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)}{3a} - \frac{2a(5B+6iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right) \\
 \frac{2aA\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
 \downarrow 4027 \\
 \frac{1}{5} \left(\frac{2 \left(\frac{30ia^4(B+iA) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} - \frac{2a^2(9A-10iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)}{3a} - \frac{2a(5B+6iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right) \\
 \frac{2aA\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
 \downarrow 218 \\
 \frac{1}{5} \left(\frac{2 \left(\frac{(15-15i)a^{5/2}(B+iA) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^2(9A-10iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)}{3a} - \frac{2a(5B+6iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right) \\
 \frac{2aA\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)}
 \end{array}$$

input `Int[((a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2), x]`

output `(-2*a*A*Sqrt[a + I*a*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) + ((-2*a*((6*I)*A + 5*B)*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (2*((15 - 15*I)*a^(5/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a^2*(9*A - (10*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a))/5`

3.166. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

3.166.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4076 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`
- rule 4081 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

3.166.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 704 vs. $2(148) = 296$.

Time = 0.13 (sec) , antiderivative size = 705, normalized size of antiderivative = 3.90

method	result
derivativedivides	$-\frac{\sqrt{a(1+i \tan(dx+c))} a \left(-72A\sqrt{ia} \sqrt{-ia} (\tan^2(dx+c)) \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} + 60iA \ln \left(\frac{2ia \tan(dx+c) + 2\sqrt{a} \tan(dx+c)}{\tan(dx+c) + i} \right) \right)}{\dots}$
default	$-\frac{\sqrt{a(1+i \tan(dx+c))} a \left(-72A\sqrt{ia} \sqrt{-ia} (\tan^2(dx+c)) \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} + 60iA \ln \left(\frac{2ia \tan(dx+c) + 2\sqrt{a} \tan(dx+c)}{\tan(dx+c) + i} \right) \right)}{\dots}$
parts	$\frac{A\sqrt{a(1+i \tan(dx+c))} a \left(5i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - ia + 3a \tan(dx+c)}{\tan(dx+c) + i} \right) \right) a (\tan^3(dx+c)) + 5\sqrt{ia} \dots}{\dots}$

```
input int((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x,method=_R
RETURNVERBOSE)
```

```
output -1/30/d*(a*(1+I*tan(d*x+c)))^(1/2)*a/tan(d*x+c)^(5/2)*(-72*A*(I*a)^(1/2)*(-
-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+60*I*A*ln(1
/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a
)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^3-15*I*(I*a)^(1/2)*2^(1/2)*ln((2*
2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x
+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3+80*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*
x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+60*B*ln(1/2*(2*I*a*tan(d*x+c)
+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a
)^(1/2)*a*tan(d*x+c)^3+30*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*
tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^3
-15*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(
d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3+24*I*A*(
I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)-3
0*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(
1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^3+20*B*(I*a)^(1/2)*(-I*a)^(
1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+12*A*(I*a)^(1/2)*(-I
*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2))/(I*a)^(1/2)/(-I*a)^(1/2)/
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
```

3.166.
$$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^2(c+dx)} dx$$

3.166.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 593 vs. $2(137) = 274$.

Time = 0.25 (sec) , antiderivative size = 593, normalized size of antiderivative = 3.28

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \frac{15 \sqrt{2} \sqrt{-\frac{(-i A^2 - 2AB + i B^2)a^3}{d^2}} (de^{(6i dx + 6i c)} - 3 de^{(4i dx + 4i c)})}{\dots}$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="fricas")`

output `1/15*(15*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*log((sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a)) - 15*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*log(-(sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*d*e^(I*d*x + I*c) - sqrt(2)*((-I*A - B)*a*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a)) - 2*sqrt(2)*((-27*I*A - 25*B)*a*e^(7*I*d*x + 7*I*c) + 3*(I*A + 5*B)*a*e^(5*I*d*x + 5*I*c) + 5*(3*I*A + 5*B)*a*e^(3*I*d*x + 3*I*c) + 15*(-I*A - B)*a*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)`

3.166.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \int \frac{(ia(\tan(c + dx) - i))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx$$

input `integrate((a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(7/2),x)`

3.166. $\int \frac{(a+ia \tan(c+dx))^{3/2} (A+B \tan(c+dx))}{\tan^{7/2}(c+dx)} dx$

output `Integral((I*a*(tan(c + d*x) - I)**(3/2)*(A + B*tan(c + d*x))/tan(c + d*x)**(7/2), x)`

3.166.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="maxima")`

output `Timed out`

3.166.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[40]Warning, replacing 40 by 37, a substitution variabl`

3.166.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) 1i)^{3/2}}{\tan(c + dx)^{7/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2))/tan(c + d*x)^(7/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2))/tan(c + d*x)^(7/2), x)`

3.167
$$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

3.167.1 Optimal result 1729
 3.167.2 Mathematica [A] (verified) 1730
 3.167.3 Rubi [A] (verified) 1730
 3.167.4 Maple [B] (verified) 1735
 3.167.5 Fricas [B] (verification not implemented) 1736
 3.167.6 Sympy [F(-1)] 1737
 3.167.7 Maxima [F(-1)] 1737
 3.167.8 Giac [F(-2)] 1738
 3.167.9 Mupad [F(-1)] 1738

3.167.1 Optimal result

Integrand size = 38, antiderivative size = 225

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx = \frac{(2 - 2i)a^{3/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$- \frac{2aA\sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{2a(8iA + 7B)\sqrt{a + ia \tan(c + dx)}}{35d \tan^{\frac{5}{2}}(c + dx)}$$

$$+ \frac{4a(19A - 21iB)\sqrt{a + ia \tan(c + dx)}}{105d \tan^{\frac{3}{2}}(c + dx)} + \frac{4a(67iA + 63B)\sqrt{a + ia \tan(c + dx)}}{105d \sqrt{\tan(c + dx)}}$$

```
output (2-2*I)*a^(3/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d+4/105*a*(67*I*A+63*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(1/2)-2/7*a*A*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(7/2)-2/35*a*(8*I*A+7*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(5/2)+4/105*a*(19*A-21*I*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(3/2)
```


3.167.2 Mathematica [A] (verified)

Time = 7.98 (sec) , antiderivative size = 416, normalized size of antiderivative = 1.85

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = -\frac{2A(a + ia \tan(c + dx))^{3/2}}{7d \tan^{7/2}(c + dx)} + 2 \left(-\frac{a(3iA+7B)(a+ia \tan(c+dx))^{3/2}}{5d \tan^{5/2}(c+dx)} + \frac{a^2(29A-21iB)(a+ia \tan(c+dx))^{3/2} - \frac{35}{4}a^2(iA+B)}{6d \tan^{3/2}(c+dx)} \left(\frac{2\sqrt{2}a \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d\sqrt{\tan(c+dx)}} \right) \sqrt{ia \tan(c+dx)} \right)$$

input `Integrate[((a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2),x]`

output `(-2*A*(a + I*a*Tan[c + d*x])^(3/2))/(7*d*Tan[c + d*x]^(7/2)) + (2*(-1/5*(a*((3*I)*A + 7*B)*(a + I*a*Tan[c + d*x])^(3/2))/(d*Tan[c + d*x]^(5/2)) + (2*((a^2*(29*A - (21*I)*B)*(a + I*a*Tan[c + d*x])^(3/2))/(6*d*Tan[c + d*x]^(3/2)) - (35*a^2*(I*A + B)*((2*Sqrt[2]*a*ArcTanH[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - (2*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x])) + (2*(-1)^(1/4)*a*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]) - (2*a*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]])))/4)/(5*a)))/(7*a)`

3.167.3 Rubi [A] (verified)

Time = 1.34 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.10, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.395$, Rules used = {3042, 4076, 27, 3042, 4081, 25, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx$$

↓ 3042

3.167. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{9/2}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan(c + dx)^{9/2}} dx \\
& \quad \downarrow 4076 \\
& \frac{2}{7} \int \frac{\sqrt{ia \tan(c + dx)a + a(a(8iA + 7B) - a(6A - 7iB) \tan(c + dx))}}{2 \tan^{7/2}(c + dx)} dx - \frac{2aA \sqrt{a + ia \tan(c + dx)}}{7d \tan^{7/2}(c + dx)} \\
& \quad \downarrow 27 \\
& \frac{1}{7} \int \frac{\sqrt{ia \tan(c + dx)a + a(a(8iA + 7B) - a(6A - 7iB) \tan(c + dx))}}{\tan^{7/2}(c + dx)} dx - \frac{2aA \sqrt{a + ia \tan(c + dx)}}{7d \tan^{7/2}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{7} \int \frac{\sqrt{ia \tan(c + dx)a + a(a(8iA + 7B) - a(6A - 7iB) \tan(c + dx))}}{\tan(c + dx)^{7/2}} dx - \frac{2aA \sqrt{a + ia \tan(c + dx)}}{7d \tan^{7/2}(c + dx)} \\
& \quad \downarrow 4081 \\
& \frac{1}{7} \left(\frac{2 \int -\frac{\sqrt{ia \tan(c + dx)a + a((19A - 21iB)a^2 + 2(8iA + 7B) \tan(c + dx)a^2)}}{\tan^{5/2}(c + dx)} dx}{5a} - \frac{2a(7B + 8iA) \sqrt{a + ia \tan(c + dx)}}{5d \tan^{5/2}(c + dx)} \right) - \\
& \quad \frac{2aA \sqrt{a + ia \tan(c + dx)}}{7d \tan^{7/2}(c + dx)} \\
& \quad \downarrow 25 \\
& \frac{1}{7} \left(-\frac{2 \int \frac{\sqrt{ia \tan(c + dx)a + a((19A - 21iB)a^2 + 2(8iA + 7B) \tan(c + dx)a^2)}}{\tan^{5/2}(c + dx)} dx}{5a} - \frac{2a(7B + 8iA) \sqrt{a + ia \tan(c + dx)}}{5d \tan^{5/2}(c + dx)} \right) - \\
& \quad \frac{2aA \sqrt{a + ia \tan(c + dx)}}{7d \tan^{7/2}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{7} \left(-\frac{2 \int \frac{\sqrt{ia \tan(c + dx)a + a((19A - 21iB)a^2 + 2(8iA + 7B) \tan(c + dx)a^2)}}{\tan(c + dx)^{5/2}} dx}{5a} - \frac{2a(7B + 8iA) \sqrt{a + ia \tan(c + dx)}}{5d \tan^{5/2}(c + dx)} \right) - \\
& \quad \frac{2aA \sqrt{a + ia \tan(c + dx)}}{7d \tan^{7/2}(c + dx)} \\
& \quad \downarrow 4081
\end{aligned}$$

3.167. $\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx$

$$\frac{1}{7} \left(\frac{2 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(67iA+63B)-2a^3(19A-21iB) \tan(c+dx))}{2 \tan^{\frac{3}{2}}(c+dx)} dx - \frac{2a^2(19A-21iB) \sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} - \frac{2a(7B+8iA) \sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right)$$

$$\frac{2aA \sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{7} \left(\frac{2 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(67iA+63B)-2a^3(19A-21iB) \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx - \frac{2a^2(19A-21iB) \sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} - \frac{2a(7B+8iA) \sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right)$$

$$\frac{2aA \sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{2 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(67iA+63B)-2a^3(19A-21iB) \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)^{3/2}} dx - \frac{2a^2(19A-21iB) \sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} - \frac{2a(7B+8iA) \sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right)$$

$$\frac{2aA \sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 4081

$$\frac{1}{7} \left(\frac{2 \left(\frac{2 \int -\frac{105a^4(A-iB)\sqrt{i \tan(c+dx)a+a} dx}{2\sqrt{\tan(c+dx)}}}{a} - \frac{2a^3(63B+67iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(19A-21iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} - \frac{2a(7B+8iA)}{5d \tan^{\frac{7}{2}}(c+dx)} \right)$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{7} \left(\frac{2 \left(\frac{-105a^3(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(63B+67iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(19A-21iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} - \frac{2a(7B+8iA)}{5d \tan^{\frac{7}{2}}(c+dx)} \right)$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{2 \left(\frac{-105a^3(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(63B+67iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(19A-21iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} - \frac{2a(7B+8iA)}{5d \tan^{\frac{7}{2}}(c+dx)} \right)$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 4027

$$\frac{1}{7} \left(\frac{2 \left(\frac{210ia^5(A-iB) \int \frac{1}{-2 \tan(c+dx)a^2 - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{3a} - \frac{2a^3(63B+67iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(19A-21iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} - \frac{2a(7B+8iA)}{5d \tan^{\frac{7}{2}}(c+dx)} \right)$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

3.167. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

$$\begin{array}{c} \downarrow 218 \\ \frac{1}{7} \left(\frac{2 \left(\frac{(105-105i)a^{7/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} - \frac{2a^3(63B+67iA)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(19A-21iB)\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} \right)}{5a} \right) \\ \frac{2aA\sqrt{a+ia\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} \end{array}$$

input `Int[((a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2), x]`

output `(-2*a*A*Sqrt[a + I*a*Tan[c + d*x]]/(7*d*Tan[c + d*x]^(7/2)) + ((-2*a*((8*I)*A + 7*B)*Sqrt[a + I*a*Tan[c + d*x]]/(5*d*Tan[c + d*x]^(5/2)) - (2*((-2*a^2*(19*A - (21*I)*B)*Sqrt[a + I*a*Tan[c + d*x]]/(3*d*Tan[c + d*x]^(3/2)) + (((-105 + 105*I)*a^(7/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a^3*((67*I)*A + 63*B)*Sqrt[a + I*a*Tan[c + d*x]]/(d*Sqrt[Tan[c + d*x]]))/(3*a)))/(5*a))/7`

3.167.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4027 Int[Sqrt[(a_) + (b_.)*tan[(e_) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*tan[(e_) + (f_.)*(x_)], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

```
rule 4076 Int[((a_) + (b_.)*tan[(e_) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

```
rule 4081 Int[((a_) + (b_.)*tan[(e_) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]
```

3.167.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 793 vs. 2(184) = 368.

Time = 0.14 (sec) , antiderivative size = 794, normalized size of antiderivative = 3.53

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} a \left(504B\sqrt{ia} \sqrt{-ia} (\tan^3(dx+c)) \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} + 536iA\sqrt{ia} \sqrt{-ia} (\tan^3(dx+c)) \right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))} a \left(504B\sqrt{ia} \sqrt{-ia} (\tan^3(dx+c)) \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} + 536iA\sqrt{ia} \sqrt{-ia} (\tan^3(dx+c)) \right)}{\dots}$
parts	$\frac{A\sqrt{a(1+i \tan(dx+c))} a \left(105i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}}{\tan(dx+c)+i} \right) a (\tan^4(dx+c)) + 420i \right)}{\dots}$

$$3.167. \int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^2(c+dx)} dx$$

```
input int((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x,method=_R
ETURNVERBOSE)
```

```
output 1/210/d*(a*(1+I*tan(d*x+c)))^(1/2)*a/tan(d*x+c)^(7/2)*(504*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+536*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-96*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+152*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-168*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+420*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^4+210*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^4-105*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^4-420*I*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^4+210*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^4+105*I*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^4-84*B*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)-60*A*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2))/(I*a)^(1/2)/(-I*a)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
```

3.167.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 638 vs. $2(171) = 342$.

Time = 0.28 (sec) , antiderivative size = 638, normalized size of antiderivative = 2.84

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx =$$

$$105 \sqrt{2} \sqrt{-\frac{(iA^2 + 2AB - iB^2)a^3}{d^2}} (de^{(8i dx + 8i c)} - 4de^{(6i dx + 6i c)} + 6de^{(4i dx + 4i c)} - 4de^{(2i dx + 2i c)} + d) \log \left(\frac{(i\sqrt{2}\sqrt{-\frac{(iA^2 + 2AB - iB^2)a^3}{d^2}} + d)}{\dots} \right)$$

```
input integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="fricas")
```

$$3.167. \quad \int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx$$

```
output -1/105*(105*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*(d*e^(8*I*d*x +
8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d
*x + 2*I*c) + d)*log((I*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*d*e
^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a)
*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2
*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a)) - 105*sqrt(2)*sqrt
(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x
+ 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)*log((-I
*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*d*e^(I*d*x + I*c) + sqrt(2
))*((-I*A - B)*a*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a)*sqrt(a/(e^(2*I*d*x + 2
*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*
e^(-I*d*x - I*c)/((-I*A - B)*a)) + 2*sqrt(2)*((211*A - 189*I*B)*a*e^(9*I*d
*x + 9*I*c) - 10*(16*A - 21*I*B)*a*e^(7*I*d*x + 7*I*c) + 14*(A + 6*I*B)*a*
e^(5*I*d*x + 5*I*c) + 70*(4*A - 3*I*B)*a*e^(3*I*d*x + 3*I*c) - 105*(A - I*
B)*a*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*
x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(8*I*d*x + 8*I*c) - 4*d*e
^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d
)
```

3.167.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(9/2),x)
```

output Timed out

3.167.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, al
gorithm="maxima")
```

3.167. $\int \frac{(a+ia \tan(c+dx))^{3/2} (A+B \tan(c+dx))}{\tan^{9/2}(c+dx)} dx$

output Timed out

3.167.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[16]Warning, replacing 16 by 40, a substitution variable

3.167.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) i)^{3/2}}{\tan(c + dx)^{9/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2))/tan(c + d*x)^(9/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2))/tan(c + d*x)^(9/2), x)`

3.168
$$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$$

3.168.1 Optimal result 1739
 3.168.2 Mathematica [A] (verified) 1740
 3.168.3 Rubi [A] (verified) 1741
 3.168.4 Maple [B] (verified) 1747
 3.168.5 Fricas [B] (verification not implemented) 1748
 3.168.6 Sympy [F(-1)] 1749
 3.168.7 Maxima [F(-1)] 1750
 3.168.8 Giac [F(-2)] 1750
 3.168.9 Mupad [F(-1)] 1750

3.168.1 Optimal result

Integrand size = 38, antiderivative size = 269

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx = \frac{(2 + 2i)a^{3/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$- \frac{2aA\sqrt{a + ia \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} - \frac{2a(10iA + 9B)\sqrt{a + ia \tan(c + dx)}}{63d \tan^{\frac{7}{2}}(c + dx)}$$

$$+ \frac{4a(11A - 12iB)\sqrt{a + ia \tan(c + dx)}}{105d \tan^{\frac{5}{2}}(c + dx)} + \frac{4a(61iA + 57B)\sqrt{a + ia \tan(c + dx)}}{315d \tan^{\frac{3}{2}}(c + dx)}$$

$$- \frac{4a(193A - 201iB)\sqrt{a + ia \tan(c + dx)}}{315d \sqrt{\tan(c + dx)}}$$

output

```
(2+2*I)*a^(3/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d-4/315*a*(193*A-201*I*B)*(a+I*a*tan(d*x+c)^(1/2))/d/tan(d*x+c)^(1/2)-2/9*a*A*(a+I*a*tan(d*x+c)^(1/2))/d/tan(d*x+c)^(9/2)-2/63*a*(10*I*A+9*B)*(a+I*a*tan(d*x+c)^(1/2))/d/tan(d*x+c)^(7/2)+4/105*a*(11*A-12*I*B)*(a+I*a*tan(d*x+c)^(1/2))/d/tan(d*x+c)^(5/2)+4/315*a*(61*I*A+57*B)*(a+I*a*tan(d*x+c)^(1/2))/d/tan(d*x+c)^(3/2)
```

3.168.2 Mathematica [A] (verified)

Time = 8.39 (sec) , antiderivative size = 470, normalized size of antiderivative = 1.75

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = -\frac{2A(a + ia \tan(c + dx))^{3/2}}{9d \tan^{9/2}(c + dx)} + \left(-\frac{3a(iA + 3B)(a + ia \tan(c + dx))^{3/2}}{7d \tan^{7/2}(c + dx)} + \frac{3a^2(17A - 9iB)(a + ia \tan(c + dx))^{3/2}}{10d \tan^{5/2}(c + dx)} + \frac{a^3(71iA + 87B)(a + ia \tan(c + dx))^{3/2}}{4d \tan^{3/2}(c + dx)} + \frac{315}{8} a^3 (A - iB) \left(\frac{2\sqrt{2}a \arctan\left(\frac{\sqrt{2}\sqrt{a + ia \tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right)}{d \sqrt{a + ia \tan(c + dx)}} \right) \right)$$

input `Integrate[((a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(11/2), x]`

output `(-2*A*(a + I*a*Tan[c + d*x])^(3/2))/(9*d*Tan[c + d*x]^(9/2)) + (2*((-3*a*(I*A + 3*B)*(a + I*a*Tan[c + d*x])^(3/2))/(7*d*Tan[c + d*x]^(7/2)) + (2*((3*a^2*(17*A - (9*I)*B)*(a + I*a*Tan[c + d*x])^(3/2))/(10*d*Tan[c + d*x]^(5/2)) + (2*((a^3*((71*I)*A + 87*B)*(a + I*a*Tan[c + d*x])^(3/2))/(4*d*Tan[c + d*x]^(3/2)) + (315*a^3*(A - I*B)*((2*sqrt[2]*a*ArcTanh[(sqrt[2]*sqrt[I*a*Tan[c + d*x]])/sqrt[a + I*a*Tan[c + d*x]])*sqrt[I*a*Tan[c + d*x]])/(d*sqrt[Tan[c + d*x]]) - (2*a^(3/2)*ArcSinh[sqrt[I*a*Tan[c + d*x]]/sqrt[a]]*sqrt[1 + I*Tan[c + d*x]]*sqrt[I*a*Tan[c + d*x]])/(d*sqrt[Tan[c + d*x]]*sqrt[a + I*a*Tan[c + d*x]]) + (2*(-1)^(1/4)*a*ArcSinh[(-1)^(1/4]*sqrt[Tan[c + d*x]])*sqrt[a + I*a*Tan[c + d*x]])/(d*sqrt[1 + I*Tan[c + d*x]]) - (2*a*sqrt[a + I*a*Tan[c + d*x]])/(d*sqrt[Tan[c + d*x]])))/8)/(5*a))/(7*a))/(9*a)`

3.168.3 Rubi [A] (verified)

Time = 1.70 (sec) , antiderivative size = 302, normalized size of antiderivative = 1.12, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.474$, Rules used = {3042, 4076, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan(c + dx)^{11/2}} dx \\
 & \quad \downarrow \text{4076} \\
 & \frac{2}{9} \int \frac{\sqrt{i \tan(c + dx)a + a(a(10iA + 9B) - a(8A - 9iB) \tan(c + dx))}}{2 \tan^{\frac{9}{2}}(c + dx)} dx - \frac{2aA \sqrt{a + ia \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{9} \int \frac{\sqrt{i \tan(c + dx)a + a(a(10iA + 9B) - a(8A - 9iB) \tan(c + dx))}}{\tan^{\frac{9}{2}}(c + dx)} dx - \frac{2aA \sqrt{a + ia \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{9} \int \frac{\sqrt{i \tan(c + dx)a + a(a(10iA + 9B) - a(8A - 9iB) \tan(c + dx))}}{\tan(c + dx)^{9/2}} dx - \frac{2aA \sqrt{a + ia \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} \\
 & \quad \downarrow \text{4081} \\
 & \frac{1}{9} \left(\frac{2 \int -\frac{3\sqrt{i \tan(c + dx)a + a((11A - 12iB)a^2 + (10iA + 9B) \tan(c + dx)a^2)}}{\tan^{\frac{7}{2}}(c + dx)} dx}{7a} - \frac{2a(9B + 10iA) \sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} \right) - \\
 & \quad \frac{2aA \sqrt{a + ia \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$\frac{1}{9} \left(\frac{6 \int \frac{\sqrt{i \tan(c+dx)a+a((11A-12iB)a^2+(10iA+9B) \tan(c+dx)a^2)} dx}{\tan^{\frac{7}{2}}(c+dx)} - \frac{2a(9B+10iA)\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} \right) -$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{9} \left(\frac{6 \int \frac{\sqrt{i \tan(c+dx)a+a((11A-12iB)a^2+(10iA+9B) \tan(c+dx)a^2)} dx}{\tan(c+dx)^{7/2}} - \frac{2a(9B+10iA)\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} \right) -$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

↓ 4081

$$\frac{1}{9} \left(\frac{6 \left(\frac{2 \int \frac{\sqrt{i \tan(c+dx)a+a(a^3(61iA+57B)-4a^3(11A-12iB) \tan(c+dx))} dx}{2 \tan^{\frac{5}{2}}(c+dx)} - \frac{2a^2(11A-12iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right)}{7a} - \frac{2a(9B+10iA)\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} \right) -$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{9} \left(\frac{6 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a(a^3(61iA+57B)-4a^3(11A-12iB) \tan(c+dx))} dx}{\tan^{\frac{5}{2}}(c+dx)} - \frac{2a^2(11A-12iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right)}{7a} - \frac{2a(9B+10iA)\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} \right) -$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

↓ 3042

3.168. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$

$$\frac{1}{9} \left(\frac{6 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(61iA+57B)-4a^3(11A-12iB) \tan(c+dx))}{\tan(c+dx)^{5/2}} dx}{5a} - \frac{2a^2(11A-12iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{5/2}(c+dx)} \right)}{7a} - \frac{2a(9B+10iA)\sqrt{a+ia \tan(c+dx)}}{7d \tan^{7/2}(c+dx)} \right)$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{9d \tan^{9/2}(c+dx)}$$

↓ 4081

$$\frac{1}{9} \left(\frac{6 \left(\frac{2 \int -\frac{\sqrt{i \tan(c+dx)a+a} ((193A-201iB)a^4+2(61iA+57B) \tan(c+dx)a^4)}{2 \tan^{3/2}(c+dx)} dx}{3a} - \frac{2a^3(57B+61iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{3/2}(c+dx)} \right)}{5a} - \frac{2a^2(11A-12iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{5/2}(c+dx)} \right)$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{9d \tan^{9/2}(c+dx)}$$

↓ 27

$$\frac{1}{9} \left(\frac{6 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} ((193A-201iB)a^4+2(61iA+57B) \tan(c+dx)a^4)}{\tan^{3/2}(c+dx)} dx}{3a} - \frac{2a^3(57B+61iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{3/2}(c+dx)} \right)}{5a} - \frac{2a^2(11A-12iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{5/2}(c+dx)} \right)$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{9d \tan^{9/2}(c+dx)}$$

↓ 3042

3.168. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{11/2}(c+dx)} dx$

$$\frac{1}{9} \left(\frac{6 \left(\int \frac{\sqrt{i \tan(c+dx)a+a} \left((193A-201iB)a^4 + 2(61iA+57B) \tan(c+dx)a^4 \right) dx}{\tan(c+dx)^{3/2}} - \frac{2a^3(57B+61iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a^2(11A-12iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right)}{7a} \right)$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

↓ 4081

$$\frac{1}{9} \left(\frac{6 \left(\frac{2 \int \frac{315a^5(iA+B)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx}{a} - \frac{2a^4(193A-201iB)\sqrt{a+ia \tan(c+dx)}}{3a d \sqrt{\tan(c+dx)}} - \frac{2a^3(57B+61iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a^2(11A-12iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right)}{7a} \right)$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{9} \left(\frac{6 \left(-\frac{315a^4(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{3a} - \frac{2a^4(193A-201iB)\sqrt{a+ia \tan(c+dx)}}{d \sqrt{\tan(c+dx)}} - \frac{2a^3(57B+61iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a^2(11A-12iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right)}{7a} \right)$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

↓ 3042

3.168. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$

$$\frac{1}{9} \left(\frac{6 \left(\frac{315a^4(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^4(193A-201iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^3(57B+61iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a^2(11A-12iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right)}{7a} \right)$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

↓ 4027

$$\frac{1}{9} \left(\frac{6 \left(\frac{630ia^6(B+iA) \int \frac{1}{-2 \tan(c+dx)a^2 - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} - \frac{2a^4(193A-201iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^3(57B+61iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a^2(11A-12iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right)}{7a} \right)$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

↓ 218

$$\frac{1}{9} \left(\frac{6 \left(\frac{2a^3(57B+61iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{(315-315i)a^{9/2}(B+iA) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^4(193A-201iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(11A-12iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right)}{7a} \right)$$

$$\frac{2aA\sqrt{a+ia \tan(c+dx)}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

3.168. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$

input `Int[((a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(11/2),x]`

output `(-2*a*A*Sqrt[a + I*a*Tan[c + d*x]]/(9*d*Tan[c + d*x]^(9/2)) + ((-2*a*((10*I)*A + 9*B)*Sqrt[a + I*a*Tan[c + d*x]]/(7*d*Tan[c + d*x]^(7/2)) - (6*((-2*a^2*(11*A - (12*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) + ((-2*a^3*((61*I)*A + 57*B)*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (((315 - 315*I)*a^(9/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a^4*(193*A - (201*I)*B)*Sqrt[a + I*a*Tan[c + d*x]]/(d*Sqrt[Tan[c + d*x]]))/(3*a)/(5*a))/(7*a))/9`

3.168.3.1 Defintions of rubi rules used

rule 277 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

```
rule 4076 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

3.168.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 882 vs. $2(220) = 440$.

Time = 0.14 (sec) , antiderivative size = 883, normalized size of antiderivative = 3.28

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} a \left(-1544 A \sqrt{ia} \sqrt{-ia} (\tan^4(dx+c)) \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - 315 i \sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a}}{\dots} \right) \right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))} a \left(-1544 A \sqrt{ia} \sqrt{-ia} (\tan^4(dx+c)) \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - 315 i \sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a}}{\dots} \right) \right)}{\dots}$
parts	Expression too large to display

```
input int((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x,method=_
RETURNVERBOSE)
```

$$3.168. \int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$$

```

output 1/630/d*(a*(1+I*tan(d*x+c)))^(1/2)*a/tan(d*x+c)^(9/2)*(-1544*A*(I*a)^(1/2)
*(-I*a)^(1/2)*tan(d*x+c)^4*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-315*I*(I*
a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))
)^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^5-288*I*B*(I*a)^(
1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+456*B
*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/
2)+1608*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^4*(a*tan(d*x+c)*(1+I*tan(d
*x+c)))^(1/2)+1260*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x
+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^5-200*I*
A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2
)-315*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*ta
n(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^5+264*A*
(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2
)+630*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(
I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^5-630*ln(1/2*(2*I*a*t
an(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/
2))*(-I*a)^(1/2)*a*tan(d*x+c)^5+1260*I*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan
(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a
*tan(d*x+c)^5+488*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3*(a*tan(d*x+c)*
(1+I*tan(d*x+c)))^(1/2)-180*B*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1..

```

3.168.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 701 vs. $2(205) = 410$.

Time = 0.26 (sec) , antiderivative size = 701, normalized size of antiderivative = 2.61

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{1/2}(c + dx)} dx =$$

$$315 \sqrt{2} \sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^3}{d^2}} (de^{(10i dx + 10i c)} - 5 de^{(8i dx + 8i c)} + 10 de^{(6i dx + 6i c)} - 10 de^{(4i dx + 4i c)} + 5 de^{(2i dx + 2i c)})$$

```

input integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x, a
lgorithm="fracas")

```

3.168. $\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{1/2}(c + dx)} dx$

output

```
-1/315*(315*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(d*e^(10*I*d*x
+ 10*I*c) - 5*d*e^(8*I*d*x + 8*I*c) + 10*d*e^(6*I*d*x + 6*I*c) - 10*d*e^(
4*I*d*x + 4*I*c) + 5*d*e^(2*I*d*x + 2*I*c) - d)*log((sqrt(2)*sqrt(-(-I*A^2
- 2*A*B + I*B^2)*a^3/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a*e^(2*
I*d*x + 2*I*c) + (-I*A - B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*
e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I
*A - B)*a)) - 315*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(d*e^(10
*I*d*x + 10*I*c) - 5*d*e^(8*I*d*x + 8*I*c) + 10*d*e^(6*I*d*x + 6*I*c) - 10
*d*e^(4*I*d*x + 4*I*c) + 5*d*e^(2*I*d*x + 2*I*c) - d)*log(-(sqrt(2)*sqrt(-
(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*d*e^(I*d*x + I*c) - sqrt(2)*((-I*A - B)*
a*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sq
rt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*
c)/((-I*A - B)*a)) + 2*sqrt(2)*((659*I*A + 633*B)*a*e^(11*I*d*x + 11*I*c)
+ 7*(-127*I*A - 159*B)*a*e^(9*I*d*x + 9*I*c) + 18*(47*I*A + 29*B)*a*e^(7*I
*d*x + 7*I*c) + 42*(27*I*A + 19*B)*a*e^(5*I*d*x + 5*I*c) + 105*(-9*I*A - 1
1*B)*a*e^(3*I*d*x + 3*I*c) + 315*(I*A + B)*a*e^(I*d*x + I*c))*sqrt(a/(e^(2
*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*
c) + 1)))/(d*e^(10*I*d*x + 10*I*c) - 5*d*e^(8*I*d*x + 8*I*c) + 10*d*e^(6*I
*d*x + 6*I*c) - 10*d*e^(4*I*d*x + 4*I*c) + 5*d*e^(2*I*d*x + 2*I*c) - d)
```

3.168.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(11/2),x)`

output `Timed out`

3.168.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x, algorithm="maxima")`

output `Timed out`

3.168.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx);OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[-85]Warning, replacing -85 by 10, a substitution varia`

3.168.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) i)^{3/2}}{\tan(c + dx)^{11/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2))/tan(c + d*x)^(11/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2))/tan(c + d*x)^(11/2), x)`

3.168. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{11/2}(c+dx)} dx$

3.169 $\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.169.1 Optimal result	1751
3.169.2 Mathematica [A] (verified)	1752
3.169.3 Rubi [A] (verified)	1753
3.169.4 Maple [B] (verified)	1759
3.169.5 Fricas [B] (verification not implemented)	1760
3.169.6 Sympy [F(-1)]	1761
3.169.7 Maxima [F]	1762
3.169.8 Giac [F(-2)]	1762
3.169.9 Mupad [F(-1)]	1762

3.169.1 Optimal result

Integrand size = 38, antiderivative size = 298

$$\begin{aligned}
 & \int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \\
 & \frac{3(-1)^{3/4}a^{5/2}(120iA + 121B) \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{64d} \\
 & + \frac{(4 + 4i)a^{5/2}(iA + B)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} \\
 & + \frac{a^2(152A - 149iB)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{64d} \\
 & + \frac{a^2(104iA + 107B)\tan^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{96d} \\
 & - \frac{a^2(8A - 11iB)\tan^{\frac{5}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{24d} \\
 & + \frac{iaB \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{4d}
 \end{aligned}$$

output
$$\frac{3}{64}(-1)^{3/4}a^{5/2}(120IA+121B)\arctan((-1)^{3/4}a^{1/2}\tan(dx+c)^{1/2}/(a+Ia\tan(dx+c))^{1/2})/d+(4+4I)a^{5/2}(IA+B)\operatorname{arctanh}((1+I)a^{1/2}\tan(dx+c)^{1/2}/(a+Ia\tan(dx+c))^{1/2})/d+1/64a^2(152A-149IB)\tan(dx+c)^{1/2}(a+Ia\tan(dx+c))^{1/2}/d+1/96a^2(104IA+107B)(a+Ia\tan(dx+c))^{1/2}\tan(dx+c)^{3/2}/d-1/24a^2(8A-11IB)(a+Ia\tan(dx+c))^{1/2}\tan(dx+c)^{5/2}/d+1/4IaB\tan(dx+c)^{5/2}(a+Ia\tan(dx+c))^{3/2}/d$$

3.169.2 Mathematica [A] (verified)

Time = 7.26 (sec) , antiderivative size = 563, normalized size of antiderivative = 1.89

$$\int \tan^{\frac{3}{2}}(c+dx)(a + ia \tan(c+dx))^{5/2}(A + B \tan(c+dx)) dx = \frac{B \tan^{\frac{3}{2}}(c+dx)(a + ia \tan(c+dx))^{5/2}}{4d} + \frac{a(8A-5iB)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}}{6d} + \frac{ia^4(40iA+43B)\sqrt{a+ia \tan(c+dx)}\left(-\frac{3}{4}(-1)^{3/4}\operatorname{arcsinh}\left(\frac{\sqrt[4]{-1}\sqrt{\tan(c+dx)}}{\sqrt{1+i \tan(c+dx)}}\right)+\frac{5}{4}\sqrt{1+i \tan(c+dx)}\right)}{4d\sqrt{1+i \tan(c+dx)}}$$

input `Integrate[Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]), x]`

output
$$\frac{(B \tan[c + d*x]^{3/2}(a + I a \tan[c + d*x])^{5/2})/(4d) + ((a(8A - (5I)B)\sqrt{\tan[c + d*x]}(a + I a \tan[c + d*x])^{5/2})/(6d) + (((-1/4I) a^4((40I)A + 43B)\sqrt{a + I a \tan[c + d*x]}((-3(-1)^{3/4}\operatorname{ArcSinh}((-1)^{1/4}\sqrt{\tan[c + d*x]})))/4 + (5\sqrt{1 + I \tan[c + d*x]}\sqrt{\tan[c + d*x]})/4 + (I/2)\sqrt{1 + I \tan[c + d*x]}\tan[c + d*x]^{3/2})/(d\sqrt{1 + I \tan[c + d*x]}) + (a((-1/4I)a^3(8A - (5I)B) - (a^3((40I)A + 43B))/4)((-4I)\sqrt{2}a\operatorname{ArcTanh}(\sqrt{2}\sqrt{I a \tan[c + d*x]})/\sqrt{a + I a \tan[c + d*x]})\sqrt{\tan[c + d*x]}/\sqrt{I a \tan[c + d*x]} + ((4I)a^{3/2}\operatorname{ArcSinh}(\sqrt{I a \tan[c + d*x]}/\sqrt{a})\sqrt{1 + I \tan[c + d*x]})\sqrt{\tan[c + d*x]}/(\sqrt{I a \tan[c + d*x]}\sqrt{a + I a \tan[c + d*x]}) + I\sqrt{\tan[c + d*x]}\sqrt{a + I a \tan[c + d*x]} + (I\sqrt{a}\operatorname{ArcSinh}(\sqrt{I a \tan[c + d*x]}/\sqrt{a})\sqrt{\tan[c + d*x]}\sqrt{a + I a \tan[c + d*x]})/(\sqrt{1 + I \tan[c + d*x]}\sqrt{I a \tan[c + d*x]})))/d)/(3a))/(4a)$$

3.169.
$$\int \tan^{\frac{3}{2}}(c+dx)(a + ia \tan(c+dx))^{5/2}(A + B \tan(c+dx)) dx$$

3.169.3 Rubi [A] (verified)

Time = 2.04 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.07, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.526$, Rules used = {3042, 4077, 27, 3042, 4077, 27, 3042, 4080, 27, 3042, 4080, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c+dx)^{3/2}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4077} \\
 & \frac{1}{4} \int \frac{1}{2} \tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a)^{3/2}(a(8A-5iB)+a(8iA+11B) \tan(c+dx)) dx + \\
 & \quad \frac{iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{8} \int \tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a)^{3/2}(a(8A-5iB)+a(8iA+11B) \tan(c+dx)) dx + \\
 & \quad \frac{iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{8} \int \tan(c+dx)^{3/2}(i \tan(c+dx)a+a)^{3/2}(a(8A-5iB)+a(8iA+11B) \tan(c+dx)) dx + \\
 & \quad \frac{iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d} \\
 & \quad \downarrow \text{4077} \\
 & \frac{1}{8} \left(\frac{1}{3} \int \frac{1}{2} \tan^{\frac{3}{2}}(c+dx) \sqrt{i \tan(c+dx)a+a} ((88A-85iB)a^2+(104iA+107B) \tan(c+dx)a^2) dx - \frac{a^2(8A-11B)}{4d} \right. \\
 & \quad \left. + \frac{iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d} \right) \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.169. $\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\frac{1}{8} \left(\frac{1}{6} \int \tan^{\frac{3}{2}}(c+dx) \sqrt{i \tan(c+dx)a+a} ((88A-85iB)a^2 + (104iA+107B) \tan(c+dx)a^2) dx - \frac{a^2(8A-11iB)}{4d} \right) \frac{iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d}$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \int \tan(c+dx)^{3/2} \sqrt{i \tan(c+dx)a+a} ((88A-85iB)a^2 + (104iA+107B) \tan(c+dx)a^2) dx - \frac{a^2(8A-11iB)}{4d} \right) \frac{iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d}$$

↓ 4080

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{\int -\frac{3}{2} \sqrt{\tan(c+dx)} \sqrt{i \tan(c+dx)a+a} (a^3(104iA+107B) - a^3(152A-149iB) \tan(c+dx)) dx}{2a} + \frac{a^2(107B+104iA) \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{3 \int \sqrt{\tan(c+dx)} \sqrt{i \tan(c+dx)a+a} (a^3(104iA+107B) - a^3(152A-149iB) \tan(c+dx)) dx}{4a} \right) \frac{iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d} \right)$$

↓ 27

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107B+104iA) \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{3 \int \sqrt{\tan(c+dx)} \sqrt{i \tan(c+dx)a+a} (a^3(104iA+107B) - a^3(152A-149iB) \tan(c+dx)) dx}{4a} \right) \frac{iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d} \right)$$

↓ 3042

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107B+104iA) \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{3 \int \sqrt{\tan(c+dx)} \sqrt{i \tan(c+dx)a+a} (a^3(104iA+107B) - a^3(152A-149iB) \tan(c+dx)) dx}{4a} \right) \frac{iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d} \right)$$

↓ 4080

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107B+104iA) \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{2d} - \frac{3 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} ((152A-149iB)a^4 + 3(120iA+121B) \tan(c+dx))}{2\sqrt{\tan(c+dx)}}}{a} \right)}{4a} \right) \frac{iaB \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{4d} \right)$$

3.169. $\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\begin{aligned} & \downarrow 27 \\ & \frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107B + 104iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - 3 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((152A-149iB)a^4+3(120iA+121B) \tan(c+dx)}{\sqrt{\tan(c+dx)}}}{2a} \right) \right) \right) \end{aligned}$$

$$\frac{iaB \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{4d}$$

$\downarrow 3042$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107B + 104iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - 3 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((152A-149iB)a^4+3(120iA+121B) \tan(c+dx)}{\sqrt{\tan(c+dx)}}}{2a} \right) \right) \right)$$

$$\frac{iaB \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{4d}$$

$\downarrow 4084$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107B + 104iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - 3 \left(\frac{512a^4(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - 3a^3(120A-121iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} \right) \right) \right)$$

$$\frac{iaB \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{4d}$$

$\downarrow 3042$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107B + 104iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - 3 \left(\frac{512a^4(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - 3a^3(120A-121iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} \right) \right) \right)$$

$$\frac{iaB \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{4d}$$

$\downarrow 4027$

3.169. $\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107B + 104iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - 3 \left(\frac{-3a^3(120A - 121iB) \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + a}}{\sqrt{\tan(c + dx)}} dx}{2a} \right) \right) \right)$$

$$\frac{iaB \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{4d}$$

↓ 218

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107B + 104iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - 3 \left(\frac{(512 - 512i)a^{9/2}(A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right)}{d} - 3a^3 \right) \right) \right)$$

$$\frac{iaB \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{4d}$$

↓ 4082

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{a^2(107B + 104iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - 3 \left(\frac{(512 - 512i)a^{9/2}(A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right)}{d} - 3a^5 \right) \right) \right)$$

$$\frac{iaB \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{4d}$$

↓ 65

3.169. $\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\frac{1}{8} \left(\frac{1}{6} \frac{a^2(107B + 104iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(\frac{(512 - 512i)a^{9/2}(A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{6a^5}{2a} \right)}{4d} \right)$$

216

$$\frac{1}{8} \left(\frac{1}{6} \frac{a^2(107B + 104iA) \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{2d} - \frac{3 \left(\frac{{}^6\sqrt{-1}a^{9/2}(120A - 121iB) \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{6a^5}{2a} \right)}{4d} \right)$$

```
input Int[Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x
]
```

```
output ((I/4)*a*B*Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^(3/2))/d + (-1/3*(a^2
*(8*A - (11*I)*B)*Tan[c + d*x]^(5/2)*Sqrt[a + I*a*Tan[c + d*x]])/d + ((a^2
*((104*I)*A + 107*B)*Tan[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]])/(2*d)
- (3*((6*(-1)^(1/4)*a^(9/2)*(120*A - (121*I)*B)*ArcTan[(-1)^(3/4)*Sqrt[a
]*Sqrt[Tan[c + d*x]]]/Sqrt[a + I*a*Tan[c + d*x]]))/d + ((512 - 512*I)*a^(9
/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]]]/Sqrt[a + I*a*Ta
n[c + d*x]]])/d)/(2*a) - (a^3*(152*A - (149*I)*B)*Sqrt[Tan[c + d*x]]*Sqrt[
a + I*a*Tan[c + d*x]]/d)/(4*a))/6)/8
```

3.169.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4077 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

```
rule 4080 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[
1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Sim
p[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*T
an[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c
- a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]
```

```
rule 4082 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 + b^2, 0] && EqQ[A*b + a*B, 0]
```

```
rule 4084 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x]
- Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[
e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

3.169.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 739 vs. $2(240) = 480$.

Time = 0.17 (sec) , antiderivative size = 740, normalized size of antiderivative = 2.48

method	result
derivativedivides	$\frac{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})^2 a^2 \left(-96B\sqrt{ia}\sqrt{-ia}(\tan^3(dx+c))\sqrt{a\tan(dx+c)(1+i\tan(dx+c))} - 128A\sqrt{ia}\sqrt{-ia} \right)}{\dots}$
default	$\frac{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})^2 a^2 \left(-96B\sqrt{ia}\sqrt{-ia}(\tan^3(dx+c))\sqrt{a\tan(dx+c)(1+i\tan(dx+c))} - 128A\sqrt{ia}\sqrt{-ia} \right)}{\dots}$
parts	Expression too large to display

```
input int(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

$$3.169. \quad \int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

output

```

1/384/d*tan(d*x+c)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)*a^2*(-96*B*(I*a)^(1/2)
*(-I*a)^(1/2)*tan(d*x+c)^3*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-128*A*(I*
a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+2
72*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)
))^(1/2)+416*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(
1/2)*tan(d*x+c)+447*I*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan
(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a-894*I*B*(I*a)^(
1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+428*B*(I*a)^(1/2)*
(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)-384*I*(I*a)^(
1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(
1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a-456*A*ln(1/2*(2*I*a*tan(d*x+c)+
2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)
^(1/2)*a+912*A*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1
/2)-768*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a+384*(I*a)^(1/2)*2^(1/2)*ln((2*
2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x
+c))/(tan(d*x+c)+I))*a-768*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*t
an(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a/(I*a)^(1/2)/
(-I*a)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)

```

3.169.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1017 vs. $2(224) = 448$.

Time = 0.32 (sec) , antiderivative size = 1017, normalized size of antiderivative = 3.41

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input

```

integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, al
gorithm="fricas")

```

output `1/384*(768*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*log((I*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 768*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*log((-I*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) + 2*sqrt(2)*(13*(56*A - 65*I*B)*a^2*e^(7*I*d*x + 7*I*c) + 3*(504*A - 425*I*B)*a^2*e^(5*I*d*x + 5*I*c) + (1096*A - 1135*I*B)*a^2*e^(3*I*d*x + 3*I*c) + 3*(104*A - 107*I*B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) - 9*sqrt(-(14400*I*A^2 + 29040*A*B - 14641*I*B^2)*a^5/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*log((sqrt(2)*((-120*I*A - 121*B)*a^2*e^(2*I*d*x + 2*I*c) + (-120*I*A - 121*B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + 2*I*sqrt(-(14400*I*A^2 + 29040*A*B - 14641*I*B^2)*a^5/d^2)*d*e^(I*...`

3.169.6 Sympy [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(3/2)*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.169.7 Maxima [F]

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \int (B \tan(dx+c) + A)(ia \tan(dx+c) + a)^{\frac{5}{2}} \tan(dx+c)^{\frac{3}{2}} dx$$

input `integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)*tan(d*x + c)^(3/2), x)`

3.169.8 Giac [F(-2)]

Exception generated.

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \text{Exception raised: TypeError}$$

input `integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:The choice was done assuming 0=[0]Warning, replacing 0 by -28, a substitution variable should perhaps be purged.Warnin`

3.169.9 Mupad [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \int \tan(c+dx)^{3/2} (A+B \tan(c+dx)) (a+a \tan(c+dx) i)^{5/2} dx$$

input `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),
x)`

output `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),
x)`

3.170 $\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.170.1 Optimal result	1764
3.170.2 Mathematica [A] (verified)	1765
3.170.3 Rubi [A] (verified)	1765
3.170.4 Maple [B] (verified)	1770
3.170.5 Fricas [B] (verification not implemented)	1771
3.170.6 Sympy [F(-1)]	1772
3.170.7 Maxima [F]	1773
3.170.8 Giac [F(-2)]	1773
3.170.9 Mupad [F(-1)]	1773

3.170.1 Optimal result

Integrand size = 38, antiderivative size = 252

$$\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx =$$

$$\frac{(-1)^{3/4} a^{5/2} (46A - 45iB) \arctan\left(\frac{(-1)^{3/4} \sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{8d}$$

$$- \frac{(4 + 4i)a^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$+ \frac{a^2(18iA + 19B) \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}}{8d}$$

$$- \frac{a^2(2A - 3iB) \tan^{3/2}(c + dx) \sqrt{a + ia \tan(c + dx)}}{4d}$$

$$+ \frac{iaB \tan^{3/2}(c + dx)(a + ia \tan(c + dx))^{3/2}}{3d}$$

output

```
-1/8*(-1)^(3/4)*a^(5/2)*(46*A-45*I*B)*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)
^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d-(4+4*I)*a^(5/2)*(A-I*B)*arctanh((1+I)*a
^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d+1/8*a^2*(18*I*A+19*B)*
tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/d-1/4*a^2*(2*A-3*I*B)*(a+I*a*tan
(d*x+c)^(1/2)*tan(d*x+c)^(3/2)/d+1/3*I*a*B*tan(d*x+c)^(3/2)*(a+I*a*tan(d*
x+c))^(3/2)/d
```

3.170.2 Mathematica [A] (verified)

Time = 6.78 (sec) , antiderivative size = 503, normalized size of antiderivative = 2.00

$$\int \sqrt{\tan(c+dx)}(a + ia \tan(c+dx))^{5/2}(A + B \tan(c+dx)) dx = \frac{B \sqrt{\tan(c+dx)}(a + ia \tan(c+dx))^{5/2}}{3d}$$

$$+ \frac{ia^3(6A-5iB)\sqrt{a+ia \tan(c+dx)}\left(-\frac{3}{4}(-1)^{3/4}\operatorname{arcsinh}\left(\sqrt[4]{-1}\sqrt{\tan(c+dx)}\right)+\frac{5}{4}\sqrt{1+i \tan(c+dx)}\sqrt{\tan(c+dx)}+\frac{1}{2}i\sqrt{1+i \tan(c+dx)}\tan^{\frac{3}{2}}(c+dx)\right)}{2d\sqrt{1+i \tan(c+dx)}}$$

input `Integrate[Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(B*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(5/2))/(3*d) + (((I/2)*a^3*(6*A - (5*I)*B)*Sqrt[a + I*a*Tan[c + d*x]]*((-3*(-1)^(3/4)*ArcSinh[(-1)^(1/4)]*Sqrt[Tan[c + d*x]]))/4 + (5*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/4 + (I/2)*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(3/2))/(d*Sqrt[1 + I*Tan[c + d*x]]) + (a*((a^2*(6*A - (5*I)*B))/2 - (I/2)*a^2*B)*(((4*I)*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[Tan[c + d*x]])/Sqrt[I*a*Tan[c + d*x]] + ((4*I)*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/(Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + I*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]] + (I*Sqrt[a]*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]]))/d)/(3*a)`

3.170.3 Rubi [A] (verified)Time = 1.57 (sec) , antiderivative size = 265, normalized size of antiderivative = 1.05, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.447$, Rules used = {3042, 4077, 27, 3042, 4077, 27, 3042, 4080, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\tan(c+dx)}(a + ia \tan(c+dx))^{5/2}(A + B \tan(c+dx)) dx$$

3.170. $\int \sqrt{\tan(c+dx)}(a + ia \tan(c+dx))^{5/2}(A + B \tan(c+dx)) dx$

$$\begin{aligned}
 & \int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{3} \int \frac{3}{2} \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^{3/2}(a(2A-iB)+a(2iA+3B) \tan(c+dx))dx + \\
 & \quad \frac{iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{4077} \\
 & \frac{1}{2} \int \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^{3/2}(a(2A-iB)+a(2iA+3B) \tan(c+dx))dx + \\
 & \quad \frac{iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2} \int \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^{3/2}(a(2A-iB)+a(2iA+3B) \tan(c+dx))dx + \\
 & \quad \frac{iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \int \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^{3/2}(a(2A-iB)+a(2iA+3B) \tan(c+dx))dx + \\
 & \quad \frac{iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{4077} \\
 & \frac{1}{2} \left(\frac{1}{2} \int \frac{1}{2} \sqrt{\tan(c+dx)} \sqrt{i \tan(c+dx)a+a} ((14A-13iB)a^2+(18iA+19B) \tan(c+dx)a^2) dx - \frac{a^2(2A-3iB)}{3d} \right) \\
 & \quad \frac{iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{2} \left(\frac{1}{4} \int \sqrt{\tan(c+dx)} \sqrt{i \tan(c+dx)a+a} ((14A-13iB)a^2+(18iA+19B) \tan(c+dx)a^2) dx - \frac{a^2(2A-3iB)}{3d} \right) \\
 & \quad \frac{iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2} \left(\frac{1}{4} \int \sqrt{\tan(c+dx)} \sqrt{i \tan(c+dx)a+a} ((14A-13iB)a^2+(18iA+19B) \tan(c+dx)a^2) dx - \frac{a^2(2A-3iB)}{3d} \right) \\
 & \quad \frac{iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{4080}
 \end{aligned}$$

3.170. $\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{\int -\frac{\sqrt{i \tan(c+dx)a+a}(a^3(18iA+19B)-a^3(46A-45iB) \tan(c+dx))}{2\sqrt{\tan(c+dx)}} dx}{a} + \frac{a^2(19B+18iA)\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}{d} \right) - \frac{iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d} \right)$$

↓ 27

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19B+18iA)\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}{d} - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a^3(18iA+19B)-a^3(46A-45iB) \tan(c+dx))}{\sqrt{\tan(c+dx)}}}{2a} \right) - \frac{iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d} \right)$$

↓ 3042

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19B+18iA)\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}{d} - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a^3(18iA+19B)-a^3(46A-45iB) \tan(c+dx))}{\sqrt{\tan(c+dx)}}}{2a} \right) - \frac{iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d} \right)$$

↓ 4084

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19B+18iA)\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}{d} - \frac{64a^3(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - a^2(45B+46iA)}{2a} \right) - \frac{iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d} \right)$$

↓ 3042

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19B+18iA)\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}{d} - \frac{64a^3(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - a^2(45B+46iA)}{2a} \right) - \frac{iaB \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}{3d} \right)$$

↓ 4027

3.170. $\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19B + 18iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{-a^2(45B + 46iA) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} \right) \right. \\ \left. - \frac{iaB \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{3d} \right) \downarrow 218$$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19B + 18iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{(64-64i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - a^2(45B + 46iA) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} \right) \right. \\ \left. - \frac{iaB \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{3d} \right) \downarrow 4082$$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19B + 18iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{(64-64i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{a^4(45B + 46iA)}{2a} \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} \right) \right. \\ \left. - \frac{iaB \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{3d} \right) \downarrow 65$$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19B + 18iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{(64-64i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^4(45B + 46iA)}{2a} \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} \right) \right. \\ \left. - \frac{iaB \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{3d} \right) \downarrow 216$$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{a^2(19B + 18iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2\sqrt{-1}a^{7/2}(45B+46iA) \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{(64-64i)a^4(45B + 46iA)}{2a} \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} \right) \right. \\ \left. - \frac{iaB \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{3d} \right)$$

input `Int[Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `((I/3)*a*B*Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(3/2))/d + (-1/2*(a^2*(2*A - (3*I)*B)*Tan[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]])/d + (-1/2*((2*(-1)^(1/4)*a^(7/2)*((46*I)*A + 45*B)*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d + ((64 - 64*I)*a^(7/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d)/a + (a^2*((18*I)*A + 19*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/d)/4)/2`

3.170.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`


```
rule 4077 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan
[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c -
a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] &&
GtQ[m, 1] && !LtQ[n, -1]
```

```
rule 4080 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[
1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Sim
p[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*T
an[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c
- a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]
```

```
rule 4082 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 + b^2, 0] && EqQ[A*b + a*B, 0]
```

```
rule 4084 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x]
- Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[
e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

3.170.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 650 vs. $2(202) = 404$.

Time = 0.15 (sec) , antiderivative size = 651, normalized size of antiderivative = 2.58

method	result
derivativedivides	$\frac{(\sqrt{\tan(dx+c)} \sqrt{a(1+i \tan(dx+c))} a^2 \left(16B\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} (\tan^2(dx+c)) - 52iB\sqrt{ia} \sqrt{-ia} \right)}{\dots}$
default	$\frac{(\sqrt{\tan(dx+c)} \sqrt{a(1+i \tan(dx+c))} a^2 \left(16B\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} (\tan^2(dx+c)) - 52iB\sqrt{ia} \sqrt{-ia} \right)}{\dots}$
parts	$\frac{A(\sqrt{\tan(dx+c)} \sqrt{a(1+i \tan(dx+c))} a^2 \left(18i\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - 4\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \right)}{\dots}$

```
input int(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_R
RETURNVERBOSE)
```

```
output -1/48/d*tan(d*x+c)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)*a^2*(16*B*(I*a)^(1/2)*
(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2-52*I*B*(I*
a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+54*
I*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)
^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a-108*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*(a*
tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+24*A*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*
x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)-48*I*(I*a)^(1/2)*2^(1/2)*ln((2*2^(
1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c)
)/(tan(d*x+c)+I))*a+57*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan
(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a-114*B*(I*a)^(1/
2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+96*I*ln(1/2*(2*I*a*t
an(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/
2))*(-I*a)^(1/2)*a-48*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*ta
n(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a-96*
ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/
2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a)/(I*a)^(1/2)/(-I*a)^(1/2)/(a*tan(d*x+c)*
(1+I*tan(d*x+c)))^(1/2)
```

3.170.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 932 vs. 2(188) = 376.

Time = 0.27 (sec) , antiderivative size = 932, normalized size of antiderivative = 3.70

$$\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/48*(96*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log((sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 96*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log(-(sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) - sqrt(2)*((-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a^2)) + 2*sqrt(2)*((66*I*A + 91*B)*a^2*e^(5*I*d*x + 5*I*c) - 2*(-54*I*A - 49*B)*a^2*e^(3*I*d*x + 3*I*c) - 3*(-14*I*A - 13*B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + 3*sqrt((2116*I*A^2 + 4140*A*B - 2025*I*B^2)*a^5/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log((sqrt(2)*((46*I*A + 45*B)*a^2*e^(2*I*d*x + 2*I*c) + (46*I*A + 45*B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + 2*sqrt((2116*I*A^2 + 4140*A*B - 2025*I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/((46*I*A + 45*B)*a^2)) - 3*sqrt((2116*I*A^2 + 4140*A*B - 2025*I*B^2)*a^5/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + ...`

3.170.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(1/2)*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.170.7 Maxima [F]

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \int (B \tan(dx+c)+A)(ia \tan(dx+c)+a)^{5/2} \sqrt{\tan(dx+c)} dx$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x+c)+A)*(I*a*tan(d*x+c)+a)^(5/2)*sqrt(tan(d*x+c)), x)`

3.170.8 Giac [F(-2)]

Exception generated.

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \text{Exception raised: TypeError}$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:The choice was done assuming 0=[0]Warning, replacing 0 by 51, a substitution variable should perhaps be purged.Warning`

3.170.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \int \sqrt{\tan(c+dx)}(A+B \tan(c+dx))(a+a \tan(c+dx) li)^{5/2} dx$$

input `int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),
x)`

output `int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),
x)`

3.171
$$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.171.1 Optimal result 1775
 3.171.2 Mathematica [B] (verified) 1776
 3.171.3 Rubi [A] (verified) 1776
 3.171.4 Maple [B] (verified) 1781
 3.171.5 Fricas [B] (verification not implemented) 1782
 3.171.6 Sympy [F] 1783
 3.171.7 Maxima [F(-1)] 1783
 3.171.8 Giac [F(-2)] 1783
 3.171.9 Mupad [F(-1)] 1784

3.171.1 Optimal result

Integrand size = 38, antiderivative size = 206

$$\int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx =$$

$$\frac{(-1)^{3/4}a^{5/2}(20iA + 23B) \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{4d}$$

$$+ \frac{(4 - 4i)a^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$- \frac{a^2(4A - 7iB)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{4d}$$

$$+ \frac{iaB\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}}{2d}$$

```
output -1/4*(-1)^(3/4)*a^(5/2)*(20*I*A+23*B)*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)
^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d+(4-4*I)*a^(5/2)*(A-I*B)*arctanh((1+I)*a
^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d-1/4*a^2*(4*A-7*I*B)*ta
n(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/d+1/2*I*a*B*tan(d*x+c)^(1/2)*(a+I*
a*tan(d*x+c))^(3/2)/d
```

3.171.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 433 vs. $2(206) = 412$.

Time = 10.69 (sec) , antiderivative size = 433, normalized size of antiderivative = 2.10

$$\int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \frac{ia^2 B \sqrt{a + ia \tan(c + dx)} \left(-\frac{3}{4}(-1)^{3/4} \operatorname{arcsinh}\left(\sqrt[4]{-1} \sqrt{\tan(c + dx)}\right)\right)}{d} + \frac{a(iaA + aB) \left(-\frac{4i\sqrt{2}a \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right) \sqrt{\tan(c + dx)}}{\sqrt{ia \tan(c + dx)}} + \frac{4ia^{3/2} \operatorname{arcsinh}\left(\frac{\sqrt{ia \tan(c + dx)}}{\sqrt{a}}\right) \sqrt{1 + i \tan(c + dx)} \sqrt{\tan(c + dx)}}{\sqrt{ia \tan(c + dx)} \sqrt{a + ia \tan(c + dx)}}\right)}{d}$$

```
input Integrate[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]], x]
```

```
output (I*a^2*B*Sqrt[a + I*a*Tan[c + d*x]]*((-3*(-1)^(3/4)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]])/4 + (5*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/4 + (I/2)*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(3/2))/(d*Sqrt[1 + I*Tan[c + d*x]]) + (a*(I*a*A + a*B)*(((4*I)*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]]*Sqrt[Tan[c + d*x]])/Sqrt[I*a*Tan[c + d*x]] + ((4*I)*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/(Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + I*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]] + (I*Sqrt[a]*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]]))/d
```

3.171.3 Rubi [A] (verified)

Time = 1.25 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.03, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4077, 27, 3042, 4077, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \quad \downarrow \quad 3042$$

3.171. $\int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$

$$\begin{aligned}
& \int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
& \quad \downarrow \text{4077} \\
& \frac{1}{2} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(4A - iB) + a(4iA + 7B) \tan(c + dx))}{2\sqrt{\tan(c + dx)}} dx + \\
& \quad \frac{iaB \sqrt{\tan(c + dx)} (a + ia \tan(c + dx))^{3/2}}{2d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{4} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(4A - iB) + a(4iA + 7B) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx + \\
& \quad \frac{iaB \sqrt{\tan(c + dx)} (a + ia \tan(c + dx))^{3/2}}{2d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{4} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(4A - iB) + a(4iA + 7B) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx + \\
& \quad \frac{iaB \sqrt{\tan(c + dx)} (a + ia \tan(c + dx))^{3/2}}{2d} \\
& \quad \downarrow \text{4077} \\
& \frac{1}{4} \left(\int \frac{\sqrt{i \tan(c + dx)a + a} (3(4A - 3iB)a^2 + (20iA + 23B) \tan(c + dx)a^2)}{2\sqrt{\tan(c + dx)}} dx - \frac{a^2(4A - 7iB) \sqrt{\tan(c + dx)} \sqrt{a}}{d} \right. \\
& \quad \left. \frac{iaB \sqrt{\tan(c + dx)} (a + ia \tan(c + dx))^{3/2}}{2d} \right) \\
& \quad \downarrow \text{27} \\
& \frac{1}{4} \left(\frac{1}{2} \int \frac{\sqrt{i \tan(c + dx)a + a} (3(4A - 3iB)a^2 + (20iA + 23B) \tan(c + dx)a^2)}{\sqrt{\tan(c + dx)}} dx - \frac{a^2(4A - 7iB) \sqrt{\tan(c + dx)} \sqrt{a}}{d} \right. \\
& \quad \left. \frac{iaB \sqrt{\tan(c + dx)} (a + ia \tan(c + dx))^{3/2}}{2d} \right) \\
& \quad \downarrow \text{3042} \\
& \frac{1}{4} \left(\frac{1}{2} \int \frac{\sqrt{i \tan(c + dx)a + a} (3(4A - 3iB)a^2 + (20iA + 23B) \tan(c + dx)a^2)}{\sqrt{\tan(c + dx)}} dx - \frac{a^2(4A - 7iB) \sqrt{\tan(c + dx)} \sqrt{a}}{d} \right. \\
& \quad \left. \frac{iaB \sqrt{\tan(c + dx)} (a + ia \tan(c + dx))^{3/2}}{2d} \right) \\
& \quad \downarrow \text{4084}
\end{aligned}$$

3.171. $\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$

$$\frac{1}{4} \left(\frac{1}{2} \left(32a^2(A - iB) \int \frac{\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx - a(20A - 23iB) \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx \right) \right. \\ \left. \frac{iaB \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}}{2d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{4} \left(\frac{1}{2} \left(32a^2(A - iB) \int \frac{\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx - a(20A - 23iB) \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx \right) \right. \\ \left. \frac{iaB \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}}{2d} \right) \\ \downarrow \text{4027}$$

$$\frac{1}{4} \left(\frac{1}{2} \left(-\frac{64ia^4(A - iB) \int \frac{1}{\frac{2 \tan(c + dx)a^2}{i \tan(c + dx)a + a} - ia} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{i \tan(c + dx)a + a}} - a(20A - 23iB) \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx \right) \right. \\ \left. \frac{iaB \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}}{2d} \right) \\ \downarrow \text{218}$$

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{(32 - 32i)a^{5/2}(A - iB) \operatorname{arctanh} \left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}} \right)}{d} - a(20A - 23iB) \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx \right) \right. \\ \left. \frac{iaB \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}}{2d} \right) \\ \downarrow \text{4082}$$

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{(32 - 32i)a^{5/2}(A - iB) \operatorname{arctanh} \left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}} \right)}{d} - \frac{a^3(20A - 23iB) \int \frac{1}{\sqrt{\tan(c + dx)}\sqrt{i \tan(c + dx)a + a}} d \tan(c + dx)}{d} \right) \right. \\ \left. \frac{iaB \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}}{2d} \right) \\ \downarrow \text{65}$$

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{(32 - 32i)a^{5/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} - \frac{2a^3(20A - 23iB) \int \frac{1}{1 - \frac{ia\tan(c+dx)}{i\tan(c+dx)a+a}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{d} \right) - \frac{iaB\sqrt{\tan(c+dx)}(a + ia\tan(c+dx))^{3/2}}{2d} \right)$$

↓ 216

$$\frac{1}{4} \left(\frac{1}{2} \left(\frac{2\sqrt[4]{-1}a^{5/2}(20A - 23iB) \operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} + \frac{(32 - 32i)a^{5/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} \right) - \frac{iaB\sqrt{\tan(c+dx)}(a + ia\tan(c+dx))^{3/2}}{2d} \right)$$

```
input Int[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]]
,x]
```

```
output ((I/2)*a*B*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(3/2))/d + (((2*(-1)^(1/4)*a^(5/2)*(20*A - (23*I)*B)*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d + ((32 - 32*I)*a^(5/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d)/2 - (a^2*(4*A - (7*I)*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]/d)/4
```

3.171.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x_, x] /; FreeQ[a, x] && !MatchQ[F_x_, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 65 Int[1/(Sqrt[(b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

3.171. $\int \frac{(a+ia\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

rule 218 $\text{Int}[(a_+) + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a) \cdot \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] \text{ ; FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

rule 3042 $\text{Int}[u_+, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

rule 4027 $\text{Int}[\text{Sqrt}[(a_+) + (b_+) \cdot \tan[(e_+) + (f_+)(x_+)]] / \text{Sqrt}[(c_+) + (d_+) \cdot \tan[(e_+) + (f_+)(x_+)]], x_Symbol] \rightarrow \text{Simp}[-2 \cdot a \cdot (b/f) \text{ Subst}[\text{Int}[1/(a \cdot c - b \cdot d - 2 \cdot a^2 \cdot x^2), x], x, \text{Sqrt}[c + d \cdot \tan[e + f \cdot x]] / \text{Sqrt}[a + b \cdot \tan[e + f \cdot x]]], x] \text{ ; FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0]$

rule 4077 $\text{Int}[(a_+) + (b_+) \cdot \tan[(e_+) + (f_+)(x_+)])^{(m_+)} \cdot ((A_+) + (B_+) \cdot \tan[(e_+) + (f_+)(x_+)])^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[b \cdot B \cdot (a + b \cdot \tan[e + f \cdot x])^{(m-1)} \cdot (c + d \cdot \tan[e + f \cdot x])^{(n+1)} / (d \cdot f \cdot (m+n)), x] + \text{Simp}[1/(d \cdot (m+n)) \text{ Int}[(a + b \cdot \tan[e + f \cdot x])^{(m-1)} \cdot (c + d \cdot \tan[e + f \cdot x])^n \cdot \text{Simp}[a \cdot A \cdot d \cdot (m+n) + B \cdot (a \cdot c \cdot (m-1) - b \cdot d \cdot (n+1)) - (B \cdot (b \cdot c - a \cdot d) \cdot (m-1) - d \cdot (A \cdot b + a \cdot B) \cdot (m+n)) \cdot \tan[e + f \cdot x], x], x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B, n\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{!LtQ}[n, -1]$

rule 4082 $\text{Int}[(a_+) + (b_+) \cdot \tan[(e_+) + (f_+)(x_+)])^{(m_+)} \cdot ((A_+) + (B_+) \cdot \tan[(e_+) + (f_+)(x_+)])^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[b \cdot (B/f) \text{ Subst}[\text{Int}[(a + b \cdot x)^{(m-1)} \cdot (c + d \cdot x)^n, x], x, \tan[e + f \cdot x]], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B, m, n\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[A \cdot b + a \cdot B, 0]$

rule 4084 $\text{Int}[(a_+) + (b_+) \cdot \tan[(e_+) + (f_+)(x_+)])^{(m_+)} \cdot ((A_+) + (B_+) \cdot \tan[(e_+) + (f_+)(x_+)])^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[(A \cdot b + a \cdot B)/b \text{ Int}[(a + b \cdot \tan[e + f \cdot x])^m \cdot (c + d \cdot \tan[e + f \cdot x])^n, x], x] - \text{Simp}[B/b \text{ Int}[(a + b \cdot \tan[e + f \cdot x])^m \cdot (c + d \cdot \tan[e + f \cdot x])^n \cdot (a - b \cdot \tan[e + f \cdot x]), x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B, m, n\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[A \cdot b + a \cdot B, 0]$

3.171.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 563 vs. $2(164) = 328$.

Time = 0.14 (sec) , antiderivative size = 564, normalized size of antiderivative = 2.74

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} (\sqrt{\tan(dx+c)}) a^2 \left(-9iB \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a+18iB\sqrt{ia}}{\dots} \right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))} (\sqrt{\tan(dx+c)}) a^2 \left(-9iB \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a+18iB\sqrt{ia}}{\dots} \right)}{\dots}$
parts	$\frac{A\sqrt{a(1+i \tan(dx+c))} (\sqrt{\tan(dx+c)}) a^2 \left(-2\sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) \sqrt{ia} \sqrt{-ia}-5 \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)}}{2\sqrt{ia}} \right) \right)}{\dots}$

```
input int((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x,method=_R
RETURNVERBOSE)
```

```
output 1/8/d*(a*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^(1/2)*a^2*(-9*I*B*ln(1/2*(2*I*
a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(
1/2))*(-I*a)^(1/2)*a+18*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*t
an(d*x+c)))^(1/2)-4*B*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+
c)))^(1/2)*tan(d*x+c)+8*I*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(
a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a
+12*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*
a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a-8*A*(I*a)^(1/2)*(-I*a)^(1/2)*(a*ta
n(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+16*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*
x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*(-I*a)^(1/2)-8*
(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+
c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a+16*ln(1/2*(2*I*a*tan(d*x+
c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I
*a)^(1/2)*a)/(I*a)^(1/2)/(-I*a)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2
)
```

$$3.171. \int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.171.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 849 vs. $2(152) = 304$.

Time = 0.28 (sec) , antiderivative size = 849, normalized size of antiderivative = 4.12

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Too large to display}$$

```
input integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="fricas")
```

```
output -1/8*(16*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log((I*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 16*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log((-I*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a^2)) + 2*sqrt(2)*((4*A - 11*I*B)*a^2*e^(3*I*d*x + 3*I*c) + (4*A - 7*I*B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + sqrt((-400*I*A^2 - 920*A*B + 529*I*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log((sqrt(2)*((20*I*A + 23*B)*a^2*e^(2*I*d*x + 2*I*c) + (20*I*A + 23*B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + 2*I*sqrt((-400*I*A^2 - 920*A*B + 529*I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c))e^(-I*d*x - I*c)/((20*I*A + 23*B)*a^2)) - sqrt((-400*I*A^2 - 920*A*B + 529*I*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log((sqrt(2)*((20*I*A + 23*B)*a^2*e^(2*I*d*x + 2*I*c) + (20*I*A + 23*B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2...))
```

3.171.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \int \frac{(ia(\tan(c + dx) - i))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

input `integrate((a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(1/2),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(5/2)*(A + B*tan(c + d*x))/sqrt(tan(c + d*x)), x)`

3.171.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="maxima")`

output `Timed out`

3.171.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx);;OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[57]Warning, replacing 57 by 18, a substitution variabl`

3.171. $\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$

3.171.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) 1i)^{5/2}}{\sqrt{\tan(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(1/2), x)`

3.172
$$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.172.1 Optimal result 1785
 3.172.2 Mathematica [B] (verified) 1786
 3.172.3 Rubi [A] (verified) 1786
 3.172.4 Maple [B] (verified) 1791
 3.172.5 Fricas [B] (verification not implemented) 1792
 3.172.6 Sympy [F] 1793
 3.172.7 Maxima [F(-1)] 1793
 3.172.8 Giac [F(-2)] 1793
 3.172.9 Mupad [F(-1)] 1794

3.172.1 Optimal result

Integrand size = 38, antiderivative size = 196

$$\int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \frac{(-1)^{3/4}a^{5/2}(2A - 5iB) \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$+ \frac{(4 + 4i)a^{5/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$+ \frac{a^2(2iA - B)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}}$$

output

```
(-1)^(3/4)*a^(5/2)*(2*A-5*I*B)*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d+(4+4*I)*a^(5/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d+a^2*(2*I*A-B)*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/d-2*a*A*(a+I*a*tan(d*x+c))^(3/2)/d/tan(d*x+c)^(1/2)
```


3.172.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 485 vs. $2(196) = 392$.

Time = 7.22 (sec) , antiderivative size = 485, normalized size of antiderivative = 2.47

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = -\frac{2A(a + ia \tan(c + dx))^{5/2}}{d\sqrt{\tan(c + dx)}} + 2 \left(\frac{2ia^3 A \sqrt{a + ia \tan(c + dx)} \left(-\frac{3}{4} (-1)^{3/4} \operatorname{arcsinh} \left(\sqrt[4]{-1} \sqrt{\tan(c + dx)} \right) + \frac{5}{4} \sqrt{1 + i \tan(c + dx)} \sqrt{\tan(c + dx)} + \frac{1}{2} i \sqrt{1 + i \tan(c + dx)} \tan^{3/2}(c + dx) \right)}{d\sqrt{1 + i \tan(c + dx)}} \right)$$

input `Integrate[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `(-2*A*(a + I*a*Tan[c + d*x])^(5/2))/(d*Sqrt[Tan[c + d*x]]) + (2*(((2*I)*a^3*A*Sqrt[a + I*a*Tan[c + d*x]]*((-3*(-1)^(3/4)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]])]/4 + (5*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/4 + (I/2)*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(3/2)))/(d*Sqrt[1 + I*Tan[c + d*x]]) + (a*(2*a^2*A + (I/2)*a^2*((5*I)*A + B))*((-4*I)*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/Sqrt[I*a*Tan[c + d*x]] + ((4*I)*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/(Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + I*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]] + (I*Sqrt[a]*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])))/d)/a`

3.172.3 Rubi [A] (verified)

Time = 1.24 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.03, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4076, 27, 3042, 4077, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.172. $\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx$

$$\begin{aligned}
& \int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx \\
& \quad \downarrow \text{4076} \\
& 2 \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(4iA + B) + a(2A + iB) \tan(c + dx))}{2\sqrt{\tan(c + dx)} \frac{2aA(a + ia \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}}} dx - \\
& \quad \downarrow \text{27} \\
& \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(4iA + B) + a(2A + iB) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(4iA + B) + a(2A + iB) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}} \\
& \quad \downarrow \text{4077} \\
& \int \frac{\sqrt{i \tan(c + dx)a + a} (3a^2(2iA + B) - a^2(2A - 5iB) \tan(c + dx))}{2\sqrt{\tan(c + dx)}} dx + \\
& \frac{a^2(-B + 2iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}} \\
& \quad \downarrow \text{27} \\
& \frac{1}{2} \int \frac{\sqrt{i \tan(c + dx)a + a} (3a^2(2iA + B) - a^2(2A - 5iB) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx + \\
& \frac{a^2(-B + 2iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \int \frac{\sqrt{i \tan(c + dx)a + a} (3a^2(2iA + B) - a^2(2A - 5iB) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx + \\
& \frac{a^2(-B + 2iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}} \\
& \quad \downarrow \text{4084}
\end{aligned}$$

3.172. $\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx$

$$\frac{1}{2} \left(8a^2(B + iA) \int \frac{\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx - a(5B + 2iA) \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx \right) + \frac{a^2(-B + 2iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}}$$

↓ 3042

$$\frac{1}{2} \left(8a^2(B + iA) \int \frac{\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx - a(5B + 2iA) \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx \right) + \frac{a^2(-B + 2iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}}$$

↓ 4027

$$\frac{1}{2} \left(\frac{16ia^4(B + iA) \int \frac{1}{\frac{2 \tan(c + dx)a^2}{i \tan(c + dx)a + a} - ia} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{i \tan(c + dx)a + a}} - a(5B + 2iA) \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx \right) + \frac{a^2(-B + 2iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}}$$

↓ 218

$$\frac{1}{2} \left(\frac{(8 - 8i)a^{5/2}(B + iA) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right)}{d} - a(5B + 2iA) \int \frac{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx \right) + \frac{a^2(-B + 2iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}}$$

↓ 4082

$$\frac{1}{2} \left(\frac{(8 - 8i)a^{5/2}(B + iA) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right)}{d} - \frac{a^3(5B + 2iA) \int \frac{1}{\sqrt{\tan(c + dx)}\sqrt{i \tan(c + dx)a + a}} d \tan(c + dx)}{d} \right) + \frac{a^2(-B + 2iA)\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}}$$

↓ 65

3.172. $\int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$

$$\frac{1}{2} \left(\frac{(8-8i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} - \frac{2a^3(5B+2iA) \int \frac{1}{1-\frac{ia\tan(c+dx)}{i\tan(c+dx)a+a}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}}{d} \right) +$$

$$\frac{a^2(-B+2iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{2aA(a+ia\tan(c+dx))^{3/2}}{d\sqrt{\tan(c+dx)}}$$

↓ 216

$$\frac{1}{2} \left(\frac{2\sqrt[4]{-1}a^{5/2}(5B+2iA)\operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} + \frac{(8-8i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} \right) +$$

$$\frac{a^2(-B+2iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{2aA(a+ia\tan(c+dx))^{3/2}}{d\sqrt{\tan(c+dx)}}$$

input `Int[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2), x]`

output `((2*(-1)^(1/4)*a^(5/2)*((2*I)*A + 5*B)*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]]]/Sqrt[a + I*a*Tan[c + d*x]])/d + ((8 - 8*I)*a^(5/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]]]/Sqrt[a + I*a*Tan[c + d*x]])/d)/2 + (a^2*((2*I)*A - B)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/d - (2*a*A*(a + I*a*Tan[c + d*x])^(3/2))/(d*Sqrt[Tan[c + d*x]])`

3.172.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

- rule 218 $\text{Int}[(a_+ + (b_-)(x_-)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a) \cdot \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] \text{ ; FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$
- rule 4027 $\text{Int}[\text{Sqrt}[(a_+ + (b_-)\tan[(e_-) + (f_-)(x_-)])]/\text{Sqrt}[(c_-) + (d_-)\tan[(e_-) + (f_-)(x_-)]], x_Symbol] \rightarrow \text{Simp}[-2*a*(b/f) \ \text{Subst}[\text{Int}[1/(a*c - b*d - 2*a^2*x^2), x], x, \text{Sqrt}[c + d*\text{Tan}[e + f*x]]/\text{Sqrt}[a + b*\text{Tan}[e + f*x]]], x] \text{ ; FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0]$
- rule 4076 $\text{Int}[(a_+ + (b_-)\tan[(e_-) + (f_-)(x_-)])^{(m_-)}((A_-) + (B_-)\tan[(e_-) + (f_-)(x_-)])^{(n_-)}, x_Symbol] \rightarrow \text{Simp}[(-a^2)*(B*c - A*d)*(a + b*\text{Tan}[e + f*x])^{(m-1)}((c + d*\text{Tan}[e + f*x])^{(n+1)})/(d*f*(b*c + a*d)*(n+1)), x] - \text{Simp}[a/(d*(b*c + a*d)*(n+1)) \ \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m-1)}(c + d*\text{Tan}[e + f*x])^{(n+1)}*\text{Simp}[A*b*d*(m-2) - B*(b*c*(m-1) + a*d*(n+1)) + (a*A*d*(m+n) - B*(a*c*(m-1) + b*d*(n+1)))*\text{Tan}[e + f*x], x], x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{LtQ}[n, -1]$
- rule 4077 $\text{Int}[(a_+ + (b_-)\tan[(e_-) + (f_-)(x_-)])^{(m_-)}((A_-) + (B_-)\tan[(e_-) + (f_-)(x_-)])^{(n_-)}, x_Symbol] \rightarrow \text{Simp}[b*B*(a + b*\text{Tan}[e + f*x])^{(m-1)}((c + d*\text{Tan}[e + f*x])^{(n+1)})/(d*f*(m+n)), x] + \text{Simp}[1/(d*(m+n)) \ \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m-1)}(c + d*\text{Tan}[e + f*x])^n*\text{Simp}[a*A*d*(m+n) + B*(a*c*(m-1) - b*d*(n+1)) - (B*(b*c - a*d)*(m-1) - d*(A*b + a*B))*(m+n))*\text{Tan}[e + f*x], x], x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{!LtQ}[n, -1]$
- rule 4082 $\text{Int}[(a_+ + (b_-)\tan[(e_-) + (f_-)(x_-)])^{(m_-)}((A_-) + (B_-)\tan[(e_-) + (f_-)(x_-)])^{(n_-)}, x_Symbol] \rightarrow \text{Simp}[b*(B/f) \ \text{Subst}[\text{Int}[(a + b*x)^{(m-1)}(c + d*x)^n, x], x, \text{Tan}[e + f*x]], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B, m, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[A*b + a*B, 0]$

rule 4084 `Int[((a_) + (b_.)*tan[(e_) + (f_.)*(x_)])^(m_)*((A_) + (B_.)*tan[(e_) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.172.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 562 vs. $2(161) = 322$.

Time = 0.16 (sec) , antiderivative size = 563, normalized size of antiderivative = 2.87

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} a^2 \left(6iA \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a \tan(dx+c) - 2i\sqrt{ia} \sqrt{2} \ln \left(\dots \right) \right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))} a^2 \left(6iA \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a \tan(dx+c) - 2i\sqrt{ia} \sqrt{2} \ln \left(\dots \right) \right)}{\dots}$
parts	$\frac{A \left(-i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}{\tan(dx+c)+i} \right) a \tan(dx+c) - i \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)}}{2\sqrt{ia}} \right) \right)}{\dots}$

input `int((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x,method=_R ETURNVERBOSE)`

output `1/2/d*(a*(1+I*tan(d*x+c)))^(1/2)*a^2*(6*I*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)-2*I*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2))*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I)*a*tan(d*x+c)+3*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)-2*B*(I*a)^(1/2))*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+4*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)-2*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2))*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I)*tan(d*x+c)*a-4*A*(I*a)^(1/2))*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-4*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c))/tan(d*x+c)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)/(I*a)^(1/2)/(-I*a)^(1/2)`

3.172.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 857 vs. $2(150) = 300$.

Time = 0.28 (sec) , antiderivative size = 857, normalized size of antiderivative = 4.37

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="fricas")
```

```
output -1/2*(4*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log((sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 4*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log(-sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) - sqrt(2)*((-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a^2)) + 2*sqrt(2)*((2*I*A + B)*a^2*e^(3*I*d*x + 3*I*c) + (2*I*A - B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + sqrt((4*I*A^2 + 20*A*B - 25*I*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log((sqrt(2)*((2*I*A + 5*B)*a^2*e^(2*I*d*x + 2*I*c) + (2*I*A + 5*B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + 2*sqrt((4*I*A^2 + 20*A*B - 25*I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c))e^(-I*d*x - I*c)/((2*I*A + 5*B)*a^2)) - sqrt((4*I*A^2 + 20*A*B - 25*I*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log((sqrt(2)*((2*I*A + 5*B)*a^2*e^(2*I*d*x + 2*I*c) + (2*I*A + 5*B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) - 2*sqrt((4*...
```

3.172.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \int \frac{(ia(\tan(c + dx) - i))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx$$

input `integrate((a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(3/2),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(5/2)*(A + B*tan(c + d*x))/tan(c + d*x)**(3/2), x)`

3.172.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="maxima")`

output `Timed out`

3.172.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx);OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[-90]Warning, replacing -90 by -93, a substitution vari`

3.172. $\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx$

3.172.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) 1i)^{5/2}}{\tan(c + dx)^{3/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(3/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(3/2), x)`

3.173
$$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.173.1 Optimal result 1795
 3.173.2 Mathematica [B] (verified) 1796
 3.173.3 Rubi [A] (verified) 1797
 3.173.4 Maple [B] (verified) 1801
 3.173.5 Fricas [B] (verification not implemented) 1802
 3.173.6 Sympy [F] 1803
 3.173.7 Maxima [F] 1804
 3.173.8 Giac [F(-2)] 1804
 3.173.9 Mupad [F(-1)] 1804

3.173.1 Optimal result

Integrand size = 38, antiderivative size = 190

$$\int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = \frac{2(-1)^{3/4}a^{5/2}B \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$+ \frac{(4 + 4i)a^{5/2}(iA + B)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$- \frac{2a^2(2iA + B)\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{3d \tan^{\frac{3}{2}}(c + dx)}$$

output

```
2*(-1)^(3/4)*a^(5/2)*B*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)^(1/2))/d+(4+4*I)*a^(5/2)*(I*A+B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)^(1/2))/d-2*a^2*(2*I*A+B)*(a+I*a*tan(d*x+c)^(1/2))/d/tan(d*x+c)^(1/2)-2/3*a*A*(a+I*a*tan(d*x+c))^(3/2)/d/tan(d*x+c)^(3/2)
```

3.173.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 554 vs. $2(190) = 380$.

Time = 7.32 (sec) , antiderivative size = 554, normalized size of antiderivative = 2.92

$$\int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = -\frac{2A(a + ia \tan(c + dx))^{5/2}}{3d \tan^{3/2}(c + dx)} + \left[-\frac{a(5iA+3B)(a+ia \tan(c+dx))^{5/2}}{d\sqrt{\tan(c+dx)}} + \frac{ia^4(5iA+3B)\sqrt{a+ia \tan(c+dx)}\left(-\frac{3}{4}(-1)^{3/4}\operatorname{arcsinh}\left(\frac{\sqrt[4]{-1}\sqrt{\tan(c+dx)}}{d\sqrt{1+i \tan(c+dx)}}\right)+\frac{5}{4}\sqrt{1+i \tan(c+dx)}\sqrt{\tan(c+dx)}\right)}{d\sqrt{1+i \tan(c+dx)}} \right] + \dots$$

```
input Integrate[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2), x]
```

```
output (-2*A*(a + I*a*Tan[c + d*x])^(5/2))/(3*d*Tan[c + d*x]^(3/2)) + (2*(-((a*((5*I)*A + 3*B)*(a + I*a*Tan[c + d*x])^(5/2))/(d*Sqrt[Tan[c + d*x]])) + (2*((I*a^4*((5*I)*A + 3*B)*Sqrt[a + I*a*Tan[c + d*x]])*(-3*(-1)^(3/4)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]])/4 + (5*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/4 + (I/2)*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(3/2)))/(d*Sqrt[1 + I*Tan[c + d*x]]) + (a*((-1/4*I)*a^3*(23*A - (15*I)*B) + a^3*((5*I)*A + 3*B))*((-4*I)*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[Tan[c + d*x]]/Sqrt[I*a*Tan[c + d*x]] + ((4*I)*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/(Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + I*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]] + (I*Sqrt[a]*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])))/d)/a)/(3*a)
```

3.173.3 Rubi [A] (verified)

Time = 1.16 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4076, 27, 3042, 4076, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx \\
 & \quad \downarrow \text{4076} \\
 & \frac{2}{3} \int \frac{3(i \tan(c + dx)a + a)^{3/2} (a(2iA + B) + iaB \tan(c + dx))}{2 \tan^{3/2}(c + dx)} dx - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{3d \tan^{3/2}(c + dx)} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(2iA + B) + iaB \tan(c + dx))}{\tan^{3/2}(c + dx)} dx - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{3d \tan^{3/2}(c + dx)} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(2iA + B) + iaB \tan(c + dx))}{\tan(c + dx)^{3/2}} dx - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{3d \tan^{3/2}(c + dx)} \\
 & \quad \downarrow \text{4076} \\
 & 2 \int -\frac{\sqrt{i \tan(c + dx)a + a} ((4A - 3iB)a^2 + B \tan(c + dx)a^2)}{2\sqrt{\tan(c + dx)}} dx - \\
 & \quad \frac{2a^2(B + 2iA)\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{3d \tan^{3/2}(c + dx)} \\
 & \quad \downarrow \text{27} \\
 & - \int \frac{\sqrt{i \tan(c + dx)a + a} ((4A - 3iB)a^2 + B \tan(c + dx)a^2)}{\sqrt{\tan(c + dx)}} dx - \\
 & \quad \frac{2a^2(B + 2iA)\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{3d \tan^{3/2}(c + dx)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& - \int \frac{\sqrt{i \tan(c+dx)a+a}((4A-3iB)a^2+B \tan(c+dx)a^2)}{\sqrt{\tan(c+dx)}} dx - \\
& \frac{2a^2(B+2iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow 4084 \\
& -4a^2(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - iaB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \\
& \frac{2a^2(B+2iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow 3042 \\
& -4a^2(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - iaB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \\
& \frac{2a^2(B+2iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow 4027 \\
& \frac{8ia^4(A-iB) \int \frac{1}{\frac{-2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} - \\
& iaB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(B+2iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \\
& \frac{2aA(a+ia \tan(c+dx))^{3/2}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow 218 \\
& -iaB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \\
& \frac{(4-4i)a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^2(B+2iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \\
& \frac{2aA(a+ia \tan(c+dx))^{3/2}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow 4082
\end{aligned}$$

3.173. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
& \frac{ia^3 B \int \frac{1}{\sqrt{\tan(c+dx)} \sqrt{ia \tan(c+dx) + a}} d \tan(c+dx)}{(4-4i)a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) - \frac{2a^2(B+2iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{3d \tan^{3/2}(c+dx)}} \\
& \quad \downarrow 65 \\
& \frac{2ia^3 B \int \frac{1}{1 - \frac{ia \tan(c+dx)}{ia \tan(c+dx) + a}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{ia \tan(c+dx) + a}}}{\frac{2a^2(B+2iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{3d \tan^{3/2}(c+dx)}} + \frac{(4-4i)a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} \\
& \quad \downarrow 216 \\
& \frac{(4-4i)a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{2(-1)^{3/4}a^{5/2}B \operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^2(B+2iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{3d \tan^{3/2}(c+dx)}
\end{aligned}$$

input `Int[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(2*(-1)^(3/4)*a^(5/2)*B*ArcTan[(-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]]/Sqrt[a + I*a*Tan[c + d*x]]]/d - ((4 - 4*I)*a^(5/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]]/Sqrt[a + I*a*Tan[c + d*x]])]/d - (2*a^2*((2*I)*A + B)*Sqrt[a + I*a*Tan[c + d*x]]/(d*Sqrt[Tan[c + d*x]]) - (2*a*A*(a + I*a*Tan[c + d*x])^(3/2))/(3*d*Tan[c + d*x]^(3/2))`

3.173.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.173.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 617 vs. 2(154) = 308.

Time = 0.16 (sec) , antiderivative size = 618, normalized size of antiderivative = 3.25

method	result
derivativedivides	$-\frac{\sqrt{a(1+i \tan(dx+c))} a^2 \left(-9iB \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a(\tan^2(dx+c))+3i\sqrt{ia} \right)}{}$
default	$-\frac{\sqrt{a(1+i \tan(dx+c))} a^2 \left(-9iB \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a(\tan^2(dx+c))+3i\sqrt{ia} \right)}{}$
parts	$-\frac{A\sqrt{a(1+i \tan(dx+c))} a^2 \left(3i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)}(1+i \tan(dx+c))-ia+3a \tan(dx+c)}{\tan(dx+c)+i} \right) a(\tan^2(dx+c))+12 \right)}{}$

input `int((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x,method=_R
ETURNVERBOSE)`

$$3.173. \int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

output

```

-1/3/d*(a*(1+I*tan(d*x+c)))^(1/2)*a^2/tan(d*x+c)^(3/2)*(-9*I*B*ln(1/2*(2*I
*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)
^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^2+3*I*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*
(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)-I*a+3*a*tan(d*x+c))/(ta
n(d*x+c)+I))*a*tan(d*x+c)^2+14*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*
tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+12*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(
d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*
tan(d*x+c)^2+6*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))
)^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^2-3*(I*a)^(1
/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/
2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2+6*B*(I*a)^(1/2)*(-I*
a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+6*ln(1/2*(2*I*a*
tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1
/2))*(-I*a)^(1/2)*a*tan(d*x+c)^2+2*A*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c
)*(1+I*tan(d*x+c)))^(1/2))/(I*a)^(1/2)/(-I*a)^(1/2)/(a*tan(d*x+c)*(1+I*ta
n(d*x+c)))^(1/2)

```

3.173.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 846 vs. $2(144) = 288$.

Time = 0.28 (sec) , antiderivative size = 846, normalized size of antiderivative = 4.45

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \text{Too large to display}$$

input

```

integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, al
gorithm="fricas")

```

output `1/6*(12*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log((I*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 12*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log((-I*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) + 4*sqrt(2)*((8*A - 3*I*B)*a^2*e^(5*I*d*x + 5*I*c) + 2*A*a^2*e^(3*I*d*x + 3*I*c) - 3*(2*A - I*B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + 3*sqrt(4*I*B^2*a^5/d^2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log((sqrt(2)*(B*a^2*e^(2*I*d*x + 2*I*c) + B*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + I*sqrt(4*I*B^2*a^5/d^2)*d*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(B*a^2)) - 3*sqrt(4*I*B^2*a^5/d^2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log((sqrt(2)*(B*a^2*e^(2*I*d*x + 2*I*c) + B*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + ...`

3.173.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \int \frac{(ia(\tan(c + dx) - i))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx$$

input `integrate((a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(5/2),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(5/2)*(A + B*tan(c + d*x))/tan(c + d*x)**(5/2), x)`

3.173.7 Maxima [F]

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{5/2}}{\tan(dx + c)^{5/2}} dx$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)/tan(d*x + c)^(5/2), x)`

3.173.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[-92]Warning, replacing -92 by -88, a substitution vari`

3.173.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) li)^{5/2}}{\tan(c + dx)^{5/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(5/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(5/2), x)`

3.173. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{5/2}(c+dx)} dx$

3.174
$$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{7/2}(c+dx)} dx$$

3.174.1 Optimal result 1805
 3.174.2 Mathematica [B] (verified) 1806
 3.174.3 Rubi [A] (verified) 1806
 3.174.4 Maple [B] (verified) 1810
 3.174.5 Fricas [B] (verification not implemented) 1811
 3.174.6 Sympy [F(-1)] 1812
 3.174.7 Maxima [F(-1)] 1812
 3.174.8 Giac [F(-2)] 1813
 3.174.9 Mupad [F(-1)] 1813

3.174.1 Optimal result

Integrand size = 38, antiderivative size = 185

$$\int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx =$$

$$\frac{(4 + 4i)a^{5/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$- \frac{2a^2(8iA + 5B)\sqrt{a + ia \tan(c + dx)}}{15d \tan^{3/2}(c + dx)}$$

$$+ \frac{2a^2(38A - 35iB)\sqrt{a + ia \tan(c + dx)}}{15d\sqrt{\tan(c + dx)}} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{5d \tan^{5/2}(c + dx)}$$

```
output (-4-4*I)*a^(5/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan
(d*x+c)^(1/2))/d+2/15*a^2*(38*A-35*I*B)*(a+I*a*tan(d*x+c)^(1/2)/d/tan(d*
x+c)^(1/2)-2/15*a^2*(8*I*A+5*B)*(a+I*a*tan(d*x+c)^(1/2)/d/tan(d*x+c)^(3/2
))-2/5*a*A*(a+I*a*tan(d*x+c)^(3/2)/d/tan(d*x+c)^(5/2)
```

3.174.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 447 vs. $2(185) = 370$.

Time = 7.69 (sec) , antiderivative size = 447, normalized size of antiderivative = 2.42

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = -\frac{2A(a + ia \tan(c + dx))^{5/2}}{5d \tan^{5/2}(c + dx)} + (iA + B) \left(\frac{4i\sqrt{2}a^2 \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} - \frac{4ia^{5/2} \operatorname{arcsinh}\left(\frac{\sqrt{ia \tan(c+dx)}}{\sqrt{a}}\right) \sqrt{1 + i \tan(c + dx)}}{d\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}} \right)$$

input `Integrate[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(-2*A*(a + I*a*Tan[c + d*x])^(5/2))/(5*d*Tan[c + d*x]^(5/2)) + (I*A + B)*((4*I)*Sqrt[2]*a^2*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - ((4*I)*a^(5/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + (5*(-1)^(3/4)*a^2*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]]*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]) - (2*a^2*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (((14*I)/3)*a^2*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - (I*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])`

3.174.3 Rubi [A] (verified)

Time = 1.02 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.04, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {3042, 4076, 27, 3042, 4076, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx$$

↓ 3042

3.174. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{7/2}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan(c + dx)^{7/2}} dx \\
& \quad \downarrow \text{4076} \\
& \frac{2}{5} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(8iA + 5B) - a(2A - 5iB) \tan(c + dx))}{2 \tan^{\frac{5}{2}}(c + dx)} dx - \\
& \quad \frac{2aA(a + ia \tan(c + dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{5} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(8iA + 5B) - a(2A - 5iB) \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx - \\
& \quad \frac{2aA(a + ia \tan(c + dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(8iA + 5B) - a(2A - 5iB) \tan(c + dx))}{\tan(c + dx)^{5/2}} dx - \\
& \quad \frac{2aA(a + ia \tan(c + dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{4076} \\
& \frac{1}{5} \left(\frac{2}{3} \int - \frac{\sqrt{i \tan(c + dx)a + a} ((38A - 35iB)a^2 + (22iA + 25B) \tan(c + dx)a^2)}{2 \tan^{\frac{3}{2}}(c + dx)} dx - \frac{2a^2(5B + 8iA) \sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right. \\
& \quad \left. \frac{2aA(a + ia \tan(c + dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c + dx)} \right) \\
& \quad \downarrow \text{27} \\
& \frac{1}{5} \left(-\frac{1}{3} \int \frac{\sqrt{i \tan(c + dx)a + a} ((38A - 35iB)a^2 + (22iA + 25B) \tan(c + dx)a^2)}{\tan^{\frac{3}{2}}(c + dx)} dx - \frac{2a^2(5B + 8iA) \sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right. \\
& \quad \left. \frac{2aA(a + ia \tan(c + dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c + dx)} \right) \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \left(-\frac{1}{3} \int \frac{\sqrt{i \tan(c + dx)a + a} ((38A - 35iB)a^2 + (22iA + 25B) \tan(c + dx)a^2)}{\tan(c + dx)^{3/2}} dx - \frac{2a^2(5B + 8iA) \sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right. \\
& \quad \left. \frac{2aA(a + ia \tan(c + dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c + dx)} \right)
\end{aligned}$$

3.174. $\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$

$$\begin{aligned}
& \downarrow 4081 \\
& \frac{1}{5} \left(\frac{1}{3} \left(\frac{2a^2(38A - 35iB)\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} - \frac{2 \int \frac{30a^3(iA+B)\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{a} \right) - \frac{2a^2(5B + 8iA)\sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right. \\
& \qquad \qquad \qquad \left. \frac{2aA(a + ia \tan(c + dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c + dx)} \right) \\
& \downarrow 27 \\
& \frac{1}{5} \left(\frac{1}{3} \left(\frac{2a^2(38A - 35iB)\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} - 60a^2(B + iA) \int \frac{\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx \right) - \frac{2a^2(5B + 8iA)\sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right. \\
& \qquad \qquad \qquad \left. \frac{2aA(a + ia \tan(c + dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c + dx)} \right) \\
& \downarrow 3042 \\
& \frac{1}{5} \left(\frac{1}{3} \left(\frac{2a^2(38A - 35iB)\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} - 60a^2(B + iA) \int \frac{\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx \right) - \frac{2a^2(5B + 8iA)\sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right. \\
& \qquad \qquad \qquad \left. \frac{2aA(a + ia \tan(c + dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c + dx)} \right) \\
& \downarrow 4027 \\
& \frac{1}{5} \left(\frac{1}{3} \left(\frac{120ia^4(B + iA) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} + \frac{2a^2(38A - 35iB)\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} \right) - \frac{2a^2(5B + 8iA)\sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right. \\
& \qquad \qquad \qquad \left. \frac{2aA(a + ia \tan(c + dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c + dx)} \right) \\
& \downarrow 218 \\
& \frac{1}{5} \left(\frac{1}{3} \left(\frac{2a^2(38A - 35iB)\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} - \frac{(60 - 60i)a^{5/2}(B + iA) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} \right) - \frac{2a^2(5B + 8iA)\sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right. \\
& \qquad \qquad \qquad \left. \frac{2aA(a + ia \tan(c + dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c + dx)} \right)
\end{aligned}$$

input `Int[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(-2*a*A*(a + I*a*Tan[c + d*x])^(3/2))/(5*d*Tan[c + d*x]^(5/2)) + ((-2*a^2*((8*I)*A + 5*B)*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) + (((-60 + 60*I)*a^(5/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d + (2*a^2*(38*A - (35*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/3)/5`

3.174.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`


```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

3.174.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 706 vs. 2(152) = 304.

Time = 0.14 (sec) , antiderivative size = 707, normalized size of antiderivative = 3.82

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} a^2 \left(-76A\sqrt{ia} \sqrt{-ia} (\tan^2(dx+c)) \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} + 60iA \ln \left(\frac{2ia \tan(dx+c) + 2\sqrt{a}}{\dots} \right) \right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))} a^2 \left(-76A\sqrt{ia} \sqrt{-ia} (\tan^2(dx+c)) \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} + 60iA \ln \left(\frac{2ia \tan(dx+c) + 2\sqrt{a}}{\dots} \right) \right)}{\dots}$
parts	$\frac{A\sqrt{a(1+i \tan(dx+c))} a^2 \left(15i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c)) - ia + 3a \tan(dx+c)}}{\tan(dx+c)+i} \right) a (\tan^3(dx+c)) + 15\sqrt{\dots} \right)}{\dots}$

```
input int((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x,method=_R
ETURNVERBOSE)
```

3.174.
$$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{7/2}(c+dx)} dx$$

output

```

-1/15/d*(a*(1+I*tan(d*x+c)))^(1/2)*a^2/tan(d*x+c)^(5/2)*(-76*A*(I*a)^(1/2)
*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+60*I*A*ln
(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)
+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^3-15*I*(I*a)^(1/2)*2^(1/2)*ln((
2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d
*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3+70*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(
d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+60*B*ln(1/2*(2*I*a*tan(d*x+
c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I
*a)^(1/2)*a*tan(d*x+c)^3+30*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+
I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)
^3-15*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*ta
n(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3+22*I*A
*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
-30*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)
^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^3+10*B*(I*a)^(1/2)*(-I*a)
^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+6*A*(I*a)^(1/2)*(-
I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2))/(I*a)^(1/2)/(-I*a)^(1/2)
/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)

```

3.174.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 608 vs. $2(141) = 282$.

Time = 0.27 (sec) , antiderivative size = 608, normalized size of antiderivative = 3.29

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \frac{2 \left(15 \sqrt{2} \sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^5}{d^2}} (de^{(6i dx + 6i c)} - 3 de^{(4i dx)} \right)}{\dots}$$

input

```

integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, al
gorithm="fricas")

```

output $2/15*(15*\sqrt{2}*\sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2}*(d*e^{(6*I*d*x + 6*I*c)} - 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} - d)*\log((\sqrt{2})*\sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2}*d*e^{(I*d*x + I*c)} + \sqrt{2}*((-I*A - B)*a^2*e^{(2*I*d*x + 2*I*c)} + (-I*A - B)*a^2)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*\sqrt{(-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1)})*e^{(-I*d*x - I*c)/((-I*A - B)*a^2)} - 15*\sqrt{2}*\sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2}*(d*e^{(6*I*d*x + 6*I*c)} - 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} - d)*\log(-(\sqrt{2})*\sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2}*d*e^{(I*d*x + I*c)} - \sqrt{2}*((-I*A - B)*a^2*e^{(2*I*d*x + 2*I*c)} + (-I*A - B)*a^2)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*\sqrt{(-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1)})*e^{(-I*d*x - I*c)/((-I*A - B)*a^2)} - 2*\sqrt{2}*(2*(-13*I*A - 10*B)*a^2*e^{(7*I*d*x + 7*I*c)} + 3*(3*I*A + 5*B)*a^2*e^{(5*I*d*x + 5*I*c)} + 20*(I*A + B)*a^2*e^{(3*I*d*x + 3*I*c)} + 15*(-I*A - B)*a^2*e^{(I*d*x + I*c)})*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*\sqrt{(-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1)})/(d*e^{(6*I*d*x + 6*I*c)} - 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} - d)$

3.174.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(7/2),x)`

output Timed out

3.174.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="maxima")`

output Timed out

3.174. $\int \frac{(a+ia \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{7/2}(c+dx)} dx$

3.174.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[-41]Warning, replacing -41 by -23, a substitution vari`

3.174.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) i)^{5/2}}{\tan(c + dx)^{7/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(7/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(7/2), x)`

3.175
$$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

3.175.1 Optimal result 1814
 3.175.2 Mathematica [B] (verified) 1815
 3.175.3 Rubi [A] (verified) 1815
 3.175.4 Maple [B] (verified) 1820
 3.175.5 Fricas [B] (verification not implemented) 1821
 3.175.6 Sympy [F(-1)] 1822
 3.175.7 Maxima [F(-1)] 1822
 3.175.8 Giac [F(-2)] 1822
 3.175.9 Mupad [F(-1)] 1823

3.175.1 Optimal result

Integrand size = 38, antiderivative size = 231

$$\int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx = \frac{(4 - 4i)a^{5/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$- \frac{2a^2(10iA + 7B)\sqrt{a + ia \tan(c + dx)}}{35d \tan^{\frac{5}{2}}(c + dx)} + \frac{2a^2(80A - 77iB)\sqrt{a + ia \tan(c + dx)}}{105d \tan^{\frac{3}{2}}(c + dx)}$$

$$+ \frac{4a^2(130iA + 133B)\sqrt{a + ia \tan(c + dx)}}{105d \sqrt{\tan(c + dx)}} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c + dx)}$$

output

```
(4-4*I)*a^(5/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d+4/105*a^2*(130*I*A+133*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(1/2)-2/35*a^2*(10*I*A+7*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(5/2)+2/105*a^2*(80*A-77*I*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(3/2)-2/7*a*A*(a+I*a*tan(d*x+c))^(3/2)/d/tan(d*x+c)^(7/2)
```

3.175.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 503 vs. $2(231) = 462$.

Time = 8.20 (sec) , antiderivative size = 503, normalized size of antiderivative = 2.18

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = -\frac{2A(a + ia \tan(c + dx))^{5/2}}{7d \tan^{7/2}(c + dx)} + 2 \left(-\frac{a(5iA+7B)(a+ia \tan(c+dx))^{5/2}}{5d \tan^{5/2}(c+dx)} - \frac{7}{2}a(A - iB) \left(\frac{4i\sqrt{2}a^2 \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{4ia^{5/2} \operatorname{arcsinh}\left(\frac{\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d\sqrt{\tan(c+dx)}} \right) \right)$$

input `Integrate[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2),x]`

output `(-2*A*(a + I*a*Tan[c + d*x])^(5/2))/(7*d*Tan[c + d*x]^(7/2)) + (2*(-1/5*(a*((5*I)*A + 7*B)*(a + I*a*Tan[c + d*x])^(5/2))/(d*Tan[c + d*x]^(5/2)) - (7*a*(A - I*B)*(((4*I)*Sqrt[2]*a^2*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])]/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - ((4*I)*a^(5/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + (5*(-1)^(3/4)*a^2*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]) - (2*a^2*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (((14*I)/3)*a^2*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - (I*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]]))/2))/(7*a)`

3.175.3 Rubi [A] (verified)

Time = 1.36 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.07, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4076, 27, 3042, 4076, 27, 3042, 4081, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx$$

3.175. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{9/2}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan(c + dx)^{9/2}} dx \\
& \quad \downarrow \text{3042} \\
& \frac{2}{7} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(10iA + 7B) - a(4A - 7iB) \tan(c + dx))}{2 \tan^{7/2}(c + dx)} dx - \\
& \quad \frac{2aA(a + ia \tan(c + dx))^{3/2}}{7d \tan^{7/2}(c + dx)} \\
& \quad \downarrow \text{4076} \\
& \frac{1}{7} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(10iA + 7B) - a(4A - 7iB) \tan(c + dx))}{\tan^{7/2}(c + dx)} dx - \\
& \quad \frac{2aA(a + ia \tan(c + dx))^{3/2}}{7d \tan^{7/2}(c + dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{7} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(10iA + 7B) - a(4A - 7iB) \tan(c + dx))}{\tan^{7/2}(c + dx)} dx - \\
& \quad \frac{2aA(a + ia \tan(c + dx))^{3/2}}{7d \tan^{7/2}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(10iA + 7B) - a(4A - 7iB) \tan(c + dx))}{\tan(c + dx)^{7/2}} dx - \\
& \quad \frac{2aA(a + ia \tan(c + dx))^{3/2}}{7d \tan^{7/2}(c + dx)} \\
& \quad \downarrow \text{4076} \\
& \frac{1}{7} \left(\frac{2}{5} \int - \frac{\sqrt{i \tan(c + dx)a + a} ((80A - 77iB)a^2 + 3(20iA + 21B) \tan(c + dx)a^2)}{2 \tan^{5/2}(c + dx)} dx - \frac{2a^2(7B + 10iA)\sqrt{a + ia \tan(c + dx)}}{5d \tan^{5/2}(c + dx)} \right. \\
& \quad \left. - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{7d \tan^{7/2}(c + dx)} \right) \\
& \quad \downarrow \text{27} \\
& \frac{1}{7} \left(-\frac{1}{5} \int \frac{\sqrt{i \tan(c + dx)a + a} ((80A - 77iB)a^2 + 3(20iA + 21B) \tan(c + dx)a^2)}{\tan^{5/2}(c + dx)} dx - \frac{2a^2(7B + 10iA)\sqrt{a + ia \tan(c + dx)}}{5d \tan^{5/2}(c + dx)} \right. \\
& \quad \left. - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{7d \tan^{7/2}(c + dx)} \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.175. $\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx$

$$\frac{1}{7} \left(-\frac{1}{5} \int \frac{\sqrt{i \tan(c+dx)a+a}((80A-77iB)a^2+3(20iA+21B)\tan(c+dx)a^2)}{\tan(c+dx)^{5/2}} dx - \frac{2a^2(7B+10iA)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{5/2}(c+dx)} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{7d \tan^{7/2}(c+dx)}$$

↓ 4081

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2a^2(80A-77iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{3/2}(c+dx)} - \frac{2 \int \frac{\sqrt{i \tan(c+dx)a+a}(a^3(130iA+133B)-a^3(80A-77iB)\tan(c+dx))}{\tan^{3/2}(c+dx)} dx}{3a} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{7d \tan^{7/2}(c+dx)} \right) - \frac{2a^3(133B+130iA)\sqrt{a+ia \tan(c+dx)}}{3ad \sqrt{\tan(c+dx)}}$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2a^2(80A-77iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{3/2}(c+dx)} - \frac{2 \int \frac{\sqrt{i \tan(c+dx)a+a}(a^3(130iA+133B)-a^3(80A-77iB)\tan(c+dx))}{\tan(c+dx)^{3/2}} dx}{3a} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{7d \tan^{7/2}(c+dx)} \right) - \frac{2a^3(133B+130iA)\sqrt{a+ia \tan(c+dx)}}{3ad \sqrt{\tan(c+dx)}}$$

↓ 4081

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2a^2(80A-77iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{3/2}(c+dx)} - \frac{2 \left(\frac{2 \int -\frac{105a^4(A-iB)\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{a} - \frac{2a^3(133B+130iA)\sqrt{a+ia \tan(c+dx)}}{d \sqrt{\tan(c+dx)}} \right)}{3a} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{7d \tan^{7/2}(c+dx)} \right) - \frac{2a^3(133B+130iA)\sqrt{a+ia \tan(c+dx)}}{3ad \sqrt{\tan(c+dx)}}$$

↓ 27

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2a^2(80A-77iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{3/2}(c+dx)} - \frac{2 \left(-210a^3(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(133B+130iA)\sqrt{a+ia \tan(c+dx)}}{d \sqrt{\tan(c+dx)}} \right)}{3a} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{7d \tan^{7/2}(c+dx)} \right) - \frac{2a^3(133B+130iA)\sqrt{a+ia \tan(c+dx)}}{3ad \sqrt{\tan(c+dx)}}$$

↓ 3042

3.175. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{9/2}(c+dx)} dx$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2a^2(80A - 77iB)\sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2 \left(-210a^3(A - iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(133B+130iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)}{3a} \right. \right.$$

$$\left. \left. \frac{2aA(a + ia \tan(c + dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c + dx)} \right) \right.$$

↓ 4027

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2a^2(80A - 77iB)\sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2 \left(\frac{420ia^5(A-iB) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d - \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} - \frac{2a^3(133B+130iA)}{d\sqrt{\tan(c+dx)}} \right)}{3a} \right. \right.$$

$$\left. \left. \frac{2aA(a + ia \tan(c + dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c + dx)} \right) \right.$$

↓ 218

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2a^2(80A - 77iB)\sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2 \left(-\frac{(210-210i)a^{7/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^3(133B+130iA)}{d\sqrt{\tan(c+dx)}} \right)}{3a} \right. \right.$$

$$\left. \left. \frac{2aA(a + ia \tan(c + dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c + dx)} \right) \right.$$

input `Int[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2),x]`

output `(-2*a*A*(a + I*a*Tan[c + d*x])^(3/2))/(7*d*Tan[c + d*x]^(7/2)) + ((-2*a^2*((10*I)*A + 7*B)*Sqrt[a + I*a*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) + ((2*a^2*(80*A - (77*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2))) - (2*(((-210 + 210*I)*a^(7/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a^3*((130*I)*A + 133*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]])))/(3*a))/5/7`

3.175.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`
- rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

3.175.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 795 vs. $2(190) = 380$.

Time = 0.14 (sec) , antiderivative size = 796, normalized size of antiderivative = 3.45

method	result
derivativedivides	$\sqrt{a(1+i \tan(dx+c))} a^2 \left(532B\sqrt{ia} \sqrt{-ia} (\tan^3(dx+c)) \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - 420iB \ln \left(\frac{2ia \tan(dx+c) + 2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))}}{\tan(dx+c)+i} \right) \right)$
default	$\sqrt{a(1+i \tan(dx+c))} a^2 \left(532B\sqrt{ia} \sqrt{-ia} (\tan^3(dx+c)) \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - 420iB \ln \left(\frac{2ia \tan(dx+c) + 2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))}}{\tan(dx+c)+i} \right) \right)$
parts	$A\sqrt{a(1+i \tan(dx+c))} a^2 \left(21i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - ia + 3a \tan(dx+c)}{\tan(dx+c)+i} \right) \right) a (\tan^4(dx+c)) + 84i$

input `int((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x,method=_RETURVERBOSE)`

output $\frac{1}{105}d*(a*(1+I*\tan(d*x+c)))^{(1/2)}*a^2/\tan(d*x+c)^{(7/2)}*(532*B*(I*a)^{(1/2)}*(-I*a)^{(1/2)}*\tan(d*x+c)^3*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-420*I*B*\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{(1/2)}*(I*a)^{(1/2)}+a)/(I*a)^{(1/2)}*(-I*a)^{(1/2)}*a*\tan(d*x+c)^4+520*I*A*(I*a)^{(1/2)}*(-I*a)^{(1/2)}*\tan(d*x+c)^3*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+160*A*(I*a)^{(1/2)}*(-I*a)^{(1/2)}*\tan(d*x+c)^2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-154*I*B*(I*a)^{(1/2)}*(-I*a)^{(1/2)}*\tan(d*x+c)^2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+420*A*\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{(1/2)}*(I*a)^{(1/2)}+a)/(I*a)^{(1/2)}*(-I*a)^{(1/2)}*a*\tan(d*x+c)^4+105*I*(I*a)^{(1/2)}*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(tan(d*x+c)+I))*a*\tan(d*x+c)^4-105*(I*a)^{(1/2)}*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(tan(d*x+c)+I))*a*\tan(d*x+c)^4+210*I*\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{(1/2)}*(I*a)^{(1/2)}+a)/(I*a)^{(1/2)}*(-I*a)^{(1/2)}*a*\tan(d*x+c)^4+210*\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{(1/2)}*(I*a)^{(1/2)}+a)/(I*a)^{(1/2)}*(-I*a)^{(1/2)}*a*\tan(d*x+c)^4-90*I*A*(I*a)^{(1/2)}*(-I*a)^{(1/2)}*\tan(d*x+c)*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-42*B*(I*a)^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)-30*A*(I*a)^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)})/(I*a)^{(1/2)}*(-I*a)^{(1/2)}/(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}$

3.175. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^9(c+dx)} dx$

3.175.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 657 vs. $2(177) = 354$.

Time = 0.27 (sec) , antiderivative size = 657, normalized size of antiderivative = 2.84

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx =$$

$$2 \left(105 \sqrt{2} \sqrt{-\frac{(iA^2 + 2AB - iB^2)a^5}{d^2}} (de^{(8i dx + 8i c)} - 4de^{(6i dx + 6i c)} + 6de^{(4i dx + 4i c)} - 4de^{(2i dx + 2i c)} + d) \log \left(\frac{(i\sqrt{2}\sqrt{-\frac{(iA^2 + 2AB - iB^2)a^5}{d^2}} + 1) e^{(I dx + I c)}}{e^{(2I dx + 2I c)} + 1} \right) + \right.$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="fricas")`

output `-2/105*(105*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)*log((I*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 105*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)*log((-I*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/((-I*A - B)*a^2)) + 2*sqrt(2)*(2*(100*A - 91*I*B)*a^2*e^(9*I*d*x + 9*I*c) - 5*(37*A - 49*I*B)*a^2*e^(7*I*d*x + 7*I*c) - 7*(5*A - 11*I*B)*a^2*e^(5*I*d*x + 5*I*c) + 245*(A - I*B)*a^2*e^(3*I*d*x + 3*I*c) - 105*(A - I*B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)`

3.175.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(9/2),x)`

output Timed out

3.175.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="maxima")`

output Timed out

3.175.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[-28]Warning, replacing -28 by 24, a substitution varia

3.175.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) 1i)^{5/2}}{\tan(c + dx)^{9/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(9/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(9/2), x)`

3.176
$$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$$

3.176.1 Optimal result 1824
 3.176.2 Mathematica [B] (verified) 1825
 3.176.3 Rubi [A] (verified) 1826
 3.176.4 Maple [B] (verified) 1831
 3.176.5 Fricas [B] (verification not implemented) 1832
 3.176.6 Sympy [F(-1)] 1833
 3.176.7 Maxima [F(-1)] 1834
 3.176.8 Giac [F(-2)] 1834
 3.176.9 Mupad [F(-1)] 1834

3.176.1 Optimal result

Integrand size = 38, antiderivative size = 277

$$\int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx = \frac{(4 + 4i)a^{5/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$- \frac{2a^2(4iA + 3B)\sqrt{a + ia \tan(c + dx)}}{21d \tan^{\frac{7}{2}}(c + dx)} + \frac{2a^2(46A - 45iB)\sqrt{a + ia \tan(c + dx)}}{105d \tan^{\frac{5}{2}}(c + dx)}$$

$$+ \frac{8a^2(59iA + 60B)\sqrt{a + ia \tan(c + dx)}}{315d \tan^{\frac{3}{2}}(c + dx)}$$

$$- \frac{8a^2(197A - 195iB)\sqrt{a + ia \tan(c + dx)}}{315d \sqrt{\tan(c + dx)}} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

output

```
(4+4*I)*a^(5/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d-8/315*a^2*(197*A-195*I*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(1/2)-2/21*a^2*(4*I*A+3*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(7/2)+2/105*a^2*(46*A-45*I*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(5/2)+8/315*a^2*(59*I*A+60*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(3/2)-2/9*a*A*(a+I*a*tan(d*x+c))^(3/2)/d/tan(d*x+c)^(9/2)
```

3.176.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 559 vs. $2(277) = 554$.

Time = 8.82 (sec) , antiderivative size = 559, normalized size of antiderivative = 2.02

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = -\frac{2A(a + ia \tan(c + dx))^{5/2}}{9d \tan^{9/2}(c + dx)} + 2 \left(-\frac{a(5iA + 9B)(a + ia \tan(c + dx))^{5/2}}{7d \tan^{7/2}(c + dx)} + \frac{a^2(53A - 45iB)(a + ia \tan(c + dx))^{5/2} - \frac{63}{4}a^2(iA + B)}{10d \tan^{5/2}(c + dx)} \right) \frac{4i\sqrt{2}a^2 \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right) \sqrt{ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}}$$

input `Integrate[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(11/2), x]`

output `(-2*A*(a + I*a*Tan[c + d*x])^(5/2))/(9*d*Tan[c + d*x]^(9/2)) + (2*(-1/7*(a*((5*I)*A + 9*B)*(a + I*a*Tan[c + d*x])^(5/2))/(d*Tan[c + d*x]^(7/2)) + (2*((a^2*(53*A - (45*I)*B)*(a + I*a*Tan[c + d*x])^(5/2))/(10*d*Tan[c + d*x]^(5/2)) - (63*a^2*(I*A + B))*((4*I)*Sqrt[2]*a^2*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - ((4*I)*a^(5/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + (5*(-1)^(3/4)*a^2*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]])*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]) - (2*a^2*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (((14*I)/3)*a^2*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - (I*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])))/4)/(7*a))/(9*a)`

3.176.3 Rubi [A] (verified)

Time = 1.68 (sec) , antiderivative size = 301, normalized size of antiderivative = 1.09, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.474$, Rules used = {3042, 4076, 27, 3042, 4076, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan(c + dx)^{11/2}} dx$$

$$\downarrow \text{4076}$$

$$\frac{2}{9} \int \frac{3(i \tan(c + dx)a + a)^{3/2} (a(4iA + 3B) - a(2A - 3iB) \tan(c + dx))}{2 \tan^{\frac{9}{2}}(c + dx)} dx - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

$$\downarrow \text{27}$$

$$\frac{1}{3} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(4iA + 3B) - a(2A - 3iB) \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

$$\downarrow \text{3042}$$

$$\frac{1}{3} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(4iA + 3B) - a(2A - 3iB) \tan(c + dx))}{\tan(c + dx)^{9/2}} dx - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

$$\downarrow \text{4076}$$

$$\frac{1}{3} \left(\frac{2}{7} \int - \frac{\sqrt{i \tan(c + dx)a + a} ((46A - 45iB)a^2 + (38iA + 39B) \tan(c + dx)a^2)}{2 \tan^{\frac{7}{2}}(c + dx)} dx - \frac{2a^2(3B + 4iA) \sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} \right)$$

$$\downarrow \text{27}$$

3.176. $\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx$

$$\frac{1}{3} \left(-\frac{1}{7} \int \frac{\sqrt{i \tan(c+dx)a+a}((46A-45iB)a^2+(38iA+39B)\tan(c+dx)a^2)}{\tan^{\frac{7}{2}}(c+dx)} dx - \frac{2a^2(3B+4iA)\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{3} \left(-\frac{1}{7} \int \frac{\sqrt{i \tan(c+dx)a+a}((46A-45iB)a^2+(38iA+39B)\tan(c+dx)a^2)}{\tan(c+dx)^{7/2}} dx - \frac{2a^2(3B+4iA)\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

↓ 4081

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A-45iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2 \int \frac{2\sqrt{i \tan(c+dx)a+a}(a^3(59iA+60B)-a^3(46A-45iB)\tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx}{5a} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c+dx)} \right)$$

↓ 27

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A-45iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{4 \int \frac{\sqrt{i \tan(c+dx)a+a}(a^3(59iA+60B)-a^3(46A-45iB)\tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx}{5a} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A-45iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{4 \int \frac{\sqrt{i \tan(c+dx)a+a}(a^3(59iA+60B)-a^3(46A-45iB)\tan(c+dx))}{\tan(c+dx)^{5/2}} dx}{5a} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c+dx)} \right)$$

↓ 4081

3.176. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A - 45iB)\sqrt{a + ia \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{4 \left(\frac{2 \int -\frac{\sqrt{i \tan(c+dx)a+a} \left((197A-195iB)a^4+2(59iA+60B) \tan(c+dx)a^4}{2 \tan^{\frac{3}{2}}(c+dx)} dx}{3a} - \frac{2a^3(60A-59iB)\sqrt{a+ia \tan(c+dx)}}{5a} \right)}{5a} \right) \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A - 45iB)\sqrt{a + ia \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{4 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} \left((197A-195iB)a^4+2(59iA+60B) \tan(c+dx)a^4}{\tan^{\frac{3}{2}}(c+dx)} dx}{3a} - \frac{2a^3(60A-59iB)\sqrt{a+ia \tan(c+dx)}}{5a} \right)}{5a} \right) \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A - 45iB)\sqrt{a + ia \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{4 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} \left((197A-195iB)a^4+2(59iA+60B) \tan(c+dx)a^4}{\tan(c+dx)^{3/2}} dx}{3a} - \frac{2a^3(60A-59iB)\sqrt{a+ia \tan(c+dx)}}{5a} \right)}{5a} \right) \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 4081

3.176. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A - 45iB)\sqrt{a + ia \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{4 \left(-\frac{2 \int \frac{315a^5(iA+B)\sqrt{i \tan(c+dx)a+a} dx}{2\sqrt{\tan(c+dx)}}}{a} - \frac{2a^4(197A-195iB)\sqrt{a+ia \tan(c+dx)}}{3a d\sqrt{\tan(c+dx)}} \right)}{5a} - 2a \right) \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A - 45iB)\sqrt{a + ia \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{4 \left(-\frac{315a^4(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^4(197A-195iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} \right)}{5a} - 2a \right) \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A - 45iB)\sqrt{a + ia \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{4 \left(-\frac{315a^4(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^4(197A-195iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} \right)}{5a} - 2a \right) \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 4027

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A - 45iB)\sqrt{a + ia \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{4 \left(-\frac{630ia^6(B+iA) \int \frac{1}{\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} - \frac{2a^4(197A-195iB)\sqrt{a+ia \tan(c+dx)}}{3a d\sqrt{\tan(c+dx)}} \right)}{5a} - 2a \right) \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

3.176. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$

$$\begin{aligned} & \downarrow 218 \\ & \frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A - 45iB)\sqrt{a + ia \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{4 \left(-\frac{2a^3(60B + 59iA)\sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{(315 - 315i)a^{9/2}(B + iA)\operatorname{arctanh}\left(\frac{1 + \sqrt{a + ia \tan(c + dx)}}{1 - \sqrt{a + ia \tan(c + dx)}}\right)}{d} \right)}{5a} \right. \right. \\ & \left. \left. + \frac{2aA(a + ia \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)} \right) \right) \end{aligned}$$

input `Int[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(11/2),x]`

output `(-2*a*A*(a + I*a*Tan[c + d*x])^(3/2))/(9*d*Tan[c + d*x]^(9/2)) + ((-2*a^2*((4*I)*A + 3*B)*Sqrt[a + I*a*Tan[c + d*x]])/(7*d*Tan[c + d*x]^(7/2)) + ((2*a^2*(46*A - (45*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) - (4*((-2*a^3*((59*I)*A + 60*B)*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (((315 - 315*I)*a^(9/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a^4*(197*A - (195*I)*B)*Sqrt[a + I*a*Tan[c + d*x]]/(d*Sqrt[Tan[c + d*x]]))/(3*a)))/(5*a))/7)/3`

3.176.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4027 Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)]]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4076 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

```
rule 4081 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

3.176.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 884 vs. 2(228) = 456.

Time = 0.16 (sec) , antiderivative size = 885, normalized size of antiderivative = 3.19

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} a^2 \left(-1576A\sqrt{ia} \sqrt{-ia} (\tan^4(dx+c)) \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - 190iA\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)} \right)}{\tan^{\frac{11}{2}}(c+dx)}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))} a^2 \left(-1576A\sqrt{ia} \sqrt{-ia} (\tan^4(dx+c)) \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - 190iA\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)} \right)}{\tan^{\frac{11}{2}}(c+dx)}$
parts	Expression too large to display

3.176.
$$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$$

```
input int((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x,method=_
RETURNVERBOSE)
```

```
output 1/315/d*(a*(1+I*tan(d*x+c)))^(1/2)*a^2/tan(d*x+c)^(9/2)*(-1576*A*(I*a)^(1/
2)*(-I*a)^(1/2)*tan(d*x+c)^4*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-190*I*A
*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
+1560*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^4*(a*tan(d*x+c)*(1+I*tan(d*x
+c)))^(1/2)+480*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3*(a*tan(d*x+c)*(1+I
*tan(d*x+c)))^(1/2)+630*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*ta
n(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^5+1
260*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*
a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^5-270*I*B*(I*a)^(1/2)*(-
I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-315*(I*a)^(
1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1
/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^5+276*A*(I*a)^(1/2)*(-
I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-315*I*(I*a)
^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(
1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^5-630*ln(1/2*(2*I*a
*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(
1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^5+1260*I*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*t
an(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)
*a*tan(d*x+c)^5+472*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3*(a*tan(d*x+c
)*(1+I*tan(d*x+c)))^(1/2)-90*B*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c))*...
```

3.176.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 722 vs. $2(213) = 426$.

Time = 0.27 (sec) , antiderivative size = 722, normalized size of antiderivative = 2.61

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx =$$

$$2 \left(315 \sqrt{2} \sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^5}{d^2}} (de^{(10i dx + 10i c)} - 5 de^{(8i dx + 8i c)} + 10 de^{(6i dx + 6i c)} - 10 de^{(4i dx + 4i c)} + 5 de^{(2i dx + 2i c)}) \right)$$

```
input integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x, a
lgorithm="fricas")
```

3.176. $\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx$

output

```

-2/315*(315*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(d*e^(10*I*d*x
+ 10*I*c) - 5*d*e^(8*I*d*x + 8*I*c) + 10*d*e^(6*I*d*x + 6*I*c) - 10*d*e^(
4*I*d*x + 4*I*c) + 5*d*e^(2*I*d*x + 2*I*c) - d)*log((sqrt(2)*sqrt(-(-I*A^2
- 2*A*B + I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a^2*e^(
2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(
(-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/
((-I*A - B)*a^2)) - 315*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(d
*e^(10*I*d*x + 10*I*c) - 5*d*e^(8*I*d*x + 8*I*c) + 10*d*e^(6*I*d*x + 6*I*c
) - 10*d*e^(4*I*d*x + 4*I*c) + 5*d*e^(2*I*d*x + 2*I*c) - d)*log(-(sqrt(2)*
sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) - sqrt(2)*((-I*A
- B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c
) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^(-
I*d*x - I*c)/((-I*A - B)*a^2)) + 2*sqrt(2)*(2*(323*I*A + 300*B)*a^2*e^(11*
I*d*x + 11*I*c) + 77*(-13*I*A - 15*B)*a^2*e^(9*I*d*x + 9*I*c) + 18*(38*I*A
+ 25*B)*a^2*e^(7*I*d*x + 7*I*c) + 42*(23*I*A + 20*B)*a^2*e^(5*I*d*x + 5*I
*c) + 1050*(-I*A - B)*a^2*e^(3*I*d*x + 3*I*c) + 315*(I*A + B)*a^2*e^(I*d*x
+ I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) +
I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(10*I*d*x + 10*I*c) - 5*d*e^(8*I*d*x +
8*I*c) + 10*d*e^(6*I*d*x + 6*I*c) - 10*d*e^(4*I*d*x + 4*I*c) + 5*d*e^(2*I
*d*x + 2*I*c) - d)

```

3.176.6 Sympy [**F(-1)**]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(11/2),x)`

output `Timed out`

3.176.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x, algorithm="maxima")`

output `Timed out`

3.176.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[35]Warning, replacing 35 by 59, a substitution variable`

3.176.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) i)^{5/2}}{\tan(c + dx)^{11/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(11/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(11/2), x)`

3.176. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{11/2}(c+dx)} dx$

$$3.177 \quad \int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$$

3.177.1 Optimal result 1835
 3.177.2 Mathematica [A] (verified) 1836
 3.177.3 Rubi [A] (verified) 1837
 3.177.4 Maple [B] (verified) 1845
 3.177.5 Fricas [B] (verification not implemented) 1846
 3.177.6 Sympy [F(-1)] 1847
 3.177.7 Maxima [F(-1)] 1848
 3.177.8 Giac [F(-2)] 1848
 3.177.9 Mupad [F(-1)] 1848

3.177.1 Optimal result

Integrand size = 38, antiderivative size = 323

$$\int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{13}{2}}(c + dx)} dx = \frac{(4 + 4i)a^{5/2}(iA + B)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$- \frac{2a^2(14iA + 11B)\sqrt{a + ia \tan(c + dx)}}{99d \tan^{\frac{9}{2}}(c + dx)} + \frac{2a^2(212A - 209iB)\sqrt{a + ia \tan(c + dx)}}{693d \tan^{\frac{7}{2}}(c + dx)}$$

$$+ \frac{4a^2(250iA + 253B)\sqrt{a + ia \tan(c + dx)}}{1155d \tan^{\frac{5}{2}}(c + dx)} - \frac{8a^2(655A - 649iB)\sqrt{a + ia \tan(c + dx)}}{3465d \tan^{\frac{3}{2}}(c + dx)}$$

$$- \frac{8a^2(2155iA + 2167B)\sqrt{a + ia \tan(c + dx)}}{3465d \sqrt{\tan(c + dx)}} - \frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

output

```
(4+4*I)*a^(5/2)*(I*A+B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d-8/3465*a^2*(2155*I*A+2167*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(1/2)-2/99*a^2*(14*I*A+11*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(9/2)+2/693*a^2*(212*A-209*I*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(7/2)+4/1155*a^2*(250*I*A+253*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(5/2)-8/3465*a^2*(655*A-649*I*B)*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(3/2)-2/11*a*A*(a+I*a*tan(d*x+c))^(3/2)/d/tan(d*x+c)^(11/2)
```

3.177.2 Mathematica [A] (verified)

Time = 9.71 (sec) , antiderivative size = 613, normalized size of antiderivative = 1.90

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{13/2}(c + dx)} dx = -\frac{2A(a + ia \tan(c + dx))^{5/2}}{11d \tan^{11/2}(c + dx)} +$$

$$2 \left(-\frac{a(5iA + 11B)(a + ia \tan(c + dx))^{5/2}}{9d \tan^{9/2}(c + dx)} + \frac{a^2(79A - 55iB)(a + ia \tan(c + dx))^{5/2}}{14d \tan^{7/2}(c + dx)} + \frac{a^3(535iA + 583B)(a + ia \tan(c + dx))^{5/2}}{20d \tan^{5/2}(c + dx)} + \frac{693a^3(A - iB)}{8} \frac{4i\sqrt{2}a^2}{\dots} \right)$$

input `Integrate[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(13/2), x]`

output `(-2*A*(a + I*a*Tan[c + d*x])^(5/2))/(11*d*Tan[c + d*x]^(11/2)) + (2*(-1/9*(a*((5*I)*A + 11*B)*(a + I*a*Tan[c + d*x])^(5/2))/(d*Tan[c + d*x]^(9/2)) + (2*((a^2*(79*A - (55*I)*B)*(a + I*a*Tan[c + d*x])^(5/2))/(14*d*Tan[c + d*x]^(7/2)) + (2*((a^3*((535*I)*A + 583*B)*(a + I*a*Tan[c + d*x])^(5/2))/(20*d*Tan[c + d*x]^(5/2)) + (693*a^3*(A - I*B)*((4*I)*Sqrt[2]*a^2*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - ((4*I)*a^(5/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x])) + (5*(-1)^(3/4)*a^2*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]])*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]) - (2*a^2*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - ((14*I)/3)*a^2*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - (I*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])))/(9*a))/(11*a)`

3.177.3 Rubi [A] (verified)

Time = 2.08 (sec) , antiderivative size = 355, normalized size of antiderivative = 1.10, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.553$, Rules used = {3042, 4076, 27, 3042, 4076, 27, 3042, 4081, 27, 3042, 4081, 25, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{13}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan(c + dx)^{13/2}} dx \\
 & \quad \downarrow \text{4076} \\
 & \frac{2}{11} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(14iA + 11B) - a(8A - 11iB) \tan(c + dx))}{2 \tan^{\frac{11}{2}}(c + dx)} dx - \\
 & \quad \frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{11} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(14iA + 11B) - a(8A - 11iB) \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx - \\
 & \quad \frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{11} \int \frac{(i \tan(c + dx)a + a)^{3/2} (a(14iA + 11B) - a(8A - 11iB) \tan(c + dx))}{\tan(c + dx)^{11/2}} dx - \\
 & \quad \frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)} \\
 & \quad \downarrow \text{4076} \\
 & \frac{1}{11} \left(\frac{2}{9} \int -\frac{\sqrt{i \tan(c + dx)a + a} ((212A - 209iB)a^2 + (184iA + 187B) \tan(c + dx)a^2)}{2 \tan^{\frac{9}{2}}(c + dx)} dx - \frac{2a^2(11B + 14iA)\sqrt{a}}{9d \tan^{\frac{9}{2}}(c + dx)} \right) \\
 & \quad \frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.177. $\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{13}{2}}(c + dx)} dx$

$$\frac{1}{11} \left(-\frac{1}{9} \int \frac{\sqrt{i \tan(c+dx)a+a}((212A-209iB)a^2+(184iA+187B)\tan(c+dx)a^2)}{\tan^{\frac{9}{2}}(c+dx)} dx - \frac{2a^2(11B+14iA)\sqrt{a}}{9d \tan^{\frac{9}{2}}(c+dx)} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{11} \left(-\frac{1}{9} \int \frac{\sqrt{i \tan(c+dx)a+a}((212A-209iB)a^2+(184iA+187B)\tan(c+dx)a^2)}{\tan(c+dx)^{9/2}} dx - \frac{2a^2(11B+14iA)\sqrt{a}}{9d \tan^{\frac{9}{2}}(c+dx)} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)}$$

↓ 4081

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2a^2(212A-209iB)\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} - \frac{2 \int \frac{3\sqrt{i \tan(c+dx)a+a}(a^3(250iA+253B)-a^3(212A-209iB)\tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx}{7a} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)} \right)$$

↓ 27

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2a^2(212A-209iB)\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} - \frac{6 \int \frac{\sqrt{i \tan(c+dx)a+a}(a^3(250iA+253B)-a^3(212A-209iB)\tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx}{7a} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2a^2(212A-209iB)\sqrt{a+ia \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} - \frac{6 \int \frac{\sqrt{i \tan(c+dx)a+a}(a^3(250iA+253B)-a^3(212A-209iB)\tan(c+dx))}{\tan(c+dx)^{7/2}} dx}{7a} \right) - \frac{2aA(a+ia \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)} \right)$$

↓ 4081

3.177. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2a^2(212A - 209iB)\sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{6 \left(2 \int \frac{\sqrt{i \tan(c+dx)a+a} \left((655A - 649iB)a^4 + 2(250iA+253B) \tan(c+dx)a^4 \right) dx}{\tan^{\frac{5}{2}}(c+dx)} \right)}{7a} \right) \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 25

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2a^2(212A - 209iB)\sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{6 \left(2 \int \frac{\sqrt{i \tan(c+dx)a+a} \left((655A - 649iB)a^4 + 2(250iA+253B) \tan(c+dx)a^4 \right) dx}{\tan^{\frac{5}{2}}(c+dx)} \right)}{7a} \right) \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2a^2(212A - 209iB)\sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{6 \left(2 \int \frac{\sqrt{i \tan(c+dx)a+a} \left((655A - 649iB)a^4 + 2(250iA+253B) \tan(c+dx)a^4 \right) dx}{\tan^{\frac{5}{2}}(c+dx)} \right)}{7a} \right) \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 4081

3.177. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$

$$\frac{1}{11} \left(\frac{1}{9} \frac{2a^2(212A - 209iB)\sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{2 \int \frac{\sqrt{i \tan(c+dx)a+a} (a^5(2155iA+2167B)-2a^5(655A-649iB) \tan(c+dx))}{2 \tan^{\frac{3}{2}}(c+dx)} dx}{6 \cdot 5a} \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{11} \left(\frac{1}{9} \frac{2a^2(212A - 209iB)\sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{2 \int \frac{\sqrt{i \tan(c+dx)a+a} (a^5(2155iA+2167B)-2a^5(655A-649iB) \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx}{6 \cdot 5a} \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \frac{2a^2(212A - 209iB)\sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{2 \left(\int \frac{\sqrt{i \tan(c+dx)a+a} (a^5(2155iA+2167B) - 2a^5(655A-649iB) \tan(c+dx)) dx}{\tan(c+dx)^{3/2}} \right)}{6 \cdot 5a} \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 4081

$$\frac{1}{11} \left(\frac{1}{9} \frac{2a^2(212A - 209iB)\sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{2 \left(\int -\frac{3465a^6(A-iB)\sqrt{i \tan(c+dx)a+a} dx}{2\sqrt{\tan(c+dx)}a} - \frac{2a^5(2167B+2155iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)}{6 \cdot 5a} \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 27

3.177. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$

$$\frac{1}{11} \left(\frac{1}{9} \frac{2a^2(212A - 209iB)\sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{2 \left(\frac{-3465a^5(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^5(2167B+2155iA)\sqrt{a+ia \tan(c+dx)}}{3a} \right)}{6 \cdot 5a} \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \frac{2a^2(212A - 209iB)\sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{2 \left(\frac{-3465a^5(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^5(2167B+2155iA)\sqrt{a+ia \tan(c+dx)}}{3a} \right)}{6 \cdot 5a} \right)$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 4027

$$\left(\frac{1}{11} \right) \left(\frac{1}{9} \right) \frac{2a^2(212A - 209iB)\sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \left(\frac{6930ia^7(A-iB) \int \frac{1}{\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} - \frac{2a^5(2167B+)}{3a} \right) \frac{1}{5a}$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 218

$$\left(\frac{1}{11} \right) \left(\frac{1}{9} \right) \frac{2a^2(212A - 209iB)\sqrt{a + ia \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \left(\frac{2a^3(253B+250iA)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{(3465-3465i)a^{11/2}(A-iB)}{2} \right) \frac{1}{5a}$$

$$\frac{2aA(a + ia \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

```
input Int[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(13/2),x]
```

3.177. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$

```
output (-2*a*A*(a + I*a*Tan[c + d*x])^(3/2))/(11*d*Tan[c + d*x]^(11/2)) + ((-2*a^
2*((14*I)*A + 11*B)*Sqrt[a + I*a*Tan[c + d*x]])/(9*d*Tan[c + d*x]^(9/2)) +
((2*a^2*(212*A - (209*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(7*d*Tan[c + d*x]
^(7/2)) - (6*((-2*a^3*((250*I)*A + 253*B)*Sqrt[a + I*a*Tan[c + d*x]])/(5*d
*Tan[c + d*x]^(5/2)) - (2*((-2*a^4*(655*A - (649*I)*B)*Sqrt[a + I*a*Tan[c
+ d*x]])/(3*d*Tan[c + d*x]^(3/2)) + (((-3465 + 3465*I)*a^(11/2)*(A - I*B)*
ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/
d - (2*a^5*((2155*I)*A + 2167*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c
+ d*x]]))/(3*a)))/(5*a)))/(7*a))/9)/11
```

3.177.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4027 Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

3.177.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 973 vs. $2(266) = 532$.

Time = 0.15 (sec) , antiderivative size = 974, normalized size of antiderivative = 3.02

method	result	size
derivativedivides	Expression too large to display	974
default	Expression too large to display	974
parts	Expression too large to display	1045

input `int((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(13/2),x,method=_RETURNVERBOSE)`

output

```

-1/3465/d*(a*(1+I*tan(d*x+c)))^(1/2)*a^2/tan(d*x+c)^(11/2)*(17336*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^5*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-13860*I*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^6-5192*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^4*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+5240*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^4*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+2090*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+13860*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^6+1610*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-3465*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^6-3036*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+17240*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^5*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+6930*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^6-2120*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+6930*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^6+3465*I*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)...

```

3.177.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 771 vs. $2(249) = 498$.

Time = 0.27 (sec) , antiderivative size = 771, normalized size of antiderivative = 2.39

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{13/2}(c + dx)} dx = \text{Too large to display}$$

input

```

integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(13/2),x, a
lgorithm="fricas")

```

output `2/3465*(3465*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(12*I*d*x + 12*I*c) - 6*d*e^(10*I*d*x + 10*I*c) + 15*d*e^(8*I*d*x + 8*I*c) - 20*d*e^(6*I*d*x + 6*I*c) + 15*d*e^(4*I*d*x + 4*I*c) - 6*d*e^(2*I*d*x + 2*I*c) + d)*log((I*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 3465*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(12*I*d*x + 12*I*c) - 6*d*e^(10*I*d*x + 10*I*c) + 15*d*e^(8*I*d*x + 8*I*c) - 20*d*e^(6*I*d*x + 6*I*c) + 15*d*e^(4*I*d*x + 4*I*c) - 6*d*e^(2*I*d*x + 2*I*c) + d)*log((-I*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*d*e^(I*d*x + I*c) + sqrt(2)*((-I*A - B)*a^2*e^(2*I*d*x + 2*I*c) + (-I*A - B)*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) + 2*sqrt(2)*(2*(3730*A - 3553*I*B)*a^2*e^(13*I*d*x + 13*I*c) - 9*(1805*A - 2013*I*B)*a^2*e^(11*I*d*x + 11*I*c) + 55*(397*A - 337*I*B)*a^2*e^(9*I*d*x + 9*I*c) + 66*(95*A - 47*I*B)*a^2*e^(7*I*d*x + 7*I*c) - 1386*(15*A - 16*I*B)*a^2*e^(5*I*d*x + 5*I*c) + 15015*(A - I*B)*a^2*e^(3*I*d*x + 3*I*c) - 3465*(A - I*B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(d*e^(12*I*d*x + 12*I*c) - 6*d*e^(10*I*d*x + 10*I*c) + 15*d*e^(8*I*d*x + ...`

3.177.6 Sympy [**F(-1)**]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{13/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(13/2),x)`

output `Timed out`

3.177.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{13/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(13/2),x, algorithm="maxima")`

output `Timed out`

3.177.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{13/2}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(13/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[-42]Warning, replacing -42 by -80, a substitution vari`

3.177.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{13/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) i)^{5/2}}{\tan(c + dx)^{13/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(13/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(13/2), x)`

3.177. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{13/2}(c+dx)} dx$

3.178
$$\int \frac{(a+ia \tan(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c+dx)\right)}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.178.1 Optimal result 1849
 3.178.2 Mathematica [B] (verified) 1850
 3.178.3 Rubi [A] (verified) 1851
 3.178.4 Maple [B] (verified) 1855
 3.178.5 Fricas [B] (verification not implemented) 1856
 3.178.6 Sympy [F] 1857
 3.178.7 Maxima [F] 1858
 3.178.8 Giac [F(-2)] 1858
 3.178.9 Mupad [F(-1)] 1858

3.178.1 Optimal result

Integrand size = 46, antiderivative size = 190

$$\int \frac{(a + ia \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx)\right)}{\tan^{\frac{5}{2}}(c + dx)} dx = \frac{2(-1)^{3/4} a^{5/2} B \arctan\left(\frac{(-1)^{3/4} \sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$+ \frac{(2 + 2i)a^{3/2}(2a + 3ib) \operatorname{Barctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}$$

$$- \frac{2a(a + 3ib)B \sqrt{a + ia \tan(c + dx)}}{d \sqrt{\tan(c + dx)}} - \frac{bB(a + ia \tan(c + dx))^{3/2}}{d \tan^{\frac{3}{2}}(c + dx)}$$

output

```
2*(-1)^(3/4)*a^(5/2)*B*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)^(1/2))/d+(2+2*I)*a^(3/2)*(2*a+3*I*b)*B*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)^(1/2))/d-2*a*(a+3*I*b)*B*(a+I*a*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(1/2)-b*B*(a+I*a*tan(d*x+c))^(3/2)/d/tan(d*x+c)^(3/2)
```


3.178.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 566 vs. $2(190) = 380$.

Time = 11.38 (sec) , antiderivative size = 566, normalized size of antiderivative = 2.98

$$\int \frac{(a + ia \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx)\right)}{\tan^{5/2}(c + dx)} dx = -\frac{bB(a + ia \tan(c + dx))^{5/2}}{ad \tan^{3/2}(c + dx)}$$

$$+ \left[-\frac{3(2a+5ib)B(a+ia \tan(c+dx))^{5/2}}{2d\sqrt{\tan(c+dx)}} + \frac{2 \left(\frac{3ia^3(2a+5ib)B\sqrt{a+ia \tan(c+dx)} \left(-\frac{3}{4}(-1)^{3/4} \operatorname{arcsinh}\left(\sqrt[4]{-1}\sqrt{\tan(c+dx)}\right) + \frac{5}{4}\sqrt{1+i \tan(c+dx)}\sqrt{\tan(c+dx)}\right)}{2d\sqrt{1+i \tan(c+dx)}} \right)}{2} \right]$$

input `Integrate[((a + I*a*Tan[c + d*x])^(5/2)*((3*b*B)/(2*a) + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `-(b*B*(a + I*a*Tan[c + d*x])^(5/2))/(a*d*Tan[c + d*x]^(3/2)) + (2*((-3*(2*a + (5*I)*b)*B*(a + I*a*Tan[c + d*x])^(5/2))/(2*d*Sqrt[Tan[c + d*x]]) + (2*(((3*I)/2)*a^3*(2*a + (5*I)*b)*B*Sqrt[a + I*a*Tan[c + d*x]]*((-3*(-1)^(3/4)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]])]/4 + (5*Sqrt[1 + I*Tan[c + d*x]])*Sqrt[Tan[c + d*x]])/4 + (I/2)*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(3/2)))/(d*Sqrt[1 + I*Tan[c + d*x]]) + (a*(((3*I)/8)*a^2*((10*I)*a - 23*b)*B + (3*a^2*(2*a + (5*I)*b)*B)/2)*((-4*I)*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[Tan[c + d*x]])/Sqrt[I*a*Tan[c + d*x]] + ((4*I)*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]])*Sqrt[Tan[c + d*x]])/(Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + I*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]] + (I*Sqrt[a]*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])))/d)/a)/(3*a)`

3.178. $\int \frac{(a+ia \tan(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c+dx)\right)}{\tan^{5/2}(c+dx)} dx$

3.178.3 Rubi [A] (verified)

Time = 1.24 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$, Rules used = {3042, 4076, 27, 3042, 4076, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan(c + dx)^{5/2}} dx \\
 & \quad \downarrow \text{4076} \\
 & \frac{2}{3} \int \frac{3(i \tan(c + dx)a + a)^{3/2} ((a + 3ib)B + ia \tan(c + dx)B)}{2 \tan^{3/2}(c + dx)} dx - \frac{bB(a + ia \tan(c + dx))^{3/2}}{d \tan^{3/2}(c + dx)} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(i \tan(c + dx)a + a)^{3/2} ((a + 3ib)B + ia \tan(c + dx)B)}{\tan^{3/2}(c + dx)} dx - \frac{bB(a + ia \tan(c + dx))^{3/2}}{d \tan^{3/2}(c + dx)} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(i \tan(c + dx)a + a)^{3/2} ((a + 3ib)B + ia \tan(c + dx)B)}{\tan(c + dx)^{3/2}} dx - \frac{bB(a + ia \tan(c + dx))^{3/2}}{d \tan^{3/2}(c + dx)} \\
 & \quad \downarrow \text{4076} \\
 & 2 \int \frac{\sqrt{i \tan(c + dx)a + a} (3a(ia - 2b)B - a^2 B \tan(c + dx))}{2\sqrt{\tan(c + dx)}} dx - \frac{bB(a + ia \tan(c + dx))^{3/2}}{d \tan^{3/2}(c + dx)} - \\
 & \quad \frac{2aB(a + 3ib)\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{\sqrt{i \tan(c + dx)a + a} (3a(ia - 2b)B - a^2 B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx - \frac{bB(a + ia \tan(c + dx))^{3/2}}{d \tan^{3/2}(c + dx)} - \\
 & \quad \frac{2aB(a + 3ib)\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.178. $\int \frac{(a + ia \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx$

$$\begin{aligned}
& \int \frac{\sqrt{i \tan(c+dx)a+a}(3a(ia-2b)B-a^2B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx - \frac{bB(a+ia \tan(c+dx))^{3/2}}{d \tan^{3/2}(c+dx)} - \\
& \quad \frac{2aB(a+3ib)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 4084 \\
& 2aB(-3b+2ia) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - iaB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \\
& \quad \frac{bB(a+ia \tan(c+dx))^{3/2}}{d \tan^{3/2}(c+dx)} - \frac{2aB(a+3ib)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 3042 \\
& 2aB(-3b+2ia) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - iaB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \\
& \quad \frac{bB(a+ia \tan(c+dx))^{3/2}}{d \tan^{3/2}(c+dx)} - \frac{2aB(a+3ib)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 4027 \\
& \frac{4ia^3B(-3b+2ia) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} - \\
& iaB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{bB(a+ia \tan(c+dx))^{3/2}}{d \tan^{3/2}(c+dx)} - \\
& \quad \frac{2aB(a+3ib)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 218 \\
& -iaB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + \\
& \frac{(2-2i)a^{3/2}B(-3b+2ia) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{bB(a+ia \tan(c+dx))^{3/2}}{d \tan^{3/2}(c+dx)} - \\
& \quad \frac{2aB(a+3ib)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 4082 \\
& 3.178. \int \frac{(a+ia \tan(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c+dx)\right)}{\tan^{5/2}(c+dx)} dx
\end{aligned}$$

$$\begin{aligned}
& \frac{ia^3 B \int \frac{1}{\sqrt{\tan(c+dx)} \sqrt{ia \tan(c+dx) + a}} d \tan(c+dx)}{d} + \\
& \frac{(2-2i)a^{3/2} B (-3b+2ia) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{bB(a+ia \tan(c+dx))^{3/2}}{d \tan^{3/2}(c+dx)} - \\
& \frac{2aB(a+3ib)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 65 \\
& \frac{2ia^3 B \int \frac{1}{1-\frac{ia \tan(c+dx)}{ia \tan(c+dx)+a}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{ia \tan(c+dx)+a}}}{d} + \frac{(2-2i)a^{3/2} B (-3b+2ia) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \\
& \frac{bB(a+ia \tan(c+dx))^{3/2}}{d \tan^{3/2}(c+dx)} - \frac{2aB(a+3ib)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 216 \\
& \frac{2(-1)^{3/4} a^{5/2} B \operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \\
& \frac{(2-2i)a^{3/2} B (-3b+2ia) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{bB(a+ia \tan(c+dx))^{3/2}}{d \tan^{3/2}(c+dx)} - \\
& \frac{2aB(a+3ib)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}
\end{aligned}$$

input `Int[((a + I*a*Tan[c + d*x])^(5/2)*((3*b*B)/(2*a) + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2), x]`

output `(2*(-1)^(3/4)*a^(5/2)*B*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]]/d + ((2 - 2*I)*a^(3/2)*((2*I)*a - 3*b)*B*ArcTanh[(((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/d - (2*a*(a + (3*I)*b)*B*Sqrt[a + I*a*Tan[c + d*x]]/(d*Sqrt[Tan[c + d*x]]) - (b*B*(a + I*a*Tan[c + d*x])^(3/2))/(d*Tan[c + d*x]^(3/2))`

3.178. $\int \frac{(a+ia \tan(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c+dx)\right)}{\tan^{5/2}(c+dx)} dx$

3.178.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.178.
$$\int \frac{(a+ia \tan(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c+dx) \right)}{\tan^{\frac{5}{2}}(c+dx)} dx$$

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.178.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 617 vs. 2(156) = 312.

Time = 0.14 (sec) , antiderivative size = 618, normalized size of antiderivative = 3.25

method	result
derivativedivides	$\frac{aB\sqrt{a(1+i\tan(dx+c))} \left(6i \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a^2 (\tan^2(dx+c)) - i\sqrt{ia} \sqrt{2} \ln \right)}{}$
default	$\frac{aB\sqrt{a(1+i \tan(dx+c))} \left(6i \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}} \right) \sqrt{-ia} a^2 (\tan^2(dx+c)) - i\sqrt{ia} \sqrt{2} \ln \right)}{}$
parts	$\frac{B \left(-i\sqrt{ia} \sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - ia + 3a \tan(dx+c)}{\tan(dx+c)+i} \right) a \tan(dx+c) - i \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}} \right) \right)}{}$

input `int((a+I*a*tan(d*x+c))^(5/2)*(3/2*b*B/a+B*tan(d*x+c))/tan(d*x+c)^(5/2),x,m
method=_RETURNVERBOSE)`

$$3.178. \int \frac{(a+ia \tan(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c+dx) \right)}{\tan^{\frac{5}{2}}(c+dx)} dx$$

output

```

1/2/d*a*B*(a*(1+I*tan(d*x+c)))^(1/2)*(6*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a^2*tan(d*x+c)^2-I*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2+2*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^2-14*I*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*b-12*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*b*tan(d*x+c)^2-(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2-2*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^2-4*(I*a)^(1/2)*(-I*a)^(1/2)*a*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)-2*b*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)*(-I*a)^(1/2))/tan(d*x+c)^(3/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)/(I*a)^(1/2)/(-I*a)^(1/2)

```

3.178.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 900 vs. $2(146) = 292$.

Time = 0.28 (sec) , antiderivative size = 900, normalized size of antiderivative = 4.74

$$\int \frac{(a + ia \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx = \text{Too large to display}$$

input

```

integrate((a+I*a*tan(d*x+c))^(5/2)*(3/2*b*B/a+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="fracas")

```

3.178.
$$\int \frac{(a + ia \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx$$

output

```

1/2*(2*sqrt(2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(
-(-4*I*B^2*a^5 + 12*B^2*a^4*b + 9*I*B^2*a^3*b^2)/d^2)*log((sqrt(2)*d*sqrt(
-(-4*I*B^2*a^5 + 12*B^2*a^4*b + 9*I*B^2*a^3*b^2)/d^2)*e^(I*d*x + I*c) + sq
rt(2)*(2*I*B*a^2 - 3*B*a*b + (2*I*B*a^2 - 3*B*a*b)*e^(2*I*d*x + 2*I*c))*sq
rt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*
d*x + 2*I*c) + 1)))e^(-I*d*x - I*c)/(2*I*B*a^2 - 3*B*a*b)) - 2*sqrt(2)*(d
*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-4*I*B^2*a^5 +
12*B^2*a^4*b + 9*I*B^2*a^3*b^2)/d^2)*log(-(sqrt(2)*d*sqrt(-(-4*I*B^2*a^5 +
12*B^2*a^4*b + 9*I*B^2*a^3*b^2)/d^2)*e^(I*d*x + I*c) - sqrt(2)*(2*I*B*a^2
- 3*B*a*b + (2*I*B*a^2 - 3*B*a*b)*e^(2*I*d*x + 2*I*c))*sqrt(a/(e^(2*I*d*x
+ 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1
))))e^(-I*d*x - I*c)/(2*I*B*a^2 - 3*B*a*b)) + 4*sqrt(2)*(B*a*b*e^(3*I*d*x
+ 3*I*c) - (I*B*a^2 - 4*B*a*b)*e^(5*I*d*x + 5*I*c) - (-I*B*a^2 + 3*B*a*b)*
e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2
*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + sqrt(4*I*B^2*a^5/d^2)*(d*e^(4*I*d*
x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log((sqrt(2)*(B*a^2*e^(2*I*d*x +
2*I*c) + B*a^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2
*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + I*sqrt(4*I*B^2*a^5/d^2)*d*e^(I*d*x
+ I*c))*e^(-I*d*x - I*c)/(B*a^2)) - sqrt(4*I*B^2*a^5/d^2)*(d*e^(4*I*d*x +
4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log((sqrt(2)*(B*a^2*e^(2*I*d*x + ...

```

3.178.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx = \frac{B \left(\int \frac{2a^3 \sqrt{ia \tan(c + dx) + a}}{\tan^{3/2}(c + dx)} dx + \int \left(-2a^3 \sqrt{ia \tan(c + dx) + a} \right) dx \right)}{1}$$

input

```

integrate((a+I*a*tan(d*x+c))**(5/2)*(3/2*b*B/a+B*tan(d*x+c))/tan(d*x+c)**(
5/2),x)

```

output

```

B*(Integral(2*a**3*sqrt(I*a*tan(c + d*x) + a)/tan(c + d*x)**(3/2), x) + In
tegral(-2*a**3*sqrt(I*a*tan(c + d*x) + a)*sqrt(tan(c + d*x)), x) + Integra
l(4*I*a**3*sqrt(I*a*tan(c + d*x) + a)/sqrt(tan(c + d*x)), x) + Integral(3*
a**2*b*sqrt(I*a*tan(c + d*x) + a)/tan(c + d*x)**(5/2), x) + Integral(-3*a
**2*b*sqrt(I*a*tan(c + d*x) + a)/sqrt(tan(c + d*x)), x) + Integral(6*I*a**2
*b*sqrt(I*a*tan(c + d*x) + a)/tan(c + d*x)**(3/2), x))/(2*a)

```

3.178.
$$\int \frac{(a + ia \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx$$

3.178.7 Maxima [F]

$$\int \frac{(a + ia \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx = \int \frac{(2B \tan(dx + c) + \frac{3Bb}{a})(ia \tan(dx + c) + a)^{5/2}}{2 \tan(dx + c)^{5/2}} dx$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(3/2*b*B/a+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="maxima")`

output `1/2*integrate((2*B*tan(d*x + c) + 3*B*b/a)*(I*a*tan(d*x + c) + a)^(5/2)/tan(d*x + c)^(5/2), x)`

3.178.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(3/2*b*B/a+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[-52]Warning, replacing -52 by -83, a substitution vari`

3.178.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx = \int \frac{(B \tan(c + dx) + \frac{3Bb}{2a})(a + a \tan(c + dx) i)^{5/2}}{\tan(c + dx)^{5/2}} dx$$

input `int(((B*tan(c + d*x) + (3*B*b)/(2*a))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(5/2),x)`

output `int(((B*tan(c + d*x) + (3*B*b)/(2*a))*(a + a*tan(c + d*x)*1i)^(5/2))/tan(c + d*x)^(5/2), x)`

3.178. $\int \frac{(a + ia \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx$

3.179
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

3.179.1 Optimal result 1859
 3.179.2 Mathematica [A] (verified) 1860
 3.179.3 Rubi [A] (verified) 1860
 3.179.4 Maple [B] (verified) 1865
 3.179.5 Fricas [B] (verification not implemented) 1866
 3.179.6 Sympy [F] 1867
 3.179.7 Maxima [F(-2)] 1867
 3.179.8 Giac [F(-2)] 1867
 3.179.9 Mupad [F(-1)] 1868

3.179.1 Optimal result

Integrand size = 38, antiderivative size = 205

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \frac{(-1)^{3/4}(2iA-B) \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ad}} - \frac{\left(\frac{1}{2} - \frac{i}{2}\right)(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ad}} + \frac{(iA-B) \tan^{\frac{3}{2}}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{(A+2iB)\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}{ad}$$

output

```
(-1)^(3/4)*(2*I*A-B)*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)^(1/2))/d/a^(1/2)+(-1/2+1/2*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)^(1/2))/d/a^(1/2)-(A+2*I*B)*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c)^(1/2))/a/d+(I*A-B)*tan(d*x+c)^(3/2)/d/(a+I*a*tan(d*x+c)^(1/2))
```

3.179.2 Mathematica [A] (verified)

Time = 2.76 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.87

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

$$= \frac{\left(\frac{1}{2} + \frac{i}{2}\right)(iA+B) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ad}}$$

$$- \frac{(-1)^{3/4}(2A+iB) \operatorname{arcsinh}\left(\sqrt[4]{-1}\sqrt{\tan(c+dx)}\right) \sqrt{1+i \tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}$$

$$+ \frac{\sqrt{\tan(c+dx)}(-A-2iB+B \tan(c+dx))}{d\sqrt{a+ia \tan(c+dx)}}$$

input `Integrate[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]], x]`

output `((1/2 + I/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/(Sqrt[a]*d) - (((-1)^(3/4)*(2*A + I*B)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]])*Sqrt[1 + I*Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]]) + (Sqrt[Tan[c + d*x]]*(-A - (2*I)*B + B*Tan[c + d*x]))/(d*Sqrt[a + I*a*Tan[c + d*x]])`

3.179.3 Rubi [A] (verified)Time = 1.23 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4078, 27, 3042, 4080, 25, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)^{3/2}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

$$\downarrow \text{4078}$$

3.179. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

$$\begin{aligned}
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\int \frac{1}{2} \sqrt{\tan(c + dx)} \sqrt{i \tan(c + dx) a + a(3a(iA - B) + 2a(A + 2iB) \tan(c + dx))} dx}{a^2} \\
& \quad \downarrow 27 \\
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\int \sqrt{\tan(c + dx)} \sqrt{i \tan(c + dx) a + a(3a(iA - B) + 2a(A + 2iB) \tan(c + dx))} dx}{2a^2} \\
& \quad \downarrow 3042 \\
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\int \sqrt{\tan(c + dx)} \sqrt{i \tan(c + dx) a + a(3a(iA - B) + 2a(A + 2iB) \tan(c + dx))} dx}{2a^2} \\
& \quad \downarrow 4080 \\
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\int -\frac{\sqrt{i \tan(c + dx) a + a(a^2(A + 2iB) - a^2(2iA - B) \tan(c + dx))}}{\sqrt{\tan(c + dx)}} dx + \frac{2a(A + 2iB) \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}}{d}}{2a^2} \\
& \quad \downarrow 25 \\
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\frac{2a(A + 2iB) \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}}{d} - \int \frac{\sqrt{i \tan(c + dx) a + a(a^2(A + 2iB) - a^2(2iA - B) \tan(c + dx))}}{\sqrt{\tan(c + dx)}} dx}{a}}{2a^2} \\
& \quad \downarrow 3042 \\
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\frac{2a(A + 2iB) \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}}{d} - \int \frac{\sqrt{i \tan(c + dx) a + a(a^2(A + 2iB) - a^2(2iA - B) \tan(c + dx))}}{\sqrt{\tan(c + dx)}} dx}{a}}{2a^2} \\
& \quad \downarrow 4084 \\
& \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
& \frac{\frac{2a(A + 2iB) \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}}{d} - \frac{a(2A + iB) \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + a}}{\sqrt{\tan(c + dx)}} dx - a^2(A - iB) \int \frac{\sqrt{i \tan(c + dx) a + a}}{\sqrt{\tan(c + dx)}} dx}{a}}{2a^2}
\end{aligned}$$

3.179. $\int \frac{\tan^{\frac{3}{2}}(c + dx)(A + B \tan(c + dx))}{\sqrt{a + ia \tan(c + dx)}} dx$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2a(A+2iB)\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}{d} - \frac{a(2A+iB) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - a^2(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{a} \\
 & \downarrow 4027 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2a(A+2iB)\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}{d} - \frac{2ia^4(A-iB) \int \frac{1}{-i \tan(c+dx)a+a} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} + a(2A+iB) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}}{a} \\
 & \downarrow 218 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2a(A+2iB)\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}{d} - \frac{a(2A+iB) \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{(1-i)a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}}{a} \\
 & \downarrow 4082 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2a(A+2iB)\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}{d} - \frac{a^3(2A+iB) \int \frac{1}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx) - \frac{(1-i)a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}}{a} \\
 & \downarrow 65 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2a(A+2iB)\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}{d} - \frac{2a^3(2A+iB) \int \frac{1}{1-i \tan(c+dx)a+a} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} - \frac{(1-i)a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}}{a} \\
 & \downarrow 216 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2a(A+2iB)\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}{d} - \frac{2^4\sqrt{-1}a^{5/2}(2A+iB) \operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) - \frac{(1-i)a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}}{a} \\
 & \downarrow 216
 \end{aligned}$$

3.179. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

input `Int[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x
]`

output `((I*A - B)*Tan[c + d*x]^(3/2))/(d*Sqrt[a + I*a*Tan[c + d*x]]) - (-(((-2*(-1)^(1/4)*a^(5/2)*(2*A + I*B)*ArcTan[(-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]]
)/Sqrt[a + I*a*Tan[c + d*x]]])/d - ((1 - I)*a^(5/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d)/a) + (2*a
*(A + (2*I)*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/d)/(2*a^2)`

3.179.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 65 `Int[1/(Sqrt[(b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Sub
st[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d
}, x] && !GtQ[c, 0]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

3.179.
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;`
`FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /;`
`FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

rule 4080 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x], x] /;`
`FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;`
`FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /;`
`FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.179.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1134 vs. $2(165) = 330$.

Time = 0.20 (sec) , antiderivative size = 1135, normalized size of antiderivative = 5.54

method	result	size
derivativedivides	Expression too large to display	1135
default	Expression too large to display	1135
parts	Expression too large to display	1199

```
input int(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x,method=_R
ETURNVERBOSE)
```

```
output 1/4/d*tan(d*x+c)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)/a*(-2*I*B*ln(1/2*(2*I*a*
tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1
/2))*(-I*a)^(1/2)*a-I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*
(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*(I*a)^(1/2)*a+B
*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-
I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*(I*a)^(1/2)*a*tan(d*x+c)^2-2*I*B*2^(1/
2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*
a*tan(d*x+c))/(tan(d*x+c)+I))*(I*a)^(1/2)*a*tan(d*x+c)-8*I*A*ln(1/2*(2*I*a
*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(
1/2))*(-I*a)^(1/2)*a*tan(d*x+c)+8*I*B*(-I*a)^(1/2)*(I*a)^(1/2)*(a*tan(d*x+
c)*(1+I*tan(d*x+c)))^(1/2)+2*A*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1
/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+
I))*a*tan(d*x+c)-4*I*B*(-I*a)^(1/2)*(I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x
+c)))^(1/2)*tan(d*x+c)^2+2*I*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1
+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c
)^2+4*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(
I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^2-B*(I*a)^(1/2)*2^(1/
2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*
a*tan(d*x+c))/(tan(d*x+c)+I))*a+4*I*A*(-I*a)^(1/2)*(I*a)^(1/2)*(a*tan(d*x+
c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^...
```

$$3.179. \quad \int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

3.179.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 807 vs. $2(155) = 310$.

Time = 0.34 (sec) , antiderivative size = 807, normalized size of antiderivative = 3.94

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx =$$

$$\left(\sqrt{2}ad\sqrt{-\frac{iA^2+2AB-iB^2}{ad^2}}e^{(idx+ic)} \log\left(\frac{i\sqrt{2}ad\sqrt{-\frac{iA^2+2AB-iB^2}{ad^2}}e^{(idx+ic)}+\sqrt{2}((iA+B)e^{(2idx+2ic)}+iA+B)}{4iA+4B}\right)\sqrt{\frac{a}{e^{(2idx+2ic)}}}} \right)$$

```
input integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="fricas")
```

```
output -1/4*(sqrt(2)*a*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)/(a*d^2))*e^(I*d*x + I*c)*log((I*sqrt(2)*a*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)/(a*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - sqrt(2)*a*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)/(a*d^2))*e^(I*d*x + I*c)*log((-I*sqrt(2)*a*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)/(a*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - a*d*sqrt((-4*I*A^2 + 4*A*B + I*B^2)/(a*d^2))*e^(I*d*x + I*c)*log(104/605*(2*sqrt(2)*((2*I*A - B)*e^(3*I*d*x + 3*I*c) + (2*I*A - B)*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + (3*I*a*d*e^(2*I*d*x + 2*I*c) - I*a*d)*sqrt((-4*I*A^2 + 4*A*B + I*B^2)/(a*d^2)))/((2*I*A - B)*e^(2*I*d*x + 2*I*c) + 2*I*A - B)) + a*d*sqrt((-4*I*A^2 + 4*A*B + I*B^2)/(a*d^2))*e^(I*d*x + I*c)*log(104/605*(2*sqrt(2)*((2*I*A - B)*e^(3*I*d*x + 3*I*c) + (2*I*A - B)*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + (-3*I*a*d*e^(2*I*d*x + 2*I*c) + I*a*d)*sqrt((-4*I*A^2 + 4*A*B + I*B^2)/(a*d^2)))/((2*I*A - B)*e^(2*I*d*x + 2*I*c) + 2*I*A - B)) + 2*sqrt(2)*((A + 3*I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + ...
```

3.179.6 Sympy [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{(A+B\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)}{\sqrt{ia(\tan(c+dx)-i)}} dx$$

input `integrate(tan(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**(3/2)/sqrt(I*a*(tan(c + d*x) - I)), x)`

3.179.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.179.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \text{Exception raised: TypeError}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[24]Warning, replacing 24 by 84, a substitution variabl`

3.179. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx$

3.179.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{\tan(c+dx)^{\frac{3}{2}}(A+B\tan(c+dx))}{\sqrt{a+a\tan(c+dx)} \operatorname{li}} dx$$

input `int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2),x)`

output `int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2), x)`

3.180
$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

3.180.1 Optimal result 1869
 3.180.2 Mathematica [A] (verified) 1870
 3.180.3 Rubi [A] (verified) 1870
 3.180.4 Maple [B] (verified) 1874
 3.180.5 Fricas [B] (verification not implemented) 1875
 3.180.6 Sympy [F] 1876
 3.180.7 Maxima [F(-2)] 1876
 3.180.8 Giac [F(-2)] 1876
 3.180.9 Mupad [B] (verification not implemented) 1877

3.180.1 Optimal result

Integrand size = 38, antiderivative size = 156

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = -\frac{2\sqrt[4]{-1}B \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ad}} - \frac{\left(\frac{1}{2} + \frac{i}{2}\right)(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ad}} + \frac{(iA-B)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}$$

output

```
-2*(-1)^(1/4)*B*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d/a^(1/2)-(1/2+1/2*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d/a^(1/2)+(I*A-B)*tan(d*x+c)^(1/2)/d/(a+I*a*tan(d*x+c))^(1/2)
```

3.180.2 Mathematica [A] (verified)

Time = 2.10 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.01

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = -\frac{\left(\frac{1}{2} + \frac{i}{2}\right)(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{\sqrt{ad}} + \frac{\left(-2\sqrt[4]{-1}B\operatorname{Arcsinh}\left(\sqrt[4]{-1}\sqrt{\tan(c+dx)}\right)\sqrt{1+i\tan(c+dx)} + (A+iB)\sqrt{\tan(c+dx)}\right)\sqrt{a+ia\tan(c+dx)}}{ad(-i+\tan(c+dx))}$$

input `Integrate[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]], x]`

output `((-1/2 - I/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/(Sqrt[a]*d) + ((-2*(-1)^(1/4)*B*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]]*Sqrt[1 + I*Tan[c + d*x]] + (A + I*B)*Sqrt[Tan[c + d*x]])*Sqrt[a + I*a*Tan[c + d*x]]/(a*d*(-I + Tan[c + d*x]))`

3.180.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4078, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx \\ & \quad \downarrow \text{4078} \\ & \frac{(-B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{\int \frac{\sqrt{i\tan(c+dx)a+a(a(iA-B)+2iaB\tan(c+dx))}}{2\sqrt{\tan(c+dx)}} dx}{a^2} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.180. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx$

$$\begin{aligned}
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA-B)+2iaB \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{2a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA-B)+2iaB \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{2a^2} \\
& \quad \downarrow \text{4084} \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{a(B + iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - 2B \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{a(B + iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - 2B \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a^2} \\
& \quad \downarrow \text{4027} \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{2ia^3(B+iA) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} - 2B \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a^2} \\
& \quad \downarrow \text{218} \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{(1-i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) - 2B \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a^2} \\
& \quad \downarrow \text{4082} \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{(1-i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) - \frac{2a^2 B \int \frac{1}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{d}}{2a^2} \\
& \quad \downarrow \text{65}
\end{aligned}$$

3.180. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

$$\frac{\frac{(-B + iA)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{(1-i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{4a^2 B \int \frac{1}{1 - \frac{ia \tan(c+dx)}{i \tan(c+dx)a+a}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d}}{2a^2}$$

↓ 216

$$\frac{\frac{(-B + iA)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{(1-i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{4\sqrt{-1}a^{3/2}B \operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d}}{2a^2}$$

input `Int[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]], x]`

output `-1/2*((4*(-1)^(1/4)*a^(3/2)*B*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d + ((1 - I)*a^(3/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d)/a^2 + ((I*A - B)*Sqrt[Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]])`

3.180.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.180.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 893 vs. $2(123) = 246$.

Time = 0.15 (sec) , antiderivative size = 894, normalized size of antiderivative = 5.73

method	result
derivativedivides	$\frac{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})\left(iB\sqrt{ia}\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)\right)}{a(\tan^2(dx+c)+1)}$
default	$\frac{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})\left(iB\sqrt{ia}\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)\right)}{a(\tan^2(dx+c)+1)}$
parts	Expression too large to display

input `int(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x,method=_RETURVERBOSE)`

output `1/4/d*tan(d*x+c)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)/a*(I*B*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2+2*I*A*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)-A*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2-I*B*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a-8*I*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)+4*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+2*B*(I*a)^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)+4*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^2-4*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+A*(I*a)^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a+4*A*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)-4*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a+4*B*(I*a)^(1/2)*(-I*a)^(1/2)*...`

3.180.
$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx$$

3.180.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 718 vs. $2(116) = 232$.

Time = 0.33 (sec) , antiderivative size = 718, normalized size of antiderivative = 4.60

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx =$$

$$\left(\sqrt{2}ad \sqrt{-\frac{-iA^2-2AB+iB^2}{ad^2}} e^{(i dx+i c)} \log \left(\frac{\sqrt{2}ad \sqrt{-\frac{-iA^2-2AB+iB^2}{ad^2}} e^{(i dx+i c)} + \sqrt{2}((iA+B)e^{(2i dx+2i c)} + iA+B)}{4iA+4B} \sqrt{\frac{a}{e^{(2i dx+2i c)}}} \right) \right)$$

```
input integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="fricas")
```

```
output -1/4*(sqrt(2)*a*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2))*e^(I*d*x + I*c)*
log((sqrt(2)*a*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2))*e^(I*d*x + I*c) +
sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*
I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(
4*I*A + 4*B)) - sqrt(2)*a*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2))*e^(I*d
*x + I*c)*log(-(sqrt(2)*a*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2))*e^(I*d
*x + I*c) - sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2
*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*
c) + 1)))/(4*I*A + 4*B)) - a*d*sqrt(-4*I*B^2/(a*d^2))*e^(I*d*x + I*c)*log(
52/605*(4*sqrt(2)*(B*e^(3*I*d*x + 3*I*c) + B*e^(I*d*x + I*c))*sqrt(a/(e^(2
*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*
c) + 1)) + (3*a*d*e^(2*I*d*x + 2*I*c) - a*d)*sqrt(-4*I*B^2/(a*d^2)))/(B*e^(
2*I*d*x + 2*I*c) + B)) + a*d*sqrt(-4*I*B^2/(a*d^2))*e^(I*d*x + I*c)*log(5
2/605*(4*sqrt(2)*(B*e^(3*I*d*x + 3*I*c) + B*e^(I*d*x + I*c))*sqrt(a/(e^(2*
I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c
) + 1)) - (3*a*d*e^(2*I*d*x + 2*I*c) - a*d)*sqrt(-4*I*B^2/(a*d^2)))/(B*e^(
2*I*d*x + 2*I*c) + B)) + 2*sqrt(2)*((-I*A + B)*e^(2*I*d*x + 2*I*c) - I*A +
B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e
^(2*I*d*x + 2*I*c) + 1))*e^(-I*d*x - I*c)/(a*d)
```

3.180.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{(A+B\tan(c+dx))\sqrt{\tan(c+dx)}}{\sqrt{ia(\tan(c+dx)-i)}} dx$$

input `integrate(tan(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(tan(c + d*x))/sqrt(I*a*(tan(c + d*x) - I)), x)`

3.180.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.180.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \text{Exception raised: TypeError}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[-7]Warning, replacing -7 by -3, a substitution variabl`

3.180. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx$

3.180.9 Mupad [B] (verification not implemented)

Time = 28.33 (sec) , antiderivative size = 4040, normalized size of antiderivative = 25.90

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \text{Too large to display}$$

```
input int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2),x)
```

```
output -(A*a^(5/2)*tan(c + d*x)^(1/2)*4i + 4*A*a^(5/2)*tan(c + d*x)^(3/2) - 4*B*a^(5/2)*tan(c + d*x)^(1/2) + B*a^(5/2)*tan(c + d*x)^(3/2)*4i + A*a^(3/2)*tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)*4i - 4*B*a^(3/2)*tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i) + (1i/8)^(1/2)*A*(-a)^(5/2)*atanh(((1i/8)^(1/2)*(-a)^(15/2)*a^(17/2)*tan(c + d*x)^(1/2)*4i - 4*(1i/8)^(1/2)*(-a)^(31/2)*tan(c + d*x)^(3/2)*(a + a*tan(c + d*x)*1i)^(1/2) - (1i/8)^(1/2)*(-a)^(31/2)*tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)*8i + 4*(1i/8)^(1/2)*(-a)^(15/2)*a^(17/2)*tan(c + d*x)^(3/2) + (1i/8)^(1/2)*(-a)^(15/2)*a^(15/2)*tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)*4i)/(a^15*(a + a*tan(c + d*x)*1i) - 2*a^(31/2)*(a + a*tan(c + d*x)*1i)^(1/2) + a^16 + a^16*tan(c + d*x)^2)*8i + 8*(1i/8)^(1/2)*B*(-a)^(5/2)*atanh(((1i/8)^(1/2)*(-a)^(15/2)*a^(17/2)*tan(c + d*x)^(1/2)*4i - 4*(1i/8)^(1/2)*(-a)^(31/2)*tan(c + d*x)^(3/2)*(a + a*tan(c + d*x)*1i)^(1/2) - (1i/8)^(1/2)*(-a)^(31/2)*tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)*8i + 4*(1i/8)^(1/2)*(-a)^(15/2)*a^(17/2)*tan(c + d*x)^(3/2) + (1i/8)^(1/2)*(-a)^(15/2)*a^(15/2)*tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)*4i)/(a^15*(a + a*tan(c + d*x)*1i) - 2*a^(31/2)*(a + a*tan(c + d*x)*1i)^(1/2) + a^16 + a^16*tan(c + d*x)^2) - A*a^2*tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)*8i - 4*A*a^2*tan(c + d*x)^(3/2)*(a + a*tan(c + d*x)*1i)^(1/2) + 8*B*a^2*tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2) - B*a^2*tan(c + d*x)^(3/2)*(a + a*tan(c + d*x)*1i)^(1/2)*4i + ...
```

3.181
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} dx$$

3.181.1 Optimal result 1878
 3.181.2 Mathematica [A] (verified) 1878
 3.181.3 Rubi [A] (verified) 1879
 3.181.4 Maple [B] (verified) 1881
 3.181.5 Fricas [B] (verification not implemented) 1882
 3.181.6 Sympy [F] 1882
 3.181.7 Maxima [F(-2)] 1883
 3.181.8 Giac [F(-2)] 1883
 3.181.9 Mupad [B] (verification not implemented) 1884

3.181.1 Optimal result

Integrand size = 38, antiderivative size = 99

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}} dx = \frac{\left(\frac{1}{2} - \frac{i}{2}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ad}} + \frac{(A + iB)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}}$$

output `(1/2-1/2*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d/a^(1/2)+(A+I*B)*tan(d*x+c)^(1/2)/d/(a+I*a*tan(d*x+c))^(1/2)`

3.181.2 Mathematica [A] (verified)

Time = 1.37 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.12

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}} dx = \frac{\sqrt{\tan(c + dx)} \left(\frac{\sqrt{2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ia \tan(c+dx)}} + \frac{2(A+iB)}{\sqrt{a+ia \tan(c+dx)}} \right)}{2d}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]),x]`

output $(\text{Sqrt}[\text{Tan}[c + d*x]] * ((\text{Sqrt}[2] * (A - I*B) * \text{ArcTanh}[(\text{Sqrt}[2] * \text{Sqrt}[I*a*\text{Tan}[c + d*x]])] / \text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]) / \text{Sqrt}[I*a*\text{Tan}[c + d*x]] + (2*(A + I*B)) / \text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])) / (2*d)$

3.181.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {3042, 4079, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{a(A - iB) \sqrt{i \tan(c + dx) a + a}}{2 \sqrt{\tan(c + dx)}} dx}{a^2} + \frac{(A + iB) \sqrt{\tan(c + dx)}}{d \sqrt{a + ia \tan(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{(A - iB) \int \frac{\sqrt{i \tan(c + dx) a + a}}{\sqrt{\tan(c + dx)}} dx}{2a} + \frac{(A + iB) \sqrt{\tan(c + dx)}}{d \sqrt{a + ia \tan(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A - iB) \int \frac{\sqrt{i \tan(c + dx) a + a}}{\sqrt{\tan(c + dx)}} dx}{2a} + \frac{(A + iB) \sqrt{\tan(c + dx)}}{d \sqrt{a + ia \tan(c + dx)}} \\
 & \quad \downarrow \text{4027} \\
 & \frac{(A + iB) \sqrt{\tan(c + dx)}}{d \sqrt{a + ia \tan(c + dx)}} - \frac{ia(A - iB) \int \frac{1}{-\frac{2 \tan(c + dx) a^2}{i \tan(c + dx) a + a} - ia} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{i \tan(c + dx) a + a}}}{d} \\
 & \quad \downarrow \text{218}
 \end{aligned}$$

3.181. $\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx$

$$\frac{\left(\frac{1}{2} - \frac{i}{2}\right)(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{\sqrt{ad}} + \frac{(A + iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]),x]`

output `((1/2 - I/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/(Sqrt[a]*d) + ((A + I*B)*Sqrt[Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]])`

3.181.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.181.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 632 vs. 2(79) = 158.

Time = 0.17 (sec) , antiderivative size = 633, normalized size of antiderivative = 6.39

method	result
derivativedivides	$-\frac{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})\left(iA\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)\right)}{a(\tan^2(dx+c))}$
default	$-\frac{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})\left(iA\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)\right)}{a(\tan^2(dx+c))}$
parts	$-\frac{A\sqrt{a(1+i\tan(dx+c))}(\sqrt{\tan(dx+c)})\left(i\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)\right)}{(\tan^2(dx+c))}$

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2),x,method=_R
ETURNVERBOSE)
```

```
output -1/4/d*tan(d*x+c)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)*(I*A*2^(1/2)*ln((2*2^(1
/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c)
)/(tan(d*x+c)+I))*a*tan(d*x+c)^2-2*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(
a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a
*tan(d*x+c)+B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*
x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2-I*A*2^(1/2
)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a
*tan(d*x+c))/(tan(d*x+c)+I))*a+4*I*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d
*x+c)))^(1/2)*tan(d*x+c)+2*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x
+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+
c)+4*I*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-B*2^(1/2)*ln((
2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d
*x+c))/(tan(d*x+c)+I))*a-4*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^
(1/2)*tan(d*x+c)+4*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2))/a
/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)/(-tan(d*x+c)+I)^2/(-I*a)^(1/2)
```

$$3.181. \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} dx$$

3.181.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 416 vs. $2(73) = 146$.

Time = 0.27 (sec) , antiderivative size = 416, normalized size of antiderivative = 4.20

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx$$

$$\left(\sqrt{2ad} \sqrt{-\frac{iA^2 + 2AB - iB^2}{ad^2}} e^{i dx + ic} \log \left(\frac{i \sqrt{2ad} \sqrt{-\frac{iA^2 + 2AB - iB^2}{ad^2}} e^{i dx + ic} + \sqrt{2}((iA + B)e^{2i dx + 2ic} + iA + B)}{4iA + 4B} \sqrt{\frac{a}{e^{2i dx + 2ic} + 1}} \right) \right)$$

```
input integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="fricas")
```

```
output 1/4*(sqrt(2)*a*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)/(a*d^2))*e^(I*d*x + I*c)*log((I*sqrt(2)*a*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)/(a*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - sqrt(2)*a*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)/(a*d^2))*e^(I*d*x + I*c)*log((-I*sqrt(2)*a*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)/(a*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) + 2*sqrt(2)*((A + I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*e^(-I*d*x - I*c)/(a*d)
```

3.181.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{ia (\tan(c + dx) - i)} \sqrt{\tan(c + dx)}} dx$$

```
input integrate((A+B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+I*a*tan(d*x+c))**(1/2),x)
```

```
output Integral((A + B*tan(c + d*x))/(sqrt(I*a*(tan(c + d*x) - I))*sqrt(tan(c + d*x))), x)
```

3.181. $\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx$

3.181.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.181.8 Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx = \text{Exception raised: NotImplementedError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Exception raised: NotImplementedError >> unable to parse Giac output: 2*((3*sqrt(abs(sageVARa))*sageVARB*sageVARa+3*i*abs(sageVARa)*sqrt(abs(sageVARa))*sageVARB+3*i*sqrt(abs(sageVARa))*sageVARa*sageVARa-3*abs(sageVARa)*sqrt(abs(sageVARa))*sa`

3.181.9 Mupad [B] (verification not implemented)

Time = 14.09 (sec) , antiderivative size = 426, normalized size of antiderivative = 4.30

$$\begin{aligned}
& \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx \\
&= \frac{B \ln \left(\frac{\sqrt{a} \sqrt{\tan(c + dx)} (2 - 2i)}{\sqrt{a + a \tan(c + dx)} \overline{1i - \sqrt{a}}} - \frac{a \tan(c + dx)}{(\sqrt{a + a \tan(c + dx)} \overline{1i - \sqrt{a}})^2} + 1i \right) \left(\frac{1}{4} + \frac{1}{4}i \right)}{\sqrt{a} d} \\
&+ \frac{A \sqrt{\tan(c + dx)} 2i}{\left(\sqrt{a + a \tan(c + dx)} \overline{1i - \sqrt{a}} \right) \left(d \overline{1i} - \frac{a d \tan(c + dx)}{(\sqrt{a + a \tan(c + dx)} \overline{1i - \sqrt{a}})^2} \right)} \\
&- \frac{2 B \sqrt{\tan(c + dx)}}{\left(\sqrt{a + a \tan(c + dx)} \overline{1i - \sqrt{a}} \right) \left(d \overline{1i} - \frac{a d \tan(c + dx)}{(\sqrt{a + a \tan(c + dx)} \overline{1i - \sqrt{a}})^2} \right)} \\
&- \frac{\sqrt{\frac{1}{8}i} B \ln \left(-\frac{a \tan(c + dx)}{(\sqrt{a + a \tan(c + dx)} \overline{1i - \sqrt{a}})^2} + \frac{2(-1)^{3/4} \sqrt{2} \sqrt{a} \sqrt{\tan(c + dx)}}{\sqrt{a + a \tan(c + dx)} \overline{1i - \sqrt{a}}} + 1i \right)}{\sqrt{a} d} \\
&+ \frac{2 \sqrt{\frac{1}{8}i} A \operatorname{atanh} \left(\frac{32 \sqrt{\frac{1}{8}i} A^2 (-a)^{9/2} \sqrt{\tan(c + dx)}}{\left(A^2 a^4 \overline{4i} - \frac{4 A^2 a^5 \tan(c + dx)}{(\sqrt{a + a \tan(c + dx)} \overline{1i - \sqrt{a}})^2} \right) \left(\sqrt{a + a \tan(c + dx)} \overline{1i - \sqrt{a}} \right)} \right)}{\sqrt{-a} d}
\end{aligned}$$

```
input int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)),x)
```

```
output (B*log((a^(1/2)*tan(c + d*x)^(1/2)*(2 - 2i))/((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2)) - a^(1/2)) - (a*tan(c + d*x))/((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2))^2 + 1i)*(1/4 + 1i/4))/(a^(1/2)*d) + (A*tan(c + d*x)^(1/2)*2i)/(((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2))*(d*1i - (a*d*tan(c + d*x))/((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2))^2)) - (2*B*tan(c + d*x)^(1/2))/(((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2))*(d*1i - (a*d*tan(c + d*x))/((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2))^2)) - ((1i/8)^(1/2)*B*log((2*(-1)^(3/4)*2^(1/2)*a^(1/2)*tan(c + d*x)^(1/2))/((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2)) - (a*tan(c + d*x))/((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2))^2 + 1i))/(a^(1/2)*d) + (2*(1i/8)^(1/2)*A*atanh((32*(1i/8)^(1/2)*A^2*(-a)^(9/2)*tan(c + d*x)^(1/2))/((A^2*a^4*4i - (4*A^2*a^5*tan(c + d*x))/((a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2))^2)*(a + a*tan(c + d*x)*1i)^(1/2) - a^(1/2))))/((-a)^(1/2)*d)
```

$$3.182 \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}} dx$$

3.182.1 Optimal result	1885
3.182.2 Mathematica [A] (verified)	1885
3.182.3 Rubi [A] (verified)	1886
3.182.4 Maple [B] (verified)	1888
3.182.5 Fricas [B] (verification not implemented)	1889
3.182.6 Sympy [F]	1890
3.182.7 Maxima [F(-2)]	1891
3.182.8 Giac [F(-2)]	1891
3.182.9 Mupad [F(-1)]	1891

3.182.1 Optimal result

Integrand size = 38, antiderivative size = 143

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}} dx = \frac{\left(\frac{1}{2} + \frac{i}{2}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ad}} + \frac{A + iB}{d\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}} - \frac{(3A + iB)\sqrt{a + ia \tan(c + dx)}}{ad\sqrt{\tan(c + dx)}}$$

output $(1/2+1/2*I)*(A-I*B)*\operatorname{arctanh}((1+I)*a^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+I*a*\tan(d*x+c))^{(1/2)})/d/a^{(1/2)}+(A+I*B)/d/\tan(d*x+c)^{(1/2)}/(a+I*a*\tan(d*x+c))^{(1/2)}-(3*A+I*B)*(a+I*a*\tan(d*x+c))^{(1/2)}/a/d/\tan(d*x+c)^{(1/2)}$

3.182.2 Mathematica [A] (verified)

Time = 1.57 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.90

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}} dx = \frac{\sqrt{2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \tan(c+dx)}{\sqrt{ia \tan(c+dx)}} + \frac{-4A+2(-3iA+B) \tan(c+dx)}{\sqrt{a+ia \tan(c+dx)}}}{2d\sqrt{\tan(c + dx)}}$$

3.182. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}} dx$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]]),x]`

output `((Sqrt[2]*(I*A + B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Tan[c + d*x])/Sqrt[I*a*Tan[c + d*x]] + (-4*A + 2*((-3*I)*A + B)*Tan[c + d*x])/Sqrt[a + I*a*Tan[c + d*x]]/(2*d*Sqrt[Tan[c + d*x]])`

3.182.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.01, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.237$, Rules used = {3042, 4079, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2} \sqrt{a + ia \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{\sqrt{i \tan(c + dx) a + a(a(3A + iB) - 2a(iA - B) \tan(c + dx))}}{2 \tan^{\frac{3}{2}}(c + dx)} dx}{a^2} + \frac{A + iB}{d \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\sqrt{i \tan(c + dx) a + a(a(3A + iB) - 2a(iA - B) \tan(c + dx))}}{\tan^{\frac{3}{2}}(c + dx)} dx}{2a^2} + \frac{A + iB}{d \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\sqrt{i \tan(c + dx) a + a(a(3A + iB) - 2a(iA - B) \tan(c + dx))}}{\tan(c + dx)^{3/2}} dx}{2a^2} + \frac{A + iB}{d \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}} \\
 & \quad \downarrow \text{4081} \\
 & \frac{2 \int \frac{a^2(iA + B) \sqrt{i \tan(c + dx) a + a}}{2 \sqrt{\tan(c + dx)}} dx}{a} - \frac{2a(3A + iB) \sqrt{a + ia \tan(c + dx)}}{d \sqrt{\tan(c + dx)}} + \frac{A + iB}{d \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}}
 \end{aligned}$$

3.182. $\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{a(B + iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a(3A+iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{A + iB}{d\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}} \\
 & \downarrow 3042 \\
 & \frac{a(B + iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a(3A+iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{A + iB}{d\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}} \\
 & \downarrow 4027 \\
 & - \frac{2ia^3(B+iA) \int \frac{1}{\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d - \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{2a^2} - \frac{2a(3A+iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \\
 & \frac{A + iB}{d\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}} \\
 & \downarrow 218 \\
 & \frac{(1-i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a(3A+iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \\
 & \frac{A + iB}{d\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}
 \end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]]),x]`

output `(A + I*B)/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + (((1 - I)*a^(3/2)*(I*A + B)*ArcTanh[(((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/d - (2*a*(3*A + I*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(2*a^2)`

3.182.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

3.182. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

3.182.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 694 vs. $2(117) = 234$.

Time = 0.18 (sec) , antiderivative size = 695, normalized size of antiderivative = 4.86

3.182.
$$\int \frac{A+B \tan(c+dx)}{\tan^3(c+dx) \sqrt{a+ia \tan(c+dx)}} dx$$

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))}}{\tan(dx+c)+i} \left(iB \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \sqrt{2} a (\tan^3(dx+c)) + 2iA\sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \right)$
default	$\frac{\sqrt{a(1+i \tan(dx+c))}}{\tan(dx+c)+i} \left(iB \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \sqrt{2} a (\tan^3(dx+c)) + 2iA\sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \right)$
parts	$\frac{A\sqrt{a(1+i \tan(dx+c))}}{\tan(dx+c)+i} \left(2i\sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i} \right) (\tan^2(dx+c)) a - \sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))-ia+3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \right)$

```
input int((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(3/2),x,method=_R
RETURNVERBOSE)
```

```
output -1/4/d*(a*(1+I*tan(d*x+c)))^(1/2)*(I*B*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d
*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*
a*tan(d*x+c)^3+2*I*A*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x
+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^2-A*ln
((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*ta
n(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^3-I*B*ln((2*2^(1/2)*(-I*a)^(
1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c
)+I))*2^(1/2)*a*tan(d*x+c)+4*I*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c
)))^(1/2)*tan(d*x+c)^2+2*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c
)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)
^2-20*I*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+A*
2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I
*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)+12*A*(-I*a)^(1/2)*(a*tan(d
*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2+4*B*(-I*a)^(1/2)*(a*tan(d*x+c)*
(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)-8*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*ta
n(d*x+c)))^(1/2))/a/tan(d*x+c)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)/
(-I*a)^(1/2)/(-tan(d*x+c)+I)^2
```

3.182.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 473 vs. 2(109) = 218.

3.182.
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}} dx$$

Time = 0.28 (sec) , antiderivative size = 473, normalized size of antiderivative = 3.31

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx$$

$$= \frac{\sqrt{2}(ade^{(3i dx + 3i c)} - ade^{(i dx + i c)}) \sqrt{-\frac{-i A^2 - 2AB + i B^2}{ad^2}} \log \left(\frac{\sqrt{2}ad \sqrt{-\frac{-i A^2 - 2AB + i B^2}{ad^2}} e^{(i dx + i c)} + \sqrt{2}((i A + B)e^{(2i dx + 2i c)} + i A + B)}{4i A + 4B} \right)}{4i A + 4B}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(3/2),x, algorithm="fricas")`

output `1/4*(sqrt(2)*(a*d*e^(3*I*d*x + 3*I*c) - a*d*e^(I*d*x + I*c))*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2))*log((sqrt(2)*a*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - sqrt(2)*(a*d*e^(3*I*d*x + 3*I*c) - a*d*e^(I*d*x + I*c))*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2))*log(-sqrt(2)*a*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2))*e^(I*d*x + I*c) - sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - 2*sqrt(2)*((5*I*A - B)*e^(4*I*d*x + 4*I*c) + 4*I*A*e^(2*I*d*x + 2*I*c) - I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(a*d*e^(3*I*d*x + 3*I*c) - a*d*e^(I*d*x + I*c))`

3.182.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{ia (\tan(c + dx) - i)} \tan^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/2)/tan(d*x+c)**(3/2),x)`

output `Integral((A + B*tan(c + d*x))/(sqrt(I*a*(tan(c + d*x) - I))*tan(c + d*x)**(3/2)), x)`

3.182.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(3/2),x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.
```

3.182.8 Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx = \text{Exception raised: NotImplementedError}$$

```
input integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(3/2),x, algorithm="giac")
```

```
output Exception raised: NotImplementedError >> unable to parse Giac output: -2*i*sageVARa*(-i)/sageVARa/sageVARd*sqrt(i*sageVARa*tan(sageVARc+sageVARd*sageVARx)+sageVARa)*sqrt(2*sageVARa^2-2*sageVARa*(sqrt(i*sageVARa*tan(sageVARc+sageVARd*sageVARx
```

3.182.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2} \sqrt{a + a \tan(c + dx)} li}$$

```
input int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + a*tan(c + d*x)*1i)^(1/2)),x)
```

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + a*tan(c + d*x)*i)^(1/2)), x)`

3.182. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} dx$

3.183 $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}} dx$

3.183.1 Optimal result 1893
 3.183.2 Mathematica [A] (verified) 1894
 3.183.3 Rubi [A] (verified) 1894
 3.183.4 Maple [B] (verified) 1898
 3.183.5 Fricas [B] (verification not implemented) 1899
 3.183.6 Sympy [F] 1900
 3.183.7 Maxima [F(-2)] 1900
 3.183.8 Giac [F(-2)] 1900
 3.183.9 Mupad [F(-1)] 1901

3.183.1 Optimal result

Integrand size = 38, antiderivative size = 191

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}} dx = \frac{\left(\frac{1}{2} + \frac{i}{2}\right) (iA + B) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ad}} + \frac{A + iB}{d \tan^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}} - \frac{(5A + 3iB)\sqrt{a + ia \tan(c + dx)}}{3ad \tan^{\frac{3}{2}}(c + dx)} + \frac{(7iA - 9B)\sqrt{a + ia \tan(c + dx)}}{3ad\sqrt{\tan(c + dx)}}$$

```
output (1/2+1/2*I)*(I*A+B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/d/a^(1/2)+1/3*(7*I*A-9*B)*(a+I*a*tan(d*x+c))^(1/2)/a/d/tan(d*x+c)^(1/2)+(A+I*B)/d/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(3/2)-1/3*(5*A+3*I*B)*(a+I*a*tan(d*x+c))^(1/2)/a/d/tan(d*x+c)^(3/2)
```

3.183.2 Mathematica [A] (verified)

Time = 3.31 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.79

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx$$

$$= \frac{-\frac{3\sqrt{2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \tan^2(c+dx)}{\sqrt{ia \tan(c+dx)}} - \frac{2(2A+(-2iA+6B) \tan(c+dx)+(7A+9iB) \tan^2(c+dx))}{\sqrt{a+ia \tan(c+dx)}}}{6d \tan^{\frac{3}{2}}(c + dx)}$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*Sqrt[a + I*a*Tan[c + d*x]]),x]`

output `((-3*Sqrt[2]*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Tan[c + d*x]^2/Sqrt[I*a*Tan[c + d*x]] - (2*(2*A + ((-2*I)*A + 6*B)*Tan[c + d*x] + (7*A + (9*I)*B)*Tan[c + d*x]^2))/Sqrt[a + I*a*Tan[c + d*x]])/(6*d*Tan[c + d*x]^(3/2))`

3.183.3 Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.04, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {3042, 4079, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2} \sqrt{a + ia \tan(c + dx)}} dx$$

↓ 4079

$$\frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(5A+3iB)-4a(iA-B) \tan(c+dx))}{2 \tan^{\frac{5}{2}}(c+dx)} dx}{a^2} + \frac{A + iB}{d \tan^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}$$

↓ 27

3.183. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} dx$

$$\begin{aligned}
& \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(5A+3iB)-4a(iA-B) \tan(c+dx)) dx}{\tan^{\frac{5}{2}}(c+dx)}}{2a^2} + \frac{A+iB}{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(5A+3iB)-4a(iA-B) \tan(c+dx)) dx}{\tan(c+dx)^{5/2}}}{2a^2} + \frac{A+iB}{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 4081 \\
& \frac{2 \int -\frac{\sqrt{i \tan(c+dx)a+a}((7iA-9B)a^2+2(5A+3iB) \tan(c+dx)a^2) dx}{2 \tan^{\frac{3}{2}}(c+dx)}}{3a} - \frac{2a(5A+3iB) \sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((7iA-9B)a^2+2(5A+3iB) \tan(c+dx)a^2) dx}{\tan^{\frac{3}{2}}(c+dx)}}{3a} - \frac{2a(5A+3iB) \sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((7iA-9B)a^2+2(5A+3iB) \tan(c+dx)a^2) dx}{\tan(c+dx)^{3/2}}}{3a} - \frac{2a(5A+3iB) \sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 4081 \\
& \frac{2 \int \frac{3a^3(A-iB) \sqrt{i \tan(c+dx)a+a} dx}{2 \sqrt{\tan(c+dx)}} - \frac{2a^2(-9B+7iA) \sqrt{a+ia \tan(c+dx)}}{d \sqrt{\tan(c+dx)}}}{3a} - \frac{2a(5A+3iB) \sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 27
\end{aligned}$$

3.183. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} dx$

$$\begin{aligned}
& - \frac{3a^2(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(-9B+7iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2a(5A+3iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \frac{2a^2}{A+iB} \\
& \frac{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{3042} \\
& - \frac{3a^2(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(-9B+7iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2a(5A+3iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \frac{2a^2}{A+iB} \\
& \frac{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{4027} \\
& - \frac{6ia^4(A-iB) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d - \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{3a} - \frac{2a^2(-9B+7iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a(5A+3iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \frac{2a^2}{A+iB} \\
& \frac{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{218} \\
& - \frac{(3-3i)a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^2(-9B+7iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a(5A+3iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \frac{2a^2}{A+iB} \\
& \frac{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{
\end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*Sqrt[a + I*a*Tan[c + d*x]]),x]`

output `(A + I*B)/(d*Tan[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]]) + ((-2*a*(5*A + (3*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (((3 - 3*I)*a^(5/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a^2*((7*I)*A - 9*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a))/(2*a^2)`

3.183. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} dx$

3.183.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`
- rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

3.183.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 739 vs. $2(155) = 310$.

Time = 0.18 (sec) , antiderivative size = 740, normalized size of antiderivative = 3.87

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} \left(60iB\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} (\tan^2(dx+c)) - 36B\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))} \left(60iB\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} (\tan^2(dx+c)) - 36B\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \right)}{\dots}$
parts	$\frac{A\sqrt{a(1+i \tan(dx+c))} \left(3i\sqrt{2} \ln \left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - ia + 3a \tan(dx+c)}{\tan(dx+c)+i} \right) (\tan^4(dx+c)) a + 36\sqrt{-ia} (\tan^2(dx+c)) \right)}{\dots}$

```
input int((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(5/2),x,method=_R
RETURNVERBOSE)
```

```
output 1/12/d*(a*(1+I*tan(d*x+c)))^(1/2)/a/tan(d*x+c)^(3/2)*(60*I*B*(-I*a)^(1/2)*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2-36*B*(-I*a)^(1/2)*(a*ta
n(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3+28*I*A*(-I*a)^(1/2)*(a*tan(d
*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3+3*B*2^(1/2)*ln((2*2^(1/2)*(-I*a
)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x
+c)+I))*a*tan(d*x+c)^4+36*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(
1/2)*tan(d*x+c)^2+3*I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(
1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^4-
6*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(
1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3+6*A*ln((2*2^(1/2)*
(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(ta
n(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^3-3*I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/
2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I
))*a*tan(d*x+c)^2-3*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+
I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2+24
*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+8*A*(-I*a
)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2))/(a*tan(d*x+c)*(1+I*tan(d*x+
c)))^(1/2)/(-tan(d*x+c)+I)^2/(-I*a)^(1/2)
```

$$3.183. \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}} dx$$

3.183.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 534 vs. $2(145) = 290$.

Time = 0.28 (sec) , antiderivative size = 534, normalized size of antiderivative = 2.80

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx =$$

$$3\sqrt{2} \left(ade^{(5i dx + 5i c)} - 2ade^{(3i dx + 3i c)} + ade^{(i dx + i c)} \right) \sqrt{-\frac{iA^2 + 2AB - iB^2}{ad^2}} \log \left(\frac{i\sqrt{2}ad\sqrt{-\frac{iA^2 + 2AB - iB^2}{ad^2}} e^{(i dx + i c)} + \sqrt{a + ia \tan(c + dx)}}{\dots} \right)$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(5/2),x, algorithm="fricas")`

output `-1/12*(3*sqrt(2)*(a*d*e^(5*I*d*x + 5*I*c) - 2*a*d*e^(3*I*d*x + 3*I*c) + a*d*e^(I*d*x + I*c))*sqrt(-(I*A^2 + 2*A*B - I*B^2)/(a*d^2))*log((I*sqrt(2)*a*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)/(a*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - 3*sqrt(2)*(a*d*e^(5*I*d*x + 5*I*c) - 2*a*d*e^(3*I*d*x + 3*I*c) + a*d*e^(I*d*x + I*c))*sqrt(-(I*A^2 + 2*A*B - I*B^2)/(a*d^2))*log((-I*sqrt(2)*a*d*sqrt(-(I*A^2 + 2*A*B - I*B^2)/(a*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) + 2*sqrt(2)*((7*A + 15*I*B)*e^(6*I*d*x + 6*I*c) - (11*A + 3*I*B)*e^(4*I*d*x + 4*I*c) - 15*(A + I*B)*e^(2*I*d*x + 2*I*c) + 3*A + 3*I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(a*d*e^(5*I*d*x + 5*I*c) - 2*a*d*e^(3*I*d*x + 3*I*c) + a*d*e^(I*d*x + I*c))`

3.183.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{ia (\tan(c + dx) - i)} \tan^{\frac{5}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/2)/tan(d*x+c)**(5/2),x)`

output `Integral((A + B*tan(c + d*x))/(sqrt(I*a*(tan(c + d*x) - I))*tan(c + d*x)**(5/2)), x)`

3.183.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.183.8 Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx = \text{Exception raised: NotImplementedError}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(5/2),x, algorithm="giac")`

output `Exception raised: NotImplementedError >> unable to parse Giac output: 2*((9*i*sageVARa*sageVARb*sageVARd+6*sageVARa*sageVARd*sageVARa)/9/sageVARa^3/sageVARd^2*sqrt(i*sageVARa*tan(sageVARc+sageVARd*sageVARx)+sageVARa)*sqrt(i*sageVARa*tan(sage`

3.183.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2} \sqrt{a + a \tan(c + dx)} \operatorname{li}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + a*tan(c + d*x)*1i)^(1/2)),x)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + a*tan(c + d*x)*1i)^(1/2)), x)`

$$3.184 \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{7}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}} dx$$

3.184.1 Optimal result	1902
3.184.2 Mathematica [A] (verified)	1903
3.184.3 Rubi [A] (verified)	1903
3.184.4 Maple [B] (verified)	1907
3.184.5 Fricas [B] (verification not implemented)	1909
3.184.6 Sympy [F]	1910
3.184.7 Maxima [F(-2)]	1910
3.184.8 Giac [F(-2)]	1910
3.184.9 Mupad [F(-1)]	1911

3.184.1 Optimal result

Integrand size = 38, antiderivative size = 237

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}} dx = -\frac{\left(\frac{1}{2} + \frac{i}{2}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ad}} + \frac{A + iB}{d \tan^{\frac{5}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}} - \frac{(7A + 5iB)\sqrt{a + ia \tan(c + dx)}}{5ad \tan^{\frac{5}{2}}(c + dx)} + \frac{(23iA - 25B)\sqrt{a + ia \tan(c + dx)}}{15ad \tan^{\frac{3}{2}}(c + dx)} + \frac{(61A + 35iB)\sqrt{a + ia \tan(c + dx)}}{15ad \sqrt{\tan(c + dx)}}$$

```
output (-1/2-1/2*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)^(1/2))/d/a^(1/2)+1/15*(61*A+35*I*B)*(a+I*a*tan(d*x+c)^(1/2)/a/d/tan(d*x+c)^(1/2)+(A+I*B)/d/(a+I*a*tan(d*x+c)^(1/2)/tan(d*x+c)^(5/2)-1/5*(7*A+5*I*B)*(a+I*a*tan(d*x+c)^(1/2)/a/d/tan(d*x+c)^(5/2)+1/15*(23*I*A-25*B)*(a+I*a*tan(d*x+c)^(1/2)/a/d/tan(d*x+c)^(3/2))
```

3.184.2 Mathematica [A] (verified)

Time = 3.82 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.73

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx$$

$$= \frac{15\sqrt{2}a(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia}\tan(c+dx)}{\sqrt{a+ia}\tan(c+dx)}\right)\tan^4(c+dx)}{(ia\tan(c+dx))^{3/2}} + \frac{2(-6A+2i(A+5iB)\tan(c+dx)+2(19A+5iB)\tan^2(c+dx)+(61iA-35B)\tan^3(c+dx))}{\sqrt{a+ia}\tan(c+dx)}$$

$$= \frac{\dots}{30d \tan^{\frac{5}{2}}(c + dx)}$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(7/2)*Sqrt[a + I*a*Tan[c + d*x]]),x]`

output `((15*Sqrt[2]*a*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Tan[c + d*x]^4)/(I*a*Tan[c + d*x]^(3/2) + (2*(-6*A + (2*I)*(A + (5*I)*B)*Tan[c + d*x] + 2*(19*A + (5*I)*B)*Tan[c + d*x]^2 + ((61*I)*A - 35*B)*Tan[c + d*x]^3))/Sqrt[a + I*a*Tan[c + d*x]])/(30*d*Tan[c + d*x]^(5/2))`

3.184.3 Rubi [A] (verified)

Time = 1.35 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.07, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.395$, Rules used = {3042, 4079, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{7/2} \sqrt{a + ia \tan(c + dx)}} dx$$

↓ 4079

$$\frac{\int \frac{\sqrt{i \tan(c+dx)a+a(a(7A+5iB)-6a(iA-B)\tan(c+dx))} dx}{2 \tan^{\frac{7}{2}}(c+dx)}}{a^2} + \frac{A + iB}{d \tan^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}$$

↓ 27

3.184. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{7}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}} dx$

$$\begin{aligned}
& \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(7A+5iB)-6a(iA-B) \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx}{2a^2} + \frac{A+iB}{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(7A+5iB)-6a(iA-B) \tan(c+dx))}{\tan(c+dx)^{7/2}} dx}{2a^2} + \frac{A+iB}{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 4081 \\
& \frac{2 \int -\frac{\sqrt{i \tan(c+dx)a+a}((23iA-25B)a^2+4(7A+5iB) \tan(c+dx)a^2)}{2 \tan^{\frac{5}{2}}(c+dx)} dx}{5a} - \frac{2a(7A+5iB) \sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 27 \\
& - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((23iA-25B)a^2+4(7A+5iB) \tan(c+dx)a^2)}{\tan^{\frac{5}{2}}(c+dx)} dx}{5a} - \frac{2a(7A+5iB) \sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 3042 \\
& - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((23iA-25B)a^2+4(7A+5iB) \tan(c+dx)a^2)}{\tan(c+dx)^{5/2}} dx}{5a} - \frac{2a(7A+5iB) \sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 4081 \\
& \frac{2 \int \frac{\sqrt{i \tan(c+dx)a+a}(a^3(61A+35iB)-2a^3(23iA-25B) \tan(c+dx))}{2 \tan^{\frac{3}{2}}(c+dx)} dx}{3a} - \frac{2a^2(-25B+23iA) \sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7A+5iB) \sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} \\
& \quad \downarrow 27
\end{aligned}$$

3.184. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{7}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} dx$

$$\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(61A+35iB)-2a^3(23iA-25B) \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx}{5a} - \frac{2a^2(-25B+23iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7A+5iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} +$$

$$\frac{A + iB \frac{2a^2}{d \tan^{\frac{5}{2}}(c+dx)} \sqrt{a+ia \tan(c+dx)}}{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}$$

↓ 3042

$$\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(61A+35iB)-2a^3(23iA-25B) \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx}{5a} - \frac{2a^2(-25B+23iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7A+5iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} +$$

$$\frac{A + iB \frac{2a^2}{d \tan^{\frac{5}{2}}(c+dx)} \sqrt{a+ia \tan(c+dx)}}{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}$$

↓ 4081

$$\frac{2 \int \frac{15a^4(iA+B)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx}{3a} - \frac{2a^3(61A+35iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(-25B+23iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7A+5iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} +$$

$$\frac{A + iB \frac{2a^2}{d \tan^{\frac{5}{2}}(c+dx)} \sqrt{a+ia \tan(c+dx)}}{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}$$

↓ 27

$$\frac{15a^3(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{3a} - \frac{2a^3(61A+35iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(-25B+23iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7A+5iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} +$$

$$\frac{A + iB \frac{2a^2}{d \tan^{\frac{5}{2}}(c+dx)} \sqrt{a+ia \tan(c+dx)}}{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}$$

↓ 3042

$$\frac{15a^3(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{3a} - \frac{2a^3(61A+35iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(-25B+23iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7A+5iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} +$$

$$\frac{A + iB \frac{2a^2}{d \tan^{\frac{5}{2}}(c+dx)} \sqrt{a+ia \tan(c+dx)}}{d \tan^{\frac{5}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}}$$

3.184. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{7}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} dx$

$$\begin{aligned}
 & \downarrow 4027 \\
 & \frac{30ia^5(B+iA) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{d} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{3a} - \frac{2a^3(61A+35iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(-25B+23iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7A+5iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
 & \frac{A+iB}{d \tan^{\frac{5}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}} \\
 & \downarrow 218 \\
 & \frac{(15-15i)a^{7/2}(B+iA) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^3(61A+35iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(-25B+23iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(7A+5iB)\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
 & \frac{A+iB}{d \tan^{\frac{5}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}}
 \end{aligned}$$

```
input Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(7/2)*Sqrt[a + I*a*Tan[c + d*x]]),x]
```

```
output (A + I*B)/(d*Tan[c + d*x]^(5/2)*Sqrt[a + I*a*Tan[c + d*x]]) + ((-2*a*(7*A + (5*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) - ((-2*a^2*((23*I)*A - 25*B)*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) + (((15 - 15*I)*a^(7/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a^3*(61*A + (35*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a))/(5*a))/(2*a^2)
```

3.184.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

$$3.184. \int \frac{A+B \tan(c+dx)}{\tan^{\frac{7}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}} dx$$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

3.184.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 814 vs. $2(193) = 386$.

Time = 0.18 (sec) , antiderivative size = 815, normalized size of antiderivative = 3.44

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} \left(-396iA\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} (\tan^3(dx+c)) + 140iB\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))} \left(-396iA\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} (\tan^3(dx+c)) + 140iB\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \right)}{\dots}$
parts	Expression too large to display

```
input int((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(7/2),x,method=_R
ETURNVERBOSE)
```

```
output 1/60/d*(a*(1+I*tan(d*x+c)))^(1/2)/a/tan(d*x+c)^(5/2)*(-396*I*A*(-I*a)^(1/2)
)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3+140*I*B*(-I*a)^(1/2)*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^4-15*A*2^(1/2)*ln((2*2^(1
/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))
/(tan(d*x+c)+I))*a*tan(d*x+c)^5+244*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(
d*x+c)))^(1/2)*tan(d*x+c)^4-15*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*t
an(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*ta
n(d*x+c)^3+30*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(
d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^4+180*B*(-
I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3+16*I*A*(-I*a
)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+15*A*ln((2*2^(1/2)
)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(
tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^3-144*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+
I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2+30*I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)
)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I)
)*a*tan(d*x+c)^4+15*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(
1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^5+
40*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+24*A*(-
I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2))/(a*tan(d*x+c)*(1+I*tan(d
*x+c)))^(1/2)/(-I*a)^(1/2)/(-tan(d*x+c)+I)^2
```

3.184. $\int \frac{A+B \tan(c+dx)}{\tan^2(c+dx)\sqrt{a+ia \tan(c+dx)}} dx$

3.184.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 596 vs. $2(181) = 362$.

Time = 0.28 (sec) , antiderivative size = 596, normalized size of antiderivative = 2.51

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx =$$

$$15\sqrt{2} \left(ade^{(7i dx + 7i c)} - 3ade^{(5i dx + 5i c)} + 3ade^{(3i dx + 3i c)} - ade^{(i dx + i c)} \right) \sqrt{\frac{-iA^2 - 2AB + iB^2}{ad^2}} \log \left(\frac{\sqrt{2ad} \sqrt{-iA^2 - 2AB + iB^2}}{\dots} \right)$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(7/2),x, algorithm="fricas")`

output

```
-1/60*(15*sqrt(2)*(a*d*e^(7*I*d*x + 7*I*c) - 3*a*d*e^(5*I*d*x + 5*I*c) + 3
*a*d*e^(3*I*d*x + 3*I*c) - a*d*e^(I*d*x + I*c))*sqrt(-(-I*A^2 - 2*A*B + I
*B^2)/(a*d^2))*log((sqrt(2)*a*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2))*e^(
I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e
^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2
*I*c) + 1)))/(4*I*A + 4*B)) - 15*sqrt(2)*(a*d*e^(7*I*d*x + 7*I*c) - 3*a*d*
e^(5*I*d*x + 5*I*c) + 3*a*d*e^(3*I*d*x + 3*I*c) - a*d*e^(I*d*x + I*c))*sqr
t(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2))*log(-(sqrt(2)*a*d*sqrt(-(-I*A^2 - 2*A
*B + I*B^2)/(a*d^2))*e^(I*d*x + I*c) - sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I
*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I
*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) + 2*sqrt(2)*((-103*I*A
+ 35*B)*e^(8*I*d*x + 8*I*c) + 6*(17*I*A - 15*B)*e^(6*I*d*x + 6*I*c) + 20*
(2*I*A - B)*e^(4*I*d*x + 4*I*c) + 30*(-5*I*A + 3*B)*e^(2*I*d*x + 2*I*c) +
15*I*A - 15*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I
*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(a*d*e^(7*I*d*x + 7*I*c) - 3*a*d*e^(5
*I*d*x + 5*I*c) + 3*a*d*e^(3*I*d*x + 3*I*c) - a*d*e^(I*d*x + I*c))
```

3.184.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{ia (\tan(c + dx) - i)} \tan^{\frac{7}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/2)/tan(d*x+c)**(7/2),x)`

output `Integral((A + B*tan(c + d*x))/(sqrt(I*a*(tan(c + d*x) - I))*tan(c + d*x)**(7/2)), x)`

3.184.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(7/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.184.8 Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx = \text{Exception raised: NotImplementedError}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(7/2),x, algorithm="giac")`

output `Exception raised: NotImplementedError >> unable to parse Giac output: 2*((-345*i*sageVARa^3*sageVARA*sageVARd^2-150*sageVARa^3*sageVARd^2*sageVARB)/(225*i)/sageVARa^6/sageVARd^3*sqrt(i*sageVARa*tan(sageVARc+sageVARd*sageVARx)+sageVARa)*sqrt`

3.184.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{7/2} \sqrt{a + a \tan(c + dx)} \operatorname{li}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(7/2)*(a + a*tan(c + d*x)*1i)^(1/2)),x)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(7/2)*(a + a*tan(c + d*x)*1i)^(1/2)), x)`

3.185
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$$

3.185.1 Optimal result 1912
 3.185.2 Mathematica [A] (verified) 1913
 3.185.3 Rubi [A] (verified) 1913
 3.185.4 Maple [B] (verified) 1917
 3.185.5 Fricas [B] (verification not implemented) 1918
 3.185.6 Sympy [F] 1919
 3.185.7 Maxima [F(-2)] 1920
 3.185.8 Giac [F(-2)] 1920
 3.185.9 Mupad [F(-1)] 1920

3.185.1 Optimal result

Integrand size = 38, antiderivative size = 203

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{2(-1)^{3/4} B \arctan\left(\frac{(-1)^{3/4} \sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{a^{3/2} d}$$

$$- \frac{\left(\frac{1}{4} - \frac{i}{4}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{a^{3/2} d}$$

$$+ \frac{(iA - B) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} + \frac{(A+3iB)\sqrt{\tan(c+dx)}}{2ad\sqrt{a+ia \tan(c+dx)}}$$

```
output 2*(-1)^(3/4)*B*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)
)^(1/2))/a^(3/2)/d+(-1/4+1/4*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(
1/2)/(a+I*a*tan(d*x+c))^(1/2))/a^(3/2)/d+1/2*(A+3*I*B)*tan(d*x+c)^(1/2)/a
/d/(a+I*a*tan(d*x+c))^(1/2)+1/3*(I*A-B)*tan(d*x+c)^(3/2)/d/(a+I*a*tan(d*x+
c))^(3/2)
```

3.185.2 Mathematica [A] (verified)

Time = 4.58 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.88

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{3}{2}}} dx = \frac{\left(\frac{1}{4} + \frac{i}{4}\right)(iA+B) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{a^{\frac{3}{2}}d} + \frac{\sqrt{a+ia \tan(c+dx)}\left(12\sqrt[4]{-1} \operatorname{Barcsinh}\left(\sqrt[4]{-1}\sqrt{\tan(c+dx)}\right)\right)(1+i \tan(c+dx))^{\frac{3}{2}} + \sqrt{\tan(c+dx)}(-3(A+B \tan(c+dx)))}{6a^2d(-i + \tan(c+dx))^2}$$

input `Integrate[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `((1/4 + I/4)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/(a^(3/2)*d) + (Sqrt[a + I*a*Tan[c + d*x]]*(12*(-1)^(1/4)*B*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]]*(1 + I*Tan[c + d*x])^(3/2) + Sqrt[Tan[c + d*x]]*(-3*(A + (3*I)*B) + ((-5*I)*A + 11*B)*Tan[c + d*x]))/(6*a^2*d*(-I + Tan[c + d*x])^2)`

3.185.3 Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4078, 27, 3042, 4078, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{3}{2}}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(c+dx)^{\frac{3}{2}}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{3}{2}}} dx \\ & \quad \downarrow \text{4078} \\ & \frac{(-B+iA) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{\frac{3}{2}}} - \frac{\int \frac{3\sqrt{\tan(c+dx)}(a(iA-B)+2iaB \tan(c+dx))}{2\sqrt{i \tan(c+dx)a+a}} dx}{3a^2} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.185. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{3}{2}}} dx$

$$\begin{aligned}
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\int \frac{\sqrt{\tan(c+dx)}(a(iA-B)+2iaB \tan(c+dx))}{\sqrt{i \tan(c+dx)a+a}} dx}{2a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\int \frac{\sqrt{\tan(c+dx)}(a(iA-B)+2iaB \tan(c+dx))}{\sqrt{i \tan(c+dx)a+a}} dx}{2a^2} \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\int -\frac{\sqrt{i \tan(c+dx)a+a}((A+3iB)a^2+4B \tan(c+dx)a^2)}{2\sqrt{\tan(c+dx)}} dx}{2a^2} - \frac{a(A+3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((A+3iB)a^2+4B \tan(c+dx)a^2)}{\sqrt{\tan(c+dx)}} dx}{2a^2} - \frac{a(A+3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((A+3iB)a^2+4B \tan(c+dx)a^2)}{\sqrt{\tan(c+dx)}} dx}{2a^2} - \frac{a(A+3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \quad \downarrow \text{4084} \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{a^2(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + 4iaB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a^2} - \frac{a(A+3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{a^2(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + 4iaB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a^2} - \frac{a(A+3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \\
 & \quad \downarrow \text{4027}
 \end{aligned}$$

3.185. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \\
 & \frac{4iaB \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + a}}{\sqrt{\tan(c + dx)}} dx - \frac{2ia^4(A - iB) \int \frac{1}{-i \tan(c + dx) a + a} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{i \tan(c + dx) a + a}}}{2a^2}}{2a^2} - \frac{a(A + 3iB) \sqrt{\tan(c + dx)}}{d \sqrt{a + ia \tan(c + dx)}} \\
 & \qquad \qquad \qquad \downarrow 218 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \\
 & \frac{4iaB \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + a}}{\sqrt{\tan(c + dx)}} dx + \frac{(1 - i)a^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{(1 + i)\sqrt{a} \sqrt{\tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right)}{d}}{2a^2} - \frac{a(A + 3iB) \sqrt{\tan(c + dx)}}{d \sqrt{a + ia \tan(c + dx)}} \\
 & \qquad \qquad \qquad \downarrow 4082 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \\
 & \frac{4ia^3B \int \frac{1}{\sqrt{\tan(c + dx)} \sqrt{i \tan(c + dx) a + a}} d \tan(c + dx) + \frac{(1 - i)a^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{(1 + i)\sqrt{a} \sqrt{\tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right)}{d}}{2a^2} - \frac{a(A + 3iB) \sqrt{\tan(c + dx)}}{d \sqrt{a + ia \tan(c + dx)}} \\
 & \qquad \qquad \qquad \downarrow 65 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \\
 & \frac{8ia^3B \int \frac{1}{1 - \frac{ia \tan(c + dx)}{i \tan(c + dx) a + a}} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{i \tan(c + dx) a + a}} + \frac{(1 - i)a^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{(1 + i)\sqrt{a} \sqrt{\tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right)}{d}}{2a^2} - \frac{a(A + 3iB) \sqrt{\tan(c + dx)}}{d \sqrt{a + ia \tan(c + dx)}} \\
 & \qquad \qquad \qquad \downarrow 216 \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \\
 & \frac{(1 - i)a^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{(1 + i)\sqrt{a} \sqrt{\tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right) - 8(-1)^{3/4} a^{5/2} B \operatorname{arctan}\left(\frac{(-1)^{3/4} \sqrt{a} \sqrt{\tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right)}{2a^2} - \frac{a(A + 3iB) \sqrt{\tan(c + dx)}}{d \sqrt{a + ia \tan(c + dx)}} \\
 & \qquad \qquad \qquad \downarrow \\
 & \frac{(-B + iA) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{a(A + 3iB) \sqrt{\tan(c + dx)}}{d \sqrt{a + ia \tan(c + dx)}}
 \end{aligned}$$

```
input Int[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]
```

3.185. $\int \frac{\tan^{\frac{3}{2}}(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{3/2}} dx$

output $((I*A - B)*\text{Tan}[c + d*x]^{(3/2)})/(3*d*(a + I*a*\text{Tan}[c + d*x]^{(3/2)}) - (((-8*(-1)^{(3/4)}*a^{(5/2)}*B*\text{ArcTan}[((-1)^{(3/4)}*\text{Sqrt}[a]*\text{Sqrt}[\text{Tan}[c + d*x]])]/\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]))/d + ((1 - I)*a^{(5/2)}*(A - I*B)*\text{ArcTanh}(((1 + I)*\text{Sqrt}[a]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]))/d)/(2*a^2) - (a*(A + (3*I)*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/(d*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])/(2*a^2)$

3.185.3.1 Defintions of rubi rules used

rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_) /; \text{FreeQ}[b, x]]$

rule 65 $\text{Int}[1/(\text{Sqrt}[(b_*)(x_)]*\text{Sqrt}[(c_) + (d_*)(x_)]), x_Symbol] \rightarrow \text{Simp}[2 \text{ Subst}[\text{Int}[1/(b - d*x^2), x], x, \text{Sqrt}[b*x]/\text{Sqrt}[c + d*x]], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ !\text{GtQ}[c, 0]$

rule 216 $\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 218 $\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4027 $\text{Int}[\text{Sqrt}[(a_*) + (b_*)*\text{tan}[(e_*) + (f_*)(x_)]]/\text{Sqrt}[(c_*) + (d_*)*\text{tan}[(e_*) + (f_*)(x_)]], x_Symbol] \rightarrow \text{Simp}[-2*a*(b/f) \text{ Subst}[\text{Int}[1/(a*c - b*d - 2*a^2*x^2), x], x, \text{Sqrt}[c + d*\text{Tan}[e + f*x]]/\text{Sqrt}[a + b*\text{Tan}[e + f*x]]], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0]$

```
rule 4078 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

```
rule 4082 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 + b^2, 0] && EqQ[A*b + a*B, 0]
```

```
rule 4084 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x]
- Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[
e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

3.185.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1214 vs. $2(160) = 320$.

Time = 0.19 (sec) , antiderivative size = 1215, normalized size of antiderivative = 5.99

method	result	size
derivativedivides	Expression too large to display	1215
default	Expression too large to display	1215
parts	Expression too large to display	1229

```
input int(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x,method=_R
ETURNVERBOSE)
```

$$3.185. \quad \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{3}{2}}} dx$$

```
output 1/24*I/d*tan(d*x+c)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)/a^2*(24*I*B*ln(1/2*(2
*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*
a)^(1/2))*(-I*a)^(1/2)*a+20*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan
(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-3*I*A*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-
I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(
d*x+c)+I))*a-3*A*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x
+c)*(1+I*tan(d*x+c))))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+
c)^3-9*I*B*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1
+I*tan(d*x+c))))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)+9*I
*A*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d
*x+c))))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2-36*I*B*(I
*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+9*B*2^(1/2)*l
n((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)-I*a+3*a*ta
n(d*x+c))/(tan(d*x+c)+I))*(I*a)^(1/2)*a*tan(d*x+c)^2+24*B*ln(1/2*(2*I*a*ta
n(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2
))*(-I*a)^(1/2)*a*tan(d*x+c)^3+44*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+
c)*(1+I*tan(d*x+c))))^(1/2)*tan(d*x+c)^2-32*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*(a
*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*tan(d*x+c)+9*A*(I*a)^(1/2)*2^(1/2)*ln(
(2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)-I*a+3*a*ta
n(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)-72*I*B*ln(1/2*(2*I*a*tan(d*x+c)+2*...
```

3.185.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 758 vs. 2(149) = 298.

Time = 0.35 (sec) , antiderivative size = 758, normalized size of antiderivative = 3.73

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx =$$

$$\left(3 \sqrt{\frac{1}{2}} a^2 d \sqrt{\frac{-i A^2 - 2 AB + i B^2}{a^3 d^2}} e^{(3i dx + 3i c)} \log \left(\frac{2i \sqrt{\frac{1}{2}} a^2 d \sqrt{\frac{-i A^2 - 2 AB + i B^2}{a^3 d^2}} e^{(i dx + i c)} + \sqrt{2} ((i A + B) e^{(2i dx + 2i c)} + i A + B) \sqrt{e^{(2i dx + 2i c)}}}{4i A + 4 B} \right) \right)$$

```
input integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, al
gorithm="fricas")
```

3.185. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

output

```

-1/12*(3*sqrt(1/2)*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2))*e^(3*I*d
*x + 3*I*c)*log((2*I*sqrt(1/2)*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^
2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sq
rt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*
d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - 3*sqrt(1/2)*a^2*d*sqrt((-I*A^2 - 2*A*
B + I*B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log((-2*I*sqrt(1/2)*a^2*d*sqrt((
-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^
(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^
(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) + 3*a^2*
d*sqrt(4*I*B^2/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(52/605*(4*sqrt(2)*(B*e^(
3*I*d*x + 3*I*c) + B*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sq
rt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) - (3*I*a^2*d*e^
(2*I*d*x + 2*I*c) - I*a^2*d)*sqrt(4*I*B^2/(a^3*d^2)))/(B*e^(2*I*d*x + 2*I*
c) + B)) - 3*a^2*d*sqrt(4*I*B^2/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(52/605*
(4*sqrt(2)*(B*e^(3*I*d*x + 3*I*c) + B*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x
+ 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)
) - (-3*I*a^2*d*e^(2*I*d*x + 2*I*c) + I*a^2*d)*sqrt(4*I*B^2/(a^3*d^2)))/(B
*e^(2*I*d*x + 2*I*c) + B)) - sqrt(2)*(2*(2*A + 5*I*B)*e^(4*I*d*x + 4*I*c)
+ 3*(A + 3*I*B)*e^(2*I*d*x + 2*I*c) - A - I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c)
+ 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))e^...

```

3.185.6 Sympy [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{3}{2}}} dx = \int \frac{(A+B \tan(c+dx)) \tan^{\frac{3}{2}}(c+dx)}{(ia(\tan(c+dx)-i))^{\frac{3}{2}}} dx$$

input `integrate(tan(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**(3/2)/(I*a*(tan(c + d*x) - I))
**(3/2), x)`

3.185.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.185.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[18]Warning, replacing 18 by 97, a substitution variable`

3.185.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \int \frac{\tan(c+dx)^{3/2}(A+B\tan(c+dx))}{(a+a\tan(c+dx)li)^{3/2}} dx$$

input `int((tan(c+d*x)^(3/2)*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*li)^(3/2),x)`

3.185. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx$

output `int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(3/2), x)`

3.185. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{3}{2}}} dx$

3.186
$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$$

3.186.1 Optimal result	1922
3.186.2 Mathematica [A] (verified)	1922
3.186.3 Rubi [A] (verified)	1923
3.186.4 Maple [B] (verified)	1925
3.186.5 Fricas [B] (verification not implemented)	1927
3.186.6 Sympy [F]	1927
3.186.7 Maxima [F(-2)]	1928
3.186.8 Giac [F(-2)]	1928
3.186.9 Mupad [F(-1)]	1928

3.186.1 Optimal result

Integrand size = 38, antiderivative size = 150

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx =$$

$$\frac{\left(\frac{1}{4} + \frac{i}{4}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{a^{3/2}d}$$

$$+ \frac{(iA - B)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} + \frac{(iA + 5B)\sqrt{\tan(c+dx)}}{6ad\sqrt{a+ia \tan(c+dx)}}$$

output `(-1/4-1/4*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/a^(3/2)/d+1/6*(I*A+5*B)*tan(d*x+c)^(1/2)/a/d/(a+I*a*tan(d*x+c))^(1/2)+1/3*(I*A-B)*tan(d*x+c)^(1/2)/d/(a+I*a*tan(d*x+c))^(3/2)`

3.186.2 Mathematica [A] (verified)

Time = 2.91 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.99

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{\sqrt{\tan(c+dx)} \left(-\frac{3i\sqrt{2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ia \tan(c+dx)}} + \frac{6(A-iB)+2}{(-i+\tan(c+dx))} \right)}{12ad}$$

input `Integrate[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])
^(3/2),x]`

output `(Sqrt[Tan[c + d*x]]*(((−3*I)*Sqrt[2]*(A − I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*T
an[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/Sqrt[I*a*Tan[c + d*x]] + (6*(A
− I*B) + 2*(I*A + 5*B)*Tan[c + d*x])/((−I + Tan[c + d*x])*Sqrt[a + I*a*Tan
[c + d*x]])))/(12*a*d)`

3.186.3 Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.237$, Rules used = {3042, 4078, 27, 3042, 4079, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+iA)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{a(iA-B)-2a(A-2iB)\tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} dx}{3a^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B+iA)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{a(iA-B)-2a(A-2iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} dx}{6a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B+iA)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{a(iA-B)-2a(A-2iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} dx}{6a^2} \\
 & \quad \downarrow \text{4079} \\
 & \frac{(-B+iA)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{3a^2(iA+B)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx}{a^2} - \frac{a(5B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}
 \end{aligned}$$

3.186. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\frac{3}{2}(B + iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{a(5B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} \\
& \downarrow 3042 \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\frac{3}{2}(B + iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{a(5B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} \\
& \downarrow 4027 \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{3ia^2(B+iA) \int \frac{1}{\frac{-2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{6a^2} - \frac{a(5B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \\
& \downarrow 218 \\
& \frac{(-B + iA)\sqrt{\tan(c + dx)}}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\left(\frac{3}{2} - \frac{3i}{2}\right)\sqrt{a}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{a(5B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}
\end{aligned}$$

input `Int[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `((I*A - B)*Sqrt[Tan[c + d*x]]/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) - (((3/2 - (3*I)/2)*Sqrt[a]*(I*A + B)*ArcTanh[(((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/d - (a*(I*A + 5*B)*Sqrt[Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]])))/(6*a^2)`

3.186.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-(A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.186.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 859 vs. $2(120) = 240$.

Time = 0.16 (sec) , antiderivative size = 860, normalized size of antiderivative = 5.73

method	result
derivativedivides	$\frac{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})\left(-3iA\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)+4A\sqrt{-\dots}\right)}{\dots}$
default	$\frac{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))})\left(-3iA\sqrt{2}\ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i}\right)+4A\sqrt{-\dots}\right)}{\dots}$
parts	Expression too large to display

input `int(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x,method=_RETURVERBOSE)`

output
$$\begin{aligned} & -1/24/d*\tan(d*x+c)^{(1/2)}*(a*(1+I*\tan(d*x+c)))^{(1/2)}/a^2*(-3*I*A*2^{(1/2)}*\ln \\ & ((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan \\ & (d*x+c))/(tan(d*x+c)+I))*a+4*A*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)) \\ &)^{(1/2)}*\tan(d*x+c)^2+3*I*B*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c) \\ &)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(tan(d*x+c)+I))*a*\tan(d*x+c) \\ & ^3-3*A*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I* \\ & a+3*a*\tan(d*x+c))/(tan(d*x+c)+I))*2^{(1/2)}*a*\tan(d*x+c)^3+12*I*B*(-I*a)^{(1/ \\ & 2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-20*I*B*(-I*a)^{(1/2)}*(a*\tan(d*x+c) \\ & *(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)^2+9*B*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/ \\ & 2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(tan(d*x+c)+I \\ &))*a*\tan(d*x+c)^2+9*I*A*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(\\ & 1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(tan(d*x+c)+I))*a*\tan(d*x+c)^2- \\ & 16*I*A*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)+9*A*2 \\ & ^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I* \\ & a+3*a*\tan(d*x+c))/(tan(d*x+c)+I))*a*\tan(d*x+c)-9*I*B*2^{(1/2)}*\ln((2*2^{(1/2)} \\ & *(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(t \\ & an(d*x+c)+I))*a*\tan(d*x+c)-3*B*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d \\ & *x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(tan(d*x+c)+I))*a-32*B*(\\ & -I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)-12*A*(-I*a)^{(\\ & 1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)})/(a*\tan(d*x+c)*(1+I*\tan(d*x+... \end{aligned}$$

3.186.
$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx$$

3.186.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 443 vs. $2(110) = 220$.

Time = 0.28 (sec) , antiderivative size = 443, normalized size of antiderivative = 2.95

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx =$$

$$\left(3\sqrt{\frac{1}{2}}a^2d\sqrt{\frac{iA^2+2AB-iB^2}{a^3d^2}}e^{(3i dx+3i c)} \log\left(\frac{2\sqrt{\frac{1}{2}}a^2d\sqrt{\frac{iA^2+2AB-iB^2}{a^3d^2}}e^{(i dx+i c)}+\sqrt{2}((iA+B)e^{(2i dx+2i c)}+iA+B)}{4iA+4B}\sqrt{\frac{a}{e^{(2i dx+2i c)}}}}\right) \right)$$

```
input integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="fricas")
```

```
output -1/12*(3*sqrt(1/2)*a^2*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log((2*sqrt(1/2)*a^2*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^3*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - 3*sqrt(1/2)*a^2*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(-(2*sqrt(1/2)*a^2*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^3*d^2))*e^(I*d*x + I*c) - sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) + sqrt(2)*(2*(-I*A - 2*B)*e^(4*I*d*x + 4*I*c) + 3*(-I*A - B)*e^(2*I*d*x + 2*I*c) - I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*e^(-3*I*d*x - 3*I*c)/(a^2*d)
```

3.186.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \int \frac{(A+B\tan(c+dx))\sqrt{\tan(c+dx)}}{(ia(\tan(c+dx)-i))^{3/2}} dx$$

```
input integrate(tan(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(3/2),x)
```

```
output Integral((A + B*tan(c + d*x))*sqrt(tan(c + d*x))/(I*a*(tan(c + d*x) - I))* (3/2), x)
```

3.186. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx$

3.186.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.186.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[-63]Warning, replacing -63 by -42, a substitution vari`

3.186.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+a\tan(c+dx)1i)^{3/2}} dx$$

input `int((tan(c+d*x)^(1/2)*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*1i)^(3/2),x)`

3.186. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx$

output `int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(3/2), x)`

3.186. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

$$3.187 \quad \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}} dx$$

3.187.1 Optimal result	1930
3.187.2 Mathematica [A] (verified)	1930
3.187.3 Rubi [A] (verified)	1931
3.187.4 Maple [B] (verified)	1933
3.187.5 Fricas [B] (verification not implemented)	1934
3.187.6 Sympy [F]	1935
3.187.7 Maxima [F(-2)]	1935
3.187.8 Giac [F(-1)]	1936
3.187.9 Mupad [F(-1)]	1936

3.187.1 Optimal result

Integrand size = 38, antiderivative size = 148

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx = \frac{\left(\frac{1}{4} - \frac{i}{4}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{a^{3/2}d} + \frac{(A + iB)\sqrt{\tan(c + dx)}}{3d(a + ia \tan(c + dx))^{3/2}} + \frac{(7A + iB)\sqrt{\tan(c + dx)}}{6ad\sqrt{a + ia \tan(c + dx)}}$$

output `(1/4-1/4*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/a^(3/2)/d+1/6*(7*A+I*B)*tan(d*x+c)^(1/2)/a/d/(a+I*a*tan(d*x+c))^(1/2)+1/3*(A+I*B)*tan(d*x+c)^(1/2)/d/(a+I*a*tan(d*x+c))^(3/2)`

3.187.2 Mathematica [A] (verified)

Time = 2.00 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.03

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx = \frac{\sqrt{\tan(c + dx)} \left(\frac{3\sqrt{2}a^2(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ia \tan(c+dx)}} + \frac{4a^3(A-iB)}{(a+ia \tan(c+dx))^{3/2}} \right)}{12a^3d}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(3/2)),x]`

3.187. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}} dx$

output $(\text{Sqrt}[\text{Tan}[c + d*x]] * ((3 * \text{Sqrt}[2] * a^2 * (A - I*B) * \text{ArcTanh}[(\text{Sqrt}[2] * \text{Sqrt}[I*a*\text{Tan}[c + d*x]])] / \text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]) / \text{Sqrt}[I*a*\text{Tan}[c + d*x]] + (4*a^3 * (A + I*B)) / (a + I*a*\text{Tan}[c + d*x])^{3/2} + (2*a^2 * (7*A + I*B)) / \text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]) / (12*a^3*d)$

3.187.3 Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.237$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{a(5A - iB) - 2a(iA - B) \tan(c + dx)}{2\sqrt{\tan(c + dx)}\sqrt{i \tan(c + dx)a + a}} dx}{3a^2} + \frac{(A + iB)\sqrt{\tan(c + dx)}}{3d(a + ia \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{a(5A - iB) - 2a(iA - B) \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{i \tan(c + dx)a + a}} dx}{6a^2} + \frac{(A + iB)\sqrt{\tan(c + dx)}}{3d(a + ia \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a(5A - iB) - 2a(iA - B) \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{i \tan(c + dx)a + a}} dx}{6a^2} + \frac{(A + iB)\sqrt{\tan(c + dx)}}{3d(a + ia \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{3a^2(A - iB)\sqrt{i \tan(c + dx)a + a}}{2\sqrt{\tan(c + dx)}} dx}{a^2} + \frac{a(7A + iB)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} + \frac{(A + iB)\sqrt{\tan(c + dx)}}{3d(a + ia \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.187. $\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx$

$$\begin{aligned} & \frac{\frac{3}{2}(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + \frac{a(7A+iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} \\ & \quad \downarrow \text{3042} \\ & \frac{\frac{3}{2}(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + \frac{a(7A+iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} \\ & \quad \downarrow \text{4027} \\ & \frac{\frac{a(7A+iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} - \frac{3ia^2(A-iB) \int \frac{1}{\frac{-2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{6a^2}}{6a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} \\ & \quad \downarrow \text{218} \\ & \frac{\left(\frac{3}{2}-\frac{3i}{2}\right)\sqrt{a}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{\frac{a(7A+iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} \end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(3/2)),x]`

output `((A + I*B)*Sqrt[Tan[c + d*x]])/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) + (((3/2 - (3*I)/2)*Sqrt[a]*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d + (a*(7*A + I*B)*Sqrt[Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]])/(6*a^2)`

3.187.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.187. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}} dx$

```
rule 4027 Int[Sqrt[(a_) + (b_.)*tan[(e_) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*tan[(e_)
+ (f_.)*(x_)]], x_Symbol] :> Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4079 Int[((a_) + (b_.)*tan[(e_) + (f_.)*(x_)])^(m_)*((A_) + (B_.)*tan[(e_) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_) + (f_.)*(x_)])^(n_), x_Symbol] :> Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.187.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 859 vs. $2(118) = 236$.

Time = 0.17 (sec) , antiderivative size = 860, normalized size of antiderivative = 5.81

method	result
derivativedivides	$\frac{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))}) \left(3iA\sqrt{2} \ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i} \right) \right) a(\tan^3(dx+c))}{\dots}$
default	$\frac{(\sqrt{\tan(dx+c)}\sqrt{a(1+i\tan(dx+c))}) \left(3iA\sqrt{2} \ln\left(\frac{2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))-ia+3a\tan(dx+c)}}{\tan(dx+c)+i} \right) \right) a(\tan^3(dx+c))}{\dots}$
parts	Expression too large to display

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(3/2),x,method=_R
ETURNVERBOSE)
```

3.187.
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}} dx$$

output $1/24/d*\tan(d*x+c)^{(1/2)}*(a*(1+I*\tan(d*x+c)))^{(1/2)}*(3*I*A*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c)))/(\tan(d*x+c)+I))*a*\tan(d*x+c)^3-9*I*B*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c)))/(\tan(d*x+c)+I))*2^{(1/2)}*a*\tan(d*x+c)^2+3*B*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c)))/(\tan(d*x+c)+I))*a*\tan(d*x+c)^3-9*I*A*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c)))/(\tan(d*x+c)+I))*2^{(1/2)}*a*\tan(d*x+c)+28*I*A*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)^2+9*A*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c)))/(\tan(d*x+c)+I))*a*\tan(d*x+c)^2+3*I*B*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c)))/(\tan(d*x+c)+I))*2^{(1/2)}*a+16*I*B*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)-9*B*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c)))/(\tan(d*x+c)+I))*a*\tan(d*x+c)-4*B*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)^2-36*I*A*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-3*A*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c)))/(\tan(d*x+c)+I))*2^{(1/2)}*a+64*A*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)+12*B*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)})/a^2/(a*\tan(d*x+c)*(1+I*\tan(d*x+c))...$

3.187.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 441 vs. $2(110) = 220$.

Time = 0.27 (sec) , antiderivative size = 441, normalized size of antiderivative = 2.98

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx = \left(3 \sqrt{\frac{1}{2} a^2 d \sqrt{\frac{-i A^2 - 2 AB + i B^2}{a^3 d^2}}} e^{(3i dx + 3i c)} \log \left(\frac{2i \sqrt{\frac{1}{2} a^2 d \sqrt{\frac{-i A^2 - 2 AB + i B^2}{a^3 d^2}}}}{\dots} \right) \right)$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="fricas")`

output `1/12*(3*sqrt(1/2)*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log((2*I*sqrt(1/2)*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2)))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B) - 3*sqrt(1/2)*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log((-2*I*sqrt(1/2)*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B) + sqrt(2)*(2*(4*A + I*B)*e^(4*I*d*x + 4*I*c) + 3*(3*A + I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))*e^(-3*I*d*x - 3*I*c)/(a^2*d)`

3.187.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{(ia (\tan(c + dx) - i))^{3/2} \sqrt{\tan(c + dx)}} dx$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+I*a*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))/((I*a*(tan(c + d*x) - I))**(3/2)*sqrt(tan(c + d*x))), x)`

3.187.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.187.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `Timed out`

3.187.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + a \tan(c + dx) 1i)^{3/2}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(3/2)),x)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(3/2)), x)`

$$3.188 \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$$

3.188.1 Optimal result	1937
3.188.2 Mathematica [A] (verified)	1937
3.188.3 Rubi [A] (verified)	1938
3.188.4 Maple [B] (verified)	1941
3.188.5 Fricas [B] (verification not implemented)	1942
3.188.6 Sympy [F]	1943
3.188.7 Maxima [F(-2)]	1943
3.188.8 Giac [F(-1)]	1944
3.188.9 Mupad [F(-1)]	1944

3.188.1 Optimal result

Integrand size = 38, antiderivative size = 194

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \frac{\left(\frac{1}{4} + \frac{i}{4}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{a^{3/2}d}$$

$$+ \frac{A + iB}{3d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}}$$

$$+ \frac{11A + 5iB}{6ad\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}} - \frac{(25A + 7iB)\sqrt{a + ia \tan(c + dx)}}{6a^2d\sqrt{\tan(c + dx)}}$$

output

```
(1/4+1/4*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/a^(3/2)/d+1/6*(11*A+5*I*B)/a/d/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2)-1/6*(25*A+7*I*B)*(a+I*a*tan(d*x+c))^(1/2)/a^2/d/tan(d*x+c)^(1/2)+1/3*(A+I*B)/d/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(3/2)
```

3.188.2 Mathematica [A] (verified)

Time = 2.67 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.86

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \frac{3\sqrt{2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \tan(c+dx)}{\sqrt{ia \tan(c+dx)}} - \frac{2i(-12A+(-39iA+9B) \tan(c+dx))}{(-i+\tan(c+dx))^{3/2}} \frac{1}{12ad\sqrt{\tan(c + dx)}}$$

3.188. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(3/2)),x]`

output `((3*Sqrt[2]*(I*A + B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Tan[c + d*x])/Sqrt[I*a*Tan[c + d*x]] - ((2*I)*(-12*A + ((-39*I)*A + 9*B)*Tan[c + d*x] + (25*A + (7*I)*B)*Tan[c + d*x]^2))/((-I + Tan[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]])/(12*a*d*Sqrt[Tan[c + d*x]])`

3.188.3 Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.03, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2}(a + ia \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{a(7A+IB)-4a(iA-B) \tan(c+dx)}{2 \tan^{\frac{3}{2}}(c+dx) \sqrt{i \tan(c+dx)a+a}} dx}{3a^2} + \frac{A + iB}{3d \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{a(7A+IB)-4a(iA-B) \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx) \sqrt{i \tan(c+dx)a+a}} dx}{6a^2} + \frac{A + iB}{3d \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a(7A+IB)-4a(iA-B) \tan(c+dx)}{\tan(c+dx)^{3/2} \sqrt{i \tan(c+dx)a+a}} dx}{6a^2} + \frac{A + iB}{3d \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{4079}
 \end{aligned}$$

3.188. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{\int \frac{\sqrt{i \tan(c+dx)a+a} \left(a^2(25A+7iB) - 2a^2(11iA-5B) \tan(c+dx) \right) dx}{2 \tan^{\frac{3}{2}}(c+dx) a^2} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}}{\frac{6a^2}{A+iB}} + \\
 & \frac{3d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}}{} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\sqrt{i \tan(c+dx)a+a} \left(a^2(25A+7iB) - 2a^2(11iA-5B) \tan(c+dx) \right) dx}{\tan^{\frac{3}{2}}(c+dx) 2a^2} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}}{\frac{6a^2}{A+iB}} + \\
 & \frac{3d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}}{} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\sqrt{i \tan(c+dx)a+a} \left(a^2(25A+7iB) - 2a^2(11iA-5B) \tan(c+dx) \right) dx}{\tan(c+dx)^{3/2} 2a^2} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}}{\frac{6a^2}{A+iB}} + \\
 & \frac{3d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}}{} \\
 & \quad \downarrow \text{4081} \\
 & \frac{2 \int \frac{3a^3(iA+B)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)} a} dx - \frac{2a^2(25A+7iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}}{\frac{6a^2}{A+iB}} + \\
 & \frac{3d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}}{} \\
 & \quad \downarrow \text{27} \\
 & \frac{3a^2(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(25A+7iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}}{\frac{6a^2}{A+iB}} + \\
 & \frac{3d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}}{} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3a^2(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(25A+7iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}}{\frac{6a^2}{A+iB}} + \\
 & \frac{3d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}}{} \\
 & \quad \downarrow \text{4027}
 \end{aligned}$$

3.188. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$

$$\frac{6ia^4(B+iA) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{2a^2} - \frac{2a^2(25A+7iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} + \frac{6a^2(A+iB)}{3d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}}$$

↓ 218

$$\frac{(3-3i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^2(25A+7iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} + \frac{6a^2(A+iB)}{3d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(3/2)),x]`

output `(A + I*B)/(3*d*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(3/2)) + ((a*(11*A + (5*I)*B))/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + (((3 - 3*I)*a^(5/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a^2*(25*A + (7*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(2*a^2))/(6*a^2)`

3.188.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4027 Int[Sqrt[(a_) + (b_.)*tan[(e_) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*tan[(e_)
+ (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4079 Int[((a_) + (b_.)*tan[(e_) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

```
rule 4081 Int[((a_) + (b_.)*tan[(e_) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

3.188.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 922 vs. $2(156) = 312$.

Time = 0.14 (sec) , antiderivative size = 923, normalized size of antiderivative = 4.76

method	result	size
derivativedivides	Expression too large to display	923
default	Expression too large to display	923
parts	Expression too large to display	967

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(3/2),x,method=_R
ETURNVERBOSE)
```

$$3.188. \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$$

```

output 1/24/d*(a*(1+I*tan(d*x+c)))^(1/2)*(3*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)
)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I)
)*a*tan(d*x+c)^4+9*I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1
+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3-3
*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^4-9*I*B*ln((2*2^(1/2)*(-
-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan
(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^2+28*I*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*
tan(d*x+c)))^(1/2)*tan(d*x+c)^3+9*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a
tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*t
an(d*x+c)^3-3*I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*ta
n(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)-256*I*A*
(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2+9*A*2^(1/2)
)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a
*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2+100*A*(-I*a)^(1/2)*(a*tan(d*x+
c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3-36*I*B*(-I*a)^(1/2)*(a*tan(d*x+c)*
(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)-3*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*
a*tan(d*x+c)+64*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d
*x+c)^2+48*I*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-204*A...

```

3.188.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 511 vs. $2(146) = 292$.

Time = 0.27 (sec) , antiderivative size = 511, normalized size of antiderivative = 2.63

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \frac{3 \sqrt{\frac{1}{2}} (a^2 de^{(5i dx + 5i c)} - a^2 de^{(3i dx + 3i c)}) \sqrt{\frac{i A^2 + 2 AB - i B^2}{a^3 d^2}} \log \left(\frac{2 \sqrt{\frac{1}{2}}}{\dots} \right)}{\dots}$$

```

input integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(3/2),x, al
gorithm="fricas")

```

output $1/12*(3*\sqrt{1/2}*(a^2*d*e^{(5*I*d*x + 5*I*c)} - a^2*d*e^{(3*I*d*x + 3*I*c)})*\sqrt{((I*A^2 + 2*A*B - I*B^2)/(a^3*d^2)}*\log((2*\sqrt{1/2}*a^2*d*\sqrt{((I*A^2 + 2*A*B - I*B^2)/(a^3*d^2)})*e^{(I*d*x + I*c)} + \sqrt{2}*((I*A + B)*e^{(2*I*d*x + 2*I*c)} + I*A + B)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1)))/(4*I*A + 4*B)) - 3*\sqrt{1/2}*(a^2*d*e^{(5*I*d*x + 5*I*c)} - a^2*d*e^{(3*I*d*x + 3*I*c)})*\sqrt{((I*A^2 + 2*A*B - I*B^2)/(a^3*d^2)}*\log(-(2*\sqrt{1/2}*a^2*d*\sqrt{((I*A^2 + 2*A*B - I*B^2)/(a^3*d^2)})*e^{(I*d*x + I*c)} - \sqrt{2}*((I*A + B)*e^{(2*I*d*x + 2*I*c)} + I*A + B)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1)))/(4*I*A + 4*B)) - \sqrt{2}*(2*(19*I*A - 4*B)*e^{(6*I*d*x + 6*I*c)} - (-25*I*A + B)*e^{(4*I*d*x + 4*I*c)} + 2*(-7*I*A + 4*B)*e^{(2*I*d*x + 2*I*c)} - I*A + B)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{((-I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} + 1)))/(a^2*d*e^{(5*I*d*x + 5*I*c)} - a^2*d*e^{(3*I*d*x + 3*I*c)})$

3.188.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{(ia(\tan(c + dx) - i))^{\frac{3}{2}} \tan^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(3/2)/(a+I*a*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))/((I*a*(tan(c + d*x) - I))**(3/2)*tan(c + d*x)**(3/2)), x)`

3.188.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is not defined.`

3.188. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$

3.188.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `Timed out`

3.188.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2} (a + a \tan(c + dx) 1i)^{3/2}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + a*tan(c + d*x)*1i)^(3/2)),x)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + a*tan(c + d*x)*1i)^(3/2)), x)`

3.189
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$$

3.189.1 Optimal result	1945
3.189.2 Mathematica [A] (verified)	1946
3.189.3 Rubi [A] (verified)	1946
3.189.4 Maple [B] (verified)	1950
3.189.5 Fricas [B] (verification not implemented)	1951
3.189.6 Sympy [F(-1)]	1952
3.189.7 Maxima [F(-2)]	1952
3.189.8 Giac [F(-1)]	1953
3.189.9 Mupad [F(-1)]	1953

3.189.1 Optimal result

Integrand size = 38, antiderivative size = 240

$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx = \frac{(\frac{1}{4} + \frac{i}{4})(iA+B)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{a^{3/2}d}$$

$$+ \frac{A+iB}{3d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} + \frac{5A+3iB}{2ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}}$$

$$- \frac{(21A+11iB)\sqrt{a+ia \tan(c+dx)}}{6a^2d \tan^{\frac{3}{2}}(c+dx)} + \frac{(39iA-25B)\sqrt{a+ia \tan(c+dx)}}{6a^2d \sqrt{\tan(c+dx)}}$$

output

```
(1/4+1/4*I)*(I*A+B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/a^(3/2)/d+1/6*(39*I*A-25*B)*(a+I*a*tan(d*x+c))^(1/2)/a^2/d/tan(d*x+c)^(1/2)+1/2*(5*A+3*I*B)/a/d/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(3/2)-1/6*(21*A+11*I*B)*(a+I*a*tan(d*x+c))^(1/2)/a^2/d/tan(d*x+c)^(3/2)+1/3*(A+I*B)/d/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(3/2)
```


3.189.2 Mathematica [A] (verified)

Time = 4.31 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.77

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = -\frac{3\sqrt{2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \tan^2(c+dx)}{\sqrt{ia \tan(c+dx)}} + \frac{8iA+24(A+iB) \tan(c+dx)}{12ad \tan^{\frac{3}{2}}(c + dx)}$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^(3/2)),x]`

output `((-3*Sqrt[2]*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Tan[c + d*x]^2)/Sqrt[I*a*Tan[c + d*x]] + ((8*I)*A + 24*(A + I*B)*Tan[c + d*x] + ((114*I)*A - 78*B)*Tan[c + d*x]^2 - 2*(39*A + (25*I)*B)*Tan[c + d*x]^3)/((-I + Tan[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]]))/(12*a*d*Tan[c + d*x]^(3/2))`

3.189.3 Rubi [A] (verified)

Time = 1.38 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.05, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.395$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2}(a + ia \tan(c + dx))^{3/2}} dx \\ & \quad \downarrow \text{4079} \\ & \frac{\int \frac{3(a(3A+iB)-2a(iA-B) \tan(c+dx))}{2 \tan^{\frac{5}{2}}(c+dx) \sqrt{i \tan(c+dx) a+a}} dx}{3a^2} + \frac{A + iB}{3d \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.189. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \frac{\int \frac{a(3A+iB)-2a(iA-B)\tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)\sqrt{i\tan(c+dx)a+a}} dx}{2a^2} + \frac{A+iB}{3d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{a(3A+iB)-2a(iA-B)\tan(c+dx)}{\tan(c+dx)^{5/2}\sqrt{i\tan(c+dx)a+a}} dx}{2a^2} + \frac{A+iB}{3d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{4079} \\
& \frac{\int \frac{\sqrt{i\tan(c+dx)a+a}(a^2(21A+11iB)-4a^2(5iA-3B)\tan(c+dx))}{2\tan^{\frac{5}{2}}(c+dx)} dx}{a^2} + \frac{a(5A+3iB)}{d\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{3d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{\sqrt{i\tan(c+dx)a+a}(a^2(21A+11iB)-4a^2(5iA-3B)\tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx}{2a^2} + \frac{a(5A+3iB)}{d\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{3d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{\sqrt{i\tan(c+dx)a+a}(a^2(21A+11iB)-4a^2(5iA-3B)\tan(c+dx))}{\tan(c+dx)^{5/2}} dx}{2a^2} + \frac{a(5A+3iB)}{d\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{3d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{4081} \\
& \frac{2\int -\frac{\sqrt{i\tan(c+dx)a+a}((39iA-25B)a^3+2(21A+11iB)\tan(c+dx)a^3)}{2\tan^{\frac{3}{2}}(c+dx)} dx}{3a} - \frac{2a^2(21A+11iB)\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \frac{a(5A+3iB)}{d\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}} + \\
& \quad \frac{2a^2}{A+iB} \\
& \quad \frac{3d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{27}
\end{aligned}$$

3.189. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^{3/2}} dx$

$$\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} \left((39iA-25B)a^3+2(21A+11iB) \tan(c+dx)a^3 \right) dx}{\tan^{\frac{3}{2}}(c+dx)} - \frac{2a^2(21A+11iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{a(5A+3iB)}{d \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}}}{2a^2} + \frac{A+iB \frac{2a^2}{3d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}}}{3d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}}$$

↓ 3042

$$\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} \left((39iA-25B)a^3+2(21A+11iB) \tan(c+dx)a^3 \right) dx}{\tan^{\frac{3}{2}}(c+dx)^{\frac{3}{2}}} - \frac{2a^2(21A+11iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{a(5A+3iB)}{d \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}}}{2a^2} + \frac{A+iB \frac{2a^2}{3d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}}}{3d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}}$$

↓ 4081

$$\frac{2 \int \frac{3a^4(A-iB)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx - \frac{2a^3(-25B+39iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(21A+11iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{a(5A+3iB)}{d \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}}}{3a} + \frac{A+iB \frac{2a^2}{3d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}}}{3d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}}$$

↓ 27

$$\frac{3a^3(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(-25B+39iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(21A+11iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{a(5A+3iB)}{d \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}}}{3a} + \frac{A+iB \frac{2a^2}{3d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}}}{3d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}}$$

↓ 3042

$$\frac{3a^3(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(-25B+39iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(21A+11iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{a(5A+3iB)}{d \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}}}{3a} + \frac{A+iB \frac{2a^2}{3d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}}}{3d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}}$$

↓ 4027

3.189. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}} dx$

$$\begin{aligned}
& \frac{6ia^5(A-iB) \int \frac{1}{\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{3a} - \frac{2a^3(-25B+39iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(21A+11iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{a(5A+3iB)}{d \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}} \\
& \frac{A+iB}{3d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}} \\
& \quad \downarrow \text{218} \\
& \frac{\frac{2a^2(21A+11iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{(3-3i)a^{7/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^3(-25B+39iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{a(5A+3iB)}{d \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}} \\
& \frac{A+iB}{3d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}}
\end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^(3/2)),x]`

output `(A + I*B)/(3*d*Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(3/2)) + ((a*(5*A + (3*I)*B))/(d*Tan[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]]) + ((-2*a^2*(21*A + (11*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (((3 - 3*I)*a^(7/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d - (2*a^3*((39*I)*A - 25*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a))/(2*a^2))/(2*a^2)`

3.189.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.189. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}} dx$

```
rule 4027 Int[Sqrt[(a_) + (b_.)*tan[(e_) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*tan[(e_)
+ (f_.)*(x_)], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4079 Int[((a_) + (b_.)*tan[(e_) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

```
rule 4081 Int[((a_) + (b_.)*tan[(e_) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

3.189.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1003 vs. $2(194) = 388$.

Time = 0.17 (sec) , antiderivative size = 1004, normalized size of antiderivative = 4.18

method	result	size
derivativedivides	Expression too large to display	1004
default	Expression too large to display	1004
parts	Expression too large to display	1052

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^(3/2),x,method=_R
ETURNVERBOSE)
```

3.189.
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$$

output

```
-1/24/d*(a*(1+I*tan(d*x+c)))^(1/2)/a^2/tan(d*x+c)^(3/2)*(3*I*B*2^(1/2)*ln(
(2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(
d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2-100*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1
+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^4-48*I*B*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d
*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*ta
n(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan
(d*x+c)^5+384*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x
+c)^3-9*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+
c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^4-9*I*A*2^(1/2
)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a
*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3+9*A*2^(1/2)*ln((2*2^(1/2)*(-I*
a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*
x+c)+I))*a*tan(d*x+c)^4+204*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))
^(1/2)*tan(d*x+c)^2+156*I*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(
1/2)*tan(d*x+c)^4-276*I*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/
2)*tan(d*x+c)^2-9*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*
tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3-32*A
*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)-16*I*A*(-I*
a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-3*A*2^(1/2)*ln((2*2^(1/2)*(-
I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(...
```

3.189.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 569 vs. 2(182) = 364.

Time = 0.29 (sec) , antiderivative size = 569, normalized size of antiderivative = 2.37

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx =$$

$$3 \sqrt{\frac{1}{2}} (a^2 de^{(7i dx + 7i c)} - 2 a^2 de^{(5i dx + 5i c)} + a^2 de^{(3i dx + 3i c)}) \sqrt{\frac{-i A^2 - 2 AB + i B^2}{a^3 d^2}} \log \left(\frac{2i \sqrt{\frac{1}{2}} a^2 d \sqrt{\frac{-i A^2 - 2 AB + i B^2}{a^3 d^2}} e^{(i dx + i c)}}{\dots} \right)$$

input

```
integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^(3/2),x, al
gorithm="fricas")
```

3.189. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$

output `-1/12*(3*sqrt(1/2)*(a^2*d*e^(7*I*d*x + 7*I*c) - 2*a^2*d*e^(5*I*d*x + 5*I*c) + a^2*d*e^(3*I*d*x + 3*I*c))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2))*log((2*I*sqrt(1/2)*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - 3*sqrt(1/2)*(a^2*d*e^(7*I*d*x + 7*I*c) - 2*a^2*d*e^(5*I*d*x + 5*I*c) + a^2*d*e^(3*I*d*x + 3*I*c))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2))*log((-2*I*sqrt(1/2)*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) + sqrt(2)*(2*(26*A + 19*I*B)*e^(8*I*d*x + 8*I*c) - (35*A + 13*I*B)*e^(6*I*d*x + 6*I*c) - 3*(23*A + 13*I*B)*e^(4*I*d*x + 4*I*c) + (19*A + 13*I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(a^2*d*e^(7*I*d*x + 7*I*c) - 2*a^2*d*e^(5*I*d*x + 5*I*c) + a^2*d*e^(3*I*d*x + 3*I*c))`

3.189.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(5/2)/(a+I*a*tan(d*x+c))**(3/2),x)`

output `Timed out`

3.189.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is not defined.`

3.189. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$

3.189.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `Timed out`

3.189.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2} (a + a \tan(c + dx) 1i)^{3/2}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + a*tan(c + d*x)*1i)^(3/2)),x)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + a*tan(c + d*x)*1i)^(3/2)), x)`

3.190
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx$$

3.190.1 Optimal result 1954
 3.190.2 Mathematica [A] (verified) 1955
 3.190.3 Rubi [A] (verified) 1955
 3.190.4 Maple [B] (verified) 1960
 3.190.5 Fracas [B] (verification not implemented) 1961
 3.190.6 Sympy [F(-1)] 1962
 3.190.7 Maxima [F(-2)] 1963
 3.190.8 Giac [F(-2)] 1963
 3.190.9 Mupad [F(-1)] 1963

3.190.1 Optimal result

Integrand size = 38, antiderivative size = 249

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx = \frac{2\sqrt[4]{-1}B \arctan\left(\frac{(-1)^{\frac{3}{4}}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{a^{\frac{5}{2}}d} + \frac{\left(\frac{1}{8} + \frac{i}{8}\right)(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{a^{\frac{5}{2}}d} + \frac{(iA-B)\tan^{\frac{5}{2}}(c+dx)}{5d(a+ia \tan(c+dx))^{\frac{5}{2}}} + \frac{(A+3iB)\tan^{\frac{3}{2}}(c+dx)}{6ad(a+ia \tan(c+dx))^{\frac{3}{2}}} - \frac{(iA-7B)\sqrt{\tan(c+dx)}}{4a^2d\sqrt{a+ia \tan(c+dx)}}$$

output `2*(-1)^(1/4)*B*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/a^(5/2)/d+(1/8+1/8*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/a^(5/2)/d-1/4*(I*A-7*B)*tan(d*x+c)^(1/2)/a^2/d/(a+I*a*tan(d*x+c))^(1/2)+1/5*(I*A-B)*tan(d*x+c)^(5/2)/d/(a+I*a*tan(d*x+c))^(5/2)+1/6*(A+3*I*B)*tan(d*x+c)^(3/2)/a/d/(a+I*a*tan(d*x+c))^(3/2)`

3.190.2 Mathematica [A] (verified)

Time = 6.08 (sec) , antiderivative size = 296, normalized size of antiderivative = 1.19

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx = \frac{240\sqrt[4]{-1}\sqrt{a}\text{Barcsinh}\left(\sqrt[4]{-1}\sqrt{\tan(c+dx)}\right) \sec^3(c+dx)(-i \cos(3(c+dx)))}{(a+ia \tan(c+dx))^{\frac{5}{2}}}$$

input `Integrate[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2), x]`

output `(240*(-1)^(1/4)*Sqrt[a]*B*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^3*((-I)*Cos[3*(c + d*x)] + Sin[3*(c + d*x)]) - (1 + I)*Sec[c + d*x]^2*Sqrt[1 + I*Tan[c + d*x]]*((-1 - I)*Sqrt[a]*(-11*A - (21*I)*B + 2*(13*A + (63*I)*B)*Cos[2*(c + d*x)] + (20*I)*(A + (6*I)*B)*Sin[2*(c + d*x)])*Sqrt[Tan[c + d*x]] + 15*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]]*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)])*Sqrt[a + I*a*Tan[c + d*x]])/(120*a^(5/2)*d*Sqrt[1 + I*Tan[c + d*x]]*(-I + Tan[c + d*x])^2*Sqrt[a + I*a*Tan[c + d*x]])`

3.190.3 Rubi [A] (verified)

Time = 1.59 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.06, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.447$, Rules used = {3042, 4078, 27, 3042, 4078, 27, 3042, 4078, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(c+dx)^{\frac{5}{2}}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx \\ & \quad \downarrow \text{4078} \\ & \frac{(-B+ia) \tan^{\frac{5}{2}}(c+dx)}{5d(a+ia \tan(c+dx))^{\frac{5}{2}}} - \frac{\int \frac{5 \tan^{\frac{3}{2}}(c+dx)(a(iA-B)+2iaB \tan(c+dx))}{2(i \tan(c+dx)a+a)^{\frac{3}{2}}} dx}{5a^2} \end{aligned}$$

3.190. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{\frac{5}{2}}} - \frac{\int \frac{\tan^{\frac{3}{2}}(c+dx)(a(iA-B)+2iaB \tan(c+dx))}{(i \tan(c+dx)a+a)^{3/2}} dx}{2a^2} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{\frac{5}{2}}} - \frac{\int \frac{\tan(c+dx)^{3/2}(a(iA-B)+2iaB \tan(c+dx))}{(i \tan(c+dx)a+a)^{3/2}} dx}{2a^2} \\
& \downarrow 4078 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{\frac{5}{2}}} - \frac{\int -\frac{3\sqrt{\tan(c+dx)}((A+3iB)a^2+4B \tan(c+dx)a^2)}{2\sqrt{i \tan(c+dx)a+a}} dx}{3a^2} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{\frac{3}{2}}} \\
& \downarrow 27 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{\frac{5}{2}}} - \frac{\int \frac{\sqrt{\tan(c+dx)}((A+3iB)a^2+4B \tan(c+dx)a^2)}{\sqrt{i \tan(c+dx)a+a}} dx}{2a^2} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{\frac{3}{2}}} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{\frac{5}{2}}} - \frac{\int \frac{\sqrt{\tan(c+dx)}((A+3iB)a^2+4B \tan(c+dx)a^2)}{\sqrt{i \tan(c+dx)a+a}} dx}{2a^2} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{\frac{3}{2}}} \\
& \downarrow 4078 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{\frac{5}{2}}} - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((iA-7B)a^3+8iB \tan(c+dx)a^3)}{2\sqrt{\tan(c+dx)}} dx}{2a^2} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{\frac{3}{2}}} \\
& \downarrow 27 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{\frac{5}{2}}} - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((iA-7B)a^3+8iB \tan(c+dx)a^3)}{\sqrt{\tan(c+dx)}} dx}{2a^2} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{\frac{3}{2}}} \\
& \downarrow 3042 \\
& \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{\frac{5}{2}}} - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((iA-7B)a^3+8iB \tan(c+dx)a^3)}{\sqrt{\tan(c+dx)}} dx}{2a^2} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{\frac{3}{2}}}
\end{aligned}$$

3.190. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx$

$$\begin{array}{c}
\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \\
\frac{a^2(-7B + iA) \sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{\int \frac{\sqrt{i \tan(c + dx)a + a} \left((iA - 7B)a^3 + 8iB \tan(c + dx)a^3 \right) dx}{\sqrt{\tan(c + dx)}}}{2a^2} - \frac{a(A + 3iB) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} \\
\hline
2a^2 \\
\downarrow 4084 \\
\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \\
\frac{a^2(-7B + iA) \sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{a^3(B + iA) \int \frac{\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx - 8a^2 B \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx}{2a^2} - \frac{a(A + 3iB) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} \\
\hline
2a^2 \\
\downarrow 3042 \\
\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \\
\frac{a^2(-7B + iA) \sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{a^3(B + iA) \int \frac{\sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx - 8a^2 B \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx}{2a^2} - \frac{a(A + 3iB) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} \\
\hline
2a^2 \\
\downarrow 4027 \\
\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \\
\frac{a^2(-7B + iA) \sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{-8a^2 B \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx - \frac{2ia^5(B + iA) \int \frac{1}{i \tan(c + dx)a + a} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{i \tan(c + dx)a + a}}}{2a^2}}{2a^2} - \frac{a(A + 3iB) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} \\
\hline
2a^2 \\
\downarrow 218 \\
\frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \\
\frac{a^2(-7B + iA) \sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{(1 - i)a^{7/2}(B + iA) \operatorname{arctanh}\left(\frac{(1 + i)\sqrt{a} \sqrt{\tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right) - 8a^2 B \int \frac{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx)a + a}}{\sqrt{\tan(c + dx)}} dx}{2a^2} - \frac{a(A + 3iB) \tan^{\frac{3}{2}}(c + dx)}{3d(a + ia \tan(c + dx))^{3/2}} \\
\hline
2a^2 \\
\downarrow 4082
\end{array}$$

3.190. $\int \frac{\tan^{\frac{5}{2}}(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{a^2(-7B + iA)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{(1-i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{8a^4 B \int \frac{1}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{2a^2} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))} \\
 & \frac{a^2(-7B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} - \frac{(1-i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{8a^4 B \int \frac{1}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{2a^2} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))} \\
 & \quad \downarrow 65 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{a^2(-7B + iA)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{(1-i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{16a^4 B \int \frac{1}{1-\frac{ia \tan(c+dx)}{i \tan(c+dx)a+a}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{2a^2} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))} \\
 & \frac{a^2(-7B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} - \frac{(1-i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{16a^4 B \int \frac{1}{1-\frac{ia \tan(c+dx)}{i \tan(c+dx)a+a}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{2a^2} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))} \\
 & \quad \downarrow 216 \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{a^2(-7B + iA)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{(1-i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{16\sqrt[4]{-1}a^{7/2}B \operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))} \\
 & \frac{a^2(-7B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} - \frac{(1-i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{16\sqrt[4]{-1}a^{7/2}B \operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))} \\
 & \frac{(-B + iA) \tan^{\frac{5}{2}}(c + dx)}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{a^2(-7B + iA)\sqrt{\tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} - \frac{(1-i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{16\sqrt[4]{-1}a^{7/2}B \operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))} \\
 & \frac{a^2(-7B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} - \frac{(1-i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{16\sqrt[4]{-1}a^{7/2}B \operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))}
 \end{aligned}$$

input `Int[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `((I*A - B)*Tan[c + d*x]^(5/2))/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) - (-1/3*(a*(A + (3*I)*B)*Tan[c + d*x]^(3/2))/(d*(a + I*a*Tan[c + d*x])^(3/2)) + (-1/2*((16*(-1)^(1/4)*a^(7/2)*B*ArcTan[(-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]]]/Sqrt[a + I*a*Tan[c + d*x]]])/d + ((1 - I)*a^(7/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]]]/Sqrt[a + I*a*Tan[c + d*x]]])/d)/a^2 + (a^2*(I*A - 7*B)*Sqrt[Tan[c + d*x]]/(d*Sqrt[a + I*a*Tan[c + d*x]]))/(2*a^2)`

3.190. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

3.190.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

```
rule 4082 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 + b^2, 0] && EqQ[A*b + a*B, 0]
```

```
rule 4084 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x]
- Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[
e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

3.190.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1529 vs. $2(198) = 396$.

Time = 0.17 (sec) , antiderivative size = 1530, normalized size of antiderivative = 6.14

method	result	size
parts	Expression too large to display	1530
derivativedivides	Expression too large to display	1532
default	Expression too large to display	1532

```
input int(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x,method=_R
ETURNVERBOSE)
```

output

```

-1/240*A/d*(a*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^(1/2)/a^3*(148*(-I*a)^(1/2)
*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3+60*I*2^(1/2)*ln((2*2
^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+
c))/(tan(d*x+c)+I))*tan(d*x+c)^3*a-15*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*t
an(d*x+c)^4*a-60*I*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*t
an(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*tan(d*x+c)*a-308*I*(
-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2+90*2^(1/2)*
ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*t
an(d*x+c))/(tan(d*x+c)+I))*tan(d*x+c)^2*a+60*I*(-I*a)^(1/2)*(a*tan(d*x+c)*
(1+I*tan(d*x+c)))^(1/2)-15*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c
)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a-220*tan(d*
x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2))/(a*tan(d*x+c)*(1+
I*tan(d*x+c)))^(1/2)/(-tan(d*x+c)+I)^4/(-I*a)^(1/2)-1/80*B/d*(a*(1+I*tan(d
*x+c)))^(1/2)*tan(d*x+c)^(1/2)*(5*I*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+
c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*(I*a)^(1/2
)*2^(1/2)*tan(d*x+c)^4*a-30*I*(I*a)^(1/2)*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2
))*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I
))*tan(d*x+c)^2*a-320*I*(-I*a)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+
c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*tan(d*x+c)^3*a+1...

```

3.190.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 777 vs. $2(185) = 370$.

Time = 0.34 (sec) , antiderivative size = 777, normalized size of antiderivative = 3.12

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx = \left(15 \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{i A^2 + 2 A B - i B^2}{a^5 d^2}} e^{(5i dx + 5i c)} \log \left(\frac{2 \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{i A^2 + 2 A B - i B^2}{a^5 d^2}} e^{(5i dx + 5i c)}}{\dots} \right) \right)$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="fricas")`

$$3.190. \quad \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx$$

output

```

1/120*(15*sqrt(1/2)*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2))*e^(5*I*d
*x + 5*I*c)*log((2*sqrt(1/2)*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2))
*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(
a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x
+ 2*I*c) + 1)))/(4*I*A + 4*B)) - 15*sqrt(1/2)*a^3*d*sqrt((I*A^2 + 2*A*B -
I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-(2*sqrt(1/2)*a^3*d*sqrt((I*A^2
+ 2*A*B - I*B^2)/(a^5*d^2))*e^(I*d*x + I*c) - sqrt(2)*((I*A + B)*e^(2*I*d
*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d
*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - 30*a^3*d*sqr
t(-4*I*B^2/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(52/605*(4*sqrt(2)*(B*e^(3*I*
d*x + 3*I*c) + B*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((
-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) + (3*a^3*d*e^(2*I*d
*x + 2*I*c) - a^3*d)*sqrt(-4*I*B^2/(a^5*d^2)))/(B*e^(2*I*d*x + 2*I*c) + B)
) + 30*a^3*d*sqrt(-4*I*B^2/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(52/605*(4*sq
rt(2)*(B*e^(3*I*d*x + 3*I*c) + B*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I
*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)) - (
3*a^3*d*e^(2*I*d*x + 2*I*c) - a^3*d)*sqrt(-4*I*B^2/(a^5*d^2)))/(B*e^(2*I*d
*x + 2*I*c) + B)) + sqrt(2)*((-23*I*A + 123*B)*e^(6*I*d*x + 6*I*c) - 6*(2*
I*A - 17*B)*e^(4*I*d*x + 4*I*c) - 2*(-4*I*A + 9*B)*e^(2*I*d*x + 2*I*c) - 3
*I*A + 3*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I...

```

3.190.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(5/2),x)`

output `Timed out`

3.190.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{\frac{5}{2}}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.190.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{\frac{5}{2}}} dx = \text{Exception raised: TypeError}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[32]Warning, replacing 32 by 64, a substitution variable`

3.190.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{\frac{5}{2}}} dx = \int \frac{\tan(c+dx)^{\frac{5}{2}}(A+B\tan(c+dx))}{(a+a\tan(c+dx)li)^{\frac{5}{2}}} dx$$

input `int((tan(c+d*x)^(5/2)*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*1i)^(5/2),x)`

3.190. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{\frac{5}{2}}} dx$

output `int((tan(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2), x)`

3.190. $\int \frac{\tan^5(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

3.191
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$$

3.191.1 Optimal result	1965
3.191.2 Mathematica [A] (verified)	1965
3.191.3 Rubi [A] (verified)	1966
3.191.4 Maple [B] (verified)	1969
3.191.5 Fricas [B] (verification not implemented)	1970
3.191.6 Sympy [F(-1)]	1971
3.191.7 Maxima [F(-2)]	1971
3.191.8 Giac [F(-2)]	1972
3.191.9 Mupad [F(-1)]	1972

3.191.1 Optimal result

Integrand size = 38, antiderivative size = 194

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx =$$

$$-\frac{(\frac{1}{8} - \frac{i}{8})(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{a^{5/2}d} + \frac{(iA-B)\tan^{\frac{3}{2}}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

$$+ \frac{(A+11iB)\sqrt{\tan(c+dx)}}{30ad(a+ia \tan(c+dx))^{3/2}} + \frac{(13A-37iB)\sqrt{\tan(c+dx)}}{60a^2d\sqrt{a+ia \tan(c+dx)}}$$

output

```
(-1/8+1/8*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/a^(5/2)/d+1/60*(13*A-37*I*B)*tan(d*x+c)^(1/2)/a^2/d/(a+I*a*tan(d*x+c))^(1/2)+1/5*(I*A-B)*tan(d*x+c)^(3/2)/d/(a+I*a*tan(d*x+c))^(5/2)+1/30*(A+11*I*B)*tan(d*x+c)^(1/2)/a/d/(a+I*a*tan(d*x+c))^(3/2)
```

3.191.2 Mathematica [A] (verified)

Time = 3.70 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.15

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx =$$

$$\frac{\sec^2(c+dx)\sqrt{\tan(c+dx)}\left(2(A+11iB+2(7A-13iB)\cos(2(c+dx))+20(iA+B)\sin(2(c+dx)))\sqrt{ia \tan(c+dx)}\right)}{120a^2d\sqrt{ia \tan(c+dx)}(-i + \dots)}$$

3.191.
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$$

input `Integrate[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `-1/120*(Sec[c + d*x]^2*Sqrt[Tan[c + d*x]]*(2*(A + (11*I)*B + 2*(7*A - (13*I)*B)*Cos[2*(c + d*x)] + 20*(I*A + B)*Sin[2*(c + d*x)])*Sqrt[I*a*Tan[c + d*x]] - 15*Sqrt[2]*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*(Cos[2*(c + d*x)] + I*Ssin[2*(c + d*x)])*Sqrt[a + I*a*Tan[c + d*x]])/(a^2*d*Sqrt[I*a*Tan[c + d*x]]*(-I + Tan[c + d*x])^2*Sqrt[a + I*a*Tan[c + d*x]])`

3.191.3 Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.06, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {3042, 4078, 27, 3042, 4078, 27, 3042, 4079, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^{3/2}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+IA) \tan^{\frac{3}{2}}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\sqrt{\tan(c+dx)}(3a(iA-B)-2a(A-4iB) \tan(c+dx))}{2(i \tan(c+dx)a+a)^{3/2}} dx}{5a^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B+IA) \tan^{\frac{3}{2}}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\sqrt{\tan(c+dx)}(3a(iA-B)-2a(A-4iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^{3/2}} dx}{10a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B+IA) \tan^{\frac{3}{2}}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\sqrt{\tan(c+dx)}(3a(iA-B)-2a(A-4iB) \tan(c+dx))}{(i \tan(c+dx)a+a)^{3/2}} dx}{10a^2} \\
 & \quad \downarrow \text{4078}
 \end{aligned}$$

3.191. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

input `Int[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `((I*A - B)*Tan[c + d*x]^(3/2))/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) - (-1/3*(a*(A + (11*I)*B)*Sqrt[Tan[c + d*x]]/(d*(a + I*a*Tan[c + d*x])^(3/2)) + ((15/2 - (15*I)/2)*a^(3/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (a^2*(13*A - (37*I)*B)*Sqrt[Tan[c + d*x]]/(d*Sqrt[a + I*a*Tan[c + d*x]]))/(6*a^2))/(10*a^2)`

3.191.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

```
rule 4079 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.191.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1085 vs. $2(156) = 312$.

Time = 0.14 (sec) , antiderivative size = 1086, normalized size of antiderivative = 5.60

method	result	size
derivativedivides	Expression too large to display	1086
default	Expression too large to display	1086
parts	Expression too large to display	1143

```
input int(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x,method=_R
ETURNVERBOSE)
```



```
output 1/240/d*tan(d*x+c)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)/a^3*(15*I*A*2^(1/2)*ln
((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan
(d*x+c))/(tan(d*x+c)+I))*a-148*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c
)))^(1/2)*tan(d*x+c)^3+60*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*
x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x
+c)+15*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))
)^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^4-212*A*(-I*a)^(1
/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2-60*I*B*(-I*a)^(1/2)
*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-52*I*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(
1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3+60*A*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan
(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2
)*a*tan(d*x+c)^3-90*I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(
1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2-
60*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^
(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3-90*B*2^(1/2)*ln((
2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d
*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2+308*I*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(
1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2+15*I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1
/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+
I))*a*tan(d*x+c)^4-60*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c...
```

3.191.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 459 vs. 2(144) = 288.

Time = 0.27 (sec) , antiderivative size = 459, normalized size of antiderivative = 2.37

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx =$$

$$\left(15 \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{-iA^2-2AB+iB^2}{a^5 d^2}} e^{(5i dx+5i c)} \log \left(\frac{2i \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{-iA^2-2AB+iB^2}{a^5 d^2}} e^{(i dx+i c)} + \sqrt{2}((iA+B)e^{(2i dx+2i c)} + iA+B) \sqrt{e^{(2i dx+2i c)}}}{4iA+4B} \right) \right)$$

```
input integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, al
gorithm="fricas")
```

3.191. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

```
output -1/120*(15*sqrt(1/2)*a^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2))*e^(5*I
*d*x + 5*I*c)*log((2*I*sqrt(1/2)*a^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^5*
d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*
sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*
I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - 15*sqrt(1/2)*a^3*d*sqrt((-I*A^2 - 2
*A*B + I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log((-2*I*sqrt(1/2)*a^3*d*sq
r t((-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)
*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I
*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - sqr
t(2)*((17*A - 23*I*B)*e^(6*I*d*x + 6*I*c) + 6*(3*A - 2*I*B)*e^(4*I*d*x + 4
*I*c) - 2*(A - 4*I*B)*e^(2*I*d*x + 2*I*c) - 3*A - 3*I*B)*sqrt(a/(e^(2*I*d*
x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) +
1)))*e^(-5*I*d*x - 5*I*c)/(a^3*d)
```

3.191.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

```
input integrate(tan(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(5/2),x)
```

```
output Timed out
```

3.191.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, al
gorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un
defined.
```

3.191. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx$

3.191.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[-83]Warning, replacing -83 by -8, a substitution varia`

3.191.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \int \frac{\tan(c+dx)^{3/2}(A+B\tan(c+dx))}{(a+a\tan(c+dx)li)^{5/2}} dx$$

input `int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2),x)`

output `int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2), x)`

$$3.192 \quad \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$$

3.192.1 Optimal result	1973
3.192.2 Mathematica [A] (verified)	1973
3.192.3 Rubi [A] (verified)	1974
3.192.4 Maple [B] (verified)	1977
3.192.5 Fricas [B] (verification not implemented)	1978
3.192.6 Sympy [F]	1979
3.192.7 Maxima [F(-2)]	1979
3.192.8 Giac [F(-2)]	1980
3.192.9 Mupad [F(-1)]	1980

3.192.1 Optimal result

Integrand size = 38, antiderivative size = 196

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx =$$

$$-\frac{\left(\frac{1}{8} + \frac{i}{8}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{a^{5/2}d} + \frac{(iA - B)\sqrt{\tan(c+dx)}}{5d(a+ia \tan(c+dx))^{5/2}}$$

$$+ \frac{(3iA + 7B)\sqrt{\tan(c+dx)}}{30ad(a+ia \tan(c+dx))^{3/2}} - \frac{(3iA - 13B)\sqrt{\tan(c+dx)}}{60a^2d\sqrt{a+ia \tan(c+dx)}}$$

```
output (-1/8-1/8*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/a^(5/2)/d-1/60*(3*I*A-13*B)*tan(d*x+c)^(1/2)/a^2/d/(a+I*a*tan(d*x+c))^(1/2)+1/5*(I*A-B)*tan(d*x+c)^(1/2)/d/(a+I*a*tan(d*x+c))^(5/2)+1/30*(3*I*A+7*B)*tan(d*x+c)^(1/2)/a/d/(a+I*a*tan(d*x+c))^(3/2)
```

3.192.2 Mathematica [A] (verified)

Time = 3.00 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.13

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{\sec^2(c+dx)\sqrt{\tan(c+dx)}\left(-2i(9A-iB+2(3A-7iB)\cos(2(c+dx)))\right)}{(a+ia \tan(c+dx))^{5/2}}$$

input `Integrate[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])5/2,x]`

output `(Sec[c + d*x]^2*Sqrt[Tan[c + d*x]]*((-2*I)*(9*A - I*B + 2*(3*A - (7*I)*B)*Cos[2*(c + d*x)] + 20*B*Sin[2*(c + d*x)])*Sqrt[I*a*Tan[c + d*x]] + 15*Sqrt[2]*(I*A + B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)])*Sqrt[a + I*a*Tan[c + d*x]])/(120*a^2*d*Sqrt[I*a*Tan[c + d*x]]*(-I + Tan[c + d*x])^2*Sqrt[a + I*a*Tan[c + d*x]])`

3.192.3 Rubi [A] (verified)

Time = 1.07 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.06, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {3042, 4078, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{4078} \\
 & \frac{(-B+iA)\sqrt{\tan(c+dx)}}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{a(iA-B)-2a(2A-3iB) \tan(c+dx)}{2\sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^{3/2}} dx}{5a^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B+iA)\sqrt{\tan(c+dx)}}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{a(iA-B)-2a(2A-3iB) \tan(c+dx)}{\sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^{3/2}} dx}{10a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B+iA)\sqrt{\tan(c+dx)}}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{a(iA-B)-2a(2A-3iB) \tan(c+dx)}{\sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^{3/2}} dx}{10a^2} \\
 & \quad \downarrow \text{4079}
 \end{aligned}$$

3.192. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\int \frac{a^2(9iA+B) - 2a^2(3A-7iB) \tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} dx}{3a^2} - \frac{a(7B+3iA)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}}$$

27

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\int \frac{a^2(9iA+B) - 2a^2(3A-7iB) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} dx}{6a^2} - \frac{a(7B+3iA)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}}$$

3042

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\int \frac{a^2(9iA+B) - 2a^2(3A-7iB) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} dx}{6a^2} - \frac{a(7B+3iA)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}}$$

4079

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\int \frac{15a^3(iA+B)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx + \frac{a^2(-13B+3iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} - \frac{a(7B+3iA)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}}$$

27

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\frac{15}{2}a(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + \frac{a^2(-13B+3iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} - \frac{a(7B+3iA)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}}$$

3042

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\frac{15}{2}a(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + \frac{a^2(-13B+3iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} - \frac{a(7B+3iA)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}}$$

4027

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\frac{a^2(-13B+3iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} - \frac{15ia^3(B+iA) \int \frac{1}{2 \tan(c+dx)a^2} dx - ia \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{6a^2}}{10a^2} - \frac{a(7B+3iA)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}}$$

218

$$\frac{(-B + iA)\sqrt{\tan(c + dx)}}{5d(a + ia \tan(c + dx))^{5/2}} - \frac{\left(\frac{15}{2} - \frac{15i}{2}\right)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) + \frac{a^2(-13B+3iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} - \frac{a(7B+3iA)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}}$$

10a²

3.192. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

input `Int[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `((I*A - B)*Sqrt[Tan[c + d*x]]/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) - (-1/3*(a*((3*I)*A + 7*B)*Sqrt[Tan[c + d*x]]/(d*(a + I*a*Tan[c + d*x])^(3/2)) + (((15/2 - (15*I)/2)*a^(3/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d + (a^2*((3*I)*A - 13*B)*Sqrt[Tan[c + d*x]]/(d*Sqrt[a + I*a*Tan[c + d*x]]))/(6*a^2))/(10*a^2)`

3.192.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b - a*B)) * (a + b*Tan[e + f*x])^m * ((c + d*Tan[e + f*x])^n / (2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1) * (c + d*Tan[e + f*x])^(n - 1) * Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n)) * Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

```
rule 4079 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.192.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1085 vs. $2(158) = 316$.

Time = 0.14 (sec) , antiderivative size = 1086, normalized size of antiderivative = 5.54

method	result	size
derivativedivides	Expression too large to display	1086
default	Expression too large to display	1086
parts	Expression too large to display	1143

```
input int(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x,method=_R
ETURNVERBOSE)
```



```
output 1/240/d*tan(d*x+c)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)/a^3*(-60*I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)-12*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3-90*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2-15*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^4-212*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2+220*I*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+15*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^4+60*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3+15*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a+60*I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3+90*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2+12*I*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2+60*I*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-60*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*...
```

3.192.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 461 vs. 2(146) = 292.

Time = 0.29 (sec) , antiderivative size = 461, normalized size of antiderivative = 2.35

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \left(15 \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{i A^2+2 AB-i B^2}{a^5 d^2}} e^{(5i dx+5i c)} \log \left(\frac{2 \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{i A^2+2 AB-i B^2}{a^5 d^2}} e^{(i dx+i c)} + \sqrt{2}((i A+B)e^{(2i dx+2i c)}+i A+B) \sqrt{\frac{a}{e^{(2i dx+2i c)}}}}{4i A+4 B} \right) \right)$$

```
input integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="fricas")
```

```
output -1/120*(15*sqrt(1/2)*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log((2*sqrt(1/2)*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2)))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - 15*sqrt(1/2)*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-(2*sqrt(1/2)*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2))*e^(I*d*x + I*c) - sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - sqrt(2)*((3*I*A + 17*B)*e^(6*I*d*x + 6*I*c) - 6*(-2*I*A - 3*B)*e^(4*I*d*x + 4*I*c) - 2*(-6*I*A + B)*e^(2*I*d*x + 2*I*c) + 3*I*A - 3*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-5*I*d*x - 5*I*c)/(a^3*d)
```

3.192.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \int \frac{(A+B \tan(c+dx)) \sqrt{\tan(c+dx)}}{(ia(\tan(c+dx)-i))^{5/2}} dx$$

```
input integrate(tan(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(5/2),x)
```

```
output Integral((A + B*tan(c + d*x))*sqrt(tan(c + d*x))/(I*a*(tan(c + d*x) - I))**(5/2), x)
```

3.192.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is not defined.
```

3.192. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

3.192.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:Non regular value [0] was discarded and replaced randomly by 0=[-27]Warning, replacing -27 by 20, a substitution varia`

3.192.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+a\tan(c+dx)1i)^{5/2}} dx$$

input `int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2),x)`

output `int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2), x)`

3.193
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}} dx$$

3.193.1 Optimal result 1981
 3.193.2 Mathematica [A] (verified) 1981
 3.193.3 Rubi [A] (verified) 1982
 3.193.4 Maple [B] (verified) 1985
 3.193.5 Fracas [B] (verification not implemented) 1986
 3.193.6 Sympy [F(-1)] 1986
 3.193.7 Maxima [F(-2)] 1987
 3.193.8 Giac [F(-1)] 1987
 3.193.9 Mupad [F(-1)] 1987

3.193.1 Optimal result

Integrand size = 38, antiderivative size = 194

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx = \frac{\left(\frac{1}{8} - \frac{i}{8}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{a^{5/2}d} + \frac{(A + iB)\sqrt{\tan(c + dx)}}{5d(a + ia \tan(c + dx))^{5/2}} + \frac{(13A + 3iB)\sqrt{\tan(c + dx)}}{30ad(a + ia \tan(c + dx))^{3/2}} + \frac{(67A - 3iB)\sqrt{\tan(c + dx)}}{60a^2d\sqrt{a + ia \tan(c + dx)}}$$

output `(1/8-1/8*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/a^(5/2)/d+1/60*(67*A-3*I*B)*tan(d*x+c)^(1/2)/a^2/d/(a+I*a*tan(d*x+c))^(1/2)+1/5*(A+I*B)*tan(d*x+c)^(1/2)/d/(a+I*a*tan(d*x+c))^(5/2)+1/30*(13*A+3*I*B)*tan(d*x+c)^(1/2)/a/d/(a+I*a*tan(d*x+c))^(3/2)`

3.193.2 Mathematica [A] (verified)

Time = 4.70 (sec) , antiderivative size = 242, normalized size of antiderivative = 1.25

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx = \frac{\sqrt{\tan(c + dx)}\left(24a^5(A + iB)\sqrt{ia \tan(c + dx)} - (a + ia \tan(c + dx))\right)}{\dots}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(5/2)),x]`

output $(\text{Sqrt}[\text{Tan}[c + d*x]]*(24*a^5*(A + I*B)*\text{Sqrt}[I*a*\text{Tan}[c + d*x]] - (a + I*a*\text{Tan}[c + d*x])*(-4*a^4*(13*A + (3*I)*B)*\text{Sqrt}[I*a*\text{Tan}[c + d*x]] + (a + I*a*\text{Tan}[c + d*x])*(2*a^3*(-67*A + (3*I)*B)*\text{Sqrt}[I*a*\text{Tan}[c + d*x]] - 15*\text{Sqrt}[2]*a^3*(A - I*B)*\text{ArcTan}[\text{H}[(\text{Sqrt}[2]*\text{Sqrt}[I*a*\text{Tan}[c + d*x]])/\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])]*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]))/(\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]*(a + I*a*\text{Tan}[c + d*x])^(5/2))$

3.193.3 Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.06, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx \\
 & \quad \downarrow 4079 \\
 & \frac{\int \frac{a(9A - iB) - 4a(iA - B) \tan(c + dx)}{2\sqrt{\tan(c + dx)}(i \tan(c + dx)a + a)^{3/2}} dx}{5a^2} + \frac{(A + iB)\sqrt{\tan(c + dx)}}{5d(a + ia \tan(c + dx))^{5/2}} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{a(9A - iB) - 4a(iA - B) \tan(c + dx)}{\sqrt{\tan(c + dx)}(i \tan(c + dx)a + a)^{3/2}} dx}{10a^2} + \frac{(A + iB)\sqrt{\tan(c + dx)}}{5d(a + ia \tan(c + dx))^{5/2}} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{a(9A - iB) - 4a(iA - B) \tan(c + dx)}{\sqrt{\tan(c + dx)}(i \tan(c + dx)a + a)^{3/2}} dx}{10a^2} + \frac{(A + iB)\sqrt{\tan(c + dx)}}{5d(a + ia \tan(c + dx))^{5/2}} \\
 & \quad \downarrow 4079 \\
 & \frac{\int \frac{a^2(41A - 9iB) - 2a^2(13iA - 3B) \tan(c + dx)}{2\sqrt{\tan(c + dx)}\sqrt{i \tan(c + dx)a + a}} dx}{3a^2} + \frac{a(13A + 3iB)\sqrt{\tan(c + dx)}}{3d(a + ia \tan(c + dx))^{3/2}} + \frac{(A + iB)\sqrt{\tan(c + dx)}}{5d(a + ia \tan(c + dx))^{5/2}} \\
 & \quad \downarrow 27
 \end{aligned}$$

3.193. $\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx$

$$\begin{aligned}
& \frac{\int \frac{a^2(41A-9iB)-2a^2(13iA-3B)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}} dx + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}}}{10a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{a^2(41A-9iB)-2a^2(13iA-3B)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}} dx + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}}}{10a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 4079 \\
& \frac{\int \frac{15a^3(A-iB)\sqrt{i\tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx + \frac{a^2(67A-3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}}}{10a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 27 \\
& \frac{\frac{15}{2}a(A-iB)\int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + \frac{a^2(67A-3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}}{10a^2} + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 3042 \\
& \frac{\frac{15}{2}a(A-iB)\int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + \frac{a^2(67A-3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}}{10a^2} + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 4027 \\
& \frac{\frac{a^2(67A-3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{15ia^3(A-iB)\int \frac{1}{\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{10a^2} + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} + \\
& \quad \frac{(A+iB)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 218 \\
& \frac{\left(\frac{15}{2}-\frac{15i}{2}\right)a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} + \frac{a^2(67A-3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}}}{10a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}}
\end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]])*(a + I*a*Tan[c + d*x])^(5/2)),x]`

$$3.193. \quad \int \frac{A+B\tan(c+dx)}{\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{5/2}} dx$$

```
output ((A + I*B)*Sqrt[Tan[c + d*x]]/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) + ((a*(1
3*A + (3*I)*B)*Sqrt[Tan[c + d*x]]/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) + ((
(15/2 - (15*I)/2)*a^(3/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c +
d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/d + (a^2*(67*A - (3*I)*B)*Sqrt[Tan[c +
d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]]))/(6*a^2)/(10*a^2)
```

3.193.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4027 Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_)
+ (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4079 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.193.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1085 vs. $2(156) = 312$.

Time = 0.15 (sec) , antiderivative size = 1086, normalized size of antiderivative = 5.60

method	result	size
derivativedivides	Expression too large to display	1086
default	Expression too large to display	1086
parts	Expression too large to display	1143

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(5/2),x,method=_R
ETURNVERBOSE)
```

```
output -1/240/d*tan(d*x+c)^(1/2)*(a*(1+I*tan(d*x+c)))^(1/2)*(15*I*A*2^(1/2)*ln((2
*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*
x+c))/(tan(d*x+c)+I))*a+60*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d
*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*
x+c)+15*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)
))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^4-60*I*B*(-I*a)^(
1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+268*I*A*(-I*a)^(1/2)*(a*tan(d*
x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3+60*A*ln((2*2^(1/2)*(-I*a)^(1/2)*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*
2^(1/2)*a*tan(d*x+c)^3-90*I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*
x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x
+c)^2-60*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x
+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3-90*B*2^(1/2
)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a
*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2+12*B*(-I*a)^(1/2)*(a*tan(d*x+c)
*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3-12*I*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(
1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2+15*I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1
/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+
I))*a*tan(d*x+c)^4-60*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(
1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)...
```


3.193.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 460 vs. $2(146) = 292$.

Time = 0.29 (sec) , antiderivative size = 460, normalized size of antiderivative = 2.37

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx = \frac{\left(15 \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{-i A^2 - 2AB + i B^2}{a^5 d^2}} e^{(5i dx + 5i c)} \log \left(\frac{2i \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{-i A^2 - 2AB + i B^2}{a^5 d^2}}}{\dots} \right) \right)}{\dots}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="fricas")`

output `1/120*(15*sqrt(1/2)*a^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log((2*I*sqrt(1/2)*a^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - 15*sqrt(1/2)*a^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log((-2*I*sqrt(1/2)*a^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) + sqrt(2)*((83*A + 3*I*B)*e^(6*I*d*x + 6*I*c) + 6*(17*A + 2*I*B)*e^(4*I*d*x + 4*I*c) + 2*(11*A + 6*I*B)*e^(2*I*d*x + 2*I*c) + 3*A + 3*I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))*e^(-5*I*d*x - 5*I*c)/(a^3*d)`

3.193.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+I*a*tan(d*x+c))**(5/2),x)`

output `Timed out`

3.193.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.193.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Timed out`

3.193.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + a \tan(c + dx) 1i)^{5/2}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(5/2)),x)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(5/2)), x)`

3.194
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$$

3.194.1 Optimal result	1988
3.194.2 Mathematica [A] (verified)	1989
3.194.3 Rubi [A] (verified)	1989
3.194.4 Maple [B] (verified)	1993
3.194.5 Fracas [B] (verification not implemented)	1994
3.194.6 Sympy [F(-1)]	1995
3.194.7 Maxima [F(-2)]	1995
3.194.8 Giac [F(-1)]	1996
3.194.9 Mupad [F(-1)]	1996

3.194.1 Optimal result

Integrand size = 38, antiderivative size = 240

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \frac{(\frac{1}{8} + \frac{i}{8})(A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{a^{5/2}d}$$

$$+ \frac{A + iB}{5d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}} + \frac{17A + 7iB}{30ad\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{3/2}}$$

$$+ \frac{151A + 41iB}{60a^2d\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}} - \frac{(317A + 67iB)\sqrt{a + ia \tan(c + dx)}}{60a^3d\sqrt{\tan(c + dx)}}$$

output

```
(1/8+1/8*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/a^(5/2)/d+1/60*(151*A+41*I*B)/a^2/d/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2)-1/60*(317*A+67*I*B)*(a+I*a*tan(d*x+c))^(1/2)/a^3/d/tan(d*x+c)^(1/2)+1/5*(A+I*B)/d/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(5/2)+1/30*(17*A+7*I*B)/a/d/tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(3/2)
```

3.194.2 Mathematica [A] (verified)

Time = 4.30 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.90

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \frac{\sec^2(c + dx) \left(\frac{15\sqrt{2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right) (-i \cos(2(c + dx)) + \sin(2(c + dx)))}{\sqrt{ia \tan(c + dx)}} \right)}{120}$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(5/2)),x]`

output `(Sec[c + d*x]^2*((15*Sqrt[2]*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*(-I)*Cos[2*(c + d*x)] + Sin[2*(c + d*x)])*Tan[c + d*x])/Sqrt[I*a*Tan[c + d*x]] + ((2*I)*((340*I)*A - 80*B + (149*A + (19*I)*B)*Tan[c + d*x] + Cos[2*(c + d*x)]*((-460*I)*A + 80*B + (466*A + (86*I)*B)*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]])/(120*a^2*d*Sqrt[Tan[c + d*x]]*(-I + Tan[c + d*x])^2)`

3.194.3 Rubi [A] (verified)

Time = 1.39 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.05, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.395$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2}(a + ia \tan(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{4079} \\ & \frac{\int \frac{a(11A + iB) - 6a(iA - B) \tan(c + dx)}{2 \tan^{\frac{3}{2}}(c + dx)(i \tan(c + dx)a + a)^{3/2}} dx}{5a^2} + \frac{A + iB}{5d\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^{5/2}} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.194. $\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{\int \frac{a(11A+iB)-6a(iA-B)\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(i\tan(c+dx)a+a)^{3/2}} dx}{10a^2} + \frac{A+iB}{5d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{5/2}} \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \frac{\int \frac{a(11A+iB)-6a(iA-B)\tan(c+dx)}{\tan(c+dx)^{3/2}(i\tan(c+dx)a+a)^{3/2}} dx}{10a^2} + \frac{A+iB}{5d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{5/2}} \\
 & \qquad \qquad \qquad \downarrow \text{4079} \\
 & \frac{\int \frac{a^2(83A+13iB)-4a^2(17iA-7B)\tan(c+dx)}{2\tan^{\frac{3}{2}}(c+dx)\sqrt{i\tan(c+dx)a+a}} dx}{\frac{10a^2}{A+iB}} + \frac{a(17A+7iB)}{3d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{3/2}} + \\
 & \qquad \qquad \qquad \frac{10a^2}{A+iB} \\
 & \qquad \qquad \qquad \frac{5d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{5/2}} \\
 & \qquad \qquad \qquad \downarrow \text{27} \\
 & \frac{\int \frac{a^2(83A+13iB)-4a^2(17iA-7B)\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} + \frac{a(17A+7iB)}{3d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{3/2}} + \\
 & \qquad \qquad \qquad \frac{10a^2}{A+iB} \\
 & \qquad \qquad \qquad \frac{5d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{5/2}} \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \frac{\int \frac{a^2(83A+13iB)-4a^2(17iA-7B)\tan(c+dx)}{\tan(c+dx)^{3/2}\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} + \frac{a(17A+7iB)}{3d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{3/2}} + \\
 & \qquad \qquad \qquad \frac{10a^2}{A+iB} \\
 & \qquad \qquad \qquad \frac{5d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{5/2}} \\
 & \qquad \qquad \qquad \downarrow \text{4079} \\
 & \frac{\int \frac{\sqrt{i\tan(c+dx)a+a}(a^3(317A+67iB)-2a^3(151iA-41B)\tan(c+dx))}{2\tan^{\frac{3}{2}}(c+dx)} dx}{6a^2} + \frac{a^2(151A+41iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}} + \frac{a(17A+7iB)}{3d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{3/2}} + \\
 & \qquad \qquad \qquad \frac{10a^2}{A+iB} \\
 & \qquad \qquad \qquad \frac{5d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{5/2}} \\
 & \qquad \qquad \qquad \downarrow \text{27}
 \end{aligned}$$

3.194. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}} dx$

$$\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(317A+67iB)-2a^3(151iA-41B) \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx}{6a^2} + \frac{a^2(151A+41iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} + \frac{a(17A+7iB)}{3d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}} + \frac{A+iB}{5d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}}$$

↓ 3042

$$\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(317A+67iB)-2a^3(151iA-41B) \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx}{6a^2} + \frac{a^2(151A+41iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} + \frac{a(17A+7iB)}{3d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}} + \frac{A+iB}{5d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}}$$

↓ 4081

$$\frac{2 \int \frac{15a^4(iA+B)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx - \frac{2a^3(317A+67iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{6a^2} + \frac{a^2(151A+41iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} + \frac{a(17A+7iB)}{3d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}} + \frac{A+iB}{5d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}}$$

↓ 27

$$\frac{15a^3(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(317A+67iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{6a^2} + \frac{a^2(151A+41iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} + \frac{a(17A+7iB)}{3d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}} + \frac{A+iB}{5d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}}$$

↓ 3042

$$\frac{15a^3(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(317A+67iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{6a^2} + \frac{a^2(151A+41iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} + \frac{a(17A+7iB)}{3d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{3/2}} + \frac{A+iB}{5d\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}}$$

↓ 4027

3.194. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{30ia^5(B+iA) \int \frac{1}{\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{2a^2} - \frac{2a^3(317A+67iB)\sqrt{a+ia \tan(c+dx)}}{6a^2 d \sqrt{\tan(c+dx)}} + \frac{a^2(151A+41iB)}{d \sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} + \frac{a(17A+7iB)}{3d \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))} \\
 & \frac{A+iB}{5d \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow \text{218} \\
 & \frac{\frac{a^2(151A+41iB)}{d \sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} + \frac{(15-15i)a^{7/2}(B+iA) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^3(317A+67iB)\sqrt{a+ia \tan(c+dx)}}{6a^2 d \sqrt{\tan(c+dx)}}}{6a^2} + \frac{a(17A+7iB)}{3d \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))} \\
 & \frac{A+iB}{5d \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^{5/2}}
 \end{aligned}$$

```
input Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(5/2)),x]
```

```
output (A + I*B)/(5*d*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(5/2)) + ((a*(17*A + (7*I)*B))/(3*d*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(3/2)) + ((a^2*(151*A + (41*I)*B))/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + (((15 - 15*I)*a^(7/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a^3*(317*A + (67*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(2*a^2)/(6*a^2)/(10*a^2))
```

3.194.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

3.194. $\int \frac{A+B \tan(c+dx)}{\tan^3(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$

```
rule 4027 Int[Sqrt[(a_) + (b_.)*tan[(e_) + (f_.)*(x_)]]/Sqrt[(c_) + (d_.)*tan[(e_)
+ (f_.)*(x_)], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4079 Int[((a_) + (b_.)*tan[(e_) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

```
rule 4081 Int[((a_) + (b_.)*tan[(e_) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

3.194.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1147 vs. $2(194) = 388$.

Time = 0.17 (sec) , antiderivative size = 1148, normalized size of antiderivative = 4.78

method	result	size
derivativedivides	Expression too large to display	1148
default	Expression too large to display	1148
parts	Expression too large to display	1188

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(5/2),x,method=_R
ETURNVERBOSE)
```

$$3.194. \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$$


```
output -1/240/d*(a*(1+I*tan(d*x+c)))^(1/2)/a^3*(15*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*
a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*
x+c)+I))*a*tan(d*x+c)+1268*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(
1/2)*tan(d*x+c)^4-4468*I*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(
1/2)*tan(d*x+c)^3-15*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1
+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^5+9
08*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3-60*I*
A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2
-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2-90*I*B*2^(1/2)*ln((2*2
^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+
c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3+60*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)
*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))
*a*tan(d*x+c)^4-5660*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*
tan(d*x+c)^2-1060*I*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*t
an(d*x+c)^2+15*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*t
an(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^5+90*A*
ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*t
an(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^3+60*I*A*2^(1/2)*ln((2*2^(
1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-I*a+3*a*tan(d*x+c)
)/(tan(d*x+c)+I))*a*tan(d*x+c)^4+268*I*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+...
```

3.194.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 529 vs. 2(182) = 364.

Time = 0.29 (sec) , antiderivative size = 529, normalized size of antiderivative = 2.20

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \frac{15 \sqrt{\frac{1}{2}}(a^3 de^{(7i dx + 7i c)} - a^3 de^{(5i dx + 5i c)}) \sqrt{\frac{iA^2 + 2AB - iB^2}{a^5 d^2}} \log \left(\frac{2 \sqrt{\dots}}{\dots} \right)}{\dots}$$

```
input integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(5/2),x, al
gorithm="fricas")
```

```
output 1/120*(15*sqrt(1/2)*(a^3*d*e^(7*I*d*x + 7*I*c) - a^3*d*e^(5*I*d*x + 5*I*c)
)*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2))*log((2*sqrt(1/2)*a^3*d*sqrt((I*A
^2 + 2*A*B - I*B^2)/(a^5*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I
*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I
*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) - 15*sqrt(1/
2)*(a^3*d*e^(7*I*d*x + 7*I*c) - a^3*d*e^(5*I*d*x + 5*I*c))*sqrt((I*A^2 + 2
*A*B - I*B^2)/(a^5*d^2))*log(-(2*sqrt(1/2)*a^3*d*sqrt((I*A^2 + 2*A*B - I*B
^2)/(a^5*d^2))*e^(I*d*x + I*c) - sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) +
I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) +
I)/(e^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) + sqrt(2)*((-463*I*A + 83*B)
*e^(8*I*d*x + 8*I*c) + (-269*I*A + 19*B)*e^(6*I*d*x + 6*I*c) - 20*(-11*I*A
+ 4*B)*e^(4*I*d*x + 4*I*c) + (29*I*A - 19*B)*e^(2*I*d*x + 2*I*c) + 3*I*A
- 3*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)
/(e^(2*I*d*x + 2*I*c) + 1)))/(a^3*d*e^(7*I*d*x + 7*I*c) - a^3*d*e^(5*I*d*x
+ 5*I*c))
```

3.194.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

```
input integrate((A+B*tan(d*x+c))/tan(d*x+c)**(3/2)/(a+I*a*tan(d*x+c))**(5/2),x)
```

```
output Timed out
```

3.194.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(5/2),x, al
gorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un
defined.
```

3.194. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$

3.194.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Timed out`

3.194.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2} (a + a \tan(c + dx) 1i)^{5/2}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + a*tan(c + d*x)*1i)^(5/2)),x)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + a*tan(c + d*x)*1i)^(5/2)), x)`

3.195
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$$

3.195.1 Optimal result	1997
3.195.2 Mathematica [A] (verified)	1998
3.195.3 Rubi [A] (verified)	1998
3.195.4 Maple [B] (verified)	2003
3.195.5 Fracas [B] (verification not implemented)	2004
3.195.6 Sympy [F(-1)]	2005
3.195.7 Maxima [F(-2)]	2005
3.195.8 Giac [F(-2)]	2006
3.195.9 Mupad [F(-1)]	2006

3.195.1 Optimal result

Integrand size = 38, antiderivative size = 286

$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx = \frac{(\frac{1}{8} + \frac{i}{8})(iA+B) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{a^{5/2}d}$$

$$+ \frac{A+iB}{5d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} + \frac{21A+11iB}{30ad \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}}$$

$$+ \frac{89A+39iB}{20a^2d \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}}$$

$$- \frac{(361A+151iB)\sqrt{a+ia \tan(c+dx)}}{60a^3d \tan^{\frac{3}{2}}(c+dx)} + \frac{(707iA-317B)\sqrt{a+ia \tan(c+dx)}}{60a^3d\sqrt{\tan(c+dx)}}$$

output

```
(1/8+1/8*I)*(I*A+B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))/a^(5/2)/d+1/60*(707*I*A-317*B)*(a+I*a*tan(d*x+c))^(1/2)/a^3/d/tan(d*x+c)^(1/2)+1/20*(89*A+39*I*B)/a^2/d/(a+I*a*tan(d*x+c))^(1/2)/tan(d*x+c)^(3/2)-1/60*(361*A+151*I*B)*(a+I*a*tan(d*x+c))^(1/2)/a^3/d/tan(d*x+c)^(3/2)+1/5*(A+I*B)/d/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(5/2)+1/30*(21*A+11*I*B)/a/d/tan(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(3/2)
```

3.195.2 Mathematica [A] (verified)

Time = 7.77 (sec) , antiderivative size = 259, normalized size of antiderivative = 0.91

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx =$$

$$i \sec^2(c + dx) \left(\frac{15\sqrt{2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)(\cos(2(c+dx))+i \sin(2(c+dx))) \tan^2(c+dx)}{\sqrt{ia \tan(c+dx)}} + \frac{\sec^2(c+dx)(-174iA+84B+(7$$

$$120a^2d \tan^{\frac{3}{2}}(c + dx)$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^(5/2)),x]`

output `((-1/120*I)*Sec[c + d*x]^2*((15*Sqrt[2]*(I*A + B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)])*Tan[c + d*x]^2)/Sqrt[I*a*Tan[c + d*x]] + (Sec[c + d*x]^2*((-174*I)*A + 84*B + ((747*I)*A - 317*B)*Cos[2*(c + d*x)] + ((-493*I)*A + 233*B)*Cos[4*(c + d*x)] - 780*A*Sin[2*(c + d*x)] - (340*I)*B*Sin[2*(c + d*x)] + 490*A*Sin[4*(c + d*x)] + (230*I)*B*Sin[4*(c + d*x)]))/Sqrt[a + I*a*Tan[c + d*x]]))/(a^2*d*Tan[c + d*x]^(3/2)*(-I + Tan[c + d*x])^2)`

3.195.3 Rubi [A] (verified)

Time = 1.75 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.07, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.474$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2}(a + ia \tan(c + dx))^{5/2}} dx$$

↓ 4079

$$\begin{aligned}
& \frac{\int \frac{a(13A+3iB)-8a(iA-B)\tan(c+dx)}{2\tan^{\frac{5}{2}}(c+dx)(i\tan(c+dx)a+a)^{3/2}} dx}{5a^2} + \frac{A+iB}{5d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{a(13A+3iB)-8a(iA-B)\tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(i\tan(c+dx)a+a)^{3/2}} dx}{10a^2} + \frac{A+iB}{5d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{a(13A+3iB)-8a(iA-B)\tan(c+dx)}{\tan(c+dx)^{5/2}(i\tan(c+dx)a+a)^{3/2}} dx}{10a^2} + \frac{A+iB}{5d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}} \\
& \quad \downarrow 4079 \\
& \frac{\int \frac{3(a^2(47A+17iB)-2a^2(21iA-11B)\tan(c+dx))}{2\tan^{\frac{5}{2}}(c+dx)\sqrt{i\tan(c+dx)a+a}} dx}{3a^2} + \frac{a(21A+11iB)}{3d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{3/2}} + \\
& \quad \frac{10a^2}{A+iB} \\
& \quad \frac{5d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}}{\downarrow 27} \\
& \frac{\int \frac{a^2(47A+17iB)-2a^2(21iA-11B)\tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)\sqrt{i\tan(c+dx)a+a}} dx}{2a^2} + \frac{a(21A+11iB)}{3d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{3/2}} + \\
& \quad \frac{10a^2}{A+iB} \\
& \quad \frac{5d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}}{\downarrow 3042} \\
& \frac{\int \frac{a^2(47A+17iB)-2a^2(21iA-11B)\tan(c+dx)}{\tan(c+dx)^{5/2}\sqrt{i\tan(c+dx)a+a}} dx}{2a^2} + \frac{a(21A+11iB)}{3d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{3/2}} + \\
& \quad \frac{10a^2}{A+iB} \\
& \quad \frac{5d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}}{\downarrow 4079} \\
& \frac{\int \frac{\sqrt{i\tan(c+dx)a+a}(a^3(361A+151iB)-4a^3(89iA-39B)\tan(c+dx))}{2\tan^{\frac{5}{2}}(c+dx)} dx}{a^2} + \frac{a^2(89A+39iB)}{d\tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}} + \frac{a(21A+11iB)}{3d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{3/2}} + \\
& \quad \frac{10a^2}{A+iB} \\
& \quad \frac{5d\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}}
\end{aligned}$$

3.195. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}} dx$

$$\int \frac{\sqrt{i \tan(c+dx)a+a} \left(a^3(361A+151iB) - 4a^3(89iA-39B) \tan(c+dx) \right) dx}{\tan^{\frac{5}{2}}(c+dx)} + \frac{a^2(89A+39iB)}{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} + \frac{a(21A+11iB)}{3d \tan^{\frac{3}{2}}(c+dx) (a+ia \tan(c+dx))^{3/2}} + \frac{A+iB}{5d \tan^{\frac{3}{2}}(c+dx) (a+ia \tan(c+dx))^{5/2}}$$

27

$$\int \frac{\sqrt{i \tan(c+dx)a+a} \left(a^3(361A+151iB) - 4a^3(89iA-39B) \tan(c+dx) \right) dx}{\tan^{\frac{5}{2}}(c+dx)} + \frac{a^2(89A+39iB)}{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} + \frac{a(21A+11iB)}{3d \tan^{\frac{3}{2}}(c+dx) (a+ia \tan(c+dx))^{3/2}} + \frac{A+iB}{5d \tan^{\frac{3}{2}}(c+dx) (a+ia \tan(c+dx))^{5/2}}$$

3042

$$2 \int - \frac{\sqrt{i \tan(c+dx)a+a} \left((707iA-317B)a^4 + 2(361A+151iB) \tan(c+dx)a^4 \right) dx}{2 \tan^{\frac{3}{2}}(c+dx)} - \frac{2a^3(361A+151iB) \sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{a^2(89A+39iB)}{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} + \frac{A+iB}{5d \tan^{\frac{3}{2}}(c+dx) (a+ia \tan(c+dx))^{5/2}}$$

4081

$$\int \frac{\sqrt{i \tan(c+dx)a+a} \left((707iA-317B)a^4 + 2(361A+151iB) \tan(c+dx)a^4 \right) dx}{\tan^{\frac{3}{2}}(c+dx)} - \frac{2a^3(361A+151iB) \sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{a^2(89A+39iB)}{d \tan^{\frac{3}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}} + \frac{A+iB}{5d \tan^{\frac{3}{2}}(c+dx) (a+ia \tan(c+dx))^{5/2}}$$

27

3042

3.195. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx) (a+ia \tan(c+dx))^{5/2}} dx$

$$\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} \left((707iA-317B)a^4+2(361A+151iB) \tan(c+dx)a^4 \right) dx}{\tan(c+dx)^{3/2}} - \frac{2a^3(361A+151iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{a^2(89A+39iB)}{d \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}}}{2a^2} + \frac{\dots}{3d \tan^{\frac{3}{2}}(c+dx)}$$

$$\frac{A+iB}{5d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}}$$

↓ 4081

$$\frac{2 \int \frac{15a^5(A-iB)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx - \frac{2a^4(-317B+707iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^3(361A+151iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{a^2(89A+39iB)}{d \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}}}{2a^2} + \frac{\dots}{3d \tan^{\frac{3}{2}}(c+dx)}$$

$$\frac{A+iB}{5d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}}$$

↓ 27

$$\frac{15a^4(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^4(-317B+707iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^3(361A+151iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{a^2(89A+39iB)}{d \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}}}{2a^2} + \frac{\dots}{3d \tan^{\frac{3}{2}}(c+dx)}$$

$$\frac{A+iB}{5d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}}$$

↓ 3042

$$\frac{15a^4(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^4(-317B+707iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^3(361A+151iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{a^2(89A+39iB)}{d \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}}}{2a^2} + \frac{\dots}{3d \tan^{\frac{3}{2}}(c+dx)}$$

$$\frac{A+iB}{5d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}}$$

↓ 4027

3.195. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{30ia^6(A-iB) \int \frac{1}{\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{3a} - \frac{2a^4(-317B+707iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a^3(361A+151iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{a^2(89A+iB)}{d \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}} \\
 & \frac{A+iB}{5d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} \\
 & \quad \downarrow \text{218} \\
 & \frac{a^2(89A+39iB)}{d \tan^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}} + \frac{-2a^3(361A+151iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{(15-15i)a^{9/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^4(-317B+707iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
 & \frac{A+iB}{5d \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}}
 \end{aligned}$$

```
input Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^(5/2)),x]
```

```
output (A + I*B)/(5*d*Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(5/2)) + ((a*(21*A + (11*I)*B))/(3*d*Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(3/2)) + ((a^2*(89*A + (39*I)*B))/(d*Tan[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]]) + ((-2*a^3*(361*A + (151*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (((15 - 15*I)*a^(9/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d - (2*a^4*((707*I)*A - 317*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a))/(2*a^2))/(10*a^2)
```

3.195.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

$$3.195. \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

3.195.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1228 vs. $2(232) = 464$.

Time = 0.17 (sec) , antiderivative size = 1229, normalized size of antiderivative = 4.30

method	result	size
derivativedivides	Expression too large to display	1229
default	Expression too large to display	1229
parts	Expression too large to display	1273

3.195.
$$\int \frac{A+B \tan (c+d x)}{\tan ^2(c+d x)\left(a+i a \tan (c+d x)\right)^{5 / 2}} d x$$

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^(5/2),x,method=_R
ETURNVERBOSE)`

output $\frac{1}{240}d*(a*(1+I*\tan(d*x+c)))^{(1/2)}*(15*I*A*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*a*\tan(d*x+c)^6-12260*I*A*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)^3+15*B*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*a*\tan(d*x+c)^6+15*I*A*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*a*\tan(d*x+c)^2+60*I*B*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*a*\tan(d*x+c)^3+60*A*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*a*\tan(d*x+c)^5-2940*I*B*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)^2-60*I*B*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*a*\tan(d*x+c)^5-90*B*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*a*\tan(d*x+c)^4-1268*B*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)^5+2828*I*A*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)^5-90*I*A*2^{(1/2)}*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*a*\tan(d*x+c)^4-60*A*\ln((2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-I*a+3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*2^...$

3.195.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 588 vs. $2(218) = 436$.

Time = 0.29 (sec) , antiderivative size = 588, normalized size of antiderivative = 2.06

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx =$$

$$15 \sqrt{\frac{1}{2}} (a^3 de^{(9i dx + 9i c)} - 2 a^3 de^{(7i dx + 7i c)} + a^3 de^{(5i dx + 5i c)}) \sqrt{\frac{-i A^2 - 2 AB + i B^2}{a^5 d^2}} \log \left(\frac{2i \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{-i A^2 - 2 AB + i B^2}{a^5 d^2}} e^{(i dx + \dots)}}{\dots} \right)$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^(5/2),x,algorithm="fricas")`

3.195. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$

output

```
-1/120*(15*sqrt(1/2)*(a^3*d*e^(9*I*d*x + 9*I*c) - 2*a^3*d*e^(7*I*d*x + 7*I
*c) + a^3*d*e^(5*I*d*x + 5*I*c))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2))*
log((2*I*sqrt(1/2)*a^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2))*e^(I*d*x
+ I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A + B)*sqrt(a/(e^(2*I
*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c)
+ 1)))/(4*I*A + 4*B)) - 15*sqrt(1/2)*(a^3*d*e^(9*I*d*x + 9*I*c) - 2*a^3*d
*e^(7*I*d*x + 7*I*c) + a^3*d*e^(5*I*d*x + 5*I*c))*sqrt((-I*A^2 - 2*A*B + I
*B^2)/(a^5*d^2))*log((-2*I*sqrt(1/2)*a^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(
a^5*d^2))*e^(I*d*x + I*c) + sqrt(2)*((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A +
B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e
^(2*I*d*x + 2*I*c) + 1)))/(4*I*A + 4*B)) + sqrt(2)*((983*A + 463*I*B)*e^(1
0*I*d*x + 10*I*c) - 2*(272*A + 97*I*B)*e^(8*I*d*x + 8*I*c) - 3*(393*A + 16
3*I*B)*e^(6*I*d*x + 6*I*c) + (381*A + 191*I*B)*e^(4*I*d*x + 4*I*c) + 2*(18
*A + 13*I*B)*e^(2*I*d*x + 2*I*c) + 3*A + 3*I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c)
+ 1))*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1)))/(a^3
*d*e^(9*I*d*x + 9*I*c) - 2*a^3*d*e^(7*I*d*x + 7*I*c) + a^3*d*e^(5*I*d*x +
5*I*c))
```

3.195.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(5/2)/(a+I*a*tan(d*x+c))**(5/2),x)`

output `Timed out`

3.195.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")`

3.195. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$

output Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.

3.195.8 Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Not invertible Error: Bad Argument Value

3.195.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2} (a + a \tan(c + dx) 1i)^{5/2}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + a*tan(c + d*x)*1i)^(5/2)),x)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + a*tan(c + d*x)*1i)^(5/2)), x)`

3.196 $\int \sqrt[3]{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$

3.196.1 Optimal result	2007
3.196.2 Mathematica [A] (verified)	2008
3.196.3 Rubi [A] (warning: unable to verify)	2008
3.196.4 Maple [A] (verified)	2011
3.196.5 Fricas [B] (verification not implemented)	2012
3.196.6 Sympy [F]	2013
3.196.7 Maxima [A] (verification not implemented)	2013
3.196.8 Giac [F]	2014
3.196.9 Mupad [B] (verification not implemented)	2014

3.196.1 Optimal result

Integrand size = 28, antiderivative size = 201

$$\int \sqrt[3]{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$= -\frac{\sqrt[3]{a}(A - iB)x}{2 \cdot 2^{2/3}} - \frac{\sqrt{3}\sqrt[3]{a}(iA + B) \arctan\left(\frac{\sqrt[3]{a+2^{2/3}}\sqrt[3]{a + ia \tan(c + dx)}}{\sqrt{3}\sqrt[3]{a}}\right)}{2^{2/3}d}$$

$$+ \frac{\sqrt[3]{a}(iA + B) \log(\cos(c + dx))}{2 \cdot 2^{2/3}d}$$

$$+ \frac{3\sqrt[3]{a}(iA + B) \log\left(\sqrt[3]{2}\sqrt[3]{a} - \sqrt[3]{a + ia \tan(c + dx)}\right)}{2 \cdot 2^{2/3}d} + \frac{3B\sqrt[3]{a + ia \tan(c + dx)}}{d}$$

output

```
-1/4*a^(1/3)*(A-I*B)*x*2^(1/3)+1/4*a^(1/3)*(I*A+B)*ln(cos(d*x+c))*2^(1/3)/
d+3/4*a^(1/3)*(I*A+B)*ln(2^(1/3)*a^(1/3)-(a+I*a*tan(d*x+c))^(1/3))*2^(1/3)
/d-1/2*a^(1/3)*(I*A+B)*arctan(1/3*(a^(1/3)+2^(2/3)*(a+I*a*tan(d*x+c))^(1/3)
))/a^(1/3)*3^(1/2))*3^(1/2)*2^(1/3)/d+3*B*(a+I*a*tan(d*x+c))^(1/3)/d
```

3.196.2 Mathematica [A] (verified)

Time = 0.81 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.93

$$\int \sqrt[3]{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$= -i\sqrt[3]{2}\sqrt[3]{a}(A - iB) \left(2\sqrt{3} \arctan \left(\frac{1 + 2^{2/3} \sqrt[3]{a + ia \tan(c + dx)}}{\sqrt[3]{a}} \right) - 2 \log \left(\sqrt[3]{2}\sqrt[3]{a} - \sqrt[3]{a + ia \tan(c + dx)} \right) \right)$$

input `Integrate[(a + I*a*Tan[c + d*x])^(1/3)*(A + B*Tan[c + d*x]),x]`

output `((-I)*2^(1/3)*a^(1/3)*(A - I*B)*(2*Sqrt[3]*ArcTan[(1 + (2^(2/3)*(a + I*a*Tan[c + d*x])^(1/3))/a^(1/3)]/Sqrt[3]) - 2*Log[2^(1/3)*a^(1/3) - (a + I*a*Tan[c + d*x])^(1/3)] + Log[2^(2/3)*a^(2/3) + 2^(1/3)*a^(1/3)*(a + I*a*Tan[c + d*x])^(1/3) + (a + I*a*Tan[c + d*x])^(2/3)]) + 12*B*(a + I*a*Tan[c + d*x])^(1/3))/(4*d)`

3.196.3 Rubi [A] (warning: unable to verify)

Time = 0.40 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.69, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4010, 3042, 3962, 69, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt[3]{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt[3]{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4010}$$

$$(A - iB) \int \sqrt[3]{i \tan(c + dx)a + adx} + \frac{3B \sqrt[3]{a + ia \tan(c + dx)}}{d}$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
& (A - iB) \int \sqrt[3]{i \tan(c + dx)a + adx} + \frac{3B \sqrt[3]{a + ia \tan(c + dx)}}{d} \\
& \quad \downarrow \text{3962} \\
& \frac{3B \sqrt[3]{a + ia \tan(c + dx)}}{d} - \frac{ia(A - iB) \int \frac{1}{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^{2/3}} d(ia \tan(c + dx))}{d} \\
& \quad \downarrow \text{69} \\
& \frac{3B \sqrt[3]{a + ia \tan(c + dx)}}{d} - \\
& \frac{ia(A - iB) \left(\frac{3 \int \frac{1}{\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)}}}{2^{2^{2/3} a^{2/3}}} d \sqrt[3]{i \tan(c + dx)a + a} + \frac{3 \int \frac{1}{-a^2 \tan^2(c + dx) + i \sqrt[3]{2} a^{4/3} \tan(c + dx) + 2^{2/3} a^{2/3}}}{2 \sqrt[3]{2} \sqrt[3]{a}} d \sqrt[3]{i \tan(c + dx)a + a} \right)}{d} \\
& \quad \downarrow \text{16} \\
& \frac{3B \sqrt[3]{a + ia \tan(c + dx)}}{d} - \\
& \frac{ia(A - iB) \left(\frac{3 \int \frac{1}{-a^2 \tan^2(c + dx) + i \sqrt[3]{2} a^{4/3} \tan(c + dx) + 2^{2/3} a^{2/3}}}{2 \sqrt[3]{2} \sqrt[3]{a}} d \sqrt[3]{i \tan(c + dx)a + a} - \frac{3 \log\left(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)}\right)}{2^{2^{2/3} a^{2/3}}} + \frac{\log(a - ia \tan(c + dx))}{2^{2^2}} \right)}{d} \\
& \quad \downarrow \text{1082} \\
& \frac{3B \sqrt[3]{a + ia \tan(c + dx)}}{d} - \\
& \frac{ia(A - iB) \left(-\frac{3 \int \frac{1}{a^2 \tan^2(c + dx) - 3(i 2^{2/3} a^{2/3} \tan(c + dx) + 1)}}{2^{2^{2/3} a^{2/3}}} - \frac{3 \log\left(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)}\right)}{2^{2^{2/3} a^{2/3}}} + \frac{\log(a - ia \tan(c + dx))}{2^{2^{2/3} a^{2/3}}} \right)}{d} \\
& \quad \downarrow \text{217} \\
& \frac{3B \sqrt[3]{a + ia \tan(c + dx)}}{d} - \\
& \frac{ia(A - iB) \left(\frac{i \sqrt{3} \operatorname{arctanh}\left(\frac{a \tan(c + dx)}{\sqrt{3}}\right)}{2^{2^{2/3} a^{2/3}}} - \frac{3 \log\left(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)}\right)}{2^{2^{2/3} a^{2/3}}} + \frac{\log(a - ia \tan(c + dx))}{2^{2^{2/3} a^{2/3}}} \right)}{d}
\end{aligned}$$

input `Int[(a + I*a*Tan[c + d*x])^(1/3)*(A + B*Tan[c + d*x]),x]`


```
output ((-I)*a*(A - I*B)*((I*Sqrt[3]*ArcTanh[(a*Tan[c + d*x])/Sqrt[3]])/(2^(2/3)*
a^(2/3)) - (3*Log[2^(1/3)*a^(1/3) - I*a*Tan[c + d*x]])/(2*2^(2/3)*a^(2/3))
+ Log[a - I*a*Tan[c + d*x]]/(2*2^(2/3)*a^(2/3)))/d + (3*B*(a + I*a*Tan[c
+ d*x])^(1/3))/d
```

3.196.3.1 Defintions of rubi rules used

```
rule 16 Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a +
b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

```
rule 69 Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(2/3)), x_Symbol] := With[
{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q^2),
x] + (-Simp[3/(2*b*q) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1
/3)], x] - Simp[3/(2*b*q^2) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)],
x])] /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3962 Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[-b/d S
ubst[Int[(a + x)^(n - 1)/(a - x), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b
, c, d, n}, x] && EqQ[a^2 + b^2, 0]
```

```
rule 4010 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]
```

3.196.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.79

method	result
derivativedivides	$3i \left(-iB(a+ia \tan(dx+c))^{\frac{1}{3}} + \frac{2^{\frac{1}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}} \right)}{6a^{\frac{2}{3}}} - \frac{2^{\frac{1}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}} (a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}} \right)}{12a^{\frac{2}{3}}} \right) / d$
default	$3i \left(-iB(a+ia \tan(dx+c))^{\frac{1}{3}} + \frac{2^{\frac{1}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}} \right)}{6a^{\frac{2}{3}}} - \frac{2^{\frac{1}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}} (a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}} \right)}{12a^{\frac{2}{3}}} \right) / d$
parts	$3iAa \left(\frac{2^{\frac{1}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}} \right)}{6a^{\frac{2}{3}}} - \frac{2^{\frac{1}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}} (a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}} \right)}{12a^{\frac{2}{3}}} - \frac{2^{\frac{1}{3}} \sqrt{3} \arctan \left(\frac{\sqrt{3} \left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}} \right)}{2 \left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}} (a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}} \right)} \right)}{12a^{\frac{2}{3}}} \right) / d$

```
input int((a+I*a*tan(d*x+c))^(1/3)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

output $3*I/d*(-I*B*(a+I*a*\tan(d*x+c))^{1/3}+(1/6*2^{1/3}/a^{2/3}*\ln((a+I*a*\tan(d*x+c))^{1/3}-2^{1/3}*a^{1/3}))-1/12*2^{1/3}/a^{2/3}*\ln((a+I*a*\tan(d*x+c))^{2/3}+2^{1/3}*a^{1/3}*(a+I*a*\tan(d*x+c))^{1/3}+2^{2/3}*a^{2/3}))-1/6*2^{1/3}/a^{2/3}*3^{1/2}*\arctan(1/3*3^{1/2}*(2^{2/3}/a^{1/3}*(a+I*a*\tan(d*x+c))^{1/3}+1)))a*(A-I*B)$

3.196.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 405 vs. $2(146) = 292$.

Time = 0.24 (sec) , antiderivative size = 405, normalized size of antiderivative = 2.01

$$\int \sqrt[3]{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$6 \cdot 2^{\frac{1}{3}} B \left(\frac{a}{e^{(2i dx + 2i c)} + 1} \right)^{\frac{1}{3}} e^{\left(\frac{2}{3}i dx + \frac{2}{3}i c\right)} + \left(\frac{1}{4}\right)^{\frac{1}{3}} (-i\sqrt{3}d - d) \left(\frac{(-iA^3 - 3A^2B + 3iAB^2 + B^3)a}{d^3} \right)^{\frac{1}{3}} \log \left(\frac{2^{\frac{1}{3}}(iA+B) \left(\frac{a}{e^{(2i dx + 2i c)} + 1} \right)^{\frac{1}{3}}}{\dots} \right)$$

input `integrate((a+I*a*tan(d*x+c))^(1/3)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output $1/2*(6*2^{1/3}*B*(a/(e^{(2*I*d*x + 2*I*c)} + 1))^{1/3}*e^{(2/3*I*d*x + 2/3*I*c)} + (1/4)^{1/3}*(-I*\sqrt{3}*d - d)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)*a/d^3)^{1/3}*\log((2^{1/3}*(I*A + B)*(a/(e^{(2*I*d*x + 2*I*c)} + 1))^{1/3}*e^{(2/3*I*d*x + 2/3*I*c)} + (1/4)^{1/3}*(I*\sqrt{3}*d + d)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)*a/d^3)^{1/3})/(I*A + B)) + (1/4)^{1/3}*(I*\sqrt{3}*d - d)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)*a/d^3)^{1/3}*\log((2^{1/3}*(I*A + B)*(a/(e^{(2*I*d*x + 2*I*c)} + 1))^{1/3}*e^{(2/3*I*d*x + 2/3*I*c)} + (1/4)^{1/3}*(-I*\sqrt{3}*d + d)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)*a/d^3)^{1/3})/(I*A + B)) + 2*(1/4)^{1/3}*d*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)*a/d^3)^{1/3}*\log((2^{1/3}*(I*A + B)*(a/(e^{(2*I*d*x + 2*I*c)} + 1))^{1/3}*e^{(2/3*I*d*x + 2/3*I*c)} - 2*(1/4)^{1/3}*d*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)*a/d^3)^{1/3})/(I*A + B)))/d$

3.196.6 Sympy [F]

$$\int \sqrt[3]{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$= \int \sqrt[3]{ia (\tan(c + dx) - i)}(A + B \tan(c + dx)) dx$$

input `integrate((a+I*a*tan(d*x+c))**(1/3)*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(1/3)*(A + B*tan(c + d*x)), x)`

3.196.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.83

$$\int \sqrt[3]{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx =$$

$$i \left(2 \sqrt[3]{32} \frac{1}{3} (A - iB) a^{\frac{4}{3}} \arctan \left(\frac{\sqrt[3]{32} \left(2^{\frac{1}{3}} a^{\frac{1}{3}} + 2(i a \tan(dx+c) + a)^{\frac{1}{3}} \right)}{6 a^{\frac{1}{3}}} \right) \right) + 2^{\frac{1}{3}} (A - iB) a^{\frac{4}{3}} \log \left(2^{\frac{2}{3}} a^{\frac{2}{3}} + 2^{\frac{1}{3}} (i a \tan(dx+c) + a)^{\frac{1}{3}} \right)$$

input `integrate((a+I*a*tan(d*x+c))^(1/3)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/4*I*(2*sqrt(3)*2^(1/3)*(A - I*B)*a^(4/3)*arctan(1/6*sqrt(3)*2^(2/3)*(2^(1/3)*a^(1/3) + 2*(I*a*tan(d*x + c) + a)^(1/3))/a^(1/3)) + 2^(1/3)*(A - I*B)*a^(4/3)*log(2^(2/3)*a^(2/3) + 2^(1/3)*(I*a*tan(d*x + c) + a)^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(2/3)) - 2*2^(1/3)*(A - I*B)*a^(4/3)*log(-2^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(1/3)) + 12*I*(I*a*tan(d*x + c) + a)^(1/3)*B*a)/(a*d)`

3.196.8 Giac [F]

$$\int \sqrt[3]{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{\frac{1}{3}} dx$$

input `integrate((a+I*a*tan(d*x+c))^(1/3)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(1/3), x)`

3.196.9 Mupad [B] (verification not implemented)

Time = 1.25 (sec) , antiderivative size = 365, normalized size of antiderivative = 1.82

$$\int \sqrt[3]{a + ia \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$= \frac{3 B (a + a \tan(c + dx) i)^{1/3}}{d} + \frac{2^{1/3} B a^{1/3} \ln \left((a (1 + \tan(c + dx) i))^{1/3} - 2^{1/3} a^{1/3} \right)}{2 d}$$

$$- \frac{\left(\frac{1}{4}i\right)^{1/3} A a^{1/3} \ln \left(A a d^2 (a + a \tan(c + dx) i)^{1/3} 9i + 18 \left(\frac{1}{4}i\right)^{1/3} A a^{4/3} d^2 \right)}{d}$$

$$- \frac{\left(\frac{1}{4}i\right)^{1/3} A a^{1/3} \ln \left(A a d^2 (a + a \tan(c + dx) i)^{1/3} 9i + 18 \left(\frac{1}{4}i\right)^{1/3} A a^{4/3} d^2 \left(-\frac{1}{2} + \frac{\sqrt{3}i}{2}\right) \right) \left(-\frac{1}{2} + \frac{\sqrt{3}i}{2}\right)}{d}$$

$$+ \frac{\left(\frac{1}{4}i\right)^{1/3} A a^{1/3} \ln \left(A a d^2 (a + a \tan(c + dx) i)^{1/3} 9i - 18 \left(\frac{1}{4}i\right)^{1/3} A a^{4/3} d^2 \left(\frac{1}{2} + \frac{\sqrt{3}i}{2}\right) \right) \left(\frac{1}{2} + \frac{\sqrt{3}i}{2}\right)}{d}$$

$$+ \frac{4^{2/3} B a^{1/3} \ln \left(\frac{9 B a (a + a \tan(c + dx) i)^{1/3}}{d} - \frac{9 2^{1/3} B a^{4/3} (-1 + \sqrt{3} i)}{2 d} \right) \left(-\frac{1}{2} + \frac{\sqrt{3}i}{2}\right)}{4 d}$$

$$- \frac{4^{2/3} B a^{1/3} \ln \left(\frac{9 B a (a + a \tan(c + dx) i)^{1/3}}{d} + \frac{9 2^{1/3} B a^{4/3} (1 + \sqrt{3} i)}{2 d} \right) \left(\frac{1}{2} + \frac{\sqrt{3}i}{2}\right)}{4 d}$$

input `int((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/3),x)`

output $(3*B*(a + a*\tan(c + d*x)*1i)^{(1/3)})/d + (2^{(1/3)}*B*a^{(1/3)}*\log((a*(\tan(c + d*x)*1i + 1))^{(1/3)} - 2^{(1/3)}*a^{(1/3)}))/(2*d) - ((1i/4)^{(1/3)}*A*a^{(1/3)}*\log(A*a*d^2*(a + a*\tan(c + d*x)*1i)^{(1/3)}*9i + 18*(1i/4)^{(1/3)}*A*a^{(4/3)}*d^2))/d - ((1i/4)^{(1/3)}*A*a^{(1/3)}*\log(A*a*d^2*(a + a*\tan(c + d*x)*1i)^{(1/3)}*9i + 18*(1i/4)^{(1/3)}*A*a^{(4/3)}*d^2*((3^{(1/2)}*1i)/2 - 1/2))*((3^{(1/2)}*1i)/2 - 1/2))/d + ((1i/4)^{(1/3)}*A*a^{(1/3)}*\log(A*a*d^2*(a + a*\tan(c + d*x)*1i)^{(1/3)}*9i - 18*(1i/4)^{(1/3)}*A*a^{(4/3)}*d^2*((3^{(1/2)}*1i)/2 + 1/2))*((3^{(1/2)}*1i)/2 + 1/2))/d + (4^{(2/3)}*B*a^{(1/3)}*\log((9*B*a*(a + a*\tan(c + d*x)*1i)^{(1/3)}))/d - (9*2^{(1/3)}*B*a^{(4/3)}*(3^{(1/2)}*1i - 1))/(2*d))*((3^{(1/2)}*1i)/2 - 1/2))/(4*d) - (4^{(2/3)}*B*a^{(1/3)}*\log((9*B*a*(a + a*\tan(c + d*x)*1i)^{(1/3)}))/d + (9*2^{(1/3)}*B*a^{(4/3)}*(3^{(1/2)}*1i + 1))/(2*d))*((3^{(1/2)}*1i)/2 + 1/2))/(4*d)$

3.197 $\int \tan^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx$

3.197.1 Optimal result	2016
3.197.2 Mathematica [A] (verified)	2017
3.197.3 Rubi [A] (warning: unable to verify)	2017
3.197.4 Maple [A] (verified)	2021
3.197.5 Fricas [B] (verification not implemented)	2023
3.197.6 Sympy [F]	2024
3.197.7 Maxima [A] (verification not implemented)	2024
3.197.8 Giac [F]	2025
3.197.9 Mupad [B] (verification not implemented)	2025

3.197.1 Optimal result

Integrand size = 36, antiderivative size = 270

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \frac{a^{2/3}(A - iB)x}{2\sqrt[3]{2}} - \frac{\sqrt{3}a^{2/3}(iA + B) \arctan\left(\frac{\sqrt[3]{a+2^{2/3}}\sqrt[3]{a + ia \tan(c + dx)}}{\sqrt{3}\sqrt[3]{a}}\right)}{\sqrt[3]{2}d} - \frac{a^{2/3}(iA + B) \log(\cos(c + dx))}{2\sqrt[3]{2}d} - \frac{3a^{2/3}(iA + B) \log\left(\sqrt[3]{2}\sqrt[3]{a} - \sqrt[3]{a + ia \tan(c + dx)}\right)}{2\sqrt[3]{2}d} - \frac{9B(a + ia \tan(c + dx))^{2/3}}{8d} + \frac{3B \tan^2(c + dx)(a + ia \tan(c + dx))^{2/3}}{8d} - \frac{3(4iA + B)(a + ia \tan(c + dx))^{5/3}}{20ad}$$

output

```
1/4*a^(2/3)*(A-I*B)*x*2^(2/3)-1/4*a^(2/3)*(I*A+B)*ln(cos(d*x+c))*2^(2/3)/d
-3/4*a^(2/3)*(I*A+B)*ln(2^(1/3)*a^(1/3)-(a+I*a*tan(d*x+c))^(1/3))*2^(2/3)/
d-1/2*a^(2/3)*(I*A+B)*arctan(1/3*(a^(1/3)+2^(2/3)*(a+I*a*tan(d*x+c))^(1/3)
)/a^(1/3)*3^(1/2))*3^(1/2)*2^(2/3)/d-9/8*B*(a+I*a*tan(d*x+c))^(2/3)/d+3/8*
B*tan(d*x+c)^2*(a+I*a*tan(d*x+c))^(2/3)/d-3/20*(4*I*A+B)*(a+I*a*tan(d*x+c)
)^(5/3)/a/d
```

3.197.2 Mathematica [A] (verified)

Time = 5.25 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.73

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx =$$

$$10 \cdot 2^{2/3} a^{2/3} (iA + B) \left(2\sqrt{3} \arctan \left(\frac{1 + \frac{2^{2/3} \sqrt[3]{a + ia \tan(c + dx)}}{\sqrt[3]{a}}}{\sqrt{3}} \right) - \log(i + \tan(c + dx)) + 3 \log \left(\sqrt[3]{2} \sqrt[3]{a} \right) \right)$$

input `Integrate[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^(2/3)*(A + B*Tan[c + d*x]),x]`

output `-1/40*(10*2^(2/3)*a^(2/3)*(I*A + B)*(2*Sqrt[3]*ArcTan[(1 + (2^(2/3)*(a + I*a*Tan[c + d*x])^(1/3))/a^(1/3)]/Sqrt[3]) - Log[I + Tan[c + d*x]] + 3*Log[2^(1/3)*a^(1/3) - (a + I*a*Tan[c + d*x])^(1/3])) + 45*B*(a + I*a*Tan[c + d*x])^(2/3) - 15*B*Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^(2/3) + (6*((4*I)*A + B)*(a + I*a*Tan[c + d*x])^(5/3))/a)/d`

3.197.3 Rubi [A] (warning: unable to verify)Time = 0.84 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.80, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.361$, Rules used = {3042, 4080, 27, 3042, 4075, 3042, 4010, 3042, 3962, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^2(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4080}$$

$$\frac{3 \int -\frac{2}{3} \tan(c + dx)(i \tan(c + dx)a + a)^{2/3}(3aB - a(4A - iB) \tan(c + dx)) dx}{8a} + \frac{3B \tan^2(c + dx)(a + ia \tan(c + dx))^{2/3}}{8d}$$

3.197. $\int \tan^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx$

$$\begin{array}{c}
\downarrow 27 \\
\frac{3B \tan^2(c+dx)(a+ia \tan(c+dx))^{2/3}}{8d} - \frac{\int \tan(c+dx)(i \tan(c+dx)a+a)^{2/3}(3aB-a(4A-iB) \tan(c+dx))dx}{4a} \\
\downarrow 3042 \\
\frac{3B \tan^2(c+dx)(a+ia \tan(c+dx))^{2/3}}{8d} - \frac{\int \tan(c+dx)(i \tan(c+dx)a+a)^{2/3}(3aB-a(4A-iB) \tan(c+dx))dx}{4a} \\
\downarrow 4075 \\
\frac{3B \tan^2(c+dx)(a+ia \tan(c+dx))^{2/3}}{8d} - \frac{\int (i \tan(c+dx)a+a)^{2/3}(a(4A-iB)+3aB \tan(c+dx))dx + \frac{3(B+4iA)(a+ia \tan(c+dx))^{5/3}}{5d}}{4a} \\
\downarrow 3042 \\
\frac{3B \tan^2(c+dx)(a+ia \tan(c+dx))^{2/3}}{8d} - \frac{\int (i \tan(c+dx)a+a)^{2/3}(a(4A-iB)+3aB \tan(c+dx))dx + \frac{3(B+4iA)(a+ia \tan(c+dx))^{5/3}}{5d}}{4a} \\
\downarrow 4010 \\
\frac{3B \tan^2(c+dx)(a+ia \tan(c+dx))^{2/3}}{8d} - \frac{4a(A-iB) \int (i \tan(c+dx)a+a)^{2/3}dx + \frac{3(B+4iA)(a+ia \tan(c+dx))^{5/3}}{5d} + \frac{9aB(a+ia \tan(c+dx))^{2/3}}{2d}}{4a} \\
\downarrow 3042 \\
\frac{3B \tan^2(c+dx)(a+ia \tan(c+dx))^{2/3}}{8d} - \frac{4a(A-iB) \int (i \tan(c+dx)a+a)^{2/3}dx + \frac{3(B+4iA)(a+ia \tan(c+dx))^{5/3}}{5d} + \frac{9aB(a+ia \tan(c+dx))^{2/3}}{2d}}{4a} \\
\downarrow 3962 \\
\frac{3B \tan^2(c+dx)(a+ia \tan(c+dx))^{2/3}}{8d} - \frac{4ia^2(A-iB) \int \frac{1}{(a-ia \tan(c+dx)) \sqrt[3]{i \tan(c+dx)a+a}} d(i \tan(c+dx)) + \frac{3(B+4iA)(a+ia \tan(c+dx))^{5/3}}{5d} + \frac{9aB(a+ia \tan(c+dx))^{2/3}}{2d}}{4a} \\
\downarrow 67
\end{array}$$

$$\frac{\frac{3B \tan^2(c+dx)(a+ia \tan(c+dx))^{2/3}}{8d} - \frac{4ia^2(A-iB) \left(-\frac{3}{2} \int \frac{1}{-a^2 \tan^2(c+dx)+i \sqrt[3]{2} a^{4/3} \tan(c+dx)+2^{2/3} a^{2/3}} d \sqrt[3]{i \tan(c+dx)} a + a - \frac{3 \int \frac{1}{\sqrt[3]{2} \sqrt[3]{a-ia \tan(c+dx)}} d \sqrt[3]{i \tan(c+dx)}}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d}}{4a}$$

↓ 16

$$\frac{\frac{3B \tan^2(c+dx)(a+ia \tan(c+dx))^{2/3}}{8d} - \frac{4ia^2(A-iB) \left(-\frac{3}{2} \int \frac{1}{-a^2 \tan^2(c+dx)+i \sqrt[3]{2} a^{4/3} \tan(c+dx)+2^{2/3} a^{2/3}} d \sqrt[3]{i \tan(c+dx)} a + a - \frac{3 \log\left(\sqrt[3]{2} \sqrt[3]{a-ia \tan(c+dx)}\right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \frac{\log(a-ia \tan(c+dx))}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d}}{4a}$$

↓ 1082

$$\frac{\frac{3B \tan^2(c+dx)(a+ia \tan(c+dx))^{2/3}}{8d} - \frac{4ia^2(A-iB) \left(\frac{3 \int \frac{1}{a^2 \tan^2(c+dx)-3} d(i^{2/3} a^{2/3} \tan(c+dx)+1)}{\sqrt[3]{2} \sqrt[3]{a}} - \frac{3 \log\left(\sqrt[3]{2} \sqrt[3]{a-ia \tan(c+dx)}\right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \frac{\log(a-ia \tan(c+dx))}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d}}{4a} + \frac{3(B+4iA)(a+ia \tan(c+dx))^{5/3}}{5d}$$

↓ 217

$$\frac{\frac{3B \tan^2(c+dx)(a+ia \tan(c+dx))^{2/3}}{8d} - \frac{4ia^2(A-iB) \left(-\frac{i \sqrt{3} \operatorname{arctanh}\left(\frac{a \tan(c+dx)}{\sqrt{3}}\right)}{\sqrt[3]{2} \sqrt[3]{a}} - \frac{3 \log\left(\sqrt[3]{2} \sqrt[3]{a-ia \tan(c+dx)}\right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \frac{\log(a-ia \tan(c+dx))}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d}}{4a} + \frac{3(B+4iA)(a+ia \tan(c+dx))^{5/3}}{5d} + \frac{9aB}{5d}$$

input `Int[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^(2/3)*(A + B*Tan[c + d*x]),x]`

output `(3*B*Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^(2/3))/(8*d) - (((-4*I)*a^2*(A - I*B)*((-I)*Sqrt[3]*ArcTanh[(a*Tan[c + d*x])/Sqrt[3]])/(2^(1/3)*a^(1/3)) - (3*Log[2^(1/3)*a^(1/3) - I*a*Tan[c + d*x]])/(2*2^(1/3)*a^(1/3)) + Log[a - I*a*Tan[c + d*x]]/(2*2^(1/3)*a^(1/3)))/d + (9*a*B*(a + I*a*Tan[c + d*x])^(2/3))/(2*d) + (3*((4*I)*A + B)*(a + I*a*Tan[c + d*x])^(5/3))/(5*d)/(4*a)`

3.197.3.1 Defintions of rubi rules used

- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 67 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(1/3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q), x] + (Simp[3/(2*b) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Simp[3/(2*b*q) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x])] /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3962 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[-b/d Subst[Int[(a + x)^(n - 1)/(a - x), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 + b^2, 0]`
- rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4080 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_)), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`

3.197.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.83

method	result
derivativedivides	$3i \left(\frac{iB(a+ia \tan(dx+c))^{\frac{8}{3}}}{8} - \frac{iBa(a+ia \tan(dx+c))^{\frac{5}{3}}}{5} - \frac{Aa(a+ia \tan(dx+c))^{\frac{5}{3}}}{5} + \frac{ia^2B(a+ia \tan(dx+c))^{\frac{2}{3}}}{2} \right) - \frac{2^{\frac{2}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}} \right)}{6a^{\frac{1}{3}}}$
default	$3i \left(\frac{iB(a+ia \tan(dx+c))^{\frac{8}{3}}}{8} - \frac{iBa(a+ia \tan(dx+c))^{\frac{5}{3}}}{5} - \frac{Aa(a+ia \tan(dx+c))^{\frac{5}{3}}}{5} + \frac{ia^2B(a+ia \tan(dx+c))^{\frac{2}{3}}}{2} \right) - \frac{2^{\frac{2}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}} \right)}{6a^{\frac{1}{3}}}$
parts	$3iA \left(-\frac{(a+ia \tan(dx+c))^{\frac{5}{3}}}{5} - \frac{2^{\frac{2}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}} \right)}{6a^{\frac{1}{3}}} - \frac{2^{\frac{2}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}} (a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} \right)}{12a^{\frac{1}{3}}} \right)$

da

```
input int(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 3*I/d/a^2*(1/8*I*B*(a+I*a*tan(d*x+c))^(8/3)-1/5*I*B*a*(a+I*a*tan(d*x+c))^(5/3)-1/5*A*a*(a+I*a*tan(d*x+c))^(5/3)+1/2*I*a^2*B*(a+I*a*tan(d*x+c))^(2/3)-(1/6*2^(2/3)/a^(1/3)*ln((a+I*a*tan(d*x+c))^(1/3)-2^(1/3)*a^(1/3))-1/12*2^(2/3)/a^(1/3)*ln((a+I*a*tan(d*x+c))^(2/3)+2^(1/3)*a^(1/3)*(a+I*a*tan(d*x+c))^(1/3)+2^(2/3)*a^(2/3))+1/6*3^(1/2)*2^(2/3)/a^(1/3)*arctan(1/3*3^(1/2)*(2^(2/3)/a^(1/3)*(a+I*a*tan(d*x+c))^(1/3)+1)))a^3*(A-I*B)
```

3.197.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 660 vs. $2(201) = 402$.

Time = 0.26 (sec) , antiderivative size = 660, normalized size of antiderivative = 2.44

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx =$$

$$3 \cdot 2^{\frac{2}{3}} (2(2iA + 3B)e^{(4i dx + 4i c)} + 2(2iA + 3B)e^{(2i dx + 2i c)} + 5B) \left(\frac{a}{e^{(2i dx + 2i c)} + 1} \right)^{\frac{2}{3}} e^{\left(\frac{4}{3}i dx + \frac{4}{3}i c\right)} - 10 \left(\frac{1}{2}\right)^{\frac{1}{3}} (de$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-1/10*(3*2^(2/3)*(2*(2*I*A + 3*B)*e^(4*I*d*x + 4*I*c) + 2*(2*I*A + 3*B)*e^(2*I*d*x + 2*I*c) + 5*B)*(a/(e^(2*I*d*x + 2*I*c) + 1))^(2/3)*e^(4/3*I*d*x + 4/3*I*c) - 10*(1/2)^(1/3)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*((I*A^3 + 3*A^2*B - 3*I*A*B^2 - B^3)*a^2/d^3)^(1/3)*log((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) + 2*(1/2)^(2/3)*d^2*((I*A^3 + 3*A^2*B - 3*I*A*B^2 - B^3)*a^2/d^3)^(2/3))/((A^2 - 2*I*A*B - B^2)*a)) + 5*(1/2)^(1/3)*((-I*sqrt(3)*d + d)*e^(4*I*d*x + 4*I*c) + 2*(-I*sqrt(3)*d + d)*e^(2*I*d*x + 2*I*c) - I*sqrt(3)*d + d)*((I*A^3 + 3*A^2*B - 3*I*A*B^2 - B^3)*a^2/d^3)^(1/3)*log((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) - (1/2)^(2/3)*(I*sqrt(3)*d^2 + d^2)*((I*A^3 + 3*A^2*B - 3*I*A*B^2 - B^3)*a^2/d^3)^(2/3))/((A^2 - 2*I*A*B - B^2)*a)) + 5*(1/2)^(1/3)*((I*sqrt(3)*d + d)*e^(4*I*d*x + 4*I*c) + 2*(I*sqrt(3)*d + d)*e^(2*I*d*x + 2*I*c) + I*sqrt(3)*d + d)*((I*A^3 + 3*A^2*B - 3*I*A*B^2 - B^3)*a^2/d^3)^(1/3)*log((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) - (1/2)^(2/3)*(-I*sqrt(3)*d^2 + d^2)*((I*A^3 + 3*A^2*B - 3*I*A*B^2 - B^3)*a^2/d^3)^(2/3))/((A^2 - 2*I*A*B - B^2)*a)))/(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)`

3.197.6 Sympy [F]

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{2/3} (A + B \tan(c + dx)) \tan^2(c + dx) dx$$

input `integrate(tan(d*x+c)**2*(a+I*a*tan(d*x+c))**(2/3)*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(2/3)*(A + B*tan(c + d*x))*tan(c + d*x)**2, x)`

3.197.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 210, normalized size of antiderivative = 0.78

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx =$$

$$i \left(20 \sqrt{3} 2^{2/3} (A - iB) a^{11/3} \arctan \left(\frac{\sqrt{3} 2^{2/3} (2^{1/3} a^{1/3} + 2(i a \tan(dx+c) + a)^{1/3})}{6 a^{1/3}} \right) - 10 \cdot 2^{2/3} (A - iB) a^{11/3} \log \left(2^{2/3} a^{2/3} + 2^{1/3} (i a \tan(dx+c) + a)^{1/3} \right) \right)$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/40*I*(20*sqrt(3)*2^(2/3)*(A - I*B)*a^(11/3)*arctan(1/6*sqrt(3)*2^(2/3)*(2^(1/3)*a^(1/3) + 2*(I*a*tan(d*x + c) + a)^(1/3))/a^(1/3)) - 10*2^(2/3)*(A - I*B)*a^(11/3)*log(2^(2/3)*a^(2/3) + 2^(1/3)*(I*a*tan(d*x + c) + a)^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(2/3)) + 20*2^(2/3)*(A - I*B)*a^(11/3)*log(-2^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(1/3)) - 15*I*(I*a*tan(d*x + c) + a)^(8/3)*B*a + 24*(I*a*tan(d*x + c) + a)^(5/3)*(A + I*B)*a^2 - 60*I*(I*a*tan(d*x + c) + a)^(2/3)*B*a^3)/(a^3*d)`

3.197.8 Giac [F]

$$\int \tan^2(c+dx)(a+ia \tan(c+dx))^{2/3}(A+B \tan(c+dx)) dx = \int (B \tan(dx+c) + A)(ia \tan(dx+c) + a)^{2/3} \tan(dx+c)^2 dx$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorith="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(2/3)*tan(d*x + c)^2, x)`

3.197.9 Mupad [B] (verification not implemented)

Time = 10.29 (sec) , antiderivative size = 436, normalized size of antiderivative = 1.61

$$\begin{aligned} & \int \tan^2(c+dx)(a+ia \tan(c+dx))^{2/3}(A+B \tan(c+dx)) dx = -\frac{3B(a+a \tan(c+dx) \operatorname{li})^{2/3}}{2d} \\ & - \frac{A(a+a \tan(c+dx) \operatorname{li})^{5/3} 3i}{5ad} + \frac{3B(a+a \tan(c+dx) \operatorname{li})^{5/3}}{5ad} \\ & - \frac{3B(a+a \tan(c+dx) \operatorname{li})^{8/3}}{8a^2d} - \frac{2^{2/3} B a^{2/3} \ln\left((a(1+\tan(c+dx) \operatorname{li}))^{1/3} - 2^{1/3} a^{1/3}\right)}{2d} \\ & + \frac{\left(\frac{1}{2}i\right)^{1/3} A a^{2/3} \ln\left((a(1+\tan(c+dx) \operatorname{li}))^{1/3} + (-1)^{1/3} 2^{1/3} a^{1/3}\right)}{d} \\ & + \frac{\left(\frac{1}{2}i\right)^{1/3} A a^{2/3} \ln\left(\frac{(-1)^{1/3} 2^{1/3} a^{1/3}}{2} - (a(1+\tan(c+dx) \operatorname{li}))^{1/3} + \frac{(-1)^{5/6} 2^{1/3} \sqrt{3} a^{1/3}}{2}\right) \left(-\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)}{d} \\ & - \frac{2^{2/3} B a^{2/3} \ln\left(\frac{9B^2 a^2 (a+a \tan(c+dx) \operatorname{li})^{1/3}}{d^2} - \frac{9 \cdot 2^{1/3} B^2 a^{7/3} \left(-\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)^2}{d^2}\right) \left(-\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)}{2d} \\ & + \frac{2^{2/3} B a^{2/3} \ln\left(\frac{9B^2 a^2 (a+a \tan(c+dx) \operatorname{li})^{1/3}}{d^2} - \frac{9 \cdot 2^{1/3} B^2 a^{7/3} \left(\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)^2}{d^2}\right) \left(\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)}{2d} \\ & - \frac{\left(\frac{1}{2}i\right)^{1/3} A a^{2/3} \ln\left((a(1+\tan(c+dx) \operatorname{li}))^{1/3} - \frac{(-1)^{1/3} 2^{1/3} a^{1/3}}{2} + \frac{(-1)^{5/6} 2^{1/3} \sqrt{3} a^{1/3}}{2}\right) \left(\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)}{d} \end{aligned}$$

3.197. $\int \tan^2(c+dx)(a+ia \tan(c+dx))^{2/3}(A+B \tan(c+dx)) dx$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(2/3),x)`

output
$$\begin{aligned} & (3*B*(a + a*\tan(c + d*x)*1i)^{(5/3)})/(5*a*d) - (A*(a + a*\tan(c + d*x)*1i)^{(5/3)*3i})/(5*a*d) - (3*B*(a + a*\tan(c + d*x)*1i)^{(2/3)})/(2*d) - (3*B*(a + a*\tan(c + d*x)*1i)^{(8/3)})/(8*a^2*d) - (2^{(2/3)}*B*a^{(2/3)}*\log((a*(\tan(c + d*x)*1i + 1))^{(1/3)} - 2^{(1/3)}*a^{(1/3)}))/(2*d) + ((1i/2)^{(1/3)}*A*a^{(2/3)}*\log((a*(\tan(c + d*x)*1i + 1))^{(1/3)} + (-1)^{(1/3)}*2^{(1/3)}*a^{(1/3)}))/d + ((1i/2)^{(1/3)}*A*a^{(2/3)}*\log((-1)^{(1/3)}*2^{(1/3)}*a^{(1/3)})/2 - (a*(\tan(c + d*x)*1i + 1))^{(1/3)} + ((-1)^{(5/6)}*2^{(1/3)}*3^{(1/2)}*a^{(1/3)})/2)*((3^{(1/2)}*1i)/2 - 1/2))/d - (2^{(2/3)}*B*a^{(2/3)}*\log((9*B^2*a^2*(a + a*\tan(c + d*x)*1i)^{(1/3)})/d^2 - (9*2^{(1/3)}*B^2*a^{(7/3)}*((3^{(1/2)}*1i)/2 - 1/2)^2)/d^2)*((3^{(1/2)}*1i)/2 - 1/2))/(2*d) + (2^{(2/3)}*B*a^{(2/3)}*\log((9*B^2*a^2*(a + a*\tan(c + d*x)*1i)^{(1/3)})/d^2 - (9*2^{(1/3)}*B^2*a^{(7/3)}*((3^{(1/2)}*1i)/2 + 1/2)^2)/d^2)*((3^{(1/2)}*1i)/2 + 1/2))/(2*d) - ((1i/2)^{(1/3)}*A*a^{(2/3)}*\log((a*(\tan(c + d*x)*1i + 1))^{(1/3)} - ((-1)^{(1/3)}*2^{(1/3)}*a^{(1/3)})/2 + ((-1)^{(5/6)}*2^{(1/3)}*3^{(1/2)}*a^{(1/3)})/2)*((3^{(1/2)}*1i)/2 + 1/2))/d \end{aligned}$$

3.198 $\int \tan(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx$

3.198.1 Optimal result	2027
3.198.2 Mathematica [A] (verified)	2028
3.198.3 Rubi [A] (warning: unable to verify)	2028
3.198.4 Maple [A] (verified)	2031
3.198.5 Fricas [B] (verification not implemented)	2033
3.198.6 Sympy [F]	2034
3.198.7 Maxima [A] (verification not implemented)	2034
3.198.8 Giac [F]	2035
3.198.9 Mupad [B] (verification not implemented)	2035

3.198.1 Optimal result

Integrand size = 34, antiderivative size = 232

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \frac{a^{2/3}(iA + B)x}{2\sqrt[3]{2}}$$

$$+ \frac{\sqrt{3}a^{2/3}(A - iB) \arctan\left(\frac{\sqrt[3]{a+2^{2/3}}\sqrt[3]{a + ia \tan(c + dx)}}{\sqrt{3}\sqrt[3]{a}}\right)}{\sqrt[3]{2}d}$$

$$+ \frac{a^{2/3}(A - iB) \log(\cos(c + dx))}{2\sqrt[3]{2}d}$$

$$+ \frac{3a^{2/3}(A - iB) \log\left(\sqrt[3]{2}\sqrt[3]{a} - \sqrt[3]{a + ia \tan(c + dx)}\right)}{2\sqrt[3]{2}d}$$

$$+ \frac{3A(a + ia \tan(c + dx))^{2/3}}{2d} - \frac{3iB(a + ia \tan(c + dx))^{5/3}}{5ad}$$

output

```
1/4*a^(2/3)*(I*A+B)*x*2^(2/3)+1/4*a^(2/3)*(A-I*B)*ln(cos(d*x+c))*2^(2/3)/d
+3/4*a^(2/3)*(A-I*B)*ln(2^(1/3)*a^(1/3)-(a+I*a*tan(d*x+c))^(1/3))*2^(2/3)/
d+1/2*a^(2/3)*(A-I*B)*arctan(1/3*(a^(1/3)+2^(2/3)*(a+I*a*tan(d*x+c))^(1/3)
)/a^(1/3)*3^(1/2))*3^(1/2)*2^(2/3)/d+3/2*A*(a+I*a*tan(d*x+c))^(2/3)/d-3/5*
I*B*(a+I*a*tan(d*x+c))^(5/3)/a/d
```

3.198.2 Mathematica [A] (verified)

Time = 1.25 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.71

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \frac{5 \cdot 2^{2/3} a^{5/3} (A - iB) \left(2\sqrt{3} \arctan \left(\frac{1 + \frac{2^{2/3} \sqrt[3]{a + ia \tan(c + dx)}}{\sqrt[3]{a}}}{\sqrt{3}} \right) - \log(i + \tan(c + dx)) \right)}{\dots}$$

```
input Integrate[Tan[c + d*x]*(a + I*a*Tan[c + d*x])^(2/3)*(A + B*Tan[c + d*x]),x
]
```

```
output (5*2^(2/3)*a^(5/3)*(A - I*B)*(2*Sqrt[3]*ArcTan[(1 + (2^(2/3)*(a + I*a*Tan[
c + d*x])^(1/3))/a^(1/3)]/Sqrt[3]] - Log[I + Tan[c + d*x]] + 3*Log[2^(1/3)
*a^(1/3) - (a + I*a*Tan[c + d*x])^(1/3)]) + 30*a*A*(a + I*a*Tan[c + d*x])^(
2/3) - (12*I)*B*(a + I*a*Tan[c + d*x])^(5/3))/(20*a*d)
```

3.198.3 Rubi [A] (warning: unable to verify)

Time = 0.54 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.74, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3042, 4075, 3042, 4010, 3042, 3962, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4075} \\ & \int (i \tan(c + dx)a + a)^{2/3}(A \tan(c + dx) - B) dx - \frac{3iB(a + ia \tan(c + dx))^{5/3}}{5ad} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\begin{aligned}
 & \int (i \tan(c + dx)a + a)^{2/3} (A \tan(c + dx) - B) dx - \frac{3iB(a + ia \tan(c + dx))^{5/3}}{5ad} \\
 & \quad \downarrow \text{4010} \\
 & -(B + iA) \int (i \tan(c + dx)a + a)^{2/3} dx + \frac{3A(a + ia \tan(c + dx))^{2/3}}{2d} - \frac{3iB(a + ia \tan(c + dx))^{5/3}}{5ad} \\
 & \quad \downarrow \text{3042} \\
 & -(B + iA) \int (i \tan(c + dx)a + a)^{2/3} dx + \frac{3A(a + ia \tan(c + dx))^{2/3}}{2d} - \frac{3iB(a + ia \tan(c + dx))^{5/3}}{5ad} \\
 & \quad \downarrow \text{3962} \\
 & \frac{ia(B + iA) \int \frac{1}{(a - ia \tan(c + dx)) \sqrt[3]{i \tan(c + dx)a + a}} d(i \tan(c + dx))}{d} + \frac{3A(a + ia \tan(c + dx))^{2/3}}{2d} - \frac{3iB(a + ia \tan(c + dx))^{5/3}}{5ad} \\
 & \quad \downarrow \text{67} \\
 & ia(B + iA) \left(-\frac{3}{2} \int \frac{1}{-a^2 \tan^2(c + dx) + i \sqrt[3]{2} a^{4/3} \tan(c + dx) + 2^{2/3} a^{2/3}} d \sqrt[3]{i \tan(c + dx)a + a} + \frac{3 \int \frac{1}{\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)}} d \sqrt[3]{i \tan(c + dx)a + a}}{2 \sqrt[3]{2} \sqrt[3]{a}} \right) \\
 & \quad \frac{3A(a + ia \tan(c + dx))^{2/3}}{2d} - \frac{3iB(a + ia \tan(c + dx))^{5/3}}{5ad} \\
 & \quad \downarrow \text{16} \\
 & ia(B + iA) \left(-\frac{3}{2} \int \frac{1}{-a^2 \tan^2(c + dx) + i \sqrt[3]{2} a^{4/3} \tan(c + dx) + 2^{2/3} a^{2/3}} d \sqrt[3]{i \tan(c + dx)a + a} - \frac{3 \log \left(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)} \right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \log \right) \\
 & \quad \frac{3A(a + ia \tan(c + dx))^{2/3}}{2d} - \frac{3iB(a + ia \tan(c + dx))^{5/3}}{5ad} \\
 & \quad \downarrow \text{1082} \\
 & ia(B + iA) \left(\frac{3 \int \frac{1}{a^2 \tan^2(c + dx) - 3} d(i 2^{2/3} a^{2/3} \tan(c + dx) + 1)}{\sqrt[3]{2} \sqrt[3]{a}} - \frac{3 \log \left(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)} \right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \frac{\log(a - ia \tan(c + dx))}{2 \sqrt[3]{2} \sqrt[3]{a}} \right) + \\
 & \quad \frac{3A(a + ia \tan(c + dx))^{2/3}}{2d} - \frac{3iB(a + ia \tan(c + dx))^{5/3}}{5ad} \\
 & \quad \downarrow \text{217}
 \end{aligned}$$

3.198. $\int \tan(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx$

$$ia(B + iA) \left(-\frac{i\sqrt{3}\operatorname{arctanh}\left(\frac{a \tan(c+dx)}{\sqrt{3}}\right)}{\sqrt[3]{2}\sqrt[3]{a}} - \frac{3 \log\left(\sqrt[3]{2}\sqrt[3]{a} - ia \tan(c+dx)\right)}{2\sqrt[3]{2}\sqrt[3]{a}} + \frac{\log(a - ia \tan(c+dx))}{2\sqrt[3]{2}\sqrt[3]{a}} \right) + \frac{3A(a + ia \tan(c + dx))^{2/3}}{2d} - \frac{3iB(a + ia \tan(c + dx))^{5/3}}{5ad}$$

input `Int[Tan[c + d*x]*(a + I*a*Tan[c + d*x])^(2/3)*(A + B*Tan[c + d*x]),x]`

output `(I*a*(I*A + B)*((-I)*Sqrt[3]*ArcTanh[(a*Tan[c + d*x])/Sqrt[3]])/(2^(1/3)*a^(1/3)) - (3*Log[2^(1/3)*a^(1/3) - I*a*Tan[c + d*x]]/(2*2^(1/3)*a^(1/3)) + Log[a - I*a*Tan[c + d*x]]/(2*2^(1/3)*a^(1/3)))/d + (3*A*(a + I*a*Tan[c + d*x])^(2/3))/(2*d) - (((3*I)/5)*B*(a + I*a*Tan[c + d*x])^(5/3))/(a*d)`

3.198.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 67 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(1/3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q), x] + (Simp[3/(2*b) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Simp[3/(2*b*q) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x]) /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3962 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[-b/d S
ubst[Int[(a + x)^(n - 1)/(a - x), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b
, c, d, n}, x] && EqQ[a^2 + b^2, 0]`

rule 4010 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp
[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e
, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B
d((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.198.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 181, normalized size of antiderivative = 0.78

method	result
derivativedivides	$\frac{-\frac{3iB(a+ia \tan(dx+c))^{\frac{5}{3}}}{5} + \frac{3aA(a+ia \tan(dx+c))^{\frac{2}{3}}}{2} + 3}{ad} \left(\frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{6a^{\frac{1}{3}}} - \frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{12a^{\frac{1}{3}}}\right)$
default	$\frac{-\frac{3iB(a+ia \tan(dx+c))^{\frac{5}{3}}}{5} + \frac{3aA(a+ia \tan(dx+c))^{\frac{2}{3}}}{2} + 3}{ad} \left(\frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{6a^{\frac{1}{3}}} - \frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{12a^{\frac{1}{3}}}\right)$
parts	$A \left(\frac{3(a+ia \tan(dx+c))^{\frac{2}{3}}}{2d} + \frac{a^{\frac{2}{3}} 2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{2d} - \frac{a^{\frac{2}{3}} 2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{4d} \right)$

input `int(tan(d*x+c)*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)`

output `3/d/a*(-1/5*I*B*(a+I*a*tan(d*x+c))^(5/3)+1/2*a*A*(a+I*a*tan(d*x+c))^(2/3)+
(1/6*2^(2/3)/a^(1/3)*ln((a+I*a*tan(d*x+c))^(1/3)-2^(1/3)*a^(1/3))-1/12*2^(
2/3)/a^(1/3)*ln((a+I*a*tan(d*x+c))^(2/3)+2^(1/3)*a^(1/3)*(a+I*a*tan(d*x+c)
)^(1/3)+2^(2/3)*a^(2/3))+1/6*3^(1/2)*2^(2/3)/a^(1/3)*arctan(1/3*3^(1/2)*(2
^(2/3)/a^(1/3)*(a+I*a*tan(d*x+c))^(1/3)+1)))*a^2*(A-I*B))`

3.198.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 571 vs. $2(168) = 336$.

Time = 0.25 (sec) , antiderivative size = 571, normalized size of antiderivative = 2.46

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{2/3}(A$$

$$3 \cdot 2^{2/3}((5A - 4iB)e^{(2i dx + 2i c)} + 5A) \left(\frac{a}{e^{(2i dx + 2i c)} + 1} \right)^{2/3} e^{(4/3 i dx + 4/3 i c)} + 10 \left(\frac{1}{2} \right)^{1/3} (de^{(2i dx + 2i c)}$$

$$+ B \tan(c + dx)) dx =$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/10*(3*2^(2/3)*((5*A - 4*I*B)*e^(2*I*d*x + 2*I*c) + 5*A)*(a/(e^(2*I*d*x + 2*I*c) + 1))^(2/3)*e^(4/3*I*d*x + 4/3*I*c) + 10*(1/2)^(1/3)*(d*e^(2*I*d*x + 2*I*c) + d)*((A^3 - 3*I*A^2*B - 3*A*B^2 + I*B^3)*a^2/d^3)^(1/3)*log((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) - 2*(1/2)^(2/3)*d^2*((A^3 - 3*I*A^2*B - 3*A*B^2 + I*B^3)*a^2/d^3)^(2/3))/((A^2 - 2*I*A*B - B^2)*a)) - 5*(1/2)^(1/3)*((I*sqrt(3)*d + d)*e^(2*I*d*x + 2*I*c) + I*sqrt(3)*d + d)*((A^3 - 3*I*A^2*B - 3*A*B^2 + I*B^3)*a^2/d^3)^(1/3)*log((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) - (1/2)^(2/3)*(I*sqrt(3)*d^2 - d^2)*((A^3 - 3*I*A^2*B - 3*A*B^2 + I*B^3)*a^2/d^3)^(2/3))/((A^2 - 2*I*A*B - B^2)*a)) - 5*(1/2)^(1/3)*((-I*sqrt(3)*d + d)*e^(2*I*d*x + 2*I*c) - I*sqrt(3)*d + d)*((A^3 - 3*I*A^2*B - 3*A*B^2 + I*B^3)*a^2/d^3)^(1/3)*log((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) - (1/2)^(2/3)*(-I*sqrt(3)*d^2 - d^2)*((A^3 - 3*I*A^2*B - 3*A*B^2 + I*B^3)*a^2/d^3)^(2/3))/((A^2 - 2*I*A*B - B^2)*a)))/(d*e^(2*I*d*x + 2*I*c) + d)`

3.198.6 Sympy [F]

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{2/3}(A + B \tan(c + dx)) \tan(c + dx) dx$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))**(2/3)*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(2/3)*(A + B*tan(c + d*x))*tan(c + d*x), x)`

3.198.7 Maxima [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.81

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \frac{10 \sqrt{3} 2^{2/3} (A - i B) a^{8/3} \arctan\left(\frac{\sqrt{3} 2^{2/3} (2^{1/3} a^{1/3} + 2(i a \tan(dx+c) + a)^{1/3})}{6 a^{1/3}}\right) - 5 \cdot 2^{2/3} (A - i B) a^{8/3} \log\left(\dots\right)}{1}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/20*(10*sqrt(3)*2^(2/3)*(A - I*B)*a^(8/3)*arctan(1/6*sqrt(3)*2^(2/3)*(2^(1/3)*a^(1/3) + 2*(I*a*tan(d*x + c) + a)^(1/3))/a^(1/3)) - 5*2^(2/3)*(A - I*B)*a^(8/3)*log(2^(2/3)*a^(2/3) + 2^(1/3)*(I*a*tan(d*x + c) + a)^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(2/3)) + 10*2^(2/3)*(A - I*B)*a^(8/3)*log(-2^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(1/3)) - 12*I*(I*a*tan(d*x + c) + a)^(5/3)*B*a + 30*(I*a*tan(d*x + c) + a)^(2/3)*A*a^2)/(a^2*d)`

3.198.8 Giac [F]

$$\int \tan(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{2/3} \tan(dx + c) dx$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(2/3)*tan(d*x + c), x)`

3.198.9 Mupad [B] (verification not implemented)

Time = 9.57 (sec) , antiderivative size = 390, normalized size of antiderivative = 1.68

$$\begin{aligned} \int \tan(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx &= \frac{3 A (a + a \tan(c + dx) i i)^{2/3}}{2 d} \\ &- \frac{B (a + a \tan(c + dx) i i)^{5/3} 3 i}{5 a d} + \frac{2^{2/3} A a^{2/3} \ln \left((a (1 + \tan(c + dx) i i))^{1/3} - 2^{1/3} a^{1/3} \right)}{2 d} \\ &+ \frac{\left(\frac{1}{2} i\right)^{1/3} B a^{2/3} \ln \left((a (1 + \tan(c + dx) i i))^{1/3} + (-1)^{1/3} 2^{1/3} a^{1/3} \right)}{d} \\ &+ \frac{\left(\frac{1}{2} i\right)^{1/3} B a^{2/3} \ln \left(\frac{(-1)^{1/3} 2^{1/3} a^{1/3}}{2} - (a (1 + \tan(c + dx) i i))^{1/3} + \frac{(-1)^{5/6} 2^{1/3} \sqrt{3} a^{1/3}}{2} \right) \left(-\frac{1}{2} + \frac{\sqrt{3} i i}{2} \right)}{d} \\ &+ \frac{2^{2/3} A a^{2/3} \ln \left(\frac{9 A^2 a^2 (a + a \tan(c + dx) i i)^{1/3}}{d^2} - \frac{9 2^{1/3} A^2 a^{7/3} \left(-\frac{1}{2} + \frac{\sqrt{3} i i}{2} \right)^2}{d^2} \right) \left(-\frac{1}{2} + \frac{\sqrt{3} i i}{2} \right)}{2 d} \\ &+ \frac{2^{2/3} A a^{2/3} \ln \left(\frac{9 A^2 a^2 (a + a \tan(c + dx) i i)^{1/3}}{d^2} - \frac{9 2^{1/3} A^2 a^{7/3} \left(\frac{1}{2} + \frac{\sqrt{3} i i}{2} \right)^2}{d^2} \right) \left(\frac{1}{2} + \frac{\sqrt{3} i i}{2} \right)}{2 d} \\ &- \frac{\left(\frac{1}{2} i\right)^{1/3} B a^{2/3} \ln \left((a (1 + \tan(c + dx) i i))^{1/3} - \frac{(-1)^{1/3} 2^{1/3} a^{1/3}}{2} + \frac{(-1)^{5/6} 2^{1/3} \sqrt{3} a^{1/3}}{2} \right) \left(\frac{1}{2} + \frac{\sqrt{3} i i}{2} \right)}{d} \end{aligned}$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*ii)^(2/3),x)`

3.198. $\int \tan(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx$

output $(3A(a + a\tan(c + dx)1i)^{2/3})/(2d) - (B(a + a\tan(c + dx)1i)^{5/3}3i)/(5a*d) + (2^{2/3}Aa^{2/3}\log((a(\tan(c + dx)1i + 1))^{1/3} - 2^{1/3}a^{1/3}))/2d + ((1i/2)^{1/3}Ba^{2/3}\log((a(\tan(c + dx)1i + 1))^{1/3} + (-1)^{1/3}2^{1/3}a^{1/3}))/d + ((1i/2)^{1/3}Ba^{2/3}\log(((-1)^{1/3}2^{1/3}a^{1/3})/2 - (a(\tan(c + dx)1i + 1))^{1/3} + ((-1)^{5/6}2^{1/3}3^{1/2}a^{1/3})/2)*((3^{1/2}1i)/2 - 1/2))/d + (2^{2/3}Aa^{2/3}\log((9A^2a^2(a + a\tan(c + dx)1i)^{1/3})/d^2 - (9*2^{1/3}A^2a^{7/3}*((3^{1/2}1i)/2 - 1/2)^2)/d^2)*((3^{1/2}1i)/2 - 1/2))/2d - (2^{2/3}Aa^{2/3}\log((9A^2a^2(a + a\tan(c + dx)1i)^{1/3})/d^2 - (9*2^{1/3}A^2a^{7/3}*((3^{1/2}1i)/2 + 1/2)^2)/d^2)*((3^{1/2}1i)/2 + 1/2))/(2*d) - ((1i/2)^{1/3}Ba^{2/3}\log((a(\tan(c + dx)1i + 1))^{1/3} - ((-1)^{1/3}2^{1/3}a^{1/3}))/2 + ((-1)^{5/6}2^{1/3}3^{1/2}a^{1/3})/2)*((3^{1/2}1i)/2 + 1/2))/d$

3.199 $\int (a+ia \tan(c+dx))^{2/3}(A+B \tan(c+dx)) dx$

3.199.1 Optimal result	2037
3.199.2 Mathematica [A] (verified)	2038
3.199.3 Rubi [A] (warning: unable to verify)	2038
3.199.4 Maple [A] (verified)	2041
3.199.5 Fracas [B] (verification not implemented)	2042
3.199.6 Sympy [F]	2042
3.199.7 Maxima [A] (verification not implemented)	2043
3.199.8 Giac [F]	2043
3.199.9 Mupad [B] (verification not implemented)	2044

3.199.1 Optimal result

Integrand size = 28, antiderivative size = 202

$$\int (a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = -\frac{a^{2/3}(A - iB)x}{2\sqrt[3]{2}} + \frac{\sqrt{3}a^{2/3}(iA + B) \arctan\left(\frac{\sqrt[3]{a+2^{2/3}}\sqrt[3]{a + ia \tan(c + dx)}}{\sqrt{3}\sqrt[3]{a}}\right)}{\sqrt[3]{2}d} + \frac{a^{2/3}(iA + B) \log(\cos(c + dx))}{2\sqrt[3]{2}d} + \frac{3a^{2/3}(iA + B) \log\left(\sqrt[3]{2}\sqrt[3]{a} - \sqrt[3]{a + ia \tan(c + dx)}\right)}{2\sqrt[3]{2}d} + \frac{3B(a + ia \tan(c + dx))^{2/3}}{2d}$$

output

```
-1/4*a^(2/3)*(A-I*B)*x*2^(2/3)+1/4*a^(2/3)*(I*A+B)*ln(cos(d*x+c))*2^(2/3)/
d+3/4*a^(2/3)*(I*A+B)*ln(2^(1/3)*a^(1/3)-(a+I*a*tan(d*x+c))^(1/3))*2^(2/3)
/d+1/2*a^(2/3)*(I*A+B)*arctan(1/3*(a^(1/3)+2^(2/3)*(a+I*a*tan(d*x+c))^(1/3
))/a^(1/3)*3^(1/2))*3^(1/2)*2^(2/3)/d+3/2*B*(a+I*a*tan(d*x+c))^(2/3)/d
```

3.199.2 Mathematica [A] (verified)

Time = 0.66 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.68

$$\int (a + ia \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx = \frac{2^{2/3} a^{2/3} (iA + B) \left(2\sqrt{3} \arctan \left(\frac{1 + \frac{2^{2/3} \sqrt[3]{a + ia \tan(c + dx)}}{\sqrt[3]{a}}}{\sqrt{3}} \right) - \log(i + \tan(c + dx)) \right)}{4d}$$

input `Integrate[(a + I*a*Tan[c + d*x])^(2/3)*(A + B*Tan[c + d*x]),x]`output `(2^(2/3)*a^(2/3)*(I*A + B)*(2*Sqrt[3]*ArcTan[(1 + (2^(2/3)*(a + I*a*Tan[c + d*x])^(1/3))/a^(1/3)]/Sqrt[3]) - Log[I + Tan[c + d*x]] + 3*Log[2^(1/3)*a^(1/3) - (a + I*a*Tan[c + d*x])^(1/3)]) + 6*B*(a + I*a*Tan[c + d*x])^(2/3))/(4*d)`**3.199.3 Rubi [A] (warning: unable to verify)**Time = 0.40 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.70, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4010, 3042, 3962, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4010} \\ & (A - iB) \int (i \tan(c + dx) a + a)^{2/3} dx + \frac{3B(a + ia \tan(c + dx))^{2/3}}{2d} \\ & \quad \downarrow \text{3042} \\ & (A - iB) \int (i \tan(c + dx) a + a)^{2/3} dx + \frac{3B(a + ia \tan(c + dx))^{2/3}}{2d} \end{aligned}$$

$$\begin{aligned}
 & \frac{3B(a + ia \tan(c + dx))^{2/3}}{2d} - \frac{ia(A - iB) \int \frac{1}{(a - ia \tan(c + dx)) \sqrt[3]{i \tan(c + dx) a + a}} d(ia \tan(c + dx))}{d} \\
 & \quad \downarrow 3962 \\
 & \frac{3B(a + ia \tan(c + dx))^{2/3}}{2d} - \frac{ia(A - iB) \left(-\frac{3}{2} \int \frac{1}{-a^2 \tan^2(c + dx) + i \sqrt[3]{2} a^{4/3} \tan(c + dx) + 2^{2/3} a^{2/3}} d \sqrt[3]{i \tan(c + dx) a + a} + \frac{3 \int \frac{1}{\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)}} d \sqrt[3]{i \tan(c + dx) a + a}}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d} \\
 & \quad \downarrow 67 \\
 & \frac{3B(a + ia \tan(c + dx))^{2/3}}{2d} - \frac{ia(A - iB) \left(-\frac{3}{2} \int \frac{1}{-a^2 \tan^2(c + dx) + i \sqrt[3]{2} a^{4/3} \tan(c + dx) + 2^{2/3} a^{2/3}} d \sqrt[3]{i \tan(c + dx) a + a} - \frac{3 \log\left(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)}\right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \log\left(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)}\right) \right)}{d} \\
 & \quad \downarrow 16 \\
 & \frac{3B(a + ia \tan(c + dx))^{2/3}}{2d} - \frac{ia(A - iB) \left(-\frac{3}{2} \int \frac{1}{-a^2 \tan^2(c + dx) + i \sqrt[3]{2} a^{4/3} \tan(c + dx) + 2^{2/3} a^{2/3}} d \sqrt[3]{i \tan(c + dx) a + a} - \frac{3 \log\left(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)}\right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \log\left(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)}\right) \right)}{d} \\
 & \quad \downarrow 1082 \\
 & \frac{3B(a + ia \tan(c + dx))^{2/3}}{2d} - \frac{ia(A - iB) \left(\frac{3 \int \frac{1}{a^2 \tan^2(c + dx) - 3} d(i 2^{2/3} a^{2/3} \tan(c + dx) + 1)}{\sqrt[3]{2} \sqrt[3]{a}} - \frac{3 \log\left(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)}\right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \frac{\log(a - ia \tan(c + dx))}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d} \\
 & \quad \downarrow 217 \\
 & \frac{3B(a + ia \tan(c + dx))^{2/3}}{2d} - \frac{ia(A - iB) \left(-\frac{i \sqrt{3} \operatorname{arctanh}\left(\frac{a \tan(c + dx)}{\sqrt{3}}\right)}{\sqrt[3]{2} \sqrt[3]{a}} - \frac{3 \log\left(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)}\right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \frac{\log(a - ia \tan(c + dx))}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d}
 \end{aligned}$$

input `Int[(a + I*a*Tan[c + d*x])^(2/3)*(A + B*Tan[c + d*x]),x]`

output `((-I)*a*(A - I*B)*(((-I)*Sqrt[3]*ArcTanh[(a*Tan[c + d*x])/Sqrt[3]])/(2^(1/3)*a^(1/3)) - (3*Log[2^(1/3)*a^(1/3) - I*a*Tan[c + d*x]]/(2*2^(1/3)*a^(1/3)) + Log[a - I*a*Tan[c + d*x]]/(2*2^(1/3)*a^(1/3))))/d + (3*B*(a + I*a*Tan[c + d*x])^(2/3))/(2*d)`

3.199. $\int (a + ia \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx$

3.199.3.1 Defintions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$
- rule 67 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))^{1/3}), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[(b*c - a*d)/b, 3]\}, \text{Simp}[-\text{Log}[\text{RemoveContent}[a + b*x, x]]/(2*b*q), x] + (\text{Simp}[3/(2*b) \text{ Subst}[\text{Int}[1/(q^2 + q*x + x^2), x], x, (c + d*x)^{1/3}], x] - \text{Simp}[3/(2*b*q) \text{ Subst}[\text{Int}[1/(q - x), x], x, (c + d*x)^{1/3}], x])] / \text{ ; FreeQ}[\{a, b, c, d\}, x] \&\& \text{PosQ}[(b*c - a*d)/b]$
- rule 217 $\text{Int}(((a_)+(b_)*(x_)^2)^{-1}), x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] \text{ ; FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \& \& (\text{LtQ}[a, 0] \text{ || } \text{LtQ}[b, 0])]$
- rule 1082 $\text{Int}(((a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}), x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S\text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] \text{ ; RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \text{ || } \text{!RationalQ}[b^2 - 4*a*c])] \text{ ; FreeQ}[\{a, b, c\}, x]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$
- rule 3962 $\text{Int}(((a_)+(b_)*\text{tan}[(c_)+(d_)*(x_)])^{(n_)}), x_Symbol] \rightarrow \text{Simp}[-b/d \text{ Subst}[\text{Int}[(a + x)^{(n - 1)}/(a - x), x], x, b*\text{Tan}[c + d*x]], x] \text{ ; FreeQ}[\{a, b, c, d, n\}, x] \&\& \text{EqQ}[a^2 + b^2, 0]$
- rule 4010 $\text{Int}(((a_)+(b_)*\text{tan}[(e_)+(f_)*(x_)])^{(m_)}*((c_)+(d_)*\text{tan}[(e_)+(f_)*(x_)])), x_Symbol] \rightarrow \text{Simp}[d*((a + b*\text{Tan}[e + f*x])^m/(f*m)), x] + \text{Simp}[(b*c + a*d)/b \text{ Int}[(a + b*\text{Tan}[e + f*x])^m, x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, m\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[a^2 + b^2, 0] \&\& \text{!LtQ}[m, 0]$

3.199.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.79

method	result
derivativedivides	$3i \left(-\frac{iB(a+ia \tan(dx+c))^{\frac{2}{3}}}{2} + \frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{6a^{\frac{1}{3}}} - \frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}}(a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}}\right)}{12a^{\frac{1}{3}}} \right) d$
default	$3i \left(-\frac{iB(a+ia \tan(dx+c))^{\frac{2}{3}}}{2} + \frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{6a^{\frac{1}{3}}} - \frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}}(a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}}\right)}{12a^{\frac{1}{3}}} \right) d$
parts	$3iAa \left(\frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{6a^{\frac{1}{3}}} - \frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}}(a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}}\right)}{12a^{\frac{1}{3}}} \right) + \frac{\sqrt{3} 2^{\frac{2}{3}} \arctan\left(\frac{\sqrt{3} \left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{2^{\frac{1}{3}} a^{\frac{1}{3}} + (a+ia \tan(dx+c))^{\frac{1}{3}}}\right)}{d}$

input `int((a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `3*I/d*(-1/2*I*B*(a+I*a*tan(d*x+c))^(2/3)+(1/6*2^(2/3)/a^(1/3)*ln((a+I*a*tan(d*x+c))^(1/3)-2^(1/3)*a^(1/3))-1/12*2^(2/3)/a^(1/3)*ln((a+I*a*tan(d*x+c))^(2/3)+2^(1/3)*a^(1/3)*(a+I*a*tan(d*x+c))^(1/3)+2^(2/3)*a^(2/3))+1/6*3^(1/2)*2^(2/3)/a^(1/3)*arctan(1/3*3^(1/2)*(2^(2/3)/a^(1/3)*(a+I*a*tan(d*x+c))^(1/3)+1)))*a*(A-I*B)`

3.199.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 489 vs. $2(146) = 292$.

Time = 0.27 (sec) , antiderivative size = 489, normalized size of antiderivative = 2.42

$$\int (a + ia \tan(c + dx))^{2/3} (A$$

$$+ B \tan(c + dx)) dx = \frac{3 \cdot 2^{\frac{2}{3}} B \left(\frac{a}{e^{(2i dx + 2i c)} + 1} \right)^{\frac{2}{3}} e^{\left(\frac{4}{3} i dx + \frac{4}{3} i c\right)} + 2 \left(\frac{1}{2}\right)^{\frac{1}{3}} d \left(\frac{(-i A^3 - 3 A^2 B + 3 i A B^2 + B^3) a^2}{d^3} \right)^{\frac{1}{3}} \log \left(\frac{2^{\frac{1}{3}} (A^2 - 2 i A B - B^2) a (a / (e^{(2 i d x + 2 i c)} + 1))^{\frac{1}{3}} e^{(2/3 i d x + 2/3 i c)} + 2 (1/2)^{\frac{2}{3}} d^2 ((-i A^3 - 3 A^2 B + 3 i A B^2 + B^3) a^2 / d^3)^{\frac{1}{3}}}{(A^2 - 2 i A B - B^2) a} \right) + (1/2)^{\frac{1}{3}} (i \sqrt{3} d - d) ((-i A^3 - 3 A^2 B + 3 i A B^2 + B^3) a^2 / d^3)^{\frac{1}{3}} \log \left(2^{\frac{1}{3}} (A^2 - 2 i A B - B^2) a (a / (e^{(2 i d x + 2 i c)} + 1))^{\frac{1}{3}} e^{(2/3 i d x + 2/3 i c)} - (1/2)^{\frac{2}{3}} (i \sqrt{3} d^2 + d^2) ((-i A^3 - 3 A^2 B + 3 i A B^2 + B^3) a^2 / d^3)^{\frac{1}{3}}}{(A^2 - 2 i A B - B^2) a} \right) + (1/2)^{\frac{1}{3}} (-i \sqrt{3} d - d) ((-i A^3 - 3 A^2 B + 3 i A B^2 + B^3) a^2 / d^3)^{\frac{1}{3}} \log \left(2^{\frac{1}{3}} (A^2 - 2 i A B - B^2) a (a / (e^{(2 i d x + 2 i c)} + 1))^{\frac{1}{3}} e^{(2/3 i d x + 2/3 i c)} - (1/2)^{\frac{2}{3}} (-i \sqrt{3} d^2 + d^2) ((-i A^3 - 3 A^2 B + 3 i A B^2 + B^3) a^2 / d^3)^{\frac{1}{3}}}{(A^2 - 2 i A B - B^2) a} \right)}{d}$$

input `integrate((a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/2*(3*2^(2/3)*B*(a/(e^(2*I*d*x + 2*I*c) + 1))^(2/3)*e^(4/3*I*d*x + 4/3*I*c) + 2*(1/2)^(1/3)*d*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)*a^2/d^3)^(1/3)*log((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) + 2*(1/2)^(2/3)*d^2*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)*a^2/d^3)^(2/3))/((A^2 - 2*I*A*B - B^2)*a)) + (1/2)^(1/3)*(I*sqrt(3)*d - d)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)*a^2/d^3)^(1/3)*log((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) - (1/2)^(2/3)*(I*sqrt(3)*d^2 + d^2)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)*a^2/d^3)^(2/3))/((A^2 - 2*I*A*B - B^2)*a)) + (1/2)^(1/3)*(-I*sqrt(3)*d - d)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)*a^2/d^3)^(1/3)*log((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) - (1/2)^(2/3)*(-I*sqrt(3)*d^2 + d^2)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)*a^2/d^3)^(2/3))/((A^2 - 2*I*A*B - B^2)*a)))/d`

3.199.6 Sympy [F]

$$\int (a + ia \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{\frac{2}{3}} (A + B \tan(c + dx)) dx$$

input `integrate((a+I*a*tan(d*x+c))**(2/3)*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(2/3)*(A + B*tan(c + d*x)), x)`

3.199. $\int (a + ia \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx$

3.199.7 Maxima [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.83

$$\int (a + ia \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx = \frac{i \left(2 \sqrt{3} 2^{2/3} (A - iB) a^{5/3} \arctan \left(\frac{\sqrt{3} 2^{2/3} (2^{1/3} a^{1/3} + 2(i a \tan(dx+c) + a)^{1/3})}{6 a^{1/3}} \right) - 2^{2/3} (A - iB) a^{5/3} \log \left(2 \right) \right)}{6 a^{1/3}}$$

input `integrate((a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/4*I*(2*sqrt(3)*2^(2/3)*(A - I*B)*a^(5/3)*arctan(1/6*sqrt(3)*2^(2/3)*(2^(1/3)*a^(1/3) + 2*(I*a*tan(d*x + c) + a)^(1/3))/a^(1/3)) - 2^(2/3)*(A - I*B)*a^(5/3)*log(2^(2/3)*a^(2/3) + 2^(1/3)*(I*a*tan(d*x + c) + a)^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(2/3)) + 2*2^(2/3)*(A - I*B)*a^(5/3)*log(-2^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(1/3)) - 6*I*(I*a*tan(d*x + c) + a)^(2/3)*B*a)/(a*d)`

3.199.8 Giac [F]

$$\int (a + ia \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(i a \tan(dx + c) + a)^{2/3} dx$$

input `integrate((a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(2/3), x)`

3.199.9 Mupad [B] (verification not implemented)

Time = 9.38 (sec) , antiderivative size = 367, normalized size of antiderivative = 1.82

$$\begin{aligned}
& \int (a + ia \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx = \frac{3B(a + a \tan(c + dx) li)^{2/3}}{2d} \\
& + \frac{2^{2/3} B a^{2/3} \ln \left((a(1 + \tan(c + dx) li))^{1/3} - 2^{1/3} a^{1/3} \right)}{2d} \\
& - \frac{\left(\frac{1}{2}i\right)^{1/3} A a^{2/3} \ln \left((a(1 + \tan(c + dx) li))^{1/3} + (-1)^{1/3} 2^{1/3} a^{1/3} \right)}{d} \\
& - \frac{\left(\frac{1}{2}i\right)^{1/3} A a^{2/3} \ln \left(\frac{(-1)^{1/3} 2^{1/3} a^{1/3}}{2} - (a(1 + \tan(c + dx) li))^{1/3} + \frac{(-1)^{5/6} 2^{1/3} \sqrt{3} a^{1/3}}{2} \right) \left(-\frac{1}{2} + \frac{\sqrt{3} li}{2} \right)}{d} \\
& + \frac{2^{2/3} B a^{2/3} \ln \left(\frac{9B^2 a^2 (a + a \tan(c + dx) li)^{1/3}}{d^2} - \frac{9 \cdot 2^{1/3} B^2 a^{7/3} \left(-\frac{1}{2} + \frac{\sqrt{3} li}{2} \right)^2}{d^2} \right) \left(-\frac{1}{2} + \frac{\sqrt{3} li}{2} \right)}{2d} \\
& - \frac{2^{2/3} B a^{2/3} \ln \left(\frac{9B^2 a^2 (a + a \tan(c + dx) li)^{1/3}}{d^2} - \frac{9 \cdot 2^{1/3} B^2 a^{7/3} \left(\frac{1}{2} + \frac{\sqrt{3} li}{2} \right)^2}{d^2} \right) \left(\frac{1}{2} + \frac{\sqrt{3} li}{2} \right)}{2d} \\
& + \frac{\left(\frac{1}{2}i\right)^{1/3} A a^{2/3} \ln \left((a(1 + \tan(c + dx) li))^{1/3} - \frac{(-1)^{1/3} 2^{1/3} a^{1/3}}{2} + \frac{(-1)^{5/6} 2^{1/3} \sqrt{3} a^{1/3}}{2} \right) \left(\frac{1}{2} + \frac{\sqrt{3} li}{2} \right)}{d}
\end{aligned}$$

input `int((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^(2/3),x)`

```

output (3*B*(a + a*tan(c + d*x)*li)^(2/3))/(2*d) + (2^(2/3)*B*a^(2/3)*log((a*(tan
(c + d*x)*li + 1))^(1/3) - 2^(1/3)*a^(1/3)))/(2*d) - ((1i/2)^(1/3)*A*a^(2/
3)*log((a*(tan(c + d*x)*li + 1))^(1/3) + (-1)^(1/3)*2^(1/3)*a^(1/3)))/d -
((1i/2)^(1/3)*A*a^(2/3)*log(((1)^(1/3)*2^(1/3)*a^(1/3))/2 - (a*(tan(c + d
*x)*li + 1))^(1/3) + ((1)^(5/6)*2^(1/3)*3^(1/2)*a^(1/3))/2)*((3^(1/2)*li
/2 - 1/2))/d + (2^(2/3)*B*a^(2/3)*log((9*B^2*a^2*(a + a*tan(c + d*x)*li)^(
1/3))/d^2 - (9*2^(1/3)*B^2*a^(7/3)*((3^(1/2)*li)/2 - 1/2)^2)/d^2)*((3^(1/2
)*li)/2 - 1/2))/(2*d) - (2^(2/3)*B*a^(2/3)*log((9*B^2*a^2*(a + a*tan(c + d
*x)*li)^(1/3))/d^2 - (9*2^(1/3)*B^2*a^(7/3)*((3^(1/2)*li)/2 + 1/2)^2)/d^2
)*((3^(1/2)*li)/2 + 1/2))/(2*d) + ((1i/2)^(1/3)*A*a^(2/3)*log((a*(tan(c + d
*x)*li + 1))^(1/3) - ((1)^(1/3)*2^(1/3)*a^(1/3))/2 + ((1)^(5/6)*2^(1/3)*
3^(1/2)*a^(1/3))/2)*((3^(1/2)*li)/2 + 1/2))/d

```

3.200 $\int \cot(c+dx)(a+ia \tan(c+dx))^{2/3}(A+B \tan(c+dx)) dx$

3.200.1 Optimal result	2045
3.200.2 Mathematica [C] (verified)	2046
3.200.3 Rubi [A] (warning: unable to verify)	2046
3.200.4 Maple [A] (verified)	2050
3.200.5 Fricas [B] (verification not implemented)	2051
3.200.6 Sympy [F]	2052
3.200.7 Maxima [A] (verification not implemented)	2053
3.200.8 Giac [F]	2053
3.200.9 Mupad [B] (verification not implemented)	2054

3.200.1 Optimal result

Integrand size = 34, antiderivative size = 289

$$\int \cot(c+dx)(a+ia \tan(c+dx))^{2/3}(A+B \tan(c+dx)) dx =$$

$$-\frac{a^{2/3}(iA+B)x}{2\sqrt[3]{2}} + \frac{\sqrt{3}a^{2/3}A \arctan\left(\frac{\sqrt[3]{a+2}\sqrt[3]{a+ia \tan(c+dx)}}{\sqrt{3}\sqrt[3]{a}}\right)}{d}$$

$$-\frac{\sqrt{3}a^{2/3}(A-iB) \arctan\left(\frac{\sqrt[3]{a+2}\sqrt[3]{a+ia \tan(c+dx)}}{\sqrt{3}\sqrt[3]{a}}\right)}{\sqrt[3]{2}d}$$

$$-\frac{a^{2/3}(A-iB) \log(\cos(c+dx))}{2\sqrt[3]{2}d} - \frac{a^{2/3}A \log(\tan(c+dx))}{2d}$$

$$+ \frac{3a^{2/3}A \log\left(\sqrt[3]{a} - \sqrt[3]{a+ia \tan(c+dx)}\right)}{2d}$$

$$-\frac{3a^{2/3}(A-iB) \log\left(\sqrt[3]{2}\sqrt[3]{a} - \sqrt[3]{a+ia \tan(c+dx)}\right)}{2\sqrt[3]{2}d}$$

output

```
-1/4*a^(2/3)*(I*A+B)*x*2^(2/3)-1/4*a^(2/3)*(A-I*B)*ln(cos(d*x+c))*2^(2/3)/
d-1/2*a^(2/3)*A*ln(tan(d*x+c))/d+3/2*a^(2/3)*A*ln(a^(1/3)-(a+I*a*tan(d*x+c)
)^(1/3))/d-3/4*a^(2/3)*(A-I*B)*ln(2^(1/3)*a^(1/3)-(a+I*a*tan(d*x+c)^(1/3)
))*2^(2/3)/d+a^(2/3)*A*arctan(1/3*(a^(1/3)+2*(a+I*a*tan(d*x+c)^(1/3))/a^(
1/3)*3^(1/2))*3^(1/2)/d-1/2*a^(2/3)*(A-I*B)*arctan(1/3*(a^(1/3)+2^(2/3)*(a
+I*a*tan(d*x+c)^(1/3))/a^(1/3)*3^(1/2))*3^(1/2)*2^(2/3)/d
```

3.200.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 4.29 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.44

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \frac{3 \left(\frac{ae^{2i(c+dx)}}{1+e^{2i(c+dx)}} \right)^{2/3} \left((A - iB) \text{Hypergeometric2F1} \left(\frac{2}{3}, 1, \frac{5}{3}, \frac{e^{2i(c+dx)}}{1+e^{2i(c+dx)}} \right) - 2A \text{Hypergeometric} \right)}{2\sqrt[3]{2d}}$$

input `Integrate[Cot[c + d*x]*(a + I*a*Tan[c + d*x])^(2/3)*(A + B*Tan[c + d*x]),x]`

output `(3*((a*E^((2*I)*(c + d*x)))/(1 + E^((2*I)*(c + d*x))))^(2/3)*((A - I*B)*Hypergeometric2F1[2/3, 1, 5/3, E^((2*I)*(c + d*x))/(1 + E^((2*I)*(c + d*x)))] - 2*A*Hypergeometric2F1[2/3, 1, 5/3, (2*E^((2*I)*(c + d*x)))/(1 + E^((2*I)*(c + d*x)))]))/(2*2^(1/3)*d)`

3.200.3 Rubi [A] (warning: unable to verify)

Time = 0.76 (sec) , antiderivative size = 218, normalized size of antiderivative = 0.75, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.382$, Rules used = {3042, 4083, 3042, 3962, 67, 16, 1082, 217, 4082, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx$$

↓ 3042

$$\int \frac{(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx))}{\tan(c + dx)} dx$$

↓ 4083

$$(B + iA) \int (i \tan(c + dx)a + a)^{2/3} dx + \frac{A \int \cot(c + dx)(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^{2/3} dx}{a}$$

↓ 3042

3.200. $\int \cot(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& (B + iA) \int (i \tan(c + dx)a + a)^{2/3} dx + \frac{A \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^{2/3}}{\tan(c + dx)} dx}{a} \\
& \quad \downarrow \text{3962} \\
& \frac{A \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^{2/3}}{\tan(c + dx)} dx}{a} - \\
& \frac{ia(B + iA) \int \frac{1}{(a - ia \tan(c + dx)) \sqrt[3]{i \tan(c + dx)a + a}} d(i a \tan(c + dx))}{d} \\
& \quad \downarrow \text{67} \\
& \frac{A \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^{2/3}}{\tan(c + dx)} dx}{a} - \\
& \frac{ia(B + iA) \left(-\frac{3}{2} \int \frac{1}{-a^2 \tan^2(c + dx) + i \sqrt[3]{2} a^{4/3} \tan(c + dx) + 2^{2/3} a^{2/3}} d \sqrt[3]{i \tan(c + dx)a + a} + \frac{3 \int \frac{1}{\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)}} d \sqrt[3]{i \tan(c + dx)a + a}}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d} \\
& \quad \downarrow \text{16} \\
& \frac{A \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^{2/3}}{\tan(c + dx)} dx}{a} - \\
& \frac{ia(B + iA) \left(-\frac{3}{2} \int \frac{1}{-a^2 \tan^2(c + dx) + i \sqrt[3]{2} a^{4/3} \tan(c + dx) + 2^{2/3} a^{2/3}} d \sqrt[3]{i \tan(c + dx)a + a} - \frac{3 \log \left(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)} \right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \log \right)}{d} \\
& \quad \downarrow \text{1082} \\
& \frac{A \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^{2/3}}{\tan(c + dx)} dx}{a} - \\
& \frac{ia(B + iA) \left(\frac{3 \int \frac{1}{a^2 \tan^2(c + dx) - 3} d(i 2^{2/3} a^{2/3} \tan(c + dx) + 1)}{\sqrt[3]{2} \sqrt[3]{a}} - \frac{3 \log \left(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)} \right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \frac{\log(a - ia \tan(c + dx))}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d} \\
& \quad \downarrow \text{217} \\
& \frac{A \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^{2/3}}{\tan(c + dx)} dx}{a} - \\
& \frac{ia(B + iA) \left(-\frac{i \sqrt{3} \operatorname{arctanh} \left(\frac{a \tan(c + dx)}{\sqrt{3}} \right)}{\sqrt[3]{2} \sqrt[3]{a}} - \frac{3 \log \left(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)} \right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \frac{\log(a - ia \tan(c + dx))}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d} \\
& \quad \downarrow \text{4082}
\end{aligned}$$

3.200. $\int \cot(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx$

$$\frac{aA \int \frac{\cot(c+dx)}{\sqrt[3]{i \tan(c+dx)a+a}} d \tan(c+dx)}{ia(B+iA) \left(-\frac{i\sqrt{3} \operatorname{arctanh}\left(\frac{a \tan(c+dx)}{\sqrt{3}}\right)}{\sqrt[3]{2}\sqrt[3]{a}} - \frac{3 \log\left(\sqrt[3]{2}\sqrt[3]{a}-ia \tan(c+dx)\right)}{2\sqrt[3]{2}\sqrt[3]{a}} + \frac{\log(a-ia \tan(c+dx))}{2\sqrt[3]{2}\sqrt[3]{a}} \right)}$$

\downarrow 67

$$\frac{aA \left(\frac{3}{2} \int \frac{1}{a^{2/3} + \sqrt[3]{i \tan(c+dx)a+a} \sqrt[3]{a+(i \tan(c+dx)a+a)^{2/3}}} d \sqrt[3]{i \tan(c+dx)a+a} - \frac{3 \int \frac{1}{\sqrt[3]{a}-\sqrt[3]{i \tan(c+dx)a+a}} d \sqrt[3]{a}}{2\sqrt[3]{a}} \right)}{ia(B+iA) \left(-\frac{i\sqrt{3} \operatorname{arctanh}\left(\frac{a \tan(c+dx)}{\sqrt{3}}\right)}{\sqrt[3]{2}\sqrt[3]{a}} - \frac{3 \log\left(\sqrt[3]{2}\sqrt[3]{a}-ia \tan(c+dx)\right)}{2\sqrt[3]{2}\sqrt[3]{a}} + \frac{\log(a-ia \tan(c+dx))}{2\sqrt[3]{2}\sqrt[3]{a}} \right)}$$

\downarrow 16

$$\frac{aA \left(\frac{3}{2} \int \frac{1}{a^{2/3} + \sqrt[3]{i \tan(c+dx)a+a} \sqrt[3]{a+(i \tan(c+dx)a+a)^{2/3}}} d \sqrt[3]{i \tan(c+dx)a+a} - \frac{\log(\tan(c+dx))}{2\sqrt[3]{a}} + \frac{3 \log\left(\sqrt[3]{a}-\sqrt[3]{a+i \tan(c+dx)a+a}\right)}{2\sqrt[3]{a}} \right)}{ia(B+iA) \left(-\frac{i\sqrt{3} \operatorname{arctanh}\left(\frac{a \tan(c+dx)}{\sqrt{3}}\right)}{\sqrt[3]{2}\sqrt[3]{a}} - \frac{3 \log\left(\sqrt[3]{2}\sqrt[3]{a}-ia \tan(c+dx)\right)}{2\sqrt[3]{2}\sqrt[3]{a}} + \frac{\log(a-ia \tan(c+dx))}{2\sqrt[3]{2}\sqrt[3]{a}} \right)}$$

\downarrow 1082

$$\frac{aA \left(-\frac{3 \int \frac{1}{-(i \tan(c+dx)a+a)^{2/3}-3} d \left(\frac{2\sqrt[3]{i \tan(c+dx)a+a}}{\sqrt[3]{a}} + 1 \right)}{\sqrt[3]{a}} - \frac{\log(\tan(c+dx))}{2\sqrt[3]{a}} + \frac{3 \log\left(\sqrt[3]{a}-\sqrt[3]{a+i \tan(c+dx)a+a}\right)}{2\sqrt[3]{a}} \right)}{ia(B+iA) \left(-\frac{i\sqrt{3} \operatorname{arctanh}\left(\frac{a \tan(c+dx)}{\sqrt{3}}\right)}{\sqrt[3]{2}\sqrt[3]{a}} - \frac{3 \log\left(\sqrt[3]{2}\sqrt[3]{a}-ia \tan(c+dx)\right)}{2\sqrt[3]{2}\sqrt[3]{a}} + \frac{\log(a-ia \tan(c+dx))}{2\sqrt[3]{2}\sqrt[3]{a}} \right)}$$

\downarrow 217

3.200. $\int \cot(c+dx)(a+ia \tan(c+dx))^{2/3}(A+B \tan(c+dx)) dx$

$$\frac{aA \left(\frac{\sqrt{3} \arctan \left(\frac{1 + \sqrt[3]{2} \sqrt[3]{a + ia \tan(c + dx)}}{\sqrt[3]{a}} \right)}{\sqrt[3]{a}} \right) - \frac{\log(\tan(c + dx))}{2\sqrt[3]{a}} + \frac{3 \log \left(\sqrt[3]{a} - \sqrt[3]{a + ia \tan(c + dx)} \right)}{2\sqrt[3]{a}}}{d}$$

$$\frac{ia(B + ia) \left(-\frac{i\sqrt{3} \operatorname{arctanh} \left(\frac{a \tan(c + dx)}{\sqrt{3}} \right)}{\sqrt[3]{2} \sqrt[3]{a}} - \frac{3 \log \left(\sqrt[3]{2} \sqrt[3]{a} - ia \tan(c + dx) \right)}{2\sqrt[3]{2} \sqrt[3]{a}} + \frac{\log(a - ia \tan(c + dx))}{2\sqrt[3]{2} \sqrt[3]{a}} \right)}{d}$$

input `Int[Cot[c + d*x]*(a + I*a*Tan[c + d*x])^(2/3)*(A + B*Tan[c + d*x]),x]`

output `((-I)*a*(I*A + B)*((-I)*Sqrt[3]*ArcTanh[(a*Tan[c + d*x])/Sqrt[3]]/(2^(1/3)*a^(1/3)) - (3*Log[2^(1/3)*a^(1/3) - I*a*Tan[c + d*x]]/(2*2^(1/3)*a^(1/3)) + Log[a - I*a*Tan[c + d*x]]/(2*2^(1/3)*a^(1/3))))/d + (a*A*((Sqrt[3]*ArcTan[(1 + (2*(a + I*a*Tan[c + d*x])^(1/3))/a^(1/3))/Sqrt[3]]/a^(1/3) - Log[Tan[c + d*x]]/(2*a^(1/3)) + (3*Log[a^(1/3) - (a + I*a*Tan[c + d*x])^(1/3)]/(2*a^(1/3)))/d`

3.200.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 67 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(1/3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q), x] + (Simp[3/(2*b) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Simp[3/(2*b*q) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x]) /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`

rule 217 `Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])]; FreeQ[{a, b, c}, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3962 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[-b/d Subst[Int[(a + x)^(n - 1)/(a - x), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 + b^2, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int[(((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.200.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 248, normalized size of antiderivative = 0.86

method	result
derivativedivides	$3a \left(\frac{2^{\frac{2}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}} \right)}{6a^{\frac{1}{3}}} - \frac{2^{\frac{2}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}} (a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}} \right)}{12a^{\frac{1}{3}}} + \frac{\sqrt{3} 2^{\frac{2}{3}} \arctan \left(\frac{\sqrt{3}}{\dots} \right)}{\dots} \right)$
default	$3a \left(\frac{2^{\frac{2}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}} \right)}{6a^{\frac{1}{3}}} - \frac{2^{\frac{2}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}} (a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}} \right)}{12a^{\frac{1}{3}}} + \frac{\sqrt{3} 2^{\frac{2}{3}} \arctan \left(\frac{\sqrt{3}}{\dots} \right)}{\dots} \right)$

```
input int(cot(d*x+c)*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 3/d*a*((1/6*2^(2/3)/a^(1/3)*ln((a+I*a*tan(d*x+c))^(1/3)-2^(1/3)*a^(1/3))-1
/12*2^(2/3)/a^(1/3)*ln((a+I*a*tan(d*x+c))^(2/3)+2^(1/3)*a^(1/3)*(a+I*a*tan
(d*x+c))^(1/3)+2^(2/3)*a^(2/3))+1/6*3^(1/2)*2^(2/3)/a^(1/3)*arctan(1/3*3^(
1/2)*(2^(2/3)/a^(1/3)*(a+I*a*tan(d*x+c))^(1/3)+1)))*(-A+I*B)+(1/3/a^(1/3)*
ln((a+I*a*tan(d*x+c))^(1/3)-a^(1/3))-1/6/a^(1/3)*ln((a+I*a*tan(d*x+c))^(2/
3)+a^(1/3)*(a+I*a*tan(d*x+c))^(1/3)+a^(2/3))+1/3*3^(1/2)/a^(1/3)*arctan(1/
3*3^(1/2)*(2/a^(1/3)*(a+I*a*tan(d*x+c))^(1/3)+1)))*A)
```

3.200.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 711 vs. 2(211) = 422.

Time = 0.27 (sec) , antiderivative size = 711, normalized size of antiderivative = 2.46

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorith
m="fricas")
```

output

```

1/2*(1/2)^(1/3)*(-I*sqrt(3) - 1)*(-(A^3 - 3*I*A^2*B - 3*A*B^2 + I*B^3)*a^2
/d^3)^(1/3)*log((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I*c) +
1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) - (1/2)^(2/3)*(I*sqrt(3)*d^2 - d^2)*(-(
A^3 - 3*I*A^2*B - 3*A*B^2 + I*B^3)*a^2/d^3)^(2/3))/((A^2 - 2*I*A*B - B^2)*
a)) + 1/2*(1/2)^(1/3)*(I*sqrt(3) - 1)*(-(A^3 - 3*I*A^2*B - 3*A*B^2 + I*B^3
)*a^2/d^3)^(1/3)*log((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I
*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) - (1/2)^(2/3)*(-I*sqrt(3)*d^2 - d^
2)*(-(A^3 - 3*I*A^2*B - 3*A*B^2 + I*B^3)*a^2/d^3)^(2/3))/((A^2 - 2*I*A*B -
B^2)*a)) + 1/2*(A^3*a^2/d^3)^(1/3)*(I*sqrt(3) - 1)*log(1/2*(2*2^(1/3)*A^2
*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) + (I*sqrt(3
)*d^2 + d^2)*(A^3*a^2/d^3)^(2/3))/(A^2*a)) + 1/2*(A^3*a^2/d^3)^(1/3)*(-I*s
qrt(3) - 1)*log(1/2*(2*2^(1/3)*A^2*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e
^(2/3*I*d*x + 2/3*I*c) + (-I*sqrt(3)*d^2 + d^2)*(A^3*a^2/d^3)^(2/3))/(A^2*
a)) + (1/2)^(1/3)*(-(A^3 - 3*I*A^2*B - 3*A*B^2 + I*B^3)*a^2/d^3)^(1/3)*log
((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2
/3*I*d*x + 2/3*I*c) - 2*(1/2)^(2/3)*d^2*(-(A^3 - 3*I*A^2*B - 3*A*B^2 + I*B
^3)*a^2/d^3)^(2/3))/((A^2 - 2*I*A*B - B^2)*a)) + (A^3*a^2/d^3)^(1/3)*log((
2^(1/3)*A^2*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c)
- (A^3*a^2/d^3)^(2/3)*d^2)/(A^2*a))

```

3.200.6 Sympy [F]

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{2/3}(A + B \tan(c + dx)) \cot(c + dx) dx$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))**(2/3)*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(2/3)*(A + B*tan(c + d*x))*cot(c + d*x), x)`

3.200.7 Maxima [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 252, normalized size of antiderivative = 0.87

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx =$$

$$2\sqrt{3}2^{2/3}(A - iB)a^{2/3} \arctan\left(\frac{\sqrt{3}2^{2/3}(2^{1/3}a^{1/3} + 2(i a \tan(dx+c)+a)^{1/3})}{6a^{1/3}}\right) - 2^{2/3}(A - iB)a^{2/3} \log\left(2^{2/3}a^{2/3} + 2^{1/3}(i a \tan(dx + c) + a)^{1/3}\right)$$

```
input integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm
m="maxima")
```

```
output -1/4*(2*sqrt(3)*2^(2/3)*(A - I*B)*a^(2/3)*arctan(1/6*sqrt(3)*2^(2/3)*(2^(1/3)*a^(1/3) + 2*(I*a*tan(d*x + c) + a)^(1/3))/a^(1/3)) - 2^(2/3)*(A - I*B)*a^(2/3)*log(2^(2/3)*a^(2/3) + 2^(1/3)*(I*a*tan(d*x + c) + a)^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(2/3)) + 2*2^(2/3)*(A - I*B)*a^(2/3)*log(-2^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(1/3)) - 4*sqrt(3)*A*a^(2/3)*arctan(1/3*sqrt(3)*(2*(I*a*tan(d*x + c) + a)^(1/3) + a^(1/3))/a^(1/3)) + 2*A*a^(2/3)*log((I*a*tan(d*x + c) + a)^(2/3) + (I*a*tan(d*x + c) + a)^(1/3)*a^(1/3) + a^(2/3)) - 4*A*a^(2/3)*log((I*a*tan(d*x + c) + a)^(1/3) - a^(1/3))/d
```

3.200.8 Giac [F]

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(i a \tan(dx + c) + a)^{2/3} \cot(dx + c) dx$$

```
input integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm
m="giac")
```

```
output integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(2/3)*cot(d*x + c),
x)
```

3.200.9 Mupad [B] (verification not implemented)

Time = 8.45 (sec) , antiderivative size = 1761, normalized size of antiderivative = 6.09

$$\int \cot(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(2/3),x)
```

```
output log(- (486*d^3*(3*A*B^2*a^9 - B^3*a^9*1i + A^2*B*a^9*3i) - (1458*a^7*d^6*(
(A^3*a^2)/d^3)^(2/3) + 243*d*(a + a*tan(c + d*x)*1i)^(1/3)*(B^2*a^8*d^3 -
5*A^2*a^8*d^3 + A*B*a^8*d^3*2i))*((A^3*a^2)/d^3)^(1/3))*((A^3*a^2)/d^3)^(2
/3) - 243*d*(a + a*tan(c + d*x)*1i)^(1/3)*(A^5*a^10 - A^4*B*a^10*4i + A^2*
B^3*a^10*2i - 5*A^3*B^2*a^10))*((A^3*a^2)/d^3)^(1/3) + log(- (486*d^3*(3*A
*B^2*a^9 - B^3*a^9*1i + A^2*B*a^9*3i) - (1458*a^7*d^6*(-(A^3*a^2 + B^3*a^2
*1i - 3*A*B^2*a^2 - A^2*B*a^2*3i)/(2*d^3))^(2/3) + 243*d*(a + a*tan(c + d*
x)*1i)^(1/3)*(B^2*a^8*d^3 - 5*A^2*a^8*d^3 + A*B*a^8*d^3*2i))*(-(A^3*a^2 +
B^3*a^2*1i - 3*A*B^2*a^2 - A^2*B*a^2*3i)/(2*d^3))^(1/3))*(-(A^3*a^2 + B^3*
a^2*1i - 3*A*B^2*a^2 - A^2*B*a^2*3i)/(2*d^3))^(2/3) - 243*d*(a + a*tan(c +
d*x)*1i)^(1/3)*(A^5*a^10 - A^4*B*a^10*4i + A^2*B^3*a^10*2i - 5*A^3*B^2*a^
10))*(-(A^3*a^2 + B^3*a^2*1i - 3*A*B^2*a^2 - A^2*B*a^2*3i)/(2*d^3))^(1/3)
+ (log(- ((3^(1/2)*1i - 1)^2*(486*d^3*(3*A*B^2*a^9 - B^3*a^9*1i + A^2*B*a^
9*3i) - ((3^(1/2)*1i - 1)*(243*d*(a + a*tan(c + d*x)*1i)^(1/3)*(B^2*a^8*d^
3 - 5*A^2*a^8*d^3 + A*B*a^8*d^3*2i) + (729*a^7*d^6*(3^(1/2)*1i - 1)^2*((A^
3*a^2)/d^3)^(2/3))/2)*((A^3*a^2)/d^3)^(1/3))/2)*((A^3*a^2)/d^3)^(2/3))/4 -
243*d*(a + a*tan(c + d*x)*1i)^(1/3)*(A^5*a^10 - A^4*B*a^10*4i + A^2*B^3*a^
10*2i - 5*A^3*B^2*a^10))*((3^(1/2)*1i - 1)*((A^3*a^2)/d^3)^(1/3))/2 - (log
(- ((3^(1/2)*1i + 1)^2*(486*d^3*(3*A*B^2*a^9 - B^3*a^9*1i + A^2*B*a^9*3i)
+ ((3^(1/2)*1i + 1)*(243*d*(a + a*tan(c + d*x)*1i)^(1/3)*(B^2*a^8*d^3 - ...
```

3.201 $\int \cot^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx$

3.201.1 Optimal result	2055
3.201.2 Mathematica [A] (verified)	2056
3.201.3 Rubi [A] (warning: unable to verify)	2056
3.201.4 Maple [A] (verified)	2061
3.201.5 Fricas [B] (verification not implemented)	2063
3.201.6 Sympy [F]	2064
3.201.7 Maxima [A] (verification not implemented)	2064
3.201.8 Giac [F]	2065
3.201.9 Mupad [B] (verification not implemented)	2065

3.201.1 Optimal result

Integrand size = 36, antiderivative size = 342

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \frac{a^{2/3}(A - iB)x}{2\sqrt[3]{2}}$$

$$+ \frac{a^{2/3}(2iA + 3B) \arctan\left(\frac{\sqrt[3]{a+2}\sqrt[3]{a + ia \tan(c + dx)}}{\sqrt{3}\sqrt[3]{a}}\right)}{\sqrt{3}d}$$

$$- \frac{\sqrt{3}a^{2/3}(iA + B) \arctan\left(\frac{\sqrt[3]{a+2}^{2/3}\sqrt[3]{a + ia \tan(c + dx)}}{\sqrt{3}\sqrt[3]{a}}\right)}{\sqrt[3]{2}d}$$

$$- \frac{a^{2/3}(iA + B) \log(\cos(c + dx))}{2\sqrt[3]{2}d} - \frac{a^{2/3}(2iA + 3B) \log(\tan(c + dx))}{6d}$$

$$+ \frac{a^{2/3}(2iA + 3B) \log\left(\sqrt[3]{a} - \sqrt[3]{a + ia \tan(c + dx)}\right)}{2d}$$

$$- \frac{3a^{2/3}(iA + B) \log\left(\sqrt[3]{2}\sqrt[3]{a} - \sqrt[3]{a + ia \tan(c + dx)}\right)}{2\sqrt[3]{2}d}$$

$$- \frac{A \cot(c + dx)(a + ia \tan(c + dx))^{2/3}}{d}$$

output $\frac{1}{4}a^{2/3}(A-I*B)*x^{2^{2/3}}-1/4*a^{2/3}*(I*A+B)*\ln(\cos(d*x+c))*2^{2/3}/d$
 $-1/6*a^{2/3}*(2*I*A+3*B)*\ln(\tan(d*x+c))/d+1/2*a^{2/3}*(2*I*A+3*B)*\ln(a^{1/3})$
 $-(a+I*a*\tan(d*x+c))^{1/3}/d-3/4*a^{2/3}*(I*A+B)*\ln(2^{1/3}*a^{1/3}-(a+I$
 $*a*\tan(d*x+c))^{1/3})*2^{2/3}/d+1/3*a^{2/3}*(2*I*A+3*B)*\arctan(1/3*(a^{1/3}$
 $+2*(a+I*a*\tan(d*x+c))^{1/3})/a^{1/3}*3^{1/2})/d*3^{1/2}-1/2*a^{2/3}*(I*A$
 $+B)*\arctan(1/3*(a^{1/3}+2^{2/3}*(a+I*a*\tan(d*x+c))^{1/3})/a^{1/3}*3^{1/2})*$
 $3^{1/2}*2^{2/3}/d-A*\cot(d*x+c)*(a+I*a*\tan(d*x+c))^{2/3}/d$

3.201.2 Mathematica [A] (verified)

Time = 3.48 (sec) , antiderivative size = 251, normalized size of antiderivative = 0.73

$$\int \cot^2(c+dx)(a+ia \tan(c+dx))^{2/3}(A$$

$$2a^2(2iA+3B) \left(2\sqrt{3} \arctan \left(\frac{\sqrt[3]{a}+2\sqrt[3]{a+ia \tan(c+dx)}}{\sqrt{3}\sqrt[3]{a}} \right) - \log(\tan(c+dx)) + 3 \log$$

$$+B \tan(c+dx)) dx =$$

input `Integrate[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^(2/3)*(A + B*Tan[c + d*x]),x]`

output $(2*a^2*((2*I)*A + 3*B)*(2*\text{Sqrt}[3]*\text{ArcTan}[(a^{1/3} + 2*(a + I*a*\text{Tan}[c + d*x])^{1/3})/(\text{Sqrt}[3]*a^{1/3})]) - \text{Log}[\text{Tan}[c + d*x]] + 3*\text{Log}[a^{1/3} - (a + I*a*\text{Tan}[c + d*x])^{1/3}]) - (3*I)*2^{2/3}*a^2*(A - I*B)*(2*\text{Sqrt}[3]*\text{ArcTan}[(1 + (2^{2/3}*(a + I*a*\text{Tan}[c + d*x])^{1/3})/a^{1/3})/\text{Sqrt}[3]] - \text{Log}[I + \text{Tan}[c + d*x]]) + 3*\text{Log}[2^{1/3}*a^{1/3} - (a + I*a*\text{Tan}[c + d*x])^{1/3}]) - 12*a^{4/3}*A*\text{Cot}[c + d*x]*(a + I*a*\text{Tan}[c + d*x])^{2/3})/(12*a^{4/3}*d)$

3.201.3 Rubi [A] (warning: unable to verify)

Time = 1.04 (sec) , antiderivative size = 267, normalized size of antiderivative = 0.78, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 4081, 27, 3042, 4083, 3042, 3962, 67, 16, 1082, 217, 4082, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.201. $\int \cot^2(c+dx)(a+ia \tan(c+dx))^{2/3}(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \int \cot^2(c+dx)(a+ia \tan(c+dx))^{2/3}(A+B \tan(c+dx)) dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a+ia \tan(c+dx))^{2/3}(A+B \tan(c+dx))}{\tan(c+dx)^2} dx \\
& \quad \downarrow \text{4081} \\
& \frac{\int \frac{1}{3} \cot(c+dx)(i \tan(c+dx)a+a)^{2/3}(a(2iA+3B)-aA \tan(c+dx)) dx}{\frac{a}{d} A \cot(c+dx)(a+ia \tan(c+dx))^{2/3}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \cot(c+dx)(i \tan(c+dx)a+a)^{2/3}(a(2iA+3B)-aA \tan(c+dx)) dx}{\frac{3a}{d} A \cot(c+dx)(a+ia \tan(c+dx))^{2/3}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{(i \tan(c+dx)a+a)^{2/3}(a(2iA+3B)-aA \tan(c+dx))}{\tan(c+dx)} dx}{3a} - \frac{A \cot(c+dx)(a+ia \tan(c+dx))^{2/3}}{d} \\
& \quad \downarrow \text{4083} \\
& \frac{(3B+2iA) \int \cot(c+dx)(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^{2/3} dx - 3a(A-iB) \int (i \tan(c+dx)a+a)^{2/3} dx}{\frac{3a}{d} A \cot(c+dx)(a+ia \tan(c+dx))^{2/3}} \\
& \quad \downarrow \text{3042} \\
& \frac{(3B+2iA) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^{2/3}}{\tan(c+dx)} dx - 3a(A-iB) \int (i \tan(c+dx)a+a)^{2/3} dx}{\frac{3a}{d} A \cot(c+dx)(a+ia \tan(c+dx))^{2/3}} \\
& \quad \downarrow \text{3962} \\
& \frac{3ia^2(A-iB) \int \frac{1}{(a-ia \tan(c+dx)) \sqrt[3]{i \tan(c+dx)a+a}} d(i \tan(c+dx))}{\frac{3a}{d} A \cot(c+dx)(a+ia \tan(c+dx))^{2/3}} + (3B+2iA) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^{2/3}}{\tan(c+dx)} dx \\
& \quad \downarrow \text{67}
\end{aligned}$$

3.201. $\int \cot^2(c+dx)(a+ia \tan(c+dx))^{2/3}(A+B \tan(c+dx)) dx$

$$\frac{3ia^2(A-iB) \left(-\frac{3}{2} \int \frac{1}{-a^2 \tan^2(c+dx) + i \sqrt[3]{2} a^{4/3} \tan(c+dx) + 2^{2/3} a^{2/3}} dx \sqrt[3]{i \tan(c+dx)a+a} - \frac{3 \int \frac{1}{\sqrt[3]{2} \sqrt[3]{a-ia \tan(c+dx)}} dx \sqrt[3]{i \tan(c+dx)a+a}}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d} = \frac{3a}{d} \frac{A \cot(c+dx)(a+ia \tan(c+dx))^{2/3}}{d}$$

↓ 16

$$\frac{3ia^2(A-iB) \left(-\frac{3}{2} \int \frac{1}{-a^2 \tan^2(c+dx) + i \sqrt[3]{2} a^{4/3} \tan(c+dx) + 2^{2/3} a^{2/3}} dx \sqrt[3]{i \tan(c+dx)a+a} - \frac{3 \log \left(\sqrt[3]{2} \sqrt[3]{a-ia \tan(c+dx)} \right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \frac{\log(a-ia \tan(c+dx))}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d} = \frac{3a}{d} \frac{A \cot(c+dx)(a+ia \tan(c+dx))^{2/3}}{d}$$

↓ 1082

$$\frac{3ia^2(A-iB) \left(\frac{3 \int \frac{1}{a^2 \tan^2(c+dx) - 3} dx (i2^{2/3} a^{2/3} \tan(c+dx) + 1)}{\sqrt[3]{2} \sqrt[3]{a}} - \frac{3 \log \left(\sqrt[3]{2} \sqrt[3]{a-ia \tan(c+dx)} \right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \frac{\log(a-ia \tan(c+dx))}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d} + (3B + 2iA) \int \frac{(a-ia \tan(c+dx))}{\tan(c+dx)} dx = \frac{3a}{d} \frac{A \cot(c+dx)(a+ia \tan(c+dx))^{2/3}}{d}$$

↓ 217

$$\frac{(3B + 2iA) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^{2/3}}{\tan(c+dx)} dx + \frac{3ia^2(A-iB) \left(-\frac{i\sqrt{3} \operatorname{arctanh} \left(\frac{a \tan(c+dx)}{\sqrt{3}} \right)}{\sqrt[3]{2} \sqrt[3]{a}} - \frac{3 \log \left(\sqrt[3]{2} \sqrt[3]{a-ia \tan(c+dx)} \right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \frac{\log(a-ia \tan(c+dx))}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d}}{d} = \frac{3a}{d} \frac{A \cot(c+dx)(a+ia \tan(c+dx))^{2/3}}{d}$$

↓ 4082

$$\frac{a^2(3B+2iA) \int \frac{\cot(c+dx)}{\sqrt[3]{i \tan(c+dx)a+a}} d \tan(c+dx) + \frac{3ia^2(A-iB) \left(-\frac{i\sqrt{3} \operatorname{arctanh} \left(\frac{a \tan(c+dx)}{\sqrt{3}} \right)}{\sqrt[3]{2} \sqrt[3]{a}} - \frac{3 \log \left(\sqrt[3]{2} \sqrt[3]{a-ia \tan(c+dx)} \right)}{2 \sqrt[3]{2} \sqrt[3]{a}} + \frac{\log(a-ia \tan(c+dx))}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{d}}{d} = \frac{3a}{d} \frac{A \cot(c+dx)(a+ia \tan(c+dx))^{2/3}}{d}$$

↓ 67

3.201. $\int \cot^2(c+dx)(a+ia \tan(c+dx))^{2/3}(A+B \tan(c+dx)) dx$

$$\begin{aligned}
 & \frac{a^2(3B+2iA) \left(\frac{3}{2} \int \frac{1}{a^{2/3} + \sqrt[3]{i \tan(c+dx)a + a} \sqrt[3]{a + (i \tan(c+dx)a+a)^{2/3}}} d \sqrt[3]{i \tan(c+dx)a + a} - \frac{3 \int \frac{1}{\sqrt[3]{a} - \sqrt[3]{i \tan(c+dx)a + a}}}{2 \sqrt[3]{a}} \right)}{d} \\
 & \frac{A \cot(c+dx)(a + ia \tan(c+dx))^{2/3}}{d} \\
 & \quad \downarrow \text{16} \\
 & \frac{a^2(3B+2iA) \left(\frac{3}{2} \int \frac{1}{a^{2/3} + \sqrt[3]{i \tan(c+dx)a + a} \sqrt[3]{a + (i \tan(c+dx)a+a)^{2/3}}} d \sqrt[3]{i \tan(c+dx)a + a} - \frac{\log(\tan(c+dx))}{2 \sqrt[3]{a}} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{i \tan(c+dx)a + a})}{2 \sqrt[3]{a}} \right)}{d} \\
 & \frac{A \cot(c+dx)(a + ia \tan(c+dx))^{2/3}}{d} \qquad \qquad \qquad 3a \\
 & \quad \downarrow \text{1082} \\
 & \frac{a^2(3B+2iA) \left(\frac{3 \int \frac{1}{-(i \tan(c+dx)a+a)^{2/3}-3} d \left(\frac{2 \sqrt[3]{i \tan(c+dx)a + a}}{\sqrt[3]{a}} + 1 \right)}{\sqrt[3]{a}} - \frac{\log(\tan(c+dx))}{2 \sqrt[3]{a}} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a + ia \tan(c+dx)})}{2 \sqrt[3]{a}} \right)}{d} \\
 & \frac{A \cot(c+dx)(a + ia \tan(c+dx))^{2/3}}{d} \qquad \qquad \qquad 3a \\
 & \quad \downarrow \text{217} \\
 & \frac{a^2(3B+2iA) \left(\frac{\sqrt{3} \arctan \left(\frac{1 + \frac{2 \sqrt[3]{a + ia \tan(c+dx)}}{\sqrt[3]{a}}}{\frac{\sqrt[3]{a}}{\sqrt{3}}} \right)}{\sqrt[3]{a}} - \frac{\log(\tan(c+dx))}{2 \sqrt[3]{a}} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a + ia \tan(c+dx)})}{2 \sqrt[3]{a}} \right)}{d} \\
 & \frac{A \cot(c+dx)(a + ia \tan(c+dx))^{2/3}}{d} \qquad \qquad \qquad 3a + \frac{3ia^2(A-iB) \left(-\frac{i\sqrt{3}a}{\dots} \right)}{3a} \\
 & \frac{A \cot(c+dx)(a + ia \tan(c+dx))^{2/3}}{d}
 \end{aligned}$$

input `Int[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^(2/3)*(A + B*Tan[c + d*x]),x]`

3.201. $\int \cot^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx$

```
output ((3*I)*a^2*(A - I*B)*((-I)*Sqrt[3]*ArcTanh[(a*Tan[c + d*x])/Sqrt[3]]/(2
^(1/3)*a^(1/3)) - (3*Log[2^(1/3)*a^(1/3) - I*a*Tan[c + d*x]]/(2*2^(1/3)*a
^(1/3)) + Log[a - I*a*Tan[c + d*x]]/(2*2^(1/3)*a^(1/3))))/d + (a^2*((2*I)*
A + 3*B)*((Sqrt[3]*ArcTan[(1 + (2*(a + I*a*Tan[c + d*x])^(1/3))/a^(1/3)]/S
qrt[3]))/a^(1/3) - Log[Tan[c + d*x]]/(2*a^(1/3)) + (3*Log[a^(1/3) - (a + I
*a*Tan[c + d*x])^(1/3)]/(2*a^(1/3))))/d)/(3*a) - (A*Cot[c + d*x]*(a + I*a
*Tan[c + d*x])^(2/3))/d
```

3.201.3.1 Defintions of rubi rules used

```
rule 16 Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a +
b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

```
rule 67 Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(1/3)), x_Symbol] := With[
{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q), x
] + (Simp[3/(2*b) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)],
x] - Simp[3/(2*b*q) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x])] /
; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 3962 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[-b/d Subst[Int[(a + x)^(n - 1)/(a - x), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 + b^2, 0]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int[(((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.201.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 294, normalized size of antiderivative = 0.86

method	result
derivativedivides	$3ia^2 \left(\frac{2^{\frac{2}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}} \right)}{6a^{\frac{1}{3}}} - \frac{2^{\frac{2}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}} (a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}} \right)}{12a^{\frac{1}{3}}} + \frac{\sqrt{3} 2^{\frac{2}{3}} \arctan \left(\frac{(a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}}{(a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}}} \right)}{a} \right)$
default	$3ia^2 \left(\frac{2^{\frac{2}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}} \right)}{6a^{\frac{1}{3}}} - \frac{2^{\frac{2}{3}} \ln \left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}} (a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}} \right)}{12a^{\frac{1}{3}}} + \frac{\sqrt{3} 2^{\frac{2}{3}} \arctan \left(\frac{(a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}}{(a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}}} \right)}{a} \right)$

```
input int(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 3*I/d*a^2*(-(1/6*2^(2/3)/a^(1/3)*ln((a+I*a*tan(d*x+c))^(1/3)-2^(1/3)*a^(1/3))-1/12*2^(2/3)/a^(1/3)*ln((a+I*a*tan(d*x+c))^(2/3)+2^(1/3)*a^(1/3)*(a+I*a*tan(d*x+c))^(1/3)+2^(2/3)*a^(2/3))+1/6*3^(1/2)*2^(2/3)/a^(1/3)*arctan(1/3*3^(1/2)*(2^(2/3)/a^(1/3)*(a+I*a*tan(d*x+c))^(1/3)+1)))*(A-I*B)/a+1/a*(1/3*I*A*(a+I*a*tan(d*x+c))^(2/3)/a/tan(d*x+c)+(2/3*A-I*B)*(1/3/a^(1/3)*ln((a+I*a*tan(d*x+c))^(1/3)-a^(1/3))-1/6/a^(1/3)*ln((a+I*a*tan(d*x+c))^(2/3)+a^(1/3)*(a+I*a*tan(d*x+c))^(1/3)+a^(2/3))+1/3*3^(1/2)/a^(1/3)*arctan(1/3*3^(1/2)*(2/a^(1/3)*(a+I*a*tan(d*x+c))^(1/3)+1)))))
```

3.201. $\int \cot^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx$

3.201.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1096 vs. $2(255) = 510$.

Time = 0.30 (sec) , antiderivative size = 1096, normalized size of antiderivative = 3.20

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm="fracas")
```

```
output -1/6*(6*2^(2/3)*(I*A*e^(2*I*d*x + 2*I*c) + I*A)*(a/(e^(2*I*d*x + 2*I*c) + 1))^(2/3)*e^(4/3*I*d*x + 4/3*I*c) - 6*(1/2)^(1/3)*(d*e^(2*I*d*x + 2*I*c) - d)*((I*A^3 + 3*A^2*B - 3*I*A*B^2 - B^3)*a^2/d^3)^(1/3)*log((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) + 2*(1/2)^(2/3)*d^2*((I*A^3 + 3*A^2*B - 3*I*A*B^2 - B^3)*a^2/d^3)^(2/3)))/((A^2 - 2*I*A*B - B^2)*a) - 3*(1/2)^(1/3)*((I*sqrt(3)*d - d)*e^(2*I*d*x + 2*I*c) - I*sqrt(3)*d + d)*((I*A^3 + 3*A^2*B - 3*I*A*B^2 - B^3)*a^2/d^3)^(1/3)*log((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) - (1/2)^(2/3)*(I*sqrt(3)*d^2 + d^2)*((I*A^3 + 3*A^2*B - 3*I*A*B^2 - B^3)*a^2/d^3)^(2/3)))/((A^2 - 2*I*A*B - B^2)*a) - 3*(1/2)^(1/3)*((-I*sqrt(3)*d - d)*e^(2*I*d*x + 2*I*c) + I*sqrt(3)*d + d)*((I*A^3 + 3*A^2*B - 3*I*A*B^2 - B^3)*a^2/d^3)^(1/3)*log((2^(1/3)*(A^2 - 2*I*A*B - B^2)*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) - (1/2)^(2/3)*(-I*sqrt(3)*d^2 + d^2)*((I*A^3 + 3*A^2*B - 3*I*A*B^2 - B^3)*a^2/d^3)^(2/3)))/((A^2 - 2*I*A*B - B^2)*a) - ((-I*sqrt(3)*d - d)*e^(2*I*d*x + 2*I*c) + I*sqrt(3)*d + d)*((-8*I*A^3 - 36*A^2*B + 54*I*A*B^2 + 27*B^3)*a^2/d^3)^(1/3)*log(1/2*(2*2^(1/3)*(4*A^2 - 12*I*A*B - 9*B^2)*a*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) + (I*sqrt(3)*d^2 - d^2)*((-8*I*A^3 - 36*A^2*B + 54*I*A*B^2 + 27*B^3)*a^2/d^3)^(2/3)))/((4*A^2 - 12*I*A*B - 9*B^2)*a) - ((I*sqrt(3)*d - d)*e^(2*I*d*x + 2*I*c) - I*sqrt(...
```

3.201.6 Sympy [F]

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{2/3}(A + B \tan(c + dx)) \cot^2(c + dx) dx$$

input `integrate(cot(d*x+c)**2*(a+I*a*tan(d*x+c))**(2/3)*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(2/3)*(A + B*tan(c + d*x))*cot(c + d*x)**2, x)`

3.201.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 298, normalized size of antiderivative = 0.87

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx =$$

$$i \left(\frac{6\sqrt{3}2^{2/3}(A-iB) \arctan\left(\frac{\sqrt{3}2^{2/3}\left(2^{1/3}a^{1/3}+2(i a \tan(dx+c)+a)^{1/3}\right)}{6a^{1/3}}\right)}{a^{1/3}} - \frac{3\cdot 2^{2/3}(A-iB) \log\left(2^{2/3}a^{2/3}+2^{1/3}(i a \tan(dx+c)+a)^{1/3}a^{1/3}+(i a \tan(dx+c)+a)\right)}{a^{1/3}} \right)$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/12*I*(6*sqrt(3)*2^(2/3)*(A - I*B)*arctan(1/6*sqrt(3)*2^(2/3)*(2^(1/3)*a^(1/3) + 2*(I*a*tan(d*x + c) + a)^(1/3))/a^(1/3))/a^(1/3) - 3*2^(2/3)*(A - I*B)*log(2^(2/3)*a^(2/3) + 2^(1/3)*(I*a*tan(d*x + c) + a)^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(2/3))/a^(1/3) + 6*2^(2/3)*(A - I*B)*log(-2^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(1/3))/a^(1/3) - 4*sqrt(3)*(2*A - 3*I*B)*arctan(1/3*sqrt(3)*(2*(I*a*tan(d*x + c) + a)^(1/3) + a^(1/3))/a^(1/3))/a^(1/3) + 2*(2*A - 3*I*B)*log((I*a*tan(d*x + c) + a)^(2/3) + (I*a*tan(d*x + c) + a)^(1/3)*a^(1/3) + a^(2/3))/a^(1/3) - 4*(2*A - 3*I*B)*log((I*a*tan(d*x + c) + a)^(1/3) - a^(1/3))/a^(1/3) - 12*I*(I*a*tan(d*x + c) + a)^(2/3)*A/(a*tan(d*x + c))*a/d`

3.201.8 Giac [F]

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{2/3} \cot(dx + c)^2 dx$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(2/3)*cot(d*x + c)^2, x)`

3.201.9 Mupad [B] (verification not implemented)

Time = 9.99 (sec) , antiderivative size = 5825, normalized size of antiderivative = 17.03

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^{2/3}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(2/3),x)`

output

```

log((18*d^3*(A^3*a^9*19i - A*B^2*a^9*27i + 45*A^2*B*a^9) - (1458*a^7*d^6*(
-((d^3*(((594*A^3*a^12 + 1458*A*B^2*a^12)*1i)/d^3 + (1458*B^3*a^12 + 486*
A^2*B*a^12)/d^3)^2 - 11664*a^10*((432*A^6*a^14 - 1458*B^6*a^14 + 15066*A^2
*B^4*a^14 - 10044*A^4*B^2*a^14)/d^6 - ((7290*A*B^5*a^14 + 3240*A^5*B*a^14
- 16470*A^3*B^3*a^14)*1i)/d^6))^(1/2)*1i - 594*A^3*a^12 + B^3*a^12*1458i -
1458*A*B^2*a^12 + A^2*B*a^12*486i)*1i)/(5832*a^10*d^3))^(2/3) - 9*d*(a +
a*tan(c + d*x)*1i)^(1/3)*(135*B^2*a^8*d^3 - 75*A^2*a^8*d^3 + A*B*a^8*d^3*1
98i))*(-(d^3*(((594*A^3*a^12 + 1458*A*B^2*a^12)*1i)/d^3 + (1458*B^3*a^12
+ 486*A^2*B*a^12)/d^3)^2 - 11664*a^10*((432*A^6*a^14 - 1458*B^6*a^14 + 15
066*A^2*B^4*a^14 - 10044*A^4*B^2*a^14)/d^6 - ((7290*A*B^5*a^14 + 3240*A^5*
B*a^14 - 16470*A^3*B^3*a^14)*1i)/d^6))^(1/2)*1i - 594*A^3*a^12 + B^3*a^12*
1458i - 1458*A*B^2*a^12 + A^2*B*a^12*486i)*1i)/(5832*a^10*d^3))^(1/3))*(-(
(d^3*(((594*A^3*a^12 + 1458*A*B^2*a^12)*1i)/d^3 + (1458*B^3*a^12 + 486*A^
2*B*a^12)/d^3)^2 - 11664*a^10*((432*A^6*a^14 - 1458*B^6*a^14 + 15066*A^2*B
^4*a^14 - 10044*A^4*B^2*a^14)/d^6 - ((7290*A*B^5*a^14 + 3240*A^5*B*a^14 -
16470*A^3*B^3*a^14)*1i)/d^6))^(1/2)*1i - 594*A^3*a^12 + B^3*a^12*1458i - 1
458*A*B^2*a^12 + A^2*B*a^12*486i)*1i)/(5832*a^10*d^3))^(2/3) + 9*d*(a + a*
tan(c + d*x)*1i)^(1/3)*(A^5*a^10*16i + 27*B^5*a^10 + A*B^4*a^10*126i + 92*
A^4*B*a^10 - 231*A^2*B^3*a^10 - A^3*B^2*a^10*208i))*(-(d^3*(((594*A^3*a^
12 + 1458*A*B^2*a^12)*1i)/d^3 + (1458*B^3*a^12 + 486*A^2*B*a^12)/d^3)^2...

```

3.202
$$\int \frac{A+B \tan(c+dx)}{\sqrt[3]{a+ia \tan(c+dx)}} dx$$

3.202.1 Optimal result 2067
 3.202.2 Mathematica [A] (verified) 2068
 3.202.3 Rubi [A] (warning: unable to verify) 2068
 3.202.4 Maple [A] (verified) 2071
 3.202.5 Fricas [B] (verification not implemented) 2072
 3.202.6 Sympy [F] 2073
 3.202.7 Maxima [A] (verification not implemented) 2073
 3.202.8 Giac [F] 2073
 3.202.9 Mupad [B] (verification not implemented) 2074

3.202.1 Optimal result

Integrand size = 28, antiderivative size = 213

$$\int \frac{A+B \tan(c+dx)}{\sqrt[3]{a+ia \tan(c+dx)}} dx = -\frac{(A-iB)x}{4\sqrt[3]{2}\sqrt[3]{a}} + \frac{\sqrt{3}(iA+B) \arctan\left(\frac{\sqrt[3]{a+2^{2/3}}\sqrt[3]{a+ia \tan(c+dx)}}{\sqrt{3}\sqrt[3]{a}}\right)}{2\sqrt[3]{2}\sqrt[3]{ad}} + \frac{(iA+B) \log(\cos(c+dx))}{4\sqrt[3]{2}\sqrt[3]{ad}} + \frac{3(iA+B) \log\left(\sqrt[3]{2}\sqrt[3]{a}-\sqrt[3]{a+ia \tan(c+dx)}\right)}{4\sqrt[3]{2}\sqrt[3]{ad}} + \frac{3(iA-B)}{2d\sqrt[3]{a+ia \tan(c+dx)}}$$

output

```
-1/8*(A-I*B)*x*2^(2/3)/a^(1/3)+1/8*(I*A+B)*ln(cos(d*x+c))*2^(2/3)/a^(1/3)/
d+3/8*(I*A+B)*ln(2^(1/3)*a^(1/3)-(a+I*a*tan(d*x+c))^(1/3))*2^(2/3)/a^(1/3)
/d+1/4*(I*A+B)*arctan(1/3*(a^(1/3)+2^(2/3)*(a+I*a*tan(d*x+c))^(1/3))/a^(1/
3)*3^(1/2))*3^(1/2)*2^(2/3)/a^(1/3)/d+3/2*(I*A-B)/d/(a+I*a*tan(d*x+c))^(1/
3)
```

3.202.2 Mathematica [A] (verified)

Time = 0.89 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.70

$$\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + ia \tan(c + dx)}} dx$$

$$\frac{2^{2/3}(iA+B) \left(2\sqrt{3} \arctan\left(\frac{\sqrt[3]{a+2^{2/3}} \sqrt[3]{a + ia \tan(c + dx)}}{\sqrt{3} \sqrt[3]{a}}\right) - \log(i + \tan(c+dx)) + 3 \log\left(\sqrt[3]{2} \sqrt[3]{a} - \sqrt[3]{a + ia \tan(c + dx)}\right) \right)}{\sqrt[3]{a}}$$

$8d$

input `Integrate[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x])^(1/3), x]`

output `((2^(2/3)*(I*A + B)*(2*Sqrt[3]*ArcTan[(a^(1/3) + 2^(2/3)*(a + I*a*Tan[c + d*x])^(1/3)]/(Sqrt[3]*a^(1/3)))] - Log[I + Tan[c + d*x]] + 3*Log[2^(1/3)*a^(1/3) - (a + I*a*Tan[c + d*x])^(1/3]))/a^(1/3) + ((12*I)*(A + I*B))/(a + I*a*Tan[c + d*x])^(1/3))/(8*d)`

3.202.3 Rubi [A] (warning: unable to verify)

Time = 0.40 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.70, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4009, 3042, 3962, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + ia \tan(c + dx)}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + ia \tan(c + dx)}} dx$$

↓ 4009

$$\frac{(A - iB) \int (i \tan(c + dx) a + a)^{2/3} dx}{2a} + \frac{3(-B + iA)}{2d \sqrt[3]{a + ia \tan(c + dx)}}$$

↓ 3042

$$\frac{(A - iB) \int (i \tan(c + dx)a + a)^{2/3} dx}{2a} + \frac{3(-B + iA)}{2d \sqrt[3]{a + ia \tan(c + dx)}}$$

↓ 3962

$$\frac{3(-B + iA)}{2d \sqrt[3]{a + ia \tan(c + dx)}} - \frac{i(A - iB) \int \frac{1}{(a - ia \tan(c + dx)) \sqrt[3]{i \tan(c + dx)a + a}} d(ia \tan(c + dx))}{2d}$$

↓ 67

$$\frac{3(-B + iA)}{2d \sqrt[3]{a + ia \tan(c + dx)}} - \frac{i(A - iB) \left(-\frac{3}{2} \int \frac{1}{-a^2 \tan^2(c + dx) + i \sqrt[3]{2a^{4/3} \tan(c + dx) + 2^{2/3} a^{2/3}}} d \sqrt[3]{i \tan(c + dx)a + a} + \frac{3 \int \frac{1}{\sqrt[3]{2 \sqrt[3]{a} - ia \tan(c + dx)}} d \sqrt[3]{i \tan(c + dx)a + a}}{2 \sqrt[3]{2 \sqrt[3]{a}}} \right)}{2d}$$

↓ 16

$$\frac{3(-B + iA)}{2d \sqrt[3]{a + ia \tan(c + dx)}} - \frac{i(A - iB) \left(-\frac{3}{2} \int \frac{1}{-a^2 \tan^2(c + dx) + i \sqrt[3]{2a^{4/3} \tan(c + dx) + 2^{2/3} a^{2/3}}} d \sqrt[3]{i \tan(c + dx)a + a} - \frac{3 \log \left(\sqrt[3]{2 \sqrt[3]{a} - ia \tan(c + dx)} \right)}{2 \sqrt[3]{2 \sqrt[3]{a}}} + \frac{\log(a - ia \tan(c + dx))}{2 \sqrt[3]{2 \sqrt[3]{a}}} \right)}{2d}$$

↓ 1082

$$\frac{3(-B + iA)}{2d \sqrt[3]{a + ia \tan(c + dx)}} - \frac{i(A - iB) \left(\frac{3 \int \frac{1}{a^2 \tan^2(c + dx) - 3} d(i 2^{2/3} a^{2/3} \tan(c + dx) + 1)}{\sqrt[3]{2 \sqrt[3]{a}}} - \frac{3 \log \left(\sqrt[3]{2 \sqrt[3]{a} - ia \tan(c + dx)} \right)}{2 \sqrt[3]{2 \sqrt[3]{a}}} + \frac{\log(a - ia \tan(c + dx))}{2 \sqrt[3]{2 \sqrt[3]{a}}} \right)}{2d}$$

↓ 217

$$\frac{3(-B + iA)}{2d \sqrt[3]{a + ia \tan(c + dx)}} - \frac{i(A - iB) \left(-\frac{i \sqrt{3} \operatorname{arctanh} \left(\frac{a \tan(c + dx)}{\sqrt{3}} \right)}{\sqrt[3]{2 \sqrt[3]{a}}} - \frac{3 \log \left(\sqrt[3]{2 \sqrt[3]{a} - ia \tan(c + dx)} \right)}{2 \sqrt[3]{2 \sqrt[3]{a}}} + \frac{\log(a - ia \tan(c + dx))}{2 \sqrt[3]{2 \sqrt[3]{a}}} \right)}{2d}$$

input `Int[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x])^(1/3),x]`

3.202. $\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + ia \tan(c + dx)}} dx$

```
output ((-1/2*I)*(A - I*B)*((-I)*Sqrt[3]*ArcTanh[(a*Tan[c + d*x])/Sqrt[3]])/(2^(
1/3)*a^(1/3)) - (3*Log[2^(1/3)*a^(1/3) - I*a*Tan[c + d*x]]/(2*2^(1/3)*a^(
1/3)) + Log[a - I*a*Tan[c + d*x]]/(2*2^(1/3)*a^(1/3)))/d + (3*(I*A - B))/
(2*d*(a + I*a*Tan[c + d*x])^(1/3))
```

3.202.3.1 Defintions of rubi rules used

```
rule 16 Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a +
b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

```
rule 67 Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(1/3)), x_Symbol] := With[
{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q), x
] + (Simp[3/(2*b) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)],
x] - Simp[3/(2*b*q) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x])] /
; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3962 Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[-b/d S
ubst[Int[(a + x)^(n - 1)/(a - x), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b
, c, d, n}, x] && EqQ[a^2 + b^2, 0]
```

```
rule 4009 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]
```

3.202.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.78

method	result
derivativedivides	$3i \left(-\frac{-\frac{A}{2} - \frac{iB}{2}}{(a+ia \tan(dx+c))^{\frac{1}{3}}} + \frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{6a^{\frac{1}{3}}} - \frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}}(a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}}\right)}{12a^{\frac{1}{3}}}\right)$
default	$3i \left(-\frac{-\frac{A}{2} - \frac{iB}{2}}{(a+ia \tan(dx+c))^{\frac{1}{3}}} + \frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{6a^{\frac{1}{3}}} - \frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}}(a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}}\right)}{12a^{\frac{1}{3}}}\right)$
parts	$3iAa \left(\frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{6a^{\frac{1}{3}}} - \frac{2^{\frac{2}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}}(a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}}\right)}{12a^{\frac{1}{3}}} + \frac{\sqrt{3} 2^{\frac{2}{3}} \arctan\left(\frac{\sqrt{3}}{\dots}\right)}{2a}\right)$

```
input int((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/3),x,method=_RETURNVERBOSE)
```

3.202. $\int \frac{A+B \tan(c+dx)}{\sqrt[3]{a+ia \tan(c+dx)}} dx$

output $3*I/d*(-(-1/2*A-1/2*I*B)/(a+I*a*\tan(d*x+c))^{(1/3)}+(1/6*2^{(2/3)}/a^{(1/3)}*\ln((a+I*a*\tan(d*x+c))^{(1/3)}-2^{(1/3)}*a^{(1/3)})-1/12*2^{(2/3)}/a^{(1/3)}*\ln((a+I*a*\tan(d*x+c))^{(2/3)}+2^{(1/3)}*a^{(1/3)}*(a+I*a*\tan(d*x+c))^{(1/3)}+2^{(2/3)}*a^{(2/3)}))+1/6*3^{(1/2)}*2^{(2/3)}/a^{(1/3)}*\arctan(1/3*3^{(1/2)}*(2^{(2/3)}/a^{(1/3)}*(a+I*a*\tan(d*x+c))^{(1/3)+1}))*((1/2*A-1/2*I*B))$

3.202.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 547 vs. $2(156) = 312$.

Time = 0.26 (sec) , antiderivative size = 547, normalized size of antiderivative = 2.57

$$\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + ia \tan(c + dx)}} dx$$

$$= \left(2 \left(\frac{1}{2} \right)^{\frac{1}{3}} ad \left(\frac{-iA^3 - 3A^2B + 3iAB^2 + B^3}{ad^3} \right)^{\frac{1}{3}} e^{(2i dx + 2i c)} \log \left(\frac{2 \left(\frac{1}{2} \right)^{\frac{2}{3}} ad^2 \left(\frac{-iA^3 - 3A^2B + 3iAB^2 + B^3}{ad^3} \right)^{\frac{2}{3}} + 2^{\frac{1}{3}} (A^2 - 2iAB - B^2)}{A^2 - 2iAB - B^2} \right) \left(\frac{e^{(2i dx + 2i c)}}{e^{(2i dx + 2i c)}} \right) \right)$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/3),x, algorithm="fracas")`

output $1/4*(2*(1/2)^{(1/3)}*a*d*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)/(a*d^3))^{(1/3)})*e^{(2*I*d*x + 2*I*c)}*\log(((2*(1/2)^{(2/3)}*a*d^2*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)/(a*d^3))^{(2/3)} + 2^{(1/3)}*(A^2 - 2*I*A*B - B^2)*(a/(e^{(2*I*d*x + 2*I*c)} + 1))^{(1/3)})*e^{(2/3*I*d*x + 2/3*I*c)})/(A^2 - 2*I*A*B - B^2)) - (1/2)^{(1/3)}*(I*sqrt(3)*a*d + a*d)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)/(a*d^3))^{(1/3)}*e^{(2*I*d*x + 2*I*c)}*\log(((2^{(1/3)}*(A^2 - 2*I*A*B - B^2)*(a/(e^{(2*I*d*x + 2*I*c)} + 1))^{(1/3)})*e^{(2/3*I*d*x + 2/3*I*c)} + (1/2)^{(2/3)}*(I*sqrt(3)*a*d^2 - a*d^2)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)/(a*d^3))^{(2/3)})/(A^2 - 2*I*A*B - B^2)) - (1/2)^{(1/3)}*(-I*sqrt(3)*a*d + a*d)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)/(a*d^3))^{(1/3)}*e^{(2*I*d*x + 2*I*c)}*\log(((2^{(1/3)}*(A^2 - 2*I*A*B - B^2)*(a/(e^{(2*I*d*x + 2*I*c)} + 1))^{(1/3)})*e^{(2/3*I*d*x + 2/3*I*c)} + (1/2)^{(2/3)}*(-I*sqrt(3)*a*d^2 - a*d^2)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)/(a*d^3))^{(2/3)})/(A^2 - 2*I*A*B - B^2)) - 3*2^{(2/3)}*((-I*A + B)*e^{(2*I*d*x + 2*I*c)} - I*A + B)*(a/(e^{(2*I*d*x + 2*I*c)} + 1))^{(2/3)}*e^{(4/3*I*d*x + 4/3*I*c)})*e^{(-2*I*d*x - 2*I*c)}/(a*d)$

3.202.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + ia \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt[3]{ia (\tan(c + dx) - i)}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/3),x)`

output `Integral((A + B*tan(c + d*x))/(I*a*(tan(c + d*x) - I))**(1/3), x)`

3.202.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.81

$$\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + ia \tan(c + dx)}} dx$$

$$= \frac{i \left(2 \sqrt{3} 2^{\frac{2}{3}} (A - i B) a^{\frac{2}{3}} \arctan \left(\frac{\sqrt{3} 2^{\frac{2}{3}} \left(2^{\frac{1}{3}} a^{\frac{1}{3}} + 2 (i a \tan(dx+c) + a)^{\frac{1}{3}} \right)}{6 a^{\frac{1}{3}}} \right) - 2^{\frac{2}{3}} (A - i B) a^{\frac{2}{3}} \log \left(2^{\frac{2}{3}} a^{\frac{2}{3}} + 2^{\frac{1}{3}} (i a \tan(dx+c) + a)^{\frac{1}{3}} \right) \right)}{6 a^{\frac{1}{3}}}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/3),x, algorithm="maxima")`

output `1/8*I*(2*sqrt(3)*2^(2/3)*(A - I*B)*a^(2/3)*arctan(1/6*sqrt(3)*2^(2/3)*(2^(1/3)*a^(1/3) + 2*(I*a*tan(d*x + c) + a)^(1/3))/a^(1/3)) - 2^(2/3)*(A - I*B)*a^(2/3)*log(2^(2/3)*a^(2/3) + 2^(1/3)*(I*a*tan(d*x + c) + a)^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(2/3)) + 2*2^(2/3)*(A - I*B)*a^(2/3)*log(-2^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(1/3)) + 12*(A + I*B)*a/(I*a*tan(d*x + c) + a)^(1/3))/(a*d)`

3.202.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + ia \tan(c + dx)}} dx = \int \frac{B \tan(dx + c) + A}{(i a \tan(dx + c) + a)^{\frac{1}{3}}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/3),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/(I*a*tan(d*x + c) + a)^(1/3), x)`

3.202. $\int \frac{A+B \tan(c+dx)}{\sqrt[3]{a + ia \tan(c + dx)}} dx$

3.202.9 Mupad [B] (verification not implemented)

Time = 8.52 (sec) , antiderivative size = 383, normalized size of antiderivative = 1.80

$$\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + ia \tan(c + dx)}} dx = \frac{A 3i}{2 d (a + a \tan(c + dx) i)^{1/3}} - \frac{3 B}{2 d (a + a \tan(c + dx) i)^{1/3}}$$

$$- \frac{\left(\frac{1}{16}i\right)^{1/3} A \ln\left((a(1 + \tan(c + dx) i))^{1/3} + (-1)^{1/3} 2^{1/3} a^{1/3}\right)}{a^{1/3} d}$$

$$+ \frac{4^{1/3} B \ln\left(18 B^2 d (a + a \tan(c + dx) i)^{1/3} - 9 4^{2/3} B^2 a^{1/3} d\right)}{4 a^{1/3} d}$$

$$+ \frac{4^{1/3} B \ln\left(18 B^2 d (a + a \tan(c + dx) i)^{1/3} - 9 4^{2/3} B^2 a^{1/3} d \left(-\frac{1}{2} + \frac{\sqrt{3} i}{2}\right)^2\right) \left(-\frac{1}{2} + \frac{\sqrt{3} i}{2}\right)}{4 a^{1/3} d}$$

$$- \frac{4^{1/3} B \ln\left(18 B^2 d (a + a \tan(c + dx) i)^{1/3} - 9 4^{2/3} B^2 a^{1/3} d \left(\frac{1}{2} + \frac{\sqrt{3} i}{2}\right)^2\right) \left(\frac{1}{2} + \frac{\sqrt{3} i}{2}\right)}{4 a^{1/3} d}$$

$$- \frac{\left(\frac{1}{16}i\right)^{1/3} A \ln\left(\frac{(-1)^{1/3} 2^{1/3} a^{1/3}}{2} - (a(1 + \tan(c + dx) i))^{1/3} + \frac{(-1)^{5/6} 2^{1/3} \sqrt{3} a^{1/3}}{2}\right) \left(-\frac{1}{2} + \frac{\sqrt{3} i}{2}\right)}{a^{1/3} d}$$

$$+ \frac{\left(\frac{1}{16}i\right)^{1/3} A \ln\left((a(1 + \tan(c + dx) i))^{1/3} - \frac{(-1)^{1/3} 2^{1/3} a^{1/3}}{2} + \frac{(-1)^{5/6} 2^{1/3} \sqrt{3} a^{1/3}}{2}\right) \left(\frac{1}{2} + \frac{\sqrt{3} i}{2}\right)}{a^{1/3} d}$$

input `int((A + B*tan(c + d*x))/(a + a*tan(c + d*x)*i)^(1/3),x)`

output `(A*3i)/(2*d*(a + a*tan(c + d*x)*i)^(1/3)) - (3*B)/(2*d*(a + a*tan(c + d*x)*i)^(1/3)) - ((i/16)^(1/3)*A*log((a*(tan(c + d*x)*i + 1))^(1/3) + (-1)^(1/3)*2^(1/3)*a^(1/3)))/(a^(1/3)*d) + (4^(1/3)*B*log(18*B^2*d*(a + a*tan(c + d*x)*i)^(1/3) - 9*4^(2/3)*B^2*a^(1/3)*d))/(4*a^(1/3)*d) + (4^(1/3)*B*log(18*B^2*d*(a + a*tan(c + d*x)*i)^(1/3) - 9*4^(2/3)*B^2*a^(1/3)*d*((3^(1/2)*i)/2 - 1/2)^2)*((3^(1/2)*i)/2 - 1/2))/(4*a^(1/3)*d) - (4^(1/3)*B*log(18*B^2*d*(a + a*tan(c + d*x)*i)^(1/3) - 9*4^(2/3)*B^2*a^(1/3)*d*((3^(1/2)*i)/2 + 1/2)^2)*((3^(1/2)*i)/2 + 1/2))/(4*a^(1/3)*d) - ((i/16)^(1/3)*A*log(((1)^(1/3)*2^(1/3)*a^(1/3))/2 - (a*(tan(c + d*x)*i + 1))^(1/3) + ((-1)^(5/6)*2^(1/3)*3^(1/2)*a^(1/3))/2)*((3^(1/2)*i)/2 - 1/2))/(a^(1/3)*d) + ((i/16)^(1/3)*A*log((a*(tan(c + d*x)*i + 1))^(1/3) - ((-1)^(1/3)*2^(1/3)*a^(1/3))/2 + ((-1)^(5/6)*2^(1/3)*3^(1/2)*a^(1/3))/2)*((3^(1/2)*i)/2 + 1/2))/(a^(1/3)*d)`

3.203 $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^{2/3}} dx$

3.203.1 Optimal result 2075
 3.203.2 Mathematica [A] (verified) 2076
 3.203.3 Rubi [A] (warning: unable to verify) 2076
 3.203.4 Maple [A] (verified) 2079
 3.203.5 Fricas [B] (verification not implemented) 2080
 3.203.6 Sympy [F] 2081
 3.203.7 Maxima [A] (verification not implemented) 2081
 3.203.8 Giac [F] 2081
 3.203.9 Mupad [B] (verification not implemented) 2082

3.203.1 Optimal result

Integrand size = 28, antiderivative size = 213

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{2/3}} dx = -\frac{(A - iB)x}{4 \cdot 2^{2/3} a^{2/3}} - \frac{\sqrt{3}(iA + B) \arctan\left(\frac{\sqrt[3]{a+2^{2/3}} \sqrt[3]{a + ia \tan(c + dx)}}{\sqrt{3} \sqrt[3]{a}}\right)}{2 \cdot 2^{2/3} a^{2/3} d} + \frac{(iA + B) \log(\cos(c + dx))}{4 \cdot 2^{2/3} a^{2/3} d} + \frac{3(iA + B) \log\left(\sqrt[3]{2} \sqrt[3]{a} - \sqrt[3]{a + ia \tan(c + dx)}\right)}{4 \cdot 2^{2/3} a^{2/3} d} + \frac{3(iA - B)}{4d(a + ia \tan(c + dx))^{2/3}}$$

output

```
-1/8*(A-I*B)*x*2^(1/3)/a^(2/3)+1/8*(I*A+B)*ln(cos(d*x+c))*2^(1/3)/a^(2/3)/d+3/8*(I*A+B)*ln(2^(1/3)*a^(1/3)-(a+I*a*tan(d*x+c))^(1/3))*2^(1/3)/a^(2/3)/d-1/4*(I*A+B)*arctan(1/3*(a^(1/3)+2^(2/3)*(a+I*a*tan(d*x+c))^(1/3))/a^(1/3)*3^(1/2))*3^(1/2)*2^(1/3)/a^(2/3)/d+3/4*(I*A-B)/d/(a+I*a*tan(d*x+c))^(2/3)
```

3.203.2 Mathematica [A] (verified)

Time = 0.79 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.93

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{2/3}} dx = \frac{i \sqrt[3]{2}^{A-iB} \left(2\sqrt{3} \arctan \left(\frac{\sqrt[3]{a+2^{2/3}} \sqrt[3]{a + ia \tan(c + dx)}}{\sqrt{3} \sqrt[3]{a}} \right) \right) - 2 \log \left(\sqrt[3]{2} \sqrt[3]{a} - \sqrt[3]{a + ia \tan(c + dx)} \right)}{a^{2/3}}$$

input `Integrate[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x])^(2/3), x]`

output `(((-I)*2^(1/3)*(A - I*B)*(2*Sqrt[3]*ArcTan[(a^(1/3) + 2^(2/3)*(a + I*a*Tan[c + d*x])^(1/3)]/(Sqrt[3]*a^(1/3))] - 2*Log[2^(1/3)*a^(1/3) - (a + I*a*Tan[c + d*x])^(1/3)] + Log[2^(2/3)*a^(2/3) + 2^(1/3)*a^(1/3)*(a + I*a*Tan[c + d*x])^(1/3) + (a + I*a*Tan[c + d*x])^(2/3)]))/a^(2/3) + ((6*I)*(A + I*B))/(a + I*a*Tan[c + d*x])^(2/3))/(8*d)`

3.203.3 Rubi [A] (warning: unable to verify)

Time = 0.42 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.70, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4009, 3042, 3962, 69, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{2/3}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{2/3}} dx \\ & \quad \downarrow \text{4009} \\ & \frac{(A - iB) \int \sqrt[3]{i \tan(c + dx)a + adx}}{2a} + \frac{3(-B + iA)}{4d(a + ia \tan(c + dx))^{2/3}} \\ & \quad \downarrow \text{3042} \\ & \frac{(A - iB) \int \sqrt[3]{i \tan(c + dx)a + adx}}{2a} + \frac{3(-B + iA)}{4d(a + ia \tan(c + dx))^{2/3}} \end{aligned}$$

3.203. $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^{2/3}} dx$

$$\begin{aligned}
& \frac{3(-B + iA)}{4d(a + ia \tan(c + dx))^{2/3}} - \frac{i(A - iB) \int \frac{1}{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^{2/3}} d(ia \tan(c + dx))}{2d} \\
& \quad \downarrow \text{3962} \\
& \frac{3(-B + iA)}{4d(a + ia \tan(c + dx))^{2/3}} - \\
& \quad \downarrow \text{69} \\
& \frac{i(A - iB) \left(\frac{3 \int \frac{1}{\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)}} d^3 \sqrt{i \tan(c + dx)a + a}}{2 \cdot 2^{2/3} a^{2/3}} + \frac{3 \int \frac{1}{-a^2 \tan^2(c + dx) + i \sqrt[3]{2} a^{4/3} \tan(c + dx) + 2^{2/3} a^{2/3}} d^3 \sqrt{i \tan(c + dx)}}{2 \sqrt[3]{2} \sqrt[3]{a}} \right)}{2d} \\
& \quad \downarrow \text{16} \\
& \frac{3(-B + iA)}{4d(a + ia \tan(c + dx))^{2/3}} - \\
& \quad \downarrow \text{1082} \\
& \frac{i(A - iB) \left(\frac{3 \int \frac{1}{-a^2 \tan^2(c + dx) + i \sqrt[3]{2} a^{4/3} \tan(c + dx) + 2^{2/3} a^{2/3}} d^3 \sqrt{i \tan(c + dx)a + a}}{2 \sqrt[3]{2} \sqrt[3]{a}} - \frac{3 \log(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)})}{2 \cdot 2^{2/3} a^{2/3}} + \frac{\log(a - ia \tan(c + dx))}{2 \cdot 2^{2/3} a^{2/3}} \right)}{2d} \\
& \quad \downarrow \text{217} \\
& \frac{3(-B + iA)}{4d(a + ia \tan(c + dx))^{2/3}} - \\
& \quad \downarrow \\
& \frac{i(A - iB) \left(\frac{i\sqrt{3} \arctan\left(\frac{a \tan(c + dx)}{\sqrt{3}}\right)}{2^{2/3} a^{2/3}} - \frac{3 \log(\sqrt[3]{2} \sqrt[3]{a - ia \tan(c + dx)})}{2 \cdot 2^{2/3} a^{2/3}} + \frac{\log(a - ia \tan(c + dx))}{2 \cdot 2^{2/3} a^{2/3}} \right)}{2d}
\end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(a + I*a*Tan[c + d*x])^(2/3),x]`

output $((-1/2*I)*(A - I*B)*((I*sqrt[3]*ArcTanh[(a*Tan[c + d*x])/sqrt[3]])/(2^(2/3)*a^(2/3)) - (3*Log[2^(1/3)*a^(1/3) - I*a*Tan[c + d*x]])/(2*2^(2/3)*a^(2/3))) + Log[a - I*a*Tan[c + d*x]]/(2*2^(2/3)*a^(2/3)))/d + (3*(I*A - B))/(4*d*(a + I*a*Tan[c + d*x])^(2/3))$

3.203.3.1 Defintions of rubi rules used

rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 69 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))^(2/3)), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[(b*c - a*d)/b, 3]\}, \text{Simp}[-\text{Log}[\text{RemoveContent}[a + b*x, x]]/(2*b*q^2), x] + (-\text{Simp}[3/(2*b*q) \text{Subst}[\text{Int}[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - \text{Simp}[3/(2*b*q^2) \text{Subst}[\text{Int}[1/(q - x), x], x, (c + d*x)^(1/3)], x])] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{PosQ}[(b*c - a*d)/b]$

rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \& \& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])$

rule 1082 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \parallel \text{!RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3962 $\text{Int}[(a_)+(b_)*\tan[(c_)+(d_)*(x_)])^{(n_)}, x_Symbol] \rightarrow \text{Simp}[-b/d \text{Subst}[\text{Int}[(a + x)^{(n-1)}/(a - x), x], x, b*\text{Tan}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \&\& \text{EqQ}[a^2 + b^2, 0]$

```
rule 4009 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]
```

3.203.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.78

method	result
derivativedivides	$3i \left(-\frac{-\frac{A}{2} - \frac{iB}{2}}{2(a+ia \tan(dx+c))^{\frac{2}{3}}} + \frac{2^{\frac{1}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{6a^{\frac{2}{3}}} - \frac{2^{\frac{1}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}}(a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}}\right)}{12a^{\frac{2}{3}}} \right) d$
default	$3i \left(-\frac{-\frac{A}{2} - \frac{iB}{2}}{2(a+ia \tan(dx+c))^{\frac{2}{3}}} + \frac{2^{\frac{1}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{6a^{\frac{2}{3}}} - \frac{2^{\frac{1}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}}(a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}}\right)}{12a^{\frac{2}{3}}} \right) d$
parts	$3iAa \left(\frac{2^{\frac{1}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{1}{3}} - 2^{\frac{1}{3}} a^{\frac{1}{3}}\right)}{6a^{\frac{2}{3}}} - \frac{2^{\frac{1}{3}} \ln\left((a+ia \tan(dx+c))^{\frac{2}{3}} + 2^{\frac{1}{3}} a^{\frac{1}{3}}(a+ia \tan(dx+c))^{\frac{1}{3}} + 2^{\frac{2}{3}} a^{\frac{2}{3}}\right)}{12a^{\frac{2}{3}}} - \frac{2^{\frac{1}{3}} \sqrt{3} \arctan\left(\frac{\sqrt{3}}{\dots}\right)}{2a} \right) d$

```
input int((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(2/3),x,method=_RETURNVERBOSE)
```

3.203. $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^{2/3}} dx$

output $3*I/d*(-1/2*(-1/2*A-1/2*I*B)/(a+I*a*\tan(d*x+c))^(2/3)+(1/6*2^(1/3)/a^(2/3)*\ln((a+I*a*\tan(d*x+c))^(1/3)-2^(1/3)*a^(1/3))-1/12*2^(1/3)/a^(2/3)*\ln((a+I*a*\tan(d*x+c))^(2/3)+2^(1/3)*a^(1/3)*(a+I*a*\tan(d*x+c))^(1/3)+2^(2/3)*a^(2/3))-1/6*2^(1/3)/a^(2/3)*3^(1/2)*\arctan(1/3*3^(1/2)*(2^(2/3)/a^(1/3)*(a+I*a*\tan(d*x+c))^(1/3)+1)))*(1/2*A-1/2*I*B)$

3.203.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 493 vs. $2(156) = 312$.

Time = 0.25 (sec) , antiderivative size = 493, normalized size of antiderivative = 2.31

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{2/3}} dx = \frac{\left(4 \left(\frac{1}{4} \right)^{\frac{1}{3}} ad \left(\frac{-iA^3 - 3A^2B + 3iAB^2 + B^3}{a^2d^3} \right)^{\frac{1}{3}} e^{(2i dx + 2i c)} \log \left(- \frac{2 \left(\frac{1}{4} \right)^{\frac{1}{3}} ad \left(\frac{-iA^3 - 3A^2B + 3iAB^2 + B^3}{a^2d^3} \right)^{\frac{1}{3}}}{\dots} \right)}{\dots}$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(2/3),x, algorithm="fracas")`

output $1/8*(4*(1/4)^(1/3)*a*d*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)/(a^2*d^3))^(1/3)*e^(2*I*d*x + 2*I*c)*\log(-(2*(1/4)^(1/3)*a*d*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)/(a^2*d^3))^(1/3) - 2^(1/3)*(I*A + B)*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c))/(I*A + B)) - 2*(1/4)^(1/3)*(-I*sqrt(3)*a*d + a*d)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)/(a^2*d^3))^(1/3)*e^(2*I*d*x + 2*I*c)*\log((2^(1/3)*(I*A + B)*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) - (1/4)^(1/3)*(I*sqrt(3)*a*d - a*d)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)/(a^2*d^3))^(1/3))/(I*A + B)) - 2*(1/4)^(1/3)*(I*sqrt(3)*a*d + a*d)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)/(a^2*d^3))^(1/3)*e^(2*I*d*x + 2*I*c)*\log((2^(1/3)*(I*A + B)*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c) - (1/4)^(1/3)*(-I*sqrt(3)*a*d - a*d)*((-I*A^3 - 3*A^2*B + 3*I*A*B^2 + B^3)/(a^2*d^3))^(1/3))/(I*A + B)) - 3*2^(1/3)*((-I*A + B)*e^(2*I*d*x + 2*I*c) - I*A + B)*(a/(e^(2*I*d*x + 2*I*c) + 1))^(1/3)*e^(2/3*I*d*x + 2/3*I*c))*e^(-2*I*d*x - 2*I*c)/(a*d)$

3.203.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{2/3}} dx = \int \frac{A + B \tan(c + dx)}{(ia (\tan(c + dx) - i))^{2/3}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(2/3),x)`

output `Integral((A + B*tan(c + d*x))/(I*a*(tan(c + d*x) - I))**(2/3), x)`

3.203.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.80

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{2/3}} dx =$$

$$i \left(2 \sqrt{3} 2^{1/3} (A - i B) a^{1/3} \arctan \left(\frac{\sqrt{3} 2^{2/3} (2^{1/3} a^{1/3} + 2 (i a \tan(dx+c) + a)^{1/3})}{6 a^{1/3}} \right) + 2^{1/3} (A - i B) a^{1/3} \log \left(2^{2/3} a^{2/3} + 2^{1/3} (i a \tan(dx+c) + a)^{1/3} \right) \right)$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(2/3),x, algorithm="maxima")`

output `-1/8*I*(2*sqrt(3)*2^(1/3)*(A - I*B)*a^(1/3)*arctan(1/6*sqrt(3)*2^(2/3)*(2^(1/3)*a^(1/3) + 2*(I*a*tan(d*x + c) + a)^(1/3))/a^(1/3)) + 2^(1/3)*(A - I*B)*a^(1/3)*log(2^(2/3)*a^(2/3) + 2^(1/3)*(I*a*tan(d*x + c) + a)^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(2/3)) - 2*2^(1/3)*(A - I*B)*a^(1/3)*log(-2^(1/3)*a^(1/3) + (I*a*tan(d*x + c) + a)^(1/3)) - 6*(A + I*B)*a/(I*a*tan(d*x + c) + a)^(2/3))/(a*d)`

3.203.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{2/3}} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^{2/3}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(2/3),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/(I*a*tan(d*x + c) + a)^(2/3), x)`

3.203. $\int \frac{A+B \tan(c+dx)}{(a+ia \tan(c+dx))^{2/3}} dx$

3.203.9 Mupad [B] (verification not implemented)

Time = 8.42 (sec) , antiderivative size = 390, normalized size of antiderivative = 1.83

$$\int \frac{A + B \tan(c + dx)}{(a + ia \tan(c + dx))^{2/3}} dx = \frac{A 3i}{4d(a + a \tan(c + dx) li)^{2/3}} - \frac{3B}{4d(a + a \tan(c + dx) li)^{2/3}}$$

$$- \frac{\left(\frac{1}{32}i\right)^{1/3} A \ln\left(A d^2 (a + a \tan(c + dx) li)^{1/3} 36i + 144 \left(\frac{1}{32}i\right)^{1/3} A a^{1/3} d^2\right)}{a^{2/3} d}$$

$$+ \frac{2^{1/3} B \ln\left(36 B d^2 (a + a \tan(c + dx) li)^{1/3} - 36 2^{1/3} B a^{1/3} d^2\right)}{4 a^{2/3} d}$$

$$- \frac{\left(\frac{1}{32}i\right)^{1/3} A \ln\left(A d^2 (a + a \tan(c + dx) li)^{1/3} 36i + 144 \left(\frac{1}{32}i\right)^{1/3} A a^{1/3} d^2 \left(-\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)\right) \left(-\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)}{a^{2/3} d}$$

$$+ \frac{\left(\frac{1}{32}i\right)^{1/3} A \ln\left(A d^2 (a + a \tan(c + dx) li)^{1/3} 36i - 144 \left(\frac{1}{32}i\right)^{1/3} A a^{1/3} d^2 \left(\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)\right) \left(\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)}{a^{2/3} d}$$

$$+ \frac{2^{1/3} B \ln\left(36 B d^2 (a + a \tan(c + dx) li)^{1/3} - 36 2^{1/3} B a^{1/3} d^2 \left(-\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)\right) \left(-\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)}{4 a^{2/3} d}$$

$$- \frac{2^{1/3} B \ln\left(36 B d^2 (a + a \tan(c + dx) li)^{1/3} + 36 2^{1/3} B a^{1/3} d^2 \left(\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)\right) \left(\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)}{4 a^{2/3} d}$$

input `int((A + B*tan(c + d*x))/(a + a*tan(c + d*x)*1i)^(2/3),x)`

```
output (A*3i)/(4*d*(a + a*tan(c + d*x)*1i)^(2/3)) - (3*B)/(4*d*(a + a*tan(c + d*x)
)*1i)^(2/3)) - ((1i/32)^(1/3)*A*log(A*d^2*(a + a*tan(c + d*x)*1i)^(1/3)*36
i + 144*(1i/32)^(1/3)*A*a^(1/3)*d^2))/(a^(2/3)*d) + (2^(1/3)*B*log(36*B*d^
2*(a + a*tan(c + d*x)*1i)^(1/3) - 36*2^(1/3)*B*a^(1/3)*d^2))/(4*a^(2/3)*d)
- ((1i/32)^(1/3)*A*log(A*d^2*(a + a*tan(c + d*x)*1i)^(1/3)*36i + 144*(1i/
32)^(1/3)*A*a^(1/3)*d^2*((3^(1/2)*1i)/2 - 1/2))*((3^(1/2)*1i)/2 - 1/2))/(a
^(2/3)*d) + ((1i/32)^(1/3)*A*log(A*d^2*(a + a*tan(c + d*x)*1i)^(1/3)*36i -
144*(1i/32)^(1/3)*A*a^(1/3)*d^2*((3^(1/2)*1i)/2 + 1/2))*((3^(1/2)*1i)/2 +
1/2))/(a^(2/3)*d) + (2^(1/3)*B*log(36*B*d^2*(a + a*tan(c + d*x)*1i)^(1/3)
- 36*2^(1/3)*B*a^(1/3)*d^2*((3^(1/2)*1i)/2 - 1/2))*((3^(1/2)*1i)/2 - 1/2)
)/(4*a^(2/3)*d) - (2^(1/3)*B*log(36*B*d^2*(a + a*tan(c + d*x)*1i)^(1/3) +
36*2^(1/3)*B*a^(1/3)*d^2*((3^(1/2)*1i)/2 + 1/2))*((3^(1/2)*1i)/2 + 1/2))/(
4*a^(2/3)*d)
```

3.204 $\int \tan^m(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.204.1 Optimal result	2083
3.204.2 Mathematica [A] (verified)	2084
3.204.3 Rubi [A] (verified)	2084
3.204.4 Maple [F]	2088
3.204.5 Fricas [F]	2088
3.204.6 Sympy [F]	2089
3.204.7 Maxima [F]	2090
3.204.8 Giac [F]	2090
3.204.9 Mupad [F(-1)]	2090

3.204.1 Optimal result

Integrand size = 34, antiderivative size = 290

$$\int \tan^m(c+dx)(a+ia \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= -\frac{2a^4(A(64+60m+19m^2+2m^3)-iB(67+60m+19m^2+2m^3)) \tan^{1+m}(c+dx)}{d(1+m)(2+m)(3+m)(4+m)}$$

$$+ \frac{8a^4(A-iB) \operatorname{Hypergeometric2F1}(1, 1+m, 2+m, i \tan(c+dx)) \tan^{1+m}(c+dx)}{d(1+m)}$$

$$+ \frac{iaB \tan^{1+m}(c+dx)(a+ia \tan(c+dx))^3}{d(4+m)}$$

$$- \frac{(A(4+m)-iB(7+m)) \tan^{1+m}(c+dx) (a^2+ia^2 \tan(c+dx))^2}{d(3+m)(4+m)}$$

$$- \frac{2(A(4+m)^2-iB(19+8m+m^2)) \tan^{1+m}(c+dx) (a^4+ia^4 \tan(c+dx))}{d(2+m)(3+m)(4+m)}$$

```
output -2*a^4*(A*(2*m^3+19*m^2+60*m+64)-I*B*(2*m^3+19*m^2+60*m+67))*tan(d*x+c)^(1+m)/d/(3+m)/(4+m)/(m^2+3*m+2)+8*a^4*(A-I*B)*hypergeom([1, 1+m], [2+m], I*tan(d*x+c))*tan(d*x+c)^(1+m)/d/(1+m)+I*a*B*tan(d*x+c)^(1+m)*(a+I*a*tan(d*x+c))^3/d/(4+m)-(A*(4+m)-I*B*(7+m))*tan(d*x+c)^(1+m)*(a^2+I*a^2*tan(d*x+c))^2/d/(3+m)/(4+m)-2*(A*(4+m)^2-I*B*(m^2+8*m+19))*tan(d*x+c)^(1+m)*(a^4+I*a^4*tan(d*x+c))/d/(4+m)/(m^2+5*m+6)
```

3.204.2 Mathematica [A] (verified)

Time = 3.10 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.60

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{a^4 \tan^{1+m}(c + dx) \left(\frac{(A - iB)(-7(6 + 5m + m^2) + 8(6 + 5m + m^2)) \operatorname{Hypergeometric2F1}(1, 1 + m, 2 + m, i \tan(c + dx)) - 4i(3 + 4m + m^2) \tan(c + dx)}{(1 + m)(2 + m)(3 + m)} \right)}{d}$$

input `Integrate[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(a^4*Tan[c + d*x]^(1 + m)*(((A - I*B)*(-7*(6 + 5*m + m^2) + 8*(6 + 5*m + m^2))*Hypergeometric2F1[1, 1 + m, 2 + m, I*Tan[c + d*x]] - (4*I)*(3 + 4*m + m^2)*Tan[c + d*x] + (2 + 3*m + m^2)*Tan[c + d*x]^2))/((1 + m)*(2 + m)*(3 + m)) + I*B*((1 + m)^(-1) + ((3*I)*Tan[c + d*x])/(2 + m) - (3*Tan[c + d*x]^2)/(3 + m) - (I*Tan[c + d*x]^3)/(4 + m)))/d`

3.204.3 Rubi [A] (verified)

Time = 1.82 (sec) , antiderivative size = 315, normalized size of antiderivative = 1.09, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {3042, 4077, 25, 3042, 4077, 27, 3042, 4077, 3042, 4075, 3042, 4020, 27, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(c + dx))^4 \tan^m(c + dx)(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int (a + ia \tan(c + dx))^4 \tan(c + dx)^m (A + B \tan(c + dx)) dx$$

$$\downarrow \text{4077}$$

$$\frac{\int -\tan^m(c + dx)(i \tan(c + dx)a + a)^3(a(iB(m + 1) - A(m + 4)) - a(iA(m + 4) + B(m + 7)) \tan(c + dx)) dx}{\frac{iaB(a + ia \tan(c + dx))^3 \tan^{m+1}(c + dx)}{d(m + 4)}} +$$

$$\downarrow \text{25}$$

$$\frac{\frac{iaB(a + ia \tan(c + dx))^3 \tan^{m+1}(c + dx)}{d(m + 4)} - \int \tan^m(c + dx)(i \tan(c + dx)a + a)^3(a(iB(m + 1) - A(m + 4)) - a(iA(m + 4) + B(m + 7)) \tan(c + dx))dx}{m + 4}}$$

↓ 3042

$$\frac{\frac{iaB(a + ia \tan(c + dx))^3 \tan^{m+1}(c + dx)}{d(m + 4)} - \int \tan(c + dx)^m(i \tan(c + dx)a + a)^3(a(iB(m + 1) - A(m + 4)) - a(iA(m + 4) + B(m + 7)) \tan(c + dx))dx}{m + 4}}$$

↓ 4077

$$\frac{\frac{iaB(a + ia \tan(c + dx))^3 \tan^{m+1}(c + dx)}{d(m + 4)} - \int \frac{2 \tan^m(c + dx)(i \tan(c + dx)a + a)^2(a^2(iB(m^2 + 6m + 5) - A(m^2 + 6m + 8)) - a^2(iA(m + 4)^2 + B(m^2 + 8m + 19)) \tan(c + dx))dx}{m + 3} + \frac{(A(m + 4) - iB(m + 4)) \tan(c + dx)}{m + 4}}{m + 4}}$$

↓ 27

$$\frac{\frac{iaB(a + ia \tan(c + dx))^3 \tan^{m+1}(c + dx)}{d(m + 4)} - 2 \int \frac{\tan^m(c + dx)(i \tan(c + dx)a + a)^2(a^2(iB(m^2 + 6m + 5) - A(m^2 + 6m + 8)) - a^2(iA(m + 4)^2 + B(m^2 + 8m + 19)) \tan(c + dx))dx}{m + 3} + \frac{(A(m + 4) - iB(m + 4)) \tan(c + dx)}{m + 4}}{m + 4}}$$

↓ 3042

$$\frac{\frac{iaB(a + ia \tan(c + dx))^3 \tan^{m+1}(c + dx)}{d(m + 4)} - 2 \int \frac{\tan(c + dx)^m(i \tan(c + dx)a + a)^2(a^2(iB(m^2 + 6m + 5) - A(m^2 + 6m + 8)) - a^2(iA(m + 4)^2 + B(m^2 + 8m + 19)) \tan(c + dx))dx}{m + 3} + \frac{(A(m + 4) - iB(m + 4)) \tan(c + dx)}{m + 4}}{m + 4}}$$

↓ 4077

$$\frac{\frac{iaB(a + ia \tan(c + dx))^3 \tan^{m+1}(c + dx)}{d(m + 4)} - 2 \left(\frac{\int \tan^m(c + dx)(i \tan(c + dx)a + a)(a^3(iB(2m^3 + 17m^2 + 44m + 29) - A(2m^3 + 17m^2 + 44m + 32)) - a^3(iA(2m^3 + 19m^2 + 60m + 64) + B(2m^3 + 19m^2 + 60m + 67)) \tan(c + dx))dx}{m + 2} + \frac{(A(m + 4) - iB(m + 4)) \tan(c + dx)}{m + 3} \right)}{m + 4}}$$

↓ 3042

$$\frac{\frac{iaB(a + ia \tan(c + dx))^3 \tan^{m+1}(c + dx)}{d(m + 4)} - 2 \left(\frac{\int \tan(c + dx)^m(i \tan(c + dx)a + a)(a^3(iB(2m^3 + 17m^2 + 44m + 29) - A(2m^3 + 17m^2 + 44m + 32)) - a^3(iA(2m^3 + 19m^2 + 60m + 64) + B(2m^3 + 19m^2 + 60m + 67)) \tan(c + dx))dx}{m + 2} + \frac{(A(m + 4) - iB(m + 4)) \tan(c + dx)}{m + 3} \right)}{m + 3}}$$

$m + 4$

3.204. $\int \tan^m(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$

$$\begin{aligned} & \downarrow 4075 \\ & \frac{iaB(a + ia \tan(c + dx))^3 \tan^{m+1}(c + dx)}{d(m + 4)} - \\ & 2 \left(\frac{\int \tan^m(c+dx) (-4(A-iB)(m+2)(m+3)(m+4)a^4 - 4(iA+B)(m+2)(m+3)(m+4) \tan(c+dx)a^4) dx + \frac{a^4(A(2m^3+19m^2+60m+64) - iB(2m^3+19m^2+60m+67))}{d(m+1)}}{m+2} \right) \end{aligned}$$

m+3

$$\begin{aligned} & \downarrow 3042 \\ & \frac{iaB(a + ia \tan(c + dx))^3 \tan^{m+1}(c + dx)}{d(m + 4)} - \\ & 2 \left(\frac{\int \tan(c+dx)^m (-4(A-iB)(m+2)(m+3)(m+4)a^4 - 4(iA+B)(m+2)(m+3)(m+4) \tan(c+dx)a^4) dx + \frac{a^4(A(2m^3+19m^2+60m+64) - iB(2m^3+19m^2+60m+67))}{d(m+1)}}{m+2} \right) \end{aligned}$$

m+3

$$\begin{aligned} & \downarrow 4020 \\ & \frac{iaB(a + ia \tan(c + dx))^3 \tan^{m+1}(c + dx)}{d(m + 4)} - \\ & 2 \left(\frac{16ia^8(m+2)^2(m+3)^2(m+4)^2(A-iB)^2 \int \frac{\tan^m(c+dx)}{4a^4(m+2)(m+3)(m+4) (4(iA+B)^2(m+2)(m+3)(m+4)a^4 + 4(A-iB)(iA+B)(m+2)(m+3)(m+4) \tan(c+dx)a^4)} dx}{m+2} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{iaB(a + ia \tan(c + dx))^3 \tan^{m+1}(c + dx)}{d(m + 4)} - \\ & 2 \left(\frac{ia^4 4^{1-m} (m+2)(m+3)(m+4)(A-iB)^2 \int \frac{4^m \tan^m(c+dx)}{4(iA+B)^2(m+2)(m+3)(m+4)a^4 + 4(A-iB)(iA+B)(m+2)(m+3)(m+4) \tan(c+dx)a^4} dx}{m+2} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 74 \\ & \frac{iaB(a + ia \tan(c + dx))^3 \tan^{m+1}(c + dx)}{d(m + 4)} - \\ & 2 \left(\frac{\frac{a^4(A(2m^3+19m^2+60m+64) - iB(2m^3+19m^2+60m+67)) \tan^{m+1}(c+dx)}{d(m+1)} - \frac{4ia^4(m+2)(m+3)(m+4)(A-iB)^2 \tan^{m+1}(c+dx) \text{Hypergeometric2F1}(1, m+1, m+2, \frac{4(A-iB)(iA+B)(m+2)(m+3)(m+4)}{4a^4(m+2)(m+3)(m+4) + 4(A-iB)(iA+B)(m+2)(m+3)(m+4) \tan(c+dx)a^4})}{m+2}}{m+2} \right) \end{aligned}$$

m+3

3.204. $\int \tan^m(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx$

input `Int[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(I*a*B*Tan[c + d*x]^(1 + m)*(a + I*a*Tan[c + d*x])^3)/(d*(4 + m)) - (((A*(4 + m) - I*B*(7 + m))*Tan[c + d*x]^(1 + m)*(a^2 + I*a^2*Tan[c + d*x])^2)/(d*(3 + m)) + (2*(((A*(4 + m)^2 - I*B*(19 + 8*m + m^2))*Tan[c + d*x]^(1 + m)*(a^4 + I*a^4*Tan[c + d*x]))/(d*(2 + m)) + ((a^4*(A*(64 + 60*m + 19*m^2 + 2*m^3) - I*B*(67 + 60*m + 19*m^2 + 2*m^3))*Tan[c + d*x]^(1 + m))/(d*(1 + m)) - ((4*I)*a^4*(A - I*B)^2*(2 + m)*(3 + m)*(4 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, ((I*A + B)*Tan[c + d*x])/(A - I*B)]*Tan[c + d*x]^(1 + m))/((I*A + B)*d*(1 + m)))/(2 + m)))/(3 + m))/(4 + m)`

3.204.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 74 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4077 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.204.4 Maple [F]

$$\int (\tan^m(dx + c))(a + ia \tan(dx + c))^4 (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x)`

3.204.5 Fracas [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + ia \tan(c + dx))^4 (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^4 \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output `integral(16*((A - I*B)*a^4*e^(10*I*d*x + 10*I*c) + (A + I*B)*a^4*e^(8*I*d*x + 8*I*c))*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m/(e^(10*I*d*x + 10*I*c) + 5*e^(8*I*d*x + 8*I*c) + 10*e^(6*I*d*x + 6*I*c) + 10*e^(4*I*d*x + 4*I*c) + 5*e^(2*I*d*x + 2*I*c) + 1), x)`

3.204.6 Sympy [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx \\ &= a^4 \left(\int A \tan^m(c + dx) dx + \int (-6A \tan^2(c + dx) \tan^m(c + dx)) dx \right. \\ & \quad + \int A \tan^4(c + dx) \tan^m(c + dx) dx + \int B \tan(c + dx) \tan^m(c + dx) dx \\ & \quad + \int (-6B \tan^3(c + dx) \tan^m(c + dx)) dx + \int B \tan^5(c + dx) \tan^m(c + dx) dx \\ & \quad + \int 4iA \tan(c + dx) \tan^m(c + dx) dx + \int (-4iA \tan^3(c + dx) \tan^m(c + dx)) dx \\ & \quad \left. + \int 4iB \tan^2(c + dx) \tan^m(c + dx) dx + \int (-4iB \tan^4(c + dx) \tan^m(c + dx)) dx \right) \end{aligned}$$

input `integrate(tan(d*x+c)**m*(a+I*a*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)`

output `a**4*(Integral(A*tan(c + d*x)**m, x) + Integral(-6*A*tan(c + d*x)**2*tan(c + d*x)**m, x) + Integral(A*tan(c + d*x)**4*tan(c + d*x)**m, x) + Integral(B*tan(c + d*x)*tan(c + d*x)**m, x) + Integral(-6*B*tan(c + d*x)**3*tan(c + d*x)**m, x) + Integral(B*tan(c + d*x)**5*tan(c + d*x)**m, x) + Integral(4*I*A*tan(c + d*x)*tan(c + d*x)**m, x) + Integral(-4*I*A*tan(c + d*x)**3*tan(c + d*x)**m, x) + Integral(4*I*B*tan(c + d*x)**2*tan(c + d*x)**m, x) + Integral(-4*I*B*tan(c + d*x)**4*tan(c + d*x)**m, x))`

3.204.7 Maxima [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^4 \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^4*tan(d*x + c)^m, x)`

3.204.8 Giac [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^4 \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^4*tan(d*x + c)^m, x)`

3.204.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \tan^m(c + dx)(a + ia \tan(c + dx))^4(A + B \tan(c + dx)) dx \\ &= \int \tan(c + dx)^m (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^4 dx \end{aligned}$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^4,x)`

output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^4, x)`

3.205 $\int \tan^m(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.205.1 Optimal result	2091
3.205.2 Mathematica [A] (verified)	2092
3.205.3 Rubi [A] (verified)	2092
3.205.4 Maple [F]	2095
3.205.5 Fricas [F]	2096
3.205.6 Sympy [F]	2096
3.205.7 Maxima [F]	2097
3.205.8 Giac [F]	2097
3.205.9 Mupad [F(-1)]	2097

3.205.1 Optimal result

Integrand size = 34, antiderivative size = 205

$$\int \tan^m(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= -\frac{a^3(A(15+11m+2m^2)-iB(17+11m+2m^2)) \tan^{1+m}(c+dx)}{d(1+m)(2+m)(3+m)}$$

$$+ \frac{4a^3(A-iB) \operatorname{Hypergeometric2F1}(1, 1+m, 2+m, i \tan(c+dx)) \tan^{1+m}(c+dx)}{d(1+m)}$$

$$+ \frac{iaB \tan^{1+m}(c+dx)(a+ia \tan(c+dx))^2}{d(3+m)}$$

$$- \frac{(A(3+m)-iB(5+m)) \tan^{1+m}(c+dx)(a^3+ia^3 \tan(c+dx))}{d(2+m)(3+m)}$$

```
output -a^3*(A*(2*m^2+11*m+15)-I*B*(2*m^2+11*m+17))*tan(d*x+c)^(1+m)/d/(3+m)/(m^2
+3*m+2)+4*a^3*(A-I*B)*hypergeom([1, 1+m], [2+m], I*tan(d*x+c))*tan(d*x+c)^(1
+m)/d/(1+m)+I*a*B*tan(d*x+c)^(1+m)*(a+I*a*tan(d*x+c))^2/d/(3+m)-(A*(3+m)-I
*B*(5+m))*tan(d*x+c)^(1+m)*(a^3+I*a^3*tan(d*x+c))/d/(2+m)/(3+m)
```

3.205.2 Mathematica [A] (verified)

Time = 2.55 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.58

$$\int \tan^m(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{a^3 \tan^{1+m}(c+dx) \left(\frac{(A-iB)(-3(2+m)+4(2+m)) \operatorname{Hypergeometric2F1}(1,1+m,2+m,i \tan(c+dx))-i(1+m) \tan(c+dx)}{(1+m)(2+m)} + iB \left(\frac{1}{1+m} + \right. \right.}{d}$$

input `Integrate[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(a^3*Tan[c + d*x]^(1 + m)*(((A - I*B)*(-3*(2 + m) + 4*(2 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, I*Tan[c + d*x]] - I*(1 + m)*Tan[c + d*x]))/((1 + m)*(2 + m)) + I*B*((1 + m)^(-1) + ((2*I)*Tan[c + d*x])/(2 + m) - Tan[c + d*x]^2/(3 + m))))/d`

3.205.3 Rubi [A] (verified)

Time = 1.26 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.30, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.324$, Rules used = {3042, 4077, 25, 3042, 4077, 3042, 4075, 3042, 4020, 27, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(c + dx))^3 \tan^m(c + dx)(A + B \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int (a + ia \tan(c + dx))^3 \tan(c + dx)^m (A + B \tan(c + dx)) dx$$

$$\downarrow 4077$$

$$\int \frac{-\tan^m(c + dx)(i \tan(c + dx)a + a)^2(a(iB(m + 1) - A(m + 3)) - a(iA(m + 3) + B(m + 5)) \tan(c + dx)) dx + iaB(a + ia \tan(c + dx))^2 \tan^{m+1}(c + dx)}{d(m + 3)} dx$$

$$\downarrow 25$$

$$\frac{\frac{iaB(a + ia \tan(c + dx))^2 \tan^{m+1}(c + dx)}{d(m + 3)} - \int \tan^m(c + dx)(i \tan(c + dx)a + a)^2(a(iB(m + 1) - A(m + 3)) - a(iA(m + 3) + B(m + 5)) \tan(c + dx))dx}{m + 3}}$$

↓ 3042

$$\frac{\frac{iaB(a + ia \tan(c + dx))^2 \tan^{m+1}(c + dx)}{d(m + 3)} - \int \tan(c + dx)^m(i \tan(c + dx)a + a)^2(a(iB(m + 1) - A(m + 3)) - a(iA(m + 3) + B(m + 5)) \tan(c + dx))dx}{m + 3}}$$

↓ 4077

$$\frac{\frac{iaB(a + ia \tan(c + dx))^2 \tan^{m+1}(c + dx)}{d(m + 3)} - \int \tan^m(c + dx)(i \tan(c + dx)a + a)(a^2(iB(2m^2 + 9m + 7) - A(2m^2 + 9m + 9)) - a^2(iA(2m^2 + 11m + 15) + B(2m^2 + 11m + 17)) \tan(c + dx))dx}{m + 2} + \frac{(A(m + 1) - B(m + 1)) \tan(c + dx)}{m + 3}}$$

↓ 3042

$$\frac{\frac{iaB(a + ia \tan(c + dx))^2 \tan^{m+1}(c + dx)}{d(m + 3)} - \int \tan(c + dx)^m(i \tan(c + dx)a + a)(a^2(iB(2m^2 + 9m + 7) - A(2m^2 + 9m + 9)) - a^2(iA(2m^2 + 11m + 15) + B(2m^2 + 11m + 17)) \tan(c + dx))dx}{m + 2} + \frac{(A(m + 1) - B(m + 1)) \tan(c + dx)}{m + 3}}$$

↓ 4075

$$\frac{\frac{iaB(a + ia \tan(c + dx))^2 \tan^{m+1}(c + dx)}{d(m + 3)} - \int \tan^m(c + dx)(-4(A - iB)(m + 2)(m + 3)a^3 - 4(iA + B)(m^2 + 5m + 6) \tan(c + dx)a^3)dx + \frac{a^3(A(2m^2 + 11m + 15) - iB(2m^2 + 11m + 17)) \tan^{m+1}(c + dx)}{d(m + 1)}}{m + 2} + \frac{(A(m + 1) - B(m + 1)) \tan(c + dx)}{m + 3}}$$

↓ 3042

$$\frac{\frac{iaB(a + ia \tan(c + dx))^2 \tan^{m+1}(c + dx)}{d(m + 3)} - \int \tan(c + dx)^m(-4(A - iB)(m + 2)(m + 3)a^3 - 4(iA + B)(m^2 + 5m + 6) \tan(c + dx)a^3)dx + \frac{a^3(A(2m^2 + 11m + 15) - iB(2m^2 + 11m + 17)) \tan^{m+1}(c + dx)}{d(m + 1)}}{m + 2} + \frac{(A(m + 1) - B(m + 1)) \tan(c + dx)}{m + 3}}$$

↓ 4020

$$\frac{\frac{iaB(a + ia \tan(c + dx))^2 \tan^{m+1}(c + dx)}{d(m + 3)} - \frac{16ia^6(m + 2)^2(m + 3)^2(A - iB)^2 \int \frac{\left(\frac{(m + 2)(m + 3) \tan(c + dx)}{m^2 + 5m + 6}\right)^m}{4a^3 \left(4(iA + B)^2(m^2 + 5m + 6)^2 a^3 + 4(A - iB)(iA + B)(m + 2)^2(m + 3)^2 \tan(c + dx)a^3\right)} dx}{d}}{m + 2} + \frac{(A(m + 1) - B(m + 1)) \tan(c + dx)}{m + 3}}$$

3.205. $\int \tan^m(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\begin{aligned} & \downarrow 27 \\ & \frac{iaB(a + ia \tan(c + dx))^2 \tan^{m+1}(c + dx)}{d(m + 3)} \\ & \frac{ia^3 4^{1-m} (m+2)^2 (m+3)^2 (A-iB)^2 \int \frac{4^m \left(\frac{(m+2)(m+3) \tan(c+dx)}{m^2+5m+6}\right)^m}{4(iA+B)^2 (m^2+5m+6)^2 a^{3+4(A-iB)(iA+B)(m+2)^2(m+3)^2 \tan(c+dx)} a^3 d(-4a^3(iA+B)(m+2)(m+3) \tan(c+dx))}{d} + a^3(A}{m+2} \\ & \hline & m + 3 \end{aligned}$$

$$\begin{aligned} & \downarrow 74 \\ & \frac{iaB(a + ia \tan(c + dx))^2 \tan^{m+1}(c + dx)}{d(m + 3)} \\ & \frac{a^3(A(2m^2+11m+15) - iB(2m^2+11m+17)) \tan^{m+1}(c+dx)}{d(m+1)} - \frac{4ia^3(m+2)^2(m+3)^2(A-iB)^2 \left(\frac{(m+2)(m+3) \tan(c+dx)}{m^2+5m+6}\right)^{m+1} \text{Hypergeometric2F1}\left(1, m+1, m+2, \frac{(m+2)(m+3) \tan(c+dx)}{m^2+5m+6}\right)}{d(m+1)(m^2+5m+6)(B+iA)} \\ & \hline & m + 3 \end{aligned}$$

input `Int[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(I*a*B*Tan[c + d*x]^(1 + m)*(a + I*a*Tan[c + d*x])^2)/(d*(3 + m)) - (((A*(3 + m) - I*B*(5 + m))*Tan[c + d*x]^(1 + m)*(a^3 + I*a^3*Tan[c + d*x]))/(d*(2 + m)) + ((a^3*(A*(15 + 11*m + 2*m^2) - I*B*(17 + 11*m + 2*m^2))*Tan[c + d*x]^(1 + m))/(d*(1 + m)) - ((4*I)*a^3*(A - I*B)^2*(2 + m)^2*(3 + m)^2*Hypergeometric2F1[1, 1 + m, 2 + m, ((I*A + B)*Tan[c + d*x])/(A - I*B)]*(((2 + m)*(3 + m)*Tan[c + d*x])/(6 + 5*m + m^2))^(1 + m))/((I*A + B)*d*(1 + m)*(6 + 5*m + m^2)))/(2 + m))/(3 + m)`

3.205.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 74 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))`

3.205. $\int \tan^m(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4077 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B))*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.205.4 Maple [F]

$$\int (\tan^m(dx + c))(a + ia \tan(dx + c))^3 (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x)`

3.205.5 Fricas [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^3 \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(8*((A - I*B)*a^3*e^(8*I*d*x + 8*I*c) + (A + I*B)*a^3*e^(6*I*d*x + 6*I*c))*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m/(e^(8*I*d*x + 8*I*c) + 4*e^(6*I*d*x + 6*I*c) + 6*e^(4*I*d*x + 4*I*c) + 4*e^(2*I*d*x + 2*I*c) + 1), x)`

3.205.6 Sympy [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= -ia^3 \left(\int ia \tan^m(c + dx) dx + \int (-3A \tan(c + dx) \tan^m(c + dx)) dx \right.$$

$$+ \int A \tan^3(c + dx) \tan^m(c + dx) dx + \int (-3B \tan^2(c + dx) \tan^m(c + dx)) dx$$

$$+ \int B \tan^4(c + dx) \tan^m(c + dx) dx + \int (-3iA \tan^2(c + dx) \tan^m(c + dx)) dx$$

$$\left. + \int iB \tan(c + dx) \tan^m(c + dx) dx + \int (-3iB \tan^3(c + dx) \tan^m(c + dx)) dx \right)$$

input `integrate(tan(d*x+c)**m*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `-I*a**3*(Integral(I*A*tan(c + d*x)**m, x) + Integral(-3*A*tan(c + d*x)*tan(c + d*x)**m, x) + Integral(A*tan(c + d*x)**3*tan(c + d*x)**m, x) + Integral(-3*B*tan(c + d*x)**2*tan(c + d*x)**m, x) + Integral(B*tan(c + d*x)**4*tan(c + d*x)**m, x) + Integral(-3*I*A*tan(c + d*x)**2*tan(c + d*x)**m, x) + Integral(I*B*tan(c + d*x)*tan(c + d*x)**m, x) + Integral(-3*I*B*tan(c + d*x)**3*tan(c + d*x)**m, x))`

3.205.7 Maxima [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^3 \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^3*tan(d*x + c)^m, x)`

3.205.8 Giac [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^3 \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^3*tan(d*x + c)^m, x)`

3.205.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \tan^m(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ &= \int \tan(c + dx)^m (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^3 dx \end{aligned}$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^3,x)`

output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^3, x)`

3.206 $\int \tan^m(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.206.1 Optimal result	2098
3.206.2 Mathematica [A] (verified)	2099
3.206.3 Rubi [A] (verified)	2099
3.206.4 Maple [F]	2102
3.206.5 Fricas [F]	2102
3.206.6 Sympy [F]	2103
3.206.7 Maxima [F]	2103
3.206.8 Giac [F]	2104
3.206.9 Mupad [F(-1)]	2104

3.206.1 Optimal result

Integrand size = 34, antiderivative size = 132

$$\int \tan^m(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{ia^2(B+(iA+B)(2+m)) \tan^{1+m}(c+dx)}{d(1+m)(2+m)}$$

$$+ \frac{2a^2(A-iB) \operatorname{Hypergeometric2F1}(1,1+m,2+m,i \tan(c+dx)) \tan^{1+m}(c+dx)}{d(1+m)}$$

$$+ \frac{iB \tan^{1+m}(c+dx)(a^2+ia^2 \tan(c+dx))}{d(2+m)}$$

```
output I*a^2*(B+(I*A+B)*(2+m))*tan(d*x+c)^(1+m)/d/(1+m)/(2+m)+2*a^2*(A-I*B)*hyper
geom([1, 1+m], [2+m], I*tan(d*x+c))*tan(d*x+c)^(1+m)/d/(1+m)+I*B*tan(d*x+c)^(
1+m)*(a^2+I*a^2*tan(d*x+c))/d/(2+m)
```

3.206.2 Mathematica [A] (verified)

Time = 1.57 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.61

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{a^2 \tan^{1+m}(c + dx)((A - 2iB)(2 + m) - 2(A - iB)(2 + m) \operatorname{Hypergeometric2F1}(1, 1 + m, 2 + m, i \tan(c + dx)))}{d(1 + m)(2 + m)}$$

input `Integrate[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `-((a^2*Tan[c + d*x]^(1 + m)*((A - (2*I)*B)*(2 + m) - 2*(A - I*B)*(2 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, I*Tan[c + d*x]] + B*(1 + m)*Tan[c + d*x]))/(d*(1 + m)*(2 + m))`

3.206.3 Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.24, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4077, 25, 3042, 4075, 3042, 4020, 27, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(c + dx))^2 \tan^m(c + dx)(A + B \tan(c + dx)) dx$$

↓ 3042

$$\int (a + ia \tan(c + dx))^2 \tan(c + dx)^m (A + B \tan(c + dx)) dx$$

↓ 4077

$$\frac{\int -\tan^m(c + dx)(i \tan(c + dx)a + a)(a(iB(m + 1) - A(m + 2)) - a(B + (iA + B)(m + 2)) \tan(c + dx)) dx}{\frac{iB(a^2 + ia^2 \tan(c + dx)) \tan^{m+1}(c + dx)}{d(m + 2)}} +$$

↓ 25

$$\frac{iB(a^2 + ia^2 \tan(c + dx)) \tan^{m+1}(c + dx)}{d(m + 2)} - \frac{\int \tan^m(c + dx)(i \tan(c + dx)a + a)(a(iB(m + 1) - A(m + 2)) - a(B + (iA + B)(m + 2)) \tan(c + dx))dx}{m + 2}$$

↓ 3042

$$\frac{iB(a^2 + ia^2 \tan(c + dx)) \tan^{m+1}(c + dx)}{d(m + 2)} - \frac{\int \tan(c + dx)^m(i \tan(c + dx)a + a)(a(iB(m + 1) - A(m + 2)) - a(B + (iA + B)(m + 2)) \tan(c + dx))dx}{m + 2}$$

↓ 4075

$$\frac{iB(a^2 + ia^2 \tan(c + dx)) \tan^{m+1}(c + dx)}{d(m + 2)} - \frac{\int \tan^m(c + dx) (-2(A - iB)(m + 2)a^2 - 2(iA + B)(m + 2) \tan(c + dx)a^2) dx - \frac{ia^2(B+(m+2)(B+iA)) \tan^{m+1}(c+dx)}{d(m+1)}}{m + 2}$$

↓ 3042

$$\frac{iB(a^2 + ia^2 \tan(c + dx)) \tan^{m+1}(c + dx)}{d(m + 2)} - \frac{\int \tan(c + dx)^m (-2(A - iB)(m + 2)a^2 - 2(iA + B)(m + 2) \tan(c + dx)a^2) dx - \frac{ia^2(B+(m+2)(B+iA)) \tan^{m+1}(c+dx)}{d(m+1)}}{m + 2}$$

↓ 4020

$$\frac{iB(a^2 + ia^2 \tan(c + dx)) \tan^{m+1}(c + dx)}{d(m + 2)} - \frac{4ia^4(m+2)^2(A-iB)^2 \int \frac{\tan^m(c+dx)}{2a^2(m+2)(2(iA+B)^2(m+2)a^2+2(A-iB)(iA+B)(m+2) \tan(c+dx)a^2)} d(-2a^2(iA+B)(m+2) \tan(c+dx))}{d} - \frac{ia^2(B+(m+2)(B+iA)) \tan^{m+1}(c+dx)}{d(m+1)}}{m + 2}$$

↓ 27

$$\frac{iB(a^2 + ia^2 \tan(c + dx)) \tan^{m+1}(c + dx)}{d(m + 2)} - \frac{ia^2 2^{1-m}(m+2)(A-iB)^2 \int \frac{2^m \tan^m(c+dx)}{2(iA+B)^2(m+2)a^2+2(A-iB)(iA+B)(m+2) \tan(c+dx)a^2} d(-2a^2(iA+B)(m+2) \tan(c+dx))}{d} - \frac{ia^2(B+(m+2)(B+iA)) \tan^{m+1}(c+dx)}{d(m+1)}}{m + 2}$$

↓ 74

$$\frac{iB(a^2 + ia^2 \tan(c + dx)) \tan^{m+1}(c + dx)}{d(m + 2)} - \frac{2ia^2(m+2)(A-iB)^2 \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, m+1, m+2, \frac{(iA+B) \tan(c+dx)}{A-iB}\right)}{d(m+1)(B+iA)} - \frac{ia^2(B+(m+2)(B+iA)) \tan^{m+1}(c+dx)}{d(m+1)}}{m + 2}$$

3.206. $\int \tan^m(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

input `Int[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(I*B*Tan[c + d*x]^(1 + m)*(a^2 + I*a^2*Tan[c + d*x]))/(d*(2 + m)) - (((-I)*a^2*(B + (I*A + B)*(2 + m))*Tan[c + d*x]^(1 + m))/(d*(1 + m)) - ((2*I)*a^2*(A - I*B)^2*(2 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, ((I*A + B)*Tan[c + d*x])/(A - I*B)]*Tan[c + d*x]^(1 + m))/((I*A + B)*d*(1 + m)))/(2 + m)`

3.206.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 74 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4077 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.206.4 Maple [F]

$$\int (\tan^m(dx + c))(a + ia \tan(dx + c))^2 (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x)`

3.206.5 Fracas [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + ia \tan(c + dx))^2 (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^2 \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(4*((A - I*B)*a^2*e^(6*I*d*x + 6*I*c) + (A + I*B)*a^2*e^(4*I*d*x + 4*I*c))*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m/(e^(6*I*d*x + 6*I*c) + 3*e^(4*I*d*x + 4*I*c) + 3*e^(2*I*d*x + 2*I*c) + 1), x)`

3.206.6 Sympy [F]

$$\begin{aligned} & \int \tan^m(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx \\ &= -a^2 \left(\int (-A \tan^m(c+dx)) dx + \int A \tan^2(c+dx) \tan^m(c+dx) dx \right. \\ & \quad + \int (-B \tan(c+dx) \tan^m(c+dx)) dx + \int B \tan^3(c+dx) \tan^m(c+dx) dx \\ & \quad \left. + \int (-2iA \tan(c+dx) \tan^m(c+dx)) dx + \int (-2iB \tan^2(c+dx) \tan^m(c+dx)) dx \right) \end{aligned}$$

input `integrate(tan(d*x+c)**m*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `-a**2*(Integral(-A*tan(c+d*x)**m,x)+Integral(A*tan(c+d*x)**2*tan(c+d*x)**m,x)+Integral(-B*tan(c+d*x)*tan(c+d*x)**m,x)+Integral(B*tan(c+d*x)**3*tan(c+d*x)**m,x)+Integral(-2*I*A*tan(c+d*x)*tan(c+d*x)**m,x)+Integral(-2*I*B*tan(c+d*x)**2*tan(c+d*x)**m,x))`

3.206.7 Maxima [F]

$$\begin{aligned} & \int \tan^m(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx \\ &= \int (B \tan(dx+c) + A)(ia \tan(dx+c) + a)^2 \tan(dx+c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x+c)+A)*(I*a*tan(d*x+c)+a)^2*tan(d*x+c)^m,x)`

3.206.8 Giac [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^2 \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^2*tan(d*x + c)^m, x)`

3.206.9 Mupad [F(-1)]

Timed out.

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^m (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^2 dx$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^2,x)`

output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^2, x)`

3.207 $\int \tan^m(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.207.1 Optimal result	2105
3.207.2 Mathematica [A] (verified)	2105
3.207.3 Rubi [A] (verified)	2106
3.207.4 Maple [F]	2108
3.207.5 Fracas [F]	2108
3.207.6 Sympy [F]	2108
3.207.7 Maxima [F]	2109
3.207.8 Giac [F]	2109
3.207.9 Mupad [F(-1)]	2109

3.207.1 Optimal result

Integrand size = 32, antiderivative size = 70

$$\int \tan^m(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{iaB \tan^{1+m}(c+dx)}{d(1+m)} + \frac{a(A-iB) \operatorname{Hypergeometric2F1}(1, 1+m, 2+m, i \tan(c+dx)) \tan^{1+m}(c+dx)}{d(1+m)}$$

output `I*a*B*tan(d*x+c)^(1+m)/d/(1+m)+a*(A-I*B)*hypergeom([1, 1+m], [2+m], I*tan(d*x+c))*tan(d*x+c)^(1+m)/d/(1+m)`

3.207.2 Mathematica [A] (verified)

Time = 0.52 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.74

$$\int \tan^m(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{a(ib + (A - ib) \operatorname{Hypergeometric2F1}(1, 1+m, 2+m, i \tan(c+dx))) \tan^{1+m}(c+dx)}{d(1+m)}$$

input `Integrate[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output $(a*(I*B + (A - I*B)*\text{Hypergeometric2F1}[1, 1 + m, 2 + m, I*\text{Tan}[c + d*x]])*T\text{an}[c + d*x]^{(1 + m)})/(d*(1 + m))$

3.207.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.39, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3042, 4075, 3042, 4020, 27, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(c + dx)) \tan^m(c + dx) (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(c + dx)) \tan(c + dx)^m (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4075} \\
 & \int \tan^m(c + dx) (a(A - iB) + a(iA + B) \tan(c + dx)) dx + \frac{iaB \tan^{m+1}(c + dx)}{d(m + 1)} \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c + dx)^m (a(A - iB) + a(iA + B) \tan(c + dx)) dx + \frac{iaB \tan^{m+1}(c + dx)}{d(m + 1)} \\
 & \quad \downarrow \text{4020} \\
 & \frac{ia^2(A - iB)^2 \int \frac{\tan^m(c + dx)}{a(iA + B)^2 + a(A - iB) \tan(c + dx)(iA + B)} d(a(iA + B) \tan(c + dx))}{d} + \frac{iaB \tan^{m+1}(c + dx)}{d(m + 1)} \\
 & \quad \downarrow \text{27} \\
 & \frac{ia(A - iB)^2 \int \frac{\tan^m(c + dx)}{a(iA + B)^2 + a(A - iB) \tan(c + dx)(iA + B)} d(a(iA + B) \tan(c + dx))}{d} + \frac{iaB \tan^{m+1}(c + dx)}{d(m + 1)} \\
 & \quad \downarrow \text{74} \\
 & \frac{ia(A - iB)^2 \tan^{m+1}(c + dx) \text{Hypergeometric2F1}\left(1, m + 1, m + 2, \frac{(iA + B) \tan(c + dx)}{A - iB}\right)}{d(m + 1)(B + iA)} + \frac{iaB \tan^{m+1}(c + dx)}{d(m + 1)}
 \end{aligned}$$

input `Int[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(I*a*B*Tan[c + d*x]^(1 + m))/(d*(1 + m)) + (I*a*(A - I*B)^2*Hypergeometric2F1[1, 1 + m, 2 + m, ((I*A + B)*Tan[c + d*x])/(A - I*B)]*Tan[c + d*x]^(1 + m))/((I*A + B)*d*(1 + m))`

3.207.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 74 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.207.4 Maple [F]

$$\int (\tan^m(dx+c))(a+ia \tan(dx+c))(A+B \tan(dx+c)) dx$$

input `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

3.207.5 Fricas [F]

$$\begin{aligned} & \int \tan^m(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx \\ &= \int (B \tan(dx+c)+A)(ia \tan(dx+c)+a) \tan(dx+c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(2*((A - I*B)*a*e^(4*I*d*x + 4*I*c) + (A + I*B)*a*e^(2*I*d*x + 2*I*c))*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m/(e^(4*I*d*x + 4*I*c) + 2*e^(2*I*d*x + 2*I*c) + 1), x)`

3.207.6 Sympy [F]

$$\begin{aligned} & \int \tan^m(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx \\ &= ia \left(\int (-iA \tan^m(c+dx)) dx + \int A \tan(c+dx) \tan^m(c+dx) dx \right. \\ & \quad \left. + \int B \tan^2(c+dx) \tan^m(c+dx) dx + \int (-iB \tan(c+dx) \tan^m(c+dx)) dx \right) \end{aligned}$$

input `integrate(tan(d*x+c)**m*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `I*a*(Integral(-I*A*tan(c + d*x)**m, x) + Integral(A*tan(c + d*x)*tan(c + d*x)**m, x) + Integral(B*tan(c + d*x)**2*tan(c + d*x)**m, x) + Integral(-I*B*tan(c + d*x)*tan(c + d*x)**m, x))`

3.207.7 Maxima [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a) \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)*tan(d*x + c)^m, x)`

3.207.8 Giac [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a) \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)*tan(d*x + c)^m, x)`

3.207.9 Mupad [F(-1)]

Timed out.

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^m (A + B \tan(c + dx)) (a + a \tan(c + dx) li) dx$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`

output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i), x)`

3.208
$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

3.208.1 Optimal result 2110
 3.208.2 Mathematica [A] (verified) 2111
 3.208.3 Rubi [A] (verified) 2111
 3.208.4 Maple [F] 2113
 3.208.5 Fricas [F] 2114
 3.208.6 Sympy [F] 2114
 3.208.7 Maxima [F(-2)] 2114
 3.208.8 Giac [F] 2115
 3.208.9 Mupad [F(-1)] 2115

3.208.1 Optimal result

Integrand size = 34, antiderivative size = 168

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= \frac{(A(1-m) - iB(1+m)) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right) \tan^{1+m}(c+dx)}{2ad(1+m)}$$

$$+ \frac{(iA - B)m \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, -\tan^2(c+dx)\right) \tan^{2+m}(c+dx)}{2ad(2+m)}$$

$$+ \frac{(A+iB) \tan^{1+m}(c+dx)}{2d(a+ia \tan(c+dx))}$$

```
output 1/2*(A*(1-m)-I*B*(1+m))*hypergeom([1, 1/2+1/2*m], [3/2+1/2*m], -tan(d*x+c)^2)
*tan(d*x+c)^(1+m)/a/d/(1+m)+1/2*(I*A-B)*m*hypergeom([1, 1+1/2*m], [2+1/2*m],
]-tan(d*x+c)^2)*tan(d*x+c)^(2+m)/a/d/(2+m)+1/2*(A+I*B)*tan(d*x+c)^(1+m)/d
/(a+I*a*tan(d*x+c))
```

3.208.2 Mathematica [A] (verified)

Time = 1.89 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.82

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= \frac{\tan^{1+m}(c+dx) \left(-\frac{(A(-1+m)+iB(1+m)) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right)}{1+m} + \frac{i(A+iB)m \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{2+m}{2}, -\tan^2(c+dx)\right)}{2+m} \right)}{2ad}$$

input `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `(Tan[c + d*x]^(1 + m)*(-(((A*(-1 + m) + I*B*(1 + m))*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2]/(1 + m)) + (I*(A + I*B)*m*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x])/(2 + m) + ((-I)*A + B)/(-I + Tan[c + d*x]))) / (2*a*d)`

3.208.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.206$, Rules used = {3042, 4079, 3042, 4021, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)^m(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$\downarrow \text{4079}$$

$$\frac{\int \tan^m(c+dx)(a(-mA+A-iB(m+1))+a(iA-B)m \tan(c+dx))dx}{2a^2} + \frac{(A+iB) \tan^{m+1}(c+dx)}{2d(a+ia \tan(c+dx))}$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
& \frac{\int \tan(c+dx)^m (a(-mA + A - iB(m+1)) + a(iA - B)m \tan(c+dx)) dx}{\frac{2a^2}{(A+iB) \tan^{m+1}(c+dx)} \cdot \frac{1}{2d(a+ia \tan(c+dx))}} + \\
& \quad \downarrow \text{4021} \\
& \frac{am(-B+iA) \int \tan^{m+1}(c+dx) dx + a(A(-m) + A - iB(m+1)) \int \tan^m(c+dx) dx}{\frac{2a^2}{(A+iB) \tan^{m+1}(c+dx)} \cdot \frac{1}{2d(a+ia \tan(c+dx))}} + \\
& \quad \downarrow \text{3042} \\
& \frac{a(A(-m) + A - iB(m+1)) \int \tan(c+dx)^m dx + am(-B+iA) \int \tan(c+dx)^{m+1} dx}{\frac{2a^2}{(A+iB) \tan^{m+1}(c+dx)} \cdot \frac{1}{2d(a+ia \tan(c+dx))}} + \\
& \quad \downarrow \text{3957} \\
& \frac{\frac{am(-B+iA) \int \frac{\tan^{m+1}(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} + \frac{a(A(-m)+A-iB(m+1)) \int \frac{\tan^m(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d}}{\frac{2a^2}{(A+iB) \tan^{m+1}(c+dx)} \cdot \frac{1}{2d(a+ia \tan(c+dx))}} + \\
& \quad \downarrow \text{278} \\
& \frac{\frac{a(A(-m)+A-iB(m+1)) \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+1}{2}, \frac{m+3}{2}, -\tan^2(c+dx)\right)}{d(m+1)} + \frac{am(-B+iA) \tan^{m+2}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+4}{2}, -\tan^2(c+dx)\right)}{d(m+2)}}{\frac{2a^2}{(A+iB) \tan^{m+1}(c+dx)} \cdot \frac{1}{2d(a+ia \tan(c+dx))}}
\end{aligned}$$

input `Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `((A + I*B)*Tan[c + d*x]^(1 + m))/(2*d*(a + I*a*Tan[c + d*x])) + ((a*(A - A*m - I*B*(1 + m))*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(1 + m))/(d*(1 + m)) + (a*(I*A - B)*m*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(2 + m))/(d*(2 + m))/(2*a^2)`

3.208.3.1 Defintions of rubi rules used

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a), x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4021 `Int[((b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Tan[e + f*x])^m, x], x] + Simp[d/b Int[(b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x] && NeQ[c^2 + d^2, 0] && !IntegerQ[2*m]`

rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.208.4 Maple [F]

$$\int \frac{(\tan^m(dx + c))(A + B \tan(dx + c))}{a + ia \tan(dx + c)} dx$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)`

3.208.5 Fricas [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{ia\tan(dx+c)+a} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="fricas")`

output `integral(1/2*((A - I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m*e^(-2*I*d*x - 2*I*c)/a, x)`

3.208.6 Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = -\frac{i\left(\int \frac{A\tan^m(c+dx)}{\tan(c+dx)-i} dx + \int \frac{B\tan(c+dx)\tan^m(c+dx)}{\tan(c+dx)-i} dx\right)}{a}$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)`

output `-I*(Integral(A*tan(c + d*x)**m/(tan(c + d*x) - I), x) + Integral(B*tan(c + d*x)*tan(c + d*x)**m/(tan(c + d*x) - I), x))/a`

3.208.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.208.8 Giac [F]

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \int \frac{(B \tan(dx+c)+A) \tan(dx+c)^m}{ia \tan(dx+c)+a} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(I*a*tan(d*x + c) + a), x)`

3.208.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \int \frac{\tan(c+dx)^m (A+B \tan(c+dx))}{a+a \tan(c+dx) li} dx$$

input `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i),x)`

output `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i), x)`

3.209
$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.209.1 Optimal result 2116
 3.209.2 Mathematica [A] (verified) 2117
 3.209.3 Rubi [A] (verified) 2117
 3.209.4 Maple [F] 2120
 3.209.5 Fricas [F] 2120
 3.209.6 Sympy [F] 2121
 3.209.7 Maxima [F(-2)] 2121
 3.209.8 Giac [F] 2121
 3.209.9 Mupad [F(-1)] 2122

3.209.1 Optimal result

Integrand size = 34, antiderivative size = 226

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$= \frac{(1-m)(A(1-m)-iB(1+m)) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right) \tan^{1+m}(c+dx)}{4a^2d(1+m)}$$

$$+ \frac{(A(2-m)-iBm) \tan^{1+m}(c+dx)}{4a^2d(1+i \tan(c+dx))}$$

$$+ \frac{m(iA(2-m)+Bm) \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, -\tan^2(c+dx)\right) \tan^{2+m}(c+dx)}{4a^2d(2+m)}$$

$$+ \frac{(A+iB) \tan^{1+m}(c+dx)}{4d(a+ia \tan(c+dx))^2}$$

```
output 1/4*(1-m)*(A*(1-m)-I*B*(1+m))*hypergeom([1, 1/2+1/2*m], [3/2+1/2*m], -tan(d*x+c)^2)*tan(d*x+c)^(1+m)/a^2/d/(1+m)+1/4*(A*(2-m)-I*B*m)*tan(d*x+c)^(1+m)/a^2/d/(1+I*tan(d*x+c))+1/4*m*(I*A*(2-m)+B*m)*hypergeom([1, 1+1/2*m], [2+1/2*m], -tan(d*x+c)^2)*tan(d*x+c)^(2+m)/a^2/d/(2+m)+1/4*(A+I*B)*tan(d*x+c)^(1+m)/d/(a+I*a*tan(d*x+c))^2
```

3.209.2 Mathematica [A] (verified)

Time = 3.22 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.77

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$= \frac{\tan^{1+m}(c+dx) \left(\frac{2a(-1+m)(A(-1+m)+iB(1+m)) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right)}{1+m} + \frac{2am(-iA(-2+m)+Bm) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right)}{1+m} \right)}{8a^3d}$$

input `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `(Tan[c + d*x]^(1 + m)*((2*a*(-1 + m)*(A*(-1 + m) + I*B*(1 + m))*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2])/(1 + m) + (2*a*m*((-I)*A*(-2 + m) + B*m)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x])/(2 + m) - (2*a*(A + I*B))/(-I + Tan[c + d*x])^2 + ((2*I)*a*A*(-2 + m) - 2*a*B*m)/(-I + Tan[c + d*x]))/(8*a^3*d)`

3.209.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3042, 4079, 3042, 4079, 27, 3042, 4021, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

↓ 3042

$$\int \frac{\tan(c+dx)^m(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

↓ 4079

$$\frac{\int \frac{\tan^m(c+dx)(a(A(3-m)-iB(m+1))-a(iA-B)(1-m) \tan(c+dx))}{i \tan(c+dx)a+a} dx}{4a^2} + \frac{(A+iB) \tan^{m+1}(c+dx)}{4d(a+ia \tan(c+dx))^2}$$

↓ 3042

$$\begin{aligned}
& \frac{\int \frac{\tan(c+dx)^m (a(A(3-m)-iB(m+1))-a(iA-B)(1-m)\tan(c+dx))}{i \tan(c+dx)a+a} dx}{4a^2} + \frac{(A+iB)\tan^{m+1}(c+dx)}{4d(a+ia\tan(c+dx))^2} \\
& \quad \downarrow \text{4079} \\
& \frac{\int 2 \tan^m(c+dx) \left((1-m)(A(1-m)-iB(m+1))a^2 + m(iA(2-m)+Bm)\tan(c+dx)a^2 \right) dx}{2a^2} + \frac{(A(2-m)-iBm)\tan^{m+1}(c+dx)}{d(1+i\tan(c+dx))} + \\
& \quad \frac{4a^2}{4d(a+ia\tan(c+dx))^2} (A+iB)\tan^{m+1}(c+dx) \\
& \quad \downarrow \text{27} \\
& \frac{\int \tan^m(c+dx) \left((1-m)(A(1-m)-iB(m+1))a^2 + m(iA(2-m)+Bm)\tan(c+dx)a^2 \right) dx}{a^2} + \frac{(A(2-m)-iBm)\tan^{m+1}(c+dx)}{d(1+i\tan(c+dx))} + \\
& \quad \frac{4a^2}{4d(a+ia\tan(c+dx))^2} (A+iB)\tan^{m+1}(c+dx) \\
& \quad \downarrow \text{3042} \\
& \frac{\int \tan(c+dx)^m \left((1-m)(A(1-m)-iB(m+1))a^2 + m(iA(2-m)+Bm)\tan(c+dx)a^2 \right) dx}{a^2} + \frac{(A(2-m)-iBm)\tan^{m+1}(c+dx)}{d(1+i\tan(c+dx))} + \\
& \quad \frac{4a^2}{4d(a+ia\tan(c+dx))^2} (A+iB)\tan^{m+1}(c+dx) \\
& \quad \downarrow \text{4021} \\
& \frac{a^2 m(Bm+iA(2-m)) \int \tan^{m+1}(c+dx) dx + a^2(1-m)(A(1-m)-iB(m+1)) \int \tan^m(c+dx) dx}{a^2} + \frac{(A(2-m)-iBm)\tan^{m+1}(c+dx)}{d(1+i\tan(c+dx))} + \\
& \quad \frac{4a^2}{4d(a+ia\tan(c+dx))^2} (A+iB)\tan^{m+1}(c+dx) \\
& \quad \downarrow \text{3042} \\
& \frac{a^2(1-m)(A(1-m)-iB(m+1)) \int \tan(c+dx)^m dx + a^2 m(Bm+iA(2-m)) \int \tan(c+dx)^{m+1} dx}{a^2} + \frac{(A(2-m)-iBm)\tan^{m+1}(c+dx)}{d(1+i\tan(c+dx))} + \\
& \quad \frac{4a^2}{4d(a+ia\tan(c+dx))^2} (A+iB)\tan^{m+1}(c+dx) \\
& \quad \downarrow \text{3957} \\
& \frac{a^2 m(Bm+iA(2-m)) \int \frac{\tan^{m+1}(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} + \frac{a^2(1-m)(A(1-m)-iB(m+1)) \int \frac{\tan^m(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} + \frac{(A(2-m)-iBm)\tan^{m+1}(c+dx)}{d(1+i\tan(c+dx))} + \\
& \quad \frac{4a^2}{4d(a+ia\tan(c+dx))^2} (A+iB)\tan^{m+1}(c+dx)
\end{aligned}$$

3.209. $\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$

↓ 278

$$\frac{\frac{a^{2(1-m)}(A(1-m)-iB(m+1)) \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+1}{2}, \frac{m+3}{2}, -\tan^2(c+dx)\right)}{d(m+1)} + \frac{a^{2m}(Bm+iA(2-m)) \tan^{m+2}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m}{2}, \frac{m+2}{2}, -\tan^2(c+dx)\right)}{d(m+2)}}{a^2} = \frac{(A+iB) \tan^{m+1}(c+dx)}{4d(a+ia \tan(c+dx))^2} 4a^2$$

input `Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `((A + I*B)*Tan[c + d*x]^(1 + m))/(4*d*(a + I*a*Tan[c + d*x])^2) + (((A*(2 - m) - I*B*m)*Tan[c + d*x]^(1 + m))/(d*(1 + I*Tan[c + d*x])) + ((a^2*(1 - m)*(A*(1 - m) - I*B*(1 + m))*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(1 + m))/(d*(1 + m)) + (a^2*m*(I*A*(2 - m) + B*m)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(2 + m))/(d*(2 + m)))/a^2)/(4*a^2)`

3.209.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4021 `Int[((b_.)*tan[(e_.) + (f_.)*(x_.)]^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[c Int[(b*Tan[e + f*x])^m, x], x] + Simp[d/b Int[(b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x] && NeQ[c^2 + d^2, 0] && !IntegerQ[2*m]`

rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)]^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)]*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]^(n_)), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.209.4 Maple [F]

$$\int \frac{(\tan^m(dx + c))(A + B \tan(dx + c))}{(a + ia \tan(dx + c))^2} dx$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x)`

output `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x)`

3.209.5 Fracas [F]

$$\int \frac{\tan^m(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^2} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)^m}{(ia \tan(dx + c) + a)^2} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="fricas")`

output `integral(1/4*((A - I*B)*e^(4*I*d*x + 4*I*c) + 2*A*e^(2*I*d*x + 2*I*c) + A + I*B)*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m*e^(-4*I*d*x - 4*I*c)/a^2, x)`

3.209.6 Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= -\frac{\int \frac{A\tan^m(c+dx)}{\tan^2(c+dx)-2i\tan(c+dx)-1} dx + \int \frac{B\tan(c+dx)\tan^m(c+dx)}{\tan^2(c+dx)-2i\tan(c+dx)-1} dx}{a^2}$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**2,x)`

output `-(Integral(A*tan(c + d*x)**m/(tan(c + d*x)**2 - 2*I*tan(c + d*x) - 1), x) + Integral(B*tan(c + d*x)*tan(c + d*x)**m/(tan(c + d*x)**2 - 2*I*tan(c + d*x) - 1), x))/a**2`

3.209.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.209.8 Giac [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{(ia\tan(dx+c)+a)^2} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(I*a*tan(d*x + c) + a)^2, x)`

3.209. $\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$

3.209.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx = \int \frac{\tan(c+dx)^m (A+B \tan(c+dx))}{(a+a \tan(c+dx) \text{ li})^2} dx$$

input `int((tan(c + d*x))^m*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^2,x)`

output `int((tan(c + d*x))^m*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^2, x)`

3.210
$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.210.1 Optimal result 2123
 3.210.2 Mathematica [A] (verified) 2124
 3.210.3 Rubi [A] (verified) 2124
 3.210.4 Maple [F] 2128
 3.210.5 Fricas [F] 2128
 3.210.6 Sympy [F] 2129
 3.210.7 Maxima [F(-2)] 2129
 3.210.8 Giac [F] 2129
 3.210.9 Mupad [F(-1)] 2130

3.210.1 Optimal result

Integrand size = 34, antiderivative size = 308

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx =$$

$$\frac{(1-m)(iB(3+m-2m^2)-A(3-7m+2m^2)) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right) \tan^{2+m}(c+dx)}{24a^3d(1+m)}$$

$$+ \frac{(2-m)m(B+iA(5-2m)+2Bm) \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, -\tan^2(c+dx)\right) \tan^{2+m}(c+dx)}{24a^3d(2+m)}$$

$$+ \frac{(A+iB) \tan^{1+m}(c+dx)}{6d(a+ia \tan(c+dx))^3} + \frac{(iB(1-2m)+A(7-2m)) \tan^{1+m}(c+dx)}{24ad(a+ia \tan(c+dx))^2}$$

$$+ \frac{(2-m)(A(5-2m)-i(B+2Bm)) \tan^{1+m}(c+dx)}{24d(a^3+ia^3 \tan(c+dx))}$$

output

```
-1/24*(1-m)*(I*B*(-2*m^2+m+3)-A*(2*m^2-7*m+3))*hypergeom([1, 1/2+1/2*m], [3/2+1/2*m], -tan(d*x+c)^2)*tan(d*x+c)^(1+m)/a^3/d/(1+m)+1/24*(2-m)*m*(B+I*A*(5-2*m)+2*B*m)*hypergeom([1, 1+1/2*m], [2+1/2*m], -tan(d*x+c)^2)*tan(d*x+c)^(2+m)/a^3/d/(2+m)+1/6*(A+I*B)*tan(d*x+c)^(1+m)/d/(a+I*a*tan(d*x+c))^3+1/24*(I*B*(1-2*m)+A*(7-2*m))*tan(d*x+c)^(1+m)/a/d/(a+I*a*tan(d*x+c))^2+1/24*(2-m)*(A*(5-2*m)-I*(2*B*m+B))*tan(d*x+c)^(1+m)/d/(a^3+I*a^3*tan(d*x+c))
```

3.210.2 Mathematica [A] (verified)

Time = 4.33 (sec) , antiderivative size = 227, normalized size of antiderivative = 0.74

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{\tan^{1+m}(c+dx) \left(\frac{(-1+m)(iB(3+m-2m^2)+A(-3+7m-2m^2)) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right) - (-2+m)m(B+2B)}{1+m} \right)}{}$$

input `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `(Tan[c + d*x]^(1 + m)*(((−1 + m)*(I*B*(3 + m − 2*m^2) + A*(−3 + 7*m − 2*m^2))*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, −Tan[c + d*x]^2])/(1 + m) − ((−2 + m)*m*(B + 2*B*m − I*A*(−5 + 2*m))*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, −Tan[c + d*x]^2]*Tan[c + d*x])/(2 + m) + ((4*I)*(A + I*B))/(-I + Tan[c + d*x])^3 + (A*(-7 + 2*m) + I*B*(-1 + 2*m))/(-I + Tan[c + d*x])^2 + ((−2 + m)*(B + 2*B*m − I*A*(−5 + 2*m)))/(-I + Tan[c + d*x]))/(24*a^3*d)`

3.210.3 Rubi [A] (verified)

Time = 1.51 (sec) , antiderivative size = 319, normalized size of antiderivative = 1.04, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.382$, Rules used = {3042, 4079, 3042, 4079, 25, 3042, 4079, 27, 3042, 4021, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

↓ 3042

$$\int \frac{\tan(c+dx)^m(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

↓ 4079

$$\int \frac{\tan^m(c+dx)(a(A(5-m)-iB(m+1))-a(iA-B)(2-m)\tan(c+dx))}{6a^2(i \tan(c+dx)a+a)^2} dx + \frac{(A+iB) \tan^{m+1}(c+dx)}{6d(a+ia \tan(c+dx))^3}$$

↓ 3042

3.210. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\frac{\int \frac{\tan(c+dx)^m (a(A(5-m)-iB(m+1))-a(iA-B)(2-m)\tan(c+dx))}{(i\tan(c+dx)a+a)^2} dx}{6a^2} + \frac{(A+iB)\tan^{m+1}(c+dx)}{6d(a+ia\tan(c+dx))^3}$$

↓ 4079

$$\frac{\int -\frac{\tan^m(c+dx)(a^2(iB(-2m^2+3m+5)-A(2m^2-9m+13))-a^2(B(1-2m)-iA(7-2m))(1-m)\tan(c+dx))}{i\tan(c+dx)a+a}}{4a^2} dx}{6a^2} + \frac{a(A(7-2m)+iB(1-2m))\tan^{m+1}(c+dx)}{4d(a+ia\tan(c+dx))^2}$$

$$\frac{(A+iB)\tan^{m+1}(c+dx)}{6d(a+ia\tan(c+dx))^3}$$

↓ 25

$$\frac{a(A(7-2m)+iB(1-2m))\tan^{m+1}(c+dx)}{4d(a+ia\tan(c+dx))^2} - \frac{\int \frac{\tan^m(c+dx)(a^2(iB(-2m^2+3m+5)-A(2m^2-9m+13))-a^2(B(1-2m)-iA(7-2m))(1-m)\tan(c+dx))}{i\tan(c+dx)a+a}}{4a^2} dx}{6a^2}$$

$$\frac{(A+iB)\tan^{m+1}(c+dx)}{6d(a+ia\tan(c+dx))^3}$$

↓ 3042

$$\frac{a(A(7-2m)+iB(1-2m))\tan^{m+1}(c+dx)}{4d(a+ia\tan(c+dx))^2} - \frac{\int \frac{\tan(c+dx)^m (a^2(iB(-2m^2+3m+5)-A(2m^2-9m+13))-a^2(B(1-2m)-iA(7-2m))(1-m)\tan(c+dx))}{i\tan(c+dx)a+a}}{4a^2} dx}{6a^2}$$

$$\frac{(A+iB)\tan^{m+1}(c+dx)}{6d(a+ia\tan(c+dx))^3}$$

↓ 4079

$$\frac{a(A(7-2m)+iB(1-2m))\tan^{m+1}(c+dx)}{4d(a+ia\tan(c+dx))^2} - \frac{\int 2\tan^m(c+dx)(a^3(1-m)(iB(-2m^2+m+3)-A(2m^2-7m+3))-a^3(2-m)m(2mB+B+iA(5-2m))\tan(c+dx))}{2a^2}}{4a^2}$$

$$\frac{(A+iB)\tan^{m+1}(c+dx)}{6d(a+ia\tan(c+dx))^3}$$

↓ 27

$$\frac{a(A(7-2m)+iB(1-2m))\tan^{m+1}(c+dx)}{4d(a+ia\tan(c+dx))^2} - \frac{\int \tan^m(c+dx)(a^3(1-m)(iB(-2m^2+m+3)-A(2m^2-7m+3))-a^3(2-m)m(2mB+B+iA(5-2m))\tan(c+dx))}{a^2}}{4a^2}$$

$$\frac{(A+iB)\tan^{m+1}(c+dx)}{6d(a+ia\tan(c+dx))^3}$$

↓ 3042

3.210. $\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

$$\frac{\frac{a(A(7-2m)+iB(1-2m)) \tan^{m+1}(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{\int \tan(c+dx)^m (a^3(1-m)(iB(-2m^2+m+3)-A(2m^2-7m+3)) - a^3(2-m)m(2mB+B+iA(5-2m)) \tan(c+dx))}{a^2}}{4a^2} = \frac{(A+iB) \tan^{m+1}(c+dx)}{6d(a+ia \tan(c+dx))^3}$$

↓ 4021

$$\frac{\frac{a(A(7-2m)+iB(1-2m)) \tan^{m+1}(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{a^3(1-m)(-A(2m^2-7m+3)+iB(-2m^2+m+3)) \int \tan^m(c+dx) dx - a^3(2-m)m(iA(5-2m)+2Bm+B) \int \tan^{m+1}(c+dx)}{a^2}}{4a^2} = \frac{(A+iB) \tan^{m+1}(c+dx)}{6d(a+ia \tan(c+dx))^3}$$

↓ 3042

$$\frac{\frac{a(A(7-2m)+iB(1-2m)) \tan^{m+1}(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{a^3(1-m)(-A(2m^2-7m+3)+iB(-2m^2+m+3)) \int \tan(c+dx)^m dx - a^3(2-m)m(iA(5-2m)+2Bm+B) \int \tan(c+dx)}{a^2}}{4a^2} = \frac{(A+iB) \tan^{m+1}(c+dx)}{6d(a+ia \tan(c+dx))^3}$$

↓ 3957

$$\frac{\frac{a(A(7-2m)+iB(1-2m)) \tan^{m+1}(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{a^3(1-m)(-A(2m^2-7m+3)+iB(-2m^2+m+3)) \int \frac{\tan^m(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx) - a^3(2-m)m(iA(5-2m)+2Bm+B) \int \tan(c+dx)}{a^2}}{4a^2} = \frac{(A+iB) \tan^{m+1}(c+dx)}{6d(a+ia \tan(c+dx))^3}$$

↓ 278

$$\frac{\frac{a(A(7-2m)+iB(1-2m)) \tan^{m+1}(c+dx)}{4d(a+ia \tan(c+dx))^2} - \frac{a^3(1-m)(-A(2m^2-7m+3)+iB(-2m^2+m+3)) \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+1}{2}, \frac{m+3}{2}, -\tan^2(c+dx)\right)}{d(m+1)}}{a^2} = \frac{(A+iB) \tan^{m+1}(c+dx)}{6d(a+ia \tan(c+dx))^3}$$

input `Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

3.210. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

```
output ((A + I*B)*Tan[c + d*x]^(1 + m))/(6*d*(a + I*a*Tan[c + d*x])^3) + ((a*(I*B
*(1 - 2*m) + A*(7 - 2*m))*Tan[c + d*x]^(1 + m))/(4*d*(a + I*a*Tan[c + d*x]
)^2) - (-(a^2*(2 - m)*(A*(5 - 2*m) - I*(B + 2*B*m))*Tan[c + d*x]^(1 + m))
/(d*(a + I*a*Tan[c + d*x]))) + ((a^3*(1 - m)*(I*B*(3 + m - 2*m^2) - A*(3 -
7*m + 2*m^2))*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2]
*Tan[c + d*x]^(1 + m))/(d*(1 + m)) - (a^3*(2 - m)*m*(B + I*A*(5 - 2*m) + 2
*B*m)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c +
d*x]^(2 + m))/(d*(2 + m))/a^2/(4*a^2)/(6*a^2)
```

3.210.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 278 Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((
c*x)^(m + 1)/(c*(m + 1))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (
-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0
] || GtQ[a, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3957 Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int
[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] &&
!IntegerQ[n]
```

```
rule 4021 Int[((b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x
_)]), x_Symbol] := Simp[c Int[(b*Tan[e + f*x])^m, x], x] + Simp[d/b Int
[(b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x] && NeQ[c^
2 + d^2, 0] && !IntegerQ[2*m]
```

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.210.4 Maple [F]

$$\int \frac{(\tan^m(dx + c))(A + B \tan(dx + c))}{(a + ia \tan(dx + c))^3} dx$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x)`

output `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x)`

3.210.5 Fracas [F]

$$\int \frac{\tan^m(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^3} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)^m}{(ia \tan(dx + c) + a)^3} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="fracas")`

output `integral(1/8*((A - I*B)*e^(6*I*d*x + 6*I*c) + (3*A - I*B)*e^(4*I*d*x + 4*I*c) + (3*A + I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m*e^(-6*I*d*x - 6*I*c)/a^3, x)`

3.210.6 Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$= \frac{i\left(\int \frac{A\tan^m(c+dx)}{\tan^3(c+dx)-3i\tan^2(c+dx)-3\tan(c+dx)+i} dx + \int \frac{B\tan(c+dx)\tan^m(c+dx)}{\tan^3(c+dx)-3i\tan^2(c+dx)-3\tan(c+dx)+i} dx\right)}{a^3}$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)`

output `I*(Integral(A*tan(c + d*x)**m/(tan(c + d*x)**3 - 3*I*tan(c + d*x)**2 - 3*tan(c + d*x) + I), x) + Integral(B*tan(c + d*x)*tan(c + d*x)**m/(tan(c + d*x)**3 - 3*I*tan(c + d*x)**2 - 3*tan(c + d*x) + I), x))/a**3`

3.210.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.210.8 Giac [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{(ia\tan(dx+c)+a)^3} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(I*a*tan(d*x + c) + a)^3, x)`

3.210. $\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

3.210.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \int \frac{\tan(c+dx)^m (A+B \tan(c+dx))}{(a+a \tan(c+dx) \text{li})^3} dx$$

input `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3,x)`

output `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3, x)`

3.211 $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

3.211.1 Optimal result 2131
 3.211.2 Mathematica [A] (verified) 2132
 3.211.3 Rubi [A] (verified) 2132
 3.211.4 Maple [F] 2137
 3.211.5 Fricas [F] 2137
 3.211.6 Sympy [F] 2138
 3.211.7 Maxima [F(-2)] 2138
 3.211.8 Giac [F] 2138
 3.211.9 Mupad [F(-1)] 2139

3.211.1 Optimal result

Integrand size = 34, antiderivative size = 386

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx =$$

$$\frac{(3-4m+m^2)(iB(1-m^2)-A(1-4m+m^2)) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right)}{48a^4d(1+m)}$$

$$- \frac{(iB(1+3m-m^2)-A(13-7m+m^2)) \tan^{1+m}(c+dx)}{48a^4d(1+i \tan(c+dx))^2}$$

$$- \frac{(2-m)(iB(2+2m-m^2)-A(8-6m+m^2)) \tan^{1+m}(c+dx)}{48a^4d(1+i \tan(c+dx))}$$

$$+ \frac{(2-m)m(B(2+2m-m^2)+iA(8-6m+m^2)) \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, -\tan^2(c+dx)\right)}{48a^4d(2+m)}$$

$$+ \frac{(A+iB) \tan^{1+m}(c+dx)}{8d(a+ia \tan(c+dx))^4} + \frac{(iB(1-m)+A(5-m)) \tan^{1+m}(c+dx)}{24ad(a+ia \tan(c+dx))^3}$$

output

```
-1/48*(m^2-4*m+3)*(I*B*(-m^2+1)-A*(m^2-4*m+1))*hypergeom([1, 1/2+1/2*m], [3/2+1/2*m], -tan(d*x+c)^2)*tan(d*x+c)^(1+m)/a^4/d/(1+m)-1/48*(I*B*(-m^2+3*m+1)-A*(m^2-7*m+13))*tan(d*x+c)^(1+m)/a^4/d/(1+I*tan(d*x+c))^2-1/48*(2-m)*(I*B*(-m^2+2*m+2)-A*(m^2-6*m+8))*tan(d*x+c)^(1+m)/a^4/d/(1+I*tan(d*x+c))+1/48*(2-m)*m*(B*(-m^2+2*m+2)+I*A*(m^2-6*m+8))*hypergeom([1, 1+1/2*m], [2+1/2*m], -tan(d*x+c)^2)*tan(d*x+c)^(2+m)/a^4/d/(2+m)+1/8*(A+I*B)*tan(d*x+c)^(1+m)/d/(a+I*a*tan(d*x+c))^4+1/24*(I*B*(1-m)+A*(5-m))*tan(d*x+c)^(1+m)/a/d/(a+I*a*tan(d*x+c))^3
```

3.211.2 Mathematica [A] (verified)

Time = 3.56 (sec) , antiderivative size = 277, normalized size of antiderivative = 0.72

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$$

$$= \frac{\tan^{1+m}(c+dx) \left(\frac{(3-4m+m^2)(iB(-1+m^2)+A(1-4m+m^2)) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right)}{1+m} + \frac{(-2+m)m(-iA(8-6m+m^2)+B(-2-2m+m^2)) \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, -\tan^2(c+dx)\right)}{2+m} + \frac{6(A+I*B)}{(-I+\tan(c+dx))^4} + \frac{2((-I)*A*(-5+m)+B*(-1+m))}{(-I+\tan(c+dx))^3} - \frac{A*(13-7m+m^2)+I*B*(-1-3m+m^2)}{(-I+\tan(c+dx))^2} + \frac{((-2+m)*(B*(2+2m-m^2)+I*A*(8-6m+m^2))}{(-I+\tan(c+dx))} \right)}{48*a^4*d}$$

input `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]`

output `(Tan[c + d*x]^(1 + m)*(((3 - 4*m + m^2)*(I*B*(-1 + m^2) + A*(1 - 4*m + m^2)))*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2])/(1 + m) + ((-2 + m)*m*((-I)*A*(8 - 6*m + m^2) + B*(-2 - 2*m + m^2))*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x])/(2 + m) + (6*(A + I*B))/(-I + Tan[c + d*x])^4 + (2*((-I)*A*(-5 + m) + B*(-1 + m)))/(-I + Tan[c + d*x])^3 - (A*(13 - 7*m + m^2) + I*B*(-1 - 3*m + m^2))/(-I + Tan[c + d*x])^2 + ((-2 + m)*(B*(2 + 2*m - m^2) + I*A*(8 - 6*m + m^2)))/(-I + Tan[c + d*x]))/(48*a^4*d)`

3.211.3 Rubi [A] (verified)

Time = 2.09 (sec) , antiderivative size = 405, normalized size of antiderivative = 1.05, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.471$, Rules used = {3042, 4079, 3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4021, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)^m(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$$

$$\downarrow \text{4079}$$

$$\begin{aligned}
 & \frac{\int \frac{\tan^m(c+dx)(a(A(7-m)-iB(m+1))-a(iA-B)(3-m)\tan(c+dx))}{(i\tan(c+dx)a+a)^3} dx}{8a^2} + \frac{(A+iB)\tan^{m+1}(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{\tan(c+dx)^m(a(A(7-m)-iB(m+1))-a(iA-B)(3-m)\tan(c+dx))}{(i\tan(c+dx)a+a)^3} dx}{8a^2} + \frac{(A+iB)\tan^{m+1}(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int -\frac{2\tan^m(c+dx)(a^2(iB(-m^2+3m+4))-A(m^2-7m+16))-a^2(B(1-m)-iA(5-m))(2-m)\tan(c+dx)}{6a^2} dx}{8a^2} + \frac{a(A(5-m)+iB(1-m))\tan^{m+1}(c+dx)}{3d(a+ia\tan(c+dx))^3} + \\
 & \quad \frac{(A+iB)\tan^{m+1}(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
 & \quad \downarrow \text{27} \\
 & \frac{a(A(5-m)+iB(1-m))\tan^{m+1}(c+dx)}{3d(a+ia\tan(c+dx))^3} - \frac{\int \frac{\tan^m(c+dx)(a^2(iB(-m^2+3m+4))-A(m^2-7m+16))-a^2(B(1-m)-iA(5-m))(2-m)\tan(c+dx)}{(i\tan(c+dx)a+a)^2} dx}{3a^2} + \\
 & \quad \frac{(A+iB)\tan^{m+1}(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a(A(5-m)+iB(1-m))\tan^{m+1}(c+dx)}{3d(a+ia\tan(c+dx))^3} - \frac{\int \frac{\tan(c+dx)^m(a^2(iB(-m^2+3m+4))-A(m^2-7m+16))-a^2(B(1-m)-iA(5-m))(2-m)\tan(c+dx)}{(i\tan(c+dx)a+a)^2} dx}{3a^2} + \\
 & \quad \frac{(A+iB)\tan^{m+1}(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
 & \quad \downarrow \text{4079} \\
 & \frac{a(A(5-m)+iB(1-m))\tan^{m+1}(c+dx)}{3d(a+ia\tan(c+dx))^3} - \frac{\int -\frac{2\tan^m(c+dx)(a^3(A(-m^3+8m^2-20m+19))-iB(m^3-4m^2+2m+7))-a^3(1-m)(B(-m^2+3m+1)+iA(m^2-}}{4a^2} dx}{8a^2} + \\
 & \quad \frac{(A+iB)\tan^{m+1}(c+dx)}{8d(a+ia\tan(c+dx))^4} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.211. $\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^4} dx$

$$\frac{a(A(5-m)+iB(1-m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^3} - \frac{(-A(m^2-7m+13)+iB(-m^2+3m+1)) \tan^{m+1}(c+dx)}{2d(1+i \tan(c+dx))^2} - \frac{\int \frac{\tan^m(c+dx) (a^3(A(-m^3+8m^2-20m+19)-iB(m^3-3m^2+2m-1)))}{(a+ia \tan(c+dx))^4} dx}{8a^2} - \frac{\int \frac{\tan^m(c+dx) (a^3(A(-m^3+8m^2-20m+19)-iB(m^3-3m^2+2m-1)))}{(a+ia \tan(c+dx))^4} dx}{3a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 3042

$$\frac{a(A(5-m)+iB(1-m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^3} - \frac{(-A(m^2-7m+13)+iB(-m^2+3m+1)) \tan^{m+1}(c+dx)}{2d(1+i \tan(c+dx))^2} - \frac{\int \frac{\tan(c+dx)^m (a^3(A(-m^3+8m^2-20m+19)-iB(m^3-3m^2+2m-1)))}{(a+ia \tan(c+dx))^4} dx}{8a^2} - \frac{\int \frac{\tan(c+dx)^m (a^3(A(-m^3+8m^2-20m+19)-iB(m^3-3m^2+2m-1)))}{(a+ia \tan(c+dx))^4} dx}{3a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 4079

$$\frac{a(A(5-m)+iB(1-m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^3} - \frac{(-A(m^2-7m+13)+iB(-m^2+3m+1)) \tan^{m+1}(c+dx)}{2d(1+i \tan(c+dx))^2} - \frac{\int -2 \tan^m(c+dx) (a^4(m^2-4m+3)(iB(1-m^2)-A(m^2-3m+1)))}{(a+ia \tan(c+dx))^4} dx}{8a^2} - \frac{\int -2 \tan^m(c+dx) (a^4(m^2-4m+3)(iB(1-m^2)-A(m^2-3m+1)))}{(a+ia \tan(c+dx))^4} dx}{3a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 27

$$\frac{a(A(5-m)+iB(1-m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^3} - \frac{(-A(m^2-7m+13)+iB(-m^2+3m+1)) \tan^{m+1}(c+dx)}{2d(1+i \tan(c+dx))^2} - \frac{\int \tan^m(c+dx) (a^4(m^2-4m+3)(iB(1-m^2)-A(m^2-3m+1)))}{(a+ia \tan(c+dx))^4} dx}{8a^2} - \frac{\int \tan^m(c+dx) (a^4(m^2-4m+3)(iB(1-m^2)-A(m^2-3m+1)))}{(a+ia \tan(c+dx))^4} dx}{3a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 3042

$$\frac{a(A(5-m)+iB(1-m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^3} - \frac{(-A(m^2-7m+13)+iB(-m^2+3m+1)) \tan^{m+1}(c+dx)}{2d(1+i \tan(c+dx))^2} - \frac{\int \tan(c+dx)^m (a^4(m^2-4m+3)(iB(1-m^2)-A(m^2-3m+1)))}{(a+ia \tan(c+dx))^4} dx}{8a^2} - \frac{\int \tan(c+dx)^m (a^4(m^2-4m+3)(iB(1-m^2)-A(m^2-3m+1)))}{(a+ia \tan(c+dx))^4} dx}{3a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 4021

3.211. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

$$\frac{a(A(5-m)+iB(1-m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^3} - \frac{(-A(m^2-7m+13)+iB(-m^2+3m+1)) \tan^{m+1}(c+dx)}{2d(1+i \tan(c+dx))^2} - \frac{a^4(m^2-4m+3)(-A(m^2-4m+1)+iB(1-m^2)) \tan^{m+1}(c+dx)}{d(a+ia \tan(c+dx))^4}$$

$8a^2$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 3042

$$\frac{a(A(5-m)+iB(1-m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^3} - \frac{(-A(m^2-7m+13)+iB(-m^2+3m+1)) \tan^{m+1}(c+dx)}{2d(1+i \tan(c+dx))^2} - \frac{a^4(m^2-4m+3)(-A(m^2-4m+1)+iB(1-m^2)) \tan^{m+1}(c+dx)}{d(a+ia \tan(c+dx))^4}$$

$8a^2$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 3957

$$\frac{a(A(5-m)+iB(1-m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^3} - \frac{(-A(m^2-7m+13)+iB(-m^2+3m+1)) \tan^{m+1}(c+dx)}{2d(1+i \tan(c+dx))^2} - \frac{a^4(m^2-4m+3)(-A(m^2-4m+1)+iB(1-m^2)) \tan^{m+1}(c+dx)}{d(a+ia \tan(c+dx))^4}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

↓ 278

$$\frac{a(A(5-m)+iB(1-m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^3} - \frac{(-A(m^2-7m+13)+iB(-m^2+3m+1)) \tan^{m+1}(c+dx)}{2d(1+i \tan(c+dx))^2} - \frac{a^3(2-m)(-A(m^2-6m+8)+iB(-m^2+2m+2)) \tan^{m+1}(c+dx)}{d(a+ia \tan(c+dx))^4}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{8d(a+ia \tan(c+dx))^4}$$

input `Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^4,x]`

```
output ((A + I*B)*Tan[c + d*x]^(1 + m))/(8*d*(a + I*a*Tan[c + d*x])^4) + ((a*(I*B
*(1 - m) + A*(5 - m))*Tan[c + d*x]^(1 + m))/(3*d*(a + I*a*Tan[c + d*x])^3)
- (((I*B*(1 + 3*m - m^2) - A*(13 - 7*m + m^2))*Tan[c + d*x]^(1 + m))/(2*d
*(1 + I*Tan[c + d*x])^2) - (-(a^3*(2 - m)*(I*B*(2 + 2*m - m^2) - A*(8 - 6
*m + m^2))*Tan[c + d*x]^(1 + m))/(d*(a + I*a*Tan[c + d*x]))) - ((a^4*(3 -
4*m + m^2)*(I*B*(1 - m^2) - A*(1 - 4*m + m^2))*Hypergeometric2F1[1, (1 + m
)/2, (3 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(1 + m))/(d*(1 + m)) - (a^4*
(2 - m)*m*(B*(2 + 2*m - m^2) + I*A*(8 - 6*m + m^2))*Hypergeometric2F1[1, (
2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(2 + m))/(d*(2 + m))/a
^2)/(2*a^2))/(3*a^2))/(8*a^2)
```

3.211.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 278 Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((
c*x)^(m + 1)/(c*(m + 1))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (
-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0
] || GtQ[a, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3957 Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int
[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] &&
!IntegerQ[n]
```

```
rule 4021 Int[((b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x
_)]), x_Symbol] := Simp[c Int[(b*Tan[e + f*x])^m, x], x] + Simp[d/b Int
[(b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x] && NeQ[c^
2 + d^2, 0] && !IntegerQ[2*m]
```

rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.211.4 Maple [F]

$$\int \frac{(\tan^m(dx + c))(A + B \tan(dx + c))}{(a + ia \tan(dx + c))^4} dx$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x)`

output `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x)`

3.211.5 Fricas [F]

$$\int \frac{\tan^m(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^4} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)^m}{(ia \tan(dx + c) + a)^4} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="fricas")`

output `integral(1/16*((A - I*B)*e^(8*I*d*x + 8*I*c) + 2*(2*A - I*B)*e^(6*I*d*x + 6*I*c) + 6*A*e^(4*I*d*x + 4*I*c) + 2*(2*A + I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m*e^(-8*I*d*x - 8*I*c)/a^4, x)`

3.211.6 Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$$

$$= \frac{\int \frac{A \tan^m(c+dx)}{\tan^4(c+dx)-4i \tan^3(c+dx)-6 \tan^2(c+dx)+4i \tan(c+dx)+1} dx + \int \frac{B \tan(c+dx) \tan^m(c+dx)}{\tan^4(c+dx)-4i \tan^3(c+dx)-6 \tan^2(c+dx)+4i \tan(c+dx)+1} dx}{a^4}$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**4,x)`

output `(Integral(A*tan(c + d*x)**m/(tan(c + d*x)**4 - 4*I*tan(c + d*x)**3 - 6*tan(c + d*x)**2 + 4*I*tan(c + d*x) + 1), x) + Integral(B*tan(c + d*x)*tan(c + d*x)**m/(tan(c + d*x)**4 - 4*I*tan(c + d*x)**3 - 6*tan(c + d*x)**2 + 4*I*tan(c + d*x) + 1), x))/a**4`

3.211.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.211.8 Giac [F]

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \int \frac{(B \tan(dx+c) + A) \tan(dx+c)^m}{(ia \tan(dx+c) + a)^4} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^4,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(I*a*tan(d*x + c) + a)^4, x)`

3.211. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx$

3.211.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^4} dx = \int \frac{\tan(c+dx)^m (A+B \tan(c+dx))}{(a+a \tan(c+dx) \text{li})^4} dx$$

input `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^4,x)`

output `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^4, x)`

3.212 $\int \tan^m(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.212.1 Optimal result	2140
3.212.2 Mathematica [F]	2141
3.212.3 Rubi [A] (verified)	2141
3.212.4 Maple [F]	2146
3.212.5 Fricas [F]	2146
3.212.6 Sympy [F(-1)]	2147
3.212.7 Maxima [F]	2147
3.212.8 Giac [F]	2147
3.212.9 Mupad [F(-1)]	2148

3.212.1 Optimal result

Integrand size = 36, antiderivative size = 316

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{4a^3(A - iB) \operatorname{AppellF1}\left(1 + m, \frac{1}{2}, 1, 2 + m, -i \tan(c + dx), i \tan(c + dx)\right) \sqrt{1 + i \tan(c + dx)}}{d(1 + m)\sqrt{a + ia \tan(c + dx)}} + \frac{2a^2(2B(19 + 17m + 4m^2) + iA(35 + 34m + 8m^2)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -m, \frac{3}{2}, 1 + i \tan(c + dx)\right) (-i \tan(c + dx))}{d(3 + 2m)(5 + 2m)} + \frac{2a^2(2iB(4 + m) - A(5 + 2m)) \tan^{1+m}(c + dx) \sqrt{a + ia \tan(c + dx)}}{d(3 + 2m)(5 + 2m)} + \frac{2iaB \tan^{1+m}(c + dx)(a + ia \tan(c + dx))^{3/2}}{d(5 + 2m)}$$

```
output 2*a^2*(2*B*(4*m^2+17*m+19)+I*A*(8*m^2+34*m+35))*hypergeom([1/2, -m], [3/2],
1+I*tan(d*x+c))*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^m/d/(3+2*m)/(5+2*m)/((
-I*tan(d*x+c))^m)+4*a^3*(A-I*B)*AppellF1(1+m,1/2,1,2+m,-I*tan(d*x+c),I*tan
(d*x+c))*(1+I*tan(d*x+c))^(1/2)*tan(d*x+c)^(1+m)/d/(1+m)/(a+I*a*tan(d*x+c)
)^(1/2)+2*a^2*(2*I*B*(4+m)-A*(5+2*m))*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^(
1+m)/d/(3+2*m)/(5+2*m)+2*I*a*B*tan(d*x+c)^(1+m)*(a+I*a*tan(d*x+c))^(3/2)/
d/(5+2*m)
```

3.212.2 Mathematica [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int \tan^m(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

input `Integrate[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Integrate[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]), x]`

3.212.3 Rubi [A] (verified)

Time = 1.68 (sec) , antiderivative size = 326, normalized size of antiderivative = 1.03, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.472$, Rules used = {3042, 4077, 27, 3042, 4077, 27, 3042, 4084, 3042, 4047, 25, 27, 152, 150, 4082, 77, 75}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(c + dx))^{5/2} \tan^m(c + dx)(A + B \tan(c + dx)) dx$$

↓ 3042

$$\int (a + ia \tan(c + dx))^{5/2} \tan(c + dx)^m (A + B \tan(c + dx)) dx$$

↓ 4077

$$2 \int -\frac{1}{2} \tan^m(c + dx)(i \tan(c + dx)a + a)^{3/2}(a(2iB(m + 1) - A(2m + 5)) - a(2B(m + 4) + iA(2m + 5)) \tan(c + dx)) dx$$

$$\frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)}$$

↓ 27

$$\frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)}$$

$$\int \tan^m(c + dx)(i \tan(c + dx)a + a)^{3/2}(a(2iB(m + 1) - A(2m + 5)) - a(2B(m + 4) + iA(2m + 5)) \tan(c + dx)) dx$$

3.212. $\int \tan^m(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)} - \\ & \frac{\int \tan(c + dx)^m (i \tan(c + dx)a + a)^{3/2} (a(2iB(m + 1) - A(2m + 5)) - a(2B(m + 4) + iA(2m + 5)) \tan(c + dx)) dx}{2m + 5} \end{aligned}$$

$$\begin{aligned} & \downarrow 4077 \\ & \frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)} - \\ & \frac{2 \int \frac{1}{2} \tan^m(c + dx) \sqrt{i \tan(c + dx)a + a} (a^2(2iB(4m^2 + 15m + 11) - A(8m^2 + 30m + 25)) - a^2(2B(4m^2 + 17m + 19) + iA(8m^2 + 34m + 35)) \tan(c + dx)) dx}{2m + 3}}{2m + 5} \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)} - \\ & \frac{\int \tan^m(c + dx) \sqrt{i \tan(c + dx)a + a} (a^2(2iB(4m^2 + 15m + 11) - A(8m^2 + 30m + 25)) - a^2(2B(4m^2 + 17m + 19) + iA(8m^2 + 34m + 35)) \tan(c + dx)) dx}{2m + 3}}{2m + 5} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)} - \\ & \frac{\int \tan(c + dx)^m \sqrt{i \tan(c + dx)a + a} (a^2(2iB(4m^2 + 15m + 11) - A(8m^2 + 30m + 25)) - a^2(2B(4m^2 + 17m + 19) + iA(8m^2 + 34m + 35)) \tan(c + dx)) dx}{2m + 3}}{2m + 5} \end{aligned}$$

$$\begin{aligned} & \downarrow 4084 \\ & \frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)} - \\ & \frac{-4a^2(4m^2 + 16m + 15)(A - iB) \int \tan^m(c + dx) \sqrt{i \tan(c + dx)a + a} dx - a(-A(8m^2 + 34m + 35) + 2iB(4m^2 + 17m + 19)) \int \tan^m(c + dx)(a - ia \tan(c + dx)) dx}{2m + 3}}{2m + 5} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)} - \\ & \frac{-4a^2(4m^2 + 16m + 15)(A - iB) \int \tan(c + dx)^m \sqrt{i \tan(c + dx)a + a} dx - a(-A(8m^2 + 34m + 35) + 2iB(4m^2 + 17m + 19)) \int \tan(c + dx)^m (a - ia \tan(c + dx)) dx}{2m + 3}}{2m + 5} \end{aligned}$$

$$\begin{aligned} & \downarrow 4047 \end{aligned}$$

3.212. $\int \tan^m(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)} - \frac{4ia^4(4m^2 + 16m + 15)(A - iB) \int \frac{\tan^m(c + dx)}{a(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a} d(ia \tan(c + dx))}{d} - a(-A(8m^2 + 34m + 35) + 2iB(4m^2 + 17m + 19)) \int \tan(c + dx)^m}{2m + 3}$$

$2m + 5$

↓ 25

$$\frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)} - \frac{4ia^4(4m^2 + 16m + 15)(A - iB) \int \frac{\tan^m(c + dx)}{a(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a} d(ia \tan(c + dx))}{d} - a(-A(8m^2 + 34m + 35) + 2iB(4m^2 + 17m + 19)) \int \tan(c + dx)^m (a - ia \tan(c + dx))}{2m + 3}$$

$2m + 5$

↓ 27

$$\frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)} - \frac{4ia^3(4m^2 + 16m + 15)(A - iB) \int \frac{\tan^m(c + dx)}{(a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a} d(ia \tan(c + dx))}{d} - a(-A(8m^2 + 34m + 35) + 2iB(4m^2 + 17m + 19)) \int \tan(c + dx)^m (a - ia \tan(c + dx))}{2m + 3}$$

$2m + 5$

↓ 152

$$\frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)} - \frac{4ia^3(4m^2 + 16m + 15)(A - iB) \sqrt{1 + i \tan(c + dx)} \int \frac{\tan^m(c + dx)}{d\sqrt{a + ia \tan(c + dx)} \int \frac{\tan^m(c + dx)}{\sqrt{i \tan(c + dx) + 1(a - ia \tan(c + dx))} d(ia \tan(c + dx))} - a(-A(8m^2 + 34m + 35) + 2iB(4m^2 + 17m + 19)) \int \tan(c + dx)^m}{2m + 3}$$

$2m + 5$

↓ 150

$$\frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)} - \frac{-a(-A(8m^2 + 34m + 35) + 2iB(4m^2 + 17m + 19)) \int \tan(c + dx)^m (a - ia \tan(c + dx))\sqrt{i \tan(c + dx)a + a} dx - \frac{4a^3(4m^2 + 16m + 15)(A - iB) \sqrt{1 + i \tan(c + dx)}}{d}}{2m + 3}$$

$2m + 5$

↓ 4082

$$\frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)} - \frac{a^3(-A(8m^2 + 34m + 35) + 2iB(4m^2 + 17m + 19)) \int \frac{\tan^m(c + dx)}{d} d \tan(c + dx) - \frac{4a^3(4m^2 + 16m + 15)(A - iB) \sqrt{1 + i \tan(c + dx)} \tan^{m+1}(c + dx) \operatorname{AppellF1}(m + 1, m + 1, 1, 1, -i \tan(c + dx))}{d(m + 1)\sqrt{a + ia \tan(c + dx)}}}{2m + 3}$$

$2m + 5$

↓ 77

3.212. $\int \tan^m(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)} - \frac{a^3(-A(8m^2 + 34m + 35) + 2iB(4m^2 + 17m + 19)) \tan^m(c + dx) (-i \tan(c + dx))^{-m} \int \frac{(-i \tan(c + dx))^m d \tan(c + dx)}{\sqrt{i \tan(c + dx) a + a}}}{d} - \frac{4a^3(4m^2 + 16m + 15)(A - iB) \sqrt{1 + i \tan(c + dx)}}{2m + 3}$$

75

$$\frac{2iaB(a + ia \tan(c + dx))^{3/2} \tan^{m+1}(c + dx)}{d(2m + 5)} - \frac{2ia^2(-A(8m^2 + 34m + 35) + 2iB(4m^2 + 17m + 19)) \sqrt{a + ia \tan(c + dx)} (-i \tan(c + dx))^{-m} \tan^m(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -m, \frac{3}{2}, i \tan(c + dx) + 1\right)}{d} - \frac{4a^3(4m^2 + 16m + 15)(A - iB) \sqrt{1 + i \tan(c + dx)}}{2m + 3}$$

input `Int[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `((2*I)*a*B*Tan[c + d*x]^(1 + m)*(a + I*a*Tan[c + d*x])^(3/2))/(d*(5 + 2*m)) - ((-2*a^2*((2*I)*B*(4 + m) - A*(5 + 2*m))*Tan[c + d*x]^(1 + m)*Sqrt[a + I*a*Tan[c + d*x]])/(d*(3 + 2*m)) + ((-4*a^3*(A - I*B)*(15 + 16*m + 4*m^2)*AppellF1[1 + m, 1/2, 1, 2 + m, (-I)*Tan[c + d*x], I*Tan[c + d*x]]*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(1 + m))/(d*(1 + m)*Sqrt[a + I*a*Tan[c + d*x]]) + ((2*I)*a^2*((2*I)*B*(19 + 17*m + 4*m^2) - A*(35 + 34*m + 8*m^2))*Hypergeometric2F1[1/2, -m, 3/2, 1 + I*Tan[c + d*x]]*Tan[c + d*x]^m*Sqrt[a + I*a*Tan[c + d*x]])/(d*((-I)*Tan[c + d*x]^m))/(3 + 2*m))/(5 + 2*m)`

3.212.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 75 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((c + d*x)^(n + 1)/(d*(n + 1)*(-d/(b*c))^m))*Hypergeometric2F1[-m, n + 1, n + 2, 1 + d*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[n] && (IntegerQ[m] || GtQ[-d/(b*c), 0])`

- rule 77 `Int[((b_.)*(x_)^(m_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[((-b)*(c/d))^IntPart[m]*((b*x)^FracPart[m]/((-d)*(x/c))^FracPart[m]) Int[((-d)*(x/c))^m*(c + d*x)^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0]`
- rule 150 `Int[((b_.)*(x_)^(m_)*((c_) + (d_.)*(x_)^(n_))*((e_) + (f_.)*(x_)^(p_)), x_] := Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])`
- rule 152 `Int[((b_.)*(x_)^(m_)*((c_) + (d_.)*(x_)^(n_))*((e_) + (f_.)*(x_)^(p_)), x_] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4047 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4077 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.212.4 Maple [F]

$$\int (\tan^m(dx + c))(a + ia \tan(dx + c))^{\frac{5}{2}} (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x)`

3.212.5 Fracas [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{\frac{5}{2}} \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output `integral(4*sqrt(2)*((A - I*B)*a^2*e^(7*I*d*x + 7*I*c) + (A + I*B)*a^2*e^(5*I*d*x + 5*I*c))*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))/(e^(6*I*d*x + 6*I*c) + 3*e^(4*I*d*x + 4*I*c) + 3*e^(2*I*d*x + 2*I*c) + 1), x)`

3.212.6 Sympy [F(-1)]

Timed out.

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**m*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.212.7 Maxima [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{5/2} \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)*tan(d*x + c)^m, x)`

3.212.8 Giac [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{5/2} \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)*tan(d*x + c)^m, x)`

3.212.9 Mupad [F(-1)]

Timed out.

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int \tan(c + dx)^m (A + B \tan(c + dx)) (a + a \tan(c + dx) i)^{5/2} dx$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),x)`output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2), x)`

3.213 $\int \tan^m(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.213.1 Optimal result	2149
3.213.2 Mathematica [F]	2150
3.213.3 Rubi [A] (verified)	2150
3.213.4 Maple [F]	2154
3.213.5 Fricas [F]	2155
3.213.6 Sympy [F]	2155
3.213.7 Maxima [F]	2155
3.213.8 Giac [F]	2156
3.213.9 Mupad [F(-1)]	2156

3.213.1 Optimal result

Integrand size = 36, antiderivative size = 227

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{2a^2(A - iB) \operatorname{AppellF1}\left(1 + m, \frac{1}{2}, 1, 2 + m, -i \tan(c + dx), i \tan(c + dx)\right) \sqrt{1 + i \tan(c + dx)}}{d(1 + m)\sqrt{a + ia \tan(c + dx)}} + \frac{2a(B + (iA + B)(3 + 2m)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -m, \frac{3}{2}, 1 + i \tan(c + dx)\right) (-i \tan(c + dx))^{-m} \tan^m(c + dx)}{d(3 + 2m)} + \frac{2iaB \tan^{1+m}(c + dx) \sqrt{a + ia \tan(c + dx)}}{d(3 + 2m)}$$

output

```
2*a*(B+(I*A+B)*(3+2*m))*hypergeom([1/2, -m], [3/2], 1+I*tan(d*x+c))*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^m/d/(3+2*m)/((-I*tan(d*x+c))^m)+2*a^2*(A-I*B)*AppellF1(1+m, 1/2, 1, 2+m, -I*tan(d*x+c), I*tan(d*x+c))*(1+I*tan(d*x+c))^(1/2)*tan(d*x+c)^(1+m)/d/(1+m)/(a+I*a*tan(d*x+c))^(1/2)+2*I*a*B*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^(1+m)/d/(3+2*m)
```

3.213.2 Mathematica [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int \tan^m(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$$

input `Integrate[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `Integrate[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]), x]`

3.213.3 Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3042, 4077, 27, 3042, 4084, 3042, 4047, 25, 27, 152, 150, 4082, 77, 75}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(c + dx))^{3/2} \tan^m(c + dx)(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(c + dx))^{3/2} \tan(c + dx)^m (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4077} \\ & \frac{2 \int -\frac{1}{2} \tan^m(c + dx) \sqrt{i \tan(c + dx) a + a(a(2iB(m + 1) - A(2m + 3)) - a(B + (iA + B)(2m + 3)) \tan(c + dx))} dx}{\frac{2iaB \sqrt{a + ia \tan(c + dx)} \tan^{m+1}(c + dx)}{d(2m + 3)}} \\ & \quad \downarrow \text{27} \\ & \frac{2iaB \sqrt{a + ia \tan(c + dx)} \tan^{m+1}(c + dx)}{d(2m + 3)} - \\ & \frac{\int \tan^m(c + dx) \sqrt{i \tan(c + dx) a + a(a(2iB(m + 1) - A(2m + 3)) - a(B + (iA + B)(2m + 3)) \tan(c + dx))} dx}{2m + 3} \end{aligned}$$

3.213. $\int \tan^m(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{2iaB\sqrt{a+ia\tan(c+dx)}\tan^{m+1}(c+dx)}{d(2m+3)} - \\
& \frac{\int \tan(c+dx)^m \sqrt{i\tan(c+dx)a+a(a(2iB(m+1)-A(2m+3))-a(B+(iA+B)(2m+3))\tan(c+dx))dx}{2m+3} \\
& \downarrow 4084 \\
& \frac{2iaB\sqrt{a+ia\tan(c+dx)}\tan^{m+1}(c+dx)}{d(2m+3)} - \\
& \frac{-2a(2m+3)(A-iB)\int \tan^m(c+dx)\sqrt{i\tan(c+dx)a+adx} - i(B+(2m+3)(B+iA))\int \tan^m(c+dx)(a-ia\tan(c+dx))}{2m+3} \\
& \downarrow 3042 \\
& \frac{2iaB\sqrt{a+ia\tan(c+dx)}\tan^{m+1}(c+dx)}{d(2m+3)} - \\
& \frac{-2a(2m+3)(A-iB)\int \tan(c+dx)^m \sqrt{i\tan(c+dx)a+adx} - i(B+(2m+3)(B+iA))\int \tan(c+dx)^m(a-ia\tan(c+dx))}{2m+3} \\
& \downarrow 4047 \\
& \frac{2iaB\sqrt{a+ia\tan(c+dx)}\tan^{m+1}(c+dx)}{d(2m+3)} - \\
& \frac{\frac{2ia^3(2m+3)(A-iB)\int -\frac{\tan^m(c+dx)}{a(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}d(ia\tan(c+dx))}{d} - i(B+(2m+3)(B+iA))\int \tan(c+dx)^m(a-ia\tan(c+dx))}{2m+3}} \\
& \downarrow 25 \\
& \frac{2iaB\sqrt{a+ia\tan(c+dx)}\tan^{m+1}(c+dx)}{d(2m+3)} - \\
& \frac{2ia^3(2m+3)(A-iB)\int \frac{\tan^m(c+dx)}{a(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}d(ia\tan(c+dx))}{d} - i(B+(2m+3)(B+iA))\int \tan(c+dx)^m(a-ia\tan(c+dx))}{2m+3}} \\
& \downarrow 27 \\
& \frac{2iaB\sqrt{a+ia\tan(c+dx)}\tan^{m+1}(c+dx)}{d(2m+3)} - \\
& \frac{2ia^2(2m+3)(A-iB)\int \frac{\tan^m(c+dx)}{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}d(ia\tan(c+dx))}{d} - i(B+(2m+3)(B+iA))\int \tan(c+dx)^m(a-ia\tan(c+dx))}{2m+3}} \\
& \downarrow 152 \\
& \frac{2iaB\sqrt{a+ia\tan(c+dx)}\tan^{m+1}(c+dx)}{d(2m+3)} - \\
& \frac{2ia^2(2m+3)(A-iB)\sqrt{1+i\tan(c+dx)}\int \frac{\tan^m(c+dx)}{\sqrt{i\tan(c+dx)+1(a-ia\tan(c+dx))}d(ia\tan(c+dx))}{d\sqrt{a+ia\tan(c+dx)}} - i(B+(2m+3)(B+iA))\int \tan(c+dx)^m}{2m+3}}
\end{aligned}$$

3.213. $\int \tan^m(c+dx)(a+ia\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx$

$$\begin{aligned} & \downarrow 150 \\ & \frac{2iaB\sqrt{a+ia\tan(c+dx)}\tan^{m+1}(c+dx)}{d(2m+3)} - \\ & \frac{-i(B+(2m+3)(B+iA))\int\tan(c+dx)^m(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+adx} - \frac{2a^2(2m+3)(A-iB)\sqrt{1+i\tan(c+dx)}}{2m+3}}{2m+3} \end{aligned}$$

$$\begin{aligned} & \downarrow 4082 \\ & \frac{2iaB\sqrt{a+ia\tan(c+dx)}\tan^{m+1}(c+dx)}{d(2m+3)} - \\ & \frac{ia^2(B+(2m+3)(B+iA))\int\frac{\tan^m(c+dx)}{\sqrt{i\tan(c+dx)a+a}}d\tan(c+dx) - \frac{2a^2(2m+3)(A-iB)\sqrt{1+i\tan(c+dx)}\tan^{m+1}(c+dx)\operatorname{AppellF1}(m+1,\frac{1}{2},1,m+2,-i\tan(c+dx))}{d(m+1)\sqrt{a+ia\tan(c+dx)}}}{2m+3} \end{aligned}$$

$$\begin{aligned} & \downarrow 77 \\ & \frac{2iaB\sqrt{a+ia\tan(c+dx)}\tan^{m+1}(c+dx)}{d(2m+3)} - \\ & \frac{ia^2(B+(2m+3)(B+iA))\tan^m(c+dx)(-i\tan(c+dx))^{-m}\int\frac{(-i\tan(c+dx))^m}{\sqrt{i\tan(c+dx)a+a}}d\tan(c+dx) - \frac{2a^2(2m+3)(A-iB)\sqrt{1+i\tan(c+dx)}\tan^{m+1}(c+dx)}{d(m+1)\sqrt{a+ia\tan(c+dx)}}}{2m+3} \end{aligned}$$

$$\begin{aligned} & \downarrow 75 \\ & \frac{2iaB\sqrt{a+ia\tan(c+dx)}\tan^{m+1}(c+dx)}{d(2m+3)} - \\ & \frac{2a^2(2m+3)(A-iB)\sqrt{1+i\tan(c+dx)}\tan^{m+1}(c+dx)\operatorname{AppellF1}(m+1,\frac{1}{2},1,m+2,-i\tan(c+dx),i\tan(c+dx)) - \frac{2a(B+(2m+3)(B+iA))\sqrt{a+ia\tan(c+dx)}}{d(m+1)\sqrt{a+ia\tan(c+dx)}}}{2m+3} \end{aligned}$$

input `Int[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `((2*I)*a*B*Tan[c + d*x]^(1 + m)*Sqrt[a + I*a*Tan[c + d*x]]/(d*(3 + 2*m)) - ((-2*a^2*(A - I*B)*(3 + 2*m)*AppellF1[1 + m, 1/2, 1, 2 + m, (-I)*Tan[c + d*x], I*Tan[c + d*x]]*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(1 + m))/(d*(1 + m)*Sqrt[a + I*a*Tan[c + d*x]]) - (2*a*(B + (I*A + B)*(3 + 2*m))*Hypergeometric2F1[1/2, -m, 3/2, 1 + I*Tan[c + d*x]]*Tan[c + d*x]^m*Sqrt[a + I*a*Tan[c + d*x]])/(d*((-I)*Tan[c + d*x])^m)/(3 + 2*m)`

3.213.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 75 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((c + d*x)^(n + 1)/(d*(n + 1)*(-d/(b*c))^m)*Hypergeometric2F1[-m, n + 1, n + 2, 1 + d*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[n] && (IntegerQ[m] || GtQ[-d/(b*c), 0])`
- rule 77 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((-b)*(c/d))^IntPart[m]*((b*x)^FracPart[m]/((-d)*(x/c))^FracPart[m]) Int[((-d)*(x/c))^m*(c + d*x)^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0]`
- rule 150 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_] := Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])`
- rule 152 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4047 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4077 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.213.4 Maple [F]

$$\int (\tan^m(dx + c))(a + ia \tan(dx + c))^{\frac{3}{2}} (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x)`

3.213.5 Fracas [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{\frac{3}{2}} \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(2*sqrt(2)*((A - I*B)*a*e^(5*I*d*x + 5*I*c) + (A + I*B)*a*e^(3*I*d*x + 3*I*c))*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))/(e^(4*I*d*x + 4*I*c) + 2*e^(2*I*d*x + 2*I*c) + 1), x)`

3.213.6 Sympy [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{\frac{3}{2}}(A + B \tan(c + dx)) \tan^m(c + dx) dx$$

input `integrate(tan(d*x+c)**m*(a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(3/2)*(A + B*tan(c + d*x))*tan(c + d*x)**m, x)`

3.213.7 Maxima [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{\frac{3}{2}} \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)*tan(d*x + c)^m, x)`

3.213.8 Giac [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{\frac{3}{2}} \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)*tan(d*x + c)^m, x)`

3.213.9 Mupad [F(-1)]

Timed out.

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int \tan(c + dx)^m (A + B \tan(c + dx)) (a + a \tan(c + dx) 1i)^{3/2} dx$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2),x)`

output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2), x)`

3.214 $\int \tan^m(c+dx) \sqrt{a + ia \tan(c + dx)} (A+B \tan(c+dx)) dx$

3.214.1 Optimal result	2157
3.214.2 Mathematica [F]	2157
3.214.3 Rubi [A] (verified)	2158
3.214.4 Maple [F]	2161
3.214.5 Fracas [F]	2162
3.214.6 Sympy [F]	2162
3.214.7 Maxima [F]	2162
3.214.8 Giac [F]	2163
3.214.9 Mupad [F(-1)]	2163

3.214.1 Optimal result

Integrand size = 36, antiderivative size = 159

$$\int \tan^m(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{a(A - iB) \operatorname{AppellF1}\left(1 + m, \frac{1}{2}, 1, 2 + m, -i \tan(c + dx), i \tan(c + dx)\right) \sqrt{1 + i \tan(c + dx)} \tan^{1+m}(c + dx)}{d(1 + m) \sqrt{a + ia \tan(c + dx)}} + \frac{2B \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -m, \frac{3}{2}, 1 + i \tan(c + dx)\right) (-i \tan(c + dx))^{-m} \tan^m(c + dx) \sqrt{a + ia \tan(c + dx)}}{d}$$

```
output 2*B*hypergeom([1/2, -m], [3/2], 1+I*tan(d*x+c))*(a+I*a*tan(d*x+c))^(1/2)*tan
(d*x+c)^m/d/((-I*tan(d*x+c))^m)+a*(A-I*B)*AppellF1(1+m, 1/2, 1, 2+m, -I*tan(d*
x+c), I*tan(d*x+c))*(1+I*tan(d*x+c))^(1/2)*tan(d*x+c)^(1+m)/d/(1+m)/(a+I*a*
tan(d*x+c))^(1/2)
```

3.214.2 Mathematica [F]

$$\int \tan^m(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \tan^m(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

input `Integrate[Tan[c + d*x]^m*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `Integrate[Tan[c + d*x]^m*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

3.214.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.306$, Rules used = {3042, 4084, 3042, 4047, 25, 27, 152, 150, 4082, 77, 75}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a + ia \tan(c + dx)} \tan^m(c + dx) (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a + ia \tan(c + dx)} \tan(c + dx)^m (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4084} \\
 & \frac{(A - iB) \int \tan^m(c + dx) \sqrt{i \tan(c + dx) a + adx} +}{a} \\
 & \quad \frac{iB \int \tan^m(c + dx) (a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + adx}}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A - iB) \int \tan(c + dx)^m \sqrt{i \tan(c + dx) a + adx} +}{a} \\
 & \quad \frac{iB \int \tan(c + dx)^m (a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + adx}}{a} \\
 & \quad \downarrow \text{4047} \\
 & \frac{ia^2(A - iB) \int -\frac{\tan^m(c+dx)}{a(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx) a+a}} d(ia \tan(c + dx))}{d} + \\
 & \quad \frac{iB \int \tan(c + dx)^m (a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + adx}}{a} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

3.214. $\int \tan^m(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \frac{iB \int \tan(c+dx)^m (a - ia \tan(c+dx)) \sqrt{i \tan(c+dx)a + adx} \, dx}{ia^2(A - iB) \int \frac{\tan^m(c+dx)}{a(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a + a}} d(ia \tan(c+dx))} \\
& \quad \downarrow \text{27} \\
& \frac{iB \int \tan(c+dx)^m (a - ia \tan(c+dx)) \sqrt{i \tan(c+dx)a + adx} \, dx}{ia(A - iB) \int \frac{\tan^m(c+dx)}{(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a + a}} d(ia \tan(c+dx))} \\
& \quad \downarrow \text{152} \\
& \frac{iB \int \tan(c+dx)^m (a - ia \tan(c+dx)) \sqrt{i \tan(c+dx)a + adx} \, dx}{ia(A - iB) \sqrt{1 + i \tan(c+dx)} \int \frac{a \tan^m(c+dx)}{\sqrt{i \tan(c+dx)+1} (a-ia \tan(c+dx))} d(ia \tan(c+dx))} \\
& \quad \downarrow \text{150} \\
& \frac{iB \int \tan(c+dx)^m (a - ia \tan(c+dx)) \sqrt{i \tan(c+dx)a + adx} \, dx}{a(A - iB) \sqrt{1 + i \tan(c+dx)} \tan^{m+1}(c+dx) \operatorname{AppellF1}(m+1, \frac{1}{2}, 1, m+2, -i \tan(c+dx), i \tan(c+dx))} + \\
& \quad \frac{a}{d(m+1) \sqrt{a + ia \tan(c+dx)}} \\
& \quad \downarrow \text{4082} \\
& \frac{iaB \int \frac{\tan^m(c+dx)}{\sqrt{i \tan(c+dx)a + a}} d \tan(c+dx)}{d} + \\
& \frac{a(A - iB) \sqrt{1 + i \tan(c+dx)} \tan^{m+1}(c+dx) \operatorname{AppellF1}(m+1, \frac{1}{2}, 1, m+2, -i \tan(c+dx), i \tan(c+dx))}{d(m+1) \sqrt{a + ia \tan(c+dx)}} \\
& \quad \downarrow \text{77} \\
& \frac{iaB \tan^m(c+dx) (-i \tan(c+dx))^{-m} \int \frac{(-i \tan(c+dx))^m}{\sqrt{i \tan(c+dx)a + a}} d \tan(c+dx)}{d} + \\
& \frac{a(A - iB) \sqrt{1 + i \tan(c+dx)} \tan^{m+1}(c+dx) \operatorname{AppellF1}(m+1, \frac{1}{2}, 1, m+2, -i \tan(c+dx), i \tan(c+dx))}{d(m+1) \sqrt{a + ia \tan(c+dx)}} \\
& \quad \downarrow \text{75} \\
& \frac{a(A - iB) \sqrt{1 + i \tan(c+dx)} \tan^{m+1}(c+dx) \operatorname{AppellF1}(m+1, \frac{1}{2}, 1, m+2, -i \tan(c+dx), i \tan(c+dx))}{d(m+1) \sqrt{a + ia \tan(c+dx)}} + \\
& \frac{2B \sqrt{a + ia \tan(c+dx)} \tan^m(c+dx) (-i \tan(c+dx))^{-m} \operatorname{Hypergeometric2F1}(\frac{1}{2}, -m, \frac{3}{2}, i \tan(c+dx) + 1)}{d}
\end{aligned}$$

input `Int[Tan[c + d*x]^m*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(a*(A - I*B)*AppellF1[1 + m, 1/2, 1, 2 + m, (-I)*Tan[c + d*x], I*Tan[c + d*x]]*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(1 + m))/(d*(1 + m)*Sqrt[a + I*a*Tan[c + d*x]]) + (2*B*Hypergeometric2F1[1/2, -m, 3/2, 1 + I*Tan[c + d*x]]*Tan[c + d*x]^m*Sqrt[a + I*a*Tan[c + d*x]])/(d*((-I)*Tan[c + d*x])^m)`

3.214.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 75 `Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((c + d*x)^(n + 1)/(d*(n + 1)*(-d/(b*c))^m)*Hypergeometric2F1[-m, n + 1, n + 2, 1 + d*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[n] && (IntegerQ[m] || GtQ[-d/(b*c), 0])`

rule 77 `Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((-b)*(c/d))^IntPart[m]*((b*x)^FracPart[m]/((-d)*(x/c))^FracPart[m]) Int[((-d)*(x/c))^m*(c + d*x)^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0]`

rule 150 `Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])`

rule 152 `Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4047 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.214.4 Maple [F]

$$\int (\tan^m(dx + c)) \sqrt{a + ia \tan(dx + c)} (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x)`

3.214.5 Fricas [F]

$$\int \tan^m(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{ia \tan(dx + c) + a \tan(dx + c)}^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(sqrt(2)*((A - I*B)*e^(3*I*d*x + 3*I*c) + (A + I*B)*e^(I*d*x + I*c)))*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))/(e^(2*I*d*x + 2*I*c) + 1), x)`

3.214.6 Sympy [F]

$$\int \tan^m(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \sqrt{ia (\tan(c + dx) - i)} (A + B \tan(c + dx)) \tan^m(c + dx) dx$$

input `integrate(tan(d*x+c)**m*(a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))*tan(c + d*x)**m, x)`

3.214.7 Maxima [F]

$$\int \tan^m(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{ia \tan(dx + c) + a \tan(dx + c)}^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)*tan(d*x + c)^m, x)`

3.214.8 Giac [F]

$$\begin{aligned} & \int \tan^m(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A) \sqrt{ia \tan(dx + c) + a} \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)*tan(d*x + c)^m, x)`

3.214.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \tan^m(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= \int \tan(c + dx)^m (A + B \tan(c + dx)) \sqrt{a + a \tan(c + dx)} li dx \end{aligned}$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2),x)`

output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2), x)`

3.215 $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

3.215.1 Optimal result 2164
 3.215.2 Mathematica [F] 2164
 3.215.3 Rubi [A] (verified) 2165
 3.215.4 Maple [F] 2169
 3.215.5 Fracas [F] 2170
 3.215.6 Sympy [F] 2170
 3.215.7 Maxima [F(-2)] 2170
 3.215.8 Giac [F] 2171
 3.215.9 Mupad [F(-1)] 2171

3.215.1 Optimal result

Integrand size = 36, antiderivative size = 214

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \frac{(A+iB) \tan^{1+m}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} + \frac{(A-iB) \operatorname{AppellF1}\left(1+m, \frac{1}{2}, 1, 2+m, -i \tan(c+dx), i \tan(c+dx)\right) \sqrt{1+i \tan(c+dx)} \tan^{1+m}(c+dx)}{2d(1+m)\sqrt{a+ia \tan(c+dx)}} + \frac{(iA-B)(1+2m) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -m, \frac{3}{2}, 1+i \tan(c+dx)\right) (-i \tan(c+dx))^{-m} \tan^m(c+dx)}{ad}$$

```
output (I*A-B)*(1+2*m)*hypergeom([1/2, -m], [3/2], 1+I*tan(d*x+c))*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^m/a/d/((-I*tan(d*x+c))^m)+(A+I*B)*tan(d*x+c)^(1+m)/d/(a+I*a*tan(d*x+c))^(1/2)+1/2*(A-I*B)*AppellF1(1+m, 1/2, 1, 2+m, -I*tan(d*x+c), I*tan(d*x+c))*(1+I*tan(d*x+c))^(1/2)*tan(d*x+c)^(1+m)/d/(1+m)/(a+I*a*tan(d*x+c))^(1/2)
```

3.215.2 Mathematica [F]

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

```
input Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]], x]
```

output `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]] , x]`

3.215.3 Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3042, 4079, 27, 3042, 4084, 3042, 4047, 25, 27, 152, 150, 4082, 77, 75}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx$$

$$\downarrow 3042$$

$$\int \frac{\tan(c+dx)^m(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx$$

$$\downarrow 4079$$

$$\frac{\int -\frac{1}{2}\tan^m(c+dx)\sqrt{i\tan(c+dx)a+a(2a(Am+iB(m+1))-a(iA-B)(2m+1)\tan(c+dx))}dx}{a^2} + \frac{(A+iB)\tan^{m+1}(c+dx)}{d\sqrt{a+ia\tan(c+dx)}}$$

$$\downarrow 27$$

$$\frac{(A+iB)\tan^{m+1}(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} - \frac{\int \tan^m(c+dx)\sqrt{i\tan(c+dx)a+a(2a(Am+iB(m+1))-a(iA-B)(2m+1)\tan(c+dx))}dx}{2a^2}$$

$$\downarrow 3042$$

$$\frac{(A+iB)\tan^{m+1}(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} - \frac{\int \tan(c+dx)^m\sqrt{i\tan(c+dx)a+a(2a(Am+iB(m+1))-a(iA-B)(2m+1)\tan(c+dx))}dx}{2a^2}$$

$$\downarrow 4084$$

$$\frac{(A+iB)\tan^{m+1}(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} - \frac{(2m+1)(A+iB)\int \tan^m(c+dx)(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+adx}-a(A-iB)\int \tan^m(c+dx)\sqrt{i\tan(c+dx)a+adx}}{2a^2}$$

3.215. $\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx$

$$\begin{array}{c}
\downarrow 3042 \\
\frac{(A + iB) \tan^{m+1}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
\frac{(2m + 1)(A + iB) \int \tan(c + dx)^m (a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + adx} - a(A - iB) \int \tan(c + dx)^m \sqrt{i \tan(c + dx) a + adx}}{2a^2} \\
\downarrow 4047 \\
\frac{(A + iB) \tan^{m+1}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
\frac{(2m + 1)(A + iB) \int \tan(c + dx)^m (a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + adx} - \frac{ia^3(A - iB) \int \frac{\tan^m(c + dx)}{a(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + adx}}}{d}}{2a^2} \\
\downarrow 25 \\
\frac{(A + iB) \tan^{m+1}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
\frac{ia^3(A - iB) \int \frac{\tan^m(c + dx)}{a(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + adx}} d(ia \tan(c + dx))}{d} + \frac{(2m + 1)(A + iB) \int \tan(c + dx)^m (a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + adx}}{2a^2} \\
\downarrow 27 \\
\frac{(A + iB) \tan^{m+1}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
\frac{ia^2(A - iB) \int \frac{\tan^m(c + dx)}{(a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + adx}} d(ia \tan(c + dx))}{d} + \frac{(2m + 1)(A + iB) \int \tan(c + dx)^m (a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + adx}}{2a^2} \\
\downarrow 152 \\
\frac{(A + iB) \tan^{m+1}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
\frac{ia^2(A - iB) \sqrt{1 + i \tan(c + dx)} \int \frac{\tan^m(c + dx)}{\sqrt{i \tan(c + dx) + 1} (a - ia \tan(c + dx))} d(ia \tan(c + dx))}{d\sqrt{a + ia \tan(c + dx)}} + \frac{(2m + 1)(A + iB) \int \tan(c + dx)^m (a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + adx}}{2a^2} \\
\downarrow 150 \\
\frac{(A + iB) \tan^{m+1}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \\
\frac{(2m + 1)(A + iB) \int \tan(c + dx)^m (a - ia \tan(c + dx)) \sqrt{i \tan(c + dx) a + adx} - \frac{a^2(A - iB) \sqrt{1 + i \tan(c + dx)} \tan^{m+1}(c + dx)}{d(m+1)}}{2a^2} \\
\downarrow 4082
\end{array}$$

3.215. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

$$\frac{\frac{(A + iB) \tan^{m+1}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{a^2(2m+1)(A+iB) \int \frac{\tan^m(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{d} - \frac{a^2(A-iB)\sqrt{1+i \tan(c+dx)} \tan^{m+1}(c+dx) \operatorname{AppellF1}(m+1, \frac{1}{2}, 1, m+2, -i \tan(c+dx), i \tan(c+dx))}{d(m+1)\sqrt{a+ia \tan(c+dx)}}}{2a^2}$$

↓ 77

$$\frac{\frac{(A + iB) \tan^{m+1}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{a^2(2m+1)(A+iB)(-i \tan(c+dx))^{-m} \tan^m(c+dx) \int \frac{(-i \tan(c+dx))^m}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{d} - \frac{a^2(A-iB)\sqrt{1+i \tan(c+dx)} \tan^{m+1}(c+dx) \operatorname{AppellF1}(m+1, \frac{1}{2}, 1, m+2, -i \tan(c+dx), i \tan(c+dx))}{d(m+1)\sqrt{a+ia \tan(c+dx)}}}{2a^2}$$

↓ 75

$$\frac{\frac{(A + iB) \tan^{m+1}(c + dx)}{d\sqrt{a + ia \tan(c + dx)}} - \frac{a^2(A-iB)\sqrt{1+i \tan(c+dx)} \tan^{m+1}(c+dx) \operatorname{AppellF1}(m+1, \frac{1}{2}, 1, m+2, -i \tan(c+dx), i \tan(c+dx))}{d(m+1)\sqrt{a+ia \tan(c+dx)}} - \frac{2ia(2m+1)(A+iB)\sqrt{a+ia \tan(c+dx)} \tan^{m+1}(c+dx)}{2a^2}}{2a^2}$$

input `Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `((A + I*B)*Tan[c + d*x]^(1 + m))/(d*Sqrt[a + I*a*Tan[c + d*x]]) - (-((a^2*(A - I*B)*AppellF1[1 + m, 1/2, 1, 2 + m, (-I)*Tan[c + d*x], I*Tan[c + d*x]]*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(1 + m))/(d*(1 + m)*Sqrt[a + I*a*Tan[c + d*x]])) - ((2*I)*a*(A + I*B)*(1 + 2*m)*Hypergeometric2F1[1/2, -m, 3/2, 1 + I*Tan[c + d*x]]*Tan[c + d*x]^m*Sqrt[a + I*a*Tan[c + d*x]])/(d*((-I)*Tan[c + d*x]^m))/(2*a^2)`

3.215.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

- rule 75 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[((c + d*x)^(n + 1)/(d*(n + 1)*(-d/(b*c))^(m))*Hypergeometric2F1[-m, n + 1, n + 2, 1 + d*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[n] && (IntegerQ[m] || GtQ[-d/(b*c), 0])`
- rule 77 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[((-b)*(c/d))^IntPart[m]*((b*x)^FracPart[m]/((-d)*(x/c))^FracPart[m]) Int[((-d)*(x/c))^m*(c + d*x)^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0]`
- rule 150 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_))*((e_) + (f_.)*(x_)^(p_)), x_] := Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])`
- rule 152 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_))*((e_) + (f_.)*(x_)^(p_)), x_] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4047 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.215.4 Maple [F]

$$\int \frac{(\tan^m(dx + c))(A + B \tan(dx + c))}{\sqrt{a + ia \tan(dx + c)}} dx$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x)`

output `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x)`

3.215.5 Fricas [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{\sqrt{ia\tan(dx+c)+a}} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="fricas")`

output `integral(1/2*sqrt(2)*((A - I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*e^(-I*d*x - I*c)/a, x)`

3.215.6 Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{(A+B\tan(c+dx))\tan^m(c+dx)}{\sqrt{ia(\tan(c+dx)-i)}} dx$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**m/sqrt(I*a*(tan(c + d*x) - I)), x)`

3.215.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is not defined.`

3.215.8 Giac [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{\sqrt{ia\tan(dx+c)+a}} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/sqrt(I*a*tan(d*x + c) + a), x)`

3.215.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{\tan(c+dx)^m (A+B\tan(c+dx))}{\sqrt{a+a\tan(c+dx)} \operatorname{li}} dx$$

input `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2),x)`

output `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2), x)`

3.216
$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$$

3.216.1 Optimal result 2172
 3.216.2 Mathematica [F] 2173
 3.216.3 Rubi [A] (verified) 2173
 3.216.4 Maple [F] 2178
 3.216.5 Fricas [F] 2178
 3.216.6 Sympy [F] 2179
 3.216.7 Maxima [F(-2)] 2179
 3.216.8 Giac [F] 2179
 3.216.9 Mupad [F(-1)] 2180

3.216.1 Optimal result

Integrand size = 36, antiderivative size = 285

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{(A+iB) \tan^{1+m}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} + \frac{(A(5-4m) - i(B+4Bm)) \tan^{1+m}(c+dx)}{6ad\sqrt{a+ia \tan(c+dx)}} + \frac{(A-iB) \operatorname{AppellF1}(1+m, \frac{1}{2}, 1, 2+m, -i \tan(c+dx), i \tan(c+dx)) \sqrt{1+i \tan(c+dx)} \tan^{1+m}(c+dx)}{4ad(1+m)\sqrt{a+ia \tan(c+dx)}} + \frac{(1+2m)(B+iA(5-4m) + 4Bm) \operatorname{Hypergeometric2F1}(\frac{1}{2}, -m, \frac{3}{2}, 1+i \tan(c+dx)) (-i \tan(c+dx))^{-m}}{6a^2d}$$

output

```
1/6*(1+2*m)*(B+I*A*(5-4*m)+4*B*m)*hypergeom([1/2, -m], [3/2], 1+I*tan(d*x+c)
)*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^m/a^2/d/((-I*tan(d*x+c))^m)+1/6*(A*(
5-4*m)-I*(4*B*m+B))*tan(d*x+c)^(1+m)/a/d/(a+I*a*tan(d*x+c))^(1/2)+1/4*(A-I
*B)*AppellF1(1+m, 1/2, 1, 2+m, -I*tan(d*x+c), I*tan(d*x+c))*(1+I*tan(d*x+c))^(1
/2)*tan(d*x+c)^(1+m)/a/d/(1+m)/(a+I*a*tan(d*x+c))^(1/2)+1/3*(A+I*B)*tan(d*
x+c)^(1+m)/d/(a+I*a*tan(d*x+c))^(3/2)
```

3.216.2 Mathematica [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx$$

input `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2), x]`

output `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2), x]`

3.216.3 Rubi [A] (verified)

Time = 1.55 (sec) , antiderivative size = 296, normalized size of antiderivative = 1.04, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.472$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4084, 3042, 4047, 25, 27, 152, 150, 4082, 77, 75}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(c+dx)^m(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{4079} \\ & \frac{\int \frac{\tan^m(c+dx)(2a(A(2-m)-iB(m+1))-a(iA-B)(1-2m)\tan(c+dx))}{2\sqrt{i\tan(c+dx)a+a}} dx}{3a^2} + \frac{(A+iB)\tan^{m+1}(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{\tan^m(c+dx)(2a(A(2-m)-iB(m+1))-a(iA-B)(1-2m)\tan(c+dx))}{\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} + \frac{(A+iB)\tan^{m+1}(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{\tan(c+dx)^m(2a(A(2-m)-iB(m+1))-a(iA-B)(1-2m)\tan(c+dx))}{\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} + \frac{(A+iB)\tan^{m+1}(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} \end{aligned}$$

3.216. $\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx$

↓ 4079

$$\frac{\int -\frac{1}{2} \tan^m(c+dx) \sqrt{i \tan(c+dx)a+a} (2a^2(iB(-4m^2-3m+1)+A(-4m^2+3m+1))-a^2(2m+1)(4mB+B+iA(5-4m)) \tan(c+dx)) dx}{a^2} + \frac{a(A(5-4m)-i(4Bm+B)) \tan^{m+1}(c+dx)}{6a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 27

$$\frac{a(A(5-4m)-i(4Bm+B)) \tan^{m+1}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{\int \tan^m(c+dx) \sqrt{i \tan(c+dx)a+a} (2a^2(iB(-4m^2-3m+1)+A(-4m^2+3m+1))-a^2(2m+1)(4mB+B+iA(5-4m)) \tan(c+dx)) dx}{2a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 3042

$$\frac{a(A(5-4m)-i(4Bm+B)) \tan^{m+1}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{\int \tan(c+dx)^m \sqrt{i \tan(c+dx)a+a} (2a^2(iB(-4m^2-3m+1)+A(-4m^2+3m+1))-a^2(2m+1)(4mB+B+iA(5-4m)) \tan(c+dx)) dx}{2a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 4084

$$\frac{a(A(5-4m)-i(4Bm+B)) \tan^{m+1}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{a(2m+1)(A(5-4m)-i(4Bm+B)) \int \tan^m(c+dx)(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx} - 3a^2(A-iB) \tan^{m+1}(c+dx)}{2a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 3042

$$\frac{a(A(5-4m)-i(4Bm+B)) \tan^{m+1}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{a(2m+1)(A(5-4m)-i(4Bm+B)) \int \tan(c+dx)^m (a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx} - 3a^2(A-iB) \tan^{m+1}(c+dx)}{2a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 4047

$$\frac{a(A(5-4m)-i(4Bm+B)) \tan^{m+1}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{a(2m+1)(A(5-4m)-i(4Bm+B)) \int \tan(c+dx)^m (a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx} - \frac{3ia^4(A-iB) \tan^{m+1}(c+dx)}{2a^2}}{2a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

3.216. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

↓ 25

$$\frac{a(A(5-4m)-i(4Bm+B)) \tan^{m+1}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{3ia^4(A-iB) \int \frac{\tan^m(c+dx)}{a(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}} d(ia \tan(c+dx))}{d} + \frac{a(2m+1)(A(5-4m)-i(4Bm+B)) \int \frac{\tan^m(c+dx)}{a(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}} d(ia \tan(c+dx))}{d} + \frac{a(2m+1)(A(5-4m)-i(4Bm+B))}{2a^2} \int \frac{\tan^m(c+dx)}{a(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}} d(ia \tan(c+dx))}{d}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 27

$$\frac{a(A(5-4m)-i(4Bm+B)) \tan^{m+1}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{3ia^3(A-iB) \int \frac{\tan^m(c+dx)}{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}} d(ia \tan(c+dx))}{d} + \frac{a(2m+1)(A(5-4m)-i(4Bm+B)) \int \frac{\tan^m(c+dx)}{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}} d(ia \tan(c+dx))}{d} + \frac{a(2m+1)(A(5-4m)-i(4Bm+B))}{2a^2} \int \frac{\tan^m(c+dx)}{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}} d(ia \tan(c+dx))}{d}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 152

$$\frac{a(A(5-4m)-i(4Bm+B)) \tan^{m+1}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{3ia^3(A-iB)\sqrt{1+i \tan(c+dx)} \int \frac{\tan^m(c+dx)}{\sqrt{i \tan(c+dx)+1(a-ia \tan(c+dx))} d(ia \tan(c+dx))}{d\sqrt{a+ia \tan(c+dx)}}}{d} + \frac{a(2m+1)(A(5-4m)-i(4Bm+B)) \int \frac{\tan^m(c+dx)}{\sqrt{i \tan(c+dx)+1(a-ia \tan(c+dx))} d(ia \tan(c+dx))}{d\sqrt{a+ia \tan(c+dx)}}}{d} + \frac{a(2m+1)(A(5-4m)-i(4Bm+B))}{2a^2} \int \frac{\tan^m(c+dx)}{\sqrt{i \tan(c+dx)+1(a-ia \tan(c+dx))} d(ia \tan(c+dx))}{d\sqrt{a+ia \tan(c+dx)}}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 150

$$\frac{a(A(5-4m)-i(4Bm+B)) \tan^{m+1}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{a(2m+1)(A(5-4m)-i(4Bm+B)) \int \tan(c+dx)^m (a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx} - \frac{3a^3(A-iB)}{2a^2} \int \tan(c+dx)^m (a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{d} + \frac{a(2m+1)(A(5-4m)-i(4Bm+B)) \int \tan(c+dx)^m (a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{d} - \frac{3a^3(A-iB)}{2a^2} \int \tan(c+dx)^m (a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+adx}}{d}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 4082

$$\frac{a(A(5-4m)-i(4Bm+B)) \tan^{m+1}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{a^3(2m+1)(A(5-4m)-i(4Bm+B)) \int \frac{\tan^m(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{d} - \frac{3a^3(A-iB)\sqrt{1+i \tan(c+dx)} \tan^{m+1}(c+dx)}{2a^2} + \frac{a^3(2m+1)(A(5-4m)-i(4Bm+B)) \int \frac{\tan^m(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{d} - \frac{3a^3(A-iB)\sqrt{1+i \tan(c+dx)} \tan^{m+1}(c+dx)}{2a^2} + \frac{a^3(2m+1)(A(5-4m)-i(4Bm+B)) \int \frac{\tan^m(c+dx)}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{d} - \frac{3a^3(A-iB)\sqrt{1+i \tan(c+dx)} \tan^{m+1}(c+dx)}{2a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 77

3.216. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

$$\frac{a(A(5-4m)-i(4Bm+B)) \tan^{m+1}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{a^3(2m+1)(A(5-4m)-i(4Bm+B))(-i \tan(c+dx))^{-m} \tan^m(c+dx) \int \frac{(-i \tan(c+dx))^m}{\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{d} - \frac{3a^3(A-i)}{2a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

↓ 75

$$\frac{a(A(5-4m)-i(4Bm+B)) \tan^{m+1}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} - \frac{3a^3(A-iB)\sqrt{1+i \tan(c+dx)} \tan^{m+1}(c+dx) \operatorname{AppellF1}\left(m+1, \frac{1}{2}, 1, m+2, -i \tan(c+dx), i \tan(c+dx)\right)}{d(m+1)\sqrt{a+ia \tan(c+dx)}} - \frac{2ia^3(2m+1)}{6a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} \quad 6a^2$$

input `Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `((A + I*B)*Tan[c + d*x]^(1 + m))/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) + ((a*(A*(5 - 4*m) - I*(B + 4*B*m))*Tan[c + d*x]^(1 + m))/(d*Sqrt[a + I*a*Tan[c + d*x]]) - ((-3*a^3*(A - I*B)*AppellF1[1 + m, 1/2, 1, 2 + m, (-I)*Tan[c + d*x], I*Tan[c + d*x]]*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(1 + m))/(d*(1 + m)*Sqrt[a + I*a*Tan[c + d*x]]) - ((2*I)*a^2*(1 + 2*m)*(A*(5 - 4*m) - I*(B + 4*B*m))*Hypergeometric2F1[1/2, -m, 3/2, 1 + I*Tan[c + d*x]]*Tan[c + d*x]^m*Sqrt[a + I*a*Tan[c + d*x]])/(d*((-I)*Tan[c + d*x])^m)/(2*a^2))/(6*a^2)`

3.216.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 75 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((c + d*x)^(n + 1)/(d*(n + 1)*(-d/(b*c))^m)*Hypergeometric2F1[-m, n + 1, n + 2, 1 + d*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[n] && (IntegerQ[m] || GtQ[-d/(b*c), 0])`

3.216. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

- rule 77 $\text{Int}[(b \cdot x)^m (c + d \cdot x)^n, x_Symbol] \rightarrow \text{Simp}[(c/d)^m \text{IntPart}[m] (b \cdot x)^{\text{FracPart}[m]} / ((-d) \cdot (x/c))^{\text{FracPart}[m]}] \text{Int}[(c/d)^m (c + d \cdot x)^n, x] /;$ FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0]
- rule 150 $\text{Int}[(b \cdot x)^m (c + d \cdot x)^n (e + f \cdot x)^p, x] \rightarrow \text{Simp}[c^n e^p (b \cdot x)^{m+1} / (b^{m+1})] \text{AppellF1}[m+1, -n, -p, m+2, (-d) \cdot (x/c), (-f) \cdot (x/e)], x] /;$ FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[p] || GtQ[e, 0]
- rule 152 $\text{Int}[(b \cdot x)^m (c + d \cdot x)^n (e + f \cdot x)^p, x] \rightarrow \text{Simp}[c^n \text{IntPart}[n] (c + d \cdot x)^{\text{FracPart}[n]} / (1 + d \cdot (x/c))^{\text{FracPart}[n]}] \text{Int}[(b \cdot x)^{m+1} (1 + d \cdot (x/c))^n (e + f \cdot x)^p, x] /;$ FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]
- rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$ FunctionOfTrigOfLinearQ[u, x]
- rule 4047 $\text{Int}[(a + b \cdot \tan[e + f \cdot x])^m (c + d \cdot \tan[e + f \cdot x])^n, x_Symbol] \rightarrow \text{Simp}[a \cdot (b/f) \text{Subst}[\text{Int}[(a + x)^{m-1} (c + (d/b) \cdot x)^n / (b^2 + a \cdot x)], x, b \cdot \tan[e + f \cdot x]], x] /;$ FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
- rule 4079 $\text{Int}[(a + b \cdot \tan[e + f \cdot x])^m (A + B \cdot \tan[e + f \cdot x])^n, x_Symbol] \rightarrow \text{Simp}[(a \cdot A + b \cdot B) (a + b \cdot \tan[e + f \cdot x])^m (c + d \cdot \tan[e + f \cdot x])^{n+1} / (2 \cdot f \cdot m \cdot (b \cdot c - a \cdot d))] + \text{Simp}[1 / (2 \cdot a \cdot m \cdot (b \cdot c - a \cdot d))] \text{Int}[(a + b \cdot \tan[e + f \cdot x])^{m+1} (c + d \cdot \tan[e + f \cdot x])^n \text{Simp}[A \cdot (b \cdot c \cdot m - a \cdot d \cdot (2 \cdot m + n + 1)) + B \cdot (a \cdot c \cdot m - b \cdot d \cdot (n + 1)) + d \cdot (A \cdot b - a \cdot B) \cdot (m + n + 1) \cdot \tan[e + f \cdot x]], x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.216.4 Maple [F]

$$\int \frac{(\tan^m(dx + c))(A + B \tan(dx + c))}{(a + ia \tan(dx + c))^{\frac{3}{2}}} dx$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x)`

output `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x)`

3.216.5 Fracas [F]

$$\int \frac{\tan^m(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)^m}{(ia \tan(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="fracas")`

output `integral(1/4*sqrt(2)*((A - I*B)*e^(4*I*d*x + 4*I*c) + 2*A*e^(2*I*d*x + 2*I*c) + A + I*B)*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*e^(-3*I*d*x - 3*I*c)/a^2, x)`

3.216.6 Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \int \frac{(A+B\tan(c+dx))\tan^m(c+dx)}{(ia(\tan(c+dx)-i))^{\frac{3}{2}}} dx$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**m/(I*a*(tan(c + d*x) - I))**(3/2), x)`

3.216.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.216.8 Giac [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{(ia\tan(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(I*a*tan(d*x + c) + a)^(3/2), x)`

3.216.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \int \frac{\tan(c+dx)^m (A+B \tan(c+dx))}{(a+a \tan(c+dx) \text{ li})^{3/2}} dx$$

input `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(3/2),x)`

output `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(3/2), x)`

3.217 $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

3.217.1 Optimal result 2181
 3.217.2 Mathematica [F] 2182
 3.217.3 Rubi [A] (verified) 2182
 3.217.4 Maple [F] 2188
 3.217.5 Fricas [F] 2188
 3.217.6 Sympy [F] 2189
 3.217.7 Maxima [F(-2)] 2189
 3.217.8 Giac [F] 2189
 3.217.9 Mupad [F(-1)] 2190

3.217.1 Optimal result

Integrand size = 36, antiderivative size = 363

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{(A+iB) \tan^{1+m}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} + \frac{(iB(1-4m)+A(11-4m)) \tan^{1+m}(c+dx)}{30ad(a+ia \tan(c+dx))^{3/2}} - \frac{(iB(13+12m-16m^2)-A(37-52m+16m^2)) \tan^{1+m}(c+dx)}{60a^2d\sqrt{a+ia \tan(c+dx)}} + \frac{(A-iB) \operatorname{AppellF1}(1+m, \frac{1}{2}, 1, 2+m, -i \tan(c+dx), i \tan(c+dx)) \sqrt{1+i \tan(c+dx)} \tan^{1+m}(c+dx)}{8a^2d(1+m)\sqrt{a+ia \tan(c+dx)}} + \frac{(1+2m)(B(13+12m-16m^2)+iA(37-52m+16m^2)) \operatorname{Hypergeometric2F1}(\frac{1}{2}, -m, \frac{3}{2}, 1+i \tan(c+dx))}{60a^3d}$$

output

```
1/60*(1+2*m)*(B*(-16*m^2+12*m+13)+I*A*(16*m^2-52*m+37))*hypergeom([1/2, -m], [3/2], 1+I*tan(d*x+c))*(a+I*a*tan(d*x+c))^(1/2)*tan(d*x+c)^m/a^3/d/((-I*tan(d*x+c))^m)-1/60*(I*B*(-16*m^2+12*m+13)-A*(16*m^2-52*m+37))*tan(d*x+c)^(1+m)/a^2/d/(a+I*a*tan(d*x+c))^(1/2)+1/8*(A-I*B)*AppellF1(1+m, 1/2, 1, 2+m, -I*tan(d*x+c), I*tan(d*x+c))*(1+I*tan(d*x+c))^(1/2)*tan(d*x+c)^(1+m)/a^2/d/(1+m)/(a+I*a*tan(d*x+c))^(1/2)+1/5*(A+I*B)*tan(d*x+c)^(1+m)/d/(a+I*a*tan(d*x+c))^(5/2)+1/30*(I*B*(1-4*m)+A*(11-4*m))*tan(d*x+c)^(1+m)/a/d/(a+I*a*tan(d*x+c))^(3/2)
```

3.217.2 Mathematica [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx$$

input `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2), x]`

output `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2), x]`

3.217.3 Rubi [A] (verified)

Time = 2.15 (sec) , antiderivative size = 381, normalized size of antiderivative = 1.05, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4084, 3042, 4047, 25, 27, 152, 150, 4082, 77, 75}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(c+dx)^m(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{4079} \\ & \frac{\int \frac{\tan^m(c+dx)(2a(A(4-m)-iB(m+1))-a(iA-B)(3-2m)\tan(c+dx))}{2(i\tan(c+dx)a+a)^{3/2}} dx}{5a^2} + \frac{(A+iB)\tan^{m+1}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{\tan^m(c+dx)(2a(A(4-m)-iB(m+1))-a(iA-B)(3-2m)\tan(c+dx))}{(i\tan(c+dx)a+a)^{3/2}} dx}{10a^2} + \frac{(A+iB)\tan^{m+1}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{\tan(c+dx)^m(2a(A(4-m)-iB(m+1))-a(iA-B)(3-2m)\tan(c+dx))}{(i\tan(c+dx)a+a)^{3/2}} dx}{10a^2} + \frac{(A+iB)\tan^{m+1}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} \end{aligned}$$

3.217. $\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx$

↓ 4079

$$\int \frac{\tan^m(c+dx) \left(2a^2 (iB(-4m^2+3m+7) - A(4m^2-13m+13)) - a^2(B(1-4m) - iA(11-4m))(1-2m) \tan(c+dx) \right)}{2\sqrt{i \tan(c+dx)a+a} \cdot 3a^2} dx + \frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} \quad 10a^2$$

↓ 27

$$\frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \int \frac{\tan^m(c+dx) \left(2a^2 (iB(-4m^2+3m+7) - A(4m^2-13m+13)) - a^2(B(1-4m) - iA(11-4m))(1-2m) \tan(c+dx) \right)}{\sqrt{i \tan(c+dx)a+a} \cdot 6a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} \quad 10a^2$$

↓ 3042

$$\frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \int \frac{\tan(c+dx)^m \left(2a^2 (iB(-4m^2+3m+7) - A(4m^2-13m+13)) - a^2(B(1-4m) - iA(11-4m))(1-2m) \tan(c+dx) \right)}{\sqrt{i \tan(c+dx)a+a} \cdot 6a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} \quad 10a^2$$

↓ 4079

$$\frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{1}{2} \tan^m(c+dx) \sqrt{i \tan(c+dx)a+a} \left(2a^3 (A(16m^3-44m^2+11m+11) + iB(16m^3-4m^2-19m+1)) - a^3(2m+1)(B(11-4m) - iA(11-4m)) \right)}{a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} \quad 10a^2$$

↓ 27

$$\frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \tan^m(c+dx) \sqrt{i \tan(c+dx)a+a} \left(2a^3 (A(16m^3-44m^2+11m+11) + iB(16m^3-4m^2-19m+1)) - a^3(2m+1)(B(11-4m) - iA(11-4m)) \right)}{2a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} \quad 10a^2$$

↓ 3042

3.217. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \tan(c+dx)^m \sqrt{i \tan(c+dx)a+a} (2a^3(A(16m^3-44m^2+11m+1)+iB(16m^3-4m^2-19m+1))-a^3(2m+1)(B))}{2a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

$10a^2$

↓ 4084

$$\frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{-15a^3(A-iB) \int \tan^m(c+dx) \sqrt{i \tan(c+dx)a+adx-a^2(2m+1)} (-A(16m^2-52m+37)+iB(-16m^2+12m+13))}{2a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

$10a^2$

↓ 3042

$$\frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{-15a^3(A-iB) \int \tan(c+dx)^m \sqrt{i \tan(c+dx)a+adx-a^2(2m+1)} (-A(16m^2-52m+37)+iB(-16m^2+12m+13))}{2a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

$10a^2$

↓ 4047

$$\frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{15ia^5(A-iB) \int -\frac{\tan^m(c+dx)}{a(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}} d(ia \tan(c+dx)) - a^2(2m+1) (-A(16m^2-52m+37)+iB)}{2a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

$10a^2$

↓ 25

$$\frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{15ia^5(A-iB) \int -\frac{\tan^m(c+dx)}{a(a-ia \tan(c+dx)) \sqrt{i \tan(c+dx)a+a}} d(ia \tan(c+dx)) - a^2(2m+1) (-A(16m^2-52m+37)+iB)}{2a^2}$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

$10a^2$

↓ 27

3.217. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{15ia^4(A-iB) \int \frac{\tan^m(c+dx)}{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}} d(ia \tan(c+dx)) - a^2(2m+1)(-A(16m^2-52m+37)+iB(-16m^2+12m+13)) \int \frac{\tan^m(c+dx)}{\sqrt{i \tan(c+dx)+1(a-ia \tan(c+dx))}} d(ia \tan(c+dx))}{2a^2} = 10a^2$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

↓ 152

$$\frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{15ia^4(A-iB)\sqrt{1+i \tan(c+dx)} \int \frac{\tan^m(c+dx)}{\sqrt{i \tan(c+dx)+1(a-ia \tan(c+dx))}} d(ia \tan(c+dx)) - a^2(2m+1)(-A(16m^2-52m+37)+iB(-16m^2+12m+13)) \int \frac{\tan^m(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} d(ia \tan(c+dx))}{2a^2} = 10a^2$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

↓ 150

$$\frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{-a^2(2m+1)(-A(16m^2-52m+37)+iB(-16m^2+12m+13)) \int \tan(c+dx)^m(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)+a}}{2a^2} = 10a^2$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

↓ 4082

$$\frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{a^4(2m+1)(-A(16m^2-52m+37)+iB(-16m^2+12m+13)) \int \frac{\tan^m(c+dx)}{\sqrt{i \tan(c+dx)+a}} d \tan(c+dx) - 15a^4(A-iB)\sqrt{1+i \tan(c+dx)} \int \frac{\tan^m(c+dx)}{\sqrt{i \tan(c+dx)+1(a-ia \tan(c+dx))}} d(ia \tan(c+dx))}{2a^2} = 10a^2$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

↓ 77

$$\frac{a(A(11-4m)+iB(1-4m)) \tan^{m+1}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{a^4(2m+1)(-A(16m^2-52m+37)+iB(-16m^2+12m+13)) \tan^m(c+dx)(-i \tan(c+dx))^{-m} \int \frac{(-i \tan(c+dx))}{\sqrt{i \tan(c+dx)+a}} d \tan(c+dx)}{d} = 10a^2$$

$$\frac{(A+iB) \tan^{m+1}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}}$$

↓ 75

3.217. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\frac{a(A(11-4m)+iB(1-4m))\tan^{m+1}(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} - \frac{a^2(-A(16m^2-52m+37)+iB(-16m^2+12m+13))\tan^{m+1}(c+dx)}{d\sqrt{a+ia\tan(c+dx)}} + \frac{2ia^3(2m+1)(-A(16m^2-52m+37)+iB(-16m^2+12m+13))\tan^{m+1}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}}$$

input `Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `((A + I*B)*Tan[c + d*x]^(1 + m))/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) + ((a*(I*B*(1 - 4*m) + A*(11 - 4*m))*Tan[c + d*x]^(1 + m))/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) - ((a^2*(I*B*(13 + 12*m - 16*m^2) - A*(37 - 52*m + 16*m^2))*Tan[c + d*x]^(1 + m))/(d*Sqrt[a + I*a*Tan[c + d*x]]) + ((-15*a^4*(A - I*B)*AppellF1[1 + m, 1/2, 1, 2 + m, (-I)*Tan[c + d*x], I*Tan[c + d*x]]*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(1 + m))/(d*(1 + m)*Sqrt[a + I*a*Tan[c + d*x]]) + ((2*I)*a^3*(1 + 2*m)*(I*B*(13 + 12*m - 16*m^2) - A*(37 - 52*m + 16*m^2))*Hypergeometric2F1[1/2, -m, 3/2, 1 + I*Tan[c + d*x]]*Tan[c + d*x]^m*Sqrt[a + I*a*Tan[c + d*x]])/(d*((-I)*Tan[c + d*x])^m)/(2*a^2)/(6*a^2)/(10*a^2))`

3.217.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 75 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((c + d*x)^(n + 1)/(d*(n + 1)*(-d/(b*c))^m)*Hypergeometric2F1[-m, n + 1, n + 2, 1 + d*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[n] && (IntegerQ[m] || GtQ[-d/(b*c), 0])`

rule 77 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((-b)*(c/d))^IntPart[m]*((b*x)^FracPart[m]/((-d)*(x/c))^FracPart[m]) Int[((-d)*(x/c))^m*(c + d*x)^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0]`

3.217. $\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx$

- rule 150 `Int[((b_.)*(x_)^(m_)*((c_) + (d_.)*(x_)^(n_)*((e_) + (f_.)*(x_)^(p_)), x_] := Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])`
- rule 152 `Int[((b_.)*(x_)^(m_)*((c_) + (d_.)*(x_)^(n_)*((e_) + (f_.)*(x_)^(p_)), x_] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4047 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`
- rule 4082 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.217.4 Maple [F]

$$\int \frac{(\tan^m(dx + c))(A + B \tan(dx + c))}{(a + ia \tan(dx + c))^{\frac{5}{2}}} dx$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x)`

output `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x)`

3.217.5 Fracas [F]

$$\int \frac{\tan^m(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)^m}{(ia \tan(dx + c) + a)^{\frac{5}{2}}} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="fricas")`

output `integral(1/8*sqrt(2)*((A - I*B)*e^(6*I*d*x + 6*I*c) + (3*A - I*B)*e^(4*I*d*x + 4*I*c) + (3*A + I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*e^(-5*I*d*x - 5*I*c)/a^3, x)`

3.217.6 Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\tan^m(c+dx)}{(ia(\tan(c+dx)-i))^{\frac{5}{2}}} dx$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**m/(I*a*(tan(c + d*x) - I))**(5/2), x)`

3.217.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.217.8 Giac [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{(ia\tan(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(I*a*tan(d*x + c) + a)^(5/2), x)`

3.217.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \int \frac{\tan(c+dx)^m (A+B \tan(c+dx))}{(a+a \tan(c+dx) \text{ li})^{5/2}} dx$$

input `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2),x)`

output `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2), x)`

3.218 $\int \tan^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$

3.218.1 Optimal result	2191
3.218.2 Mathematica [F]	2191
3.218.3 Rubi [A] (verified)	2192
3.218.4 Maple [F]	2195
3.218.5 Fricas [F]	2196
3.218.6 Sympy [F]	2196
3.218.7 Maxima [F]	2196
3.218.8 Giac [F]	2197
3.218.9 Mupad [F(-1)]	2197

3.218.1 Optimal result

Integrand size = 34, antiderivative size = 167

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \frac{(A - iB) \operatorname{AppellF1}(1 + m, 1 - n, 1, 2 + m, -i \tan(c + dx), i \tan(c + dx))(1 + i \tan(c + dx))^{-n} \tan^{1+m}(c + dx)}{d(1 + m)} + \frac{iB \operatorname{Hypergeometric2F1}(1 + m, 1 - n, 2 + m, -i \tan(c + dx))(1 + i \tan(c + dx))^{-n} \tan^{1+m}(c + dx)(a + i \tan(c + dx))}{d(1 + m)}$$

output $(A-I*B)*\operatorname{AppellF1}(1+m,1-n,1,2+m,-I*\tan(d*x+c),I*\tan(d*x+c))*\tan(d*x+c)^{(1+m)}*(a+I*a*\tan(d*x+c))^n/d/(1+m)/((1+I*\tan(d*x+c))^n)+I*B*\operatorname{hypergeom}([1+m,1-n],[2+m],[-I*\tan(d*x+c))*\tan(d*x+c)^{(1+m)}*(a+I*a*\tan(d*x+c))^n/d/(1+m)/((1+I*\tan(d*x+c))^n)$

3.218.2 Mathematica [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \tan^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

input `Integrate[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `Integrate[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

3.218.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.324$, Rules used = {3042, 4084, 3042, 4047, 25, 27, 152, 150, 4082, 76, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c + dx)^m(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4084} \\
 & \frac{(A - iB) \int \tan^m(c + dx)(i \tan(c + dx)a + a)^n dx + iB \int \tan^m(c + dx)(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A - iB) \int \tan(c + dx)^m(i \tan(c + dx)a + a)^n dx + iB \int \tan(c + dx)^m(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n dx}{a} \\
 & \quad \downarrow \text{4047} \\
 & \frac{ia^2(A - iB) \int -\frac{\tan^m(c+dx)(i \tan(c+dx)a+a)^{n-1}}{a(a-ia \tan(c+dx))} d(ia \tan(c + dx))}{a} + \\
 & \frac{iB \int \tan(c + dx)^m(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n dx}{a} \\
 & \quad \downarrow \text{25} \\
 & \frac{iB \int \tan(c + dx)^m(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n dx}{a} - \\
 & \frac{ia^2(A - iB) \int \frac{\tan^m(c+dx)(i \tan(c+dx)a+a)^{n-1}}{a(a-ia \tan(c+dx))} d(ia \tan(c + dx))}{d}
 \end{aligned}$$

3.218. $\int \tan^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{iB \int \tan(c+dx)^m (a - ia \tan(c+dx)) (i \tan(c+dx) a + a)^n dx}{ia(A - iB) \int \frac{\tan^m(c+dx) (i \tan(c+dx) a + a)^{n-1} d(ia \tan(c+dx))}{a - ia \tan(c+dx)}} \\
 & \downarrow 152 \\
 & \frac{iB \int \tan(c+dx)^m (a - ia \tan(c+dx)) (i \tan(c+dx) a + a)^n dx}{i(A - iB) (1 + i \tan(c+dx))^{-n} (a + ia \tan(c+dx))^n \int \frac{(i \tan(c+dx) + 1)^{n-1} \tan^m(c+dx) d(ia \tan(c+dx))}{a - ia \tan(c+dx)}} \\
 & \downarrow 150 \\
 & \frac{iB \int \tan(c+dx)^m (a - ia \tan(c+dx)) (i \tan(c+dx) a + a)^n dx + (A - iB) \tan^{m+1}(c+dx) (1 + i \tan(c+dx))^{-n} (a + ia \tan(c+dx))^n \text{AppellF1}(m+1, 1-n, 1, m+2, -i \tan(c+dx))}{d(m+1)} \\
 & \downarrow 4082 \\
 & \frac{iaB \int \tan^m(c+dx) (i \tan(c+dx) a + a)^{n-1} d \tan(c+dx) + (A - iB) \tan^{m+1}(c+dx) (1 + i \tan(c+dx))^{-n} (a + ia \tan(c+dx))^n \text{AppellF1}(m+1, 1-n, 1, m+2, -i \tan(c+dx))}{d(m+1)} \\
 & \downarrow 76 \\
 & \frac{iB (1 + i \tan(c+dx))^{-n} (a + ia \tan(c+dx))^n \int (i \tan(c+dx) + 1)^{n-1} \tan^m(c+dx) d \tan(c+dx) + (A - iB) \tan^{m+1}(c+dx) (1 + i \tan(c+dx))^{-n} (a + ia \tan(c+dx))^n \text{AppellF1}(m+1, 1-n, 1, m+2, -i \tan(c+dx))}{d(m+1)} \\
 & \downarrow 74 \\
 & \frac{(A - iB) \tan^{m+1}(c+dx) (1 + i \tan(c+dx))^{-n} (a + ia \tan(c+dx))^n \text{AppellF1}(m+1, 1-n, 1, m+2, -i \tan(c+dx)) + iB \tan^{m+1}(c+dx) (1 + i \tan(c+dx))^{-n} (a + ia \tan(c+dx))^n \text{Hypergeometric2F1}(m+1, 1-n, m+2, -i \tan(c+dx))}{d(m+1)}
 \end{aligned}$$

input `Int[Tan[c + d*x]^m*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

3.218. $\int \tan^m(c+dx)(a + ia \tan(c+dx))^n(A + B \tan(c+dx)) dx$


```
output ((A - I*B)*AppellF1[1 + m, 1 - n, 1, 2 + m, (-I)*Tan[c + d*x], I*Tan[c + d
*x]]*Tan[c + d*x]^(1 + m)*(a + I*a*Tan[c + d*x])^n)/(d*(1 + m)*(1 + I*Tan[
c + d*x])^n) + (I*B*Hypergeometric2F1[1 + m, 1 - n, 2 + m, (-I)*Tan[c + d*
x]]*Tan[c + d*x]^(1 + m)*(a + I*a*Tan[c + d*x])^n)/(d*(1 + m)*(1 + I*Tan[c
+ d*x])^n)
```

3.218.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 74 Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x
)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x]
/; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0]
&& !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))
```

```
rule 76 Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[c^IntPart
[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*
(x/c))^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !Integer
Q[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0] && ((RationalQ[m] && !(EqQ[n, -2
^(-1)] && EqQ[c^2 - d^2, 0])) || !RationalQ[n])
```

```
rule 150 Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_
] := Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2
, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !In
tegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])
```

```
rule 152 Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_
] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n])
Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x], x] /; FreeQ[{b, c, d, e, f, m,
n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]
```

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4047 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.218.4 Maple [F]

$$\int (\tan^m(dx + c)) (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.218.5 Fracas [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(((A - I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*(2*a*e^(2*I*d*x + 2*I*c))/(e^(2*I*d*x + 2*I*c) + 1))^n*((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))^m/(e^(2*I*d*x + 2*I*c) + 1), x)`

3.218.6 Sympy [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (ia(\tan(c + dx) - i))^n(A + B \tan(c + dx)) \tan^m(c + dx) dx$$

input `integrate(tan(d*x+c)**m*(a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x))*tan(c + d*x)**m, x)`

3.218.7 Maxima [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*tan(d*x + c)^m, x)`

3.218.8 Giac [F]

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*tan(d*x + c)^m, x)`

3.218.9 Mupad [F(-1)]

Timed out.

$$\int \tan^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^m (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^n dx$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n,x)`

output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n, x)`

3.219 $\int \tan^3(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.219.1 Optimal result	2198
3.219.2 Mathematica [A] (verified)	2199
3.219.3 Rubi [A] (verified)	2199
3.219.4 Maple [F]	2203
3.219.5 Fracas [F]	2203
3.219.6 Sympy [F]	2203
3.219.7 Maxima [F]	2204
3.219.8 Giac [F]	2204
3.219.9 Mupad [F(-1)]	2204

3.219.1 Optimal result

Integrand size = 34, antiderivative size = 245

$$\int \tan^3(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \frac{2(iBn - A(3+n))(a+ia \tan(c+dx))^n}{dn(2+n)(3+n)}$$

$$+ \frac{(A-iB) \operatorname{Hypergeometric2F1}\left(1, n, 1+n, \frac{1}{2}(1+i \tan(c+dx))\right) (a+ia \tan(c+dx))^n}{2dn}$$

$$- \frac{(iBn - A(3+n)) \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(2+n)(3+n)}$$

$$+ \frac{B \tan^3(c+dx)(a+ia \tan(c+dx))^n}{d(3+n)}$$

$$- \frac{(An(3+n) - iB(6+3n+n^2))(a+ia \tan(c+dx))^{1+n}}{ad(1+n)(2+n)(3+n)}$$

```
output 2*(I*B*n-A*(3+n))*(a+I*a*tan(d*x+c))^n/d/n/(2+n)/(3+n)+1/2*(A-I*B)*hypergeometric([1, n],[1+n],1/2+1/2*I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/n-(I*B*n-A*(3+n))*tan(d*x+c)^2*(a+I*a*tan(d*x+c))^n/d/(2+n)/(3+n)+B*tan(d*x+c)^3*(a+I*a*tan(d*x+c))^n/d/(3+n)-(A*n*(3+n)-I*B*(n^2+3*n+6))*(a+I*a*tan(d*x+c))^(1+n)/a/d/(3+n)/(n^2+3*n+2)
```

3.219.2 Mathematica [A] (verified)

Time = 4.14 (sec) , antiderivative size = 189, normalized size of antiderivative = 0.77

$$\int \tan^3(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \frac{(a + ia \tan(c + dx))^n (2iBn(8 + 5n + n^2) - 2A(6 + 8n + 5n^2 + n^3) + (A - iB)(6 + 11n + 6n^2 + n^3)) \operatorname{Hy}}{}$$

input `Integrate[Tan[c + d*x]^3*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `((a + I*a*Tan[c + d*x])^n*((2*I)*B*n*(8 + 5*n + n^2) - 2*A*(6 + 8*n + 5*n^2 + n^3) + (A - I*B)*(6 + 11*n + 6*n^2 + n^3))*Hypergeometric2F1[1, n, 1 + n, (1 + I*Tan[c + d*x])/2] - 2*n*(I*A*n*(3 + n) + B*(6 + 3*n + n^2))*Tan[c + d*x] + 2*n*(1 + n)*((-I)*B*n + A*(3 + n))*Tan[c + d*x]^2 + 2*B*n*(2 + 3*n + n^2)*Tan[c + d*x]^3)/(2*d*n*(1 + n)*(2 + n)*(3 + n))`

3.219.3 Rubi [A] (verified)

Time = 1.28 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.06, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.382$, Rules used = {3042, 4080, 25, 3042, 4080, 25, 3042, 4075, 3042, 4010, 3042, 3962, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^3(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^3(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4080}$$

$$\int \frac{-\tan^2(c + dx)(i \tan(c + dx)a + a)^n(3aB + a(iBn - A(n + 3)) \tan(c + dx)) dx}{a(n + 3)} + \frac{B \tan^3(c + dx)(a + ia \tan(c + dx))^n}{d(n + 3)}$$

$$\downarrow \text{25}$$

$$\begin{aligned}
 & \frac{B \tan^3(c+dx)(a+ia \tan(c+dx))^n}{d(n+3)} - \frac{\int \tan^2(c+dx)(i \tan(c+dx)a+a)^n(3aB+a(iBn-A(n+3))) \tan(c+dx) dx}{a(n+3)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{B \tan^3(c+dx)(a+ia \tan(c+dx))^n}{d(n+3)} - \frac{\int \tan(c+dx)^2(i \tan(c+dx)a+a)^n(3aB+a(iBn-A(n+3))) \tan(c+dx) dx}{a(n+3)} \\
 & \quad \downarrow \text{4080} \\
 & \frac{B \tan^3(c+dx)(a+ia \tan(c+dx))^n}{d(n+3)} - \frac{\int -\tan(c+dx)(i \tan(c+dx)a+a)^n(2a^2(iBn-A(n+3))-a^2(iAn(n+3)+B(n^2+3n+6))) \tan(c+dx) dx}{a(n+2)} + \frac{a(-A(n+3)+iBn) \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} \\
 & \quad \downarrow \text{25} \\
 & \frac{B \tan^3(c+dx)(a+ia \tan(c+dx))^n}{d(n+3)} - \frac{a(-A(n+3)+iBn) \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \frac{\int \tan(c+dx)(i \tan(c+dx)a+a)^n(2a^2(iBn-A(n+3))-a^2(iAn(n+3)+B(n^2+3n+6))) \tan(c+dx) dx}{a(n+2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{B \tan^3(c+dx)(a+ia \tan(c+dx))^n}{d(n+3)} - \frac{a(-A(n+3)+iBn) \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \frac{\int \tan(c+dx)(i \tan(c+dx)a+a)^n(2a^2(iBn-A(n+3))-a^2(iAn(n+3)+B(n^2+3n+6))) \tan(c+dx) dx}{a(n+2)} \\
 & \quad \downarrow \text{4075} \\
 & \frac{B \tan^3(c+dx)(a+ia \tan(c+dx))^n}{d(n+3)} - \frac{a(-A(n+3)+iBn) \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \frac{\int (i \tan(c+dx)a+a)^n((iAn(n+3)+B(n^2+3n+6))a^2+2(iBn-A(n+3)) \tan(c+dx)a^2) dx}{a(n+2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{B \tan^3(c+dx)(a+ia \tan(c+dx))^n}{d(n+3)} - \frac{a(-A(n+3)+iBn) \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \frac{\int (i \tan(c+dx)a+a)^n((iAn(n+3)+B(n^2+3n+6))a^2+2(iBn-A(n+3)) \tan(c+dx)a^2) dx}{a(n+2)} \\
 & \quad \downarrow \\
 & \frac{B \tan^3(c+dx)(a+ia \tan(c+dx))^n}{d(n+3)} - \frac{a(-A(n+3)+iBn) \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \frac{\int (i \tan(c+dx)a+a)^n((iAn(n+3)+B(n^2+3n+6))a^2+2(iBn-A(n+3)) \tan(c+dx)a^2) dx}{a(n+2)}
 \end{aligned}$$

3.219. $\int \tan^3(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

$$\begin{aligned}
 & \downarrow 4010 \\
 & \frac{B \tan^3(c + dx)(a + ia \tan(c + dx))^n}{d(n + 3)} - \\
 & \frac{\frac{a(-A(n+3)+iBn) \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \frac{a^2(n^2+5n+6)(B+iA) \int (i \tan(c+dx)a+a)^n dx + \frac{2a^2(-A(n+3)+iBn)(a+ia \tan(c+dx))^n}{dn} - \frac{a(A}{a(n+2)}}{a(n+3)} \\
 & \downarrow 3042 \\
 & \frac{B \tan^3(c + dx)(a + ia \tan(c + dx))^n}{d(n + 3)} - \\
 & \frac{\frac{a(-A(n+3)+iBn) \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \frac{a^2(n^2+5n+6)(B+iA) \int (i \tan(c+dx)a+a)^n dx + \frac{2a^2(-A(n+3)+iBn)(a+ia \tan(c+dx))^n}{dn} - \frac{a(A}{a(n+2)}}{a(n+3)} \\
 & \downarrow 3962 \\
 & \frac{B \tan^3(c + dx)(a + ia \tan(c + dx))^n}{d(n + 3)} - \\
 & \frac{\frac{a(-A(n+3)+iBn) \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \frac{ia^3(n^2+5n+6)(B+iA) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{a-ia \tan(c+dx)} d(ia \tan(c+dx)) + \frac{2a^2(-A(n+3)+iBn)(a+ia \tan(c+dx))^n}{dn}}{a(n+2)}}{a(n+3)} \\
 & \downarrow 78 \\
 & \frac{B \tan^3(c + dx)(a + ia \tan(c + dx))^n}{d(n + 3)} - \\
 & \frac{\frac{a(-A(n+3)+iBn) \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \frac{ia^2(n^2+5n+6)(B+iA)(a+ia \tan(c+dx))^n \operatorname{Hypergeometric2F1}\left(1, n, n+1, \frac{i \tan(c+dx)a+a}{2a}\right) + 2a^2}{2dn}}{a(n+2)}}{a(n+3)}
 \end{aligned}$$

input `Int[Tan[c + d*x]^3*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `(B*Tan[c + d*x]^3*(a + I*a*Tan[c + d*x])^n)/(d*(3 + n)) - ((a*(I*B*n - A*(3 + n))*Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^n)/(d*(2 + n)) - ((2*a^2*(I*B*n - A*(3 + n))*(a + I*a*Tan[c + d*x])^n)/(d*n) - ((I/2)*a^2*(I*A + B)*(6 + 5*n + n^2)*Hypergeometric2F1[1, n, 1 + n, (a + I*a*Tan[c + d*x])/(2*a)]*(a + I*a*Tan[c + d*x])^n)/(d*n) - (a*(A*n*(3 + n) - I*B*(6 + 3*n + n^2))*(a + I*a*Tan[c + d*x])^(1 + n))/(d*(1 + n)))/(a*(2 + n))/(a*(3 + n))`

3.219.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3962 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[-b/d Subst[Int[(a + x)^(n - 1)/(a - x), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 + b^2, 0]`
- rule 4010 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`
- rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`
- rule 4080 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`

3.219.4 Maple [F]

$$\int (\tan^3(dx + c)) (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^3*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^3*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.219.5 Fricas [F]

$$\begin{aligned} & \int \tan^3(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^3 dx \end{aligned}$$

input `integrate(tan(d*x+c)^3*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(((I*A + B)*e^(8*I*d*x + 8*I*c) - 2*(I*A + 2*B)*e^(6*I*d*x + 6*I*c) + 6*B*e^(4*I*d*x + 4*I*c) - 2*(-I*A + 2*B)*e^(2*I*d*x + 2*I*c) - I*A + B)*(2*a*e^(2*I*d*x + 2*I*c)/(e^(2*I*d*x + 2*I*c) + 1))^n/(e^(8*I*d*x + 8*I*c) + 4*e^(6*I*d*x + 6*I*c) + 6*e^(4*I*d*x + 4*I*c) + 4*e^(2*I*d*x + 2*I*c) + 1), x)`

3.219.6 Sympy [F]

$$\begin{aligned} & \int \tan^3(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (ia(\tan(c + dx) - i))^n (A + B \tan(c + dx)) \tan^3(c + dx) dx \end{aligned}$$

input `integrate(tan(d*x+c)**3*(a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x))*tan(c + d*x)**3, x)`

3.219.7 Maxima [F]

$$\int \tan^3(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^3 dx$$

input `integrate(tan(d*x+c)^3*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*tan(d*x + c)^3, x)`

3.219.8 Giac [F]

$$\int \tan^3(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^3 dx$$

input `integrate(tan(d*x+c)^3*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*tan(d*x + c)^3, x)`

3.219.9 Mupad [F(-1)]

Timed out.

$$\int \tan^3(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^3 (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^n dx$$

input `int(tan(c + d*x)^3*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n,x)`

output `int(tan(c + d*x)^3*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n, x)`

3.220 $\int \tan^2(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.220.1 Optimal result	2205
3.220.2 Mathematica [A] (verified)	2206
3.220.3 Rubi [A] (verified)	2206
3.220.4 Maple [F]	2209
3.220.5 Fricas [F]	2209
3.220.6 Sympy [F]	2210
3.220.7 Maxima [F]	2210
3.220.8 Giac [F]	2210
3.220.9 Mupad [F(-1)]	2211

3.220.1 Optimal result

Integrand size = 34, antiderivative size = 164

$$\int \tan^2(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= -\frac{2B(a+ia \tan(c+dx))^n}{dn(2+n)}$$

$$+ \frac{(iA+B) \operatorname{Hypergeometric2F1}\left(1, n, 1+n, \frac{1}{2}(1+i \tan(c+dx))\right) (a+ia \tan(c+dx))^n}{2dn}$$

$$+ \frac{B \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(2+n)} - \frac{(Bn+iA(2+n))(a+ia \tan(c+dx))^{1+n}}{ad(1+n)(2+n)}$$

```
output -2*B*(a+I*a*tan(d*x+c))^n/d/n/(2+n)+1/2*(I*A+B)*hypergeom([1, n],[1+n],1/2
+1/2*I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/n+B*tan(d*x+c)^2*(a+I*a*tan(d*x+
c))^n/d/(2+n)-(B*n+I*A*(2+n))*(a+I*a*tan(d*x+c))^(1+n)/a/d/(1+n)/(2+n)
```

3.220.2 Mathematica [A] (verified)

Time = 1.48 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.84

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \frac{(a + ia \tan(c + dx))^n (-4B - 4iAn - 4Bn - 2iAn^2 - 2Bn^2 + (iA + B)(2 + 3n + n^2) \text{Hypergeometric2F1}[1, n, 1 + n, (1 + I \tan[c + d*x])/2] + 2*n*((-I)*B*n + A*(2 + n))*\tan[c + d*x] + 2*B*n*(1 + n)*\tan[c + d*x]^2)}{2*d*n*(1 + n)*(2 + n)}$$

input `Integrate[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`output `((a + I*a*Tan[c + d*x])^n*(-4*B - (4*I)*A*n - 4*B*n - (2*I)*A*n^2 - 2*B*n^2 + (I*A + B)*(2 + 3*n + n^2)*Hypergeometric2F1[1, n, 1 + n, (1 + I*Tan[c + d*x])/2] + 2*n*((-I)*B*n + A*(2 + n))*Tan[c + d*x] + 2*B*n*(1 + n)*Tan[c + d*x]^2))/(2*d*n*(1 + n)*(2 + n))`**3.220.3 Rubi [A] (verified)**Time = 0.79 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3042, 4080, 25, 3042, 4075, 3042, 4010, 3042, 3962, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^2(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4080}$$

$$\frac{\int -\tan(c + dx)(i \tan(c + dx)a + a)^n(2aB + a(iBn - A(n + 2)) \tan(c + dx)) dx}{a(n + 2)} + \frac{B \tan^2(c + dx)(a + ia \tan(c + dx))^n}{d(n + 2)}$$

$$\downarrow \text{25}$$

$$\begin{aligned}
& \frac{B \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \\
& \frac{\int \tan(c+dx)(i \tan(c+dx)a+a)^n(2aB+a(iBn-A(n+2)) \tan(c+dx))dx}{a(n+2)} \\
& \quad \downarrow \text{3042} \\
& \frac{B \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \\
& \frac{\int \tan(c+dx)(i \tan(c+dx)a+a)^n(2aB+a(iBn-A(n+2)) \tan(c+dx))dx}{a(n+2)} \\
& \quad \downarrow \text{4075} \\
& \frac{B \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \\
& \frac{\int (i \tan(c+dx)a+a)^n(2aB \tan(c+dx)-a(iBn-A(n+2)))dx + \frac{(Bn+iA(n+2))(a+ia \tan(c+dx))^{n+1}}{d(n+1)}}{a(n+2)} \\
& \quad \downarrow \text{3042} \\
& \frac{B \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \\
& \frac{\int (i \tan(c+dx)a+a)^n(2aB \tan(c+dx)-a(iBn-A(n+2)))dx + \frac{(Bn+iA(n+2))(a+ia \tan(c+dx))^{n+1}}{d(n+1)}}{a(n+2)} \\
& \quad \downarrow \text{4010} \\
& \frac{B \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \\
& \frac{a(n+2)(A-iB) \int (i \tan(c+dx)a+a)^n dx + \frac{(Bn+iA(n+2))(a+ia \tan(c+dx))^{n+1}}{d(n+1)} + \frac{2aB(a+ia \tan(c+dx))^n}{dn}}{a(n+2)} \\
& \quad \downarrow \text{3042} \\
& \frac{B \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \\
& \frac{a(n+2)(A-iB) \int (i \tan(c+dx)a+a)^n dx + \frac{(Bn+iA(n+2))(a+ia \tan(c+dx))^{n+1}}{d(n+1)} + \frac{2aB(a+ia \tan(c+dx))^n}{dn}}{a(n+2)} \\
& \quad \downarrow \text{3962} \\
& \frac{B \tan^2(c+dx)(a+ia \tan(c+dx))^n}{d(n+2)} - \\
& \frac{ia^2(n+2)(A-iB) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{a-ia \tan(c+dx)} d(i \tan(c+dx)) + \frac{(Bn+iA(n+2))(a+ia \tan(c+dx))^{n+1}}{d(n+1)} + \frac{2aB(a+ia \tan(c+dx))^n}{dn}}{a(n+2)} \\
& \quad \downarrow \text{78}
\end{aligned}$$

3.220. $\int \tan^2(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

$$\frac{B \tan^2(c + dx)(a + ia \tan(c + dx))^n}{d(n + 2)} - \frac{ia(n+2)(A - iB)(a + ia \tan(c + dx))^n \operatorname{Hypergeometric2F1}\left(1, n, n+1, \frac{i \tan(c + dx)a + a}{2a}\right)}{2dn} + \frac{(Bn + iA(n+2))(a + ia \tan(c + dx))^{n+1}}{d(n+1)} + \frac{2aB(a + ia \tan(c + dx))^n}{dn}$$

$$a(n + 2)$$

input `Int[Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

output `(B*Tan[c + d*x]^2*(a + I*a*Tan[c + d*x])^n)/(d*(2 + n)) - ((2*a*B*(a + I*a*Tan[c + d*x])^n)/(d*n) - ((I/2)*a*(A - I*B)*(2 + n)*Hypergeometric2F1[1, n, 1 + n, (a + I*a*Tan[c + d*x])/(2*a)]*(a + I*a*Tan[c + d*x])^n)/(d*n) + ((B*n + I*A*(2 + n))*(a + I*a*Tan[c + d*x])^(1 + n))/(d*(1 + n)))/(a*(2 + n))`

3.220.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3962 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[-b/d Subst[Int[(a + x)^(n - 1)/(a - x), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 + b^2, 0]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4080 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`

3.220.4 Maple [F]

$$\int (\tan^2(dx + c)) (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.220.5 Fricas [F]

$$\begin{aligned} & \int \tan^2(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^2 dx \end{aligned}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(-(A - I*B)*e^(6*I*d*x + 6*I*c) - (A - 3*I*B)*e^(4*I*d*x + 4*I*c) - (A + 3*I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*(2*a*e^(2*I*d*x + 2*I*c)/(e^(2*I*d*x + 2*I*c) + 1))^n/(e^(6*I*d*x + 6*I*c) + 3*e^(4*I*d*x + 4*I*c) + 3*e^(2*I*d*x + 2*I*c) + 1), x)`

3.220.6 Sympy [F]

$$\begin{aligned} & \int \tan^2(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (ia(\tan(c + dx) - i))^n(A + B \tan(c + dx)) \tan^2(c + dx) dx \end{aligned}$$

input `integrate(tan(d*x+c)**2*(a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x))*tan(c + d*x)**2, x)`

3.220.7 Maxima [F]

$$\begin{aligned} & \int \tan^2(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^2 dx \end{aligned}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*tan(d*x + c)^2, x)`

3.220.8 Giac [F]

$$\begin{aligned} & \int \tan^2(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^2 dx \end{aligned}$$

input `integrate(tan(d*x+c)^2*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*tan(d*x + c)^2, x)`

3.220.9 Mupad [F(-1)]

Timed out.

$$\int \tan^2(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^2 (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^n dx$$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n,x)`output `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n, x)`

3.221 $\int \tan(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.221.1 Optimal result	2212
3.221.2 Mathematica [A] (verified)	2212
3.221.3 Rubi [A] (verified)	2213
3.221.4 Maple [F]	2215
3.221.5 Fricas [F]	2215
3.221.6 Sympy [F]	2216
3.221.7 Maxima [F]	2216
3.221.8 Giac [F]	2216
3.221.9 Mupad [F(-1)]	2217

3.221.1 Optimal result

Integrand size = 32, antiderivative size = 111

$$\int \tan(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \frac{A(a+ia \tan(c+dx))^n}{dn} - \frac{(A-iB) \operatorname{Hypergeometric2F1}\left(1, n, 1+n, \frac{1}{2}(1+i \tan(c+dx))\right) (a+ia \tan(c+dx))^n}{2dn} - \frac{iB(a+ia \tan(c+dx))^{1+n}}{ad(1+n)}$$

```
output A*(a+I*a*tan(d*x+c))^n/d/n-1/2*(A-I*B)*hypergeom([1, n], [1+n], 1/2+1/2*I*ta
n(d*x+c))*(a+I*a*tan(d*x+c))^n/d/n-I*B*(a+I*a*tan(d*x+c))^(1+n)/a/d/(1+n)
```

3.221.2 Mathematica [A] (verified)

Time = 0.81 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.78

$$\int \tan(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \frac{(a+ia \tan(c+dx))^n \left(-((A-iB)(1+n) \operatorname{Hypergeometric2F1}\left(1, n, 1+n, \frac{1}{2}(1+i \tan(c+dx))\right)) + 2(A+B \tan(c+dx)) \right)}{2dn(1+n)}$$

input `Integrate[Tan[c + d*x]*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `((a + I*a*Tan[c + d*x])^n*(-((A - I*B)*(1 + n)*Hypergeometric2F1[1, n, 1 + n, (1 + I*Tan[c + d*x])/2]) + 2*(A + A*n - I*B*n + B*n*Tan[c + d*x]))/(2*d*n*(1 + n))`

3.221.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.219$, Rules used = {3042, 4075, 3042, 4010, 3042, 3962, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4075} \\
 & \int (i \tan(c + dx)a + a)^n(A \tan(c + dx) - B) dx - \frac{iB(a + ia \tan(c + dx))^{n+1}}{ad(n + 1)} \\
 & \quad \downarrow \text{3042} \\
 & \int (i \tan(c + dx)a + a)^n(A \tan(c + dx) - B) dx - \frac{iB(a + ia \tan(c + dx))^{n+1}}{ad(n + 1)} \\
 & \quad \downarrow \text{4010} \\
 & -(B + iA) \int (i \tan(c + dx)a + a)^n dx + \frac{A(a + ia \tan(c + dx))^n}{dn} - \frac{iB(a + ia \tan(c + dx))^{n+1}}{ad(n + 1)} \\
 & \quad \downarrow \text{3042} \\
 & -(B + iA) \int (i \tan(c + dx)a + a)^n dx + \frac{A(a + ia \tan(c + dx))^n}{dn} - \frac{iB(a + ia \tan(c + dx))^{n+1}}{ad(n + 1)} \\
 & \quad \downarrow \text{3962}
 \end{aligned}$$

$$\frac{ia(B + iA) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{a-ia \tan(c+dx)} d(ia \tan(c + dx))}{d} + \frac{A(a + ia \tan(c + dx))^n}{dn} - \frac{iB(a + ia \tan(c + dx))^{n+1}}{ad(n + 1)}$$

↓ 78

$$\frac{i(B + iA)(a + ia \tan(c + dx))^n \text{Hypergeometric2F1}\left(1, n, n + 1, \frac{i \tan(c+dx)a+a}{2a}\right)}{dn} + \frac{A(a + ia \tan(c + dx))^n}{dn} - \frac{iB(a + ia \tan(c + dx))^{n+1}}{ad(n + 1)}$$

input `Int[Tan[c + d*x]*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `(A*(a + I*a*Tan[c + d*x])^n)/(d*n) + ((I/2)*(I*A + B)*Hypergeometric2F1[1, n, 1 + n, (a + I*a*Tan[c + d*x])/(2*a)]*(a + I*a*Tan[c + d*x])^n)/(d*n) - (I*B*(a + I*a*Tan[c + d*x])^(1 + n))/(a*d*(1 + n))`

3.221.3.1 Defintions of rubi rules used

rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b *c - a*d)^n*(a + b*x)^(m + 1)/(b^(n + 1)*(m + 1))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3962 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[-b/d Subst[Int[(a + x)^(n - 1)/(a - x), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 + b^2, 0]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.221.4 Maple [F]

$$\int \tan(dx + c) (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.221.5 Fracas [F]

$$\begin{aligned} & \int \tan(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c) dx \end{aligned}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(((-I*A - B)*e^(4*I*d*x + 4*I*c) + 2*B*e^(2*I*d*x + 2*I*c) + I*A - B)*(2*a*e^(2*I*d*x + 2*I*c)/(e^(2*I*d*x + 2*I*c) + 1))^n/(e^(4*I*d*x + 4*I*c) + 2*e^(2*I*d*x + 2*I*c) + 1), x)`

3.221.6 Sympy [F]

$$\begin{aligned} & \int \tan(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (ia(\tan(c + dx) - i))^n(A + B \tan(c + dx)) \tan(c + dx) dx \end{aligned}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x))*tan(c + d*x), x)`

3.221.7 Maxima [F]

$$\begin{aligned} & \int \tan(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c) dx \end{aligned}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*tan(d*x + c), x)`

3.221.8 Giac [F]

$$\begin{aligned} & \int \tan(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c) dx \end{aligned}$$

input `integrate(tan(d*x+c)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*tan(d*x + c), x)`

3.221.9 Mupad [F(-1)]

Timed out.

$$\int \tan(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx) (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^n dx$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n,x)`output `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n, x)`

3.222 $\int (a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$

3.222.1 Optimal result	2218
3.222.2 Mathematica [A] (verified)	2218
3.222.3 Rubi [A] (verified)	2219
3.222.4 Maple [F]	2220
3.222.5 Fricas [F]	2220
3.222.6 Sympy [F]	2221
3.222.7 Maxima [F]	2221
3.222.8 Giac [F]	2222
3.222.9 Mupad [F(-1)]	2222

3.222.1 Optimal result

Integrand size = 26, antiderivative size = 78

$$\int (a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \frac{B(a + ia \tan(c + dx))^n}{dn} - \frac{(iA + B) \operatorname{Hypergeometric2F1}\left(1, n, 1 + n, \frac{1}{2}(1 + i \tan(c + dx))\right) (a + ia \tan(c + dx))^n}{2dn}$$

```
output B*(a+I*a*tan(d*x+c))^n/d/n-1/2*(I*A+B)*hypergeom([1, n],[1+n],1/2+1/2*I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/n
```

3.222.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.77

$$\int (a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \frac{(2B - (iA + B) \operatorname{Hypergeometric2F1}\left(1, n, 1 + n, \frac{1}{2}(1 + i \tan(c + dx))\right)) (a + ia \tan(c + dx))^n}{2dn}$$

```
input Integrate[(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]
```

```
output ((2*B - (I*A + B)*Hypergeometric2F1[1, n, 1 + n, (1 + I*Tan[c + d*x])/2])* (a + I*a*Tan[c + d*x])^n)/(2*d*n)
```

3.222.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.08, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$, Rules used = {3042, 4010, 3042, 3962, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4010} \\
 & (A - iB) \int (i \tan(c + dx)a + a)^n dx + \frac{B(a + ia \tan(c + dx))^n}{dn} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \int (i \tan(c + dx)a + a)^n dx + \frac{B(a + ia \tan(c + dx))^n}{dn} \\
 & \quad \downarrow \text{3962} \\
 & \frac{B(a + ia \tan(c + dx))^n}{dn} - \frac{ia(A - iB) \int \frac{(i \tan(c + dx)a + a)^{n-1} d(ia \tan(c + dx))}{a - ia \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{78} \\
 & \frac{B(a + ia \tan(c + dx))^n}{dn} - \frac{i(A - iB)(a + ia \tan(c + dx))^n \operatorname{Hypergeometric2F1}\left(1, n, n + 1, \frac{i \tan(c + dx)a + a}{2a}\right)}{2dn}
 \end{aligned}$$

input `Int[(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `(B*(a + I*a*Tan[c + d*x])^n)/(d*n) - ((I/2)*(A - I*B)*Hypergeometric2F1[1, n, 1 + n, (a + I*a*Tan[c + d*x])/(2*a)]*(a + I*a*Tan[c + d*x])^n)/(d*n)`

3.222.3.1 Defintions of rubi rules used

rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3962 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[-b/d Subst[Int[(a + x)^(n - 1)/(a - x), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 + b^2, 0]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

3.222.4 Maple [F]

$$\int (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.222.5 Fracas [F]

$$\begin{aligned} & \int (a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n dx \end{aligned}$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(((A - I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*(2*a*e^(2*I*d*x + 2*I*c))/(e^(2*I*d*x + 2*I*c) + 1))^n/(e^(2*I*d*x + 2*I*c) + 1), x)`

3.222.6 Sympy [F]

$$\begin{aligned} & \int (a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (ia(\tan(c + dx) - i))^n (A + B \tan(c + dx)) dx \end{aligned}$$

input `integrate((a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x)), x)`

3.222.7 Maxima [F]

$$\begin{aligned} & \int (a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(i a \tan(dx + c) + a)^n dx \end{aligned}$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n, x)`

3.222.8 Giac [F]

$$\int (a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n dx$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n, x)`

3.222.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^n dx$$

input `int((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n,x)`

output `int((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n, x)`

3.223 $\int \cot(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.223.1 Optimal result	2223
3.223.2 Mathematica [A] (verified)	2223
3.223.3 Rubi [A] (verified)	2224
3.223.4 Maple [F]	2226
3.223.5 Fricas [F]	2226
3.223.6 Sympy [F]	2227
3.223.7 Maxima [F]	2227
3.223.8 Giac [F]	2227
3.223.9 Mupad [F(-1)]	2228

3.223.1 Optimal result

Integrand size = 32, antiderivative size = 97

$$\int \cot(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \frac{(A-iB) \operatorname{Hypergeometric2F1}\left(1, n, 1+n, \frac{1}{2}(1+i \tan(c+dx))\right) (a+ia \tan(c+dx))^n}{2dn}$$

$$- \frac{A \operatorname{Hypergeometric2F1}\left(1, n, 1+n, 1+i \tan(c+dx)\right) (a+ia \tan(c+dx))^n}{dn}$$

```
output 1/2*(A-I*B)*hypergeom([1, n],[1+n],1/2+1/2*I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/n-A*hypergeom([1, n],[1+n],1+I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/n
```

3.223.2 Mathematica [A] (verified)

Time = 1.22 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.28

$$\int \cot(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \frac{(a+ia \tan(c+dx))^n \left((iA+B)n \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{1}{2}(1+i \tan(c+dx))\right) \right) (-i + \tan(c+dx))}{4}$$

```
input Integrate[Cot[c + d*x]*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]
```

output $((a + I*a*\text{Tan}[c + d*x])^n*((I*A + B)*n*\text{Hypergeometric2F1}[1, 1 + n, 2 + n, (1 + I*\text{Tan}[c + d*x])/2]*(-I + \text{Tan}[c + d*x]) - (2*I)*(((-I)*A + B)*(1 + n) + 2*A*n*\text{Hypergeometric2F1}[1, 1 + n, 2 + n, 1 + I*\text{Tan}[c + d*x]])*(-I + \text{Tan}[c + d*x]))) / (4*d*n*(1 + n))$

3.223.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.06, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.219$, Rules used = {3042, 4083, 3042, 3962, 78, 4082, 75}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\tan(c + dx)} dx \\
 & \quad \downarrow 4083 \\
 & (B + iA) \int (i \tan(c + dx)a + a)^n dx + \frac{A \int \cot(c + dx)(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n dx}{a} \\
 & \quad \downarrow 3042 \\
 & (B + iA) \int (i \tan(c + dx)a + a)^n dx + \frac{A \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n}{\tan(c + dx)} dx}{a} \\
 & \quad \downarrow 3962 \\
 & \frac{A \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n}{\tan(c + dx)} dx}{a} - \frac{ia(B + iA) \int \frac{(i \tan(c + dx)a + a)^{n-1}}{a - ia \tan(c + dx)} d(i \tan(c + dx))}{d} \\
 & \quad \downarrow 78 \\
 & \frac{A \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n}{\tan(c + dx)} dx}{a} - \\
 & \frac{i(B + iA)(a + ia \tan(c + dx))^n \text{Hypergeometric2F1}\left(1, n, n + 1, \frac{i \tan(c + dx)a + a}{2a}\right)}{2dn} \\
 & \quad \downarrow 4082
 \end{aligned}$$

3.223. $\int \cot(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$

$$\frac{aA \int \cot(c+dx)(i \tan(c+dx)a+a)^{n-1} d \tan(c+dx)}{d} - \frac{i(B+iA)(a+ia \tan(c+dx))^n \operatorname{Hypergeometric2F1}\left(1, n, n+1, \frac{i \tan(c+dx)a+a}{2a}\right)}{2dn}$$

↓ 75

$$\frac{i(B+iA)(a+ia \tan(c+dx))^n \operatorname{Hypergeometric2F1}\left(1, n, n+1, \frac{i \tan(c+dx)a+a}{2a}\right)}{2dn} - \frac{A(a+ia \tan(c+dx))^n \operatorname{Hypergeometric2F1}(1, n, n+1, i \tan(c+dx)+1)}{dn}$$

input `Int[Cot[c + d*x]*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `-((A*Hypergeometric2F1[1, n, 1 + n, 1 + I*Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*n)) - ((I/2)*(I*A + B)*Hypergeometric2F1[1, n, 1 + n, (a + I*a*Tan[c + d*x])/(2*a)]*(a + I*a*Tan[c + d*x])^n)/(d*n)`

3.223.3.1 Defintions of rubi rules used

rule 75 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((c + d*x)^(n + 1)/(d*(n + 1)*(-d/(b*c))^m)*Hypergeometric2F1[-m, n + 1, n + 2, 1 + d*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[n] && (IntegerQ[m] || GtQ[-d/(b*c), 0])`

rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3962 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[-b/d Subst[Int[(a + x)^(n - 1)/(a - x), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 + b^2, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.223.4 Maple [F]

$$\int \cot(dx + c) (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(cot(d*x+c)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(cot(d*x+c)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.223.5 Fracas [F]

$$\begin{aligned} & \int \cot(c + dx) (a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A) (ia \tan(dx + c) + a)^n \cot(dx + c) dx \end{aligned}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A - B)*(2*a*e^(2*I*d*x + 2*I*c))/(e^(2*I*d*x + 2*I*c) + 1))^n/(e^(2*I*d*x + 2*I*c) - 1), x)`

3.223.6 Sympy [F]

$$\begin{aligned} & \int \cot(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (ia(\tan(c + dx) - i))^n (A + B \tan(c + dx)) \cot(c + dx) dx \end{aligned}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x))*cot(c + d*x), x)`

3.223.7 Maxima [F]

$$\begin{aligned} & \int \cot(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c) dx \end{aligned}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*cot(d*x + c), x)`

3.223.8 Giac [F]

$$\begin{aligned} & \int \cot(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c) dx \end{aligned}$$

input `integrate(cot(d*x+c)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*cot(d*x + c), x)`

3.223.9 Mupad [F(-1)]

Timed out.

$$\int \cot(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx) (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^n dx$$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n,x)`output `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n, x)`

3.224 $\int \cot^2(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.224.1 Optimal result	2229
3.224.2 Mathematica [A] (verified)	2230
3.224.3 Rubi [A] (verified)	2230
3.224.4 Maple [F]	2233
3.224.5 Fricas [F]	2233
3.224.6 Sympy [F]	2234
3.224.7 Maxima [F]	2234
3.224.8 Giac [F]	2234
3.224.9 Mupad [F(-1)]	2235

3.224.1 Optimal result

Integrand size = 34, antiderivative size = 131

$$\int \cot^2(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= -\frac{A \cot(c+dx)(a+ia \tan(c+dx))^n}{d}$$

$$+ \frac{(iA+B) \operatorname{Hypergeometric2F1}\left(1, n, 1+n, \frac{1}{2}(1+i \tan(c+dx))\right)(a+ia \tan(c+dx))^n}{2dn}$$

$$- \frac{(B+iAn) \operatorname{Hypergeometric2F1}\left(1, n, 1+n, 1+i \tan(c+dx)\right)(a+ia \tan(c+dx))^n}{dn}$$

```
output -A*cot(d*x+c)*(a+I*a*tan(d*x+c))^n/d+1/2*(I*A+B)*hypergeom([1, n],[1+n],1/2+1/2*I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/n-(B+I*A*n)*hypergeom([1, n],[1+n],1+I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/n
```

3.224.2 Mathematica [A] (verified)

Time = 2.92 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.19

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx =$$

$$\frac{(a + ia \tan(c + dx))^n (2(-iA + B)(1 + n) + 4Bn \operatorname{Hypergeometric2F1}(1, 1 + n, 2 + n, 1 + i \tan(c + dx)))}{d}$$

input `Integrate[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `-1/4*((a + I*a*Tan[c + d*x])^n*(2*((-I)*A + B)*(1 + n) + 4*B*n*Hypergeometric2F1[1, 1 + n, 2 + n, 1 + I*Tan[c + d*x]]*(1 + I*Tan[c + d*x]) + (A - I*B)*n*Hypergeometric2F1[1, 1 + n, 2 + n, (1 + I*Tan[c + d*x])/2]*(-I + Tan[c + d*x]) + 4*A*n*Hypergeometric2F1[2, 1 + n, 2 + n, 1 + I*Tan[c + d*x]]*(-I + Tan[c + d*x]))) / (d*n*(1 + n))`

3.224.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.10, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4081, 3042, 4083, 3042, 3962, 78, 4082, 75}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\tan(c + dx)^2} dx$$

$$\downarrow 4081$$

$$\frac{\int \cot(c + dx)(i \tan(c + dx)a + a)^n(a(B + iAn) - aA(1 - n) \tan(c + dx)) dx}{A \cot(c + dx) \frac{a}{d} + (a + ia \tan(c + dx))^n}$$

$$\downarrow 3042$$

$$\frac{\int \frac{(i \tan(c + dx)a + a)^n(a(B + iAn) - aA(1 - n) \tan(c + dx))}{\tan(c + dx)} dx}{a} - \frac{A \cot(c + dx)(a + ia \tan(c + dx))^n}{d}$$

3.224. $\int \cot^2(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$

↓ 4083

$$\frac{(B + iAn) \int \cot(c + dx)(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n dx - a(A - iB) \int (i \tan(c + dx)a + a)^n dx}{\frac{A \cot(c + dx)(a + ia \tan(c + dx))^n}{d}}$$

↓ 3042

$$\frac{(B + iAn) \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n}{\tan(c + dx)} dx - a(A - iB) \int (i \tan(c + dx)a + a)^n dx}{\frac{A \cot(c + dx)(a + ia \tan(c + dx))^n}{d}}$$

↓ 3962

$$\frac{ia^2(A - iB) \int \frac{(i \tan(c + dx)a + a)^{n-1}}{a - ia \tan(c + dx)} d(ia \tan(c + dx)) + (B + iAn) \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n}{\tan(c + dx)} dx}{\frac{A \cot(c + dx)(a + ia \tan(c + dx))^n}{d}}$$

↓ 78

$$\frac{(B + iAn) \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n}{\tan(c + dx)} dx + \frac{ia(A - iB)(a + ia \tan(c + dx))^n \operatorname{Hypergeometric2F1}\left(1, n, n + 1, \frac{i \tan(c + dx)a + a}{2a}\right)}{2dn}}{\frac{A \cot(c + dx)(a + ia \tan(c + dx))^n}{d}}$$

↓ 4082

$$\frac{a^2(B + iAn) \int \cot(c + dx)(i \tan(c + dx)a + a)^{n-1} d \tan(c + dx) + \frac{ia(A - iB)(a + ia \tan(c + dx))^n \operatorname{Hypergeometric2F1}\left(1, n, n + 1, \frac{i \tan(c + dx)a + a}{2a}\right)}{2dn}}{\frac{A \cot(c + dx)(a + ia \tan(c + dx))^n}{d}}$$

↓ 75

$$\frac{\frac{ia(A - iB)(a + ia \tan(c + dx))^n \operatorname{Hypergeometric2F1}\left(1, n, n + 1, \frac{i \tan(c + dx)a + a}{2a}\right)}{2dn} - \frac{a(B + iAn)(a + ia \tan(c + dx))^n \operatorname{Hypergeometric2F1}\left(1, n, n + 1, \frac{i \tan(c + dx)a + a}{2a}\right)}{dn}}{\frac{A \cot(c + dx)(a + ia \tan(c + dx))^n}{d}}$$

input `Int[Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output $-\left(\frac{A \cot(c + dx)(a + I a \tan(c + dx))^n}{d}\right) + \left(-\frac{(a(B + I A n) \operatorname{Hypergeometric2F1}[1, n, 1 + n, 1 + I \tan(c + dx)](a + I a \tan(c + dx))^n}{(d n)}\right) + \left(\frac{(I/2) a (A - I B) \operatorname{Hypergeometric2F1}[1, n, 1 + n, (a + I a \tan(c + dx))]}{(2 a)}\right) \frac{(a + I a \tan(c + dx))^n}{(d n)} / a$

3.224.3.1 Defintions of rubi rules used

rule 75 $\operatorname{Int}[(b \cdot x)^m ((c) + (d \cdot x)^n), x_Symbol] \rightarrow \operatorname{Simp}[(c + dx)^{n+1} / (d(n+1) (-d/(b \cdot c))^m) \operatorname{Hypergeometric2F1}[-m, n+1, n+2, 1 + d(x/c)], x] /;$ $\operatorname{FreeQ}\{b, c, d, m, n, x\} \ \&\& \ !\operatorname{IntegerQ}[n] \ \&\& \ (\operatorname{IntegerQ}[m] \ || \ \operatorname{GtQ}[-d/(b \cdot c), 0])$

rule 78 $\operatorname{Int}[(a) + (b \cdot x)^m ((c) + (d \cdot x)^n), x_Symbol] \rightarrow \operatorname{Simp}[(b \cdot c - a \cdot d)^n (a + b \cdot x)^{m+1} / (b^{n+1} (m+1)) \operatorname{Hypergeometric2F1}[-n, m+1, m+2, (-d)(a + b \cdot x)/(b \cdot c - a \cdot d)], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, m, x\} \ \&\& \ !\operatorname{IntegerQ}[m] \ \&\& \ \operatorname{IntegerQ}[n]$

rule 3042 $\operatorname{Int}[u, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /;$ $\operatorname{FunctionOfTrigOfLinearQ}[u, x]$

rule 3962 $\operatorname{Int}[(a) + (b \cdot x) \tan[(c) + (d \cdot x)], x_Symbol] \rightarrow \operatorname{Simp}[-b/d \operatorname{Subst}[\operatorname{Int}[(a + x)^{n-1} / (a - x), x], x, b \cdot \tan(c + dx)], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, n, x\} \ \&\& \ \operatorname{EqQ}[a^2 + b^2, 0]$

rule 4081 $\operatorname{Int}[(a) + (b \cdot x) \tan[(e) + (f \cdot x)]^m ((A) + (B \cdot x) \tan[(e) + (f \cdot x)]), x_Symbol] \rightarrow \operatorname{Simp}[(A \cdot d - B \cdot c) (a + b \cdot \tan[e + f \cdot x])^m ((c + d \cdot \tan[e + f \cdot x])^{n+1} / (f(n+1)(c^2 + d^2))), x] - \operatorname{Simp}[1 / (a(n+1)(c^2 + d^2)) \operatorname{Int}[(a + b \cdot \tan[e + f \cdot x])^m (c + d \cdot \tan[e + f \cdot x])^{n+1} \operatorname{Simp}[A(b \cdot d \cdot m - a \cdot c(n+1)) - B(b \cdot c \cdot m + a \cdot d(n+1)) - a(B \cdot c - A \cdot d)(m + n + 1) \tan[e + f \cdot x], x], x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, e, f, A, B, m, x\} \ \&\& \ \operatorname{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \operatorname{EqQ}[a^2 + b^2, 0] \ \&\& \ \operatorname{LtQ}[n, -1]$

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int((((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]))/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.224.4 Maple [F]

$$\int (\cot^2(dx + c)) (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.224.5 Fracas [F]

$$\begin{aligned} & \int \cot^2(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^2 dx \end{aligned}$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(-(A - I*B)*e^(4*I*d*x + 4*I*c) + 2*A*e^(2*I*d*x + 2*I*c) + A + I*B)*(2*a*e^(2*I*d*x + 2*I*c)/(e^(2*I*d*x + 2*I*c) + 1))^n/(e^(4*I*d*x + 4*I*c) - 2*e^(2*I*d*x + 2*I*c) + 1), x)`

3.224.6 Sympy [F]

$$\begin{aligned} & \int \cot^2(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (ia(\tan(c + dx) - i))^n(A + B \tan(c + dx)) \cot^2(c + dx) dx \end{aligned}$$

input `integrate(cot(d*x+c)**2*(a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x))*cot(c + d*x)**2, x)`

3.224.7 Maxima [F]

$$\begin{aligned} & \int \cot^2(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^2 dx \end{aligned}$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*cot(d*x + c)^2, x)`

3.224.8 Giac [F]

$$\begin{aligned} & \int \cot^2(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^2 dx \end{aligned}$$

input `integrate(cot(d*x+c)^2*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*cot(d*x + c)^2, x)`

3.224.9 Mupad [F(-1)]

Timed out.

$$\int \cot^2(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^2 (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^n dx$$

input `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n,x)`output `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n, x)`

3.225 $\int \cot^3(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.225.1 Optimal result	2236
3.225.2 Mathematica [A] (verified)	2237
3.225.3 Rubi [A] (verified)	2237
3.225.4 Maple [F]	2240
3.225.5 Fricas [F]	2241
3.225.6 Sympy [F]	2241
3.225.7 Maxima [F]	2241
3.225.8 Giac [F]	2242
3.225.9 Mupad [F(-1)]	2242

3.225.1 Optimal result

Integrand size = 34, antiderivative size = 185

$$\int \cot^3(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= -\frac{(2B+iAn) \cot(c+dx)(a+ia \tan(c+dx))^n}{2d} - \frac{A \cot^2(c+dx)(a+ia \tan(c+dx))^n}{2d}$$

$$- \frac{(A-iB) \operatorname{Hypergeometric2F1}\left(1, n, 1+n, \frac{1}{2}(1+i \tan(c+dx))\right)(a+ia \tan(c+dx))^n}{2dn}$$

$$- \frac{(2iBn-A(2-n+n^2)) \operatorname{Hypergeometric2F1}\left(1, n, 1+n, 1+i \tan(c+dx)\right)(a+ia \tan(c+dx))^n}{2dn}$$

```
output -1/2*(2*B+I*A*n)*cot(d*x+c)*(a+I*a*tan(d*x+c))^n/d-1/2*A*cot(d*x+c)^2*(a+I
*a*tan(d*x+c))^n/d-1/2*(A-I*B)*hypergeom([1, n],[1+n],1/2+1/2*I*tan(d*x+c)
)*(a+I*a*tan(d*x+c))^n/d/n-1/2*(2*I*B*n-A*(n^2-n+2))*hypergeom([1, n],[1+n
],1+I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/n
```

3.225.2 Mathematica [A] (verified)

Time = 3.17 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.19

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \frac{(a + ia \tan(c + dx))^n \left(-i(A - iB)n \operatorname{Hypergeometric2F1} \left(1, 1 + n, 2 + n, \frac{1}{2}(1 + i \tan(c + dx)) \right) \right) (-i + \tan(c + dx))}{4d(1 + n)}$$

input `Integrate[Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `((a + I*a*Tan[c + d*x])^n*((-I)*(A - I*B)*n*Hypergeometric2F1[1, 1 + n, 2 + n, (1 + I*Tan[c + d*x])/2]*(-I + Tan[c + d*x]) + 2*(A + I*B + A*n + I*B*n + 2*A*n*Hypergeometric2F1[3, 1 + n, 2 + n, 1 + I*Tan[c + d*x]]) + 2*A*n*Hypergeometric2F1[1, 1 + n, 2 + n, 1 + I*Tan[c + d*x]]*(1 + I*Tan[c + d*x]) + (2*I)*A*n*Hypergeometric2F1[3, 1 + n, 2 + n, 1 + I*Tan[c + d*x]]*Tan[c + d*x] - 2*B*n*Hypergeometric2F1[2, 1 + n, 2 + n, 1 + I*Tan[c + d*x]]*(-I + Tan[c + d*x])))/(4*d*n*(1 + n))`

3.225.3 Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.11, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.324$, Rules used = {3042, 4081, 3042, 4081, 3042, 4083, 3042, 3962, 78, 4082, 75}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\tan(c + dx)^3} dx$$

$$\downarrow \text{4081}$$

$$\int \frac{\cot^2(c + dx)(i \tan(c + dx)a + a)^n(a(2B + iAn) - aA(2 - n) \tan(c + dx)) dx}{2a}$$

$$\frac{A \cot^2(c + dx)(a + ia \tan(c + dx))^n}{2d}$$

$$\downarrow \text{3042}$$

$$\frac{\int \frac{(i \tan(c+dx)a+a)^n (a(2B+iAn)-aA(2-n) \tan(c+dx))}{\tan(c+dx)^2} dx}{2a} - \frac{A \cot^2(c+dx)(a+ia \tan(c+dx))^n}{2d}$$

↓ 4081

$$\frac{\int \cot(c+dx)(i \tan(c+dx)a+a)^n (a^2(2iBn-A(n^2-n+2))-a^2(1-n)(2B+iAn) \tan(c+dx)) dx}{a} - \frac{a(2B+iAn) \cot(c+dx)(a+ia \tan(c+dx))^n}{d}$$

$$\frac{A \cot^2(c+dx)(a+ia \tan(c+dx))^n}{2d}$$

↓ 3042

$$\frac{\int \frac{(i \tan(c+dx)a+a)^n (a^2(2iBn-A(n^2-n+2))-a^2(1-n)(2B+iAn) \tan(c+dx))}{\tan(c+dx)} dx}{a} - \frac{a(2B+iAn) \cot(c+dx)(a+ia \tan(c+dx))^n}{d}$$

$$\frac{A \cot^2(c+dx)(a+ia \tan(c+dx))^n}{2d}$$

↓ 4083

$$\frac{a(-A(n^2-n+2)+2iBn) \int \cot(c+dx)(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n dx - 2a^2(B+iA) \int (i \tan(c+dx)a+a)^n dx}{a} - \frac{a(2B+iAn) \cot(c+dx)(a+ia \tan(c+dx))^n}{d}$$

$$\frac{A \cot^2(c+dx)(a+ia \tan(c+dx))^n}{2d}$$

↓ 3042

$$\frac{a(-A(n^2-n+2)+2iBn) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\tan(c+dx)} dx - 2a^2(B+iA) \int (i \tan(c+dx)a+a)^n dx}{a} - \frac{a(2B+iAn) \cot(c+dx)(a+ia \tan(c+dx))^n}{d}$$

$$\frac{A \cot^2(c+dx)(a+ia \tan(c+dx))^n}{2d}$$

↓ 3962

$$\frac{2ia^3(B+iA) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{a-ia \tan(c+dx)} d(ia \tan(c+dx)) + a(-A(n^2-n+2)+2iBn) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\tan(c+dx)} dx}{a} - \frac{a(2B+iAn) \cot(c+dx)(a+ia \tan(c+dx))^n}{d}$$

$$\frac{A \cot^2(c+dx)(a+ia \tan(c+dx))^n}{2d}$$

↓ 78

$$\frac{a(-A(n^2-n+2)+2iBn) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\tan(c+dx)} dx + \frac{ia^2(B+iA)(a+ia \tan(c+dx))^n \text{Hypergeometric2F1}\left(1, n, n+1, \frac{i \tan(c+dx)a+a}{2a}\right)}{dn}}{a} - \frac{a(2B+iAn) \cot(c+dx)(a+ia \tan(c+dx))^n}{d}$$

$$\frac{A \cot^2(c+dx)(a+ia \tan(c+dx))^n}{2d}$$

3.225. $\int \cot^3(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

↓ 4082

$$\frac{a^3(-A(n^2-n+2)+2iBn) \int \cot(c+dx)(i \tan(c+dx)a+a)^{n-1} d \tan(c+dx)}{d} + \frac{ia^2(B+iA)(a+ia \tan(c+dx))^n \text{Hypergeometric2F1}\left(1, n, n+1, \frac{i \tan(c+dx)a+a}{2a}\right)}{dn} - \frac{a}{2a}$$

$$\frac{A \cot^2(c+dx)(a+ia \tan(c+dx))^n}{2d}$$

↓ 75

$$\frac{ia^2(B+iA)(a+ia \tan(c+dx))^n \text{Hypergeometric2F1}\left(1, n, n+1, \frac{i \tan(c+dx)a+a}{2a}\right)}{dn} - \frac{a^2(-A(n^2-n+2)+2iBn)(a+ia \tan(c+dx))^n \text{Hypergeometric2F1}(1, n, n+1, i \tan(c+dx))}{a dn} - \frac{a}{2a}$$

$$\frac{A \cot^2(c+dx)(a+ia \tan(c+dx))^n}{2d}$$

```
input Int[Cot[c + d*x]^3*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]
```

```
output -1/2*(A*Cot[c + d*x]^2*(a + I*a*Tan[c + d*x])^n)/d + (-((a*(2*B + I*A*n)*Cot[c + d*x]*(a + I*a*Tan[c + d*x])^n)/d) + (-((a^2*((2*I)*B*n - A*(2 - n + n^2))*Hypergeometric2F1[1, n, 1 + n, 1 + I*Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*n)) + (I*a^2*(I*A + B)*Hypergeometric2F1[1, n, 1 + n, (a + I*a*Tan[c + d*x])/(2*a)]*(a + I*a*Tan[c + d*x])^n)/(d*n))/a)/(2*a)
```

3.225.3.1 Defintions of rubi rules used

```
rule 75 Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((c + d*x)^(n + 1)/(d*(n + 1)*(-d/(b*c))^m)*Hypergeometric2F1[-m, n + 1, n + 2, 1 + d*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[n] && (IntegerQ[m] || GtQ[-d/(b*c), 0])
```

```
rule 78 Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

3.225. $\int \cot^3(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

rule 3962 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[-b/d Subst[Int[(a + x)^(n - 1)/(a - x), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[a^2 + b^2, 0]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4083 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(A*b + a*B)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m, x], x] - Simp[(B*c - A*d)/(b*c + a*d) Int[(a + b*Tan[e + f*x])^m*((a - b*Tan[e + f*x])/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.225.4 Maple [F]

$$\int (\cot^3(dx + c)) (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.225.5 Fricas [F]

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^3 dx$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(((-I*A - B)*e^(6*I*d*x + 6*I*c) + (-3*I*A - B)*e^(4*I*d*x + 4*I*c) + (-3*I*A + B)*e^(2*I*d*x + 2*I*c) - I*A + B)*(2*a*e^(2*I*d*x + 2*I*c)/(e^(2*I*d*x + 2*I*c) + 1))^n/(e^(6*I*d*x + 6*I*c) - 3*e^(4*I*d*x + 4*I*c) + 3*e^(2*I*d*x + 2*I*c) - 1), x)`

3.225.6 Sympy [F]

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (ia(\tan(c + dx) - i))^n(A + B \tan(c + dx)) \cot^3(c + dx) dx$$

input `integrate(cot(d*x+c)**3*(a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x))*cot(c + d*x)**3, x)`

3.225.7 Maxima [F]

$$\int \cot^3(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^3 dx$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*cot(d*x + c)^3, x)`

3.225.8 Giac [F]

$$\begin{aligned} & \int \cot^3(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^3 dx \end{aligned}$$

input `integrate(cot(d*x+c)^3*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*cot(d*x + c)^3, x)`

3.225.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \cot^3(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int \cot(c + dx)^3 (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^n dx \end{aligned}$$

input `int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n,x)`

output `int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n, x)`

3.226 $\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.226.1 Optimal result	2243
3.226.2 Mathematica [F]	2244
3.226.3 Rubi [A] (warning: unable to verify)	2244
3.226.4 Maple [F]	2250
3.226.5 Fricas [F]	2251
3.226.6 Sympy [F(-1)]	2251
3.226.7 Maxima [F]	2251
3.226.8 Giac [F]	2252
3.226.9 Mupad [F(-1)]	2252

3.226.1 Optimal result

Integrand size = 36, antiderivative size = 383

$$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= -\frac{2(2iAn(5+2n)+B(15+10n+4n^2))\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(1+2n)(3+2n)(5+2n)}$$

$$+ \frac{2(iA+B) \operatorname{AppellF1}\left(\frac{1}{2}, 1-n, 1, \frac{3}{2}, -i \tan(c+dx), i \tan(c+dx)\right) (1+i \tan(c+dx))^{-n} \sqrt{\tan(c+dx)}}{d}$$

$$- \frac{2(4Bn(9+8n+2n^2)+iA(15+36n+32n^2+8n^3)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1-n, \frac{3}{2}, -i \tan(c+dx)\right)}{d(1+2n)(3+2n)(5+2n)}$$

$$- \frac{2(2iBn-A(5+2n)) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(3+2n)(5+2n)}$$

$$+ \frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(5+2n)}$$

output

```
-2*(2*I*A*n*(5+2*n)+B*(4*n^2+10*n+15))*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))
^n/d/(5+2*n)/(4*n^2+8*n+3)+2*(I*A+B)*AppellF1(1/2,1-n,1,3/2,-I*tan(d*x+c),
I*tan(d*x+c))*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n/d/((1+I*tan(d*x+c))^n)
-2*(4*B*n*(2*n^2+8*n+9)+I*A*(8*n^3+32*n^2+36*n+15))*hypergeom([1/2, 1-n],[
3/2],-I*tan(d*x+c))*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n/d/(5+2*n)/(4*n^2
+8*n+3)/((1+I*tan(d*x+c))^n)-2*(2*I*B*n-A*(5+2*n))*tan(d*x+c)^(3/2)*(a+I*a
*tan(d*x+c))^n/d/(3+2*n)/(5+2*n)+2*B*tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^n
/d/(5+2*n)
```

3.226.2 Mathematica [F]

$$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

input `Integrate[Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

output `Integrate[Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

3.226.3 Rubi [A] (warning: unable to verify)

Time = 2.06 (sec) , antiderivative size = 421, normalized size of antiderivative = 1.10, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.611$, Rules used = {3042, 4080, 27, 3042, 4080, 27, 3042, 4080, 27, 3042, 4084, 3042, 4047, 25, 27, 148, 27, 334, 333, 4082, 76, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c+dx)^{5/2}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4080}$$

$$\frac{2 \int -\frac{1}{2} \tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a)^n(5aB+a(2iBn-A(2n+5)) \tan(c+dx)) dx}{a(2n+5)} +$$

$$\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)}$$

$$\downarrow \text{27}$$

$$\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{\int \tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a)^n(5aB+a(2iBn-A(2n+5)) \tan(c+dx))dx}{a(2n+5)}$$

↓ 3042

$$\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{\int \tan(c+dx)^{3/2}(i \tan(c+dx)a+a)^n(5aB+a(2iBn-A(2n+5)) \tan(c+dx))dx}{a(2n+5)}$$

↓ 4080

$$\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2 \int -\frac{1}{2} \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^n(3a^2(2iBn-A(2n+5))-a^2(2iAn(2n+5)+B(4n^2+10n+15)) \tan(c+dx))dx}{a(2n+3)} + \frac{2a(-A(2n+5)+2iBn) \tan(c+dx)}{d(2n+3)}$$

a(2n+5)

↓ 27

$$\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{\int \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^n(3a^2(2iBn-A(2n+5))-a^2(2iAn(2n+5)+B(4n^2+10n+15)) \tan(c+dx))dx}{a(2n+3)}$$

a(2n+5)

↓ 3042

$$\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{\int \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^n(3a^2(2iBn-A(2n+5))-a^2(2iAn(2n+5)+B(4n^2+10n+15)) \tan(c+dx))dx}{a(2n+3)}$$

a(2n+5)

↓ 4080

$$\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2 \int \frac{(i \tan(c+dx)a+a)^n((2iAn(2n+5)+B(4n^2+10n+15))a^3+(4iBn(2n^2+8n+9))-A(8n^3+32n^2+24n+8)) \sqrt{\tan(c+dx)}}{a(2n+1)}dx}{a(2n+1)}$$

a(2n+5)

↓ 27

3.226. $\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

$$\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{\int \frac{(i \tan(c+dx)a+a)^n ((2iAn(2n+5)+B(4n^2+10n+15))a^3+(4iBn(2n^2+8n+9)-A(8n^3+32n^2))\sqrt{\tan(c+dx)}}{a(2n+1)} dx}{a(2n+5)}$$

3042

$$\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{\int \frac{(i \tan(c+dx)a+a)^n ((2iAn(2n+5)+B(4n^2+10n+15))a^3+(4iBn(2n^2+8n+9)-A(8n^3+32n^2))\sqrt{\tan(c+dx)}}{a(2n+1)} dx}{a(2n+5)}$$

4084

$$\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{a^3(8n^3+36n^2+46n+15)(B+iA) \int \frac{(i \tan(c+dx)a+a)^n dx - a^2(4Bn(2n^2+8n+9)+iA(8n^3+32n^2))\sqrt{\tan(c+dx)}}{a(2n+1)} dx}{a(2n+5)}$$

3042

$$\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{a^3(8n^3+36n^2+46n+15)(B+iA) \int \frac{(i \tan(c+dx)a+a)^n dx - a^2(4Bn(2n^2+8n+9)+iA(8n^3+32n^2))\sqrt{\tan(c+dx)}}{a(2n+1)} dx}{a(2n+5)}$$

4047

$$\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{ia^5(8n^3+36n^2+46n+15)(B+iA) \int \frac{(i \tan(c+dx)a+a)^{n-1} d(ia \tan(c+dx))}{a\sqrt{\tan(c+dx)}(a-ia \tan(c+dx))} dx - a^2(4Bn)}{a(2n+5)}$$

25

$$\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{ia^5(8n^3+36n^2+46n+15)(B+iA) \int \frac{(i \tan(c+dx)a+a)^{n-1} d(ia \tan(c+dx))}{a\sqrt{\tan(c+dx)}(a-ia \tan(c+dx))} dx - a^2(4Bn)}{a(2n+5)}$$

3.226. $\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

$$\begin{aligned} & \downarrow 27 \\ & \frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{ia^4(8n^3+36n^2+46n+15)(B+iA) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)(a-ia \tan(c+dx))}} d(ia \tan(c+dx))}{d} - a^2(4Bn) \\ & \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{a^2(4Bn)}{a(2n+1)} \end{aligned}$$

$$\begin{aligned} & \downarrow 148 \\ & \frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a^5(8n^3+36n^2+46n+15)(B+iA) \int \frac{(a-ia^3 \tan^2(c+dx))^{n-1}}{a(ia^2 \tan^2(c+dx)+1)} d\sqrt{\tan(c+dx)}}{d} - a^2(4Bn(2n^2+8n)) \\ & \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{a^2(4Bn(2n^2+8n))}{a(2n+1)} \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a^4(8n^3+36n^2+46n+15)(B+iA) \int \frac{(a-ia^3 \tan^2(c+dx))^{n-1}}{ia^2 \tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} - a^2(4Bn(2n^2+8n)) \\ & \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{a^2(4Bn(2n^2+8n))}{a(2n+1)} \end{aligned}$$

$$\begin{aligned} & \downarrow 334 \\ & \frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a^3(8n^3+36n^2+46n+15)(B+iA)(1-ia^2 \tan^2(c+dx))^{-n}(a-ia^3 \tan^2(c+dx))^n \int \frac{(1-ia^2 \tan^2(c+dx))}{ia^2 \tan^2(c+dx)}}{d} \\ & \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{a^2(4Bn(2n^2+8n))}{a(2n+1)} \end{aligned}$$

$$\begin{aligned} & \downarrow 333 \\ & \frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2ia^4(8n^3+36n^2+46n+15)(B+iA) \tan(c+dx)(1-ia^2 \tan^2(c+dx))^{-n}(a-ia^3 \tan^2(c+dx))^n \int \frac{(1-ia^2 \tan^2(c+dx))}{ia^2 \tan^2(c+dx)}}{d} \\ & \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{a^2(4Bn(2n^2+8n))}{a(2n+1)} \end{aligned}$$

$$\downarrow 4082$$

3.226. $\int \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

$$\frac{2B \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n}{d(2n + 5)} - \frac{2ia^4(8n^3 + 36n^2 + 46n + 15)(B + ia) \tan(c + dx)(1 - ia^2 \tan^2(c + dx))^{-n}(a - ia^3 \tan^2(c + dx))^n}{d}$$

$$\frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2ia^4(8n^3+36n^2+46n+15)(B+ia) \tan(c+dx)(1-ia^2 \tan^2(c+dx))^{-n}(a-ia^3 \tan^2(c+dx))^n}{d}$$

↓ 76

$$\frac{2B \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n}{d(2n + 5)} - \frac{2ia^4(8n^3 + 36n^2 + 46n + 15)(B + ia) \tan(c + dx)(1 - ia^2 \tan^2(c + dx))^{-n}(a - ia^3 \tan^2(c + dx))^n}{d}$$

$$\frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2ia^4(8n^3+36n^2+46n+15)(B+ia) \tan(c+dx)(1-ia^2 \tan^2(c+dx))^{-n}(a-ia^3 \tan^2(c+dx))^n}{d}$$

↓ 74

$$\frac{2B \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n}{d(2n + 5)} - \frac{2ia^4(8n^3 + 36n^2 + 46n + 15)(B + ia) \tan(c + dx)(1 - ia^2 \tan^2(c + dx))^{-n}(a - ia^3 \tan^2(c + dx))^n}{d}$$

$$\frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2ia^4(8n^3+36n^2+46n+15)(B+ia) \tan(c+dx)(1-ia^2 \tan^2(c+dx))^{-n}(a-ia^3 \tan^2(c+dx))^n}{d}$$

input `Int[Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `(2*B*Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^n)/(d*(5 + 2*n)) - ((2*a*((2*I)*B*n - A*(5 + 2*n))*Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^n)/(d*(3 + 2*n)) - ((-2*a^2*((2*I)*A*n*(5 + 2*n) + B*(15 + 10*n + 4*n^2))*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + 2*n)) + ((-2*a^3*(4*B*n*(9 + 8*n + 2*n^2) + I*A*(15 + 36*n + 32*n^2 + 8*n^3))*Hypergeometric2F1[1/2, 1 - n, 3/2, (-I)*Tan[c + d*x]]*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + I*Tan[c + d*x])^n) + ((2*I)*a^4*(I*A + B)*(15 + 46*n + 36*n^2 + 8*n^3)*AppellF1[1/2, 1, 1 - n, 3/2, (-I)*a^2*Tan[c + d*x]^2, I*a^2*Tan[c + d*x]^2]*Tan[c + d*x]*(a - I*a^3*Tan[c + d*x]^2)^n)/(d*(1 - I*a^2*Tan[c + d*x]^2)^n))/(a*(1 + 2*n))/(a*(3 + 2*n))/(a*(5 + 2*n))`

3.226.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 74 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)]) && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0]))`
- rule 76 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0] && ((RationalQ[m] && !(EqQ[n, -2^(-1)]) && EqQ[c^2 - d^2, 0])) || !RationalQ[n]`
- rule 148 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_.)*((e_) + (f_.)*(x_))^(p_.), x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c + d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x]] /; FreeQ[{b, c, d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`
- rule 333 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`
- rule 334 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4047 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4080 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.226.4 Maple [F]

$$\int \left(\tan^{\frac{5}{2}}(dx + c) \right) (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.226. $\int \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$

3.226.5 Fracas [F]

$$\int \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^{\frac{5}{2}} dx$$

input `integrate(tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(-((A - I*B)*e^(6*I*d*x + 6*I*c) - (A - 3*I*B)*e^(4*I*d*x + 4*I*c) - (A + 3*I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*(2*a*e^(2*I*d*x + 2*I*c)/(e^(2*I*d*x + 2*I*c) + 1))^n*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))/(e^(6*I*d*x + 6*I*c) + 3*e^(4*I*d*x + 4*I*c) + 3*e^(2*I*d*x + 2*I*c) + 1), x)`

3.226.6 Sympy [F(-1)]

Timed out.

$$\int \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(5/2)*(a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.226.7 Maxima [F]

$$\int \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^{\frac{5}{2}} dx$$

input `integrate(tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*tan(d*x + c)^(5/2), x)`

3.226.8 Giac [F]

$$\begin{aligned} & \int \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate(tan(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*tan(d*x + c)^(5/2), x)`

3.226.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \tan^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int \tan(c + dx)^{5/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) i)^n dx \end{aligned}$$

input `int(tan(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n,x)`

output `int(tan(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n, x)`

3.227 $\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.227.1 Optimal result	2253
3.227.2 Mathematica [F]	2254
3.227.3 Rubi [A] (warning: unable to verify)	2254
3.227.4 Maple [F]	2260
3.227.5 Fricas [F]	2260
3.227.6 Sympy [F(-1)]	2260
3.227.7 Maxima [F]	2261
3.227.8 Giac [F]	2261
3.227.9 Mupad [F(-1)]	2261

3.227.1 Optimal result

Integrand size = 36, antiderivative size = 291

$$\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= -\frac{2(2iBn - A(3 + 2n))\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(1+2n)(3+2n)}$$

$$- \frac{2(A-iB) \operatorname{AppellF1}\left(\frac{1}{2}, 1-n, 1, \frac{3}{2}, -i \tan(c+dx), i \tan(c+dx)\right) (1+i \tan(c+dx))^{-n} \sqrt{\tan(c+dx)}}{d}$$

$$+ \frac{2(2An(3+2n) - iB(3+6n+4n^2)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1-n, \frac{3}{2}, -i \tan(c+dx)\right) (1+i \tan(c+dx))^{-n}}{d(1+2n)(3+2n)}$$

$$+ \frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(3+2n)}$$

```
output -2*(2*I*B*n-A*(3+2*n))*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n/d/(4*n^2+8*n+
3)-2*(A-I*B)*AppellF1(1/2,1-n,1,3/2,-I*tan(d*x+c),I*tan(d*x+c))*tan(d*x+c)
^(1/2)*(a+I*a*tan(d*x+c))^n/d/((1+I*tan(d*x+c))^n)+2*(2*A*n*(3+2*n)-I*B*(4
*n^2+6*n+3))*hypergeom([1/2, 1-n], [3/2], -I*tan(d*x+c))*tan(d*x+c)^(1/2)*(a
+I*a*tan(d*x+c))^n/d/(4*n^2+8*n+3)/((1+I*tan(d*x+c))^n)+2*B*tan(d*x+c)^(3/
2)*(a+I*a*tan(d*x+c))^n/d/(3+2*n)
```

3.227.2 Mathematica [F]

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

input `Integrate[Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

output `Integrate[Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

3.227.3 Rubi [A] (warning: unable to verify)

Time = 1.46 (sec) , antiderivative size = 330, normalized size of antiderivative = 1.13, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.528$, Rules used = {3042, 4080, 27, 3042, 4080, 27, 3042, 4084, 3042, 4047, 25, 27, 148, 27, 334, 333, 4082, 76, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^{3/2}(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4080}$$

$$2 \int \frac{-\frac{1}{2} \sqrt{\tan(c + dx)}(i \tan(c + dx)a + a)^n(3aB + a(2iBn - A(2n + 3)) \tan(c + dx)) dx}{a(2n + 3)} +$$

$$\frac{2B \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n}{d(2n + 3)}$$

$$\downarrow \text{27}$$

$$\begin{aligned}
 & \frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \\
 & \frac{\int \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^n(3aB+a(2iBn-A(2n+3)) \tan(c+dx))dx}{a(2n+3)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \\
 & \frac{\int \sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^n(3aB+a(2iBn-A(2n+3)) \tan(c+dx))dx}{a(2n+3)} \\
 & \quad \downarrow \text{4080} \\
 & \frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \\
 & \frac{2 \int -\frac{(i \tan(c+dx)a+a)^n(a^2(2iBn-A(2n+3))-a^2(2iAn(2n+3)+B(4n^2+6n+3)) \tan(c+dx))}{2\sqrt{\tan(c+dx)}} dx}{a(2n+1)} + \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))}{d(2n+1)} \\
 & \quad \downarrow \text{27} \\
 & \frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \\
 & \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{\int \frac{(i \tan(c+dx)a+a)^n(a^2(2iBn-A(2n+3))-a^2(2iAn(2n+3)+B(4n^2+6n+3)) \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \\
 & \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{\int \frac{(i \tan(c+dx)a+a)^n(a^2(2iBn-A(2n+3))-a^2(2iAn(2n+3)+B(4n^2+6n+3)) \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)} \\
 & \quad \downarrow \text{4084} \\
 & \frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \\
 & \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{a(2An(2n+3)-iB(4n^2+6n+3)) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - a^2(4n^2+8n+)}{a(2n+1)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.227. $\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{a(2An(2n+3)-iB(4n^2+6n+3)) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - a^2(4n^2+8n+3)}{a(2n+1)}$$

4047

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{a(2An(2n+3)-iB(4n^2+6n+3)) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - \frac{ia^4(4n^2+8n+3)}{a(2n+1)}}{a(2n+3)}$$

25

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{ia^4(4n^2+8n+3)(A-iB) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{a\sqrt{\tan(c+dx)}(a-ia \tan(c+dx))} d(ia \tan(c+dx))}{d} + a(2An(2n+3)-iB)}{a(2n+1)}$$

27

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{ia^3(4n^2+8n+3)(A-iB) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)}(a-ia \tan(c+dx))} d(ia \tan(c+dx))}{d} + a(2An(2n+3)-iB)}{a(2n+1)}$$

148

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{a(2An(2n+3)-iB(4n^2+6n+3)) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - \frac{2a^4(4n^2+8n+3)}{a(2n+1)}}{a(2n+3)}$$

27

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{a(2An(2n+3)-iB(4n^2+6n+3)) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(4n^2+8n+3)}{a(2n+1)}}{a(2n+3)}$$

3.227. $\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

↓ 334

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{a(2An(2n+3)-iB(4n^2+6n+3)) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(4n^2+8n+3)}{a(2n+3)}}{a(2n+3)}$$

↓ 333

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{a(2An(2n+3)-iB(4n^2+6n+3)) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - \frac{2ia^3(4n^2+8n+3)}{a(2n+3)}}{a(2n+3)}$$

↓ 4082

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{a^3(2An(2n+3)-iB(4n^2+6n+3)) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)}} d \tan(c+dx) - \frac{2ia^3(4n^2+8n+3)(A-)}{a(2n+3)}}{a(2n+3)}$$

↓ 76

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{a^2(2An(2n+3)-iB(4n^2+6n+3))(1+i \tan(c+dx))^{-n}(a+ia \tan(c+dx))^n \int \frac{(i \tan(c+dx)+1)^{n-1}}{\sqrt{\tan(c+dx)}}}{a(2n+3)}$$

↓ 74

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{2a^2(2An(2n+3)-iB(4n^2+6n+3))\sqrt{\tan(c+dx)}(1+i \tan(c+dx))^{-n}(a+ia \tan(c+dx))^n \text{Hypergeometric}}{a(2n+3)}$$

input `Int[Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

3.227. $\int \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$


```
output (2*B*Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^n)/(d*(3 + 2*n)) - ((2*a*((
2*I)*B*n - A*(3 + 2*n))*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1
+ 2*n)) - ((2*a^2*(2*A*n*(3 + 2*n) - I*B*(3 + 6*n + 4*n^2))*Hypergeometri
c2F1[1/2, 1 - n, 3/2, (-I)*Tan[c + d*x]]*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c
+ d*x])^n)/(d*(1 + I*Tan[c + d*x])^n) - ((2*I)*a^3*(A - I*B)*(3 + 8*n + 4
*n^2)*AppellF1[1/2, 1, 1 - n, 3/2, (-I)*a^2*Tan[c + d*x]^2, I*a^2*Tan[c +
d*x]^2]*Tan[c + d*x]*(a - I*a^3*Tan[c + d*x]^2)^n)/(d*(1 - I*a^2*Tan[c + d
*x]^2)^n))/(a*(1 + 2*n)))/(a*(3 + 2*n))
```

3.227.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 74 Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x
)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x]
/; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0]
&& !(EqQ[n, -2^(-1)]) && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0]))
```

```
rule 76 Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[c^IntPart
[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*
(x/c))^n, x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !Integer
Q[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0] && ((RationalQ[m] && !(EqQ[n, -2
^(-1)]) && EqQ[c^2 - d^2, 0])) || !RationalQ[n]
```

```
rule 148 Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_),
x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c
+ d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x] /; FreeQ[{b, c,
d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]
```

```
rule 333 Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Sim
p[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; F
reeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a,
0]) && (IntegerQ[q] || GtQ[c, 0])
```

- rule 334 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4047 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4080 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`
- rule 4082 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`
- rule 4084 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.227.4 Maple [F]

$$\int \left(\tan^{\frac{3}{2}}(dx + c) \right) (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.227.5 Fracas [F]

$$\begin{aligned} & \int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output `integral(((−I*A − B)*e^(4*I*d*x + 4*I*c) + 2*B*e^(2*I*d*x + 2*I*c) + I*A − B)*(2*a*e^(2*I*d*x + 2*I*c)/(e^(2*I*d*x + 2*I*c) + 1))^n*sqrt((−I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))/(e^(4*I*d*x + 4*I*c) + 2*e^(2*I*d*x + 2*I*c) + 1), x)`

3.227.6 Sympy [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(3/2)*(a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.227.7 Maxima [F]

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^{\frac{3}{2}} dx$$

input `integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*tan(d*x + c)^(3/2), x)`

3.227.8 Giac [F]

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \tan(dx + c)^{\frac{3}{2}} dx$$

input `integrate(tan(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*tan(d*x + c)^(3/2), x)`

3.227.9 Mupad [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^{3/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) i)^n dx$$

input `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n,x)`

output `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n, x)`

3.227. $\int \tan^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$

3.228 $\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$

3.228.1 Optimal result	2262
3.228.2 Mathematica [F]	2263
3.228.3 Rubi [A] (warning: unable to verify)	2263
3.228.4 Maple [F]	2268
3.228.5 Fracas [F]	2268
3.228.6 Sympy [F]	2269
3.228.7 Maxima [F]	2269
3.228.8 Giac [F]	2269
3.228.9 Mupad [F(-1)]	2270

3.228.1 Optimal result

Integrand size = 36, antiderivative size = 215

$$\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \frac{2B\sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^n}{d(1 + 2n)}$$

$$- \frac{2(iA + B) \operatorname{AppellF1}\left(\frac{1}{2}, 1 - n, 1, \frac{3}{2}, -i \tan(c + dx), i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n} \sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{2(2Bn + iA(1 + 2n)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1 - n, \frac{3}{2}, -i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n} \sqrt{\tan(c + dx)}}{d(1 + 2n)}$$

```
output 2*B*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n/d/(1+2*n)-2*(I*A+B)*AppellF1(1/2
,1-n,1,3/2,-I*tan(d*x+c),I*tan(d*x+c))*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))
^n/d/((1+I*tan(d*x+c))^n)+2*(2*B*n+I*A*(1+2*n))*hypergeom([1/2, 1-n],[3/2
,-I*tan(d*x+c))*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n/d/(1+2*n)/((1+I*tan(
d*x+c))^n)
```

3.228.2 Mathematica [F]

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

input `Integrate[Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `Integrate[Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

3.228.3 Rubi [A] (warning: unable to verify)

Time = 0.99 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.15, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 4080, 27, 3042, 4084, 3042, 4047, 25, 27, 148, 27, 334, 333, 4082, 76, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$\downarrow 3042$$

$$\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$\downarrow 4080$$

$$\frac{2 \int -\frac{(i \tan(c+dx)a+a)^n(aB-a(2nA+A-2iBn) \tan(c+dx))}{2\sqrt{\tan(c+dx)}} dx}{a(2n+1)} + \frac{2B \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)}$$

$$\downarrow 27$$

$$\frac{2B \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n}{d(2n+1)} - \frac{\int \frac{(i \tan(c+dx)a+a)^n(aB-a(2nA+A-2iBn) \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)}$$

$$\downarrow 3042$$

3.228. $\int \sqrt{\tan(c+dx)}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

$$\begin{aligned}
 & \frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \frac{\int \frac{(i\tan(c+dx)a+a)^n(aB-a(2nA+A-2iBn)\tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)} \\
 & \quad \downarrow 4084 \\
 & \frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \\
 & \frac{a(2n+1)(B+iA)\int \frac{(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - (2Bn+iA(2n+1))\int \frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)} \\
 & \quad \downarrow 3042 \\
 & \frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \\
 & \frac{a(2n+1)(B+iA)\int \frac{(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - (2Bn+iA(2n+1))\int \frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)} \\
 & \quad \downarrow 4047 \\
 & \frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \\
 & \frac{ia^3(2n+1)(B+iA)\int -\frac{(i\tan(c+dx)a+a)^{n-1}}{a\sqrt{\tan(c+dx)}(a-ia\tan(c+dx))}d(ia\tan(c+dx)) - (2Bn+iA(2n+1))\int \frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)} \\
 & \quad \downarrow 25 \\
 & \frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \\
 & \frac{ia^3(2n+1)(B+iA)\int \frac{(i\tan(c+dx)a+a)^{n-1}}{a\sqrt{\tan(c+dx)}(a-ia\tan(c+dx))}d(ia\tan(c+dx)) - (2Bn+iA(2n+1))\int \frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)} \\
 & \quad \downarrow 27 \\
 & \frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \\
 & \frac{ia^2(2n+1)(B+iA)\int \frac{(i\tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)}(a-ia\tan(c+dx))}d(ia\tan(c+dx)) - (2Bn+iA(2n+1))\int \frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)} \\
 & \quad \downarrow 148 \\
 & \frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \\
 & \frac{2a^3(2n+1)(B+iA)\int \frac{(a-ia^3\tan^2(c+dx))^{n-1}}{a(ia^2\tan^2(c+dx)+1)}d\sqrt{\tan(c+dx)} - (2Bn+iA(2n+1))\int \frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)}
 \end{aligned}$$

3.228. $\int \sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n(A+B\tan(c+dx)) dx$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \\
 & \frac{2a^2(2n+1)(B+iA) \int \frac{(a-ia^3\tan^2(c+dx))^{n-1}}{ia^2\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{a(2n+1)} - (2Bn+iA(2n+1)) \int \frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx \\
 & \downarrow 334 \\
 & \frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \\
 & \frac{2a(2n+1)(B+iA)(1-ia^2\tan^2(c+dx))^{-n}(a-ia^3\tan^2(c+dx))^n \int \frac{(1-ia^2\tan^2(c+dx))^{n-1}}{ia^2\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{a(2n+1)} - (2Bn+iA(2n+1)) \int \frac{(a-ia\tan(c+dx))^n}{\sqrt{\tan(c+dx)}} dx \\
 & \downarrow 333 \\
 & \frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \\
 & \frac{2ia^2(2n+1)(B+iA)\tan(c+dx)(1-ia^2\tan^2(c+dx))^{-n}(a-ia^3\tan^2(c+dx))^n \operatorname{AppellF1}\left(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2\tan^2(c+dx), ia^2\tan^2(c+dx)\right)}{a(2n+1)} - (2Bn+iA(2n+1)) \int \frac{(a-ia\tan(c+dx))^n}{\sqrt{\tan(c+dx)}} dx \\
 & \downarrow 4082 \\
 & \frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \\
 & \frac{2ia^2(2n+1)(B+iA)\tan(c+dx)(1-ia^2\tan^2(c+dx))^{-n}(a-ia^3\tan^2(c+dx))^n \operatorname{AppellF1}\left(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2\tan^2(c+dx), ia^2\tan^2(c+dx)\right)}{a(2n+1)} - \frac{a^2(2Bn+iA(2n+1)) \int \frac{(a-ia\tan(c+dx))^n}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)} \\
 & \downarrow 76 \\
 & \frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \\
 & \frac{2ia^2(2n+1)(B+iA)\tan(c+dx)(1-ia^2\tan^2(c+dx))^{-n}(a-ia^3\tan^2(c+dx))^n \operatorname{AppellF1}\left(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2\tan^2(c+dx), ia^2\tan^2(c+dx)\right)}{a(2n+1)} - \frac{a(2Bn+iA(2n+1)) \int \frac{(a-ia\tan(c+dx))^n}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)} \\
 & \downarrow 74 \\
 & \frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \\
 & \frac{2ia^2(2n+1)(B+iA)\tan(c+dx)(1-ia^2\tan^2(c+dx))^{-n}(a-ia^3\tan^2(c+dx))^n \operatorname{AppellF1}\left(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2\tan^2(c+dx), ia^2\tan^2(c+dx)\right)}{a(2n+1)} - \frac{2a(2Bn+iA(2n+1)) \int \frac{(a-ia\tan(c+dx))^n}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)}
 \end{aligned}$$

3.228. $\int \sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n(A+B\tan(c+dx)) dx$

input `Int[Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `(2*B*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + 2*n)) - ((-2*a*(2*B*n + I*A*(1 + 2*n))*Hypergeometric2F1[1/2, 1 - n, 3/2, (-I)*Tan[c + d*x]]*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + I*Tan[c + d*x])^n) + ((2*I)*a^2*(I*A + B)*(1 + 2*n)*AppellF1[1/2, 1, 1 - n, 3/2, (-I)*a^2*Tan[c + d*x]^2, I*a^2*Tan[c + d*x]^2]*Tan[c + d*x]*(a - I*a^3*Tan[c + d*x]^2)^n)/(d*(1 - I*a^2*Tan[c + d*x]^2)^n)/(a*(1 + 2*n))`

3.228.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 74 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))`

rule 76 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0] && ((RationalQ[m] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0])) || !RationalQ[n])`

rule 148 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_.)*((e_) + (f_.)*(x_))^(p_.), x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c + d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x]] /; FreeQ[{b, c, d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`

- rule 333 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`
- rule 334 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4047 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4080 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`
- rule 4082 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.228.4 Maple [F]

$$\int \left(\sqrt{\tan(dx + c)} \right) (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.228.5 Fracas [F]

$$\begin{aligned} & \int \sqrt{\tan(c + dx)} (a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A) (ia \tan(dx + c) + a)^n \sqrt{\tan(dx + c)} dx \end{aligned}$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output `integral(((A - I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*(2*a*e^(2*I*d*x + 2*I*c))/(e^(2*I*d*x + 2*I*c) + 1))^n*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))/(e^(2*I*d*x + 2*I*c) + 1), x)`

3.228.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n(A+B\tan(c+dx))dx \\ &= \int (ia(\tan(c+dx)-i))^n(A+B\tan(c+dx))\sqrt{\tan(c+dx)}dx \end{aligned}$$

input `integrate(tan(d*x+c)**(1/2)*(a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x))*sqrt(tan(c + d*x)), x)`

3.228.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n(A+B\tan(c+dx))dx \\ &= \int (B\tan(dx+c)+A)(ia\tan(dx+c)+a)^n\sqrt{\tan(dx+c)}dx \end{aligned}$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*sqrt(tan(d*x + c)), x)`

3.228.8 Giac [F]

$$\begin{aligned} & \int \sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n(A+B\tan(c+dx))dx \\ &= \int (B\tan(dx+c)+A)(ia\tan(dx+c)+a)^n\sqrt{\tan(dx+c)}dx \end{aligned}$$

input `integrate(tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*sqrt(tan(d*x + c)), x)`

3.228.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \sqrt{\tan(c + dx)}(A + B \tan(c + dx)) (a + a \tan(c + dx) \text{ li})^n dx$$

input `int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n,x)`output `int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n, x)`

3.229
$$\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.229.1 Optimal result 2271
 3.229.2 Mathematica [F] 2271
 3.229.3 Rubi [A] (warning: unable to verify) 2272
 3.229.4 Maple [F] 2275
 3.229.5 Fracas [F] 2276
 3.229.6 Sympy [F] 2276
 3.229.7 Maxima [F] 2276
 3.229.8 Giac [F] 2277
 3.229.9 Mupad [F(-1)] 2277

3.229.1 Optimal result

Integrand size = 36, antiderivative size = 158

$$\int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \frac{2(A - iB) \operatorname{AppellF1}\left(\frac{1}{2}, 1 - n, 1, \frac{3}{2}, -i \tan(c + dx), i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n} \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))}{d} + \frac{2iB \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1 - n, \frac{3}{2}, -i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n} \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))}{d}$$

output

```
2*(A-I*B)*AppellF1(1/2,1-n,1,3/2,-I*tan(d*x+c),I*tan(d*x+c))*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n/d/((1+I*tan(d*x+c))^n)+2*I*B*hypergeom([1/2, 1-n],[3/2],-I*tan(d*x+c))*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n/d/((1+I*tan(d*x+c))^n)
```

3.229.2 Mathematica [F]

$$\int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

input `Integrate[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `Integrate[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]], x]`

3.229.3 Rubi [A] (warning: unable to verify)

Time = 0.67 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.11, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.361$, Rules used = {3042, 4084, 3042, 4047, 25, 27, 148, 27, 334, 333, 4082, 76, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
 & \quad \downarrow \text{4084} \\
 & (A - iB) \int \frac{(i \tan(c + dx)a + a)^n}{\sqrt{\tan(c + dx)}} dx + \frac{iB \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n}{\sqrt{\tan(c + dx)}} dx}{a} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \int \frac{(i \tan(c + dx)a + a)^n}{\sqrt{\tan(c + dx)}} dx + \frac{iB \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n}{\sqrt{\tan(c + dx)}} dx}{a} \\
 & \quad \downarrow \text{4047} \\
 & \frac{ia^2(A - iB) \int -\frac{(i \tan(c + dx)a + a)^{n-1}}{a \sqrt{\tan(c + dx)}(a - ia \tan(c + dx))} d(ia \tan(c + dx))}{d} + \frac{iB \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n}{\sqrt{\tan(c + dx)}} dx}{a} \\
 & \quad \downarrow \text{25} \\
 & \frac{iB \int \frac{(a - ia \tan(c + dx))(i \tan(c + dx)a + a)^n}{\sqrt{\tan(c + dx)}} dx}{a} - \frac{ia^2(A - iB) \int \frac{(i \tan(c + dx)a + a)^{n-1}}{a \sqrt{\tan(c + dx)}(a - ia \tan(c + dx))} d(ia \tan(c + dx))}{d} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.229. $\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$

$$\begin{aligned}
& \frac{iB \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n dx}{\sqrt{\tan(c+dx)}}}{a} - \frac{ia(A-iB) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)}(a-ia \tan(c+dx))} d(ia \tan(c+dx))}{d} \\
& \quad \downarrow 148 \\
& \frac{2a^2(A-iB) \int \frac{(a-ia^3 \tan^2(c+dx))^{n-1}}{a(ia^2 \tan^2(c+dx)+1)} d\sqrt{\tan(c+dx)}}{d} + \frac{iB \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n dx}{\sqrt{\tan(c+dx)}}}{a} \\
& \quad \downarrow 27 \\
& \frac{2a(A-iB) \int \frac{(a-ia^3 \tan^2(c+dx))^{n-1}}{ia^2 \tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} + \frac{iB \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n dx}{\sqrt{\tan(c+dx)}}}{a} \\
& \quad \downarrow 334 \\
& \frac{2(A-iB) (a-ia^3 \tan^2(c+dx))^n (1-ia^2 \tan^2(c+dx))^{-n} \int \frac{(1-ia^2 \tan^2(c+dx))^{n-1}}{ia^2 \tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} + \\
& \quad \frac{iB \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n dx}{\sqrt{\tan(c+dx)}}}{a} \\
& \quad \downarrow 333 \\
& \frac{iB \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n dx}{\sqrt{\tan(c+dx)}}}{a} + \\
& \frac{2ia(A-iB) \tan(c+dx) (a-ia^3 \tan^2(c+dx))^n (1-ia^2 \tan^2(c+dx))^{-n} \text{AppellF1}(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2 \tan^2(c+dx))}{d} \\
& \quad \downarrow 4082 \\
& \frac{iaB \int \frac{(i \tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)}} d \tan(c+dx)}{d} + \\
& \frac{2ia(A-iB) \tan(c+dx) (a-ia^3 \tan^2(c+dx))^n (1-ia^2 \tan^2(c+dx))^{-n} \text{AppellF1}(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2 \tan^2(c+dx))}{d} \\
& \quad \downarrow 76 \\
& \frac{iB(1+i \tan(c+dx))^{-n} (a+ia \tan(c+dx))^n \int \frac{(i \tan(c+dx)+1)^{n-1}}{\sqrt{\tan(c+dx)}} d \tan(c+dx)}{d} + \\
& \frac{2ia(A-iB) \tan(c+dx) (1-ia^2 \tan^2(c+dx))^{-n} (a-ia^3 \tan^2(c+dx))^n \text{AppellF1}(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2 \tan^2(c+dx))}{d} \\
& \quad \downarrow 74 \\
& \frac{2ia(A-iB) \tan(c+dx) (1-ia^2 \tan^2(c+dx))^{-n} (a-ia^3 \tan^2(c+dx))^n \text{AppellF1}(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2 \tan^2(c+dx))}{d} \\
& \frac{2iB \sqrt{\tan(c+dx)} (1+i \tan(c+dx))^{-n} (a+ia \tan(c+dx))^n \text{Hypergeometric2F1}(\frac{1}{2}, 1-n, \frac{3}{2}, -i \tan(c+dx))}{d}
\end{aligned}$$

3.229. $\int \frac{(a+ia \tan(c+dx))^n (A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

input `Int[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `((2*I)*B*Hypergeometric2F1[1/2, 1 - n, 3/2, (-I)*Tan[c + d*x]]*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + I*Tan[c + d*x])^n) + ((2*I)*a*(A - I*B)*AppellF1[1/2, 1, 1 - n, 3/2, (-I)*a^2*Tan[c + d*x]^2, I*a^2*Tan[c + d*x]^2]*Tan[c + d*x]*(a - I*a^3*Tan[c + d*x]^2)^n)/(d*(1 - I*a^2*Tan[c + d*x]^2)^n)`

3.229.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 74 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))`

rule 76 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0] && ((RationalQ[m] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0])) || !RationalQ[n])`

rule 148 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_.))*((e_) + (f_.)*(x_)^(p_.)), x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c + d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x] /; FreeQ[{b, c, d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`

rule 333 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 334 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a)^FracPart[p]) Int[(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4047 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4082 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.229.4 Maple [F]

$$\int \frac{(a + ia \tan(dx + c))^n (A + B \tan(dx + c))}{\sqrt{\tan(dx + c)}} dx$$

input `int((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x)`

output `int((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x)`

3.229.5 Fricas [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n}{\sqrt{\tan(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="fricas")`

output `integral(((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A - B)*(2*a*e^(2*I*d*x + 2*I*c))/(e^(2*I*d*x + 2*I*c) + 1))^n*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))/(e^(2*I*d*x + 2*I*c) - 1), x)`

3.229.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \int \frac{(ia(\tan(c + dx) - i))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

input `integrate((a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c))/tan(d*x+c)**(1/2),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x))/sqrt(tan(c + d*x))), x)`

3.229.7 Maxima [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n}{\sqrt{\tan(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n/sqrt(tan(d*x + c)), x)`

3.229.8 Giac [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n}{\sqrt{\tan(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n/sqrt(tan(d*x + c)), x)`

3.229.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) li)^n}{\sqrt{\tan(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n)/tan(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n)/tan(c + d*x)^(1/2), x)`

3.230
$$\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.230.1 Optimal result 2278
 3.230.2 Mathematica [F] 2278
 3.230.3 Rubi [A] (warning: unable to verify) 2279
 3.230.4 Maple [F] 2284
 3.230.5 Fricas [F] 2284
 3.230.6 Sympy [F] 2284
 3.230.7 Maxima [F(-1)] 2285
 3.230.8 Giac [F] 2285
 3.230.9 Mupad [F(-1)] 2285

3.230.1 Optimal result

Integrand size = 36, antiderivative size = 194

$$\int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = -\frac{2A(a + ia \tan(c + dx))^n}{d\sqrt{\tan(c + dx)}} + \frac{2(iA + B) \operatorname{AppellF1}\left(\frac{1}{2}, 1 - n, 1, \frac{3}{2}, -i \tan(c + dx), i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n} \sqrt{\tan(c + dx)}}{d} - \frac{2iA(1 - 2n) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1 - n, \frac{3}{2}, -i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n} \sqrt{\tan(c + dx)}(a + ia \tan(c + dx))}{d}$$

```
output -2*A*(a+I*a*tan(d*x+c))^n/d/tan(d*x+c)^(1/2)+2*(I*A+B)*AppellF1(1/2,1-n,1,
3/2,-I*tan(d*x+c),I*tan(d*x+c))*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n/d/((
1+I*tan(d*x+c))^n)-2*I*A*(1-2*n)*hypergeom([1/2, 1-n],[3/2],-I*tan(d*x+c))
*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n/d/((1+I*tan(d*x+c))^n)
```

3.230.2 Mathematica [F]

$$\int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

input `Integrate[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `Integrate[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2), x]`

3.230.3 Rubi [A] (warning: unable to verify)

Time = 0.95 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.13, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 4081, 27, 3042, 4084, 3042, 4047, 25, 27, 148, 27, 334, 333, 4082, 76, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx \\
 & \quad \downarrow \text{4081} \\
 & \frac{2 \int \frac{(i \tan(c+dx)a+a)^n (a(B+2iAn)-aA(1-2n) \tan(c+dx))}{2\sqrt{\tan(c+dx)}} dx}{a} - \frac{2A(a + ia \tan(c + dx))^n}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{(i \tan(c+dx)a+a)^n (a(B+2iAn)-aA(1-2n) \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{a} - \frac{2A(a + ia \tan(c + dx))^n}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{(i \tan(c+dx)a+a)^n (a(B+2iAn)-aA(1-2n) \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{a} - \frac{2A(a + ia \tan(c + dx))^n}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{4084} \\
 & \frac{a(B + iA) \int \frac{(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - iA(1 - 2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{2A(a + ia \tan(c + dx))^n} - \\
 & \quad \frac{a}{d\sqrt{\tan(c + dx)}}
 \end{aligned}$$

3.230. $\int \frac{(a+ia \tan(c+dx))^n (A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

$$\begin{array}{c}
\downarrow 3042 \\
\frac{a(B+iA) \int \frac{(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - iA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{2A(a+ia \tan(c+dx))^n} \\
\frac{a}{d\sqrt{\tan(c+dx)}} \\
\downarrow 4047 \\
\frac{ia^3(B+iA) \int -\frac{(i \tan(c+dx)a+a)^{n-1}}{a\sqrt{\tan(c+dx)}(a-ia \tan(c+dx))} d(ia \tan(c+dx))}{d} - iA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{2A(a+ia \tan(c+dx))^n} \\
\frac{a}{d\sqrt{\tan(c+dx)}} \\
\downarrow 25 \\
\frac{ia^3(B+iA) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{a\sqrt{\tan(c+dx)}(a-ia \tan(c+dx))} d(ia \tan(c+dx))}{d} - iA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{2A(a+ia \tan(c+dx))^n} \\
\frac{a}{d\sqrt{\tan(c+dx)}} \\
\downarrow 27 \\
\frac{ia^2(B+iA) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)}(a-ia \tan(c+dx))} d(ia \tan(c+dx))}{d} - iA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{2A(a+ia \tan(c+dx))^n} \\
\frac{a}{d\sqrt{\tan(c+dx)}} \\
\downarrow 148 \\
\frac{2a^3(B+iA) \int \frac{(a-ia^3 \tan^2(c+dx))^{n-1}}{a(ia^2 \tan^2(c+dx)+1)} d\sqrt{\tan(c+dx)}}{d} - iA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{2A(a+ia \tan(c+dx))^n} \\
\frac{a}{d\sqrt{\tan(c+dx)}} \\
\downarrow 27 \\
\frac{2a^2(B+iA) \int \frac{(a-ia^3 \tan^2(c+dx))^{n-1}}{ia^2 \tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} - iA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{2A(a+ia \tan(c+dx))^n} \\
\frac{a}{d\sqrt{\tan(c+dx)}} \\
\downarrow 334
\end{array}$$

3.230. $\int \frac{(a+ia \tan(c+dx))^n (A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

$$\frac{2a(B+iA)(1-ia^2 \tan^2(c+dx))^{-n}(a-ia^3 \tan^2(c+dx))^n \int \frac{(1-ia^2 \tan^2(c+dx))^{n-1}}{ia^2 \tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - iA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{d}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{d\sqrt{\tan(c + dx)}}$$

↓ 333

$$\frac{2ia^2(B+iA) \tan(c+dx)(1-ia^2 \tan^2(c+dx))^{-n}(a-ia^3 \tan^2(c+dx))^n \text{AppellF1}(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2 \tan^2(c+dx), ia^2 \tan^2(c+dx)) - iA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{d}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{d\sqrt{\tan(c + dx)}}$$

a

↓ 4082

$$\frac{2ia^2(B+iA) \tan(c+dx)(1-ia^2 \tan^2(c+dx))^{-n}(a-ia^3 \tan^2(c+dx))^n \text{AppellF1}(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2 \tan^2(c+dx), ia^2 \tan^2(c+dx)) - ia^2 A(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{d}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{d\sqrt{\tan(c + dx)}}$$

a

↓ 76

$$\frac{2ia^2(B+iA) \tan(c+dx)(1-ia^2 \tan^2(c+dx))^{-n}(a-ia^3 \tan^2(c+dx))^n \text{AppellF1}(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2 \tan^2(c+dx), ia^2 \tan^2(c+dx)) - iaA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{d}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{d\sqrt{\tan(c + dx)}}$$

a

↓ 74

$$\frac{2ia^2(B+iA) \tan(c+dx)(1-ia^2 \tan^2(c+dx))^{-n}(a-ia^3 \tan^2(c+dx))^n \text{AppellF1}(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2 \tan^2(c+dx), ia^2 \tan^2(c+dx)) - 2iaA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{d}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{d\sqrt{\tan(c + dx)}}$$

a

input `Int[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2), x]`

3.230. $\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$


```
output (-2*A*(a + I*a*Tan[c + d*x])^n)/(d*Sqrt[Tan[c + d*x]]) + (((-2*I)*a*A*(1 -
2*n)*Hypergeometric2F1[1/2, 1 - n, 3/2, (-I)*Tan[c + d*x]]*Sqrt[Tan[c + d
*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + I*Tan[c + d*x])^n) + ((2*I)*a^2*(I*
A + B)*AppellF1[1/2, 1, 1 - n, 3/2, (-I)*a^2*Tan[c + d*x]^2, I*a^2*Tan[c +
d*x]^2]*Tan[c + d*x]*(a - I*a^3*Tan[c + d*x]^2)^n)/(d*(1 - I*a^2*Tan[c +
d*x]^2)^n)/a
```

3.230.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 74 Int[((b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[c^n*((b*x
)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x]
/; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0]
&& !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))
```

```
rule 76 Int[((b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[c^IntPart
[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*
(x/c))^n, x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !Integer
Q[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0] && ((RationalQ[m] && !(EqQ[n, -2
^(-1)] && EqQ[c^2 - d^2, 0])) || !RationalQ[n])
```

```
rule 148 Int[((b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)),
x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c
+ d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x] /; FreeQ[{b, c,
d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]
```

```
rule 333 Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Sim
p[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; F
reeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a,
0]) && (IntegerQ[q] || GtQ[c, 0])
```

rule 334 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4047 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.230.4 Maple [F]

$$\int \frac{(a + ia \tan(dx + c))^n (A + B \tan(dx + c))}{\tan(dx + c)^{\frac{3}{2}}} dx$$

input `int((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x)`

output `int((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x)`

3.230.5 Fricas [F]

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n}{\tan(dx + c)^{\frac{3}{2}}} dx \end{aligned}$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="fricas")`

output `integral(-((A - I*B)*e^(4*I*d*x + 4*I*c) + 2*A*e^(2*I*d*x + 2*I*c) + A + I*B)*(2*a*e^(2*I*d*x + 2*I*c)/(e^(2*I*d*x + 2*I*c) + 1))^n*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))/(e^(4*I*d*x + 4*I*c) - 2*e^(2*I*d*x + 2*I*c) + 1), x)`

3.230.6 Sympy [F]

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(ia(\tan(c + dx) - i))^n (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \end{aligned}$$

input `integrate((a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c))/tan(d*x+c)**(3/2),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x))/tan(c + d*x)**(3/2), x)`

3.230.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="maxima")
```

```
output Timed out
```

3.230.8 Giac [F]

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n}{\tan(dx + c)^{\frac{3}{2}}} dx \end{aligned}$$

```
input integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="giac")
```

```
output integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n/tan(d*x + c)^(3/2), x)
```

3.230.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) i)^n}{\tan(c + dx)^{3/2}} dx \end{aligned}$$

```
input int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n)/tan(c + d*x)^(3/2),x)
```

```
output int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n)/tan(c + d*x)^(3/2), x)
```

3.230. $\int \frac{(a+ia \tan(c+dx))^n (A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

3.231
$$\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.231.1 Optimal result 2286
 3.231.2 Mathematica [F] 2287
 3.231.3 Rubi [A] (warning: unable to verify) 2287
 3.231.4 Maple [F] 2293
 3.231.5 Fricas [F] 2293
 3.231.6 Sympy [F] 2293
 3.231.7 Maxima [F] 2294
 3.231.8 Giac [F] 2294
 3.231.9 Mupad [F(-1)] 2294

3.231.1 Optimal result

Integrand size = 36, antiderivative size = 247

$$\int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= -\frac{2A(a + ia \tan(c + dx))^n}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2(3B + 2iAn)(a + ia \tan(c + dx))^n}{3d \sqrt{\tan(c + dx)}}$$

$$- \frac{2(A - iB) \operatorname{AppellF1}\left(\frac{1}{2}, 1 - n, 1, \frac{3}{2}, -i \tan(c + dx), i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n} \sqrt{\tan(c + dx)}}{d}$$

$$- \frac{2(1 - 2n)(3iB - 2An) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1 - n, \frac{3}{2}, -i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n} \sqrt{\tan(c + dx)}}{3d}$$

output

```
-2/3*(3*B+2*I*A*n)*(a+I*a*tan(d*x+c))^n/d/tan(d*x+c)^(1/2)-2*(A-I*B)*Appel
lF1(1/2,1-n,1,3/2,-I*tan(d*x+c),I*tan(d*x+c))*tan(d*x+c)^(1/2)*(a+I*a*tan(
d*x+c))^n/d/((1+I*tan(d*x+c))^n)-2/3*(1-2*n)*(3*I*B-2*A*n)*hypergeom([1/2,
1-n],[3/2],[-I*tan(d*x+c))*tan(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n/d/((1+I*t
an(d*x+c))^n)-2/3*A*(a+I*a*tan(d*x+c))^n/d/tan(d*x+c)^(3/2)
```

3.231.2 Mathematica [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

input `Integrate[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2), x]`

output `Integrate[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2), x]`

3.231.3 Rubi [A] (warning: unable to verify)

Time = 1.40 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.13, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.528$, Rules used = {3042, 4081, 27, 3042, 4081, 27, 3042, 4084, 3042, 4047, 25, 27, 148, 27, 334, 333, 4082, 76, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx$$

$$\downarrow \text{4081}$$

$$\frac{2 \int \frac{(i \tan(c+dx)a+a)^n (a(3B+2iAn)-aA(3-2n) \tan(c+dx))}{2 \tan^{\frac{3}{2}}(c+dx)} dx}{3a} - \frac{2A(a + ia \tan(c + dx))^n}{3d \tan^{\frac{3}{2}}(c + dx)}$$

$$\downarrow \text{27}$$

$$\frac{\int \frac{(i \tan(c+dx)a+a)^n (a(3B+2iAn)-aA(3-2n) \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx}{3a} - \frac{2A(a + ia \tan(c + dx))^n}{3d \tan^{\frac{3}{2}}(c + dx)}$$

3.231. $\int \frac{(a+ia \tan(c+dx))^n (A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \int \frac{(i \tan(c+dx)a+a)^n (a(3B+2iAn)-aA(3-2n) \tan(c+dx))}{\tan(c+dx)^{3/2}} dx - \frac{2A(a+ia \tan(c+dx))^n}{3d \tan^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2 \int \frac{(i \tan(c+dx)a+a)^n (a^2(6iBn-A(4n^2-2n+3))-a^2(1-2n)(3B+2iAn) \tan(c+dx))}{2\sqrt{\tan(c+dx)}} dx}{a} - \frac{2a(3B+2iAn)(a+ia \tan(c+dx))^n}{d\sqrt{\tan(c+dx)}} \\
 & \quad \downarrow \text{4081} \\
 & \frac{3a}{2A(a+ia \tan(c+dx))^n} - \frac{3d \tan^{\frac{3}{2}}(c+dx)}{3d \tan^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(i \tan(c+dx)a+a)^n (a^2(6iBn-A(4n^2-2n+3))-a^2(1-2n)(3B+2iAn) \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx - \frac{2a(3B+2iAn)(a+ia \tan(c+dx))^n}{d\sqrt{\tan(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3a}{2A(a+ia \tan(c+dx))^n} - \frac{3d \tan^{\frac{3}{2}}(c+dx)}{3d \tan^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow \text{4084} \\
 & \frac{-3a^2(A-iB) \int \frac{(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{a} - \frac{2a(3B+2iAn)(a+ia \tan(c+dx))^n}{d\sqrt{\tan(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3a}{2A(a+ia \tan(c+dx))^n} - \frac{3d \tan^{\frac{3}{2}}(c+dx)}{3d \tan^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow \text{4047} \\
 & \frac{-3a^2(A-iB) \int \frac{(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{a} - \frac{2a(3B+2iAn)(a+ia \tan(c+dx))^n}{d\sqrt{\tan(c+dx)}} \\
 & \quad \downarrow \\
 & \frac{3a}{2A(a+ia \tan(c+dx))^n} - \frac{3d \tan^{\frac{3}{2}}(c+dx)}{3d \tan^{\frac{3}{2}}(c+dx)}
 \end{aligned}$$

3.231. $\int \frac{(a+ia \tan(c+dx))^n (A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

$$\frac{3ia^4(A-iB) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{a \sqrt{\tan(c+dx)(a-ia \tan(c+dx))}} d(ia \tan(c+dx))}{a} - a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - \frac{2a(3B+2iAn)(a+ia \tan(c+dx))}{d \sqrt{\tan(c+dx)}}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{3d \tan^{\frac{3}{2}}(c + dx)} \quad 3a$$

↓ 25

$$\frac{3ia^4(A-iB) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{a \sqrt{\tan(c+dx)(a-ia \tan(c+dx))}} d(ia \tan(c+dx))}{a} - a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - \frac{2a(3B+2iAn)(a+ia \tan(c+dx))}{d \sqrt{\tan(c+dx)}}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{3d \tan^{\frac{3}{2}}(c + dx)} \quad 3a$$

↓ 27

$$\frac{3ia^3(A-iB) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)(a-ia \tan(c+dx))}} d(ia \tan(c+dx))}{a} - a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - \frac{2a(3B+2iAn)(a+ia \tan(c+dx))}{d \sqrt{\tan(c+dx)}}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{3d \tan^{\frac{3}{2}}(c + dx)} \quad 3a$$

↓ 148

$$\frac{6a^4(A-iB) \int \frac{(a-ia^3 \tan^2(c+dx))^{n-1}}{a(ia^2 \tan^2(c+dx)+1)} d\sqrt{\tan(c+dx)}}{a} - a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - \frac{2a(3B+2iAn)(a+ia \tan(c+dx))}{d \sqrt{\tan(c+dx)}}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{3d \tan^{\frac{3}{2}}(c + dx)} \quad 3a$$

↓ 27

$$\frac{6a^3(A-iB) \int \frac{(a-ia^3 \tan^2(c+dx))^{n-1}}{ia^2 \tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{a} - a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - \frac{2a(3B+2iAn)(a+ia \tan(c+dx))}{d \sqrt{\tan(c+dx)}}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{3d \tan^{\frac{3}{2}}(c + dx)} \quad 3a$$

↓ 334

3.231. $\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

$$\frac{6a^2(A-iB)(a-ia^3 \tan^2(c+dx))^n (1-ia^2 \tan^2(c+dx))^{-n} \int \frac{(1-ia^2 \tan^2(c+dx))^{n-1}}{ia^2 \tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} - \frac{a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx))}{\sqrt{\tan(c+dx)}}}{a}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{3d \tan^{\frac{3}{2}}(c + dx)} \qquad 3a$$

↓ 333

$$-a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - \frac{6ia^3(A-iB) \tan(c+dx)(a-ia^3 \tan^2(c+dx))^n (1-ia^2 \tan^2(c+dx))^{-n} \text{AppellF1}(\frac{1}{2}, 1, 1-n, \frac{1}{2}, 1, 1-n, -ia^2 \tan^2(c+dx))}{d}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{3d \tan^{\frac{3}{2}}(c + dx)} \qquad 3a$$

↓ 4082

$$- \frac{a^3(1-2n)(-2An+3iB) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)}} d \tan(c+dx)}{d} - \frac{6ia^3(A-iB) \tan(c+dx)(a-ia^3 \tan^2(c+dx))^n (1-ia^2 \tan^2(c+dx))^{-n} \text{AppellF1}(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2 \tan^2(c+dx))}{d}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{3d \tan^{\frac{3}{2}}(c + dx)} \qquad 3a$$

↓ 76

$$- \frac{a^2(1-2n)(-2An+3iB)(1+i \tan(c+dx))^{-n}(a+ia \tan(c+dx))^n \int \frac{(i \tan(c+dx)+1)^{n-1}}{\sqrt{\tan(c+dx)}} d \tan(c+dx)}{d} - \frac{6ia^3(A-iB) \tan(c+dx)(1-ia^2 \tan^2(c+dx))^{-n}(a-ia^3 \tan^2(c+dx))}{a}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{3d \tan^{\frac{3}{2}}(c + dx)} \qquad 3a$$

↓ 74

$$- \frac{2a^2(1-2n)(-2An+3iB)\sqrt{\tan(c+dx)}(1+i \tan(c+dx))^{-n}(a+ia \tan(c+dx))^n \text{Hypergeometric2F1}(\frac{1}{2}, 1-n, \frac{3}{2}, -i \tan(c+dx))}{d} - \frac{6ia^3(A-iB) \tan(c+dx)(1-ia^2 \tan^2(c+dx))}{a}$$

$$\frac{2A(a + ia \tan(c + dx))^n}{3d \tan^{\frac{3}{2}}(c + dx)} \qquad 3a$$

input Int[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]

3.231. $\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

```
output (-2*A*(a + I*a*Tan[c + d*x])^n)/(3*d*Tan[c + d*x]^(3/2)) + ((-2*a*(3*B + (
2*I)*A^n)*(a + I*a*Tan[c + d*x])^n)/(d*Sqrt[Tan[c + d*x]]) + ((-2*a^2*(1 -
2*n)*((3*I)*B - 2*A^n)*Hypergeometric2F1[1/2, 1 - n, 3/2, (-I)*Tan[c + d*
x]]*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + I*Tan[c + d*x])^n
) - ((6*I)*a^3*(A - I*B)*AppellF1[1/2, 1, 1 - n, 3/2, (-I)*a^2*Tan[c + d*x
]^2, I*a^2*Tan[c + d*x]^2]*Tan[c + d*x]*(a - I*a^3*Tan[c + d*x]^2)^n)/(d*(
1 - I*a^2*Tan[c + d*x]^2)^n)/a)/(3*a)
```

3.231.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 74 Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x
)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x]
/; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0]
&& !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))
```

```
rule 76 Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[c^IntPart
[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*
(x/c))^n, x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !Integer
Q[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0] && ((RationalQ[m] && !(EqQ[n, -2
^(-1)] && EqQ[c^2 - d^2, 0])) || !RationalQ[n])
```

```
rule 148 Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_),
x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c
+ d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x] /; FreeQ[{b, c,
d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]
```

```
rule 333 Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Sim
p[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; F
reeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a,
0]) && (IntegerQ[q] || GtQ[c, 0])
```

rule 334 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4047 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

3.231.4 Maple [F]

$$\int \frac{(a + ia \tan(dx + c))^n (A + B \tan(dx + c))}{\tan(dx + c)^{\frac{5}{2}}} dx$$

input `int((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x)`

output `int((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x)`

3.231.5 Fricas [F]

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n}{\tan(dx + c)^{\frac{5}{2}}} dx \end{aligned}$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="fricas")`

output `integral(((-I*A - B)*e^(6*I*d*x + 6*I*c) + (-3*I*A - B)*e^(4*I*d*x + 4*I*c) + (-3*I*A + B)*e^(2*I*d*x + 2*I*c) - I*A + B)*(2*a*e^(2*I*d*x + 2*I*c)/(e^(2*I*d*x + 2*I*c) + 1))^n*sqrt((-I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) + 1))/(e^(6*I*d*x + 6*I*c) - 3*e^(4*I*d*x + 4*I*c) + 3*e^(2*I*d*x + 2*I*c) - 1), x)`

3.231.6 Sympy [F]

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(ia(\tan(c + dx) - i))^n (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \end{aligned}$$

input `integrate((a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c))/tan(d*x+c)**(5/2),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x))/tan(c + d*x)**(5/2), x)`

3.231.7 Maxima [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(i a \tan(dx + c) + a)^n}{\tan(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n/tan(d*x + c)^(5/2), x)`

3.231.8 Giac [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(i a \tan(dx + c) + a)^n}{\tan(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n/tan(d*x + c)^(5/2), x)`

3.231.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) li)^n}{\tan(c + dx)^{5/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n)/tan(c + d*x)^(5/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n)/tan(c + d*x)^(5/2), x
)`

3.231. $\int \frac{(a+ia \tan(c+dx))^n (A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

3.232 $\int \tan^2(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

3.232.1 Optimal result	2296
3.232.2 Mathematica [A] (verified)	2296
3.232.3 Rubi [A] (verified)	2297
3.232.4 Maple [A] (verified)	2299
3.232.5 Fricas [A] (verification not implemented)	2299
3.232.6 Sympy [A] (verification not implemented)	2300
3.232.7 Maxima [A] (verification not implemented)	2300
3.232.8 Giac [B] (verification not implemented)	2301
3.232.9 Mupad [B] (verification not implemented)	2301

3.232.1 Optimal result

Integrand size = 29, antiderivative size = 87

$$\int \tan^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= -((aA - bB)x) + \frac{(Ab + aB) \log(\cos(c + dx))}{d}$$

$$+ \frac{(aA - bB) \tan(c + dx)}{d} + \frac{(Ab + aB) \tan^2(c + dx)}{2d} + \frac{bB \tan^3(c + dx)}{3d}$$

```
output - (A*a-B*b)*x+(A*b+B*a)*ln(cos(d*x+c))/d+(A*a-B*b)*tan(d*x+c)/d+1/2*(A*b+B*a)*tan(d*x+c)^2/d+1/3*b*B*tan(d*x+c)^3/d
```

3.232.2 Mathematica [A] (verified)

Time = 0.62 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.99

$$\int \tan^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{(-6aA + 6bB) \arctan(\tan(c + dx)) + 6(Ab + aB) \log(\cos(c + dx)) + 6(aA - bB) \tan(c + dx) + 3(Ab + aB) \tan^2(c + dx)}{6d}$$

```
input Integrate[Tan[c + d*x]^2*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]
```

output $((-6*a*A + 6*b*B)*ArcTan[Tan[c + d*x]] + 6*(A*b + a*B)*Log[Cos[c + d*x]] + 6*(a*A - b*B)*Tan[c + d*x] + 3*(A*b + a*B)*Tan[c + d*x]^2 + 2*b*B*Tan[c + d*x]^3)/(6*d)$

3.232.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 4075, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx \\ & \quad \downarrow 3042 \\ & \int \tan(c + dx)^2(a + b \tan(c + dx))(A + B \tan(c + dx)) dx \\ & \quad \downarrow 4075 \\ & \int \tan^2(c + dx)(aA - bB + (Ab + aB) \tan(c + dx)) dx + \frac{bB \tan^3(c + dx)}{3d} \\ & \quad \downarrow 3042 \\ & \int \tan(c + dx)^2(aA - bB + (Ab + aB) \tan(c + dx)) dx + \frac{bB \tan^3(c + dx)}{3d} \\ & \quad \downarrow 4011 \\ & \int \tan(c + dx)(-Ab - aB + (aA - bB) \tan(c + dx)) dx + \frac{(aB + Ab) \tan^2(c + dx)}{2d} + \frac{bB \tan^3(c + dx)}{3d} \\ & \quad \downarrow 3042 \\ & \int \tan(c + dx)(-Ab - aB + (aA - bB) \tan(c + dx)) dx + \frac{(aB + Ab) \tan^2(c + dx)}{2d} + \frac{bB \tan^3(c + dx)}{3d} \\ & \quad \downarrow 4008 \\ & -(aB + Ab) \int \tan(c + dx) dx + \frac{(aB + Ab) \tan^2(c + dx)}{2d} + \frac{(aA - bB) \tan(c + dx)}{d} - x(aA - bB) + \frac{bB \tan^3(c + dx)}{3d} \\ & \quad \downarrow 3042 \end{aligned}$$

3.232. $\int \tan^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\begin{aligned}
 & -(aB + Ab) \int \tan(c + dx) dx + \frac{(aB + Ab) \tan^2(c + dx)}{2d} + \frac{(aA - bB) \tan(c + dx)}{d} - x(aA - bB) + \\
 & \quad \frac{bB \tan^3(c + dx)}{3d} \\
 & \quad \downarrow \text{3956} \\
 & \frac{(aB + Ab) \tan^2(c + dx)}{2d} + \frac{(aA - bB) \tan(c + dx)}{d} + \frac{(aB + Ab) \log(\cos(c + dx))}{d} - x(aA - bB) + \\
 & \quad \frac{bB \tan^3(c + dx)}{3d}
 \end{aligned}$$

input `Int[Tan[c + d*x]^2*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `-((a*A - b*B)*x) + ((A*b + a*B)*Log[Cos[c + d*x]])/d + ((a*A - b*B)*Tan[c + d*x])/d + ((A*b + a*B)*Tan[c + d*x]^2)/(2*d) + (b*B*Tan[c + d*x]^3)/(3*d)`

3.232.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

3.232.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.01

method	result
norman	$(-aA + Bb)x + \frac{(aA - Bb)\tan(dx+c)}{d} + \frac{(Ab + Ba)(\tan^2(dx+c))}{2d} + \frac{bB(\tan^3(dx+c))}{3d} - \frac{(Ab + Ba)\ln(1 + \tan^2(dx+c))}{2d}$
parts	$\frac{(Ab + Ba)\left(\frac{\tan^2(dx+c)}{2} - \frac{\ln(1 + \tan^2(dx+c))}{2}\right)}{d} + \frac{Bb\left(\frac{\tan^3(dx+c)}{3} - \tan(dx+c) + \arctan(\tan(dx+c))\right)}{d} + \frac{aA(\tan(dx+c))}{d}$
derivativedivides	$\frac{\frac{bB(\tan^3(dx+c))}{3} + \frac{Ab(\tan^2(dx+c))}{2} + \frac{B(\tan^2(dx+c))a}{2} + A \tan(dx+c)a - bB \tan(dx+c) + \frac{(-Ab - Ba)\ln(1 + \tan^2(dx+c))}{2}}{d} + (-a)$
default	$\frac{\frac{bB(\tan^3(dx+c))}{3} + \frac{Ab(\tan^2(dx+c))}{2} + \frac{B(\tan^2(dx+c))a}{2} + A \tan(dx+c)a - bB \tan(dx+c) + \frac{(-Ab - Ba)\ln(1 + \tan^2(dx+c))}{2}}{d} + (-a)$
parallelrisch	$-\frac{-2bB(\tan^3(dx+c)) + 6Axad - 3Ab(\tan^2(dx+c)) - 6Bbdx - 3B(\tan^2(dx+c))a + 3A \ln(1 + \tan^2(dx+c))b - 6A \tan(dx+c)}{6d}$
risch	$-iAbx - iBax - Aax + Bbx - \frac{2iAbc}{d} - \frac{2iaBc}{d} + \frac{2i(-3iAb e^{4i(dx+c)} - 3iBa e^{4i(dx+c)} + 3Aa e^{4i(dx+c)})}{d}$

```
input int(tan(d*x+c)^2*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output (-A*a+B*b)*x+(A*a-B*b)*tan(d*x+c)/d+1/2*(A*b+B*a)*tan(d*x+c)^2/d+1/3*b*B*tan(d*x+c)^3/d-1/2*(A*b+B*a)/d*ln(1+tan(d*x+c)^2)
```

3.232.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.98

$$\int \tan^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2 B b \tan(dx + c)^3 - 6 (A a - B b) dx + 3 (B a + A b) \tan(dx + c)^2 + 3 (B a + A b) \log\left(\frac{1}{\tan(dx+c)^2 + 1}\right) + 6 (A a - B b) \tan(dx + c)}{6 d}$$

3.232. $\int \tan^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/6*(2*B*b*tan(d*x + c)^3 - 6*(A*a - B*b)*d*x + 3*(B*a + A*b)*tan(d*x + c)^2 + 3*(B*a + A*b)*log(1/(tan(d*x + c)^2 + 1)) + 6*(A*a - B*b)*tan(d*x + c))/d`

3.232.6 Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.56

$$\int \tan^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \begin{cases} -Aax + \frac{Aa \tan(c+dx)}{d} - \frac{Ab \log(\tan^2(c+dx)+1)}{2d} + \frac{Ab \tan^2(c+dx)}{2d} - \frac{Ba \log(\tan^2(c+dx)+1)}{2d} + \frac{Ba \tan^2(c+dx)}{2d} + Bbx + B \\ x(A + B \tan(c))(a + b \tan(c)) \tan^2(c) \end{cases}$$

input `integrate(tan(d*x+c)**2*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Piecewise((-A*a*x + A*a*tan(c + d*x)/d - A*b*log(tan(c + d*x)**2 + 1)/(2*d) + A*b*tan(c + d*x)**2/(2*d) - B*a*log(tan(c + d*x)**2 + 1)/(2*d) + B*a*tan(c + d*x)**2/(2*d) + B*b*x + B*b*tan(c + d*x)**3/(3*d) - B*b*tan(c + d*x))/d, Ne(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))*tan(c)**2, True))`

3.232.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.99

$$\int \tan^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2 B b \tan(dx + c)^3 + 3 (B a + A b) \tan(dx + c)^2 - 6 (A a - B b)(dx + c) - 3 (B a + A b) \log(\tan(dx + c)^2 + 1)}{6 d}$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/6*(2*B*b*tan(d*x + c)^3 + 3*(B*a + A*b)*tan(d*x + c)^2 - 6*(A*a - B*b)*(d*x + c) - 3*(B*a + A*b)*log(tan(d*x + c)^2 + 1) + 6*(A*a - B*b)*tan(d*x + c))/d`

3.232. $\int \tan^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

3.232.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 937 vs. $2(83) = 166$.

Time = 0.85 (sec) , antiderivative size = 937, normalized size of antiderivative = 10.77

$$\int \tan^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

```
-1/6*(6*A*a*d*x*tan(d*x)^3*tan(c)^3 - 6*B*b*d*x*tan(d*x)^3*tan(c)^3 - 3*B*
a*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2
+ tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^3*tan(c)^3 - 3*A*b*log(4*(tan(d*x)
^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + t
an(c)^2 + 1))*tan(d*x)^3*tan(c)^3 - 18*A*a*d*x*tan(d*x)^2*tan(c)^2 + 18*B*
b*d*x*tan(d*x)^2*tan(c)^2 - 3*B*a*tan(d*x)^3*tan(c)^3 - 3*A*b*tan(d*x)^3*t
an(c)^3 + 9*B*a*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d
*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^2*tan(c)^2 + 9*A*b*l
og(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 +
tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^2*tan(c)^2 + 6*A*a*tan(d*x)^3*tan(c)^
2 - 6*B*b*tan(d*x)^3*tan(c)^2 + 6*A*a*tan(d*x)^2*tan(c)^3 - 6*B*b*tan(d*x)
^2*tan(c)^3 + 18*A*a*d*x*tan(d*x)*tan(c) - 18*B*b*d*x*tan(d*x)*tan(c) - 3*
B*a*tan(d*x)^3*tan(c) - 3*A*b*tan(d*x)^3*tan(c) + 3*B*a*tan(d*x)^2*tan(c)^
2 + 3*A*b*tan(d*x)^2*tan(c)^2 - 3*B*a*tan(d*x)*tan(c)^3 - 3*A*b*tan(d*x)*t
an(c)^3 + 2*B*b*tan(d*x)^3 - 9*B*a*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)
*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)*t
an(c) - 9*A*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)
)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)*tan(c) - 12*A*a*tan(d*
x)^2*tan(c) + 18*B*b*tan(d*x)^2*tan(c) - 12*A*a*tan(d*x)*tan(c)^2 + 18*B*b
*tan(d*x)*tan(c)^2 + 2*B*b*tan(c)^3 - 6*A*a*d*x + 6*B*b*d*x + 3*B*a*tan...
```

3.232.9 Mupad [B] (verification not implemented)

Time = 7.28 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.97

$$\int \tan^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{\tan(c + dx) (Aa - Bb) - \ln(\tan(c + dx)^2 + 1) \left(\frac{Ab}{2} + \frac{Ba}{2}\right) + \tan(c + dx)^2 \left(\frac{Ab}{2} + \frac{Ba}{2}\right) - dx(Aa - Bb)}{d}$$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)`

output $(\tan(c + d*x)*(A*a - B*b) - \log(\tan(c + d*x)^2 + 1)*((A*b)/2 + (B*a)/2) + \tan(c + d*x)^2*((A*b)/2 + (B*a)/2) - d*x*(A*a - B*b) + (B*b*\tan(c + d*x)^3)/3)/d$

3.233 $\int \tan(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

3.233.1 Optimal result	2303
3.233.2 Mathematica [A] (verified)	2303
3.233.3 Rubi [A] (verified)	2304
3.233.4 Maple [A] (verified)	2305
3.233.5 Fricas [A] (verification not implemented)	2306
3.233.6 Sympy [A] (verification not implemented)	2306
3.233.7 Maxima [A] (verification not implemented)	2307
3.233.8 Giac [B] (verification not implemented)	2307
3.233.9 Mupad [B] (verification not implemented)	2308

3.233.1 Optimal result

Integrand size = 27, antiderivative size = 65

$$\int \tan(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= -((Ab+aB)x) - \frac{(aA-bB) \log(\cos(c+dx))}{d} + \frac{(Ab+aB) \tan(c+dx)}{d} + \frac{bB \tan^2(c+dx)}{2d}$$

output `-(A*b+B*a)*x-(A*a-B*b)*ln(cos(d*x+c))/d+(A*b+B*a)*tan(d*x+c)/d+1/2*b*B*tan(d*x+c)^2/d`

3.233.2 Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.03

$$\int \tan(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{-2(Ab+aB) \arctan(\tan(c+dx)) + 2(-aA+bB) \log(\cos(c+dx)) + 2(Ab+aB) \tan(c+dx) + bB \tan^2(c+dx)}{2d}$$

input `Integrate[Tan[c + d*x]*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(-2*(A*b + a*B)*ArcTan[Tan[c + d*x]] + 2*(-(a*A) + b*B)*Log[Cos[c + d*x]] + 2*(A*b + a*B)*Tan[c + d*x] + b*B*Tan[c + d*x]^2)/(2*d)`

3.233.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4075, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(c+dx)(a+b\tan(c+dx))(A+B\tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c+dx)(a+b\tan(c+dx))(A+B\tan(c+dx)) dx \\
 & \quad \downarrow \text{4075} \\
 & \int \tan(c+dx)(aA-bB+(Ab+aB)\tan(c+dx)) dx + \frac{bB\tan^2(c+dx)}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c+dx)(aA-bB+(Ab+aB)\tan(c+dx)) dx + \frac{bB\tan^2(c+dx)}{2d} \\
 & \quad \downarrow \text{4008} \\
 & (aA-bB) \int \tan(c+dx) dx + \frac{(aB+Ab)\tan(c+dx)}{d} - x(aB+Ab) + \frac{bB\tan^2(c+dx)}{2d} \\
 & \quad \downarrow \text{3042} \\
 & (aA-bB) \int \tan(c+dx) dx + \frac{(aB+Ab)\tan(c+dx)}{d} - x(aB+Ab) + \frac{bB\tan^2(c+dx)}{2d} \\
 & \quad \downarrow \text{3956} \\
 & \frac{(aB+Ab)\tan(c+dx)}{d} - \frac{(aA-bB)\log(\cos(c+dx))}{d} - x(aB+Ab) + \frac{bB\tan^2(c+dx)}{2d}
 \end{aligned}$$

input `Int[Tan[c + d*x]*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `-((A*b + a*B)*x) - ((a*A - b*B)*Log[Cos[c + d*x]])/d + ((A*b + a*B)*Tan[c + d*x])/d + (b*B*Tan[c + d*x]^2)/(2*d)`

3.233.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_. + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.233.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.06

method	result
norman	$(-Ab - Ba)x + \frac{(Ab+Ba)\tan(dx+c)}{d} + \frac{bB(\tan^2(dx+c))}{2d} + \frac{(aA-Bb)\ln(1+\tan^2(dx+c))}{2d}$
derivativedivides	$\frac{\frac{bB(\tan^2(dx+c))}{2} + A\tan(dx+c)b + B\tan(dx+c)a + \frac{(aA-Bb)\ln(1+\tan^2(dx+c))}{2}}{d} + (-Ab-Ba)\arctan(\tan(dx+c))$
default	$\frac{\frac{bB(\tan^2(dx+c))}{2} + A\tan(dx+c)b + B\tan(dx+c)a + \frac{(aA-Bb)\ln(1+\tan^2(dx+c))}{2}}{d} + (-Ab-Ba)\arctan(\tan(dx+c))$
parts	$\frac{(Ab+Ba)(\tan(dx+c)-\arctan(\tan(dx+c)))}{d} + \frac{aA\ln(1+\tan^2(dx+c))}{2d} + \frac{Bb\left(\frac{\tan^2(dx+c)}{2} - \frac{\ln(1+\tan^2(dx+c))}{2}\right)}{d}$
parallelrisch	$\frac{-2Abdx - 2Bxad + bB(\tan^2(dx+c)) + aA\ln(1+\tan^2(dx+c)) + 2A\tan(dx+c)b - B\ln(1+\tan^2(dx+c))b + 2B\tan(dx+c)a}{2d}$
risch	$-Abx - Bax + iAax - iBbx + \frac{2iaAc}{d} - \frac{2iBbc}{d} + \frac{2i(-iBbe^{2i(dx+c)} + Abe^{2i(dx+c)} + Ba e^{2i(dx+c)} + Ab + B)}{d(e^{2i(dx+c)} + 1)^2}$

input `int(tan(d*x+c)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

3.233. $\int \tan(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

output $(-A*b-B*a)*x+(A*b+B*a)*\tan(d*x+c)/d+1/2*b*B*\tan(d*x+c)^2/d+1/2*(A*a-B*b)/d*\ln(1+\tan(d*x+c)^2)$

3.233.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.02

$$\int \tan(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{Bb \tan(dx + c)^2 - 2(Ba + Ab)dx - (Aa - Bb) \log\left(\frac{1}{\tan(dx+c)^2+1}\right) + 2(Ba + Ab) \tan(dx + c)}{2d}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output $1/2*(B*b*\tan(d*x + c)^2 - 2*(B*a + A*b)*d*x - (A*a - B*b)*\log(1/(\tan(d*x + c)^2 + 1)) + 2*(B*a + A*b)*\tan(d*x + c))/d$

3.233.6 Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.60

$$\int \tan(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \frac{Aa \log(\tan^2(c+dx)+1)}{2d} - Abx + \frac{Ab \tan(c+dx)}{d} - Bax + \frac{Ba \tan(c+dx)}{d} - \frac{Bb \log(\tan^2(c+dx)+1)}{2d} + \frac{Bb \tan^2(c+dx)}{2d} & \text{for } d \\ x(A + B \tan(c))(a + b \tan(c)) \tan(c) & \text{otherwise} \end{cases}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Piecewise((A*a*log(tan(c + d*x)**2 + 1)/(2*d) - A*b*x + A*b*tan(c + d*x)/d - B*a*x + B*a*tan(c + d*x)/d - B*b*log(tan(c + d*x)**2 + 1)/(2*d) + B*b*tan(c + d*x)**2/(2*d), Ne(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))*tan(c), True))`

3.233.7 Maxima [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.02

$$\int \tan(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{Bb \tan(dx + c)^2 - 2(Ba + Ab)(dx + c) + (Aa - Bb) \log(\tan(dx + c)^2 + 1) + 2(Ba + Ab) \tan(dx + c)}{2d}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(B*b*tan(d*x + c)^2 - 2*(B*a + A*b)*(d*x + c) + (A*a - B*b)*log(tan(d*x + c)^2 + 1) + 2*(B*a + A*b)*tan(d*x + c))/d`

3.233.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 556 vs. 2(63) = 126.

Time = 0.51 (sec) , antiderivative size = 556, normalized size of antiderivative = 8.55

$$\int \tan(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx =$$

$$\frac{2Badx \tan(dx)^2 \tan(c)^2 + 2Abdx \tan(dx)^2 \tan(c)^2 + Aa \log\left(\frac{4(\tan(dx)^2 \tan(c)^2 - 2 \tan(dx) \tan(c) + 1)}{\tan(dx)^2 \tan(c)^2 + \tan(dx)^2 + \tan(c)^2 + 1}\right) \tan(dx)}{2d}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

```
-1/2*(2*B*a*d*x*tan(d*x)^2*tan(c)^2 + 2*A*b*d*x*tan(d*x)^2*tan(c)^2 + A*a*
log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 +
tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^2*tan(c)^2 - B*b*log(4*(tan(d*x)^2*t
an(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c
)^2 + 1))*tan(d*x)^2*tan(c)^2 - 4*B*a*d*x*tan(d*x)*tan(c) - 4*A*b*d*x*tan(
d*x)*tan(c) - B*b*tan(d*x)^2*tan(c)^2 - 2*A*a*log(4*(tan(d*x)^2*tan(c)^2 -
2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))
*tan(d*x)*tan(c) + 2*B*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) +
1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)*tan(c) + 2*
B*a*tan(d*x)^2*tan(c) + 2*A*b*tan(d*x)^2*tan(c) + 2*B*a*tan(d*x)*tan(c)^2
+ 2*A*b*tan(d*x)*tan(c)^2 + 2*B*a*d*x + 2*A*b*d*x - B*b*tan(d*x)^2 - B*b*t
an(c)^2 + A*a*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x
)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1)) - B*b*log(4*(tan(d*x)^2*tan(c)^
2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 +
1)) - 2*B*a*tan(d*x) - 2*A*b*tan(d*x) - 2*B*a*tan(c) - 2*A*b*tan(c) - B*b)
/(d*tan(d*x)^2*tan(c)^2 - 2*d*tan(d*x)*tan(c) + d)
```

3.233.9 Mupad [B] (verification not implemented)

Time = 8.43 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.97

$$\int \tan(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{\tan(c + dx)(Ab + Ba) + \ln(\tan(c + dx)^2 + 1) \left(\frac{Aa}{2} - \frac{Bb}{2}\right) - dx(Ab + Ba) + \frac{Bb \tan(c + dx)^2}{2}}{d}$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)`

output `(tan(c + d*x)*(A*b + B*a) + log(tan(c + d*x)^2 + 1)*((A*a)/2 - (B*b)/2) - d*x*(A*b + B*a) + (B*b*tan(c + d*x)^2)/2)/d`

3.234 $\int (a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

3.234.1 Optimal result	2309
3.234.2 Mathematica [A] (verified)	2309
3.234.3 Rubi [A] (verified)	2310
3.234.4 Maple [A] (verified)	2311
3.234.5 Fricas [A] (verification not implemented)	2311
3.234.6 Sympy [B] (verification not implemented)	2312
3.234.7 Maxima [A] (verification not implemented)	2312
3.234.8 Giac [B] (verification not implemented)	2313
3.234.9 Mupad [B] (verification not implemented)	2313

3.234.1 Optimal result

Integrand size = 21, antiderivative size = 42

$$\int (a + b \tan(c + dx))(A + B \tan(c + dx)) dx = (aA - bB)x - \frac{(Ab + aB) \log(\cos(c + dx))}{d} + \frac{bB \tan(c + dx)}{d}$$

output `(A*a-B*b)*x-(A*b+B*a)*ln(cos(d*x+c))/d+b*B*tan(d*x+c)/d`

3.234.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.40

$$\int (a + b \tan(c + dx))(A + B \tan(c + dx)) dx = aAx - \frac{bB \arctan(\tan(c + dx))}{d} - \frac{Ab \log(\cos(c + dx))}{d} - \frac{aB \log(\cos(c + dx))}{d} + \frac{bB \tan(c + dx)}{d}$$

input `Integrate[(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `a*A*x - (b*B*ArcTan[Tan[c + d*x]])/d - (A*b*Log[Cos[c + d*x]])/d - (a*B*Log[Cos[c + d*x]])/d + (b*B*Tan[c + d*x])/d`

3.234.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \tan(c + dx))(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + b \tan(c + dx))(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4008} \\ & (aB + Ab) \int \tan(c + dx) dx + x(aA - bB) + \frac{bB \tan(c + dx)}{d} \\ & \quad \downarrow \text{3042} \\ & (aB + Ab) \int \tan(c + dx) dx + x(aA - bB) + \frac{bB \tan(c + dx)}{d} \\ & \quad \downarrow \text{3956} \\ & -\frac{(aB + Ab) \log(\cos(c + dx))}{d} + x(aA - bB) + \frac{bB \tan(c + dx)}{d} \end{aligned}$$

input `Int[(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(a*A - b*B)*x - ((A*b + a*B)*Log[Cos[c + d*x]])/d + (b*B*Tan[c + d*x])/d`

3.234.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

```
rule 4008 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f),
x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e,
f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]
```

3.234.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.12

method	result
norman	$(aA - Bb)x + \frac{bB \tan(dx+c)}{d} + \frac{(Ab+Ba) \ln(1+\tan^2(dx+c))}{2d}$
derivativedivides	$\frac{bB \tan(dx+c) + \frac{(Ab+Ba) \ln(1+\tan^2(dx+c))}{2}}{d} + (aA - Bb) \arctan(\tan(dx+c))$
default	$\frac{bB \tan(dx+c) + \frac{(Ab+Ba) \ln(1+\tan^2(dx+c))}{2}}{d} + (aA - Bb) \arctan(\tan(dx+c))$
parts	$Aax + \frac{(Ab+Ba) \ln(1+\tan^2(dx+c))}{2d} + \frac{Bb(\tan(dx+c) - \arctan(\tan(dx+c)))}{d}$
parallelrisch	$\frac{2Axad - 2Bbdx + A \ln(1+\tan^2(dx+c))b + B \ln(1+\tan^2(dx+c))a + 2bB \tan(dx+c)}{2d}$
risch	$iAbx + iBax + Aax - Bbx + \frac{2iAbc}{d} + \frac{2iaBc}{d} + \frac{2iBb}{d(e^{2i(dx+c)}+1)} - \frac{\ln(e^{2i(dx+c)}+1)Ab}{d} - \frac{a \ln(e^{2i(dx+c)}+1)}{d}$

```
input int((a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output (A*a-B*b)*x+b*B*tan(d*x+c)/d+1/2*(A*b+B*a)/d*ln(1+tan(d*x+c)^2)
```

3.234.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.19

$$\int (a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2(Aa - Bb)dx + 2Bb \tan(dx + c) - (Ba + Ab) \log\left(\frac{1}{\tan(dx+c)^2 + 1}\right)}{2d}$$

```
input integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fracas")
```

```
output 1/2*(2*(A*a - B*b)*d*x + 2*B*b*tan(d*x + c) - (B*a + A*b)*log(1/(tan(d*x +
c)^2 + 1)))/d
```

3.234.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(36) = 72$.

Time = 0.08 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.74

$$\int (a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \begin{cases} Aax + \frac{Ab \log(\tan^2(c+dx)+1)}{2d} + \frac{Ba \log(\tan^2(c+dx)+1)}{2d} - Bbx + \frac{Bb \tan(c+dx)}{d} & \text{for } d \neq 0 \\ x(A + B \tan(c))(a + b \tan(c)) & \text{otherwise} \end{cases}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Piecewise((A*a*x + A*b*log(tan(c + d*x)**2 + 1)/(2*d) + B*a*log(tan(c + d*x)**2 + 1)/(2*d) - B*b*x + B*b*tan(c + d*x)/d, Ne(d, 0)), (x*(A + B*tan(c))*a + b*tan(c)), True))`

3.234.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.19

$$\int (a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2 B b \tan(dx + c) + 2(Aa - Bb)(dx + c) + (Ba + Ab) \log(\tan(dx + c)^2 + 1)}{2d}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(2*B*b*tan(d*x + c) + 2*(A*a - B*b)*(d*x + c) + (B*a + A*b)*log(tan(d*x + c)^2 + 1))/d`

3.234.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 289 vs. $2(42) = 84$.

Time = 0.39 (sec) , antiderivative size = 289, normalized size of antiderivative = 6.88

$$\int (a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2 A a d x \tan(dx) \tan(c) - 2 B b d x \tan(dx) \tan(c) - B a \log\left(\frac{4(\tan(dx)^2 \tan(c)^2 - 2 \tan(dx) \tan(c) + 1)}{\tan(dx)^2 \tan(c)^2 + \tan(dx)^2 + \tan(c)^2 + 1}\right) \tan(dx) \tan(c)}{d}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `1/2*(2*A*a*d*x*tan(d*x)*tan(c) - 2*B*b*d*x*tan(d*x)*tan(c) - B*a*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)*tan(c) - A*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)*tan(c) - 2*A*a*d*x + 2*B*b*d*x + B*a*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1)) + A*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1)) - 2*B*b*tan(d*x) - 2*B*b*tan(c))/(d*tan(d*x)*tan(c) - d)`

3.234.9 Mupad [B] (verification not implemented)

Time = 7.18 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.31

$$\int (a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{B b \tan(c + dx) + \frac{A b \ln(\tan(c + dx)^2 + 1)}{2} + \frac{B a \ln(\tan(c + dx)^2 + 1)}{2} + A a d x - B b d x}{d}$$

input `int((A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)`

output `(B*b*tan(c + d*x) + (A*b*log(tan(c + d*x)^2 + 1))/2 + (B*a*log(tan(c + d*x)^2 + 1))/2 + A*a*d*x - B*b*d*x)/d`

3.235 $\int \cot(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

3.235.1 Optimal result	2314
3.235.2 Mathematica [A] (verified)	2314
3.235.3 Rubi [A] (verified)	2315
3.235.4 Maple [A] (verified)	2317
3.235.5 Fricas [A] (verification not implemented)	2317
3.235.6 Sympy [B] (verification not implemented)	2318
3.235.7 Maxima [A] (verification not implemented)	2318
3.235.8 Giac [A] (verification not implemented)	2319
3.235.9 Mupad [B] (verification not implemented)	2319

3.235.1 Optimal result

Integrand size = 27, antiderivative size = 37

$$\int \cot(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= (Ab+aB)x - \frac{bB \log(\cos(c+dx))}{d} + \frac{aA \log(\sin(c+dx))}{d}$$

output $(A*b+B*a)*x-b*B*\ln(\cos(d*x+c))/d+a*A*\ln(\sin(d*x+c))/d$

3.235.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.32

$$\int \cot(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= Abx + aBx + \frac{aA \log(\cos(c+dx))}{d} - \frac{bB \log(\cos(c+dx))}{d} + \frac{aA \log(\tan(c+dx))}{d}$$

input $\text{Integrate}[\text{Cot}[c + d*x]*(a + b*\text{Tan}[c + d*x])*(A + B*\text{Tan}[c + d*x]),x]$

output $A*b*x + a*B*x + (a*A*\text{Log}[\text{Cos}[c + d*x]])/d - (b*B*\text{Log}[\text{Cos}[c + d*x]])/d + (a*A*\text{Log}[\text{Tan}[c + d*x]])/d$

3.235. $\int \cot(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

3.235.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.296$, Rules used = {3042, 4072, 3042, 3956, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan(c+dx)} dx \\
 & \quad \downarrow \text{4072} \\
 & \int \cot(c+dx)(aA+(Ab+aB) \tan(c+dx)) dx + bB \int \tan(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{aA+(Ab+aB) \tan(c+dx)}{\tan(c+dx)} dx + bB \int \tan(c+dx) dx \\
 & \quad \downarrow \text{3956} \\
 & \int \frac{aA+(Ab+aB) \tan(c+dx)}{\tan(c+dx)} dx - \frac{bB \log(\cos(c+dx))}{d} \\
 & \quad \downarrow \text{4014} \\
 & aA \int \cot(c+dx) dx + x(aB+Ab) - \frac{bB \log(\cos(c+dx))}{d} \\
 & \quad \downarrow \text{3042} \\
 & aA \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx + x(aB+Ab) - \frac{bB \log(\cos(c+dx))}{d} \\
 & \quad \downarrow \text{25} \\
 & -aA \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx + x(aB+Ab) - \frac{bB \log(\cos(c+dx))}{d} \\
 & \quad \downarrow \text{3956} \\
 & x(aB+Ab) + \frac{aA \log(-\sin(c+dx))}{d} - \frac{bB \log(\cos(c+dx))}{d}
 \end{aligned}$$

input `Int[Cot[c + d*x]*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(A*b + a*B)*x - (b*B*Log[Cos[c + d*x]])/d + (a*A*Log[-Sin[c + d*x]])/d`

3.235.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4072 `Int[((((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*(d/b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c - a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0]`

3.235.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.27

method	result	size
parallelrisch	$\frac{(-aA+Bb)\ln(\sec^2(dx+c))+2aA\ln(\tan(dx+c))+2dx(Ab+Ba)}{2d}$	47
norman	$(Ab+Ba)x + \frac{aA\ln(\tan(dx+c))}{d} - \frac{(aA-Bb)\ln(1+\tan^2(dx+c))}{2d}$	48
derivativedivides	$\frac{(-aA+Bb)\ln(1+\tan^2(dx+c))}{2} + \frac{(Ab+Ba)\arctan(\tan(dx+c))+aA\ln(\tan(dx+c))}{d}$	52
default	$\frac{(-aA+Bb)\ln(1+\tan^2(dx+c))}{2} + \frac{(Ab+Ba)\arctan(\tan(dx+c))+aA\ln(\tan(dx+c))}{d}$	52
risch	$Abx + Bax - iAax + iBbx - \frac{2iaAc}{d} + \frac{2iBbc}{d} + \frac{aA\ln(e^{2i(dx+c)}-1)}{d} - \frac{\ln(e^{2i(dx+c)}+1)Bb}{d}$	77

input `int(cot(d*x+c)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`output `1/2*((-A*a+B*b)*ln(sec(d*x+c)^2)+2*a*A*ln(tan(d*x+c))+2*d*x*(A*b+B*a))/d`**3.235.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.59

$$\int \cot(c+dx)(a+b\tan(c+dx))(A+B\tan(c+dx))dx$$

$$= \frac{2(Ba+Ab)dx + Aa \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) - Bb \log\left(\frac{1}{\tan(dx+c)^2+1}\right)}{2d}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fracas")`output `1/2*(2*(B*a + A*b)*d*x + A*a*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1)) - B*b*log(1/(tan(d*x + c)^2 + 1)))/d`

3.235.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 78 vs. $2(34) = 68$.

Time = 0.21 (sec) , antiderivative size = 78, normalized size of antiderivative = 2.11

$$\int \cot(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \begin{cases} -\frac{Aa \log(\tan^2(c+dx)+1)}{2d} + \frac{Aa \log(\tan(c+dx))}{d} + Abx + Bax + \frac{Bb \log(\tan^2(c+dx)+1)}{2d} & \text{for } d \neq 0 \\ x(A + B \tan(c))(a + b \tan(c)) \cot(c) & \text{otherwise} \end{cases}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Piecewise((-A*a*log(tan(c + d*x)**2 + 1)/(2*d) + A*a*log(tan(c + d*x))/d + A*b*x + B*a*x + B*b*log(tan(c + d*x)**2 + 1)/(2*d), Ne(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))*cot(c), True))`

3.235.7 Maxima [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.41

$$\int \cot(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2Aa \log(\tan(dx + c)) + 2(Ba + Ab)(dx + c) - (Aa - Bb) \log(\tan(dx + c)^2 + 1)}{2d}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(2*A*a*log(tan(d*x + c)) + 2*(B*a + A*b)*(d*x + c) - (A*a - B*b)*log(tan(d*x + c)^2 + 1))/d`

3.235.8 Giac [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.43

$$\int \cot(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2 A a \log(|\tan(dx + c)|) + 2 (B a + A b)(dx + c) - (A a - B b) \log(\tan(dx + c)^2 + 1)}{2 d}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `1/2*(2*A*a*log(abs(tan(d*x + c))) + 2*(B*a + A*b)*(d*x + c) - (A*a - B*b)*log(tan(d*x + c)^2 + 1))/d`

3.235.9 Mupad [B] (verification not implemented)

Time = 7.38 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.86

$$\int \cot(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{A a \ln(\tan(c + dx))}{d} - \frac{\ln(\tan(c + dx) - i) (A + B i) (a + b i)}{2 d}$$

$$+ \frac{\ln(\tan(c + dx) + i) (A - B i) (b + a i) i}{2 d}$$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)`

output `(log(tan(c + d*x) + 1i)*(A - B*1i)*(a*1i + b)*1i)/(2*d) - (log(tan(c + d*x) - 1i)*(A + B*1i)*(a + b*1i))/(2*d) + (A*a*log(tan(c + d*x)))/d`

3.236 $\int \cot^2(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

3.236.1 Optimal result	2320
3.236.2 Mathematica [C] (verified)	2320
3.236.3 Rubi [A] (verified)	2321
3.236.4 Maple [A] (verified)	2323
3.236.5 Fricas [A] (verification not implemented)	2323
3.236.6 Sympy [B] (verification not implemented)	2324
3.236.7 Maxima [A] (verification not implemented)	2324
3.236.8 Giac [B] (verification not implemented)	2325
3.236.9 Mupad [B] (verification not implemented)	2325

3.236.1 Optimal result

Integrand size = 29, antiderivative size = 43

$$\int \cot^2(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= -((aA - bB)x) - \frac{aA \cot(c+dx)}{d} + \frac{(Ab + aB) \log(\sin(c+dx))}{d}$$

output

```
-(A*a-B*b)*x-a*A*cot(d*x+c)/d+(A*b+B*a)*ln(sin(d*x+c))/d
```

3.236.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.05 (sec) , antiderivative size = 88, normalized size of antiderivative = 2.05

$$\int \cot^2(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= bBx - \frac{aA \cot(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(c+dx)\right)}{d}$$

$$+ \frac{Ab \log(\cos(c+dx))}{d} + \frac{aB \log(\cos(c+dx))}{d}$$

$$+ \frac{Ab \log(\tan(c+dx))}{d} + \frac{aB \log(\tan(c+dx))}{d}$$

input `Integrate[Cot[c + d*x]^2*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `b*B*x - (a*A*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2])/d + (A*b*Log[Cos[c + d*x]])/d + (a*B*Log[Cos[c + d*x]])/d + (A*b*Log[Tan[c + d*x]])/d + (a*B*Log[Tan[c + d*x]])/d`

3.236.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.05, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.241$, Rules used = {3042, 4074, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan(c + dx)^2} dx \\
 & \quad \downarrow \text{4074} \\
 & \int \cot(c + dx)(Ab + aB - (aA - bB) \tan(c + dx)) dx - \frac{aA \cot(c + dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\tan(c + dx)} dx - \frac{aA \cot(c + dx)}{d} \\
 & \quad \downarrow \text{4014} \\
 & (aB + Ab) \int \cot(c + dx) dx - (x(aA - bB)) - \frac{aA \cot(c + dx)}{d} \\
 & \quad \downarrow \text{3042} \\
 & (aB + Ab) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - (x(aA - bB)) - \frac{aA \cot(c + dx)}{d} \\
 & \quad \downarrow \text{25} \\
 & -(aB + Ab) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - (x(aA - bB)) - \frac{aA \cot(c + dx)}{d} \\
 & \quad \downarrow \text{3956}
 \end{aligned}$$

3.236. $\int \cot^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\frac{(aB + Ab) \log(-\sin(c + dx))}{d} - (x(aA - bB)) - \frac{aA \cot(c + dx)}{d}$$

input `Int[Cot[c + d*x]^2*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `-((a*A - b*B)*x) - (a*A*Cot[c + d*x])/d + ((A*b + a*B)*Log[-Sin[c + d*x]])/d`

3.236.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4074 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.236.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.51

method	result
parallelrisch	$\frac{(-Ab-Ba)\ln(\sec^2(dx+c))+(2Ab+2Ba)\ln(\tan(dx+c))-2aA\cot(dx+c)-2dx(aA-Bb)}{2d}$
derivativedivides	$\frac{(-Ab-Ba)\ln\left(\frac{1+\tan^2(dx+c)}{2}\right)+(-aA+Bb)\arctan(\tan(dx+c))+(Ab+Ba)\ln(\tan(dx+c))-\frac{aA}{\tan(dx+c)}}{d}$
default	$\frac{(-Ab-Ba)\ln\left(\frac{1+\tan^2(dx+c)}{2}\right)+(-aA+Bb)\arctan(\tan(dx+c))+(Ab+Ba)\ln(\tan(dx+c))-\frac{aA}{\tan(dx+c)}}{d}$
norman	$\frac{(-aA+Bb)x\tan(dx+c)-\frac{aA}{d}}{\tan(dx+c)}+\frac{(Ab+Ba)\ln(\tan(dx+c))}{d}-\frac{(Ab+Ba)\ln(1+\tan^2(dx+c))}{2d}$
risch	$-iAbx - iBax - Aax + Bbx - \frac{2iAbc}{d} - \frac{2iaBc}{d} - \frac{2iaA}{d(e^{2i(dx+c)}-1)} + \frac{\ln(e^{2i(dx+c)}-1)Ab}{d} + \frac{a\ln(e^{2i(dx+c)}-1)}{d}$

input `int(cot(d*x+c)^2*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/2*((-A*b-B*a)*ln(sec(d*x+c)^2)+(2*A*b+2*B*a)*ln(tan(d*x+c))-2*a*A*cot(d*x+c)-2*d*x*(A*a-B*b))/d`

3.236.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.70

$$\int \cot^2(c+dx)(a+b\tan(c+dx))(A+B\tan(c+dx))dx$$

$$= -\frac{2(Aa-Bb)dx\tan(dx+c)-(Ba+Ab)\log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right)\tan(dx+c)+2Aa}{2d\tan(dx+c)}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x,algorithm="fricas")`

output `-1/2*(2*(A*a-B*b)*d*x*tan(d*x+c)-(B*a+A*b)*log(tan(d*x+c)^2/(tan(d*x+c)^2+1))*tan(d*x+c)+2*A*a)/(d*tan(d*x+c))`

3.236.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. 2(36) = 72.

Time = 0.32 (sec) , antiderivative size = 121, normalized size of antiderivative = 2.81

$$\int \cot^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \tilde{\infty} Aax \\ x(A + B \tan(c))(a + b \tan(c)) \cot^2(c) \\ \tilde{\infty} Aax \\ -Aax - \frac{Aa}{d \tan(c+dx)} - \frac{Ab \log(\tan^2(c+dx)+1)}{2d} + \frac{Ab \log(\tan(c+dx))}{d} - \frac{Ba \log(\tan^2(c+dx)+1)}{2d} + \frac{Ba \log(\tan(c+dx))}{d} + Bbx \end{cases}$$

input `integrate(cot(d*x+c)**2*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Piecewise((zoo*A*a*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*cot(c)**2, Eq(d, 0)), (zoo*A*a*x, Eq(c, -d*x)), (-A*a*x - A*a/(d*tan(c + d*x)) - A*b*log(tan(c + d*x)**2 + 1)/(2*d) + A*b*log(tan(c + d*x))/d - B*a*log(tan(c + d*x)**2 + 1)/(2*d) + B*a*log(tan(c + d*x))/d + B*b*x, True))`

3.236.7 Maxima [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.58

$$\int \cot^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx =$$

$$\frac{2(Aa - Bb)(dx + c) + (Ba + Ab) \log(\tan(dx + c)^2 + 1) - 2(Ba + Ab) \log(\tan(dx + c)) + \frac{2Aa}{\tan(dx+c)}}{2d}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(2*(A*a - B*b)*(d*x + c) + (B*a + A*b)*log(tan(d*x + c)^2 + 1) - 2*(B*a + A*b)*log(tan(d*x + c)) + 2*A*a/tan(d*x + c))/d`

3.236.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. $2(43) = 86$.

Time = 0.54 (sec) , antiderivative size = 119, normalized size of antiderivative = 2.77

$$\int \cot^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{Aa \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 2(Aa - Bb)(dx + c) - 2(Ba + Ab) \log\left(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 1\right) + 2(Ba + Ab) \log}{2d}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `1/2*(A*a*tan(1/2*d*x + 1/2*c) - 2*(A*a - B*b)*(d*x + c) - 2*(B*a + A*b)*log(tan(1/2*d*x + 1/2*c)^2 + 1) + 2*(B*a + A*b)*log(abs(tan(1/2*d*x + 1/2*c))) - (2*B*a*tan(1/2*d*x + 1/2*c) + 2*A*b*tan(1/2*d*x + 1/2*c) + A*a)/tan(1/2*d*x + 1/2*c))/d`

3.236.9 Mupad [B] (verification not implemented)

Time = 8.48 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.02

$$\int \cot^2(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx)) (Ab + Ba)}{d} - \frac{\ln(\tan(c + dx) + 1i) (A - B 1i) (b + a 1i)}{2d}$$

$$- \frac{Aa \cot(c + dx)}{d} + \frac{\ln(\tan(c + dx) - i) (A + B 1i) (a + b 1i) 1i}{2d}$$

input `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)`

output `(log(tan(c + d*x))*(A*b + B*a))/d + (log(tan(c + d*x) - 1i)*(A + B*1i)*(a + b*1i)*1i)/(2*d) - (log(tan(c + d*x) + 1i)*(A - B*1i)*(a*1i + b))/(2*d) - (A*a*cot(c + d*x))/d`

3.237 $\int \cot^3(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

3.237.1 Optimal result	2326
3.237.2 Mathematica [C] (verified)	2326
3.237.3 Rubi [A] (verified)	2327
3.237.4 Maple [A] (verified)	2329
3.237.5 Fricas [A] (verification not implemented)	2330
3.237.6 Sympy [B] (verification not implemented)	2330
3.237.7 Maxima [A] (verification not implemented)	2331
3.237.8 Giac [B] (verification not implemented)	2331
3.237.9 Mupad [B] (verification not implemented)	2332

3.237.1 Optimal result

Integrand size = 29, antiderivative size = 66

$$\int \cot^3(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= -((Ab+aB)x) - \frac{(Ab+aB) \cot(c+dx)}{d} - \frac{aA \cot^2(c+dx)}{2d} - \frac{(aA-bB) \log(\sin(c+dx))}{d}$$

output `-(A*b+B*a)*x-(A*b+B*a)*cot(d*x+c)/d-1/2*a*A*cot(d*x+c)^2/d-(A*a-B*b)*ln(sin(d*x+c))/d`

3.237.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.50 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.17

$$\int \cot^3(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx =$$

$$\frac{aA \cot^2(c+dx) + 2(Ab+aB) \cot(c+dx) \text{Hypergeometric2F1}(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(c+dx)) + 2(aA-bB) \log(\sin(c+dx))}{2d}$$

input `Integrate[Cot[c + d*x]^3*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output $-1/2*(a*A*\text{Cot}[c + d*x]^2 + 2*(A*b + a*B)*\text{Cot}[c + d*x]*\text{Hypergeometric2F1}[-1/2, 1, 1/2, -\text{Tan}[c + d*x]^2] + 2*(a*A - b*B)*(\text{Log}[\text{Cos}[c + d*x]] + \text{Log}[\text{Tan}[c + d*x]]))/d$

3.237.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 4074, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^3(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan(c + dx)^3} dx \\
 & \quad \downarrow 4074 \\
 & \int \cot^2(c + dx)(Ab + aB - (aA - bB) \tan(c + dx)) dx - \frac{aA \cot^2(c + dx)}{2d} \\
 & \quad \downarrow 3042 \\
 & \int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\tan(c + dx)^2} dx - \frac{aA \cot^2(c + dx)}{2d} \\
 & \quad \downarrow 4012 \\
 & \int -\cot(c + dx)(aA - bB + (Ab + aB) \tan(c + dx)) dx - \frac{(aB + Ab) \cot(c + dx)}{d} - \frac{aA \cot^2(c + dx)}{2d} \\
 & \quad \downarrow 25 \\
 & -\int \cot(c + dx)(aA - bB + (Ab + aB) \tan(c + dx)) dx - \frac{(aB + Ab) \cot(c + dx)}{d} - \frac{aA \cot^2(c + dx)}{2d} \\
 & \quad \downarrow 3042 \\
 & -\int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\tan(c + dx)} dx - \frac{(aB + Ab) \cot(c + dx)}{d} - \frac{aA \cot^2(c + dx)}{2d} \\
 & \quad \downarrow 4014 \\
 & -(aA - bB) \int \cot(c + dx) dx - \frac{(aB + Ab) \cot(c + dx)}{d} - x(aB + Ab) - \frac{aA \cot^2(c + dx)}{2d}
 \end{aligned}$$

3.237. $\int \cot^3(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& -(aA - bB) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - \frac{(aB + Ab) \cot(c + dx)}{d} - x(aB + Ab) - \frac{aA \cot^2(c + dx)}{2d} \\
& \downarrow \text{25} \\
& (aA - bB) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - \frac{(aB + Ab) \cot(c + dx)}{d} - x(aB + Ab) - \frac{aA \cot^2(c + dx)}{2d} \\
& \downarrow \text{3956} \\
& -\frac{(aB + Ab) \cot(c + dx)}{d} - \frac{(aA - bB) \log(-\sin(c + dx))}{d} - x(aB + Ab) - \frac{aA \cot^2(c + dx)}{2d}
\end{aligned}$$

input `Int[Cot[c + d*x]^3*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `-((A*b + a*B)*x) - ((A*b + a*B)*Cot[c + d*x])/d - (a*A*Cot[c + d*x]^2)/(2*d) - ((a*A - b*B)*Log[-Sin[c + d*x]])/d`

3.237.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

3.237.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.35

method	result
derivativedivides	$\frac{(aA - Bb) \ln\left(\frac{1 + \tan^2(dx+c)}{2}\right) + (-Ab - Ba) \arctan(\tan(dx+c)) - \frac{Ab+Ba}{\tan(dx+c)} + (-aA+Bb) \ln(\tan(dx+c)) - \frac{aA}{2 \tan(dx+c)^2}}{d}$
default	$\frac{(aA - Bb) \ln\left(\frac{1 + \tan^2(dx+c)}{2}\right) + (-Ab - Ba) \arctan(\tan(dx+c)) - \frac{Ab+Ba}{\tan(dx+c)} + (-aA+Bb) \ln(\tan(dx+c)) - \frac{aA}{2 \tan(dx+c)^2}}{d}$
parallelrisc	$\frac{-A(\cot^2(dx+c))a - 2Abdx - 2Bxad - 2A \cot(dx+c)b - 2aA \ln(\tan(dx+c)) + A \ln(\sec^2(dx+c))a - 2B \cot(dx+c)a + 2B \ln(\tan(dx+c))}{2d}$
norman	$\frac{(-Ab - Ba)x(\tan^2(dx+c)) - \frac{aA}{2d} - \frac{(Ab+Ba) \tan(dx+c)}{d}}{\tan(dx+c)^2} - \frac{(aA - Bb) \ln(\tan(dx+c))}{d} + \frac{(aA - Bb) \ln(1 + \tan^2(dx+c))}{2d}$
risc	$-Abx - Bax + iAax - iBbx + \frac{2iaAc}{d} - \frac{2iBbc}{d} - \frac{2i(iAa e^{2i(dx+c)} + Ab e^{2i(dx+c)} + Ba e^{2i(dx+c)} - Ab - Ba)}{d(e^{2i(dx+c)} - 1)^2}$

```
input int(cot(d*x+c)^3*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE
)
```

```
output 1/d*(1/2*(A*a-B*b)*ln(1+tan(d*x+c)^2)+(-A*b-B*a)*arctan(tan(d*x+c))-
(A*b+B*a)/tan(d*x+c)+(-A*a+B*b)*ln(tan(d*x+c))-1/2*a*A/tan(d*x+c)^2)
```


3.237.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.44

$$\int \cot^3(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{(Aa - Bb) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^2 + (2(Ba + Ab)dx + Aa) \tan(dx+c)^2 + Aa + 2(Ba + Ab)}{2d \tan(dx+c)^2}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-1/2*((A*a - B*b)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^2 + (2*(B*a + A*b)*d*x + A*a)*tan(d*x + c)^2 + A*a + 2*(B*a + A*b)*tan(d*x + c))/(d*tan(d*x + c)^2)`

3.237.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 148 vs. 2(56) = 112.

Time = 0.56 (sec) , antiderivative size = 148, normalized size of antiderivative = 2.24

$$\int \cot^3(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \begin{cases} \tilde{\infty} Aax \\ x(A + B \tan(c))(a + b \tan(c)) \cot^3(c) \\ \tilde{\infty} Aax \\ \frac{Aa \log(\tan^2(c+dx)+1)}{2d} - \frac{Aa \log(\tan(c+dx))}{d} - \frac{Aa}{2d \tan^2(c+dx)} - Abx - \frac{Ab}{d \tan(c+dx)} - Bax - \frac{Ba}{d \tan(c+dx)} - \frac{Bb \log(\tan^2(c+dx))}{2d} \end{cases}$$

input `integrate(cot(d*x+c)**3*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Piecewise((zoo*A*a*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))*cot(c)**3, Eq(d, 0)), (zoo*A*a*x, Eq(c, -d*x)), (A*a*log(tan(c + d*x)**2 + 1)/(2*d) - A*a*log(tan(c + d*x))/d - A*a/(2*d*tan(c + d*x)**2) - A*b*x - A*b/(d*tan(c + d*x)) - B*a*x - B*a/(d*tan(c + d*x)) - B*b*log(tan(c + d*x)**2 + 1)/(2*d) + B*b*log(tan(c + d*x))/d, True))`

3.237.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.30

$$\int \cot^3(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{2(Ba + Ab)(dx + c) - (Aa - Bb) \log(\tan(dx + c)^2 + 1) + 2(Aa - Bb) \log(\tan(dx + c)) + \frac{Aa + 2(Ba + Ab) \tan(dx + c)}{\tan(dx + c)}}{2d}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(2*(B*a + A*b)*(d*x + c) - (A*a - B*b)*log(tan(d*x + c)^2 + 1) + 2*(A*a - B*b)*log(tan(d*x + c)) + (A*a + 2*(B*a + A*b)*tan(d*x + c))/tan(d*x + c)^2)/d`

3.237.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. 2(64) = 128.

Time = 0.63 (sec) , antiderivative size = 179, normalized size of antiderivative = 2.71

$$\int \cot^3(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{Aa \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 4Ba \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 4Ab \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 8(Ba + Ab)(dx + c) - 8(Aa - Bb) \log(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + 1) + 8(Aa - Bb) \log(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)) - (12Aa \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 12Bb \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 4Ba \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 4Ab \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - Aa) / \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2}{d}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `-1/8*(A*a*tan(1/2*d*x + 1/2*c)^2 - 4*B*a*tan(1/2*d*x + 1/2*c) - 4*A*b*tan(1/2*d*x + 1/2*c) + 8*(B*a + A*b)*(d*x + c) - 8*(A*a - B*b)*log(tan(1/2*d*x + 1/2*c)^2 + 1) + 8*(A*a - B*b)*log(abs(tan(1/2*d*x + 1/2*c))) - (12*A*a*tan(1/2*d*x + 1/2*c)^2 - 12*B*b*tan(1/2*d*x + 1/2*c)^2 - 4*B*a*tan(1/2*d*x + 1/2*c) - 4*A*b*tan(1/2*d*x + 1/2*c) - A*a)/tan(1/2*d*x + 1/2*c)^2)/d`

3.237.9 Mupad [B] (verification not implemented)

Time = 7.28 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.64

$$\int \cot^3(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= -\frac{\ln(\tan(c + dx)) (Aa - Bb)}{d} - \frac{\cot(c + dx)^2 \left(\frac{Aa}{2} + \tan(c + dx) (Ab + Ba)\right)}{d}$$

$$+ \frac{\ln(\tan(c + dx) - i) (A + B i) (a + b i)}{2d}$$

$$- \frac{\ln(\tan(c + dx) + i) (A - B i) (b + a i) i}{2d}$$

input `int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)`output `(log(tan(c + d*x) - 1i)*(A + B*1i)*(a + b*1i))/(2*d) - (cot(c + d*x)^2*((A*a)/2 + tan(c + d*x)*(A*b + B*a)))/d - (log(tan(c + d*x))*(A*a - B*b))/d - (log(tan(c + d*x) + 1i)*(A - B*1i)*(a*1i + b)*1i)/(2*d)`

3.238 $\int \cot^4(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

3.238.1 Optimal result	2333
3.238.2 Mathematica [C] (verified)	2333
3.238.3 Rubi [A] (verified)	2334
3.238.4 Maple [A] (verified)	2337
3.238.5 Fricas [A] (verification not implemented)	2337
3.238.6 Sympy [B] (verification not implemented)	2338
3.238.7 Maxima [A] (verification not implemented)	2338
3.238.8 Giac [B] (verification not implemented)	2339
3.238.9 Mupad [B] (verification not implemented)	2339

3.238.1 Optimal result

Integrand size = 29, antiderivative size = 87

$$\int \cot^4(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= (aA - bB)x + \frac{(aA - bB) \cot(c+dx)}{d} - \frac{(Ab + aB) \cot^2(c+dx)}{2d}$$

$$- \frac{aA \cot^3(c+dx)}{3d} - \frac{(Ab + aB) \log(\sin(c+dx))}{d}$$

```
output (A*a-B*b)*x+(A*a-B*b)*cot(d*x+c)/d-1/2*(A*b+B*a)*cot(d*x+c)^2/d-1/3*a*A*co
t(d*x+c)^3/d-(A*b+B*a)*ln(sin(d*x+c))/d
```

3.238.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.10 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.16

$$\int \cot^4(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx =$$

$$\frac{2aA \cot^3(c+dx) \text{Hypergeometric2F1}(-\frac{3}{2}, 1, -\frac{1}{2}, -\tan^2(c+dx)) + 6bB \cot(c+dx) \text{Hypergeometric2F1}(\frac{3}{2}, 1, \frac{5}{2}, -\tan^2(c+dx))}{d}$$

input `Integrate[Cot[c + d*x]^4*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `-1/6*(2*a*A*Cot[c + d*x]^3*Hypergeometric2F1[-3/2, 1, -1/2, -Tan[c + d*x]^2] + 6*b*B*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2] + 3*(A*b + a*B)*(Cot[c + d*x]^2 + 2*(Log[Cos[c + d*x]] + Log[Tan[c + d*x]])))/d`

3.238.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$, Rules used = {3042, 4074, 3042, 4012, 25, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^4(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan(c + dx)^4} dx \\
 & \quad \downarrow \text{4074} \\
 & \int \cot^3(c + dx)(Ab + aB - (aA - bB) \tan(c + dx)) dx - \frac{aA \cot^3(c + dx)}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\tan(c + dx)^3} dx - \frac{aA \cot^3(c + dx)}{3d} \\
 & \quad \downarrow \text{4012} \\
 & \int -\cot^2(c + dx)(aA - bB + (Ab + aB) \tan(c + dx)) dx - \frac{(aB + Ab) \cot^2(c + dx)}{2d} - \frac{aA \cot^3(c + dx)}{3d} \\
 & \quad \downarrow \text{25} \\
 & -\int \cot^2(c + dx)(aA - bB + (Ab + aB) \tan(c + dx)) dx - \frac{(aB + Ab) \cot^2(c + dx)}{2d} - \frac{aA \cot^3(c + dx)}{3d} \\
 & \quad \downarrow \text{3042} \\
 & -\int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\tan(c + dx)^2} dx - \frac{(aB + Ab) \cot^2(c + dx)}{2d} - \frac{aA \cot^3(c + dx)}{3d}
 \end{aligned}$$

3.238. $\int \cot^4(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \downarrow 4012 \\
& - \int \cot(c+dx)(Ab+aB-(aA-bB)\tan(c+dx))dx - \frac{(aB+Ab)\cot^2(c+dx)}{2d} + \\
& \quad \frac{(aA-bB)\cot(c+dx)}{d} - \frac{aA\cot^3(c+dx)}{3d} \\
& \downarrow 3042 \\
& - \int \frac{Ab+aB-(aA-bB)\tan(c+dx)}{\tan(c+dx)} dx - \frac{(aB+Ab)\cot^2(c+dx)}{2d} + \frac{(aA-bB)\cot(c+dx)}{d} - \\
& \quad \frac{aA\cot^3(c+dx)}{3d} \\
& \downarrow 4014 \\
& -(aB+Ab) \int \cot(c+dx)dx - \frac{(aB+Ab)\cot^2(c+dx)}{2d} + \frac{(aA-bB)\cot(c+dx)}{d} + x(aA-bB) - \\
& \quad \frac{aA\cot^3(c+dx)}{3d} \\
& \downarrow 3042 \\
& -(aB+Ab) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx - \frac{(aB+Ab)\cot^2(c+dx)}{2d} + \frac{(aA-bB)\cot(c+dx)}{d} + \\
& \quad x(aA-bB) - \frac{aA\cot^3(c+dx)}{3d} \\
& \downarrow 25 \\
& (aB+Ab) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx - \frac{(aB+Ab)\cot^2(c+dx)}{2d} + \frac{(aA-bB)\cot(c+dx)}{d} + \\
& \quad x(aA-bB) - \frac{aA\cot^3(c+dx)}{3d} \\
& \downarrow 3956 \\
& -\frac{(aB+Ab)\cot^2(c+dx)}{2d} + \frac{(aA-bB)\cot(c+dx)}{d} - \frac{(aB+Ab)\log(-\sin(c+dx))}{d} + x(aA-bB) - \\
& \quad \frac{aA\cot^3(c+dx)}{3d}
\end{aligned}$$

input `Int[Cot[c + d*x]^4*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(a*A - b*B)*x + ((a*A - b*B)*Cot[c + d*x])/d - ((A*b + a*B)*Cot[c + d*x]^2)/(2*d) - (a*A*Cot[c + d*x]^3)/(3*d) - ((A*b + a*B)*Log[-Sin[c + d*x]])/d`

3.238.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`
- rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`
- rule 4074 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.238.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.22

method	result
derivativedivides	$\frac{(Ab+Ba) \ln\left(\frac{1+\tan^2(dx+c)}{2}\right) + (aA-Bb) \arctan(\tan(dx+c)) + (-Ab-Ba) \ln(\tan(dx+c)) - \frac{Ab+Ba}{2 \tan(dx+c)^2} - \frac{-aA+Bb}{\tan(dx+c)} - \frac{a}{3 \tan(dx+c)}}{d}$
default	$\frac{(Ab+Ba) \ln\left(\frac{1+\tan^2(dx+c)}{2}\right) + (aA-Bb) \arctan(\tan(dx+c)) + (-Ab-Ba) \ln(\tan(dx+c)) - \frac{Ab+Ba}{2 \tan(dx+c)^2} - \frac{-aA+Bb}{\tan(dx+c)} - \frac{a}{3 \tan(dx+c)}}{d}$
norman	$\frac{(aA-Bb) \left(\frac{\tan^2(dx+c)}{d}\right) + (aA-Bb)x \tan^3(dx+c) - \frac{aA}{3d} - \frac{(Ab+Ba) \tan(dx+c)}{2d}}{\tan(dx+c)^3} - \frac{(Ab+Ba) \ln(\tan(dx+c))}{d} + \frac{(Ab+Ba) \ln(\tan(dx+c))}{d}$
parallelrisch	$\frac{-2A(\cot^3(dx+c))^a - 3Ab(\cot^2(dx+c)) + 6Axad - 3Ba(\cot^2(dx+c)) - 6Bbdx + 6aA \cot(dx+c) - 6A \ln(\tan(dx+c))b + 3A}{6d}$
risch	$iAbx + iBax + Aax - Bbx + \frac{2iAbc}{d} + \frac{2iaBc}{d} - \frac{2i(3iAb e^{4i(dx+c)} + 3iBa e^{4i(dx+c)} - 6Aa e^{4i(dx+c)} + 3Bb)}$

```
input int(cot(d*x+c)^4*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/d*(1/2*(A*b+B*a)*ln(1+tan(d*x+c)^2)+(A*a-B*b)*arctan(tan(d*x+c))+(-A*b-B*a)*ln(tan(d*x+c))-1/2*(A*b+B*a)/tan(d*x+c)^2-(-A*a+B*b)/tan(d*x+c)-1/3*A*a/tan(d*x+c)^3)
```

3.238.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.39

$$\int \cot^4(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{3(Ba + Ab) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^3 - 3(2(Aa - Bb)dx - Ba - Ab) \tan(dx+c)^3 - 6(Aa - Bb)dx}{6d \tan(dx+c)^3}$$

```
input integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output -1/6*(3*(B*a + A*b)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^3 - 3*(2*(A*a - B*b)*d*x - B*a - A*b)*tan(d*x + c)^3 - 6*(A*a - B*b)*tan(d*x + c)^2 + 2*A*a + 3*(B*a + A*b)*tan(d*x + c))/(d*tan(d*x + c)^3)
```

3.238. $\int \cot^4(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

3.238.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 178 vs. 2(75) = 150.

Time = 0.85 (sec) , antiderivative size = 178, normalized size of antiderivative = 2.05

$$\int \cot^4(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \tilde{\infty} A a x \\ x(A + B \tan(c))(a + b \tan(c)) \cot^4(c) \\ \tilde{\infty} A a x \\ A a x + \frac{A a}{d \tan(c + dx)} - \frac{A a}{3 d \tan^3(c + dx)} + \frac{A b \log(\tan^2(c + dx) + 1)}{2 d} - \frac{A b \log(\tan(c + dx))}{d} - \frac{A b}{2 d \tan^2(c + dx)} + \frac{B a \log(\tan^2(c + dx) + 1)}{2 d} \end{cases}$$

input `integrate(cot(d*x+c)**4*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Piecewise((zoo*A*a*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))*cot(c)**4, Eq(d, 0)), (zoo*A*a*x, Eq(c, -d*x)), (A*a*x + A*a/(d*tan(c + d*x)) - A*a/(3*d*tan(c + d*x)**3) + A*b*log(tan(c + d*x)**2 + 1)/(2*d) - A*b*log(tan(c + d*x))/d - A*b/(2*d*tan(c + d*x)**2) + B*a*log(tan(c + d*x)**2 + 1)/(2*d) - B*a*log(tan(c + d*x))/d - B*a/(2*d*tan(c + d*x)**2) - B*b*x - B*b/(d*tan(c + d*x)), True))`

3.238.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.20

$$\int \cot^4(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{6(Aa - Bb)(dx + c) + 3(Ba + Ab) \log(\tan(dx + c)^2 + 1) - 6(Ba + Ab) \log(\tan(dx + c)) + \frac{6(Aa - Bb) \tan(dx + c)}{d}}{6d}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/6*(6*(A*a - B*b)*(d*x + c) + 3*(B*a + A*b)*log(tan(d*x + c)^2 + 1) - 6*(B*a + A*b)*log(tan(d*x + c)) + (6*(A*a - B*b)*tan(d*x + c)^2 - 2*A*a - 3*(B*a + A*b)*tan(d*x + c))/tan(d*x + c)^3)/d`

3.238.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 237 vs. 2(83) = 166.

Time = 0.76 (sec) , antiderivative size = 237, normalized size of antiderivative = 2.72

$$\int \cot^4(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{Aa \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 3Ba \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 3Ab \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 15Aa \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 12Bbt}{d}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `1/24*(A*a*tan(1/2*d*x + 1/2*c)^3 - 3*B*a*tan(1/2*d*x + 1/2*c)^2 - 3*A*b*tan(1/2*d*x + 1/2*c)^2 - 15*A*a*tan(1/2*d*x + 1/2*c) + 12*B*b*tan(1/2*d*x + 1/2*c) + 24*(A*a - B*b)*(d*x + c) + 24*(B*a + A*b)*log(tan(1/2*d*x + 1/2*c)^2 + 1) - 24*(B*a + A*b)*log(abs(tan(1/2*d*x + 1/2*c))) + (44*B*a*tan(1/2*d*x + 1/2*c)^3 + 44*A*b*tan(1/2*d*x + 1/2*c)^3 + 15*A*a*tan(1/2*d*x + 1/2*c)^2 - 12*B*b*tan(1/2*d*x + 1/2*c)^2 - 3*B*a*tan(1/2*d*x + 1/2*c) - 3*A*b*tan(1/2*d*x + 1/2*c) - A*a)/tan(1/2*d*x + 1/2*c)^3)/d`

3.238.9 Mupad [B] (verification not implemented)

Time = 7.54 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.46

$$\int \cot^4(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= -\frac{\cot(c + dx)^3 ((Bb - Aa) \tan(c + dx)^2 + \left(\frac{Ab}{2} + \frac{Ba}{2}\right) \tan(c + dx) + \frac{Aa}{3})}{d} - \frac{\ln(\tan(c + dx)) (Ab + Ba)}{d} - \frac{\ln(\tan(c + dx) - i) (A + B i) (a + b i) i}{2d} + \frac{\ln(\tan(c + dx) + i) (A - B i) (b + a i)}{2d}$$

input `int(cot(c + d*x)^4*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)`

output `(log(tan(c + d*x) + i)*(A - B*i)*(a*i + b))/(2*d) - (log(tan(c + d*x))*(A*b + B*a))/d - (log(tan(c + d*x) - i)*(A + B*i)*(a + b*i)*i)/(2*d) - (cot(c + d*x)^3*((A*a)/3 + tan(c + d*x)*((A*b)/2 + (B*a)/2) - tan(c + d*x)^2*(A*a - B*b))/d`

3.239 $\int \cot^5(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

3.239.1 Optimal result	2340
3.239.2 Mathematica [C] (verified)	2340
3.239.3 Rubi [A] (verified)	2341
3.239.4 Maple [A] (verified)	2344
3.239.5 Fricas [A] (verification not implemented)	2345
3.239.6 Sympy [B] (verification not implemented)	2345
3.239.7 Maxima [A] (verification not implemented)	2346
3.239.8 Giac [B] (verification not implemented)	2346
3.239.9 Mupad [B] (verification not implemented)	2347

3.239.1 Optimal result

Integrand size = 29, antiderivative size = 108

$$\int \cot^5(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= (Ab+aB)x + \frac{(Ab+aB) \cot(c+dx)}{d} + \frac{(aA-bB) \cot^2(c+dx)}{2d}$$

$$- \frac{(Ab+aB) \cot^3(c+dx)}{3d} - \frac{aA \cot^4(c+dx)}{4d} + \frac{(aA-bB) \log(\sin(c+dx))}{d}$$

```
output (A*b+B*a)*x+(A*b+B*a)*cot(d*x+c)/d+1/2*(A*a-B*b)*cot(d*x+c)^2/d-1/3*(A*b+B
*a)*cot(d*x+c)^3/d-1/4*a*A*cot(d*x+c)^4/d+(A*a-B*b)*ln(sin(d*x+c))/d
```

3.239.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.25 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.93

$$\int \cot^5(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx =$$

$$\frac{4(Ab+aB) \cot^3(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, -\tan^2(c+dx)\right) + 3((-2aA+2bB) \cot^2(c+dx) + \dots}{12d}$$

input `Integrate[Cot[c + d*x]^5*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output
$$-1/12*(4*(A*b + a*B)*Cot[c + d*x]^3*Hypergeometric2F1[-3/2, 1, -1/2, -Tan[c + d*x]^2] + 3*((-2*a*A + 2*b*B)*Cot[c + d*x]^2 + a*A*Cot[c + d*x]^4 - 4*(a*A - b*B)*(Log[Cos[c + d*x]] + Log[Tan[c + d*x]])))/d$$

3.239.3 Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.02, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.517$, Rules used = {3042, 4074, 3042, 4012, 25, 3042, 4012, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^5(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan(c + dx)^5} dx \\
 & \quad \downarrow \text{4074} \\
 & \int \cot^4(c + dx)(Ab + aB - (aA - bB) \tan(c + dx)) dx - \frac{aA \cot^4(c + dx)}{4d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\tan(c + dx)^4} dx - \frac{aA \cot^4(c + dx)}{4d} \\
 & \quad \downarrow \text{4012} \\
 & \int -\cot^3(c + dx)(aA - bB + (Ab + aB) \tan(c + dx)) dx - \frac{(aB + Ab) \cot^3(c + dx)}{3d} - \frac{aA \cot^4(c + dx)}{4d} \\
 & \quad \downarrow \text{25} \\
 & -\int \cot^3(c + dx)(aA - bB + (Ab + aB) \tan(c + dx)) dx - \frac{(aB + Ab) \cot^3(c + dx)}{3d} - \frac{aA \cot^4(c + dx)}{4d} \\
 & \quad \downarrow \text{3042} \\
 & -\int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\tan(c + dx)^3} dx - \frac{(aB + Ab) \cot^3(c + dx)}{3d} - \frac{aA \cot^4(c + dx)}{4d}
 \end{aligned}$$

3.239. $\int \cot^5(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \downarrow 4012 \\
& - \int \cot^2(c+dx)(Ab+aB-(aA-bB)\tan(c+dx))dx - \frac{(aB+Ab)\cot^3(c+dx)}{3d} + \\
& \quad \frac{(aA-bB)\cot^2(c+dx)}{2d} - \frac{aA\cot^4(c+dx)}{4d} \\
& \downarrow 3042 \\
& - \int \frac{Ab+aB-(aA-bB)\tan(c+dx)}{\tan(c+dx)^2} dx - \frac{(aB+Ab)\cot^3(c+dx)}{3d} + \frac{(aA-bB)\cot^2(c+dx)}{2d} - \\
& \quad \frac{aA\cot^4(c+dx)}{4d} \\
& \downarrow 4012 \\
& - \int -\cot(c+dx)(aA-bB+(Ab+aB)\tan(c+dx))dx - \frac{(aB+Ab)\cot^3(c+dx)}{3d} + \\
& \quad \frac{(aA-bB)\cot^2(c+dx)}{2d} + \frac{(aB+Ab)\cot(c+dx)}{d} - \frac{aA\cot^4(c+dx)}{4d} \\
& \downarrow 25 \\
& \int \cot(c+dx)(aA-bB+(Ab+aB)\tan(c+dx))dx - \frac{(aB+Ab)\cot^3(c+dx)}{3d} + \\
& \quad \frac{(aA-bB)\cot^2(c+dx)}{2d} + \frac{(aB+Ab)\cot(c+dx)}{d} - \frac{aA\cot^4(c+dx)}{4d} \\
& \downarrow 3042 \\
& \int \frac{aA-bB+(Ab+aB)\tan(c+dx)}{\tan(c+dx)} dx - \frac{(aB+Ab)\cot^3(c+dx)}{3d} + \frac{(aA-bB)\cot^2(c+dx)}{2d} + \\
& \quad \frac{(aB+Ab)\cot(c+dx)}{d} - \frac{aA\cot^4(c+dx)}{4d} \\
& \downarrow 4014 \\
& (aA-bB) \int \cot(c+dx)dx - \frac{(aB+Ab)\cot^3(c+dx)}{3d} + \frac{(aA-bB)\cot^2(c+dx)}{2d} + \\
& \quad \frac{(aB+Ab)\cot(c+dx)}{d} + x(aB+Ab) - \frac{aA\cot^4(c+dx)}{4d} \\
& \downarrow 3042 \\
& (aA-bB) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx - \frac{(aB+Ab)\cot^3(c+dx)}{3d} + \frac{(aA-bB)\cot^2(c+dx)}{2d} + \\
& \quad \frac{(aB+Ab)\cot(c+dx)}{d} + x(aB+Ab) - \frac{aA\cot^4(c+dx)}{4d} \\
& \downarrow 25
\end{aligned}$$

$$\begin{aligned}
& -(aA - bB) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - \frac{(aB + Ab) \cot^3(c + dx)}{3d} + \frac{(aA - bB) \cot^2(c + dx)}{2d} + \\
& \quad \frac{(aB + Ab) \cot(c + dx)}{d} + x(aB + Ab) - \frac{aA \cot^4(c + dx)}{4d} \\
& \qquad \qquad \qquad \downarrow \text{3956} \\
& -\frac{(aB + Ab) \cot^3(c + dx)}{3d} + \frac{(aA - bB) \cot^2(c + dx)}{2d} + \frac{(aB + Ab) \cot(c + dx)}{d} + \\
& \quad \frac{(aA - bB) \log(-\sin(c + dx))}{d} + x(aB + Ab) - \frac{aA \cot^4(c + dx)}{4d}
\end{aligned}$$

input `Int[Cot[c + d*x]^5*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]), x]`

output `(A*b + a*B)*x + ((A*b + a*B)*Cot[c + d*x])/d + ((a*A - b*B)*Cot[c + d*x]^2)/(2*d) - ((A*b + a*B)*Cot[c + d*x]^3)/(3*d) - (a*A*Cot[c + d*x]^4)/(4*d) + ((a*A - b*B)*Log[-Sin[c + d*x]])/d`

3.239.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)
)*(x_)], x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

3.239.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.15

method	result
derivativedivides	$\frac{(-aA+Bb)\ln\left(\frac{1+\tan^2(dx+c)}{2}\right) + (Ab+Ba)\arctan(\tan(dx+c)) - \frac{-Ab-Ba}{\tan(dx+c)} - \frac{Ab+Ba}{3\tan(dx+c)^3} - \frac{-aA+Bb}{2\tan(dx+c)^2} + (aA-Bb)\ln(\tan(dx+c))}{d}$
default	$\frac{(-aA+Bb)\ln\left(\frac{1+\tan^2(dx+c)}{2}\right) + (Ab+Ba)\arctan(\tan(dx+c)) - \frac{-Ab-Ba}{\tan(dx+c)} - \frac{Ab+Ba}{3\tan(dx+c)^3} - \frac{-aA+Bb}{2\tan(dx+c)^2} + (aA-Bb)\ln(\tan(dx+c))}{d}$
norman	$\frac{(Ab+Ba)\left(\frac{\tan^3(dx+c)}{d}\right) + (Ab+Ba)x(\tan^4(dx+c)) - \frac{aA}{4d} - \frac{(Ab+Ba)\tan(dx+c)}{3d} + \frac{(aA-Bb)\left(\frac{\tan^2(dx+c)}{2d}\right)}{\tan(dx+c)^4} + \frac{(aA-Bb)\ln(\tan(dx+c))}{d}}{\tan(dx+c)^4}$
parallelrisch	$-3A(\cot^4(dx+c))a - 4Ab(\cot^3(dx+c)) - 4Ba(\cot^3(dx+c)) + 6A(\cot^2(dx+c))a + 12Abdx - 6Bb(\cot^2(dx+c)) + 12Bxad + 3A^2$
risch	$Abx + Bax - iAax + iBbx - \frac{2iaAc}{d} + \frac{2iBbc}{d} - \frac{2(-6iAbe^{6i(dx+c)} - 6iBa e^{6i(dx+c)} + 6Aa e^{6i(dx+c)} - 3B^2)}{d}$

```
input int(cot(d*x+c)^5*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE
)
```

```
output 1/d*(1/2*(-A*a+B*b)*ln(1+tan(d*x+c)^2)+(A*b+B*a)*arctan(tan(d*x+c))-(-A*b-
B*a)/tan(d*x+c)-1/3*(A*b+B*a)/tan(d*x+c)^3-1/2*(-A*a+B*b)/tan(d*x+c)^2+(A
a-B*b)*ln(tan(d*x+c))-1/4*a*A/tan(d*x+c)^4)
```

$$3.239. \quad \int \cot^5(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

3.239.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.28

$$\int \cot^5(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{6(Aa - Bb) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^4 + 3(4(Ba + Ab)dx + 3Aa - 2Bb) \tan(dx+c)^4 + 12(Ba + Ab) \tan(dx+c)^4}{12d \tan(dx+c)^4}$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/12*(6*(A*a - B*b)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^4 + 3*(4*(B*a + A*b)*d*x + 3*A*a - 2*B*b)*tan(d*x + c)^4 + 12*(B*a + A*b)*tan(d*x + c)^3 + 6*(A*a - B*b)*tan(d*x + c)^2 - 3*A*a - 4*(B*a + A*b)*tan(d*x + c))/(d*tan(d*x + c)^4)`

3.239.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 209 vs. 2(95) = 190.

Time = 1.12 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.94

$$\int \cot^5(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \tilde{\infty} Aax \\ x(A + B \tan(c))(a + b \tan(c)) \cot^5(c) \\ \tilde{\infty} Aax \\ -\frac{Aa \log(\tan^2(c+dx)+1)}{2d} + \frac{Aa \log(\tan(c+dx))}{d} + \frac{Aa}{2d \tan^2(c+dx)} - \frac{Aa}{4d \tan^4(c+dx)} + Abx + \frac{Ab}{d \tan(c+dx)} - \frac{Ab}{3d \tan^3(c+dx)} + \dots \end{cases}$$

input `integrate(cot(d*x+c)**5*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Piecewise((zoo*A*a*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))*cot(c)**5, Eq(d, 0)), (zoo*A*a*x, Eq(c, -d*x)), (-A*a*log(tan(c + d*x)**2 + 1)/(2*d) + A*a*log(tan(c + d*x))/d + A*a/(2*d*tan(c + d*x)**2) - A*a/(4*d*tan(c + d*x)**4) + A*b*x + A*b/(d*tan(c + d*x)) - A*b/(3*d*tan(c + d*x)**3) + B*a*x + B*a/(d*tan(c + d*x)) - B*a/(3*d*tan(c + d*x)**3) + B*b*log(tan(c + d*x)**2 + 1)/(2*d) - B*b*log(tan(c + d*x))/d - B*b/(2*d*tan(c + d*x)**2), True))`

3.239. $\int \cot^5(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

3.239.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.13

$$\int \cot^5(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{12(Ba + Ab)(dx + c) - 6(Aa - Bb) \log(\tan(dx + c)^2 + 1) + 12(Aa - Bb) \log(\tan(dx + c)) + \frac{12(Ba + Ab)}{12d}}{12d}$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/12*(12*(B*a + A*b)*(d*x + c) - 6*(A*a - B*b)*log(tan(d*x + c)^2 + 1) + 12*(A*a - B*b)*log(tan(d*x + c)) + (12*(B*a + A*b)*tan(d*x + c)^3 + 6*(A*a - B*b)*tan(d*x + c)^2 - 3*A*a - 4*(B*a + A*b)*tan(d*x + c))/tan(d*x + c)^4)/d`

3.239.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 299 vs. 2(102) = 204.

Time = 0.90 (sec) , antiderivative size = 299, normalized size of antiderivative = 2.77

$$\int \cot^5(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx =$$

$$\frac{3Aa \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^4 - 8Ba \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 8Ab \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 36Aa \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + 24Ab \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 3Aa}{\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^4} + \frac{12(Ba + Ab)}{12d}$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `-1/192*(3*A*a*tan(1/2*d*x + 1/2*c)^4 - 8*B*a*tan(1/2*d*x + 1/2*c)^3 - 8*A*b*tan(1/2*d*x + 1/2*c)^3 - 36*A*a*tan(1/2*d*x + 1/2*c)^2 + 24*B*b*tan(1/2*d*x + 1/2*c)^2 + 120*B*a*tan(1/2*d*x + 1/2*c) + 120*A*b*tan(1/2*d*x + 1/2*c) - 192*(B*a + A*b)*(d*x + c) + 192*(A*a - B*b)*log(tan(1/2*d*x + 1/2*c)^2 + 1) - 192*(A*a - B*b)*log(abs(tan(1/2*d*x + 1/2*c))) + (400*A*a*tan(1/2*d*x + 1/2*c)^4 - 400*B*b*tan(1/2*d*x + 1/2*c)^4 - 120*B*a*tan(1/2*d*x + 1/2*c)^3 - 120*A*b*tan(1/2*d*x + 1/2*c)^3 - 36*A*a*tan(1/2*d*x + 1/2*c)^2 + 24*B*b*tan(1/2*d*x + 1/2*c)^2 + 8*B*a*tan(1/2*d*x + 1/2*c) + 8*A*b*tan(1/2*d*x + 1/2*c) + 3*A*a)/tan(1/2*d*x + 1/2*c)^4)/d`

3.239.9 Mupad [B] (verification not implemented)

Time = 7.87 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.34

$$\int \cot^5(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \frac{\ln(\tan(c + dx))(Aa - Bb)}{d} - \frac{\cot(c + dx)^4((-Ab - Ba) \tan(c + dx)^3 + (\frac{Bb}{2} - \frac{Aa}{2}) \tan(c + dx)^2 + (\frac{Ab}{3} + \frac{Ba}{3}) \tan(c + dx) + \frac{Aa}{4})}{d} - \frac{\ln(\tan(c + dx) - i)(A + B i)(a + b i)}{2d} + \frac{\ln(\tan(c + dx) + i)(A - B i)(b + a i) i}{2d}$$

input `int(cot(c + d*x)^5*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)`output `(log(tan(c + d*x))*(A*a - B*b))/d - (cot(c + d*x)^4*((A*a)/4 + tan(c + d*x))*((A*b)/3 + (B*a)/3) - tan(c + d*x)^3*(A*b + B*a) - tan(c + d*x)^2*((A*a)/2 - (B*b)/2))/d - (log(tan(c + d*x) - 1i)*(A + B*1i)*(a + b*1i))/(2*d) + (log(tan(c + d*x) + 1i)*(A - B*1i)*(a*1i + b)*1i)/(2*d)`

3.240 $\int \tan^2(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.240.1 Optimal result	2348
3.240.2 Mathematica [C] (verified)	2349
3.240.3 Rubi [A] (verified)	2349
3.240.4 Maple [A] (verified)	2352
3.240.5 Fricas [A] (verification not implemented)	2353
3.240.6 Sympy [A] (verification not implemented)	2353
3.240.7 Maxima [A] (verification not implemented)	2354
3.240.8 Giac [B] (verification not implemented)	2354
3.240.9 Mupad [B] (verification not implemented)	2355

3.240.1 Optimal result

Integrand size = 31, antiderivative size = 148

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= -((a^2A - Ab^2 - 2abB)x) + \frac{(2aAb + a^2B - b^2B) \log(\cos(c + dx))}{d}$$

$$- \frac{b(Ab + aB) \tan(c + dx)}{d} - \frac{B(a + b \tan(c + dx))^2}{d}$$

$$+ \frac{(4Ab - aB)(a + b \tan(c + dx))^3}{12b^2d} + \frac{B \tan(c + dx)(a + b \tan(c + dx))^3}{4bd}$$

output $-(A*a^2-A*b^2-2*B*a*b)*x+(2*A*a*b+B*a^2-B*b^2)*\ln(\cos(d*x+c))/d-b*(A*b+B*a)*\tan(d*x+c)/d-1/2*B*(a+b*\tan(d*x+c))^2/d+1/12*(4*A*b-B*a)*(a+b*\tan(d*x+c))^3/b^2/d+1/4*B*\tan(d*x+c)*(a+b*\tan(d*x+c))^3/b/d$

3.240.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.22 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.49

$$\int \tan^2(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx = \frac{B \tan(c+dx)(a+b \tan(c+dx))^3}{4bd} + \frac{(4Ab-aB)(a+b \tan(c+dx))^3}{3bd} + \frac{2((Ab-aB)(i(a+ib)^2 \log(i-\tan(c+dx))-i(a-ib)^2 \log(i+\tan(c+dx))-2b^2 \tan(c+dx))-B((ia-b)^3 \log(i-\tan(c+dx))+B((ia-b)^3 \log(i+\tan(c+dx))))}{4bd}$$

input `Integrate[Tan[c + d*x]^2*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output $(B \tan[c + d*x] * (a + b \tan[c + d*x])^3) / (4 * b * d) + (((4 * A * b - a * B) * (a + b \tan[c + d*x])^3) / (3 * b * d) + (2 * ((A * b - a * B) * (I * (a + I * b)^2 * \text{Log}[I - \tan[c + d*x]] - I * (a - I * b)^2 * \text{Log}[I + \tan[c + d*x]] - 2 * b^2 * \tan[c + d*x]) - B * ((I * a - b)^3 * \text{Log}[I - \tan[c + d*x]] - (I * a + b)^3 * \text{Log}[I + \tan[c + d*x]] + 6 * a * b^2 * \tan[c + d*x] + b^3 * \tan[c + d*x]^2))) / d) / (4 * b)$

3.240.3 Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4090, 25, 3042, 4113, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^2(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

↓ 3042

$$\int \tan(c+dx)^2(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

↓ 4090

$$\int \frac{-(a+b \tan(c+dx))^2(-((4Ab-aB) \tan^2(c+dx)) + 4bB \tan(c+dx) + aB) dx}{\frac{4b}{B \tan(c+dx)(a+b \tan(c+dx))^3}} +$$

↓ 25

$$\begin{aligned}
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^3}{4bd} - \frac{\int (a+b \tan(c+dx))^2 \left(-((4Ab-aB) \tan^2(c+dx)) + 4bB \tan(c+dx) + aB \right) dx}{4b} \\
& \quad \downarrow \text{3042} \\
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^3}{4bd} - \frac{\int (a+b \tan(c+dx))^2 \left(-((4Ab-aB) \tan(c+dx)^2) + 4bB \tan(c+dx) + aB \right) dx}{4b} \\
& \quad \downarrow \text{4113} \\
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^3}{4bd} - \frac{\int (a+b \tan(c+dx))^2 (4Ab + 4B \tan(c+dx)b) dx - \frac{(4Ab-aB)(a+b \tan(c+dx))^3}{3bd}}{4b} \\
& \quad \downarrow \text{3042} \\
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^3}{4bd} - \frac{\int (a+b \tan(c+dx))^2 (4Ab + 4B \tan(c+dx)b) dx - \frac{(4Ab-aB)(a+b \tan(c+dx))^3}{3bd}}{4b} \\
& \quad \downarrow \text{4011} \\
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^3}{4bd} - \frac{\int (a+b \tan(c+dx))(4b(aA-bB) + 4b(Ab+aB) \tan(c+dx)) dx - \frac{(4Ab-aB)(a+b \tan(c+dx))^3}{3bd} + \frac{2bB(a+b \tan(c+dx))^2}{d}}{4b} \\
& \quad \downarrow \text{3042} \\
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^3}{4bd} - \frac{\int (a+b \tan(c+dx))(4b(aA-bB) + 4b(Ab+aB) \tan(c+dx)) dx - \frac{(4Ab-aB)(a+b \tan(c+dx))^3}{3bd} + \frac{2bB(a+b \tan(c+dx))^2}{d}}{4b} \\
& \quad \downarrow \text{4008} \\
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^3}{4bd} - \frac{4b(a^2B + 2aAb - b^2B) \int \tan(c+dx) dx + 4bx(a^2A - 2abB - Ab^2) + \frac{4b^2(aB+Ab) \tan(c+dx)}{d} - \frac{(4Ab-aB)(a+b \tan(c+dx))^2}{3bd}}{4b} \\
& \quad \downarrow \text{3042} \\
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^3}{4bd} - \frac{4b(a^2B + 2aAb - b^2B) \int \tan(c+dx) dx + 4bx(a^2A - 2abB - Ab^2) + \frac{4b^2(aB+Ab) \tan(c+dx)}{d} - \frac{(4Ab-aB)(a+b \tan(c+dx))^2}{3bd}}{4b}
\end{aligned}$$

3.240. $\int \tan^2(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

$$\begin{aligned} & \downarrow 3956 \\ & \frac{B \tan(c+dx)(a+b \tan(c+dx))^3}{4bd} - \\ & \frac{-\frac{4b(a^2B+2aAb-b^2B) \log(\cos(c+dx))}{d} + 4bx(a^2A - 2abB - Ab^2) + \frac{4b^2(aB+Ab) \tan(c+dx)}{d} - \frac{(4Ab-aB)(a+b \tan(c+dx))^3}{3bd} + \frac{2bB}{4b}}{4b} \end{aligned}$$

input `Int[Tan[c + d*x]^2*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]), x]`

output `(B*Tan[c + d*x]*(a + b*Tan[c + d*x])^3)/(4*b*d) - (4*b*(a^2*A - A*b^2 - 2*a*b*B)*x - (4*b*(2*a*A*b + a^2*B - b^2*B)*Log[Cos[c + d*x]])/d + (4*b^2*(A*b + a*B)*Tan[c + d*x])/d + (2*b*B*(a + b*Tan[c + d*x])^2)/d - ((4*A*b - a*B)*(a + b*Tan[c + d*x])^3)/(3*b*d))/(4*b)`

3.240.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

```
rule 4090 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4113 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

3.240.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.00

method	result
parts	$\frac{(Ab^2+2Bab)\left(\frac{\tan^3(dx+c)}{3}-\tan(dx+c)+\arctan(\tan(dx+c))\right)}{d} + \frac{(2Aab+Ba^2)\left(\frac{\tan^2(dx+c)}{2}-\frac{\ln(1+\tan^2(dx+c))}{2}\right)}{d}$
norman	$(-Aa^2 + Ab^2 + 2Bab)x + \frac{(Aa^2 - Ab^2 - 2Bab)\tan(dx+c)}{d} + \frac{(2Aab + Ba^2 - Bb^2)(\tan^2(dx+c))}{2d} + \frac{Bb^2(\tan^2(dx+c))}{2d}$
derivativedivides	$\frac{Bb^2(\tan^4(dx+c))}{4} + \frac{Ab^2(\tan^3(dx+c))}{3} + \frac{2Bab(\tan^3(dx+c))}{3} + Aab(\tan^2(dx+c)) + \frac{B(\tan^2(dx+c))a^2}{2} - \frac{Bb^2(\tan^2(dx+c))}{2} + A$
default	$\frac{Bb^2(\tan^4(dx+c))}{4} + \frac{Ab^2(\tan^3(dx+c))}{3} + \frac{2Bab(\tan^3(dx+c))}{3} + Aab(\tan^2(dx+c)) + \frac{B(\tan^2(dx+c))a^2}{2} - \frac{Bb^2(\tan^2(dx+c))}{2} + A$
parallelrisc	$-3Bb^2(\tan^4(dx+c)) - 4Ab^2(\tan^3(dx+c)) - 8Bab(\tan^3(dx+c)) + 12Axa^2d - 12Ab^2dx - 12Aab(\tan^2(dx+c)) - 24Bab$
risc	$iBb^2x - \frac{2ia^2Bc}{d} - \frac{4iAabc}{d} - Aa^2x + Ab^2x + 2Babx - iBa^2x + \frac{2iBb^2c}{d} - 2iAabx + \frac{-4iA$

```
input int(tan(d*x+c)^2*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

3.240. $\int \tan^2(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

output $(A*b^2+2*B*a*b)/d*(1/3*\tan(dx+c)^3-\tan(dx+c)+\arctan(\tan(dx+c)))+(2*A*a*b+B*a^2)/d*(1/2*\tan(dx+c)^2-1/2*\ln(1+\tan(dx+c)^2))+A*a^2/d*(\tan(dx+c)-\arctan(\tan(dx+c)))+B*b^2/d*(1/4*\tan(dx+c)^4-1/2*\tan(dx+c)^2+1/2*\ln(1+\tan(dx+c)^2))$

3.240.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.99

$$\int \tan^2(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

$$= \frac{3Bb^2 \tan(dx+c)^4 + 4(2Bab + Ab^2) \tan(dx+c)^3 - 12(Aa^2 - 2Bab - Ab^2)dx + 6(Ba^2 + 2Aab - Bb^2)}{12d}$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output $1/12*(3*B*b^2*\tan(dx+c)^4 + 4*(2*B*a*b + A*b^2)*\tan(dx+c)^3 - 12*(A*a^2 - 2*B*a*b - A*b^2)*dx + 6*(B*a^2 + 2*A*a*b - B*b^2)*\tan(dx+c)^2 + 6*(B*a^2 + 2*A*a*b - B*b^2)*\log(1/(\tan(dx+c)^2 + 1)) + 12*(A*a^2 - 2*B*a*b - A*b^2)*\tan(dx+c))/d$

3.240.6 Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.66

$$\int \tan^2(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

$$= \begin{cases} -Aa^2x + \frac{Aa^2 \tan(c+dx)}{d} - \frac{Aab \log(\tan^2(c+dx)+1)}{d} + \frac{Aab \tan^2(c+dx)}{d} + Ab^2x + \frac{Ab^2 \tan^3(c+dx)}{3d} - \frac{Ab^2 \tan(c+dx)}{d} - \frac{Ba^2}{d} \\ x(A+B\tan(c))(a+b\tan(c))^2 \tan^2(c) \end{cases}$$

input `integrate(tan(d*x+c)**2*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `Piecewise((-A*a**2*x + A*a**2*tan(c + d*x)/d - A*a*b*log(tan(c + d*x)**2 + 1)/d + A*a*b*tan(c + d*x)**2/d + A*b**2*x + A*b**2*tan(c + d*x)**3/(3*d) - A*b**2*tan(c + d*x)/d - B*a**2*log(tan(c + d*x)**2 + 1)/(2*d) + B*a**2*tan(c + d*x)**2/(2*d) + 2*B*a*b*x + 2*B*a*b*tan(c + d*x)**3/(3*d) - 2*B*a*b*tan(c + d*x)/d + B*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + B*b**2*tan(c + d*x)**4/(4*d) - B*b**2*tan(c + d*x)**2/(2*d), Ne(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**2*tan(c)**2, True))`

3.240.7 Maxima [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.99

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{3 B b^2 \tan(dx + c)^4 + 4(2 B a b + A b^2) \tan(dx + c)^3 + 6(B a^2 + 2 A a b - B b^2) \tan(dx + c)^2 - 12(A a^2 - 2$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/12*(3*B*b^2*tan(d*x + c)^4 + 4*(2*B*a*b + A*b^2)*tan(d*x + c)^3 + 6*(B*a^2 + 2*A*a*b - B*b^2)*tan(d*x + c)^2 - 12*(A*a^2 - 2*B*a*b - A*b^2)*(d*x + c) - 6*(B*a^2 + 2*A*a*b - B*b^2)*log(tan(d*x + c)^2 + 1) + 12*(A*a^2 - 2*B*a*b - A*b^2)*tan(d*x + c))/d`

3.240.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2078 vs. 2(141) = 282.

Time = 1.84 (sec) , antiderivative size = 2078, normalized size of antiderivative = 14.04

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

```
-1/12*(12*A*a^2*d*x*tan(d*x)^4*tan(c)^4 - 24*B*a*b*d*x*tan(d*x)^4*tan(c)^4
- 12*A*b^2*d*x*tan(d*x)^4*tan(c)^4 - 6*B*a^2*log(4*(tan(d*x)^2*tan(c)^2 -
2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))
*tan(d*x)^4*tan(c)^4 - 12*A*a*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*ta
n(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^4*ta
n(c)^4 + 6*B*b^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(
d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^4*tan(c)^4 - 48*A*a
^2*d*x*tan(d*x)^3*tan(c)^3 + 96*B*a*b*d*x*tan(d*x)^3*tan(c)^3 + 48*A*b^2*d
*x*tan(d*x)^3*tan(c)^3 - 6*B*a^2*tan(d*x)^4*tan(c)^4 - 12*A*a*b*tan(d*x)^4
*tan(c)^4 + 9*B*b^2*tan(d*x)^4*tan(c)^4 + 24*B*a^2*log(4*(tan(d*x)^2*tan(c)
)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2
+ 1))*tan(d*x)^3*tan(c)^3 + 48*A*a*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*
x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)
^3*tan(c)^3 - 24*B*b^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)
/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^3*tan(c)^3 +
12*A*a^2*tan(d*x)^4*tan(c)^3 - 24*B*a*b*tan(d*x)^4*tan(c)^3 - 12*A*b^2*tan
(d*x)^4*tan(c)^3 + 12*A*a^2*tan(d*x)^3*tan(c)^4 - 24*B*a*b*tan(d*x)^3*tan(
c)^4 - 12*A*b^2*tan(d*x)^3*tan(c)^4 + 72*A*a^2*d*x*tan(d*x)^2*tan(c)^2 - 1
44*B*a*b*d*x*tan(d*x)^2*tan(c)^2 - 72*A*b^2*d*x*tan(d*x)^2*tan(c)^2 - 6*B*
a^2*tan(d*x)^4*tan(c)^2 - 12*A*a*b*tan(d*x)^4*tan(c)^2 + 6*B*b^2*tan(d*...
```

3.240.9 Mupad [B] (verification not implemented)

Time = 7.33 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.02

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= x(-Aa^2 + 2Bab + Ab^2) + \frac{\tan(c + dx)^3 \left(\frac{Ab^2}{3} + \frac{2Bab}{3} \right)}{d}$$

$$- \frac{\tan(c + dx)(-Aa^2 + 2Bab + Ab^2)}{d} - \frac{\ln(\tan(c + dx)^2 + 1) \left(\frac{Ba^2}{2} + Aab - \frac{Bb^2}{2} \right)}{d}$$

$$+ \frac{\tan(c + dx)^2 \left(\frac{Ba^2}{2} + Aab - \frac{Bb^2}{2} \right)}{d} + \frac{Bb^2 \tan(c + dx)^4}{4d}$$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)`

output $x*(A*b^2 - A*a^2 + 2*B*a*b) + (\tan(c + d*x)^3*((A*b^2)/3 + (2*B*a*b)/3))/d$
 $- (\tan(c + d*x)*(A*b^2 - A*a^2 + 2*B*a*b))/d - (\log(\tan(c + d*x)^2 + 1)*$
 $(B*a^2)/2 - (B*b^2)/2 + A*a*b))/d + (\tan(c + d*x)^2*((B*a^2)/2 - (B*b^2)/2$
 $+ A*a*b))/d + (B*b^2*\tan(c + d*x)^4)/(4*d)$

3.241 $\int \tan(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.241.1 Optimal result	2357
3.241.2 Mathematica [C] (verified)	2357
3.241.3 Rubi [A] (verified)	2358
3.241.4 Maple [A] (verified)	2360
3.241.5 Fricas [A] (verification not implemented)	2361
3.241.6 Sympy [A] (verification not implemented)	2361
3.241.7 Maxima [A] (verification not implemented)	2362
3.241.8 Giac [B] (verification not implemented)	2362
3.241.9 Mupad [B] (verification not implemented)	2363

3.241.1 Optimal result

Integrand size = 29, antiderivative size = 112

$$\int \tan(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= -((2aAb + a^2B - b^2B) x) - \frac{(a^2A - Ab^2 - 2abB) \log(\cos(c+dx))}{d}$$

$$+ \frac{b(aA - bB) \tan(c+dx)}{d} + \frac{A(a+b \tan(c+dx))^2}{2d} + \frac{B(a+b \tan(c+dx))^3}{3bd}$$

```
output - (2*A*a*b+B*a^2-B*b^2)*x - (A*a^2-A*b^2-2*B*a*b)*ln(cos(d*x+c))/d + b*(A*a-B*b)
        *tan(d*x+c)/d + 1/2*A*(a+b*tan(d*x+c))^2/d + 1/3*B*(a+b*tan(d*x+c))^3/b/d
```

3.241.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.91 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.54

$$\int \tan(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{2B(a+b \tan(c+dx))^3 + 3(aA + bB) (i((a+ib)^2 \log(i - \tan(c+dx)) - (a-ib)^2 \log(i + \tan(c+dx))) -$$

input `Integrate[Tan[c + d*x]*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output $(2*B*(a + b*\text{Tan}[c + d*x])^3 + 3*(a*A + b*B)*(I*((a + I*b)^2*\text{Log}[I - \text{Tan}[c + d*x]] - (a - I*b)^2*\text{Log}[I + \text{Tan}[c + d*x]]) - 2*b^2*\text{Tan}[c + d*x]) + 3*A*(I*a - b)^3*\text{Log}[I - \text{Tan}[c + d*x]] - (I*a + b)^3*\text{Log}[I + \text{Tan}[c + d*x]] + 6*a*b^2*\text{Tan}[c + d*x] + b^3*\text{Tan}[c + d*x]^2)/(6*b*d)$

3.241.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 4075, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4075} \\
 & \int (A \tan(c + dx) - B)(a + b \tan(c + dx))^2 dx + \frac{B(a + b \tan(c + dx))^3}{3bd} \\
 & \quad \downarrow \text{3042} \\
 & \int (A \tan(c + dx) - B)(a + b \tan(c + dx))^2 dx + \frac{B(a + b \tan(c + dx))^3}{3bd} \\
 & \quad \downarrow \text{4011} \\
 & \int (a + b \tan(c + dx))(-Ab - aB + (aA - bB) \tan(c + dx)) dx + \frac{A(a + b \tan(c + dx))^2}{2d} + \\
 & \quad \frac{B(a + b \tan(c + dx))^3}{3bd} \\
 & \quad \downarrow \text{3042} \\
 & \int (a + b \tan(c + dx))(-Ab - aB + (aA - bB) \tan(c + dx)) dx + \frac{A(a + b \tan(c + dx))^2}{2d} + \\
 & \quad \frac{B(a + b \tan(c + dx))^3}{3bd}
 \end{aligned}$$

3.241. $\int \tan(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \downarrow 4008 \\
& (a^2A - 2abB - Ab^2) \int \tan(c + dx) dx - x(a^2B + 2aAb - b^2B) + \frac{b(aA - bB) \tan(c + dx)}{d} + \\
& \quad \frac{A(a + b \tan(c + dx))^2}{2d} + \frac{B(a + b \tan(c + dx))^3}{3bd} \\
& \downarrow 3042 \\
& (a^2A - 2abB - Ab^2) \int \tan(c + dx) dx - x(a^2B + 2aAb - b^2B) + \frac{b(aA - bB) \tan(c + dx)}{d} + \\
& \quad \frac{A(a + b \tan(c + dx))^2}{2d} + \frac{B(a + b \tan(c + dx))^3}{3bd} \\
& \downarrow 3956 \\
& -\frac{(a^2A - 2abB - Ab^2) \log(\cos(c + dx))}{d} - x(a^2B + 2aAb - b^2B) + \frac{b(aA - bB) \tan(c + dx)}{d} + \\
& \quad \frac{A(a + b \tan(c + dx))^2}{2d} + \frac{B(a + b \tan(c + dx))^3}{3bd}
\end{aligned}$$

input `Int[Tan[c + d*x]*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `-((2*a*A*b + a^2*B - b^2*B)*x) - ((a^2*A - A*b^2 - 2*a*b*B)*Log[Cos[c + d*x]])/d + (b*(a*A - b*B)*Tan[c + d*x])/d + (A*(a + b*Tan[c + d*x])^2)/(2*d) + (B*(a + b*Tan[c + d*x])^3)/(3*b*d)`

3.241.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

```
rule 4011 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

3.241.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.07

method	result
norman	$(-2Aab - B a^2 + B b^2) x + \frac{(2Aab+B a^2-B b^2) \tan(dx+c)}{d} + \frac{B b^2 (\tan^3(dx+c))}{3d} + \frac{b(Ab+2Ba)(\tan^2(dx+c))}{2d}$
parts	$\frac{(A b^2+2Bab) \left(\frac{\tan^2(dx+c)}{2} - \frac{\ln(1+\tan^2(dx+c))}{2} \right)}{d} + \frac{(2Aab+B a^2)(\tan(dx+c)-\arctan(\tan(dx+c)))}{d} + \frac{A \ln(1+\tan^2(dx+c))}{2d}$
derivativedivides	$\frac{\frac{B b^2 (\tan^3(dx+c))}{3} + \frac{A b^2 (\tan^2(dx+c))}{2} + Bab(\tan^2(dx+c)) + 2Aab \tan(dx+c) + B \tan(dx+c) a^2 - B b^2 \tan(dx+c) + \frac{(A a^2 - A b^2) \ln(1 + \tan^2(dx+c))}{2}}{d}$
default	$\frac{\frac{B b^2 (\tan^3(dx+c))}{3} + \frac{A b^2 (\tan^2(dx+c))}{2} + Bab(\tan^2(dx+c)) + 2Aab \tan(dx+c) + B \tan(dx+c) a^2 - B b^2 \tan(dx+c) + \frac{(A a^2 - A b^2) \ln(1 + \tan^2(dx+c))}{2}}{d}$
parallelrisch	$\frac{2B b^2 (\tan^3(dx+c)) - 12Aabd x + 3A b^2 (\tan^2(dx+c)) - 6B x a^2 d + 6B b^2 dx + 6Bab (\tan^2(dx+c)) + 3A \ln(1 + \tan^2(dx+c)) a^2}{6d}$
risch	$-2Aabx - B a^2 x + B b^2 x + \frac{2i(-3iA b^2 e^{4i(dx+c)} - 6iBab e^{4i(dx+c)} + 6Aab e^{4i(dx+c)} + 3B a^2 e^{4i(dx+c)} - 6B b^2)}{6d}$

```
input int(tan(d*x+c)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output (-2*A*a*b-B*a^2+B*b^2)*x+(2*A*a*b+B*a^2-B*b^2)/d*tan(d*x+c)+1/3*B*b^2/d*tan
(d*x+c)^3+1/2*b*(A*b+2*B*a)/d*tan(d*x+c)^2+1/2*(A*a^2-A*b^2-2*B*a*b)/d*ln
(1+tan(d*x+c)^2)
```

3.241. $\int \tan(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.241.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.06

$$\int \tan(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{2 B b^2 \tan(dx + c)^3 - 6 (B a^2 + 2 A a b - B b^2) dx + 3 (2 B a b + A b^2) \tan(dx + c)^2 - 3 (A a^2 - 2 B a b - A b^2)}{6 d}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/6*(2*B*b^2*tan(d*x + c)^3 - 6*(B*a^2 + 2*A*a*b - B*b^2)*d*x + 3*(2*B*a*b + A*b^2)*tan(d*x + c)^2 - 3*(A*a^2 - 2*B*a*b - A*b^2)*log(1/(tan(d*x + c)^2 + 1)) + 6*(B*a^2 + 2*A*a*b - B*b^2)*tan(d*x + c))/d`

3.241.6 Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.71

$$\int \tan(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \frac{A a^2 \log(\tan^2(c+dx)+1)}{2d} - 2 A a b x + \frac{2 A a b \tan(c+dx)}{d} - \frac{A b^2 \log(\tan^2(c+dx)+1)}{2d} + \frac{A b^2 \tan^2(c+dx)}{2d} - B a^2 x + \frac{B a^2 \tan(c+dx)}{d} \\ x(A + B \tan(c))(a + b \tan(c))^2 \tan(c) \end{cases}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `Piecewise((A*a**2*log(tan(c + d*x)**2 + 1)/(2*d) - 2*A*a*b*x + 2*A*a*b*tan(c + d*x)/d - A*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + A*b**2*tan(c + d*x)**2/(2*d) - B*a**2*x + B*a**2*tan(c + d*x)/d - B*a*b*log(tan(c + d*x)**2 + 1)/d + B*a*b*tan(c + d*x)**2/d + B*b**2*x + B*b**2*tan(c + d*x)**3/(3*d) - B*b**2*tan(c + d*x)/d, Ne(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**2*tan(c), True))`

3.241.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.07

$$\int \tan(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{2 B b^2 \tan(dx + c)^3 + 3(2 B a b + A b^2) \tan(dx + c)^2 - 6(B a^2 + 2 A a b - B b^2)(dx + c) + 3(A a^2 - 2 B a b - A b^2) \log(\tan(dx + c)^2 + 1)}{6 d}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/6*(2*B*b^2*tan(d*x + c)^3 + 3*(2*B*a*b + A*b^2)*tan(d*x + c)^2 - 6*(B*a^2 + 2*A*a*b - B*b^2)*(d*x + c) + 3*(A*a^2 - 2*B*a*b - A*b^2)*log(tan(d*x + c)^2 + 1) + 6*(B*a^2 + 2*A*a*b - B*b^2)*tan(d*x + c))/d`

3.241.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1389 vs. 2(108) = 216.

Time = 1.05 (sec) , antiderivative size = 1389, normalized size of antiderivative = 12.40

$$\int \tan(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

```

-1/6*(6*B*a^2*d*x*tan(d*x)^3*tan(c)^3 + 12*A*a*b*d*x*tan(d*x)^3*tan(c)^3 -
6*B*b^2*d*x*tan(d*x)^3*tan(c)^3 + 3*A*a^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*
tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*ta
n(d*x)^3*tan(c)^3 - 6*B*a*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c)
+ 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^3*tan(c)
^3 - 3*A*b^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)
^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^3*tan(c)^3 - 18*B*a^2*d
*x*tan(d*x)^2*tan(c)^2 - 36*A*a*b*d*x*tan(d*x)^2*tan(c)^2 + 18*B*b^2*d*x*t
an(d*x)^2*tan(c)^2 - 6*B*a*b*tan(d*x)^3*tan(c)^3 - 3*A*b^2*tan(d*x)^3*tan(
c)^3 - 9*A*a^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*
x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^2*tan(c)^2 + 18*B*a*b
*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2
+ tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^2*tan(c)^2 + 9*A*b^2*log(4*(tan(d*x)
)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 +
tan(c)^2 + 1))*tan(d*x)^2*tan(c)^2 + 6*B*a^2*tan(d*x)^3*tan(c)^2 + 12*A*a*
b*tan(d*x)^3*tan(c)^2 - 6*B*b^2*tan(d*x)^3*tan(c)^2 + 6*B*a^2*tan(d*x)^2*t
an(c)^3 + 12*A*a*b*tan(d*x)^2*tan(c)^3 - 6*B*b^2*tan(d*x)^2*tan(c)^3 + 18*
B*a^2*d*x*tan(d*x)*tan(c) + 36*A*a*b*d*x*tan(d*x)*tan(c) - 18*B*b^2*d*x*ta
n(d*x)*tan(c) - 6*B*a*b*tan(d*x)^3*tan(c) - 3*A*b^2*tan(d*x)^3*tan(c) + 6*
B*a*b*tan(d*x)^2*tan(c)^2 + 3*A*b^2*tan(d*x)^2*tan(c)^2 - 6*B*a*b*tan(d...

```

3.241.9 Mupad [B] (verification not implemented)

Time = 7.32 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.08

$$\begin{aligned}
 & \int \tan(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 &= \frac{\tan(c + dx)^2 \left(\frac{Ab^2}{2} + B a b \right)}{d} - x (B a^2 + 2 A a b - B b^2) \\
 &+ \frac{\tan(c + dx) (B a^2 + 2 A a b - B b^2)}{d} \\
 &- \frac{\ln(\tan(c + dx)^2 + 1) \left(-\frac{A a^2}{2} + B a b + \frac{A b^2}{2} \right)}{d} + \frac{B b^2 \tan(c + dx)^3}{3 d}
 \end{aligned}$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)`

output `(tan(c + d*x)^2*((A*b^2)/2 + B*a*b))/d - x*(B*a^2 - B*b^2 + 2*A*a*b) + (tan(c + d*x)*(B*a^2 - B*b^2 + 2*A*a*b))/d - (log(tan(c + d*x)^2 + 1))*((A*b^2)/2 - (A*a^2)/2 + B*a*b))/d + (B*b^2*tan(c + d*x)^3)/(3*d)`

3.241. $\int \tan(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.242 $\int (a + b \tan(c + dx))^2 (A + B \tan(c + dx)) dx$

3.242.1 Optimal result	2364
3.242.2 Mathematica [C] (verified)	2364
3.242.3 Rubi [A] (verified)	2365
3.242.4 Maple [A] (verified)	2367
3.242.5 Fricas [A] (verification not implemented)	2367
3.242.6 Sympy [A] (verification not implemented)	2368
3.242.7 Maxima [A] (verification not implemented)	2368
3.242.8 Giac [B] (verification not implemented)	2369
3.242.9 Mupad [B] (verification not implemented)	2369

3.242.1 Optimal result

Integrand size = 23, antiderivative size = 87

$$\int (a + b \tan(c + dx))^2 (A + B \tan(c + dx)) dx$$

$$= (a^2 A - Ab^2 - 2abB) x - \frac{(2aAb + a^2 B - b^2 B) \log(\cos(c + dx))}{d}$$

$$+ \frac{b(Ab + aB) \tan(c + dx)}{d} + \frac{B(a + b \tan(c + dx))^2}{2d}$$

output `(A*a^2-A*b^2-2*B*a*b)*x-(2*A*a*b+B*a^2-B*b^2)*ln(cos(d*x+c))/d+b*(A*b+B*a)*tan(d*x+c)/d+1/2*B*(a+b*tan(d*x+c))^2/d`

3.242.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.48 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.10

$$\int (a + b \tan(c + dx))^2 (A + B \tan(c + dx)) dx$$

$$= \frac{(a + ib)^2 (-iA + B) \log(i - \tan(c + dx)) + (a - ib)^2 (iA + B) \log(i + \tan(c + dx)) + 2b(Ab + 2aB) \tan(c + dx)}{2d}$$

input `Integrate[(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output $((a + I*b)^2*((-I)*A + B)*\text{Log}[I - \text{Tan}[c + d*x]] + (a - I*b)^2*(I*A + B)*\text{Log}[I + \text{Tan}[c + d*x]] + 2*b*(A*b + 2*a*B)*\text{Tan}[c + d*x] + b^2*B*\text{Tan}[c + d*x]^2)/(2*d)$

3.242.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \tan(c + dx))^2 (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + b \tan(c + dx))^2 (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4011} \\ & \int (a + b \tan(c + dx))(aA - bB + (Ab + aB) \tan(c + dx)) dx + \frac{B(a + b \tan(c + dx))^2}{2d} \\ & \quad \downarrow \text{3042} \\ & \int (a + b \tan(c + dx))(aA - bB + (Ab + aB) \tan(c + dx)) dx + \frac{B(a + b \tan(c + dx))^2}{2d} \\ & \quad \downarrow \text{4008} \\ & (a^2B + 2aAb - b^2B) \int \tan(c + dx) dx + x(a^2A - 2abB - Ab^2) + \frac{b(aB + Ab) \tan(c + dx)}{d} + \\ & \quad \frac{B(a + b \tan(c + dx))^2}{2d} \\ & \quad \downarrow \text{3042} \\ & (a^2B + 2aAb - b^2B) \int \tan(c + dx) dx + x(a^2A - 2abB - Ab^2) + \frac{b(aB + Ab) \tan(c + dx)}{d} + \\ & \quad \frac{B(a + b \tan(c + dx))^2}{2d} \\ & \quad \downarrow \text{3956} \end{aligned}$$

$$-\frac{(a^2B + 2aAb - b^2B) \log(\cos(c + dx))}{d} + x \frac{(a^2A - 2abB - Ab^2)}{B(a + b \tan(c + dx))^2} + \frac{b(aB + Ab) \tan(c + dx)}{d} +$$

input `Int[(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(a^2*A - A*b^2 - 2*a*b*B)*x - ((2*a*A*b + a^2*B - b^2*B)*Log[Cos[c + d*x]])/d + (b*(A*b + a*B)*Tan[c + d*x])/d + (B*(a + b*Tan[c + d*x])^2)/(2*d)`

3.242.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

3.242.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.03

method	result
norman	$(A a^2 - A b^2 - 2Bab) x + \frac{b(Ab+2Ba) \tan(dx+c)}{d} + \frac{B b^2 (\tan^2(dx+c))}{2d} + \frac{(2Aab+B a^2 - B b^2) \ln(1+\tan^2(dx+c))}{2d}$
derivativedivides	$\frac{\frac{B b^2 (\tan^2(dx+c))}{2} + A b^2 \tan(dx+c) + 2Bab \tan(dx+c) + \frac{(2Aab+B a^2 - B b^2) \ln(1+\tan^2(dx+c))}{2}}{d} + (A a^2 - A b^2 - 2Bab) \arctan(\frac{\tan(dx+c)}{1})$
default	$\frac{\frac{B b^2 (\tan^2(dx+c))}{2} + A b^2 \tan(dx+c) + 2Bab \tan(dx+c) + \frac{(2Aab+B a^2 - B b^2) \ln(1+\tan^2(dx+c))}{2}}{d} + (A a^2 - A b^2 - 2Bab) \arctan(\frac{\tan(dx+c)}{1})$
parts	$A a^2 x + \frac{(A b^2 + 2Bab)(\tan(dx+c) - \arctan(\tan(dx+c)))}{d} + \frac{(2Aab + B a^2) \ln(1 + \tan^2(dx+c))}{2d} + \frac{B b^2 \left(\frac{\tan^2(dx+c)}{2} \right)}{d}$
parallelrisch	$\frac{2Ax a^2 d - 2A b^2 dx - 4Babd x + B b^2 (\tan^2(dx+c)) + 2A \ln(1 + \tan^2(dx+c)) ab + 2A b^2 \tan(dx+c) + B \ln(1 + \tan^2(dx+c)) a^2}{2d}$
risch	$\frac{4iAabc}{d} - \frac{2iB b^2 c}{d} + \frac{2ia^2 Bc}{d} + A a^2 x - A b^2 x - 2Babx + \frac{2ib(Ab e^{2i(dx+c)} + 2Ba e^{2i(dx+c)} - iBb e^{2i(dx+c)})}{d(e^{2i(dx+c)} + 1)^2}$

input `int((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `(A*a^2-A*b^2-2*B*a*b)*x+b*(A*b+2*B*a)/d*tan(d*x+c)+1/2*B*b^2/d*tan(d*x+c)^2+1/2*(2*A*a*b+B*a^2-B*b^2)/d*ln(1+tan(d*x+c)^2)`

3.242.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.05

$$\int (a + b \tan(c + dx))^2 (A + B \tan(c + dx)) dx$$

$$= \frac{B b^2 \tan(dx+c)^2 + 2(A a^2 - 2Bab - A b^2) dx - (B a^2 + 2Aab - B b^2) \log\left(\frac{1}{\tan(dx+c)^2 + 1}\right) + 2(2Bab + A b^2) \tan(dx+c)}{2d}$$

input `integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/2*(B*b^2*tan(d*x + c)^2 + 2*(A*a^2 - 2*B*a*b - A*b^2)*d*x - (B*a^2 + 2*A*a*b - B*b^2)*log(1/(tan(d*x + c)^2 + 1)) + 2*(2*B*a*b + A*b^2)*tan(d*x + c))/d`

3.242.6 Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.64

$$\int (a + b \tan(c + dx))^2 (A + B \tan(c + dx)) dx$$

$$= \begin{cases} Aa^2x + \frac{Aab \log(\tan^2(c+dx)+1)}{d} - Ab^2x + \frac{Ab^2 \tan(c+dx)}{d} + \frac{Ba^2 \log(\tan^2(c+dx)+1)}{2d} - 2Babx + \frac{2Bab \tan(c+dx)}{d} - \frac{Bb^2}{d} \\ x(A + B \tan(c)) (a + b \tan(c))^2 \end{cases}$$

input `integrate((a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`output `Piecewise((A*a**2*x + A*a*b*log(tan(c + d*x)**2 + 1)/d - A*b**2*x + A*b**2*tan(c + d*x)/d + B*a**2*log(tan(c + d*x)**2 + 1)/(2*d) - 2*B*a*b*x + 2*B*a*b*tan(c + d*x)/d - B*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + B*b**2*tan(c + d*x)**2/(2*d), Ne(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**2, True))`**3.242.7 Maxima [A] (verification not implemented)**

Time = 0.38 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.05

$$\int (a + b \tan(c + dx))^2 (A + B \tan(c + dx)) dx$$

$$= \frac{Bb^2 \tan(dx + c)^2 + 2(Aa^2 - 2Bab - Ab^2)(dx + c) + (Ba^2 + 2Aab - Bb^2) \log(\tan(dx + c)^2 + 1) + 2(2Aab + Ab^2) \tan(dx + c)}{2d}$$

input `integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`output `1/2*(B*b^2*tan(d*x + c)^2 + 2*(A*a^2 - 2*B*a*b - A*b^2)*(d*x + c) + (B*a^2 + 2*A*a*b - B*b^2)*log(tan(d*x + c)^2 + 1) + 2*(2*B*a*b + A*b^2)*tan(d*x + c))/d`

3.242.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 811 vs. 2(85) = 170.

Time = 0.70 (sec) , antiderivative size = 811, normalized size of antiderivative = 9.32

$$\int (a + b \tan(c + dx))^2 (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

```

1/2*(2*A*a^2*d*x*tan(d*x)^2*tan(c)^2 - 4*B*a*b*d*x*tan(d*x)^2*tan(c)^2 - 2
*A*b^2*d*x*tan(d*x)^2*tan(c)^2 - B*a^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(
d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*
x)^2*tan(c)^2 - 2*A*a*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1
)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^2*tan(c)^2 +
B*b^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan
(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^2*tan(c)^2 - 4*A*a^2*d*x*tan(
d*x)*tan(c) + 8*B*a*b*d*x*tan(d*x)*tan(c) + 4*A*b^2*d*x*tan(d*x)*tan(c) +
B*b^2*tan(d*x)^2*tan(c)^2 + 2*B*a^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x
)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)*
tan(c) + 4*A*a*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(
d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)*tan(c) - 2*B*b^2*lo
g(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + t
an(d*x)^2 + tan(c)^2 + 1))*tan(d*x)*tan(c) - 4*B*a*b*tan(d*x)^2*tan(c) - 2
*A*b^2*tan(d*x)^2*tan(c) - 4*B*a*b*tan(d*x)*tan(c)^2 - 2*A*b^2*tan(d*x)*ta
n(c)^2 + 2*A*a^2*d*x - 4*B*a*b*d*x - 2*A*b^2*d*x + B*b^2*tan(d*x)^2 + B*b^
2*tan(c)^2 - B*a^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(ta
n(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1)) - 2*A*a*b*log(4*(tan(d*x)^
2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + ta
n(c)^2 + 1)) + B*b^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1...

```

3.242.9 Mupad [B] (verification not implemented)

Time = 8.35 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.05

$$\int (a + b \tan(c + dx))^2 (A + B \tan(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx)^2 + 1) \left(\frac{B a^2}{2} + A a b - \frac{B b^2}{2} \right)}{d} - x (-A a^2 + 2 B a b + A b^2)$$

$$+ \frac{\tan(c + dx) (A b^2 + 2 B a b)}{d} + \frac{B b^2 \tan(c + dx)^2}{2 d}$$

input `int((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)`

output `(log(tan(c + d*x)^2 + 1)*((B*a^2)/2 - (B*b^2)/2 + A*a*b))/d - x*(A*b^2 - A*a^2 + 2*B*a*b) + (tan(c + d*x)*(A*b^2 + 2*B*a*b))/d + (B*b^2*tan(c + d*x)^2)/(2*d)`

3.243 $\int \cot(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.243.1 Optimal result	2371
3.243.2 Mathematica [C] (verified)	2371
3.243.3 Rubi [A] (verified)	2372
3.243.4 Maple [A] (verified)	2374
3.243.5 Fricas [A] (verification not implemented)	2374
3.243.6 Sympy [A] (verification not implemented)	2375
3.243.7 Maxima [A] (verification not implemented)	2375
3.243.8 Giac [A] (verification not implemented)	2376
3.243.9 Mupad [B] (verification not implemented)	2376

3.243.1 Optimal result

Integrand size = 29, antiderivative size = 70

$$\int \cot(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= (2aAb + a^2B - b^2B) x - \frac{b(Ab + 2aB) \log(\cos(c+dx))}{d}$$

$$+ \frac{a^2A \log(\sin(c+dx))}{d} + \frac{b^2B \tan(c+dx)}{d}$$

```
output (2*A*a*b+B*a^2-B*b^2)*x-b*(A*b+2*B*a)*ln(cos(d*x+c))/d+a^2*A*ln(sin(d*x+c)
)/d+b^2*B*tan(d*x+c)/d
```

3.243.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.32 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.33

$$\int \cot(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx =$$

$$\frac{(a+ib)^2(A+iB) \log(i-\tan(c+dx)) - 2a^2A \log(\tan(c+dx)) + (a-ib)^2(A-iB) \log(i+\tan(c+dx))}{2d}$$

input `Integrate[Cot[c + d*x]*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output
$$\frac{-1/2*((a + I*b)^2*(A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]] - 2*a^2*A*\text{Log}[\text{Tan}[c + d*x]] + (a - I*b)^2*(A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]] - 2*b*B*(a + b*\text{Tan}[c + d*x]))}{d}$$

3.243.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.241$, Rules used = {3042, 4089, 3042, 4107, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx \\ & \quad \downarrow 3042 \\ & \int \frac{(a + b \tan(c + dx))^2(A + B \tan(c + dx))}{\tan(c + dx)} dx \\ & \quad \downarrow 4089 \\ & \int \cot(c + dx) (Aa^2 + b(Ab + 2aB) \tan^2(c + dx) + (Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx + \\ & \quad \quad \quad \frac{b^2B \tan(c + dx)}{d} \\ & \quad \downarrow 3042 \\ & \int \frac{Aa^2 + b(Ab + 2aB) \tan(c + dx)^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx)}{\tan(c + dx)} dx + \frac{b^2B \tan(c + dx)}{d} \\ & \quad \downarrow 4107 \\ & a^2A \int \cot(c + dx) dx + b(2aB + Ab) \int \tan(c + dx) dx + x(a^2B + 2aAb - b^2B) + \frac{b^2B \tan(c + dx)}{d} \\ & \quad \downarrow 3042 \\ & a^2A \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + b(2aB + Ab) \int \tan(c + dx) dx + x(a^2B + 2aAb - b^2B) + \\ & \quad \quad \quad \frac{b^2B \tan(c + dx)}{d} \\ & \quad \downarrow 25 \end{aligned}$$

3.243. $\int \cot(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$a^2(-A) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + b(2aB + Ab) \int \tan(c + dx) dx + x(a^2B + 2aAb - b^2B) + \frac{b^2B \tan(c + dx)}{d}$$

↓ 3956

$$x(a^2B + 2aAb - b^2B) + \frac{a^2A \log(-\sin(c + dx))}{d} - \frac{b(2aB + Ab) \log(\cos(c + dx))}{d} + \frac{b^2B \tan(c + dx)}{d}$$

input `Int[Cot[c + d*x]*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(2*a*A*b + a^2*B - b^2*B)*x - (b*(A*b + 2*a*B)*Log[Cos[c + d*x]])/d + (a^2*A*Log[-Sin[c + d*x]])/d + (b^2*B*Tan[c + d*x])/d`

3.243.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4089 `Int[(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b^2*B*(Tan[e + f*x]/(d*f)), x] + Simp[1/d Int[(a^2*A*d - b^2*B*c + (2*a*A*b + B*(a^2 - b^2))*d*Tan[e + f*x] + (A*b^2*d - b*B*(b*c - 2*a*d))*Tan[e + f*x]^2)/(c + d*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4107 `Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2/tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[B*x, x] + (Simp[A Int[1/Tan[e + f*x], x], x] + Simp[C Int[Tan[e + f*x], x], x]) /; FreeQ[{e, f, A, B, C}, x] && NeQ[A, C]`

3.243.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.14

method	result
parallelrisc	$\frac{(-Aa^2 + Ab^2 + 2Bab) \ln(\sec^2(dx+c)) + 2Aa^2 \ln(\tan(dx+c)) + 2Bb^2 \tan(dx+c) + 4d(Aab + \frac{1}{2}Ba^2 - \frac{1}{2}Bb^2)x}{2d}$
norman	$(2Aab + Ba^2 - Bb^2)x + \frac{b^2B \tan(dx+c)}{d} + \frac{Aa^2 \ln(\tan(dx+c))}{d} - \frac{(Aa^2 - Ab^2 - 2Bab) \ln(1 + \tan^2(dx+c))}{2d}$
derivativedivides	$- \frac{(Aa^2 - Ab^2 - 2Bab) \ln(\cot^2(dx+c)+1)}{2} + \frac{(2Aab + Ba^2 - Bb^2)(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c))) - \frac{Bb^2}{\cot(dx+c)} + b(Ab + 2Ba) \ln(\cot(dx+c))}{d}$
default	$- \frac{(Aa^2 - Ab^2 - 2Bab) \ln(\cot^2(dx+c)+1)}{2} + \frac{(2Aab + Ba^2 - Bb^2)(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c))) - \frac{Bb^2}{\cot(dx+c)} + b(Ab + 2Ba) \ln(\cot(dx+c))}{d}$
risc	$2Aabx + Ba^2x - Bb^2x + iAb^2x - iAa^2x + \frac{4iBabc}{d} + 2iBabx - \frac{2ia^2Ac}{d} + \frac{2iAb^2c}{d} + \frac{2i}{d(e^{2i(dx+c)} + 1)}$

input `int(cot(d*x+c)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output $1/2*((-Aa^2 + Ab^2 + 2Bab) \ln(\sec^2(dx+c)^2) + 2Aa^2 \ln(\tan(dx+c)) + 2Bb^2 \tan(dx+c) + 4d(Aab + 1/2Ba^2 - 1/2Bb^2)x)/d$

3.243.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.31

$$\int \cot(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{Aa^2 \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) + 2Bb^2 \tan(dx+c) + 2(Ba^2 + 2Aab - Bb^2)dx - (2Bab + Ab^2) \log\left(\frac{1}{\tan(dx+c)^2+1}\right)}{2d}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output $1/2*(Aa^2 \log(\tan(dx+c)^2/(\tan(dx+c)^2+1)) + 2Bb^2 \tan(dx+c) + 2*(Ba^2 + 2Aab - Bb^2)*dx - (2Bab + Ab^2) \log(1/(\tan(dx+c)^2+1)))/d$

3.243. $\int \cot(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.243.6 Sympy [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.84

$$\int \cot(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \begin{cases} -\frac{Aa^2 \log(\tan^2(c+dx)+1)}{2d} + \frac{Aa^2 \log(\tan(c+dx))}{d} + 2Aabx + \frac{Ab^2 \log(\tan^2(c+dx)+1)}{2d} + Ba^2x + \frac{Bab \log(\tan^2(c+dx)+1)}{d} - \\ x(A + B \tan(c))(a + b \tan(c))^2 \cot(c) \end{cases}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`output `Piecewise((-A*a**2*log(tan(c + d*x)**2 + 1)/(2*d) + A*a**2*log(tan(c + d*x)))/d + 2*A*a*b*x + A*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + B*a**2*x + B*a*b*log(tan(c + d*x)**2 + 1)/d - B*b**2*x + B*b**2*tan(c + d*x)/d, Ne(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**2*cot(c), True))`**3.243.7 Maxima [A] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.21

$$\int \cot(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{2Aa^2 \log(\tan(dx + c)) + 2Bb^2 \tan(dx + c) + 2(Ba^2 + 2Aab - Bb^2)(dx + c) - (Aa^2 - 2Bab - Ab^2) \log(\tan(dx + c)^2 + 1)}{2d}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`output `1/2*(2*A*a^2*log(tan(d*x + c)) + 2*B*b^2*tan(d*x + c) + 2*(B*a^2 + 2*A*a*b - B*b^2)*(d*x + c) - (A*a^2 - 2*B*a*b - A*b^2)*log(tan(d*x + c)^2 + 1))/d`

3.243.8 Giac [A] (verification not implemented)

Time = 0.70 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.23

$$\int \cot(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{2 A a^2 \log(|\tan(dx + c)|) + 2 B b^2 \tan(dx + c) + 2 (B a^2 + 2 A a b - B b^2)(dx + c) - (A a^2 - 2 B a b - A b^2) \log(\tan(dx + c)^2 + 1)}{2 d}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `1/2*(2*A*a^2*log(abs(tan(d*x + c))) + 2*B*b^2*tan(d*x + c) + 2*(B*a^2 + 2*A*a*b - B*b^2)*(d*x + c) - (A*a^2 - 2*B*a*b - A*b^2)*log(tan(d*x + c)^2 + 1))/d`

3.243.9 Mupad [B] (verification not implemented)

Time = 7.51 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.29

$$\int \cot(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{A a^2 \ln(\tan(c + dx))}{d} + \frac{\ln(\tan(c + dx) + 1i) (A - B 1i) (b + a 1i)^2}{2 d}$$

$$+ \frac{B b^2 \tan(c + dx)}{d} + \frac{\ln(\tan(c + dx) - i) (A + B 1i) (-b + a 1i)^2}{2 d}$$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)`

output `(A*a^2*log(tan(c + d*x)))/d + (log(tan(c + d*x) + 1i)*(A - B*1i)*(a*1i + b)^2)/(2*d) + (B*b^2*tan(c + d*x))/d + (log(tan(c + d*x) - 1i)*(A + B*1i)*(a*1i - b)^2)/(2*d)`

3.244 $\int \cot^2(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.244.1 Optimal result	2377
3.244.2 Mathematica [C] (verified)	2377
3.244.3 Rubi [A] (verified)	2378
3.244.4 Maple [A] (verified)	2380
3.244.5 Fricas [A] (verification not implemented)	2380
3.244.6 Sympy [B] (verification not implemented)	2381
3.244.7 Maxima [A] (verification not implemented)	2381
3.244.8 Giac [A] (verification not implemented)	2382
3.244.9 Mupad [B] (verification not implemented)	2382

3.244.1 Optimal result

Integrand size = 31, antiderivative size = 72

$$\begin{aligned} & \int \cot^2(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx \\ &= -((a^2A - Ab^2 - 2abB) x) - \frac{a^2A \cot(c+dx)}{d} \\ & \quad - \frac{b^2B \log(\cos(c+dx))}{d} + \frac{a(2Ab + aB) \log(\sin(c+dx))}{d} \end{aligned}$$

```
output -(A*a^2-A*b^2-2*B*a*b)*x-a^2*A*cot(d*x+c)/d-b^2*B*ln(cos(d*x+c))/d+a*(2*A*
b+B*a)*ln(sin(d*x+c))/d
```

3.244.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.39

$$\begin{aligned} & \int \cot^2(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx \\ &= \frac{-2a^2A \cot(c+dx) + i(a+ib)^2(A+iB) \log(i - \tan(c+dx)) + 2a(2Ab + aB) \log(\tan(c+dx)) - (a - ib)}{2d} \end{aligned}$$

input `Integrate[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output $(-2*a^2*A*Cot[c + d*x] + I*(a + I*b)^2*(A + I*B)*Log[I - Tan[c + d*x]] + 2*a*(2*A*b + a*B)*Log[Tan[c + d*x]] - (a - I*b)^2*(I*A + B)*Log[I + Tan[c + d*x]])/(2*d)$

3.244.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.226$, Rules used = {3042, 4087, 3042, 4107, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^2(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{(a + b \tan(c + dx))^2(A + B \tan(c + dx))}{\tan(c + dx)^2} dx \\
 & \quad \downarrow 4087 \\
 & \int \cot(c + dx) (b^2 B \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)) dx - \\
 & \quad \quad \quad \frac{a^2 A \cot(c + dx)}{d} \\
 & \quad \downarrow 3042 \\
 & \int \frac{b^2 B \tan(c + dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)}{\tan(c + dx)} dx - \frac{a^2 A \cot(c + dx)}{d} \\
 & \quad \downarrow 4107 \\
 & a(aB + 2Ab) \int \cot(c + dx) dx + b^2 B \int \tan(c + dx) dx - x(a^2 A - 2abB - Ab^2) - \frac{a^2 A \cot(c + dx)}{d} \\
 & \quad \downarrow 3042 \\
 & a(aB + 2Ab) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + b^2 B \int \tan(c + dx) dx - x(a^2 A - 2abB - Ab^2) - \\
 & \quad \quad \quad \frac{a^2 A \cot(c + dx)}{d} \\
 & \quad \downarrow 25
 \end{aligned}$$

3.244. $\int \cot^2(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\begin{aligned}
 & -a(aB + 2Ab) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + b^2B \int \tan(c + dx) dx - x(a^2A - 2abB - Ab^2) - \\
 & \quad \frac{a^2A \cot(c + dx)}{d} \\
 & \quad \downarrow \text{3956} \\
 & -x(a^2A - 2abB - Ab^2) - \frac{a^2A \cot(c + dx)}{d} + \frac{a(aB + 2Ab) \log(-\sin(c + dx))}{d} - \\
 & \quad \frac{b^2B \log(\cos(c + dx))}{d}
 \end{aligned}$$

input `Int[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `-((a^2*A - A*b^2 - 2*a*b*B)*x) - (a^2*A*Cot[c + d*x])/d - (b^2*B*Log[Cos[c + d*x]])/d + (a*(2*A*b + a*B)*Log[-Sin[c + d*x]])/d`

3.244.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4087 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(B*c - A*d)*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`

```
rule 4107 Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2
)/tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[B*x, x] + (Simp[A Int[1/Tan[
e + f*x], x], x] + Simp[C Int[Tan[e + f*x], x], x]) /; FreeQ[{e, f, A, B,
C}, x] && NeQ[A, C]
```

3.244.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.17

method	result
derivativedivides	$\frac{A^2(-\cot(dx+c)-dx-c)+B^2 a^2 \ln(\sin(dx+c))+2Aab \ln(\sin(dx+c))+2Bab(dx+c)+A b^2(dx+c)-B b^2 \ln(\cos(dx+c))}{d}$
default	$\frac{A^2(-\cot(dx+c)-dx-c)+B^2 a^2 \ln(\sin(dx+c))+2Aab \ln(\sin(dx+c))+2Bab(dx+c)+A b^2(dx+c)-B b^2 \ln(\cos(dx+c))}{d}$
parallelrisc	$\frac{(-2Aab-B a^2+B b^2) \ln(\sec^2(dx+c))+(4Aab+2B a^2) \ln(\tan(dx+c))-2A a^2 \cot(dx+c)-2dx(A a^2-A b^2-2Bab)}{2d}$
norman	$\frac{(-A a^2+A b^2+2Bab)x \tan(dx+c)-\frac{A a^2}{d} + \frac{a(2Ab+Ba) \ln(\tan(dx+c))}{d} - \frac{(2Aab+B a^2-B b^2) \ln(1+\tan^2(dx+c))}{2d}}{\tan(dx+c)}$
risc	$-\frac{2ia^2Bc}{d} + iB b^2x + \frac{2iB b^2c}{d} - A a^2x + A b^2x + 2Babx - \frac{4iAabc}{d} - \frac{2iA a^2}{d(e^{2i(dx+c)}-1)} - 2iAabx$

```
input int(cot(d*x+c)^2*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

```
output 1/d*(A*a^2*(-cot(d*x+c)-d*x-c)+B*a^2*ln(sin(d*x+c))+2*A*a*b*ln(sin(d*x+c))
+2*B*a*b*(d*x+c)+A*b^2*(d*x+c)-B*b^2*ln(cos(d*x+c)))
```

3.244.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.56

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$\frac{Bb^2 \log\left(\frac{1}{\tan(dx+c)^2+1}\right) \tan(dx+c) + 2(Aa^2 - 2Bab - Ab^2)dx \tan(dx+c) + 2Aa^2 - (Ba^2 + 2Aab) \log(\tan(dx+c))}{2d \tan(dx+c)}$$

```
input integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="f
ricas")
```

output
$$\frac{-1/2*(B*b^2*\log(1/(\tan(d*x + c)^2 + 1))*\tan(d*x + c) + 2*(A*a^2 - 2*B*a*b - A*b^2)*d*x*\tan(d*x + c) + 2*A*a^2 - (B*a^2 + 2*A*a*b)*\log(\tan(d*x + c)^2 / (\tan(d*x + c)^2 + 1))*\tan(d*x + c))/(d*\tan(d*x + c))$$

3.244.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 167 vs. $2(66) = 132$.

Time = 0.46 (sec) , antiderivative size = 167, normalized size of antiderivative = 2.32

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \tilde{\infty} A a^2 x \\ x(A + B \tan(c))(a + b \tan(c))^2 \cot^2(c) \\ \tilde{\infty} A a^2 x \\ -A a^2 x - \frac{A a^2}{d \tan(c+dx)} - \frac{A a b \log(\tan^2(c+dx)+1)}{d} + \frac{2 A a b \log(\tan(c+dx))}{d} + A b^2 x - \frac{B a^2 \log(\tan^2(c+dx)+1)}{2d} + \frac{B a^2 \log(\tan(c+dx))}{d} \end{cases}$$

input `integrate(cot(d*x+c)**2*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `Piecewise((zoo*A*a**2*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**2*cot(c)**2, Eq(d, 0)), (zoo*A*a**2*x, Eq(c, -d*x)), (-A*a**2*x - A*a**2/(d*tan(c + d*x)) - A*a*b*log(tan(c + d*x)**2 + 1)/d + 2*A*a*b*log(tan(c + d*x))/d + A*b**2*x - B*a**2*log(tan(c + d*x)**2 + 1)/(2*d) + B*a**2*log(tan(c + d*x))/d + 2*B*a*b*x + B*b**2*log(tan(c + d*x)**2 + 1)/(2*d), True))`

3.244.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.29

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$\frac{2(Aa^2 - 2Bab - Ab^2)(dx + c) + (Ba^2 + 2Aab - Bb^2) \log(\tan(dx + c)^2 + 1) - 2(Ba^2 + 2Aab) \log(\tan(c + dx))}{2d}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

3.244. $\int \cot^2(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

output
$$\frac{-1/2*(2*(A*a^2 - 2*B*a*b - A*b^2)*(d*x + c) + (B*a^2 + 2*A*a*b - B*b^2)*\log(\tan(d*x + c)^2 + 1) - 2*(B*a^2 + 2*A*a*b)*\log(\tan(d*x + c)) + 2*A*a^2/\tan(d*x + c))/d}$$

3.244.8 Giac [A] (verification not implemented)

Time = 0.87 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.64

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{2(Aa^2 - 2Bab - Ab^2)(dx + c) + (Ba^2 + 2Aab - Bb^2) \log(\tan(dx + c)^2 + 1) - 2(Ba^2 + 2Aab) \log(\tan(dx + c))}{2d}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output
$$\frac{-1/2*(2*(A*a^2 - 2*B*a*b - A*b^2)*(d*x + c) + (B*a^2 + 2*A*a*b - B*b^2)*\log(\tan(d*x + c)^2 + 1) - 2*(B*a^2 + 2*A*a*b)*\log(\tan(d*x + c)) + 2*(B*a^2 + 2*A*a*b)*\tan(d*x + c) + 2*A*a*b*\tan(d*x + c) + A*a^2)/\tan(d*x + c))/d}$$

3.244.9 Mupad [B] (verification not implemented)

Time = 7.53 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.39

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{\ln(\tan(c + dx)) (Ba^2 + 2Aba) - \ln(\tan(c + dx) - i) (-B + A i) (-b + a i)^2}{d} - \frac{\ln(\tan(c + dx) + i) (B + A i) (b + a i)^2}{2d} - \frac{Aa^2 \cot(c + dx)}{d}$$

input `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)`

output
$$\frac{(\log(\tan(c + d*x))*(B*a^2 + 2*A*a*b))/d - (\log(\tan(c + d*x) - 1i)*(A*1i - B)*(a*1i - b)^2)/(2*d) + (\log(\tan(c + d*x) + 1i)*(A*1i + B)*(a*1i + b)^2)/(2*d) - (A*a^2*\cot(c + d*x))/d}$$

3.244. $\int \cot^2(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.245 $\int \cot^3(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.245.1 Optimal result	2383
3.245.2 Mathematica [C] (verified)	2383
3.245.3 Rubi [A] (verified)	2384
3.245.4 Maple [A] (verified)	2387
3.245.5 Fricas [A] (verification not implemented)	2387
3.245.6 Sympy [B] (verification not implemented)	2388
3.245.7 Maxima [A] (verification not implemented)	2388
3.245.8 Giac [B] (verification not implemented)	2389
3.245.9 Mupad [B] (verification not implemented)	2389

3.245.1 Optimal result

Integrand size = 31, antiderivative size = 88

$$\int \cot^3(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= (b^2B - a(2Ab + aB)) x - \frac{a(2Ab + aB) \cot(c+dx)}{d}$$

$$- \frac{a^2A \cot^2(c+dx)}{2d} - \frac{(a^2A - Ab^2 - 2abB) \log(\sin(c+dx))}{d}$$

```
output (B*b^2-a*(2*A*b+B*a))*x-a*(2*A*b+B*a)*cot(d*x+c)/d-1/2*a^2*A*cot(d*x+c)^2/d-(A*a^2-A*b^2-2*B*a*b)*ln(sin(d*x+c))/d
```

3.245.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.40

$$\int \cot^3(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{-2a(2Ab + aB) \cot(c+dx) - a^2A \cot^2(c+dx) + (a+ib)^2(A+iB) \log(i - \tan(c+dx)) - 2(a^2A - Ab^2)}{2d}$$

input `Integrate[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output $(-2*a*(2*A*b + a*B)*\text{Cot}[c + d*x] - a^2*A*\text{Cot}[c + d*x]^2 + (a + I*b)^2*(A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]] - 2*(a^2*A - A*b^2 - 2*a*b*B)*\text{Log}[\text{Tan}[c + d*x]] + (a - I*b)^2*(A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]])/(2*d)$

3.245.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.03, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 4087, 3042, 4111, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^3(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))^2(A + B \tan(c + dx))}{\tan(c + dx)^3} dx \\
 & \quad \downarrow \text{4087} \\
 & \int \cot^2(c + dx) (b^2 B \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)) dx - \\
 & \quad \quad \quad \frac{a^2 A \cot^2(c + dx)}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{b^2 B \tan(c + dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)}{\tan(c + dx)^2} dx - \frac{a^2 A \cot^2(c + dx)}{2d} \\
 & \quad \downarrow \text{4111} \\
 & \int -\cot(c + dx) (Aa^2 - 2bBa - Ab^2 - (b^2 B - a(2Ab + aB)) \tan(c + dx)) dx - \\
 & \quad \quad \quad \frac{a^2 A \cot^2(c + dx)}{2d} - \frac{a(aB + 2Ab) \cot(c + dx)}{d} \\
 & \quad \downarrow \text{25} \\
 & - \int \cot(c + dx) (Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2 B) \tan(c + dx)) dx - \frac{a^2 A \cot^2(c + dx)}{2d} - \\
 & \quad \quad \quad \frac{a(aB + 2Ab) \cot(c + dx)}{d}
 \end{aligned}$$

3.245. $\int \cot^3(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \downarrow 3042 \\
& - \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx)}{\tan(c + dx)} dx - \frac{a^2 A \cot^2(c + dx)}{2d} - \\
& \quad \frac{a(aB + 2Ab) \cot(c + dx)}{d} \\
& \downarrow 4014 \\
& -(a^2 A - 2abB - Ab^2) \int \cot(c + dx) dx - x(a^2 B + 2aAb - b^2 B) - \frac{a^2 A \cot^2(c + dx)}{2d} - \\
& \quad \frac{a(aB + 2Ab) \cot(c + dx)}{d} \\
& \downarrow 3042 \\
& -(a^2 A - 2abB - Ab^2) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - x(a^2 B + 2aAb - b^2 B) - \frac{a^2 A \cot^2(c + dx)}{2d} - \\
& \quad \frac{a(aB + 2Ab) \cot(c + dx)}{d} \\
& \downarrow 25 \\
& (a^2 A - 2abB - Ab^2) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - x(a^2 B + 2aAb - b^2 B) - \frac{a^2 A \cot^2(c + dx)}{2d} - \\
& \quad \frac{a(aB + 2Ab) \cot(c + dx)}{d} \\
& \downarrow 3956 \\
& - \frac{(a^2 A - 2abB - Ab^2) \log(-\sin(c + dx))}{d} - x(a^2 B + 2aAb - b^2 B) - \frac{a^2 A \cot^2(c + dx)}{2d} - \\
& \quad \frac{a(aB + 2Ab) \cot(c + dx)}{d}
\end{aligned}$$

input `Int[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `-((2*a*A*b + a^2*B - b^2*B)*x) - (a*(2*A*b + a*B)*Cot[c + d*x])/d - (a^2*A *Cot[c + d*x]^2)/(2*d) - ((a^2*A - A*b^2 - 2*a*b*B)*Log[-Sin[c + d*x]])/d`

3.245.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`
- rule 4087 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(B*c - A*d)*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`
- rule 4111 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.245.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.22

method	result
derivativedivides	$\frac{A a^2 \left(-\frac{\cot^2(dx+c)}{2} - \ln(\sin(dx+c)) \right) + B a^2 (-\cot(dx+c) - dx - c) + 2Aab(-\cot(dx+c) - dx - c) + 2Bab \ln(\sin(dx+c))}{d}$
default	$\frac{A a^2 \left(-\frac{\cot^2(dx+c)}{2} - \ln(\sin(dx+c)) \right) + B a^2 (-\cot(dx+c) - dx - c) + 2Aab(-\cot(dx+c) - dx - c) + 2Bab \ln(\sin(dx+c))}{d}$
parallelrisc	$\frac{(A a^2 - A b^2 - 2Bab) \ln(\sec^2(dx+c)) + (-2A a^2 + 2A b^2 + 4Bab) \ln(\tan(dx+c)) - A(\cot^2(dx+c)) a^2 + (-4Aab - 2B a^2) \cot(dx+c)}{2d}$
norman	$\frac{(-2Aab - B a^2 + B b^2)x(\tan^2(dx+c)) - \frac{A a^2}{2d} - \frac{a(2Ab+Ba)\tan(dx+c)}{d}}{\tan(dx+c)^2} - \frac{(A a^2 - A b^2 - 2Bab) \ln(\tan(dx+c))}{d} + \frac{(A a^2 - B b^2) \ln(\sin(dx+c))}{d}$
risc	$-2Aabx - B a^2x + B b^2x - \frac{2iA b^2c}{d} - iA b^2x + iA a^2x - \frac{4iBabc}{d} - \frac{2ia(iAa e^{2i(dx+c)} + 2Ab e^{2i(dx+c)})}{d(e^{2i(dx+c)} + 1)}$

input `int(cot(d*x+c)^3*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*(A*a^2*(-1/2*cot(d*x+c)^2-ln(sin(d*x+c)))+B*a^2*(-cot(d*x+c)-d*x-c)+2*A*a*b*(-cot(d*x+c)-d*x-c)+2*B*a*b*ln(sin(d*x+c))+A*b^2*ln(sin(d*x+c))+B*b^2*(d*x+c))`

3.245.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.39

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{(Aa^2 - 2Bab - Ab^2) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2 + 1}\right) \tan(dx+c)^2 + Aa^2 + (Aa^2 + 2(Ba^2 + 2Aab - Bb^2)dx) \tan(dx+c)}{2d \tan(dx+c)^2}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-1/2*((A*a^2 - 2*B*a*b - A*b^2)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^2 + A*a^2 + (A*a^2 + 2*(B*a^2 + 2*A*a*b - B*b^2)*d*x)*tan(d*x + c)^2 + 2*(B*a^2 + 2*A*a*b)*tan(d*x + c))/(d*tan(d*x + c)^2)`

3.245. $\int \cot^3(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.245.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 214 vs. 2(78) = 156.

Time = 0.82 (sec) , antiderivative size = 214, normalized size of antiderivative = 2.43

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \tilde{\infty} A a^2 x \\ x(A + B \tan(c))(a + b \tan(c))^2 \cot^3(c) \\ \tilde{\infty} A a^2 x \\ \frac{A a^2 \log(\tan^2(c + dx) + 1)}{2d} - \frac{A a^2 \log(\tan(c + dx))}{d} - \frac{A a^2}{2d \tan^2(c + dx)} - 2 A a b x - \frac{2 A a b}{d \tan(c + dx)} - \frac{A b^2 \log(\tan^2(c + dx) + 1)}{2d} + \frac{A b^2 \log(\tan(c + dx))}{d} \end{cases}$$

input `integrate(cot(d*x+c)**3*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `Piecewise((zoo*A*a**2*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**2*cot(c)**3, Eq(d, 0)), (zoo*A*a**2*x, Eq(c, -d*x)), (A*a**2*log(tan(c + d*x)**2 + 1)/(2*d) - A*a**2*log(tan(c + d*x))/d - A*a**2/(2*d*tan(c + d*x)**2) - 2*A*a*b*x - 2*A*a*b/(d*tan(c + d*x)) - A*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + A*b**2*log(tan(c + d*x))/d - B*a**2*x - B*a**2/(d*tan(c + d*x)) - B*a*b*log(tan(c + d*x)**2 + 1)/d + 2*B*a*b*log(tan(c + d*x))/d + B*b**2*x, True))`

3.245.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.36

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$\frac{2(Ba^2 + 2Aab - Bb^2)(dx + c) - (Aa^2 - 2Bab - Ab^2) \log(\tan(dx + c)^2 + 1) + 2(Aa^2 - 2Bab - Ab^2) \log(\tan(dx + c))}{2d}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(2*(B*a^2 + 2*A*a*b - B*b^2)*(d*x + c) - (A*a^2 - 2*B*a*b - A*b^2)*log(tan(d*x + c)^2 + 1) + 2*(A*a^2 - 2*B*a*b - A*b^2)*log(tan(d*x + c)) + (A*a^2 + 2*(B*a^2 + 2*A*a*b)*tan(d*x + c))/tan(d*x + c)^2)/d`

3.245. $\int \cot^3(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.245.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 237 vs. 2(86) = 172.

Time = 1.10 (sec) , antiderivative size = 237, normalized size of antiderivative = 2.69

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$Aa^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 4Ba^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 8Aab \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 8(Ba^2 + 2Aab - Bb^2)(dx +$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `-1/8*(A*a^2*tan(1/2*d*x + 1/2*c)^2 - 4*B*a^2*tan(1/2*d*x + 1/2*c) - 8*A*a*b*tan(1/2*d*x + 1/2*c) + 8*(B*a^2 + 2*A*a*b - B*b^2)*(d*x + c) - 8*(A*a^2 - 2*B*a*b - A*b^2)*log(tan(1/2*d*x + 1/2*c)^2 + 1) + 8*(A*a^2 - 2*B*a*b - A*b^2)*log(abs(tan(1/2*d*x + 1/2*c))) - (12*A*a^2*tan(1/2*d*x + 1/2*c)^2 - 24*B*a*b*tan(1/2*d*x + 1/2*c)^2 - 12*A*b^2*tan(1/2*d*x + 1/2*c)^2 - 4*B*a^2*tan(1/2*d*x + 1/2*c) - 8*A*a*b*tan(1/2*d*x + 1/2*c) - A*a^2)/tan(1/2*d*x + 1/2*c)^2)/d`

3.245.9 Mupad [B] (verification not implemented)

Time = 8.04 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.44

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx))(-Aa^2 + 2Bab + Ab^2)}{d}$$

$$- \frac{\cot(c + dx)^2 \left(\frac{Aa^2}{2} + \tan(c + dx)(Ba^2 + 2Aba) \right)}{d}$$

$$- \frac{\ln(\tan(c + dx) + i)(A - B i)(b + a i)^2}{2d}$$

$$- \frac{\ln(\tan(c + dx) - i)(A + B i)(-b + a i)^2}{2d}$$

input `int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)`

output $(\log(\tan(c + dx))(A^2b - A^2a + 2Bab))/d - (\cot(c + dx)^2((A^2a^2)/2 + \tan(c + dx)(B^2a^2 + 2Aab)))/d - (\log(\tan(c + dx) + i)(A - Bi)(a + bi)^2)/(2d) - (\log(\tan(c + dx) - i)(A + Bi)(a - bi)^2)/(2d)$

3.246 $\int \cot^4(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.246.1 Optimal result	2391
3.246.2 Mathematica [C] (verified)	2391
3.246.3 Rubi [A] (verified)	2392
3.246.4 Maple [A] (verified)	2395
3.246.5 Fricas [A] (verification not implemented)	2396
3.246.6 Sympy [B] (verification not implemented)	2396
3.246.7 Maxima [A] (verification not implemented)	2397
3.246.8 Giac [B] (verification not implemented)	2397
3.246.9 Mupad [B] (verification not implemented)	2398

3.246.1 Optimal result

Integrand size = 31, antiderivative size = 118

$$\int \cot^4(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= (a^2A - Ab^2 - 2abB)x + \frac{(a^2A - Ab^2 - 2abB) \cot(c+dx)}{d} - \frac{a(2Ab + aB) \cot^2(c+dx)}{2d}$$

$$- \frac{a^2A \cot^3(c+dx)}{3d} + \frac{(b^2B - a(2Ab + aB)) \log(\sin(c+dx))}{d}$$

```
output (A*a^2-A*b^2-2*B*a*b)*x+(A*a^2-A*b^2-2*B*a*b)*cot(d*x+c)/d-1/2*a*(2*A*b+B*a)*cot(d*x+c)^2/d-1/3*a^2*A*cot(d*x+c)^3/d+(B*b^2-a*(2*A*b+B*a))*ln(sin(d*x+c))/d
```

3.246.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.25 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.29

$$\int \cot^4(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{6(a^2A - Ab^2 - 2abB) \cot(c+dx) - 3a(2Ab + aB) \cot^2(c+dx) - 2a^2A \cot^3(c+dx) + 3(a+ib)^2(-iA +$$

input `Integrate[Cot[c + d*x]^4*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output $(6*(a^2*A - A*b^2 - 2*a*b*B)*\text{Cot}[c + d*x] - 3*a*(2*A*b + a*B)*\text{Cot}[c + d*x]^2 - 2*a^2*A*\text{Cot}[c + d*x]^3 + 3*(a + I*b)^2*((-I)*A + B)*\text{Log}[I - \text{Tan}[c + d*x]] - 6*(2*a*A*b + a^2*B - b^2*B)*\text{Log}[\text{Tan}[c + d*x]] + 3*(a - I*b)^2*(I*A + B)*\text{Log}[I + \text{Tan}[c + d*x]])/(6*d)$

3.246.3 Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 4087, 3042, 4111, 25, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \tan(c + dx))^2(A + B \tan(c + dx))}{\tan(c + dx)^4} dx$$

$$\downarrow 4087$$

$$\int \cot^3(c + dx) (b^2 B \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)) dx - \frac{a^2 A \cot^3(c + dx)}{3d}$$

$$\downarrow 3042$$

$$\int \frac{b^2 B \tan(c + dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)}{\tan(c + dx)^3} dx - \frac{a^2 A \cot^3(c + dx)}{3d}$$

$$\downarrow 4111$$

$$\int -\cot^2(c + dx) (Aa^2 - 2bBa - Ab^2 - (b^2 B - a(2Ab + aB)) \tan(c + dx)) dx - \frac{a^2 A \cot^3(c + dx)}{3d} - \frac{a(aB + 2Ab) \cot^2(c + dx)}{2d}$$

$$\downarrow 25$$

$$\begin{aligned}
& - \int \cot^2(c+dx) (Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c+dx)) dx - \\
& \quad \frac{a^2 A \cot^3(c+dx)}{3d} - \frac{a(aB + 2Ab) \cot^2(c+dx)}{2d} \\
& \quad \downarrow \text{3042} \\
& - \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c+dx)}{\tan(c+dx)^2} dx - \frac{a^2 A \cot^3(c+dx)}{3d} - \\
& \quad \frac{a(aB + 2Ab) \cot^2(c+dx)}{2d} \\
& \quad \downarrow \text{4012} \\
& - \int \cot(c+dx) (Ba^2 + 2Aba - b^2B - (Aa^2 - 2bBa - Ab^2) \tan(c+dx)) dx + \\
& \frac{(a^2 A - 2abB - Ab^2) \cot(c+dx)}{d} - \frac{a^2 A \cot^3(c+dx)}{3d} - \frac{a(aB + 2Ab) \cot^2(c+dx)}{2d} \\
& \quad \downarrow \text{3042} \\
& - \int \frac{Ba^2 + 2Aba - b^2B - (Aa^2 - 2bBa - Ab^2) \tan(c+dx)}{\tan(c+dx)} dx + \\
& \frac{(a^2 A - 2abB - Ab^2) \cot(c+dx)}{d} - \frac{a^2 A \cot^3(c+dx)}{3d} - \frac{a(aB + 2Ab) \cot^2(c+dx)}{2d} \\
& \quad \downarrow \text{4014} \\
& -(a^2 B + 2aAb - b^2 B) \int \cot(c+dx) dx + \frac{(a^2 A - 2abB - Ab^2) \cot(c+dx)}{d} + \\
& x(a^2 A - 2abB - Ab^2) - \frac{a^2 A \cot^3(c+dx)}{3d} - \frac{a(aB + 2Ab) \cot^2(c+dx)}{2d} \\
& \quad \downarrow \text{3042} \\
& -(a^2 B + 2aAb - b^2 B) \int -\tan\left(c+dx + \frac{\pi}{2}\right) dx + \frac{(a^2 A - 2abB - Ab^2) \cot(c+dx)}{d} + \\
& x(a^2 A - 2abB - Ab^2) - \frac{a^2 A \cot^3(c+dx)}{3d} - \frac{a(aB + 2Ab) \cot^2(c+dx)}{2d} \\
& \quad \downarrow \text{25} \\
& (a^2 B + 2aAb - b^2 B) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + \frac{(a^2 A - 2abB - Ab^2) \cot(c+dx)}{d} + \\
& x(a^2 A - 2abB - Ab^2) - \frac{a^2 A \cot^3(c+dx)}{3d} - \frac{a(aB + 2Ab) \cot^2(c+dx)}{2d} \\
& \quad \downarrow \text{3956} \\
& \frac{(a^2 A - 2abB - Ab^2) \cot(c+dx)}{d} - \frac{(a^2 B + 2aAb - b^2 B) \log(-\sin(c+dx))}{d} + \\
& x(a^2 A - 2abB - Ab^2) - \frac{a^2 A \cot^3(c+dx)}{3d} - \frac{a(aB + 2Ab) \cot^2(c+dx)}{2d}
\end{aligned}$$

3.246. $\int \cot^4(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

input `Int[Cot[c + d*x]^4*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(a^2*A - A*b^2 - 2*a*b*B)*x + ((a^2*A - A*b^2 - 2*a*b*B)*Cot[c + d*x])/d - (a*(2*A*b + a*B)*Cot[c + d*x]^2)/(2*d) - (a^2*A*Cot[c + d*x]^3)/(3*d) - (2*a*A*b + a^2*B - b^2*B)*Log[-Sin[c + d*x]]/d`

3.246.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

```
rule 4087 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(- (B*c - A*d))*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

```
rule 4111 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]
```

3.246.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.15

method	result
derivativedivides	$A a^2 \left(-\frac{\cot^3(dx+c)}{3} + \cot(dx+c) + dx+c \right) + B a^2 \left(-\frac{\cot^2(dx+c)}{2} - \ln(\sin(dx+c)) \right) + 2Aab \left(-\frac{\cot^2(dx+c)}{2} - \ln(\sin(dx+c)) \right) + \frac{d}{d}$
default	$A a^2 \left(-\frac{\cot^3(dx+c)}{3} + \cot(dx+c) + dx+c \right) + B a^2 \left(-\frac{\cot^2(dx+c)}{2} - \ln(\sin(dx+c)) \right) + 2Aab \left(-\frac{\cot^2(dx+c)}{2} - \ln(\sin(dx+c)) \right) + \frac{d}{d}$
parallelrisch	$\frac{3(2Aab + B a^2 - B b^2) \ln(\sec^2(dx+c)) + 6(-2Aab - B a^2 + B b^2) \ln(\tan(dx+c)) - 2A(\cot^3(dx+c))a^2 + 3(-2Aab - B a^2)(\cot^2(dx+c))}{6d}$
norman	$\frac{(A a^2 - A b^2 - 2Bab) (\tan^2(dx+c))}{d} + \frac{(A a^2 - A b^2 - 2Bab)x (\tan^3(dx+c)) - \frac{A a^2}{3d} - \frac{a(2Ab+Ba) \tan(dx+c)}{2d}}{\tan(dx+c)^3} - \frac{(2Aab + B a^2 - B b^2)}{d}$
risch	$\frac{4iAabc}{d} + \frac{2ia^2Bc}{d} - iB b^2 x + A a^2 x - A b^2 x - 2Babx - \frac{2iB b^2 c}{d} - \frac{2i(6iAab e^{4i(dx+c)} + 3iB a^2 e^{4i(dx+c)})}{d}$

```
input int(cot(d*x+c)^4*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

3.246. $\int \cot^4(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

output $1/d*(A*a^2*(-1/3*\cot(d*x+c)^3+\cot(d*x+c)+d*x+c)+B*a^2*(-1/2*\cot(d*x+c)^2-\ln(\sin(d*x+c)))+2*A*a*b*(-1/2*\cot(d*x+c)^2-\ln(\sin(d*x+c)))+2*B*a*b*(-\cot(d*x+c)-d*x-c)+A*b^2*(-\cot(d*x+c)-d*x-c)+B*b^2*\ln(\sin(d*x+c)))$

3.246.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.33

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \frac{3(Ba^2 + 2Aab - Bb^2) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^3 + 3(Ba^2 + 2Aab - 2(Aa^2 - 2Bab - Ab^2)dx) + 6d \tan(dx+c)}{6d \tan(dx+c)}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output $-1/6*(3*(B*a^2 + 2*A*a*b - B*b^2)*\log(\tan(d*x + c)^2/(\tan(d*x + c)^2 + 1)) * \tan(d*x + c)^3 + 3*(B*a^2 + 2*A*a*b - 2*(A*a^2 - 2*B*a*b - A*b^2)*d*x)*\tan(d*x + c)^3 + 2*A*a^2 - 6*(A*a^2 - 2*B*a*b - A*b^2)*\tan(d*x + c)^2 + 3*(B*a^2 + 2*A*a*b)*\tan(d*x + c))/(d*\tan(d*x + c)^3)$

3.246.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 260 vs. 2(107) = 214.

Time = 1.13 (sec) , antiderivative size = 260, normalized size of antiderivative = 2.20

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \begin{cases} \tilde{\infty} Aa^2x \\ x(A + B \tan(c))(a + b \tan(c))^2 \cot^4(c) \\ \tilde{\infty} Aa^2x \\ Aa^2x + \frac{Aa^2}{d \tan(c+dx)} - \frac{Aa^2}{3d \tan^3(c+dx)} + \frac{Aab \log(\tan^2(c+dx)+1)}{d} - \frac{2Aab \log(\tan(c+dx))}{d} - \frac{Aab}{d \tan^2(c+dx)} - Ab^2x - \frac{Ab^2}{d \tan(c+dx)} \end{cases}$$

input `integrate(cot(d*x+c)**4*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `Piecewise((zoo*A*a**2*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**2*cot(c)**4, Eq(d, 0)), (zoo*A*a**2*x, Eq(c, -d*x)), (A*a**2*x + A*a**2/(d*tan(c + d*x)) - A*a**2/(3*d*tan(c + d*x)**3) + A*a*b*log(tan(c + d*x)**2 + 1)/d - 2*A*a*b*log(tan(c + d*x))/d - A*a*b/(d*tan(c + d*x)**2) - A*b**2*x - A*b**2/(d*tan(c + d*x)) + B*a**2*log(tan(c + d*x)**2 + 1)/(2*d) - B*a**2*log(tan(c + d*x))/d - B*a**2/(2*d*tan(c + d*x)**2) - 2*B*a*b*x - 2*B*a*b/(d*tan(c + d*x)) - B*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + B*b**2*log(tan(c + d*x))/d, True))`

3.246.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.26

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{6(Aa^2 - 2Bab - Ab^2)(dx + c) + 3(Ba^2 + 2Aab - Bb^2) \log(\tan(dx + c)^2 + 1) - 6(Ba^2 + 2Aab - Bb^2)}{6d}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/6*(6*(A*a^2 - 2*B*a*b - A*b^2)*(d*x + c) + 3*(B*a^2 + 2*A*a*b - B*b^2)*log(tan(d*x + c)^2 + 1) - 6*(B*a^2 + 2*A*a*b - B*b^2)*log(tan(d*x + c)) - (2*A*a^2 - 6*(A*a^2 - 2*B*a*b - A*b^2)*tan(d*x + c)^2 + 3*(B*a^2 + 2*A*a*b)*tan(d*x + c))/tan(d*x + c)^3)/d`

3.246.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 334 vs. 2(114) = 228.

Time = 1.43 (sec) , antiderivative size = 334, normalized size of antiderivative = 2.83

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{Aa^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 3Ba^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 6Aab \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 15Aa^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 24}{6d}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output
$$\frac{1}{24}(A^2 \tan^3(\frac{1}{2}dx + \frac{1}{2}c) - 3B A^2 \tan^2(\frac{1}{2}dx + \frac{1}{2}c) - 6A a b \tan(\frac{1}{2}dx + \frac{1}{2}c) + 15A^2 a \tan(\frac{1}{2}dx + \frac{1}{2}c) + 24B a b \tan(\frac{1}{2}dx + \frac{1}{2}c) + 12A b^2 \tan(\frac{1}{2}dx + \frac{1}{2}c) + 24(A^2 - 2B a b - A b^2)(dx + c) + 24(B A^2 + 2A a b - B b^2) \log(\tan(\frac{1}{2}dx + \frac{1}{2}c)^2 + 1) - 24(B A^2 + 2A a b - B b^2) \log(\text{abs}(\tan(\frac{1}{2}dx + \frac{1}{2}c))) + (44B A^2 \tan^3(\frac{1}{2}dx + \frac{1}{2}c) + 88A a b \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 - 44B b^2 \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 + 15A^2 a \tan^2(\frac{1}{2}dx + \frac{1}{2}c) - 24B a b \tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 12A b^2 \tan(\frac{1}{2}dx + \frac{1}{2}c)^2 - 3B A^2 \tan(\frac{1}{2}dx + \frac{1}{2}c) - 6A a b \tan(\frac{1}{2}dx + \frac{1}{2}c) - A^2) / \tan(\frac{1}{2}dx + \frac{1}{2}c)^3) / d$$

3.246.9 Mupad [B] (verification not implemented)

Time = 7.95 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.32

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$\frac{\cot(c + dx)^3 \left(\frac{A a^2}{3} + \tan(c + dx)^2 (-A a^2 + 2 B a b + A b^2) + \tan(c + dx) \left(\frac{B a^2}{2} + A b a \right) \right)}{d} - \frac{\ln(\tan(c + dx)) (B a^2 + 2 A a b - B b^2)}{d} + \frac{\ln(\tan(c + dx) - i) (-B + A i) (-b + a i)^2}{2d} - \frac{\ln(\tan(c + dx) + i) (B + A i) (b + a i)^2}{2d}$$

input `int(cot(c + d*x)^4*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)`

output
$$(\log(\tan(c + d*x) - i) * (A * i - B) * (a * i - b)^2) / (2 * d) - (\log(\tan(c + d*x)) * (B * a^2 - B * b^2 + 2 * A * a * b)) / d - (\cot(c + d*x)^3 * ((A * a^2) / 3 + \tan(c + d*x)^2 * (A * b^2 - A * a^2 + 2 * B * a * b) + \tan(c + d*x) * ((B * a^2) / 2 + A * a * b))) / d - (\log(\tan(c + d*x) + i) * (A * i + B) * (a * i + b)^2) / (2 * d)$$

3.247 $\int \cot^5(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.247.1 Optimal result	2399
3.247.2 Mathematica [C] (verified)	2400
3.247.3 Rubi [A] (verified)	2400
3.247.4 Maple [A] (verified)	2404
3.247.5 Fricas [A] (verification not implemented)	2405
3.247.6 Sympy [B] (verification not implemented)	2405
3.247.7 Maxima [A] (verification not implemented)	2406
3.247.8 Giac [B] (verification not implemented)	2407
3.247.9 Mupad [B] (verification not implemented)	2407

3.247.1 Optimal result

Integrand size = 31, antiderivative size = 151

$$\begin{aligned} & \int \cot^5(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx \\ &= (2aAb + a^2B - b^2B) x - \frac{(b^2B - a(2Ab + aB)) \cot(c+dx)}{d} \\ & \quad + \frac{(a^2A - Ab^2 - 2abB) \cot^2(c+dx)}{2d} - \frac{a(2Ab + aB) \cot^3(c+dx)}{3d} \\ & \quad - \frac{a^2A \cot^4(c+dx)}{4d} + \frac{(a^2A - Ab^2 - 2abB) \log(\sin(c+dx))}{d} \end{aligned}$$

output

```
(2*A*a*b+B*a^2-B*b^2)*x-(B*b^2-a*(2*A*b+B*a))*cot(d*x+c)/d+1/2*(A*a^2-A*b^2-2*B*a*b)*cot(d*x+c)^2/d-1/3*a*(2*A*b+B*a)*cot(d*x+c)^3/d-1/4*a^2*A*cot(d*x+c)^4/d+(A*a^2-A*b^2-2*B*a*b)*ln(sin(d*x+c))/d
```

3.247.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.10 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.19

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{12(2aAb + a^2B - b^2B) \cot(c + dx) + 6(a^2A - Ab^2 - 2abB) \cot^2(c + dx) - 4a(2Ab + aB) \cot^3(c + dx) - \dots}{(12*d)}$$

input `Integrate[Cot[c + d*x]^5*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output $(12*(2*a*A*b + a^2*B - b^2*B)*\text{Cot}[c + d*x] + 6*(a^2*A - A*b^2 - 2*a*b*B)*\text{Cot}[c + d*x]^2 - 4*a*(2*A*b + a*B)*\text{Cot}[c + d*x]^3 - 3*a^2*A*\text{Cot}[c + d*x]^4 - 6*((a + I*b)^2*(A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]] + (-2*a^2*A + 2*A*b^2 + 4*a*b*B)*\text{Log}[\text{Tan}[c + d*x]] + (a - I*b)^2*(A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]]))/ (12*d)$

3.247.3 Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 4087, 3042, 4111, 25, 3042, 4012, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^2(A + B \tan(c + dx))}{\tan(c + dx)^5} dx$$

$$\downarrow \text{4087}$$

$$\int \cot^4(c + dx) (b^2 B \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)) dx - \frac{a^2 A \cot^4(c + dx)}{4d}$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
& \int \frac{b^2 B \tan(c+dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2Ab + aB)}{\tan(c+dx)^4} dx - \frac{a^2 A \cot^4(c+dx)}{4d} \\
& \quad \downarrow 4111 \\
& \int -\cot^3(c+dx) (Aa^2 - 2bBa - Ab^2 - (b^2 B - a(2Ab + aB)) \tan(c+dx)) dx - \\
& \quad \frac{a^2 A \cot^4(c+dx)}{4d} - \frac{a(aB + 2Ab) \cot^3(c+dx)}{3d} \\
& \quad \downarrow 25 \\
& - \int \cot^3(c+dx) (Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2 B) \tan(c+dx)) dx - \\
& \quad \frac{a^2 A \cot^4(c+dx)}{4d} - \frac{a(aB + 2Ab) \cot^3(c+dx)}{3d} \\
& \quad \downarrow 3042 \\
& - \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2 B) \tan(c+dx)}{\tan(c+dx)^3} dx - \frac{a^2 A \cot^4(c+dx)}{4d} - \\
& \quad \frac{a(aB + 2Ab) \cot^3(c+dx)}{3d} \\
& \quad \downarrow 4012 \\
& - \int \cot^2(c+dx) (Ba^2 + 2Aba - b^2 B - (Aa^2 - 2bBa - Ab^2) \tan(c+dx)) dx + \\
& \quad \frac{(a^2 A - 2abB - Ab^2) \cot^2(c+dx)}{2d} - \frac{a^2 A \cot^4(c+dx)}{4d} - \frac{a(aB + 2Ab) \cot^3(c+dx)}{3d} \\
& \quad \downarrow 3042 \\
& - \int \frac{Ba^2 + 2Aba - b^2 B - (Aa^2 - 2bBa - Ab^2) \tan(c+dx)}{\tan(c+dx)^2} dx + \\
& \quad \frac{(a^2 A - 2abB - Ab^2) \cot^2(c+dx)}{2d} - \frac{a^2 A \cot^4(c+dx)}{4d} - \frac{a(aB + 2Ab) \cot^3(c+dx)}{3d} \\
& \quad \downarrow 4012 \\
& - \int -\cot(c+dx) (Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2 B) \tan(c+dx)) dx + \\
& \quad \frac{(a^2 A - 2abB - Ab^2) \cot^2(c+dx)}{2d} + \frac{(a^2 B + 2aAb - b^2 B) \cot(c+dx)}{d} - \frac{a^2 A \cot^4(c+dx)}{4d} - \\
& \quad \frac{a(aB + 2Ab) \cot^3(c+dx)}{3d} \\
& \quad \downarrow 25
\end{aligned}$$

$$\begin{aligned}
& \int \cot(c+dx) (Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c+dx)) dx + \\
& \frac{(a^2A - 2abB - Ab^2) \cot^2(c+dx)}{2d} + \frac{(a^2B + 2aAb - b^2B) \cot(c+dx)}{d} - \frac{a^2A \cot^4(c+dx)}{4d} - \\
& \frac{a(aB + 2Ab) \cot^3(c+dx)}{3d} \\
& \quad \downarrow \text{3042} \\
& \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c+dx)}{\tan(c+dx)} dx + \frac{(a^2A - 2abB - Ab^2) \cot^2(c+dx)}{2d} + \\
& \frac{(a^2B + 2aAb - b^2B) \cot(c+dx)}{d} - \frac{a^2A \cot^4(c+dx)}{4d} - \frac{a(aB + 2Ab) \cot^3(c+dx)}{3d} \\
& \quad \downarrow \text{4014} \\
& \frac{(a^2A - 2abB - Ab^2) \int \cot(c+dx) dx + (a^2A - 2abB - Ab^2) \cot^2(c+dx)}{2d} + \\
& \frac{(a^2B + 2aAb - b^2B) \cot(c+dx)}{d} + x(a^2B + 2aAb - b^2B) - \frac{a^2A \cot^4(c+dx)}{4d} - \\
& \frac{a(aB + 2Ab) \cot^3(c+dx)}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{(a^2A - 2abB - Ab^2) \int -\tan\left(c+dx + \frac{\pi}{2}\right) dx + (a^2A - 2abB - Ab^2) \cot^2(c+dx)}{2d} + \\
& \frac{(a^2B + 2aAb - b^2B) \cot(c+dx)}{d} + x(a^2B + 2aAb - b^2B) - \frac{a^2A \cot^4(c+dx)}{4d} - \\
& \frac{a(aB + 2Ab) \cot^3(c+dx)}{3d} \\
& \quad \downarrow \text{25} \\
& -(a^2A - 2abB - Ab^2) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + \frac{(a^2A - 2abB - Ab^2) \cot^2(c+dx)}{2d} + \\
& \frac{(a^2B + 2aAb - b^2B) \cot(c+dx)}{d} + x(a^2B + 2aAb - b^2B) - \frac{a^2A \cot^4(c+dx)}{4d} - \\
& \frac{a(aB + 2Ab) \cot^3(c+dx)}{3d} \\
& \quad \downarrow \text{3956} \\
& \frac{(a^2A - 2abB - Ab^2) \cot^2(c+dx)}{2d} + \frac{(a^2B + 2aAb - b^2B) \cot(c+dx)}{d} + \\
& \frac{(a^2A - 2abB - Ab^2) \log(-\sin(c+dx))}{d} + x(a^2B + 2aAb - b^2B) - \frac{a^2A \cot^4(c+dx)}{4d} - \\
& \frac{a(aB + 2Ab) \cot^3(c+dx)}{3d}
\end{aligned}$$

3.247. $\int \cot^5(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

input `Int[Cot[c + d*x]^5*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(2*a*A*b + a^2*B - b^2*B)*x + ((2*a*A*b + a^2*B - b^2*B)*Cot[c + d*x])/d + ((a^2*A - A*b^2 - 2*a*b*B)*Cot[c + d*x]^2)/(2*d) - (a*(2*A*b + a*B)*Cot[c + d*x]^3)/(3*d) - (a^2*A*Cot[c + d*x]^4)/(4*d) + ((a^2*A - A*b^2 - 2*a*b*B)*Log[-Sin[c + d*x]])/d`

3.247.3.1 Defintions of rubi rules used

rule 25 `Int[-(F*x_), x_Symbol] := Simp[Identity[-1] Int[F*x, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

```
rule 4087 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(- (B*c - A*d))*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

```
rule 4111 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]
```

3.247.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.07

method	result
derivativedivides	$A a^2 \left(-\frac{\cot^4(dx+c)}{4} + \frac{\cot^2(dx+c)}{2} + \ln(\sin(dx+c)) \right) + B a^2 \left(-\frac{\cot^3(dx+c)}{3} + \cot(dx+c) + dx+c \right) + 2Aab \left(-\frac{\cot^3(dx+c)}{3} + \cot(dx+c) + dx+c \right)$
default	$A a^2 \left(-\frac{\cot^4(dx+c)}{4} + \frac{\cot^2(dx+c)}{2} + \ln(\sin(dx+c)) \right) + B a^2 \left(-\frac{\cot^3(dx+c)}{3} + \cot(dx+c) + dx+c \right) + 2Aab \left(-\frac{\cot^3(dx+c)}{3} + \cot(dx+c) + dx+c \right)$
parallelrisch	$\frac{6(-A a^2 + A b^2 + 2Bab) \ln(\sec^2(dx+c)) + 12(A a^2 - A b^2 - 2Bab) \ln(\tan(dx+c)) - 3A(\cot^4(dx+c))a^2 + 4(-2Aab - B a^2)}{12d} \left(\frac{(2Aab + B a^2 - B b^2)(\tan^3(dx+c))}{d} + (2Aab + B a^2 - B b^2)x(\tan^4(dx+c)) - \frac{A a^2}{4d} + \frac{(A a^2 - A b^2 - 2Bab)(\tan^2(dx+c))}{2d} - \frac{a(2Ab + B a^2)}{2d} \right)$
norman	$\frac{(2Aab + B a^2 - B b^2)(\tan^3(dx+c))}{d} + (2Aab + B a^2 - B b^2)x(\tan^4(dx+c)) - \frac{A a^2}{4d} + \frac{(A a^2 - A b^2 - 2Bab)(\tan^2(dx+c))}{2d} - \frac{a(2Ab + B a^2)}{2d}$
risch	$2Aabx + B a^2x - B b^2x + \frac{2iA b^2c}{d} + iA b^2x + \frac{4iBabc}{d} - iA a^2x + 2iBabx - \frac{2ia^2Ac}{d} - \frac{2i(6iE)}{d}$

```
input int(cot(d*x+c)^5*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

output $1/d*(A*a^2*(-1/4*\cot(d*x+c)^4+1/2*\cot(d*x+c)^2+\ln(\sin(d*x+c)))+B*a^2*(-1/3*\cot(d*x+c)^3+\cot(d*x+c)+d*x+c)+2*A*a*b*(-1/3*\cot(d*x+c)^3+\cot(d*x+c)+d*x+c)+2*B*a*b*(-1/2*\cot(d*x+c)^2-\ln(\sin(d*x+c)))+A*b^2*(-1/2*\cot(d*x+c)^2-\ln(\sin(d*x+c)))+B*b^2*(-\cot(d*x+c)-d*x-c))$

3.247.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.26

$$\int \cot^5(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

$$= \frac{6(Aa^2 - 2Bab - Ab^2) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^4 + 3(3Aa^2 - 4Bab - 2Ab^2 + 4(Ba^2 + 2Aab - B$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output $1/12*(6*(A*a^2 - 2*B*a*b - A*b^2)*\log(\tan(d*x + c)^2/(\tan(d*x + c)^2 + 1)) * \tan(d*x + c)^4 + 3*(3*A*a^2 - 4*B*a*b - 2*A*b^2 + 4*(B*a^2 + 2*A*a*b - B*b^2)*d*x)*\tan(d*x + c)^4 + 12*(B*a^2 + 2*A*a*b - B*b^2)*\tan(d*x + c)^3 - 3*A*a^2 + 6*(A*a^2 - 2*B*a*b - A*b^2)*\tan(d*x + c)^2 - 4*(B*a^2 + 2*A*a*b)*\tan(d*x + c))/(d*\tan(d*x + c)^4)$

3.247.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 313 vs. $2(136) = 272$.

Time = 1.85 (sec) , antiderivative size = 313, normalized size of antiderivative = 2.07

$$\int \cot^5(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

$$= \begin{cases} \tilde{\infty}Aa^2x \\ x(A+B\tan(c))(a+b\tan(c))^2\cot^5(c) \\ \tilde{\infty}Aa^2x \\ -\frac{Aa^2\log(\tan^2(c+dx)+1)}{2d} + \frac{Aa^2\log(\tan(c+dx))}{d} + \frac{Aa^2}{2d\tan^2(c+dx)} - \frac{Aa^2}{4d\tan^4(c+dx)} + 2Aabx + \frac{2Aab}{d\tan(c+dx)} - \frac{2Aab}{3d\tan^3(c+dx)} \end{cases}$$

input `integrate(cot(d*x+c)**5*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `Piecewise((zoo*A*a**2*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**2*cot(c)**5, Eq(d, 0)), (zoo*A*a**2*x, Eq(c, -d*x)), (-A*a**2*log(tan(c + d*x)**2 + 1)/(2*d) + A*a**2*log(tan(c + d*x))/d + A*a**2/(2*d*tan(c + d*x)**2) - A*a**2/(4*d*tan(c + d*x)**4) + 2*A*a*b*x + 2*A*a*b/(d*tan(c + d*x)) - 2*A*a*b/(3*d*tan(c + d*x)**3) + A*b**2*log(tan(c + d*x)**2 + 1)/(2*d) - A*b**2*log(tan(c + d*x))/d - A*b**2/(2*d*tan(c + d*x)**2) + B*a**2*x + B*a**2/(d*tan(c + d*x)) - B*a**2/(3*d*tan(c + d*x)**3) + B*a*b*log(tan(c + d*x)**2 + 1)/d - 2*B*a*b*log(tan(c + d*x))/d - B*a*b/(d*tan(c + d*x)**2) - B*b**2*x - B*b**2/(d*tan(c + d*x)), True))`

3.247.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.16

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{12(Ba^2 + 2Aab - Bb^2)(dx + c) - 6(Aa^2 - 2Bab - Ab^2) \log(\tan(dx + c)^2 + 1) + 12(Aa^2 - 2Bab - Ab^2)}{12d}$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/12*(12*(B*a^2 + 2*A*a*b - B*b^2)*(d*x + c) - 6*(A*a^2 - 2*B*a*b - A*b^2)*log(tan(d*x + c)^2 + 1) + 12*(A*a^2 - 2*B*a*b - A*b^2)*log(tan(d*x + c)) + (12*(B*a^2 + 2*A*a*b - B*b^2)*tan(d*x + c)^3 - 3*A*a^2 + 6*(A*a^2 - 2*B*a*b - A*b^2)*tan(d*x + c)^2 - 4*(B*a^2 + 2*A*a*b)*tan(d*x + c))/tan(d*x + c)^4/d`

3.247.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 435 vs. $2(145) = 290$.

Time = 1.18 (sec) , antiderivative size = 435, normalized size of antiderivative = 2.88

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$3 A a^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 - 8 B a^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 16 A a b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 36 A a^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

$$\begin{aligned} & -1/192*(3*A*a^2*\tan(1/2*d*x + 1/2*c)^4 - 8*B*a^2*\tan(1/2*d*x + 1/2*c)^3 - \\ & 16*A*a*b*\tan(1/2*d*x + 1/2*c)^3 - 36*A*a^2*\tan(1/2*d*x + 1/2*c)^2 + 48*B*a \\ & *b*\tan(1/2*d*x + 1/2*c)^2 + 24*A*b^2*\tan(1/2*d*x + 1/2*c)^2 + 120*B*a^2*ta \\ & n(1/2*d*x + 1/2*c) + 240*A*a*b*\tan(1/2*d*x + 1/2*c) - 96*B*b^2*\tan(1/2*d*x \\ & + 1/2*c) - 192*(B*a^2 + 2*A*a*b - B*b^2)*(d*x + c) + 192*(A*a^2 - 2*B*a*b \\ & - A*b^2)*\log(\tan(1/2*d*x + 1/2*c)^2 + 1) - 192*(A*a^2 - 2*B*a*b - A*b^2)* \\ & \log(\text{abs}(\tan(1/2*d*x + 1/2*c))) + (400*A*a^2*\tan(1/2*d*x + 1/2*c)^4 - 800*B \\ & *a*b*\tan(1/2*d*x + 1/2*c)^4 - 400*A*b^2*\tan(1/2*d*x + 1/2*c)^4 - 120*B*a^2 \\ & * \tan(1/2*d*x + 1/2*c)^3 - 240*A*a*b*\tan(1/2*d*x + 1/2*c)^3 + 96*B*b^2*\tan(\\ & 1/2*d*x + 1/2*c)^3 - 36*A*a^2*\tan(1/2*d*x + 1/2*c)^2 + 48*B*a*b*\tan(1/2*d* \\ & x + 1/2*c)^2 + 24*A*b^2*\tan(1/2*d*x + 1/2*c)^2 + 8*B*a^2*\tan(1/2*d*x + 1/2 \\ & *c) + 16*A*a*b*\tan(1/2*d*x + 1/2*c) + 3*A*a^2)/\tan(1/2*d*x + 1/2*c)^4)/d \end{aligned}$$
3.247.9 Mupad [B] (verification not implemented)

Time = 8.04 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.21

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$\frac{\cot(c + dx)^4 \left(\frac{Aa^2}{4} + \tan(c + dx)^2 \left(-\frac{Aa^2}{2} + B a b + \frac{A b^2}{2} \right) - \tan(c + dx)^3 (B a^2 + 2 A a b - B b^2) + \tan(c + dx)^4 (A + B \tan(c + dx)) \right)}{d}$$

$$- \frac{\ln(\tan(c + dx)) (-A a^2 + 2 B a b + A b^2)}{d}$$

$$+ \frac{\ln(\tan(c + dx) + i) (A - B i) (b + a i)^2}{2 d}$$

$$+ \frac{\ln(\tan(c + dx) - i) (A + B i) (-b + a i)^2}{2 d}$$

3.247. $\int \cot^5(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

input `int(cot(c + d*x)^5*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)`

output `(log(tan(c + d*x) + 1i)*(A - B*1i)*(a*1i + b)^2)/(2*d) - (log(tan(c + d*x))*(A*b^2 - A*a^2 + 2*B*a*b))/d - (cot(c + d*x)^4*((A*a^2)/4 + tan(c + d*x)^2*((A*b^2)/2 - (A*a^2)/2 + B*a*b) - tan(c + d*x)^3*(B*a^2 - B*b^2 + 2*A*a*b) + tan(c + d*x)*((B*a^2)/3 + (2*A*a*b)/3)))/d + (log(tan(c + d*x) - 1i) *(A + B*1i)*(a*1i - b)^2)/(2*d)`

3.248 $\int \tan^2(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.248.1 Optimal result	2409
3.248.2 Mathematica [C] (verified)	2410
3.248.3 Rubi [A] (verified)	2410
3.248.4 Maple [A] (verified)	2414
3.248.5 Fricas [A] (verification not implemented)	2414
3.248.6 Sympy [B] (verification not implemented)	2415
3.248.7 Maxima [A] (verification not implemented)	2415
3.248.8 Giac [B] (verification not implemented)	2416
3.248.9 Mupad [B] (verification not implemented)	2417

3.248.1 Optimal result

Integrand size = 31, antiderivative size = 201

$$\int \tan^2(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= -((a^3A - 3aAb^2 - 3a^2bB + b^3B) x) + \frac{(3a^2Ab - Ab^3 + a^3B - 3ab^2B) \log(\cos(c+dx))}{d} - \frac{b(2aAb + a^2B - b^2B) \tan(c+dx)}{d} - \frac{(Ab + aB)(a + b \tan(c+dx))^2}{2d} - \frac{B(a + b \tan(c+dx))^3}{3d} + \frac{(5Ab - aB)(a + b \tan(c+dx))^4}{20b^2d} + \frac{B \tan(c+dx)(a + b \tan(c+dx))^4}{5bd}$$

```
output - (A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*x+(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*ln(
cos(d*x+c))/d-b*(2*A*a*b+B*a^2-B*b^2)*tan(d*x+c)/d-1/2*(A*b+B*a)*(a+b*tan(
d*x+c))^2/d-1/3*B*(a+b*tan(d*x+c))^3/d+1/20*(5*A*b-B*a)*(a+b*tan(d*x+c))^4
/b^2/d+1/5*B*tan(d*x+c)*(a+b*tan(d*x+c))^4/b/d
```


3.248.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.37 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.20

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{3(5Ab - aB)(a + b \tan(c + dx))^4}{b} + 12B \tan(c + dx)(a + b \tan(c + dx))^4 - 30(Ab - aB) ((ia - b)^3 \log(i - \tan(c + dx)) + \dots)$$

input `Integrate[Tan[c + d*x]^2*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output $((3*(5*A*b - a*B)*(a + b*\text{Tan}[c + d*x])^4)/b + 12*B*\text{Tan}[c + d*x]*(a + b*\text{Tan}[c + d*x])^4 - 30*(A*b - a*B)*((I*a - b)^3*\text{Log}[I - \text{Tan}[c + d*x]] - (I*a + b)^3*\text{Log}[I + \text{Tan}[c + d*x]] + 6*a*b^2*\text{Tan}[c + d*x] + b^3*\text{Tan}[c + d*x]^2) + 10*B*((3*I)*(a + I*b)^4*\text{Log}[I - \text{Tan}[c + d*x]] - (3*I)*(a - I*b)^4*\text{Log}[I + \text{Tan}[c + d*x]] + 6*b^2*(-6*a^2 + b^2)*\text{Tan}[c + d*x] - 12*a*b^3*\text{Tan}[c + d*x]^2 - 2*b^4*\text{Tan}[c + d*x]^3))/(60*b*d)$

3.248.3 Rubi [A] (verified)

Time = 1.06 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.07, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 4090, 25, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^2(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4090}$$

$$\int \frac{-(a + b \tan(c + dx))^3 (-((5Ab - aB) \tan^2(c + dx)) + 5bB \tan(c + dx) + aB) dx}{\frac{5b}{B \tan(c + dx)(a + b \tan(c + dx))^4} + 5bd} +$$

$$\downarrow \text{25}$$

3.248. $\int \tan^2(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^4}{5bd} - \frac{\int (a+b \tan(c+dx))^3 \left(-((5Ab-aB) \tan^2(c+dx)) + 5bB \tan(c+dx) + aB \right) dx}{5b} \\
& \quad \downarrow \text{3042} \\
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^4}{5bd} - \frac{\int (a+b \tan(c+dx))^3 \left(-((5Ab-aB) \tan(c+dx)^2) + 5bB \tan(c+dx) + aB \right) dx}{5b} \\
& \quad \downarrow \text{4113} \\
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^4}{5bd} - \frac{\int (a+b \tan(c+dx))^3 (5Ab + 5B \tan(c+dx)b) dx - \frac{(5Ab-aB)(a+b \tan(c+dx))^4}{4bd}}{5b} \\
& \quad \downarrow \text{3042} \\
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^4}{5bd} - \frac{\int (a+b \tan(c+dx))^3 (5Ab + 5B \tan(c+dx)b) dx - \frac{(5Ab-aB)(a+b \tan(c+dx))^4}{4bd}}{5b} \\
& \quad \downarrow \text{4011} \\
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^4}{5bd} - \frac{\int (a+b \tan(c+dx))^2 (5b(aA-bB) + 5b(Ab+aB) \tan(c+dx)) dx - \frac{(5Ab-aB)(a+b \tan(c+dx))^4}{4bd} + \frac{5bB(a+b \tan(c+dx))^3}{3d}}{5b} \\
& \quad \downarrow \text{3042} \\
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^4}{5bd} - \frac{\int (a+b \tan(c+dx))^2 (5b(aA-bB) + 5b(Ab+aB) \tan(c+dx)) dx - \frac{(5Ab-aB)(a+b \tan(c+dx))^4}{4bd} + \frac{5bB(a+b \tan(c+dx))^3}{3d}}{5b} \\
& \quad \downarrow \text{4011} \\
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^4}{5bd} - \frac{\int (a+b \tan(c+dx)) (5b(Aa^2 - 2bBa - Ab^2) + 5b(Ba^2 + 2Aba - b^2B) \tan(c+dx)) dx - \frac{(5Ab-aB)(a+b \tan(c+dx))^4}{4bd}}{5b} \\
& \quad \downarrow \text{3042} \\
& \frac{B \tan(c+dx)(a+b \tan(c+dx))^4}{5bd} - \frac{\int (a+b \tan(c+dx)) (5b(Aa^2 - 2bBa - Ab^2) + 5b(Ba^2 + 2Aba - b^2B) \tan(c+dx)) dx - \frac{(5Ab-aB)(a+b \tan(c+dx))^4}{4bd}}{5b}
\end{aligned}$$

3.248. $\int \tan^2(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\begin{array}{c}
\downarrow 4008 \\
\frac{B \tan(c+dx)(a+b \tan(c+dx))^4}{5bd} - \\
\frac{5b(a^3B+3a^2Ab-3ab^2B-Ab^3) \int \tan(c+dx)dx + \frac{5b^2(a^2B+2aAb-b^2B) \tan(c+dx)}{d} + 5bx(a^3A-3a^2bB-3aAb^2+b^3B)}{5b} \\
\downarrow 3042 \\
\frac{B \tan(c+dx)(a+b \tan(c+dx))^4}{5bd} - \\
\frac{5b(a^3B+3a^2Ab-3ab^2B-Ab^3) \int \tan(c+dx)dx + \frac{5b^2(a^2B+2aAb-b^2B) \tan(c+dx)}{d} + 5bx(a^3A-3a^2bB-3aAb^2+b^3B)}{5b} \\
\downarrow 3956 \\
\frac{B \tan(c+dx)(a+b \tan(c+dx))^4}{5bd} - \\
\frac{\frac{5b^2(a^2B+2aAb-b^2B) \tan(c+dx)}{d} - \frac{5b(a^3B+3a^2Ab-3ab^2B-Ab^3) \log(\cos(c+dx))}{d} + 5bx(a^3A-3a^2bB-3aAb^2+b^3B) - \frac{(5Ab-a^2B)}{d}}{5b}
\end{array}$$

input `Int[Tan[c + d*x]^2*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(B*Tan[c + d*x]*(a + b*Tan[c + d*x])^4)/(5*b*d) - (5*b*(a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*x - (5*b*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*Log[Cos[c + d*x]])/d + (5*b^2*(2*a*A*b + a^2*B - b^2*B)*Tan[c + d*x])/d + (5*b*(A*b + a*B)*(a + b*Tan[c + d*x])^2)/(2*d) + (5*b*B*(a + b*Tan[c + d*x])^3)/(3*d) - ((5*A*b - a*B)*(a + b*Tan[c + d*x])^4)/(4*b*d))/(5*b)`

3.248.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4011 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4090 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

3.248.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.03

method	result
parts	$\frac{(A b^3 + 3 B a b^2) \left(\frac{\tan^4(dx+c)}{4} - \frac{\tan^2(dx+c)}{2} + \frac{\ln(1+\tan^2(dx+c))}{2} \right)}{d} + \frac{(3 A a b^2 + 3 B a^2 b) \left(\frac{\tan^3(dx+c)}{3} - \tan(dx+c) \right)}{d}$
norman	$(-A a^3 + 3 A a b^2 + 3 B a^2 b - B b^3) x + \frac{(A a^3 - 3 A a b^2 - 3 B a^2 b + B b^3) \tan(dx+c)}{d} + \frac{(3 A a^2 b - A b^3 + B a^3) \arctan(\tan(dx+c))}{d}$
derivativedivides	$\frac{B b^3 (\tan^5(dx+c))}{5} + \frac{A b^3 (\tan^4(dx+c))}{4} + \frac{3 B a b^2 (\tan^4(dx+c))}{4} + A a b^2 (\tan^3(dx+c)) + B a^2 b (\tan^3(dx+c)) - \frac{B b^3 (\tan^3(dx+c))}{3}$
default	$\frac{B b^3 (\tan^5(dx+c))}{5} + \frac{A b^3 (\tan^4(dx+c))}{4} + \frac{3 B a b^2 (\tan^4(dx+c))}{4} + A a b^2 (\tan^3(dx+c)) + B a^2 b (\tan^3(dx+c)) - \frac{B b^3 (\tan^3(dx+c))}{3}$
parallelrisch	$- \frac{-12 B b^3 (\tan^5(dx+c)) - 15 A b^3 (\tan^4(dx+c)) - 45 B a b^2 (\tan^4(dx+c)) - 60 A a b^2 (\tan^3(dx+c)) - 60 B a^2 b (\tan^3(dx+c))}{d}$
risch	$-A a^3 x + 3 A a b^2 x + 3 B a^2 b x - B b^3 x + \frac{2 i A b^3 c}{d} + \frac{6 i B a b^2 c}{d} - 3 i A a^2 b x + i A b^3 x - \frac{6 i A a^2 b c}{d}$

input `int(tan(d*x+c)^2*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output $(A*b^3+3*B*a*b^2)/d*(1/4*\tan(d*x+c)^4-1/2*\tan(d*x+c)^2+1/2*\ln(1+\tan(d*x+c)^2))+ (3*A*a*b^2+3*B*a^2*b)/d*(1/3*\tan(d*x+c)^3-\tan(d*x+c)+\arctan(\tan(d*x+c)))+ (3*A*a^2*b+B*a^3)/d*(1/2*\tan(d*x+c)^2-1/2*\ln(1+\tan(d*x+c)^2))+ A*a^3/d*(\tan(d*x+c)-\arctan(\tan(d*x+c)))+ B*b^3/d*(1/5*\tan(d*x+c)^5-1/3*\tan(d*x+c)^3+\tan(d*x+c)-\arctan(\tan(d*x+c)))$

3.248.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.06

$$\int \tan^2(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{12 B b^3 \tan(dx+c)^5 + 15 (3 B a b^2 + A b^3) \tan(dx+c)^4 + 20 (3 B a^2 b + 3 A a b^2 - B b^3) \tan(dx+c)^3 - 60 (A a^3 + B a^2 b + 3 A a b^2 + B b^3) \tan(dx+c)^2 + 60 (A a^2 b + B a b^2) \tan(dx+c) - 60 A a^2 b \arctan(\tan(dx+c))}{d}$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output $\frac{1}{60}(12Bb^3 \tan(dx+c)^5 + 15(3Bab^2 + Ab^3) \tan(dx+c)^4 + 20(3Ba^2b + 3Aab^2 - Bb^3) \tan(dx+c)^3 - 60(Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3) dx + 30(Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3) \tan(dx+c)^2 + 30(Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3) \log(1/(\tan(dx+c)^2 + 1)) + 60(Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3) \tan(dx+c))/d$

3.248.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 384 vs. $2(185) = 370$.

Time = 0.18 (sec) , antiderivative size = 384, normalized size of antiderivative = 1.91

$$\int \tan^2(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \begin{cases} -Aa^3x + \frac{Aa^3 \tan(c+dx)}{d} - \frac{3Aa^2b \log(\tan^2(c+dx)+1)}{2d} + \frac{3Aa^2b \tan^2(c+dx)}{2d} + 3Aab^2x + \frac{Aab^2 \tan^3(c+dx)}{d} - \frac{3Aab^2 \tan(c+dx)}{d} \\ x(A+B \tan(c))(a+b \tan(c))^3 \tan^2(c) \end{cases}$$

input `integrate(tan(d*x+c)**2*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)), x)`

output `Piecewise((-A*a**3*x + A*a**3*tan(c + d*x)/d - 3*A*a**2*b*log(tan(c + d*x)**2 + 1)/(2*d) + 3*A*a**2*b*tan(c + d*x)**2/(2*d) + 3*A*a*b**2*x + A*a*b**2*tan(c + d*x)**3/d - 3*A*a*b**2*tan(c + d*x)/d + A*b**3*log(tan(c + d*x)**2 + 1)/(2*d) + A*b**3*tan(c + d*x)**4/(4*d) - A*b**3*tan(c + d*x)**2/(2*d) - B*a**3*log(tan(c + d*x)**2 + 1)/(2*d) + B*a**3*tan(c + d*x)**2/(2*d) + 3*B*a**2*b*x + B*a**2*b*tan(c + d*x)**3/d - 3*B*a**2*b*tan(c + d*x)/d + 3*B*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + 3*B*a*b**2*tan(c + d*x)**4/(4*d) - 3*B*a*b**2*tan(c + d*x)**2/(2*d) - B*b**3*x + B*b**3*tan(c + d*x)**5/(5*d) - B*b**3*tan(c + d*x)**3/(3*d) + B*b**3*tan(c + d*x)/d, Ne(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**3*tan(c)**2, True))`

3.248.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.06

$$\int \tan^2(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{12Bb^3 \tan(dx+c)^5 + 15(3Bab^2 + Ab^3) \tan(dx+c)^4 + 20(3Ba^2b + 3Aab^2 - Bb^3) \tan(dx+c)^3 + 30(Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3) \tan(dx+c)^2 + 30(Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3) \tan(dx+c) + 30(Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3) \log(\tan(dx+c)^2 + 1) + 60(Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3) \tan(dx+c)}{d}$$

3.248. $\int \tan^2(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/60*(12*B*b^3*tan(d*x + c)^5 + 15*(3*B*a*b^2 + A*b^3)*tan(d*x + c)^4 + 20*(3*B*a^2*b + 3*A*a*b^2 - B*b^3)*tan(d*x + c)^3 + 30*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*tan(d*x + c)^2 - 60*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*(d*x + c) - 30*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*log(tan(d*x + c)^2 + 1) + 60*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*tan(d*x + c))/d`

3.248.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3757 vs. $2(192) = 384$.

Time = 3.79 (sec) , antiderivative size = 3757, normalized size of antiderivative = 18.69

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

```

-1/60*(60*A*a^3*d*x*tan(d*x)^5*tan(c)^5 - 180*B*a^2*b*d*x*tan(d*x)^5*tan(c)
)^5 - 180*A*a*b^2*d*x*tan(d*x)^5*tan(c)^5 + 60*B*b^3*d*x*tan(d*x)^5*tan(c)
^5 - 30*B*a^3*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)
)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^5*tan(c)^5 - 90*A*a^2*
b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2
+ tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^5*tan(c)^5 + 90*B*a*b^2*log(4*(tan
(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^
2 + tan(c)^2 + 1))*tan(d*x)^5*tan(c)^5 + 30*A*b^3*log(4*(tan(d*x)^2*tan(c)
^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 +
1))*tan(d*x)^5*tan(c)^5 - 300*A*a^3*d*x*tan(d*x)^4*tan(c)^4 + 900*B*a^2*b
*d*x*tan(d*x)^4*tan(c)^4 + 900*A*a*b^2*d*x*tan(d*x)^4*tan(c)^4 - 300*B*b^3
*d*x*tan(d*x)^4*tan(c)^4 - 30*B*a^3*tan(d*x)^5*tan(c)^5 - 90*A*a^2*b*tan(d
*x)^5*tan(c)^5 + 135*B*a*b^2*tan(d*x)^5*tan(c)^5 + 45*A*b^3*tan(d*x)^5*tan
(c)^5 + 150*B*a^3*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan
(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^4*tan(c)^4 + 450*A
*a^2*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan
(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^4*tan(c)^4 - 450*B*a*b^2*log(
4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan
(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^4*tan(c)^4 - 150*A*b^3*log(4*(tan(d*x)^2
*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + ...

```

3.248.9 Mupad [B] (verification not implemented)

Time = 7.31 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.08

$$\begin{aligned}
 & \int \tan^2(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx \\
 &= \frac{\tan(c + dx) (Aa^3 + Bb^3 - 3ab(Ab + Ba))}{d} \\
 & - \frac{\tan(c + dx)^3 \left(\frac{Bb^3}{3} - ab(Ab + Ba) \right)}{d} - x(Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3) \\
 & + \frac{\ln(\tan(c + dx)^2 + 1) \left(-\frac{Ba^3}{2} - \frac{3Aa^2b}{2} + \frac{3Bab^2}{2} + \frac{Ab^3}{2} \right)}{d} \\
 & + \frac{\tan(c + dx)^4 \left(\frac{Ab^3}{4} + \frac{3Bab^2}{4} \right)}{d} \\
 & - \frac{\tan(c + dx)^2 \left(-\frac{Ba^3}{2} - \frac{3Aa^2b}{2} + \frac{3Bab^2}{2} + \frac{Ab^3}{2} \right)}{d} + \frac{Bb^3 \tan(c + dx)^5}{5d}
 \end{aligned}$$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`

3.248. $\int \tan^2(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

output $(\tan(c + dx)(Aa^3 + Bb^3 - 3ab(Ab + Ba)))/d - (\tan(c + dx)^3((Bb^3)/3 - ab(Ab + Ba)))/d - x(Aa^3 + Bb^3 - 3Aa^2b - 3Ba^2b) + (\log(\tan(c + dx)^2 + 1)((Ab^3)/2 - (Ba^3)/2 - (3Aa^2b)/2 + (3Bab^2)/2))/d + (\tan(c + dx)^4((Ab^3)/4 + (3Bab^2)/4))/d - (\tan(c + dx)^2((Ab^3)/2 - (Ba^3)/2 - (3Aa^2b)/2 + (3Bab^2)/2))/d + (Bb^3 \tan(c + dx)^5)/(5d)$

3.249 $\int \tan(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.249.1 Optimal result	2419
3.249.2 Mathematica [C] (verified)	2420
3.249.3 Rubi [A] (verified)	2420
3.249.4 Maple [A] (verified)	2423
3.249.5 Fricas [A] (verification not implemented)	2424
3.249.6 Sympy [B] (verification not implemented)	2424
3.249.7 Maxima [A] (verification not implemented)	2425
3.249.8 Giac [B] (verification not implemented)	2425
3.249.9 Mupad [B] (verification not implemented)	2426

3.249.1 Optimal result

Integrand size = 29, antiderivative size = 165

$$\int \tan(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= -((3a^2Ab - Ab^3 + a^3B - 3ab^2B) x) - \frac{(a^3A - 3aAb^2 - 3a^2bB + b^3B) \log(\cos(c+dx))}{d}$$

$$+ \frac{b(a^2A - Ab^2 - 2abB) \tan(c+dx)}{d} + \frac{(aA - bB)(a+b \tan(c+dx))^2}{2d}$$

$$+ \frac{A(a+b \tan(c+dx))^3}{3d} + \frac{B(a+b \tan(c+dx))^4}{4bd}$$

output

```
-(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*x-(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*ln(
cos(d*x+c))/d+b*(A*a^2-A*b^2-2*B*a*b)*tan(d*x+c)/d+1/2*(A*a-B*b)*(a+b*tan(
d*x+c))^2/d+1/3*A*(a+b*tan(d*x+c))^3/d+1/4*B*(a+b*tan(d*x+c))^4/b/d
```

3.249.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.58 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.27

$$\int \tan(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{-6iA(a + ib)^4 \log(i - \tan(c + dx)) + 6iA(a - ib)^4 \log(i + \tan(c + dx)) - 12Ab^2(-6a^2 + b^2) \tan(c + dx)}{}$$

input `Integrate[Tan[c + d*x]*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output $((-6*I)*A*(a + I*b)^4*\text{Log}[I - \text{Tan}[c + d*x]] + (6*I)*A*(a - I*b)^4*\text{Log}[I + \text{Tan}[c + d*x]] - 12*A*b^2*(-6*a^2 + b^2)*\text{Tan}[c + d*x] + 24*a*A*b^3*\text{Tan}[c + d*x]^2 + 4*A*b^4*\text{Tan}[c + d*x]^3 + 3*B*(a + b*\text{Tan}[c + d*x])^4 - 6*(a*A + b*B)*((I*a - b)^3*\text{Log}[I - \text{Tan}[c + d*x]] - (I*a + b)^3*\text{Log}[I + \text{Tan}[c + d*x]] + 6*a*b^2*\text{Tan}[c + d*x] + b^3*\text{Tan}[c + d*x]^2))/(12*b*d)$

3.249.3 Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 4075, 3042, 4011, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4075}$$

$$\int (A \tan(c + dx) - B)(a + b \tan(c + dx))^3 dx + \frac{B(a + b \tan(c + dx))^4}{4bd}$$

$$\downarrow \text{3042}$$

$$\int (A \tan(c + dx) - B)(a + b \tan(c + dx))^3 dx + \frac{B(a + b \tan(c + dx))^4}{4bd}$$

$$\begin{aligned}
& \downarrow 4011 \\
& \int (a + b \tan(c + dx))^2 (-Ab - aB + (aA - bB) \tan(c + dx)) dx + \frac{A(a + b \tan(c + dx))^3}{3d} + \\
& \quad \frac{B(a + b \tan(c + dx))^4}{4bd} \\
& \downarrow 3042 \\
& \int (a + b \tan(c + dx))^2 (-Ab - aB + (aA - bB) \tan(c + dx)) dx + \frac{A(a + b \tan(c + dx))^3}{3d} + \\
& \quad \frac{B(a + b \tan(c + dx))^4}{4bd} \\
& \downarrow 4011 \\
& \int (a + b \tan(c + dx)) (-Ba^2 - 2Aba + b^2B + (Aa^2 - 2bBa - Ab^2) \tan(c + dx)) dx + \\
& \quad \frac{(aA - bB)(a + b \tan(c + dx))^2}{2d} + \frac{A(a + b \tan(c + dx))^3}{3d} + \frac{B(a + b \tan(c + dx))^4}{4bd} \\
& \downarrow 3042 \\
& \int (a + b \tan(c + dx)) (-Ba^2 - 2Aba + b^2B + (Aa^2 - 2bBa - Ab^2) \tan(c + dx)) dx + \\
& \quad \frac{(aA - bB)(a + b \tan(c + dx))^2}{2d} + \frac{A(a + b \tan(c + dx))^3}{3d} + \frac{B(a + b \tan(c + dx))^4}{4bd} \\
& \downarrow 4008 \\
& (a^3A - 3a^2bB - 3aAb^2 + b^3B) \int \tan(c + dx) dx + \frac{b(a^2A - 2abB - Ab^2) \tan(c + dx)}{d} - \\
& x(a^3B + 3a^2Ab - 3ab^2B - Ab^3) + \frac{(aA - bB)(a + b \tan(c + dx))^2}{2d} + \frac{A(a + b \tan(c + dx))^3}{3d} + \\
& \quad \frac{B(a + b \tan(c + dx))^4}{4bd} \\
& \downarrow 3042 \\
& (a^3A - 3a^2bB - 3aAb^2 + b^3B) \int \tan(c + dx) dx + \frac{b(a^2A - 2abB - Ab^2) \tan(c + dx)}{d} - \\
& x(a^3B + 3a^2Ab - 3ab^2B - Ab^3) + \frac{(aA - bB)(a + b \tan(c + dx))^2}{2d} + \frac{A(a + b \tan(c + dx))^3}{3d} + \\
& \quad \frac{B(a + b \tan(c + dx))^4}{4bd} \\
& \downarrow 3956
\end{aligned}$$

$$\frac{b(a^2A - 2abB - Ab^2) \tan(c + dx) - (a^3A - 3a^2bB - 3aAb^2 + b^3B) \log(\cos(c + dx))}{d} - x(a^3B + 3a^2Ab - 3ab^2B - Ab^3) + \frac{(aA - bB)(a + b \tan(c + dx))^2}{2d} + \frac{A(a + b \tan(c + dx))^3}{3d} + \frac{B(a + b \tan(c + dx))^4}{4bd}$$

input `Int[Tan[c + d*x]*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `-((3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*x) - ((a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*Log[Cos[c + d*x]])/d + (b*(a^2*A - A*b^2 - 2*a*b*B)*Tan[c + d*x])/d + ((a*A - b*B)*(a + b*Tan[c + d*x])^2)/(2*d) + (A*(a + b*Tan[c + d*x])^3)/(3*d) + (B*(a + b*Tan[c + d*x])^4)/(4*b*d)`

3.249.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

3.249.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.09

method	result
norman	$(-3Aa^2b + Ab^3 - Ba^3 + 3Bab^2)x + \frac{(3Aa^2b - Ab^3 + Ba^3 - 3Bab^2) \tan(dx+c)}{d} + \frac{Bb^3 \tan^4(dx+c)}{4d}$
parts	$\frac{(Ab^3 + 3Bab^2) \left(\frac{\tan^3(dx+c)}{3} - \tan(dx+c) + \arctan(\tan(dx+c)) \right)}{d} + \frac{(3Aab^2 + 3Ba^2b) \left(\frac{\tan^2(dx+c)}{2} - \frac{\ln(1+\tan^2(dx+c))}{2} \right)}{d}$
derivativedivides	$\frac{Bb^3 \tan^4(dx+c)}{4} + \frac{Ab^3 \tan^3(dx+c)}{3} + Bab^2 \tan^3(dx+c) + \frac{3Aab^2 \tan^2(dx+c)}{2} + \frac{3Ba^2b \tan^2(dx+c)}{2} - \frac{Bb^3 \tan^2(dx+c)}{2}$
default	$\frac{Bb^3 \tan^4(dx+c)}{4} + \frac{Ab^3 \tan^3(dx+c)}{3} + Bab^2 \tan^3(dx+c) + \frac{3Aab^2 \tan^2(dx+c)}{2} + \frac{3Ba^2b \tan^2(dx+c)}{2} - \frac{Bb^3 \tan^2(dx+c)}{2}$
parallelrisch	$\frac{3Bb^3 \tan^4(dx+c) + 4Ab^3 \tan^3(dx+c) + 12Bab^2 \tan^3(dx+c) - 36Aa^2bdx + 12Ab^3dx + 18Aab^2 \tan^2(dx+c) - 12Bx}{d}$
risch	$-\frac{6iBa^2bc}{d} - 3iBa^2bx + \frac{2i(9Aa^2b - 12Bab^2 + 3Ba^3 - 4Ab^3 - 18Bab^2e^{6i(dx+c)} + 9Aa^2be^{6i(dx+c)} + 27Aa^2be^{4i(dx+c)})}{d}$

```
input int(tan(d*x+c)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output (-3*A*a^2*b+A*b^3-B*a^3+3*B*a*b^2)*x+(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)/d*tan(d*x+c)+1/4*B*b^3/d*tan(d*x+c)^4+1/2*b*(3*A*a*b+3*B*a^2-B*b^2)/d*tan(d*x+c)^2+1/3*b^2*(A*b+3*B*a)/d*tan(d*x+c)^3+1/2*(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)/d*ln(1+tan(d*x+c)^2)
```

3.249. $\int \tan(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.249.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.08

$$\int \tan(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{3 B b^3 \tan(dx + c)^4 + 4(3 B a b^2 + A b^3) \tan(dx + c)^3 - 12(B a^3 + 3 A a^2 b - 3 B a b^2 - A b^3) dx + 6(3 B a^2 b}{}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/12*(3*B*b^3*tan(d*x + c)^4 + 4*(3*B*a*b^2 + A*b^3)*tan(d*x + c)^3 - 12*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*d*x + 6*(3*B*a^2*b + 3*A*a*b^2 - B*b^3)*tan(d*x + c)^2 - 6*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*log(1/(tan(d*x + c)^2 + 1)) + 12*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*tan(d*x + c))/d`

3.249.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 311 vs. 2(151) = 302.

Time = 0.16 (sec) , antiderivative size = 311, normalized size of antiderivative = 1.88

$$\int \tan(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \frac{A a^3 \log(\tan^2(c+dx)+1)}{2d} - 3 A a^2 b x + \frac{3 A a^2 b \tan(c+dx)}{d} - \frac{3 A a b^2 \log(\tan^2(c+dx)+1)}{2d} + \frac{3 A a b^2 \tan^2(c+dx)}{2d} + A b^3 x + \frac{A b^3 \tan^3(c+dx)}{3d} \\ x(A + B \tan(c))(a + b \tan(c))^3 \tan(c) \end{cases}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `Piecewise((A*a**3*log(tan(c + d*x)**2 + 1)/(2*d) - 3*A*a**2*b*x + 3*A*a**2*b*tan(c + d*x)/d - 3*A*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + 3*A*a*b**2*tan(c + d*x)**2/(2*d) + A*b**3*x + A*b**3*tan(c + d*x)**3/(3*d) - A*b**3*tan(c + d*x)/d - B*a**3*x + B*a**3*tan(c + d*x)/d - 3*B*a**2*b*log(tan(c + d*x)**2 + 1)/(2*d) + 3*B*a**2*b*tan(c + d*x)**2/(2*d) + 3*B*a*b**2*x + B*a*b**2*tan(c + d*x)**3/d - 3*B*a*b**2*tan(c + d*x)/d + B*b**3*log(tan(c + d*x)**2 + 1)/(2*d) + B*b**3*tan(c + d*x)**4/(4*d) - B*b**3*tan(c + d*x)**2/(2*d), Ne(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**3*tan(c), True))`

3.249. $\int \tan(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.249.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.08

$$\int \tan(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{3 B b^3 \tan(dx + c)^4 + 4(3 B a b^2 + A b^3) \tan(dx + c)^3 + 6(3 B a^2 b + 3 A a b^2 - B b^3) \tan(dx + c)^2 - 12(B a^3 + 3 A a^2 b - 3 B a b^2 - A b^3) \tan(dx + c) + 6(A a^3 - 3 B a^2 b - 3 A a b^2 + B b^3) \log(\tan(dx + c)^2 + 1) + 12(B a^3 + 3 A a^2 b - 3 B a b^2 - A b^3) \tan(dx + c)}{d}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/12*(3*B*b^3*tan(d*x + c)^4 + 4*(3*B*a*b^2 + A*b^3)*tan(d*x + c)^3 + 6*(3*B*a^2*b + 3*A*a*b^2 - B*b^3)*tan(d*x + c)^2 - 12*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*(d*x + c) + 6*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*log(tan(d*x + c)^2 + 1) + 12*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*tan(d*x + c))/d`

3.249.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2670 vs. 2(159) = 318.

Time = 2.18 (sec) , antiderivative size = 2670, normalized size of antiderivative = 16.18

$$\int \tan(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

```

-1/12*(12*B*a^3*d*x*tan(d*x)^4*tan(c)^4 + 36*A*a^2*b*d*x*tan(d*x)^4*tan(c)
^4 - 36*B*a*b^2*d*x*tan(d*x)^4*tan(c)^4 - 12*A*b^3*d*x*tan(d*x)^4*tan(c)^4
+ 6*A*a^3*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2
*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^4*tan(c)^4 - 18*B*a^2*b*l
og(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 +
tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^4*tan(c)^4 - 18*A*a*b^2*log(4*(tan(d*
x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 +
tan(c)^2 + 1))*tan(d*x)^4*tan(c)^4 + 6*B*b^3*log(4*(tan(d*x)^2*tan(c)^2 -
2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))
*tan(d*x)^4*tan(c)^4 - 48*B*a^3*d*x*tan(d*x)^3*tan(c)^3 - 144*A*a^2*b*d*x*
tan(d*x)^3*tan(c)^3 + 144*B*a*b^2*d*x*tan(d*x)^3*tan(c)^3 + 48*A*b^3*d*x*t
an(d*x)^3*tan(c)^3 - 18*B*a^2*b*tan(d*x)^4*tan(c)^4 - 18*A*a*b^2*tan(d*x)^
4*tan(c)^4 + 9*B*b^3*tan(d*x)^4*tan(c)^4 - 24*A*a^3*log(4*(tan(d*x)^2*tan(
c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2
+ 1))*tan(d*x)^3*tan(c)^3 + 72*B*a^2*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan
(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d
*x)^3*tan(c)^3 + 72*A*a*b^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c)
+ 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^3*tan(c)
^3 - 24*B*b^3*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)
^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^3*tan(c)^3 + 12*B*a...

```

3.249.9 Mupad [B] (verification not implemented)

Time = 8.24 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.10

$$\begin{aligned}
 & \int \tan(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx \\
 &= x(-B a^3 - 3 A a^2 b + 3 B a b^2 + A b^3) - \frac{\tan(c + dx)^2 \left(\frac{B b^3}{2} - \frac{3 a b (A b + B a)}{2} \right)}{d} \\
 & \quad - \frac{\tan(c + dx) (-B a^3 - 3 A a^2 b + 3 B a b^2 + A b^3)}{d} \\
 & \quad + \frac{\ln(\tan(c + dx)^2 + 1) \left(\frac{A a^3}{2} - \frac{3 B a^2 b}{2} - \frac{3 A a b^2}{2} + \frac{B b^3}{2} \right)}{d} \\
 & \quad + \frac{\tan(c + dx)^3 \left(\frac{A b^3}{3} + B a b^2 \right)}{d} + \frac{B b^3 \tan(c + dx)^4}{4 d}
 \end{aligned}$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`

output $x*(A*b^3 - B*a^3 - 3*A*a^2*b + 3*B*a*b^2) - (\tan(c + d*x)^2*((B*b^3)/2 - (3*a*b*(A*b + B*a))/2))/d - (\tan(c + d*x)*(A*b^3 - B*a^3 - 3*A*a^2*b + 3*B*a*b^2))/d + (\log(\tan(c + d*x)^2 + 1)*((A*a^3)/2 + (B*b^3)/2 - (3*A*a*b^2)/2 - (3*B*a^2*b)/2))/d + (\tan(c + d*x)^3*((A*b^3)/3 + B*a*b^2))/d + (B*b^3*\tan(c + d*x)^4)/(4*d)$

3.250 $\int (a + b \tan(c + dx))^3 (A + B \tan(c + dx)) dx$

3.250.1 Optimal result	2428
3.250.2 Mathematica [C] (verified)	2428
3.250.3 Rubi [A] (verified)	2429
3.250.4 Maple [A] (verified)	2431
3.250.5 Fricas [A] (verification not implemented)	2431
3.250.6 Sympy [A] (verification not implemented)	2432
3.250.7 Maxima [A] (verification not implemented)	2432
3.250.8 Giac [B] (verification not implemented)	2433
3.250.9 Mupad [B] (verification not implemented)	2434

3.250.1 Optimal result

Integrand size = 23, antiderivative size = 140

$$\int (a + b \tan(c + dx))^3 (A + B \tan(c + dx)) dx$$

$$= (a^3 A - 3aAb^2 - 3a^2bB + b^3B) x - \frac{(3a^2Ab - Ab^3 + a^3B - 3ab^2B) \log(\cos(c + dx))}{d}$$

$$+ \frac{b(2aAb + a^2B - b^2B) \tan(c + dx)}{d}$$

$$+ \frac{(Ab + aB)(a + b \tan(c + dx))^2}{2d} + \frac{B(a + b \tan(c + dx))^3}{3d}$$

```
output (A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*x-(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*ln(cos(d*x+c))/d+b*(2*A*a*b+B*a^2-B*b^2)*tan(d*x+c)/d+1/2*(A*b+B*a)*(a+b*tan(d*x+c))^2/d+1/3*B*(a+b*tan(d*x+c))^3/d
```

3.250.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.04 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.93

$$\int (a + b \tan(c + dx))^3 (A + B \tan(c + dx)) dx$$

$$= \frac{3(a + ib)^3(-iA + B) \log(i - \tan(c + dx)) + 3(a - ib)^3(iA + B) \log(i + \tan(c + dx)) + 6b(3aAb + 3a^2B \cdot \dots}{6d}$$

input `Integrate[(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output $(3*(a + I*b)^3*((-I)*A + B)*\text{Log}[I - \text{Tan}[c + d*x]] + 3*(a - I*b)^3*(I*A + B)*\text{Log}[I + \text{Tan}[c + d*x]] + 6*b*(3*a*A*b + 3*a^2*B - b^2*B)*\text{Tan}[c + d*x] + 3*b^2*(A*b + 3*a*B)*\text{Tan}[c + d*x]^2 + 2*b^3*B*\text{Tan}[c + d*x]^3)/(6*d)$

3.250.3 Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.348$, Rules used = {3042, 4011, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \tan(c + dx))^3 (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + b \tan(c + dx))^3 (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4011} \\ & \int (a + b \tan(c + dx))^2 (aA - bB + (Ab + aB) \tan(c + dx)) dx + \frac{B(a + b \tan(c + dx))^3}{3d} \\ & \quad \downarrow \text{3042} \\ & \int (a + b \tan(c + dx))^2 (aA - bB + (Ab + aB) \tan(c + dx)) dx + \frac{B(a + b \tan(c + dx))^3}{3d} \\ & \quad \downarrow \text{4011} \\ & \int (a + b \tan(c + dx)) (Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx + \\ & \quad \frac{(aB + Ab)(a + b \tan(c + dx))^2}{2d} + \frac{B(a + b \tan(c + dx))^3}{3d} \\ & \quad \downarrow \text{3042} \\ & \int (a + b \tan(c + dx)) (Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx + \\ & \quad \frac{(aB + Ab)(a + b \tan(c + dx))^2}{2d} + \frac{B(a + b \tan(c + dx))^3}{3d} \\ & \quad \downarrow \text{4008} \end{aligned}$$

3.250. $\int (a + b \tan(c + dx))^3 (A + B \tan(c + dx)) dx$

$$\begin{aligned}
& (a^3B + 3a^2Ab - 3ab^2B - Ab^3) \int \tan(c + dx)dx + \frac{b(a^2B + 2aAb - b^2B) \tan(c + dx)}{d} + \\
& x(a^3A - 3a^2bB - 3aAb^2 + b^3B) + \frac{(aB + Ab)(a + b \tan(c + dx))^2}{2d} + \frac{B(a + b \tan(c + dx))^3}{3d} \\
& \quad \downarrow \text{3042} \\
& (a^3B + 3a^2Ab - 3ab^2B - Ab^3) \int \tan(c + dx)dx + \frac{b(a^2B + 2aAb - b^2B) \tan(c + dx)}{d} + \\
& x(a^3A - 3a^2bB - 3aAb^2 + b^3B) + \frac{(aB + Ab)(a + b \tan(c + dx))^2}{2d} + \frac{B(a + b \tan(c + dx))^3}{3d} \\
& \quad \downarrow \text{3956} \\
& \frac{b(a^2B + 2aAb - b^2B) \tan(c + dx)}{d} - \frac{(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \log(\cos(c + dx))}{d} + \\
& x(a^3A - 3a^2bB - 3aAb^2 + b^3B) + \frac{(aB + Ab)(a + b \tan(c + dx))^2}{2d} + \frac{B(a + b \tan(c + dx))^3}{3d}
\end{aligned}$$

input `Int[(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*x - ((3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*Log[Cos[c + d*x]])/d + (b*(2*a*A*b + a^2*B - b^2*B)*Tan[c + d*x])/d + ((A*b + a*B)*(a + b*Tan[c + d*x])^2)/(2*d) + (B*(a + b*Tan[c + d*x])^3)/(3*d)`

3.250.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)]*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

```
rule 4011 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

3.250.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.01

method	result
norman	$(Aa^3 - 3Aab^2 - 3Ba^2b + Bb^3)x + \frac{b(3Aab+3Ba^2-Bb^2)\tan(dx+c)}{d} + \frac{Bb^3(\tan^3(dx+c))}{3d} + \frac{b^2(Ab-3Ba^2)}{3d}$
parts	$Aa^3x + \frac{(Ab^3+3Ba^2b^2)\left(\frac{\tan^2(dx+c)}{2} - \frac{\ln(1+\tan^2(dx+c))}{2}\right)}{d} + \frac{(3Aab^2+3Ba^2b)(\tan(dx+c)-\arctan(\tan(dx+c)))}{d}$
derivativedivides	$\frac{Bb^3(\tan^3(dx+c))}{3} + \frac{Ab^3(\tan^2(dx+c))}{2} + \frac{3Ba^2b^2(\tan^2(dx+c))}{2} + 3Aab^2\tan(dx+c) + 3Ba^2b\tan(dx+c) - Bb^3\tan(dx+c) + \frac{3b^2(Ab-3Ba^2)}{3d}$
default	$\frac{Bb^3(\tan^3(dx+c))}{3} + \frac{Ab^3(\tan^2(dx+c))}{2} + \frac{3Ba^2b^2(\tan^2(dx+c))}{2} + 3Aab^2\tan(dx+c) + 3Ba^2b\tan(dx+c) - Bb^3\tan(dx+c) + \frac{3b^2(Ab-3Ba^2)}{3d}$
parallelrisch	$\frac{2Bb^3(\tan^3(dx+c)) + 6Aa^3d - 18Aab^2dx + 3Ab^3(\tan^2(dx+c)) - 18Ba^2bdx + 6Bb^3dx + 9Ba^2b^2(\tan^2(dx+c)) + 9A\ln(1+\tan^2(dx+c))}{d}$
risch	$Aa^3x - 3Aab^2x - 3Ba^2bx + Bb^3x + 3iAa^2bx + iBa^3x - 3iBa^2bx + \frac{6iAa^2bc}{d} - \frac{6iBa^2b^2c}{d}$

```
input int((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output (A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*x+b*(3*A*a*b+3*B*a^2-B*b^2)/d*tan(d*x+c)
+1/3*B*b^3/d*tan(d*x+c)^3+1/2*b^2*(A*b+3*B*a)/d*tan(d*x+c)^2+1/2*(3*A*a^2*
b-A*b^3+B*a^3-3*B*a*b^2)/d*ln(1+tan(d*x+c)^2)
```

3.250.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.01

$$\int (a + b \tan(c + dx))^3 (A + B \tan(c + dx)) dx$$

$$= \frac{2Bb^3 \tan(dx+c)^3 + 6(Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3)dx + 3(3Bab^2 + Ab^3) \tan(dx+c)^2 - 3(Ba^3 + 3Bab^2)}{6d}$$

input `integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output $\frac{1}{6}*(2*B*b^3*\tan(d*x + c)^3 + 6*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*d*x + 3*(3*B*a*b^2 + A*b^3)*\tan(d*x + c)^2 - 3*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*\log(1/(\tan(d*x + c)^2 + 1)) + 6*(3*B*a^2*b + 3*A*a*b^2 - B*b^3)*\tan(d*x + c))/d$

3.250.6 Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.71

$$\int (a + b \tan(c + dx))^3 (A + B \tan(c + dx)) dx$$

$$= \begin{cases} Aa^3x + \frac{3Aa^2b \log(\tan^2(c+dx)+1)}{2d} - 3Aab^2x + \frac{3Aab^2 \tan(c+dx)}{d} - \frac{Ab^3 \log(\tan^2(c+dx)+1)}{2d} + \frac{Ab^3 \tan^2(c+dx)}{2d} + \frac{Ba^3 \log(\tan^2(c+dx)+1)}{2d} \\ x(A + B \tan(c)) (a + b \tan(c))^3 \end{cases}$$

input `integrate((a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `Piecewise((A*a**3*x + 3*A*a**2*b*log(tan(c + d*x)**2 + 1)/(2*d) - 3*A*a*b**2*x + 3*A*a*b**2*tan(c + d*x)/d - A*b**3*log(tan(c + d*x)**2 + 1)/(2*d) + A*b**3*tan(c + d*x)**2/(2*d) + B*a**3*log(tan(c + d*x)**2 + 1)/(2*d) - 3*B*a**2*b*x + 3*B*a**2*b*tan(c + d*x)/d - 3*B*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + 3*B*a*b**2*tan(c + d*x)**2/(2*d) + B*b**3*x + B*b**3*tan(c + d*x)**3/(3*d) - B*b**3*tan(c + d*x)/d, Ne(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**3, True))`

3.250.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.02

$$\int (a + b \tan(c + dx))^3 (A + B \tan(c + dx)) dx$$

$$= \frac{2 B b^3 \tan(dx + c)^3 + 3 (3 B a b^2 + A b^3) \tan(dx + c)^2 + 6 (A a^3 - 3 B a^2 b - 3 A a b^2 + B b^3) (dx + c) + 3 (B a^3 + 3 A a^2 b - 3 A a b^2 + B b^3) \tan(dx + c) + 3 (A a^3 + 3 A a^2 b - 3 A a b^2 + B b^3) \log(\tan^2(dx + c) + 1) + 3 (3 A a b^2 + A b^3) \tan(dx + c) + 3 (3 A a b^2 + A b^3) \log(\tan^2(dx + c) + 1) + 3 (3 A a b^2 + A b^3) \tan(dx + c) + 3 (3 A a b^2 + A b^3) \log(\tan^2(dx + c) + 1)}{6 d}$$

input `integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

3.250. $\int (a + b \tan(c + dx))^3 (A + B \tan(c + dx)) dx$

output $1/6*(2*B*b^3*\tan(dx + c)^3 + 3*(3*B*a*b^2 + A*b^3)*\tan(dx + c)^2 + 6*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*(dx + c) + 3*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*\log(\tan(dx + c)^2 + 1) + 6*(3*B*a^2*b + 3*A*a*b^2 - B*b^3)*\tan(dx + c))/d$

3.250.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1751 vs. $2(136) = 272$.

Time = 1.40 (sec) , antiderivative size = 1751, normalized size of antiderivative = 12.51

$$\int (a + b \tan(c + dx))^3 (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output $1/6*(6*A*a^3*d*x*\tan(dx)^3*\tan(c)^3 - 18*B*a^2*b*d*x*\tan(dx)^3*\tan(c)^3 - 18*A*a*b^2*d*x*\tan(dx)^3*\tan(c)^3 + 6*B*b^3*d*x*\tan(dx)^3*\tan(c)^3 - 3*B*a^3*\log(4*(\tan(dx)^2*\tan(c)^2 - 2*\tan(dx)*\tan(c) + 1)/(\tan(dx)^2*\tan(c)^2 + \tan(dx)^2 + \tan(c)^2 + 1))*\tan(dx)^3*\tan(c)^3 - 9*A*a^2*b*\log(4*(\tan(dx)^2*\tan(c)^2 - 2*\tan(dx)*\tan(c) + 1)/(\tan(dx)^2*\tan(c)^2 + \tan(dx)^2 + \tan(c)^2 + 1))*\tan(dx)^3*\tan(c)^3 + 9*B*a*b^2*\log(4*(\tan(dx)^2*\tan(c)^2 - 2*\tan(dx)*\tan(c) + 1)/(\tan(dx)^2*\tan(c)^2 + \tan(dx)^2 + \tan(c)^2 + 1))*\tan(dx)^3*\tan(c)^3 + 3*A*b^3*\log(4*(\tan(dx)^2*\tan(c)^2 - 2*\tan(dx)*\tan(c) + 1)/(\tan(dx)^2*\tan(c)^2 + \tan(dx)^2 + \tan(c)^2 + 1))*\tan(dx)^3*\tan(c)^3 - 18*A*a^3*d*x*\tan(dx)^2*\tan(c)^2 + 54*B*a^2*b*d*x*\tan(dx)^2*\tan(c)^2 + 54*A*a*b^2*d*x*\tan(dx)^2*\tan(c)^2 - 18*B*b^3*d*x*\tan(dx)^2*\tan(c)^2 + 9*B*a*b^2*\tan(dx)^3*\tan(c)^3 + 3*A*b^3*\tan(dx)^3*\tan(c)^3 + 9*B*a^3*\log(4*(\tan(dx)^2*\tan(c)^2 - 2*\tan(dx)*\tan(c) + 1)/(\tan(dx)^2*\tan(c)^2 + \tan(dx)^2 + \tan(c)^2 + 1))*\tan(dx)^2*\tan(c)^2 + 27*A*a^2*b*\log(4*(\tan(dx)^2*\tan(c)^2 - 2*\tan(dx)*\tan(c) + 1)/(\tan(dx)^2*\tan(c)^2 + \tan(dx)^2 + \tan(c)^2 + 1))*\tan(dx)^2*\tan(c)^2 - 27*B*a*b^2*\log(4*(\tan(dx)^2*\tan(c)^2 - 2*\tan(dx)*\tan(c) + 1)/(\tan(dx)^2*\tan(c)^2 + \tan(dx)^2 + \tan(c)^2 + 1))*\tan(dx)^2*\tan(c)^2 - 9*A*b^3*\log(4*(\tan(dx)^2*\tan(c)^2 - 2*\tan(dx)*\tan(c) + 1)/(\tan(dx)^2*\tan(c)^2 + \tan(dx)^2 + \tan(c)^2 + 1))*\tan(dx)^2*\tan(c)^2 - 18*B*a^2*b*\tan(dx)^3*\tan(c)^2 - 18*A*a*b^2*\tan(dx)...$

3.250.9 Mupad [B] (verification not implemented)

Time = 8.27 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.01

$$\int (a + b \tan(c + dx))^3 (A + B \tan(c + dx)) dx$$

$$= x (A a^3 - 3 B a^2 b - 3 A a b^2 + B b^3)$$

$$- \frac{\ln(\tan(c + dx)^2 + 1) \left(-\frac{B a^3}{2} - \frac{3 A a^2 b}{2} + \frac{3 B a b^2}{2} + \frac{A b^3}{2} \right)}{d}$$

$$+ \frac{\tan(c + dx)^2 \left(\frac{A b^3}{2} + \frac{3 B a b^2}{2} \right)}{d}$$

$$- \frac{\tan(c + dx) (B b^3 - 3 a b (A b + B a))}{d} + \frac{B b^3 \tan(c + dx)^3}{3 d}$$

input `int((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`output `x*(A*a^3 + B*b^3 - 3*A*a*b^2 - 3*B*a^2*b) - (log(tan(c + d*x)^2 + 1)*((A*b^3)/2 - (B*a^3)/2 - (3*A*a^2*b)/2 + (3*B*a*b^2)/2))/d + (tan(c + d*x)^2*((A*b^3)/2 + (3*B*a*b^2)/2))/d - (tan(c + d*x)*(B*b^3 - 3*a*b*(A*b + B*a)))/d + (B*b^3*tan(c + d*x)^3)/(3*d)`

3.251 $\int \cot(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.251.1 Optimal result	2435
3.251.2 Mathematica [C] (verified)	2435
3.251.3 Rubi [A] (verified)	2436
3.251.4 Maple [A] (verified)	2439
3.251.5 Fricas [A] (verification not implemented)	2440
3.251.6 Sympy [A] (verification not implemented)	2440
3.251.7 Maxima [A] (verification not implemented)	2441
3.251.8 Giac [A] (verification not implemented)	2441
3.251.9 Mupad [B] (verification not implemented)	2442

3.251.1 Optimal result

Integrand size = 29, antiderivative size = 117

$$\int \cot(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= (3a^2Ab - Ab^3 + a^3B - 3ab^2B) x - \frac{b(3aAb + 3a^2B - b^2B) \log(\cos(c+dx))}{d}$$

$$+ \frac{a^3A \log(\sin(c+dx))}{d} + \frac{b^2(Ab + 2aB) \tan(c+dx)}{d} + \frac{bB(a+b \tan(c+dx))^2}{2d}$$

```
output (3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*x-b*(3*A*a*b+3*B*a^2-B*b^2)*ln(cos(d*x+c))
/d+a^3*A*ln(sin(d*x+c))/d+b^2*(A*b+2*B*a)*tan(d*x+c)/d+1/2*b*B*(a+b*tan(d*x+c))^2/d
```

3.251.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.63 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.98

$$\int \cot(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{-(a+ib)^3(A+iB) \log(i-\tan(c+dx)) + 2a^3A \log(\tan(c+dx)) - (a-ib)^3(A-iB) \log(i+\tan(c+dx))}{2d}$$

input `Integrate[Cot[c + d*x]*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(-((a + I*b)^3*(A + I*B)*Log[I - Tan[c + d*x]]) + 2*a^3*A*Log[Tan[c + d*x]] - (a - I*b)^3*(A - I*B)*Log[I + Tan[c + d*x]] + 2*b^2*(A*b + 2*a*B)*Tan[c + d*x] + b*B*(a + b*Tan[c + d*x])^2)/(2*d)`

3.251.3 Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.02, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.379$, Rules used = {3042, 4090, 27, 3042, 4120, 25, 3042, 4107, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))^3(A + B \tan(c + dx))}{\tan(c + dx)} dx \\
 & \quad \downarrow \text{4090} \\
 & \frac{1}{2} \int 2 \cot(c + dx)(a + b \tan(c + dx)) (Aa^2 + b(Ab + 2aB) \tan^2(c + dx) + (Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx + \\
 & \quad \frac{bB(a + b \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{27} \\
 & \int \cot(c + dx)(a + b \tan(c + dx)) (Aa^2 + b(Ab + 2aB) \tan^2(c + dx) + (Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx + \\
 & \quad \frac{bB(a + b \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx)) (Aa^2 + b(Ab + 2aB) \tan(c + dx))^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx)}{\tan(c + dx) \frac{bB(a + b \tan(c + dx))^2}{2d}} dx + \\
 & \quad \downarrow \text{4120}
 \end{aligned}$$

3.251. $\int \cot(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& - \int -\cot(c + dx) (Aa^3 + b(3Ba^2 + 3Aba - b^2B) \tan^2(c + dx) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)) dx + \\
& \quad \frac{b^2(2aB + Ab) \tan(c + dx)}{d} + \frac{bB(a + b \tan(c + dx))^2}{2d} \\
& \quad \downarrow 25 \\
& \int \cot(c + dx) (Aa^3 + b(3Ba^2 + 3Aba - b^2B) \tan^2(c + dx) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)) dx + \\
& \quad \frac{b^2(2aB + Ab) \tan(c + dx)}{d} + \frac{bB(a + b \tan(c + dx))^2}{2d} \\
& \quad \downarrow 3042 \\
& \int \frac{Aa^3 + b(3Ba^2 + 3Aba - b^2B) \tan(c + dx)^2 + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\tan(c + dx)} dx + \\
& \quad \frac{b^2(2aB + Ab) \tan(c + dx)}{d} + \frac{bB(a + b \tan(c + dx))^2}{2d} \\
& \quad \downarrow 4107 \\
& a^3 A \int \cot(c + dx) dx + b(3a^2 B + 3aAb - b^2 B) \int \tan(c + dx) dx + \\
& \quad x(a^3 B + 3a^2 Ab - 3ab^2 B - Ab^3) + \frac{b^2(2aB + Ab) \tan(c + dx)}{d} + \frac{bB(a + b \tan(c + dx))^2}{2d} \\
& \quad \downarrow 3042 \\
& a^3 A \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + b(3a^2 B + 3aAb - b^2 B) \int \tan(c + dx) dx + \\
& \quad x(a^3 B + 3a^2 Ab - 3ab^2 B - Ab^3) + \frac{b^2(2aB + Ab) \tan(c + dx)}{d} + \frac{bB(a + b \tan(c + dx))^2}{2d} \\
& \quad \downarrow 25 \\
& a^3(-A) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + b(3a^2 B + 3aAb - b^2 B) \int \tan(c + dx) dx + \\
& \quad x(a^3 B + 3a^2 Ab - 3ab^2 B - Ab^3) + \frac{b^2(2aB + Ab) \tan(c + dx)}{d} + \frac{bB(a + b \tan(c + dx))^2}{2d} \\
& \quad \downarrow 3956 \\
& \frac{a^3 A \log(-\sin(c + dx))}{d} - \frac{b(3a^2 B + 3aAb - b^2 B) \log(\cos(c + dx))}{d} + \\
& \quad x(a^3 B + 3a^2 Ab - 3ab^2 B - Ab^3) + \frac{b^2(2aB + Ab) \tan(c + dx)}{d} + \frac{bB(a + b \tan(c + dx))^2}{2d}
\end{aligned}$$

input `Int[Cot[c + d*x]*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

```
output (3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*x - (b*(3*a*A*b + 3*a^2*B - b^2*B)
*Log[Cos[c + d*x]])/d + (a^3*A*Log[-Sin[c + d*x]])/d + (b^2*(A*b + 2*a*B)*
Tan[c + d*x])/d + (b*B*(a + b*Tan[c + d*x])^2)/(2*d)
```

3.251.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4090 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_.)]*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4107 Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_.)] + (C_.)*tan[(e_.) + (f_.)*(x_.)]^2
)/tan[(e_.) + (f_.)*(x_.)], x_Symbol] := Simp[B*x, x] + (Simp[A Int[1/Tan[
e + f*x], x], x] + Simp[C Int[Tan[e + f*x], x], x]) /; FreeQ[{e, f, A, B,
C}, x] && NeQ[A, C]
```

```
rule 4120 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f_)*
*(x_)])^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

3.251.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.03

method	result
parallelrisch	$\frac{(-A a^3 + 3A a b^2 + 3B a^2 b - B b^3) \ln(\sec^2(dx+c)) + 2A a^3 \ln(\tan(dx+c)) + B b^3 (\tan^2(dx+c)) + (2A b^3 + 6B a b^2) \tan(dx+c)}{2d}$
norman	$(3A a^2 b - A b^3 + B a^3 - 3B a b^2) x + \frac{b^2 (A b + 3B a) \tan(dx+c)}{d} + \frac{B b^3 (\tan^2(dx+c))}{2d} + \frac{A a^3 \ln(\tan(dx+c))}{d}$
derivativedivides	$\frac{\frac{B b^3 (\tan^2(dx+c))}{2} + A b^3 \tan(dx+c) + 3B a b^2 \tan(dx+c) + \frac{(-A a^3 + 3A a b^2 + 3B a^2 b - B b^3) \ln(1 + \tan^2(dx+c))}{2}}{d} + (3A a^2 b - A b^3)$
default	$\frac{\frac{B b^3 (\tan^2(dx+c))}{2} + A b^3 \tan(dx+c) + 3B a b^2 \tan(dx+c) + \frac{(-A a^3 + 3A a b^2 + 3B a^2 b - B b^3) \ln(1 + \tan^2(dx+c))}{2}}{d} + (3A a^2 b - A b^3)$
risch	$\frac{6i B a^2 b c}{d} + 3i B a^2 b x + \frac{2i b^2 (-i B b e^{2i(dx+c)} + A b e^{2i(dx+c)} + 3B a e^{2i(dx+c)} + A b + 3B a)}{d(e^{2i(dx+c)} + 1)^2} - \frac{2i B b^3 c}{d} + 3A a^2 b x$

```
input int(cot(d*x+c)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
)
```

```
output 1/2*((-A*a^3+3*A*a*b^2+3*B*a^2*b-B*b^3)*ln(sec(d*x+c)^2)+2*A*a^3*ln(tan(d*
x+c))+B*b^3*tan(d*x+c)^2+(2*A*b^3+6*B*a*b^2)*tan(d*x+c)+6*(A*a^2*b-1/3*A*b
^3+1/3*B*a^3-B*a*b^2)*d*x)/d
```

3.251. $\int \cot(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.251.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.14

$$\int \cot(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{Bb^3 \tan(dx + c)^2 + Aa^3 \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) + 2(Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3)dx - (3Ba^2b + 3Aab^2 - Bb^3)}{2d}$$

```
input integrate(cot(d*x+c)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output 1/2*(B*b^3*tan(d*x + c)^2 + A*a^3*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))
+ 2*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*d*x - (3*B*a^2*b + 3*A*a*b^2
- B*b^3)*log(1/(tan(d*x + c)^2 + 1)) + 2*(3*B*a*b^2 + A*b^3)*tan(d*x + c))
/d
```

3.251.6 Sympy [A] (verification not implemented)

Time = 0.46 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.74

$$\int \cot(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \begin{cases} -\frac{Aa^3 \log(\tan^2(c+dx)+1)}{2d} + \frac{Aa^3 \log(\tan(c+dx))}{d} + 3Aa^2bx + \frac{3Aab^2 \log(\tan^2(c+dx)+1)}{2d} - Ab^3x + \frac{Ab^3 \tan(c+dx)}{d} + Ba^3 \\ x(A + B \tan(c))(a + b \tan(c))^3 \cot(c) \end{cases}$$

```
input integrate(cot(d*x+c)*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)
```

```
output Piecewise((-A*a**3*log(tan(c + d*x)**2 + 1)/(2*d) + A*a**3*log(tan(c + d*x
))/d + 3*A*a**2*b*x + 3*A*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) - A*b**3*x
+ A*b**3*tan(c + d*x)/d + B*a**3*x + 3*B*a**2*b*log(tan(c + d*x)**2 + 1)/
(2*d) - 3*B*a*b**2*x + 3*B*a*b**2*tan(c + d*x)/d - B*b**3*log(tan(c + d*x)
**2 + 1)/(2*d) + B*b**3*tan(c + d*x)**2/(2*d), Ne(d, 0)), (x*(A + B*tan(c)
)*(a + b*tan(c))**3*cot(c), True))
```

3.251.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.06

$$\int \cot(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{Bb^3 \tan(dx + c)^2 + 2Aa^3 \log(\tan(dx + c)) + 2(Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3)(dx + c) - (Aa^3 - 3Ba^2b + 3Aab^2 - Bb^3)}{2d}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(B*b^3*tan(d*x + c)^2 + 2*A*a^3*log(tan(d*x + c)) + 2*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*(d*x + c) - (A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*log(tan(d*x + c)^2 + 1) + 2*(3*B*a*b^2 + A*b^3)*tan(d*x + c))/d`

3.251.8 Giac [A] (verification not implemented)

Time = 1.13 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.10

$$\int \cot(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{Bb^3 \tan(dx + c)^2 + 2Aa^3 \log(|\tan(dx + c)|) + 6Bab^2 \tan(dx + c) + 2Ab^3 \tan(dx + c) + 2(Ba^3 + 3Aa^2b - 3Aab^2 - Bb^3)(dx + c) - (Aa^3 - 3Ba^2b + 3Aab^2 - Bb^3)}{2d}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `1/2*(B*b^3*tan(d*x + c)^2 + 2*A*a^3*log(abs(tan(d*x + c))) + 6*B*a*b^2*tan(d*x + c) + 2*A*b^3*tan(d*x + c) + 2*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*(d*x + c) - (A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*log(tan(d*x + c)^2 + 1))/d`

3.251.9 Mupad [B] (verification not implemented)

Time = 8.48 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.01

$$\int \cot(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{\tan(c + dx)(Ab^3 + 3Bab^2)}{d} + \frac{Aa^3 \ln(\tan(c + dx))}{d}$$

$$+ \frac{Bb^3 \tan(c + dx)^2}{2d} - \frac{\ln(\tan(c + dx) + 1i)(A - B1i)(b + a1i)^3 1i}{2d}$$

$$- \frac{\ln(\tan(c + dx) - 1i)(A + B1i)(-b + a1i)^3 1i}{2d}$$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`output `(tan(c + d*x)*(A*b^3 + 3*B*a*b^2))/d + (A*a^3*log(tan(c + d*x)))/d - (log(tan(c + d*x) + 1i)*(A - B*1i)*(a*1i + b)^3*1i)/(2*d) - (log(tan(c + d*x) - 1i)*(A + B*1i)*(a*1i - b)^3*1i)/(2*d) + (B*b^3*tan(c + d*x)^2)/(2*d)`

3.252 $\int \cot^2(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.252.1 Optimal result	2443
3.252.2 Mathematica [C] (verified)	2443
3.252.3 Rubi [A] (verified)	2444
3.252.4 Maple [A] (verified)	2447
3.252.5 Fricas [A] (verification not implemented)	2447
3.252.6 Sympy [A] (verification not implemented)	2448
3.252.7 Maxima [A] (verification not implemented)	2448
3.252.8 Giac [A] (verification not implemented)	2449
3.252.9 Mupad [B] (verification not implemented)	2449

3.252.1 Optimal result

Integrand size = 31, antiderivative size = 119

$$\int \cot^2(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= -((a^3A - 3aAb^2 - 3a^2bB + b^3B)x) - \frac{b^2(Ab + 3aB) \log(\cos(c+dx))}{d} + \frac{a^2(3Ab + aB) \log(\sin(c+dx))}{d} + \frac{b^2(aA + bB) \tan(c+dx)}{d} - \frac{aA \cot(c+dx)(a + b \tan(c+dx))^2}{d}$$

output $-(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*x-b^2*(A*b+3*B*a)*\ln(\cos(d*x+c))/d+a^2*(3*A*b+B*a)*\ln(\sin(d*x+c))/d+b^2*(A*a+B*b)*\tan(d*x+c)/d-a*A*\cot(d*x+c)*(a+b*\tan(d*x+c))^2/d$

3.252.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.59 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.95

$$\int \cot^2(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{-2a^3A \cot(c+dx) + i(a+ib)^3(A+ib) \log(i - \tan(c+dx)) + 2a^2(3Ab + aB) \log(\tan(c+dx)) + (ia + b^2 \tan^2(c+dx))}{2d}$$

3.252. $\int \cot^2(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

input `Integrate[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output $(-2*a^3*A*Cot[c + d*x] + I*(a + I*b)^3*(A + I*B)*Log[I - Tan[c + d*x]] + 2*a^2*(3*A*b + a*B)*Log[Tan[c + d*x]] + (I*a + b)^3*(A - I*B)*Log[I + Tan[c + d*x]] + 2*b^3*B*Tan[c + d*x])/(2*d)$

3.252.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.02, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 4088, 3042, 4120, 25, 3042, 4107, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^2(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))^3(A + B \tan(c + dx))}{\tan(c + dx)^2} dx \\
 & \quad \downarrow \text{4088} \\
 & \int \cot(c + dx)(a + b \tan(c + dx)) (b(aA + bB) \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(3Ab + aB)) dx - \\
 & \quad \frac{aA \cot(c + dx)(a + b \tan(c + dx))^2}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx)) (b(aA + bB) \tan(c + dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(3Ab + aB))}{\tan(c + dx) \frac{aA \cot(c + dx)(a + b \tan(c + dx))^2}{d}} dx - \\
 & \quad \downarrow \text{4120} \\
 & - \int -\cot(c + dx) ((3Ab + aB)a^2 + b^2(Ab + 3aB) \tan^2(c + dx) - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)) dx + \\
 & \quad \frac{b^2(aA + bB) \tan(c + dx)}{d} - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^2}{d}
 \end{aligned}$$

3.252. $\int \cot^2(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \int \cot(c + dx) \left((3Ab + aB)a^2 + b^2(Ab + 3aB) \tan^2(c + dx) - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx) \right) dx + \\
& \frac{b^2(aA + bB) \tan(c + dx)}{d} - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^2}{d} \\
& \quad \downarrow \text{25} \\
& \int \frac{(3Ab + aB)a^2 + b^2(Ab + 3aB) \tan^2(c + dx) - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\tan(c + dx)} dx + \\
& \frac{b^2(aA + bB) \tan(c + dx)}{d} - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^2}{d} \\
& \quad \downarrow \text{3042} \\
& a^2(aB + 3Ab) \int \cot(c + dx) dx + b^2(3aB + Ab) \int \tan(c + dx) dx - \\
& x(a^3A - 3a^2bB - 3aAb^2 + b^3B) + \frac{b^2(aA + bB) \tan(c + dx)}{d} - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^2}{d} \\
& \quad \downarrow \text{4107} \\
& a^2(aB + 3Ab) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + b^2(3aB + Ab) \int \tan(c + dx) dx - \\
& x(a^3A - 3a^2bB - 3aAb^2 + b^3B) + \frac{b^2(aA + bB) \tan(c + dx)}{d} - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^2}{d} \\
& \quad \downarrow \text{3042} \\
& -\left(a^2(aB + 3Ab) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx\right) + b^2(3aB + Ab) \int \tan(c + dx) dx - \\
& x(a^3A - 3a^2bB - 3aAb^2 + b^3B) + \frac{b^2(aA + bB) \tan(c + dx)}{d} - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^2}{d} \\
& \quad \downarrow \text{25} \\
& \frac{a^2(aB + 3Ab) \log(-\sin(c + dx))}{d} - x(a^3A - 3a^2bB - 3aAb^2 + b^3B) + \frac{b^2(aA + bB) \tan(c + dx)}{d} - \\
& \frac{b^2(3aB + Ab) \log(\cos(c + dx))}{d} - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^2}{d} \\
& \quad \downarrow \text{3956}
\end{aligned}$$

input `Int[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `-((a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*x) - (b^2*(A*b + 3*a*B)*Log[Cos[c + d*x]])/d + (a^2*(3*A*b + a*B)*Log[-Sin[c + d*x]])/d + (b^2*(a*A + b*B)*Tan[c + d*x])/d - (a*A*Cot[c + d*x]*(a + b*Tan[c + d*x])^2)/d`

3.252. $\int \cot^2(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.252.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`
- rule 4107 `Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[B*x, x] + (Simp[A Int[1/Tan[e + f*x], x], x] + Simp[C Int[Tan[e + f*x], x], x]) /; FreeQ[{e, f, A, B, C}, x] && NeQ[A, C]`
- rule 4120 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]`

3.252.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.99

method	result
parallelrisch	$\frac{(-3Aa^2b+Ab^3-Ba^3+3Bab^2)\ln(\sec^2(dx+c))+(6Aa^2b+2Ba^3)\ln(\tan(dx+c))-2A\cot(dx+c)a^3+2Bb^3\tan(dx+c)-2Aa^3+3Aab^2+3Ba^2b-Bb^3}{2d}$
derivativedivides	$\frac{Bb^3\tan(dx+c)+\frac{(-3Aa^2b+Ab^3-Ba^3+3Bab^2)\ln(1+\tan^2(dx+c))}{2}+(-Aa^3+3Aab^2+3Ba^2b-Bb^3)\arctan(\tan(dx+c))}{d}$
default	$\frac{Bb^3\tan(dx+c)+\frac{(-3Aa^2b+Ab^3-Ba^3+3Bab^2)\ln(1+\tan^2(dx+c))}{2}+(-Aa^3+3Aab^2+3Ba^2b-Bb^3)\arctan(\tan(dx+c))}{d}$
norman	$\frac{(-Aa^3+3Aab^2+3Ba^2b-Bb^3)x\tan(dx+c)+\frac{Bb^3(\tan^2(dx+c))}{d}-\frac{Aa^3}{d}}{\tan(dx+c)}+\frac{a^2(3Ab+Ba)\ln(\tan(dx+c))}{d}-\frac{(3Aa^2b-Aa^3)}{d}$
risch	$-Aa^3x+3Aab^2x+3Ba^2bx-Bb^3x-3iAa^2bx-iBa^3x+3iBab^2x+iAb^3x-\frac{2ia^3}{d}$

input `int(cot(d*x+c)^2*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/2*((-3*A*a^2*b+A*b^3-B*a^3+3*B*a*b^2)*ln(sec(d*x+c)^2)+(6*A*a^2*b+2*B*a^3)*ln(tan(d*x+c))-2*A*cot(d*x+c)*a^3+2*B*b^3*tan(d*x+c)-2*d*x*(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3))/d`

3.252.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.22

$$\int \cot^2(c+dx)(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx$$

$$= \frac{2Bb^3\tan(dx+c)^2-2Aa^3-2(Aa^3-3Ba^2b-3Aab^2+Bb^3)dx\tan(dx+c)+(Ba^3+3Aa^2b)\log\left(\frac{\tan(dx+c)}{\tan(dx+c)+1}\right)}{2d\tan(dx+c)}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output `1/2*(2*B*b^3*tan(d*x+c)^2-2*A*a^3-2*(A*a^3-3*B*a^2*b-3*A*a*b^2+B*b^3)*d*x*tan(d*x+c)+(B*a^3+3*A*a^2*b)*log(tan(d*x+c)^2/(tan(d*x+c)+1))*tan(d*x+c)-(3*B*a*b^2+A*b^3)*log(1/(tan(d*x+c)^2+1))*tan(d*x+c))/(d*tan(d*x+c))`

3.252. $\int \cot^2(c+dx)(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx$

3.252.6 Sympy [A] (verification not implemented)

Time = 0.77 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.87

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \tilde{\infty} A a^3 x \\ x(A + B \tan(c))(a + b \tan(c))^3 \cot^2(c) \\ \tilde{\infty} A a^3 x \\ -A a^3 x - \frac{A a^3}{d \tan(c+dx)} - \frac{3A a^2 b \log(\tan^2(c+dx)+1)}{2d} + \frac{3A a^2 b \log(\tan(c+dx))}{d} + 3A a b^2 x + \frac{A b^3 \log(\tan^2(c+dx)+1)}{2d} - \frac{B a^3 \log(\tan^2(c+dx)+1)}{2d} \end{cases}$$

input `integrate(cot(d*x+c)**2*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`output `Piecewise((zoo*A*a**3*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**3*cot(c)**2, Eq(d, 0)), (zoo*A*a**3*x, Eq(c, -d*x)), (-A*a**3*x - A*a**3/(d*tan(c + d*x)) - 3*A*a**2*b*log(tan(c + d*x)**2 + 1)/(2*d) + 3*A*a**2*b*log(tan(c + d*x))/d + 3*A*a*b**2*x + A*b**3*log(tan(c + d*x)**2 + 1)/(2*d) - B*a**3*log(tan(c + d*x)**2 + 1)/(2*d) + B*a**3*log(tan(c + d*x))/d + 3*B*a**2*b*x + 3*B*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) - B*b**3*x + B*b**3*tan(c + d*x)/d, True))`**3.252.7 Maxima [A] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.05

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{2 B b^3 \tan(dx + c) - \frac{2 A a^3}{\tan(dx+c)} - 2(A a^3 - 3 B a^2 b - 3 A a b^2 + B b^3)(dx + c) - (B a^3 + 3 A a^2 b - 3 B a b^2 - A b^3) \log(\tan(dx + c)^2 + 1) + 2(B a^3 + 3 A a^2 b) \log(\tan(dx + c))}{2 d}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`output `1/2*(2*B*b^3*tan(d*x + c) - 2*A*a^3/tan(d*x + c) - 2*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*(d*x + c) - (B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*log(tan(d*x + c)^2 + 1) + 2*(B*a^3 + 3*A*a^2*b)*log(tan(d*x + c)))/d`

3.252.8 Giac [A] (verification not implemented)

Time = 1.44 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.28

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{2 B b^3 \tan(dx + c) - 2(Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3)(dx + c) - (Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3) \log(\tan(dx + c))}{2d}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `1/2*(2*B*b^3*tan(d*x + c) - 2*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*(d*x + c) - (B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*log(tan(d*x + c)^2 + 1) + 2*(B*a^3 + 3*A*a^2*b)*log(abs(tan(d*x + c))) - 2*(B*a^3*tan(d*x + c) + 3*A*a^2*b*tan(d*x + c) + A*a^3)/tan(d*x + c))/d`

3.252.9 Mupad [B] (verification not implemented)

Time = 7.70 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.96

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx)) (B a^3 + 3 A b a^2)}{d} - \frac{A a^3 \cot(c + dx)}{d}$$

$$+ \frac{B b^3 \tan(c + dx)}{d} + \frac{\ln(\tan(c + dx) - i) (A + B i) (a + b i)^3 i}{2d}$$

$$- \frac{\ln(\tan(c + dx) + i) (A - B i) (a - b i)^3 i}{2d}$$

input `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`

output `(log(tan(c + d*x))*(B*a^3 + 3*A*a^2*b))/d + (log(tan(c + d*x) - 1i)*(A + B*1i)*(a + b*1i)^3*1i)/(2*d) - (log(tan(c + d*x) + 1i)*(A - B*1i)*(a - b*1i)^3*1i)/(2*d) - (A*a^3*cot(c + d*x))/d + (B*b^3*tan(c + d*x))/d`

3.253 $\int \cot^3(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.253.1 Optimal result	2450
3.253.2 Mathematica [C] (verified)	2450
3.253.3 Rubi [A] (verified)	2451
3.253.4 Maple [A] (verified)	2454
3.253.5 Fricas [A] (verification not implemented)	2455
3.253.6 Sympy [B] (verification not implemented)	2455
3.253.7 Maxima [A] (verification not implemented)	2456
3.253.8 Giac [A] (verification not implemented)	2456
3.253.9 Mupad [B] (verification not implemented)	2457

3.253.1 Optimal result

Integrand size = 31, antiderivative size = 127

$$\int \cot^3(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= -((3a^2Ab - Ab^3 + a^3B - 3ab^2B)x) - \frac{a^2(2Ab + aB) \cot(c+dx)}{d} - \frac{b^3B \log(\cos(c+dx))}{d}$$

$$- \frac{a(a^2A - 3Ab^2 - 3abB) \log(\sin(c+dx))}{d} - \frac{aA \cot^2(c+dx)(a+b \tan(c+dx))^2}{2d}$$

```
output - (3*A*a^2*b - A*b^3 + B*a^3 - 3*B*a*b^2)*x - a^2*(2*A*b + B*a)*cot(d*x+c)/d - b^3*B*ln
(cos(d*x+c))/d - a*(A*a^2 - 3*A*b^2 - 3*B*a*b)*ln(sin(d*x+c))/d - 1/2*a*A*cot(d*x+
c)^2*(a+b*tan(d*x+c))^2/d
```

3.253.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.45 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.99

$$\int \cot^3(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{-2a^2(3Ab + aB) \cot(c+dx) - a^3A \cot^2(c+dx) + (a+ib)^3(A+iB) \log(i - \tan(c+dx)) - 2a(a^2A - 3Ab^2 - 3abB) \log(\sin(c+dx))}{2d}$$

input `Integrate[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output $(-2*a^2*(3*A*b + a*B)*\text{Cot}[c + d*x] - a^3*A*\text{Cot}[c + d*x]^2 + (a + I*b)^3*(A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]] - 2*a*(a^2*A - 3*A*b^2 - 3*a*b*B)*\text{Log}[\text{Tan}[c + d*x]] + (a - I*b)^3*(A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]])/(2*d)$

3.253.3 Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.02, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4088, 27, 3042, 4118, 25, 3042, 4107, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^3(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))^3(A + B \tan(c + dx))}{\tan(c + dx)^3} dx \\
 & \quad \downarrow \text{4088} \\
 & \frac{1}{2} \int 2 \cot^2(c + dx)(a + b \tan(c + dx)) (b^2 B \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)) dx - \\
 & \quad \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{27} \\
 & \int \cot^2(c + dx)(a + b \tan(c + dx)) (b^2 B \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)) dx - \\
 & \quad \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx)) (b^2 B \tan^2(c + dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB))}{\tan(c + dx)^2} dx - \\
 & \quad \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} \\
 & \quad \downarrow \text{4118}
 \end{aligned}$$

3.253. $\int \cot^3(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \int -\cot(c + dx) \left(-B \tan^2(c + dx) b^3 + a(Aa^2 - 3bBa - 3Ab^2) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) \right) dx - \\
& \frac{a^2(aB + 2Ab) \cot(c + dx)}{d} - \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} \\
& \quad \downarrow 25 \\
& - \int \cot(c + dx) \left(-B \tan^2(c + dx) b^3 + a(Aa^2 - 3bBa - 3Ab^2) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) \right) dx - \\
& \frac{a^2(aB + 2Ab) \cot(c + dx)}{d} - \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} \\
& \quad \downarrow 3042 \\
& - \int \frac{-B \tan(c + dx)^2 b^3 + a(Aa^2 - 3bBa - 3Ab^2) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\tan(c + dx)} dx - \\
& \frac{a^2(aB + 2Ab) \cot(c + dx)}{d} - \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} \\
& \quad \downarrow 4107 \\
& -a(a^2A - 3abB - 3Ab^2) \int \cot(c + dx) dx + b^3B \int \tan(c + dx) dx - \frac{a^2(aB + 2Ab) \cot(c + dx)}{d} - \\
& x(a^3B + 3a^2Ab - 3ab^2B - Ab^3) - \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} \\
& \quad \downarrow 3042 \\
& -a(a^2A - 3abB - 3Ab^2) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + b^3B \int \tan(c + dx) dx - \\
& \frac{a^2(aB + 2Ab) \cot(c + dx)}{d} - x(a^3B + 3a^2Ab - 3ab^2B - Ab^3) - \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} \\
& \quad \downarrow 25 \\
& a(a^2A - 3abB - 3Ab^2) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + b^3B \int \tan(c + dx) dx - \\
& \frac{a^2(aB + 2Ab) \cot(c + dx)}{d} - x(a^3B + 3a^2Ab - 3ab^2B - Ab^3) - \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} \\
& \quad \downarrow 3956 \\
& \frac{a(a^2A - 3abB - 3Ab^2) \log(-\sin(c + dx))}{d} - \frac{a^2(aB + 2Ab) \cot(c + dx)}{d} - \\
& x(a^3B + 3a^2Ab - 3ab^2B - Ab^3) - \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^2}{2d} - \frac{b^3B \log(\cos(c + dx))}{d}
\end{aligned}$$

input `Int[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output $-\left(\left(3a^2Ab - Ab^3 + a^3B - 3ab^2B\right)x\right) - \left(a^2\left(2Ab + aB\right)\cot\left[c + dx\right]\right)/d - \left(b^3B\log\left[\cos\left[c + dx\right]\right]\right)/d - \left(a\left(a^2A - 3Ab^2 - 3abB\right)\log\left[-\sin\left[c + dx\right]\right]\right)/d - \left(aA\cot\left[c + dx\right]^2\left(a + b\tan\left[c + dx\right]\right)^2\right)/\left(2d\right)$

3.253.3.1 Defintions of rubi rules used

rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$

rule 27 $\text{Int}[(a_)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)(Gx_)] /; \text{FreeQ}[b, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3956 $\text{Int}[\tan[(c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\log[\text{RemoveContent}[\cos[c + dx], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

rule 4088 $\text{Int}[\left(\left(a_.\right) + \left(b_.\right)\tan\left[\left(e_.\right) + \left(f_.\right)\left(x_.\right)\right]\right)^m \left(\left(A_.\right) + \left(B_.\right)\tan\left[\left(e_.\right) + \left(f_.\right)\left(x_.\right)\right]\right)^n, x_Symbol] \rightarrow \text{Simp}[\left(b*c - a*d\right)\left(B*c - A*d\right)\left(a + b\tan\left[e + f*x\right]\right)^{m-1} \left(c + d\tan\left[e + f*x\right]\right)^{n+1} / \left(d*f*(n+1)*(c^2 + d^2)\right), x] - \text{Simp}\left[1 / \left(d*(n+1)*(c^2 + d^2)\right) \text{Int}\left[\left(a + b\tan\left[e + f*x\right]\right)^{m-2} \left(c + d\tan\left[e + f*x\right]\right)^{n+1} \text{Simp}\left[a*A*d*(b*d*(m-1) - a*c*(n+1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m-1) + a*d*(n+1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n+1)*\tan\left[e + f*x\right] - b*(d*(A*b*c + a*B*c - a*A*d)*(m+n) - b*B*(c^2*(m-1) - d^2*(n+1)))\right]*\tan\left[e + f*x\right]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ (\text{IntegerQ}[m] \ || \ \text{IntegersQ}[2*m, 2*n])$

rule 4107 $\text{Int}[\left(\left(A_.\right) + \left(B_.\right)\tan\left[\left(e_.\right) + \left(f_.\right)\left(x_.\right)\right] + \left(C_.\right)\tan\left[\left(e_.\right) + \left(f_.\right)\left(x_.\right)\right]^2\right) / \tan\left[\left(e_.\right) + \left(f_.\right)\left(x_.\right)\right], x_Symbol] \rightarrow \text{Simp}[B*x, x] + \left(\text{Simp}[A \text{ Int}[1/\tan\left[e + f*x\right], x], x] + \text{Simp}[C \text{ Int}[\tan\left[e + f*x\right], x], x]\right) /; \text{FreeQ}[\{e, f, A, B, C\}, x] \ \&\& \ \text{NeQ}[A, C]$

```
rule 4118 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.
)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f
_.)*(x_)])^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]
```

3.253.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.07

method	result
parallelrisch	$\frac{(A a^3 - 3A a b^2 - 3B a^2 b + B b^3) \ln(\sec^2(dx+c)) + (-2A a^3 + 6A a b^2 + 6B a^2 b) \ln(\tan(dx+c)) - A(\cot^2(dx+c)) a^3 + (-6A a^3 + 6A a b^2 + 6B a^2 b) \ln(\tan(dx+c))}{2d}$
derivativedivides	$\frac{(A a^3 - 3A a b^2 - 3B a^2 b + B b^3) \ln(1 + \tan^2(dx+c))}{2} + \frac{(-3A a^2 b + A b^3 - B a^3 + 3B a b^2) \arctan(\tan(dx+c)) - \frac{A a^3}{2 \tan(dx+c)^2} - \frac{a^2(3A a b^2 + 6B a^2 b)}{2 \tan(dx+c)}}{d}$
default	$\frac{(A a^3 - 3A a b^2 - 3B a^2 b + B b^3) \ln(1 + \tan^2(dx+c))}{2} + \frac{(-3A a^2 b + A b^3 - B a^3 + 3B a b^2) \arctan(\tan(dx+c)) - \frac{A a^3}{2 \tan(dx+c)^2} - \frac{a^2(3A a b^2 + 6B a^2 b)}{2 \tan(dx+c)}}{d}$
norman	$\frac{(-3A a^2 b + A b^3 - B a^3 + 3B a b^2) x (\tan^2(dx+c)) - \frac{A a^3}{2d} - \frac{a^2(3A a b^2 + 6B a^2 b) \tan(dx+c)}{d}}{\tan(dx+c)^2} + \frac{(A a^3 - 3A a b^2 - 3B a^2 b + B b^3) \ln(1 + \tan^2(dx+c))}{2d}$
risch	$-\frac{6iB a^2 b c}{d} - 3iB a^2 b x + \frac{2iB b^3 c}{d} - \frac{6iA a b^2 c}{d} - 3A a^2 b x + A b^3 x - B a^3 x + 3B a b^2 x - \frac{2ia^2(iA a^2 b^2 + 6iA a b^2 c + 6iA a^2 b^2 c + 6iA a^2 b^2 c + 6iA a^2 b^2 c)}{d}$

```
input int(cot(d*x+c)^3*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

```
output 1/2*((A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*ln(sec(d*x+c)^2)+(-2*A*a^3+6*A*a*b^
2+6*B*a^2*b)*ln(tan(d*x+c))-A*cot(d*x+c)^2*a^3+(-6*A*a^2*b-2*B*a^3)*cot(d*
x+c)-6*(A*a^2*b-1/3*A*b^3+1/3*B*a^3-B*a*b^2)*d*x)/d
```

3.253. $\int \cot^3(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.253.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.28

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx =$$

$$\frac{Bb^3 \log\left(\frac{1}{\tan(dx+c)^2+1}\right) \tan(dx+c)^2 + Aa^3 + (Aa^3 - 3Ba^2b - 3Aab^2) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^2}{2d \tan(dx+c)}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-1/2*(B*b^3*log(1/(tan(d*x + c)^2 + 1))*tan(d*x + c)^2 + A*a^3 + (A*a^3 - 3*B*a^2*b - 3*A*a*b^2)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^2 + (A*a^3 + 2*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*d*x)*tan(d*x + c)^2 + 2*(B*a^3 + 3*A*a^2*b)*tan(d*x + c))/(d*tan(d*x + c)^2)`

3.253.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 262 vs. 2(121) = 242.

Time = 1.08 (sec) , antiderivative size = 262, normalized size of antiderivative = 2.06

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \tilde{\infty} Aa^3 x \\ x(A + B \tan(c))(a + b \tan(c))^3 \cot^3(c) \\ \tilde{\infty} Aa^3 x \\ \frac{Aa^3 \log(\tan^2(c+dx)+1)}{2d} - \frac{Aa^3 \log(\tan(c+dx))}{d} - \frac{Aa^3}{2d \tan^2(c+dx)} - 3Aa^2bx - \frac{3Aa^2b}{d \tan(c+dx)} - \frac{3Aab^2 \log(\tan^2(c+dx)+1)}{2d} + \frac{3Aab^2}{2d} \end{cases}$$

input `integrate(cot(d*x+c)**3*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `Piecewise((zoo*A*a**3*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**3*cot(c)**3, Eq(d, 0)), (zoo*A*a**3*x, Eq(c, -d*x)), (A*a**3*log(tan(c + d*x)**2 + 1)/(2*d) - A*a**3*log(tan(c + d*x))/d - A*a**3/(2*d*tan(c + d*x)**2) - 3*A*a**2*b*x - 3*A*a**2*b/(d*tan(c + d*x)) - 3*A*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + 3*A*a*b**2*log(tan(c + d*x))/d + A*b**3*x - B*a**3*x - B*a**3/(d*tan(c + d*x)) - 3*B*a**2*b*log(tan(c + d*x)**2 + 1)/(2*d) + 3*B*a**2*b*log(tan(c + d*x))/d + 3*B*a*b**2*x + B*b**3*log(tan(c + d*x)**2 + 1)/(2*d), True))`

3.253.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.12

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{2(Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3)(dx + c) - (Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3) \log(\tan(dx + c)^2 + 1) + \dots}{2d}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(2*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*(d*x + c) - (A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*log(tan(d*x + c)^2 + 1) + 2*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2)*log(tan(d*x + c)) + (A*a^3 + 2*(B*a^3 + 3*A*a^2*b)*tan(d*x + c)))/tan(d*x + c)^2/d`

3.253.8 Giac [A] (verification not implemented)

Time = 1.40 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.52

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{2(Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3)(dx + c) - (Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3) \log(\tan(dx + c)^2 + 1) + \dots}{2d}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output
$$-1/2*(2*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*(d*x + c) - (A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*\log(\tan(d*x + c)^2 + 1) + 2*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2)*\log(\text{abs}(\tan(d*x + c))) - (3*A*a^3*\tan(d*x + c)^2 - 9*B*a^2*b*\tan(d*x + c)^2 - 9*A*a*b^2*\tan(d*x + c)^2 - 2*B*a^3*\tan(d*x + c) - 6*A*a^2*b*\tan(d*x + c) - A*a^3)/\tan(d*x + c)^2)/d$$

3.253.9 Mupad [B] (verification not implemented)

Time = 8.15 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.06

$$\begin{aligned} & \int \cot^3(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ &= \frac{\ln(\tan(c + dx))(-Aa^3 + 3Ba^2b + 3Aab^2)}{d} \\ & \quad - \frac{\cot(c + dx)^2 \left(\tan(c + dx)(Ba^3 + 3Aba^2) + \frac{Aa^3}{2} \right)}{d} \\ & \quad + \frac{\ln(\tan(c + dx) + i)(A - B i)(b + a i)^3 i}{2d} \\ & \quad + \frac{\ln(\tan(c + dx) - i)(A + B i)(-b + a i)^3 i}{2d} \end{aligned}$$

input `int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`

output
$$(\log(\tan(c + d*x))*(3*A*a*b^2 - A*a^3 + 3*B*a^2*b))/d - (\cot(c + d*x)^2*(\tan(c + d*x)*(B*a^3 + 3*A*a^2*b) + (A*a^3)/2))/d + (\log(\tan(c + d*x) + 1i)*(A - B*1i)*(a*1i + b)^3*1i)/(2*d) + (\log(\tan(c + d*x) - 1i)*(A + B*1i)*(a*1i - b)^3*1i)/(2*d)$$

3.254 $\int \cot^4(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.254.1 Optimal result	2458
3.254.2 Mathematica [C] (verified)	2459
3.254.3 Rubi [A] (verified)	2459
3.254.4 Maple [A] (verified)	2463
3.254.5 Fricas [A] (verification not implemented)	2464
3.254.6 Sympy [B] (verification not implemented)	2464
3.254.7 Maxima [A] (verification not implemented)	2465
3.254.8 Giac [B] (verification not implemented)	2466
3.254.9 Mupad [B] (verification not implemented)	2466

3.254.1 Optimal result

Integrand size = 31, antiderivative size = 154

$$\int \cot^4(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= (a^3A - 3aAb^2 - 3a^2bB + b^3B) x + \frac{a(3a^2A - 8Ab^2 - 9abB) \cot(c+dx)}{3d}$$

$$- \frac{a^2(5Ab + 3aB) \cot^2(c+dx)}{6d} - \frac{(3a^2Ab - Ab^3 + a^3B - 3ab^2B) \log(\sin(c+dx))}{d}$$

$$- \frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^2}{3d}$$

```
output (A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*x+1/3*a*(3*A*a^2-8*A*b^2-9*B*a*b)*cot(d*x+c)/d-1/6*a^2*(5*A*b+3*B*a)*cot(d*x+c)^2/d-(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*ln(sin(d*x+c))/d-1/3*a*A*cot(d*x+c)^3*(a+b*tan(d*x+c))^2/d
```

3.254.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.28 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.06

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{6a(a^2A - 3Ab^2 - 3abB) \cot(c + dx) - 3a^2(3Ab + aB) \cot^2(c + dx) - 2a^3A \cot^3(c + dx) + 3(a + ib)^3(-i \dots)}{6d}$$

input `Integrate[Cot[c + d*x]^4*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(6*a*(a^2*A - 3*A*b^2 - 3*a*b*B)*Cot[c + d*x] - 3*a^2*(3*A*b + a*B)*Cot[c + d*x]^2 - 2*a^3*A*Cot[c + d*x]^3 + 3*(a + I*b)^3*((-I)*A + B)*Log[I - Tan[c + d*x]] - 6*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*Log[Tan[c + d*x]] + 3*(a - I*b)^3*(I*A + B)*Log[I + Tan[c + d*x]])/(6*d)`

3.254.3 Rubi [A] (verified)

Time = 1.02 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.05, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 4088, 3042, 4118, 25, 3042, 4111, 27, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^3(A + B \tan(c + dx))}{\tan(c + dx)^4} dx$$

$$\downarrow \text{4088}$$

$$\frac{1}{3} \int \cot^3(c + dx)(a + b \tan(c + dx)) (-b(aA - 3bB) \tan^2(c + dx) - 3(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(5Ab + 3aB)) dx - \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{3} \int \frac{(a + b \tan(c + dx)) (-b(aA - 3bB) \tan(c + dx)^2 - 3(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(5Ab + 3aB))}{\tan(c + dx)^3} dx - \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d}$$

↓ 4118

$$\frac{1}{3} \left(\int -\cot^2(c + dx) (b^2(aA - 3bB) \tan^2(c + dx) + 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(3Aa^2 - 9bBa - 8Ab^2)) dx + \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d} \right)$$

↓ 25

$$\frac{1}{3} \left(- \int \cot^2(c + dx) (b^2(aA - 3bB) \tan^2(c + dx) + 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(3Aa^2 - 9bBa - 8Ab^2)) dx + \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d} \right)$$

↓ 3042

$$\frac{1}{3} \left(- \int \frac{b^2(aA - 3bB) \tan(c + dx)^2 + 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(3Aa^2 - 9bBa - 8Ab^2)}{\tan(c + dx)^2} dx + \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d} \right)$$

↓ 4111

$$\frac{1}{3} \left(- \int 3 \cot(c + dx) (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)) dx + \frac{a(3a^2A - 9bBa - 8Ab^2)}{3} + \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d} \right)$$

↓ 27

$$\frac{1}{3} \left(-3 \int \cot(c + dx) (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)) dx + \frac{a(3a^2A - 9bBa - 8Ab^2)}{3} + \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d} \right)$$

↓ 3042

$$\frac{1}{3} \left(-3 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\tan(c + dx)} dx + \frac{a(3a^2A - 9abB - 8Ab^2)}{d} \right. \\ \left. \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d} \right. \\ \downarrow 4014$$

$$\frac{1}{3} \left(-3 \left((a^3B + 3a^2Ab - 3ab^2B - Ab^3) \int \cot(c + dx) dx - x(a^3A - 3a^2bB - 3aAb^2 + b^3B) \right) + \frac{a(3a^2A - 9abB - 8Ab^2)}{d} \right. \\ \left. \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d} \right. \\ \downarrow 3042$$

$$\frac{1}{3} \left(-3 \left((a^3B + 3a^2Ab - 3ab^2B - Ab^3) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - x(a^3A - 3a^2bB - 3aAb^2 + b^3B) \right) + \frac{a(3a^2A - 9abB - 8Ab^2)}{d} \right. \\ \left. \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d} \right. \\ \downarrow 25$$

$$\frac{1}{3} \left(-3 \left(-(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - (x(a^3A - 3a^2bB - 3aAb^2 + b^3B)) \right) + \frac{a(3a^2A - 9abB - 8Ab^2)}{d} \right. \\ \left. \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d} \right. \\ \downarrow 3956$$

$$\frac{1}{3} \left(\frac{a(3a^2A - 9abB - 8Ab^2) \cot(c + dx)}{d} - \frac{a^2(3aB + 5Ab) \cot^2(c + dx)}{2d} - 3 \left(\frac{(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \log|a + b \tan(c + dx)|}{d} \right) \right. \\ \left. \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^2}{3d} \right)$$

input `Int[Cot[c + d*x]^4*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `((a*(3*a^2*A - 8*A*b^2 - 9*a*b*B)*Cot[c + d*x])/d - (a^2*(5*A*b + 3*a*B)*Cot[c + d*x]^2)/(2*d) - 3*(-((a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*x) + ((3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*Log[-Sin[c + d*x]]/d))/3 - (a*A*Cot[c + d*x]^3*(a + b*Tan[c + d*x])^2)/(3*d)`

3.254.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`
- rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

```
rule 4111 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]
```

```
rule 4118 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.
)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-(b*c - a*d))*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]
```

3.254.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.12

method	result
parallelrisch	$\frac{3(3A^2b - Ab^3 + Ba^3 - 3Bab^2) \ln(\sec^2(dx+c)) + 6(-3A^2b + Ab^3 - Ba^3 + 3Bab^2) \ln(\tan(dx+c)) - 2A(\cot^3(dx+c))a^3 - B(\cot^3(dx+c))b^3}{6d}$
derivativedivides	$\frac{(3A^2b - Ab^3 + Ba^3 - 3Bab^2) \ln(1 + \tan^2(dx+c))}{2} + (Aa^3 - 3Aab^2 - 3Ba^2b + Bb^3) \arctan(\tan(dx+c)) + (-3A^2b + Ab^3 - Ba^3 + 3Bab^2) \tan(dx+c)}{d}$
default	$\frac{(3A^2b - Ab^3 + Ba^3 - 3Bab^2) \ln(1 + \tan^2(dx+c))}{2} + (Aa^3 - 3Aab^2 - 3Ba^2b + Bb^3) \arctan(\tan(dx+c)) + (-3A^2b + Ab^3 - Ba^3 + 3Bab^2) \tan(dx+c)}{d}$
norman	$\frac{(Aa^3 - 3Aab^2 - 3Ba^2b + Bb^3)x(\tan^3(dx+c)) + \frac{a(Aa^2 - 3Aab^2 - 3Bab)}{d}(\tan^2(dx+c)) - \frac{Aa^3}{3d} - \frac{a^2(3Ab + Ba)\tan(dx+c)}{2d}}{\tan(dx+c)^3} - \frac{3A^2b - Ab^3 + Ba^3 - 3Bab^2}{2d}$
risch	$Aa^3x - 3Aab^2x - 3Ba^2bx + Bb^3x - \frac{6iBab^2c}{d} + 3iAa^2bx + iBa^3x - 3iBab^2x - \frac{2iAb^3}{d}$

```
input int(cot(d*x+c)^4*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

$$3.254. \quad \int \cot^4(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

output $1/6*(3*(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*\ln(\sec(dx+c)^2)+6*(-3*A*a^2*b+A*b^3-B*a^3+3*B*a*b^2)*\ln(\tan(dx+c))-2*A*\cot(dx+c)^3*a^3+3*(-3*A*a^2*b-B*a^3)*\cot(dx+c)^2+6*a*\cot(dx+c)*(A*a^2-3*A*b^2-3*B*a*b)+6*d*x*(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3))/d$

3.254.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.18

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{3(Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^3 + 2Aa^3 + 3(Ba^3 + 3Aa^2b - 2(Aa^3 - 3Aa^2b - 3Aa*b^2 - B*b^3))}{d}$$

input `integrate(cot(dx+c)^4*(a+b*tan(dx+c))^3*(A+B*tan(dx+c)),x, algorithm="fricas")`

output $-1/6*(3*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*\log(\tan(dx + c)^2/(\tan(dx + c)^2 + 1))*\tan(dx + c)^3 + 2*A*a^3 + 3*(B*a^3 + 3*A*a^2*b - 2*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*d*x)*\tan(dx + c)^3 - 6*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2)*\tan(dx + c)^2 + 3*(B*a^3 + 3*A*a^2*b)*\tan(dx + c))/(d*\tan(dx + c)^3)$

3.254.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 332 vs. 2(150) = 300.

Time = 1.81 (sec) , antiderivative size = 332, normalized size of antiderivative = 2.16

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \begin{cases} \tilde{\omega}Aa^3x \\ x(A + B \tan(c))(a + b \tan(c))^3 \cot^4(c) \\ \tilde{\omega}Aa^3x \\ Aa^3x + \frac{Aa^3}{d \tan(c+dx)} - \frac{Aa^3}{3d \tan^3(c+dx)} + \frac{3Aa^2b \log(\tan^2(c+dx)+1)}{2d} - \frac{3Aa^2b \log(\tan(c+dx))}{d} - \frac{3Aa^2b}{2d \tan^2(c+dx)} - 3Aab^2x - \end{cases}$$

3.254. $\int \cot^4(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

input `integrate(cot(d*x+c)**4*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `Piecewise((zoo*A*a**3*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**3*cot(c)**4, Eq(d, 0)), (zoo*A*a**3*x, Eq(c, -d*x)), (A*a**3*x + A*a**3/(d*tan(c + d*x)) - A*a**3/(3*d*tan(c + d*x)**3) + 3*A*a**2*b*log(tan(c + d*x)**2 + 1)/(2*d) - 3*A*a**2*b*log(tan(c + d*x))/d - 3*A*a**2*b/(2*d*tan(c + d*x)**2) - 3*A*a*b**2*x - 3*A*a*b**2/(d*tan(c + d*x)) - A*b**3*log(tan(c + d*x)**2 + 1)/(2*d) + A*b**3*log(tan(c + d*x))/d + B*a**3*log(tan(c + d*x)**2 + 1)/(2*d) - B*a**3*log(tan(c + d*x))/d - B*a**3/(2*d*tan(c + d*x)**2) - 3*B*a**2*b*x - 3*B*a**2*b/(d*tan(c + d*x)) - 3*B*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) + 3*B*a*b**2*log(tan(c + d*x))/d + B*b**3*x, True))`

3.254.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.17

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{6(Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3)(dx + c) + 3(Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3) \log(\tan(dx + c)^2 + 1) - 6(Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3) \tan(dx + c) + 3(Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3) \tan(dx + c)^2 + 3(Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3) \tan(dx + c)^3}{d}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/6*(6*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*(d*x + c) + 3*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*log(tan(d*x + c)^2 + 1) - 6*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*log(tan(d*x + c)) - (2*A*a^3 - 6*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2)*tan(d*x + c)^2 + 3*(B*a^3 + 3*A*a^2*b)*tan(d*x + c))/tan(d*x + c)^3)/d`

3.254. $\int \cot^4(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.254.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 390 vs. $2(148) = 296$.

Time = 0.97 (sec) , antiderivative size = 390, normalized size of antiderivative = 2.53

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{Aa^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 3Ba^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 9Aa^2b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 15Aa^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 36A^2a^2b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 15A^2a^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 36A^2a^2b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 36A^2a^2b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 24(Aa^3 - 3Ba^2b - 3A^2a^2b - 3A^2a^2b + B^2b^3)(dx + c) + 24(Ba^3 + 3A^2a^2b - 3Ba^2b^2 - A^2b^3) \log(\tan^2\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1) - 24(Ba^3 + 3A^2a^2b - 3Ba^2b^2 - A^2b^3) \log(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)) + (44Ba^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 132A^2a^2b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 132Ba^2b^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 44A^2b^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 15A^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 36Ba^2b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 36A^2a^2b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 3Ba^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 9A^2a^2b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - A^3/\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3)/d$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output $\frac{1}{24}(Aa^3 \tan^3\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 3Ba^3 \tan^2\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 9A^2a^2b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 15A^2a^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 36A^2a^2b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 36A^2a^2b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 24(Aa^3 - 3Ba^2b - 3A^2a^2b - 3A^2a^2b + B^2b^3)(dx + c) + 24(Ba^3 + 3A^2a^2b - 3Ba^2b^2 - A^2b^3) \log(\tan^2\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 1) - 24(Ba^3 + 3A^2a^2b - 3Ba^2b^2 - A^2b^3) \log(\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)) + (44Ba^3 \tan^3\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 132A^2a^2b \tan^3\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 132Ba^2b^2 \tan^3\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 44A^2b^3 \tan^3\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 15A^3 \tan^2\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 36Ba^2b \tan^2\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 36A^2a^2b \tan^2\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 3Ba^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 9A^2a^2b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - A^3/\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3)/d$

3.254.9 Mupad [B] (verification not implemented)

Time = 8.17 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.10

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx))(-Ba^3 - 3Aa^2b + 3Bab^2 + Ab^3)}{d} - \frac{\cot(c + dx)^3 \left(\tan(c + dx) \left(\frac{Ba^3}{2} + \frac{3Aba^2}{2} \right) + \frac{Aa^3}{3} + \tan(c + dx)^2(-Aa^3 + 3Ba^2b + 3Aab^2) \right)}{d} - \frac{\ln(\tan(c + dx) - i)(A + Bli)(a + bli)^3 li}{2d} + \frac{\ln(\tan(c + dx) + li)(A - Bli)(a - bli)^3 li}{2d}$$

3.254. $\int \cot^4(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

input `int(cot(c + d*x)^4*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`

output `(log(tan(c + d*x))*(A*b^3 - B*a^3 - 3*A*a^2*b + 3*B*a*b^2))/d - (cot(c + d*x)^3*(tan(c + d*x)*((B*a^3)/2 + (3*A*a^2*b)/2) + (A*a^3)/3 + tan(c + d*x)^2*(3*A*a*b^2 - A*a^3 + 3*B*a^2*b)))/d - (log(tan(c + d*x) - 1i)*(A + B*1i)*(a + b*1i)^3*1i)/(2*d) + (log(tan(c + d*x) + 1i)*(A - B*1i)*(a - b*1i)^3*1i)/(2*d)`

3.255 $\int \cot^5(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.255.1 Optimal result	2468
3.255.2 Mathematica [C] (verified)	2469
3.255.3 Rubi [A] (verified)	2469
3.255.4 Maple [A] (verified)	2474
3.255.5 Fricas [A] (verification not implemented)	2475
3.255.6 Sympy [B] (verification not implemented)	2475
3.255.7 Maxima [A] (verification not implemented)	2476
3.255.8 Giac [B] (verification not implemented)	2476
3.255.9 Mupad [B] (verification not implemented)	2477

3.255.1 Optimal result

Integrand size = 31, antiderivative size = 191

$$\int \cot^5(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= (3a^2Ab - Ab^3 + a^3B - 3ab^2B) x + \frac{(3a^2Ab - Ab^3 + a^3B - 3ab^2B) \cot(c+dx)}{d}$$

$$+ \frac{a(2a^2A - 5Ab^2 - 6abB) \cot^2(c+dx)}{4d} - \frac{a^2(3Ab + 2aB) \cot^3(c+dx)}{6d}$$

$$+ \frac{(a^3A - 3aAb^2 - 3a^2bB + b^3B) \log(\sin(c+dx))}{d} - \frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^2}{4d}$$

output

```
(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*x+(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*cot(d*x+c)/d+1/4*a*(2*A*a^2-5*A*b^2-6*B*a*b)*cot(d*x+c)^2/d-1/6*a^2*(3*A*b+2*B*a)*cot(d*x+c)^3/d+(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*ln(sin(d*x+c))/d-1/4*a*A*cot(d*x+c)^4*(a+b*tan(d*x+c))^2/d
```

3.255.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.78 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.04

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{12(3a^2Ab - Ab^3 + a^3B - 3ab^2B) \cot(c + dx) + 6a(a^2A - 3Ab^2 - 3abB) \cot^2(c + dx) - 4a^2(3Ab + aB) \cot^3(c + dx) + 12a^2b^2(A + B) \cot^4(c + dx) - 6a^2b^2(A + B) \cot^5(c + dx) + 12a^2b^2(A + B) \tan(c + dx) - 6a^2b^2(A + B) \tan^2(c + dx) + 12a^2b^2(A + B) \tan^3(c + dx) - 6a^2b^2(A + B) \tan^4(c + dx) + 12a^2b^2(A + B) \tan^5(c + dx)}{12d}$$

input `Integrate[Cot[c + d*x]^5*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output $(12*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*\text{Cot}[c + d*x] + 6*a*(a^2*A - 3*A*b^2 - 3*a*b*B)*\text{Cot}[c + d*x]^2 - 4*a^2*(3*A*b + a*B)*\text{Cot}[c + d*x]^3 - 3*a^3*A*\text{Cot}[c + d*x]^4 - 6*(a + I*b)^3*(A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]] + 12*(a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*\text{Log}[\text{Tan}[c + d*x]] - 6*(a - I*b)^3*(A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]])/(12*d)$

3.255.3 Rubi [A] (verified)

Time = 1.29 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.07, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.548$, Rules used = {3042, 4088, 27, 3042, 4118, 25, 3042, 4111, 27, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^3(A + B \tan(c + dx))}{\tan(c + dx)^5} dx$$

$$\downarrow \text{4088}$$

$$\frac{1}{4} \int 2 \cot^4(c + dx)(a + b \tan(c + dx)) (-b(aA - 2bB) \tan^2(c + dx) - 2(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(3Ab + 2aB)) dx - \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d}$$

3.255. $\int \cot^5(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\downarrow 27$$

$$\frac{1}{2} \int \frac{\cot^4(c+dx)(a+b\tan(c+dx))(-b(aA-2bB)\tan^2(c+dx) - 2(Aa^2-2bBa-Ab^2)\tan(c+dx) + a(3Ab+2aB)) dx - aA \cot^4(c+dx)(a+b\tan(c+dx))^2}{4d}$$

$$\downarrow 3042$$

$$\frac{1}{2} \int \frac{(a+b\tan(c+dx))(-b(aA-2bB)\tan(c+dx)^2 - 2(Aa^2-2bBa-Ab^2)\tan(c+dx) + a(3Ab+2aB))}{\tan(c+dx)^4} dx - \frac{aA \cot^4(c+dx)(a+b\tan(c+dx))^2}{4d}$$

$$\downarrow 4118$$

$$\frac{1}{2} \left(\int -\cot^3(c+dx)(b^2(aA-2bB)\tan^2(c+dx) + 2(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx) + a(2Aa^2-6bBa-5Ab^2)) dx - \frac{aA \cot^4(c+dx)(a+b\tan(c+dx))^2}{4d} \right)$$

$$\downarrow 25$$

$$\frac{1}{2} \left(- \int \cot^3(c+dx)(b^2(aA-2bB)\tan^2(c+dx) + 2(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx) + a(2Aa^2-6bBa-5Ab^2)) dx - \frac{aA \cot^4(c+dx)(a+b\tan(c+dx))^2}{4d} \right)$$

$$\downarrow 3042$$

$$\frac{1}{2} \left(- \int \frac{b^2(aA-2bB)\tan(c+dx)^2 + 2(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx) + a(2Aa^2-6bBa-5Ab^2)}{\tan(c+dx)^3} dx - \frac{aA \cot^4(c+dx)(a+b\tan(c+dx))^2}{4d} \right)$$

$$\downarrow 4111$$

$$\frac{1}{2} \left(- \int 2 \cot^2(c+dx)(Ba^3+3Aba^2-3b^2Ba-Ab^3 - (Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx)) dx + \frac{a(2a^2A-6bBa-5Ab^2)}{4d} - \frac{aA \cot^4(c+dx)(a+b\tan(c+dx))^2}{4d} \right)$$

$$\downarrow 27$$

$$\frac{1}{2} \left(-2 \int \cot^2(c+dx) (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx)) dx + \frac{a(2a^2A}{4d} \cot^4(c+dx)(a+b \tan(c+dx))^2 \right)$$

↓ 3042

$$\frac{1}{2} \left(-2 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx)}{\tan(c+dx)^2} dx + \frac{a(2a^2A - 6abB - 5Ab^2)}{2d} \cot^4(c+dx)(a+b \tan(c+dx))^2 \right)$$

↓ 4012

$$\frac{1}{2} \left(-2 \left(\int -\cot(c+dx) (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)) dx - \frac{(a^3B + 3a^2Ab - 3ab^2B - Ab^3)}{d} \cot(c+dx) \right) + \frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^2}{4d} \right)$$

↓ 25

$$\frac{1}{2} \left(-2 \left(- \int \cot(c+dx) (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)) dx - \frac{(a^3B + 3a^2Ab - 3ab^2B - Ab^3)}{d} \cot(c+dx) \right) + \frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^2}{4d} \right)$$

↓ 3042

$$\frac{1}{2} \left(-2 \left(- \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)}{\tan(c+dx)} dx - \frac{(a^3B + 3a^2Ab - 3ab^2B - Ab^3)}{d} \cot(c+dx) \right) + \frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^2}{4d} \right)$$

↓ 4014

$$\frac{1}{2} \left(-2 \left(-(a^3A - 3a^2bB - 3aAb^2 + b^3B) \int \cot(c+dx) dx - \frac{(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \cot(c+dx)}{d} - (x(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \cot(c+dx) + \frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^2}{4d}) \right) \right)$$

↓ 3042

$$\frac{1}{2} \left(-2 \left(-(a^3 A - 3a^2 b B - 3a A b^2 + b^3 B) \int -\tan \left(c + dx + \frac{\pi}{2} \right) dx - \frac{(a^3 B + 3a^2 A b - 3a b^2 B - A b^3) \cot(c + dx)}{d} \right. \right. \\ \left. \left. \frac{a A \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d} \right) \right. \\ \left. \downarrow 25 \right.$$

$$\frac{1}{2} \left(-2 \left((a^3 A - 3a^2 b B - 3a A b^2 + b^3 B) \int \tan \left(\frac{1}{2}(2c + \pi) + dx \right) dx - \frac{(a^3 B + 3a^2 A b - 3a b^2 B - A b^3) \cot(c + dx)}{d} \right. \right. \\ \left. \left. \frac{a A \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d} \right) \right. \\ \left. \downarrow 3956 \right.$$

$$\frac{1}{2} \left(\frac{a(2a^2 A - 6abB - 5Ab^2) \cot^2(c + dx)}{2d} - \frac{a^2(2aB + 3Ab) \cot^3(c + dx)}{3d} - 2 \left(-\frac{(a^3 B + 3a^2 A b - 3a b^2 B - A b^3) \cot(c + dx)}{d} \right. \right. \\ \left. \left. \frac{a A \cot^4(c + dx)(a + b \tan(c + dx))^2}{4d} \right) \right.$$

input `Int[Cot[c + d*x]^5*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `((a*(2*a^2*A - 5*A*b^2 - 6*a*b*B)*Cot[c + d*x]^2)/(2*d) - (a^2*(3*A*b + 2*a*B)*Cot[c + d*x]^3)/(3*d) - 2*(-((3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*x) - ((3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*Cot[c + d*x])/d - ((a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*Log[-Sin[c + d*x]]/d))/2 - (a*A*Cot[c + d*x]^4*(a + b*Tan[c + d*x])^2)/(4*d)`

3.255.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)])^(n_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4111 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)] + (C_.)*tan[(e_.) + (f_.)*(x_.)]^2), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`


```
rule 4118 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.
)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f
_.)*(x_)])^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]
```

3.255.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.09

method	result
parallelrisch	$\frac{6(-Aa^3+3Aab^2+3Ba^2b-Bb^3)\ln(\sec^2(dx+c))+12(Aa^3-3Aab^2-3Ba^2b+Bb^3)\ln(\tan(dx+c))-3A(\cot^4(dx+c))a^3}{d}$
derivativedivides	$\frac{(-Aa^3+3Aab^2+3Ba^2b-Bb^3)\ln(1+\tan^2(dx+c))}{2} + (3Aa^2b-Ab^3+Ba^3-3Ba^2b^2)\arctan(\tan(dx+c)) - \frac{3Aa^2b+Ab^3-Ba^3}{\tan(dx+c)}$
default	$\frac{(-Aa^3+3Aab^2+3Ba^2b-Bb^3)\ln(1+\tan^2(dx+c))}{2} + (3Aa^2b-Ab^3+Ba^3-3Ba^2b^2)\arctan(\tan(dx+c)) - \frac{3Aa^2b+Ab^3-Ba^3}{\tan(dx+c)}$
norman	$\frac{(3Aa^2b-Ab^3+Ba^3-3Ba^2b^2)(\tan^3(dx+c))}{d} + (3Aa^2b-Ab^3+Ba^3-3Ba^2b^2)x(\tan^4(dx+c)) - \frac{Aa^3}{4d} + \frac{a(Aa^2-3Aa^2b^2-3Bab)}{2d}$
risch	$\frac{6iAab^2c}{d} - \frac{2iBb^3c}{d} + 3iBa^2bx - \frac{2i(12Aa^2b-9Ba^2b^2+4Ba^3-6iAa^3e^{2i(dx+c)}-6iAa^3e^{6i(dx+c)}+6iAa^3e^{4i(dx+c)})}{d}$

```
input int(cot(d*x+c)^5*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/12*(6*(-A*a^3+3*A*a*b^2+3*B*a^2*b-B*b^3)*ln(sec(d*x+c)^2)+12*(A*a^3-3*A*
a*b^2-3*B*a^2*b+B*b^3)*ln(tan(d*x+c))-3*A*cot(d*x+c)^4*a^3+4*(-3*A*a^2*b-B
*a^3)*cot(d*x+c)^3+6*a*cot(d*x+c)^2*(A*a^2-3*A*b^2-3*B*a*b)+12*cot(d*x+c)*
(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)+36*(A*a^2*b-1/3*A*b^3+1/3*B*a^3-B*a*b^2)
*d*x)/d
```

3.255. $\int \cot^5(c+dx)(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx$

3.255.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.18

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{6(Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^4 + 3(3Aa^3 - 6Ba^2b - 6Aab^2 + 4(Ba^3$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/12*(6*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^4 + 3*(3*A*a^3 - 6*B*a^2*b - 6*A*a*b^2 + 4*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*d*x)*tan(d*x + c)^4 - 3*A*a^3 + 12*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*tan(d*x + c)^3 + 6*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2)*tan(d*x + c)^2 - 4*(B*a^3 + 3*A*a^2*b)*tan(d*x + c))/(d*tan(d*x + c)^4)`

3.255.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 400 vs. 2(187) = 374.

Time = 2.31 (sec) , antiderivative size = 400, normalized size of antiderivative = 2.09

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \tilde{\infty} Aa^3 x \\ x(A + B \tan(c))(a + b \tan(c))^3 \cot^5(c) \\ \tilde{\infty} Aa^3 x \\ -\frac{Aa^3 \log(\tan^2(c+dx)+1)}{2d} + \frac{Aa^3 \log(\tan(c+dx))}{d} + \frac{Aa^3}{2d \tan^2(c+dx)} - \frac{Aa^3}{4d \tan^4(c+dx)} + 3Aa^2bx + \frac{3Aa^2b}{d \tan(c+dx)} - \frac{Aa^2b}{d \tan^3(c+dx)} \end{cases}$$

input `integrate(cot(d*x+c)**5*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

```
output Piecewise((zoo*A*a**3*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**3*cot(c)**5, Eq(d, 0)), (zoo*A*a**3*x, Eq(c, -d*x)), (-A*a**3*log(tan(c + d*x)**2 + 1)/(2*d) + A*a**3*log(tan(c + d*x))/d + A*a**3/(2*d*tan(c + d*x)**2) - A*a**3/(4*d*tan(c + d*x)**4) + 3*A*a**2*b*x + 3*A*a**2*b/(d*tan(c + d*x)) - A*a**2*b/(d*tan(c + d*x)**3) + 3*A*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) - 3*A*a*b**2*log(tan(c + d*x))/d - 3*A*a*b**2/(2*d*tan(c + d*x)**2) - A*b**3*x - A*b**3/(d*tan(c + d*x)) + B*a**3*x + B*a**3/(d*tan(c + d*x)) - B*a**3/(3*d*tan(c + d*x)**3) + 3*B*a**2*b*log(tan(c + d*x)**2 + 1)/(2*d) - 3*B*a**2*b*log(tan(c + d*x))/d - 3*B*a**2*b/(2*d*tan(c + d*x)**2) - 3*B*a*b**2*x - 3*B*a*b**2/(d*tan(c + d*x)) - B*b**3*log(tan(c + d*x)**2 + 1)/(2*d) + B*b**3*log(tan(c + d*x))/d, True))
```

3.255.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.13

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{12(Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3)(dx + c) - 6(Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3) \log(\tan(dx + c)^2 + 1) + \dots}{\dots}$$

```
input integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")
```

```
output 1/12*(12*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*(d*x + c) - 6*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*log(tan(d*x + c)^2 + 1) + 12*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 - A*b^3)*log(tan(d*x + c)) - (3*A*a^3 - 12*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*tan(d*x + c)^3 - 6*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2)*tan(d*x + c)^2 + 4*(B*a^3 + 3*A*a^2*b)*tan(d*x + c))/tan(d*x + c)^4/d
```

3.255.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 528 vs. 2(185) = 370.

Time = 1.09 (sec) , antiderivative size = 528, normalized size of antiderivative = 2.76

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx =$$

$$\frac{3Aa^3 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^4 - 8Ba^3 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 24Aa^2b \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 36Aa^3 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + \dots}{\dots}$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `-1/192*(3*A*a^3*tan(1/2*d*x + 1/2*c)^4 - 8*B*a^3*tan(1/2*d*x + 1/2*c)^3 - 24*A*a^2*b*tan(1/2*d*x + 1/2*c)^3 - 36*A*a^3*tan(1/2*d*x + 1/2*c)^2 + 72*B*a^2*b*tan(1/2*d*x + 1/2*c)^2 + 72*A*a*b^2*tan(1/2*d*x + 1/2*c)^2 + 120*B*a^3*tan(1/2*d*x + 1/2*c) + 360*A*a^2*b*tan(1/2*d*x + 1/2*c) - 288*B*a*b^2*tan(1/2*d*x + 1/2*c) - 96*A*b^3*tan(1/2*d*x + 1/2*c) - 192*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*(d*x + c) + 192*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*log(tan(1/2*d*x + 1/2*c)^2 + 1) - 192*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*log(abs(tan(1/2*d*x + 1/2*c))) + (400*A*a^3*tan(1/2*d*x + 1/2*c)^4 - 1200*B*a^2*b*tan(1/2*d*x + 1/2*c)^4 - 1200*A*a*b^2*tan(1/2*d*x + 1/2*c)^4 + 400*B*b^3*tan(1/2*d*x + 1/2*c)^4 - 120*B*a^3*tan(1/2*d*x + 1/2*c)^3 - 360*A*a^2*b*tan(1/2*d*x + 1/2*c)^3 + 288*B*a*b^2*tan(1/2*d*x + 1/2*c)^3 + 96*A*b^3*tan(1/2*d*x + 1/2*c)^3 - 36*A*a^3*tan(1/2*d*x + 1/2*c)^2 + 72*B*a^2*b*tan(1/2*d*x + 1/2*c)^2 + 72*A*a*b^2*tan(1/2*d*x + 1/2*c)^2 + 8*B*a^3*tan(1/2*d*x + 1/2*c) + 24*A*a^2*b*tan(1/2*d*x + 1/2*c) + 3*A*a^3)/tan(1/2*d*x + 1/2*c)^4)/d`

3.255.9 Mupad [B] (verification not implemented)

Time = 7.83 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.07

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx)) (A a^3 - 3 B a^2 b - 3 A a b^2 + B b^3)}{d}$$

$$- \frac{\cot(c + dx)^4 \left(\tan(c + dx) \left(\frac{B a^3}{3} + A b a^2 \right) + \frac{A a^3}{4} + \tan(c + dx)^2 \left(-\frac{A a^3}{2} + \frac{3 B a^2 b}{2} + \frac{3 A a b^2}{2} \right) + \tan(c + dx) \left(\frac{B a^3}{3} + A b a^2 \right) + \frac{A a^3}{4} \right)}{d}$$

$$- \frac{\ln(\tan(c + dx) + 1i) (A - B 1i) (b + a 1i)^3 1i}{2 d}$$

$$- \frac{\ln(\tan(c + dx) - 1i) (A + B 1i) (-b + a 1i)^3 1i}{2 d}$$

input `int(cot(c + d*x)^5*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`

output $(\log(\tan(c + dx))(Aa^3 + Bb^3 - 3Aab^2 - 3Ba^2b))/d - (\cot(c + dx))^4(\tan(c + dx)((Ba^3)/3 + Aa^2b) + (Aa^3)/4 + \tan(c + dx)^2((3Aab^2)/2 - (Aa^3)/2 + (3Ba^2b)/2) + \tan(c + dx)^3(Ab^3 - Ba^3 - 3Aa^2b + 3Bab^2))/d - (\log(\tan(c + dx) + 1i)(A - B1i)(a1i + b)^31i)/(2*d) - (\log(\tan(c + dx) - 1i)(A + B1i)(a1i - b)^31i)/(2*d)$

3.256 $\int \cot^6(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.256.1 Optimal result	2479
3.256.2 Mathematica [C] (verified)	2480
3.256.3 Rubi [A] (verified)	2480
3.256.4 Maple [A] (verified)	2485
3.256.5 Fricas [A] (verification not implemented)	2486
3.256.6 Sympy [B] (verification not implemented)	2486
3.256.7 Maxima [A] (verification not implemented)	2487
3.256.8 Giac [B] (verification not implemented)	2488
3.256.9 Mupad [B] (verification not implemented)	2489

3.256.1 Optimal result

Integrand size = 31, antiderivative size = 233

$$\int \cot^6(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= -((a^3A - 3aAb^2 - 3a^2bB + b^3B)x) - \frac{(a^3A - 3aAb^2 - 3a^2bB + b^3B) \cot(c+dx)}{d}$$

$$+ \frac{(3a^2Ab - Ab^3 + a^3B - 3ab^2B) \cot^2(c+dx)}{2d}$$

$$+ \frac{a(5a^2A - 12Ab^2 - 15abB) \cot^3(c+dx)}{15d} - \frac{a^2(7Ab + 5aB) \cot^4(c+dx)}{20d}$$

$$+ \frac{(3a^2Ab - Ab^3 + a^3B - 3ab^2B) \log(\sin(c+dx))}{d} - \frac{aA \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d}$$

output

```
-(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*x-(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*cot
(d*x+c)/d+1/2*(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*cot(d*x+c)^2/d+1/15*a*(5*A
*a^2-12*A*b^2-15*B*a*b)*cot(d*x+c)^3/d-1/20*a^2*(7*A*b+5*B*a)*cot(d*x+c)^4
/d+(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*ln(sin(d*x+c))/d-1/5*a*A*cot(d*x+c)^5
*(a+b*tan(d*x+c))^2/d
```

3.256.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.24 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.02

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{-60(a^3A - 3aAb^2 - 3a^2bB + b^3B) \cot(c + dx) + 30(3a^2Ab - Ab^3 + a^3B - 3ab^2B) \cot^2(c + dx) + 20a(a^2A - 3aAb + a^2B) \cot^3(c + dx) + 15a^2(3Ab - Ab^2 + aB) \cot^4(c + dx) + 12a^3A \cot^5(c + dx) + (30I)(a + Ib)^3(A + IB) \log[I - \tan(c + dx)] + 60(3a^2Ab - Ab^3 + a^3B - 3a^2bB) \log[\tan(c + dx)] + 30(Ia + b)^3(A - IB) \log[I + \tan(c + dx)]}{60d}$$

input `Integrate[Cot[c + d*x]^6*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output $(-60*(a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*\text{Cot}[c + d*x] + 30*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*\text{Cot}[c + d*x]^2 + 20*a*(a^2*A - 3*A*b^2 - 3*a*b*B)*\text{Cot}[c + d*x]^3 - 15*a^2*(3*A*b + a*B)*\text{Cot}[c + d*x]^4 - 12*a^3*A*\text{Cot}[c + d*x]^5 + (30*I)*(a + I*b)^3*(A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]] + 60*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*\text{Log}[\text{Tan}[c + d*x]] + 30*(I*a + b)^3*(A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]])/(60*d)$

3.256.3 Rubi [A] (verified)

Time = 1.57 (sec) , antiderivative size = 242, normalized size of antiderivative = 1.04, number of steps used = 18, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.581$, Rules used = {3042, 4088, 3042, 4118, 25, 3042, 4111, 27, 3042, 4012, 25, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^3(A + B \tan(c + dx))}{\tan(c + dx)^6} dx$$

$$\downarrow \text{4088}$$

$$\frac{1}{5} \int \cot^5(c+dx)(a+b \tan(c+dx)) \left(-b(3aA-5bB) \tan^2(c+dx) - 5(Aa^2-2bBa-Ab^2) \tan(c+dx) + a(7Ab+5aB) \right) dx - \frac{aA \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d}$$

↓ 3042

$$\frac{1}{5} \int \frac{(a+b \tan(c+dx)) \left(-b(3aA-5bB) \tan^2(c+dx) - 5(Aa^2-2bBa-Ab^2) \tan(c+dx) + a(7Ab+5aB) \right)}{\tan(c+dx)^5} dx - \frac{aA \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d}$$

↓ 4118

$$\frac{1}{5} \left(\int -\cot^4(c+dx) \left(b^2(3aA-5bB) \tan^2(c+dx) + 5(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx) + a(5Aa^2-15Ba^2-12Ab^2) \right) dx + \frac{aA \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d} \right)$$

↓ 25

$$\frac{1}{5} \left(- \int \cot^4(c+dx) \left(b^2(3aA-5bB) \tan^2(c+dx) + 5(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx) + a(5Aa^2-15Ba^2-12Ab^2) \right) dx + \frac{aA \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(- \int \frac{b^2(3aA-5bB) \tan^2(c+dx) + 5(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx) + a(5Aa^2-15Ba^2-12Ab^2)}{\tan(c+dx)^4} dx + \frac{aA \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d} \right)$$

↓ 4111

$$\frac{1}{5} \left(- \int 5 \cot^3(c+dx) \left(Ba^3+3Aba^2-3b^2Ba-Ab^3 - (Aa^3-3bBa^2-3Ab^2a+b^3B) \tan(c+dx) \right) dx + \frac{a(5a^2A+5Ba^2+5Ab^2)}{5d} + \frac{aA \cot^5(c+dx)(a+b \tan(c+dx))^2}{5d} \right)$$

↓ 27

$$\frac{1}{5} \left(-5 \int \cot^3(c+dx) (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx)) dx + \frac{a(5a^2A}{5d} \cot^5(c+dx)(a+b \tan(c+dx))^2 \right)$$

↓ 3042

$$\frac{1}{5} \left(-5 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx)}{\tan(c+dx)^3} dx + \frac{a(5a^2A - 15abB - 12}{3d} \cot^5(c+dx)(a+b \tan(c+dx))^2 \right)$$

↓ 4012

$$\frac{1}{5} \left(-5 \left(\int -\cot^2(c+dx) (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)) dx - \frac{(a^3}{5d} \cot^5(c+dx)(a+b \tan(c+dx))^2 \right) \right)$$

↓ 25

$$\frac{1}{5} \left(-5 \left(- \int \cot^2(c+dx) (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)) dx - \frac{(a^3}{5d} \cot^5(c+dx)(a+b \tan(c+dx))^2 \right) \right)$$

↓ 3042

$$\frac{1}{5} \left(-5 \left(- \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)}{\tan(c+dx)^2} dx - \frac{(a^3B + 3a^2Ab - 3}{5d} \cot^5(c+dx)(a+b \tan(c+dx))^2 \right) \right)$$

↓ 4012

$$\frac{1}{5} \left(-5 \left(- \int \cot(c+dx) (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx)) dx - \frac{(a^3E}{5d} \cot^5(c+dx)(a+b \tan(c+dx))^2 \right) \right)$$

↓ 3042

3.256. $\int \cot^6(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\frac{1}{5} \left(-5 \left(- \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\tan(c + dx)} dx - \frac{(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \cot^2(c + dx)}{2d} + \frac{(a^3A - 3a^2Ab + 3ab^2A - Ab^3) \cot^2(c + dx)}{2d} \right) \right. \\ \left. \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^2}{5d} \right) \downarrow 4014$$

$$\frac{1}{5} \left(-5 \left(-(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \int \cot(c + dx) dx - \frac{(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \cot^2(c + dx)}{2d} + \frac{(a^3A - 3a^2Ab + 3ab^2A - Ab^3) \cot^2(c + dx)}{2d} \right) \right. \\ \left. \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^2}{5d} \right) \downarrow 3042$$

$$\frac{1}{5} \left(-5 \left(-(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - \frac{(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \cot^2(c + dx)}{2d} + \frac{(a^3A - 3a^2Ab + 3ab^2A - Ab^3) \cot^2(c + dx)}{2d} \right) \right. \\ \left. \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^2}{5d} \right) \downarrow 25$$

$$\frac{1}{5} \left(-5 \left((a^3B + 3a^2Ab - 3ab^2B - Ab^3) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - \frac{(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \cot^2(c + dx)}{2d} + \frac{(a^3A - 3a^2Ab + 3ab^2A - Ab^3) \cot^2(c + dx)}{2d} \right) \right. \\ \left. \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^2}{5d} \right) \downarrow 3956$$

$$\frac{1}{5} \left(\frac{a(5a^2A - 15abB - 12Ab^2) \cot^3(c + dx)}{3d} - \frac{a^2(5aB + 7Ab) \cot^4(c + dx)}{4d} - 5 \left(- \frac{(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \cot^2(c + dx)}{2d} + \frac{(a^3A - 3a^2Ab + 3ab^2A - Ab^3) \cot^2(c + dx)}{2d} \right) \right. \\ \left. \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^2}{5d} \right)$$

input `Int[Cot[c + d*x]^6*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `((a*(5*a^2*A - 12*A*b^2 - 15*a*b*B)*Cot[c + d*x]^3)/(3*d) - (a^2*(7*A*b + 5*a*B)*Cot[c + d*x]^4)/(4*d) - 5*((a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*x + ((a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*Cot[c + d*x])/d - ((3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*Cot[c + d*x]^2)/(2*d) - ((3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*Log[-Sin[c + d*x]])/d)/5 - (a*A*Cot[c + d*x]^5*(a + b*Tan[c + d*x])^2)/(5*d)`

3.256. $\int \cot^6(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.256.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`
- rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`
- rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

```
rule 4111 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]
```

```
rule 4118 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.
)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-(b*c - a*d))*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]
```

3.256.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.04

method	result
parallelrisch	$(-90A^2b + 30Ab^3 - 30Ba^3 + 90Ba^2b^2) \ln(\sec^2(dx+c)) + (180A^2b - 60Ab^3 + 60Ba^3 - 180Ba^2b^2) \ln(\tan(dx+c)) - 12A^3$
derivativedivides	$\frac{(-3A^2b + Ab^3 - Ba^3 + 3Ba^2b^2) \ln(1 + \tan^2(dx+c))}{2} + (-A^3 + 3Aa^2b + 3Ba^2b - Bb^3) \arctan(\tan(dx+c)) - \frac{-3A^2b + Ab^3 - B$
default	$\frac{(-3A^2b + Ab^3 - Ba^3 + 3Ba^2b^2) \ln(1 + \tan^2(dx+c))}{2} + (-A^3 + 3Aa^2b + 3Ba^2b - Bb^3) \arctan(\tan(dx+c)) - \frac{-3A^2b + Ab^3 - B$
norman	$\frac{(-A^3 + 3Aa^2b + 3Ba^2b - Bb^3)x(\tan^5(dx+c)) - \frac{Aa^3}{5d} + \frac{(3A^2b - Ab^3 + Ba^3 - 3Ba^2b^2)(\tan^3(dx+c))}{2d}}{\tan(dx+c)^5} - \frac{(A^3 - 3Aa^2b - 3Ba^2$
risch	$-A^3x + 3Aa^2b^2x + 3Ba^2bx - Bb^3x + \frac{2iAb^3c}{d} - 3iA^2bx - iBa^3x + 3iBa^2bx - \frac{2i(-$

```
input int(cot(d*x+c)^6*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

3.256. $\int \cot^6(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

output $1/60*((-90*A*a^2*b+30*A*b^3-30*B*a^3+90*B*a*b^2)*\ln(\sec(d*x+c)^2)+(180*A*a^2*b-60*A*b^3+60*B*a^3-180*B*a*b^2)*\ln(\tan(d*x+c))-12*A*\cot(d*x+c)^5*a^3+(-45*A*a^2*b-15*B*a^3)*\cot(d*x+c)^4+20*a*\cot(d*x+c)^3*(A*a^2-3*A*b^2-3*B*a*b)+(90*A*a^2*b-30*A*b^3+30*B*a^3-90*B*a*b^2)*\cot(d*x+c)^2+(-60*A*a^3+180*A*a*b^2+180*B*a^2*b-60*B*b^3)*\cot(d*x+c)-60*d*x*(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3))/d$

3.256.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.14

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{30(Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^5 + 15(3Ba^3 + 9Aa^2b - 6Bab^2 - 2Ab^3)}{d}$$

input `integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output $1/60*(30*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*\log(\tan(d*x + c)^2/(\tan(d*x + c)^2 + 1))*\tan(d*x + c)^5 + 15*(3*B*a^3 + 9*A*a^2*b - 6*B*a*b^2 - 2*A*b^3 - 4*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*d*x)*\tan(d*x + c)^5 - 60*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*\tan(d*x + c)^4 - 12*A*a^3 + 30*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*\tan(d*x + c)^3 + 20*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2)*\tan(d*x + c)^2 - 15*(B*a^3 + 3*A*a^2*b)*\tan(d*x + c))/(d*\tan(d*x + c)^5)$

3.256.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 471 vs. 2(231) = 462.

Time = 4.17 (sec) , antiderivative size = 471, normalized size of antiderivative = 2.02

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \tilde{\infty} Aa^3x \\ x(A + B \tan(c))(a + b \tan(c))^3 \cot^6(c) \\ \tilde{\infty} Aa^3x \\ -Aa^3x - \frac{Aa^3}{d \tan(c+dx)} + \frac{Aa^3}{3d \tan^3(c+dx)} - \frac{Aa^3}{5d \tan^5(c+dx)} - \frac{3Aa^2b \log(\tan^2(c+dx)+1)}{2d} + \frac{3Aa^2b \log(\tan(c+dx))}{d} + \frac{3Aa^2b}{2d \tan^2(c+dx)} \end{cases}$$

3.256. $\int \cot^6(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

input `integrate(cot(d*x+c)**6*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `Piecewise((zoo*A*a**3*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**3*cot(c)**6, Eq(d, 0)), (zoo*A*a**3*x, Eq(c, -d*x)), (-A*a**3*x - A*a**3/(d*tan(c + d*x)) + A*a**3/(3*d*tan(c + d*x)**3) - A*a**3/(5*d*tan(c + d*x)**5) - 3*A*a**2*b*log(tan(c + d*x)**2 + 1)/(2*d) + 3*A*a**2*b*log(tan(c + d*x))/d + 3*A*a**2*b/(2*d*tan(c + d*x)**2) - 3*A*a**2*b/(4*d*tan(c + d*x)**4) + 3*A*a*b**2*x + 3*A*a*b**2/(d*tan(c + d*x)) - A*a*b**2/(d*tan(c + d*x)**3) + A*b**3*log(tan(c + d*x)**2 + 1)/(2*d) - A*b**3*log(tan(c + d*x))/d - A*b**3/(2*d*tan(c + d*x)**2) - B*a**3*log(tan(c + d*x)**2 + 1)/(2*d) + B*a**3*log(tan(c + d*x))/d + B*a**3/(2*d*tan(c + d*x)**2) - B*a**3/(4*d*tan(c + d*x)**4) + 3*B*a**2*b*x + 3*B*a**2*b/(d*tan(c + d*x)) - B*a**2*b/(d*tan(c + d*x)**3) + 3*B*a*b**2*log(tan(c + d*x)**2 + 1)/(2*d) - 3*B*a*b**2*log(tan(c + d*x))/d - 3*B*a*b**2/(2*d*tan(c + d*x)**2) - B*b**3*x - B*b**3/(d*tan(c + d*x)), True))`

3.256.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.07

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{60(Aa^3 - 3Ba^2b - 3Aab^2 + Bb^3)(dx + c) + 30(Ba^3 + 3Aa^2b - 3Bab^2 - Ab^3) \log(\tan(dx + c)^2 + 1)}{1}$$

input `integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/60*(60*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*(d*x + c) + 30*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*log(tan(d*x + c)^2 + 1) - 60*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*log(tan(d*x + c)) + (60*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*tan(d*x + c)^4 + 12*A*a^3 - 30*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*tan(d*x + c)^3 - 20*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2)*tan(d*x + c)^2 + 15*(B*a^3 + 3*A*a^2*b)*tan(d*x + c))/tan(d*x + c)^5/d`

3.256.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 670 vs. $2(225) = 450$.

Time = 1.12 (sec) , antiderivative size = 670, normalized size of antiderivative = 2.88

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{6 A a^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5 - 15 B a^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 - 45 A a^2 b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 - 70 A a^3 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3}{\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^5}$$

input `integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

```
1/960*(6*A*a^3*tan(1/2*d*x + 1/2*c)^5 - 15*B*a^3*tan(1/2*d*x + 1/2*c)^4 -
45*A*a^2*b*tan(1/2*d*x + 1/2*c)^4 - 70*A*a^3*tan(1/2*d*x + 1/2*c)^3 + 120*
B*a^2*b*tan(1/2*d*x + 1/2*c)^3 + 120*A*a*b^2*tan(1/2*d*x + 1/2*c)^3 + 180*
B*a^3*tan(1/2*d*x + 1/2*c)^2 + 540*A*a^2*b*tan(1/2*d*x + 1/2*c)^2 - 360*B*
a*b^2*tan(1/2*d*x + 1/2*c)^2 - 120*A*b^3*tan(1/2*d*x + 1/2*c)^2 + 660*A*a^
3*tan(1/2*d*x + 1/2*c) - 1800*B*a^2*b*tan(1/2*d*x + 1/2*c) - 1800*A*a*b^2*
tan(1/2*d*x + 1/2*c) + 480*B*b^3*tan(1/2*d*x + 1/2*c) - 960*(A*a^3 - 3*B*a
^2*b - 3*A*a*b^2 + B*b^3)*(d*x + c) - 960*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 -
A*b^3)*log(tan(1/2*d*x + 1/2*c)^2 + 1) + 960*(B*a^3 + 3*A*a^2*b - 3*B*a*b
^2 - A*b^3)*log(abs(tan(1/2*d*x + 1/2*c))) - (2192*B*a^3*tan(1/2*d*x + 1/2
*c)^5 + 6576*A*a^2*b*tan(1/2*d*x + 1/2*c)^5 - 6576*B*a*b^2*tan(1/2*d*x + 1
/2*c)^5 - 2192*A*b^3*tan(1/2*d*x + 1/2*c)^5 + 660*A*a^3*tan(1/2*d*x + 1/2*
c)^4 - 1800*B*a^2*b*tan(1/2*d*x + 1/2*c)^4 - 1800*A*a*b^2*tan(1/2*d*x + 1/
2*c)^4 + 480*B*b^3*tan(1/2*d*x + 1/2*c)^4 - 180*B*a^3*tan(1/2*d*x + 1/2*c)
^3 - 540*A*a^2*b*tan(1/2*d*x + 1/2*c)^3 + 360*B*a*b^2*tan(1/2*d*x + 1/2*c)
^3 + 120*A*b^3*tan(1/2*d*x + 1/2*c)^3 - 70*A*a^3*tan(1/2*d*x + 1/2*c)^2 +
120*B*a^2*b*tan(1/2*d*x + 1/2*c)^2 + 120*A*a*b^2*tan(1/2*d*x + 1/2*c)^2 +
15*B*a^3*tan(1/2*d*x + 1/2*c) + 45*A*a^2*b*tan(1/2*d*x + 1/2*c) + 6*A*a^3)
/tan(1/2*d*x + 1/2*c)^5)/d
```

3.256.9 Mupad [B] (verification not implemented)

Time = 8.43 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.02

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx =$$

$$\frac{\cot(c + dx)^5 \left(\tan(c + dx) \left(\frac{Ba^3}{4} + \frac{3Aba^2}{4} \right) + \frac{Aa^3}{5} + \tan(c + dx)^2 \left(-\frac{Aa^3}{3} + Ba^2b + Aab^2 \right) + \tan(c + dx) \left(\frac{Aa^2b}{2} + \frac{Aab^2}{2} \right) \right)}{d} + \frac{\ln(\tan(c + dx)) (-Ba^3 - 3Aa^2b + 3Bab^2 + Ab^3)}{d}$$

$$+ \frac{\ln(\tan(c + dx) - i) (A + B i) (a + b i)^3 i}{2d}$$

$$- \frac{\ln(\tan(c + dx) + i) (A - B i) (a - b i)^3 i}{2d}$$

input `int(cot(c + d*x)^6*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`output `(log(tan(c + d*x) - 1i)*(A + B*1i)*(a + b*1i)^3*1i)/(2*d) - (log(tan(c + d*x))*(A*b^3 - B*a^3 - 3*A*a^2*b + 3*B*a*b^2))/d - (cot(c + d*x)^5*(tan(c + d*x)*((B*a^3)/4 + (3*A*a^2*b)/4) + (A*a^3)/5 + tan(c + d*x)^2*(A*a*b^2 - (A*a^3)/3 + B*a^2*b) + tan(c + d*x)^4*(A*a^3 + B*b^3 - 3*A*a*b^2 - 3*B*a^2*b) + tan(c + d*x)^3*((A*b^3)/2 - (B*a^3)/2 - (3*A*a^2*b)/2 + (3*B*a*b^2)/2))/d - (log(tan(c + d*x) + 1i)*(A - B*1i)*(a - b*1i)^3*1i)/(2*d)`

3.257 $\int \tan^2(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.257.1 Optimal result	2490
3.257.2 Mathematica [C] (verified)	2491
3.257.3 Rubi [A] (verified)	2491
3.257.4 Maple [A] (verified)	2495
3.257.5 Fricas [A] (verification not implemented)	2496
3.257.6 Sympy [B] (verification not implemented)	2496
3.257.7 Maxima [A] (verification not implemented)	2497
3.257.8 Giac [B] (verification not implemented)	2498
3.257.9 Mupad [B] (verification not implemented)	2499

3.257.1 Optimal result

Integrand size = 31, antiderivative size = 263

$$\begin{aligned} & \int \tan^2(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx \\ &= -((a^4A - 6a^2Ab^2 + Ab^4 - 4a^3bB + 4ab^3B) x) \\ & \quad + \frac{(4a^3Ab - 4aAb^3 + a^4B - 6a^2b^2B + b^4B) \log(\cos(c+dx))}{d} \\ & \quad - \frac{b(3a^2Ab - Ab^3 + a^3B - 3ab^2B) \tan(c+dx)}{d} \\ & \quad - \frac{(2aAb + a^2B - b^2B) (a+b \tan(c+dx))^2}{2d} \\ & \quad - \frac{(Ab + aB)(a+b \tan(c+dx))^3}{3d} - \frac{B(a+b \tan(c+dx))^4}{4d} \\ & \quad + \frac{(6Ab - aB)(a+b \tan(c+dx))^5}{30b^2d} + \frac{B \tan(c+dx)(a+b \tan(c+dx))^5}{6bd} \end{aligned}$$

```
output - (A*a^4-6*A*a^2*b^2+A*b^4-4*B*a^3*b+4*B*a*b^3)*x+(4*A*a^3*b-4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*ln(cos(d*x+c))/d-b*(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*tan(d*x+c)/d-1/2*(2*A*a*b+B*a^2-B*b^2)*(a+b*tan(d*x+c))^2/d-1/3*(A*b+B*a)*(a+b*tan(d*x+c))^3/d-1/4*B*(a+b*tan(d*x+c))^4/d+1/30*(6*A*b-B*a)*(a+b*tan(d*x+c))^5/b^2/d+1/6*B*tan(d*x+c)*(a+b*tan(d*x+c))^5/b/d
```

3.257.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.10 (sec) , antiderivative size = 290, normalized size of antiderivative = 1.10

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{2(6Ab - aB)(a + b \tan(c + dx))^5}{b} + 10B \tan(c + dx)(a + b \tan(c + dx))^5 + 10(Ab - aB)(3i(a + ib)^4 \log(i - \tan(c + dx)) + \dots)$$

input `Integrate[Tan[c + d*x]^2*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `((2*(6*A*b - a*B)*(a + b*Tan[c + d*x])^5)/b + 10*B*Tan[c + d*x]*(a + b*Tan[c + d*x])^5 + 10*(A*b - a*B)*((3*I)*(a + I*b)^4*Log[I - Tan[c + d*x]] - (3*I)*(a - I*b)^4*Log[I + Tan[c + d*x]] + 6*b^2*(-6*a^2 + b^2)*Tan[c + d*x] - 12*a*b^3*Tan[c + d*x]^2 - 2*b^4*Tan[c + d*x]^3) + 5*B*((6*I)*(a + I*b)^5*Log[I - Tan[c + d*x]] - 6*(I*a + b)^5*Log[I + Tan[c + d*x]] - 60*a*b^2*(2*a^2 - b^2)*Tan[c + d*x] + 6*b^3*(-10*a^2 + b^2)*Tan[c + d*x]^2 - 20*a*b^4*Tan[c + d*x]^3 - 3*b^5*Tan[c + d*x]^4))/(60*b*d)`

3.257.3 Rubi [A] (verified)

Time = 1.37 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.05, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 4090, 25, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^2(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4090}$$

$$\begin{aligned}
& \frac{\int -(a + b \tan(c + dx))^4 \left(-((6Ab - aB) \tan^2(c + dx)) + 6bB \tan(c + dx) + aB \right) dx}{\frac{6b}{B \tan(c + dx)(a + b \tan(c + dx))^5} + \frac{6bd}{6bd}} \\
& \quad \downarrow 25 \\
& \frac{\int (a + b \tan(c + dx))^4 \left(-((6Ab - aB) \tan^2(c + dx)) + 6bB \tan(c + dx) + aB \right) dx}{\frac{6bd}{B \tan(c + dx)(a + b \tan(c + dx))^5} - \frac{6bd}{6bd}} \\
& \quad \downarrow 3042 \\
& \frac{\int (a + b \tan(c + dx))^4 \left(-((6Ab - aB) \tan^2(c + dx)^2) + 6bB \tan(c + dx) + aB \right) dx}{\frac{6bd}{B \tan(c + dx)(a + b \tan(c + dx))^5} - \frac{6bd}{6bd}} \\
& \quad \downarrow 4113 \\
& \frac{\int (a + b \tan(c + dx))^4 (6Ab + 6B \tan(c + dx)b) dx - \frac{(6Ab - aB)(a + b \tan(c + dx))^5}{5bd}}{\frac{6bd}{B \tan(c + dx)(a + b \tan(c + dx))^5} - \frac{6bd}{6bd}} \\
& \quad \downarrow 3042 \\
& \frac{\int (a + b \tan(c + dx))^4 (6Ab + 6B \tan(c + dx)b) dx - \frac{(6Ab - aB)(a + b \tan(c + dx))^5}{5bd}}{\frac{6bd}{B \tan(c + dx)(a + b \tan(c + dx))^5} - \frac{6bd}{6bd}} \\
& \quad \downarrow 4011 \\
& \frac{\int (a + b \tan(c + dx))^3 (6b(aA - bB) + 6b(Ab + aB) \tan(c + dx)) dx - \frac{(6Ab - aB)(a + b \tan(c + dx))^5}{5bd} + \frac{3bB(a + b \tan(c + dx))^4}{2d}}{\frac{6bd}{B \tan(c + dx)(a + b \tan(c + dx))^5} - \frac{6bd}{6bd}} \\
& \quad \downarrow 3042 \\
& \frac{\int (a + b \tan(c + dx))^3 (6b(aA - bB) + 6b(Ab + aB) \tan(c + dx)) dx - \frac{(6Ab - aB)(a + b \tan(c + dx))^5}{5bd} + \frac{3bB(a + b \tan(c + dx))^4}{2d}}{\frac{6bd}{B \tan(c + dx)(a + b \tan(c + dx))^5} - \frac{6bd}{6bd}} \\
& \quad \downarrow 4011 \\
& \frac{\int (a + b \tan(c + dx))^2 (6b(Aa^2 - 2bBa - Ab^2) + 6b(Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx - \frac{(6Ab - aB)(a + b \tan(c + dx))^5}{5bd}}{\frac{6bd}{B \tan(c + dx)(a + b \tan(c + dx))^5} - \frac{6bd}{6bd}}
\end{aligned}$$

3.257. $\int \tan^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{B \tan(c + dx)(a + b \tan(c + dx))^5}{6bd} - \\ & \frac{\int (a + b \tan(c + dx))^2 (6b(Aa^2 - 2bBa - Ab^2) + 6b(Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx - \frac{(6Ab - aB)(a + b \tan(c + dx))}{5bd}}{6b} \end{aligned}$$

$$\begin{aligned} & \downarrow 4011 \\ & \frac{B \tan(c + dx)(a + b \tan(c + dx))^5}{6bd} - \\ & \frac{\int (a + b \tan(c + dx)) (6b(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + 6b(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)) dx + \frac{3b}{6b}}{6b} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{B \tan(c + dx)(a + b \tan(c + dx))^5}{6bd} - \\ & \frac{\int (a + b \tan(c + dx)) (6b(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + 6b(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)) dx + \frac{3b}{6b}}{6b} \end{aligned}$$

$$\begin{aligned} & \downarrow 4008 \\ & \frac{B \tan(c + dx)(a + b \tan(c + dx))^5}{6bd} - \\ & \frac{6b(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) \int \tan(c + dx) dx + \frac{3b(a^2B + 2aAb - b^2B)(a + b \tan(c + dx))^2}{d} + \frac{6b^2(a^3B + 3a^2Ab - 3Ab^3)}{d}}{6b} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{B \tan(c + dx)(a + b \tan(c + dx))^5}{6bd} - \\ & \frac{6b(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) \int \tan(c + dx) dx + \frac{3b(a^2B + 2aAb - b^2B)(a + b \tan(c + dx))^2}{d} + \frac{6b^2(a^3B + 3a^2Ab - 3Ab^3)}{d}}{6b} \end{aligned}$$

$$\begin{aligned} & \downarrow 3956 \\ & \frac{B \tan(c + dx)(a + b \tan(c + dx))^5}{6bd} - \\ & \frac{\frac{3b(a^2B + 2aAb - b^2B)(a + b \tan(c + dx))^2}{d} + \frac{6b^2(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \tan(c + dx)}{d} - \frac{6b(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) \log(\cos(c + dx))}{d}}{6b} \end{aligned}$$

input `Int[Tan[c + d*x]^2*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

```
output (B*Tan[c + d*x]*(a + b*Tan[c + d*x])^5)/(6*b*d) - (6*b*(a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*a^3*b*B + 4*a*b^3*B)*x - (6*b*(4*a^3*A*b - 4*a*A*b^3 + a^4*B - 6*a^2*b^2*B + b^4*B)*Log[Cos[c + d*x]])/d + (6*b^2*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*Tan[c + d*x])/d + (3*b*(2*a*A*b + a^2*B - b^2*B)*(a + b*Tan[c + d*x])^2)/d + (2*b*(A*b + a*B)*(a + b*Tan[c + d*x])^3)/d + (3*b*B*(a + b*Tan[c + d*x])^4)/(2*d) - ((6*A*b - a*B)*(a + b*Tan[c + d*x])^5)/(5*b*d))/(6*b)
```

3.257.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4008 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]
```

```
rule 4011 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]
```

```
rule 4090 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4113 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

3.257.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 274, normalized size of antiderivative = 1.04

method	result
parts	$\frac{(Ab^4+4Bab^3)\left(\frac{\tan^5(dx+c)}{5}-\frac{\tan^3(dx+c)}{3}+\tan(dx+c)-\arctan(\tan(dx+c))\right)}{d} + \frac{(4Aab^3+6Ba^2b^2)\left(\frac{\tan^4(dx+c)}{4}\right)}{d}$
norman	$(-Aa^4 + 6Aa^2b^2 - Ab^4 + 4Ba^3b - 4Bab^3)x + \frac{(Aa^4 - 6Aa^2b^2 + Ab^4 - 4Ba^3b + 4Bab^3)\tan(dx+c)}{d}$
derivativedivides	$\frac{Bb^4(\tan^6(dx+c))}{6} + \frac{Ab^4(\tan^5(dx+c))}{5} + \frac{4Ba^3b^3(\tan^5(dx+c))}{5} + Aab^3(\tan^4(dx+c)) + \frac{3Ba^2b^2(\tan^4(dx+c))}{2} - \frac{Bb^4(\tan^4(dx+c))}{4}$
default	$\frac{Bb^4(\tan^6(dx+c))}{6} + \frac{Ab^4(\tan^5(dx+c))}{5} + \frac{4Ba^3b^3(\tan^5(dx+c))}{5} + Aab^3(\tan^4(dx+c)) + \frac{3Ba^2b^2(\tan^4(dx+c))}{2} - \frac{Bb^4(\tan^4(dx+c))}{4}$
parallelrisc	$-60A^4b^4 \tan(dx+c) + 360A^2a^2b^2 \tan(dx+c) + 240Ba^3b \tan(dx+c) - 240Ba^3b^3 \tan(dx+c) + 20A^4b^4 (\tan^3(dx+c)) - 30B$
risc	Expression too large to display

```
input int(tan(d*x+c)^2*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

3.257. $\int \tan^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

output $(A*b^4+4*B*a*b^3)/d*(1/5*\tan(dx+c)^5-1/3*\tan(dx+c)^3+\tan(dx+c)-\arctan(\tan(dx+c)))+(4*A*a*b^3+6*B*a^2*b^2)/d*(1/4*\tan(dx+c)^4-1/2*\tan(dx+c)^2+1/2*\ln(1+\tan(dx+c)^2))+(6*A*a^2*b^2+4*B*a^3*b)/d*(1/3*\tan(dx+c)^3-\tan(dx+c)+\arctan(\tan(dx+c)))+(4*A*a^3*b+B*a^4)/d*(1/2*\tan(dx+c)^2-1/2*\ln(1+\tan(dx+c)^2))+A*a^4/d*(\tan(dx+c)-\arctan(\tan(dx+c)))+B*b^4/d*(1/6*\tan(dx+c)^6-1/4*\tan(dx+c)^4+1/2*\tan(dx+c)^2-1/2*\ln(1+\tan(dx+c)^2))$

3.257.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.10

$$\int \tan^2(c+dx)(a+b\tan(c+dx))^4(A+B\tan(c+dx))dx$$

$$= \frac{10Bb^4\tan(dx+c)^6 + 12(4Bab^3 + Ab^4)\tan(dx+c)^5 + 15(6Ba^2b^2 + 4Aab^3 - Bb^4)\tan(dx+c)^4 + 20$$

input `integrate(tan(dx+c)^2*(a+b*tan(dx+c))^4*(A+B*tan(dx+c)),x, algorithm="fracas")`

output $1/60*(10*B*b^4*\tan(dx+c)^6 + 12*(4*B*a*b^3 + A*b^4)*\tan(dx+c)^5 + 15*(6*B*a^2*b^2 + 4*A*a*b^3 - B*b^4)*\tan(dx+c)^4 + 20*(4*B*a^3*b + 6*A*a^2*b^2 - 4*B*a*b^3 - A*b^4)*\tan(dx+c)^3 - 60*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*dx + 30*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*\tan(dx+c)^2 + 30*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*\log(1/(\tan(dx+c)^2 + 1)) + 60*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*\tan(dx+c))/d$

3.257.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 536 vs. $2(248) = 496$.

Time = 0.25 (sec) , antiderivative size = 536, normalized size of antiderivative = 2.04

$$\int \tan^2(c+dx)(a+b\tan(c+dx))^4(A+B\tan(c+dx))dx$$

$$= \begin{cases} -Aa^4x + \frac{Aa^4\tan(c+dx)}{d} - \frac{2Aa^3b\log(\tan^2(c+dx)+1)}{d} + \frac{2Aa^3b\tan^2(c+dx)}{d} + 6Aa^2b^2x + \frac{2Aa^2b^2\tan^3(c+dx)}{d} - \frac{6Aa^2b^2\tan^5(c+dx)}{d} \\ x(A+B\tan(c))(a+b\tan(c))^4\tan^2(c) \end{cases}$$

3.257. $\int \tan^2(c+dx)(a+b\tan(c+dx))^4(A+B\tan(c+dx))dx$

input `integrate(tan(d*x+c)**2*(a+b*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)`

output `Piecewise((-A*a**4*x + A*a**4*tan(c + d*x)/d - 2*A*a**3*b*log(tan(c + d*x)**2 + 1)/d + 2*A*a**3*b*tan(c + d*x)**2/d + 6*A*a**2*b**2*x + 2*A*a**2*b**2*tan(c + d*x)**3/d - 6*A*a**2*b**2*tan(c + d*x)/d + 2*A*a*b**3*log(tan(c + d*x)**2 + 1)/d + A*a*b**3*tan(c + d*x)**4/d - 2*A*a*b**3*tan(c + d*x)**2/d - A*b**4*x + A*b**4*tan(c + d*x)**5/(5*d) - A*b**4*tan(c + d*x)**3/(3*d) + A*b**4*tan(c + d*x)/d - B*a**4*log(tan(c + d*x)**2 + 1)/(2*d) + B*a**4*tan(c + d*x)**2/(2*d) + 4*B*a**3*b*x + 4*B*a**3*b*tan(c + d*x)**3/(3*d) - 4*B*a**3*b*tan(c + d*x)/d + 3*B*a**2*b**2*log(tan(c + d*x)**2 + 1)/d + 3*B*a**2*b**2*tan(c + d*x)**4/(2*d) - 3*B*a**2*b**2*tan(c + d*x)**2/d - 4*B*a*b**3*x + 4*B*a*b**3*tan(c + d*x)**5/(5*d) - 4*B*a*b**3*tan(c + d*x)**3/(3*d) + 4*B*a*b**3*tan(c + d*x)/d - B*b**4*log(tan(c + d*x)**2 + 1)/(2*d) + B*b**4*tan(c + d*x)**6/(6*d) - B*b**4*tan(c + d*x)**4/(4*d) + B*b**4*tan(c + d*x)**2/(2*d), Ne(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**4*tan(c)**2, True))`

3.257.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 290, normalized size of antiderivative = 1.10

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{10 B b^4 \tan(dx + c)^6 + 12 (4 B a b^3 + A b^4) \tan(dx + c)^5 + 15 (6 B a^2 b^2 + 4 A a b^3 - B b^4) \tan(dx + c)^4 + 20$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/60*(10*B*b^4*tan(d*x + c)^6 + 12*(4*B*a*b^3 + A*b^4)*tan(d*x + c)^5 + 15*(6*B*a^2*b^2 + 4*A*a*b^3 - B*b^4)*tan(d*x + c)^4 + 20*(4*B*a^3*b + 6*A*a^2*b^2 - 4*B*a*b^3 - A*b^4)*tan(d*x + c)^3 + 30*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*tan(d*x + c)^2 - 60*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*(d*x + c) - 30*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*log(tan(d*x + c)^2 + 1) + 60*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*tan(d*x + c))/d`

3.257. $\int \tan^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

3.257.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6042 vs. $2(252) = 504$.

Time = 7.74 (sec) , antiderivative size = 6042, normalized size of antiderivative = 22.97

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")
```

```
output -1/60*(60*A*a^4*d*x*tan(d*x)^6*tan(c)^6 - 240*B*a^3*b*d*x*tan(d*x)^6*tan(c)^6 - 360*A*a^2*b^2*d*x*tan(d*x)^6*tan(c)^6 + 240*B*a*b^3*d*x*tan(d*x)^6*tan(c)^6 + 60*A*b^4*d*x*tan(d*x)^6*tan(c)^6 - 30*B*a^4*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^6*tan(c)^6 - 120*A*a^3*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^6*tan(c)^6 + 180*B*a^2*b^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^6*tan(c)^6 + 120*A*a*b^3*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^6*tan(c)^6 - 30*B*b^4*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^6*tan(c)^6 - 360*A*a^4*d*x*tan(d*x)^5*tan(c)^5 + 1440*B*a^3*b*d*x*tan(d*x)^5*tan(c)^5 + 2160*A*a^2*b^2*d*x*tan(d*x)^5*tan(c)^5 - 1440*B*a*b^3*d*x*tan(d*x)^5*tan(c)^5 - 360*A*b^4*d*x*tan(d*x)^5*tan(c)^5 - 30*B*a^4*tan(d*x)^6*tan(c)^6 - 120*A*a^3*b*tan(d*x)^6*tan(c)^6 + 270*B*a^2*b^2*tan(d*x)^6*tan(c)^6 + 180*A*a*b^3*tan(d*x)^6*tan(c)^6 - 55*B*b^4*tan(d*x)^6*tan(c)^6 + 180*B*a^4*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^5*tan(c)^5 + 720*A*a^3*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + ...
```

3.257.9 Mupad [B] (verification not implemented)

Time = 7.98 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.14

$$\begin{aligned}
& \int \tan^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx \\
&= \frac{\tan(c + dx) (A a^4 + A b^4 + 4 B a b^3 - 2 a^2 b (3 A b + 2 B a))}{d} \\
&\quad - \frac{\tan(c + dx)^3 \left(\frac{A b^4}{3} + \frac{4 B a b^3}{3} - \frac{2 a^2 b (3 A b + 2 B a)}{3} \right)}{d} \\
&\quad - x (A a^4 - 4 B a^3 b - 6 A a^2 b^2 + 4 B a b^3 + A b^4) + \frac{\tan(c + dx)^5 \left(\frac{A b^4}{5} + \frac{4 B a b^3}{5} \right)}{d} \\
&\quad - \frac{\ln(\tan(c + dx)^2 + 1) \left(\frac{B a^4}{2} + 2 A a^3 b - 3 B a^2 b^2 - 2 A a b^3 + \frac{B b^4}{2} \right)}{d} \\
&\quad - \frac{\tan(c + dx)^4 \left(\frac{B b^4}{4} - \frac{a b^2 (2 A b + 3 B a)}{2} \right)}{d} \\
&\quad + \frac{\tan(c + dx)^2 \left(\frac{B a^4}{2} + \frac{B b^4}{2} + 2 A a^3 b - a b^2 (2 A b + 3 B a) \right)}{d} + \frac{B b^4 \tan(c + dx)^6}{6 d}
\end{aligned}$$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^4,x)`

```

output (tan(c + d*x)*(A*a^4 + A*b^4 + 4*B*a*b^3 - 2*a^2*b*(3*A*b + 2*B*a)))/d - (
tan(c + d*x)^3*((A*b^4)/3 + (4*B*a*b^3)/3 - (2*a^2*b*(3*A*b + 2*B*a))/3))/
d - x*(A*a^4 + A*b^4 - 6*A*a^2*b^2 + 4*B*a*b^3 - 4*B*a^3*b) + (tan(c + d*x
)^5*((A*b^4)/5 + (4*B*a*b^3)/5))/d - (log(tan(c + d*x)^2 + 1)*((B*a^4)/2 +
(B*b^4)/2 - 3*B*a^2*b^2 - 2*A*a*b^3 + 2*A*a^3*b))/d - (tan(c + d*x)^4*(B
*b^4)/4 - (a*b^2*(2*A*b + 3*B*a))/2))/d + (tan(c + d*x)^2*((B*a^4)/2 + (B
*b^4)/2 + 2*A*a^3*b - a*b^2*(2*A*b + 3*B*a)))/d + (B*b^4*tan(c + d*x)^6)/(6
*d)

```

3.258 $\int \tan(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.258.1 Optimal result	2500
3.258.2 Mathematica [C] (verified)	2501
3.258.3 Rubi [A] (verified)	2501
3.258.4 Maple [A] (verified)	2504
3.258.5 Fricas [A] (verification not implemented)	2505
3.258.6 Sympy [B] (verification not implemented)	2506
3.258.7 Maxima [A] (verification not implemented)	2507
3.258.8 Giac [B] (verification not implemented)	2507
3.258.9 Mupad [B] (verification not implemented)	2508

3.258.1 Optimal result

Integrand size = 29, antiderivative size = 226

$$\int \tan(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= -((4a^3Ab - 4aAb^3 + a^4B - 6a^2b^2B + b^4B) x) - \frac{(a^4A - 6a^2Ab^2 + Ab^4 - 4a^3bB + 4ab^3B) \log(\cos(c+dx))}{d} + \frac{b(a^3A - 3aAb^2 - 3a^2bB + b^3B) \tan(c+dx)}{d} + \frac{(a^2A - Ab^2 - 2abB)(a+b \tan(c+dx))^2}{2d} + \frac{(aA - bB)(a+b \tan(c+dx))^3}{3d} + \frac{A(a+b \tan(c+dx))^4}{4d} + \frac{B(a+b \tan(c+dx))^5}{5bd}$$

output

```
-(4*A*a^3*b-4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*x-(A*a^4-6*A*a^2*b^2+A*b^4-4*B*a^3*b+4*B*a*b^3)*ln(cos(d*x+c))/d+b*(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*tan(d*x+c)/d+1/2*(A*a^2-A*b^2-2*B*a*b)*(a+b*tan(d*x+c))^2/d+1/3*(A*a-B*b)*(a+b*tan(d*x+c))^3/d+1/4*A*(a+b*tan(d*x+c))^4/d+1/5*B*(a+b*tan(d*x+c))^5/b/d
```

3.258.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.54 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.14

$$\int \tan(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{12B(a + b \tan(c + dx))^5 + 10(aA + bB)(3i(a + ib)^4 \log(i - \tan(c + dx)) - 3i(a - ib)^4 \log(i + \tan(c + dx)))}{60bd}$$

input `Integrate[Tan[c + d*x]*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output $(12*B*(a + b*\text{Tan}[c + d*x])^5 + 10*(a*A + b*B)*((3*I)*(a + I*b)^4*\text{Log}[I - \text{Tan}[c + d*x]] - (3*I)*(a - I*b)^4*\text{Log}[I + \text{Tan}[c + d*x]] + 6*b^2*(-6*a^2 + b^2)*\text{Tan}[c + d*x] - 12*a*b^3*\text{Tan}[c + d*x]^2 - 2*b^4*\text{Tan}[c + d*x]^3) - 5*A*((6*I)*(a + I*b)^5*\text{Log}[I - \text{Tan}[c + d*x]] - 6*(I*a + b)^5*\text{Log}[I + \text{Tan}[c + d*x]] - 60*a*b^2*(2*a^2 - b^2)*\text{Tan}[c + d*x] + 6*b^3*(-10*a^2 + b^2)*\text{Tan}[c + d*x]^2 - 20*a*b^4*\text{Tan}[c + d*x]^3 - 3*b^5*\text{Tan}[c + d*x]^4))/(60*b*d)$

3.258.3 Rubi [A] (verified)

Time = 1.01 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$, Rules used = {3042, 4075, 3042, 4011, 3042, 4011, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \tan(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow 4075$$

$$\int (A \tan(c + dx) - B)(a + b \tan(c + dx))^4 dx + \frac{B(a + b \tan(c + dx))^5}{5bd}$$

$$\downarrow 3042$$

$$\begin{aligned}
& \int (A \tan(c + dx) - B)(a + b \tan(c + dx))^4 dx + \frac{B(a + b \tan(c + dx))^5}{5bd} \\
& \quad \downarrow \text{4011} \\
& \int (a + b \tan(c + dx))^3 (-Ab - aB + (aA - bB) \tan(c + dx)) dx + \frac{A(a + b \tan(c + dx))^4}{4d} + \\
& \quad \frac{B(a + b \tan(c + dx))^5}{5bd} \\
& \quad \downarrow \text{3042} \\
& \int (a + b \tan(c + dx))^3 (-Ab - aB + (aA - bB) \tan(c + dx)) dx + \frac{A(a + b \tan(c + dx))^4}{4d} + \\
& \quad \frac{B(a + b \tan(c + dx))^5}{5bd} \\
& \quad \downarrow \text{4011} \\
& \int (a + b \tan(c + dx))^2 (-Ba^2 - 2Aba + b^2B + (Aa^2 - 2bBa - Ab^2) \tan(c + dx)) dx + \\
& \quad \frac{(aA - bB)(a + b \tan(c + dx))^3}{3d} + \frac{A(a + b \tan(c + dx))^4}{4d} + \frac{B(a + b \tan(c + dx))^5}{5bd} \\
& \quad \downarrow \text{3042} \\
& \int (a + b \tan(c + dx))^2 (-Ba^2 - 2Aba + b^2B + (Aa^2 - 2bBa - Ab^2) \tan(c + dx)) dx + \\
& \quad \frac{(aA - bB)(a + b \tan(c + dx))^3}{3d} + \frac{A(a + b \tan(c + dx))^4}{4d} + \frac{B(a + b \tan(c + dx))^5}{5bd} \\
& \quad \downarrow \text{4011} \\
& \int (a + b \tan(c + dx)) (-Ba^3 - 3Aba^2 + 3b^2Ba + Ab^3 + (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)) dx + \\
& \quad \frac{(a^2A - 2abB - Ab^2)(a + b \tan(c + dx))^2}{2d} + \frac{(aA - bB)(a + b \tan(c + dx))^3}{3d} + \\
& \quad \frac{A(a + b \tan(c + dx))^4}{4d} + \frac{B(a + b \tan(c + dx))^5}{5bd} \\
& \quad \downarrow \text{3042} \\
& \int (a + b \tan(c + dx)) (-Ba^3 - 3Aba^2 + 3b^2Ba + Ab^3 + (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)) dx + \\
& \quad \frac{(a^2A - 2abB - Ab^2)(a + b \tan(c + dx))^2}{2d} + \frac{(aA - bB)(a + b \tan(c + dx))^3}{3d} + \\
& \quad \frac{A(a + b \tan(c + dx))^4}{4d} + \frac{B(a + b \tan(c + dx))^5}{5bd} \\
& \quad \downarrow \text{4008}
\end{aligned}$$

3.258. $\int \tan(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

$$\begin{aligned} & (a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) \int \tan(c + dx)dx + \\ & \frac{(a^2A - 2abB - Ab^2)(a + b \tan(c + dx))^2}{2d} + \frac{b(a^3A - 3a^2bB - 3aAb^2 + b^3B) \tan(c + dx)}{d} - \\ & x(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) + \frac{(aA - bB)(a + b \tan(c + dx))^3}{3d} + \\ & \frac{A(a + b \tan(c + dx))^4}{4d} + \frac{B(a + b \tan(c + dx))^5}{5bd} \end{aligned}$$

↓ 3042

$$\begin{aligned} & (a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) \int \tan(c + dx)dx + \\ & \frac{(a^2A - 2abB - Ab^2)(a + b \tan(c + dx))^2}{2d} + \frac{b(a^3A - 3a^2bB - 3aAb^2 + b^3B) \tan(c + dx)}{d} - \\ & x(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) + \frac{(aA - bB)(a + b \tan(c + dx))^3}{3d} + \\ & \frac{A(a + b \tan(c + dx))^4}{4d} + \frac{B(a + b \tan(c + dx))^5}{5bd} \end{aligned}$$

↓ 3956

$$\begin{aligned} & \frac{(a^2A - 2abB - Ab^2)(a + b \tan(c + dx))^2}{2d} + \frac{b(a^3A - 3a^2bB - 3aAb^2 + b^3B) \tan(c + dx)}{d} - \\ & \frac{(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) \log(\cos(c + dx))}{d} - \\ & x(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) + \frac{(aA - bB)(a + b \tan(c + dx))^3}{3d} + \\ & \frac{A(a + b \tan(c + dx))^4}{4d} + \frac{B(a + b \tan(c + dx))^5}{5bd} \end{aligned}$$

input `Int[Tan[c + d*x]*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `-((4*a^3*A*b - 4*a*A*b^3 + a^4*B - 6*a^2*b^2*B + b^4*B)*x) - ((a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*a^3*b*B + 4*a*b^3*B)*Log[Cos[c + d*x]])/d + (b*(a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*Tan[c + d*x])/d + ((a^2*A - A*b^2 - 2*a*b*B)*(a + b*Tan[c + d*x])^2)/(2*d) + ((a*A - b*B)*(a + b*Tan[c + d*x])^3)/(3*d) + (A*(a + b*Tan[c + d*x])^4)/(4*d) + (B*(a + b*Tan[c + d*x])^5)/(5*b*d)`

3.258.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.258.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 242, normalized size of antiderivative = 1.07

method	result
parts	$\frac{(Ab^4+4Bab^3)\left(\frac{\tan^4(dx+c)}{4}-\frac{\tan^2(dx+c)}{2}+\frac{\ln(1+\tan^2(dx+c))}{2}\right)}{d} + \frac{(4Aab^3+6Ba^2b^2)\left(\frac{\tan^3(dx+c)}{3}-\tan(dx+c)\right)}{d}$
norman	$(-4Aa^3b+4Aab^3-Ba^4+6Ba^2b^2-Bb^4)x + \frac{(4Aa^3b-4Aab^3+Ba^4-6Ba^2b^2+Bb^4)\tan(dx+c)}{d}$
derivativedivides	$\frac{Bb^4\left(\frac{\tan^5(dx+c)}{5}+\frac{Ab^4\left(\frac{\tan^4(dx+c)}{4}+Ba^3\left(\tan^4(dx+c)\right)+\frac{4Aab^3\left(\frac{\tan^3(dx+c)}{3}+2Ba^2b^2\left(\tan^3(dx+c)\right)-\frac{Bb^4\left(\frac{\tan^3(dx+c)}{3}\right)}{d}\right)}{d}\right)}{d}$
default	$\frac{Bb^4\left(\frac{\tan^5(dx+c)}{5}+\frac{Ab^4\left(\frac{\tan^4(dx+c)}{4}+Ba^3\left(\tan^4(dx+c)\right)+\frac{4Aab^3\left(\frac{\tan^3(dx+c)}{3}+2Ba^2b^2\left(\tan^3(dx+c)\right)-\frac{Bb^4\left(\frac{\tan^3(dx+c)}{3}\right)}{d}\right)}{d}\right)}{d}$
parallelrisch	$60Bb^4\tan(dx+c)-60Bxa^4d+360Ba^2b^2dx-240Aa^3bdx+240Aab^3dx+240Aa^3b\tan(dx+c)-240Aab^3\tan(dx+c)-360Bb^4$
risch	$-\frac{12iAa^2b^2c}{d} + \frac{8iBab^3c}{d} + \frac{2i(60Aa^3b-80Aab^3-120Ba^2b^2+15Ba^4+23Bb^4-270iAa^2b^2e^{6i(dx+c)}-180iBa^3be^{6i(dx+c)})}{d}$

input `int(tan(d*x+c)*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output $(A*b^4+4*B*a*b^3)/d*(1/4*\tan(d*x+c)^4-1/2*\tan(d*x+c)^2+1/2*\ln(1+\tan(d*x+c)^2))+ (4*A*a*b^3+6*B*a^2*b^2)/d*(1/3*\tan(d*x+c)^3-\tan(d*x+c)+\arctan(\tan(d*x+c)))+ (6*A*a^2*b^2+4*B*a^3*b)/d*(1/2*\tan(d*x+c)^2-1/2*\ln(1+\tan(d*x+c)^2))+ (4*A*a^3*b+B*a^4)/d*(\tan(d*x+c)-\arctan(\tan(d*x+c)))+ 1/2/d*A*\ln(1+\tan(d*x+c)^2)*a^4+B*b^4/d*(1/5*\tan(d*x+c)^5-1/3*\tan(d*x+c)^3+\tan(d*x+c)-\arctan(\tan(d*x+c)))$

3.258.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.08

$$\int \tan(c+dx)(a+b\tan(c+dx))^4(A+B\tan(c+dx))dx$$

$$= \frac{12Bb^4\tan(dx+c)^5+15(4Bab^3+Ab^4)\tan(dx+c)^4+20(6Ba^2b^2+4Aab^3-Bb^4)\tan(dx+c)^3-60Aa^4b^2\tan(dx+c)^2+60Aa^4b^2}{d}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fricas")`


```
output 1/60*(12*B*b^4*tan(d*x + c)^5 + 15*(4*B*a*b^3 + A*b^4)*tan(d*x + c)^4 + 20
*(6*B*a^2*b^2 + 4*A*a*b^3 - B*b^4)*tan(d*x + c)^3 - 60*(B*a^4 + 4*A*a^3*b
- 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*d*x + 30*(4*B*a^3*b + 6*A*a^2*b^2 - 4*B
*a*b^3 - A*b^4)*tan(d*x + c)^2 - 30*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B
*a*b^3 + A*b^4)*log(1/(tan(d*x + c)^2 + 1)) + 60*(B*a^4 + 4*A*a^3*b - 6*B
a^2*b^2 - 4*A*a*b^3 + B*b^4)*tan(d*x + c))/d
```

3.258.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 437 vs. $2(214) = 428$.

Time = 0.19 (sec) , antiderivative size = 437, normalized size of antiderivative = 1.93

$$\int \tan(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \frac{Aa^4 \log(\tan^2(c+dx)+1)}{2d} - 4Aa^3bx + \frac{4Aa^3b \tan(c+dx)}{d} - \frac{3Aa^2b^2 \log(\tan^2(c+dx)+1)}{d} + \frac{3Aa^2b^2 \tan^2(c+dx)}{d} + 4Aab^3x + \frac{4Aa^2b^2 \tan(c+dx)}{d} \\ x(A + B \tan(c))(a + b \tan(c))^4 \tan(c) \end{cases}$$

```
input integrate(tan(d*x+c)*(a+b*tan(d*x+c))**4*(A+B*tan(d*x+c)), x)
```

```
output Piecewise((A*a**4*log(tan(c + d*x)**2 + 1)/(2*d) - 4*A*a**3*b*x + 4*A*a**3
*b*tan(c + d*x)/d - 3*A*a**2*b**2*log(tan(c + d*x)**2 + 1)/d + 3*A*a**2*b*
**2*tan(c + d*x)**2/d + 4*A*a*b**3*x + 4*A*a*b**3*tan(c + d*x)**3/(3*d) - 4
*A*a*b**3*tan(c + d*x)/d + A*b**4*log(tan(c + d*x)**2 + 1)/(2*d) + A*b**4*
tan(c + d*x)**4/(4*d) - A*b**4*tan(c + d*x)**2/(2*d) - B*a**4*x + B*a**4*t
an(c + d*x)/d - 2*B*a**3*b*log(tan(c + d*x)**2 + 1)/d + 2*B*a**3*b*tan(c +
d*x)**2/d + 6*B*a**2*b**2*x + 2*B*a**2*b**2*tan(c + d*x)**3/d - 6*B*a**2*
b**2*tan(c + d*x)/d + 2*B*a*b**3*log(tan(c + d*x)**2 + 1)/d + B*a*b**3*tan
(c + d*x)**4/d - 2*B*a*b**3*tan(c + d*x)**2/d - B*b**4*x + B*b**4*tan(c +
d*x)**5/(5*d) - B*b**4*tan(c + d*x)**3/(3*d) + B*b**4*tan(c + d*x)/d, Ne(d
, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**4*tan(c), True))
```

3.258.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.09

$$\int \tan(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{12 B b^4 \tan(dx + c)^5 + 15(4 B a b^3 + A b^4) \tan(dx + c)^4 + 20(6 B a^2 b^2 + 4 A a b^3 - B b^4) \tan(dx + c)^3 + 30$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/60*(12*B*b^4*tan(d*x + c)^5 + 15*(4*B*a*b^3 + A*b^4)*tan(d*x + c)^4 + 20*(6*B*a^2*b^2 + 4*A*a*b^3 - B*b^4)*tan(d*x + c)^3 + 30*(4*B*a^3*b + 6*A*a^2*b^2 - 4*B*a*b^3 - A*b^4)*tan(d*x + c)^2 - 60*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*(d*x + c) + 30*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)^2 + 1) + 60*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*tan(d*x + c))/d`

3.258.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4489 vs. 2(218) = 436.

Time = 4.84 (sec) , antiderivative size = 4489, normalized size of antiderivative = 19.86

$$\int \tan(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

```
-1/60*(60*B*a^4*d*x*tan(d*x)^5*tan(c)^5 + 240*A*a^3*b*d*x*tan(d*x)^5*tan(c)^5 - 360*B*a^2*b^2*d*x*tan(d*x)^5*tan(c)^5 - 240*A*a*b^3*d*x*tan(d*x)^5*tan(c)^5 + 60*B*b^4*d*x*tan(d*x)^5*tan(c)^5 + 30*A*a^4*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^5*tan(c)^5 - 120*B*a^3*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^5*tan(c)^5 - 180*A*a^2*b^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^5*tan(c)^5 + 120*B*a*b^3*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^5*tan(c)^5 + 30*A*b^4*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^5*tan(c)^5 - 300*B*a^4*d*x*tan(d*x)^4*tan(c)^4 - 1200*A*a^3*b*d*x*tan(d*x)^4*tan(c)^4 + 1800*B*a^2*b^2*d*x*tan(d*x)^4*tan(c)^4 + 1200*A*a*b^3*d*x*tan(d*x)^4*tan(c)^4 - 300*B*b^4*d*x*tan(d*x)^4*tan(c)^4 - 120*B*a^3*b*tan(d*x)^5*tan(c)^5 - 180*A*a^2*b^2*tan(d*x)^5*tan(c)^5 + 180*B*a*b^3*tan(d*x)^5*tan(c)^5 + 45*A*b^4*tan(d*x)^5*tan(c)^5 - 150*A*a^4*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^4*tan(c)^4 + 600*B*a^3*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^4*tan(c)^4 + 900*A...
```

3.258.9 Mupad [B] (verification not implemented)

Time = 8.52 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.11

$$\int \tan(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{\tan(c + dx) (B a^4 + B b^4 + 4 A a^3 b - 2 a b^2 (2 A b + 3 B a))}{d}$$

$$- \frac{\tan(c + dx)^2 \left(\frac{A b^4}{2} + 2 B a b^3 - a^2 b (3 A b + 2 B a) \right)}{d}$$

$$- x (B a^4 + 4 A a^3 b - 6 B a^2 b^2 - 4 A a b^3 + B b^4) + \frac{\tan(c + dx)^4 \left(\frac{A b^4}{4} + B a b^3 \right)}{d}$$

$$+ \frac{\ln(\tan(c + dx)^2 + 1) \left(\frac{A a^4}{2} - 2 B a^3 b - 3 A a^2 b^2 + 2 B a b^3 + \frac{A b^4}{2} \right)}{d}$$

$$- \frac{\tan(c + dx)^3 \left(\frac{B b^4}{3} - \frac{2 a b^2 (2 A b + 3 B a)}{3} \right)}{d} + \frac{B b^4 \tan(c + dx)^5}{5 d}$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^4,x)`

3.258. $\int \tan(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

output $(\tan(c + dx)(B^4a^4 + B^4b^4 + 4A^3a^3b - 2a^2b^2(2Ab + 3Ba)))/d - (\tan(c + dx)^2((A^4b^4)/2 + 2B^3a^3b^3 - a^2b^2(3Ab + 2Ba)))/d - x(B^4a^4 + B^4b^4 - 6B^2a^2b^2 - 4A^3a^3b^3 + 4A^3a^3b) + (\tan(c + dx)^4((A^4b^4)/4 + B^3a^3b^3))/d + (\log(\tan(c + dx)^2 + 1)((A^4a^4)/2 + (A^4b^4)/2 - 3A^3a^2b^2 + 2B^3a^3b^3 - 2B^3a^3b))/d - (\tan(c + dx)^3((B^4b^4)/3 - (2a^2b^2(2Ab + 3Ba))/3))/d + (B^4b^4 \tan(c + dx)^5)/(5d)$

3.259 $\int (a + b \tan(c + dx))^4 (A + B \tan(c + dx)) dx$

3.259.1 Optimal result	2510
3.259.2 Mathematica [C] (verified)	2511
3.259.3 Rubi [A] (verified)	2511
3.259.4 Maple [A] (verified)	2514
3.259.5 Fricas [A] (verification not implemented)	2514
3.259.6 Sympy [A] (verification not implemented)	2515
3.259.7 Maxima [A] (verification not implemented)	2515
3.259.8 Giac [B] (verification not implemented)	2516
3.259.9 Mupad [B] (verification not implemented)	2517

3.259.1 Optimal result

Integrand size = 23, antiderivative size = 202

$$\begin{aligned} & \int (a + b \tan(c + dx))^4 (A + B \tan(c + dx)) dx \\ &= (a^4 A - 6a^2 Ab^2 + Ab^4 - 4a^3 bB + 4ab^3 B) x \\ & \quad - \frac{(4a^3 Ab - 4aAb^3 + a^4 B - 6a^2 b^2 B + b^4 B) \log(\cos(c + dx))}{d} \\ & \quad + \frac{b(3a^2 Ab - Ab^3 + a^3 B - 3ab^2 B) \tan(c + dx)}{d} \\ & \quad + \frac{(2aAb + a^2 B - b^2 B) (a + b \tan(c + dx))^2}{2d} \\ & \quad + \frac{(Ab + aB) (a + b \tan(c + dx))^3}{3d} + \frac{B(a + b \tan(c + dx))^4}{4d} \end{aligned}$$

output

```
(A*a^4-6*A*a^2*b^2+Ab^4-4*B*a^3*b+4*B*a*b^3)*x-(4*A*a^3*b-4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*ln(cos(d*x+c))/d+b*(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*tan(d*x+c)/d+1/2*(2*A*a*b+B*a^2-B*b^2)*(a+b*tan(d*x+c))^2/d+1/3*(A*b+B*a)*(a+b*tan(d*x+c))^3/d+1/4*B*(a+b*tan(d*x+c))^4/d
```

3.259.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.99 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.19

$$\int (a + b \tan(c + dx))^4 (A + B \tan(c + dx)) dx$$

$$= \frac{-2(Ab - aB) (3i(a + ib)^4 \log(i - \tan(c + dx)) - 3i(a - ib)^4 \log(i + \tan(c + dx)) + 6b^2(-6a^2 + b^2) \tan(c + dx))}{12bd}$$

input `Integrate[(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output
$$\frac{(-2*(A*b - a*B)*((3*I)*(a + I*b)^4*\text{Log}[I - \text{Tan}[c + d*x]] - (3*I)*(a - I*b)^4*\text{Log}[I + \text{Tan}[c + d*x]] + 6*b^2*(-6*a^2 + b^2)*\text{Tan}[c + d*x] - 12*a*b^3*\text{Tan}[c + d*x]^2 - 2*b^4*\text{Tan}[c + d*x]^3) + B*(6*((-I)*a + b)^5*\text{Log}[I - \text{Tan}[c + d*x]] + 6*(I*a + b)^5*\text{Log}[I + \text{Tan}[c + d*x]] + 60*a*b^2*(2*a^2 - b^2)*\text{Tan}[c + d*x] - 6*b^3*(-10*a^2 + b^2)*\text{Tan}[c + d*x]^2 + 20*a*b^4*\text{Tan}[c + d*x]^3 + 3*b^5*\text{Tan}[c + d*x]^4))/(12*b*d)}$$

3.259.3 Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.435$, Rules used = {3042, 4011, 3042, 4011, 3042, 4011, 3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \tan(c + dx))^4 (A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int (a + b \tan(c + dx))^4 (A + B \tan(c + dx)) dx$$

$$\downarrow \text{4011}$$

$$\int (a + b \tan(c + dx))^3 (aA - bB + (Ab + aB) \tan(c + dx)) dx + \frac{B(a + b \tan(c + dx))^4}{4d}$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
& \int (a + b \tan(c + dx))^3 (aA - bB + (Ab + aB) \tan(c + dx)) dx + \frac{B(a + b \tan(c + dx))^4}{4d} \\
& \quad \downarrow 4011 \\
& \int (a + b \tan(c + dx))^2 (Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx + \\
& \quad \frac{(aB + Ab)(a + b \tan(c + dx))^3}{3d} + \frac{B(a + b \tan(c + dx))^4}{4d} \\
& \quad \downarrow 3042 \\
& \int (a + b \tan(c + dx))^2 (Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx + \\
& \quad \frac{(aB + Ab)(a + b \tan(c + dx))^3}{3d} + \frac{B(a + b \tan(c + dx))^4}{4d} \\
& \quad \downarrow 4011 \\
& \int (a + b \tan(c + dx)) (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)) dx + \\
& \quad \frac{(a^2B + 2aAb - b^2B)(a + b \tan(c + dx))^2}{2d} + \frac{(aB + Ab)(a + b \tan(c + dx))^3}{3d} + \\
& \quad \frac{B(a + b \tan(c + dx))^4}{4d} \\
& \quad \downarrow 3042 \\
& \int (a + b \tan(c + dx)) (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)) dx + \\
& \quad \frac{(a^2B + 2aAb - b^2B)(a + b \tan(c + dx))^2}{2d} + \frac{(aB + Ab)(a + b \tan(c + dx))^3}{3d} + \\
& \quad \frac{B(a + b \tan(c + dx))^4}{4d} \\
& \quad \downarrow 4008 \\
& \frac{(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) \int \tan(c + dx) dx + (a^2B + 2aAb - b^2B)(a + b \tan(c + dx))^2}{2d} + \frac{b(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \tan(c + dx)}{d} + \\
& \quad x(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) + \frac{(aB + Ab)(a + b \tan(c + dx))^3}{3d} + \\
& \quad \frac{B(a + b \tan(c + dx))^4}{4d} \\
& \quad \downarrow 3042
\end{aligned}$$

3.259. $\int (a + b \tan(c + dx))^4 (A + B \tan(c + dx)) dx$

$$\begin{aligned}
& (a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) \int \tan(c + dx) dx + \\
& \frac{(a^2B + 2aAb - b^2B)(a + b \tan(c + dx))^2}{2d} + \frac{b(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \tan(c + dx)}{d} + \\
& x(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) + \frac{(aB + Ab)(a + b \tan(c + dx))^3}{3d} + \\
& \frac{B(a + b \tan(c + dx))^4}{4d} \\
& \quad \downarrow \text{3956} \\
& \frac{(a^2B + 2aAb - b^2B)(a + b \tan(c + dx))^2}{2d} + \frac{b(a^3B + 3a^2Ab - 3ab^2B - Ab^3) \tan(c + dx)}{d} - \\
& \frac{(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) \log(\cos(c + dx))}{d} + \\
& x(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) + \frac{(aB + Ab)(a + b \tan(c + dx))^3}{3d} + \\
& \frac{B(a + b \tan(c + dx))^4}{4d}
\end{aligned}$$

input `Int[(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*a^3*b*B + 4*a*b^3*B)*x - ((4*a^3*A*b - 4*a*A*b^3 + a^4*B - 6*a^2*b^2*B + b^4*B)*Log[Cos[c + d*x]])/d + (b*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*Tan[c + d*x])/d + ((2*a*A*b + a^2*B - b^2*B)*(a + b*Tan[c + d*x])^2)/(2*d) + ((A*b + a*B)*(a + b*Tan[c + d*x])^3)/(3*d) + (B*(a + b*Tan[c + d*x])^4)/(4*d)`

3.259.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4008 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f), x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]`


```
rule 4011 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

3.259.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.00

method	result
norman	$(Aa^4 - 6Aa^2b^2 + Ab^4 - 4Ba^3b + 4Bab^3)x + \frac{b(6Aa^2b - Ab^3 + 4Ba^3 - 4Bab^2) \tan(dx+c)}{d} + \frac{Bb^4}{d}$
parts	$Aa^4x + \frac{(Ab^4 + 4Bab^3) \left(\frac{\tan^3(dx+c)}{3} - \tan(dx+c) + \arctan(\tan(dx+c)) \right)}{d} + \frac{(4Aab^3 + 6Ba^2b^2) \left(\frac{\tan^2(dx+c)}{2} - \tan(dx+c) \right)}{d}$
derivativedivides	$\frac{Bb^4 \frac{\tan^4(dx+c)}{4} + Ab^4 \frac{\tan^3(dx+c)}{3} + 4Bab^3 \frac{\tan^3(dx+c)}{3} + 2Aab^3 \tan^2(dx+c) + 3Ba^2b^2 \tan^2(dx+c) - \frac{Bb^4 \tan^2(dx+c)}{2}}{d}$
default	$\frac{Bb^4 \frac{\tan^4(dx+c)}{4} + Ab^4 \frac{\tan^3(dx+c)}{3} + 4Bab^3 \frac{\tan^3(dx+c)}{3} + 2Aab^3 \tan^2(dx+c) + 3Ba^2b^2 \tan^2(dx+c) - \frac{Bb^4 \tan^2(dx+c)}{2}}{d}$
parallelrisch	$\frac{3Bb^4 \tan^4(dx+c) + 4Aab^4 \tan^3(dx+c) + 16Bab^3 \tan^3(dx+c) + 12Aa^4d - 72Aa^2b^2dx + 12Ab^4dx + 24Aab^3 \tan^2(dx+c)}{d}$
risch	$-\frac{a^4 \ln(e^{2i(dx+c)} + 1)B}{d} - \frac{\ln(e^{2i(dx+c)} + 1)Bb^4}{d} + Aa^4x + \frac{8iAa^3bc}{d} + \frac{2ia^4Bc}{d} + iBb^4x - \frac{12iBa^2b^2c}{d} +$

```
input int((a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output (A*a^4-6*A*a^2*b^2+A*b^4-4*B*a^3*b+4*B*a*b^3)*x+b*(6*A*a^2*b-A*b^3+4*B*a^3
-4*B*a*b^2)/d*tan(d*x+c)+1/4*B*b^4/d*tan(d*x+c)^4+1/2*b^2*(4*A*a*b+6*B*a^2
-B*b^2)/d*tan(d*x+c)^2+1/3*b^3*(A*b+4*B*a)/d*tan(d*x+c)^3+1/2*(4*A*a^3*b-4
*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)/d*ln(1+tan(d*x+c)^2)
```

3.259.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.00

$$\int (a + b \tan(c + dx))^4 (A + B \tan(c + dx)) dx$$

$$= \frac{3Bb^4 \tan(dx+c)^4 + 4(4Bab^3 + Ab^4) \tan(dx+c)^3 + 12(Aa^4 - 4Ba^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4)dx + \dots}{d}$$

3.259. $\int (a + b \tan(c + dx))^4 (A + B \tan(c + dx)) dx$

input `integrate((a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output
$$\frac{1}{12} * (3 * B * b^4 * \tan(dx + c)^4 + 4 * (4 * B * a * b^3 + A * b^4) * \tan(dx + c)^3 + 12 * (A * a^4 - 4 * B * a^3 * b - 6 * A * a^2 * b^2 + 4 * B * a * b^3 + A * b^4) * dx + 6 * (6 * B * a^2 * b^2 + 4 * A * a * b^3 - B * b^4) * \tan(dx + c)^2 - 6 * (B * a^4 + 4 * A * a^3 * b - 6 * B * a^2 * b^2 - 4 * A * a * b^3 + B * b^4) * \log(1 / (\tan(dx + c)^2 + 1)) + 12 * (4 * B * a^3 * b + 6 * A * a^2 * b^2 - 4 * B * a * b^3 - A * b^4) * \tan(dx + c)) / d$$

3.259.6 Sympy [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 347, normalized size of antiderivative = 1.72

$$\int (a + b \tan(c + dx))^4 (A + B \tan(c + dx)) dx$$

$$= \begin{cases} Aa^4x + \frac{2Aa^3b \log(\tan^2(c+dx)+1)}{d} - 6Aa^2b^2x + \frac{6Aa^2b^2 \tan(c+dx)}{d} - \frac{2Aab^3 \log(\tan^2(c+dx)+1)}{d} + \frac{2Aab^3 \tan^2(c+dx)}{d} + Ab^4x \\ x(A + B \tan(c)) (a + b \tan(c))^4 \end{cases}$$

input `integrate((a+b*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)`

output `Piecewise((A*a**4*x + 2*A*a**3*b*log(tan(c + d*x)**2 + 1)/d - 6*A*a**2*b**2*x + 6*A*a**2*b**2*tan(c + d*x)/d - 2*A*a*b**3*log(tan(c + d*x)**2 + 1)/d + 2*A*a*b**3*tan(c + d*x)**2/d + A*b**4*x + A*b**4*tan(c + d*x)**3/(3*d) - A*b**4*tan(c + d*x)/d + B*a**4*log(tan(c + d*x)**2 + 1)/(2*d) - 4*B*a**3*b*x + 4*B*a**3*b*tan(c + d*x)/d - 3*B*a**2*b**2*log(tan(c + d*x)**2 + 1)/d + 3*B*a**2*b**2*tan(c + d*x)**2/d + 4*B*a*b**3*x + 4*B*a*b**3*tan(c + d*x)**3/(3*d) - 4*B*a*b**3*tan(c + d*x)/d + B*b**4*log(tan(c + d*x)**2 + 1)/(2*d) + B*b**4*tan(c + d*x)**4/(4*d) - B*b**4*tan(c + d*x)**2/(2*d), Ne(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**4, True))`

3.259.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.00

$$\int (a + b \tan(c + dx))^4 (A + B \tan(c + dx)) dx$$

$$= \frac{3Bb^4 \tan(dx + c)^4 + 4(4Bab^3 + Ab^4) \tan(dx + c)^3 + 6(6Ba^2b^2 + 4Aab^3 - Bb^4) \tan(dx + c)^2 + 12(Aa^4 + 4Aab^3 + 6Aa^2b^2 + 4Ab^4) \tan(dx + c) + 12Aa^4x + 12Aa^3b \log(\tan^2(dx + c) + 1) + 12Aa^2b^2 \tan(dx + c) + 12Aab^3 \tan^2(dx + c) + 12Ab^4 \tan^3(dx + c) + 12Bb^4 \log(\tan^2(dx + c) + 1) + 12Bb^4 \tan(dx + c)}{d}$$

```
input integrate((a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")
```

```
output 1/12*(3*B*b^4*tan(d*x + c)^4 + 4*(4*B*a*b^3 + A*b^4)*tan(d*x + c)^3 + 6*(6
*B*a^2*b^2 + 4*A*a*b^3 - B*b^4)*tan(d*x + c)^2 + 12*(A*a^4 - 4*B*a^3*b - 6
*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*(d*x + c) + 6*(B*a^4 + 4*A*a^3*b - 6*B*a^2
*b^2 - 4*A*a*b^3 + B*b^4)*log(tan(d*x + c)^2 + 1) + 12*(4*B*a^3*b + 6*A*a^
2*b^2 - 4*B*a*b^3 - A*b^4)*tan(d*x + c))/d
```

3.259.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3133 vs. $2(196) = 392$.

Time = 2.81 (sec) , antiderivative size = 3133, normalized size of antiderivative = 15.51

$$\int (a + b \tan(c + dx))^4 (A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")
```

```
output 1/12*(12*A*a^4*d*x*tan(d*x)^4*tan(c)^4 - 48*B*a^3*b*d*x*tan(d*x)^4*tan(c)^
4 - 72*A*a^2*b^2*d*x*tan(d*x)^4*tan(c)^4 + 48*B*a*b^3*d*x*tan(d*x)^4*tan(c
)^4 + 12*A*b^4*d*x*tan(d*x)^4*tan(c)^4 - 6*B*a^4*log(4*(tan(d*x)^2*tan(c)^
2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 +
1))*tan(d*x)^4*tan(c)^4 - 24*A*a^3*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*
x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)
^4*tan(c)^4 + 36*B*a^2*b^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c)
+ 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^4*tan(c)^
4 + 24*A*a*b^3*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*
x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^4*tan(c)^4 - 6*B*b^4*
log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 +
tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^4*tan(c)^4 - 48*A*a^4*d*x*tan(d*x)^3
*tan(c)^3 + 192*B*a^3*b*d*x*tan(d*x)^3*tan(c)^3 + 288*A*a^2*b^2*d*x*tan(d*
x)^3*tan(c)^3 - 192*B*a*b^3*d*x*tan(d*x)^3*tan(c)^3 - 48*A*b^4*d*x*tan(d*x
)^3*tan(c)^3 + 36*B*a^2*b^2*tan(d*x)^4*tan(c)^4 + 24*A*a*b^3*tan(d*x)^4*ta
n(c)^4 - 9*B*b^4*tan(d*x)^4*tan(c)^4 + 24*B*a^4*log(4*(tan(d*x)^2*tan(c)^2
- 2*tan(d*x)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1
))*tan(d*x)^3*tan(c)^3 + 96*A*a^3*b*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x
)*tan(c) + 1)/(tan(d*x)^2*tan(c)^2 + tan(d*x)^2 + tan(c)^2 + 1))*tan(d*x)^
3*tan(c)^3 - 144*B*a^2*b^2*log(4*(tan(d*x)^2*tan(c)^2 - 2*tan(d*x)*tan(...
```

3.259.9 Mupad [B] (verification not implemented)

Time = 7.08 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.01

$$\begin{aligned}
& \int (a + b \tan(c + dx))^4 (A + B \tan(c + dx)) dx \\
&= x (Aa^4 - 4Ba^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4) \\
&\quad - \frac{\tan(c + dx) (Ab^4 + 4Bab^3 - 2a^2b(3Ab + 2Ba))}{d} + \frac{\tan(c + dx)^3 \left(\frac{Ab^4}{3} + \frac{4Bab^3}{3}\right)}{d} \\
&\quad + \frac{\ln(\tan(c + dx)^2 + 1) \left(\frac{Ba^4}{2} + 2Aa^3b - 3Ba^2b^2 - 2Aab^3 + \frac{Bb^4}{2}\right)}{d} \\
&\quad - \frac{\tan(c + dx)^2 \left(\frac{Bb^4}{2} - ab^2(2Ab + 3Ba)\right)}{d} + \frac{Bb^4 \tan(c + dx)^4}{4d}
\end{aligned}$$

input `int((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^4,x)`output `x*(A*a^4 + A*b^4 - 6*A*a^2*b^2 + 4*B*a*b^3 - 4*B*a^3*b) - (tan(c + d*x)*(A*b^4 + 4*B*a*b^3 - 2*a^2*b*(3*A*b + 2*B*a)))/d + (tan(c + d*x)^3*((A*b^4)/3 + (4*B*a*b^3)/3))/d + (log(tan(c + d*x)^2 + 1)*((B*a^4)/2 + (B*b^4)/2 - 3*B*a^2*b^2 - 2*A*a*b^3 + 2*A*a^3*b))/d - (tan(c + d*x)^2*((B*b^4)/2 - a*b^2*(2*A*b + 3*B*a)))/d + (B*b^4*tan(c + d*x)^4)/(4*d)`

3.260 $\int \cot(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.260.1 Optimal result	2518
3.260.2 Mathematica [C] (verified)	2519
3.260.3 Rubi [A] (verified)	2519
3.260.4 Maple [A] (verified)	2523
3.260.5 Fricas [A] (verification not implemented)	2524
3.260.6 Sympy [A] (verification not implemented)	2524
3.260.7 Maxima [A] (verification not implemented)	2525
3.260.8 Giac [A] (verification not implemented)	2525
3.260.9 Mupad [B] (verification not implemented)	2526

3.260.1 Optimal result

Integrand size = 29, antiderivative size = 172

$$\begin{aligned} & \int \cot(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx \\ &= (4a^3Ab - 4aAb^3 + a^4B - 6a^2b^2B + b^4B) x \\ & \quad - \frac{b(6a^2Ab - Ab^3 + 4a^3B - 4ab^2B) \log(\cos(c+dx))}{d} \\ & \quad + \frac{a^4A \log(\sin(c+dx))}{d} + \frac{b^2(3aAb + 3a^2B - b^2B) \tan(c+dx)}{d} \\ & \quad + \frac{b(Ab + 2aB)(a+b \tan(c+dx))^2}{2d} + \frac{bB(a+b \tan(c+dx))^3}{3d} \end{aligned}$$

```
output (4*A*a^3*b-4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*x-b*(6*A*a^2*b-A*b^3+4*B*a^3-4*B*a*b^2)*ln(cos(d*x+c))/d+a^4*A*ln(sin(d*x+c))/d+b^2*(3*A*a*b+3*B*a^2-B*b^2)*tan(d*x+c)/d+1/2*b*(A*b+2*B*a)*(a+b*tan(d*x+c))^2/d+1/3*b*B*(a+b*tan(d*x+c))^3/d
```

3.260.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.52 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.87

$$\int \cot(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{-3(a + ib)^4(A + iB) \log(i - \tan(c + dx)) + 6a^4A \log(\tan(c + dx)) - 3(a - ib)^4(A - iB) \log(i + \tan(c + dx))}{6d}$$

input `Integrate[Cot[c + d*x]*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(-3*(a + I*b)^4*(A + I*B)*Log[I - Tan[c + d*x]] + 6*a^4*A*Log[Tan[c + d*x]] - 3*(a - I*b)^4*(A - I*B)*Log[I + Tan[c + d*x]] + 6*b^2*(3*a*A*b + 3*a^2*B - b^2*B)*Tan[c + d*x] + 3*b*(A*b + 2*a*B)*(a + b*Tan[c + d*x])^2 + 2*b*B*(a + b*Tan[c + d*x])^3)/(6*d)`

3.260.3 Rubi [A] (verified)

Time = 1.08 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.01, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.483$, Rules used = {3042, 4090, 27, 3042, 4130, 27, 3042, 4120, 25, 3042, 4107, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^4(A + B \tan(c + dx))}{\tan(c + dx)} dx$$

$$\downarrow \text{4090}$$

$$\frac{1}{3} \int 3 \cot(c + dx)(a + b \tan(c + dx))^2 (Aa^2 + b(Ab + 2aB) \tan^2(c + dx) + (Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx + \frac{bB(a + b \tan(c + dx))^3}{3d}$$

$$\downarrow \text{27}$$

$$\begin{aligned}
& \int \cot(c+dx)(a+b \tan(c+dx))^2 (Aa^2 + b(Ab + 2aB) \tan^2(c+dx) + (Ba^2 + 2Aba - b^2B) \tan(c+dx)) dx + \\
& \quad \frac{bB(a+b \tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a+b \tan(c+dx))^2 (Aa^2 + b(Ab + 2aB) \tan(c+dx)^2 + (Ba^2 + 2Aba - b^2B) \tan(c+dx))}{\tan(c+dx) \frac{bB(a+b \tan(c+dx))^3}{3d}} dx + \\
& \quad \downarrow \text{4130} \\
& \frac{1}{2} \int 2 \cot(c+dx)(a+b \tan(c+dx)) (Aa^3 + b(3Ba^2 + 3Aba - b^2B) \tan^2(c+dx) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)) dx + \\
& \quad \frac{b(2aB + Ab)(a+b \tan(c+dx))^2}{2d} + \frac{bB(a+b \tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{27} \\
& \int \cot(c+dx)(a+b \tan(c+dx)) (Aa^3 + b(3Ba^2 + 3Aba - b^2B) \tan^2(c+dx) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)) dx + \\
& \quad \frac{b(2aB + Ab)(a+b \tan(c+dx))^2}{2d} + \frac{bB(a+b \tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a+b \tan(c+dx)) (Aa^3 + b(3Ba^2 + 3Aba - b^2B) \tan(c+dx)^2 + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx))}{\tan(c+dx) \frac{b(2aB + Ab)(a+b \tan(c+dx))^2}{2d} + \frac{bB(a+b \tan(c+dx))^3}{3d}} \\
& \quad \downarrow \text{4120} \\
& - \int -\cot(c+dx) (Aa^4 + b(4Ba^3 + 6Aba^2 - 4b^2Ba - Ab^3) \tan^2(c+dx) + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \tan(c+dx) + \\
& \quad \frac{b^2(3a^2B + 3aAb - b^2B) \tan(c+dx)}{d} + \frac{b(2aB + Ab)(a+b \tan(c+dx))^2}{2d} + \frac{bB(a+b \tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{25} \\
& \int \cot(c+dx) (Aa^4 + b(4Ba^3 + 6Aba^2 - 4b^2Ba - Ab^3) \tan^2(c+dx) + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \tan(c+dx) + \\
& \quad \frac{b^2(3a^2B + 3aAb - b^2B) \tan(c+dx)}{d} + \frac{b(2aB + Ab)(a+b \tan(c+dx))^2}{2d} + \frac{bB(a+b \tan(c+dx))^3}{3d}
\end{aligned}$$

3.260. $\int \cot(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

↓ 3042

$$\int \frac{Aa^4 + b(4Ba^3 + 6Aba^2 - 4b^2Ba - Ab^3) \tan(c + dx)^2 + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \tan(c + dx)}{\tan(c + dx)} \\ \frac{b^2(3a^2B + 3aAb - b^2B) \tan(c + dx)}{d} + \frac{b(2aB + Ab)(a + b \tan(c + dx))^2}{2d} + \frac{bB(a + b \tan(c + dx))^3}{3d}$$

↓ 4107

$$a^4A \int \cot(c + dx) dx + b(4a^3B + 6a^2Ab - 4ab^2B - Ab^3) \int \tan(c + dx) dx + \\ \frac{b^2(3a^2B + 3aAb - b^2B) \tan(c + dx)}{d} + x(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) + \\ \frac{b(2aB + Ab)(a + b \tan(c + dx))^2}{2d} + \frac{bB(a + b \tan(c + dx))^3}{3d}$$

↓ 3042

$$a^4A \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + b(4a^3B + 6a^2Ab - 4ab^2B - Ab^3) \int \tan(c + dx) dx + \\ \frac{b^2(3a^2B + 3aAb - b^2B) \tan(c + dx)}{d} + x(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) + \\ \frac{b(2aB + Ab)(a + b \tan(c + dx))^2}{2d} + \frac{bB(a + b \tan(c + dx))^3}{3d}$$

↓ 25

$$a^4(-A) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx + b(4a^3B + 6a^2Ab - 4ab^2B - Ab^3) \int \tan(c + dx) dx + \\ \frac{b^2(3a^2B + 3aAb - b^2B) \tan(c + dx)}{d} + x(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) + \\ \frac{b(2aB + Ab)(a + b \tan(c + dx))^2}{2d} + \frac{bB(a + b \tan(c + dx))^3}{3d}$$

↓ 3956

$$\frac{a^4A \log(-\sin(c + dx))}{d} + \frac{b^2(3a^2B + 3aAb - b^2B) \tan(c + dx)}{d} - \\ \frac{b(4a^3B + 6a^2Ab - 4ab^2B - Ab^3) \log(\cos(c + dx))}{d} + x(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) + \\ \frac{b(2aB + Ab)(a + b \tan(c + dx))^2}{2d} + \frac{bB(a + b \tan(c + dx))^3}{3d}$$

input `Int[Cot[c + d*x]*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`


```
output (4*a^3*A*b - 4*a*A*b^3 + a^4*B - 6*a^2*b^2*B + b^4*B)*x - (b*(6*a^2*A*b -
A*b^3 + 4*a^3*B - 4*a*b^2*B)*Log[Cos[c + d*x]])/d + (a^4*A*Log[-Sin[c + d*
x]])/d + (b^2*(3*a*A*b + 3*a^2*B - b^2*B)*Tan[c + d*x])/d + (b*(A*b + 2*a*
B)*(a + b*Tan[c + d*x])^2)/(2*d) + (b*B*(a + b*Tan[c + d*x])^3)/(3*d)
```

3.260.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3956 Int[tan[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4090 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4107 Int[((A_) + (B_)*tan[(e_) + (f_)*(x_)]) + (C_)*tan[(e_) + (f_)*(x_)^2
)/tan[(e_) + (f_)*(x_)], x_Symbol] := Simp[B*x, x] + (Simp[A Int[1/Tan[
e + f*x], x], x] + Simp[C Int[Tan[e + f*x], x], x]) /; FreeQ[{e, f, A, B,
C}, x] && NeQ[A, C]
```

```
rule 4120 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f_)*
*(x_)])^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

```
rule 4130 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_)
+ (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_)
+ (f_)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

3.260.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.00

method	result
parallelrisch	$\frac{(-3Aa^4 + 18Aa^2b^2 - 3Ab^4 + 12Ba^3b - 12Ba^2b^3) \ln(\sec^2(dx+c)) + 6Aa^4 \ln(\tan(dx+c)) + 2Bb^4(\tan^3(dx+c)) + (3Ab^4 + 12Ba^3b - 12Ba^2b^3) \tan(dx+c)}{6d}$
norman	$(4Aa^3b - 4Aa^2b^2 + Ba^4 - 6Ba^2b^2 + Bb^4)x + \frac{b^2(4Aab + 6Ba^2 - Bb^2) \tan(dx+c)}{d} + \frac{Bb^4(\tan^3(dx+c))}{3d}$
derivativedivides	$\frac{Bb^4(\tan^3(dx+c))}{3} + \frac{Ab^4(\tan^2(dx+c))}{2} + 2Ba^3b^3(\tan^2(dx+c)) + 4Aa^3b^3 \tan(dx+c) + 6Ba^2b^2 \tan(dx+c) - Bb^4 \tan(dx+c) + \dots$
default	$\frac{Bb^4(\tan^3(dx+c))}{3} + \frac{Ab^4(\tan^2(dx+c))}{2} + 2Ba^3b^3(\tan^2(dx+c)) + 4Aa^3b^3 \tan(dx+c) + 6Ba^2b^2 \tan(dx+c) - Bb^4 \tan(dx+c) + \dots$
risch	$\frac{8iBa^3bc}{d} - \frac{2iAb^4c}{d} - iAa^4x + \frac{Aa^4 \ln(e^{2i(dx+c)} - 1)}{d} + \frac{\ln(e^{2i(dx+c)} + 1)Ab^4}{d} + 4Aa^3bx - 4Aa^2b^2x - \dots$

```
input int(cot(d*x+c)*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE
)
```

3.260. $\int \cot(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

output $\frac{1}{6} * ((-3 * A * a^4 + 18 * A * a^2 * b^2 - 3 * A * b^4 + 12 * B * a^3 * b - 12 * B * a * b^3) * \ln(\sec(dx+c)^2) + 6 * A * a^4 * \ln(\tan(dx+c)) + 2 * B * b^4 * \tan(dx+c)^3 + (3 * A * b^4 + 12 * B * a * b^3) * \tan(dx+c)^2 + (24 * A * a * b^3 + 36 * B * a^2 * b^2 - 6 * B * b^4) * \tan(dx+c) + 24 * d * (A * a^3 * b - A * a * b^3 + 1/4 * B * a^4 - 3/2 * B * a^2 * b^2 + 1/4 * B * b^4) * x) / d$

3.260.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.08

$$\int \cot(c + dx)(a + b \tan(c + dx))^4 (A + B \tan(c + dx)) dx$$

$$= \frac{2 B b^4 \tan(dx + c)^3 + 3 A a^4 \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) + 6 (B a^4 + 4 A a^3 b - 6 B a^2 b^2 - 4 A a b^3 + B b^4) dx + 3 (4 B a^4 + 12 B a^2 b^2 - 6 B b^4) \tan(dx + c)}{d}$$

input `integrate(cot(dx+c)*(a+b*tan(dx+c))^4*(A+B*tan(dx+c)),x, algorithm="fricas")`

output $\frac{1}{6} * (2 * B * b^4 * \tan(dx + c)^3 + 3 * A * a^4 * \log(\tan(dx + c)^2 / (\tan(dx + c)^2 + 1)) + 6 * (B * a^4 + 4 * A * a^3 * b - 6 * B * a^2 * b^2 - 4 * A * a * b^3 + B * b^4) * dx + 3 * (4 * B * a^4 * b^3 + A * b^4) * \tan(dx + c)^2 - 3 * (4 * B * a^3 * b + 6 * A * a^2 * b^2 - 4 * B * a * b^3 - A * b^4) * \log(1 / (\tan(dx + c)^2 + 1)) + 6 * (6 * B * a^2 * b^2 + 4 * A * a * b^3 - B * b^4) * \tan(dx + c)) / d$

3.260.6 Sympy [A] (verification not implemented)

Time = 0.75 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.69

$$\int \cot(c + dx)(a + b \tan(c + dx))^4 (A + B \tan(c + dx)) dx$$

$$= \begin{cases} -\frac{A a^4 \log(\tan^2(c+dx)+1)}{2d} + \frac{A a^4 \log(\tan(c+dx))}{d} + 4 A a^3 b x + \frac{3 A a^2 b^2 \log(\tan^2(c+dx)+1)}{d} - 4 A a b^3 x + \frac{4 A a b^3 \tan(c+dx)}{d} \\ x(A + B \tan(c)) (a + b \tan(c))^4 \cot(c) \end{cases}$$

input `integrate(cot(dx+c)*(a+b*tan(dx+c))**4*(A+B*tan(dx+c)),x)`

output `Piecewise((-A*a**4*log(tan(c + d*x)**2 + 1)/(2*d) + A*a**4*log(tan(c + d*x)))/d + 4*A*a**3*b*x + 3*A*a**2*b**2*log(tan(c + d*x)**2 + 1)/d - 4*A*a*b**3*x + 4*A*a*b**3*tan(c + d*x)/d - A*b**4*log(tan(c + d*x)**2 + 1)/(2*d) + A*b**4*tan(c + d*x)**2/(2*d) + B*a**4*x + 2*B*a**3*b*log(tan(c + d*x)**2 + 1)/d - 6*B*a**2*b**2*x + 6*B*a**2*b**2*tan(c + d*x)/d - 2*B*a*b**3*log(tan(c + d*x)**2 + 1)/d + 2*B*a*b**3*tan(c + d*x)**2/d + B*b**4*x + B*b**4*tan(c + d*x)**3/(3*d) - B*b**4*tan(c + d*x)/d, Ne(d, 0)), (x*(A + B*tan(c))* (a + b*tan(c))**4*cot(c), True))`

3.260.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.02

$$\int \cot(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{2 B b^4 \tan(dx + c)^3 + 6 A a^4 \log(\tan(dx + c)) + 3(4 B a b^3 + A b^4) \tan(dx + c)^2 + 6(B a^4 + 4 A a^3 b - 6 B a^2 b^2 - 4 A a b^3 + B b^4) \tan(dx + c) - 3(A a^4 - 4 B a^3 b - 6 A a^2 b^2 + 4 B a b^3 + A b^4) \log(\tan(dx + c)^2 + 1) + 6(6 B a^2 b^2 + 4 A a b^3 - B b^4) \tan(dx + c)}{d}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/6*(2*B*b^4*tan(d*x + c)^3 + 6*A*a^4*log(tan(d*x + c)) + 3*(4*B*a*b^3 + A*b^4)*tan(d*x + c)^2 + 6*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*(d*x + c) - 3*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)^2 + 1) + 6*(6*B*a^2*b^2 + 4*A*a*b^3 - B*b^4)*tan(d*x + c))/d`

3.260.8 Giac [A] (verification not implemented)

Time = 1.74 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.11

$$\int \cot(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{2 B b^4 \tan(dx + c)^3 + 12 B a b^3 \tan(dx + c)^2 + 3 A b^4 \tan(dx + c)^2 + 6 A a^4 \log(|\tan(dx + c)|) + 36 B a^2 b^2 \tan(dx + c) - 3(A a^4 - 4 B a^3 b - 6 A a^2 b^2 + 4 B a b^3 + A b^4) \log(\tan(dx + c)^2 + 1) + 6(6 B a^2 b^2 + 4 A a b^3 - B b^4) \tan(dx + c)}{d}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

3.260. $\int \cot(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

output $1/6*(2*B*b^4*\tan(d*x + c)^3 + 12*B*a*b^3*\tan(d*x + c)^2 + 3*A*b^4*\tan(d*x + c)^2 + 6*A*a^4*\log(\text{abs}(\tan(d*x + c))) + 36*B*a^2*b^2*\tan(d*x + c) + 24*A*a*b^3*\tan(d*x + c) - 6*B*b^4*\tan(d*x + c) + 6*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*(d*x + c) - 3*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*\log(\tan(d*x + c)^2 + 1))/d$

3.260.9 Mupad [B] (verification not implemented)

Time = 7.47 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.88

$$\int \cot(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{\tan(c + dx)^2 \left(\frac{Ab^4}{2} + 2Bab^3 \right)}{d} - \frac{\tan(c + dx) (Bb^4 - 2ab^2(2Ab + 3Ba))}{d}$$

$$+ \frac{Aa^4 \ln(\tan(c + dx))}{d} - \frac{\ln(\tan(c + dx) + 1i) (A - B1i) (b + a1i)^4}{2d}$$

$$- \frac{\ln(\tan(c + dx) - i) (A + B1i) (-b + a1i)^4}{2d} + \frac{Bb^4 \tan(c + dx)^3}{3d}$$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^4,x)`

output $(\tan(c + d*x)^2*((A*b^4)/2 + 2*B*a*b^3))/d - (\tan(c + d*x)*(B*b^4 - 2*a*b^2*(2*A*b + 3*B*a)))/d + (A*a^4*\log(\tan(c + d*x)))/d - (\log(\tan(c + d*x) + 1i)*(A - B*1i)*(a*1i + b)^4)/(2*d) - (\log(\tan(c + d*x) - 1i)*(A + B*1i)*(a*1i - b)^4)/(2*d) + (B*b^4*\tan(c + d*x)^3)/(3*d)$

3.261 $\int \cot^2(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.261.1 Optimal result	2527
3.261.2 Mathematica [C] (verified)	2528
3.261.3 Rubi [A] (verified)	2528
3.261.4 Maple [A] (verified)	2532
3.261.5 Fricas [A] (verification not implemented)	2533
3.261.6 Sympy [A] (verification not implemented)	2533
3.261.7 Maxima [A] (verification not implemented)	2534
3.261.8 Giac [A] (verification not implemented)	2534
3.261.9 Mupad [B] (verification not implemented)	2535

3.261.1 Optimal result

Integrand size = 31, antiderivative size = 175

$$\int \cot^2(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= -((a^4A - 6a^2Ab^2 + Ab^4 - 4a^3bB + 4ab^3B)x) - \frac{b^2(4aAb + 6a^2B - b^2B) \log(\cos(c+dx))}{d}$$

$$+ \frac{a^3(4Ab + aB) \log(\sin(c+dx))}{d} + \frac{b^2(a^2A + Ab^2 + 3abB) \tan(c+dx)}{d}$$

$$+ \frac{b(2aA + bB)(a+b \tan(c+dx))^2}{2d} - \frac{aA \cot(c+dx)(a+b \tan(c+dx))^3}{d}$$

output

```
-(A*a^4-6*A*a^2*b^2+A*b^4-4*B*a^3*b+4*B*a*b^3)*x-b^2*(4*A*a*b+6*B*a^2-B*b^2)*ln(cos(d*x+c))/d+a^3*(4*A*b+B*a)*ln(sin(d*x+c))/d+b^2*(A*a^2+A*b^2+3*B*a*b)*tan(d*x+c)/d+1/2*b*(2*A*a+B*b)*(a+b*tan(d*x+c))^2/d-a*A*cot(d*x+c)*(a+b*tan(d*x+c))^3/d
```

3.261.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.11 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.77

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{-2a^4 A \cot(c + dx) + i(a + ib)^4(A + iB) \log(i - \tan(c + dx)) + 2a^3(4Ab + aB) \log(\tan(c + dx)) - (a - b)^4(A + B \tan(c + dx))}{2d}$$

input `Integrate[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(-2*a^4*A*Cot[c + d*x] + I*(a + I*b)^4*(A + I*B)*Log[I - Tan[c + d*x]] + 2*a^3*(4*A*b + a*B)*Log[Tan[c + d*x]] - (a - I*b)^4*(I*A + B)*Log[I + Tan[c + d*x]] + 2*b^3*(A*b + 4*a*B)*Tan[c + d*x] + b^4*B*Tan[c + d*x]^2)/(2*d)`

3.261.3 Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.01, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 4088, 3042, 4130, 27, 3042, 4120, 25, 3042, 4107, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \tan(c + dx))^4(A + B \tan(c + dx))}{\tan(c + dx)^2} dx$$

$$\downarrow 4088$$

$$\int \cot(c + dx)(a + b \tan(c + dx))^2 (b(2aA + bB) \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(4Ab + aB)) dx - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^3}{d}$$

$$\downarrow 3042$$

$$\begin{aligned}
& \int \frac{(a + b \tan(c + dx))^2 (b(2aA + bB) \tan(c + dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(4Ab + aB))}{\tan(c + dx)} dx - \\
& \quad \frac{aA \cot(c + dx)(a + b \tan(c + dx))^3}{d} \\
& \quad \downarrow 4130 \\
& \quad \frac{1}{2} \int 2 \cot(c + dx)(a + b \tan(c + dx)) ((4Ab + aB)a^2 + b(Aa^2 + 3bBa + Ab^2) \tan^2(c + dx) - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)) dx + \\
& \quad \frac{b(2aA + bB)(a + b \tan(c + dx))^2}{2d} - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^3}{d} \\
& \quad \downarrow 27 \\
& \quad \int \cot(c + dx)(a + b \tan(c + dx)) ((4Ab + aB)a^2 + b(Aa^2 + 3bBa + Ab^2) \tan^2(c + dx) - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)) dx + \\
& \quad \frac{b(2aA + bB)(a + b \tan(c + dx))^2}{2d} - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^3}{d} \\
& \quad \downarrow 3042 \\
& \quad \int \frac{(a + b \tan(c + dx)) ((4Ab + aB)a^2 + b(Aa^2 + 3bBa + Ab^2) \tan(c + dx)^2 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx))}{\tan(c + dx)} dx - \\
& \quad \frac{b(2aA + bB)(a + b \tan(c + dx))^2}{2d} - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^3}{d} \\
& \quad \downarrow 4120 \\
& \quad - \int -\cot(c + dx) ((4Ab + aB)a^3 + b^2(6Ba^2 + 4Aba - b^2B) \tan^2(c + dx) - (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4) \tan(c + dx) + \\
& \quad \frac{b^2(a^2A + 3abB + Ab^2) \tan(c + dx)}{d} + \frac{b(2aA + bB)(a + b \tan(c + dx))^2}{2d} - \\
& \quad \frac{aA \cot(c + dx)(a + b \tan(c + dx))^3}{d} \\
& \quad \downarrow 25 \\
& \quad \int \cot(c + dx) ((4Ab + aB)a^3 + b^2(6Ba^2 + 4Aba - b^2B) \tan^2(c + dx) - (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4) \tan(c + dx) + \\
& \quad \frac{b^2(a^2A + 3abB + Ab^2) \tan(c + dx)}{d} + \frac{b(2aA + bB)(a + b \tan(c + dx))^2}{2d} - \\
& \quad \frac{aA \cot(c + dx)(a + b \tan(c + dx))^3}{d} \\
& \quad \downarrow 3042
\end{aligned}$$

3.261. $\int \cot^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \int \frac{(4Ab + aB)a^3 + b^2(6Ba^2 + 4Aba - b^2B) \tan(c + dx)^2 - (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4) \tan(c + dx)}{\tan(c + dx)} \\
& \quad - \frac{b^2(a^2A + 3abB + Ab^2) \tan(c + dx)}{d} + \frac{b(2aA + bB)(a + b \tan(c + dx))^2}{aA \cot(c + dx)(a + b \tan(c + dx))^3} - \\
& \quad \quad \quad \downarrow \text{4107} \\
& \quad a^3(aB + 4Ab) \int \cot(c + dx) dx + b^2(6a^2B + 4aAb - b^2B) \int \tan(c + dx) dx + \\
& \quad \frac{b^2(a^2A + 3abB + Ab^2) \tan(c + dx)}{2d} - x(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) + \\
& \quad \quad \frac{b(2aA + bB)(a + b \tan(c + dx))^2}{2d} - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^3}{d} \\
& \quad \quad \quad \downarrow \text{3042} \\
& \quad a^3(aB + 4Ab) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx + b^2(6a^2B + 4aAb - b^2B) \int \tan(c + dx) dx + \\
& \quad \frac{b^2(a^2A + 3abB + Ab^2) \tan(c + dx)}{2d} - x(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) + \\
& \quad \quad \frac{b(2aA + bB)(a + b \tan(c + dx))^2}{2d} - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^3}{d} \\
& \quad \quad \quad \downarrow \text{25} \\
& \quad -\left(a^3(aB + 4Ab) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx\right) + b^2(6a^2B + 4aAb - b^2B) \int \tan(c + dx) dx + \\
& \quad \frac{b^2(a^2A + 3abB + Ab^2) \tan(c + dx)}{2d} - x(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) + \\
& \quad \quad \frac{b(2aA + bB)(a + b \tan(c + dx))^2}{2d} - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^3}{d} \\
& \quad \quad \quad \downarrow \text{3956} \\
& \quad \frac{a^3(aB + 4Ab) \log(-\sin(c + dx))}{d} + \frac{b^2(a^2A + 3abB + Ab^2) \tan(c + dx)}{d} - \\
& \quad \frac{b^2(6a^2B + 4aAb - b^2B) \log(\cos(c + dx))}{d} - x(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) + \\
& \quad \quad \frac{b(2aA + bB)(a + b \tan(c + dx))^2}{2d} - \frac{aA \cot(c + dx)(a + b \tan(c + dx))^3}{d}
\end{aligned}$$

input `Int[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

```
output 
$$-\left((a^4A - 6a^2Ab^2 + Ab^4 - 4a^3bB + 4ab^3B)x\right) - (b^2(4aAb + 6a^2B - b^2B)\text{Log}[\text{Cos}[c + dx]])/d + (a^3(4Ab + aB)\text{Log}[-\text{Sin}[c + dx]])/d + (b^2(a^2A + Ab^2 + 3abB)\text{Tan}[c + dx])/d + (b(2aA + bB)(a + b\text{Tan}[c + dx])^2)/(2d) - (aA\text{Cot}[c + dx](a + b\text{Tan}[c + dx])^3)/d$$

```

3.261.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + dx], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4088 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])
```

```
rule 4107 Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[B*x, x] + (Simp[A Int[1/Tan[e + f*x], x], x] + Simp[C Int[Tan[e + f*x], x], x]) /; FreeQ[{e, f, A, B, C}, x] && NeQ[A, C]
```

```
rule 4120 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f_)*
(x_)])^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

```
rule 4130 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_)
+ (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_)
+ (f_)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && (!IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

3.261.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.91

method	result
parallelrisch	$\frac{(-4A^3b+4Aa^3-Ba^4+6Ba^2b^2-Bb^4) \ln(\sec^2(dx+c))+(8A^3b+2Ba^4) \ln(\tan(dx+c))+Bb^4(\tan^2(dx+c))+(2A^3b+2Aa^3-Ba^4) \ln(1+\tan^2(dx+c))}{2d}$
derivativedivides	$\frac{\frac{Bb^4(\tan^2(dx+c))}{2}+Ab^4 \tan(dx+c)+4Ba^3 \tan(dx+c)+\frac{(-4A^3b+4Aa^3-Ba^4+6Ba^2b^2-Bb^4) \ln(1+\tan^2(dx+c))}{2}}{d}+(-A^3b+2Aa^3-Ba^4) \ln(1+\tan^2(dx+c))$
default	$\frac{\frac{Bb^4(\tan^2(dx+c))}{2}+Ab^4 \tan(dx+c)+4Ba^3 \tan(dx+c)+\frac{(-4A^3b+4Aa^3-Ba^4+6Ba^2b^2-Bb^4) \ln(1+\tan^2(dx+c))}{2}}{d}+(-A^3b+2Aa^3-Ba^4) \ln(1+\tan^2(dx+c))$
norman	$\frac{(-A^4+6A^2a^2b^2-A^4b^4+4Ba^3b-4Ba^3b^3)x \tan(dx+c)+\frac{b^3(Ab+4Ba)(\tan^2(dx+c))}{d}-\frac{A^4a^4}{d}+\frac{Bb^4(\tan^3(dx+c))}{2d}}{\tan(dx+c)}+a^3(4Ab^3+3A^2a^2b^2-3A^2a^2b^2b^3)$
risch	$\frac{a^4 \ln(e^{2i(dx+c)}-1)B}{d}+\frac{12iBa^2b^2c}{d}-\frac{2ia^4Bc}{d}+\frac{8iAab^3c}{d}-iBb^4x+\frac{\ln(e^{2i(dx+c)}+1)Bb^4}{d}-\frac{2iBb^4c}{d}+\frac{2a^3(4Ab^3+3A^2a^2b^2-3A^2a^2b^2b^3)}{d}$

```
input int(cot(d*x+c)^2*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

$$3.261. \quad \int \cot^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

output $\frac{1}{2} * ((-4 * A * a^3 * b + 4 * A * a * b^3 - B * a^4 + 6 * B * a^2 * b^2 - B * b^4) * \ln(\sec(dx+c)^2) + (8 * A * a^3 * b + 2 * B * a^4) * \ln(\tan(dx+c)) + B * b^4 * \tan(dx+c)^2 + (2 * A * b^4 + 8 * B * a * b^3) * \tan(dx+c) - 2 * A * \cot(dx+c) * a^4 - 2 * d * x * (A * a^4 - 6 * A * a^2 * b^2 + A * b^4 - 4 * B * a^3 * b + 4 * B * a * b^3)) / d$

3.261.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.10

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{Bb^4 \tan(dx + c)^3 - 2Aa^4 + (Ba^4 + 4Aa^3b) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx + c) - (6Ba^2b^2 + 4Aab^3 - Bb^4) \log(\tan(dx+c))}{d}$$

input `integrate(cot(dx+c)^2*(a+b*tan(dx+c))^4*(A+B*tan(dx+c)),x, algorithm="fricas")`

output $\frac{1}{2} * (B * b^4 * \tan(dx + c)^3 - 2 * A * a^4 + (B * a^4 + 4 * A * a^3 * b) * \log(\tan(dx + c)^2 / (\tan(dx + c)^2 + 1)) * \tan(dx + c) - (6 * B * a^2 * b^2 + 4 * A * a * b^3 - B * b^4) * \log(1 / (\tan(dx + c)^2 + 1)) * \tan(dx + c) + 2 * (4 * B * a * b^3 + A * b^4) * \tan(dx + c)^2 + (B * b^4 - 2 * (A * a^4 - 4 * B * a^3 * b - 6 * A * a^2 * b^2 + 4 * B * a * b^3 + A * b^4) * d * x) * \tan(dx + c)) / (d * \tan(dx + c))$

3.261.6 Sympy [A] (verification not implemented)

Time = 1.00 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.65

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \tilde{\infty} A a^4 x \\ x(A + B \tan(c))(a + b \tan(c))^4 \cot^2(c) \\ \tilde{\infty} A a^4 x \end{cases}$$

$$- A a^4 x - \frac{A a^4}{d \tan(c+dx)} - \frac{2 A a^3 b \log(\tan^2(c+dx)+1)}{d} + \frac{4 A a^3 b \log(\tan(c+dx))}{d} + 6 A a^2 b^2 x + \frac{2 A a b^3 \log(\tan^2(c+dx)+1)}{d} - A b^4 x$$

input `integrate(cot(dx+c)**2*(a+b*tan(dx+c))**4*(A+B*tan(dx+c)),x)`

3.261. $\int \cot^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

output `Piecewise((zoo*A*a**4*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**4*cot(c)**2, Eq(d, 0)), (zoo*A*a**4*x, Eq(c, -d*x)), (-A*a**4*x - A*a**4/(d*tan(c + d*x)) - 2*A*a**3*b*log(tan(c + d*x)**2 + 1)/d + 4*A*a**3*b*log(tan(c + d*x))/d + 6*A*a**2*b**2*x + 2*A*a*b**3*log(tan(c + d*x)**2 + 1)/d - A*b**4*x + A*b**4*tan(c + d*x)/d - B*a**4*log(tan(c + d*x)**2 + 1)/(2*d) + B*a**4*log(tan(c + d*x))/d + 4*B*a**3*b*x + 3*B*a**2*b**2*log(tan(c + d*x)**2 + 1)/d - 4*B*a*b**3*x + 4*B*a*b**3*tan(c + d*x)/d - B*b**4*log(tan(c + d*x)**2 + 1)/(2*d) + B*b**4*tan(c + d*x)**2/(2*d), True))`

3.261.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.94

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{Bb^4 \tan(dx + c)^2 - \frac{2Aa^4}{\tan(dx+c)} - 2(Aa^4 - 4Ba^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4)(dx + c) - (Ba^4 + 4Aa^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4) \log(\tan(dx + c)^2 + 1) + 2(Ba^4 + 4Aa^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4) \tan(dx + c)}{d}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(B*b^4*tan(d*x + c)^2 - 2*A*a^4/tan(d*x + c) - 2*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*(d*x + c) - (B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*log(tan(d*x + c)^2 + 1) + 2*(B*a^4 + 4*A*a^3*b)*log(tan(d*x + c)) + 2*(4*B*a*b^3 + A*b^4)*tan(d*x + c))/d`

3.261.8 Giac [A] (verification not implemented)

Time = 1.28 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.11

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{Bb^4 \tan(dx + c)^2 + 8Bab^3 \tan(dx + c) + 2Ab^4 \tan(dx + c) - 2(Aa^4 - 4Ba^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4) \log(\tan(dx + c)^2 + 1) + 2(Ba^4 + 4Aa^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4) \tan(dx + c)}{d}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

3.261. $\int \cot^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

output `1/2*(B*b^4*tan(d*x + c)^2 + 8*B*a*b^3*tan(d*x + c) + 2*A*b^4*tan(d*x + c) - 2*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*(d*x + c) - (B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*log(tan(d*x + c)^2 + 1) + 2*(B*a^4 + 4*A*a^3*b)*log(abs(tan(d*x + c))) - 2*(B*a^4*tan(d*x + c) + 4*A*a^3*b*tan(d*x + c) + A*a^4)/tan(d*x + c))/d`

3.261.9 Mupad [B] (verification not implemented)

Time = 7.98 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.81

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{\tan(c + dx) (A b^4 + 4 B a b^3)}{d} + \frac{\ln(\tan(c + dx)) (B a^4 + 4 A b a^3)}{d}$$

$$+ \frac{\ln(\tan(c + dx) - i) (-B + A i) (-b + a i)^4}{2 d}$$

$$- \frac{\ln(\tan(c + dx) + i) (B + A i) (b + a i)^4}{2 d} - \frac{A a^4 \cot(c + dx)}{d} + \frac{B b^4 \tan(c + dx)^2}{2 d}$$

input `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^4,x)`

output `(tan(c + d*x)*(A*b^4 + 4*B*a*b^3))/d + (log(tan(c + d*x))*(B*a^4 + 4*A*a^3*b))/d + (log(tan(c + d*x) - 1i)*(A*1i - B)*(a*1i - b)^4)/(2*d) - (log(tan(c + d*x) + 1i)*(A*1i + B)*(a*1i + b)^4)/(2*d) - (A*a^4*cot(c + d*x))/d + (B*b^4*tan(c + d*x)^2)/(2*d)`

3.262 $\int \cot^3(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.262.1 Optimal result	2536
3.262.2 Mathematica [C] (verified)	2537
3.262.3 Rubi [A] (verified)	2537
3.262.4 Maple [A] (verified)	2541
3.262.5 Fricas [A] (verification not implemented)	2542
3.262.6 Sympy [A] (verification not implemented)	2542
3.262.7 Maxima [A] (verification not implemented)	2543
3.262.8 Giac [A] (verification not implemented)	2543
3.262.9 Mupad [B] (verification not implemented)	2544

3.262.1 Optimal result

Integrand size = 31, antiderivative size = 186

$$\int \cot^3(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= -((4a^3Ab - 4aAb^3 + a^4B - 6a^2b^2B + b^4B) x) - \frac{b^3(Ab + 4aB) \log(\cos(c+dx))}{d}$$

$$- \frac{a^2(a^2A - 6Ab^2 - 4abB) \log(\sin(c+dx))}{d} + \frac{b^2(3aAb + a^2B + b^2B) \tan(c+dx)}{d}$$

$$- \frac{a(5Ab + 2aB) \cot(c+dx)(a+b \tan(c+dx))^2}{2d} - \frac{aA \cot^2(c+dx)(a+b \tan(c+dx))^3}{2d}$$

output

```
-(4*A*a^3*b-4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*x-b^3*(A*b+4*B*a)*ln(cos(d*x+c))/d-a^2*(A*a^2-6*A*b^2-4*B*a*b)*ln(sin(d*x+c))/d+b^2*(3*A*a*b+B*a^2+B*b^2)*tan(d*x+c)/d-1/2*a*(5*A*b+2*B*a)*cot(d*x+c)*(a+b*tan(d*x+c))^2/d-1/2*a*A*cot(d*x+c)^2*(a+b*tan(d*x+c))^3/d
```

3.262.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.70 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.75

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{-2a^3(4Ab + aB) \cot(c + dx) - a^4 A \cot^2(c + dx) + (a + ib)^4(A + iB) \log(i - \tan(c + dx)) - 2a^2(a^2 A - 6a^2 B) \cot^2(c + dx) + (a - ib)^4(A - iB) \log(i + \tan(c + dx)) + 2b^4 B \tan^2(c + dx)}{2d}$$

input `Integrate[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(-2*a^3*(4*A*b + a*B)*Cot[c + d*x] - a^4*A*Cot[c + d*x]^2 + (a + I*b)^4*(A + I*B)*Log[I - Tan[c + d*x]] - 2*a^2*(a^2*A - 6*A*b^2 - 4*a*b*B)*Log[Tan[c + d*x]] + (a - I*b)^4*(A - I*B)*Log[I + Tan[c + d*x]] + 2*b^4*B*Tan[c + d*x])/ (2*d)`

3.262.3 Rubi [A] (verified)

Time = 1.20 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.03, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 4088, 3042, 4128, 27, 3042, 4120, 25, 3042, 4107, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^4(A + B \tan(c + dx))}{\tan(c + dx)^3} dx$$

$$\downarrow \text{4088}$$

$$\frac{1}{2} \int \cot^2(c + dx)(a + b \tan(c + dx))^2 (b(aA + 2bB) \tan^2(c + dx) - 2(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(5Ab + 2aB)) dx - \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^3}{2d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{2} \int \frac{(a + b \tan(c + dx))^2 (b(aA + 2bB) \tan(c + dx)^2 - 2(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(5Ab + 2aB))}{\tan(c + dx)^2} dx - \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^3}{2d}$$

↓ 4128

$$\frac{1}{2} \left(\int -2 \cot(c + dx)(a + b \tan(c + dx)) (-b(Ba^2 + 3Aba + b^2B) \tan^2(c + dx) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)) \right) \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^3}{2d}$$

↓ 27

$$\frac{1}{2} \left(-2 \int \cot(c + dx)(a + b \tan(c + dx)) (-b(Ba^2 + 3Aba + b^2B) \tan^2(c + dx) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)) \right) \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^3}{2d}$$

↓ 3042

$$\frac{1}{2} \left(-2 \int \frac{(a + b \tan(c + dx)) (-b(Ba^2 + 3Aba + b^2B) \tan(c + dx)^2 + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx))}{\tan(c + dx)} \right) \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^3}{2d}$$

↓ 4120

$$\frac{1}{2} \left(-2 \left(- \int -\cot(c + dx) (-((Ab + 4aB) \tan^2(c + dx)b^3) + a^2(Aa^2 - 4bBa - 6Ab^2) + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3)) \right) \right) \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^3}{2d}$$

↓ 25

$$\frac{1}{2} \left(-2 \left(\int \cot(c + dx) (-((Ab + 4aB) \tan^2(c + dx)b^3) + a^2(Aa^2 - 4bBa - 6Ab^2) + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3)) \right) \right) \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^3}{2d}$$

↓ 3042

$$\frac{1}{2} \left(-2 \left(\int \frac{-((Ab + 4aB) \tan(c + dx)^2 b^3) + a^2(Aa^2 - 4bBa - 6Ab^2) + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B)}{\tan(c + dx)} \right. \right. \\ \left. \left. \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^3}{2d} \right) \right. \\ \left. \downarrow 4107 \right.$$

$$\frac{1}{2} \left(-2 \left(a^2(a^2A - 4abB - 6Ab^2) \int \cot(c + dx) dx - (b^3(4aB + Ab) \int \tan(c + dx) dx) \right) - \frac{b^2(a^2B + 3aAb + b^2B)}{d} \right. \\ \left. \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^3}{2d} \right) \\ \left. \downarrow 3042 \right.$$

$$\frac{1}{2} \left(-2 \left(a^2(a^2A - 4abB - 6Ab^2) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - (b^3(4aB + Ab) \int \tan(c + dx) dx) \right) - \frac{b^2(a^2B + 3aAb + b^2B)}{d} \right. \\ \left. \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^3}{2d} \right) \\ \left. \downarrow 25 \right.$$

$$\frac{1}{2} \left(-2 \left(-a^2(a^2A - 4abB - 6Ab^2) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - (b^3(4aB + Ab) \int \tan(c + dx) dx) \right) - \frac{b^2(a^2B + 3aAb + b^2B)}{d} \right. \\ \left. \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^3}{2d} \right) \\ \left. \downarrow 3956 \right.$$

$$\frac{1}{2} \left(-2 \left(-\frac{b^2(a^2B + 3aAb + b^2B) \tan(c + dx)}{d} + \frac{a^2(a^2A - 4abB - 6Ab^2) \log(-\sin(c + dx))}{d} \right) + x(a^4B + 4a^3Ab - b^4B) \right. \\ \left. \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^3}{2d} \right)$$

input `Int[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `-1/2*(a*A*Cot[c + d*x]^2*(a + b*Tan[c + d*x])^3)/d + (-((a*(5*A*b + 2*a*B)*Cot[c + d*x]*(a + b*Tan[c + d*x])^2)/d - 2*((4*a^3*A*b - 4*a*A*b^3 + a^4*B - 6*a^2*b^2*B + b^4*B)*x + (b^3*(A*b + 4*a*B)*Log[Cos[c + d*x]])/d + (a^2*(a^2*A - 6*A*b^2 - 4*a*b*B)*Log[-Sin[c + d*x]])/d - (b^2*(3*a*A*b + a^2*B + b^2*B)*Tan[c + d*x])/d))/2`

3.262. $\int \cot^3(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

3.262.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`
- rule 4107 `Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2 /tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[B*x, x] + (Simp[A Int[1/Tan[e + f*x], x], x] + Simp[C Int[Tan[e + f*x], x], x]) /; FreeQ[{e, f, A, B, C}, x] && NeQ[A, C]`

rule 4120 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]`

rule 4128 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

3.262.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.90

method	result
parallelrisch	$\frac{(Aa^4 - 6Aa^2b^2 + Ab^4 - 4Ba^3b + 4Ba^3b^3) \ln(\sec^2(dx+c)) + (-2Aa^4 + 12Aa^2b^2 + 8Ba^3b) \ln(\tan(dx+c)) - A(\cot^2(dx+c))}{2d}$
derivativedivides	$Bb^4 \tan(dx+c) + \frac{(Aa^4 - 6Aa^2b^2 + Ab^4 - 4Ba^3b + 4Ba^3b^3) \ln(1 + \tan^2(dx+c))}{2} + \frac{(-4Aa^3b + 4Aab^3 - Ba^4 + 6Ba^2b^2 - Bb^4) \arctan(\tan(dx+c))}{d}$
default	$Bb^4 \tan(dx+c) + \frac{(Aa^4 - 6Aa^2b^2 + Ab^4 - 4Ba^3b + 4Ba^3b^3) \ln(1 + \tan^2(dx+c))}{2} + \frac{(-4Aa^3b + 4Aab^3 - Ba^4 + 6Ba^2b^2 - Bb^4) \arctan(\tan(dx+c))}{d}$
norman	$\frac{(-4Aa^3b + 4Aab^3 - Ba^4 + 6Ba^2b^2 - Bb^4)x(\tan^2(dx+c)) + \frac{Bb^4(\tan^3(dx+c))}{d} - \frac{Aa^4}{2d} - \frac{a^3(4Ab+Ba)\tan(dx+c)}{d}}{\tan(dx+c)^2} + \frac{(Aa^4 - 6Aa^2b^2 + Ab^4 - 4Ba^3b + 4Ba^3b^3) \ln(1 + \tan^2(dx+c))}{2}$
risch	$\frac{2iAb^4c}{d} + iAa^4x + 4iBa^3b^3x - \frac{Aa^4 \ln(e^{2i(dx+c)} - 1)}{d} - \frac{\ln(e^{2i(dx+c)} + 1)Ab^4}{d} + \frac{6a^2 \ln(e^{2i(dx+c)} - 1)Ab^2}{d}$

input `int(cot(d*x+c)^3*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

$$3.262. \quad \int \cot^3(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

output $\frac{1}{2}((Aa^4 - 6Aa^2b^2 + Ab^4 - 4Ba^3b + 4B^2a^2b^3) \ln(\sec(dx+c)^2) + (-2Aa^4 + 12Aa^2b^2 + 8B^2a^3b) \ln(\tan(dx+c)) - A \cot(dx+c)^2 a^4 + (-8Aa^3b - 2B^2a^4) \cot(dx+c) + 2B^2b^4 \tan(dx+c) - 8d(Aa^3b - Aa^2b^3 + \frac{1}{4}Ba^4 - \frac{3}{2}Ba^2b^2 + \frac{1}{4}B^2b^4)x) / d$

3.262.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.07

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{2Bb^4 \tan(dx + c)^3 - Aa^4 - (Aa^4 - 4Ba^3b - 6Aa^2b^2) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx + c)^2 - (4Bab^3 + Ab^4)}{d}$$

input `integrate(cot(dx+c)^3*(a+b*tan(dx+c))^4*(A+B*tan(dx+c)),x, algorithm="fricas")`

output $\frac{1}{2}((2B^2b^4 \tan(dx + c)^3 - Aa^4 - (Aa^4 - 4Ba^3b - 6Aa^2b^2) \log(\frac{\tan(dx + c)^2}{\tan(dx + c)^2 + 1}) \tan(dx + c)^2 - (4B^2a^2b^3 + Ab^4) \log(1/(\tan(dx + c)^2 + 1)) \tan(dx + c)^2 - (Aa^4 + 2(Ba^4 + 4Aa^3b - 6B^2a^2b^2 - 4Aa^2b^3 + B^2b^4) dx) \tan(dx + c)^2 - 2(Ba^4 + 4Aa^3b) \tan(dx + c)) / (d \tan(dx + c)^2)$

3.262.6 Sympy [A] (verification not implemented)

Time = 1.79 (sec) , antiderivative size = 309, normalized size of antiderivative = 1.66

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \tilde{\infty} Aa^4 x \\ x(A + B \tan(c))(a + b \tan(c))^4 \cot^3(c) \\ \tilde{\infty} Aa^4 x \end{cases}$$

$$= \frac{Aa^4 \log(\tan^2(c+dx)+1)}{2d} - \frac{Aa^4 \log(\tan(c+dx))}{d} - \frac{Aa^4}{2d \tan^2(c+dx)} - 4Aa^3bx - \frac{4Aa^3b}{d \tan(c+dx)} - \frac{3Aa^2b^2 \log(\tan^2(c+dx)+1)}{d} + \dots$$

input `integrate(cot(dx+c)**3*(a+b*tan(dx+c))**4*(A+B*tan(dx+c)),x)`

3.262. $\int \cot^3(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

output `Piecewise((zoo*A**4*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**4*cot(c)**3, Eq(d, 0)), (zoo*A**4*x, Eq(c, -d*x)), (A**4*log(tan(c + d*x)**2 + 1)/(2*d) - A**4*log(tan(c + d*x))/d - A**4/(2*d*tan(c + d*x)**2) - 4*A**3*b*x - 4*A**3*b/(d*tan(c + d*x)) - 3*A**2*b**2*log(tan(c + d*x)**2 + 1)/d + 6*A**2*b**2*log(tan(c + d*x))/d + 4*A*a*b**3*x + A*b**4*log(tan(c + d*x)**2 + 1)/(2*d) - B**4*x - B**4/(d*tan(c + d*x)) - 2*B**3*b*log(tan(c + d*x)**2 + 1)/d + 4*B**3*b*log(tan(c + d*x))/d + 6*B**2*b**2*x + 2*B*a*b**3*log(tan(c + d*x)**2 + 1)/d - B*b**4*x + B*b**4*tan(c + d*x)/d, True))`

3.262.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.93

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{2 B b^4 \tan(dx + c) - 2 (B a^4 + 4 A a^3 b - 6 B a^2 b^2 - 4 A a b^3 + B b^4)(dx + c) + (A a^4 - 4 B a^3 b - 6 A a^2 b^2 + 4 A a b^3 + B b^4) \log(\tan(dx + c)^2 + 1) - 2 (A a^4 - 4 B a^3 b - 6 A a^2 b^2) \log(\tan(dx + c)) - (A a^4 + 2 (B a^4 + 4 A a^3 b) \tan(dx + c)) / \tan(dx + c)^2}{d}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(2*B*b^4*tan(d*x + c) - 2*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*(d*x + c) + (A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)^2 + 1) - 2*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2)*log(tan(d*x + c)) - (A*a^4 + 2*(B*a^4 + 4*A*a^3*b)*tan(d*x + c))/tan(d*x + c)^2)/d`

3.262.8 Giac [A] (verification not implemented)

Time = 1.39 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.20

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{2 B b^4 \tan(dx + c) - 2 (B a^4 + 4 A a^3 b - 6 B a^2 b^2 - 4 A a b^3 + B b^4)(dx + c) + (A a^4 - 4 B a^3 b - 6 A a^2 b^2 + 4 A a b^3 + B b^4) \log(\tan(dx + c)^2 + 1) - 2 (A a^4 - 4 B a^3 b - 6 A a^2 b^2) \log(\tan(dx + c)) - (A a^4 + 2 (B a^4 + 4 A a^3 b) \tan(dx + c)) / \tan(dx + c)^2}{d}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

3.262. $\int \cot^3(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

output $1/2*(2*B*b^4*\tan(d*x + c) - 2*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*(d*x + c) + (A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*\log(\tan(d*x + c)^2 + 1) - 2*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2)*\log(\tan(d*x + c)) + (3*A*a^4*\tan(d*x + c)^2 - 12*B*a^3*b*\tan(d*x + c)^2 - 18*A*a^2*b^2*\tan(d*x + c)^2 - 2*B*a^4*\tan(d*x + c) - 8*A*a^3*b*\tan(d*x + c) - A*a^4)/\tan(d*x + c)^2)/d$

3.262.9 Mupad [B] (verification not implemented)

Time = 7.40 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.80

$$\begin{aligned} & \int \cot^3(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx \\ &= \frac{\ln(\tan(c + dx))(-Aa^4 + 4Ba^3b + 6Aa^2b^2)}{d} \\ & \quad - \frac{\cot(c + dx)^2 \left(\tan(c + dx)(Ba^4 + 4Aba^3) + \frac{Aa^4}{2} \right)}{d} \\ & \quad + \frac{\ln(\tan(c + dx) + 1i)(A - B1i)(b + a1i)^4}{2d} + \frac{Bb^4 \tan(c + dx)}{d} \\ & \quad + \frac{\ln(\tan(c + dx) - 1i)(A + B1i)(-b + a1i)^4}{2d} \end{aligned}$$

input `int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^4,x)`

output $(\log(\tan(c + d*x))*(6*A*a^2*b^2 - A*a^4 + 4*B*a^3*b))/d - (\cot(c + d*x)^2*(\tan(c + d*x)*(B*a^4 + 4*A*a^3*b) + (A*a^4)/2))/d + (\log(\tan(c + d*x) + 1i))*(A - B*1i)*(a*1i + b)^4/(2*d) + (B*b^4*\tan(c + d*x))/d + (\log(\tan(c + d*x) - 1i)*(A + B*1i)*(a*1i - b)^4)/(2*d)$

3.263 $\int \cot^4(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.263.1 Optimal result	2545
3.263.2 Mathematica [C] (verified)	2546
3.263.3 Rubi [A] (verified)	2546
3.263.4 Maple [A] (verified)	2550
3.263.5 Fricas [A] (verification not implemented)	2551
3.263.6 Sympy [A] (verification not implemented)	2551
3.263.7 Maxima [A] (verification not implemented)	2552
3.263.8 Giac [A] (verification not implemented)	2552
3.263.9 Mupad [B] (verification not implemented)	2553

3.263.1 Optimal result

Integrand size = 31, antiderivative size = 187

$$\int \cot^4(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= (a^4A - 6a^2Ab^2 + Ab^4 - 4a^3bB + 4ab^3B) x + \frac{a^2(a^2A - 3Ab^2 - 3abB) \cot(c+dx)}{d}$$

$$- \frac{b^4B \log(\cos(c+dx))}{d} - \frac{a(4a^2Ab - 4Ab^3 + a^3B - 6ab^2B) \log(\sin(c+dx))}{d}$$

$$- \frac{a(2Ab + aB) \cot^2(c+dx)(a+b \tan(c+dx))^2}{2d} - \frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^3}{3d}$$

output

```
(A*a^4-6*A*a^2*b^2+A*b^4-4*B*a^3*b+4*B*a*b^3)*x+a^2*(A*a^2-3*A*b^2-3*B*a*b)*cot(d*x+c)/d-b^4*B*ln(cos(d*x+c))/d-a*(4*A*a^2*b-4*A*b^3+B*a^3-6*B*a*b^2)*ln(sin(d*x+c))/d-1/2*a*(2*A*b+B*a)*cot(d*x+c)^2*(a+b*tan(d*x+c))^2/d-1/3*a*A*cot(d*x+c)^3*(a+b*tan(d*x+c))^3/d
```


3.263.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.17 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.89

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{6a^2(a^2A - 6Ab^2 - 4abB) \cot(c + dx) - 3a^3(4Ab + aB) \cot^2(c + dx) - 2a^4A \cot^3(c + dx) + 3(a + ib)^4(-$$

input `Integrate[Cot[c + d*x]^4*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output $(6a^2(a^2A - 6Ab^2 - 4abB) \cot[c + d*x] - 3a^3(4Ab + aB) \cot[c + d*x]^2 - 2a^4A \cot[c + d*x]^3 + 3(a + I*b)^4((-I)*A + B) \text{Log}[I - \text{Tan}[c + d*x]] - 6a*(4a^2A*b - 4A*b^3 + a^3*B - 6a*b^2*B) \text{Log}[\text{Tan}[c + d*x]] + 3(a - I*b)^4(I*A + B) \text{Log}[I + \text{Tan}[c + d*x]])/(6*d)$

3.263.3 Rubi [A] (verified)

Time = 1.17 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.01, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4118, 3042, 4107, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \tan(c + dx))^4(A + B \tan(c + dx))}{\tan(c + dx)^4} dx$$

$$\downarrow 4088$$

$$\frac{1}{3} \int 3 \cot^3(c + dx)(a + b \tan(c + dx))^2 (b^2 B \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)) dx - \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^3}{3d}$$

$$\downarrow 27$$

$$\begin{aligned}
& \int \cot^3(c+dx)(a+b \tan(c+dx))^2 (b^2 B \tan^2(c+dx) - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2Ab + aB)) dx - \\
& \frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a+b \tan(c+dx))^2 (b^2 B \tan^2(c+dx) - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2Ab + aB))}{\tan(c+dx)^3} dx - \\
& \frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{4128} \\
& \frac{1}{2} \int -2 \cot^2(c+dx)(a+b \tan(c+dx)) (-B \tan^2(c+dx)b^3 + a(Aa^2 - 3bBa - 3Ab^2) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)) dx - \\
& \frac{a(aB + 2Ab) \cot^2(c+dx)(a+b \tan(c+dx))^2}{2d} - \frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{27} \\
& - \int \cot^2(c+dx)(a+b \tan(c+dx)) (-B \tan^2(c+dx)b^3 + a(Aa^2 - 3bBa - 3Ab^2) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)) dx - \\
& \frac{a(aB + 2Ab) \cot^2(c+dx)(a+b \tan(c+dx))^2}{2d} - \frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{3042} \\
& - \int \frac{(a+b \tan(c+dx)) (-B \tan(c+dx)^2 b^3 + a(Aa^2 - 3bBa - 3Ab^2) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx))}{\tan(c+dx)^2} dx - \\
& \frac{a(aB + 2Ab) \cot^2(c+dx)(a+b \tan(c+dx))^2}{2d} - \frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{4118} \\
& - \int \cot(c+dx) (-B \tan^2(c+dx)b^4 + a(Ba^3 + 4Aba^2 - 6b^2Ba - 4Ab^3) - (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4) \tan(c+dx) + \\
& \frac{a^2(a^2A - 3abB - 3Ab^2) \cot(c+dx)}{d} - \frac{a(aB + 2Ab) \cot^2(c+dx)(a+b \tan(c+dx))^2}{2d} - \\
& \frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.263. $\int \cot^4(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& - \int \frac{-B \tan(c+dx)^2 b^4 + a(Ba^3 + 4Aba^2 - 6b^2Ba - 4Ab^3) - (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4) \tan(c+dx)}{\tan(c+dx)} \\
& \frac{a^2(a^2A - 3abB - 3Ab^2) \cot(c+dx)}{d} - \frac{a(aB + 2Ab) \cot^2(c+dx)(a + b \tan(c+dx))^2}{aA \cot^3(c+dx)(a + b \tan(c+dx))^{2d}} \\
& \frac{2d}{3d} \\
& \quad \downarrow \text{4107} \\
& -a(a^3B + 4a^2Ab - 6ab^2B - 4Ab^3) \int \cot(c+dx)dx + b^4B \int \tan(c+dx)dx + \\
& \frac{a^2(a^2A - 3abB - 3Ab^2) \cot(c+dx)}{2d} + x(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) - \\
& \frac{a(aB + 2Ab) \cot^2(c+dx)(a + b \tan(c+dx))^2}{2d} - \frac{aA \cot^3(c+dx)(a + b \tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{3042} \\
& -a(a^3B + 4a^2Ab - 6ab^2B - 4Ab^3) \int -\tan\left(c+dx + \frac{\pi}{2}\right)dx + b^4B \int \tan(c+dx)dx + \\
& \frac{a^2(a^2A - 3abB - 3Ab^2) \cot(c+dx)}{2d} + x(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) - \\
& \frac{a(aB + 2Ab) \cot^2(c+dx)(a + b \tan(c+dx))^2}{2d} - \frac{aA \cot^3(c+dx)(a + b \tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{25} \\
& a(a^3B + 4a^2Ab - 6ab^2B - 4Ab^3) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right)dx + b^4B \int \tan(c+dx)dx + \\
& \frac{a^2(a^2A - 3abB - 3Ab^2) \cot(c+dx)}{2d} + x(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) - \\
& \frac{a(aB + 2Ab) \cot^2(c+dx)(a + b \tan(c+dx))^2}{2d} - \frac{aA \cot^3(c+dx)(a + b \tan(c+dx))^3}{3d} \\
& \quad \downarrow \text{3956} \\
& \frac{a^2(a^2A - 3abB - 3Ab^2) \cot(c+dx)}{d} - \frac{a(a^3B + 4a^2Ab - 6ab^2B - 4Ab^3) \log(-\sin(c+dx))}{d} + \\
& x(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) - \frac{a(aB + 2Ab) \cot^2(c+dx)(a + b \tan(c+dx))^2}{2d} - \\
& \frac{aA \cot^3(c+dx)(a + b \tan(c+dx))^3}{3d} - \frac{b^4B \log(\cos(c+dx))}{d}
\end{aligned}$$

input `Int[Cot[c + d*x]^4*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

```
output (a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*a^3*b*B + 4*a*b^3*B)*x + (a^2*(a^2*A - 3*
A*b^2 - 3*a*b*B)*Cot[c + d*x])/d - (b^4*B*Log[Cos[c + d*x]])/d - (a*(4*a^2
*A*b - 4*A*b^3 + a^3*B - 6*a*b^2*B)*Log[-Sin[c + d*x]])/d - (a*(2*A*b + a*
B)*Cot[c + d*x]^2*(a + b*Tan[c + d*x])^2)/(2*d) - (a*A*Cot[c + d*x]^3*(a +
b*Tan[c + d*x])^3)/(3*d)
```

3.263.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4088 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])^((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x
])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2))
Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*
(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n
+ 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[
e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n
+ 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] &
& LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])
```

```
rule 4107 Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2
]/tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[B*x, x] + (Simp[A Int[1/Tan[
e + f*x], x], x] + Simp[C Int[Tan[e + f*x], x], x]) /; FreeQ[{e, f, A, B,
C}, x] && NeQ[A, C]
```

```
rule 4118 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
)*(x_)]^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)
)*(x_)]^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]
```

```
rule 4128 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

3.263.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.04

method	result
derivativedivides	$\frac{(4A^3b - 4Aab^3 + Ba^4 - 6Ba^2b^2 + Bb^4) \ln(1 + \tan^2(dx+c))}{2} + \frac{(Aa^4 - 6Aa^2b^2 + Ab^4 - 4Ba^3b + 4Bab^3) \arctan(\tan(dx+c)) - \frac{1}{3}}{d}$
default	$\frac{(4A^3b - 4Aab^3 + Ba^4 - 6Ba^2b^2 + Bb^4) \ln(1 + \tan^2(dx+c))}{2} + \frac{(Aa^4 - 6Aa^2b^2 + Ab^4 - 4Ba^3b + 4Bab^3) \arctan(\tan(dx+c)) - \frac{1}{3}}{d}$
parallelrisch	$3(4A^3b - 4Aab^3 + Ba^4 - 6Ba^2b^2 + Bb^4) \ln(\sec^2(dx+c)) + 6(-4A^3b + 4Aab^3 - Ba^4 + 6Ba^2b^2) \ln(\tan(dx+c)) - 2A(\cot(dx+c) - \frac{1}{3})$
norman	$\frac{(Aa^4 - 6Aa^2b^2 + Ab^4 - 4Ba^3b + 4Bab^3)x(\tan^3(dx+c)) + \frac{a^2(Aa^2 - 6Aa^2b^2 - 4Bab)(\tan^2(dx+c))}{d} - \frac{Aa^4}{3d} - \frac{a^3(4Ab + Ba)\tan(dx+c)}{2d}}{\tan(dx+c)^3}$
risch	$-\frac{a^4 \ln(e^{2i(dx+c)} - 1)B}{d} - \frac{\ln(e^{2i(dx+c)} + 1)Bb^4}{d} + Aa^4x - \frac{2ia^2(-6Aa^2e^{4i(dx+c)} + 18Ab^2e^{4i(dx+c)} + 12Babe^{4i(dx+c)})}{d}$

```
input int(cot(d*x+c)^4*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

$$3.263. \quad \int \cot^4(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

output $1/d*(1/2*(4*A*a^3*b-4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*\ln(1+\tan(dx+c)^2)+(A*a^4-6*A*a^2*b^2+A*b^4-4*B*a^3*b+4*B*a*b^3)*\arctan(\tan(dx+c))-1/3*A*a^4/\tan(dx+c)^3-1/2*a^3*(4*A*b+B*a)/\tan(dx+c)^2-a*(4*A*a^2*b-4*A*b^3+B*a^3-6*B*a*b^2)*\ln(\tan(dx+c))+a^2*(A*a^2-6*A*b^2-4*B*a*b)/\tan(dx+c))$

3.263.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.19

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$\frac{3 B b^4 \log\left(\frac{1}{\tan(dx+c)^2+1}\right) \tan(dx+c)^3 + 2 A a^4 + 3 (B a^4 + 4 A a^3 b - 6 B a^2 b^2 - 4 A a b^3) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right)}{d}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output $-1/6*(3*B*b^4*\log(1/(\tan(dx+c)^2+1))*\tan(dx+c)^3+2*A*a^4+3*(B*a^4+4*A*a^3*b-6*B*a^2*b^2-4*A*a*b^3)*\log(\tan(dx+c)^2/(\tan(dx+c)^2+1))*\tan(dx+c)^3+3*(B*a^4+4*A*a^3*b-2*(A*a^4-4*B*a^3*b-6*A*a^2*b^2+4*B*a*b^3+A*b^4)*dx)*\tan(dx+c)^3-6*(A*a^4-4*B*a^3*b-6*A*a^2*b^2)*\tan(dx+c)^2+3*(B*a^4+4*A*a^3*b)*\tan(dx+c))/d*\tan(dx+c)^3$

3.263.6 Sympy [A] (verification not implemented)

Time = 2.29 (sec) , antiderivative size = 369, normalized size of antiderivative = 1.97

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \tilde{\infty} A a^4 x \\ x(A + B \tan(c))(a + b \tan(c))^4 \cot^4(c) \\ \tilde{\infty} A a^4 x \end{cases}$$

$$A a^4 x + \frac{A a^4}{d \tan(c+dx)} - \frac{A a^4}{3d \tan^3(c+dx)} + \frac{2 A a^3 b \log(\tan^2(c+dx)+1)}{d} - \frac{4 A a^3 b \log(\tan(c+dx))}{d} - \frac{2 A a^3 b}{d \tan^2(c+dx)} - 6 A a^2 b^2 x -$$

input `integrate(cot(d*x+c)**4*(a+b*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)`

3.263. $\int \cot^4(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

output `Piecewise((zoo*A*a**4*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**4*cot(c)**4, Eq(d, 0)), (zoo*A*a**4*x, Eq(c, -d*x)), (A*a**4*x + A*a**4/(d*tan(c + d*x)) - A*a**4/(3*d*tan(c + d*x)**3) + 2*A*a**3*b*log(tan(c + d*x)**2 + 1)/d - 4*A*a**3*b*log(tan(c + d*x))/d - 2*A*a**3*b/(d*tan(c + d*x)**2) - 6*A*a**2*b**2*x - 6*A*a**2*b**2/(d*tan(c + d*x)) - 2*A*a*b**3*log(tan(c + d*x)**2 + 1)/d + 4*A*a*b**3*log(tan(c + d*x))/d + A*b**4*x + B*a**4*log(tan(c + d*x)**2 + 1)/(2*d) - B*a**4*log(tan(c + d*x))/d - B*a**4/(2*d*tan(c + d*x)**2) - 4*B*a**3*b*x - 4*B*a**3*b/(d*tan(c + d*x)) - 3*B*a**2*b**2*log(tan(c + d*x)**2 + 1)/d + 6*B*a**2*b**2*log(tan(c + d*x))/d + 4*B*a*b**3*x + B*b**4*log(tan(c + d*x)**2 + 1)/(2*d), True))`

3.263.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.08

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{6(Aa^4 - 4Ba^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4)(dx + c) + 3(Ba^4 + 4Aa^3b - 6Ba^2b^2 - 4Aab^3 + Bb^4) \log(\tan(c + dx))}{1}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/6*(6*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*(d*x + c) + 3*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*log(tan(d*x + c))^2 + 1) - 6*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3)*log(tan(d*x + c)) - (2*A*a^4 - 6*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2)*tan(d*x + c))^2 + 3*(B*a^4 + 4*A*a^3*b)*tan(d*x + c))/tan(d*x + c)^3/d`

3.263.8 Giac [A] (verification not implemented)

Time = 1.51 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.50

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{6(Aa^4 - 4Ba^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4)(dx + c) + 3(Ba^4 + 4Aa^3b - 6Ba^2b^2 - 4Aab^3 + Bb^4) \log(\tan(c + dx))}{1}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output
$$\frac{1}{6} \cdot (6 \cdot (A \cdot a^4 - 4 \cdot B \cdot a^3 \cdot b - 6 \cdot A \cdot a^2 \cdot b^2 + 4 \cdot B \cdot a \cdot b^3 + A \cdot b^4) \cdot (d \cdot x + c) + 3 \cdot (B \cdot a^4 + 4 \cdot A \cdot a^3 \cdot b - 6 \cdot B \cdot a^2 \cdot b^2 - 4 \cdot A \cdot a \cdot b^3 + B \cdot b^4) \cdot \log(\tan(d \cdot x + c)^2 + 1) - 6 \cdot (B \cdot a^4 + 4 \cdot A \cdot a^3 \cdot b - 6 \cdot B \cdot a^2 \cdot b^2 - 4 \cdot A \cdot a \cdot b^3) \cdot \log(\text{abs}(\tan(d \cdot x + c)))) + (11 \cdot B \cdot a^4 \cdot \tan(d \cdot x + c)^3 + 44 \cdot A \cdot a^3 \cdot b \cdot \tan(d \cdot x + c)^3 - 66 \cdot B \cdot a^2 \cdot b^2 \cdot \tan(d \cdot x + c)^3 - 44 \cdot A \cdot a \cdot b^3 \cdot \tan(d \cdot x + c)^3 + 6 \cdot A \cdot a^4 \cdot \tan(d \cdot x + c)^2 - 24 \cdot B \cdot a^3 \cdot b \cdot \tan(d \cdot x + c)^2 - 36 \cdot A \cdot a^2 \cdot b^2 \cdot \tan(d \cdot x + c)^2 - 3 \cdot B \cdot a^4 \cdot \tan(d \cdot x + c) - 12 \cdot A \cdot a^3 \cdot b \cdot \tan(d \cdot x + c) - 2 \cdot A \cdot a^4) / \tan(d \cdot x + c)^3) / d$$

3.263.9 Mupad [B] (verification not implemented)

Time = 8.02 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.95

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= -\frac{\ln(\tan(c + dx))(B a^4 + 4 A a^3 b - 6 B a^2 b^2 - 4 A a b^3)}{d}$$

$$-\frac{\cot(c + dx)^3 \left(\tan(c + dx) \left(\frac{B a^4}{2} + 2 A b a^3 \right) + \frac{A a^4}{3} + \tan(c + dx)^2 (-A a^4 + 4 B a^3 b + 6 A a^2 b^2) \right)}{d}$$

$$-\frac{\ln(\tan(c + dx) - i) (-B + A i) (-b + a i)^4}{2 d}$$

$$+\frac{\ln(\tan(c + dx) + i) (B + A i) (b + a i)^4}{2 d}$$

input `int(cot(c + d*x)^4*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^4,x)`

output
$$(\log(\tan(c + d \cdot x) + 1i) \cdot (A \cdot 1i + B) \cdot (a \cdot 1i + b)^4) / (2 \cdot d) - (\cot(c + d \cdot x)^3 \cdot (\tan(c + d \cdot x) \cdot ((B \cdot a^4) / 2 + 2 \cdot A \cdot a^3 \cdot b) + (A \cdot a^4) / 3 + \tan(c + d \cdot x)^2 \cdot (6 \cdot A \cdot a^2 \cdot b^2 - A \cdot a^4 + 4 \cdot B \cdot a^3 \cdot b))) / d - (\log(\tan(c + d \cdot x) - 1i) \cdot (A \cdot 1i - B) \cdot (a \cdot 1i - b)^4) / (2 \cdot d) - (\log(\tan(c + d \cdot x)) \cdot (B \cdot a^4 - 6 \cdot B \cdot a^2 \cdot b^2 - 4 \cdot A \cdot a \cdot b^3 + 4 \cdot A \cdot a^3 \cdot b)) / d$$

3.264 $\int \cot^5(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.264.1 Optimal result	2554
3.264.2 Mathematica [C] (verified)	2555
3.264.3 Rubi [A] (verified)	2555
3.264.4 Maple [A] (verified)	2560
3.264.5 Fricas [A] (verification not implemented)	2561
3.264.6 Sympy [B] (verification not implemented)	2561
3.264.7 Maxima [A] (verification not implemented)	2562
3.264.8 Giac [B] (verification not implemented)	2563
3.264.9 Mupad [B] (verification not implemented)	2564

3.264.1 Optimal result

Integrand size = 31, antiderivative size = 225

$$\int \cot^5(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= (4a^3Ab - 4aAb^3 + a^4B - 6a^2b^2B + b^4B) x$$

$$+ \frac{a(24a^2Ab - 19Ab^3 + 6a^3B - 34ab^2B) \cot(c+dx)}{6d}$$

$$+ \frac{a^2(6a^2A - 13Ab^2 - 16abB) \cot^2(c+dx)}{12d}$$

$$+ \frac{(a^4A - 6a^2Ab^2 + Ab^4 - 4a^3bB + 4ab^3B) \log(\sin(c+dx))}{12d}$$

$$- \frac{a(7Ab + 4aB) \cot^3(c+dx)(a+b \tan(c+dx))^2}{12d} - \frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^3}{4d}$$

output

```
(4*A*a^3*b-4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*x+1/6*a*(24*A*a^2*b-19*A*b^3+6*B*a^3-34*B*a*b^2)*cot(d*x+c)/d+1/12*a^2*(6*A*a^2-13*A*b^2-16*B*a*b)*cot(d*x+c)^2/d+(A*a^4-6*A*a^2*b^2+A*b^4-4*B*a^3*b+4*B*a*b^3)*ln(sin(d*x+c))/d-1/12*a*(7*A*b+4*B*a)*cot(d*x+c)^3*(a+b*tan(d*x+c))^2/d-1/4*a*A*cot(d*x+c)^4*(a+b*tan(d*x+c))^3/d
```

3.264.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.99 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.94

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{12a(4a^2Ab - 4Ab^3 + a^3B - 6ab^2B) \cot(c + dx) + 6a^2(a^2A - 6Ab^2 - 4abB) \cot^2(c + dx) - 4a^3(4Ab + aB) \cot^3(c + dx) + 6a^4(4Ab + aB) \cot^4(c + dx) - 6a^5(4Ab + aB) \cot^5(c + dx)}{12d}$$

input `Integrate[Cot[c + d*x]^5*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output $(12*a*(4*a^2*A*b - 4*A*b^3 + a^3*B - 6*a*b^2*B)*\text{Cot}[c + d*x] + 6*a^2*(a^2*A - 6*A*b^2 - 4*a*b*B)*\text{Cot}[c + d*x]^2 - 4*a^3*(4*A*b + a*B)*\text{Cot}[c + d*x]^3 - 3*a^4*A*\text{Cot}[c + d*x]^4 - 6*(a + I*b)^4*(A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]] + 12*(a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*a^3*b*B + 4*a*b^3*B)*\text{Log}[\text{Tan}[c + d*x]] - 6*(a - I*b)^4*(A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]])/(12*d)$

3.264.3 Rubi [A] (verified)

Time = 1.55 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.06, number of steps used = 15, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 4088, 3042, 4128, 27, 3042, 4118, 3042, 4111, 27, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^4(A + B \tan(c + dx))}{\tan(c + dx)^5} dx$$

$$\downarrow \text{4088}$$

$$\frac{1}{4} \int \cot^4(c + dx)(a + b \tan(c + dx))^2 (-b(aA - 4bB) \tan^2(c + dx) - 4(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(7Ab + 4aB)) dx - \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^3}{4d}$$

$$\downarrow \text{3042}$$

$$\frac{1}{4} \int \frac{(a + b \tan(c + dx))^2 (-b(aA - 4bB) \tan(c + dx)^2 - 4(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(7Ab + 4aB))}{\tan(c + dx)^4} dx - \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^3}{4d}$$

↓ 4128

$$\frac{1}{4} \left(\frac{1}{3} \int -2 \cot^3(c + dx)(a + b \tan(c + dx)) (b(2Ba^2 + 5Aba - 6b^2B) \tan^2(c + dx) + 6(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)) \right) \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^3}{4d}$$

↓ 27

$$\frac{1}{4} \left(-\frac{2}{3} \int \cot^3(c + dx)(a + b \tan(c + dx)) (b(2Ba^2 + 5Aba - 6b^2B) \tan^2(c + dx) + 6(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)) \right) \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^3}{4d}$$

↓ 3042

$$\frac{1}{4} \left(-\frac{2}{3} \int \frac{(a + b \tan(c + dx)) (b(2Ba^2 + 5Aba - 6b^2B) \tan(c + dx)^2 + 6(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(7Ab + 4aB))}{\tan(c + dx)^3} \right) \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^3}{4d}$$

↓ 4118

$$\frac{1}{4} \left(-\frac{2}{3} \left(\int \cot^2(c + dx) (b^2(2Ba^2 + 5Aba - 6b^2B) \tan^2(c + dx) - 6(Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4) \tan(c + dx) + a(7Ab + 4aB)) \right) \right) \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^3}{4d}$$

↓ 3042

$$\frac{1}{4} \left(-\frac{2}{3} \left(\int \frac{b^2(2Ba^2 + 5Aba - 6b^2B) \tan(c + dx)^2 - 6(Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4) \tan(c + dx) + a(7Ab + 4aB)}{\tan(c + dx)^2} \right) \right) \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^3}{4d}$$

↓ 4111

$$\frac{1}{4} \left(-\frac{2}{3} \left(\int -6 \cot(c+dx) (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4 + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \tan(c+dx)) dx \right. \right. \\ \left. \left. \frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^3}{4d} \right) \right. \\ \left. \downarrow 27 \right.$$

$$\frac{1}{4} \left(-\frac{2}{3} \left(-6 \int \cot(c+dx) (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4 + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \tan(c+dx)) dx \right. \right. \\ \left. \left. \frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^3}{4d} \right) \right. \\ \left. \downarrow 3042 \right.$$

$$\frac{1}{4} \left(-\frac{2}{3} \left(-6 \int \frac{Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4 + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \tan(c+dx)}{\tan(c+dx)} dx \right. \right. \\ \left. \left. \frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^3}{4d} \right) \right. \\ \left. \downarrow 4014 \right.$$

$$\frac{1}{4} \left(-\frac{2}{3} \left(-6 \left((a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) \int \cot(c+dx) dx + x(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) \right) \right. \right. \\ \left. \left. \frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^3}{4d} \right) \right. \\ \left. \downarrow 3042 \right.$$

$$\frac{1}{4} \left(-\frac{2}{3} \left(-6 \left((a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx + x(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) \right) \right. \right. \\ \left. \left. \frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^3}{4d} \right) \right. \\ \left. \downarrow 25 \right.$$

$$\frac{1}{4} \left(-\frac{2}{3} \left(-6 \left(x(a^4B + 4a^3Ab - 6a^2b^2B - 4aAb^3 + b^4B) - (a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) \int \tan\left(\frac{1}{2}(2c+dx)\right) dx \right) \right. \right. \\ \left. \left. \frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^3}{4d} \right) \right. \\ \left. \downarrow 3956 \right.$$

$$\frac{1}{4} \left(-\frac{2}{3} \left(-\frac{a^2(6a^2A - 16abB - 13Ab^2) \cot^2(c + dx)}{2d} - \frac{a(6a^3B + 24a^2Ab - 34ab^2B - 19Ab^3) \cot(c + dx)}{d} - 6 \left(\frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^3}{4d} \right) \right) \right)$$

input `Int[Cot[c + d*x]^5*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `-1/4*(a*A*Cot[c + d*x]^4*(a + b*Tan[c + d*x])^3)/d + ((-2*(-((a*(24*a^2*A*b - 19*A*b^3 + 6*a^3*B - 34*a*b^2*B)*Cot[c + d*x])/d) - (a^2*(6*a^2*A - 13*A*b^2 - 16*a*b*B)*Cot[c + d*x]^2)/(2*d) - 6*((4*a^3*A*b - 4*a*A*b^3 + a^4*B - 6*a^2*b^2*B + b^4*B)*x + ((a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*a^3*b*B + 4*a*b^3*B)*Log[-Sin[c + d*x]]/d)))/3 - (a*(7*A*b + 4*a*B)*Cot[c + d*x]^3*(a + b*Tan[c + d*x])^2)/(3*d))/4`

3.264.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

```
rule 4088 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x
])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2))
Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*
(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n
+ 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[
e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n
+ 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] &
& LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])
```

```
rule 4111 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]
```

```
rule 4118 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.
)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-(b*c - a*d))*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]
```

```
rule 4128 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2, x_Symbol] :> Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

3.264.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.07

method	result
derivativedivides	$\frac{(-A a^4 + 6A a^2 b^2 - A b^4 + 4B a^3 b - 4B a b^3) \ln(1 + \tan^2(dx+c))}{2} + (4A a^3 b - 4A a b^3 + B a^4 - 6B a^2 b^2 + B b^4) \arctan(\tan(dx+c)) +$
default	$\frac{(-A a^4 + 6A a^2 b^2 - A b^4 + 4B a^3 b - 4B a b^3) \ln(1 + \tan^2(dx+c))}{2} + (4A a^3 b - 4A a b^3 + B a^4 - 6B a^2 b^2 + B b^4) \arctan(\tan(dx+c)) +$
parallelrisch	$6(-A a^4 + 6A a^2 b^2 - A b^4 + 4B a^3 b - 4B a b^3) \ln(\sec^2(dx+c)) + 12(A a^4 - 6A a^2 b^2 + A b^4 - 4B a^3 b + 4B a b^3) \ln(\tan(dx+c)) -$
norman	$(4A a^3 b - 4A a b^3 + B a^4 - 6B a^2 b^2 + B b^4) x (\tan^4(dx+c)) + \frac{a(4A a^2 b - 4A b^3 + B a^3 - 6B a b^2)}{d} \frac{(\tan^3(dx+c))}{\tan(dx+c)^4} - \frac{A a^4}{4d} + \frac{a^2(A a^2 - 6B a b^2 + B b^4)}{d \tan(dx+c)^4}$
risch	$\frac{8iB a^3 b c}{d} - iA a^4 x - 4iB a b^3 x - \frac{4ia(8A a^2 b - 9B a b^2 + 2B a^3 - 3iA a^3 e^{2i(dx+c)} - 3iA a^3 e^{6i(dx+c)} + 3iA a^3 e^{4i(dx+c)})}{d}$

```
input int(cot(d*x+c)^5*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)), x, method=_RETURNVERBOSE)
```

```
output 1/d*(1/2*(-A*a^4+6*A*a^2*b^2-A*b^4+4*B*a^3*b-4*B*a*b^3)*ln(1+tan(d*x+c)^2)
+(4*A*a^3*b-4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*arctan(tan(d*x+c))+(A*a^4-6
*A*a^2*b^2+A*b^4-4*B*a^3*b+4*B*a*b^3)*ln(tan(d*x+c))-1/4*A*a^4/tan(d*x+c)^
4-1/3*a^3*(4*A*b+B*a)/tan(d*x+c)^3+a*(4*A*a^2*b-4*A*b^3+B*a^3-6*B*a*b^2)/t
an(d*x+c)+1/2*a^2*(A*a^2-6*A*b^2-4*B*a*b)/tan(d*x+c)^2)
```

3.264. $\int \cot^5(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

3.264.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.11

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{6(Aa^4 - 4Ba^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^4 - 3Aa^4 + 3(3Aa^4 - 8Ba^3b$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/12*(6*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^4 - 3*A*a^4 + 3*(3*A*a^4 - 8*B*a^3*b - 12*A*a^2*b^2 + 4*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*d*x)*tan(d*x + c)^4 + 12*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3) *tan(d*x + c)^3 + 6*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2)*tan(d*x + c)^2 - 4*(B*a^4 + 4*A*a^3*b)*tan(d*x + c))/(d*tan(d*x + c)^4)`

3.264.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 459 vs. 2(226) = 452.

Time = 4.17 (sec) , antiderivative size = 459, normalized size of antiderivative = 2.04

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \tilde{\infty}Aa^4x \\ x(A + B \tan(c))(a + b \tan(c))^4 \cot^5(c) \\ \tilde{\infty}Aa^4x \\ -\frac{Aa^4 \log(\tan^2(c+dx)+1)}{2d} + \frac{Aa^4 \log(\tan(c+dx))}{d} + \frac{Aa^4}{2d \tan^2(c+dx)} - \frac{Aa^4}{4d \tan^4(c+dx)} + 4Aa^3bx + \frac{4Aa^3b}{d \tan(c+dx)} - \frac{4Aa^3b}{3d \tan^3(c+dx)} \end{cases}$$

input `integrate(cot(d*x+c)**5*(a+b*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)`


```
output Piecewise((zoo*A*a**4*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**4*cot(c)**5, Eq(d, 0)), (zoo*A*a**4*x, Eq(c, -d*x)), (-A*a**4*log(tan(c + d*x)**2 + 1)/(2*d) + A*a**4*log(tan(c + d*x))/d + A*a**4/(2*d*tan(c + d*x)**2) - A*a**4/(4*d*tan(c + d*x)**4) + 4*A*a**3*b*x + 4*A*a**3*b/(d*tan(c + d*x)) - 4*A*a**3*b/(3*d*tan(c + d*x)**3) + 3*A*a**2*b**2*log(tan(c + d*x)**2 + 1)/d - 6*A*a**2*b**2*log(tan(c + d*x))/d - 3*A*a**2*b**2/(d*tan(c + d*x)**2) - 4*A*a*b**3*x - 4*A*a*b**3/(d*tan(c + d*x)) - A*b**4*log(tan(c + d*x)**2 + 1)/(2*d) + A*b**4*log(tan(c + d*x))/d + B*a**4*x + B*a**4/(d*tan(c + d*x)) - B*a**4/(3*d*tan(c + d*x)**3) + 2*B*a**3*b*log(tan(c + d*x)**2 + 1)/d - 4*B*a**3*b*log(tan(c + d*x))/d - 2*B*a**3*b/(d*tan(c + d*x)**2) - 6*B*a**2*b**2*x - 6*B*a**2*b**2/(d*tan(c + d*x)) - 2*B*a*b**3*log(tan(c + d*x)**2 + 1)/d + 4*B*a*b**3*log(tan(c + d*x))/d + B*b**4*x, True))
```

3.264.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.09

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{12(Ba^4 + 4Aa^3b - 6Ba^2b^2 - 4Aab^3 + Bb^4)(dx + c) - 6(Aa^4 - 4Ba^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4) \log(\tan(c + dx)^2 + 1)}{d}$$

```
input integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")
```

```
output 1/12*(12*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*(d*x + c) - 6*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)^2 + 1) + 12*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)) - (3*A*a^4 - 12*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3)*tan(d*x + c)^3 - 6*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2)*tan(d*x + c)^2 + 4*(B*a^4 + 4*A*a^3*b)*tan(d*x + c))/tan(d*x + c)^4/d
```

3.264.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 584 vs. $2(217) = 434$.

Time = 1.68 (sec) , antiderivative size = 584, normalized size of antiderivative = 2.60

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$3 Aa^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 - 8 Ba^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 32 Aa^3 b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 - 36 Aa^4 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

```
-1/192*(3*A*a^4*tan(1/2*d*x + 1/2*c)^4 - 8*B*a^4*tan(1/2*d*x + 1/2*c)^3 -
32*A*a^3*b*tan(1/2*d*x + 1/2*c)^3 - 36*A*a^4*tan(1/2*d*x + 1/2*c)^2 + 96*B
*a^3*b*tan(1/2*d*x + 1/2*c)^2 + 144*A*a^2*b^2*tan(1/2*d*x + 1/2*c)^2 + 120
*B*a^4*tan(1/2*d*x + 1/2*c) + 480*A*a^3*b*tan(1/2*d*x + 1/2*c) - 576*B*a^2
*b^2*tan(1/2*d*x + 1/2*c) - 384*A*a*b^3*tan(1/2*d*x + 1/2*c) - 192*(B*a^4
+ 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*(d*x + c) + 192*(A*a^4 - 4*
B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*log(tan(1/2*d*x + 1/2*c)^2 + 1)
- 192*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*log(abs(tan(1
/2*d*x + 1/2*c))) + (400*A*a^4*tan(1/2*d*x + 1/2*c)^4 - 1600*B*a^3*b*tan(1
/2*d*x + 1/2*c)^4 - 2400*A*a^2*b^2*tan(1/2*d*x + 1/2*c)^4 + 1600*B*a*b^3*t
an(1/2*d*x + 1/2*c)^4 + 400*A*b^4*tan(1/2*d*x + 1/2*c)^4 - 120*B*a^4*tan(1
/2*d*x + 1/2*c)^3 - 480*A*a^3*b*tan(1/2*d*x + 1/2*c)^3 + 576*B*a^2*b^2*tan
(1/2*d*x + 1/2*c)^3 + 384*A*a*b^3*tan(1/2*d*x + 1/2*c)^3 - 36*A*a^4*tan(1/
2*d*x + 1/2*c)^2 + 96*B*a^3*b*tan(1/2*d*x + 1/2*c)^2 + 144*A*a^2*b^2*tan(1
/2*d*x + 1/2*c)^2 + 8*B*a^4*tan(1/2*d*x + 1/2*c) + 32*A*a^3*b*tan(1/2*d*x
+ 1/2*c) + 3*A*a^4)/tan(1/2*d*x + 1/2*c)^4)/d
```

3.264.9 Mupad [B] (verification not implemented)

Time = 8.51 (sec) , antiderivative size = 218, normalized size of antiderivative = 0.97

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{\ln(\tan(c + dx)) (Aa^4 - 4Ba^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4)}{d} - \frac{\cot(c + dx)^4 \left(\tan(c + dx) \left(\frac{Ba^4}{3} + \frac{4Aba^3}{3} \right) + \frac{Aa^4}{4} - \tan(c + dx)^3 (Ba^4 + 4Aa^3b - 6Ba^2b^2 - 4Aab^3 + Ab^4) \right)}{d}$$

$$- \frac{\ln(\tan(c + dx) + i) (A - Bi) (b + ai)^4}{2d}$$

$$- \frac{\ln(\tan(c + dx) - i) (A + Bi) (-b + ai)^4}{2d}$$

input `int(cot(c + d*x)^5*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^4,x)`output `(log(tan(c + d*x))*(A*a^4 + A*b^4 - 6*A*a^2*b^2 + 4*B*a*b^3 - 4*B*a^3*b))/d - (cot(c + d*x)^4*(tan(c + d*x)*((B*a^4)/3 + (4*A*a^3*b)/3) + (A*a^4)/4 - tan(c + d*x)^3*(B*a^4 - 6*B*a^2*b^2 - 4*A*a*b^3 + 4*A*a^3*b) + tan(c + d*x)^2*(3*A*a^2*b^2 - (A*a^4)/2 + 2*B*a^3*b))/d - (log(tan(c + d*x) + i)*(A - B*i)*(a*i + b)^4)/(2*d) - (log(tan(c + d*x) - i)*(A + B*i)*(a*i - b)^4)/(2*d)`

3.265 $\int \cot^6(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.265.1 Optimal result	2565
3.265.2 Mathematica [C] (verified)	2566
3.265.3 Rubi [A] (verified)	2566
3.265.4 Maple [A] (verified)	2572
3.265.5 Fricas [A] (verification not implemented)	2572
3.265.6 Sympy [A] (verification not implemented)	2573
3.265.7 Maxima [A] (verification not implemented)	2574
3.265.8 Giac [B] (verification not implemented)	2574
3.265.9 Mupad [B] (verification not implemented)	2575

3.265.1 Optimal result

Integrand size = 31, antiderivative size = 273

$$\int \cot^6(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= -((a^4A - 6a^2Ab^2 + Ab^4 - 4a^3bB + 4ab^3B) x$$

$$- \frac{(a^4A - 6a^2Ab^2 + Ab^4 - 4a^3bB + 4ab^3B) \cot(c+dx)}{d}$$

$$+ \frac{a(40a^2Ab - 28Ab^3 + 10a^3B - 55ab^2B) \cot^2(c+dx)}{20d}$$

$$+ \frac{a^2(10a^2A - 18Ab^2 - 25abB) \cot^3(c+dx)}{30d}$$

$$+ \frac{(4a^3Ab - 4aAb^3 + a^4B - 6a^2b^2B + b^4B) \log(\sin(c+dx))}{d}$$

$$- \frac{a(8Ab + 5aB) \cot^4(c+dx)(a+b \tan(c+dx))^2}{20d} - \frac{aA \cot^5(c+dx)(a+b \tan(c+dx))^3}{5d}$$

output

```
-(A*a^4-6*A*a^2*b^2+A*b^4-4*B*a^3*b+4*B*a*b^3)*x-(A*a^4-6*A*a^2*b^2+A*b^4-4*B*a^3*b+4*B*a*b^3)*cot(d*x+c)/d+1/20*a*(40*A*a^2*b-28*A*b^3+10*B*a^3-55*B*a*b^2)*cot(d*x+c)^2/d+1/30*a^2*(10*A*a^2-18*A*b^2-25*B*a*b)*cot(d*x+c)^3/d+(4*A*a^3*b-4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*ln(sin(d*x+c))/d-1/20*a*(8*A*b+5*B*a)*cot(d*x+c)^4*(a+b*tan(d*x+c))^2/d-1/5*a*A*cot(d*x+c)^5*(a+b*tan(d*x+c))^3/d
```

3.265.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.72 (sec) , antiderivative size = 257, normalized size of antiderivative = 0.94

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{-60(a^4A - 6a^2Ab^2 + Ab^4 - 4a^3bB + 4ab^3B) \cot(c + dx) + 30a(4a^2Ab - 4Ab^3 + a^3B - 6ab^2B) \cot^2(c + dx) + \dots}{60}$$

input `Integrate[Cot[c + d*x]^6*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(-60*(a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*a^3*b*B + 4*a*b^3*B)*Cot[c + d*x] + 30*a*(4*a^2*A*b - 4*A*b^3 + a^3*B - 6*a*b^2*B)*Cot[c + d*x]^2 + 20*a^2*(a^2*A - 6*A*b^2 - 4*a*b*B)*Cot[c + d*x]^3 - 15*a^3*(4*A*b + a*B)*Cot[c + d*x]^4 - 12*a^4*A*Cot[c + d*x]^5 + (30*I)*(a + I*b)^4*(A + I*B)*Log[I - Tan[c + d*x]] + 60*(4*a^3*A*b - 4*a*A*b^3 + a^4*B - 6*a^2*b^2*B + b^4*B)*Log[Tan[c + d*x]] - 30*(a - I*b)^4*(I*A + B)*Log[I + Tan[c + d*x]])/(60*d)`

3.265.3 Rubi [A] (verified)

Time = 1.91 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.05, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.548$, Rules used = {3042, 4088, 3042, 4128, 27, 3042, 4118, 3042, 4111, 27, 3042, 4012, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^4(A + B \tan(c + dx))}{\tan(c + dx)^6} dx$$

$$\downarrow \text{4088}$$

$$\frac{1}{5} \int \cot^5(c+dx)(a+b \tan(c+dx))^2 (-b(2aA-5bB) \tan^2(c+dx) - 5(Aa^2-2bBa-Ab^2) \tan(c+dx) + a(8Ab+5aB)) dx - \frac{aA \cot^5(c+dx)(a+b \tan(c+dx))^3}{5d}$$

↓ 3042

$$\frac{1}{5} \int \frac{(a+b \tan(c+dx))^2 (-b(2aA-5bB) \tan(c+dx)^2 - 5(Aa^2-2bBa-Ab^2) \tan(c+dx) + a(8Ab+5aB))}{\tan(c+dx)^5} dx - \frac{aA \cot^5(c+dx)(a+b \tan(c+dx))^3}{5d}$$

↓ 4128

$$\frac{1}{5} \left(\frac{1}{4} \int -2 \cot^4(c+dx)(a+b \tan(c+dx)) (b(5Ba^2+12Aba-10b^2B) \tan^2(c+dx) + 10(Ba^3+3Aba^2-3b^2Ba)) dx - \frac{aA \cot^5(c+dx)(a+b \tan(c+dx))^3}{5d} \right)$$

↓ 27

$$\frac{1}{5} \left(-\frac{1}{2} \int \cot^4(c+dx)(a+b \tan(c+dx)) (b(5Ba^2+12Aba-10b^2B) \tan^2(c+dx) + 10(Ba^3+3Aba^2-3b^2Ba)) dx - \frac{aA \cot^5(c+dx)(a+b \tan(c+dx))^3}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(-\frac{1}{2} \int \frac{(a+b \tan(c+dx)) (b(5Ba^2+12Aba-10b^2B) \tan(c+dx)^2 + 10(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx))}{\tan(c+dx)^4} dx - \frac{aA \cot^5(c+dx)(a+b \tan(c+dx))^3}{5d} \right)$$

↓ 4118

$$\frac{1}{5} \left(\frac{1}{2} \left(\frac{a^2(10a^2A-25abB-18Ab^2) \cot^3(c+dx)}{3d} - \int \cot^3(c+dx) (b^2(5Ba^2+12Aba-10b^2B) \tan^2(c+dx) - 10(Ba^3+3Aba^2-3b^2Ba)) dx \right) - \frac{aA \cot^5(c+dx)(a+b \tan(c+dx))^3}{5d} \right)$$

↓ 3042

3.265. $\int \cot^6(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

$$\frac{1}{5} \left(\frac{1}{2} \left(\frac{a^2(10a^2A - 25abB - 18Ab^2) \cot^3(c + dx)}{3d} - \int \frac{b^2(5Ba^2 + 12Aba - 10b^2B) \tan(c + dx)^2 - 10(Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4 + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \tan(c + dx))}{5d} dx + \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^3}{5d} \right) \right)$$

↓ 4111

$$\frac{1}{5} \left(\frac{1}{2} \left(- \int -10 \cot^2(c + dx) (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4 + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \tan(c + dx))}{5d} dx + \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^3}{5d} \right) \right)$$

↓ 27

$$\frac{1}{5} \left(\frac{1}{2} \left(10 \int \cot^2(c + dx) (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4 + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \tan(c + dx))}{5d} dx + \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^3}{5d} \right) \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{2} \left(10 \int \frac{Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4 + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \tan(c + dx)}{\tan(c + dx)^2} dx + \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^3}{5d} \right) \right)$$

↓ 4012

$$\frac{1}{5} \left(\frac{1}{2} \left(10 \left(\int \cot(c + dx) (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B - (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4) \tan(c + dx))}{5d} dx + \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^3}{5d} \right) \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{1}{2} \left(10 \left(\int \frac{Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B - (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4) \tan(c + dx)}{\tan(c + dx)} dx + \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^3}{5d} \right) \right)$$

↓ 4014

3.265. $\int \cot^6(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

$$\frac{1}{5} \left(\frac{1}{2} \left(10 \left((a^4 B + 4a^3 Ab - 6a^2 b^2 B - 4aAb^3 + b^4 B) \int \cot(c + dx) dx - \frac{(a^4 A - 4a^3 bB - 6a^2 Ab^2 + 4ab^3 B + Ab^4)}{d} \right. \right. \right. \\ \left. \left. \left. \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^3}{5d} \right) \right) \right. \\ \left. \downarrow 3042 \right.$$

$$\frac{1}{5} \left(\frac{1}{2} \left(10 \left((a^4 B + 4a^3 Ab - 6a^2 b^2 B - 4aAb^3 + b^4 B) \int -\tan\left(c + dx + \frac{\pi}{2}\right) dx - \frac{(a^4 A - 4a^3 bB - 6a^2 Ab^2 + 4ab^3 B + Ab^4)}{d} \right. \right. \right. \\ \left. \left. \left. \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^3}{5d} \right) \right) \right. \\ \left. \downarrow 25 \right.$$

$$\frac{1}{5} \left(\frac{1}{2} \left(10 \left(-(a^4 B + 4a^3 Ab - 6a^2 b^2 B - 4aAb^3 + b^4 B) \int \tan\left(\frac{1}{2}(2c + \pi) + dx\right) dx - \frac{(a^4 A - 4a^3 bB - 6a^2 Ab^2 + 4ab^3 B + Ab^4)}{d} \right. \right. \right. \\ \left. \left. \left. \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^3}{5d} \right) \right) \right. \\ \left. \downarrow 3956 \right.$$

$$\frac{1}{5} \left(\frac{1}{2} \left(\frac{a^2(10a^2 A - 25abB - 18Ab^2) \cot^3(c + dx)}{3d} + \frac{a(10a^3 B + 40a^2 Ab - 55ab^2 B - 28Ab^3) \cot^2(c + dx)}{2d} + 10 \left(- \right. \right. \right. \\ \left. \left. \left. \frac{aA \cot^5(c + dx)(a + b \tan(c + dx))^3}{5d} \right) \right) \right.$$

input `Int[Cot[c + d*x]^6*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `-1/5*(a*A*Cot[c + d*x]^5*(a + b*Tan[c + d*x])^3)/d + (((a*(40*a^2*A*b - 28*A*b^3 + 10*a^3*B - 55*a*b^2*B)*Cot[c + d*x]^2)/(2*d) + (a^2*(10*a^2*A - 18*A*b^2 - 25*a*b*B)*Cot[c + d*x]^3)/(3*d) + 10*(-((a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*a^3*b*B + 4*a*b^3*B)*x) - ((a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*a^3*b*B + 4*a*b^3*B)*Cot[c + d*x])/d + ((4*a^3*A*b - 4*a*A*b^3 + a^4*B - 6*a^2*b^2*B + b^4*B)*Log[-Sin[c + d*x]])/d))/2 - (a*(8*A*b + 5*a*B)*Cot[c + d*x]^4*(a + b*Tan[c + d*x])^2)/(4*d))/5`

3.265.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`
- rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`
- rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4111 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

rule 4118 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(-(b*c - a*d))*(c^2*C - B*c*d + A*d^2)*((c + d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

3.265.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.04

method	result
parallelrisch	$(-120A a^3 b + 120A a b^3 - 30B a^4 + 180B a^2 b^2 - 30B b^4) \ln(\sec^2(dx+c)) + (240A a^3 b - 240A a b^3 + 60B a^4 - 360B a^2 b^2 + 60B b^4) \ln(\tan(dx+c))$
derivativedivides	$\frac{(-4A a^3 b + 4A a b^3 - B a^4 + 6B a^2 b^2 - B b^4) \ln(1 + \tan^2(dx+c))}{2} + (-A a^4 + 6A a^2 b^2 - A b^4 + 4B a^3 b - 4B a b^3) \arctan(\tan(dx+c))$
default	$\frac{(-4A a^3 b + 4A a b^3 - B a^4 + 6B a^2 b^2 - B b^4) \ln(1 + \tan^2(dx+c))}{2} + (-A a^4 + 6A a^2 b^2 - A b^4 + 4B a^3 b - 4B a b^3) \arctan(\tan(dx+c))$
norman	$(-A a^4 + 6A a^2 b^2 - A b^4 + 4B a^3 b - 4B a b^3) x (\tan^5(dx+c)) - \frac{A a^4}{5d} - \frac{(A a^4 - 6A a^2 b^2 + A b^4 - 4B a^3 b + 4B a b^3) (\tan^4(dx+c))}{d} + \frac{a}{\tan(dx+c)^5}$
risch	$\frac{a^4 \ln(e^{2i(dx+c)} - 1) B}{d} + \frac{12iB a^2 b^2 c}{d} - \frac{2ia^4 Bc}{d} - \frac{2iB b^4 c}{d} - iB b^4 x + \frac{8iA a b^3 c}{d} - A a^4 x - iB a^4 x - \frac{4a}{\tan(dx+c)^5}$

input `int(cot(d*x+c)^6*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/60*((-120*A*a^3*b+120*A*a*b^3-30*B*a^4+180*B*a^2*b^2-30*B*b^4)*ln(sec(d*x+c)^2)+(240*A*a^3*b-240*A*a*b^3+60*B*a^4-360*B*a^2*b^2+60*B*b^4)*ln(tan(d*x+c))-12*A*cot(d*x+c)^5*a^4+(-60*A*a^3*b-15*B*a^4)*cot(d*x+c)^4+20*a^2*cot(d*x+c)^3*(A*a^2-6*A*b^2-4*B*a*b)+(120*A*a^3*b-120*A*a*b^3+30*B*a^4-180*B*a^2*b^2)*cot(d*x+c)^2+(-60*A*a^4+360*A*a^2*b^2-60*A*b^4+240*B*a^3*b-240*B*a*b^3)*cot(d*x+c)-60*d*x*(A*a^4-6*A*a^2*b^2+A*b^4-4*B*a^3*b+4*B*a*b^3))/d`

3.265.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.10

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{30(Ba^4 + 4Aa^3b - 6Ba^2b^2 - 4Aab^3 + Bb^4) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^5 + 15(3Ba^4 + 12Aa^3b - 12Aa^2b^2 - 4Aab^3 + Bb^4) \tan(dx+c)^4 + 15(3Ba^4 + 12Aa^3b - 12Aa^2b^2 - 4Aab^3 + Bb^4) \tan(dx+c)^3 + 15(3Ba^4 + 12Aa^3b - 12Aa^2b^2 - 4Aab^3 + Bb^4) \tan(dx+c)^2 + 15(3Ba^4 + 12Aa^3b - 12Aa^2b^2 - 4Aab^3 + Bb^4) \tan(dx+c) + 15(3Ba^4 + 12Aa^3b - 12Aa^2b^2 - 4Aab^3 + Bb^4)}{d}$$

input `integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fracas")`

```
output 1/60*(30*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*log(tan(d*x
+ c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^5 + 15*(3*B*a^4 + 12*A*a^3*b -
12*B*a^2*b^2 - 8*A*a*b^3 - 4*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3
+ A*b^4)*d*x)*tan(d*x + c)^5 - 12*A*a^4 - 60*(A*a^4 - 4*B*a^3*b - 6*A*a^2*
b^2 + 4*B*a*b^3 + A*b^4)*tan(d*x + c)^4 + 30*(B*a^4 + 4*A*a^3*b - 6*B*a^2*
b^2 - 4*A*a*b^3)*tan(d*x + c)^3 + 20*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2)*tan
(d*x + c)^2 - 15*(B*a^4 + 4*A*a^3*b)*tan(d*x + c))/(d*tan(d*x + c)^5)
```

3.265.6 Sympy [A] (verification not implemented)

Time = 5.55 (sec) , antiderivative size = 546, normalized size of antiderivative = 2.00

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \begin{cases} \tilde{\infty} A a^4 x \\ x(A + B \tan(c))(a + b \tan(c))^4 \cot^6(c) \\ \tilde{\infty} A a^4 x \\ -A a^4 x - \frac{A a^4}{d \tan(c+dx)} + \frac{A a^4}{3d \tan^3(c+dx)} - \frac{A a^4}{5d \tan^5(c+dx)} - \frac{2A a^3 b \log(\tan^2(c+dx)+1)}{d} + \frac{4A a^3 b \log(\tan(c+dx))}{d} + \frac{2A a^3 b}{d \tan^2(c+dx)} \end{cases}$$

```
input integrate(cot(d*x+c)**6*(a+b*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)
```

```
output Piecewise((zoo*A*a**4*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**4*cot(c)**6, Eq(d, 0)), (zoo*A*a**4*x, Eq(c, -d*x)), (-A*a**4*x - A*a**4/(d*tan(c + d*x)) + A*a**4/(3*d*tan(c + d*x)**3) - A*a**4/(5*d*tan(c + d*x)**5) - 2*A*a**3*b*log(tan(c + d*x)**2 + 1)/d + 4*A*a**3*b*log(tan(c + d*x))/d + 2*A*a**3*b/(d*tan(c + d*x)**2) - A*a**3*b/(d*tan(c + d*x)**4) + 6*A*a**2*b**2*x + 6*A*a**2*b**2/(d*tan(c + d*x)) - 2*A*a**2*b**2/(d*tan(c + d*x)**3) + 2*A*a*b**3*log(tan(c + d*x)**2 + 1)/d - 4*A*a*b**3*log(tan(c + d*x))/d - 2*A*a*b**3/(d*tan(c + d*x)**2) - A*b**4*x - A*b**4/(d*tan(c + d*x)) - B*a**4*log(tan(c + d*x)**2 + 1)/(2*d) + B*a**4*log(tan(c + d*x))/d + B*a**4/(2*d*tan(c + d*x)**2) - B*a**4/(4*d*tan(c + d*x)**4) + 4*B*a**3*b*x + 4*B*a**3*b/(d*tan(c + d*x)) - 4*B*a**3*b/(3*d*tan(c + d*x)**3) + 3*B*a**2*b**2*log(tan(c + d*x)**2 + 1)/d - 6*B*a**2*b**2*log(tan(c + d*x))/d - 3*B*a**2*b**2/(d*tan(c + d*x)**2) - 4*B*a*b**3*x - 4*B*a*b**3/(d*tan(c + d*x)) - B*b**4*log(tan(c + d*x)**2 + 1)/(2*d) + B*b**4*log(tan(c + d*x))/d, True))
```

3.265.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.06

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$60(Aa^4 - 4Ba^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4)(dx + c) + 30(Ba^4 + 4Aa^3b - 6Ba^2b^2 - 4Aab^3 + Bb^4) \log(\tan(dx + c)^2 + 1) - 60(Ba^4 + 4Aa^3b - 6Ba^2b^2 - 4Aa^2b^2 - 4Aa^2b^2 - 4Aa^2b^2 + Bb^4) \log(\tan(dx + c)) + (12Aa^4 + 60(Aa^4 - 4Ba^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4) \tan(dx + c)^4 - 30(Ba^4 + 4Aa^3b - 6Ba^2b^2 - 4Aa^2b^2 - 4Aa^2b^2 + Bb^4) \tan(dx + c)^3 - 20(Aa^4 - 4Ba^3b - 6Aa^2b^2) \tan(dx + c)^2 + 15(Ba^4 + 4Aa^3b) \tan(dx + c)) / \tan(dx + c)^5 / d$$

input `integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/60*(60*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*(d*x + c) + 30*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*log(tan(d*x + c)^2 + 1) - 60*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*log(tan(d*x + c)) + (12*A*a^4 + 60*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*tan(d*x + c)^4 - 30*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3) *tan(d*x + c)^3 - 20*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2)*tan(d*x + c)^2 + 15*(B*a^4 + 4*A*a^3*b)*tan(d*x + c))/tan(d*x + c)^5/d`

3.265.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 763 vs. 2(265) = 530.

Time = 1.72 (sec) , antiderivative size = 763, normalized size of antiderivative = 2.79

$$\int \cot^6(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^6*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

```

1/960*(6*A*a^4*tan(1/2*d*x + 1/2*c)^5 - 15*B*a^4*tan(1/2*d*x + 1/2*c)^4 -
60*A*a^3*b*tan(1/2*d*x + 1/2*c)^4 - 70*A*a^4*tan(1/2*d*x + 1/2*c)^3 + 160*
B*a^3*b*tan(1/2*d*x + 1/2*c)^3 + 240*A*a^2*b^2*tan(1/2*d*x + 1/2*c)^3 + 18
0*B*a^4*tan(1/2*d*x + 1/2*c)^2 + 720*A*a^3*b*tan(1/2*d*x + 1/2*c)^2 - 720*
B*a^2*b^2*tan(1/2*d*x + 1/2*c)^2 - 480*A*a*b^3*tan(1/2*d*x + 1/2*c)^2 + 66
0*A*a^4*tan(1/2*d*x + 1/2*c) - 2400*B*a^3*b*tan(1/2*d*x + 1/2*c) - 3600*A*
a^2*b^2*tan(1/2*d*x + 1/2*c) + 1920*B*a*b^3*tan(1/2*d*x + 1/2*c) + 480*A*b
^4*tan(1/2*d*x + 1/2*c) - 960*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3
+ A*b^4)*(d*x + c) - 960*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B
*b^4)*log(tan(1/2*d*x + 1/2*c)^2 + 1) + 960*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b
^2 - 4*A*a*b^3 + B*b^4)*log(abs(tan(1/2*d*x + 1/2*c))) - (2192*B*a^4*tan(1
/2*d*x + 1/2*c)^5 + 8768*A*a^3*b*tan(1/2*d*x + 1/2*c)^5 - 13152*B*a^2*b^2*
tan(1/2*d*x + 1/2*c)^5 - 8768*A*a*b^3*tan(1/2*d*x + 1/2*c)^5 + 2192*B*b^4*
tan(1/2*d*x + 1/2*c)^5 + 660*A*a^4*tan(1/2*d*x + 1/2*c)^4 - 2400*B*a^3*b*t
an(1/2*d*x + 1/2*c)^4 - 3600*A*a^2*b^2*tan(1/2*d*x + 1/2*c)^4 + 1920*B*a*b
^3*tan(1/2*d*x + 1/2*c)^4 + 480*A*b^4*tan(1/2*d*x + 1/2*c)^4 - 180*B*a^4*t
an(1/2*d*x + 1/2*c)^3 - 720*A*a^3*b*tan(1/2*d*x + 1/2*c)^3 + 720*B*a^2*b^2
*tan(1/2*d*x + 1/2*c)^3 + 480*A*a*b^3*tan(1/2*d*x + 1/2*c)^3 - 70*A*a^4*ta
n(1/2*d*x + 1/2*c)^2 + 160*B*a^3*b*tan(1/2*d*x + 1/2*c)^2 + 240*A*a^2*b^2*
tan(1/2*d*x + 1/2*c)^2 + 15*B*a^4*tan(1/2*d*x + 1/2*c) + 60*A*a^3*b*tan...

```

3.265.9 Mupad [B] (verification not implemented)

Time = 8.62 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.96

$$\begin{aligned}
& \int \cot^6(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx \\
&= \frac{\ln(\tan(c + dx))(Ba^4 + 4Aa^3b - 6Ba^2b^2 - 4Aab^3 + Bb^4)}{\cot(c + dx)^5 \left(\tan(c + dx) \left(\frac{Ba^4}{4} + Aba^3 \right) + \frac{Aa^4}{5} - \tan(c + dx)^3 \left(\frac{Ba^4}{2} + 2Aa^3b - 3Ba^2b^2 - 2Aab^3 + Bb^4 \right) \right)} \\
&+ \frac{\ln(\tan(c + dx) - i)(-B + A li)(-b + a li)^4}{2d} \\
&- \frac{\ln(\tan(c + dx) + li)(B + A li)(b + a li)^4}{2d}
\end{aligned}$$

input `int(cot(c + d*x)^6*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^4,x)`

output $(\log(\tan(c + dx)) * (B*a^4 + B*b^4 - 6*B*a^2*b^2 - 4*A*a*b^3 + 4*A*a^3*b)) / d - (\cot(c + dx)^5 * (\tan(c + dx) * ((B*a^4)/4 + A*a^3*b) + (A*a^4)/5 - \tan(c + dx)^3 * ((B*a^4)/2 - 3*B*a^2*b^2 - 2*A*a*b^3 + 2*A*a^3*b) + \tan(c + dx)^2 * (2*A*a^2*b^2 - (A*a^4)/3 + (4*B*a^3*b)/3) + \tan(c + dx)^4 * (A*a^4 + A*b^4 - 6*A*a^2*b^2 + 4*B*a*b^3 - 4*B*a^3*b)) / d + (\log(\tan(c + dx) - 1i) * (A*1i - B) * (a*1i - b)^4) / (2*d) - (\log(\tan(c + dx) + 1i) * (A*1i + B) * (a*1i + b)^4) / (2*d)$

3.266 $\int \cot^7(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.266.1 Optimal result	2577
3.266.2 Mathematica [C] (verified)	2578
3.266.3 Rubi [A] (verified)	2578
3.266.4 Maple [A] (verified)	2584
3.266.5 Fricas [A] (verification not implemented)	2585
3.266.6 Sympy [A] (verification not implemented)	2586
3.266.7 Maxima [A] (verification not implemented)	2586
3.266.8 Giac [B] (verification not implemented)	2587
3.266.9 Mupad [B] (verification not implemented)	2588

3.266.1 Optimal result

Integrand size = 31, antiderivative size = 323

$$\int \cot^7(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$$

$$= -\left(\frac{(4a^3Ab - 4aAb^3 + a^4B - 6a^2b^2B + b^4B)x}{(4a^3Ab - 4aAb^3 + a^4B - 6a^2b^2B + b^4B) \cot(c+dx)} - \frac{d}{(a^4A - 6a^2Ab^2 + Ab^4 - 4a^3bB + 4ab^3B) \cot^2(c+dx)} + \frac{2d}{a(20a^2Ab - 13Ab^3 + 5a^3B - 27ab^2B) \cot^3(c+dx)} + \frac{15d}{a^2(5a^2A - 8Ab^2 - 12abB) \cot^4(c+dx)} - \frac{20d}{(a^4A - 6a^2Ab^2 + Ab^4 - 4a^3bB + 4ab^3B) \log(\sin(c+dx))} - \frac{d}{a(3Ab + 2aB) \cot^5(c+dx)(a+b \tan(c+dx))^2} - \frac{aA \cot^6(c+dx)(a+b \tan(c+dx))^3}{6d}\right)$$

output

```

-(4*A*a^3*b-4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*x-(4*A*a^3*b-4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*cot(d*x+c)/d-1/2*(A*a^4-6*A*a^2*b^2+A*b^4-4*B*a^3*b+4*B*a*b^3)*cot(d*x+c)^2/d+1/15*a*(20*A*a^2*b-13*A*b^3+5*B*a^3-27*B*a*b^2)*cot(d*x+c)^3/d+1/20*a^2*(5*A*a^2-8*A*b^2-12*B*a*b)*cot(d*x+c)^4/d-(A*a^4-6*A*a^2*b^2+A*b^4-4*B*a^3*b+4*B*a*b^3)*ln(sin(d*x+c))/d-1/10*a*(3*A*b+2*B*a)*cot(d*x+c)^5*(a+b*tan(d*x+c))^2/d-1/6*a*A*cot(d*x+c)^6*(a+b*tan(d*x+c))^3/d
    
```


3.266.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.38 (sec) , antiderivative size = 299, normalized size of antiderivative = 0.93

$$\int \cot^7(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$= \frac{-60(4a^3Ab - 4aAb^3 + a^4B - 6a^2b^2B + b^4B) \cot(c + dx) - 30(a^4A - 6a^2Ab^2 + Ab^4 - 4a^3bB + 4ab^3B) \cot^2(c + dx) + 15a^2(2a^2A - 6Aab^2 - 4a^3bB) \cot^3(c + dx) - 12a^3(4Aab + a^2B) \cot^4(c + dx) - 10a^4A \cot^5(c + dx) + 30(a + I*b)^4(A + I*B) \operatorname{Log}[I - \tan(c + dx)] - 60(a^4A - 6a^2Aab^2 + Ab^4 - 4a^3bB + 4a^2b^2B) \operatorname{Log}[\tan(c + dx)] + 30(a - I*b)^4(A - I*B) \operatorname{Log}[I + \tan(c + dx)]}{(60*d)}$$

input `Integrate[Cot[c + d*x]^7*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output $(-60*(4*a^3*A*b - 4*a*A*b^3 + a^4*B - 6*a^2*b^2*B + b^4*B)*\operatorname{Cot}[c + d*x] - 30*(a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*a^3*b*B + 4*a*b^3*B)*\operatorname{Cot}[c + d*x]^2 + 20*a*(4*a^2*A*b - 4*A*b^3 + a^3*B - 6*a*b^2*B)*\operatorname{Cot}[c + d*x]^3 + 15*a^2*(a^2*A - 6*A*b^2 - 4*a*b*B)*\operatorname{Cot}[c + d*x]^4 - 12*a^3*(4*A*b + a*B)*\operatorname{Cot}[c + d*x]^5 - 10*a^4*A*\operatorname{Cot}[c + d*x]^6 + 30*(a + I*b)^4*(A + I*B)*\operatorname{Log}[I - \tan[c + d*x]] - 60*(a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*a^3*b*B + 4*a*b^3*B)*\operatorname{Log}[\tan[c + d*x]] + 30*(a - I*b)^4*(A - I*B)*\operatorname{Log}[I + \tan[c + d*x]])/(60*d)$

3.266.3 Rubi [A] (verified)

Time = 2.29 (sec) , antiderivative size = 338, normalized size of antiderivative = 1.05, number of steps used = 21, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.677$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4118, 3042, 4111, 27, 3042, 4012, 3042, 4012, 25, 3042, 4014, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^7(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \tan(c + dx))^4(A + B \tan(c + dx))}{\tan(c + dx)^7} dx$$

$$\downarrow 4088$$

$$\frac{1}{6} \int 3 \cot^6(c+dx)(a+b \tan(c+dx))^2 (-b(aA-2bB) \tan^2(c+dx) - 2(Aa^2-2bBa-Ab^2) \tan(c+dx) + a(3Ab+2aB)) dx - \frac{aA \cot^6(c+dx)(a+b \tan(c+dx))^3}{6d}$$

↓ 27

$$\frac{1}{2} \int \cot^6(c+dx)(a+b \tan(c+dx))^2 (-b(aA-2bB) \tan^2(c+dx) - 2(Aa^2-2bBa-Ab^2) \tan(c+dx) + a(3Ab+2aB)) dx - \frac{aA \cot^6(c+dx)(a+b \tan(c+dx))^3}{6d}$$

↓ 3042

$$\frac{1}{2} \int \frac{(a+b \tan(c+dx))^2 (-b(aA-2bB) \tan(c+dx)^2 - 2(Aa^2-2bBa-Ab^2) \tan(c+dx) + a(3Ab+2aB))}{\tan(c+dx)^6} dx - \frac{aA \cot^6(c+dx)(a+b \tan(c+dx))^3}{6d}$$

↓ 4128

$$\frac{1}{2} \left(\frac{1}{5} \int -2 \cot^5(c+dx)(a+b \tan(c+dx)) (b(3Ba^2+7Aba-5b^2B) \tan^2(c+dx) + 5(Ba^3+3Aba^2-3b^2Ba-Ab^3)) dx - \frac{aA \cot^6(c+dx)(a+b \tan(c+dx))^3}{6d} \right)$$

↓ 27

$$\frac{1}{2} \left(-\frac{2}{5} \int \cot^5(c+dx)(a+b \tan(c+dx)) (b(3Ba^2+7Aba-5b^2B) \tan^2(c+dx) + 5(Ba^3+3Aba^2-3b^2Ba-Ab^3)) dx - \frac{aA \cot^6(c+dx)(a+b \tan(c+dx))^3}{6d} \right)$$

↓ 3042

$$\frac{1}{2} \left(-\frac{2}{5} \int \frac{(a+b \tan(c+dx)) (b(3Ba^2+7Aba-5b^2B) \tan(c+dx)^2 + 5(Ba^3+3Aba^2-3b^2Ba-Ab^3)) \tan(c+dx)}{\tan(c+dx)^5} dx - \frac{aA \cot^6(c+dx)(a+b \tan(c+dx))^3}{6d} \right)$$

↓ 4118

$$\frac{1}{2} \left(-\frac{2}{5} \left(\int \cot^4(c+dx) (b^2(3Ba^2 + 7Aba - 5b^2B) \tan^2(c+dx) - 5(Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4) \tan(c+dx) + a \frac{aA \cot^6(c+dx)(a+b \tan(c+dx))^3}{6d} \right) \right)$$

↓ 3042

$$\frac{1}{2} \left(-\frac{2}{5} \left(\int \frac{b^2(3Ba^2 + 7Aba - 5b^2B) \tan(c+dx)^2 - 5(Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4) \tan(c+dx) + a \frac{aA \cot^6(c+dx)(a+b \tan(c+dx))^3}{6d}}{\tan(c+dx)^4} \right) \right)$$

↓ 4111

$$\frac{1}{2} \left(-\frac{2}{5} \left(\int -5 \cot^3(c+dx) (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4 + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \tan(c+dx) + a \frac{aA \cot^6(c+dx)(a+b \tan(c+dx))^3}{6d} \right) \right)$$

↓ 27

$$\frac{1}{2} \left(-\frac{2}{5} \left(-5 \int \cot^3(c+dx) (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4 + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \tan(c+dx) + a \frac{aA \cot^6(c+dx)(a+b \tan(c+dx))^3}{6d} \right) \right)$$

↓ 3042

$$\frac{1}{2} \left(-\frac{2}{5} \left(-5 \int \frac{Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4 + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \tan(c+dx)}{\tan(c+dx)^3} dx + a \frac{aA \cot^6(c+dx)(a+b \tan(c+dx))^3}{6d} \right) \right)$$

↓ 4012

$$\frac{1}{2} \left(-\frac{2}{5} \left(-5 \left(\int \cot^2(c+dx) (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B - (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4) \tan(c+dx) + a \frac{aA \cot^6(c+dx)(a+b \tan(c+dx))^3}{6d} \right) \right) \right)$$

↓ 3042

$$\frac{1}{2} \left(-\frac{2}{5} \left(-5 \left(\int \frac{Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B - (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4) \tan(c + dx)}{\tan(c + dx)^2} \right) \right) \right) \frac{aA \cot^6(c + dx)(a + b \tan(c + dx))^3}{6d}$$

↓ 4012

$$\frac{1}{2} \left(-\frac{2}{5} \left(-5 \left(\int -\cot(c + dx) (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4 + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \right) \right) \right) \frac{aA \cot^6(c + dx)(a + b \tan(c + dx))^3}{6d}$$

↓ 25

$$\frac{1}{2} \left(-\frac{2}{5} \left(-5 \left(- \int \cot(c + dx) (Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4 + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \right) \right) \right) \frac{aA \cot^6(c + dx)(a + b \tan(c + dx))^3}{6d}$$

↓ 3042

$$\frac{1}{2} \left(-\frac{2}{5} \left(-5 \left(- \int \frac{Aa^4 - 4bBa^3 - 6Ab^2a^2 + 4b^3Ba + Ab^4 + (Ba^4 + 4Aba^3 - 6b^2Ba^2 - 4Ab^3a + b^4B) \tan(c + dx)}{\tan(c + dx)} \right) \right) \right) \frac{aA \cot^6(c + dx)(a + b \tan(c + dx))^3}{6d}$$

↓ 4014

$$\frac{1}{2} \left(-\frac{2}{5} \left(-5 \left(-(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) \int \cot(c + dx) dx - \frac{(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4)}{2d} \right) \right) \right) \frac{aA \cot^6(c + dx)(a + b \tan(c + dx))^3}{6d}$$

↓ 3042

$$\frac{1}{2} \left(-\frac{2}{5} \left(-5 \left(-(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4) \int -\tan \left(c + dx + \frac{\pi}{2} \right) dx - \frac{(a^4A - 4a^3bB - 6a^2Ab^2 + 4ab^3B + Ab^4)}{2d} \right) \right) \right) \frac{aA \cot^6(c + dx)(a + b \tan(c + dx))^3}{6d}$$

↓ 25

$$\frac{1}{2} \left(-\frac{2}{5} \left(-5 \left((a^4 A - 4a^3 b B - 6a^2 A b^2 + 4ab^3 B + Ab^4) \int \tan \left(\frac{1}{2}(2c + \pi) + dx \right) dx - \frac{(a^4 A - 4a^3 b B - 6a^2 A b^2 + 4ab^3 B + Ab^4)}{6d} \right) \right) \right)$$

$$\frac{aA \cot^6(c + dx)(a + b \tan(c + dx))^3}{6d}$$

↓ 3956

$$\frac{1}{2} \left(-\frac{2}{5} \left(-\frac{a^2(5a^2 A - 12ab B - 8Ab^2) \cot^4(c + dx)}{4d} - \frac{a(5a^3 B + 20a^2 A b - 27ab^2 B - 13Ab^3) \cot^3(c + dx)}{3d} - 5 \left(-\frac{aA \cot^6(c + dx)(a + b \tan(c + dx))^3}{6d} \right) \right) \right)$$

```
input Int[Cot[c + d*x]^7*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]
```

```
output -1/6*(a*A*Cot[c + d*x]^6*(a + b*Tan[c + d*x])^3)/d + ((-2*(-1/3*(a*(20*a^2
*A*b - 13*A*b^3 + 5*a^3*B - 27*a*b^2*B)*Cot[c + d*x]^3)/d - (a^2*(5*a^2*A
- 8*A*b^2 - 12*a*b*B)*Cot[c + d*x]^4)/(4*d) - 5*(-((4*a^3*A*b - 4*a*A*b^3
+ a^4*B - 6*a^2*b^2*B + b^4*B)*x) - ((4*a^3*A*b - 4*a*A*b^3 + a^4*B - 6*a^
2*b^2*B + b^4*B)*Cot[c + d*x])/d - ((a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*a^3*b
*B + 4*a*b^3*B)*Cot[c + d*x]^2)/(2*d) - ((a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*
a^3*b*B + 4*a*b^3*B)*Log[-Sin[c + d*x]]/d)))/5 - (a*(3*A*b + 2*a*B)*Cot[c
+ d*x]^5*(a + b*Tan[c + d*x])^2)/(5*d))/2
```

3.266.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4111 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

```
rule 4118 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
)*(x_)]^(n)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]
```

```
rule 4128 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]^(n)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

3.266.4 Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 330, normalized size of antiderivative = 1.02

method	result
parallelrisch	$(30A^4 - 180A^2b^2 + 30Ab^4 - 120Ba^3b + 120Bab^3) \ln(\sec^2(dx+c)) + (-60A^4 + 360A^2b^2 - 60Ab^4 + 240Ba^3b - 240Bab^3) \arctan(\tan(dx+c)) -$
derivativedivides	$\frac{(A^4 - 6A^2b^2 + Ab^4 - 4Ba^3b + 4Bab^3) \ln(1 + \tan^2(dx+c))}{2} + (-4A^3b + 4Aab^3 - Ba^4 + 6Ba^2b^2 - Bb^4) \arctan(\tan(dx+c)) -$
default	$\frac{(A^4 - 6A^2b^2 + Ab^4 - 4Ba^3b + 4Bab^3) \ln(1 + \tan^2(dx+c))}{2} + (-4A^3b + 4Aab^3 - Ba^4 + 6Ba^2b^2 - Bb^4) \arctan(\tan(dx+c)) -$
norman	$(-4A^3b + 4Aab^3 - Ba^4 + 6Ba^2b^2 - Bb^4)x(\tan^6(dx+c)) - \frac{Aa^4}{6d} - \frac{(4A^3b - 4Aab^3 + Ba^4 - 6Ba^2b^2 + Bb^4)(\tan^5(dx+c))}{d} -$
risch	Expression too large to display

```
input int(cot(d*x+c)^7*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x,method=_RETURNVERBO
SE)
```

3.266. $\int \cot^7(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

```
output 1/60*((30*A*a^4-180*A*a^2*b^2+30*A*b^4-120*B*a^3*b+120*B*a*b^3)*ln(sec(d*x
+c)^2)+(-60*A*a^4+360*A*a^2*b^2-60*A*b^4+240*B*a^3*b-240*B*a*b^3)*ln(tan(d
*x+c))-10*A*cot(d*x+c)^6*a^4+(-48*A*a^3*b-12*B*a^4)*cot(d*x+c)^5+15*a^2*co
t(d*x+c)^4*(A*a^2-6*A*b^2-4*B*a*b)+(80*A*a^3*b-80*A*a*b^3+20*B*a^4-120*B*a
^2*b^2)*cot(d*x+c)^3+(-30*A*a^4+180*A*a^2*b^2-30*A*b^4+120*B*a^3*b-120*B*a
*b^3)*cot(d*x+c)^2+(-240*A*a^3*b+240*A*a*b^3-60*B*a^4+360*B*a^2*b^2-60*B*b
^4)*cot(d*x+c)-240*d*(A*a^3*b-A*a*b^3+1/4*B*a^4-3/2*B*a^2*b^2+1/4*B*b^4)*x
)/d
```

3.266.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 350, normalized size of antiderivative = 1.08

$$\int \cot^7(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$\frac{30(Aa^4 - 4Ba^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^6 + 5(11Aa^4 - 36Ba^3b - 54Aa^2b^2 + 24Bab^3 + 6Aa^2b^2 - 4Aa^2b^2 + 4Aa^2b^2 + 12(Ba^4 + 4Aa^3b - 6Ba^2b^2 - 4Aa^2b^3 + Bb^4)*dx)*\tan(dx+c)^6 + 60*(Ba^4 + 4Aa^3b - 6Ba^2b^2 - 4Aa^2b^3 + Bb^4)*\tan(dx+c)^5 + 10*Aa^4 + 30*(Aa^4 - 4Ba^3b - 6Aa^2b^2 + 4Aa^2b^2 + 4Aa^2b^2 + 12(Ba^4 + 4Aa^3b - 6Ba^2b^2 - 4Aa^2b^3 + Bb^4)*\tan(dx+c)^3 - 15*(Aa^4 - 4Ba^3b - 6Aa^2b^2)*\tan(dx+c)^2 + 12*(Ba^4 + 4Aa^3b)*\tan(dx+c))/(d*\tan(dx+c)^6)}$$

```
input integrate(cot(d*x+c)^7*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="f
ricas")
```

```
output -1/60*(30*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*log(tan(d*
x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^6 + 5*(11*A*a^4 - 36*B*a^3*b -
54*A*a^2*b^2 + 24*B*a*b^3 + 6*A*b^4 + 12*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2
- 4*A*a^2*b^3 + B*b^4)*d*x)*tan(d*x + c)^6 + 60*(B*a^4 + 4*A*a^3*b - 6*B*a^
2*b^2 - 4*A*a^2*b^3 + B*b^4)*tan(d*x + c)^5 + 10*A*a^4 + 30*(A*a^4 - 4*B*a^3
*b - 6*A*a^2*b^2 + 4*B*a^2*b^2 + 4*A*a^2*b^2 + 12*(B*a^4 + 4*A*a^3*b - 6*B
*a^2*b^2 - 4*A*a^2*b^3 + B*b^4)*tan(d*x + c)^3 - 15*(A*a^4 - 4*B*a^3*b - 6*A
*a^2*b^2)*tan(d*x + c)^2 + 12*(B*a^4 + 4*A*a^3*b)*tan(d*x + c))/(d*tan(d*x
+ c)^6)
```


3.266.6 Sympy [A] (verification not implemented)

Time = 12.58 (sec) , antiderivative size = 643, normalized size of antiderivative = 1.99

$$\int \cot^7(c+dx)(a+b\tan(c+dx))^4(A+B\tan(c+dx))dx$$

$$= \begin{cases} \tilde{\infty}Aa^4x \\ x(A+B\tan(c))(a+b\tan(c))^4\cot^7(c) \\ \tilde{\infty}Aa^4x \\ \frac{Aa^4\log(\tan^2(c+dx)+1)}{2d} - \frac{Aa^4\log(\tan(c+dx))}{d} - \frac{Aa^4}{2d\tan^2(c+dx)} + \frac{Aa^4}{4d\tan^4(c+dx)} - \frac{Aa^4}{6d\tan^6(c+dx)} - 4Aa^3bx - \frac{4Aa^3b}{d\tan(c+dx)} \end{cases}$$

input `integrate(cot(d*x+c)**7*(a+b*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)`

output `Piecewise((zoo*A*a**4*x, Eq(c, 0) & Eq(d, 0)), (x*(A + B*tan(c))*(a + b*tan(c))**4*cot(c)**7, Eq(d, 0)), (zoo*A*a**4*x, Eq(c, -d*x)), (A*a**4*log(tan(c + d*x)**2 + 1)/(2*d) - A*a**4*log(tan(c + d*x))/d - A*a**4/(2*d*tan(c + d*x)**2) + A*a**4/(4*d*tan(c + d*x)**4) - A*a**4/(6*d*tan(c + d*x)**6) - 4*A*a**3*b*x - 4*A*a**3*b/(d*tan(c + d*x)) + 4*A*a**3*b/(3*d*tan(c + d*x)**3) - 4*A*a**3*b/(5*d*tan(c + d*x)**5) - 3*A*a**2*b**2*log(tan(c + d*x)**2 + 1)/d + 6*A*a**2*b**2*log(tan(c + d*x))/d + 3*A*a**2*b**2/(d*tan(c + d*x)**2) - 3*A*a**2*b**2/(2*d*tan(c + d*x)**4) + 4*A*a*b**3*x + 4*A*a*b**3/(d*tan(c + d*x)) - 4*A*a*b**3/(3*d*tan(c + d*x)**3) + A*b**4*log(tan(c + d*x)**2 + 1)/(2*d) - A*b**4*log(tan(c + d*x))/d - A*b**4/(2*d*tan(c + d*x)**2) - B*a**4*x - B*a**4/(d*tan(c + d*x)) + B*a**4/(3*d*tan(c + d*x)**3) - B*a**4/(5*d*tan(c + d*x)**5) - 2*B*a**3*b*log(tan(c + d*x)**2 + 1)/d + 4*B*a**3*b*log(tan(c + d*x))/d + 2*B*a**3*b/(d*tan(c + d*x)**2) - B*a**3*b/(d*tan(c + d*x)**4) + 6*B*a**2*b**2*x + 6*B*a**2*b**2/(d*tan(c + d*x)) - 2*B*a**2*b**2/(d*tan(c + d*x)**3) + 2*B*a*b**3*log(tan(c + d*x)**2 + 1)/d - 4*B*a*b**3*log(tan(c + d*x))/d - 2*B*a*b**3/(d*tan(c + d*x)**2) - B*b**4*x - B*b**4/(d*tan(c + d*x)), True))`

3.266.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 333, normalized size of antiderivative = 1.03

$$\int \cot^7(c+dx)(a+b\tan(c+dx))^4(A+B\tan(c+dx))dx =$$

$$\frac{60(Ba^4 + 4Aa^3b - 6Ba^2b^2 - 4Aab^3 + Bb^4)(dx+c) - 30(Aa^4 - 4Ba^3b - 6Aa^2b^2 + 4Bab^3 + Ab^4)}{1}$$

input `integrate(cot(d*x+c)^7*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/60*(60*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*(d*x + c) - 30*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)^2 + 1) + 60*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)) + (60*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*tan(d*x + c)^5 + 10*A*a^4 + 30*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*tan(d*x + c)^4 - 20*(B*a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3)*tan(d*x + c)^3 - 15*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2)*tan(d*x + c)^2 + 12*(B*a^4 + 4*A*a^3*b)*tan(d*x + c))/tan(d*x + c)^6)/d`

3.266.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 943 vs. $2(313) = 626$.

Time = 1.85 (sec) , antiderivative size = 943, normalized size of antiderivative = 2.92

$$\int \cot^7(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^7*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output

```

-1/1920*(5*A*a^4*tan(1/2*d*x + 1/2*c)^6 - 12*B*a^4*tan(1/2*d*x + 1/2*c)^5
- 48*A*a^3*b*tan(1/2*d*x + 1/2*c)^5 - 60*A*a^4*tan(1/2*d*x + 1/2*c)^4 + 12
0*B*a^3*b*tan(1/2*d*x + 1/2*c)^4 + 180*A*a^2*b^2*tan(1/2*d*x + 1/2*c)^4 +
140*B*a^4*tan(1/2*d*x + 1/2*c)^3 + 560*A*a^3*b*tan(1/2*d*x + 1/2*c)^3 - 48
0*B*a^2*b^2*tan(1/2*d*x + 1/2*c)^3 - 320*A*a*b^3*tan(1/2*d*x + 1/2*c)^3 +
435*A*a^4*tan(1/2*d*x + 1/2*c)^2 - 1440*B*a^3*b*tan(1/2*d*x + 1/2*c)^2 - 2
160*A*a^2*b^2*tan(1/2*d*x + 1/2*c)^2 + 960*B*a*b^3*tan(1/2*d*x + 1/2*c)^2
+ 240*A*b^4*tan(1/2*d*x + 1/2*c)^2 - 1320*B*a^4*tan(1/2*d*x + 1/2*c) - 528
0*A*a^3*b*tan(1/2*d*x + 1/2*c) + 7200*B*a^2*b^2*tan(1/2*d*x + 1/2*c) + 480
0*A*a*b^3*tan(1/2*d*x + 1/2*c) - 960*B*b^4*tan(1/2*d*x + 1/2*c) + 1920*(B*
a^4 + 4*A*a^3*b - 6*B*a^2*b^2 - 4*A*a*b^3 + B*b^4)*(d*x + c) - 1920*(A*a^4
- 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*log(tan(1/2*d*x + 1/2*c)^2
+ 1) + 1920*(A*a^4 - 4*B*a^3*b - 6*A*a^2*b^2 + 4*B*a*b^3 + A*b^4)*log(abs
(tan(1/2*d*x + 1/2*c))) - (4704*A*a^4*tan(1/2*d*x + 1/2*c)^6 - 18816*B*a^3
*b*tan(1/2*d*x + 1/2*c)^6 - 28224*A*a^2*b^2*tan(1/2*d*x + 1/2*c)^6 + 18816
*B*a*b^3*tan(1/2*d*x + 1/2*c)^6 + 4704*A*b^4*tan(1/2*d*x + 1/2*c)^6 - 1320
*B*a^4*tan(1/2*d*x + 1/2*c)^5 - 5280*A*a^3*b*tan(1/2*d*x + 1/2*c)^5 + 7200
*B*a^2*b^2*tan(1/2*d*x + 1/2*c)^5 + 4800*A*a*b^3*tan(1/2*d*x + 1/2*c)^5 -
960*B*b^4*tan(1/2*d*x + 1/2*c)^5 - 435*A*a^4*tan(1/2*d*x + 1/2*c)^4 + 1440
*B*a^3*b*tan(1/2*d*x + 1/2*c)^4 + 2160*A*a^2*b^2*tan(1/2*d*x + 1/2*c)^4...

```

3.266.9 Mupad [B] (verification not implemented)

Time = 8.11 (sec) , antiderivative size = 307, normalized size of antiderivative = 0.95

$$\int \cot^7(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx =$$

$$\frac{\cot(c + dx)^6 \left(\tan(c + dx) \left(\frac{B a^4}{5} + \frac{4 A b a^3}{5} \right) + \frac{A a^4}{6} - \tan(c + dx)^3 \left(\frac{B a^4}{3} + \frac{4 A a^3 b}{3} - 2 B a^2 b^2 - \frac{4 A a b^3}{3} \right) \right)}{\ln(\tan(c + dx)) (A a^4 - 4 B a^3 b - 6 A a^2 b^2 + 4 B a b^3 + A b^4)}$$

$$+ \frac{\ln(\tan(c + dx) + i) (A - B i) (b + a i)^4}{2 d}$$

$$+ \frac{\ln(\tan(c + dx) - i) (A + B i) (-b + a i)^4}{2 d}$$

input `int(cot(c + d*x)^7*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^4,x)`

output $(\log(\tan(c + dx) + 1i) \cdot (A - B \cdot 1i) \cdot (a \cdot 1i + b)^4) / (2 \cdot d) - (\log(\tan(c + dx)) \cdot (A \cdot a^4 + A \cdot b^4 - 6 \cdot A \cdot a^2 \cdot b^2 + 4 \cdot B \cdot a \cdot b^3 - 4 \cdot B \cdot a^3 \cdot b)) / d - (\cot(c + dx) \cdot \tan^6(c + dx) \cdot ((B \cdot a^4) / 5 + (4 \cdot A \cdot a^3 \cdot b) / 5) + (A \cdot a^4) / 6 - \tan^3(c + dx) \cdot ((B \cdot a^4) / 3 - 2 \cdot B \cdot a^2 \cdot b^2 - (4 \cdot A \cdot a \cdot b^3) / 3 + (4 \cdot A \cdot a^3 \cdot b) / 3) + \tan^2(c + dx) \cdot ((3 \cdot A \cdot a^2 \cdot b^2) / 2 - (A \cdot a^4) / 4 + B \cdot a^3 \cdot b) + \tan^4(c + dx) \cdot ((A \cdot a^4) / 2 + (A \cdot b^4) / 2 - 3 \cdot A \cdot a^2 \cdot b^2 + 2 \cdot B \cdot a \cdot b^3 - 2 \cdot B \cdot a^3 \cdot b) + \tan^5(c + dx) \cdot (B \cdot a^4 + B \cdot b^4 - 6 \cdot B \cdot a^2 \cdot b^2 - 4 \cdot A \cdot a \cdot b^3 + 4 \cdot A \cdot a^3 \cdot b)) / d + (\log(\tan(c + dx) - 1i) \cdot (A + B \cdot 1i) \cdot (a \cdot 1i - b)^4) / (2 \cdot d)$

3.267
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.267.1 Optimal result 2590
 3.267.2 Mathematica [C] (verified) 2590
 3.267.3 Rubi [A] (verified) 2591
 3.267.4 Maple [A] (verified) 2594
 3.267.5 Fricas [A] (verification not implemented) 2595
 3.267.6 Sympy [C] (verification not implemented) 2595
 3.267.7 Maxima [A] (verification not implemented) 2596
 3.267.8 Giac [A] (verification not implemented) 2597
 3.267.9 Mupad [B] (verification not implemented) 2597

3.267.1 Optimal result

Integrand size = 31, antiderivative size = 127

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = -\frac{(Ab-aB)x}{a^2+b^2} + \frac{(aA+bB) \log(\cos(c+dx))}{(a^2+b^2)d} - \frac{a^3(Ab-aB) \log(a+b \tan(c+dx))}{b^3(a^2+b^2)d} + \frac{(Ab-aB) \tan(c+dx)}{b^2d} + \frac{B \tan^2(c+dx)}{2bd}$$

output

```
-(A*b-B*a)*x/(a^2+b^2)+(A*a+B*b)*ln(cos(d*x+c))/(a^2+b^2)/d-a^3*(A*b-B*a)*ln(a+b*tan(d*x+c))/b^3/(a^2+b^2)/d+(A*b-B*a)*tan(d*x+c)/b^2/d+1/2*B*tan(d*x+c)^2/b/d
```

3.267.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.56 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.09

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{-\frac{b(A+iB) \log(i-\tan(c+dx))}{a+ib} - \frac{b(A-iB) \log(i+\tan(c+dx))}{a-ib} + \frac{2a^3(-Ab+aB) \log(a+b \tan(c+dx))}{b^2(a^2+b^2)} + \frac{2(Ab-aB) \tan(c+dx)}{b} + B \tan^2(c+dx)}{2bd}$$

input `Integrate[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(-((b*(A + I*B)*Log[I - Tan[c + d*x]])/(a + I*b)) - (b*(A - I*B)*Log[I + Tan[c + d*x]])/(a - I*b) + (2*a^3*(-(A*b) + a*B)*Log[a + b*Tan[c + d*x]])/(b^2*(a^2 + b^2)) + (2*(A*b - a*B)*Tan[c + d*x])/b + B*Tan[c + d*x]^2)/(2*b*d)`

3.267.3 Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.13, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4090, 27, 3042, 4130, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^3(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx \\
 & \quad \downarrow \text{4090} \\
 & \frac{\int -\frac{2\tan(c+dx)((Ab-aB)\tan^2(c+dx)+bB\tan(c+dx)+aB)}{a+b\tan(c+dx)} dx}{2b} + \frac{B\tan^2(c+dx)}{2bd} \\
 & \quad \downarrow \text{27} \\
 & \frac{B\tan^2(c+dx)}{2bd} - \frac{\int \frac{\tan(c+dx)((Ab-aB)\tan^2(c+dx)+bB\tan(c+dx)+aB)}{a+b\tan(c+dx)} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{B\tan^2(c+dx)}{2bd} - \frac{\int \frac{\tan(c+dx)((Ab-aB)\tan(c+dx)^2+bB\tan(c+dx)+aB)}{a+b\tan(c+dx)} dx}{b} \\
 & \quad \downarrow \text{4130} \\
 & \frac{B\tan^2(c+dx)}{2bd} - \frac{\int \frac{A\tan(c+dx)b^2+(-Ba^2+Ab+a+b^2B)\tan^2(c+dx)+a(Ab-aB)}{a+b\tan(c+dx)} dx}{b} - \frac{(Ab-aB)\tan(c+dx)}{bd} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.267. $\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$

$$\begin{aligned}
 & \frac{B \tan^2(c + dx)}{2bd} - \frac{\int \frac{A \tan(c+dx)b^2 + (-Ba^2 + Aba + b^2B) \tan(c+dx)^2 + a(Ab - aB)}{a+b \tan(c+dx)} dx}{b} - \frac{(Ab - aB) \tan(c+dx)}{bd} \\
 & \quad \downarrow \text{4109} \\
 & \frac{B \tan^2(c + dx)}{2bd} - \frac{\frac{b^2(aA+bB) \int \tan(c+dx) dx}{a^2+b^2} + \frac{a^3(Ab - aB) \int \frac{\tan^2(c+dx)+1}{a+b \tan(c+dx)} dx}{b} + \frac{b^2x(Ab - aB)}{a^2+b^2}}{b} - \frac{(Ab - aB) \tan(c+dx)}{bd} \\
 & \quad \downarrow \text{3042} \\
 & \frac{B \tan^2(c + dx)}{2bd} - \frac{\frac{b^2(aA+bB) \int \tan(c+dx) dx}{a^2+b^2} + \frac{a^3(Ab - aB) \int \frac{\tan(c+dx)^2+1}{a+b \tan(c+dx)} dx}{b} + \frac{b^2x(Ab - aB)}{a^2+b^2}}{b} - \frac{(Ab - aB) \tan(c+dx)}{bd} \\
 & \quad \downarrow \text{3956} \\
 & \frac{B \tan^2(c + dx)}{2bd} - \frac{\frac{a^3(Ab - aB) \int \frac{\tan(c+dx)^2+1}{a+b \tan(c+dx)} dx}{a^2+b^2} - \frac{b^2(aA+bB) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{b^2x(Ab - aB)}{a^2+b^2}}{b} - \frac{(Ab - aB) \tan(c+dx)}{bd} \\
 & \quad \downarrow \text{4100} \\
 & \frac{B \tan^2(c + dx)}{2bd} - \frac{\frac{a^3(Ab - aB) \int \frac{1}{a+b \tan(c+dx)} d(b \tan(c+dx))}{bd(a^2+b^2)} - \frac{b^2(aA+bB) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{b^2x(Ab - aB)}{a^2+b^2}}{b} - \frac{(Ab - aB) \tan(c+dx)}{bd} \\
 & \quad \downarrow \text{16} \\
 & \frac{B \tan^2(c + dx)}{2bd} - \frac{-\frac{b^2(aA+bB) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{b^2x(Ab - aB)}{a^2+b^2} + \frac{a^3(Ab - aB) \log(a+b \tan(c+dx))}{bd(a^2+b^2)}}{b} - \frac{(Ab - aB) \tan(c+dx)}{bd}
 \end{aligned}$$

input `Int[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(B*Tan[c + d*x]^2)/(2*b*d) - (((b^2*(A*b - a*B)*x)/(a^2 + b^2) - (b^2*(a*A + b*B)*Log[Cos[c + d*x]]))/(a^2 + b^2)*d) + (a^3*(A*b - a*B)*Log[a + b*Tan[c + d*x]])/(b*(a^2 + b^2)*d)/b - ((A*b - a*B)*Tan[c + d*x])/(b*d)/b`

3.267.3.1 Defintions of rubi rules used

- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`
- rule 4100 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]`
- rule 4109 `Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C, 0]`


```
rule 4130 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && (!IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

3.267.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{\frac{bB(\tan^2(dx+c))}{2} + A \tan(dx+c)b - B \tan(dx+c)a}{b^2} + \frac{(-aA - Bb) \ln(1 + \tan^2(dx+c))}{2} + \frac{(-Ab + Ba) \arctan(\tan(dx+c))}{a^2 + b^2} - \frac{a^3(Ab - Ba) \ln(a + b \tan(dx+c))}{b^3(a^2 + b^2)}$
default	$\frac{\frac{bB(\tan^2(dx+c))}{2} + A \tan(dx+c)b - B \tan(dx+c)a}{b^2} + \frac{(-aA - Bb) \ln(1 + \tan^2(dx+c))}{2} + \frac{(-Ab + Ba) \arctan(\tan(dx+c))}{a^2 + b^2} - \frac{a^3(Ab - Ba) \ln(a + b \tan(dx+c))}{b^3(a^2 + b^2)}$
norman	$\frac{(Ab - Ba) \tan(dx+c)}{b^2 d} - \frac{(Ab - Ba)x}{a^2 + b^2} + \frac{B(\tan^2(dx+c))}{2bd} - \frac{(aA + Bb) \ln(1 + \tan^2(dx+c))}{2d(a^2 + b^2)} - \frac{a^3(Ab - Ba) \ln(a + b \tan(dx+c))}{b^3(a^2 + b^2)d}$
parallelrisch	$- \frac{2A b^4 dx - 2Ba b^3 dx - B a^2 b^2 (\tan^2(dx+c)) - B b^4 (\tan^2(dx+c)) + A \ln(1 + \tan^2(dx+c)) a b^3 + 2A \ln(a + b \tan(dx+c)) a^3 b}{d}$
risch	$- \frac{x B}{ib - a} + \frac{2i B a^2 c}{b^3 d} + \frac{2i a^3 A x}{b^2(a^2 + b^2)} + \frac{2i(-i B b e^{2i(dx+c)} + A b e^{2i(dx+c)} - B a e^{2i(dx+c)} + Ab - Ba)}{d b^2 (e^{2i(dx+c)} + 1)^2} - \frac{2i B x}{b} - \frac{2i a^4}{b^3(a^2 + b^2)}$

```
input int(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE
)
```

```
output 1/d*(1/b^2*(1/2*B*tan(d*x+c)^2*b+A*tan(d*x+c)*b-B*tan(d*x+c)*a)+1/(a^2+b^2
)*(1/2*(-A*a-B*b)*ln(1+tan(d*x+c)^2)+(-A*b+B*a)*arctan(tan(d*x+c)))-1/b^3*
a^3*(A*b-B*a)/(a^2+b^2)*ln(a+b*tan(d*x+c)))
```

3.267.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.50

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{2(Bab^3 - Ab^4)dx + (Ba^2b^2 + Bb^4)\tan(dx+c)^2 + (Ba^4 - Aa^3b)\log\left(\frac{b^2\tan(dx+c)^2 + 2ab\tan(dx+c) + a^2}{\tan(dx+c)^2 + 1}\right) - (Ba^4 - Aa^3b)}{2(a^2b^3 + b^5)d}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output `1/2*(2*(B*a*b^3 - A*b^4)*d*x + (B*a^2*b^2 + B*b^4)*tan(d*x + c)^2 + (B*a^4 - A*a^3*b)*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - (B*a^4 - A*a^3*b - A*a*b^3 - B*b^4)*log(1/(tan(d*x + c)^2 + 1)) - 2*(B*a^3*b - A*a^2*b^2 + B*a*b^3 - A*b^4)*tan(d*x + c))/((a^2*b^3 + b^5)*d)`

3.267.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.95 (sec) , antiderivative size = 1297, normalized size of antiderivative = 10.21

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)**3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Piecewise((zoo*x*(A + B*tan(c))*tan(c)**2, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), ((-A*log(tan(c + d*x)**2 + 1)/(2*d) + A*tan(c + d*x)**2/(2*d) + B*x + B*tan(c + d*x)**3/(3*d) - B*tan(c + d*x)/d)/a, Eq(b, 0)), (-3*A*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 3*I*A*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) + I*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + A*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 2*A*tan(c + d*x)**2/(2*b*d*tan(c + d*x) - 2*I*b*d) + 3*A/(2*b*d*tan(c + d*x) - 2*I*b*d) - 3*I*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - 3*B*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) - 2*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 2*I*B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) - 2*I*b*d) + B*tan(c + d*x)**3/(2*b*d*tan(c + d*x) - 2*I*b*d) + I*B*tan(c + d*x)**2/(2*b*d*tan(c + d*x) - 2*I*b*d) + 3*I*B/(2*b*d*tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (-3*A*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) - 3*I*A*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - I*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + A*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2*I*b*d) + 2*A*tan(c + d*x)**2/(2*b*d*tan(c + d*x) + 2*I*b*d) + 3*A/(2*b*d*tan(c + d*x) + 2*I*b*d) + 3*I*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) - 3*B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - 2*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) - 2*I*B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2*I*b*d)...`

3.267.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.02

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$= \frac{2(Ba-Ab)(dx+c)}{a^2+b^2} + \frac{2(Ba^4-Aa^3b) \log(b \tan(dx+c)+a)}{a^2b^3+b^5} - \frac{(Aa+Bb) \log(\tan(dx+c)^2+1)}{a^2+b^2} + \frac{Bb \tan(dx+c)^2 - 2(Ba-Ab) \tan(dx+c)}{b^2}$$

$2d$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(2*(B*a - A*b)*(d*x + c)/(a^2 + b^2) + 2*(B*a^4 - A*a^3*b)*log(b*tan(d*x + c) + a)/(a^2*b^3 + b^5) - (A*a + B*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) + (B*b*tan(d*x + c)^2 - 2*(B*a - A*b)*tan(d*x + c))/b^2)/d`

3.267.8 Giac [A] (verification not implemented)

Time = 0.65 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.06

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{2(Ba-Ab)(dx+c)}{a^2+b^2} - \frac{(Aa+Bb)\log(\tan(dx+c)^2+1)}{a^2+b^2} + \frac{2(Ba^4-Aa^3b)\log(|b\tan(dx+c)+a|)}{a^2b^3+b^5} + \frac{Bb\tan(dx+c)^2-2Ba\tan(dx+c)+2Ab\tan(dx+c)}{b^2}$$

$$2d$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `1/2*(2*(B*a - A*b)*(d*x + c)/(a^2 + b^2) - (A*a + B*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) + 2*(B*a^4 - A*a^3*b)*log(abs(b*tan(d*x + c) + a))/(a^2*b^3 + b^5) + (B*b*tan(d*x + c)^2 - 2*B*a*tan(d*x + c) + 2*A*b*tan(d*x + c))/b^2)/d`

3.267.9 Mupad [B] (verification not implemented)

Time = 7.61 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.13

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{\tan(c+dx) \left(\frac{A}{b} - \frac{Ba}{b^2}\right)}{d}$$

$$- \frac{\ln(\tan(c+dx) - i) (-B + A i)}{2d (-b + a i)}$$

$$+ \frac{\ln(a + b\tan(c+dx)) (Ba^4 - Aa^3b)}{d (a^2b^3 + b^5)}$$

$$- \frac{\ln(\tan(c+dx) + i) (A - B i)}{2d (a - b i)}$$

$$+ \frac{B\tan(c+dx)^2}{2bd}$$

input `int((tan(c + d*x)^3*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `(tan(c + d*x)*(A/b - (B*a)/b^2))/d - (log(tan(c + d*x) - 1i)*(A*1i - B))/(2*d*(a*1i - b)) + (log(a + b*tan(c + d*x))*(B*a^4 - A*a^3*b))/(d*(b^5 + a^2*b^3)) - (log(tan(c + d*x) + 1i)*(A - B*1i))/(2*d*(a - b*1i)) + (B*tan(c + d*x)^2)/(2*b*d)`

3.268 $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.268.1 Optimal result 2598
 3.268.2 Mathematica [C] (verified) 2598
 3.268.3 Rubi [A] (verified) 2599
 3.268.4 Maple [A] (verified) 2601
 3.268.5 Fricas [A] (verification not implemented) 2602
 3.268.6 Sympy [C] (verification not implemented) 2602
 3.268.7 Maxima [A] (verification not implemented) 2603
 3.268.8 Giac [A] (verification not implemented) 2604
 3.268.9 Mupad [B] (verification not implemented) 2604

3.268.1 Optimal result

Integrand size = 31, antiderivative size = 101

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = -\frac{(aA+bB)x}{a^2+b^2} - \frac{(Ab-aB) \log(\cos(c+dx))}{(a^2+b^2)d} + \frac{a^2(Ab-aB) \log(a+b \tan(c+dx))}{b^2(a^2+b^2)d} + \frac{B \tan(c+dx)}{bd}$$

output `-(A*a+B*b)*x/(a^2+b^2)-(A*b-B*a)*ln(cos(d*x+c))/(a^2+b^2)/d+a^2*(A*b-B*a)*ln(a+b*tan(d*x+c))/b^2/(a^2+b^2)/d+B*tan(d*x+c)/b/d`

3.268.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.66 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{\frac{i(A+ib) \log(i-\tan(c+dx))}{a+ib} - \frac{(iA+B) \log(i+\tan(c+dx))}{a-ib} + \frac{2a^2(Ab-aB) \log(a+b \tan(c+dx))}{b^2(a^2+b^2)} + \frac{2B \tan(c+dx)}{b}}{2d}$$

input `Integrate[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `((I*(A + I*B)*Log[I - Tan[c + d*x]])/(a + I*b) - ((I*A + B)*Log[I + Tan[c + d*x]])/(a - I*b) + (2*a^2*(A*b - a*B)*Log[a + b*Tan[c + d*x]])/(b^2*(a^2 + b^2)) + (2*B*Tan[c + d*x])/b)/(2*d)`

3.268.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.07, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 4089, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^2(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx \\
 & \quad \downarrow \text{4089} \\
 & \frac{\int -\frac{((Ab-aB)\tan^2(c+dx))+bB\tan(c+dx)+aB}{a+b\tan(c+dx)} dx}{b} + \frac{B\tan(c+dx)}{bd} \\
 & \quad \downarrow \text{25} \\
 & \frac{B\tan(c+dx)}{bd} - \frac{\int -\frac{((Ab-aB)\tan^2(c+dx))+bB\tan(c+dx)+aB}{a+b\tan(c+dx)} dx}{b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{B\tan(c+dx)}{bd} - \frac{\int -\frac{((Ab-aB)\tan(c+dx)^2)+bB\tan(c+dx)+aB}{a+b\tan(c+dx)} dx}{b} \\
 & \quad \downarrow \text{4109} \\
 & \frac{B\tan(c+dx)}{bd} - \frac{a^2(Ab-aB)\int \frac{\tan^2(c+dx)+1}{a+b\tan(c+dx)} dx}{a^2+b^2} - \frac{b(Ab-aB)\int \tan(c+dx) dx}{a^2+b^2} + \frac{bx(aA+bB)}{a^2+b^2} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.268. $\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$

$$\begin{aligned} & \frac{B \tan(c+dx)}{bd} - \frac{\frac{a^2(Ab-aB) \int \frac{\tan(c+dx)^2+1}{a+b \tan(c+dx)} dx}{a^2+b^2} - \frac{b(Ab-aB) \int \tan(c+dx) dx}{a^2+b^2} + \frac{bx(aA+bB)}{a^2+b^2}}{b} \\ & \quad \downarrow \text{3956} \\ & \frac{B \tan(c+dx)}{bd} - \frac{\frac{a^2(Ab-aB) \int \frac{\tan(c+dx)^2+1}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{b(Ab-aB) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{bx(aA+bB)}{a^2+b^2}}{b} \\ & \quad \downarrow \text{4100} \\ & \frac{B \tan(c+dx)}{bd} - \frac{\frac{a^2(Ab-aB) \int \frac{1}{a+b \tan(c+dx)} d(b \tan(c+dx))}{bd(a^2+b^2)} + \frac{b(Ab-aB) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{bx(aA+bB)}{a^2+b^2}}{b} \\ & \quad \downarrow \text{16} \\ & \frac{B \tan(c+dx)}{bd} - \frac{\frac{a^2(Ab-aB) \log(a+b \tan(c+dx))}{bd(a^2+b^2)} + \frac{b(Ab-aB) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{bx(aA+bB)}{a^2+b^2}}{b} \end{aligned}$$

input `Int[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `-(((b*(a*A + b*B)*x)/(a^2 + b^2) + (b*(A*b - a*B)*Log[Cos[c + d*x]])/((a^2 + b^2)*d) - (a^2*(A*b - a*B)*Log[a + b*Tan[c + d*x]])/(b*(a^2 + b^2)*d))/b) + (B*Tan[c + d*x])/(b*d)`

3.268.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_.)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4089 `Int[(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]))/((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[b^2 *B*(Tan[e + f*x]/(d*f)), x] + Simp[1/d Int[(a^2*A*d - b^2*B*c + (2*a*A*b + B*(a^2 - b^2))*d*Tan[e + f*x] + (A*b^2*d - b*B*(b*c - 2*a*d))*Tan[e + f*x]^2)/(c + d*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4100 `Int[(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]`

rule 4109 `Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C, 0]`

3.268.4 Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{\frac{\tan(dx+c)B}{b} + \frac{(Ab-Ba)\ln(1+\tan^2(dx+c))}{2} + \frac{(-aA-Bb)\arctan(\tan(dx+c))}{a^2+b^2} + \frac{a^2(Ab-Ba)\ln(a+b\tan(dx+c))}{b^2(a^2+b^2)}}{d}$
default	$\frac{\frac{\tan(dx+c)B}{b} + \frac{(Ab-Ba)\ln(1+\tan^2(dx+c))}{2} + \frac{(-aA-Bb)\arctan(\tan(dx+c))}{a^2+b^2} + \frac{a^2(Ab-Ba)\ln(a+b\tan(dx+c))}{b^2(a^2+b^2)}}{d}$
norman	$\frac{B\tan(dx+c)}{bd} - \frac{(aA+Bb)x}{a^2+b^2} + \frac{a^2(Ab-Ba)\ln(a+b\tan(dx+c))}{b^2(a^2+b^2)d} + \frac{(Ab-Ba)\ln(1+\tan^2(dx+c))}{2d(a^2+b^2)}$
parallelrisch	$\frac{-2Aa b^2 dx - 2B b^3 dx + A \ln(1+\tan^2(dx+c)) b^3 + 2A \ln(a+b\tan(dx+c)) a^2 b - B \ln(1+\tan^2(dx+c)) a b^2 - 2B \ln(a+b\tan(dx+c)) b^2}{2d(a^2+b^2)b^2}$
risch	$-\frac{ixB}{ib-a} + \frac{xA}{ib-a} + \frac{2iAx}{b} + \frac{2iAc}{bd} - \frac{2iaBx}{b^2} - \frac{2iBac}{b^2d} - \frac{2ia^2Ax}{(a^2+b^2)b} - \frac{2ia^2Ac}{d(a^2+b^2)b} + \frac{2ia^3Bx}{b^2(a^2+b^2)} + \frac{2ia^3Bc}{d(a^2+b^2)b^2}$

input `int(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

$$3.268. \int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

output $1/d*(\tan(dx+c)*B/b+1/(a^2+b^2)*(1/2*(A*b-B*a)*\ln(1+\tan(dx+c)^2)+(-A*a-B*b)*\arctan(\tan(dx+c))))+1/b^2*a^2*(A*b-B*a)/(a^2+b^2)*\ln(a+b*\tan(dx+c))$

3.268.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.48

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{2(Aab^2+Bb^3)dx+(Ba^3-Aa^2b)\log\left(\frac{b^2\tan(dx+c)^2+2ab\tan(dx+c)+a^2}{\tan(dx+c)^2+1}\right)-(Ba^3-Aa^2b+Bab^2-Ab^3)\log}{2(a^2b^2+b^4)d}$$

input `integrate(tan(dx+c)^2*(A+B*tan(dx+c))/(a+b*tan(dx+c)),x, algorithm="fricas")`

output $-1/2*(2*(A*a*b^2+B*b^3)*dx+(B*a^3-A*a^2*b)*\log((b^2*\tan(dx+c)^2+2*a*b*\tan(dx+c)+a^2)/(\tan(dx+c)^2+1))-(B*a^3-A*a^2*b+B*a*b^2-A*b^3)*\log(1/(\tan(dx+c)^2+1))-2*(B*a^2*b+B*b^3)*\tan(dx+c))/((a^2*b^2+b^4)*d)$

3.268.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.69 (sec) , antiderivative size = 1015, normalized size of antiderivative = 10.05

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

input `integrate(tan(dx+c)**2*(A+B*tan(dx+c))/(a+b*tan(dx+c)),x)`

output `Piecewise((zoo*x*(A + B*tan(c))*tan(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (-A*x + A*tan(c + d*x)/d - B*log(tan(c + d*x)**2 + 1)/(2*d) + B*tan(c + d*x)**2/(2*d))/a, Eq(b, 0)), (I*A*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + A*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) + A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*A*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*A/(2*b*d*tan(c + d*x) - 2*I*b*d) - 3*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 3*I*B*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) + I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 2*B*tan(c + d*x)**2/(2*b*d*tan(c + d*x) - 2*I*b*d) + 3*B/(2*b*d*tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (-I*A*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + A*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) + A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*A*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*A/(2*b*d*tan(c + d*x) + 2*I*b*d) - 3*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) - 3*I*B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2*I*b*d) + 2*B*tan(c + d*x)**2/(2*b*d*tan(c + d*x) + 2*I*b*d) + 3*B/(2*b*d*tan(c + d*x) + 2*I*b*d), Eq(a, I*b)), (x*(A + B*tan(c))*tan(c)**2/(a + b*tan(c)), Eq(d, 0)), (2*A*a**2*b*log(a/b + tan(c + d*x))/(2*a**2*b...`

3.268.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.08

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{a + b \tan(c + dx)} dx$$

$$= -\frac{\frac{2(Aa+Bb)(dx+c)}{a^2+b^2} + \frac{2(Ba^3-Aa^2b) \log(b \tan(dx+c)+a)}{a^2b^2+b^4} + \frac{(Ba-Ab) \log(\tan(dx+c)^2+1)}{a^2+b^2} - \frac{2B \tan(dx+c)}{b}}{2d}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(2*(A*a + B*b)*(d*x + c)/(a^2 + b^2) + 2*(B*a^3 - A*a^2*b)*log(b*tan(d*x + c) + a)/(a^2*b^2 + b^4) + (B*a - A*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) - 2*B*tan(d*x + c)/b)/d`

3.268.8 Giac [A] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.09

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{a + b \tan(c + dx)} dx$$

$$= -\frac{\frac{2(Aa+Bb)(dx+c)}{a^2+b^2} + \frac{(Ba-Ab) \log(\tan(dx+c)^2+1)}{a^2+b^2} + \frac{2(Ba^3-Aa^2b) \log(|b \tan(dx+c)+a|)}{a^2b^2+b^4} - \frac{2B \tan(dx+c)}{b}}{2d}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `-1/2*(2*(A*a + B*b)*(d*x + c)/(a^2 + b^2) + (B*a - A*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) + 2*(B*a^3 - A*a^2*b)*log(abs(b*tan(d*x + c) + a))/(a^2*b^2 + b^4) - 2*B*tan(d*x + c)/b)/d`

3.268.9 Mupad [B] (verification not implemented)

Time = 7.84 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.16

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{a + b \tan(c + dx)} dx = \frac{B \tan(c + dx)}{bd} + \frac{\ln(\tan(c + dx) + 1i)(A - B 1i)}{2d(b + a 1i)}$$

$$- \frac{\ln(a + b \tan(c + dx))(B a^3 - A a^2 b)}{d(a^2 b^2 + b^4)}$$

$$+ \frac{\ln(\tan(c + dx) - 1i)(-B + A 1i)}{2d(a + b 1i)}$$

input `int((tan(c + d*x)^2*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `(log(tan(c + d*x) + 1i)*(A - B*1i))/(2*d*(a*1i + b)) - (log(a + b*tan(c + d*x))*(B*a^3 - A*a^2*b))/(d*(b^4 + a^2*b^2)) + (B*tan(c + d*x))/(b*d) + (log(tan(c + d*x) - 1i)*(A*1i - B))/(2*d*(a + b*1i))`

3.269 $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.269.1 Optimal result 2605
 3.269.2 Mathematica [C] (verified) 2605
 3.269.3 Rubi [A] (verified) 2606
 3.269.4 Maple [A] (verified) 2608
 3.269.5 Fricas [A] (verification not implemented) 2608
 3.269.6 Sympy [C] (verification not implemented) 2609
 3.269.7 Maxima [A] (verification not implemented) 2610
 3.269.8 Giac [A] (verification not implemented) 2611
 3.269.9 Mupad [B] (verification not implemented) 2611

3.269.1 Optimal result

Integrand size = 29, antiderivative size = 80

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{a + b \tan(c + dx)} dx = \frac{(Ab - aB)x}{a^2 + b^2} - \frac{B \log(\cos(c + dx))}{bd} - \frac{a(Ab - aB) \log(a \cos(c + dx) + b \sin(c + dx))}{b(a^2 + b^2)d}$$

output `(A*b-B*a)*x/(a^2+b^2)-B*ln(cos(d*x+c))/b/d-a*(A*b-B*a)*ln(a*cos(d*x+c)+b*sin(d*x+c))/b/(a^2+b^2)/d`

3.269.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.22

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{a + b \tan(c + dx)} dx = \frac{(a - ib)b(A + iB) \log(i - \tan(c + dx)) + (a + ib)b(A - iB) \log(i + \tan(c + dx)) + 2a(-Ab + aB) \log(a + b \tan(c + dx))}{2b(a^2 + b^2)d}$$

input `Integrate[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output $((a - I*b)*b*(A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]] + (a + I*b)*b*(A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]] + 2*a*(-(A*b) + a*B)*\text{Log}[a + b*\text{Tan}[c + d*x]])/(2*b*(a^2 + b^2)*d)$

3.269.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.94, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 4072, 27, 3042, 3956, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx \\ & \quad \downarrow 3042 \\ & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx \\ & \quad \downarrow 4072 \\ & \frac{\int \frac{(Ab-aB)\tan(c+dx)}{a+b\tan(c+dx)} dx}{b} + \frac{B \int \tan(c+dx) dx}{b} \\ & \quad \downarrow 27 \\ & \frac{(Ab-aB) \int \frac{\tan(c+dx)}{a+b\tan(c+dx)} dx}{b} + \frac{B \int \tan(c+dx) dx}{b} \\ & \quad \downarrow 3042 \\ & \frac{(Ab-aB) \int \frac{\tan(c+dx)}{a+b\tan(c+dx)} dx}{b} + \frac{B \int \tan(c+dx) dx}{b} \\ & \quad \downarrow 3956 \\ & \frac{(Ab-aB) \int \frac{\tan(c+dx)}{a+b\tan(c+dx)} dx}{b} - \frac{B \log(\cos(c+dx))}{bd} \\ & \quad \downarrow 4014 \\ & \frac{(Ab-aB) \left(\frac{bx}{a^2+b^2} - \frac{a \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a^2+b^2} \right)}{b} - \frac{B \log(\cos(c+dx))}{bd} \\ & \quad \downarrow 3042 \end{aligned}$$

$$\frac{(Ab - aB) \left(\frac{bx}{a^2 + b^2} - \frac{a \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2 + b^2} \right)}{b} - \frac{B \log(\cos(c + dx))}{bd}$$

↓ 4013

$$\frac{(Ab - aB) \left(\frac{bx}{a^2 + b^2} - \frac{a \log(a \cos(c+dx) + b \sin(c+dx))}{d(a^2 + b^2)} \right)}{b} - \frac{B \log(\cos(c + dx))}{bd}$$

input `Int[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `-((B*Log[Cos[c + d*x]])/(b*d)) + ((A*b - a*B)*((b*x)/(a^2 + b^2) - (a*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d)))/b`

3.269.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*(x_), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

```
rule 4072 Int[(((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*(d/b) Int[Tan[e + f*x], x], x] + Simp[1/b Int[Simp[A*b*c + (A*b*d + B*(b*c - a*d))*Tan[e + f*x], x]/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0]
```

3.269.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.09

method	result
derivativedivides	$\frac{\frac{(aA+Bb)\ln(1+\tan^2(dx+c))}{2} + (Ab-Ba)\arctan(\tan(dx+c)) - \frac{a(Ab-Ba)\ln(a+b\tan(dx+c))}{(a^2+b^2)b}}{d}$
default	$\frac{\frac{(aA+Bb)\ln(1+\tan^2(dx+c))}{2} + (Ab-Ba)\arctan(\tan(dx+c)) - \frac{a(Ab-Ba)\ln(a+b\tan(dx+c))}{(a^2+b^2)b}}{d}$
norman	$\frac{(Ab-Ba)x}{a^2+b^2} + \frac{(aA+Bb)\ln(1+\tan^2(dx+c))}{2d(a^2+b^2)} - \frac{a(Ab-Ba)\ln(a+b\tan(dx+c))}{(a^2+b^2)bd}$
parallelrisch	$\frac{2Ab^2dx - 2Babdx + A\ln(1+\tan^2(dx+c))ab - 2A\ln(a+b\tan(dx+c))ab + B\ln(1+\tan^2(dx+c))b^2 + 2B\ln(a+b\tan(dx+c))}{2(a^2+b^2)bd}$
risch	$\frac{xB}{ib-a} + \frac{ixA}{ib-a} + \frac{2iaAx}{a^2+b^2} + \frac{2iaAc}{d(a^2+b^2)} - \frac{2ia^2Bx}{b(a^2+b^2)} - \frac{2ia^2Bc}{bd(a^2+b^2)} + \frac{2iBx}{b} + \frac{2iBc}{bd} - \frac{a\ln\left(e^{2i(dx+c)} - \frac{ib+a}{ib-a}\right)A}{d(a^2+b^2)}$

```
input int(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/d*(1/(a^2+b^2)*(1/2*(A*a+B*b)*ln(1+tan(d*x+c)^2)+(A*b-B*a)*arctan(tan(d*x+c)))-a*(A*b-B*a)/(a^2+b^2)/b*ln(a+b*tan(d*x+c)))
```

3.269.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.38

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{2(Bab - Ab^2)dx - (Ba^2 - Aab) \log\left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2 + 1}\right) + (Ba^2 + Bb^2) \log\left(\frac{1}{\tan(dx+c)^2 + 1}\right)}{2(a^2b + b^3)d}$$

3.269. $\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output `-1/2*(2*(B*a*b - A*b^2)*d*x - (B*a^2 - A*a*b)*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) + (B*a^2 + B*b^2)*log(1/(tan(d*x + c)^2 + 1)))/((a^2*b + b^3)*d)`

3.269.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.55 (sec) , antiderivative size = 700, normalized size of antiderivative = 8.75

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \left\{ \begin{array}{l} \tilde{\infty}x(A+B\tan(c)) \\ \frac{A \log(\tan^2(c+dx)+1)}{2d} - Bx + \frac{B \tan(c+dx)}{d} \\ \frac{A dx \tan(c+dx)}{2bd \tan(c+dx)-2ibd} - \frac{iA dx}{2bd \tan(c+dx)-2ibd} - \frac{A}{2bd \tan(c+dx)-2ibd} + \frac{iB dx \tan(c+dx)}{2bd \tan(c+dx)-2ibd} + \frac{B dx}{2bd \tan(c+dx)-2ibd} + \frac{B \log(\tan^2(c+dx)+1)}{2bd \tan(c+dx)-2ibd} \\ \frac{A dx \tan(c+dx)}{2bd \tan(c+dx)+2ibd} + \frac{iA dx}{2bd \tan(c+dx)+2ibd} - \frac{A}{2bd \tan(c+dx)+2ibd} - \frac{iB dx \tan(c+dx)}{2bd \tan(c+dx)+2ibd} + \frac{B dx}{2bd \tan(c+dx)+2ibd} + \frac{B \log(\tan^2(c+dx)+1)}{2bd \tan(c+dx)+2ibd} \\ \frac{x(A+B\tan(c)) \tan(c)}{a+b\tan(c)} \\ -\frac{2Aab \log(\frac{a}{b}+\tan(c+dx))}{2a^2bd+2b^3d} + \frac{Aab \log(\tan^2(c+dx)+1)}{2a^2bd+2b^3d} + \frac{2Ab^2 dx}{2a^2bd+2b^3d} + \frac{2Ba^2 \log(\frac{a}{b}+\tan(c+dx))}{2a^2bd+2b^3d} - \frac{2Babd x}{2a^2bd+2b^3d} + \frac{Bb^2 \log(\tan^2(c+dx)+1)}{2a^2bd+2b^3d} \end{array} \right.$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Piecewise((zoo*x*(A + B*tan(c)), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), ((A*log(tan(c + d*x)**2 + 1)/(2*d) - B*x + B*tan(c + d*x)/d)/a, Eq(b, 0)), (A*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*A*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) - A/(2*b*d*tan(c + d*x) - 2*I*b*d) + I*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + B*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) + B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*B/(2*b*d*tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (A*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*A*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - A/(2*b*d*tan(c + d*x) + 2*I*b*d) - I*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) + B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*B/(2*b*d*tan(c + d*x) + 2*I*b*d), Eq(a, I*b)), (x*(A + B*tan(c))*tan(c)/(a + b*tan(c)), Eq(d, 0)), (-2*A*a*b*log(a/b + tan(c + d*x))/(2*a**2*b*d + 2*b**3*d) + A*a*b*log(tan(c + d*x)**2 + 1)/(2*a**2*b*d + 2*b**3*d) + 2*A*b**2*d*x/(2*a**2*b*d + 2*b**3*d) + 2*B*a**2*log(a/b + tan(c + d*x))/(2*a**2*b*d + 2*b**3*d) - 2*B*a*b*d*x/(2*a**2*b*d + 2*b**3*d) + B*b**2*log(tan(c + d*x)**2 + 1)/(2*a**2*b*d + 2*b**3*d), True))`

3.269.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.18

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= -\frac{\frac{2(Ba-Ab)(dx+c)}{a^2+b^2} - \frac{2(Ba^2-Aab)\log(b\tan(dx+c)+a)}{a^2b+b^3} - \frac{(Aa+Bb)\log(\tan(dx+c)^2+1)}{a^2+b^2}}{2d}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(2*(B*a - A*b)*(d*x + c)/(a^2 + b^2) - 2*(B*a^2 - A*a*b)*log(b*tan(d*x + c) + a)/(a^2*b + b^3) - (A*a + B*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2))/d`

3.269.8 Giac [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.19

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= -\frac{\frac{2(Ba-Ab)(dx+c)}{a^2+b^2} - \frac{(Aa+Bb)\log(\tan(dx+c)^2+1)}{a^2+b^2} - \frac{2(Ba^2-Aab)\log(|b\tan(dx+c)+a|)}{a^2b+b^3}}{2d}$$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")
```

```
output -1/2*(2*(B*a - A*b)*(d*x + c)/(a^2 + b^2) - (A*a + B*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) - 2*(B*a^2 - A*a*b)*log(abs(b*tan(d*x + c) + a))/(a^2*b + b^3))/d
```

3.269.9 Mupad [B] (verification not implemented)

Time = 7.71 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.25

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{\ln(\tan(c+dx)-i)(-B+Ai)}{2d(-b+Ai)} + \frac{\ln(\tan(c+dx)+i)(A-Bi)}{2d(a-bi)} - \frac{a \ln(a+b\tan(c+dx))(Ab-Ba)}{bd(a^2+b^2)}$$

```
input int((tan(c + d*x)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)),x)
```

```
output (log(tan(c + d*x) - 1i)*(A*1i - B))/(2*d*(a*1i - b)) + (log(tan(c + d*x) + 1i)*(A - B*1i))/(2*d*(a - b*1i)) - (a*log(a + b*tan(c + d*x))*(A*b - B*a))/(b*d*(a^2 + b^2))
```

3.270 $\int \frac{A+B \tan(c+dx)}{a+b \tan(c+dx)} dx$

3.270.1 Optimal result	2612
3.270.2 Mathematica [A] (verified)	2612
3.270.3 Rubi [A] (verified)	2613
3.270.4 Maple [A] (verified)	2614
3.270.5 Fricas [A] (verification not implemented)	2615
3.270.6 Sympy [C] (verification not implemented)	2615
3.270.7 Maxima [A] (verification not implemented)	2616
3.270.8 Giac [A] (verification not implemented)	2616
3.270.9 Mupad [B] (verification not implemented)	2617

3.270.1 Optimal result

Integrand size = 23, antiderivative size = 58

$$\int \frac{A + B \tan(c + dx)}{a + b \tan(c + dx)} dx = \frac{(aA + bB)x}{a^2 + b^2} + \frac{(Ab - aB) \log(a \cos(c + dx) + b \sin(c + dx))}{(a^2 + b^2) d}$$

output $(A*a+B*b)*x/(a^2+b^2)+(A*b-B*a)*\ln(a*\cos(d*x+c)+b*\sin(d*x+c))/(a^2+b^2)/d$

3.270.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.14

$$\int \frac{A + B \tan(c + dx)}{a + b \tan(c + dx)} dx = \frac{2(aA + bB) \arctan(\tan(c + dx)) - (Ab - aB) (\log(\sec^2(c + dx)) - 2 \log(a + b \tan(c + dx)))}{2(a^2 + b^2) d}$$

input `Integrate[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x]),x]`

output $(2*(a*A + b*B)*ArcTan[Tan[c + d*x]] - (A*b - a*B)*(Log[Sec[c + d*x]^2] - 2 *Log[a + b*Tan[c + d*x]]))/(2*(a^2 + b^2)*d)$

3.270.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{a + b \tan(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{a + b \tan(c + dx)} dx \\
 & \quad \downarrow \text{4014} \\
 & \frac{(Ab - aB) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{x(aA + bB)}{a^2 + b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(Ab - aB) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{x(aA + bB)}{a^2 + b^2} \\
 & \quad \downarrow \text{4013} \\
 & \frac{(Ab - aB) \log(a \cos(c + dx) + b \sin(c + dx))}{d(a^2 + b^2)} + \frac{x(aA + bB)}{a^2 + b^2}
 \end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x]),x]`

output `((a*A + b*B)*x)/(a^2 + b^2) + ((A*b - a*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d)`

3.270.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

3.270.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.41

method	result
derivativedivides	$\frac{\frac{(-Ab+Ba)\ln(1+\tan^2(dx+c))}{2} + (aA+Bb)\arctan(\tan(dx+c)) + \frac{(Ab-Ba)\ln(a+b\tan(dx+c))}{a^2+b^2}}{d}$
default	$\frac{\frac{(-Ab+Ba)\ln(1+\tan^2(dx+c))}{2} + (aA+Bb)\arctan(\tan(dx+c)) + \frac{(Ab-Ba)\ln(a+b\tan(dx+c))}{a^2+b^2}}{d}$
norman	$\frac{(aA+Bb)x}{a^2+b^2} + \frac{(Ab-Ba)\ln(a+b\tan(dx+c))}{d(a^2+b^2)} - \frac{(Ab-Ba)\ln(1+\tan^2(dx+c))}{2d(a^2+b^2)}$
parallelrisch	$-\frac{-2Axad-2Bbdx+A\ln(1+\tan^2(dx+c))b-2A\ln(a+b\tan(dx+c))b-B\ln(1+\tan^2(dx+c))a+2B\ln(a+b\tan(dx+c))a}{2d(a^2+b^2)}$
risch	$\frac{ixB}{ib-a} - \frac{xA}{ib-a} - \frac{2iAbx}{a^2+b^2} + \frac{2iBxa}{a^2+b^2} - \frac{2iAbc}{d(a^2+b^2)} + \frac{2iBac}{d(a^2+b^2)} + \frac{\ln\left(e^{2i(dx+c)} - \frac{ib+a}{ib-a}\right)Ab}{d(a^2+b^2)} - \frac{\ln\left(e^{2i(dx+c)} - \frac{ib+a}{ib-a}\right)}{d(a^2+b^2)}$

input `int((A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*(1/(a^2+b^2)*(1/2*(-A*b+B*a)*ln(1+tan(d*x+c)^2)+(A*a+B*b)*arctan(tan(d*x+c)))+(A*b-B*a)/(a^2+b^2)*ln(a+b*tan(d*x+c)))`

3.270.
$$\int \frac{A+B \tan(c+dx)}{a+b \tan(c+dx)} dx$$

3.270.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.31

$$\int \frac{A + B \tan(c + dx)}{a + b \tan(c + dx)} dx = \frac{2(Aa + Bb)dx - (Ba - Ab) \log\left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2 + 1}\right)}{2(a^2 + b^2)d}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fracas")`output `1/2*(2*(A*a + B*b)*d*x - (B*a - A*b)*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)))/((a^2 + b^2)*d)`**3.270.6 Sympy [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.47 (sec) , antiderivative size = 524, normalized size of antiderivative = 9.03

$$\int \frac{A + B \tan(c + dx)}{a + b \tan(c + dx)} dx = \begin{cases} \frac{\infty x(A+B \tan(c))}{\tan(c)} \\ \frac{Ax + \frac{B \log(\tan^2(c+dx)+1)}{2d}}{a} \\ \frac{iAdx \tan(c+dx)}{2bd \tan(c+dx)-2ibd} + \frac{Adx}{2bd \tan(c+dx)-2ibd} + \frac{iA}{2bd \tan(c+dx)-2ibd} + \frac{Bdx \tan(c+dx)}{2bd \tan(c+dx)-2ibd} - \frac{iBdx}{2bd \tan(c+dx)-2ibd} - \frac{B}{2bd \tan(c+dx)} \\ - \frac{iAdx \tan(c+dx)}{2bd \tan(c+dx)+2ibd} + \frac{Adx}{2bd \tan(c+dx)+2ibd} - \frac{iA}{2bd \tan(c+dx)+2ibd} + \frac{Bdx \tan(c+dx)}{2bd \tan(c+dx)+2ibd} + \frac{iBdx}{2bd \tan(c+dx)+2ibd} - \frac{B}{2bd \tan(c+dx)} \\ \frac{x(A+B \tan(c))}{a+b \tan(c)} \\ \frac{2Aadx}{2a^2d+2b^2d} + \frac{2Ab \log(\frac{a}{b} + \tan(c+dx))}{2a^2d+2b^2d} - \frac{Ab \log(\tan^2(c+dx)+1)}{2a^2d+2b^2d} - \frac{2Ba \log(\frac{a}{b} + \tan(c+dx))}{2a^2d+2b^2d} + \frac{Ba \log(\tan^2(c+dx)+1)}{2a^2d+2b^2d} + \frac{2Bbdx}{2a^2d+2b^2d} \end{cases}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Piecewise((zoo*x*(A + B*tan(c))/tan(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), ((A*x + B*log(tan(c + d*x)**2 + 1)/(2*d))/a, Eq(b, 0)), (I*A*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + A*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) + I*A/(2*b*d*tan(c + d*x) - 2*I*b*d) + B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*B*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) - B/(2*b*d*tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (-I*A*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + A*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - I*A/(2*b*d*tan(c + d*x) + 2*I*b*d) + B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - B/(2*b*d*tan(c + d*x) + 2*I*b*d), Eq(a, I*b)), (x*(A + B*tan(c))/(a + b*tan(c)), Eq(d, 0)), (2*A*a*d*x/(2*a**2*d + 2*b**2*d) + 2*A*b*log(a/b + tan(c + d*x))/(2*a**2*d + 2*b**2*d) - A*b*log(tan(c + d*x)**2 + 1)/(2*a**2*d + 2*b**2*d) - 2*B*a*log(a/b + tan(c + d*x))/(2*a**2*d + 2*b**2*d) + B*a*log(tan(c + d*x)**2 + 1)/(2*a**2*d + 2*b**2*d) + 2*B*b*d*x/(2*a**2*d + 2*b**2*d), True))`

3.270.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.52

$$\int \frac{A + B \tan(c + dx)}{a + b \tan(c + dx)} dx = \frac{\frac{2(Aa+Bb)(dx+c)}{a^2+b^2} - \frac{2(Ba-Ab)\log(b\tan(dx+c)+a)}{a^2+b^2} + \frac{(Ba-Ab)\log(\tan(dx+c)^2+1)}{a^2+b^2}}{2d}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(2*(A*a + B*b)*(d*x + c)/(a^2 + b^2) - 2*(B*a - A*b)*log(b*tan(d*x + c) + a)/(a^2 + b^2) + (B*a - A*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2))/d`

3.270.8 Giac [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.62

$$\int \frac{A + B \tan(c + dx)}{a + b \tan(c + dx)} dx = \frac{\frac{2(Aa+Bb)(dx+c)}{a^2+b^2} + \frac{(Ba-Ab)\log(\tan(dx+c)^2+1)}{a^2+b^2} - \frac{2(Bab-Ab^2)\log(|b\tan(dx+c)+a|)}{a^2b+b^3}}{2d}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

3.270. $\int \frac{A+B \tan(c+dx)}{a+b \tan(c+dx)} dx$

output $1/2*(2*(A*a + B*b)*(d*x + c)/(a^2 + b^2) + (B*a - A*b)*\log(\tan(d*x + c)^2 + 1)/(a^2 + b^2) - 2*(B*a*b - A*b^2)*\log(\text{abs}(b*\tan(d*x + c) + a))/(a^2*b + b^3))/d$

3.270.9 Mupad [B] (verification not implemented)

Time = 7.80 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.60

$$\int \frac{A + B \tan(c + dx)}{a + b \tan(c + dx)} dx = \frac{\ln(a + b \tan(c + dx)) (Ab - Ba)}{d (a^2 + b^2)} - \frac{\ln(\tan(c + dx) + 1i) (A - B 1i)}{2d (b + a 1i)} - \frac{\ln(\tan(c + dx) - 1i) (-B + A 1i)}{2d (a + b 1i)}$$

input `int((A + B*tan(c + d*x))/(a + b*tan(c + d*x)),x)`

output $(\log(a + b*\tan(c + d*x))*(A*b - B*a))/(d*(a^2 + b^2)) - (\log(\tan(c + d*x) + 1i)*(A - B*1i))/(2*d*(a*1i + b)) - (\log(\tan(c + d*x) - 1i)*(A*1i - B))/(2*d*(a + b*1i))$

3.271 $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.271.1 Optimal result 2618
 3.271.2 Mathematica [C] (verified) 2618
 3.271.3 Rubi [A] (verified) 2619
 3.271.4 Maple [A] (verified) 2621
 3.271.5 Fricas [A] (verification not implemented) 2621
 3.271.6 Sympy [C] (verification not implemented) 2622
 3.271.7 Maxima [A] (verification not implemented) 2622
 3.271.8 Giac [A] (verification not implemented) 2623
 3.271.9 Mupad [B] (verification not implemented) 2623

3.271.1 Optimal result

Integrand size = 29, antiderivative size = 80

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = -\frac{(Ab-aB)x}{a^2+b^2} + \frac{A \log(\sin(c+dx))}{ad} - \frac{b(Ab-aB) \log(a \cos(c+dx)+b \sin(c+dx))}{a(a^2+b^2)d}$$

output `-(A*b-B*a)*x/(a^2+b^2)+A*ln(sin(d*x+c))/a/d-b*(A*b-B*a)*ln(a*cos(d*x+c)+b*sin(d*x+c))/a/(a^2+b^2)/d`

3.271.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.43 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.41

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = -\frac{(A+iB) \log(i-\tan(c+dx))}{a+ib} - \frac{2A \log(\tan(c+dx))}{a} + \frac{(A-iB) \log(i+\tan(c+dx))}{a-ib} + \frac{2b(Ab-aB) \log(a+b \tan(c+dx))}{a(a^2+b^2)}$$

input `Integrate[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output
$$\frac{-1/2*((A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]]/(a + I*b) - (2*A*\text{Log}[\text{Tan}[c + d*x]])/a + ((A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]]/(a - I*b) + (2*b*(A*b - a*B)*\text{Log}[a + b*\text{Tan}[c + d*x]])/(a*(a^2 + b^2)))/d$$

3.271.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.02, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.207$, Rules used = {3042, 4094, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx \\ & \quad \downarrow 3042 \\ & \int \frac{A+B\tan(c+dx)}{\tan(c+dx)(a+b\tan(c+dx))} dx \\ & \quad \downarrow 4094 \\ & -\frac{b(Ab-aB) \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} + \frac{A \int \cot(c+dx) dx}{a} - \frac{x(Ab-aB)}{a^2+b^2} \\ & \quad \downarrow 3042 \\ & -\frac{b(Ab-aB) \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} + \frac{A \int -\tan(c+dx+\frac{\pi}{2}) dx}{a} - \frac{x(Ab-aB)}{a^2+b^2} \\ & \quad \downarrow 25 \\ & -\frac{b(Ab-aB) \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} - \frac{A \int \tan(\frac{1}{2}(2c+\pi)+dx) dx}{a} - \frac{x(Ab-aB)}{a^2+b^2} \\ & \quad \downarrow 3956 \\ & -\frac{b(Ab-aB) \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} - \frac{x(Ab-aB)}{a^2+b^2} + \frac{A \log(-\sin(c+dx))}{ad} \\ & \quad \downarrow 4013 \\ & -\frac{b(Ab-aB) \log(a \cos(c+dx) + b \sin(c+dx))}{ad(a^2+b^2)} - \frac{x(Ab-aB)}{a^2+b^2} + \frac{A \log(-\sin(c+dx))}{ad} \end{aligned}$$

3.271.
$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

input `Int[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `-(((A*b - a*B)*x)/(a^2 + b^2)) + (A*Log[-Sin[c + d*x]])/(a*d) - (b*(A*b - a*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d)`

3.271.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4094 `Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(B*(b*c + a*d) + A*(a*c - b*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[b*((A*b - a*B)/((b*c - a*d)*(a^2 + b^2))) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] + Simp[d*((B*c - A*d)/((b*c - a*d)*(c^2 + d^2))) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.271.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.19

method	result
parallelrisch	$\frac{(-2Ab^2+2Bab)\ln(a+b\tan(dx+c))+(-Aa^2-Bab)\ln(\sec^2(dx+c))+2A(a^2+b^2)\ln(\tan(dx+c))-2adx(Ab-Ba)}{2(a^2+b^2)ad}$
derivativedivides	$\frac{\frac{(-aA-Bb)\ln(1+\tan^2(dx+c))}{2}+(-Ab+Ba)\arctan(\tan(dx+c))+\frac{A\ln(\tan(dx+c))}{a}-\frac{(Ab-Ba)b\ln(a+b\tan(dx+c))}{(a^2+b^2)a}}{d}$
default	$\frac{\frac{(-aA-Bb)\ln(1+\tan^2(dx+c))}{2}+(-Ab+Ba)\arctan(\tan(dx+c))+\frac{A\ln(\tan(dx+c))}{a}-\frac{(Ab-Ba)b\ln(a+b\tan(dx+c))}{(a^2+b^2)a}}{d}$
norman	$-\frac{(Ab-Ba)x}{a^2+b^2}+\frac{A\ln(\tan(dx+c))}{ad}-\frac{(aA+Bb)\ln(1+\tan^2(dx+c))}{2d(a^2+b^2)}-\frac{(Ab-Ba)b\ln(a+b\tan(dx+c))}{(a^2+b^2)ad}$
risch	$-\frac{xB}{ib-a}-\frac{ixA}{ib-a}-\frac{2ixA}{a}-\frac{2iAc}{ad}+\frac{2ib^2Ax}{a(a^2+b^2)}+\frac{2ib^2Ac}{ad(a^2+b^2)}-\frac{2ibBx}{a^2+b^2}-\frac{2ibBc}{d(a^2+b^2)}+\frac{A\ln(e^{2i(dx+c)}-1)}{ad}$

input `int(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{2}*((-2A*b^2+2B*a*b)*\ln(a+b*\tan(d*x+c))+(-A*a^2-B*a*b)*\ln(\sec(d*x+c)^2)+2*A*(a^2+b^2)*\ln(\tan(d*x+c))-2*a*d*x*(A*b-B*a))/(a^2+b^2)/a/d$$

3.271.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.48

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{2(Ba^2 - Aab)dx + (Aa^2 + Ab^2) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) + (Bab - Ab^2) \log\left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2+1}\right)}{2(a^3 + ab^2)d}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fracas")`

output
$$\frac{1}{2}*(2*(B*a^2 - A*a*b)*d*x + (A*a^2 + A*b^2)*\log(\tan(d*x + c)^2/(\tan(d*x + c)^2 + 1)) + (B*a*b - A*b^2)*\log((b^2*\tan(d*x + c)^2 + 2*a*b*\tan(d*x + c) + a^2)/(\tan(d*x + c)^2 + 1)))/((a^3 + a*b^2)*d)$$

3.271.
$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

3.271.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.05 (sec) , antiderivative size = 952, normalized size of antiderivative = 11.90

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Piecewise((zoo*x*(A + B*tan(c))*cot(c)/tan(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), ((-A*log(tan(c + d*x)**2 + 1)/(2*d) + A*log(tan(c + d*x))/d + B*x)/a, Eq(b, 0)), ((-A*x - A/(d*tan(c + d*x)) - B*log(tan(c + d*x)**2 + 1)/(2*d) + B*log(tan(c + d*x))/d)/b, Eq(a, 0)), (A*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*A*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - A*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 2*I*A*log(tan(c + d*x))*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 2*A*log(tan(c + d*x))/(2*b*d*tan(c + d*x) - 2*I*b*d) + A/(2*b*d*tan(c + d*x) - 2*I*b*d) + I*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + B*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) + I*B/(2*b*d*tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (A*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*A*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) - A*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2*I*b*d) - 2*I*A*log(tan(c + d*x))*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + 2*A*log(tan(c + d*x))/(2*b*d*tan(c + d*x) + 2*I*b*d) + A/(2*b*d*tan(c + d*x) + 2*I*b*d) - I*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - I*B/(2*b*d*tan(c + d*x) + 2*I*b*d), Eq(a, I*b)), (x*(A + B*tan(c))*cot(c)/(a + b*tan(c)), Eq(d, 0)), (-A*a**2*log(tan(c + d*x)**2 + 1)/(2*a**3*d + 2*a*b**2*d) + 2*A*a**2*log(tan(c + d*x))/(2*a**3*d + ...`

3.271.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.34

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{2(Ba-Ab)(dx+c)}{a^2+b^2} + \frac{2(Bab-Ab^2)\log(b\tan(dx+c)+a)}{a^3+ab^2} - \frac{(Aa+Bb)\log(\tan(dx+c)^2+1)}{a^2+b^2} + \frac{2A\log(\tan(dx+c))}{a}$$

$2d$

3.271. $\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(2*(B*a - A*b)*(d*x + c)/(a^2 + b^2) + 2*(B*a*b - A*b^2)*log(b*tan(d*x + c) + a)/(a^3 + a*b^2) - (A*a + B*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) + 2*A*log(tan(d*x + c))/a)/d`

3.271.8 Giac [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.41

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{\frac{2(Ba-Ab)(dx+c)}{a^2+b^2} - \frac{(Aa+Bb)\log(\tan(dx+c)^2+1)}{a^2+b^2} + \frac{2(Bab^2-Ab^3)\log(|b\tan(dx+c)+a|)}{a^3b+ab^3} + \frac{2A\log(|\tan(dx+c)|)}{a}}{2d}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `1/2*(2*(B*a - A*b)*(d*x + c)/(a^2 + b^2) - (A*a + B*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) + 2*(B*a*b^2 - A*b^3)*log(abs(b*tan(d*x + c) + a))/(a^3*b + a*b^3) + 2*A*log(abs(tan(d*x + c)))/a)/d`

3.271.9 Mupad [B] (verification not implemented)

Time = 9.40 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.44

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{A \ln(\tan(c+dx))}{ad} - \frac{\ln(\tan(c+dx) - i) (-B + A i)}{2d (-b + a i)} - \frac{\ln(\tan(c+dx) + i) (A - B i)}{2d (a - b i)} - \frac{b \ln(a + b \tan(c+dx)) (Ab - Ba)}{ad (a^2 + b^2)}$$

input `int((cot(c + d*x)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `(A*log(tan(c + d*x)))/(a*d) - (log(tan(c + d*x) - 1i)*(A*1i - B))/(2*d*(a*1i - b)) - (log(tan(c + d*x) + 1i)*(A - B*1i))/(2*d*(a - b*1i)) - (b*log(a + b*tan(c + d*x))*(A*b - B*a))/(a*d*(a^2 + b^2))`

3.271. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.272 $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.272.1 Optimal result 2625
 3.272.2 Mathematica [C] (verified) 2625
 3.272.3 Rubi [A] (verified) 2626
 3.272.4 Maple [A] (verified) 2628
 3.272.5 Fricas [A] (verification not implemented) 2629
 3.272.6 Sympy [C] (verification not implemented) 2629
 3.272.7 Maxima [A] (verification not implemented) 2630
 3.272.8 Giac [A] (verification not implemented) 2631
 3.272.9 Mupad [B] (verification not implemented) 2631

3.272.1 Optimal result

Integrand size = 31, antiderivative size = 103

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = -\frac{(aA+bB)x}{a^2+b^2} - \frac{A \cot(c+dx)}{ad} - \frac{(Ab-aB) \log(\sin(c+dx))}{a^2d} + \frac{b^2(Ab-aB) \log(a \cos(c+dx)+b \sin(c+dx))}{a^2(a^2+b^2)d}$$

```
output - (A*a+B*b)*x/(a^2+b^2)-A*cot(d*x+c)/a/d-(A*b-B*a)*ln(sin(d*x+c))/a^2/d+b^2
*(A*b-B*a)*ln(a*cos(d*x+c)+b*sin(d*x+c))/a^2/(a^2+b^2)/d
```

3.272.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.95 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.34

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{-\frac{2A \cot(c+dx)}{a} + \frac{i(A+iB) \log(i-\tan(c+dx))}{a+ib} + \frac{2(-Ab+aB) \log(\tan(c+dx))}{a^2} - \frac{(iA+B) \log(i+\tan(c+dx))}{a-ib} + \frac{2b^2(Ab-aB) \log(a+b \tan(c+dx))}{a^2(a^2+b^2)}}{2d}$$

input `Integrate[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `((-2*A*Cot[c + d*x])/a + (I*(A + I*B)*Log[I - Tan[c + d*x]])/(a + I*b) + (2*(-(A*b) + a*B)*Log[Tan[c + d*x]])/a^2 - ((I*A + B)*Log[I + Tan[c + d*x]])/(a - I*b) + (2*b^2*(A*b - a*B)*Log[a + b*Tan[c + d*x]])/(a^2*(a^2 + b^2)))/(2*d)`

3.272.3 Rubi [A] (verified)

Time = 0.68 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.08, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.258$, Rules used = {3042, 4092, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^2(a+b \tan(c+dx))} dx \\
 & \quad \downarrow \text{4092} \\
 & -\frac{\int \frac{\cot(c+dx)(Ab \tan^2(c+dx)+aA \tan(c+dx)+Ab-aB)}{a+b \tan(c+dx)} dx}{a} - \frac{A \cot(c+dx)}{ad} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\int \frac{Ab \tan(c+dx)^2+aA \tan(c+dx)+Ab-aB}{\tan(c+dx)(a+b \tan(c+dx))} dx}{a} - \frac{A \cot(c+dx)}{ad} \\
 & \quad \downarrow \text{4134} \\
 & -\frac{b^2(Ab-aB) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{(Ab-aB) \int \cot(c+dx) dx}{a} + \frac{ax(aA+bB)}{a^2+b^2} - \frac{A \cot(c+dx)}{ad} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{b^2(Ab-aB) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{(Ab-aB) \int -\tan(c+dx+\frac{\pi}{2}) dx}{a} + \frac{ax(aA+bB)}{a^2+b^2} - \frac{A \cot(c+dx)}{ad} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

3.272. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\begin{aligned}
& -\frac{b^2(Ab-aB) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{(Ab-aB) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx}{a} + \frac{ax(aA+bB)}{a^2+b^2} - \frac{A \cot(c+dx)}{ad} \\
& \quad \downarrow \text{3956} \\
& -\frac{b^2(Ab-aB) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{ax(aA+bB)}{a^2+b^2} + \frac{(Ab-aB) \log(-\sin(c+dx))}{ad} - \frac{A \cot(c+dx)}{ad} \\
& \quad \downarrow \text{4013} \\
& -\frac{b^2(Ab-aB) \log(a \cos(c+dx)+b \sin(c+dx))}{ad(a^2+b^2)} + \frac{ax(aA+bB)}{a^2+b^2} + \frac{(Ab-aB) \log(-\sin(c+dx))}{ad} - \frac{A \cot(c+dx)}{ad}
\end{aligned}$$

input `Int[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `-((A*Cot[c + d*x])/(a*d)) - ((a*(a*A + b*B)*x)/(a^2 + b^2) + ((A*b - a*B)*Log[-Sin[c + d*x]])/(a*d) - (b^2*(A*b - a*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d)/a`

3.272.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

```
rule 4092 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)
/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^
2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*
B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n +
2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n
+ 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1]
&& (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m]
|| (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4134 Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^
2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/
((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)
*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Sim
p[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*
x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x]
&& NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

3.272.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.18

method	result
parallelrisch	$\frac{(2Ab^3 - 2Ba^2b^2) \ln(a + b \tan(dx + c)) + (Aa^2b - Ba^3) \ln(\sec^2(dx + c)) - 2(a^2 + b^2)(Ab - Ba) \ln(\tan(dx + c)) - 2(A(a^2 + b^2) \tan(dx + c) - a^2d)}{2a^2d(a^2 + b^2)}$
derivativedivides	$\frac{\frac{(Ab - Ba) \ln(1 + \tan^2(dx + c))}{2} + (-aA - Bb) \arctan(\tan(dx + c))}{a^2 + b^2} - \frac{A}{a \tan(dx + c)} + \frac{(-Ab + Ba) \ln(\tan(dx + c))}{a^2} + \frac{(Ab - Ba)b^2 \ln(a + b \tan(dx + c))}{(a^2 + b^2)a^2}$
default	$\frac{\frac{(Ab - Ba) \ln(1 + \tan^2(dx + c))}{2} + (-aA - Bb) \arctan(\tan(dx + c))}{a^2 + b^2} - \frac{A}{a \tan(dx + c)} + \frac{(-Ab + Ba) \ln(\tan(dx + c))}{a^2} + \frac{(Ab - Ba)b^2 \ln(a + b \tan(dx + c))}{(a^2 + b^2)a^2}$
norman	$-\frac{A}{ad} - \frac{(aA + Bb)x \tan(dx + c)}{a^2 + b^2} + \frac{(Ab - Ba)b^2 \ln(a + b \tan(dx + c))}{a^2d(a^2 + b^2)} - \frac{(Ab - Ba) \ln(\tan(dx + c))}{a^2d} + \frac{(Ab - Ba) \ln(1 + \tan^2(dx + c))}{2d(a^2 + b^2)}$
risch	$-\frac{ixB}{ib - a} + \frac{xA}{ib - a} + \frac{2iAbx}{a^2} + \frac{2iAbc}{a^2d} - \frac{2ixB}{a} - \frac{2iBc}{ad} - \frac{2ib^3Ax}{a^2(a^2 + b^2)} - \frac{2ib^3Ac}{a^2d(a^2 + b^2)} + \frac{2ib^2Bx}{a(a^2 + b^2)} + \frac{2ib^2Bc}{ad(a^2 + b^2)}$

3.272.
$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{a + b \tan(c + dx)} dx$$

```
input int(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/2*((2*A*b^3-2*B*a*b^2)*ln(a+b*tan(d*x+c))+(A*a^2*b-B*a^3)*ln(sec(d*x+c)^2)-2*(a^2+b^2)*(A*b-B*a)*ln(tan(d*x+c))-2*(A*(a^2+b^2)*cot(d*x+c)+a*d*x*(A*a+B*b))*a)/a^2/d/(a^2+b^2)
```

3.272.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.72

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{2 A a^3 + 2 A a b^2 + 2 (A a^3 + B a^2 b) d x \tan(dx+c) - (B a^3 - A a^2 b + B a b^2 - A b^3) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)}{2 (a^4 + a^2 b^2) d \tan(dx+c)}$$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")
```

```
output -1/2*(2*A*a^3 + 2*A*a*b^2 + 2*(A*a^3 + B*a^2*b)*d*x*tan(d*x + c) - (B*a^3 - A*a^2*b + B*a*b^2 - A*b^3)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c) + (B*a*b^2 - A*b^3)*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1))*tan(d*x + c))/((a^4 + a^2*b^2)*d*tan(d*x + c))
```

3.272.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.00 (sec) , antiderivative size = 2071, normalized size of antiderivative = 20.11

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)**2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)
```

output `Piecewise((zoo*A*x, Eq(a, 0) & Eq(b, 0) & Eq(c, 0) & Eq(d, 0)), ((-A*x - A/(d*tan(c + d*x)) - B*log(tan(c + d*x)**2 + 1)/(2*d) + B*log(tan(c + d*x))/d)/a, Eq(b, 0)), ((A*log(tan(c + d*x)**2 + 1)/(2*d) - A*log(tan(c + d*x))/d - A/(2*d*tan(c + d*x)**2) - B*x - B/(d*tan(c + d*x)))/b, Eq(a, 0)), (-3*A*d*x*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - 3*I*A*d*x*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - I*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) + A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) + 2*I*A*log(tan(c + d*x))*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - 2*A*log(tan(c + d*x))*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - 3*A*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - 2*I*A/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) + I*B*d*x*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - B*d*x*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) + 2*B*log(tan(c + d*x))*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) + 2*I*B*log(tan(c + d*x))*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) + I*B*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x))...`

3.272.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.27

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{a + b \tan(c + dx)} dx =$$

$$\frac{\frac{2(Aa+Bb)(dx+c)}{a^2+b^2} + \frac{2(Bab^2-Ab^3)\log(b\tan(dx+c)+a)}{a^4+a^2b^2} + \frac{(Ba-Ab)\log(\tan(dx+c)^2+1)}{a^2+b^2} - \frac{2(Ba-Ab)\log(\tan(dx+c))}{a^2} + \frac{2A}{a\tan(dx+c)}}{2d}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(2*(A*a + B*b)*(d*x + c)/(a^2 + b^2) + 2*(B*a*b^2 - A*b^3)*log(b*tan(d*x + c) + a)/(a^4 + a^2*b^2) + (B*a - A*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) - 2*(B*a - A*b)*log(tan(d*x + c))/a^2 + 2*A/(a*tan(d*x + c)))/d`

3.272.8 Giac [A] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.52

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{\frac{2(Aa+Bb)(dx+c)}{a^2+b^2} + \frac{(Ba-Ab)\log(\tan(dx+c)^2+1)}{a^2+b^2} + \frac{2(Bab^3-Ab^4)\log(|b\tan(dx+c)+a|)}{a^4b+a^2b^3} - \frac{2(Ba-Ab)\log(|\tan(dx+c)|)}{a^2} + \frac{2(Ba\tan(c+dx))}{a^2}}{2d}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `-1/2*(2*(A*a + B*b)*(d*x + c)/(a^2 + b^2) + (B*a - A*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) + 2*(B*a*b^3 - A*b^4)*log(abs(b*tan(d*x + c) + a))/(a^4*b + a^2*b^3) - 2*(B*a - A*b)*log(abs(tan(d*x + c)))/a^2 + 2*(B*a*tan(d*x + c) - A*b*tan(d*x + c) + A*a)/(a^2*tan(d*x + c)))/d`

3.272.9 Mupad [B] (verification not implemented)

Time = 9.86 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.36

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{\ln(a+b\tan(c+dx))(Ab^3-BAb^2)}{d(a^4+a^2b^2)} - \frac{\ln(\tan(c+dx))(Ab-BA)}{a^2d} + \frac{\ln(\tan(c+dx)+1i)(A-B1i)}{2d(b+a1i)} - \frac{A\cot(c+dx)}{ad} + \frac{\ln(\tan(c+dx)-i)(-B+A1i)}{2d(a+b1i)}$$

input `int((cot(c + d*x)^2*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `(log(a + b*tan(c + d*x))*(A*b^3 - B*a*b^2))/(d*(a^4 + a^2*b^2)) - (log(tan(c + d*x))*(A*b - B*a))/(a^2*d) + (log(tan(c + d*x) + 1i)*(A - B*1i))/(2*d*(a*1i + b)) - (A*cot(c + d*x))/(a*d) + (log(tan(c + d*x) - 1i)*(A*1i - B))/(2*d*(a + b*1i))`

3.273 $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.273.1 Optimal result	2632
3.273.2 Mathematica [C] (verified)	2632
3.273.3 Rubi [A] (verified)	2633
3.273.4 Maple [A] (verified)	2636
3.273.5 Fricas [A] (verification not implemented)	2637
3.273.6 Sympy [C] (verification not implemented)	2637
3.273.7 Maxima [A] (verification not implemented)	2638
3.273.8 Giac [A] (verification not implemented)	2639
3.273.9 Mupad [B] (verification not implemented)	2639

3.273.1 Optimal result

Integrand size = 31, antiderivative size = 137

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{(Ab-aB)x}{a^2+b^2} + \frac{(Ab-aB) \cot(c+dx)}{a^2d} - \frac{A \cot^2(c+dx)}{2ad} - \frac{(a^2A-Ab^2+abB) \log(\sin(c+dx))}{a^3d} - \frac{b^3(Ab-aB) \log(a \cos(c+dx)+b \sin(c+dx))}{a^3(a^2+b^2)d}$$

output $(A*b-B*a)*x/(a^2+b^2)+(A*b-B*a)*\cot(d*x+c)/a^2/d-1/2*A*\cot(d*x+c)^2/a/d-(A*a^2-A*b^2+B*a*b)*\ln(\sin(d*x+c))/a^3/d-b^3*(A*b-B*a)*\ln(a*\cos(d*x+c)+b*\sin(d*x+c))/a^3/(a^2+b^2)/d$

3.273.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.49 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.19

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{2(Ab-aB) \cot(c+dx)}{a^2} - \frac{A \cot^2(c+dx)}{a} + \frac{(A+iB) \log(i-\tan(c+dx))}{a+ib} - \frac{2(a^2A-Ab^2+abB) \log(\tan(c+dx))}{a^3} + \frac{(A-iB) \log(i+\tan(c+dx))}{a-ib}$$

$2d$

input `Integrate[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output $((2*(A*b - a*B)*\text{Cot}[c + d*x])/a^2 - (A*\text{Cot}[c + d*x]^2)/a + ((A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]])/(a + I*b) - (2*(a^2*A - A*b^2 + a*b*B)*\text{Log}[\text{Tan}[c + d*x]])/a^3 + ((A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]])/(a - I*b) + (2*b^3*(-(A*b) + a*B)*\text{Log}[a + b*\text{Tan}[c + d*x]])/(a^3*(a^2 + b^2)))/(2*d)$

3.273.3 Rubi [A] (verified)

Time = 1.07 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.12, number of steps used = 12, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 4092, 27, 3042, 4132, 25, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B\tan(c+dx)}{\tan(c+dx)^3(a+b\tan(c+dx))} dx \\
 & \quad \downarrow \text{4092} \\
 & -\frac{\int \frac{2\cot^2(c+dx)(Ab\tan^2(c+dx)+aA\tan(c+dx)+Ab-aB)}{a+b\tan(c+dx)} dx}{2a} - \frac{A\cot^2(c+dx)}{2ad} \\
 & \quad \downarrow \text{27} \\
 & -\frac{\int \frac{\cot^2(c+dx)(Ab\tan^2(c+dx)+aA\tan(c+dx)+Ab-aB)}{a+b\tan(c+dx)} dx}{a} - \frac{A\cot^2(c+dx)}{2ad} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\int \frac{Ab\tan(c+dx)^2+aA\tan(c+dx)+Ab-aB}{\tan(c+dx)^2(a+b\tan(c+dx))} dx}{a} - \frac{A\cot^2(c+dx)}{2ad} \\
 & \quad \downarrow \text{4132} \\
 & -\frac{\int -\frac{\cot(c+dx)(Aa^2+B\tan(c+dx)a^2+bBa-Ab^2-b(Ab-aB)\tan^2(c+dx))}{a+b\tan(c+dx)} dx}{a} - \frac{(Ab-aB)\cot(c+dx)}{ad} - \frac{A\cot^2(c+dx)}{2ad} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

3.273. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$

$$\begin{aligned}
 & \frac{\int \frac{\cot(c+dx)(Aa^2+B \tan(c+dx)a^2+bBa-Ab^2-b(Ab-aB) \tan^2(c+dx))}{a+b \tan(c+dx)} dx}{a} - \frac{(Ab-aB) \cot(c+dx)}{ad} - \frac{A \cot^2(c+dx)}{2ad} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{Aa^2+B \tan(c+dx)a^2+bBa-Ab^2-b(Ab-aB) \tan(c+dx)^2}{\tan(c+dx)(a+b \tan(c+dx))} dx}{a} - \frac{(Ab-aB) \cot(c+dx)}{ad} - \frac{A \cot^2(c+dx)}{2ad} \\
 & \quad \downarrow \text{4134} \\
 & \frac{\frac{(a^2A+abB-Ab^2)}{a} \int \cot(c+dx) dx + \frac{b^3(Ab-aB) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^2x(Ab-aB)}{a^2+b^2}}{a} - \frac{(Ab-aB) \cot(c+dx)}{ad} - \frac{A \cot^2(c+dx)}{2ad} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{(a^2A+abB-Ab^2)}{a} \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx + \frac{b^3(Ab-aB) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^2x(Ab-aB)}{a^2+b^2}}{a} - \frac{(Ab-aB) \cot(c+dx)}{ad} \\
 & \quad \frac{A \cot^2(c+dx)}{2ad} \\
 & \quad \downarrow \text{25} \\
 & \frac{\frac{(a^2A+abB-Ab^2)}{a} \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx + \frac{b^3(Ab-aB) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^2x(Ab-aB)}{a^2+b^2}}{a} - \frac{(Ab-aB) \cot(c+dx)}{ad} \\
 & \quad \frac{A \cot^2(c+dx)}{2ad} \\
 & \quad \downarrow \text{3956} \\
 & \frac{\frac{b^3(Ab-aB) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{(a^2A+abB-Ab^2) \log(-\sin(c+dx))}{ad} - \frac{a^2x(Ab-aB)}{a^2+b^2}}{a} - \frac{(Ab-aB) \cot(c+dx)}{ad} \\
 & \quad \frac{A \cot^2(c+dx)}{2ad} \\
 & \quad \downarrow \text{4013} \\
 & \frac{\frac{(a^2A+abB-Ab^2) \log(-\sin(c+dx))}{ad} - \frac{a^2x(Ab-aB)}{a^2+b^2} + \frac{b^3(Ab-aB) \log(a \cos(c+dx)+b \sin(c+dx))}{ad(a^2+b^2)}}{a} - \frac{(Ab-aB) \cot(c+dx)}{ad} \\
 & \quad \frac{A \cot^2(c+dx)}{2ad}
 \end{aligned}$$

input `Int[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

3.273. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

output
$$-1/2*(A*\cot[c + d*x]^2)/(a*d) - (-(((A*b - a*B)*\cot[c + d*x])/(a*d)) + (-((a^2*(A*b - a*B)*x)/(a^2 + b^2)) + ((a^2*A - A*b^2 + a*b*B)*\log[-\sin[c + d*x]])/(a*d) + (b^3*(A*b - a*B)*\log[a*\cos[c + d*x] + b*\sin[c + d*x]])/(a*(a^2 + b^2)*d))/a/a$$

3.273.3.1 Defintions of rubi rules used

rule 25 $\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[F_x, x], x]$

rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3956 $\text{Int}[\tan[(c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\log[\text{RemoveContent}[\cos[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

rule 4013 $\text{Int}[(c_ + (d_)*\tan[(e_.) + (f_.)*(x_)])/(a_ + (b_)*\tan[(e_.) + (f_.)*(x_)]), x_Symbol] \rightarrow \text{Simp}[(c/(b*f))*\log[\text{RemoveContent}[a*\cos[e + f*x] + b*\sin[e + f*x], x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[a*c + b*d, 0]$

rule 4092 $\text{Int}[(a_ + (b_)*\tan[(e_.) + (f_.)*(x_)])^{(m_)}*((A_.) + (B_)*\tan[(e_.) + (f_.)*(x_)])^{(n_)}, x_Symbol] \rightarrow \text{Simp}[b*(A*b - a*B)*(a + b*\tan[e + f*x])^{(m+1)}*((c + d*\tan[e + f*x])^{(n+1)})/(f*(m+1)*(b*c - a*d)*(a^2 + b^2)), x] + \text{Simp}[1/((m+1)*(b*c - a*d)*(a^2 + b^2)) \text{ Int}[(a + b*\tan[e + f*x])^{(m+1)}*(c + d*\tan[e + f*x])^n * \text{Simp}[b*B*(b*c*(m+1) + a*d*(n+1)) + A*(a*(b*c - a*d)*(m+1) - b^2*d*(m+n+2)) - (A*b - a*B)*(b*c - a*d)*(m+1)*\tan[e + f*x] - b*d*(A*b - a*B)*(m+n+2)*\tan[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ (\text{IntegerQ}[m] \ || \ \text{IntegersQ}[2*m, 2*n]) \ \&\& \ !(\ \text{ILtQ}[n, -1] \ \&\& \ (\ !\text{IntegerQ}[m] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{NeQ}[a, 0])))$

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0]))
```

```
rule 4134 Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]*(x_))), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

3.273.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.11

method	result
derivativedivides	$\frac{(aA+Bb) \ln\left(\frac{1+\tan^2(dx+c)}{2}\right) + (Ab-Ba) \arctan(\tan(dx+c))}{a^2+b^2} - \frac{A}{2a \tan(dx+c)^2} - \frac{-Ab+Ba}{a^2 \tan(dx+c)} + \frac{(-A a^2 + A b^2 - Bab) \ln(\tan(dx+c))}{a^3 d}$
default	$\frac{(aA+Bb) \ln\left(\frac{1+\tan^2(dx+c)}{2}\right) + (Ab-Ba) \arctan(\tan(dx+c))}{a^2+b^2} - \frac{A}{2a \tan(dx+c)^2} - \frac{-Ab+Ba}{a^2 \tan(dx+c)} + \frac{(-A a^2 + A b^2 - Bab) \ln(\tan(dx+c))}{a^3 d}$
parallelrisch	$\frac{(-2A b^4 + 2B a b^3) \ln(a+b \tan(dx+c)) + (A a^4 + B a^3 b) \ln(\sec^2(dx+c)) + (-2A a^4 + 2A b^4 - 2B a^3 b - 2B a b^3) \ln(\tan(dx+c))}{2(a^2+b^2)a^3 d}$
norman	$\frac{(Ab-Ba) \tan(dx+c)}{a^2 d} + \frac{(Ab-Ba)x \tan^2(dx+c)}{a^2+b^2} - \frac{A}{2ad} + \frac{(aA+Bb) \ln(1+\tan^2(dx+c))}{2d(a^2+b^2)} - \frac{(A a^2 - A b^2 + Bab) \ln(\tan(dx+c))}{a^3 d}$
risch	$\frac{x B}{i b - a} - \frac{2i A b^2 x}{a^3} + \frac{2i A c}{a d} + \frac{2i B b c}{a^2 d} - \frac{2i b^3 B c}{a^2 d (a^2 + b^2)} - \frac{2i (i A a e^{2i(dx+c)} - A b e^{2i(dx+c)} + B a e^{2i(dx+c)} + A b - B a)}{a^2 d (e^{2i(dx+c)} - 1)^2}$

3.273.
$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

```
input int(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/d*(1/(a^2+b^2)*(1/2*(A*a+B*b)*ln(1+tan(d*x+c)^2)+(A*b-B*a)*arctan(tan(d*x+c)))-1/2/a*A/tan(d*x+c)^2-(-A*b+B*a)/a^2/tan(d*x+c)+1/a^3*(-A*a^2+A*b^2-B*a*b)*ln(tan(d*x+c))-(A*b-B*a)*b^3/(a^2+b^2)/a^3*ln(a+b*tan(d*x+c)))
```

3.273.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.71

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{Aa^4 + Aa^2b^2 + (Aa^4 + Ba^3b + Bab^3 - Ab^4) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^2 - (Bab^3 - Ab^4) \log\left(\frac{b^2 \tan(dx+c)^2}{\tan(dx+c)^2+1}\right)}{a^2+b^2}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")
```

```
output -1/2*(A*a^4 + A*a^2*b^2 + (A*a^4 + B*a^3*b + B*a*b^3 - A*b^4)*log(tan(d*x+c)^2/(tan(d*x+c)^2+1))*tan(d*x+c)^2 - (B*a*b^3 - A*b^4)*log((b^2*tan(d*x+c)^2+2*a*b*tan(d*x+c)+a^2)/(tan(d*x+c)^2+1))*tan(d*x+c)^2 + (A*a^4 + A*a^2*b^2 + 2*(B*a^4 - A*a^3*b)*d*x)*tan(d*x+c)^2 + 2*(B*a^4 - A*a^3*b + B*a^2*b^2 - A*a*b^3)*tan(d*x+c))/((a^5 + a^3*b^2)*d*tan(d*x+c)^2)
```

3.273.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.85 (sec) , antiderivative size = 2599, normalized size of antiderivative = 18.97

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)**3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)
```

output `Piecewise((zoo*A*x, Eq(a, 0) & Eq(b, 0) & Eq(c, 0) & Eq(d, 0)), ((A*log(tan(c + d*x)**2 + 1)/(2*d) - A*log(tan(c + d*x))/d - A/(2*d*tan(c + d*x)**2) - B*x - B/(d*tan(c + d*x)))/a, Eq(b, 0)), ((A*x + A/(d*tan(c + d*x)) - A/(3*d*tan(c + d*x)**3) + B*log(tan(c + d*x)**2 + 1)/(2*d) - B*log(tan(c + d*x))/d - B/(2*d*tan(c + d*x)**2))/b, Eq(a, 0)), (-3*I*A*d*x*tan(c + d*x)**3/(2*a*d*tan(c + d*x)**3 + 2*I*a*d*tan(c + d*x)**2) + 3*A*d*x*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**3 + 2*I*a*d*tan(c + d*x)**2) + 2*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**3/(2*a*d*tan(c + d*x)**3 + 2*I*a*d*tan(c + d*x)**2) + 2*I*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**3 + 2*I*a*d*tan(c + d*x)**2) - 4*A*log(tan(c + d*x))*tan(c + d*x)**3/(2*a*d*tan(c + d*x)**3 + 2*I*a*d*tan(c + d*x)**2) - 4*I*A*log(tan(c + d*x))*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**3 + 2*I*a*d*tan(c + d*x)**2) - 3*I*A*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**3 + 2*I*a*d*tan(c + d*x)**2) + A*tan(c + d*x)/(2*a*d*tan(c + d*x)**3 + 2*I*a*d*tan(c + d*x)**2) - I*A/(2*a*d*tan(c + d*x)**3 + 2*I*a*d*tan(c + d*x)**2) - 3*B*d*x*tan(c + d*x)**3/(2*a*d*tan(c + d*x)**3 + 2*I*a*d*tan(c + d*x)**2) - 3*I*B*d*x*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**3 + 2*I*a*d*tan(c + d*x)**2) - I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**3/(2*a*d*tan(c + d*x)**3 + 2*I*a*d*tan(c + d*x)**2) + B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**3 + 2*I*a*d*tan(c + d*x)**2) + 2*I*B*log(tan(c + d*x))*tan(c + d*x)**3/(2*a*d*tan(c + d...`

3.273.7 Maxima [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.15

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx =$$

$$\frac{\frac{2(Ba-Ab)(dx+c)}{a^2+b^2} - \frac{2(Bab^3-Ab^4) \log(b \tan(dx+c)+a)}{a^5+a^3b^2} - \frac{(Aa+Bb) \log(\tan(dx+c)^2+1)}{a^2+b^2} + \frac{2(Aa^2+Bab-Ab^2) \log(\tan(dx+c))}{a^3} + \frac{Aa}{2d}}$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(2*(B*a - A*b)*(d*x + c)/(a^2 + b^2) - 2*(B*a*b^3 - A*b^4)*log(b*tan(d*x + c) + a)/(a^5 + a^3*b^2) - (A*a + B*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) + 2*(A*a^2 + B*a*b - A*b^2)*log(tan(d*x + c))/a^3 + (A*a + 2*(B*a - A*b)*tan(d*x + c))/(a^2*tan(d*x + c)^2))/d`

3.273.8 Giac [A] (verification not implemented)

Time = 0.70 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.56

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{\frac{2(Ba-Ab)(dx+c)}{a^2+b^2} - \frac{(Aa+Bb)\log(\tan(dx+c)^2+1)}{a^2+b^2} - \frac{2(Bab^4-Ab^5)\log(|b\tan(dx+c)+a|)}{a^5b+a^3b^3} + \frac{2(Aa^2+Bab-Ab^2)\log(|\tan(dx+c)|)}{a^3}}{2d}$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="gias")`

output `-1/2*(2*(B*a - A*b)*(d*x + c)/(a^2 + b^2) - (A*a + B*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) - 2*(B*a*b^4 - A*b^5)*log(abs(b*tan(d*x + c) + a))/(a^5*b + a^3*b^3) + 2*(A*a^2 + B*a*b - A*b^2)*log(abs(tan(d*x + c)))/a^3 - (3*A*a^2*tan(d*x + c)^2 + 3*B*a*b*tan(d*x + c)^2 - 3*A*b^2*tan(d*x + c)^2 - 2*B*a^2*tan(d*x + c) + 2*A*a*b*tan(d*x + c) - A*a^2)/(a^3*tan(d*x + c)^2))/d`

3.273.9 Mupad [B] (verification not implemented)

Time = 10.90 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.28

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = -\frac{\cot(c+dx)^2 \left(\frac{A}{2a} - \frac{\tan(c+dx)(Ab-Ba)}{a^2} \right)}{d} + \frac{\ln(\tan(c+dx) - i)(-B + A i)}{2d(-b + a i)} - \frac{\ln(\tan(c+dx))(Aa^2 + Bab - Ab^2)}{a^3 d} - \frac{\ln(a + b\tan(c+dx))(Ab^4 - Bab^3)}{d(a^5 + a^3b^2)} + \frac{\ln(\tan(c+dx) + i)(A - B i)}{2d(a - b i)}$$

input `int((cot(c + d*x)^3*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `(log(tan(c + d*x) - 1i)*(A*1i - B))/(2*d*(a*1i - b)) - (cot(c + d*x)^2*(A/(2*a) - (tan(c + d*x)*(A*b - B*a))/a^2))/d - (log(tan(c + d*x))*(A*a^2 - A*b^2 + B*a*b))/(a^3*d) - (log(a + b*tan(c + d*x))*(A*b^4 - B*a*b^3))/(d*(a^5 + a^3*b^2)) + (log(tan(c + d*x) + 1i)*(A - B*1i))/(2*d*(a - b*1i))`

3.274 $\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.274.1 Optimal result	2640
3.274.2 Mathematica [C] (verified)	2640
3.274.3 Rubi [A] (verified)	2641
3.274.4 Maple [A] (verified)	2645
3.274.5 Fricas [A] (verification not implemented)	2646
3.274.6 Sympy [C] (verification not implemented)	2646
3.274.7 Maxima [A] (verification not implemented)	2647
3.274.8 Giac [A] (verification not implemented)	2648
3.274.9 Mupad [B] (verification not implemented)	2648

3.274.1 Optimal result

Integrand size = 31, antiderivative size = 169

$$\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{(aA+bB)x}{a^2+b^2} + \frac{(a^2A-Ab^2+abB) \cot(c+dx)}{a^3d} + \frac{(Ab-aB) \cot^2(c+dx)}{2a^2d} - \frac{A \cot^3(c+dx)}{3ad} + \frac{(a^2-b^2)(Ab-aB) \log(\sin(c+dx))}{a^4d} + \frac{b^4(Ab-aB) \log(a \cos(c+dx)+b \sin(c+dx))}{a^4(a^2+b^2)d}$$

output $(A*a+B*b)*x/(a^2+b^2)+(A*a^2-A*b^2+B*a*b)*\cot(d*x+c)/a^3/d+1/2*(A*b-B*a)*\cot(d*x+c)^2/a^2/d-1/3*A*\cot(d*x+c)^3/a/d+(a^2-b^2)*(A*b-B*a)*\ln(\sin(d*x+c))/a^4/d+b^4*(A*b-B*a)*\ln(a*\cos(d*x+c)+b*\sin(d*x+c))/a^4/(a^2+b^2)/d$

3.274.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.95 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.15

$$\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{6(a^2A-Ab^2+abB) \cot(c+dx)}{a^3} + \frac{3(Ab-aB) \cot^2(c+dx)}{a^2} - \frac{2A \cot^3(c+dx)}{a} + \frac{3(-iA+B) \log(i-\tan(c+dx))}{a+ib} + \frac{6(a-b)(a+b)(Ab-aB) \log(\tan(c+dx))}{a^4} + \frac{6d}{6d}$$

input `Integrate[(Cot[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output $((6*(a^2*A - A*b^2 + a*b*B)*\text{Cot}[c + d*x])/a^3 + (3*(A*b - a*B)*\text{Cot}[c + d*x]^2)/a^2 - (2*A*\text{Cot}[c + d*x]^3)/a + (3*((-I)*A + B)*\text{Log}[I - \text{Tan}[c + d*x]])/(a + I*b) + (6*(a - b)*(a + b)*(A*b - a*B)*\text{Log}[\text{Tan}[c + d*x]])/a^4 + (3*(I*A + B)*\text{Log}[I + \text{Tan}[c + d*x]])/(a - I*b) + (6*b^4*(A*b - a*B)*\text{Log}[a + b*\text{Tan}[c + d*x]])/(a^4*(a^2 + b^2)))/(6*d)$

3.274.3 Rubi [A] (verified)

Time = 1.45 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.14, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx \\ & \quad \downarrow 3042 \\ & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^4(a+b \tan(c+dx))} dx \\ & \quad \downarrow 4092 \\ & \frac{\int \frac{3 \cot^3(c+dx)(Ab \tan^2(c+dx)+aA \tan(c+dx)+Ab-aB)}{a+b \tan(c+dx)} dx}{3a} - \frac{A \cot^3(c+dx)}{3ad} \\ & \quad \downarrow 27 \\ & \frac{\int \frac{\cot^3(c+dx)(Ab \tan^2(c+dx)+aA \tan(c+dx)+Ab-aB)}{a+b \tan(c+dx)} dx}{a} - \frac{A \cot^3(c+dx)}{3ad} \\ & \quad \downarrow 3042 \\ & \frac{\int \frac{Ab \tan(c+dx)^2+aA \tan(c+dx)+Ab-aB}{\tan(c+dx)^3(a+b \tan(c+dx))} dx}{a} - \frac{A \cot^3(c+dx)}{3ad} \\ & \quad \downarrow 4132 \\ & \frac{\int -\frac{2 \cot^2(c+dx)(Aa^2+B \tan(c+dx)a^2+bBa-Ab^2-b(Ab-aB) \tan^2(c+dx))}{a+b \tan(c+dx)} dx}{2a} - \frac{(Ab-aB) \cot^2(c+dx)}{2ad} - \frac{A \cot^3(c+dx)}{3ad} \end{aligned}$$

3.274. $\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\begin{array}{c}
 \int \frac{\cot^2(c+dx)(Aa^2+B \tan(c+dx)a^2+bBa-Ab^2-b(Ab-aB) \tan^2(c+dx))}{a+b \tan(c+dx)} dx - \frac{(Ab-aB) \cot^2(c+dx)}{2ad} - \frac{A \cot^3(c+dx)}{3ad} \\
 \downarrow 27 \\
 \int \frac{Aa^2+B \tan(c+dx)a^2+bBa-Ab^2-b(Ab-aB) \tan(c+dx)^2}{\tan(c+dx)^2(a+b \tan(c+dx))} dx - \frac{(Ab-aB) \cot^2(c+dx)}{2ad} - \frac{A \cot^3(c+dx)}{3ad} \\
 \downarrow 3042 \\
 \int \frac{\cot(c+dx)(A \tan(c+dx)a^3+b(Aa^2+bBa-Ab^2) \tan^2(c+dx)+(a^2-b^2)(Ab-aB))}{a+b \tan(c+dx)} dx - \frac{(a^2A+abB-Ab^2) \cot(c+dx)}{ad} - \frac{(Ab-aB) \cot^2(c+dx)}{2ad} \\
 \downarrow 4132 \\
 \frac{A \cot^3(c+dx)}{3ad} \\
 \downarrow 3042 \\
 \int \frac{A \tan(c+dx)a^3+b(Aa^2+bBa-Ab^2) \tan(c+dx)^2+(a^2-b^2)(Ab-aB)}{\tan(c+dx)(a+b \tan(c+dx))} dx - \frac{(a^2A+abB-Ab^2) \cot(c+dx)}{ad} - \frac{(Ab-aB) \cot^2(c+dx)}{2ad} \\
 \downarrow 4134 \\
 \frac{A \cot^3(c+dx)}{3ad} \\
 \downarrow 3042 \\
 \frac{(a^2-b^2)(Ab-aB) \int \cot(c+dx) dx}{a} + \frac{b^4(Ab-aB) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^3x(aA+bB)}{a^2+b^2} - \frac{(a^2A+abB-Ab^2) \cot(c+dx)}{ad} - \frac{(Ab-aB) \cot^2(c+dx)}{2ad} \\
 \downarrow 3042 \\
 \frac{(a^2-b^2)(Ab-aB) \int -\tan(c+dx+\frac{\pi}{2}) dx}{a} + \frac{b^4(Ab-aB) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^3x(aA+bB)}{a^2+b^2} - \frac{(a^2A+abB-Ab^2) \cot(c+dx)}{ad} - \frac{(Ab-aB) \cot^2(c+dx)}{2ad} \\
 \downarrow 25 \\
 \frac{A \cot^3(c+dx)}{3ad}
 \end{array}$$

3.274. $\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\begin{aligned}
 & \frac{\frac{(a^2-b^2)(Ab-aB) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx}{a} + \frac{b^4(Ab-aB) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^3x(aA+bB)}{a^2+b^2}}{a} - \frac{(a^2A+abB-Ab^2) \cot(c+dx)}{ad} - \frac{(Ab-aB) \cot^2(c+dx)}{2ad} \\
 & \frac{A \cot^3(c+dx)}{3ad} \\
 & \quad \downarrow \text{3956} \\
 & \frac{\frac{b^4(Ab-aB) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{(a^2-b^2)(Ab-aB) \log(-\sin(c+dx))}{ad} + \frac{a^3x(aA+bB)}{a^2+b^2}}{a} - \frac{(a^2A+abB-Ab^2) \cot(c+dx)}{ad} - \frac{(Ab-aB) \cot^2(c+dx)}{2ad} \\
 & \frac{A \cot^3(c+dx)}{3ad} \\
 & \quad \downarrow \text{4013} \\
 & \frac{\frac{(a^2-b^2)(Ab-aB) \log(-\sin(c+dx))}{ad} + \frac{b^4(Ab-aB) \log(a \cos(c+dx)+b \sin(c+dx))}{ad(a^2+b^2)} + \frac{a^3x(aA+bB)}{a^2+b^2}}{a} - \frac{(a^2A+abB-Ab^2) \cot(c+dx)}{ad} - \frac{(Ab-aB) \cot^2(c+dx)}{2ad} \\
 & \frac{A \cot^3(c+dx)}{3ad}
 \end{aligned}$$

```
input Int[(Cot[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]
```

```
output -1/3*(A*Cot[c + d*x]^3)/(a*d) - (-1/2*((A*b - a*B)*Cot[c + d*x]^2)/(a*d) +
(-(((a^2*A - A*b^2 + a*b*B)*Cot[c + d*x]))/(a*d)) - ((a^3*(a*A + b*B)*x)/(
a^2 + b^2) + ((a^2 - b^2)*(A*b - a*B)*Log[-Sin[c + d*x]])/(a*d) + (b^4*(A*
b - a*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))/a/a/a
```

3.274.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4134 Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^
2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/
((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)
*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Sim
p[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*
x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x]
&& NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

3.274.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.12

method	result
derivativedivides	$\frac{(-Ab+Ba)\ln\left(\frac{1+\tan^2(dx+c)}{2}\right) + (aA+Bb)\arctan(\tan(dx+c))}{a^2+b^2} - \frac{-Ab+Ba}{2a^2\tan(dx+c)^2} - \frac{-Aa^2+Ab^2-Bab}{a^3\tan(dx+c)} + \frac{(Aa^2b-Ab^3-Ba^3+Ba^2b^2)}{a^4}$
default	$\frac{(-Ab+Ba)\ln\left(\frac{1+\tan^2(dx+c)}{2}\right) + (aA+Bb)\arctan(\tan(dx+c))}{a^2+b^2} - \frac{-Ab+Ba}{2a^2\tan(dx+c)^2} - \frac{-Aa^2+Ab^2-Bab}{a^3\tan(dx+c)} + \frac{(Aa^2b-Ab^3-Ba^3+Ba^2b^2)}{a^4}$
parallelrisch	$(6Ab^5-6Bab^4)\ln(a+b\tan(dx+c)) + (-3Aa^4b+3Ba^5)\ln(\sec^2(dx+c)) + (6Aa^4b-6Ab^5-6Ba^5+6Bab^4)\ln(\tan(dx+c))$
norman	$\frac{(aA+Bb)x(\tan^3(dx+c))}{a^2+b^2} + \frac{(Aa^2-Ab^2+Bab)(\tan^2(dx+c))}{a^3d} - \frac{A}{3ad} + \frac{(Ab-Ba)\tan(dx+c)}{2a^2d} + \frac{(Ab-Ba)(a^2-b^2)\ln(\tan(dx+c))}{a^4d}$
risch	$-\frac{2iBb^2x}{a^3} - \frac{xA}{ib-a} + \frac{2ixB}{a} - \frac{2iBb^2c}{a^3d} - \frac{2ib^5Ac}{(a^2+b^2)a^4d} + \frac{2ib^4Bx}{(a^2+b^2)a^3} + \frac{2iBc}{ad} - \frac{2i(-3iAabe^{4i(dx+c)}+3iBa^2e^{4i(dx+c)})}{a^4d}$

```
input int(cot(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/d*(1/(a^2+b^2)*(1/2*(-A*b+B*a)*ln(1+tan(d*x+c)^2)+(A*a+B*b)*arctan(tan(d
*x+c)))-1/2*(-A*b+B*a)/a^2/tan(d*x+c)^2-(-A*a^2+A*b^2-B*a*b)/a^3/tan(d*x+c
)+1/a^4*(A*a^2*b-A*b^3-B*a^3+B*a*b^2)*ln(tan(d*x+c))-1/3/a*A/tan(d*x+c)^3+
(A*b-B*a)*b^4/(a^2+b^2)/a^4*ln(a+b*tan(d*x+c)))
```

$$3.274. \int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.274.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 293, normalized size of antiderivative = 1.73

$$\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx =$$

$$\frac{2Aa^5 + 2Aa^3b^2 + 3(Ba^5 - Aa^4b - Bab^4 + Ab^5) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2+1}\right) \tan(dx+c)^3 + 3(Bab^4 - Ab^5) \log}{-}$$

input `integrate(cot(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output `-1/6*(2*A*a^5 + 2*A*a^3*b^2 + 3*(B*a^5 - A*a^4*b - B*a*b^4 + A*b^5)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1))*tan(d*x + c)^3 + 3*(B*a*b^4 - A*b^5)*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1))*tan(d*x + c)^3 + 3*(B*a^5 - A*a^4*b + B*a^3*b^2 - A*a^2*b^3 - 2*(A*a^5 + B*a^4*b)*d*x)*tan(d*x + c)^3 - 6*(A*a^5 + B*a^4*b + B*a^2*b^3 - A*a*b^4)*tan(d*x + c)^2 + 3*(B*a^5 - A*a^4*b + B*a^3*b^2 - A*a^2*b^3)*tan(d*x + c))/(a^6 + a^4*b^2)*d*tan(d*x + c)^3`

3.274.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 4.20 (sec) , antiderivative size = 3016, normalized size of antiderivative = 17.85

$$\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)**4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Piecewise((zoo*A*x, Eq(a, 0) & Eq(b, 0) & Eq(c, 0) & Eq(d, 0)), ((A*x + A/(d*tan(c + d*x)) - A/(3*d*tan(c + d*x)**3) + B*log(tan(c + d*x)**2 + 1)/(2*d) - B*log(tan(c + d*x))/d - B/(2*d*tan(c + d*x)**2))/a, Eq(b, 0)), ((-A*log(tan(c + d*x)**2 + 1)/(2*d) + A*log(tan(c + d*x))/d + A/(2*d*tan(c + d*x)**2) - A/(4*d*tan(c + d*x)**4) + B*x + B/(d*tan(c + d*x)) - B/(3*d*tan(c + d*x)**3))/b, Eq(a, 0)), (15*A*d*x*tan(c + d*x)**4/(6*a*d*tan(c + d*x)**4 + 6*I*a*d*tan(c + d*x)**3) + 15*I*A*d*x*tan(c + d*x)**3/(6*a*d*tan(c + d*x)**4 + 6*I*a*d*tan(c + d*x)**3) + 6*I*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**4/(6*a*d*tan(c + d*x)**4 + 6*I*a*d*tan(c + d*x)**3) - 6*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**3/(6*a*d*tan(c + d*x)**4 + 6*I*a*d*tan(c + d*x)**3) - 12*I*A*log(tan(c + d*x))*tan(c + d*x)**4/(6*a*d*tan(c + d*x)**4 + 6*I*a*d*tan(c + d*x)**3) + 12*A*log(tan(c + d*x))*tan(c + d*x)**3/(6*a*d*tan(c + d*x)**4 + 6*I*a*d*tan(c + d*x)**3) + 15*A*tan(c + d*x)**3/(6*a*d*tan(c + d*x)**4 + 6*I*a*d*tan(c + d*x)**3) + 9*I*A*tan(c + d*x)**2/(6*a*d*tan(c + d*x)**4 + 6*I*a*d*tan(c + d*x)**3) + A*tan(c + d*x)/(6*a*d*tan(c + d*x)**4 + 6*I*a*d*tan(c + d*x)**3) - 2*I*A/(6*a*d*tan(c + d*x)**4 + 6*I*a*d*tan(c + d*x)**3) - 9*I*B*d*x*tan(c + d*x)**4/(6*a*d*tan(c + d*x)**4 + 6*I*a*d*tan(c + d*x)**3) + 9*B*d*x*tan(c + d*x)**3/(6*a*d*tan(c + d*x)**4 + 6*I*a*d*tan(c + d*x)**3) + 6*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**4/(6*a*d*tan(c + d*x)**4 + 6*I*a*d*tan(c + d*x)**3) + 6*I*B*log(tan(c + d...`

3.274.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.18

$$\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$= \frac{\frac{6(Aa+Bb)(dx+c)}{a^2+b^2} - \frac{6(Bab^4-Ab^5) \log(b \tan(dx+c)+a)}{a^6+a^4b^2} + \frac{3(Ba-Ab) \log(\tan(dx+c)^2+1)}{a^2+b^2} - \frac{6(Ba^3-Aa^2b-Bab^2+Ab^3) \log(\tan(dx+c))}{a^4}}{6d}$$

input `integrate(cot(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/6*(6*(A*a + B*b)*(d*x + c)/(a^2 + b^2) - 6*(B*a*b^4 - A*b^5)*log(b*tan(d*x + c) + a)/(a^6 + a^4*b^2) + 3*(B*a - A*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) - 6*(B*a^3 - A*a^2*b - B*a*b^2 + A*b^3)*log(tan(d*x + c))/a^4 - (2*A*a^2 - 6*(A*a^2 + B*a*b - A*b^2)*tan(d*x + c)^2 + 3*(B*a^2 - A*a*b)*tan(d*x + c))/(a^3*tan(d*x + c)^3))/d`

3.274. $\int \frac{\cot^4(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.274.8 Giac [A] (verification not implemented)

Time = 0.87 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.69

$$\int \frac{\cot^4(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{6(Aa+Bb)(dx+c)}{a^2+b^2} + \frac{3(Ba-Ab)\log(\tan(dx+c)^2+1)}{a^2+b^2} - \frac{6(Bab^5-Ab^6)\log(|b\tan(dx+c)+a|)}{a^6b+a^4b^3} - \frac{6(Ba^3-Aa^2b- Bab^2+Ab^3)\log(|\tan(dx+c)|)}{a^4}$$

```
input integrate(cot(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")
```

```
output 1/6*(6*(A*a + B*b)*(d*x + c)/(a^2 + b^2) + 3*(B*a - A*b)*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) - 6*(B*a*b^5 - A*b^6)*log(abs(b*tan(d*x + c) + a))/(a^6 * b + a^4*b^3) - 6*(B*a^3 - A*a^2*b - B*a*b^2 + A*b^3)*log(abs(tan(d*x + c)))/a^4 + (11*B*a^3*tan(d*x + c)^3 - 11*A*a^2*b*tan(d*x + c)^3 - 11*B*a*b^2*tan(d*x + c)^3 + 11*A*b^3*tan(d*x + c)^3 + 6*A*a^3*tan(d*x + c)^2 + 6*B*a^2*b*tan(d*x + c)^2 - 6*A*a*b^2*tan(d*x + c)^2 - 3*B*a^3*tan(d*x + c) + 3*A*a^2*b*tan(d*x + c) - 2*A*a^3)/(a^4*tan(d*x + c)^3))/d
```

3.274.9 Mupad [B] (verification not implemented)

Time = 10.06 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.23

$$\int \frac{\cot^4(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{\cot(c+dx)^3 \left(\frac{\tan(c+dx)^2 (Aa^2+Bab-Ab^2)}{a^3} - \frac{A}{3a} + \frac{\tan(c+dx)(Ab-Ba)}{2a^2} \right)}{d} + \frac{\ln(a+b\tan(c+dx))(Ab^5-Bab^4)}{d(a^6+a^4b^2)} - \frac{\ln(\tan(c+dx))(Ba^3-Aa^2b-Bab^2+Ab^3)}{a^4d} - \frac{\ln(\tan(c+dx)+1i)(A-B1i)}{2d(b+a1i)} - \frac{\ln(\tan(c+dx)-1i)(-B+A1i)}{2d(a+b1i)}$$

```
input int((cot(c + d*x)^4*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)),x)
```

output $(\cot(c + d*x))^3 * ((\tan(c + d*x))^2 * (A*a^2 - A*b^2 + B*a*b)) / a^3 - A / (3*a) +$
 $(\tan(c + d*x) * (A*b - B*a)) / (2*a^2)) / d + (\log(a + b*\tan(c + d*x)) * (A*b^5 -$
 $B*a*b^4)) / (d*(a^6 + a^4*b^2)) - (\log(\tan(c + d*x)) * (A*b^3 + B*a^3 - A*a^2$
 $*b - B*a*b^2)) / (a^4*d) - (\log(\tan(c + d*x) + 1i) * (A - B*1i)) / (2*d*(a*1i +$
 $b)) - (\log(\tan(c + d*x) - 1i) * (A*1i - B)) / (2*d*(a + b*1i))$

3.275 $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.275.1 Optimal result 2650
 3.275.2 Mathematica [C] (verified) 2651
 3.275.3 Rubi [A] (verified) 2651
 3.275.4 Maple [A] (verified) 2655
 3.275.5 Fricas [B] (verification not implemented) 2656
 3.275.6 Sympy [C] (verification not implemented) 2656
 3.275.7 Maxima [A] (verification not implemented) 2657
 3.275.8 Giac [A] (verification not implemented) 2658
 3.275.9 Mupad [B] (verification not implemented) 2658

3.275.1 Optimal result

Integrand size = 31, antiderivative size = 208

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= -\frac{(2aAb - a^2B + b^2B)x}{(a^2 + b^2)^2} + \frac{(a^2A - Ab^2 + 2abB) \log(\cos(c+dx))}{(a^2 + b^2)^2 d}$$

$$+ \frac{a^2(a^2Ab + 3Ab^3 - 2a^3B - 4ab^2B) \log(a+b \tan(c+dx))}{b^3(a^2 + b^2)^2 d}$$

$$- \frac{(aAb - 2a^2B - b^2B) \tan(c+dx)}{b^2(a^2 + b^2) d} + \frac{a(Ab - aB) \tan^2(c+dx)}{b(a^2 + b^2) d(a+b \tan(c+dx))}$$

output

```
-(2*A*a*b-B*a^2+B*b^2)*x/(a^2+b^2)^2+(A*a^2-A*b^2+2*B*a*b)*ln(cos(d*x+c))/
(a^2+b^2)^2/d+a^2*(A*a^2*b+3*A*b^3-2*B*a^3-4*B*a*b^2)*ln(a+b*tan(d*x+c))/b
^3/(a^2+b^2)^2/d-(A*a*b-2*B*a^2-B*b^2)*tan(d*x+c)/b^2/(a^2+b^2)/d+a*(A*b-B
*a)*tan(d*x+c)^2/b/(a^2+b^2)/d/(a+b*tan(d*x+c))
```

3.275.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.00 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.93

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \frac{\frac{(A+iB) \log(i-\tan(c+dx))}{(a+ib)^2} + \frac{(A-iB) \log(i+\tan(c+dx))}{(a-ib)^2} + \frac{2a^2(-a^2Ab-3Ab^3+2a^3B+4ab^2B) \log(a+b \tan(c+dx))}{b^3(a^2+b^2)^2} + \frac{2a^2(-aAb+2a^2B)}{b^3(a^2+b^2)(a+b \tan(c+dx))}}{2d}$$

```
input Integrate[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]
```

```
output -1/2*(((A + I*B)*Log[I - Tan[c + d*x]])/(a + I*b)^2 + ((A - I*B)*Log[I + Tan[c + d*x]])/(a - I*b)^2 + (2*a^2*(-(a^2*A*b) - 3*A*b^3 + 2*a^3*B + 4*a*b^2*B)*Log[a + b*Tan[c + d*x]])/(b^3*(a^2 + b^2)^2) + (2*a^2*(-(a*A*b) + 2*a^2*B + b^2*B))/(b^3*(a^2 + b^2)*(a + b*Tan[c + d*x])) - (2*B*Tan[c + d*x]^2)/(b*(a + b*Tan[c + d*x])))/d
```

3.275.3 Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.08, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 4088, 25, 3042, 4130, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(c+dx)^3(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\ & \quad \downarrow \text{4088} \\ & \frac{\int -\frac{\tan(c+dx)((-2Ba^2+Aba-b^2B) \tan^2(c+dx)-b(Ab-aB) \tan(c+dx)+2a(Ab-aB))}{a+b \tan(c+dx)} dx}{b(a^2+b^2)} + \\ & \quad \frac{a(Ab-aB) \tan^2(c+dx)}{bd(a^2+b^2)(a+b \tan(c+dx))} \end{aligned}$$

$$\begin{aligned}
 & \downarrow 25 \\
 & \frac{a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \\
 & \frac{\int \frac{\tan(c+dx)((-2Ba^2 + Aba - b^2B) \tan^2(c+dx) - b(Ab - aB) \tan(c+dx) + 2a(Ab - aB))}{a + b \tan(c+dx)} dx}{b(a^2 + b^2)} \\
 & \downarrow 3042 \\
 & \frac{a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \\
 & \frac{\int \frac{\tan(c+dx)((-2Ba^2 + Aba - b^2B) \tan^2(c+dx)^2 - b(Ab - aB) \tan(c+dx) + 2a(Ab - aB))}{a + b \tan(c+dx)} dx}{b(a^2 + b^2)} \\
 & \downarrow 4130 \\
 & \frac{a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \\
 & \frac{\int -\frac{((aA + bB) \tan(c+dx)b^2) + (a^2 + b^2)(Ab - 2aB) \tan^2(c+dx) + a(-2Ba^2 + Aba - b^2B)}{a + b \tan(c+dx)} dx}{b} + \frac{(-2a^2B + aAb - b^2B) \tan(c+dx)}{bd} \\
 & \frac{b(a^2 + b^2)}{b(a^2 + b^2)} \\
 & \downarrow 25 \\
 & \frac{a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \\
 & \frac{(-2a^2B + aAb - b^2B) \tan(c+dx)}{bd} - \frac{\int -\frac{((aA + bB) \tan(c+dx)b^2) + (a^2 + b^2)(Ab - 2aB) \tan^2(c+dx) + a(-2Ba^2 + Aba - b^2B)}{a + b \tan(c+dx)} dx}{b} \\
 & \frac{b(a^2 + b^2)}{b(a^2 + b^2)} \\
 & \downarrow 3042 \\
 & \frac{a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \\
 & \frac{(-2a^2B + aAb - b^2B) \tan(c+dx)}{bd} - \frac{\int -\frac{((aA + bB) \tan(c+dx)b^2) + (a^2 + b^2)(Ab - 2aB) \tan^2(c+dx) + a(-2Ba^2 + Aba - b^2B)}{a + b \tan(c+dx)} dx}{b} \\
 & \frac{b(a^2 + b^2)}{b(a^2 + b^2)} \\
 & \downarrow 4109 \\
 & \frac{a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \\
 & \frac{(-2a^2B + aAb - b^2B) \tan(c+dx)}{bd} - \frac{b^2(a^2A + 2abB - Ab^2) \int \tan(c+dx) dx}{a^2 + b^2} + \frac{a^2(-2a^3B + a^2Ab - 4ab^2B + 3Ab^3) \int \frac{\tan^2(c+dx) + 1}{a + b \tan(c+dx)} dx}{a^2 + b^2} - \frac{b^2x(a^2(-B) + 2aAb + b^2)}{a^2 + b^2} \\
 & \frac{b(a^2 + b^2)}{b(a^2 + b^2)} \\
 & \downarrow 3042
 \end{aligned}$$

3.275. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\frac{\frac{a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \frac{(-2a^2B + aAb - b^2B) \tan(c + dx)}{bd}}{\frac{b^2(a^2A + 2abB - Ab^2) \int \tan(c + dx) dx}{a^2 + b^2} + \frac{a^2(-2a^3B + a^2Ab - 4ab^2B + 3Ab^3) \int \frac{\tan(c + dx)^2 + 1}{a + b \tan(c + dx)} dx}{a^2 + b^2} - \frac{b^2x(a^2(-B) + 2aAb + b^2B)}{a^2 + b^2}}$$

↓ 3956

$$\frac{\frac{a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \frac{(-2a^2B + aAb - b^2B) \tan(c + dx)}{bd}}{\frac{a^2(-2a^3B + a^2Ab - 4ab^2B + 3Ab^3) \int \frac{\tan(c + dx)^2 + 1}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{b^2(a^2A + 2abB - Ab^2) \log(\cos(c + dx))}{d(a^2 + b^2)} - \frac{b^2x(a^2(-B) + 2aAb + b^2B)}{a^2 + b^2}}$$

↓ 4100

$$\frac{\frac{a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \frac{(-2a^2B + aAb - b^2B) \tan(c + dx)}{bd}}{\frac{a^2(-2a^3B + a^2Ab - 4ab^2B + 3Ab^3) \int \frac{1}{a + b \tan(c + dx)} d(b \tan(c + dx))}{bd(a^2 + b^2)} + \frac{b^2(a^2A + 2abB - Ab^2) \log(\cos(c + dx))}{d(a^2 + b^2)} - \frac{b^2x(a^2(-B) + 2aAb + b^2B)}{a^2 + b^2}}$$

↓ 16

$$\frac{\frac{a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \frac{(-2a^2B + aAb - b^2B) \tan(c + dx)}{bd}}{\frac{b^2(a^2A + 2abB - Ab^2) \log(\cos(c + dx))}{d(a^2 + b^2)} - \frac{b^2x(a^2(-B) + 2aAb + b^2B)}{a^2 + b^2} + \frac{a^2(-2a^3B + a^2Ab - 4ab^2B + 3Ab^3) \log(a + b \tan(c + dx))}{bd(a^2 + b^2)}}$$

input `Int[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `(a*(A*b - a*B)*Tan[c + d*x]^2)/(b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])) - ((-((b^2*(2*a*A*b - a^2*B + b^2*B)*x)/(a^2 + b^2)) + (b^2*(a^2*A - A*b^2 + 2*a*b*B)*Log[Cos[c + d*x]])/((a^2 + b^2)*d) + (a^2*(a^2*A*b + 3*A*b^3 - 2*a^3*B - 4*a*b^2*B)*Log[a + b*Tan[c + d*x]])/(b*(a^2 + b^2)*d))/b + ((a*A*b - 2*a^2*B - b^2*B)*Tan[c + d*x])/(b*d)/(b*(a^2 + b^2))`

3.275.3.1 Defintions of rubi rules used

- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((A*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`
- rule 4100 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]`
- rule 4109 `Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C, 0]`

```
rule 4130 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
) + (f_.)*(x_)^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

3.275.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.83

method	result
derivativedivides	$\frac{\tan(dx+c)B}{b^2} + \frac{(-Aa^2+Ab^2-2Bab)\ln(1+\tan^2(dx+c)) + (-2Aab+B a^2 - B b^2) \arctan(\tan(dx+c))}{(a^2+b^2)^2} + \frac{a^2(Aa^2b+3Ab^3-2Ba^3-4Bb^3)}{b^3(a^2+b^2)}$
default	$\frac{\tan(dx+c)B}{b^2} + \frac{(-Aa^2+Ab^2-2Bab)\ln(1+\tan^2(dx+c)) + (-2Aab+B a^2 - B b^2) \arctan(\tan(dx+c))}{(a^2+b^2)^2} + \frac{a^2(Aa^2b+3Ab^3-2Ba^3-4Bb^3)}{b^3(a^2+b^2)}$
norman	$\frac{B(\tan^2(dx+c))}{bd} + \frac{(Aa^2b-2Ba^3-Bab^2)a}{db^3(a^2+b^2)} - \frac{a(2Aab-B a^2 + B b^2)x}{a^4+2a^2b^2+b^4} - \frac{b(2Aab-B a^2 + B b^2)x \tan(dx+c)}{a^4+2a^2b^2+b^4} + \frac{a^2(Aa^2b+3Ab^3-2Ba^3-4Bb^3)}{(a^2+b^2)^2}$
parallelrisch	$-\frac{2B\ln(1+\tan^2(dx+c))a^2b^4 - 2B(\tan^2(dx+c))a^4b^2 - 4B(\tan^2(dx+c))a^2b^4 - A\ln(1+\tan^2(dx+c))\tan(dx+c)b^6 + A\ln(1+\tan^2(dx+c))}{(a^2+b^2)^2}$
risch	$-\frac{x B}{2i b a - a^2 + b^2} + \frac{2i(-2i B a^3 b e^{2i(dx+c)} - 2i B a b^3 e^{2i(dx+c)} + 2B a^4 e^{2i(dx+c)} - B b^4 e^{2i(dx+c)} + 2B a^4 + 2B a^2 b^2 + B b^4 - (e^{2i(dx+c)} + 1)(ib+a)(-ib+a)^2(-ib e^{2i(dx+c)} + a e^{2i(dx+c)} + ib+a)b^2 d)}{(e^{2i(dx+c)} + 1)(ib+a)(-ib+a)^2(-ib e^{2i(dx+c)} + a e^{2i(dx+c)} + ib+a)b^2 d}$

```
input int (tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBO
SE)
```

```
output 1/d*(tan(d*x+c)*B/b^2+1/(a^2+b^2)^2*(1/2*(-A*a^2+A*b^2-2*B*a*b)*ln(1+tan(d
*x+c)^2)+(-2*A*a*b+B*a^2-B*b^2)*arctan(tan(d*x+c)))+1/b^3*a^2*(A*a^2*b+3*A
*b^3-2*B*a^3-4*B*a*b^2)/(a^2+b^2)^2*ln(a+b*tan(d*x+c))+1/b^3*a^3*(A*b-B*a)
/(a^2+b^2)/(a+b*tan(d*x+c)))
```

$$3.275. \int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.275.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 434 vs. $2(209) = 418$.

Time = 0.34 (sec) , antiderivative size = 434, normalized size of antiderivative = 2.09

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx =$$

$$2Ba^4b^2 - 2Aa^3b^3 - 2(Ba^3b^3 - 2Aa^2b^4 - Bab^5)dx - 2(Ba^4b^2 + 2Ba^2b^4 + Bb^6)\tan(dx+c)^2 + (2B$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fracas")`

output `-1/2*(2*B*a^4*b^2 - 2*A*a^3*b^3 - 2*(B*a^3*b^3 - 2*A*a^2*b^4 - B*a*b^5)*d*x - 2*(B*a^4*b^2 + 2*B*a^2*b^4 + B*b^6)*tan(d*x + c)^2 + (2*B*a^6 - A*a^5*b + 4*B*a^4*b^2 - 3*A*a^3*b^3 + (2*B*a^5*b - A*a^4*b^2 + 4*B*a^3*b^3 - 3*A*a^2*b^4)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - (2*B*a^6 - A*a^5*b + 4*B*a^4*b^2 - 2*A*a^3*b^3 + 2*B*a^2*b^4 - A*a*b^5 + (2*B*a^5*b - A*a^4*b^2 + 4*B*a^3*b^3 - 2*A*a^2*b^4 + 2*B*a*b^5 - A*b^6)*tan(d*x + c))*log(1/(tan(d*x + c)^2 + 1)) - 2*(2*B*a^5*b - A*a^4*b^2 + 2*B*a^3*b^3 + B*a*b^5 + (B*a^2*b^4 - 2*A*a*b^5 - B*b^6)*d*x)*tan(d*x + c))/((a^4*b^4 + 2*a^2*b^6 + b^8)*d*tan(d*x + c) + (a^5*b^3 + 2*a^3*b^5 + a*b^7)*d)`

3.275.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.40 (sec) , antiderivative size = 4534, normalized size of antiderivative = 21.80

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)**3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Piecewise((zoo*x*(A + B*tan(c))*tan(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (-A*log(tan(c + d*x)**2 + 1)/(2*d) + A*tan(c + d*x)**2/(2*d) + B*x + B*tan(c + d*x)**3/(3*d) - B*tan(c + d*x)/d)/a**2, Eq(b, 0)), (3*I*A*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 6*A*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 3*I*A*d*x/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 2*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 4*I*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 2*A*log(tan(c + d*x)**2 + 1)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 5*I*A*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 4*A/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 9*B*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 18*I*B*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 9*B*d*x/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 4*I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 8*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 4*I*B*log(tan(c + d*x)**2 + 1)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*t...`

3.275.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.06

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{\frac{2(Ba^2-2Aab-Bb^2)(dx+c)}{a^4+2a^2b^2+b^4} - \frac{2(2Ba^5-Aa^4b+4Ba^3b^2-3Aa^2b^3)\log(b\tan(dx+c)+a)}{a^4b^3+2a^2b^5+b^7} - \frac{(Aa^2+2Bab-Ab^2)\log(\tan(dx+c)^2+1)}{a^4+2a^2b^2+b^4}}{2d} - \frac{1}{a^3b^3}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `1/2*(2*(B*a^2 - 2*A*a*b - B*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) - 2*(2*B*a^5 - A*a^4*b + 4*B*a^3*b^2 - 3*A*a^2*b^3)*log(b*tan(d*x + c) + a)/(a^4*b^3 + 2*a^2*b^5 + b^7) - (A*a^2 + 2*B*a*b - A*b^2)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) - 2*(B*a^4 - A*a^3*b)/(a^3*b^3 + a*b^5 + (a^2*b^4 + b^6)*tan(d*x + c)) + 2*B*tan(d*x + c)/b^2)/d`

3.275. $\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

3.275.8 Giac [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 290, normalized size of antiderivative = 1.39

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{2(Ba^2-2Aab-Bb^2)(dx+c)}{a^4+2a^2b^2+b^4} - \frac{(Aa^2+2Bab-Ab^2)\log(\tan(dx+c)^2+1)}{a^4+2a^2b^2+b^4} - \frac{2(2Ba^5-Aa^4b+4Ba^3b^2-3Aa^2b^3)\log(|b\tan(dx+c)+a|)}{a^4b^3+2a^2b^5+b^7} + \frac{2B}{2d}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output $\frac{1}{2}*(2*(B*a^2 - 2*A*a*b - B*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) - (A*a^2 + 2*B*a*b - A*b^2)*\log(\tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) - 2*(2*B*a^5 - A*a^4*b + 4*B*a^3*b^2 - 3*A*a^2*b^3)*\log(\text{abs}(b*\tan(d*x + c) + a))/(a^4*b^3 + 2*a^2*b^5 + b^7) + 2*B*\tan(d*x + c)/b^2 + 2*(2*B*a^5*b*\tan(d*x + c) - A*a^4*b^2*\tan(d*x + c) + 4*B*a^3*b^3*\tan(d*x + c) - 3*A*a^2*b^4*\tan(d*x + c) + B*a^6 + 3*B*a^4*b^2 - 2*A*a^3*b^3)/((a^4*b^3 + 2*a^2*b^5 + b^7)*(b*\tan(d*x + c) + a)))/d$

3.275.9 Mupad [B] (verification not implemented)

Time = 8.33 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.01

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{B\tan(c+dx)}{b^2d} - \frac{\ln(a+b\tan(c+dx))(2Ba^5 - Aa^4b + 4Ba^3b^2 - 3Aa^2b^3)}{d(a^4b^3 + 2a^2b^5 + b^7)}$$

$$- \frac{\ln(\tan(c+dx) - i)(A + B i)}{2d(a^2 + ab2i - b^2)} - \frac{\ln(\tan(c+dx) + i)(B + A i)}{2d(a^2 i + 2ab - b^2 i)}$$

$$- \frac{a^2(Ba^2 - Aab)}{bd(\tan(c+dx)b^3 + ab^2)(a^2 + b^2)}$$

input `int((tan(c + d*x)^3*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`

output $(B*\tan(c + d*x))/(b^2*d) - (\log(a + b*\tan(c + d*x))*(2*B*a^5 - 3*A*a^2*b^3 + 4*B*a^3*b^2 - A*a^4*b))/(d*(b^7 + 2*a^2*b^5 + a^4*b^3)) - (\log(\tan(c + d*x) - i)*(A + B*i))/(2*d*(a*b*2i + a^2 - b^2)) - (\log(\tan(c + d*x) + i)*(A*i + B))/(2*d*(2*a*b + a^2*i - b^2*i)) - (a^2*(B*a^2 - A*a*b))/(b*d*(a*b^2 + b^3*\tan(c + d*x))*(a^2 + b^2))$

3.275. $\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

3.276
$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.276.1 Optimal result 2659
 3.276.2 Mathematica [C] (verified) 2659
 3.276.3 Rubi [A] (verified) 2660
 3.276.4 Maple [A] (verified) 2663
 3.276.5 Fricas [B] (verification not implemented) 2663
 3.276.6 Sympy [C] (verification not implemented) 2664
 3.276.7 Maxima [A] (verification not implemented) 2665
 3.276.8 Giac [A] (verification not implemented) 2665
 3.276.9 Mupad [B] (verification not implemented) 2666

3.276.1 Optimal result

Integrand size = 31, antiderivative size = 157

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = -\frac{(a^2A - Ab^2 + 2abB)x}{(a^2 + b^2)^2} - \frac{(2aAb - a^2B + b^2B) \log(\cos(c+dx))}{(a^2 + b^2)^2 d} - \frac{a(2Ab^3 - a(a^2 + 3b^2)B) \log(a+b \tan(c+dx))}{b^2(a^2 + b^2)^2 d} - \frac{a^2(Ab - aB)}{b^2(a^2 + b^2)d(a+b \tan(c+dx))}$$

```
output - (A*a^2-A*b^2+2*B*a*b)*x/(a^2+b^2)^2-(2*A*a*b-B*a^2+B*b^2)*ln(cos(d*x+c))/
(a^2+b^2)^2/d-a*(2*A*b^3-a*(a^2+3*b^2)*B)*ln(a+b*tan(d*x+c))/b^2/(a^2+b^2)^2/d-a^2*(A*b-B*a)/b^2/(a^2+b^2)/d/(a+b*tan(d*x+c))
```

3.276.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.89 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.93

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \frac{\frac{i(A+iB) \log(i-\tan(c+dx))}{(a+ib)^2} - \frac{i(A-iB) \log(i+\tan(c+dx))}{(a-ib)^2} + \frac{2a \left((-2Ab+a(3+\frac{a^2}{b^2})B) \log(a+b \tan(c+dx)) + \frac{a(a^2+b^2)(-Ab+aB)}{b^2(a+b \tan(c+dx))} \right)}{(a^2+b^2)^2}}{2d}$$

3.276.
$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

input `Integrate[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `((I*(A + I*B)*Log[I - Tan[c + d*x]])/(a + I*b)^2 - (I*(A - I*B)*Log[I + Tan[c + d*x]])/(a - I*b)^2 + (2*a*((-2*A*b + a*(3 + a^2/b^2)*B)*Log[a + b*Tan[c + d*x]] + (a*(a^2 + b^2)*(-A*b) + a*B))/(b^2*(a + b*Tan[c + d*x])))/(a^2 + b^2)^2)/(2*d)`

3.276.3 Rubi [A] (verified)

Time = 0.79 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.09, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 4087, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c + dx)^2(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{4087} \\
 & \frac{\int -\frac{((a^2+b^2)B \tan^2(c+dx)) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{a+b \tan(c+dx)} dx}{b(a^2 + b^2)} - \frac{a^2(Ab - aB)}{b^2 d (a^2 + b^2) (a + b \tan(c + dx))} \\
 & \quad \downarrow \text{25} \\
 & -\frac{\int -\frac{((a^2+b^2)B \tan^2(c+dx)) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{a+b \tan(c+dx)} dx}{b(a^2 + b^2)} - \frac{a^2(Ab - aB)}{b^2 d (a^2 + b^2) (a + b \tan(c + dx))} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\int -\frac{((a^2+b^2)B \tan(c+dx)^2) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{a+b \tan(c+dx)} dx}{b(a^2 + b^2)} - \frac{a^2(Ab - aB)}{b^2 d (a^2 + b^2) (a + b \tan(c + dx))} \\
 & \quad \downarrow \text{4109} \\
 & -\frac{b(a^2(-B)+2aAb+b^2B) \int \tan(c+dx) dx}{a^2+b^2} + \frac{a(2Ab^3-aB(a^2+3b^2)) \int \frac{\tan^2(c+dx)+1}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{bx(a^2A+2abB-Ab^2)}{a^2+b^2} \\
 & \quad \frac{b(a^2 + b^2)}{a^2(Ab - aB)} \\
 & \quad \frac{b^2 d (a^2 + b^2) (a + b \tan(c + dx))}
 \end{aligned}$$

3.276. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{-\frac{b(a^2(-B)+2aAb+b^2B) \int \tan(c+dx) dx}{a^2+b^2} + \frac{a(2Ab^3-aB(a^2+3b^2)) \int \frac{\tan(c+dx)^2+1}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{bx(a^2A+2abB-Ab^2)}{a^2+b^2}}{\frac{b(a^2+b^2)}{a^2(Ab-aB)}} \\
& \frac{\frac{b(a^2+b^2)}{a^2(Ab-aB)}}{b^2d(a^2+b^2)(a+b \tan(c+dx))} \\
& \downarrow 3956 \\
& \frac{\frac{a(2Ab^3-aB(a^2+3b^2)) \int \frac{\tan(c+dx)^2+1}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{b(a^2(-B)+2aAb+b^2B) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{bx(a^2A+2abB-Ab^2)}{a^2+b^2}}{\frac{b(a^2+b^2)}{a^2(Ab-aB)}} \\
& \frac{\frac{b(a^2+b^2)}{a^2(Ab-aB)}}{b^2d(a^2+b^2)(a+b \tan(c+dx))} \\
& \downarrow 4100 \\
& \frac{\frac{a(2Ab^3-aB(a^2+3b^2)) \int \frac{1}{a+b \tan(c+dx)} d(b \tan(c+dx))}{bd(a^2+b^2)} + \frac{b(a^2(-B)+2aAb+b^2B) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{bx(a^2A+2abB-Ab^2)}{a^2+b^2}}{\frac{b(a^2+b^2)}{a^2(Ab-aB)}} \\
& \frac{\frac{b(a^2+b^2)}{a^2(Ab-aB)}}{b^2d(a^2+b^2)(a+b \tan(c+dx))} \\
& \downarrow 16 \\
& \frac{\frac{a^2(Ab-aB)}{b^2d(a^2+b^2)(a+b \tan(c+dx))}}{\frac{b(a^2(-B)+2aAb+b^2B) \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{bx(a^2A+2abB-Ab^2)}{a^2+b^2} + \frac{a(2Ab^3-aB(a^2+3b^2)) \log(a+b \tan(c+dx))}{bd(a^2+b^2)}} \\
& \frac{b(a^2+b^2)}{b^2d(a^2+b^2)(a+b \tan(c+dx))}
\end{aligned}$$

input `Int[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `-(((b*(a^2*A - A*b^2 + 2*a*b*B)*x)/(a^2 + b^2) + (b*(2*a*A*b - a^2*B + b^2*B)*Log[Cos[c + d*x]])/((a^2 + b^2)*d) + (a*(2*A*b^3 - a*(a^2 + 3*b^2)*B)*Log[a + b*Tan[c + d*x]])/(b*(a^2 + b^2)*d))/(b*(a^2 + b^2))) - (a^2*(A*b - a*B))/(b^2*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))`

3.276.3.1 Defintions of rubi rules used

- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4087 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(- (B*c - A*d))*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`
- rule 4100 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]`
- rule 4109 `Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C, 0]`

3.276.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.99

method	result
derivativedivides	$\frac{\frac{(2Aab - B a^2 + B b^2) \ln(1 + \tan^2(dx+c))}{2} + (-A a^2 + A b^2 - 2Bab) \arctan(\tan(dx+c))}{(a^2 + b^2)^2} - \frac{a^2(Ab - Ba)}{b^2(a^2 + b^2)(a + b \tan(dx+c))} - \frac{a(2A b^3 - B a^3)}{d}$
default	$\frac{\frac{(2Aab - B a^2 + B b^2) \ln(1 + \tan^2(dx+c))}{2} + (-A a^2 + A b^2 - 2Bab) \arctan(\tan(dx+c))}{(a^2 + b^2)^2} - \frac{a^2(Ab - Ba)}{b^2(a^2 + b^2)(a + b \tan(dx+c))} - \frac{a(2A b^3 - B a^3)}{d}$
norman	$\frac{a(A a^2 - A b^2 + 2Bab)x}{a^4 + 2a^2b^2 + b^4} - \frac{b(A a^2 - A b^2 + 2Bab)x \tan(dx+c)}{a^4 + 2a^2b^2 + b^4} - \frac{(Aab - B a^2)a}{d b^2(a^2 + b^2)} + \frac{(2Aab - B a^2 + B b^2) \ln(1 + \tan^2(dx+c))}{2d(a^4 + 2a^2b^2 + b^4)} - \frac{a^2 b^2 \ln(a + b \tan(dx+c))}{a + b \tan(dx+c)}$
parallelrisch	$-2A a^2 b^3 + 2B a^5 - 2Ax \tan(dx+c) a^2 b^3 d - 4Bx \tan(dx+c) a b^4 d + 2B \ln(a + b \tan(dx+c)) a^5 - 2A a^4 b + 2B a^3 b^2 + B \ln(1 + \tan^2(dx+c))$
risch	$-\frac{2ia^4 Bx}{(a^4 + 2a^2b^2 + b^4)b^2} + \frac{x A}{2iba - a^2 + b^2} + \frac{2iBc}{d b^2} + \frac{4iabAc}{(a^4 + 2a^2b^2 + b^4)d} + \frac{2ia^2 A}{(ib+a)d(-ib+a)^2(-ib e^{2i(dx+c)} + a e^{2i(dx+c)})}$

input `int(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(1/(a^2+b^2)^2*(1/2*(2*A*a*b-B*a^2+B*b^2)*ln(1+tan(d*x+c)^2)+(-A*a^2+A*b^2-2*B*a*b)*arctan(tan(d*x+c)))-a^2*(A*b-B*a)/b^2/(a^2+b^2)/(a+b*tan(d*x+c))-a*(2*A*b^3-B*a^3-3*B*a*b^2)/(a^2+b^2)^2/b^2*ln(a+b*tan(d*x+c)))`

3.276.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 311 vs. 2(155) = 310.

Time = 0.31 (sec) , antiderivative size = 311, normalized size of antiderivative = 1.98

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{2Ba^3b^2 - 2Aa^2b^3 - 2(Aa^3b^2 + 2Ba^2b^3 - Aab^4)dx + (Ba^5 + 3Ba^3b^2 - 2Aa^2b^3 + (Ba^4b + 3Ba^2b^3 - 2Aa^2b^2 + 2Aab^2 + Bb^2) \ln(a+b \tan(c+dx)))}{(a+b \tan(c+dx))^2}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fracas")`

3.276.
$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

```
output 1/2*(2*B*a^3*b^2 - 2*A*a^2*b^3 - 2*(A*a^3*b^2 + 2*B*a^2*b^3 - A*a*b^4)*d*x
+ (B*a^5 + 3*B*a^3*b^2 - 2*A*a^2*b^3 + (B*a^4*b + 3*B*a^2*b^3 - 2*A*a*b^4
)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d
*x + c)^2 + 1)) - (B*a^5 + 2*B*a^3*b^2 + B*a*b^4 + (B*a^4*b + 2*B*a^2*b^3
+ B*b^5)*tan(d*x + c))*log(1/(tan(d*x + c)^2 + 1)) - 2*(B*a^4*b - A*a^3*b^
2 + (A*a^2*b^3 + 2*B*a*b^4 - A*b^5)*d*x)*tan(d*x + c))/((a^4*b^3 + 2*a^2*b
^5 + b^7)*d*tan(d*x + c) + (a^5*b^2 + 2*a^3*b^4 + a*b^6)*d)
```

3.276.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.05 (sec) , antiderivative size = 3485, normalized size of antiderivative = 22.20

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)**2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)
```

```
output Piecewise((zoo*x*(A + B*tan(c)), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), ((-A*x +
A*tan(c + d*x)/d - B*log(tan(c + d*x)**2 + 1)/(2*d) + B*tan(c + d*x)**2/(
2*d))/a**2, Eq(b, 0)), (A*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 -
8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 2*I*A*d*x*tan(c + d*x)/(4*b**2*d*tan
(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - A*d*x/(4*b**2*d*tan(c
+ d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 3*A*tan(c + d*x)/(4*b**
2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 2*I*A/(4*b**2*
d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 3*I*B*d*x*tan(c
+ d*x)**2/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d)
+ 6*B*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x)
- 4*b**2*d) - 3*I*B*d*x/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*
x) - 4*b**2*d) + 2*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2/(4*b**2*d*ta
n(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 4*I*B*log(tan(c + d*
x)**2 + 1)*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x
) - 4*b**2*d) - 2*B*log(tan(c + d*x)**2 + 1)/(4*b**2*d*tan(c + d*x)**2 - 8
*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 5*I*B*tan(c + d*x)/(4*b**2*d*tan(c +
d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 4*B/(4*b**2*d*tan(c + d*x)
**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d), Eq(a, -I*b)), (A*d*x*tan(c + d*
x)**2/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 2*
I*A*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x)...
```

3.276.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.25

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx =$$

$$\frac{\frac{2(Aa^2+2Bab-Ab^2)(dx+c)}{a^4+2a^2b^2+b^4} - \frac{2(Ba^4+3Ba^2b^2-2Aab^3)\log(b\tan(dx+c)+a)}{a^4b^2+2a^2b^4+b^6} + \frac{(Ba^2-2Aab-Bb^2)\log(\tan(dx+c)^2+1)}{a^4+2a^2b^2+b^4} - \frac{2}{a^3b^2+ab^4}}{2d}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `-1/2*(2*(A*a^2 + 2*B*a*b - A*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) - 2*(B*a^4 + 3*B*a^2*b^2 - 2*A*a*b^3)*log(b*tan(d*x + c) + a)/(a^4*b^2 + 2*a^2*b^4 + b^6) + (B*a^2 - 2*A*a*b - B*b^2)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) - 2*(B*a^3 - A*a^2*b)/(a^3*b^2 + a*b^4 + (a^2*b^3 + b^5)*tan(d*x + c)))/d`

3.276.8 Giac [A] (verification not implemented)

Time = 0.58 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.55

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx =$$

$$\frac{\frac{2(Aa^2+2Bab-Ab^2)(dx+c)}{a^4+2a^2b^2+b^4} + \frac{(Ba^2-2Aab-Bb^2)\log(\tan(dx+c)^2+1)}{a^4+2a^2b^2+b^4} - \frac{2(Ba^4+3Ba^2b^2-2Aab^3)\log(|b\tan(dx+c)+a|)}{a^4b^2+2a^2b^4+b^6} + \frac{2(Ba^4\tan(dx+c)+a)}{a^3b^2+ab^4}}{2d}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output `-1/2*(2*(A*a^2 + 2*B*a*b - A*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) + (B*a^2 - 2*A*a*b - B*b^2)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) - 2*(B*a^4 + 3*B*a^2*b^2 - 2*A*a*b^3)*log(abs(b*tan(d*x + c) + a))/(a^4*b^2 + 2*a^2*b^4 + b^6) + 2*(B*a^4*tan(d*x + c) + 3*B*a^2*b^2*tan(d*x + c) - 2*A*a*b^3*tan(d*x + c) + A*a^4 + 2*B*a^3*b - A*a^2*b^2)/((a^4*b + 2*a^2*b^3 + b^5)*(b*tan(d*x + c) + a)))/d`

3.276.9 Mupad [B] (verification not implemented)

Time = 7.84 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.05

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{\ln(\tan(c+dx)+1i)(B+A1i)}{2d(-a^2+ab2i+b^2)} + \frac{\ln(\tan(c+dx)-1i)(A+B1i)}{2d(-a^21i+2ab+b^21i)}$$

$$- \frac{a^2(Ab-BA)}{b^2d(a^2+b^2)(a+b\tan(c+dx))}$$

$$+ \frac{a\ln(a+b\tan(c+dx))(Ba^3+3Bab^2-2Ab^3)}{b^2d(a^2+b^2)^2}$$

input `int((tan(c + d*x)^2*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`output `(log(tan(c + d*x) + 1i)*(A*1i + B))/(2*d*(a*b*2i - a^2 + b^2)) + (log(tan(c + d*x) - 1i)*(A + B*1i))/(2*d*(2*a*b - a^2*1i + b^2*1i)) - (a^2*(A*b - B*a))/(b^2*d*(a^2 + b^2)*(a + b*tan(c + d*x))) + (a*log(a + b*tan(c + d*x)))*(B*a^3 - 2*A*b^3 + 3*B*a*b^2)/(b^2*d*(a^2 + b^2)^2)`

$$3.277 \quad \int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.277.1 Optimal result	2667
3.277.2 Mathematica [C] (verified)	2667
3.277.3 Rubi [A] (verified)	2668
3.277.4 Maple [A] (verified)	2670
3.277.5 Fricas [A] (verification not implemented)	2670
3.277.6 Sympy [C] (verification not implemented)	2671
3.277.7 Maxima [A] (verification not implemented)	2672
3.277.8 Giac [B] (verification not implemented)	2672
3.277.9 Mupad [B] (verification not implemented)	2673

3.277.1 Optimal result

Integrand size = 29, antiderivative size = 115

$$\begin{aligned} & \int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\ &= \frac{(2aAb - a^2B + b^2B)x}{(a^2 + b^2)^2} - \frac{(a^2A - Ab^2 + 2abB) \log(a \cos(c+dx) + b \sin(c+dx))}{(a^2 + b^2)^2 d} \\ & \quad + \frac{a(Ab - aB)}{b(a^2 + b^2) d(a + b \tan(c+dx))} \end{aligned}$$

output $(2Aa^2b - Bb^2a^2 + Bb^2)x / (a^2 + b^2)^2 - (Aa^2 - Ab^2 + 2Bab) \ln(a \cos(dx+c) + b \sin(dx+c)) / (a^2 + b^2)^2 d + a(Ab - Ba) / (b(a^2 + b^2) d(a + b \tan(dx+c)))$

3.277.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.27 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.22

$$\begin{aligned} & \int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\ &= \frac{(A+iB) \log(i - \tan(c+dx))}{(a+ib)^2} + \frac{(A-iB) \log(i + \tan(c+dx))}{(a-ib)^2} + \frac{2 \left((-a^2A + Ab^2 - 2abB) \log(a + b \tan(c+dx)) - \frac{a(a^2 + b^2)(-Ab + aB)}{b(a + b \tan(c+dx))} \right)}{(a^2 + b^2)^2} \\ & \quad \frac{1}{2d} \end{aligned}$$

3.277. $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

input `Integrate[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `((A + I*B)*Log[I - Tan[c + d*x]])/(a + I*b)^2 + ((A - I*B)*Log[I + Tan[c + d*x]])/(a - I*b)^2 + (2*((-a^2*A) + A*b^2 - 2*a*b*B)*Log[a + b*Tan[c + d*x]] - (a*(a^2 + b^2)*(-A*b) + a*B))/(b*(a + b*Tan[c + d*x]))/(a^2 + b^2)^2/(2*d)`

3.277.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.10, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.207$, Rules used = {3042, 4074, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{4074} \\
 & \frac{\int \frac{Ab-aB+(aA+bB) \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{a(Ab-aB)}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{Ab-aB+(aA+bB) \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{a(Ab-aB)}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{4014} \\
 & \frac{x(a^2(-B)+2aAb+b^2B)}{a^2+b^2} - \frac{(a^2A+2abB-Ab^2) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{a(Ab-aB)}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{x(a^2(-B)+2aAb+b^2B)}{a^2+b^2} - \frac{(a^2A+2abB-Ab^2) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{a(Ab-aB)}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{4013}
 \end{aligned}$$

3.277. $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\frac{a(Ab - aB)}{bd(a^2 + b^2)(a + b \tan(c + dx))} + \frac{\frac{x(a^2(-B) + 2aAb + b^2B)}{a^2 + b^2} - \frac{(a^2A + 2abB - Ab^2) \log(a \cos(c + dx) + b \sin(c + dx))}{d(a^2 + b^2)}}{a^2 + b^2}$$

input `Int[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `((2*a*A*b - a^2*B + b^2*B)*x)/(a^2 + b^2) - ((a^2*A - A*b^2 + 2*a*b*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]]/((a^2 + b^2)*d))/(a^2 + b^2) + (a*(A*b - a*B))/(b*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))`

3.277.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4074 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.277.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.26

method	result
derivativedivides	$\frac{\frac{(A a^2 - A b^2 + 2Bab) \ln(1 + \tan^2(dx+c))}{2} + \frac{(2Aab - B a^2 + B b^2) \arctan(\tan(dx+c))}{(a^2+b^2)^2} + \frac{a(Ab-Ba)}{(a^2+b^2)b(a+b \tan(dx+c))} - \frac{(A a^2 - A b^2 + 2Bab)}{(a^2+b^2)^2}}{d}$
default	$\frac{\frac{(A a^2 - A b^2 + 2Bab) \ln(1 + \tan^2(dx+c))}{2} + \frac{(2Aab - B a^2 + B b^2) \arctan(\tan(dx+c))}{(a^2+b^2)^2} + \frac{a(Ab-Ba)}{(a^2+b^2)b(a+b \tan(dx+c))} - \frac{(A a^2 - A b^2 + 2Bab)}{(a^2+b^2)^2}}{d}$
norman	$\frac{\frac{a(2Aab - B a^2 + B b^2)x}{a^4 + 2a^2b^2 + b^4} + \frac{b(2Aab - B a^2 + B b^2)x \tan(dx+c)}{a^4 + 2a^2b^2 + b^4} + \frac{a(Ab-Ba)}{(a^2+b^2)bd}}{a+b \tan(dx+c)} + \frac{(A a^2 - A b^2 + 2Bab) \ln(1 + \tan^2(dx+c))}{2d(a^4 + 2a^2b^2 + b^4)} - \frac{(A a^2 - A b^2 + 2Bab)}{(a^2+b^2)^2}$
parallelrisch	$\frac{2A a^3 b + 2A a b^3 - 2B a^2 b^2 - 2B a^4 - 2A \ln(a+b \tan(dx+c)) a^3 b + A \ln(1 + \tan^2(dx+c)) \tan(dx+c) a^2 b^2 - 2A \ln(a+b \tan(dx+c))}{(a^2+b^2)^2}$
risch	$\frac{x B}{2i b a - a^2 + b^2} + \frac{i x A}{2i b a - a^2 + b^2} + \frac{2i a^2 A x}{a^4 + 2a^2 b^2 + b^4} - \frac{2i A b^2 x}{a^4 + 2a^2 b^2 + b^4} + \frac{4i B a b x}{a^4 + 2a^2 b^2 + b^4} + \frac{2i a^2 A c}{d(a^4 + 2a^2 b^2 + b^4)} - \frac{2i A c}{d(a^4 + 2a^2 b^2 + b^4)}$

input `int(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(1/(a^2+b^2)^2*(1/2*(A*a^2-A*b^2+2*B*a*b)*ln(1+tan(d*x+c)^2)+(2*A*a*b-B*a^2+B*b^2)*arctan(tan(d*x+c)))+a*(A*b-B*a)/(a^2+b^2)/b/(a+b*tan(d*x+c))- (A*a^2-A*b^2+2*B*a*b)/(a^2+b^2)^2*ln(a+b*tan(d*x+c)))`

3.277.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.92

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \frac{2Ba^2b - 2Aab^2 + 2(Ba^3 - 2Aa^2b - Bab^2)dx + (Aa^3 + 2Ba^2b - Aab^2 + (Aa^2b + 2Bab^2 - Ab^3) \tan(c+dx))}{2((a^4b + 2a^2b^3 + b^5)d \tan(c+dx) + (Aa^3 + 2Ba^2b - Aab^2 + (Aa^2b + 2Bab^2 - Ab^3) \tan(c+dx))}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

output
$$-1/2*(2*B*a^2*b - 2*A*a*b^2 + 2*(B*a^3 - 2*A*a^2*b - B*a*b^2)*d*x + (A*a^3 + 2*B*a^2*b - A*a*b^2 + (A*a^2*b + 2*B*a*b^2 - A*b^3)*\tan(d*x + c))*\log((b^2*\tan(d*x + c)^2 + 2*a*b*\tan(d*x + c) + a^2)/(\tan(d*x + c)^2 + 1)) - 2*(B*a^3 - A*a^2*b - (B*a^2*b - 2*A*a*b^2 - B*b^3)*d*x)*\tan(d*x + c))/((a^4*b + 2*a^2*b^3 + b^5)*d*\tan(d*x + c) + (a^5 + 2*a^3*b^2 + a*b^4)*d)$$

3.277.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.84 (sec) , antiderivative size = 2987, normalized size of antiderivative = 25.97

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Piecewise((zoo*x*(A + B*tan(c))/tan(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), ((A*log(tan(c + d*x)**2 + 1)/(2*d) - B*x + B*tan(c + d*x)/d)/a**2, Eq(b, 0)), (I*A*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 2*A*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - I*A*d*x/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + I*A*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + B*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 2*I*B*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - B*d*x/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 3*B*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 2*I*B/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d), Eq(a, -I*b)), (-I*A*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 2*A*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + I*A*d*x/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - I*A*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + B*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 2*I*B*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - B*d*x/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4...`

3.277.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.61

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx =$$

$$-\frac{\frac{2(Ba^2-2Aab-Bb^2)(dx+c)}{a^4+2a^2b^2+b^4} + \frac{2(Aa^2+2Bab-Ab^2)\log(b\tan(dx+c)+a)}{a^4+2a^2b^2+b^4} - \frac{(Aa^2+2Bab-Ab^2)\log(\tan(dx+c)^2+1)}{a^4+2a^2b^2+b^4} + \frac{2(Ba^2-2Aab-Bb^2)}{a^3b+ab^3+(a^2b^2+b^4)}}{2d}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `-1/2*(2*(B*a^2 - 2*A*a*b - B*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) + 2*(A*a^2 + 2*B*a*b - A*b^2)*log(b*tan(d*x + c) + a)/(a^4 + 2*a^2*b^2 + b^4) - (A*a^2 + 2*B*a*b - A*b^2)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) + 2*(B*a^2 - A*a*b)/(a^3*b + a*b^3 + (a^2*b^2 + b^4)*tan(d*x + c)))/d`

3.277.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 241 vs. 2(117) = 234.

Time = 0.43 (sec) , antiderivative size = 241, normalized size of antiderivative = 2.10

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx =$$

$$-\frac{\frac{2(Ba^2-2Aab-Bb^2)(dx+c)}{a^4+2a^2b^2+b^4} - \frac{(Aa^2+2Bab-Ab^2)\log(\tan(dx+c)^2+1)}{a^4+2a^2b^2+b^4} + \frac{2(Aa^2b+2Bab^2-Ab^3)\log(|b\tan(dx+c)+a|)}{a^4b+2a^2b^3+b^5} - \frac{2(Aa^2b^2\tan(dx+c))}{a^4b+2a^2b^3+b^5}}{2d}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output `-1/2*(2*(B*a^2 - 2*A*a*b - B*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) - (A*a^2 + 2*B*a*b - A*b^2)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) + 2*(A*a^2*b + 2*B*a*b^2 - A*b^3)*log(abs(b*tan(d*x + c) + a))/(a^4*b + 2*a^2*b^3 + b^5) - 2*(A*a^2*b^2*tan(d*x + c) + 2*B*a*b^3*tan(d*x + c) - A*b^4*tan(d*x + c) - B*a^4 + 2*A*a^3*b + B*a^2*b^2)/((a^4*b + 2*a^2*b^3 + b^5)*(b*tan(d*x + c) + a)))/d`

3.277. $\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

3.277.9 Mupad [B] (verification not implemented)

Time = 8.19 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.42

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \frac{a(Ab-Ba)}{bd(a^2+b^2)(a+b\tan(c+dx))} + \frac{\ln(\tan(c+dx)-i)(A+B1i)}{2d(a^2+ab2i-b^2)} + \frac{\ln(\tan(c+dx)+1i)(B+A1i)}{2d(a^21i+2ab-b^21i)} - \frac{\ln(a+b\tan(c+dx))\left(\frac{A}{a^2+b^2} - \frac{2b(Ab-Ba)}{(a^2+b^2)^2}\right)}{d}$$

input `int((tan(c + d*x)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`output `(log(tan(c + d*x) - 1i)*(A + B*1i))/(2*d*(a*b*2i + a^2 - b^2)) - (log(a + b*tan(c + d*x))*(A/(a^2 + b^2) - (2*b*(A*b - B*a))/(a^2 + b^2)^2))/d + (log(tan(c + d*x) + 1i)*(A*1i + B))/(2*d*(2*a*b + a^2*1i - b^2*1i)) + (a*(A*b - B*a))/(b*d*(a^2 + b^2)*(a + b*tan(c + d*x)))`

3.278 $\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^2} dx$

3.278.1 Optimal result	2674
3.278.2 Mathematica [C] (verified)	2674
3.278.3 Rubi [A] (verified)	2675
3.278.4 Maple [A] (verified)	2677
3.278.5 Fricas [A] (verification not implemented)	2677
3.278.6 Sympy [C] (verification not implemented)	2678
3.278.7 Maxima [A] (verification not implemented)	2679
3.278.8 Giac [B] (verification not implemented)	2679
3.278.9 Mupad [B] (verification not implemented)	2680

3.278.1 Optimal result

Integrand size = 23, antiderivative size = 111

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^2} dx = \frac{(a^2 A - Ab^2 + 2abB) x}{(a^2 + b^2)^2} + \frac{(2aAb - a^2 B + b^2 B) \log(a \cos(c + dx) + b \sin(c + dx))}{(a^2 + b^2)^2 d} - \frac{Ab - aB}{(a^2 + b^2) d(a + b \tan(c + dx))}$$

```
output (A*a^2-A*b^2+2*B*a*b)*x/(a^2+b^2)^2+(2*A*a*b-B*a^2+B*b^2)*ln(a*cos(d*x+c)+
b*sin(d*x+c))/(a^2+b^2)^2/d+(-A*b+B*a)/(a^2+b^2)/d/(a+b*tan(d*x+c))
```

3.278.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.33 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.71

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^2} dx = \frac{B((-ia-b) \log(i - \tan(c+dx)) + i(a+ib) \log(i + \tan(c+dx)) + 2b \log(a + b \tan(c+dx)))}{a^2 + b^2} - (Ab - aB) \left(\frac{i \log(i - \tan(c+dx))}{(a+ib)^2} - \frac{i \log(i + \tan(c+dx))}{(a-ib)^2} \right)$$

input `Integrate[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^2,x]`

output `((B*(((-1)*a - b)*Log[I - Tan[c + d*x]] + I*(a + I*b)*Log[I + Tan[c + d*x]] + 2*b*Log[a + b*Tan[c + d*x]]))/(a^2 + b^2) - (A*b - a*B)*((I*Log[I - Tan[c + d*x]])/(a + I*b)^2 - (I*Log[I + Tan[c + d*x]])/(a - I*b)^2 + (2*b*(-2*a*Log[a + b*Tan[c + d*x]] + (a^2 + b^2)/(a + b*Tan[c + d*x])))/(a^2 + b^2)^2))/(2*b*d)`

3.278.3 Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.10, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {3042, 4012, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{4012} \\
 & \frac{\int \frac{aA + bB - (Ab - aB) \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} - \frac{Ab - aB}{d(a^2 + b^2)(a + b \tan(c + dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{aA + bB - (Ab - aB) \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} - \frac{Ab - aB}{d(a^2 + b^2)(a + b \tan(c + dx))} \\
 & \quad \downarrow \text{4014} \\
 & \frac{(a^2(-B) + 2aAb + b^2B) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{x(a^2A + 2abB - Ab^2)}{a^2 + b^2} - \frac{Ab - aB}{d(a^2 + b^2)(a + b \tan(c + dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(a^2(-B) + 2aAb + b^2B) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{x(a^2A + 2abB - Ab^2)}{a^2 + b^2} - \frac{Ab - aB}{d(a^2 + b^2)(a + b \tan(c + dx))}
 \end{aligned}$$

3.278. $\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^2} dx$

$$\frac{\frac{(a^2(-B)+2aAb+b^2B) \log(a \cos(c+dx)+b \sin(c+dx))}{d(a^2+b^2)} + \frac{x(a^2A+2abB-Ab^2)}{a^2+b^2}}{a^2+b^2} - \frac{Ab-aB}{d(a^2+b^2)(a+b \tan(c+dx))}$$

input `Int[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^2,x]`

output `((a^2*A - A*b^2 + 2*a*b*B)*x)/(a^2 + b^2) + ((2*a*A*b - a^2*B + b^2*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]]/((a^2 + b^2)*d))/(a^2 + b^2) - (A*b - a*B)/((a^2 + b^2)*d*(a + b*Tan[c + d*x]))`

3.278.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4013 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

3.278.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.27

method	result
derivativedivides	$\frac{\frac{(-2Aab+B a^2-B b^2) \ln(1+\tan^2(dx+c))}{2} + (A a^2-A b^2+2Bab) \arctan(\tan(dx+c))}{(a^2+b^2)^2} - \frac{Ab-Ba}{(a^2+b^2)(a+b \tan(dx+c))} + \frac{(2Aab-B a^2+B b^2) \ln(a+b \tan(dx+c))}{(a^2+b^2)}$
default	$\frac{\frac{(-2Aab+B a^2-B b^2) \ln(1+\tan^2(dx+c))}{2} + (A a^2-A b^2+2Bab) \arctan(\tan(dx+c))}{(a^2+b^2)^2} - \frac{Ab-Ba}{(a^2+b^2)(a+b \tan(dx+c))} + \frac{(2Aab-B a^2+B b^2) \ln(a+b \tan(dx+c))}{(a^2+b^2)}$
norman	$\frac{a(A a^2-A b^2+2Bab)x}{a^4+2a^2b^2+b^4} + \frac{b(A a^2-A b^2+2Bab)x \tan(dx+c)}{a^4+2a^2b^2+b^4} + \frac{(Ab-Ba)b \tan(dx+c)}{ad(a^2+b^2)} + \frac{(2Aab-B a^2+B b^2) \ln(a+b \tan(dx+c))}{d(a^4+2a^2b^2+b^4)}$
parallelrisch	$- \frac{-2A b^4 \tan(dx+c) - B \ln(1+\tan^2(dx+c)) \tan(dx+c) a^3 b + 2B \ln(a+b \tan(dx+c)) \tan(dx+c) a^3 b - 2A a^2 b^2 \tan(dx+c) + \dots}{a+b \tan(dx+c)}$
risch	$\frac{ixB}{2iba-a^2+b^2} - \frac{xA}{2iba-a^2+b^2} - \frac{4iabAx}{a^4+2a^2b^2+b^4} + \frac{2ia^2Bx}{a^4+2a^2b^2+b^4} - \frac{2iBb^2x}{a^4+2a^2b^2+b^4} - \frac{4iabAc}{(a^4+2a^2b^2+b^4)d} + \frac{2i}{(a^4+2a^2b^2+b^4)}$

input `int((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(1/(a^2+b^2)^2*(1/2*(-2*A*a*b+B*a^2-B*b^2)*ln(1+tan(d*x+c)^2)+(A*a^2-A*b^2+2*B*a*b)*arctan(tan(d*x+c)))-(A*b-B*a)/(a^2+b^2)/(a+b*tan(d*x+c))+(2*A*a*b-B*a^2+B*b^2)/(a^2+b^2)^2*ln(a+b*tan(d*x+c)))`

3.278.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 222, normalized size of antiderivative = 2.00

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{2 Bab^2 - 2 Ab^3 + 2 (Aa^3 + 2 Ba^2b - Aab^2)dx - (Ba^3 - 2 Aa^2b - Bab^2 + (Ba^2b - 2 Aab^2 - Bb^3) \tan(dx+c))}{2((a^4b + 2 a^2b^3 + b^5)d \tan(dx+c) + \dots)}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

output $1/2*(2*B*a*b^2 - 2*A*b^3 + 2*(A*a^3 + 2*B*a^2*b - A*a*b^2)*d*x - (B*a^3 - 2*A*a^2*b - B*a*b^2 + (B*a^2*b - 2*A*a*b^2 - B*b^3)*\tan(d*x + c))*\log((b^2 * \tan(d*x + c)^2 + 2*a*b*\tan(d*x + c) + a^2)/(\tan(d*x + c)^2 + 1)) - 2*(B*a^2*b - A*a*b^2 - (A*a^2*b + 2*B*a*b^2 - A*b^3)*d*x)*\tan(d*x + c))/((a^4*b + 2*a^2*b^3 + b^5)*d*\tan(d*x + c) + (a^5 + 2*a^3*b^2 + a*b^4)*d)$

3.278.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.83 (sec) , antiderivative size = 2878, normalized size of antiderivative = 25.93

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Piecewise((zoo*x*(A + B*tan(c))/tan(c)**2, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), ((A*x + B*log(tan(c + d*x)**2 + 1)/(2*d))/a**2, Eq(b, 0)), (-A*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 2*I*A*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + A*d*x/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - A*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 2*I*A/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + I*B*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 2*B*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - I*B*d*x/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + I*B*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 - 8*I*b**2*d*tan(c + d*x) - 4*b**2*d), Eq(a, -I*b)), (-A*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 2*I*A*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + A*d*x/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - A*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - 2*I*A/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) - I*B*d*x*tan(c + d*x)**2/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + 2*B*d*x*tan(c + d*x)/(4*b**2*d*tan(c + d*x)**2 + 8*I*b**2*d*tan(c + d*x) - 4*b**2*d) + I*B*d*x/(4...`

3.278.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.59

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{2(Aa^2 + 2Bab - Ab^2)(dx+c)}{a^4 + 2a^2b^2 + b^4} - \frac{2(Ba^2 - 2Aab - Bb^2) \log(b \tan(dx+c) + a)}{a^4 + 2a^2b^2 + b^4} + \frac{(Ba^2 - 2Aab - Bb^2) \log(\tan(dx+c)^2 + 1)}{a^4 + 2a^2b^2 + b^4} + \frac{2(Ba - Ab)}{a^3 + ab^2 + (a^2b + b^3)}$$

$2d$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`output `1/2*(2*(A*a^2 + 2*B*a*b - A*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) - 2*(B*a^2 - 2*A*a*b - B*b^2)*log(b*tan(d*x + c) + a)/(a^4 + 2*a^2*b^2 + b^4) + (B*a^2 - 2*A*a*b - B*b^2)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) + 2*(B*a - A*b)/(a^3 + a*b^2 + (a^2*b + b^3)*tan(d*x + c)))/d`**3.278.8 Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 234 vs. 2(111) = 222.

Time = 0.40 (sec) , antiderivative size = 234, normalized size of antiderivative = 2.11

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{2(Aa^2 + 2Bab - Ab^2)(dx+c)}{a^4 + 2a^2b^2 + b^4} + \frac{(Ba^2 - 2Aab - Bb^2) \log(\tan(dx+c)^2 + 1)}{a^4 + 2a^2b^2 + b^4} - \frac{2(Ba^2b - 2Aab^2 - Bb^3) \log(|b \tan(dx+c) + a|)}{a^4b + 2a^2b^3 + b^5} + \frac{2(Ba^2b \tan(dx+c) - Ab^3)}{a^4b + 2a^2b^3 + b^5}$$

$2d$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")`output `1/2*(2*(A*a^2 + 2*B*a*b - A*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) + (B*a^2 - 2*A*a*b - B*b^2)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) - 2*(B*a^2*b - 2*A*a*b^2 - B*b^3)*log(abs(b*tan(d*x + c) + a))/(a^4*b + 2*a^2*b^3 + b^5) + 2*(B*a^2*b*tan(d*x + c) - 2*A*a*b^2*tan(d*x + c) - B*b^3*tan(d*x + c) + 2*B*a^3 - 3*A*a^2*b - A*b^3)/((a^4 + 2*a^2*b^2 + b^4)*(b*tan(d*x + c) + a)))/d`

3.278.9 Mupad [B] (verification not implemented)

Time = 8.47 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.38

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^2} dx = \frac{\ln(a + b \tan(c + dx)) (-B a^2 + 2 A a b + B b^2)}{d (a^2 + b^2)^2} - \frac{A b - B a}{d (a^2 + b^2) (a + b \tan(c + dx))} - \frac{\ln(\tan(c + dx) + 1i) (B + A 1i)}{2 d (-a^2 + a b 2i + b^2)} - \frac{\ln(\tan(c + dx) - i) (A + B 1i)}{2 d (-a^2 1i + 2 a b + b^2 1i)}$$

input `int((A + B*tan(c + d*x))/(a + b*tan(c + d*x))^2,x)`output `(log(a + b*tan(c + d*x))*(B*b^2 - B*a^2 + 2*A*a*b))/(d*(a^2 + b^2)^2) - (A*b - B*a)/(d*(a^2 + b^2)*(a + b*tan(c + d*x))) - (log(tan(c + d*x) + 1i)*(A*1i + B))/(2*d*(a*b*2i - a^2 + b^2)) - (log(tan(c + d*x) - 1i)*(A + B*1i))/(2*d*(2*a*b - a^2*1i + b^2*1i))`

3.279 $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.279.1 Optimal result 2681
 3.279.2 Mathematica [C] (verified) 2681
 3.279.3 Rubi [A] (verified) 2682
 3.279.4 Maple [A] (verified) 2685
 3.279.5 Fricas [B] (verification not implemented) 2685
 3.279.6 Sympy [C] (verification not implemented) 2686
 3.279.7 Maxima [A] (verification not implemented) 2687
 3.279.8 Giac [B] (verification not implemented) 2687
 3.279.9 Mupad [B] (verification not implemented) 2688

3.279.1 Optimal result

Integrand size = 29, antiderivative size = 137

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= -\frac{(2aAb - a^2B + b^2B)x}{(a^2 + b^2)^2} + \frac{A \log(\sin(c+dx))}{a^2d}$$

$$- \frac{b(3a^2Ab + Ab^3 - 2a^3B) \log(a \cos(c+dx) + b \sin(c+dx))}{a^2(a^2 + b^2)^2d}$$

$$+ \frac{b(Ab - aB)}{a(a^2 + b^2)d(a + b \tan(c+dx))}$$

output

```
-(2*A*a*b-B*a^2+B*b^2)*x/(a^2+b^2)^2+A*ln(sin(d*x+c))/a^2/d-b*(3*A*a^2*b+A
*b^3-2*B*a^3)*ln(a*cos(d*x+c)+b*sin(d*x+c))/a^2/(a^2+b^2)^2/d+b*(A*b-B*a)/
a/(a^2+b^2)/d/(a+b*tan(d*x+c))
```

3.279.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.96 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.34

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{-\frac{a(a-ib)(A+iB) \log(i-\tan(c+dx))}{2(a+ib)} + \frac{A(a^2+b^2) \log(\tan(c+dx))}{a} - \frac{a(a+ib)(A-iB) \log(i+\tan(c+dx))}{2(a-ib)} + \frac{b(-3a^2Ab - Ab^3 + 2a^3B) \log(a + b \tan(c+dx))}{a(a^2+b^2)}}{a(a^2 + b^2)d}$$

3.279. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

input `Integrate[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output $(-1/2*(a*(a - I*b)*(A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]])/(a + I*b) + (A*(a^2 + b^2)*\text{Log}[\text{Tan}[c + d*x]])/a - (a*(a + I*b)*(A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]])/(2*(a - I*b)) + (b*(-3*a^2*A*b - A*b^3 + 2*a^3*B)*\text{Log}[a + b*\text{Tan}[c + d*x]])/(a*(a^2 + b^2)) + (b*(A*b - a*B))/(a + b*\text{Tan}[c + d*x])/(a*(a^2 + b^2)*d)$

3.279.3 Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.18, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 4092, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)(a+b \tan(c+dx))^2} dx \\ & \quad \downarrow \text{4092} \\ & \frac{\int \frac{\cot(c+dx)(b(Ab-aB) \tan^2(c+dx)-a(Ab-aB) \tan(c+dx)+A(a^2+b^2))}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{b(Ab-aB)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{b(Ab-aB) \tan(c+dx)^2-a(Ab-aB) \tan(c+dx)+A(a^2+b^2)}{\tan(c+dx)(a+b \tan(c+dx))} dx}{a(a^2+b^2)} + \frac{b(Ab-aB)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\ & \quad \downarrow \text{4134} \\ & \frac{\frac{A(a^2+b^2)}{a} \int \cot(c+dx) dx - \frac{b(-2a^3B+3a^2Ab+Ab^3)}{a(a^2+b^2)} \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx - \frac{ax(a^2(-B)+2aAb+b^2B)}{a^2+b^2}}{a(a^2+b^2)} + \frac{b(Ab-aB)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.279. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\begin{aligned}
 & \frac{\frac{A(a^2+b^2) \int -\tan(c+dx+\frac{\pi}{2})dx}{a} - \frac{b(-2a^3B+3a^2Ab+Ab^3) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{ax(a^2(-B)+2aAb+b^2B)}{a^2+b^2}}{a(a^2+b^2)} + \\
 & \quad \frac{b(Ab-aB)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{-\frac{A(a^2+b^2) \int \tan(\frac{1}{2}(2c+\pi)+dx) dx}{a} - \frac{b(-2a^3B+3a^2Ab+Ab^3) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{ax(a^2(-B)+2aAb+b^2B)}{a^2+b^2}}{a(a^2+b^2)} + \\
 & \quad \frac{b(Ab-aB)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{3956} \\
 & \frac{-\frac{b(-2a^3B+3a^2Ab+Ab^3) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{ax(a^2(-B)+2aAb+b^2B)}{a^2+b^2} + \frac{A(a^2+b^2) \log(-\sin(c+dx))}{ad}}{a(a^2+b^2)} + \\
 & \quad \frac{b(Ab-aB)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{4013} \\
 & \frac{\frac{b(Ab-aB)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \frac{-\frac{ax(a^2(-B)+2aAb+b^2B)}{a^2+b^2} + \frac{A(a^2+b^2) \log(-\sin(c+dx))}{ad} - \frac{b(-2a^3B+3a^2Ab+Ab^3) \log(a \cos(c+dx)+b \sin(c+dx))}{ad(a^2+b^2)}}{a(a^2+b^2)}}{a(a^2+b^2)}
 \end{aligned}$$

input `Int[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `((-(a*(2*a*A*b - a^2*B + b^2*B)*x)/(a^2 + b^2)) + (A*(a^2 + b^2)*Log[-Sin[c + d*x]])/(a*d) - (b*(3*a^2*A*b + A*b^3 - 2*a^3*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))/(a*(a^2 + b^2)) + (b*(A*b - a*B))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))`

3.279.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

3.279. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4134 `Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2/(((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.279.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.19

method	result
derivativedivides	$\frac{\frac{(-A a^2 + A b^2 - 2Bab) \ln(1 + \tan^2(dx+c))}{2} + \frac{(-2Aab + B a^2 - B b^2) \arctan(\tan(dx+c))}{(a^2+b^2)^2} + \frac{A \ln(\tan(dx+c))}{a^2} - \frac{b(3A a^2 b + A b^3 - 2B a^3) \ln(1 + \tan^2(dx+c))}{(a^2+b^2)^2}}{d}$
default	$\frac{\frac{(-A a^2 + A b^2 - 2Bab) \ln(1 + \tan^2(dx+c))}{2} + \frac{(-2Aab + B a^2 - B b^2) \arctan(\tan(dx+c))}{(a^2+b^2)^2} + \frac{A \ln(\tan(dx+c))}{a^2} - \frac{b(3A a^2 b + A b^3 - 2B a^3) \ln(1 + \tan^2(dx+c))}{(a^2+b^2)^2}}{d}$
parallelrisch	$-6(A a^2 b + \frac{1}{3} A b^3 - \frac{2}{3} B a^3) b(a+b \tan(dx+c)) \ln(a+b \tan(dx+c)) - a^2(a+b \tan(dx+c))(A a^2 - A b^2 + 2Bab) \ln(\sec^2(dx+c))$
norman	$\frac{\frac{a(2Aab - B a^2 + B b^2)x}{a^4 + 2a^2b^2 + b^4} - \frac{b(2Aab - B a^2 + B b^2)x \tan(dx+c)}{a^4 + 2a^2b^2 + b^4} - \frac{(A b^2 - Bab) b \tan(dx+c)}{d a^2 (a^2 + b^2)}}{a+b \tan(dx+c)} + \frac{A \ln(\tan(dx+c))}{a^2 d} - \frac{(A a^2 - A b^2 + 2Bab) \ln(\sec^2(dx+c))}{2d}$
risch	$-\frac{x B}{2i b a - a^2 + b^2} + \frac{2i b^4 A c}{a^2 d (a^4 + 2a^2 b^2 + b^4)} + \frac{6i A b^2 x}{a^4 + 2a^2 b^2 + b^4} - \frac{i x A}{2i b a - a^2 + b^2} + \frac{6i A b^2 c}{d (a^4 + 2a^2 b^2 + b^4)} - \frac{2i x A}{a^2} + \frac{2i b^4 A c}{a^2 (a^4 + 2a^2 b^2 + b^4)}$

input `int(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(1/(a^2+b^2)^2*(1/2*(-A*a^2+A*b^2-2*B*a*b)*ln(1+tan(d*x+c)^2)+(-2*A*a*b+B*a^2-B*b^2)*arctan(tan(d*x+c)))+1/a^2*A*ln(tan(d*x+c))-b*(3*A*a^2*b+A*b^3-2*B*a^3)/(a^2+b^2)^2/a^2*ln(a+b*tan(d*x+c))+(A*b-B*a)*b/(a^2+b^2)/a/(a+b*tan(d*x+c))`

3.279.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 323 vs. 2(137) = 274.

Time = 0.30 (sec) , antiderivative size = 323, normalized size of antiderivative = 2.36

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \frac{2 B a^2 b^3 - 2 A a b^4 - 2 (B a^5 - 2 A a^4 b - B a^3 b^2) dx - (A a^5 + 2 A a^3 b^2 + A a b^4 + (A a^4 b + 2 A a^2 b^3 + A b^5) \tan(c+dx))}{(a+b \tan(c+dx))^2}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

3.279.
$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

```
output -1/2*(2*B*a^2*b^3 - 2*A*a*b^4 - 2*(B*a^5 - 2*A*a^4*b - B*a^3*b^2)*d*x - (A
*a^5 + 2*A*a^3*b^2 + A*a*b^4 + (A*a^4*b + 2*A*a^2*b^3 + A*b^5)*tan(d*x + c
))*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1)) - (2*B*a^4*b - 3*A*a^3*b^2 - A
*a*b^4 + (2*B*a^3*b^2 - 3*A*a^2*b^3 - A*b^5)*tan(d*x + c))*log((b^2*tan(d*
x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - 2*(B*a^3*b^2
- A*a^2*b^3 + (B*a^4*b - 2*A*a^3*b^2 - B*a^2*b^3)*d*x)*tan(d*x + c)/((a^6
*b + 2*a^4*b^3 + a^2*b^5)*d*tan(d*x + c) + (a^7 + 2*a^5*b^2 + a^3*b^4)*d)
```

3.279.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.80 (sec) , antiderivative size = 4488, normalized size of antiderivative = 32.76

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)
```

```
output Piecewise((zoo*x*(A + B*tan(c))*cot(c)/tan(c)**2, Eq(a, 0) & Eq(b, 0) & Eq
(d, 0)), ((-A*log(tan(c + d*x)**2 + 1)/(2*d) + A*log(tan(c + d*x))/d + B*x
)/a**2, Eq(b, 0)), ((A*log(tan(c + d*x)**2 + 1)/(2*d) - A*log(tan(c + d*x)
))/d - A/(2*d*tan(c + d*x)**2) - B*x - B/(d*tan(c + d*x)))/b**2, Eq(a, 0)),
(3*I*A*d*x*tan(c + d*x)**2/(4*a**2*d*tan(c + d*x)**2 + 8*I*a**2*d*tan(c +
d*x) - 4*a**2*d) - 6*A*d*x*tan(c + d*x)/(4*a**2*d*tan(c + d*x)**2 + 8*I*a
**2*d*tan(c + d*x) - 4*a**2*d) - 3*I*A*d*x/(4*a**2*d*tan(c + d*x)**2 + 8*I
*a**2*d*tan(c + d*x) - 4*a**2*d) - 2*A*log(tan(c + d*x)**2 + 1)*tan(c + d*
x)**2/(4*a**2*d*tan(c + d*x)**2 + 8*I*a**2*d*tan(c + d*x) - 4*a**2*d) - 4*
I*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(4*a**2*d*tan(c + d*x)**2 + 8*I*
a**2*d*tan(c + d*x) - 4*a**2*d) + 2*A*log(tan(c + d*x)**2 + 1)/(4*a**2*d*t
an(c + d*x)**2 + 8*I*a**2*d*tan(c + d*x) - 4*a**2*d) + 4*A*log(tan(c + d*x
))*tan(c + d*x)**2/(4*a**2*d*tan(c + d*x)**2 + 8*I*a**2*d*tan(c + d*x) - 4
*a**2*d) + 8*I*A*log(tan(c + d*x))*tan(c + d*x)/(4*a**2*d*tan(c + d*x)**2
+ 8*I*a**2*d*tan(c + d*x) - 4*a**2*d) - 4*A*log(tan(c + d*x))/(4*a**2*d*ta
n(c + d*x)**2 + 8*I*a**2*d*tan(c + d*x) - 4*a**2*d) + 3*I*A*tan(c + d*x)/(
4*a**2*d*tan(c + d*x)**2 + 8*I*a**2*d*tan(c + d*x) - 4*a**2*d) - 4*A/(4*a*
**2*d*tan(c + d*x)**2 + 8*I*a**2*d*tan(c + d*x) - 4*a**2*d) + B*d*x*tan(c +
d*x)**2/(4*a**2*d*tan(c + d*x)**2 + 8*I*a**2*d*tan(c + d*x) - 4*a**2*d) +
2*I*B*d*x*tan(c + d*x)/(4*a**2*d*tan(c + d*x)**2 + 8*I*a**2*d*tan(c + ...
```

3.279.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.52

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{2(Ba^2-2Aab-Bb^2)(dx+c)}{a^4+2a^2b^2+b^4} + \frac{2(2Ba^3b-3Aa^2b^2-Ab^4) \log(b \tan(dx+c)+a)}{a^6+2a^4b^2+a^2b^4} - \frac{(Aa^2+2Bab-Ab^2) \log(\tan(dx+c)^2+1)}{a^4+2a^2b^2+b^4} - \frac{2(Ba^2-2Aab-Bb^2)}{a^4+a^2b^2+(a^3b^2+c^2)}$$

$$2d$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `1/2*(2*(B*a^2 - 2*A*a*b - B*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) + 2*(2*B*a^3*b - 3*A*a^2*b^2 - A*b^4)*log(b*tan(d*x + c) + a)/(a^6 + 2*a^4*b^2 + a^2*b^4) - (A*a^2 + 2*B*a*b - A*b^2)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) - 2*(B*a*b - A*b^2)/(a^4 + a^2*b^2 + (a^3*b + a*b^3)*tan(d*x + c)) + 2*A*log(tan(d*x + c))/a^2)/d`

3.279.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 279 vs. 2(137) = 274.

Time = 0.66 (sec) , antiderivative size = 279, normalized size of antiderivative = 2.04

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{2(Ba^2-2Aab-Bb^2)(dx+c)}{a^4+2a^2b^2+b^4} - \frac{(Aa^2+2Bab-Ab^2) \log(\tan(dx+c)^2+1)}{a^4+2a^2b^2+b^4} + \frac{2(2Ba^3b^2-3Aa^2b^3-Ab^5) \log(|b \tan(dx+c)+a|)}{a^6b+2a^4b^3+a^2b^5} + \frac{2A \log(|\tan(dx+c)|)}{a^2}$$

$$2d$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output `1/2*(2*(B*a^2 - 2*A*a*b - B*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) - (A*a^2 + 2*B*a*b - A*b^2)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) + 2*(2*B*a^3*b^2 - 3*A*a^2*b^3 - A*b^5)*log(abs(b*tan(d*x + c) + a))/(a^6*b + 2*a^4*b^3 + a^2*b^5) + 2*A*log(abs(tan(d*x + c)))/a^2 - 2*(2*B*a^3*b^2*tan(d*x + c) - 3*A*a^2*b^3*tan(d*x + c) - A*b^5*tan(d*x + c) + 3*B*a^4*b - 4*A*a^3*b^2 + B*a^2*b^3 - 2*A*a*b^4)/((a^6 + 2*a^4*b^2 + a^2*b^4)*(b*tan(d*x + c) + a)))/d`

3.279. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.279.9 Mupad [B] (verification not implemented)

Time = 9.16 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.31

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{A \ln(\tan(c+dx))}{a^2 d} - \frac{\ln(\tan(c+dx) - i)(A + B i)}{2 d (a^2 + a b 2i - b^2)}$$

$$- \frac{\ln(\tan(c+dx) + i)(B + A i)}{2 d (a^2 i + 2 a b - b^2 i)} + \frac{A b^2 - B a b}{a d (a^2 + b^2)(a + b \tan(c+dx))}$$

$$- \frac{b \ln(a + b \tan(c+dx))(-2 B a^3 + 3 A a^2 b + A b^3)}{a^2 d (a^2 + b^2)^2}$$

input `int((cot(c + d*x)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`output `(A*log(tan(c + d*x)))/(a^2*d) - (log(tan(c + d*x) - 1i)*(A + B*1i))/(2*d*(a*b*2i + a^2 - b^2)) - (log(tan(c + d*x) + 1i)*(A*1i + B))/(2*d*(2*a*b + a^2*1i - b^2*1i)) + (A*b^2 - B*a*b)/(a*d*(a^2 + b^2)*(a + b*tan(c + d*x))) - (b*log(a + b*tan(c + d*x))*(A*b^3 - 2*B*a^3 + 3*A*a^2*b))/(a^2*d*(a^2 + b^2)^2)`

3.280 $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.280.1 Optimal result 2689
 3.280.2 Mathematica [C] (verified) 2690
 3.280.3 Rubi [A] (verified) 2690
 3.280.4 Maple [A] (verified) 2694
 3.280.5 Fricas [B] (verification not implemented) 2694
 3.280.6 Sympy [C] (verification not implemented) 2695
 3.280.7 Maxima [A] (verification not implemented) 2696
 3.280.8 Giac [A] (verification not implemented) 2697
 3.280.9 Mupad [B] (verification not implemented) 2697

3.280.1 Optimal result

Integrand size = 31, antiderivative size = 192

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= -\frac{(a^2A - Ab^2 + 2abB)x}{(a^2 + b^2)^2} - \frac{(2Ab - aB) \log(\sin(c+dx))}{a^3d}$$

$$+ \frac{b^2(4a^2Ab + 2Ab^3 - 3a^3B - ab^2B) \log(a \cos(c+dx) + b \sin(c+dx))}{a^3(a^2 + b^2)^2d}$$

$$- \frac{b(a^2A + 2Ab^2 - abB)}{a^2(a^2 + b^2)d(a+b \tan(c+dx))} - \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))}$$

output

```
-(A*a^2-A*b^2+2*B*a*b)*x/(a^2+b^2)^2-(2*A*b-B*a)*ln(sin(d*x+c))/a^3/d+b^2*(4*A*a^2*b+2*A*b^3-3*B*a^3-B*a*b^2)*ln(a*cos(d*x+c)+b*sin(d*x+c))/a^3/(a^2+b^2)^2/d-b*(A*a^2+2*A*b^2-B*a*b)/a^2/(a^2+b^2)/d/(a+b*tan(d*x+c))-A*cot(d*x+c)/a/d/(a+b*tan(d*x+c))
```


3.280.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.77 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.01

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{-\frac{2A \cot(c+dx)}{a^2} + \frac{i(A+iB) \log(i-\tan(c+dx))}{(a+ib)^2} + \frac{2(-2Ab+aB) \log(\tan(c+dx))}{a^3} - \frac{(iA+B) \log(i+\tan(c+dx))}{(a-ib)^2} - \frac{2b^2(-4a^2Ab-2Ab^3+3a^3B)}{a^3(c+dx)}}{2d}$$

input `Integrate[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `((-2*A*Cot[c + d*x])/a^2 + (I*(A + I*B)*Log[I - Tan[c + d*x]])/(a + I*b)^2 + (2*(-2*A*b + a*B)*Log[Tan[c + d*x]])/a^3 - ((I*A + B)*Log[I + Tan[c + d*x]])/(a - I*b)^2 - (2*b^2*(-4*a^2*A*b - 2*A*b^3 + 3*a^3*B + a*b^2*B)*Log[a + b*Tan[c + d*x]])/(a^3*(a^2 + b^2)^2) + (2*b^2*(-(A*b) + a*B))/(a^2*(a^2 + b^2)*(a + b*Tan[c + d*x])))/(2*d)`

3.280.3 Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.16, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 4092, 3042, 4132, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^2(a + b \tan(c + dx))^2} dx$$

↓ 4092

$$-\frac{\int \frac{\cot(c+dx)(2Ab \tan^2(c+dx)+aA \tan(c+dx)+2Ab-aB)}{(a+b \tan(c+dx))^2} dx}{a} - \frac{A \cot(c + dx)}{ad(a + b \tan(c + dx))}$$

↓ 3042

3.280. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\begin{aligned}
& \frac{\int \frac{2Ab \tan(c+dx)^2 + aA \tan(c+dx) + 2Ab - aB}{\tan(c+dx)(a+b \tan(c+dx))^2} dx}{a} - \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))} \\
& \quad \downarrow 4132 \\
& \frac{\int \frac{\cot(c+dx) \left((aA+bB) \tan(c+dx)a^2 + b(Aa^2 - bBa + 2Ab^2) \tan^2(c+dx) + (a^2+b^2)(2Ab-aB) \right)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{b(a^2A - abB + 2Ab^2)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
& \quad \downarrow 3042 \\
& \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))} \\
& \quad \downarrow 4134 \\
& \frac{\int \frac{(aA+bB) \tan(c+dx)a^2 + b(Aa^2 - bBa + 2Ab^2) \tan(c+dx)^2 + (a^2+b^2)(2Ab-aB)}{\tan(c+dx)(a+b \tan(c+dx))} dx}{a(a^2+b^2)} + \frac{b(a^2A - abB + 2Ab^2)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
& \quad \downarrow 4134 \\
& \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))} \\
& \quad \downarrow 3042 \\
& \frac{(a^2+b^2)(2Ab-aB) \int \cot(c+dx) dx}{a} - \frac{b^2(-3a^3B + 4a^2Ab - ab^2B + 2Ab^3) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^2x(a^2A + 2abB - Ab^2)}{a^2+b^2} + \frac{b(a^2A - abB + 2Ab^2)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
& \quad \downarrow 3042 \\
& \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))} \\
& \quad \downarrow 25 \\
& \frac{(a^2+b^2)(2Ab-aB) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx}{a} - \frac{b^2(-3a^3B + 4a^2Ab - ab^2B + 2Ab^3) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^2x(a^2A + 2abB - Ab^2)}{a^2+b^2} + \frac{b(a^2A - abB + 2Ab^2)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
& \quad \downarrow 3956 \\
& \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))} \\
& \quad \downarrow 3956 \\
& \frac{(a^2+b^2)(2Ab-aB) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx}{a} - \frac{b^2(-3a^3B + 4a^2Ab - ab^2B + 2Ab^3) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^2x(a^2A + 2abB - Ab^2)}{a^2+b^2} + \frac{b(a^2A - abB + 2Ab^2)}{ad(a^2+b^2)(a+b \tan(c+dx))}
\end{aligned}$$

3.280. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\begin{aligned}
 & -\frac{b^2(-3a^3B+4a^2Ab-ab^2B+2Ab^3) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{(a^2+b^2)(2Ab-aB) \log(-\sin(c+dx))}{ad} + \frac{a^2x(a^2A+2abB-Ab^2)}{a^2+b^2} \\
 & + \frac{b(a^2A-abB+2Ab^2)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
 & - \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{4013} \\
 & -\frac{b(a^2A-abB+2Ab^2)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \frac{(a^2+b^2)(2Ab-aB) \log(-\sin(c+dx))}{ad} + \frac{a^2x(a^2A+2abB-Ab^2)}{a^2+b^2} - \frac{b^2(-3a^3B+4a^2Ab-ab^2B+2Ab^3) \log(a \cos(c+dx)+b \sin(c+dx))}{ad(a^2+b^2)} \\
 & - \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))}
 \end{aligned}$$

input `Int[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `-((A*Cot[c + d*x])/(a*d*(a + b*Tan[c + d*x]))) - (((a^2*(a^2*A - A*b^2 + 2*a*b*B)*x)/(a^2 + b^2) + ((a^2 + b^2)*(2*A*b - a*B)*Log[-Sin[c + d*x]])/(a*d) - (b^2*(4*a^2*A*b + 2*A*b^3 - 3*a^3*B - a*b^2*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))/(a*(a^2 + b^2)) + (b*(a^2*A + 2*A*b^2 - a*b*B))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))/a`

3.280.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

3.280. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4134 `Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/(a^2 + b^2)*(c^2 + d^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.280.4 Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.02

method	result
derivativedivides	$\frac{\frac{(2Aab - B a^2 + B b^2) \ln(1 + \tan^2(dx+c))}{2} + (-A a^2 + A b^2 - 2Bab) \arctan(\tan(dx+c))}{(a^2 + b^2)^2} - \frac{A}{a^2 \tan(dx+c)} + \frac{(-2Ab + Ba) \ln(\tan(dx+c))}{a^3} + \dots$
default	$\frac{\frac{(2Aab - B a^2 + B b^2) \ln(1 + \tan^2(dx+c))}{2} + (-A a^2 + A b^2 - 2Bab) \arctan(\tan(dx+c))}{(a^2 + b^2)^2} - \frac{A}{a^2 \tan(dx+c)} + \frac{(-2Ab + Ba) \ln(\tan(dx+c))}{a^3} + \dots$
parallelrisch	$4(A a^2 b + \frac{1}{2} A b^3 - \frac{3}{4} B a^3 - \frac{1}{4} B a b^2) b^2 (a + b \tan(dx+c)) \ln(a + b \tan(dx+c)) + a^3 (a + b \tan(dx+c)) (Aab - \frac{1}{2} B a^2 + \frac{1}{2} B b^2) \ln(\dots)$
norman	$\frac{(A a^2 b + 2A b^3 - B a b^2) b (\tan^2(dx+c))}{d a^3 (a^2 + b^2)} - \frac{A}{ad} - \frac{a(A a^2 - A b^2 + 2Bab) x \tan(dx+c)}{a^4 + 2a^2 b^2 + b^4} - \frac{b(A a^2 - A b^2 + 2Bab) x (\tan^2(dx+c))}{a^4 + 2a^2 b^2 + b^4} + \frac{b^2(4A \dots)}{\tan(dx+c)(a + b \tan(dx+c))}$
risch	$-\frac{2iBc}{a^2 d} + \frac{x A}{2iba - a^2 + b^2} - \frac{4ib^5 Ax}{(a^4 + 2a^2 b^2 + b^4) a^3} - \frac{2i(A a^4 e^{2i(dx+c)} - 2A b^4 e^{2i(dx+c)} + B a b^3 e^{2i(dx+c)} - 2iA a^3 b e^{2i(dx+c)} - \dots)}{(e^{2i(dx+c)} - 1)(ib+a)(-ib+a)^2 (-ib e^{2i(dx+c)} - \dots)}$

input `int(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(1/(a^2+b^2)^2*(1/2*(2*A*a*b-B*a^2+B*b^2)*ln(1+tan(d*x+c)^2)+(-A*a^2+A*b^2-2*B*a*b)*arctan(tan(d*x+c)))-1/a^2*A/tan(d*x+c)+(-2*A*b+B*a)/a^3*ln(tan(d*x+c))+b^2*(4*A*a^2*b+2*A*b^3-3*B*a^3-B*a*b^2)/(a^2+b^2)^2/a^3*ln(a+b*tan(d*x+c))-(A*b-B*a)*b^2/(a^2+b^2)/a^2/(a+b*tan(d*x+c))`

3.280.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 465 vs. 2(190) = 380.

Time = 0.32 (sec) , antiderivative size = 465, normalized size of antiderivative = 2.42

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx = \frac{2Aa^6 + 4Aa^4b^2 + 2Aa^2b^4 + 2(Ba^3b^3 - Aa^2b^4 + (Aa^5b + 2Ba^4b^2 - Aa^3b^3)dx) \tan(dx + c)^2 - ((Ba^5b^3 - \dots))}{\dots}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

3.280.
$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

output

```
-1/2*(2*A*a^6 + 4*A*a^4*b^2 + 2*A*a^2*b^4 + 2*(B*a^3*b^3 - A*a^2*b^4 + (A*
a^5*b + 2*B*a^4*b^2 - A*a^3*b^3)*d*x)*tan(d*x + c)^2 - ((B*a^5*b - 2*A*a^4
*b^2 + 2*B*a^3*b^3 - 4*A*a^2*b^4 + B*a*b^5 - 2*A*b^6)*tan(d*x + c)^2 + (B*
a^6 - 2*A*a^5*b + 2*B*a^4*b^2 - 4*A*a^3*b^3 + B*a^2*b^4 - 2*A*a*b^5)*tan(d
*x + c))*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1)) + ((3*B*a^3*b^3 - 4*A*a^
2*b^4 + B*a*b^5 - 2*A*b^6)*tan(d*x + c)^2 + (3*B*a^4*b^2 - 4*A*a^3*b^3 + B
*a^2*b^4 - 2*A*a*b^5)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*
x + c) + a^2)/(tan(d*x + c)^2 + 1)) + 2*(A*a^5*b + 2*A*a^3*b^3 - B*a^2*b^4
+ 2*A*a*b^5 + (A*a^6 + 2*B*a^5*b - A*a^4*b^2)*d*x)*tan(d*x + c))/((a^7*b
+ 2*a^5*b^3 + a^3*b^5)*d*tan(d*x + c)^2 + (a^8 + 2*a^6*b^2 + a^4*b^4)*d*ta
n(d*x + c))
```

3.280.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.92 (sec) , antiderivative size = 8145, normalized size of antiderivative = 42.42

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)**2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Piecewise(((-A*x - A/(d*tan(c + d*x)) - B*log(tan(c + d*x)**2 + 1)/(2*d) + B*log(tan(c + d*x))/d)/a**2, Eq(b, 0)), ((A*x + A/(d*tan(c + d*x)) - A/(3*d*tan(c + d*x)**3) + B*log(tan(c + d*x)**2 + 1)/(2*d) - B*log(tan(c + d*x)))/d - B/(2*d*tan(c + d*x)**2))/b**2, Eq(a, 0)), (-9*A*d*x*tan(c + d*x)**3/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x)) - 18*I*A*d*x*tan(c + d*x)**2/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x)) + 9*A*d*x*tan(c + d*x)/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x)) - 4*I*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**3/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x)) + 8*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x)) + 4*I*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x)) + 8*I*A*log(tan(c + d*x))*tan(c + d*x)**3/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x)) - 16*A*log(tan(c + d*x))*tan(c + d*x)**2/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x)) - 8*I*A*log(tan(c + d*x))*tan(c + d*x)/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x)) - 9*A*tan(c + d*x)**2/(4*a**2*d*tan(c + d*x)**3 + 8*I*a**2*d*tan(c + d*x)**2 - 4*a**2*d*tan(c + d*x)) - 14*I*A*tan(c + d*x)/(4*a**2*d*tan(c + d...`

3.280.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.36

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \frac{2(Aa^2+2Bab-Ab^2)(dx+c)}{a^4+2a^2b^2+b^4} + \frac{2(3Ba^3b^2-4Aa^2b^3+Bab^4-2Ab^5) \log(b \tan(dx+c)+a)}{a^7+2a^5b^2+a^3b^4} + \frac{(Ba^2-2Aab-Bb^2) \log(\tan(dx+c)^2+1)}{a^4+2a^2b^2+b^4} + \frac{2}{2d}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `-1/2*(2*(A*a^2 + 2*B*a*b - A*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) + 2*(3*B*a^3*b^2 - 4*A*a^2*b^3 + B*a*b^4 - 2*A*b^5)*log(b*tan(d*x + c) + a)/(a^7 + 2*a^5*b^2 + a^3*b^4) + (B*a^2 - 2*A*a*b - B*b^2)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) + 2*(A*a^3 + A*a*b^2 + (A*a^2*b - B*a*b^2 + 2*A*b^3)*tan(d*x + c))/((a^4*b + a^2*b^3)*tan(d*x + c)^2 + (a^5 + a^3*b^2)*tan(d*x + c)) - 2*(B*a - 2*A*b)*log(tan(d*x + c))/a^3)/d`

3.280. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.280.8 Giac [A] (verification not implemented)

Time = 0.87 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.89

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx =$$

$$\frac{2(Aa^2+2Bab-Ab^2)(dx+c)}{a^4+2a^2b^2+b^4} + \frac{(Ba^2-2Aab-Bb^2)\log(\tan(dx+c)^2+1)}{a^4+2a^2b^2+b^4} + \frac{2(3Ba^3b^3-4Aa^2b^4+Bab^5-2Ab^6)\log(|b\tan(dx+c)+a|)}{a^7b+2a^5b^3+a^3b^5} +$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output
$$-1/2*(2*(A*a^2 + 2*B*a*b - A*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) + (B*a^2 - 2*A*a*b - B*b^2)*\log(\tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) + 2*(3*B*a^3*b^3 - 4*A*a^2*b^4 + B*a*b^5 - 2*A*b^6)*\log(\text{abs}(b*\tan(d*x + c) + a)))/(a^7*b + 2*a^5*b^3 + a^3*b^5) + (B*a^4*b*\tan(d*x + c)^2 - 2*A*a^3*b^2*\tan(d*x + c)^2 - B*a^2*b^3*\tan(d*x + c)^2 + B*a^5*\tan(d*x + c) - 3*B*a^3*b^2*\tan(d*x + c) + 6*A*a^2*b^3*\tan(d*x + c) - 2*B*a*b^4*\tan(d*x + c) + 4*A*b^5*\tan(d*x + c) + 2*A*a^5 + 4*A*a^3*b^2 + 2*A*a*b^4)/((a^6 + 2*a^4*b^2 + a^2*b^4)*(b*\tan(d*x + c)^2 + a*\tan(d*x + c))) - 2*(B*a - 2*A*b)*\log(\text{abs}(\tan(d*x + c)))/a^3)/d$$

3.280.9 Mupad [B] (verification not implemented)

Time = 10.85 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.20

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{b^2 \ln(a+b\tan(c+dx))(-3Ba^3+4Aa^2b-Bab^2+2Ab^3)}{a^3d(a^2+b^2)^2}$$

$$- \frac{\ln(\tan(c+dx))(2Ab-Ba)}{a^3d} + \frac{\ln(\tan(c+dx)+i)(B+Ai)}{2d(-a^2+ab2i+b^2)}$$

$$+ \frac{\ln(\tan(c+dx)-i)(A+Bli)}{2d(-a^2li+2ab+b^2li)} - \frac{\frac{A}{a} + \frac{\tan(c+dx)(Aa^2b-Bab^2+2Ab^3)}{a^2(a^2+b^2)}}{d(b\tan(c+dx)^2+a\tan(c+dx))}$$

input `int((cot(c+d*x))^2*(A+B*tan(c+d*x)))/(a+b*tan(c+d*x))^2,x)`

output

```
(log(tan(c + d*x) + 1i)*(A*1i + B))/(2*d*(a*b*2i - a^2 + b^2)) - (log(tan(
c + d*x))*(2*A*b - B*a))/(a^3*d) - (A/a + (tan(c + d*x)*(2*A*b^3 + A*a^2*b
- B*a*b^2))/(a^2*(a^2 + b^2)))/(d*(a*tan(c + d*x) + b*tan(c + d*x)^2)) +
(log(tan(c + d*x) - 1i)*(A + B*1i))/(2*d*(2*a*b - a^2*1i + b^2*1i)) + (b^2
*log(a + b*tan(c + d*x))*(2*A*b^3 - 3*B*a^3 + 4*A*a^2*b - B*a*b^2))/(a^3*d
*(a^2 + b^2)^2)
```

3.281 $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.281.1 Optimal result 2699
 3.281.2 Mathematica [C] (verified) 2700
 3.281.3 Rubi [A] (verified) 2700
 3.281.4 Maple [A] (verified) 2705
 3.281.5 Fricas [B] (verification not implemented) 2705
 3.281.6 Sympy [C] (verification not implemented) 2706
 3.281.7 Maxima [A] (verification not implemented) 2707
 3.281.8 Giac [A] (verification not implemented) 2708
 3.281.9 Mupad [B] (verification not implemented) 2709

3.281.1 Optimal result

Integrand size = 31, antiderivative size = 250

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{(2aAb - a^2B + b^2B)x}{(a^2 + b^2)^2} - \frac{(a^2A - 3Ab^2 + 2abB) \log(\sin(c+dx))}{a^4d}$$

$$- \frac{b^3(5a^2Ab + 3Ab^3 - 4a^3B - 2ab^2B) \log(a \cos(c+dx) + b \sin(c+dx))}{a^4(a^2 + b^2)^2d}$$

$$+ \frac{b(2a^2Ab + 3Ab^3 - a^3B - 2ab^2B)}{a^3(a^2 + b^2)d(a+b \tan(c+dx))} + \frac{(3Ab - 2aB) \cot(c+dx)}{2a^2d(a+b \tan(c+dx))} - \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))}$$

```
output (2*A*a*b-B*a^2+B*b^2)*x/(a^2+b^2)^2-(A*a^2-3*A*b^2+2*B*a*b)*ln(sin(d*x+c))
/a^4/d-b^3*(5*A*a^2*b+3*A*b^3-4*B*a^3-2*B*a*b^2)*ln(a*cos(d*x+c)+b*sin(d*x
+c))/a^4/(a^2+b^2)^2/d+b*(2*A*a^2*b+3*A*b^3-B*a^3-2*B*a*b^2)/a^3/(a^2+b^2)
/d/(a+b*tan(d*x+c))+1/2*(3*A*b-2*B*a)*cot(d*x+c)/a^2/d/(a+b*tan(d*x+c))-1/
2*A*cot(d*x+c)^2/a/d/(a+b*tan(d*x+c))
```

3.281.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.96 (sec) , antiderivative size = 220, normalized size of antiderivative = 0.88

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{-\frac{2(-2Ab+aB) \cot(c+dx)}{a^3} - \frac{A \cot^2(c+dx)}{a^2} + \frac{(A+iB) \log(i-\tan(c+dx))}{(a+ib)^2} - \frac{2(a^2A-3Ab^2+2abB) \log(\tan(c+dx))}{a^4} + \frac{(A-iB) \log(i+\tan(c+dx))}{(a-ib)^2}}{2d}$$

input `Integrate[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `((-2*(-2*A*b + a*B)*Cot[c + d*x])/a^3 - (A*Cot[c + d*x]^2)/a^2 + ((A + I*B)*Log[I - Tan[c + d*x]])/(a + I*b)^2 - (2*(a^2*A - 3*A*b^2 + 2*a*b*B)*Log[Tan[c + d*x]])/a^4 + ((A - I*B)*Log[I + Tan[c + d*x]])/(a - I*b)^2 + (2*b^3*(-5*a^2*A*b - 3*A*b^3 + 4*a^3*B + 2*a*b^2*B)*Log[a + b*Tan[c + d*x]])/(a^4*(a^2 + b^2)^2) + (2*b^3*(A*b - a*B))/(a^3*(a^2 + b^2)*(a + b*Tan[c + d*x])))/(2*d)`

3.281.3 Rubi [A] (verified)

Time = 1.70 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.15, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 4092, 3042, 4132, 27, 3042, 4132, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

↓ 3042

$$\int \frac{A+B \tan(c+dx)}{\tan(c+dx)^3(a+b \tan(c+dx))^2} dx$$

↓ 4092

$$\frac{\int \frac{\cot^2(c+dx)(3Ab \tan^2(c+dx)+2aA \tan(c+dx)+3Ab-2aB)}{(a+b \tan(c+dx))^2} dx}{2a} - \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))}$$

↓ 3042

3.281. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\begin{aligned}
 & \frac{\int \frac{3Ab \tan(c+dx)^2 + 2aA \tan(c+dx) + 3Ab - 2aB}{\tan(c+dx)^2 (a+b \tan(c+dx))^2} dx}{2a} - \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))} \\
 & \quad \downarrow 4132 \\
 & \frac{\int -\frac{2 \cot(c+dx) (Aa^2 + B \tan(c+dx)a^2 + 2bBa - 3Ab^2 - b(3Ab - 2aB) \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx}{a} - \frac{(3Ab - 2aB) \cot(c+dx)}{ad(a+b \tan(c+dx))} \\
 & \quad \frac{2a}{2ad(a+b \tan(c+dx))} \\
 & \quad \downarrow 27 \\
 & \frac{2 \int \frac{\cot(c+dx) (Aa^2 + B \tan(c+dx)a^2 + 2bBa - 3Ab^2 - b(3Ab - 2aB) \tan^2(c+dx))}{(a+b \tan(c+dx))^2} dx}{a} - \frac{(3Ab - 2aB) \cot(c+dx)}{ad(a+b \tan(c+dx))} \\
 & \quad \frac{2a}{2ad(a+b \tan(c+dx))} \\
 & \quad \downarrow 3042 \\
 & \frac{2 \int \frac{Aa^2 + B \tan(c+dx)a^2 + 2bBa - 3Ab^2 - b(3Ab - 2aB) \tan(c+dx)^2}{\tan(c+dx)(a+b \tan(c+dx))^2} dx}{a} - \frac{(3Ab - 2aB) \cot(c+dx)}{ad(a+b \tan(c+dx))} - \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))} \\
 & \quad \downarrow 4132 \\
 & \frac{2 \left(\int \frac{\cot(c+dx) \left(-((Ab - aB) \tan(c+dx)a^3) - b(-Ba^3 + 2Aba^2 - 2b^2Ba + 3Ab^3) \tan^2(c+dx) + (a^2 + b^2)(Aa^2 + 2bBa - 3Ab^2) \right)}{a+b \tan(c+dx)} dx}{a(a^2 + b^2)} - \frac{b(a^3(-B) + 2a^2Ab - 2ab^2B + 3Ab^3)}{ad(a^2 + b^2)(a+b \tan(c+dx))} \right)}{a} \\
 & \quad \frac{2a}{2ad(a+b \tan(c+dx))} \\
 & \quad \downarrow 3042 \\
 & \frac{2 \left(\int \frac{-((Ab - aB) \tan(c+dx)a^3) - b(-Ba^3 + 2Aba^2 - 2b^2Ba + 3Ab^3) \tan(c+dx)^2 + (a^2 + b^2)(Aa^2 + 2bBa - 3Ab^2)}{\tan(c+dx)(a+b \tan(c+dx))} dx - \frac{b(a^3(-B) + 2a^2Ab - 2ab^2B + 3Ab^3)}{ad(a^2 + b^2)(a+b \tan(c+dx))} \right)}{a} \\
 & \quad \frac{2a}{2ad(a+b \tan(c+dx))} \\
 & \quad \downarrow 4134
 \end{aligned}$$

3.281. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$2 \left(\frac{\frac{(a^2+b^2)(a^2A+2abB-3Ab^2)}{a} \int \cot(c+dx) dx + \frac{b^3(-4a^3B+5a^2Ab-2ab^2B+3Ab^3)}{a(a^2+b^2)} \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx - \frac{a^3x(a^2(-B)+2aAb+b^2B)}{a^2+b^2} \right) - \frac{b(a^3(-B)+2a^2Ab-)}{ad(a^2+b^2)(a+b)}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))}$$

↓ 3042

$$2 \left(\frac{\frac{(a^2+b^2)(a^2A+2abB-3Ab^2)}{a} \int -\tan(c+dx+\frac{\pi}{2}) dx + \frac{b^3(-4a^3B+5a^2Ab-2ab^2B+3Ab^3)}{a(a^2+b^2)} \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx - \frac{a^3x(a^2(-B)+2aAb+b^2B)}{a^2+b^2} \right) - \frac{b(a^3(-B)+2a^2Ab-)}{ad(a^2+b^2)}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))}$$

↓ 25

$$2 \left(\frac{\frac{(a^2+b^2)(a^2A+2abB-3Ab^2)}{a} \int \tan(\frac{1}{2}(2c+\pi)+dx) dx + \frac{b^3(-4a^3B+5a^2Ab-2ab^2B+3Ab^3)}{a(a^2+b^2)} \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx - \frac{a^3x(a^2(-B)+2aAb+b^2B)}{a^2+b^2} \right) - \frac{b(a^3(-B)+2a^2Ab-)}{ad(a^2+b^2)}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))}$$

↓ 3956

$$2 \left(\frac{\frac{b^3(-4a^3B+5a^2Ab-2ab^2B+3Ab^3)}{a(a^2+b^2)} \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx + \frac{(a^2+b^2)(a^2A+2abB-3Ab^2)}{ad} \log(-\sin(c+dx)) - \frac{a^3x(a^2(-B)+2aAb+b^2B)}{a^2+b^2} \right) - \frac{b(a^3(-B)+2a^2Ab-)}{ad(a^2+b^2)(a+b)}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))}$$

↓ 4013

3.281. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\frac{2 \left(\frac{(a^2+b^2)(a^2A+2abB-3Ab^2) \log(-\sin(c+dx))}{ad} - \frac{a^3x(a^2(-B)+2aAb+b^2B)}{a^2+b^2} + \frac{b^3(-4a^3B+5a^2Ab-2ab^2B+3Ab^3) \log(a \cos(c+dx)+b \sin(c+dx))}{ad(a^2+b^2)} - \frac{b(a^3(-B)+2aAb+b^2B)}{ad(a^2+b^2)} \right)}{a(a^2+b^2)} = \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))}$$

```
input Int[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]
```

```
output -1/2*(A*Cot[c + d*x]^2)/(a*d*(a + b*Tan[c + d*x])) - (((3*A*b - 2*a*B)*Cot[c + d*x])/(a*d*(a + b*Tan[c + d*x]))) + (2*((-(a^3*(2*a*A*b - a^2*B + b^2*B)*x)/(a^2 + b^2)) + ((a^2 + b^2)*(a^2*A - 3*A*b^2 + 2*a*b*B)*Log[-Sin[c + d*x]])/(a*d) + (b^3*(5*a^2*A*b + 3*A*b^3 - 4*a^3*B - 2*a*b^2*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))/(a*(a^2 + b^2)) - (b*(2*a^2*A*b + 3*A*b^3 - a^3*B - 2*a*b^2*B))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])))/a/(2*a)
```

3.281.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4013 Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

3.281. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4134 `Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/(a^2 + b^2)*(c^2 + d^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.281.4 Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 227, normalized size of antiderivative = 0.91

method	result
derivativedivides	$\frac{(A a^2 - A b^2 + 2Bab) \ln(1 + \tan^2(dx+c)) + (2Aab - B a^2 + B b^2) \arctan(\tan(dx+c))}{(a^2+b^2)^2} - \frac{A}{2a^2 \tan(dx+c)^2} - \frac{-2Ab+Ba}{a^3 \tan(dx+c)} + \frac{(-A a^2 + 3A b^2 + 2Bab)}{d}$
default	$\frac{(A a^2 - A b^2 + 2Bab) \ln(1 + \tan^2(dx+c)) + (2Aab - B a^2 + B b^2) \arctan(\tan(dx+c))}{(a^2+b^2)^2} - \frac{A}{2a^2 \tan(dx+c)^2} - \frac{-2Ab+Ba}{a^3 \tan(dx+c)} + \frac{(-A a^2 + 3A b^2 + 2Bab)}{d}$
parallelrisc	$-10b^3 (A a^2 b + \frac{3}{5} A b^3 - \frac{4}{5} B a^3 - \frac{2}{5} B a b^2) (a+b \tan(dx+c)) \ln(a+b \tan(dx+c)) + a^4 (a+b \tan(dx+c)) (A a^2 - A b^2 + 2Bab) \ln(a+b \tan(dx+c))$
norman	$\frac{a(2Aab - B a^2 + B b^2)x(\tan^2(dx+c))}{a^4 + 2a^2b^2 + b^4} + \frac{b(2Aab - B a^2 + B b^2)x(\tan^3(dx+c))}{a^4 + 2a^2b^2 + b^4} - \frac{A}{2ad} + \frac{(3Ab - 2Ba) \tan(dx+c)}{2a^2d} - \frac{(2A a^2 b^2 + 3A b^4 - B a^2 b^2)}{d a^4} \tan(dx+c)^2 (a+b \tan(dx+c))$
risc	$\frac{x B}{2iba - a^2 + b^2} - \frac{4ib^5 Bx}{a^3(a^4 + 2a^2b^2 + b^4)} - \frac{4ib^5 Bc}{a^3d(a^4 + 2a^2b^2 + b^4)} + \frac{10ib^4 Ac}{a^2d(a^4 + 2a^2b^2 + b^4)} + \frac{6ib^6 Ac}{a^4d(a^4 + 2a^2b^2 + b^4)} + \frac{2iAc}{a^2d}$

input `int(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(1/(a^2+b^2)^2*(1/2*(A*a^2-A*b^2+2*B*a*b)*ln(1+tan(d*x+c)^2)+(2*A*a*b-B*a^2+B*b^2)*arctan(tan(d*x+c)))-1/2/a^2*A/tan(d*x+c)^2-(-2*A*b+B*a)/a^3/tan(d*x+c)+(-A*a^2+3*A*b^2-2*B*a*b)/a^4*ln(tan(d*x+c))-b^3*(5*A*a^2*b+3*A*b^3-4*B*a^3-2*B*a*b^2)/(a^2+b^2)^2/a^4*ln(a+b*tan(d*x+c))+(A*b-B*a)*b^3/(a^2+b^2)/a^3/(a+b*tan(d*x+c))`

3.281.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 590 vs. 2(246) = 492.

Time = 0.35 (sec) , antiderivative size = 590, normalized size of antiderivative = 2.36

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \frac{Aa^7 + 2Aa^5b^2 + Aa^3b^4 + (Aa^6b + 2Aa^4b^3 - 2Ba^3b^4 + 3Aa^2b^5 + 2(Ba^6b - 2Aa^5b^2 - Ba^4b^3)dx) \tan(c+dx)}{d(a+b \tan(c+dx))^2}$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

3.281. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$


```
output -1/2*(A*a^7 + 2*A*a^5*b^2 + A*a^3*b^4 + (A*a^6*b + 2*A*a^4*b^3 - 2*B*a^3*b^4 + 3*A*a^2*b^5 + 2*(B*a^6*b - 2*A*a^5*b^2 - B*a^4*b^3)*d*x)*tan(d*x + c)^3 + (A*a^7 + 2*B*a^6*b - 2*A*a^5*b^2 + 4*B*a^4*b^3 - 7*A*a^3*b^4 + 4*B*a^2*b^5 - 6*A*a*b^6 + 2*(B*a^7 - 2*A*a^6*b - B*a^5*b^2)*d*x)*tan(d*x + c)^2 + ((A*a^6*b + 2*B*a^5*b^2 - A*a^4*b^3 + 4*B*a^3*b^4 - 5*A*a^2*b^5 + 2*B*a*b^6 - 3*A*b^7)*tan(d*x + c)^3 + (A*a^7 + 2*B*a^6*b - A*a^5*b^2 + 4*B*a^4*b^3 - 5*A*a^3*b^4 + 2*B*a^2*b^5 - 3*A*a*b^6)*tan(d*x + c)^2)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1)) - ((4*B*a^3*b^4 - 5*A*a^2*b^5 + 2*B*a*b^6 - 3*A*b^7)*tan(d*x + c)^3 + (4*B*a^4*b^3 - 5*A*a^3*b^4 + 2*B*a^2*b^5 - 3*A*a*b^6)*tan(d*x + c)^2)*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) + (2*B*a^7 - 3*A*a^6*b + 4*B*a^5*b^2 - 6*A*a^4*b^3 + 2*B*a^3*b^4 - 3*A*a^2*b^5)*tan(d*x + c))/((a^8*b + 2*a^6*b^3 + a^4*b^5)*d*tan(d*x + c)^3 + (a^9 + 2*a^7*b^2 + a^5*b^4)*d*tan(d*x + c)^2)
```

3.281.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 4.01 (sec) , antiderivative size = 9896, normalized size of antiderivative = 39.58

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)**3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)
```

output `Piecewise(((A*log(tan(c + d*x)**2 + 1)/(2*d) - A*log(tan(c + d*x))/d - A/(2*d*tan(c + d*x)**2) - B*x - B/(d*tan(c + d*x)))/a**2, Eq(b, 0)), ((-A*log(tan(c + d*x)**2 + 1)/(2*d) + A*log(tan(c + d*x))/d + A/(2*d*tan(c + d*x)**2) - A/(4*d*tan(c + d*x)**4) + B*x + B/(d*tan(c + d*x)) - B/(3*d*tan(c + d*x)**3))/b**2, Eq(a, 0)), (-15*I*A*d*x*tan(c + d*x)**4/(4*a**2*d*tan(c + d*x)**4 + 8*I*a**2*d*tan(c + d*x)**3 - 4*a**2*d*tan(c + d*x)**2) + 30*A*d*x*tan(c + d*x)**3/(4*a**2*d*tan(c + d*x)**4 + 8*I*a**2*d*tan(c + d*x)**3 - 4*a**2*d*tan(c + d*x)**2) + 15*I*A*d*x*tan(c + d*x)**2/(4*a**2*d*tan(c + d*x)**4 + 8*I*a**2*d*tan(c + d*x)**3 - 4*a**2*d*tan(c + d*x)**2) + 8*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**4/(4*a**2*d*tan(c + d*x)**4 + 8*I*a**2*d*tan(c + d*x)**3 - 4*a**2*d*tan(c + d*x)**2) + 16*I*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**3/(4*a**2*d*tan(c + d*x)**4 + 8*I*a**2*d*tan(c + d*x)**3 - 4*a**2*d*tan(c + d*x)**2) - 8*A*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2/(4*a**2*d*tan(c + d*x)**4 + 8*I*a**2*d*tan(c + d*x)**3 - 4*a**2*d*tan(c + d*x)**2) - 16*A*log(tan(c + d*x))*tan(c + d*x)**4/(4*a**2*d*tan(c + d*x)**4 + 8*I*a**2*d*tan(c + d*x)**3 - 4*a**2*d*tan(c + d*x)**2) - 32*I*A*log(tan(c + d*x))*tan(c + d*x)**3/(4*a**2*d*tan(c + d*x)**4 + 8*I*a**2*d*tan(c + d*x)**3 - 4*a**2*d*tan(c + d*x)**2) + 16*A*log(tan(c + d*x))*tan(c + d*x)**2/(4*a**2*d*tan(c + d*x)**4 + 8*I*a**2*d*tan(c + d*x)**3 - 4*a**2*d*tan(c + d*x)**2) - 15*I*A*tan(c + d*x)**3/(4*a**2*d*tan(c + d*x)**4 + ...`

3.281.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 325, normalized size of antiderivative = 1.30

$$\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx =$$

$$\frac{2(Ba^2 - 2Aab - Bb^2)(dx + c)}{a^4 + 2a^2b^2 + b^4} - \frac{2(4Ba^3b^3 - 5Aa^2b^4 + 2Bab^5 - 3Ab^6) \log(b \tan(dx + c) + a)}{a^8 + 2a^6b^2 + a^4b^4} - \frac{(Aa^2 + 2Bab - Ab^2) \log(\tan(dx + c)^2 + 1)}{a^4 + 2a^2b^2 + b^4} +$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output
$$-1/2*(2*(B*a^2 - 2*A*a*b - B*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) - 2*(4*B*a^3*b^3 - 5*A*a^2*b^4 + 2*B*a*b^5 - 3*A*b^6)*\log(b*\tan(d*x + c) + a)/(a^8 + 2*a^6*b^2 + a^4*b^4) - (A*a^2 + 2*B*a*b - A*b^2)*\log(\tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) + (A*a^4 + A*a^2*b^2 + 2*(B*a^3*b - 2*A*a^2*b^2 + 2*B*a*b^3 - 3*A*b^4)*\tan(d*x + c)^2 + (2*B*a^4 - 3*A*a^3*b + 2*B*a^2*b^2 - 3*A*a*b^3)*\tan(d*x + c))/((a^5*b + a^3*b^3)*\tan(d*x + c)^3 + (a^6 + a^4*b^2)*\tan(d*x + c)^2) + 2*(A*a^2 + 2*B*a*b - 3*A*b^2)*\log(\tan(d*x + c))/a^4)/d$$

3.281.8 Giac [A] (verification not implemented)

Time = 0.91 (sec) , antiderivative size = 402, normalized size of antiderivative = 1.61

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \frac{2(Ba^2-2Aab-Bb^2)(dx+c)}{a^4+2a^2b^2+b^4} - \frac{(Aa^2+2Bab-Ab^2)\log(\tan(dx+c)^2+1)}{a^4+2a^2b^2+b^4} - \frac{2(4Ba^3b^4-5Aa^2b^5+2Bab^6-3Ab^7)\log(|b\tan(dx+c)+a|)}{a^8b+2a^6b^3+a^4b^5} + \dots$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output
$$-1/2*(2*(B*a^2 - 2*A*a*b - B*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) - (A*a^2 + 2*B*a*b - A*b^2)*\log(\tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) - 2*(4*B*a^3*b^4 - 5*A*a^2*b^5 + 2*B*a*b^6 - 3*A*b^7)*\log(\text{abs}(b*\tan(d*x + c) + a))/(a^8*b + 2*a^6*b^3 + a^4*b^5) + 2*(4*B*a^3*b^4*\tan(d*x + c) - 5*A*a^2*b^5*\tan(d*x + c) + 2*B*a*b^6*\tan(d*x + c) - 3*A*b^7*\tan(d*x + c) + 5*B*a^4*b^3 - 6*A*a^3*b^4 + 3*B*a^2*b^5 - 4*A*a*b^6)/((a^8 + 2*a^6*b^2 + a^4*b^4)*(b*\tan(d*x + c) + a)) + 2*(A*a^2 + 2*B*a*b - 3*A*b^2)*\log(\text{abs}(\tan(d*x + c))))/a^4 - (3*A*a^2*\tan(d*x + c)^2 + 6*B*a*b*\tan(d*x + c)^2 - 9*A*b^2*\tan(d*x + c)^2 - 2*B*a^2*\tan(d*x + c) + 4*A*a*b*\tan(d*x + c) - A*a^2)/(a^4*\tan(d*x + c)^2))/d$$

3.281.9 Mupad [B] (verification not implemented)

Time = 11.86 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.14

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{\frac{\tan(c+dx)(3Ab-2Ba)}{2a^2} - \frac{A}{2a} + \frac{\tan(c+dx)^2(-Ba^3b+2Aa^2b^2-2Bab^3+3Ab^4)}{a^3(a^2+b^2)}}{d(b \tan(c+dx)^3 + a \tan(c+dx)^2)}$$

$$- \frac{\ln(\tan(c+dx))(Aa^2 + 2Bab - 3Ab^2)}{a^4d} + \frac{\ln(\tan(c+dx) - i)(A + B i)}{2d(a^2 + ab2i - b^2)}$$

$$- \frac{\ln(a + b \tan(c+dx))(-4Ba^3b^3 + 5Aa^2b^4 - 2Bab^5 + 3Ab^6)}{d(a^8 + 2a^6b^2 + a^4b^4)}$$

$$+ \frac{\ln(\tan(c+dx) + i)(B + A i)}{2d(a^2 i + 2ab - b^2 i)}$$

input `int((cot(c + d*x)^3*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`output `((tan(c + d*x)*(3*A*b - 2*B*a))/(2*a^2) - A/(2*a) + (tan(c + d*x)^2*(3*A*b^4 + 2*A*a^2*b^2 - 2*B*a*b^3 - B*a^3*b))/(a^3*(a^2 + b^2)))/(d*(a*tan(c + d*x)^2 + b*tan(c + d*x)^3)) - (log(tan(c + d*x))*(A*a^2 - 3*A*b^2 + 2*B*a*b))/(a^4*d) + (log(tan(c + d*x) - i)*(A + B*i))/(2*d*(a*b*2i + a^2 - b^2)) - (log(a + b*tan(c + d*x))*(3*A*b^6 + 5*A*a^2*b^4 - 4*B*a^3*b^3 - 2*B*a*b^5))/(d*(a^8 + a^4*b^4 + 2*a^6*b^2)) + (log(tan(c + d*x) + i)*(A*i + B))/(2*d*(2*a*b + a^2*i - b^2*i))`

$$3.282 \quad \int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

3.282.1 Optimal result	2710
3.282.2 Mathematica [C] (verified)	2711
3.282.3 Rubi [A] (verified)	2711
3.282.4 Maple [A] (verified)	2717
3.282.5 Fricas [B] (verification not implemented)	2717
3.282.6 Sympy [F(-2)]	2718
3.282.7 Maxima [A] (verification not implemented)	2719
3.282.8 Giac [A] (verification not implemented)	2719
3.282.9 Mupad [B] (verification not implemented)	2720

3.282.1 Optimal result

Integrand size = 31, antiderivative size = 331

$$\begin{aligned} & \int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\ &= \frac{(a^3A - 3aAb^2 + 3a^2bB - b^3B)x}{(a^2+b^2)^3} + \frac{(3a^2Ab - Ab^3 - a^3B + 3ab^2B) \log(\cos(c+dx))}{(a^2+b^2)^3 d} \\ &+ \frac{a^2(a^4Ab + 3a^2Ab^3 + 6Ab^5 - 3a^5B - 9a^3b^2B - 10ab^4B) \log(a+b \tan(c+dx))}{b^4(a^2+b^2)^3 d} \\ &- \frac{(a^3Ab + 3aAb^3 - 3a^4B - 6a^2b^2B - b^4B) \tan(c+dx)}{b^3(a^2+b^2)^2 d} \\ &+ \frac{a(Ab - aB) \tan^3(c+dx)}{2b(a^2+b^2)d(a+b \tan(c+dx))^2} + \frac{a(a^2Ab + 5Ab^3 - 3a^3B - 7ab^2B) \tan^2(c+dx)}{2b^2(a^2+b^2)^2 d(a+b \tan(c+dx))} \end{aligned}$$

output $(A*a^3-3*A*a*b^2+3*B*a^2*b-B*b^3)*x/(a^2+b^2)^3+(3*A*a^2*b-A*b^3-B*a^3+3*B*a*b^2)*\ln(\cos(d*x+c))/(a^2+b^2)^3/d+a^2*(A*a^4*b+3*A*a^2*b^3+6*A*b^5-3*B*a^5-9*B*a^3*b^2-10*B*a*b^4)*\ln(a+b*\tan(d*x+c))/b^4/(a^2+b^2)^3/d-(A*a^3*b+3*A*a*b^3-3*B*a^4-6*B*a^2*b^2-B*b^4)*\tan(d*x+c)/b^3/(a^2+b^2)^2/d+1/2*a*(A*b-B*a)*\tan(d*x+c)^3/b/(a^2+b^2)/d/(a+b*\tan(d*x+c))^2+1/2*a*(A*a^2*b+5*A*b^3-3*B*a^3-7*B*a*b^2)*\tan(d*x+c)^2/b^2/(a^2+b^2)^2/d/(a+b*\tan(d*x+c))$

3.282.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 5.22 (sec) , antiderivative size = 275, normalized size of antiderivative = 0.83

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{\frac{(A+iB) \log(i-\tan(c+dx))}{(-ia+b)^3} + \frac{(A-iB) \log(i+\tan(c+dx))}{(ia+b)^3} + \frac{2a^2(a^4Ab+3a^2Ab^3+6Ab^5-3a^5B-9a^3b^2B-10ab^4B) \log(a+b \tan(c+dx))}{b^4(a^2+b^2)^3} + \frac{a^2}{b^4(a^2+b^2)^3}}{2d}$$

input `Integrate[(Tan[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

output
$$\left(\frac{(A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]]}{(-I)*a + b} + \frac{(A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]]}{I*a + b} \right) / (I*a + b)^3 + \frac{(2*a^2*(a^4*A*b + 3*a^2*A*b^3 + 6*A*b^5 - 3*a^5*B - 9*a^3*b^2*B - 10*a*b^4*B)*\text{Log}[a + b*\text{Tan}[c + d*x]]}{(b^4*(a^2 + b^2)^3) + (a^3*(-(a*A*b) + 3*a^2*B + 2*b^2*B)) / (b^4*(a^2 + b^2)*(a + b*\text{Tan}[c + d*x])^2) + (2*B*\text{Tan}[c + d*x]^3) / (b*(a + b*\text{Tan}[c + d*x])^2) - (2*a^2*(-2*a^3*A*b - 4*a*A*b^3 + 6*a^4*B + 11*a^2*b^2*B + 3*b^4*B)) / (b^4*(a^2 + b^2)^2*(a + b*\text{Tan}[c + d*x]))}{(2*d)}$$

3.282.3 Rubi [A] (verified)

Time = 1.84 (sec) , antiderivative size = 363, normalized size of antiderivative = 1.10, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 4088, 25, 3042, 4128, 27, 3042, 4130, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$\downarrow 3042$$

$$\int \frac{\tan(c+dx)^4(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$\downarrow 4088$$

$$\begin{aligned}
& \frac{\int -\frac{\tan^2(c+dx)((-3Ba^2+Aba-2b^2B)\tan^2(c+dx)-2b(Ab-aB)\tan(c+dx)+3a(Ab-aB))}{(a+b\tan(c+dx))^2} dx}{2b(a^2+b^2)} + \\
& \quad \frac{a(Ab-aB)\tan^3(c+dx)}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} \\
& \quad \downarrow 25 \\
& \frac{\int \frac{\tan^2(c+dx)((-3Ba^2+Aba-2b^2B)\tan^2(c+dx)-2b(Ab-aB)\tan(c+dx)+3a(Ab-aB))}{(a+b\tan(c+dx))^2} dx}{2b(a^2+b^2)} - \\
& \quad \frac{a(Ab-aB)\tan^3(c+dx)}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\tan(c+dx)^2((-3Ba^2+Aba-2b^2B)\tan(c+dx)^2-2b(Ab-aB)\tan(c+dx)+3a(Ab-aB))}{(a+b\tan(c+dx))^2} dx}{2b(a^2+b^2)} - \\
& \quad \frac{a(Ab-aB)\tan^3(c+dx)}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} \\
& \quad \downarrow 4128 \\
& \frac{\int \frac{2\tan(c+dx)((Aa^2+2bBa-Ab^2)\tan(c+dx)b^2+(-3Ba^4+Aba^3-6b^2Ba^2+3Ab^3a-b^4B)\tan^2(c+dx)+a(-3Ba^3+Ab^2-7b^2Ba+5Ab^3))}{(a+b\tan(c+dx))b(a^2+b^2)} dx}{2b(a^2+b^2)} - \frac{a(-3a^3B+a^2)}{bd(a^2+b^2)} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{\tan(c+dx)((Aa^2+2bBa-Ab^2)\tan(c+dx)b^2+(-3Ba^4+Aba^3-6b^2Ba^2+3Ab^3a-b^4B)\tan^2(c+dx)+a(-3Ba^3+Ab^2-7b^2Ba+5Ab^3))}{(a+b\tan(c+dx))b(a^2+b^2)} dx}{2b(a^2+b^2)} - \frac{a(-3a^3B+a^2)}{bd(a^2+b^2)} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\tan(c+dx)((Aa^2+2bBa-Ab^2)\tan(c+dx)b^2+(-3Ba^4+Aba^3-6b^2Ba^2+3Ab^3a-b^4B)\tan^2(c+dx)+a(-3Ba^3+Ab^2-7b^2Ba+5Ab^3))}{(a+b\tan(c+dx))b(a^2+b^2)} dx}{2b(a^2+b^2)} - \frac{a(-3a^3B+a^2)}{bd(a^2+b^2)} \\
& \quad \downarrow 4130
\end{aligned}$$

3.282. $\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

$$\frac{\frac{a(Ab - aB) \tan^3(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \int \frac{-((-Ba^2 + 2Aba + b^2B) \tan(c + dx)b^3) + (a^2 + b^2)^2 (Ab - 3aB) \tan^2(c + dx) + a(-3Ba^4 + Aba^3 - 6b^2Ba^2 + 3Ab^3a - b^4B)}{a + b \tan(c + dx)} dx}{b(a^2 + b^2)} + \frac{(-3a^4B + a^3Ab - 6a^2b^2B + 3aAb^3 - b^4B)}{bd}$$

$$2b(a^2 + b^2)$$

↓ 25

$$\frac{\frac{a(Ab - aB) \tan^3(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \int \frac{-((-Ba^2 + 2Aba + b^2B) \tan(c + dx)b^3) + (a^2 + b^2)^2 (Ab - 3aB) \tan^2(c + dx) + a(-3Ba^4 + Aba^3 - 6b^2Ba^2 + 3Ab^3a - b^4B)}{a + b \tan(c + dx)} dx}{b(a^2 + b^2)} + \frac{(-3a^4B + a^3Ab - 6a^2b^2B + 3aAb^3 - b^4B)}{bd} \tan(c + dx)$$

$$2b(a^2 + b^2)$$

↓ 3042

$$\frac{\frac{a(Ab - aB) \tan^3(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \int \frac{-((-Ba^2 + 2Aba + b^2B) \tan(c + dx)b^3) + (a^2 + b^2)^2 (Ab - 3aB) \tan(c + dx) + a(-3Ba^4 + Aba^3 - 6b^2Ba^2 + 3Ab^3a - b^4B)}{a + b \tan(c + dx)} dx}{b(a^2 + b^2)} + \frac{(-3a^4B + a^3Ab - 6a^2b^2B + 3aAb^3 - b^4B)}{bd} \tan(c + dx)$$

$$2b(a^2 + b^2)$$

↓ 4109

$$\frac{\frac{a(Ab - aB) \tan^3(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \int \frac{-((-Ba^2 + 2Aba + b^2B) \tan(c + dx)b^3) + (a^2 + b^2)^2 (Ab - 3aB) \tan(c + dx) + a(-3Ba^4 + Aba^3 - 6b^2Ba^2 + 3Ab^3a - b^4B)}{a + b \tan(c + dx)} dx}{b(a^2 + b^2)} + \frac{a^2(-3a^5B + a^4Ab - 9a^3b^2B + 3a^2Ab^3 - 10ab^4B + 6a^2b^5)}{b(a^2 + b^2)} + \frac{b^3(a^3(-B) + 3a^2Ab + 3ab^2B - Ab^3) \int \tan(c + dx) dx}{a^2 + b^2}$$

$$2b(a^2 + b^2)$$

↓ 3042

$$\frac{\frac{a(Ab - aB) \tan^3(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \int \frac{-((-Ba^2 + 2Aba + b^2B) \tan(c + dx)b^3) + (a^2 + b^2)^2 (Ab - 3aB) \tan(c + dx) + a(-3Ba^4 + Aba^3 - 6b^2Ba^2 + 3Ab^3a - b^4B)}{a + b \tan(c + dx)} dx}{b(a^2 + b^2)} + \frac{a^2(-3a^5B + a^4Ab - 9a^3b^2B + 3a^2Ab^3 - 10ab^4B + 6a^2b^5)}{b(a^2 + b^2)} + \frac{b^3(a^3(-B) + 3a^2Ab + 3ab^2B - Ab^3) \int \tan(c + dx) dx}{a^2 + b^2}$$

$$2b(a^2 + b^2)$$

↓ 3956

3.282. $\int \frac{\tan^4(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$

$$\begin{array}{c}
\frac{a(Ab - aB) \tan^3(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \\
\frac{(-3a^4B + a^3Ab - 6a^2b^2B + 3aAb^3 - b^4B) \tan(c + dx)}{bd} - \frac{a^2(-3a^5B + a^4Ab - 9a^3b^2B + 3a^2Ab^3 - 10ab^4B + 6Ab^5) \int \frac{\tan(c+dx)^2 + 1}{a+b \tan(c+dx)} dx}{a^2 + b^2} + \frac{b^3(a^3(-B) + 3a^2Ab + 3ab^2B)}{b} \\
\hline
\frac{b(a^2 + b^2)}{2b(a^2 + b^2)} \\
\downarrow 4100 \\
\frac{a(Ab - aB) \tan^3(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \\
\frac{(-3a^4B + a^3Ab - 6a^2b^2B + 3aAb^3 - b^4B) \tan(c + dx)}{bd} - \frac{a^2(-3a^5B + a^4Ab - 9a^3b^2B + 3a^2Ab^3 - 10ab^4B + 6Ab^5) \int \frac{1}{a+b \tan(c+dx)} d(b \tan(c+dx))}{bd(a^2 + b^2)} + \frac{b^3(a^3(-B) + 3a^2Ab + 3ab^2B)}{b} \\
\hline
\frac{b(a^2 + b^2)}{2b(a^2 + b^2)} \\
\downarrow 16 \\
\frac{a(Ab - aB) \tan^3(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \\
\frac{(-3a^4B + a^3Ab - 6a^2b^2B + 3aAb^3 - b^4B) \tan(c + dx)}{bd} - \frac{b^3(a^3(-B) + 3a^2Ab + 3ab^2B - Ab^3) \log(\cos(c + dx))}{d(a^2 + b^2)} + \frac{b^3x(a^3A + 3a^2bB - 3aAb^2 - b^3B)}{a^2 + b^2} + \frac{a^2(-3a^5B + a^4Ab - 9a^3b^2B + 3a^2Ab^3 - 10ab^4B + 6Ab^5)}{b} \\
\hline
\frac{b(a^2 + b^2)}{2b(a^2 + b^2)}
\end{array}$$

input `Int[(Tan[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

output `(a*(A*b - a*B)*Tan[c + d*x]^3)/(2*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2 - ((a*(a^2*A*b + 5*A*b^3 - 3*a^3*B - 7*a*b^2*B)*Tan[c + d*x]^2)/(b*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))) + (2*(-((b^3*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B)*x)/(a^2 + b^2) + (b^3*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B)*Log[Cos[c + d*x]])/((a^2 + b^2)*d) + (a^2*(a^4*A*b + 3*a^2*A*b^3 + 6*A*b^5 - 3*a^5*B - 9*a^3*b^2*B - 10*a*b^4*B)*Log[a + b*Tan[c + d*x]])/(b*(a^2 + b^2)*d))/b) + ((a^3*A*b + 3*a*A*b^3 - 3*a^4*B - 6*a^2*b^2*B - b^4*B)*Tan[c + d*x])/(b*d))/((b*(a^2 + b^2)))/(2*b*(a^2 + b^2))`

3.282.3.1 Defintions of rubi rules used

- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_.)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_.)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`
- rule 4100 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]`

rule 4109 `Int[((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C, 0]`

rule 4128 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)]^(m_))*((c_) + (d_)*tan[(e_) + (f_)*(x_)]^(n_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4130 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)]^(m_))*((c_) + (d_)*tan[(e_) + (f_)*(x_)]^(n_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.282.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.79

method	result
derivativedivides	$\frac{\tan(dx+c)B}{b^3} + \frac{(-3Aa^2b+Ab^3+Ba^3-3Bab^2)\ln(1+\tan^2(dx+c))}{2} + \frac{(Aa^3-3Aab^2+3Ba^2b-Bb^3)\arctan(\tan(dx+c))}{(a^2+b^2)^3} + \frac{a^2(Aa^4b+...)}{(a^2+b^2)^3}$
default	$\frac{\tan(dx+c)B}{b^3} + \frac{(-3Aa^2b+Ab^3+Ba^3-3Bab^2)\ln(1+\tan^2(dx+c))}{2} + \frac{(Aa^3-3Aab^2+3Ba^2b-Bb^3)\arctan(\tan(dx+c))}{(a^2+b^2)^3} + \frac{a^2(Aa^4b+...)}{(a^2+b^2)^3}$
norman	$\frac{B(\tan^3(dx+c))}{bd} + \frac{(Aa^3-3Aab^2+3Ba^2b-Bb^3)a^2x}{(a^4+2a^2b^2+b^4)(a^2+b^2)} + \frac{b^2(Aa^3-3Aab^2+3Ba^2b-Bb^3)x(\tan^2(dx+c))}{(a^4+2a^2b^2+b^4)(a^2+b^2)} + \frac{a(2Aa^4b+4Aa^2b^3-6Bb^3a)}{db^3(a^2+b^2)}$
parallelrisc	Expression too large to display
risc	Expression too large to display

input `int(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/d*(tan(d*x+c)*B/b^3+1/(a^2+b^2)^3*(1/2*(-3*A*a^2*b+A*b^3+B*a^3-3*B*a*b^2)*ln(1+tan(d*x+c)^2)+(A*a^3-3*A*a*b^2+3*B*a^2*b-B*b^3)*arctan(tan(d*x+c)))+1/b^4*a^2*(A*a^4*b+3*A*a^2*b^3+6*A*b^5-3*B*a^5-9*B*a^3*b^2-10*B*a*b^4)/(a^2+b^2)^3*ln(a+b*tan(d*x+c))-1/2/b^4*a^4*(A*b-B*a)/(a^2+b^2)/(a+b*tan(d*x+c))^2+1/b^4*a^3*(2*A*a^2*b+4*A*b^3-3*B*a^3-5*B*a*b^2)/(a^2+b^2)^2/(a+b*tan(d*x+c)))`

3.282.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 890 vs. 2(328) = 656.

Time = 0.38 (sec) , antiderivative size = 890, normalized size of antiderivative = 2.69

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \frac{3Ba^7b^2 - Aa^6b^3 + 9Ba^5b^4 - 7Aa^4b^5 - 2(Ba^6b^3 + 3Ba^4b^5 + 3Ba^2b^7 + Bb^9)\tan(dx+c)^3 - 2(Aa^5b^4}{(a+b\tan(c+dx))^3}$$

input `integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="fricas")`

3.282. $\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

```
output -1/2*(3*B*a^7*b^2 - A*a^6*b^3 + 9*B*a^5*b^4 - 7*A*a^4*b^5 - 2*(B*a^6*b^3 +
3*B*a^4*b^5 + 3*B*a^2*b^7 + B*b^9)*tan(d*x + c)^3 - 2*(A*a^5*b^4 + 3*B*a^
4*b^5 - 3*A*a^3*b^6 - B*a^2*b^7)*d*x - (9*B*a^7*b^2 - 3*A*a^6*b^3 + 23*B*a
^5*b^4 - 9*A*a^4*b^5 + 12*B*a^3*b^6 + 4*B*a*b^8 + 2*(A*a^3*b^6 + 3*B*a^2*b
^7 - 3*A*a*b^8 - B*b^9)*d*x)*tan(d*x + c)^2 + (3*B*a^9 - A*a^8*b + 9*B*a^7
*b^2 - 3*A*a^6*b^3 + 10*B*a^5*b^4 - 6*A*a^4*b^5 + (3*B*a^7*b^2 - A*a^6*b^3
+ 9*B*a^5*b^4 - 3*A*a^4*b^5 + 10*B*a^3*b^6 - 6*A*a^2*b^7)*tan(d*x + c)^2
+ 2*(3*B*a^8*b - A*a^7*b^2 + 9*B*a^6*b^3 - 3*A*a^5*b^4 + 10*B*a^4*b^5 - 6*
A*a^3*b^6)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^
2)/(tan(d*x + c)^2 + 1)) - (3*B*a^9 - A*a^8*b + 9*B*a^7*b^2 - 3*A*a^6*b^3
+ 9*B*a^5*b^4 - 3*A*a^4*b^5 + 3*B*a^3*b^6 - A*a^2*b^7 + (3*B*a^7*b^2 - A*a
^6*b^3 + 9*B*a^5*b^4 - 3*A*a^4*b^5 + 9*B*a^3*b^6 - 3*A*a^2*b^7 + 3*B*a*b^8
- A*b^9)*tan(d*x + c)^2 + 2*(3*B*a^8*b - A*a^7*b^2 + 9*B*a^6*b^3 - 3*A*a^
5*b^4 + 9*B*a^4*b^5 - 3*A*a^3*b^6 + 3*B*a^2*b^7 - A*a*b^8)*tan(d*x + c))*l
og(1/(tan(d*x + c)^2 + 1)) - 2*(3*B*a^8*b - A*a^7*b^2 + 6*B*a^6*b^3 - 3*A*
a^5*b^4 - 2*B*a^4*b^5 + 4*A*a^3*b^6 + B*a^2*b^7 + 2*(A*a^4*b^5 + 3*B*a^3*b
^6 - 3*A*a^2*b^7 - B*a*b^8)*d*x)*tan(d*x + c))/((a^6*b^6 + 3*a^4*b^8 + 3*a
^2*b^10 + b^12)*d*tan(d*x + c)^2 + 2*(a^7*b^5 + 3*a^5*b^7 + 3*a^3*b^9 + a*
b^11)*d*tan(d*x + c) + (a^8*b^4 + 3*a^6*b^6 + 3*a^4*b^8 + a^2*b^10)*d)
```

3.282.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Exception raised: AttributeError}$$

```
input integrate(tan(d*x+c)**4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)
```

```
output Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'
```

3.282.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 389, normalized size of antiderivative = 1.18

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{2(Aa^3+3Ba^2b-3Aab^2-Bb^3)(dx+c)}{a^6+3a^4b^2+3a^2b^4+b^6} - \frac{2(3Ba^7-Aa^6b+9Ba^5b^2-3Aa^4b^3+10Ba^3b^4-6Aa^2b^5)\log(b\tan(dx+c)+a)}{a^6b^4+3a^4b^6+3a^2b^8+b^{10}} + \frac{(Ba^3-3Aa^2b-3Bab^2+Ab^3)\log(\tan(dx+c)^2+1)}{a^6+3a^4b^2+3a^2b^4+b^6} - \frac{2(3Ba^7-Aa^6b+9Ba^5b^2-3Aa^4b^3+10Ba^3b^4-6Aa^2b^5)\log(\tan(dx+c)^2+1)}{a^6+3a^4b^2+3a^2b^4+b^6} - \frac{(5Ba^7-3Aa^6b+9Ba^5b^2-7Aa^4b^3+2(3Ba^6b-2Aa^5b^2+5Ba^4b^3-4Aa^3b^4)\tan(dx+c))/(a^6b^4+2a^4b^6+a^2b^8+(a^4b^6+2a^2b^8+b^{10})\tan(dx+c)^2+2(a^5b^5+2a^3b^7+ab^9)\tan(dx+c))+2B\tan(dx+c)/b^3}{d}$$

```
input integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="maxima")
```

```
output 1/2*(2*(A*a^3 + 3*B*a^2*b - 3*A*a*b^2 - B*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - 2*(3*B*a^7 - A*a^6*b + 9*B*a^5*b^2 - 3*A*a^4*b^3 + 10*B*a^3*b^4 - 6*A*a^2*b^5)*log(b*tan(d*x + c) + a)/(a^6*b^4 + 3*a^4*b^6 + 3*a^2*b^8 + b^10) + (B*a^3 - 3*A*a^2*b - 3*B*a*b^2 + A*b^3)*log(tan(d*x + c)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - (5*B*a^7 - 3*A*a^6*b + 9*B*a^5*b^2 - 7*A*a^4*b^3 + 2*(3*B*a^6*b - 2*A*a^5*b^2 + 5*B*a^4*b^3 - 4*A*a^3*b^4)*tan(d*x + c))/(a^6*b^4 + 2*a^4*b^6 + a^2*b^8 + (a^4*b^6 + 2*a^2*b^8 + b^10)*tan(d*x + c)^2 + 2*(a^5*b^5 + 2*a^3*b^7 + a*b^9)*tan(d*x + c)) + 2*B*tan(d*x + c)/b^3)/d
```

3.282.8 Giac [A] (verification not implemented)

Time = 1.12 (sec) , antiderivative size = 505, normalized size of antiderivative = 1.53

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{2(Aa^3+3Ba^2b-3Aab^2-Bb^3)(dx+c)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{(Ba^3-3Aa^2b-3Bab^2+Ab^3)\log(\tan(dx+c)^2+1)}{a^6+3a^4b^2+3a^2b^4+b^6} - \frac{2(3Ba^7-Aa^6b+9Ba^5b^2-3Aa^4b^3+10Ba^3b^4-6Aa^2b^5)\log(\tan(dx+c)^2+1)}{a^6+3a^4b^2+3a^2b^4+b^6}$$

```
input integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="giac")
```

output $\frac{1}{2} \cdot (2 \cdot (A \cdot a^3 + 3 \cdot B \cdot a^2 \cdot b - 3 \cdot A \cdot a \cdot b^2 - B \cdot b^3) \cdot (d \cdot x + c) / (a^6 + 3 \cdot a^4 \cdot b^2 + 3 \cdot a^2 \cdot b^4 + b^6) + (B \cdot a^3 - 3 \cdot A \cdot a^2 \cdot b - 3 \cdot B \cdot a \cdot b^2 + A \cdot b^3) \cdot \log(\tan(d \cdot x + c)^2 + 1) / (a^6 + 3 \cdot a^4 \cdot b^2 + 3 \cdot a^2 \cdot b^4 + b^6) - 2 \cdot (3 \cdot B \cdot a^7 - A \cdot a^6 \cdot b + 9 \cdot B \cdot a^5 \cdot b^2 - 3 \cdot A \cdot a^4 \cdot b^3 + 10 \cdot B \cdot a^3 \cdot b^4 - 6 \cdot A \cdot a^2 \cdot b^5) \cdot \log(\text{abs}(b \cdot \tan(d \cdot x + c) + a)) / (a^6 \cdot b^4 + 3 \cdot a^4 \cdot b^6 + 3 \cdot a^2 \cdot b^8 + b^{10}) + 2 \cdot B \cdot \tan(d \cdot x + c) / b^3 + (9 \cdot B \cdot a^7 \cdot b^2 \cdot \tan(d \cdot x + c)^2 - 3 \cdot A \cdot a^6 \cdot b^3 \cdot \tan(d \cdot x + c)^2 + 27 \cdot B \cdot a^5 \cdot b^4 \cdot \tan(d \cdot x + c)^2 - 9 \cdot A \cdot a^4 \cdot b^5 \cdot \tan(d \cdot x + c)^2 + 30 \cdot B \cdot a^3 \cdot b^6 \cdot \tan(d \cdot x + c)^2 - 18 \cdot A \cdot a^2 \cdot b^7 \cdot \tan(d \cdot x + c)^2 + 12 \cdot B \cdot a^8 \cdot b \cdot \tan(d \cdot x + c) - 2 \cdot A \cdot a^7 \cdot b^2 \cdot \tan(d \cdot x + c) + 38 \cdot B \cdot a^6 \cdot b^3 \cdot \tan(d \cdot x + c) - 6 \cdot A \cdot a^5 \cdot b^4 \cdot \tan(d \cdot x + c) + 50 \cdot B \cdot a^4 \cdot b^5 \cdot \tan(d \cdot x + c) - 28 \cdot A \cdot a^3 \cdot b^6 \cdot \tan(d \cdot x + c) + 4 \cdot B \cdot a^9 + 13 \cdot B \cdot a^7 \cdot b^2 + A \cdot a^6 \cdot b^3 + 21 \cdot B \cdot a^5 \cdot b^4 - 11 \cdot A \cdot a^4 \cdot b^5) / ((a^6 \cdot b^4 + 3 \cdot a^4 \cdot b^6 + 3 \cdot a^2 \cdot b^8 + b^{10}) \cdot (b \cdot \tan(d \cdot x + c) + a)^2) / d$

3.282.9 Mupad [B] (verification not implemented)

Time = 9.00 (sec) , antiderivative size = 335, normalized size of antiderivative = 1.01

$$\int \frac{\tan^4(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{B \tan(c + dx)}{b^3 d} + \frac{\ln(\tan(c + dx) - i)(-B + A i)}{2 d (-a^3 - a^2 b 3i + 3 a b^2 + b^3 i)} + \frac{\ln(\tan(c + dx) + i)(A - B i)}{2 d (-a^3 i - 3 a^2 b + a b^2 3i + b^3)}$$

$$- \frac{\frac{5 B a^7 - 3 A a^6 b + 9 B a^5 b^2 - 7 A a^4 b^3}{2 b (a^4 + 2 a^2 b^2 + b^4)} + \frac{\tan(c + dx) (3 B a^6 - 2 A a^5 b + 5 B a^4 b^2 - 4 A a^3 b^3)}{a^4 + 2 a^2 b^2 + b^4}}{d (a^2 b^3 + 2 a b^4 \tan(c + dx) + b^5 \tan(c + dx)^2)}$$

$$+ \frac{a^2 \ln(a + b \tan(c + dx)) (-3 B a^5 + A a^4 b - 9 B a^3 b^2 + 3 A a^2 b^3 - 10 B a b^4 + 6 A b^5)}{b^4 d (a^2 + b^2)^3}$$

input `int((tan(c + d*x)^4*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3,x)`

output $(\log(\tan(c + d \cdot x) - i) \cdot (A \cdot i - B)) / (2 \cdot d \cdot (3 \cdot a \cdot b^2 - a^2 \cdot b \cdot 3i - a^3 + b^3 \cdot 1i)) - ((5 \cdot B \cdot a^7 - 7 \cdot A \cdot a^4 \cdot b^3 + 9 \cdot B \cdot a^5 \cdot b^2 - 3 \cdot A \cdot a^6 \cdot b) / (2 \cdot b \cdot (a^4 + b^4 + 2 \cdot a^2 \cdot b^2))) + (\tan(c + d \cdot x) \cdot (3 \cdot B \cdot a^6 - 4 \cdot A \cdot a^3 \cdot b^3 + 5 \cdot B \cdot a^4 \cdot b^2 - 2 \cdot A \cdot a^5 \cdot b)) / (a^4 + b^4 + 2 \cdot a^2 \cdot b^2) / (d \cdot (a^2 \cdot b^3 + b^5 \cdot \tan(c + d \cdot x)^2 + 2 \cdot a \cdot b^4 \cdot \tan(c + d \cdot x))) + (\log(\tan(c + d \cdot x) + i) \cdot (A - B \cdot 1i)) / (2 \cdot d \cdot (a \cdot b^2 \cdot 3i - 3 \cdot a^2 \cdot b - a^3 \cdot 1i + b^3)) + (B \cdot \tan(c + d \cdot x)) / (b^3 \cdot d) + (a^2 \cdot \log(a + b \cdot \tan(c + d \cdot x))) \cdot (6 \cdot A \cdot b^5 - 3 \cdot B \cdot a^5 + 3 \cdot A \cdot a^2 \cdot b^3 - 9 \cdot B \cdot a^3 \cdot b^2 + A \cdot a^4 \cdot b - 10 \cdot B \cdot a \cdot b^4)) / (b^4 \cdot d \cdot (a^2 + b^2)^3)$

3.283 $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

3.283.1 Optimal result 2721
 3.283.2 Mathematica [C] (verified) 2722
 3.283.3 Rubi [A] (verified) 2722
 3.283.4 Maple [A] (verified) 2726
 3.283.5 Fricas [B] (verification not implemented) 2727
 3.283.6 Sympy [F(-2)] 2727
 3.283.7 Maxima [A] (verification not implemented) 2728
 3.283.8 Giac [A] (verification not implemented) 2728
 3.283.9 Mupad [B] (verification not implemented) 2729

3.283.1 Optimal result

Integrand size = 31, antiderivative size = 250

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= -\frac{(3a^2Ab - Ab^3 - a^3B + 3ab^2B)x}{(a^2 + b^2)^3} + \frac{(a^3A - 3aAb^2 + 3a^2bB - b^3B) \log(\cos(c+dx))}{(a^2 + b^2)^3 d}$$

$$+ \frac{a(a^2Ab^3 - 3Ab^5 + a^5B + 3a^3b^2B + 6ab^4B) \log(a+b \tan(c+dx))}{b^3(a^2 + b^2)^3 d}$$

$$+ \frac{a(Ab - aB) \tan^2(c+dx)}{2b(a^2 + b^2) d(a+b \tan(c+dx))^2} - \frac{a^2(2Ab^3 - a(a^2 + 3b^2) B)}{b^3(a^2 + b^2)^2 d(a+b \tan(c+dx))}$$

output

```
-(3*A*a^2*b-A*b^3-B*a^3+3*B*a*b^2)*x/(a^2+b^2)^3+(A*a^3-3*A*a*b^2+3*B*a^2*b-B*b^3)*ln(cos(d*x+c))/(a^2+b^2)^3/d+a*(A*a^2*b^3-3*A*b^5+B*a^5+3*B*a^3*b^2+6*B*a*b^4)*ln(a+b*tan(d*x+c))/b^3/(a^2+b^2)^3/d+1/2*a*(A*b-B*a)*tan(d*x+c)^2/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^2-a^2*(2*A*b^3-a*(a^2+3*b^2)*B)/b^3/(a^2+b^2)^2/d/(a+b*tan(d*x+c))
```


3.283.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.37 (sec) , antiderivative size = 235, normalized size of antiderivative = 0.94

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= -\frac{(A+iB) \log(i-\tan(c+dx))}{2(a+ib)^3 d} - \frac{(A-iB) \log(i+\tan(c+dx))}{2(a-ib)^3 d}$$

$$+ \frac{a(a^2 Ab^3 - 3Ab^5 + a^5 B + 3a^3 b^2 B + 6ab^4 B) \log(a+b \tan(c+dx))}{b^3 (a^2 + b^2)^3 d}$$

$$+ \frac{a^3 (Ab - aB)}{2b^3 (a^2 + b^2) d (a+b \tan(c+dx))^2} - \frac{a^2 (a^2 Ab + 3Ab^3 - 2a^3 B - 4ab^2 B)}{b^3 (a^2 + b^2)^2 d (a+b \tan(c+dx))}$$

input `Integrate[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

output `-1/2*((A + I*B)*Log[I - Tan[c + d*x]])/((a + I*b)^3*d) - ((A - I*B)*Log[I + Tan[c + d*x]])/(2*(a - I*b)^3*d) + (a*(a^2*A*b^3 - 3*A*b^5 + a^5*B + 3*a^3*b^2*B + 6*a*b^4*B)*Log[a + b*Tan[c + d*x]])/(b^3*(a^2 + b^2)^3*d) + (a^3*(A*b - a*B))/(2*b^3*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) - (a^2*(a^2*A*b + 3*A*b^3 - 2*a^3*B - 4*a*b^2*B))/(b^3*(a^2 + b^2)^2*d*(a + b*Tan[c + d*x]))`

3.283.3 Rubi [A] (verified)

Time = 1.28 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.14, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4088, 27, 3042, 4118, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)^3(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$\downarrow \text{4088}$$

3.283. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\begin{aligned}
 & \int \frac{-\frac{2 \tan(c+dx) \left(-((a^2+b^2)B \tan^2(c+dx) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)) \right)}{(a+b \tan(c+dx))^2} dx}{2b(a^2+b^2)} + \\
 & \quad \frac{a(Ab-aB) \tan^2(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow 27 \\
 & \quad \frac{a(Ab-aB) \tan^2(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \\
 & \int \frac{\frac{\tan(c+dx) \left(-((a^2+b^2)B \tan^2(c+dx) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)) \right)}{(a+b \tan(c+dx))^2} dx}{b(a^2+b^2)} \\
 & \quad \downarrow 3042 \\
 & \quad \frac{a(Ab-aB) \tan^2(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \\
 & \int \frac{\frac{\tan(c+dx) \left(-((a^2+b^2)B \tan^2(c+dx) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)) \right)}{(a+b \tan(c+dx))^2} dx}{b(a^2+b^2)} \\
 & \quad \downarrow 4118 \\
 & \quad \frac{a(Ab-aB) \tan^2(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \\
 & \int \frac{\frac{(Aa^2+2bBa-Ab^2) \tan(c+dx)b^2 - (a^2+b^2)^2 B \tan^2(c+dx) + a(2Ab^3-a(a^2+3b^2)B)}{a+b \tan(c+dx)} dx}{b(a^2+b^2)} + \frac{a^2(2Ab^3-aB(a^2+3b^2))}{b^2d(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \downarrow 3042 \\
 & \quad \frac{a(Ab-aB) \tan^2(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \\
 & \int \frac{\frac{(Aa^2+2bBa-Ab^2) \tan(c+dx)b^2 - (a^2+b^2)^2 B \tan^2(c+dx) + a(2Ab^3-a(a^2+3b^2)B)}{a+b \tan(c+dx)} dx}{b(a^2+b^2)} + \frac{a^2(2Ab^3-aB(a^2+3b^2))}{b^2d(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \downarrow 4109 \\
 & \quad \frac{a(Ab-aB) \tan^2(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \\
 & \frac{b^2(a^3A+3a^2bB-3aAb^2-b^3B) \int \tan(c+dx) dx}{a^2+b^2} - \frac{a(a^5B+3a^3b^2B+a^2Ab^3+6ab^4B-3Ab^5) \int \frac{\tan^2(c+dx)+1}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{b^2x(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{a^2+b^2} + \frac{a^2(2)}{b^2d(a^2+b^2)} \\
 & \quad \downarrow 3042
 \end{aligned}$$

3.283. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{\frac{a(Ab - aB) \tan^2(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{b^2(a^3A + 3a^2bB - 3aAb^2 - b^3B) \int \tan(c + dx) dx}{a^2 + b^2} - \frac{a(a^5B + 3a^3b^2B + a^2Ab^3 + 6ab^4B - 3Ab^5) \int \frac{\tan(c + dx)^2 + 1}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{b^2x(a^3(-B) + 3a^2Ab + 3ab^2B - Ab^3)}{a^2 + b^2}}{b(a^2 + b^2)} + \frac{a^2(2)}{b^2d(a^2 + b^2)}$$

↓ 3956

$$\frac{\frac{a(Ab - aB) \tan^2(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(a^5B + 3a^3b^2B + a^2Ab^3 + 6ab^4B - 3Ab^5) \int \frac{\tan(c + dx)^2 + 1}{a + b \tan(c + dx)} dx}{a^2 + b^2} - \frac{b^2(a^3A + 3a^2bB - 3aAb^2 - b^3B) \log(\cos(c + dx))}{d(a^2 + b^2)} + \frac{b^2x(a^3(-B) + 3a^2Ab + 3ab^2B - Ab^3)}{a^2 + b^2}}{b(a^2 + b^2)} + \frac{a^2(2)}{b^2d(a^2 + b^2)}$$

↓ 4100

$$\frac{\frac{a(Ab - aB) \tan^2(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(a^5B + 3a^3b^2B + a^2Ab^3 + 6ab^4B - 3Ab^5) \int \frac{1}{a + b \tan(c + dx)} d(b \tan(c + dx))}{bd(a^2 + b^2)} - \frac{b^2(a^3A + 3a^2bB - 3aAb^2 - b^3B) \log(\cos(c + dx))}{d(a^2 + b^2)} + \frac{b^2x(a^3(-B) + 3a^2Ab + 3ab^2B - Ab^3)}{a^2 + b^2}}{b(a^2 + b^2)}$$

↓ 16

$$\frac{\frac{a(Ab - aB) \tan^2(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{b^2(a^3A + 3a^2bB - 3aAb^2 - b^3B) \log(\cos(c + dx))}{d(a^2 + b^2)} + \frac{b^2x(a^3(-B) + 3a^2Ab + 3ab^2B - Ab^3)}{a^2 + b^2} - \frac{a(a^5B + 3a^3b^2B + a^2Ab^3 + 6ab^4B - 3Ab^5)}{bd(a^2 + b^2)}}{b(a^2 + b^2)} + \frac{a^2(2Ab^3 - aB(a^2 + 3b^2))}{b^2d(a^2 + b^2)(a + b \tan(c + dx))}$$

input `Int[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

output `(a*(A*b - a*B)*Tan[c + d*x]^2)/(2*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) - (((b^2*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B)*x)/(a^2 + b^2) - (b^2*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B)*Log[Cos[c + d*x]])/((a^2 + b^2)*d) - (a*(a^2*A*b^3 - 3*A*b^5 + a^5*B + 3*a^3*b^2*B + 6*a*b^4*B)*Log[a + b*Tan[c + d*x]])/(b*(a^2 + b^2)*d))/(b*(a^2 + b^2)) + (a^2*(2*A*b^3 - a*(a^2 + 3*b^2)*B))/(b^2*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(b*(a^2 + b^2))`

3.283.3.1 Defintions of rubi rules used

- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_.)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`
- rule 4100 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_.)]^2), x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]`
- rule 4109 `Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_.)] + (C_.)*tan[(e_.) + (f_.)*(x_.)]^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C, 0]`

```
rule 4118 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.) + (f
_.)*(x_)])^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]
```

3.283.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 242, normalized size of antiderivative = 0.97

method	result
derivativedivides	$\frac{\frac{(-A a^3 + 3A a b^2 - 3B a^2 b + B b^3) \ln(1 + \tan^2(dx+c))}{2} + \frac{(-3A a^2 b + A b^3 + B a^3 - 3B a b^2) \arctan(\tan(dx+c))}{(a^2 + b^2)^3} + \frac{a(A a^2 b^3 - 3A b^5 + B a^5)}{d}$
default	$\frac{\frac{(-A a^3 + 3A a b^2 - 3B a^2 b + B b^3) \ln(1 + \tan^2(dx+c))}{2} + \frac{(-3A a^2 b + A b^3 + B a^3 - 3B a b^2) \arctan(\tan(dx+c))}{(a^2 + b^2)^3} + \frac{a(A a^2 b^3 - 3A b^5 + B a^5)}{d}$
norman	$\frac{a^2(A a^3 b + 5A a b^3 - 3B a^4 - 7B a^2 b^2)}{2d b^3(a^4 + 2a^2 b^2 + b^4)} - \frac{(3A a^2 b - A b^3 - B a^3 + 3B a b^2) a^2 x}{(a^4 + 2a^2 b^2 + b^4)(a^2 + b^2)} - \frac{b^2(3A a^2 b - A b^3 - B a^3 + 3B a b^2) x (\tan^2(dx+c))}{(a^4 + 2a^2 b^2 + b^4)(a^2 + b^2)} - \frac{a}{(a+b \tan(dx+c))^2}$
parallelrisch	Expression too large to display
risch	$-\frac{x B}{3i a^2 b - i b^3 - a^3 + 3a b^2} - \frac{2i a^6 B x}{b^3(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)} - \frac{12i a^2 b B c}{d(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)} + \frac{2i B c}{b^3 d} - \frac{2i a^3 A x}{a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6}$

```
input int(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x,method=_RETURNVERBO
SE)
```

```
output 1/d*(1/(a^2+b^2)^3*(1/2*(-A*a^3+3*A*a*b^2-3*B*a^2*b+B*b^3)*ln(1+tan(d*x+c)
^2)+(-3*A*a^2*b+A*b^3+B*a^3-3*B*a*b^2)*arctan(tan(d*x+c)))+a*(A*a^2*b^3-3*
A*b^5+B*a^5+3*B*a^3*b^2+6*B*a*b^4)/(a^2+b^2)^3/b^3*ln(a+b*tan(d*x+c))-a^2*
(A*a^2*b+3*A*b^3-2*B*a^3-4*B*a*b^2)/b^3/(a^2+b^2)^2/(a+b*tan(d*x+c))+1/2*a
^3*(A*b-B*a)/b^3/(a^2+b^2)/(a+b*tan(d*x+c))^2)
```

3.283.
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

3.283.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 666 vs. $2(245) = 490$.

Time = 0.34 (sec) , antiderivative size = 666, normalized size of antiderivative = 2.66

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{Ba^6b^2 + Aa^5b^3 + 7Ba^4b^4 - 5Aa^3b^5 + 2(Ba^5b^3 - 3Aa^4b^4 - 3Ba^3b^5 + Aa^2b^6)dx - (3Ba^6b^2 - Aa^5b^3 + 9$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="fracas")
```

```
output 1/2*(B*a^6*b^2 + A*a^5*b^3 + 7*B*a^4*b^4 - 5*A*a^3*b^5 + 2*(B*a^5*b^3 - 3*A*a^4*b^4 - 3*B*a^3*b^5 + A*a^2*b^6)*d*x - (3*B*a^6*b^2 - A*a^5*b^3 + 9*B*a^4*b^4 - 7*A*a^3*b^5 - 2*(B*a^3*b^5 - 3*A*a^2*b^6 - 3*B*a*b^7 + A*b^8)*d*x)*tan(d*x + c)^2 + (B*a^8 + 3*B*a^6*b^2 + A*a^5*b^3 + 6*B*a^4*b^4 - 3*A*a^3*b^5 + (B*a^6*b^2 + 3*B*a^4*b^4 + A*a^3*b^5 + 6*B*a^2*b^6 - 3*A*a*b^7)*tan(d*x + c)^2 + 2*(B*a^7*b + 3*B*a^5*b^3 + A*a^4*b^4 + 6*B*a^3*b^5 - 3*A*a^2*b^6)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - (B*a^8 + 3*B*a^6*b^2 + 3*B*a^4*b^4 + B*a^2*b^6 + (B*a^6*b^2 + 3*B*a^4*b^4 + 3*B*a^2*b^6 + B*b^8)*tan(d*x + c)^2 + 2*(B*a^7*b + 3*B*a^5*b^3 + 3*B*a^3*b^5 + B*a*b^7)*tan(d*x + c))*log(1/(tan(d*x + c)^2 + 1)) - 2*(B*a^7*b + 3*B*a^5*b^3 - 3*A*a^4*b^4 - 4*B*a^3*b^5 + 3*A*a^2*b^6 - 2*(B*a^4*b^4 - 3*A*a^3*b^5 - 3*B*a^2*b^6 + A*a*b^7)*d*x)*tan(d*x + c))/((a^6*b^5 + 3*a^4*b^7 + 3*a^2*b^9 + b^11)*d*tan(d*x + c)^2 + 2*(a^7*b^4 + 3*a^5*b^6 + 3*a^3*b^8 + a*b^10)*d*tan(d*x + c) + (a^8*b^3 + 3*a^6*b^5 + 3*a^4*b^7 + a^2*b^9)*d)
```

3.283.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Exception raised: AttributeError}$$

```
input integrate(tan(d*x+c)**3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)
```

```
output Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'
```

3.283. $\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

3.283.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 366, normalized size of antiderivative = 1.46

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{2(Ba^3-3Aa^2b-3Bab^2+Ab^3)(dx+c)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{2(Ba^6+3Ba^4b^2+3Aa^3b^3+6Ba^2b^4-3Aab^5)\log(b\tan(dx+c)+a)}{a^6b^3+3a^4b^5+3a^2b^7+b^9} - \frac{(Aa^3+3Ba^2b-3Aab^2-Bb^3)\log(\tan(dx+c)^2+1)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{2(2Ba^5b-Aa^4b^2+4Ba^3b^3-3Aa^2b^4)\tan(dx+c)}{a^6b^3+2a^4b^5+a^2b^7+(a^4b^5+2a^2b^7+b^9)\tan(dx+c)^2} + \frac{2(a^5b^4+2a^3b^6+ab^8)\tan(dx+c)}{d}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="maxima")
```

```
output 1/2*(2*(B*a^3 - 3*A*a^2*b - 3*B*a*b^2 + A*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + 2*(B*a^6 + 3*B*a^4*b^2 + A*a^3*b^3 + 6*B*a^2*b^4 - 3*A*a*b^5)*log(b*tan(d*x + c) + a)/(a^6*b^3 + 3*a^4*b^5 + 3*a^2*b^7 + b^9) - (A*a^3 + 3*B*a^2*b - 3*A*a*b^2 - B*b^3)*log(tan(d*x + c)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + (3*B*a^6 - A*a^5*b + 7*B*a^4*b^2 - 5*A*a^3*b^3 + 2*(2*B*a^5*b - A*a^4*b^2 + 4*B*a^3*b^3 - 3*A*a^2*b^4)*tan(d*x + c))/(a^6*b^3 + 2*a^4*b^5 + a^2*b^7 + (a^4*b^5 + 2*a^2*b^7 + b^9)*tan(d*x + c)^2 + 2*(a^5*b^4 + 2*a^3*b^6 + a*b^8)*tan(d*x + c))/d
```

3.283.8 Giac [A] (verification not implemented)

Time = 0.85 (sec) , antiderivative size = 458, normalized size of antiderivative = 1.83

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{2(Ba^3-3Aa^2b-3Bab^2+Ab^3)(dx+c)}{a^6+3a^4b^2+3a^2b^4+b^6} - \frac{(Aa^3+3Ba^2b-3Aab^2-Bb^3)\log(\tan(dx+c)^2+1)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{2(Ba^6+3Ba^4b^2+3Aa^3b^3+6Ba^2b^4-3Aab^5)\log(b\tan(dx+c)+a)}{a^6b^3+3a^4b^5+3a^2b^7+b^9} + \frac{2(2Ba^5b-Aa^4b^2+4Ba^3b^3-3Aa^2b^4)\tan(dx+c)}{a^6b^3+2a^4b^5+a^2b^7+(a^4b^5+2a^2b^7+b^9)\tan(dx+c)^2} + \frac{2(a^5b^4+2a^3b^6+ab^8)\tan(dx+c)}{d}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="giac")
```

output $\frac{1}{2} \cdot (2 \cdot (B \cdot a^3 - 3 \cdot A \cdot a^2 \cdot b - 3 \cdot B \cdot a \cdot b^2 + A \cdot b^3) \cdot (d \cdot x + c) / (a^6 + 3 \cdot a^4 \cdot b^2 + 3 \cdot a^2 \cdot b^4 + b^6) - (A \cdot a^3 + 3 \cdot B \cdot a^2 \cdot b - 3 \cdot A \cdot a \cdot b^2 - B \cdot b^3) \cdot \log(\tan(d \cdot x + c)^2 + 1) / (a^6 + 3 \cdot a^4 \cdot b^2 + 3 \cdot a^2 \cdot b^4 + b^6) + 2 \cdot (B \cdot a^6 + 3 \cdot B \cdot a^4 \cdot b^2 + A \cdot a^3 \cdot b^3 + 6 \cdot B \cdot a^2 \cdot b^4 - 3 \cdot A \cdot a \cdot b^5) \cdot \log(\text{abs}(b \cdot \tan(d \cdot x + c) + a)) / (a^6 \cdot b^3 + 3 \cdot a^4 \cdot b^5 + 3 \cdot a^2 \cdot b^7 + b^9) - (3 \cdot B \cdot a^6 \cdot b \cdot \tan(d \cdot x + c)^2 + 9 \cdot B \cdot a^4 \cdot b^3 \cdot \tan(d \cdot x + c)^2 + 3 \cdot A \cdot a^3 \cdot b^4 \cdot \tan(d \cdot x + c)^2 + 18 \cdot B \cdot a^2 \cdot b^5 \cdot \tan(d \cdot x + c)^2 - 9 \cdot A \cdot a \cdot b^6 \cdot \tan(d \cdot x + c)^2 + 2 \cdot B \cdot a^7 \cdot \tan(d \cdot x + c) + 2 \cdot A \cdot a^6 \cdot b \cdot \tan(d \cdot x + c) + 6 \cdot B \cdot a^5 \cdot b^2 \cdot \tan(d \cdot x + c) + 14 \cdot A \cdot a^4 \cdot b^3 \cdot \tan(d \cdot x + c) + 28 \cdot B \cdot a^3 \cdot b^4 \cdot \tan(d \cdot x + c) - 12 \cdot A \cdot a^2 \cdot b^5 \cdot \tan(d \cdot x + c) + A \cdot a^7 - B \cdot a^6 \cdot b + 9 \cdot A \cdot a^5 \cdot b^2 + 11 \cdot B \cdot a^4 \cdot b^3 - 4 \cdot A \cdot a^3 \cdot b^4) / ((a^6 \cdot b^2 + 3 \cdot a^4 \cdot b^4 + 3 \cdot a^2 \cdot b^6 + b^8) \cdot (b \cdot \tan(d \cdot x + c) + a)^2)) / d$

3.283.9 Mupad [B] (verification not implemented)

Time = 8.27 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.23

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{\frac{3Ba^6 - Aa^5b + 7Ba^4b^2 - 5Aa^3b^3}{2b^3(a^4 + 2a^2b^2 + b^4)} - \frac{a^2 \tan(c+dx)(-2Ba^3 + Aa^2b - 4Bab^2 + 3Ab^3)}{b^2(a^4 + 2a^2b^2 + b^4)}}{d(a^2 + 2ab \tan(c+dx) + b^2 \tan(c+dx)^2)}$$

$$+ \frac{\ln(\tan(c+dx) - i)(-B + A i)}{2d(-a^3 i + 3a^2b + ab^2 3i - b^3)} + \frac{\ln(\tan(c+dx) + 1i)(A - B i)}{2d(-a^3 + a^2b 3i + 3ab^2 - b^3 i)}$$

$$+ \frac{a \ln(a + b \tan(c+dx))(Ba^5 + 3Ba^3b^2 + Aa^2b^3 + 6Bab^4 - 3Ab^5)}{b^3 d(a^2 + b^2)^3}$$

input `int((tan(c + d*x))^3*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3,x`

output $((3 \cdot B \cdot a^6 - 5 \cdot A \cdot a^3 \cdot b^3 + 7 \cdot B \cdot a^4 \cdot b^2 - A \cdot a^5 \cdot b) / (2 \cdot b^3 \cdot (a^4 + b^4 + 2 \cdot a^2 \cdot b^2)) - (a^2 \cdot \tan(c + d \cdot x) \cdot (3 \cdot A \cdot b^3 - 2 \cdot B \cdot a^3 + A \cdot a^2 \cdot b - 4 \cdot B \cdot a \cdot b^2)) / (b^2 \cdot (a^4 + b^4 + 2 \cdot a^2 \cdot b^2))) / (d \cdot (a^2 + b^2 \cdot \tan(c + d \cdot x)^2 + 2 \cdot a \cdot b \cdot \tan(c + d \cdot x))) + (\log(\tan(c + d \cdot x) - 1i) \cdot (A \cdot 1i - B)) / (2 \cdot d \cdot (a \cdot b^2 \cdot 3i + 3 \cdot a^2 \cdot b - a^3 \cdot 1i - b^3)) + (\log(\tan(c + d \cdot x) + 1i) \cdot (A - B \cdot 1i)) / (2 \cdot d \cdot (3 \cdot a \cdot b^2 + a^2 \cdot b \cdot 3i - a^3 - b^3 \cdot 1i)) + (a \cdot \log(a + b \cdot \tan(c + d \cdot x)) \cdot (B \cdot a^5 - 3 \cdot A \cdot b^5 + A \cdot a^2 \cdot b^3 + 3 \cdot B \cdot a^3 \cdot b^2 + 6 \cdot B \cdot a \cdot b^4)) / (b^3 \cdot d \cdot (a^2 + b^2)^3)$

3.284 $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

3.284.1 Optimal result 2730
 3.284.2 Mathematica [C] (verified) 2731
 3.284.3 Rubi [A] (verified) 2731
 3.284.4 Maple [A] (verified) 2734
 3.284.5 Fricas [B] (verification not implemented) 2735
 3.284.6 Sympy [F(-2)] 2736
 3.284.7 Maxima [A] (verification not implemented) 2736
 3.284.8 Giac [B] (verification not implemented) 2737
 3.284.9 Mupad [B] (verification not implemented) 2737

3.284.1 Optimal result

Integrand size = 31, antiderivative size = 189

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= -\frac{(a^3A - 3aAb^2 + 3a^2bB - b^3B)x}{(a^2 + b^2)^3}$$

$$- \frac{(3a^2Ab - Ab^3 - a^3B + 3ab^2B) \log(a \cos(c+dx) + b \sin(c+dx))}{(a^2 + b^2)^3 d}$$

$$- \frac{a^2(Ab - aB)}{2b^2(a^2 + b^2)d(a + b \tan(c+dx))^2} + \frac{a(2Ab^3 - a(a^2 + 3b^2)B)}{b^2(a^2 + b^2)^2 d(a + b \tan(c+dx))}$$

output

```
-(A*a^3-3*A*a*b^2+3*B*a^2*b-B*b^3)*x/(a^2+b^2)^3-(3*A*a^2*b-A*b^3-B*a^3+3*
B*a*b^2)*ln(a*cos(d*x+c)+b*sin(d*x+c))/(a^2+b^2)^3/d-1/2*a^2*(A*b-B*a)/b^2
/(a^2+b^2)/d/(a+b*tan(d*x+c))^2+a*(2*A*b^3-a*(a^2+3*b^2)*B)/b^2/(a^2+b^2)^
2/d/(a+b*tan(d*x+c))
```

3.284.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 5.75 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.52

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$-\frac{Ab+aB}{b(a+b\tan(c+dx))^2} - \frac{2B\tan(c+dx)}{(a+b\tan(c+dx))^2} + B\left(\frac{i\log(i-\tan(c+dx))}{(a+ib)^2} - \frac{i\log(i+\tan(c+dx))}{(a-ib)^2} + \frac{2b(-2a\log(a+b\tan(c+dx))+\frac{a^2+b^2}{a+b\tan(c+dx)})}{(a^2+b^2)^2}\right)$$

input `Integrate[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

output `(-((A*b + a*B)/(b*(a + b*Tan[c + d*x])^2)) - (2*B*Tan[c + d*x])/(a + b*Tan[c + d*x])^2 + B*((I*Log[I - Tan[c + d*x]])/(a + I*b)^2 - (I*Log[I + Tan[c + d*x]])/(a - I*b)^2 + (2*b*(-2*a*Log[a + b*Tan[c + d*x]] + (a^2 + b^2)/(a + b*Tan[c + d*x])))/(a^2 + b^2)^2) + (A*b - a*B)*((I*Log[I - Tan[c + d*x]])/(a + I*b)^3 - Log[I + Tan[c + d*x]]/(I*a + b)^3 + (b*((-6*a^2 + 2*b^2)*Log[a + b*Tan[c + d*x]] + ((a^2 + b^2)*(5*a^2 + b^2 + 4*a*b*Tan[c + d*x])))/(a + b*Tan[c + d*x])^2))/(a^2 + b^2)^3)/(2*b*d)`

3.284.3 Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.14, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 4087, 25, 3042, 4111, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)^2(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$\downarrow \text{4087}$$

3.284. $\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{\int -\frac{((a^2+b^2)B \tan^2(c+dx)) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{(a+b \tan(c+dx))^2} dx}{b(a^2+b^2)} - \frac{a^2(Ab-aB)}{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow 25 \\
 & \frac{\int -\frac{((a^2+b^2)B \tan^2(c+dx)) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{(a+b \tan(c+dx))^2} dx}{b(a^2+b^2)} - \frac{a^2(Ab-aB)}{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow 3042 \\
 & \frac{\int -\frac{((a^2+b^2)B \tan(c+dx)^2) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{(a+b \tan(c+dx))^2} dx}{b(a^2+b^2)} - \frac{a^2(Ab-aB)}{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow 4111 \\
 & \frac{\int \frac{b(Aa^2+2bBa-Ab^2) - b(-Ba^2+2Aba+b^2B) \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2+b^2} - \frac{a(2Ab^3-aB(a^2+3b^2))}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \frac{b(a^2+b^2)}{a^2(Ab-aB)} \\
 & \quad \frac{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{b(Aa^2+2bBa-Ab^2) - b(-Ba^2+2Aba+b^2B) \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2+b^2} - \frac{a(2Ab^3-aB(a^2+3b^2))}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \frac{b(a^2+b^2)}{a^2(Ab-aB)} \\
 & \quad \frac{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow 4014 \\
 & \frac{b(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{a^2+b^2} \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx + \frac{bx(a^3A+3a^2bB-3aAb^2-b^3B)}{a^2+b^2} - \frac{a(2Ab^3-aB(a^2+3b^2))}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \frac{b(a^2+b^2)}{a^2(Ab-aB)} \\
 & \quad \frac{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow 3042 \\
 & \frac{b(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{a^2+b^2} \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx + \frac{bx(a^3A+3a^2bB-3aAb^2-b^3B)}{a^2+b^2} - \frac{a(2Ab^3-aB(a^2+3b^2))}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \frac{b(a^2+b^2)}{a^2(Ab-aB)} \\
 & \quad \frac{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow 4013
 \end{aligned}$$

3.284. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{\frac{a^2(Ab - aB)}{2b^2d(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{b(a^3(-B) + 3a^2Ab + 3ab^2B - Ab^3) \log(a \cos(c + dx) + b \sin(c + dx))}{d(a^2 + b^2)} + \frac{bx(a^3A + 3a^2bB - 3aAb^2 - b^3B)}{a^2 + b^2}}{a^2 + b^2} - \frac{a(2Ab^3 - aB(a^2 + 3b^2))}{bd(a^2 + b^2)(a + b \tan(c + dx))}$$

$$b(a^2 + b^2)$$

input `Int[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

output `-1/2*(a^2*(A*b - a*B))/(b^2*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) - (((b*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B)*x)/(a^2 + b^2) + (b*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d))/(a^2 + b^2) - (a*(2*A*b^3 - a*(a^2 + 3*b^2)*B))/(b*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(b*(a^2 + b^2))`

3.284.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

```
rule 4087 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(- (B*c - A*d))*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

```
rule 4111 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]
```

3.284.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.18

method	result
derivativedivides	$\frac{\frac{(3A a^2 b - A b^3 - B a^3 + 3B a b^2)}{2} \ln(1 + \tan^2(dx+c)) + (-A a^3 + 3A a b^2 - 3B a^2 b + B b^3) \arctan(\tan(dx+c))}{(a^2+b^2)^3} - \frac{a^2(Ab-Ba)}{2b^2(a^2+b^2)(a+b \tan(dx+c))} \frac{d}{d}$
default	$\frac{\frac{(3A a^2 b - A b^3 - B a^3 + 3B a b^2)}{2} \ln(1 + \tan^2(dx+c)) + (-A a^3 + 3A a b^2 - 3B a^2 b + B b^3) \arctan(\tan(dx+c))}{(a^2+b^2)^3} - \frac{a^2(Ab-Ba)}{2b^2(a^2+b^2)(a+b \tan(dx+c))} \frac{d}{d}$
norman	$-\frac{(2A a b^3 - B a^4 - 3B a^2 b^2)(\tan^2(dx+c))}{2a d(a^4 + 2a^2 b^2 + b^4)} - \frac{a(A a^3 - A a b^2 + 2B a^2 b)}{2d b(a^4 + 2a^2 b^2 + b^4)} - \frac{(A a^3 - 3A a b^2 + 3B a^2 b - B b^3) a^2 x}{(a^4 + 2a^2 b^2 + b^4)(a^2 + b^2)} - \frac{b^2(A a^3 - 3A a b^2 + 3B a^2 b - B b^3)}{(a^4 + 2a^2 b^2 + b^4)(a+b \tan(dx+c))^2}$
risch	$-\frac{i x B}{3i a^2 b - i b^3 - a^3 + 3a b^2} + \frac{x A}{3i a^2 b - i b^3 - a^3 + 3a b^2} + \frac{6i A a^2 b x}{a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6} - \frac{2i A b^3 x}{a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6} - \frac{2i B x}{a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6}$
parallelrisch	$-4A x \tan(dx+c) a^4 b^3 d + 12A x \tan(dx+c) a^2 b^5 d - 12B x \tan(dx+c) a^3 b^4 d + 4B x \tan(dx+c) a b^6 d + 6B \ln(1 + \tan^2(dx+c)) t$

```
input int(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x,method=_RETURNVERBOSE)
```

$$3.284. \int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

output $1/d*(1/(a^2+b^2)^3*(1/2*(3*A*a^2*b-A*b^3-B*a^3+3*B*a*b^2)*\ln(1+\tan(dx+c)^2)+(-A*a^3+3*A*a*b^2-3*B*a^2*b+B*b^3)*\arctan(\tan(dx+c)))-1/2*a^2*(A*b-B*a)/b^2/(a^2+b^2)/(a+b*\tan(dx+c))^2-(3*A*a^2*b-A*b^3-B*a^3+3*B*a*b^2)/(a^2+b^2)^3*\ln(a+b*\tan(dx+c))+a*(2*A*b^3-B*a^3-3*B*a*b^2)/(a^2+b^2)^2/b^2/(a+b*\tan(dx+c)))$

3.284.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 478 vs. $2(184) = 368$.

Time = 0.28 (sec) , antiderivative size = 478, normalized size of antiderivative = 2.53

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{Ba^5 - 3Aa^4b - 5Ba^3b^2 + 3Aa^2b^3 - 2(Aa^5 + 3Ba^4b - 3Aa^3b^2 - Ba^2b^3)dx + (Ba^5 + Aa^4b + 7Ba^3b^2 - 5Aa^2b^3 - 3Aa^3b^2 - Bba^4)dx + (Ba^5 - 3Aa^4b - 3Ba^3b^2 + Aa^2b^3 + (Ba^3b^2 - 3Aa^2b^3 - 3Ba^2b^4 + Ab^5)dx + 2*(Ba^4b - 3Aa^3b^2 - 3Ba^2b^3 + Aab^4)dx) \log((b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2)/(\tan(dx+c)^2 + 1)) + 2*(Aa^5 + 3Ba^4b - 3Aa^3b^2 - 3Ba^2b^3 + 2Aa^2b^4 - 2*(Aa^4b + 3Ba^3b^2 - 3Aa^2b^3 - Bba^4)dx) \tan(dx+c)}{(a^6b^2 + 3a^4b^4 + 3a^2b^6 + b^8)d \tan(dx+c)^2 + 2*(a^7b + 3a^5b^3 + 3a^3b^5 + ab^7)dx \tan(dx+c) + (a^8 + 3a^6b^2 + 3a^4b^4 + a^2b^6)d}$$

input `integrate(tan(dx+c)^2*(A+B*tan(dx+c))/(a+b*tan(dx+c))^3,x, algorithm="fracas")`

output $1/2*(B*a^5 - 3*A*a^4*b - 5*B*a^3*b^2 + 3*A*a^2*b^3 - 2*(A*a^5 + 3*B*a^4*b - 3*A*a^3*b^2 - B*a^2*b^3)*d*x + (B*a^5 + A*a^4*b + 7*B*a^3*b^2 - 5*A*a^2*b^3 - 2*(A*a^3*b^2 + 3*B*a^2*b^3 - 3*A*a*b^4 - B*b^5)*d*x)*\tan(dx + c)^2 + (B*a^5 - 3*A*a^4*b - 3*B*a^3*b^2 + A*a^2*b^3 + (B*a^3*b^2 - 3*A*a^2*b^3 - 3*B*a*b^4 + A*b^5)*\tan(dx + c))^2 + 2*(B*a^4*b - 3*A*a^3*b^2 - 3*B*a^2*b^3 + A*a*b^4)*\tan(dx + c)*\log((b^2*\tan(dx + c)^2 + 2*a*b*\tan(dx + c) + a^2)/(\tan(dx + c)^2 + 1)) + 2*(A*a^5 + 3*B*a^4*b - 3*A*a^3*b^2 - 3*B*a^2*b^3 + 2*A*a^2*b^4 - 2*(A*a^4*b + 3*B*a^3*b^2 - 3*A*a^2*b^3 - B*a*b^4)*d*x)*\tan(dx + c)/((a^6*b^2 + 3*a^4*b^4 + 3*a^2*b^6 + b^8)*d*\tan(dx + c)^2 + 2*(a^7*b + 3*a^5*b^3 + 3*a^3*b^5 + a*b^7)*d*\tan(dx + c) + (a^8 + 3*a^6*b^2 + 3*a^4*b^4 + a^2*b^6)*d)$

3.284.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Exception raised: AttributeError}$$

```
input integrate(tan(d*x+c)**2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)
```

```
output Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'
```

3.284.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 333, normalized size of antiderivative = 1.76

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx =$$

$$\frac{2(Aa^3+3Ba^2b-3Aab^2-Bb^3)(dx+c)}{a^6+3a^4b^2+3a^2b^4+b^6} - \frac{2(Ba^3-3Aa^2b-3Bab^2+Ab^3)\log(b\tan(dx+c)+a)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{(Ba^3-3Aa^2b-3Bab^2+Ab^3)\log(\tan(dx+c))}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{2d}{2d}$$

```
input integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="maxima")
```

```
output -1/2*(2*(A*a^3 + 3*B*a^2*b - 3*A*a*b^2 - B*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - 2*(B*a^3 - 3*A*a^2*b - 3*B*a*b^2 + A*b^3)*log(b*tan(d*x + c) + a)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + (B*a^3 - 3*A*a^2*b - 3*B*a*b^2 + A*b^3)*log(tan(d*x + c)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + (B*a^5 + A*a^4*b + 5*B*a^3*b^2 - 3*A*a^2*b^3 + 2*(B*a^4*b + 3*B*a^2*b^3 - 2*A*a*b^4)*tan(d*x + c))/(a^6*b^2 + 2*a^4*b^4 + a^2*b^6 + (a^4*b^4 + 2*a^2*b^6 + b^8)*tan(d*x + c)^2 + 2*(a^5*b^3 + 2*a^3*b^5 + a*b^7)*tan(d*x + c)))/d
```

3.284.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 410 vs. $2(184) = 368$.

Time = 0.70 (sec) , antiderivative size = 410, normalized size of antiderivative = 2.17

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx =$$

$$\frac{2(Aa^3+3Ba^2b-3Aab^2-Bb^3)(dx+c)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{(Ba^3-3Aa^2b-3Bab^2+Ab^3)\log(\tan(dx+c)^2+1)}{a^6+3a^4b^2+3a^2b^4+b^6} - \frac{2(Ba^3b-3Aa^2b^2-3Bab^3+Ab^4)\log(|b\tan(dx+c)+a|)}{a^6b+3a^4b^3+3a^2b^5+b^7}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="giac")`

output
$$\begin{aligned} & -1/2*(2*(A*a^3 + 3*B*a^2*b - 3*A*a*b^2 - B*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2 \\ & + 3*a^2*b^4 + b^6) + (B*a^3 - 3*A*a^2*b - 3*B*a*b^2 + A*b^3)*\log(\tan(d*x \\ & + c)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - 2*(B*a^3*b - 3*A*a^2*b^2 \\ & - 3*B*a*b^3 + A*b^4)*\log(\text{abs}(b*\tan(d*x + c) + a))/(a^6*b + 3*a^4*b^3 + 3* \\ & a^2*b^5 + b^7) + (3*B*a^3*b^4*\tan(d*x + c)^2 - 9*A*a^2*b^5*\tan(d*x + c)^2 \\ & - 9*B*a*b^6*\tan(d*x + c)^2 + 3*A*b^7*\tan(d*x + c)^2 + 2*B*a^6*b*\tan(d*x + \\ & c) + 14*B*a^4*b^3*\tan(d*x + c) - 22*A*a^3*b^4*\tan(d*x + c) - 12*B*a^2*b^5* \\ & \tan(d*x + c) + 2*A*a*b^6*\tan(d*x + c) + B*a^7 + A*a^6*b + 9*B*a^5*b^2 - 11 \\ & *A*a^4*b^3 - 4*B*a^3*b^4)/((a^6*b^2 + 3*a^4*b^4 + 3*a^2*b^6 + b^8)*(b*\tan(d*x \\ & + c) + a)^2))/d \end{aligned}$$

3.284.9 Mupad [B] (verification not implemented)

Time = 7.82 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.48

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{\ln(a+b\tan(c+dx))(Ba^3-3Aa^2b-3Bab^2+Ab^3)}{d(a^2+b^2)^3}$$

$$- \frac{\ln(\tan(c+dx)-i)(-B+Ali)}{2d(-a^3-a^2b^3i+3ab^2+b^3i)} - \frac{\ln(\tan(c+dx)+i)(A-Bli)}{2d(-a^3li-3a^2b+ab^2^3i+b^3)}$$

$$- \frac{\frac{a(Ba^4+Aa^3b+5Ba^2b^2-3Aab^3)}{2b^2(a^4+2a^2b^2+b^4)} + \frac{\tan(c+dx)(Ba^4+3Ba^2b^2-2Aab^3)}{b(a^4+2a^2b^2+b^4)}}{d(a^2+2ab\tan(c+dx)+b^2\tan(c+dx)^2)}$$

input `int((tan(c + d*x)^2*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3,x)`

output `(log(a + b*tan(c + d*x))*(A*b^3 + B*a^3 - 3*A*a^2*b - 3*B*a*b^2))/(d*(a^2 + b^2)^3) - (log(tan(c + d*x) - 1i)*(A*1i - B))/(2*d*(3*a*b^2 - a^2*b*3i - a^3 + b^3*1i)) - (log(tan(c + d*x) + 1i)*(A - B*1i))/(2*d*(a*b^2*3i - 3*a^2*b - a^3*1i + b^3)) - ((a*(B*a^4 + 5*B*a^2*b^2 - 3*A*a*b^3 + A*a^3*b))/(2*b^2*(a^4 + b^4 + 2*a^2*b^2)) + (tan(c + d*x)*(B*a^4 + 3*B*a^2*b^2 - 2*A*a*b^3))/(b*(a^4 + b^4 + 2*a^2*b^2)))/(d*(a^2 + b^2*tan(c + d*x)^2 + 2*a*b*tan(c + d*x)))`

3.285 $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

3.285.1 Optimal result 2739
 3.285.2 Mathematica [C] (verified) 2740
 3.285.3 Rubi [A] (verified) 2740
 3.285.4 Maple [A] (verified) 2743
 3.285.5 Fricas [B] (verification not implemented) 2743
 3.285.6 Sympy [F(-2)] 2744
 3.285.7 Maxima [A] (verification not implemented) 2744
 3.285.8 Giac [B] (verification not implemented) 2745
 3.285.9 Mupad [B] (verification not implemented) 2746

3.285.1 Optimal result

Integrand size = 29, antiderivative size = 179

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{(3a^2Ab - Ab^3 - a^3B + 3ab^2B)x}{(a^2 + b^2)^3}$$

$$- \frac{(a^3A - 3aAb^2 + 3a^2bB - b^3B) \log(a \cos(c+dx) + b \sin(c+dx))}{(a^2 + b^2)^3 d}$$

$$+ \frac{a(Ab - aB)}{2b(a^2 + b^2)d(a + b \tan(c+dx))^2} + \frac{a^2A - Ab^2 + 2abB}{(a^2 + b^2)^2 d(a + b \tan(c+dx))}$$

```
output (3*A*a^2*b-A*b^3-B*a^3+3*B*a*b^2)*x/(a^2+b^2)^3-(A*a^3-3*A*a*b^2+3*B*a^2*b
-B*b^3)*ln(a*cos(d*x+c)+b*sin(d*x+c))/(a^2+b^2)^3/d+1/2*a*(A*b-B*a)/b/(a^2
+b^2)/d/(a+b*tan(d*x+c))^2+(A*a^2-A*b^2+2*B*a*b)/(a^2+b^2)^2/d/(a+b*tan(d*
x+c))
```

3.285.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.97 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.05

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{\frac{(A+iB)\log(i-\tan(c+dx))}{(a+ib)^3} + \frac{(A-iB)\log(i+\tan(c+dx))}{(a-ib)^3} - \frac{2(a^3A-3aAb^2+3a^2bB-b^3B)\log(a+b\tan(c+dx))}{(a^2+b^2)^3} + \frac{a(Ab-aB)}{b(a^2+b^2)(a+b\tan(c+dx))^2}}{2d}$$

input `Integrate[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

output `((A + I*B)*Log[I - Tan[c + d*x]])/(a + I*b)^3 + ((A - I*B)*Log[I + Tan[c + d*x]])/(a - I*b)^3 - (2*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B)*Log[a + b*Tan[c + d*x]])/(a^2 + b^2)^3 + (a*(A*b - a*B))/(b*(a^2 + b^2)*(a + b*Tan[c + d*x])^2) + (2*(a^2*A - A*b^2 + 2*a*b*B))/((a^2 + b^2)^2*(a + b*Tan[c + d*x]))/(2*d)`

3.285.3 Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.12, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 4074, 3042, 4012, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$\downarrow \text{4074}$$

$$\frac{\int \frac{Ab-aB+(aA+bB)\tan(c+dx)}{(a+b\tan(c+dx))^2} dx}{a^2+b^2} + \frac{a(Ab-aB)}{2bd(a^2+b^2)(a+b\tan(c+dx))^2}$$

$$\downarrow \text{3042}$$

3.285. $\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{\int \frac{Ab - aB + (aA + bB) \tan(c + dx)}{(a + b \tan(c + dx))^2} dx}{a^2 + b^2} + \frac{a(Ab - aB)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} \\
 & \quad \downarrow 4012 \\
 & \frac{\int \frac{-Ba^2 + 2Aba + b^2B + (Aa^2 + 2bBa - Ab^2) \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{a^2A + 2abB - Ab^2}{d(a^2 + b^2)(a + b \tan(c + dx))} + \\
 & \quad \frac{a^2 + b^2}{a(Ab - aB)} \\
 & \quad \frac{2bd(a^2 + b^2)(a + b \tan(c + dx))^2}{} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{-Ba^2 + 2Aba + b^2B + (Aa^2 + 2bBa - Ab^2) \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{a^2A + 2abB - Ab^2}{d(a^2 + b^2)(a + b \tan(c + dx))} + \\
 & \quad \frac{a^2 + b^2}{a(Ab - aB)} \\
 & \quad \frac{2bd(a^2 + b^2)(a + b \tan(c + dx))^2}{} \\
 & \quad \downarrow 4014 \\
 & \frac{x \left(\frac{a^3(-B) + 3a^2Ab + 3ab^2B - Ab^3}{a^2 + b^2} \right) - \left(\frac{a^3A + 3a^2bB - 3aAb^2 - b^3B}{a^2 + b^2} \right) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{a^2A + 2abB - Ab^2}{d(a^2 + b^2)(a + b \tan(c + dx))} + \\
 & \quad \frac{a^2 + b^2}{a(Ab - aB)} \\
 & \quad \frac{2bd(a^2 + b^2)(a + b \tan(c + dx))^2}{} \\
 & \quad \downarrow 3042 \\
 & \frac{x \left(\frac{a^3(-B) + 3a^2Ab + 3ab^2B - Ab^3}{a^2 + b^2} \right) - \left(\frac{a^3A + 3a^2bB - 3aAb^2 - b^3B}{a^2 + b^2} \right) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{a^2A + 2abB - Ab^2}{d(a^2 + b^2)(a + b \tan(c + dx))} + \\
 & \quad \frac{a^2 + b^2}{a(Ab - aB)} \\
 & \quad \frac{2bd(a^2 + b^2)(a + b \tan(c + dx))^2}{} \\
 & \quad \downarrow 4013 \\
 & \frac{a^2A + 2abB - Ab^2}{d(a^2 + b^2)(a + b \tan(c + dx))} + \frac{a(Ab - aB)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} + \\
 & \quad \frac{x \left(\frac{a^3(-B) + 3a^2Ab + 3ab^2B - Ab^3}{a^2 + b^2} \right) - \left(\frac{a^3A + 3a^2bB - 3aAb^2 - b^3B}{a^2 + b^2} \right) \log(a \cos(c + dx) + b \sin(c + dx))}{a^2 + b^2} \\
 & \quad \frac{a^2 + b^2}{}
 \end{aligned}$$

input `Int[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

3.285. $\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$

```
output (a*(A*b - a*B))/(2*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + (((3*a^2*A*b
- A*b^3 - a^3*B + 3*a*b^2*B)*x)/(a^2 + b^2) - ((a^3*A - 3*a*A*b^2 + 3*a^2
*b*B - b^3*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d))/(a^2
+ b^2) + (a^2*A - A*b^2 + 2*a*b*B)/((a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(
a^2 + b^2)
```

3.285.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4012 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]
```

```
rule 4013 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*
(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Si
n[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

3.285.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.19

method	result
derivativedivides	$\frac{\frac{(A a^3 - 3A a b^2 + 3B a^2 b - B b^3) \ln(1 + \tan^2(dx+c))}{2} + (3A a^2 b - A b^3 - B a^3 + 3B a b^2) \arctan(\tan(dx+c))}{(a^2+b^2)^3} + \frac{a(Ab-Ba)}{2(a^2+b^2)b(a+b \tan(dx+c))} \frac{d}{d}$
default	$\frac{(A a^3 - 3A a b^2 + 3B a^2 b - B b^3) \ln(1 + \tan^2(dx+c))}{2} + (3A a^2 b - A b^3 - B a^3 + 3B a b^2) \arctan(\tan(dx+c))}{(a^2+b^2)^3} + \frac{a(Ab-Ba)}{2(a^2+b^2)b(a+b \tan(dx+c))} \frac{d}{d}$
norman	$\frac{(A a^2 b^2 - A b^4 + 2B a b^3) \tan(dx+c)}{db(a^4+2a^2b^2+b^4)} + \frac{(3A a^2 b - A b^3 - B a^3 + 3B a b^2) a^2 x}{(a^4+2a^2b^2+b^4)(a^2+b^2)} + \frac{b^2(3A a^2 b - A b^3 - B a^3 + 3B a b^2) x (\tan^2(dx+c))}{(a^4+2a^2b^2+b^4)(a^2+b^2)} + \frac{a(3A a^2 b^2 - A b^4 + 2B a b^3)}{(a+b \tan(dx+c))^2}$
risch	$\frac{x B}{3i a^2 b - i b^3 - a^3 + 3a b^2} + \frac{i x A}{3i a^2 b - i b^3 - a^3 + 3a b^2} + \frac{2i a^3 A x}{a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6} - \frac{6i a b^2 A x}{a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6} + \frac{6i a^2 b B x}{a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6}$
parallelrisch	$\frac{12A x \tan(dx+c) a^3 b^4 d - 4A x \tan(dx+c) a b^6 d + 12B x \tan(dx+c) a^2 b^5 d + 6A \ln(a+b \tan(dx+c)) a^3 b^4 - 6B \ln(a+b \tan(dx+c)) a^2 b^5}{(a+b \tan(dx+c))^3}$

input `int(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} \left(\frac{1}{(a^2+b^2)^3} \left(\frac{1}{2} (A a^3 - 3A a b^2 + 3B a^2 b - B b^3) \ln(1 + \tan^2(dx+c)) + (3A a^2 b - A b^3 - B a^3 + 3B a b^2) \arctan(\tan(dx+c)) \right) + \frac{1}{2} \frac{a(Ab-Ba)}{(a^2+b^2)b(a+b \tan(dx+c))} \right) + \frac{(A a^2 b^2 - A b^4 + 2B a b^3) \tan(dx+c)}{db(a^4+2a^2b^2+b^4)} + \frac{(3A a^2 b - A b^3 - B a^3 + 3B a b^2) a^2 x}{(a^4+2a^2b^2+b^4)(a^2+b^2)} + \frac{b^2(3A a^2 b - A b^3 - B a^3 + 3B a b^2) x (\tan^2(dx+c))}{(a^4+2a^2b^2+b^4)(a^2+b^2)} + \frac{a(3A a^2 b^2 - A b^4 + 2B a b^3)}{(a+b \tan(dx+c))^2}$$

3.285.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 488 vs. 2(176) = 352.

Time = 0.28 (sec) , antiderivative size = 488, normalized size of antiderivative = 2.73

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx = \frac{3Ba^4b - 5Aa^3b^2 - 3Ba^2b^3 + Aab^4 + 2(Ba^5 - 3Aa^4b - 3Ba^3b^2 + Aa^2b^3)dx - (Ba^4b - 3Aa^3b^2 - 5Aa^2b^3)}{(a+b \tan(c+dx))^3}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="fricas")`

3.285.
$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

```
output -1/2*(3*B*a^4*b - 5*A*a^3*b^2 - 3*B*a^2*b^3 + A*a*b^4 + 2*(B*a^5 - 3*A*a^4
*b - 3*B*a^3*b^2 + A*a^2*b^3)*d*x - (B*a^4*b - 3*A*a^3*b^2 - 5*B*a^2*b^3 +
3*A*a*b^4 - 2*(B*a^3*b^2 - 3*A*a^2*b^3 - 3*B*a*b^4 + A*b^5)*d*x)*tan(d*x
+ c)^2 + (A*a^5 + 3*B*a^4*b - 3*A*a^3*b^2 - B*a^2*b^3 + (A*a^3*b^2 + 3*B*a
^2*b^3 - 3*A*a*b^4 - B*b^5)*tan(d*x + c)^2 + 2*(A*a^4*b + 3*B*a^3*b^2 - 3*
A*a^2*b^3 - B*a*b^4)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x
+ c) + a^2)/(tan(d*x + c)^2 + 1)) - 2*(B*a^5 - 2*A*a^4*b - 3*B*a^3*b^2 +
3*A*a^2*b^3 + 2*B*a*b^4 - A*b^5 - 2*(B*a^4*b - 3*A*a^3*b^2 - 3*B*a^2*b^3 +
A*a*b^4)*d*x)*tan(d*x + c))/((a^6*b^2 + 3*a^4*b^4 + 3*a^2*b^6 + b^8)*d*ta
n(d*x + c)^2 + 2*(a^7*b + 3*a^5*b^3 + 3*a^3*b^5 + a*b^7)*d*tan(d*x + c) +
(a^8 + 3*a^6*b^2 + 3*a^4*b^4 + a^2*b^6)*d)
```

3.285.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx = \text{Exception raised: AttributeError}$$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)
```

```
output Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'
```

3.285.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 330, normalized size of antiderivative = 1.84

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx =$$

$$\frac{2(Ba^3 - 3Aa^2b - 3Bab^2 + Ab^3)(dx+c)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} + \frac{2(Aa^3 + 3Ba^2b - 3Aab^2 - Bb^3) \log(b \tan(dx+c)+a)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} - \frac{(Aa^3 + 3Ba^2b - 3Aab^2 - Bb^3) \log(\tan(dx+c))}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6}$$

$2d$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="max
ima")
```

3.285. $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

output
$$\begin{aligned} & -1/2*(2*(B*a^3 - 3*A*a^2*b - 3*B*a*b^2 + A*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2 \\ & + 3*a^2*b^4 + b^6) + 2*(A*a^3 + 3*B*a^2*b - 3*A*a*b^2 - B*b^3)*\log(b*\tan(\\ & d*x + c) + a)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - (A*a^3 + 3*B*a^2*b - 3 \\ & *A*a*b^2 - B*b^3)*\log(\tan(d*x + c)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b \\ & ^6) + (B*a^4 - 3*A*a^3*b - 3*B*a^2*b^2 + A*a*b^3 - 2*(A*a^2*b^2 + 2*B*a*b^ \\ & 3 - A*b^4)*\tan(d*x + c))/(a^6*b + 2*a^4*b^3 + a^2*b^5 + (a^4*b^3 + 2*a^2*b \\ & ^5 + b^7)*\tan(d*x + c)^2 + 2*(a^5*b^2 + 2*a^3*b^4 + a*b^6)*\tan(d*x + c))/ \\ & d \end{aligned}$$

3.285.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 410 vs. $2(176) = 352$.

Time = 0.58 (sec) , antiderivative size = 410, normalized size of antiderivative = 2.29

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \frac{2(Ba^3-3Aa^2b-3Bab^2+Ab^3)(dx+c)}{a^6+3a^4b^2+3a^2b^4+b^6} - \frac{(Aa^3+3Ba^2b-3Aab^2-Bb^3)\log(\tan(dx+c)^2+1)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{2(Aa^3b+3Ba^2b^2-3Aab^3-Bb^4)\log(|b\tan(dx+c)+a|)}{a^6b+3a^4b^3+3a^2b^5+b^7}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="giac")`

output
$$\begin{aligned} & -1/2*(2*(B*a^3 - 3*A*a^2*b - 3*B*a*b^2 + A*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2 \\ & + 3*a^2*b^4 + b^6) - (A*a^3 + 3*B*a^2*b - 3*A*a*b^2 - B*b^3)*\log(\tan(d*x \\ & + c)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + 2*(A*a^3*b + 3*B*a^2*b^2 \\ & - 3*A*a*b^3 - B*b^4)*\log(\text{abs}(b*\tan(d*x + c) + a))/(a^6*b + 3*a^4*b^3 + 3* \\ & a^2*b^5 + b^7) - (3*A*a^3*b^3*\tan(d*x + c)^2 + 9*B*a^2*b^4*\tan(d*x + c)^2 \\ & - 9*A*a*b^5*\tan(d*x + c)^2 - 3*B*b^6*\tan(d*x + c)^2 + 8*A*a^4*b^2*\tan(d*x \\ & + c) + 22*B*a^3*b^3*\tan(d*x + c) - 18*A*a^2*b^4*\tan(d*x + c) - 2*B*a*b^5*\tan \\ & (d*x + c) - 2*A*b^6*\tan(d*x + c) - B*a^6 + 6*A*a^5*b + 11*B*a^4*b^2 - 7* \\ & A*a^3*b^3 - A*a*b^5)/((a^6*b + 3*a^4*b^3 + 3*a^2*b^5 + b^7)*(b*\tan(d*x + c) \\ & + a)^2))/d \end{aligned}$$

3.285.
$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

3.285.9 Mupad [B] (verification not implemented)

Time = 7.88 (sec) , antiderivative size = 282, normalized size of antiderivative = 1.58

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{\frac{\tan(c+dx)(Aa^2b+2Bab^2-Ab^3)}{a^4+2a^2b^2+b^4} - \frac{Ba^4-3Aa^3b-3Ba^2b^2+Aab^3}{2b(a^4+2a^2b^2+b^4)}}{d(a^2+2ab\tan(c+dx)+b^2\tan(c+dx)^2)}$$

$$- \frac{\ln(a+b\tan(c+dx))\left(\frac{Aa+3Bb}{(a^2+b^2)^2} - \frac{4b^2(Aa+Bb)}{(a^2+b^2)^3}\right)}{d}$$

$$- \frac{\ln(\tan(c+dx)-i)(-B+Ai)}{2d(-a^3i+3a^2b+ab^23i-b^3)} - \frac{\ln(\tan(c+dx)+i)(A-Bi)}{2d(-a^3+a^2b3i+3ab^2-b^3i)}$$

input `int((tan(c + d*x)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3,x)`output `((tan(c + d*x)*(A*a^2*b - A*b^3 + 2*B*a*b^2))/(a^4 + b^4 + 2*a^2*b^2) - (B*a^4 - 3*B*a^2*b^2 + A*a*b^3 - 3*A*a^3*b)/(2*b*(a^4 + b^4 + 2*a^2*b^2)))/(d*(a^2 + b^2*tan(c + d*x)^2 + 2*a*b*tan(c + d*x))) - (log(a + b*tan(c + d*x))*((A*a + 3*B*b)/(a^2 + b^2)^2 - (4*b^2*(A*a + B*b))/(a^2 + b^2)^3))/d - (log(tan(c + d*x) - 1i)*(A*1i - B))/(2*d*(a*b^2*3i + 3*a^2*b - a^3*1i - b^3)) - (log(tan(c + d*x) + 1i)*(A - B*1i))/(2*d*(3*a*b^2 + a^2*b*3i - a^3 - b^3*1i))`

3.286 $\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^3} dx$

3.286.1 Optimal result 2747
 3.286.2 Mathematica [C] (verified) 2748
 3.286.3 Rubi [A] (verified) 2748
 3.286.4 Maple [A] (verified) 2751
 3.286.5 Fricas [B] (verification not implemented) 2751
 3.286.6 Sympy [F(-2)] 2752
 3.286.7 Maxima [A] (verification not implemented) 2752
 3.286.8 Giac [B] (verification not implemented) 2753
 3.286.9 Mupad [B] (verification not implemented) 2754

3.286.1 Optimal result

Integrand size = 23, antiderivative size = 175

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{(a^3 A - 3aAb^2 + 3a^2bB - b^3 B) x}{(a^2 + b^2)^3} + \frac{(3a^2 Ab - Ab^3 - a^3 B + 3ab^2 B) \log(a \cos(c + dx) + b \sin(c + dx))}{(a^2 + b^2)^3 d}$$

$$- \frac{Ab - aB}{2(a^2 + b^2) d(a + b \tan(c + dx))^2} - \frac{2aAb - a^2 B + b^2 B}{(a^2 + b^2)^2 d(a + b \tan(c + dx))}$$

output `(A*a^3-3*A*a*b^2+3*B*a^2*b-B*b^3)*x/(a^2+b^2)^3+(3*A*a^2*b-A*b^3-B*a^3+3*B*a*b^2)*ln(a*cos(d*x+c)+b*sin(d*x+c))/(a^2+b^2)^3/d+1/2*(-A*b+B*a)/(a^2+b^2)/d/(a+b*tan(d*x+c))^2+(-2*A*a*b+B*a^2-B*b^2)/(a^2+b^2)^2/d/(a+b*tan(d*x+c))`

3.286.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.04 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.39

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^3} dx =$$

$$\frac{B \left(\frac{i \log(i - \tan(c + dx))}{(a + ib)^2} - \frac{i \log(i + \tan(c + dx))}{(a - ib)^2} + \frac{2b \left(-2a \log(a + b \tan(c + dx)) + \frac{a^2 + b^2}{a + b \tan(c + dx)} \right)}{(a^2 + b^2)^2} \right) + (Ab - aB) \left(\frac{i \log(i - \tan(c + dx))}{(a + ib)^3} \right)}{2bd}$$

input `Integrate[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^3,x]`

output `-1/2*(B*((I*Log[I - Tan[c + d*x]])/(a + I*b)^2 - (I*Log[I + Tan[c + d*x]])/(a - I*b)^2 + (2*b*(-2*a*Log[a + b*Tan[c + d*x]] + (a^2 + b^2)/(a + b*Tan[c + d*x])))/(a^2 + b^2)^2) + (A*b - a*B)*((I*Log[I - Tan[c + d*x]])/(a + I*b)^3 - Log[I + Tan[c + d*x]]/(I*a + b)^3 + (b*((-6*a^2 + 2*b^2)*Log[a + b*Tan[c + d*x]] + ((a^2 + b^2)*(5*a^2 + b^2 + 4*a*b*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2))/(a^2 + b^2)^3)/(b*d)`

3.286.3 Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.13, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.348$, Rules used = {3042, 4012, 3042, 4012, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^3} dx$$

$$\downarrow \text{4012}$$

$$\frac{\int \frac{aA + bB - (Ab - aB) \tan(c + dx)}{(a + b \tan(c + dx))^2} dx}{a^2 + b^2} - \frac{Ab - aB}{2d(a^2 + b^2)(a + b \tan(c + dx))^2}$$

$$\begin{aligned}
& \int \frac{aA+bB-(Ab-aB)\tan(c+dx)}{(a+b\tan(c+dx))^2} dx \quad \downarrow \text{3042} \\
& \frac{Ab-aB}{2d(a^2+b^2)(a+b\tan(c+dx))^2} \\
& \downarrow \text{4012} \\
& \frac{\int \frac{Aa^2+2bBa-Ab^2-(-Ba^2+2Aba+b^2B)\tan(c+dx)}{a+b\tan(c+dx)} dx}{a^2+b^2} - \frac{a^2(-B)+2aAb+b^2B}{d(a^2+b^2)(a+b\tan(c+dx))} \quad \downarrow \text{3042} \\
& \frac{Ab-aB}{2d(a^2+b^2)(a+b\tan(c+dx))^2} \\
& \downarrow \text{3042} \\
& \frac{\int \frac{Aa^2+2bBa-Ab^2-(-Ba^2+2Aba+b^2B)\tan(c+dx)}{a+b\tan(c+dx)} dx}{a^2+b^2} - \frac{a^2(-B)+2aAb+b^2B}{d(a^2+b^2)(a+b\tan(c+dx))} \quad \downarrow \text{4014} \\
& \frac{\frac{(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a^2+b^2} + \frac{x(a^3A+3a^2bB-3aAb^2-b^3B)}{a^2+b^2}}{a^2+b^2} - \frac{a^2(-B)+2aAb+b^2B}{d(a^2+b^2)(a+b\tan(c+dx))} \\
& \frac{a^2+b^2}{Ab-aB} \\
& \frac{Ab-aB}{2d(a^2+b^2)(a+b\tan(c+dx))^2} \quad \downarrow \text{3042} \\
& \frac{(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a^2+b^2} + \frac{x(a^3A+3a^2bB-3aAb^2-b^3B)}{a^2+b^2} - \frac{a^2(-B)+2aAb+b^2B}{d(a^2+b^2)(a+b\tan(c+dx))} \\
& \frac{a^2+b^2}{Ab-aB} \\
& \frac{Ab-aB}{2d(a^2+b^2)(a+b\tan(c+dx))^2} \quad \downarrow \text{4013} \\
& \frac{(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \log(a \cos(c+dx)+b \sin(c+dx))}{d(a^2+b^2)} + \frac{x(a^3A+3a^2bB-3aAb^2-b^3B)}{a^2+b^2} - \frac{a^2(-B)+2aAb+b^2B}{d(a^2+b^2)(a+b\tan(c+dx))} \\
& \frac{a^2+b^2}{Ab-aB} \\
& \frac{Ab-aB}{2d(a^2+b^2)(a+b\tan(c+dx))^2}
\end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^3,x]`

```
output -1/2*(A*b - a*B)/((a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + (((a^3*A - 3*a*
A*b^2 + 3*a^2*b*B - b^3*B)*x)/(a^2 + b^2) + ((3*a^2*A*b - A*b^3 - a^3*B +
3*a*b^2*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d))/(a^2 + b
^2) - (2*a*A*b - a^2*B + b^2*B)/((a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(a^2
+ b^2)
```

3.286.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4012 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]
```

```
rule 4013 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*
(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Si
n[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

3.286.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.19

method	result
derivativedivides	$\frac{\frac{(-3Aa^2b+Ab^3+B a^3-3Ba b^2) \ln(1+\tan^2(dx+c))}{2} + (Aa^3-3Aa b^2+3B a^2b-B b^3) \arctan(\tan(dx+c))}{(a^2+b^2)^3} + \frac{(3Aa^2b-Ab^3-B a^3+3Ba^2b)}{d(a^2+b^2)}$
default	$\frac{\frac{(-3Aa^2b+Ab^3+B a^3-3Ba b^2) \ln(1+\tan^2(dx+c))}{2} + (Aa^3-3Aa b^2+3B a^2b-B b^3) \arctan(\tan(dx+c))}{(a^2+b^2)^3} + \frac{(3Aa^2b-Ab^3-B a^3+3Ba^2b)}{d(a^2+b^2)}$
norman	$\frac{(Aa^3-3Aa b^2+3B a^2b-B b^3)a^2x}{(a^4+2a^2b^2+b^4)(a^2+b^2)} + \frac{b^2(Aa^3-3Aa b^2+3B a^2b-B b^3)x(\tan^2(dx+c))}{(a^4+2a^2b^2+b^4)(a^2+b^2)} - \frac{3Aa^2b^2+Ab^4-2Ba^3b}{2bd(a^4+2a^2b^2+b^4)} + \frac{b(2Aa b^2-B a^2b)}{2da(a^4+2a^2b^2+b^4)} - \frac{3Aa^2b^2+Ab^4-2Ba^3b}{2bd(a^4+2a^2b^2+b^4)} + \frac{b(2Aa b^2-B a^2b)}{2da(a^4+2a^2b^2+b^4)} - \frac{3Aa^2b^2+Ab^4-2Ba^3b}{2bd(a^4+2a^2b^2+b^4)} + \frac{b(2Aa b^2-B a^2b)}{2da(a^4+2a^2b^2+b^4)} - \frac{3Aa^2b^2+Ab^4-2Ba^3b}{2bd(a^4+2a^2b^2+b^4)} + \frac{b(2Aa b^2-B a^2b)}{2da(a^4+2a^2b^2+b^4)}$
risch	$\frac{ixB}{3ia^2b-ib^3-a^3+3ab^2} - \frac{x A}{3ia^2b-ib^3-a^3+3ab^2} - \frac{6iAa^2bx}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{2iAb^3x}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{2iBxa^3}{a^6+3a^4b^2+3a^2b^4+b^6}$
parallelrisch	Expression too large to display

input `int((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`output
$$\frac{1}{d} \left(\frac{1}{(a^2+b^2)^3} \left(\frac{1}{2} (-3Aa^2b+Ab^3+B a^3-3Ba b^2) \ln(1+\tan^2(dx+c)) + (Aa^3-3Aa b^2+3B a^2b-B b^3) \arctan(\tan(dx+c)) \right) + \frac{(3Aa^2b-Ab^3-B a^3+3Ba^2b)}{d(a^2+b^2)} \right) - \frac{1}{2} \frac{(Aa^3-3Aa b^2+3B a^2b-B b^3)}{(a^2+b^2)^2} \frac{\ln(a+b \tan(dx+c))}{(a+b \tan(dx+c))} - \frac{1}{2} \frac{(Aa^3-3Aa b^2+3B a^2b-B b^3)}{(a^2+b^2)^2} \frac{\arctan(\tan(dx+c))}{(a+b \tan(dx+c))}$$
3.286.5 Fracas [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 482 vs. $2(171) = 342$.

Time = 0.27 (sec) , antiderivative size = 482, normalized size of antiderivative = 2.75

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{5Ba^3b^2 - 7Aa^2b^3 - Bab^4 - Ab^5 + 2(Aa^5 + 3Ba^4b - 3Aa^3b^2 - Ba^2b^3)dx - (3Ba^3b^2 - 5Aa^2b^3 - 3Ba^2b^2)}{(a + b \tan(c + dx))^3}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="fricas")`

```
output 1/2*(5*B*a^3*b^2 - 7*A*a^2*b^3 - B*a*b^4 - A*b^5 + 2*(A*a^5 + 3*B*a^4*b -
3*A*a^3*b^2 - B*a^2*b^3)*d*x - (3*B*a^3*b^2 - 5*A*a^2*b^3 - 3*B*a*b^4 + A*
b^5 - 2*(A*a^3*b^2 + 3*B*a^2*b^3 - 3*A*a*b^4 - B*b^5)*d*x)*tan(d*x + c)^2
- (B*a^5 - 3*A*a^4*b - 3*B*a^3*b^2 + A*a^2*b^3 + (B*a^3*b^2 - 3*A*a^2*b^3
- 3*B*a*b^4 + A*b^5)*tan(d*x + c)^2 + 2*(B*a^4*b - 3*A*a^3*b^2 - 3*B*a^2*b
^3 + A*a*b^4)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) +
a^2)/(tan(d*x + c)^2 + 1)) - 2*(2*B*a^4*b - 3*A*a^3*b^2 - 3*B*a^2*b^3 + 3
*A*a*b^4 + B*b^5 - 2*(A*a^4*b + 3*B*a^3*b^2 - 3*A*a^2*b^3 - B*a*b^4)*d*x)*
tan(d*x + c))/((a^6*b^2 + 3*a^4*b^4 + 3*a^2*b^6 + b^8)*d*tan(d*x + c)^2 +
2*(a^7*b + 3*a^5*b^3 + 3*a^3*b^5 + a*b^7)*d*tan(d*x + c) + (a^8 + 3*a^6*b^
2 + 3*a^4*b^4 + a^2*b^6)*d)
```

3.286.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^3} dx = \text{Exception raised: AttributeError}$$

```
input integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)
```

```
output Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'
```

3.286.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 321, normalized size of antiderivative = 1.83

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{2(Aa^3 + 3Ba^2b - 3Aab^2 - Bb^3)(dx+c)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} - \frac{2(Ba^3 - 3Aa^2b - 3Bab^2 + Ab^3) \log(b \tan(dx+c) + a)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} + \frac{(Ba^3 - 3Aa^2b - 3Bab^2 + Ab^3) \log(\tan(dx+c))}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6}$$

$2d$

```
input integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="maxima")
```

output $\frac{1}{2} * (2 * (A * a^3 + 3 * B * a^2 * b - 3 * A * a * b^2 - B * b^3) * (d * x + c) / (a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) - 2 * (B * a^3 - 3 * A * a^2 * b - 3 * B * a * b^2 + A * b^3) * \log(b * \tan(d * x + c) + a) / (a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) + (B * a^3 - 3 * A * a^2 * b - 3 * B * a * b^2 + A * b^3) * \log(\tan(d * x + c)^2 + 1) / (a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) + (3 * B * a^3 - 5 * A * a^2 * b - B * a * b^2 - A * b^3 + 2 * (B * a^2 * b - 2 * A * a * b^2 - B * b^3) * \tan(d * x + c)) / (a^6 + 2 * a^4 * b^2 + a^2 * b^4 + (a^4 * b^2 + 2 * a^2 * b^4 + b^6) * \tan(d * x + c)^2 + 2 * (a^5 * b + 2 * a^3 * b^3 + a * b^5) * \tan(d * x + c))) / d$

3.286.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 409 vs. $2(171) = 342$.

Time = 0.54 (sec) , antiderivative size = 409, normalized size of antiderivative = 2.34

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{2(Aa^3 + 3Ba^2b - 3Aab^2 - Bb^3)(dx+c)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} + \frac{(Ba^3 - 3Aa^2b - 3Bab^2 + Ab^3) \log(\tan(dx+c)^2 + 1)}{a^6 + 3a^4b^2 + 3a^2b^4 + b^6} - \frac{2(Ba^3b - 3Aa^2b^2 - 3Bab^3 + Ab^4) \log(|b \tan(dx+c) + a|)}{a^6b + 3a^4b^3 + 3a^2b^5 + b^7}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="giac")`

output $\frac{1}{2} * (2 * (A * a^3 + 3 * B * a^2 * b - 3 * A * a * b^2 - B * b^3) * (d * x + c) / (a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) + (B * a^3 - 3 * A * a^2 * b - 3 * B * a * b^2 + A * b^3) * \log(\tan(d * x + c)^2 + 1) / (a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) - 2 * (B * a^3 * b - 3 * A * a^2 * b^2 - 3 * B * a * b^3 + A * b^4) * \log(\text{abs}(b * \tan(d * x + c) + a)) / (a^6 * b + 3 * a^4 * b^3 + 3 * a^2 * b^5 + b^7) + (3 * B * a^3 * b^2 * \tan(d * x + c)^2 - 9 * A * a^2 * b^3 * \tan(d * x + c)^2 - 9 * B * a * b^4 * \tan(d * x + c)^2 + 3 * A * b^5 * \tan(d * x + c)^2 + 8 * B * a^4 * b * \tan(d * x + c) - 22 * A * a^3 * b^2 * \tan(d * x + c) - 18 * B * a^2 * b^3 * \tan(d * x + c) + 2 * A * a * b^4 * \tan(d * x + c) - 2 * B * b^5 * \tan(d * x + c) + 6 * B * a^5 - 14 * A * a^4 * b - 7 * B * a^3 * b^2 - 3 * A * a^2 * b^3 - B * a * b^4 - A * b^5) / ((a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) * (b * \tan(d * x + c) + a)^2)) / d$

3.286.9 Mupad [B] (verification not implemented)

Time = 8.25 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.59

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^3} dx = \frac{\ln(a + b \tan(c + dx)) \left(\frac{3Ab - Ba}{(a^2 + b^2)^2} - \frac{4b^2(Ab - Ba)}{(a^2 + b^2)^3} \right)}{d} - \frac{\frac{-3Ba^3 + 5Aa^2b + BAb^2 + Ab^3}{2(a^4 + 2a^2b^2 + b^4)} + \frac{\tan(c + dx)(-Ba^2b + 2Aab^2 + Bb^3)}{a^4 + 2a^2b^2 + b^4}}{d(a^2 + 2ab \tan(c + dx) + b^2 \tan(c + dx)^2)} + \frac{\ln(\tan(c + dx) - i)(-B + A i)}{2d(-a^3 - a^2b^3i + 3ab^2 + b^3i)} + \frac{\ln(\tan(c + dx) + i)(A - B i)}{2d(-a^3 i - 3a^2b + ab^2^3i + b^3)}$$

input `int((A + B*tan(c + d*x))/(a + b*tan(c + d*x))^3,x)`output `(log(a + b*tan(c + d*x))*((3*A*b - B*a)/(a^2 + b^2)^2 - (4*b^2*(A*b - B*a))/(a^2 + b^2)^3))/d - ((A*b^3 - 3*B*a^3 + 5*A*a^2*b + B*a*b^2)/(2*(a^4 + b^4 + 2*a^2*b^2)) + (tan(c + d*x)*(B*b^3 + 2*A*a*b^2 - B*a^2*b))/(a^4 + b^4 + 2*a^2*b^2))/(d*(a^2 + b^2*tan(c + d*x)^2 + 2*a*b*tan(c + d*x))) + (log(tan(c + d*x) - 1i)*(A*1i - B))/(2*d*(3*a*b^2 - a^2*b^3i - a^3 + b^3*1i)) + (log(tan(c + d*x) + 1i)*(A - B*1i))/(2*d*(a*b^2*3i - 3*a^2*b - a^3*1i + b^3))`

3.287 $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

3.287.1 Optimal result 2755
 3.287.2 Mathematica [C] (verified) 2756
 3.287.3 Rubi [A] (verified) 2756
 3.287.4 Maple [A] (verified) 2760
 3.287.5 Fricas [B] (verification not implemented) 2761
 3.287.6 Sympy [F(-2)] 2761
 3.287.7 Maxima [A] (verification not implemented) 2762
 3.287.8 Giac [B] (verification not implemented) 2762
 3.287.9 Mupad [B] (verification not implemented) 2763

3.287.1 Optimal result

Integrand size = 29, antiderivative size = 215

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= -\frac{(3a^2Ab - Ab^3 - a^3B + 3ab^2B)x}{(a^2 + b^2)^3} + \frac{A \log(\sin(c+dx))}{a^3d}$$

$$- \frac{b(6a^4Ab + 3a^2Ab^3 + Ab^5 - 3a^5B + a^3b^2B) \log(a \cos(c+dx) + b \sin(c+dx))}{a^3(a^2 + b^2)^3d}$$

$$+ \frac{b(Ab - aB)}{2a(a^2 + b^2)d(a+b \tan(c+dx))^2} + \frac{b(3a^2Ab + Ab^3 - 2a^3B)}{a^2(a^2 + b^2)^2d(a+b \tan(c+dx))}$$

output

```
-(3*A*a^2*b-A*b^3-B*a^3+3*B*a*b^2)*x/(a^2+b^2)^3+A*ln(sin(d*x+c))/a^3/d-b*
(6*A*a^4*b+3*A*a^2*b^3+A*b^5-3*B*a^5+B*a^3*b^2)*ln(a*cos(d*x+c)+b*sin(d*x+
c))/a^3/(a^2+b^2)^3/d+1/2*b*(A*b-B*a)/a/(a^2+b^2)/d/(a+b*tan(d*x+c))^2+b*(
3*A*a^2*b+A*b^3-2*B*a^3)/a^2/(a^2+b^2)^2/d/(a+b*tan(d*x+c))
```

3.287.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.88 (sec) , antiderivative size = 254, normalized size of antiderivative = 1.18

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{-\frac{a(a-ib)(A+iB)\log(i-\tan(c+dx))}{(a+ib)^2} + \frac{2A(a^2+b^2)\log(\tan(c+dx))}{a^2} - \frac{a(a+ib)(A-iB)\log(i+\tan(c+dx))}{(a-ib)^2} - \frac{2b(6a^4Ab+3a^2Ab^3+Ab^5-3a^5B)}{a^2(a^2+b^2)}}{2a(a^2+b^2)d}$$

input `Integrate[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

output
$$\begin{aligned} & -((a*(a - I*b)*(A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]])/(a + I*b)^2) + (2*A*(a^2 + b^2)*\text{Log}[\text{Tan}[c + d*x]])/a^2 - (a*(a + I*b)*(A - I*B)*\text{Log}[I + \text{Tan}[c + d*x]])/(a - I*b)^2 - (2*b*(6*a^4*A*b + 3*a^2*A*b^3 + A*b^5 - 3*a^5*B + a^3*b^2*B)*\text{Log}[a + b*\text{Tan}[c + d*x]])/(a^2*(a^2 + b^2)^2) + (b*(A*b - a*B))/(a + b*\text{Tan}[c + d*x])^2 + (4*a*b*(A*b - a*B))/((a^2 + b^2)*(a + b*\text{Tan}[c + d*x])) + (2*A*b^2)/(a^2 + a*b*\text{Tan}[c + d*x])/(2*a*(a^2 + b^2)*d) \end{aligned}$$

3.287.3 Rubi [A] (verified)

Time = 1.34 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.20, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.379$, Rules used = {3042, 4092, 27, 3042, 4132, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)(a + b \tan(c + dx))^3} dx$$

↓ 4092

$$\int \frac{2 \cot(c + dx)(b(Ab - aB) \tan^2(c + dx) - a(Ab - aB) \tan(c + dx) + A(a^2 + b^2))}{(a + b \tan(c + dx))^2} dx + \frac{b(Ab - aB)}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 27

3.287. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{\int \frac{\cot(c+dx)(b(Ab-aB)\tan^2(c+dx)-a(Ab-aB)\tan(c+dx)+A(a^2+b^2))}{(a+b\tan(c+dx))^2} dx}{a(a^2+b^2)} + \frac{b(Ab-aB)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 3042

$$\frac{\int \frac{b(Ab-aB)\tan(c+dx)^2-a(Ab-aB)\tan(c+dx)+A(a^2+b^2)}{\tan(c+dx)(a+b\tan(c+dx))^2} dx}{a(a^2+b^2)} + \frac{b(Ab-aB)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 4132

$$\frac{\int \frac{\cot(c+dx)\left(-((-Ba^2+2Aba+b^2B)\tan(c+dx)a^2)+A(a^2+b^2)^2+b(-2Ba^3+3Aba^2+Ab^3)\tan^2(c+dx)\right)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} + \frac{b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)(a+b\tan(c+dx))} +$$

$$\frac{a(a^2+b^2)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2} \frac{b(Ab-aB)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 3042

$$\frac{\int \frac{-((-Ba^2+2Aba+b^2B)\tan(c+dx)a^2)+A(a^2+b^2)^2+b(-2Ba^3+3Aba^2+Ab^3)\tan(c+dx)^2}{\tan(c+dx)(a+b\tan(c+dx))} dx}{a(a^2+b^2)} + \frac{b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)(a+b\tan(c+dx))} +$$

$$\frac{a(a^2+b^2)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2} \frac{b(Ab-aB)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 4134

$$\frac{A(a^2+b^2)^2 \int \cot(c+dx) dx}{a} - \frac{b(-3a^5B+6a^4Ab+a^3b^2B+3a^2Ab^3+Ab^5) \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^2x(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{a^2+b^2}$$

$$\frac{a(a^2+b^2)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2} \frac{b(Ab-aB)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 3042

$$\frac{A(a^2+b^2)^2 \int -\tan(c+dx+\frac{\pi}{2}) dx}{a} - \frac{b(-3a^5B+6a^4Ab+a^3b^2B+3a^2Ab^3+Ab^5) \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^2x(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{a^2+b^2}$$

$$\frac{a(a^2+b^2)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2} \frac{b(Ab-aB)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 25

3.287. $\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

$$\frac{\frac{A(a^2+b^2)^2 \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx}{a} - \frac{b(-3a^5B+6a^4Ab+a^3b^2B+3a^2Ab^3+Ab^5) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^2x(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{a^2+b^2}}{a(a^2+b^2)} + \frac{b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)(a+b \tan(c+dx))} = \frac{b(Ab-aB)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} + \frac{b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \frac{A(a^2+b^2)^2 \log(-\sin(c+dx))}{ad} - \frac{a^2x(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{a^2+b^2}$$

↓ 3956

$$\frac{b(-3a^5B+6a^4Ab+a^3b^2B+3a^2Ab^3+Ab^5) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{A(a^2+b^2)^2 \log(-\sin(c+dx))}{ad} - \frac{a^2x(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{a^2+b^2} + \frac{b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)(a+b \tan(c+dx))} = \frac{b(Ab-aB)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} + \frac{b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \frac{A(a^2+b^2)^2 \log(-\sin(c+dx))}{ad} - \frac{a^2x(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{a^2+b^2}$$

↓ 4013

$$\frac{b(Ab-aB)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} + \frac{b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \frac{A(a^2+b^2)^2 \log(-\sin(c+dx))}{ad} - \frac{a^2x(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{a^2+b^2} - \frac{b(-3a^5B+6a^4Ab+a^3b^2B+3a^2Ab^3+Ab^5) \log(a \cos(c+dx))}{ad(a^2+b^2)}$$

```
input Int[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]
```

```
output (b*(A*b - a*B))/(2*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + (((-(a^2*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B)*x)/(a^2 + b^2)) + (A*(a^2 + b^2)^2*Log[-Sin[c + d*x]]/(a*d) - (b*(6*a^4*A*b + 3*a^2*A*b^3 + A*b^5 - 3*a^5*B + a^3*b^2*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]]/(a*(a^2 + b^2)*d))/(a*(a^2 + b^2)) + (b*(3*a^2*A*b + A*b^3 - 2*a^3*B))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])))/(a*(a^2 + b^2))
```

3.287.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

3.287. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4134 Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^
2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/
((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)
*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Sim
p[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*
x])/(c + d*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x]
&& NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

3.287.4 Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.13

method	result
derivativedivides	$\frac{(-A a^3 + 3A a b^2 - 3B a^2 b + B b^3) \ln(1 + \tan^2(dx+c))}{2(a^2+b^2)^3} + \frac{(-3A a^2 b + A b^3 + B a^3 - 3B a b^2) \arctan(\tan(dx+c))}{(a^2+b^2)^3} + \frac{A \ln(\tan(dx+c))}{a^3} + \frac{b}{(a^2+b^2)d}$
default	$\frac{(-A a^3 + 3A a b^2 - 3B a^2 b + B b^3) \ln(1 + \tan^2(dx+c))}{2(a^2+b^2)^3} + \frac{(-3A a^2 b + A b^3 + B a^3 - 3B a b^2) \arctan(\tan(dx+c))}{(a^2+b^2)^3} + \frac{A \ln(\tan(dx+c))}{a^3} + \frac{b}{(a^2+b^2)d}$
parallelrisch	$-12b(a+b \tan(dx+c))^2 (A a^4 b + \frac{1}{2} A a^2 b^3 + \frac{1}{6} A b^5 - \frac{1}{2} B a^5 + \frac{1}{6} B a^3 b^2) \ln(a+b \tan(dx+c)) - a^3 (a+b \tan(dx+c))^2 (A a^3 - 3A a b^2)$
norman	$\frac{(3A a^2 b - A b^3 - B a^3 + 3B a b^2) a^2 x}{(a^4 + 2a^2 b^2 + b^4)(a^2 + b^2)} - \frac{b^2 (3A a^2 b - A b^3 - B a^3 + 3B a b^2) x (\tan^2(dx+c))}{(a^4 + 2a^2 b^2 + b^4)(a^2 + b^2)} - \frac{b (4A a^2 b^2 + 2A b^4 - 3B a^3 b - B a b^3) \tan(dx+c)}{d a^2 (a^4 + 2a^2 b^2 + b^4)} - \frac{b^2}{(a+b \tan(dx+c))^2}$
risch	$-\frac{x B}{3i a^2 b - i b^3 - a^3 + 3a b^2} - \frac{6i a^2 b B x}{a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6} + \frac{2i B b^3 c}{d(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)} + \frac{2i b^6 A x}{a^3(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)} - \frac{b}{d(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)}$

```
input int(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x,method=_RETURNVERBOSE
)
```

```
output 1/d*(1/(a^2+b^2)^3*(1/2*(-A*a^3+3*A*a*b^2-3*B*a^2*b+B*b^3)*ln(1+tan(d*x+c)
^2)+(-3*A*a^2*b+A*b^3+B*a^3-3*B*a*b^2)*arctan(tan(d*x+c)))+1/a^3*A*ln(tan(
d*x+c))+b*(3*A*a^2*b+A*b^3-2*B*a^3)/(a^2+b^2)^2/a^2/(a+b*tan(d*x+c))-b*(6*
A*a^4*b+3*A*a^2*b^3+A*b^5-3*B*a^5+B*a^3*b^2)/(a^2+b^2)^3/a^3*ln(a+b*tan(d*
x+c))+1/2*(A*b-B*a)*b/(a^2+b^2)/a/(a+b*tan(d*x+c))^2)
```

$$3.287. \int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

3.287.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 683 vs. $2(213) = 426$.

Time = 0.33 (sec) , antiderivative size = 683, normalized size of antiderivative = 3.18

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx = \frac{7Ba^5b^3 - 9Aa^4b^4 + Ba^3b^5 - 3Aa^2b^6 - 2(Ba^8 - 3Aa^7b - 3Ba^6b^2 + Aa^5b^3)dx - (5Ba^5b^3 - 7Aa^4b^4 -$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="fricas")`

output `-1/2*(7*B*a^5*b^3 - 9*A*a^4*b^4 + B*a^3*b^5 - 3*A*a^2*b^6 - 2*(B*a^8 - 3*A*a^7*b - 3*B*a^6*b^2 + A*a^5*b^3)*d*x - (5*B*a^5*b^3 - 7*A*a^4*b^4 - B*a^3*b^5 - A*a^2*b^6 + 2*(B*a^6*b^2 - 3*A*a^5*b^3 - 3*B*a^4*b^4 + A*a^3*b^5)*d*x)*tan(d*x + c)^2 - (A*a^8 + 3*A*a^6*b^2 + 3*A*a^4*b^4 + A*a^2*b^6 + (A*a^6*b^2 + 3*A*a^4*b^4 + 3*A*a^2*b^6 + A*b^8))*tan(d*x + c)^2 + 2*(A*a^7*b + 3*A*a^5*b^3 + 3*A*a^3*b^5 + A*a*b^7)*tan(d*x + c))*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1)) - (3*B*a^7*b - 6*A*a^6*b^2 - B*a^5*b^3 - 3*A*a^4*b^4 - A*a^2*b^6 + (3*B*a^5*b^3 - 6*A*a^4*b^4 - B*a^3*b^5 - 3*A*a^2*b^6 - A*b^8))*tan(d*x + c)^2 + 2*(3*B*a^6*b^2 - 6*A*a^5*b^3 - B*a^4*b^4 - 3*A*a^3*b^5 - A*a*b^7)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - 2*(3*B*a^6*b^2 - 4*A*a^5*b^3 - 3*B*a^4*b^4 + 3*A*a^3*b^5 + A*a*b^7 + 2*(B*a^7*b - 3*A*a^6*b^2 - 3*B*a^5*b^3 + A*a^4*b^4)*d*x)*tan(d*x + c))/((a^9*b^2 + 3*a^7*b^4 + 3*a^5*b^6 + a^3*b^8)*d*tan(d*x + c)^2 + 2*(a^10*b + 3*a^8*b^3 + 3*a^6*b^5 + a^4*b^7)*d*tan(d*x + c) + (a^11 + 3*a^9*b^2 + 3*a^7*b^4 + a^5*b^6)*d)`

3.287.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx = \text{Exception raised: AttributeError}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)`

output `Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'`

3.287. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

3.287.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 372, normalized size of antiderivative = 1.73

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{2(Ba^3-3Aa^2b-3Bab^2+Ab^3)(dx+c)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{2(3Ba^5b-6Aa^4b^2-Ba^3b^3-3Aa^2b^4-Ab^6)\log(b\tan(dx+c)+a)}{a^9+3a^7b^2+3a^5b^4+a^3b^6} - \frac{(Aa^3+3Ba^2b-3Aab^2-Bb^3)\log(\tan(dx+c)^2+1)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{(5Ba^4b-7Aa^3b^2+Ba^2b^3-3Aa^2b^4+2(2Ba^3b^2-3Aa^2b^3-Ab^5)\tan(dx+c))/(a^8+2a^6b^2+a^4b^4+(a^6b^2+2a^4b^4+a^2b^6)\tan(dx+c)^2+2(a^7b+a^5b^3+a^3b^5)\tan(dx+c))+2A\log(\tan(dx+c))/a^3}{d}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="maxima")`

output `1/2*(2*(B*a^3 - 3*A*a^2*b - 3*B*a*b^2 + A*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + 2*(3*B*a^5*b - 6*A*a^4*b^2 - B*a^3*b^3 - 3*A*a^2*b^4 - A*b^6)*log(b*tan(d*x + c) + a)/(a^9 + 3*a^7*b^2 + 3*a^5*b^4 + a^3*b^6) - (A*a^3 + 3*B*a^2*b - 3*A*a*b^2 - B*b^3)*log(tan(d*x + c)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - (5*B*a^4*b - 7*A*a^3*b^2 + B*a^2*b^3 - 3*A*a*b^4 + 2*(2*B*a^3*b^2 - 3*A*a^2*b^3 - A*b^5)*tan(d*x + c))/(a^8 + 2*a^6*b^2 + a^4*b^4 + (a^6*b^2 + 2*a^4*b^4 + a^2*b^6)*tan(d*x + c)^2 + 2*(a^7*b + 2*a^5*b^3 + a^3*b^5)*tan(d*x + c)) + 2*A*log(tan(d*x + c))/a^3)/d`

3.287.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 479 vs. 2(213) = 426.

Time = 1.00 (sec) , antiderivative size = 479, normalized size of antiderivative = 2.23

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{2(Ba^3-3Aa^2b-3Bab^2+Ab^3)(dx+c)}{a^6+3a^4b^2+3a^2b^4+b^6} - \frac{(Aa^3+3Ba^2b-3Aab^2-Bb^3)\log(\tan(dx+c)^2+1)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{2(3Ba^5b^2-6Aa^4b^3-Ba^3b^4-3Aa^2b^5-Ab^7)\log(\tan(dx+c)^2+1)}{a^9b+3a^7b^3+3a^5b^5+a^3b^7}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="giac")`

output $\frac{1}{2} \cdot (2 \cdot (B \cdot a^3 - 3 \cdot A \cdot a^2 \cdot b - 3 \cdot B \cdot a \cdot b^2 + A \cdot b^3) \cdot (d \cdot x + c) / (a^6 + 3 \cdot a^4 \cdot b^2 + 3 \cdot a^2 \cdot b^4 + b^6) - (A \cdot a^3 + 3 \cdot B \cdot a^2 \cdot b - 3 \cdot A \cdot a \cdot b^2 - B \cdot b^3) \cdot \log(\tan(d \cdot x + c)^2 + 1) / (a^6 + 3 \cdot a^4 \cdot b^2 + 3 \cdot a^2 \cdot b^4 + b^6) + 2 \cdot (3 \cdot B \cdot a^5 \cdot b^2 - 6 \cdot A \cdot a^4 \cdot b^3 - B \cdot a^3 \cdot b^4 - 3 \cdot A \cdot a^2 \cdot b^5 - A \cdot b^7) \cdot \log(\text{abs}(b \cdot \tan(d \cdot x + c) + a)) / (a^9 \cdot b + 3 \cdot a^7 \cdot b^3 + 3 \cdot a^5 \cdot b^5 + a^3 \cdot b^7) + 2 \cdot A \cdot \log(\text{abs}(\tan(d \cdot x + c))) / a^3 - (9 \cdot B \cdot a^5 \cdot b^3 \cdot \tan(d \cdot x + c)^2 - 18 \cdot A \cdot a^4 \cdot b^4 \cdot \tan(d \cdot x + c)^2 - 3 \cdot B \cdot a^3 \cdot b^5 \cdot \tan(d \cdot x + c)^2 - 9 \cdot A \cdot a^2 \cdot b^6 \cdot \tan(d \cdot x + c)^2 - 3 \cdot A \cdot b^8 \cdot \tan(d \cdot x + c)^2 + 22 \cdot B \cdot a^6 \cdot b^2 \cdot \tan(d \cdot x + c) - 42 \cdot A \cdot a^5 \cdot b^3 \cdot \tan(d \cdot x + c) - 2 \cdot B \cdot a^4 \cdot b^4 \cdot \tan(d \cdot x + c) - 26 \cdot A \cdot a^3 \cdot b^5 \cdot \tan(d \cdot x + c) - 8 \cdot A \cdot a \cdot b^7 \cdot \tan(d \cdot x + c) + 14 \cdot B \cdot a^7 \cdot b - 25 \cdot A \cdot a^6 \cdot b^2 + 3 \cdot B \cdot a^5 \cdot b^3 - 19 \cdot A \cdot a^4 \cdot b^4 + B \cdot a^3 \cdot b^5 - 6 \cdot A \cdot a^2 \cdot b^6) / ((a^9 + 3 \cdot a^7 \cdot b^2 + 3 \cdot a^5 \cdot b^4 + a^3 \cdot b^6) \cdot (b \cdot \tan(d \cdot x + c) + a)^2) / d$

3.287.9 Mupad [B] (verification not implemented)

Time = 10.32 (sec) , antiderivative size = 315, normalized size of antiderivative = 1.47

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{-5 B a^3 b + 7 A a^2 b^2 - B a b^3 + 3 A b^4}{2 a (a^4 + 2 a^2 b^2 + b^4)} + \frac{\tan(c+dx) (-2 B a^3 b^2 + 3 A a^2 b^3 + A b^5)}{a^2 (a^4 + 2 a^2 b^2 + b^4)} + \frac{A \ln(\tan(c+dx))}{a^3 d}$$

$$+ \frac{\ln(\tan(c+dx) - i) (-B + A i)}{2 d (-a^3 i + 3 a^2 b + a b^2 3i - b^3)} + \frac{\ln(\tan(c+dx) + i) (A - B i)}{2 d (-a^3 + a^2 b 3i + 3 a b^2 - b^3 i)}$$

$$- \frac{b \ln(a + b \tan(c+dx)) (-3 B a^5 + 6 A a^4 b + B a^3 b^2 + 3 A a^2 b^3 + A b^5)}{a^3 d (a^2 + b^2)^3}$$

input `int((cot(c + d*x)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3,x)`

output $((3 \cdot A \cdot b^4 + 7 \cdot A \cdot a^2 \cdot b^2 - B \cdot a \cdot b^3 - 5 \cdot B \cdot a^3 \cdot b) / (2 \cdot a \cdot (a^4 + b^4 + 2 \cdot a^2 \cdot b^2)) + (\tan(c + d \cdot x) \cdot (A \cdot b^5 + 3 \cdot A \cdot a^2 \cdot b^3 - 2 \cdot B \cdot a^3 \cdot b^2)) / (a^2 \cdot (a^4 + b^4 + 2 \cdot a^2 \cdot b^2))) / (d \cdot (a^2 + b^2 \cdot \tan(c + d \cdot x)^2 + 2 \cdot a \cdot b \cdot \tan(c + d \cdot x))) + (A \cdot \log(\tan(c + d \cdot x))) / (a^3 \cdot d) + (\log(\tan(c + d \cdot x) - i) \cdot (A \cdot i - B)) / (2 \cdot d \cdot (a \cdot b^2 \cdot 3i + 3 \cdot a^2 \cdot b - a^3 \cdot i - b^3)) + (\log(\tan(c + d \cdot x) + i) \cdot (A - B \cdot i)) / (2 \cdot d \cdot (3 \cdot a \cdot b^2 + a^2 \cdot b \cdot 3i - a^3 - b^3 \cdot i)) - (b \cdot \log(a + b \cdot \tan(c + d \cdot x)) \cdot (A \cdot b^5 - 3 \cdot B \cdot a^5 + 3 \cdot A \cdot a^2 \cdot b^3 + B \cdot a^3 \cdot b^2 + 6 \cdot A \cdot a^4 \cdot b)) / (a^3 \cdot d \cdot (a^2 + b^2)^3)$

3.288 $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

3.288.1 Optimal result 2764
 3.288.2 Mathematica [C] (verified) 2765
 3.288.3 Rubi [A] (verified) 2765
 3.288.4 Maple [A] (verified) 2770
 3.288.5 Fricas [B] (verification not implemented) 2770
 3.288.6 Sympy [F(-2)] 2771
 3.288.7 Maxima [A] (verification not implemented) 2772
 3.288.8 Giac [A] (verification not implemented) 2772
 3.288.9 Mupad [B] (verification not implemented) 2773

3.288.1 Optimal result

Integrand size = 31, antiderivative size = 287

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= -\frac{(a^3 A - 3aAb^2 + 3a^2bB - b^3 B)x}{(a^2 + b^2)^3} - \frac{(3Ab - aB) \log(\sin(c+dx))}{a^4 d}$$

$$+ \frac{b^2(10a^4 Ab + 9a^2 Ab^3 + 3Ab^5 - 6a^5 B - 3a^3 b^2 B - ab^4 B) \log(a \cos(c+dx) + b \sin(c+dx))}{a^4 (a^2 + b^2)^3 d}$$

$$- \frac{b(2a^2 A + 3Ab^2 - abB)}{2a^2 (a^2 + b^2) d(a + b \tan(c+dx))^2} - \frac{A \cot(c+dx)}{ad(a + b \tan(c+dx))^2}$$

$$- \frac{b(a^4 A + 6a^2 Ab^2 + 3Ab^4 - 3a^3 bB - ab^3 B)}{a^3 (a^2 + b^2)^2 d(a + b \tan(c+dx))}$$

output

```
-(A*a^3-3*A*a*b^2+3*B*a^2*b-B*b^3)*x/(a^2+b^2)^3-(3*A*b-B*a)*ln(sin(dx+c))
/a^4/d+b^2*(10*A*a^4*b+9*A*a^2*b^3+3*A*b^5-6*B*a^5-3*B*a^3*b^2-B*a*b^4)*l
n(a*cos(dx+c)+b*sin(dx+c))/a^4/(a^2+b^2)^3/d-1/2*b*(2*A*a^2+3*A*b^2-B*a*
b)/a^2/(a^2+b^2)/d/(a+b*tan(dx+c))^2-A*cot(dx+c)/a/d/(a+b*tan(dx+c))^2-
b*(A*a^4+6*A*a^2*b^2+3*A*b^4-3*B*a^3*b-B*a*b^3)/a^3/(a^2+b^2)^2/d/(a+b*tan
(dx+c))
```

3.288.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.46 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.00

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= -\frac{A \cot(c+dx)}{a^3 d} + \frac{(A+iB) \log(i-\tan(c+dx))}{2(ia-b)^3 d}$$

$$- \frac{(3Ab-aB) \log(\tan(c+dx))}{a^4 d} - \frac{(iA+B) \log(i+\tan(c+dx))}{2(a-ib)^3 d}$$

$$+ \frac{b^2(10a^4 Ab + 9a^2 Ab^3 + 3Ab^5 - 6a^5 B - 3a^3 b^2 B - ab^4 B) \log(a+b \tan(c+dx))}{a^4 (a^2+b^2)^3 d}$$

$$- \frac{b^2(Ab-aB)}{2a^2 (a^2+b^2) d(a+b \tan(c+dx))^2} - \frac{b^2(4a^2 Ab + 2Ab^3 - 3a^3 B - ab^2 B)}{a^3 (a^2+b^2)^2 d(a+b \tan(c+dx))}$$

input `Integrate[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

output `-((A*Cot[c + d*x])/(a^3*d)) + ((A + I*B)*Log[I - Tan[c + d*x]])/(2*(I*a - b)^3*d) - ((3*A*b - a*B)*Log[Tan[c + d*x]])/(a^4*d) - ((I*A + B)*Log[I + Tan[c + d*x]])/(2*(a - I*b)^3*d) + (b^2*(10*a^4*A*b + 9*a^2*A*b^3 + 3*A*b^5 - 6*a^5*B - 3*a^3*b^2*B - a*b^4*B)*Log[a + b*Tan[c + d*x]])/(a^4*(a^2 + b^2)^3*d) - (b^2*(A*b - a*B))/(2*a^2*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) - (b^2*(4*a^2*A*b + 2*A*b^3 - 3*a^3*B - a*b^2*B))/(a^3*(a^2 + b^2)^2*d*(a + b*Tan[c + d*x]))`

3.288.3 Rubi [A] (verified)

Time = 1.78 (sec) , antiderivative size = 333, normalized size of antiderivative = 1.16, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 4092, 3042, 4132, 27, 3042, 4132, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

↓ 3042

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^2 (a + b \tan(c + dx))^3} dx \\
 & \quad \downarrow \text{4092} \\
 & \frac{\int \frac{\cot(c+dx)(3Ab \tan^2(c+dx) + aA \tan(c+dx) + 3Ab - aB)}{(a+b \tan(c+dx))^3} dx}{a} - \frac{A \cot(c + dx)}{ad(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{3Ab \tan(c+dx)^2 + aA \tan(c+dx) + 3Ab - aB}{\tan(c+dx)(a+b \tan(c+dx))^3} dx}{a} - \frac{A \cot(c + dx)}{ad(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{4132} \\
 & \frac{\int \frac{2 \cot(c+dx) \left((aA+bB) \tan(c+dx)a^2 + b(2Aa^2 - bBa + 3Ab^2) \tan^2(c+dx) + (a^2+b^2)(3Ab - aB) \right)}{(a+b \tan(c+dx))^2} dx}{2a(a^2+b^2)} + \frac{b(2a^2A - abB + 3Ab^2)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \frac{A \cot(c + dx)}{ad(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\cot(c+dx) \left((aA+bB) \tan(c+dx)a^2 + b(2Aa^2 - bBa + 3Ab^2) \tan^2(c+dx) + (a^2+b^2)(3Ab - aB) \right)}{(a+b \tan(c+dx))^2} dx}{a(a^2+b^2)} + \frac{b(2a^2A - abB + 3Ab^2)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \frac{A \cot(c + dx)}{ad(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{(aA+bB) \tan(c+dx)a^2 + b(2Aa^2 - bBa + 3Ab^2) \tan(c+dx)^2 + (a^2+b^2)(3Ab - aB)}{\tan(c+dx)(a+b \tan(c+dx))^2} dx}{a(a^2+b^2)} + \frac{b(2a^2A - abB + 3Ab^2)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \frac{A \cot(c + dx)}{ad(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{4132} \\
 & \frac{\int \frac{\cot(c+dx) \left((Aa^2 + 2bBa - Ab^2) \tan(c+dx)a^3 + b(Aa^4 - 3bBa^3 + 6Ab^2a^2 - b^3Ba + 3Ab^4) \tan^2(c+dx) + (a^2+b^2)^2(3Ab - aB) \right)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{b(Aa^4 - 3a^3bB + 6a^2Ab^2 - ab^3)}{ad(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \frac{A \cot(c + dx)}{ad(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.288. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{\int \frac{(Aa^2+2bBa-Ab^2) \tan(c+dx)a^3+b(Aa^4-3bBa^3+6Ab^2a^2-b^3Ba+3Ab^4) \tan(c+dx)^2+(a^2+b^2)^2(3Ab-aB)}{\tan(c+dx)(a+b \tan(c+dx))} dx + \frac{b(a^4A-3a^3bB+6a^2Ab^2-ab^3B+3Ab^4)}{ad(a^2+b^2)(a+b \tan(c+dx))}}{a(a^2+b^2)}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^2} \quad a$$

↓ 4134

$$\frac{\frac{(a^2+b^2)^2(3Ab-aB) \int \cot(c+dx) dx}{a} - \frac{b^2(-6a^5B+10a^4Ab-3a^3b^2B+9a^2Ab^3-ab^4B+3Ab^5) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^3x(a^3A+3a^2bB-3aAb^2-b^3B)}{a^2+b^2}}{a(a^2+b^2)} + \frac{b(a^4A-3a^3bB+6a^2Ab^2-ab^3B+3Ab^4)}{aa}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^2} \quad a$$

↓ 3042

$$\frac{\frac{(a^2+b^2)^2(3Ab-aB) \int -\tan(c+dx+\frac{\pi}{2}) dx}{a} - \frac{b^2(-6a^5B+10a^4Ab-3a^3b^2B+9a^2Ab^3-ab^4B+3Ab^5) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^3x(a^3A+3a^2bB-3aAb^2-b^3B)}{a^2+b^2}}{a(a^2+b^2)} + \frac{b(a^4A-3a^3bB+6a^2Ab^2-ab^3B+3Ab^4)}{aa}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^2} \quad a$$

↓ 25

$$\frac{\frac{(a^2+b^2)^2(3Ab-aB) \int \tan(\frac{1}{2}(2c+\pi)+dx) dx}{a} - \frac{b^2(-6a^5B+10a^4Ab-3a^3b^2B+9a^2Ab^3-ab^4B+3Ab^5) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^3x(a^3A+3a^2bB-3aAb^2-b^3B)}{a^2+b^2}}{a(a^2+b^2)} + \frac{b(a^4A-3a^3bB+6a^2Ab^2-ab^3B+3Ab^4)}{aa}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^2} \quad a$$

↓ 3956

$$\frac{\frac{b^2(-6a^5B+10a^4Ab-3a^3b^2B+9a^2Ab^3-ab^4B+3Ab^5) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{(a^2+b^2)^2(3Ab-aB) \log(-\sin(c+dx))}{ad} + \frac{a^3x(a^3A+3a^2bB-3aAb^2-b^3B)}{a^2+b^2}}{a(a^2+b^2)} + \frac{b(a^4A-3a^3bB+6a^2Ab^2-ab^3B+3Ab^4)}{aa}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^2} \quad a$$

↓ 4013

3.288. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{b(2a^2A-abB+3Ab^2)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2} + \frac{b(a^4A-3a^3bB+6a^2Ab^2-ab^3B+3Ab^4)}{ad(a^2+b^2)(a+b\tan(c+dx))} + \frac{(a^2+b^2)^2(3Ab-aB)\log(-\sin(c+dx))}{ad} + \frac{a^3x(a^3A+3a^2bB-3aAb^2-b^3B)}{a^2+b^2} - \frac{b^2}{a(a^2+b^2)}$$

$$\frac{A \cot(c+dx)}{ad(a+b\tan(c+dx))^2}$$

input `Int[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

output `-((A*Cot[c + d*x])/(a*d*(a + b*Tan[c + d*x])^2)) - ((b*(2*a^2*A + 3*A*b^2 - a*b*B))/(2*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + (((a^3*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B)*x)/(a^2 + b^2) + ((a^2 + b^2)^2*(3*A*b - a*B)*Log[-Sin[c + d*x]])/(a*d) - (b^2*(10*a^4*A*b + 9*a^2*A*b^3 + 3*A*b^5 - 6*a^5*B - 3*a^3*b^2*B - a*b^4*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))/(a*(a^2 + b^2)) + (b*(a^4*A + 6*a^2*A*b^2 + 3*A*b^4 - 3*a^3*b*B - a*b^3*B))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(a*(a^2 + b^2))/a`

3.288.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4134 `Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/(a^2 + b^2)*(c^2 + d^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.288.4 Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.01

method	result
derivativedivides	$\frac{(3Aa^2b - Ab^3 - Ba^3 + 3Bab^2) \ln(1 + \tan^2(dx+c)) + (-Aa^3 + 3Aab^2 - 3Ba^2b + Bb^3) \arctan(\tan(dx+c))}{(a^2+b^2)^3} - \frac{A}{a^3 \tan(dx+c)} + \frac{(-3Ab+B^2)}{a^3 \tan(dx+c)}$
default	$\frac{(3Aa^2b - Ab^3 - Ba^3 + 3Bab^2) \ln(1 + \tan^2(dx+c)) + (-Aa^3 + 3Aab^2 - 3Ba^2b + Bb^3) \arctan(\tan(dx+c))}{(a^2+b^2)^3} - \frac{A}{a^3 \tan(dx+c)} + \frac{(-3Ab+B^2)}{a^3 \tan(dx+c)}$
parallelrisch	$20(Aa^4b + \frac{9}{10}Aa^2b^3 + \frac{3}{10}Ab^5 - \frac{3}{5}Ba^5 - \frac{3}{10}Ba^3b^2 - \frac{1}{10}Bab^4)b^2(a+b \tan(dx+c))^2 \ln(a+b \tan(dx+c)) + 3(Aa^2b - \frac{1}{3}Ab^3 - \frac{1}{3}Ba^3)$
norman	$\frac{b(3Aa^4b + 11Aa^2b^3 + 6Ab^5 - 4Ba^3b^2 - 2Bab^4)(\tan^2(dx+c))}{da^3(a^4+2a^2b^2+b^4)} - \frac{A}{ad} - \frac{b^2(Aa^3 - 3Aab^2 + 3Ba^2b - Bb^3)x(\tan^3(dx+c))}{(a^4+2a^2b^2+b^4)(a^2+b^2)} + \frac{b^2(4Aa^4b + 4Aa^2b^3 + 4Ab^5 - 4Ba^4 - 4Ba^2b^2 - 4Bab^4)}{(a^4+2a^2b^2+b^4)(a^2+b^2)}$
risch	Expression too large to display

input `int(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} \left(\frac{1}{(a^2+b^2)^3} \left(\frac{1}{2} (3Aa^2b - Ab^3 - Ba^3 + 3Bab^2) \ln(1 + \tan^2(dx+c)) + (-Aa^3 + 3Aab^2 - 3Ba^2b + Bb^3) \arctan(\tan(dx+c)) \right) - \frac{1}{a^3} \frac{A}{\tan(dx+c)} + \frac{(-3Ab+B^2)}{a^3 \tan(dx+c)} \right) - \frac{b^2(4Aa^4b + 4Aa^2b^3 + 4Ab^5 - 4Ba^4 - 4Ba^2b^2 - 4Bab^4)}{(a^4+2a^2b^2+b^4)(a^2+b^2)}$$

3.288.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 917 vs. 2(283) = 566.

Time = 0.37 (sec) , antiderivative size = 917, normalized size of antiderivative = 3.20

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx = \frac{2Aa^9 + 6Aa^7b^2 + 6Aa^5b^4 + 2Aa^3b^6 + (7Ba^5b^4 - 9Aa^4b^5 + Ba^3b^6 - 3Aa^2b^7 + 2(Aa^7b^2 + 3Ba^6b^3 - \dots)}{\dots}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="fricas")`

output `-1/2*(2*A*a^9 + 6*A*a^7*b^2 + 6*A*a^5*b^4 + 2*A*a^3*b^6 + (7*B*a^5*b^4 - 9*A*a^4*b^5 + B*a^3*b^6 - 3*A*a^2*b^7 + 2*(A*a^7*b^2 + 3*B*a^6*b^3 - 3*A*a^5*b^4 - B*a^4*b^5)*d*x)*tan(d*x + c)^3 + 2*(A*a^7*b^2 + 4*B*a^6*b^3 - 2*A*a^5*b^4 - 3*B*a^4*b^5 + 6*A*a^3*b^6 - B*a^2*b^7 + 3*A*a*b^8 + 2*(A*a^8*b + 3*B*a^7*b^2 - 3*A*a^6*b^3 - B*a^5*b^4)*d*x)*tan(d*x + c)^2 - ((B*a^7*b^2 - 3*A*a^6*b^3 + 3*B*a^5*b^4 - 9*A*a^4*b^5 + 3*B*a^3*b^6 - 9*A*a^2*b^7 + B*a*b^8 - 3*A*b^9)*tan(d*x + c)^3 + 2*(B*a^8*b - 3*A*a^7*b^2 + 3*B*a^6*b^3 - 9*A*a^5*b^4 + 3*B*a^4*b^5 - 9*A*a^3*b^6 + B*a^2*b^7 - 3*A*a*b^8)*tan(d*x + c)^2 + (B*a^9 - 3*A*a^8*b + 3*B*a^7*b^2 - 9*A*a^6*b^3 + 3*B*a^5*b^4 - 9*A*a^4*b^5 + B*a^3*b^6 - 3*A*a^2*b^7)*tan(d*x + c))*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1)) + ((6*B*a^5*b^4 - 10*A*a^4*b^5 + 3*B*a^3*b^6 - 9*A*a^2*b^7 + B*a*b^8 - 3*A*b^9)*tan(d*x + c)^3 + 2*(6*B*a^6*b^3 - 10*A*a^5*b^4 + 3*B*a^4*b^5 - 9*A*a^3*b^6 + B*a^2*b^7 - 3*A*a*b^8)*tan(d*x + c)^2 + (6*B*a^7*b^2 - 10*A*a^6*b^3 + 3*B*a^5*b^4 - 9*A*a^4*b^5 + B*a^3*b^6 - 3*A*a^2*b^7)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) + (4*A*a^8*b + 12*A*a^6*b^3 - 9*B*a^5*b^4 + 23*A*a^4*b^5 - 3*B*a^3*b^6 + 9*A*a^2*b^7 + 2*(A*a^9 + 3*B*a^8*b - 3*A*a^7*b^2 - B*a^6*b^3)*d*x)*tan(d*x + c))/((a^10*b^2 + 3*a^8*b^4 + 3*a^6*b^6 + a^4*b^8)*d*tan(d*x + c)^3 + 2*(a^11*b + 3*a^9*b^3 + 3*a^7*b^5 + a^5*b^7)*d*tan(d*x + c)^2 + (a^12 + 3*a^10*b^2 + 3*a^8*b^4 + a^6*b^6)*d*tan(d*x + c))`

3.288.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx = \text{Exception raised: AttributeError}$$

input `integrate(cot(d*x+c)**2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)`

output `Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'`

3.288.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 454, normalized size of antiderivative = 1.58

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx =$$

$$\frac{2(Aa^3+3Ba^2b-3Aab^2-Bb^3)(dx+c)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{2(6Ba^5b^2-10Aa^4b^3+3Ba^3b^4-9Aa^2b^5+Bab^6-3Ab^7) \log(b \tan(dx+c)+a)}{a^{10}+3a^8b^2+3a^6b^4+a^4b^6} + \frac{(Ba^3-3Aa^2b-3Aab^2-Bb^3) \log(\tan(dx+c)^2+1)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{(2Aa^6+4Aa^4b^2+2Aa^2b^4+2Aa^4b^2-3Ba^3b^3+6Aa^2b^4-Ba^2b^5+3Ab^6) \tan(dx+c)^2 + (4Aa^5b-7Ba^4b^2+17Aa^3b^3-3Ba^2b^4+9Aa^2b^5) \tan(dx+c)}{(a^7b^2+2a^5b^4+a^3b^6) \tan(dx+c)^3 + 2(a^8b+2a^6b^3+a^4b^5) \tan(dx+c)^2 + (a^9+2a^7b^2+a^5b^4) \tan(dx+c)} - \frac{2(Ba-3Aa^2b) \log(\tan(dx+c))}{a^4} / d$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="maxima")`

output `-1/2*(2*(A*a^3 + 3*B*a^2*b - 3*A*a*b^2 - B*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + 2*(6*B*a^5*b^2 - 10*A*a^4*b^3 + 3*B*a^3*b^4 - 9*A*a^2*b^5 + B*a*b^6 - 3*A*b^7)*log(b*tan(d*x + c) + a)/(a^10 + 3*a^8*b^2 + 3*a^6*b^4 + a^4*b^6) + (B*a^3 - 3*A*a^2*b - 3*B*a*b^2 + A*b^3)*log(tan(d*x + c)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + (2*A*a^6 + 4*A*a^4*b^2 + 2*A*a^2*b^4 + 2*(A*a^4*b^2 - 3*B*a^3*b^3 + 6*A*a^2*b^4 - B*a^2*b^5 + 3*A*b^6)*tan(d*x + c)^2 + (4*A*a^5*b - 7*B*a^4*b^2 + 17*A*a^3*b^3 - 3*B*a^2*b^4 + 9*A*a^2*b^5)*tan(d*x + c))/((a^7*b^2 + 2*a^5*b^4 + a^3*b^6)*tan(d*x + c)^3 + 2*(a^8*b + 2*a^6*b^3 + a^4*b^5)*tan(d*x + c)^2 + (a^9 + 2*a^7*b^2 + a^5*b^4)*tan(d*x + c)) - 2*(B*a - 3*A*b)*log(tan(d*x + c))/a^4)/d`

3.288.8 Giac [A] (verification not implemented)

Time = 0.92 (sec) , antiderivative size = 560, normalized size of antiderivative = 1.95

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx =$$

$$\frac{2(Aa^3+3Ba^2b-3Aab^2-Bb^3)(dx+c)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{(Ba^3-3Aa^2b-3Aab^2+Bb^3) \log(\tan(dx+c)^2+1)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{2(6Ba^5b^3-10Aa^4b^4+3Ba^3b^5-9Aa^2b^6+3Aab^7-Bb^8) \tan(dx+c)}{a^{10}b+3a^8b^3+3a^6b^5+a^4b^7+b^9} + \frac{(2Aa^6+4Aa^4b^2+2Aa^2b^4+2Aa^4b^2-3Ba^3b^3+6Aa^2b^4-Ba^2b^5+3Ab^6) \tan(dx+c)^2 + (4Aa^5b-7Ba^4b^2+17Aa^3b^3-3Ba^2b^4+9Aa^2b^5) \tan(dx+c)}{(a^7b^2+2a^5b^4+a^3b^6) \tan(dx+c)^3 + 2(a^8b+2a^6b^3+a^4b^5) \tan(dx+c)^2 + (a^9+2a^7b^2+a^5b^4) \tan(dx+c)} - \frac{2(Ba-3Aa^2b) \log(\tan(dx+c))}{a^4} / d$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="giac")`

output

$$\begin{aligned}
& -1/2*(2*(A*a^3 + 3*B*a^2*b - 3*A*a*b^2 - B*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2 \\
& + 3*a^2*b^4 + b^6) + (B*a^3 - 3*A*a^2*b - 3*B*a*b^2 + A*b^3)*\log(\tan(d*x \\
& + c)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + 2*(6*B*a^5*b^3 - 10*A*a^4 \\
& *b^4 + 3*B*a^3*b^5 - 9*A*a^2*b^6 + B*a*b^7 - 3*A*b^8)*\log(\text{abs}(b*\tan(d*x + \\
& c) + a))/(a^{10}*b + 3*a^8*b^3 + 3*a^6*b^5 + a^4*b^7) - (18*B*a^5*b^4*\tan(d \\
& *x + c)^2 - 30*A*a^4*b^5*\tan(d*x + c)^2 + 9*B*a^3*b^6*\tan(d*x + c)^2 - 27* \\
& A*a^2*b^7*\tan(d*x + c)^2 + 3*B*a*b^8*\tan(d*x + c)^2 - 9*A*b^9*\tan(d*x + c) \\
& ^2 + 42*B*a^6*b^3*\tan(d*x + c) - 68*A*a^5*b^4*\tan(d*x + c) + 26*B*a^4*b^5* \\
& \tan(d*x + c) - 66*A*a^3*b^6*\tan(d*x + c) + 8*B*a^2*b^7*\tan(d*x + c) - 22*A \\
& *a*b^8*\tan(d*x + c) + 25*B*a^7*b^2 - 39*A*a^6*b^3 + 19*B*a^5*b^4 - 41*A*a^4 \\
& *b^5 + 6*B*a^3*b^6 - 14*A*a^2*b^7)/((a^{10} + 3*a^8*b^2 + 3*a^6*b^4 + a^4*b \\
& ^6)*(b*\tan(d*x + c) + a)^2) - 2*(B*a - 3*A*b)*\log(\text{abs}(\tan(d*x + c)))/a^4 + \\
& 2*(B*a*\tan(d*x + c) - 3*A*b*\tan(d*x + c) + A*a)/(a^4*\tan(d*x + c))/d
\end{aligned}$$

3.288.9 Mupad [B] (verification not implemented)

Time = 13.09 (sec) , antiderivative size = 380, normalized size of antiderivative = 1.32

$$\begin{aligned}
& \int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx \\
& = \frac{b^2 \ln(a + b \tan(c + dx)) (-6 B a^5 + 10 A a^4 b - 3 B a^3 b^2 + 9 A a^2 b^3 - B a b^4 + 3 A b^5)}{a^4 d (a^2 + b^2)^3} \\
& \quad - \frac{\ln(\tan(c + dx) - i) (-B + A i)}{2 d (-a^3 - a^2 b 3i + 3 a b^2 + b^3 i)} \\
& \quad - \frac{\ln(\tan(c + dx)) (3 A b - B a)}{a^4 d} - \frac{\ln(\tan(c + dx) + i) (A - B i)}{2 d (-a^3 i - 3 a^2 b + a b^2 3i + b^3)} \\
& \quad - \frac{\frac{A}{a} + \frac{\tan(c+dx)^2 (A a^4 b^2 - 3 B a^3 b^3 + 6 A a^2 b^4 - B a b^5 + 3 A b^6)}{a^3 (a^4 + 2 a^2 b^2 + b^4)}}{d (a^2 \tan(c + dx) + 2 a b \tan(c + dx)^2 + b^2 \tan(c + dx)^3)} + \frac{\tan(c+dx) (4 A a^4 b - 7 B a^3 b^2 + 17 A a^2 b^3 - 3 B a b^4 + 9 A b^5)}{2 a^2 (a^4 + 2 a^2 b^2 + b^4)}
\end{aligned}$$

input `int((cot(c + d*x)^2*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3,x)`

output $(b^2 \log(a + b \tan(c + dx)) (3Ab^5 - 6B^2a^5 + 9A^2a^2b^3 - 3B^2a^3b^2 + 10A^2a^4b - B^2ab^4)) / (a^4 d (a^2 + b^2)^3) - (\log(\tan(c + dx)) - 1) (A^2i - B) / (2d (3a^2b^2 - a^2b^3i - a^3 + b^3i)) - (\log(\tan(c + dx)) (3Ab - B^2a)) / (a^4 d) - (\log(\tan(c + dx)) + 1) (A - B^2i) / (2d (a^2b^2 - 3a^2b - a^3i + b^3)) - (A/a + (\tan(c + dx))^2 (3Ab^6 + 6A^2a^2b^4 + A^2a^4b^2 - 3B^2a^3b^3 - B^2ab^5)) / (a^3 (a^4 + b^4 + 2a^2b^2)) + (\tan(c + dx) (9Ab^5 + 17A^2a^2b^3 - 7B^2a^3b^2 + 4A^2a^4b - 3B^2ab^4)) / (2a^2 (a^4 + b^4 + 2a^2b^2)) / (d (a^2 \tan(c + dx) + b^2 \tan(c + dx)^3 + 2ab \tan(c + dx)^2))$

3.288. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

3.289 $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

3.289.1 Optimal result 2775
 3.289.2 Mathematica [C] (verified) 2776
 3.289.3 Rubi [A] (verified) 2776
 3.289.4 Maple [A] (verified) 2782
 3.289.5 Fricas [B] (verification not implemented) 2782
 3.289.6 Sympy [F(-2)] 2783
 3.289.7 Maxima [A] (verification not implemented) 2784
 3.289.8 Giac [B] (verification not implemented) 2784
 3.289.9 Mupad [B] (verification not implemented) 2785

3.289.1 Optimal result

Integrand size = 31, antiderivative size = 352

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{(3a^2Ab - Ab^3 - a^3B + 3ab^2B)x}{(a^2 + b^2)^3} - \frac{(a^2A - 6Ab^2 + 3abB) \log(\sin(c+dx))}{a^5d}$$

$$- \frac{b^3(15a^4Ab + 17a^2Ab^3 + 6Ab^5 - 10a^5B - 9a^3b^2B - 3ab^4B) \log(a \cos(c+dx) + b \sin(c+dx))}{a^5(a^2 + b^2)^3d}$$

$$+ \frac{b(5a^2Ab + 6Ab^3 - 2a^3B - 3ab^2B)}{2a^3(a^2 + b^2)d(a+b \tan(c+dx))^2} + \frac{(2Ab - aB) \cot(c+dx)}{a^2d(a+b \tan(c+dx))^2}$$

$$- \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^2} + \frac{b(3a^4Ab + 11a^2Ab^3 + 6Ab^5 - a^5B - 6a^3b^2B - 3ab^4B)}{a^4(a^2 + b^2)^2d(a+b \tan(c+dx))}$$

output

```
(3*A*a^2*b-A*b^3-B*a^3+3*B*a*b^2)*x/(a^2+b^2)^3-(A*a^2-6*A*b^2+3*B*a*b)*ln
(sin(d*x+c))/a^5/d-b^3*(15*A*a^4*b+17*A*a^2*b^3+6*A*b^5-10*B*a^5-9*B*a^3*b
^2-3*B*a*b^4)*ln(a*cos(d*x+c)+b*sin(d*x+c))/a^5/(a^2+b^2)^3/d+1/2*b*(5*A*a
^2*b+6*A*b^3-2*B*a^3-3*B*a*b^2)/a^3/(a^2+b^2)/d/(a+b*tan(d*x+c))^2+(2*A*b-
B*a)*cot(d*x+c)/a^2/d/(a+b*tan(d*x+c))^2-1/2*A*cot(d*x+c)^2/a/d/(a+b*tan(d
*x+c))^2+b*(3*A*a^4*b+11*A*a^2*b^3+6*A*b^5-B*a^5-6*B*a^3*b^2-3*B*a*b^4)/a^
4/(a^2+b^2)^2/d/(a+b*tan(d*x+c))
```

3.289.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.69 (sec) , antiderivative size = 320, normalized size of antiderivative = 0.91

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{(3Ab - aB)\cot(c+dx)}{a^4d} - \frac{A\cot^2(c+dx)}{2a^3d} + \frac{(A+iB)\log(i-\tan(c+dx))}{2(a+ib)^3d}$$

$$- \frac{(a^2A - 6Ab^2 + 3abB)\log(\tan(c+dx))}{a^5d} + \frac{(A-iB)\log(i+\tan(c+dx))}{2(a-ib)^3d}$$

$$- \frac{b^3(15a^4Ab + 17a^2Ab^3 + 6Ab^5 - 10a^5B - 9a^3b^2B - 3ab^4B)\log(a+b\tan(c+dx))}{a^5(a^2+b^2)^3d}$$

$$+ \frac{b^3(Ab - aB)}{2a^3(a^2+b^2)d(a+b\tan(c+dx))^2} + \frac{b^3(5a^2Ab + 3Ab^3 - 4a^3B - 2ab^2B)}{a^4(a^2+b^2)^2d(a+b\tan(c+dx))}$$

input `Integrate[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

output `((3*A*b - a*B)*Cot[c + d*x])/(a^4*d) - (A*Cot[c + d*x]^2)/(2*a^3*d) + ((A + I*B)*Log[I - Tan[c + d*x]])/(2*(a + I*b)^3*d) - ((a^2*A - 6*A*b^2 + 3*a*b*B)*Log[Tan[c + d*x]])/(a^5*d) + ((A - I*B)*Log[I + Tan[c + d*x]])/(2*(a - I*b)^3*d) - (b^3*(15*a^4*A*b + 17*a^2*A*b^3 + 6*A*b^5 - 10*a^5*B - 9*a^3*b^2*B - 3*a*b^4*B)*Log[a + b*Tan[c + d*x]])/(a^5*(a^2 + b^2)^3*d) + (b^3*(A*b - a*B))/(2*a^3*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + (b^3*(5*a^2*A*b + 3*A*b^3 - 4*a^3*B - 2*a*b^2*B))/(a^4*(a^2 + b^2)^2*d*(a + b*Tan[c + d*x]))`

3.289.3 Rubi [A] (verified)

Time = 2.38 (sec) , antiderivative size = 406, normalized size of antiderivative = 1.15, number of steps used = 17, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.548$, Rules used = {3042, 4092, 27, 3042, 4132, 25, 3042, 4132, 27, 3042, 4132, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

3.289. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^3 (a + b \tan(c + dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & - \frac{\int \frac{2 \cot^2(c + dx)(2Ab \tan^2(c + dx) + aA \tan(c + dx) + 2Ab - aB)}{(a + b \tan(c + dx))^3} dx}{2a} - \frac{A \cot^2(c + dx)}{2ad(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{4092} \\
 & - \frac{\int \frac{\cot^2(c + dx)(2Ab \tan^2(c + dx) + aA \tan(c + dx) + 2Ab - aB)}{(a + b \tan(c + dx))^3} dx}{a} - \frac{A \cot^2(c + dx)}{2ad(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{27} \\
 & - \frac{\int \frac{2Ab \tan(c + dx)^2 + aA \tan(c + dx) + 2Ab - aB}{\tan(c + dx)^2 (a + b \tan(c + dx))^3} dx}{a} - \frac{A \cot^2(c + dx)}{2ad(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{\int \frac{\cot(c + dx)(Aa^2 + B \tan(c + dx)a^2 + 3bBa - 6Ab^2 - 3b(2Ab - aB) \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx}{a} - \frac{(2Ab - aB) \cot(c + dx)}{ad(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{4132} \\
 & - \frac{\int \frac{\cot(c + dx)(Aa^2 + B \tan(c + dx)a^2 + 3bBa - 6Ab^2 - 3b(2Ab - aB) \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx}{a} - \frac{(2Ab - aB) \cot(c + dx)}{ad(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{25} \\
 & - \frac{\int \frac{\cot(c + dx)(Aa^2 + B \tan(c + dx)a^2 + 3bBa - 6Ab^2 - 3b(2Ab - aB) \tan^2(c + dx))}{(a + b \tan(c + dx))^3} dx}{a} - \frac{(2Ab - aB) \cot(c + dx)}{ad(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{\int \frac{Aa^2 + B \tan(c + dx)a^2 + 3bBa - 6Ab^2 - 3b(2Ab - aB) \tan(c + dx)^2}{\tan(c + dx)(a + b \tan(c + dx))^3} dx}{a} - \frac{(2Ab - aB) \cot(c + dx)}{ad(a + b \tan(c + dx))^2} - \frac{A \cot^2(c + dx)}{2ad(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{4132} \\
 & - \frac{\int \frac{2 \cot(c + dx) \left(-((Ab - aB) \tan(c + dx)a^3) - b(-2Ba^3 + 5Aba^2 - 3b^2Ba + 6Ab^3) \tan^2(c + dx) + (a^2 + b^2)(Aa^2 + 3bBa - 6Ab^2) \right)}{(a + b \tan(c + dx))^2}{2a(a^2 + b^2)} dx}{a} - \frac{b(-2a^3B + 5a^2Ab - 3ab^2B + 6Aa^2b^2)}{2ad(a^2 + b^2)(a + b \tan(c + dx))} \\
 & \quad \downarrow \\
 & - \frac{A \cot^2(c + dx)}{2ad(a + b \tan(c + dx))^2}
 \end{aligned}$$

3.289. $\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$

↓ 27

$$\int \frac{\cot(c+dx) \left(-((Ab-aB) \tan(c+dx)a^3) - b(-2Ba^3+5Aba^2-3b^2Ba+6Ab^3) \tan^2(c+dx) + (a^2+b^2)(Aa^2+3bBa-6Ab^2) \right)}{a(a^2+b^2)^2} dx - \frac{b(-2a^3B+5a^2Ab-3ab^2B+6Ab^3)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^2}$$

↓ 3042

$$\int \frac{-((Ab-aB) \tan(c+dx)a^3) - b(-2Ba^3+5Aba^2-3b^2Ba+6Ab^3) \tan(c+dx)^2 + (a^2+b^2)(Aa^2+3bBa-6Ab^2)}{a(a^2+b^2)^2} dx - \frac{b(-2a^3B+5a^2Ab-3ab^2B+6Ab^3)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} - \frac{(2Ab)}{ad(a^2+b^2)}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^2}$$

↓ 4132

$$\int \frac{\cot(c+dx) \left(-((-Ba^2+2Aba+b^2B) \tan(c+dx)a^4) - b(-Ba^5+3Aba^4-6b^2Ba^3+11Ab^3a^2-3b^4Ba+6Ab^5) \tan^2(c+dx) + (a^2+b^2)^2(Aa^2+3bBa-6Ab^2) \right)}{a(a^2+b^2)^2} dx - \frac{b(a^5(-B)+3Aba^4-6b^2Ba^3+11Ab^3a^2-3b^4Ba+6Ab^5)}{2ad(a^2+b^2)^2(a+b \tan(c+dx))^2}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^2}$$

↓ 3042

$$\int \frac{-((-Ba^2+2Aba+b^2B) \tan(c+dx)a^4) - b(-Ba^5+3Aba^4-6b^2Ba^3+11Ab^3a^2-3b^4Ba+6Ab^5) \tan(c+dx)^2 + (a^2+b^2)^2(Aa^2+3bBa-6Ab^2)}{a(a^2+b^2)^2} dx - \frac{b(a^5(-B)+3Aba^4-6b^2Ba^3+11Ab^3a^2-3b^4Ba+6Ab^5)}{2ad(a^2+b^2)^2(a+b \tan(c+dx))^2}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^2}$$

↓ 4134

3.289. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{(a^2+b^2)^2(a^2A+3abB-6Ab^2) \int \cot(c+dx) dx + \frac{b^3(-10a^5B+15a^4Ab-9a^3b^2B+17a^2Ab^3-3ab^4B+6Ab^5) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx - a^4x(a^3(-B)+3a^2Ab+3ab^2B)}{a(a^2+b^2)}}{a(a^2+b^2)}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^2}$$

↓ 3042

$$\frac{(a^2+b^2)^2(a^2A+3abB-6Ab^2) \int -\tan(c+dx+\frac{\pi}{2}) dx + \frac{b^3(-10a^5B+15a^4Ab-9a^3b^2B+17a^2Ab^3-3ab^4B+6Ab^5) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx - a^4x(a^3(-B)+3a^2Ab+3ab^2B)}{a(a^2+b^2)}}{a(a^2+b^2)}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^2}$$

↓ 25

$$\frac{(a^2+b^2)^2(a^2A+3abB-6Ab^2) \int \tan(\frac{1}{2}(2c+\pi)+dx) dx + \frac{b^3(-10a^5B+15a^4Ab-9a^3b^2B+17a^2Ab^3-3ab^4B+6Ab^5) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx - a^4x(a^3(-B)+3a^2Ab+3ab^2B)}{a(a^2+b^2)}}{a(a^2+b^2)}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^2}$$

↓ 3956

$$\frac{b^3(-10a^5B+15a^4Ab-9a^3b^2B+17a^2Ab^3-3ab^4B+6Ab^5) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx + \frac{(a^2+b^2)^2(a^2A+3abB-6Ab^2) \log(-\sin(c+dx)) - a^4x(a^3(-B)+3a^2Ab+3ab^2B)}{ad}}{a(a^2+b^2)}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^2}$$

↓ 4013

3.289. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{\frac{(a^2+b^2)^2 (a^2 A+3abB-6Ab^2) \log(-\sin(c+dx))}{ad} - \frac{a^4 x (a^3(-B)+3a^2 Ab+3ab^2 B-Ab^3)}{a^2+b^2} + \frac{b^3 (-10a^5 B+15a^4 Ab-9a^3 b^2 B+17a^2 Ab^3-3ab^4 B+6Ab^5) \log(a \cos(c+dx))}{ad(a^2+b^2)}}{a(a^2+b^2)} = \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^2}$$

```
input Int[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]
```

```
output -1/2*(A*Cot[c + d*x]^2)/(a*d*(a + b*Tan[c + d*x])^2) - (-(((2*A*b - a*B)*Cot[c + d*x])/(a*d*(a + b*Tan[c + d*x])^2)) + (-1/2*(b*(5*a^2*A*b + 6*A*b^3 - 2*a^3*B - 3*a*b^2*B))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + ((-(a^4*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B)*x)/(a^2 + b^2)) + ((a^2 + b^2)^2*(a^2*A - 6*A*b^2 + 3*a*b*B)*Log[-Sin[c + d*x]])/(a*d) + (b^3*(15*a^4*A*b + 17*a^2*A*b^3 + 6*A*b^5 - 10*a^5*B - 9*a^3*b^2*B - 3*a*b^4*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))/(a*(a^2 + b^2)) - (b*(3*a^4*A*b + 11*a^2*A*b^3 + 6*A*b^5 - a^5*B - 6*a^3*b^2*B - 3*a*b^4*B))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(a*(a^2 + b^2)))/a/a
```

3.289.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4132 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4134 `Int[((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2)/(((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.289.4 Maple [A] (verified)

Time = 0.71 (sec) , antiderivative size = 320, normalized size of antiderivative = 0.91

method	result
derivativedivides	$\frac{(A a^3 - 3A a b^2 + 3B a^2 b - B b^3) \ln(1 + \tan^2(dx+c)) + (3A a^2 b - A b^3 - B a^3 + 3B a b^2) \arctan(\tan(dx+c))}{(a^2 + b^2)^3} - \frac{A}{2a^3 \tan(dx+c)^2} - \frac{-3Ab+}{a^4 \tan(dx+c)}$
default	$\frac{(A a^3 - 3A a b^2 + 3B a^2 b - B b^3) \ln(1 + \tan^2(dx+c)) + (3A a^2 b - A b^3 - B a^3 + 3B a b^2) \arctan(\tan(dx+c))}{(a^2 + b^2)^3} - \frac{A}{2a^3 \tan(dx+c)^2} - \frac{-3Ab+}{a^4 \tan(dx+c)}$
parallelrisch	$-30b^3 (A a^4 b + \frac{17}{15} A a^2 b^3 + \frac{2}{5} A b^5 - \frac{2}{3} B a^5 - \frac{3}{5} B a^3 b^2 - \frac{1}{5} B a b^4) (a+b \tan(dx+c))^2 \ln(a+b \tan(dx+c)) + a^5 (a+b \tan(dx+c))$
norman	$\frac{(2Ab-Ba) \tan(dx+c)}{a^2 d} + \frac{b^2 (3A a^2 b - A b^3 - B a^3 + 3B a b^2) x (\tan^4(dx+c))}{(a^4 + 2a^2 b^2 + b^4) (a^2 + b^2)} + \frac{(3A a^2 b - A b^3 - B a^3 + 3B a b^2) a^2 x (\tan^2(dx+c))}{(a^4 + 2a^2 b^2 + b^4) (a^2 + b^2)} - \frac{A}{2ad}$
risch	Expression too large to display

input `int(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} \left(\frac{1}{(a^2+b^2)^3} \left(\frac{1}{2} (A a^3 - 3A a b^2 + 3B a^2 b - B b^3) \ln(1 + \tan^2(dx+c)) + (3A a^2 b - A b^3 - B a^3 + 3B a b^2) \arctan(\tan(dx+c)) \right) - \frac{1}{2} \frac{A}{a^3 \tan(dx+c)^2} - \frac{(-3A b + B a)}{a^4 \tan(dx+c)} + \frac{(-A a^2 + 6A b^2 - 3B a b)}{a^5 \ln(\tan(dx+c))} + b^3 \frac{(5A a^2 b + 3A b^3 - 4B a^3 - 2B a b^2)}{(a^2 + b^2)^2 a^4} \frac{1}{(a+b \tan(dx+c))} - b^3 \frac{(15A a^4 b + 17A a^2 b^3 + 6A b^5 - 10B a^5 - 9B a^3 b^2 - 3B a b^4)}{(a^2 + b^2)^3 a^5 \ln(a+b \tan(dx+c))} + \frac{1}{2} \frac{(A b - B a) b^3}{(a^2 + b^2) a^3} \frac{1}{(a+b \tan(dx+c))^2} \right)$$

3.289.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1065 vs. 2(346) = 692.

Time = 0.40 (sec) , antiderivative size = 1065, normalized size of antiderivative = 3.03

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx = \frac{A a^{10} + 3 A a^8 b^2 + 3 A a^6 b^4 + A a^4 b^6 + (A a^8 b^2 + 3 A a^6 b^4 - 9 B a^5 b^5 + 14 A a^4 b^6 - 3 B a^3 b^7 + 6 A a^2 b^8 + 2 B a b^9 - 3 B^2 a^2 b^6 + 3 B^2 a b^8 - 3 B^2 b^{10})}{(a^2 + b^2)^3}$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="fricas")`

output `-1/2*(A*a^10 + 3*A*a^8*b^2 + 3*A*a^6*b^4 + A*a^4*b^6 + (A*a^8*b^2 + 3*A*a^6*b^4 - 9*B*a^5*b^5 + 14*A*a^4*b^6 - 3*B*a^3*b^7 + 6*A*a^2*b^8 + 2*(B*a^8*b^2 - 3*A*a^7*b^3 - 3*B*a^6*b^4 + A*a^5*b^5)*d*x)*tan(d*x + c)^4 + 2*(A*a^9*b + B*a^8*b^2 - 2*B*a^6*b^4 + 6*B*a^4*b^6 - 11*A*a^3*b^7 + 3*B*a^2*b^8 - 6*A*a*b^9 + 2*(B*a^9*b - 3*A*a^8*b^2 - 3*B*a^7*b^3 + A*a^6*b^4)*d*x)*tan(d*x + c)^3 + (A*a^10 + 4*B*a^9*b - 8*A*a^8*b^2 + 12*B*a^7*b^3 - 30*A*a^6*b^4 + 23*B*a^5*b^5 - 45*A*a^4*b^6 + 9*B*a^3*b^7 - 18*A*a^2*b^8 + 2*(B*a^10 - 3*A*a^9*b - 3*B*a^8*b^2 + A*a^7*b^3)*d*x)*tan(d*x + c)^2 + ((A*a^8*b^2 + 3*B*a^7*b^3 - 3*A*a^6*b^4 + 9*B*a^5*b^5 - 15*A*a^4*b^6 + 9*B*a^3*b^7 - 17*A*a^2*b^8 + 3*B*a*b^9 - 6*A*b^10)*tan(d*x + c)^4 + 2*(A*a^9*b + 3*B*a^8*b^2 - 3*A*a^7*b^3 + 9*B*a^6*b^4 - 15*A*a^5*b^5 + 9*B*a^4*b^6 - 17*A*a^3*b^7 + 3*B*a^2*b^8 - 6*A*a*b^9)*tan(d*x + c)^3 + (A*a^10 + 3*B*a^9*b - 3*A*a^8*b^2 + 9*B*a^7*b^3 - 15*A*a^6*b^4 + 9*B*a^5*b^5 - 17*A*a^4*b^6 + 3*B*a^3*b^7 - 6*A*a^2*b^8)*tan(d*x + c)^2)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1)) - ((10*B*a^5*b^5 - 15*A*a^4*b^6 + 9*B*a^3*b^7 - 17*A*a^2*b^8 + 3*B*a*b^9 - 6*A*b^10)*tan(d*x + c)^4 + 2*(10*B*a^6*b^4 - 15*A*a^5*b^5 + 9*B*a^4*b^6 - 17*A*a^3*b^7 + 3*B*a^2*b^8 - 6*A*a*b^9)*tan(d*x + c)^3 + (10*B*a^7*b^3 - 15*A*a^6*b^4 + 9*B*a^5*b^5 - 17*A*a^4*b^6 + 3*B*a^3*b^7 - 6*A*a^2*b^8)*tan(d*x + c)^2)*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) + 2*(B*a^10 - 2*A*a^9*b + 3*B*a^8*b^2 - 6*A*a^7*b^3 + 3*B...`

3.289.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Exception raised: AttributeError}$$

input `integrate(cot(d*x+c)**3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)`

output `Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'`

3.289.7 Maxima [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 541, normalized size of antiderivative = 1.54

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx =$$

$$\frac{2(Ba^3-3Aa^2b-3Bab^2+Ab^3)(dx+c)}{a^6+3a^4b^2+3a^2b^4+b^6} - \frac{2(10Ba^5b^3-15Aa^4b^4+9Ba^3b^5-17Aa^2b^6+3Bab^7-6Ab^8)\log(b\tan(dx+c)+a)}{a^{11}+3a^9b^2+3a^7b^4+a^5b^6} - \frac{(Aa^3+3Ba^2b+3Ab^3)\log(\tan(dx+c)+1)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{(Aa^7+2Aa^5b^2+Aa^3b^4+2(Ba^5b^2-3Aa^4b^3+6Ba^3b^4-11Aa^2b^5+3Ba^2b^6-6Aa^2b^7)\tan(dx+c)^3+(4Ba^6b-11Aa^5b^2+17Ba^4b^3-33Aa^3b^4+9Ba^2b^5-18Aa^2b^6)\tan(dx+c)^2+2(Ba^7-2Aa^6b+2Ba^5b^2-4Aa^4b^3+Ba^3b^4-2Aa^2b^5)\tan(dx+c))}{(a^8b^2+2a^6b^4+a^4b^6)\tan(dx+c)^4+2(a^9b+2a^7b^3+a^5b^5)\tan(dx+c)^3+(a^{10}+2a^8b^2+a^6b^4)\tan(dx+c)^2+2(Aa^2+3Ba^2b-6Aa^2b^2)\log(\tan(dx+c))/a^5}/d$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="maxima")
```

```
output -1/2*(2*(B*a^3 - 3*A*a^2*b - 3*B*a*b^2 + A*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - 2*(10*B*a^5*b^3 - 15*A*a^4*b^4 + 9*B*a^3*b^5 - 17*A*a^2*b^6 + 3*B*a*b^7 - 6*A*b^8)*log(b*tan(d*x + c) + a)/(a^11 + 3*a^9*b^2 + 3*a^7*b^4 + a^5*b^6) - (A*a^3 + 3*B*a^2*b - 3*A*a*b^2 - B*b^3)*log(tan(d*x + c)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + (A*a^7 + 2*A*a^5*b^2 + A*a^3*b^4 + 2*(B*a^5*b^2 - 3*A*a^4*b^3 + 6*B*a^3*b^4 - 11*A*a^2*b^5 + 3*B*a*b^6 - 6*A*b^7)*tan(d*x + c)^3 + (4*B*a^6*b - 11*A*a^5*b^2 + 17*B*a^4*b^3 - 33*A*a^3*b^4 + 9*B*a^2*b^5 - 18*A*a*b^6)*tan(d*x + c)^2 + 2*(B*a^7 - 2*A*a^6*b + 2*B*a^5*b^2 - 4*A*a^4*b^3 + B*a^3*b^4 - 2*A*a^2*b^5)*tan(d*x + c))/(a^8*b^2 + 2*a^6*b^4 + a^4*b^6)*tan(d*x + c)^4 + 2*(a^9*b + 2*a^7*b^3 + a^5*b^5)*tan(d*x + c)^3 + (a^10 + 2*a^8*b^2 + a^6*b^4)*tan(d*x + c)^2 + 2*(A*a^2 + 3*B*a*b - 6*A*b^2)*log(tan(d*x + c))/a^5)/d
```

3.289.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 812 vs. 2(346) = 692.

Time = 1.24 (sec) , antiderivative size = 812, normalized size of antiderivative = 2.31

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx =$$

$$\frac{4(Ba^3-3Aa^2b-3Bab^2+Ab^3)(dx+c)}{a^6+3a^4b^2+3a^2b^4+b^6} - \frac{2(Aa^3+3Ba^2b-3Aab^2-Bb^3)\log(\tan(dx+c)^2+1)}{a^6+3a^4b^2+3a^2b^4+b^6} - \frac{4(10Ba^5b^4-15Aa^4b^5+9Ba^3b^6-17Aa^2b^7+3Bab^8-6Ab^9)\log(b\tan(dx+c)+a)}{a^{11}b+3a^9b^3+3a^7b^5+a^5b^7+b^9} + \frac{(Aa^3+3Ba^2b+3Ab^3)\log(\tan(dx+c)+1)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{(Aa^7+2Aa^5b^2+Aa^3b^4+2(Ba^5b^2-3Aa^4b^3+6Ba^3b^4-11Aa^2b^5+3Ba^2b^6-6Aa^2b^7)\tan(dx+c)^3+(4Ba^6b-11Aa^5b^2+17Ba^4b^3-33Aa^3b^4+9Ba^2b^5-18Aa^2b^6)\tan(dx+c)^2+2(Ba^7-2Aa^6b+2Ba^5b^2-4Aa^4b^3+Ba^3b^4-2Aa^2b^5)\tan(dx+c))}{(a^8b^2+2a^6b^4+a^4b^6)\tan(dx+c)^4+2(a^9b+2a^7b^3+a^5b^5)\tan(dx+c)^3+(a^{10}+2a^8b^2+a^6b^4)\tan(dx+c)^2+2(Aa^2+3Ba^2b-6Aa^2b^2)\log(\tan(dx+c))/a^5}/d$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="giac")
```

$$3.289. \int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

output

$$\begin{aligned}
& -1/4*(4*(B*a^3 - 3*A*a^2*b - 3*B*a*b^2 + A*b^3)*(d*x + c)/(a^6 + 3*a^4*b^2 \\
& + 3*a^2*b^4 + b^6) - 2*(A*a^3 + 3*B*a^2*b - 3*A*a*b^2 - B*b^3)*\log(\tan(d*x \\
& + c)^2 + 1)/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - 4*(10*B*a^5*b^4 - 15*A \\
& *a^4*b^5 + 9*B*a^3*b^6 - 17*A*a^2*b^7 + 3*B*a*b^8 - 6*A*b^9)*\log(\text{abs}(b*\tan \\
& (d*x + c) + a))/(a^{11}*b + 3*a^9*b^3 + 3*a^7*b^5 + a^5*b^7) - (3*A*a^7*b^2* \\
& \tan(d*x + c)^4 + 9*B*a^6*b^3*\tan(d*x + c)^4 - 9*A*a^5*b^4*\tan(d*x + c)^4 - \\
& 3*B*a^4*b^5*\tan(d*x + c)^4 + 6*A*a^8*b*\tan(d*x + c)^3 + 14*B*a^7*b^2*\tan(\\
& d*x + c)^3 - 6*A*a^6*b^3*\tan(d*x + c)^3 - 34*B*a^5*b^4*\tan(d*x + c)^3 + 56 \\
& *A*a^4*b^5*\tan(d*x + c)^3 - 36*B*a^3*b^6*\tan(d*x + c)^3 + 68*A*a^2*b^7*\tan \\
& (d*x + c)^3 - 12*B*a*b^8*\tan(d*x + c)^3 + 24*A*b^9*\tan(d*x + c)^3 + 3*A*a^ \\
& 9*\tan(d*x + c)^2 + B*a^8*b*\tan(d*x + c)^2 + 13*A*a^7*b^2*\tan(d*x + c)^2 - \\
& 45*B*a^6*b^3*\tan(d*x + c)^2 + 88*A*a^5*b^4*\tan(d*x + c)^2 - 52*B*a^4*b^5*t \\
& \tan(d*x + c)^2 + 102*A*a^3*b^6*\tan(d*x + c)^2 - 18*B*a^2*b^7*\tan(d*x + c)^2 \\
& + 36*A*a*b^8*\tan(d*x + c)^2 - 4*B*a^9*\tan(d*x + c) + 8*A*a^8*b*\tan(d*x + \\
& c) - 12*B*a^7*b^2*\tan(d*x + c) + 24*A*a^6*b^3*\tan(d*x + c) - 12*B*a^5*b^4* \\
& \tan(d*x + c) + 24*A*a^4*b^5*\tan(d*x + c) - 4*B*a^3*b^6*\tan(d*x + c) + 8*A* \\
& a^2*b^7*\tan(d*x + c) - 2*A*a^9 - 6*A*a^7*b^2 - 6*A*a^5*b^4 - 2*A*a^3*b^6)/ \\
& ((a^{10} + 3*a^8*b^2 + 3*a^6*b^4 + a^4*b^6)*(b*\tan(d*x + c)^2 + a*\tan(d*x + \\
& c))^2) + 4*(A*a^2 + 3*B*a*b - 6*A*b^2)*\log(\text{abs}(\tan(d*x + c)))/a^5/d
\end{aligned}$$

3.289.9 Mupad [B] (verification not implemented)

Time = 14.25 (sec) , antiderivative size = 434, normalized size of antiderivative = 1.23

$$\begin{aligned}
& \int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx \\
& = \frac{\frac{\tan(c+dx)(2Ab-Ba)}{a^2} - \frac{A}{2a} + \frac{\tan(c+dx)^3(-Ba^5b^2+3Aa^4b^3-6Ba^3b^4+11Aa^2b^5-3Ba^2b^6+6Ab^7)}{a^4(a^4+2a^2b^2+b^4)} + \frac{\tan(c+dx)^2(-4Ba^5b+11Aa^4b^2)}{2a^4}}{d(a^2\tan(c+dx)^2+2ab\tan(c+dx)^3+b^2\tan(c+dx)^4)} \\
& + \frac{\ln(a+b\tan(c+dx))\left(\frac{A}{a^3} - \frac{Aa+3Bb}{(a^2+b^2)^2} - \frac{6Ab^2}{a^5} + \frac{3Bb}{a^4} + \frac{4b^2(Aa+Bb)}{(a^2+b^2)^3}\right)}{d} \\
& - \frac{\ln(\tan(c+dx)-i)(-B+Ai)}{2d(-a^3+3a^2b+ab^2-3b^3)} - \frac{\ln(\tan(c+dx))(Aa^2+3Bab-6Ab^2)}{a^5d} \\
& - \frac{\ln(\tan(c+dx)+i)(A-Bi)}{2d(-a^3+a^2b+3ab^2-b^3)}
\end{aligned}$$

input `int((cot(c + d*x))^3*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3,x`

output

$$\begin{aligned} & ((\tan(c + dx) * (2 * A * b - B * a)) / a^2 - A / (2 * a) + (\tan(c + dx)^3 * (6 * A * b^7 + 1 \\ & 1 * A * a^2 * b^5 + 3 * A * a^4 * b^3 - 6 * B * a^3 * b^4 - B * a^5 * b^2 - 3 * B * a * b^6)) / (a^4 * (a^4 \\ & 4 + b^4 + 2 * a^2 * b^2)) + (\tan(c + dx)^2 * (18 * A * b^6 + 33 * A * a^2 * b^4 + 11 * A * a^4 * b^2 \\ & 4 * b^2 - 17 * B * a^3 * b^3 - 9 * B * a * b^5 - 4 * B * a^5 * b)) / (2 * a^3 * (a^4 + b^4 + 2 * a^2 * b^2))) \\ & / (d * (a^2 * \tan(c + dx)^2 + b^2 * \tan(c + dx)^4 + 2 * a * b * \tan(c + dx)^3)) \\ & + (\log(a + b * \tan(c + dx)) * (A / a^3 - (A * a + 3 * B * b) / (a^2 + b^2)^2 - (6 * A * b^2) / a^5 \\ & + (3 * B * b) / a^4 + (4 * b^2 * (A * a + B * b)) / (a^2 + b^2)^3)) / d - (\log(\tan(c + dx) - 1i) * (A * 1i - B)) \\ & / (2 * d * (a * b^2 * 3i + 3 * a^2 * b - a^3 * 1i - b^3)) - (\log(\tan(c + dx)) * (A * a^2 - 6 * A * b^2 + 3 * B * a * b)) \\ & / (a^5 * d) - (\log(\tan(c + dx) + 1i) * (A - B * 1i)) / (2 * d * (3 * a * b^2 + a^2 * b * 3i - a^3 - b^3 * 1i)) \end{aligned}$$

3.290
$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

3.290.1 Optimal result 2787
 3.290.2 Mathematica [C] (verified) 2788
 3.290.3 Rubi [A] (verified) 2788
 3.290.4 Maple [A] (verified) 2794
 3.290.5 Fricas [B] (verification not implemented) 2794
 3.290.6 Sympy [F(-2)] 2795
 3.290.7 Maxima [A] (verification not implemented) 2796
 3.290.8 Giac [B] (verification not implemented) 2796
 3.290.9 Mupad [B] (verification not implemented) 2797

3.290.1 Optimal result

Integrand size = 31, antiderivative size = 351

$$\begin{aligned} & \int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx \\ &= \frac{(a^4A - 6a^2Ab^2 + Ab^4 + 4a^3bB - 4ab^3B)x}{(a^2 + b^2)^4} \\ &+ \frac{(4a^3Ab - 4aAb^3 - a^4B + 6a^2b^2B - b^4B) \log(\cos(c+dx))}{(a^2 + b^2)^4 d} \\ &+ \frac{a(4a^2Ab^5 - 4Ab^7 + a^7B + 4a^5b^2B + 5a^3b^4B + 10ab^6B) \log(a+b \tan(c+dx))}{b^4 (a^2 + b^2)^4 d} \\ &+ \frac{a(Ab - aB) \tan^3(c+dx)}{3b(a^2 + b^2) d(a+b \tan(c+dx))^3} + \frac{a(2Ab^3 - a(a^2 + 3b^2)B) \tan^2(c+dx)}{2b^2(a^2 + b^2)^2 d(a+b \tan(c+dx))^2} \\ &+ \frac{a^2(a^2Ab^3 - 3Ab^5 + a^5B + 3a^3b^2B + 6ab^4B)}{b^4(a^2 + b^2)^3 d(a+b \tan(c+dx))} \end{aligned}$$

output

```
(A*a^4-6*A*a^2*b^2+A*b^4+4*B*a^3*b-4*B*a*b^3)*x/(a^2+b^2)^4+(4*A*a^3*b-4*A*a*b^3-B*a^4+6*B*a^2*b^2-B*b^4)*ln(cos(d*x+c))/(a^2+b^2)^4/d+a*(4*A*a^2*b^5-4*A*b^7+B*a^7+4*B*a^5*b^2+5*B*a^3*b^4+10*B*a*b^6)*ln(a+b*tan(d*x+c))/b^4/(a^2+b^2)^4/d+1/3*a*(A*b-B*a)*tan(d*x+c)^3/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^3+1/2*a*(2*A*b^3-a*(a^2+3*b^2)*B)*tan(d*x+c)^2/b^2/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^2+a^2*(A*a^2*b^3-3*A*b^5+B*a^5+3*B*a^3*b^2+6*B*a*b^4)/b^4/(a^2+b^2)^3/d/(a+b*tan(d*x+c))
```

3.290.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.56 (sec) , antiderivative size = 307, normalized size of antiderivative = 0.87

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

$$= \frac{\frac{3(-iA+B) \log(i-\tan(c+dx))}{(a+ib)^4} + \frac{3(iA+B) \log(i+\tan(c+dx))}{(a-ib)^4} + \frac{6a(4a^2Ab^5-4Ab^7+a^7B+4a^5b^2B+5a^3b^4B+10ab^6B) \log(a+b \tan(c+dx))}{b^4(a^2+b^2)^4}}{6d}$$

input `Integrate[(Tan[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]`

output `((3*((-I)*A + B)*Log[I - Tan[c + d*x]])/(a + I*b)^4 + (3*(I*A + B)*Log[I + Tan[c + d*x]])/(a - I*b)^4 + (6*a*(4*a^2*A*b^5 - 4*A*b^7 + a^7*B + 4*a^5*b^2*B + 5*a^3*b^4*B + 10*a*b^6*B)*Log[a + b*Tan[c + d*x]])/(b^4*(a^2 + b^2)^4) + (2*a^4*(-(A*b) + a*B))/(b^4*(a^2 + b^2)*(a + b*Tan[c + d*x])^3) + (3*a^3*(2*a^2*A*b + 4*A*b^3 - 3*a^3*B - 5*a*b^2*B))/(b^4*(a^2 + b^2)^2*(a + b*Tan[c + d*x])^2) + (6*a^2*(-(a^4*A*b) - 3*a^2*A*b^3 - 6*A*b^5 + 3*a^5*B + 9*a^3*b^2*B + 10*a*b^4*B))/(b^4*(a^2 + b^2)^3*(a + b*Tan[c + d*x]))/(6*d)`

3.290.3 Rubi [A] (verified)

Time = 1.86 (sec) , antiderivative size = 402, normalized size of antiderivative = 1.15, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4118, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

↓ 3042

$$\int \frac{\tan(c+dx)^4(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

↓ 4088

$$\begin{aligned}
 & \int \frac{-3 \tan^2(c+dx) \left(-((a^2+b^2)B \tan^2(c+dx)) - b(Ab-aB) \tan(c+dx) + a(Ab-aB) \right) dx}{(a+b \tan(c+dx))^3} + \\
 & \quad \frac{3b(a^2+b^2)}{a(Ab-aB) \tan^3(c+dx)} \\
 & \quad \frac{3bd(a^2+b^2)(a+b \tan(c+dx))^3}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} \\
 & \quad \downarrow 27 \\
 & \quad \frac{a(Ab-aB) \tan^3(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} - \\
 & \int \frac{\tan^2(c+dx) \left(-((a^2+b^2)B \tan^2(c+dx)) - b(Ab-aB) \tan(c+dx) + a(Ab-aB) \right) dx}{(a+b \tan(c+dx))^3} \\
 & \quad \frac{b(a^2+b^2)}{b(a^2+b^2)} \\
 & \quad \downarrow 3042 \\
 & \quad \frac{a(Ab-aB) \tan^3(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} - \\
 & \int \frac{\tan(c+dx)^2 \left(-((a^2+b^2)B \tan(c+dx)^2) - b(Ab-aB) \tan(c+dx) + a(Ab-aB) \right) dx}{(a+b \tan(c+dx))^3} \\
 & \quad \frac{b(a^2+b^2)}{b(a^2+b^2)} \\
 & \quad \downarrow 4128 \\
 & \quad \frac{a(Ab-aB) \tan^3(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} - \\
 & \int \frac{2 \tan(c+dx) \left((Aa^2+2bBa-Ab^2) \tan(c+dx)b^2 - (a^2+b^2)^2 B \tan^2(c+dx) + a(2Ab^3-a(a^2+3b^2)B) \right) dx}{(a+b \tan(c+dx))^2} \\
 & \quad \frac{2b(a^2+b^2)}{2b(a^2+b^2)} - \frac{a(2Ab^3-aB(a^2+3b^2)) \tan^2(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \frac{b(a^2+b^2)}{b(a^2+b^2)} \\
 & \quad \downarrow 27 \\
 & \quad \frac{a(Ab-aB) \tan^3(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} - \\
 & \int \frac{\tan(c+dx) \left((Aa^2+2bBa-Ab^2) \tan(c+dx)b^2 - (a^2+b^2)^2 B \tan^2(c+dx) + a(2Ab^3-a(a^2+3b^2)B) \right) dx}{(a+b \tan(c+dx))^2} \\
 & \quad \frac{b(a^2+b^2)}{b(a^2+b^2)} - \frac{a(2Ab^3-aB(a^2+3b^2)) \tan^2(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \frac{b(a^2+b^2)}{b(a^2+b^2)} \\
 & \quad \downarrow 3042 \\
 & \quad \frac{a(Ab-aB) \tan^3(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} - \\
 & \int \frac{\tan(c+dx) \left((Aa^2+2bBa-Ab^2) \tan(c+dx)b^2 - (a^2+b^2)^2 B \tan^2(c+dx)^2 + a(2Ab^3-a(a^2+3b^2)B) \right) dx}{(a+b \tan(c+dx))^2} \\
 & \quad \frac{b(a^2+b^2)}{b(a^2+b^2)} - \frac{a(2Ab^3-aB(a^2+3b^2)) \tan^2(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \frac{b(a^2+b^2)}{b(a^2+b^2)} \\
 & \quad \downarrow 4118
 \end{aligned}$$

3.290. $\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

$$\frac{a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} - \frac{\int -\left(\frac{(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \tan(c + dx)b^3}{a + b \tan(c + dx)} + (a^2 + b^2)^3 B \tan^2(c + dx) + a(Ba^5 + 3b^2Ba^3 + Ab^3a^2 + 6b^4Ba - 3Ab^5)\right) dx}{b(a^2 + b^2)} - \frac{a^2(a^5B + 3a^3b^2B + a^2Ab^3 + 6ab^4B - b^2d(a^2 + b^2)(a + b \tan(c + dx)))}{b^2d(a^2 + b^2)(a + b \tan(c + dx))}$$

$$b(a^2 + b^2)$$

↓ 25

$$\frac{a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} - \frac{\int -\left(\frac{(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \tan(c + dx)b^3}{a + b \tan(c + dx)} + (a^2 + b^2)^3 B \tan^2(c + dx) + a(Ba^5 + 3b^2Ba^3 + Ab^3a^2 + 6b^4Ba - 3Ab^5)\right) dx}{b(a^2 + b^2)} - \frac{a^2(a^5B + 3a^3b^2B + a^2Ab^3 + 6ab^4B - b^2d(a^2 + b^2)(a + b \tan(c + dx)))}{b^2d(a^2 + b^2)(a + b \tan(c + dx))}$$

$$b(a^2 + b^2)$$

↓ 3042

$$\frac{a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} - \frac{\int -\left(\frac{(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \tan(c + dx)b^3}{a + b \tan(c + dx)} + (a^2 + b^2)^3 B \tan^2(c + dx) + a(Ba^5 + 3b^2Ba^3 + Ab^3a^2 + 6b^4Ba - 3Ab^5)\right) dx}{b(a^2 + b^2)} - \frac{a^2(a^5B + 3a^3b^2B + a^2Ab^3 + 6ab^4B - b^2d(a^2 + b^2)(a + b \tan(c + dx)))}{b^2d(a^2 + b^2)(a + b \tan(c + dx))}$$

$$b(a^2 + b^2)$$

↓ 4109

$$\frac{a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} - \frac{b^3(a^4(-B) + 4a^3Ab + 6a^2b^2B - 4aAb^3 - b^4B) \int \tan(c + dx) dx}{a^2 + b^2} + \frac{a(a^7B + 4a^5b^2B + 5a^3b^4B + 4a^2Ab^5 + 10ab^6B - 4Ab^7) \int \frac{\tan^2(c + dx) + 1}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{b^3x(a^4A + 4a^3bB - 6a^2b^2d(a^2 + b^2)(a + b \tan(c + dx)))}{a^2}$$

$$b(a^2 + b^2)$$

↓ 3042

$$\frac{a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} - \frac{b^3(a^4(-B) + 4a^3Ab + 6a^2b^2B - 4aAb^3 - b^4B) \int \tan(c + dx) dx}{a^2 + b^2} + \frac{a(a^7B + 4a^5b^2B + 5a^3b^4B + 4a^2Ab^5 + 10ab^6B - 4Ab^7) \int \frac{\tan(c + dx)^2 + 1}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{b^3x(a^4A + 4a^3bB - 6a^2b^2d(a^2 + b^2)(a + b \tan(c + dx)))}{a^2}$$

$$b(a^2 + b^2)$$

↓ 3956

3.290. $\int \frac{\tan^4(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^4} dx$

$$\frac{a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} - \frac{a(a^7B + 4a^5b^2B + 5a^3b^4B + 4a^2Ab^5 + 10ab^6B - 4Ab^7) \int \frac{\tan(c+dx)^2 + 1}{a+b \tan(c+dx)} dx}{a^2 + b^2} + \frac{b^3(a^4(-B) + 4a^3Ab + 6a^2b^2B - 4aAb^3 - b^4B) \log(\cos(c+dx))}{d(a^2 + b^2)} + \frac{b^3x(a^4A + 4a^3bB - 6a^2b^2B^2 - 4ab^3B + Ab^4)}{a^2 + b^2}$$

$$\frac{a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} - \frac{a(a^7B + 4a^5b^2B + 5a^3b^4B + 4a^2Ab^5 + 10ab^6B - 4Ab^7) \int \frac{1}{a+b \tan(c+dx)} d(b \tan(c+dx))}{bd(a^2 + b^2)} + \frac{b^3(a^4(-B) + 4a^3Ab + 6a^2b^2B - 4aAb^3 - b^4B) \log(\cos(c+dx))}{d(a^2 + b^2)} + \frac{b^3x(a^4A + 4a^3bB - 6a^2b^2B^2 - 4ab^3B + Ab^4)}{a^2 + b^2}$$

$$\frac{a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} - \frac{a^2(a^5B + 3a^3b^2B + a^2Ab^3 + 6ab^4B - 3Ab^5)}{b^2d(a^2 + b^2)(a + b \tan(c + dx))} + \frac{b^3(a^4(-B) + 4a^3Ab + 6a^2b^2B - 4aAb^3 - b^4B) \log(\cos(c+dx))}{d(a^2 + b^2)} + \frac{b^3x(a^4A + 4a^3bB - 6a^2b^2B^2 - 4ab^3B + Ab^4)}{a^2 + b^2} + \frac{a(a^7B + 4a^5b^2B + 5a^3b^4B + 4a^2Ab^5 + 10ab^6B - 4Ab^7)}{b^2d(a^2 + b^2)(a + b \tan(c + dx))}$$

4100

16

```
input Int[(Tan[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]
```

```
output (a*(A*b - a*B)*Tan[c + d*x]^3)/(3*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^3 - (-1/2*(a*(2*A*b^3 - a*(a^2 + 3*b^2)*B)*Tan[c + d*x]^2)/(b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + (-(((b^3*(a^4*A - 6*a^2*A*b^2 + A*b^4 + 4*a^3*b*B - 4*a*b^3*B)*x)/(a^2 + b^2) + (b^3*(4*a^3*A*b - 4*a*A*b^3 - a^4*B + 6*a^2*b^2*B - b^4*B)*Log[Cos[c + d*x]])/((a^2 + b^2)*d) + (a*(4*a^2*A*b^5 - 4*A*b^7 + a^7*B + 4*a^5*b^2*B + 5*a^3*b^4*B + 10*a*b^6*B)*Log[a + b*Tan[c + d*x]])/(b*(a^2 + b^2)*d))/(b*(a^2 + b^2))) - (a^2*(a^2*A*b^3 - 3*A*b^5 + a^5*B + 3*a^3*b^2*B + 6*a*b^4*B))/(b^2*(a^2 + b^2)*d*(a + b*Tan[c + d*x])))/(b*(a^2 + b^2))/(b*(a^2 + b^2))
```

3.290.3.1 Defintions of rubi rules used

- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_.)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_.)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`
- rule 4100 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]`

rule 4109 `Int[((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*A + b*B - a*C)*(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C, 0]`

rule 4118 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^n*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-(b*c - a*d))*(c^2*C - B*c*d + A*d^2)*((c + d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`

rule 4128 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^m*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^n*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

3.290.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 343, normalized size of antiderivative = 0.98

method	result
derivativedivides	$\frac{(-4A a^3 b + 4A a b^3 + B a^4 - 6B a^2 b^2 + B b^4) \ln(1 + \tan^2(dx+c))}{2(a^2+b^2)^4} + \frac{(A a^4 - 6A a^2 b^2 + A b^4 + 4B a^3 b - 4B a b^3) \arctan(\tan(dx+c))}{(a^2+b^2)^4} + \frac{a(4A a^3 b - 4A a b^3 + B a^4 - 6B a^2 b^2 + B b^4)}{(a^2+b^2)^4}$
default	$\frac{(-4A a^3 b + 4A a b^3 + B a^4 - 6B a^2 b^2 + B b^4) \ln(1 + \tan^2(dx+c))}{2(a^2+b^2)^4} + \frac{(A a^4 - 6A a^2 b^2 + A b^4 + 4B a^3 b - 4B a b^3) \arctan(\tan(dx+c))}{(a^2+b^2)^4} + \frac{a(4A a^3 b - 4A a b^3 + B a^4 - 6B a^2 b^2 + B b^4)}{(a^2+b^2)^4}$
norman	$\frac{(A a^4 - 6A a^2 b^2 + A b^4 + 4B a^3 b - 4B a b^3) a^3 x}{(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)(a^2 + b^2)} + \frac{b^3 (A a^4 - 6A a^2 b^2 + A b^4 + 4B a^3 b - 4B a b^3) x (\tan^3(dx+c))}{(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)(a^2 + b^2)} - \frac{a^3 (2A a^5 b + 4A a^3 b^3 + 2A a b^5)}{6d b^4 (a^2 + b^2)}$
risch	Expression too large to display
parallelrisc	Expression too large to display

input `int(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `1/d*(1/(a^2+b^2)^4*(1/2*(-4*A*a^3*b+4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*ln(1+tan(d*x+c)^2)+(A*a^4-6*A*a^2*b^2+A*b^4+4*B*a^3*b-4*B*a*b^3)*arctan(tan(d*x+c)))+a*(4*A*a^2*b^5-4*A*b^7+B*a^7+4*B*a^5*b^2+5*B*a^3*b^4+10*B*a*b^6)/(a^2+b^2)^4/b^4*ln(a+b*tan(d*x+c))-a^2*(A*a^4*b+3*A*a^2*b^3+6*A*b^5-3*B*a^5-9*B*a^3*b^2-10*B*a*b^4)/b^4/(a^2+b^2)^3/(a+b*tan(d*x+c))-1/3*a^4*(A*b-B*a)/b^4/(a^2+b^2)/(a+b*tan(d*x+c))^3+1/2*a^3*(2*A*a^2*b+4*A*b^3-3*B*a^3-5*B*a*b^2)/b^4/(a^2+b^2)^2/(a+b*tan(d*x+c))^2)`

3.290.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1113 vs. 2(346) = 692.

Time = 0.36 (sec) , antiderivative size = 1113, normalized size of antiderivative = 3.17

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="fracas")`

```
output 1/6*(3*B*a^9*b^2 + 6*B*a^7*b^4 + 18*A*a^6*b^5 + 47*B*a^5*b^6 - 26*A*a^4*b^
7 - (11*B*a^8*b^3 - 2*A*a^7*b^4 + 42*B*a^6*b^5 - 6*A*a^5*b^6 + 75*B*a^4*b^
7 - 48*A*a^3*b^8 - 6*(A*a^4*b^7 + 4*B*a^3*b^8 - 6*A*a^2*b^9 - 4*B*a*b^10 +
A*b^11)*d*x)*tan(d*x + c)^3 + 6*(A*a^7*b^4 + 4*B*a^6*b^5 - 6*A*a^5*b^6 -
4*B*a^4*b^7 + A*a^3*b^8)*d*x - 3*(5*B*a^9*b^2 + 18*B*a^7*b^4 + 2*A*a^6*b^5
+ 37*B*a^5*b^6 - 30*A*a^4*b^7 - 20*B*a^3*b^8 + 12*A*a^2*b^9 - 6*(A*a^5*b^
6 + 4*B*a^4*b^7 - 6*A*a^3*b^8 - 4*B*a^2*b^9 + A*a*b^10)*d*x)*tan(d*x + c)^
2 + 3*(B*a^11 + 4*B*a^9*b^2 + 5*B*a^7*b^4 + 4*A*a^6*b^5 + 10*B*a^5*b^6 - 4
*A*a^4*b^7 + (B*a^8*b^3 + 4*B*a^6*b^5 + 5*B*a^4*b^7 + 4*A*a^3*b^8 + 10*B*a
^2*b^9 - 4*A*a*b^10)*tan(d*x + c)^3 + 3*(B*a^9*b^2 + 4*B*a^7*b^4 + 5*B*a^5
*b^6 + 4*A*a^4*b^7 + 10*B*a^3*b^8 - 4*A*a^2*b^9)*tan(d*x + c)^2 + 3*(B*a^1
0*b + 4*B*a^8*b^3 + 5*B*a^6*b^5 + 4*A*a^5*b^6 + 10*B*a^4*b^7 - 4*A*a^3*b^8
)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d
*x + c)^2 + 1)) - 3*(B*a^11 + 4*B*a^9*b^2 + 6*B*a^7*b^4 + 4*B*a^5*b^6 + B
a^3*b^8 + (B*a^8*b^3 + 4*B*a^6*b^5 + 6*B*a^4*b^7 + 4*B*a^2*b^9 + B*b^11)*t
an(d*x + c)^3 + 3*(B*a^9*b^2 + 4*B*a^7*b^4 + 6*B*a^5*b^6 + 4*B*a^3*b^8 + B
*a*b^10)*tan(d*x + c)^2 + 3*(B*a^10*b + 4*B*a^8*b^3 + 6*B*a^6*b^5 + 4*B*a^
4*b^7 + B*a^2*b^9)*tan(d*x + c))*log(1/(tan(d*x + c)^2 + 1)) - 3*(2*B*a^10
*b + 5*B*a^8*b^3 + 2*A*a^7*b^4 + 12*B*a^6*b^5 - 22*A*a^5*b^6 - 35*B*a^4*b^
7 + 20*A*a^3*b^8 - 6*(A*a^6*b^5 + 4*B*a^5*b^6 - 6*A*a^4*b^7 - 4*B*a^3*b...
```

3.290.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx = \text{Exception raised: AttributeError}$$

```
input integrate(tan(d*x+c)**4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**4,x)
```

```
output Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'
```

3.290.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 583, normalized size of antiderivative = 1.66

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$$

$$= \frac{6(Aa^4+4Ba^3b-6Aa^2b^2-4Bab^3+Ab^4)(dx+c)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} + \frac{6(Ba^8+4Ba^6b^2+5Ba^4b^4+4Aa^3b^5+10Ba^2b^6-4Aab^7)\log(b\tan(dx+c)+a)}{a^8b^4+4a^6b^6+6a^4b^8+4a^2b^{10}+b^{12}} + \frac{3(Ba^4-4Aa^3b+6Aa^2b^2-4Aab^3+Bb^4)}{a^8b^4+4a^6b^6+6a^4b^8+4a^2b^{10}+b^{12}}$$

```
input integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="maxima")
```

```
output 1/6*(6*(A*a^4 + 4*B*a^3*b - 6*A*a^2*b^2 - 4*B*a*b^3 + A*b^4)*(d*x + c)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + 6*(B*a^8 + 4*B*a^6*b^2 + 5*B*a^4*b^4 + 4*A*a^3*b^5 + 10*B*a^2*b^6 - 4*A*a*b^7)*log(b*tan(d*x + c) + a)/(a^8*b^4 + 4*a^6*b^6 + 6*a^4*b^8 + 4*a^2*b^10 + b^12) + 3*(B*a^4 - 4*A*a^3*b - 6*B*a^2*b^2 + 4*A*a*b^3 + B*b^4)*log(tan(d*x + c)^2 + 1)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + (11*B*a^9 - 2*A*a^8*b + 34*B*a^7*b^2 - 4*A*a^6*b^3 + 47*B*a^5*b^4 - 26*A*a^4*b^5 + 6*(3*B*a^7*b^2 - A*a^6*b^3 + 9*B*a^5*b^4 - 3*A*a^4*b^5 + 10*B*a^3*b^6 - 6*A*a^2*b^7)*tan(d*x + c)^2 + 3*(9*B*a^8*b - 2*A*a^7*b^2 + 28*B*a^6*b^3 - 6*A*a^5*b^4 + 35*B*a^4*b^5 - 20*A*a^3*b^6)*tan(d*x + c))/(a^9*b^4 + 3*a^7*b^6 + 3*a^5*b^8 + a^3*b^10 + (a^6*b^7 + 3*a^4*b^9 + 3*a^2*b^11 + b^13)*tan(d*x + c)^3 + 3*(a^7*b^6 + 3*a^5*b^8 + 3*a^3*b^10 + a*b^12)*tan(d*x + c)^2 + 3*(a^8*b^5 + 3*a^6*b^7 + 3*a^4*b^9 + a^2*b^11)*tan(d*x + c))/d
```

3.290.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 719 vs. 2(346) = 692.

Time = 1.26 (sec) , antiderivative size = 719, normalized size of antiderivative = 2.05

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$$

$$= \frac{6(Aa^4+4Ba^3b-6Aa^2b^2-4Bab^3+Ab^4)(dx+c)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} + \frac{3(Ba^4-4Aa^3b-6Ba^2b^2+4Aab^3+Bb^4)\log(\tan(dx+c)^2+1)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} + \frac{6(Ba^8+4Ba^6b^2+5Ba^4b^4+4Aa^3b^5+10Ba^2b^6-4Aab^7)}{a^8b^4+4a^6b^6+6a^4b^8+4a^2b^{10}+b^{12}}$$

```
input integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="giac")
```

$$3.290. \int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$$

output

```

1/6*(6*(A*a^4 + 4*B*a^3*b - 6*A*a^2*b^2 - 4*B*a*b^3 + A*b^4)*(d*x + c)/(a^
8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + 3*(B*a^4 - 4*A*a^3*b - 6*B*
a^2*b^2 + 4*A*a*b^3 + B*b^4)*log(tan(d*x + c)^2 + 1)/(a^8 + 4*a^6*b^2 + 6*
a^4*b^4 + 4*a^2*b^6 + b^8) + 6*(B*a^8 + 4*B*a^6*b^2 + 5*B*a^4*b^4 + 4*A*a^
3*b^5 + 10*B*a^2*b^6 - 4*A*a*b^7)*log(abs(b*tan(d*x + c) + a))/(a^8*b^4 +
4*a^6*b^6 + 6*a^4*b^8 + 4*a^2*b^10 + b^12) - (11*B*a^8*b^2*tan(d*x + c)^3
+ 44*B*a^6*b^4*tan(d*x + c)^3 + 55*B*a^4*b^6*tan(d*x + c)^3 + 44*A*a^3*b^7
*tan(d*x + c)^3 + 110*B*a^2*b^8*tan(d*x + c)^3 - 44*A*a*b^9*tan(d*x + c)^3
+ 15*B*a^9*b*tan(d*x + c)^2 + 6*A*a^8*b^2*tan(d*x + c)^2 + 60*B*a^7*b^3*t
an(d*x + c)^2 + 24*A*a^6*b^4*tan(d*x + c)^2 + 51*B*a^5*b^5*tan(d*x + c)^2
+ 186*A*a^4*b^6*tan(d*x + c)^2 + 270*B*a^3*b^7*tan(d*x + c)^2 - 96*A*a^2*b
^8*tan(d*x + c)^2 + 6*B*a^10*tan(d*x + c) + 6*A*a^9*b*tan(d*x + c) + 21*B*
a^8*b^2*tan(d*x + c) + 24*A*a^7*b^3*tan(d*x + c) - 24*B*a^6*b^4*tan(d*x +
c) + 210*A*a^5*b^5*tan(d*x + c) + 225*B*a^4*b^6*tan(d*x + c) - 72*A*a^3*b^
7*tan(d*x + c) + 2*A*a^10 - B*a^9*b + 6*A*a^8*b^2 - 26*B*a^7*b^3 + 74*A*a^
6*b^4 + 63*B*a^5*b^5 - 18*A*a^4*b^6)/((a^8*b^3 + 4*a^6*b^5 + 6*a^4*b^7 + 4
*a^2*b^9 + b^11)*(b*tan(d*x + c) + a)^3))/d

```

3.290.9 Mupad [B] (verification not implemented)

Time = 8.81 (sec) , antiderivative size = 486, normalized size of antiderivative = 1.38

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

$$= \frac{11 B a^9 - 2 A a^8 b + 34 B a^7 b^2 - 4 A a^6 b^3 + 47 B a^5 b^4 - 26 A a^4 b^5}{6 b^4 (a^6 + 3 a^4 b^2 + 3 a^2 b^4 + b^6)} + \frac{\tan(c+dx)^2 (3 B a^7 - A a^6 b + 9 B a^5 b^2 - 3 A a^4 b^3 + 10 B a^3 b^4 - 6 A a^2 b^5)}{b^2 (a^6 + 3 a^4 b^2 + 3 a^2 b^4 + b^6)} + \frac{d (a^3 + 3 a^2 b \tan(c+dx) + 3 a b^2 \tan(c+dx)^2 + b^3 \tan(c+dx)^3)}{b^4 d (a^2 + b^2)^4}$$

$$+ \frac{\ln(\tan(c+dx) - i) (A + B i)}{2 d (a^4 i - 4 a^3 b - a^2 b^2 6 i + 4 a b^3 + b^4 i)} + \frac{\ln(\tan(c+dx) + i) (B + A i)}{2 d (a^4 - a^3 b 4 i - 6 a^2 b^2 + a b^3 4 i + b^4)}$$

$$+ \frac{a \ln(a + b \tan(c+dx)) (B a^7 + 4 B a^5 b^2 + 5 B a^3 b^4 + 4 A a^2 b^5 + 10 B a b^6 - 4 A b^7)}{b^4 d (a^2 + b^2)^4}$$

input `int((tan(c + d*x)^4*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^4,x)`

output

$$\begin{aligned} & ((11*B*a^9 - 26*A*a^4*b^5 - 4*A*a^6*b^3 + 47*B*a^5*b^4 + 34*B*a^7*b^2 - 2* \\ & A*a^8*b)/(6*b^4*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2)) + (\tan(c + d*x)^2*(3* \\ & B*a^7 - 6*A*a^2*b^5 - 3*A*a^4*b^3 + 10*B*a^3*b^4 + 9*B*a^5*b^2 - A*a^6*b)) \\ & / (b^2*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2)) + (\tan(c + d*x)*(9*B*a^8 - 20*A \\ & *a^3*b^5 - 6*A*a^5*b^3 + 35*B*a^4*b^4 + 28*B*a^6*b^2 - 2*A*a^7*b))/(2*b^3* \\ & (a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2)))/(d*(a^3 + b^3*\tan(c + d*x)^3 + 3*a*b \\ & ^2*\tan(c + d*x)^2 + 3*a^2*b*\tan(c + d*x))) + (\log(\tan(c + d*x) - 1i)*(A + \\ & B*1i))/(2*d*(4*a*b^3 - 4*a^3*b + a^4*1i + b^4*1i - a^2*b^2*6i)) + (\log(\tan \\ & (c + d*x) + 1i)*(A*1i + B))/(2*d*(a*b^3*4i - a^3*b*4i + a^4 + b^4 - 6*a^2* \\ & b^2)) + (a*\log(a + b*\tan(c + d*x))*(B*a^7 - 4*A*b^7 + 4*A*a^2*b^5 + 5*B*a^ \\ & 3*b^4 + 4*B*a^5*b^2 + 10*B*a*b^6))/(b^4*d*(a^2 + b^2)^4) \end{aligned}$$

3.291
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

3.291.1 Optimal result 2799
 3.291.2 Mathematica [C] (verified) 2800
 3.291.3 Rubi [A] (verified) 2800
 3.291.4 Maple [A] (verified) 2805
 3.291.5 Fricas [B] (verification not implemented) 2805
 3.291.6 Sympy [F(-2)] 2806
 3.291.7 Maxima [A] (verification not implemented) 2807
 3.291.8 Giac [B] (verification not implemented) 2807
 3.291.9 Mupad [B] (verification not implemented) 2808

3.291.1 Optimal result

Integrand size = 31, antiderivative size = 298

$$\begin{aligned} & \int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx \\ &= -\frac{(4a^3Ab - 4aAb^3 - a^4B + 6a^2b^2B - b^4B)x}{(a^2+b^2)^4} \\ & \quad + \frac{(a^4A - 6a^2Ab^2 + Ab^4 + 4a^3bB - 4ab^3B) \log(a \cos(c+dx) + b \sin(c+dx))}{(a^2+b^2)^4 d} \\ & \quad + \frac{a(Ab - aB) \tan^2(c+dx)}{3b(a^2+b^2)d(a+b \tan(c+dx))^3} + \frac{a^2(a^2Ab - 5Ab^3 + 2a^3B + 8ab^2B)}{6b^3(a^2+b^2)^2 d(a+b \tan(c+dx))^2} \\ & \quad - \frac{a(a^4Ab + 5a^2Ab^3 - 8Ab^5 + 2a^5B + 7a^3b^2B + 17ab^4B)}{3b^3(a^2+b^2)^3 d(a+b \tan(c+dx))} \end{aligned}$$

output

```
- (4*A*a^3*b-4*A*a*b^3-B*a^4+6*B*a^2*b^2-B*b^4)*x/(a^2+b^2)^4+(A*a^4-6*A*a^2*b^2+A*b^4+4*B*a^3*b-4*B*a*b^3)*ln(a*cos(d*x+c)+b*sin(d*x+c))/(a^2+b^2)^4/d+1/3*a*(A*b-B*a)*tan(d*x+c)^2/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^3+1/6*a^2*(A*a^2*b-5*A*b^3+2*B*a^3+8*B*a*b^2)/b^3/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^2-1/3*a*(A*a^4*b+5*A*a^2*b^3-8*A*b^5+2*B*a^5+7*B*a^3*b^2+17*B*a*b^4)/b^3/(a^2+b^2)^3/d/(a+b*tan(d*x+c))
```

3.291.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.37 (sec) , antiderivative size = 465, normalized size of antiderivative = 1.56

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx = -\frac{B\tan^2(c+dx)}{bd(a+b\tan(c+dx))^3} + \frac{(6aAb^3+6b^4B)\left(-\frac{i\log(i-\tan(c+dx))}{2(a+ib)^4} + \frac{i\log(i+\tan(c+dx))}{2(a-ib)^4} + \frac{4a(a-b)b(a+b)\log(a+b\tan(c+dx))}{(a^2+b^2)^4}\right)}{b} - \frac{(-Ab-2aB)\tan(c+dx)}{2bd(a+b\tan(c+dx))^3} - \frac{aAb+2a^2B-2b^2B}{3bd(a+b\tan(c+dx))^3} + \dots$$

input `Integrate[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]`

output `-(B*Tan[c + d*x]^2)/(b*d*(a + b*Tan[c + d*x])^3) - (-1/2*((-(A*b) - 2*a*B)*Tan[c + d*x])/(b*d*(a + b*Tan[c + d*x])^3) - (-1/3*(a*A*b + 2*a^2*B - 2*b^2*B)/(b*d*(a + b*Tan[c + d*x])^3) + (((6*a*A*b^3 + 6*b^4*B)*((-1/2*I)*Log[I - Tan[c + d*x]])/(a + I*b)^4 + ((I/2)*Log[I + Tan[c + d*x]])/(a - I*b)^4 + (4*a*(a - b)*b*(a + b)*Log[a + b*Tan[c + d*x]])/(a^2 + b^2)^4 - b/(3*(a^2 + b^2)*(a + b*Tan[c + d*x])^3) - (a*b)/((a^2 + b^2)^2*(a + b*Tan[c + d*x])^2) - (b*(3*a^2 - b^2))/((a^2 + b^2)^3*(a + b*Tan[c + d*x])))/b - 6*A*b^2*(-1/2*Log[I - Tan[c + d*x]])/(I*a - b)^3 + Log[I + Tan[c + d*x]]/(2*(I*a + b)^3) + (b*(3*a^2 - b^2)*Log[a + b*Tan[c + d*x]])/(a^2 + b^2)^3 - b/(2*(a^2 + b^2)*(a + b*Tan[c + d*x])^2) - (2*a*b)/((a^2 + b^2)^2*(a + b*Tan[c + d*x])))/(3*b*d)/(2*b))/b`

3.291.3 Rubi [A] (verified)

Time = 1.55 (sec) , antiderivative size = 346, normalized size of antiderivative = 1.16, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 4088, 25, 3042, 4118, 25, 3042, 4111, 27, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx \quad \downarrow \quad 3042$$

3.291. $\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$

$$\begin{aligned}
 & \int \frac{\tan(c+dx)^3(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx \\
 & \quad \downarrow 4088 \\
 & \int \frac{\tan(c+dx) \left(-((2Ba^2+Aba+3b^2B) \tan^2(c+dx)) - 3b(Ab-aB) \tan(c+dx) + 2a(Ab-aB) \right)}{(a+b \tan(c+dx))^3} dx + \\
 & \quad \frac{3b(a^2+b^2)}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} + \\
 & \quad \frac{a(Ab-aB) \tan^2(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} \\
 & \quad \downarrow 25 \\
 & \int \frac{\tan(c+dx) \left(-((2Ba^2+Aba+3b^2B) \tan^2(c+dx)) - 3b(Ab-aB) \tan(c+dx) + 2a(Ab-aB) \right)}{(a+b \tan(c+dx))^3} dx \\
 & \quad \frac{3b(a^2+b^2)}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} - \\
 & \quad \frac{a(Ab-aB) \tan^2(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} - \\
 & \quad \downarrow 3042 \\
 & \int \frac{\tan(c+dx) \left(-((2Ba^2+Aba+3b^2B) \tan^2(c+dx)) - 3b(Ab-aB) \tan(c+dx) + 2a(Ab-aB) \right)}{(a+b \tan(c+dx))^3} dx \\
 & \quad \frac{3b(a^2+b^2)}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} - \\
 & \quad \frac{a(Ab-aB) \tan^2(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} - \\
 & \quad \downarrow 4118 \\
 & \int \frac{-3(Aa^2+2bBa-Ab^2) \tan(c+dx)b^2 + (a^2+b^2) \left((2Ba^2+Aba+3b^2B) \tan^2(c+dx) + a(2Ba^3+Aba^2+8b^2Ba-5Ab^3) \right)}{(a+b \tan(c+dx))^2} dx - \frac{a^2(2a^3B+a^2Ab+8ab^2B-5Ab^3)}{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \frac{3b(a^2+b^2)}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} - \\
 & \quad \downarrow 25 \\
 & \int \frac{-3(Aa^2+2bBa-Ab^2) \tan(c+dx)b^2 + (a^2+b^2) \left((2Ba^2+Aba+3b^2B) \tan^2(c+dx) + a(2Ba^3+Aba^2+8b^2Ba-5Ab^3) \right)}{(a+b \tan(c+dx))^2} dx - \frac{a^2(2a^3B+a^2Ab+8ab^2B-5Ab^3)}{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \frac{3b(a^2+b^2)}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} - \\
 & \quad \downarrow 3042 \\
 & \int \frac{-3(Aa^2+2bBa-Ab^2) \tan(c+dx)b^2 + (a^2+b^2) \left((2Ba^2+Aba+3b^2B) \tan^2(c+dx) + a(2Ba^3+Aba^2+8b^2Ba-5Ab^3) \right)}{(a+b \tan(c+dx))^2} dx - \frac{a^2(2a^3B+a^2Ab+8ab^2B-5Ab^3)}{2b^2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \frac{3b(a^2+b^2)}{3bd(a^2+b^2)(a+b \tan(c+dx))^3} - \\
 & \quad \downarrow 4111
 \end{aligned}$$

3.291. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

$$\frac{\int \frac{a(Ab - aB) \tan^2(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} dx - \frac{3 \left((-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) b^2 + (Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) \tan(c + dx) b^2 \right)}{a^2 + b^2} dx - \frac{a(2a^5B + a^4Ab + 7a^3b^2B + 5a^2Ab^3 + 17ab^4B - 8Ab^5)}{bd(a^2 + b^2)(a + b \tan(c + dx))}}{b(a^2 + b^2)} - \frac{a^2(2a^3B + a^2b^2)}{2b^2d(a^2 + b^2)}}{3b(a^2 + b^2)}$$

27

$$\frac{\int \frac{a(Ab - aB) \tan^2(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} dx - \frac{3 \left((-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) b^2 + (Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) \tan(c + dx) b^2 \right)}{a^2 + b^2} dx - \frac{a(2a^5B + a^4Ab + 7a^3b^2B + 5a^2Ab^3 + 17ab^4B - 8Ab^5)}{bd(a^2 + b^2)(a + b \tan(c + dx))}}{b(a^2 + b^2)} - \frac{a^2(2a^3B + a^2b^2)}{2b^2d(a^2 + b^2)}}{3b(a^2 + b^2)}$$

3042

$$\frac{\int \frac{a(Ab - aB) \tan^2(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} dx - \frac{3 \left((-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) b^2 + (Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) \tan(c + dx) b^2 \right)}{a^2 + b^2} dx - \frac{a(2a^5B + a^4Ab + 7a^3b^2B + 5a^2Ab^3 + 17ab^4B - 8Ab^5)}{bd(a^2 + b^2)(a + b \tan(c + dx))}}{b(a^2 + b^2)} - \frac{a^2(2a^3B + a^2b^2)}{2b^2d(a^2 + b^2)}}{3b(a^2 + b^2)}$$

4014

$$\frac{\int \frac{a(Ab - aB) \tan^2(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} dx - \frac{3 \left(\frac{b^2x(a^4(-B) + 4a^3Ab + 6a^2b^2B - 4aAb^3 - b^4B)}{a^2 + b^2} - \frac{b^2(a^4A + 4a^3bB - 6a^2Ab^2 - 4ab^3B + Ab^4)}{a^2 + b^2} \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx \right)}{a^2 + b^2} dx - \frac{a(2a^5B + a^4Ab + 7a^3b^2B + 5a^2Ab^3 + 17ab^4B - 8Ab^5)}{bd(a^2 + b^2)(a + b \tan(c + dx))}}{b(a^2 + b^2)} - \frac{a^2(2a^3B + a^2b^2)}{2b^2d(a^2 + b^2)}}{3b(a^2 + b^2)}$$

3042

$$\frac{\int \frac{a(Ab - aB) \tan^2(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} dx - \frac{3 \left(\frac{b^2x(a^4(-B) + 4a^3Ab + 6a^2b^2B - 4aAb^3 - b^4B)}{a^2 + b^2} - \frac{b^2(a^4A + 4a^3bB - 6a^2Ab^2 - 4ab^3B + Ab^4)}{a^2 + b^2} \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx \right)}{a^2 + b^2} dx - \frac{a(2a^5B + a^4Ab + 7a^3b^2B + 5a^2Ab^3 + 17ab^4B - 8Ab^5)}{bd(a^2 + b^2)(a + b \tan(c + dx))}}{b(a^2 + b^2)} - \frac{a^2(2a^3B + a^2b^2)}{2b^2d(a^2 + b^2)}}{3b(a^2 + b^2)}$$

4013

3.291. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

$$\frac{a^2(2a^3B+a^2Ab+8ab^2B-5Ab^3)}{2b^2d(a^2+b^2)(a+b\tan(c+dx))^2} - \frac{a(Ab-aB)\tan^2(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^3} - \frac{b^2(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4)\log(a\cos(c+dx)+b\sin(c+dx))}{d(a^2+b^2)} - \frac{b^2(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4)}{a^2+b^2}$$

$$\frac{3b(a^2+b^2)}{b(a^2+b^2)}$$

```
input Int[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]
```

```
output (a*(A*b - a*B)*Tan[c + d*x]^2)/(3*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^3 - (-1/2*(a^2*(a^2*A*b - 5*A*b^3 + 2*a^3*B + 8*a*b^2*B))/(b^2*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) - ((-3*((b^2*(4*a^3*A*b - 4*a*A*b^3 - a^4*B + 6*a^2*b^2*B - b^4*B)*x)/(a^2 + b^2) - (b^2*(a^4*A - 6*a^2*A*b^2 + A*b^4 + 4*a^3*b*B - 4*a*b^3*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a^2 + b^2)*d))/(a^2 + b^2) - (a*(a^4*A*b + 5*a^2*A*b^3 - 8*A*b^5 + 2*a^5*B + 7*a^3*b^2*B + 17*a*b^4*B))/(b*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(b*(a^2 + b^2)))/(3*b*(a^2 + b^2))
```

3.291.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4013 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

rule 4014 $\text{Int}[\frac{(c + d)\tan(e + f x)}{(a + b)\tan(e + f x) + (f x)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(a c + b d)(x/(a^2 + b^2)), x] + \text{Simp}[(b c - a d)/(a^2 + b^2) \text{Int}[(b - a \tan[e + f x])/(a + b \tan[e + f x]), x], x] /;$
 $\text{FreeQ}\{a, b, c, d, e, f\}, x\} \ \&\& \ \text{NeQ}[b c - a d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[a c + b d, 0]$

rule 4088 $\text{Int}[\frac{(a + b)\tan(e + f x)^m (A + B)\tan(e + f x) + (f x)^n (c + d)\tan(e + f x)}{(b c - a d)(B c - A d)(a + b \tan[e + f x])^{m-1} (c + d \tan[e + f x])^{n+1} (d^2 f^2 (n+1)(c^2 + d^2))}, x] - \text{Simp}[1/(d(n+1)(c^2 + d^2)) \text{Int}[(a + b \tan[e + f x])^{m-2} (c + d \tan[e + f x])^{n+1} \text{Simp}[a A d (b d (m-1) - a c (n+1)) + (b B c - (A b + a B) d)(b c (m-1) + a d (n+1)) - d((a A - b B)(b c - a d) + (A b + a B)(a c + b d))(n+1) \tan[e + f x] - b(d(A b c + a B c - a A d)(m+n) - b B (c^2 (m-1) - d^2 (n+1))) \tan[e + f x]^2, x], x], x] /;$
 $\text{FreeQ}\{a, b, c, d, e, f, A, B\}, x\} \ \&\& \ \text{NeQ}[b c - a d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ (\text{IntegerQ}[m] \ || \ \text{IntegersQ}[2 m, 2 n])$

rule 4111 $\text{Int}[\frac{(a + b)\tan(e + f x)^m (A + B)\tan(e + f x) + (f x)^n (C + D)\tan(e + f x)^2}{(A b^2 - a b B + a^2 C)(a + b \tan[e + f x])^{m+1} (b f (m+1)(a^2 + b^2))}, x] + \text{Simp}[1/(a^2 + b^2) \text{Int}[(a + b \tan[e + f x])^{m+1} \text{Simp}[b B + a(A - C) - (A b - a B - b C) \tan[e + f x], x], x], x] /;$
 $\text{FreeQ}\{a, b, e, f, A, B, C\}, x\} \ \&\& \ \text{NeQ}[A b^2 - a b B + a^2 C, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$

rule 4118 $\text{Int}[\frac{(a + b)\tan(e + f x)^n (A + B)\tan(e + f x) + (C + D)\tan(e + f x)^2}{(b c - a d)(c^2 C - B c d + A d^2)(c + d \tan[e + f x])^{n+1} (d^2 f^2 (n+1)(c^2 + d^2))}, x] + \text{Simp}[1/(d(c^2 + d^2)) \text{Int}[(c + d \tan[e + f x])^{n+1} \text{Simp}[a d (A c - c C + B d) + b(c^2 C - B c d + A d^2) + d(A b c + a B c - b c C - a A d + b B d + a C d) \tan[e + f x] + b C (c^2 + d^2) \tan[e + f x]^2, x], x], x] /;$
 $\text{FreeQ}\{a, b, c, d, e, f, A, B, C\}, x\} \ \&\& \ \text{NeQ}[b c - a d, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{LtQ}[n, -1]$

3.291.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 318, normalized size of antiderivative = 1.07

method	result
derivativedivides	$\frac{(-A a^4 + 6A a^2 b^2 - A b^4 - 4B a^3 b + 4B a b^3) \ln(1 + \tan^2(dx+c)) + (-4A a^3 b + 4A a b^3 + B a^4 - 6B a^2 b^2 + B b^4) \arctan(\tan(dx+c))}{(a^2 + b^2)^4} + \frac{A a^6 b + 8A a^4 b^3 - 9A a^2 b^5 + 2B a^7 + 6B a^5 b^2 + 20B a^3 b^4}{6a^2 d(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)} (\tan^3(dx+c)) + \frac{(A a^6 + 6A a^4 b^2 - 3A a^2 b^4 + 8B a^3 b^3) (\tan^2(dx+c))}{2ad(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)} - \frac{a(3A a^5 b + 6A a^3 b^3 - 3A a b^5 + 4B a^4 b^2 + 4B a^2 b^4)}{3(a^2 + b^2)^4}$
default	$\frac{(-A a^4 + 6A a^2 b^2 - A b^4 - 4B a^3 b + 4B a b^3) \ln(1 + \tan^2(dx+c)) + (-4A a^3 b + 4A a b^3 + B a^4 - 6B a^2 b^2 + B b^4) \arctan(\tan(dx+c))}{(a^2 + b^2)^4} + \frac{A a^6 b + 8A a^4 b^3 - 9A a^2 b^5 + 2B a^7 + 6B a^5 b^2 + 20B a^3 b^4}{6a^2 d(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)} (\tan^3(dx+c)) + \frac{(A a^6 + 6A a^4 b^2 - 3A a^2 b^4 + 8B a^3 b^3) (\tan^2(dx+c))}{2ad(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)} - \frac{a(3A a^5 b + 6A a^3 b^3 - 3A a b^5 + 4B a^4 b^2 + 4B a^2 b^4)}{3(a^2 + b^2)^4}$
norman	$\frac{A a^6 b + 8A a^4 b^3 - 9A a^2 b^5 + 2B a^7 + 6B a^5 b^2 + 20B a^3 b^4}{6a^2 d(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)} (\tan^3(dx+c)) + \frac{(A a^6 + 6A a^4 b^2 - 3A a^2 b^4 + 8B a^3 b^3) (\tan^2(dx+c))}{2ad(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)} - \frac{a(3A a^5 b + 6A a^3 b^3 - 3A a b^5 + 4B a^4 b^2 + 4B a^2 b^4)}{3(a^2 + b^2)^4}$
risch	Expression too large to display
parallelrisch	Expression too large to display

input `int (tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} \left(\frac{1}{(a^2 + b^2)^4} \left(\frac{1}{2} (-A a^4 + 6A a^2 b^2 - A b^4 - 4B a^3 b + 4B a b^3) \ln(1 + \tan^2(dx+c)) + (-4A a^3 b + 4A a b^3 + B a^4 - 6B a^2 b^2 + B b^4) \arctan(\tan(dx+c)) \right) + \frac{A a^6 b + 8A a^4 b^3 - 9A a^2 b^5 + 2B a^7 + 6B a^5 b^2 + 20B a^3 b^4}{6a^2 d(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)} (\tan^3(dx+c)) - \frac{a(3A a^5 b + 6A a^3 b^3 - 3A a b^5 + 4B a^4 b^2 + 4B a^2 b^4)}{3(a^2 + b^2)^4} \right) - \frac{1}{2} \frac{a^2 (A a^2 b^3 - 3A a b^5 + B a^5 + 3B a^3 b^2 + 6B a^2 b^4)}{(a^2 + b^2)^3} - \frac{b^3}{(a+b \tan(dx+c))} - \frac{1}{2} \frac{a^2 (A a^2 b^3 + 3A a b^5 - 2B a^3 b^2 - 4B a^2 b^4)}{b^3 (a^2 + b^2)^2} - \frac{1}{3} \frac{a^3 (A b - B a)}{b^3 (a^2 + b^2)} - \frac{a}{(a+b \tan(dx+c))^3}$$

3.291.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 813 vs. 2(289) = 578.

Time = 0.29 (sec) , antiderivative size = 813, normalized size of antiderivative = 2.73

$$\int \frac{\tan^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^4} dx$$

$$= \frac{3 A a^7 + 18 B a^6 b - 30 A a^5 b^2 - 26 B a^4 b^3 + 11 A a^3 b^4 + (2 B a^7 + A a^6 b + 6 B a^5 b^2 + 18 A a^4 b^3 + 48 B a^3 b^4 - 30 A a^2 b^4 - 26 B a b^5 + 11 A b^6 + 6 B a^7 + 18 B a^6 b - 30 A a^5 b^2 - 26 B a^4 b^3 + 11 A a^3 b^4 + 2 B a^7 + A a^6 b + 6 B a^5 b^2 + 18 A a^4 b^3 + 48 B a^3 b^4 - 30 A a^2 b^4 - 26 B a b^5 + 11 A b^6)}{3 a^2 d (a^6 + 3 a^4 b^2 + 3 a^2 b^4 + b^6)} (\tan^3(dx+c)) + \frac{(A a^6 + 6 A a^4 b^2 - 3 A a^2 b^4 + 8 B a^3 b^3) (\tan^2(dx+c))}{2 a d (a^6 + 3 a^4 b^2 + 3 a^2 b^4 + b^6)} - \frac{a (3 A a^5 b + 6 A a^3 b^3 - 3 A a b^5 + 4 B a^4 b^2 + 4 B a^2 b^4)}{3 (a^2 + b^2)^4}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="fricas")`

3.291.
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

```
output 1/6*(3*A*a^7 + 18*B*a^6*b - 30*A*a^5*b^2 - 26*B*a^4*b^3 + 11*A*a^3*b^4 + (
2*B*a^7 + A*a^6*b + 6*B*a^5*b^2 + 18*A*a^4*b^3 + 48*B*a^3*b^4 - 27*A*a^2*b
^5 + 6*(B*a^4*b^3 - 4*A*a^3*b^4 - 6*B*a^2*b^5 + 4*A*a*b^6 + B*b^7)*d*x)*ta
n(d*x + c)^3 + 6*(B*a^7 - 4*A*a^6*b - 6*B*a^5*b^2 + 4*A*a^4*b^3 + B*a^3*b^
4)*d*x + 3*(A*a^7 - 2*B*a^6*b + 16*A*a^5*b^2 + 30*B*a^4*b^3 - 23*A*a^3*b^4
- 12*B*a^2*b^5 + 6*A*a*b^6 + 6*(B*a^5*b^2 - 4*A*a^4*b^3 - 6*B*a^3*b^4 + 4
*A*a^2*b^5 + B*a*b^6)*d*x)*tan(d*x + c)^2 + 3*(A*a^7 + 4*B*a^6*b - 6*A*a^5
*b^2 - 4*B*a^4*b^3 + A*a^3*b^4 + (A*a^4*b^3 + 4*B*a^3*b^4 - 6*A*a^2*b^5 -
4*B*a*b^6 + A*b^7)*tan(d*x + c)^3 + 3*(A*a^5*b^2 + 4*B*a^4*b^3 - 6*A*a^3*b
^4 - 4*B*a^2*b^5 + A*a*b^6)*tan(d*x + c)^2 + 3*(A*a^6*b + 4*B*a^5*b^2 - 6*
A*a^4*b^3 - 4*B*a^3*b^4 + A*a^2*b^5)*tan(d*x + c))*log((b^2*tan(d*x + c)^2
+ 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - 3*(2*B*a^7 - 9*A*a^6*
b - 22*B*a^5*b^2 + 26*A*a^4*b^3 + 20*B*a^3*b^4 - 9*A*a^2*b^5 - 6*(B*a^6*b
- 4*A*a^5*b^2 - 6*B*a^4*b^3 + 4*A*a^3*b^4 + B*a^2*b^5)*d*x)*tan(d*x + c))/
((a^8*b^3 + 4*a^6*b^5 + 6*a^4*b^7 + 4*a^2*b^9 + b^11)*d*tan(d*x + c)^3 + 3
*(a^9*b^2 + 4*a^7*b^4 + 6*a^5*b^6 + 4*a^3*b^8 + a*b^10)*d*tan(d*x + c)^2 +
3*(a^10*b + 4*a^8*b^3 + 6*a^6*b^5 + 4*a^4*b^7 + a^2*b^9)*d*tan(d*x + c) +
(a^11 + 4*a^9*b^2 + 6*a^7*b^4 + 4*a^5*b^6 + a^3*b^8)*d)
```

3.291.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx = \text{Exception raised: AttributeError}$$

```
input integrate(tan(d*x+c)**3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**4,x)
```

```
output Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'
```

3.291.7 Maxima [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 550, normalized size of antiderivative = 1.85

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$$

$$= \frac{6(Ba^4-4Aa^3b-6Ba^2b^2+4Aab^3+Bb^4)(dx+c)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} + \frac{6(Aa^4+4Ba^3b-6Aa^2b^2-4Bab^3+Ab^4)\log(b\tan(dx+c)+a)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} - \frac{3(Aa^4+4Ba^3b-6Aa^2b^2-4Bab^3+Ab^4)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="maxima")
```

```
output 1/6*(6*(B*a^4 - 4*A*a^3*b - 6*B*a^2*b^2 + 4*A*a*b^3 + B*b^4)*(d*x + c)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + 6*(A*a^4 + 4*B*a^3*b - 6*A*a^2*b^2 - 4*B*a*b^3 + A*b^4)*log(b*tan(d*x + c) + a)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) - 3*(A*a^4 + 4*B*a^3*b - 6*A*a^2*b^2 - 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)^2 + 1)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) - (2*B*a^8 + A*a^7*b + 4*B*a^6*b^2 + 14*A*a^5*b^3 + 26*B*a^4*b^4 - 11*A*a^3*b^5 + 6*(B*a^6*b^2 + 3*B*a^4*b^4 + A*a^3*b^5 + 6*B*a^2*b^6 - 3*A*a*b^7)*tan(d*x + c)^2 + 3*(2*B*a^7*b + A*a^6*b^2 + 6*B*a^5*b^3 + 8*A*a^4*b^4 + 20*B*a^3*b^5 - 9*A*a^2*b^6)*tan(d*x + c))/(a^9*b^3 + 3*a^7*b^5 + 3*a^5*b^7 + a^3*b^9 + (a^6*b^6 + 3*a^4*b^8 + 3*a^2*b^10 + b^12)*tan(d*x + c)^3 + 3*(a^7*b^5 + 3*a^5*b^7 + 3*a^3*b^9 + a*b^11)*tan(d*x + c)^2 + 3*(a^8*b^4 + 3*a^6*b^6 + 3*a^4*b^8 + a^2*b^10)*tan(d*x + c)))/d
```

3.291.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 670 vs. 2(289) = 578.

Time = 0.98 (sec) , antiderivative size = 670, normalized size of antiderivative = 2.25

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$$

$$= \frac{6(Ba^4-4Aa^3b-6Ba^2b^2+4Aab^3+Bb^4)(dx+c)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} - \frac{3(Aa^4+4Ba^3b-6Aa^2b^2-4Bab^3+Ab^4)\log(\tan(dx+c)^2+1)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} + \frac{6(Aa^4b+4Ba^3b^2-6Aa^2b^3-4Bab^4+Ab^5)}{a^8b+4a^6b^2+6a^4b^3+4a^2b^4+b^5}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="giac")
```

output

```

1/6*(6*(B*a^4 - 4*A*a^3*b - 6*B*a^2*b^2 + 4*A*a*b^3 + B*b^4)*(d*x + c)/(a^
8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) - 3*(A*a^4 + 4*B*a^3*b - 6*A*
a^2*b^2 - 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)^2 + 1)/(a^8 + 4*a^6*b^2 + 6*
a^4*b^4 + 4*a^2*b^6 + b^8) + 6*(A*a^4*b + 4*B*a^3*b^2 - 6*A*a^2*b^3 - 4*B*
a*b^4 + A*b^5)*log(abs(b*tan(d*x + c) + a))/(a^8*b + 4*a^6*b^3 + 6*a^4*b^5
+ 4*a^2*b^7 + b^9) - (11*A*a^4*b^6*tan(d*x + c)^3 + 44*B*a^3*b^7*tan(d*x
+ c)^3 - 66*A*a^2*b^8*tan(d*x + c)^3 - 44*B*a*b^9*tan(d*x + c)^3 + 11*A*b^
10*tan(d*x + c)^3 + 6*B*a^8*b^2*tan(d*x + c)^2 + 24*B*a^6*b^4*tan(d*x + c)
^2 + 39*A*a^5*b^5*tan(d*x + c)^2 + 186*B*a^4*b^6*tan(d*x + c)^2 - 210*A*a^
3*b^7*tan(d*x + c)^2 - 96*B*a^2*b^8*tan(d*x + c)^2 + 15*A*a*b^9*tan(d*x +
c)^2 + 6*B*a^9*b*tan(d*x + c) + 3*A*a^8*b^2*tan(d*x + c) + 24*B*a^7*b^3*ta
n(d*x + c) + 60*A*a^6*b^4*tan(d*x + c) + 210*B*a^5*b^5*tan(d*x + c) - 201*
A*a^4*b^6*tan(d*x + c) - 72*B*a^3*b^7*tan(d*x + c) + 6*A*a^2*b^8*tan(d*x +
c) + 2*B*a^10 + A*a^9*b + 6*B*a^8*b^2 + 26*A*a^7*b^3 + 74*B*a^6*b^4 - 63*
A*a^5*b^5 - 18*B*a^4*b^6)/((a^8*b^3 + 4*a^6*b^5 + 6*a^4*b^7 + 4*a^2*b^9 +
b^11)*(b*tan(d*x + c) + a)^3))/d

```

3.291.9 Mupad [B] (verification not implemented)

Time = 9.10 (sec) , antiderivative size = 470, normalized size of antiderivative = 1.58

$$\begin{aligned}
& \int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx \\
&= \frac{\ln(a+b\tan(c+dx)) \left(\frac{A}{(a^2+b^2)^2} - \frac{4b(2Ab-Ba)}{(a^2+b^2)^3} + \frac{8b^3(Ab-Ba)}{(a^2+b^2)^4} \right)}{d} \\
&\quad - \frac{a^2(2Ba^6+Aa^5b+4Ba^4b^2+14Aa^3b^3+26Ba^2b^4-11Aab^5)}{6b^3(a^6+3a^4b^2+3a^2b^4+b^6)} + \frac{\tan(c+dx)^2(Ba^6+3Ba^4b^2+Aa^3b^3+6Ba^2b^4-3Aab^5)}{b(a^6+3a^4b^2+3a^2b^4+b^6)} + \frac{a\tan(c+dx)}{b^3} \\
&\quad - \frac{\ln(\tan(c+dx)+1i)(B+Ali)}{2d(a^41i+4a^3b-a^2b^26i-4ab^3+b^41i)} - \frac{\ln(\tan(c+dx)-i)(A+B1i)}{2d(a^4+a^3b4i-6a^2b^2-ab^34i+b^4)}
\end{aligned}$$

input `int((tan(c + d*x))^3*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^4,x)`

output

```
(log(a + b*tan(c + d*x))*(A/(a^2 + b^2)^2 - (4*b*(2*A*b - B*a))/(a^2 + b^2)^3 + (8*b^3*(A*b - B*a))/(a^2 + b^2)^4))/d - ((a^2*(2*B*a^6 + 14*A*a^3*b^3 + 26*B*a^2*b^4 + 4*B*a^4*b^2 - 11*A*a*b^5 + A*a^5*b))/(6*b^3*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2)) + (tan(c + d*x)^2*(B*a^6 + A*a^3*b^3 + 6*B*a^2*b^4 + 3*B*a^4*b^2 - 3*A*a*b^5))/(b*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2)) + (a*tan(c + d*x)*(2*B*a^6 + 8*A*a^3*b^3 + 20*B*a^2*b^4 + 6*B*a^4*b^2 - 9*A*a*b^5 + A*a^5*b))/(2*b^2*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2)))/(d*(a^3 + b^3*tan(c + d*x)^3 + 3*a*b^2*tan(c + d*x)^2 + 3*a^2*b*tan(c + d*x))) - (log(tan(c + d*x) + 1i)*(A*1i + B))/(2*d*(4*a^3*b - 4*a*b^3 + a^4*1i + b^4*1i - a^2*b^2*6i)) - (log(tan(c + d*x) - 1i)*(A + B*1i))/(2*d*(a^3*b*4i - a*b^3*4i + a^4 + b^4 - 6*a^2*b^2))
```


3.292 $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

3.292.1 Optimal result 2810
 3.292.2 Mathematica [C] (verified) 2811
 3.292.3 Rubi [A] (verified) 2811
 3.292.4 Maple [A] (verified) 2815
 3.292.5 Fricas [B] (verification not implemented) 2816
 3.292.6 Sympy [F(-2)] 2817
 3.292.7 Maxima [B] (verification not implemented) 2817
 3.292.8 Giac [B] (verification not implemented) 2818
 3.292.9 Mupad [B] (verification not implemented) 2819

3.292.1 Optimal result

Integrand size = 31, antiderivative size = 261

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

$$= -\frac{(a^4A - 6a^2Ab^2 + Ab^4 + 4a^3bB - 4ab^3B)x}{(a^2 + b^2)^4}$$

$$- \frac{(4a^3Ab - 4aAb^3 - a^4B + 6a^2b^2B - b^4B) \log(a \cos(c+dx) + b \sin(c+dx))}{(a^2 + b^2)^4 d}$$

$$- \frac{a^2(Ab - aB)}{3b^2(a^2 + b^2)d(a+b \tan(c+dx))^3}$$

$$+ \frac{a(2Ab^3 - a(a^2 + 3b^2)B)}{2b^2(a^2 + b^2)^2 d(a+b \tan(c+dx))^2} + \frac{3a^2Ab - Ab^3 - a^3B + 3ab^2B}{(a^2 + b^2)^3 d(a+b \tan(c+dx))}$$

```
output - (A*a^4-6*A*a^2*b^2+A*b^4+4*B*a^3*b-4*B*a*b^3)*x/(a^2+b^2)^4-(4*A*a^3*b-4*
A*a*b^3-B*a^4+6*B*a^2*b^2-B*b^4)*ln(a*cos(d*x+c)+b*sin(d*x+c))/(a^2+b^2)^4
/d-1/3*a^2*(A*b-B*a)/b^2/(a^2+b^2)/d/(a+b*tan(d*x+c))^3+1/2*a*(2*A*b^3-a*(
a^2+3*b^2)*B)/b^2/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^2+(3*A*a^2*b-A*b^3-B*a^3+
3*B*a*b^2)/(a^2+b^2)^3/d/(a+b*tan(d*x+c))
```

3.292.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.32 (sec) , antiderivative size = 411, normalized size of antiderivative = 1.57

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^4} dx = -\frac{B \tan(c + dx)}{2bd(a + b \tan(c + dx))^3} + \frac{(6Ab^3 - 6ab^2B) \left(-\frac{i \log(i - \tan(c + dx))}{2(a + ib)^4} + \frac{i \log(i + \tan(c + dx))}{2(a - ib)^4} + \frac{4a(a - b)b(a + b) \log(a + b \tan(c + dx))}{(a^2 + b^2)^4} - \frac{b}{3(a^2 + b^2)(a + b \tan(c + dx))^3} \right) + \frac{2Ab + aB}{3bd(a + b \tan(c + dx))^3} + \dots$$

input `Integrate[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]`

output `-1/2*(B*Tan[c + d*x])/(b*d*(a + b*Tan[c + d*x])^3) - ((2*A*b + a*B)/(3*b*d*(a + b*Tan[c + d*x])^3) + (((6*A*b^3 - 6*a*b^2*B)*((-1/2*I)*Log[I - Tan[c + d*x]])/(a + I*b)^4 + ((I/2)*Log[I + Tan[c + d*x]])/(a - I*b)^4 + (4*a*(a - b)*b*(a + b)*Log[a + b*Tan[c + d*x]]/(a^2 + b^2)^4 - b/(3*(a^2 + b^2)*(a + b*Tan[c + d*x])^3) - (a*b)/((a^2 + b^2)^2*(a + b*Tan[c + d*x])) - (b*(3*a^2 - b^2))/((a^2 + b^2)^3*(a + b*Tan[c + d*x]))) / b + 6*b*B*(-1/2*Log[I - Tan[c + d*x]]/(I*a - b)^3 + Log[I + Tan[c + d*x]]/(2*(I*a + b)^3) + (b*(3*a^2 - b^2)*Log[a + b*Tan[c + d*x]]/(a^2 + b^2)^3 - b/(2*(a^2 + b^2)*(a + b*Tan[c + d*x])^2) - (2*a*b)/((a^2 + b^2)^2*(a + b*Tan[c + d*x]))) / (3*b*d)) / (2*b)`

3.292.3 Rubi [A] (verified)

Time = 1.32 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.15, number of steps used = 11, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4087, 25, 3042, 4111, 3042, 4012, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^4} dx \xrightarrow{3042} \int \frac{\tan(c + dx)^2(A + B \tan(c + dx))}{(a + b \tan(c + dx))^4} dx$$

$$\begin{array}{c}
\int \frac{-((a^2+b^2)B \tan^2(c+dx)) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{b(a^2+b^2)(a+b \tan(c+dx))^3} dx \\
\downarrow 4087 \\
\int \frac{-((a^2+b^2)B \tan^2(c+dx)) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{b(a^2+b^2)(a+b \tan(c+dx))^3} dx - \frac{a^2(Ab-aB)}{3b^2d(a^2+b^2)(a+b \tan(c+dx))^3} \\
\downarrow 25 \\
\int \frac{-((a^2+b^2)B \tan^2(c+dx)) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{b(a^2+b^2)(a+b \tan(c+dx))^3} dx - \frac{a^2(Ab-aB)}{3b^2d(a^2+b^2)(a+b \tan(c+dx))^3} \\
\downarrow 3042 \\
\int \frac{-((a^2+b^2)B \tan(c+dx)^2) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{b(a^2+b^2)(a+b \tan(c+dx))^3} dx - \frac{a^2(Ab-aB)}{3b^2d(a^2+b^2)(a+b \tan(c+dx))^3} \\
\downarrow 4111 \\
\int \frac{b(Aa^2+2bBa-Ab^2) - b(-Ba^2+2Aba+b^2B) \tan(c+dx)}{(a+b \tan(c+dx))^2} dx - \frac{a(2Ab^3-aB(a^2+3b^2))}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} \\
\frac{b(a^2+b^2)}{a^2(Ab-aB)} \\
\frac{3b^2d(a^2+b^2)(a+b \tan(c+dx))^3} \\
\downarrow 3042 \\
\int \frac{b(Aa^2+2bBa-Ab^2) - b(-Ba^2+2Aba+b^2B) \tan(c+dx)}{(a+b \tan(c+dx))^2} dx - \frac{a(2Ab^3-aB(a^2+3b^2))}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} \\
\frac{b(a^2+b^2)}{a^2(Ab-aB)} \\
\frac{3b^2d(a^2+b^2)(a+b \tan(c+dx))^3} \\
\downarrow 4012 \\
\int \frac{b(Aa^3+3bBa^2-3Ab^2a-b^3B) - b(-Ba^3+3Aba^2+3b^2Ba-Ab^3) \tan(c+dx)}{a+b \tan(c+dx)} dx - \frac{b(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{d(a^2+b^2)(a+b \tan(c+dx))} \\
\frac{b(a^2+b^2)}{a^2+b^2} \\
\frac{a^2(Ab-aB)}{3b^2d(a^2+b^2)(a+b \tan(c+dx))^3} \\
\downarrow 3042
\end{array}$$

3.292. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

$$\begin{aligned}
& \frac{\int \frac{b(Aa^3+3bBa^2-3Ab^2a-b^3B)-b(-Ba^3+3Aba^2+3b^2Ba-Ab^3)\tan(c+dx)}{a+b\tan(c+dx)} dx - \frac{b(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{d(a^2+b^2)(a+b\tan(c+dx))}}{a^2+b^2} - \frac{a(2Ab^3-aB(a^2+3b^2))}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} \\
& \frac{b(a^2+b^2)}{a^2(Ab-aB)} \\
& \frac{3b^2d(a^2+b^2)(a+b\tan(c+dx))^3}{\downarrow 4014} \\
& \frac{\frac{b(a^4(-B)+4a^3Ab+6a^2b^2B-4aAb^3-b^4B)}{a^2+b^2} \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx + \frac{bx(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4)}{a^2+b^2} - \frac{b(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{d(a^2+b^2)(a+b\tan(c+dx))}}{a^2+b^2} - \frac{a(2Ab^3-aB(a^2+3b^2))}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} \\
& \frac{b(a^2+b^2)}{a^2(Ab-aB)} \\
& \frac{3b^2d(a^2+b^2)(a+b\tan(c+dx))^3}{\downarrow 3042} \\
& \frac{\frac{b(a^4(-B)+4a^3Ab+6a^2b^2B-4aAb^3-b^4B)}{a^2+b^2} \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx + \frac{bx(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4)}{a^2+b^2} - \frac{b(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{d(a^2+b^2)(a+b\tan(c+dx))}}{a^2+b^2} - \frac{a(2Ab^3-aB(a^2+3b^2))}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} \\
& \frac{b(a^2+b^2)}{a^2(Ab-aB)} \\
& \frac{3b^2d(a^2+b^2)(a+b\tan(c+dx))^3}{\downarrow 4013} \\
& \frac{a^2(Ab-aB)}{3b^2d(a^2+b^2)(a+b\tan(c+dx))^3} \\
& \frac{b(a^4(-B)+4a^3Ab+6a^2b^2B-4aAb^3-b^4B)}{d(a^2+b^2)} \log(a \cos(c+dx)+b \sin(c+dx)) + \frac{bx(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4)}{a^2+b^2} - \frac{b(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{d(a^2+b^2)(a+b\tan(c+dx))}}{a^2+b^2} - \frac{a(2Ab^3-aB(a^2+3b^2))}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} \\
& \frac{b(a^2+b^2)}{a^2(Ab-aB)}
\end{aligned}$$

input `Int[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]`

```
output -1/3*(a^2*(A*b - a*B))/(b^2*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^3) - (-1/2*
(a*(2*A*b^3 - a*(a^2 + 3*b^2)*B))/(b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2)
+ (((b*(a^4*A - 6*a^2*A*b^2 + A*b^4 + 4*a^3*b*B - 4*a*b^3*B)*x)/(a^2 + b^
2) + (b*(4*a^3*A*b - 4*a*A*b^3 - a^4*B + 6*a^2*b^2*B - b^4*B)*Log[a*Cos[c
+ d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d))/(a^2 + b^2) - (b*(3*a^2*A*b - A
*b^3 - a^3*B + 3*a*b^2*B))/((a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(a^2 + b^
2))/(b*(a^2 + b^2))
```

3.292.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4012 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]
```

```
rule 4013 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*
(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Si
n[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

```
rule 4014 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.
)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]
```

```
rule 4087 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(- (B*c - A*d))*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

```
rule 4111 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]
```

3.292.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 301, normalized size of antiderivative = 1.15

method	result
derivativedivides	$\frac{(4Aa^3b - 4Aab^3 - Ba^4 + 6Ba^2b^2 - Bb^4) \ln(1 + \tan^2(dx+c)) + (-Aa^4 + 6Aa^2b^2 - Ab^4 - 4Ba^3b + 4Ba^3b^3) \arctan(\tan(dx+c))}{(a^2 + b^2)^4} - \frac{3b^2(a^2 + b^2)}{3b^2(a^2 + b^2)}$
default	$\frac{(4Aa^3b - 4Aab^3 - Ba^4 + 6Ba^2b^2 - Bb^4) \ln(1 + \tan^2(dx+c)) + (-Aa^4 + 6Aa^2b^2 - Ab^4 - 4Ba^3b + 4Ba^3b^3) \arctan(\tan(dx+c))}{(a^2 + b^2)^4} - \frac{3b^2(a^2 + b^2)}{3b^2(a^2 + b^2)}$
norman	$-\frac{(8Aa^3b^3 - Ba^6 - 6Ba^4b^2 + 3Ba^2b^4)(\tan^2(dx+c))}{2ad(a^6 + 3a^4b^2 + 3a^2b^4 + b^6)} - \frac{a(Aa^5 - 3Aa^3b^2 + 3Ba^4b - Ba^2b^3)}{3db(a^6 + 3a^4b^2 + 3a^2b^4 + b^6)} - \frac{(Aa^4 - 6Aa^2b^2 + Ab^4 + 4Ba^3b - 4Ba^3b^3)}{(a^6 + 3a^4b^2 + 3a^2b^4 + b^6)(a^2 + b^2)}$
risch	Expression too large to display
parallelrisch	Expression too large to display

```
input int(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x,method=_RETURNVERBOSE)
```

3.292.
$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

output $\frac{1}{d} \left(\frac{1}{(a^2+b^2)^4} \left(\frac{1}{2} (4Aa^3b - 4Aab^3 - Ba^4 + 6Ba^2b^2 - Bb^4) \ln(1 + \tan(dx+c)^2) + (-Aa^4 + 6Aa^2b^2 - Ab^4 - 4Bb^3a + 4Bab^3) \arctan(\tan(dx+c)) \right) - \frac{1}{3} a^2 \frac{(Ab - Ba)}{b^2} \frac{1}{(a^2+b^2)} \frac{1}{(a+b \tan(dx+c))^3} + \frac{(3Aa^2b - Ab^3 - Ba^3 + 3Bab^2)}{(a^2+b^2)^3} \frac{1}{(a+b \tan(dx+c))} - \frac{(4Aa^3b - 4Aab^3 - Ba^4 + 6Ba^2b^2 - Bb^4)}{(a^2+b^2)^4} \ln(a+b \tan(dx+c)) + \frac{1}{2} a \frac{(2Ab^3 - Ba^3 - 3Bab^2)}{(a^2+b^2)^2} \frac{1}{b^2} \frac{1}{(a+b \tan(dx+c))^2} \right)$

3.292.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 836 vs. $2(253) = 506$.

Time = 0.33 (sec) , antiderivative size = 836, normalized size of antiderivative = 3.20

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

$$= \frac{3Ba^7 - 12Aa^6b - 30Ba^5b^2 + 30Aa^4b^3 + 11Ba^3b^4 - 2Aa^2b^5 + (Ba^6b + 2Aa^5b^2 + 18Ba^4b^3 - 30Aa^3b^4$$

input `integrate(tan(dx+c)^2*(A+B*tan(dx+c))/(a+b*tan(dx+c))^4,x, algorithm="fricas")`

output $\frac{1}{6} (3Ba^7 - 12Aa^6b - 30Ba^5b^2 + 30Aa^4b^3 + 11Ba^3b^4 - 2Aa^2b^5 + (Ba^6b + 2Aa^5b^2 + 18Ba^4b^3 - 30Aa^3b^4 - 27Ba^2b^5 + 12Aa^2b^6 - 6(Aa^4b^3 + 4Ba^3b^4 - 6Aa^2b^5 - 4Bab^6 + Ab^7)dx) \tan(dx+c)^3 - 6(Aa^7 + 4Ba^6b - 6Aa^5b^2 - 4Ba^4b^3 + Aa^3b^4)dx + 3(Ba^7 + 2Aa^6b + 16Ba^5b^2 - 24Aa^4b^3 - 23Ba^3b^4 + 16Aa^2b^5 + 6Bab^6 - 2Ab^7 - 6(Aa^5b^2 + 4Ba^4b^3 - 6Aa^3b^4 - 4Ba^2b^5 + Aab^6)dx) \tan(dx+c)^2 + 3(Ba^7 - 4Aa^6b - 6Ba^5b^2 + 4Aa^4b^3 + Ba^3b^4 + (Ba^4b^3 - 4Aa^3b^4 - 6Ba^2b^5 + 4Aab^6 + Bb^7) \tan(dx+c)^3 + 3(Ba^5b^2 - 4Aa^4b^3 - 6Ba^3b^4 + 4Aa^2b^5 + Bab^6) \tan(dx+c)^2 + 3(Ba^6b - 4Aa^5b^2 - 6Ba^4b^3 + 4Aa^3b^4 + Ba^2b^5) \tan(dx+c)) \log((b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2) / (\tan(dx+c)^2 + 1)) + 3(2Aa^7 + 9Ba^6b - 16Aa^5b^2 - 26Ba^4b^3 + 24Aa^3b^4 + 9Ba^2b^5 - 2Aab^6 - 6(Aa^6b + 4Ba^5b^2 - 6Aa^4b^3 - 4Ba^3b^4 + Aa^2b^5)dx) \tan(dx+c) / ((a^8b^3 + 4a^6b^5 + 6a^4b^7 + 4a^2b^9 + b^11)dx \tan(dx+c)^3 + 3(a^9b^2 + 4a^7b^4 + 6a^5b^6 + 4a^3b^8 + ab^10)dx \tan(dx+c)^2 + 3(a^10b + 4a^8b^3 + 6a^6b^5 + 4a^4b^7 + a^2b^9)dx \tan(dx+c) + (a^11 + 4a^9b^2 + 6a^7b^4 + 4a^5b^6 + a^3b^8)dx)$

3.292. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

3.292.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx = \text{Exception raised: AttributeError}$$

```
input integrate(tan(d*x+c)**2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**4,x)
```

```
output Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'
```

3.292.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 526 vs. $2(253) = 506$.

Time = 0.40 (sec) , antiderivative size = 526, normalized size of antiderivative = 2.02

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx = \frac{6(Aa^4+4Ba^3b-6Aa^2b^2-4Bab^3+Ab^4)(dx+c)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} - \frac{6(Ba^4-4Aa^3b-6Ba^2b^2+4Aab^3+Bb^4)\log(b\tan(dx+c)+a)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} + \frac{3(Ba^4-4Aa^3b-6Ba^2b^2+4Aab^3+Bb^4)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8}$$

```
input integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="maxima")
```

```
output -1/6*(6*(A*a^4 + 4*B*a^3*b - 6*A*a^2*b^2 - 4*B*a*b^3 + A*b^4)*(d*x + c)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) - 6*(B*a^4 - 4*A*a^3*b - 6*B*a^2*b^2 + 4*A*a*b^3 + B*b^4)*log(b*tan(d*x + c) + a)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + 3*(B*a^4 - 4*A*a^3*b - 6*B*a^2*b^2 + 4*A*a*b^3 + B*b^4)*log(tan(d*x + c)^2 + 1)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + (B*a^7 + 2*A*a^6*b + 14*B*a^5*b^2 - 20*A*a^4*b^3 - 11*B*a^3*b^4 + 2*A*a^2*b^5 + 6*(B*a^3*b^4 - 3*A*a^2*b^5 - 3*B*a*b^6 + A*b^7)*tan(d*x + c)^2 + 3*(B*a^6*b + 8*B*a^4*b^3 - 14*A*a^3*b^4 - 9*B*a^2*b^5 + 2*A*a*b^6)*tan(d*x + c))/(a^9*b^2 + 3*a^7*b^4 + 3*a^5*b^6 + a^3*b^8 + (a^6*b^5 + 3*a^4*b^7 + 3*a^2*b^9 + b^11)*tan(d*x + c)^3 + 3*(a^7*b^4 + 3*a^5*b^6 + 3*a^3*b^8 + a*b^10)*tan(d*x + c)^2 + 3*(a^8*b^3 + 3*a^6*b^5 + 3*a^4*b^7 + a^2*b^9)*tan(d*x + c))/d
```


3.292.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 632 vs. $2(253) = 506$.

Time = 0.83 (sec) , antiderivative size = 632, normalized size of antiderivative = 2.42

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx =$$

$$\frac{6(Aa^4+4Ba^3b-6Aa^2b^2-4Bab^3+Ab^4)(dx+c)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} + \frac{3(Ba^4-4Aa^3b-6Ba^2b^2+4Aab^3+Bb^4) \log(\tan(dx+c)^2+1)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} - \frac{6(Ba^4b-4Aa^3b^2-6Aa^2b^3+3Ab^4)}{a^8b^2+4a^6b^4+6a^4b^6+4a^2b^8+b^{10}}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="giac")`

output `-1/6*(6*(A*a^4 + 4*B*a^3*b - 6*A*a^2*b^2 - 4*B*a*b^3 + A*b^4)*(d*x + c)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + 3*(B*a^4 - 4*A*a^3*b - 6*B*a^2*b^2 + 4*A*a*b^3 + B*b^4)*log(tan(d*x + c)^2 + 1)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) - 6*(B*a^4*b - 4*A*a^3*b^2 - 6*B*a^2*b^3 + 4*A*a*b^4 + B*b^5)*log(abs(b*tan(d*x + c) + a))/(a^8*b + 4*a^6*b^3 + 6*a^4*b^5 + 4*a^2*b^7 + b^9) + (11*B*a^4*b^5*tan(d*x + c)^3 - 44*A*a^3*b^6*tan(d*x + c)^3 - 66*B*a^2*b^7*tan(d*x + c)^3 + 44*A*a*b^8*tan(d*x + c)^3 + 11*B*b^9*tan(d*x + c)^3 + 39*B*a^5*b^4*tan(d*x + c)^2 - 150*A*a^4*b^5*tan(d*x + c)^2 - 210*B*a^3*b^6*tan(d*x + c)^2 + 120*A*a^2*b^7*tan(d*x + c)^2 + 15*B*a*b^8*tan(d*x + c)^2 + 6*A*b^9*tan(d*x + c)^2 + 3*B*a^8*b*tan(d*x + c) + 60*B*a^6*b^3*tan(d*x + c) - 174*A*a^5*b^4*tan(d*x + c) - 201*B*a^4*b^5*tan(d*x + c) + 96*A*a^3*b^6*tan(d*x + c) + 6*B*a^2*b^7*tan(d*x + c) + 6*A*a*b^8*tan(d*x + c) + B*a^9 + 2*A*a^8*b + 26*B*a^7*b^2 - 62*A*a^6*b^3 - 63*B*a^5*b^4 + 26*A*a^4*b^5 + 2*A*a^2*b^7)/((a^8*b^2 + 4*a^6*b^4 + 6*a^4*b^6 + 4*a^2*b^8 + b^10)*(b*tan(d*x + c) + a)^3))/d`

3.292.9 Mupad [B] (verification not implemented)

Time = 7.87 (sec) , antiderivative size = 446, normalized size of antiderivative = 1.71

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$$

$$= \frac{\ln(a+b\tan(c+dx)) \left(\frac{B}{(a^2+b^2)^2} - \frac{4b(Aa+2Bb)}{(a^2+b^2)^3} + \frac{8b^3(Aa+Bb)}{(a^2+b^2)^4} \right)}{d} - \frac{\frac{\tan(c+dx)^2 (Ba^3b^2-3Aa^2b^3-3Bab^4+Ab^5)}{a^6+3a^4b^2+3a^2b^4+b^6} + \frac{a(Ba^6+2Aa^5b+14Ba^4b^2-20Aa^3b^3-11Ba^2b^4+2Aab^5)}{6b^2(a^6+3a^4b^2+3a^2b^4+b^6)} + \frac{\tan(c+dx)(Ba^6+8Aa^5b+14Ba^4b^2-20Aa^3b^3-11Ba^2b^4+2Aab^5)}{2b(a^6+3a^4b^2+3a^2b^4+b^6)}}{d(a^3+3a^2b\tan(c+dx)+3ab^2\tan(c+dx)^2+b^3\tan(c+dx)^3)} - \frac{\ln(\tan(c+dx)-i)(A+B1i)}{2d(a^41i-4a^3b-a^2b^26i+4ab^3+b^41i)} - \frac{\ln(\tan(c+dx)+1i)(B+A1i)}{2d(a^4-a^3b4i-6a^2b^2+ab^34i+b^4)}$$

input `int((tan(c + d*x))^2*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^4,x)`output `(log(a + b*tan(c + d*x))*(B/(a^2 + b^2)^2 - (4*b*(A*a + 2*B*b))/(a^2 + b^2)^3 + (8*b^3*(A*a + B*b))/(a^2 + b^2)^4))/d - ((tan(c + d*x))^2*(A*b^5 - 3*A*a^2*b^3 + B*a^3*b^2 - 3*B*a*b^4))/(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2) + (a*(B*a^6 - 20*A*a^3*b^3 - 11*B*a^2*b^4 + 14*B*a^4*b^2 + 2*A*a*b^5 + 2*A*a^5*b))/(6*b^2*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2)) + (tan(c + d*x)*(B*a^6 - 14*A*a^3*b^3 - 9*B*a^2*b^4 + 8*B*a^4*b^2 + 2*A*a*b^5))/(2*b*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2)))/(d*(a^3 + b^3*tan(c + d*x)^3 + 3*a*b^2*tan(c + d*x)^2 + 3*a^2*b*tan(c + d*x))) - (log(tan(c + d*x) - 1i)*(A + B*1i))/(2*d*(4*a*b^3 - 4*a^3*b + a^4*1i + b^4*1i - a^2*b^2*6i)) - (log(tan(c + d*x) + 1i)*(A*1i + B))/(2*d*(a*b^3*4i - a^3*b*4i + a^4 + b^4 - 6*a^2*b^2))`

3.293 $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

3.293.1 Optimal result 2820
 3.293.2 Mathematica [C] (verified) 2821
 3.293.3 Rubi [A] (verified) 2821
 3.293.4 Maple [A] (verified) 2824
 3.293.5 Fricas [B] (verification not implemented) 2825
 3.293.6 Sympy [F(-2)] 2826
 3.293.7 Maxima [B] (verification not implemented) 2826
 3.293.8 Giac [B] (verification not implemented) 2827
 3.293.9 Mupad [B] (verification not implemented) 2828

3.293.1 Optimal result

Integrand size = 29, antiderivative size = 250

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

$$= \frac{(4a^3Ab - 4aAb^3 - a^4B + 6a^2b^2B - b^4B)x}{(a^2 + b^2)^4}$$

$$- \frac{(a^4A - 6a^2Ab^2 + Ab^4 + 4a^3bB - 4ab^3B) \log(a \cos(c+dx) + b \sin(c+dx))}{(a^2 + b^2)^4 d}$$

$$+ \frac{a(Ab - aB)}{3b(a^2 + b^2) d(a + b \tan(c+dx))^3}$$

$$+ \frac{a^2A - Ab^2 + 2abB}{2(a^2 + b^2)^2 d(a + b \tan(c+dx))^2} + \frac{a^3A - 3aAb^2 + 3a^2bB - b^3B}{(a^2 + b^2)^3 d(a + b \tan(c+dx))}$$

output

```
(4*A*a^3*b-4*A*a*b^3-B*a^4+6*B*a^2*b^2-B*b^4)*x/(a^2+b^2)^4-(A*a^4-6*A*a^2*b^2+A*b^4+4*B*a^3*b-4*B*a*b^3)*ln(a*cos(d*x+c)+b*sin(d*x+c))/(a^2+b^2)^4/d+1/3*a*(A*b-B*a)/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^3+1/2*(A*a^2-A*b^2+2*B*a*b)/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^2+(A*a^3-3*A*a*b^2+3*B*a^2*b-B*b^3)/(a^2+b^2)^3/d/(a+b*tan(d*x+c))
```

3.293.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.32 (sec) , antiderivative size = 248, normalized size of antiderivative = 0.99

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^4} dx$$

$$= \frac{\frac{3(A+ib) \log(i - \tan(c+dx))}{(a+ib)^4} + \frac{3(A-ib) \log(i + \tan(c+dx))}{(a-ib)^4} - \frac{6(a^4 A - 6a^2 Ab^2 + Ab^4 + 4a^3 bB - 4ab^3 B) \log(a + b \tan(c+dx))}{(a^2 + b^2)^4} + \frac{2a(Ab - a^2 B)}{b(a^2 + b^2)(a + b \tan(c + dx))}}{6d}$$

```
input Integrate[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]
```

```
output ((3*(A + I*B)*Log[I - Tan[c + d*x]])/(a + I*b)^4 + (3*(A - I*B)*Log[I + Tan[c + d*x]])/(a - I*b)^4 - (6*(a^4*A - 6*a^2*A*b^2 + A*b^4 + 4*a^3*b*B - 4*a*b^3*B)*Log[a + b*Tan[c + d*x]])/(a^2 + b^2)^4 + (2*a*(A*b - a*B))/(b*(a^2 + b^2)*(a + b*Tan[c + d*x])^3) + (3*(a^2*A - A*b^2 + 2*a*b*B))/((a^2 + b^2)^2*(a + b*Tan[c + d*x])^2) + (6*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B))/((a^2 + b^2)^3*(a + b*Tan[c + d*x]))/(6*d)
```

3.293.3 Rubi [A] (verified)

Time = 1.17 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.13, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 4074, 3042, 4012, 3042, 4012, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^4} dx$$

↓ 3042

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^4} dx$$

↓ 4074

$$\frac{\int \frac{Ab - aB + (aA + bB) \tan(c + dx)}{(a + b \tan(c + dx))^3} dx}{a^2 + b^2} + \frac{a(Ab - aB)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3}$$

↓ 3042

$$\frac{\int \frac{Ab - aB + (aA + bB) \tan(c + dx)}{(a + b \tan(c + dx))^3} dx}{a^2 + b^2} + \frac{a(Ab - aB)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3}$$

↓ 4012

$$\frac{\int \frac{-Ba^2 + 2Aba + b^2B + (Aa^2 + 2bBa - Ab^2) \tan(c + dx)}{(a + b \tan(c + dx))^2} dx}{a^2 + b^2} + \frac{a^2A + 2abB - Ab^2}{2d(a^2 + b^2)(a + b \tan(c + dx))^2} +$$

$$\frac{a^2 + b^2}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} \frac{a(Ab - aB)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3}$$

↓ 3042

$$\frac{\int \frac{-Ba^2 + 2Aba + b^2B + (Aa^2 + 2bBa - Ab^2) \tan(c + dx)}{(a + b \tan(c + dx))^2} dx}{a^2 + b^2} + \frac{a^2A + 2abB - Ab^2}{2d(a^2 + b^2)(a + b \tan(c + dx))^2} +$$

$$\frac{a^2 + b^2}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} \frac{a(Ab - aB)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3}$$

↓ 4012

$$\frac{\int \frac{-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3 + (Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{a^3A + 3a^2bB - 3aAb^2 - b^3B}{d(a^2 + b^2)(a + b \tan(c + dx))} + \frac{a^2A + 2abB - Ab^2}{2d(a^2 + b^2)(a + b \tan(c + dx))^2} +$$

$$\frac{a^2 + b^2}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} \frac{a(Ab - aB)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3}$$

↓ 3042

$$\frac{\int \frac{-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3 + (Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{a^3A + 3a^2bB - 3aAb^2 - b^3B}{d(a^2 + b^2)(a + b \tan(c + dx))} + \frac{a^2A + 2abB - Ab^2}{2d(a^2 + b^2)(a + b \tan(c + dx))^2} +$$

$$\frac{a^2 + b^2}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} \frac{a(Ab - aB)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3}$$

↓ 4014

$$\frac{x(a^4(-B) + 4a^3Ab + 6a^2b^2B - 4aAb^3 - b^4B)}{a^2 + b^2} - \frac{(a^4A + 4a^3bB - 6a^2Ab^2 - 4ab^3B + Ab^4) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} + \frac{a^3A + 3a^2bB - 3aAb^2 - b^3B}{d(a^2 + b^2)(a + b \tan(c + dx))} + \frac{a^2A + 2abB - Ab^2}{2d(a^2 + b^2)(a + b \tan(c + dx))^2} +$$

$$\frac{a^2 + b^2}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} \frac{a(Ab - aB)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3}$$

↓ 3042

3.293. $\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^4} dx$

$$\frac{x(a^4(-B)+4a^3Ab+6a^2b^2B-4aAb^3-b^4B)}{a^2+b^2} - \frac{(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{a^3A+3a^2bB-3aAb^2-b^3B}{d(a^2+b^2)(a+b \tan(c+dx))} + \frac{a^2A+2abB-Ab^2}{2d(a^2+b^2)(a+b \tan(c+dx))}$$

$$\frac{a(Ab - aB)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3}$$

↓ 4013

$$\frac{a(Ab - aB)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^3} + \frac{x(a^4(-B)+4a^3Ab+6a^2b^2B-4aAb^3-b^4B)}{a^2+b^2} - \frac{(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4) \log(a \cos(c + dx))}{d(a^2+b^2)}$$

$$\frac{a^2A+2abB-Ab^2}{2d(a^2+b^2)(a+b \tan(c+dx))^2} + \frac{a^3A+3a^2bB-3aAb^2-b^3B}{d(a^2+b^2)(a+b \tan(c+dx))} + \frac{x(a^4(-B)+4a^3Ab+6a^2b^2B-4aAb^3-b^4B)}{a^2+b^2} - \frac{(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4) \log(a \cos(c + dx))}{d(a^2+b^2)}$$

$$a^2 + b^2$$

```
input Int[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]
```

```
output (a*(A*b - a*B))/(3*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^3) + ((a^2*A - A*b^2 + 2*a*b*B)/(2*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + (((4*a^3*A*b - 4*a*A*b^3 - a^4*B + 6*a^2*b^2*B - b^4*B)*x)/(a^2 + b^2) - ((a^4*A - 6*a^2*A*b^2 + A*b^4 + 4*a^3*b*B - 4*a*b^3*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a^2 + b^2)*d)/(a^2 + b^2) + (a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B)/(a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(a^2 + b^2))/(a^2 + b^2)
```

3.293.3.1 Defintions of rubi rules used

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]
```

```
rule 4012 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]
```

3.293. $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

rule 4074 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.293.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.15

method	result
derivativedivides	$\frac{(A a^4 - 6A a^2 b^2 + A b^4 + 4B a^3 b - 4B a b^3) \ln(1 + \tan^2(dx+c)) + (4A a^3 b - 4A a b^3 - B a^4 + 6B a^2 b^2 - B b^4) \arctan(\tan(dx+c))}{(a^2 + b^2)^4} + \frac{1}{3(a^2 + b^2)}$
default	$\frac{(A a^4 - 6A a^2 b^2 + A b^4 + 4B a^3 b - 4B a b^3) \ln(1 + \tan^2(dx+c)) + (4A a^3 b - 4A a b^3 - B a^4 + 6B a^2 b^2 - B b^4) \arctan(\tan(dx+c))}{(a^2 + b^2)^4} + \frac{1}{3(a^2 + b^2)}$
norman	$\frac{(4A a^3 b - 4A a b^3 - B a^4 + 6B a^2 b^2 - B b^4) a^3 x}{(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)(a^2 + b^2)} + \frac{b^3 (4A a^3 b - 4A a b^3 - B a^4 + 6B a^2 b^2 - B b^4) x (\tan^3(dx+c))}{(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)(a^2 + b^2)} + \frac{a (9A a^4 b^2 - 8A a^2 b^4 - A b^6)}{6b^2 (a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)}$
risch	Expression too large to display
parallelrisch	Expression too large to display

input `int(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x,method=_RETURNVERBOSE)`

3.293.
$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

output $1/d*(1/(a^2+b^2)^4*(1/2*(A*a^4-6*A*a^2*b^2+A*b^4+4*B*a^3*b-4*B*a*b^3)*\ln(1+\tan(dx+c)^2)+(4*A*a^3*b-4*A*a*b^3-B*a^4+6*B*a^2*b^2-B*b^4)*\arctan(\tan(dx+c)))+1/3*a*(A*b-B*a)/(a^2+b^2)/b/(a+b*\tan(dx+c))^3+1/2*(A*a^2-A*b^2+2*B*a*b)/(a^2+b^2)^2/(a+b*\tan(dx+c))^2+(A*a^3-3*A*a*b^2+3*B*a^2*b-B*b^3)/(a^2+b^2)^3/(a+b*\tan(dx+c))-(A*a^4-6*A*a^2*b^2+A*b^4+4*B*a^3*b-4*B*a*b^3)/(a^2+b^2)^4*\ln(a+b*\tan(dx+c)))$

3.293.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 838 vs. $2(245) = 490$.

Time = 0.30 (sec) , antiderivative size = 838, normalized size of antiderivative = 3.35

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx =$$

$$\frac{12Ba^6b - 27Aa^5b^2 - 30Ba^4b^3 + 18Aa^3b^4 + 2Ba^2b^5 + Aab^6 - (2Ba^5b^2 - 11Aa^4b^3 - 30Ba^3b^4 + 30Aa^2b^5 + 12Ba^2b^6 - 3Aab^7 - 6*(Ba^4b^3 - 4Aa^3b^4 - 6Ba^2b^5 + 4Aa^2b^6 + Bb^7)*dx)*\tan(dx+c)^3 + 6*(Ba^7 - 4Aa^6b - 6Ba^5b^2 + 4Aa^4b^3 + Ba^3b^4)*dx - 3*(2Ba^6b - 9Aa^5b^2 - 24Ba^4b^3 + 26Aa^3b^4 + 16Ba^2b^5 - 9Aa^2b^6 - 2Bb^7 - 6*(Ba^5b^2 - 4Aa^4b^3 - 6Ba^3b^4 + 4Aa^2b^5 + Ba^2b^6)*dx)*\tan(dx+c)^2 + 3*(Aa^7 + 4Ba^6b - 6Aa^5b^2 - 4Ba^4b^3 + Aa^3b^4 + (Aa^4b^3 + 4Ba^3b^4 - 6Aa^2b^5 - 4Ba^2b^6 + Ab^7)*\tan(dx+c)^3 + 3*(Aa^5b^2 + 4Ba^4b^3 - 6Aa^3b^4 - 4Ba^2b^5 + Aa^2b^6)*\tan(dx+c)^2 + 3*(Aa^6b + 4Ba^5b^2 - 6Aa^4b^3 - 4Ba^3b^4 + Aa^2b^5)*\tan(dx+c))*\log((b^2*\tan(dx+c)^2 + 2*a*b*\tan(dx+c) + a^2)/(\tan(dx+c)^2 + 1)) - 3*(2Ba^7 - 6Aa^6b - 16Ba^5b^2 + 23Aa^4b^3 + 24Ba^3b^4 - 16Aa^2b^5 - 2Ba^2b^6 - Ab^7 - 6*(Ba^6b - 4Aa^5b^2 - 6Ba^4b^3 + 4Aa^3b^4 + Ba^2b^5)*dx)*\tan(dx+c))/((a^8*b^3 + 4a^6*b^5 + 6a^4*b^7 + 4a^2*b^9 + b^11)*d*\tan(dx+c)^3 + 3*(a^9*b^2 + 4a^7*b^4 + 6a^5*b^6 + 4a^3*b^8 + a*b^10)*d*\tan(dx+c)^2 + 3*(a^10*b + 4a^8*b^3 + 6a^6*b^5 + 4a^4*b^7 + a^2*b^9)*d*\tan(dx+c) + (a^11 + 4a^9*b^2 + 6a^7*b^4 + 4a^5*b^6 + a^3*b^8)*d)$$

input `integrate(tan(dx+c)*(A+B*tan(dx+c))/(a+b*tan(dx+c))^4,x, algorithm="fricas")`

output $-1/6*(12*B*a^6*b - 27*A*a^5*b^2 - 30*B*a^4*b^3 + 18*A*a^3*b^4 + 2*B*a^2*b^5 + A*a*b^6 - (2*B*a^5*b^2 - 11*A*a^4*b^3 - 30*B*a^3*b^4 + 30*A*a^2*b^5 + 12*B*a^2*b^6 - 3*A*b^7 - 6*(B*a^4*b^3 - 4*A*a^3*b^4 - 6*B*a^2*b^5 + 4*A*a^2*b^6 + B*b^7)*dx)*\tan(dx+c)^3 + 6*(B*a^7 - 4*A*a^6*b - 6*B*a^5*b^2 + 4*A*a^4*b^3 + B*a^3*b^4)*dx - 3*(2*B*a^6*b - 9*A*a^5*b^2 - 24*B*a^4*b^3 + 26*A*a^3*b^4 + 16*B*a^2*b^5 - 9*A*a^2*b^6 - 2*B*b^7 - 6*(B*a^5*b^2 - 4*A*a^4*b^3 - 6*B*a^3*b^4 + 4*A*a^2*b^5 + B*a^2*b^6)*dx)*\tan(dx+c)^2 + 3*(A*a^7 + 4*B*a^6*b - 6*A*a^5*b^2 - 4*B*a^4*b^3 + A*a^3*b^4 + (A*a^4*b^3 + 4*B*a^3*b^4 - 6*A*a^2*b^5 - 4*B*a^2*b^6 + A*b^7)*\tan(dx+c)^3 + 3*(A*a^5*b^2 + 4*B*a^4*b^3 - 6*A*a^3*b^4 - 4*B*a^2*b^5 + A*a^2*b^6)*\tan(dx+c)^2 + 3*(A*a^6*b + 4*B*a^5*b^2 - 6*A*a^4*b^3 - 4*B*a^3*b^4 + A*a^2*b^5)*\tan(dx+c))*\log((b^2*\tan(dx+c)^2 + 2*a*b*\tan(dx+c) + a^2)/(\tan(dx+c)^2 + 1)) - 3*(2*B*a^7 - 6*A*a^6*b - 16*B*a^5*b^2 + 23*A*a^4*b^3 + 24*B*a^3*b^4 - 16*A*a^2*b^5 - 2*B*a^2*b^6 - A*b^7 - 6*(B*a^6*b - 4*A*a^5*b^2 - 6*B*a^4*b^3 + 4*A*a^3*b^4 + B*a^2*b^5)*dx)*\tan(dx+c))/((a^8*b^3 + 4*a^6*b^5 + 6*a^4*b^7 + 4*a^2*b^9 + b^11)*d*\tan(dx+c)^3 + 3*(a^9*b^2 + 4*a^7*b^4 + 6*a^5*b^6 + 4*a^3*b^8 + a*b^10)*d*\tan(dx+c)^2 + 3*(a^10*b + 4*a^8*b^3 + 6*a^6*b^5 + 4*a^4*b^7 + a^2*b^9)*d*\tan(dx+c) + (a^11 + 4*a^9*b^2 + 6*a^7*b^4 + 4*a^5*b^6 + a^3*b^8)*d)$

3.293. $\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$

3.293.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx = \text{Exception raised: AttributeError}$$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**4,x)
```

```
output Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'
```

3.293.7 Maxima [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 523 vs. $2(245) = 490$.

Time = 0.37 (sec) , antiderivative size = 523, normalized size of antiderivative = 2.09

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx =$$

$$\frac{6(Ba^4-4Aa^3b-6Ba^2b^2+4Aab^3+Bb^4)(dx+c)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} + \frac{6(Aa^4+4Ba^3b-6Aa^2b^2-4Bab^3+Ab^4)\log(b\tan(dx+c)+a)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} - \frac{3(Aa^4+4Ba^3b-6Aa^2b^2-4Bab^3+Ab^4)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8}$$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="maxima")
```

```
output -1/6*(6*(B*a^4 - 4*A*a^3*b - 6*B*a^2*b^2 + 4*A*a*b^3 + B*b^4)*(d*x + c)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + 6*(A*a^4 + 4*B*a^3*b - 6*A*a^2*b^2 - 4*B*a*b^3 + A*b^4)*log(b*tan(d*x + c) + a)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) - 3*(A*a^4 + 4*B*a^3*b - 6*A*a^2*b^2 - 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)^2 + 1)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + (2*B*a^6 - 11*A*a^5*b - 20*B*a^4*b^2 + 14*A*a^3*b^3 + 2*B*a^2*b^4 + A*a*b^5 - 6*(A*a^3*b^3 + 3*B*a^2*b^4 - 3*A*a*b^5 - B*b^6)*tan(d*x + c)^2 - 3*(5*A*a^4*b^2 + 14*B*a^3*b^3 - 12*A*a^2*b^4 - 2*B*a*b^5 - A*b^6)*tan(d*x + c))/(a^9*b + 3*a^7*b^3 + 3*a^5*b^5 + a^3*b^7 + (a^6*b^4 + 3*a^4*b^6 + 3*a^2*b^8 + b^10)*tan(d*x + c)^3 + 3*(a^7*b^3 + 3*a^5*b^5 + 3*a^3*b^7 + a*b^9)*tan(d*x + c)^2 + 3*(a^8*b^2 + 3*a^6*b^4 + 3*a^4*b^6 + a^2*b^8)*tan(d*x + c))/d
```

3.293. $\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$

3.293.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 638 vs. $2(245) = 490$.

Time = 0.69 (sec) , antiderivative size = 638, normalized size of antiderivative = 2.55

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx =$$

$$\frac{6(Ba^4-4Aa^3b-6Ba^2b^2+4Aab^3+Bb^4)(dx+c)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} - \frac{3(Aa^4+4Ba^3b-6Aa^2b^2-4Bab^3+Ab^4)\log(\tan(dx+c)^2+1)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} + \frac{6(Aa^4b+4Ba^3b^2-6Aa^2b^3+4Aab^4+Bb^5)}{a^8b+4a^6b^3+6a^4b^5+4a^2b^7+b^9}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="giac")`

output

```
-1/6*(6*(B*a^4 - 4*A*a^3*b - 6*B*a^2*b^2 + 4*A*a*b^3 + B*b^4)*(d*x + c)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) - 3*(A*a^4 + 4*B*a^3*b - 6*A*a^2*b^2 - 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)^2 + 1)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + 6*(A*a^4*b + 4*B*a^3*b^2 - 6*A*a^2*b^3 - 4*B*a*b^4 + A*b^5)*log(abs(b*tan(d*x + c) + a))/(a^8*b + 4*a^6*b^3 + 6*a^4*b^5 + 4*a^2*b^7 + b^9) - (11*A*a^4*b^4*tan(d*x + c)^3 + 44*B*a^3*b^5*tan(d*x + c)^3 - 66*A*a^2*b^6*tan(d*x + c)^3 - 44*B*a*b^7*tan(d*x + c)^3 + 11*A*b^8*tan(d*x + c)^3 + 39*A*a^5*b^3*tan(d*x + c)^2 + 150*B*a^4*b^4*tan(d*x + c)^2 - 210*A*a^3*b^5*tan(d*x + c)^2 - 120*B*a^2*b^6*tan(d*x + c)^2 + 15*A*a*b^7*tan(d*x + c)^2 - 6*B*b^8*tan(d*x + c)^2 + 48*A*a^6*b^2*tan(d*x + c) + 174*B*a^5*b^3*tan(d*x + c) - 219*A*a^4*b^4*tan(d*x + c) - 96*B*a^3*b^5*tan(d*x + c) - 6*A*a^2*b^6*tan(d*x + c) - 6*B*a*b^7*tan(d*x + c) - 3*A*b^8*tan(d*x + c) - 2*B*a^8 + 22*A*a^7*b + 62*B*a^6*b^2 - 69*A*a^5*b^3 - 26*B*a^4*b^4 - 4*A*a^3*b^5 - 2*B*a^2*b^6 - A*a*b^7)/((a^8*b + 4*a^6*b^3 + 6*a^4*b^5 + 4*a^2*b^7 + b^9)*(b*tan(d*x + c) + a)^3))/d
```

3.293.9 Mupad [B] (verification not implemented)

Time = 8.45 (sec) , antiderivative size = 447, normalized size of antiderivative = 1.79

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx =$$

$$\frac{\frac{\tan(c+dx)(-5Aa^4b-14Ba^3b^2+12Aa^2b^3+2Bab^4+Ab^5)}{2(a^6+3a^4b^2+3a^2b^4+b^6)} + \frac{2Ba^6-11Aa^5b-20Ba^4b^2+14Aa^3b^3+2Ba^2b^4+Aab^5}{6b(a^6+3a^4b^2+3a^2b^4+b^6)} + \frac{\tan(c+dx)}{d(a^3+3a^2b\tan(c+dx)+3ab^2\tan(c+dx)^2+b^3\tan(c+dx)^3)} + \frac{\ln(a+b\tan(c+dx))\left(\frac{A}{(a^2+b^2)^2} - \frac{4b(2Ab-Ba)}{(a^2+b^2)^3} + \frac{8b^3(Ab-Ba)}{(a^2+b^2)^4}\right)}{d} + \frac{\ln(\tan(c+dx)+1i)(B+A1i)}{2d(a^41i+4a^3b-a^2b^26i-4ab^3+b^41i)} + \frac{\ln(\tan(c+dx)-i)(A+B1i)}{2d(a^4+a^3b4i-6a^2b^2-ab^34i+b^4)}$$

input `int((tan(c + d*x)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^4,x)`output `(log(tan(c + d*x) + 1i)*(A*1i + B))/(2*d*(4*a^3*b - 4*a*b^3 + a^4*1i + b^4*1i - a^2*b^2*6i)) - (log(a + b*tan(c + d*x))*(A/(a^2 + b^2)^2 - (4*b*(2*A*b - B*a))/(a^2 + b^2)^3 + (8*b^3*(A*b - B*a))/(a^2 + b^2)^4))/d - ((tan(c + d*x)*(A*b^5 + 12*A*a^2*b^3 - 14*B*a^3*b^2 - 5*A*a^4*b + 2*B*a*b^4))/(2*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2)) + (2*B*a^6 + 14*A*a^3*b^3 + 2*B*a^2*b^4 - 20*B*a^4*b^2 + A*a*b^5 - 11*A*a^5*b)/(6*b*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2)) + (tan(c + d*x)^2*(B*b^5 - A*a^3*b^2 - 3*B*a^2*b^3 + 3*A*a*b^4))/(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2))/(d*(a^3 + b^3*tan(c + d*x)^3 + 3*a*b^2*tan(c + d*x)^2 + 3*a^2*b*tan(c + d*x))) + (log(tan(c + d*x) - 1i)*(A + B*1i))/(2*d*(a^3*b*4i - a*b^3*4i + a^4 + b^4 - 6*a^2*b^2))`

3.294 $\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^4} dx$

3.294.1 Optimal result 2829
 3.294.2 Mathematica [C] (verified) 2830
 3.294.3 Rubi [A] (verified) 2830
 3.294.4 Maple [A] (verified) 2833
 3.294.5 Fricas [B] (verification not implemented) 2834
 3.294.6 Sympy [F(-2)] 2835
 3.294.7 Maxima [B] (verification not implemented) 2836
 3.294.8 Giac [B] (verification not implemented) 2836
 3.294.9 Mupad [B] (verification not implemented) 2837

3.294.1 Optimal result

Integrand size = 23, antiderivative size = 247

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^4} dx$$

$$= \frac{(a^4 A - 6a^2 Ab^2 + Ab^4 + 4a^3 b B - 4ab^3 B) x}{(a^2 + b^2)^4}$$

$$+ \frac{(4a^3 Ab - 4aAb^3 - a^4 B + 6a^2 b^2 B - b^4 B) \log(a \cos(c + dx) + b \sin(c + dx))}{(a^2 + b^2)^4 d}$$

$$- \frac{Ab - aB}{3(a^2 + b^2) d(a + b \tan(c + dx))^3}$$

$$- \frac{2aAb - a^2 B + b^2 B}{2(a^2 + b^2)^2 d(a + b \tan(c + dx))^2} - \frac{3a^2 Ab - Ab^3 - a^3 B + 3ab^2 B}{(a^2 + b^2)^3 d(a + b \tan(c + dx))}$$

output

```
(A*a^4-6*A*a^2*b^2+A*b^4+4*B*a^3*b-4*B*a*b^3)*x/(a^2+b^2)^4+(4*A*a^3*b-4*A*a*b^3-B*a^4+6*B*a^2*b^2-B*b^4)*ln(a*cos(d*x+c)+b*sin(d*x+c))/(a^2+b^2)^4/d+1/3*(-A*b+B*a)/(a^2+b^2)/d/(a+b*tan(d*x+c))^3+1/2*(-2*A*a*b+B*a^2-B*b^2)/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^2+(-3*A*a^2*b+A*b^3+B*a^3-3*B*a*b^2)/(a^2+b^2)^3/d/(a+b*tan(d*x+c))
```

3.294.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.28 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.32

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^4} dx =$$

$$\frac{(Ab - aB) \left(\frac{3i \log(i - \tan(c + dx))}{(a + ib)^4} - \frac{3i \log(i + \tan(c + dx))}{(a - ib)^4} - \frac{24a(a - b)b(a + b) \log(a + b \tan(c + dx))}{(a^2 + b^2)^4} + \frac{2b}{(a^2 + b^2)(a + b \tan(c + dx))^3} + \frac{6bd}{(a^2 + b^2)(a + b \tan(c + dx))^2} + \frac{4ab}{(a^2 + b^2)^2(a + b \tan(c + dx))} \right)}{2bd}$$

input `Integrate[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^4,x]`

output `-1/6*((A*b - a*B)*(((3*I)*Log[I - Tan[c + d*x]])/(a + I*b)^4 - ((3*I)*Log[I + Tan[c + d*x]])/(a - I*b)^4 - (24*a*(a - b)*b*(a + b)*Log[a + b*Tan[c + d*x]])/(a^2 + b^2)^4 + (2*b)/((a^2 + b^2)*(a + b*Tan[c + d*x])^3) + (6*a*b)/((a^2 + b^2)^2*(a + b*Tan[c + d*x])^2) + (6*b*(3*a^2 - b^2))/((a^2 + b^2)^3*(a + b*Tan[c + d*x])))/(b*d) - (B*(Log[I - Tan[c + d*x]]/(I*a - b)^3 - Log[I + Tan[c + d*x]]/(I*a + b)^3 - (2*b*(3*a^2 - b^2)*Log[a + b*Tan[c + d*x]])/(a^2 + b^2)^3 + b/((a^2 + b^2)*(a + b*Tan[c + d*x])^2) + (4*a*b)/((a^2 + b^2)^2*(a + b*Tan[c + d*x]))))/(2*b*d)`

3.294.3 Rubi [A] (verified)

Time = 1.15 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.13, number of steps used = 10, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.435$, Rules used = {3042, 4012, 3042, 4012, 3042, 4012, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^4} dx$$

$$\downarrow \text{4012}$$

3.294. $\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^4} dx$

$$\begin{aligned}
 & \frac{\int \frac{aA+bB-(Ab-aB)\tan(c+dx)}{(a+b\tan(c+dx))^3} dx}{a^2+b^2} - \frac{Ab-aB}{3d(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{aA+bB-(Ab-aB)\tan(c+dx)}{(a+b\tan(c+dx))^3} dx}{a^2+b^2} - \frac{Ab-aB}{3d(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \quad \downarrow \text{4012} \\
 & \frac{\int \frac{Aa^2+2bBa-Ab^2-(-Ba^2+2Aba+b^2B)\tan(c+dx)}{(a+b\tan(c+dx))^2} dx}{a^2+b^2} - \frac{a^2(-B)+2aAb+b^2B}{2d(a^2+b^2)(a+b\tan(c+dx))^2} \\
 & \quad \frac{a^2+b^2}{Ab-aB} \\
 & \quad \frac{3d(a^2+b^2)(a+b\tan(c+dx))^3}{3d(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{Aa^2+2bBa-Ab^2-(-Ba^2+2Aba+b^2B)\tan(c+dx)}{(a+b\tan(c+dx))^2} dx}{a^2+b^2} - \frac{a^2(-B)+2aAb+b^2B}{2d(a^2+b^2)(a+b\tan(c+dx))^2} \\
 & \quad \frac{a^2+b^2}{Ab-aB} \\
 & \quad \frac{3d(a^2+b^2)(a+b\tan(c+dx))^3}{3d(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \quad \downarrow \text{4012} \\
 & \frac{\int \frac{Aa^3+3bBa^2-3Ab^2a-b^3B-(-Ba^3+3Aba^2+3b^2Ba-Ab^3)\tan(c+dx)}{a+b\tan(c+dx)} dx}{a^2+b^2} - \frac{a^3(-B)+3a^2Ab+3ab^2B-Ab^3}{d(a^2+b^2)(a+b\tan(c+dx))} - \frac{a^2(-B)+2aAb+b^2B}{2d(a^2+b^2)(a+b\tan(c+dx))^2} \\
 & \quad \frac{a^2+b^2}{Ab-aB} \\
 & \quad \frac{3d(a^2+b^2)(a+b\tan(c+dx))^3}{3d(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{Aa^3+3bBa^2-3Ab^2a-b^3B-(-Ba^3+3Aba^2+3b^2Ba-Ab^3)\tan(c+dx)}{a+b\tan(c+dx)} dx}{a^2+b^2} - \frac{a^3(-B)+3a^2Ab+3ab^2B-Ab^3}{d(a^2+b^2)(a+b\tan(c+dx))} - \frac{a^2(-B)+2aAb+b^2B}{2d(a^2+b^2)(a+b\tan(c+dx))^2} \\
 & \quad \frac{a^2+b^2}{Ab-aB} \\
 & \quad \frac{3d(a^2+b^2)(a+b\tan(c+dx))^3}{3d(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \quad \downarrow \text{4014}
 \end{aligned}$$

3.294. $\int \frac{A+B\tan(c+dx)}{(a+b\tan(c+dx))^4} dx$

$$\frac{\frac{(a^4(-B)+4a^3Ab+6a^2b^2B-4aAb^3-b^4B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx + \frac{x(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4)}{a^2+b^2}}{a^2+b^2} - \frac{a^3(-B)+3a^2Ab+3ab^2B-Ab^3}{d(a^2+b^2)(a+b \tan(c+dx))} - \frac{a^2(-B)+2aAb+b^2B}{2d(a^2+b^2)(a+b \tan(c+dx))}}{a^2+b^2} = \frac{Ab - aB}{3d(a^2 + b^2)(a + b \tan(c + dx))^3}$$

↓ 3042

$$\frac{\frac{(a^4(-B)+4a^3Ab+6a^2b^2B-4aAb^3-b^4B) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx + \frac{x(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4)}{a^2+b^2}}{a^2+b^2} - \frac{a^3(-B)+3a^2Ab+3ab^2B-Ab^3}{d(a^2+b^2)(a+b \tan(c+dx))} - \frac{a^2(-B)+2aAb+b^2B}{2d(a^2+b^2)(a+b \tan(c+dx))}}{a^2+b^2} = \frac{Ab - aB}{3d(a^2 + b^2)(a + b \tan(c + dx))^3}$$

↓ 4013

$$\frac{\frac{(a^4(-B)+4a^3Ab+6a^2b^2B-4aAb^3-b^4B) \log(a \cos(c+dx)+b \sin(c+dx)) + \frac{x(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4)}{a^2+b^2}}{d(a^2+b^2)} - \frac{a^3(-B)+3a^2Ab+3ab^2B-Ab^3}{d(a^2+b^2)(a+b \tan(c+dx))} - \frac{a^2(-B)+2aAb+b^2B}{2d(a^2+b^2)(a+b \tan(c+dx))}}{a^2+b^2} = \frac{Ab - aB}{3d(a^2 + b^2)(a + b \tan(c + dx))^3}$$

input `Int[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^4,x]`

output `-1/3*(A*b - a*B)/((a^2 + b^2)*d*(a + b*Tan[c + d*x])^3) + (-1/2*(2*a*A*b - a^2*B + b^2*B)/((a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + (((a^4*A - 6*a^2*A*b^2 + A*b^4 + 4*a^3*b*B - 4*a*b^3*B)*x)/(a^2 + b^2) + ((4*a^3*A*b - 4*a*A*b^3 - a^4*B + 6*a^2*b^2*B - b^4*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d))/(a^2 + b^2) - (3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B)/((a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(a^2 + b^2)/(a^2 + b^2)`

3.294.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4013 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/(a_. + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/(a_. + (b_.)*tan[(e_.) + (f_.)*(x_)])*(x_)), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

3.294.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.15

method	result
derivativedivides	$\frac{(-4Aa^3b+4Aab^3+Ba^4-6Ba^2b^2+Bb^4)}{2} \ln(1+\tan^2(dx+c)) + \frac{(Aa^4-6Aa^2b^2+Ab^4+4Ba^3b-4Bab^3)}{(a^2+b^2)^4} \arctan(\tan(dx+c)) - \frac{3Aa^2}{(a^2+b^2)}$
default	$\frac{(-4Aa^3b+4Aab^3+Ba^4-6Ba^2b^2+Bb^4)}{2} \ln(1+\tan^2(dx+c)) + \frac{(Aa^4-6Aa^2b^2+Ab^4+4Ba^3b-4Bab^3)}{(a^2+b^2)^4} \arctan(\tan(dx+c)) - \frac{3Aa^2}{(a^2+b^2)}$
norman	$\frac{(Aa^4-6Aa^2b^2+Ab^4+4Ba^3b-4Bab^3)a^3x}{(a^6+3a^4b^2+3a^2b^4+b^6)(a^2+b^2)} + \frac{b^3(Aa^4-6Aa^2b^2+Ab^4+4Ba^3b-4Bab^3)x(\tan^3(dx+c))}{(a^6+3a^4b^2+3a^2b^4+b^6)(a^2+b^2)} - \frac{20Aa^4b^3+6Aa^2b^5+2Aa^2b^7}{6b^2d(a^6+3a^4b^2+3a^2b^4+b^6)}$
risch	Expression too large to display
parallelrisch	Expression too large to display

3.294. $\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^4} dx$

input `int((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `1/d*(1/(a^2+b^2)^4*(1/2*(-4*A*a^3*b+4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*ln(1+tan(d*x+c)^2)+(A*a^4-6*A*a^2*b^2+A*b^4+4*B*a^3*b-4*B*a*b^3)*arctan(tan(d*x+c)))-(3*A*a^2*b-A*b^3-B*a^3+3*B*a*b^2)/(a^2+b^2)^3/(a+b*tan(d*x+c))+4*A*a^3*b-4*A*a*b^3-B*a^4+6*B*a^2*b^2-B*b^4)/(a^2+b^2)^4*ln(a+b*tan(d*x+c))-1/3*(A*b-B*a)/(a^2+b^2)/(a+b*tan(d*x+c))^3-1/2*(2*A*a*b-B*a^2+B*b^2)/(a^2+b^2)^2/(a+b*tan(d*x+c))^2)`

3.294.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 815 vs. $2(239) = 478$.

Time = 0.30 (sec) , antiderivative size = 815, normalized size of antiderivative = 3.30

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^4} dx$$

$$= \frac{27 Ba^5 b^2 - 48 Aa^4 b^3 - 18 Ba^3 b^4 - 6 Aa^2 b^5 - Bab^6 - 2 Ab^7 - (11 Ba^4 b^3 - 26 Aa^3 b^4 - 30 Ba^2 b^5 + 18 Aab^6)}{(a + b \tan(c + dx))^4}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="fracas")`

```
output 1/6*(27*B*a^5*b^2 - 48*A*a^4*b^3 - 18*B*a^3*b^4 - 6*A*a^2*b^5 - B*a*b^6 -
2*A*b^7 - (11*B*a^4*b^3 - 26*A*a^3*b^4 - 30*B*a^2*b^5 + 18*A*a*b^6 + 3*B*b
^7 - 6*(A*a^4*b^3 + 4*B*a^3*b^4 - 6*A*a^2*b^5 - 4*B*a*b^6 + A*b^7)*d*x)*ta
n(d*x + c)^3 + 6*(A*a^7 + 4*B*a^6*b - 6*A*a^5*b^2 - 4*B*a^4*b^3 + A*a^3*b^
4)*d*x - 3*(9*B*a^5*b^2 - 20*A*a^4*b^3 - 26*B*a^3*b^4 + 22*A*a^2*b^5 + 9*B
*a*b^6 - 2*A*b^7 - 6*(A*a^5*b^2 + 4*B*a^4*b^3 - 6*A*a^3*b^4 - 4*B*a^2*b^5
+ A*a*b^6)*d*x)*tan(d*x + c)^2 - 3*(B*a^7 - 4*A*a^6*b - 6*B*a^5*b^2 + 4*A*
a^4*b^3 + B*a^3*b^4 + (B*a^4*b^3 - 4*A*a^3*b^4 - 6*B*a^2*b^5 + 4*A*a*b^6 +
B*b^7)*tan(d*x + c)^3 + 3*(B*a^5*b^2 - 4*A*a^4*b^3 - 6*B*a^3*b^4 + 4*A*a^
2*b^5 + B*a*b^6)*tan(d*x + c)^2 + 3*(B*a^6*b - 4*A*a^5*b^2 - 6*B*a^4*b^3 +
4*A*a^3*b^4 + B*a^2*b^5)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*ta
n(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - 3*(6*B*a^6*b - 12*A*a^5*b^2 - 23
*B*a^4*b^3 + 30*A*a^3*b^4 + 16*B*a^2*b^5 - 2*A*a*b^6 + B*b^7 - 6*(A*a^6*b
+ 4*B*a^5*b^2 - 6*A*a^4*b^3 - 4*B*a^3*b^4 + A*a^2*b^5)*d*x)*tan(d*x + c))/
((a^8*b^3 + 4*a^6*b^5 + 6*a^4*b^7 + 4*a^2*b^9 + b^11)*d*tan(d*x + c)^3 + 3
*(a^9*b^2 + 4*a^7*b^4 + 6*a^5*b^6 + 4*a^3*b^8 + a*b^10)*d*tan(d*x + c)^2 +
3*(a^10*b + 4*a^8*b^3 + 6*a^6*b^5 + 4*a^4*b^7 + a^2*b^9)*d*tan(d*x + c) +
(a^11 + 4*a^9*b^2 + 6*a^7*b^4 + 4*a^5*b^6 + a^3*b^8)*d)
```

3.294.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^4} dx = \text{Exception raised: AttributeError}$$

```
input integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))**4,x)
```

```
output Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'
```

3.294.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 514 vs. $2(239) = 478$.

Time = 0.53 (sec) , antiderivative size = 514, normalized size of antiderivative = 2.08

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^4} dx$$

$$= \frac{6(Aa^4 + 4Ba^3b - 6Aa^2b^2 - 4Bab^3 + Ab^4)(dx+c)}{a^8 + 4a^6b^2 + 6a^4b^4 + 4a^2b^6 + b^8} - \frac{6(Ba^4 - 4Aa^3b - 6Ba^2b^2 + 4Aab^3 + Bb^4) \log(b \tan(dx+c) + a)}{a^8 + 4a^6b^2 + 6a^4b^4 + 4a^2b^6 + b^8} + \frac{3(Ba^4 - 4Aa^3b - 6Ba^2b^2)}{a^8 + 4a^6b^2}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="maxima")`

output

$$\frac{1}{6} \cdot \frac{(6(Aa^4 + 4Ba^3b - 6Aa^2b^2 - 4Bab^3 + Ab^4)(dx+c) - 6(Ba^4 - 4Aa^3b - 6Ba^2b^2 + 4Aab^3 + Bb^4) \log(b \tan(dx+c) + a) + 3(Ba^4 - 4Aa^3b - 6Ba^2b^2 + 4Aab^3 + Bb^4) \log(\tan(dx+c)^2 + 1) + 11Ba^5 - 26Aa^4b - 14Ba^3b^2 - 4Aa^2b^3 - Bb^5) \tan(dx+c)^2 + 3(5Ba^4b - 14Aa^3b^2 - 12Ba^2b^3 + 2Aab^4 - Bb^5) \tan(dx+c))}{(a^9 + 3a^7b^2 + 3a^5b^4 + a^3b^6 + (a^6b^3 + 3a^4b^5 + 3a^2b^7 + b^9) \tan(dx+c)^3 + 3(a^7b^2 + 3a^5b^4 + 3a^3b^6 + ab^8) \tan(dx+c)^2 + 3(a^8b + 3a^6b^3 + 3a^4b^5 + a^2b^7) \tan(dx+c))} / d$$
3.294.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 630 vs. $2(239) = 478$.

Time = 0.67 (sec) , antiderivative size = 630, normalized size of antiderivative = 2.55

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^4} dx$$

$$= \frac{6(Aa^4 + 4Ba^3b - 6Aa^2b^2 - 4Bab^3 + Ab^4)(dx+c)}{a^8 + 4a^6b^2 + 6a^4b^4 + 4a^2b^6 + b^8} + \frac{3(Ba^4 - 4Aa^3b - 6Ba^2b^2 + 4Aab^3 + Bb^4) \log(\tan(dx+c)^2 + 1)}{a^8 + 4a^6b^2 + 6a^4b^4 + 4a^2b^6 + b^8} - \frac{6(Ba^4b - 4Aa^3b^2 - 6Ba^2b^3)}{a^8b + 4a^6b^3}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="giac")`

output $\frac{1}{6} \cdot (6 \cdot (A \cdot a^4 + 4 \cdot B \cdot a^3 \cdot b - 6 \cdot A \cdot a^2 \cdot b^2 - 4 \cdot B \cdot a \cdot b^3 + A \cdot b^4) \cdot (d \cdot x + c) / (a^8 + 4 \cdot a^6 \cdot b^2 + 6 \cdot a^4 \cdot b^4 + 4 \cdot a^2 \cdot b^6 + b^8) + 3 \cdot (B \cdot a^4 - 4 \cdot A \cdot a^3 \cdot b - 6 \cdot B \cdot a^2 \cdot b^2 + 4 \cdot A \cdot a \cdot b^3 + B \cdot b^4) \cdot \log(\tan(d \cdot x + c)^2 + 1) / (a^8 + 4 \cdot a^6 \cdot b^2 + 6 \cdot a^4 \cdot b^4 + 4 \cdot a^2 \cdot b^6 + b^8) - 6 \cdot (B \cdot a^4 \cdot b - 4 \cdot A \cdot a^3 \cdot b^2 - 6 \cdot B \cdot a^2 \cdot b^3 + 4 \cdot A \cdot a \cdot b^4 + B \cdot b^5) \cdot \log(\text{abs}(b \cdot \tan(d \cdot x + c) + a)) / (a^8 \cdot b + 4 \cdot a^6 \cdot b^3 + 6 \cdot a^4 \cdot b^5 + 4 \cdot a^2 \cdot b^7 + b^9) + (11 \cdot B \cdot a^4 \cdot b^3 \cdot \tan(d \cdot x + c)^3 - 44 \cdot A \cdot a^3 \cdot b^4 \cdot \tan(d \cdot x + c)^3 - 66 \cdot B \cdot a^2 \cdot b^5 \cdot \tan(d \cdot x + c)^3 + 44 \cdot A \cdot a \cdot b^6 \cdot \tan(d \cdot x + c)^3 + 11 \cdot B \cdot b^7 \cdot \tan(d \cdot x + c)^3 + 39 \cdot B \cdot a^5 \cdot b^2 \cdot \tan(d \cdot x + c)^2 - 150 \cdot A \cdot a^4 \cdot b^3 \cdot \tan(d \cdot x + c)^2 - 210 \cdot B \cdot a^3 \cdot b^4 \cdot \tan(d \cdot x + c)^2 + 120 \cdot A \cdot a^2 \cdot b^5 \cdot \tan(d \cdot x + c)^2 + 15 \cdot B \cdot a \cdot b^6 \cdot \tan(d \cdot x + c)^2 + 6 \cdot A \cdot b^7 \cdot \tan(d \cdot x + c)^2 + 48 \cdot B \cdot a^6 \cdot b \cdot \tan(d \cdot x + c) - 174 \cdot A \cdot a^5 \cdot b^2 \cdot \tan(d \cdot x + c) - 219 \cdot B \cdot a^4 \cdot b^3 \cdot \tan(d \cdot x + c) + 96 \cdot A \cdot a^3 \cdot b^4 \cdot \tan(d \cdot x + c) - 6 \cdot B \cdot a^2 \cdot b^5 \cdot \tan(d \cdot x + c) + 6 \cdot A \cdot a \cdot b^6 \cdot \tan(d \cdot x + c) - 3 \cdot B \cdot b^7 \cdot \tan(d \cdot x + c) + 22 \cdot B \cdot a^7 - 70 \cdot A \cdot a^6 \cdot b - 69 \cdot B \cdot a^5 \cdot b^2 + 14 \cdot A \cdot a^4 \cdot b^3 - 4 \cdot B \cdot a^3 \cdot b^4 - 6 \cdot A \cdot a^2 \cdot b^5 - B \cdot a \cdot b^6 - 2 \cdot A \cdot b^7) / ((a^8 + 4 \cdot a^6 \cdot b^2 + 6 \cdot a^4 \cdot b^4 + 4 \cdot a^2 \cdot b^6 + b^8) \cdot (b \cdot \tan(d \cdot x + c) + a)^3) / d$

3.294.9 Mupad [B] (verification not implemented)

Time = 8.43 (sec) , antiderivative size = 442, normalized size of antiderivative = 1.79

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^4} dx =$$

$$\frac{-11 B a^5 + 26 A a^4 b + 14 B a^3 b^2 + 4 A a^2 b^3 + B a b^4 + 2 A b^5}{6 (a^6 + 3 a^4 b^2 + 3 a^2 b^4 + b^6)} + \frac{\tan(c + dx) (-5 B a^4 b + 14 A a^3 b^2 + 12 B a^2 b^3 - 2 A a b^4 + B b^5)}{2 (a^6 + 3 a^4 b^2 + 3 a^2 b^4 + b^6)} - \frac{\tan(c + dx)^2}{2 (a^6 + 3 a^4 b^2 + 3 a^2 b^4 + b^6)}$$

$$- \frac{d (a^3 + 3 a^2 b \tan(c + dx) + 3 a b^2 \tan(c + dx)^2 + b^3 \tan(c + dx)^3)}{\ln(a + b \tan(c + dx)) \left(\frac{B}{(a^2 + b^2)^2} - \frac{4 b (A a + 2 B b)}{(a^2 + b^2)^3} + \frac{8 b^3 (A a + B b)}{(a^2 + b^2)^4} \right)}$$

$$+ \frac{d \ln(\tan(c + dx) - i) (A + B i)}{2 d (a^4 i - 4 a^3 b - a^2 b^2 6 i + 4 a b^3 + b^4 i)} + \frac{\ln(\tan(c + dx) + i) (B + A i)}{2 d (a^4 - a^3 b 4 i - 6 a^2 b^2 + a b^3 4 i + b^4)}$$

input `int((A + B*tan(c + d*x))/(a + b*tan(c + d*x))^4,x)`

output $(\log(\tan(c + dx) - 1i) \cdot (A + B \cdot 1i)) / (2 \cdot d \cdot (4 \cdot a \cdot b^3 - 4 \cdot a^3 \cdot b + a^4 \cdot 1i + b^4 \cdot 1i - a^2 \cdot b^2 \cdot 6i)) - (\log(a + b \cdot \tan(c + dx)) \cdot (B / (a^2 + b^2)^2 - (4 \cdot b \cdot (A \cdot a + 2 \cdot B \cdot b)) / (a^2 + b^2)^3 + (8 \cdot b^3 \cdot (A \cdot a + B \cdot b)) / (a^2 + b^2)^4)) / d - ((2 \cdot A \cdot b^5 - 11 \cdot B \cdot a^5 + 4 \cdot A \cdot a^2 \cdot b^3 + 14 \cdot B \cdot a^3 \cdot b^2 + 26 \cdot A \cdot a^4 \cdot b + B \cdot a \cdot b^4) / (6 \cdot (a^6 + b^6 + 3 \cdot a^2 \cdot b^4 + 3 \cdot a^4 \cdot b^2))) + (\tan(c + dx) \cdot (B \cdot b^5 + 14 \cdot A \cdot a^3 \cdot b^2 + 12 \cdot B \cdot a^2 \cdot b^3 - 2 \cdot A \cdot a \cdot b^4 - 5 \cdot B \cdot a^4 \cdot b)) / (2 \cdot (a^6 + b^6 + 3 \cdot a^2 \cdot b^4 + 3 \cdot a^4 \cdot b^2)) - (\tan(c + dx)^2 \cdot (A \cdot b^5 - 3 \cdot A \cdot a^2 \cdot b^3 + B \cdot a^3 \cdot b^2 - 3 \cdot B \cdot a \cdot b^4)) / (a^6 + b^6 + 3 \cdot a^2 \cdot b^4 + 3 \cdot a^4 \cdot b^2)) / (d \cdot (a^3 + b^3 \cdot \tan(c + dx)^3 + 3 \cdot a \cdot b^2 \cdot \tan(c + dx)^2 + 3 \cdot a^2 \cdot b \cdot \tan(c + dx))) + (\log(\tan(c + dx) + 1i) \cdot (A \cdot 1i + B)) / (2 \cdot d \cdot (a \cdot b^3 \cdot 4i - a^3 \cdot b \cdot 4i + a^4 + b^4 - 6 \cdot a^2 \cdot b^2))$

3.295 $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

3.295.1 Optimal result 2839
 3.295.2 Mathematica [C] (verified) 2840
 3.295.3 Rubi [A] (verified) 2840
 3.295.4 Maple [A] (verified) 2845
 3.295.5 Fricas [B] (verification not implemented) 2845
 3.295.6 Sympy [F(-2)] 2846
 3.295.7 Maxima [A] (verification not implemented) 2847
 3.295.8 Giac [B] (verification not implemented) 2847
 3.295.9 Mupad [B] (verification not implemented) 2848

3.295.1 Optimal result

Integrand size = 29, antiderivative size = 302

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

$$= -\frac{(4a^3Ab - 4aAb^3 - a^4B + 6a^2b^2B - b^4B)x}{(a^2 + b^2)^4} + \frac{A \log(\sin(c+dx))}{a^4d}$$

$$- \frac{b(10a^6Ab + 5a^4Ab^3 + 4a^2Ab^5 + Ab^7 - 4a^7B + 4a^5b^2B) \log(a \cos(c+dx) + b \sin(c+dx))}{a^4(a^2 + b^2)^4d}$$

$$+ \frac{b(Ab - aB)}{3a(a^2 + b^2)d(a + b \tan(c+dx))^3} + \frac{b(3a^2Ab + Ab^3 - 2a^3B)}{2a^2(a^2 + b^2)^2d(a + b \tan(c+dx))^2}$$

$$+ \frac{b(6a^4Ab + 3a^2Ab^3 + Ab^5 - 3a^5B + a^3b^2B)}{a^3(a^2 + b^2)^3d(a + b \tan(c+dx))}$$

```
output - (4*A*a^3*b-4*A*a*b^3-B*a^4+6*B*a^2*b^2-B*b^4)*x/(a^2+b^2)^4+A*ln(sin(d*x+c))
/a^4/d-b*(10*A*a^6*b+5*A*a^4*b^3+4*A*a^2*b^5+A*b^7-4*B*a^7+4*B*a^5*b^2)
*ln(a*cos(d*x+c)+b*sin(d*x+c))/a^4/(a^2+b^2)^4/d+1/3*b*(A*b-B*a)/a/(a^2+b^2)
/d/(a+b*tan(d*x+c))^3+1/2*b*(3*A*a^2*b+A*b^3-2*B*a^3)/a^2/(a^2+b^2)^2/d/
(a+b*tan(d*x+c))^2+b*(6*A*a^4*b+3*A*a^2*b^3+A*b^5-3*B*a^5+B*a^3*b^2)/a^3/(
a^2+b^2)^3/d/(a+b*tan(d*x+c))
```

3.295.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.54 (sec) , antiderivative size = 308, normalized size of antiderivative = 1.02

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^4} dx$$

$$= \frac{3(-a^4(a-ib)^4(A+iB) \log(i-\tan(c+dx))+2A(a^2+b^2)^4 \log(\tan(c+dx))-a^4(a+ib)^4(A-iB) \log(i+\tan(c+dx))-2b(10a^6 Ab+5a^4 Ab^3+4a^2 Ab^5+A^2 b^7))}{a^2(a^2+b^2)^2} + \frac{6a^2(a^2-b^2)}{a^2(a^2+b^2)^2}$$

input `Integrate[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]`

output `((3*(-(a^4*(a - I*b)^4*(A + I*B)*Log[I - Tan[c + d*x]]) + 2*A*(a^2 + b^2)^4*Log[Tan[c + d*x]] - a^4*(a + I*b)^4*(A - I*B)*Log[I + Tan[c + d*x]] - 2*b*(10*a^6*A*b + 5*a^4*A*b^3 + 4*a^2*A*b^5 + A*b^7 - 4*a^7*B + 4*a^5*b^2*B)*Log[a + b*Tan[c + d*x]]))/(a^2*(a^2 + b^2)^2) + (2*a*b*(a^2 + b^2)*(A*b - a*B))/(a + b*Tan[c + d*x])^3 + (3*b*(3*a^2*A*b + A*b^3 - 2*a^3*B))/(a + b*Tan[c + d*x])^2 + (6*b*(6*a^4*A*b + 3*a^2*A*b^3 + A*b^5 - 3*a^5*B + a^3*b^2*B))/(a*(a^2 + b^2)*(a + b*Tan[c + d*x]))/(6*a^2*(a^2 + b^2)^2*d)`

3.295.3 Rubi [A] (verified)

Time = 1.89 (sec) , antiderivative size = 358, normalized size of antiderivative = 1.19, number of steps used = 14, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.483$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^4} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)(a + b \tan(c + dx))^4} dx$$

↓ 4092

$$\frac{\int \frac{3 \cot(c+dx)(b(Ab-aB) \tan^2(c+dx)-a(Ab-aB) \tan(c+dx)+A(a^2+b^2))}{(a+b \tan(c+dx))^3} dx}{3a(a^2+b^2)} + \frac{b(Ab-aB)}{3ad(a^2+b^2)(a+b \tan(c+dx))^3}$$

3.295. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

$$\begin{aligned}
& \int \frac{\cot(c+dx)(b(Ab-aB)\tan^2(c+dx)-a(Ab-aB)\tan(c+dx)+A(a^2+b^2))}{a(a^2+b^2)(a+b\tan(c+dx))^3} dx + \frac{b(Ab-aB)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
& \quad \downarrow 27 \\
& \int \frac{b(Ab-aB)\tan(c+dx)^2-a(Ab-aB)\tan(c+dx)+A(a^2+b^2)}{\tan(c+dx)(a+b\tan(c+dx))^3} dx + \frac{b(Ab-aB)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
& \quad \downarrow 3042 \\
& \int \frac{2\cot(c+dx)\left(-((-Ba^2+2Aba+b^2B)\tan(c+dx)a^2)+A(a^2+b^2)^2+b(-2Ba^3+3Aba^2+Ab^3)\tan^2(c+dx)\right)}{\frac{(a+b\tan(c+dx))^2}{2a(a^2+b^2)}} dx + \frac{b(-2a^3B+3a^2Ab+Ab^3)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2} + \\
& \quad \frac{a(a^2+b^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
& \quad \downarrow 27 \\
& \int \frac{\cot(c+dx)\left(-((-Ba^2+2Aba+b^2B)\tan(c+dx)a^2)+A(a^2+b^2)^2+b(-2Ba^3+3Aba^2+Ab^3)\tan^2(c+dx)\right)}{\frac{(a+b\tan(c+dx))^2}{a(a^2+b^2)}} dx + \frac{b(-2a^3B+3a^2Ab+Ab^3)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2} + \\
& \quad \frac{a(a^2+b^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
& \quad \downarrow 3042 \\
& \int \frac{-((-Ba^2+2Aba+b^2B)\tan(c+dx)a^2)+A(a^2+b^2)^2+b(-2Ba^3+3Aba^2+Ab^3)\tan(c+dx)^2}{\frac{\tan(c+dx)(a+b\tan(c+dx))^2}{a(a^2+b^2)}} dx + \frac{b(-2a^3B+3a^2Ab+Ab^3)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2} + \\
& \quad \frac{a(a^2+b^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
& \quad \downarrow 4132 \\
& \int \frac{\cot(c+dx)\left(-((-Ba^3+3Aba^2+3b^2Ba-Ab^3)\tan(c+dx)a^3)+A(a^2+b^2)^3+b(-3Ba^5+6Aba^4+b^2Ba^3+3Ab^3a^2+Ab^5)\tan^2(c+dx)\right)}{\frac{a+b\tan(c+dx)}{a(a^2+b^2)}} dx + \frac{b(-3a^5B+6a^4Ab+a^3B)}{ad(a^2+b^2)(a+b\tan(c+dx))} + \\
& \quad \frac{a(a^2+b^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
& \quad \downarrow 3042
\end{aligned}$$

3.295. $\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$

$$\frac{\int \frac{-((-Ba^3+3Aba^2+3b^2Ba-Ab^3)\tan(c+dx)a^3)+A(a^2+b^2)^3+b(-3Ba^5+6Aba^4+b^2Ba^3+3Ab^3a^2+Ab^5)\tan(c+dx)^2}{\tan(c+dx)(a+b\tan(c+dx))} dx}{a(a^2+b^2)} + \frac{b(-3a^5B+6a^4Ab+a^3b^2B+3a^2Ab^3)}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

$$\frac{b(Ab-aB)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3}$$

↓ 4134

$$\frac{A(a^2+b^2)^3 \int \cot(c+dx) dx}{a} - \frac{b(-4a^7B+10a^6Ab+4a^5b^2B+5a^4Ab^3+4a^2Ab^5+Ab^7) \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^3x(a^4(-B)+4a^3Ab+6a^2b^2B-4aAb^3-b^4B)}{a^2+b^2} + \frac{b(-3a^5B+6a^4Ab+a^3b^2B+3a^2Ab^3)}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

$$\frac{b(Ab-aB)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3}$$

↓ 3042

$$\frac{A(a^2+b^2)^3 \int -\tan(c+dx+\frac{\pi}{2}) dx}{a} - \frac{b(-4a^7B+10a^6Ab+4a^5b^2B+5a^4Ab^3+4a^2Ab^5+Ab^7) \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^3x(a^4(-B)+4a^3Ab+6a^2b^2B-4aAb^3-b^4B)}{a^2+b^2} + \frac{b(-3a^5B+6a^4Ab+a^3b^2B+3a^2Ab^3)}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

$$\frac{b(Ab-aB)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3}$$

↓ 25

$$\frac{A(a^2+b^2)^3 \int \tan(\frac{1}{2}(2c+\pi)+dx) dx}{a} - \frac{b(-4a^7B+10a^6Ab+4a^5b^2B+5a^4Ab^3+4a^2Ab^5+Ab^7) \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^3x(a^4(-B)+4a^3Ab+6a^2b^2B-4aAb^3-b^4B)}{a^2+b^2} + \frac{b(-3a^5B+6a^4Ab+a^3b^2B+3a^2Ab^3)}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

$$\frac{b(Ab-aB)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3}$$

↓ 3956

$$\frac{b(-4a^7B+10a^6Ab+4a^5b^2B+5a^4Ab^3+4a^2Ab^5+Ab^7) \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} + \frac{A(a^2+b^2)^3 \log(-\sin(c+dx))}{ad} - \frac{a^3x(a^4(-B)+4a^3Ab+6a^2b^2B-4aAb^3-b^4B)}{a^2+b^2} + \frac{b(-3a^5B+6a^4Ab+a^3b^2B+3a^2Ab^3)}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

$$\frac{b(Ab-aB)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3}$$

↓ 4013

3.295. $\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$

$$\frac{b(Ab - aB)}{3ad(a^2 + b^2)(a + b \tan(c + dx))^3} + \frac{A(a^2 + b^2)^3 \log(-\sin(c + dx)) - a^3 x(a^4(-B) + 4a^3 Ab + 6a^2 b^2 B - 4aAb^3 - b^4 B)}{ad} + \frac{b(-3a^5 B + 6a^4 Ab + a^3 b^2 B + 3a^2 Ab^3 + Ab^5)}{ad(a^2 + b^2)(a + b \tan(c + dx))} + \frac{b(-2a^3 B + 3a^2 Ab + Ab^3)}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2} + \frac{a(a^2 + b^2)}{a(a^2 + b^2)}$$

```
input Int[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]
```

```
output (b*(A*b - a*B))/(3*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^3) + ((b*(3*a^2*A*b + A*b^3 - 2*a^3*B))/(2*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + (((-((a^3*(4*a^3*A*b - 4*a*A*b^3 - a^4*B + 6*a^2*b^2*B - b^4*B)*x)/(a^2 + b^2)) + (A*(a^2 + b^2)^3*Log[-Sin[c + d*x]])/(a*d) - (b*(10*a^6*A*b + 5*a^4*A*b^3 + 4*a^2*A*b^5 + A*b^7 - 4*a^7*B + 4*a^5*b^2*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))/(a*(a^2 + b^2)) + (b*(6*a^4*A*b + 3*a^2*A*b^3 + A*b^5 - 3*a^5*B + a^3*b^2*B))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(a*(a^2 + b^2)))/(a*(a^2 + b^2))
```

3.295.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3956 Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 4013 Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

3.295. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4134 `Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/(a^2 + b^2)*(c^2 + d^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.295.4 Maple [A] (verified)

Time = 0.68 (sec) , antiderivative size = 335, normalized size of antiderivative = 1.11

method	result
derivativedivides	$\frac{(-A a^4 + 6A a^2 b^2 - A b^4 - 4B a^3 b + 4B a b^3) \ln(1 + \tan^2(dx+c)) + (-4A a^3 b + 4A a b^3 + B a^4 - 6B a^2 b^2 + B b^4) \arctan(\tan(dx+c))}{(a^2 + b^2)^4} + A \ln$
default	$\frac{(-A a^4 + 6A a^2 b^2 - A b^4 - 4B a^3 b + 4B a b^3) \ln(1 + \tan^2(dx+c)) + (-4A a^3 b + 4A a b^3 + B a^4 - 6B a^2 b^2 + B b^4) \arctan(\tan(dx+c))}{(a^2 + b^2)^4} + A \ln$
parallelrisch	$-20(A a^6 b + \frac{1}{2} A a^4 b^3 + \frac{2}{5} A a^2 b^5 + \frac{1}{10} A b^7 - \frac{2}{5} B a^7 + \frac{2}{5} B a^5 b^2) b(a + b \tan(dx+c))^3 \ln(a + b \tan(dx+c)) - a^4(a + b \tan(dx+c))$
norman	$-\frac{(4A a^3 b - 4A a b^3 - B a^4 + 6B a^2 b^2 - B b^4) a^3 x}{(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)(a^2 + b^2)} - \frac{b^3(4A a^3 b - 4A a b^3 - B a^4 + 6B a^2 b^2 - B b^4) x (\tan^3(dx+c))}{(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)(a^2 + b^2)} - \frac{b(10A a^4 b^2 + 9A a^2 b^4 + 10A a^2 b^4 + 9A a^2 b^4)}{a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6}$
risch	Expression too large to display

input `int(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x,method=_RETURNVERBOSE)`

output `1/d*(1/(a^2+b^2)^4*(1/2*(-A*a^4+6*A*a^2*b^2-A*b^4-4*B*a^3*b+4*B*a*b^3)*ln(1+tan(d*x+c)^2)+(-4*A*a^3*b+4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*arctan(tan(d*x+c)))+1/a^4*A*ln(tan(d*x+c))+1/2*b*(3*A*a^2*b+A*b^3-2*B*a^3)/(a^2+b^2)^2/a^2/(a+b*tan(d*x+c))^2+b*(6*A*a^4*b+3*A*a^2*b^3+A*b^5-3*B*a^5+B*a^3*b^2)/(a^2+b^2)^3/a^3/(a+b*tan(d*x+c))-b*(10*A*a^6*b+5*A*a^4*b^3+4*A*a^2*b^5+A*b^7-4*B*a^7+4*B*a^5*b^2)/(a^2+b^2)^4/a^4*ln(a+b*tan(d*x+c))+1/3*(A*b-B*a)*b/(a^2+b^2)/a/(a+b*tan(d*x+c))^3)`

3.295.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1126 vs. 2(299) = 598.

Time = 0.38 (sec) , antiderivative size = 1126, normalized size of antiderivative = 3.73

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^4} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="fricas")`

3.295.
$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

```
output -1/6*(48*B*a^8*b^3 - 75*A*a^7*b^4 + 6*B*a^6*b^5 - 42*A*a^5*b^6 + 2*B*a^4*b^7 - 11*A*a^3*b^8 - (26*B*a^7*b^4 - 47*A*a^6*b^5 - 18*B*a^5*b^6 - 6*A*a^4*b^7 - 3*A*a^2*b^9 + 6*(B*a^8*b^3 - 4*A*a^7*b^4 - 6*B*a^6*b^5 + 4*A*a^5*b^6 + B*a^4*b^7)*d*x)*tan(d*x + c)^3 - 6*(B*a^11 - 4*A*a^10*b - 6*B*a^9*b^2 + 4*A*a^8*b^3 + B*a^7*b^4)*d*x - 3*(20*B*a^8*b^3 - 35*A*a^7*b^4 - 22*B*a^6*b^5 + 12*A*a^5*b^6 + 2*B*a^4*b^7 + 5*A*a^3*b^8 + 2*A*a*b^10 + 6*(B*a^9*b^2 - 4*A*a^8*b^3 - 6*B*a^7*b^4 + 4*A*a^6*b^5 + B*a^5*b^6)*d*x)*tan(d*x + c)^2 - 3*(A*a^11 + 4*A*a^9*b^2 + 6*A*a^7*b^4 + 4*A*a^5*b^6 + A*a^3*b^8 + (A*a^8*b^3 + 4*A*a^6*b^5 + 6*A*a^4*b^7 + 4*A*a^2*b^9 + A*b^11)*tan(d*x + c)^3 + 3*(A*a^9*b^2 + 4*A*a^7*b^4 + 6*A*a^5*b^6 + 4*A*a^3*b^8 + A*a*b^10)*tan(d*x + c)^2 + 3*(A*a^10*b + 4*A*a^8*b^3 + 6*A*a^6*b^5 + 4*A*a^4*b^7 + A*a^2*b^9)*tan(d*x + c))*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1)) - 3*(4*B*a^10*b - 10*A*a^9*b^2 - 4*B*a^8*b^3 - 5*A*a^7*b^4 - 4*A*a^5*b^6 - A*a^3*b^8 + (4*B*a^7*b^4 - 10*A*a^6*b^5 - 4*B*a^5*b^6 - 5*A*a^4*b^7 - 4*A*a^2*b^9 - A*b^11)*tan(d*x + c)^3 + 3*(4*B*a^8*b^3 - 10*A*a^7*b^4 - 4*B*a^6*b^5 - 5*A*a^5*b^6 - 4*A*a^3*b^8 - A*a*b^10)*tan(d*x + c)^2 + 3*(4*B*a^9*b^2 - 10*A*a^8*b^3 - 4*B*a^7*b^4 - 5*A*a^6*b^5 - 4*A*a^4*b^7 - A*a^2*b^9)*tan(d*x + c))*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - 3*(12*B*a^9*b^2 - 20*A*a^8*b^3 - 30*B*a^7*b^4 + 37*A*a^6*b^5 + 2*B*a^5*b^6 + 18*A*a^4*b^7 + 5*A*a^2*b^9 + 6*(B*a^10*b - 4*A*a^9*b^2 - 6*B*a^8*b...
```

3.295.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx = \text{Exception raised: AttributeError}$$

```
input integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**4,x)
```

```
output Exception raised: AttributeError >> 'NoneType' object has no attribute 'primitive'
```

3.295.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 580, normalized size of antiderivative = 1.92

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

$$= \frac{6(Ba^4-4Aa^3b-6Ba^2b^2+4Aab^3+Bb^4)(dx+c)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} + \frac{6(4Ba^7b-10Aa^6b^2-4Ba^5b^3-5Aa^4b^4-4Aa^2b^6-Ab^8) \log(b \tan(dx+c)+a)}{a^{12}+4a^{10}b^2+6a^8b^4+4a^6b^6+a^4b^8} - \frac{3(Aa^4+4Aa^3b-6Aa^2b^2+4Aab^3+Bb^4)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="maxima")`

output `1/6*(6*(B*a^4 - 4*A*a^3*b - 6*B*a^2*b^2 + 4*A*a*b^3 + B*b^4)*(d*x + c)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + 6*(4*B*a^7*b - 10*A*a^6*b^2 - 4*B*a^5*b^3 - 5*A*a^4*b^4 - 4*A*a^2*b^6 - A*b^8)*log(b*tan(d*x + c) + a))/(a^12 + 4*a^10*b^2 + 6*a^8*b^4 + 4*a^6*b^6 + a^4*b^8) - 3*(A*a^4 + 4*B*a^3*b - 6*A*a^2*b^2 - 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)^2 + 1)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) - (26*B*a^7*b - 47*A*a^6*b^2 + 4*B*a^5*b^3 - 34*A*a^4*b^4 + 2*B*a^3*b^5 - 11*A*a^2*b^6 + 6*(3*B*a^5*b^3 - 6*A*a^4*b^4 - B*a^3*b^5 - 3*A*a^2*b^6 - A*b^8))*tan(d*x + c)^2 + 3*(14*B*a^6*b^2 - 27*A*a^5*b^3 - 2*B*a^4*b^4 - 16*A*a^3*b^5 - 5*A*a*b^7)*tan(d*x + c))/(a^12 + 3*a^10*b^2 + 3*a^8*b^4 + a^6*b^6 + (a^9*b^3 + 3*a^7*b^5 + 3*a^5*b^7 + a^3*b^9)*tan(d*x + c)^3 + 3*(a^10*b^2 + 3*a^8*b^4 + 3*a^6*b^6 + a^4*b^8)*tan(d*x + c)^2 + 3*(a^11*b + 3*a^9*b^3 + 3*a^7*b^5 + a^5*b^7)*tan(d*x + c)) + 6*A*log(tan(d*x + c))/a^4)/d`

3.295.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 722 vs. 2(299) = 598.

Time = 0.99 (sec) , antiderivative size = 722, normalized size of antiderivative = 2.39

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

$$= \frac{6(Ba^4-4Aa^3b-6Ba^2b^2+4Aab^3+Bb^4)(dx+c)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} - \frac{3(Aa^4+4Ba^3b-6Aa^2b^2-4Bab^3+Ab^4) \log(\tan(dx+c)^2+1)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} + \frac{6(4Ba^7b^2-10Aa^6b^3-4Aa^5b^4+4Aa^3b^5+Bb^6)}{a^{12}+4a^{10}b^2+6a^8b^4+4a^6b^6+a^4b^8}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="giac")`

3.295. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

output

$$\frac{1}{6} \frac{(6(Ba^4 - 4Aa^3b - 6B^2a^2b^2 + 4A^2ab^3 + B^2b^4)(dx + c)/(a^8 + 4a^6b^2 + 6a^4b^4 + 4a^2b^6 + b^8) - 3(Aa^4 + 4B^3a^3b - 6A^2a^2b^2 - 4B^2ab^3 + Ab^4) \log(\tan(dx + c)^2 + 1)/(a^8 + 4a^6b^2 + 6a^4b^4 + 4a^2b^6 + b^8) + 6(4B^7a^7b^2 - 10A^6a^6b^3 - 4B^5a^5b^4 - 5A^4a^4b^5 - 4A^2a^2b^7 - Ab^9) \log(\operatorname{abs}(b \tan(dx + c) + a))/(a^{12}b + 4a^{10}b^3 + 6a^8b^5 + 4a^6b^7 + a^4b^9) + 6A \log(\operatorname{abs}(\tan(dx + c)))}{a^4 - (44B^7a^7b^4 \tan(dx + c)^3 - 110A^6a^6b^5 \tan(dx + c)^3 - 44B^5a^5b^6 \tan(dx + c)^3 - 55A^4a^4b^7 \tan(dx + c)^3 - 44A^2a^2b^9 \tan(dx + c)^3 - 11Ab^{11} \tan(dx + c)^3 + 150B^8a^8b^3 \tan(dx + c)^2 - 366A^7a^7b^4 \tan(dx + c)^2 - 120B^6a^6b^5 \tan(dx + c)^2 - 219A^5a^5b^6 \tan(dx + c)^2 - 6B^4a^4b^7 \tan(dx + c)^2 - 156A^3a^3b^8 \tan(dx + c)^2 - 39A^2a^2b^{10} \tan(dx + c)^2 + 174B^9a^9b^2 \tan(dx + c) - 411A^8a^8b^3 \tan(dx + c) - 96B^7a^7b^4 \tan(dx + c) - 294A^6a^6b^5 \tan(dx + c) - 6B^5a^5b^6 \tan(dx + c) - 195A^4a^4b^7 \tan(dx + c) - 48A^2a^2b^9 \tan(dx + c) + 70B^10a^{10}b - 157A^9a^9b^2 - 14B^8a^8b^3 - 136A^7a^7b^4 + 6B^6a^6b^5 - 89A^5a^5b^6 + 2B^4a^4b^7 - 22A^3a^3b^8)/(a^{12} + 4a^{10}b^2 + 6a^8b^4 + 4a^6b^6 + a^4b^8)(b \tan(dx + c) + a)^3)/d$$

3.295.9 Mupad [B] (verification not implemented)

Time = 11.61 (sec) , antiderivative size = 484, normalized size of antiderivative = 1.60

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^4} dx$$

$$= \frac{\frac{-26Ba^5b + 47Aa^4b^2 - 4Ba^3b^3 + 34Aa^2b^4 - 2Bab^5 + 11Ab^6}{6a(a^6 + 3a^4b^2 + 3a^2b^4 + b^6)} + \frac{\tan(c + dx)^2(-3Ba^5b^3 + 6Aa^4b^4 + Ba^3b^5 + 3Aa^2b^6 + Ab^8)}{a^3(a^6 + 3a^4b^2 + 3a^2b^4 + b^6)} + \frac{\tan(c + dx)}{a^3(a^6 + 3a^4b^2 + 3a^2b^4 + b^6)}}{d(a^3 + 3a^2b \tan(c + dx) + 3ab^2 \tan(c + dx)^2 + b^3 \tan(c + dx)^3)} + \frac{A \ln(\tan(c + dx))}{a^4 d} - \frac{\ln(\tan(c + dx) + i)(B + A i)}{2d(a^4 i + 4a^3 b - a^2 b^2 6i - 4ab^3 + b^4 i)} - \frac{\ln(\tan(c + dx) - i)(A + B i)}{2d(a^4 + a^3 b 4i - 6a^2 b^2 - a b^3 4i + b^4)} - \frac{b \ln(a + b \tan(c + dx))(-4Ba^7 + 10Aa^6 b + 4Ba^5 b^2 + 5Aa^4 b^3 + 4Aa^2 b^5 + Ab^7)}{a^4 d(a^2 + b^2)^4}$$

input `int((cot(c + d*x)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^4,x)`

output
$$\begin{aligned} & ((11Ab^6 + 34A^2b^4 + 47A^4b^2 - 4B^3b^3 - 2B^5b^5 - 26B^7b^7) / (6a(a^6 + b^6 + 3a^2b^4 + 3a^4b^2)) + (\tan(c + dx)^2(Ab^8 + 3A^2b^6 + 6A^4b^4 + B^3b^5 - 3B^5b^3)) / (a^3(a^6 + b^6 + 3a^2b^4 + 3a^4b^2)) + (\tan(c + dx)(5Ab^7 + 16A^2b^5 + 27A^4b^3 + 2B^3b^4 - 14B^5b^2)) / (2a^2(a^6 + b^6 + 3a^2b^4 + 3a^4b^2))) / (d(a^3 + b^3 \tan(c + dx)^3 + 3ab^2 \tan(c + dx)^2 + 3a^2b \tan(c + dx))) + (A \log(\tan(c + dx))) / (a^4 d) - (\log(\tan(c + dx) + 1i)(A + B)) / (2d(4a^3b - 4ab^3 + a^4 1i + b^4 1i - a^2 b^2 6i)) - (\log(\tan(c + dx) - 1i)(A + B 1i)) / (2d(a^3 b 4i - ab^3 4i + a^4 + b^4 - 6a^2 b^2)) - (b \log(a + b \tan(c + dx))(Ab^7 - 4B^3b^4 + 4A^2b^5 + 5A^4b^3 + 4B^5b^2 + 10A^6b)) / (a^4 d (a^2 + b^2)^4) \end{aligned}$$

3.296 $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

3.296.1 Optimal result 2850
 3.296.2 Mathematica [C] (verified) 2851
 3.296.3 Rubi [A] (verified) 2851
 3.296.4 Maple [A] (verified) 2857
 3.296.5 Fricas [B] (verification not implemented) 2857
 3.296.6 Sympy [F(-2)] 2858
 3.296.7 Maxima [A] (verification not implemented) 2859
 3.296.8 Giac [B] (verification not implemented) 2859
 3.296.9 Mupad [B] (verification not implemented) 2860

3.296.1 Optimal result

Integrand size = 31, antiderivative size = 399

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

$$= -\frac{(a^4 A - 6a^2 Ab^2 + Ab^4 + 4a^3 bB - 4ab^3 B)x}{(a^2 + b^2)^4} - \frac{(4Ab - aB) \log(\sin(c+dx))}{a^5 d}$$

$$+ \frac{b^2(20a^6 Ab + 24a^4 Ab^3 + 16a^2 Ab^5 + 4Ab^7 - 10a^7 B - 5a^5 b^2 B - 4a^3 b^4 B - ab^6 B) \log(a \cos(c+dx) + b \sin(c+dx))}{a^5 (a^2 + b^2)^4 d}$$

$$- \frac{b(3a^2 A + 4Ab^2 - abB)}{3a^2 (a^2 + b^2) d(a+b \tan(c+dx))^3} - \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3}$$

$$- \frac{b(2a^4 A + 8a^2 Ab^2 + 4Ab^4 - 3a^3 bB - ab^3 B)}{2a^3 (a^2 + b^2)^2 d(a+b \tan(c+dx))^2}$$

$$- \frac{b(a^6 A + 13a^4 Ab^2 + 12a^2 Ab^4 + 4Ab^6 - 6a^5 bB - 3a^3 b^3 B - ab^5 B)}{a^4 (a^2 + b^2)^3 d(a+b \tan(c+dx))}$$

output

```
-(A*a^4-6*A*a^2*b^2+A*b^4+4*B*a^3*b-4*B*a*b^3)*x/(a^2+b^2)^4-(4*A*b-B*a)*ln(sin(dx+c))/a^5/d+b^2*(20*A*a^6*b+24*A*a^4*b^3+16*A*a^2*b^5+4*A*b^7-10*B*a^7-5*B*a^5*b^2-4*B*a^3*b^4-B*a*b^6)*ln(a*cos(dx+c)+b*sin(dx+c))/a^5/(a^2+b^2)^4/d-1/3*b*(3*A*a^2+4*A*b^2-B*a*b)/a^2/(a^2+b^2)/d/(a+b*tan(dx+c))^3-A*cot(dx+c)/a/d/(a+b*tan(dx+c))^3-1/2*b*(2*A*a^4+8*A*a^2*b^2+4*A*b^4-3*B*a^3*b-B*a*b^3)/a^3/(a^2+b^2)^2/d/(a+b*tan(dx+c))^2-b*(A*a^6+13*A*a^4*b^2+12*A*a^2*b^4+4*A*b^6-6*B*a^5*b-3*B*a^3*b^3-B*a*b^5)/a^4/(a^2+b^2)^3/d/(a+b*tan(dx+c))
```

3.296.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.58 (sec) , antiderivative size = 357, normalized size of antiderivative = 0.89

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

$$= \frac{-\frac{6A \cot(c+dx)}{a^4} + \frac{3i(A+iB) \log(i-\tan(c+dx))}{(a+ib)^4} + \frac{6(-4Ab+aB) \log(\tan(c+dx))}{a^5} - \frac{3(iA+B) \log(i+\tan(c+dx))}{(a-ib)^4} - \frac{6b^2(-20a^6Ab-24a^4Ab)}{a^5}}$$

input `Integrate[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]`

output $((-6*A*\text{Cot}[c + d*x])/a^4 + ((3*I)*(A + I*B)*\text{Log}[I - \text{Tan}[c + d*x]])/(a + I*b)^4 + (6*(-4*A*b + a*B)*\text{Log}[\text{Tan}[c + d*x]])/a^5 - (3*(I*A + B)*\text{Log}[I + \text{Tan}[c + d*x]])/(a - I*b)^4 - (6*b^2*(-20*a^6*A*b - 24*a^4*A*b^3 - 16*a^2*A*b^5 - 4*A*b^7 + 10*a^7*B + 5*a^5*b^2*B + 4*a^3*b^4*B + a*b^6*B)*\text{Log}[a + b*\text{Tan}[c + d*x]])/(a^5*(a^2 + b^2)^4) + (2*b^2*(-(A*b) + a*B))/(a^2*(a^2 + b^2)*(a + b*\text{Tan}[c + d*x])^3) + (3*b^2*(-4*a^2*A*b - 2*A*b^3 + 3*a^3*B + a*b^2*B))/(a^3*(a^2 + b^2)^2*(a + b*\text{Tan}[c + d*x])^2) + (6*b^2*(-10*a^4*A*b - 9*a^2*A*b^3 - 3*A*b^5 + 6*a^5*B + 3*a^3*b^2*B + a*b^4*B))/(a^4*(a^2 + b^2)^3*(a + b*\text{Tan}[c + d*x]))/(6*d)$

3.296.3 Rubi [A] (verified)

Time = 2.47 (sec) , antiderivative size = 459, normalized size of antiderivative = 1.15, number of steps used = 16, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.516$, Rules used = {3042, 4092, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A+B \tan(c+dx)}{\tan(c+dx)^2(a+b \tan(c+dx))^4} dx$$

$$\downarrow \text{4092}$$

3.296. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

$$\begin{aligned}
 & \frac{\int \frac{\cot(c+dx)(4Ab \tan^2(c+dx)+aA \tan(c+dx)+4Ab-aB)}{(a+b \tan(c+dx))^4} dx}{a} - \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{4Ab \tan(c+dx)^2+aA \tan(c+dx)+4Ab-aB}{\tan(c+dx)(a+b \tan(c+dx))^4} dx}{a} - \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3} \\
 & \quad \downarrow \text{4132} \\
 & \frac{\int \frac{3 \cot(c+dx)((aA+bB) \tan(c+dx)a^2+b(3Aa^2-bBa+4Ab^2) \tan^2(c+dx)+(a^2+b^2)(4Ab-aB))}{(a+b \tan(c+dx))^3} dx}{3a(a^2+b^2)} + \frac{b(3a^2A-abB+4Ab^2)}{3ad(a^2+b^2)(a+b \tan(c+dx))^3} \\
 & \quad \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\cot(c+dx)((aA+bB) \tan(c+dx)a^2+b(3Aa^2-bBa+4Ab^2) \tan^2(c+dx)+(a^2+b^2)(4Ab-aB))}{(a+b \tan(c+dx))^3} dx}{a(a^2+b^2)} + \frac{b(3a^2A-abB+4Ab^2)}{3ad(a^2+b^2)(a+b \tan(c+dx))^3} \\
 & \quad \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{(aA+bB) \tan(c+dx)a^2+b(3Aa^2-bBa+4Ab^2) \tan(c+dx)^2+(a^2+b^2)(4Ab-aB)}{\tan(c+dx)(a+b \tan(c+dx))^3} dx}{a(a^2+b^2)} + \frac{b(3a^2A-abB+4Ab^2)}{3ad(a^2+b^2)(a+b \tan(c+dx))^3} \\
 & \quad \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3} \\
 & \quad \downarrow \text{4132} \\
 & \frac{\int \frac{2 \cot(c+dx)((Aa^2+2bBa-Ab^2) \tan(c+dx)a^3+b(2Aa^4-3bBa^3+8Ab^2a^2-b^3Ba+4Ab^4) \tan^2(c+dx)+(a^2+b^2)^2(4Ab-aB))}{(a+b \tan(c+dx))^2} dx}{2a(a^2+b^2)} + \frac{b(2a^4A-3a^3bB+8a^2Ab^2-b^3Ba+4Ab^4)}{2ad(a^2+b^2)(a+b \tan(c+dx))^3} \\
 & \quad \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3} \\
 & \quad \downarrow \text{27} \\
 & \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3}
 \end{aligned}$$

3.296. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

$$\int \frac{\cot(c+dx) \left((Aa^2+2bBa-Ab^2) \tan(c+dx)a^3 + b(2Aa^4-3bBa^3+8Ab^2a^2-b^3Ba+4Ab^4) \tan^2(c+dx) + (a^2+b^2)^2(4Ab-aB) \right)}{(a+b \tan(c+dx))^2} dx + \frac{b(2a^4A-3a^3bB+8a^2Ab^2-ab^3B+4Ab^4)}{2ad(a^2+b^2)(a+b \tan(c+dx))}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3} \quad a$$

↓ 3042

$$\int \frac{(Aa^2+2bBa-Ab^2) \tan(c+dx)a^3 + b(2Aa^4-3bBa^3+8Ab^2a^2-b^3Ba+4Ab^4) \tan(c+dx)^2 + (a^2+b^2)^2(4Ab-aB)}{\tan(c+dx)(a+b \tan(c+dx))^2} dx + \frac{b(2a^4A-3a^3bB+8a^2Ab^2-ab^3B+4Ab^4)}{2ad(a^2+b^2)(a+b \tan(c+dx))}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3} \quad a$$

↓ 4132

$$\int \frac{\cot(c+dx) \left((Aa^3+3bBa^2-3Ab^2a-b^3B) \tan(c+dx)a^4 + b(Aa^6-6bBa^5+13Ab^2a^4-3b^3Ba^3+12Ab^4a^2-b^5Ba+4Ab^6) \tan^2(c+dx) + (a^2+b^2)^3(4Ab-aB) \right)}{a+b \tan(c+dx)} dx + \frac{b(a^6A-6a^5B)}{a(a^2+b^2)}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3} \quad a$$

↓ 3042

$$\int \frac{(Aa^3+3bBa^2-3Ab^2a-b^3B) \tan(c+dx)a^4 + b(Aa^6-6bBa^5+13Ab^2a^4-3b^3Ba^3+12Ab^4a^2-b^5Ba+4Ab^6) \tan(c+dx)^2 + (a^2+b^2)^3(4Ab-aB)}{\tan(c+dx)(a+b \tan(c+dx))} dx + \frac{b(a^6A-6a^5B)}{a(a^2+b^2)}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3} \quad a$$

↓ 4134

$$\frac{(a^2+b^2)^3(4Ab-aB)}{a} \int \cot(c+dx) dx - \frac{b^2(-10a^7B+20a^6Ab-5a^5b^2B+24a^4Ab^3-4a^3b^4B+16a^2Ab^5-ab^6B+4Ab^7)}{a(a^2+b^2)} \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx + \frac{a^4x(a^4A+4a^3bB-6a^2b^2B)}{a^2}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3}$$

3.296. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

↓ 3042

$$\frac{\frac{(a^2+b^2)^3(4Ab-aB) \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx}{a} - \frac{b^2(-10a^7B+20a^6Ab-5a^5b^2B+24a^4Ab^3-4a^3b^4B+16a^2Ab^5-ab^6B+4Ab^7) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^4x(a^4A+4a^3bB)}{a(a^2+b^2)}}{a(a^2+b^2)}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3}$$

↓ 25

$$\frac{\frac{(a^2+b^2)^3(4Ab-aB) \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx}{a} - \frac{b^2(-10a^7B+20a^6Ab-5a^5b^2B+24a^4Ab^3-4a^3b^4B+16a^2Ab^5-ab^6B+4Ab^7) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^4x(a^4A+4a^3bB)}{a(a^2+b^2)}}{a(a^2+b^2)}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3}$$

↓ 3956

$$\frac{\frac{b^2(-10a^7B+20a^6Ab-5a^5b^2B+24a^4Ab^3-4a^3b^4B+16a^2Ab^5-ab^6B+4Ab^7) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{(a^2+b^2)^3(4Ab-aB) \log(-\sin(c+dx))}{ad} + \frac{a^4x(a^4A+4a^3bB)}{a(a^2+b^2)}}{a(a^2+b^2)}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3}$$

↓ 4013

$$\frac{\frac{b(3a^2A-abB+4Ab^2)}{3ad(a^2+b^2)(a+b \tan(c+dx))^3} + \frac{b(2a^4A-3a^3bB+8a^2Ab^2-ab^3B+4Ab^4)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} + \frac{b(a^6A-6a^5bB+13a^4Ab^2-3a^3b^3B+12a^2Ab^4-ab^5B+4Ab^6)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \frac{(a^2+b^2)^3(4Aa^4+4Aa^3bB)}{a(a^2+b^2)}}{a(a^2+b^2)}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^3}$$

input `Int[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]`

3.296. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

```

output  -((A*Cot[c + d*x])/(a*d*(a + b*Tan[c + d*x])^3)) - ((b*(3*a^2*A + 4*A*b^2
- a*b*B))/(3*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^3) + ((b*(2*a^4*A + 8*a^
2*A*b^2 + 4*A*b^4 - 3*a^3*b*B - a*b^3*B))/(2*a*(a^2 + b^2)*d*(a + b*Tan[c
+ d*x])^2) + (((a^4*(a^4*A - 6*a^2*A*b^2 + A*b^4 + 4*a^3*b*B - 4*a*b^3*B)*
x)/(a^2 + b^2) + ((a^2 + b^2)^3*(4*A*b - a*B)*Log[-Sin[c + d*x]])/(a*d) -
(b^2*(20*a^6*A*b + 24*a^4*A*b^3 + 16*a^2*A*b^5 + 4*A*b^7 - 10*a^7*B - 5*a^
5*b^2*B - 4*a^3*b^4*B - a*b^6*B)*Log[a*Cos[c + d*x] + b*SIN[c + d*x]])/(a*
(a^2 + b^2)*d))/(a*(a^2 + b^2)) + (b*(a^6*A + 13*a^4*A*b^2 + 12*a^2*A*b^4
+ 4*A*b^6 - 6*a^5*b*B - 3*a^3*b^3*B - a*b^5*B))/(a*(a^2 + b^2)*d*(a + b*Ta
n[c + d*x])))/(a*(a^2 + b^2)))/(a*(a^2 + b^2))/a

```

3.296.3.1 Defintions of rubi rules used

```

rule 25  Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]

```

```

rule 27  Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]

```

```

rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

```

rule 3956 Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]

```

```

rule 4013 Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*
(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Si
n[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]

```

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4134 `Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/(a^2 + b^2)*(c^2 + d^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.296.4 Maple [A] (verified)

Time = 1.34 (sec) , antiderivative size = 400, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{(4Aa^3b - 4Aab^3 - Ba^4 + 6Ba^2b^2 - Bb^4) \ln(1 + \tan^2(dx+c)) + (-Aa^4 + 6Aa^2b^2 - Ab^4 - 4Ba^3b + 4Bab^3) \arctan(\tan(dx+c))}{(a^2 + b^2)^4} - \frac{a^4 \tan(dx+c)}{(a^2 + b^2)^4}$
default	$\frac{(4Aa^3b - 4Aab^3 - Ba^4 + 6Ba^2b^2 - Bb^4) \ln(1 + \tan^2(dx+c)) + (-Aa^4 + 6Aa^2b^2 - Ab^4 - 4Ba^3b + 4Bab^3) \arctan(\tan(dx+c))}{(a^2 + b^2)^4} - \frac{a^4 \tan(dx+c)}{(a^2 + b^2)^4}$
norman	$\frac{b(6Aa^6b + 33Aa^4b^3 + 35Aa^2b^5 + 12Ab^7 - 10Ba^5b^2 - 9Ba^3b^4 - 3Ba^6b^6)(\tan^2(dx+c))}{da^3(a^6 + 3a^4b^2 + 3a^2b^4 + b^6)} - \frac{A}{ad} - \frac{b^3(Aa^4 - 6Aa^2b^2 + Ab^4 + 4Ba^3b - 4Bab^3)}{(a^6 + 3a^4b^2 + 3a^2b^4 + b^6)(a^2 + b^2)}$
parallelrisch	Expression too large to display
risch	Expression too large to display

```
input int(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x,method=_RETURNVERBOSE)
```

```
output 1/d*(1/(a^2+b^2)^4*(1/2*(4*A*a^3*b-4*A*a*b^3-B*a^4+6*B*a^2*b^2-B*b^4)*ln(1+tan(d*x+c)^2)+(-A*a^4+6*A*a^2*b^2-A*b^4-4*B*a^3*b+4*B*a*b^3)*arctan(tan(d*x+c)))-1/a^4*A/tan(d*x+c)+(-4*A*b+B*a)/a^5*ln(tan(d*x+c))-1/2*b^2*(4*A*a^2*b+2*A*a*b^3-3*B*a^3-B*a*b^2)/(a^2+b^2)^2/a^3/(a+b*tan(d*x+c))^2-b^2*(10*A*a^4*b+9*A*a^2*b^3+3*A*b^5-6*B*a^5-3*B*a^3*b^2-B*a*b^4)/(a^2+b^2)^3/a^4/(a+b*tan(d*x+c))+b^2*(20*A*a^6*b+24*A*a^4*b^3+16*A*a^2*b^5+4*A*b^7-10*B*a^7-5*B*a^5*b^2-4*B*a^3*b^4-B*a*b^6)/(a^2+b^2)^4/a^5*ln(a+b*tan(d*x+c))-1/3*(A*b-B*a)*b^2/(a^2+b^2)/a^2/(a+b*tan(d*x+c))^3)
```

3.296.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1510 vs. 2(393) = 786.

Time = 0.44 (sec) , antiderivative size = 1510, normalized size of antiderivative = 3.78

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="fracas")
```



```
output -1/6*(6*A*a^12 + 24*A*a^10*b^2 + 36*A*a^8*b^4 + 24*A*a^6*b^6 + 6*A*a^4*b^8
+ (47*B*a^7*b^5 - 74*A*a^6*b^6 + 6*B*a^5*b^7 - 42*A*a^4*b^8 + 3*B*a^3*b^9
- 12*A*a^2*b^10 + 6*(A*a^9*b^3 + 4*B*a^8*b^4 - 6*A*a^7*b^5 - 4*B*a^6*b^6
+ A*a^5*b^7)*d*x)*tan(d*x + c)^4 + 3*(2*A*a^9*b^3 + 35*B*a^8*b^4 - 46*A*a^
7*b^5 - 12*B*a^6*b^6 + 8*A*a^5*b^7 - 5*B*a^4*b^8 + 20*A*a^3*b^9 - 2*B*a^2*
b^10 + 8*A*a*b^11 + 6*(A*a^10*b^2 + 4*B*a^9*b^3 - 6*A*a^8*b^4 - 4*B*a^7*b^
5 + A*a^6*b^6)*d*x)*tan(d*x + c)^3 + 3*(6*A*a^10*b^2 + 20*B*a^9*b^3 - 6*A*
a^8*b^4 - 37*B*a^7*b^5 + 80*A*a^6*b^6 - 18*B*a^5*b^7 + 68*A*a^4*b^8 - 5*B*
a^3*b^9 + 20*A*a^2*b^10 + 6*(A*a^11*b + 4*B*a^10*b^2 - 6*A*a^9*b^3 - 4*B*a
^8*b^4 + A*a^7*b^5)*d*x)*tan(d*x + c)^2 - 3*((B*a^9*b^3 - 4*A*a^8*b^4 + 4*
B*a^7*b^5 - 16*A*a^6*b^6 + 6*B*a^5*b^7 - 24*A*a^4*b^8 + 4*B*a^3*b^9 - 16*A
*a^2*b^10 + B*a*b^11 - 4*A*b^12)*tan(d*x + c)^4 + 3*(B*a^10*b^2 - 4*A*a^9*
b^3 + 4*B*a^8*b^4 - 16*A*a^7*b^5 + 6*B*a^6*b^6 - 24*A*a^5*b^7 + 4*B*a^4*b^
8 - 16*A*a^3*b^9 + B*a^2*b^10 - 4*A*a*b^11)*tan(d*x + c)^3 + 3*(B*a^11*b -
4*A*a^10*b^2 + 4*B*a^9*b^3 - 16*A*a^8*b^4 + 6*B*a^7*b^5 - 24*A*a^6*b^6 +
4*B*a^5*b^7 - 16*A*a^4*b^8 + B*a^3*b^9 - 4*A*a^2*b^10)*tan(d*x + c)^2 + (B
*a^12 - 4*A*a^11*b + 4*B*a^10*b^2 - 16*A*a^9*b^3 + 6*B*a^8*b^4 - 24*A*a^7*
b^5 + 4*B*a^6*b^6 - 16*A*a^5*b^7 + B*a^4*b^8 - 4*A*a^3*b^9)*tan(d*x + c))*
log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1)) + 3*((10*B*a^7*b^5 - 20*A*a^6*b^6
+ 5*B*a^5*b^7 - 24*A*a^4*b^8 + 4*B*a^3*b^9 - 16*A*a^2*b^10 + B*a*b^11 ...
```

3.296.6 Sympy [**F(-2)**]

Exception generated.

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^4} dx = \text{Exception raised: AttributeError}$$

```
input integrate(cot(d*x+c)**2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**4,x)
```

```
output Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'
```

3.296.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 698, normalized size of antiderivative = 1.75

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx =$$

$$\frac{6(Aa^4+4Ba^3b-6Aa^2b^2-4Bab^3+Ab^4)(dx+c)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} + \frac{6(10Ba^7b^2-20Aa^6b^3+5Ba^5b^4-24Aa^4b^5+4Ba^3b^6-16Aa^2b^7+Bab^8-4Ab^9) \log(b \tan(dx+c)+a)}{a^{13}+4a^{11}b^2+6a^9b^4+4a^7b^6+a^5b^8}$$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="maxima")
```

```
output -1/6*(6*(A*a^4 + 4*B*a^3*b - 6*A*a^2*b^2 - 4*B*a*b^3 + A*b^4)*(d*x + c)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + 6*(10*B*a^7*b^2 - 20*A*a^6*b^3 + 5*B*a^5*b^4 - 24*A*a^4*b^5 + 4*B*a^3*b^6 - 16*A*a^2*b^7 + B*a*b^8 - 4*A*b^9)*log(b*tan(d*x + c) + a)/(a^13 + 4*a^11*b^2 + 6*a^9*b^4 + 4*a^7*b^6 + a^5*b^8) + 3*(B*a^4 - 4*A*a^3*b - 6*B*a^2*b^2 + 4*A*a*b^3 + B*b^4)*log(tan(d*x + c)^2 + 1)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + (6*A*a^9 + 18*A*a^7*b^2 + 18*A*a^5*b^4 + 6*A*a^3*b^6 + 6*(A*a^6*b^3 - 6*B*a^5*b^4 + 13*A*a^4*b^5 - 3*B*a^3*b^6 + 12*A*a^2*b^7 - B*a*b^8 + 4*A*b^9)*tan(d*x + c)^3 + 3*(6*A*a^7*b^2 - 27*B*a^6*b^3 + 62*A*a^5*b^4 - 16*B*a^4*b^5 + 60*A*a^3*b^6 - 5*B*a^2*b^7 + 20*A*a*b^8)*tan(d*x + c)^2 + (18*A*a^8*b - 47*B*a^7*b^2 + 128*A*a^6*b^3 - 34*B*a^5*b^4 + 130*A*a^4*b^5 - 11*B*a^3*b^6 + 44*A*a^2*b^7)*tan(d*x + c))/(a^10*b^3 + 3*a^8*b^5 + 3*a^6*b^7 + a^4*b^9)*tan(d*x + c)^4 + 3*(a^11*b^2 + 3*a^9*b^4 + 3*a^7*b^6 + a^5*b^8)*tan(d*x + c)^3 + 3*(a^12*b + 3*a^10*b^3 + 3*a^8*b^5 + a^6*b^7)*tan(d*x + c)^2 + (a^13 + 3*a^11*b^2 + 3*a^9*b^4 + a^7*b^6)*tan(d*x + c)) - 6*(B*a - 4*A*b)*log(tan(d*x + c))/a^5)/d
```

3.296.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 846 vs. 2(393) = 786.

Time = 1.31 (sec) , antiderivative size = 846, normalized size of antiderivative = 2.12

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx =$$

$$\frac{6(Aa^4+4Ba^3b-6Aa^2b^2-4Bab^3+Ab^4)(dx+c)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} + \frac{3(Ba^4-4Aa^3b-6Aa^2b^2+4Aab^3+Bb^4) \log(\tan(dx+c)^2+1)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} + \frac{6(10Ba^7b^3-20Aa^6b^4-12Aa^5b^5+6Aa^4b^6-4Aa^3b^7+2Aa^2b^8+Bab^9-4Ab^{10}) \log(b \tan(dx+c)+a)}{a^{13}+4a^{11}b^2+6a^9b^4+4a^7b^6+a^5b^8}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="giac")`

output
$$\begin{aligned} & -1/6*(6*(A*a^4 + 4*B*a^3*b - 6*A*a^2*b^2 - 4*B*a*b^3 + A*b^4)*(d*x + c)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + 3*(B*a^4 - 4*A*a^3*b - 6*B*a^2*b^2 + 4*A*a*b^3 + B*b^4)*\log(\tan(d*x + c)^2 + 1)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + 6*(10*B*a^7*b^3 - 20*A*a^6*b^4 + 5*B*a^5*b^5 - 24*A*a^4*b^6 + 4*B*a^3*b^7 - 16*A*a^2*b^8 + B*a*b^9 - 4*A*b^{10})*\log(\text{abs}(b*\tan(d*x + c) + a))/(a^{13}*b + 4*a^{11}*b^3 + 6*a^9*b^5 + 4*a^7*b^7 + a^5*b^9) - (110*B*a^7*b^5*\tan(d*x + c)^3 - 220*A*a^6*b^6*\tan(d*x + c)^3 + 55*B*a^5*b^7*\tan(d*x + c)^3 - 264*A*a^4*b^8*\tan(d*x + c)^3 + 44*B*a^3*b^9*\tan(d*x + c)^3 - 176*A*a^2*b^{10}*\tan(d*x + c)^3 + 11*B*a*b^{11}*\tan(d*x + c)^3 - 44*A*b^{12}*\tan(d*x + c)^3 + 366*B*a^8*b^4*\tan(d*x + c)^2 - 720*A*a^7*b^5*\tan(d*x + c)^2 + 219*B*a^6*b^6*\tan(d*x + c)^2 - 906*A*a^5*b^7*\tan(d*x + c)^2 + 156*B*a^4*b^8*\tan(d*x + c)^2 - 600*A*a^3*b^9*\tan(d*x + c)^2 + 39*B*a^2*b^{10}*\tan(d*x + c)^2 - 150*A*a*b^{11}*\tan(d*x + c)^2 + 411*B*a^9*b^3*\tan(d*x + c) - 792*A*a^8*b^4*\tan(d*x + c) + 294*B*a^7*b^5*\tan(d*x + c) - 1050*A*a^6*b^6*\tan(d*x + c) + 195*B*a^5*b^7*\tan(d*x + c) - 696*A*a^4*b^8*\tan(d*x + c) + 48*B*a^3*b^9*\tan(d*x + c) - 174*A*a^2*b^{10}*\tan(d*x + c) + 157*B*a^{10}*b^2 - 294*A*a^9*b^3 + 136*B*a^8*b^4 - 414*A*a^7*b^5 + 89*B*a^6*b^6 - 278*A*a^5*b^7 + 22*B*a^4*b^8 - 70*A*a^3*b^9)/(a^{13} + 4*a^{11}*b^2 + 6*a^9*b^4 + 4*a^7*b^6 + a^5*b^8)*(b*\tan(d*x + c) + a)^3 - 6*(B*a - 4*A*b)*\log(\text{abs}(\tan(d*x + c)))/a^5 + 6*(B*a*\tan(d*x + c) - 4*A*b*\tan(d*x + c) + A*a)/(a^5*t...$$

3.296.9 Mupad [B] (verification not implemented)

Time = 12.85 (sec) , antiderivative size = 576, normalized size of antiderivative = 1.44

$$\begin{aligned} & \int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx \\ & = \frac{b^2 \ln(a+b\tan(c+dx))(-10Ba^7+20Aa^6b-5Ba^5b^2+24Aa^4b^3-4Ba^3b^4+16Aa^2b^5-Bab^6+a^5d(a^2+b^2)^4)}{a^5d(a^2+b^2)^4} \\ & \quad - \frac{\ln(\tan(c+dx))(4Ab-Ba)}{a^5d} - \frac{\ln(\tan(c+dx)-i)(A+Bi)}{2d(a^4bi-4a^3b-a^2b^26i+4ab^3+b^4li)} \\ & \quad - \frac{\ln(\tan(c+dx)+i)(B+Ai)}{2d(a^4-a^3b4i-6a^2b^2+ab^34i+b^4)} \\ & \quad - \frac{\frac{A}{a} + \frac{\tan(c+dx)^3(Aa^6b^3-6Ba^5b^4+13Aa^4b^5-3Ba^3b^6+12Aa^2b^7-Bab^8+4Ab^9)}{a^4(a^6+3a^4b^2+3a^2b^4+b^6)}}{\tan(c+dx)^2 \frac{(6Aa^6b^2-27Ba^5b^3+62Aa^4b^4-2a^3(a^6+3a^4b^2+3a^2b^4+b^6))}{2a^3(a^6+3a^4b^2+3a^2b^4+b^6)}} + \frac{\tan(c+dx)^2(6Aa^6b^2-27Ba^5b^3+62Aa^4b^4-2a^3(a^6+3a^4b^2+3a^2b^4+b^6))}{2a^3(a^6+3a^4b^2+3a^2b^4+b^6)} \\ & \quad d(a^3\tan(c+dx)+3a^2b\tan(c+dx))^2 + \end{aligned}$$

3.296. $\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$

input `int((cot(c + d*x))^2*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^4,x)`

output `(b^2*log(a + b*tan(c + d*x))*(4*A*b^7 - 10*B*a^7 + 16*A*a^2*b^5 + 24*A*a^4*b^3 - 4*B*a^3*b^4 - 5*B*a^5*b^2 + 20*A*a^6*b - B*a*b^6))/(a^5*d*(a^2 + b^2)^4) - (log(tan(c + d*x))*(4*A*b - B*a))/(a^5*d) - (log(tan(c + d*x) - 1i)*(A + B*1i))/(2*d*(4*a*b^3 - 4*a^3*b + a^4*1i + b^4*1i - a^2*b^2*6i)) - (log(tan(c + d*x) + 1i)*(A*1i + B))/(2*d*(a*b^3*4i - a^3*b*4i + a^4 + b^4 - 6*a^2*b^2)) - (A/a + (tan(c + d*x))^3*(4*A*b^9 + 12*A*a^2*b^7 + 13*A*a^4*b^5 + A*a^6*b^3 - 3*B*a^3*b^6 - 6*B*a^5*b^4 - B*a*b^8))/(a^4*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2)) + (tan(c + d*x))^2*(20*A*b^8 + 60*A*a^2*b^6 + 62*A*a^4*b^4 + 6*A*a^6*b^2 - 16*B*a^3*b^5 - 27*B*a^5*b^3 - 5*B*a*b^7))/(2*a^3*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2)) + (tan(c + d*x)*(44*A*b^7 + 130*A*a^2*b^5 + 128*A*a^4*b^3 - 34*B*a^3*b^4 - 47*B*a^5*b^2 + 18*A*a^6*b - 11*B*a*b^6))/(6*a^2*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2))/(d*(a^3*tan(c + d*x) + b^3*tan(c + d*x)^4 + 3*a^2*b*tan(c + d*x)^2 + 3*a*b^2*tan(c + d*x)^3))`

3.297 $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

3.297.1 Optimal result 2862
 3.297.2 Mathematica [C] (verified) 2863
 3.297.3 Rubi [A] (verified) 2864
 3.297.4 Maple [A] (verified) 2870
 3.297.5 Fricas [B] (verification not implemented) 2871
 3.297.6 Sympy [F(-2)] 2872
 3.297.7 Maxima [A] (verification not implemented) 2873
 3.297.8 Giac [A] (verification not implemented) 2874
 3.297.9 Mupad [B] (verification not implemented) 2875

3.297.1 Optimal result

Integrand size = 31, antiderivative size = 477

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

$$= \frac{(4a^3Ab - 4aAb^3 - a^4B + 6a^2b^2B - b^4B)x}{(a^2 + b^2)^4} - \frac{(a^2A - 10Ab^2 + 4abB) \log(\sin(c+dx))}{a^6d}$$

$$- \frac{b^3(35a^6Ab + 56a^4Ab^3 + 39a^2Ab^5 + 10Ab^7 - 20a^7B - 24a^5b^2B - 16a^3b^4B - 4ab^6B) \log(a \cos(c+dx))}{a^6(a^2 + b^2)^4d}$$

$$+ \frac{b(9a^2Ab + 10Ab^3 - 3a^3B - 4ab^2B)}{3a^3(a^2 + b^2)d(a+b \tan(c+dx))^3} + \frac{(5Ab - 2aB) \cot(c+dx)}{2a^2d(a+b \tan(c+dx))^3}$$

$$- \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3} + \frac{b(7a^4Ab + 19a^2Ab^3 + 10Ab^5 - 2a^5B - 8a^3b^2B - 4ab^4B)}{2a^4(a^2 + b^2)^2d(a+b \tan(c+dx))^2}$$

$$+ \frac{b(4a^6Ab + 27a^4Ab^3 + 29a^2Ab^5 + 10Ab^7 - a^7B - 13a^5b^2B - 12a^3b^4B - 4ab^6B)}{a^5(a^2 + b^2)^3d(a+b \tan(c+dx))}$$

output

```
(4*A*a^3*b-4*A*a*b^3-B*a^4+6*B*a^2*b^2-B*b^4)*x/(a^2+b^2)^4-(A*a^2-10*A*b^2+4*B*a*b)*ln(sin(d*x+c))/a^6/d-b^3*(35*A*a^6*b+56*A*a^4*b^3+39*A*a^2*b^5+10*A*b^7-20*B*a^7-24*B*a^5*b^2-16*B*a^3*b^4-4*B*a*b^6)*ln(a*cos(d*x+c))+b*ln(a*cos(d*x+c))/a^6/(a^2+b^2)^4/d+1/3*b*(9*A*a^2*b+10*A*b^3-3*B*a^3-4*B*a*b^2)/a^3/(a^2+b^2)/d/(a+b*tan(d*x+c))^3+1/2*(5*A*b-2*B*a)*cot(d*x+c)/a^2/d/(a+b*tan(d*x+c))^3-1/2*A*cot(d*x+c)^2/a/d/(a+b*tan(d*x+c))^3+1/2*b*(7*A*a^4*b+19*A*a^2*b^3+10*A*b^5-2*B*a^5-8*B*a^3*b^2-4*B*a*b^4)/a^4/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^2+b*(4*A*a^6*b+27*A*a^4*b^3+29*A*a^2*b^5+10*A*b^7-B*a^7-13*B*a^5*b^2-12*B*a^3*b^4-4*B*a*b^6)/a^5/(a^2+b^2)^3/d/(a+b*tan(d*x+c))
```

3.297. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

3.297.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.73 (sec) , antiderivative size = 417, normalized size of antiderivative = 0.87

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

$$= \frac{(4Ab - aB) \cot(c+dx)}{a^5 d} - \frac{A \cot^2(c+dx)}{2a^4 d} + \frac{(A+iB) \log(i - \tan(c+dx))}{2(a+ib)^4 d}$$

$$- \frac{(a^2 A - 10Ab^2 + 4abB) \log(\tan(c+dx))}{a^6 d} + \frac{(A-iB) \log(i + \tan(c+dx))}{2(a-ib)^4 d}$$

$$- \frac{b^3(35a^6 Ab + 56a^4 Ab^3 + 39a^2 Ab^5 + 10Ab^7 - 20a^7 B - 24a^5 b^2 B - 16a^3 b^4 B - 4ab^6 B) \log(a+b \tan(c+dx))}{a^6 (a^2 + b^2)^4 d}$$

$$+ \frac{b^3(Ab - aB)}{3a^3 (a^2 + b^2) d(a+b \tan(c+dx))^3} + \frac{b^3(5a^2 Ab + 3Ab^3 - 4a^3 B - 2ab^2 B)}{2a^4 (a^2 + b^2)^2 d(a+b \tan(c+dx))^2}$$

$$+ \frac{b^3(15a^4 Ab + 17a^2 Ab^3 + 6Ab^5 - 10a^5 B - 9a^3 b^2 B - 3ab^4 B)}{a^5 (a^2 + b^2)^3 d(a+b \tan(c+dx))}$$

input `Integrate[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]`

output `((4*A*b - a*B)*Cot[c + d*x])/(a^5*d) - (A*Cot[c + d*x]^2)/(2*a^4*d) + ((A + I*B)*Log[I - Tan[c + d*x]])/(2*(a + I*b)^4*d) - ((a^2*A - 10*A*b^2 + 4*a*b*B)*Log[Tan[c + d*x]])/(a^6*d) + ((A - I*B)*Log[I + Tan[c + d*x]])/(2*(a - I*b)^4*d) - (b^3*(35*a^6*A*b + 56*a^4*A*b^3 + 39*a^2*A*b^5 + 10*A*b^7 - 20*a^7*B - 24*a^5*b^2*B - 16*a^3*b^4*B - 4*a*b^6*B)*Log[a + b*Tan[c + d*x]])/(a^6*(a^2 + b^2)^4*d) + (b^3*(A*b - a*B))/(3*a^3*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^3) + (b^3*(5*a^2*A*b + 3*A*b^3 - 4*a^3*B - 2*a*b^2*B))/(2*a^4*(a^2 + b^2)^2*d*(a + b*Tan[c + d*x])^2) + (b^3*(15*a^4*A*b + 17*a^2*A*b^3 + 6*A*b^5 - 10*a^5*B - 9*a^3*b^2*B - 3*a*b^4*B))/(a^5*(a^2 + b^2)^3*d*(a + b*Tan[c + d*x]))`

3.297.3 Rubi [A] (verified)

Time = 3.10 (sec) , antiderivative size = 545, normalized size of antiderivative = 1.14, number of steps used = 19, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.613$, Rules used = {3042, 4092, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^3(a+b \tan(c+dx))^4} dx \\
 & \quad \downarrow \text{4092} \\
 & -\frac{\int \frac{\cot^2(c+dx)(5Ab \tan^2(c+dx)+2aA \tan(c+dx)+5Ab-2aB)}{(a+b \tan(c+dx))^4} dx}{2a} - \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\int \frac{5Ab \tan(c+dx)^2+2aA \tan(c+dx)+5Ab-2aB}{\tan(c+dx)^2(a+b \tan(c+dx))^4} dx}{2a} - \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3} \\
 & \quad \downarrow \text{4132} \\
 & -\frac{\int -\frac{2 \cot(c+dx)(Aa^2+B \tan(c+dx)a^2+4bBa-10Ab^2-2b(5Ab-2aB) \tan^2(c+dx))}{(a+b \tan(c+dx))^4} dx}{a} - \frac{(5Ab-2aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^3} \\
 & \quad \downarrow \text{27} \\
 & -\frac{2 \int \frac{\cot(c+dx)(Aa^2+B \tan(c+dx)a^2+4bBa-10Ab^2-2b(5Ab-2aB) \tan^2(c+dx))}{(a+b \tan(c+dx))^4} dx}{a} - \frac{(5Ab-2aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^3} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2 \int \frac{Aa^2+B \tan(c+dx)a^2+4bBa-10Ab^2-2b(5Ab-2aB) \tan(c+dx)^2}{\tan(c+dx)(a+b \tan(c+dx))^4} dx}{a} - \frac{(5Ab-2aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^3} - \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3}
 \end{aligned}$$

3.297. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

↓ 4132

$$2 \left(\frac{\int \frac{3 \cot(c+dx) \left(-((Ab-aB) \tan(c+dx)a^3) - b(-3Ba^3+9Aba^2-4b^2Ba+10Ab^3) \tan^2(c+dx) + (a^2+b^2)(Aa^2+4bBa-10Ab^2) \right)}{(a+b \tan(c+dx))^3} dx}{3a(a^2+b^2)} - \frac{b(-3a^3B+9a^2Ab-4ab^2B)}{3ad(a^2+b^2)(a+b \tan(c+dx))} \right)$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3} \quad 2a$$

↓ 27

$$2 \left(\frac{\int \frac{\cot(c+dx) \left(-((Ab-aB) \tan(c+dx)a^3) - b(-3Ba^3+9Aba^2-4b^2Ba+10Ab^3) \tan^2(c+dx) + (a^2+b^2)(Aa^2+4bBa-10Ab^2) \right)}{(a+b \tan(c+dx))^3} dx}{a(a^2+b^2)} - \frac{b(-3a^3B+9a^2Ab-4ab^2B)}{3ad(a^2+b^2)(a+b \tan(c+dx))} \right)$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3} \quad 2a$$

↓ 3042

$$2 \left(\frac{\int \frac{-((Ab-aB) \tan(c+dx)a^3) - b(-3Ba^3+9Aba^2-4b^2Ba+10Ab^3) \tan(c+dx)^2 + (a^2+b^2)(Aa^2+4bBa-10Ab^2)}{\tan(c+dx)(a+b \tan(c+dx))^3} dx}{a(a^2+b^2)} - \frac{b(-3a^3B+9a^2Ab-4ab^2B+10Ab^3)}{3ad(a^2+b^2)(a+b \tan(c+dx))^3} \right)$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3} \quad 2a$$

↓ 4132

$$2 \left(\frac{\int \frac{2 \cot(c+dx) \left(-((-Ba^2+2Aba+b^2B) \tan(c+dx)a^4) - b(-2Ba^5+7Aba^4-8b^2Ba^3+19Ab^3a^2-4b^4Ba+10Ab^5) \tan^2(c+dx) + (a^2+b^2)^2(Aa^2+4bBa-10Ab^2) \right)}{(a+b \tan(c+dx))^2} dx}{2a(a^2+b^2)} - \frac{b(-3a^3B+9a^2Ab-4ab^2B+10Ab^3)}{3ad(a^2+b^2)(a+b \tan(c+dx))} \right)$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3} \quad 2a$$

↓ 27

$$2 \left(\frac{\int \frac{2 \cot(c+dx) \left(-((-Ba^2+2Aba+b^2B) \tan(c+dx)a^4) - b(-2Ba^5+7Aba^4-8b^2Ba^3+19Ab^3a^2-4b^4Ba+10Ab^5) \tan^2(c+dx) + (a^2+b^2)^2(Aa^2+4bBa-10Ab^2) \right)}{(a+b \tan(c+dx))^2} dx}{2a(a^2+b^2)} - \frac{b(-3a^3B+9a^2Ab-4ab^2B+10Ab^3)}{3ad(a^2+b^2)(a+b \tan(c+dx))} \right)$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3} \quad 2a$$

3.297. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

$$2 \left(\int \frac{\cot(c+dx) \left(- \left((-Ba^2 + 2Aba + b^2 B) \tan(c+dx)a^4 \right) - b \left(-2Ba^5 + 7Aba^4 - 8b^2Ba^3 + 19Ab^3a^2 - 4b^4Ba + 10Ab^5 \right) \tan^2(c+dx) + (a^2+b^2)^2 (Aa^2 + 4bBa - 10Ab^2) \right)}{\frac{(a+b \tan(c+dx))^2}{a(a^2+b^2)}} dx \right) \frac{1}{a(a^2+b^2)}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3} \quad 2a$$

↓ 3042

$$2 \left(\int \frac{- \left((-Ba^2 + 2Aba + b^2 B) \tan(c+dx)a^4 \right) - b \left(-2Ba^5 + 7Aba^4 - 8b^2Ba^3 + 19Ab^3a^2 - 4b^4Ba + 10Ab^5 \right) \tan(c+dx)^2 + (a^2+b^2)^2 (Aa^2 + 4bBa - 10Ab^2)}{\frac{\tan(c+dx)(a+b \tan(c+dx))^2}{a(a^2+b^2)}} dx \right) \frac{1}{a(a^2+b^2)}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3} \quad 2a$$

↓ 4132

$$2 \left(\int \frac{\cot(c+dx) \left(- \left((-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \tan(c+dx)a^5 \right) - b \left(-Ba^7 + 4Aba^6 - 13b^2Ba^5 + 27Ab^3a^4 - 12b^4Ba^3 + 29Ab^5a^2 - 4b^6Ba + 10Ab^7 \right) \tan^2(c+dx) + (a^2+b^2)^2 (Aa^3 + 3Aba^2 + 3b^2Ba - Ab^3) \right)}{\frac{a+b \tan(c+dx)}{a(a^2+b^2)}} dx \right) \frac{1}{a(a^2+b^2)}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3}$$

↓ 3042

$$2 \left(\int \frac{- \left((-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \tan(c+dx)a^5 \right) - b \left(-Ba^7 + 4Aba^6 - 13b^2Ba^5 + 27Ab^3a^4 - 12b^4Ba^3 + 29Ab^5a^2 - 4b^6Ba + 10Ab^7 \right) \tan(c+dx)^2 + (a^2+b^2)^3 (Aa^3 + 3Aba^2 + 3b^2Ba - Ab^3)}{\frac{\tan(c+dx)(a+b \tan(c+dx))}{a(a^2+b^2)}} dx \right) \frac{1}{a(a^2+b^2)}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3}$$

3.297. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

↓ 4134

$$2 \left(\frac{(a^2+b^2)^3 (a^2A+4abB-10Ab^2) \int \cot(c+dx) dx}{a} + \frac{b^3 (-20a^7B+35a^6Ab-24a^5b^2B+56a^4Ab^3-16a^3b^4B+39a^2Ab^5-4ab^6B+10Ab^7) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a(a^2+b^2)}{a(a^2+b^2)} \right)$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3}$$

↓ 3042

$$2 \left(\frac{(a^2+b^2)^3 (a^2A+4abB-10Ab^2) \int -\tan(c+dx+\frac{\pi}{2}) dx}{a} + \frac{b^3 (-20a^7B+35a^6Ab-24a^5b^2B+56a^4Ab^3-16a^3b^4B+39a^2Ab^5-4ab^6B+10Ab^7) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a(a^2+b^2)}{a(a^2+b^2)} \right)$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3}$$

↓ 25

$$2 \left(-\frac{(a^2+b^2)^3 (a^2A+4abB-10Ab^2) \int \tan(\frac{1}{2}(2c+\pi)+dx) dx}{a} + \frac{b^3 (-20a^7B+35a^6Ab-24a^5b^2B+56a^4Ab^3-16a^3b^4B+39a^2Ab^5-4ab^6B+10Ab^7) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a(a^2+b^2)}{a(a^2+b^2)} \right)$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^3}$$

↓ 3956

3.297. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

$$2 \left(\frac{b^3(-20a^7B+35a^6Ab-24a^5b^2B+56a^4Ab^3-16a^3b^4B+39a^2Ab^5-4ab^6B+10Ab^7) \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx + \frac{(a^2+b^2)^3(a^2A+4abB-10Ab^2) \log(-\sin(c+dx))}{ad}}{a(a^2+b^2)} \right)$$

$$\frac{A \cot^2(c + dx)}{2ad(a + b \tan(c + dx))^3}$$

↓ 4013

$$2 \left(\frac{\frac{(a^2+b^2)^3(a^2A+4abB-10Ab^2) \log(-\sin(c+dx))}{ad} - \frac{a^5x(a^4(-B)+4a^3Ab+6a^2b^2B-4aAb^3-b^4B)}{a^2+b^2} + \frac{b^3(-20a^7B+35a^6Ab-24a^5b^2B+56a^4Ab^3-16a^3b^4B+39a^2Ab^5-4ab^6B+10Ab^7)}{ad(a^2+b^2)}}{a(a^2+b^2)} \right)$$

$$\frac{A \cot^2(c + dx)}{2ad(a + b \tan(c + dx))^3}$$

input `Int[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]`

output `-1/2*(A*Cot[c + d*x]^2)/(a*d*(a + b*Tan[c + d*x])^3) - (((5*A*b - 2*a*B)*Cot[c + d*x])/(a*d*(a + b*Tan[c + d*x])^3)) + (2*(-1/3*(b*(9*a^2*A*b + 10*A*b^3 - 3*a^3*B - 4*a*b^2*B)))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^3) + (-1/2*(b*(7*a^4*A*b + 19*a^2*A*b^3 + 10*A*b^5 - 2*a^5*B - 8*a^3*b^2*B - 4*a*b^4*B)))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + (((-((a^5*(4*a^3*A*b - 4*a*A*b^3 - a^4*B + 6*a^2*b^2*B - b^4*B)*x)/(a^2 + b^2)) + ((a^2 + b^2)^3*(a^2*A - 10*A*b^2 + 4*a*b*B)*Log[-Sin[c + d*x]])/(a*d) + (b^3*(35*a^6*A*b + 56*a^4*A*b^3 + 39*a^2*A*b^5 + 10*A*b^7 - 20*a^7*B - 24*a^5*b^2*B - 16*a^3*b^4*B - 4*a*b^6*B)*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))/(a*(a^2 + b^2)) - (b*(4*a^6*A*b + 27*a^4*A*b^3 + 29*a^2*A*b^5 + 10*A*b^7 - a^7*B - 13*a^5*b^2*B - 12*a^3*b^4*B - 4*a*b^6*B))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])))/(a*(a^2 + b^2)))/(a*(a^2 + b^2)))/a)/(2*a)`

3.297.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`
- rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`
- rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4134 Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^
2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]*(c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/
((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)
*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Sim
p[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f
*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x]
&& NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

3.297.4 Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 429, normalized size of antiderivative = 0.90

method	result
derivativedivides	$\frac{(A a^4 - 6A a^2 b^2 + A b^4 + 4B a^3 b - 4B a b^3) \ln(1 + \tan^2(dx+c)) + (4A a^3 b - 4A a b^3 - B a^4 + 6B a^2 b^2 - B b^4) \arctan(\tan(dx+c))}{(a^2 + b^2)^4} - \frac{A}{2a^4 \tan(dx+c)}$
default	$\frac{(A a^4 - 6A a^2 b^2 + A b^4 + 4B a^3 b - 4B a b^3) \ln(1 + \tan^2(dx+c)) + (4A a^3 b - 4A a b^3 - B a^4 + 6B a^2 b^2 - B b^4) \arctan(\tan(dx+c))}{(a^2 + b^2)^4} - \frac{A}{2a^4 \tan(dx+c)}$
parallelrisch	$-70b^3(a+b \tan(dx+c))^3 (A a^6 b + \frac{8}{5} A a^4 b^3 + \frac{39}{35} A a^2 b^5 + \frac{2}{7} A b^7 - \frac{4}{7} B a^7 - \frac{24}{35} B a^5 b^2 - \frac{16}{35} B a^3 b^4 - \frac{4}{35} B a b^6) \ln(a+b \tan(dx+c))$
norman	$\frac{b^3 (4A a^3 b - 4A a b^3 - B a^4 + 6B a^2 b^2 - B b^4) x (\tan^5(dx+c))}{(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)(a^2 + b^2)} + \frac{(4A a^3 b - 4A a b^3 - B a^4 + 6B a^2 b^2 - B b^4) a^3 x (\tan^2(dx+c))}{(a^6 + 3a^4 b^2 + 3a^2 b^4 + b^6)(a^2 + b^2)} - \frac{A}{2ad} + \frac{5A}{2a^4 \tan(dx+c)}$
risch	Expression too large to display

3.297. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

```
input int(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x,method=_RETURNVERBOSE)
```

```
output 1/d*(1/(a^2+b^2)^4*(1/2*(A*a^4-6*A*a^2*b^2+A*b^4+4*B*a^3*b-4*B*a*b^3)*ln(1+tan(d*x+c)^2)+(4*A*a^3*b-4*A*a*b^3-B*a^4+6*B*a^2*b^2-B*b^4)*arctan(tan(d*x+c)))-1/2/a^4*A/tan(d*x+c)^2-(-4*A*b+B*a)/a^5/tan(d*x+c)+(-A*a^2+10*A*b^2-4*B*a*b)/a^6*ln(tan(d*x+c))+1/2*b^3*(5*A*a^2*b+3*A*b^3-4*B*a^3-2*B*a*b^2)/(a^2+b^2)^2/a^4/(a+b*tan(d*x+c))^2+b^3*(15*A*a^4*b+17*A*a^2*b^3+6*A*b^5-10*B*a^5-9*B*a^3*b^2-3*B*a*b^4)/(a^2+b^2)^3/a^5/(a+b*tan(d*x+c))-b^3*(35*A*a^6*b+56*A*a^4*b^3+39*A*a^2*b^5+10*A*b^7-20*B*a^7-24*B*a^5*b^2-16*B*a^3*b^4-4*B*a*b^6)/(a^2+b^2)^4/a^6*ln(a+b*tan(d*x+c))+1/3*(A*b-B*a)*b^3/(a^2+b^2)/a^3/(a+b*tan(d*x+c))^3)
```

3.297.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1732 vs. $2(467) = 934$.

Time = 0.53 (sec) , antiderivative size = 1732, normalized size of antiderivative = 3.63

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="fricas")
```

```
output -1/6*(3*A*a^13 + 12*A*a^11*b^2 + 18*A*a^9*b^4 + 12*A*a^7*b^6 + 3*A*a^5*b^8
+ (3*A*a^10*b^3 + 12*A*a^8*b^5 - 74*B*a^7*b^6 + 125*A*a^6*b^7 - 42*B*a^5*
b^8 + 102*A*a^4*b^9 - 12*B*a^3*b^10 + 30*A*a^2*b^11 + 6*(B*a^10*b^3 - 4*A*
a^9*b^4 - 6*B*a^8*b^5 + 4*A*a^7*b^6 + B*a^6*b^7)*d*x)*tan(d*x + c)^5 + 3*(
3*A*a^11*b^2 + 2*B*a^10*b^3 + 4*A*a^9*b^4 - 46*B*a^8*b^5 + 63*A*a^7*b^6 +
8*B*a^6*b^7 - 10*A*a^5*b^8 + 20*B*a^4*b^9 - 48*A*a^3*b^10 + 8*B*a^2*b^11 -
20*A*a*b^12 + 6*(B*a^11*b^2 - 4*A*a^10*b^3 - 6*B*a^9*b^4 + 4*A*a^8*b^5 +
B*a^7*b^6)*d*x)*tan(d*x + c)^4 + 3*(3*A*a^12*b + 6*B*a^11*b^2 - 11*A*a^10*
b^3 - 6*B*a^9*b^4 - 32*A*a^8*b^5 + 80*B*a^7*b^6 - 177*A*a^6*b^7 + 68*B*a^5
*b^8 - 165*A*a^4*b^9 + 20*B*a^3*b^10 - 50*A*a^2*b^11 + 6*(B*a^12*b - 4*A*a
^11*b^2 - 6*B*a^10*b^3 + 4*A*a^9*b^4 + B*a^8*b^5)*d*x)*tan(d*x + c)^3 + (3
*A*a^13 + 18*B*a^12*b - 51*A*a^11*b^2 + 72*B*a^10*b^3 - 234*A*a^9*b^4 + 21
6*B*a^8*b^5 - 513*A*a^7*b^6 + 162*B*a^6*b^7 - 399*A*a^5*b^8 + 44*B*a^4*b^9
- 110*A*a^3*b^10 + 6*(B*a^13 - 4*A*a^12*b - 6*B*a^11*b^2 + 4*A*a^10*b^3 +
B*a^9*b^4)*d*x)*tan(d*x + c)^2 + 3*((A*a^10*b^3 + 4*B*a^9*b^4 - 6*A*a^8*b
^5 + 16*B*a^7*b^6 - 34*A*a^6*b^7 + 24*B*a^5*b^8 - 56*A*a^4*b^9 + 16*B*a^3*
b^10 - 39*A*a^2*b^11 + 4*B*a*b^12 - 10*A*b^13)*tan(d*x + c)^5 + 3*(A*a^11*
b^2 + 4*B*a^10*b^3 - 6*A*a^9*b^4 + 16*B*a^8*b^5 - 34*A*a^7*b^6 + 24*B*a^6*
b^7 - 56*A*a^5*b^8 + 16*B*a^4*b^9 - 39*A*a^3*b^10 + 4*B*a^2*b^11 - 10*A*a*
b^12)*tan(d*x + c)^4 + 3*(A*a^12*b + 4*B*a^11*b^2 - 6*A*a^10*b^3 + 16*B...
```

3.297.6 Sympy [F(-2)]

Exception generated.

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx = \text{Exception raised: AttributeError}$$

```
input integrate(cot(d*x+c)**3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**4,x)
```

```
output Exception raised: AttributeError >> 'NoneType' object has no attribute 'pr
imitive'
```

3.297.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 815, normalized size of antiderivative = 1.71

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx = \frac{6(Ba^4-4Aa^3b-6Ba^2b^2+4Aab^3+Bb^4)(dx+c)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} - \frac{6(20Ba^7b^3-35Aa^6b^4+24Ba^5b^5-56Aa^4b^6+16Ba^3b^7-39Aa^2b^8+4Bab^9-10Ab^{10}) \log(b \tan(dx+c)+a)}{a^{14}+4a^{12}b^2+6a^{10}b^4+4a^8b^6+a^6b^8}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="maxima")
```

```
output -1/6*(6*(B*a^4 - 4*A*a^3*b - 6*B*a^2*b^2 + 4*A*a*b^3 + B*b^4)*(d*x + c)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) - 6*(20*B*a^7*b^3 - 35*A*a^6*b^4 + 24*B*a^5*b^5 - 56*A*a^4*b^6 + 16*B*a^3*b^7 - 39*A*a^2*b^8 + 4*B*a*b^9 - 10*A*b^10)*log(b*tan(d*x + c) + a)/(a^14 + 4*a^12*b^2 + 6*a^10*b^4 + 4*a^8*b^6 + a^6*b^8) - 3*(A*a^4 + 4*B*a^3*b - 6*A*a^2*b^2 - 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)^2 + 1)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) + (3*A*a^10 + 9*A*a^8*b^2 + 9*A*a^6*b^4 + 3*A*a^4*b^6 + 6*(B*a^7*b^3 - 4*A*a^6*b^4 + 13*B*a^5*b^5 - 27*A*a^4*b^6 + 12*B*a^3*b^7 - 29*A*a^2*b^8 + 4*B*a*b^9 - 10*A*b^10)*tan(d*x + c)^4 + 3*(6*B*a^8*b^2 - 23*A*a^7*b^3 + 62*B*a^6*b^4 - 134*A*a^5*b^5 + 60*B*a^4*b^6 - 145*A*a^3*b^7 + 20*B*a^2*b^8 - 50*A*a*b^9)*tan(d*x + c)^3 + (18*B*a^9*b - 63*A*a^8*b^2 + 128*B*a^7*b^3 - 296*A*a^6*b^4 + 130*B*a^5*b^5 - 319*A*a^4*b^6 + 44*B*a^3*b^7 - 110*A*a^2*b^8)*tan(d*x + c)^2 + 3*(2*B*a^10 - 5*A*a^9*b + 6*B*a^8*b^2 - 15*A*a^7*b^3 + 6*B*a^6*b^4 - 15*A*a^5*b^5 + 2*B*a^4*b^6 - 5*A*a^3*b^7)*tan(d*x + c))/(a^11*b^3 + 3*a^9*b^5 + 3*a^7*b^7 + a^5*b^9)*tan(d*x + c)^5 + 3*(a^12*b^2 + 3*a^10*b^4 + 3*a^8*b^6 + a^6*b^8)*tan(d*x + c)^4 + 3*(a^13*b + 3*a^11*b^3 + 3*a^9*b^5 + a^7*b^7)*tan(d*x + c)^3 + (a^14 + 3*a^12*b^2 + 3*a^10*b^4 + a^8*b^6)*tan(d*x + c)^2) + 6*(A*a^2 + 4*B*a*b - 10*A*b^2)*log(tan(d*x + c))/a^6)/d
```


3.297.8 Giac [A] (verification not implemented)

Time = 1.11 (sec) , antiderivative size = 903, normalized size of antiderivative = 1.89

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx =$$

$$\frac{6(Ba^4-4Aa^3b-6Ba^2b^2+4Aab^3+Bb^4)(dx+c)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} - \frac{3(Aa^4+4Ba^3b-6Aa^2b^2-4Bab^3+Ab^4) \log(\tan(dx+c)^2+1)}{a^8+4a^6b^2+6a^4b^4+4a^2b^6+b^8} - \frac{6(20Ba^7b^4-35Aa^6b^5+24B^2a^5b^6-56A^2a^4b^7+16B^2a^3b^8-39A^2a^2b^9+4B^2a^10b-10A^2b^{11}) \log(\tan(dx+c))}{a^{14}+4a^{12}b^2+6a^{10}b^4+4a^8b^6+a^6b^8+220A^2a^7b^6 \tan(dx+c)^3-385A^2a^6b^7 \tan(dx+c)^3+264A^2a^5b^8 \tan(dx+c)^3-616A^2a^4b^9 \tan(dx+c)^3+176A^2a^3b^{10} \tan(dx+c)^3-429A^2a^2b^{11} \tan(dx+c)^3+44A^2a^12b \tan(dx+c)^3-110A^2a^{13}b \tan(dx+c)^3+720A^2a^8b^5 \tan(dx+c)^2-1245A^2a^7b^6 \tan(dx+c)^2+906A^2a^6b^7 \tan(dx+c)^2-2040A^2a^5b^8 \tan(dx+c)^2+600A^2a^4b^9 \tan(dx+c)^2-1425A^2a^3b^{10} \tan(dx+c)^2+150A^2a^2b^{11} \tan(dx+c)^2-366A^2a^12b \tan(dx+c)^2+792A^2a^9b^4 \tan(dx+c)-1350A^2a^8b^5 \tan(dx+c)+1050A^2a^7b^6 \tan(dx+c)-2271A^2a^6b^7 \tan(dx+c)+696A^2a^5b^8 \tan(dx+c)-1596A^2a^4b^9 \tan(dx+c)+174A^2a^3b^{10} \tan(dx+c)-411A^2a^2b^{11} \tan(dx+c)+294A^2a^{10}b^3-492A^2a^9b^4+414A^2a^8b^5-853A^2a^7b^6+278A^2a^6b^7-606A^2a^5b^8+70A^2a^4b^9-157A^2a^3b^{10}}{((a^{14}+4a^{12}b^2+6a^{10}b^4+4a^8b^6+a^6b^8)(b \tan(dx+c)+a)^3)+6(A^2a^2+4A^2a^3b-10A^2a^2b^2) \log(\tan(dx+c))}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="giac")
```

```
output -1/6*(6*(B*a^4 - 4*A*a^3*b - 6*B*a^2*b^2 + 4*A*a*b^3 + B*b^4)*(d*x + c)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) - 3*(A*a^4 + 4*B*a^3*b - 6*A*a^2*b^2 - 4*B*a*b^3 + A*b^4)*log(tan(d*x + c)^2 + 1)/(a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8) - 6*(20*B*a^7*b^4 - 35*A*a^6*b^5 + 24*B*a^5*b^6 - 56*A*a^4*b^7 + 16*B*a^3*b^8 - 39*A*a^2*b^9 + 4*B*a*b^10 - 10*A*b^11)*log(abs(b*tan(d*x + c) + a))/(a^14*b + 4*a^12*b^3 + 6*a^10*b^5 + 4*a^8*b^7 + a^6*b^9) + (220*B*a^7*b^6*tan(d*x + c)^3 - 385*A*a^6*b^7*tan(d*x + c)^3 + 264*B*a^5*b^8*tan(d*x + c)^3 - 616*A*a^4*b^9*tan(d*x + c)^3 + 176*B*a^3*b^10*tan(d*x + c)^3 - 429*A*a^2*b^11*tan(d*x + c)^3 + 44*B*a*b^12*tan(d*x + c)^3 - 110*A*b^13*tan(d*x + c)^3 + 720*B*a^8*b^5*tan(d*x + c)^2 - 1245*A*a^7*b^6*tan(d*x + c)^2 + 906*B*a^6*b^7*tan(d*x + c)^2 - 2040*A*a^5*b^8*tan(d*x + c)^2 + 600*B*a^4*b^9*tan(d*x + c)^2 - 1425*A*a^3*b^10*tan(d*x + c)^2 + 150*B*a^2*b^11*tan(d*x + c)^2 - 366*A*a*b^12*tan(d*x + c)^2 + 792*B*a^9*b^4*tan(d*x + c) - 1350*A*a^8*b^5*tan(d*x + c) + 1050*B*a^7*b^6*tan(d*x + c) - 2271*A*a^6*b^7*tan(d*x + c) + 696*B*a^5*b^8*tan(d*x + c) - 1596*A*a^4*b^9*tan(d*x + c) + 174*B*a^3*b^10*tan(d*x + c) - 411*A*a^2*b^11*tan(d*x + c) + 294*B*a^10*b^3 - 492*A*a^9*b^4 + 414*B*a^8*b^5 - 853*A*a^7*b^6 + 278*B*a^6*b^7 - 606*A*a^5*b^8 + 70*B*a^4*b^9 - 157*A*a^3*b^10)/((a^14 + 4*a^12*b^2 + 6*a^10*b^4 + 4*a^8*b^6 + a^6*b^8)*(b*tan(d*x + c) + a)^3) + 6*(A*a^2 + 4*B*a*b - 10*A*b^2)*log(abs(tan(d*x + c)))/a^6 - 3*(3*A*a^2*tan...
```

3.297.9 Mupad [B] (verification not implemented)

Time = 14.99 (sec) , antiderivative size = 664, normalized size of antiderivative = 1.39

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$$

$$= \frac{\frac{\tan(c+dx)(5Ab-2Ba)}{2a^2} - \frac{A}{2a} + \frac{\tan(c+dx)^4(-Ba^7b^3+4Aa^6b^4-13Ba^5b^5+27Aa^4b^6-12Ba^3b^7+29Aa^2b^8-4Bab^9+10Ab^{10})}{a^5(a^6+3a^4b^2+3a^2b^4+b^6)}}{d(a^3\tan(c+dx) - \frac{\ln(\tan(c+dx))(Aa^2+4Bab-10Ab^2)}{a^6d} + \frac{\ln(\tan(c+dx)+1i)(B+A1i)}{2d(a^41i+4a^3b-a^2b^26i-4ab^3+b^41i)} + \frac{\ln(\tan(c+dx)-1i)(A+B1i)}{2d(a^4+a^3b4i-6a^2b^2-a^2b^34i+b^4)} - \frac{\ln(a+b\tan(c+dx))(-20Ba^7b^3+35Aa^6b^4-24Ba^5b^5+56Aa^4b^6-16Ba^3b^7+39Aa^2b^8-4Bab^9+10Ab^{10})}{d(a^{14}+4a^{12}b^2+6a^{10}b^4+4a^8b^6+a^6b^8)}}$$

input `int((cot(c + d*x)^3*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^4,x)`

```
output ((tan(c + d*x)*(5*A*b - 2*B*a))/(2*a^2) - A/(2*a) + (tan(c + d*x)^4*(10*A*
b^10 + 29*A*a^2*b^8 + 27*A*a^4*b^6 + 4*A*a^6*b^4 - 12*B*a^3*b^7 - 13*B*a^5
*b^5 - B*a^7*b^3 - 4*B*a*b^9))/(a^5*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2)) +
(tan(c + d*x)^3*(50*A*b^9 + 145*A*a^2*b^7 + 134*A*a^4*b^5 + 23*A*a^6*b^3
- 60*B*a^3*b^6 - 62*B*a^5*b^4 - 6*B*a^7*b^2 - 20*B*a*b^8))/(2*a^4*(a^6 + b
^6 + 3*a^2*b^4 + 3*a^4*b^2)) + (tan(c + d*x)^2*(110*A*b^8 + 319*A*a^2*b^6
+ 296*A*a^4*b^4 + 63*A*a^6*b^2 - 130*B*a^3*b^5 - 128*B*a^5*b^3 - 44*B*a*b
^7 - 18*B*a^7*b))/(6*a^3*(a^6 + b^6 + 3*a^2*b^4 + 3*a^4*b^2)))/(d*(a^3*tan(
c + d*x)^2 + b^3*tan(c + d*x)^5 + 3*a^2*b*tan(c + d*x)^3 + 3*a*b^2*tan(c +
d*x)^4)) - (log(tan(c + d*x))*(A*a^2 - 10*A*b^2 + 4*B*a*b))/(a^6*d) + (lo
g(tan(c + d*x) + 1i)*(A*1i + B))/(2*d*(4*a^3*b - 4*a*b^3 + a^4*1i + b^4*1i
- a^2*b^2*6i)) + (log(tan(c + d*x) - 1i)*(A + B*1i))/(2*d*(a^3*b*4i - a*b
^3*4i + a^4 + b^4 - 6*a^2*b^2)) - (log(a + b*tan(c + d*x))*(10*A*b^10 + 39
*A*a^2*b^8 + 56*A*a^4*b^6 + 35*A*a^6*b^4 - 16*B*a^3*b^7 - 24*B*a^5*b^5 - 2
0*B*a^7*b^3 - 4*B*a*b^9))/(d*(a^14 + a^6*b^8 + 4*a^8*b^6 + 6*a^10*b^4 + 4*
a^12*b^2))
```

$$3.298 \quad \int \frac{\tan^3(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.298.1 Optimal result	2876
3.298.2 Mathematica [A] (verified)	2876
3.298.3 Rubi [A] (verified)	2877
3.298.4 Maple [A] (verified)	2878
3.298.5 Fricas [A] (verification not implemented)	2879
3.298.6 Sympy [B] (verification not implemented)	2879
3.298.7 Maxima [A] (verification not implemented)	2879
3.298.8 Giac [B] (verification not implemented)	2880
3.298.9 Mupad [B] (verification not implemented)	2880

3.298.1 Optimal result

Integrand size = 34, antiderivative size = 29

$$\int \frac{\tan^3(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{B \log(\cos(c+dx))}{d} + \frac{B \tan^2(c+dx)}{2d}$$

output `B*ln(cos(d*x+c))/d+1/2*B*tan(d*x+c)^2/d`

3.298.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.90

$$\int \frac{\tan^3(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{B(2 \log(\cos(c+dx)) + \tan^2(c+dx))}{2d}$$

input `Integrate[(Tan[c + d*x]^3*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(B*(2*Log[Cos[c + d*x]] + Tan[c + d*x]^2))/(2*d)`

3.298.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.147$, Rules used = {2011, 3042, 3954, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^3(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \tan^3(c+dx) dx$$

$$\downarrow \text{3042}$$

$$B \int \tan(c+dx)^3 dx$$

$$\downarrow \text{3954}$$

$$B \left(\frac{\tan^2(c+dx)}{2d} - \int \tan(c+dx) dx \right)$$

$$\downarrow \text{3042}$$

$$B \left(\frac{\tan^2(c+dx)}{2d} - \int \tan(c+dx) dx \right)$$

$$\downarrow \text{3956}$$

$$B \left(\frac{\tan^2(c+dx)}{2d} + \frac{\log(\cos(c+dx))}{d} \right)$$

input `Int[(Tan[c + d*x]^3*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `B*(Log[Cos[c + d*x]]/d + Tan[c + d*x]^2/(2*d))`

3.298.3.1 Defintions of rubi rules used

```
rule 2011 Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :>
  Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
  a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3954 Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[b*((b*Tan[c + d
*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x],
x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

```
rule 3956 Int[tan[(c_) + (d_)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

3.298.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.03

method	result	size
derivativedivides	$\frac{B \left(\frac{\tan^2(dx+c)}{2} - \frac{\ln(1+\tan^2(dx+c))}{2} \right)}{d}$	30
default	$\frac{B \left(\frac{\tan^2(dx+c)}{2} - \frac{\ln(1+\tan^2(dx+c))}{2} \right)}{d}$	30
parallelrisch	$-\frac{-B(\tan^2(dx+c)) + B \ln(1+\tan^2(dx+c))}{2d}$	31
norman	$\frac{B(\tan^2(dx+c))}{2d} - \frac{B \ln(1+\tan^2(dx+c))}{2d}$	33
risch	$-iBx - \frac{2iBc}{d} + \frac{2B e^{2i(dx+c)}}{d(e^{2i(dx+c)}+1)^2} + \frac{B \ln(e^{2i(dx+c)}+1)}{d}$	60

```
input int(tan(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

```
output 1/d*B*(1/2*tan(d*x+c)^2-1/2*ln(1+tan(d*x+c)^2))
```

3.298.
$$\int \frac{\tan^3(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.298.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{\tan^3(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{B\tan(dx+c)^2 + B\log\left(\frac{1}{\tan(dx+c)^2+1}\right)}{2d}$$

input `integrate(tan(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output `1/2*(B*tan(d*x + c)^2 + B*log(1/(tan(d*x + c)^2 + 1)))/d`

3.298.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. 2(24) = 48.

Time = 0.57 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.83

$$\int \frac{\tan^3(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \begin{cases} -\frac{B\log(\tan^2(c+dx)+1)}{2d} + \frac{B\tan^2(c+dx)}{2d} & \text{for } d \neq 0 \\ \frac{x(Ba+Bb\tan(c))\tan^3(c)}{a+b\tan(c)} & \text{otherwise} \end{cases}$$

input `integrate(tan(d*x+c)**3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Piecewise((-B*log(tan(c + d*x)**2 + 1)/(2*d) + B*tan(c + d*x)**2/(2*d), Ne(d, 0)), (x*(B*a + B*b*tan(c))*tan(c)**3/(a + b*tan(c)), True))`

3.298.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.03

$$\int \frac{\tan^3(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{B\tan(dx+c)^2 - B\log(\tan(dx+c)^2+1)}{2d}$$

input `integrate(tan(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(B*tan(d*x + c)^2 - B*log(tan(d*x + c)^2 + 1))/d`

3.298. $\int \frac{\tan^3(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$

3.298.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 187 vs. $2(27) = 54$.

Time = 0.57 (sec) , antiderivative size = 187, normalized size of antiderivative = 6.45

$$\int \frac{\tan^3(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{B \log \left(\left| -\frac{\cos(dx+c)+1}{\cos(dx+c)-1} - \frac{\cos(dx+c)-1}{\cos(dx+c)+1} + 2 \right| \right) - B \log \left(\left| -\frac{\cos(dx+c)+1}{\cos(dx+c)-1} - \frac{\cos(dx+c)-1}{\cos(dx+c)+1} - 2 \right| \right) + \frac{B \left(\frac{\cos(dx+c)+1}{\cos(dx+c)-1} + \frac{\cos(dx+c)-1}{\cos(dx+c)+1} \right)}{\frac{\cos(dx+c)+1}{\cos(dx+c)-1} + \frac{\cos(dx+c)-1}{\cos(dx+c)+1}}}{2d}$$

input `integrate(tan(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `-1/2*(B*log(abs(-(cos(d*x + c) + 1)/(cos(d*x + c) - 1) - (cos(d*x + c) - 1)/(cos(d*x + c) + 1) + 2)) - B*log(abs(-(cos(d*x + c) + 1)/(cos(d*x + c) - 1) - (cos(d*x + c) - 1)/(cos(d*x + c) + 1) - 2)) + (B*((cos(d*x + c) + 1)/(cos(d*x + c) - 1) + (cos(d*x + c) - 1)/(cos(d*x + c) + 1)) + 6*B)/((cos(d*x + c) + 1)/(cos(d*x + c) - 1) + (cos(d*x + c) - 1)/(cos(d*x + c) + 1) + 2))/d`

3.298.9 Mupad [B] (verification not implemented)

Time = 7.51 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.97

$$\int \frac{\tan^3(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = -\frac{B (\ln (\tan(c+dx)^2 + 1) - \tan(c+dx)^2)}{2d}$$

input `int((tan(c + d*x)^3*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `-(B*(log(tan(c + d*x)^2 + 1) - tan(c + d*x)^2))/(2*d)`

$$3.299 \quad \int \frac{\tan^2(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.299.1 Optimal result	2881
3.299.2 Mathematica [A] (verified)	2881
3.299.3 Rubi [A] (verified)	2882
3.299.4 Maple [A] (verified)	2883
3.299.5 Fricas [A] (verification not implemented)	2883
3.299.6 Sympy [B] (verification not implemented)	2884
3.299.7 Maxima [A] (verification not implemented)	2884
3.299.8 Giac [A] (verification not implemented)	2884
3.299.9 Mupad [B] (verification not implemented)	2885

3.299.1 Optimal result

Integrand size = 34, antiderivative size = 16

$$\int \frac{\tan^2(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = -Bx + \frac{B \tan(c+dx)}{d}$$

output `-B*x+B*tan(d*x+c)/d`

3.299.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.56

$$\int \frac{\tan^2(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = B \left(-\frac{\arctan(\tan(c+dx))}{d} + \frac{\tan(c+dx)}{d} \right)$$

input `Integrate[(Tan[c + d*x]^2*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `B*(-(ArcTan[Tan[c + d*x]]/d) + Tan[c + d*x]/d)`

3.299.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2011, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^2(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \tan^2(c+dx) dx$$

$$\downarrow \text{3042}$$

$$B \int \tan(c+dx)^2 dx$$

$$\downarrow \text{3954}$$

$$B \left(\frac{\tan(c+dx)}{d} - \int 1 dx \right)$$

$$\downarrow \text{24}$$

$$B \left(\frac{\tan(c+dx)}{d} - x \right)$$

input `Int[(Tan[c + d*x]^2*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `B*(-x + Tan[c + d*x]/d)`

3.299.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

3.299.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

method	result	size
norman	$-Bx + \frac{B \tan(dx+c)}{d}$	17
parallelrisch	$-\frac{Bdx - B \tan(dx+c)}{d}$	20
derivativedivides	$\frac{B(\tan(dx+c) - \arctan(\tan(dx+c)))}{d}$	22
default	$\frac{B(\tan(dx+c) - \arctan(\tan(dx+c)))}{d}$	22
risch	$-Bx + \frac{2iB}{d(e^{2i(dx+c)} + 1)}$	26

input `int(tan(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `-B*x+B*tan(d*x+c)/d`

3.299.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

$$\int \frac{\tan^2(c+dx)(aB + bB \tan(c+dx))}{a + b \tan(c+dx)} dx = -\frac{Bdx - B \tan(dx+c)}{d}$$

input `integrate(tan(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fracas")`

output `-(B*d*x - B*tan(d*x + c))/d`

3.299.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 36 vs. $2(12) = 24$.

Time = 0.40 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.25

$$\int \frac{\tan^2(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \begin{cases} -Bx + \frac{B\tan(c+dx)}{d} & \text{for } d \neq 0 \\ \frac{x(Ba+Bb\tan(c))\tan^2(c)}{a+b\tan(c)} & \text{otherwise} \end{cases}$$

input `integrate(tan(d*x+c)**2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Piecewise((-B*x + B*tan(c + d*x)/d, Ne(d, 0)), (x*(B*a + B*b*tan(c))*tan(c)**2/(a + b*tan(c)), True))`

3.299.7 Maxima [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int \frac{\tan^2(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = -\frac{(dx+c)B - B\tan(dx+c)}{d}$$

input `integrate(tan(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `-((d*x + c)*B - B*tan(d*x + c))/d`

3.299.8 Giac [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.38

$$\int \frac{\tan^2(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = -\frac{(dx+c)B - B\tan(dx+c)}{d}$$

input `integrate(tan(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `-((d*x + c)*B - B*tan(d*x + c))/d`

3.299.9 Mupad [B] (verification not implemented)

Time = 7.98 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\tan^2(c + dx)(aB + bB \tan(c + dx))}{a + b \tan(c + dx)} dx = \frac{B \tan(c + dx)}{d} - Bx$$

input `int((tan(c + d*x)^2*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `(B*tan(c + d*x))/d - B*x`

$$\mathbf{3.300} \quad \int \frac{\tan(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.300.1 Optimal result	2886
3.300.2 Mathematica [A] (verified)	2886
3.300.3 Rubi [A] (verified)	2887
3.300.4 Maple [A] (verified)	2888
3.300.5 Fricas [A] (verification not implemented)	2888
3.300.6 Sympy [B] (verification not implemented)	2889
3.300.7 Maxima [A] (verification not implemented)	2889
3.300.8 Giac [B] (verification not implemented)	2889
3.300.9 Mupad [B] (verification not implemented)	2890

3.300.1 Optimal result

Integrand size = 32, antiderivative size = 13

$$\int \frac{\tan(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = -\frac{B \log(\cos(c+dx))}{d}$$

output `-B*ln(cos(d*x+c))/d`

3.300.2 Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\tan(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = -\frac{B \log(\cos(c+dx))}{d}$$

input `Integrate[(Tan[c + d*x]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `-((B*Log[Cos[c + d*x]])/d)`

3.300.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {2011, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

↓ 2011

$$B \int \tan(c+dx) dx$$

↓ 3042

$$B \int \tan(c+dx) dx$$

↓ 3956

$$-\frac{B \log(\cos(c+dx))}{d}$$

input `Int[(Tan[c + d*x]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `-((B*Log[Cos[c + d*x]])/d)`

3.300.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]`

3.300.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.38

method	result	size
derivativedivides	$\frac{B \ln(1+\tan^2(dx+c))}{2d}$	18
default	$\frac{B \ln(1+\tan^2(dx+c))}{2d}$	18
norman	$\frac{B \ln(1+\tan^2(dx+c))}{2d}$	18
parallelrisc	$\frac{B \ln(1+\tan^2(dx+c))}{2d}$	18
risc	$iBx + \frac{2iBc}{d} - \frac{B \ln(e^{2i(dx+c)}+1)}{d}$	33

```
input int(tan(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/2*B/d*ln(1+tan(d*x+c)^2)
```

3.300.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.46

$$\int \frac{\tan(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = -\frac{B \log\left(\frac{1}{\tan(dx+c)^2+1}\right)}{2d}$$

```
input integrate(tan(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fracas")
```

```
output -1/2*B*log(1/(tan(d*x + c)^2 + 1))/d
```

3.300.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. $2(12) = 24$.

Time = 0.33 (sec) , antiderivative size = 37, normalized size of antiderivative = 2.85

$$\int \frac{\tan(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \begin{cases} \frac{B \log(\tan^2(c+dx)+1)}{2d} & \text{for } d \neq 0 \\ \frac{x(Ba+Bb\tan(c))\tan(c)}{a+b\tan(c)} & \text{otherwise} \end{cases}$$

input `integrate(tan(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Piecewise((B*log(tan(c+d*x)**2+1)/(2*d), Ne(d, 0)), (x*(B*a+B*b*tan(c))*tan(c)/(a+b*tan(c)), True))`

3.300.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

$$\int \frac{\tan(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{B \log(\tan(dx+c)^2+1)}{2d}$$

input `integrate(tan(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*B*log(tan(d*x+c)^2+1)/d`

3.300.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 99 vs. $2(13) = 26$.

Time = 0.33 (sec) , antiderivative size = 99, normalized size of antiderivative = 7.62

$$\int \frac{\tan(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{B \log\left(\left|-\frac{\cos(dx+c)+1}{\cos(dx+c)-1} - \frac{\cos(dx+c)-1}{\cos(dx+c)+1} + 2\right|\right) - B \log\left(\left|-\frac{\cos(dx+c)+1}{\cos(dx+c)-1} - \frac{\cos(dx+c)-1}{\cos(dx+c)+1} - 2\right|\right)}{2d}$$

input `integrate(tan(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `1/2*(B*log(abs(-(cos(d*x + c) + 1)/(cos(d*x + c) - 1) - (cos(d*x + c) - 1)/(cos(d*x + c) + 1) + 2)) - B*log(abs(-(cos(d*x + c) + 1)/(cos(d*x + c) - 1) - (cos(d*x + c) - 1)/(cos(d*x + c) + 1) - 2)))/d`

3.300.9 Mupad [B] (verification not implemented)

Time = 7.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.31

$$\int \frac{\tan(c + dx)(aB + bB \tan(c + dx))}{a + b \tan(c + dx)} dx = \frac{B \ln(\tan(c + dx)^2 + 1)}{2d}$$

input `int((tan(c + d*x)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `(B*log(tan(c + d*x)^2 + 1))/(2*d)`

3.301 $\int \frac{aB+bB \tan(c+dx)}{a+b \tan(c+dx)} dx$

3.301.1 Optimal result 2891
 3.301.2 Mathematica [A] (verified) 2891
 3.301.3 Rubi [A] (verified) 2892
 3.301.4 Maple [A] (verified) 2893
 3.301.5 Fricas [A] (verification not implemented) 2893
 3.301.6 Sympy [A] (verification not implemented) 2893
 3.301.7 Maxima [C] (verification not implemented) 2894
 3.301.8 Giac [C] (verification not implemented) 2894
 3.301.9 Mupad [B] (verification not implemented) 2894

3.301.1 Optimal result

Integrand size = 26, antiderivative size = 3

$$\int \frac{aB + bB \tan(c + dx)}{a + b \tan(c + dx)} dx = Bx$$

output

```
B*x
```

3.301.2 Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{aB + bB \tan(c + dx)}{a + b \tan(c + dx)} dx = Bx$$

input

```
Integrate[(a*B + b*B*Tan[c + d*x])/(a + b*Tan[c + d*x]),x]
```

output

```
B*x
```

3.301.3 Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {2011, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \tan(c + dx)}{a + b \tan(c + dx)} dx$$

↓ 2011

$$B \int 1 dx$$

↓ 24

$$Bx$$

input `Int[(a*B + b*B*Tan[c + d*x])/(a + b*Tan[c + d*x]),x]`

output `B*x`

3.301.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

3.301.4 Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
default	Bx	4
norman	Bx	4
risch	Bx	4
derivativedivides	$\frac{B \arctan(\tan(dx+c))}{d}$	13

input `int((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`output `B*x`**3.301.5 Fracas [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{aB + bB \tan(c + dx)}{a + b \tan(c + dx)} dx = Bx$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`output `B*x`**3.301.6 Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int \frac{aB + bB \tan(c + dx)}{a + b \tan(c + dx)} dx = Bx$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`output `B*x`

3.301.7 Maxima [C] (verification not implemented)

Result contains higher order function than in optimal. Order 3 vs. order 1.

Time = 0.36 (sec) , antiderivative size = 10, normalized size of antiderivative = 3.33

$$\int \frac{aB + bB \tan(c + dx)}{a + b \tan(c + dx)} dx = \frac{(dx + c)B}{d}$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `(d*x + c)*B/d`

3.301.8 Giac [C] (verification not implemented)

Result contains higher order function than in optimal. Order 3 vs. order 1.

Time = 0.32 (sec) , antiderivative size = 10, normalized size of antiderivative = 3.33

$$\int \frac{aB + bB \tan(c + dx)}{a + b \tan(c + dx)} dx = \frac{(dx + c)B}{d}$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `(d*x + c)*B/d`

3.301.9 Mupad [B] (verification not implemented)

Time = 7.31 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \frac{aB + bB \tan(c + dx)}{a + b \tan(c + dx)} dx = Bx$$

input `int((B*a + B*b*tan(c + d*x))/(a + b*tan(c + d*x)),x)`

output `B*x`

3.302
$$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.302.1 Optimal result 2895
 3.302.2 Mathematica [B] (verified) 2895
 3.302.3 Rubi [A] (verified) 2896
 3.302.4 Maple [A] (verified) 2897
 3.302.5 Fricas [A] (verification not implemented) 2897
 3.302.6 Sympy [B] (verification not implemented) 2898
 3.302.7 Maxima [B] (verification not implemented) 2898
 3.302.8 Giac [B] (verification not implemented) 2898
 3.302.9 Mupad [B] (verification not implemented) 2899

3.302.1 Optimal result

Integrand size = 32, antiderivative size = 12

$$\int \frac{\cot(c + dx)(aB + bB \tan(c + dx))}{a + b \tan(c + dx)} dx = \frac{B \log(\sin(c + dx))}{d}$$

output `B*ln(sin(d*x+c))/d`

3.302.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 25 vs. 2(12) = 24.

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.08

$$\int \frac{\cot(c + dx)(aB + bB \tan(c + dx))}{a + b \tan(c + dx)} dx = B \left(\frac{\log(\cos(c + dx))}{d} + \frac{\log(\tan(c + dx))}{d} \right)$$

input `Integrate[(Cot[c + d*x]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `B*(Log[Cos[c + d*x]]/d + Log[Tan[c + d*x]]/d)`

3.302.3 Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2011, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx \\ & \quad \downarrow \text{2011} \\ & B \int \cot(c+dx) dx \\ & \quad \downarrow \text{3042} \\ & B \int -\tan\left(c+dx+\frac{\pi}{2}\right) dx \\ & \quad \downarrow \text{25} \\ & -B \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx \\ & \quad \downarrow \text{3956} \\ & \frac{B \log(-\sin(c+dx))}{d} \end{aligned}$$

input `Int[(Cot[c + d*x]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(B*Log[-Sin[c + d*x]])/d`

3.302.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

3.302.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

method	result	size
derivativedivides	$\frac{B \ln(\sin(dx+c))}{d}$	13
default	$\frac{B \ln(\sin(dx+c))}{d}$	13
parallelrisc	$\frac{B(2 \ln(\tan(dx+c)) - \ln(\sec^2(dx+c)))}{2d}$	28
norman	$\frac{B \ln(\tan(dx+c))}{d} - \frac{B \ln(1+\tan^2(dx+c))}{2d}$	31
risc	$-iBx - \frac{2iBc}{d} + \frac{B \ln(e^{2i(dx+c)} - 1)}{d}$	32

input `int(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `B*ln(sin(d*x+c))/d`

3.302.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.67

$$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{B \log\left(-\frac{1}{2} \cos(2dx+2c) + \frac{1}{2}\right)}{2d}$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output `1/2*B*log(-1/2*cos(2*d*x + 2*c) + 1/2)/d`

3.302. $\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.302.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 49 vs. $2(10) = 20$.

Time = 0.50 (sec) , antiderivative size = 49, normalized size of antiderivative = 4.08

$$\int \frac{\cot(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \begin{cases} -\frac{B \log(\tan^2(c+dx)+1)}{2d} + \frac{B \log(\tan(c+dx))}{d} & \text{for } d \neq 0 \\ \frac{x(Ba+Bb\tan(c))\cot(c)}{a+b\tan(c)} & \text{otherwise} \end{cases}$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Piecewise((-B*log(tan(c + d*x)**2 + 1)/(2*d) + B*log(tan(c + d*x))/d, Ne(d, 0)), (x*(B*a + B*b*tan(c))*cot(c)/(a + b*tan(c)), True))`

3.302.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(12) = 24$.

Time = 0.35 (sec) , antiderivative size = 29, normalized size of antiderivative = 2.42

$$\begin{aligned} & \int \frac{\cot(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx \\ &= -\frac{B \log(\tan(dx+c)^2+1) - 2B \log(\tan(dx+c))}{2d} \end{aligned}$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(B*log(tan(d*x + c)^2 + 1) - 2*B*log(tan(d*x + c)))/d`

3.302.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 59 vs. $2(12) = 24$.

Time = 0.33 (sec) , antiderivative size = 59, normalized size of antiderivative = 4.92

$$\begin{aligned} & \int \frac{\cot(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx \\ &= \frac{B \log\left(\frac{|\cos(dx+c)+1|}{|\cos(dx+c)+1|}\right) - 2B \log\left(\left|-\frac{\cos(dx+c)-1}{\cos(dx+c)+1} + 1\right|\right)}{2d} \end{aligned}$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `1/2*(B*log(abs(-cos(d*x + c) + 1)/abs(cos(d*x + c) + 1)) - 2*B*log(abs(-cos(d*x + c) - 1)/(cos(d*x + c) + 1) + 1))/d`

3.302.9 Mupad [B] (verification not implemented)

Time = 7.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 2.25

$$\int \frac{\cot(c + dx)(aB + bB \tan(c + dx))}{a + b \tan(c + dx)} dx = -\frac{B (\ln(\tan(c + dx)^2 + 1) - 2 \ln(\tan(c + dx)))}{2d}$$

input `int((cot(c + d*x)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `-(B*(log(tan(c + d*x)^2 + 1) - 2*log(tan(c + d*x))))/(2*d)`

3.303 $\int \frac{\cot^2(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.303.1 Optimal result 2900
 3.303.2 Mathematica [C] (verified) 2900
 3.303.3 Rubi [A] (verified) 2901
 3.303.4 Maple [A] (verified) 2902
 3.303.5 Fricas [B] (verification not implemented) 2902
 3.303.6 Sympy [B] (verification not implemented) 2903
 3.303.7 Maxima [A] (verification not implemented) 2903
 3.303.8 Giac [B] (verification not implemented) 2904
 3.303.9 Mupad [B] (verification not implemented) 2904

3.303.1 Optimal result

Integrand size = 34, antiderivative size = 17

$$\int \frac{\cot^2(c + dx)(aB + bB \tan(c + dx))}{a + b \tan(c + dx)} dx = -Bx - \frac{B \cot(c + dx)}{d}$$

output `-B*x-B*cot(d*x+c)/d`

3.303.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.02 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.76

$$\int \frac{\cot^2(c + dx)(aB + bB \tan(c + dx))}{a + b \tan(c + dx)} dx = -\frac{B \cot(c + dx) \text{Hypergeometric2F1}(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(c + dx))}{d}$$

input `Integrate[(Cot[c + d*x]^2*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `-((B*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2])/d)`

3.303.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2011, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^2(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \cot^2(c+dx) dx$$

$$\downarrow \text{3042}$$

$$B \int \tan\left(c+dx+\frac{\pi}{2}\right)^2 dx$$

$$\downarrow \text{3954}$$

$$B\left(-\int 1 dx - \frac{\cot(c+dx)}{d}\right)$$

$$\downarrow \text{24}$$

$$B\left(-\frac{\cot(c+dx)}{d} - x\right)$$

input `Int[(Cot[c + d*x]^2*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `B*(-x - Cot[c + d*x]/d)`

3.303.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

3.303.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.29

method	result	size
derivativedivides	$\frac{B(-\cot(dx+c)-dx-c)}{d}$	22
default	$\frac{B(-\cot(dx+c)-dx-c)}{d}$	22
risch	$-Bx - \frac{2iB}{d(e^{2i(dx+c)}-1)}$	26
parallelrisch	$-\frac{B(\tan(dx+c)dx+1)}{\tan(dx+c)d}$	26
norman	$-\frac{B}{d} - \frac{Bx \tan(dx+c)}{\tan(dx+c)}$	27

input `int(cot(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*B*(-cot(d*x+c)-d*x-c)`

3.303.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. $2(17) = 34$.

Time = 0.25 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.47

$$\int \frac{\cot^2(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = -\frac{Bdx \sin(2dx+2c) + B \cos(2dx+2c) + B}{d \sin(2dx+2c)}$$

input `integrate(cot(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fracas")`

output $-(B*d*x*\sin(2*d*x + 2*c) + B*\cos(2*d*x + 2*c) + B)/(d*\sin(2*d*x + 2*c))$

3.303.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. $2(14) = 28$.

Time = 0.42 (sec) , antiderivative size = 37, normalized size of antiderivative = 2.18

$$\int \frac{\cot^2(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \begin{cases} -Bx - \frac{B\cot(c+dx)}{d} & \text{for } d \neq 0 \\ \frac{x(Ba+Bb\tan(c))\cot^2(c)}{a+b\tan(c)} & \text{otherwise} \end{cases}$$

input `integrate(cot(d*x+c)**2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Piecewise((-B*x - B*cot(c + d*x)/d, Ne(d, 0)), (x*(B*a + B*b*tan(c))*cot(c)**2/(a + b*tan(c)), True))`

3.303.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.35

$$\int \frac{\cot^2(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = -\frac{(dx+c)B + \frac{B}{\tan(dx+c)}}{d}$$

input `integrate(cot(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output $-((d*x + c)*B + B/\tan(d*x + c))/d$

3.303.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(17) = 34$.

Time = 0.36 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.29

$$\int \frac{\cot^2(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= -\frac{2(dx+c)B - B\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + \frac{B}{\tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)}}{2d}$$

input `integrate(cot(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `-1/2*(2*(d*x + c)*B - B*tan(1/2*d*x + 1/2*c) + B/tan(1/2*d*x + 1/2*c))/d`

3.303.9 Mupad [B] (verification not implemented)

Time = 7.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

$$\int \frac{\cot^2(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = -\frac{B(\cot(c+dx) + dx)}{d}$$

input `int((cot(c + d*x)^2*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `-(B*(cot(c + d*x) + d*x))/d`

3.304
$$\int \frac{\cot^3(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.304.1 Optimal result 2905
 3.304.2 Mathematica [A] (verified) 2905
 3.304.3 Rubi [A] (verified) 2906
 3.304.4 Maple [A] (verified) 2908
 3.304.5 Fricas [A] (verification not implemented) 2908
 3.304.6 Sympy [B] (verification not implemented) 2909
 3.304.7 Maxima [A] (verification not implemented) 2909
 3.304.8 Giac [B] (verification not implemented) 2910
 3.304.9 Mupad [B] (verification not implemented) 2910

3.304.1 Optimal result

Integrand size = 34, antiderivative size = 30

$$\int \frac{\cot^3(c + dx)(aB + bB \tan(c + dx))}{a + b \tan(c + dx)} dx = -\frac{B \cot^2(c + dx)}{2d} - \frac{B \log(\sin(c + dx))}{d}$$

output `-1/2*B*cot(d*x+c)^2/d-B*ln(sin(d*x+c))/d`

3.304.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.17

$$\int \frac{\cot^3(c + dx)(aB + bB \tan(c + dx))}{a + b \tan(c + dx)} dx = -\frac{B(\cot^2(c + dx) + 2 \log(\cos(c + dx)) + 2 \log(\tan(c + dx)))}{2d}$$

input `Integrate[(Cot[c + d*x]^3*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `-1/2*(B*(Cot[c + d*x]^2 + 2*Log[Cos[c + d*x]] + 2*Log[Tan[c + d*x]]))/d`

3.304.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2011, 3042, 25, 3954, 25, 3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^3(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \cot^3(c+dx) dx \\
 & \quad \downarrow \text{3042} \\
 & B \int -\tan\left(c+dx+\frac{\pi}{2}\right)^3 dx \\
 & \quad \downarrow \text{25} \\
 & -B \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right)^3 dx \\
 & \quad \downarrow \text{3954} \\
 & -B\left(\frac{\cot^2(c+dx)}{2d} - \int -\cot(c+dx) dx\right) \\
 & \quad \downarrow \text{25} \\
 & -B\left(\int \cot(c+dx) dx + \frac{\cot^2(c+dx)}{2d}\right) \\
 & \quad \downarrow \text{3042} \\
 & -B\left(\int -\tan\left(c+dx+\frac{\pi}{2}\right) dx + \frac{\cot^2(c+dx)}{2d}\right) \\
 & \quad \downarrow \text{25} \\
 & -B\left(\frac{\cot^2(c+dx)}{2d} - \int \tan\left(\frac{1}{2}(2c+\pi)+dx\right) dx\right) \\
 & \quad \downarrow \text{3956} \\
 & -B\left(\frac{\cot^2(c+dx)}{2d} + \frac{\log(-\sin(c+dx))}{d}\right)
 \end{aligned}$$

input `Int[(Cot[c + d*x]^3*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `-(B*(Cot[c + d*x]^2/(2*d) + Log[-Sin[c + d*x]]/d))`

3.304.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3954 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d
x])^(n - 1)/(d(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x]
, x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3956 `Int[tan[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]`

3.304.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.87

method	result	size
derivativedivides	$\frac{B \left(-\frac{\cot^2(dx+c)}{2} - \ln(\sin(dx+c)) \right)}{d}$	26
default	$\frac{B \left(-\frac{\cot^2(dx+c)}{2} - \ln(\sin(dx+c)) \right)}{d}$	26
parallelrisc	$-\frac{B(2 \ln(\tan(dx+c)) - \ln(\sec^2(dx+c)) + \cot^2(dx+c))}{2d}$	36
norman	$-\frac{B}{2d \tan(dx+c)^2} - \frac{B \ln(\tan(dx+c))}{d} + \frac{B \ln(1+\tan^2(dx+c))}{2d}$	46
risc	$iBx + \frac{2iBc}{d} + \frac{2Be^{2i(dx+c)}}{d(e^{2i(dx+c)}-1)^2} - \frac{B \ln(e^{2i(dx+c)}-1)}{d}$	61

```
input int(cot(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

```
output 1/d*B*(-1/2*cot(d*x+c)^2-ln(sin(d*x+c)))
```

3.304.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.77

$$\int \frac{\cot^3(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$= -\frac{(B \cos(2dx+2c) - B) \log\left(-\frac{1}{2} \cos(2dx+2c) + \frac{1}{2}\right) - 2B}{2(d \cos(2dx+2c) - d)}$$

```
input integrate(cot(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm=
"fracas")
```

```
output -1/2*((B*cos(2*d*x + 2*c) - B)*log(-1/2*cos(2*d*x + 2*c) + 1/2) - 2*B)/(d*
cos(2*d*x + 2*c) - d)
```

3.304.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. $2(26) = 52$.

Time = 0.92 (sec) , antiderivative size = 76, normalized size of antiderivative = 2.53

$$\int \frac{\cot^3(c + dx)(aB + bB \tan(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \begin{cases} \tilde{\infty} Bx & \text{for } c = 0 \wedge d = 0 \\ \frac{x(Ba + Bb \tan(c)) \cot^3(c)}{a + b \tan(c)} & \text{for } d = 0 \\ \tilde{\infty} Bx & \text{for } c = -dx \\ \frac{B \log(\tan^2(c + dx) + 1)}{2d} - \frac{B \log(\tan(c + dx))}{d} - \frac{B}{2d \tan^2(c + dx)} & \text{otherwise} \end{cases}$$

input `integrate(cot(d*x+c)**3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Piecewise((zoo*B*x, Eq(c, 0) & Eq(d, 0)), (x*(B*a + B*b*tan(c))*cot(c)**3/(a + b*tan(c)), Eq(d, 0)), (zoo*B*x, Eq(c, -d*x)), (B*log(tan(c + d*x)**2 + 1)/(2*d) - B*log(tan(c + d*x))/d - B/(2*d*tan(c + d*x)**2), True))`

3.304.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.33

$$\int \frac{\cot^3(c + dx)(aB + bB \tan(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \frac{B \log(\tan(dx + c)^2 + 1) - 2B \log(\tan(dx + c)) - \frac{B}{\tan(dx + c)^2}}{2d}$$

input `integrate(cot(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(B*log(tan(d*x + c)^2 + 1) - 2*B*log(tan(d*x + c)) - B/tan(d*x + c)^2)/d`

3.304.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 124 vs. 2(28) = 56.

Time = 0.39 (sec) , antiderivative size = 124, normalized size of antiderivative = 4.13

$$\int \frac{\cot^3(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx =$$

$$\frac{4B \log\left(\frac{|-\cos(dx+c)+1|}{|\cos(dx+c)+1|}\right) - 8B \log\left(\left|-\frac{\cos(dx+c)-1}{\cos(dx+c)+1} + 1\right|\right) - \frac{\left(B + \frac{4B(\cos(dx+c)-1)}{\cos(dx+c)+1}\right)(\cos(dx+c)+1)}{\cos(dx+c)-1} - \frac{B(\cos(dx+c)-1)}{\cos(dx+c)+1}}{8d}$$

input `integrate(cot(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `-1/8*(4*B*log(abs(-cos(d*x + c) + 1)/abs(cos(d*x + c) + 1)) - 8*B*log(abs(-cos(d*x + c) - 1)/(cos(d*x + c) + 1) + 1)) - (B + 4*B*(cos(d*x + c) - 1)/(cos(d*x + c) + 1))*(cos(d*x + c) + 1)/(cos(d*x + c) - 1) - B*(cos(d*x + c) - 1)/(cos(d*x + c) + 1))/d`

3.304.9 Mupad [B] (verification not implemented)

Time = 7.27 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.23

$$\int \frac{\cot^3(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{B(\cot(c+dx)^2 - \ln(\tan(c+dx)^2 + 1) + 2\ln(\tan(c+dx)))}{2d}$$

input `int((cot(c + d*x)^3*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `-(B*(2*log(tan(c + d*x)) - log(tan(c + d*x)^2 + 1) + cot(c + d*x)^2))/(2*d)`

3.305 $\int \frac{\cot^4(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.305.1 Optimal result 2911
 3.305.2 Mathematica [C] (verified) 2911
 3.305.3 Rubi [A] (verified) 2912
 3.305.4 Maple [A] (verified) 2913
 3.305.5 Fricas [B] (verification not implemented) 2914
 3.305.6 Sympy [A] (verification not implemented) 2914
 3.305.7 Maxima [A] (verification not implemented) 2915
 3.305.8 Giac [B] (verification not implemented) 2915
 3.305.9 Mupad [B] (verification not implemented) 2915

3.305.1 Optimal result

Integrand size = 34, antiderivative size = 31

$$\int \frac{\cot^4(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = Bx + \frac{B \cot(c+dx)}{d} - \frac{B \cot^3(c+dx)}{3d}$$

output `B*x+B*cot(d*x+c)/d-1/3*B*cot(d*x+c)^3/d`

3.305.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.02 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.10

$$\int \frac{\cot^4(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = -\frac{B \cot^3(c+dx) \text{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, -\tan^2(c+dx)\right)}{3d}$$

input `Integrate[(Cot[c + d*x]^4*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `-1/3*(B*Cot[c + d*x]^3*Hypergeometric2F1[-3/2, 1, -1/2, -Tan[c + d*x]^2])/d`

3.305.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2011, 3042, 3954, 3042, 3954, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot^4(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx \\ & \quad \downarrow \text{2011} \\ & B \int \cot^4(c+dx) dx \\ & \quad \downarrow \text{3042} \\ & B \int \tan\left(c+dx+\frac{\pi}{2}\right)^4 dx \\ & \quad \downarrow \text{3954} \\ & B\left(-\int \cot^2(c+dx) dx - \frac{\cot^3(c+dx)}{3d}\right) \\ & \quad \downarrow \text{3042} \\ & B\left(-\int \tan\left(c+dx+\frac{\pi}{2}\right)^2 dx - \frac{\cot^3(c+dx)}{3d}\right) \\ & \quad \downarrow \text{3954} \\ & B\left(\int 1 dx - \frac{\cot^3(c+dx)}{3d} + \frac{\cot(c+dx)}{d}\right) \\ & \quad \downarrow \text{24} \\ & B\left(-\frac{\cot^3(c+dx)}{3d} + \frac{\cot(c+dx)}{d} + x\right) \end{aligned}$$

input `Int[(Cot[c + d*x]^4*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `B*(x + Cot[c + d*x]/d - Cot[c + d*x]^3/(3*d))`

3.305.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

3.305.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

method	result	size
derivativedivides	$\frac{B \left(-\frac{\cot^3(dx+c)}{3} + \cot(dx+c) + dx+c \right)}{d}$	27
default	$\frac{B \left(-\frac{\cot^3(dx+c)}{3} + \cot(dx+c) + dx+c \right)}{d}$	27
parallelrisch	$-\frac{B(\cot^3(dx+c) - 3dx - 3\cot(dx+c))}{3d}$	28
norman	$\frac{Bx(\tan^3(dx+c)) + \frac{B(\tan^2(dx+c))}{d} - \frac{B}{3d}}{\tan(dx+c)^3}$	41
risch	$Bx + \frac{4iB(3e^{4i(dx+c)} - 3e^{2i(dx+c)} + 2)}{3d(e^{2i(dx+c)} - 1)^3}$	49

input `int(cot(d*x+c)^4*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)), x, method=_RETURNVERBOSE)`

output `1/d*B*(-1/3*cot(d*x+c)^3+cot(d*x+c)+d*x+c)`

3.305.
$$\int \frac{\cot^4(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.305.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 90 vs. $2(29) = 58$.

Time = 0.26 (sec) , antiderivative size = 90, normalized size of antiderivative = 2.90

$$\int \frac{\cot^4(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{4B\cos(2dx+2c)^2 + 2B\cos(2dx+2c) + 3(Bdx\cos(2dx+2c) - Bdx)\sin(2dx+2c) - 2B}{3(d\cos(2dx+2c) - d)\sin(2dx+2c)}$$

input `integrate(cot(d*x+c)^4*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output `1/3*(4*B*cos(2*d*x + 2*c)^2 + 2*B*cos(2*d*x + 2*c) + 3*(B*d*x*cos(2*d*x + 2*c) - B*d*x)*sin(2*d*x + 2*c) - 2*B)/((d*cos(2*d*x + 2*c) - d)*sin(2*d*x + 2*c))`

3.305.6 Sympy [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.58

$$\int \frac{\cot^4(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \begin{cases} Bx - \frac{B\cot^3(c+dx)}{3d} + \frac{B\cot(c+dx)}{d} & \text{for } d \neq 0 \\ \frac{x(Ba+Bb\tan(c))\cot^4(c)}{a+b\tan(c)} & \text{otherwise} \end{cases}$$

input `integrate(cot(d*x+c)**4*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Piecewise((B*x - B*cot(c + d*x)**3/(3*d) + B*cot(c + d*x)/d, Ne(d, 0)), (x*(B*a + B*b*tan(c))*cot(c)**4/(a + b*tan(c)), True))`

3.305.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.23

$$\int \frac{\cot^4(c + dx)(aB + bB \tan(c + dx))}{a + b \tan(c + dx)} dx = \frac{3(dx + c)B + \frac{3B \tan(dx+c)^2 - B}{\tan(dx+c)^3}}{3d}$$

input `integrate(cot(d*x+c)^4*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/3*(3*(d*x + c)*B + (3*B*tan(d*x + c)^2 - B)/tan(d*x + c)^3)/d`

3.305.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 69 vs. 2(29) = 58.

Time = 0.41 (sec) , antiderivative size = 69, normalized size of antiderivative = 2.23

$$\int \frac{\cot^4(c + dx)(aB + bB \tan(c + dx))}{a + b \tan(c + dx)} dx$$

$$= \frac{B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 24(dx + c)B - 15B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + \frac{15B \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - B}{\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3}}{24d}$$

input `integrate(cot(d*x+c)^4*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `1/24*(B*tan(1/2*d*x + 1/2*c)^3 + 24*(d*x + c)*B - 15*B*tan(1/2*d*x + 1/2*c) + (15*B*tan(1/2*d*x + 1/2*c)^2 - B)/tan(1/2*d*x + 1/2*c)^3)/d`

3.305.9 Mupad [B] (verification not implemented)

Time = 7.14 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03

$$\int \frac{\cot^4(c + dx)(aB + bB \tan(c + dx))}{a + b \tan(c + dx)} dx = Bx - \frac{\frac{B}{3} - B \tan(c + dx)^2}{d \tan(c + dx)^3}$$

input `int((cot(c + d*x)^4*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `B*x - (B/3 - B*tan(c + d*x)^2)/(d*tan(c + d*x)^3)`

3.306
$$\int \frac{\tan^4(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.306.1 Optimal result 2916
 3.306.2 Mathematica [C] (verified) 2916
 3.306.3 Rubi [A] (verified) 2917
 3.306.4 Maple [A] (verified) 2921
 3.306.5 Fricas [A] (verification not implemented) 2921
 3.306.6 Sympy [C] (verification not implemented) 2922
 3.306.7 Maxima [A] (verification not implemented) 2923
 3.306.8 Giac [A] (verification not implemented) 2924
 3.306.9 Mupad [B] (verification not implemented) 2924

3.306.1 Optimal result

Integrand size = 34, antiderivative size = 102

$$\int \frac{\tan^4(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \frac{aBx}{a^2+b^2} + \frac{bB \log(\cos(c+dx))}{(a^2+b^2)d} + \frac{a^4B \log(a+b \tan(c+dx))}{b^3(a^2+b^2)d} - \frac{aB \tan(c+dx)}{b^2d} + \frac{B \tan^2(c+dx)}{2bd}$$

```
output a*B*x/(a^2+b^2)+b*B*ln(cos(d*x+c))/(a^2+b^2)/d+a^4*B*ln(a+b*tan(d*x+c))/b^3/(a^2+b^2)/d-a*B*tan(d*x+c)/b^2/d+1/2*B*tan(d*x+c)^2/b/d
```

3.306.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.39 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.06

$$\int \frac{\tan^4(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \frac{B \left(\frac{\log(i-\tan(c+dx))}{ia-b} - \frac{\log(i+\tan(c+dx))}{ia+b} + \frac{2a^4 \log(a+b \tan(c+dx))}{b^3(a^2+b^2)} - \frac{2a \tan(c+dx)}{b^2} + \frac{\tan^2(c+dx)}{b} \right)}{2d}$$

input `Integrate[(Tan[c + d*x]^4*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2, x]`

output `(B*(Log[I - Tan[c + d*x]]/(I*a - b) - Log[I + Tan[c + d*x]]/(I*a + b) + (2*a^4*Log[a + b*Tan[c + d*x]])/(b^3*(a^2 + b^2)) - (2*a*Tan[c + d*x])/b^2 + Tan[c + d*x]^2/b))/(2*d)`

3.306.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.13, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.382$, Rules used = {2011, 3042, 4049, 27, 3042, 4130, 25, 3042, 4110, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^4(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{\tan^4(c+dx)}{a+b\tan(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{\tan(c+dx)^4}{a+b\tan(c+dx)} dx \\
 & \quad \downarrow \text{4049} \\
 & B \left(\frac{\int -\frac{2\tan(c+dx)(a\tan^2(c+dx)+b\tan(c+dx)+a)}{a+b\tan(c+dx)} dx}{2b} + \frac{\tan^2(c+dx)}{2bd} \right) \\
 & \quad \downarrow \text{27} \\
 & B \left(\frac{\tan^2(c+dx)}{2bd} - \frac{\int \frac{\tan(c+dx)(a\tan^2(c+dx)+b\tan(c+dx)+a)}{a+b\tan(c+dx)} dx}{b} \right) \\
 & \quad \downarrow \text{3042} \\
 & B \left(\frac{\tan^2(c+dx)}{2bd} - \frac{\int \frac{\tan(c+dx)(a\tan(c+dx)^2+b\tan(c+dx)+a)}{a+b\tan(c+dx)} dx}{b} \right)
 \end{aligned}$$

$$\begin{array}{c}
\downarrow 4130 \\
B \left(\frac{\tan^2(c+dx)}{2bd} - \frac{\int -\frac{a^2+(a^2-b^2)\tan^2(c+dx)}{a+b\tan(c+dx)} dx}{b} + \frac{a\tan(c+dx)}{bd} \right) \\
\downarrow 25 \\
B \left(\frac{\tan^2(c+dx)}{2bd} - \frac{\frac{a\tan(c+dx)}{bd} - \int \frac{a^2+(a^2-b^2)\tan^2(c+dx)}{a+b\tan(c+dx)} dx}{b} \right) \\
\downarrow 3042 \\
B \left(\frac{\tan^2(c+dx)}{2bd} - \frac{\frac{a\tan(c+dx)}{bd} - \int \frac{a^2+(a^2-b^2)\tan(c+dx)^2}{a+b\tan(c+dx)} dx}{b} \right) \\
\downarrow 4110 \\
B \left(\frac{\tan^2(c+dx)}{2bd} - \frac{\frac{a\tan(c+dx)}{bd} - \frac{-b^3 \int \tan(c+dx) dx}{a^2+b^2} + \frac{a^4 \int \frac{\tan^2(c+dx)+1}{a+b\tan(c+dx)} dx}{a^2+b^2} + \frac{ab^2x}{a^2+b^2}}{b} \right) \\
\downarrow 3042 \\
B \left(\frac{\tan^2(c+dx)}{2bd} - \frac{\frac{a\tan(c+dx)}{bd} - \frac{-b^3 \int \tan(c+dx) dx}{a^2+b^2} + \frac{a^4 \int \frac{\tan(c+dx)^2+1}{a+b\tan(c+dx)} dx}{a^2+b^2} + \frac{ab^2x}{a^2+b^2}}{b} \right) \\
\downarrow 3956 \\
B \left(\frac{\tan^2(c+dx)}{2bd} - \frac{\frac{a\tan(c+dx)}{bd} - \frac{\frac{a^4 \int \frac{\tan(c+dx)^2+1}{a+b\tan(c+dx)} dx}{a^2+b^2} + \frac{ab^2x}{a^2+b^2} + \frac{b^3 \log(\cos(c+dx))}{d(a^2+b^2)}}{b} \right) \\
\downarrow 4100 \\
B \left(\frac{\tan^2(c+dx)}{2bd} - \frac{\frac{a\tan(c+dx)}{bd} - \frac{\frac{a^4 \int \frac{1}{a+b\tan(c+dx)} d(b\tan(c+dx))}{bd(a^2+b^2)} + \frac{ab^2x}{a^2+b^2} + \frac{b^3 \log(\cos(c+dx))}{d(a^2+b^2)}}{b} \right)
\end{array}$$

3.306. $\int \frac{\tan^4(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

$$B \left(\frac{\tan^2(c+dx)}{2bd} - \frac{\frac{a \tan(c+dx)}{bd} - \frac{\frac{ab^2x}{a^2+b^2} + \frac{b^3 \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{a^4 \log(a+b \tan(c+dx))}{bd(a^2+b^2)}}{b}}{b} \right)$$

input `Int[(Tan[c + d*x]^4*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `B*(Tan[c + d*x]^2/(2*b*d) - (-(((a*b^2*x)/(a^2 + b^2) + (b^3*Log[Cos[c + d*x]]))/(a^2 + b^2)*d) + (a^4*Log[a + b*Tan[c + d*x]])/(b*(a^2 + b^2)*d))/b) + (a*Tan[c + d*x])/(b*d))/b`

3.306.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_.)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_.)*(Gx_)] /; FreeQ[b, x]`

rule 2011 `Int[(u_.)*((a_.) + (b_.)*(v_))^(m_.)*((c_.) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4049 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m - 2)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n - 1))), x] + Simp[1/(d*(m + n - 1)) Int[(a + b*Tan[e + f*x])^(m - 3)*(c + d*Tan[e + f*x])^n*Simp[a^3*d*(m + n - 1) - b^2*(b*c*(m - 2) + a*d*(1 + n)) + b*d*(m + n - 1)*(3*a^2 - b^2)*Tan[e + f*x] - b^2*(b*c*(m - 2) - a*d*(3*m + 2*n - 4))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && GtQ[m, 2] && (GeQ[n, -1] || IntegerQ[m]) && !(IGtQ[n, 2] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4100 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]`

rule 4110 `Int[((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[a*(A - C)*(x/(a^2 + b^2)), x] + (Simp[(a^2*C + A*b^2)/(a^2 + b^2) Int[(1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[b*((A - C)/(a^2 + b^2)) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, C}, x] && NeQ[a^2*C + A*b^2, 0] && NeQ[a^2 + b^2, 0] && NeQ[A, C]`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.306.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.91

method	result
derivativedivides	$\frac{B \left(-\frac{b(\tan^2(dx+c))}{2} + a \tan(dx+c) + \frac{-b \ln(1+\tan^2(dx+c))}{2} + a \arctan(\tan(dx+c)) + \frac{a^4 \ln(a+b \tan(dx+c))}{b^3(a^2+b^2)} \right)}{d}$
default	$\frac{B \left(-\frac{b(\tan^2(dx+c))}{2} + a \tan(dx+c) + \frac{-b \ln(1+\tan^2(dx+c))}{2} + a \arctan(\tan(dx+c)) + \frac{a^4 \ln(a+b \tan(dx+c))}{b^3(a^2+b^2)} \right)}{d}$
parallelrisch	$-\frac{-2Ba^3b^3dx - Ba^2b^2(\tan^2(dx+c)) - Bb^4(\tan^2(dx+c)) + B \ln(1+\tan^2(dx+c))b^4 - 2B \ln(a+b \tan(dx+c))a^4 + 2Ba^3b \tan(dx+c)}{2b^3d(a^2+b^2)}$
norman	$\frac{\frac{Ba^2x}{a^2+b^2} + \frac{Ba^3}{db^3} + \frac{bBax \tan(dx+c)}{a^2+b^2} + \frac{B(\tan^3(dx+c))}{2d} - \frac{Ba(\tan^2(dx+c))}{2bd}}{a+b \tan(dx+c)} + \frac{a^4 B \ln(a+b \tan(dx+c))}{b^3(a^2+b^2)d} - \frac{Bb \ln(1+\tan^2(dx+c))}{2d(a^2+b^2)}$
risch	$-\frac{x B}{ib-a} + \frac{2i B a^2 x}{b^3} + \frac{2i B a^2 c}{b^3 d} - \frac{2i B x}{b} - \frac{2i B c}{bd} - \frac{2ia^4 B x}{b^3(a^2+b^2)} - \frac{2ia^4 B c}{b^3 d(a^2+b^2)} + \frac{2B(-ia e^{2i(dx+c)} + b e^{2i(dx+c)})}{b^2 d(e^{2i(dx+c)} + 1)^2}$

input `int (tan(d*x+c)^4*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNV ERBOSE)`

output `1/d*B*(-1/b^2*(-1/2*b*tan(d*x+c)^2+a*tan(d*x+c))+1/(a^2+b^2)*(-1/2*b*ln(1+tan(d*x+c)^2)+a*arctan(tan(d*x+c)))+1/b^3*a^4/(a^2+b^2)*ln(a+b*tan(d*x+c)))`

3.306.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.41

$$\int \frac{\tan^4(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{2 Bab^3 dx + Ba^4 \log \left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2 + 1} \right) + (Ba^2 b^2 + Bb^4) \tan(dx+c)^2 - (Ba^4 - Bb^4) \log \left(\frac{1}{\tan(dx+c)} \right)}{2(a^2 b^3 + b^5) d}$$

input `integrate(tan(d*x+c)^4*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm m="fricas")`

output $1/2*(2*B*a*b^3*d*x + B*a^4*\log((b^2*\tan(d*x + c)^2 + 2*a*b*\tan(d*x + c) + a^2)/(\tan(d*x + c)^2 + 1)) + (B*a^2*b^2 + B*b^4)*\tan(d*x + c)^2 - (B*a^4 - B*b^4)*\log(1/(\tan(d*x + c)^2 + 1)) - 2*(B*a^3*b + B*a*b^3)*\tan(d*x + c))/((a^2*b^3 + b^5)*d)$

3.306.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.47 (sec) , antiderivative size = 782, normalized size of antiderivative = 7.67

$$\int \frac{\tan^4(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \left\{ \begin{array}{l} \tilde{\infty} Bx \tan^3(c) \\ B \left(x + \frac{\tan^3(c+dx)}{3d} - \frac{\tan(c+dx)}{d} \right) \\ \frac{3iBdx \tan(c+dx)}{2bd \tan(c+dx) - 2ibd} - \frac{3Bdx}{2bd \tan(c+dx) - 2ibd} - \frac{2B \log(\tan^2(c+dx)+1) \tan(c+dx)}{2bd \tan(c+dx) - 2ibd} + \frac{2iB \log(\tan^2(c+dx)+1)}{2bd \tan(c+dx) - 2ibd} + \frac{B \tan^3(c+dx)}{2bd \tan(c+dx) - 2ibd} \\ \frac{3iBdx \tan(c+dx)}{2bd \tan(c+dx) + 2ibd} - \frac{3Bdx}{2bd \tan(c+dx) + 2ibd} - \frac{2B \log(\tan^2(c+dx)+1) \tan(c+dx)}{2bd \tan(c+dx) + 2ibd} - \frac{2iB \log(\tan^2(c+dx)+1)}{2bd \tan(c+dx) + 2ibd} + \frac{B \tan^3(c+dx)}{2bd \tan(c+dx) + 2ibd} \\ \frac{x(Ba+Bb \tan(c)) \tan^4(c)}{(a+b \tan(c))^2} \\ \frac{2Ba^4 \log(\frac{a}{b} + \tan(c+dx))}{2a^2b^3d+2b^5d} - \frac{2Ba^3b \tan(c+dx)}{2a^2b^3d+2b^5d} + \frac{Ba^2b^2 \tan^2(c+dx)}{2a^2b^3d+2b^5d} + \frac{2Bab^3 dx}{2a^2b^3d+2b^5d} - \frac{2Bab^3 \tan(c+dx)}{2a^2b^3d+2b^5d} - \frac{Bb^4 \log(\tan^2(c+dx)+1)}{2a^2b^3d+2b^5d} \end{array} \right.$$

input `integrate(tan(d*x+c)**4*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Piecewise((zoo*B*x*tan(c)**3, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (B*(x + tan(c + d*x)**3/(3*d) - tan(c + d*x)/d)/a, Eq(b, 0)), (-3*I*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - 3*B*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) - 2*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 2*I*B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) - 2*I*b*d) + B*tan(c + d*x)**3/(2*b*d*tan(c + d*x) - 2*I*b*d) + I*B*tan(c + d*x)**2/(2*b*d*tan(c + d*x) - 2*I*b*d) - B*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 4*I*B/(2*b*d*tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (3*I*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) - 3*B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - 2*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) - 2*I*B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2*I*b*d) + B*tan(c + d*x)**3/(2*b*d*tan(c + d*x) + 2*I*b*d) - I*B*tan(c + d*x)**2/(2*b*d*tan(c + d*x) + 2*I*b*d) - B*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) - 4*I*B/(2*b*d*tan(c + d*x) + 2*I*b*d), Eq(a, I*b)), (x*(B*a + B*b*tan(c))*tan(c)**4/(a + b*tan(c))**2, Eq(d, 0)), (2*B*a**4*log(a/b + tan(c + d*x))/(2*a**2*b**3*d + 2*b**5*d) - 2*B*a**3*b*tan(c + d*x)/(2*a**2*b**3*d + 2*b**5*d) + B*a**2*b**2*tan(c + d*x)**2/(2*a**2*b**3*d + 2*b**5*d) + 2*B*a*b**3*d*x/(2*a**2*b**3*d + 2*b**5*d) - 2*B*a*b**3*tan(c + d*x)/(2*a**2*b**3*d + 2*b**5*d) - B*b**4*log(tan(c + d*x)**2 + 1)/(2*a**2*b**3*d + 2*b**5*d) + B*b**4*tan(c + d*x)**2/(2*a**2*b**3*d + 2*b**5*d), True))`

3.306.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.02

$$\int \frac{\tan^4(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{\frac{2Ba^4 \log(b \tan(dx+c)+a)}{a^2 b^3 + b^5} + \frac{2(dx+c)Ba}{a^2 + b^2} - \frac{Bb \log(\tan(dx+c)^2 + 1)}{a^2 + b^2} + \frac{Bb \tan(dx+c)^2 - 2Ba \tan(dx+c)}{b^2}}{2d}$$

input `integrate(tan(d*x+c)^4*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `1/2*(2*B*a^4*log(b*tan(d*x + c) + a)/(a^2*b^3 + b^5) + 2*(d*x + c)*B*a/(a^2 + b^2) - B*b*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) + (B*b*tan(d*x + c)^2 - 2*B*a*tan(d*x + c))/b^2)/d`

3.306.8 Giac [A] (verification not implemented)

Time = 0.89 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.03

$$\int \frac{\tan^4(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{\frac{2Ba^4 \log(|b\tan(dx+c)+a|)}{a^2b^3+b^5} + \frac{2(dx+c)Ba}{a^2+b^2} - \frac{Bb \log(\tan(dx+c)^2+1)}{a^2+b^2} + \frac{Bb \tan(dx+c)^2 - 2Ba \tan(dx+c)}{b^2}}{2d}$$

input `integrate(tan(d*x+c)^4*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm m="giac")`

output `1/2*(2*B*a^4*log(abs(b*tan(d*x + c) + a))/(a^2*b^3 + b^5) + 2*(d*x + c)*B*a/(a^2 + b^2) - B*b*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) + (B*b*tan(d*x + c))^2 - 2*B*a*tan(d*x + c))/b^2)/d`

3.306.9 Mupad [B] (verification not implemented)

Time = 8.16 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.12

$$\int \frac{\tan^4(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \frac{B \tan(c+dx)^2}{2bd} - \frac{B \ln(\tan(c+dx) + 1i)}{2d(b+a1i)}$$

$$- \frac{Ba \tan(c+dx)}{b^2 d}$$

$$+ \frac{Ba^4 \ln(a+b\tan(c+dx))}{b^3 d (a^2+b^2)}$$

$$- \frac{B \ln(\tan(c+dx) - i) 1i}{2d(a+b1i)}$$

input `int((tan(c + d*x)^4*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`

output `(B*tan(c + d*x)^2)/(2*b*d) - (B*log(tan(c + d*x) + 1i))/(2*d*(a*1i + b)) - (B*log(tan(c + d*x) - 1i)*1i)/(2*d*(a + b*1i)) - (B*a*tan(c + d*x))/(b^2*d) + (B*a^4*log(a + b*tan(c + d*x)))/(b^3*d*(a^2 + b^2))`

3.307 $\int \frac{\tan^3(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.307.1 Optimal result	2925
3.307.2 Mathematica [C] (verified)	2925
3.307.3 Rubi [A] (verified)	2926
3.307.4 Maple [A] (verified)	2929
3.307.5 Fricas [A] (verification not implemented)	2929
3.307.6 Sympy [C] (verification not implemented)	2930
3.307.7 Maxima [A] (verification not implemented)	2931
3.307.8 Giac [A] (verification not implemented)	2931
3.307.9 Mupad [B] (verification not implemented)	2932

3.307.1 Optimal result

Integrand size = 34, antiderivative size = 83

$$\int \frac{\tan^3(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = -\frac{bBx}{a^2+b^2} + \frac{aB \log(\cos(c+dx))}{(a^2+b^2)d} - \frac{a^3B \log(a+b \tan(c+dx))}{b^2(a^2+b^2)d} + \frac{B \tan(c+dx)}{bd}$$

output `-b*B*x/(a^2+b^2)+a*B*ln(cos(d*x+c))/(a^2+b^2)/d-a^3*B*ln(a+b*tan(d*x+c))/b^2/(a^2+b^2)/d+B*tan(d*x+c)/b/d`

3.307.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.39 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.11

$$\int \frac{\tan^3(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = -\frac{B \left(\frac{\log(i-\tan(c+dx))}{a+ib} + \frac{\log(i+\tan(c+dx))}{a-ib} + \frac{2a^3 \log(a+b \tan(c+dx))}{b^2(a^2+b^2)} - \frac{2 \tan(c+dx)}{b} \right)}{2d}$$

input `Integrate[(Tan[c + d*x]^3*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2, x]`

output
$$\frac{-1/2*(B*(\text{Log}[I - \text{Tan}[c + d*x]]/(a + I*b) + \text{Log}[I + \text{Tan}[c + d*x]]/(a - I*b) + (2*a^3*\text{Log}[a + b*\text{Tan}[c + d*x]])/(b^2*(a^2 + b^2)) - (2*\text{Tan}[c + d*x])/b)}{d}$$

3.307.3 Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.07, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {2011, 3042, 4049, 25, 3042, 4109, 3042, 3956, 4100, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^3(c+dx)(aB + bB \tan(c+dx))}{(a + b \tan(c+dx))^2} dx \\ & \quad \downarrow \text{2011} \\ & B \int \frac{\tan^3(c+dx)}{a + b \tan(c+dx)} dx \\ & \quad \downarrow \text{3042} \\ & B \int \frac{\tan(c+dx)^3}{a + b \tan(c+dx)} dx \\ & \quad \downarrow \text{4049} \\ & B \left(\frac{\int \frac{-a \tan^2(c+dx) + b \tan(c+dx) + a}{a + b \tan(c+dx)} dx}{b} + \frac{\tan(c+dx)}{bd} \right) \\ & \quad \downarrow \text{25} \\ & B \left(\frac{\tan(c+dx)}{bd} - \frac{\int \frac{a \tan^2(c+dx) + b \tan(c+dx) + a}{a + b \tan(c+dx)} dx}{b} \right) \\ & \quad \downarrow \text{3042} \\ & B \left(\frac{\tan(c+dx)}{bd} - \frac{\int \frac{a \tan(c+dx)^2 + b \tan(c+dx) + a}{a + b \tan(c+dx)} dx}{b} \right) \\ & \quad \downarrow \text{4109} \end{aligned}$$

$$\begin{aligned}
& B \left(\frac{\tan(c+dx)}{bd} - \frac{ab \int \tan(c+dx) dx}{a^2+b^2} + \frac{a^3 \int \frac{\tan^2(c+dx)+1}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{b^2 x}{a^2+b^2} \right) \\
& \quad \downarrow \text{3042} \\
& B \left(\frac{\tan(c+dx)}{bd} - \frac{ab \int \tan(c+dx) dx}{a^2+b^2} + \frac{a^3 \int \frac{\tan(c+dx)^2+1}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{b^2 x}{a^2+b^2} \right) \\
& \quad \downarrow \text{3956} \\
& B \left(\frac{\tan(c+dx)}{bd} - \frac{a^3 \int \frac{\tan(c+dx)^2+1}{a+b \tan(c+dx)} dx}{a^2+b^2} - \frac{ab \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{b^2 x}{a^2+b^2} \right) \\
& \quad \downarrow \text{4100} \\
& B \left(\frac{\tan(c+dx)}{bd} - \frac{a^3 \int \frac{1}{a+b \tan(c+dx)} d(b \tan(c+dx))}{bd(a^2+b^2)} - \frac{ab \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{b^2 x}{a^2+b^2} \right) \\
& \quad \downarrow \text{16} \\
& B \left(\frac{\tan(c+dx)}{bd} - \frac{ab \log(\cos(c+dx))}{d(a^2+b^2)} + \frac{b^2 x}{a^2+b^2} + \frac{a^3 \log(a+b \tan(c+dx))}{bd(a^2+b^2)} \right)
\end{aligned}$$

input `Int[(Tan[c + d*x]^3*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `B*(-(((b^2*x)/(a^2 + b^2) - (a*b*Log[Cos[c + d*x]]))/((a^2 + b^2)*d) + (a^3 *Log[a + b*Tan[c + d*x]])/(b*(a^2 + b^2)*d))/b) + Tan[c + d*x]/(b*d))`

3.307.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_.))^(m_.)*((c_) + (d_.)*(v_.))^(n_.), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplrQ[c + d*x,
a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4049 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_.)])^(n_.), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m - 2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d*f*(m + n - 1))), x] + Simp[1/(d*(m + n - 1))
Int[(a + b*Tan[e + f*x])^(m - 3)*(c + d*Tan[e + f*x])^n*Simp[a^3*d*(m + n
- 1) - b^2*(b*c*(m - 2) + a*d*(1 + n)) + b*d*(m + n - 1)*(3*a^2 - b^2)*Tan[
e + f*x] - b^2*(b*c*(m - 2) - a*d*(3*m + 2*n - 4))*Tan[e + f*x]^2, x], x],
x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2
, 0] && NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && GtQ[m, 2] && (GeQ[n, -1] || I
ntegerQ[m]) && !(IGtQ[n, 2] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0]))
)`

rule 4100 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_.)*((A_) + (C_.)*tan[(e_.) +
(f_.)*(x_.)]^2), x_Symbol] := Simp[A/(b*f) Subst[Int[(a + x)^m, x], x, b*
Tan[e + f*x]], x] /; FreeQ[{a, b, e, f, A, C, m}, x] && EqQ[A, C]`

rule 4109 `Int[((A_) + (B_.)*tan[(e_.) + (f_.)*(x_.)] + (C_.)*tan[(e_.) + (f_.)*(x_.)]^2
) / ((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(a*A + b*B - a
C)(x/(a^2 + b^2)), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(
1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]), x], x] - Simp[(A*b - a*B - b*C)/(
a^2 + b^2) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, e, f, A, B, C}, x] &
& NeQ[A*b^2 - a*b*B + a^2*C, 0] && NeQ[a^2 + b^2, 0] && NeQ[A*b - a*B - b*C
, 0]`

3.307.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.96

method	result
derivativedivides	$\frac{B \left(\frac{\tan(dx+c)}{b} + \frac{-\frac{a \ln(1+\tan^2(dx+c))}{2} - b \arctan(\tan(dx+c)) - \frac{a^3 \ln(a+b \tan(dx+c))}{b^2(a^2+b^2)}}{a^2+b^2} \right)}{d}$
default	$\frac{B \left(\frac{\tan(dx+c)}{b} + \frac{-\frac{a \ln(1+\tan^2(dx+c))}{2} - b \arctan(\tan(dx+c)) - \frac{a^3 \ln(a+b \tan(dx+c))}{b^2(a^2+b^2)}}{a^2+b^2} \right)}{d}$
parallelrisch	$-\frac{2Bb^3dx + B \ln(1+\tan^2(dx+c))a^2 + 2B \ln(a+b \tan(dx+c))a^3 - 2Ba^2b \tan(dx+c) - 2Bb^3 \tan(dx+c)}{2(a^2+b^2)b^2d}$
norman	$\frac{\frac{B(\tan^2(dx+c))}{d} - \frac{Ba^2}{db^2} - \frac{bBa^2x}{a^2+b^2} - \frac{b^2Bx \tan(dx+c)}{a^2+b^2}}{a+b \tan(dx+c)} - \frac{Ba \ln(1+\tan^2(dx+c))}{2d(a^2+b^2)} - \frac{a^3 B \ln(a+b \tan(dx+c))}{b^2(a^2+b^2)d}$
risch	$-\frac{ixB}{ib-a} - \frac{2iaBx}{b^2} - \frac{2iBac}{b^2d} + \frac{2ia^3Bx}{b^2(a^2+b^2)} + \frac{2ia^3Bc}{d(a^2+b^2)b^2} + \frac{2iB}{db(e^{2i(dx+c)}+1)} + \frac{\ln(e^{2i(dx+c)}+1)Ba}{b^2d} - \frac{a^3 \ln(e^{2i(dx+c)}+1)}{b^2d}$

input `int(tan(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNV ERBOSE)`

output `1/d*B*(tan(d*x+c)/b+1/(a^2+b^2)*(-1/2*a*ln(1+tan(d*x+c)^2)-b*arctan(tan(d*x+c)))-1/b^2*a^3/(a^2+b^2)*ln(a+b*tan(d*x+c)))`

3.307.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.43

$$\int \frac{\tan^3(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx =$$

$$-\frac{2Bb^3dx + Ba^3 \log\left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2 + 1}\right) - (Ba^3 + Bab^2) \log\left(\frac{1}{\tan(dx+c)^2 + 1}\right) - 2(Ba^2b + Bb^3) \tan(dx+c)}{2(a^2b^2 + b^4)d}$$

input `integrate(tan(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

output `-1/2*(2*B*b^3*d*x + B*a^3*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - (B*a^3 + B*a*b^2)*log(1/(tan(d*x + c)^2 + 1)) - 2*(B*a^2*b + B*b^3)*tan(d*x + c))/((a^2*b^2 + b^4)*d)`

3.307.
$$\int \frac{\tan^3(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.307.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.04 (sec) , antiderivative size = 660, normalized size of antiderivative = 7.95

$$\int \frac{\tan^3(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \begin{cases} \tilde{\infty} Bx \tan^2(c) \\ B \left(\frac{-\frac{\log(\tan^2(c+dx)+1)}{2d} + \frac{\tan^2(c+dx)}{2d}}{a} \right) \\ -\frac{3Bdx \tan(c+dx)}{2bd \tan(c+dx)-2ibd} + \frac{3iBdx}{2bd \tan(c+dx)-2ibd} + \frac{iB \log(\tan^2(c+dx)+1) \tan(c+dx)}{2bd \tan(c+dx)-2ibd} + \frac{B \log(\tan^2(c+dx)+1)}{2bd \tan(c+dx)-2ibd} + \frac{2B \tan^2(c+dx)}{2bd \tan(c+dx)-2ibd} \\ -\frac{3Bdx \tan(c+dx)}{2bd \tan(c+dx)+2ibd} - \frac{3iBdx}{2bd \tan(c+dx)+2ibd} - \frac{iB \log(\tan^2(c+dx)+1) \tan(c+dx)}{2bd \tan(c+dx)+2ibd} + \frac{B \log(\tan^2(c+dx)+1)}{2bd \tan(c+dx)+2ibd} + \frac{2B \tan^2(c+dx)}{2bd \tan(c+dx)+2ibd} \\ \frac{x(Ba+Bb \tan(c)) \tan^3(c)}{(a+b \tan(c))^2} \\ -\frac{2Ba^3 \log(\frac{a}{b} + \tan(c+dx))}{2a^2b^2d+2b^4d} + \frac{2Ba^2b \tan(c+dx)}{2a^2b^2d+2b^4d} - \frac{Bab^2 \log(\tan^2(c+dx)+1)}{2a^2b^2d+2b^4d} - \frac{2Bb^3dx}{2a^2b^2d+2b^4d} + \frac{2Bb^3 \tan(c+dx)}{2a^2b^2d+2b^4d} \end{cases}$$

input `integrate(tan(d*x+c)**3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Piecewise((zoo*B*x*tan(c)**2, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (B*(-log(tan(c + d*x)**2 + 1)/(2*d) + tan(c + d*x)**2/(2*d))/a, Eq(b, 0)), (-3*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 3*I*B*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) + I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 2*B*tan(c + d*x)**2/(2*b*d*tan(c + d*x) - 2*I*b*d) + 2*I*B*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 5*B/(2*b*d*tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (-3*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) - 3*I*B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2*I*b*d) + 2*B*tan(c + d*x)**2/(2*b*d*tan(c + d*x) + 2*I*b*d) - 2*I*B*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + 5*B/(2*b*d*tan(c + d*x) + 2*I*b*d), Eq(a, I*b)), (x*(B*a + B*b*tan(c))*tan(c)**3/(a + b*tan(c))**2, Eq(d, 0)), (-2*B*a**3*log(a/b + tan(c + d*x))/(2*a**2*b**2*d + 2*b**4*d) + 2*B*a**2*b*tan(c + d*x)/(2*a**2*b**2*d + 2*b**4*d) - B*a*b**2*log(tan(c + d*x)**2 + 1)/(2*a**2*b**2*d + 2*b**4*d) - 2*B*b**3*d*x/(2*a**2*b**2*d + 2*b**4*d) + 2*B*b**3*tan(c + d*x)/(2*a**2*b**2*d + 2*b**4*d), True))`

3.307.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.07

$$\int \frac{\tan^3(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= -\frac{\frac{2Ba^3 \log(b\tan(dx+c)+a)}{a^2b^2+b^4} + \frac{2(dx+c)Bb}{a^2+b^2} + \frac{Ba \log(\tan(dx+c)^2+1)}{a^2+b^2} - \frac{2B\tan(dx+c)}{b}}{2d}$$

```
input integrate(tan(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")
```

```
output -1/2*(2*B*a^3*log(b*tan(d*x + c) + a)/(a^2*b^2 + b^4) + 2*(d*x + c)*B*b/(a^2 + b^2) + B*a*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) - 2*B*tan(d*x + c)/b)/d
```

3.307.8 Giac [A] (verification not implemented)

Time = 0.67 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.08

$$\int \frac{\tan^3(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= -\frac{\frac{2Ba^3 \log(|b\tan(dx+c)+a|)}{a^2b^2+b^4} + \frac{2(dx+c)Bb}{a^2+b^2} + \frac{Ba \log(\tan(dx+c)^2+1)}{a^2+b^2} - \frac{2B\tan(dx+c)}{b}}{2d}$$

```
input integrate(tan(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")
```

```
output -1/2*(2*B*a^3*log(abs(b*tan(d*x + c) + a))/(a^2*b^2 + b^4) + 2*(d*x + c)*B*b/(a^2 + b^2) + B*a*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) - 2*B*tan(d*x + c)/b)/d
```

3.307.9 Mupad [B] (verification not implemented)

Time = 7.21 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.18

$$\int \frac{\tan^3(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \frac{B\tan(c+dx)}{bd} - \frac{B\ln(\tan(c+dx)+1i)}{2d(a-b1i)} - \frac{Ba^3\ln(a+b\tan(c+dx))}{b^2d(a^2+b^2)} - \frac{B\ln(\tan(c+dx)-i)1i}{2d(-b+a1i)}$$

input `int((tan(c + d*x)^3*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`output `(B*tan(c + d*x))/(b*d) - (B*log(tan(c + d*x) + 1i))/(2*d*(a - b*1i)) - (B*log(tan(c + d*x) - 1i)*1i)/(2*d*(a*1i - b)) - (B*a^3*log(a + b*tan(c + d*x)))/(b^2*d*(a^2 + b^2))`

3.308 $\int \frac{\tan^2(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.308.1 Optimal result 2933
 3.308.2 Mathematica [C] (verified) 2933
 3.308.3 Rubi [A] (verified) 2934
 3.308.4 Maple [A] (verified) 2936
 3.308.5 Fricas [A] (verification not implemented) 2937
 3.308.6 Sympy [C] (verification not implemented) 2937
 3.308.7 Maxima [A] (verification not implemented) 2938
 3.308.8 Giac [A] (verification not implemented) 2938
 3.308.9 Mupad [B] (verification not implemented) 2939

3.308.1 Optimal result

Integrand size = 34, antiderivative size = 81

$$\int \frac{\tan^2(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = -\frac{aBx}{b^2} + \frac{a^3Bx}{b^2(a^2+b^2)} - \frac{B \log(\cos(c+dx))}{bd} + \frac{a^2B \log(a \cos(c+dx) + b \sin(c+dx))}{b(a^2+b^2)d}$$

output

```
-a*B*x/b^2+a^3*B*x/b^2/(a^2+b^2)-B*ln(cos(d*x+c))/b/d+a^2*B*ln(a*cos(d*x+c)+b*sin(d*x+c))/b/(a^2+b^2)/d
```

3.308.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.98

$$\int \frac{\tan^2(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \frac{B(b(ia+b) \log(i-\tan(c+dx)) + b(-ia+b) \log(i+\tan(c+dx)) + 2a^2 \log(a+b \tan(c+dx)))}{2b(a^2+b^2)d}$$

input

```
Integrate[(Tan[c + d*x]^2*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2, x]
```

output $(B*(b*(I*a + b)*\text{Log}[I - \text{Tan}[c + d*x]] + b*((-I)*a + b)*\text{Log}[I + \text{Tan}[c + d*x]] + 2*a^2*\text{Log}[a + b*\text{Tan}[c + d*x]])/(2*b*(a^2 + b^2)*d)$

3.308.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.95, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2011, 3042, 4024, 3042, 3956, 3965, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^2(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx \\ & \quad \downarrow \text{2011} \\ & B \int \frac{\tan^2(c+dx)}{a+b\tan(c+dx)} dx \\ & \quad \downarrow \text{3042} \\ & B \int \frac{\tan(c+dx)^2}{a+b\tan(c+dx)} dx \\ & \quad \downarrow \text{4024} \\ & B \left(\frac{a^2 \int \frac{1}{a+b\tan(c+dx)} dx}{b^2} + \frac{\int \tan(c+dx) dx}{b} - \frac{ax}{b^2} \right) \\ & \quad \downarrow \text{3042} \\ & B \left(\frac{a^2 \int \frac{1}{a+b\tan(c+dx)} dx}{b^2} + \frac{\int \tan(c+dx) dx}{b} - \frac{ax}{b^2} \right) \\ & \quad \downarrow \text{3956} \\ & B \left(\frac{a^2 \int \frac{1}{a+b\tan(c+dx)} dx}{b^2} - \frac{ax}{b^2} - \frac{\log(\cos(c+dx))}{bd} \right) \\ & \quad \downarrow \text{3965} \\ & B \left(\frac{a^2 \left(\frac{b \int \frac{b-a\tan(c+dx)}{a+b\tan(c+dx)} dx}{a^2+b^2} + \frac{ax}{a^2+b^2} \right)}{b^2} - \frac{ax}{b^2} - \frac{\log(\cos(c+dx))}{bd} \right) \end{aligned}$$

$$\begin{array}{c}
 \downarrow \text{3042} \\
 B \left(\frac{a^2 \left(\frac{b \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2+b^2} + \frac{ax}{a^2+b^2} \right)}{b^2} - \frac{ax}{b^2} - \frac{\log(\cos(c+dx))}{bd} \right) \\
 \downarrow \text{4013} \\
 B \left(\frac{a^2 \left(\frac{b \log(a \cos(c+dx)+b \sin(c+dx))}{d(a^2+b^2)} + \frac{ax}{a^2+b^2} \right)}{b^2} - \frac{ax}{b^2} - \frac{\log(\cos(c+dx))}{bd} \right)
 \end{array}$$

```
input Int[(Tan[c + d*x]^2*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]
```

```
output B*(-((a*x)/b^2) - Log[Cos[c + d*x]]/(b*d) + (a^2*((a*x)/(a^2 + b^2) + (b*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d)))/b^2)
```

3.308.3.1 Defintions of rubi rules used

```
rule 2011 Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :=
  Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x, a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3956 Int[tan[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

```
rule 3965 Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[a*(x/(a^2 + b^2)), x] + Simp[b/(a^2 + b^2) Int[(b - a*Tan[c + d*x])/(a + b*Tan[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]
```

```
rule 4013 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)]) / ((a_) + (b_)*tan[(e_) + (f_)*
(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Si
n[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

```
rule 4024 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])^2 / ((a_) + (b_)*tan[(e_) + (
f_)*(x_)]), x_Symbol] := Simp[d*(2*b*c - a*d)*(x/b^2), x] + (Simp[d^2/b I
nt[Tan[e + f*x], x], x] + Simp[(b*c - a*d)^2/b^2 Int[1/(a + b*Tan[e + f*x
]), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2
+ b^2, 0]
```

3.308.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.73

method	result	size
parallelrisch	$\frac{-2Babdx + B \ln(1 + \tan^2(dx+c))b^2 + 2B \ln(a+b \tan(dx+c))a^2}{2bd(a^2+b^2)}$	59
derivativedivides	$\frac{B \left(\frac{b \ln(1 + \tan^2(dx+c))}{2} - \frac{a \arctan(\tan(dx+c))}{a^2+b^2} + \frac{a^2 \ln(a+b \tan(dx+c))}{(a^2+b^2)b} \right)}{d}$	69
default	$\frac{B \left(\frac{b \ln(1 + \tan^2(dx+c))}{2} - \frac{a \arctan(\tan(dx+c))}{a^2+b^2} + \frac{a^2 \ln(a+b \tan(dx+c))}{(a^2+b^2)b} \right)}{d}$	69
norman	$\frac{-\frac{B a^2 x}{a^2+b^2} - \frac{b B a x \tan(dx+c)}{a^2+b^2}}{a+b \tan(dx+c)} + \frac{B a^2 \ln(a+b \tan(dx+c))}{bd(a^2+b^2)} + \frac{B b \ln(1 + \tan^2(dx+c))}{2d(a^2+b^2)}$	111
risch	$\frac{x B}{ib-a} + \frac{2i B x}{b} + \frac{2i B c}{bd} - \frac{2ia^2 B x}{b(a^2+b^2)} - \frac{2ia^2 B c}{bd(a^2+b^2)} - \frac{\ln(e^{2i(dx+c)}+1) B}{bd} + \frac{a^2 \ln\left(e^{2i(dx+c)} - \frac{ib+a}{ib-a}\right) B}{bd(a^2+b^2)}$	147

```
input int(tan(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNV
ERBOSE)
```

```
output 1/2*(-2*B*a*b*d*x+B*ln(1+tan(d*x+c)^2)*b^2+2*B*ln(a+b*tan(d*x+c))*a^2)/b/d
/(a^2+b^2)
```

3.308.
$$\int \frac{\tan^2(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.308.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= -\frac{2Babdx - Ba^2 \log\left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2 + 1}\right) + (Ba^2 + Bb^2) \log\left(\frac{1}{\tan(dx+c)^2 + 1}\right)}{2(a^2b + b^3)d}$$

input `integrate(tan(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm m="fricas")`

output `-1/2*(2*B*a*b*d*x - B*a^2*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) + (B*a^2 + B*b^2)*log(1/(tan(d*x + c)^2 + 1)))/(a^2*b + b^3)*d`

3.308.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.74 (sec) , antiderivative size = 442, normalized size of antiderivative = 5.46

$$\int \frac{\tan^2(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \left\{ \begin{array}{l} \tilde{\infty} Bx \tan(c) \\ \frac{B\left(-x + \frac{\tan(c+dx)}{d}\right)}{a} \\ \frac{iBdx \tan(c+dx)}{2bd \tan(c+dx) - 2ibd} + \frac{Bdx}{2bd \tan(c+dx) - 2ibd} + \frac{B \log(\tan^2(c+dx)+1) \tan(c+dx)}{2bd \tan(c+dx) - 2ibd} - \frac{iB \log(\tan^2(c+dx)+1)}{2bd \tan(c+dx) - 2ibd} - \frac{iB}{2bd \tan(c+dx) - 2ibd} \\ -\frac{iBdx \tan(c+dx)}{2bd \tan(c+dx) + 2ibd} + \frac{Bdx}{2bd \tan(c+dx) + 2ibd} + \frac{B \log(\tan^2(c+dx)+1) \tan(c+dx)}{2bd \tan(c+dx) + 2ibd} + \frac{iB \log(\tan^2(c+dx)+1)}{2bd \tan(c+dx) + 2ibd} + \frac{iB}{2bd \tan(c+dx) + 2ibd} \\ \frac{x(Ba+Bb \tan(c)) \tan^2(c)}{(a+b \tan(c))^2} \\ \frac{2Ba^2 \log\left(\frac{a}{b} + \tan(c+dx)\right)}{2a^2bd + 2b^3d} - \frac{2Babdx}{2a^2bd + 2b^3d} + \frac{Bb^2 \log(\tan^2(c+dx)+1)}{2a^2bd + 2b^3d} \end{array} \right.$$

input `integrate(tan(d*x+c)**2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Piecewise((zoo*B*x*tan(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (B*(-x + tan(c + d*x))/d)/a, Eq(b, 0)), (I*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + B*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) + B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*B/(2*b*d*tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (-I*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) + B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*B/(2*b*d*tan(c + d*x) + 2*I*b*d), Eq(a, I*b)), (x*(B*a + B*b*tan(c))*tan(c)**2/(a + b*tan(c))**2, Eq(d, 0)), (2*B*a**2*log(a/b + tan(c + d*x))/(2*a**2*b*d + 2*b**3*d) - 2*B*a*b*d*x/(2*a**2*b*d + 2*b**3*d) + B*b**2*log(tan(c + d*x)**2 + 1)/(2*a**2*b*d + 2*b**3*d), True))`

3.308.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.93

$$\int \frac{\tan^2(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{\frac{2Ba^2 \log(b \tan(dx+c)+a)}{a^2b+b^3} - \frac{2(dx+c)Ba}{a^2+b^2} + \frac{Bb \log(\tan(dx+c)^2+1)}{a^2+b^2}}{2d}$$

input `integrate(tan(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `1/2*(2*B*a^2*log(b*tan(d*x + c) + a)/(a^2*b + b^3) - 2*(d*x + c)*B*a/(a^2 + b^2) + B*b*log(tan(d*x + c)^2 + 1)/(a^2 + b^2))/d`

3.308.8 Giac [A] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.94

$$\int \frac{\tan^2(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{\frac{2Ba^2 \log(b \tan(dx+c)+a)}{a^2b+b^3} - \frac{2(dx+c)Ba}{a^2+b^2} + \frac{Bb \log(\tan(dx+c)^2+1)}{a^2+b^2}}{2d}$$

input `integrate(tan(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm
m="giac")`

output `1/2*(2*B*a^2*log(abs(b*tan(d*x + c) + a))/(a^2*b + b^3) - 2*(d*x + c)*B*a/
(a^2 + b^2) + B*b*log(tan(d*x + c)^2 + 1)/(a^2 + b^2))/d`

3.308.9 Mupad [B] (verification not implemented)

Time = 7.36 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00

$$\int \frac{\tan^2(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \frac{B \ln(\tan(c+dx) + 1i)}{2d(b+a1i)} + \frac{B a^2 \ln(a+b\tan(c+dx))}{bd(a^2+b^2)} + \frac{B \ln(\tan(c+dx) - i) 1i}{2d(a+b1i)}$$

input `int((tan(c + d*x)^2*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`

output `(B*log(tan(c + d*x) - 1i)*1i)/(2*d*(a + b*1i)) + (B*log(tan(c + d*x) + 1i)
)/(2*d*(a*1i + b)) + (B*a^2*log(a + b*tan(c + d*x)))/(b*d*(a^2 + b^2))`

3.309
$$\int \frac{\tan(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.309.1 Optimal result 2940
 3.309.2 Mathematica [C] (verified) 2940
 3.309.3 Rubi [A] (verified) 2941
 3.309.4 Maple [A] (verified) 2942
 3.309.5 Fricas [A] (verification not implemented) 2943
 3.309.6 Sympy [C] (verification not implemented) 2943
 3.309.7 Maxima [A] (verification not implemented) 2944
 3.309.8 Giac [A] (verification not implemented) 2944
 3.309.9 Mupad [B] (verification not implemented) 2945

3.309.1 Optimal result

Integrand size = 32, antiderivative size = 48

$$\int \frac{\tan(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^2} dx = \frac{bBx}{a^2 + b^2} - \frac{aB \log(a \cos(c + dx) + b \sin(c + dx))}{(a^2 + b^2) d}$$

output `b*B*x/(a^2+b^2)-a*B*ln(a*cos(d*x+c)+b*sin(d*x+c))/(a^2+b^2)/d`

3.309.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.40

$$\int \frac{\tan(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^2} dx = \frac{B(2(-ia + b)(c + dx) + 2ia \arctan(\tan(c + dx)) - a \log((a \cos(c + dx) + b \sin(c + dx))^2))}{2(a^2 + b^2) d}$$

input `Integrate[(Tan[c + d*x]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `(B*(2*((-I)*a + b)*(c + d*x) + (2*I)*a*ArcTan[Tan[c + d*x]] - a*Log[(a*Cos[c + d*x] + b*Sin[c + d*x])^2]))/(2*(a^2 + b^2)*d)`

3.309.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {2011, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(c+dx)(aB + bB \tan(c+dx))}{(a + b \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{\tan(c+dx)}{a + b \tan(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{\tan(c+dx)}{a + b \tan(c+dx)} dx \\
 & \quad \downarrow \text{4014} \\
 & B \left(\frac{bx}{a^2 + b^2} - \frac{a \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2 + b^2} \right) \\
 & \quad \downarrow \text{3042} \\
 & B \left(\frac{bx}{a^2 + b^2} - \frac{a \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2 + b^2} \right) \\
 & \quad \downarrow \text{4013} \\
 & B \left(\frac{bx}{a^2 + b^2} - \frac{a \log(a \cos(c+dx) + b \sin(c+dx))}{d(a^2 + b^2)} \right)
 \end{aligned}$$

input `Int[(Tan[c + d*x]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `B*((b*x)/(a^2 + b^2) - (a*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d))`

3.309.3.1 Defintions of rubi rules used

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplrQ[c + d*x,
a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*
(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Si
n[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*
(x_)])*(x_)), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a
*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[a*c + b*d, 0]`

3.309.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.06

method	result	size
parallelrisch	$\frac{2Bbdx+B \ln(1+\tan^2(dx+c))a-2B \ln(a+b \tan(dx+c))a}{2d(a^2+b^2)}$	51
derivativedivides	$\frac{B \left(\frac{\frac{a \ln(1+\tan^2(dx+c))}{2} + b \arctan(\tan(dx+c))}{a^2+b^2} - \frac{a \ln(a+b \tan(dx+c))}{a^2+b^2} \right)}{d}$	64
default	$\frac{B \left(\frac{\frac{a \ln(1+\tan^2(dx+c))}{2} + b \arctan(\tan(dx+c))}{a^2+b^2} - \frac{a \ln(a+b \tan(dx+c))}{a^2+b^2} \right)}{d}$	64
risch	$\frac{ixB}{ib-a} + \frac{2iBxa}{a^2+b^2} + \frac{2iBac}{d(a^2+b^2)} - \frac{\ln(e^{2i(dx+c)} - \frac{ib+a}{ib-a})Ba}{d(a^2+b^2)}$	95
norman	$\frac{\frac{bBax}{a^2+b^2} + \frac{b^2 Bx \tan(dx+c)}{a^2+b^2}}{a+b \tan(dx+c)} + \frac{Ba \ln(1+\tan^2(dx+c))}{2d(a^2+b^2)} - \frac{Ba \ln(a+b \tan(dx+c))}{d(a^2+b^2)}$	105

3.309. $\int \frac{\tan(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

```
input int(tan(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNVER
BOSE)
```

```
output 1/2*(2*B*b*d*x+B*ln(1+tan(d*x+c)^2)*a-2*B*ln(a+b*tan(d*x+c))*a)/d/(a^2+b^2
)
```

3.309.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.35

$$\int \frac{\tan(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \frac{2Bbdx - Ba \log\left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2 + 1}\right)}{2(a^2 + b^2)d}$$

```
input integrate(tan(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm=
"fricas")
```

```
output 1/2*(2*B*b*d*x - B*a*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(
tan(d*x + c)^2 + 1)))/((a^2 + b^2)*d)
```

3.309.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.58 (sec) , antiderivative size = 282, normalized size of antiderivative = 5.88

$$\int \frac{\tan(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \begin{cases} \tilde{\infty} Bx & \text{for } a = 0 \wedge b = 0 \wedge d = 0 \\ \frac{B \log(\tan^2(c+dx)+1)}{2ad} & \text{for } b = 0 \\ \frac{Bdx \tan(c+dx)}{2bd \tan(c+dx)-2ibd} - \frac{iBdx}{2bd \tan(c+dx)-2ibd} - \frac{B}{2bd \tan(c+dx)-2ibd} & \text{for } a = -ib \\ \frac{Bdx \tan(c+dx)}{2bd \tan(c+dx)+2ibd} + \frac{iBdx}{2bd \tan(c+dx)+2ibd} - \frac{B}{2bd \tan(c+dx)+2ibd} & \text{for } a = ib \\ \frac{x(Ba+Bb \tan(c)) \tan(c)}{(a+b \tan(c))^2} & \text{for } d = 0 \\ -\frac{2Ba \log\left(\frac{a}{b} + \tan(c+dx)\right)}{2a^2d+2b^2d} + \frac{Ba \log(\tan^2(c+dx)+1)}{2a^2d+2b^2d} + \frac{2Bbdx}{2a^2d+2b^2d} & \text{otherwise} \end{cases}$$

input `integrate(tan(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Piecewise((zoo*B*x, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (B*log(tan(c + d*x)**2 + 1)/(2*a*d), Eq(b, 0)), (B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*B*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) - B/(2*b*d*tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - B/(2*b*d*tan(c + d*x) + 2*I*b*d), Eq(a, I*b)), (x*(B*a + B*b*tan(c))*tan(c)/(a + b*tan(c))**2, Eq(d, 0)), (-2*B*a*log(a/b + tan(c + d*x))/(2*a**2*d + 2*b**2*d) + B*a*log(tan(c + d*x)**2 + 1)/(2*a**2*d + 2*b**2*d) + 2*B*b*d*x/(2*a**2*d + 2*b**2*d), True))`

3.309.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.48

$$\int \frac{\tan(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{\frac{2(dx+c)Bb}{a^2+b^2} - \frac{2Ba\log(b\tan(dx+c)+a)}{a^2+b^2} + \frac{Ba\log(\tan(dx+c)^2+1)}{a^2+b^2}}{2d}$$

input `integrate(tan(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `1/2*(2*(d*x + c)*B*b/(a^2 + b^2) - 2*B*a*log(b*tan(d*x + c) + a)/(a^2 + b^2) + B*a*log(tan(d*x + c)^2 + 1)/(a^2 + b^2))/d`

3.309.8 Giac [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.58

$$\int \frac{\tan(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= -\frac{\frac{2Bab\log(|b\tan(dx+c)+a|)}{a^2b+b^3} - \frac{2(dx+c)Bb}{a^2+b^2} - \frac{Ba\log(\tan(dx+c)^2+1)}{a^2+b^2}}{2d}$$

input `integrate(tan(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output
$$-1/2*(2*B*a*b*\log(\text{abs}(b*\tan(dx + c) + a))/(a^2*b + b^3) - 2*(dx + c)*B*b/(a^2 + b^2) - B*a*\log(\tan(dx + c)^2 + 1)/(a^2 + b^2))/d$$

3.309.9 Mupad [B] (verification not implemented)

Time = 8.75 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.65

$$\int \frac{\tan(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^2} dx = \frac{B \ln(\tan(c + dx) + 1i)}{2d(a - b1i)} - \frac{Ba \ln(a + b \tan(c + dx))}{d(a^2 + b^2)} + \frac{B \ln(\tan(c + dx) - i) 1i}{2d(-b + a1i)}$$

input
$$\text{int}((\tan(c + d*x)*(B*a + B*b*\tan(c + d*x)))/(a + b*\tan(c + d*x))^2,x)$$

output
$$(B*\log(\tan(c + d*x) + 1i))/(2*d*(a - b*1i)) + (B*\log(\tan(c + d*x) - 1i)*1i)/(2*d*(a*1i - b)) - (B*a*\log(a + b*\tan(c + d*x)))/(d*(a^2 + b^2))$$

3.310 $\int \frac{aB+bB \tan(c+dx)}{(a+b \tan(c+dx))^2} dx$

3.310.1 Optimal result	2946
3.310.2 Mathematica [C] (verified)	2946
3.310.3 Rubi [A] (verified)	2947
3.310.4 Maple [A] (verified)	2948
3.310.5 Fricas [A] (verification not implemented)	2949
3.310.6 Sympy [C] (verification not implemented)	2949
3.310.7 Maxima [A] (verification not implemented)	2950
3.310.8 Giac [A] (verification not implemented)	2950
3.310.9 Mupad [B] (verification not implemented)	2951

3.310.1 Optimal result

Integrand size = 26, antiderivative size = 47

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^2} dx = \frac{aBx}{a^2 + b^2} + \frac{bB \log(a \cos(c + dx) + b \sin(c + dx))}{(a^2 + b^2) d}$$

output `a*B*x/(a^2+b^2)+b*B*ln(a*cos(d*x+c)+b*sin(d*x+c))/(a^2+b^2)/d`

3.310.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.03 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.64

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^2} dx = \frac{B((-ia - b) \log(i - \tan(c + dx)) + i(a + ib) \log(i + \tan(c + dx)) + 2b \log(a + b \tan(c + dx)))}{2(a^2 + b^2) d}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(a + b*Tan[c + d*x])^2,x]`

output `(B*(((-I)*a - b)*Log[I - Tan[c + d*x]] + I*(a + I*b)*Log[I + Tan[c + d*x]] + 2*b*Log[a + b*Tan[c + d*x]]))/(2*(a^2 + b^2)*d)`

3.310.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$, Rules used = {2011, 3042, 3965, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{1}{a + b \tan(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{a + b \tan(c + dx)} dx \\
 & \quad \downarrow \text{3965} \\
 & B \left(\frac{b \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2 + b^2} + \frac{ax}{a^2 + b^2} \right) \\
 & \quad \downarrow \text{3042} \\
 & B \left(\frac{b \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a^2 + b^2} + \frac{ax}{a^2 + b^2} \right) \\
 & \quad \downarrow \text{4013} \\
 & B \left(\frac{b \log(a \cos(c + dx) + b \sin(c + dx))}{d(a^2 + b^2)} + \frac{ax}{a^2 + b^2} \right)
 \end{aligned}$$

input `Int[(a*B + b*B*Tan[c + d*x])/(a + b*Tan[c + d*x])^2,x]`

output `B*((a*x)/(a^2 + b^2) + (b*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d))`

3.310.3.1 Defintions of rubi rules used

```
rule 2011 Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :=
  Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
  a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3965 Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[a*(x/(a^
2 + b^2)), x] + Simp[b/(a^2 + b^2) Int[(b - a*Tan[c + d*x])/(a + b*Tan[c
+ d*x]), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0]
```

```
rule 4013 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*
(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Si
n[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]
```

3.310.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.09

method	result	size
parallelrisch	$-\frac{-2Bxad+B\ln(1+\tan^2(dx+c))b-2Bb\ln(a+b\tan(dx+c))}{2d(a^2+b^2)}$	51
derivativdivides	$\frac{B\left(\frac{-\frac{b\ln(1+\tan^2(dx+c))}{2}+a\arctan(\tan(dx+c))+\frac{b\ln(a+b\tan(dx+c))}{a^2+b^2}}{a^2+b^2}\right)}{d}$	63
default	$\frac{B\left(\frac{-\frac{b\ln(1+\tan^2(dx+c))}{2}+a\arctan(\tan(dx+c))+\frac{b\ln(a+b\tan(dx+c))}{a^2+b^2}}{a^2+b^2}\right)}{d}$	63
risch	$-\frac{xB}{ib-a}-\frac{2ibBx}{a^2+b^2}-\frac{2ibBc}{d(a^2+b^2)}+\frac{b\ln\left(e^{2i(dx+c)}-\frac{ib+a}{ib-a}\right)B}{d(a^2+b^2)}$	93
norman	$\frac{\frac{B}{a^2+b^2}+\frac{bBax\tan(dx+c)}{a^2+b^2}}{a+b\tan(dx+c)}+\frac{Bb\ln(a+b\tan(dx+c))}{d(a^2+b^2)}-\frac{Bb\ln(1+\tan^2(dx+c))}{2d(a^2+b^2)}$	104

input `int((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output
$$-1/2*(-2*B*x*a*d+B*\ln(1+\tan(d*x+c)^2)*b-2*B*b*\ln(a+b*\tan(d*x+c)))/d/(a^2+b^2)$$

3.310.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.36

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^2} dx = \frac{2Badx + Bb \log\left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2 + 1}\right)}{2(a^2 + b^2)d}$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

output
$$1/2*(2*B*a*d*x + B*b*\log((b^2*\tan(d*x + c)^2 + 2*a*b*\tan(d*x + c) + a^2)/(\tan(d*x + c)^2 + 1)))/((a^2 + b^2)*d)$$

3.310.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.57 (sec) , antiderivative size = 272, normalized size of antiderivative = 5.79

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^2} dx = \begin{cases} \frac{\infty Bx}{\tan(c)} & \text{for } a = 0 \wedge b = 0 \wedge d = 0 \\ \frac{Bx}{a} & \text{for } b = 0 \\ \frac{iBdx \tan(c+dx)}{2bd \tan(c+dx) - 2ibd} + \frac{Bdx}{2bd \tan(c+dx) - 2ibd} + \frac{iB}{2bd \tan(c+dx) - 2ibd} & \text{for } a = -ib \\ -\frac{iBdx \tan(c+dx)}{2bd \tan(c+dx) + 2ibd} + \frac{Bdx}{2bd \tan(c+dx) + 2ibd} - \frac{iB}{2bd \tan(c+dx) + 2ibd} & \text{for } a = ib \\ \frac{x(Ba + Bb \tan(c))}{(a + b \tan(c))^2} & \text{for } d = 0 \\ \frac{2Badx}{2a^2d + 2b^2d} + \frac{2Bb \log\left(\frac{a}{b} + \tan(c+dx)\right)}{2a^2d + 2b^2d} - \frac{Bb \log(\tan^2(c+dx) + 1)}{2a^2d + 2b^2d} & \text{otherwise} \end{cases}$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Piecewise((zoo*B*x/tan(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (B*x/a, Eq(b, 0)), (I*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + B*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) + I*B/(2*b*d*tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (-I*B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) - I*B/(2*b*d*tan(c + d*x) + 2*I*b*d), Eq(a, I*b)), (x*(B*a + B*b*tan(c))/(a + b*tan(c))**2, Eq(d, 0)), (2*B*a*d*x/(2*a**2*d + 2*b**2*d) + 2*B*b*log(a/b + tan(c + d*x))/(2*a**2*d + 2*b**2*d) - B*b*log(tan(c + d*x)**2 + 1)/(2*a**2*d + 2*b**2*d), True))`

3.310.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.53

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^2} dx = \frac{\frac{2(dx+c)Ba}{a^2+b^2} + \frac{2Bb \log(b \tan(dx+c)+a)}{a^2+b^2} - \frac{Bb \log(\tan(dx+c)^2+1)}{a^2+b^2}}{2d}$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `1/2*(2*(d*x + c)*B*a/(a^2 + b^2) + 2*B*b*log(b*tan(d*x + c) + a)/(a^2 + b^2) - B*b*log(tan(d*x + c)^2 + 1)/(a^2 + b^2))/d`

3.310.8 Giac [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.64

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^2} dx = \frac{\frac{2Bb^2 \log(|b \tan(dx+c)+a|)}{a^2b+b^3} + \frac{2(dx+c)Ba}{a^2+b^2} - \frac{Bb \log(\tan(dx+c)^2+1)}{a^2+b^2}}{2d}$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output `1/2*(2*B*b^2*log(abs(b*tan(d*x + c) + a))/(a^2*b + b^3) + 2*(d*x + c)*B*a/(a^2 + b^2) - B*b*log(tan(d*x + c)^2 + 1)/(a^2 + b^2))/d`

3.310.9 Mupad [B] (verification not implemented)

Time = 8.08 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.62

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^2} dx = \frac{B b \ln(a + b \tan(c + dx))}{d (a^2 + b^2)} - \frac{B \ln(\tan(c + dx) + 1i)}{2 d (b + a 1i)} - \frac{B \ln(\tan(c + dx) - i) 1i}{2 d (a + b 1i)}$$

input `int((B*a + B*b*tan(c + d*x))/(a + b*tan(c + d*x))^2,x)`output `(B*b*log(a + b*tan(c + d*x)))/(d*(a^2 + b^2)) - (B*log(tan(c + d*x) + 1i))/(2*d*(a*1i + b)) - (B*log(tan(c + d*x) - 1i)*1i)/(2*d*(a + b*1i))`

3.311
$$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.311.1 Optimal result 2952
 3.311.2 Mathematica [C] (verified) 2952
 3.311.3 Rubi [A] (verified) 2953
 3.311.4 Maple [A] (verified) 2955
 3.311.5 Fricas [A] (verification not implemented) 2955
 3.311.6 Sympy [C] (verification not implemented) 2956
 3.311.7 Maxima [A] (verification not implemented) 2957
 3.311.8 Giac [A] (verification not implemented) 2957
 3.311.9 Mupad [B] (verification not implemented) 2958

3.311.1 Optimal result

Integrand size = 32, antiderivative size = 69

$$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = -\frac{bBx}{a^2+b^2} + \frac{B \log(\sin(c+dx))}{ad} - \frac{b^2B \log(a \cos(c+dx)+b \sin(c+dx))}{a(a^2+b^2)d}$$

output `-b*B*x/(a^2+b^2)+B*ln(sin(d*x+c))/a/d-b^2*B*ln(a*cos(d*x+c)+b*sin(d*x+c))/a/(a^2+b^2)/d`

3.311.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.14

$$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \frac{B(a(a+ib) \log(i-\cot(c+dx))+a(a-ib) \log(i+\cot(c+dx))+2b^2 \log(b+a \cot(c+dx)))}{2a(a^2+b^2)d}$$

input `Integrate[(Cot[c+d*x]*(a*B+b*B*Tan[c+d*x]))/(a+b*Tan[c+d*x])^2,x]`

output
$$\frac{-1/2*(B*(a*(a + I*b)*Log[I - Cot[c + d*x]] + a*(a - I*b)*Log[I + Cot[c + d*x]] + 2*b^2*Log[b + a*Cot[c + d*x]))}{(a*(a^2 + b^2)*d)}$$

3.311.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.01, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.219$, Rules used = {2011, 3042, 4054, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot(c+dx)(aB + bB \tan(c+dx))}{(a + b \tan(c+dx))^2} dx \\ & \quad \downarrow \text{2011} \\ & B \int \frac{\cot(c+dx)}{a + b \tan(c+dx)} dx \\ & \quad \downarrow \text{3042} \\ & B \int \frac{1}{\tan(c+dx)(a + b \tan(c+dx))} dx \\ & \quad \downarrow \text{4054} \\ & B \left(-\frac{b^2 \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2 + b^2)} + \frac{\int \cot(c+dx) dx}{a} - \frac{bx}{a^2 + b^2} \right) \\ & \quad \downarrow \text{3042} \\ & B \left(-\frac{b^2 \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2 + b^2)} + \frac{\int -\tan(c+dx + \frac{\pi}{2}) dx}{a} - \frac{bx}{a^2 + b^2} \right) \\ & \quad \downarrow \text{25} \\ & B \left(-\frac{b^2 \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2 + b^2)} - \frac{\int \tan(\frac{1}{2}(2c + \pi) + dx) dx}{a} - \frac{bx}{a^2 + b^2} \right) \\ & \quad \downarrow \text{3956} \\ & B \left(-\frac{b^2 \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2 + b^2)} - \frac{bx}{a^2 + b^2} + \frac{\log(-\sin(c+dx))}{ad} \right) \end{aligned}$$

$$\downarrow 4013$$

$$B\left(-\frac{b^2 \log(a \cos(c + dx) + b \sin(c + dx))}{ad(a^2 + b^2)} - \frac{bx}{a^2 + b^2} + \frac{\log(-\sin(c + dx))}{ad}\right)$$

input `Int[(Cot[c + d*x]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `B*(-((b*x)/(a^2 + b^2)) + Log[-Sin[c + d*x]]/(a*d) - (b^2*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))`

3.311.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4054 `Int[1/(((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[(a*c - b*d)*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[b^2/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[d^2/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.311.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.16

method	result	size
parallelrisch	$\frac{B(-2abdx+2\ln(\tan(dx+c))a^2+2\ln(\tan(dx+c))b^2-\ln(\sec^2(dx+c))a^2-2b^2\ln(a+b\tan(dx+c)))}{2ad(a^2+b^2)}$	80
derivativedivides	$\frac{B\left(\frac{\ln(\tan(dx+c))}{a} + \frac{-\frac{a\ln(1+\tan^2(dx+c))}{2} - b\arctan(\tan(dx+c)) - \frac{b^2\ln(a+b\tan(dx+c))}{a(a^2+b^2)}}{a^2+b^2}\right)}{d}$	81
default	$\frac{B\left(\frac{\ln(\tan(dx+c))}{a} + \frac{-\frac{a\ln(1+\tan^2(dx+c))}{2} - b\arctan(\tan(dx+c)) - \frac{b^2\ln(a+b\tan(dx+c))}{a(a^2+b^2)}}{a^2+b^2}\right)}{d}$	81
norman	$\frac{-\frac{bBax}{a^2+b^2} - \frac{b^2Bx\tan(dx+c)}{a^2+b^2}}{a+b\tan(dx+c)} + \frac{B\ln(\tan(dx+c))}{ad} - \frac{Ba\ln(1+\tan^2(dx+c))}{2d(a^2+b^2)} - \frac{Bb^2\ln(a+b\tan(dx+c))}{ad(a^2+b^2)}$	127
risch	$-\frac{ixB}{ib-a} - \frac{2ixB}{a} - \frac{2iBc}{ad} + \frac{2ib^2Bx}{a(a^2+b^2)} + \frac{2ib^2Bc}{ad(a^2+b^2)} + \frac{\ln(e^{2i(dx+c)}-1)B}{ad} - \frac{b^2\ln\left(e^{2i(dx+c)} - \frac{ib+a}{ib-a}\right)B}{ad(a^2+b^2)}$	149

input `int(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/2/a/d/(a^2+b^2)*B*(-2*a*b*d*x+2*ln(tan(d*x+c))*a^2+2*ln(tan(d*x+c))*b^2-ln(sec(d*x+c)^2)*a^2-2*b^2*ln(a+b*tan(d*x+c)))`

3.311.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.51

$$\int \frac{\cot(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= -\frac{2Babdx + Bb^2 \log\left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2 + 1}\right) - (Ba^2 + Bb^2) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2 + 1}\right)}{2(a^3 + ab^2)d}$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

output `-1/2*(2*B*a*b*d*x + B*b^2*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)) - (B*a^2 + B*b^2)*log(tan(d*x + c)^2/(tan(d*x + c)^2 + 1)))/(a^3 + a*b^2)*d`

3.311. $\int \frac{\cot(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

3.311.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.43 (sec) , antiderivative size = 672, normalized size of antiderivative = 9.74

$$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \left\{ \begin{array}{l} \frac{\tilde{\infty} Bx \cot(c)}{\tan(c)} \\ B \left(-\frac{\log(\tan^2(c+dx)+1)}{2d} + \frac{\log(\tan(c+dx))}{d} \right) \\ \frac{B \left(-x - \frac{1}{d \tan(c+dx)} \right)}{a} \\ \frac{B \left(-x - \frac{1}{d \tan(c+dx)} \right)}{b} \\ \frac{Bdx \tan(c+dx)}{2bd \tan(c+dx)-2ibd} - \frac{iBdx}{2bd \tan(c+dx)-2ibd} - \frac{iB \log(\tan^2(c+dx)+1) \tan(c+dx)}{2bd \tan(c+dx)-2ibd} - \frac{B \log(\tan^2(c+dx)+1)}{2bd \tan(c+dx)-2ibd} + \frac{2iB \log(\tan(c+dx)) \tan(c+dx)}{2bd \tan(c+dx)-2ibd} \\ \frac{Bdx \tan(c+dx)}{2bd \tan(c+dx)+2ibd} + \frac{iBdx}{2bd \tan(c+dx)+2ibd} + \frac{iB \log(\tan^2(c+dx)+1) \tan(c+dx)}{2bd \tan(c+dx)+2ibd} - \frac{B \log(\tan^2(c+dx)+1)}{2bd \tan(c+dx)+2ibd} - \frac{2iB \log(\tan(c+dx)) \tan(c+dx)}{2bd \tan(c+dx)+2ibd} \\ \frac{x(Ba+Bb \tan(c)) \cot(c)}{(a+b \tan(c))^2} \\ -\frac{Ba^2 \log(\tan^2(c+dx)+1)}{2a^3d+2ab^2d} + \frac{2Ba^2 \log(\tan(c+dx))}{2a^3d+2ab^2d} - \frac{2Babd}{2a^3d+2ab^2d} - \frac{2Bb^2 \log(\frac{a}{b} + \tan(c+dx))}{2a^3d+2ab^2d} + \frac{2Bb^2 \log(\tan(c+dx))}{2a^3d+2ab^2d} \end{array} \right.$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Piecewise((zoo*B*x*cot(c)/tan(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (B*(-log(tan(c + d*x)**2 + 1)/(2*d) + log(tan(c + d*x))/d)/a, Eq(b, 0)), (B*(-x - 1/(d*tan(c + d*x)))/b, Eq(a, 0)), (B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*B*d*x/(2*b*d*tan(c + d*x) - 2*I*b*d) - I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) - B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 2*I*B*log(tan(c + d*x))*tan(c + d*x)/(2*b*d*tan(c + d*x) - 2*I*b*d) + 2*B*log(tan(c + d*x))/(2*b*d*tan(c + d*x) - 2*I*b*d) + B/(2*b*d*tan(c + d*x) - 2*I*b*d), Eq(a, -I*b)), (B*d*x*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*B*d*x/(2*b*d*tan(c + d*x) + 2*I*b*d) + I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) - B*log(tan(c + d*x)**2 + 1)/(2*b*d*tan(c + d*x) + 2*I*b*d) - 2*I*B*log(tan(c + d*x))*tan(c + d*x)/(2*b*d*tan(c + d*x) + 2*I*b*d) + 2*B*log(tan(c + d*x))/(2*b*d*tan(c + d*x) + 2*I*b*d) + B/(2*b*d*tan(c + d*x) + 2*I*b*d), Eq(a, I*b)), (x*(B*a + B*b*tan(c))*cot(c)/(a + b*tan(c))**2, Eq(d, 0)), (-B*a**2*log(tan(c + d*x)**2 + 1)/(2*a**3*d + 2*a*b**2*d) + 2*B*a**2*log(tan(c + d*x))/(2*a**3*d + 2*a*b**2*d) - 2*B*a*b*d*x/(2*a**3*d + 2*a*b**2*d) - 2*B*b**2*log(a/b + tan(c + d*x))/(2*a**3*d + 2*a*b**2*d) + 2*B*b**2*log(tan(c + d*x))/(2*a**3*d + 2*a*b**2*d), True))`

3.311. $\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.311.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.28

$$\int \frac{\cot(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= -\frac{\frac{2Bb^2\log(b\tan(dx+c)+a)}{a^3+ab^2} + \frac{2(dx+c)Bb}{a^2+b^2} + \frac{Ba\log(\tan(dx+c)^2+1)}{a^2+b^2} - \frac{2B\log(\tan(dx+c))}{a}}{2d}$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `-1/2*(2*B*b^2*log(b*tan(d*x + c) + a)/(a^3 + a*b^2) + 2*(d*x + c)*B*b/(a^2 + b^2) + B*a*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) - 2*B*log(tan(d*x + c))/a)/d`

3.311.8 Giac [A] (verification not implemented)

Time = 0.46 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.33

$$\int \frac{\cot(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= -\frac{\frac{2Bb^3\log(|b\tan(dx+c)+a|)}{a^3b+ab^3} + \frac{2(dx+c)Bb}{a^2+b^2} + \frac{Ba\log(\tan(dx+c)^2+1)}{a^2+b^2} - \frac{2B\log(|\tan(dx+c)|)}{a}}{2d}$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output `-1/2*(2*B*b^3*log(abs(b*tan(d*x + c) + a))/(a^3*b + a*b^3) + 2*(d*x + c)*B*b/(a^2 + b^2) + B*a*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) - 2*B*log(abs(tan(d*x + c))))/a)/d`

3.311.9 Mupad [B] (verification not implemented)

Time = 7.40 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.43

$$\int \frac{\cot(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \frac{B \ln(\tan(c+dx))}{ad} - \frac{B \ln(\tan(c+dx)+1i)}{2d(a-b1i)} - \frac{Bb^2 \ln(a+b\tan(c+dx))}{ad(a^2+b^2)} - \frac{B \ln(\tan(c+dx)-i)1i}{2d(-b+a1i)}$$

input `int((cot(c + d*x)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`output `(B*log(tan(c + d*x)))/(a*d) - (B*log(tan(c + d*x) + 1i))/(2*d*(a - b*1i)) - (B*log(tan(c + d*x) - 1i)*1i)/(2*d*(a*1i - b)) - (B*b^2*log(a + b*tan(c + d*x)))/(a*d*(a^2 + b^2))`

3.312 $\int \frac{\cot^2(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.312.1 Optimal result 2959
 3.312.2 Mathematica [C] (verified) 2959
 3.312.3 Rubi [A] (verified) 2960
 3.312.4 Maple [A] (verified) 2962
 3.312.5 Fricas [A] (verification not implemented) 2963
 3.312.6 Sympy [C] (verification not implemented) 2964
 3.312.7 Maxima [A] (verification not implemented) 2964
 3.312.8 Giac [A] (verification not implemented) 2965
 3.312.9 Mupad [B] (verification not implemented) 2965

3.312.1 Optimal result

Integrand size = 34, antiderivative size = 85

$$\int \frac{\cot^2(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = -\frac{aBx}{a^2+b^2} - \frac{B \cot(c+dx)}{ad} - \frac{bB \log(\sin(c+dx))}{a^2d} + \frac{b^3B \log(a \cos(c+dx)+b \sin(c+dx))}{a^2(a^2+b^2)d}$$

output `-a*B*x/(a^2+b^2)-B*cot(d*x+c)/a/d-b*B*ln(sin(d*x+c))/a^2/d+b^3*B*ln(a*cos(d*x+c)+b*sin(d*x+c))/a^2/(a^2+b^2)/d`

3.312.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.40 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.14

$$\int \frac{\cot^2(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = -\frac{B \left(\frac{\cot(c+dx)}{a} - \frac{\log(i-\cot(c+dx))}{2(ia+b)} + \frac{\log(i+\cot(c+dx))}{2(ia-b)} - \frac{b^3 \log(b+a \cot(c+dx))}{a^2(a^2+b^2)} \right)}{d}$$

input `Integrate[(Cot[c + d*x]^2*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2, x]`

output $-\left(\frac{B(\cot[c + dx]/a - \text{Log}[I - \cot[c + dx]]/(2(Ia + b)) + \text{Log}[I + \cot[c + dx]]/(2(Ia - b)) - (b^3 \text{Log}[b + a \cot[c + dx]])/(a^2(a^2 + b^2))}{d}\right)$

3.312.3 Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.08, number of steps used = 9, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {2011, 3042, 4052, 3042, 4134, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^2(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^2} dx$$

↓ 2011

$$B \int \frac{\cot^2(c + dx)}{a + b \tan(c + dx)} dx$$

↓ 3042

$$B \int \frac{1}{\tan(c + dx)^2(a + b \tan(c + dx))} dx$$

↓ 4052

$$B \left(-\frac{\int \frac{\cot(c+dx)(b \tan^2(c+dx) + a \tan(c+dx) + b)}{a + b \tan(c+dx)} dx}{a} - \frac{\cot(c + dx)}{ad} \right)$$

↓ 3042

$$B \left(-\frac{\int \frac{b \tan(c+dx)^2 + a \tan(c+dx) + b}{\tan(c+dx)(a + b \tan(c+dx))} dx}{a} - \frac{\cot(c + dx)}{ad} \right)$$

↓ 4134

$$B \left(-\frac{\frac{b^3 \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{b \int \cot(c+dx) dx}{a} + \frac{a^2 x}{a^2+b^2} - \frac{\cot(c + dx)}{ad} \right)$$

↓ 3042

$$\begin{aligned}
& B \left(\frac{-\frac{b^3 \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{b \int -\tan(c+dx+\frac{\pi}{2}) dx}{a} + \frac{a^2 x}{a^2+b^2} - \frac{\cot(c+dx)}{ad}}{a} \right) \\
& \quad \downarrow \text{25} \\
& B \left(\frac{-\frac{b^3 \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{b \int \tan(\frac{1}{2}(2c+\pi)+dx) dx}{a} + \frac{a^2 x}{a^2+b^2} - \frac{\cot(c+dx)}{ad}}{a} \right) \\
& \quad \downarrow \text{3956} \\
& B \left(\frac{-\frac{b^3 \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{a^2 x}{a^2+b^2} + \frac{b \log(-\sin(c+dx))}{ad} - \frac{\cot(c+dx)}{ad}}{a} \right) \\
& \quad \downarrow \text{4013} \\
& B \left(\frac{-\frac{a^2 x}{a^2+b^2} - \frac{b^3 \log(a \cos(c+dx)+b \sin(c+dx))}{ad(a^2+b^2)} + \frac{b \log(-\sin(c+dx))}{ad} - \frac{\cot(c+dx)}{ad}}{a} \right)
\end{aligned}$$

input `Int[(Cot[c + d*x]^2*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `B*(-(Cot[c + d*x]/(a*d)) - ((a^2*x)/(a^2 + b^2) + (b*Log[-Sin[c + d*x]])/(a*d) - (b^3*Log[a*Cos[c + d*x] + b*Sin[c + d*x]]/(a*(a^2 + b^2)*d))/a)`

3.312.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_.)])/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4052 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(a^2 + b^2)*(b*c - a*d))), x] + Simp[1/((m + 1)*(a^2 + b^2)*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2) - b*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b^2*d*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && LtQ[m, -1] && (LtQ[n, 0] || IntegerQ[m]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4134 `Int[((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_.)] + (C_.)*tan[(e_.) + (f_.)*(x_.)]^2)/(((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)]*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)])), x_Symbol] := Simp[(a*(A*c - c*C + B*d) + b*(B*c - A*d + C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 - a*b*B + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C - B*c*d + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.312.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.12

method	result
derivativedivides	$\frac{B \left(-\frac{1}{a \tan(dx+c)} - \frac{b \ln(\tan(dx+c))}{a^2} + \frac{b \ln(1+\tan^2(dx+c))}{2} - \frac{a \arctan(\tan(dx+c))}{a^2+b^2} + \frac{b^3 \ln(a+b \tan(dx+c))}{a^2(a^2+b^2)} \right)}{d}$
default	$\frac{B \left(-\frac{1}{a \tan(dx+c)} - \frac{b \ln(\tan(dx+c))}{a^2} + \frac{b \ln(1+\tan^2(dx+c))}{2} - \frac{a \arctan(\tan(dx+c))}{a^2+b^2} + \frac{b^3 \ln(a+b \tan(dx+c))}{a^2(a^2+b^2)} \right)}{d}$
parallelrisc	$-\frac{\left(x a^3 d + \ln(\tan(dx+c)) a^2 b + \ln(\tan(dx+c)) b^3 - \frac{b \ln(\sec^2(dx+c)) a^2}{2} - b^3 \ln(a+b \tan(dx+c)) + a^3 \cot(dx+c) + a b^2 \cot(dx+c) \right)}{a^2 d (a^2 + b^2)}$
norman	$\frac{\frac{B b^2 (\tan^2(dx+c))}{d a^2} - \frac{B}{d} - \frac{B a^2 x \tan(dx+c)}{a^2+b^2} - \frac{b B a x (\tan^2(dx+c))}{a^2+b^2}}{\tan(dx+c)(a+b \tan(dx+c))} + \frac{B b^3 \ln(a+b \tan(dx+c))}{(a^2+b^2)a^2 d} - \frac{B b \ln(\tan(dx+c))}{a^2 d} + \frac{B b \ln(e^{2i(dx+c)}-1)}{a^2 d}$
risc	$\frac{x B}{i b - a} + \frac{2 i B b x}{a^2} + \frac{2 i B b c}{a^2 d} - \frac{2 i b^3 B x}{a^2 (a^2 + b^2)} - \frac{2 i b^3 B c}{a^2 d (a^2 + b^2)} - \frac{2 i B}{d a (e^{2 i (d x + c)} - 1)} - \frac{\ln(e^{2 i (d x + c)} - 1) B b}{a^2 d} + \frac{b^3 \ln(e^{2 i (d x + c)} - 1)}{a^2 d}$

input `int(cot(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNV ERBOSE)`

output `1/d*B*(-1/a/tan(d*x+c)-b/a^2*ln(tan(d*x+c))+1/(a^2+b^2)*(1/2*b*ln(1+tan(d*x+c)^2)-a*arctan(tan(d*x+c)))+b^3/a^2/(a^2+b^2)*ln(a+b*tan(d*x+c)))`

3.312.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.73

$$\int \frac{\cot^2(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \frac{2 B a^3 dx \tan(dx+c) - B b^3 \log\left(\frac{b^2 \tan(dx+c)^2 + 2 a b \tan(dx+c) + a^2}{\tan(dx+c)^2 + 1}\right) \tan(dx+c) + 2 B a^3 + 2 B a b^2 + (B a^2 b + 2 B a^2 b^2)}{2 (a^4 + a^2 b^2) d \tan(dx+c)}$$

input `integrate(cot(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

output `-1/2*(2*B*a^3*d*x*tan(d*x+c)-B*b^3*log((b^2*tan(d*x+c)^2+2*a*b*tan(d*x+c)+a^2)/(tan(d*x+c)^2+1))*tan(d*x+c)+2*B*a^3+2*B*a*b^2+(B*a^2*b+B*b^3)*log(tan(d*x+c)^2/(tan(d*x+c)^2+1))*tan(d*x+c)/((a^4+a^2*b^2)*d*tan(d*x+c))`

3.312.
$$\int \frac{\cot^2(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.312.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.65 (sec) , antiderivative size = 1137, normalized size of antiderivative = 13.38

$$\int \frac{\cot^2(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)**2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Piecewise((zoo*B*x, Eq(a, 0) & Eq(b, 0) & Eq(c, 0) & Eq(d, 0)), (B*(-x - c
ot(c + d*x)/d)/a, Eq(b, 0)), (B*(log(tan(c + d*x)**2 + 1)/(2*d) - log(tan(
c + d*x))/d - 1/(2*d*tan(c + d*x)**2))/b, Eq(a, 0)), (-3*B*d*x*tan(c + d*x
)**2/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - 3*I*B*d*x*tan(c + d*
x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - I*B*log(tan(c + d*x)**
2 + 1)*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan(c + d*x)) + B*
log(tan(c + d*x)**2 + 1)*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 + 2*I*a*d*tan
(c + d*x)) + 2*I*B*log(tan(c + d*x))*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**
2 + 2*I*a*d*tan(c + d*x)) - 2*B*log(tan(c + d*x))*tan(c + d*x)/(2*a*d*tan(
c + d*x)**2 + 2*I*a*d*tan(c + d*x)) - 3*B*tan(c + d*x)/(2*a*d*tan(c + d*x)
2 + 2*I*a*d*tan(c + d*x)) - 2*I*B/(2*a*d*tan(c + d*x)2 + 2*I*a*d*tan(c
+ d*x)), Eq(b, -I*a)), (-3*B*d*x*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**2 -
2*I*a*d*tan(c + d*x)) + 3*I*B*d*x*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 - 2
*I*a*d*tan(c + d*x)) + I*B*log(tan(c + d*x)**2 + 1)*tan(c + d*x)**2/(2*a*d
*tan(c + d*x)**2 - 2*I*a*d*tan(c + d*x)) + B*log(tan(c + d*x)**2 + 1)*tan(
c + d*x)/(2*a*d*tan(c + d*x)**2 - 2*I*a*d*tan(c + d*x)) - 2*I*B*log(tan(c
+ d*x))*tan(c + d*x)**2/(2*a*d*tan(c + d*x)**2 - 2*I*a*d*tan(c + d*x)) - 2
*B*log(tan(c + d*x))*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 - 2*I*a*d*tan(c +
d*x)) - 3*B*tan(c + d*x)/(2*a*d*tan(c + d*x)**2 - 2*I*a*d*tan(c + d*x)) +
2*I*B/(2*a*d*tan(c + d*x)**2 - 2*I*a*d*tan(c + d*x)), Eq(b, I*a)), (zoo...`

3.312.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.24

$$\int \frac{\cot^2(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{\frac{2Bb^3 \log(b \tan(dx+c)+a)}{a^4+a^2b^2} - \frac{2(dx+c)Ba}{a^2+b^2} + \frac{Bb \log(\tan(dx+c)^2+1)}{a^2+b^2} - \frac{2Bb \log(\tan(dx+c))}{a^2} - \frac{2B}{a \tan(dx+c)}}{2d}$$

3.312. $\int \frac{\cot^2(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

input `integrate(cot(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm m="maxima")`

output $\frac{1}{2}*(2*B*b^3*\log(b*\tan(d*x + c) + a)/(a^4 + a^2*b^2) - 2*(d*x + c)*B*a/(a^2 + b^2) + B*b*\log(\tan(d*x + c)^2 + 1)/(a^2 + b^2) - 2*B*b*\log(\tan(d*x + c)))/a^2 - 2*B/(a*\tan(d*x + c)))/d$

3.312.8 Giac [A] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.44

$$\int \frac{\cot^2(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{\frac{2Bb^4 \log(|b \tan(dx+c)+a|)}{a^4b+a^2b^3} - \frac{2(dx+c)Ba}{a^2+b^2} + \frac{Bb \log(\tan(dx+c)^2+1)}{a^2+b^2} - \frac{2Bb \log(|\tan(dx+c)|)}{a^2} + \frac{2(Bb \tan(dx+c)-Ba)}{a^2 \tan(dx+c)}}{2d}$$

input `integrate(cot(d*x+c)^2*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm m="giac")`

output $\frac{1}{2}*(2*B*b^4*\log(\text{abs}(b*\tan(d*x + c) + a))/(a^4*b + a^2*b^3) - 2*(d*x + c)*B*a/(a^2 + b^2) + B*b*\log(\tan(d*x + c)^2 + 1)/(a^2 + b^2) - 2*B*b*\log(\text{abs}(\tan(d*x + c)))/a^2 + 2*(B*b*\tan(d*x + c) - B*a)/(a^2*\tan(d*x + c)))/d$

3.312.9 Mupad [B] (verification not implemented)

Time = 7.56 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.33

$$\int \frac{\cot^2(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^2} dx = \frac{B \ln(\tan(c + dx) + 1i)}{2d(b + a 1i)} - \frac{B \cot(c + dx)}{ad} - \frac{Bb \ln(\tan(c + dx))}{a^2 d} + \frac{Bb^3 \ln(a + b \tan(c + dx))}{a^2 d(a^2 + b^2)} + \frac{B \ln(\tan(c + dx) - i) 1i}{2d(a + b 1i)}$$

input `int((cot(c + d*x)^2*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`

output `(B*log(tan(c + d*x) - 1i)*1i)/(2*d*(a + b*1i)) + (B*log(tan(c + d*x) + 1i))/(2*d*(a*1i + b)) - (B*cot(c + d*x))/(a*d) - (B*b*log(tan(c + d*x)))/(a^2 *d) + (B*b^3*log(a + b*tan(c + d*x)))/(a^2*d*(a^2 + b^2))`

3.313 $\int \frac{\cot^3(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.313.1 Optimal result 2967
 3.313.2 Mathematica [C] (verified) 2967
 3.313.3 Rubi [A] (verified) 2968
 3.313.4 Maple [A] (verified) 2972
 3.313.5 Fricas [A] (verification not implemented) 2972
 3.313.6 Sympy [C] (verification not implemented) 2973
 3.313.7 Maxima [A] (verification not implemented) 2974
 3.313.8 Giac [A] (verification not implemented) 2974
 3.313.9 Mupad [B] (verification not implemented) 2975

3.313.1 Optimal result

Integrand size = 34, antiderivative size = 112

$$\int \frac{\cot^3(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \frac{bBx}{a^2+b^2} + \frac{bB \cot(c+dx)}{a^2d} - \frac{B \cot^2(c+dx)}{2ad} - \frac{(a^2-b^2)B \log(\sin(c+dx))}{a^3d} - \frac{b^4B \log(a \cos(c+dx)+b \sin(c+dx))}{a^3(a^2+b^2)d}$$

```
output b*B*x/(a^2+b^2)+b*B*cot(d*x+c)/a^2/d-1/2*B*cot(d*x+c)^2/a/d-(a^2-b^2)*B*ln
(sin(d*x+c))/a^3/d-b^4*B*ln(a*cos(d*x+c)+b*sin(d*x+c))/a^3/(a^2+b^2)/d
```

3.313.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.65 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.96

$$\int \frac{\cot^3(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = -\frac{B \left(-\frac{2b \cot(c+dx)}{a^2} + \frac{\cot^2(c+dx)}{a} - \frac{\log(i-\cot(c+dx))}{a-ib} - \frac{\log(i+\cot(c+dx))}{a+ib} + \frac{2b^4 \log(b+a \cot(c+dx))}{a^3(a^2+b^2)} \right)}{2d}$$

input `Integrate[(Cot[c + d*x]^3*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2, x]`

output `-1/2*(B*((-2*b*Cot[c + d*x])/a^2 + Cot[c + d*x]^2/a - Log[I - Cot[c + d*x]]/(a - I*b) - Log[I + Cot[c + d*x]]/(a + I*b) + (2*b^4*Log[b + a*Cot[c + d*x]])/(a^3*(a^2 + b^2))))/d`

3.313.3 Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.12, number of steps used = 13, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.382$, Rules used = {2011, 3042, 4052, 27, 3042, 4132, 25, 3042, 4135, 3042, 25, 3956, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^3(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{\cot^3(c+dx)}{a+b\tan(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{\tan(c+dx)^3(a+b\tan(c+dx))} dx \\
 & \quad \downarrow \text{4052} \\
 & B \left(-\frac{\int \frac{2\cot^2(c+dx)(b\tan^2(c+dx)+a\tan(c+dx)+b)}{a+b\tan(c+dx)} dx}{2a} - \frac{\cot^2(c+dx)}{2ad} \right) \\
 & \quad \downarrow \text{27} \\
 & B \left(-\frac{\int \frac{\cot^2(c+dx)(b\tan^2(c+dx)+a\tan(c+dx)+b)}{a+b\tan(c+dx)} dx}{a} - \frac{\cot^2(c+dx)}{2ad} \right) \\
 & \quad \downarrow \text{3042} \\
 & B \left(-\frac{\int \frac{b\tan(c+dx)^2+a\tan(c+dx)+b}{\tan(c+dx)^2(a+b\tan(c+dx))} dx}{a} - \frac{\cot^2(c+dx)}{2ad} \right)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 4132 \\
 B & \left(-\frac{\int -\frac{\cot(c+dx)(a^2-b^2-b^2 \tan^2(c+dx))}{a+b \tan(c+dx)} dx}{a} - \frac{b \cot(c+dx)}{ad} - \frac{\cot^2(c+dx)}{2ad} \right) \\
 & \downarrow 25 \\
 B & \left(-\frac{\int \frac{\cot(c+dx)(a^2-b^2-b^2 \tan^2(c+dx))}{a+b \tan(c+dx)} dx}{a} - \frac{b \cot(c+dx)}{ad} - \frac{\cot^2(c+dx)}{2ad} \right) \\
 & \downarrow 3042 \\
 B & \left(-\frac{\int \frac{a^2-b^2-b^2 \tan(c+dx)^2}{\tan(c+dx)(a+b \tan(c+dx))} dx}{a} - \frac{b \cot(c+dx)}{ad} - \frac{\cot^2(c+dx)}{2ad} \right) \\
 & \downarrow 4135 \\
 B & \left(-\frac{\frac{(a^2-b^2) \int \cot(c+dx) dx}{a} + \frac{b^4 \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^2 bx}{a^2+b^2}}{a} - \frac{b \cot(c+dx)}{ad} - \frac{\cot^2(c+dx)}{2ad} \right) \\
 & \downarrow 3042 \\
 B & \left(-\frac{\frac{(a^2-b^2) \int -\tan(c+dx+\frac{\pi}{2}) dx}{a} + \frac{b^4 \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^2 bx}{a^2+b^2}}{a} - \frac{b \cot(c+dx)}{ad} - \frac{\cot^2(c+dx)}{2ad} \right) \\
 & \downarrow 25 \\
 B & \left(-\frac{\frac{(a^2-b^2) \int \tan(\frac{1}{2}(2c+\pi)+dx) dx}{a} + \frac{b^4 \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} - \frac{a^2 bx}{a^2+b^2}}{a} - \frac{b \cot(c+dx)}{ad} - \frac{\cot^2(c+dx)}{2ad} \right) \\
 & \downarrow 3956 \\
 B & \left(-\frac{\frac{b^4 \int \frac{b-a \tan(c+dx)}{a+b \tan(c+dx)} dx}{a(a^2+b^2)} + \frac{(a^2-b^2) \log(-\sin(c+dx))}{ad} - \frac{a^2 bx}{a^2+b^2}}{a} - \frac{b \cot(c+dx)}{ad} - \frac{\cot^2(c+dx)}{2ad} \right)
 \end{aligned}$$

3.313. $\int \frac{\cot^3(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$B \left(\frac{\frac{(a^2-b^2) \log(-\sin(c+dx))}{ad} - \frac{a^2bx}{a^2+b^2} + \frac{b^4 \log(a \cos(c+dx)+b \sin(c+dx))}{ad(a^2+b^2)}}{a} - \frac{b \cot(c+dx)}{ad} - \frac{\cot^2(c+dx)}{2ad} \right)$$

input `Int[(Cot[c + d*x]^3*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `B*(-1/2*Cot[c + d*x]^2/(a*d) - ((b*Cot[c + d*x])/(a*d)) + (-((a^2*b*x)/(a^2 + b^2)) + ((a^2 - b^2)*Log[-Sin[c + d*x]])/(a*d) + (b^4*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/(a*(a^2 + b^2)*d))/a/a`

3.313.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4052 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(a^2 + b^2)*(b*c - a*d))), x] + Simp[1/((m + 1)*(a^2 + b^2)*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2) - b*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b^2*d*(m + n + 2)*Tan[e + f*x]^2, x], x], x] / ; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && LtQ[m, -1] && (LtQ[n, 0] || IntegerQ[m]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] / ; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4135 `Int[((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2)/(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*(A*c - c*C) - b*(A*d - C*d))*(x/((a^2 + b^2)*(c^2 + d^2))), x] + (Simp[(A*b^2 + a^2*C)/((b*c - a*d)*(a^2 + b^2)) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] - Simp[(c^2*C + A*d^2)/((b*c - a*d)*(c^2 + d^2)) Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x]), x], x]) / ; FreeQ[{a, b, c, d, e, f, A, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.313.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.03

method	result
derivativedivides	$B \left(-\frac{1}{2a \tan(dx+c)^2} + \frac{(-a^2+b^2) \ln(\tan(dx+c))}{a^3} + \frac{b}{a^2 \tan(dx+c)} + \frac{\frac{a \ln(1+\tan^2(dx+c))}{2} + b \arctan(\tan(dx+c))}{a^2+b^2} - \frac{b^4 \ln(a+b \tan(dx+c))}{a^3(a^2+b^2)} \right)$
default	$B \left(-\frac{1}{2a \tan(dx+c)^2} + \frac{(-a^2+b^2) \ln(\tan(dx+c))}{a^3} + \frac{b}{a^2 \tan(dx+c)} + \frac{\frac{a \ln(1+\tan^2(dx+c))}{2} + b \arctan(\tan(dx+c))}{a^2+b^2} - \frac{b^4 \ln(a+b \tan(dx+c))}{a^3(a^2+b^2)} \right)$
parallelrisch	$-\frac{B(-2x a^3 b d + 2 \ln(\tan(dx+c)) a^4 - 2 \ln(\tan(dx+c)) b^4 - \ln(\sec^2(dx+c)) a^4 + 2 \ln(a+b \tan(dx+c)) b^4 - 2 a^3 b \cot(dx+c) - 2 a^2 b^2 \tan(dx+c))}{2(a^2+b^2)a^3 d}$
norman	$\frac{\frac{b^2 B x (\tan^3(dx+c))}{a^2+b^2} + \frac{B b^2 (\tan^2(dx+c))}{d a^2} + \frac{b B a x (\tan^2(dx+c))}{a^2+b^2} - \frac{B}{2d} + \frac{B b \tan(dx+c)}{2 a d}}{\tan(dx+c)^2 (a+b \tan(dx+c))} + \frac{B a \ln(1+\tan^2(dx+c))}{2d(a^2+b^2)} - \frac{(a^2-b^2) B}{2d(a^2+b^2)}$
risch	$\frac{ixB}{ib-a} + \frac{2ixB}{a} + \frac{2iBc}{ad} - \frac{2iBb^2x}{a^3} - \frac{2iBb^2c}{a^3d} + \frac{2ib^4Bx}{(a^2+b^2)a^3} + \frac{2ib^4Bc}{(a^2+b^2)a^3d} + \frac{2iB(-ia e^{2i(dx+c)} + b e^{2i(dx+c)} - 1)}{d a^2 (e^{2i(dx+c)} - 1)^2}$

input `int(cot(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNV ERBOSE)`

output `1/d*B*(-1/2/a/tan(d*x+c)^2+1/a^3*(-a^2+b^2)*ln(tan(d*x+c))+b/a^2/tan(d*x+c)+1/(a^2+b^2)*(1/2*a*ln(1+tan(d*x+c)^2)+b*arctan(tan(d*x+c)))-b^4/a^3/(a^2 +b^2)*ln(a+b*tan(d*x+c)))`

3.313.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.71

$$\int \frac{\cot^3(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx =$$

$$\frac{Bb^4 \log\left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2 + 1}\right) \tan(dx+c)^2 + Ba^4 + Ba^2b^2 + (Ba^4 - Bb^4) \log\left(\frac{\tan(dx+c)^2}{\tan(dx+c)^2 + 1}\right) \tan(dx+c)}{2(a^5 + a^3b^2)d \tan(dx+c)}$$

input `integrate(cot(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorith m="fricas")`

3.313.
$$\int \frac{\cot^3(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

output
$$-1/2*(B*b^4*\log((b^2*\tan(dx + c)^2 + 2*a*b*\tan(dx + c) + a^2)/(\tan(dx + c)^2 + 1))*\tan(dx + c)^2 + B*a^4 + B*a^2*b^2 + (B*a^4 - B*b^4)*\log(\tan(dx + c)^2/(\tan(dx + c)^2 + 1))*\tan(dx + c)^2 - (2*B*a^3*b*dx - B*a^4 - B*a^2*b^2)*\tan(dx + c)^2 - 2*(B*a^3*b + B*a*b^3)*\tan(dx + c))/((a^5 + a^3*b^2)*d*\tan(dx + c)^2)$$

3.313.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 3.83 (sec) , antiderivative size = 1401, normalized size of antiderivative = 12.51

$$\int \frac{\cot^3(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate(cot(dx+c)**3*(B*a+b*B*tan(dx+c))/(a+b*tan(dx+c))**2,x)`

output `Piecewise((zoo*B*x, Eq(a, 0) & Eq(b, 0) & Eq(c, 0) & Eq(d, 0)), (B*(log(tan(c + dx)**2 + 1)/(2*d) - log(tan(c + dx))/d - 1/(2*d*tan(c + dx)**2)))/a, Eq(b, 0)), (B*(x + 1/(d*tan(c + dx)) - 1/(3*d*tan(c + dx)**3))/b, Eq(a, 0)), (-3*I*B*d*x*tan(c + dx)**3/(2*a*d*tan(c + dx)**3 + 2*I*a*d*tan(c + dx)**2) + 3*B*d*x*tan(c + dx)**2/(2*a*d*tan(c + dx)**3 + 2*I*a*d*tan(c + dx)**2) + 2*B*log(tan(c + dx)**2 + 1)*tan(c + dx)**3/(2*a*d*tan(c + dx)**3 + 2*I*a*d*tan(c + dx)**2) + 2*I*B*log(tan(c + dx)**2 + 1)*tan(c + dx)**2/(2*a*d*tan(c + dx)**3 + 2*I*a*d*tan(c + dx)**2) - 4*B*log(tan(c + dx))*tan(c + dx)**3/(2*a*d*tan(c + dx)**3 + 2*I*a*d*tan(c + dx)**2) - 4*I*B*log(tan(c + dx))*tan(c + dx)**2/(2*a*d*tan(c + dx)**3 + 2*I*a*d*tan(c + dx)**2) - 3*I*B*tan(c + dx)**2/(2*a*d*tan(c + dx)**3 + 2*I*a*d*tan(c + dx)**2) + B*tan(c + dx)/(2*a*d*tan(c + dx)**3 + 2*I*a*d*tan(c + dx)**2) - I*B/(2*a*d*tan(c + dx)**3 + 2*I*a*d*tan(c + dx)**2), Eq(b, -I*a)), (3*I*B*d*x*tan(c + dx)**3/(2*a*d*tan(c + dx)**3 - 2*I*a*d*tan(c + dx)**2) + 3*B*d*x*tan(c + dx)**2/(2*a*d*tan(c + dx)**3 - 2*I*a*d*tan(c + dx)**2) + 2*B*log(tan(c + dx)**2 + 1)*tan(c + dx)**3/(2*a*d*tan(c + dx)**3 - 2*I*a*d*tan(c + dx)**2) - 2*I*B*log(tan(c + dx)**2 + 1)*tan(c + dx)**2/(2*a*d*tan(c + dx)**3 - 2*I*a*d*tan(c + dx)**2) - 4*B*log(tan(c + dx))*tan(c + dx)**3/(2*a*d*tan(c + dx)**3 - 2*I*a*d*tan(c + dx)**2) + 4*I*B*log(tan(c + dx))*tan(c + dx)**2/(2*a*d*tan(c + dx)**3...`

3.313.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.16

$$\int \frac{\cot^3(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx =$$

$$\frac{\frac{2Bb^4 \log(b\tan(dx+c)+a)}{a^5+a^3b^2} - \frac{2(dx+c)Bb}{a^2+b^2} - \frac{Ba \log(\tan(dx+c)^2+1)}{a^2+b^2} + \frac{2(Ba^2-Bb^2) \log(\tan(dx+c))}{a^3} - \frac{2Bb \tan(dx+c)-Ba}{a^2 \tan(dx+c)^2}}{2d}$$

input `integrate(cot(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm m="maxima")`

output `-1/2*(2*B*b^4*log(b*tan(d*x + c) + a)/(a^5 + a^3*b^2) - 2*(d*x + c)*B*b/(a^2 + b^2) - B*a*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) + 2*(B*a^2 - B*b^2)*log(tan(d*x + c))/a^3 - (2*B*b*tan(d*x + c) - B*a)/(a^2*tan(d*x + c)^2))/d`

3.313.8 Giac [A] (verification not implemented)

Time = 0.60 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.47

$$\int \frac{\cot^3(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx =$$

$$\frac{\frac{2Bb^5 \log(|b\tan(dx+c)+a|)}{a^5b+a^3b^3} - \frac{2(dx+c)Bb}{a^2+b^2} - \frac{Ba \log(\tan(dx+c)^2+1)}{a^2+b^2} + \frac{2(Ba^2-Bb^2) \log(|\tan(dx+c)|)}{a^3} - \frac{3Ba^2 \tan(dx+c)^2 - 3Bb^2 \tan(dx+c)}{a^3 \tan(dx+c)^2}}{2d}$$

input `integrate(cot(d*x+c)^3*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm m="giac")`

output `-1/2*(2*B*b^5*log(abs(b*tan(d*x + c) + a))/(a^5*b + a^3*b^3) - 2*(d*x + c)*B*b/(a^2 + b^2) - B*a*log(tan(d*x + c)^2 + 1)/(a^2 + b^2) + 2*(B*a^2 - B*b^2)*log(abs(tan(d*x + c)))/a^3 - (3*B*a^2*tan(d*x + c)^2 - 3*B*b^2*tan(d*x + c)^2 + 2*B*a*b*tan(d*x + c) - B*a^2)/(a^3*tan(d*x + c)^2))/d`

3.313.9 Mupad [B] (verification not implemented)

Time = 7.30 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.28

$$\int \frac{\cot^3(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = -\frac{\cot(c+dx)^2 \left(\frac{B}{2a} - \frac{Bb\tan(c+dx)}{a^2} \right)}{d}$$

$$+ \frac{B \ln(\tan(c+dx) + 1i)}{2d(a-b1i)}$$

$$- \frac{B \ln(\tan(c+dx)) (a^2 - b^2)}{a^3 d}$$

$$- \frac{Bb^4 \ln(a+b\tan(c+dx))}{d(a^5 + a^3b^2)}$$

$$+ \frac{B \ln(\tan(c+dx) - i) 1i}{2d(-b+a1i)}$$

input `int((cot(c + d*x)^3*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`output `(B*log(tan(c + d*x) + 1i))/(2*d*(a - b*1i)) - (cot(c + d*x)^2*(B/(2*a) - (B*b*tan(c + d*x))/a^2))/d + (B*log(tan(c + d*x) - 1i)*1i)/(2*d*(a*1i - b)) - (B*log(tan(c + d*x))*(a^2 - b^2))/(a^3*d) - (B*b^4*log(a + b*tan(c + d*x)))/(d*(a^5 + a^3*b^2))`

3.314 $\int \frac{3+\tan(c+dx)}{2-\tan(c+dx)} dx$

3.314.1 Optimal result	2976
3.314.2 Mathematica [B] (verified)	2976
3.314.3 Rubi [A] (verified)	2977
3.314.4 Maple [A] (verified)	2978
3.314.5 Fricas [A] (verification not implemented)	2979
3.314.6 Sympy [B] (verification not implemented)	2979
3.314.7 Maxima [A] (verification not implemented)	2979
3.314.8 Giac [A] (verification not implemented)	2980
3.314.9 Mupad [B] (verification not implemented)	2980

3.314.1 Optimal result

Integrand size = 21, antiderivative size = 25

$$\int \frac{3 + \tan(c + dx)}{2 - \tan(c + dx)} dx = x - \frac{\log(2 \cos(c + dx) - \sin(c + dx))}{d}$$

output `x-ln(2*cos(d*x+c)-sin(d*x+c))/d`

3.314.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 62 vs. 2(25) = 50.

Time = 0.06 (sec) , antiderivative size = 62, normalized size of antiderivative = 2.48

$$\int \frac{3 + \tan(c + dx)}{2 - \tan(c + dx)} dx = \frac{\arctan(\tan(c + dx))}{d} + \frac{\log(5 - 4(2 - \tan(c + dx)) + (2 - \tan(c + dx))^2)}{2d} - \frac{\log(2 - \tan(c + dx))}{d}$$

input `Integrate[(3 + Tan[c + d*x])/(2 - Tan[c + d*x]),x]`

output `ArcTan[Tan[c + d*x]]/d + Log[5 - 4*(2 - Tan[c + d*x]) + (2 - Tan[c + d*x])^2]/(2*d) - Log[2 - Tan[c + d*x]]/d`

3.314.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$, Rules used = {3042, 4014, 25, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(c+dx)+3}{2-\tan(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)+3}{2-\tan(c+dx)} dx \\
 & \quad \downarrow \text{4014} \\
 & x - \int -\frac{2\tan(c+dx)+1}{2-\tan(c+dx)} dx \\
 & \quad \downarrow \text{25} \\
 & \int \frac{2\tan(c+dx)+1}{2-\tan(c+dx)} dx + x \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{2\tan(c+dx)+1}{2-\tan(c+dx)} dx + x \\
 & \quad \downarrow \text{4013} \\
 & x - \frac{\log(2\cos(c+dx) - \sin(c+dx))}{d}
 \end{aligned}$$

input `Int[(3 + Tan[c + d*x])/(2 - Tan[c + d*x]),x]`

output `x - Log[2*Cos[c + d*x] - Sin[c + d*x]]/d`

3.314.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

3.314.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.32

method	result	size
norman	$x - \frac{\ln(-2+\tan(dx+c))}{d} + \frac{\ln(1+\tan^2(dx+c))}{2d}$	33
risch	$ix + x + \frac{2ic}{d} - \frac{\ln(e^{2i(dx+c)} + \frac{3}{5} - \frac{4i}{5})}{d}$	33
parallelrisc	$\frac{2dx + \ln(1+\tan^2(dx+c)) - 2\ln(-2+\tan(dx+c))}{2d}$	33
derivativdivides	$-\frac{\ln(-2+\tan(dx+c)) + \frac{\ln(1+\tan^2(dx+c))}{2}}{d} + \arctan(\tan(dx+c))$	37
default	$-\frac{\ln(-2+\tan(dx+c)) + \frac{\ln(1+\tan^2(dx+c))}{2}}{d} + \arctan(\tan(dx+c))$	37

input `int((3+tan(d*x+c))/(2-tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `x-1/d*ln(-2+tan(d*x+c))+1/2/d*ln(1+tan(d*x+c)^2)`

3.314.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.76

$$\int \frac{3 + \tan(c + dx)}{2 - \tan(c + dx)} dx = \frac{2 dx - \log\left(\frac{\tan(dx+c)^2 - 4 \tan(dx+c) + 4}{\tan(dx+c)^2 + 1}\right)}{2 d}$$

input `integrate((3+tan(d*x+c))/(2-tan(d*x+c)),x, algorithm="fracas")`

output `1/2*(2*d*x - log((tan(d*x + c)^2 - 4*tan(d*x + c) + 4)/(tan(d*x + c)^2 + 1)))/d`

3.314.6 Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. 2(19) = 38.

Time = 0.09 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.56

$$\int \frac{3 + \tan(c + dx)}{2 - \tan(c + dx)} dx = \begin{cases} x - \frac{\log(\tan(c+dx)-2)}{d} + \frac{\log(\tan^2(c+dx)+1)}{2d} & \text{for } d \neq 0 \\ \frac{x(\tan(c)+3)}{2-\tan(c)} & \text{otherwise} \end{cases}$$

input `integrate((3+tan(d*x+c))/(2-tan(d*x+c)),x)`

output `Piecewise((x - log(tan(c + d*x) - 2)/d + log(tan(c + d*x)**2 + 1)/(2*d), Ne(d, 0)), (x*(tan(c) + 3)/(2 - tan(c)), True))`

3.314.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.40

$$\int \frac{3 + \tan(c + dx)}{2 - \tan(c + dx)} dx = \frac{2 dx + 2c + \log(\tan(dx+c)^2 + 1) - 2 \log(\tan(dx+c) - 2)}{2 d}$$

input `integrate((3+tan(d*x+c))/(2-tan(d*x+c)),x, algorithm="maxima")`

output `1/2*(2*d*x + 2*c + log(tan(d*x + c)^2 + 1) - 2*log(tan(d*x + c) - 2))/d`

3.314.8 Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.44

$$\int \frac{3 + \tan(c + dx)}{2 - \tan(c + dx)} dx = \frac{2dx + 2c + \log(\tan(dx + c)^2 + 1) - 2 \log(|\tan(dx + c) - 2|)}{2d}$$

input `integrate((3+tan(d*x+c))/(2-tan(d*x+c)),x, algorithm="giac")`output `1/2*(2*d*x + 2*c + log(tan(d*x + c)^2 + 1) - 2*log(abs(tan(d*x + c) - 2)))/d`**3.314.9 Mupad [B] (verification not implemented)**

Time = 7.68 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.96

$$\int \frac{3 + \tan(c + dx)}{2 - \tan(c + dx)} dx = -\frac{\ln(\tan(c + dx) - 2)}{d} + \frac{\ln(\tan(c + dx) - i) \left(\frac{1}{2} - \frac{1}{2}i\right)}{d} + \frac{\ln(\tan(c + dx) + 1i) \left(\frac{1}{2} + \frac{1}{2}i\right)}{d}$$

input `int(-(tan(c + d*x) + 3)/(tan(c + d*x) - 2),x)`output `(log(tan(c + d*x) - 1i)*(1/2 - 1i/2))/d - log(tan(c + d*x) - 2)/d + (log(tan(c + d*x) + 1i)*(1/2 + 1i/2))/d`

$$\mathbf{3.315} \quad \int \frac{\frac{bB}{a} + B \tan(c+dx)}{a+b \tan(c+dx)} dx$$

3.315.1 Optimal result	2981
3.315.2 Mathematica [A] (verified)	2981
3.315.3 Rubi [A] (verified)	2982
3.315.4 Maple [A] (verified)	2983
3.315.5 Fricas [A] (verification not implemented)	2984
3.315.6 Sympy [C] (verification not implemented)	2984
3.315.7 Maxima [A] (verification not implemented)	2985
3.315.8 Giac [A] (verification not implemented)	2985
3.315.9 Mupad [B] (verification not implemented)	2985

3.315.1 Optimal result

Integrand size = 28, antiderivative size = 58

$$\int \frac{\frac{bB}{a} + B \tan(c+dx)}{a+b \tan(c+dx)} dx = \frac{2bBx}{a^2 + b^2} - \frac{\left(a - \frac{b^2}{a}\right) B \log(a \cos(c+dx) + b \sin(c+dx))}{(a^2 + b^2) d}$$

output `2*b*B*x/(a^2+b^2)-(a-b^2/a)*B*ln(a*cos(d*x+c)+b*sin(d*x+c))/(a^2+b^2)/d`

3.315.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.12

$$\int \frac{\frac{bB}{a} + B \tan(c+dx)}{a+b \tan(c+dx)} dx = \frac{B(4ab \arctan(\tan(c+dx)) + (a^2 - b^2) (\log(\sec^2(c+dx)) - 2 \log(a + b \tan(c+dx))))}{2a(a^2 + b^2) d}$$

input `Integrate[((b*B)/a + B*Tan[c + d*x])/(a + b*Tan[c + d*x]),x]`

output `(B*(4*a*b*ArcTan[Tan[c + d*x]] + (a^2 - b^2)*(Log[Sec[c + d*x]^2] - 2*Log[a + b*Tan[c + d*x]])))/(2*a*(a^2 + b^2)*d)`

$$3.315. \quad \int \frac{\frac{bB}{a} + B \tan(c+dx)}{a+b \tan(c+dx)} dx$$

3.315.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\frac{bB}{a} + B \tan(c + dx)}{a + b \tan(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\frac{bB}{a} + B \tan(c + dx)}{a + b \tan(c + dx)} dx \\
 & \quad \downarrow \text{4014} \\
 & \frac{2bBx}{a^2 + b^2} - \frac{B \left(a - \frac{b^2}{a} \right) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2bBx}{a^2 + b^2} - \frac{B \left(a - \frac{b^2}{a} \right) \int \frac{b - a \tan(c + dx)}{a + b \tan(c + dx)} dx}{a^2 + b^2} \\
 & \quad \downarrow \text{4013} \\
 & \frac{2bBx}{a^2 + b^2} - \frac{B \left(a - \frac{b^2}{a} \right) \log(a \cos(c + dx) + b \sin(c + dx))}{d(a^2 + b^2)}
 \end{aligned}$$

input `Int[((b*B)/a + B*Tan[c + d*x])/(a + b*Tan[c + d*x]),x]`

output `(2*b*B*x)/(a^2 + b^2) - ((a - b^2/a)*B*Log[a*Cos[c + d*x] + b*Sin[c + d*x]])/((a^2 + b^2)*d)`

3.315.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4013 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

3.315.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.47

method	result
derivativedivides	$\frac{B \left(\frac{(a^2 - b^2) \ln(1 + \tan^2(dx + c))}{2} + 2ab \arctan(\tan(dx + c)) - \frac{(a^2 - b^2) \ln(a + b \tan(dx + c))}{a^2 + b^2} \right)}{da}$
default	$\frac{B \left(\frac{(a^2 - b^2) \ln(1 + \tan^2(dx + c))}{2} + 2ab \arctan(\tan(dx + c)) - \frac{(a^2 - b^2) \ln(a + b \tan(dx + c))}{a^2 + b^2} \right)}{da}$
norman	$\frac{2bBx}{a^2 + b^2} + \frac{B(a^2 - b^2) \ln(1 + \tan^2(dx + c))}{2ad(a^2 + b^2)} - \frac{B(a^2 - b^2) \ln(a + b \tan(dx + c))}{ad(a^2 + b^2)}$
parallelrisch	$\frac{4Babdx + B \ln(1 + \tan^2(dx + c))a^2 - B \ln(1 + \tan^2(dx + c))b^2 - 2B \ln(a + b \tan(dx + c))a^2 + 2Bb^2 \ln(a + b \tan(dx + c))}{2ad(a^2 + b^2)}$
risch	$-\frac{x B b}{a(i b - a)} + \frac{i x B}{i b - a} + \frac{2 i B x a}{a^2 + b^2} - \frac{2 i b^2 B x}{a(a^2 + b^2)} + \frac{2 i B a c}{d(a^2 + b^2)} - \frac{2 i b^2 B c}{a d(a^2 + b^2)} - \frac{\ln\left(e^{2i(dx+c)} - \frac{ib+a}{ib-a}\right) B a}{d(a^2 + b^2)} + \frac{b^2 \ln\left(e^{2i(dx+c)} - \frac{ib+a}{ib-a}\right)}{a d(a^2 + b^2)}$

input `int((b*B/a+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d/a*B*(1/(a^2+b^2)*(1/2*(a^2-b^2)*ln(1+tan(d*x+c)^2)+2*a*b*arctan(tan(d*x+c)))-(a^2-b^2)/(a^2+b^2)*ln(a+b*tan(d*x+c)))`

3.315.
$$\int \frac{\frac{bB}{a} + B \tan(c+dx)}{a + b \tan(c+dx)} dx$$

3.315.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.34

$$\int \frac{\frac{bB}{a} + B \tan(c + dx)}{a + b \tan(c + dx)} dx = \frac{4 Babdx - (Ba^2 - Bb^2) \log\left(\frac{b^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + a^2}{\tan(dx+c)^2 + 1}\right)}{2(a^3 + ab^2)d}$$

input `integrate((b*B/a+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fracas")`output `1/2*(4*B*a*b*d*x - (B*a^2 - B*b^2)*log((b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2)/(tan(d*x + c)^2 + 1)))/(a^3 + a*b^2)*d`**3.315.6 Sympy [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.45 (sec) , antiderivative size = 235, normalized size of antiderivative = 4.05

$$\int \frac{\frac{bB}{a} + B \tan(c + dx)}{a + b \tan(c + dx)} dx = \begin{cases} \text{NaN} \\ \frac{B \log(\tan^2(c+dx)+1)}{2ad} \\ -\frac{B}{bd \tan(c+dx) - ibd} \\ -\frac{B}{bd \tan(c+dx) + ibd} \\ \frac{x(B \tan(c) + \frac{Bb}{a})}{a + b \tan(c)} \\ -\frac{2Ba^2 \log(\frac{a}{b} + \tan(c+dx))}{2a^3d + 2ab^2d} + \frac{Ba^2 \log(\tan^2(c+dx)+1)}{2a^3d + 2ab^2d} + \frac{4Babdx}{2a^3d + 2ab^2d} + \frac{2Bb^2 \log(\frac{a}{b} + \tan(c+dx))}{2a^3d + 2ab^2d} - \frac{Bb^2 \log(\tan^2(c+dx)+1)}{2a^3d + 2ab^2d} \end{cases}$$

input `integrate((b*B/a+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`output `Piecewise((nan, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (B*log(tan(c + d*x)**2 + 1)/(2*a*d), Eq(b, 0)), (-B/(b*d*tan(c + d*x) - I*b*d), Eq(a, -I*b)), (-B/(b*d*tan(c + d*x) + I*b*d), Eq(a, I*b)), (x*(B*tan(c) + B*b/a)/(a + b*tan(c)), Eq(d, 0)), (-2*B*a**2*log(a/b + tan(c + d*x))/(2*a**3*d + 2*a*b**2*d) + B*a**2*log(tan(c + d*x)**2 + 1)/(2*a**3*d + 2*a*b**2*d) + 4*B*a*b*d*x/(2*a**3*d + 2*a*b**2*d) + 2*B*b**2*log(a/b + tan(c + d*x))/(2*a**3*d + 2*a*b**2*d) - B*b**2*log(tan(c + d*x)**2 + 1)/(2*a**3*d + 2*a*b**2*d), True))`

3.315. $\int \frac{\frac{bB}{a} + B \tan(c+dx)}{a + b \tan(c+dx)} dx$

3.315.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.64

$$\int \frac{\frac{bB}{a} + B \tan(c + dx)}{a + b \tan(c + dx)} dx = \frac{\frac{4(dx+c)Bb}{a^2+b^2} - \frac{2(Ba^2-Bb^2) \log(b \tan(dx+c)+a)}{a^3+ab^2} + \frac{(Ba^2-Bb^2) \log(\tan(dx+c)^2+1)}{a^3+ab^2}}{2d}$$

input `integrate((b*B/a+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`output `1/2*(4*(d*x + c)*B*b/(a^2 + b^2) - 2*(B*a^2 - B*b^2)*log(b*tan(d*x + c) + a)/(a^3 + a*b^2) + (B*a^2 - B*b^2)*log(tan(d*x + c)^2 + 1)/(a^3 + a*b^2))/d`**3.315.8 Giac [A] (verification not implemented)**

Time = 0.36 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.71

$$\int \frac{\frac{bB}{a} + B \tan(c + dx)}{a + b \tan(c + dx)} dx = \frac{\frac{4(dx+c)Bb}{a^2+b^2} + \frac{(Ba^2-Bb^2) \log(\tan(dx+c)^2+1)}{a^3+ab^2} - \frac{2(Ba^2b-Bb^3) \log(|b \tan(dx+c)+a|)}{a^3b+ab^3}}{2d}$$

input `integrate((b*B/a+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`output `1/2*(4*(d*x + c)*B*b/(a^2 + b^2) + (B*a^2 - B*b^2)*log(tan(d*x + c)^2 + 1)/(a^3 + a*b^2) - 2*(B*a^2*b - B*b^3)*log(abs(b*tan(d*x + c) + a))/(a^3*b + a*b^3))/d`**3.315.9 Mupad [B] (verification not implemented)**

Time = 8.68 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.93

$$\int \frac{\frac{bB}{a} + B \tan(c + dx)}{a + b \tan(c + dx)} dx = -\frac{\ln(\tan(c + dx) - i) (Bb + Ba \operatorname{li})}{2d (ab - a^2 \operatorname{li})} - \frac{\ln(\tan(c + dx) + i) (Ba + Bb \operatorname{li})}{2d (-a^2 + ab \operatorname{li})} - \frac{B \ln(a + b \tan(c + dx)) (a^2 - b^2)}{ad (a^2 + b^2)}$$

3.315. $\int \frac{\frac{bB}{a} + B \tan(c + dx)}{a + b \tan(c + dx)} dx$

input `int((B*tan(c + d*x) + (B*b)/a)/(a + b*tan(c + d*x)),x)`

output `- (log(tan(c + d*x) - 1i)*(B*a*1i + B*b))/(2*d*(a*b - a^2*1i)) - (log(tan(c + d*x) + 1i)*(B*a + B*b*1i))/(2*d*(a*b*1i - a^2)) - (B*log(a + b*tan(c + d*x))*(a^2 - b^2))/(a*d*(a^2 + b^2))`

3.316 $\int \frac{a+b \tan(c+dx)}{(b+a \tan(c+dx))^2} dx$

3.316.1 Optimal result	2987
3.316.2 Mathematica [C] (verified)	2987
3.316.3 Rubi [A] (verified)	2988
3.316.4 Maple [A] (verified)	2990
3.316.5 Fricas [A] (verification not implemented)	2990
3.316.6 Sympy [C] (verification not implemented)	2991
3.316.7 Maxima [A] (verification not implemented)	2992
3.316.8 Giac [A] (verification not implemented)	2992
3.316.9 Mupad [B] (verification not implemented)	2993

3.316.1 Optimal result

Integrand size = 23, antiderivative size = 101

$$\int \frac{a + b \tan(c + dx)}{(b + a \tan(c + dx))^2} dx = -\frac{a(a^2 - 3b^2)x}{(a^2 + b^2)^2} + \frac{b(3a^2 - b^2) \log(b \cos(c + dx) + a \sin(c + dx))}{(a^2 + b^2)^2 d} - \frac{a^2 - b^2}{(a^2 + b^2) d (b + a \tan(c + dx))}$$

```
output -a*(a^2-3*b^2)*x/(a^2+b^2)^2+b*(3*a^2-b^2)*ln(b*cos(d*x+c)+a*sin(d*x+c))/(a^2+b^2)^2/d+(-a^2+b^2)/(a^2+b^2)/d/(b+a*tan(d*x+c))
```

3.316.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.58 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.85

$$\int \frac{a + b \tan(c + dx)}{(b + a \tan(c + dx))^2} dx = \frac{b(-((a+ib) \log(i - \tan(c+dx))) - (a-ib) \log(i + \tan(c+dx))) + 2a \log(b + a \tan(c+dx))}{a^2 + b^2} + (a - b)(a + b) \left(\frac{i \log(i - \tan(c+dx))}{(a - ib)^2} - \frac{i \log(i + \tan(c+dx))}{(a + ib)^2} \right)$$

$2ad$

```
input Integrate[(a + b*Tan[c + d*x])/(b + a*Tan[c + d*x])^2,x]
```

```
output ((b*(-((a + I*b)*Log[I - Tan[c + d*x]]) - (a - I*b)*Log[I + Tan[c + d*x]]
+ 2*a*Log[b + a*Tan[c + d*x]]))/(a^2 + b^2) + (a - b)*(a + b)*((I*Log[I -
Tan[c + d*x]])/(a - I*b)^2 - (I*Log[I + Tan[c + d*x]])/(a + I*b)^2 + (2*a*
(2*b*Log[b + a*Tan[c + d*x]] - (a^2 + b^2)/(b + a*Tan[c + d*x])))/(a^2 + b
^2)^2))/(2*a*d)
```

3.316.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.11, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {3042, 4012, 3042, 4014, 3042, 4013}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \tan(c + dx)}{(a \tan(c + dx) + b)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{a + b \tan(c + dx)}{(a \tan(c + dx) + b)^2} dx \\
 & \quad \downarrow \text{4012} \\
 & \frac{\int \frac{2ab - (a^2 - b^2) \tan(c + dx)}{b + a \tan(c + dx)} dx}{a^2 + b^2} - \frac{a^2 - b^2}{d(a^2 + b^2)(a \tan(c + dx) + b)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{2ab - (a^2 - b^2) \tan(c + dx)}{b + a \tan(c + dx)} dx}{a^2 + b^2} - \frac{a^2 - b^2}{d(a^2 + b^2)(a \tan(c + dx) + b)} \\
 & \quad \downarrow \text{4014} \\
 & \frac{b(3a^2 - b^2) \int \frac{a - b \tan(c + dx)}{b + a \tan(c + dx)} dx}{a^2 + b^2} - \frac{ax(a^2 - 3b^2)}{a^2 + b^2} - \frac{a^2 - b^2}{d(a^2 + b^2)(a \tan(c + dx) + b)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b(3a^2 - b^2) \int \frac{a - b \tan(c + dx)}{b + a \tan(c + dx)} dx}{a^2 + b^2} - \frac{ax(a^2 - 3b^2)}{a^2 + b^2} - \frac{a^2 - b^2}{d(a^2 + b^2)(a \tan(c + dx) + b)} \\
 & \quad \downarrow \text{4013}
 \end{aligned}$$

3.316. $\int \frac{a + b \tan(c + dx)}{(b + a \tan(c + dx))^2} dx$

$$\frac{\frac{b(3a^2-b^2) \log(a \sin(c+dx)+b \cos(c+dx))}{d(a^2+b^2)} - \frac{ax(a^2-3b^2)}{a^2+b^2}}{a^2+b^2} - \frac{a^2-b^2}{d(a^2+b^2)(a \tan(c+dx)+b)}$$

input `Int[(a + b*Tan[c + d*x])/(b + a*Tan[c + d*x])^2,x]`

output `(-((a*(a^2 - 3*b^2)*x)/(a^2 + b^2)) + (b*(3*a^2 - b^2)*Log[b*Cos[c + d*x] + a*Sin[c + d*x]])/((a^2 + b^2)*d))/(a^2 + b^2) - (a^2 - b^2)/((a^2 + b^2)*d*(b + a*Tan[c + d*x]))`

3.316.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4013 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c/(b*f))*Log[RemoveContent[a*Cos[e + f*x] + b*Sin[e + f*x], x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[a*c + b*d, 0]`

rule 4014 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(a*c + b*d)*(x/(a^2 + b^2)), x] + Simp[(b*c - a*d)/(a^2 + b^2) Int[(b - a*Tan[e + f*x])/(a + b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[a*c + b*d, 0]`

3.316.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.24

method	result
derivativedivides	$-\frac{a^2-b^2}{(a^2+b^2)(b+a \tan(dx+c))} + \frac{b(3a^2-b^2) \ln(b+a \tan(dx+c))}{(a^2+b^2)^2} + \frac{(-3a^2b+b^3) \ln(1+\tan^2(dx+c))}{2(a^2+b^2)^2} + \frac{(-a^3+3ab^2) \arctan(\tan(dx+c))}{(a^2+b^2)^2}$
default	$-\frac{a^2-b^2}{(a^2+b^2)(b+a \tan(dx+c))} + \frac{b(3a^2-b^2) \ln(b+a \tan(dx+c))}{(a^2+b^2)^2} + \frac{(-3a^2b+b^3) \ln(1+\tan^2(dx+c))}{2(a^2+b^2)^2} + \frac{(-a^3+3ab^2) \arctan(\tan(dx+c))}{(a^2+b^2)^2}$
norman	$\frac{(a^2-b^2)a \tan(dx+c)}{bd(a^2+b^2)} - \frac{a^2(a^2-3b^2)x \tan(dx+c)}{a^4+2a^2b^2+b^4} - \frac{ba(a^2-3b^2)x}{a^4+2a^2b^2+b^4} + \frac{b(3a^2-b^2) \ln(b+a \tan(dx+c))}{d(a^4+2a^2b^2+b^4)} - \frac{b(3a^2-b^2) \ln(1+\tan^2(dx+c))}{2d(a^4+2a^2b^2+b^4)}$
parallelrisch	$-\frac{2x \tan(dx+c)a^4bd-6x \tan(dx+c)a^2b^3d+3 \ln(1+\tan^2(dx+c)) \tan(dx+c)a^3b^2-\ln(1+\tan^2(dx+c)) \tan(dx+c)ab^4-6x^2 \tan^2(dx+c)a^4bd}{d^2(a^4+2a^2b^2+b^4)}$
risch	$\frac{ixb}{2iba+a^2-b^2} - \frac{xa}{2iba+a^2-b^2} - \frac{6ib a^2 x}{a^4+2a^2b^2+b^4} + \frac{2ib^3 x}{a^4+2a^2b^2+b^4} - \frac{6ib a^2 c}{d(a^4+2a^2b^2+b^4)} + \frac{2ib^3 c}{d(a^4+2a^2b^2+b^4)} - \frac{(-i)}{(-i)}$

input `int((a+b*tan(d*x+c))/(b+a*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*(-(a^2-b^2)/(a^2+b^2)/(b+a*tan(d*x+c))+b*(3*a^2-b^2)/(a^2+b^2)^2*ln(b+a*tan(d*x+c))+1/(a^2+b^2)^2*(1/2*(-3*a^2*b+b^3)*ln(1+tan(d*x+c)^2)+(-a^3+3*a*b^2)*arctan(tan(d*x+c))))`

3.316.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.89

$$\int \frac{a + b \tan(c + dx)}{(b + a \tan(c + dx))^2} dx =$$

$$-\frac{2a^4 - 2a^2b^2 + 2(a^3b - 3ab^3)dx - (3a^2b^2 - b^4 + (3a^3b - ab^3) \tan(dx + c)) \log\left(\frac{a^2 \tan(dx+c)^2 + 2ab \tan(dx+c) + b^2}{\tan(dx+c)^2 + 1}\right)}{2((a^5 + 2a^3b^2 + ab^4)d \tan(dx + c) + (a^4b + 2a^2b^3 + b^4))}$$

input `integrate((a+b*tan(d*x+c))/(b+a*tan(d*x+c))^2,x, algorithm="fricas")`

```
output -1/2*(2*a^4 - 2*a^2*b^2 + 2*(a^3*b - 3*a*b^3)*d*x - (3*a^2*b^2 - b^4 + (3*
a^3*b - a*b^3)*tan(d*x + c))*log((a^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c)
+ b^2)/(tan(d*x + c)^2 + 1)) - 2*(a^3*b - a*b^3 - (a^4 - 3*a^2*b^2)*d*x)*t
an(d*x + c))/((a^5 + 2*a^3*b^2 + a*b^4)*d*tan(d*x + c) + (a^4*b + 2*a^2*b^
3 + b^5)*d)
```

3.316.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.75 (sec) , antiderivative size = 1348, normalized size of antiderivative = 13.35

$$\int \frac{a + b \tan(c + dx)}{(b + a \tan(c + dx))^2} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))/(b+a*tan(d*x+c))**2,x)
```

```
output Piecewise((zoo*x*tan(c), Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (log(tan(c + d*x)
)**2 + 1)/(2*b*d), Eq(a, 0)), (I/(2*a*d*tan(c + d*x)**2 - 4*I*a*d*tan(c +
d*x) - 2*a*d), Eq(b, -I*a)), (-I/(2*a*d*tan(c + d*x)**2 + 4*I*a*d*tan(c +
d*x) - 2*a*d), Eq(b, I*a)), (x*(a + b*tan(c))/(a*tan(c) + b)**2, Eq(d, 0))
, (-2*a**4*d*x*tan(c + d*x)/(2*a**5*d*tan(c + d*x) + 2*a**4*b*d + 4*a**3*b
**2*d*tan(c + d*x) + 4*a**2*b**3*d + 2*a*b**4*d*tan(c + d*x) + 2*b**5*d) -
2*a**4/(2*a**5*d*tan(c + d*x) + 2*a**4*b*d + 4*a**3*b**2*d*tan(c + d*x) +
4*a**2*b**3*d + 2*a*b**4*d*tan(c + d*x) + 2*b**5*d) - 2*a**3*b*d*x/(2*a**
5*d*tan(c + d*x) + 2*a**4*b*d + 4*a**3*b**2*d*tan(c + d*x) + 4*a**2*b**3*d
+ 2*a*b**4*d*tan(c + d*x) + 2*b**5*d) + 6*a**3*b*log(tan(c + d*x) + b/a)*
tan(c + d*x)/(2*a**5*d*tan(c + d*x) + 2*a**4*b*d + 4*a**3*b**2*d*tan(c + d
*x) + 4*a**2*b**3*d + 2*a*b**4*d*tan(c + d*x) + 2*b**5*d) - 3*a**3*b*log(t
an(c + d*x)**2 + 1)*tan(c + d*x)/(2*a**5*d*tan(c + d*x) + 2*a**4*b*d + 4*a
**3*b**2*d*tan(c + d*x) + 4*a**2*b**3*d + 2*a*b**4*d*tan(c + d*x) + 2*b**5
*d) + 6*a**2*b**2*d*x*tan(c + d*x)/(2*a**5*d*tan(c + d*x) + 2*a**4*b*d + 4
*a**3*b**2*d*tan(c + d*x) + 4*a**2*b**3*d + 2*a*b**4*d*tan(c + d*x) + 2*b*
**5*d) + 6*a**2*b**2*log(tan(c + d*x) + b/a)/(2*a**5*d*tan(c + d*x) + 2*a**
4*b*d + 4*a**3*b**2*d*tan(c + d*x) + 4*a**2*b**3*d + 2*a*b**4*d*tan(c + d*
*x) + 2*b**5*d) - 3*a**2*b**2*log(tan(c + d*x)**2 + 1)/(2*a**5*d*tan(c + d*
*x) + 2*a**4*b*d + 4*a**3*b**2*d*tan(c + d*x) + 4*a**2*b**3*d + 2*a*b**4...
```

3.316.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.59

$$\int \frac{a + b \tan(c + dx)}{(b + a \tan(c + dx))^2} dx =$$

$$-\frac{\frac{2(a^3 - 3ab^2)(dx+c)}{a^4 + 2a^2b^2 + b^4} - \frac{2(3a^2b - b^3) \log(a \tan(dx+c) + b)}{a^4 + 2a^2b^2 + b^4} + \frac{(3a^2b - b^3) \log(\tan(dx+c)^2 + 1)}{a^4 + 2a^2b^2 + b^4} + \frac{2(a^2 - b^2)}{a^2b + b^3 + (a^3 + ab^2) \tan(dx+c)}}{2d}$$

input `integrate((a+b*tan(d*x+c))/(b+a*tan(d*x+c))^2,x, algorithm="maxima")`output `-1/2*(2*(a^3 - 3*a*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) - 2*(3*a^2*b - b^3)*log(a*tan(d*x + c) + b)/(a^4 + 2*a^2*b^2 + b^4) + (3*a^2*b - b^3)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) + 2*(a^2 - b^2)/(a^2*b + b^3 + (a^3 + a*b^2)*tan(d*x + c)))/d`**3.316.8 Giac [A] (verification not implemented)**

Time = 0.42 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.97

$$\int \frac{a + b \tan(c + dx)}{(b + a \tan(c + dx))^2} dx =$$

$$-\frac{\frac{2(a^3 - 3ab^2)(dx+c)}{a^4 + 2a^2b^2 + b^4} + \frac{(3a^2b - b^3) \log(\tan(dx+c)^2 + 1)}{a^4 + 2a^2b^2 + b^4} - \frac{2(3a^3b - ab^3) \log(|a \tan(dx+c) + b|)}{a^5 + 2a^3b^2 + ab^4} + \frac{2(3a^3b \tan(dx+c) - ab^3 \tan(dx+c) + a^4)}{(a^4 + 2a^2b^2 + b^4)(a \tan(dx+c))}}{2d}$$

input `integrate((a+b*tan(d*x+c))/(b+a*tan(d*x+c))^2,x, algorithm="giac")`output `-1/2*(2*(a^3 - 3*a*b^2)*(d*x + c)/(a^4 + 2*a^2*b^2 + b^4) + (3*a^2*b - b^3)*log(tan(d*x + c)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) - 2*(3*a^3*b - a*b^3)*log(abs(a*tan(d*x + c) + b))/(a^5 + 2*a^3*b^2 + a*b^4) + 2*(3*a^3*b*tan(d*x + c) - a*b^3*tan(d*x + c) + a^4 + 3*a^2*b^2 - 2*b^4)/((a^4 + 2*a^2*b^2 + b^4)*(a*tan(d*x + c) + b)))/d`

3.316.9 Mupad [B] (verification not implemented)

Time = 7.81 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.50

$$\int \frac{a + b \tan(c + dx)}{(b + a \tan(c + dx))^2} dx = \frac{b \ln(b + a \tan(c + dx)) (3a^2 - b^2)}{d(a^2 + b^2)^2} - \frac{\ln(\tan(c + dx) + 1i) (a - b 1i)}{2d(-a^2 1i + 2ab + b^2 1i)} - \frac{a^2 - b^2}{d(a^2 + b^2)(b + a \tan(c + dx))} - \frac{\ln(\tan(c + dx) - 1i) (a + b 1i)}{2d(a^2 1i + 2ab - b^2 1i)}$$

input `int((a + b*tan(c + d*x))/(b + a*tan(c + d*x))^2,x)`output `(b*log(b + a*tan(c + d*x))*(3*a^2 - b^2))/(d*(a^2 + b^2)^2) - (log(tan(c + d*x) + 1i)*(a - b*1i))/(2*d*(2*a*b - a^2*1i + b^2*1i)) - (a^2 - b^2)/(d*(a^2 + b^2)*(b + a*tan(c + d*x))) - (log(tan(c + d*x) - 1i)*(a + b*1i))/(2*d*(2*a*b + a^2*1i - b^2*1i))`

3.317 $\int \tan^3(c+dx) \sqrt{a + b \tan(c + dx)} (A+B \tan(c+dx)) dx$

3.317.1 Optimal result	2994
3.317.2 Mathematica [A] (verified)	2995
3.317.3 Rubi [A] (warning: unable to verify)	2995
3.317.4 Maple [B] (verified)	3000
3.317.5 Fricas [B] (verification not implemented)	3001
3.317.6 Sympy [F]	3002
3.317.7 Maxima [F]	3003
3.317.8 Giac [F(-1)]	3003
3.317.9 Mupad [B] (verification not implemented)	3004

3.317.1 Optimal result

Integrand size = 33, antiderivative size = 233

$$\int \tan^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\sqrt{a - ib}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}}\right)}{d} + \frac{\sqrt{a + ib}(A + iB) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a + ib}}\right)}{d}$$

$$- \frac{2A\sqrt{a + b \tan(c + dx)}}{d} - \frac{2(14aAb - 8a^2B + 35b^2B)(a + b \tan(c + dx))^{3/2}}{105b^3d}$$

$$+ \frac{2(7Ab - 4aB) \tan(c + dx)(a + b \tan(c + dx))^{3/2}}{35b^2d}$$

$$+ \frac{2B \tan^2(c + dx)(a + b \tan(c + dx))^{3/2}}{7bd}$$

output

```
(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))*(a-I*b)^(1/2)/d+(A+I
*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))*(a+I*b)^(1/2)/d-2*A*(a+b
*tan(d*x+c))^(1/2)/d-2/105*(14*A*a*b-8*B*a^2+35*B*b^2)*(a+b*tan(d*x+c))^(3
/2)/b^3/d+2/35*(7*A*b-4*B*a)*tan(d*x+c)*(a+b*tan(d*x+c))^(3/2)/b^2/d+2/7*B
*tan(d*x+c)^2*(a+b*tan(d*x+c))^(3/2)/b/d
```

3.317.2 Mathematica [A] (verified)

Time = 2.97 (sec) , antiderivative size = 212, normalized size of antiderivative = 0.91

$$\int \tan^3(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$= \frac{\sqrt{a-ib}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a-ib}}\right)}{d} + \frac{\sqrt{a+ib}(A+iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a+ib}}\right)}{d}$$

$$+ \frac{2\sqrt{a+b\tan(c+dx)}(-14a^2Ab-105Ab^3+8a^3B-35ab^2B-b(-7aAb+4a^2B+35b^2B)\tan(c+dx))}{105b^3d}$$

input `Integrate[Tan[c + d*x]^3*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`output `(Sqrt[a - I*b]*(A - I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/d + (Sqrt[a + I*b]*(A + I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/d + (2*Sqrt[a + b*Tan[c + d*x]]*(-14*a^2*A*b - 105*A*b^3 + 8*a^3*B - 35*a*b^2*B - b*(-7*a*A*b + 4*a^2*B + 35*b^2*B)*Tan[c + d*x] + 3*b^2*(7*A*b + a*B)*Tan[c + d*x]^2 + 15*b^3*B*Tan[c + d*x]^3))/(105*b^3*d)`**3.317.3 Rubi [A] (warning: unable to verify)**Time = 1.46 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.12, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 4090, 27, 3042, 4130, 27, 3042, 4113, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^3(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c+dx)^3\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow \text{4090}$$

$$\frac{2\int -\frac{1}{2}\tan(c+dx)\sqrt{a+b\tan(c+dx)}(-((7Ab-4aB)\tan^2(c+dx))+7bB\tan(c+dx)+4aB)dx}{\frac{7b}{2B\tan^2(c+dx)(a+b\tan(c+dx))^{3/2}} + 7bd}$$

$$3.317. \quad \int \tan^3(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\begin{array}{c}
 \downarrow 27 \\
 \frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \\
 \frac{\int \tan(c+dx) \sqrt{a+b \tan(c+dx)} \left(-((7Ab-4aB) \tan^2(c+dx)) + 7bB \tan(c+dx) + 4aB \right) dx}{7b} \\
 \downarrow 3042 \\
 \frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \\
 \frac{\int \tan(c+dx) \sqrt{a+b \tan(c+dx)} \left(-((7Ab-4aB) \tan^2(c+dx)^2) + 7bB \tan(c+dx) + 4aB \right) dx}{7b} \\
 \downarrow 4130 \\
 \frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \\
 \frac{2 \int \frac{1}{2} \sqrt{a+b \tan(c+dx)} (35A \tan(c+dx)b^2 + (-8Ba^2+14Aba+35b^2B) \tan^2(c+dx) + 2a(7Ab-4aB)) dx}{5b} - \frac{2(7Ab-4aB) \tan(c+dx)(a+b \tan(c+dx))}{5bd} \\
 \downarrow 27 \\
 \frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \\
 \frac{\int \sqrt{a+b \tan(c+dx)} (35A \tan(c+dx)b^2 + (-8Ba^2+14Aba+35b^2B) \tan^2(c+dx) + 2a(7Ab-4aB)) dx}{5b} - \frac{2(7Ab-4aB) \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} \\
 \downarrow 3042 \\
 \frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \\
 \frac{\int \sqrt{a+b \tan(c+dx)} (35A \tan(c+dx)b^2 + (-8Ba^2+14Aba+35b^2B) \tan(c+dx)^2 + 2a(7Ab-4aB)) dx}{5b} - \frac{2(7Ab-4aB) \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} \\
 \downarrow 4113 \\
 \frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \\
 \frac{\int \sqrt{a+b \tan(c+dx)} (35Ab^2 \tan(c+dx) - 35b^2B) dx + \frac{2(-8a^2B+14aAb+35b^2B)(a+b \tan(c+dx))^{3/2}}{3bd}}{5b} - \frac{2(7Ab-4aB) \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} \\
 \downarrow 3042 \\
 \frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \\
 \frac{\int \sqrt{a+b \tan(c+dx)} (35Ab^2 \tan(c+dx) - 35b^2B) dx + \frac{2(-8a^2B+14aAb+35b^2B)(a+b \tan(c+dx))^{3/2}}{3bd}}{5b} - \frac{2(7Ab-4aB) \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} \\
 \downarrow 4011
 \end{array}$$

3.317. $\int \tan^3(c+dx) \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$

$$\frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \frac{\int \frac{35b^2(aA-bB) \tan(c+dx) - 35b^2(Ab+aB)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{2(-8a^2B+14aAb+35b^2B)(a+b \tan(c+dx))^{3/2}}{3bd} + \frac{70Ab^2 \sqrt{a+b \tan(c+dx)}}{d}}{5b} - \frac{2(7Ab-4aB) \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd}$$

↓ 3042

$$\frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \frac{\int \frac{35b^2(aA-bB) \tan(c+dx) - 35b^2(Ab+aB)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{2(-8a^2B+14aAb+35b^2B)(a+b \tan(c+dx))^{3/2}}{3bd} + \frac{70Ab^2 \sqrt{a+b \tan(c+dx)}}{d}}{5b} - \frac{2(7Ab-4aB) \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd}$$

↓ 4022

$$\frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \frac{2(7Ab-4aB) \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} + \frac{35}{2} b^2(-b+ia)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx - \frac{35}{2} b^2(b+ia)(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \frac{2(-8a^2B)}{5b}$$

↓ 3042

$$\frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \frac{2(7Ab-4aB) \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} + \frac{35}{2} b^2(-b+ia)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx - \frac{35}{2} b^2(b+ia)(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \frac{2(-8a^2B)}{5b}$$

↓ 4020

$$\frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \frac{2(7Ab-4aB) \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} + \frac{35ib^2(b+ia)(A-iB) \int \frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} - \frac{35ib^2(-b+ia)(A+iB) \int \frac{1}{(i \tan(c+dx) + 1) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d}$$

↓ 25

$$\frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \frac{2(7Ab-4aB) \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} + \frac{35ib^2(b+ia)(A-iB) \int \frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \frac{35ib^2(-b+ia)(A+iB) \int \frac{1}{(i \tan(c+dx) + 1) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d}$$

↓ 73

$$\frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \frac{2(7Ab-4aB) \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} + \frac{35b(-b+ia)(A+iB) \int \frac{1}{-i \tan^2(c+dx) - \frac{ia}{b} + 1} d \sqrt{a+b \tan(c+dx)}}{d} - \frac{35b(b+ia)(A-iB) \int \frac{1}{i \tan^2(c+dx) + \frac{ia}{b}} d \sqrt{a+b \tan(c+dx)}}{d}$$

3.317. $\int \tan^3(c+dx) \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$

$$\begin{aligned}
 & \downarrow 221 \\
 & \frac{2B \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}}{7bd} - \\
 & \frac{-\frac{2(7Ab-4aB) \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} + \frac{2(-8a^2B+14aAb+35b^2B)(a+b \tan(c+dx))^{3/2}}{3bd} - \frac{35b^2(b+ia)(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} + \frac{35b^2(-b+ia)}{5b}}{7b}
 \end{aligned}$$

input `Int[Tan[c + d*x]^3*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(2*B*Tan[c + d*x]^2*(a + b*Tan[c + d*x])^(3/2))/(7*b*d) - ((-2*(7*A*b - 4*a*B)*Tan[c + d*x]*(a + b*Tan[c + d*x])^(3/2))/(5*b*d) + ((-35*b^2*(I*a + b)*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d) + (35*(I*a - b)*b^2*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d) + (70*A*b^2*Sqrt[a + b*Tan[c + d*x]])/d + (2*(14*a*A*b - 8*a^2*B + 35*b^2*B)*(a + b*Tan[c + d*x])^(3/2))/(3*b*d))/(5*b))/(7*b)`

3.317.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]^m * ((c_.) + (d_.)\tan[(e_.) + (f_.)x]), x_Symbol] \rightarrow \text{Simp}[d*(a + b*\tan[e + f*x])^m/(f*m), x] + \text{Int}[(a + b*\tan[e + f*x])^{m-1} * \text{Simp}[a*c - b*d + (b*c + a*d)*\tan[e + f*x], x], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]

rule 4020 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]^m * ((c_.) + (d_.)\tan[(e_.) + (f_.)x]), x_Symbol] \rightarrow \text{Simp}[c*(d/f) \text{Subst}[\text{Int}[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*\tan[e + f*x]], x] /;$ FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]

rule 4022 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]^m * ((c_.) + (d_.)\tan[(e_.) + (f_.)x]), x_Symbol] \rightarrow \text{Simp}[(c + I*d)/2 \text{Int}[(a + b*\tan[e + f*x])^m * (1 - I*\tan[e + f*x]), x], x] + \text{Simp}[(c - I*d)/2 \text{Int}[(a + b*\tan[e + f*x])^m * (1 + I*\tan[e + f*x]), x], x] /;$ FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]

rule 4090 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]^m * ((A_.) + (B_.)\tan[(e_.) + (f_.)x])^n * ((c_.) + (d_.)\tan[(e_.) + (f_.)x]), x_Symbol] \rightarrow \text{Simp}[b*B*(a + b*\tan[e + f*x])^{m-1} * ((c + d*\tan[e + f*x])^{n+1}/(d*f*(m+n))), x] + \text{Simp}[1/(d*(m+n)) \text{Int}[(a + b*\tan[e + f*x])^{m-2} * (c + d*\tan[e + f*x])^n * \text{Simp}[a^2*A*d*(m+n) - b*B*(b*c*(m-1) + a*d*(n+1)) + d*(m+n)*(2*a*A*b + B*(a^2 - b^2))*\tan[e + f*x] - (b*B*(b*c - a*d)*(m-1) - b*(A*b + a*B)*d*(m+n))*\tan[e + f*x]^2, x], x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

rule 4113 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]^m * ((A_.) + (B_.)\tan[(e_.) + (f_.)x]) + (C_.)\tan[(e_.) + (f_.)x]^2, x_Symbol] \rightarrow \text{Simp}[C*(a + b*\tan[e + f*x])^{m+1}/(b*f*(m+1)), x] + \text{Int}[(a + b*\tan[e + f*x])^m * \text{Simp}[A - C + B*\tan[e + f*x], x], x] /;$ FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]

```
rule 4130 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

3.317.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 936 vs. 2(201) = 402.

Time = 0.40 (sec) , antiderivative size = 937, normalized size of antiderivative = 4.02

method	result
parts	$2A \left(\frac{(a+b \tan(dx+c))^{\frac{5}{2}}}{5} - \frac{a(a+b \tan(dx+c))^{\frac{3}{2}}}{3} - \sqrt{a+b \tan(dx+c)} b^2 - b^2 \left(-\frac{\sqrt{2\sqrt{a^2+b^2}+2a} \ln \left(\frac{b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)}}{8} \right)}{8} \right) \right)$
derivativedivides	Expression too large to display
default	Expression too large to display

```
input int((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^3*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

3.317. $\int \tan^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

output

```

2*A/d/b^2*(1/5*(a+b*tan(d*x+c))^(5/2)-1/3*a*(a+b*tan(d*x+c))^(3/2)-(a+b*tan(d*x+c))^(1/2)*b^2-b^2*(-1/8*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))+1/2*(a-(a^2+b^2)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))+1/8*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))+1/2*((a^2+b^2)^(1/2)-a)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))))+B*(2/7/d/b^3*(a+b*tan(d*x+c))^(7/2)-4/5/d/b^3*(a+b*tan(d*x+c))^(5/2)*a+2/3/d/b^3*a^2*(a+b*tan(d*x+c))^(3/2)-2/3/d/b*(a+b*tan(d*x+c))^(3/2)-1/4/d/b*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))+1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))+1/4/d/b*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))+1/4/d/b*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))-1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))-1/4/d/b*(2*(a^2+b^2)...

```

3.317.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1312 vs. $2(195) = 390$.

Time = 0.27 (sec) , antiderivative size = 1312, normalized size of antiderivative = 5.63

$$\int \tan^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input

```

integrate((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^3*(A+B*tan(d*x+c)),x, algorithm="fricas")

```


output

```
-1/210*(105*b^3*d*sqrt(-(2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A
*B^3))*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(
2*(A^3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) + (B*d^3*sqrt
(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3))*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d
^4) + (2*A^2*B*a + (A^3 - A*B^2)*b)*d)*sqrt(-(2*A*B*b + d^2*sqrt(-(4*A^2*B
^2*a^2 + 4*(A^3*B - A*B^3))*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2
- B^2)*a)/d^2)) - 105*b^3*d*sqrt(-(2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*
(A^3*B - A*B^3))*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d
^2)*log(-(2*(A^3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) -
(B*d^3*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3))*a*b + (A^4 - 2*A^2*B^2 + B
^4)*b^2)/d^4) + (2*A^2*B*a + (A^3 - A*B^2)*b)*d)*sqrt(-(2*A*B*b + d^2*sqrt
(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3))*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^
4) - (A^2 - B^2)*a)/d^2)) - 105*b^3*d*sqrt(-(2*A*B*b - d^2*sqrt(-(4*A^2*B^
2*a^2 + 4*(A^3*B - A*B^3))*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 -
B^2)*a)/d^2)*log(-(2*(A^3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x +
c) + a) + (B*d^3*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3))*a*b + (A^4 - 2*A
^2*B^2 + B^4)*b^2)/d^4) - (2*A^2*B*a + (A^3 - A*B^2)*b)*d)*sqrt(-(2*A*B*b
- d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3))*a*b + (A^4 - 2*A^2*B^2 + B^
4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)) + 105*b^3*d*sqrt(-(2*A*B*b - d^2*sqrt(
-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3))*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/...
```

3.317.6 Sympy [F]

$$\int \tan^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} \tan^3(c + dx) dx$$

input `integrate((a+b*tan(d*x+c))**(1/2)*tan(d*x+c)**3*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))*tan(c + d*x)**3, x)`

3.317.7 Maxima [F]

$$\int \tan^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \tan(dx + c)^3 dx$$

input `integrate((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*tan(d*x + c)^3, x)`

3.317.8 Giac [F(-1)]

Timed out.

$$\int \tan^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.317.9 Mupad [B] (verification not implemented)

Time = 66.52 (sec) , antiderivative size = 1093, normalized size of antiderivative = 4.69

$$\int \tan^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \frac{2 A (a + b \tan(c + dx))^{5/2}}{5 b^2 d} - \sqrt{a + b \tan(c + dx)} \left(2 a \left(\frac{2 B (a^2 + b^2)}{b^3 d} - \frac{4 B a^2}{b^3 d} \right) + \frac{8 B a^3}{b^3 d} - \frac{4 B a (a^2 + b^2)}{b^3 d} \right) - \left(\frac{2 A (a^2 + b^2)}{b^2 d} - \frac{2 A a^2}{b^2 d} \right) \sqrt{a + b \tan(c + dx)} - \left(\frac{2 B (a^2 + b^2)}{3 b^3 d} - \frac{4 B a^2}{3 b^3 d} \right) (a + b \tan(c + dx))^{3/2} + \frac{2 B (a + b \tan(c + dx))^{7/2}}{7 b^3 d} - \frac{2 A a (a + b \tan(c + dx))^{3/2}}{3 b^2 d} - \frac{4 B a (a + b \tan(c + dx))^{5/2}}{5 b^3 d} + \operatorname{atan} \left(\frac{d^3 \left(\frac{16 (B^2 b^4 - B^2 a^2 b^2) \sqrt{a + b \tan(c + dx)}}{d^2} + \frac{16 a b^2 (\sqrt{-B^4 b^2 d^4 + B^2 a d^2}) \sqrt{a + b \tan(c + dx)}}{d^4} \right)}{8 (B^3 a^2 b^3 + B^3 b^5)} \right) \sqrt{-\frac{\sqrt{-B^4 b^2 d^4 + B^2 a d^2}}{4 d^4}}$$

input `int(tan(c + d*x)^3*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`

output `atan((d^3*((16*(B^2*b^4 - B^2*a^2*b^2)*(a + b*tan(c + d*x))^(1/2))/d^2 + (16*a*b^2*((-B^4*b^2*d^4)^(1/2) + B^2*a*d^2)*(a + b*tan(c + d*x))^(1/2))/d^4)*(-((-B^4*b^2*d^4)^(1/2) + B^2*a*d^2)/(4*d^4))^(1/2)*1i)/(8*(B^3*b^5 + B^3*a^2*b^3)))*(-((-B^4*b^2*d^4)^(1/2) + B^2*a*d^2)/(4*d^4))^(1/2)*2i - ((2*B*(a^2 + b^2))/(3*b^3*d) - (4*B*a^2)/(3*b^3*d))*(a + b*tan(c + d*x))^(3/2) + atan((d^3*((16*(B^2*b^4 - B^2*a^2*b^2)*(a + b*tan(c + d*x))^(1/2))/d^2 - (16*a*b^2*((-B^4*b^2*d^4)^(1/2) - B^2*a*d^2)*(a + b*tan(c + d*x))^(1/2))/d^4)*((-B^4*b^2*d^4)^(1/2) - B^2*a*d^2)/(4*d^4))^(1/2)*1i)/(8*(B^3*b^5 + B^3*a^2*b^3)))*((-B^4*b^2*d^4)^(1/2) - B^2*a*d^2)/(4*d^4))^(1/2)*2i - (a + b*tan(c + d*x))^(1/2)*(2*a*((2*B*(a^2 + b^2))/(b^3*d) - (4*B*a^2)/(b^3*d)) + (8*B*a^3)/(b^3*d) - (4*B*a*(a^2 + b^2))/(b^3*d)) - atan((A^2*b^4*((-A^4*b^2*d^4)^(1/2)/(4*d^4) + (A^2*a)/(4*d^2))^(1/2)*(a + b*tan(c + d*x))^(1/2)*32i)/((16*A*b^4*((-A^4*b^2*d^4)^(1/2))/d^3 + (16*A*a^2*b^2*((-A^4*b^2*d^4)^(1/2))/d^3) + (a*b^2*((-A^4*b^2*d^4)^(1/2))/(4*d^4) + (A^2*a)/(4*d^2))^(1/2)*(a + b*tan(c + d*x))^(1/2)*(-A^4*b^2*d^4)^(1/2)*32i)/((16*A*b^4*((-A^4*b^2*d^4)^(1/2))/d + (16*A*a^2*b^2*((-A^4*b^2*d^4)^(1/2))/d))*((-A^4*b^2*d^4)^(1/2) + A^2*a*d^2)/(4*d^4))^(1/2)*2i + atan((A^2*b^4*((A^2*a)/(4*d^2) - (-A^4*b^2*d^4)^(1/2)/(4*d^4))^(1/2)*(a + b*tan(c + d*x))^(1/2)*32i)/((16*A*b^4*((-A^4*b^2*d^4)^(1/2))/d^3 + (16*A*a^2*b^2*((-A^4*b^2*d^4)^(1/2))/d^3) - (a*b^2*((A^2*a)/(4*d^2) - (-A^4*b^2*d^4)^(1/2)/(4*d^4))^(1/2)*(a ...`

3.318 $\int \tan^2(c+dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.318.1 Optimal result	3005
3.318.2 Mathematica [A] (verified)	3006
3.318.3 Rubi [A] (warning: unable to verify)	3006
3.318.4 Maple [B] (verified)	3010
3.318.5 Fricas [B] (verification not implemented)	3011
3.318.6 Sympy [F]	3012
3.318.7 Maxima [F]	3013
3.318.8 Giac [F(-1)]	3013
3.318.9 Mupad [B] (verification not implemented)	3014

3.318.1 Optimal result

Integrand size = 33, antiderivative size = 186

$$\int \tan^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\sqrt{a - ib}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}}\right)}{d} - \frac{\sqrt{a + ib}(iA - B) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a + ib}}\right)}{d} - \frac{2B \sqrt{a + b \tan(c + dx)}}{d} + \frac{2(5Ab - 2aB)(a + b \tan(c + dx))^{3/2}}{15b^2d} + \frac{2B \tan(c + dx)(a + b \tan(c + dx))^{3/2}}{5bd}$$

output

```
(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))*(a-I*b)^(1/2)/d-(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))*(a+I*b)^(1/2)/d-2*B*(a+b*tan(d*x+c))^(1/2)/d+2/15*(5*A*b-2*B*a)*(a+b*tan(d*x+c))^(3/2)/b^2/d+2/5*B*tan(d*x+c)*(a+b*tan(d*x+c))^(3/2)/b/d
```

3.318.2 Mathematica [A] (verified)

Time = 2.09 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.91

$$\int \tan^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{15\sqrt{a - ib}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}}\right) + 15\sqrt{a + ib}(-iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a + ib}}\right) + 2\sqrt{a + b \tan(c + dx)}}{15d}$$

input `Integrate[Tan[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`output `(15*Sqrt[a - I*b]*(I*A + B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] + 15*Sqrt[a + I*b]*((-I)*A + B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + (2*Sqrt[a + b*Tan[c + d*x]]*(5*a*A*b - 2*a^2*B - 15*b^2*B + b*(5*A*b + a*B)*Tan[c + d*x] + 3*b^2*B*Tan[c + d*x]^2))/b^2)/(15*d)`**3.318.3 Rubi [A] (warning: unable to verify)**Time = 1.01 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.97, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 4090, 27, 3042, 4113, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^2 \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$\downarrow \text{4090}$$

$$\frac{2 \int -\frac{1}{2} \sqrt{a + b \tan(c + dx)} (-(5Ab - 2aB) \tan^2(c + dx) + 5bB \tan(c + dx) + 2aB) dx}{\frac{5b}{2B \tan(c + dx) (a + b \tan(c + dx))^{3/2}}} +$$

$$\frac{5bd}{5bd}$$

$$\downarrow \text{27}$$

$$\begin{aligned}
& \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} - \\
& \frac{\int \sqrt{a+b \tan(c+dx)}(-((5Ab-2aB) \tan^2(c+dx)) + 5bB \tan(c+dx) + 2aB) dx}{5b} \\
& \quad \downarrow \text{3042} \\
& \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} - \\
& \frac{\int \sqrt{a+b \tan(c+dx)}(-((5Ab-2aB) \tan(c+dx)^2) + 5bB \tan(c+dx) + 2aB) dx}{5b} \\
& \quad \downarrow \text{4113} \\
& \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} - \\
& \frac{\int \sqrt{a+b \tan(c+dx)}(5Ab + 5B \tan(c+dx)b)dx - \frac{2(5Ab-2aB)(a+b \tan(c+dx))^{3/2}}{3bd}}{5b} \\
& \quad \downarrow \text{3042} \\
& \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} - \\
& \frac{\int \sqrt{a+b \tan(c+dx)}(5Ab + 5B \tan(c+dx)b)dx - \frac{2(5Ab-2aB)(a+b \tan(c+dx))^{3/2}}{3bd}}{5b} \\
& \quad \downarrow \text{4011} \\
& \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} - \\
& \frac{\int \frac{5b(aA-bB)+5b(Ab+aB) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx - \frac{2(5Ab-2aB)(a+b \tan(c+dx))^{3/2}}{3bd} + \frac{10bB \sqrt{a+b \tan(c+dx)}}{d}}{5b} \\
& \quad \downarrow \text{3042} \\
& \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} - \\
& \frac{\int \frac{5b(aA-bB)+5b(Ab+aB) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx - \frac{2(5Ab-2aB)(a+b \tan(c+dx))^{3/2}}{3bd} + \frac{10bB \sqrt{a+b \tan(c+dx)}}{d}}{5b} \\
& \quad \downarrow \text{4022} \\
& \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} - \\
& \frac{\frac{5}{2}b(a+ib)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{5}{2}b(a-ib)(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx - \frac{2(5Ab-2aB)(a+b \tan(c+dx))^{3/2}}{3bd} + 10bB \sqrt{a+b \tan(c+dx)}}{5b} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

$$\begin{aligned}
 & \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} - \\
 & \frac{\frac{5}{2}b(a+ib)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{5}{2}b(a-ib)(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx - \frac{2(5Ab-2aB)(a+b \tan(c+dx))^{3/2}}{3bd} + 10}{5b} \\
 & \quad \downarrow 4020 \\
 & \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} - \\
 & \frac{\frac{5ib(a-ib)(A-iB) \int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} - \frac{5ib(a+ib)(A+iB) \int -\frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d}}{5b} \\
 & \quad \downarrow 25 \\
 & \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} - \\
 & \frac{-\frac{5ib(a-ib)(A-iB) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \frac{5ib(a+ib)(A+iB) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d}}{5b} \\
 & \quad \downarrow 73 \\
 & \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} - \\
 & \frac{\frac{5(a+ib)(A+iB) \int -\frac{1}{i \tan^2(c+dx) - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{d} + \frac{5(a-ib)(A-iB) \int \frac{1}{i \tan^2(c+dx) + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{d} - \frac{2(5Ab-2aB)(a+b \tan(c+dx))^{3/2}}{3bd}}{5b} \\
 & \quad \downarrow 221 \\
 & \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{3/2}}{5bd} - \\
 & \frac{\frac{5b\sqrt{a-ib}(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{5b\sqrt{a+ib}(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} - \frac{2(5Ab-2aB)(a+b \tan(c+dx))^{3/2}}{3bd} + \frac{10bB\sqrt{a+b \tan(c+dx)}}{d}}{5b}
 \end{aligned}$$

input `Int[Tan[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(2*B*Tan[c + d*x]*(a + b*Tan[c + d*x])^(3/2))/(5*b*d) - ((5*Sqrt[a - I*b]*b*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d + (5*Sqrt[a + I*b]*b*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d + (10*b*B*Sqrt[a + b*Tan[c + d*x]])/d - (2*(5*A*b - 2*a*B)*(a + b*Tan[c + d*x])^(3/2))/(3*b*d))/(5*b)`

3.318.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`


```
rule 4090 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4113 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

3.318.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 872 vs. 2(158) = 316.

Time = 0.13 (sec) , antiderivative size = 873, normalized size of antiderivative = 4.69

method	result
parts	$A \left(\frac{2(a+b \tan(dx+c))^{\frac{3}{2}}}{3db} - \frac{\sqrt{2\sqrt{a^2+b^2}+2a} a \ln(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}})}{4db} \right) +$
derivativedivides	Expression too large to display
default	Expression too large to display

```
input int((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^2*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

$$3.318. \quad \int \tan^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

output

```
A*(2/3/d/b*(a+b*tan(d*x+c))^(3/2)-1/4/d/b*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a*
ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^
2+b^2)^(1/2))+1/4/d/b*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*ln(b*t
an(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)
^(1/2))-1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/
2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))+1/4/d/b*(
2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2
)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))-1/4/d/b*(2*(a^2+b^2)^(1/2)+2*
a)^(1/2)*(a^2+b^2)^(1/2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)
^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))+1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)
*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b
^2)^(1/2)-2*a)^(1/2)))+2*B/d/b^2*(1/5*(a+b*tan(d*x+c))^(5/2)-1/3*a*(a+b*ta
n(d*x+c))^(3/2)-(a+b*tan(d*x+c))^(1/2)*b^2-b^2*(-1/8*(2*(a^2+b^2)^(1/2)+2*
a)^(1/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(
1/2)+(a^2+b^2)^(1/2))+1/2*(a-(a^2+b^2)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/
2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^
2+b^2)^(1/2)-2*a)^(1/2))+1/8*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*ln((a+b*tan(d*x+
c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))+1/
2*((a^2+b^2)^(1/2)-a)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(
1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)...
```

3.318.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1265 vs. $2(152) = 304$.

Time = 0.27 (sec) , antiderivative size = 1265, normalized size of antiderivative = 6.80

$$\int \tan^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input

```
integrate((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^2*(A+B*tan(d*x+c)),x, algorithm
m="fricas")
```

```
output 1/30*(15*b^2*d*sqrt((2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)
)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2*(A
^3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) + (A*d^3*sqrt(-(
4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4)
- (2*A*B^2*a + (A^2*B - B^3)*b)*d)*sqrt((2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^
2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2
)*a)/d^2)) - 15*b^2*d*sqrt((2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B
- A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log
(-(2*(A^3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) - (A*d^3*
sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2
)/d^4) - (2*A*B^2*a + (A^2*B - B^3)*b)*d)*sqrt((2*A*B*b + d^2*sqrt(-(4*A^2
*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^
2 - B^2)*a)/d^2)) - 15*b^2*d*sqrt((2*A*B*b - d^2*sqrt(-(4*A^2*B^2*a^2 + 4*
(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d
^2)*log(-(2*(A^3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) +
(A*d^3*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B
^4)*b^2)/d^4) + (2*A*B^2*a + (A^2*B - B^3)*b)*d)*sqrt((2*A*B*b - d^2*sqrt(
-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4
) - (A^2 - B^2)*a)/d^2)) + 15*b^2*d*sqrt((2*A*B*b - d^2*sqrt(-(4*A^2*B^2*a
^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - ...
```

3.318.6 Sympy [F]

$$\int \tan^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} \tan^2(c + dx) dx$$

```
input integrate((a+b*tan(d*x+c))**(1/2)*tan(d*x+c)**2*(A+B*tan(d*x+c)),x)
```

```
output Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))*tan(c + d*x)**2, x)
```

3.318.7 Maxima [F]

$$\int \tan^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \tan(dx + c)^2 dx$$

input `integrate((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*tan(d*x + c)^2, x)`

3.318.8 Giac [F(-1)]

Timed out.

$$\int \tan^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.318.9 Mupad [B] (verification not implemented)

Time = 25.43 (sec) , antiderivative size = 938, normalized size of antiderivative = 5.04

$$\begin{aligned}
& \int \tan^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx \\
&= \operatorname{atanh} \left(\frac{d^3 \left(\frac{16(A^2 b^4 - A^2 a^2 b^2) \sqrt{a + b \tan(c + dx)}}{d^2} + \frac{16 a b^2 (\sqrt{-A^4 b^2 d^4 + A^2 a d^2}) \sqrt{a + b \tan(c + dx)}}{d^4} \right) \sqrt{-\frac{\sqrt{-A^4 b^2 d^4 + A^2 a d^2}}{d^4}}}{16(A^3 a^2 b^3 + A^3 b^5)} \right) \\
&\quad - \left(\frac{2 B (a^2 + b^2)}{b^2 d} - \frac{2 B a^2}{b^2 d} \right) \sqrt{a + b \tan(c + dx)} \\
&\quad + \operatorname{atanh} \left(\frac{d^3 \left(\frac{16(A^2 b^4 - A^2 a^2 b^2) \sqrt{a + b \tan(c + dx)}}{d^2} - \frac{16 a b^2 (\sqrt{-A^4 b^2 d^4 - A^2 a d^2}) \sqrt{a + b \tan(c + dx)}}{d^4} \right) \sqrt{\frac{\sqrt{-A^4 b^2 d^4 - A^2 a d^2}}{d^4}}}{16(A^3 a^2 b^3 + A^3 b^5)} \right) \\
&\quad + \frac{2 A (a + b \tan(c + dx))^{3/2}}{3 b d} + \frac{2 B (a + b \tan(c + dx))^{5/2}}{5 b^2 d} - \frac{2 B a (a + b \tan(c + dx))^{3/2}}{3 b^2 d} \\
&\quad - \operatorname{atan} \left(\frac{B^2 b^4 \sqrt{\frac{\sqrt{-B^4 b^2 d^4}}{4 d^4}} + \frac{B^2 a}{4 d^2} \sqrt{a + b \tan(c + dx)}}{\frac{16 B b^4 \sqrt{-B^4 b^2 d^4}}{d^3} + \frac{16 B a^2 b^2 \sqrt{-B^4 b^2 d^4}}{d^3}} + \frac{a b^2 \sqrt{\frac{\sqrt{-B^4 b^2 d^4}}{4 d^4}} + \frac{B^2 a}{4 d^2} \sqrt{a + b \tan(c + dx)}}{\frac{16 B b^4 \sqrt{-B^4 b^2 d^4}}{d} + \frac{16 B a^2 b^2 \sqrt{-B^4 b^2 d^4}}{d}} \right)
\end{aligned}$$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`

output

$$\begin{aligned} & \operatorname{atanh}\left(\frac{d^3 \left((16A^2b^4 - A^2a^2b^2)(a + b \tan(c + dx))^{1/2} \right)}{d^2 + (16ab^2(-A^4b^2d^4)^{1/2} + A^2ad^2)(a + b \tan(c + dx))^{1/2}}\right) / d^4 \\ & \left(-\left((-A^4b^2d^4)^{1/2} + A^2ad^2 \right) / d^4 \right)^{1/2} / (16(A^3b^5 + A^3a^2b^3)) \\ & \left(-\left((-A^4b^2d^4)^{1/2} + A^2ad^2 \right) / d^4 \right)^{1/2} - \left((2B(a^2 + b^2)) / (b^2d) - (2Ba^2) / (b^2d) \right) (a + b \tan(c + dx))^{1/2} \\ & + \operatorname{atanh}\left(\frac{d^3 \left((16A^2b^4 - A^2a^2b^2)(a + b \tan(c + dx))^{1/2} \right)}{d^2 - (16ab^2(-A^4b^2d^4)^{1/2} - A^2ad^2)(a + b \tan(c + dx))^{1/2}}\right) / d^4 \\ & \left(\left((-A^4b^2d^4)^{1/2} - A^2ad^2 \right) / d^4 \right)^{1/2} / (16(A^3b^5 + A^3a^2b^3)) \\ & \left(\left((-A^4b^2d^4)^{1/2} - A^2ad^2 \right) / d^4 \right)^{1/2} - \operatorname{atan}\left(\frac{B^2b^4(-B^4b^2d^4)^{1/2}}{4d^4} + \frac{B^2a}{4d^2}\right)^{1/2} (a + b \tan(c + dx))^{1/2} \\ & * 32i / \left(\frac{16Bb^4(-B^4b^2d^4)^{1/2}}{d^3} + \frac{16Ba^2b^2(-B^4b^2d^4)^{1/2}}{d^3} + (ab^2(-B^4b^2d^4)^{1/2} / (4d^4) + (B^2a) / (4d^2))^{1/2} (a + b \tan(c + dx))^{1/2} \right. \\ & \left. * (-B^4b^2d^4)^{1/2} * 32i \right) / \left(\frac{16Bb^4(-B^4b^2d^4)^{1/2}}{d} + \frac{16Ba^2b^2(-B^4b^2d^4)^{1/2}}{d} \right) \\ & \left(\left((-B^4b^2d^4)^{1/2} + B^2ad^2 \right) / (4d^4) \right)^{1/2} * 2i + \operatorname{atan}\left(\frac{B^2b^4((B^2a) / (4d^2) - (-B^4b^2d^4)^{1/2} / (4d^4))^{1/2}}{(16Bb^4(-B^4b^2d^4)^{1/2}) / d^3} + \frac{16Ba^2b^2(-B^4b^2d^4)^{1/2}}{d^3} - (ab^2((B^2a) / (4d^2) - (-B^4b^2d^4)^{1/2} / (4d^4))^{1/2} (a + b \tan(c + dx))^{1/2} * 32i}{(16Bb^4(-B^4b^2d^4)^{1/2}) / d} + \frac{16Ba^2b^2(-B^4b^2d^4)^{1/2}}{d} \right) \\ & \left(-\left((-B^4b^2d^4)^{1/2} - B^2ad^2 \right) \dots \right) \end{aligned}$$

3.319 $\int \tan(c+dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.319.1 Optimal result	3016
3.319.2 Mathematica [A] (verified)	3016
3.319.3 Rubi [A] (warning: unable to verify)	3017
3.319.4 Maple [B] (verified)	3020
3.319.5 Fricas [B] (verification not implemented)	3021
3.319.6 Sympy [F]	3022
3.319.7 Maxima [F]	3023
3.319.8 Giac [F(-1)]	3023
3.319.9 Mupad [B] (verification not implemented)	3024

3.319.1 Optimal result

Integrand size = 31, antiderivative size = 146

$$\int \tan(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= -\frac{\sqrt{a - ib}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}}\right) - \sqrt{a + ib}(A + iB) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a + ib}}\right)}{d} + \frac{2A\sqrt{a + b \tan(c + dx)}}{d} + \frac{2B(a + b \tan(c + dx))^{3/2}}{3bd}$$

```
output -(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))*(a-I*b)^(1/2)/d-(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))*(a+I*b)^(1/2)/d+2*A*(a+b*tan(d*x+c))^(1/2)/d+2/3*B*(a+b*tan(d*x+c))^(3/2)/b/d
```

3.319.2 Mathematica [A] (verified)

Time = 0.55 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.96

$$\int \tan(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{-3\sqrt{a - ib}b(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}}\right) - 3\sqrt{a + ib}b(A + iB) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a + ib}}\right) + 2\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{3bd}$$

input `Integrate[Tan[c + d*x]*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(-3*Sqrt[a - I*b]*b*(A - I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] - 3*Sqrt[a + I*b]*b*(A + I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + 2*Sqrt[a + b*Tan[c + d*x]]*(3*A*b + a*B + b*B*Tan[c + d*x]))/(3*b*d)`

3.319.3 Rubi [A] (warning: unable to verify)

Time = 0.73 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.99, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4075, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4075} \\
 & \int (A \tan(c + dx) - B) \sqrt{a + b \tan(c + dx)} dx + \frac{2B(a + b \tan(c + dx))^{3/2}}{3bd} \\
 & \quad \downarrow \text{3042} \\
 & \int (A \tan(c + dx) - B) \sqrt{a + b \tan(c + dx)} dx + \frac{2B(a + b \tan(c + dx))^{3/2}}{3bd} \\
 & \quad \downarrow \text{4011} \\
 & \int \frac{-Ab - aB + (aA - bB) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2A \sqrt{a + b \tan(c + dx)}}{d} + \frac{2B(a + b \tan(c + dx))^{3/2}}{3bd} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{-Ab - aB + (aA - bB) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2A \sqrt{a + b \tan(c + dx)}}{d} + \frac{2B(a + b \tan(c + dx))^{3/2}}{3bd} \\
 & \quad \downarrow \text{4022}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{2}(-b+ia)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2}(b+ia)(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \\
& \quad \frac{2A\sqrt{a+b \tan(c+dx)}}{d} + \frac{2B(a+b \tan(c+dx))^{3/2}}{3bd} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2}(-b+ia)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2}(b+ia)(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \\
& \quad \frac{2A\sqrt{a+b \tan(c+dx)}}{d} + \frac{2B(a+b \tan(c+dx))^{3/2}}{3bd} \\
& \quad \downarrow \text{4020} \\
& \frac{i(b+ia)(A-iB) \int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} - \\
& \frac{i(-b+ia)(A+iB) \int -\frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \frac{2A\sqrt{a+b \tan(c+dx)}}{d} + \\
& \quad \frac{2B(a+b \tan(c+dx))^{3/2}}{3bd} \\
& \quad \downarrow \text{25} \\
& \frac{i(b+ia)(A-iB) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \\
& \frac{i(-b+ia)(A+iB) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \frac{2A\sqrt{a+b \tan(c+dx)}}{d} + \\
& \quad \frac{2B(a+b \tan(c+dx))^{3/2}}{3bd} \\
& \quad \downarrow \text{73} \\
& \frac{(-b+ia)(A+iB) \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} - \\
& \frac{(b+ia)(A-iB) \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \frac{2A\sqrt{a+b \tan(c+dx)}}{d} + \\
& \quad \frac{2B(a+b \tan(c+dx))^{3/2}}{3bd} \\
& \quad \downarrow \text{221} \\
& \frac{(b+ia)(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} + \frac{(-b+ia)(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} + \\
& \quad \frac{2A\sqrt{a+b \tan(c+dx)}}{d} + \frac{2B(a+b \tan(c+dx))^{3/2}}{3bd}
\end{aligned}$$

input `Int[Tan[c + d*x]*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

3.319. $\int \tan(c+dx)\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$

```
output -(((I*a + b)*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*
d)) + ((I*a - b)*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I
*b]*d) + (2*A*Sqrt[a + b*Tan[c + d*x]])/d + (2*B*(a + b*Tan[c + d*x])^(3/2
))/ (3*b*d)
```

3.319.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4011 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

3.319.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 828 vs. 2(122) = 244.

Time = 0.11 (sec) , antiderivative size = 829, normalized size of antiderivative = 5.68

method	result
parts	$A \left(\frac{2\sqrt{a+b \tan(dx+c)}}{2\sqrt{a+b \tan(dx+c)}} - \frac{\sqrt{2\sqrt{a^2+b^2+2a} \ln(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2+2a}+\sqrt{a^2+b^2}})}{4} + \frac{(a-\sqrt{a^2+b^2}) \arctan(\dots)}{4} \right)$
derivativedivides	$\frac{2B(a+b \tan(dx+c))^{\frac{3}{2}}}{3bd} + \frac{2A\sqrt{a+b \tan(dx+c)}}{d} - \frac{\ln(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2+2a}+\sqrt{a^2+b^2}})}{4d} A$
default	$\frac{2B(a+b \tan(dx+c))^{\frac{3}{2}}}{3bd} + \frac{2A\sqrt{a+b \tan(dx+c)}}{d} - \frac{\ln(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2+2a}+\sqrt{a^2+b^2}})}{4d} A$

```
input int((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

$$3.319. \int \tan(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

output

```
A/d*(2*(a+b*tan(d*x+c))^(1/2)-1/4*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*ln(b*tan(d
*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/
2)))+(a-(a^2+b^2)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d
*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))
+1/4*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(
1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))+((a^2+b^2)^(1/2)-a)/(2*(a
^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(
d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)))+B*(2/3/d/b*(a+b*tan(d*x+c))
^(3/2)-1/4/d/b*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a*ln(b*tan(d*x+c)+a+(a+b*tan(
d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))+1/4/d/b*(2*(a
^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c)
)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))-1/d*b/(2*(a^2+b^2)^(
1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(
1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))+1/4/d/b*(2*(a^2+b^2)^(1/2)+2*a)^(1/2
)*a*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a
-(a^2+b^2)^(1/2))-1/4/d/b*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*ln
((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+
b^2)^(1/2))+1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)
+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)))
```

3.319.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1228 vs. 2(116) = 232.

Time = 0.29 (sec) , antiderivative size = 1228, normalized size of antiderivative = 8.41

$$\int \tan(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input

```
integrate((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)*(A+B*tan(d*x+c)),x, algorithm=
"fricas")
```

output

```

1/6*(3*b*d*sqrt(-(2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a
*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2*(A^3*
B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) + (B*d^3*sqrt(-(4*A
^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) + (
2*A^2*B*a + (A^3 - A*B^2)*b)*d)*sqrt(-(2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2
+ 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*
a)/d^2)) - 3*b*d*sqrt(-(2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*
B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2
*(A^3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) - (B*d^3*sqrt
(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^
4) + (2*A^2*B*a + (A^3 - A*B^2)*b)*d)*sqrt(-(2*A*B*b + d^2*sqrt(-(4*A^2*B^
2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 -
B^2)*a)/d^2)) - 3*b*d*sqrt(-(2*A*B*b - d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*
B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*l
og(-(2*(A^3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) + (B*d^
3*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b
^2)/d^4) - (2*A^2*B*a + (A^3 - A*B^2)*b)*d)*sqrt(-(2*A*B*b - d^2*sqrt(-(4*
A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) -
(A^2 - B^2)*a)/d^2)) + 3*b*d*sqrt(-(2*A*B*b - d^2*sqrt(-(4*A^2*B^2*a^2 + 4
*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*...

```

3.319.6 Sympy [F]

$$\int \tan(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} \tan(c + dx) dx$$

input `integrate((a+b*tan(d*x+c))**(1/2)*tan(d*x+c)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))*tan(c + d*x), x)`

3.319.7 Maxima [F]

$$\int \tan(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \tan(dx + c) dx$$

input `integrate((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*tan(d*x + c), x)`

3.319.8 Giac [F(-1)]

Timed out.

$$\int \tan(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.319.9 Mupad [B] (verification not implemented)

Time = 14.08 (sec) , antiderivative size = 864, normalized size of antiderivative = 5.92

$$\begin{aligned}
& \int \tan(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx \\
&= \operatorname{atanh} \left(\frac{d^3 \left(\frac{16 (B^2 b^4 - B^2 a^2 b^2) \sqrt{a + b \tan(c + dx)}}{d^2} + \frac{16 a b^2 (\sqrt{-B^4 b^2 d^4 + B^2 a d^2}) \sqrt{a + b \tan(c + dx)}}{d^4} \right) \sqrt{-\frac{\sqrt{-B^4 b^2 d^4 + B^2 a d^2}}{d^4}}}{16 (B^3 a^2 b^3 + B^3 b^5)} \right) \\
&+ \operatorname{atanh} \left(\frac{d^3 \left(\frac{16 (B^2 b^4 - B^2 a^2 b^2) \sqrt{a + b \tan(c + dx)}}{d^2} - \frac{16 a b^2 (\sqrt{-B^4 b^2 d^4 - B^2 a d^2}) \sqrt{a + b \tan(c + dx)}}{d^4} \right) \sqrt{\frac{\sqrt{-B^4 b^2 d^4 - B^2 a d^2}}{d^4}}}{16 (B^3 a^2 b^3 + B^3 b^5)} \right) \\
&- 2 \operatorname{atanh} \left(\frac{32 A^2 b^4 \sqrt{\frac{\sqrt{-A^4 b^2 d^4}}{4 d^4}} + \frac{A^2 a}{4 d^2} \sqrt{a + b \tan(c + dx)}}{\frac{16 A b^4 \sqrt{-A^4 b^2 d^4}}{d^3} + \frac{16 A a^2 b^2 \sqrt{-A^4 b^2 d^4}}{d^3}} \right) \\
&+ \frac{32 a b^2 \sqrt{\frac{\sqrt{-A^4 b^2 d^4}}{4 d^4}} + \frac{A^2 a}{4 d^2} \sqrt{a + b \tan(c + dx)} \sqrt{-A^4 b^2 d^4}}{\frac{16 A b^4 \sqrt{-A^4 b^2 d^4}}{d} + \frac{16 A a^2 b^2 \sqrt{-A^4 b^2 d^4}}{d}} \sqrt{\frac{\sqrt{-A^4 b^2 d^4} + A^2 a d^2}{4 d^4}} \\
&+ 2 \operatorname{atanh} \left(\frac{32 A^2 b^4 \sqrt{\frac{A^2 a}{4 d^2} - \frac{\sqrt{-A^4 b^2 d^4}}{4 d^4}} \sqrt{a + b \tan(c + dx)}}{\frac{16 A b^4 \sqrt{-A^4 b^2 d^4}}{d^3} + \frac{16 A a^2 b^2 \sqrt{-A^4 b^2 d^4}}{d^3}} \right) \\
&- \frac{32 a b^2 \sqrt{\frac{A^2 a}{4 d^2} - \frac{\sqrt{-A^4 b^2 d^4}}{4 d^4}} \sqrt{a + b \tan(c + dx)} \sqrt{-A^4 b^2 d^4}}{\frac{16 A b^4 \sqrt{-A^4 b^2 d^4}}{d} + \frac{16 A a^2 b^2 \sqrt{-A^4 b^2 d^4}}{d}} \sqrt{\frac{\sqrt{-A^4 b^2 d^4} - A^2 a d^2}{4 d^4}} \\
&+ \frac{2 A \sqrt{a + b \tan(c + dx)}}{d} + \frac{2 B (a + b \tan(c + dx))^{3/2}}{3 b d}
\end{aligned}$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`

output

$$\begin{aligned} & \operatorname{atanh}\left(\frac{d^3 \left((16(B^2 b^4 - B^2 a^2 b^2)(a + b \tan(c + dx))^{1/2})/d^2 + \right.}{\left. (16 a b^2 ((-B^4 b^2 d^4)^{1/2} + B^2 a d^2)(a + b \tan(c + dx))^{1/2})/d^4 \right)} \right) \\ & \left(- \left((-B^4 b^2 d^4)^{1/2} + B^2 a d^2 \right) / d^4 \right)^{1/2} \Big/ \left(16(B^3 b^5 + B^3 a^2 b^3) \right) \\ & \left(- \left((-B^4 b^2 d^4)^{1/2} + B^2 a d^2 \right) / d^4 \right)^{1/2} + \operatorname{atanh}\left(\frac{d^3 \left((16 \right.}{\left. (B^2 b^4 - B^2 a^2 b^2)(a + b \tan(c + dx))^{1/2})/d^2 - (16 a b^2 ((-B^4 \right.}{\left. b^2 d^4)^{1/2} - B^2 a d^2)(a + b \tan(c + dx))^{1/2})/d^4 \right)} \right) \\ & \left(\left((-B^4 b^2 d^4)^{1/2} - B^2 a d^2 \right) / d^4 \right)^{1/2} \Big/ \left(16(B^3 b^5 + B^3 a^2 b^3) \right) \\ & \left(\left((-B^4 b^2 d^4)^{1/2} - B^2 a d^2 \right) / d^4 \right)^{1/2} - 2 \operatorname{atanh}\left(\frac{32 A^2 b^4 ((-A^4 b^2 d^4)^{1/2})}{(4 d^4)} + \frac{(A^2 a)}{(4 d^2)} \right)^{1/2} \\ & \left(a + b \tan(c + dx) \right)^{1/2} \Big/ \left((16 A b^4 (-A^4 b^2 d^4)^{1/2}) / d^3 + (16 A a^2 b^2 (-A^4 b^2 d^4)^{1/2}) / d^3 \right) \\ & + (32 a b^2 ((-A^4 b^2 d^4)^{1/2}) / (4 d^4) + (A^2 a) / (4 d^2))^{1/2} \\ & \left(a + b \tan(c + dx) \right)^{1/2} \left((-A^4 b^2 d^4)^{1/2} \right) \Big/ \left((16 A b^4 (-A^4 b^2 d^4)^{1/2}) / d + \right. \\ & \left. (16 A a^2 b^2 (-A^4 b^2 d^4)^{1/2}) / d \right) \left(\left((-A^4 b^2 d^4)^{1/2} + A^2 a d^2 \right) / (4 d^4) \right)^{1/2} \\ & + 2 \operatorname{atanh}\left(\frac{32 A^2 b^4 ((A^2 a) / (4 d^2) - (-A^4 b^2 d^4)^{1/2})}{(4 d^4)} \right)^{1/2} \\ & \left(a + b \tan(c + dx) \right)^{1/2} \Big/ \left((16 A b^4 (-A^4 b^2 d^4)^{1/2}) / d^3 + \right. \\ & \left. (16 A a^2 b^2 (-A^4 b^2 d^4)^{1/2}) / d^3 - (32 a b^2 ((A^2 a) / (4 d^2) - \right. \\ & \left. (-A^4 b^2 d^4)^{1/2}) / (4 d^4) \right)^{1/2} \left(a + b \tan(c + dx) \right)^{1/2} \left((-A^4 b^2 d^4)^{1/2} \right) \\ & \Big/ \left((16 A b^4 (-A^4 b^2 d^4)^{1/2}) / d + (16 A a^2 b^2 (-A^4 b^2 d^4)^{1/2}) / d \right) \\ & \left(- \left((-A^4 b^2 d^4)^{1/2} - A^2 a d^2 \right) / (4 d^4) \right)^{1/2} + (2 A (a + b \tan(c + dx))^{1/2}) / d + (2 B (a + b \tan(c + d \dots \end{aligned}$$

3.320 $\int \sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx$

3.320.1 Optimal result	3026
3.320.2 Mathematica [A] (verified)	3026
3.320.3 Rubi [A] (warning: unable to verify)	3027
3.320.4 Maple [B] (verified)	3030
3.320.5 Fricas [B] (verification not implemented)	3031
3.320.6 Sympy [F]	3031
3.320.7 Maxima [F(-2)]	3032
3.320.8 Giac [F(-1)]	3032
3.320.9 Mupad [B] (verification not implemented)	3033

3.320.1 Optimal result

Integrand size = 25, antiderivative size = 122

$$\int \sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$= -\frac{\sqrt{a - ib}(iA + B)\operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}}\right)}{d}$$

$$+ \frac{\sqrt{a + ib}(iA - B)\operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a + ib}}\right)}{d} + \frac{2B\sqrt{a + b \tan(c + dx)}}{d}$$

output `-(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))*(a-I*b)^(1/2)/d+(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))*(a+I*b)^(1/2)/d+2*B*(a+b*tan(d*x+c))^(1/2)/d`

3.320.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.98

$$\int \sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx$$

$$= \frac{-i\sqrt{a - ib}(A - iB)\operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}}\right) + i\sqrt{a + ib}(A + iB)\operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a + ib}}\right) + 2B\sqrt{a + b \tan(c + dx)}}{d}$$

input `Integrate[Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `((-I)*Sqrt[a - I*b]*(A - I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] + I*Sqrt[a + I*b]*(A + I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + 2*B*Sqrt[a + b*Tan[c + d*x]])/d`

3.320.3 Rubi [A] (warning: unable to verify)

Time = 0.56 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.84, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$, Rules used = {3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4011} \\
 & \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2B \sqrt{a + b \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2B \sqrt{a + b \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{4022} \\
 & \frac{1}{2}(a + ib)(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a - ib)(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \\
 & \quad \frac{2B \sqrt{a + b \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2}(a + ib)(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a - ib)(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \\
 & \quad \frac{2B \sqrt{a + b \tan(c + dx)}}{d}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 4020 \\
& \frac{i(a-ib)(A-ib) \int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} - \\
& \frac{i(a+ib)(A+ib) \int -\frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \frac{2B\sqrt{a+b \tan(c+dx)}}{d} \\
& \downarrow 25 \\
& -\frac{i(a-ib)(A-ib) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \\
& \frac{i(a+ib)(A+ib) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \frac{2B\sqrt{a+b \tan(c+dx)}}{d} \\
& \downarrow 73 \\
& \frac{(a+ib)(A+ib) \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \\
& \frac{(a-ib)(A-ib) \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \frac{2B\sqrt{a+b \tan(c+dx)}}{d} \\
& \downarrow 221 \\
& \frac{\sqrt{a-ib}(A-ib) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{\sqrt{a+ib}(A+ib) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} + \\
& \frac{2B\sqrt{a+b \tan(c+dx)}}{d}
\end{aligned}$$

input `Int[Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[a - I*b]*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d + (Sqrt[a + I*b]*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d + (2*B*Sqrt[a + b*Tan[c + d*x]])/d`

3.320.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
 Q[u, x]`
- rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
 (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
 [(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
 , x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
 0] && GtQ[m, 0]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) +
 (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
 c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
 b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
 (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
 1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
 *(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
 - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

3.320.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 811 vs. 2(102) = 204.

Time = 0.09 (sec) , antiderivative size = 812, normalized size of antiderivative = 6.66

method	result
parts	$\frac{\ln\left(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}\right) A \sqrt{2\sqrt{a^2+b^2}+2a}}{4db} - \frac{\ln\left(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}\right) A \sqrt{2\sqrt{a^2+b^2}+2a}}{4db}$
derivativedivides	$\frac{2B\sqrt{a+b \tan(dx+c)}}{d} - \frac{\ln\left(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}\right) A \sqrt{2\sqrt{a^2+b^2}+2a}}{4db} \sqrt{a^2+b^2}$
default	$\frac{2B\sqrt{a+b \tan(dx+c)}}{d} - \frac{\ln\left(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}\right) A \sqrt{2\sqrt{a^2+b^2}+2a}}{4db} \sqrt{a^2+b^2}$

input `int((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 1/4/d/b*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2) \\ & + (a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-1/4/d/b*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2) \\ & *(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2)) \\ & *A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)+1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2) \\ & * \arctan\left(\frac{2*(a+b*\tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2)}{2*(a^2+b^2)^(1/2)-2*a} \right) \\ & / (2*(a^2+b^2)^(1/2)-2*a)^(1/2) * A-1/4/d/b*\ln((a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2) \\ & - b*\tan(d*x+c)-a-(a^2+b^2)^(1/2)) * A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2) * a+1/4/d/b*\ln((a+b*\tan(d*x+c))^(1/2) \\ & *(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*\tan(d*x+c)-a-(a^2+b^2)^(1/2)) * A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2) * \\ & (a^2+b^2)^(1/2)-1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2) * \arctan\left(\frac{2*(a^2+b^2)^(1/2)+2*a}{2*(a+b*\tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2)} \right) \\ & / (2*(a^2+b^2)^(1/2)-2*a)^(1/2) * A+B/d*(2*(a+b*\tan(d*x+c))^(1/2)-1/4*(2*(a^2+b^2)^(1/2)+2*a)^(1/2) * \ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2) \\ & *(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))+(a-(a^2+b^2)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2) * \arctan\left(\frac{2*(a+b*\tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2)}{2*(a^2+b^2)^(1/2)-2*a} \right) \\ & / (2*(a^2+b^2)^(1/2)-2*a)^(1/2) \\ & +1/4*(2*(a^2+b^2)^(1/2)+2*a)^(1/2) * \ln((a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*\tan(d*x+c)-a-(a^2+b^2)^(1/2)) \\ & +((a^2+b^2)^(1/2)-a)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2) * \arctan\left(\frac{2*(a^2+b^2)^(1/2)+2*a}{2*(a+b*\tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2)} \right) \\ & / (2*(a^2+b^2)^(1/2)-2*a)^(1/2) \end{aligned}$$

3.320. $\int \sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx$

3.320.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1199 vs. $2(97) = 194$.

Time = 0.33 (sec) , antiderivative size = 1199, normalized size of antiderivative = 9.83

$$\int \sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output

```
-1/2*(d*sqrt((2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3))*a*b +
(A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2*(A^3*B +
A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) + (A*d^3*sqrt(-(4*A^2*B
^2*a^2 + 4*(A^3*B - A*B^3))*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (2*A*
B^2*a + (A^2*B - B^3)*b)*d)*sqrt((2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(
A^3*B - A*B^3))*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^
2)) - d*sqrt((2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3))*a*b +
(A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2*(A^3*B +
A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) - (A*d^3*sqrt(-(4*A^2*B
^2*a^2 + 4*(A^3*B - A*B^3))*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (2*A*
B^2*a + (A^2*B - B^3)*b)*d)*sqrt((2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(
A^3*B - A*B^3))*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^
2)) - d*sqrt((2*A*B*b - d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3))*a*b +
(A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2*(A^3*B +
A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) + (A*d^3*sqrt(-(4*A^2*B
^2*a^2 + 4*(A^3*B - A*B^3))*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) + (2*A*
B^2*a + (A^2*B - B^3)*b)*d)*sqrt((2*A*B*b - d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(
A^3*B - A*B^3))*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^
2)) + d*sqrt((2*A*B*b - d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3))*a*b +
(A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2*(A^3*B...
```

3.320.6 Sympy [F]

$$\int \sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x)), x)`

3.320.7 Maxima [F(-2)]

Exception generated.

$$\int \sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is`

3.320.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.320.9 Mupad [B] (verification not implemented)

Time = 10.81 (sec) , antiderivative size = 845, normalized size of antiderivative = 6.93

$$\begin{aligned}
& \int \sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx \\
&= 2 \operatorname{atanh} \left(\frac{32 B^2 b^4 \sqrt{\frac{B^2 a}{4 d^2} - \frac{\sqrt{-B^4 b^2 d^4}}{4 d^4}} \sqrt{a + b \tan(c + dx)}}{\frac{16 B b^4 \sqrt{-B^4 b^2 d^4}}{d^3} + \frac{16 B a^2 b^2 \sqrt{-B^4 b^2 d^4}}{d^3}} \right) \\
&\quad - \frac{32 a b^2 \sqrt{\frac{B^2 a}{4 d^2} - \frac{\sqrt{-B^4 b^2 d^4}}{4 d^4}} \sqrt{a + b \tan(c + dx)} \sqrt{-B^4 b^2 d^4}}{\frac{16 B b^4 \sqrt{-B^4 b^2 d^4}}{d} + \frac{16 B a^2 b^2 \sqrt{-B^4 b^2 d^4}}{d}} \sqrt{\frac{\sqrt{-B^4 b^2 d^4} - B^2 a d^2}{4 d^4}} \\
&\quad - \operatorname{atanh} \left(\frac{d^3 \left(\frac{16 (A^2 b^4 - A^2 a^2 b^2) \sqrt{a + b \tan(c + dx)}}{d^2} - \frac{16 a b^2 (\sqrt{-A^4 b^2 d^4} - A^2 a d^2) \sqrt{a + b \tan(c + dx)}}{d^4} \right) \sqrt{\frac{\sqrt{-A^4 b^2 d^4} - A^2 a d^2}{d^4}}}{16 (A^3 a^2 b^3 + A^3 b^5)} \right) \\
&\quad - 2 \operatorname{atanh} \left(\frac{32 B^2 b^4 \sqrt{\frac{\sqrt{-B^4 b^2 d^4}}{4 d^4} + \frac{B^2 a}{4 d^2}} \sqrt{a + b \tan(c + dx)}}{\frac{16 B b^4 \sqrt{-B^4 b^2 d^4}}{d^3} + \frac{16 B a^2 b^2 \sqrt{-B^4 b^2 d^4}}{d^3}} \right) \\
&\quad + \frac{32 a b^2 \sqrt{\frac{\sqrt{-B^4 b^2 d^4}}{4 d^4} + \frac{B^2 a}{4 d^2}} \sqrt{a + b \tan(c + dx)} \sqrt{-B^4 b^2 d^4}}{\frac{16 B b^4 \sqrt{-B^4 b^2 d^4}}{d} + \frac{16 B a^2 b^2 \sqrt{-B^4 b^2 d^4}}{d}} \sqrt{\frac{\sqrt{-B^4 b^2 d^4} + B^2 a d^2}{4 d^4}} \\
&\quad - \operatorname{atanh} \left(\frac{d^3 \left(\frac{16 (A^2 b^4 - A^2 a^2 b^2) \sqrt{a + b \tan(c + dx)}}{d^2} + \frac{16 a b^2 (\sqrt{-A^4 b^2 d^4} + A^2 a d^2) \sqrt{a + b \tan(c + dx)}}{d^4} \right) \sqrt{\frac{\sqrt{-A^4 b^2 d^4} + A^2 a d^2}{d^4}}}{16 (A^3 a^2 b^3 + A^3 b^5)} \right) \\
&\quad + \frac{2 B \sqrt{a + b \tan(c + dx)}}{d}
\end{aligned}$$

input `int((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`

output

$$\begin{aligned}
& 2*\operatorname{atanh}\left(\frac{32*B^2*b^4*((B^2*a)/(4*d^2) - (-B^4*b^2*d^4)^{(1/2)}/(4*d^4))^{(1/2)}}{(16*B*b^4*(-B^4*b^2*d^4)^{(1/2)})/d^3 + (16*B*a^2*b^2*(-B^4*b^2*d^4)^{(1/2)})/d^3} - \frac{32*a*b^2*((B^2*a)/(4*d^2) - (-B^4*b^2*d^4)^{(1/2)}/(4*d^4))^{(1/2)}}{(16*B*b^4*(-B^4*b^2*d^4)^{(1/2)})/d + (16*B*a^2*b^2*(-B^4*b^2*d^4)^{(1/2)})/d}\right) \\
& * \frac{(a + b*\tan(c + d*x))^{(1/2)}}{(16*B*b^4*(-B^4*b^2*d^4)^{(1/2)})/d^3 + (16*B*a^2*b^2*(-B^4*b^2*d^4)^{(1/2)})/d^3} - \frac{32*a*b^2*((B^2*a)/(4*d^2) - (-B^4*b^2*d^4)^{(1/2)}/(4*d^4))^{(1/2)}}{(16*B*b^4*(-B^4*b^2*d^4)^{(1/2)})/d + (16*B*a^2*b^2*(-B^4*b^2*d^4)^{(1/2)})/d} \\
& * \frac{(a + b*\tan(c + d*x))^{(1/2)}}{(16*B*b^4*(-B^4*b^2*d^4)^{(1/2)})/d + (16*B*a^2*b^2*(-B^4*b^2*d^4)^{(1/2)})/d} * \frac{(-B^4*b^2*d^4)^{(1/2)}}{(16*B*b^4*(-B^4*b^2*d^4)^{(1/2)})/d + (16*B*a^2*b^2*(-B^4*b^2*d^4)^{(1/2)})/d} \\
& * \left(\frac{(-B^4*b^2*d^4)^{(1/2)} - B^2*a*d^2}{(4*d^4)} \right)^{(1/2)} - \operatorname{atanh}\left(\frac{d^3*((16*(A^2*b^4 - A^2*a^2*b^2)*(a + b*\tan(c + d*x))^{(1/2)})/d^2 - (16*a*b^2*((-A^4*b^2*d^4)^{(1/2)} - A^2*a*d^2)*(a + b*\tan(c + d*x))^{(1/2)})/d^4)}{(16*(A^3*b^5 + A^3*a^2*b^3))}\right) \\
& * \left(\frac{(-A^4*b^2*d^4)^{(1/2)} - A^2*a*d^2}{d^4} \right)^{(1/2)} / \left(\frac{(-A^4*b^2*d^4)^{(1/2)} - A^2*a*d^2}{d^4} \right)^{(1/2)} / (16*(A^3*b^5 + A^3*a^2*b^3)) * \left(\frac{(-A^4*b^2*d^4)^{(1/2)} - A^2*a*d^2}{d^4} \right)^{(1/2)} \\
& - 2*\operatorname{atanh}\left(\frac{32*B^2*b^4*((-B^4*b^2*d^4)^{(1/2)}/(4*d^4) + (B^2*a)/(4*d^2))^{(1/2)}}{(16*B*b^4*(-B^4*b^2*d^4)^{(1/2)})/d^3 + (16*B*a^2*b^2*(-B^4*b^2*d^4)^{(1/2)})/d^3} + \frac{32*a*b^2*((-B^4*b^2*d^4)^{(1/2)}/(4*d^4) + (B^2*a)/(4*d^2))^{(1/2)}}{(16*B*b^4*(-B^4*b^2*d^4)^{(1/2)})/d + (16*B*a^2*b^2*(-B^4*b^2*d^4)^{(1/2)})/d}\right) \\
& * \frac{(a + b*\tan(c + d*x))^{(1/2)}}{(16*B*b^4*(-B^4*b^2*d^4)^{(1/2)})/d + (16*B*a^2*b^2*(-B^4*b^2*d^4)^{(1/2)})/d} * \frac{(-B^4*b^2*d^4)^{(1/2)}}{(16*B*b^4*(-B^4*b^2*d^4)^{(1/2)})/d + (16*B*a^2*b^2*(-B^4*b^2*d^4)^{(1/2)})/d} \\
& * \left(\frac{(-B^4*b^2*d^4)^{(1/2)} + B^2*a*d^2}{(4*d^4)} \right)^{(1/2)} - \operatorname{atanh}\left(\frac{d^3*((16*(A^2*b^4 - A^2*a^2*b^2)*(a + b*\tan(c + d*x))^{(1/2)})/d^2 + (16*a*b^2*((-A^4*b^2*d^4)^{(1/2)} + A^2*a*d^2)*(a + b*\tan(c + d*x))^{(1/2)})/d^4)}{(16*(A^3*b^5 + A^3*a^2*b^3))}\right) \\
& * \left(\frac{(-A^4*b^2*d^4)^{(1/2)} + A^2*a*d^2}{d^4} \right)^{(1/2)} / \left(\frac{(-A^4*b^2*d^4)^{(1/2)} + A^2*a*d^2}{d^4} \right)^{(1/2)} / (16*(A^3*b^5 + A^3*a^2*b^3)) * \left(\frac{(-A^4*b^2*d^4)^{(1/2)} + A^2*a*d^2}{d^4} \right)^{(1/2)} \\
& + (2*B*(a + b*\tan(c + d*x))^{(1/2)})/d
\end{aligned}$$

3.321 $\int \cot(c+dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.321.1 Optimal result	3035
3.321.2 Mathematica [A] (verified)	3035
3.321.3 Rubi [A] (warning: unable to verify)	3036
3.321.4 Maple [B] (verified)	3039
3.321.5 Fricas [B] (verification not implemented)	3040
3.321.6 Sympy [F]	3041
3.321.7 Maxima [F]	3042
3.321.8 Giac [F(-1)]	3042
3.321.9 Mupad [B] (verification not implemented)	3042

3.321.1 Optimal result

Integrand size = 31, antiderivative size = 131

$$\int \cot(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= -\frac{2\sqrt{a}A \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{d} + \frac{\sqrt{a - ib}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d}$$

$$+ \frac{\sqrt{a + ib}(A + iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d}$$

```
output -2*A*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))*a^(1/2)/d+(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))*(a-I*b)^(1/2)/d+(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))*(a+I*b)^(1/2)/d
```

3.321.2 Mathematica [A] (verified)

Time = 0.66 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.67

$$\int \cot(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx =$$

$$-\frac{2\sqrt{a}A \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right) - \frac{(A(b^2+a\sqrt{-b^2})+b(a-\sqrt{-b^2})B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-\sqrt{-b^2}}}\right)}{\sqrt{-b^2}\sqrt{a-\sqrt{-b^2}}} + \frac{(A(b^2-a\sqrt{-b^2})+b(a+\sqrt{-b^2})B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+\sqrt{-b^2}}}\right)}{\sqrt{-b^2}\sqrt{a+\sqrt{-b^2}}}}{d}$$

input `Integrate[Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output $-\left(\frac{2\sqrt{a}A\operatorname{ArcTanh}\left[\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a}}\right] - \left(\frac{A(b^2 + a\sqrt{-b^2}) + b(a - \sqrt{-b^2})B}{\sqrt{a - \sqrt{-b^2}}}\right)\operatorname{ArcTanh}\left[\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a - \sqrt{-b^2}}}\right]}{\sqrt{-b^2}\sqrt{a - \sqrt{-b^2}}}\right) + \left(\frac{A(b^2 - a\sqrt{-b^2}) + b(a + \sqrt{-b^2})B}{\sqrt{a + \sqrt{-b^2}}}\right)\operatorname{ArcTanh}\left[\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a + \sqrt{-b^2}}}\right]}{\sqrt{-b^2}\sqrt{a + \sqrt{-b^2}}}\right)/d$

3.321.3 Rubi [A] (warning: unable to verify)

Time = 0.99 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.90, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 4095, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\tan(c+dx)}dx \\ & \quad \downarrow \text{4095} \\ & \int \frac{Ab+aB-(aA-bB)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}}dx + aA \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b\tan(c+dx)}}dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{Ab+aB-(aA-bB)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}}dx + aA \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}}dx \\ & \quad \downarrow \text{4022} \\ & -\frac{1}{2}(a+ib)(-B+iA) \int \frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}}dx + \frac{1}{2}(a-ib)(B+iA) \int \frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}}dx + \\ & \quad aA \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}}dx \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{2}(a+ib)(-B+iA) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}(a-ib)(B+iA) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \\
& \quad aA \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx \\
& \quad \downarrow 4020 \\
& \frac{i(a-ib)(B+iA) \int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \\
& \frac{i(a+ib)(-B+iA) \int -\frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \\
& \quad aA \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx \\
& \quad \downarrow 25 \\
& \frac{i(a-ib)(B+iA) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} - \\
& \frac{i(a+ib)(-B+iA) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \\
& \quad aA \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx \\
& \quad \downarrow 73 \\
& \frac{(a+ib)(-B+iA) \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \\
& \frac{(a-ib)(B+iA) \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + aA \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx \\
& \quad \downarrow 221 \\
& aA \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + \frac{\sqrt{a-ib}(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} - \\
& \quad \frac{\sqrt{a+ib}(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} \\
& \quad \downarrow 4117 \\
& \frac{aA \int \frac{\cot(c+dx)}{\sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{d} + \frac{\sqrt{a-ib}(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} - \\
& \quad \frac{\sqrt{a+ib}(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} \\
& \quad \downarrow 73
\end{aligned}$$

$$\frac{2aA \int \frac{1}{\frac{a+b \tan(c+dx)}{b} - \frac{a}{b}} d\sqrt{a+b \tan(c+dx)}}{bd} + \frac{\sqrt{a-ib}(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} - \frac{\sqrt{a+ib}(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d}$$

↓ 221

$$\frac{\sqrt{a-ib}(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} - \frac{\sqrt{a+ib}(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} - \frac{2\sqrt{a}A \operatorname{Arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{d}$$

input `Int[Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[a - I*b]*(I*A + B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d - (Sqrt[a + I*b]*(I*A - B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d - (2*Sqrt[a]*A*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/d`

3.321.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4095 `Int((((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*Sqrt[(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])]/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[Simp[A*(a*c + b*d) + B*(b*c - a*d) - (A*(b*c - a*d) - B*(a*c + b*d))*Tan[e + f*x], x]/Sqrt[c + d*Tan[e + f*x]], x] - Simp[(b*c - a*d)*((B*a - A*b)/(a^2 + b^2)) Int[(1 + Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*Sqrt[c + d*Tan[e + f*x])], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

3.321.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 975 vs. 2(107) = 214.

Time = 0.24 (sec) , antiderivative size = 976, normalized size of antiderivative = 7.45

method	result
derivativedivides	$\frac{\ln\left(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}\right) A \sqrt{2\sqrt{a^2+b^2}+2a}}{4d} - \frac{\ln\left(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}\right) A \sqrt{2\sqrt{a^2+b^2}+2a}}{4d}$
default	$\frac{\ln\left(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}\right) A \sqrt{2\sqrt{a^2+b^2}+2a}}{4d} - \frac{\ln\left(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}\right) A \sqrt{2\sqrt{a^2+b^2}+2a}}{4d}$

3.321. $\int \cot(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

input `int(cot(d*x+c)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/4/d*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/4/d/b*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*B*(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/4/d/b*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*(a^2+b^2)^(1/2)-1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a+1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B-1/4/d*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/4/d/b*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*B*(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/4/d/b*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*(a^2+b^2)^(1/2)+1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-...`

3.321.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1218 vs. 2(101) = 202.

Time = 0.53 (sec) , antiderivative size = 2452, normalized size of antiderivative = 18.72

$$\int \cot(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output

```

[-1/2*(d*sqrt(-(2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b
+ (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2*(A^3*B
+ A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) + (B*d^3*sqrt(-(4*A^2
*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) + (2*
A^2*B*a + (A^3 - A*B^2)*b)*d)*sqrt(-(2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 +
4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)
/d^2)) - d*sqrt(-(2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a
*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2*(A^3*
B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) - (B*d^3*sqrt(-(4*A
^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) + (
2*A^2*B*a + (A^3 - A*B^2)*b)*d)*sqrt(-(2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2
+ 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*
a)/d^2)) - d*sqrt(-(2*A*B*b - d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)
)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2*(A^
3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) + (B*d^3*sqrt(-(4
*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) -
(2*A^2*B*a + (A^3 - A*B^2)*b)*d)*sqrt(-(2*A*B*b - d^2*sqrt(-(4*A^2*B^2*a^
2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2
)*a)/d^2)) + d*sqrt(-(2*A*B*b - d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^
3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(...

```

3.321.6 Sympy [F]

$$\int \cot(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} \cot(c + dx) dx$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))*cot(c + d*x), x)`

3.321.7 Maxima [F]

$$\int \cot(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \cot(dx + c) dx$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*cot(d*x + c), x)`

3.321.8 Giac [F(-1)]

Timed out.

$$\int \cot(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.321.9 Mupad [B] (verification not implemented)

Time = 10.09 (sec) , antiderivative size = 9785, normalized size of antiderivative = 74.69

$$\int \cot(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`

```

output (A*a^(1/2)*atan(((A*a^(1/2))*((32*(a + b*tan(c + d*x))^(1/2)*(A^4*b^12 + B^
4*b^12 + 2*A^2*B^2*b^12 + 3*A^4*a^4*b^8 + 2*B^4*a^2*b^10 + B^4*a^4*b^8 + 6
*A^2*B^2*a^2*b^10 - 8*A^3*B*a^3*b^9))/d^4 + (A*a^(1/2))*((32*(3*A^3*a^2*b^1
0*d^2 + 3*A^3*a^4*b^8*d^2 + B^3*a^3*b^9*d^2 + B^3*a*b^11*d^2 - 15*A^2*B*a*
b^11*d^2 - 9*A*B^2*a^2*b^10*d^2 - 9*A*B^2*a^4*b^8*d^2 - 15*A^2*B*a^3*b^9*d
^2))/d^5 + (A*a^(1/2))*((32*(a + b*tan(c + d*x))^(1/2)*(10*B^2*a^3*b^8*d^2
- 18*A^2*a^3*b^8*d^2 + 16*A*B*b^11*d^2 - 6*A^2*a*b^10*d^2 + 6*B^2*a*b^10*d
^2 + 24*A*B*a^2*b^9*d^2))/d^4 - (A*a^(1/2))*((32*(12*A*a*b^10*d^4 + 12*A*a^
3*b^8*d^4))/d^5 - (32*A*a^(1/2)*(16*b^10*d^4 + 24*a^2*b^8*d^4)*(a + b*tan(
c + d*x))^(1/2))/d^5))/d))/d)*i)/d + (A*a^(1/2))*((32*(a + b*tan(c + d
*x))^(1/2)*(A^4*b^12 + B^4*b^12 + 2*A^2*B^2*b^12 + 3*A^4*a^4*b^8 + 2*B^4*a
^2*b^10 + B^4*a^4*b^8 + 6*A^2*B^2*a^2*b^10 - 8*A^3*B*a^3*b^9))/d^4 - (A*a^
(1/2))*((32*(3*A^3*a^2*b^10*d^2 + 3*A^3*a^4*b^8*d^2 + B^3*a^3*b^9*d^2 + B^3
*a*b^11*d^2 - 15*A^2*B*a*b^11*d^2 - 9*A*B^2*a^2*b^10*d^2 - 9*A*B^2*a^4*b^8
*d^2 - 15*A^2*B*a^3*b^9*d^2))/d^5 - (A*a^(1/2))*((32*(a + b*tan(c + d*x))^(
1/2)*(10*B^2*a^3*b^8*d^2 - 18*A^2*a^3*b^8*d^2 + 16*A*B*b^11*d^2 - 6*A^2*a*
b^10*d^2 + 6*B^2*a*b^10*d^2 + 24*A*B*a^2*b^9*d^2))/d^4 + (A*a^(1/2))*((32*(
12*A*a*b^10*d^4 + 12*A*a^3*b^8*d^4))/d^5 + (32*A*a^(1/2)*(16*b^10*d^4 + 24
*a^2*b^8*d^4)*(a + b*tan(c + d*x))^(1/2))/d^5))/d))/d)*i)/d)/((64*(A^
5*a*b^12 + A^5*a^3*b^10 + A^2*B^3*a^2*b^11 + A^2*B^3*a^4*b^9 + 3*A^3*B^...

```

3.322 $\int \cot^2(c+dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.322.1 Optimal result	3044
3.322.2 Mathematica [A] (verified)	3045
3.322.3 Rubi [A] (warning: unable to verify)	3045
3.322.4 Maple [B] (verified)	3050
3.322.5 Fricas [B] (verification not implemented)	3051
3.322.6 Sympy [F]	3052
3.322.7 Maxima [F]	3053
3.322.8 Giac [F(-1)]	3053
3.322.9 Mupad [B] (verification not implemented)	3053

3.322.1 Optimal result

Integrand size = 33, antiderivative size = 167

$$\int \cot^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= -\frac{(Ab + 2aB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ad}} + \frac{\sqrt{a - ib}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d}$$

$$- \frac{\sqrt{a + ib}(iA - B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d} - \frac{A \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{d}$$

output

```
-(A*b+2*B*a)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/d/a^(1/2)+(I*A+B)*arc
tanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))*(a-I*b)^(1/2)/d-(I*A-B)*arctanh
((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))*(a+I*b)^(1/2)/d-A*cot(d*x+c)*(a+b*t
an(d*x+c))^(1/2)/d
```

3.322.2 Mathematica [A] (verified)

Time = 2.65 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.41

$$\int \cot^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{(Ab+2aB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{a}} + \frac{(A(b^2+a\sqrt{-b^2})+b(a-\sqrt{-b^2})B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-\sqrt{-b^2}}}\right)}{\sqrt{a-\sqrt{-b^2}}} + \frac{(A(b^2-a\sqrt{-b^2})+b(a+\sqrt{-b^2})B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+\sqrt{-b^2}}}\right)}{\sqrt{a+\sqrt{-b^2}}}$$

```
input Integrate[Cot[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]
```

```
output (-((A*b + 2*a*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]]/Sqrt[a]) + ((A*(b^2 + a*Sqrt[-b^2]) + b*(a - Sqrt[-b^2]))*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - Sqrt[-b^2]]]/Sqrt[a - Sqrt[-b^2]] + ((A*(b^2 - a*Sqrt[-b^2]) + b*(a + Sqrt[-b^2]))*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + Sqrt[-b^2]]]/Sqrt[a + Sqrt[-b^2]] - A*b*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/b)/d
```

3.322.3 Rubi [A] (warning: unable to verify)

Time = 1.23 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.93, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 4091, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx))}{\tan(c + dx)^2} dx$$

$$\downarrow \text{4091}$$

$$- \int \frac{\cot(c + dx) (-Ab \tan^2(c + dx) - 2(aA - bB) \tan(c + dx) + Ab + 2aB)}{2\sqrt{a + b \tan(c + dx)}} dx - \frac{A \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{d}$$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{1}{2} \int \frac{\cot(c+dx) (-Ab \tan^2(c+dx) - 2(aA - bB) \tan(c+dx) + Ab + 2aB)}{\sqrt{a+b \tan(c+dx)}} dx - \\
& \quad \frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} \\
& \downarrow 3042 \\
& \frac{1}{2} \int \frac{-Ab \tan(c+dx)^2 - 2(aA - bB) \tan(c+dx) + Ab + 2aB}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx - \\
& \quad \frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} \\
& \downarrow 4136 \\
& \frac{1}{2} \left(\int \frac{2(aA - bB + (Ab + aB) \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx + (2aB + Ab) \int \frac{\cot(c+dx) (\tan^2(c+dx) + 1)}{\sqrt{a+b \tan(c+dx)}} dx \right) - \\
& \quad \frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} \\
& \downarrow 27 \\
& \frac{1}{2} \left((2aB + Ab) \int \frac{\cot(c+dx) (\tan^2(c+dx) + 1)}{\sqrt{a+b \tan(c+dx)}} dx - 2 \int \frac{aA - bB + (Ab + aB) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx \right) - \\
& \quad \frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} \\
& \downarrow 3042 \\
& \frac{1}{2} \left((2aB + Ab) \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx - 2 \int \frac{aA - bB + (Ab + aB) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx \right) - \\
& \quad \frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} \\
& \downarrow 4022 \\
& - \frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \\
& \frac{1}{2} \left((2aB + Ab) \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{1}{2} (a+ib)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2} (a-i) \right) \right) \\
& \downarrow 3042
\end{aligned}$$

$$\begin{aligned}
& -\frac{A \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left((2aB+Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{1}{2}(a+ib)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}(a-ib)(A-iB) \int \frac{1+i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx \right) \right) \\
& \quad \downarrow 4020 \\
& -\frac{A \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left((2aB+Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{i(a-ib)(A-iB) \int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \frac{i(a+ib)(A+iB) \int \frac{1}{(1+i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} \right) \right) \\
& \quad \downarrow 25 \\
& -\frac{A \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left((2aB+Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{i(a+ib)(A+iB) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \frac{i(a-ib)(A-iB) \int \frac{1}{(i \tan(c+dx)-1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} \right) \right) \\
& \quad \downarrow 73 \\
& -\frac{A \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left((2aB+Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{(a+ib)(A+iB) \int \frac{1}{-i \frac{\tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \frac{(a-ib)(A-iB) \int \frac{1}{i \frac{\tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} \right) \right) \\
& \quad \downarrow 221 \\
& -\frac{A \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left((2aB+Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{\sqrt{a-ib}(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{\sqrt{a+ib}(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} \right) \right) \\
& \quad \downarrow 4117 \\
& -\frac{A \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left(\frac{(2aB+Ab) \int \frac{\cot(c+dx)}{\sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{d} - 2 \left(\frac{\sqrt{a-ib}(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{\sqrt{a+ib}(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} \right) \right) \\
& \quad \downarrow 73
\end{aligned}$$

$$\begin{aligned}
 & -\frac{A \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{d} + \\
 & \frac{1}{2} \left(\frac{2(2aB+Ab) \int \frac{1}{\frac{a+b \tan(c+dx)}{b} - \frac{a}{b}} d\sqrt{a+b \tan(c+dx)}}{bd} - 2 \left(\frac{\sqrt{a-ib}(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{\sqrt{a+ib}(A+iB)}{d} \right) \right) \\
 & \quad \downarrow \text{221} \\
 & -\frac{A \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{d} + \\
 & \frac{1}{2} \left(-\frac{2(2aB+Ab) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ad}} - 2 \left(\frac{\sqrt{a-ib}(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{\sqrt{a+ib}(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} \right) \right)
 \end{aligned}$$

input `Int[Cot[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(-2*((Sqrt[a - I*b]*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d + (Sqrt[a + I*b]*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d) - (2*(A*b + 2*a*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/(Sqrt[a]*d))/2 - (A*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/d`

3.322.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4091 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`


```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.322.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1028 vs. $2(141) = 282$.

Time = 0.22 (sec) , antiderivative size = 1029, normalized size of antiderivative = 6.16

method	result	size
derivativedivides	Expression too large to display	1029
default	Expression too large to display	1029

```
input int(cot(d*x+c)^2*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

output

```

1/4/d/b*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(
1/2)+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)-1/4/
d/b*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)
+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+1/4/d*ln(b*tan(d*x+c)+
a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*B*
(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*
(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2
*a)^(1/2))*B*(a^2+b^2)^(1/2)-1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2
*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-
2*a)^(1/2))*A-1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))
^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*a-1
/4/d/b*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)
)-a-(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)+1/4/d
/b*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-
(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-1/4/d*ln((a+b*tan(d*x+c)
))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*B*(
2*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*
(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*
a)^(1/2))*B*(a^2+b^2)^(1/2)+1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2
*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)...

```

3.322.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1282 vs. $2(135) = 270$.

Time = 0.88 (sec) , antiderivative size = 2579, normalized size of antiderivative = 15.44

$$\int \cot^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input

```

integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm
m="fricas")

```

output

```
[1/2*(a*d*sqrt((2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b
+ (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2*(A^3*B
+ A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) + (A*d^3*sqrt(-(4*A^2
*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (2*
A*B^2*a + (A^2*B - B^3)*b)*d)*sqrt((2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4
*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/
d^2))*tan(d*x + c) - a*d*sqrt((2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3
*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*
log(-(2*(A^3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) - (A*d
^3*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*
b^2)/d^4) - (2*A*B^2*a + (A^2*B - B^3)*b)*d)*sqrt((2*A*B*b + d^2*sqrt(-(4*
A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) -
(A^2 - B^2)*a)/d^2))*tan(d*x + c) - a*d*sqrt((2*A*B*b - d^2*sqrt(-(4*A^2*B
^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2
- B^2)*a)/d^2)*log(-(2*(A^3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x +
c) + a) + (A*d^3*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*
A^2*B^2 + B^4)*b^2)/d^4) + (2*A*B^2*a + (A^2*B - B^3)*b)*d)*sqrt((2*A*B*b
- d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^
4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2))*tan(d*x + c) + a*d*sqrt((2*A*B*b - d^2
*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)...
```

3.322.6 Sympy [F]

$$\int \cot^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} \cot^2(c + dx) dx$$

input `integrate(cot(d*x+c)**2*(a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))*cot(c + d*x)**2, x)`

3.322.7 Maxima [F]

$$\int \cot^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \cot(dx + c)^2 dx$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*cot(d*x + c)^2, x)`

3.322.8 Giac [F(-1)]

Timed out.

$$\int \cot^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.322.9 Mupad [B] (verification not implemented)

Time = 8.83 (sec) , antiderivative size = 10987, normalized size of antiderivative = 65.79

$$\int \cot^2(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`

output $(\operatorname{atan}(\frac{((16(a + b \tan(c + dx))^{1/2}(3A^4b^{12} + 2B^4b^{12} + 3A^2B^2b^{12} + 3A^4a^2b^{10} + 2A^4a^4b^8 + 6B^4a^4b^8 + 29A^2B^2a^2b^{10} - 4AB^3ab^{11} + 8A^3Bab^{11} + 20AB^3a^3b^9 - 4A^3B^3a^3b^9))}{d^4} + ((Ab + 2Ba)((8(12B^3a^2b^{10}d^2 - 20A^3a^3b^9d^2 + 12B^3a^4b^8d^2 + 28A^2Bb^{12}d^2 - 20A^3aab^{11}d^2 + 60AB^2aab^{11}d^2 + 60AB^2a^3b^9d^2 - 8A^2B^2a^2b^{10}d^2 - 36A^2B^2a^4b^8d^2))}{d^5} - (((16(a + b \tan(c + dx))^{1/2}(36B^2a^3b^8d^2 - 20A^2a^3b^8d^2 + 32ABb^{11}d^2 - 8A^2aab^{10}d^2 + 12B^2aab^{10}d^2 + 64AB^2a^2b^9d^2))}{d^4} + ((Ab + 2Ba)((8(32A^2b^{11}d^4 + 48B^2a^3b^8d^4 + 32A^2a^2b^9d^4 + 48B^2a^3b^8d^4))}{d^5} - (8(Ab + 2Ba)(32b^{10}d^4 + 48a^2b^8d^4)(a + b \tan(c + dx))^{1/2})/(a^{1/2}d^5)))/(2a^{1/2}d)))(Ab + 2Ba))/(2a^{1/2}d)))/(2a^{1/2}d))(Ab + 2Ba)1i)/(2a^{1/2}d) + (((16(a + b \tan(c + dx))^{1/2}(3A^4b^{12} + 2B^4b^{12} + 3A^2B^2b^{12} + 3A^4a^2b^{10} + 2A^4a^4b^8 + 6B^4a^4b^8 + 29A^2B^2a^2b^{10} - 4AB^3ab^{11} + 8A^3Bab^{11} + 20AB^3a^3b^9 - 4A^3B^3a^3b^9))}{d^4} - ((Ab + 2Ba)((8(12B^3a^2b^{10}d^2 - 20A^3a^3b^9d^2 + 12B^3a^4b^8d^2 + 28A^2Bb^{12}d^2 - 20A^3aab^{11}d^2 + 60AB^2aab^{11}d^2 + 60AB^2a^3b^9d^2 - 8A^2B^2a^2b^{10}d^2 - 36A^2B^2a^4b^8d^2))}{d^5} + (((16(a + b \tan(c + dx))^{1/2}(36B^2a^3b^8d^2 - 20A^2a^3b^8d^2 + 32ABb^{11}d^2 - 8A^2aab^{10}d^2 + 12B^2aab^{10}d^2 + \dots$

3.323 $\int \cot^3(c+dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.323.1 Optimal result	3055
3.323.2 Mathematica [A] (verified)	3056
3.323.3 Rubi [A] (warning: unable to verify)	3056
3.323.4 Maple [B] (verified)	3062
3.323.5 Fricas [B] (verification not implemented)	3063
3.323.6 Sympy [F]	3064
3.323.7 Maxima [F]	3065
3.323.8 Giac [F(-1)]	3065
3.323.9 Mupad [B] (verification not implemented)	3065

3.323.1 Optimal result

Integrand size = 33, antiderivative size = 219

$$\int \cot^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{(8a^2 A + Ab^2 - 4abB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{4a^{3/2}d}$$

$$- \frac{\sqrt{a - ib}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d} - \frac{\sqrt{a + ib}(A + iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d}$$

$$- \frac{(Ab + 4aB) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{4ad} - \frac{A \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d}$$

output

```
1/4*(8*A*a^2+A*b^2-4*B*a*b)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/a^(3/2)
)/d-(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))*(a-I*b)^(1/2)/d-
(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))*(a+I*b)^(1/2)/d-1/4*
(A*b+4*B*a)*cot(d*x+c)*(a+b*tan(d*x+c))^(1/2)/a/d-1/2*A*cot(d*x+c)^2*(a+b*
tan(d*x+c))^(1/2)/d
```

3.323.2 Mathematica [A] (verified)

Time = 4.92 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.24

$$\int \cot^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{(8a^2A + Ab^2 - 4abB) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a}}\right)}{a^{3/2}} + \frac{4(-aAb + Ab\sqrt{-b^2 + b^2B + a\sqrt{-b^2}B}) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - \sqrt{-b^2}}}\right)}{\sqrt{a - \sqrt{-b^2}}} - \frac{4(aAb + Ab\sqrt{-b^2} - b^2B + a)}{4d}$$

input `Integrate[Cot[c + d*x]^3*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`output `((8*a^2*A + A*b^2 - 4*a*b*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/a^(3/2) + ((4*(-(a*A*b) + A*b*Sqrt[-b^2] + b^2*B + a*Sqrt[-b^2])*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - Sqrt[-b^2]]])/Sqrt[a - Sqrt[-b^2]] - (4*(a*A*b + A*b*Sqrt[-b^2] - b^2*B + a*Sqrt[-b^2])*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + Sqrt[-b^2]]])/Sqrt[a + Sqrt[-b^2]] - (b*Cot[c + d*x]*(A*b + 4*a*B + 2*a*A*Cot[c + d*x])*Sqrt[a + b*Tan[c + d*x]])/a)/b)/(4*d)`**3.323.3 Rubi [A] (warning: unable to verify)**Time = 1.75 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.00, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 4091, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx))}{\tan(c + dx)^3} dx$$

$$\downarrow \text{4091}$$

$$\begin{aligned}
& -\frac{1}{2} \int -\frac{\cot^2(c+dx) (-3Ab \tan^2(c+dx) - 4(aA - bB) \tan(c+dx) + Ab + 4aB)}{2\sqrt{a+b \tan(c+dx)}} dx - \\
& \quad \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} \\
& \quad \downarrow 27 \\
& \frac{1}{4} \int \frac{\cot^2(c+dx) (-3Ab \tan^2(c+dx) - 4(aA - bB) \tan(c+dx) + Ab + 4aB)}{\sqrt{a+b \tan(c+dx)}} dx - \\
& \quad \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} \\
& \quad \downarrow 3042 \\
& \frac{1}{4} \int \frac{-3Ab \tan(c+dx)^2 - 4(aA - bB) \tan(c+dx) + Ab + 4aB}{\tan(c+dx)^2 \sqrt{a+b \tan(c+dx)}} dx - \\
& \quad \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} \\
& \quad \downarrow 4132 \\
& \frac{1}{4} \left(-\frac{\int \frac{\cot(c+dx)(8Aa^2 - 4bBa + 8(Ab+aB) \tan(c+dx)a + Ab^2 + b(Ab+4aB) \tan^2(c+dx))}{2\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{(4aB + Ab) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} \right. \\
& \quad \left. \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} \right) \\
& \quad \downarrow 27 \\
& \frac{1}{4} \left(-\frac{\int \frac{\cot(c+dx)(8Aa^2 - 4bBa + 8(Ab+aB) \tan(c+dx)a + Ab^2 + b(Ab+4aB) \tan^2(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{(4aB + Ab) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} \right. \\
& \quad \left. \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} \right) \\
& \quad \downarrow 3042 \\
& \frac{1}{4} \left(-\frac{\int \frac{8Aa^2 - 4bBa + 8(Ab+aB) \tan(c+dx)a + Ab^2 + b(Ab+4aB) \tan(c+dx)^2}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{(4aB + Ab) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} \right) \\
& \quad \left. \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} \right) \\
& \quad \downarrow 4136
\end{aligned}$$

$$\frac{1}{4} \left(-\frac{(8a^2A - 4abB + Ab^2) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx + \int \frac{8(aAb+aB)-a(aA-bB) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{(4aB + Ab) \cot(c+dx)}{ad} \right) - \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d}$$

↓ 27

$$\frac{1}{4} \left(-\frac{(8a^2A - 4abB + Ab^2) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx + 8 \int \frac{a(Ab+aB)-a(aA-bB) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{(4aB + Ab) \cot(c+dx)}{ad} \right) - \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d}$$

↓ 3042

$$\frac{1}{4} \left(-\frac{(8a^2A - 4abB + Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 8 \int \frac{a(Ab+aB)-a(aA-bB) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{(4aB + Ab) \cot(c+dx)}{ad} \right) - \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d}$$

↓ 4022

$$-\frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} + \frac{1}{4} \left(-\frac{(4aB + Ab) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} - \frac{(8a^2A - 4abB + Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 8 \left(\frac{1}{2} a(a-b) \int \frac{1}{\sqrt{a+b \tan(c+dx)}} dx \right)}{2a} \right) - \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d}$$

↓ 3042

$$-\frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} + \frac{1}{4} \left(-\frac{(4aB + Ab) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} - \frac{(8a^2A - 4abB + Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 8 \left(\frac{1}{2} a(a-b) \int \frac{1}{\sqrt{a+b \tan(c+dx)}} dx \right)}{2a} \right) - \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d}$$

↓ 4020

$$-\frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} + \frac{1}{4} \left(-\frac{(4aB + Ab) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} - \frac{(8a^2A - 4abB + Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 8 \left(\frac{1}{2} a(a-b) \int \frac{1}{\sqrt{a+b \tan(c+dx)}} dx \right)}{2a} \right) - \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d}$$

$$\begin{aligned} & \downarrow 25 \\ & -\frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2d} + \\ \frac{1}{4} \left(-\frac{(4aB+Ab) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad} - \frac{(8a^2A-4abB+Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 8 \left(-\frac{ia(a-b)}{2a} \right)}{2a} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 73 \\ & -\frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2d} + \\ \frac{1}{4} \left(-\frac{(4aB+Ab) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad} - \frac{(8a^2A-4abB+Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 8 \left(\frac{a(a-ib)}{2a} \right)}{2a} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 221 \\ & -\frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2d} + \\ \frac{1}{4} \left(-\frac{(4aB+Ab) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad} - \frac{(8a^2A-4abB+Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 8 \left(\frac{a\sqrt{a-ib}}{2a} \right)}{2a} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 4117 \\ & -\frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2d} + \\ \frac{1}{4} \left(-\frac{(4aB+Ab) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad} - \frac{(8a^2A-4abB+Ab^2) \int \frac{\cot(c+dx)}{\sqrt{a+b \tan(c+dx)}} d \tan(c+dx) + 8 \left(\frac{a\sqrt{a-ib}(B+iA) \arctan(\frac{a+b \tan(c+dx)}{b})}{d} \right)}{2a} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 73 \\ & -\frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2d} + \\ \frac{1}{4} \left(-\frac{(4aB+Ab) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad} - \frac{2(8a^2A-4abB+Ab^2) \int \frac{1}{\frac{a+b \tan(c+dx)}{b} - \frac{a}{b}} d\sqrt{a+b \tan(c+dx)}}{bd} + 8 \left(\frac{a\sqrt{a-ib}(B+iA) \arctan(\frac{a+b \tan(c+dx)}{b})}{d} \right)}{2a} \right) \end{aligned}$$

$$\frac{1}{4} \left(\frac{A \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} + \frac{(4aB + Ab) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{ad} - \frac{2(8a^2A - 4abB + Ab^2) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a}}\right)}{\sqrt{ad}} + 8 \frac{a\sqrt{a - ib}(B + iA)}{2a} \right)$$

input `Int[Cot[c + d*x]^3*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `-1/2*(A*Cot[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]])/d + (-1/2*(8*((a*Sqrt[a - I*b])*(I*A + B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d - (a*Sqrt[a + I*b]*(I*A - B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d - (2*(8*a^2*A + A*b^2 - 4*a*b*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]]/(Sqrt[a]*d))/a - ((A*b + 4*a*B)*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]]/(a*d))/4`

3.323.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.323. $\int \cot^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4091 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.323.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1144 vs. $2(185) = 370$.

Time = 0.25 (sec) , antiderivative size = 1145, normalized size of antiderivative = 5.23

method	result	size
derivativedivides	Expression too large to display	1145
default	Expression too large to display	1145

```
input int(cot(d*x+c)^3*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

output

```

-1/4/d*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1
/2)+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/4/d/b*ln(b*tan(d*x+
c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))
*B*(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/4/d/b*ln(b*tan(d*x+c)+a
+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*B*(
2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2
*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-
2*a)^(1/2))*A*(a^2+b^2)^(1/2)+1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*
(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2
*a)^(1/2))*A*a-1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+
c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B+
1/4/d*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)
-a-(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/4/d/b*ln((a+b*tan(d*
x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*
B*(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/4/d/b*ln((a+b*tan(d*x+c)
)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*B*(2
*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2
*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2
*a)^(1/2))*A*(a^2+b^2)^(1/2)-1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*
(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)...

```

3.323.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1338 vs. $2(179) = 358$.

Time = 3.17 (sec) , antiderivative size = 2691, normalized size of antiderivative = 12.29

$$\int \cot^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input

```

integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm
m="fricas")

```

```

output [1/8*(4*a^2*d*sqrt(-(2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)
)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2*(A
^3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) + (B*d^3*sqrt(-(
4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4)
+ (2*A^2*B*a + (A^3 - A*B^2)*b)*d)*sqrt(-(2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a
^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B
^2)*a)/d^2))*tan(d*x + c)^2 - 4*a^2*d*sqrt(-(2*A*B*b + d^2*sqrt(-(4*A^2*B^2
*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 -
B^2)*a)/d^2)*log(-(2*(A^3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c
) + a) - (B*d^3*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^
2*B^2 + B^4)*b^2)/d^4) + (2*A^2*B*a + (A^3 - A*B^2)*b)*d)*sqrt(-(2*A*B*b +
d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4
)*b^2)/d^4) - (A^2 - B^2)*a)/d^2))*tan(d*x + c)^2 - 4*a^2*d*sqrt(-(2*A*B*b
- d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B
^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2*(A^3*B + A*B^3)*a + (A^4 - B^4
)*b)*sqrt(b*tan(d*x + c) + a) + (B*d^3*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A
*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (2*A^2*B*a + (A^3 - A*B^2)
*b)*d)*sqrt(-(2*A*B*b - d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b +
(A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2))*tan(d*x + c)^2 +
4*a^2*d*sqrt(-(2*A*B*b - d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*...

```

3.323.6 Sympy [F]

$$\int \cot^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} \cot^3(c + dx) dx$$

```

input integrate(cot(d*x+c)**3*(a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)

```

```

output Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))*cot(c + d*x)**3, x)

```

3.323.7 Maxima [F]

$$\int \cot^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \cot(dx + c)^3 dx$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*cot(d*x + c)^3, x)`

3.323.8 Giac [F(-1)]

Timed out.

$$\int \cot^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.323.9 Mupad [B] (verification not implemented)

Time = 9.46 (sec) , antiderivative size = 14195, normalized size of antiderivative = 64.82

$$\int \cot^3(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`

output $(\operatorname{atan}(\frac{(2A^3b^{14}d^2 + 2A^3a^2b^{12}d^2 - 96A^3a^4b^{10}d^2 - 96A^3a^6b^8d^2 - 160B^3a^3b^{11}d^2 - 160B^3a^5b^9d^2 + 48A^2Bab^{13}d^2 - 192AB^2a^2b^{12}d^2 + 96AB^2a^4b^{10}d^2 + 288AB^2a^6b^8d^2 + 528A^2Bba^3b^{11}d^2 + 480A^2Bba^5b^9d^2)}{(8a^2d^5)} + (\frac{(64Aab^{12}d^4 + 448Aa^3b^{10}d^4 + 384Aa^5b^8d^4 - 256Bba^2b^{11}d^4 - 256Bba^4b^9d^4)}{(8a^2d^5)} - \frac{(512a^2b^{10}d^4 + 768a^4b^8d^4)(a + b\tan(c + dx))^{1/2}(64A^2a^7 + A^2a^3b^4 + 16A^2a^5b^2 + 16B^2a^5b^2 - 64ABba^6b - 8ABba^4b^3)^{1/2}}{(64a^5d^5)})) \frac{(64A^2a^7 + A^2a^3b^4 + 16A^2a^5b^2 + 16B^2a^5b^2 - 64ABba^6b - 8ABba^4b^3)^{1/2}}{(8a^3d)} - ((a + b\tan(c + dx))^{1/2} \frac{(128B^2a^3b^{10}d^2 - 576A^2a^5b^8d^2 - 256A^2a^3b^{10}d^2 + 320B^2a^5b^8d^2 - 4A^2ab^{12}d^2 + 544ABba^2b^{11}d^2 + 1024ABba^4b^9d^2)}{(8a^2d^4)} * (64A^2a^7 + A^2a^3b^4 + 16A^2a^5b^2 + 16B^2a^5b^2 - 64ABba^6b - 8ABba^4b^3)^{1/2}}{(8a^3d)} * (64A^2a^7 + A^2a^3b^4 + 16A^2a^5b^2 + 16B^2a^5b^2 - 64ABba^6b - 8ABba^4b^3)^{1/2}}{(8a^3d)} - ((a + b\tan(c + dx))^{1/2} \frac{(A^2B^2b^{14} - A^4b^{14} + 17A^4a^2b^{12} + 16A^4a^4b^{10} + 96A^4a^6b^8 + 48B^4a^2b^{12} + 48B^4a^4b^{10} + 32B^4a^6b^8 + 95A^2B^2a^2b^{12} + 448A^2B^2a^4b^{10} - 8AB^3a^3b^{13} + 4A^3Bba^3b^{13} - 120AB^3a^3b^{11} + 64AB^3a^5b^9 - 8A^3Bba^3b^{11} - 320A^3Bba^5b^9)}{(8a^2d^4)} * (64A^2a^7 + A^2a^3b^4 \dots$

3.324 $\int \cot^4(c+dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.324.1 Optimal result	3067
3.324.2 Mathematica [B] (verified)	3068
3.324.3 Rubi [A] (warning: unable to verify)	3068
3.324.4 Maple [B] (verified)	3077
3.324.5 Fricas [B] (verification not implemented)	3078
3.324.6 Sympy [F]	3078
3.324.7 Maxima [F]	3079
3.324.8 Giac [F(-1)]	3079
3.324.9 Mupad [B] (verification not implemented)	3080

3.324.1 Optimal result

Integrand size = 33, antiderivative size = 279

$$\int \cot^4(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{(8a^2 Ab - Ab^3 + 16a^3 B + 2ab^2 B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{8a^{5/2}d}$$

$$- \frac{\sqrt{a - ib}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d} + \frac{\sqrt{a + ib}(iA - B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d}$$

$$+ \frac{(8a^2 A + Ab^2 - 2abB) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{8a^2 d}$$

$$- \frac{(Ab + 6aB) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{12ad} - \frac{A \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{3d}$$

output

```
1/8*(8*A*a^2*b-A*b^3+16*B*a^3+2*B*a*b^2)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/a^(5/2)/d-(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))*(a-I*b)^(1/2)/d+(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))*(a+I*b)^(1/2)/d+1/8*(8*A*a^2+A*b^2-2*B*a*b)*cot(d*x+c)*(a+b*tan(d*x+c))^(1/2)/a^2/d-1/12*(A*b+6*B*a)*cot(d*x+c)^2*(a+b*tan(d*x+c))^(1/2)/a/d-1/3*A*cot(d*x+c)^3*(a+b*tan(d*x+c))^(1/2)/d
```

3.324.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 566 vs. $2(279) = 558$.

Time = 6.49 (sec) , antiderivative size = 566, normalized size of antiderivative = 2.03

$$\int \cot^4(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= 2b^4 \left(\frac{5A \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{16a^{5/2}b} + \frac{(Ab+aB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ab^4}} - \frac{3(Ab+aB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{8a^{5/2}b^2} - \frac{(aA-bB)}{\dots} \right)$$

input `Integrate[Cot[c + d*x]^4*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output

```
(2*b^4*((5*A*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/(16*a^(5/2)*b) + (
(A*b + a*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]]/(Sqrt[a]*b^4) - (3*
(A*b + a*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]]/(8*a^(5/2)*b^2) - (
(a*A - b*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]]/(2*a^(3/2)*b^3) + (
(a*A*b - A*b*Sqrt[-b^2] - b^2*B - a*Sqrt[-b^2]*B)*ArcTanh[Sqrt[a + b*Tan[c
+ d*x]]/Sqrt[a - Sqrt[-b^2]]])/(2*b^4*Sqrt[-b^2]*Sqrt[a - Sqrt[-b^2]]) -
((a*A*b + A*b*Sqrt[-b^2] - b^2*B + a*Sqrt[-b^2]*B)*ArcTanh[Sqrt[a + b*Tan[
c + d*x]]/Sqrt[a + Sqrt[-b^2]]])/(2*(-b^2)^(5/2)*Sqrt[a + Sqrt[-b^2]]) - (
5*A*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]]/(16*a^2*b^2) + (3*(A*b + a*B)*C
ot[c + d*x]*Sqrt[a + b*Tan[c + d*x]]/(8*a^2*b^3) + ((a*A - b*B)*Cot[c + d
*x]*Sqrt[a + b*Tan[c + d*x]]/(2*a*b^4) + (5*A*Cot[c + d*x]^2*Sqrt[a + b*T
an[c + d*x]]/(24*a*b^3) - ((A*b + a*B)*Cot[c + d*x]^2*Sqrt[a + b*Tan[c +
d*x]]/(4*a*b^4) - (A*Cot[c + d*x]^3*Sqrt[a + b*Tan[c + d*x]]/(6*b^4))))/d
```

3.324.3 Rubi [A] (warning: unable to verify)

Time = 2.31 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.03, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4091, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^4(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

3.324. $\int \cot^4(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan(c+dx)^4} dx \\
& \quad \downarrow \text{3042} \\
& -\frac{1}{3} \int -\frac{\cot^3(c+dx)(-5Ab \tan^2(c+dx)-6(aA-bB) \tan(c+dx)+Ab+6aB)}{2\sqrt{a+b \tan(c+dx)}} dx - \\
& \quad \frac{A \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{6} \int \frac{\cot^3(c+dx)(-5Ab \tan^2(c+dx)-6(aA-bB) \tan(c+dx)+Ab+6aB)}{\sqrt{a+b \tan(c+dx)}} dx - \\
& \quad \frac{A \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{6} \int \frac{-5Ab \tan(c+dx)^2-6(aA-bB) \tan(c+dx)+Ab+6aB}{\tan(c+dx)^3 \sqrt{a+b \tan(c+dx)}} dx - \\
& \quad \frac{A \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} \\
& \quad \downarrow \text{4132} \\
& \frac{1}{6} \left(-\frac{\int \frac{3 \cot^2(c+dx)(8Aa^2-2bBa+8(Ab+aB) \tan(c+dx)a+Ab^2+b(Ab+6aB) \tan^2(c+dx))}{2\sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{(6aB+Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2ad} \right. \\
& \quad \left. \frac{A \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} \right) \\
& \quad \downarrow \text{27} \\
& \frac{1}{6} \left(-\frac{3 \int \frac{\cot^2(c+dx)(8Aa^2-2bBa+8(Ab+aB) \tan(c+dx)a+Ab^2+b(Ab+6aB) \tan^2(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx}{4a} - \frac{(6aB+Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2ad} \right. \\
& \quad \left. \frac{A \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

$$\frac{1}{6} \left(\frac{3 \int \frac{8Aa^2 - 2bBa + 8(Ab + aB) \tan(c + dx)a + Ab^2 + b(Ab + 6aB) \tan(c + dx)^2}{\tan(c + dx)^2 \sqrt{a + b \tan(c + dx)}} dx}{4a} - \frac{(6aB + Ab) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2ad} \right)$$

$$\frac{A \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{3d}$$

↓ 4132

$$\frac{1}{6} \left(\frac{3 \left(\int - \frac{\cot(c + dx) (16Ba^3 + 8Aba^2 - 16(aA - bB) \tan(c + dx)a^2 + 2b^2Ba - Ab^3 - b(8Aa^2 - 2bBa + Ab^2) \tan^2(c + dx))}{2\sqrt{a + b \tan(c + dx)}} dx}{a} - \frac{(8a^2A - 2abB + Ab^2) \cot(c + dx)}{ad} \right)}{4a}$$

$$\frac{A \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{3d}$$

↓ 27

$$\frac{1}{6} \left(\frac{3 \left(\int \frac{\cot(c + dx) (16Ba^3 + 8Aba^2 - 16(aA - bB) \tan(c + dx)a^2 + 2b^2Ba - Ab^3 - b(8Aa^2 - 2bBa + Ab^2) \tan^2(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx}{2a} - \frac{(8a^2A - 2abB + Ab^2) \cot(c + dx)}{ad} \right)}{4a}$$

$$\frac{A \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{3 \left(\int \frac{16Ba^3 + 8Aba^2 - 16(aA - bB) \tan(c + dx)a^2 + 2b^2Ba - Ab^3 - b(8Aa^2 - 2bBa + Ab^2) \tan(c + dx)^2}{\tan(c + dx) \sqrt{a + b \tan(c + dx)}} dx}{2a} - \frac{(8a^2A - 2abB + Ab^2) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{ad} \right)}{4a}$$

$$\frac{A \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{3d}$$

↓ 4136

$$\frac{1}{6} \left(\frac{3 \left(\int -\frac{16((aA-bB)a^2+(Ab+aB)\tan(c+dx)a^2)}{\sqrt{a+b\tan(c+dx)}} dx + (16a^3B+8a^2Ab+2ab^2B-Ab^3) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b\tan(c+dx)}} dx - (8a^2A-2abB+Ab^2) \cot(c+dx) \right)}{4a} \right)$$

$$\frac{A \cot^3(c+dx) \sqrt{a+b\tan(c+dx)}}{3d}$$

↓ 27

$$\frac{1}{6} \left(\frac{3 \left(\frac{(16a^3B+8a^2Ab+2ab^2B-Ab^3) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b\tan(c+dx)}} dx - 16 \int \frac{(aA-bB)a^2+(Ab+aB)\tan(c+dx)a^2}{\sqrt{a+b\tan(c+dx)}} dx - (8a^2A-2abB+Ab^2) \cot(c+dx)}{4a} \right)}{4a}$$

$$\frac{A \cot^3(c+dx) \sqrt{a+b\tan(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{3 \left(\frac{(16a^3B+8a^2Ab+2ab^2B-Ab^3) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - 16 \int \frac{(aA-bB)a^2+(Ab+aB)\tan(c+dx)a^2}{\sqrt{a+b\tan(c+dx)}} dx - (8a^2A-2abB+Ab^2) \cot(c+dx)}{4a} \right)}{4a}$$

$$\frac{A \cot^3(c+dx) \sqrt{a+b\tan(c+dx)}}{3d}$$

↓ 4022

$$- \frac{A \cot^3(c+dx) \sqrt{a+b\tan(c+dx)}}{3d} +$$

$$\frac{1}{6} \left(\frac{(6aB+Ab) \cot^2(c+dx) \sqrt{a+b\tan(c+dx)}}{2ad} - \frac{3 \left(-\frac{(8a^2A-2abB+Ab^2) \cot(c+dx) \sqrt{a+b\tan(c+dx)}}{ad} + \frac{(16a^3B+8a^2Ab+2ab^2B-Ab^3) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b\tan(c+dx)}} dx - 16 \int \frac{(aA-bB)a^2+(Ab+aB)\tan(c+dx)a^2}{\sqrt{a+b\tan(c+dx)}} dx - (8a^2A-2abB+Ab^2) \cot(c+dx)}{4a} \right)}{4a}$$

↓ 3042

$$\frac{1}{6} \left(-\frac{(6aB + Ab) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2ad} - \frac{A \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{3d} + 3 \left(-\frac{(8a^2A - 2abB + Ab^2) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{ad} + \frac{(16a^3B + 8a^2Ab + \dots)}{\dots} \right) \right)$$

↓ 4020

$$\frac{1}{6} \left(-\frac{(6aB + Ab) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2ad} - \frac{A \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{3d} + 3 \left(-\frac{(8a^2A - 2abB + Ab^2) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{ad} + \frac{(16a^3B + 8a^2Ab + \dots)}{\dots} \right) \right)$$

↓ 25

$$\frac{1}{6} \left(-\frac{(6aB + Ab) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2ad} - \frac{A \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{3d} + 3 \left(-\frac{(8a^2A - 2abB + Ab^2) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{ad} + \frac{(16a^3B + 8a^2Ab + \dots)}{\dots} \right) \right)$$

↓ 73

$$\frac{1}{6} \left(-\frac{(6aB + Ab) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2ad} - \frac{A \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{3d} + 3 \left(-\frac{(8a^2A - 2abB + Ab^2) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{ad} + \frac{(16a^3B + 8a^2Ab + 2a^2A)}{2ad} \right) \right)$$

↓ 221

$$\frac{1}{6} \left(-\frac{(6aB + Ab) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2ad} - \frac{A \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{3d} + 3 \left(-\frac{(8a^2A - 2abB + Ab^2) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{ad} + \frac{(16a^3B + 8a^2Ab + 2a^2A)}{2ad} \right) \right)$$

↓ 4117

$$\frac{1}{6} \left(-\frac{(6aB + Ab) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2ad} - \frac{A \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{3d} + 3 \left(-\frac{(8a^2A - 2abB + Ab^2) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{ad} + \frac{(16a^3B + 8a^2Ab + 2a^2A)}{2ad} \right) \right)$$

↓ 73

$$\begin{aligned}
 & -\frac{A \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \\
 & \left(\frac{1}{6} \frac{(6aB + Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2ad} - \frac{3 \left(-\frac{(8a^2A - 2abB + Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} + \frac{2(16a^3B + 8a^2Ab + 3a^2)}{6ad} \right)}{6} \right) \\
 & \quad \downarrow 221 \\
 & -\frac{A \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \\
 & \left(\frac{1}{6} \frac{(6aB + Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2ad} - \frac{3 \left(-\frac{(8a^2A - 2abB + Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} + \frac{2(16a^3B + 8a^2Ab + 3a^2)}{6ad} \right)}{6} \right)
 \end{aligned}$$

```
input Int[Cot[c + d*x]^4*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]
```

```
output -1/3*(A*Cot[c + d*x]^3*Sqrt[a + b*Tan[c + d*x]])/d + (-1/2*((A*b + 6*a*B)*Cot[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]])/(a*d) - (3*((-16*((a^2*Sqrt[a - I*b])*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d + (a^2*Sqrt[a + I*b])*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d - (2*(8*a^2*A*b - A*b^3 + 16*a^3*B + 2*a*b^2*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/(Sqrt[a]*d))/(2*a) - ((8*a^2*A + A*b^2 - 2*a*b*B)*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/(a*d))/(4*a))/6
```

3.324.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4091 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.324.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1294 vs. $2(241) = 482$.

Time = 0.23 (sec) , antiderivative size = 1295, normalized size of antiderivative = 4.64

method	result	size
derivativedivides	Expression too large to display	1295
default	Expression too large to display	1295

```
input int(cot(d*x+c)^4*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output -1/4/d/b*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(
(1/2)+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)+1/4
/d/b*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2
)+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-1/4/d*ln(b*tan(d*x+c)
+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*B
*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2
*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-
2*a)^(1/2))*B*(a^2+b^2)^(1/2)+1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((
2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)
-2*a)^(1/2))*A+1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c)
)^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*a+
1/4/d/b*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)
-a-(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)-1/4/
d/b*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a
-(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+1/4/d*ln((a+b*tan(d*x+c)
)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*B*
(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2
*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2
*a)^(1/2))*B*(a^2+b^2)^(1/2)-1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((
2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)...
```

3.324.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1376 vs. $2(235) = 470$.

Time = 9.35 (sec) , antiderivative size = 2769, normalized size of antiderivative = 9.92

$$\int \cot^4(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm
m="fricas")`

output `[-1/48*(24*a^3*d*sqrt((2*A*B*b + d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B
^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2*
(A^3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) + (A*d^3*sqrt(
-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4
) - (2*A*B^2*a + (A^2*B - B^3)*b)*d)*sqrt((2*A*B*b + d^2*sqrt(-(4*A^2*B^2*
a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B
^2)*a)/d^2))*tan(d*x + c)^3 - 24*a^3*d*sqrt((2*A*B*b + d^2*sqrt(-(4*A^2*B^2
2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 -
B^2)*a)/d^2)*log(-(2*(A^3*B + A*B^3)*a + (A^4 - B^4)*b)*sqrt(b*tan(d*x +
c) + a) - (A*d^3*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A
^2*B^2 + B^4)*b^2)/d^4) - (2*A*B^2*a + (A^2*B - B^3)*b)*d)*sqrt((2*A*B*b +
d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4
) *b^2)/d^4) - (A^2 - B^2)*a)/d^2))*tan(d*x + c)^3 - 24*a^3*d*sqrt((2*A*B*b
- d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B
^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2)*log(-(2*(A^3*B + A*B^3)*a + (A^4 - B^4
) *b)*sqrt(b*tan(d*x + c) + a) + (A*d^3*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A
*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) + (2*A*B^2*a + (A^2*B - B^3)
*b)*d)*sqrt((2*A*B*b - d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a*b +
(A^4 - 2*A^2*B^2 + B^4)*b^2)/d^4) - (A^2 - B^2)*a)/d^2))*tan(d*x + c)^3 +
24*a^3*d*sqrt((2*A*B*b - d^2*sqrt(-(4*A^2*B^2*a^2 + 4*(A^3*B - A*B^3)*a...`

3.324.6 Sympy [F]

$$\begin{aligned} & \int \cot^4(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= \int (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} \cot^4(c + dx) dx \end{aligned}$$

input `integrate(cot(d*x+c)**4*(a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))*cot(c + d*x)**4, x)`

3.324.7 Maxima [F]

$$\begin{aligned} & \int \cot^4(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \cot(dx + c)^4 dx \end{aligned}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*cot(d*x + c)^4, x)`

3.324.8 Giac [F(-1)]

Timed out.

$$\int \cot^4(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.324.9 Mupad [B] (verification not implemented)

Time = 9.61 (sec) , antiderivative size = 16796, normalized size of antiderivative = 60.20

$$\int \cot^4(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(cot(c + d*x)^4*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)
```

```
output atan(((((((224*A*a^4*b^11*d^4 - 32*A*a^2*b^13*d^4 + 256*A*a^6*b^9*d^4 + 64
*B*a^3*b^12*d^4 + 448*B*a^5*b^10*d^4 + 384*B*a^7*b^8*d^4)/(a^4*d^5) - ((20
48*a^4*b^10*d^4 + 3072*a^6*b^8*d^4)*(a + b*tan(c + d*x))^(1/2)*((B^2*a)/(4
*d^2) - (A^2*a)/(4*d^2) - (2*A^2*B^2*b^2*d^4 - B^4*b^2*d^4 - 4*A^2*B^2*a^2
*d^4 - A^4*b^2*d^4 + 4*A*B^3*a*b*d^4 - 4*A^3*B*a*b*d^4)^(1/2)/(4*d^4) + (A
*B*b)/(2*d^2))^(1/2))/(4*a^4*d^4))*((B^2*a)/(4*d^2) - (A^2*a)/(4*d^2) - (2
*A^2*B^2*b^2*d^4 - B^4*b^2*d^4 - 4*A^2*B^2*a^2*d^4 - A^4*b^2*d^4 + 4*A*B^3
*a*b*d^4 - 4*A^3*B*a*b*d^4)^(1/2)/(4*d^4) + (A*B*b)/(2*d^2))^(1/2) + ((a +
b*tan(c + d*x))^(1/2)*(16*B^2*a^3*b^12*d^2 - 512*A^2*a^5*b^10*d^2 - 1280*
A^2*a^7*b^8*d^2 - 64*A^2*a^3*b^12*d^2 + 1024*B^2*a^5*b^10*d^2 + 2304*B^2*a
^7*b^8*d^2 + 4*A^2*a*b^14*d^2 - 16*A*B*a^2*b^13*d^2 + 2048*A*B*a^4*b^11*d
^2 + 4096*A*B*a^6*b^9*d^2))/(4*a^4*d^4))*((B^2*a)/(4*d^2) - (A^2*a)/(4*d^2)
- (2*A^2*B^2*b^2*d^4 - B^4*b^2*d^4 - 4*A^2*B^2*a^2*d^4 - A^4*b^2*d^4 + 4*
A*B^3*a*b*d^4 - 4*A^3*B*a*b*d^4)^(1/2)/(4*d^4) + (A*B*b)/(2*d^2))^(1/2) -
(16*A^3*a^3*b^13*d^2 - 144*A^3*a^5*b^11*d^2 - 160*A^3*a^7*b^9*d^2 - 2*B^3*
a^2*b^14*d^2 - 2*B^3*a^4*b^12*d^2 + 96*B^3*a^6*b^10*d^2 + 96*B^3*a^8*b^8*d
^2 - (A^2*B*b^16*d^2)/2 + 2*A*B^2*a*b^15*d^2 + 50*A*B^2*a^3*b^13*d^2 + 528
*A*B^2*a^5*b^11*d^2 + 480*A*B^2*a^7*b^9*d^2 - (49*A^2*B*a^2*b^14*d^2)/2 +
168*A^2*B*a^4*b^12*d^2 - 96*A^2*B*a^6*b^10*d^2 - 288*A^2*B*a^8*b^8*d^2)/(a
^4*d^5))*((B^2*a)/(4*d^2) - (A^2*a)/(4*d^2) - (2*A^2*B^2*b^2*d^4 - B^4*...
```

3.325 $\int \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.325.1 Optimal result	3081
3.325.2 Mathematica [A] (verified)	3082
3.325.3 Rubi [A] (warning: unable to verify)	3082
3.325.4 Maple [B] (verified)	3086
3.325.5 Fricas [B] (verification not implemented)	3087
3.325.6 Sympy [F]	3088
3.325.7 Maxima [F]	3089
3.325.8 Giac [F(-1)]	3089
3.325.9 Mupad [B] (verification not implemented)	3089

3.325.1 Optimal result

Integrand size = 33, antiderivative size = 214

$$\int \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{(a-ib)^{3/2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d} - \frac{(a+ib)^{3/2}(iA-B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d} - \frac{2(Ab+aB)\sqrt{a+b \tan(c+dx)}}{d} - \frac{2B(a+b \tan(c+dx))^{3/2}}{3d} + \frac{2(7Ab-2aB)(a+b \tan(c+dx))^{5/2}}{35b^2d} + \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd}$$

output $(a-I*b)^{(3/2)}*(I*A+B)*\operatorname{arctanh}((a+b*\tan(d*x+c))^{(1/2)}/(a-I*b)^{(1/2)})/d-(a+I*b)^{(3/2)}*(I*A-B)*\operatorname{arctanh}((a+b*\tan(d*x+c))^{(1/2)}/(a+I*b)^{(1/2)})/d-2*(A*b+B*a)*(a+b*\tan(d*x+c))^{(1/2)}/d-2/3*B*(a+b*\tan(d*x+c))^{(3/2)}/d+2/35*(7*A*b-2*B*a)*(a+b*\tan(d*x+c))^{(5/2)}/b^2/d+2/7*B*\tan(d*x+c)*(a+b*\tan(d*x+c))^{(5/2)}/b/d$

3.325.2 Mathematica [A] (verified)

Time = 2.80 (sec) , antiderivative size = 252, normalized size of antiderivative = 1.18

$$\int \tan^2(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx = \frac{2(7Ab-2aB)(a+b\tan(c+dx))^{5/2}}{b} + 10B\tan(c+dx)(a+b\tan(c+dx))^{5/2} + \frac{35}{3}b(A-iB)\left(3\sqrt{\dots}\right)$$

```
input Integrate[Tan[c + d*x]^2*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x
]
```

```
output ((2*(7*A*b - 2*a*B)*(a + b*Tan[c + d*x])^(5/2))/b + 10*B*Tan[c + d*x]*(a +
b*Tan[c + d*x])^(5/2) + (35*b*(A - I*B)*(3*Sqrt[a - I*b]*(I*a + b)*ArcTan
h[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] - I*Sqrt[a + b*Tan[c + d*x]]*(4*
a - (3*I)*b + b*Tan[c + d*x]))/3 + (35*b*(A + I*B)*(3*Sqrt[a + I*b]*((-I)
*a + b)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + I*Sqrt[a + b*Tan
[c + d*x]]*(4*a + (3*I)*b + b*Tan[c + d*x]))/3)/(35*b*d)
```

3.325.3 Rubi [A] (warning: unable to verify)Time = 1.28 (sec) , antiderivative size = 209, normalized size of antiderivative = 0.98, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 4090, 27, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^2(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx$$

$$\downarrow 3042$$

$$\int \tan(c+dx)^2(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx$$

$$\downarrow 4090$$

$$\frac{2 \int -\frac{1}{2}(a+b\tan(c+dx))^{3/2} \left(-((7Ab-2aB)\tan^2(c+dx)) + 7bB\tan(c+dx) + 2aB \right) dx}{7b} + \frac{2B\tan(c+dx)(a+b\tan(c+dx))^{5/2}}{7bd}$$

3.325. $\int \tan^2(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx$

$$\begin{array}{c}
\downarrow 27 \\
\frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd} - \\
\frac{\int (a+b \tan(c+dx))^{3/2} (-((7Ab-2aB) \tan^2(c+dx)) + 7bB \tan(c+dx) + 2aB) dx}{7b} \\
\downarrow 3042 \\
\frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd} - \\
\frac{\int (a+b \tan(c+dx))^{3/2} (-((7Ab-2aB) \tan(c+dx)^2) + 7bB \tan(c+dx) + 2aB) dx}{7b} \\
\downarrow 4113 \\
\frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd} - \\
\frac{\int (a+b \tan(c+dx))^{3/2} (7Ab + 7B \tan(c+dx)b) dx - \frac{2(7Ab-2aB)(a+b \tan(c+dx))^{5/2}}{5bd}}{7b} \\
\downarrow 3042 \\
\frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd} - \\
\frac{\int (a+b \tan(c+dx))^{3/2} (7Ab + 7B \tan(c+dx)b) dx - \frac{2(7Ab-2aB)(a+b \tan(c+dx))^{5/2}}{5bd}}{7b} \\
\downarrow 4011 \\
\frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd} - \\
\frac{\int \sqrt{a+b \tan(c+dx)} (7b(aA-bB) + 7b(Ab+aB) \tan(c+dx)) dx - \frac{2(7Ab-2aB)(a+b \tan(c+dx))^{5/2}}{5bd} + \frac{14bB(a+b \tan(c+dx))^{5/2}}{3d}}{7b} \\
\downarrow 3042 \\
\frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd} - \\
\frac{\int \sqrt{a+b \tan(c+dx)} (7b(aA-bB) + 7b(Ab+aB) \tan(c+dx)) dx - \frac{2(7Ab-2aB)(a+b \tan(c+dx))^{5/2}}{5bd} + \frac{14bB(a+b \tan(c+dx))^{5/2}}{3d}}{7b} \\
\downarrow 4011 \\
\frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd} - \\
\frac{\int \frac{7b(Aa^2-2bBa-Ab^2)+7b(Ba^2+2Aba-b^2B) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx - \frac{2(7Ab-2aB)(a+b \tan(c+dx))^{5/2}}{5bd} + \frac{14b(aB+Ab)\sqrt{a+b \tan(c+dx)}}{d} + \frac{14bB(a+b \tan(c+dx))^{5/2}}{3d}}{7b} \\
\downarrow 3042
\end{array}$$

3.325. $\int \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\frac{\frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd} - \int \frac{7b(Aa^2-2bBa-Ab^2)+7b(Ba^2+2Aba-b^2B) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx - \frac{2(7Ab-2aB)(a+b \tan(c+dx))^{5/2}}{5bd} + \frac{14b(aB+Ab)\sqrt{a+b \tan(c+dx)}}{d} + \frac{14bB(a+b \tan(c+dx))^{3/2}}{5bd}}{7b} \xrightarrow{4022}$$

$$\frac{\frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd} - \frac{7}{2}b(a+ib)^2(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{7}{2}b(a-ib)^2(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx - \frac{2(7Ab-2aB)(a+b \tan(c+dx))^{5/2}}{5bd}}{7b} \xrightarrow{3042}$$

$$\frac{\frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd} - \frac{7}{2}b(a+ib)^2(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{7}{2}b(a-ib)^2(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx - \frac{2(7Ab-2aB)(a+b \tan(c+dx))^{5/2}}{5bd}}{7b} \xrightarrow{4020}$$

$$\frac{\frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd} - \frac{7ib(a-ib)^2(A-iB) \int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} - \frac{7ib(a+ib)^2(A+iB) \int -\frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d}}{7b} \xrightarrow{25}$$

$$\frac{\frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd} - \frac{7ib(a-ib)^2(A-iB) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \frac{7ib(a+ib)^2(A+iB) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d}}{7b} \xrightarrow{73}$$

$$\frac{\frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd} - \frac{7(a+ib)^2(A+iB) \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{d} + \frac{7(a-ib)^2(A-iB) \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{d} - \frac{2(7Ab-2aB)(a+b \tan(c+dx))^{5/2}}{5bd}}{7b} \xrightarrow{221}$$

$$\frac{\frac{2B \tan(c+dx)(a+b \tan(c+dx))^{5/2}}{7bd} - \frac{7b(a-ib)^{3/2}(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{7b(a+ib)^{3/2}(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} - \frac{2(7Ab-2aB)(a+b \tan(c+dx))^{5/2}}{5bd} + \frac{14b(aB+Ab)\sqrt{a+b \tan(c+dx)}}{d}}{7b}$$

3.325. $\int \tan^2(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

input `Int[Tan[c + d*x]^2*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(2*B*Tan[c + d*x]*(a + b*Tan[c + d*x])^(5/2))/(7*b*d) - ((7*(a - I*b)^(3/2)*b*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d + (7*(a + I*b)^(3/2)*b*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d + (14*b*(A*b + a*B)*Sqrt[a + b*Tan[c + d*x]])/d + (14*b*B*(a + b*Tan[c + d*x])^(3/2))/(3*d) - (2*(7*A*b - 2*a*B)*(a + b*Tan[c + d*x])^(5/2))/(5*b*d))/(7*b)`

3.325.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n)), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

3.325.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1696 vs. $2(182) = 364$.

Time = 0.15 (sec) , antiderivative size = 1697, normalized size of antiderivative = 7.93

method	result	size
parts	Expression too large to display	1697
derivativedivides	Expression too large to display	1717
default	Expression too large to display	1717

input `int(tan(d*x+c)^2*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)`

output `A*(2/5/b/d*(a+b*tan(d*x+c))^(5/2)-2*b*(a+b*tan(d*x+c))^(1/2)/d+1/4/b/d*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-1/4/b/d*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2+1/4*b/d*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*b/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a+b/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*(a^2+b^2)^(1/2)-1/4/b/d*ln(b*tan(d*x+c)+a-(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+1/4/b/d*ln(b*tan(d*x+c)+a-(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2-1/4*b/d*ln(b*tan(d*x+c)+a-(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*b/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)-(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a+b/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)-(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*(a^2+b^2)^(1/2))+B*(2/7/d/b^2*(a+b*tan(d*x+c))^(7/2)-2/5/d/b^2*(a+b*tan(d*...`

3.325.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3148 vs. 2(176) = 352.

Time = 0.48 (sec) , antiderivative size = 3148, normalized size of antiderivative = 14.71

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm
m="fricas")`

output

```
-1/210*(105*b^2*d*sqrt((6*A*B*a^2*b - 2*A*B*b^3 - (A^2 - B^2)*a^3 + 3*(A^2
- B^2)*a*b^2 + d^2*sqrt(-(4*A^2*B^2*a^6 + 12*(A^3*B - A*B^3)*a^5*b + 3*(3
*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 - 40*(A^3*B - A*B^3)*a^3*b^3 - 6*(A^4 -
8*A^2*B^2 + B^4)*a^2*b^4 + 12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2*B^2 +
B^4)*b^6)/d^4))/d^2)*log((2*(A^3*B + A*B^3)*a^5 + 3*(A^4 - B^4)*a^4*b - 4*
(A^3*B + A*B^3)*a^3*b^2 + 2*(A^4 - B^4)*a^2*b^3 - 6*(A^3*B + A*B^3)*a*b^4
- (A^4 - B^4)*b^5)*sqrt(b*tan(d*x + c) + a) + ((A*a - B*b)*d^3*sqrt(-(4*A^
2*B^2*a^6 + 12*(A^3*B - A*B^3)*a^5*b + 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*
b^2 - 40*(A^3*B - A*B^3)*a^3*b^3 - 6*(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 + 12*
(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2*B^2 + B^4)*b^6)/d^4) - (2*A*B^2*a^4 +
(5*A^2*B - 3*B^3)*a^3*b + 3*(A^3 - 3*A*B^2)*a^2*b^2 - (7*A^2*B - B^3)*a*b
^3 - (A^3 - A*B^2)*b^4)*d)*sqrt((6*A*B*a^2*b - 2*A*B*b^3 - (A^2 - B^2)*a^3
+ 3*(A^2 - B^2)*a*b^2 + d^2*sqrt(-(4*A^2*B^2*a^6 + 12*(A^3*B - A*B^3)*a^5
*b + 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 - 40*(A^3*B - A*B^3)*a^3*b^3 -
6*(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 + 12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A
^2*B^2 + B^4)*b^6)/d^4))/d^2)) - 105*b^2*d*sqrt((6*A*B*a^2*b - 2*A*B*b^3 -
(A^2 - B^2)*a^3 + 3*(A^2 - B^2)*a*b^2 + d^2*sqrt(-(4*A^2*B^2*a^6 + 12*(A^
3*B - A*B^3)*a^5*b + 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 - 40*(A^3*B -
A*B^3)*a^3*b^3 - 6*(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 + 12*(A^3*B - A*B^3)*a*
b^5 + (A^4 - 2*A^2*B^2 + B^4)*b^6)/d^4))/d^2)*log((2*(A^3*B + A*B^3)*a^...
```

3.325.6 Sympy [F]

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx))(a + b \tan(c + dx))^{\frac{3}{2}} \tan^2(c + dx) dx$$

input `integrate(tan(d*x+c)**2*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)*tan(c + d*x)**2, x)`

3.325.7 Maxima [F]

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{3/2} \tan(dx + c)^2 dx$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)*tan(d*x + c)^2, x)`

3.325.8 Giac [F(-1)]

Timed out.

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.325.9 Mupad [B] (verification not implemented)

Time = 82.10 (sec) , antiderivative size = 2993, normalized size of antiderivative = 13.99

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2),x)`

output

$$\begin{aligned} & \log((16A^3ab^3(a^2 + b^2)^2)/d^3 - (((16b^2*(((-A^4b^2d^4(3a^2 - b^2)^2)^{(1/2)} - A^2a^3d^2 + 3A^2ab^2d^2)/d^4)^{(1/2)}*(Ab^3 + Aa^2b \\ & + a*d*(((-A^4b^2d^4(3a^2 - b^2)^2)^{(1/2)} - A^2a^3d^2 + 3A^2ab^2d^2)/d^4)^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)}))/d + (16A^2b^2*(a + b*\tan(c \\ & + d*x))^{(1/2)}*(a^4 + b^4 - 6a^2b^2))/d^2)*(((-A^4b^2d^4(3a^2 - b^2)^2)^{(1/2)} - A^2a^3d^2 + 3A^2ab^2d^2)/d^4)^{(1/2)})/2)*((6A^4a^2b^4d \\ & ^4 - A^4b^6d^4 - 9A^4a^4b^2d^4)^{(1/2)}/(4d^4) - (A^2a^3)/(4d^2) + (3A^2ab^2)/(4d^2))^{(1/2)} - \log((16A^3ab^3(a^2 + b^2)^2)/d^3 - (((1 \\ & 6b^2*(-((-A^4b^2d^4(3a^2 - b^2)^2)^{(1/2)} + A^2a^3d^2 - 3A^2ab^2d^2)/d^4)^{(1/2)}*(Ab^3 + Aa^2b - a*d*(-((-A^4b^2d^4(3a^2 - b^2)^2)^{(\\ & 1/2)} + A^2a^3d^2 - 3A^2ab^2d^2)/d^4)^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)} \\ &)))/d - (16A^2b^2*(a + b*\tan(c + d*x))^{(1/2)}*(a^4 + b^4 - 6a^2b^2))/d^2)*(-((-A^4b^2d^4(3a^2 - b^2)^2)^{(1/2)} + A^2a^3d^2 - 3A^2ab^2d^2 \\ &)/d^4)^{(1/2)})/2)*(-((6A^4a^2b^4d^4 - A^4b^6d^4 - 9A^4a^4b^2d^4)^{(1/2)} + A^2a^3d^2 - 3A^2ab^2d^2)/(4d^4))^{(1/2)} - \log((16A^3ab^3 \\ & (a^2 + b^2)^2)/d^3 - (((16b^2*(((-A^4b^2d^4(3a^2 - b^2)^2)^{(1/2)} - A^2a^3d^2 + 3A^2ab^2d^2)/d^4)^{(1/2)}*(Ab^3 + Aa^2b - a*d*(((-A^4b^2 \\ & *d^4(3a^2 - b^2)^2)^{(1/2)} - A^2a^3d^2 + 3A^2ab^2d^2)/d^4)^{(1/2)}*(a \\ & + b*\tan(c + d*x))^{(1/2)}))/d - (16A^2b^2*(a + b*\tan(c + d*x))^{(1/2)}*(a^4 \\ & + b^4 - 6a^2b^2))/d^2)*(((-A^4b^2d^4(3a^2 - b^2)^2)^{(1/2)} - A^2*... \end{aligned}$$

3.326 $\int \tan(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.326.1 Optimal result	3091
3.326.2 Mathematica [A] (verified)	3092
3.326.3 Rubi [A] (warning: unable to verify)	3092
3.326.4 Maple [B] (verified)	3096
3.326.5 Fricas [B] (verification not implemented)	3097
3.326.6 Sympy [F]	3097
3.326.7 Maxima [F]	3098
3.326.8 Giac [F(-1)]	3098
3.326.9 Mupad [B] (verification not implemented)	3099

3.326.1 Optimal result

Integrand size = 31, antiderivative size = 175

$$\int \tan(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx =$$

$$\frac{(a - ib)^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d}$$

$$- \frac{(a + ib)^{3/2}(A + iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d} + \frac{2(aA - bB)\sqrt{a + b \tan(c + dx)}}{d}$$

$$+ \frac{2A(a + b \tan(c + dx))^{3/2}}{3d} + \frac{2B(a + b \tan(c + dx))^{5/2}}{5bd}$$

output

```
-(a-I*b)^(3/2)*(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d-(a+
I*b)^(3/2)*(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d+2*(A*a-
B*b)*(a+b*tan(d*x+c))^(1/2)/d+2/3*A*(a+b*tan(d*x+c))^(3/2)/d+2/5*B*(a+b*ta
n(d*x+c))^(5/2)/b/d
```

3.326.2 Mathematica [A] (verified)

Time = 1.39 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.10

$$\int \tan(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{6B(a+b \tan(c+dx))^{5/2}}{b} + 5(A-iB) \left(-3(a-ib)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}} \right) + \sqrt{a+b \tan(c+dx)} \right)$$

input `Integrate[Tan[c + d*x]*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`output `((6*B*(a + b*Tan[c + d*x])^(5/2))/b + 5*(A - I*B)*(-3*(a - I*b)^(3/2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] + Sqrt[a + b*Tan[c + d*x]]*(4*a - (3*I)*b + b*Tan[c + d*x])) + 5*(A + I*B)*(-3*(a + I*b)^(3/2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + Sqrt[a + b*Tan[c + d*x]]*(4*a + (3*I)*b + b*Tan[c + d*x])))/(15*d)`**3.326.3 Rubi [A] (warning: unable to verify)**Time = 0.92 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.91, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 4075, 3042, 4011, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx \\ & \quad \downarrow \text{4075} \\ & \int (A \tan(c+dx) - B)(a+b \tan(c+dx))^{3/2} dx + \frac{2B(a+b \tan(c+dx))^{5/2}}{5bd} \\ & \quad \downarrow \text{3042} \\ & \int (A \tan(c+dx) - B)(a+b \tan(c+dx))^{3/2} dx + \frac{2B(a+b \tan(c+dx))^{5/2}}{5bd} \end{aligned}$$

$$\begin{aligned}
& \downarrow 4011 \\
& \int \sqrt{a + b \tan(c + dx)}(-Ab - aB + (aA - bB) \tan(c + dx))dx + \frac{2A(a + b \tan(c + dx))^{3/2}}{3d} + \\
& \quad \frac{2B(a + b \tan(c + dx))^{5/2}}{5bd} \\
& \downarrow 3042 \\
& \int \sqrt{a + b \tan(c + dx)}(-Ab - aB + (aA - bB) \tan(c + dx))dx + \frac{2A(a + b \tan(c + dx))^{3/2}}{3d} + \\
& \quad \frac{2B(a + b \tan(c + dx))^{5/2}}{5bd} \\
& \downarrow 4011 \\
& \int \frac{-Ba^2 - 2Aba + b^2B + (Aa^2 - 2bBa - Ab^2) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2(aA - bB)\sqrt{a + b \tan(c + dx)}}{d} + \\
& \quad \frac{2A(a + b \tan(c + dx))^{3/2}}{3d} + \frac{2B(a + b \tan(c + dx))^{5/2}}{5bd} \\
& \downarrow 3042 \\
& \int \frac{-Ba^2 - 2Aba + b^2B + (Aa^2 - 2bBa - Ab^2) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2(aA - bB)\sqrt{a + b \tan(c + dx)}}{d} + \\
& \quad \frac{2A(a + b \tan(c + dx))^{3/2}}{3d} + \frac{2B(a + b \tan(c + dx))^{5/2}}{5bd} \\
& \downarrow 4022 \\
& \frac{1}{2}(a + ib)^2(-B + iA) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx - \frac{1}{2}(a - ib)^2(B + iA) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \\
& \quad \frac{2(aA - bB)\sqrt{a + b \tan(c + dx)}}{d} + \frac{2A(a + b \tan(c + dx))^{3/2}}{3d} + \frac{2B(a + b \tan(c + dx))^{5/2}}{5bd} \\
& \downarrow 3042 \\
& \frac{1}{2}(a + ib)^2(-B + iA) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx - \frac{1}{2}(a - ib)^2(B + iA) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \\
& \quad \frac{2(aA - bB)\sqrt{a + b \tan(c + dx)}}{d} + \frac{2A(a + b \tan(c + dx))^{3/2}}{3d} + \frac{2B(a + b \tan(c + dx))^{5/2}}{5bd} \\
& \downarrow 4020 \\
& \frac{i(a - ib)^2(B + iA) \int -\frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} - \\
& \frac{i(a + ib)^2(-B + iA) \int -\frac{1}{(i \tan(c + dx) + 1)\sqrt{a + b \tan(c + dx)}} d(-i \tan(c + dx))}{2d} + \\
& \quad \frac{2(aA - bB)\sqrt{a + b \tan(c + dx)}}{d} + \frac{2A(a + b \tan(c + dx))^{3/2}}{3d} + \frac{2B(a + b \tan(c + dx))^{5/2}}{5bd}
\end{aligned}$$

$$\begin{aligned}
& \downarrow 25 \\
& \frac{i(a-ib)^2(B+iA) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \\
& \frac{i(a+ib)^2(-B+iA) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \\
& \frac{2(aA-bB)\sqrt{a+b \tan(c+dx)}}{d} + \frac{2A(a+b \tan(c+dx))^{3/2}}{3d} + \frac{2B(a+b \tan(c+dx))^{5/2}}{5bd} \\
& \downarrow 73 \\
& \frac{(a+ib)^2(-B+iA) \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} - \\
& \frac{(a-ib)^2(B+iA) \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \frac{2(aA-bB)\sqrt{a+b \tan(c+dx)}}{d} + \\
& \frac{2A(a+b \tan(c+dx))^{3/2}}{3d} + \frac{2B(a+b \tan(c+dx))^{5/2}}{5bd} \\
& \downarrow 221 \\
& -\frac{(a-ib)^{3/2}(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{(a+ib)^{3/2}(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} + \\
& \frac{2(aA-bB)\sqrt{a+b \tan(c+dx)}}{d} + \frac{2A(a+b \tan(c+dx))^{3/2}}{3d} + \frac{2B(a+b \tan(c+dx))^{5/2}}{5bd}
\end{aligned}$$

input `Int[Tan[c + d*x]*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `-(((a - I*b)^(3/2)*(I*A + B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d) + ((a + I*b)^(3/2)*(I*A - B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d + (2*(a*A - b*B)*Sqrt[a + b*Tan[c + d*x]])/d + (2*A*(a + b*Tan[c + d*x])^(3/2))/(3*d) + (2*B*(a + b*Tan[c + d*x])^(5/2))/(5*b*d)`

3.326.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

- rule 221 $\text{Int}[(a + b(x^2)^{-1}), x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) \cdot \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] \text{ ; FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b]$
- rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$
- rule 4011 $\text{Int}[(a + b \tan(e + f(x)))^m \cdot (c + d \tan(e + f(x))) \cdot (f(x))], x_Symbol] \rightarrow \text{Simp}[d \cdot (a + b \tan[e + f*x])^m / (f*m), x] + \text{Int}[(a + b \tan[e + f*x])^{m-1} \cdot \text{Simp}[a*c - b*d + (b*c + a*d) \cdot \tan[e + f*x], x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[m, 0]$
- rule 4020 $\text{Int}[(a + b \tan(e + f(x)))^m \cdot (c + d \tan(e + f(x))) \cdot (f(x))], x_Symbol] \rightarrow \text{Simp}[c \cdot (d/f) \text{ Subst}[\text{Int}[(a + (b/d)x)^m / (d^2 + c*x), x], x, d \cdot \tan[e + f*x]], x] \text{ ; FreeQ}\{a, b, c, d, e, f, m, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[c^2 + d^2, 0]$
- rule 4022 $\text{Int}[(a + b \tan(e + f(x)))^m \cdot (c + d \tan(e + f(x))) \cdot (f(x))], x_Symbol] \rightarrow \text{Simp}[(c + I*d)/2 \text{ Int}[(a + b \tan[e + f*x])^m \cdot (1 - I \cdot \tan[e + f*x]), x], x] + \text{Simp}[(c - I*d)/2 \text{ Int}[(a + b \tan[e + f*x])^m \cdot (1 + I \cdot \tan[e + f*x]), x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, m, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0] \ \&\& \ \text{!IntegerQ}[m]$
- rule 4075 $\text{Int}[(a + b \tan(e + f(x)))^m \cdot (A + B \tan(e + f(x))) \cdot (c + d \tan(e + f(x))) \cdot (f(x))], x_Symbol] \rightarrow \text{Simp}[B \cdot d \cdot (a + b \tan[e + f*x])^{m+1} / (b*f*(m+1)), x] + \text{Int}[(a + b \tan[e + f*x])^m \cdot \text{Simp}[A*c - B*d + (B*c + A*d) \cdot \tan[e + f*x], x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, A, B, m, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{!LeQ}[m, -1]$

3.326.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1667 vs. $2(147) = 294$.

Time = 0.11 (sec) , antiderivative size = 1668, normalized size of antiderivative = 9.53

method	result	size
parts	Expression too large to display	1668
derivativedivides	Expression too large to display	1686
default	Expression too large to display	1686

```
input int(tan(d*x+c)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

```
output A*(2/3/d*(a+b*tan(d*x+c))^(3/2)+2/d*(a+b*tan(d*x+c))^(1/2)*a+1/4/d*ln(b*ta
n(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(
1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)-1/2/d*ln(b*tan(d*x+c)
+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(
2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2
*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-
2*a)^(1/2))*(a^2+b^2)-1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan
(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2
))*(a^2+b^2)^(1/2)*a+2/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(
d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2
))*a^2-1/4/d*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(
d*x+c)-a-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)+1/
2/d*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a
-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+1/d/(2*(a^2+b^2)^(1/2)-2
*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/
(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*(a^2+b^2)+1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2
)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b
^2)^(1/2)-2*a)^(1/2))*(a^2+b^2)^(1/2)*a-2/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*
arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^
2)^(1/2)-2*a)^(1/2))*a^2)+B*(2/5/b/d*(a+b*tan(d*x+c))^(5/2)-2*b*(a+b*ta...
```

3.326.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3102 vs. 2(141) = 282.

Time = 0.46 (sec) , antiderivative size = 3102, normalized size of antiderivative = 17.73

$$\int \tan(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm=
"fricas")
```

```
output -1/30*(15*b*d*sqrt(-(6*A*B*a^2*b - 2*A*B*b^3 - (A^2 - B^2)*a^3 + 3*(A^2 -
B^2)*a*b^2 + d^2*sqrt(-(4*A^2*B^2*a^6 + 12*(A^3*B - A*B^3)*a^5*b + 3*(3*A^
4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 - 40*(A^3*B - A*B^3)*a^3*b^3 - 6*(A^4 - 8*
A^2*B^2 + B^4)*a^2*b^4 + 12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2*B^2 + B^4
)*b^6)/d^4))/d^2)*log((2*(A^3*B + A*B^3)*a^5 + 3*(A^4 - B^4)*a^4*b - 4*(A^
3*B + A*B^3)*a^3*b^2 + 2*(A^4 - B^4)*a^2*b^3 - 6*(A^3*B + A*B^3)*a*b^4 - (
A^4 - B^4)*b^5)*sqrt(b*tan(d*x + c) + a) + ((B*a + A*b)*d^3*sqrt(-(4*A^2*B
^2*a^6 + 12*(A^3*B - A*B^3)*a^5*b + 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2
- 40*(A^3*B - A*B^3)*a^3*b^3 - 6*(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 + 12*(A^
3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2*B^2 + B^4)*b^6)/d^4) + (2*A^2*B*a^4 + (3
*A^3 - 5*A*B^2)*a^3*b - 3*(3*A^2*B - B^3)*a^2*b^2 - (A^3 - 7*A*B^2)*a*b^3
+ (A^2*B - B^3)*b^4)*d)*sqrt(-(6*A*B*a^2*b - 2*A*B*b^3 - (A^2 - B^2)*a^3 +
3*(A^2 - B^2)*a*b^2 + d^2*sqrt(-(4*A^2*B^2*a^6 + 12*(A^3*B - A*B^3)*a^5*b
+ 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 - 40*(A^3*B - A*B^3)*a^3*b^3 - 6
*(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 + 12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2
*B^2 + B^4)*b^6)/d^4))/d^2)) - 15*b*d*sqrt(-(6*A*B*a^2*b - 2*A*B*b^3 - (A^
2 - B^2)*a^3 + 3*(A^2 - B^2)*a*b^2 + d^2*sqrt(-(4*A^2*B^2*a^6 + 12*(A^3*B
- A*B^3)*a^5*b + 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 - 40*(A^3*B - A*B^
3)*a^3*b^3 - 6*(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 + 12*(A^3*B - A*B^3)*a*b^5
+ (A^4 - 2*A^2*B^2 + B^4)*b^6)/d^4))/d^2)*log((2*(A^3*B + A*B^3)*a^5 + ...
```

3.326.6 SymPy [F]

$$\int \tan(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx))(a + b \tan(c + dx))^{\frac{3}{2}} \tan(c + dx) dx$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)*tan(c + d*x), x)`

3.326.7 Maxima [F]

$$\int \tan(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{3}{2}} \tan(dx + c) dx$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)*tan(d*x + c), x)`

3.326.8 Giac [F(-1)]

Timed out.

$$\int \tan(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.326.9 Mupad [B] (verification not implemented)

Time = 34.89 (sec) , antiderivative size = 2868, normalized size of antiderivative = 16.39

$$\int \tan(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2),x)
```

```
output log((16*B^3*a*b^3*(a^2 + b^2)^2)/d^3 - (((16*b^2*((-B^4*b^2*d^4*(3*a^2 -
b^2)^2)^(1/2) - B^2*a^3*d^2 + 3*B^2*a*b^2*d^2)/d^4)^(1/2)*(B*b^3 + B*a^2*b
+ a*d*((-B^4*b^2*d^4*(3*a^2 - b^2)^2)^(1/2) - B^2*a^3*d^2 + 3*B^2*a*b^2*
d^2)/d^4)^(1/2)*(a + b*tan(c + d*x))^(1/2)))/d + (16*B^2*b^2*(a + b*tan(c
+ d*x))^(1/2)*(a^4 + b^4 - 6*a^2*b^2))/d^2)*((( -B^4*b^2*d^4*(3*a^2 - b^2)^
2)^(1/2) - B^2*a^3*d^2 + 3*B^2*a*b^2*d^2)/d^4)^(1/2))/2)*((6*B^4*a^2*b^4*d
^4 - B^4*b^6*d^4 - 9*B^4*a^4*b^2*d^4)^(1/2)/(4*d^4) - (B^2*a^3)/(4*d^2) +
(3*B^2*a*b^2)/(4*d^2))^(1/2) - log((16*B^3*a*b^3*(a^2 + b^2)^2)/d^3 - (((1
6*b^2*((-B^4*b^2*d^4*(3*a^2 - b^2)^2)^(1/2) + B^2*a^3*d^2 - 3*B^2*a*b^2*
d^2)/d^4)^(1/2)*(B*b^3 + B*a^2*b - a*d*((-B^4*b^2*d^4*(3*a^2 - b^2)^2)^(
1/2) + B^2*a^3*d^2 - 3*B^2*a*b^2*d^2)/d^4)^(1/2)*(a + b*tan(c + d*x))^(1/2
)))/d - (16*B^2*b^2*(a + b*tan(c + d*x))^(1/2)*(a^4 + b^4 - 6*a^2*b^2))/d^
2)*((( -B^4*b^2*d^4*(3*a^2 - b^2)^2)^(1/2) + B^2*a^3*d^2 - 3*B^2*a*b^2*d^
2)/d^4)^(1/2))/2)*(((6*B^4*a^2*b^4*d^4 - B^4*b^6*d^4 - 9*B^4*a^4*b^2*d^4)^(
1/2) + B^2*a^3*d^2 - 3*B^2*a*b^2*d^2)/(4*d^4))^(1/2) - log((16*B^3*a*b^3*
(a^2 + b^2)^2)/d^3 - (((16*b^2*((-B^4*b^2*d^4*(3*a^2 - b^2)^2)^(1/2) - B^
2*a^3*d^2 + 3*B^2*a*b^2*d^2)/d^4)^(1/2)*(B*b^3 + B*a^2*b - a*d*((-B^4*b^2
*d^4*(3*a^2 - b^2)^2)^(1/2) - B^2*a^3*d^2 + 3*B^2*a*b^2*d^2)/d^4)^(1/2)*(a
+ b*tan(c + d*x))^(1/2)))/d - (16*B^2*b^2*(a + b*tan(c + d*x))^(1/2)*(a^4
+ b^4 - 6*a^2*b^2))/d^2)*((( -B^4*b^2*d^4*(3*a^2 - b^2)^2)^(1/2) - B^2*...
```

3.327 $\int (a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx$

3.327.1 Optimal result	3100
3.327.2 Mathematica [A] (verified)	3100
3.327.3 Rubi [A] (warning: unable to verify)	3101
3.327.4 Maple [B] (verified)	3104
3.327.5 Fricas [B] (verification not implemented)	3105
3.327.6 Sympy [F]	3105
3.327.7 Maxima [F(-2)]	3106
3.327.8 Giac [F(-1)]	3106
3.327.9 Mupad [B] (verification not implemented)	3106

3.327.1 Optimal result

Integrand size = 25, antiderivative size = 150

$$\int (a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx =$$

$$-\frac{(a - ib)^{3/2} (iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}}\right)}{d}$$

$$+ \frac{(a + ib)^{3/2} (iA - B) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a + ib}}\right)}{d}$$

$$+ \frac{2(Ab + aB) \sqrt{a + b \tan(c + dx)}}{d} + \frac{2B(a + b \tan(c + dx))^{3/2}}{3d}$$

output $-(a-I*b)^{(3/2)}*(I*A+B)*\operatorname{arctanh}((a+b*\tan(d*x+c))^{(1/2)}/(a-I*b)^{(1/2)})/d+(a+I*b)^{(3/2)}*(I*A-B)*\operatorname{arctanh}((a+b*\tan(d*x+c))^{(1/2)}/(a+I*b)^{(1/2)})/d+2*(A*b+B*a)*(a+b*\tan(d*x+c))^{(1/2)}/d+2/3*B*(a+b*\tan(d*x+c))^{(3/2)}/d$

3.327.2 Mathematica [A] (verified)

Time = 0.55 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.93

$$\int (a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx =$$

$$\frac{-3i(a - ib)^{3/2} (A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}}\right) + 3i(a + ib)^{3/2} (A + iB) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a + ib}}\right)}{3d}$$

input `Integrate[(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `((-3*I)*(a - I*b)^(3/2)*(A - I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] + (3*I)*(a + I*b)^(3/2)*(A + I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + 2*Sqrt[a + b*Tan[c + d*x]]*(3*A*b + 4*a*B + b*B*Tan[c + d*x]))/(3*d)`

3.327.3 Rubi [A] (warning: unable to verify)

Time = 0.76 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.87, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.440$, Rules used = {3042, 4011, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4011} \\
 & \int \sqrt{a + b \tan(c + dx)} (aA - bB + (Ab + aB) \tan(c + dx)) dx + \frac{2B(a + b \tan(c + dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{a + b \tan(c + dx)} (aA - bB + (Ab + aB) \tan(c + dx)) dx + \frac{2B(a + b \tan(c + dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{4011} \\
 & \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2(aB + Ab) \sqrt{a + b \tan(c + dx)}}{d} + \\
 & \quad \frac{2B(a + b \tan(c + dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2(aB + Ab) \sqrt{a + b \tan(c + dx)}}{d} + \\
 & \quad \frac{2B(a + b \tan(c + dx))^{3/2}}{3d}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 4022 \\
& \frac{1}{2}(a-ib)^2(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}(a+ib)^2(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \\
& \quad \frac{2(aB+Ab)\sqrt{a+b \tan(c+dx)}}{d} + \frac{2B(a+b \tan(c+dx))^{3/2}}{3d} \\
& \downarrow 3042 \\
& \frac{1}{2}(a-ib)^2(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}(a+ib)^2(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \\
& \quad \frac{2(aB+Ab)\sqrt{a+b \tan(c+dx)}}{d} + \frac{2B(a+b \tan(c+dx))^{3/2}}{3d} \\
& \downarrow 4020 \\
& \frac{i(a-ib)^2(A-iB) \int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} - \\
& \frac{i(a+ib)^2(A+iB) \int -\frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \\
& \quad \frac{2(aB+Ab)\sqrt{a+b \tan(c+dx)}}{d} + \frac{2B(a+b \tan(c+dx))^{3/2}}{3d} \\
& \downarrow 25 \\
& \frac{i(a-ib)^2(A-iB) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \\
& \frac{i(a+ib)^2(A+iB) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \\
& \quad \frac{2(aB+Ab)\sqrt{a+b \tan(c+dx)}}{d} + \frac{2B(a+b \tan(c+dx))^{3/2}}{3d} \\
& \downarrow 73 \\
& \frac{(a-ib)^2(A-iB) \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \\
& \frac{(a+ib)^2(A+iB) \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \frac{2(aB+Ab)\sqrt{a+b \tan(c+dx)}}{d} + \\
& \quad \frac{2B(a+b \tan(c+dx))^{3/2}}{3d} \\
& \downarrow 221 \\
& \frac{(a-ib)^{3/2}(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{(a+ib)^{3/2}(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} + \\
& \quad \frac{2(aB+Ab)\sqrt{a+b \tan(c+dx)}}{d} + \frac{2B(a+b \tan(c+dx))^{3/2}}{3d}
\end{aligned}$$

input `Int[(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `((a - I*b)^(3/2)*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d + ((a + I*b)^(3/2)*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d + (2*(A*b + a*B)*Sqrt[a + b*Tan[c + d*x]])/d + (2*B*(a + b*Tan[c + d*x])^(3/2))/(3*d)`

3.327.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

3.327.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1656 vs. $2(126) = 252$.

Time = 0.09 (sec) , antiderivative size = 1657, normalized size of antiderivative = 11.05

method	result	size
parts	Expression too large to display	1657
derivativedivides	Expression too large to display	1665
default	Expression too large to display	1665

```
input int((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output A*(2*b*(a+b*tan(d*x+c))^(1/2)/d-1/4/b/d*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))
^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(a^2+b^2)^(1/2)*(2*(
a^2+b^2)^(1/2)+2*a)^(1/2)*a+1/4/b/d*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/
2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(
1/2)*a^2-1/4*b/d*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1
/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b/d/(2*(a^2+
b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+
2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*(a^2+b^2)^(1/2)+2*b/d/(2*(a^2+b
^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2
*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a+1/4/d/b*ln((a+b*tan(d*x+c))^(1
/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*(2*(a^2+
b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*a-1/4/d/b*ln((a+b*tan(d*x+c))^(1/2)*
(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*(2*(a^2+b^2)
^(1/2)+2*a)^(1/2)*a^2+1/4/d*b*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)
+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+
1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-
2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*(a^2+b^2)^(1/2)-2
/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2
*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a)+B*(2/3/d*(a+b*t
an(d*x+c))^(3/2)+2/d*(a+b*tan(d*x+c))^(1/2)*a+1/4/d*ln(b*tan(d*x+c)+a+(...
```

3.327.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3058 vs. $2(120) = 240$.

Time = 0.47 (sec) , antiderivative size = 3058, normalized size of antiderivative = 20.39

$$\int (a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output

```
1/6*(3*d*sqrt((6*A*B*a^2*b - 2*A*B*b^3 - (A^2 - B^2)*a^3 + 3*(A^2 - B^2)*a
*b^2 + d^2*sqrt(-(4*A^2*B^2*a^6 + 12*(A^3*B - A*B^3)*a^5*b + 3*(3*A^4 - 14
*A^2*B^2 + 3*B^4)*a^4*b^2 - 40*(A^3*B - A*B^3)*a^3*b^3 - 6*(A^4 - 8*A^2*B^
2 + B^4)*a^2*b^4 + 12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2*B^2 + B^4)*b^6)
/d^4))/d^2)*log((2*(A^3*B + A*B^3)*a^5 + 3*(A^4 - B^4)*a^4*b - 4*(A^3*B +
A*B^3)*a^3*b^2 + 2*(A^4 - B^4)*a^2*b^3 - 6*(A^3*B + A*B^3)*a*b^4 - (A^4 -
B^4)*b^5)*sqrt(b*tan(d*x + c) + a) + ((A*a - B*b)*d^3*sqrt(-(4*A^2*B^2*a^6
+ 12*(A^3*B - A*B^3)*a^5*b + 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 - 40*
(A^3*B - A*B^3)*a^3*b^3 - 6*(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 + 12*(A^3*B -
A*B^3)*a*b^5 + (A^4 - 2*A^2*B^2 + B^4)*b^6)/d^4) - (2*A*B^2*a^4 + (5*A^2*B
- 3*B^3)*a^3*b + 3*(A^3 - 3*A*B^2)*a^2*b^2 - (7*A^2*B - B^3)*a*b^3 - (A^3
- A*B^2)*b^4)*d)*sqrt((6*A*B*a^2*b - 2*A*B*b^3 - (A^2 - B^2)*a^3 + 3*(A^2
- B^2)*a*b^2 + d^2*sqrt(-(4*A^2*B^2*a^6 + 12*(A^3*B - A*B^3)*a^5*b + 3*(3
*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 - 40*(A^3*B - A*B^3)*a^3*b^3 - 6*(A^4 -
8*A^2*B^2 + B^4)*a^2*b^4 + 12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2*B^2 +
B^4)*b^6)/d^4))/d^2)) - 3*d*sqrt((6*A*B*a^2*b - 2*A*B*b^3 - (A^2 - B^2)*a^
3 + 3*(A^2 - B^2)*a*b^2 + d^2*sqrt(-(4*A^2*B^2*a^6 + 12*(A^3*B - A*B^3)*a^
5*b + 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 - 40*(A^3*B - A*B^3)*a^3*b^3
- 6*(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 + 12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*
A^2*B^2 + B^4)*b^6)/d^4))/d^2)*log((2*(A^3*B + A*B^3)*a^5 + 3*(A^4 - B^...
```

3.327.6 Sympy [F]

$$\int (a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx)) (a + b \tan(c + dx))^{\frac{3}{2}} dx$$

input `integrate((a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2), x)`

3.327.7 Maxima [F(-2)]

Exception generated.

$$\int (a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is`

3.327.8 Giac [F(-1)]

Timed out.

$$\int (a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.327.9 Mupad [B] (verification not implemented)

Time = 20.84 (sec) , antiderivative size = 2823, normalized size of antiderivative = 18.82

$$\int (a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2),x)`

output $\log\left(\frac{(16b^2((-A^4b^2d^4(3a^2 - b^2)^2)^{1/2} - A^2a^3d^2 + 3A^2ab^2d^2)/d^4)^{1/2}(Ab^3 + Aa^2b - a*d*((-A^4b^2d^4(3a^2 - b^2)^2)^{1/2} - A^2a^3d^2 + 3A^2ab^2d^2)/d^4)^{1/2}(a + b*\tan(c + dx))^{1/2}}{d} - \frac{(16A^2b^2(a + b*\tan(c + dx))^{1/2}(a^4 + b^4 - 6a^2b^2))/d^2 * ((-A^4b^2d^4(3a^2 - b^2)^2)^{1/2} - A^2a^3d^2 + 3A^2ab^2d^2)/d^4)^{1/2}}{2} - \frac{(16A^3ab^3(a^2 + b^2)^2)/d^3 * ((6A^4a^2b^4d^4 - A^4b^6d^4 - 9A^4a^4b^2d^4)^{1/2}/(4d^4) - (A^2a^3)/(4d^2) + (3A^2ab^2)/(4d^2))^{1/2}}{2} - \log\left(\frac{(16b^2((-A^4b^2d^4(3a^2 - b^2)^2)^{1/2} - A^2a^3d^2 + 3A^2ab^2d^2)/d^4)^{1/2}(Ab^3 + Aa^2b + a*d*((-A^4b^2d^4(3a^2 - b^2)^2)^{1/2} - A^2a^3d^2 + 3A^2ab^2d^2)/d^4)^{1/2}(a + b*\tan(c + dx))^{1/2}}{d} + \frac{(16A^2b^2(a + b*\tan(c + dx))^{1/2}(a^4 + b^4 - 6a^2b^2))/d^2 * ((-A^4b^2d^4(3a^2 - b^2)^2)^{1/2} - A^2a^3d^2 + 3A^2ab^2d^2)/d^4)^{1/2}}{2} - \frac{(16A^3ab^3(a^2 + b^2)^2)/d^3 * ((6A^4a^2b^4d^4 - A^4b^6d^4 - 9A^4a^4b^2d^4)^{1/2} - A^2a^3d^2 + 3A^2ab^2d^2)/(4d^4))^{1/2}}{2} - \log\left(\frac{(16b^2(-((-A^4b^2d^4(3a^2 - b^2)^2)^{1/2} + A^2a^3d^2 - 3A^2ab^2d^2)/d^4)^{1/2}(Ab^3 + Aa^2b + a*d*(-((-A^4b^2d^4(3a^2 - b^2)^2)^{1/2} + A^2a^3d^2 - 3A^2ab^2d^2)/d^4)^{1/2}(a + b*\tan(c + dx))^{1/2}}{d} + \frac{(16A^2b^2(a + b*\tan(c + dx))^{1/2}(a^4 + b^4 - 6a^2b^2))/d^2 * (-((-A^4b^2d^4(3a^2 - b^2)^2)^{1/2} + A^2a^3d^2 - 3A^2ab^2d^2)/d^4)^{1/2}}{2} \dots$

3.328 $\int \cot(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.328.1 Optimal result	3108
3.328.2 Mathematica [A] (verified)	3109
3.328.3 Rubi [A] (warning: unable to verify)	3109
3.328.4 Maple [B] (verified)	3114
3.328.5 Fricas [B] (verification not implemented)	3115
3.328.6 Sympy [F]	3115
3.328.7 Maxima [F]	3115
3.328.8 Giac [F(-1)]	3116
3.328.9 Mupad [B] (verification not implemented)	3116

3.328.1 Optimal result

Integrand size = 31, antiderivative size = 152

$$\int \cot(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx =$$

$$-\frac{2a^{3/2}A \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{d} + \frac{(a-ib)^{3/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d}$$

$$+ \frac{(a+ib)^{3/2}(A+iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d} + \frac{2bB\sqrt{a+b \tan(c+dx)}}{d}$$

output

```
-2*a^(3/2)*A*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/d+(a-I*b)^(3/2)*(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d+(a+I*b)^(3/2)*(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d+2*b*B*(a+b*tan(d*x+c))^(1/2)/d
```

3.328.2 Mathematica [A] (verified)

Time = 0.41 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.95

$$\int \cot(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{-2a^{3/2} A \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right) + (a - ib)^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right) + (a + B \tan(c+dx))}{d}$$

input `Integrate[Cot[c + d*x]*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`output `(-2*a^(3/2)*A*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]] + (a - I*b)^(3/2)*(A - I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] + (a + I*b)^(3/2)*(A + I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + 2*b*B*Sqrt[a + b*Tan[c + d*x]])/d`**3.328.3 Rubi [A] (warning: unable to verify)**Time = 1.30 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.91, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 4090, 27, 3042, 4136, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan(c + dx)} dx \\ & \quad \downarrow \text{4090} \\ & 2 \int \frac{\cot(c + dx) (Aa^2 + b(Ab + 2aB) \tan^2(c + dx) + (Ba^2 + 2Aba - b^2B) \tan(c + dx))}{\frac{2\sqrt{a + b \tan(c + dx)}}{2bB\sqrt{a + b \tan(c + dx)}}} dx + \\ & \quad \downarrow \text{27} \end{aligned}$$

$$\begin{aligned}
& \int \frac{\cot(c+dx) (Aa^2 + b(Ab + 2aB) \tan^2(c+dx) + (Ba^2 + 2Aba - b^2B) \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx + \\
& \quad \frac{2bB \sqrt{a+b \tan(c+dx)}}{d} \\
& \quad \downarrow \text{3042} \\
& \int \frac{Aa^2 + b(Ab + 2aB) \tan(c+dx)^2 + (Ba^2 + 2Aba - b^2B) \tan(c+dx)}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + \\
& \quad \frac{2bB \sqrt{a+b \tan(c+dx)}}{d} \\
& \quad \downarrow \text{4136} \\
& \int \frac{Ba^2 + 2Aba - b^2B - (Aa^2 - 2bBa - Ab^2) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \\
& a^2 A \int \frac{\cot(c+dx) (\tan^2(c+dx) + 1)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{2bB \sqrt{a+b \tan(c+dx)}}{d} \\
& \quad \downarrow \text{3042} \\
& \int \frac{Ba^2 + 2Aba - b^2B - (Aa^2 - 2bBa - Ab^2) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \\
& a^2 A \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + \frac{2bB \sqrt{a+b \tan(c+dx)}}{d} \\
& \quad \downarrow \text{4022} \\
& a^2 A \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2} (a+ib)^2 (-B+iA) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \\
& \quad \frac{1}{2} (a-ib)^2 (B+iA) \int \frac{i \tan(c+dx) + 1}{\sqrt{a+b \tan(c+dx)}} dx + \frac{2bB \sqrt{a+b \tan(c+dx)}}{d} \\
& \quad \downarrow \text{3042} \\
& a^2 A \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2} (a+ib)^2 (-B+iA) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \\
& \quad \frac{1}{2} (a-ib)^2 (B+iA) \int \frac{i \tan(c+dx) + 1}{\sqrt{a+b \tan(c+dx)}} dx + \frac{2bB \sqrt{a+b \tan(c+dx)}}{d} \\
& \quad \downarrow \text{4020} \\
& a^2 A \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + \\
& \quad \frac{i(a-ib)^2 (B+iA) \int -\frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \\
& \quad \frac{i(a+ib)^2 (-B+iA) \int -\frac{1}{(i \tan(c+dx)+1) \sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \frac{2bB \sqrt{a+b \tan(c+dx)}}{d}
\end{aligned}$$

3.328. $\int \cot(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \downarrow 25 \\
& \frac{a^2 A \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - i(a-ib)^2(B+iA) \int \frac{1}{(1-i\tan(c+dx))\sqrt{a+b\tan(c+dx)}} d(i\tan(c+dx))}{2d} - \\
& \frac{i(a+ib)^2(-B+iA) \int \frac{1}{(i\tan(c+dx)+1)\sqrt{a+b\tan(c+dx)}} d(-i\tan(c+dx))}{2d} + \frac{2bB\sqrt{a+b\tan(c+dx)}}{d} \\
& \downarrow 73 \\
& \frac{a^2 A \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - (a+ib)^2(-B+iA) \int \frac{1}{-\frac{i\tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b\tan(c+dx)}}{bd} + \\
& \frac{(a-ib)^2(B+iA) \int \frac{1}{\frac{i\tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b\tan(c+dx)}}{bd} + \frac{2bB\sqrt{a+b\tan(c+dx)}}{d} \\
& \downarrow 221 \\
& a^2 A \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx + \frac{(a-ib)^{3/2}(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} - \\
& \frac{(a+ib)^{3/2}(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} + \frac{2bB\sqrt{a+b\tan(c+dx)}}{d} \\
& \downarrow 4117 \\
& \frac{a^2 A \int \frac{\cot(c+dx)}{\sqrt{a+b\tan(c+dx)}} d\tan(c+dx)}{d} + \frac{(a-ib)^{3/2}(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} - \\
& \frac{(a+ib)^{3/2}(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} + \frac{2bB\sqrt{a+b\tan(c+dx)}}{d} \\
& \downarrow 73 \\
& \frac{2a^2 A \int \frac{1}{\frac{a+b\tan(c+dx)}{b} - \frac{a}{b}} d\sqrt{a+b\tan(c+dx)}}{bd} + \frac{(a-ib)^{3/2}(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} - \\
& \frac{(a+ib)^{3/2}(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} + \frac{2bB\sqrt{a+b\tan(c+dx)}}{d} \\
& \downarrow 221 \\
& -\frac{2a^{3/2} A \operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a}}\right)}{d} + \frac{(a-ib)^{3/2}(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} - \\
& \frac{(a+ib)^{3/2}(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} + \frac{2bB\sqrt{a+b\tan(c+dx)}}{d}
\end{aligned}$$

input `Int[Cot[c + d*x]*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `((a - I*b)^(3/2)*(I*A + B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d - ((a + I*b)^(3/2)*(I*A - B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d - (2*a^(3/2)*A*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/d + (2*b*B*Sqrt[a + b*Tan[c + d*x]])/d`

3.328.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n)), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4136 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.328.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1653 vs. $2(126) = 252$.

Time = 0.24 (sec) , antiderivative size = 1654, normalized size of antiderivative = 10.88

method	result	size
derivativdivides	Expression too large to display	1654
default	Expression too large to display	1654

```
input int(cot(d*x+c)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

```
output 2*b*B*(a+b*tan(d*x+c))^(1/2)/d-1/d*b^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arcta
n(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1
/2)-2*a)^(1/2))*A+1/d*b^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan
(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2
))*A+1/4/d*b*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan
(d*x+c)-a-(a^2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/4/d*b*ln(b*ta
n(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(
1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/4/d*b*ln((a+b*tan(d*x+c))^(1/2)*(
2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*B*(2*(a^2+b^2
)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*a-1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arc
tan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(
1/2)-2*a)^(1/2))*A*(a^2+b^2)^(1/2)*a+1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*ar
ctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)
^(1/2)-2*a)^(1/2))*A*(a^2+b^2)^(1/2)*a-1/4/d*b*ln(b*tan(d*x+c)+a+(a+b*tan(
d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*B*(2*(a^2+b^2
)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*a-1/4/d*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+
c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1
/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)+1/2/d*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1
/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*
a)^(1/2)*a+1/4/d*b*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)...
```

3.328.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3075 vs. $2(120) = 240$.

Time = 2.67 (sec) , antiderivative size = 6165, normalized size of antiderivative = 40.56

$$\int \cot(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.328.6 Sympy [F]

$$\int \cot(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx))(a + b \tan(c + dx))^{\frac{3}{2}} \cot(c + dx) dx$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)*cot(c + d*x), x)`

3.328.7 Maxima [F]

$$\int \cot(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{3}{2}} \cot(dx + c) dx$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)*cot(d*x + c), x)`

3.328. $\int \cot(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.328.8 Giac [F(-1)]

Timed out.

$$\int \cot(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output Timed out

3.328.9 Mupad [B] (verification not implemented)

Time = 11.80 (sec) , antiderivative size = 20255, normalized size of antiderivative = 133.26

$$\int \cot(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2),x)`

output
$$\begin{aligned} & (2*B*b*(a + b*\tan(c + d*x))^{(1/2)})/d - \operatorname{atan}\left(\frac{(32*(4*B*a*b^{11}*d^4 + 12*A*a^2*b^{10}*d^4 + 12*A*a^4*b^8*d^4 + 4*B*a^3*b^9*d^4))/d^5 - (32*(16*b^{10}*d^4 + 24*a^2*b^8*d^4)*(a + b*\tan(c + d*x))^{(1/2)} - ((8*A^2*a^3*d^2 - 8*B^2*a^3*d^2 + 16*A*B*b^3*d^2 - 24*A^2*a*b^2*d^2 + 24*B^2*a*b^2*d^2 - 48*A*B*a^2*b*d^2)^2/64 - d^4*(A^4*a^6 + A^4*b^6 + B^4*a^6 + B^4*b^6 + 2*A^2*B^2*a^6 + 2*A^2*B^2*b^6 + 3*A^4*a^2*b^4 + 3*A^4*a^4*b^2 + 3*B^4*a^2*b^4 + 3*B^4*a^4*b^2 + 6*A^2*B^2*a^2*b^4 + 6*A^2*B^2*a^4*b^2))^{(1/2)} - A^2*a^3*d^2 + B^2*a^3*d^2 - 2*A*B*b^3*d^2 + 3*A^2*a*b^2*d^2 - 3*B^2*a*b^2*d^2 + 6*A*B*a^2*b*d^2)/(4*d^4))^{(1/2)}}{d^4} * (-((8*A^2*a^3*d^2 - 8*B^2*a^3*d^2 + 16*A*B*b^3*d^2 - 24*A^2*a*b^2*d^2 + 24*B^2*a*b^2*d^2 - 48*A*B*a^2*b*d^2)^2/64 - d^4*(A^4*a^6 + A^4*b^6 + B^4*a^6 + B^4*b^6 + 2*A^2*B^2*a^6 + 2*A^2*B^2*b^6 + 3*A^4*a^2*b^4 + 3*A^4*a^4*b^2 + 3*B^4*a^2*b^4 + 3*B^4*a^4*b^2 + 6*A^2*B^2*a^2*b^4 + 6*A^2*B^2*a^4*b^2))^{(1/2)} - A^2*a^3*d^2 + B^2*a^3*d^2 - 2*A*B*b^3*d^2 + 3*A^2*a*b^2*d^2 - 3*B^2*a*b^2*d^2 + 6*A*B*a^2*b*d^2)/(4*d^4))^{(1/2)}\right) \\ & - (32*(a + b*\tan(c + d*x))^{(1/2)}*(28*A^2*a^3*b^{10}*d^2 - 18*A^2*a^5*b^8*d^2 - 28*B^2*a^3*b^{10}*d^2 + 10*B^2*a^5*b^8*d^2 - 16*A*B*b^{13}*d^2 + 22*A^2*a*b^{12}*d^2 - 22*B^2*a*b^{12}*d^2 + 16*A*B*a^2*b^{11}*d^2 + 64*A*B*a^4*b^9*d^2))/d^4 * (-((8*A^2*a^3*d^2 - 8*B^2*a^3*d^2 + 16*A*B*b^3*d^2 - 24*A^2*a*b^2*d^2 + 24*B^2*a*b^2*d^2 - 48*A*B*a^2*b*d^2)^2/64 - d^4*(A^4*a^6 + A^4*b^6 + B^4*a^6 + B^4*b^6 + 2*A^2*B^2*a^6 + 2*A^2*B^2*b^6 + 3*A^4*a^2*b^4 + 3*A... \end{aligned}$$

3.329 $\int \cot^2(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.329.1 Optimal result	3117
3.329.2 Mathematica [A] (verified)	3118
3.329.3 Rubi [A] (warning: unable to verify)	3118
3.329.4 Maple [B] (verified)	3123
3.329.5 Fricas [B] (verification not implemented)	3124
3.329.6 Sympy [F]	3125
3.329.7 Maxima [F]	3125
3.329.8 Giac [F(-1)]	3125
3.329.9 Mupad [B] (verification not implemented)	3126

3.329.1 Optimal result

Integrand size = 33, antiderivative size = 169

$$\int \cot^2(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx =$$

$$\frac{\sqrt{a}(3Ab+2aB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{d}$$

$$+ \frac{(a-ib)^{3/2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d}$$

$$- \frac{(a+ib)^{3/2}(iA-B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d} - \frac{aA \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{d}$$

```
output (a-I*b)^(3/2)*(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d-(a+I
*b)^(3/2)*(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d-(3*A*b+2
*B*a)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))*a^(1/2)/d-a*A*cot(d*x+c)*(a+
b*tan(d*x+c))^(1/2)/d
```

3.329.2 Mathematica [A] (verified)

Time = 0.64 (sec) , antiderivative size = 282, normalized size of antiderivative = 1.67

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{-\sqrt{a}(3Ab + 2aB)\operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a}}\right) + (a - ib)^{3/2}(iA + B)\operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a-ib}}\right)}{d}$$

input `Integrate[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(-(Sqrt[a]*(3*A*b + 2*a*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]]) + (a - I*b)^(3/2)*(I*A + B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] - I*a*A*Sqrt[a + I*b]*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + A*Sqrt[a + I*b]*b*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + a*Sqrt[a + I*b]*B*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + I*Sqrt[a + I*b]*b*B*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] - a*A*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/d`

3.329.3 Rubi [A] (warning: unable to verify)

Time = 1.40 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.93, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 4088, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan(c + dx)^2} dx$$

$$\downarrow \text{4088}$$

$$\int \frac{\cot(c+dx) (-b(aA - 2bB) \tan^2(c+dx) - 2(Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(3Ab + 2aB))}{2\sqrt{a+b\tan(c+dx)} \frac{aA \cot(c+dx) \sqrt{a+b\tan(c+dx)}}{d}} dx -$$

↓ 27

$$\frac{1}{2} \int \frac{\cot(c+dx) (-b(aA - 2bB) \tan^2(c+dx) - 2(Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(3Ab + 2aB))}{\sqrt{a+b\tan(c+dx)} \frac{aA \cot(c+dx) \sqrt{a+b\tan(c+dx)}}{d}} dx -$$

↓ 3042

$$\frac{1}{2} \int \frac{-b(aA - 2bB) \tan(c+dx)^2 - 2(Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(3Ab + 2aB)}{\tan(c+dx) \sqrt{a+b\tan(c+dx)} \frac{aA \cot(c+dx) \sqrt{a+b\tan(c+dx)}}{d}} dx -$$

↓ 4136

$$\frac{1}{2} \left(\int -\frac{2(Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c+dx))}{\sqrt{a+b\tan(c+dx)} \frac{aA \cot(c+dx) \sqrt{a+b\tan(c+dx)}}{d}} dx + a(2aB + 3Ab) \int \frac{\cot(c+dx) (\tan^2(c+dx) + 1)}{\sqrt{a+b\tan(c+dx)}} dx - \right)$$

↓ 27

$$\frac{1}{2} \left(a(2aB + 3Ab) \int \frac{\cot(c+dx) (\tan^2(c+dx) + 1)}{\sqrt{a+b\tan(c+dx)} \frac{aA \cot(c+dx) \sqrt{a+b\tan(c+dx)}}{d}} dx - 2 \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx - \right)$$

↓ 3042

$$\frac{1}{2} \left(a(2aB + 3Ab) \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx) \sqrt{a+b\tan(c+dx)} \frac{aA \cot(c+dx) \sqrt{a+b\tan(c+dx)}}{d}} dx - 2 \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx - \right)$$

↓ 4022

$$\begin{aligned}
& -\frac{aA \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left(a(2aB+3Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{1}{2} (a-ib)^2 (A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2} (a-ib)^2 (A-iB) \int \frac{1}{\sqrt{a+b \tan(c+dx)}} dx \right) \right) \\
& \quad \downarrow \text{3042} \\
& -\frac{aA \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left(a(2aB+3Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{1}{2} (a-ib)^2 (A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2} (a-ib)^2 (A-iB) \int \frac{1}{\sqrt{a+b \tan(c+dx)}} dx \right) \right) \\
& \quad \downarrow \text{4020} \\
& -\frac{aA \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left(a(2aB+3Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{i(a-ib)^2 (A-iB) \int -\frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} dx}{2d} \right) \right) \\
& \quad \downarrow \text{25} \\
& -\frac{aA \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left(a(2aB+3Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{i(a+ib)^2 (A+iB) \int \frac{1}{(i \tan(c+dx)+1) \sqrt{a+b \tan(c+dx)}} dx}{2d} \right) \right) \\
& \quad \downarrow \text{73} \\
& -\frac{aA \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left(a(2aB+3Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{(a-ib)^2 (A-iB) \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} dx \sqrt{a+b \tan(c+dx)}}{bd} \right) \right) \\
& \quad \downarrow \text{221} \\
& -\frac{aA \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left(a(2aB+3Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{(a-ib)^{3/2} (A-iB) \arctan \left(\frac{\tan(c+dx)}{\sqrt{a-ib}} \right)}{d} + \frac{(a+ib)^{3/2}}{d} \right) \right) \\
& \quad \downarrow \text{4117}
\end{aligned}$$

$$\begin{aligned}
& -\frac{aA \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left(\frac{a(2aB+3Ab) \int \frac{\cot(c+dx)}{\sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{d} - 2 \left(\frac{(a-ib)^{3/2}(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{(a+ib)^{3/2}(A+iB)}{d} \right) \right) \\
& \quad \downarrow \text{73} \\
& -\frac{aA \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left(\frac{2a(2aB+3Ab) \int \frac{1}{\frac{a+b \tan(c+dx)}{b} - \frac{a}{b}} d \sqrt{a+b \tan(c+dx)}}{bd} - 2 \left(\frac{(a-ib)^{3/2}(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{(a+ib)^{3/2}(A+iB)}{d} \right) \right) \\
& \quad \downarrow \text{221} \\
& -\frac{aA \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \\
\frac{1}{2} & \left(-\frac{2\sqrt{a}(2aB+3Ab) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{d} - 2 \left(\frac{(a-ib)^{3/2}(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{(a+ib)^{3/2}(A+iB)}{d} \right) \right)
\end{aligned}$$

input `Int[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(-2*(((a - I*b)^(3/2)*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d + ((a + I*b)^(3/2)*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d) - (2*Sqrt[a]*(3*A*b + 2*a*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/d)/2 - (a*A*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/d`

3.329.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
 Q[u, x]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
 (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
 c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
 b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
 (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
 1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
 *(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
 - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`
- rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
 (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
 mp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x
])^n)/(d*f*(n + 1)*(c^2 + d^2)), x] - Simp[1/(d*(n + 1)*(c^2 + d^2))
 Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*
 (b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n
 + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[
 e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n
 + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] &&
 NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] &
 & LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2]), x_Symbol] :=
  Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
  FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2])/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.329.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1688 vs. $2(143) = 286$.

Time = 0.24 (sec) , antiderivative size = 1689, normalized size of antiderivative = 9.99

method	result	size
derivativedivides	Expression too large to display	1689
default	Expression too large to display	1689

```
input int(cot(d*x+c)^2*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

output
$$\begin{aligned}
& -1/4/d/b*\ln((a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*\tan(d*x+c)-a-(a^2+b^2)^(1/2))*A*(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-1 \\
& /d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*\tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*(a^2+b^2)^(1/2)*a+ \\
& 1/4/d/b*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*A*(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+1 \\
& /d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan((2*(a+b*\tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*(a^2+b^2)^(1/2)*a+1 \\
& /d*b^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan((2*(a+b*\tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B-1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*\tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*b^2-1/4/d*b*\ln((a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*\tan(d*x+c)-a-(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/4/d*b*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/4/d/b*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2-1/4/d*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*B*(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/2/d*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)...
\end{aligned}$$

3.329.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3116 vs. $2(137) = 274$.

Time = 3.34 (sec) , antiderivative size = 6248, normalized size of antiderivative = 36.97

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.329.6 Sympy [F]

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx))(a + b \tan(c + dx))^{\frac{3}{2}} \cot^2(c + dx) dx$$

input `integrate(cot(d*x+c)**2*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)*cot(c + d*x)**2, x)`

3.329.7 Maxima [F]

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{3}{2}} \cot(dx + c)^2 dx$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^2, x)`

3.329.8 Giac [F(-1)]

Timed out.

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.329.9 Mupad [B] (verification not implemented)

Time = 9.84 (sec) , antiderivative size = 21319, normalized size of antiderivative = 126.15

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2),x)
```

```
output (a^(1/2)*atan(((a^(1/2))*((16*(a + b*tan(c + d*x))^(1/2)*(2*A^4*b^16 + 2*B^4*b^16 + 4*A^2*B^2*b^16 - A^4*a^2*b^14 + 66*A^4*a^4*b^12 - A^4*a^6*b^10 + 2*A^4*a^8*b^8 + 8*B^4*a^2*b^14 + 16*B^4*a^4*b^12 - 16*B^4*a^6*b^10 + 6*B^4*a^8*b^8 + 25*A^2*B^2*a^2*b^14 - 130*A^2*B^2*a^4*b^12 + 145*A^2*B^2*a^6*b^10 + 12*A*B^3*a^3*b^13 - 104*A*B^3*a^5*b^11 + 44*A*B^3*a^7*b^9 - 84*A^3*B*a^3*b^13 + 144*A^3*B*a^5*b^11 - 12*A^3*B*a^7*b^9))/d^4 + (a^(1/2)*(3*A*b + 2*B*a))*((8*(100*A^3*a^2*b^13*d^2 + 44*A^3*a^4*b^11*d^2 - 56*A^3*a^6*b^9*d^2 - 92*B^3*a^3*b^12*d^2 - 84*B^3*a^5*b^10*d^2 + 12*B^3*a^7*b^8*d^2 + 4*B^3*a*b^14*d^2 - 92*A^2*B*a*b^14*d^2 - 216*A*B^2*a^2*b^13*d^2 - 48*A*B^2*a^4*b^11*d^2 + 168*A*B^2*a^6*b^9*d^2 + 208*A^2*B*a^3*b^12*d^2 + 264*A^2*B*a^5*b^10*d^2 - 36*A^2*B*a^7*b^8*d^2))/d^5 - (a^(1/2)*(3*A*b + 2*B*a))*((16*(a + b*tan(c + d*x))^(1/2)*(92*A^2*a^3*b^10*d^2 - 20*A^2*a^5*b^8*d^2 - 56*B^2*a^3*b^10*d^2 + 36*B^2*a^5*b^8*d^2 - 32*A*B*b^13*d^2 + 44*A^2*a*b^12*d^2 - 44*B^2*a*b^12*d^2 + 32*A*B*a^2*b^11*d^2 + 176*A*B*a^4*b^9*d^2))/d^4 + (a^(1/2)*(3*A*b + 2*B*a))*((8*(80*A*a*b^11*d^4 + 80*A*a^3*b^9*d^4 + 48*B*a^2*b^10*d^4 + 48*B*a^4*b^8*d^4))/d^5 - (8*a^(1/2)*(3*A*b + 2*B*a)*(32*b^10*d^4 + 48*a^2*b^8*d^4)*(a + b*tan(c + d*x))^(1/2))/d^5))/(2*d)))/(2*d)))/(2*d)))*(3*A*b + 2*B*a)*1i)/(2*d) + (a^(1/2))*((16*(a + b*tan(c + d*x))^(1/2)*(2*A^4*b^16 + 2*B^4*b^16 + 4*A^2*B^2*b^16 - A^4*a^2*b^14 + 66*A^4*a^4*b^12 - A^4*a^6*b^10 + 2*A^4*a^8*b^8 + 8*B^4*a^2*b^14 + 16*B^4*a^4*b^12 - 16*B...
```

3.330 $\int \cot^3(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.330.1 Optimal result	3127
3.330.2 Mathematica [A] (verified)	3128
3.330.3 Rubi [A] (warning: unable to verify)	3128
3.330.4 Maple [B] (verified)	3134
3.330.5 Fricas [B] (verification not implemented)	3135
3.330.6 Sympy [F]	3136
3.330.7 Maxima [F]	3136
3.330.8 Giac [F(-1)]	3136
3.330.9 Mupad [B] (verification not implemented)	3137

3.330.1 Optimal result

Integrand size = 33, antiderivative size = 219

$$\int \cot^3(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{(8a^2A - 3Ab^2 - 12abB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right) + B \tan(c+dx)}{4\sqrt{ad}}$$

$$- \frac{(a-ib)^{3/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d}$$

$$- \frac{(a+ib)^{3/2}(A+iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d}$$

$$- \frac{(5Ab+4aB) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{4d}$$

$$- \frac{aA \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d}$$

output

```
-(a-I*b)^(3/2)*(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d-(a+
I*b)^(3/2)*(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d+1/4*(8*
A*a^2-3*A*b^2-12*B*a*b)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/d/a^(1/2)-
1/4*(5*A*b+4*B*a)*cot(d*x+c)*(a+b*tan(d*x+c))^(1/2)/d-1/2*a*A*cot(d*x+c)^2
*(a+b*tan(d*x+c))^(1/2)/d
```

3.330.2 Mathematica [A] (verified)

Time = 2.70 (sec) , antiderivative size = 195, normalized size of antiderivative = 0.89

$$\int \cot^3(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx = \frac{(8a^2A - 3Ab^2 - 12abB) \operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a}}\right) - \sqrt{a}\left(4(a-ib)^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a}}\right) + 4(a+ib)^{3/2}(A+iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a}}\right)\right)}{4\sqrt{a}d}$$

input `Integrate[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `((8*a^2*A - 3*A*b^2 - 12*a*b*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]] - Sqrt[a]*(4*(a - I*b)^(3/2)*(A - I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] + 4*(a + I*b)^(3/2)*(A + I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]]) + Cot[c + d*x]*(5*A*b + 4*a*B + 2*a*A*Cot[c + d*x])*Sqrt[a + b*Tan[c + d*x]])/(4*Sqrt[a]*d)`

3.330.3 Rubi [A] (warning: unable to verify)

Time = 1.85 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.00, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 4088, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^3(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\tan(c+dx)^3} dx$$

$$\downarrow \text{4088}$$

$$\frac{1}{2} \int \frac{\cot^2(c+dx) (-b(3aA - 4bB) \tan^2(c+dx) - 4(Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(5Ab + 4aB))}{2\sqrt{a+b\tan(c+dx)} \left(\frac{aA \cot^2(c+dx) \sqrt{a+b\tan(c+dx)}}{2d} \right)} dx -$$

$$\downarrow 27$$

$$\frac{1}{4} \int \frac{\cot^2(c+dx) (-b(3aA - 4bB) \tan^2(c+dx) - 4(Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(5Ab + 4aB))}{\sqrt{a+b \tan(c+dx)}} dx - \frac{aA \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d}$$

$$\downarrow 3042$$

$$\frac{1}{4} \int \frac{-b(3aA - 4bB) \tan(c+dx)^2 - 4(Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(5Ab + 4aB)}{\tan(c+dx)^2 \sqrt{a+b \tan(c+dx)}} dx - \frac{aA \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d}$$

$$\downarrow 4132$$

$$\frac{1}{4} \left(\frac{\int \frac{\cot(c+dx)(ab(5Ab+4aB) \tan^2(c+dx) + 8a(Ba^2+2Aba-b^2B) \tan(c+dx) + a(8Aa^2-12bBa-3Ab^2))}{2\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{(4aB + 5Ab) \cot(c+dx)}{a} \right) - \frac{aA \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d}$$

$$\downarrow 27$$

$$\frac{1}{4} \left(\frac{\int \frac{\cot(c+dx)(ab(5Ab+4aB) \tan^2(c+dx) + 8a(Ba^2+2Aba-b^2B) \tan(c+dx) + a(8Aa^2-12bBa-3Ab^2))}{\sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{(4aB + 5Ab) \cot(c+dx)}{a} \right) - \frac{aA \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d}$$

$$\downarrow 3042$$

$$\frac{1}{4} \left(\frac{\int \frac{ab(5Ab+4aB) \tan(c+dx)^2 + 8a(Ba^2+2Aba-b^2B) \tan(c+dx) + a(8Aa^2-12bBa-3Ab^2)}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{(4aB + 5Ab) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} \right) - \frac{aA \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d}$$

$$\downarrow 4136$$

$$\begin{aligned}
& \frac{1}{4} \left(\frac{\int \frac{8(a(Ba^2+2Aba-b^2B)-a(Aa^2-2bBa-Ab^2)\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx + a(8a^2A - 12abB - 3Ab^2) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b\tan(c+dx)}} dx}{2a} \right. \\
& \qquad \qquad \qquad \frac{aA \cot^2(c+dx) \sqrt{a+b\tan(c+dx)}}{2d} \\
& \qquad \qquad \qquad \downarrow 27 \\
& \frac{1}{4} \left(\frac{8 \int \frac{a(Ba^2+2Aba-b^2B)-a(Aa^2-2bBa-Ab^2)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + a(8a^2A - 12abB - 3Ab^2) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b\tan(c+dx)}} dx}{2a} \right. \\
& \qquad \qquad \qquad \frac{aA \cot^2(c+dx) \sqrt{a+b\tan(c+dx)}}{2d} \\
& \qquad \qquad \qquad \downarrow 3042 \\
& \frac{1}{4} \left(\frac{8 \int \frac{a(Ba^2+2Aba-b^2B)-a(Aa^2-2bBa-Ab^2)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + a(8a^2A - 12abB - 3Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx}{2a} \right. \\
& \qquad \qquad \qquad \frac{aA \cot^2(c+dx) \sqrt{a+b\tan(c+dx)}}{2d} \\
& \qquad \qquad \qquad \downarrow 4022 \\
& \qquad \qquad \qquad - \frac{aA \cot^2(c+dx) \sqrt{a+b\tan(c+dx)}}{2d} + \\
& \frac{1}{4} \left(- \frac{(4aB + 5Ab) \cot(c+dx) \sqrt{a+b\tan(c+dx)}}{d} - \frac{a(8a^2A - 12abB - 3Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx + 8 \left(\frac{1}{2} \right)}{2a} \right. \\
& \qquad \qquad \qquad \downarrow 3042 \\
& \qquad \qquad \qquad - \frac{aA \cot^2(c+dx) \sqrt{a+b\tan(c+dx)}}{2d} + \\
& \frac{1}{4} \left(- \frac{(4aB + 5Ab) \cot(c+dx) \sqrt{a+b\tan(c+dx)}}{d} - \frac{a(8a^2A - 12abB - 3Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx + 8 \left(\frac{1}{2} \right)}{2a} \right. \\
& \qquad \qquad \qquad \downarrow 4020 \\
& \qquad \qquad \qquad - \frac{aA \cot^2(c+dx) \sqrt{a+b\tan(c+dx)}}{2d} + \\
& \frac{1}{4} \left(- \frac{(4aB + 5Ab) \cot(c+dx) \sqrt{a+b\tan(c+dx)}}{d} - \frac{a(8a^2A - 12abB - 3Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx + 8 \left(\frac{1}{2} \right)}{2a} \right)
\end{aligned}$$

3.330. $\int \cot^3(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx$

$$\begin{aligned} & \downarrow 25 \\ & -\frac{aA \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2d} + \\ \frac{1}{4} \left(-\frac{(4aB+5Ab) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{d} - \frac{a(8a^2A-12abB-3Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 8 \left(\frac{a}{b} \right)}{d} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 73 \\ & -\frac{aA \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2d} + \\ \frac{1}{4} \left(-\frac{(4aB+5Ab) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{d} - \frac{a(8a^2A-12abB-3Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 8 \left(\frac{a}{b} \right)}{d} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 221 \\ & -\frac{aA \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2d} + \\ \frac{1}{4} \left(-\frac{(4aB+5Ab) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{d} - \frac{a(8a^2A-12abB-3Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 8 \left(\frac{a}{b} \right)}{d} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 4117 \\ & -\frac{aA \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2d} + \\ \frac{1}{4} \left(-\frac{(4aB+5Ab) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{d} - \frac{a(8a^2A-12abB-3Ab^2) \int \frac{\cot(c+dx)}{\sqrt{a+b \tan(c+dx)}} d \tan(c+dx) + 8 \left(\frac{a(a-ib)^{3/2}(B-2a)}{2a} \right)}{d} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 73 \\ & -\frac{aA \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2d} + \\ \frac{1}{4} \left(-\frac{(4aB+5Ab) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{d} - \frac{2a(8a^2A-12abB-3Ab^2) \int \frac{1}{\frac{a+b \tan(c+dx)}{b} - \frac{a}{b}} d \sqrt{a+b \tan(c+dx)}}{bd} + 8 \left(\frac{a(a-ib)^{3/2}(B-2a)}{2a} \right) \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 221 \\ & -\frac{aA \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} + \\ \frac{1}{4} & \left(-\frac{(4aB+5Ab) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} - \frac{2\sqrt{a}(8a^2A-12abB-3Ab^2) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{d} + 8 \left(\frac{a(a-ib)^3}{2} \right) \right) \end{aligned}$$

input `Int[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `-1/2*(a*A*Cot[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]])/d + (-1/2*(8*((a*(a - I*b)^(3/2)*(I*A + B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d - (a*(a + I*b)^(3/2)*(I*A - B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d - (2*Sqrt[a]*(8*a^2*A - 3*A*b^2 - 12*a*b*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/d)/a - ((5*A*b + 4*a*B)*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/d)/4`

3.330.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.330.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1804 vs. $2(185) = 370$.

Time = 0.26 (sec) , antiderivative size = 1805, normalized size of antiderivative = 8.24

method	result	size
derivativedivides	Expression too large to display	1805
default	Expression too large to display	1805

```
input int(cot(d*x+c)^3*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

output
$$-5/4/d/\tan(dx+c)^2A(a+b\tan(dx+c))^{3/2}+2a^{3/2}A\operatorname{arctanh}((a+b\tan(dx+c))^{1/2}/a^{1/2})/d+1/d*b/(2*(a^2+b^2)^{1/2}-2a)^{1/2}*\operatorname{arctan}((2*(a+b\tan(dx+c))^{1/2}+(2*(a^2+b^2)^{1/2}+2a)^{1/2})/(2*(a^2+b^2)^{1/2}-2a)^{1/2})*B*(a^2+b^2)^{1/2}+1/d/(2*(a^2+b^2)^{1/2}-2a)^{1/2}*\operatorname{arctan}(((2*(a^2+b^2)^{1/2}+2a)^{1/2}-2*(a+b\tan(dx+c))^{1/2})/(2*(a^2+b^2)^{1/2}-2a)^{1/2})*A*(a^2+b^2)^{1/2}*a-1/d/(2*(a^2+b^2)^{1/2}-2a)^{1/2}*\operatorname{arctan}((2*(a+b\tan(dx+c))^{1/2}+(2*(a^2+b^2)^{1/2}+2a)^{1/2})/(2*(a^2+b^2)^{1/2}-2a)^{1/2})*A*(a^2+b^2)^{1/2}*a+1/4/d/b*\ln((a+b\tan(dx+c))^{1/2}*(2*(a^2+b^2)^{1/2}+2a)^{1/2}-b*\tan(dx+c)-a-(a^2+b^2)^{1/2})*B*(2*(a^2+b^2)^{1/2}+2a)^{1/2}*a^2-1/d*b/(2*(a^2+b^2)^{1/2}-2a)^{1/2}*\operatorname{arctan}(((2*(a^2+b^2)^{1/2}+2a)^{1/2}-2*(a+b\tan(dx+c))^{1/2})/(2*(a^2+b^2)^{1/2}-2a)^{1/2})*B*(a^2+b^2)^{1/2}+2/d*b/(2*(a^2+b^2)^{1/2}-2a)^{1/2}*\operatorname{arctan}(((2*(a^2+b^2)^{1/2}+2a)^{1/2}-2*(a+b\tan(dx+c))^{1/2})/(2*(a^2+b^2)^{1/2}-2a)^{1/2})*B*a-2/d*b/(2*(a^2+b^2)^{1/2}-2a)^{1/2}*\operatorname{arctan}((2*(a+b\tan(dx+c))^{1/2}+(2*(a^2+b^2)^{1/2}+2a)^{1/2})/(2*(a^2+b^2)^{1/2}-2a)^{1/2})*B*a-1/4/d/b*\ln(b*\tan(dx+c)+a+(a+b\tan(dx+c))^{1/2}*(2*(a^2+b^2)^{1/2}+2a)^{1/2}+(a^2+b^2)^{1/2})*B*(2*(a^2+b^2)^{1/2}+2a)^{1/2}*a^2-1/4/d/b*\ln((a+b\tan(dx+c))^{1/2}*(2*(a^2+b^2)^{1/2}+2a)^{1/2}-b*\tan(dx+c)-a-(a^2+b^2)^{1/2})*B*(2*(a^2+b^2)^{1/2}+2a)^{1/2}*(a^2+b^2)^{1/2}*a+1/4/d/b*\ln(b*\tan(dx+c)+a+(a+b\tan(dx+c))^{1/2}*(2*(a^2+b^2)^{1/2}+2a)^{1/2}+(a^2+b^2)^{1/2})*B*(2*(...$$

3.330.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3170 vs. $2(179) = 358$.

Time = 6.20 (sec) , antiderivative size = 6357, normalized size of antiderivative = 29.03

$$\int \cot^3(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx = \text{Too large to display}$$

input `integrate(cot(dx+c)^3*(a+b*tan(dx+c))^(3/2)*(A+B*tan(dx+c)),x, algorithm="fracas")`

output Too large to include

3.330.6 Sympy [F]

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx))(a + b \tan(c + dx))^{\frac{3}{2}} \cot^3(c + dx) dx$$

input `integrate(cot(d*x+c)**3*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)*cot(c + d*x)**3, x)`

3.330.7 Maxima [F]

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{3}{2}} \cot(dx + c)^3 dx$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^3, x)`

3.330.8 Giac [F(-1)]

Timed out.

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.330.9 Mupad [B] (verification not implemented)

Time = 9.95 (sec) , antiderivative size = 23016, normalized size of antiderivative = 105.10

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2),x)
```

```
output atan((((932*A^3*a^3*b^12*d^2 + 1344*A^3*a^5*b^10*d^2 - 192*A^3*a^7*b^8*d^2 + 1600*B^3*a^2*b^13*d^2 + 704*B^3*a^4*b^11*d^2 - 896*B^3*a^6*b^9*d^2 + 348*A^2*B*b^15*d^2 - 604*A^3*a*b^14*d^2 + 1760*A*B^2*a*b^14*d^2 - 3232*A*B^2*a^3*b^12*d^2 - 4416*A*B^2*a^5*b^10*d^2 + 576*A*B^2*a^7*b^8*d^2 - 3780*A^2*B*a^2*b^13*d^2 - 1440*A^2*B*a^4*b^11*d^2 + 2688*A^2*B*a^6*b^9*d^2)/(2*d^5) - (((384*A*b^12*d^4 + 1280*B*a*b^11*d^4 - 384*A*a^2*b^10*d^4 - 768*A*a^4*b^8*d^4 + 1280*B*a^3*b^9*d^4)/(2*d^5) + ((512*b^10*d^4 + 768*a^2*b^8*d^4)*(a + b*tan(c + d*x))^(1/2)*(((8*A^2*a^3*d^2 - 8*B^2*a^3*d^2 + 16*A*B*b^3*d^2 - 24*A^2*a*b^2*d^2 + 24*B^2*a*b^2*d^2 - 48*A*B*a^2*b*d^2)^2/64 - d^4*(A^4*a^6 + A^4*b^6 + B^4*a^6 + B^4*b^6 + 2*A^2*B^2*a^6 + 2*A^2*B^2*b^6 + 3*A^4*a^2*b^4 + 3*A^4*a^4*b^2 + 3*B^4*a^2*b^4 + 3*B^4*a^4*b^2 + 6*A^2*B^2*a^2*b^4 + 6*A^2*B^2*a^4*b^2)))^(1/2) + A^2*a^3*d^2 - B^2*a^3*d^2 + 2*A*B*b^3*d^2 - 3*A^2*a*b^2*d^2 + 3*B^2*a*b^2*d^2 - 6*A*B*a^2*b*d^2)/(4*d^4))^(1/2))/d^4)*(((8*A^2*a^3*d^2 - 8*B^2*a^3*d^2 + 16*A*B*b^3*d^2 - 24*A^2*a*b^2*d^2 + 24*B^2*a*b^2*d^2 - 48*A*B*a^2*b*d^2)^2/64 - d^4*(A^4*a^6 + A^4*b^6 + B^4*a^6 + B^4*b^6 + 2*A^2*B^2*a^6 + 2*A^2*B^2*b^6 + 3*A^4*a^2*b^4 + 3*A^4*a^4*b^2 + 3*B^4*a^2*b^4 + 3*B^4*a^4*b^2 + 6*A^2*B^2*a^2*b^4 + 6*A^2*B^2*a^4*b^2)))^(1/2) + A^2*a^3*d^2 - B^2*a^3*d^2 + 2*A*B*b^3*d^2 - 3*A^2*a*b^2*d^2 + 3*B^2*a*b^2*d^2 - 6*A*B*a^2*b*d^2)/(4*d^4))^(1/2) + ((a + b*tan(c + d*x))^(1/2)*(1088*A^2*a^3*b^10*d^2 - 576*A^2*a^5*b^8*d^2 - 1472*B^2*a^3*...
```


3.331 $\int \cot^4(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.331.1 Optimal result	3138
3.331.2 Mathematica [A] (verified)	3139
3.331.3 Rubi [A] (warning: unable to verify)	3139
3.331.4 Maple [B] (verified)	3147
3.331.5 Fricas [B] (verification not implemented)	3148
3.331.6 Sympy [F]	3149
3.331.7 Maxima [F(-1)]	3149
3.331.8 Giac [F(-1)]	3149
3.331.9 Mupad [B] (verification not implemented)	3150

3.331.1 Optimal result

Integrand size = 33, antiderivative size = 278

$$\int \cot^4(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{(24a^2Ab + Ab^3 + 16a^3B - 6ab^2B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right) + B \tan(c+dx)}{8a^{3/2}d} - \frac{(a-ib)^{3/2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d} + \frac{(a+ib)^{3/2}(iA-B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d} + \frac{(8a^2A - Ab^2 - 10abB) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{8ad} - \frac{(7Ab + 6aB) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{12d} - \frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d}$$

```
output 1/8*(24*A*a^2*b+A*b^3+16*B*a^3-6*B*a*b^2)*arctanh((a+b*tan(d*x+c))^(1/2)/a
^(1/2))/a^(3/2)/d-(a-I*b)^(3/2)*(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-
I*b)^(1/2))/d+(a+I*b)^(3/2)*(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)
^(1/2))/d+1/8*(8*A*a^2-A*b^2-10*B*a*b)*cot(d*x+c)*(a+b*tan(d*x+c))^(1/2)/a
/d-1/12*(7*A*b+6*B*a)*cot(d*x+c)^2*(a+b*tan(d*x+c))^(1/2)/d-1/3*a*A*cot(d*
x+c)^3*(a+b*tan(d*x+c))^(1/2)/d
```

3.331.2 Mathematica [A] (verified)

Time = 5.97 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.87

$$\int \cot^4(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx = \frac{3(24a^2Ab + Ab^3 + 16a^3B - 6ab^2B) \operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a}}\right) + \sqrt{a}(-24ia(a-ib))^{3/2}(A+B\tan(c+dx))}{(24a^{3/2}d)}$$

```
input Integrate[Cot[c + d*x]^4*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x
]
```

```
output (3*(24*a^2*A*b + A*b^3 + 16*a^3*B - 6*a*b^2*B)*ArcTanh[Sqrt[a + b*Tan[c +
d*x]]/Sqrt[a]] + Sqrt[a]*((-24*I)*a*(a - I*b)^(3/2)*(A - I*B)*ArcTanh[Sqrt
[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] + (24*I)*a*(a + I*b)^(3/2)*(A + I*B)*A
rcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] - Cot[c + d*x]*(-24*a^2*A +
3*A*b^2 + 30*a*b*B + 2*a*(7*A*b + 6*a*B)*Cot[c + d*x] + 8*a^2*A*Cot[c + d
*x]^2)*Sqrt[a + b*Tan[c + d*x]])/(24*a^(3/2)*d)
```

3.331.3 Rubi [A] (warning: unable to verify)Time = 2.41 (sec) , antiderivative size = 282, normalized size of antiderivative = 1.01, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4088, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^4(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\tan(c+dx)^4} dx$$

$$\downarrow 4088$$

$$\frac{1}{3} \int \frac{\cot^3(c+dx) (-b(5aA - 6bB) \tan^2(c+dx) - 6(Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(7Ab + 6aB))}{2\sqrt{a+b \tan(c+dx)}} dx -$$

$$\frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d}$$

↓ 27

$$\frac{1}{6} \int \frac{\cot^3(c+dx) (-b(5aA - 6bB) \tan^2(c+dx) - 6(Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(7Ab + 6aB))}{\sqrt{a+b \tan(c+dx)}} dx -$$

$$\frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \int \frac{-b(5aA - 6bB) \tan(c+dx)^2 - 6(Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(7Ab + 6aB)}{\tan(c+dx)^3 \sqrt{a+b \tan(c+dx)}} dx -$$

$$\frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d}$$

↓ 4132

$$\frac{1}{6} \left(\int \frac{3 \cot^2(c+dx) (ab(7Ab+6aB) \tan^2(c+dx) + 8a(Ba^2+2Aba-b^2B) \tan(c+dx) + a(8Aa^2-10bBa-Ab^2))}{2\sqrt{a+b \tan(c+dx)}} dx - \frac{(6aB + 7Ab) \cot^2(c+dx)}{2a} \right)$$

$$\frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d}$$

↓ 27

$$\frac{1}{6} \left(\frac{3 \int \frac{\cot^2(c+dx) (ab(7Ab+6aB) \tan^2(c+dx) + 8a(Ba^2+2Aba-b^2B) \tan(c+dx) + a(8Aa^2-10bBa-Ab^2))}{\sqrt{a+b \tan(c+dx)}} dx}{4a} - \frac{(6aB + 7Ab) \cot^2(c+dx)}{2a} \right)$$

$$\frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{3 \int \frac{ab(7Ab+6aB) \tan(c+dx)^2 + 8a(Ba^2+2Aba-b^2B) \tan(c+dx) + a(8Aa^2-10bBa-Ab^2)}{\tan(c+dx)^2 \sqrt{a+b \tan(c+dx)}} dx}{4a} - \frac{(6aB + 7Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} \right)$$

$$\frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d}$$

↓ 4132

3.331. $\int \cot^4(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\frac{1}{6} \left(\frac{3 \left(\int -\frac{\cot(c+dx)(-16(Aa^2-2bBa-Ab^2)\tan(c+dx)a^2-b(8Aa^2-10bBa-Ab^2)\tan^2(c+dx)a+(16Ba^3+24Aba^2-6b^2Ba+Ab^3)a)}{2\sqrt{a+b}\tan(c+dx)} dx - (8a^2A-10abB) \right)}{4a} \right)$$

$$\frac{aA \cot^3(c+dx)\sqrt{a+b}\tan(c+dx)}{3d}$$

↓ 27

$$\frac{1}{6} \left(\frac{3 \left(\int \frac{\cot(c+dx)(-16(Aa^2-2bBa-Ab^2)\tan(c+dx)a^2-b(8Aa^2-10bBa-Ab^2)\tan^2(c+dx)a+(16Ba^3+24Aba^2-6b^2Ba+Ab^3)a)}{\sqrt{a+b}\tan(c+dx)} dx - (8a^2A-10abB) \right)}{4a} \right)$$

$$\frac{aA \cot^3(c+dx)\sqrt{a+b}\tan(c+dx)}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{3 \left(\int \frac{-16(Aa^2-2bBa-Ab^2)\tan(c+dx)a^2-b(8Aa^2-10bBa-Ab^2)\tan(c+dx)^2a+(16Ba^3+24Aba^2-6b^2Ba+Ab^3)a}{\tan(c+dx)\sqrt{a+b}\tan(c+dx)} dx - (8a^2A-10abB-Ab^2)\cot(c+dx) \right)}{4a} \right)$$

$$\frac{aA \cot^3(c+dx)\sqrt{a+b}\tan(c+dx)}{3d}$$

↓ 4136

$$\frac{1}{6} \left(\frac{3 \left(\int -\frac{16((Aa^2-2bBa-Ab^2)a^2+(Ba^2+2Aba-b^2B)\tan(c+dx)a^2)}{\sqrt{a+b}\tan(c+dx)} dx + a(16a^3B+24a^2Ab-6ab^2B+Ab^3) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b}\tan(c+dx)} dx - (8a^2A-10abB-Ab^2)\cot(c+dx) \right)}{4a} \right)$$

$$\frac{aA \cot^3(c+dx)\sqrt{a+b}\tan(c+dx)}{3d}$$

↓ 27

3.331. $\int \cot^4(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx$

$$\frac{1}{6} \left(\frac{3 \left(\frac{a(16a^3B+24a^2Ab-6ab^2B+Ab^3) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx - 16 \int \frac{(Aa^2-2bBa-Ab^2)a^2 + (Ba^2+2Aba-b^2B) \tan(c+dx)a^2}{\sqrt{a+b \tan(c+dx)}} dx}{2a} \right) - (8a^2A)}{4a} \right)$$

$$\frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d}$$

↓ 3042

$$\frac{1}{6} \left(\frac{3 \left(\frac{a(16a^3B+24a^2Ab-6ab^2B+Ab^3) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 16 \int \frac{(Aa^2-2bBa-Ab^2)a^2 + (Ba^2+2Aba-b^2B) \tan(c+dx)a^2}{\sqrt{a+b \tan(c+dx)}} dx}{2a} \right) - (8a^2A)}{4a} \right)$$

$$\frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d}$$

↓ 4022

$$- \frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} +$$

$$\frac{1}{6} \left(\frac{(6aB+7Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} - \frac{3 \left(- \frac{(8a^2A-10abB-Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \frac{a(16a^3B+24a^2Ab-6ab^2B+Ab^3)}{2a} \right)}{4a} \right)$$

↓ 3042

$$- \frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} +$$

$$\frac{1}{6} \left(\frac{(6aB+7Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} - \frac{3 \left(- \frac{(8a^2A-10abB-Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \frac{a(16a^3B+24a^2Ab-6ab^2B+Ab^3)}{2a} \right)}{4a} \right)$$

↓ 4020

3.331. $\int \cot^4(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\frac{1}{6} \left(-\frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \frac{(6aB+7Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} - \frac{3 \left(-\frac{(8a^2A-10abB-Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \frac{a(16a^3B+24a^2)}{\dots} \right)}{\dots} \right)$$

25

$$\frac{1}{6} \left(-\frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \frac{(6aB+7Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} - \frac{3 \left(-\frac{(8a^2A-10abB-Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \frac{a(16a^3B+24a^2)}{\dots} \right)}{\dots} \right)$$

73

$$\frac{1}{6} \left(-\frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \frac{(6aB+7Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} - \frac{3 \left(-\frac{(8a^2A-10abB-Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \frac{a(16a^3B+24a^2)}{\dots} \right)}{\dots} \right)$$

221

$$\frac{1}{6} \left(-\frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \frac{(6aB+7Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} - 3 \left(-\frac{(8a^2A-10abB-Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \frac{a(16a^3B+24a^2)}{\dots} \right) \right)$$

4117

$$\frac{1}{6} \left(-\frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \frac{(6aB+7Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} - 3 \left(-\frac{(8a^2A-10abB-Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \frac{a(16a^3B+24a^2)}{\dots} \right) \right)$$

73

$$\frac{1}{6} \left(-\frac{aA \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \frac{(6aB+7Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} - 3 \left(-\frac{(8a^2A-10abB-Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \frac{2a(16a^3B+24a^2)}{\dots} \right) \right)$$

221

3.331. $\int \cot^4(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\frac{1}{6} \left(-\frac{(6aB + 7Ab) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} - \frac{aA \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{3d} + \frac{(8a^2A - 10abB - Ab^2) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{d} + \frac{2\sqrt{a}(16a^3B + \dots)}{\dots} \right)$$

```
input Int[Cot[c + d*x]^4*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]
```

```
output -1/3*(a*A*Cot[c + d*x]^3*Sqrt[a + b*Tan[c + d*x]])/d + (-1/2*((7*A*b + 6*a*B)*Cot[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]])/d - (3*((-16*((a^2*(a - I*b)^(3/2)*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d + (a^2*(a + I*b)^(3/2)*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d - (2*Sqrt[a]*(24*a^2*A*b + A*b^3 + 16*a^3*B - 6*a*b^2*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/d)/(2*a) - ((8*a^2*A - A*b^2 - 10*a*b*B)*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/d)/(4*a))/6
```

3.331.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

3.331. $\int \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.331.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1984 vs. $2(240) = 480$.

Time = 0.22 (sec) , antiderivative size = 1985, normalized size of antiderivative = 7.14

method	result	size
derivativedivides	Expression too large to display	1985
default	Expression too large to display	1985

```
input int(cot(d*x+c)^4*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

output $2/d*a^{(3/2)}*\operatorname{arctanh}((a+b*\tan(dx+c))^{(1/2)}/a^{(1/2)})*B-1/3/d/\tan(dx+c)^3*A$
 $*(a+b*\tan(dx+c))^{(3/2)}+3/d*b*A*a^{(1/2)}*\operatorname{arctanh}((a+b*\tan(dx+c))^{(1/2)}/a^{(1/2)})$
 $-1/4/d/b*\ln((a+b*\tan(dx+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-b*\tan(dx+c)-a-(a^2+b^2)^{(1/2)})$
 $*A*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*a^2+1/d*b/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}$
 $*\operatorname{arctan}(((2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-2*(a+b*\tan(dx+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})$
 $*A*(a^2+b^2)^{(1/2)}-2/d*b/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}$
 $*\operatorname{arctan}(((2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-2*(a+b*\tan(dx+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})$
 $*A*a+1/d/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}$
 $*\operatorname{arctan}(((2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-2*(a+b*\tan(dx+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})$
 $*B*(a^2+b^2)^{(1/2)}*a+1/4/d/b*\ln((a+b*\tan(dx+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-b*\tan(dx+c)-a-(a^2+b^2)^{(1/2)})$
 $*A*(a^2+b^2)^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*a-1/4/d/b*\ln(b*\tan(dx+c)+a+(a+b*\tan(dx+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}+(a^2+b^2)^{(1/2)})$
 $*A*(a^2+b^2)^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*a-5/4/d/b/\tan(dx+c)^3*(a+b*\tan(dx+c))^{(5/2)}$
 $*B-1/8/d/\tan(dx+c)^3/a*(a+b*\tan(dx+c))^{(5/2)}*A+1/8/d/\tan(dx+c)^3*(a+b*\tan(dx+c))^{(1/2)}$
 $*A*a+1/8/d*b^3/a^{(3/2)}*\operatorname{arctanh}((a+b*\tan(dx+c))^{(1/2)}/a^{(1/2)})$
 $*A-3/4/d*b^2/a^{(1/2)}*\operatorname{arctanh}((a+b*\tan(dx+c))^{(1/2)}/a^{(1/2)})$
 $*B+1/4/d*\ln(b*\tan(dx+c)+a+(a+b*\tan(dx+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}+(a^2+b^2)^{(1/2)})$
 $*B*(a^2+b^2)^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-1/2/d*\ln(b*\tan(dx+c)+a+(a+b*\tan(dx+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}+...$

3.331.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3218 vs. $2(234) = 468$.

Time = 15.33 (sec) , antiderivative size = 6452, normalized size of antiderivative = 23.21

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(dx+c)^4*(a+b*tan(dx+c))^(3/2)*(A+B*tan(dx+c)),x, algorithm="fracas")`

output Too large to include

3.331.6 Sympy [F]

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx))(a + b \tan(c + dx))^{\frac{3}{2}} \cot^4(c + dx) dx$$

input `integrate(cot(d*x+c)**4*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)*cot(c + d*x)**4, x)`

3.331.7 Maxima [F(-1)]

Timed out.

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `Timed out`

3.331.8 Giac [F(-1)]

Timed out.

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.331.9 Mupad [B] (verification not implemented)

Time = 10.79 (sec) , antiderivative size = 25789, normalized size of antiderivative = 92.77

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(cot(c + d*x)^4*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2),x)
```

```
output atan(((((((256*A*a*b^13*d^4 + 5376*A*a^3*b^11*d^4 + 5120*A*a^5*b^9*d^4 - 1
536*B*a^2*b^12*d^4 + 1536*B*a^4*b^10*d^4 + 3072*B*a^6*b^8*d^4)/(8*a^2*d^5)
- ((2048*a^2*b^10*d^4 + 3072*a^4*b^8*d^4)*(a + b*tan(c + d*x))^(1/2))*(-(
(8*A^2*a^3*d^2 - 8*B^2*a^3*d^2 + 16*A*B*b^3*d^2 - 24*A^2*a*b^2*d^2 + 24*B^
2*a*b^2*d^2 - 48*A*B*a^2*b*d^2)^2/64 - d^4*(A^4*a^6 + A^4*b^6 + B^4*a^6 +
B^4*b^6 + 2*A^2*B^2*a^6 + 2*A^2*B^2*b^6 + 3*A^4*a^2*b^4 + 3*A^4*a^4*b^2 +
3*B^4*a^2*b^4 + 3*B^4*a^4*b^2 + 6*A^2*B^2*a^2*b^4 + 6*A^2*B^2*a^4*b^2))^(1
/2) + A^2*a^3*d^2 - B^2*a^3*d^2 + 2*A*B*b^3*d^2 - 3*A^2*a*b^2*d^2 + 3*B^2*
a*b^2*d^2 - 6*A*B*a^2*b*d^2)/(4*d^4))^(1/2))/(4*a^2*d^4))*(-(((8*A^2*a^3*d
^2 - 8*B^2*a^3*d^2 + 16*A*B*b^3*d^2 - 24*A^2*a*b^2*d^2 + 24*B^2*a*b^2*d^2
- 48*A*B*a^2*b*d^2)^2/64 - d^4*(A^4*a^6 + A^4*b^6 + B^4*a^6 + B^4*b^6 + 2*
A^2*B^2*a^6 + 2*A^2*B^2*b^6 + 3*A^4*a^2*b^4 + 3*A^4*a^4*b^2 + 3*B^4*a^2*b^
4 + 3*B^4*a^4*b^2 + 6*A^2*B^2*a^2*b^4 + 6*A^2*B^2*a^4*b^2))^(1/2) + A^2*a^
3*d^2 - B^2*a^3*d^2 + 2*A*B*b^3*d^2 - 3*A^2*a*b^2*d^2 + 3*B^2*a*b^2*d^2 -
6*A*B*a^2*b*d^2)/(4*d^4))^(1/2) + ((a + b*tan(c + d*x))^(1/2)*(3008*A^2*a^
3*b^12*d^2 + 5888*A^2*a^5*b^10*d^2 - 1280*A^2*a^7*b^8*d^2 - 2672*B^2*a^3*b
^12*d^2 - 4352*B^2*a^5*b^10*d^2 + 2304*B^2*a^7*b^8*d^2 + 4*A^2*a*b^14*d^2
- 2096*A*B*a^2*b^13*d^2 + 1024*A*B*a^4*b^11*d^2 + 11264*A*B*a^6*b^9*d^2))/
(4*a^2*d^4))*(-(((8*A^2*a^3*d^2 - 8*B^2*a^3*d^2 + 16*A*B*b^3*d^2 - 24*A^2*
a*b^2*d^2 + 24*B^2*a*b^2*d^2 - 48*A*B*a^2*b*d^2)^2/64 - d^4*(A^4*a^6 + ...
```

3.332 $\int \tan^2(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.332.1 Optimal result	3151
3.332.2 Mathematica [A] (verified)	3152
3.332.3 Rubi [A] (warning: unable to verify)	3152
3.332.4 Maple [B] (verified)	3157
3.332.5 Fricas [B] (verification not implemented)	3158
3.332.6 Sympy [F]	3159
3.332.7 Maxima [F(-1)]	3160
3.332.8 Giac [F(-1)]	3160
3.332.9 Mupad [F(-1)]	3160

3.332.1 Optimal result

Integrand size = 33, antiderivative size = 252

$$\int \tan^2(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \frac{(a-ib)^{5/2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d} - \frac{(a+ib)^{5/2}(iA-B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d} - \frac{2(2aAb+a^2B-b^2B)\sqrt{a+b \tan(c+dx)}}{d} - \frac{2(Ab+aB)(a+b \tan(c+dx))^{3/2}}{3d} - \frac{2B(a+b \tan(c+dx))^{5/2}}{5d} + \frac{2(9Ab-2aB)(a+b \tan(c+dx))^{7/2}}{63b^2d} + \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{7/2}}{9bd}$$

output

```
(a-I*b)^(5/2)*(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d-(a+I
*b)^(5/2)*(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d-2*(2*A*a
*b+B*a^2-B*b^2)*(a+b*tan(d*x+c))^(1/2)/d-2/3*(A*b+B*a)*(a+b*tan(d*x+c))^(3
/2)/d-2/5*B*(a+b*tan(d*x+c))^(5/2)/d+2/63*(9*A*b-2*B*a)*(a+b*tan(d*x+c))^(
7/2)/b^2/d+2/9*B*tan(d*x+c)*(a+b*tan(d*x+c))^(7/2)/b/d
```

3.332.2 Mathematica [A] (verified)

Time = 4.97 (sec) , antiderivative size = 296, normalized size of antiderivative = 1.17

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{2(9Ab - 2aB)(a + b \tan(c + dx))^{7/2}}{b} + 14B \tan(c + dx)(a + b \tan(c + dx))^{7/2} - \frac{63}{2}ib(A - iB) \left(\frac{2}{5}\right)$$

input `Integrate[Tan[c + d*x]^2*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `((2*(9*A*b - 2*a*B)*(a + b*Tan[c + d*x])^(7/2))/b + 14*B*Tan[c + d*x]*(a + b*Tan[c + d*x])^(7/2) - ((63*I)/2)*b*(A - I*B)*((2*(a + b*Tan[c + d*x])^(5/2))/5 + (2*(a - I*b)*(-3*(a - I*b)^(3/2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] + Sqrt[a + b*Tan[c + d*x]]*(4*a - (3*I)*b + b*Tan[c + d*x])))/3) + ((63*I)/2)*b*(A + I*B)*((2*(a + b*Tan[c + d*x])^(5/2))/5 + (2*(a + I*b)*(-3*(a + I*b)^(3/2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + Sqrt[a + b*Tan[c + d*x]]*(4*a + (3*I)*b + b*Tan[c + d*x])))/3))/(63*b*d)`

3.332.3 Rubi [A] (warning: unable to verify)

Time = 1.61 (sec) , antiderivative size = 246, normalized size of antiderivative = 0.98, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 4090, 27, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^2(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4090}$$

$$\begin{aligned}
& \frac{2 \int -\frac{1}{2}(a + b \tan(c + dx))^{5/2} (-(9Ab - 2aB) \tan^2(c + dx) + 9bB \tan(c + dx) + 2aB) dx}{9b} + \\
& \quad \frac{2B \tan(c + dx)(a + b \tan(c + dx))^{7/2}}{9bd} \\
& \quad \downarrow 27 \\
& \quad \frac{2B \tan(c + dx)(a + b \tan(c + dx))^{7/2}}{9bd} - \\
& \frac{\int (a + b \tan(c + dx))^{5/2} (-(9Ab - 2aB) \tan^2(c + dx) + 9bB \tan(c + dx) + 2aB) dx}{9b} \\
& \quad \downarrow 3042 \\
& \quad \frac{2B \tan(c + dx)(a + b \tan(c + dx))^{7/2}}{9bd} - \\
& \frac{\int (a + b \tan(c + dx))^{5/2} (-(9Ab - 2aB) \tan^2(c + dx) + 9bB \tan(c + dx) + 2aB) dx}{9b} \\
& \quad \downarrow 4113 \\
& \quad \frac{2B \tan(c + dx)(a + b \tan(c + dx))^{7/2}}{9bd} - \\
& \frac{\int (a + b \tan(c + dx))^{5/2} (9Ab + 9B \tan(c + dx)b) dx - \frac{2(9Ab - 2aB)(a + b \tan(c + dx))^{7/2}}{7bd}}{9b} \\
& \quad \downarrow 3042 \\
& \quad \frac{2B \tan(c + dx)(a + b \tan(c + dx))^{7/2}}{9bd} - \\
& \frac{\int (a + b \tan(c + dx))^{5/2} (9Ab + 9B \tan(c + dx)b) dx - \frac{2(9Ab - 2aB)(a + b \tan(c + dx))^{7/2}}{7bd}}{9b} \\
& \quad \downarrow 4011 \\
& \quad \frac{2B \tan(c + dx)(a + b \tan(c + dx))^{7/2}}{9bd} - \\
& \frac{\int (a + b \tan(c + dx))^{3/2} (9b(aA - bB) + 9b(Ab + aB) \tan(c + dx)) dx - \frac{2(9Ab - 2aB)(a + b \tan(c + dx))^{7/2}}{7bd} + \frac{18bB(a + b \tan(c + dx))^{7/2}}{5d}}{9b} \\
& \quad \downarrow 3042 \\
& \quad \frac{2B \tan(c + dx)(a + b \tan(c + dx))^{7/2}}{9bd} - \\
& \frac{\int (a + b \tan(c + dx))^{3/2} (9b(aA - bB) + 9b(Ab + aB) \tan(c + dx)) dx - \frac{2(9Ab - 2aB)(a + b \tan(c + dx))^{7/2}}{7bd} + \frac{18bB(a + b \tan(c + dx))^{7/2}}{5d}}{9b} \\
& \quad \downarrow 4011
\end{aligned}$$

3.332. $\int \tan^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\frac{2B \tan(c + dx)(a + b \tan(c + dx))^{7/2}}{9bd} - \frac{\int \sqrt{a + b \tan(c + dx)}(9b(Aa^2 - 2bBa - Ab^2) + 9b(Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx - \frac{2(9Ab - 2aB)(a + b \tan(c + dx))}{7bd}}{9b}$$

↓ 3042

$$\frac{2B \tan(c + dx)(a + b \tan(c + dx))^{7/2}}{9bd} - \frac{\int \sqrt{a + b \tan(c + dx)}(9b(Aa^2 - 2bBa - Ab^2) + 9b(Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx - \frac{2(9Ab - 2aB)(a + b \tan(c + dx))}{7bd}}{9b}$$

↓ 4011

$$\frac{2B \tan(c + dx)(a + b \tan(c + dx))^{7/2}}{9bd} - \frac{\int \frac{9b(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + 9b(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{18b(a^2B + 2aAb - b^2B) \sqrt{a + b \tan(c + dx)}}{d} - \frac{2(9Ab - 2aB)(a + b \tan(c + dx))}{7bd}}{9b}$$

↓ 3042

$$\frac{2B \tan(c + dx)(a + b \tan(c + dx))^{7/2}}{9bd} - \frac{\int \frac{9b(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + 9b(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{18b(a^2B + 2aAb - b^2B) \sqrt{a + b \tan(c + dx)}}{d} - \frac{2(9Ab - 2aB)(a + b \tan(c + dx))}{7bd}}{9b}$$

↓ 4022

$$\frac{2B \tan(c + dx)(a + b \tan(c + dx))^{7/2}}{9bd} - \frac{\frac{9}{2}b(a + ib)^3(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{9}{2}b(a - ib)^3(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \frac{18b(a^2B + 2aAb - b^2B) \sqrt{a + b \tan(c + dx)}}{d}}{9b}$$

↓ 3042

$$\frac{2B \tan(c + dx)(a + b \tan(c + dx))^{7/2}}{9bd} - \frac{\frac{9}{2}b(a + ib)^3(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{9}{2}b(a - ib)^3(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \frac{18b(a^2B + 2aAb - b^2B) \sqrt{a + b \tan(c + dx)}}{d}}{9b}$$

↓ 4020

$$\frac{2B \tan(c + dx)(a + b \tan(c + dx))^{7/2}}{9bd} - \frac{\frac{9ib(a - ib)^3(A - iB) \int -\frac{1}{(1 - i \tan(c + dx)) \sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx)) - \frac{9ib(a + ib)^3(A + iB) \int -\frac{1}{(i \tan(c + dx) + 1) \sqrt{a + b \tan(c + dx)}} d(-i \tan(c + dx))}{2d}}{9b} +$$

↓ 25

3.332. $\int \tan^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\begin{aligned}
 & \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{7/2}}{9bd} - \frac{9ib(a-ib)^3(A-ib) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \frac{9ib(a+ib)^3(A+ib) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \dots \\
 & \quad \downarrow \text{73} \\
 & \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{7/2}}{9bd} - \frac{9bd}{9(a+ib)^3(A+ib) \int \frac{1}{-i \tan^2(\frac{c+dx}{b}) - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}} + \frac{9(a-ib)^3(A-ib) \int \frac{1}{i \tan^2(\frac{c+dx}{b}) + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{d} + \frac{18b(a^2B+2aAb-b^2)}{9b} \\
 & \quad \downarrow \text{221} \\
 & \frac{2B \tan(c+dx)(a+b \tan(c+dx))^{7/2}}{9bd} - \frac{9bd}{18b(a^2B+2aAb-b^2B)\sqrt{a+b \tan(c+dx)}} + \frac{9b(a-ib)^{5/2}(A-ib) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{9b(a+ib)^{5/2}(A+ib) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} - \frac{2(9Ab-2aB)}{9b}
 \end{aligned}$$

input `Int[Tan[c + d*x]^2*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(2*B*Tan[c + d*x]*(a + b*Tan[c + d*x])^(7/2))/(9*b*d) - ((9*(a - I*b)^(5/2)*b*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d + (9*(a + I*b)^(5/2)*b*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d + (18*b*(2*a*A*b + a^2*B - b^2*B)*Sqrt[a + b*Tan[c + d*x]])/d + (6*b*(A*b + a*B)*(a + b*Tan[c + d*x])^(3/2))/d + (18*b*B*(a + b*Tan[c + d*x])^(5/2))/(5*d) - (2*(9*A*b - 2*a*B)*(a + b*Tan[c + d*x])^(7/2))/(7*b*d))/(9*b)`

3.332.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
 Q[u, x]`
- rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
 (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
 [(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
 , x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
 0] && GtQ[m, 0]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) +
 (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
 c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
 b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
 (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
 1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
 *(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
 - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

```
rule 4090 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4113 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

3.332.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2437 vs. $2(216) = 432$.

Time = 0.18 (sec) , antiderivative size = 2438, normalized size of antiderivative = 9.67

method	result	size
parts	Expression too large to display	2438
derivativedivides	Expression too large to display	2469
default	Expression too large to display	2469

```
input int(tan(d*x+c)^2*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

output

```
A*(2/7/b/d*(a+b*tan(d*x+c))^(7/2)-2/3*b*(a+b*tan(d*x+c))^(3/2)/d-4*b/d*(a+b*tan(d*x+c))^(1/2)*a+1/4/b/d*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*a^2-1/4*b/d*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)-1/4/b/d*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3+3/4*b/d*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+2*b/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*(a^2+b^2)^(1/2)*a-3*b/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^2+b^3/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))-1/4/b/d*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*a^2+1/4*b/d*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)+1/4/b/d*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3-3/...
```

3.332.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4963 vs. $2(210) = 420$.

Time = 0.86 (sec) , antiderivative size = 4963, normalized size of antiderivative = 19.69

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input

```
integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output 1/630*(315*b^2*d*sqrt((10*A*B*a^4*b - 20*A*B*a^2*b^3 + 2*A*B*b^5 - (A^2 -
B^2)*a^5 + 10*(A^2 - B^2)*a^3*b^2 - 5*(A^2 - B^2)*a*b^4 + d^2*sqrt(-(4*A^2
*B^2*a^10 + 20*(A^3*B - A*B^3)*a^9*b + 5*(5*A^4 - 26*A^2*B^2 + 5*B^4)*a^8*
b^2 - 240*(A^3*B - A*B^3)*a^7*b^3 - 20*(5*A^4 - 32*A^2*B^2 + 5*B^4)*a^6*b^
4 + 504*(A^3*B - A*B^3)*a^5*b^5 + 10*(11*A^4 - 62*A^2*B^2 + 11*B^4)*a^4*b^
6 - 240*(A^3*B - A*B^3)*a^3*b^7 - 20*(A^4 - 7*A^2*B^2 + B^4)*a^2*b^8 + 20*
(A^3*B - A*B^3)*a*b^9 + (A^4 - 2*A^2*B^2 + B^4)*b^10)/d^4))/d^2)*log(-(2*(
A^3*B + A*B^3)*a^9 + 5*(A^4 - B^4)*a^8*b - 16*(A^3*B + A*B^3)*a^7*b^2 - 28
*(A^3*B + A*B^3)*a^5*b^4 - 14*(A^4 - B^4)*a^4*b^5 - 8*(A^4 - B^4)*a^2*b^7
+ 10*(A^3*B + A*B^3)*a*b^8 + (A^4 - B^4)*b^9)*sqrt(b*tan(d*x + c) + a) + (
(A*a^2 - 2*B*a*b - A*b^2)*d^3*sqrt(-(4*A^2*B^2*a^10 + 20*(A^3*B - A*B^3)*a
^9*b + 5*(5*A^4 - 26*A^2*B^2 + 5*B^4)*a^8*b^2 - 240*(A^3*B - A*B^3)*a^7*b^
3 - 20*(5*A^4 - 32*A^2*B^2 + 5*B^4)*a^6*b^4 + 504*(A^3*B - A*B^3)*a^5*b^5
+ 10*(11*A^4 - 62*A^2*B^2 + 11*B^4)*a^4*b^6 - 240*(A^3*B - A*B^3)*a^3*b^7
- 20*(A^4 - 7*A^2*B^2 + B^4)*a^2*b^8 + 20*(A^3*B - A*B^3)*a*b^9 + (A^4 - 2
*A^2*B^2 + B^4)*b^10)/d^4) - (2*A*B^2*a^7 + (9*A^2*B - 5*B^3)*a^6*b + 2*(5
*A^3 - 16*A*B^2)*a^5*b^2 - 5*(11*A^2*B - 3*B^3)*a^4*b^3 - 10*(2*A^3 - 5*A*
B^2)*a^3*b^4 + (31*A^2*B - 11*B^3)*a^2*b^5 + 2*(A^3 - 6*A*B^2)*a*b^6 - (A^
2*B - B^3)*b^7)*d)*sqrt((10*A*B*a^4*b - 20*A*B*a^2*b^3 + 2*A*B*b^5 - (A^2
- B^2)*a^5 + 10*(A^2 - B^2)*a^3*b^2 - 5*(A^2 - B^2)*a*b^4 + d^2*sqrt(-(...
```

3.332.6 Sympy [F]

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx))(a + b \tan(c + dx))^{5/2} \tan^2(c + dx) dx$$

```
input integrate(tan(d*x+c)**2*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)
```

```
output Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(5/2)*tan(c + d*x)**2,
x)
```

3.332.7 Maxima [F(-1)]

Timed out.

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `Timed out`

3.332.8 Giac [F(-1)]

Timed out.

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.332.9 Mupad [F(-1)]

Timed out.

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Hanged}$$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)`

output `\text{Hanged}`

3.333 $\int \tan(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.333.1 Optimal result	3161
3.333.2 Mathematica [A] (verified)	3162
3.333.3 Rubi [A] (warning: unable to verify)	3162
3.333.4 Maple [B] (verified)	3166
3.333.5 Fricas [B] (verification not implemented)	3167
3.333.6 Sympy [F]	3168
3.333.7 Maxima [F]	3169
3.333.8 Giac [F(-1)]	3169
3.333.9 Mupad [B] (verification not implemented)	3169

3.333.1 Optimal result

Integrand size = 31, antiderivative size = 213

$$\int \tan(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx =$$

$$\frac{(a-ib)^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d} - \frac{(a+ib)^{5/2}(A+iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d}$$

$$+ \frac{2(a^2A - Ab^2 - 2abB)\sqrt{a+b \tan(c+dx)}}{d} + \frac{2(aA - bB)(a+b \tan(c+dx))^{3/2}}{3d}$$

$$+ \frac{2A(a+b \tan(c+dx))^{5/2}}{5d} + \frac{2B(a+b \tan(c+dx))^{7/2}}{7bd}$$

output $-(a-I*b)^{(5/2)}*(A-I*B)*\operatorname{arctanh}((a+b*\tan(d*x+c))^{(1/2)}/(a-I*b)^{(1/2)})/d-(a+I*b)^{(5/2)}*(A+I*B)*\operatorname{arctanh}((a+b*\tan(d*x+c))^{(1/2)}/(a+I*b)^{(1/2)})/d+2*(A*a^2-A*b^2-2*B*a*b)*(a+b*\tan(d*x+c))^{(1/2)}/d+2/3*(A*a-B*b)*(a+b*\tan(d*x+c))^{(3/2)}/d+2/5*A*(a+b*\tan(d*x+c))^{(5/2)}/d+2/7*B*(a+b*\tan(d*x+c))^{(7/2)}/b/d$

3.333.2 Mathematica [A] (verified)

Time = 1.74 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.21

$$\int \tan(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{\frac{4B(a+b \tan(c+dx))^{7/2}}{b} - 7i(iA + B) \left(\frac{2}{5}(a + b \tan(c + dx))^{5/2} + \frac{2}{3}(a - ib) \left(-3(a - ib)^{3/2} \arctan \left(\frac{a + b \tan(c + dx)}{a - ib} \right) \right) \right)}{14d}$$

input `Integrate[Tan[c + d*x]*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`output `((4*B*(a + b*Tan[c + d*x])^(7/2))/b - (7*I)*(I*A + B)*((2*(a + b*Tan[c + d*x])^(5/2))/5 + (2*(a - I*b)*(-3*(a - I*b)^(3/2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] + Sqrt[a + b*Tan[c + d*x]]*(4*a - (3*I)*b + b*Tan[c + d*x])))/3) - (7*I)*(I*A - B)*((2*(a + b*Tan[c + d*x])^(5/2))/5 + (2*(a + I*b)*(-3*(a + I*b)^(3/2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + Sqrt[a + b*Tan[c + d*x]]*(4*a + (3*I)*b + b*Tan[c + d*x])))/3))/(14*d)`**3.333.3 Rubi [A] (warning: unable to verify)**Time = 1.16 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.01, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 4075, 3042, 4011, 3042, 4011, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4075} \\ & \int (A \tan(c + dx) - B)(a + b \tan(c + dx))^{5/2} dx + \frac{2B(a + b \tan(c + dx))^{7/2}}{7bd} \\ & \quad \downarrow \text{3042} \\ & \int (A \tan(c + dx) - B)(a + b \tan(c + dx))^{5/2} dx + \frac{2B(a + b \tan(c + dx))^{7/2}}{7bd} \end{aligned}$$

$$\begin{aligned}
& \int (a + b \tan(c + dx))^{3/2} (-Ab - aB + (aA - bB) \tan(c + dx)) dx + \frac{2A(a + b \tan(c + dx))^{5/2}}{5d} + \\
& \quad \frac{2B(a + b \tan(c + dx))^{7/2}}{7bd} \\
& \quad \downarrow \text{4011} \\
& \int (a + b \tan(c + dx))^{3/2} (-Ab - aB + (aA - bB) \tan(c + dx)) dx + \frac{2A(a + b \tan(c + dx))^{5/2}}{5d} + \\
& \quad \frac{2B(a + b \tan(c + dx))^{7/2}}{7bd} \\
& \quad \downarrow \text{3042} \\
& \int \sqrt{a + b \tan(c + dx)} (-Ba^2 - 2Aba + b^2B + (Aa^2 - 2bBa - Ab^2) \tan(c + dx)) dx + \\
& \quad \frac{2(aA - bB)(a + b \tan(c + dx))^{3/2}}{3d} + \frac{2A(a + b \tan(c + dx))^{5/2}}{5d} + \frac{2B(a + b \tan(c + dx))^{7/2}}{7bd} \\
& \quad \downarrow \text{4011} \\
& \int \sqrt{a + b \tan(c + dx)} (-Ba^2 - 2Aba + b^2B + (Aa^2 - 2bBa - Ab^2) \tan(c + dx)) dx + \\
& \quad \frac{2(aA - bB)(a + b \tan(c + dx))^{3/2}}{3d} + \frac{2A(a + b \tan(c + dx))^{5/2}}{5d} + \frac{2B(a + b \tan(c + dx))^{7/2}}{7bd} \\
& \quad \downarrow \text{3042} \\
& \int \frac{-Ba^3 - 3Aba^2 + 3b^2Ba + Ab^3 + (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \\
& \quad \frac{2(a^2A - 2abB - Ab^2) \sqrt{a + b \tan(c + dx)}}{d} + \frac{2(aA - bB)(a + b \tan(c + dx))^{3/2}}{3d} + \\
& \quad \frac{2A(a + b \tan(c + dx))^{5/2}}{5d} + \frac{2B(a + b \tan(c + dx))^{7/2}}{7bd} \\
& \quad \downarrow \text{4011} \\
& \int \frac{-Ba^3 - 3Aba^2 + 3b^2Ba + Ab^3 + (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \\
& \quad \frac{2(a^2A - 2abB - Ab^2) \sqrt{a + b \tan(c + dx)}}{d} + \frac{2(aA - bB)(a + b \tan(c + dx))^{3/2}}{3d} + \\
& \quad \frac{2A(a + b \tan(c + dx))^{5/2}}{5d} + \frac{2B(a + b \tan(c + dx))^{7/2}}{7bd} \\
& \quad \downarrow \text{3042} \\
& \int \frac{-Ba^3 - 3Aba^2 + 3b^2Ba + Ab^3 + (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \\
& \quad \frac{2(a^2A - 2abB - Ab^2) \sqrt{a + b \tan(c + dx)}}{d} + \frac{2(aA - bB)(a + b \tan(c + dx))^{3/2}}{3d} + \\
& \quad \frac{2A(a + b \tan(c + dx))^{5/2}}{5d} + \frac{2B(a + b \tan(c + dx))^{7/2}}{7bd} \\
& \quad \downarrow \text{4022}
\end{aligned}$$

$$\begin{aligned}
& -\frac{1}{2}(-b+ia)^3(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}(b+ia)^3(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \\
& \quad \frac{2(a^2A-2abB-Ab^2)}{d} \frac{1}{\sqrt{a+b \tan(c+dx)}} + \frac{2(aA-bB)(a+b \tan(c+dx))^{3/2}}{3d} + \\
& \quad \frac{2A(a+b \tan(c+dx))^{5/2}}{5d} + \frac{2B(a+b \tan(c+dx))^{7/2}}{7bd} \\
& \quad \downarrow \text{3042} \\
& -\frac{1}{2}(-b+ia)^3(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}(b+ia)^3(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \\
& \quad \frac{2(a^2A-2abB-Ab^2)}{d} \frac{1}{\sqrt{a+b \tan(c+dx)}} + \frac{2(aA-bB)(a+b \tan(c+dx))^{3/2}}{3d} + \\
& \quad \frac{2A(a+b \tan(c+dx))^{5/2}}{5d} + \frac{2B(a+b \tan(c+dx))^{7/2}}{7bd} \\
& \quad \downarrow \text{4020} \\
& \quad \frac{i(b+ia)^3(A-iB) \int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \\
& \quad \frac{i(-b+ia)^3(A+iB) \int -\frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \\
& \quad \frac{2(a^2A-2abB-Ab^2)}{d} \frac{1}{\sqrt{a+b \tan(c+dx)}} + \frac{2(aA-bB)(a+b \tan(c+dx))^{3/2}}{3d} + \\
& \quad \frac{2A(a+b \tan(c+dx))^{5/2}}{5d} + \frac{2B(a+b \tan(c+dx))^{7/2}}{7bd} \\
& \quad \downarrow \text{25} \\
& \quad \frac{i(b+ia)^3(A-iB) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} - \\
& \quad \frac{i(-b+ia)^3(A+iB) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \\
& \quad \frac{2(a^2A-2abB-Ab^2)}{d} \frac{1}{\sqrt{a+b \tan(c+dx)}} + \frac{2(aA-bB)(a+b \tan(c+dx))^{3/2}}{3d} + \\
& \quad \frac{2A(a+b \tan(c+dx))^{5/2}}{5d} + \frac{2B(a+b \tan(c+dx))^{7/2}}{7bd} \\
& \quad \downarrow \text{73} \\
& \quad \frac{(-b+ia)^3(A+iB) \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \\
& \quad \frac{(b+ia)^3(A-iB) \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \\
& \quad \frac{2(a^2A-2abB-Ab^2)}{d} \frac{1}{\sqrt{a+b \tan(c+dx)}} + \frac{2(aA-bB)(a+b \tan(c+dx))^{3/2}}{3d} + \\
& \quad \frac{2A(a+b \tan(c+dx))^{5/2}}{5d} + \frac{2B(a+b \tan(c+dx))^{7/2}}{7bd}
\end{aligned}$$

3.333. $\int \tan(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\begin{aligned}
 & \downarrow 221 \\
 & \frac{2(a^2A - 2abB - Ab^2)\sqrt{a + b \tan(c + dx)}}{d} + \frac{(b + ia)^3(A - iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \\
 & \frac{(-b + ia)^3(A + iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} + \frac{2(aA - bB)(a + b \tan(c + dx))^{3/2}}{3d} + \\
 & \frac{2A(a + b \tan(c + dx))^{5/2}}{5d} + \frac{2B(a + b \tan(c + dx))^{7/2}}{7bd}
 \end{aligned}$$

input `Int[Tan[c + d*x]*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `((I*a + b)^3*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]]/(Sqrt[a - I*b]*d) - ((I*a - b)^3*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]]/(Sqrt[a + I*b]*d) + (2*(a^2*A - A*b^2 - 2*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/d + (2*(a*A - b*B)*(a + b*Tan[c + d*x])^(3/2))/(3*d) + (2*A*(a + b*Tan[c + d*x])^(5/2))/(5*d) + (2*B*(a + b*Tan[c + d*x])^(7/2))/(7*b*d)`

3.333.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.333.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2406 vs. $2(181) = 362$.

Time = 0.12 (sec) , antiderivative size = 2407, normalized size of antiderivative = 11.30

method	result	size
parts	Expression too large to display	2407
derivativedivides	Expression too large to display	2426
default	Expression too large to display	2426

input `int(tan(d*x+c)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 2/5*A*(a+b*\tan(d*x+c))^{5/2}/d+2/3/d*A*a*(a+b*\tan(d*x+c))^{3/2}+2/d*(a+b*\tan(d*x+c))^{1/2} \\ & *A*a^2-2/d*b^2*(a+b*\tan(d*x+c))^{1/2}*A-3/4/d*\ln(b*\tan(d*x+c))+a+(a+b*\tan(d*x+c))^{1/2} \\ & *(2*(a^2+b^2)^{1/2}+2*a)^{1/2}+(a^2+b^2)^{1/2})*A*(2*(a^2+b^2)^{1/2}+2*a)^{1/2} \\ & *(2*(a^2+b^2)^{1/2}+2*a)^{1/2}+a+(a+b*\tan(d*x+c))^{1/2}*(2*(a^2+b^2)^{1/2}+2*a)^{1/2} \\ & *(2*(a^2+b^2)^{1/2}+2*a)^{1/2}+3/4*A/d*\ln(b*\tan(d*x+c))+a-(a+b*\tan(d*x+c))^{1/2} \\ & *(2*(a^2+b^2)^{1/2}+2*a)^{1/2}+(a^2+b^2)^{1/2})*(2*(a^2+b^2)^{1/2}+2*a)^{1/2} \\ & *a^2-1/4*A/d*\ln(b*\tan(d*x+c))+a-(a+b*\tan(d*x+c))^{1/2}*(2*(a^2+b^2)^{1/2}+2*a)^{1/2} \\ & *(2*(a^2+b^2)^{1/2}+2*a)^{1/2}*(2*(a^2+b^2)^{1/2}+2*a)^{1/2} \\ & *b^2+1/d/(2*(a^2+b^2)^{1/2}-2*a)^{1/2}*\arctan((2*(a+b*\tan(d*x+c))^{1/2}+(2*(a^2+b^2)^{1/2}+2*a)^{1/2}))/ \\ & (2*(a^2+b^2)^{1/2}-2*a)^{1/2})*A*a^3+A/d/(2*(a^2+b^2)^{1/2}-2*a)^{1/2}*\arctan((2*(a+b*\tan(d*x+c))^{1/2}- \\ & (2*(a^2+b^2)^{1/2}+2*a)^{1/2}))/((2*(a^2+b^2)^{1/2}-2*a)^{1/2})*a^3-3/d*b^2/(2*(a^2+b^2)^{1/2}-2*a)^{1/2} \\ & *\arctan((2*(a+b*\tan(d*x+c))^{1/2}+(2*(a^2+b^2)^{1/2}+2*a)^{1/2}))/((2*(a^2+b^2)^{1/2}-2*a)^{1/2}) \\ & *A*a-3*A/d/(2*(a^2+b^2)^{1/2}-2*a)^{1/2}*\arctan((2*(a+b*\tan(d*x+c))^{1/2}- \\ & (2*(a^2+b^2)^{1/2}+2*a)^{1/2}))/((2*(a^2+b^2)^{1/2}-2*a)^{1/2}) \\ & *a*b^2+1/2/d*\ln(b*\tan(d*x+c))+a+(a+b*\tan(d*x+c))^{1/2}*(2*(a^2+b^2)^{1/2}+2*a)^{1/2} \\ & *(2*(a^2+b^2)^{1/2}+2*a)^{1/2}+(a^2+b^2)^{1/2})*A*(2*(a^2+b^2)^{1/2}+2*a)^{1/2} \\ & *(a^2+b^2)^{1/2})*a-1/d/(2*(a^2+b^2)^{1/2}-2*a)^{1/2}*\arctan((2*(a+b*\tan(d*x+c))^{1/2}+(2*(a^2+b^2)^{1/2}+2*a)^{1/2}))/ \\ & (2*(a^2+b^2)^{1/2}-2*a)^{1/2})*A*(a^2+b^2)^{1/2} \dots \end{aligned}$$

3.333.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4916 vs. $2(175) = 350$.

Time = 0.84 (sec) , antiderivative size = 4916, normalized size of antiderivative = 23.08

$$\int \tan(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/210*(105*b*d*sqrt(-(10*A*B*a^4*b - 20*A*B*a^2*b^3 + 2*A*B*b^5 - (A^2 - B^2)*a^5 + 10*(A^2 - B^2)*a^3*b^2 - 5*(A^2 - B^2)*a*b^4 + d^2*sqrt(-(4*A^2*B^2*a^10 + 20*(A^3*B - A*B^3)*a^9*b + 5*(5*A^4 - 26*A^2*B^2 + 5*B^4)*a^8*b^2 - 240*(A^3*B - A*B^3)*a^7*b^3 - 20*(5*A^4 - 32*A^2*B^2 + 5*B^4)*a^6*b^4 + 504*(A^3*B - A*B^3)*a^5*b^5 + 10*(11*A^4 - 62*A^2*B^2 + 11*B^4)*a^4*b^6 - 240*(A^3*B - A*B^3)*a^3*b^7 - 20*(A^4 - 7*A^2*B^2 + B^4)*a^2*b^8 + 20*(A^3*B - A*B^3)*a*b^9 + (A^4 - 2*A^2*B^2 + B^4)*b^10)/d^4))/d^2)*log(-(2*(A^3*B + A*B^3)*a^9 + 5*(A^4 - B^4)*a^8*b - 16*(A^3*B + A*B^3)*a^7*b^2 - 28*(A^3*B + A*B^3)*a^5*b^4 - 14*(A^4 - B^4)*a^4*b^5 - 8*(A^4 - B^4)*a^2*b^7 + 10*(A^3*B + A*B^3)*a*b^8 + (A^4 - B^4)*b^9)*sqrt(b*tan(d*x + c) + a) + ((B*a^2 + 2*A*a*b - B*b^2)*d^3*sqrt(-(4*A^2*B^2*a^10 + 20*(A^3*B - A*B^3)*a^9*b + 5*(5*A^4 - 26*A^2*B^2 + 5*B^4)*a^8*b^2 - 240*(A^3*B - A*B^3)*a^7*b^3 - 20*(5*A^4 - 32*A^2*B^2 + 5*B^4)*a^6*b^4 + 504*(A^3*B - A*B^3)*a^5*b^5 + 10*(11*A^4 - 62*A^2*B^2 + 11*B^4)*a^4*b^6 - 240*(A^3*B - A*B^3)*a^3*b^7 - 20*(A^4 - 7*A^2*B^2 + B^4)*a^2*b^8 + 20*(A^3*B - A*B^3)*a*b^9 + (A^4 - 2*A^2*B^2 + B^4)*b^10)/d^4) + (2*A^2*B*a^7 + (5*A^3 - 9*A*B^2)*a^6*b - 2*(16*A^2*B - 5*B^3)*a^5*b^2 - 5*(3*A^3 - 11*A*B^2)*a^4*b^3 + 10*(5*A^2*B - 2*B^3)*a^3*b^4 + (11*A^3 - 31*A*B^2)*a^2*b^5 - 2*(6*A^2*B - B^3)*a*b^6 - (A^3 - A*B^2)*b^7)*d)*sqrt(-(10*A*B*a^4*b - 20*A*B*a^2*b^3 + 2*A*B*b^5 - (A^2 - B^2)*a^5 + 10*(A^2 - B^2)*a^3*b^2 - 5*(A^2 - B^2)*a*b^4 + d^2*sqrt(-(...`

3.333.6 Sympy [F]

$$\int \tan(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx))(a + b \tan(c + dx))^{5/2} \tan(c + dx) dx$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(5/2)*tan(c + d*x), x)`

3.333.7 Maxima [F]

$$\int \tan(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{5/2} \tan(dx + c) dx$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)*tan(d*x + c), x)`

3.333.8 Giac [F(-1)]

Timed out.

$$\int \tan(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.333.9 Mupad [B] (verification not implemented)

Time = 94.32 (sec) , antiderivative size = 3932, normalized size of antiderivative = 18.46

$$\int \tan(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)`

output

$$\begin{aligned} & \log(- (((((-A^4*b^2*d^4*(5*a^4 + b^4 - 10*a^2*b^2)^2)^{(1/2)} + A^2*a^5*d^2 \\ & - 10*A^2*a^3*b^2*d^2 + 5*A^2*a*b^4*d^2)/d^4)^{(1/2)}*(32*A*b^6 - 32*A*a^4*b \\ & ^2 + 32*a*b^2*d*((-A^4*b^2*d^4*(5*a^4 + b^4 - 10*a^2*b^2)^2)^{(1/2)} + A^2* \\ & a^5*d^2 - 10*A^2*a^3*b^2*d^2 + 5*A^2*a*b^4*d^2)/d^4)^{(1/2)}*(a + b*\tan(c + \\ & d*x))^{(1/2)}))/ (2*d) - (16*A^2*b^2*(a + b*\tan(c + d*x))^{(1/2)}*(a^6 - b^6 + \\ & 15*a^2*b^4 - 15*a^4*b^2))/d^2)* ((((-A^4*b^2*d^4*(5*a^4 + b^4 - 10*a^2*b^2)^2)^{(1/2)} + A^2*a^5*d^2 - 10*A^2*a^3*b^2*d^2 + 5*A^2*a*b^4*d^2)/d^4)^{(1/2)}) \\ & /2 - (8*A^3*a*b^2*(a^2 - 3*b^2)*(a^2 + b^2)^3)/d^3)* ((20*A^4*a^2*b^8*d^4 - \\ & A^4*b^10*d^4 - 110*A^4*a^4*b^6*d^4 + 100*A^4*a^6*b^4*d^4 - 25*A^4*a^8*b^2 \\ & *d^4)^{(1/2)}/(4*d^4) + (A^2*a^5)/(4*d^2) - (5*A^2*a^3*b^2)/(2*d^2) + (5*A^2 \\ & *a*b^4)/(4*d^2))^{(1/2)} - \log((((((-A^4*b^2*d^4*(5*a^4 + b^4 - 10*a^2*b^2)^2)^{(1/2)} + A^2*a^5*d^2 - 10*A^2*a^3*b^2*d^2 + 5*A^2*a*b^4*d^2)/d^4)^{(1/2)} \\ & *(32*A*a^4*b^2 - 32*A*b^6 + 32*a*b^2*d*((-A^4*b^2*d^4*(5*a^4 + b^4 - 10*a^2*b^2)^2)^{(1/2)} + A^2*a^5*d^2 - 10*A^2*a^3*b^2*d^2 + 5*A^2*a*b^4*d^2)/d^4 \\ &)^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)}))/ (2*d) - (16*A^2*b^2*(a + b*\tan(c + d* \\ & x))^{(1/2)}*(a^6 - b^6 + 15*a^2*b^4 - 15*a^4*b^2))/d^2)* ((((-A^4*b^2*d^4*(5*a^4 + b^4 - 10*a^2*b^2)^2)^{(1/2)} + A^2*a^5*d^2 - 10*A^2*a^3*b^2*d^2 + 5*A^2 \\ & *a*b^4*d^2)/d^4)^{(1/2)})/2 - (8*A^3*a*b^2*(a^2 - 3*b^2)*(a^2 + b^2)^3)/d^3) \\ & * (((20*A^4*a^2*b^8*d^4 - A^4*b^10*d^4 - 110*A^4*a^4*b^6*d^4 + 100*A^4*a^6* \\ & b^4*d^4 - 25*A^4*a^8*b^2*d^4)^{(1/2)} + A^2*a^5*d^2 - 10*A^2*a^3*b^2*d^2 \dots \end{aligned}$$

3.334 $\int (a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx$

3.334.1 Optimal result	3171
3.334.2 Mathematica [A] (verified)	3172
3.334.3 Rubi [A] (warning: unable to verify)	3172
3.334.4 Maple [B] (verified)	3176
3.334.5 Fricas [B] (verification not implemented)	3177
3.334.6 Sympy [F]	3177
3.334.7 Maxima [F(-2)]	3178
3.334.8 Giac [F(-1)]	3178
3.334.9 Mupad [B] (verification not implemented)	3178

3.334.1 Optimal result

Integrand size = 25, antiderivative size = 188

$$\int (a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx =$$

$$-\frac{(a - ib)^{5/2} (iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}}\right)}{d}$$

$$+ \frac{(a + ib)^{5/2} (iA - B) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a + ib}}\right)}{d}$$

$$+ \frac{2(2aAb + a^2B - b^2B) \sqrt{a + b \tan(c + dx)}}{d}$$

$$+ \frac{2(Ab + aB)(a + b \tan(c + dx))^{3/2}}{3d} + \frac{2B(a + b \tan(c + dx))^{5/2}}{5d}$$

output

```
-(a-I*b)^(5/2)*(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d+(a+
I*b)^(5/2)*(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d+2*(2*A*
a*b+B*a^2-B*b^2)*(a+b*tan(d*x+c))^(1/2)/d+2/3*(A*b+B*a)*(a+b*tan(d*x+c))^(
3/2)/d+2/5*B*(a+b*tan(d*x+c))^(5/2)/d
```

3.334.2 Mathematica [A] (verified)

Time = 1.17 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.24

$$\int (a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx = \frac{i \left((A - iB) \left(\frac{2}{5} (a + b \tan(c + dx))^{5/2} + \frac{2}{3} (a - ib) \left(-3(a - ib)^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}} \right) \right) \right) \right)}{\dots}$$

input `Integrate[(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`output `((I/2)*((A - I*B)*((2*(a + b*Tan[c + d*x])^(5/2))/5 + (2*(a - I*b)*(-3*(a - I*b)^(3/2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] + Sqrt[a + b*Tan[c + d*x]]*(4*a - (3*I)*b + b*Tan[c + d*x])))/3) - (A + I*B)*((2*(a + b*Tan[c + d*x])^(5/2))/5 + (2*(a + I*b)*(-3*(a + I*b)^(3/2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + Sqrt[a + b*Tan[c + d*x]]*(4*a + (3*I)*b + b*Tan[c + d*x])))/3))/d`**3.334.3 Rubi [A] (warning: unable to verify)**Time = 0.99 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.90, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.520$, Rules used = {3042, 4011, 3042, 4011, 3042, 4011, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4011} \\ & \int (a + b \tan(c + dx))^{3/2} (aA - bB + (Ab + aB) \tan(c + dx)) dx + \frac{2B(a + b \tan(c + dx))^{5/2}}{5d} \\ & \quad \downarrow \text{3042} \\ & \int (a + b \tan(c + dx))^{3/2} (aA - bB + (Ab + aB) \tan(c + dx)) dx + \frac{2B(a + b \tan(c + dx))^{5/2}}{5d} \end{aligned}$$

$$\begin{aligned}
& \int \sqrt{a + b \tan(c + dx)} (Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx + \\
& \quad \frac{2(aB + Ab)(a + b \tan(c + dx))^{3/2}}{3d} + \frac{2B(a + b \tan(c + dx))^{5/2}}{5d} \\
& \quad \downarrow 4011 \\
& \int \sqrt{a + b \tan(c + dx)} (Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx + \\
& \quad \frac{2(aB + Ab)(a + b \tan(c + dx))^{3/2}}{3d} + \frac{2B(a + b \tan(c + dx))^{5/2}}{5d} \\
& \quad \downarrow 3042 \\
& \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \\
& \quad \frac{2(a^2B + 2aAb - b^2B) \sqrt{a + b \tan(c + dx)}}{d} + \frac{2(aB + Ab)(a + b \tan(c + dx))^{3/2}}{3d} + \\
& \quad \frac{2B(a + b \tan(c + dx))^{5/2}}{5d} \\
& \quad \downarrow 4011 \\
& \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \\
& \quad \frac{2(a^2B + 2aAb - b^2B) \sqrt{a + b \tan(c + dx)}}{d} + \frac{2(aB + Ab)(a + b \tan(c + dx))^{3/2}}{3d} + \\
& \quad \frac{2B(a + b \tan(c + dx))^{5/2}}{5d} \\
& \quad \downarrow 3042 \\
& \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \\
& \quad \frac{2(a^2B + 2aAb - b^2B) \sqrt{a + b \tan(c + dx)}}{d} + \frac{2(aB + Ab)(a + b \tan(c + dx))^{3/2}}{3d} + \\
& \quad \frac{2B(a + b \tan(c + dx))^{5/2}}{5d} \\
& \quad \downarrow 4022 \\
& \frac{1}{2}(a - ib)^3(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a + ib)^3(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \\
& \quad \frac{2(a^2B + 2aAb - b^2B) \sqrt{a + b \tan(c + dx)}}{d} + \frac{2(aB + Ab)(a + b \tan(c + dx))^{3/2}}{3d} + \\
& \quad \frac{2B(a + b \tan(c + dx))^{5/2}}{5d} \\
& \quad \downarrow 3042 \\
& \frac{1}{2}(a - ib)^3(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a + ib)^3(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \\
& \quad \frac{2(a^2B + 2aAb - b^2B) \sqrt{a + b \tan(c + dx)}}{d} + \frac{2(aB + Ab)(a + b \tan(c + dx))^{3/2}}{3d} + \\
& \quad \frac{2B(a + b \tan(c + dx))^{5/2}}{5d} \\
& \quad \downarrow 4020
\end{aligned}$$

$$\begin{aligned}
& \frac{i(a-ib)^3(A-iB) \int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} - \\
& \frac{i(a+ib)^3(A+iB) \int -\frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \\
& \frac{2(a^2B+2aAb-b^2B) \sqrt{a+b \tan(c+dx)}}{d} + \frac{2(aB+Ab)(a+b \tan(c+dx))^{3/2}}{3d} + \\
& \frac{2B(a+b \tan(c+dx))^{5/2}}{5d} \\
& \quad \downarrow \text{25} \\
& \frac{i(a-ib)^3(A-iB) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \\
& \frac{i(a+ib)^3(A+iB) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \\
& \frac{2(a^2B+2aAb-b^2B) \sqrt{a+b \tan(c+dx)}}{d} + \frac{2(aB+Ab)(a+b \tan(c+dx))^{3/2}}{3d} + \\
& \frac{2B(a+b \tan(c+dx))^{5/2}}{5d} \\
& \quad \downarrow \text{73} \\
& \frac{(a-ib)^3(A-iB) \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \\
& \frac{(a+ib)^3(A+iB) \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \\
& \frac{2(a^2B+2aAb-b^2B) \sqrt{a+b \tan(c+dx)}}{d} + \frac{2(aB+Ab)(a+b \tan(c+dx))^{3/2}}{3d} + \\
& \frac{2B(a+b \tan(c+dx))^{5/2}}{5d} \\
& \quad \downarrow \text{221} \\
& \frac{2(a^2B+2aAb-b^2B) \sqrt{a+b \tan(c+dx)}}{d} + \frac{(a-ib)^{5/2}(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \\
& \frac{(a+ib)^{5/2}(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} + \frac{2(aB+Ab)(a+b \tan(c+dx))^{3/2}}{3d} + \\
& \frac{2B(a+b \tan(c+dx))^{5/2}}{5d}
\end{aligned}$$

input `Int[(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

```
output ((a - I*b)^(5/2)*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d + ((a + I
*b)^(5/2)*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d + (2*(2*a*A*b +
a^2*B - b^2*B)*Sqrt[a + b*Tan[c + d*x]])/d + (2*(A*b + a*B)*(a + b*Tan[c +
d*x])^(3/2))/(3*d) + (2*B*(a + b*Tan[c + d*x])^(5/2))/(5*d)
```

3.334.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4011 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

3.334.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2386 vs. $2(160) = 320$.

Time = 0.11 (sec) , antiderivative size = 2387, normalized size of antiderivative = 12.70

method	result	size
parts	Expression too large to display	2387
derivativedivides	Expression too large to display	2405
default	Expression too large to display	2405

```
input int((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output A*(2/3*b*(a+b*tan(d*x+c))^(3/2)/d+4*b/d*(a+b*tan(d*x+c))^(1/2)*a-1/4/b/d*ln
(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2
+b^2)^(1/2))*2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*a^2+1/4*b/d*ln(
b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b
^2)^(1/2))*2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)+1/4/b/d*ln(b*tan(
d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1
/2))*2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3-3/4*b/d*ln(b*tan(d*x+c)+a+(a+b*tan(
d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*2*(a^2+b^2)^(
1/2)+2*a)^(1/2)*a-2*b/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(
d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)
)*(a^2+b^2)^(1/2)*a+3*b/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan
(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2
))*a^2-b^3/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2
)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))+1/4/b/d*ln
((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+
b^2)^(1/2))*2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*a^2-1/4*b/d*ln((
a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^
2)^(1/2))*2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)-1/4/b/d*ln((a+b*ta
n(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/
2))*2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3+3/4*b/d*ln((a+b*tan(d*x+c))^(1/2)...
```

3.334.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4855 vs. $2(154) = 308$.

Time = 0.82 (sec) , antiderivative size = 4855, normalized size of antiderivative = 25.82

$$\int (a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output

```
-1/30*(15*d*sqrt((10*A*B*a^4*b - 20*A*B*a^2*b^3 + 2*A*B*b^5 - (A^2 - B^2)*
a^5 + 10*(A^2 - B^2)*a^3*b^2 - 5*(A^2 - B^2)*a*b^4 + d^2*sqrt(-(4*A^2*B^2*
a^10 + 20*(A^3*B - A*B^3)*a^9*b + 5*(5*A^4 - 26*A^2*B^2 + 5*B^4)*a^8*b^2 -
240*(A^3*B - A*B^3)*a^7*b^3 - 20*(5*A^4 - 32*A^2*B^2 + 5*B^4)*a^6*b^4 + 5
04*(A^3*B - A*B^3)*a^5*b^5 + 10*(11*A^4 - 62*A^2*B^2 + 11*B^4)*a^4*b^6 - 2
40*(A^3*B - A*B^3)*a^3*b^7 - 20*(A^4 - 7*A^2*B^2 + B^4)*a^2*b^8 + 20*(A^3*
B - A*B^3)*a*b^9 + (A^4 - 2*A^2*B^2 + B^4)*b^10)/d^4)/d^2)*log(-(2*(A^3*B
+ A*B^3)*a^9 + 5*(A^4 - B^4)*a^8*b - 16*(A^3*B + A*B^3)*a^7*b^2 - 28*(A^3
*B + A*B^3)*a^5*b^4 - 14*(A^4 - B^4)*a^4*b^5 - 8*(A^4 - B^4)*a^2*b^7 + 10*
(A^3*B + A*B^3)*a*b^8 + (A^4 - B^4)*b^9)*sqrt(b*tan(d*x + c) + a) + ((A*a^
2 - 2*B*a*b - A*b^2)*d^3*sqrt(-(4*A^2*B^2*a^10 + 20*(A^3*B - A*B^3)*a^9*b
+ 5*(5*A^4 - 26*A^2*B^2 + 5*B^4)*a^8*b^2 - 240*(A^3*B - A*B^3)*a^7*b^3 - 2
0*(5*A^4 - 32*A^2*B^2 + 5*B^4)*a^6*b^4 + 504*(A^3*B - A*B^3)*a^5*b^5 + 10*
(11*A^4 - 62*A^2*B^2 + 11*B^4)*a^4*b^6 - 240*(A^3*B - A*B^3)*a^3*b^7 - 20*
(A^4 - 7*A^2*B^2 + B^4)*a^2*b^8 + 20*(A^3*B - A*B^3)*a*b^9 + (A^4 - 2*A^2*
B^2 + B^4)*b^10)/d^4) - (2*A*B^2*a^7 + (9*A^2*B - 5*B^3)*a^6*b + 2*(5*A^3
- 16*A*B^2)*a^5*b^2 - 5*(11*A^2*B - 3*B^3)*a^4*b^3 - 10*(2*A^3 - 5*A*B^2)*
a^3*b^4 + (31*A^2*B - 11*B^3)*a^2*b^5 + 2*(A^3 - 6*A*B^2)*a*b^6 - (A^2*B -
B^3)*b^7)*d)*sqrt((10*A*B*a^4*b - 20*A*B*a^2*b^3 + 2*A*B*b^5 - (A^2 - B^2)
)*a^5 + 10*(A^2 - B^2)*a^3*b^2 - 5*(A^2 - B^2)*a*b^4 + d^2*sqrt(-(4*A^2...
```

3.334.6 Sympy [F]

$$\int (a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2} dx$$

input `integrate((a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(5/2), x)`

3.334.7 Maxima [F(-2)]

Exception generated.

$$\int (a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is`

3.334.8 Giac [F(-1)]

Timed out.

$$\int (a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.334.9 Mupad [B] (verification not implemented)

Time = 41.53 (sec) , antiderivative size = 3863, normalized size of antiderivative = 20.55

$$\int (a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)`

output

$$\begin{aligned} & \log(- (((((-B^4*b^2*d^4*(5*a^4 + b^4 - 10*a^2*b^2)^2)^{(1/2)} + B^2*a^5*d^2 \\ & - 10*B^2*a^3*b^2*d^2 + 5*B^2*a*b^4*d^2)/d^4)^{(1/2)}*(32*B*b^6 - 32*B*a^4*b \\ & ^2 + 32*a*b^2*d*((-B^4*b^2*d^4*(5*a^4 + b^4 - 10*a^2*b^2)^2)^{(1/2)} + B^2* \\ & a^5*d^2 - 10*B^2*a^3*b^2*d^2 + 5*B^2*a*b^4*d^2)/d^4)^{(1/2)}*(a + b*\tan(c + \\ & d*x))^{(1/2)}))/ (2*d) - (16*B^2*b^2*(a + b*\tan(c + d*x))^{(1/2)}*(a^6 - b^6 + \\ & 15*a^2*b^4 - 15*a^4*b^2))/d^2)* ((((-B^4*b^2*d^4*(5*a^4 + b^4 - 10*a^2*b^2)^2)^{(1/2)} + B^2*a^5*d^2 - 10*B^2*a^3*b^2*d^2 + 5*B^2*a*b^4*d^2)/d^4)^{(1/2)}) \\ & /2 - (8*B^3*a*b^2*(a^2 - 3*b^2)*(a^2 + b^2)^3)/d^3)* ((20*B^4*a^2*b^8*d^4 - \\ & B^4*b^10*d^4 - 110*B^4*a^4*b^6*d^4 + 100*B^4*a^6*b^4*d^4 - 25*B^4*a^8*b^2 \\ & *d^4)^{(1/2)}/(4*d^4) + (B^2*a^5)/(4*d^2) - (5*B^2*a^3*b^2)/(2*d^2) + (5*B^2 \\ & *a*b^4)/(4*d^2))^{(1/2)} - \log((((((-B^4*b^2*d^4*(5*a^4 + b^4 - 10*a^2*b^2)^2)^{(1/2)} + B^2*a^5*d^2 - 10*B^2*a^3*b^2*d^2 + 5*B^2*a*b^4*d^2)/d^4)^{(1/2)} \\ & *(32*B*a^4*b^2 - 32*B*b^6 + 32*a*b^2*d*((-B^4*b^2*d^4*(5*a^4 + b^4 - 10*a^2*b^2)^2)^{(1/2)} + B^2*a^5*d^2 - 10*B^2*a^3*b^2*d^2 + 5*B^2*a*b^4*d^2)/d^4 \\ &)^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)}))/ (2*d) - (16*B^2*b^2*(a + b*\tan(c + d* \\ & x))^{(1/2)}*(a^6 - b^6 + 15*a^2*b^4 - 15*a^4*b^2))/d^2)* ((((-B^4*b^2*d^4*(5*a^4 + b^4 - 10*a^2*b^2)^2)^{(1/2)} + B^2*a^5*d^2 - 10*B^2*a^3*b^2*d^2 + 5*B^2 \\ & *a*b^4*d^2)/d^4)^{(1/2)})/2 - (8*B^3*a*b^2*(a^2 - 3*b^2)*(a^2 + b^2)^3)/d^3) \\ & * (((20*B^4*a^2*b^8*d^4 - B^4*b^10*d^4 - 110*B^4*a^4*b^6*d^4 + 100*B^4*a^6* \\ & b^4*d^4 - 25*B^4*a^8*b^2*d^4)^{(1/2)} + B^2*a^5*d^2 - 10*B^2*a^3*b^2*d^2 \dots \end{aligned}$$

3.335 $\int \cot(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.335.1 Optimal result	3180
3.335.2 Mathematica [A] (verified)	3181
3.335.3 Rubi [A] (warning: unable to verify)	3181
3.335.4 Maple [B] (verified)	3187
3.335.5 Fricas [B] (verification not implemented)	3188
3.335.6 Sympy [F]	3189
3.335.7 Maxima [F]	3189
3.335.8 Giac [F(-1)]	3189
3.335.9 Mupad [B] (verification not implemented)	3190

3.335.1 Optimal result

Integrand size = 31, antiderivative size = 182

$$\int \cot(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx =$$

$$-\frac{2a^{5/2}A \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{d} + \frac{(a-ib)^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d}$$

$$+ \frac{(a+ib)^{5/2}(A+iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d}$$

$$+ \frac{2b(Ab+2aB)\sqrt{a+b \tan(c+dx)}}{d} + \frac{2bB(a+b \tan(c+dx))^{3/2}}{3d}$$

output

```
-2*a^(5/2)*A*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/d+(a-I*b)^(5/2)*(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d+(a+I*b)^(5/2)*(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d+2*b*(A*b+2*B*a)*(a+b*tan(d*x+c))^(1/2)/d+2/3*b*B*(a+b*tan(d*x+c))^(3/2)/d
```

3.335.2 Mathematica [A] (verified)

Time = 1.26 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.97

$$\int \cot(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{2\left(-3a^{5/2}A \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right) + \frac{3}{2}(a - ib)^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right) + B \tan(c+dx)\right)}{3d}$$

input `Integrate[Cot[c + d*x]*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`output `(2*(-3*a^(5/2)*A*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]] + (3*(a - I*b)^(5/2)*(A - I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]]))/2 + (3*(a + I*b)^(5/2)*(A + I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/2 + 3*b*(A*b + 2*a*B)*Sqrt[a + b*Tan[c + d*x]] + b*B*(a + b*Tan[c + d*x])^(3/2))/(3*d)`**3.335.3 Rubi [A] (warning: unable to verify)**Time = 1.64 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.93, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.581$, Rules used = {3042, 4090, 27, 3042, 4130, 27, 3042, 4136, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan(c + dx)} dx \\ & \quad \downarrow \text{4090} \\ & \frac{2}{3} \int \frac{3}{2} \cot(c + dx) \sqrt{a + b \tan(c + dx)} (Aa^2 + b(Ab + 2aB) \tan^2(c + dx) + (Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx + \frac{2bB(a + b \tan(c + dx))^{3/2}}{3d} \end{aligned}$$

$$\begin{aligned}
& \downarrow 27 \\
& \int \cot(c + dx) \sqrt{a + b \tan(c + dx)} (Aa^2 + b(Ab + 2aB) \tan^2(c + dx) + (Ba^2 + 2Aba - b^2B) \tan(c + dx)) dx + \\
& \quad \frac{2bB(a + b \tan(c + dx))^{3/2}}{3d} \\
& \downarrow 3042 \\
& \int \frac{\sqrt{a + b \tan(c + dx)} (Aa^2 + b(Ab + 2aB) \tan(c + dx)^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx))}{\tan(c + dx) \frac{2bB(a + b \tan(c + dx))^{3/2}}{3d}} dx + \\
& \downarrow 4130 \\
& 2 \int \frac{\cot(c + dx) (Aa^3 + b(3Ba^2 + 3Aba - b^2B) \tan^2(c + dx) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx))}{\frac{2\sqrt{a + b \tan(c + dx)}}{d} + \frac{2bB(a + b \tan(c + dx))^{3/2}}{3d}} dx + \\
& \downarrow 27 \\
& \int \frac{\cot(c + dx) (Aa^3 + b(3Ba^2 + 3Aba - b^2B) \tan^2(c + dx) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx))}{\frac{\sqrt{a + b \tan(c + dx)}}{d} + \frac{2bB(a + b \tan(c + dx))^{3/2}}{3d}} dx + \\
& \downarrow 3042 \\
& \int \frac{Aa^3 + b(3Ba^2 + 3Aba - b^2B) \tan(c + dx)^2 + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\tan(c + dx) \frac{\sqrt{a + b \tan(c + dx)}}{d} + \frac{2bB(a + b \tan(c + dx))^{3/2}}{3d}} dx + \\
& \downarrow 4136 \\
& a^3 A \int \frac{\cot(c + dx) (\tan^2(c + dx) + 1)}{\sqrt{a + b \tan(c + dx)}} dx + \\
& \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\frac{\sqrt{a + b \tan(c + dx)}}{d} + \frac{2bB(a + b \tan(c + dx))^{3/2}}{3d}} dx + \\
& \downarrow 3042
\end{aligned}$$

3.335. $\int \cot(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& a^3 A \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx + \\
& \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + \\
& \frac{2b(2aB + Ab)\sqrt{a+b\tan(c+dx)}}{d} + \frac{2bB(a+b\tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{4022} \\
& a^3 A \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - \frac{1}{2}(a+ib)^3(-B+iA) \int \frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + \\
& \frac{1}{2}(a-ib)^3(B+iA) \int \frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}} dx + \frac{2b(2aB + Ab)\sqrt{a+b\tan(c+dx)}}{d} + \\
& \frac{2bB(a+b\tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{3042} \\
& a^3 A \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - \frac{1}{2}(a+ib)^3(-B+iA) \int \frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + \\
& \frac{1}{2}(a-ib)^3(B+iA) \int \frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}} dx + \frac{2b(2aB + Ab)\sqrt{a+b\tan(c+dx)}}{d} + \\
& \frac{2bB(a+b\tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{4020} \\
& a^3 A \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx + \\
& \frac{i(a-ib)^3(B+iA) \int -\frac{1}{(1-i\tan(c+dx))\sqrt{a+b\tan(c+dx)}} d(i\tan(c+dx))}{2d} + \\
& \frac{i(a+ib)^3(-B+iA) \int -\frac{1}{(i\tan(c+dx)+1)\sqrt{a+b\tan(c+dx)}} d(-i\tan(c+dx))}{2d} + \\
& \frac{2b(2aB + Ab)\sqrt{a+b\tan(c+dx)}}{d} + \frac{2bB(a+b\tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{25} \\
& a^3 A \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - \\
& \frac{i(a-ib)^3(B+iA) \int \frac{1}{(1-i\tan(c+dx))\sqrt{a+b\tan(c+dx)}} d(i\tan(c+dx))}{2d} - \\
& \frac{i(a+ib)^3(-B+iA) \int \frac{1}{(i\tan(c+dx)+1)\sqrt{a+b\tan(c+dx)}} d(-i\tan(c+dx))}{2d} + \\
& \frac{2b(2aB + Ab)\sqrt{a+b\tan(c+dx)}}{d} + \frac{2bB(a+b\tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{73}
\end{aligned}$$

3.335. $\int \cot(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$

$$\begin{aligned}
& \frac{a^3 A \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - (a+ib)^3(-B+iA) \int \frac{1}{-\frac{i\tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b\tan(c+dx)}}{bd} + \\
& \frac{(a-ib)^3(B+iA) \int \frac{1}{\frac{i\tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b\tan(c+dx)}}{bd} + \frac{2b(2aB+Ab)\sqrt{a+b\tan(c+dx)}}{d} + \\
& \frac{2bB(a+b\tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{221} \\
& \frac{a^3 A \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx + \frac{(a-ib)^{5/2}(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} -}{d} - \\
& \frac{(a+ib)^{5/2}(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right) + \frac{2b(2aB+Ab)\sqrt{a+b\tan(c+dx)}}{d} +}{d} + \\
& \frac{2bB(a+b\tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{4117} \\
& \frac{a^3 A \int \frac{\cot(c+dx)}{\sqrt{a+b\tan(c+dx)}} d\tan(c+dx) + \frac{(a-ib)^{5/2}(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} -}{d} - \\
& \frac{(a+ib)^{5/2}(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right) + \frac{2b(2aB+Ab)\sqrt{a+b\tan(c+dx)}}{d} +}{d} + \\
& \frac{2bB(a+b\tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{73} \\
& \frac{2a^3 A \int \frac{1}{\frac{a+b\tan(c+dx)}{b} - \frac{a}{b}} d\sqrt{a+b\tan(c+dx)} + \frac{(a-ib)^{5/2}(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} -}{bd} - \\
& \frac{(a+ib)^{5/2}(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right) + \frac{2b(2aB+Ab)\sqrt{a+b\tan(c+dx)}}{d} +}{d} + \\
& \frac{2bB(a+b\tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{221} \\
& -\frac{2a^{5/2} A \operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a}}\right)}{d} + \frac{(a-ib)^{5/2}(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} - \\
& \frac{(a+ib)^{5/2}(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right) + \frac{2b(2aB+Ab)\sqrt{a+b\tan(c+dx)}}{d} +}{d} + \\
& \frac{2bB(a+b\tan(c+dx))^{3/2}}{3d}
\end{aligned}$$

input `Int[Cot[c + d*x]*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `((a - I*b)^(5/2)*(I*A + B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d - ((a + I*b)^(5/2)*(I*A - B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d - (2*a^(5/2)*A*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/d + (2*b*(A*b + 2*a*B)*Sqrt[a + b*Tan[c + d*x]])/d + (2*b*B*(a + b*Tan[c + d*x])^(3/2))/(3*d)`

3.335.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n)), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.335.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2372 vs. $2(152) = 304$.

Time = 0.25 (sec) , antiderivative size = 2373, normalized size of antiderivative = 13.04

method	result	size
derivativedivides	Expression too large to display	2373
default	Expression too large to display	2373

```
input int(cot(d*x+c)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

output $\frac{2}{d*b^2*(a+b*\tan(d*x+c))^{1/2}}*A-2*a^{5/2}*A*\operatorname{arctanh}((a+b*\tan(d*x+c))^{1/2})/a^{1/2})/d+2/3*b*B*(a+b*\tan(d*x+c))^{3/2}/d-2/d*b/(2*(a^2+b^2)^{1/2}-2*a)^{1/2}*\operatorname{arctan}((2*(a+b*\tan(d*x+c))^{1/2}+(2*(a^2+b^2)^{1/2}+2*a)^{1/2})/(2*(a^2+b^2)^{1/2}-2*a)^{1/2})*B*(a^2+b^2)^{1/2}*a+1/4/d/b*\ln((a+b*\tan(d*x+c))^{1/2}*(2*(a^2+b^2)^{1/2}+2*a)^{1/2}-b*\tan(d*x+c)-a-(a^2+b^2)^{1/2})*B*(2*(a^2+b^2)^{1/2}+2*a)^{1/2}*(a^2+b^2)^{1/2}*a^2-1/4/d/b*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^{1/2}*(2*(a^2+b^2)^{1/2}+2*a)^{1/2}+(a^2+b^2)^{1/2})*B*(2*(a^2+b^2)^{1/2}+2*a)^{1/2}*(a^2+b^2)^{1/2}*a^2+2/d*b/(2*(a^2+b^2)^{1/2}-2*a)^{1/2}*\operatorname{arctan}(((2*(a^2+b^2)^{1/2}+2*a)^{1/2}-2*(a+b*\tan(d*x+c))^{1/2})/(2*(a^2+b^2)^{1/2}-2*a)^{1/2})*B*(a^2+b^2)^{1/2}*a-3/d*b^2/(2*(a^2+b^2)^{1/2}-2*a)^{1/2}*\operatorname{arctan}(((2*(a^2+b^2)^{1/2}+2*a)^{1/2}-2*(a+b*\tan(d*x+c))^{1/2})/(2*(a^2+b^2)^{1/2}-2*a)^{1/2})*A*a-3/d*b/(2*(a^2+b^2)^{1/2}-2*a)^{1/2}*\operatorname{arctan}(((2*(a^2+b^2)^{1/2}+2*a)^{1/2}-2*(a+b*\tan(d*x+c))^{1/2})/(2*(a^2+b^2)^{1/2}-2*a)^{1/2})*B*a^2-1/4/d/b*\ln((a+b*\tan(d*x+c))^{1/2}*(2*(a^2+b^2)^{1/2}+2*a)^{1/2}-b*\tan(d*x+c)-a-(a^2+b^2)^{1/2})*B*(2*(a^2+b^2)^{1/2}+2*a)^{1/2}*a^3+1/4/d/b*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^{1/2}*(2*(a^2+b^2)^{1/2}+2*a)^{1/2}+(a^2+b^2)^{1/2})*B*(2*(a^2+b^2)^{1/2}+2*a)^{1/2}*a^3-1/4/d*b^2*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^{1/2}*(2*(a^2+b^2)^{1/2}+2*a)^{1/2}+(a^2+b^2)^{1/2})*A*(2*(a^2+b^2)^{1/2}+2*a)^{1/2}+3/4/d*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^{1/2}*(2*(a^2+b^2)^{1/2}+2*a)^{1/2}+(a^2+b^2)^{1/2}...$

3.335.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4872 vs. $2(146) = 292$.

Time = 10.91 (sec) , antiderivative size = 9757, normalized size of antiderivative = 53.61

$$\int \cot(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.335.6 Sympy [F]

$$\int \cot(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx))(a + b \tan(c + dx))^{5/2} \cot(c + dx) dx$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(5/2)*cot(c + d*x), x)`

3.335.7 Maxima [F]

$$\int \cot(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{5/2} \cot(dx + c) dx$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)*cot(d*x + c), x)`

3.335.8 Giac [F(-1)]

Timed out.

$$\int \cot(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.335.9 Mupad [B] (verification not implemented)

Time = 14.54 (sec) , antiderivative size = 29441, normalized size of antiderivative = 161.76

$$\int \cot(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)
```

```
output ((2*A*b^2 - 2*B*a*b)/d + (6*B*a*b)/d)*(a + b*tan(c + d*x))^(1/2) - atan(((
((32*(3*A^3*a^2*b^16*d^2 + 48*A^3*a^4*b^14*d^2 - 30*A^3*a^6*b^12*d^2 - 72*
A^3*a^8*b^10*d^2 + 3*A^3*a^10*b^8*d^2 + 6*B^3*a^5*b^13*d^2 + 8*B^3*a^7*b^1
1*d^2 + 3*B^3*a^9*b^9*d^2 - B^3*a*b^17*d^2 - A^2*B*a*b^17*d^2 + 3*A*B^2*a^
2*b^16*d^2 - 32*A*B^2*a^4*b^14*d^2 + 46*A*B^2*a^6*b^12*d^2 + 72*A*B^2*a^8*
b^10*d^2 - 9*A*B^2*a^10*b^8*d^2 - 16*A^2*B*a^3*b^15*d^2 + 150*A^2*B*a^5*b^
13*d^2 + 96*A^2*B*a^7*b^11*d^2 - 69*A^2*B*a^9*b^9*d^2))/d^5 - (((32*(4*A*a
*b^12*d^4 + 16*A*a^3*b^10*d^4 + 12*A*a^5*b^8*d^4 + 8*B*a^2*b^11*d^4 + 8*B*
a^4*b^9*d^4))/d^5 - (32*(16*b^10*d^4 + 24*a^2*b^8*d^4)*(a + b*tan(c + d*x)
)^(1/2)*(((8*B^2*a^5*d^2 - 8*A^2*a^5*d^2 + 80*A^2*a^3*b^2*d^2 - 80*B^2*a^
3*b^2*d^2 + 16*A*B*b^5*d^2 - 40*A^2*a*b^4*d^2 + 40*B^2*a*b^4*d^2 - 160*A*B
*a^2*b^3*d^2 + 80*A*B*a^4*b*d^2)^2/64 - d^4*(A^4*a^10 + A^4*b^10 + B^4*a^1
0 + B^4*b^10 + 2*A^2*B^2*a^10 + 2*A^2*B^2*b^10 + 5*A^4*a^2*b^8 + 10*A^4*a^
4*b^6 + 10*A^4*a^6*b^4 + 5*A^4*a^8*b^2 + 5*B^4*a^2*b^8 + 10*B^4*a^4*b^6 +
10*B^4*a^6*b^4 + 5*B^4*a^8*b^2 + 10*A^2*B^2*a^2*b^8 + 20*A^2*B^2*a^4*b^6 +
20*A^2*B^2*a^6*b^4 + 10*A^2*B^2*a^8*b^2)))^(1/2) + A^2*a^5*d^2 - B^2*a^5*d
^2 - 10*A^2*a^3*b^2*d^2 + 10*B^2*a^3*b^2*d^2 - 2*A*B*b^5*d^2 + 5*A^2*a*b^4
*d^2 - 5*B^2*a*b^4*d^2 + 20*A*B*a^2*b^3*d^2 - 10*A*B*a^4*b*d^2)/(4*d^4))^(
1/2))/d^4)*(((8*B^2*a^5*d^2 - 8*A^2*a^5*d^2 + 80*A^2*a^3*b^2*d^2 - 80*B^2
*a^3*b^2*d^2 + 16*A*B*b^5*d^2 - 40*A^2*a*b^4*d^2 + 40*B^2*a*b^4*d^2 - 1...
```

3.336 $\int \cot^2(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.336.1 Optimal result	3191
3.336.2 Mathematica [B] (verified)	3192
3.336.3 Rubi [A] (warning: unable to verify)	3192
3.336.4 Maple [B] (verified)	3198
3.336.5 Fricas [B] (verification not implemented)	3199
3.336.6 Sympy [F(-1)]	3200
3.336.7 Maxima [F]	3200
3.336.8 Giac [F(-1)]	3200
3.336.9 Mupad [B] (verification not implemented)	3201

3.336.1 Optimal result

Integrand size = 33, antiderivative size = 196

$$\int \cot^2(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx =$$

$$\frac{a^{3/2}(5Ab+2aB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{d}$$

$$+ \frac{(a-ib)^{5/2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d}$$

$$- \frac{(a+ib)^{5/2}(iA-B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d}$$

$$+ \frac{b(aA+2bB)\sqrt{a+b \tan(c+dx)}}{d} - \frac{aA \cot(c+dx)(a+b \tan(c+dx))^{3/2}}{d}$$

output

```
-a^(3/2)*(5*A*b+2*B*a)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/d+(a-I*b)^(5/2)*(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d-(a+I*b)^(5/2)*(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d+b*(A*a+2*B*b)*(a+b*tan(d*x+c))^(1/2)/d-a*A*cot(d*x+c)*(a+b*tan(d*x+c))^(3/2)/d
```

3.336.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 400 vs. $2(196) = 392$.

Time = 1.13 (sec) , antiderivative size = 400, normalized size of antiderivative = 2.04

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{2bB \cot(c + dx)(a + b \tan(c + dx))^{3/2}}{d} + 2 \left(-\frac{b(Ab + 4aB) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{d} - 2 \left(-\frac{a^{5/2}(5Ab + 2aB) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a}}\right)}{4d} + \frac{i\sqrt{a - ib}\left(\frac{1}{4}ia\right)}{d} \right) \right)$$

input `Integrate[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(2*b*B*Cot[c + d*x]*(a + b*Tan[c + d*x])^(3/2))/d + 2*(-((b*(A*b + 4*a*B)*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/d) - 2*(-((-1/4*(a^(5/2)*(5*A*b + 2*a*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/d + (I*Sqrt[a - I*b]*((I/4)*a*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B) - (a*(a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B))/4)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/((-a + I*b)*d) - (I*Sqrt[a + I*b]*((-1/4*I)*a*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B) - (a*(a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B))/4)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/((-a - I*b)*d))/a + ((a^2*A - 2*A*b^2 - 6*a*b*B)*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/(4*d))`

3.336.3 Rubi [A] (warning: unable to verify)

Time = 1.78 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.95, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 4088, 27, 3042, 4130, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

$$\begin{aligned}
& \int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan(c + dx)^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\frac{1}{2} \cot(c + dx) \sqrt{a + b \tan(c + dx)} (b(aA + 2bB) \tan^2(c + dx) - 2(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(5Ab + 2aB))}{aA \cot(c + dx)(a + b \tan(c + dx))^{3/2}} dx - \\
& \quad \downarrow \text{4088} \\
& \int \frac{\frac{1}{2} \cot(c + dx) \sqrt{a + b \tan(c + dx)} (b(aA + 2bB) \tan^2(c + dx) - 2(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(5Ab + 2aB))}{aA \cot(c + dx)(a + b \tan(c + dx))^{3/2}} dx - \\
& \quad \downarrow \text{27} \\
& \int \frac{\frac{1}{2} \cot(c + dx) \sqrt{a + b \tan(c + dx)} (b(aA + 2bB) \tan^2(c + dx) - 2(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(5Ab + 2aB))}{aA \cot(c + dx)(a + b \tan(c + dx))^{3/2}} dx - \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \int \frac{\sqrt{a + b \tan(c + dx)} (b(aA + 2bB) \tan(c + dx)^2 - 2(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(5Ab + 2aB))}{aA \cot(c + dx)(a + b \tan(c + dx))^{3/2}} dx - \\
& \quad \downarrow \text{4130} \\
& \frac{1}{2} \left(2 \int \frac{\cot(c + dx) ((5Ab + 2aB)a^2 - b(Aa^2 - 6bBa - 2Ab^2) \tan^2(c + dx) - 2(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx))}{2\sqrt{a + b \tan(c + dx)}} dx - \right. \\
& \quad \left. \frac{aA \cot(c + dx)(a + b \tan(c + dx))^{3/2}}{d} \right) \\
& \quad \downarrow \text{27} \\
& \frac{1}{2} \left(\int \frac{\cot(c + dx) ((5Ab + 2aB)a^2 - b(Aa^2 - 6bBa - 2Ab^2) \tan^2(c + dx) - 2(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx - \right. \\
& \quad \left. \frac{aA \cot(c + dx)(a + b \tan(c + dx))^{3/2}}{d} \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

$$\frac{1}{2} \left(\int \frac{(5Ab + 2aB)a^2 - b(Aa^2 - 6bBa - 2Ab^2) \tan(c + dx)^2 - 2(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\tan(c + dx) \sqrt{a + b \tan(c + dx)}} dx + \frac{aA \cot(c + dx)(a + b \tan(c + dx))^{3/2}}{d} \right)$$

↓ 4136

$$\frac{1}{2} \left(a^2(2aB + 5Ab) \int \frac{\cot(c + dx) (\tan^2(c + dx) + 1)}{\sqrt{a + b \tan(c + dx)}} dx + \int -\frac{2(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3Ab^2a + b^3B)) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{aA \cot(c + dx)(a + b \tan(c + dx))^{3/2}}{d} \right)$$

↓ 27

$$\frac{1}{2} \left(a^2(2aB + 5Ab) \int \frac{\cot(c + dx) (\tan^2(c + dx) + 1)}{\sqrt{a + b \tan(c + dx)}} dx - 2 \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3Ab^2a + b^3B)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{aA \cot(c + dx)(a + b \tan(c + dx))^{3/2}}{d} \right)$$

↓ 3042

$$\frac{1}{2} \left(a^2(2aB + 5Ab) \int \frac{\tan(c + dx)^2 + 1}{\tan(c + dx) \sqrt{a + b \tan(c + dx)}} dx - 2 \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3Ab^2a + b^3B)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{aA \cot(c + dx)(a + b \tan(c + dx))^{3/2}}{d} \right)$$

↓ 4022

$$\frac{1}{2} \left(a^2(2aB + 5Ab) \int \frac{\tan(c + dx)^2 + 1}{\tan(c + dx) \sqrt{a + b \tan(c + dx)}} dx - 2 \left(\frac{1}{2}(a - ib)^3(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a + ib)^3(A + iB) \int \frac{-i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx \right) + \frac{aA \cot(c + dx)(a + b \tan(c + dx))^{3/2}}{d} \right)$$

↓ 3042

$$\frac{1}{2} \left(a^2(2aB + 5Ab) \int \frac{\tan(c + dx)^2 + 1}{\tan(c + dx) \sqrt{a + b \tan(c + dx)}} dx - 2 \left(\frac{1}{2}(a - ib)^3(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a + ib)^3(A + iB) \int \frac{-i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx \right) + \frac{aA \cot(c + dx)(a + b \tan(c + dx))^{3/2}}{d} \right)$$

↓ 4020

$$\begin{aligned}
& -\frac{aA \cot(c+dx)(a+b \tan(c+dx))^{3/2}}{d} + \\
\frac{1}{2} & \left(a^2(2aB+5Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{i(a-ib)^3(A-iB) \int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} dx}{2d} \right) \right) \\
& \quad \downarrow 25 \\
& -\frac{aA \cot(c+dx)(a+b \tan(c+dx))^{3/2}}{d} + \\
\frac{1}{2} & \left(a^2(2aB+5Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{i(a+ib)^3(A+iB) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} dx}{2d} \right) \right) \\
& \quad \downarrow 73 \\
& -\frac{aA \cot(c+dx)(a+b \tan(c+dx))^{3/2}}{d} + \\
\frac{1}{2} & \left(a^2(2aB+5Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{(a-ib)^3(A-iB) \int \frac{1}{i \tan^2 \frac{(c+dx)}{b} + \frac{ia}{b} + 1} dx \sqrt{a+b \tan(c+dx)}}{bd} \right) \right) \\
& \quad \downarrow 221 \\
& -\frac{aA \cot(c+dx)(a+b \tan(c+dx))^{3/2}}{d} + \\
\frac{1}{2} & \left(a^2(2aB+5Ab) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 2 \left(\frac{(a-ib)^{5/2}(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{(a+ib)^5}{d} \right) \right) \\
& \quad \downarrow 4117 \\
& -\frac{aA \cot(c+dx)(a+b \tan(c+dx))^{3/2}}{d} + \\
\frac{1}{2} & \left(\frac{a^2(2aB+5Ab) \int \frac{\cot(c+dx)}{\sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{d} - 2 \left(\frac{(a-ib)^{5/2}(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{(a+ib)^{5/2}(A+ib)}{d} \right) \right) \\
& \quad \downarrow 73 \\
& -\frac{aA \cot(c+dx)(a+b \tan(c+dx))^{3/2}}{d} + \\
\frac{1}{2} & \left(\frac{2a^2(2aB+5Ab) \int \frac{1}{\frac{a+b \tan(c+dx)}{b} - \frac{a}{b}} d \sqrt{a+b \tan(c+dx)}}{bd} - 2 \left(\frac{(a-ib)^{5/2}(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{(a+ib)^5}{d} \right) \right) \\
& \quad \downarrow 221
\end{aligned}$$

$$-\frac{aA \cot(c+dx)(a+b \tan(c+dx))^{3/2}}{d} + \frac{1}{2} \left(-\frac{2a^{3/2}(2aB+5Ab) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{d} - 2 \left(\frac{(a-ib)^{5/2}(A-iB) \operatorname{arctan}\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d} + \frac{(a+ib)^{5/2}(A+iB) \operatorname{arctan}\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d} \right) \right)$$

input `Int[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `-((a*A*Cot[c + d*x]*(a + b*Tan[c + d*x])^(3/2))/d) + (-2*(((a - I*b)^(5/2)*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d) + ((a + I*b)^(5/2)*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d) - (2*a^(3/2)*(5*A*b + 2*a*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]]/d) + (2*b*(a*A + 2*b*B)*Sqrt[a + b*Tan[c + d*x]]/d)/2`

3.336.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((A*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(GtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.336.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2389 vs. $2(168) = 336$.

Time = 0.24 (sec) , antiderivative size = 2390, normalized size of antiderivative = 12.19

method	result	size
derivativedivides	Expression too large to display	2390
default	Expression too large to display	2390

```
input int(cot(d*x+c)^2*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

output $-2/d*a^{(5/2)}*\operatorname{arctanh}((a+b*\tan(dx+c))^{(1/2)}/a^{(1/2)})*B-1/d/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\operatorname{arctan}((2*(a+b*\tan(dx+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*B*a^3-1/d*b^3/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\operatorname{arctan}(((2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-2*(a+b*\tan(dx+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*A+1/d*b^3/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\operatorname{arctan}((2*(a+b*\tan(dx+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*A+1/4/d*b^2*\ln((a+b*\tan(dx+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-b*\tan(dx+c)-a-(a^2+b^2)^{(1/2)})*B*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-1/4/d*b^2*\ln(b*\tan(dx+c)+a+(a+b*\tan(dx+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}+(a^2+b^2)^{(1/2)})*B*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-3/4/d*\ln((a+b*\tan(dx+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-b*\tan(dx+c)-a-(a^2+b^2)^{(1/2)})*B*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*a^2+1/d/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\operatorname{arctan}(((2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-2*(a+b*\tan(dx+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*B*a^3+3/4/d*\ln(b*\tan(dx+c)+a+(a+b*\tan(dx+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}+(a^2+b^2)^{(1/2)})*B*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*a^2+3/d*b/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\operatorname{arctan}(((2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-2*(a+b*\tan(dx+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*A*a^2-3/d*b^2/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\operatorname{arctan}(((2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-2*(a+b*\tan(dx+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*B*a-1/d*b^2/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\operatorname{arctan}((2*(a+b*\tan(dx+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})$

3.336.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4900 vs. $2(162) = 324$.

Time = 13.31 (sec) , antiderivative size = 9815, normalized size of antiderivative = 50.08

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(dx+c)^2*(a+b*tan(dx+c))^(5/2)*(A+B*tan(dx+c)),x, algorithm="fracas")`

output `Too large to include`

3.336. $\int \cot^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.336.6 Sympy [F(-1)]

Timed out.

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**2*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.336.7 Maxima [F]

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{5/2} \cot(dx + c)^2 dx$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^2, x)`

3.336.8 Giac [F(-1)]

Timed out.

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.336.9 Mupad [B] (verification not implemented)

Time = 11.72 (sec) , antiderivative size = 31186, normalized size of antiderivative = 159.11

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)
```

```
output (2*B*b^2*(a + b*tan(c + d*x))^(1/2))/d - atan((((8*(176*A^3*a^5*b^13*d^2
- 400*A^3*a^3*b^15*d^2 + 488*A^3*a^7*b^11*d^2 - 92*A^3*a^9*b^9*d^2 + 12*B^
3*a^2*b^16*d^2 + 192*B^3*a^4*b^14*d^2 - 120*B^3*a^6*b^12*d^2 - 288*B^3*a^8
*b^10*d^2 + 12*B^3*a^10*b^8*d^2 + 4*A^3*a*b^17*d^2 + 4*A*B^2*a*b^17*d^2 +
464*A*B^2*a^3*b^15*d^2 - 920*A*B^2*a^5*b^13*d^2 - 1104*A*B^2*a^7*b^11*d^2
+ 276*A*B^2*a^9*b^9*d^2 + 172*A^2*B*a^2*b^16*d^2 - 1468*A^2*B*a^4*b^14*d^2
- 776*A^2*B*a^6*b^12*d^2 + 828*A^2*B*a^8*b^10*d^2 - 36*A^2*B*a^10*b^8*d^2
))/d^5 - (((8*(16*B*a*b^12*d^4 + 128*A*a^2*b^11*d^4 + 128*A*a^4*b^9*d^4 +
64*B*a^3*b^10*d^4 + 48*B*a^5*b^8*d^4))/d^5 - (16*(32*b^10*d^4 + 48*a^2*b^8
*d^4)*(a + b*tan(c + d*x))^(1/2)*(((8*B^2*a^5*d^2 - 8*A^2*a^5*d^2 + 80*A^
2*a^3*b^2*d^2 - 80*B^2*a^3*b^2*d^2 + 16*A*B*b^5*d^2 - 40*A^2*a*b^4*d^2 + 4
0*B^2*a*b^4*d^2 - 160*A*B*a^2*b^3*d^2 + 80*A*B*a^4*b*d^2)^2/64 - d^4*(A^4*
a^10 + A^4*b^10 + B^4*a^10 + B^4*b^10 + 2*A^2*B^2*a^10 + 2*A^2*B^2*b^10 +
5*A^4*a^2*b^8 + 10*A^4*a^4*b^6 + 10*A^4*a^6*b^4 + 5*A^4*a^8*b^2 + 5*B^4*a^
2*b^8 + 10*B^4*a^4*b^6 + 10*B^4*a^6*b^4 + 5*B^4*a^8*b^2 + 10*A^2*B^2*a^2*b
^8 + 20*A^2*B^2*a^4*b^6 + 20*A^2*B^2*a^6*b^4 + 10*A^2*B^2*a^8*b^2))^1/2)
- A^2*a^5*d^2 + B^2*a^5*d^2 + 10*A^2*a^3*b^2*d^2 - 10*B^2*a^3*b^2*d^2 + 2*
A*B*b^5*d^2 - 5*A^2*a*b^4*d^2 + 5*B^2*a*b^4*d^2 - 20*A*B*a^2*b^3*d^2 + 10*
A*B*a^4*b*d^2)/(4*d^4))^1/2))/d^4)*(((8*B^2*a^5*d^2 - 8*A^2*a^5*d^2 + 80
*A^2*a^3*b^2*d^2 - 80*B^2*a^3*b^2*d^2 + 16*A*B*b^5*d^2 - 40*A^2*a*b^4*d...
```


3.337 $\int \cot^3(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.337.1 Optimal result	3202
3.337.2 Mathematica [B] (verified)	3203
3.337.3 Rubi [A] (warning: unable to verify)	3203
3.337.4 Maple [B] (verified)	3209
3.337.5 Fricas [B] (verification not implemented)	3210
3.337.6 Sympy [F(-1)]	3210
3.337.7 Maxima [F(-1)]	3210
3.337.8 Giac [F(-1)]	3211
3.337.9 Mupad [B] (verification not implemented)	3211

3.337.1 Optimal result

Integrand size = 33, antiderivative size = 220

$$\int \cot^3(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \frac{\sqrt{a}(8a^2A - 15Ab^2 - 20abB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right) + B \tan(c+dx)}{4d} - \frac{(a-ib)^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d} - \frac{(a+ib)^{5/2}(A+iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d} - \frac{a(7Ab + 4aB) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{4d} - \frac{aA \cot^2(c+dx)(a+b \tan(c+dx))^{3/2}}{2d}$$

output

```
-(a-I*b)^(5/2)*(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d-(a+I*b)^(5/2)*(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d+1/4*(8*A*a^2-15*A*b^2-20*B*a*b)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))*a^(1/2)/d-1/4*a*(7*A*b+4*B*a)*cot(d*x+c)*(a+b*tan(d*x+c))^(1/2)/d-1/2*a*A*cot(d*x+c)^2*(a+b*tan(d*x+c))^(3/2)/d
```

3.337.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 448 vs. $2(220) = 440$.

Time = 2.55 (sec) , antiderivative size = 448, normalized size of antiderivative = 2.04

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx =$$

$$-\sqrt{a}(8a^2A - 15Ab^2 - 20abB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right) + 4(a - ib)^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right) +$$

input `Integrate[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `-1/4*(-(Sqrt[a]*(8*a^2*A - 15*A*b^2 - 20*a*b*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]]) + 4*(a - I*b)^(5/2)*(A - I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] + 4*a^2*A*Sqrt[a + I*b]*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + (8*I)*a*A*Sqrt[a + I*b]*b*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] - 4*A*Sqrt[a + I*b]*b^2*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + (4*I)*a^2*Sqrt[a + I*b]*B*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] - 8*a*Sqrt[a + I*b]*b*B*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] - (4*I)*Sqrt[a + I*b]*b^2*B*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + 9*a*A*b*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]] + 4*a^2*B*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]] + 2*a^2*A*Cot[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]))/d`

3.337.3 Rubi [A] (warning: unable to verify)

Time = 1.84 (sec) , antiderivative size = 214, normalized size of antiderivative = 0.97, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

↓ 3042

$$\begin{aligned}
& \int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan(c + dx)^3} dx \\
& \quad \downarrow 4088 \\
& \frac{1}{2} \int \frac{1}{2} \cot^2(c + dx) \sqrt{a + b \tan(c + dx)} (-b(aA - 4bB) \tan^2(c + dx) - 4(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(7Ab + 4aB)) dx - \\
& \quad \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^{3/2}}{2d} \\
& \quad \downarrow 27 \\
& \frac{1}{4} \int \cot^2(c + dx) \sqrt{a + b \tan(c + dx)} (-b(aA - 4bB) \tan^2(c + dx) - 4(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(7Ab + 4aB)) dx - \\
& \quad \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^{3/2}}{2d} \\
& \quad \downarrow 3042 \\
& \frac{1}{4} \int \frac{\sqrt{a + b \tan(c + dx)} (-b(aA - 4bB) \tan(c + dx)^2 - 4(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(7Ab + 4aB))}{\tan(c + dx)^2} dx - \\
& \quad \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^{3/2}}{2d} \\
& \quad \downarrow 4128 \\
& \frac{1}{4} \left(\int - \frac{\cot(c + dx) (b(4Ba^2 + 9Aba - 8b^2B) \tan^2(c + dx) + 8(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(8Aa^2 - 20bE))}{2\sqrt{a + b \tan(c + dx)}} \right. \\
& \quad \left. \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^{3/2}}{2d} \right) \\
& \quad \downarrow 27 \\
& \frac{1}{4} \left(-\frac{1}{2} \int \frac{\cot(c + dx) (b(4Ba^2 + 9Aba - 8b^2B) \tan^2(c + dx) + 8(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(8Aa^2 - 20bE))}{\sqrt{a + b \tan(c + dx)}} \right. \\
& \quad \left. \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^{3/2}}{2d} \right) \\
& \quad \downarrow 3042 \\
& \frac{1}{4} \left(-\frac{1}{2} \int \frac{b(4Ba^2 + 9Aba - 8b^2B) \tan(c + dx)^2 + 8(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(8Aa^2 - 20bE)}{\tan(c + dx) \sqrt{a + b \tan(c + dx)}} \right. \\
& \quad \left. \frac{aA \cot^2(c + dx)(a + b \tan(c + dx))^{3/2}}{2d} \right)
\end{aligned}$$

3.337. $\int \cot^3(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

↓ 4136

$$\frac{1}{4} \left(\frac{1}{2} \left(-a(8a^2A - 20abB - 15Ab^2) \int \frac{\cot(c+dx) (\tan^2(c+dx) + 1)}{\sqrt{a+b\tan(c+dx)}} dx - \int \frac{8(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (a+b)\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx \right) \right. \\ \left. - \frac{aA \cot^2(c+dx)(a+b\tan(c+dx))^{3/2}}{2d} \right)$$

↓ 27

$$\frac{1}{4} \left(\frac{1}{2} \left(-a(8a^2A - 20abB - 15Ab^2) \int \frac{\cot(c+dx) (\tan^2(c+dx) + 1)}{\sqrt{a+b\tan(c+dx)}} dx - 8 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (a+b)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx \right) \right. \\ \left. - \frac{aA \cot^2(c+dx)(a+b\tan(c+dx))^{3/2}}{2d} \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{2} \left(-a(8a^2A - 20abB - 15Ab^2) \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - 8 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (a+b)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx \right) \right. \\ \left. - \frac{aA \cot^2(c+dx)(a+b\tan(c+dx))^{3/2}}{2d} \right)$$

↓ 4022

$$- \frac{aA \cot^2(c+dx)(a+b\tan(c+dx))^{3/2}}{2d} + \\ \frac{1}{4} \left(-\frac{a(4aB + 7Ab) \cot(c+dx)\sqrt{a+b\tan(c+dx)}}{d} + \frac{1}{2} \left(-a(8a^2A - 20abB - 15Ab^2) \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx \right) \right)$$

↓ 3042

$$- \frac{aA \cot^2(c+dx)(a+b\tan(c+dx))^{3/2}}{2d} + \\ \frac{1}{4} \left(-\frac{a(4aB + 7Ab) \cot(c+dx)\sqrt{a+b\tan(c+dx)}}{d} + \frac{1}{2} \left(-a(8a^2A - 20abB - 15Ab^2) \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx \right) \right)$$

↓ 4020

$$- \frac{aA \cot^2(c+dx)(a+b\tan(c+dx))^{3/2}}{2d} + \\ \frac{1}{4} \left(-\frac{a(4aB + 7Ab) \cot(c+dx)\sqrt{a+b\tan(c+dx)}}{d} + \frac{1}{2} \left(-a(8a^2A - 20abB - 15Ab^2) \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx \right) \right)$$

↓ 25

$$\begin{aligned}
& -\frac{aA \cot^2(c+dx)(a+b \tan(c+dx))^{3/2}}{2d} + \\
\frac{1}{4} & \left(-\frac{a(4aB+7Ab) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \frac{1}{2} \left(-a(8a^2A-20abB-15Ab^2) \int \frac{\tan(c+dx)^2 +}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} \right. \right. \\
& \quad \downarrow 73 \\
& -\frac{aA \cot^2(c+dx)(a+b \tan(c+dx))^{3/2}}{2d} + \\
\frac{1}{4} & \left(-\frac{a(4aB+7Ab) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \frac{1}{2} \left(-a(8a^2A-20abB-15Ab^2) \int \frac{\tan(c+dx)^2 +}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} \right. \right. \\
& \quad \downarrow 221 \\
& -\frac{aA \cot^2(c+dx)(a+b \tan(c+dx))^{3/2}}{2d} + \\
\frac{1}{4} & \left(-\frac{a(4aB+7Ab) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \frac{1}{2} \left(-a(8a^2A-20abB-15Ab^2) \int \frac{\tan(c+dx)^2 +}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} \right. \right. \\
& \quad \downarrow 4117 \\
& -\frac{aA \cot^2(c+dx)(a+b \tan(c+dx))^{3/2}}{2d} + \\
\frac{1}{4} & \left(-\frac{a(4aB+7Ab) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \frac{1}{2} \left(-\frac{a(8a^2A-20abB-15Ab^2) \int \frac{\cot(c+dx)}{\sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{d} \right. \right. \\
& \quad \downarrow 73 \\
& -\frac{aA \cot^2(c+dx)(a+b \tan(c+dx))^{3/2}}{2d} + \\
\frac{1}{4} & \left(-\frac{a(4aB+7Ab) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \frac{1}{2} \left(-\frac{2a(8a^2A-20abB-15Ab^2) \int \frac{1}{\frac{a+b \tan(c+dx)}{b} - \frac{a}{b}} d \sqrt{a+b \tan(c+dx)}}{bd} \right. \right. \\
& \quad \downarrow 221 \\
& -\frac{aA \cot^2(c+dx)(a+b \tan(c+dx))^{3/2}}{2d} + \\
\frac{1}{4} & \left(-\frac{a(4aB+7Ab) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} + \frac{1}{2} \left(\frac{2\sqrt{a}(8a^2A-20abB-15Ab^2) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{d} \right. \right.
\end{aligned}$$

input `Int[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

```
output -1/2*(a*A*Cot[c + d*x]^2*(a + b*Tan[c + d*x])^(3/2))/d + ((-8*(((a - I*b)^(5/2)*(I*A + B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]]))/d - ((a + I*b)^(5/2)*(I*A - B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]]))/d + (2*Sqrt[a]*(8*a^2*A - 15*A*b^2 - 20*a*b*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/d)/2 - (a*(7*A*b + 4*a*B)*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/d)/4
```

3.337.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4136 `Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.337.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2487 vs. $2(186) = 372$.

Time = 0.27 (sec) , antiderivative size = 2488, normalized size of antiderivative = 11.31

method	result	size
derivativdivides	Expression too large to display	2488
default	Expression too large to display	2488

```
input int(cot(d*x+c)^3*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 2*a^(5/2)*A*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/d+2/d*b/(2*(a^2+b^2)^(
1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(
1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*(a^2+b^2)^(1/2)*a-1/4/d/b*ln((a+b*t
an(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1
/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*a^2+1/4/d/b*ln(b*tan(
d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1
/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*a^2-2/d*b/(2*(a^2+b^2
)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c)
)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*(a^2+b^2)^(1/2)*a+3/d*b^2/(2*(a^
2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d
*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a+3/d*b/(2*(a^2+b^2)^(1/2)-
2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))
/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*a^2+1/4/d/b*ln((a+b*tan(d*x+c))^(1/2)*(2
*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*B*(2*(a^2+b^2
)^(1/2)+2*a)^(1/2)*a^3-1/4/d/b*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*
(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2
)*a^3+1/4/d*b^2*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2
)+2*a)^(1/2)+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-3/4/d*ln(b*t
an(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)
^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2-1/d/(2*(a^2+b^2)^(1/2)-2*a)...
```


3.337.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4935 vs. $2(180) = 360$.

Time = 16.38 (sec) , antiderivative size = 9888, normalized size of antiderivative = 44.95

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm m="fracas")`

output Too large to include

3.337.6 Sympy [F(-1)]

Timed out.

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**3*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output Timed out

3.337.7 Maxima [F(-1)]

Timed out.

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm m="maxima")`

output Timed out

3.337.8 Giac [F(-1)]

Timed out.

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

```
input integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")
```

```
output Timed out
```

3.337.9 Mupad [B] (verification not implemented)

Time = 12.57 (sec) , antiderivative size = 32561, normalized size of antiderivative = 148.00

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)
```

```
output atan(-(((3708*A^3*a^2*b^16*d^2 - 6912*A^3*a^4*b^14*d^2 - 5820*A^3*a^6*b^12*d^2 + 4608*A^3*a^8*b^10*d^2 - 192*A^3*a^10*b^8*d^2 - 6400*B^3*a^3*b^15*d^2 + 2816*B^3*a^5*b^13*d^2 + 7808*B^3*a^7*b^11*d^2 - 1472*B^3*a^9*b^9*d^2 + 64*B^3*a*b^17*d^2 - 1856*A^2*B*a*b^17*d^2 - 7552*A*B^2*a^2*b^16*d^2 + 23488*A*B^2*a^4*b^14*d^2 + 16256*A*B^2*a^6*b^12*d^2 - 14208*A*B^2*a^8*b^10*d^2 + 576*A*B^2*a^10*b^8*d^2 + 20504*A^2*B*a^3*b^15*d^2 - 5000*A^2*B*a^5*b^13*d^2 - 22944*A^2*B*a^7*b^11*d^2 + 4416*A^2*B*a^9*b^9*d^2)/(2*d^5) - (((1664*A*a*b^12*d^4 + 896*A*a^3*b^10*d^4 - 768*A*a^5*b^8*d^4 + 2048*B*a^2*b^11*d^4 + 2048*B*a^4*b^9*d^4)/(2*d^5) - ((512*b^10*d^4 + 768*a^2*b^8*d^4)*(a + b*tan(c + d*x))^(1/2))*(((8*B^2*a^5*d^2 - 8*A^2*a^5*d^2 + 80*A^2*a^3*b^2*d^2 - 80*B^2*a^3*b^2*d^2 + 16*A*B*b^5*d^2 - 40*A^2*a*b^4*d^2 + 40*B^2*a*b^4*d^2 - 160*A*B*a^2*b^3*d^2 + 80*A*B*a^4*b*d^2)^2/64 - d^4*(A^4*a^10 + A^4*b^10 + B^4*a^10 + B^4*b^10 + 2*A^2*B^2*a^10 + 2*A^2*B^2*b^10 + 5*A^4*a^2*b^8 + 10*A^4*a^4*b^6 + 10*A^4*a^6*b^4 + 5*A^4*a^8*b^2 + 5*B^4*a^2*b^8 + 10*B^4*a^4*b^6 + 10*B^4*a^6*b^4 + 5*B^4*a^8*b^2 + 10*A^2*B^2*a^2*b^8 + 20*A^2*B^2*a^4*b^6 + 20*A^2*B^2*a^6*b^4 + 10*A^2*B^2*a^8*b^2)))^(1/2) + A^2*a^5*d^2 - B^2*a^5*d^2 - 10*A^2*a^3*b^2*d^2 + 10*B^2*a^3*b^2*d^2 - 2*A*B*b^5*d^2 + 5*A^2*a*b^4*d^2 - 5*B^2*a*b^4*d^2 + 20*A*B*a^2*b^3*d^2 - 10*A*B*a^4*b*d^2)/(4*d^4))^(1/2))/d^4)*(((8*B^2*a^5*d^2 - 8*A^2*a^5*d^2 + 80*A^2*a^3*b^2*d^2 - 80*B^2*a^3*b^2*d^2 + 16*A*B*b^5*d^2 - 40*A^2*a*b^4*d^2 + 40*...
```

3.338 $\int \cot^4(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.338.1 Optimal result	3212
3.338.2 Mathematica [A] (verified)	3213
3.338.3 Rubi [A] (warning: unable to verify)	3213
3.338.4 Maple [B] (verified)	3220
3.338.5 Fricas [B] (verification not implemented)	3221
3.338.6 Sympy [F(-1)]	3222
3.338.7 Maxima [F(-1)]	3222
3.338.8 Giac [F(-1)]	3222
3.338.9 Mupad [B] (verification not implemented)	3223

3.338.1 Optimal result

Integrand size = 33, antiderivative size = 277

$$\int \cot^4(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \frac{(40a^2Ab - 5Ab^3 + 16a^3B - 30ab^2B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right) + B \tan(c+dx)}{8\sqrt{a}d} - \frac{(a-ib)^{5/2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d} + \frac{(a+ib)^{5/2}(iA-B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d} + \frac{(8a^2A - 11Ab^2 - 18abB) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{8d} - \frac{a(3Ab + 2aB) \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{4d} - \frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^{3/2}}{3d}$$

output

```
-(a-I*b)^(5/2)*(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d+(a+I*b)^(5/2)*(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d+1/8*(40*A*a^2*b-5*A*b^3+16*B*a^3-30*B*a*b^2)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/d/a^(1/2)+1/8*(8*A*a^2-11*A*b^2-18*B*a*b)*cot(d*x+c)*(a+b*tan(d*x+c))^(1/2)/d-1/4*a*(3*A*b+2*B*a)*cot(d*x+c)^2*(a+b*tan(d*x+c))^(1/2)/d-1/3*a*A*cot(d*x+c)^3*(a+b*tan(d*x+c))^(3/2)/d
```

3.338.2 Mathematica [A] (verified)

Time = 6.49 (sec) , antiderivative size = 548, normalized size of antiderivative = 1.98

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx =$$

$$\frac{2bB \cot^3(c + dx)(a + b \tan(c + dx))^{3/2}}{3d}$$

$$-\frac{2}{3} \left(\frac{3Ab^2 \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{5d} \right) - \frac{2}{5} \left(\frac{(6Ab^2 - 5a(aA - 2bB)) \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{4d} \right)$$

input `Integrate[Cot[c + d*x]^4*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(-2*b*B*Cot[c + d*x]^3*(a + b*Tan[c + d*x])^(3/2))/(3*d) - (2*((3*A*b^2*Cot[c + d*x]^3*Sqrt[a + b*Tan[c + d*x]])/(5*d) - (2*((6*A*b^2 - 5*a*(a*A - 2*b*B))*Cot[c + d*x]^3*Sqrt[a + b*Tan[c + d*x]])/(4*d) - ((15*a*(13*a*A*b + 6*a^2*B - 8*b^2*B))*Cot[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]])/(16*d) - ((-45*a^(5/2)*(40*a^2*A*b - 5*A*b^3 + 16*a^3*B - 30*a*b^2*B))*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/(16*d) + (I*Sqrt[a - I*b]*(((45*I)/2)*a^3*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B) - (45*a^3*(a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B))/2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/((-a + I*b)*d) - (I*Sqrt[a + I*b]*(((45*I)/2)*a^3*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B) - (45*a^3*(a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B))/2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/((-a - I*b)*d))/a + (45*a^2*(8*a^2*A - 11*A*b^2 - 18*a*b*B)*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/(16*d))/(2*a))/(3*a))/5)/3`

3.338.3 Rubi [A] (warning: unable to verify)

Time = 2.46 (sec) , antiderivative size = 276, normalized size of antiderivative = 1.00, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.338. $\int \cot^4(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \int \cot^4(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\tan(c+dx)^4} dx \\
& \quad \downarrow \text{4088} \\
& \frac{1}{3} \int \frac{\frac{3}{2} \cot^3(c+dx) \sqrt{a+b\tan(c+dx)} (-b(aA-2bB)\tan^2(c+dx) - 2(Aa^2-2bBa-Ab^2)\tan(c+dx) + a(3Ab+2aB)) dx - aA \cot^3(c+dx)(a+b\tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{2} \int \frac{\cot^3(c+dx) \sqrt{a+b\tan(c+dx)} (-b(aA-2bB)\tan^2(c+dx) - 2(Aa^2-2bBa-Ab^2)\tan(c+dx) + a(3Ab+2aB)) dx - aA \cot^3(c+dx)(a+b\tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2} \int \frac{\sqrt{a+b\tan(c+dx)} (-b(aA-2bB)\tan(c+dx)^2 - 2(Aa^2-2bBa-Ab^2)\tan(c+dx) + a(3Ab+2aB)) dx - aA \cot^3(c+dx)(a+b\tan(c+dx))^{3/2}}{3d} \\
& \quad \downarrow \text{4128} \\
& \frac{1}{2} \left(\frac{1}{2} \int -\frac{\cot^2(c+dx) (b(6Ba^2+13Aba-8b^2B)\tan^2(c+dx) + 8(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx) + aA \cot^3(c+dx)(a+b\tan(c+dx))^{3/2}}{2\sqrt{a+b\tan(c+dx)}} \right. \\
& \quad \left. \frac{1}{4} \int \frac{\cot^2(c+dx) (b(6Ba^2+13Aba-8b^2B)\tan^2(c+dx) + 8(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx) + aA \cot^3(c+dx)(a+b\tan(c+dx))^{3/2}}{\sqrt{a+b\tan(c+dx)}} \right) \\
& \quad \downarrow \text{27} \\
& \frac{1}{2} \left(-\frac{1}{4} \int \frac{\cot^2(c+dx) (b(6Ba^2+13Aba-8b^2B)\tan^2(c+dx) + 8(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx) + aA \cot^3(c+dx)(a+b\tan(c+dx))^{3/2}}{\sqrt{a+b\tan(c+dx)}} \right. \\
& \quad \left. \frac{1}{4} \int \frac{\cot^2(c+dx) (b(6Ba^2+13Aba-8b^2B)\tan^2(c+dx) + 8(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx) + aA \cot^3(c+dx)(a+b\tan(c+dx))^{3/2}}{\sqrt{a+b\tan(c+dx)}} \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.338. $\int \cot^4(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$

$$\frac{1}{2} \left(-\frac{1}{4} \int \frac{b(6Ba^2 + 13Aba - 8b^2B) \tan(c+dx)^2 + 8(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx) + a(8Aa^2 - 18bBa + 11Ab^2)}{\tan(c+dx)^2 \sqrt{a+b \tan(c+dx)}} \right) \\ \frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^{3/2}}{3d} \\ \downarrow \text{4132}$$

$$\frac{1}{2} \left(\frac{1}{4} \left(\int -\frac{\cot(c+dx)(-ab(8Aa^2 - 18bBa - 11Ab^2) \tan^2(c+dx) - 16a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) + a(16Ba^3 + 40Aba^2 - 30b^2Ba - 5Ab^3))}{2\sqrt{a+b \tan(c+dx)}} \right) \right) \\ \frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^{3/2}}{3d} \\ \downarrow \text{27}$$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{(8a^2A - 18abB - 11Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} - \int \frac{\cot(c+dx)(-ab(8Aa^2 - 18bBa - 11Ab^2) \tan^2(c+dx) - 16a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) + a(16Ba^3 + 40Aba^2 - 30b^2Ba - 5Ab^3))}{2\sqrt{a+b \tan(c+dx)}} \right) \right) \\ \frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^{3/2}}{3d} \\ \downarrow \text{3042}$$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{(8a^2A - 18abB - 11Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} - \int \frac{-ab(8Aa^2 - 18bBa - 11Ab^2) \tan(c+dx)^2 - 16a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) + a(16Ba^3 + 40Aba^2 - 30b^2Ba - 5Ab^3)}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} \right) \right) \\ \frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^{3/2}}{3d} \\ \downarrow \text{4136}$$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{(8a^2A - 18abB - 11Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} - \int -\frac{16(a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + a(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3))}{\sqrt{a+b \tan(c+dx)}} \right) \right) \\ \frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^{3/2}}{3d} \\ \downarrow \text{27}$$

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{(8a^2A - 18abB - 11Ab^2) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{d} - \frac{a(16a^3B + 40a^2Ab - 30ab^2B - 5Ab^3) \int \frac{\cot(c + dx)}{\tan(c + dx)} dx}{\frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^{3/2}}{3d}} \right) \right)$$

↓ 3042

$$\frac{1}{2} \left(\frac{1}{4} \left(\frac{(8a^2A - 18abB - 11Ab^2) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{d} - \frac{a(16a^3B + 40a^2Ab - 30ab^2B - 5Ab^3) \int \frac{\cot(c + dx)}{\tan(c + dx)} dx}{\frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^{3/2}}{3d}} \right) \right)$$

↓ 4022

$$- \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^{3/2}}{3d} + \frac{1}{2} \left(- \frac{a(2aB + 3Ab) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} + \frac{1}{4} \left(\frac{(8a^2A - 18abB - 11Ab^2) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{d} \right) \right)$$

↓ 3042

$$- \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^{3/2}}{3d} + \frac{1}{2} \left(- \frac{a(2aB + 3Ab) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} + \frac{1}{4} \left(\frac{(8a^2A - 18abB - 11Ab^2) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{d} \right) \right)$$

↓ 4020

$$- \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^{3/2}}{3d} + \frac{1}{2} \left(- \frac{a(2aB + 3Ab) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} + \frac{1}{4} \left(\frac{(8a^2A - 18abB - 11Ab^2) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{d} \right) \right)$$

↓ 25

$$- \frac{aA \cot^3(c + dx)(a + b \tan(c + dx))^{3/2}}{3d} + \frac{1}{2} \left(- \frac{a(2aB + 3Ab) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} + \frac{1}{4} \left(\frac{(8a^2A - 18abB - 11Ab^2) \cot(c + dx) \sqrt{a + b \tan(c + dx)}}{d} \right) \right)$$

$$\begin{aligned} & \downarrow 73 \\ & -\frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^{3/2}}{3d} + \\ & \frac{1}{2} \left(-\frac{a(2aB+3Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} + \frac{1}{4} \left(\frac{(8a^2A-18abB-11Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} \right) \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 221 \\ & -\frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^{3/2}}{3d} + \\ & \frac{1}{2} \left(-\frac{a(2aB+3Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} + \frac{1}{4} \left(\frac{(8a^2A-18abB-11Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} \right) \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 4117 \\ & -\frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^{3/2}}{3d} + \\ & \frac{1}{2} \left(-\frac{a(2aB+3Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} + \frac{1}{4} \left(\frac{(8a^2A-18abB-11Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} \right) \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 73 \\ & -\frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^{3/2}}{3d} + \\ & \frac{1}{2} \left(-\frac{a(2aB+3Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} + \frac{1}{4} \left(\frac{(8a^2A-18abB-11Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} \right) \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 221 \\ & -\frac{aA \cot^3(c+dx)(a+b \tan(c+dx))^{3/2}}{3d} + \\ & \frac{1}{2} \left(-\frac{a(2aB+3Ab) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} + \frac{1}{4} \left(\frac{(8a^2A-18abB-11Ab^2) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{d} \right) \right) \end{aligned}$$

input `Int[Cot[c + d*x]^4*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `-1/3*(a*A*Cot[c + d*x]^3*(a + b*Tan[c + d*x])^(3/2))/d + (-1/2*(a*(3*A*b + 2*a*B)*Cot[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]])/d + (-1/2*(-16*((a - I*b)^(5/2)*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]]))/d + (a*(a + I*b)^(5/2)*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d - (2*Sqrt[a]*(40*a^2*A*b - 5*A*b^3 + 16*a^3*B - 30*a*b^2*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/d/a + ((8*a^2*A - 11*A*b^2 - 18*a*b*B)*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/d)/4)/2`

3.338.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.338.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2667 vs. $2(239) = 478$.

Time = 0.25 (sec) , antiderivative size = 2668, normalized size of antiderivative = 9.63

method	result	size
derivativedivides	Expression too large to display	2668
default	Expression too large to display	2668

```
input int(cot(d*x+c)^4*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

output

```
-11/8/d/tan(d*x+c)^3*(a+b*tan(d*x+c))^(5/2)*A+2/d*a^(5/2)*arctanh((a+b*tan
(d*x+c))^(1/2)/a^(1/2))*B+1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b
*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(
1/2))*B*a^3+1/d*b^3/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1
/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A-
1/d*b^3/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*
(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A-1/4/d*b^2*ln(
(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b
^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/4/d*b^2*ln(b*tan(d*x+c)+a+(a+
b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*B*(2*(a
^2+b^2)^(1/2)+2*a)^(1/2)+3/4/d*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2
)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/
2)*a^2-1/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(
1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*a^3-3/4/d*
ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^
2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2-3/d*b/(2*(a^2+b^2)^(1/2
)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2
))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a^2+3/d*b^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/
2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2
+b^2)^(1/2)-2*a)^(1/2))*B*a+1/d*b^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arcta...
```

3.338.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4974 vs. $2(233) = 466$.

Time = 25.11 (sec) , antiderivative size = 9965, normalized size of antiderivative = 35.97

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.338.6 Sympy [F(-1)]

Timed out.

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**4*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.338.7 Maxima [F(-1)]

Timed out.

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `Timed out`

3.338.8 Giac [F(-1)]

Timed out.

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^4*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.338.9 Mupad [B] (verification not implemented)

Time = 11.68 (sec) , antiderivative size = 33949, normalized size of antiderivative = 122.56

$$\int \cot^4(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(cot(c + d*x)^4*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)
```

```
output (atan((((((a + b*tan(c + d*x))^(1/2)*(153*A^4*b^20 + 128*B^4*b^20 + 231*A^
2*B^2*b^20 - 7*A^4*a^2*b^18 + 9895*A^4*a^4*b^16 - 27465*A^4*a^6*b^14 + 263
20*A^4*a^8*b^12 - 832*A^4*a^10*b^10 + 128*A^4*a^12*b^8 - 132*B^4*a^2*b^18
+ 16380*B^4*a^4*b^16 - 25596*B^4*a^6*b^14 + 21060*B^4*a^8*b^12 - 4032*B^4*
a^10*b^10 + 384*B^4*a^12*b^8 + 6811*A^2*B^2*a^2*b^18 - 61315*A^2*B^2*a^4*b
^16 + 184661*A^2*B^2*a^6*b^14 - 121620*A^2*B^2*a^8*b^12 + 23296*A^2*B^2*a^
10*b^10 - 300*A*B^3*a*b^19 + 600*A^3*B*a*b^19 + 17860*A*B^3*a^3*b^17 - 917
00*A*B^3*a^5*b^15 + 110172*A*B^3*a^7*b^13 - 43520*A*B^3*a^9*b^11 + 4352*A*
B^3*a^11*b^9 - 12860*A^3*B*a^3*b^17 + 79680*A^3*B*a^5*b^15 - 126700*A^3*B*
a^7*b^13 + 40960*A^3*B*a^9*b^11 - 1280*A^3*B*a^11*b^9))/(64*d^4) + (((3225
*A^3*a^3*b^15*d^2 - 1088*A^3*a^5*b^13*d^2 - 3984*A^3*a^7*b^11*d^2 + 736*A^
3*a^9*b^9*d^2 + 1854*B^3*a^2*b^16*d^2 - 3456*B^3*a^4*b^14*d^2 - 2910*B^3*a
^6*b^12*d^2 + 2304*B^3*a^8*b^10*d^2 - 96*B^3*a^10*b^8*d^2 + (295*A^2*B*b^1
8*d^2)/2 - 407*A^3*a*b^17*d^2 + 1178*A*B^2*a*b^17*d^2 - 10572*A*B^2*a^3*b^
15*d^2 + 1930*A*B^2*a^5*b^13*d^2 + 11472*A*B^2*a^7*b^11*d^2 - 2208*A*B^2*a
^9*b^9*d^2 - 4716*A^2*B*a^2*b^16*d^2 + (22193*A^2*B*a^4*b^14*d^2)/2 + 8568
*A^2*B*a^6*b^12*d^2 - 7104*A^2*B*a^8*b^10*d^2 + 288*A^2*B*a^10*b^8*d^2)/(1
6*d^5) + (((a + b*tan(c + d*x))^(1/2)*(320*A^2*a^3*b^12*d^2 - 19456*A^2*a
^5*b^10*d^2 + 1280*A^2*a^7*b^8*d^2 - 2320*B^2*a^3*b^12*d^2 + 16896*B^2*a^5
*b^10*d^2 - 2304*B^2*a^7*b^8*d^2 - 2048*A*B*b^15*d^2 + 4764*A^2*a*b^14*...
```

3.339 $\int \cot^5(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.339.1 Optimal result	3224
3.339.2 Mathematica [A] (verified)	3225
3.339.3 Rubi [A] (warning: unable to verify)	3226
3.339.4 Maple [B] (verified)	3235
3.339.5 Fricas [B] (verification not implemented)	3236
3.339.6 Sympy [F(-1)]	3237
3.339.7 Maxima [F(-1)]	3237
3.339.8 Giac [F(-1)]	3237
3.339.9 Mupad [B] (verification not implemented)	3238

3.339.1 Optimal result

Integrand size = 33, antiderivative size = 342

$$\int \cot^5(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx =$$

$$\frac{(128a^4A - 240a^2Ab^2 - 5Ab^4 - 320a^3bB + 40ab^3B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{64a^{3/2}d}$$

$$+ \frac{(a-ib)^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{d}$$

$$+ \frac{(a+ib)^{5/2}(A+iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{d}$$

$$+ \frac{(144a^2Ab - 5Ab^3 + 64a^3B - 88ab^2B) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{64ad}$$

$$+ \frac{(48a^2A - 59Ab^2 - 104abB) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{96d}$$

$$- \frac{a(11Ab + 8aB) \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{24d}$$

$$- \frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^{3/2}}{4d}$$

output
$$-1/64*(128*A*a^4-240*A*a^2*b^2-5*A*b^4-320*B*a^3*b+40*B*a*b^3)*\operatorname{arctanh}((a+b*\tan(dx+c))^{1/2}/a^{1/2})/a^{3/2}/d+(a-I*b)^{5/2}*(A-I*B)*\operatorname{arctanh}((a+b*\tan(dx+c))^{1/2}/(a-I*b)^{1/2})/d+(a+I*b)^{5/2}*(A+I*B)*\operatorname{arctanh}((a+b*\tan(dx+c))^{1/2}/(a+I*b)^{1/2})/d+1/64*(144*A*a^2*b-5*A*b^3+64*B*a^3-88*B*a*b^2)*\cot(dx+c)*(a+b*\tan(dx+c))^{1/2}/a/d+1/96*(48*A*a^2-59*A*b^2-104*B*a*b)*\cot(dx+c)^2*(a+b*\tan(dx+c))^{1/2}/d-1/24*a*(11*A*b+8*B*a)*\cot(dx+c)^3*(a+b*\tan(dx+c))^{1/2}/d-1/4*a*A*\cot(dx+c)^4*(a+b*\tan(dx+c))^{3/2}/d$$

3.339.2 Mathematica [A] (verified)

Time = 6.58 (sec) , antiderivative size = 622, normalized size of antiderivative = 1.82

$$\int \cot^5(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))dx =$$

$$\frac{2bB \cot^4(c+dx)(a+b\tan(c+dx))^{3/2}}{5d}$$

$$-\frac{2}{5} \left(\frac{b(5Ab+2aB) \cot^4(c+dx) \sqrt{a+b\tan(c+dx)}}{7d} \right) - \frac{2}{7} \left(-\frac{(35a^2A-40Ab^2-72abB) \cot^4(c+dx) \sqrt{a+b\tan(c+dx)}}{16d} \right)$$

input `Integrate[Cot[c + d*x]^5*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output $(-2*b*B*Cot[c + d*x]^4*(a + b*Tan[c + d*x])^(3/2))/(5*d) - (2*((b*(5*A*b + 2*a*B)*Cot[c + d*x]^4*Sqrt[a + b*Tan[c + d*x]])/(7*d) - (2*(-1/16*((35*a^2*A - 40*A*b^2 - 72*a*b*B)*Cot[c + d*x]^4*Sqrt[a + b*Tan[c + d*x]])/d - ((7*a*(85*a*A*b + 40*a^2*B - 48*b^2*B)*Cot[c + d*x]^3*Sqrt[a + b*Tan[c + d*x]])/(24*d) - ((35*a^2*(48*a^2*A - 59*A*b^2 - 104*a*b*B)*Cot[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]])/(32*d) - (-(((-105*a^(5/2)*(128*a^4*A - 240*a^2*A*b^2 - 5*A*b^4 - 320*a^3*b*B + 40*a*b^3*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/(32*d) + (I*Sqrt[a - I*b]*(210*a^4*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B) + (210*I)*a^4*(a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B))*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/((-a + I*b)*d) - (I*Sqrt[a + I*b]*(210*a^4*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B) - (210*I)*a^4*(a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B))*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/((-a - I*b)*d))/a - (105*a^2*(144*a^2*A*b - 5*A*b^3 + 64*a^3*B - 88*a*b^2*B)*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/(32*d))/(2*a))/(3*a))/(4*a))/7)/5$

3.339.3 Rubi [A] (warning: unable to verify)

Time = 3.04 (sec) , antiderivative size = 357, normalized size of antiderivative = 1.04, number of steps used = 26, number of rules used = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.758$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan(c + dx)^5} dx$$

$$\downarrow 4088$$

$$\frac{1}{4} \int \frac{1}{2} \cot^4(c + dx) \sqrt{a + b \tan(c + dx)} (-b(5aA - 8bB) \tan^2(c + dx) - 8(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(11Ab + 8aB)) dx - \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}}{4d}$$

$$\downarrow 27$$

$$\frac{1}{8} \int \frac{\cot^4(c + dx) \sqrt{a + b \tan(c + dx)} (-b(5aA - 8bB) \tan^2(c + dx) - 8(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(11Ab + 8aB))}{aA \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}} dx$$

$$\downarrow \text{3042}$$

$$\frac{1}{8} \int \frac{\sqrt{a + b \tan(c + dx)} (-b(5aA - 8bB) \tan(c + dx)^2 - 8(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(11Ab + 8aB))}{aA \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}} dx$$

$$\downarrow \text{4128}$$

$$\frac{1}{8} \left(\frac{1}{3} \int - \frac{\cot^3(c + dx) (b(40Ba^2 + 85Aba - 48b^2B) \tan^2(c + dx) + 48(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx))}{2\sqrt{a + b \tan(c + dx)}} dx \right)$$

$$\downarrow \text{27}$$

$$\frac{1}{8} \left(-\frac{1}{6} \int \frac{\cot^3(c + dx) (b(40Ba^2 + 85Aba - 48b^2B) \tan^2(c + dx) + 48(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx \right)$$

$$\downarrow \text{3042}$$

$$\frac{1}{8} \left(-\frac{1}{6} \int \frac{b(40Ba^2 + 85Aba - 48b^2B) \tan(c + dx)^2 + 48(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(48Aa^2 - 48Ab^2)}{\tan(c + dx)^3 \sqrt{a + b \tan(c + dx)}} dx \right)$$

$$\downarrow \text{4132}$$

$$\frac{1}{8} \left(\frac{1}{6} \left(\int - \frac{3 \cot^2(c + dx) (-ab(48Aa^2 - 104bBa - 59Ab^2) \tan^2(c + dx) - 64a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx) + a(64Ba^3 + 144Aba^2 - 88b^2B))}{2\sqrt{a + b \tan(c + dx)}} dx \right) \right)$$

$$\downarrow \text{27}$$

3.339. $\int \cot^5(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{(48a^2A - 104abB - 59Ab^2) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} - 3 \int \frac{\cot^2(c+dx)(-ab(48Aa^2-104bBa-59Ab^2) \tan^2(c+dx) + 64a^2)}{4d} \right) - \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}}{4d} \right) \downarrow 3042$$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{(48a^2A - 104abB - 59Ab^2) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} - 3 \int \frac{-ab(48Aa^2-104bBa-59Ab^2) \tan(c+dx)^2 - 64a^2}{4d} \right) - \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}}{4d} \right) \downarrow 4132$$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{(48a^2A - 104abB - 59Ab^2) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} - 3 \int \frac{\cot(c+dx)(128Aa^5 - 320bBa^4 - 240Ab^2a^3 + 40b^3Ba^2)}{4d} \right) - \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}}{4d} \right) \downarrow 27$$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{(48a^2A - 104abB - 59Ab^2) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} - 3 \int \frac{\cot(c+dx)(128(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx) + 64a^2)}{4d} \right) - \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}}{4d} \right) \downarrow 3042$$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{(48a^2A - 104abB - 59Ab^2) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} - 3 \left(\frac{\int \frac{128(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) a^2 + b}{\sqrt{a + b \tan(c + dx)}} dx \right) \right) - \frac{aA \cot^4(c + dx) (a + b \tan(c + dx))^{3/2}}{4d} \right) \downarrow 4136$$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{(48a^2A - 104abB - 59Ab^2) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} - 3 \left(\frac{\int \frac{128(a^2(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) - a^2(Aa^3 - 3b^2Ba - Ab^3)) \tan(c + dx) a^2 + b}{\sqrt{a + b \tan(c + dx)}} dx \right) \right) - \frac{aA \cot^4(c + dx) (a + b \tan(c + dx))^{3/2}}{4d} \right) \downarrow 27$$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{(48a^2A - 104abB - 59Ab^2) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} - 3 \left(\frac{128 \int \frac{a^2(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) - a^2(Aa^3 - 3b^2Ba - Ab^3)}{\sqrt{a + b \tan(c + dx)}} dx}{\sqrt{a + b \tan(c + dx)}} \right) \right) - \frac{aA \cot^4(c + dx) (a + b \tan(c + dx))^{3/2}}{4d} \right) \downarrow 3042$$

$$\frac{1}{8} \left(\frac{1}{6} \left(\frac{(48a^2A - 104abB - 59Ab^2) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} - 3 \left(\frac{128 \int \frac{a^2(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) - a^2(Aa^3 - 3b^2Ba - Ab^3)}{\sqrt{a + b \tan(c + dx)}} dx}{\sqrt{a + b \tan(c + dx)}} \right) \right) - \frac{aA \cot^4(c + dx) (a + b \tan(c + dx))^{3/2}}{4d} \right) \downarrow 4022$$

$$-\frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^{3/2}}{4d} + \frac{1}{8} \left(-\frac{a(8aB+11Ab) \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \frac{1}{6} \left(\frac{(48a^2A-104abB-59Ab^2) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} \right) \right)$$

↓ 3042

$$-\frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^{3/2}}{4d} + \frac{1}{8} \left(-\frac{a(8aB+11Ab) \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \frac{1}{6} \left(\frac{(48a^2A-104abB-59Ab^2) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} \right) \right)$$

↓ 4020

$$-\frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^{3/2}}{4d} + \frac{1}{8} \left(-\frac{a(8aB+11Ab) \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \frac{1}{6} \left(\frac{(48a^2A-104abB-59Ab^2) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} \right) \right)$$

↓ 25

$$-\frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^{3/2}}{4d} + \frac{1}{8} \left(-\frac{a(8aB+11Ab) \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \frac{1}{6} \left(\frac{(48a^2A-104abB-59Ab^2) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} \right) \right)$$

↓ 73

$$\begin{aligned}
 & -\frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^{3/2}}{4d} + \\
 & \left(\frac{1}{8} - \frac{a(8aB+11Ab) \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \frac{1}{6} \left(\frac{(48a^2A-104abB-59Ab^2) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} \right) \right)
 \end{aligned}$$

↓ 221

$$\begin{aligned}
 & -\frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^{3/2}}{4d} + \\
 & \left(\frac{1}{8} - \frac{a(8aB+11Ab) \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \frac{1}{6} \left(\frac{(48a^2A-104abB-59Ab^2) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} \right) \right)
 \end{aligned}$$

↓ 4117

$$\begin{aligned}
 & -\frac{aA \cot^4(c+dx)(a+b \tan(c+dx))^{3/2}}{4d} + \\
 & \left(\frac{1}{8} - \frac{a(8aB+11Ab) \cot^3(c+dx) \sqrt{a+b \tan(c+dx)}}{3d} + \frac{1}{6} \left(\frac{(48a^2A-104abB-59Ab^2) \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2d} \right) \right)
 \end{aligned}$$

↓ 73

$$\begin{array}{c}
 \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}}{4d} + \\
 \left(\frac{1}{8} \left[-\frac{a(8aB + 11Ab) \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{3d} + \frac{1}{6} \left[\frac{(48a^2A - 104abB - 59Ab^2) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} \right. \right. \right. \\
 \left. \left. \left. \downarrow 221 \right. \right. \right. \\
 \frac{aA \cot^4(c + dx)(a + b \tan(c + dx))^{3/2}}{4d} + \\
 \left(\frac{1}{8} \left[-\frac{a(8aB + 11Ab) \cot^3(c + dx) \sqrt{a + b \tan(c + dx)}}{3d} + \frac{1}{6} \left[\frac{(48a^2A - 104abB - 59Ab^2) \cot^2(c + dx) \sqrt{a + b \tan(c + dx)}}{2d} \right. \right. \right.
 \end{array}$$

input `Int[Cot[c + d*x]^5*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `-1/4*(a*A*Cot[c + d*x]^4*(a + b*Tan[c + d*x])^(3/2))/d + (-1/3*(a*(11*A*b + 8*a*B)*Cot[c + d*x]^3*Sqrt[a + b*Tan[c + d*x]])/d + (((48*a^2*A - 59*A*b^2 - 104*a*b*B)*Cot[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]])/(2*d) - (3*(-1/2*(128*((a^2*(a - I*b)^(5/2)*(I*A + B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d - (a^2*(a + I*b)^(5/2)*(I*A - B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d) - (2*Sqrt[a]*(128*a^4*A - 240*a^2*A*b^2 - 5*A*b^4 - 320*a^3*b*B + 40*a*b^3*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/d)/a - ((144*a^2*A*b - 5*A*b^3 + 64*a^3*B - 88*a*b^2*B)*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/d)/(4*a))/6)/8`

3.339.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.339.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2913 vs. $2(300) = 600$.

Time = 0.30 (sec) , antiderivative size = 2914, normalized size of antiderivative = 8.52

method	result	size
derivativedivides	Expression too large to display	2914
default	Expression too large to display	2914

```
input int(cot(d*x+c)^5*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

output `-55/24/d/b/tan(d*x+c)^4*B*(a+b*tan(d*x+c))^(3/2)*a^2-2*a^(5/2)*A*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/d-2/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctanh((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*(a^2+b^2)^(1/2)*a+1/4/d/b*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*a^2-1/4/d/b*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*a^2+2/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctanh(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*(a^2+b^2)^(1/2)*a-3/d*b^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctanh(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a-3/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctanh(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*a^2-1/4/d/b*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3+1/4/d/b*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3+5/64/d*b^4/a^(3/2)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))*A+55/192/d/tan(d*x+c)^4*A*(a+b*tan(d*x+c))^(3/2)*a-5/64/d/tan(d*x+c)^4*(a+b*tan(d*x+c))^(1/2)*A*a^2-5/64/d/tan(d*x+c)^4/a*(a+b*tan(d*x+c))^(7/2)*A-1/4/d*b^2*ln(b*tan(d*x+c)+a...`

3.339.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5045 vs. 2(294) = 588.

Time = 55.78 (sec) , antiderivative size = 10107, normalized size of antiderivative = 29.55

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `Too large to include`

3.339.6 Sympy [F(-1)]

Timed out.

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**5*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.339.7 Maxima [F(-1)]

Timed out.

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `Timed out`

3.339.8 Giac [F(-1)]

Timed out.

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^5*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.339.9 Mupad [B] (verification not implemented)

Time = 12.43 (sec) , antiderivative size = 36736, normalized size of antiderivative = 107.42

$$\int \cot^5(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(cot(c + d*x)^5*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)
```

```
output atan(((((((10240*A*a*b^14*d^4 + 436224*A*a^3*b^12*d^4 + 229376*A*a^5*b^10*
d^4 - 196608*A*a^7*b^8*d^4 - 81920*B*a^2*b^13*d^4 + 442368*B*a^4*b^11*d^4
+ 524288*B*a^6*b^9*d^4)/(512*a^2*d^5) - ((131072*a^2*b^10*d^4 + 196608*a^4
*b^8*d^4)*(a + b*tan(c + d*x))^(1/2)*((((8*B^2*a^5*d^2 - 8*A^2*a^5*d^2 + 8
0*A^2*a^3*b^2*d^2 - 80*B^2*a^3*b^2*d^2 + 16*A*B*b^5*d^2 - 40*A^2*a*b^4*d^2
+ 40*B^2*a*b^4*d^2 - 160*A*B*a^2*b^3*d^2 + 80*A*B*a^4*b*d^2)^2/64 - d^4*(
A^4*a^10 + A^4*b^10 + B^4*a^10 + B^4*b^10 + 2*A^2*B^2*a^10 + 2*A^2*B^2*b^1
0 + 5*A^4*a^2*b^8 + 10*A^4*a^4*b^6 + 10*A^4*a^6*b^4 + 5*A^4*a^8*b^2 + 5*B^
4*a^2*b^8 + 10*B^4*a^4*b^6 + 10*B^4*a^6*b^4 + 5*B^4*a^8*b^2 + 10*A^2*B^2*a
^2*b^8 + 20*A^2*B^2*a^4*b^6 + 20*A^2*B^2*a^6*b^4 + 10*A^2*B^2*a^8*b^2)))^(1
/2) + A^2*a^5*d^2 - B^2*a^5*d^2 - 10*A^2*a^3*b^2*d^2 + 10*B^2*a^3*b^2*d^2
- 2*A*B*b^5*d^2 + 5*A^2*a*b^4*d^2 - 5*B^2*a*b^4*d^2 + 20*A*B*a^2*b^3*d^2 -
10*A*B*a^4*b*d^2)/(4*d^4))^(1/2))/(256*a^2*d^4))*((((8*B^2*a^5*d^2 - 8*A^
2*a^5*d^2 + 80*A^2*a^3*b^2*d^2 - 80*B^2*a^3*b^2*d^2 + 16*A*B*b^5*d^2 - 40*
A^2*a*b^4*d^2 + 40*B^2*a*b^4*d^2 - 160*A*B*a^2*b^3*d^2 + 80*A*B*a^4*b*d^2)
^2/64 - d^4*(A^4*a^10 + A^4*b^10 + B^4*a^10 + B^4*b^10 + 2*A^2*B^2*a^10 +
2*A^2*B^2*b^10 + 5*A^4*a^2*b^8 + 10*A^4*a^4*b^6 + 10*A^4*a^6*b^4 + 5*A^4*a
^8*b^2 + 5*B^4*a^2*b^8 + 10*B^4*a^4*b^6 + 10*B^4*a^6*b^4 + 5*B^4*a^8*b^2 +
10*A^2*B^2*a^2*b^8 + 20*A^2*B^2*a^4*b^6 + 20*A^2*B^2*a^6*b^4 + 10*A^2*B^2
*a^8*b^2)))^(1/2) + A^2*a^5*d^2 - B^2*a^5*d^2 - 10*A^2*a^3*b^2*d^2 + 10*...
```

3.340 $\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{5/2} dx$

3.340.1 Optimal result	3239
3.340.2 Mathematica [A] (verified)	3239
3.340.3 Rubi [A] (warning: unable to verify)	3240
3.340.4 Maple [B] (verified)	3243
3.340.5 Fricas [B] (verification not implemented)	3244
3.340.6 Sympy [F]	3245
3.340.7 Maxima [F(-2)]	3246
3.340.8 Giac [F(-1)]	3246
3.340.9 Mupad [B] (verification not implemented)	3246

3.340.1 Optimal result

Integrand size = 27, antiderivative size = 151

$$\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{5/2} dx = \frac{(ia - b)(a - ib)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}}\right) - (a + ib)^{5/2}(ia + b) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a + ib}}\right)}{d} - \frac{2b(a^2 + b^2) \sqrt{a + b \tan(c + dx)}}{d} + \frac{2b(a + b \tan(c + dx))^{5/2}}{5d}$$

```
output (I*a-b)*(a-I*b)^(5/2)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d-(a+I
*b)^(5/2)*(I*a+b)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d-2*b*(a^2
+b^2)*(a+b*tan(d*x+c))^(1/2)/d+2/5*b*(a+b*tan(d*x+c))^(5/2)/d
```

3.340.2 Mathematica [A] (verified)

Time = 1.42 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.28

$$\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{5/2} dx = \frac{\cos(c + dx)(a - b \tan(c + dx)) \left(5i(a - ib)^{5/2}(a + ib) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}}\right) - 5i(a - ib)(a + b \tan(c + dx))^{5/2} \right)}{5d(a \cos(c + dx) - b \sin(c + dx))}$$

input `Integrate[(-a + b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(5/2),x]`

output `(Cos[c + d*x]*(a - b*Tan[c + d*x])*((5*I)*(a - I*b)^(5/2)*(a + I*b)*ArcTan
h[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] - (5*I)*(a - I*b)*(a + I*b)^(5/2
)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]] + 2*b*Sqrt[a + b*Tan[c +
d*x]]*(-4*a^2 - 5*b^2 + 2*a*b*Tan[c + d*x] + b^2*Tan[c + d*x]^2)))/(5*d*(
a*Cos[c + d*x] - b*Sin[c + d*x]))`

3.340.3 Rubi [A] (warning: unable to verify)

Time = 0.69 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.80, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 4011, 27, 3042, 3963, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (b \tan(c + dx) - a)(a + b \tan(c + dx))^{5/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (b \tan(c + dx) - a)(a + b \tan(c + dx))^{5/2} dx \\
 & \quad \downarrow \text{4011} \\
 & \int (-a^2 - b^2)(a + b \tan(c + dx))^{3/2} dx + \frac{2b(a + b \tan(c + dx))^{5/2}}{5d} \\
 & \quad \downarrow \text{27} \\
 & \frac{2b(a + b \tan(c + dx))^{5/2}}{5d} - (a^2 + b^2) \int (a + b \tan(c + dx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{2b(a + b \tan(c + dx))^{5/2}}{5d} - (a^2 + b^2) \int (a + b \tan(c + dx))^{3/2} dx \\
 & \quad \downarrow \text{3963} \\
 & \frac{2b(a + b \tan(c + dx))^{5/2}}{5d} - (a^2 + b^2) \left(\int \frac{a^2 + 2b \tan(c + dx)a - b^2}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2b\sqrt{a + b \tan(c + dx)}}{d} \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{2b(a + b \tan(c + dx))^{5/2}}{5d} - (a^2 + b^2) \left(\int \frac{a^2 + 2b \tan(c + dx)a - b^2}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2b\sqrt{a + b \tan(c + dx)}}{d} \right) \\
& \quad \downarrow 4022 \\
& \frac{2b(a + b \tan(c + dx))^{5/2}}{5d} - \\
& (a^2 + b^2) \left(\frac{1}{2}(a - ib)^2 \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a + ib)^2 \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2b\sqrt{a + b \tan(c + dx)}}{d} \right) \\
& \quad \downarrow 3042 \\
& \frac{2b(a + b \tan(c + dx))^{5/2}}{5d} - \\
& (a^2 + b^2) \left(\frac{1}{2}(a - ib)^2 \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a + ib)^2 \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2b\sqrt{a + b \tan(c + dx)}}{d} \right) \\
& \quad \downarrow 4020 \\
& \frac{2b(a + b \tan(c + dx))^{5/2}}{5d} - \\
& (a^2 + b^2) \left(\frac{i(a - ib)^2 \int -\frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} - \frac{i(a + ib)^2 \int -\frac{1}{(i \tan(c + dx) + 1)\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} \right) \\
& \quad \downarrow 25 \\
& \frac{2b(a + b \tan(c + dx))^{5/2}}{5d} - \\
& (a^2 + b^2) \left(-\frac{i(a - ib)^2 \int \frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} + \frac{i(a + ib)^2 \int \frac{1}{(i \tan(c + dx) + 1)\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} \right) \\
& \quad \downarrow 73 \\
& \frac{2b(a + b \tan(c + dx))^{5/2}}{5d} - \\
& (a^2 + b^2) \left(\frac{(a - ib)^2 \int \frac{1}{\frac{i \tan^2(c + dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)}}{bd} + \frac{(a + ib)^2 \int \frac{1}{-\frac{i \tan^2(c + dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)}}{bd} \right) \\
& \quad \downarrow 221 \\
& \frac{2b(a + b \tan(c + dx))^{5/2}}{5d} - \\
& (a^2 + b^2) \left(\frac{(a - ib)^{3/2} \arctan\left(\frac{\tan(c + dx)}{\sqrt{a - ib}}\right)}{d} + \frac{(a + ib)^{3/2} \arctan\left(\frac{\tan(c + dx)}{\sqrt{a + ib}}\right)}{d} + \frac{2b\sqrt{a + b \tan(c + dx)}}{d} \right)
\end{aligned}$$

input `Int[(-a + b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(5/2),x]`

output `(2*b*(a + b*Tan[c + d*x])^(5/2))/(5*d) - (a^2 + b^2)*(((a - I*b)^(3/2)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/d + ((a + I*b)^(3/2)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/d + (2*b*Sqrt[a + b*Tan[c + d*x]])/d)`

3.340.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n_), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3963 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Int[(a^2 - b^2 + 2*a*b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(n - 2), x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 + b^2, 0] && GtQ[n, 1]`

```
rule 4011 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

3.340.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1374 vs. $2(127) = 254$.

Time = 0.08 (sec) , antiderivative size = 1375, normalized size of antiderivative = 9.11

method	result	size
derivativedivides	Expression too large to display	1375
default	Expression too large to display	1375
parts	Expression too large to display	2381

```
input int((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))^(5/2),x,method=_RETURNVERBOSE)
```

output $2/5*b*(a+b*\tan(d*x+c))^(5/2)/d-2/d*b*(a+b*\tan(d*x+c))^(1/2)*a^2-2/d*b^3*(a+b*\tan(d*x+c))^(1/2)+1/4/d/b*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2)*(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3+1/4/d*b*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2)*(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^4+1/4/d*b^3*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan((2*(a+b*\tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*(a^2+b^2)^(1/2)*a^2+1/d*b^3/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan((2*(a+b*\tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*(a^2+b^2)^(1/2)-2/d*b/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan((2*(a+b*\tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^3-2/d*b^3/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan((2*(a+b*\tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a-1/4/d/b*\ln((a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*\tan(d*x+c)-a-(a^2+b^2)^(1/2))*(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3-1/4/d*b*\ln((a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*\tan(d*x+c)-a-(a^2+b^2)^(1/2))*(a^2+b^2)^(1/2)*(2*(a^...$

3.340.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1472 vs. $2(121) = 242$.

Time = 0.29 (sec) , antiderivative size = 1472, normalized size of antiderivative = 9.75

$$\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{5/2} dx = \text{Too large to display}$$

input `integrate((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))^(5/2),x, algorithm="fracas")`

```

output -1/10*(5*d*sqrt(-(a^7 - a^5*b^2 - 5*a^3*b^4 - 3*a*b^6 + d^2*sqrt(-(9*a^12*
b^2 + 30*a^10*b^4 + 31*a^8*b^6 + 4*a^6*b^8 - 9*a^4*b^10 - 2*a^2*b^12 + b^1
4)/d^4))/d^2)*log(-(3*a^10*b + 11*a^8*b^3 + 14*a^6*b^5 + 6*a^4*b^7 - a^2*b
^9 - b^11)*sqrt(b*tan(d*x + c) + a) + (a*d^3*sqrt(-(9*a^12*b^2 + 30*a^10*b
^4 + 31*a^8*b^6 + 4*a^6*b^8 - 9*a^4*b^10 - 2*a^2*b^12 + b^14)/d^4) + (3*a^
6*b^2 + 5*a^4*b^4 + a^2*b^6 - b^8)*d)*sqrt(-(a^7 - a^5*b^2 - 5*a^3*b^4 - 3
*a*b^6 + d^2*sqrt(-(9*a^12*b^2 + 30*a^10*b^4 + 31*a^8*b^6 + 4*a^6*b^8 - 9*
a^4*b^10 - 2*a^2*b^12 + b^14)/d^4))/d^2)) - 5*d*sqrt(-(a^7 - a^5*b^2 - 5*a
^3*b^4 - 3*a*b^6 + d^2*sqrt(-(9*a^12*b^2 + 30*a^10*b^4 + 31*a^8*b^6 + 4*a^
6*b^8 - 9*a^4*b^10 - 2*a^2*b^12 + b^14)/d^4))/d^2)*log(-(3*a^10*b + 11*a^8
*b^3 + 14*a^6*b^5 + 6*a^4*b^7 - a^2*b^9 - b^11)*sqrt(b*tan(d*x + c) + a) -
(a*d^3*sqrt(-(9*a^12*b^2 + 30*a^10*b^4 + 31*a^8*b^6 + 4*a^6*b^8 - 9*a^4*b
^10 - 2*a^2*b^12 + b^14)/d^4) + (3*a^6*b^2 + 5*a^4*b^4 + a^2*b^6 - b^8)*d)
*sqrt(-(a^7 - a^5*b^2 - 5*a^3*b^4 - 3*a*b^6 + d^2*sqrt(-(9*a^12*b^2 + 30*a
^10*b^4 + 31*a^8*b^6 + 4*a^6*b^8 - 9*a^4*b^10 - 2*a^2*b^12 + b^14)/d^4))/d
^2)) - 5*d*sqrt(-(a^7 - a^5*b^2 - 5*a^3*b^4 - 3*a*b^6 - d^2*sqrt(-(9*a^12*
b^2 + 30*a^10*b^4 + 31*a^8*b^6 + 4*a^6*b^8 - 9*a^4*b^10 - 2*a^2*b^12 + b^1
4)/d^4))/d^2)*log(-(3*a^10*b + 11*a^8*b^3 + 14*a^6*b^5 + 6*a^4*b^7 - a^2*b
^9 - b^11)*sqrt(b*tan(d*x + c) + a) + (a*d^3*sqrt(-(9*a^12*b^2 + 30*a^10*b
^4 + 31*a^8*b^6 + 4*a^6*b^8 - 9*a^4*b^10 - 2*a^2*b^12 + b^14)/d^4) - (3...

```

3.340.6 Sympy [F]

$$\begin{aligned}
& \int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{5/2} dx = \\
& - \int a^3 \sqrt{a + b \tan(c + dx)} dx - \int \left(-b^3 \sqrt{a + b \tan(c + dx)} \tan^3(c + dx) \right) dx \\
& - \int \left(-ab^2 \sqrt{a + b \tan(c + dx)} \tan^2(c + dx) \right) dx \\
& - \int a^2 b \sqrt{a + b \tan(c + dx)} \tan(c + dx) dx
\end{aligned}$$

```

input integrate((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))**(5/2),x)

```

```

output -Integral(a**3*sqrt(a + b*tan(c + d*x)), x) - Integral(-b**3*sqrt(a + b*tan(c + d*x))*tan(c + d*x)**3, x) - Integral(-a*b**2*sqrt(a + b*tan(c + d*x))*tan(c + d*x)**2, x) - Integral(a**2*b*sqrt(a + b*tan(c + d*x))*tan(c + d*x), x)

```

3.340.7 Maxima [F(-2)]

Exception generated.

$$\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{5/2} dx = \text{Exception raised: ValueError}$$

input `integrate((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is`

3.340.8 Giac [F(-1)]

Timed out.

$$\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{5/2} dx = \text{Timed out}$$

input `integrate((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Timed out`

3.340.9 Mupad [B] (verification not implemented)

Time = 32.67 (sec) , antiderivative size = 3441, normalized size of antiderivative = 22.79

$$\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{5/2} dx = \text{Too large to display}$$

input `int(-(a + b*tan(c + d*x))^(5/2)*(a - b*tan(c + d*x)),x)`

output $\log((8a^3b^3(3a^2 - b^2)(a^2 + b^2)^3)/d^3 - (((((-a^4b^2d^4(5a^4 + b^4 - 10a^2b^2)^2)^{(1/2)} - a^7d^2 - 5a^3b^4d^2 + 10a^5b^2d^2)/d^4)^{(1/2)}(64a^2b^5 + 64a^4b^3 + 32ab^2d*((-a^4b^2d^4(5a^4 + b^4 - 10a^2b^2)^2)^{(1/2)} - a^7d^2 - 5a^3b^4d^2 + 10a^5b^2d^2)/d^4)^{(1/2)}(a + b\tan(c + dx))^{(1/2)}))/2d) + (16a^2b^2(a + b\tan(c + dx))^{(1/2)}(a^6 - b^6 + 15a^2b^4 - 15a^4b^2))/d^2 * (((-a^4b^2d^4(5a^4 + b^4 - 10a^2b^2)^2)^{(1/2)} - a^7d^2 - 5a^3b^4d^2 + 10a^5b^2d^2)/d^4)^{(1/2)}/2) * ((20a^6b^8d^4 - a^4b^10d^4 - 110a^8b^6d^4 + 100a^10b^4d^4 - 25a^12b^2d^4)^{(1/2)}/(4d^4) - a^7/(4d^2) - (5a^3b^4)/(4d^2) + (5a^5b^2)/(2d^2))^{(1/2)} - \log((8a^3b^3(3a^2 - b^2)(a^2 + b^2)^3)/d^3 - ((((-((-a^4b^2d^4(5a^4 + b^4 - 10a^2b^2)^2)^{(1/2)} + a^7d^2 + 5a^3b^4d^2 - 10a^5b^2d^2)/d^4)^{(1/2)}(64a^2b^5 + 64a^4b^3 - 32ab^2d*((-a^4b^2d^4(5a^4 + b^4 - 10a^2b^2)^2)^{(1/2)} + a^7d^2 + 5a^3b^4d^2 - 10a^5b^2d^2)/d^4)^{(1/2)}(a + b\tan(c + dx))^{(1/2)}))/2d) - (16a^2b^2(a + b\tan(c + dx))^{(1/2)}(a^6 - b^6 + 15a^2b^4 - 15a^4b^2))/d^2 * (((-a^4b^2d^4(5a^4 + b^4 - 10a^2b^2)^2)^{(1/2)} + a^7d^2 + 5a^3b^4d^2 - 10a^5b^2d^2)/d^4)^{(1/2)}/2) * (-a^7d^2 + (20a^6b^8d^4 - a^4b^10d^4 - 110a^8b^6d^4 + 100a^10b^4d^4 - 25a^12b^2d^4)^{(1/2)} + 5a^3b^4d^2 - 10a^5b^2d^2)/(4d^4))^{(1/2)} - \log((8a^3b^3(3a^2 - b^2)(a^2 + b^2)^3)/d^3 - (((((-a^4b^2d^4(5a^4 ...$

3.341 $\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{3/2} dx$

3.341.1 Optimal result	3248
3.341.2 Mathematica [C] (verified)	3249
3.341.3 Rubi [A] (warning: unable to verify)	3249
3.341.4 Maple [B] (verified)	3254
3.341.5 Fricas [B] (verification not implemented)	3255
3.341.6 Sympy [F]	3256
3.341.7 Maxima [F(-2)]	3256
3.341.8 Giac [F(-1)]	3256
3.341.9 Mupad [B] (verification not implemented)	3257

3.341.1 Optimal result

Integrand size = 27, antiderivative size = 408

$$\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{3/2} dx =$$

$$-\frac{b(a^2 + b^2) \operatorname{arctanh}\left(\frac{\sqrt{a + \sqrt{a^2 + b^2}} - \sqrt{2}\sqrt{a + b \tan(c + dx)}}{\sqrt{a - \sqrt{a^2 + b^2}}}\right)}{\sqrt{2}\sqrt{a - \sqrt{a^2 + b^2}}d}$$

$$+ \frac{b(a^2 + b^2) \operatorname{arctanh}\left(\frac{\sqrt{a + \sqrt{a^2 + b^2}} + \sqrt{2}\sqrt{a + b \tan(c + dx)}}{\sqrt{a - \sqrt{a^2 + b^2}}}\right)}{\sqrt{2}\sqrt{a - \sqrt{a^2 + b^2}}d}$$

$$- \frac{b(a^2 + b^2) \log\left(a + \sqrt{a^2 + b^2} + b \tan(c + dx) - \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}}\sqrt{a + b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}}d}$$

$$+ \frac{b(a^2 + b^2) \log\left(a + \sqrt{a^2 + b^2} + b \tan(c + dx) + \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}}\sqrt{a + b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}}d}$$

$$+ \frac{2b(a + b \tan(c + dx))^{3/2}}{3d}$$

output
$$\begin{aligned} & -1/2*b*(a^2+b^2)*\operatorname{arctanh}(((a+(a^2+b^2)^{(1/2)})^{(1/2)}-2^{(1/2)}*(a+b*\tan(dx+c))^{(1/2)})/(a-(a^2+b^2)^{(1/2)})^{(1/2)})/d*2^{(1/2)}/(a-(a^2+b^2)^{(1/2)})^{(1/2)}+ \\ & 1/2*b*(a^2+b^2)*\operatorname{arctanh}(((a+(a^2+b^2)^{(1/2)})^{(1/2)}+2^{(1/2)}*(a+b*\tan(dx+c))^{(1/2)})/(a-(a^2+b^2)^{(1/2)})^{(1/2)})/d*2^{(1/2)}/(a-(a^2+b^2)^{(1/2)})^{(1/2)}-1/4 \\ & *b*(a^2+b^2)*\ln(a+(a^2+b^2)^{(1/2)}-2^{(1/2)}*(a+(a^2+b^2)^{(1/2)})^{(1/2)}*(a+b*\tan(dx+c))^{(1/2)}+b*\tan(dx+c))/d*2^{(1/2)}/(a+(a^2+b^2)^{(1/2)})^{(1/2)}+1/4*b*(\\ & a^2+b^2)*\ln(a+(a^2+b^2)^{(1/2)}+2^{(1/2)}*(a+(a^2+b^2)^{(1/2)})^{(1/2)}*(a+b*\tan(dx+c))^{(1/2)}+b*\tan(dx+c))/d*2^{(1/2)}/(a+(a^2+b^2)^{(1/2)})^{(1/2)}+2/3*b*(a+b* \\ & \tan(dx+c))^{(3/2)}/d \end{aligned}$$

3.341.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.51 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.45

$$\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{3/2} dx = \frac{(a - b \tan(c + dx)) \left(3i\sqrt{a - ib}(a^2 + b^2) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}}\right) \cos(c + dx) - 3i\sqrt{a + ib}(a^2 + b^2) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a + ib}}\right) \cos(c + dx) \right)}{3d(a \cos(c + dx) + b \sin(c + dx))}$$

input `Integrate[(-a + b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(3/2), x]`

output
$$\begin{aligned} & ((a - b*\tan[c + d*x])*((3*I)*\sqrt{a - I*b}*(a^2 + b^2)*\operatorname{ArcTanh}[\sqrt{a + b* \\ & \tan[c + d*x]}/\sqrt{a - I*b}]*\cos[c + d*x] - (3*I)*\sqrt{a + I*b}*(a^2 + b^2) \\ &)*\operatorname{ArcTanh}[\sqrt{a + b*\tan[c + d*x]}/\sqrt{a + I*b}]*\cos[c + d*x] + 2*b*(a*\cos[c + d*x] + b*\sin[c + d*x])* \\ & \sqrt{a + b*\tan[c + d*x]})/(3*d*(a*\cos[c + d*x] - b*\sin[c + d*x])) \end{aligned}$$

3.341.3 Rubi [A] (warning: unable to verify)

Time = 0.72 (sec) , antiderivative size = 432, normalized size of antiderivative = 1.06, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.481$, Rules used = {3042, 4011, 27, 3042, 3966, 483, 1449, 1142, 25, 27, 1083, 219, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.341. $\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{3/2} dx$

$$\begin{aligned}
 & \int (b \tan(c + dx) - a)(a + b \tan(c + dx))^{3/2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (b \tan(c + dx) - a)(a + b \tan(c + dx))^{3/2} dx \\
 & \quad \downarrow \text{4011} \\
 & \int (-a^2 - b^2) \sqrt{a + b \tan(c + dx)} dx + \frac{2b(a + b \tan(c + dx))^{3/2}}{3d} \\
 & \quad \downarrow \text{27} \\
 & \frac{2b(a + b \tan(c + dx))^{3/2}}{3d} - (a^2 + b^2) \int \sqrt{a + b \tan(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{2b(a + b \tan(c + dx))^{3/2}}{3d} - (a^2 + b^2) \int \sqrt{a + b \tan(c + dx)} dx \\
 & \quad \downarrow \text{3966} \\
 & \frac{2b(a + b \tan(c + dx))^{3/2}}{3d} - \frac{b(a^2 + b^2) \int \frac{\sqrt{a + b \tan(c + dx)}}{\tan^2(c + dx) b^2 + b^2} d(b \tan(c + dx))}{d} \\
 & \quad \downarrow \text{483} \\
 & \frac{2b(a + b \tan(c + dx))^{3/2}}{3d} - \frac{2b(a^2 + b^2) \int \frac{b^2 \tan^2(c + dx)}{b^4 \tan^4(c + dx) - 2ab^2 \tan^2(c + dx) + a^2 + b^2} d\sqrt{a + b \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{1449} \\
 & \frac{2b(a + b \tan(c + dx))^{3/2}}{3d} - \\
 & 2b(a^2 + b^2) \left(\frac{\int \frac{\sqrt{a + b \tan(c + dx)}}{b^2 \tan^2(c + dx) - \sqrt{2b\sqrt{a + \sqrt{a^2 + b^2} \tan(c + dx)} + \sqrt{a^2 + b^2}} d\sqrt{a + b \tan(c + dx)}}{2\sqrt{2}\sqrt{\sqrt{a^2 + b^2} + a}} - \frac{\int \frac{\sqrt{a + b \tan(c + dx)}}{b^2 \tan^2(c + dx) + \sqrt{2b\sqrt{a + \sqrt{a^2 + b^2} \tan(c + dx)} + \sqrt{a^2 + b^2}} d\sqrt{a + b \tan(c + dx)}}{2\sqrt{2}\sqrt{\sqrt{a^2 + b^2} + a}} \right) \\
 & \quad \downarrow \text{1142} \\
 & \frac{2b(a + b \tan(c + dx))^{3/2}}{3d} - \\
 & 2b(a^2 + b^2) \left(\frac{\sqrt{\sqrt{a^2 + b^2} + a} \int \frac{1}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}} \sqrt{a + b \tan(c + dx)}} d\sqrt{a + b \tan(c + dx)}}{\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}(\sqrt{a + \sqrt{a^2 + b^2}} - \sqrt{2}\sqrt{a + b \tan(c + dx)})}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}}} d\sqrt{a + b \tan(c + dx)}}{2\sqrt{2}\sqrt{\sqrt{a^2 + b^2} + a}} \right)
 \end{aligned}$$

3.341. $\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{3/2} dx$

$$\begin{array}{c}
 \downarrow 25 \\
 \frac{2b(a + b \tan(c + dx))^{3/2}}{3d} - \\
 2b(a^2 + b^2) \left(\frac{\int \frac{\sqrt{a^2 + b^2} + a}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2} \sqrt{a + \sqrt{a^2 + b^2}} \sqrt{a + b \tan(c + dx)}} dx - \frac{1}{2} \int \frac{\sqrt{2} (\sqrt{a + \sqrt{a^2 + b^2}} - \sqrt{2} \sqrt{a + b \tan(c + dx)})}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2} \sqrt{a + \sqrt{a^2 + b^2}}} dx \right) \\
 \frac{2\sqrt{2} \sqrt{a^2 + b^2} + a}
 \end{array}$$

$$\begin{array}{c}
 \downarrow 27 \\
 \frac{2b(a + b \tan(c + dx))^{3/2}}{3d} - \\
 2b(a^2 + b^2) \left(\frac{\int \frac{\sqrt{a^2 + b^2} + a}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2} \sqrt{a + \sqrt{a^2 + b^2}} \sqrt{a + b \tan(c + dx)}} dx - \int \frac{\sqrt{a + \sqrt{a^2 + b^2}} - \sqrt{2} \sqrt{a + b \tan(c + dx)}}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2} \sqrt{a + \sqrt{a^2 + b^2}}} dx \right) \\
 \frac{2\sqrt{2} \sqrt{a^2 + b^2} + a}
 \end{array}$$

$$\begin{array}{c}
 \downarrow 1083 \\
 \frac{2b(a + b \tan(c + dx))^{3/2}}{3d} - \\
 2b(a^2 + b^2) \left(\frac{-\sqrt{2} \sqrt{a^2 + b^2} + a \int \frac{1}{2(a - \sqrt{a^2 + b^2}) - b^2 \tan^2(c + dx)} d(2\sqrt{a + b \tan(c + dx)} - \sqrt{2} \sqrt{a + \sqrt{a^2 + b^2}}) - \int \frac{\sqrt{a + \sqrt{a^2 + b^2}} - \sqrt{2} \sqrt{a + b \tan(c + dx)}}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2} \sqrt{a + \sqrt{a^2 + b^2}}} dx \right) \\
 \frac{2\sqrt{2} \sqrt{a^2 + b^2} + a}
 \end{array}$$

$$\begin{array}{c}
 \downarrow 219 \\
 \frac{2b(a + b \tan(c + dx))^{3/2}}{3d} - \\
 2b(a^2 + b^2) \left(\frac{\int \frac{\sqrt{a + \sqrt{a^2 + b^2}} - \sqrt{2} \sqrt{a + b \tan(c + dx)}}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2} \sqrt{a + \sqrt{a^2 + b^2}} \sqrt{a + b \tan(c + dx)}} dx - \frac{\sqrt{a^2 + b^2} + a \operatorname{arctanh} \left(\frac{2\sqrt{a + b \tan(c + dx)} - \sqrt{2} \sqrt{a + \sqrt{a^2 + b^2}}}{\sqrt{2} \sqrt{a - \sqrt{a^2 + b^2}}} \right)}{\sqrt{a - \sqrt{a^2 + b^2}}} \right) \\
 \frac{2\sqrt{2} \sqrt{a^2 + b^2} + a}
 \end{array}$$

\downarrow 1103

3.341. $\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{3/2} dx$

$$2b(a^2 + b^2) \left(\frac{2b(a + b \tan(c + dx))^{3/2}}{3d} - \frac{\frac{1}{2} \log\left(-\sqrt{2}\sqrt{a^2+b^2+a}\sqrt{a+b \tan(c+dx)}+\sqrt{a^2+b^2+b^2 \tan^2(c+dx)}\right)}{2\sqrt{2}\sqrt{a^2+b^2+a}} - \frac{\sqrt{\sqrt{a^2+b^2+a}} \operatorname{arctanh}\left(\frac{2\sqrt{a+b \tan(c+dx)}-\sqrt{2}\sqrt{\sqrt{a^2+b^2+a}}}{\sqrt{2}\sqrt{a-\sqrt{a^2+b^2+a}}}\right)}{\sqrt{a-\sqrt{a^2+b^2+a}}}\right)$$

input `Int[(-a + b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(3/2),x]`

output `(-2*b*(a^2 + b^2)*((-((Sqrt[a + Sqrt[a^2 + b^2]]*ArcTanh[(-(Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]) + 2*Sqrt[a + b*Tan[c + d*x]])/(Sqrt[2]*Sqrt[a - Sqrt[a^2 + b^2]])]/Sqrt[a - Sqrt[a^2 + b^2]]) + Log[Sqrt[a^2 + b^2] + b^2*Tan[c + d*x]^2 - Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]*Sqrt[a + b*Tan[c + d*x]]]/2)/(2*Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]) - ((Sqrt[a + Sqrt[a^2 + b^2]]*ArcTanh[(Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]] + 2*Sqrt[a + b*Tan[c + d*x]])/(Sqrt[2]*Sqrt[a - Sqrt[a^2 + b^2]])]/Sqrt[a - Sqrt[a^2 + b^2]]) + Log[Sqrt[a^2 + b^2] + b^2*Tan[c + d*x]^2 + Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]*Sqrt[a + b*Tan[c + d*x]]/2)/(2*Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]))/d + (2*b*(a + b*Tan[c + d*x])^(3/2))/(3*d)`

3.341.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 483 `Int[Sqrt[(c_) + (d_.)*(x_)]/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[2*d Subst[Int[x^2/(b*c^2 + a*d^2 - 2*b*c*x^2 + b*x^4), x], x, Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x]`

rule 1083 $\text{Int}[(a_.) + (b_.)x + (c_.)x^2]^{-1}, x_Symbol] \rightarrow \text{Simp}[-2 \text{ Subst}[\text{Int}[1/\text{Simp}[b^2 - 4ac - x^2], x], x, b + 2cx], x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d_.) + (e_.)x]/(a_.) + (b_.)x + (c_.)x^2, x_Symbol] \rightarrow \text{Simp}[d(\text{Log}[\text{RemoveContent}[a + bx + cx^2], x])/b], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2cd - be, 0]$

rule 1142 $\text{Int}[(d_.) + (e_.)x]/(a_.) + (b_.)x + (c_.)x^2, x_Symbol] \rightarrow \text{Simp}[(2cd - be)/(2c) \text{ Int}[1/(a + bx + cx^2), x], x] + \text{Simp}[e/(2c) \text{ Int}[(b + 2cx)/(a + bx + cx^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

rule 1449 $\text{Int}[x^m]/(a_.) + (b_.)x^2 + (c_.)x^4, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a/c, 2]\}, \text{With}[\{r = \text{Rt}[2q - b/c, 2]\}, \text{Simp}[1/(2cr) \text{ Int}[x^{m-1}/(q - rx + x^2), x], x] - \text{Simp}[1/(2cr) \text{ Int}[x^{m-1}/(q + rx + x^2), x], x]]] /; \text{FreeQ}[\{a, b, c\}, x] \&\& \text{NeQ}[b^2 - 4ac, 0] \&\& \text{GeQ}[m, 1] \&\& \text{LtQ}[m, 3] \&\& \text{NegQ}[b^2 - 4ac]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 3966 $\text{Int}[(a_.) + (b_.)\tan[(c_.) + (d_.)x]]^{n_}, x_Symbol] \rightarrow \text{Simp}[b/d \text{ Subst}[\text{Int}[(a + x)^n/(b^2 + x^2), x], x, b\tan[c + dx]], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \&\& \text{NeQ}[a^2 + b^2, 0]$

rule 4011 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]]^{m_}((c_.) + (d_.)\tan[(e_.) + (f_.)x]), x_Symbol] \rightarrow \text{Simp}[d((a + b\tan[e + fx])^m/(f^m)), x] + \text{Int}[(a + b\tan[e + fx])^{m-1} \text{Simp}[a^2c - b^2d + (b^2c + a^2d)\tan[e + fx], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b^2c - a^2d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{GtQ}[m, 0]$

3.341.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 985 vs. 2(333) = 666.

Time = 0.09 (sec) , antiderivative size = 986, normalized size of antiderivative = 2.42

method	result
derivativedivides	$\frac{2b(a+b \tan(dx+c))^{\frac{3}{2}}}{3d} + \frac{\ln(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}) \sqrt{2\sqrt{a^2+b^2}+2a} \sqrt{a^2+b^2} a^2}{4bd}$
default	$\frac{2b(a+b \tan(dx+c))^{\frac{3}{2}}}{3d} + \frac{\ln(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}) \sqrt{2\sqrt{a^2+b^2}+2a} \sqrt{a^2+b^2} a^2}{4bd}$
parts	Expression too large to display

input `int((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & 2/3*b*(a+b*\tan(d*x+c))^(3/2)/d+1/4/b/d*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)* \\ & (2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)* \\ & (a^2+b^2)^(1/2)*a^2+1/4*b/d*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)* \\ & (2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)* \\ & (a^2+b^2)^(1/2)-b/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan(((2*(a+b*\tan(d*x+c))^(1/2)+ \\ & (2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)))*a^2-b^3/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)* \\ & \arctan(((2*(a+b*\tan(d*x+c))^(1/2)+2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))- \\ & 1/4/b/d*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)* \\ & (2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+ \\ & (a^2+b^2)^(1/2)* \\ & (2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3-1/4*b/d*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)* \\ & (2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)* \\ & a-1/4/b/d*\ln((a+b*\tan(d*x+c))^(1/2)* \\ & (2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*\tan(d*x+c)-a-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)* \\ & (a^2+b^2)^(1/2)*a^2-1/4*b/d*\ln((a+b*\tan(d*x+c))^(1/2)* \\ & (2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*\tan(d*x+c)-a-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)* \\ & (a^2+b^2)^(1/2)+b/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)* \\ & (2*(a+b*\tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))* \\ & a^2+b^3/d/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*\tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))+ \\ & 1/4/b/d*\ln((a+b*\tan(d*x+c))^(1/2)* \\ & (2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*\tan(d*x+c)-a-(a^2+... \end{aligned}$$

3.341. $\int(-a+b \tan(c+dx))(a+b \tan(c+dx))^{3/2} dx$

3.341.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 956 vs. $2(335) = 670$.

Time = 0.28 (sec) , antiderivative size = 956, normalized size of antiderivative = 2.34

$$\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{3/2} dx = \frac{3d \sqrt{-\frac{a^5 + 2a^3b^2 + ab^4 + d^2 \sqrt{-\frac{a^8b^2 + 4a^6b^4 + 6a^4b^6 + 4a^2b^8 + b^{10}}{d^4}}}{d^2}}}{\log\left(d^3 \sqrt{-\frac{a^5 + 2a^3b^2 + ab^4 + d^2 \sqrt{-\frac{a^8b^2 + 4a^6b^4 + 6a^4b^6 + 4a^2b^8 + b^{10}}{d^4}}}{d^2}}\right)} + \dots$$

input `integrate((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))^(3/2),x, algorithm="fricas")`

output

```
1/6*(3*d*sqrt(-(a^5 + 2*a^3*b^2 + a*b^4 + d^2*sqrt(-(a^8*b^2 + 4*a^6*b^4 + 6*a^4*b^6 + 4*a^2*b^8 + b^10)/d^4))/d^2)*log(d^3*sqrt(-(a^5 + 2*a^3*b^2 + a*b^4 + d^2*sqrt(-(a^8*b^2 + 4*a^6*b^4 + 6*a^4*b^6 + 4*a^2*b^8 + b^10)/d^4))/d^2)*sqrt(-(a^8*b^2 + 4*a^6*b^4 + 6*a^4*b^6 + 4*a^2*b^8 + b^10)/d^4) + (a^6*b + 3*a^4*b^3 + 3*a^2*b^5 + b^7)*sqrt(b*tan(d*x + c) + a)) - 3*d*sqrt(-(a^5 + 2*a^3*b^2 + a*b^4 + d^2*sqrt(-(a^8*b^2 + 4*a^6*b^4 + 6*a^4*b^6 + 4*a^2*b^8 + b^10)/d^4))/d^2)*log(-d^3*sqrt(-(a^5 + 2*a^3*b^2 + a*b^4 + d^2*sqrt(-(a^8*b^2 + 4*a^6*b^4 + 6*a^4*b^6 + 4*a^2*b^8 + b^10)/d^4))/d^2)*sqrt(-(a^8*b^2 + 4*a^6*b^4 + 6*a^4*b^6 + 4*a^2*b^8 + b^10)/d^4) + (a^6*b + 3*a^4*b^3 + 3*a^2*b^5 + b^7)*sqrt(b*tan(d*x + c) + a)) - 3*d*sqrt(-(a^5 + 2*a^3*b^2 + a*b^4 - d^2*sqrt(-(a^8*b^2 + 4*a^6*b^4 + 6*a^4*b^6 + 4*a^2*b^8 + b^10)/d^4))/d^2)*log(d^3*sqrt(-(a^5 + 2*a^3*b^2 + a*b^4 - d^2*sqrt(-(a^8*b^2 + 4*a^6*b^4 + 6*a^4*b^6 + 4*a^2*b^8 + b^10)/d^4))/d^2)*sqrt(-(a^8*b^2 + 4*a^6*b^4 + 6*a^4*b^6 + 4*a^2*b^8 + b^10)/d^4) + (a^6*b + 3*a^4*b^3 + 3*a^2*b^5 + b^7)*sqrt(b*tan(d*x + c) + a)) + 3*d*sqrt(-(a^5 + 2*a^3*b^2 + a*b^4 - d^2*sqrt(-(a^8*b^2 + 4*a^6*b^4 + 6*a^4*b^6 + 4*a^2*b^8 + b^10)/d^4))/d^2)*log(-d^3*sqrt(-(a^5 + 2*a^3*b^2 + a*b^4 - d^2*sqrt(-(a^8*b^2 + 4*a^6*b^4 + 6*a^4*b^6 + 4*a^2*b^8 + b^10)/d^4))/d^2)*sqrt(-(a^8*b^2 + 4*a^6*b^4 + 6*a^4*b^6 + 4*a^2*b^8 + b^10)/d^4) + (a^6*b + 3*a^4*b^3 + 3*a^2*b^5 + b^7)*sqrt(b*tan(d*x + c) + a)) + 4*(b^2*tan(d*x + c) + a*b)*sqrt(b*tan(...
```

3.341.6 Sympy [F]

$$\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{3/2} dx =$$

$$- \int a^2 \sqrt{a + b \tan(c + dx)} dx - \int \left(-b^2 \sqrt{a + b \tan(c + dx)} \tan^2(c + dx) \right) dx$$

input `integrate((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))**(3/2),x)`

output `-Integral(a**2*sqrt(a + b*tan(c + d*x)), x) - Integral(-b**2*sqrt(a + b*tan(c + d*x))*tan(c + d*x)**2, x)`

3.341.7 Maxima [F(-2)]

Exception generated.

$$\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{3/2} dx = \text{Exception raised: ValueError}$$

input `integrate((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is`

3.341.8 Giac [F(-1)]

Timed out.

$$\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{3/2} dx = \text{Timed out}$$

input `integrate((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `Timed out`

3.341.9 Mupad [B] (verification not implemented)

Time = 19.90 (sec) , antiderivative size = 2529, normalized size of antiderivative = 6.20

$$\int (-a + b \tan(c + dx))(a + b \tan(c + dx))^{3/2} dx = \text{Too large to display}$$

```
input int(-(a + b*tan(c + d*x))^(3/2)*(a - b*tan(c + d*x)),x)
```

```
output log((((16*b^4*(a + b*tan(c + d*x))^(1/2)*(a^4 + b^4 - 6*a^2*b^2))/d^2 + (1
6*a*b^2*(((b^6*d^4*(3*a^2 - b^2)^2)^(1/2) - 3*a*b^4*d^2 + a^3*b^2*d^2)/d^
4)^(1/2)*(a^2*b + b^3 - d*(((b^6*d^4*(3*a^2 - b^2)^2)^(1/2) - 3*a*b^4*d^2
+ a^3*b^2*d^2)/d^4)^(1/2)*(a + b*tan(c + d*x))^(1/2)))/d)*(((b^6*d^4*(3*
a^2 - b^2)^2)^(1/2) - 3*a*b^4*d^2 + a^3*b^2*d^2)/d^4)^(1/2))/2 - (8*b^5*(a
^2 - b^2)*(a^2 + b^2)^2)/d^3)*((6*a^2*b^8*d^4 - b^10*d^4 - 9*a^4*b^6*d^4)^
(1/2)/(4*d^4) - (3*a*b^4)/(4*d^2) + (a^3*b^2)/(4*d^2))^(1/2) - log(- (((16
*b^4*(a + b*tan(c + d*x))^(1/2)*(a^4 + b^4 - 6*a^2*b^2))/d^2 - (16*a*b^2*(
-((b^6*d^4*(3*a^2 - b^2)^2)^(1/2) + 3*a*b^4*d^2 - a^3*b^2*d^2)/d^4)^(1/2)
*(a^2*b + b^3 + d*(-((b^6*d^4*(3*a^2 - b^2)^2)^(1/2) + 3*a*b^4*d^2 - a^3*
b^2*d^2)/d^4)^(1/2)*(a + b*tan(c + d*x))^(1/2)))/d)*(-((b^6*d^4*(3*a^2 -
b^2)^2)^(1/2) + 3*a*b^4*d^2 - a^3*b^2*d^2)/d^4)^(1/2))/2 - (8*b^5*(a^2 - b
^2)*(a^2 + b^2)^2)/d^3)*(-((6*a^2*b^8*d^4 - b^10*d^4 - 9*a^4*b^6*d^4)^(1/2
) + 3*a*b^4*d^2 - a^3*b^2*d^2)/(4*d^4))^(1/2) - log(- (((16*b^4*(a + b*tan
(c + d*x))^(1/2)*(a^4 + b^4 - 6*a^2*b^2))/d^2 - (16*a*b^2*(((b^6*d^4*(3*a
^2 - b^2)^2)^(1/2) - 3*a*b^4*d^2 + a^3*b^2*d^2)/d^4)^(1/2)*(a^2*b + b^3 +
d*(((b^6*d^4*(3*a^2 - b^2)^2)^(1/2) - 3*a*b^4*d^2 + a^3*b^2*d^2)/d^4)^(1/
2)*(a + b*tan(c + d*x))^(1/2)))/d)*(((b^6*d^4*(3*a^2 - b^2)^2)^(1/2) - 3*
a*b^4*d^2 + a^3*b^2*d^2)/d^4)^(1/2))/2 - (8*b^5*(a^2 - b^2)*(a^2 + b^2)^2
/d^3)*(((6*a^2*b^8*d^4 - b^10*d^4 - 9*a^4*b^6*d^4)^(1/2) - 3*a*b^4*d^2 ...
```


3.342 $\int (-a + b \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx$

3.342.1 Optimal result	3258
3.342.2 Mathematica [C] (verified)	3259
3.342.3 Rubi [A] (warning: unable to verify)	3259
3.342.4 Maple [B] (verified)	3264
3.342.5 Fricas [B] (verification not implemented)	3265
3.342.6 Sympy [F]	3265
3.342.7 Maxima [F(-2)]	3266
3.342.8 Giac [F(-1)]	3266
3.342.9 Mupad [B] (verification not implemented)	3267

3.342.1 Optimal result

Integrand size = 27, antiderivative size = 422

$$\begin{aligned}
 & \int (-a + b \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx \\
 = & -\frac{b\sqrt{a^2 + b^2} \operatorname{arctanh}\left(\frac{\sqrt{a + \sqrt{a^2 + b^2}} - \sqrt{2}\sqrt{a + b \tan(c + dx)}}{\sqrt{a - \sqrt{a^2 + b^2}}}\right)}{\sqrt{2}\sqrt{a - \sqrt{a^2 + b^2}}d} \\
 & + \frac{b\sqrt{a^2 + b^2} \operatorname{arctanh}\left(\frac{\sqrt{a + \sqrt{a^2 + b^2}} + \sqrt{2}\sqrt{a + b \tan(c + dx)}}{\sqrt{a - \sqrt{a^2 + b^2}}}\right)}{\sqrt{2}\sqrt{a - \sqrt{a^2 + b^2}}d} \\
 & + \frac{b\sqrt{a^2 + b^2} \log\left(a + \sqrt{a^2 + b^2} + b \tan(c + dx) - \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}}\sqrt{a + b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}}d} \\
 & - \frac{b\sqrt{a^2 + b^2} \log\left(a + \sqrt{a^2 + b^2} + b \tan(c + dx) + \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}}\sqrt{a + b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}}d} \\
 & + \frac{2b\sqrt{a + b \tan(c + dx)}}{d}
 \end{aligned}$$

output
$$\begin{aligned} & -1/2*b*\operatorname{arctanh}\left(\left(\frac{a+(a^2+b^2)^{1/2}}{a-(a^2+b^2)^{1/2}}\right)^{1/2}-2^{1/2}\frac{(a+b*\tan(dx+c))^{1/2}}{(a-(a^2+b^2)^{1/2})^{1/2}}\right)/ \\ & \left(\frac{a+(a^2+b^2)^{1/2}}{a-(a^2+b^2)^{1/2}}\right)^{1/2}\frac{(a^2+b^2)^{1/2}/d*2^{1/2}}{(a-(a^2+b^2)^{1/2})^{1/2}} \\ & +1/2*b*\operatorname{arctanh}\left(\left(\frac{a+(a^2+b^2)^{1/2}}{a-(a^2+b^2)^{1/2}}\right)^{1/2}+2^{1/2}\frac{(a+b*\tan(dx+c))^{1/2}}{(a-(a^2+b^2)^{1/2})^{1/2}}\right)/ \\ & \left(\frac{a+(a^2+b^2)^{1/2}}{a-(a^2+b^2)^{1/2}}\right)^{1/2}\frac{(a^2+b^2)^{1/2}/d*2^{1/2}}{(a-(a^2+b^2)^{1/2})^{1/2}} \\ & +1/4*b*\ln\left(\frac{a+(a^2+b^2)^{1/2}-2^{1/2}\frac{(a+(a^2+b^2)^{1/2})^{1/2}}{(a+b*\tan(dx+c))^{1/2}}+b*\tan(dx+c)}{a+(a^2+b^2)^{1/2}}\right) \\ & -1/4*b*\ln\left(\frac{a+(a^2+b^2)^{1/2}+2^{1/2}\frac{(a+(a^2+b^2)^{1/2})^{1/2}}{(a+b*\tan(dx+c))^{1/2}}+b*\tan(dx+c)}{a+(a^2+b^2)^{1/2}}\right) \\ & +2*b*(a+b*\tan(dx+c))^{1/2}/d \end{aligned}$$

3.342.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.37

$$\int (-a + b \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx$$

$$= \frac{\cos(c + dx)(a - b \tan(c + dx)) \left(i\sqrt{a - ib}(a + ib) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}}\right) - i(a - ib)\sqrt{a + ib} \operatorname{arctanh}\left(\frac{\sqrt{a - b \tan(c + dx)}}{\sqrt{a + ib}}\right) \right)}{d(a \cos(c + dx) - b \sin(c + dx))}$$

input `Integrate[(-a + b*Tan[c + d*x])*Sqrt[a + b*Tan[c + d*x]],x]`

output
$$\begin{aligned} & (\operatorname{Cos}[c + d*x]*(a - b*\operatorname{Tan}[c + d*x])*(I*\operatorname{Sqrt}[a - I*b]*(a + I*b)*\operatorname{ArcTanh}[\operatorname{Sqrt}[\\ & [a + b*\operatorname{Tan}[c + d*x]]/\operatorname{Sqrt}[a - I*b]] - I*(a - I*b)*\operatorname{Sqrt}[a + I*b]*\operatorname{ArcTanh}[\operatorname{Sqrt}[\\ & [a + b*\operatorname{Tan}[c + d*x]]/\operatorname{Sqrt}[a + I*b]] + 2*b*\operatorname{Sqrt}[a + b*\operatorname{Tan}[c + d*x]])/(d* \\ & (a*\operatorname{Cos}[c + d*x] - b*\operatorname{Sin}[c + d*x])) \end{aligned}$$

3.342.3 Rubi [A] (warning: unable to verify)

Time = 0.73 (sec) , antiderivative size = 453, normalized size of antiderivative = 1.07, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.481$, Rules used = {3042, 4011, 27, 3042, 3966, 484, 1407, 1142, 25, 27, 1083, 219, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (b \tan(c + dx) - a) \sqrt{a + b \tan(c + dx)} dx$$

3.342. $\int (-a + b \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \int (b \tan(c + dx) - a) \sqrt{a + b \tan(c + dx)} dx \\
 & \downarrow 4011 \\
 & \int \frac{-a^2 - b^2}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2b\sqrt{a + b \tan(c + dx)}}{d} \\
 & \downarrow 27 \\
 & \frac{2b\sqrt{a + b \tan(c + dx)}}{d} - (a^2 + b^2) \int \frac{1}{\sqrt{a + b \tan(c + dx)}} dx \\
 & \downarrow 3042 \\
 & \frac{2b\sqrt{a + b \tan(c + dx)}}{d} - (a^2 + b^2) \int \frac{1}{\sqrt{a + b \tan(c + dx)}} dx \\
 & \downarrow 3966 \\
 & \frac{2b\sqrt{a + b \tan(c + dx)}}{d} - \frac{b(a^2 + b^2) \int \frac{1}{\sqrt{a + b \tan(c + dx)} (\tan^2(c + dx) b^2 + b^2)} d(b \tan(c + dx))}{d} \\
 & \downarrow 484 \\
 & \frac{2b\sqrt{a + b \tan(c + dx)}}{d} - \frac{2b(a^2 + b^2) \int \frac{1}{b^4 \tan^4(c + dx) - 2ab^2 \tan^2(c + dx) + a^2 + b^2} d\sqrt{a + b \tan(c + dx)}}{d} \\
 & \downarrow 1407 \\
 & \frac{2b\sqrt{a + b \tan(c + dx)}}{d} - \\
 & 2b(a^2 + b^2) \left(\frac{\int \frac{\sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}} - \sqrt{a + b \tan(c + dx)}}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}} \sqrt{a + b \tan(c + dx)}} d\sqrt{a + b \tan(c + dx)}}{2\sqrt{2}\sqrt{a^2 + b^2} \sqrt{\sqrt{a^2 + b^2} + a}} + \frac{\int \frac{\sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}} + \sqrt{a + b \tan(c + dx)}}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} + \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}} \sqrt{a + b \tan(c + dx)}}}{2\sqrt{2}\sqrt{a^2 + b^2} \sqrt{\sqrt{a^2 + b^2} - a}} \right) \\
 & \downarrow 1142 \\
 & \frac{2b\sqrt{a + b \tan(c + dx)}}{d} - \\
 & 2b(a^2 + b^2) \left(\frac{\sqrt{a^2 + b^2} + a \int \frac{1}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}} \sqrt{a + b \tan(c + dx)}} d\sqrt{a + b \tan(c + dx)}}{\sqrt{2}} - \frac{1}{2} \int \frac{\sqrt{2}(\sqrt{a + \sqrt{a^2 + b^2}} - \sqrt{2}\sqrt{a + b \tan(c + dx)})}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}} \sqrt{a + b \tan(c + dx)}}}{2\sqrt{2}\sqrt{a^2 + b^2} \sqrt{\sqrt{a^2 + b^2} + a}} \right) \\
 & \downarrow 25
 \end{aligned}$$

3.342. $\int (-a + b \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx$

$$2b(a^2 + b^2) \left(\frac{\frac{2b\sqrt{a + b \tan(c + dx)}}{d}}{\frac{\sqrt{\sqrt{a^2 + b^2} + a} \int \frac{1}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}} \sqrt{a + b \tan(c + dx)}}{\sqrt{2}} d\sqrt{a + b \tan(c + dx)} + \frac{1}{2} \int \frac{\sqrt{2}(\sqrt{a + \sqrt{a^2 + b^2}} - \sqrt{2}\sqrt{a + b \tan(c + dx)})}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}} \sqrt{a + b \tan(c + dx)}}{\sqrt{2}} dx} \right)$$

27

$$2b(a^2 + b^2) \left(\frac{\frac{2b\sqrt{a + b \tan(c + dx)}}{d}}{\frac{\sqrt{\sqrt{a^2 + b^2} + a} \int \frac{1}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}} \sqrt{a + b \tan(c + dx)}}{\sqrt{2}} d\sqrt{a + b \tan(c + dx)} + \int \frac{\sqrt{a + \sqrt{a^2 + b^2}} - \sqrt{2}\sqrt{a + b \tan(c + dx)}}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}} \sqrt{a + b \tan(c + dx)}}{\sqrt{2}} dx} \right)$$

1083

$$2b(a^2 + b^2) \left(\frac{\frac{2b\sqrt{a + b \tan(c + dx)}}{d}}{\frac{\int \frac{\sqrt{a + \sqrt{a^2 + b^2}} - \sqrt{2}\sqrt{a + b \tan(c + dx)}}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}} \sqrt{a + b \tan(c + dx)}}{\sqrt{2}} d\sqrt{a + b \tan(c + dx)} - \sqrt{2}\sqrt{\sqrt{a^2 + b^2} + a} \int \frac{1}{2(a - \sqrt{a^2 + b^2}) - b^2 \tan^2(c + dx)} d(2\sqrt{a + b \tan(c + dx)})}{2\sqrt{2}\sqrt{a^2 + b^2} \sqrt{\sqrt{a^2 + b^2} + a}}$$

219

$$2b(a^2 + b^2) \left(\frac{\frac{2b\sqrt{a + b \tan(c + dx)}}{d}}{\frac{\int \frac{\sqrt{a + \sqrt{a^2 + b^2}} - \sqrt{2}\sqrt{a + b \tan(c + dx)}}{b^2 \tan^2(c + dx) + \sqrt{a^2 + b^2} - \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}} \sqrt{a + b \tan(c + dx)}}{\sqrt{2}} d\sqrt{a + b \tan(c + dx)} - \frac{\sqrt{\sqrt{a^2 + b^2} + a} \operatorname{arctanh} \left(\frac{2\sqrt{a + b \tan(c + dx)} - \sqrt{2}\sqrt{\sqrt{a^2 + b^2} + a}}{\sqrt{2}\sqrt{a - \sqrt{a^2 + b^2}}} \right)}{\sqrt{a - \sqrt{a^2 + b^2}}}}{2\sqrt{2}\sqrt{a^2 + b^2} \sqrt{\sqrt{a^2 + b^2} + a}}$$

1103

$$2b(a^2 + b^2) \left(\frac{2b\sqrt{a + b \tan(c + dx)}}{d} - \frac{\frac{\sqrt{\sqrt{a^2+b^2}+a} \operatorname{arctanh}\left(\frac{2\sqrt{a+b \tan(c+dx)} - \sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}}{\sqrt{2}\sqrt{a-\sqrt{a^2+b^2}}}\right)}{\sqrt{a-\sqrt{a^2+b^2}}} - \frac{1}{2} \log\left(-\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}\sqrt{a+b \tan(c+dx)} + \sqrt{a^2+b^2} + b^2 \tan^2(c+dx)\right)}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{\sqrt{a^2+b^2}+a}} \right)$$

d

input `Int[(-a + b*Tan[c + d*x])*Sqrt[a + b*Tan[c + d*x]],x]`

output `(-2*b*(a^2 + b^2)*((-((Sqrt[a + Sqrt[a^2 + b^2]]*ArcTanh[(-(Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]) + 2*Sqrt[a + b*Tan[c + d*x]])/(Sqrt[2]*Sqrt[a - Sqrt[a^2 + b^2]])]/Sqrt[a - Sqrt[a^2 + b^2]]) - Log[Sqrt[a^2 + b^2] + b^2*Tan[c + d*x]^2 - Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]*Sqrt[a + b*Tan[c + d*x]]]/2)/(2*Sqrt[2]*Sqrt[a^2 + b^2]*Sqrt[a + Sqrt[a^2 + b^2]]) + (-((Sqrt[a + Sqrt[a^2 + b^2]]*ArcTanh[(Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]] + 2*Sqrt[a + b*Tan[c + d*x]])/(Sqrt[2]*Sqrt[a - Sqrt[a^2 + b^2]])]/Sqrt[a - Sqrt[a^2 + b^2]]) + Log[Sqrt[a^2 + b^2] + b^2*Tan[c + d*x]^2 + Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]*Sqrt[a + b*Tan[c + d*x]]]/2)/(2*Sqrt[2]*Sqrt[a^2 + b^2]*Sqrt[a + Sqrt[a^2 + b^2]]))/d + (2*b*Sqrt[a + b*Tan[c + d*x]])/d`

3.342.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 484 `Int[1/(Sqrt[(c_) + (d_.)*(x_)]*((a_) + (b_.)*(x_)^2)), x_Symbol] := Simp[2*d Subst[Int[1/(b*c^2 + a*d^2 - 2*b*c*x^2 + b*x^4), x], x, Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x]`

rule 1083 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1142 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`

rule 1407 `Int[((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(-1), x_Symbol] := With[{q = Rt[a/c, 2]}, With[{r = Rt[2*q - b/c, 2]}, Simp[1/(2*c*q*r) Int[(r - x)/(q - r*x + x^2), x], x] + Simp[1/(2*c*q*r) Int[(r + x)/(q + r*x + x^2), x], x]]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && NegQ[b^2 - 4*a*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3966 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[(a + x)^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[a^2 + b^2, 0]`

rule 4011 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

3.342.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 813 vs. 2(341) = 682.

Time = 0.07 (sec) , antiderivative size = 814, normalized size of antiderivative = 1.93

method	result
parts	$b \left(\frac{1}{2\sqrt{a+b \tan(dx+c)}} - \frac{\sqrt{2\sqrt{a^2+b^2}+2a} \ln \left(\frac{b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}}{4} \right)}{4} \right) + \frac{(a-\sqrt{a^2+b^2}) \arctan \left(\frac{b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}}{4} \right)}{4}$
derivativedivides	Expression too large to display
default	Expression too large to display

input `int((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & b/d*(2*(a+b*\tan(d*x+c))^{(1/2)}-1/4*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*\ln(b*\tan(d \\ & *x+c)+a+(a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}+(a^2+b^2)^{(1/ \\ & 2)))+(a-(a^2+b^2)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan((2*(a+b*\tan(d \\ & *x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}) \\ & +1/4*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*\ln((a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(\\ & 1/2)}+2*a)^{(1/2)}-b*\tan(d*x+c)-a-(a^2+b^2)^{(1/2)})+((a^2+b^2)^{(1/2)}-a)/(2*(a \\ & ^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan(((2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-2*(a+b*\tan(\\ & d*x+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}))-1/4/b/d*\ln(b*\tan(d*x+c)+a+(\\ & a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}+(a^2+b^2)^{(1/2)})*(2*(a \\ & ^2+b^2)^{(1/2)}+2*a)^{(1/2)}*a^2+1/4/b/d*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^{(1 \\ & /2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}+(a^2+b^2)^{(1/2)})*(a^2+b^2)^{(1/2)}*(2*(a^2 \\ & +b^2)^{(1/2)}+2*a)^{(1/2)}*a-b/d/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan((2*(a+b* \\ & \tan(d*x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(\\ & 1/2)})*a+1/4/d/b*\ln((a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-b* \\ & \tan(d*x+c)-a-(a^2+b^2)^{(1/2)})*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*a^2-1/4/d/b*\ln \\ & ((a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-b*\tan(d*x+c)-a-(a^2+ \\ & b^2)^{(1/2)})*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*(a^2+b^2)^{(1/2)}*a+1/d*b/(2*(a^2+ \\ & b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan(((2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-2*(a+b*\tan(d*x \\ & +c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*a \end{aligned}$$

3.342.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 719 vs. $2(343) = 686$.

Time = 0.27 (sec) , antiderivative size = 719, normalized size of antiderivative = 1.70

$$\int (-a + b \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx =$$

$$d \sqrt{-\frac{a^3 + ab^2 + d^2 \sqrt{-\frac{a^4 b^2 + 2a^2 b^4 + b^6}{d^4}}}{d^2}} \log \left((a^4 b + 2a^2 b^3 + b^5) \sqrt{b \tan(dx + c) + a} + \left(ad^3 \sqrt{-\frac{a^4 b^2 + 2a^2 b^4 + b^6}{d^4}} + \right) \right)$$

```
input integrate((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))^(1/2),x, algorithm="fricas")
```

```
output -1/2*(d*sqrt(-(a^3 + a*b^2 + d^2*sqrt(-(a^4*b^2 + 2*a^2*b^4 + b^6)/d^4))/d
^2)*log((a^4*b + 2*a^2*b^3 + b^5)*sqrt(b*tan(d*x + c) + a) + (a*d^3*sqrt(-
(a^4*b^2 + 2*a^2*b^4 + b^6)/d^4) + (a^2*b^2 + b^4)*d)*sqrt(-(a^3 + a*b^2 +
d^2*sqrt(-(a^4*b^2 + 2*a^2*b^4 + b^6)/d^4))/d^2)) - d*sqrt(-(a^3 + a*b^2
+ d^2*sqrt(-(a^4*b^2 + 2*a^2*b^4 + b^6)/d^4))/d^2)*log((a^4*b + 2*a^2*b^3
+ b^5)*sqrt(b*tan(d*x + c) + a) - (a*d^3*sqrt(-(a^4*b^2 + 2*a^2*b^4 + b^6)
/d^4) + (a^2*b^2 + b^4)*d)*sqrt(-(a^3 + a*b^2 + d^2*sqrt(-(a^4*b^2 + 2*a^2
*b^4 + b^6)/d^4))/d^2)) - d*sqrt(-(a^3 + a*b^2 - d^2*sqrt(-(a^4*b^2 + 2*a^
2*b^4 + b^6)/d^4))/d^2)*log((a^4*b + 2*a^2*b^3 + b^5)*sqrt(b*tan(d*x + c)
+ a) + (a*d^3*sqrt(-(a^4*b^2 + 2*a^2*b^4 + b^6)/d^4) - (a^2*b^2 + b^4)*d)*
sqrt(-(a^3 + a*b^2 - d^2*sqrt(-(a^4*b^2 + 2*a^2*b^4 + b^6)/d^4))/d^2)) + d
*sqrt(-(a^3 + a*b^2 - d^2*sqrt(-(a^4*b^2 + 2*a^2*b^4 + b^6)/d^4))/d^2)*log
((a^4*b + 2*a^2*b^3 + b^5)*sqrt(b*tan(d*x + c) + a) - (a*d^3*sqrt(-(a^4*b^
2 + 2*a^2*b^4 + b^6)/d^4) - (a^2*b^2 + b^4)*d)*sqrt(-(a^3 + a*b^2 - d^2*sq
rt(-(a^4*b^2 + 2*a^2*b^4 + b^6)/d^4))/d^2)) - 4*sqrt(b*tan(d*x + c) + a)*b
)/d
```

3.342.6 Sympy [F]

$$\int (-a + b \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx$$

$$= - \int a \sqrt{a + b \tan(c + dx)} dx - \int \left(-b \sqrt{a + b \tan(c + dx)} \tan(c + dx) \right) dx$$

```
input integrate((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))**(1/2),x)
```

3.342. $\int (-a + b \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx$

output `-Integral(a*sqrt(a + b*tan(c + d*x)), x) - Integral(-b*sqrt(a + b*tan(c + d*x))*tan(c + d*x), x)`

3.342.7 Maxima [F(-2)]

Exception generated.

$$\int (-a + b \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx = \text{Exception raised: ValueError}$$

input `integrate((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is`

3.342.8 Giac [F(-1)]

Timed out.

$$\int (-a + b \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx = \text{Timed out}$$

input `integrate((-a+b*tan(d*x+c))*(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.342.9 Mupad [B] (verification not implemented)

Time = 10.21 (sec) , antiderivative size = 581, normalized size of antiderivative = 1.38

$$\begin{aligned}
& \int (-a + b \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx \\
&= \operatorname{atanh} \left(\frac{d^3 \left(\frac{16(a^2 b^4 - a^4 b^2) \sqrt{a + b \tan(c + dx)}}{d^2} + \frac{16 a b^2 (a^3 + 1 i b a^2) \sqrt{a + b \tan(c + dx)}}{d^2} \right) \sqrt{-\frac{a^3 + 1 i b a^2}{d^2}}}{16 (a^5 b^3 + a^3 b^5)} \right) \sqrt{-\frac{a^3 + 1 i b a^2}{d^2}} \\
&+ \operatorname{atanh} \left(\frac{d^3 \sqrt{-\frac{a^3 + a^2 b 1 i}{d^2}} \left(\frac{16(a^2 b^4 - a^4 b^2) \sqrt{a + b \tan(c + dx)}}{d^2} - \frac{16 a b^2 (-a^3 + a^2 b 1 i) \sqrt{a + b \tan(c + dx)}}{d^2} \right)}{16 (a^5 b^3 + a^3 b^5)} \right) \sqrt{\frac{-a^3 + a^2 b 1 i}{d^2}} \\
&+ \frac{2 b \sqrt{a + b \tan(c + dx)}}{d} - \operatorname{atan} \left(\frac{b^6 \sqrt{\frac{a b^2}{4 d^2} - \frac{b^3 1 i}{4 d^2}} \sqrt{a + b \tan(c + dx)} 32 i}{\frac{b^8 16 i}{d} + \frac{a^2 b^6 16 i}{d}} \right. \\
&\quad \left. + \frac{32 a b^5 \sqrt{\frac{a b^2}{4 d^2} - \frac{b^3 1 i}{4 d^2}} \sqrt{a + b \tan(c + dx)}}{\frac{b^8 16 i}{d} + \frac{a^2 b^6 16 i}{d}} \right) \sqrt{\frac{a b^2 - b^3 1 i}{4 d^2}} 2 i \\
&+ \operatorname{atan} \left(\frac{b^6 \sqrt{\frac{a b^2}{4 d^2} + \frac{b^3 1 i}{4 d^2}} \sqrt{a + b \tan(c + dx)} 32 i}{\frac{b^8 16 i}{d} + \frac{a^2 b^6 16 i}{d}} \right. \\
&\quad \left. - \frac{32 a b^5 \sqrt{\frac{a b^2}{4 d^2} + \frac{b^3 1 i}{4 d^2}} \sqrt{a + b \tan(c + dx)}}{\frac{b^8 16 i}{d} + \frac{a^2 b^6 16 i}{d}} \right) \sqrt{\frac{b^3 1 i + a b^2}{4 d^2}} 2 i
\end{aligned}$$

input `int(-(a + b*tan(c + d*x))^(1/2)*(a - b*tan(c + d*x)),x)`

output

$$\begin{aligned} & \operatorname{atan}\left(\frac{b^6((b^3 \cdot 1i)/(4d^2) + (a \cdot b^2)/(4d^2))^{1/2} \cdot (a + b \cdot \tan(c + dx))^{1/2} \cdot 32i}{(b^8 \cdot 16i)/d + (a^2 \cdot b^6 \cdot 16i)/d} - \frac{(32 \cdot a \cdot b^5 \cdot ((b^3 \cdot 1i)/(4d^2) + (a \cdot b^2)/(4d^2))^{1/2} \cdot (a + b \cdot \tan(c + dx))^{1/2})}{(b^8 \cdot 16i)/d + (a^2 \cdot b^6 \cdot 16i)/d}\right) \\ & - \operatorname{atan}\left(\frac{b^6 \cdot ((a \cdot b^2)/(4d^2) - (b^3 \cdot 1i)/(4d^2))^{1/2} \cdot (a + b \cdot \tan(c + dx))^{1/2} \cdot 32i}{(b^8 \cdot 16i)/d + (a^2 \cdot b^6 \cdot 16i)/d} + \frac{(32 \cdot a \cdot b^5 \cdot ((a \cdot b^2)/(4d^2) - (b^3 \cdot 1i)/(4d^2))^{1/2} \cdot (a + b \cdot \tan(c + dx))^{1/2})}{(b^8 \cdot 16i)/d + (a^2 \cdot b^6 \cdot 16i)/d}\right) \\ & + \frac{(a \cdot b^2 - b^3 \cdot 1i)/(4d^2)^{1/2} \cdot 2i + \operatorname{atanh}\left(\frac{d^3 \cdot ((16 \cdot (a^2 \cdot b^4 - a^4 \cdot b^2) \cdot (a + b \cdot \tan(c + dx))^{1/2})/d^2 + (16 \cdot a \cdot b^2 \cdot (a^2 \cdot b \cdot 1i + a^3) \cdot (a + b \cdot \tan(c + dx))^{1/2})/d^2 \cdot (-a^2 \cdot b \cdot 1i + a^3)/d^2)^{1/2}}{16 \cdot (a^3 \cdot b^5 + a^5 \cdot b^3)}\right) \cdot (-a^2 \cdot b \cdot 1i + a^3)/d^2)^{1/2}}{16 \cdot (a^3 \cdot b^5 + a^5 \cdot b^3)} \\ & + \frac{\operatorname{atanh}\left(\frac{d^3 \cdot ((a^2 \cdot b \cdot 1i - a^3)/d^2)^{1/2} \cdot ((16 \cdot (a^2 \cdot b^4 - a^4 \cdot b^2) \cdot (a + b \cdot \tan(c + dx))^{1/2})/d^2 - (16 \cdot a \cdot b^2 \cdot (a^2 \cdot b \cdot 1i - a^3) \cdot (a + b \cdot \tan(c + dx))^{1/2})/d^2)}{16 \cdot (a^3 \cdot b^5 + a^5 \cdot b^3)}\right) \cdot ((a^2 \cdot b \cdot 1i - a^3)/d^2)^{1/2}}{16 \cdot (a^3 \cdot b^5 + a^5 \cdot b^3)} + (2 \cdot b \cdot (a + b \cdot \tan(c + dx))^{1/2})/d \end{aligned}$$

3.343
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

3.343.1 Optimal result 3269
 3.343.2 Mathematica [A] (verified) 3270
 3.343.3 Rubi [A] (warning: unable to verify) 3270
 3.343.4 Maple [B] (verified) 3274
 3.343.5 Fracas [B] (verification not implemented) 3275
 3.343.6 Sympy [F] 3276
 3.343.7 Maxima [F] 3277
 3.343.8 Giac [F(-1)] 3277
 3.343.9 Mupad [B] (verification not implemented) 3277

3.343.1 Optimal result

Integrand size = 33, antiderivative size = 213

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = \frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ibd}} + \frac{(A+iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ibd}} - \frac{2(10aAb-8a^2B+15b^2B)\sqrt{a+b \tan(c+dx)}}{15b^3d} + \frac{2(5Ab-4aB)\tan(c+dx)\sqrt{a+b \tan(c+dx)}}{15b^2d} + \frac{2B \tan^2(c+dx)\sqrt{a+b \tan(c+dx)}}{5bd}$$

```
output (A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d/(a-I*b)^(1/2)+(A+I
*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d/(a+I*b)^(1/2)-2/15*(10
*A*a*b-8*B*a^2+15*B*b^2)*(a+b*tan(d*x+c))^(1/2)/b^3/d+2/15*(5*A*b-4*B*a)*(
a+b*tan(d*x+c))^(1/2)*tan(d*x+c)/b^2/d+2/5*B*(a+b*tan(d*x+c))^(1/2)*tan(d*
x+c)^2/b/d
```

3.343.2 Mathematica [A] (verified)

Time = 4.34 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.80

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

$$= \frac{15(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib}} + \frac{15(A+iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ib}} + \frac{2\sqrt{a+b \tan(c+dx)}(-10aAb+8a^2B-15b^2B+b(5Ab-4a^2))}{15d b^3}$$

input `Integrate[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `((15*(A - I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/Sqrt[a - I*b] + (15*(A + I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/Sqrt[a + I*b] + (2*Sqrt[a + b*Tan[c + d*x]]*(-10*a*A*b + 8*a^2*B - 15*b^2*B + b*(5*A*b - 4*a*B)*Tan[c + d*x] + 3*b^2*B*Tan[c + d*x]^2))/b^3)/(15*d)`

3.343.3 Rubi [A] (warning: unable to verify)

Time = 1.19 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.04, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 4090, 27, 3042, 4130, 27, 3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

↓ 3042

$$\int \frac{\tan(c+dx)^3(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

↓ 4090

$$2 \int \frac{\tan(c+dx)((5Ab-4aB) \tan^2(c+dx)+5bB \tan(c+dx)+4aB)}{2\sqrt{a+b \tan(c+dx)}} dx + \frac{2B \tan^2(c+dx)\sqrt{a+b \tan(c+dx)}}{5bd}$$

↓ 27

3.343. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\begin{aligned}
& \frac{2B \tan^2(c+dx) \sqrt{a+b \tan(c+dx)}}{5bd} - \frac{\int \frac{\tan(c+dx) \left(-((5Ab-4aB) \tan^2(c+dx) + 5bB \tan(c+dx) + 4aB) \right)}{\sqrt{a+b \tan(c+dx)}} dx}{5b} \\
& \quad \downarrow 3042 \\
& \frac{2B \tan^2(c+dx) \sqrt{a+b \tan(c+dx)}}{5bd} - \frac{\int \frac{\tan(c+dx) \left(-((5Ab-4aB) \tan(c+dx)^2 + 5bB \tan(c+dx) + 4aB) \right)}{\sqrt{a+b \tan(c+dx)}} dx}{5b} \\
& \quad \downarrow 4130 \\
& \frac{2B \tan^2(c+dx) \sqrt{a+b \tan(c+dx)}}{5bd} - \\
& \frac{2 \int \frac{15A \tan(c+dx)b^2 + (-8Ba^2 + 10Aba + 15b^2B) \tan^2(c+dx) + 2a(5Ab-4aB)}{2\sqrt{a+b \tan(c+dx)}} dx}{3b} - \frac{2(5Ab-4aB) \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} \\
& \quad \downarrow 5b \\
& \quad \downarrow 27 \\
& \frac{2B \tan^2(c+dx) \sqrt{a+b \tan(c+dx)}}{5bd} - \\
& \frac{\int \frac{15A \tan(c+dx)b^2 + (-8Ba^2 + 10Aba + 15b^2B) \tan^2(c+dx) + 2a(5Ab-4aB)}{\sqrt{a+b \tan(c+dx)}} dx}{3b} - \frac{2(5Ab-4aB) \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} \\
& \quad \downarrow 5b \\
& \quad \downarrow 3042 \\
& \frac{2B \tan^2(c+dx) \sqrt{a+b \tan(c+dx)}}{5bd} - \\
& \frac{\int \frac{15A \tan(c+dx)b^2 + (-8Ba^2 + 10Aba + 15b^2B) \tan(c+dx)^2 + 2a(5Ab-4aB)}{\sqrt{a+b \tan(c+dx)}} dx}{3b} - \frac{2(5Ab-4aB) \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} \\
& \quad \downarrow 5b \\
& \quad \downarrow 4113 \\
& \frac{2B \tan^2(c+dx) \sqrt{a+b \tan(c+dx)}}{5bd} - \\
& \frac{\int \frac{15Ab^2 \tan(c+dx) - 15b^2B}{\sqrt{a+b \tan(c+dx)}} dx + \frac{2(-8a^2B + 10aAb + 15b^2B)}{bd} \frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{3b} - \frac{2(5Ab-4aB) \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} \\
& \quad \downarrow 5b \\
& \quad \downarrow 3042 \\
& \frac{2B \tan^2(c+dx) \sqrt{a+b \tan(c+dx)}}{5bd} - \\
& \frac{\int \frac{15Ab^2 \tan(c+dx) - 15b^2B}{\sqrt{a+b \tan(c+dx)}} dx + \frac{2(-8a^2B + 10aAb + 15b^2B)}{bd} \frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{3b} - \frac{2(5Ab-4aB) \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} \\
& \quad \downarrow 5b \\
& \quad \downarrow 4022
\end{aligned}$$

3.343. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\begin{aligned}
 & \frac{2B \tan^2(c+dx) \sqrt{a+b \tan(c+dx)}}{5bd} - \\
 & \frac{-\frac{2(5Ab-4aB) \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} + \frac{\frac{15}{2} b^2 (-B+iA) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx - \frac{15}{2} b^2 (B+iA) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \frac{2(-8a^2 B+10aAb+15b^2 B)}{bd}}{5b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2B \tan^2(c+dx) \sqrt{a+b \tan(c+dx)}}{5bd} - \\
 & \frac{-\frac{2(5Ab-4aB) \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} + \frac{\frac{15}{2} b^2 (-B+iA) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx - \frac{15}{2} b^2 (B+iA) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \frac{2(-8a^2 B+10aAb+15b^2 B)}{bd}}{5b} \\
 & \quad \downarrow \text{4020} \\
 & \frac{2B \tan^2(c+dx) \sqrt{a+b \tan(c+dx)}}{5bd} - \\
 & \frac{-\frac{2(5Ab-4aB) \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} + \frac{15ib^2 (B+iA) \int -\frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx)) - \frac{15ib^2 (-B+iA) \int -\frac{1}{(i \tan(c+dx)+1) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{3b}}{5b} \\
 & \quad \downarrow \text{25} \\
 & \frac{2B \tan^2(c+dx) \sqrt{a+b \tan(c+dx)}}{5bd} - \\
 & \frac{-\frac{2(5Ab-4aB) \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} + \frac{15ib^2 (B+iA) \int \frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx)) + \frac{15ib^2 (-B+iA) \int \frac{1}{(i \tan(c+dx)+1) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{3b}}{5b} \\
 & \quad \downarrow \text{73} \\
 & \frac{2B \tan^2(c+dx) \sqrt{a+b \tan(c+dx)}}{5bd} - \\
 & \frac{-\frac{2(5Ab-4aB) \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} + \frac{15b(-B+iA) \int \frac{1}{-i \tan^2(c+dx) - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)} + \frac{15b(B+iA) \int \frac{1}{i \tan^2(c+dx) + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{3b}}{5b} \\
 & \quad \downarrow \text{221} \\
 & \frac{2B \tan^2(c+dx) \sqrt{a+b \tan(c+dx)}}{5bd} - \\
 & \frac{-\frac{2(5Ab-4aB) \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} + \frac{2(-8a^2 B+10aAb+15b^2 B) \sqrt{a+b \tan(c+dx)}}{bd} - \frac{15b^2 (B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{3b} + \frac{15b^2 (-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{3b}}{5b}
 \end{aligned}$$

input `Int[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

3.343. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

```
output (2*B*Tan[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]]/(5*b*d) - ((-2*(5*A*b - 4*a*
B)*Tan[c + d*x]*Sqrt[a + b*Tan[c + d*x]]/(3*b*d) + ((-15*b^2*(I*A + B)*Ar
cTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d) + (15*b^2*(I*A - B)*Ar
cTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d) + (2*(10*a*A*b - 8*a^2
*B + 15*b^2*B)*Sqrt[a + b*Tan[c + d*x]]/(b*d))/(3*b))/(5*b)
```

3.343.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```


rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.343.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2014 vs. $2(183) = 366$.

Time = 0.18 (sec) , antiderivative size = 2015, normalized size of antiderivative = 9.46

method	result	size
parts	Expression too large to display	2015
derivativedivides	Expression too large to display	4107
default	Expression too large to display	4107

3.343.
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

```
input int(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output 2*A/d/b^2*(1/3*(a+b*tan(d*x+c))^(3/2)-a*(a+b*tan(d*x+c))^(1/2)-b^2*(1/4/(a
^2+b^2)^(1/2)*(-1/2*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*ln(b*tan(d*x+c)+a+(a+b*t
an(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))+2*((a^2+b^
2)^(1/2)-a)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)
+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)))+1/4/(a^2+b
^2)^(1/2)*(1/2*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*ln(b*tan(d*x+c)+a-(a+b*tan(d*
x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))+2*((a^2+b^2)^(1
/2)-a)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)-(2*(
a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)))))+B*(2/5/d/b^3*
(a+b*tan(d*x+c))^(5/2)-4/3/d/b^3*a*(a+b*tan(d*x+c))^(3/2)+2/d/b^3*a^2*(a+b
*tan(d*x+c))^(1/2)-2/d/b*(a+b*tan(d*x+c))^(1/2)+1/4/d/b/(a^2+b^2)*ln(b*tan
(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(
1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2+1/4/d*b/(a^2+b^2)*ln(b*tan(d*x+c)+
a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2
*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/4/d/b/(a^2+b^2)^(3/2)*ln(b*tan(d*x+c)+a+(a+b
*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+
b^2)^(1/2)+2*a)^(1/2)*a^3-1/4/d*b/(a^2+b^2)^(3/2)*ln(b*tan(d*x+c)+a+(a+b*t
an(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^
2)^(1/2)+2*a)^(1/2)*a-1/d/b/(a^2+b^2)^(1/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*
arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2...
```

3.343.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1767 vs. 2(177) = 354.

Time = 0.30 (sec) , antiderivative size = 1767, normalized size of antiderivative = 8.30

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorith
m="fricas")
```

output

```
-1/30*(15*b^3*d*sqrt(((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*log((2*(A^3*B + A*B^3)*a - (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) + ((B*a^3 - A*a^2*b + B*a*b^2 - A*b^3)*d^3*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) - (2*A^2*B*a^2 - (A^3 - 3*A*B^2)*a*b - (A^2*B - B^3)*b^2)*d)*sqrt(((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))) - 15*b^3*d*sqrt(((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*log((2*(A^3*B + A*B^3)*a - (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) - ((B*a^3 - A*a^2*b + B*a*b^2 - A*b^3)*d^3*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) - (2*A^2*B*a^2 - (A^3 - 3*A*B^2)*a*b - (A^2*B - B^3)*b^2)*d)*sqrt(((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))) - 15*b^3*d*sqrt(-(a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) - 2*A*B*b - (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*log(...
```

3.343.6 Sympy [F]

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = \int \frac{(A+B \tan(c+dx)) \tan^3(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$$

input `integrate(tan(d*x+c)**3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**3/sqrt(a + b*tan(c + d*x)), x)`

3.343.7 Maxima [F]

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^3}{\sqrt{b\tan(dx+c)+a}} dx$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm m="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^3/sqrt(b*tan(d*x + c) + a), x)`

3.343.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm m="giac")`

output `Timed out`

3.343.9 Mupad [B] (verification not implemented)

Time = 16.40 (sec) , antiderivative size = 3054, normalized size of antiderivative = 14.34

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Too large to display}$$

input `int((tan(c + d*x)^3*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x)`

output

$$\begin{aligned} & \operatorname{atan}\left(\frac{B^2 b^2 \left(-16 B^4 b^2 d^4\right)^{1/2}}{16\left(a^2 d^4 + b^2 d^4\right)} - \frac{B^2 a d^2}{4\left(a^2 d^4 + b^2 d^4\right)}\right)^{1/2} (a + b \tan(c + dx))^{1/2} * 32i / \left(\frac{16 B^3 a b^3 d^3}{a^2 d^4 + b^2 d^4} - \frac{4 B b^3 d^2 \left(-16 B^4 b^2 d^4\right)^{1/2}}{a^2 d^5 + b^2 d^5}\right) + \\ & \frac{a b^2 \left(-16 B^4 b^2 d^4\right)^{1/2}}{16\left(a^2 d^4 + b^2 d^4\right)} - \frac{B^2 a d^2}{4\left(a^2 d^4 + b^2 d^4\right)}\right)^{1/2} (a + b \tan(c + dx))^{1/2} * (-16 B^4 b^2 d^4)^{1/2} * 8i / \left(\frac{16 B^3 a b^5 d^5}{a^2 d^4 + b^2 d^4} - \frac{4 B b^5 d^4 \left(-16 B^4 b^2 d^4\right)^{1/2}}{a^2 d^5 + b^2 d^5}\right) + \\ & \frac{16 B^3 a^3 b^3 d^5}{a^2 d^4 + b^2 d^4} - \frac{4 B a^2 b^3 d^4 \left(-16 B^4 b^2 d^4\right)^{1/2}}{a^2 d^5 + b^2 d^5}\right) - \frac{B^2 a^2 b^2 d^2 \left(-16 B^4 b^2 d^4\right)^{1/2}}{16\left(a^2 d^4 + b^2 d^4\right)} - \frac{B^2 a d^2}{4\left(a^2 d^4 + b^2 d^4\right)}\right)^{1/2} (a + b \tan(c + dx))^{1/2} * 32i / \left(\frac{16 B^3 a b^5 d^5}{a^2 d^4 + b^2 d^4} - \frac{4 B b^5 d^4 \left(-16 B^4 b^2 d^4\right)^{1/2}}{a^2 d^5 + b^2 d^5}\right) + \\ & \frac{16 B^3 a^3 b^3 d^5}{a^2 d^4 + b^2 d^4} - \frac{4 B a^2 b^3 d^4 \left(-16 B^4 b^2 d^4\right)^{1/2}}{a^2 d^5 + b^2 d^5}\right) * \left(\frac{-16 B^4 b^2 d^4}{16\left(a^2 d^4 + b^2 d^4\right)} - \frac{B^2 a d^2}{4\left(a^2 d^4 + b^2 d^4\right)}\right)^{1/2} * 2i - \operatorname{atan}\left(\frac{A^2 b^2 \left(A^2 a d^2\right)}{4\left(a^2 d^4 + b^2 d^4\right)} - \frac{-16 A^4 b^2 d^4}{16\left(a^2 d^4 + b^2 d^4\right)}\right)^{1/2} (a + b \tan(c + dx))^{1/2} * 32i / \left(\frac{16 A^3 b^2}{d} - \frac{16 A^3 a^2 b^2 d^3}{a^2 d^4 + b^2 d^4} + \frac{4 A a b^2 d^2 \left(-16 A^4 b^2 d^4\right)^{1/2}}{a^2 d^5 + b^2 d^5}\right) + \frac{a b^2 \left(A^2 a d^2\right)}{4\left(a^2 d^4 + b^2 d^4\right)} - \frac{-16 A^4 b^2 d^4}{16\left(a^2 d^4 + b^2 d^4\right)}\right)^{1/2} (a + b \tan(c + dx))^{1/2} * (-16 A^4 b^2 d^4)^{1/2} * \dots \end{aligned}$$

3.344
$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

3.344.1 Optimal result 3279
 3.344.2 Mathematica [A] (verified) 3280
 3.344.3 Rubi [A] (warning: unable to verify) 3280
 3.344.4 Maple [B] (verified) 3284
 3.344.5 Fracas [B] (verification not implemented) 3285
 3.344.6 Sympy [F] 3285
 3.344.7 Maxima [F] 3286
 3.344.8 Giac [F(-1)] 3286
 3.344.9 Mupad [B] (verification not implemented) 3286

3.344.1 Optimal result

Integrand size = 33, antiderivative size = 166

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = \frac{(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ibd}} - \frac{(iA-B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ibd}} + \frac{2(3Ab-2aB)\sqrt{a+b \tan(c+dx)}}{3b^2d} + \frac{2B \tan(c+dx)\sqrt{a+b \tan(c+dx)}}{3bd}$$

output

```
(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d/(a-I*b)^(1/2)-(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d/(a+I*b)^(1/2)+2/3*(3*A*b-2*B*a)*(a+b*tan(d*x+c))^(1/2)/b^2/d+2/3*B*(a+b*tan(d*x+c))^(1/2)*tan(d*x+c)/b/d
```

3.344.2 Mathematica [A] (verified)

Time = 1.69 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.84

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

$$= \frac{3(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib}} + \frac{3(-iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ib}} + \frac{2\sqrt{a+b\tan(c+dx)}(3Ab-2aB+bB\tan(c+dx))}{b^2}$$

$$\qquad\qquad\qquad 3d$$

input `Integrate[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `((3*(I*A + B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/Sqrt[a - I*b] + (3*((-I)*A + B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/Sqrt[a + I*b] + (2*Sqrt[a + b*Tan[c + d*x]]*(3*A*b - 2*a*B + b*B*Tan[c + d*x]))/b^2)/(3*d)`

3.344.3 Rubi [A] (warning: unable to verify)Time = 0.83 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.95, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 4090, 27, 3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

$$\downarrow 3042$$

$$\int \frac{\tan(c+dx)^2(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

$$\downarrow 4090$$

$$\frac{2 \int -\frac{((3Ab-2aB)\tan^2(c+dx)+3bB\tan(c+dx)+2aB)}{2\sqrt{a+b\tan(c+dx)}} dx}{3b} + \frac{2B\tan(c+dx)\sqrt{a+b\tan(c+dx)}}{3bd}$$

$$\downarrow 27$$

3.344. $\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

$$\begin{aligned}
 & \frac{2B \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} - \frac{\int \frac{-((3Ab-2aB) \tan^2(c+dx) + 3bB \tan(c+dx) + 2aB) dx}{\sqrt{a+b \tan(c+dx)}}}{3b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2B \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} - \frac{\int \frac{-((3Ab-2aB) \tan(c+dx)^2 + 3bB \tan(c+dx) + 2aB) dx}{\sqrt{a+b \tan(c+dx)}}}{3b} \\
 & \quad \downarrow \text{4113} \\
 & \frac{2B \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} - \frac{\int \frac{3Ab+3B \tan(c+dx)b}{\sqrt{a+b \tan(c+dx)}} dx - \frac{2(3Ab-2aB) \sqrt{a+b \tan(c+dx)}}{bd}}{3b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2B \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} - \frac{\int \frac{3Ab+3B \tan(c+dx)b}{\sqrt{a+b \tan(c+dx)}} dx - \frac{2(3Ab-2aB) \sqrt{a+b \tan(c+dx)}}{bd}}{3b} \\
 & \quad \downarrow \text{4022} \\
 & \frac{2B \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} - \frac{\frac{3}{2}b(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{3}{2}b(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx - \frac{2(3Ab-2aB) \sqrt{a+b \tan(c+dx)}}{bd}}{3b} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2B \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} - \frac{\frac{3}{2}b(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{3}{2}b(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx - \frac{2(3Ab-2aB) \sqrt{a+b \tan(c+dx)}}{bd}}{3b} \\
 & \quad \downarrow \text{4020} \\
 & \frac{2B \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} - \frac{\frac{3ib(A-iB) \int -\frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} - \frac{3ib(A+iB) \int -\frac{1}{(i \tan(c+dx)+1) \sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} - \frac{2(3Ab-2aB) \sqrt{a+b \tan(c+dx)}}{bd}}{3b} \\
 & \quad \downarrow \text{25} \\
 & \frac{2B \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} - \frac{\frac{3ib(A-iB) \int \frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \frac{3ib(A+iB) \int \frac{1}{(i \tan(c+dx)+1) \sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} - \frac{2(3Ab-2aB) \sqrt{a+b \tan(c+dx)}}{bd}}{3b} \\
 & \quad \downarrow \text{73}
 \end{aligned}$$

3.344. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\frac{3(A+iB) \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)} + \frac{3bd}{3b} \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)} - \frac{2(3Ab-2aB)\sqrt{a+b \tan(c+dx)}}{bd}}{3b}$$

↓ 221

$$\frac{2B \tan(c+dx)\sqrt{a+b \tan(c+dx)}}{3b} - \frac{3b(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} + \frac{3b(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} - \frac{2(3Ab-2aB)\sqrt{a+b \tan(c+dx)}}{bd}$$

input `Int[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `(2*B*Tan[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/(3*b*d) - ((3*b*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d) + (3*b*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d) - (2*(3*A*b - 2*a*B)*Sqrt[a + b*Tan[c + d*x]]/(b*d))/(3*b)`

3.344.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

3.344.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1948 vs. $2(140) = 280$.

Time = 0.12 (sec) , antiderivative size = 1949, normalized size of antiderivative = 11.74

method	result	size
parts	Expression too large to display	1949
derivativedivides	Expression too large to display	4040
default	Expression too large to display	4040

```
input int(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2), x, method=_RETURNV
ERBOSE)
```

```
output A*(2/d/b*(a+b*tan(d*x+c))^(1/2)-1/4/d/b/(a^2+b^2)*ln(b*tan(d*x+c)+a+(a+b*t
an(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^
2)^(1/2)+2*a)^(1/2)*a^2-1/4/d*b/(a^2+b^2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c
))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2
)+2*a)^(1/2)+1/4/d/b/(a^2+b^2)^(3/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/
2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(
1/2)*a^3+1/4/d*b/(a^2+b^2)^(3/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)
*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1
/2)*a+1/d/b/(a^2+b^2)^(1/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*t
an(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1
/2))*a^2+1/d*b/(a^2+b^2)^(1/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+
b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)
^(1/2))-1/d/b/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b
*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(
1/2))*a^4-3/d*b/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(
a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*
a)^(1/2))*a^2-2/d*b^3/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan
((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/
2)-2*a)^(1/2))+1/4/d/b/(a^2+b^2)*ln(b*tan(d*x+c)+a-(a+b*tan(d*x+c))^(1/2)*
(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(...
```

3.344.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1725 vs. $2(134) = 268$.

Time = 0.31 (sec) , antiderivative size = 1725, normalized size of antiderivative = 10.39

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm
m="fricas")
```

```
output 1/6*(3*b^2*d*sqrt(-(a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^
3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A
*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*log((2*(A^3*B + A*B^3)*a - (A^4 -
B^4)*b)*sqrt(b*tan(d*x + c) + a) + ((A*a^3 + B*a^2*b + A*a*b^2 + B*b^3)*d
^3*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*
b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + (2*A*B^2*a^2 - (3*A^2*B - B^3)*a*b +
(A^3 - A*B^2)*b^2)*d)*sqrt(-(a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^
3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d
^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))) - 3*b^2*d*sqrt(-(a^2
+ b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2
+ B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a
^2 + b^2)*d^2))*log((2*(A^3*B + A*B^3)*a - (A^4 - B^4)*b)*sqrt(b*tan(d*x +
c) + a) - ((A*a^3 + B*a^2*b + A*a*b^2 + B*b^3)*d^3*sqrt(-(4*A^2*B^2*a^2 -
4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 +
b^4)*d^4)) + (2*A*B^2*a^2 - (3*A^2*B - B^3)*a*b + (A^3 - A*B^2)*b^2)*d)*sq
rt(-(a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 -
2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B
^2)*a)/((a^2 + b^2)*d^2))) - 3*b^2*d*sqrt(((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^
2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2
*b^2 + b^4)*d^4)) - 2*A*B*b - (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*log((2*...
```

3.344.6 Sympy [F]

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(A+B\tan(c+dx))\tan^2(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx$$

```
input integrate(tan(d*x+c)**2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)
```

3.344. $\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**2/sqrt(a + b*tan(c + d*x)), x)`

3.344.7 Maxima [F]

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)^2}{\sqrt{b \tan(dx + c) + a}} dx$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^2/sqrt(b*tan(d*x + c) + a), x)`

3.344.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.344.9 Mupad [B] (verification not implemented)

Time = 11.91 (sec) , antiderivative size = 2981, normalized size of antiderivative = 17.96

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `int((tan(c + d*x)^2*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x)`

output

$$\begin{aligned}
& 2*\operatorname{atanh}\left(\frac{32*A^2*b^2*(-16*A^4*b^2*d^4)^{(1/2)} / (16*(a^2*d^4 + b^2*d^4)) - (A^2*a*d^2) / (4*(a^2*d^4 + b^2*d^4))}{(4*(a^2*d^4 + b^2*d^4))^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)}}\right) / \left(\frac{16*A^3*a*b^3*d^3}{(a^2*d^4 + b^2*d^4)} - \frac{4*A*b^3*d^2*(-16*A^4*b^2*d^4)^{(1/2)}}{(a^2*d^5 + b^2*d^5)} + \frac{8*a*b^2*(-16*A^4*b^2*d^4)^{(1/2)} / (16*(a^2*d^4 + b^2*d^4)) - (A^2*a*d^2) / (4*(a^2*d^4 + b^2*d^4))}{(4*(a^2*d^4 + b^2*d^4))^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)}}\right) * (-16*A^4*b^2*d^4)^{(1/2)} / \left(\frac{16*A^3*a*b^5*d^5}{(a^2*d^4 + b^2*d^4)} - \frac{4*A*b^5*d^4*(-16*A^4*b^2*d^4)^{(1/2)}}{(a^2*d^5 + b^2*d^5)} + \frac{16*A^3*a^3*b^3*d^5}{(a^2*d^4 + b^2*d^4)} - \frac{4*A*a^2*b^3*d^4*(-16*A^4*b^2*d^4)^{(1/2)}}{(a^2*d^5 + b^2*d^5)} - \frac{32*A^2*a^2*b^2*d^2*(-16*A^4*b^2*d^4)^{(1/2)} / (16*(a^2*d^4 + b^2*d^4)) - (A^2*a*d^2) / (4*(a^2*d^4 + b^2*d^4))}{(4*(a^2*d^4 + b^2*d^4))^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)}}\right) / \left(\frac{16*A^3*a*b^5*d^5}{(a^2*d^4 + b^2*d^4)} - \frac{4*A*b^5*d^4*(-16*A^4*b^2*d^4)^{(1/2)}}{(a^2*d^5 + b^2*d^5)} + \frac{16*A^3*a^3*b^3*d^5}{(a^2*d^4 + b^2*d^4)} - \frac{4*A*a^2*b^3*d^4*(-16*A^4*b^2*d^4)^{(1/2)}}{(a^2*d^5 + b^2*d^5)}\right) * \left(\frac{-16*A^4*b^2*d^4)^{(1/2)} / (16*(a^2*d^4 + b^2*d^4)) - (A^2*a*d^2) / (4*(a^2*d^4 + b^2*d^4))}{(4*(a^2*d^4 + b^2*d^4))^{(1/2)}} - \operatorname{atan}\left(\frac{a*b^2*(-16*B^4*b^2*d^4)^{(1/2)} / (16*(a^2*d^4 + b^2*d^4)) + (B^2*a*d^2) / (4*(a^2*d^4 + b^2*d^4))}{(4*(a^2*d^4 + b^2*d^4))^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)}}\right) * (-16*B^4*b^2*d^4)^{(1/2)} * i\right) / \left(\frac{16*B^3*a^2*b^4*d^5}{(a^2*d^4 + b^2*d^4)} - 16*B^3*a^2*b^2*d - 16*B^3*b^4*d + \frac{16*B^3*a^4*b^2*d^5}{(a^2*d^4 + b^2*d^4)} + \frac{4*B*a^3*b^2*d^4*(-16*B^4*b^2*d^4)^{(1/2)}}{(a^2*d^5 + b^2*d^5)} + \frac{4*B*a*b^4*d^4*(-16*B^4*b^2*d^4)^{(1/2)}}{(a^2*d^5 + b^2*d^5)} - (B^2*b\dots
\end{aligned}$$

$$3.345 \quad \int \frac{\tan(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

3.345.1 Optimal result	3288
3.345.2 Mathematica [A] (verified)	3288
3.345.3 Rubi [A] (warning: unable to verify)	3289
3.345.4 Maple [B] (verified)	3291
3.345.5 Fricas [B] (verification not implemented)	3292
3.345.6 Sympy [F]	3293
3.345.7 Maxima [F]	3294
3.345.8 Giac [F(-1)]	3294
3.345.9 Mupad [B] (verification not implemented)	3294

3.345.1 Optimal result

Integrand size = 31, antiderivative size = 124

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = -\frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ibd}} - \frac{(A+iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ibd}} + \frac{2B\sqrt{a+b \tan(c+dx)}}{bd}$$

output `-(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d/(a-I*b)^(1/2)-(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d/(a+I*b)^(1/2)+2*B*(a+b*tan(d*x+c))^(1/2)/b/d`

3.345.2 Mathematica [A] (verified)

Time = 0.56 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.95

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = -\frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib}} + \frac{(A+iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ib}} - \frac{2B\sqrt{a+b \tan(c+dx)}}{b}$$

3.345. $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

input `Integrate[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `-((((A - I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/Sqrt[a - I*b] + ((A + I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/Sqrt[a + I*b] - (2*B*Sqrt[a + b*Tan[c + d*x]])/b)/d)`

3.345.3 Rubi [A] (warning: unable to verify)

Time = 0.57 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.88, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 4075, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx \\
 & \quad \downarrow \text{4075} \\
 & \int \frac{A\tan(c+dx)-B}{\sqrt{a+b\tan(c+dx)}} dx + \frac{2B\sqrt{a+b\tan(c+dx)}}{bd} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A\tan(c+dx)-B}{\sqrt{a+b\tan(c+dx)}} dx + \frac{2B\sqrt{a+b\tan(c+dx)}}{bd} \\
 & \quad \downarrow \text{4022} \\
 & \frac{1}{2}(-B+iA) \int \frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx - \frac{1}{2}(B+iA) \int \frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}} dx + \\
 & \quad \frac{2B\sqrt{a+b\tan(c+dx)}}{bd} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2}(-B+iA) \int \frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx - \frac{1}{2}(B+iA) \int \frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}} dx + \\
 & \quad \frac{2B\sqrt{a+b\tan(c+dx)}}{bd}
 \end{aligned}$$

3.345. $\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

$$\begin{aligned}
& \downarrow 4020 \\
& \frac{i(B+iA) \int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} - \\
& \frac{i(-B+iA) \int -\frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \frac{2B\sqrt{a+b \tan(c+dx)}}{bd} \\
& \downarrow 25 \\
& \frac{i(B+iA) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \\
& \frac{i(-B+iA) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} + \frac{2B\sqrt{a+b \tan(c+dx)}}{bd} \\
& \downarrow 73 \\
& \frac{(-B+iA) \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} - \\
& \frac{(B+iA) \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \frac{2B\sqrt{a+b \tan(c+dx)}}{bd} \\
& \downarrow 221 \\
& -\frac{(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} + \frac{(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} + \frac{2B\sqrt{a+b \tan(c+dx)}}{bd}
\end{aligned}$$

input `Int[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `-(((I*A + B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d)) + ((I*A - B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d) + (2*B*Sqrt[a + b*Tan[c + d*x]])/(b*d)`

3.345.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

3.345. $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.345.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1909 vs. $2(104) = 208$.

Time = 0.12 (sec) , antiderivative size = 1910, normalized size of antiderivative = 15.40

method	result	size
parts	Expression too large to display	1910
derivativedivides	Expression too large to display	3997
default	Expression too large to display	3997

input `int(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

$$3.345. \quad \int \frac{\tan(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

output

```

A/d*(1/2/(a^2+b^2)^(1/2)*(-1/2*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))
+2*((a^2+b^2)^(1/2)-a)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)))
+1/2/(a^2+b^2)^(1/2)*(1/2*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*ln(b*tan(d*x+c)+a-(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))+2*((a^2+b^2)^(1/2)-a)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)-(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))))+B*(
2/d/b*(a+b*tan(d*x+c))^(1/2)-1/4/d/b/(a^2+b^2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2-1/4/d*b/(a^2+b^2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/4/d/b/(a^2+b^2)^(3/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3+1/4/d*b/(a^2+b^2)^(3/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+1/d/b/(a^2+b^2)^(1/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)))*a^2+1/d*b/(a^2+b^2)^(1/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)...

```

3.345.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1706 vs. $2(98) = 196$.

Time = 0.30 (sec) , antiderivative size = 1706, normalized size of antiderivative = 13.76

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Too large to display}$$

input

```

integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm=
"fricas")

```

output $1/2*(b*d*\sqrt{((a^2 + b^2)*d^2*\sqrt{-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2})/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*\log((2*(A^3*B + A*B^3)*a - (A^4 - B^4)*b)*\sqrt{b*\tan(d*x + c) + a} + ((B*a^3 - A*a^2*b + B*a*b^2 - A*b^3)*d^3*\sqrt{-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2})/((a^4 + 2*a^2*b^2 + b^4)*d^4)) - (2*A^2*B*a^2 - (A^3 - 3*A*B^2)*a*b - (A^2*B - B^3)*b^2)*d)*\sqrt{((a^2 + b^2)*d^2*\sqrt{-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2})/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))} - b*d*\sqrt{((a^2 + b^2)*d^2*\sqrt{-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2})/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*\log((2*(A^3*B + A*B^3)*a - (A^4 - B^4)*b)*\sqrt{b*\tan(d*x + c) + a} - ((B*a^3 - A*a^2*b + B*a*b^2 - A*b^3)*d^3*\sqrt{-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2})/((a^4 + 2*a^2*b^2 + b^4)*d^4)) - (2*A^2*B*a^2 - (A^3 - 3*A*B^2)*a*b - (A^2*B - B^3)*b^2)*d)*\sqrt{((a^2 + b^2)*d^2*\sqrt{-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2})/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))} - b*d*\sqrt{-(a^2 + b^2)*d^2*\sqrt{-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2})/((a^4 + 2*a^2*b^2 + b^4)*d^4))} - 2*A*B*b - (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*\log((2*(A^3*B + A*B^3)...$

3.345.6 Sympy [F]

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(A+B\tan(c+dx))\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2), x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)/sqrt(a + b*tan(c + d*x)), x)`

3.345.7 Maxima [F]

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)}{\sqrt{b\tan(dx+c)+a}} dx$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)/sqrt(b*tan(d*x + c) + a), x)`

3.345.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.345.9 Mupad [B] (verification not implemented)

Time = 11.28 (sec) , antiderivative size = 2930, normalized size of antiderivative = 23.63

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Too large to display}$$

input `int((tan(c + d*x)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x)`

output

$$\begin{aligned}
& 2*\operatorname{atanh}((32*B^2*b^2*((-16*B^4*b^2*d^4)^{(1/2)}/(16*(a^2*d^4 + b^2*d^4)) - (B^2*a*d^2)/(4*(a^2*d^4 + b^2*d^4)))^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)})/((16*B^3*a*b^3*d^3)/(a^2*d^4 + b^2*d^4) - (4*B*b^3*d^2*(-16*B^4*b^2*d^4)^{(1/2)})/(a^2*d^5 + b^2*d^5)) + (8*a*b^2*((-16*B^4*b^2*d^4)^{(1/2)}/(16*(a^2*d^4 + b^2*d^4)) - (B^2*a*d^2)/(4*(a^2*d^4 + b^2*d^4)))^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)}*(-16*B^4*b^2*d^4)^{(1/2)})/((16*B^3*a*b^5*d^5)/(a^2*d^4 + b^2*d^4) - (4*B*b^5*d^4*(-16*B^4*b^2*d^4)^{(1/2)})/(a^2*d^5 + b^2*d^5)) + (16*B^3*a^3*b^3*d^5)/(a^2*d^4 + b^2*d^4) - (4*B*a^2*b^3*d^4*(-16*B^4*b^2*d^4)^{(1/2)})/(a^2*d^5 + b^2*d^5) - (32*B^2*a^2*b^2*d^2*((-16*B^4*b^2*d^4)^{(1/2)}/(16*(a^2*d^4 + b^2*d^4)) - (B^2*a*d^2)/(4*(a^2*d^4 + b^2*d^4)))^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)})/((16*B^3*a*b^5*d^5)/(a^2*d^4 + b^2*d^4) - (4*B*b^5*d^4*(-16*B^4*b^2*d^4)^{(1/2)})/(a^2*d^5 + b^2*d^5)) + (16*B^3*a^3*b^3*d^5)/(a^2*d^4 + b^2*d^4) - (4*B*a^2*b^3*d^4*(-16*B^4*b^2*d^4)^{(1/2)})/(a^2*d^5 + b^2*d^5)))*((-16*B^4*b^2*d^4)^{(1/2)}/(16*(a^2*d^4 + b^2*d^4)) - (B^2*a*d^2)/(4*(a^2*d^4 + b^2*d^4)))^{(1/2)} - 2*\operatorname{atanh}((8*a*b^2*((-16*A^4*b^2*d^4)^{(1/2)}/(16*(a^2*d^4 + b^2*d^4)) + (A^2*a*d^2)/(4*(a^2*d^4 + b^2*d^4)))^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)}*(-16*A^4*b^2*d^4)^{(1/2)})/((16*A^3*a^2*b^4*d^5)/(a^2*d^4 + b^2*d^4) - 16*A^3*a^2*b^2*d - 16*A^3*b^4*d + (16*A^3*a^4*b^2*d^5)/(a^2*d^4 + b^2*d^4) + (4*A*a^3*b^2*d^4*(-16*A^4*b^2*d^4)^{(1/2)})/(a^2*d^5 + b^2*d^5)) + (4*A*a*b^4*d^4*(-16*A^4*b^2*d^4)^{(1/2)})/(a^2*d^5 + b^2*d^5)) - (32*...
\end{aligned}$$

3.346 $\int \frac{A+B \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$

3.346.1 Optimal result	3296
3.346.2 Mathematica [A] (verified)	3296
3.346.3 Rubi [A] (warning: unable to verify)	3297
3.346.4 Maple [B] (verified)	3299
3.346.5 Fricas [B] (verification not implemented)	3300
3.346.6 Sympy [F]	3300
3.346.7 Maxima [F(-2)]	3301
3.346.8 Giac [F(-1)]	3301
3.346.9 Mupad [B] (verification not implemented)	3301

3.346.1 Optimal result

Integrand size = 25, antiderivative size = 102

$$\int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = -\frac{(iA + B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a - ibd}} + \frac{(iA - B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a + ibd}}$$

output `-(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d/(a-I*b)^(1/2)+(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d/(a+I*b)^(1/2)`

3.346.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.99

$$\int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \frac{i \left(-\frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib}} + \frac{(A+iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ib}} \right)}{d}$$

input `Integrate[(A + B*Tan[c + d*x])/Sqrt[a + b*Tan[c + d*x]],x]`

output $(I*(-((A - I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/Sqrt[a - I*b]) + ((A + I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/Sqrt[a + I*b])/d$

3.346.3 Rubi [A] (warning: unable to verify)

Time = 0.44 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.81, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$, Rules used = {3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx \\ & \quad \downarrow \text{4022} \\ & \frac{1}{2}(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \frac{1}{2}(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx \\ & \quad \downarrow \text{4020} \\ & \frac{i(A - iB) \int -\frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} - \\ & \frac{i(A + iB) \int -\frac{1}{(i \tan(c + dx) + 1)\sqrt{a + b \tan(c + dx)}} d(-i \tan(c + dx))}{2d} \\ & \quad \downarrow \text{25} \\ & \frac{i(A + iB) \int \frac{1}{(i \tan(c + dx) + 1)\sqrt{a + b \tan(c + dx)}} d(-i \tan(c + dx))}{2d} - \\ & \frac{i(A - iB) \int \frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} \\ & \quad \downarrow \text{73} \end{aligned}$$

3.346. $\int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx$

$$\frac{(A + iB) \int \frac{1}{-i \frac{\tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)}}{bd} + \frac{(A - iB) \int \frac{1}{i \frac{\tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)}}{bd}$$

↓ 221

$$\frac{(A - iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} + \frac{(A + iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}}$$

input `Int[(A + B*Tan[c + d*x])/Sqrt[a + b*Tan[c + d*x]],x]`

output `((A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d) + ((A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d)`

3.346.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n_), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

3.346.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1891 vs. $2(84) = 168$.

Time = 0.50 (sec) , antiderivative size = 1892, normalized size of antiderivative = 18.55

method	result	size
parts	Expression too large to display	1892
derivativedivides	Expression too large to display	3976
default	Expression too large to display	3976

```
input int((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x,method=_RETURNVERBOSE)
```

```
output A*(1/4/d/b/(a^2+b^2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)
^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2+1/4/d
*b/(a^2+b^2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2
*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/4/d/b/(a^2+b^2)
^(3/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1
/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3-1/4/d*b/(a^2+b^2)^(
3/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2
)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-1/d/b/(a^2+b^2)^(1/2)/(
2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b
^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^2-1/d*b/(a^2+b^2)^(1/2
)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b
^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))+1/d/b/(a^2+b^2)^(3/2)
/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b
^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^4+3/d*b/(a^2+b^2)^(3
/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2
+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^2+2/d*b^3/(a^2+b
^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2
*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))-1/4/d/b/(a^2+b
^2)*ln(b*tan(d*x+c)+a-(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)
+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2-1/4/d*b/(a^2+b^2)*1...
```

3.346.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1669 vs. $2(79) = 158$.

Time = 0.28 (sec) , antiderivative size = 1669, normalized size of antiderivative = 16.36

$$\int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

```
input integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="fracas")
```

```
output -1/2*sqrt(-((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b +
(A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (
A^2 - B^2)*a)/((a^2 + b^2)*d^2))*log((2*(A^3*B + A*B^3)*a - (A^4 - B^4)*b)
*sqrt(b*tan(d*x + c) + a) + ((A*a^3 + B*a^2*b + A*a*b^2 + B*b^3)*d^3*sqrt(
-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a
^4 + 2*a^2*b^2 + b^4)*d^4)) + (2*A*B^2*a^2 - (3*A^2*B - B^3)*a*b + (A^3 -
A*B^2)*b^2)*d)*sqrt(-((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*
B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2
*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))) + 1/2*sqrt(-((a^2 + b^2)*d^2*s
qrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)
/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^
2))*log((2*(A^3*B + A*B^3)*a - (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) - (
(A*a^3 + B*a^2*b + A*a*b^2 + B*b^3)*d^3*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B -
A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) +
(2*A*B^2*a^2 - (3*A^2*B - B^3)*a*b + (A^3 - A*B^2)*b^2)*d)*sqrt(-((a^2 +
b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 +
B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2
+ b^2)*d^2))) + 1/2*sqrt(((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B
- A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)
) - 2*A*B*b - (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*log((2*(A^3*B + A*B^3)*...
```

3.346.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx$$

```
input integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)
```

```
output Integral((A + B*tan(c + d*x))/sqrt(a + b*tan(c + d*x)), x)
```

3.346. $\int \frac{A+B \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$

3.346.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is`

3.346.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.346.9 Mupad [B] (verification not implemented)

Time = 10.17 (sec) , antiderivative size = 2909, normalized size of antiderivative = 28.52

$$\int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `int((A + B*tan(c + d*x))/(a + b*tan(c + d*x))^(1/2),x)`

output

$$\begin{aligned}
& 2*\operatorname{atanh}((8*a*b^2*(-(-16*A^4*b^2*d^4)^{(1/2)}/(16*(a^2*d^4 + b^2*d^4)) - (A^2*a*d^2)/(4*(a^2*d^4 + b^2*d^4))))^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)}*(-16*A^4*b^2*d^4)^{(1/2)})/((16*A^3*a*b^5*d^5)/(a^2*d^4 + b^2*d^4) + (4*A*b^5*d^4*(-16*A^4*b^2*d^4)^{(1/2)})/(a^2*d^5 + b^2*d^5) + (16*A^3*a^3*b^3*d^5)/(a^2*d^4 + b^2*d^4) + (4*A*a^2*b^3*d^4*(-16*A^4*b^2*d^4)^{(1/2)})/(a^2*d^5 + b^2*d^5)) - (32*A^2*b^2*(-(-16*A^4*b^2*d^4)^{(1/2)}/(16*(a^2*d^4 + b^2*d^4)) - (A^2*a*d^2)/(4*(a^2*d^4 + b^2*d^4))))^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)})/((16*A^3*a*b^3*d^3)/(a^2*d^4 + b^2*d^4) + (4*A*b^3*d^2*(-16*A^4*b^2*d^4)^{(1/2)})/(a^2*d^5 + b^2*d^5)) + (32*A^2*a^2*b^2*d^2*(-(-16*A^4*b^2*d^4)^{(1/2)}/(16*(a^2*d^4 + b^2*d^4)) - (A^2*a*d^2)/(4*(a^2*d^4 + b^2*d^4))))^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)})/((16*A^3*a*b^5*d^5)/(a^2*d^4 + b^2*d^4) + (4*A*b^5*d^4*(-16*A^4*b^2*d^4)^{(1/2)})/(a^2*d^5 + b^2*d^5) + (16*A^3*a^3*b^3*d^5)/(a^2*d^4 + b^2*d^4) + (4*A*a^2*b^3*d^4*(-16*A^4*b^2*d^4)^{(1/2)})/(a^2*d^5 + b^2*d^5))) * (-(-16*A^4*b^2*d^4)^{(1/2)}/(16*(a^2*d^4 + b^2*d^4)) - (A^2*a*d^2)/(4*(a^2*d^4 + b^2*d^4)))^{(1/2)} - 2*\operatorname{atanh}((8*a*b^2*(-(-16*B^4*b^2*d^4)^{(1/2)}/(16*(a^2*d^4 + b^2*d^4)) + (B^2*a*d^2)/(4*(a^2*d^4 + b^2*d^4))))^{(1/2)}*(a + b*\tan(c + d*x))^{(1/2)}*(-16*B^4*b^2*d^4)^{(1/2)})/((16*B^3*a^2*b^4*d^5)/(a^2*d^4 + b^2*d^4) - 16*B^3*a^2*b^2*d - 16*B^3*b^4*d + (16*B^3*a^4*b^2*d^5)/(a^2*d^4 + b^2*d^4) + (4*B*a^3*b^2*d^4*(-16*B^4*b^2*d^4)^{(1/2)})/(a^2*d^5 + b^2*d^5) + (4*B*a*b^4*d^4*(-16*B^4*b^2*d^4)^{(1/2)})/(a^2*d^5 + b^2*d^5))...
\end{aligned}$$

$$3.347 \quad \int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

3.347.1 Optimal result	3303
3.347.2 Mathematica [A] (verified)	3303
3.347.3 Rubi [A] (warning: unable to verify)	3304
3.347.4 Maple [B] (verified)	3307
3.347.5 Fricas [B] (verification not implemented)	3308
3.347.6 Sympy [F]	3309
3.347.7 Maxima [F]	3310
3.347.8 Giac [F(-1)]	3310
3.347.9 Mupad [B] (verification not implemented)	3310

3.347.1 Optimal result

Integrand size = 31, antiderivative size = 131

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = -\frac{2A \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ad}} + \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib}d} + \frac{(A+iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ib}d}$$

```
output -2*A*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/d/a^(1/2)+(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d/(a-I*b)^(1/2)+(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d/(a+I*b)^(1/2)
```

3.347.2 Mathematica [A] (verified)

Time = 0.81 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.30

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = \frac{2A \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{a}} + \frac{(-Ab+\sqrt{-b^2}B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-\sqrt{-b^2}}}\right)}{\sqrt{a-\sqrt{-b^2}}d} - \frac{(Ab+\sqrt{-b^2}B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+\sqrt{-b^2}}}\right)}{\sqrt{a+\sqrt{-b^2}}d}$$

3.347. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

input `Integrate[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `-(((2*A*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/Sqrt[a] + (((-(A*b) + Sqrt[-b^2]*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - Sqrt[-b^2]]])/Sqrt[a - Sqrt[-b^2]] - ((A*b + Sqrt[-b^2]*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + Sqrt[-b^2]]])/Sqrt[a + Sqrt[-b^2]]))/b)/d)`

3.347.3 Rubi [A] (warning: unable to verify)

Time = 0.94 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.90, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 4096, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A+B\tan(c+dx)}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx \\
 & \quad \downarrow \text{4096} \\
 & \int \frac{B-A\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + A \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b\tan(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B-A\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + A \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx \\
 & \quad \downarrow \text{4022} \\
 & -\frac{1}{2}(-B+iA) \int \frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}(B+iA) \int \frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}} dx + \\
 & \quad A \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{2}(-B+iA) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}(B+iA) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \\
& \quad A \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx \\
& \quad \downarrow 4020 \\
& \quad \frac{i(B+iA) \int -\frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{+} \\
& \quad \frac{i(-B+iA) \int -\frac{2d}{(i \tan(c+dx)+1) \sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{+} \\
& \quad A \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx \\
& \quad \downarrow 25 \\
& \quad \frac{i(B+iA) \int \frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{-} \\
& \quad \frac{i(-B+iA) \int \frac{2d}{(i \tan(c+dx)+1) \sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{+} \\
& \quad A \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx \\
& \quad \downarrow 73 \\
& \quad \frac{(-B+iA) \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{+} \\
& \quad \frac{(B+iA) \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + A \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx \\
& \quad \downarrow 221 \\
& \quad A \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + \frac{(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \\
& \quad \frac{(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} \\
& \quad \downarrow 4117 \\
& \quad \frac{A \int \frac{\cot(c+dx)}{\sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{d} + \frac{(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} \\
& \quad \downarrow 73
\end{aligned}$$

$$\begin{aligned}
& \frac{2A \int \frac{1}{\frac{a+b \tan(c+dx)}{b} - \frac{a}{b}} d\sqrt{a+b \tan(c+dx)}}{bd} + \frac{(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \\
& \frac{(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} \\
& \quad \downarrow \text{221} \\
& \frac{(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} - \frac{2A \operatorname{Arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ad}}
\end{aligned}$$

input `Int[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `((I*A + B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]]/(Sqrt[a - I*b]*d) - ((I*A - B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]]/(Sqrt[a + I*b]*d) - (2*A*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/(Sqrt[a]*d)`

3.347.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

3.347. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

```
rule 4096 Int((((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_
.)*(x_)])^(n_))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[
1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[a*A + b*B - (A*b - a*B)*Tan
[e + f*x], x], x] + Simp[b*(A*b - a*B)/(a^2 + b^2) Int[(c + d*Tan[e
+ f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

3.347.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4001 vs. $2(107) = 214$.

Time = 0.28 (sec) , antiderivative size = 4002, normalized size of antiderivative = 30.55

method	result	size
derivativedivides	Expression too large to display	4002
default	Expression too large to display	4002

```
input int(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x,method=_RETURNVER
BOSE)
```

output

```

1/d/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a^3+1/4/d/(a^2+b^2)^(3/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2-1/4/d/(a^2+b^2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+2/d/(a^2+b^2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a^2-1/d/(a^2+b^2)^(1/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a-1/d/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a^3-1/4/d/(a^2+b^2)^(3/2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2+1/d/(a^2+b^2)^(1/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a+1/4/d/(a^2+b^2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-2/d/(a^2+b^2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a^2+2/d*b^3/(a^2+b^2)^(...

```

3.347.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1721 vs. $2(101) = 202$.

Time = 0.82 (sec) , antiderivative size = 3458, normalized size of antiderivative = 26.40

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = \text{Too large to display}$$

input

```

integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="fricas")

```

output

```

[-1/2*(a*d*sqrt(((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*
a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*
b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*log((2*(A^3*B + A*B^3)*a - (A^4 - B^
4)*b)*sqrt(b*tan(d*x + c) + a) + ((B*a^3 - A*a^2*b + B*a*b^2 - A*b^3)*d^3*
sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2
)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) - (2*A^2*B*a^2 - (A^3 - 3*A*B^2)*a*b - (A
^2*B - B^3)*b^2)*d)*sqrt(((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B
- A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4))
+ 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))) - a*d*sqrt(((a^2 + b^2)*d^
2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b
^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)
*d^2))*log((2*(A^3*B + A*B^3)*a - (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a)
- ((B*a^3 - A*a^2*b + B*a*b^2 - A*b^3)*d^3*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B
- A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)
) - (2*A^2*B*a^2 - (A^3 - 3*A*B^2)*a*b - (A^2*B - B^3)*b^2)*d)*sqrt(((a^2
+ b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2
+ B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a
^2 + b^2)*d^2))) - a*d*sqrt(-((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^
3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d
^4)) - 2*A*B*b - (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*log((2*(A^3*B + A*B^...

```

3.347.6 Sympy [F]

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = \int \frac{(A+B \tan(c+dx)) \cot(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2), x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)/sqrt(a + b*tan(c + d*x)), x)`

3.347.7 Maxima [F]

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx = \int \frac{(B \tan(dx + c) + A) \cot(dx + c)}{\sqrt{b \tan(dx + c) + a}} dx$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)/sqrt(b*tan(d*x + c) + a), x)`

3.347.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.347.9 Mupad [B] (verification not implemented)

Time = 12.12 (sec) , antiderivative size = 7099, normalized size of antiderivative = 54.19

$$\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `int((cot(c + d*x)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x)`

output

$$\begin{aligned} & - \operatorname{atan}\left(\frac{((32(12A^2Bb^9d^2 + 3A^3ab^8d^2 - 9AB^2ab^8d^2))/d^5 - ((32(16Ab^{10}d^4 - 4B^2a^2b^9d^4 + 12A^2b^8d^4))/d^5 - (32(16b^{10}d^4 + 24a^2b^8d^4))(a + b\tan(c + dx))^{1/2}) \cdot (-((8A^2ad^2 - 8B^2ad^2 + 16ABbd^2)^{2/4} - (16a^2d^4 + 16b^2d^4)(A^4 + 2A^2B^2 + B^4))^{1/2} - 4A^2ad^2 + 4B^2ad^2 - 8ABbd^2)/(16(a^2d^4 + b^2d^4))^{1/2})}{d^4} \cdot (-((8A^2ad^2 - 8B^2ad^2 + 16ABbd^2)^{2/4} - (16a^2d^4 + 16b^2d^4)(A^4 + 2A^2B^2 + B^4))^{1/2} - 4A^2ad^2 + 4B^2ad^2 - 8ABbd^2)/(16(a^2d^4 + b^2d^4))^{1/2} + (32(a + b\tan(c + dx))^{1/2})(16ABbd^9d^2 + 18A^2ab^8d^2 - 10B^2ab^8d^2))/d^4} \cdot (-((8A^2ad^2 - 8B^2ad^2 + 16ABbd^2)^{2/4} - (16a^2d^4 + 16b^2d^4)(A^4 + 2A^2B^2 + B^4))^{1/2} - 4A^2ad^2 + 4B^2ad^2 - 8ABbd^2)/(16(a^2d^4 + b^2d^4))^{1/2}) \cdot (-((8A^2ad^2 - 8B^2ad^2 + 16ABbd^2)^{2/4} - (16a^2d^4 + 16b^2d^4)(A^4 + 2A^2B^2 + B^4))^{1/2} - 4A^2ad^2 + 4B^2ad^2 - 8ABbd^2)/(16(a^2d^4 + b^2d^4))^{1/2}) \cdot (-((8A^2ad^2 - 8B^2ad^2 + 16ABbd^2)^{2/4} - (16a^2d^4 + 16b^2d^4)(A^4 + 2A^2B^2 + B^4))^{1/2} - 4A^2ad^2 + 4B^2ad^2 - 8ABbd^2)/(16(a^2d^4 + b^2d^4))^{1/2}) + (32(3A^4b^8 + B^4b^8)(a + b\tan(c + dx))^{1/2})/d^4} \cdot (-((8A^2ad^2 - 8B^2ad^2 + 16ABbd^2)^{2/4} - (16a^2d^4 + 16b^2d^4)(A^4 + 2A^2B^2 + B^4))^{1/2} - 4A^2ad^2 + 4B^2ad^2 - 8ABbd^2)/(16(a^2d^4 + b^2d^4))^{1/2}) \cdot (-((8A^2ad^2 - 8B^2ad^2 + 16ABbd^2)^{2/4} - (16a^2d^4 + 16b^2d^4)(A^4 + 2A^2B^2 + B^4))^{1/2} - 4A^2ad^2 + 4B^2ad^2 - 8ABbd^2)/(16(a^2d^4 + b^2d^4))^{1/2}) + (32(12A^2Bb^9d^2 + 3A^3ab^8d^2 - 9AB^2ab^8d^2))/d^5 - ((32(16Ab^{10}d^4 - 4B^2a^2b^9d^4 + 12A^2b^8d^4))/d^5 + (32(16b^{10}d^4 + 24a^2b^8d^4))(a + b\tan(c + dx))^{1/2})/d^5 \end{aligned}$$

3.348 $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

3.348.1 Optimal result 3312
 3.348.2 Mathematica [A] (verified) 3313
 3.348.3 Rubi [A] (warning: unable to verify) 3313
 3.348.4 Maple [B] (verified) 3318
 3.348.5 Fracas [B] (verification not implemented) 3319
 3.348.6 Sympy [F] 3319
 3.348.7 Maxima [F] 3320
 3.348.8 Giac [F(-1)] 3320
 3.348.9 Mupad [B] (verification not implemented) 3320

3.348.1 Optimal result

Integrand size = 33, antiderivative size = 169

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = \frac{(Ab-2aB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{a^{3/2}d} + \frac{(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib}d} - \frac{(iA-B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ib}d} - \frac{A \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad}$$

output

```
(A*b-2*B*a)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/a^(3/2)/d+(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d/(a-I*b)^(1/2)-(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d/(a+I*b)^(1/2)-A*cot(d*x+c)*(a+b*tan(d*x+c))^(1/2)/a/d
```

3.348.2 Mathematica [A] (verified)

Time = 3.14 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.19

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

$$= \frac{b(Ab-2aB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{a^{3/2}} + \frac{(A\sqrt{-b^2}+bB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-\sqrt{-b^2}}}\right)}{\sqrt{a-\sqrt{-b^2}}} - \frac{(A\sqrt{-b^2}-bB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+\sqrt{-b^2}}}\right)}{\sqrt{a+\sqrt{-b^2}}}$$

bd

input `Integrate[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `((b*(A*b - 2*a*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/a^(3/2) + ((A*Sqrt[-b^2] + b*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - Sqrt[-b^2]]])/Sqrt[a - Sqrt[-b^2]] - ((A*Sqrt[-b^2] - b*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + Sqrt[-b^2]]])/Sqrt[a + Sqrt[-b^2]] - (A*b*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]]/a)/(b*d)`

3.348.3 Rubi [A] (warning: unable to verify)

Time = 1.27 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.97, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 4092, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

↓ 3042

$$\int \frac{A+B \tan(c+dx)}{\tan(c+dx)^2 \sqrt{a+b \tan(c+dx)}} dx$$

↓ 4092

$$\frac{\int \frac{\cot(c+dx)(Ab \tan^2(c+dx)+2aA \tan(c+dx)+Ab-2aB)}{2\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad}$$

↓ 27

3.348. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\begin{aligned}
& \frac{\int \frac{\cot(c+dx)(Ab \tan^2(c+dx)+2aA \tan(c+dx)+Ab-2aB)}{\sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{Ab \tan(c+dx)^2+2aA \tan(c+dx)+Ab-2aB}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} \\
& \quad \downarrow 4136 \\
& \frac{\int \frac{2(aA+aB \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx + (Ab-2aB) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} \\
& \quad \downarrow 27 \\
& \frac{2 \int \frac{aA+aB \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + (Ab-2aB) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} \\
& \quad \downarrow 3042 \\
& \frac{2 \int \frac{aA+aB \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + (Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} \\
& \quad \downarrow 4022 \\
& \frac{(Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 2 \left(\frac{1}{2} a(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2} a(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx \right)}{2a} - \frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} \\
& \quad \downarrow 3042 \\
& \frac{(Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 2 \left(\frac{1}{2} a(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2} a(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx \right)}{2a} - \frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} \\
& \quad \downarrow 4020
\end{aligned}$$

3.348. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{(Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 2 \left(\frac{ad}{ia(A-iB) \int \frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))} - \frac{ia(A+iB) \int \frac{1}{(1+i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} \right)}{2a}$$

25

$$\frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{(Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 2 \left(\frac{ad}{ia(A+iB) \int \frac{1}{(i \tan(c+dx)+1) \sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))} - \frac{ia(A-iB) \int \frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} \right)}{2a}$$

73

$$\frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{(Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 2 \left(\frac{ad}{a(A+iB) \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}} + \frac{a(A-iB) \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} \right)}{2a}$$

221

$$\frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{(Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 2 \left(\frac{ad}{a(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)} + \frac{a(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a-ib}} \right)}{2a}$$

4117

$$\frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{\frac{(Ab-2aB) \int \frac{\cot(c+dx)}{\sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{d} + 2 \left(\frac{ad}{a(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)} + \frac{a(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a-ib}} \right)}{2a}$$

73

$$\frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{2(Ab-2aB) \int \frac{1}{\frac{a+b \tan(c+dx)}{b} - \frac{a}{b}} d\sqrt{a+b \tan(c+dx)} + 2 \left(\frac{ad}{a(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)} + \frac{a(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a-ib}} \right)}{2a}$$

221

3.348. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\frac{\frac{A \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} - \frac{2(Ab-2aB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ad}} + 2 \left(\frac{a(A-iB) \operatorname{arctan}\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} + \frac{a(A+iB) \operatorname{arctan}\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} \right)}{2a}$$

input `Int[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `-1/2*(2*((a*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d) + (a*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d)) - (2*(A*b - 2*a*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]]/(Sqrt[a]*d))/a - (A*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/(a*d)`

3.348.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 $\text{Int}[(a + (b \tan(e + f x)) + (c + (d \tan(e + f x)) + (f x)))^m, x_Symbol] := \text{Simp}[c(d/f) \text{Subst}[\text{Int}[(a + (b/d)x)^m/(d^2 + c x), x], x, d \tan[e + f x]], x] /; \text{FreeQ}\{a, b, c, d, e, f, m\}, x] \&\& \text{NeQ}[b c - a d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{EqQ}[c^2 + d^2, 0]$

rule 4022 $\text{Int}[(a + (b \tan(e + f x)) + (c + (d \tan(e + f x)) + (f x)))^m, x_Symbol] := \text{Simp}[(c + I d)/2 \text{Int}[(a + b \tan[e + f x])^m (1 - I \tan[e + f x]), x], x] + \text{Simp}[(c - I d)/2 \text{Int}[(a + b \tan[e + f x])^m (1 + I \tan[e + f x]), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m\}, x] \&\& \text{NeQ}[b c - a d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{IntegerQ}[m]$

rule 4092 $\text{Int}[(a + (b \tan(e + f x)) + (c + (d \tan(e + f x)) + (f x)))^m (A + (B \tan(e + f x)) + (f x))^n, x_Symbol] := \text{Simp}[b(A b - a B)(a + b \tan[e + f x])^{m+1} (c + d \tan[e + f x])^{n+1} / (f(m+1)(b c - a d)(a^2 + b^2)), x] + \text{Simp}[1 / ((m+1)(b c - a d)(a^2 + b^2)) \text{Int}[(a + b \tan[e + f x])^{m+1} (c + d \tan[e + f x])^n \text{Simp}[b B(b c(m+1) + a d(n+1)) + A(a(b c - a d)(m+1) - b^2 d(m+n+2)) - (A b - a B)(b c - a d)(m+1) \tan[e + f x] - b d(A b - a B)(m+n+2) \tan[e + f x]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, n\}, x] \&\& \text{NeQ}[b c - a d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{LtQ}[m, -1] \&\& (\text{IntegerQ}[m] || \text{IntegersQ}[2 m, 2 n]) \&\& !(\text{ILtQ}[n, -1] \&\& (! \text{IntegerQ}[m] || (\text{EqQ}[c, 0] \&\& \text{NeQ}[a, 0])))$

rule 4117 $\text{Int}[(a + (b \tan(e + f x)) + (c + (d \tan(e + f x)) + (f x)))^m (c + (d \tan(e + f x)) + (f x))^n (A + (C \tan(e + f x)) + (f x))^2, x_Symbol] := \text{Simp}[A/f \text{Subst}[\text{Int}[(a + b x)^m (c + d x)^n, x], x, \tan[e + f x]], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, C, m, n\}, x] \&\& \text{EqQ}[A, C]$

rule 4136 $\text{Int}[(c + (d \tan(e + f x)) + (f x))^n (A + (B \tan(e + f x)) + (f x)) + (C \tan(e + f x)) + (f x))^2 / ((a + (b \tan(e + f x)) + (f x))), x_Symbol] := \text{Simp}[1/(a^2 + b^2) \text{Int}[(c + d \tan[e + f x])^n \text{Simp}[b B + a(A - C) + (a B - b(A - C)) \tan[e + f x], x], x], x] + \text{Simp}[(A b^2 - a b B + a^2 C)/(a^2 + b^2) \text{Int}[(c + d \tan[e + f x])^n ((1 + \tan[e + f x]^2)/(a + b \tan[e + f x])), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C, n\}, x] \&\& \text{NeQ}[b c - a d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& ! \text{GtQ}[n, 0] \&\& ! \text{LeQ}[n, -1]$

3.348.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4056 vs. $2(143) = 286$.

Time = 0.22 (sec) , antiderivative size = 4057, normalized size of antiderivative = 24.01

method	result	size
derivativdivides	Expression too large to display	4057
default	Expression too large to display	4057

```
input int(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output -1/4/d*b^2/(a^2+b^2)^(3/2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*
a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1
/4/d/(a^2+b^2)^(3/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)
^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2+2/d
/(a^2+b^2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(
1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*a^2-1/4/d/
(a^2+b^2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*
x+c)-a-(a^2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2/d/(a^2+b^2)/(2
*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(
1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*a^2-1/4/d/(a^2+b^2)^(3/
2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-
(a^2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2-1/d/(a^2+b^2)^(3/2)/(
2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*
tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*a^3+1/d/(a^2+b^2)^(1/2
)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a
+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*a-1/d*b/(a^2+b^2)^(
1/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2
*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A-2/d*b^3/(a^2+b^2
)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*
(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A+1/4/d*b^2/...
```

3.348.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1783 vs. $2(139) = 278$.

Time = 1.99 (sec) , antiderivative size = 3582, normalized size of antiderivative = 21.20

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm
m="fricas")
```

```
output [1/2*(a^2*d*sqrt(-(a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)
)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*
B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*log((2*(A^3*B + A*B^3)*a - (A^4 -
B^4)*b)*sqrt(b*tan(d*x + c) + a) + ((A*a^3 + B*a^2*b + A*a*b^2 + B*b^3)*d^
3*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b
^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + (2*A*B^2*a^2 - (3*A^2*B - B^3)*a*b +
(A^3 - A*B^2)*b^2)*d)*sqrt(-(a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3
*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^
4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*tan(d*x + c) - a^2*d*sq
rt(-(a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 -
2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B
^2)*a)/((a^2 + b^2)*d^2))*log((2*(A^3*B + A*B^3)*a - (A^4 - B^4)*b)*sqrt(b
*tan(d*x + c) + a) - ((A*a^3 + B*a^2*b + A*a*b^2 + B*b^3)*d^3*sqrt(-(4*A^2
*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*
a^2*b^2 + b^4)*d^4)) + (2*A*B^2*a^2 - (3*A^2*B - B^3)*a*b + (A^3 - A*B^2)*
b^2)*d)*sqrt(-(a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*
b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b
+ (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*tan(d*x + c) - a^2*d*sqrt(((a^2 + b^2)
)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^
4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) - 2*A*B*b - (A^2 - B^2)*a)/((a^2...
```

3.348.6 Sympy [F]

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(A+B\tan(c+dx))\cot^2(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx$$

```
input integrate(cot(d*x+c)**2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)
```

3.348. $\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**2/sqrt(a + b*tan(c + d*x)), x)`

3.348.7 Maxima [F]

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx = \int \frac{(B \tan(dx + c) + A) \cot(dx + c)^2}{\sqrt{b \tan(dx + c) + a}} dx$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^2/sqrt(b*tan(d*x + c) + a), x)`

3.348.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.348.9 Mupad [B] (verification not implemented)

Time = 9.96 (sec) , antiderivative size = 9790, normalized size of antiderivative = 57.93

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `int((cot(c + d*x)^2*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x)`

output

```

atan(((((((8*(32*A*a*b^11*d^4 + 16*A*a^3*b^9*d^4 - 64*B*a^2*b^10*d^4 - 48*
B*a^4*b^8*d^4))/(a^2*d^5) - (16*(32*a^2*b^10*d^4 + 48*a^4*b^8*d^4)*(a + b*
tan(c + d*x))^(1/2)*(((8*A^2*a*d^2 - 8*B^2*a*d^2 + 16*A*B*b*d^2)^2/4 - (1
6*a^2*d^4 + 16*b^2*d^4)*(A^4 + 2*A^2*B^2 + B^4))^(1/2) - 4*A^2*a*d^2 + 4*B
^2*a*d^2 - 8*A*B*b*d^2)/(16*(a^2*d^4 + b^2*d^4)))^(1/2))/(a^2*d^4))*(((8*
A^2*a*d^2 - 8*B^2*a*d^2 + 16*A*B*b*d^2)^2/4 - (16*a^2*d^4 + 16*b^2*d^4)*(A
^4 + 2*A^2*B^2 + B^4))^(1/2) - 4*A^2*a*d^2 + 4*B^2*a*d^2 - 8*A*B*b*d^2)/(1
6*(a^2*d^4 + b^2*d^4)))^(1/2) - (16*(a + b*tan(c + d*x))^(1/2)*(20*A^2*a^3
*b^8*d^2 - 36*B^2*a^3*b^8*d^2 - 4*A^2*a*b^10*d^2 + 48*A*B*a^2*b^9*d^2))/(a
^2*d^4))*(((8*A^2*a*d^2 - 8*B^2*a*d^2 + 16*A*B*b*d^2)^2/4 - (16*a^2*d^4 +
16*b^2*d^4)*(A^4 + 2*A^2*B^2 + B^4))^(1/2) - 4*A^2*a*d^2 + 4*B^2*a*d^2 -
8*A*B*b*d^2)/(16*(a^2*d^4 + b^2*d^4)))^(1/2) + (8*(4*A^3*b^11*d^2 + 16*A^3
*a^2*b^9*d^2 + 12*B^3*a^3*b^8*d^2 + 12*A^2*B*a*b^10*d^2 - 48*A*B^2*a^2*b^9
*d^2 - 36*A^2*B*a^3*b^8*d^2))/(a^2*d^5))*(((8*A^2*a*d^2 - 8*B^2*a*d^2 + 1
6*A*B*b*d^2)^2/4 - (16*a^2*d^4 + 16*b^2*d^4)*(A^4 + 2*A^2*B^2 + B^4))^(1/2
) - 4*A^2*a*d^2 + 4*B^2*a*d^2 - 8*A*B*b*d^2)/(16*(a^2*d^4 + b^2*d^4)))^(1/
2) - (16*(a + b*tan(c + d*x))^(1/2)*(A^2*B^2*b^10 - A^4*b^10 + 2*A^4*a^2*b
^8 + 6*B^4*a^2*b^8 - 4*A*B^3*a*b^9 + 4*A^3*B*a*b^9))/(a^2*d^4))*(((8*A^2*
a*d^2 - 8*B^2*a*d^2 + 16*A*B*b*d^2)^2/4 - (16*a^2*d^4 + 16*b^2*d^4)*(A^4 +
2*A^2*B^2 + B^4))^(1/2) - 4*A^2*a*d^2 + 4*B^2*a*d^2 - 8*A*B*b*d^2)/(16...

```


3.349 $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

3.349.1 Optimal result	3322
3.349.2 Mathematica [A] (verified)	3323
3.349.3 Rubi [A] (warning: unable to verify)	3323
3.349.4 Maple [B] (verified)	3329
3.349.5 Fracas [B] (verification not implemented)	3330
3.349.6 Sympy [F]	3330
3.349.7 Maxima [F]	3331
3.349.8 Giac [F(-1)]	3331
3.349.9 Mupad [B] (verification not implemented)	3331

3.349.1 Optimal result

Integrand size = 33, antiderivative size = 224

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = \frac{(8a^2A - 3Ab^2 + 4abB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{4a^{5/2}d} - \frac{(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib}d} - \frac{(A + iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ib}d} + \frac{(3Ab - 4aB) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{4a^2d} - \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2ad}$$

output `1/4*(8*A*a^2-3*A*b^2+4*B*a*b)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/a^(5/2)/d-(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d/(a-I*b)^(1/2)-(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d/(a+I*b)^(1/2)+1/4*(3*A*b-4*B*a)*cot(d*x+c)*(a+b*tan(d*x+c))^(1/2)/a^2/d-1/2*A*cot(d*x+c)^2*(a+b*tan(d*x+c))^(1/2)/a/d`

3.349.2 Mathematica [A] (verified)

Time = 6.32 (sec) , antiderivative size = 360, normalized size of antiderivative = 1.61

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

$$= 2b^3 \left(\frac{A \operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ab^3}} - \frac{3A \operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a}}\right)}{8a^{5/2}b} + \frac{B \operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a}}\right)}{2a^{3/2}b^2} - \frac{(A\sqrt{-b^2}+bB) \operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a}}\right)}{2b^3\sqrt{-b^2}\sqrt{a}} \right)$$

```
input Integrate[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x
]
```

```
output (2*b^3*((A*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/(Sqrt[a]*b^3) - (3*A
*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/(8*a^(5/2)*b) + (B*ArcTanh[Sqr
t[a + b*Tan[c + d*x]]/Sqrt[a]])/(2*a^(3/2)*b^2) - ((A*Sqrt[-b^2] + b*B)*Ar
cTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - Sqrt[-b^2]]])/(2*b^3*Sqrt[-b^2]*Sq
rt[a - Sqrt[-b^2]]) - (b*(A*Sqrt[-b^2] - b*B)*ArcTanh[Sqrt[a + b*Tan[c + d
*x]]/Sqrt[a + Sqrt[-b^2]]])/(2*(-b^2)^(5/2)*Sqrt[a + Sqrt[-b^2]]) + (3*A*C
ot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/(8*a^2*b^2) - (B*Cot[c + d*x]*Sqrt[a
+ b*Tan[c + d*x]])/(2*a*b^3) - (A*Cot[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]]
)/(4*a*b^3))/d
```

3.349.3 Rubi [A] (warning: unable to verify)Time = 1.70 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.03, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A+B\tan(c+dx)}{\tan(c+dx)^3\sqrt{a+b\tan(c+dx)}} dx$$

3.349. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

$$\begin{array}{c}
\downarrow 4092 \\
\frac{\int \frac{\cot^2(c+dx)(3Ab \tan^2(c+dx)+4aA \tan(c+dx)+3Ab-4aB)}{2\sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2ad} \\
\downarrow 27 \\
\frac{\int \frac{\cot^2(c+dx)(3Ab \tan^2(c+dx)+4aA \tan(c+dx)+3Ab-4aB)}{\sqrt{a+b \tan(c+dx)}} dx}{4a} - \frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2ad} \\
\downarrow 3042 \\
\frac{\int \frac{3Ab \tan(c+dx)^2+4aA \tan(c+dx)+3Ab-4aB}{\tan(c+dx)^2\sqrt{a+b \tan(c+dx)}} dx}{4a} - \frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2ad} \\
\downarrow 4132 \\
\frac{\int \frac{\cot(c+dx)(8Aa^2+8B \tan(c+dx)a^2+4bBa-3Ab^2-b(3Ab-4aB) \tan^2(c+dx))}{2\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{(3Ab-4aB) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad} \\
\frac{4a}{2ad} \frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2ad} \\
\downarrow 27 \\
\frac{\int \frac{\cot(c+dx)(8Aa^2+8B \tan(c+dx)a^2+4bBa-3Ab^2-b(3Ab-4aB) \tan^2(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{(3Ab-4aB) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad} \\
\frac{4a}{2ad} \frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2ad} \\
\downarrow 3042 \\
\frac{\int \frac{8Aa^2+8B \tan(c+dx)a^2+4bBa-3Ab^2-b(3Ab-4aB) \tan(c+dx)^2}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{(3Ab-4aB) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad} \\
\frac{4a}{2ad} \frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2ad} \\
\downarrow 4136 \\
\frac{(8a^2A+4abB-3Ab^2) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx + \int \frac{8(a^2B-a^2A \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx}{2a} - \frac{(3Ab-4aB) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad} \\
\frac{4a}{2ad} \frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2ad} \\
\downarrow 27
\end{array}$$

3.349. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\frac{(8a^2A+4abB-3Ab^2) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx + 8 \int \frac{a^2B-a^2A \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx - \frac{(3Ab-4aB) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad}}{2a} - \frac{4a}{2ad} \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2ad}$$

↓ 3042

$$\frac{(8a^2A+4abB-3Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 8 \int \frac{a^2B-a^2A \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx - \frac{(3Ab-4aB) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad}}{2a} - \frac{4a}{2ad} \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2ad}$$

↓ 4022

$$\frac{(3Ab-4aB) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} + \frac{(8a^2A+4abB-3Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 8 \left(\frac{1}{2} a^2(B+iA) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2} a^2(-E) \right)}{2a} - \frac{4a}{2ad} \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2ad}$$

↓ 3042

$$\frac{(3Ab-4aB) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} + \frac{(8a^2A+4abB-3Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 8 \left(\frac{1}{2} a^2(B+iA) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2} a^2(-E) \right)}{2a} - \frac{4a}{2ad} \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2ad}$$

↓ 4020

$$\frac{(3Ab-4aB) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} + \frac{(8a^2A+4abB-3Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 8 \left(\frac{ia^2(B+iA) \int -\frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} dx}{2d} \right)}{2a} - \frac{4a}{2ad} \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2ad}$$

↓ 25

$$\frac{(3Ab-4aB) \cot(c+dx) \sqrt{a+b \tan(c+dx)}}{ad} + \frac{(8a^2A+4abB-3Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + 8 \left(-\frac{ia^2(B+iA) \int \frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} dx}{2d} \right)}{2a} - \frac{4a}{2ad} \frac{A \cot^2(c+dx) \sqrt{a+b \tan(c+dx)}}{2ad}$$

↓ 73

3.349. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\begin{aligned}
 & -\frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2ad} - \frac{(3Ab-4aB) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad} + \frac{(8a^2A+4abB-3Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 8 \left(\frac{a^2(B+iA) \int \frac{1}{i \tan^2(\frac{c+dx}{b} + \frac{ia}{b} + 1)} d\sqrt{a+b \tan(c+dx)}}{bd} \right)}{4a} \\
 & \quad \downarrow \text{221} \\
 & -\frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2ad} - \frac{(3Ab-4aB) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad} + \frac{(8a^2A+4abB-3Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 8 \left(\frac{a^2(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{a^2(-B+iA)}{2a} \right)}{4a} \\
 & \quad \downarrow \text{4117} \\
 & -\frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2ad} - \frac{(3Ab-4aB) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad} + \frac{(8a^2A+4abB-3Ab^2) \int \frac{\cot(c+dx)}{\sqrt{a+b \tan(c+dx)}} d \tan(c+dx) + 8 \left(\frac{a^2(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{a^2(-B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} \right)}{4a} \\
 & \quad \downarrow \text{73} \\
 & -\frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2ad} - \frac{(3Ab-4aB) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad} + \frac{2(8a^2A+4abB-3Ab^2) \int \frac{1}{\frac{a+b \tan(c+dx)}{b} - \frac{a}{b}} d\sqrt{a+b \tan(c+dx)} + 8 \left(\frac{a^2(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{a^2(-B+iA)}{2a} \right)}{4a} \\
 & \quad \downarrow \text{221} \\
 & -\frac{A \cot^2(c+dx)\sqrt{a+b \tan(c+dx)}}{2ad} - \frac{(3Ab-4aB) \cot(c+dx)\sqrt{a+b \tan(c+dx)}}{ad} + \frac{2(8a^2A+4abB-3Ab^2) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right) + 8 \left(\frac{a^2(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{a^2(-B+iA)}{2a} \right)}{4a}
 \end{aligned}$$

input `Int[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

3.349. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

```
output -1/2*(A*Cot[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]]/(a*d) - ((8*((a^2*(I*A +
B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d) - (a^2*(I*A - B)*
ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d)) - (2*(8*a^2*A - 3*A
*b^2 + 4*a*b*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]]/(Sqrt[a]*d))/(2
*a) - ((3*A*b - 4*a*B)*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]]/(a*d))/(4*a)
```

3.349.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.349.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4174 vs. $2(190) = 380$.

Time = 0.24 (sec) , antiderivative size = 4175, normalized size of antiderivative = 18.64

method	result	size
derivativdivides	Expression too large to display	4175
default	Expression too large to display	4175

```
input int(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output -1/d/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+
c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*
a^3-1/4/d/(a^2+b^2)^(3/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2
+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^
2+1/4/d/(a^2+b^2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-
b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2/d/(a^2
+b^2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-
2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a^2+1/d/(a^2+b^
2)^(1/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2
*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a+1/d/(a^2+b
^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1
/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a^3+1/4/d/(
a^2+b^2)^(3/2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*t
an(d*x+c)-a-(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2-1/d/(a^2+
b^2)^(1/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(
1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a-1/4/d/(a
^2+b^2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(
1/2)+(a^2+b^2)^(1/2))*A*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+2/d/(a^2+b^2)/(2*(
a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1
/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*A*a^2-2/d*b^3/(a^2+b^2)^...
```


3.349.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1836 vs. $2(184) = 368$.

Time = 7.51 (sec) , antiderivative size = 3688, normalized size of antiderivative = 16.46

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm
m="fricas")
```

```
output [1/8*(4*a^3*d*sqrt(((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*log((2*(A^3*B + A*B^3)*a - (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) + ((B*a^3 - A*a^2*b + B*a*b^2 - A*b^3)*d^3*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) - (2*A^2*B*a^2 - (A^3 - 3*A*B^2)*a*b - (A^2*B - B^3)*b^2)*d)*sqrt(((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*tan(d*x + c)^2 - 4*a^3*d*sqrt(((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*log((2*(A^3*B + A*B^3)*a - (A^4 - B^4)*b)*sqrt(b*tan(d*x + c) + a) - ((B*a^3 - A*a^2*b + B*a*b^2 - A*b^3)*d^3*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) - (2*A^2*B*a^2 - (A^3 - 3*A*B^2)*a*b - (A^2*B - B^3)*b^2)*d)*sqrt(((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*A*B*b + (A^2 - B^2)*a)/((a^2 + b^2)*d^2))*tan(d*x + c)^2 - 4*a^3*d*sqrt(-((a^2 + b^2)*d^2*sqrt(-(4*A^2*B^2*a^2 - 4*(A^3*B - A*B^3)*a*b + (A^4 - 2*A^2*B^2 + B^4)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) - 2*A*B*b - (A^2 - B^2)*a...
```

3.349.6 Sympy [F]

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(A+B\tan(c+dx))\cot^3(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx$$

```
input integrate(cot(d*x+c)**3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)
```

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**3/sqrt(a + b*tan(c + d*x)), x)`

3.349.7 Maxima [F]

$$\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx = \int \frac{(B \tan(dx + c) + A) \cot(dx + c)^3}{\sqrt{b \tan(dx + c) + a}} dx$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^3/sqrt(b*tan(d*x + c) + a), x)`

3.349.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.349.9 Mupad [B] (verification not implemented)

Time = 10.91 (sec) , antiderivative size = 13182, normalized size of antiderivative = 58.85

$$\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `int((cot(c + d*x)^3*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x)`

output

$$\begin{aligned}
& - \left((5A^2b^2 - 4B^2ab)(a + b\tan(c + dx))^{1/2} \right) / (4a) - \left((3A^2b^2 - 4B^2ab)(a + b\tan(c + dx))^{3/2} \right) / (4a^2) / (d(a + b\tan(c + dx))^2 + a^2d - 2ad(a + b\tan(c + dx))) - \operatorname{atan}\left(\frac{(640A^2a^4b^{10}d^4 - 384A^2a^2b^{12}d^4 + 768A^2a^6b^8d^4 + 512B^2a^3b^{11}d^4 + 256B^2a^5b^9d^4)}{(2a^4d^5) + ((512a^4b^{10}d^4 + 768a^6b^8d^4)(a + b\tan(c + dx))^{1/2}) - ((8A^2ad^2 - 8B^2ad^2 + 16ABbd^2)^{2/4} - (16a^2d^4 + 16b^2d^4)(A^4 + 2A^2B^2 + B^4))^{1/2} - 4A^2ad^2 + 4B^2ad^2 - 8ABbd^2}{(16(a^2d^4 + b^2d^4))^{1/2}} \right) / (a^4d^4) * \left(-((8A^2ad^2 - 8B^2ad^2 + 16ABbd^2)^{2/4} - (16a^2d^4 + 16b^2d^4)(A^4 + 2A^2B^2 + B^4))^{1/2} - 4A^2ad^2 + 4B^2ad^2 - 8ABbd^2 \right) / (16(a^2d^4 + b^2d^4))^{1/2} - ((a + b\tan(c + dx))^{1/2} * (576A^2a^5b^8d^2 - 192A^2a^3b^{10}d^2 + 64B^2a^3b^{10}d^2 - 320B^2a^5b^8d^2 + 36A^2ab^{12}d^2 - 96ABa^2b^{11}d^2 + 768ABa^4b^9d^2)) / (a^4d^4) * \left(-((8A^2ad^2 - 8B^2ad^2 + 16ABbd^2)^{2/4} - (16a^2d^4 + 16b^2d^4)(A^4 + 2A^2B^2 + B^4))^{1/2} - 4A^2ad^2 + 4B^2ad^2 - 8ABbd^2 \right) / (16(a^2d^4 + b^2d^4))^{1/2} + (64B^3a^2b^{11}d^2 - 192A^3a^5b^8d^2 + 256B^3a^4b^9d^2 + 36A^2Bb^{13}d^2 + 36A^3ab^{12}d^2 - 96AB^2ab^{12}d^2 - 384AB^2a^3b^{10}d^2 + 576AB^2a^5b^8d^2 + 96A^2B^2ab^{11}d^2 - 768A^2B^2a^4b^9d^2) / (2a^4d^5) * \left(-((8A^2ad^2 - 8B^2ad^2 + 16ABbd^2)^{2/4} - (16a^2d^4 + 16b^2d^4)(A^4 + 2A^2B^2 + B^4))^{1/2} - 4A^2ad^2 + 4B^2ad^2 - 8ABbd^2 \right) / (16(a^2d^4 + b^2d^4))^{1/2}
\end{aligned}$$

3.350
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.350.1 Optimal result 3333
 3.350.2 Mathematica [C] (verified) 3334
 3.350.3 Rubi [A] (warning: unable to verify) 3334
 3.350.4 Maple [B] (verified) 3339
 3.350.5 Fracas [B] (verification not implemented) 3340
 3.350.6 Sympy [F] 3341
 3.350.7 Maxima [F(-1)] 3342
 3.350.8 Giac [F(-1)] 3342
 3.350.9 Mupad [B] (verification not implemented) 3342

3.350.1 Optimal result

Integrand size = 33, antiderivative size = 264

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{3/2}d} + \frac{(A+iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a+ib)^{3/2}d} + \frac{2a(Ab-aB) \tan^2(c+dx)}{b(a^2+b^2)d\sqrt{a+b \tan(c+dx)}} + \frac{2(6a^2Ab+3Ab^3-8a^3B-5ab^2B)\sqrt{a+b \tan(c+dx)}}{3b^3(a^2+b^2)d} - \frac{2(3aAb-4a^2B-b^2B)\tan(c+dx)\sqrt{a+b \tan(c+dx)}}{3b^2(a^2+b^2)d}$$

```
output (A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(3/2)/d+(A+I
*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(3/2)/d+2/3*(6*A
*a^2*b+3*A*b^3-8*B*a^3-5*B*a*b^2)*(a+b*tan(d*x+c))^(1/2)/b^3/(a^2+b^2)/d-2
/3*(3*A*a*b-4*B*a^2-B*b^2)*(a+b*tan(d*x+c))^(1/2)*tan(d*x+c)/b^2/(a^2+b^2)
/d+2*a*(A*b-B*a)*tan(d*x+c)^2/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^(1/2)
```

3.350.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 3.80 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.14

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \frac{3iA \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ib}} \right) + \frac{2(6aAb-}{b^2\sqrt{a+}}$$

input `Integrate[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `((3*I)*A*(ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]]/Sqrt[a - I*b] - ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]]/Sqrt[a + I*b]) + (2*(6*a*A*b - 8*a^2*B + 3*b^2*B))/(b^2*Sqrt[a + b*Tan[c + d*x]]) + ((3*I)*(a*A + b*B)*((a + I*b)*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Tan[c + d*x])/(a - I*b)] - (a - I*b)*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Tan[c + d*x])/(a + I*b)]))/((a^2 + b^2)*Sqrt[a + b*Tan[c + d*x]]) + (2*(3*A*b - 4*a*B)*Tan[c + d*x])/(b*Sqrt[a + b*Tan[c + d*x]]) + (2*B*Tan[c + d*x]^2)/Sqrt[a + b*Tan[c + d*x]])/(3*b*d)`

3.350.3 Rubi [A] (warning: unable to verify)

Time = 1.38 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.03, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 4088, 27, 3042, 4130, 27, 3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$$

↓ 3042

$$\int \frac{\tan(c+dx)^3(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$$

↓ 4088

$$\begin{aligned}
& \frac{2 \int -\frac{\tan(c+dx)((-4Ba^2+3Aba-b^2B)\tan^2(c+dx)-b(Ab-aB)\tan(c+dx)+4a(Ab-aB))}{2\sqrt{a+b\tan(c+dx)}} dx}{b(a^2+b^2)} + \\
& \frac{2a(Ab-aB)\tan^2(c+dx)}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \\
& \quad \downarrow \text{27} \\
& \frac{2a(Ab-aB)\tan^2(c+dx)}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \\
& \frac{\int \frac{\tan(c+dx)((-4Ba^2+3Aba-b^2B)\tan^2(c+dx)-b(Ab-aB)\tan(c+dx)+4a(Ab-aB))}{\sqrt{a+b\tan(c+dx)}} dx}{b(a^2+b^2)} \\
& \quad \downarrow \text{3042} \\
& \frac{2a(Ab-aB)\tan^2(c+dx)}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \\
& \frac{\int \frac{\tan(c+dx)((-4Ba^2+3Aba-b^2B)\tan(c+dx)^2-b(Ab-aB)\tan(c+dx)+4a(Ab-aB))}{\sqrt{a+b\tan(c+dx)}} dx}{b(a^2+b^2)} \\
& \quad \downarrow \text{4130} \\
& \frac{2a(Ab-aB)\tan^2(c+dx)}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \\
& \frac{2 \int -\frac{-3(aA+bB)\tan(c+dx)b^2+(-8Ba^3+6Aba^2-5b^2Ba+3Ab^3)\tan^2(c+dx)+2a(-4Ba^2+3Aba-b^2B)}{2\sqrt{a+b\tan(c+dx)}} dx}{3b} + \frac{2(-4a^2B+3aAb-b^2B)\tan(c+dx)\sqrt{a+b\tan(c+dx)}}{3bd} \\
& \quad \downarrow \text{27} \\
& \frac{2a(Ab-aB)\tan^2(c+dx)}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \\
& \frac{2(-4a^2B+3aAb-b^2B)\tan(c+dx)\sqrt{a+b\tan(c+dx)}}{3bd} - \frac{\int \frac{-3(aA+bB)\tan(c+dx)b^2+(-8Ba^3+6Aba^2-5b^2Ba+3Ab^3)\tan^2(c+dx)+2a(-4Ba^2+3Aba-b^2B)}{\sqrt{a+b\tan(c+dx)}} dx}{3b} \\
& \quad \downarrow \text{3042} \\
& \frac{2a(Ab-aB)\tan^2(c+dx)}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \\
& \frac{2(-4a^2B+3aAb-b^2B)\tan(c+dx)\sqrt{a+b\tan(c+dx)}}{3bd} - \frac{\int \frac{-3(aA+bB)\tan(c+dx)b^2+(-8Ba^3+6Aba^2-5b^2Ba+3Ab^3)\tan(c+dx)+2a(-4Ba^2+3Aba-b^2B)}{\sqrt{a+b\tan(c+dx)}} dx}{3b} \\
& \quad \downarrow \text{4113}
\end{aligned}$$

3.350. $\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{2a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} - \\
 & \frac{2(-4a^2B + 3aAb - b^2B) \tan(c + dx) \sqrt{a + b \tan(c + dx)}}{3bd} - \frac{\int \frac{-3(Ab - aB)b^2 - 3(aA + bB) \tan(c + dx)b^2}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2(-8a^3B + 6a^2Ab - 5ab^2B + 3Ab^3) \sqrt{a + b \tan(c + dx)}}{bd}}{3b} \\
 & \frac{2a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2)} - \\
 & \frac{2(-4a^2B + 3aAb - b^2B) \tan(c + dx) \sqrt{a + b \tan(c + dx)}}{3bd} - \frac{\int \frac{-3(Ab - aB)b^2 - 3(aA + bB) \tan(c + dx)b^2}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2(-8a^3B + 6a^2Ab - 5ab^2B + 3Ab^3) \sqrt{a + b \tan(c + dx)}}{bd}}{3b} \\
 & \frac{2a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} - \\
 & \frac{2(-4a^2B + 3aAb - b^2B) \tan(c + dx) \sqrt{a + b \tan(c + dx)}}{3bd} - \frac{-\frac{3}{2}b^2(b + ia)(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{3}{2}b^2(a + ib)(B + iA) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2(-8a^3B + 6a^2Ab - 5ab^2B + 3Ab^3) \sqrt{a + b \tan(c + dx)}}{bd}}{3b} \\
 & \frac{2a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} - \\
 & \frac{2(-4a^2B + 3aAb - b^2B) \tan(c + dx) \sqrt{a + b \tan(c + dx)}}{3bd} - \frac{-\frac{3}{2}b^2(b + ia)(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{3}{2}b^2(a + ib)(B + iA) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2(-8a^3B + 6a^2Ab - 5ab^2B + 3Ab^3) \sqrt{a + b \tan(c + dx)}}{bd}}{3b} \\
 & \frac{2a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} - \\
 & \frac{2(-4a^2B + 3aAb - b^2B) \tan(c + dx) \sqrt{a + b \tan(c + dx)}}{3bd} - \frac{3ib^2(a + ib)(B + iA) \int \frac{1}{(1 - i \tan(c + dx)) \sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} + \frac{3ib^2(b + ia)(A + iB) \int \frac{1}{(1 + i \tan(c + dx)) \sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d}}{3b} \\
 & \frac{2a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} - \\
 & \frac{2(-4a^2B + 3aAb - b^2B) \tan(c + dx) \sqrt{a + b \tan(c + dx)}}{3bd} - \frac{3ib^2(a + ib)(B + iA) \int \frac{1}{(1 - i \tan(c + dx)) \sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} - \frac{3ib^2(b + ia)(A + iB) \int \frac{1}{(1 + i \tan(c + dx)) \sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d}}{3b} \\
 & \frac{2a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} - \\
 & \frac{2(-4a^2B + 3aAb - b^2B) \tan(c + dx) \sqrt{a + b \tan(c + dx)}}{3bd} - \frac{3ib^2(a + ib)(B + iA) \int \frac{1}{(1 - i \tan(c + dx)) \sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} - \frac{3ib^2(b + ia)(A + iB) \int \frac{1}{(1 + i \tan(c + dx)) \sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d}}{3b}
 \end{aligned}$$

3.350. $\int \frac{\tan^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx$

$$\frac{2a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} - \frac{2(-4a^2B + 3aAb - b^2B) \tan(c + dx) \sqrt{a + b \tan(c + dx)}}{3bd} - \frac{3b(b+ia)(A+iB) \int \frac{1}{-i \tan^2(c+dx) - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \frac{3b(a+ib)(B+iA) \int \frac{1}{i \tan^2(c+dx) + \frac{ib}{b} - 1} d\sqrt{a+b \tan(c+dx)}}{3bd}$$

↓ 221

$$\frac{2a(Ab - aB) \tan^2(c + dx)}{bd(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} - \frac{2(-8a^3B + 6a^2Ab - 5ab^2B + 3Ab^3) \sqrt{a + b \tan(c + dx)}}{bd} + \frac{3b^2(a+ib)(B+iA) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{3bd\sqrt{a-ib}} - \frac{3b^2(a-ib)(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{3bd\sqrt{a+ib}}$$

```
input Int[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]
```

```
output (2*a*(A*b - a*B)*Tan[c + d*x]^2)/(b*(a^2 + b^2)*d*sqrt[a + b*Tan[c + d*x]]
) - ((2*(3*a*A*b - 4*a^2*B - b^2*B)*Tan[c + d*x]*sqrt[a + b*Tan[c + d*x]])
/(3*b*d) - ((3*(a + I*b)*b^2*(I*A + B)*ArcTan[Tan[c + d*x]/sqrt[a - I*b]])
/(sqrt[a - I*b]*d) - (3*b^2*(I*a + b)*(A + I*B)*ArcTan[Tan[c + d*x]/sqrt[a
+ I*b]])/(sqrt[a + I*b]*d) + (2*(6*a^2*A*b + 3*A*b^3 - 8*a^3*B - 5*a*b^2*
B)*sqrt[a + b*Tan[c + d*x]])/(b*d))/(3*b))/(b*(a^2 + b^2))
```

3.350.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

3.350. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

```
rule 4130 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
) + (f_.)*(x_)^2), x_Symbol] :> Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

3.350.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3753 vs. $2(236) = 472$.

Time = 0.18 (sec) , antiderivative size = 3754, normalized size of antiderivative = 14.22

method	result	size
parts	Expression too large to display	3754
derivativedivides	Expression too large to display	8025
default	Expression too large to display	8025

```
input int(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,method=_RETURNV
ERBOSE)
```

output

```
A*(2/d/b^2*(a+b*tan(d*x+c))^(1/2)-1/4/d/(a^2+b^2)^2*ln(b*tan(d*x+c)+a+(a+b
*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+
b^2)^(1/2)+2*a)^(1/2)*a^2-1/4/d*b^2/(a^2+b^2)^2*ln(b*tan(d*x+c)+a+(a+b*tan
(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)
^(1/2)+2*a)^(1/2)+1/2/d/(a^2+b^2)^(5/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))
^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2
*a)^(1/2)*a^3+1/2/d*b^2/(a^2+b^2)^(5/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))
^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2
*a)^(1/2)*a+1/d/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a
+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a
)^(1/2))*a^2-1/d/(a^2+b^2)^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*
tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(
1/2))*a^3+1/d*b^2/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*
(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2
*a)^(1/2))-1/d*b^2/(a^2+b^2)^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+
b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a
)^(1/2))*a-2/d*b^2/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*
(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2
*a)^(1/2))*a^2-2/d*b^4/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arcta
n((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)...
```

3.350.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4421 vs. $2(229) = 458$.

Time = 0.71 (sec) , antiderivative size = 4421, normalized size of antiderivative = 16.75

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="fricas")`

```

output 1/6*(3*((a^2*b^4 + b^6)*d*tan(d*x + c) + (a^3*b^3 + a*b^5)*d)*sqrt((6*A*B*
a^2*b - 2*A*B*b^3 + (A^2 - B^2)*a^3 - 3*(A^2 - B^2)*a*b^2 + (a^6 + 3*a^4*b
^2 + 3*a^2*b^4 + b^6)*d^2*sqrt(-(4*A^2*B^2*a^6 - 12*(A^3*B - A*B^3)*a^5*b
+ 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 + 40*(A^3*B - A*B^3)*a^3*b^3 - 6*
(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 - 12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2*
B^2 + B^4)*b^6)/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8
+ 6*a^2*b^10 + b^12)*d^4)))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2))*lo
g(-(2*(A^3*B + A*B^3)*a^3 - 3*(A^4 - B^4)*a^2*b - 6*(A^3*B + A*B^3)*a*b^2
+ (A^4 - B^4)*b^3)*sqrt(b*tan(d*x + c) + a) + ((B*a^8 - 2*A*a^7*b + 2*B*a^
6*b^2 - 6*A*a^5*b^3 - 6*A*a^3*b^5 - 2*B*a^2*b^6 - 2*A*a*b^7 - B*b^8)*d^3*s
qrt(-(4*A^2*B^2*a^6 - 12*(A^3*B - A*B^3)*a^5*b + 3*(3*A^4 - 14*A^2*B^2 + 3
*B^4)*a^4*b^2 + 40*(A^3*B - A*B^3)*a^3*b^3 - 6*(A^4 - 8*A^2*B^2 + B^4)*a^2
*b^4 - 12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2*B^2 + B^4)*b^6)/((a^12 + 6*
a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 + b^12)*d^4))
- (2*A^2*B*a^5 - (3*A^3 - 7*A*B^2)*a^4*b - 2*(7*A^2*B - 3*B^3)*a^3*b^2 +
4*(A^3 - 4*A*B^2)*a^2*b^3 + 2*(4*A^2*B - B^3)*a*b^4 - (A^3 - A*B^2)*b^5)*d
)*sqrt((6*A*B*a^2*b - 2*A*B*b^3 + (A^2 - B^2)*a^3 - 3*(A^2 - B^2)*a*b^2 +
(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*sqrt(-(4*A^2*B^2*a^6 - 12*(A^3*B -
A*B^3)*a^5*b + 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 + 40*(A^3*B - A*B^3
)*a^3*b^3 - 6*(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 - 12*(A^3*B - A*B^3)*a*b^...

```

3.350.6 Sympy [F]

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \int \frac{(A+B \tan(c+dx)) \tan^3(c+dx)}{(a+b \tan(c+dx))^{\frac{3}{2}}} dx$$

```
input integrate(tan(d*x+c)**3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)
```

```
output Integral((A + B*tan(c + d*x))*tan(c + d*x)**3/(a + b*tan(c + d*x))**(3/2),
x)
```

3.350.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\tan^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm m="maxima")`

output `Timed out`

3.350.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm m="giac")`

output `Timed out`

3.350.9 Mupad [B] (verification not implemented)

Time = 23.82 (sec) , antiderivative size = 5811, normalized size of antiderivative = 22.01

$$\int \frac{\tan^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `int((tan(c + d*x)^3*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x)`

output

$$\begin{aligned}
& (\log(24A^3a^3b^6d^2 - (((((96A^4a^2b^4d^4 - 16A^4b^6d^4 - 144A^4a^4b^2d^4)^{(1/2)} + 4A^2a^3d^2 - 12A^2ab^2d^2)/(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{(1/2)} * (((((96A^4a^2b^4d^4 - 16A^4b^6d^4 - 144A^4a^4b^2d^4)^{(1/2)} + 4A^2a^3d^2 - 12A^2ab^2d^2)/(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{(1/2)} * (a + b \tan(c + dx))^{(1/2)} * (64a^5b^{12}d^5 + 320a^3b^{10}d^5 + 640a^5b^8d^5 + 640a^7b^6d^5 + 320a^9b^4d^5 + 64a^{11}b^2d^5)))/4 - 32Ab^{12}d^4 - 96Aa^2b^{10}d^4 - 64Aa^4b^8d^4 + 64Aa^6b^6d^4 + 96Aa^8b^4d^4 + 32Aa^{10}b^2d^4)/4 + (a + b \tan(c + dx))^{(1/2)} * (16A^2b^{10}d^3 + 32A^2a^2b^8d^3 - 32A^2a^6b^4d^3 - 16A^2a^8b^2d^3)) * (((96A^4a^2b^4d^4 - 16A^4b^6d^4 - 144A^4a^4b^2d^4)^{(1/2)} + 4A^2a^3d^2 - 12A^2ab^2d^2)/(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{(1/2)})/4 \\
& + 24A^3a^5b^4d^2 + 8A^3a^7b^2d^2 + 8A^3a^8b^8d^2 * (((96A^4a^2b^4d^4 - 16A^4b^6d^4 - 144A^4a^4b^2d^4)^{(1/2)} + 4A^2a^3d^2 - 12A^2ab^2d^2)/(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{(1/2)})/4 \\
& + (\log(24A^3a^3b^6d^2 - ((((-(96A^4a^2b^4d^4 - 16A^4b^6d^4 - 144A^4a^4b^2d^4)^{(1/2)} - 4A^2a^3d^2 + 12A^2ab^2d^2)/(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{(1/2)} * ((((-(96A^4a^2b^4d^4 - 16A^4b^6d^4 - 144A^4a^4b^2d^4)^{(1/2)} - 4A^2a^3d^2 + 12A^2ab^2d^2)/(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{(1/2)} * (a...
\end{aligned}$$

3.351
$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.351.1 Optimal result 3344
 3.351.2 Mathematica [C] (verified) 3344
 3.351.3 Rubi [A] (warning: unable to verify) 3345
 3.351.4 Maple [B] (verified) 3348
 3.351.5 Fricas [B] (verification not implemented) 3349
 3.351.6 Sympy [F] 3350
 3.351.7 Maxima [F(-1)] 3351
 3.351.8 Giac [F(-1)] 3351
 3.351.9 Mupad [B] (verification not implemented) 3351

3.351.1 Optimal result

Integrand size = 33, antiderivative size = 167

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{3/2}d} - \frac{(iA-B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a+ib)^{3/2}d} - \frac{2a^2(Ab-aB)}{b^2(a^2+b^2)d\sqrt{a+b \tan(c+dx)}} + \frac{2B\sqrt{a+b \tan(c+dx)}}{b^2d}$$

```
output (I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(3/2)/d-(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(3/2)/d-2*a^2*(A*b-B*a)/b^2/(a^2+b^2)/d/(a+b*tan(d*x+c))^(1/2)+2*B*(a+b*tan(d*x+c))^(1/2)/b^2/d
```

3.351.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.41 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.49

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{iB \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ib}} \right)}{b\sqrt{a+b \tan(c+dx)}} + \frac{-2Ab+2A^2}{b^2\sqrt{a+b \tan(c+dx)}}$$

3.351.
$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

input `Integrate[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `(I*B*(ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]]/Sqrt[a - I*b] - ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]]/Sqrt[a + I*b]) + (-2*A*b + 4*a*B)/(b*Sqrt[a + b*Tan[c + d*x]]) + ((A*b - a*B)*((-I)*a + b)*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Tan[c + d*x])/(a - I*b)] + (I*a + b)*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Tan[c + d*x])/(a + I*b)])/((a^2 + b^2)*Sqrt[a + b*Tan[c + d*x]]) + (2*B*Tan[c + d*x])/Sqrt[a + b*Tan[c + d*x]]/(b*d)`

3.351.3 Rubi [A] (warning: unable to verify)

Time = 0.94 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.11, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 4087, 25, 3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^2(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{4087} \\
 & \frac{\int -\frac{((a^2+b^2)B \tan^2(c+dx))-b(Ab-aB) \tan(c+dx)+a(Ab-aB)}{\sqrt{a+b \tan(c+dx)}} dx}{b(a^2+b^2)} - \frac{2a^2(Ab-aB)}{b^2d(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int -\frac{((a^2+b^2)B \tan^2(c+dx))-b(Ab-aB) \tan(c+dx)+a(Ab-aB)}{\sqrt{a+b \tan(c+dx)}} dx}{b(a^2+b^2)} - \frac{2a^2(Ab-aB)}{b^2d(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int -\frac{((a^2+b^2)B \tan(c+dx)^2)-b(Ab-aB) \tan(c+dx)+a(Ab-aB)}{\sqrt{a+b \tan(c+dx)}} dx}{b(a^2+b^2)} - \frac{2a^2(Ab-aB)}{b^2d(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow \text{4113}
 \end{aligned}$$

3.351. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{\int \frac{b(aA+bB)-b(Ab-aB)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx - \frac{2B(a^2+b^2)\sqrt{a+b\tan(c+dx)}}{bd}}{b(a^2+b^2)} - \frac{2a^2(Ab-aB)}{b^2d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{b(aA+bB)-b(Ab-aB)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx - \frac{2B(a^2+b^2)\sqrt{a+b\tan(c+dx)}}{bd}}{b(a^2+b^2)} - \frac{2a^2(Ab-aB)}{b^2d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \\
 & \quad \downarrow \text{4022} \\
 & \frac{\frac{2a^2(Ab-aB)}{b^2d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \frac{\frac{1}{2}b(a-ib)(A+iB)\int \frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}b(a+ib)(A-ib)\int \frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}} dx - \frac{2B(a^2+b^2)\sqrt{a+b\tan(c+dx)}}{bd}}{b(a^2+b^2)}}{b(a^2+b^2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\frac{2a^2(Ab-aB)}{b^2d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \frac{\frac{1}{2}b(a-ib)(A+iB)\int \frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}b(a+ib)(A-ib)\int \frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}} dx - \frac{2B(a^2+b^2)\sqrt{a+b\tan(c+dx)}}{bd}}{b(a^2+b^2)}}{b(a^2+b^2)} \\
 & \quad \downarrow \text{4020} \\
 & \frac{\frac{2a^2(Ab-aB)}{b^2d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \frac{ib(a+ib)(A-ib)\int -\frac{1}{(1-i\tan(c+dx))\sqrt{a+b\tan(c+dx)}}d(i\tan(c+dx))}{2d} - \frac{ib(a-ib)(A+iB)\int -\frac{1}{(i\tan(c+dx)+1)\sqrt{a+b\tan(c+dx)}}d(-i\tan(c+dx))}{2d}}{b(a^2+b^2)}}{b(a^2+b^2)} \\
 & \quad \downarrow \text{25} \\
 & \frac{\frac{2a^2(Ab-aB)}{b^2d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \frac{ib(a+ib)(A-ib)\int -\frac{1}{(1-i\tan(c+dx))\sqrt{a+b\tan(c+dx)}}d(i\tan(c+dx))}{2d} + \frac{ib(a-ib)(A+iB)\int \frac{1}{(i\tan(c+dx)+1)\sqrt{a+b\tan(c+dx)}}d(-i\tan(c+dx))}{2d}}{b(a^2+b^2)}}{b(a^2+b^2)} \\
 & \quad \downarrow \text{73} \\
 & \frac{\frac{2a^2(Ab-aB)}{b^2d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \frac{(a-ib)(A+iB)\int -\frac{1}{i\tan^2(c+dx)-\frac{ia}{b}+1}d\sqrt{a+b\tan(c+dx)}}{d} + \frac{(a+ib)(A-ib)\int \frac{1}{i\tan^2(c+dx)+\frac{ia}{b}+1}d\sqrt{a+b\tan(c+dx)}}{d}}{b(a^2+b^2)}}{b(a^2+b^2)} \\
 & \quad \downarrow \text{221}
 \end{aligned}$$

3.351. $\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$

$$\frac{2a^2(Ab - aB)}{b^2 d (a^2 + b^2) \sqrt{a + b \tan(c + dx)}} - \frac{-\frac{2B(a^2 + b^2)\sqrt{a + b \tan(c + dx)}}{bd} + \frac{b(a + ib)(A - iB) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a - ib}}\right)}{d\sqrt{a - ib}} + \frac{b(a - ib)(A + iB) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a + ib}}\right)}{d\sqrt{a + ib}}}{b(a^2 + b^2)}$$

input `Int[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `(-2*a^2*(A*b - a*B))/(b^2*(a^2 + b^2)*d*sqrt[a + b*Tan[c + d*x]]) - (((a + I*b)*b*(A - I*B)*ArcTan[Tan[c + d*x]/sqrt[a - I*b]])/(sqrt[a - I*b]*d) + ((a - I*b)*b*(A + I*B)*ArcTan[Tan[c + d*x]/sqrt[a + I*b]])/(sqrt[a + I*b]*d) - (2*(a^2 + b^2)*B*sqrt[a + b*Tan[c + d*x]])/(b*d)/(b*(a^2 + b^2))`

3.351.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

```
rule 4087 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f
_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[
(- (B*c - A*d))*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(
c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1
)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*
c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2
)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b
*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

```
rule 4113 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

3.351.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3712 vs. $2(145) = 290$.

Time = 0.18 (sec) , antiderivative size = 3713, normalized size of antiderivative = 22.23

method	result	size
parts	Expression too large to display	3713
derivativedivides	Expression too large to display	7982
default	Expression too large to display	7982

```
input int(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,method=_RETURNV
ERBOSE)
```

output

```
A*(-1/4/d/b/(a^2+b^2)^2*ln(b*tan(d*x+c))+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3-1/4/d*b/(a^2+b^2)^2*ln(b*tan(d*x+c))+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+1/4/d/b/(a^2+b^2)^(5/2)*ln(b*tan(d*x+c))+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^4-1/4/d*b^3/(a^2+b^2)^(5/2)*ln(b*tan(d*x+c))+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/d/b/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^3+1/d*b/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a+1/d*b/(a^2+b^2)^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^2-1/d/b/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^5+1/d*b^3/(a^2+b^2)^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))-3/d*b^3/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a-4/d*b/(...
```

3.351.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4379 vs. $2(139) = 278$.

Time = 0.70 (sec) , antiderivative size = 4379, normalized size of antiderivative = 26.22

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="fricas")`

output

```
-1/2*((a^2*b^3 + b^5)*d*tan(d*x + c) + (a^3*b^2 + a*b^4)*d)*sqrt(-(6*A*B*
a^2*b - 2*A*B*b^3 + (A^2 - B^2)*a^3 - 3*(A^2 - B^2)*a*b^2 + (a^6 + 3*a^4*b
^2 + 3*a^2*b^4 + b^6)*d^2*sqrt(-(4*A^2*B^2*a^6 - 12*(A^3*B - A*B^3)*a^5*b
+ 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 + 40*(A^3*B - A*B^3)*a^3*b^3 - 6*
(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 - 12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2*
B^2 + B^4)*b^6)/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8
+ 6*a^2*b^10 + b^12)*d^4)))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2))*lo
g(-(2*(A^3*B + A*B^3)*a^3 - 3*(A^4 - B^4)*a^2*b - 6*(A^3*B + A*B^3)*a*b^2
+ (A^4 - B^4)*b^3)*sqrt(b*tan(d*x + c) + a) + ((A*a^8 + 2*B*a^7*b + 2*A*a^
6*b^2 + 6*B*a^5*b^3 + 6*B*a^3*b^5 - 2*A*a^2*b^6 + 2*B*a*b^7 - A*b^8)*d^3*s
qrt(-(4*A^2*B^2*a^6 - 12*(A^3*B - A*B^3)*a^5*b + 3*(3*A^4 - 14*A^2*B^2 + 3
*B^4)*a^4*b^2 + 40*(A^3*B - A*B^3)*a^3*b^3 - 6*(A^4 - 8*A^2*B^2 + B^4)*a^2
*b^4 - 12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2*B^2 + B^4)*b^6)/((a^12 + 6*
a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 + b^12)*d^4))
+ (2*A*B^2*a^5 - (7*A^2*B - 3*B^3)*a^4*b + 2*(3*A^3 - 7*A*B^2)*a^3*b^2 +
4*(4*A^2*B - B^3)*a^2*b^3 - 2*(A^3 - 4*A*B^2)*a*b^4 - (A^2*B - B^3)*b^5)*d
)*sqrt(-(6*A*B*a^2*b - 2*A*B*b^3 + (A^2 - B^2)*a^3 - 3*(A^2 - B^2)*a*b^2 +
(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*sqrt(-(4*A^2*B^2*a^6 - 12*(A^3*B
- A*B^3)*a^5*b + 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 + 40*(A^3*B - A*B
^3)*a^3*b^3 - 6*(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 - 12*(A^3*B - A*B^3)*a*b...
```

3.351.6 Sympy [F]

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \int \frac{(A+B \tan(c+dx)) \tan^2(c+dx)}{(a+b \tan(c+dx))^{\frac{3}{2}}} dx$$

input `integrate(tan(d*x+c)**2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**2/(a + b*tan(c + d*x))**(3/2), x)`

3.351.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm m="maxima")`

output `Timed out`

3.351.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm m="giac")`

output `Timed out`

3.351.9 Mupad [B] (verification not implemented)

Time = 16.12 (sec) , antiderivative size = 5768, normalized size of antiderivative = 34.54

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `int((tan(c + d*x)^2*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x)`

output

$$\begin{aligned}
& (\log(\left(\left(\left(a + b \tan(c + dx)\right)^{1/2}\right) \cdot \left(16A^2b^{10}d^3 + 32A^2a^2b^8d^3 - 32A^2a^6b^4d^3 - 16A^2a^8b^2d^3\right) + \left(\left(\left(96A^4a^2b^4d^4 - 16A^4\right.\right.\right. \\
& \left.\left.\left.\cdot b^6d^4 - 144A^4a^4b^2d^4\right)^{1/2} - 4A^2a^3d^2 + 12A^2a^2b^2d^2\right) / \left(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4\right)\right)^{1/2} \cdot \left(64A^4a^2b^{11}d^4 - \left(\left(\left(96A^4a^2b^4d^4 - 16A^4\right.\right.\right. \\
& \left.\left.\left.\cdot b^6d^4 - 144A^4a^4b^2d^4\right)^{1/2} - 4A^2a^3d^2 + 12A^2a^2b^2d^2\right) / \left(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4\right)\right)^{1/2} \cdot \left(a + b \tan(c + dx)\right)^{1/2} \cdot \left(64a^4b^{12}d^5 + 320a^3b^{10}d^5 + 640a^5b^8d^5 + 640a^7b^6d^5 + 320a^9b^4d^5 + 64a^{11}b^2d^5\right) / 4 + 256A^4a^3b^9d^4 + 384A^4a^5b^7d^4 + 256A^4a^7b^5d^4 + 64A^4a^9b^3d^4) / 4) \cdot \left(\left(\left(96A^4a^2b^4d^4 - 16A^4\right.\right.\right. \\
& \left.\left.\left.\cdot b^6d^4 - 144A^4a^4b^2d^4\right)^{1/2} - 4A^2a^3d^2 + 12A^2a^2b^2d^2\right) / \left(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4\right)\right)^{1/2} / 4 - 8A^3b^9d^2 - 24A^3a^2b^7d^2 - 24A^3a^4b^5d^2 - 8A^3a^6b^3d^2) \cdot \left(\left(\left(96A^4a^2b^4d^4 - 16A^4\right.\right.\right. \\
& \left.\left.\left.\cdot b^6d^4 - 144A^4a^4b^2d^4\right)^{1/2} - 4A^2a^3d^2 + 12A^2a^2b^2d^2\right) / \left(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4\right)\right)^{1/2} / 4 + \log(\left(\left(\left(a + b \tan(c + dx)\right)^{1/2}\right) \cdot \left(16A^2b^{10}d^3 + 32A^2a^2b^8d^3 - 32A^2a^6b^4d^3 - 16A^2a^8b^2d^3\right) + \left(\left(\left(96A^4a^2b^4d^4 - 16A^4\right.\right.\right. \\
& \left.\left.\left.\cdot b^6d^4 - 144A^4a^4b^2d^4\right)^{1/2} + 4A^2a^3d^2 - 12A^2a^2b^2d^2\right) / \left(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4\right)\right)^{1/2} \cdot \left(64A^4a^2b^{11}d^4 - \left(\left(\left(96A^4a^2b^4d^4 - 16A^4\right.\right.\right. \\
& \left.\left.\left.\cdot b^6d^4 - 144A^4a^4b^2d^4\right)^{1/2} + 4A^2a^3d^2 - 12A^2a^2b^2d^2\right) / \left(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4\right)\right)^{1/2} + 4A^2 \dots
\end{aligned}$$

3.352
$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.352.1 Optimal result 3353
 3.352.2 Mathematica [A] (verified) 3353
 3.352.3 Rubi [A] (warning: unable to verify) 3354
 3.352.4 Maple [B] (verified) 3357
 3.352.5 Fricas [B] (verification not implemented) 3358
 3.352.6 Sympy [F] 3358
 3.352.7 Maxima [F] 3359
 3.352.8 Giac [F(-1)] 3359
 3.352.9 Mupad [B] (verification not implemented) 3359

3.352.1 Optimal result

Integrand size = 31, antiderivative size = 141

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = -\frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{3/2}d} - \frac{(A+iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a+ib)^{3/2}d} + \frac{2a(Ab-aB)}{b(a^2+b^2)d\sqrt{a+b \tan(c+dx)}}$$

output `-(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(3/2)/d-(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(3/2)/d+2*a*(A*B-B*A)/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^(1/2)`

3.352.2 Mathematica [A] (verified)

Time = 1.52 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.62

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{b(A(b^2-a\sqrt{-b^2})-b(a+\sqrt{-b^2})B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-\sqrt{-b^2}}}\right)}{\sqrt{-b^2}\sqrt{a-\sqrt{-b^2}}} - \frac{b(A(b^2+a\sqrt{-b^2})+b(a-\sqrt{-b^2})B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+\sqrt{-b^2}}}\right)}{\sqrt{-b^2}\sqrt{a+\sqrt{-b^2}}} + \frac{2a(Ab-aB)}{b(a^2+b^2)d\sqrt{a+b \tan(c+dx)}}$$

input `Integrate[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output $((b*(A*(b^2 - a*\text{Sqrt}[-b^2]) - b*(a + \text{Sqrt}[-b^2])*B)*\text{ArcTanh}[\text{Sqrt}[a + b*\text{Tan}[c + d*x]]/\text{Sqrt}[a - \text{Sqrt}[-b^2]]]) / (\text{Sqrt}[-b^2]*\text{Sqrt}[a - \text{Sqrt}[-b^2]]) - (b*(A*(b^2 + a*\text{Sqrt}[-b^2]) + b*(-a + \text{Sqrt}[-b^2])*B)*\text{ArcTanh}[\text{Sqrt}[a + b*\text{Tan}[c + d*x]]/\text{Sqrt}[a + \text{Sqrt}[-b^2]]]) / (\text{Sqrt}[-b^2]*\text{Sqrt}[a + \text{Sqrt}[-b^2]]) + (2*a*(A*b - a*B))/\text{Sqrt}[a + b*\text{Tan}[c + d*x]] / (b*(a^2 + b^2)*d)$

3.352.3 Rubi [A] (warning: unable to verify)

Time = 0.64 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.07, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 4074, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{4074} \\ & \frac{\int \frac{Ab-aB+(aA+bB)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx}{a^2+b^2} + \frac{2a(Ab-aB)}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{Ab-aB+(aA+bB)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx}{a^2+b^2} + \frac{2a(Ab-aB)}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \\ & \quad \downarrow \text{4022} \\ & \frac{2a(Ab-aB)}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \\ & \frac{\frac{1}{2}(b+ia)(A+iB) \int \frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx - \frac{1}{2}(-b+ia)(A-iB) \int \frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}} dx}{a^2+b^2} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\begin{aligned}
 & \frac{2a(Ab - aB)}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \\
 & \frac{\frac{1}{2}(b + ia)(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx - \frac{1}{2}(-b + ia)(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} \\
 & \quad \downarrow \text{4020} \\
 & \frac{2a(Ab - aB)}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \\
 & \frac{i(-b + ia)(A - iB) \int -\frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx)) - \frac{i(b + ia)(A + iB) \int -\frac{1}{(i \tan(c + dx) + 1)\sqrt{a + b \tan(c + dx)}} d(-i \tan(c + dx))}{2d}}{a^2 + b^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{2a(Ab - aB)}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \\
 & \frac{i(-b + ia)(A - iB) \int -\frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx)) + \frac{i(b + ia)(A + iB) \int \frac{1}{(i \tan(c + dx) + 1)\sqrt{a + b \tan(c + dx)}} d(-i \tan(c + dx))}{2d}}{a^2 + b^2} \\
 & \quad \downarrow \text{73} \\
 & \frac{2a(Ab - aB)}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \\
 & \frac{(b + ia)(A + iB) \int -\frac{1}{\frac{i \tan^2(c + dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)} - \frac{(-b + ia)(A - iB) \int \frac{1}{\frac{i \tan^2(c + dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)}}{bd}}{a^2 + b^2} \\
 & \quad \downarrow \text{221} \\
 & \frac{2a(Ab - aB)}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{(b + ia)(A + iB) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a + ib}}\right) - (-b + ia)(A - iB) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a - ib}}\right)}{d\sqrt{a + ib} \quad d\sqrt{a - ib}} \quad a^2 + b^2
 \end{aligned}$$

input `Int[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `(-(((I*a - b)*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d)) + ((I*a + b)*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d))/(a^2 + b^2) + (2*a*(A*b - a*B))/(b*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])`

3.352.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`
- rule 4074 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.352.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3687 vs. $2(121) = 242$.

Time = 0.10 (sec) , antiderivative size = 3688, normalized size of antiderivative = 26.16

method	result	size
parts	Expression too large to display	3688
derivativedivides	Expression too large to display	7956
default	Expression too large to display	7956

```
input int(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,method=_RETURNVER
BOSE)
```

```
output A*(1/4/d/(a^2+b^2)^2*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)
^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2+1/4/d
*b^2/(a^2+b^2)^2*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/
2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/2/d/(a^2+b^
2)^(5/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(
1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3-1/2/d*b^2/(a^2+b^
2)^(5/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(
1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-1/d/(a^2+b^2)^(3/2)
/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^
2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^2+1/d/(a^2+b^2)^2/(2
*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(
1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^3-1/d*b^2/(a^2+b^2)^(3/
2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+
b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))+1/d*b^2/(a^2+b^2)^2/
(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)
)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a+2/d*b^2/(a^2+b^2)^(5/
2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+
b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^2+2/d*b^4/(a^2+b^2)
^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*
(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))-1/4/d/(a^2+b...
```

3.352.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4329 vs. $2(115) = 230$.

Time = 0.70 (sec) , antiderivative size = 4329, normalized size of antiderivative = 30.70

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm=
"fricas")
```

```
output -1/2*((a^2*b^2 + b^4)*d*tan(d*x + c) + (a^3*b + a*b^3)*d)*sqrt((6*A*B*a^2
*b - 2*A*B*b^3 + (A^2 - B^2)*a^3 - 3*(A^2 - B^2)*a*b^2 + (a^6 + 3*a^4*b^2
+ 3*a^2*b^4 + b^6)*d^2*sqrt(-(4*A^2*B^2*a^6 - 12*(A^3*B - A*B^3)*a^5*b + 3
*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 + 40*(A^3*B - A*B^3)*a^3*b^3 - 6*(A^
4 - 8*A^2*B^2 + B^4)*a^2*b^4 - 12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2*B^2
+ B^4)*b^6))/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 +
6*a^2*b^10 + b^12)*d^4))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2))*log(-
(2*(A^3*B + A*B^3)*a^3 - 3*(A^4 - B^4)*a^2*b - 6*(A^3*B + A*B^3)*a*b^2 + (
A^4 - B^4)*b^3)*sqrt(b*tan(d*x + c) + a) + ((B*a^8 - 2*A*a^7*b + 2*B*a^6*b
^2 - 6*A*a^5*b^3 - 6*A*a^3*b^5 - 2*B*a^2*b^6 - 2*A*a*b^7 - B*b^8)*d^3*sqrt
(-(4*A^2*B^2*a^6 - 12*(A^3*B - A*B^3)*a^5*b + 3*(3*A^4 - 14*A^2*B^2 + 3*B^
4)*a^4*b^2 + 40*(A^3*B - A*B^3)*a^3*b^3 - 6*(A^4 - 8*A^2*B^2 + B^4)*a^2*b^
4 - 12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2*B^2 + B^4)*b^6))/((a^12 + 6*a^1
0*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 + b^12)*d^4)) -
(2*A^2*B*a^5 - (3*A^3 - 7*A*B^2)*a^4*b - 2*(7*A^2*B - 3*B^3)*a^3*b^2 + 4*(
A^3 - 4*A*B^2)*a^2*b^3 + 2*(4*A^2*B - B^3)*a*b^4 - (A^3 - A*B^2)*b^5)*d)*s
qrt((6*A*B*a^2*b - 2*A*B*b^3 + (A^2 - B^2)*a^3 - 3*(A^2 - B^2)*a*b^2 + (a^
6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*sqrt(-(4*A^2*B^2*a^6 - 12*(A^3*B - A*
B^3)*a^5*b + 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 + 40*(A^3*B - A*B^3)*a
^3*b^3 - 6*(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 - 12*(A^3*B - A*B^3)*a*b^5 + ...
```

3.352.6 Sympy [F]

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(A+B\tan(c+dx))\tan(c+dx)}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx$$

```
input integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)
```

3.352. $\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)/(a + b*tan(c + d*x))**(3/2), x)`

3.352.7 Maxima [F]

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)}{(b \tan(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)/(b*tan(d*x + c) + a)^(3/2), x)`

3.352.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `Timed out`

3.352.9 Mupad [B] (verification not implemented)

Time = 15.34 (sec) , antiderivative size = 5742, normalized size of antiderivative = 40.72

$$\int \frac{\tan(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `int((tan(c + d*x)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x)`

output

$$\begin{aligned}
& (\log(- (((((96*A^4*a^2*b^4*d^4 - 16*A^4*b^6*d^4 - 144*A^4*a^4*b^2*d^4)^(1/2) + 4*A^2*a^3*d^2 - 12*A^2*a*b^2*d^2)/(a^6*d^4 + b^6*d^4 + 3*a^2*b^4*d^4 + 3*a^4*b^2*d^4))^(1/2)*(32*A*b^12*d^4 + (((96*A^4*a^2*b^4*d^4 - 16*A^4*b^6*d^4 - 144*A^4*a^4*b^2*d^4)^(1/2) + 4*A^2*a^3*d^2 - 12*A^2*a*b^2*d^2)/(a^6*d^4 + b^6*d^4 + 3*a^2*b^4*d^4 + 3*a^4*b^2*d^4))^(1/2)*(a + b*\tan(c + d*x))^(1/2)*(64*a*b^12*d^5 + 320*a^3*b^10*d^5 + 640*a^5*b^8*d^5 + 640*a^7*b^6*d^5 + 320*a^9*b^4*d^5 + 64*a^11*b^2*d^5))/4 + 96*A*a^2*b^10*d^4 + 64*A*a^4*b^8*d^4 - 64*A*a^6*b^6*d^4 - 96*A*a^8*b^4*d^4 - 32*A*a^10*b^2*d^4))/4 + (a + b*\tan(c + d*x))^(1/2)*(16*A^2*b^10*d^3 + 32*A^2*a^2*b^8*d^3 - 32*A^2*a^6*b^4*d^3 - 16*A^2*a^8*b^2*d^3))*(((96*A^4*a^2*b^4*d^4 - 16*A^4*b^6*d^4 - 144*A^4*a^4*b^2*d^4)^(1/2) + 4*A^2*a^3*d^2 - 12*A^2*a*b^2*d^2)/(a^6*d^4 + b^6*d^4 + 3*a^2*b^4*d^4 + 3*a^4*b^2*d^4))^(1/2))/4 - 24*A^3*a^3*b^6*d^2 - 24*A^3*a^5*b^4*d^2 - 8*A^3*a^7*b^2*d^2 - 8*A^3*a*b^8*d^2)*(((96*A^4*a^2*b^4*d^4 - 16*A^4*b^6*d^4 - 144*A^4*a^4*b^2*d^4)^(1/2) + 4*A^2*a^3*d^2 - 12*A^2*a*b^2*d^2)/(a^6*d^4 + b^6*d^4 + 3*a^2*b^4*d^4 + 3*a^4*b^2*d^4))^(1/2))/4 + (\log(- ((((-((96*A^4*a^2*b^4*d^4 - 16*A^4*b^6*d^4 - 144*A^4*a^4*b^2*d^4)^(1/2) - 4*A^2*a^3*d^2 + 12*A^2*a*b^2*d^2)/(a^6*d^4 + b^6*d^4 + 3*a^2*b^4*d^4 + 3*a^4*b^2*d^4))^(1/2)*(32*A*b^12*d^4 + (((96*A^4*a^2*b^4*d^4 - 16*A^4*b^6*d^4 - 144*A^4*a^4*b^2*d^4)^(1/2) - 4*A^2*a^3*d^2 + 12*A^2*a*b^2*d^2)/(a^6*d^4 + b^6*d^4 + 3*a^2*b^4*d^4 + 3*a^4*b^2*d^4))^(1/2)*(a ...
\end{aligned}$$

3.353 $\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx$

3.353.1 Optimal result	3361
3.353.2 Mathematica [C] (verified)	3361
3.353.3 Rubi [A] (warning: unable to verify)	3362
3.353.4 Maple [B] (verified)	3364
3.353.5 Fricas [B] (verification not implemented)	3365
3.353.6 Sympy [F]	3366
3.353.7 Maxima [F(-2)]	3367
3.353.8 Giac [F(-1)]	3367
3.353.9 Mupad [B] (verification not implemented)	3367

3.353.1 Optimal result

Integrand size = 25, antiderivative size = 138

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = -\frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a - ib)^{3/2}d} + \frac{(iA - B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a + ib)^{3/2}d} - \frac{2(Ab - aB)}{(a^2 + b^2) d \sqrt{a + b \tan(c + dx)}}$$

output `-(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(3/2)/d+(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(3/2)/d-2*(A*b-B*a)/(a^2+b^2)/d/(a+b*tan(d*x+c))^(1/2)`

3.353.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.23 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.82

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \frac{i \left(\frac{(A-iB) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b \tan(c+dx)}{a-ib}\right)}{a-ib} - \frac{(A+iB) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b \tan(c+dx)}{a+ib}\right)}{a+ib} \right)}{d \sqrt{a + b \tan(c + dx)}}$$

input `Integrate[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^(3/2),x]`

output $(I*((A - I*B)*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*\text{Tan}[c + d*x])/(a - I*b)])/(a - I*b) - ((A + I*B)*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*\text{Tan}[c + d*x])/(a + I*b)])/(a + I*b))/(d*\text{Sqrt}[a + b*\text{Tan}[c + d*x]])$

3.353.3 Rubi [A] (warning: unable to verify)

Time = 0.61 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.04, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$, Rules used = {3042, 4012, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx \\ & \quad \downarrow 3042 \\ & \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx \\ & \quad \downarrow 4012 \\ & \frac{\int \frac{aA + bB - (Ab - aB) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} - \frac{2(Ab - aB)}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} \\ & \quad \downarrow 3042 \\ & \frac{\int \frac{aA + bB - (Ab - aB) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} - \frac{2(Ab - aB)}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} \\ & \quad \downarrow 4022 \\ & -\frac{2(Ab - aB)}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} + \\ & \frac{\frac{1}{2}(a - ib)(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a + ib)(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} \\ & \quad \downarrow 3042 \\ & -\frac{2(Ab - aB)}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} + \\ & \frac{\frac{1}{2}(a - ib)(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a + ib)(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} \\ & \quad \downarrow 4020 \end{aligned}$$

$$\begin{aligned}
 & \frac{-\frac{2(Ab - aB)}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{i(a+ib)(A-iB) \int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} - \frac{i(a-ib)(A+iB) \int -\frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d}}{a^2 + b^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{-\frac{2(Ab - aB)}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{i(a-ib)(A+iB) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} - \frac{i(a+ib)(A-iB) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d}}{a^2 + b^2} \\
 & \quad \downarrow \text{73} \\
 & \frac{-\frac{2(Ab - aB)}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{(a-ib)(A+iB) \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \frac{(a+ib)(A-iB) \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd}}{a^2 + b^2} \\
 & \quad \downarrow \text{221} \\
 & -\frac{2(Ab - aB)}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{(a+ib)(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} + \frac{(a-ib)(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}}
 \end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^(3/2),x]`

output `((a + I*b)*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]]/(Sqrt[a - I*b]*d) + ((a - I*b)*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]]/(Sqrt[a + I*b]*d))/(a^2 + b^2) - (2*(A*b - a*B))/((a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])`

3.353.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

3.353. $\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx$

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4020 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

3.353.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3683 vs. $2(118) = 236$.

Time = 0.09 (sec) , antiderivative size = 3684, normalized size of antiderivative = 26.70

method	result	size
parts	Expression too large to display	3684
derivativedivides	Expression too large to display	7951
default	Expression too large to display	7951

input `int((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

$$3.353. \quad \int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx$$

output

```
A*(1/4/d/b/(a^2+b^2)^2*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3+1/4/d*b/(a^2+b^2)^2*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^-1/4/d/b/(a^2+b^2)^(5/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^4+1/4/d*b^3/(a^2+b^2)^(5/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/d/b/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^3-1/d*b/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^-1/d*b/(a^2+b^2)^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^2+1/d/b/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^5-1/d*b^3/(a^2+b^2)^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))+3/d*b^3/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a+4/d*b/(a...
```

3.353.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4318 vs. $2(113) = 226$.

Time = 0.69 (sec) , antiderivative size = 4318, normalized size of antiderivative = 31.29

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="fracas")`

```
output 1/2*((a^2*b + b^3)*d*tan(d*x + c) + (a^3 + a*b^2)*d)*sqrt(-(6*A*B*a^2*b -
2*A*B*b^3 + (A^2 - B^2)*a^3 - 3*(A^2 - B^2)*a*b^2 + (a^6 + 3*a^4*b^2 + 3*
a^2*b^4 + b^6)*d^2*sqrt(-(4*A^2*B^2*a^6 - 12*(A^3*B - A*B^3)*a^5*b + 3*(3*
A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 + 40*(A^3*B - A*B^3)*a^3*b^3 - 6*(A^4 -
8*A^2*B^2 + B^4)*a^2*b^4 - 12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2*B^2 + B
^4)*b^6))/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^
2*b^10 + b^12)*d^4)))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2))*log(-(2*(
A^3*B + A*B^3)*a^3 - 3*(A^4 - B^4)*a^2*b - 6*(A^3*B + A*B^3)*a*b^2 + (A^4
- B^4)*b^3)*sqrt(b*tan(d*x + c) + a) + ((A*a^8 + 2*B*a^7*b + 2*A*a^6*b^2 +
6*B*a^5*b^3 + 6*B*a^3*b^5 - 2*A*a^2*b^6 + 2*B*a*b^7 - A*b^8)*d^3*sqrt(-(4
*A^2*B^2*a^6 - 12*(A^3*B - A*B^3)*a^5*b + 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a
^4*b^2 + 40*(A^3*B - A*B^3)*a^3*b^3 - 6*(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 -
12*(A^3*B - A*B^3)*a*b^5 + (A^4 - 2*A^2*B^2 + B^4)*b^6))/((a^12 + 6*a^10*b^
2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 + b^12)*d^4)) + (2*A
*B^2*a^5 - (7*A^2*B - 3*B^3)*a^4*b + 2*(3*A^3 - 7*A*B^2)*a^3*b^2 + 4*(4*A^
2*B - B^3)*a^2*b^3 - 2*(A^3 - 4*A*B^2)*a*b^4 - (A^2*B - B^3)*b^5)*d)*sqrt(
-(6*A*B*a^2*b - 2*A*B*b^3 + (A^2 - B^2)*a^3 - 3*(A^2 - B^2)*a*b^2 + (a^6 +
3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*sqrt(-(4*A^2*B^2*a^6 - 12*(A^3*B - A*B^3
)*a^5*b + 3*(3*A^4 - 14*A^2*B^2 + 3*B^4)*a^4*b^2 + 40*(A^3*B - A*B^3)*a^3*
b^3 - 6*(A^4 - 8*A^2*B^2 + B^4)*a^2*b^4 - 12*(A^3*B - A*B^3)*a*b^5 + (A...
```

3.353.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{\frac{3}{2}}} dx$$

```
input integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)
```

```
output Integral((A + B*tan(c + d*x))/(a + b*tan(c + d*x))**(3/2), x)
```

3.353.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is`

3.353.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `Timed out`

3.353.9 Mupad [B] (verification not implemented)

Time = 14.92 (sec) , antiderivative size = 5737, normalized size of antiderivative = 41.57

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `int((A + B*tan(c + d*x))/(a + b*tan(c + d*x))^(3/2),x)`

output $(\log(\left(\frac{(a + b \tan(c + dx))^{1/2} (16A^2b^{10}d^3 + 32A^2a^2b^8d^3 - 32A^2a^6b^4d^3 - 16A^2a^8b^2d^3) - (((96A^4a^2b^4d^4 - 16A^4b^6d^4 - 144A^4a^4b^2d^4)^{1/2} - 4A^2a^3d^2 + 12A^2ab^2d^2)}{(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{1/2} * (((96A^4a^2b^4d^4 - 16A^4b^6d^4 - 144A^4a^4b^2d^4)^{1/2} - 4A^2a^3d^2 + 12A^2ab^2d^2)}{(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{1/2} * (a + b \tan(c + dx))^{1/2} (64a^5b^8d^5 + 640a^7b^6d^5 + 320a^9b^4d^5 + 64a^{11}b^2d^5)}{4} + 64A^2a^9b^3d^4)/4 * (((96A^4a^2b^4d^4 - 16A^4b^6d^4 - 144A^4a^4b^2d^4)^{1/2} - 4A^2a^3d^2 + 12A^2ab^2d^2)}{(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{1/2}}/4 + 8A^3b^9d^2 + 24A^3a^2b^7d^2 + 24A^3a^4b^5d^2 + 8A^3a^6b^3d^2) * (((96A^4a^2b^4d^4 - 16A^4b^6d^4 - 144A^4a^4b^2d^4)^{1/2} - 4A^2a^3d^2 + 12A^2ab^2d^2)}{(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{1/2}}/4 + (\log(\left(\frac{(a + b \tan(c + dx))^{1/2} (16A^2b^{10}d^3 + 32A^2a^2b^8d^3 - 32A^2a^6b^4d^3 - 16A^2a^8b^2d^3) - ((-((96A^4a^2b^4d^4 - 16A^4b^6d^4 - 144A^4a^4b^2d^4)^{1/2} + 4A^2a^3d^2 - 12A^2ab^2d^2)}{(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{1/2} * (((-((96A^4a^2b^4d^4 - 16A^4b^6d^4 - 144A^4a^4b^2d^4)^{1/2} + 4A^2a^3d^2 - 12A^2a...$

3.354 $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

3.354.1 Optimal result 3369
 3.354.2 Mathematica [A] (verified) 3369
 3.354.3 Rubi [A] (warning: unable to verify) 3370
 3.354.4 Maple [B] (verified) 3374
 3.354.5 Fricas [B] (verification not implemented) 3375
 3.354.6 Sympy [F] 3375
 3.354.7 Maxima [F] 3375
 3.354.8 Giac [F(-1)] 3376
 3.354.9 Mupad [B] (verification not implemented) 3376

3.354.1 Optimal result

Integrand size = 31, antiderivative size = 171

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx =$$

$$-\frac{2A \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{a^{3/2}d} + \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{3/2}d}$$

$$+ \frac{(A+iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a+ib)^{3/2}d} + \frac{2b(Ab-aB)}{a(a^2+b^2)d\sqrt{a+b \tan(c+dx)}}$$

```
output -2*A*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/a^(3/2)/d+(A-I*B)*arctanh((a+
b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(3/2)/d+(A+I*B)*arctanh((a+b*ta
n(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(3/2)/d+2*b*(A*b-B*a)/a/(a^2+b^2)/d
/(a+b*tan(d*x+c))^(1/2)
```

3.354.2 Mathematica [A] (verified)

Time = 1.34 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.09

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = -\frac{2A(a^2+b^2) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{a}} + \frac{a(a+ib)(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib} a(a^2+b^2)}$$

input `Integrate[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `((-2*A*(a^2 + b^2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]]/Sqrt[a] + (a*(a + I*b)*(A - I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]]/Sqrt[a - I*b] + (a*(a - I*b)*(A + I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/Sqrt[a + I*b] + (2*b*(A*b - a*B))/Sqrt[a + b*Tan[c + d*x]])/(a*(a^2 + b^2)*d)`

3.354.3 Rubi [A] (warning: unable to verify)

Time = 1.36 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.14, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.516$, Rules used = {3042, 4092, 27, 3042, 4136, 25, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)(a+b \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow 4092 \\
 & \frac{2 \int \frac{\cot(c+dx)(b(Ab-aB) \tan^2(c+dx)-a(Ab-aB) \tan(c+dx)+A(a^2+b^2))}{2\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(Ab-aB)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{\cot(c+dx)(b(Ab-aB) \tan^2(c+dx)-a(Ab-aB) \tan(c+dx)+A(a^2+b^2))}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(Ab-aB)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{b(Ab-aB) \tan(c+dx)^2-a(Ab-aB) \tan(c+dx)+A(a^2+b^2)}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(Ab-aB)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 4136
 \end{aligned}$$

3.354. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \frac{A(a^2 + b^2) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx + \int -\frac{a(Ab-aB)+a(aA+bB) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{\frac{a(a^2 + b^2)}{2b(Ab - aB)}} + \\
& \frac{ad(a^2 + b^2) \sqrt{a + b \tan(c + dx)}}{\downarrow 25} \\
& \frac{A(a^2 + b^2) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx - \int \frac{a(Ab-aB)+a(aA+bB) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{\frac{a(a^2 + b^2)}{2b(Ab - aB)}} + \\
& \frac{ad(a^2 + b^2) \sqrt{a + b \tan(c + dx)}}{\downarrow 3042} \\
& \frac{A(a^2 + b^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - \int \frac{a(Ab-aB)+a(aA+bB) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{\frac{a(a^2 + b^2)}{2b(Ab - aB)}} + \\
& \frac{ad(a^2 + b^2) \sqrt{a + b \tan(c + dx)}}{\downarrow 4022} \\
& \frac{2b(Ab - aB)}{ad(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} + \\
& \frac{A(a^2 + b^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2}a(b + ia)(A + iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}a(-b + ia)(A - iB) \int \frac{i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2 + b^2)}}{\downarrow 3042} \\
& \frac{2b(Ab - aB)}{ad(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} + \\
& \frac{A(a^2 + b^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2}a(b + ia)(A + iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}a(-b + ia)(A - iB) \int \frac{i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2 + b^2)}}{\downarrow 4020} \\
& \frac{2b(Ab - aB)}{ad(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} + \\
& \frac{A(a^2 + b^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + \frac{ia(-b+ia)(A-iB) \int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \frac{ia(b+ia)(A+iB) \int -\frac{1}{(1+i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d}}{a(a^2 + b^2)}}{\downarrow 25}
\end{aligned}$$

3.354. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

$$\frac{A(a^2 + b^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - \frac{2b(Ab - aB)}{ad(a^2 + b^2)\sqrt{a+b\tan(c+dx)}} + \frac{ia(-b+ia)(A-iB) \int \frac{1}{(1-i\tan(c+dx))\sqrt{a+b\tan(c+dx)}} d(i\tan(c+dx))}{2d} - \frac{ia(b+ia)(A+iB) \int \frac{1}{(1+i\tan(c+dx))\sqrt{a+b\tan(c+dx)}} d(i\tan(c+dx))}{2d}}{a(a^2 + b^2)}$$

↓ 73

$$\frac{A(a^2 + b^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - \frac{2b(Ab - aB)}{ad(a^2 + b^2)\sqrt{a+b\tan(c+dx)}} + \frac{a(b+ia)(A+iB) \int \frac{1}{-\frac{i\tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a+b\tan(c+dx)}}{bd} - \frac{a(-b+ia)(A-iB) \int \frac{1}{\frac{i\tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a+b\tan(c+dx)}}{bd}}{a(a^2 + b^2)}$$

↓ 221

$$\frac{A(a^2 + b^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx + \frac{2b(Ab - aB)}{ad(a^2 + b^2)\sqrt{a+b\tan(c+dx)}} + \frac{a(-b+ia)(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{a(b+ia)(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}}}{a(a^2 + b^2)}$$

↓ 4117

$$\frac{A(a^2+b^2) \int \frac{\cot(c+dx)}{\sqrt{a+b\tan(c+dx)}} d\tan(c+dx)}{d} + \frac{2b(Ab - aB)}{ad(a^2 + b^2)\sqrt{a+b\tan(c+dx)}} + \frac{a(-b+ia)(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{a(b+ia)(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}}}{a(a^2 + b^2)}$$

↓ 73

$$\frac{2A(a^2+b^2) \int \frac{1}{\frac{a+b\tan(c+dx)}{b} - \frac{a}{b}} d\sqrt{a+b\tan(c+dx)}}{bd} + \frac{2b(Ab - aB)}{ad(a^2 + b^2)\sqrt{a+b\tan(c+dx)}} + \frac{a(-b+ia)(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{a(b+ia)(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}}}{a(a^2 + b^2)}$$

↓ 221

$$\frac{-\frac{2A(a^2+b^2) \operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ad}} + \frac{2b(Ab - aB)}{ad(a^2 + b^2)\sqrt{a+b\tan(c+dx)}} + \frac{a(-b+ia)(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{a(b+ia)(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}}}{a(a^2 + b^2)}$$

input `Int[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

```
output ((a*(I*a - b)*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]]/(Sqrt[a - I*b]
*d) - (a*(I*a + b)*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]]/(Sqrt[a +
I*b]*d) - (2*A*(a^2 + b^2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]]/(Sqr
rt[a]*d))/(a*(a^2 + b^2)) + (2*b*(A*b - a*B))/(a*(a^2 + b^2)*d*Sqrt[a + b*
Tan[c + d*x]])
```

3.354.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

3.354.
$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

```
rule 4092 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)
/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^
2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*
B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2
)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n
+ 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1]
&& (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m]
|| (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

```
rule 4136 Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.354.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 7981 vs. $2(145) = 290$.

Time = 0.23 (sec) , antiderivative size = 7982, normalized size of antiderivative = 46.68

method	result	size
derivativedivides	Expression too large to display	7982
default	Expression too large to display	7982

```
input int(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,method=_RETURNVER
BOSE)
```

$$3.354. \quad \int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

output result too large to display

3.354.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4402 vs. 2(139) = 278.

Time = 4.11 (sec) , antiderivative size = 8820, normalized size of antiderivative = 51.58

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="fricas")`

output Too large to include

3.354.6 Sympy [F]

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(A+B\tan(c+dx))\cot(c+dx)}{(a+b\tan(c+dx))^{3/2}} dx$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)/(a + b*tan(c + d*x))**(3/2), x)`

3.354.7 Maxima [F]

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)}{(b\tan(dx+c)+a)^{3/2}} dx$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)/(b*tan(d*x + c) + a)^(3/2), x)`

3.354. $\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$

3.354.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="giac")`

output Timed out

3.354.9 Mupad [B] (verification not implemented)

Time = 15.35 (sec) , antiderivative size = 26139, normalized size of antiderivative = 152.86

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

input `int((cot(c + d*x)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x)`

output `atan(-(((8*A^2*a^3*d^2 - 8*B^2*a^3*d^2 - 16*A*B*b^3*d^2 - 24*A^2*a*b^2*d^2 + 24*B^2*a*b^2*d^2 + 48*A*B*a^2*b*d^2)^2/4 - (A^4 + 2*A^2*B^2 + B^4)*(16*a^6*d^4 + 16*b^6*d^4 + 48*a^2*b^4*d^4 + 48*a^4*b^2*d^4))^(1/2) - 4*A^2*a^3*d^2 + 4*B^2*a^3*d^2 + 8*A*B*b^3*d^2 + 12*A^2*a*b^2*d^2 - 12*B^2*a*b^2*d^2 - 24*A*B*a^2*b*d^2)/(16*(a^6*d^4 + b^6*d^4 + 3*a^2*b^4*d^4 + 3*a^4*b^2*d^4))^(1/2)*((a + b*tan(c + d*x))^(1/2)*(256*A^2*a^8*b^26*d^7 + 1472*A^2*a^10*b^24*d^7 + 3712*A^2*a^12*b^22*d^7 + 6272*A^2*a^14*b^20*d^7 + 9856*A^2*a^16*b^18*d^7 + 14336*A^2*a^18*b^16*d^7 + 15232*A^2*a^20*b^14*d^7 + 10112*A^2*a^22*b^12*d^7 + 3712*A^2*a^24*b^10*d^7 + 576*A^2*a^26*b^8*d^7 + 832*B^2*a^10*b^24*d^7 + 5504*B^2*a^12*b^22*d^7 + 15232*B^2*a^14*b^20*d^7 + 22400*B^2*a^16*b^18*d^7 + 17920*B^2*a^18*b^16*d^7 + 6272*B^2*a^20*b^14*d^7 - 896*B^2*a^22*b^12*d^7 - 1408*B^2*a^24*b^10*d^7 - 320*B^2*a^26*b^8*d^7 - 512*A*B*a^9*b^25*d^7 - 1792*A*B*a^11*b^23*d^7 + 1792*A*B*a^13*b^21*d^7 + 19712*A*B*a^15*b^19*d^7 + 44800*A*B*a^17*b^17*d^7 + 51968*A*B*a^19*b^15*d^7 + 34048*A*B*a^21*b^13*d^7 + 12032*A*B*a^23*b^11*d^7 + 1792*A*B*a^25*b^9*d^7) + (-(((8*A^2*a^3*d^2 - 8*B^2*a^3*d^2 - 16*A*B*b^3*d^2 - 24*A^2*a*b^2*d^2 + 24*B^2*a*b^2*d^2 + 48*A*B*a^2*b*d^2)^2/4 - (A^4 + 2*A^2*B^2 + B^4)*(16*a^6*d^4 + 16*b^6*d^4 + 48*a^2*b^4*d^4 + 48*a^4*b^2*d^4))^(1/2) - 4*A^2*a^3*d^2 + 4*B^2*a^3*d^2 + 8*A*B*b^3*d^2 + 12*A^2*a*b^2*d^2 - 12*B^2*a*b^2*d^2 - 24*A*B*a^2*b*d^2)/(16*(a^6*d^4 + b^6*d^4 + 3*a^2*b^4*d^4 + 3*a^4*b...`

3.355 $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

3.355.1 Optimal result 3377
 3.355.2 Mathematica [A] (verified) 3378
 3.355.3 Rubi [A] (warning: unable to verify) 3378
 3.355.4 Maple [B] (verified) 3384
 3.355.5 Fracas [B] (verification not implemented) 3384
 3.355.6 Sympy [F] 3385
 3.355.7 Maxima [F] 3385
 3.355.8 Giac [F(-1)] 3385
 3.355.9 Mupad [B] (verification not implemented) 3386

3.355.1 Optimal result

Integrand size = 33, antiderivative size = 219

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{(3Ab - 2aB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{a^{5/2}d} + \frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{3/2}d} - \frac{(iA - B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a+ib)^{3/2}d} - \frac{b(a^2A + 3Ab^2 - 2abB)}{a^2(a^2 + b^2)d\sqrt{a+b \tan(c+dx)}} - \frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}}$$

```
output (3*A*b-2*B*a)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/a^(5/2)/d+(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(3/2)/d-(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(3/2)/d-b*(A*a^2+3*A*b^2-2*B*a*b)/a^2/(a^2+b^2)/d/(a+b*tan(d*x+c))^(1/2)-A*cot(d*x+c)/a/d/(a+b*tan(d*x+c))^(1/2)
```


3.355.2 Mathematica [A] (verified)

Time = 4.17 (sec) , antiderivative size = 208, normalized size of antiderivative = 0.95

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{(3Ab-2aB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{a}} + a^2 \left(\frac{(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{3/2}} \right)$$

input `Integrate[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `((3*A*b - 2*a*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]]/Sqrt[a] + a^2 *(((I*A + B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/(a - I*b)^(3/2) + (((-I)*A + B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/(a + I*b)^(3/2)) - (b*(a^2*A + 3*A*b^2 - 2*a*b*B))/((a^2 + b^2)*Sqrt[a + b*Tan[c + d*x]]) - (a*A*Cot[c + d*x])/Sqrt[a + b*Tan[c + d*x]]/(a^2*d)`

3.355.3 Rubi [A] (warning: unable to verify)

Time = 1.82 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.16, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^2(a+b \tan(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{4092} \\ & -\frac{\int \frac{\cot(c+dx)(3Ab \tan^2(c+dx)+2aA \tan(c+dx)+3Ab-2aB)}{2(a+b \tan(c+dx))^{3/2}} dx}{a} - \frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} \\ & \quad \downarrow \text{27} \end{aligned}$$

3.355. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{\int \frac{\cot(c+dx)(3Ab \tan^2(c+dx)+2aA \tan(c+dx)+3Ab-2aB)}{(a+b \tan(c+dx))^{3/2}} dx}{2a} - \frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{3Ab \tan(c+dx)^2+2aA \tan(c+dx)+3Ab-2aB}{\tan(c+dx)(a+b \tan(c+dx))^{3/2}} dx}{2a} - \frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow \text{4132} \\
 & \frac{2 \int \frac{\cot(c+dx)(2(aA+bB) \tan(c+dx)a^2+b(Aa^2-2bBa+3Ab^2) \tan^2(c+dx)+(a^2+b^2)(3Ab-2aB))}{2\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \frac{2a}{ad\sqrt{a+b \tan(c+dx)}} \frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\cot(c+dx)(2(aA+bB) \tan(c+dx)a^2+b(Aa^2-2bBa+3Ab^2) \tan^2(c+dx)+(a^2+b^2)(3Ab-2aB))}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \frac{2a}{ad\sqrt{a+b \tan(c+dx)}} \frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{2(aA+bB) \tan(c+dx)a^2+b(Aa^2-2bBa+3Ab^2) \tan(c+dx)^2+(a^2+b^2)(3Ab-2aB)}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \frac{2a}{ad\sqrt{a+b \tan(c+dx)}} \frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow \text{4136} \\
 & \frac{(a^2+b^2)(3Ab-2aB) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx + \int \frac{2(a^2(aA+bB)-a^2(Ab-aB) \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \frac{2a}{ad\sqrt{a+b \tan(c+dx)}} \frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.355. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

$$\frac{(a^2+b^2)(3Ab-2aB) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx + 2 \int \frac{a^2(aA+bB)-a^2(Ab-aB) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}$$

$$\frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}}$$

↓ 3042

$$\frac{(a^2+b^2)(3Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 2 \int \frac{a^2(aA+bB)-a^2(Ab-aB) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}$$

$$\frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}}$$

↓ 4022

$$\frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}}$$

$$\frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(3Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 2 \left(\frac{1}{2} a^2(a-ib)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2} a^2(a+ib)(A-iB) \int \frac{1+i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx \right)}{a(a^2+b^2)}$$

2a

↓ 3042

$$\frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}}$$

$$\frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(3Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 2 \left(\frac{1}{2} a^2(a-ib)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2} a^2(a+ib)(A-iB) \int \frac{1+i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx \right)}{a(a^2+b^2)}$$

2a

↓ 4020

$$\frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}}$$

$$\frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(3Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 2 \left(\frac{ia^2(a+ib)(A-iB) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2d} d(i \tan(c+dx)) \right)}{a(a^2+b^2)}$$

2a

↓ 25

$$\frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}}$$

$$\frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(3Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 2 \left(\frac{ia^2(a-ib)(A+iB) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2d} d(-i \tan(c+dx)) \right)}{a(a^2+b^2)}$$

2a

3.355. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

$$\begin{aligned} & \downarrow 73 \\ & -\frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} - \\ & \frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(3Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 2 \left(\frac{a^2(a-ib)(A+iB) \int \frac{1}{-i \tan^2(c+dx) - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} \right)}{a(a^2+b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 221 \\ & -\frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} - \\ & \frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(3Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 2 \left(\frac{a^2(a+ib)(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} + \frac{a^2(a-ib)(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} \right)}{a(a^2+b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 4117 \\ & -\frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} - \\ & \frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(3Ab-2aB) \int \frac{\cot(c+dx)}{\sqrt{a+b \tan(c+dx)}} d \tan(c+dx) + 2 \left(\frac{a^2(a+ib)(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} + \frac{a^2(a-ib)(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} \right)}{a(a^2+b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 73 \\ & -\frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} - \\ & \frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{2(a^2+b^2)(3Ab-2aB) \int \frac{1}{\frac{a+b \tan(c+dx)}{b} - \frac{a}{b}} d\sqrt{a+b \tan(c+dx)}}{a(a^2+b^2)} + 2 \left(\frac{a^2(a+ib)(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} + \frac{a^2(a-ib)(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 221 \\ & -\frac{A \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} - \\ & \frac{2b(a^2A-2abB+3Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{2(a^2+b^2)(3Ab-2aB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ad}} + 2 \left(\frac{a^2(a+ib)(A-iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} + \frac{a^2(a-ib)(A+iB) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} \right) \end{aligned}$$

input `Int[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2), x]`

```
output -((A*Cot[c + d*x])/(a*d*Sqrt[a + b*Tan[c + d*x]])) - ((2*((a^2*(a + I*b)*(
A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d) + (a^2*(a -
I*b)*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d)) - (
2*(a^2 + b^2)*(3*A*b - 2*a*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/(
Sqrt[a]*d))/(a*(a^2 + b^2)) + (2*b*(a^2*A + 3*A*b^2 - 2*a*b*B))/(a*(a^2 +
b^2)*d*Sqrt[a + b*Tan[c + d*x]]))/(2*a)
```

3.355.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

rule 4022 $\text{Int}[(a + b \tan(e + f x))^m ((c + d \tan(e + f x) + (f x)))]$, x_Symbol] \rightarrow $\text{Simp}[(c + I d)/2 \text{Int}[(a + b \tan(e + f x))^m (1 - I \tan(e + f x)), x], x] + \text{Simp}[(c - I d)/2 \text{Int}[(a + b \tan(e + f x))^m (1 + I \tan(e + f x)), x], x] /;$ FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]

rule 4092 $\text{Int}[(a + b \tan(e + f x))^m ((A + B \tan(e + f x) + (f x)))]$, x_Symbol] \rightarrow $\text{Simp}[b*(A*b - a*B)*(a + b \tan(e + f x))^{m+1} ((c + d \tan(e + f x))^{n+1} / (f*(m+1)*(b*c - a*d)*(a^2 + b^2))), x] + \text{Simp}[1/((m+1)*(b*c - a*d)*(a^2 + b^2)) \text{Int}[(a + b \tan(e + f x))^{m+1} (c + d \tan(e + f x))^n \text{Simp}[b*B*(b*c*(m+1) + a*d*(n+1)) + A*(a*(b*c - a*d)*(m+1) - b^2*d*(m+n+2)) - (A*b - a*B)*(b*c - a*d)*(m+1)*\tan(e + f x) - b*d*(A*b - a*B)*(m+n+2)*\tan(e + f x)^2, x], x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

rule 4117 $\text{Int}[(a + b \tan(e + f x))^m ((c + d \tan(e + f x) + (f x)))]$, x_Symbol] \rightarrow $\text{Simp}[A/f \text{Subst}[\text{Int}[(a + b*x)^m (c + d*x)^n, x], x, \tan(e + f*x)], x] /;$ FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]

rule 4132 $\text{Int}[(a + b \tan(e + f x))^m ((c + d \tan(e + f x) + (f x)))]$, x_Symbol] \rightarrow $\text{Simp}[(A*b^2 - a*(b*B - a*C))*(a + b \tan(e + f x))^{m+1} ((c + d \tan(e + f x))^{n+1} / (f*(m+1)*(b*c - a*d)*(a^2 + b^2))), x] + \text{Simp}[1/((m+1)*(b*c - a*d)*(a^2 + b^2)) \text{Int}[(a + b \tan(e + f x))^{m+1} (c + d \tan(e + f x))^n \text{Simp}[A*(a*(b*c - a*d)*(m+1) - b^2*d*(m+n+2)) + (b*B - a*C)*(b*c*(m+1) + a*d*(n+1)) - (m+1)*(b*c - a*d)*(A*b - a*B - b*C)*\tan(e + f x) - d*(A*b^2 - a*(b*B - a*C))*(m+n+2)*\tan(e + f x)^2, x], x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2])/(a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.355.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 8042 vs. $2(191) = 382$.

Time = 0.24 (sec) , antiderivative size = 8043, normalized size of antiderivative = 36.73

method	result	size
derivativedivides	Expression too large to display	8043
default	Expression too large to display	8043

```
input int(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output result too large to display
```

3.355.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4530 vs. $2(186) = 372$.

Time = 17.37 (sec) , antiderivative size = 9075, normalized size of antiderivative = 41.44

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorith
m="fricas")
```

```
output Too large to include
```

3.355. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

3.355.6 Sympy [F]

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(A+B\tan(c+dx))\cot^2(c+dx)}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx$$

input `integrate(cot(d*x+c)**2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**2/(a + b*tan(c + d*x))**(3/2), x)`

3.355.7 Maxima [F]

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^2}{(b\tan(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm m="maxima")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^2/(b*tan(d*x + c) + a)^(3/2), x)`

3.355.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm m="giac")`

output `Timed out`

3.355.9 Mupad [B] (verification not implemented)

Time = 12.65 (sec) , antiderivative size = 38368, normalized size of antiderivative = 175.20

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

```
input int((cot(c + d*x)^2*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x)
```

```
output ((2*(A*b^3 - B*a*b^2))/(a*b^2 + a^3) - ((a + b*tan(c + d*x))*(3*A*b^3 + A*
a^2*b - 2*B*a*b^2))/(a*(a*b^2 + a^3)))/(d*(a + b*tan(c + d*x))^(3/2) - a*d
*(a + b*tan(c + d*x))^(1/2)) + atan(-((((((8*A^2*a^3*d^2 - 8*B^2*a^3*d^2 -
16*A*B*b^3*d^2 - 24*A^2*a*b^2*d^2 + 24*B^2*a*b^2*d^2 + 48*A*B*a^2*b*d^2)^
2/4 - (A^4 + 2*A^2*B^2 + B^4)*(16*a^6*d^4 + 16*b^6*d^4 + 48*a^2*b^4*d^4 +
48*a^4*b^2*d^4))^(1/2) - 4*A^2*a^3*d^2 + 4*B^2*a^3*d^2 + 8*A*B*b^3*d^2 + 1
2*A^2*a*b^2*d^2 - 12*B^2*a*b^2*d^2 - 24*A*B*a^2*b*d^2)/(16*(a^6*d^4 + b^6
d^4 + 3*a^2*b^4*d^4 + 3*a^4*b^2*d^4)))^(1/2)*((a + b*tan(c + d*x))^(1/2)*
(576*A^2*a^15*b^28*d^7 + 5184*A^2*a^17*b^26*d^7 + 21568*A^2*a^19*b^24*d^7
+ 53888*A^2*a^21*b^22*d^7 + 87808*A^2*a^23*b^20*d^7 + 94976*A^2*a^25*b^18*
d^7 + 66304*A^2*a^27*b^16*d^7 + 27008*A^2*a^29*b^14*d^7 + 4288*A^2*a^31*b^
12*d^7 - 832*A^2*a^33*b^10*d^7 - 320*A^2*a^35*b^8*d^7 + 256*B^2*a^17*b^26*
d^7 + 1472*B^2*a^19*b^24*d^7 + 3712*B^2*a^21*b^22*d^7 + 6272*B^2*a^23*b^20
*d^7 + 9856*B^2*a^25*b^18*d^7 + 14336*B^2*a^27*b^16*d^7 + 15232*B^2*a^29*b
^14*d^7 + 10112*B^2*a^31*b^12*d^7 + 3712*B^2*a^33*b^10*d^7 + 576*B^2*a^35*
b^8*d^7 - 768*A*B*a^16*b^27*d^7 - 6400*A*B*a^18*b^25*d^7 - 25856*A*B*a^20*
b^23*d^7 - 66304*A*B*a^22*b^21*d^7 - 116480*A*B*a^24*b^19*d^7 - 141568*A*B
*a^26*b^17*d^7 - 116480*A*B*a^28*b^15*d^7 - 61696*A*B*a^30*b^13*d^7 - 1894
4*A*B*a^32*b^11*d^7 - 2560*A*B*a^34*b^9*d^7) - (((8*A^2*a^3*d^2 - 8*B^2*a
^3*d^2 - 16*A*B*b^3*d^2 - 24*A^2*a*b^2*d^2 + 24*B^2*a*b^2*d^2 + 48*A*B*...
```

3.356
$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.356.1 Optimal result 3387
 3.356.2 Mathematica [A] (verified) 3388
 3.356.3 Rubi [A] (warning: unable to verify) 3388
 3.356.4 Maple [B] (verified) 3395
 3.356.5 Fracas [B] (verification not implemented) 3395
 3.356.6 Sympy [F] 3396
 3.356.7 Maxima [F(-1)] 3396
 3.356.8 Giac [F(-1)] 3396
 3.356.9 Mupad [B] (verification not implemented) 3397

3.356.1 Optimal result

Integrand size = 33, antiderivative size = 285

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{(8a^2A - 15Ab^2 + 12abB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{4a^{7/2}d} - \frac{(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a - ib)^{3/2}d} - \frac{(A + iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a + ib)^{3/2}d} + \frac{b(7a^2Ab + 15Ab^3 - 4a^3B - 12ab^2B)}{4a^3(a^2 + b^2)d\sqrt{a + b \tan(c+dx)}} + \frac{(5Ab - 4aB) \cot(c+dx)}{4a^2d\sqrt{a + b \tan(c+dx)}} - \frac{A \cot^2(c+dx)}{2ad\sqrt{a + b \tan(c+dx)}}$$

```
output 1/4*(8*A*a^2-15*A*b^2+12*B*a*b)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/a^(7/2)/d-(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(3/2)/d-(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(3/2)/d+1/4*b*(7*A*a^2*b+15*A*b^3-4*B*a^3-12*B*a*b^2)/a^3/(a^2+b^2)/d/(a+b*tan(d*x+c))^(1/2)+1/4*(5*A*b-4*B*a)*cot(d*x+c)/a^2/d/(a+b*tan(d*x+c))^(1/2)-1/2*A*cot(d*x+c)^2/a/d/(a+b*tan(d*x+c))^(1/2)
```

3.356.2 Mathematica [A] (verified)

Time = 6.28 (sec) , antiderivative size = 409, normalized size of antiderivative = 1.44

$$\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = -\frac{A \cot^2(c + dx)}{2ad\sqrt{a + b \tan(c + dx)}} - \frac{\left(\frac{(a^2 + b^2)(8a^2A - 15Ab^2 + 12abB) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a}}\right)}{4\sqrt{ad}}\right) + \frac{i\sqrt{a - ib}(a^3(Ab - aB) - ia^3(aA + bB)) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a}}\right)}{(-a + ib)d}}{a(a^2 + b^2)} - \frac{(5Ab - 4aB) \cot(c + dx)}{2ad\sqrt{a + b \tan(c + dx)}}$$

2a

input `Integrate[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2), x]`

output `-1/2*(A*Cot[c + d*x]^2)/(a*d*Sqrt[a + b*Tan[c + d*x]]) - (-1/2*((5*A*b - 4*a*B)*Cot[c + d*x])/(a*d*Sqrt[a + b*Tan[c + d*x]]) - ((2*(((a^2 + b^2)*(8*a^2*A - 15*A*b^2 + 12*a*b*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/(4*Sqrt[a]*d) + (I*Sqrt[a - I*b]*(a^3*(A*b - a*B) - I*a^3*(a*A + b*B))*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/((-a + I*b)*d) - (I*Sqrt[a + I*b]*(a^3*(A*b - a*B) + I*a^3*(a*A + b*B))*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/((-a - I*b)*d)))/(a*(a^2 + b^2)) + (2*((b^2*(-8*a^2*A + 15*A*b^2 - 12*a*b*B))/4 - a*(-2*a^2*b*B - (3*a*b*(5*A*b - 4*a*B))/4)))/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])/a)/(2*a)`

3.356.3 Rubi [A] (warning: unable to verify)

Time = 2.35 (sec) , antiderivative size = 324, normalized size of antiderivative = 1.14, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^3(a + b \tan(c + dx))^{3/2}} dx$$

$$\begin{aligned}
 & \int \frac{\cot^2(c+dx)(5Ab \tan^2(c+dx)+4aA \tan(c+dx)+5Ab-4aB)}{2(a+b \tan(c+dx))^{3/2}} dx - \frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 4092 \\
 & \int \frac{\cot^2(c+dx)(5Ab \tan^2(c+dx)+4aA \tan(c+dx)+5Ab-4aB)}{(a+b \tan(c+dx))^{3/2}} dx - \frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 27 \\
 & \int \frac{5Ab \tan(c+dx)^2+4aA \tan(c+dx)+5Ab-4aB}{\tan(c+dx)^2(a+b \tan(c+dx))^{3/2}} dx - \frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 3042 \\
 & \int \frac{\cot(c+dx)(8Aa^2+8B \tan(c+dx)a^2+12bBa-15Ab^2-3b(5Ab-4aB) \tan^2(c+dx))}{2(a+b \tan(c+dx))^{3/2}} dx - \frac{(5Ab-4aB) \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 4132 \\
 & \frac{4a}{2ad\sqrt{a+b \tan(c+dx)}} \frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 27 \\
 & \int \frac{\cot(c+dx)(8Aa^2+8B \tan(c+dx)a^2+12bBa-15Ab^2-3b(5Ab-4aB) \tan^2(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx - \frac{(5Ab-4aB) \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 3042 \\
 & \int \frac{8Aa^2+8B \tan(c+dx)a^2+12bBa-15Ab^2-3b(5Ab-4aB) \tan(c+dx)^2}{\tan(c+dx)(a+b \tan(c+dx))^{3/2}} dx - \frac{(5Ab-4aB) \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 4132 \\
 & \frac{2 \int \frac{\cot(c+dx)(-8(Ab-aB) \tan(c+dx)a^3-b(-4Ba^3+7Aba^2-12b^2Ba+15Ab^3) \tan^2(c+dx)+(a^2+b^2)(8Aa^2+12bBa-15Ab^2))}{2\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-4a^3B+7a^2Ab-12ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow \\
 & \frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}}
 \end{aligned}$$

3.356. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

↓ 27

$$\frac{\int \frac{\cot(c+dx)(-8(Ab-aB)\tan(c+dx)a^3 - b(-4Ba^3+7Aba^2-12b^2Ba+15Ab^3)\tan^2(c+dx) + (a^2+b^2)(8Aa^2+12bBa-15Ab^2))}{\sqrt{a+b\tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}$$

$$\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b\tan(c+dx)}} \quad 4a$$

↓ 3042

$$\frac{\int \frac{-8(Ab-aB)\tan(c+dx)a^3 - b(-4Ba^3+7Aba^2-12b^2Ba+15Ab^3)\tan(c+dx)^2 + (a^2+b^2)(8Aa^2+12bBa-15Ab^2)}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}$$

$$\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b\tan(c+dx)}} \quad 4a$$

↓ 4136

$$\frac{\int -\frac{8((Ab-aB)a^3+(aA+bB)\tan(c+dx)a^3)}{\sqrt{a+b\tan(c+dx)}} dx + (a^2+b^2)(8a^2A+12abB-15Ab^2) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b\tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}$$

$$\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b\tan(c+dx)}} \quad 4a$$

↓ 27

$$\frac{(a^2+b^2)(8a^2A+12abB-15Ab^2) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b\tan(c+dx)}} dx - 8 \int \frac{(Ab-aB)a^3+(aA+bB)\tan(c+dx)a^3}{\sqrt{a+b\tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}$$

$$\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b\tan(c+dx)}} \quad 4a$$

↓ 3042

$$\frac{(a^2+b^2)(8a^2A+12abB-15Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - 8 \int \frac{(Ab-aB)a^3+(aA+bB)\tan(c+dx)a^3}{\sqrt{a+b\tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}$$

$$\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b\tan(c+dx)}} \quad 4a$$

3.356. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$

$$\begin{aligned} & \downarrow 4022 \\ & -\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}} - \\ & -\frac{(5Ab-4aB) \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} + \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(8a^2A+12abB-15Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 8\left(\frac{1}{2}a^3(b+ia)(A+ib)\right)}{2a} \frac{1}{a(a^2+b^2)} \end{aligned}$$

4a

$$\begin{aligned} & \downarrow 3042 \\ & -\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}} - \\ & -\frac{(5Ab-4aB) \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} + \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(8a^2A+12abB-15Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 8\left(\frac{1}{2}a^3(b+ia)(A+ib)\right)}{2a} \frac{1}{a(a^2+b^2)} \end{aligned}$$

4a

$$\begin{aligned} & \downarrow 4020 \\ & -\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}} - \\ & -\frac{(5Ab-4aB) \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} + \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(8a^2A+12abB-15Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 8\left(-\frac{ia^3(-b+ia)(A-ib)}{a(a^2+b^2)}\right)}{2a} \frac{1}{a(a^2+b^2)} \end{aligned}$$

4a

$$\begin{aligned} & \downarrow 25 \\ & -\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}} - \\ & -\frac{(5Ab-4aB) \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} + \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(8a^2A+12abB-15Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 8\left(\frac{ia^3(-b+ia)(A-ib)}{a(a^2+b^2)}\right)}{2a} \frac{1}{a(a^2+b^2)} \end{aligned}$$

4a

$$\begin{aligned} & \downarrow 73 \\ & -\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}} - \\ & -\frac{(5Ab-4aB) \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} + \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(8a^2A+12abB-15Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 8\left(\frac{a^3(b+ia)(A+ib)}{a(a^2+b^2)}\right)}{2a} \frac{1}{a(a^2+b^2)} \end{aligned}$$

2a

4a

$$\begin{aligned} & \downarrow 221 \\ & -\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}} - \\ & -\frac{(5Ab-4aB) \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} + \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(8a^2A+12abB-15Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 8\left(\frac{a^3(b+ia)(A+ib)}{a(a^2+b^2)}\right)}{2a} \frac{1}{a(a^2+b^2)} \end{aligned}$$

3.356. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

$$\frac{-\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}} - \frac{(5Ab-4aB) \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} + \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(8a^2A+12abB-15Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - 8 \left(\frac{a^3(b+ia)(A+iB)}{d} \right)}{2a}}{4a}$$

4117

$$\frac{-\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}} - \frac{(5Ab-4aB) \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} + \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)(8a^2A+12abB-15Ab^2) \int \frac{\cot(c+dx)}{\sqrt{a+b \tan(c+dx)}} d \tan(c+dx) - 8 \left(\frac{a^3(b+ia)(A+iB)}{d} \right)}{2a}}{4a}$$

73

$$\frac{-\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}} - \frac{(5Ab-4aB) \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} + \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{2(a^2+b^2)(8a^2A+12abB-15Ab^2) \int \frac{\frac{1}{a+b \tan(c+dx)} - \frac{a}{b}}{bd} d\sqrt{a+b \tan(c+dx)} - 8 \left(\frac{a^3(b+ia)(A+iB)}{d} \right)}{2a}}{4a}$$

221

$$\frac{-\frac{A \cot^2(c+dx)}{2ad\sqrt{a+b \tan(c+dx)}} - \frac{(5Ab-4aB) \cot(c+dx)}{ad\sqrt{a+b \tan(c+dx)}} + \frac{2b(-4a^3B+7a^2Ab-12ab^2B+15Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{2(a^2+b^2)(8a^2A+12abB-15Ab^2) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right) - 8 \left(\frac{a^3(b+ia)(A+iB)}{d} \right)}{\sqrt{ad}}}{2a}}{4a}$$

```
input Int[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]
```

```
output -1/2*(A*Cot[c + d*x]^2)/(a*d*Sqrt[a + b*Tan[c + d*x]]) - (((5*A*b - 4*A*B)*Cot[c + d*x])/(a*d*Sqrt[a + b*Tan[c + d*x]])) + ((-8*(-(a^3*(I*a - b)*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d)) + (a^3*(I*a + b)*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d)) - (2*(a^2 + b^2)*(8*a^2*A - 15*A*b^2 + 12*a*b*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/(Sqrt[a]*d)/(a*(a^2 + b^2)) - (2*b*(7*a^2*A*b + 15*A*b^3 - 4*a^3*B - 12*a*b^2*B))/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])/(2*a))/(4*a)
```

3.356.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 73 Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```


rule 4022 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]]^{(m_.)}((c_.) + (d_.)\tan[(e_.) + (f_.)x]), x_Symbol] \rightarrow \text{Simp}[(c + I*d)/2 \text{ Int}[(a + b*\tan[e + f*x])^m(1 - I*\tan[e + f*x]), x], x] + \text{Simp}[(c - I*d)/2 \text{ Int}[(a + b*\tan[e + f*x])^m(1 + I*\tan[e + f*x]), x], x] /;$ FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]

rule 4092 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]]^{(m_.)}((A_.) + (B_.)\tan[(e_.) + (f_.)x])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[b*(A*b - a*B)*(a + b*\tan[e + f*x])^{(m+1)}((c + d*\tan[e + f*x])^{(n+1)}/(f*(m+1)*(b*c - a*d)*(a^2 + b^2))), x] + \text{Simp}[1/((m+1)*(b*c - a*d)*(a^2 + b^2)) \text{ Int}[(a + b*\tan[e + f*x])^{(m+1)}(c + d*\tan[e + f*x])^n \text{Simp}[b*B*(b*c*(m+1) + a*d*(n+1)) + A*(a*(b*c - a*d)*(m+1) - b^2*d*(m+n+2)) - (A*b - a*B)*(b*c - a*d)*(m+1)*\tan[e + f*x] - b*d*(A*b - a*B)*(m+n+2)*\tan[e + f*x]^2, x], x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

rule 4117 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]]^{(m_.)}((c_.) + (d_.)\tan[(e_.) + (f_.)x])^{(n_.)}((A_.) + (C_.)\tan[(e_.) + (f_.)x])^2, x_Symbol] \rightarrow \text{Simp}[A/f \text{ Subst}[\text{Int}[(a + b*x)^m(c + d*x)^n, x], x, \tan[e + f*x]], x] /;$ FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]

rule 4132 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x]]^{(m_.)}((c_.) + (d_.)\tan[(e_.) + (f_.)x])^{(n_.)}((A_.) + (B_.)\tan[(e_.) + (f_.)x] + (C_.)\tan[(e_.) + (f_.)x])^2, x_Symbol] \rightarrow \text{Simp}[(A*b^2 - a*(b*B - a*C))*(a + b*\tan[e + f*x])^{(m+1)}((c + d*\tan[e + f*x])^{(n+1)}/(f*(m+1)*(b*c - a*d)*(a^2 + b^2))), x] + \text{Simp}[1/((m+1)*(b*c - a*d)*(a^2 + b^2)) \text{ Int}[(a + b*\tan[e + f*x])^{(m+1)}(c + d*\tan[e + f*x])^n \text{Simp}[A*(a*(b*c - a*d)*(m+1) - b^2*d*(m+n+2)) + (b*B - a*C)*(b*c*(m+1) + a*d*(n+1)) - (m+1)*(b*c - a*d)*(A*b - a*B - b*C)*\tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m+n+2)*\tan[e + f*x]^2, x], x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2])/(a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.356.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 8163 vs. $2(247) = 494$.

Time = 0.24 (sec) , antiderivative size = 8164, normalized size of antiderivative = 28.65

method	result	size
derivativedivides	Expression too large to display	8164
default	Expression too large to display	8164

```
input int(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output result too large to display
```

3.356.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4601 vs. $2(241) = 482$.

Time = 42.42 (sec) , antiderivative size = 9219, normalized size of antiderivative = 32.35

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorith
m="fracas")
```

```
output Too large to include
```

3.356.6 Sympy [F]

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(A+B\tan(c+dx))\cot^3(c+dx)}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx$$

input `integrate(cot(d*x+c)**3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**3/(a + b*tan(c + d*x))**(3/2), x)`

3.356.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm m="maxima")`

output `Timed out`

3.356.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm m="giac")`

output `Timed out`

3.356.9 Mupad [B] (verification not implemented)

Time = 12.50 (sec) , antiderivative size = 42371, normalized size of antiderivative = 148.67

$$\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

```
input int((cot(c + d*x)^3*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x)
```

```
output ((2*(A*b^4 - B*a*b^3))/(a*(a^2 + b^2)) + ((a + b*tan(c + d*x))^2*(15*A*b^4
+ 7*A*a^2*b^2 - 12*B*a*b^3 - 4*B*a^3*b))/(4*a^3*(a^2 + b^2)) - ((a + b*ta
n(c + d*x))*(25*A*b^4 + 9*A*a^2*b^2 - 20*B*a*b^3 - 4*B*a^3*b))/(4*a^2*(a^2
+ b^2)))/(d*(a + b*tan(c + d*x))^(5/2) - 2*a*d*(a + b*tan(c + d*x))^(3/2)
+ a^2*d*(a + b*tan(c + d*x))^(1/2)) + atan((((a + b*tan(c + d*x))^(1/2))*
704643072*A^4*a^29*b^20*d^5 - 290979840*A^4*a^23*b^26*d^5 - 465043456*A^4*
a^25*b^24*d^5 - 37224448*A^4*a^27*b^22*d^5 - 58982400*A^4*a^21*b^28*d^5 +
767033344*A^4*a^31*b^18*d^5 + 238551040*A^4*a^33*b^16*d^5 + 1572864*A^4*a^
35*b^14*d^5 + 92536832*A^4*a^37*b^12*d^5 + 96468992*A^4*a^39*b^10*d^5 + 25
165824*A^4*a^41*b^8*d^5 + 37748736*B^4*a^23*b^26*d^5 + 226492416*B^4*a^25*
b^24*d^5 + 536870912*B^4*a^27*b^22*d^5 + 587202560*B^4*a^29*b^20*d^5 + 176
160768*B^4*a^31*b^18*d^5 - 234881024*B^4*a^33*b^16*d^5 - 234881024*B^4*a^3
5*b^14*d^5 - 50331648*B^4*a^37*b^12*d^5 + 20971520*B^4*a^39*b^10*d^5 + 838
8608*B^4*a^41*b^8*d^5 - 94371840*A*B^3*a^22*b^27*d^5 - 364904448*A*B^3*a^2
4*b^25*d^5 + 37748736*A*B^3*a^26*b^23*d^5 + 2554331136*A*B^3*a^28*b^21*d^5
+ 5989466112*A*B^3*a^30*b^19*d^5 + 6606028800*A*B^3*a^32*b^17*d^5 + 37874
56512*A*B^3*a^34*b^15*d^5 + 918552576*A*B^3*a^36*b^13*d^5 - 56623104*A*B^3
*a^38*b^11*d^5 - 50331648*A*B^3*a^40*b^9*d^5 + 330301440*A^3*B*a^22*b^27*d
^5 + 1915748352*A^3*B*a^24*b^25*d^5 + 4279238656*A^3*B*a^26*b^23*d^5 + 405
9037696*A^3*B*a^28*b^21*d^5 + 154140672*A^3*B*a^30*b^19*d^5 - 282591232...
```

3.357 $\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

3.357.1 Optimal result 3398
 3.357.2 Mathematica [C] (verified) 3399
 3.357.3 Rubi [A] (warning: unable to verify) 3400
 3.357.4 Maple [B] (verified) 3406
 3.357.5 Fricas [B] (verification not implemented) 3407
 3.357.6 Sympy [F] 3407
 3.357.7 Maxima [F(-1)] 3407
 3.357.8 Giac [F(-1)] 3408
 3.357.9 Mupad [B] (verification not implemented) 3408

3.357.1 Optimal result

Integrand size = 33, antiderivative size = 371

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = -\frac{(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{5/2}d} + \frac{(iA-B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a+ib)^{5/2}d} + \frac{2a(Ab-aB) \tan^3(c+dx)}{3b(a^2+b^2)d(a+b \tan(c+dx))^{3/2}} + \frac{2a(a^2Ab+3Ab^3-2a^3B-4ab^2B) \tan^2(c+dx)}{b^2(a^2+b^2)^2 d \sqrt{a+b \tan(c+dx)}} + \frac{2(8a^4Ab+17a^2Ab^3+3Ab^5-16a^5B-30a^3b^2B-8ab^4B) \sqrt{a+b \tan(c+dx)}}{3b^4(a^2+b^2)^2 d} - \frac{2(4a^3Ab+10aAb^3-8a^4B-15a^2b^2B-b^4B) \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3b^3(a^2+b^2)^2 d}$$

```
output -(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(5/2)/d+(I*
A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(5/2)/d+2/3*(8*
A*a^4*b+17*A*a^2*b^3+3*A*b^5-16*B*a^5-30*B*a^3*b^2-8*B*a*b^4)*(a+b*tan(d*x
+c))^(1/2)/b^4/(a^2+b^2)^2/d-2/3*(4*A*a^3*b+10*A*a*b^3-8*B*a^4-15*B*a^2*b^
2-B*b^4)*(a+b*tan(d*x+c))^(1/2)*tan(d*x+c)/b^3/(a^2+b^2)^2/d+2*a*(A*a^2*b+
3*A*b^3-2*B*a^3-4*B*a*b^2)*tan(d*x+c)^2/b^2/(a^2+b^2)^2/d/(a+b*tan(d*x+c))
^(1/2)+2/3*a*(A*b-B*a)*tan(d*x+c)^3/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)
```

3.357.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.39 (sec) , antiderivative size = 450, normalized size of antiderivative = 1.21

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{2B \tan^3(c+dx)}{3bd(a+b \tan(c+dx))^{3/2}}$$

$$\begin{aligned}
 & \left(\frac{3(Ab-2aB) \tan^2(c+dx)}{bd(a+b \tan(c+dx))^{3/2}} + \right. \\
 & \quad \left. \frac{3(4aAb-8a^2B+b^2B) \tan(c+dx)}{2bd(a+b \tan(c+dx))^{3/2}} - \right. \\
 & \quad \left. \frac{2(8a^2Ab+Ab^3-16a^3B+2ab^2B)}{3b(a+b \tan(c+dx))^{3/2}} + \frac{\left(-\frac{3Ab^5}{2} + \frac{3}{2}ab^4B\right) \left(-\frac{\text{Hypergeometric2F1}}{3(a+b)}\right)}{2} \right)
 \end{aligned}$$

input `Integrate[(Tan[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]`

$$3.357. \quad \int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

output $(2*B*\text{Tan}[c + d*x]^3)/(3*b*d*(a + b*\text{Tan}[c + d*x])^{(3/2)}) + (2*((3*(A*b - 2*a*B)*\text{Tan}[c + d*x]^2)/(b*d*(a + b*\text{Tan}[c + d*x])^{(3/2)}) + (2*((3*(4*a*A*b - 8*a^2*B + b^2*B)*\text{Tan}[c + d*x]))/(2*b*d*(a + b*\text{Tan}[c + d*x])^{(3/2)}) - (3*((-2*(8*a^2*A*b + A*b^3 - 16*a^3*B + 2*a*b^2*B)))/(3*b*(a + b*\text{Tan}[c + d*x])^{(3/2)}) + (2*((((-3*A*b^5)/2 + (3*a*b^4*B)/2)*(-1/3*\text{Hypergeometric2F1}[-3/2, 1, -1/2, (a + b*\text{Tan}[c + d*x])/(a - I*b)])/((I*a + b)*(a + b*\text{Tan}[c + d*x])^{(3/2)}) + \text{Hypergeometric2F1}[-3/2, 1, -1/2, (a + b*\text{Tan}[c + d*x])/(a + I*b)]/(3*(I*a - b)*(a + b*\text{Tan}[c + d*x])^{(3/2)})))/b - (3*b^3*B*(-\text{Hypergeometric2F1}[-1/2, 1, 1/2, (a + b*\text{Tan}[c + d*x])/(a - I*b)]/((I*a + b)*\text{Sqrt}[a + b*\text{Tan}[c + d*x]])) + \text{Hypergeometric2F1}[-1/2, 1, 1/2, (a + b*\text{Tan}[c + d*x])/(a + I*b)]/((I*a - b)*\text{Sqrt}[a + b*\text{Tan}[c + d*x]]))/2))/(3*b)))/(4*b*d))/b)/(3*b)$

3.357.3 Rubi [A] (warning: unable to verify)

Time = 2.17 (sec) , antiderivative size = 395, normalized size of antiderivative = 1.06, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4130, 27, 3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{\tan(c+dx)^4(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

↓ 4088

$$\frac{2 \int -\frac{3 \tan^2(c+dx)((-2Ba^2+Aba-b^2B) \tan^2(c+dx)-b(Ab-aB) \tan(c+dx)+2a(Ab-aB))}{2(a+b \tan(c+dx))^{3/2}} dx}{3b(a^2+b^2)} + \frac{2a(Ab-aB) \tan^3(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^{3/2}}$$

↓ 27

$$\frac{2a(Ab-aB) \tan^3(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^{3/2}} - \frac{\int \frac{\tan^2(c+dx)((-2Ba^2+Aba-b^2B) \tan^2(c+dx)-b(Ab-aB) \tan(c+dx)+2a(Ab-aB))}{(a+b \tan(c+dx))^{3/2}} dx}{b(a^2+b^2)}$$

3.357. $\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\begin{array}{c}
 \downarrow 3042 \\
 \frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \\
 \int \frac{\tan(c+dx)^2((-2Ba^2 + Aba - b^2B) \tan(c+dx)^2 - b(Ab - aB) \tan(c+dx) + 2a(Ab - aB))}{(a + b \tan(c+dx))^{3/2}} dx \\
 \frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \\
 \downarrow 4128 \\
 2 \int \frac{\tan(c+dx)((Aa^2 + 2bBa - Ab^2) \tan(c+dx)b^2 + (-8Ba^4 + 4Aba^3 - 15b^2Ba^2 + 10Ab^3a - b^4B) \tan^2(c+dx) + 4a(-2Ba^3 + Aba^2 - 4b^2Ba + 3Ab^3))}{2\sqrt{a+b \tan(c+dx)} b(a^2+b^2)} dx - \frac{2a(-2a^3B - b^4)}{b(a^2+b^2)} \\
 \downarrow 27 \\
 \frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \\
 \int \frac{\tan(c+dx)((Aa^2 + 2bBa - Ab^2) \tan(c+dx)b^2 + (-8Ba^4 + 4Aba^3 - 15b^2Ba^2 + 10Ab^3a - b^4B) \tan^2(c+dx) + 4a(-2Ba^3 + Aba^2 - 4b^2Ba + 3Ab^3))}{\sqrt{a+b \tan(c+dx)} b(a^2+b^2)} dx - \frac{2a(-2a^3B - b^4)}{bd(a^2+b^2)} \\
 \downarrow 3042 \\
 \frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \\
 \int \frac{\tan(c+dx)((Aa^2 + 2bBa - Ab^2) \tan(c+dx)b^2 + (-8Ba^4 + 4Aba^3 - 15b^2Ba^2 + 10Ab^3a - b^4B) \tan^2(c+dx) + 4a(-2Ba^3 + Aba^2 - 4b^2Ba + 3Ab^3))}{\sqrt{a+b \tan(c+dx)} b(a^2+b^2)} dx - \frac{2a(-2a^3B - b^4)}{bd(a^2+b^2)} \\
 \downarrow 4130 \\
 \frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \\
 2 \int \frac{-3(-Ba^2 + 2Aba + b^2B) \tan(c+dx)b^3 + (-16Ba^5 + 8Aba^4 - 30b^2Ba^3 + 17Ab^3a^2 - 8b^4Ba + 3Ab^5) \tan^2(c+dx) + 2a(-8Ba^4 + 4Aba^3 - 15b^2Ba^2 + 10Ab^3a - b^4B)}{2\sqrt{a+b \tan(c+dx)} 3b} dx \\
 \frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \\
 \downarrow 27 \\
 \frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \\
 \frac{2(-8a^4B + 4a^3Ab - 15a^2b^2B + 10aAb^3 - b^4B) \tan(c+dx) \sqrt{a+b \tan(c+dx)}}{3bd} - \int \frac{-3(-Ba^2 + 2Aba + b^2B) \tan(c+dx)b^3 + (-16Ba^5 + 8Aba^4 - 30b^2Ba^3 + 17Ab^3a^2 - 8b^4Ba + 3Ab^5)}{\sqrt{a+b \tan(c+dx)} 3b} dx \\
 \frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} -
 \end{array}$$

3.357. $\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

↓ 3042

$$\frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2(-8a^4B + 4a^3Ab - 15a^2b^2B + 10aAb^3 - b^4B) \tan(c + dx) \sqrt{a + b \tan(c + dx)}}{3bd} - \frac{\int \frac{-3(-Ba^2 + 2Aba + b^2B) \tan(c + dx)b^3 + (-16Ba^5 + 8Aba^4 - 30b^2Ba^3 + 17Ab^3a^2 - 8b^4B)}{\sqrt{a + b \tan(c + dx)}} dx}{3b}$$

$$b(a^2 + b^2)$$

↓ 4113

$$\frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2(-8a^4B + 4a^3Ab - 15a^2b^2B + 10aAb^3 - b^4B) \tan(c + dx) \sqrt{a + b \tan(c + dx)}}{3bd} - \frac{\int \frac{3b^3(Aa^2 + 2bBa - Ab^2) - 3b^3(-Ba^2 + 2Aba + b^2B) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx}{3b} + \frac{2(-16a^5B + 8a^4Ab)}{3b}$$

$$b(a^2 + b^2)$$

↓ 3042

$$\frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2(-8a^4B + 4a^3Ab - 15a^2b^2B + 10aAb^3 - b^4B) \tan(c + dx) \sqrt{a + b \tan(c + dx)}}{3bd} - \frac{\int \frac{3b^3(Aa^2 + 2bBa - Ab^2) - 3b^3(-Ba^2 + 2Aba + b^2B) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx}{3b} + \frac{2(-16a^5B + 8a^4Ab)}{3b}$$

$$b(a^2 + b^2)$$

↓ 4022

$$\frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2a(-2a^3B + a^2Ab - 4ab^2B + 3Ab^3) \tan^2(c + dx)}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{2(-8a^4B + 4a^3Ab - 15a^2b^2B + 10aAb^3 - b^4B) \tan(c + dx) \sqrt{a + b \tan(c + dx)}}{3bd} - \frac{\frac{3}{2}b^3(a - ib)^2(A + iB) \int \frac{1 - i}{\sqrt{a + b \tan(c + dx)}} dx}{3b}$$

$$b(a^2 + b^2)$$

↓ 3042

$$\frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2a(-2a^3B + a^2Ab - 4ab^2B + 3Ab^3) \tan^2(c + dx)}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{2(-8a^4B + 4a^3Ab - 15a^2b^2B + 10aAb^3 - b^4B) \tan(c + dx) \sqrt{a + b \tan(c + dx)}}{3bd} - \frac{\frac{3}{2}b^3(a - ib)^2(A + iB) \int \frac{1 - i}{\sqrt{a + b \tan(c + dx)}} dx}{3b}$$

$$b(a^2 + b^2)$$

↓ 4020

3.357. $\int \frac{\tan^4(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx$

$$\frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2a(-2a^3B + a^2Ab - 4ab^2B + 3Ab^3) \tan^2(c + dx)}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{2(-8a^4B + 4a^3Ab - 15a^2b^2B + 10aAb^3 - b^4B) \tan(c + dx)\sqrt{a + b \tan(c + dx)}}{3bd} - \frac{3ib^3(a + ib)^2(A - iB) \int \frac{1}{\sqrt{a + b \tan(c + dx)}} dx}{b(a^2 + b^2)}$$

↓ 25

$$\frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2a(-2a^3B + a^2Ab - 4ab^2B + 3Ab^3) \tan^2(c + dx)}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{2(-8a^4B + 4a^3Ab - 15a^2b^2B + 10aAb^3 - b^4B) \tan(c + dx)\sqrt{a + b \tan(c + dx)}}{3bd} - \frac{3ib^3(a + ib)^2(A - iB) \int \frac{1}{\sqrt{a + b \tan(c + dx)}} dx}{b(a^2 + b^2)}$$

↓ 73

$$\frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2a(-2a^3B + a^2Ab - 4ab^2B + 3Ab^3) \tan^2(c + dx)}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{2(-8a^4B + 4a^3Ab - 15a^2b^2B + 10aAb^3 - b^4B) \tan(c + dx)\sqrt{a + b \tan(c + dx)}}{3bd} - \frac{3b^2(a - ib)^2(A + iB) \int \frac{1}{\sqrt{a + b \tan(c + dx)}} dx}{b(a^2 + b^2)}$$

↓ 221

$$\frac{2a(Ab - aB) \tan^3(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2a(-2a^3B + a^2Ab - 4ab^2B + 3Ab^3) \tan^2(c + dx)}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{2(-8a^4B + 4a^3Ab - 15a^2b^2B + 10aAb^3 - b^4B) \tan(c + dx)\sqrt{a + b \tan(c + dx)}}{3bd} - \frac{2(-16a^5B + 8a^4Ab - 30a^3b^2B + 16a^2b^3B - 8ab^4B) \sqrt{a + b \tan(c + dx)}}{b(a^2 + b^2)}$$

input `Int[(Tan[c + d*x]^4*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]`

output `(2*a*(A*b - a*B)*Tan[c + d*x]^3)/(3*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) - ((-2*a*(a^2*A*b + 3*A*b^3 - 2*a^3*B - 4*a*b^2*B)*Tan[c + d*x]^2)/(b*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]) + ((2*(4*a^3*A*b + 10*a*A*b^3 - 8*a^4*B - 15*a^2*b^2*B - b^4*B)*Tan[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/(3*b*d) - ((3*(a + I*b)^2*b^3*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d) + (3*(a - I*b)^2*b^3*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d) + (2*(8*a^4*A*b + 17*a^2*A*b^3 + 3*A*b^5 - 16*a^5*B - 30*a^3*b^2*B - 8*a*b^4*B)*Sqrt[a + b*Tan[c + d*x]])/(b*d))/(3*b))/(b*(a^2 + b^2))`

3.357. $\int \frac{\tan^4(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

3.357.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(GtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.357.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4591 vs. $2(339) = 678$.

Time = 0.20 (sec) , antiderivative size = 4592, normalized size of antiderivative = 12.38

method	result	size
parts	Expression too large to display	4592
derivativedivides	Expression too large to display	12953
default	Expression too large to display	12953

```
input int(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2), x, method=_RETURNV
ERBOSE)
```

```
output A*(-3/4/d*b^3/(a^2+b^2)^(7/2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)
+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*
a+1/d/b/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)
^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))
*a^4-1/d/b/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b
^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/
2))*a^6-4/d*b/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^
2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(
1/2))*a^4-1/d*b^3/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((
2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-
2*a)^(1/2))*a^2-1/4/d/b/(a^2+b^2)^(7/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))
^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*
a)^(1/2)*a^5+1/2/d*b/(a^2+b^2)^(7/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(
1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a
)^(1/2)*a^3+3/4/d*b^3/(a^2+b^2)^(7/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(
1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a
)^(1/2)*a-1/d/b/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a
+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a
)^(1/2))*a^4+1/d/b/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2
*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)+2*a)^(1/2))
```

3.357.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7358 vs. $2(334) = 668$.

Time = 2.20 (sec) , antiderivative size = 7358, normalized size of antiderivative = 19.83

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="fricas")`

output Too large to include

3.357.6 Sympy [F]

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\tan^4(c+dx)}{(a+b\tan(c+dx))^{5/2}} dx$$

input `integrate(tan(d*x+c)**4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**4/(a + b*tan(c + d*x))**(5/2), x)`

3.357.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\tan^4(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output Timed out

3.357.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^4(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^4*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Timed out`

3.357.9 Mupad [B] (verification not implemented)

Time = 47.63 (sec) , antiderivative size = 9547, normalized size of antiderivative = 25.73

$$\int \frac{\tan^4(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `int((tan(c + d*x)^4*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)`

output `(log((((a + b*tan(c + d*x))^(1/2)*(320*A^2*a^4*b^14*d^3 - 16*A^2*b^18*d^3 + 1024*A^2*a^6*b^12*d^3 + 1440*A^2*a^8*b^10*d^3 + 1024*A^2*a^10*b^8*d^3 + 320*A^2*a^12*b^6*d^3 - 16*A^2*a^16*b^2*d^3) - (((320*A^4*a^2*b^8*d^4 - 16*A^4*b^10*d^4 - 1760*A^4*a^4*b^6*d^4 + 1600*A^4*a^6*b^4*d^4 - 400*A^4*a^8*b^2*d^4)^(1/2) - 4*A^2*a^5*d^2 + 40*A^2*a^3*b^2*d^2 - 20*A^2*a*b^4*d^2)/(a^10*d^4 + b^10*d^4 + 5*a^2*b^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^(1/2)*((((320*A^4*a^2*b^8*d^4 - 16*A^4*b^10*d^4 - 1760*A^4*a^4*b^6*d^4 + 1600*A^4*a^6*b^4*d^4 - 400*A^4*a^8*b^2*d^4)^(1/2) - 4*A^2*a^5*d^2 + 40*A^2*a^3*b^2*d^2 - 20*A^2*a*b^4*d^2)/(a^10*d^4 + b^10*d^4 + 5*a^2*b^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^(1/2)*(a + b*tan(c + d*x))^(1/2)*(64*a*b^22*d^5 + 640*a^3*b^20*d^5 + 2880*a^5*b^18*d^5 + 7680*a^7*b^16*d^5 + 13440*a^9*b^14*d^5 + 16128*a^11*b^12*d^5 + 13440*a^13*b^10*d^5 + 7680*a^15*b^8*d^5 + 2880*a^17*b^6*d^5 + 640*a^19*b^4*d^5 + 64*a^21*b^2*d^5))/4 - 32*A*b^21*d^4 - 160*A*a^2*b^19*d^4 - 128*A*a^4*b^17*d^4 + 896*A*a^6*b^15*d^4 + 3136*A*a^8*b^13*d^4 + 4928*A*a^10*b^11*d^4 + 4480*A*a^12*b^9*d^4 + 2432*A*a^14*b^7*d^4 + 736*A*a^16*b^5*d^4 + 96*A*a^18*b^3*d^4))/4)*(((320*A^4*a^2*b^8*d^4 - 16*A^4*b^10*d^4 - 1760*A^4*a^4*b^6*d^4 + 1600*A^4*a^6*b^4*d^4 - 400*A^4*a^8*b^2*d^4)^(1/2) - 4*A^2*a^5*d^2 + 40*A^2*a^3*b^2*d^2 - 20*A^2*a*b^4*d^2)/(a^10*d^4 + b^10*d^4 + 5*a^2*b^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^(1/2))/4 + 96*A^3*a^3...`

3.358
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

3.358.1 Optimal result 3409
 3.358.2 Mathematica [C] (verified) 3410
 3.358.3 Rubi [A] (warning: unable to verify) 3410
 3.358.4 Maple [B] (verified) 3415
 3.358.5 Fracas [B] (verification not implemented) 3416
 3.358.6 Sympy [F] 3416
 3.358.7 Maxima [F(-1)] 3416
 3.358.8 Giac [F(-1)] 3417
 3.358.9 Mupad [B] (verification not implemented) 3417

3.358.1 Optimal result

Integrand size = 33, antiderivative size = 261

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{5/2}d} + \frac{(A+iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a+ib)^{5/2}d} + \frac{2a(Ab-aB) \tan^2(c+dx)}{3b(a^2+b^2)d(a+b \tan(c+dx))^{3/2}} - \frac{2a^2(a^2Ab+7Ab^3-4a^3B-10ab^2B)}{3b^3(a^2+b^2)^2d\sqrt{a+b \tan(c+dx)}} - \frac{2(aAb-4a^2B-3b^2B)\sqrt{a+b \tan(c+dx)}}{3b^3(a^2+b^2)d}$$

```
output (A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(5/2)/d+(A+I
*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(5/2)/d-2/3*a^2*
(A*a^2*b+7*A*b^3-4*B*a^3-10*B*a*b^2)/b^3/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^(1
/2)-2/3*(A*a*b-4*B*a^2-3*B*b^2)*(a+b*tan(d*x+c))^(1/2)/b^3/(a^2+b^2)/d+2/3
*a*(A*b-B*a)*tan(d*x+c)^2/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)
```


3.358.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 3.64 (sec) , antiderivative size = 309, normalized size of antiderivative = 1.18

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx =$$

$$-2(a-ib)(a+ib)(-2aAb+8a^2B+b^2B)-b^2(aA+bB)\left(i(a+ib)\operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{a+b \tan(c+dx)}{a+b \tan(c+dx)}\right)\right)$$

input `Integrate[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]`

output `-1/3*(-2*(a - I*b)*(a + I*b)*(-2*a*A*b + 8*a^2*B + b^2*B) - b^2*(a*A + b*B))*(I*(a + I*b)*Hypergeometric2F1[-3/2, 1, -1/2, (a + b*Tan[c + d*x])/(a - I*b)] - (I*a + b)*Hypergeometric2F1[-3/2, 1, -1/2, (a + b*Tan[c + d*x])/(a + I*b)]) - 6*(a - I*b)*(a + I*b)*b*(-(A*b) + 4*a*B)*Tan[c + d*x] - 6*(a - I*b)*(a + I*b)*b^2*B*Tan[c + d*x]^2 + 3*A*b^2*(I*(a + I*b)*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Tan[c + d*x])/(a - I*b)] - (I*a + b)*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Tan[c + d*x])/(a + I*b)])*(a + b*Tan[c + d*x])/(b^3*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2))`

3.358.3 Rubi [A] (warning: unable to verify)

Time = 1.45 (sec) , antiderivative size = 298, normalized size of antiderivative = 1.14, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 4088, 27, 3042, 4118, 3042, 4113, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)^3(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

$$\downarrow \text{4088}$$

3.358. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
& 2 \int -\frac{\tan(c+dx)((-4Ba^2+Aba-3b^2B)\tan^2(c+dx)-3b(Ab-aB)\tan(c+dx)+4a(Ab-aB))}{2(a+b\tan(c+dx))^{3/2}} dx \\
& \quad + \frac{3b(a^2+b^2)}{2a(Ab-aB)\tan^2(c+dx)} \\
& \quad \frac{2a(Ab-aB)\tan^2(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} \\
& \quad \downarrow 27 \\
& \quad \frac{2a(Ab-aB)\tan^2(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \\
& \int \frac{\tan(c+dx)((-4Ba^2+Aba-3b^2B)\tan^2(c+dx)-3b(Ab-aB)\tan(c+dx)+4a(Ab-aB))}{(a+b\tan(c+dx))^{3/2}} dx \\
& \quad \frac{3b(a^2+b^2)}{3b(a^2+b^2)} \\
& \quad \downarrow 3042 \\
& \quad \frac{2a(Ab-aB)\tan^2(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \\
& \int \frac{\tan(c+dx)((-4Ba^2+Aba-3b^2B)\tan^2(c+dx)^2-3b(Ab-aB)\tan(c+dx)+4a(Ab-aB))}{(a+b\tan(c+dx))^{3/2}} dx \\
& \quad \frac{3b(a^2+b^2)}{3b(a^2+b^2)} \\
& \quad \downarrow 4118 \\
& \quad \frac{2a(Ab-aB)\tan^2(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \\
& \int \frac{3(Aa^2+2bBa-Ab^2)\tan(c+dx)b^2+(a^2+b^2)(-4Ba^2+Aba-3b^2B)\tan^2(c+dx)+a(-4Ba^3+Ab^2-10b^2Ba+7Ab^3)}{\sqrt{a+b\tan(c+dx)}b(a^2+b^2)} dx \\
& \quad + \frac{2a^2(-4a^3B+a^2Ab-10ab^2B+7Ab^3)}{b^2d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \\
& \quad \frac{3b(a^2+b^2)}{3b(a^2+b^2)} \\
& \quad \downarrow 3042 \\
& \quad \frac{2a(Ab-aB)\tan^2(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \\
& \int \frac{3(Aa^2+2bBa-Ab^2)\tan(c+dx)b^2+(a^2+b^2)(-4Ba^2+Aba-3b^2B)\tan^2(c+dx)+a(-4Ba^3+Ab^2-10b^2Ba+7Ab^3)}{\sqrt{a+b\tan(c+dx)}b(a^2+b^2)} dx \\
& \quad + \frac{2a^2(-4a^3B+a^2Ab-10ab^2B+7Ab^3)}{b^2d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \\
& \quad \frac{3b(a^2+b^2)}{3b(a^2+b^2)} \\
& \quad \downarrow 4113 \\
& \quad \frac{2a(Ab-aB)\tan^2(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \\
& \int \frac{3(-Ba^2+2Aba+b^2B)b^2+3(Aa^2+2bBa-Ab^2)\tan(c+dx)b^2}{\sqrt{a+b\tan(c+dx)}b(a^2+b^2)} dx + \frac{2(a^2+b^2)(-4a^2B+aAb-3b^2B)\sqrt{a+b\tan(c+dx)}}{bd} \\
& \quad + \frac{2a^2(-4a^3B+a^2Ab-10ab^2B+7Ab^3)}{b^2d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \\
& \quad \frac{3b(a^2+b^2)}{3b(a^2+b^2)} \\
& \quad \downarrow 3042
\end{aligned}$$

3.358. $\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

$$\frac{\frac{2a(Ab - aB) \tan^2(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \int \frac{3(-Ba^2 + 2Aba + b^2B)b^2 + 3(Aa^2 + 2bBa - Ab^2) \tan(c + dx)b^2}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2(a^2 + b^2)(-4a^2B + aAb - 3b^2B)}{bd} \sqrt{a + b \tan(c + dx)}}{b(a^2 + b^2)} + \frac{2a^2(-4a^3B + a^2Ab - 10ab^2B + 7Ab^3)}{b^2d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}}$$

↓ 4022

$$\frac{\frac{2a(Ab - aB) \tan^2(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2a^2(-4a^3B + a^2Ab - 10ab^2B + 7Ab^3)}{b^2d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{\frac{3}{2}b^2(a - ib)^2(-B + iA) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx - \frac{3}{2}b^2(a + ib)^2(B + iA) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2(a^2 + b^2)(-4a^2B + aAb - 3b^2B)}{bd} \sqrt{a + b \tan(c + dx)}}{b(a^2 + b^2)}}{3b(a^2 + b^2)}$$

↓ 3042

$$\frac{\frac{2a(Ab - aB) \tan^2(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2a^2(-4a^3B + a^2Ab - 10ab^2B + 7Ab^3)}{b^2d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{\frac{3}{2}b^2(a - ib)^2(-B + iA) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx - \frac{3}{2}b^2(a + ib)^2(B + iA) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx + \frac{2(a^2 + b^2)(-4a^2B + aAb - 3b^2B)}{bd} \sqrt{a + b \tan(c + dx)}}{b(a^2 + b^2)}}{3b(a^2 + b^2)}$$

↓ 4020

$$\frac{\frac{2a(Ab - aB) \tan^2(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2a^2(-4a^3B + a^2Ab - 10ab^2B + 7Ab^3)}{b^2d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{\frac{3ib^2(a + ib)^2(B + iA) \int -\frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} - \frac{3ib^2(a - ib)^2(-B + iA) \int -\frac{1}{(i \tan(c + dx) + 1)\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d}}{b(a^2 + b^2)}}{3b(a^2 + b^2)}$$

↓ 25

$$\frac{\frac{2a(Ab - aB) \tan^2(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2a^2(-4a^3B + a^2Ab - 10ab^2B + 7Ab^3)}{b^2d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{\frac{3ib^2(a + ib)^2(B + iA) \int \frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} + \frac{3ib^2(a - ib)^2(-B + iA) \int \frac{1}{(i \tan(c + dx) + 1)\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d}}{b(a^2 + b^2)}}{3b(a^2 + b^2)}$$

↓ 73

$$\frac{\frac{2a(Ab - aB) \tan^2(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2a^2(-4a^3B + a^2Ab - 10ab^2B + 7Ab^3)}{b^2d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{\frac{3b(a - ib)^2(-B + iA) \int -\frac{1}{\frac{i \tan^2(c + dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)}}{d} - \frac{3b(a + ib)^2(B + iA) \int \frac{1}{\frac{i \tan^2(c + dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)}}{d}}{b(a^2 + b^2)}}{3b(a^2 + b^2)}$$

↓ 221

3.358. $\int \frac{\tan^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx$

$$\frac{2a(Ab - aB) \tan^2(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2(a^2 + b^2)(-4a^2B + aAb - 3b^2B)\sqrt{a + b \tan(c + dx)}}{bd} - \frac{3b^2(a + ib)^2(B + iA) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a - ib}}\right)}{d\sqrt{a - ib}} + \frac{3b^2(a - ib)^2(-B + iA)}{d\sqrt{a - ib}}$$

$$\frac{2a^2(-4a^3B + a^2Ab - 10ab^2B + 7Ab^3)}{b^2d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{2(a^2 + b^2)(-4a^2B + aAb - 3b^2B)\sqrt{a + b \tan(c + dx)}}{bd} - \frac{3b^2(a + ib)^2(B + iA) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a - ib}}\right)}{d\sqrt{a - ib}} + \frac{3b^2(a - ib)^2(-B + iA)}{d\sqrt{a - ib}}$$

$$3b(a^2 + b^2)$$

input `Int[(Tan[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]`

output `(2*a*(A*b - a*B)*Tan[c + d*x]^2)/(3*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) - ((2*a^2*(a^2*A*b + 7*A*b^3 - 4*a^3*B - 10*a*b^2*B))/(b^2*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]) + ((-3*(a + I*b)^2*b^2*(I*A + B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d) + (3*(a - I*b)^2*b^2*(I*A - B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d) + (2*(a^2 + b^2)*(a*A*b - 4*a^2*B - 3*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(b*d))/(b*(a^2 + b^2))/(3*b*(a^2 + b^2))`

3.358.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4118 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c + d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`

3.358.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4547 vs. $2(231) = 462$.

Time = 0.14 (sec) , antiderivative size = 4548, normalized size of antiderivative = 17.43

method	result	size
parts	Expression too large to display	4548
derivativedivides	Expression too large to display	12907
default	Expression too large to display	12907

```
input int(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output A*(-6/d*a^2/(a^2+b^2)^2/(a+b*tan(d*x+c))^(1/2)+2/3/d/b^2*a^3/(a^2+b^2)/(a+
b*tan(d*x+c))^(3/2)+1/d/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arct
an(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(
1/2)-2*a)^(1/2))*a^5+3/4/d/(a^2+b^2)^(7/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+
c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2
)+2*a)^(1/2)*a^4-1/2/d/(a^2+b^2)^3*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2
))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(
1/2)*a^3-1/d/(a^2+b^2)^3/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(
d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2
))*a^4+1/2/d/(a^2+b^2)^3*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(
1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3+1/
d/(a^2+b^2)^3/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a
)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^4+2/d/(
a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1
/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^3-1/d/
(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(
1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^5-3/4
/d/(a^2+b^2)^(7/2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2
)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^4-1/4/d*b
^4/(a^2+b^2)^(7/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)...
```

3.358.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7284 vs. $2(227) = 454$.

Time = 2.20 (sec) , antiderivative size = 7284, normalized size of antiderivative = 27.91

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="fricas")`

output Too large to include

3.358.6 Sympy [F]

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\tan^3(c+dx)}{(a+b\tan(c+dx))^{5/2}} dx$$

input `integrate(tan(d*x+c)**3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**3/(a + b*tan(c + d*x))**(5/2), x)`

3.358.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output Timed out

3.358.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

```
input integrate(tan(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm
m="giac")
```

```
output Timed out
```

3.358.9 Mupad [B] (verification not implemented)

Time = 37.09 (sec) , antiderivative size = 9498, normalized size of antiderivative = 36.39

$$\int \frac{\tan^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

```
input int((tan(c + d*x)^3*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)
```

```
output (log((((320*B^4*a^2*b^8*d^4 - 16*B^4*b^10*d^4 - 1760*B^4*a^4*b^6*d^4 + 16
00*B^4*a^6*b^4*d^4 - 400*B^4*a^8*b^2*d^4)^(1/2) - 4*B^2*a^5*d^2 + 40*B^2*a
^3*b^2*d^2 - 20*B^2*a*b^4*d^2)/(a^10*d^4 + b^10*d^4 + 5*a^2*b^8*d^4 + 10*a
^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^(1/2)*((a + b*tan(c + d*x))^(
1/2)*(320*B^2*a^4*b^14*d^3 - 16*B^2*b^18*d^3 + 1024*B^2*a^6*b^12*d^3 + 14
40*B^2*a^8*b^10*d^3 + 1024*B^2*a^10*b^8*d^3 + 320*B^2*a^12*b^6*d^3 - 16*B^
2*a^16*b^2*d^3) - (((320*B^4*a^2*b^8*d^4 - 16*B^4*b^10*d^4 - 1760*B^4*a^4
*b^6*d^4 + 1600*B^4*a^6*b^4*d^4 - 400*B^4*a^8*b^2*d^4)^(1/2) - 4*B^2*a^5*d
^2 + 40*B^2*a^3*b^2*d^2 - 20*B^2*a*b^4*d^2)/(a^10*d^4 + b^10*d^4 + 5*a^2*b
^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^(1/2)*((((320*
B^4*a^2*b^8*d^4 - 16*B^4*b^10*d^4 - 1760*B^4*a^4*b^6*d^4 + 1600*B^4*a^6*b^
4*d^4 - 400*B^4*a^8*b^2*d^4)^(1/2) - 4*B^2*a^5*d^2 + 40*B^2*a^3*b^2*d^2 -
20*B^2*a*b^4*d^2)/(a^10*d^4 + b^10*d^4 + 5*a^2*b^8*d^4 + 10*a^4*b^6*d^4 +
10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^(1/2)*(a + b*tan(c + d*x))^(1/2)*(64*a*b^
22*d^5 + 640*a^3*b^20*d^5 + 2880*a^5*b^18*d^5 + 7680*a^7*b^16*d^5 + 13440*
a^9*b^14*d^5 + 16128*a^11*b^12*d^5 + 13440*a^13*b^10*d^5 + 7680*a^15*b^8*d
^5 + 2880*a^17*b^6*d^5 + 640*a^19*b^4*d^5 + 64*a^21*b^2*d^5))/4 - 32*B*b^2
1*d^4 - 160*B*a^2*b^19*d^4 - 128*B*a^4*b^17*d^4 + 896*B*a^6*b^15*d^4 + 313
6*B*a^8*b^13*d^4 + 4928*B*a^10*b^11*d^4 + 4480*B*a^12*b^9*d^4 + 2432*B*a^1
4*b^7*d^4 + 736*B*a^16*b^5*d^4 + 96*B*a^18*b^3*d^4))/4 + 96*B^3*a^3...
```

3.358. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

3.359
$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

3.359.1 Optimal result 3418
 3.359.2 Mathematica [C] (verified) 3418
 3.359.3 Rubi [A] (warning: unable to verify) 3419
 3.359.4 Maple [B] (verified) 3422
 3.359.5 Fricas [B] (verification not implemented) 3423
 3.359.6 Sympy [F] 3424
 3.359.7 Maxima [F(-1)] 3424
 3.359.8 Giac [F(-1)] 3424
 3.359.9 Mupad [B] (verification not implemented) 3425

3.359.1 Optimal result

Integrand size = 33, antiderivative size = 198

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{5/2}d} - \frac{(iA-B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a+ib)^{5/2}d} - \frac{2a^2(Ab-aB)}{3b^2(a^2+b^2)d(a+b \tan(c+dx))^{3/2}} + \frac{2a(2Ab^3-a(a^2+3b^2)B)}{b^2(a^2+b^2)^2d\sqrt{a+b \tan(c+dx)}}$$

```
output (I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(5/2)/d-(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(5/2)/d+2*a*(2*A*b^3-a*(a^2+3*b^2)*B)/b^2/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^(1/2)-2/3*a^2*(A*b-B*a)/b^2/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)
```

3.359.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.16 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.31

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{2(a-ib)(a+ib)(Ab+2aB)+b(Ab-aB)\left(i(a+ib)\operatorname{Hypergeometric2F1}\left(-\frac{3}{2},1,-\frac{1}{2},\frac{a+b \tan(c+dx)}{a-ib}\right)-i(a-ib)\operatorname{Hypergeometric2F1}\left(-\frac{3}{2},1,-\frac{1}{2},\frac{a+b \tan(c+dx)}{a+ib}\right)\right)}{(a^2+b^2)^2d\sqrt{a+b \tan(c+dx)}}$$

3.359.
$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

input `Integrate[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]`

output `-1/3*(2*(a - I*b)*(a + I*b)*(A*b + 2*a*B) + b*(A*b - a*B)*(I*(a + I*b)*Hypergeometric2F1[-3/2, 1, -1/2, (a + b*Tan[c + d*x])/(a - I*b)] - (I*a + b)*Hypergeometric2F1[-3/2, 1, -1/2, (a + b*Tan[c + d*x])/(a + I*b)]) + 6*(a - I*b)*(a + I*b)*b*B*Tan[c + d*x] + 3*b*B*(I*(a + I*b)*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Tan[c + d*x])/(a - I*b)] - (I*a + b)*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Tan[c + d*x])/(a + I*b)])*(a + b*Tan[c + d*x]))/(b^2*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2))`

3.359.3 Rubi [A] (warning: unable to verify)

Time = 1.08 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.14, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3042, 4087, 25, 3042, 4111, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^2(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{4087} \\
 & \frac{\int \frac{-((a^2+b^2)B \tan^2(c+dx))-b(Ab-aB) \tan(c+dx)+a(Ab-aB)}{(a+b \tan(c+dx))^{3/2}} dx}{b(a^2+b^2)} - \frac{2a^2(Ab-aB)}{3b^2d(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{-((a^2+b^2)B \tan^2(c+dx))-b(Ab-aB) \tan(c+dx)+a(Ab-aB)}{(a+b \tan(c+dx))^{3/2}} dx}{b(a^2+b^2)} - \frac{2a^2(Ab-aB)}{3b^2d(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{-((a^2+b^2)B \tan(c+dx)^2)-b(Ab-aB) \tan(c+dx)+a(Ab-aB)}{(a+b \tan(c+dx))^{3/2}} dx}{b(a^2+b^2)} - \frac{2a^2(Ab-aB)}{3b^2d(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow \text{4111}
 \end{aligned}$$

3.359. $\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{\int \frac{b(Aa^2+2bBa-Ab^2)-b(-Ba^2+2Aba+b^2B)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx - \frac{2a(2Ab^3-aB(a^2+3b^2))}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{\frac{b(a^2+b^2)}{2a^2(Ab-aB)} - \frac{3b^2d(a^2+b^2)(a+b\tan(c+dx))^{3/2}}{3b^2d(a^2+b^2)(a+b\tan(c+dx))^{3/2}}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{b(Aa^2+2bBa-Ab^2)-b(-Ba^2+2Aba+b^2B)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx - \frac{2a(2Ab^3-aB(a^2+3b^2))}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{\frac{b(a^2+b^2)}{2a^2(Ab-aB)} - \frac{3b^2d(a^2+b^2)(a+b\tan(c+dx))^{3/2}}{3b^2d(a^2+b^2)(a+b\tan(c+dx))^{3/2}}} \\
 & \quad \downarrow \text{4022} \\
 & \frac{-\frac{2a(2Ab^3-aB(a^2+3b^2))}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{\frac{1}{2}b(a-ib)^2(A+iB)\int \frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}b(a+ib)^2(A-iB)\int \frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}} dx}{a^2+b^2}}{\frac{b(a^2+b^2)}{2a^2(Ab-aB)} - \frac{3b^2d(a^2+b^2)(a+b\tan(c+dx))^{3/2}}{3b^2d(a^2+b^2)(a+b\tan(c+dx))^{3/2}}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{-\frac{2a(2Ab^3-aB(a^2+3b^2))}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{\frac{1}{2}b(a-ib)^2(A+iB)\int \frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}b(a+ib)^2(A-iB)\int \frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}} dx}{a^2+b^2}}{\frac{b(a^2+b^2)}{2a^2(Ab-aB)} - \frac{3b^2d(a^2+b^2)(a+b\tan(c+dx))^{3/2}}{3b^2d(a^2+b^2)(a+b\tan(c+dx))^{3/2}}} \\
 & \quad \downarrow \text{4020} \\
 & \frac{-\frac{2a(2Ab^3-aB(a^2+3b^2))}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{\frac{ib(a+ib)^2(A-iB)\int -\frac{1}{(1-i\tan(c+dx))\sqrt{a+b\tan(c+dx)}} d(i\tan(c+dx))}{2d} - \frac{ib(a-ib)^2(A+iB)\int -\frac{1}{(i\tan(c+dx)+1)\sqrt{a+b\tan(c+dx)}} d(i\tan(c+dx))}{2d}}{a^2+b^2}}{\frac{b(a^2+b^2)}{2a^2(Ab-aB)} - \frac{3b^2d(a^2+b^2)(a+b\tan(c+dx))^{3/2}}{3b^2d(a^2+b^2)(a+b\tan(c+dx))^{3/2}}} \\
 & \quad \downarrow \text{25} \\
 & \frac{-\frac{2a(2Ab^3-aB(a^2+3b^2))}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{\frac{ib(a-ib)^2(A+iB)\int \frac{1}{(i\tan(c+dx)+1)\sqrt{a+b\tan(c+dx)}} d(-i\tan(c+dx))}{2d} - \frac{ib(a+ib)^2(A-iB)\int \frac{1}{(1-i\tan(c+dx))\sqrt{a+b\tan(c+dx)}} d(i\tan(c+dx))}{2d}}{a^2+b^2}}{\frac{b(a^2+b^2)}{2a^2(Ab-aB)} - \frac{3b^2d(a^2+b^2)(a+b\tan(c+dx))^{3/2}}{3b^2d(a^2+b^2)(a+b\tan(c+dx))^{3/2}}} \\
 & \quad \downarrow \text{73}
 \end{aligned}$$

3.359. $\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{2a^2(Ab - aB)}{3b^2d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{(a - ib)^2(A + iB) \int \frac{1}{-i \tan^2 \frac{(c + dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)}}{d\sqrt{a + b \tan(c + dx)}} + \frac{(a + ib)^2(A - iB) \int \frac{1}{i \tan^2 \frac{(c + dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)}}{d\sqrt{a + b \tan(c + dx)}} \\
 & - \frac{2a(2Ab^3 - aB(a^2 + 3b^2))}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{2a^2(Ab - aB)}{b(a^2 + b^2)} \\
 & \quad \downarrow \text{221} \\
 & \frac{2a^2(Ab - aB)}{3b^2d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{b(a - ib)^2(A + iB) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a + ib}}\right)}{d\sqrt{a + ib}} + \frac{b(a + ib)^2(A - iB) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a - ib}}\right)}{d\sqrt{a - ib}} \\
 & - \frac{2a(2Ab^3 - aB(a^2 + 3b^2))}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{2a^2(Ab - aB)}{b(a^2 + b^2)}
 \end{aligned}$$

```
input Int[(Tan[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]
```

```
output (-2*a^2*(A*b - a*B))/(3*b^2*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) - ((
((a + I*b)^2*b*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b
]*d) + ((a - I*b)^2*b*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[
a + I*b]*d))/(a^2 + b^2) - (2*a*(2*A*b^3 - a*(a^2 + 3*b^2)*B))/(b*(a^2 + b
^2)*d*Sqrt[a + b*Tan[c + d*x]]))/(b*(a^2 + b^2))
```

3.359.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 73 Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

3.359. $\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx$

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4087 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(- (B*c - A*d)*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`

rule 4111 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.359.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4491 vs. $2(174) = 348$.

Time = 0.12 (sec) , antiderivative size = 4492, normalized size of antiderivative = 22.69

method	result	size
parts	Expression too large to display	4492
derivativedivides	Expression too large to display	12849
default	Expression too large to display	12849

3.359.
$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

input `int(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x,method=_RETURNV
ERBOSE)`

output `A*(3/4/d*b^3/(a^2+b^2)^(7/2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+
2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a
-1/d/b/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(
1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*
a^4+1/d/b/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^
2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2
)*)a^6+4/d*b/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2
+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(
1/2))*a^4+1/d*b^3/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2
*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2
*a)^(1/2))*a^2+1/4/d/b/(a^2+b^2)^(7/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1
/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*
a)^(1/2)*a^5-1/2/d*b/(a^2+b^2)^(7/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1
/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a
^(1/2))*a^3-3/4/d*b^3/(a^2+b^2)^(7/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1
/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a
^(1/2))*a+1/d/b/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+
b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)
^(1/2))*a^4-1/d/b/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*
(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)...`

3.359.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7218 vs. 2(168) = 336.

Time = 2.23 (sec) , antiderivative size = 7218, normalized size of antiderivative = 36.45

$$\int \frac{\tan^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm
m="fricas")`

output Too large to include

3.359.6 Sympy [F]

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\tan^2(c+dx)}{(a+b\tan(c+dx))^{5/2}} dx$$

input `integrate(tan(d*x+c)**2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**2/(a + b*tan(c + d*x))**(5/2), x)`

3.359.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `Timed out`

3.359.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Timed out`

3.359.9 Mupad [B] (verification not implemented)

Time = 27.68 (sec) , antiderivative size = 9468, normalized size of antiderivative = 47.82

$$\int \frac{\tan^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

```
input int((tan(c + d*x)^2*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)
```

```
output (log((((a + b*tan(c + d*x))^(1/2)*(320*A^2*a^4*b^14*d^3 - 16*A^2*b^18*d^3
+ 1024*A^2*a^6*b^12*d^3 + 1440*A^2*a^8*b^10*d^3 + 1024*A^2*a^10*b^8*d^3 +
320*A^2*a^12*b^6*d^3 - 16*A^2*a^16*b^2*d^3) + (((320*A^4*a^2*b^8*d^4 - 16
*A^4*b^10*d^4 - 1760*A^4*a^4*b^6*d^4 + 1600*A^4*a^6*b^4*d^4 - 400*A^4*a^8
*b^2*d^4)^(1/2) - 4*A^2*a^5*d^2 + 40*A^2*a^3*b^2*d^2 - 20*A^2*a*b^4*d^2)/(a
^10*d^4 + b^10*d^4 + 5*a^2*b^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a
^8*b^2*d^4))^(1/2)*(896*A*a^6*b^15*d^4 - 32*A*b^21*d^4 - 160*A*a^2*b^19*d
4 - 128*A*a^4*b^17*d^4 - (((320*A^4*a^2*b^8*d^4 - 16*A^4*b^10*d^4 - 1760*
A^4*a^4*b^6*d^4 + 1600*A^4*a^6*b^4*d^4 - 400*A^4*a^8*b^2*d^4)^(1/2) - 4*A
^2*a^5*d^2 + 40*A^2*a^3*b^2*d^2 - 20*A^2*a*b^4*d^2)/(a^10*d^4 + b^10*d^4 +
5*a^2*b^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^(1/2)*(a
+ b*tan(c + d*x))^(1/2)*(64*a*b^22*d^5 + 640*a^3*b^20*d^5 + 2880*a^5*b^18
*d^5 + 7680*a^7*b^16*d^5 + 13440*a^9*b^14*d^5 + 16128*a^11*b^12*d^5 + 1344
0*a^13*b^10*d^5 + 7680*a^15*b^8*d^5 + 2880*a^17*b^6*d^5 + 640*a^19*b^4*d^5
+ 64*a^21*b^2*d^5))/4 + 3136*A*a^8*b^13*d^4 + 4928*A*a^10*b^11*d^4 + 4480
*A*a^12*b^9*d^4 + 2432*A*a^14*b^7*d^4 + 736*A*a^16*b^5*d^4 + 96*A*a^18*b^3
*d^4))/4)*(((320*A^4*a^2*b^8*d^4 - 16*A^4*b^10*d^4 - 1760*A^4*a^4*b^6*d^4
+ 1600*A^4*a^6*b^4*d^4 - 400*A^4*a^8*b^2*d^4)^(1/2) - 4*A^2*a^5*d^2 + 40*A
^2*a^3*b^2*d^2 - 20*A^2*a*b^4*d^2)/(a^10*d^4 + b^10*d^4 + 5*a^2*b^8*d^4 +
10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^(1/2))/4 - 96*A^3*a^3...
```


3.360
$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

3.360.1 Optimal result 3426
 3.360.2 Mathematica [A] (verified) 3426
 3.360.3 Rubi [A] (warning: unable to verify) 3427
 3.360.4 Maple [B] (verified) 3430
 3.360.5 Fricas [B] (verification not implemented) 3431
 3.360.6 Sympy [F] 3432
 3.360.7 Maxima [F(-1)] 3432
 3.360.8 Giac [F(-1)] 3432
 3.360.9 Mupad [B] (verification not implemented) 3433

3.360.1 Optimal result

Integrand size = 31, antiderivative size = 188

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx =$$

$$-\frac{(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{5/2}d} - \frac{(A+iB)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a+ib)^{5/2}d}$$

$$+ \frac{2a(Ab-aB)}{3b(a^2+b^2)d(a+b \tan(c+dx))^{3/2}} + \frac{2(a^2A-Ab^2+2abB)}{(a^2+b^2)^2d\sqrt{a+b \tan(c+dx)}}$$

output

```
-(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(5/2)/d-(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(5/2)/d+2*(A*a^2-A*b^2+2*B*a*b)/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^(1/2)+2/3*a*(A*b-B*a)/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)
```

3.360.2 Mathematica [A] (verified)

Time = 3.87 (sec) , antiderivative size = 325, normalized size of antiderivative = 1.73

$$\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{3b(-a^2(A\sqrt{-b^2+bB})+b^2(A\sqrt{-b^2+bB})+2ab(Ab-\sqrt{-b^2}B))\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-\sqrt{-b^2}}}\right)}{\sqrt{-b^2}\sqrt{a-\sqrt{-b^2}}}$$

input `Integrate[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]`

output
$$\frac{((3*b*(-(a^2*(A*\sqrt{-b^2} + b*B)) + b^2*(A*\sqrt{-b^2} + b*B) + 2*a*b*(A*b - \sqrt{-b^2}*B))*\text{ArcTanh}[\sqrt{a + b*\text{Tan}[c + d*x]}/\sqrt{a - \sqrt{-b^2}}])}{(\sqrt{-b^2}*\sqrt{a - \sqrt{-b^2}}) - (3*b*(2*a*A*b^2 + a^2*A*\sqrt{-b^2} + A*(-b^2)^{3/2} - a^2*b*B + b^3*B + 2*a*b*\sqrt{-b^2}*B)*\text{ArcTanh}[\sqrt{a + b*\text{Tan}[c + d*x]}/\sqrt{a + \sqrt{-b^2}}])}{(\sqrt{-b^2}*\sqrt{a + \sqrt{-b^2}}) + (2*a*(a^2 + b^2)*(A*b - a*B))/(a + b*\text{Tan}[c + d*x])^{3/2} + (6*b*(a^2*A - A*b^2 + 2*a*b*B))/\sqrt{a + b*\text{Tan}[c + d*x]}}/(3*b*(a^2 + b^2)^2*d)$$

3.360.3 Rubi [A] (warning: unable to verify)

Time = 0.93 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.13, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4074, 3042, 4012, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{4074} \\ & \frac{\int \frac{Ab-aB+(aA+bB) \tan(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx}{a^2+b^2} + \frac{2a(Ab-aB)}{3bd(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\ & \quad \downarrow \text{3042} \\ & \frac{\int \frac{Ab-aB+(aA+bB) \tan(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx}{a^2+b^2} + \frac{2a(Ab-aB)}{3bd(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\ & \quad \downarrow \text{4012} \\ & \frac{\int \frac{-Ba^2+2Aba+b^2B+(Aa^2+2bBa-Ab^2) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{a^2+b^2} + \frac{2(a^2A+2abB-Ab^2)}{d(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \\ & \quad \frac{2a(Ab-aB)}{3bd(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \end{aligned}$$

3.360. $\int \frac{\tan(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \int \frac{-Ba^2+2Aba+b^2B+(Aa^2+2bBa-Ab^2)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}\frac{a^2+b^2}{a^2+b^2}} dx + \frac{2(a^2A+2abB-Ab^2)}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \\
 & \frac{a^2+b^2}{2a(Ab-aB)} \\
 & \frac{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}}{\downarrow 3042} \\
 & \frac{2a(Ab-aB)}{\downarrow 4022} \\
 & \frac{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}}{+} + \frac{2(a^2A+2abB-Ab^2)}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \\
 & \frac{\frac{1}{2}(a-ib)^2(-B+iA)\int\frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}}dx - \frac{1}{2}(a+ib)^2(B+iA)\int\frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}}dx}{a^2+b^2} \\
 & \frac{a^2+b^2}{\downarrow 3042} \\
 & \frac{2a(Ab-aB)}{\downarrow 4020} \\
 & \frac{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}}{+} + \frac{2(a^2A+2abB-Ab^2)}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \\
 & \frac{\frac{1}{2}(a-ib)^2(-B+iA)\int\frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}}dx - \frac{1}{2}(a+ib)^2(B+iA)\int\frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}}dx}{a^2+b^2} \\
 & \frac{a^2+b^2}{\downarrow 4020} \\
 & \frac{2a(Ab-aB)}{\downarrow 25} \\
 & \frac{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}}{+} + \frac{2(a^2A+2abB-Ab^2)}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \\
 & \frac{\frac{i(a-ib)^2(-B+iA)\int\frac{1-i\tan(c+dx)+1}{2d\sqrt{a+b\tan(c+dx)}}d(-i\tan(c+dx)) - \frac{i(a+ib)^2(B+iA)\int\frac{1}{(1-i\tan(c+dx))\sqrt{a+b\tan(c+dx)}}d}{a^2+b^2}}{a^2+b^2} \\
 & \frac{a^2+b^2}{\downarrow 73} \\
 & \frac{2a(Ab-aB)}{\downarrow 221} \\
 & \frac{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}}{+} + \frac{2(a^2A+2abB-Ab^2)}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \\
 & \frac{(a-ib)^2(-B+iA)\int\frac{1}{b}\frac{1-i\tan^2(c+dx)}{bd}d\sqrt{a+b\tan(c+dx)} - \frac{(a+ib)^2(B+iA)\int\frac{1}{b}\frac{1+i\tan^2(c+dx)}{bd}d\sqrt{a+b\tan(c+dx)}}{a^2+b^2}
 \end{aligned}$$

3.360. $\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

$$\frac{2(a^2A+2abB-Ab^2)}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{2a(Ab-aB)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{(a-ib)^2(-B+iA)\arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} - \frac{(a+ib)^2(B+iA)\arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}}$$

$$a^2 + b^2$$

input `Int[(Tan[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]`

output `(2*a*(A*b - a*B))/(3*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) + ((-(((a + I*b)^2*(I*A + B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d)) + ((a - I*b)^2*(I*A - B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d))/(a^2 + b^2) + (2*(a^2*A - A*b^2 + 2*a*b*B))/((a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])/(a^2 + b^2)`

3.360.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

3.360.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4483 vs. $2(164) = 328$.

Time = 0.13 (sec) , antiderivative size = 4484, normalized size of antiderivative = 23.85

method	result	size
parts	Expression too large to display	4484
derivativedivides	Expression too large to display	12841
default	Expression too large to display	12841

```
input int(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x,method=_RETURNVER
BOSE)
```

output $A*(2/d*a^2/(a^2+b^2)^2/(a+b*\tan(d*x+c))^{(1/2)}-1/d/(a^2+b^2)^{(7/2)}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan(((2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-2*(a+b*\tan(d*x+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*a^{5-3/4}/d/(a^2+b^2)^{(7/2)}*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}+(a^2+b^2)^{(1/2)})*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*a^{4+1/2}/d/(a^2+b^2)^3*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}+(a^2+b^2)^{(1/2)}))*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*a^{3+1/d}/(a^2+b^2)^3/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan((2*(a+b*\tan(d*x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*a^{4-1/2}/d/(a^2+b^2)^3*\ln((a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-b*\tan(d*x+c)-a-(a^2+b^2)^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*a^{3-1/d}/(a^2+b^2)^3/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan(((2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-2*(a+b*\tan(d*x+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*a^{4-2/d}/(a^2+b^2)^{(5/2)}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan((2*(a+b*\tan(d*x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*a^{3+1/d}/(a^2+b^2)^{(7/2)}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan((2*(a+b*\tan(d*x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*a^{5+3/4}/d/(a^2+b^2)^{(7/2)}*\ln((a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-b*\tan(d*x+c)-a-(a^2+b^2)^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*a^{4+1/4}/d*b^4/(a^2+b^2)^{(7/2)}*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}+(a^2+b^2)^{(1/2)}*(2*(a^2+b^2)^{...$

3.360.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7209 vs. $2(158) = 316$.

Time = 2.26 (sec) , antiderivative size = 7209, normalized size of antiderivative = 38.35

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="fricas")`

output Too large to include

3.360.6 Sympy [F]

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\tan(c+dx)}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)/(a + b*tan(c + d*x))**(5/2), x)`

3.360.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `Timed out`

3.360.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Timed out`

3.360.9 Mupad [B] (verification not implemented)

Time = 26.56 (sec) , antiderivative size = 9464, normalized size of antiderivative = 50.34

$$\int \frac{\tan(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

```
input int((tan(c + d*x)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)
```

```
output (log((((320*B^4*a^2*b^8*d^4 - 16*B^4*b^10*d^4 - 1760*B^4*a^4*b^6*d^4 + 16
00*B^4*a^6*b^4*d^4 - 400*B^4*a^8*b^2*d^4)^(1/2) - 4*B^2*a^5*d^2 + 40*B^2*a
^3*b^2*d^2 - 20*B^2*a*b^4*d^2)/(a^10*d^4 + b^10*d^4 + 5*a^2*b^8*d^4 + 10*a
^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^(1/2)*((a + b*tan(c + d*x))^(
1/2)*(320*B^2*a^4*b^14*d^3 - 16*B^2*b^18*d^3 + 1024*B^2*a^6*b^12*d^3 + 14
40*B^2*a^8*b^10*d^3 + 1024*B^2*a^10*b^8*d^3 + 320*B^2*a^12*b^6*d^3 - 16*B^
2*a^16*b^2*d^3) + (((320*B^4*a^2*b^8*d^4 - 16*B^4*b^10*d^4 - 1760*B^4*a^4
*b^6*d^4 + 1600*B^4*a^6*b^4*d^4 - 400*B^4*a^8*b^2*d^4)^(1/2) - 4*B^2*a^5*d
^2 + 40*B^2*a^3*b^2*d^2 - 20*B^2*a*b^4*d^2)/(a^10*d^4 + b^10*d^4 + 5*a^2*b
^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^(1/2)*(896*B*a^
6*b^15*d^4 - 32*B*b^21*d^4 - 160*B*a^2*b^19*d^4 - 128*B*a^4*b^17*d^4 - (((
(320*B^4*a^2*b^8*d^4 - 16*B^4*b^10*d^4 - 1760*B^4*a^4*b^6*d^4 + 1600*B^4*a
^6*b^4*d^4 - 400*B^4*a^8*b^2*d^4)^(1/2) - 4*B^2*a^5*d^2 + 40*B^2*a^3*b^2*d
^2 - 20*B^2*a*b^4*d^2)/(a^10*d^4 + b^10*d^4 + 5*a^2*b^8*d^4 + 10*a^4*b^6*d
^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^(1/2)*(a + b*tan(c + d*x))^(1/2)*(64
*a*b^22*d^5 + 640*a^3*b^20*d^5 + 2880*a^5*b^18*d^5 + 7680*a^7*b^16*d^5 + 1
3440*a^9*b^14*d^5 + 16128*a^11*b^12*d^5 + 13440*a^13*b^10*d^5 + 7680*a^15*
b^8*d^5 + 2880*a^17*b^6*d^5 + 640*a^19*b^4*d^5 + 64*a^21*b^2*d^5))/4 + 313
6*B*a^8*b^13*d^4 + 4928*B*a^10*b^11*d^4 + 4480*B*a^12*b^9*d^4 + 2432*B*a^1
4*b^7*d^4 + 736*B*a^16*b^5*d^4 + 96*B*a^18*b^3*d^4))/4))/4 - 96*B^3*a^3...
```


3.361 $\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^{5/2}} dx$

3.361.1 Optimal result	3434
3.361.2 Mathematica [C] (verified)	3434
3.361.3 Rubi [A] (warning: unable to verify)	3435
3.361.4 Maple [B] (verified)	3438
3.361.5 Fricas [B] (verification not implemented)	3439
3.361.6 Sympy [F]	3440
3.361.7 Maxima [F(-2)]	3440
3.361.8 Giac [F(-1)]	3440
3.361.9 Mupad [B] (verification not implemented)	3441

3.361.1 Optimal result

Integrand size = 25, antiderivative size = 185

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = -\frac{(iA + B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a - ib)^{5/2}d} + \frac{(iA - B)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a + ib)^{5/2}d} - \frac{2(Ab - aB)}{3(a^2 + b^2)d(a + b \tan(c + dx))^{3/2}} - \frac{2(2aAb - a^2B + b^2B)}{(a^2 + b^2)^2 d \sqrt{a + b \tan(c + dx)}}$$

```
output - (I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(5/2)/d+(I*
A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(5/2)/d-2*(2*A*
a*b-B*a^2+B*b^2)/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^(1/2)-2/3*(A*b-B*a)/(a^2+b
^2)/d/(a+b*tan(d*x+c))^(3/2)
```

3.361.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.27 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.62

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \frac{i \left(-\frac{(A-ib) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{a+b \tan(c+dx)}{a-ib}\right)}{a-ib} + \frac{(A+ib) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{a+b \tan(c+dx)}{a+ib}\right)}{a+ib} \right)}{3d(a + b \tan(c + dx))^{3/2}}$$

input `Integrate[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^(5/2),x]`

output `((-1/3*I)*(-(((A - I*B)*Hypergeometric2F1[-3/2, 1, -1/2, (a + b*Tan[c + d*x])/(a - I*b)])/(a - I*b)))/(a - I*b)) + ((A + I*B)*Hypergeometric2F1[-3/2, 1, -1/2, (a + b*Tan[c + d*x])/(a + I*b)])/(a + I*b))/(d*(a + b*Tan[c + d*x])^(3/2))`

3.361.3 Rubi [A] (warning: unable to verify)

Time = 0.93 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.11, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.440$, Rules used = {3042, 4012, 3042, 4012, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{4012} \\
 & \frac{\int \frac{aA + bB - (Ab - aB) \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx}{a^2 + b^2} - \frac{2(Ab - aB)}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{aA + bB - (Ab - aB) \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx}{a^2 + b^2} - \frac{2(Ab - aB)}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{4012} \\
 & \frac{\int \frac{Aa^2 + 2bBa - Ab^2 - (-Ba^2 + 2Aba + b^2B) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} - \frac{2(a^2(-B) + 2aAb + b^2B)}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a^2 + b^2}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2(Ab - aB)}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{\int \frac{Aa^2+2bBa-Ab^2 - (-Ba^2+2Aba+b^2B) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{a^2+b^2} - \frac{2(a^2(-B)+2aAb+b^2B)}{d(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \frac{a^2+b^2}{2(Ab-aB)} \\
 & \quad \frac{2(Ab-aB)}{3d(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 4022 \\
 & - \frac{2(Ab-aB)}{3d(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \\
 & \frac{2(a^2(-B)+2aAb+b^2B)}{d(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{\frac{1}{2}(a-ib)^2(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}(a+ib)^2(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx}{a^2+b^2} \\
 & \quad \frac{a^2+b^2}{a^2+b^2} \\
 & \quad \downarrow 3042 \\
 & - \frac{2(Ab-aB)}{3d(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \\
 & \frac{2(a^2(-B)+2aAb+b^2B)}{d(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{\frac{1}{2}(a-ib)^2(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}(a+ib)^2(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx}{a^2+b^2} \\
 & \quad \frac{a^2+b^2}{a^2+b^2} \\
 & \quad \downarrow 4020 \\
 & - \frac{2(Ab-aB)}{3d(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \\
 & \frac{2(a^2(-B)+2aAb+b^2B)}{d(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{i(a+ib)^2(A-iB) \int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx)) - i(a-ib)^2(A+iB) \int -\frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}}}{a^2+b^2} \\
 & \quad \frac{a^2+b^2}{a^2+b^2} \\
 & \quad \downarrow 25 \\
 & - \frac{2(Ab-aB)}{3d(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \\
 & \frac{2(a^2(-B)+2aAb+b^2B)}{d(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{i(a-ib)^2(A+iB) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx)) - i(a+ib)^2(A-iB) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}}}{a^2+b^2} \\
 & \quad \frac{a^2+b^2}{a^2+b^2} \\
 & \quad \downarrow 73 \\
 & - \frac{2(Ab-aB)}{3d(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \\
 & \frac{2(a^2(-B)+2aAb+b^2B)}{d(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a-ib)^2(A+iB) \int \frac{1}{-i \tan^2(c+dx) - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)} - (a+ib)^2(A-iB) \int \frac{1}{i \tan^2(c+dx) + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{a^2+b^2} \\
 & \quad \frac{a^2+b^2}{a^2+b^2} \\
 & \quad \downarrow 221
 \end{aligned}$$

3.361. $\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^{5/2}} dx$

$$-\frac{2(Ab - aB)}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{\frac{2(a^2(-B) + 2aAb + b^2B)}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{(a - ib)^2(A + iB) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a + ib}}\right)}{d\sqrt{a + ib}} + \frac{(a + ib)^2(A - iB) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a - ib}}\right)}{d\sqrt{a - ib}}}{a^2 + b^2}$$

input `Int[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^(5/2), x]`

output `(-2*(A*b - a*B))/(3*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) + (((a + I*b)^2*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]]/(Sqrt[a - I*b]*d) + ((a - I*b)^2*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]]/(Sqrt[a + I*b]*d))/(a^2 + b^2) - (2*(2*a*A*b - a^2*B + b^2*B))/((a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])/(a^2 + b^2)`

3.361.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

3.361.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4472 vs. $2(161) = 322$.

Time = 0.11 (sec) , antiderivative size = 4473, normalized size of antiderivative = 24.18

method	result	size
parts	Expression too large to display	4473
derivativedivides	Expression too large to display	12836
default	Expression too large to display	12836

```
input int((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x,method=_RETURNVERBOSE)
```

output

```
A*(-1/4/d/b/(a^2+b^2)^(7/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^5+1/2/d*b/(a^2+b^2)^(7/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3+3/4/d*b^3/(a^2+b^2)^(7/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-1/d/b/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^4+1/d/b/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^6+4/d*b/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^4-2/d*b/(a^2+b^2)^3/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^3+1/4/d/b/(a^2+b^2)^3*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^4-2/d*b^3/(a^2+b^2)^3/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a+1/d*b^3/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)...
```

3.361.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7181 vs. $2(156) = 312$.

Time = 2.21 (sec) , antiderivative size = 7181, normalized size of antiderivative = 38.82

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="fracas")`

output Too large to include

3.361.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))/(a + b*tan(c + d*x))**(5/2), x)`

3.361.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is`

3.361.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Timed out`

3.361.9 Mupad [B] (verification not implemented)

Time = 27.14 (sec) , antiderivative size = 9457, normalized size of antiderivative = 51.12

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

```
input int((A + B*tan(c + d*x))/(a + b*tan(c + d*x))^(5/2),x)
```

```
output (log((((a + b*tan(c + d*x))^(1/2)*(320*A^2*a^4*b^14*d^3 - 16*A^2*b^18*d^3
+ 1024*A^2*a^6*b^12*d^3 + 1440*A^2*a^8*b^10*d^3 + 1024*A^2*a^10*b^8*d^3 +
320*A^2*a^12*b^6*d^3 - 16*A^2*a^16*b^2*d^3) - (((320*A^4*a^2*b^8*d^4 - 16
*A^4*b^10*d^4 - 1760*A^4*a^4*b^6*d^4 + 1600*A^4*a^6*b^4*d^4 - 400*A^4*a^8
*b^2*d^4)^(1/2) - 4*A^2*a^5*d^2 + 40*A^2*a^3*b^2*d^2 - 20*A^2*a*b^4*d^2)/(a
^10*d^4 + b^10*d^4 + 5*a^2*b^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a
^8*b^2*d^4))^(1/2)*(((320*A^4*a^2*b^8*d^4 - 16*A^4*b^10*d^4 - 1760*A^4*a
^4*b^6*d^4 + 1600*A^4*a^6*b^4*d^4 - 400*A^4*a^8*b^2*d^4)^(1/2) - 4*A^2*a^5
*d^2 + 40*A^2*a^3*b^2*d^2 - 20*A^2*a*b^4*d^2)/(a^10*d^4 + b^10*d^4 + 5*a^2
*b^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^(1/2)*(a + b
tan(c + d*x))^(1/2)*(64*a*b^22*d^5 + 640*a^3*b^20*d^5 + 2880*a^5*b^18*d^5
+ 7680*a^7*b^16*d^5 + 13440*a^9*b^14*d^5 + 16128*a^11*b^12*d^5 + 13440*a^1
3*b^10*d^5 + 7680*a^15*b^8*d^5 + 2880*a^17*b^6*d^5 + 640*a^19*b^4*d^5 + 64
*a^21*b^2*d^5))/4 - 32*A*b^21*d^4 - 160*A*a^2*b^19*d^4 - 128*A*a^4*b^17*d^
4 + 896*A*a^6*b^15*d^4 + 3136*A*a^8*b^13*d^4 + 4928*A*a^10*b^11*d^4 + 4480
*A*a^12*b^9*d^4 + 2432*A*a^14*b^7*d^4 + 736*A*a^16*b^5*d^4 + 96*A*a^18*b^3
*d^4))/4)*(((320*A^4*a^2*b^8*d^4 - 16*A^4*b^10*d^4 - 1760*A^4*a^4*b^6*d^4
+ 1600*A^4*a^6*b^4*d^4 - 400*A^4*a^8*b^2*d^4)^(1/2) - 4*A^2*a^5*d^2 + 40*A
^2*a^3*b^2*d^2 - 20*A^2*a*b^4*d^2)/(a^10*d^4 + b^10*d^4 + 5*a^2*b^8*d^4 +
10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^(1/2))/4 + 96*A^3*a^3...
```


3.362
$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

3.362.1 Optimal result 3442
 3.362.2 Mathematica [A] (verified) 3443
 3.362.3 Rubi [A] (warning: unable to verify) 3443
 3.362.4 Maple [B] (verified) 3449
 3.362.5 Fracas [B] (verification not implemented) 3449
 3.362.6 Sympy [F] 3450
 3.362.7 Maxima [F] 3450
 3.362.8 Giac [F(-1)] 3450
 3.362.9 Mupad [B] (verification not implemented) 3451

3.362.1 Optimal result

Integrand size = 31, antiderivative size = 224

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = -\frac{2A \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{a^{5/2}d} + \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{5/2}d} + \frac{(A+iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a+ib)^{5/2}d} + \frac{2b(Ab-aB)}{3a(a^2+b^2)d(a+b \tan(c+dx))^{3/2}} + \frac{2b(3a^2Ab+Ab^3-2a^3B)}{a^2(a^2+b^2)^2d\sqrt{a+b \tan(c+dx)}}$$

```
output -2*A*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/a^(5/2)/d+(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(5/2)/d+(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(5/2)/d+2*b*(3*A*a^2*b+A*b^3-2*B*a^3)/a^2/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^(1/2)+2/3*b*(A*b-B*a)/a/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)
```

3.362.2 Mathematica [A] (verified)

Time = 5.47 (sec) , antiderivative size = 242, normalized size of antiderivative = 1.08

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{2 \left(-\frac{3A(a^2+b^2) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{a^{3/2}} + \frac{3a(a+ib)(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{2(a-ib)^{3/2}} \right)}{1}$$

input `Integrate[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]`

output `(2*((-3*A*(a^2 + b^2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/a^(3/2) + (3*a*(a + I*b)*(A - I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/(2*(a - I*b)^(3/2)) + (3*a*(a - I*b)*(A + I*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/(2*(a + I*b)^(3/2)) + (b*(A*b - a*B))/(a + b*Tan[c + d*x])^(3/2) + (3*b*(3*a^2*A*b + A*b^3 - 2*a^3*B))/(a*(a^2 + b^2)*Sqrt[a + b*Tan[c + d*x]])))/(3*a*(a^2 + b^2)*d)`

3.362.3 Rubi [A] (warning: unable to verify)

Time = 1.87 (sec) , antiderivative size = 272, normalized size of antiderivative = 1.21, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.613$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4136, 25, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A+B \tan(c+dx)}{\tan(c+dx)(a+b \tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{4092} \\ & \frac{2 \int \frac{3 \cot(c+dx)(b(Ab-aB) \tan^2(c+dx) - a(Ab-aB) \tan(c+dx) + A(a^2+b^2))}{2(a+b \tan(c+dx))^{3/2}} dx}{\frac{3a(a^2+b^2)}{2b(Ab-aB)}} + \\ & \quad \frac{3ad(a^2+b^2)}{(a+b \tan(c+dx))^{3/2}} \end{aligned}$$

3.362. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \int \frac{\cot(c+dx)(b(Ab-aB)\tan^2(c+dx)-a(Ab-aB)\tan(c+dx)+A(a^2+b^2))}{a(a^2+b^2)(a+b\tan(c+dx))^{3/2}} dx + \frac{2b(Ab-aB)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} \\
& \quad \downarrow 27 \\
& \int \frac{b(Ab-aB)\tan(c+dx)^2-a(Ab-aB)\tan(c+dx)+A(a^2+b^2)}{a(a^2+b^2)\tan(c+dx)(a+b\tan(c+dx))^{3/2}} dx + \frac{2b(Ab-aB)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} \\
& \quad \downarrow 3042 \\
& \int \frac{\cot(c+dx)\left(-((-Ba^2+2Aba+b^2B)\tan(c+dx)a^2)+A(a^2+b^2)^2+b(-2Ba^3+3Aba^2+Ab^3)\tan^2(c+dx)\right)}{2\sqrt{a+b\tan(c+dx)}a(a^2+b^2)} dx + \frac{2b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \\
& \quad \frac{a(a^2+b^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} \\
& \quad \downarrow 27 \\
& \int \frac{\cot(c+dx)\left(-((-Ba^2+2Aba+b^2B)\tan(c+dx)a^2)+A(a^2+b^2)^2+b(-2Ba^3+3Aba^2+Ab^3)\tan^2(c+dx)\right)}{\sqrt{a+b\tan(c+dx)}a(a^2+b^2)} dx + \frac{2b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \\
& \quad \frac{a(a^2+b^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} \\
& \quad \downarrow 3042 \\
& \int \frac{-((-Ba^2+2Aba+b^2B)\tan(c+dx)a^2)+A(a^2+b^2)^2+b(-2Ba^3+3Aba^2+Ab^3)\tan(c+dx)^2}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}a(a^2+b^2)} dx + \frac{2b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \\
& \quad \frac{a(a^2+b^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} \\
& \quad \downarrow 4136 \\
& \int -\frac{(-Ba^2+2Aba+b^2B)a^2+(Aa^2+2bBa-Ab^2)\tan(c+dx)a^2}{\sqrt{a+b\tan(c+dx)}a(a^2+b^2)} dx + A(a^2+b^2)^2 \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b\tan(c+dx)}} dx + \frac{2b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \\
& \quad \frac{a(a^2+b^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} \\
& \quad \downarrow 25
\end{aligned}$$

3.362. $\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

$$\frac{A(a^2+b^2)^2 \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx - \int \frac{(-Ba^2+2Aba+b^2B)a^2+(Aa^2+2bBa-Ab^2) \tan(c+dx)a^2}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} +$$

$$\frac{a(a^2+b^2)}{2b(Ab-aB)} \\ \frac{2b(Ab-aB)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}$$

↓ 3042

$$\frac{A(a^2+b^2)^2 \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - \int \frac{(-Ba^2+2Aba+b^2B)a^2+(Aa^2+2bBa-Ab^2) \tan(c+dx)a^2}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} +$$

$$\frac{a(a^2+b^2)}{2b(Ab-aB)} \\ \frac{2b(Ab-aB)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}$$

↓ 4022

$$\frac{2b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{A(a^2+b^2)^2 \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2}a^2(a-ib)^2(-B+iA) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}a^2(a+ib)^2(B+iA) \int \frac{1+i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)}$$

↓ 3042

$$\frac{2b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{A(a^2+b^2)^2 \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2}a^2(a-ib)^2(-B+iA) \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}a^2(a+ib)^2(B+iA) \int \frac{1+i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)}$$

↓ 4020

$$\frac{2b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{A(a^2+b^2)^2 \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + \frac{ia^2(a-ib)^2(-B+iA) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d}}{a(a^2+b^2)}$$

↓ 25

$$\frac{2b(-2a^3B+3a^2Ab+Ab^3)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{A(a^2+b^2)^2 \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - \frac{ia^2(a-ib)^2(-B+iA) \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d}}{a(a^2+b^2)}$$

↓ 73

3.362. $\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\frac{\frac{2b(Ab - aB)}{3ad(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-2a^3B + 3a^2Ab + Ab^3)}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{A(a^2 + b^2)^2 \int \frac{\tan(c + dx)^2 + 1}{\tan(c + dx)\sqrt{a + b \tan(c + dx)}} dx - \frac{a^2(a - ib)^2(-B + iA) \int \frac{1}{b} d\sqrt{a + b \tan(c + dx)} - \frac{a^2(a + ib)^2}{bd} + \frac{a^2(a + ib)^2(-B + iA) \int \frac{1}{b} d\sqrt{a + b \tan(c + dx)}}{a(a^2 + b^2)}}{a(a^2 + b^2)}$$

↓ 221

$$\frac{\frac{2b(Ab - aB)}{3ad(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-2a^3B + 3a^2Ab + Ab^3)}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{A(a^2 + b^2)^2 \int \frac{\tan(c + dx)^2 + 1}{\tan(c + dx)\sqrt{a + b \tan(c + dx)}} dx - \frac{a^2(a - ib)^2(-B + iA) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a + ib}}\right)}{d\sqrt{a + ib}} + \frac{a^2(a + ib)^2(B + iA) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a - ib}}\right)}{d\sqrt{a - ib}}}{a(a^2 + b^2)}$$

↓ 4117

$$\frac{\frac{2b(Ab - aB)}{3ad(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-2a^3B + 3a^2Ab + Ab^3)}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{A(a^2 + b^2)^2 \int \frac{\cot(c + dx)}{\sqrt{a + b \tan(c + dx)}} d \tan(c + dx) - \frac{a^2(a - ib)^2(-B + iA) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a + ib}}\right)}{d\sqrt{a + ib}} + \frac{a^2(a + ib)^2(B + iA) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a - ib}}\right)}{d\sqrt{a - ib}}}{a(a^2 + b^2)}$$

↓ 73

$$\frac{\frac{2b(Ab - aB)}{3ad(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-2a^3B + 3a^2Ab + Ab^3)}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{2A(a^2 + b^2)^2 \int \frac{1}{a + b \tan(c + dx)} d\sqrt{a + b \tan(c + dx)} - \frac{a^2(a - ib)^2(-B + iA) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a + ib}}\right)}{d\sqrt{a + ib}} + \frac{a^2(a + ib)^2(B + iA) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a - ib}}\right)}{d\sqrt{a - ib}}}{a(a^2 + b^2)}$$

↓ 221

$$\frac{\frac{2b(Ab - aB)}{3ad(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-2a^3B + 3a^2Ab + Ab^3)}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{a^2(a - ib)^2(-B + iA) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a + ib}}\right)}{d\sqrt{a + ib}} + \frac{a^2(a + ib)^2(B + iA) \arctan\left(\frac{\tan(c + dx)}{\sqrt{a - ib}}\right)}{d\sqrt{a - ib}} - \frac{2A(a^2 + b^2)^2 \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a}}\right)}{\sqrt{ad}}}{a(a^2 + b^2)}$$

```
input Int[(Cot[c + d*x]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2), x]
```

3.362. $\int \frac{\cot(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx$

```
output (2*b*(A*b - a*B))/(3*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) + (((a^2*
(a + I*b)^2*(I*A + B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d
) - (a^2*(a - I*b)^2*(I*A - B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a
+ I*b]*d) - (2*A*(a^2 + b^2)^2*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])
/(Sqrt[a]*d))/(a*(a^2 + b^2)) + (2*b*(3*a^2*A*b + A*b^3 - 2*a^3*B))/(a*(a^
2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]))/(a*(a^2 + b^2))
```

3.362.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2])/(a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.362.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 12869 vs. $2(194) = 388$.

Time = 0.23 (sec) , antiderivative size = 12870, normalized size of antiderivative = 57.46

method	result	size
derivativeldivides	Expression too large to display	12870
default	Expression too large to display	12870

```
input int(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x,method=_RETURNVER
BOSE)
```

```
output result too large to display
```

3.362.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7342 vs. $2(189) = 378$.

Time = 19.73 (sec) , antiderivative size = 14700, normalized size of antiderivative = 65.62

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm=
"fracas")
```

```
output Too large to include
```

3.362. $\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

3.362.6 Sympy [F]

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\cot(c+dx)}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)**(5/2), x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)/(a + b*tan(c + d*x))**(5/2), x)`

3.362.7 Maxima [F]

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)}{(b\tan(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2), x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)/(b*tan(d*x + c) + a)^(5/2), x)`

3.362.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cot(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2), x, algorithm="giac")`

output `Timed out`

3.362.9 Mupad [B] (verification not implemented)

Time = 18.99 (sec) , antiderivative size = 45681, normalized size of antiderivative = 203.93

$$\int \frac{\cot(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

```
input int((cot(c + d*x)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)
```

```
output atan(-((((((8*A^2*a^5*d^2 - 8*B^2*a^5*d^2 - 80*A^2*a^3*b^2*d^2 + 80*B^2*a^3*b^2*d^2 + 16*A*B*b^5*d^2 + 40*A^2*a*b^4*d^2 - 40*B^2*a*b^4*d^2 - 160*A*B*a^2*b^3*d^2 + 80*A*B*a^4*b*d^2)^2/4 - (A^4 + 2*A^2*B^2 + B^4)*(16*a^10*d^4 + 16*b^10*d^4 + 80*a^2*b^8*d^4 + 160*a^4*b^6*d^4 + 160*a^6*b^4*d^4 + 80*a^8*b^2*d^4))^(1/2) - 4*A^2*a^5*d^2 + 4*B^2*a^5*d^2 + 40*A^2*a^3*b^2*d^2 - 40*B^2*a^3*b^2*d^2 - 8*A*B*b^5*d^2 - 20*A^2*a*b^4*d^2 + 20*B^2*a*b^4*d^2 + 80*A*B*a^2*b^3*d^2 - 40*A*B*a^4*b*d^2)/(16*(a^10*d^4 + b^10*d^4 + 5*a^2*b^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^(1/2))*((a + b*tan(c + d*x))^(1/2)*(256*A^2*a^15*b^44*d^7 + 4608*A^2*a^17*b^42*d^7 + 40512*A^2*a^19*b^40*d^7 + 224768*A^2*a^21*b^38*d^7 + 864768*A^2*a^23*b^36*d^7 + 2419200*A^2*a^25*b^34*d^7 + 5055232*A^2*a^27*b^32*d^7 + 8007168*A^2*a^29*b^30*d^7 + 9664512*A^2*a^31*b^28*d^7 + 8859136*A^2*a^33*b^26*d^7 + 6095232*A^2*a^35*b^24*d^7 + 3095040*A^2*a^37*b^22*d^7 + 1164800*A^2*a^39*b^20*d^7 + 376320*A^2*a^41*b^18*d^7 + 154368*A^2*a^43*b^16*d^7 + 76288*A^2*a^45*b^14*d^7 + 28416*A^2*a^47*b^12*d^7 + 6144*A^2*a^49*b^10*d^7 + 576*A^2*a^51*b^8*d^7 - 1344*B^2*a^19*b^40*d^7 - 15872*B^2*a^21*b^38*d^7 - 81408*B^2*a^23*b^36*d^7 - 225792*B^2*a^25*b^34*d^7 - 302848*B^2*a^27*b^32*d^7 + 139776*B^2*a^29*b^30*d^7 + 1537536*B^2*a^31*b^28*d^7 + 3587584*B^2*a^33*b^26*d^7 + 5106816*B^2*a^35*b^24*d^7 + 5051904*B^2*a^37*b^22*d^7 + 3587584*B^2*a^39*b^20*d^7 + 1817088*B^2*a^41*b^18*d^7 + 628992*B^2*a^43*b^16*d^7 + 1...
```

3.363 $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

3.363.1 Optimal result 3452
 3.363.2 Mathematica [A] (verified) 3453
 3.363.3 Rubi [A] (warning: unable to verify) 3453
 3.363.4 Maple [B] (verified) 3460
 3.363.5 Fricas [B] (verification not implemented) 3460
 3.363.6 Sympy [F] 3461
 3.363.7 Maxima [F(-1)] 3461
 3.363.8 Giac [F(-1)] 3461
 3.363.9 Mupad [B] (verification not implemented) 3462

3.363.1 Optimal result

Integrand size = 33, antiderivative size = 289

$$\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{(5Ab - 2aB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{a^{7/2}d} + \frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{5/2}d} - \frac{(iA - B) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a+ib)^{5/2}d} - \frac{b(3a^2A + 5Ab^2 - 2abB)}{3a^2(a^2 + b^2)d(a+b \tan(c+dx))^{3/2}} - \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} - \frac{b(a^4A + 10a^2Ab^2 + 5Ab^4 - 6a^3bB - 2ab^3B)}{a^3(a^2 + b^2)^2d\sqrt{a+b \tan(c+dx)}}$$

output

```
(5*A*b-2*B*a)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/a^(7/2)/d+(I*A+B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(5/2)/d-(I*A-B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(5/2)/d-b*(A*a^4+10*A*a^2*b^2+5*A*b^4-6*B*a^3*b-2*B*a*b^3)/a^3/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^(1/2)-1/3*b*(3*A*a^2+5*A*b^2-2*B*a*b)/a^2/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)-A*cot(d*x+c)/a/d/(a+b*tan(d*x+c))^(3/2)
```

3.363.2 Mathematica [A] (verified)

Time = 5.43 (sec) , antiderivative size = 306, normalized size of antiderivative = 1.06

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx = \frac{b(-3a^2A - 5Ab^2 + 2abB)}{(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{3aA \cot(c + dx)}{(a + b \tan(c + dx))^{3/2}} + \frac{3 \left(\frac{(a^2 + b^2)^2 (5Ab - 2aB) \arctan\left(\frac{a + b \tan(c + dx)}{\sqrt{a}}\right)}{\sqrt{a}} \right)}{(a + b \tan(c + dx))^{3/2}}$$

input `Integrate[(Cot[c + d*x]^2*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2), x]`

output `((b*(-3*a^2*A - 5*A*b^2 + 2*a*b*B))/((a^2 + b^2)*(a + b*Tan[c + d*x])^(3/2)) - (3*a*A*Cot[c + d*x])/(a + b*Tan[c + d*x])^(3/2) + (3*((a^2 + b^2)^2*(5*A*b - 2*a*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/Sqrt[a] + (a^3*(a + I*b)^2*(I*A + B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/Sqrt[a - I*b] + (a^3*(a - I*b)^2*((-I)*A + B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/Sqrt[a + I*b] - (b*(a^4*A + 10*a^2*A*b^2 + 5*A*b^4 - 6*a^3*b*B - 2*a*b^3*B))/Sqrt[a + b*Tan[c + d*x]]))/(a*(a^2 + b^2)^2)/(3*a^2*d)`

3.363.3 Rubi [A] (warning: unable to verify)

Time = 2.37 (sec) , antiderivative size = 343, normalized size of antiderivative = 1.19, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^2(a + b \tan(c + dx))^{5/2}} dx$$

↓ 4092

$$-\frac{\int \frac{\cot(c + dx)(5Ab \tan^2(c + dx) + 2aA \tan(c + dx) + 5Ab - 2aB)}{2(a + b \tan(c + dx))^{5/2}} dx}{a} - \frac{A \cot(c + dx)}{ad(a + b \tan(c + dx))^{3/2}}$$

3.363. $\int \frac{\cot^2(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx$

$$\begin{aligned}
 & \int \frac{\cot(c+dx)(5Ab \tan^2(c+dx)+2aA \tan(c+dx)+5Ab-2aB)}{(a+b \tan(c+dx))^{5/2}} dx - \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 27 \\
 & \int \frac{5Ab \tan(c+dx)^2+2aA \tan(c+dx)+5Ab-2aB}{\tan(c+dx)(a+b \tan(c+dx))^{5/2}} dx - \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 3042 \\
 & 2 \int \frac{3 \cot(c+dx)(2(aA+bB) \tan(c+dx)a^2+b(3Aa^2-2bBa+5Ab^2) \tan^2(c+dx)+(a^2+b^2)(5Ab-2aB))}{2(a+b \tan(c+dx))^{3/2} 3a(a^2+b^2)} dx + \frac{2b(3a^2A-2abB+5Ab^2)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 4132 \\
 & \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 27 \\
 & \int \frac{\cot(c+dx)(2(aA+bB) \tan(c+dx)a^2+b(3Aa^2-2bBa+5Ab^2) \tan^2(c+dx)+(a^2+b^2)(5Ab-2aB))}{(a+b \tan(c+dx))^{3/2} a(a^2+b^2)} dx + \frac{2b(3a^2A-2abB+5Ab^2)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 3042 \\
 & \int \frac{2(aA+bB) \tan(c+dx)a^2+b(3Aa^2-2bBa+5Ab^2) \tan(c+dx)^2+(a^2+b^2)(5Ab-2aB)}{\tan(c+dx)(a+b \tan(c+dx))^{3/2} a(a^2+b^2)} dx + \frac{2b(3a^2A-2abB+5Ab^2)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 4132 \\
 & 2 \int \frac{\cot(c+dx)(2(Aa^2+2bBa-Ab^2) \tan(c+dx)a^3+b(Aa^4-6bBa^3+10Ab^2a^2-2b^3Ba+5Ab^4) \tan^2(c+dx)+(a^2+b^2)^2(5Ab-2aB))}{2\sqrt{a+b \tan(c+dx)} a(a^2+b^2)} dx + \frac{2b(a^4A-6a^3bB+10a^2Ab^2)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 27 \\
 & \frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}}
 \end{aligned}$$

3.363. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\int \frac{\cot(c+dx) \left(2(Aa^2+2bBa-Ab^2) \tan(c+dx)a^3 + b(Aa^4-6bBa^3+10Ab^2a^2-2b^3Ba+5Ab^4) \tan^2(c+dx) + (a^2+b^2)^2(5Ab-2aB) \right) dx}{\frac{\sqrt{a+b \tan(c+dx)}}{a(a^2+b^2)}} + \frac{2b(a^4A-6a^3bB+10a^2Ab^2-2ab^3B+5Ab^4)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}}$$

2a

↓ 3042

$$\int \frac{2(Aa^2+2bBa-Ab^2) \tan(c+dx)a^3 + b(Aa^4-6bBa^3+10Ab^2a^2-2b^3Ba+5Ab^4) \tan(c+dx)^2 + (a^2+b^2)^2(5Ab-2aB)}{\frac{\tan(c+dx)\sqrt{a+b \tan(c+dx)}}{a(a^2+b^2)}} dx + \frac{2b(a^4A-6a^3bB+10a^2Ab^2-2ab^3B+5Ab^4)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}}$$

2a

↓ 4136

$$\frac{(a^2+b^2)^2(5Ab-2aB) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx + \int \frac{2(a^3(Aa^2+2bBa-Ab^2)-a^3(-Ba^2+2Aba+b^2B) \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(a^4A-6a^3bB+10a^2Ab^2-2ab^3B+5Ab^4)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}}$$

2a

↓ 27

$$\frac{(a^2+b^2)^2(5Ab-2aB) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx + 2 \int \frac{a^3(Aa^2+2bBa-Ab^2)-a^3(-Ba^2+2Aba+b^2B) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(a^4A-6a^3bB+10a^2Ab^2-2ab^3B+5Ab^4)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}}$$

2a

↓ 3042

$$\frac{(a^2+b^2)^2(5Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + 2 \int \frac{a^3(Aa^2+2bBa-Ab^2)-a^3(-Ba^2+2Aba+b^2B) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(a^4A-6a^3bB+10a^2Ab^2-2ab^3B+5Ab^4)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}$$

$$\frac{A \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}}$$

2a

↓ 4022

3.363. $\int \frac{\cot^2(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\frac{2b(3a^2A-2abB+5Ab^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{-\frac{A \cot(c+dx)}{ad(a+b\tan(c+dx))^{3/2}}}{2a} + \frac{2b(a^4A-6a^3bB+10a^2Ab^2-2ab^3B+5Ab^4)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{(a^2+b^2)^2(5Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx + 2\left(\frac{1}{2}a^3(a-ib)^2(A+ia^3(a+ib)^2(A-ib))\right)}{a(a^2+b^2)}$$

↓ 3042

$$\frac{2b(3a^2A-2abB+5Ab^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{-\frac{A \cot(c+dx)}{ad(a+b\tan(c+dx))^{3/2}}}{2a} + \frac{2b(a^4A-6a^3bB+10a^2Ab^2-2ab^3B+5Ab^4)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{(a^2+b^2)^2(5Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx + 2\left(\frac{1}{2}a^3(a-ib)^2(A+ia^3(a+ib)^2(A-ib))\right)}{a(a^2+b^2)}$$

↓ 4020

$$\frac{2b(3a^2A-2abB+5Ab^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{-\frac{A \cot(c+dx)}{ad(a+b\tan(c+dx))^{3/2}}}{2a} + \frac{2b(a^4A-6a^3bB+10a^2Ab^2-2ab^3B+5Ab^4)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{(a^2+b^2)^2(5Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx + 2\left(\frac{ia^3(a+ib)^2(A-ib)}{a(a^2+b^2)}\right)}{2a}$$

↓ 25

$$\frac{2b(3a^2A-2abB+5Ab^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{-\frac{A \cot(c+dx)}{ad(a+b\tan(c+dx))^{3/2}}}{2a} + \frac{2b(a^4A-6a^3bB+10a^2Ab^2-2ab^3B+5Ab^4)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{(a^2+b^2)^2(5Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx + 2\left(\frac{ia^3(a-ib)^2(A+ia^3(a+ib)^2(A-ib))}{a(a^2+b^2)}\right)}{2a}$$

↓ 73

$$\frac{2b(3a^2A-2abB+5Ab^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{-\frac{A \cot(c+dx)}{ad(a+b\tan(c+dx))^{3/2}}}{2a} + \frac{2b(a^4A-6a^3bB+10a^2Ab^2-2ab^3B+5Ab^4)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{(a^2+b^2)^2(5Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx + 2\left(\frac{a^3(a-ib)^2(A+ia^3(a+ib)^2(A-ib))}{a(a^2+b^2)}\right)}{2a}$$

↓ 221

3.363. $\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

$$\frac{2b(3a^2A-2abB+5Ab^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{A \cot(c+dx)}{ad(a+b\tan(c+dx))^{3/2}} + \frac{2b(a^4A-6a^3bB+10a^2Ab^2-2ab^3B+5Ab^4)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{(a^2+b^2)^2(5Ab-2aB) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx + 2 \left(\frac{a^3(a+ib)^2(A-ib)}{a(a^2+b^2)} \right)}{2a(a^2+b^2)}$$

↓ 4117

$$\frac{2b(3a^2A-2abB+5Ab^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{A \cot(c+dx)}{ad(a+b\tan(c+dx))^{3/2}} + \frac{2b(a^4A-6a^3bB+10a^2Ab^2-2ab^3B+5Ab^4)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{(a^2+b^2)^2(5Ab-2aB) \int \frac{\cot(c+dx)}{\sqrt{a+b\tan(c+dx)}} d\tan(c+dx) + 2 \left(\frac{a^3(a+ib)^2(A-ib)}{a(a^2+b^2)} \right)}{2a(a^2+b^2)}$$

↓ 73

$$\frac{2b(3a^2A-2abB+5Ab^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{A \cot(c+dx)}{ad(a+b\tan(c+dx))^{3/2}} + \frac{2b(a^4A-6a^3bB+10a^2Ab^2-2ab^3B+5Ab^4)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{2(a^2+b^2)^2(5Ab-2aB) \int \frac{1}{\frac{a+b\tan(c+dx)}{bd} - \frac{a}{b}} d\sqrt{a+b\tan(c+dx)} + 2 \left(\frac{a^3(a+ib)^2(A-ib)}{a(a^2+b^2)} \right)}{2a(a^2+b^2)}$$

↓ 221

$$\frac{2b(3a^2A-2abB+5Ab^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{A \cot(c+dx)}{ad(a+b\tan(c+dx))^{3/2}} + \frac{2b(a^4A-6a^3bB+10a^2Ab^2-2ab^3B+5Ab^4)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{2(a^2+b^2)^2(5Ab-2aB) \operatorname{arctanh}\left(\frac{\sqrt{a+b\tan(c+dx)}}{\sqrt{a}}\right) + 2 \left(\frac{a^3(a+ib)^2(A-ib)}{a(a^2+b^2)} \right)}{2a(a^2+b^2)}$$

input `Int[(Cot[c + d*x])^2*(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^(5/2),x]`


```
output -((A*Cot[c + d*x])/(a*d*(a + b*Tan[c + d*x])^(3/2))) - ((2*b*(3*a^2*A + 5*
A*b^2 - 2*a*b*B))/(3*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) + ((2*((a
^3*(a + I*b)^2*(A - I*B)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b
]*d) + (a^3*(a - I*b)^2*(A + I*B)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqr
t[a + I*b]*d)) - (2*(a^2 + b^2)^2*(5*A*b - 2*a*B)*ArcTanh[Sqrt[a + b*Tan[c
+ d*x]]/Sqrt[a]])/(Sqrt[a]*d)/(a*(a^2 + b^2)) + (2*b*(a^4*A + 10*a^2*A*b
^2 + 5*A*b^4 - 6*a^3*b*B - 2*a*b^3*B))/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c +
d*x]]))/(a*(a^2 + b^2))/(2*a)
```

3.363.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 73 Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

rule 4022 $\text{Int}(((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^{(m_.)}*((c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_.)]), x_Symbol] \rightarrow \text{Simp}[(c + I*d)/2 \text{ Int}[(a + b*\tan[e + f*x])^m*(1 - I*\tan[e + f*x]), x], x] + \text{Simp}[(c - I*d)/2 \text{ Int}[(a + b*\tan[e + f*x])^m*(1 + I*\tan[e + f*x]), x], x] /;$ FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]

rule 4092 $\text{Int}(((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^{(m_.)}*((A_.) + (B_.)*\tan[(e_.) + (f_.)*(x_.)]*(c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[b*(A*b - a*B)*(a + b*\tan[e + f*x])^{(m+1)}*((c + d*\tan[e + f*x])^{(n+1)})/(f*(m+1)*(b*c - a*d)*(a^2 + b^2)), x] + \text{Simp}[1/((m+1)*(b*c - a*d)*(a^2 + b^2)) \text{ Int}[(a + b*\tan[e + f*x])^{(m+1)}*(c + d*\tan[e + f*x])^n*\text{Simp}[b*B*(b*c*(m+1) + a*d*(n+1)) + A*(a*(b*c - a*d)*(m+1) - b^2*d*(m+n+2)) - (A*b - a*B)*(b*c - a*d)*(m+1)*\tan[e + f*x] - b*d*(A*b - a*B)*(m+n+2)*\tan[e + f*x]^2, x], x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

rule 4117 $\text{Int}(((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^{(m_.)}*((c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_.)])^{(n_.)}*((A_.) + (C_.)*\tan[(e_.) + (f_.)*(x_.)]^2), x_Symbol] \rightarrow \text{Simp}[A/f \text{ Subst}[\text{Int}[(a + b*x)^m*(c + d*x)^n, x], x, \tan[e + f*x]], x] /;$ FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]

rule 4132 $\text{Int}(((a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_.)])^{(m_.)}*((c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_.)])^{(n_.)}*((A_.) + (B_.)*\tan[(e_.) + (f_.)*(x_.)] + (C_.)*\tan[(e_.) + (f_.)*(x_.)]^2), x_Symbol] \rightarrow \text{Simp}[(A*b^2 - a*(b*B - a*C))*(a + b*\tan[e + f*x])^{(m+1)}*((c + d*\tan[e + f*x])^{(n+1)})/(f*(m+1)*(b*c - a*d)*(a^2 + b^2)), x] + \text{Simp}[1/((m+1)*(b*c - a*d)*(a^2 + b^2)) \text{ Int}[(a + b*\tan[e + f*x])^{(m+1)}*(c + d*\tan[e + f*x])^n*\text{Simp}[A*(a*(b*c - a*d)*(m+1) - b^2*d*(m+n+2)) + (b*B - a*C)*(b*c*(m+1) + a*d*(n+1)) - (m+1)*(b*c - a*d)*(A*b - a*B - b*C)*\tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m+n+2)*\tan[e + f*x]^2, x], x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.363.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 12967 vs. $2(257) = 514$.

Time = 0.26 (sec) , antiderivative size = 12968, normalized size of antiderivative = 44.87

method	result	size
derivativedivides	Expression too large to display	12968
default	Expression too large to display	12968

```
input int(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output result too large to display
```

3.363.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7533 vs. $2(252) = 504$.

Time = 57.60 (sec) , antiderivative size = 15082, normalized size of antiderivative = 52.19

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorith
m="fracas")
```

```
output Too large to include
```

3.363.6 Sympy [F]

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\cot^2(c+dx)}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx$$

input `integrate(cot(d*x+c)**2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**2/(a + b*tan(c + d*x))**(5/2), x)`

3.363.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm m="maxima")`

output `Timed out`

3.363.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^2*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm m="giac")`

output `Timed out`

3.363.9 Mupad [B] (verification not implemented)

Time = 15.51 (sec) , antiderivative size = 67465, normalized size of antiderivative = 233.44

$$\int \frac{\cot^2(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

```
input int((cot(c + d*x)^2*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)
```

```
output ((2*(a + b*tan(c + d*x))*(5*A*b^5 + 11*A*a^2*b^3 - 8*B*a^3*b^2 - 2*B*a*b^4
))/((3*(a*b^2 + a^3)^2) + (2*(A*b^3 - B*a*b^2))/(3*a*(a^2 + b^2))) - ((a + b
*tan(c + d*x))^2*(5*A*b^5 + 10*A*a^2*b^3 - 6*B*a^3*b^2 + A*a^4*b - 2*B*a*b
^4))/(a^3*(a^2 + b^2)^2)/(d*(a + b*tan(c + d*x))^(5/2) - a*d*(a + b*tan(c
+ d*x))^(3/2)) + atan((((((8*A^2*a^5*d^2 - 8*B^2*a^5*d^2 - 80*A^2*a^3*b
^2*d^2 + 80*B^2*a^3*b^2*d^2 + 16*A*B*b^5*d^2 + 40*A^2*a*b^4*d^2 - 40*B^2*a
b^4*d^2 - 160*A*B*a^2*b^3*d^2 + 80*A*B*a^4*b*d^2)^2/4 - (A^4 + 2*A^2*B^2 +
B^4)*(16*a^10*d^4 + 16*b^10*d^4 + 80*a^2*b^8*d^4 + 160*a^4*b^6*d^4 + 160*
a^6*b^4*d^4 + 80*a^8*b^2*d^4))^(1/2) - 4*A^2*a^5*d^2 + 4*B^2*a^5*d^2 + 40*
A^2*a^3*b^2*d^2 - 40*B^2*a^3*b^2*d^2 - 8*A*B*b^5*d^2 - 20*A^2*a*b^4*d^2 +
20*B^2*a*b^4*d^2 + 80*A*B*a^2*b^3*d^2 - 40*A*B*a^4*b*d^2)/(16*(a^10*d^4 +
b^10*d^4 + 5*a^2*b^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4
))))^(1/2)*((((((8*A^2*a^5*d^2 - 8*B^2*a^5*d^2 - 80*A^2*a^3*b^2*d^2 + 80*B
^2*a^3*b^2*d^2 + 16*A*B*b^5*d^2 + 40*A^2*a*b^4*d^2 - 40*B^2*a*b^4*d^2 - 160
*A*B*a^2*b^3*d^2 + 80*A*B*a^4*b*d^2)^2/4 - (A^4 + 2*A^2*B^2 + B^4)*(16*a^1
0*d^4 + 16*b^10*d^4 + 80*a^2*b^8*d^4 + 160*a^4*b^6*d^4 + 160*a^6*b^4*d^4 +
80*a^8*b^2*d^4))^(1/2) - 4*A^2*a^5*d^2 + 4*B^2*a^5*d^2 + 40*A^2*a^3*b^2*d
^2 - 40*B^2*a^3*b^2*d^2 - 8*A*B*b^5*d^2 - 20*A^2*a*b^4*d^2 + 20*B^2*a*b^4*
d^2 + 80*A*B*a^2*b^3*d^2 - 40*A*B*a^4*b*d^2)/(16*(a^10*d^4 + b^10*d^4 + 5*
a^2*b^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))))^(1/2)*...
```

3.364
$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

3.364.1 Optimal result 3463
 3.364.2 Mathematica [A] (verified) 3464
 3.364.3 Rubi [A] (warning: unable to verify) 3464
 3.364.4 Maple [B] (verified) 3472
 3.364.5 Fricas [B] (verification not implemented) 3472
 3.364.6 Sympy [F] 3473
 3.364.7 Maxima [F(-1)] 3473
 3.364.8 Giac [F(-1)] 3473
 3.364.9 Mupad [B] (verification not implemented) 3474

3.364.1 Optimal result

Integrand size = 33, antiderivative size = 364

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{(8a^2A - 35Ab^2 + 20abB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{4a^{9/2}d}$$

$$- \frac{(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a - ib)^{5/2}d} - \frac{(A + iB) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a + ib)^{5/2}d}$$

$$+ \frac{b(27a^2Ab + 35Ab^3 - 12a^3B - 20ab^2B)}{12a^3(a^2 + b^2)d(a + b \tan(c+dx))^{3/2}} + \frac{(7Ab - 4aB) \cot(c+dx)}{4a^2d(a + b \tan(c+dx))^{3/2}}$$

$$- \frac{A \cot^2(c+dx)}{2ad(a + b \tan(c+dx))^{3/2}} + \frac{b(11a^4Ab + 62a^2Ab^3 + 35Ab^5 - 4a^5B - 40a^3b^2B - 20ab^4B)}{4a^4(a^2 + b^2)^2d\sqrt{a + b \tan(c+dx)}}$$

output

```
1/4*(8*A*a^2-35*A*b^2+20*B*a*b)*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/a^(9/2)/d-(A-I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(5/2)/d-(A+I*B)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(5/2)/d+1/4*b*(11*A*a^4*b+62*A*a^2*b^3+35*A*b^5-4*B*a^5-40*B*a^3*b^2-20*B*a*b^4)/a^4/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^(1/2)+1/12*b*(27*A*a^2*b+35*A*b^3-12*B*a^3-20*B*a*b^2)/a^3/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)+1/4*(7*A*b-4*B*a)*cot(d*x+c)/a^2/d/(a+b*tan(d*x+c))^(3/2)-1/2*A*cot(d*x+c)^2/a/d/(a+b*tan(d*x+c))^(3/2)
```

3.364.2 Mathematica [A] (verified)

Time = 6.36 (sec) , antiderivative size = 593, normalized size of antiderivative = 1.63

$$\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = -\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} - \frac{(7Ab-4aB) \cot(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} - \frac{2\left(\frac{1}{4}b^2(-8a^2A+35Ab^2-20abB)-a(-2a^2bB-\frac{5}{4}ab(7Ab-4aB))\right)}{3a(a^2+b^2)d(a+b \tan(c+dx))^{3/2}} + \frac{2\left(\frac{3(a^2+b^2)^2(8a^2A-35Ab^2+20abB)}{8\sqrt{ad}}\right) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{2}$$

input `Integrate[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]`

output `-1/2*(A*Cot[c + d*x]^2)/(a*d*(a + b*Tan[c + d*x])^(3/2)) - (-1/2*((7*A*b - 4*a*B)*Cot[c + d*x])/(a*d*(a + b*Tan[c + d*x])^(3/2)) - ((2*((b^2*(-8*a^2 *A + 35*A*b^2 - 20*a*b*B))/4 - a*(-2*a^2*b*B - (5*a*b*(7*A*b - 4*a*B))/4)))/(3*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) + (2*((2*((3*(a^2 + b^2)^2*(8*a^2*A - 35*A*b^2 + 20*a*b*B)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/(8*Sqrt[a]*d) + (I*Sqrt[a - I*b]*(((3*I)/2)*a^4*(a^2*A - A*b^2 + 2*a*b*B) + (3*a^4*(2*a*A*b - a^2*B + b^2*B))/2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/((-a + I*b)*d) - (I*Sqrt[a + I*b]*(((3*I)/2)*a^4*(a^2*A - A*b^2 + 2*a*b*B) + (3*a^4*(2*a*A*b - a^2*B + b^2*B))/2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/((-a - I*b)*d)))/(a*(a^2 + b^2)) + (2*((-3*b^2*(a^2 + b^2)*(8*a^2*A - 35*A*b^2 + 20*a*b*B))/8 - a*(3*a^3*b*(A*b - a*B) - (3*a*b*(27*a^2*A*b + 35*A*b^3 - 12*a^3*B - 20*a*b^2*B))/8)))/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])))/(3*a*(a^2 + b^2)))/a/(2*a)`

3.364.3 Rubi [A] (warning: unable to verify)

Time = 3.08 (sec) , antiderivative size = 423, normalized size of antiderivative = 1.16, number of steps used = 26, number of rules used = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.758$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.364. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^3(a+b \tan(c+dx))^{5/2}} dx \\
& \quad \downarrow \text{4092} \\
& \frac{\int \frac{\cot^2(c+dx)(7Ab \tan^2(c+dx)+4aA \tan(c+dx)+7Ab-4aB)}{2(a+b \tan(c+dx))^{5/2}} dx}{2a} - \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{\cot^2(c+dx)(7Ab \tan^2(c+dx)+4aA \tan(c+dx)+7Ab-4aB)}{(a+b \tan(c+dx))^{5/2}} dx}{4a} - \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{7Ab \tan(c+dx)^2+4aA \tan(c+dx)+7Ab-4aB}{\tan(c+dx)^2(a+b \tan(c+dx))^{5/2}} dx}{4a} - \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{4132} \\
& \frac{\int -\frac{\cot(c+dx)(8Aa^2+8B \tan(c+dx)a^2+20bBa-35Ab^2-5b(7Ab-4aB) \tan^2(c+dx))}{2(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{(7Ab-4aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} \\
& \quad \frac{4a}{2ad(a+b \tan(c+dx))^{3/2}} \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{27} \\
& \frac{\int \frac{\cot(c+dx)(8Aa^2+8B \tan(c+dx)a^2+20bBa-35Ab^2-5b(7Ab-4aB) \tan^2(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx}{2a} - \frac{(7Ab-4aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} \\
& \quad \frac{4a}{2ad(a+b \tan(c+dx))^{3/2}} \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{8Aa^2+8B \tan(c+dx)a^2+20bBa-35Ab^2-5b(7Ab-4aB) \tan(c+dx)^2}{\tan(c+dx)(a+b \tan(c+dx))^{5/2}} dx}{2a} - \frac{(7Ab-4aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} \\
& \quad \frac{4a}{2ad(a+b \tan(c+dx))^{3/2}} \frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{4132}
\end{aligned}$$

3.364. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$2 \int \frac{3 \cot(c+dx) \left(-8(Ab-aB) \tan(c+dx)a^3 - b \left(-12Ba^3 + 27Aba^2 - 20b^2Ba + 35Ab^3 \right) \tan^2(c+dx) + (a^2+b^2) \left(8Aa^2 + 20bBa - 35Ab^2 \right) \right)}{2(a+b \tan(c+dx))^{3/2} \cdot 3a(a^2+b^2)} dx - \frac{2b \left(-12a^3B + 27a^2Ab - 20ab^2B + 35Ab^3 \right)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} \quad 4a$$

↓ 27

$$\int \frac{\cot(c+dx) \left(-8(Ab-aB) \tan(c+dx)a^3 - b \left(-12Ba^3 + 27Aba^2 - 20b^2Ba + 35Ab^3 \right) \tan^2(c+dx) + (a^2+b^2) \left(8Aa^2 + 20bBa - 35Ab^2 \right) \right)}{(a+b \tan(c+dx))^{3/2} \cdot a(a^2+b^2)} dx - \frac{2b \left(-12a^3B + 27a^2Ab - 20ab^2B + 35Ab^3 \right)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} \quad 4a$$

↓ 3042

$$\int \frac{-8(Ab-aB) \tan(c+dx)a^3 - b \left(-12Ba^3 + 27Aba^2 - 20b^2Ba + 35Ab^3 \right) \tan(c+dx)^2 + (a^2+b^2) \left(8Aa^2 + 20bBa - 35Ab^2 \right)}{\tan(c+dx)(a+b \tan(c+dx))^{3/2} \cdot a(a^2+b^2)} dx - \frac{2b \left(-12a^3B + 27a^2Ab - 20ab^2B + 35Ab^3 \right)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} \quad 4a$$

↓ 4132

$$2 \int \frac{\cot(c+dx) \left(-8 \left(-Ba^2 + 2Aba + b^2B \right) \tan(c+dx)a^4 - b \left(-4Ba^5 + 11Aba^4 - 40b^2Ba^3 + 62Ab^3a^2 - 20b^4Ba + 35Ab^5 \right) \tan^2(c+dx) + (a^2+b^2)^2 \left(8Aa^2 + 20bBa - 35Ab^2 \right) \right)}{2\sqrt{a+b \tan(c+dx)} \cdot a(a^2+b^2)} dx - \frac{2b \left(-12a^3B + 27a^2Ab - 20ab^2B + 35Ab^3 \right)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} \quad 4a$$

↓ 27

$$\int \frac{\cot(c+dx) \left(-8 \left(-Ba^2 + 2Aba + b^2B \right) \tan(c+dx)a^4 - b \left(-4Ba^5 + 11Aba^4 - 40b^2Ba^3 + 62Ab^3a^2 - 20b^4Ba + 35Ab^5 \right) \tan^2(c+dx) + (a^2+b^2)^2 \left(8Aa^2 + 20bBa - 35Ab^2 \right) \right)}{\sqrt{a+b \tan(c+dx)} \cdot a(a^2+b^2)} dx - \frac{2b \left(-12a^3B + 27a^2Ab - 20ab^2B + 35Ab^3 \right)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} \quad 4a$$

3.364. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

↓ 3042

$$\frac{\int \frac{-8(-Ba^2+2Aba+b^2B)\tan(c+dx)a^4 - b(-4Ba^5+11Aba^4-40b^2Ba^3+62Ab^3a^2-20b^4Ba+35Ab^5)\tan(c+dx)^2 + (a^2+b^2)^2(8Aa^2+20bBa-35Ab^2)}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - \frac{2b(-4a^5B+11a^4Ab-4a^3A^2+20a^2bB-35aAb^2)}{a(a^2+b^2)}}{a(a^2+b^2)} = \frac{2a}{4a}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b\tan(c+dx))^{3/2}}$$

↓ 4136

$$\frac{(a^2+b^2)^2(8a^2A+20abB-35Ab^2) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b\tan(c+dx)}} dx + \frac{8((-Ba^2+2Aba+b^2B)a^4 + (Aa^2+2bBa-Ab^2)\tan(c+dx)a^4)}{\sqrt{a+b\tan(c+dx)}} dx - \frac{2b(-4a^5B+11a^4Ab-4a^3A^2+20a^2bB-35aAb^2)}{a(a^2+b^2)}}{a(a^2+b^2)} = \frac{2a}{4a}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b\tan(c+dx))^{3/2}}$$

↓ 27

$$\frac{(a^2+b^2)^2(8a^2A+20abB-35Ab^2) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b\tan(c+dx)}} dx - \frac{8((-Ba^2+2Aba+b^2B)a^4 + (Aa^2+2bBa-Ab^2)\tan(c+dx)a^4)}{\sqrt{a+b\tan(c+dx)}} dx - \frac{2b(-4a^5B+11a^4Ab-4a^3A^2+20a^2bB-35aAb^2)}{a(a^2+b^2)}}{a(a^2+b^2)} = \frac{2a}{4a}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b\tan(c+dx))^{3/2}}$$

↓ 3042

$$\frac{(a^2+b^2)^2(8a^2A+20abB-35Ab^2) \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - \frac{8((-Ba^2+2Aba+b^2B)a^4 + (Aa^2+2bBa-Ab^2)\tan(c+dx)a^4)}{\sqrt{a+b\tan(c+dx)}} dx - \frac{2b(-4a^5B+11a^4Ab-4a^3A^2+20a^2bB-35aAb^2)}{a(a^2+b^2)}}{a(a^2+b^2)} = \frac{2a}{4a}$$

$$\frac{A \cot^2(c+dx)}{2ad(a+b\tan(c+dx))^{3/2}}$$

↓ 4022

3.364. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} - \frac{2b(-4a^5B+11a^4Ab-40a^3b^2B+62a^2Ab^3-20ab^4B+35Ab^5)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)^2(8a^2A+20aB)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} - \frac{(7Ab-4aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} + \frac{2b(-12a^3B+27a^2Ab-20ab^2B+35Ab^3)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}$$

4a

↓ 3042

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} - \frac{2b(-4a^5B+11a^4Ab-40a^3b^2B+62a^2Ab^3-20ab^4B+35Ab^5)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)^2(8a^2A+20aB)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} - \frac{(7Ab-4aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} + \frac{2b(-12a^3B+27a^2Ab-20ab^2B+35Ab^3)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}$$

4a

↓ 4020

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} - \frac{2b(-4a^5B+11a^4Ab-40a^3b^2B+62a^2Ab^3-20ab^4B+35Ab^5)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)^2(8a^2A+20aB)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} - \frac{(7Ab-4aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} + \frac{2b(-12a^3B+27a^2Ab-20ab^2B+35Ab^3)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}$$

↓ 25

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} - \frac{2b(-4a^5B+11a^4Ab-40a^3b^2B+62a^2Ab^3-20ab^4B+35Ab^5)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)^2(8a^2A+20aB)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} - \frac{(7Ab-4aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} + \frac{2b(-12a^3B+27a^2Ab-20ab^2B+35Ab^3)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}$$

↓ 73

$$\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} - \frac{2b(-4a^5B+11a^4Ab-40a^3b^2B+62a^2Ab^3-20ab^4B+35Ab^5)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)^2(8a^2A+20aB)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} - \frac{(7Ab-4aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} + \frac{2b(-12a^3B+27a^2Ab-20ab^2B+35Ab^3)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}$$

↓ 221

3.364. $\int \frac{\cot^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & -\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} - \\
 & -\frac{(7Ab-4aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} + \frac{-2b(-12a^3B+27a^2Ab-20ab^2B+35Ab^3)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \frac{-2b(-4a^5B+11a^4Ab-40a^3b^2B+62a^2Ab^3-20ab^4B+35Ab^5)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)^2(8a^2A+2)}{(a^2+b^2)^2(8a^2A+2)}
 \end{aligned}$$

4a

4117

$$\begin{aligned}
 & -\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} - \\
 & -\frac{(7Ab-4aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} + \frac{-2b(-12a^3B+27a^2Ab-20ab^2B+35Ab^3)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \frac{-2b(-4a^5B+11a^4Ab-40a^3b^2B+62a^2Ab^3-20ab^4B+35Ab^5)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{(a^2+b^2)^2(8a^2A+2)}{(a^2+b^2)^2(8a^2A+2)}
 \end{aligned}$$

4a

73

$$\begin{aligned}
 & -\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} - \\
 & -\frac{(7Ab-4aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} + \frac{-2b(-12a^3B+27a^2Ab-20ab^2B+35Ab^3)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \frac{-2b(-4a^5B+11a^4Ab-40a^3b^2B+62a^2Ab^3-20ab^4B+35Ab^5)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{2(a^2+b^2)^2(8a^2A+2)}{(a^2+b^2)^2(8a^2A+2)}
 \end{aligned}$$

4a

221

$$\begin{aligned}
 & -\frac{A \cot^2(c+dx)}{2ad(a+b \tan(c+dx))^{3/2}} - \\
 & -\frac{(7Ab-4aB) \cot(c+dx)}{ad(a+b \tan(c+dx))^{3/2}} + \frac{-2b(-12a^3B+27a^2Ab-20ab^2B+35Ab^3)}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \frac{-2b(-4a^5B+11a^4Ab-40a^3b^2B+62a^2Ab^3-20ab^4B+35Ab^5)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{2(a^2+b^2)^2(8a^2A+2)}{(a^2+b^2)^2(8a^2A+2)}
 \end{aligned}$$

4a

```
input Int[(Cot[c + d*x]^3*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]
```

output
$$-1/2*(A*\cot[c + d*x]^2)/(a*d*(a + b*\tan[c + d*x])^(3/2)) - (-(((7*A*b - 4*a*B)*\cot[c + d*x])/(a*d*(a + b*\tan[c + d*x])^(3/2))) + ((-2*b*(27*a^2*A*b + 35*A*b^3 - 12*a^3*B - 20*a*b^2*B))/(3*a*(a^2 + b^2)*d*(a + b*\tan[c + d*x])^(3/2)) + ((-8*(-(a^4*(a + I*b)^2*(I*A + B)*\text{ArcTan}[\tan[c + d*x]/\text{Sqrt}[a - I*b]])/(\text{Sqrt}[a - I*b]*d)) + (a^4*(a - I*b)^2*(I*A - B)*\text{ArcTan}[\tan[c + d*x]/\text{Sqrt}[a + I*b]])/(\text{Sqrt}[a + I*b]*d)) - (2*(a^2 + b^2)^2*(8*a^2*A - 35*A*b^2 + 20*a*b*B)*\text{ArcTanh}[\text{Sqrt}[a + b*\tan[c + d*x]]/\text{Sqrt}[a]])/(\text{Sqrt}[a]*d))/(a*(a^2 + b^2)) - (2*b*(11*a^4*A*b + 62*a^2*A*b^3 + 35*A*b^5 - 4*a^5*B - 40*a^3*b^2*B - 20*a*b^4*B))/(a*(a^2 + b^2)*d*\text{Sqrt}[a + b*\tan[c + d*x]])))/(a*(a^2 + b^2)))/(2*a))/(4*a)$$

3.364.3.1 Defintions of rubi rules used

rule 25 $\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$

rule 27 $\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x)] /; \text{FreeQ}[b, x]$

rule 73 $\text{Int}[(a_ + (b_)*(x_))^(m_)*((c_ + (d_)*(x_))^(n_), x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \quad \text{Subst}[\text{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 221 $\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4020 $\text{Int}[(a_ + (b_)*\tan[(e_ + (f_)*(x_))]^(m_)*((c_ + (d_)*\tan[(e_ + (f_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[c*(d/f) \quad \text{Subst}[\text{Int}[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*\tan[e + f*x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{NeQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[c^2 + d^2, 0]$

rule 4022 $\text{Int}[(a_. + (b_.)\tan[(e_.) + (f_.)(x_.)]^{(m_.)}((c_.) + (d_.)\tan[(e_.) + (f_.)(x_.)]), x_Symbol] \rightarrow \text{Simp}[(c + I*d)/2 \text{ Int}[(a + b*\text{Tan}[e + f*x])^m*(1 - I*\text{Tan}[e + f*x]), x], x] + \text{Simp}[(c - I*d)/2 \text{ Int}[(a + b*\text{Tan}[e + f*x])^m*(1 + I*\text{Tan}[e + f*x]), x], x] /;$ FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]

rule 4092 $\text{Int}[(a_. + (b_.)\tan[(e_.) + (f_.)(x_.)]^{(m_.)}((A_.) + (B_.)\tan[(e_.) + (f_.)(x_.)]^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[b*(A*b - a*B)*(a + b*\text{Tan}[e + f*x])^{(m+1)}*((c + d*\text{Tan}[e + f*x])^{(n+1)}/(f*(m+1)*(b*c - a*d)*(a^2 + b^2))), x] + \text{Simp}[1/((m+1)*(b*c - a*d)*(a^2 + b^2)) \text{ Int}[(a + b*\text{Tan}[e + f*x])^{(m+1)}*(c + d*\text{Tan}[e + f*x])^n*\text{Simp}[b*B*(b*c*(m+1) + a*d*(n+1)) + A*(a*(b*c - a*d)*(m+1) - b^2*d*(m+n+2)) - (A*b - a*B)*(b*c - a*d)*(m+1)*\text{Tan}[e + f*x] - b*d*(A*b - a*B)*(m+n+2)*\text{Tan}[e + f*x]^2, x], x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

rule 4117 $\text{Int}[(a_. + (b_.)\tan[(e_.) + (f_.)(x_.)]^{(m_.)}((c_.) + (d_.)\tan[(e_.) + (f_.)(x_.)]^{(n_.)}((A_) + (C_.)\tan[(e_.) + (f_.)(x_.)]^2), x_Symbol] \rightarrow \text{Simp}[A/f \text{ Subst}[\text{Int}[(a + b*x)^m*(c + d*x)^n, x], x, \text{Tan}[e + f*x]], x] /;$ FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]

rule 4132 $\text{Int}[(a_. + (b_.)\tan[(e_.) + (f_.)(x_.)]^{(m_.)}((c_.) + (d_.)\tan[(e_.) + (f_.)(x_.)]^{(n_.)}((A_.) + (B_.)\tan[(e_.) + (f_.)(x_.)] + (C_.)\tan[(e_.) + (f_.)(x_.)]^2), x_Symbol] \rightarrow \text{Simp}[(A*b^2 - a*(b*B - a*C))*(a + b*\text{Tan}[e + f*x])^{(m+1)}*((c + d*\text{Tan}[e + f*x])^{(n+1)}/(f*(m+1)*(b*c - a*d)*(a^2 + b^2))), x] + \text{Simp}[1/((m+1)*(b*c - a*d)*(a^2 + b^2)) \text{ Int}[(a + b*\text{Tan}[e + f*x])^{(m+1)}*(c + d*\text{Tan}[e + f*x])^n*\text{Simp}[A*(a*(b*c - a*d)*(m+1) - b^2*d*(m+n+2)) + (b*B - a*C)*(b*c*(m+1) + a*d*(n+1)) - (m+1)*(b*c - a*d)*(A*b - a*B - b*C)*\text{Tan}[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m+n+2)*\text{Tan}[e + f*x]^2, x], x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2])/(a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.364.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 13091 vs. $2(322) = 644$.

Time = 0.27 (sec) , antiderivative size = 13092, normalized size of antiderivative = 35.97

method	result	size
derivativdivides	Expression too large to display	13092
default	Expression too large to display	13092

```
input int(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output result too large to display
```

3.364.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7654 vs. $2(316) = 632$.

Time = 111.45 (sec) , antiderivative size = 15325, normalized size of antiderivative = 42.10

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorith
m="fracas")
```

```
output Too large to include
```

3.364. $\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

3.364.6 Sympy [F]

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\cot^3(c+dx)}{(a+b\tan(c+dx))^{5/2}} dx$$

input `integrate(cot(d*x+c)**3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**3/(a + b*tan(c + d*x))**(5/2), x)`

3.364.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm m="maxima")`

output `Timed out`

3.364.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cot^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^3*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm m="giac")`

output `Timed out`

3.364.9 Mupad [B] (verification not implemented)

Time = 14.67 (sec) , antiderivative size = 71314, normalized size of antiderivative = 195.92

$$\int \frac{\cot^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

```
input int((cot(c + d*x)^3*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)
```

```
output (((a + b*tan(c + d*x))^3*(35*A*b^6 + 62*A*a^2*b^4 + 11*A*a^4*b^2 - 40*B*a^3*b^3 - 20*B*a*b^5 - 4*B*a^5*b))/(4*(a^8 + a^4*b^4 + 2*a^6*b^2)) - ((a + b*tan(c + d*x))^2*(175*A*b^6 + 310*A*a^2*b^4 + 39*A*a^4*b^2 - 208*B*a^3*b^3 - 100*B*a*b^5 - 12*B*a^5*b))/(12*(a^7 + a^3*b^4 + 2*a^5*b^2)) + (2*(A*b^4 - B*a*b^3))/(3*a*(a^2 + b^2)) + (2*(a + b*tan(c + d*x))*(7*A*b^6 + 13*A*a^2*b^4 - 10*B*a^3*b^3 - 4*B*a*b^5))/(3*a^2*(a^4 + b^4 + 2*a^2*b^2)))/(d*(a + b*tan(c + d*x))^(7/2) - 2*a*d*(a + b*tan(c + d*x))^(5/2) + a^2*d*(a + b*tan(c + d*x))^(3/2)) + atan((((a + b*tan(c + d*x))^(1/2)*(321126400*A^4*a^28*b^44*d^5 + 2422210560*A^4*a^30*b^42*d^5 + 1411383296*A^4*a^32*b^40*d^5 - 54989422592*A^4*a^34*b^38*d^5 - 325864914944*A^4*a^36*b^36*d^5 - 1011294928896*A^4*a^38*b^34*d^5 - 2054783238144*A^4*a^40*b^32*d^5 - 2923490705408*A^4*a^42*b^30*d^5 - 2962565365760*A^4*a^44*b^28*d^5 - 2094150975488*A^4*a^46*b^26*d^5 - 943762440192*A^4*a^48*b^24*d^5 - 175655354368*A^4*a^50*b^22*d^5 + 74523344896*A^4*a^52*b^20*d^5 + 62081990656*A^4*a^54*b^18*d^5 + 17307795456*A^4*a^56*b^16*d^5 + 1629487104*A^4*a^58*b^14*d^5 + 44302336*A^4*a^60*b^12*d^5 + 104857600*A^4*a^62*b^10*d^5 + 25165824*A^4*a^64*b^8*d^5 - 104857600*B^4*a^30*b^42*d^5 - 838860800*B^4*a^32*b^40*d^5 - 838860800*B^4*a^34*b^38*d^5 + 17624465408*B^4*a^36*b^36*d^5 + 114621939712*B^4*a^38*b^34*d^5 + 382445027328*B^4*a^40*b^32*d^5 + 842753114112*B^4*a^42*b^30*d^5 + 1327925035008*B^4*a^44*b^28*d^5 + 1546246946816*B^4*a^46*b^26*d^5 + 1344...
```

3.365 $\int \frac{aB+bB \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$

3.365.1 Optimal result 3475
 3.365.2 Mathematica [C] (verified) 3476
 3.365.3 Rubi [A] (warning: unable to verify) 3476
 3.365.4 Maple [B] (verified) 3480
 3.365.5 Fricas [A] (verification not implemented) 3481
 3.365.6 Sympy [F] 3482
 3.365.7 Maxima [F(-2)] 3482
 3.365.8 Giac [F(-1)] 3483
 3.365.9 Mupad [B] (verification not implemented) 3483

3.365.1 Optimal result

Integrand size = 28, antiderivative size = 362

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx$$

$$= \frac{bB \operatorname{arctanh}\left(\frac{\sqrt{a+\sqrt{a^2+b^2}}-\sqrt{2}\sqrt{a+b \tan(c+dx)}}{\sqrt{a-\sqrt{a^2+b^2}}}\right)}{\sqrt{2}\sqrt{a-\sqrt{a^2+b^2}}d} - \frac{bB \operatorname{arctanh}\left(\frac{\sqrt{a+\sqrt{a^2+b^2}}+\sqrt{2}\sqrt{a+b \tan(c+dx)}}{\sqrt{a-\sqrt{a^2+b^2}}}\right)}{\sqrt{2}\sqrt{a-\sqrt{a^2+b^2}}d}$$

$$+ \frac{bB \log\left(a + \sqrt{a^2 + b^2} + b \tan(c + dx) - \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}}\sqrt{a + b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}}d}$$

$$- \frac{bB \log\left(a + \sqrt{a^2 + b^2} + b \tan(c + dx) + \sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}}\sqrt{a + b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{a + \sqrt{a^2 + b^2}}d}$$

output

```
1/2*b*B*arctanh(((a+(a^2+b^2)^(1/2))^(1/2)-2^(1/2)*(a+b*tan(d*x+c))^(1/2))
/(a-(a^2+b^2)^(1/2))^(1/2))/d*2^(1/2)/(a-(a^2+b^2)^(1/2))^(1/2)-1/2*b*B*ar
ctanh(((a+(a^2+b^2)^(1/2))^(1/2)+2^(1/2)*(a+b*tan(d*x+c))^(1/2))/(a-(a^2+b
^2)^(1/2))^(1/2))/d*2^(1/2)/(a-(a^2+b^2)^(1/2))^(1/2)+1/4*b*B*ln(a+(a^2+b^
2)^(1/2)-2^(1/2)*(a+(a^2+b^2)^(1/2))^(1/2)*(a+b*tan(d*x+c))^(1/2)+b*tan(d*
x+c))/d*2^(1/2)/(a+(a^2+b^2)^(1/2))^(1/2)-1/4*b*B*ln(a+(a^2+b^2)^(1/2)+2^(
1/2)*(a+(a^2+b^2)^(1/2))^(1/2)*(a+b*tan(d*x+c))^(1/2)+b*tan(d*x+c))/d*2^(1
/2)/(a+(a^2+b^2)^(1/2))^(1/2)
```

3.365.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.24

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx$$

$$= -\frac{iB \left(\sqrt{a - ib} \operatorname{arctanh} \left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a - ib}} \right) - \sqrt{a + ib} \operatorname{arctanh} \left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a + ib}} \right) \right)}{d}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/Sqrt[a + b*Tan[c + d*x]],x]`

output `((-I)*B*(Sqrt[a - I*b]*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] - Sqrt[a + I*b]*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]]))/d`

3.365.3 Rubi [A] (warning: unable to verify)

Time = 0.61 (sec) , antiderivative size = 403, normalized size of antiderivative = 1.11, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.393$, Rules used = {2011, 3042, 3966, 483, 1449, 1142, 25, 27, 1083, 219, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx$$

$$\downarrow \text{2011}$$

$$B \int \sqrt{a + b \tan(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$B \int \sqrt{a + b \tan(c + dx)} dx$$

$$\downarrow \text{3966}$$

$$\frac{bB \int \frac{\sqrt{a + b \tan(c + dx)}}{\tan^2(c + dx) b^2 + b^2} d(b \tan(c + dx))}{d}$$

$$\downarrow \text{483}$$

$$2bB \int \frac{b^2 \tan^2(c+dx)}{b^4 \tan^4(c+dx) - 2ab^2 \tan^2(c+dx) + a^2 + b^2} d\sqrt{a + b \tan(c + dx)}$$

↓ 1449

$$2bB \left(\frac{\int \frac{\sqrt{a+b \tan(c+dx)}}{b^2 \tan^2(c+dx) - \sqrt{2}b\sqrt{a+\sqrt{a^2+b^2}} \tan(c+dx) + \sqrt{a^2+b^2}} d\sqrt{a+b \tan(c+dx)}}{2\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}} - \frac{\int \frac{\sqrt{a+b \tan(c+dx)}}{b^2 \tan^2(c+dx) + \sqrt{2}b\sqrt{a+\sqrt{a^2+b^2}} \tan(c+dx) + \sqrt{a^2+b^2}} d\sqrt{a+b \tan(c+dx)}}{2\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}} \right)$$

↓ 1142

$$2bB \left(\frac{\sqrt{\sqrt{a^2+b^2}+a} \int \frac{1}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}} \sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}(\sqrt{a+\sqrt{a^2+b^2}} - \sqrt{2}\sqrt{a+b \tan(c+dx)})}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}} \sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{2\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}} \right)$$

↓ 25

$$2bB \left(\frac{\sqrt{\sqrt{a^2+b^2}+a} \int \frac{1}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}} \sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{\sqrt{2}} - \frac{1}{2} \int \frac{\sqrt{2}(\sqrt{a+\sqrt{a^2+b^2}} - \sqrt{2}\sqrt{a+b \tan(c+dx)})}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}} \sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{2\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}} \right)$$

↓ 27

$$2bB \left(\frac{\sqrt{\sqrt{a^2+b^2}+a} \int \frac{1}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}} \sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{\sqrt{2}} - \frac{\int \frac{\sqrt{a+\sqrt{a^2+b^2}} - \sqrt{2}\sqrt{a+b \tan(c+dx)}}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}} \sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{\sqrt{2}}}{2\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}} \right)$$

↓ 1083

$$2bB \left(\frac{-\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a} \int \frac{1}{2(a-\sqrt{a^2+b^2}) - b^2 \tan^2(c+dx)} d(2\sqrt{a+b \tan(c+dx)} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}})}{2\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}} - \frac{\int \frac{\sqrt{a+\sqrt{a^2+b^2}} - \sqrt{2}\sqrt{a+b \tan(c+dx)}}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}} \sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{\sqrt{2}}}{2\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}} \right)$$

↓ 219

3.365. $\int \frac{aB + bB \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$

$$2bB \left(\frac{\int \frac{\sqrt{a+\sqrt{a^2+b^2}}-\sqrt{2}\sqrt{a+b \tan(c+dx)}}{b^2 \tan^2(c+dx)+\sqrt{a^2+b^2}-\sqrt{2}\sqrt{a+\sqrt{a^2+b^2}}\sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)} - \frac{\sqrt{\sqrt{a^2+b^2}+a} \operatorname{arctanh}\left(\frac{2\sqrt{a+b \tan(c+dx)}-\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}}{\sqrt{2}\sqrt{a-\sqrt{a^2+b^2}}}\right)}{\sqrt{a-\sqrt{a^2+b^2}}}}{2\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}} \right)$$

↓ 1103

$$2bB \left(\frac{\frac{1}{2} \log\left(-\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}\sqrt{a+b \tan(c+dx)}+\sqrt{a^2+b^2}+b^2 \tan^2(c+dx)\right) - \frac{\sqrt{\sqrt{a^2+b^2}+a} \operatorname{arctanh}\left(\frac{2\sqrt{a+b \tan(c+dx)}-\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}}{\sqrt{2}\sqrt{a-\sqrt{a^2+b^2}}}\right)}{\sqrt{a-\sqrt{a^2+b^2}}}}{2\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}} \right)$$

input `Int[(a*B + b*B*Tan[c + d*x])/Sqrt[a + b*Tan[c + d*x]],x]`

output `(2*b*B*((-(Sqrt[a + Sqrt[a^2 + b^2]]*ArcTanh[(-(Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]) + 2*Sqrt[a + b*Tan[c + d*x]])/(Sqrt[2]*Sqrt[a - Sqrt[a^2 + b^2]])])/Sqrt[a - Sqrt[a^2 + b^2]]) + Log[Sqrt[a^2 + b^2] + b^2*Tan[c + d*x]^2 - Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]*Sqrt[a + b*Tan[c + d*x]]]/2)/(2*Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]) - ((Sqrt[a + Sqrt[a^2 + b^2]]*ArcTanh[(Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]] + 2*Sqrt[a + b*Tan[c + d*x]])/(Sqrt[2]*Sqrt[a - Sqrt[a^2 + b^2]])])/Sqrt[a - Sqrt[a^2 + b^2]]) + Log[Sqrt[a^2 + b^2] + b^2*Tan[c + d*x]^2 + Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]*Sqrt[a + b*Tan[c + d*x]]]/2)/(2*Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]])))/d`

3.365.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

3.365. $\int \frac{aB+bB \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$

- rule 483 `Int[Sqrt[(c_) + (d_)*(x_)]/((a_) + (b_)*(x_)^2), x_Symbol] := Simp[2*d
Subst[Int[x^2/(b*c^2 + a*d^2 - 2*b*c*x^2 + b*x^4), x], x, Sqrt[c + d*x]], x
] /; FreeQ[{a, b, c, d}, x]`
- rule 1083 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int
1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x]`
- rule 1103 `Int[((d_) + (e_)*(x_)]/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1142 `Int[((d_) + (e_)*(x_)]/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`
- rule 1449 `Int[(x_)^(m_)]/((a_) + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol] := With[{q =
Rt[a/c, 2]}, With[{r = Rt[2*q - b/c, 2]}, Simp[1/(2*c*r) Int[x^(m - 1)/(q
- r*x + x^2), x], x] - Simp[1/(2*c*r) Int[x^(m - 1)/(q + r*x + x^2), x],
x]]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && GeQ[m, 1] && LtQ[m,
3] && NegQ[b^2 - 4*a*c]`
- rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x
] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x
, a + b*x])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`
- rule 3966 `Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Su
bst[Int[(a + x)^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b, c
, d, n}, x] && NeQ[a^2 + b^2, 0]`

3.365.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 661 vs. 2(291) = 582.

Time = 0.12 (sec) , antiderivative size = 662, normalized size of antiderivative = 1.83

method	result
derivativedivides	$-\frac{\ln\left(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}\right) B\sqrt{a^2+b^2} \sqrt{2\sqrt{a^2+b^2}+2a}}{4db} + \frac{B(a^2+b^2) \arctan\left(\frac{\sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}}{d}\right)}{d}$
default	$-\frac{\ln\left(b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}\right) B\sqrt{a^2+b^2} \sqrt{2\sqrt{a^2+b^2}+2a}}{4db} + \frac{B(a^2+b^2) \arctan\left(\frac{\sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}}{d}\right)}{d}$
parts	Expression too large to display

input `int((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & -1/4/d/b*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2) \\
 & + (a^2+b^2)^(1/2))*B*(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/d \\
 & *B/b*(a^2+b^2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan((2*(a+b*\tan(d*x+c))^(1/2) \\
 & + (2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))+1/4/d/b* \\
 & \ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2) \\
 & *B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-1/d*B/b*a^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2) \\
 & *\arctan((2*(a+b*\tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2) \\
 & +1/4/d/b*\ln((a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*\tan(d*x+c)-a-(a^2+b^2)^(1/2) \\
 & *B*(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/d*B/b*(a^2+b^2)/(2*(a^2+b^2)^(1/2)-2*a) \\
 & ^{(1/2)*\arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*\tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2) \\
 & -1/4/d/b*\ln((a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*\tan(d*x+c)-a-(a^2+b^2)^(1/2) \\
 & *B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+1/d*B/b*a^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan(((2*(a^2+b^2)^(1/2) \\
 & +2*a)^(1/2)-2*(a+b*\tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))
 \end{aligned}$$

3.365.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 399, normalized size of antiderivative = 1.10

$$\begin{aligned}
\int \frac{aB + bB \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = & -\frac{1}{2} \sqrt{-\frac{B^2 a + \sqrt{-\frac{B^4 b^2}{d^4}} d^2}{d^2}} \log \left(\sqrt{b \tan(dx + c) + a} B^3 b \right. \\
& \left. + \sqrt{-\frac{B^4 b^2}{d^4}} d^3 \sqrt{-\frac{B^2 a + \sqrt{-\frac{B^4 b^2}{d^4}} d^2}{d^2}} \right) \\
& + \frac{1}{2} \sqrt{-\frac{B^2 a + \sqrt{-\frac{B^4 b^2}{d^4}} d^2}{d^2}} \log \left(\sqrt{b \tan(dx + c) + a} B^3 b \right. \\
& \left. - \sqrt{-\frac{B^4 b^2}{d^4}} d^3 \sqrt{-\frac{B^2 a + \sqrt{-\frac{B^4 b^2}{d^4}} d^2}{d^2}} \right) \\
& + \frac{1}{2} \sqrt{-\frac{B^2 a - \sqrt{-\frac{B^4 b^2}{d^4}} d^2}{d^2}} \log \left(\sqrt{b \tan(dx + c) + a} B^3 b \right. \\
& \left. + \sqrt{-\frac{B^4 b^2}{d^4}} d^3 \sqrt{-\frac{B^2 a - \sqrt{-\frac{B^4 b^2}{d^4}} d^2}{d^2}} \right) \\
& - \frac{1}{2} \sqrt{-\frac{B^2 a - \sqrt{-\frac{B^4 b^2}{d^4}} d^2}{d^2}} \log \left(\sqrt{b \tan(dx + c) + a} B^3 b \right. \\
& \left. - \sqrt{-\frac{B^4 b^2}{d^4}} d^3 \sqrt{-\frac{B^2 a - \sqrt{-\frac{B^4 b^2}{d^4}} d^2}{d^2}} \right)
\end{aligned}$$

```
input integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="fracas")
```



```
output -1/2*sqrt(-(B^2*a + sqrt(-B^4*b^2/d^4)*d^2)/d^2)*log(sqrt(b*tan(d*x + c) +
a)*B^3*b + sqrt(-B^4*b^2/d^4)*d^3*sqrt(-(B^2*a + sqrt(-B^4*b^2/d^4)*d^2)/
d^2)) + 1/2*sqrt(-(B^2*a + sqrt(-B^4*b^2/d^4)*d^2)/d^2)*log(sqrt(b*tan(d*x
+ c) + a)*B^3*b - sqrt(-B^4*b^2/d^4)*d^3*sqrt(-(B^2*a + sqrt(-B^4*b^2/d^4
)*d^2)/d^2)) + 1/2*sqrt(-(B^2*a - sqrt(-B^4*b^2/d^4)*d^2)/d^2)*log(sqrt(b*
tan(d*x + c) + a)*B^3*b + sqrt(-B^4*b^2/d^4)*d^3*sqrt(-(B^2*a - sqrt(-B^4*
b^2/d^4)*d^2)/d^2)) - 1/2*sqrt(-(B^2*a - sqrt(-B^4*b^2/d^4)*d^2)/d^2)*log(
sqrt(b*tan(d*x + c) + a)*B^3*b - sqrt(-B^4*b^2/d^4)*d^3*sqrt(-(B^2*a - sqr
t(-B^4*b^2/d^4)*d^2)/d^2))
```

3.365.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = B \int \sqrt{a + b \tan(c + dx)} dx$$

```
input integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)
```

```
output B*Integral(sqrt(a + b*tan(c + d*x)), x)
```

3.365.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Exception raised: ValueError}$$

```
input integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="maxima
")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more
details)Is
```

3.365.8 Giac [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output Timed out

3.365.9 Mupad [B] (verification not implemented)

Time = 9.98 (sec) , antiderivative size = 3033, normalized size of antiderivative = 8.38

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `int((B*a + B*b*tan(c + d*x))/(a + b*tan(c + d*x))^(1/2),x)`

output `2*atanh((8*a*b^2*(a + b*tan(c + d*x))^(1/2)*(- (-16*B^4*a^4*b^2*d^4)^(1/2) / (16*(a^2*d^4 + b^2*d^4)) - (B^2*a^3*d^2)/(4*(a^2*d^4 + b^2*d^4)))^(1/2)*(-16*B^4*a^4*b^2*d^4)^(1/2))/((16*B^3*a^4*b^5*d^5)/(a^2*d^4 + b^2*d^4) + (16*B^3*a^6*b^3*d^5)/(a^2*d^4 + b^2*d^4) + (4*B*a^3*b^3*d^4*(-16*B^4*a^4*b^2*d^4)^(1/2))/(a^2*d^5 + b^2*d^5) + (4*B*a*b^5*d^4*(-16*B^4*a^4*b^2*d^4)^(1/2))/(a^2*d^5 + b^2*d^5)) - (32*B^2*a^2*b^2*(a + b*tan(c + d*x))^(1/2)*(- (-16*B^4*a^4*b^2*d^4)^(1/2) / (16*(a^2*d^4 + b^2*d^4)) - (B^2*a^3*d^2)/(4*(a^2*d^4 + b^2*d^4)))^(1/2))/((16*B^3*a^4*b^3*d^3)/(a^2*d^4 + b^2*d^4) + (4*B*a*b^3*d^2*(-16*B^4*a^4*b^2*d^4)^(1/2))/(a^2*d^5 + b^2*d^5)) + (32*B^2*a^4*b^2*d^2*(a + b*tan(c + d*x))^(1/2)*(- (-16*B^4*a^4*b^2*d^4)^(1/2) / (16*(a^2*d^4 + b^2*d^4)) - (B^2*a^3*d^2)/(4*(a^2*d^4 + b^2*d^4)))^(1/2))/((16*B^3*a^4*b^5*d^5)/(a^2*d^4 + b^2*d^4) + (16*B^3*a^6*b^3*d^5)/(a^2*d^4 + b^2*d^4) + (4*B*a^3*b^3*d^4*(-16*B^4*a^4*b^2*d^4)^(1/2))/(a^2*d^5 + b^2*d^5) + (4*B*a*b^5*d^4*(-16*B^4*a^4*b^2*d^4)^(1/2))/(a^2*d^5 + b^2*d^5)))*(- (-16*B^4*a^4*b^2*d^4)^(1/2) / (16*(a^2*d^4 + b^2*d^4)) - (B^2*a^3*d^2)/(4*(a^2*d^4 + b^2*d^4)))^(1/2) - 2*atanh((8*a*b^2*((-16*B^4*b^6*d^4)^(1/2) / (16*(a^2*d^4 + b^2*d^4)) + (B^2*a*b^2*d^2)/(4*(a^2*d^4 + b^2*d^4)))^(1/2)*(a + b*tan(c + d*x))^(1/2)*(-16*B^4*b^6*d^4)^(1/2))/((16*B^3*a^2*b^7*d^5)/(a^2*d^4 + b^2*d^4) - 16*B^3*a^2*b^5*d - 16*B^3*b^7*d + (16*B^3*a^4*b^5*d^5)/(a^2*d^4 + b^2*d^4) + (4*B*a^3*b^3*d^4*(-16*B^4*b^6*d^4)^(1/2))/(a^2*d^5 + b^...`

3.366 $\int \frac{aB+bB \tan(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx$

3.366.1 Optimal result	3484
3.366.2 Mathematica [C] (verified)	3485
3.366.3 Rubi [A] (warning: unable to verify)	3485
3.366.4 Maple [B] (verified)	3489
3.366.5 Fricas [B] (verification not implemented)	3490
3.366.6 Sympy [F]	3491
3.366.7 Maxima [F(-2)]	3491
3.366.8 Giac [F(-1)]	3492
3.366.9 Mupad [B] (verification not implemented)	3492

3.366.1 Optimal result

Integrand size = 28, antiderivative size = 406

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \frac{bB \operatorname{arctanh}\left(\frac{\sqrt{a+\sqrt{a^2+b^2}}-\sqrt{2}\sqrt{a+b \tan(c+dx)}}{\sqrt{a-\sqrt{a^2+b^2}}}\right)}{\sqrt{2}\sqrt{a^2+b^2}\sqrt{a-\sqrt{a^2+b^2}}d} - \frac{bB \operatorname{arctanh}\left(\frac{\sqrt{a+\sqrt{a^2+b^2}}+\sqrt{2}\sqrt{a+b \tan(c+dx)}}{\sqrt{a-\sqrt{a^2+b^2}}}\right)}{\sqrt{2}\sqrt{a^2+b^2}\sqrt{a-\sqrt{a^2+b^2}}d} - \frac{bB \log\left(a + \sqrt{a^2+b^2} + b \tan(c + dx) - \sqrt{2}\sqrt{a + \sqrt{a^2+b^2}}\sqrt{a + b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{a + \sqrt{a^2+b^2}}d} + \frac{bB \log\left(a + \sqrt{a^2+b^2} + b \tan(c + dx) + \sqrt{2}\sqrt{a + \sqrt{a^2+b^2}}\sqrt{a + b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{a + \sqrt{a^2+b^2}}d}$$

```
output 1/2*b*B*arctanh(((a+(a^2+b^2)^(1/2))^(1/2)-2^(1/2)*(a+b*tan(d*x+c))^(1/2))
/(a-(a^2+b^2)^(1/2))^(1/2))/d*2^(1/2)/(a^2+b^2)^(1/2)/(a-(a^2+b^2)^(1/2))^(
(1/2)-1/2*b*B*arctanh(((a+(a^2+b^2)^(1/2))^(1/2)+2^(1/2)*(a+b*tan(d*x+c))^(
1/2))/(a-(a^2+b^2)^(1/2))^(1/2))/d*2^(1/2)/(a^2+b^2)^(1/2)/(a-(a^2+b^2)^(
1/2))^(1/2)-1/4*b*B*ln(a+(a^2+b^2)^(1/2)-2^(1/2)*(a+(a^2+b^2)^(1/2))^(1/2)
*(a+b*tan(d*x+c))^(1/2)+b*tan(d*x+c))/d*2^(1/2)/(a^2+b^2)^(1/2)/(a+(a^2+b
^2)^(1/2))^(1/2)+1/4*b*B*ln(a+(a^2+b^2)^(1/2)+2^(1/2)*(a+(a^2+b^2)^(1/2))^(
1/2)*(a+b*tan(d*x+c))^(1/2)+b*tan(d*x+c))/d*2^(1/2)/(a^2+b^2)^(1/2)/(a+(a
^2+b^2)^(1/2))^(1/2)
```

3.366.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.04 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.22

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = -\frac{iB \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ib}} \right)}{d}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(a + b*Tan[c + d*x])^(3/2),x]`

output `((-I)*B*(ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]]/Sqrt[a - I*b] - ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]]/Sqrt[a + I*b]))/d`

3.366.3 Rubi [A] (warning: unable to verify)

Time = 0.66 (sec) , antiderivative size = 426, normalized size of antiderivative = 1.05, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.393$, Rules used = {2011, 3042, 3966, 484, 1407, 1142, 25, 27, 1083, 219, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx \\ & \quad \downarrow \text{2011} \\ & B \int \frac{1}{\sqrt{a + b \tan(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & B \int \frac{1}{\sqrt{a + b \tan(c + dx)}} dx \\ & \quad \downarrow \text{3966} \\ & \frac{bB \int \frac{1}{\sqrt{a + b \tan(c + dx)} (\tan^2(c + dx) b^2 + b^2)} d(b \tan(c + dx))}{d} \\ & \quad \downarrow \text{484} \end{aligned}$$

$$\begin{aligned}
 & \frac{2bB \int \frac{1}{b^4 \tan^4(c+dx) - 2ab^2 \tan^2(c+dx) + a^2 + b^2} d\sqrt{a+b \tan(c+dx)}}{d} \\
 & \quad \downarrow 1407 \\
 & 2bB \left(\frac{\int \frac{\sqrt{2}\sqrt{a+\sqrt{a^2+b^2}} - \sqrt{a+b \tan(c+dx)}}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}}\sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{\sqrt{a^2+b^2}+a}} + \frac{\int \frac{\sqrt{2}\sqrt{a+\sqrt{a^2+b^2}} + \sqrt{a+b \tan(c+dx)}}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} + \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}}\sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{\sqrt{a^2+b^2}+a}} \right) \\
 & \quad \downarrow 1142 \\
 & 2bB \left(\frac{\int \frac{\sqrt{\sqrt{a^2+b^2}+a} \int \frac{1}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}}\sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{\sqrt{2}}}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{\sqrt{a^2+b^2}+a}} - \frac{1}{2} \int \frac{\sqrt{2}(\sqrt{a+\sqrt{a^2+b^2}} - \sqrt{2}\sqrt{a+b \tan(c+dx)})}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}}\sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{\sqrt{a^2+b^2}+a}} \right) \\
 & \quad \downarrow 25 \\
 & 2bB \left(\frac{\int \frac{\sqrt{\sqrt{a^2+b^2}+a} \int \frac{1}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}}\sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{\sqrt{2}}}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{\sqrt{a^2+b^2}+a}} + \frac{1}{2} \int \frac{\sqrt{2}(\sqrt{a+\sqrt{a^2+b^2}} - \sqrt{2}\sqrt{a+b \tan(c+dx)})}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}}\sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{\sqrt{a^2+b^2}+a}} \right) \\
 & \quad \downarrow 27 \\
 & 2bB \left(\frac{\int \frac{\sqrt{\sqrt{a^2+b^2}+a} \int \frac{1}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}}\sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{\sqrt{2}}}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{\sqrt{a^2+b^2}+a}} + \frac{\int \frac{\sqrt{a+\sqrt{a^2+b^2}} - \sqrt{2}\sqrt{a+b \tan(c+dx)}}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}}\sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{\sqrt{2}}}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{\sqrt{a^2+b^2}+a}} \right) \\
 & \quad \downarrow 1083 \\
 & 2bB \left(\frac{\int \frac{\sqrt{a+\sqrt{a^2+b^2}} - \sqrt{2}\sqrt{a+b \tan(c+dx)}}{b^2 \tan^2(c+dx) + \sqrt{a^2+b^2} - \sqrt{2}\sqrt{a+\sqrt{a^2+b^2}}\sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)}}{\sqrt{2}} - \sqrt{2}\sqrt{\sqrt{a^2+b^2}+a} \int \frac{1}{2(a-\sqrt{a^2+b^2}) - b^2 \tan^2(c+dx)} d(2\sqrt{a+b \tan(c+dx)})}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{\sqrt{a^2+b^2}+a}} \right) \\
 & \quad \downarrow 219
 \end{aligned}$$

3.366. $\int \frac{aB+bB \tan(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx$

$$2bB \left(\frac{\int \frac{\sqrt{a+\sqrt{a^2+b^2}}-\sqrt{2}\sqrt{a+b \tan(c+dx)}}{b^2 \tan^2(c+dx)+\sqrt{a^2+b^2}-\sqrt{2}\sqrt{a+\sqrt{a^2+b^2}}\sqrt{a+b \tan(c+dx)}} d\sqrt{a+b \tan(c+dx)} - \frac{\sqrt{\sqrt{a^2+b^2}+a} \operatorname{arctanh}\left(\frac{2\sqrt{a+b \tan(c+dx)}-\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}}{\sqrt{2}\sqrt{a-\sqrt{a^2+b^2}}}\right)}{\sqrt{a-\sqrt{a^2+b^2}}}}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{\sqrt{a^2+b^2}+a}} \right) +$$

↓ 1103

$$2bB \left(-\frac{\frac{\sqrt{\sqrt{a^2+b^2}+a} \operatorname{arctanh}\left(\frac{2\sqrt{a+b \tan(c+dx)}-\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}}{\sqrt{2}\sqrt{a-\sqrt{a^2+b^2}}}\right)}{\sqrt{a-\sqrt{a^2+b^2}}}}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{\sqrt{a^2+b^2}+a}} - \frac{1}{2} \log\left(-\sqrt{2}\sqrt{\sqrt{a^2+b^2}+a}\sqrt{a+b \tan(c+dx)}+\sqrt{a^2+b^2}+b^2 \tan^2(c+dx)\right)}{2\sqrt{2}\sqrt{a^2+b^2}\sqrt{\sqrt{a^2+b^2}+a}} \right) +$$

input `Int[(a*B + b*B*Tan[c + d*x])/(a + b*Tan[c + d*x])^(3/2), x]`

output `(2*b*B*((-((Sqrt[a + Sqrt[a^2 + b^2]]*ArcTanh[(-(Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]) + 2*Sqrt[a + b*Tan[c + d*x]])/(Sqrt[2]*Sqrt[a - Sqrt[a^2 + b^2]])])]/Sqrt[a - Sqrt[a^2 + b^2]]) - Log[Sqrt[a^2 + b^2] + b^2*Tan[c + d*x]^2 - Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]*Sqrt[a + b*Tan[c + d*x]]/2)/(2*Sqrt[2]*Sqrt[a^2 + b^2]*Sqrt[a + Sqrt[a^2 + b^2]]) + (-((Sqrt[a + Sqrt[a^2 + b^2]]*ArcTanh[(Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]] + 2*Sqrt[a + b*Tan[c + d*x]])/(Sqrt[2]*Sqrt[a - Sqrt[a^2 + b^2]])])]/Sqrt[a - Sqrt[a^2 + b^2]]) + Log[Sqrt[a^2 + b^2] + b^2*Tan[c + d*x]^2 + Sqrt[2]*Sqrt[a + Sqrt[a^2 + b^2]]*Sqrt[a + b*Tan[c + d*x]]/2)/(2*Sqrt[2]*Sqrt[a^2 + b^2]*Sqrt[a + Sqrt[a^2 + b^2]])))/d`

3.366.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

- rule 219 $\text{Int}[(a_ + (b_ \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])) \cdot \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[a, 2])], x] \text{ ; FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 484 $\text{Int}[1/(\text{Sqrt}[(c_ + (d_ \cdot x)] \cdot ((a_ + (b_ \cdot x)^2))), x_Symbol] \rightarrow \text{Simp}[2 \cdot d \ \text{Subst}[\text{Int}[1/(b \cdot c^2 + a \cdot d^2 - 2 \cdot b \cdot c \cdot x^2 + b \cdot x^4), x], x, \text{Sqrt}[c + d \cdot x]], x] \text{ ; FreeQ}\{a, b, c, d\}, x]$
- rule 1083 $\text{Int}[(a_ + (b_ \cdot x) + (c_ \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[-2 \ \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4 \cdot a \cdot c - x^2, x], x], x, b + 2 \cdot c \cdot x], x] \text{ ; FreeQ}\{a, b, c, x\}$
- rule 1103 $\text{Int}[(d_ + (e_ \cdot x))/(a_ + (b_ \cdot x) + (c_ \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] \text{ ; FreeQ}\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$
- rule 1142 $\text{Int}[(d_ + (e_ \cdot x))/(a_ + (b_ \cdot x) + (c_ \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[(2 \cdot c \cdot d - b \cdot e)/(2 \cdot c) \ \text{Int}[1/(a + b \cdot x + c \cdot x^2), x], x] + \text{Simp}[e/(2 \cdot c) \ \text{Int}[(b + 2 \cdot c \cdot x)/(a + b \cdot x + c \cdot x^2), x], x] \text{ ; FreeQ}\{a, b, c, d, e\}, x]$
- rule 1407 $\text{Int}[(a_ + (b_ \cdot x)^2 + (c_ \cdot x)^4)^{-1}, x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[a/c, 2]\}, \text{With}\{r = \text{Rt}[2 \cdot q - b/c, 2]\}, \text{Simp}[1/(2 \cdot c \cdot q \cdot r) \ \text{Int}[(r - x)/(q - r \cdot x + x^2), x], x] + \text{Simp}[1/(2 \cdot c \cdot q \cdot r) \ \text{Int}[(r + x)/(q + r \cdot x + x^2), x], x]] \text{ ; FreeQ}\{a, b, c, x\} \ \&\& \ \text{NeQ}[b^2 - 4 \cdot a \cdot c, 0] \ \&\& \ \text{NegQ}[b^2 - 4 \cdot a \cdot c]$
- rule 2011 $\text{Int}[(u_ \cdot ((a_ + (b_ \cdot v))^m) \cdot ((c_ + (d_ \cdot v))^n), x_Symbol] \rightarrow \text{Simp}[(b/d)^m \ \text{Int}[u \cdot (c + d \cdot v)^{m+n}, x], x] \text{ ; FreeQ}\{a, b, c, d, n\}, x] \ \&\& \ \text{EqQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ (!\text{IntegerQ}[n] \ || \ \text{SimplerQ}[c + d \cdot x, a + b \cdot x])$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$

```
rule 3966 Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[b/d Subst[Int[(a + x)^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[a^2 + b^2, 0]
```

3.366.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1574 vs. $2(327) = 654$.

Time = 0.12 (sec) , antiderivative size = 1575, normalized size of antiderivative = 3.88

method	result	size
derivativdivides	Expression too large to display	1575
default	Expression too large to display	1575
parts	Expression too large to display	3686

```
input int((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,method=_RETURNVERBOSE)
```

```
output 1/4/d/b/(a^2+b^2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2)*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2+1/4/d*b/(a^2+b^2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2)*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/4/d/b/(a^2+b^2)^(3/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2)*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3-1/4/d*b/(a^2+b^2)^(3/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2)*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+3/d*b/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*a^2+2/d*b^3/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B-1/d/b/(a^2+b^2)^(1/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*a^2-1/d*b/(a^2+b^2)^(1/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B+1/d/b/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*B*a^4-1/4/d/b/(a^2+b^2)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*B*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2-1/...
```


3.366.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 805 vs. $2(329) = 658$.

Time = 0.26 (sec) , antiderivative size = 805, normalized size of antiderivative = 1.98

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \frac{1}{2} \sqrt{-\frac{(a^2 + b^2) \sqrt{-\frac{B^4 b^2}{(a^4 + 2a^2 b^2 + b^4) d^4} d^2} + B^2 a}{(a^2 + b^2) d^2}} \log \left(\sqrt{b \tan(dx + c) + a} B^3 b \right)$$

$$+ \left(B^2 b^2 d + (a^3 + ab^2) \sqrt{-\frac{B^4 b^2}{(a^4 + 2a^2 b^2 + b^4) d^4} d^3} \right) \sqrt{-\frac{(a^2 + b^2) \sqrt{-\frac{B^4 b^2}{(a^4 + 2a^2 b^2 + b^4) d^4} d^2} + B^2 a}{(a^2 + b^2) d^2}}$$

$$- \frac{1}{2} \sqrt{-\frac{(a^2 + b^2) \sqrt{-\frac{B^4 b^2}{(a^4 + 2a^2 b^2 + b^4) d^4} d^2} + B^2 a}{(a^2 + b^2) d^2}} \log \left(\sqrt{b \tan(dx + c) + a} B^3 b \right)$$

$$- \left(B^2 b^2 d + (a^3 + ab^2) \sqrt{-\frac{B^4 b^2}{(a^4 + 2a^2 b^2 + b^4) d^4} d^3} \right) \sqrt{-\frac{(a^2 + b^2) \sqrt{-\frac{B^4 b^2}{(a^4 + 2a^2 b^2 + b^4) d^4} d^2} + B^2 a}{(a^2 + b^2) d^2}}$$

$$+ \frac{1}{2} \sqrt{\frac{(a^2 + b^2) \sqrt{-\frac{B^4 b^2}{(a^4 + 2a^2 b^2 + b^4) d^4} d^2} - B^2 a}{(a^2 + b^2) d^2}} \log \left(\sqrt{b \tan(dx + c) + a} B^3 b \right)$$

$$+ \left(B^2 b^2 d - (a^3 + ab^2) \sqrt{-\frac{B^4 b^2}{(a^4 + 2a^2 b^2 + b^4) d^4} d^3} \right) \sqrt{\frac{(a^2 + b^2) \sqrt{-\frac{B^4 b^2}{(a^4 + 2a^2 b^2 + b^4) d^4} d^2} - B^2 a}{(a^2 + b^2) d^2}}$$

$$- \frac{1}{2} \sqrt{\frac{(a^2 + b^2) \sqrt{-\frac{B^4 b^2}{(a^4 + 2a^2 b^2 + b^4) d^4} d^2} - B^2 a}{(a^2 + b^2) d^2}} \log \left(\sqrt{b \tan(dx + c) + a} B^3 b \right)$$

$$- \left(B^2 b^2 d - (a^3 + ab^2) \sqrt{-\frac{B^4 b^2}{(a^4 + 2a^2 b^2 + b^4) d^4} d^3} \right) \sqrt{\frac{(a^2 + b^2) \sqrt{-\frac{B^4 b^2}{(a^4 + 2a^2 b^2 + b^4) d^4} d^2} - B^2 a}{(a^2 + b^2) d^2}}$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="fracas")`

output $1/2*\sqrt{-((a^2 + b^2)*\sqrt{-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)})*d^2 + B^2*a}/((a^2 + b^2)*d^2))*\log(\sqrt{b*\tan(dx + c) + a}*B^3*b + (B^2*b^2*d + (a^3 + a*b^2)*\sqrt{-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)})*d^3)*\sqrt{-((a^2 + b^2)*\sqrt{-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)})*d^2 + B^2*a}/((a^2 + b^2)*d^2))) - 1/2*\sqrt{-((a^2 + b^2)*\sqrt{-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)})*d^2 + B^2*a}/((a^2 + b^2)*d^2))*\log(\sqrt{b*\tan(dx + c) + a}*B^3*b - (B^2*b^2*d + (a^3 + a*b^2)*\sqrt{-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)})*d^3)*\sqrt{-((a^2 + b^2)*\sqrt{-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)})*d^2 + B^2*a}/((a^2 + b^2)*d^2))) + 1/2*\sqrt{((a^2 + b^2)*\sqrt{-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)})*d^2 - B^2*a}/((a^2 + b^2)*d^2))*\log(\sqrt{b*\tan(dx + c) + a}*B^3*b + (B^2*b^2*d - (a^3 + a*b^2)*\sqrt{-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)})*d^3)*\sqrt{((a^2 + b^2)*\sqrt{-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)})*d^2 - B^2*a}/((a^2 + b^2)*d^2))) - 1/2*\sqrt{((a^2 + b^2)*\sqrt{-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)})*d^2 - B^2*a}/((a^2 + b^2)*d^2))*\log(\sqrt{b*\tan(dx + c) + a}*B^3*b - (B^2*b^2*d - (a^3 + a*b^2)*\sqrt{-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)})*d^3)*\sqrt{((a^2 + b^2)*\sqrt{-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)})*d^2 - B^2*a}/((a^2 + b^2)*d^2)))$

3.366.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = B \int \frac{1}{\sqrt{a + b \tan(c + dx)}} dx$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)`

output `B*Integral(1/sqrt(a + b*tan(c + d*x)), x)`

3.366.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is

3.366.8 Giac [**F(-1)**]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="giac")`

output Timed out

3.366.9 Mupad [**B**] (verification not implemented)

Time = 12.47 (sec) , antiderivative size = 6453, normalized size of antiderivative = 15.89

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `int((B*a + B*b*tan(c + d*x))/(a + b*tan(c + d*x))^(3/2),x)`

output $\log\left(\left(a + b \tan(c + dx)\right)^{1/2} \left(16B^2a^2b^{10}d^3 + 32B^2a^4b^8d^3 - 32B^2a^8b^4d^3 - 16B^2a^{10}b^2d^3\right) - \left(\left(\left(8B^2a^5d^2 - 24B^2a^3b^2d^2\right)^{2/4} - B^4a^4(16a^6d^4 + 16b^6d^4 + 48a^2b^4d^4 + 48a^4b^2d^4)\right)^{1/2} - 4B^2a^5d^2 + 12B^2a^3b^2d^2\right) / \left(16(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4)\right)^{1/2} \right) \left(a + b \tan(c + dx)\right)^{1/2} \left(\left(\left(8B^2a^5d^2 - 24B^2a^3b^2d^2\right)^{2/4} - B^4a^4(16a^6d^4 + 16b^6d^4 + 48a^2b^4d^4 + 48a^4b^2d^4)\right)^{1/2} - 4B^2a^5d^2 + 12B^2a^3b^2d^2\right) / \left(16(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4)\right)^{1/2} \right) \left(64ab^{12}d^5 + 320a^3b^{10}d^5 + 640a^5b^8d^5 + 640a^7b^6d^5 + 320a^9b^4d^5 + 64a^{11}b^2d^5\right) + 64B^2a^2b^{11}d^4 + 256B^2a^4b^9d^4 + 384B^2a^6b^7d^4 + 256B^2a^8b^5d^4 + 64B^2a^{10}b^3d^4 \left(\left(\left(8B^2a^5d^2 - 24B^2a^3b^2d^2\right)^{2/4} - B^4a^4(16a^6d^4 + 16b^6d^4 + 48a^2b^4d^4 + 48a^4b^2d^4)\right)^{1/2} - 4B^2a^5d^2 + 12B^2a^3b^2d^2\right) / \left(16(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4)\right)^{1/2} + 8B^3a^3b^9d^2 + 24B^3a^5b^7d^2 + 24B^3a^7b^5d^2 + 8B^3a^9b^3d^2 \left(\left(\left(8B^2a^5d^2 - 24B^2a^3b^2d^2\right)^{2/4} - B^4a^4(16a^6d^4 + 16b^6d^4 + 48a^2b^4d^4 + 48a^4b^2d^4)\right)^{1/2} - 4B^2a^5d^2 + 12B^2a^3b^2d^2\right) / \left(16(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4)\right)^{1/2} \right) - \log\left(8B^3a^3b^9d^2 - \left(-\left(96B^4a^6b^4d^4 - 16B^4a^4b^6d^4 - 144B^4a^8b^2d^4\right)^{1/2} + 4B^2a^5d^2 - 12B^2a^3b^2d^2\right) / \left(16a^6 \dots\right.\right.$

3.367
$$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.367.1 Optimal result	3494
3.367.2 Mathematica [A] (verified)	3494
3.367.3 Rubi [A] (warning: unable to verify)	3495
3.367.4 Maple [B] (verified)	3498
3.367.5 Fricas [B] (verification not implemented)	3499
3.367.6 Sympy [F]	3500
3.367.7 Maxima [F(-1)]	3501
3.367.8 Giac [F(-1)]	3501
3.367.9 Mupad [B] (verification not implemented)	3501

3.367.1 Optimal result

Integrand size = 34, antiderivative size = 119

$$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = -\frac{2B \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ad}} + \frac{B \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ibd}} + \frac{B \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ibd}}$$

output `-2*B*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/d/a^(1/2)+B*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d/(a-I*b)^(1/2)+B*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d/(a+I*b)^(1/2)`

3.367.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.94

$$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{B \left(-\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{a}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ib}} \right)}{d}$$

input `Integrate[(Cot[c + d*x]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2), x]`

output $(B*((-2*\text{ArcTanh}[\text{Sqrt}[a + b*\text{Tan}[c + d*x]]/\text{Sqrt}[a]])/\text{Sqrt}[a] + \text{ArcTanh}[\text{Sqrt}[a + b*\text{Tan}[c + d*x]]/\text{Sqrt}[a - I*b]]/\text{Sqrt}[a - I*b] + \text{ArcTanh}[\text{Sqrt}[a + b*\text{Tan}[c + d*x]]/\text{Sqrt}[a + I*b]]/\text{Sqrt}[a + I*b]))/d$

3.367.3 Rubi [A] (warning: unable to verify)

Time = 0.88 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.91, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {2011, 3042, 4057, 25, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$$

$$\downarrow 2011$$

$$B \int \frac{\cot(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx$$

$$\downarrow 3042$$

$$B \int \frac{1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx$$

$$\downarrow 4057$$

$$B \left(\int -\frac{\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b\tan(c+dx)}} dx \right)$$

$$\downarrow 25$$

$$B \left(\int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b\tan(c+dx)}} dx - \int \frac{\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx \right)$$

$$\downarrow 3042$$

$$B \left(\int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - \int \frac{\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx \right)$$

$$\downarrow 4022$$

$$B \left(\int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b\tan(c+dx)}} dx - \frac{1}{2}i \int \frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}i \int \frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}} dx \right)$$

$$\begin{aligned}
& \downarrow 3042 \\
& B\left(-\frac{1}{2}i \int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}i \int \frac{i \tan(c+dx)+1}{\sqrt{a+b \tan(c+dx)}} dx + \int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx\right) \\
& \downarrow 4020 \\
& B\left(\int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx - \frac{\int -\frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} - \frac{\int -\frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d}\right) \\
& \downarrow 25 \\
& B\left(\int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + \frac{\int \frac{1}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} + \frac{\int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d}\right) \\
& \downarrow 73 \\
& B\left(-\frac{i \int \frac{1}{-\frac{i \tan^2(c+dx)}{b}-\frac{ia}{b}+1} d\sqrt{a+b \tan(c+dx)}}{bd} + \frac{i \int \frac{1}{\frac{i \tan^2(c+dx)}{b}+\frac{ia}{b}+1} d\sqrt{a+b \tan(c+dx)}}{bd} + \int \frac{\tan(c+dx)}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx\right) \\
& \downarrow 221 \\
& B\left(\int \frac{\tan(c+dx)^2+1}{\tan(c+dx)\sqrt{a+b \tan(c+dx)}} dx + \frac{i \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{i \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}}\right) \\
& \downarrow 4117 \\
& B\left(\frac{\int \frac{\cot(c+dx)}{\sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{d} + \frac{i \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{i \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}}\right) \\
& \downarrow 73 \\
& B\left(\frac{2 \int \frac{1}{\frac{a+b \tan(c+dx)}{b}-\frac{a}{b}} d\sqrt{a+b \tan(c+dx)}}{bd} + \frac{i \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{i \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}}\right) \\
& \downarrow 221 \\
& B\left(\frac{i \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{i \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}} - \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ad}}\right)
\end{aligned}$$

input `Int[(Cot[c + d*x]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `B*((I*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d) - (I*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d) - (2*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/(Sqrt[a]*d))`

3.367.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 2011 `Int[(u_.)*((a_.) + (b_.)*(v_)^(m_.))*((c_.) + (d_.)*(v_)^(n_.)), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4057 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)/((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(c^2 + d^2) Int[(a + b*Tan[e + f*x])^m*(c - d*Tan[e + f*x]), x], x] + Simp[d^2/(c^2 + d^2) Int[(a + b*Tan[e + f*x])^m*((1 + Tan[e + f*x]^2)/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

3.367.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 373 vs. 2(97) = 194.

Time = 0.34 (sec) , antiderivative size = 374, normalized size of antiderivative = 3.14

method	result
default	$2Bb^2 \left(\frac{\sqrt{2\sqrt{a^2+b^2}+2a} \ln\left(\frac{b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}}{2}\right) + 2(a-\sqrt{a^2+b^2}) \arctan\left(\frac{2\sqrt{a+b \tan(dx+c)}+\sqrt{2\sqrt{a^2+b^2}+2a}}{\sqrt{2\sqrt{a^2+b^2}-2a}}\right)}{4\sqrt{a^2+b^2}} \right)$

input `int(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

$$3.367. \int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

output $2*B/d*b^2*(1/b^2*(1/4/(a^2+b^2)^{(1/2)}*(1/2*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}+(a^2+b^2)^{(1/2)}))+2*(a-(a^2+b^2)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan((2*(a+b*\tan(d*x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}))+1/4/(a^2+b^2)^{(1/2)}*(-1/2*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}*ln((a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-b*\tan(d*x+c)-a-(a^2+b^2)^{(1/2)}))+2*((a^2+b^2)^{(1/2)}-a)/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan(((2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-2*(a+b*\tan(d*x+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}))-1/b^2/a^{(1/2)}*\operatorname{arctanh}((a+b*\tan(d*x+c))^{(1/2)}/a^{(1/2)})$

3.367.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 829 vs. 2(93) = 186.

Time = 0.29 (sec) , antiderivative size = 1674, normalized size of antiderivative = 14.07

$$\int \frac{\cot(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="fracas")`

output

```

[-1/2*(a*d*sqrt(((a^2 + b^2)*sqrt(-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*
d^2 + B^2*a)/((a^2 + b^2)*d^2))*log(sqrt(b*tan(d*x + c) + a)*B^3 + ((a^2 +
b^2)*sqrt(-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^3 - B^2*a*d)*sqrt(((a
^2 + b^2)*sqrt(-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^2 + B^2*a)/((a^2
+ b^2)*d^2))) - a*d*sqrt(((a^2 + b^2)*sqrt(-B^4*b^2/((a^4 + 2*a^2*b^2 + b^
4)*d^4))*d^2 + B^2*a)/((a^2 + b^2)*d^2))*log(sqrt(b*tan(d*x + c) + a)*B^3
- ((a^2 + b^2)*sqrt(-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^3 - B^2*a*d)
*sqrt(((a^2 + b^2)*sqrt(-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^2 + B^2*
a)/((a^2 + b^2)*d^2))) - a*d*sqrt(-((a^2 + b^2)*sqrt(-B^4*b^2/((a^4 + 2*a^
2*b^2 + b^4)*d^4))*d^2 - B^2*a)/((a^2 + b^2)*d^2))*log(sqrt(b*tan(d*x + c)
+ a)*B^3 + ((a^2 + b^2)*sqrt(-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^3
+ B^2*a*d)*sqrt(-((a^2 + b^2)*sqrt(-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))
*d^2 - B^2*a)/((a^2 + b^2)*d^2))) + a*d*sqrt(-((a^2 + b^2)*sqrt(-B^4*b^2/(
(a^4 + 2*a^2*b^2 + b^4)*d^4))*d^2 - B^2*a)/((a^2 + b^2)*d^2))*log(sqrt(b*t
an(d*x + c) + a)*B^3 - ((a^2 + b^2)*sqrt(-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)
*d^4))*d^3 + B^2*a*d)*sqrt(-((a^2 + b^2)*sqrt(-B^4*b^2/((a^4 + 2*a^2*b^2 +
b^4)*d^4))*d^2 - B^2*a)/((a^2 + b^2)*d^2))) - 2*B*sqrt(a)*log((b*tan(d*x
+ c) - 2*sqrt(b*tan(d*x + c) + a)*sqrt(a) + 2*a)/tan(d*x + c)))/(a*d), -1/
2*(a*d*sqrt(((a^2 + b^2)*sqrt(-B^4*b^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^2
+ B^2*a)/((a^2 + b^2)*d^2))*log(sqrt(b*tan(d*x + c) + a)*B^3 + ((a^2 + ...

```

3.367.6 Sympy [F]

$$\int \frac{\cot(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = B \int \frac{\cot(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)`

output `B*Integral(cot(c + d*x)/sqrt(a + b*tan(c + d*x)), x)`

3.367.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\cot(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `Timed out`

3.367.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cot(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `Timed out`

3.367.9 Mupad [B] (verification not implemented)

Time = 10.11 (sec) , antiderivative size = 2142, normalized size of antiderivative = 18.00

$$\int \frac{\cot(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `int((cot(c + d*x)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x)`

output

```

- atan(((((((32*(16*B*b^10*d^2 + 12*B*a^2*b^8*d^2))/d^3 - (32*(16*b^10*d^4
+ 24*a^2*b^8*d^4)*(a + b*tan(c + d*x))^(1/2)*(B^2/(4*(a*d^2 - b*d^2*1i)))
^(1/2))/d^4)*(B^2/(4*(a*d^2 - b*d^2*1i)))^(1/2) + (576*B^2*a*b^8*(a + b*ta
n(c + d*x))^(1/2))/d^2)*(B^2/(4*(a*d^2 - b*d^2*1i)))^(1/2) - (96*B^3*a*b^8
)/d^3)*(B^2/(4*(a*d^2 - b*d^2*1i)))^(1/2) - (96*B^4*b^8*(a + b*tan(c + d*x
))^(1/2))/d^4)*(B^2/(4*(a*d^2 - b*d^2*1i)))^(1/2)*1i - (((((32*(16*B*b^10*
d^2 + 12*B*a^2*b^8*d^2))/d^3 + (32*(16*b^10*d^4 + 24*a^2*b^8*d^4)*(a + b*ta
n(c + d*x))^(1/2)*(B^2/(4*(a*d^2 - b*d^2*1i)))^(1/2))/d^4)*(B^2/(4*(a*d^2
- b*d^2*1i)))^(1/2) - (576*B^2*a*b^8*(a + b*tan(c + d*x))^(1/2))/d^2)*(B^
2/(4*(a*d^2 - b*d^2*1i)))^(1/2) - (96*B^3*a*b^8)/d^3)*(B^2/(4*(a*d^2 - b*d
^2*1i)))^(1/2) + (96*B^4*b^8*(a + b*tan(c + d*x))^(1/2))/d^4)*(B^2/(4*(a*d
^2 - b*d^2*1i)))^(1/2)*1i)/(((((((32*(16*B*b^10*d^2 + 12*B*a^2*b^8*d^2))/d^
3 - (32*(16*b^10*d^4 + 24*a^2*b^8*d^4)*(a + b*tan(c + d*x))^(1/2)*(B^2/(4*
(a*d^2 - b*d^2*1i)))^(1/2))/d^4)*(B^2/(4*(a*d^2 - b*d^2*1i)))^(1/2) + (576
*B^2*a*b^8*(a + b*tan(c + d*x))^(1/2))/d^2)*(B^2/(4*(a*d^2 - b*d^2*1i)))^(
1/2) - (96*B^3*a*b^8)/d^3)*(B^2/(4*(a*d^2 - b*d^2*1i)))^(1/2) - (96*B^4*b^
8*(a + b*tan(c + d*x))^(1/2))/d^4)*(B^2/(4*(a*d^2 - b*d^2*1i)))^(1/2) + ((
(((32*(16*B*b^10*d^2 + 12*B*a^2*b^8*d^2))/d^3 + (32*(16*b^10*d^4 + 24*a^2*
b^8*d^4)*(a + b*tan(c + d*x))^(1/2)*(B^2/(4*(a*d^2 - b*d^2*1i)))^(1/2))/d^
4)*(B^2/(4*(a*d^2 - b*d^2*1i)))^(1/2) - (576*B^2*a*b^8*(a + b*tan(c + d...

```

3.368 $\int \frac{aB+bB \tan(c+dx)}{(a+b \tan(c+dx))^{5/2}} dx$

3.368.1 Optimal result	3503
3.368.2 Mathematica [C] (verified)	3503
3.368.3 Rubi [A] (warning: unable to verify)	3504
3.368.4 Maple [B] (verified)	3507
3.368.5 Fricas [B] (verification not implemented)	3508
3.368.6 Sympy [F]	3508
3.368.7 Maxima [F(-2)]	3509
3.368.8 Giac [F(-1)]	3509
3.368.9 Mupad [B] (verification not implemented)	3509

3.368.1 Optimal result

Integrand size = 28, antiderivative size = 123

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = -\frac{iB \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a - ib)^{3/2}d} + \frac{iB \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a + ib)^{3/2}d} - \frac{2bB}{(a^2 + b^2) d \sqrt{a + b \tan(c + dx)}}$$

output `-I*B*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(3/2)/d+I*B*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(3/2)/d-2*b*B/(a^2+b^2)/d/(a+b*tan(d*x+c))^(1/2)`

3.368.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.14 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.86

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \frac{B\left(i(a + ib) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b \tan(c+dx)}{a-ib}\right) + (-ia - b) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b \tan(c+dx)}{a+ib}\right)\right)}{(a^2 + b^2) d \sqrt{a + b \tan(c + dx)}}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(a + b*Tan[c + d*x])^(5/2),x]`

output $(B*(I*(a + I*b)*\text{Hypergeometric2F1}[-1/2, 1, 1/2, (a + b*\text{Tan}[c + d*x])/(a - I*b)] + ((-I)*a - b)*\text{Hypergeometric2F1}[-1/2, 1, 1/2, (a + b*\text{Tan}[c + d*x])/(a + I*b)]))/((a^2 + b^2)*d*\text{Sqrt}[a + b*\text{Tan}[c + d*x]])$

3.368.3 Rubi [A] (warning: unable to verify)

Time = 0.59 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.02, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {2011, 3042, 3964, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx \\
 & \quad \downarrow 2011 \\
 & B \int \frac{1}{(a + b \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow 3042 \\
 & B \int \frac{1}{(a + b \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow 3964 \\
 & B \left(\frac{\int \frac{a - b \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} - \frac{2b}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} \right) \\
 & \quad \downarrow 3042 \\
 & B \left(\frac{\int \frac{a - b \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} - \frac{2b}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} \right) \\
 & \quad \downarrow 4022 \\
 & B \left(-\frac{2b}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} + \frac{\frac{1}{2}(a - ib) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a + ib) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} \right) \\
 & \quad \downarrow 3042
 \end{aligned}$$

$$B \left(-\frac{2b}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} + \frac{\frac{1}{2}(a - ib) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a + ib) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} \right)$$

↓ 4020

$$B \left(-\frac{2b}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} + \frac{\frac{i(a+ib) \int -\frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d} - \frac{i(a-ib) \int -\frac{1}{(i \tan(c+dx)+1) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d}}{a^2 + b^2} \right)$$

↓ 25

$$B \left(-\frac{2b}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} + \frac{\frac{i(a-ib) \int \frac{1}{(i \tan(c+dx)+1) \sqrt{a+b \tan(c+dx)}} d(-i \tan(c+dx))}{2d} - \frac{i(a+ib) \int \frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}} d(i \tan(c+dx))}{2d}}{a^2 + b^2} \right)$$

↓ 73

$$B \left(-\frac{2b}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} + \frac{\frac{(a-ib) \int \frac{1}{-i \tan^2(c+dx) - \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd} + \frac{(a+ib) \int \frac{1}{i \tan^2(c+dx) + \frac{ia}{b} + 1} d\sqrt{a+b \tan(c+dx)}}{bd}}{a^2 + b^2} \right)$$

↓ 221

$$B \left(-\frac{2b}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} + \frac{\frac{(a+ib) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} + \frac{(a-ib) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}}}{a^2 + b^2} \right)$$

input `Int[(a*B + b*B*Tan[c + d*x])/(a + b*Tan[c + d*x])^(5/2), x]`

output `B*(((a + I*b)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d) + ((a - I*b)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d))/(a^2 + b^2) - (2*b)/((a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])`

3.368.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x
] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x
, a + b*x])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`
- rule 3964 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((a +
b*Tan[c + d*x])^(n + 1)/(d*(n + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2)
Int[(a - b*Tan[c + d*x])*(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a,
b, c, d}, x] && NeQ[a^2 + b^2, 0] && LtQ[n, -1]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

3.368.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1954 vs. $2(103) = 206$.

Time = 0.09 (sec) , antiderivative size = 1955, normalized size of antiderivative = 15.89

method	result	size
derivativedivides	Expression too large to display	1955
default	Expression too large to display	1955
parts	Expression too large to display	4475

input `int((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/4/d*B/b/(a^2+b^2)^2*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2) \\ &)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3+1/4/ \\ & d*B*b/(a^2+b^2)^2*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1 \\ & /2)+2*a)^(1/2)+(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-1/4/d*B/b/ \\ & (a^2+b^2)^(5/2)*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2) \\ &)+2*a)^(1/2)+(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^4+1/4/d*B*b^ \\ & 3/(a^2+b^2)^(5/2)*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1 \\ & /2)+2*a)^(1/2)+(a^2+b^2)^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+4/d*B*b/(a^2 \\ & +b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan((2*(a+b*\tan(d*x+c))^(1/2) \\ & +(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^3+3/d*B*b \\ & ^3/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan((2*(a+b*\tan(d*x+c) \\ &)^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a-1/ \\ & d*B/b/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan((2*(a+b*\tan(d*x \\ & +c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a \\ & ^3-1/d*B*b/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan((2*(a+b*\ta \\ & n(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/ \\ & 2))*a-1/d*B*b/(a^2+b^2)^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan((2*(a+b*\tan \\ & (d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2) \\ &))*a^2+1/d*B/b/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan((2*(a+ \\ & b*\tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2... \end{aligned}$$

3.368.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2174 vs. $2(97) = 194$.

Time = 0.31 (sec) , antiderivative size = 2174, normalized size of antiderivative = 17.67

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="fracas")`

output `-1/2*(4*sqrt(b*tan(d*x + c) + a)*B*b + ((a^2*b + b^3)*d*tan(d*x + c) + (a^3 + a*b^2)*d)*sqrt(-(B^2*a^3 - 3*B^2*a*b^2 + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*sqrt(-(9*B^4*a^4*b^2 - 6*B^4*a^2*b^4 + B^4*b^6)/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 + b^12)*d^4)))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2))*log(-(3*B^3*a^2*b - B^3*b^3)*sqrt(b*tan(d*x + c) + a) + ((a^8 + 2*a^6*b^2 - 2*a^2*b^6 - b^8)*d^3*sqrt(-(9*B^4*a^4*b^2 - 6*B^4*a^2*b^4 + B^4*b^6)/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 + b^12)*d^4)) + 2*(3*B^2*a^3*b^2 - B^2*a*b^4)*d)*sqrt(-(B^2*a^3 - 3*B^2*a*b^2 + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*sqrt(-(9*B^4*a^4*b^2 - 6*B^4*a^2*b^4 + B^4*b^6)/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 + b^12)*d^4)))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2))) - ((a^2*b + b^3)*d*tan(d*x + c) + (a^3 + a*b^2)*d)*sqrt(-(B^2*a^3 - 3*B^2*a*b^2 + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*sqrt(-(9*B^4*a^4*b^2 - 6*B^4*a^2*b^4 + B^4*b^6)/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 + b^12)*d^4)))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2))*log(-(3*B^3*a^2*b - B^3*b^3)*sqrt(b*tan(d*x + c) + a) - ((a^8 + 2*a^6*b^2 - 2*a^2*b^6 - b^8)*d^3*sqrt(-(9*B^4*a^4*b^2 - 6*B^4*a^2*b^4 + B^4*b^6)/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 + b^12)*d^4)) + 2*(3*B^2*a^3*b^2 - B^2*a*b^4)*d)*sqrt(-(B^2*a^3 - 3*B^2*a*b^2 + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*sqrt(-(9*B^4*a^4*b^2 - 6*B^4*a^2*b^4 + B^4*b^6)/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 + b^12)*d^4)))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2)))`

3.368.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = B \int \frac{1}{a\sqrt{a + b \tan(c + dx)} + b\sqrt{a + b \tan(c + dx)} \tan(c + dx)} dx$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

3.368. $\int \frac{aB+bB \tan(c+dx)}{(a+b \tan(c+dx))^{5/2}} dx$

output `B*Integral(1/(a*sqrt(a + b*tan(c + d*x)) + b*sqrt(a + b*tan(c + d*x))*tan(c + d*x)), x)`

3.368.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is`

3.368.8 Giac [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Timed out`

3.368.9 Mupad [B] (verification not implemented)

Time = 21.21 (sec) , antiderivative size = 9618, normalized size of antiderivative = 78.20

$$\int \frac{aB + bB \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `int((B*a + B*b*tan(c + d*x))/(a + b*tan(c + d*x))^(5/2),x)`

output $(\log(16*B^3*a^4*b^15*d^2 - (((320*B^4*a^6*b^8*d^4 - 16*B^4*a^4*b^10*d^4 - 1760*B^4*a^8*b^6*d^4 + 1600*B^4*a^10*b^4*d^4 - 400*B^4*a^12*b^2*d^4)^{(1/2)} - 4*B^2*a^7*d^2 - 20*B^2*a^3*b^4*d^2 + 40*B^2*a^5*b^2*d^2)/(a^{10*d^4} + b^{10*d^4} + 5*a^2*b^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^{(1/2)} * (((320*B^4*a^6*b^8*d^4 - 16*B^4*a^4*b^10*d^4 - 1760*B^4*a^8*b^6*d^4 + 1600*B^4*a^10*b^4*d^4 - 400*B^4*a^12*b^2*d^4)^{(1/2)} - 4*B^2*a^7*d^2 - 20*B^2*a^3*b^4*d^2 + 40*B^2*a^5*b^2*d^2)/(a^{10*d^4} + b^{10*d^4} + 5*a^2*b^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^{(1/2)} * (((320*B^4*a^6*b^8*d^4 - 16*B^4*a^4*b^10*d^4 - 1760*B^4*a^8*b^6*d^4 + 1600*B^4*a^10*b^4*d^4 - 400*B^4*a^12*b^2*d^4)^{(1/2)} - 4*B^2*a^7*d^2 - 20*B^2*a^3*b^4*d^2 + 40*B^2*a^5*b^2*d^2)/(a^{10*d^4} + b^{10*d^4} + 5*a^2*b^8*d^4 + 10*a^4*b^6*d^4 + 10*a^6*b^4*d^4 + 5*a^8*b^2*d^4))^{(1/2)} * (a + b*\tan(c + d*x))^{(1/2)} * (64*a*b^{22*d^5} + 640*a^3*b^{20*d^5} + 2880*a^5*b^{18*d^5} + 7680*a^7*b^{16*d^5} + 13440*a^9*b^{14*d^5} + 16128*a^{11}*b^{12*d^5} + 13440*a^{13}*b^{10*d^5} + 7680*a^{15}*b^8*d^5 + 2880*a^{17}*b^6*d^5 + 640*a^{19}*b^4*d^5 + 64*a^{21}*b^2*d^5))/4 - 32*B*a*b^{21*d^4} - 160*B*a^3*b^{19*d^4} - 128*B*a^5*b^{17*d^4} + 896*B*a^7*b^{15*d^4} + 3136*B*a^9*b^{13*d^4} + 4928*B*a^{11}*b^{11*d^4} + 4480*B*a^{13}*b^9*d^4 + 2432*B*a^{15}*b^7*d^4 + 736*B*a^{17}*b^5*d^4 + 96*B*a^{19}*b^3*d^4))/4 - (a + b*\tan(c + d*x))^{(1/2)} * (320*B^2*a^6*b^{14*d^3} - 16*B^2*a^2*b^{18*d^3} + 1024*B^2*a^8*b^{12*d^3} + 1440*B^2*a^{10}*b^{10*d^3} + 1024*B^2*a^{12}*b^8*d^3 + 320*B^2...$

3.369
$$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

3.369.1 Optimal result	3511
3.369.2 Mathematica [A] (verified)	3511
3.369.3 Rubi [A] (warning: unable to verify)	3512
3.369.4 Maple [B] (verified)	3516
3.369.5 Fricas [B] (verification not implemented)	3517
3.369.6 Sympy [F]	3518
3.369.7 Maxima [F(-1)]	3519
3.369.8 Giac [F(-1)]	3519
3.369.9 Mupad [B] (verification not implemented)	3519

3.369.1 Optimal result

Integrand size = 34, antiderivative size = 154

$$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx =$$

$$-\frac{2B \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{a^{3/2}d} + \frac{B \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{3/2}d}$$

$$+ \frac{B \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a+ib)^{3/2}d} + \frac{2b^2 B}{a(a^2+b^2)d\sqrt{a+b \tan(c+dx)}}$$

output `-2*B*arctanh((a+b*tan(d*x+c))^(1/2)/a^(1/2))/a^(3/2)/d+B*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(3/2)/d+B*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(3/2)/d+2*b^2*B/a/(a^2+b^2)/d/(a+b*tan(d*x+c))^(1/2)`

3.369.2 Mathematica [A] (verified)

Time = 1.25 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.08

$$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{B \left(-\frac{2(a^2+b^2) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{a}} + \frac{a(a+ib) \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib}} \right)}{a(a^2+b^2)d}$$

input `Integrate[(Cot[c + d*x]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]`

output `(B*((-2*(a^2 + b^2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/Sqrt[a] + (a*(a + I*b)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/Sqrt[a - I*b] + (a*(a - I*b)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/Sqrt[a + I*b] + (2*b^2)/Sqrt[a + b*Tan[c + d*x]]))/(a*(a^2 + b^2)*d)`

3.369.3 Rubi [A] (warning: unable to verify)

Time = 1.24 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.14, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2011, 3042, 4052, 27, 3042, 4136, 25, 3042, 4022, 3042, 4020, 25, 73, 221, 4117, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{\cot(c+dx)}{(a+b\tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{\tan(c+dx)(a+b\tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{4052} \\
 & B \left(\frac{2 \int \frac{\cot(c+dx)(a^2-b\tan(c+dx)a+b^2+b^2\tan^2(c+dx))}{2\sqrt{a+b\tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b^2}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \right) \\
 & \quad \downarrow \text{27} \\
 & B \left(\frac{\int \frac{\cot(c+dx)(a^2-b\tan(c+dx)a+b^2+b^2\tan^2(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b^2}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& B \left(\frac{\int \frac{a^2 - b \tan(c+dx) a + b^2 + b^2 \tan(c+dx)^2}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{a(a^2 + b^2)} + \frac{2b^2}{ad(a^2 + b^2) \sqrt{a+b \tan(c+dx)}} \right) \\
& \quad \downarrow 4136 \\
& B \left(\frac{(a^2 + b^2) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx + \int -\frac{\tan(c+dx)a^2 + ba}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2 + b^2)} + \frac{2b^2}{ad(a^2 + b^2) \sqrt{a+b \tan(c+dx)}} \right) \\
& \quad \downarrow 25 \\
& B \left(\frac{(a^2 + b^2) \int \frac{\cot(c+dx)(\tan^2(c+dx)+1)}{\sqrt{a+b \tan(c+dx)}} dx - \int \frac{\tan(c+dx)a^2 + ba}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2 + b^2)} + \frac{2b^2}{ad(a^2 + b^2) \sqrt{a+b \tan(c+dx)}} \right) \\
& \quad \downarrow 3042 \\
& B \left(\frac{(a^2 + b^2) \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx - \int \frac{\tan(c+dx)a^2 + ba}{\sqrt{a+b \tan(c+dx)}} dx}{a(a^2 + b^2)} + \frac{2b^2}{ad(a^2 + b^2) \sqrt{a+b \tan(c+dx)}} \right) \\
& \quad \downarrow 4022 \\
& B \left(\frac{2b^2}{ad(a^2 + b^2) \sqrt{a+b \tan(c+dx)}} + \frac{(a^2 + b^2) \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2} a(b + ia) \int \frac{1 - i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2} a}{a(a^2 + b^2)} \right) \\
& \quad \downarrow 3042 \\
& B \left(\frac{2b^2}{ad(a^2 + b^2) \sqrt{a+b \tan(c+dx)}} + \frac{(a^2 + b^2) \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2} a(b + ia) \int \frac{1 - i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2} a}{a(a^2 + b^2)} \right) \\
& \quad \downarrow 4020 \\
& B \left(\frac{2b^2}{ad(a^2 + b^2) \sqrt{a+b \tan(c+dx)}} + \frac{(a^2 + b^2) \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx) \sqrt{a+b \tan(c+dx)}} dx + \frac{ia(-b+ia) \int -\frac{1}{(1-i \tan(c+dx)) \sqrt{a+b \tan(c+dx)}}}{2d}}{a(a^2 + b^2)} \right) \\
& \quad \downarrow 25
\end{aligned}$$

$$B \left(\frac{2b^2}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{(a^2 + b^2) \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a + b \tan(c+dx)}} dx - \frac{ia(-b+ia) \int \frac{1}{(1-i \tan(c+dx))\sqrt{a + b \tan(c+dx)}} dx}{2d}}{a(a^2 + b^2)} \right)$$

↓ 73

$$B \left(\frac{2b^2}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{(a^2 + b^2) \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a + b \tan(c+dx)}} dx - \frac{a(b+ia) \int \frac{1}{-i \tan^2(c+dx) - \frac{ia}{b} + 1}}{bd} d\sqrt{a + b \tan(c+dx)}}{a(a^2 + b^2)} \right)$$

↓ 221

$$B \left(\frac{2b^2}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{(a^2 + b^2) \int \frac{\tan(c+dx)^2 + 1}{\tan(c+dx)\sqrt{a + b \tan(c+dx)}} dx + \frac{a(-b+ia) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{a(b+ia) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}}}{a(a^2 + b^2)} \right)$$

↓ 4117

$$B \left(\frac{2b^2}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{\frac{(a^2 + b^2) \int \frac{\cot(c+dx)}{\sqrt{a + b \tan(c+dx)}} d \tan(c+dx)}{d} + \frac{a(-b+ia) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{a(b+ia) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}}}{a(a^2 + b^2)} \right)$$

↓ 73

$$B \left(\frac{2b^2}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{\frac{2(a^2 + b^2) \int \frac{1}{\frac{a + b \tan(c+dx)}{b} - \frac{a}{b}} d\sqrt{a + b \tan(c+dx)}}{bd} + \frac{a(-b+ia) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{a(b+ia) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}}}{a(a^2 + b^2)} \right)$$

↓ 221

$$B \left(\frac{2b^2}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{-\frac{2(a^2 + b^2) \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ad}} + \frac{a(-b+ia) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{a(b+ia) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}}}{a(a^2 + b^2)} \right)$$

input `Int[(Cot[c + d*x]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2), x]`

```
output B*(((a*(I*a - b)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d) - (
a*(I*a + b)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d) - (2*(a^
2 + b^2)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a]])/(Sqrt[a]*d))/(a*(a^2 +
b^2)) + (2*b^2)/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])
```

3.369.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 73 Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n_), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 2011 Int[(u_.)*((a_) + (b_.)*(v_)^(m_.))*((c_) + (d_.)*(v_)^(n_.)), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x
] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x
, a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

```
rule 4052 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m + 1)*((c
+ d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(a^2 + b^2)*(b*c - a*d))), x] + Simp[1
/((m + 1)*(a^2 + b^2)*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c +
d*Tan[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2) - b*(b*c -
a*d)*(m + 1)*Tan[e + f*x] - b^2*d*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /
; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0]
&& NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && LtQ[m, -1] && (LtQ[n, 0] || Integ
erQ[m]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

```
rule 4136 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.369.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1778 vs. 2(130) = 260.

Time = 0.33 (sec) , antiderivative size = 1779, normalized size of antiderivative = 11.55

method	result	size
default	Expression too large to display	1779

3.369.
$$\int \frac{\cot(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$$

input `int(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x,method=_RETURNVERBOSE)`

output `B*(-1/4/d/(a^2+b^2)^2*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^-1/4/d*b^2/(a^2+b^2)^2*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/2/d/(a^2+b^2)^(5/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3+1/2/d*b^2/(a^2+b^2)^(5/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^-2/d*b^4/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))+1/d/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^-1/d/(a^2+b^2)^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^3+1/d*b^2/(a^2+b^2)^(3/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))-1/d*b^2/(a^2+b^2)^2/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^-2/d*b^2/(a^2+b^2)^(5/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^2+1/4/d/(a^2+...`

3.369.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2281 vs. 2(126) = 252.

Time = 0.39 (sec) , antiderivative size = 4578, normalized size of antiderivative = 29.73

$$\int \frac{\cot(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x,algorithm="fricas")`

output

```
[1/2*(4*sqrt(b*tan(d*x + c) + a)*B*a*b^2 + ((a^4*b + a^2*b^3)*d*tan(d*x +
c) + (a^5 + a^3*b^2)*d)*sqrt((B^2*a^3 - 3*B^2*a*b^2 + (a^6 + 3*a^4*b^2 + 3
*a^2*b^4 + b^6)*d^2*sqrt(-(9*B^4*a^4*b^2 - 6*B^4*a^2*b^4 + B^4*b^6)/((a^12
+ 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 + b^12)*
d^4)))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2))*log(-(3*B^3*a^2 - B^3*b^
2)*sqrt(b*tan(d*x + c) + a) + (2*(a^7 + 3*a^5*b^2 + 3*a^3*b^4 + a*b^6)*d^3
*sqrt(-(9*B^4*a^4*b^2 - 6*B^4*a^2*b^4 + B^4*b^6)/((a^12 + 6*a^10*b^2 + 15*
a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 + b^12)*d^4)) - (3*B^2*a^4
- 4*B^2*a^2*b^2 + B^2*b^4)*d)*sqrt((B^2*a^3 - 3*B^2*a*b^2 + (a^6 + 3*a^4*b
^2 + 3*a^2*b^4 + b^6)*d^2*sqrt(-(9*B^4*a^4*b^2 - 6*B^4*a^2*b^4 + B^4*b^6)/
((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 +
b^12)*d^4)))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2))) - ((a^4*b + a^2*b
^3)*d*tan(d*x + c) + (a^5 + a^3*b^2)*d)*sqrt((B^2*a^3 - 3*B^2*a*b^2 + (a^6
+ 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*sqrt(-(9*B^4*a^4*b^2 - 6*B^4*a^2*b^4 +
B^4*b^6)/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a
^2*b^10 + b^12)*d^4)))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2))*log(-(3*
B^3*a^2 - B^3*b^2)*sqrt(b*tan(d*x + c) + a) - (2*(a^7 + 3*a^5*b^2 + 3*a^3*
b^4 + a*b^6)*d^3*sqrt(-(9*B^4*a^4*b^2 - 6*B^4*a^2*b^4 + B^4*b^6)/((a^12 +
6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 + b^12)*d^4
)) - (3*B^2*a^4 - 4*B^2*a^2*b^2 + B^2*b^4)*d)*sqrt((B^2*a^3 - 3*B^2*a*b...
```

3.369.6 Sympy [F]

$$\int \frac{\cot(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = B \int \frac{\cot(c+dx)}{a\sqrt{a+b \tan(c+dx)} + b\sqrt{a+b \tan(c+dx)} \tan(c+dx)}$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output `B*Integral(cot(c + d*x)/(a*sqrt(a + b*tan(c + d*x)) + b*sqrt(a + b*tan(c + d*x))*tan(c + d*x)), x)`

3.369.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\cot(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `Timed out`

3.369.8 Giac [F(-1)]

Timed out.

$$\int \frac{\cot(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Timed out`

3.369.9 Mupad [B] (verification not implemented)

Time = 14.09 (sec) , antiderivative size = 7172, normalized size of antiderivative = 46.57

$$\int \frac{\cot(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `int((cot(c + d*x)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)`

output $(\log(\frac{((96B^4a^2b^4d^4 - 16B^4b^6d^4 - 144B^4a^4b^2d^4)^{1/2} + 4B^2a^3d^2 - 12B^2ab^2d^2)/(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{1/2} * (((((96B^4a^2b^4d^4 - 16B^4b^6d^4 - 144B^4a^4b^2d^4)^{1/2} + 4B^2a^3d^2 - 12B^2ab^2d^2)/(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{1/2} * (512B^8a^8b^28d^8 - (((96B^4a^2b^4d^4 - 16B^4b^6d^4 - 144B^4a^4b^2d^4)^{1/2} + 4B^2a^3d^2 - 12B^2ab^2d^2)/(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{1/2} * (a + b \tan(c + dx))^{1/2} * (512a^9b^28d^9 + 5376a^{11}b^{26}d^9 + 25344a^{13}b^{24}d^9 + 70656a^{15}b^{22}d^9 + 129024a^{17}b^{20}d^9 + 161280a^{19}b^{18}d^9 + 139776a^{21}b^{16}d^9 + 82944a^{23}b^{14}d^9 + 32256a^{25}b^{12}d^9 + 7424a^{27}b^{10}d^9 + 768a^{29}b^8d^9)) / 4 + 5248B^8a^{10}b^{26}d^8 + 23936B^8a^{12}b^{24}d^8 + 64000B^8a^{14}b^{22}d^8 + 111104B^8a^{16}b^{20}d^8 + 130816B^8a^{18}b^{18}d^8 + 105728B^8a^{20}b^{16}d^8 + 57856B^8a^{22}b^{14}d^8 + 20480B^8a^{24}b^{12}d^8 + 4224B^8a^{26}b^{10}d^8 + 384B^8a^{28}b^8d^8)) / 4 + (a + b \tan(c + dx))^{1/2} * (256B^2a^8b^{26}d^7 + 1472B^2a^{10}b^{24}d^7 + 3712B^2a^{12}b^{22}d^7 + 6272B^2a^{14}b^{20}d^7 + 9856B^2a^{16}b^{18}d^7 + 14336B^2a^{18}b^{16}d^7 + 15232B^2a^{20}b^{14}d^7 + 10112B^2a^{22}b^{12}d^7 + 3712B^2a^{24}b^{10}d^7 + 576B^2a^{26}b^8d^7)) * (((96B^4a^2b^4d^4 - 16B^4b^6d^4 - 144B^4a^4b^2d^4)^{1/2} + 4B^2a^3d^2 - 12B^2ab^2d^2)/(a^6d^4 + b^6d^4 + 3a^2b^4d^4 + 3a^4b^2d^4))^{1/2}) / 4 - \dots$

$$3.370 \quad \int \frac{-a+b \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$$

3.370.1 Optimal result	3521
3.370.2 Mathematica [A] (verified)	3521
3.370.3 Rubi [A] (warning: unable to verify)	3522
3.370.4 Maple [B] (verified)	3524
3.370.5 Fricas [B] (verification not implemented)	3525
3.370.6 Sympy [F]	3525
3.370.7 Maxima [F(-2)]	3526
3.370.8 Giac [F(-1)]	3526
3.370.9 Mupad [B] (verification not implemented)	3526

3.370.1 Optimal result

Integrand size = 27, antiderivative size = 102

$$\begin{aligned} & \int \frac{-a+b \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx \\ &= \frac{(ia-b)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ibd}} - \frac{(ia+b)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ibd}} \end{aligned}$$

output $(I*a-b)*\operatorname{arctanh}((a+b*\tan(d*x+c))^{1/2}/(a-I*b)^{1/2})/d/(a-I*b)^{1/2}-(I*a+b)*\operatorname{arctanh}((a+b*\tan(d*x+c))^{1/2}/(a+I*b)^{1/2})/d/(a+I*b)^{1/2}$

3.370.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.07

$$\begin{aligned} & \int \frac{-a+b \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx \\ &= \frac{i\left((a+ib)^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right) - (a-ib)^{3/2}\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)\right)}{\sqrt{a-ib}\sqrt{a+ibd}} \end{aligned}$$

input $\operatorname{Integrate}[(-a + b*\operatorname{Tan}[c + d*x])/Sqrt[a + b*\operatorname{Tan}[c + d*x]],x]$

$$3.370. \quad \int \frac{-a+b \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$$

output $(I*((a + I*b)^{(3/2)}*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]] - (a - I*b)^{(3/2)}*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/(Sqrt[a - I*b]*Sqrt[a + I*b]*d)$

3.370.3 Rubi [A] (warning: unable to verify)

Time = 0.41 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.83, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.259$, Rules used = {3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{b \tan(c + dx) - a}{\sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{b \tan(c + dx) - a}{\sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4022} \\
 & -\frac{1}{2}(a - ib) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx - \frac{1}{2}(a + ib) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{2}(a - ib) \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx - \frac{1}{2}(a + ib) \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4020} \\
 & \frac{i(a - ib) \int -\frac{1}{(i \tan(c + dx) + 1)\sqrt{a + b \tan(c + dx)}} d(-i \tan(c + dx))}{2d} - \\
 & \frac{i(a + ib) \int -\frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} \\
 & \quad \downarrow \text{25} \\
 & \frac{i(a + ib) \int \frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} - \\
 & \frac{i(a - ib) \int \frac{1}{(i \tan(c + dx) + 1)\sqrt{a + b \tan(c + dx)}} d(-i \tan(c + dx))}{2d} \\
 & \quad \downarrow \text{73}
 \end{aligned}$$

$$\frac{(a - ib) \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)}}{bd} - \frac{(a + ib) \int \frac{1}{\frac{i \tan^2(c+dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)}}{bd}$$

↓ 221

$$-\frac{(a + ib) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a-ib}}\right)}{d\sqrt{a-ib}} - \frac{(a - ib) \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}}$$

input `Int[(-a + b*Tan[c + d*x])/Sqrt[a + b*Tan[c + d*x]],x]`

output `-((a + I*b)*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]]/(Sqrt[a - I*b]*d)) - ((a - I*b)*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]]/(Sqrt[a + I*b]*d))`

3.370.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

3.370.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1892 vs. $2(84) = 168$.

Time = 0.09 (sec) , antiderivative size = 1893, normalized size of antiderivative = 18.56

method	result	size
parts	Expression too large to display	1893
derivativedivides	Expression too large to display	1905
default	Expression too large to display	1905

```
input int((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x,method=_RETURNVERBOSE)
```

```
output b/d*(1/2/(a^2+b^2)^(1/2)*(-1/2*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*ln(b*tan(d*x+
c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))
+2*((a^2+b^2)^(1/2)-a)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*
x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)))
+1/2/(a^2+b^2)^(1/2)*(1/2*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*ln(b*tan(d*x+c)+a-
(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))+2*((
a^2+b^2)^(1/2)-a)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))
^(1/2)-(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))))-a*(
1/4/d/b/(a^2+b^2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1
/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2+1/4/d*b/
(a^2+b^2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)
^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-1/4/d/b/(a^2+b^2)^(3
/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)
+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3-1/4/d*b/(a^2+b^2)^(3/2)
)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(
a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-1/d/b/(a^2+b^2)^(1/2)/(2*(
a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1
/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^2-1/d*b/(a^2+b^2)^(1/2)/(
2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)
^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))+1/d/b/(a^2+b^2)^(3/2)...
```

3.370.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1115 vs. $2(78) = 156$.

Time = 0.26 (sec) , antiderivative size = 1115, normalized size of antiderivative = 10.93

$$\int \frac{-a + b \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

```
input integrate((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="fricas")
```

```
output 1/2*sqrt(-((a^2 + b^2)*d^2*sqrt(-(9*a^4*b^2 - 6*a^2*b^4 + b^6)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + a^3 - 3*a*b^2)/((a^2 + b^2)*d^2))*log(-(3*a^4*b + 2*a^2*b^3 - b^5)*sqrt(b*tan(d*x + c) + a) + ((a^4 - b^4)*d^3*sqrt(-(9*a^4*b^2 - 6*a^2*b^4 + b^6)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*(3*a^3*b^2 - a*b^4)*d)*sqrt(-((a^2 + b^2)*d^2*sqrt(-(9*a^4*b^2 - 6*a^2*b^4 + b^6)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + a^3 - 3*a*b^2)/((a^2 + b^2)*d^2))) - 1/2*sqrt(-((a^2 + b^2)*d^2*sqrt(-(9*a^4*b^2 - 6*a^2*b^4 + b^6)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + a^3 - 3*a*b^2)/((a^2 + b^2)*d^2))*log(-(3*a^4*b + 2*a^2*b^3 - b^5)*sqrt(b*tan(d*x + c) + a) - ((a^4 - b^4)*d^3*sqrt(-(9*a^4*b^2 - 6*a^2*b^4 + b^6)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + 2*(3*a^3*b^2 - a*b^4)*d)*sqrt(-((a^2 + b^2)*d^2*sqrt(-(9*a^4*b^2 - 6*a^2*b^4 + b^6)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) + a^3 - 3*a*b^2)/((a^2 + b^2)*d^2))) - 1/2*sqrt(((a^2 + b^2)*d^2*sqrt(-(9*a^4*b^2 - 6*a^2*b^4 + b^6)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) - a^3 + 3*a*b^2)/((a^2 + b^2)*d^2))*log(-(3*a^4*b + 2*a^2*b^3 - b^5)*sqrt(b*tan(d*x + c) + a) + ((a^4 - b^4)*d^3*sqrt(-(9*a^4*b^2 - 6*a^2*b^4 + b^6)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) - 2*(3*a^3*b^2 - a*b^4)*d)*sqrt(((a^2 + b^2)*d^2*sqrt(-(9*a^4*b^2 - 6*a^2*b^4 + b^6)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) - a^3 + 3*a*b^2)/((a^2 + b^2)*d^2))) + 1/2*sqrt(((a^2 + b^2)*d^2*sqrt(-(9*a^4*b^2 - 6*a^2*b^4 + b^6)/((a^4 + 2*a^2*b^2 + b^4)*d^4)) - a^3 + 3*a*b^2)/((a^2 + b^2)*d^2))*log(-(3*a^4*b + 2*a^2*b^3 - b^5)*sqrt(b*tan(d*x + c) + a) - ...
```

3.370.6 Sympy [F]

$$\int \frac{-a + b \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = - \int \frac{a}{\sqrt{a + b \tan(c + dx)}} dx - \int \left(- \frac{b \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} \right) dx$$

```
input integrate((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)
```

3.370. $\int \frac{-a + b \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx$

output `-Integral(a/sqrt(a + b*tan(c + d*x)), x) - Integral(-b*tan(c + d*x)/sqrt(a + b*tan(c + d*x)), x)`

3.370.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{-a + b \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Exception raised: ValueError}$$

input `integrate((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is`

3.370.8 Giac [F(-1)]

Timed out.

$$\int \frac{-a + b \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.370.9 Mupad [B] (verification not implemented)

Time = 10.32 (sec) , antiderivative size = 2731, normalized size of antiderivative = 26.77

$$\int \frac{-a + b \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `int(-(a - b*tan(c + d*x))/(a + b*tan(c + d*x))^(1/2),x)`

output $2*\operatorname{atanh}\left(\frac{32*a^2*b^2*((-16*a^4*b^2*d^4)^{1/2}/(16*(a^2*d^4 + b^2*d^4)) - (a^3*d^2)/(4*(a^2*d^4 + b^2*d^4)))^{1/2}*(a + b*\tan(c + d*x))^{1/2}}{(16*a^4*b^3*d^3)/(a^2*d^4 + b^2*d^4) - (4*a*b^3*d^2*(-16*a^4*b^2*d^4)^{1/2})/(a^2*d^5 + b^2*d^5)} + (8*a*b^2*((-16*a^4*b^2*d^4)^{1/2}/(16*(a^2*d^4 + b^2*d^4)) - (a^3*d^2)/(4*(a^2*d^4 + b^2*d^4)))^{1/2}*(a + b*\tan(c + d*x))^{1/2}*(-16*a^4*b^2*d^4)^{1/2}\right) / \left(\frac{16*a^4*b^5*d^5}{a^2*d^4 + b^2*d^4} + \frac{16*a^6*b^3*d^5}{a^2*d^4 + b^2*d^4} - \frac{4*a^3*b^3*d^4*(-16*a^4*b^2*d^4)^{1/2}}{a^2*d^5 + b^2*d^5} - \frac{4*a*b^5*d^4*(-16*a^4*b^2*d^4)^{1/2}}{a^2*d^5 + b^2*d^5} \right) - \left(\frac{32*a^4*b^2*d^2*((-16*a^4*b^2*d^4)^{1/2}/(16*(a^2*d^4 + b^2*d^4)) - (a^3*d^2)/(4*(a^2*d^4 + b^2*d^4)))^{1/2}*(a + b*\tan(c + d*x))^{1/2}}{(16*a^4*b^5*d^5)/(a^2*d^4 + b^2*d^4) + \frac{16*a^6*b^3*d^5}{a^2*d^4 + b^2*d^4} - \frac{4*a^3*b^3*d^4*(-16*a^4*b^2*d^4)^{1/2}}{a^2*d^5 + b^2*d^5} - \frac{4*a*b^5*d^4*(-16*a^4*b^2*d^4)^{1/2}}{a^2*d^5 + b^2*d^5}} \right) - 2*\operatorname{atanh}\left(\frac{32*a^2*b^4*d^2*((-16*b^6*d^4)^{1/2}/(16*(a^2*d^4 + b^2*d^4)) + (a*b^2*d^2)/(4*(a^2*d^4 + b^2*d^4)))^{1/2}*(a + b*\tan(c + d*x))^{1/2}}{(16*a^2*b^7*d^5)/(a^2*d^4 + b^2*d^4) - 16*a^2*b^5*d - 16*b^7*d + \frac{16*a^4*b^5*d^5}{a^2*d^4 + b^2*d^4} + \frac{4*a*b^5*d^4*(-16*b^6*d^4)^{1/2}}{a^2*d^5 + b^2*d^5} + \frac{4*a^3*b^3*d^4*(-16*b^6*d^4)^{1/2}}{a^2*d^5 + b^2*d^5}} - \frac{32*b^4*((-16*b^6*d^4)^{1/2}/(16*(a^2*d^4 + b^2*d^4)) + (a*b^2*d^2)/(4*(a^2*d^4 + b^2*d^4)))^{1/2}}{(16*(a^2*d^4 + b^2*d^4)) + (a*b^2*d^2)/(4*(a^2*d^4 + b^2*d^4))} \right)$

3.371
$$\int \frac{-a+b \tan(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx$$

3.371.1 Optimal result 3528
 3.371.2 Mathematica [C] (verified) 3528
 3.371.3 Rubi [A] (warning: unable to verify) 3529
 3.371.4 Maple [B] (verified) 3532
 3.371.5 Fricas [B] (verification not implemented) 3533
 3.371.6 Sympy [F] 3533
 3.371.7 Maxima [F(-2)] 3534
 3.371.8 Giac [F(-1)] 3534
 3.371.9 Mupad [B] (verification not implemented) 3535

3.371.1 Optimal result

Integrand size = 27, antiderivative size = 132

$$\int \frac{-a+b \tan(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx = \frac{(ia-b)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{3/2}d} - \frac{(ia+b)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a+ib)^{3/2}d} + \frac{4ab}{(a^2+b^2)d\sqrt{a+b \tan(c+dx)}}$$

```
output (I*a-b)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(3/2)/d-(I*a+b)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(3/2)/d+4*a*b/(a^2+b^2)/d/(a+b*tan(d*x+c))^(1/2)
```

3.371.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.43 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.17

$$\int \frac{-a+b \tan(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx = \frac{i \cos(c+dx) \left((a+ib)^2 \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b \tan(c+dx)}{a-ib}\right) - (a-ib)^2 \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b \tan(c+dx)}{a+ib}\right) \right)}{(a-ib)(a+ib)d(a \cos(c+dx) - b \sin(c+dx))\sqrt{a+b \tan(c+dx)}}$$

input `Integrate[(-a + b*Tan[c + d*x])/(a + b*Tan[c + d*x])^(3/2),x]`

output `((-I)*Cos[c + d*x]*((a + I*b)^2*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Tan[c + d*x])/(a - I*b)] - (a - I*b)^2*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Tan[c + d*x])/(a + I*b)])*(a - b*Tan[c + d*x])/((a - I*b)*(a + I*b)*d*(a *Cos[c + d*x] - b*Sin[c + d*x])*Sqrt[a + b*Tan[c + d*x]])`

3.371.3 Rubi [A] (warning: unable to verify)

Time = 0.61 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.98, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.370$, Rules used = {3042, 4012, 25, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{b \tan(c + dx) - a}{(a + b \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{b \tan(c + dx) - a}{(a + b \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{4012} \\
 & \frac{\int -\frac{a^2 - 2b \tan(c + dx)a - b^2}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} + \frac{4ab}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} \\
 & \quad \downarrow \text{25} \\
 & \frac{4ab}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} - \frac{\int \frac{a^2 - 2b \tan(c + dx)a - b^2}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{4ab}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} - \frac{\int \frac{a^2 - 2b \tan(c + dx)a - b^2}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} \\
 & \quad \downarrow \text{4022} \\
 & \frac{4ab}{d(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} - \frac{\frac{1}{2}(a - ib)^2 \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a + ib)^2 \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & \frac{4ab}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} - \frac{\frac{1}{2}(a - ib)^2 \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a + ib)^2 \int \frac{i \tan(c + dx) + 1}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} \\
 & \downarrow \text{4020} \\
 & \frac{4ab}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} - \frac{i(a + ib)^2 \int -\frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx)) - \frac{1}{(i \tan(c + dx) + 1)\sqrt{a + b \tan(c + dx)}} d(-i \tan(c + dx))}{2d} \frac{1}{a^2 + b^2} \\
 & \downarrow \text{25} \\
 & \frac{4ab}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} - \frac{i(a - ib)^2 \int \frac{1}{(i \tan(c + dx) + 1)\sqrt{a + b \tan(c + dx)}} d(-i \tan(c + dx)) - \frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} \frac{1}{a^2 + b^2} \\
 & \downarrow \text{73} \\
 & \frac{4ab}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} - \frac{(a - ib)^2 \int -\frac{1}{\frac{i \tan^2(c + dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)} + \frac{(a + ib)^2 \int \frac{1}{\frac{i \tan^2(c + dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)}}{bd} \frac{1}{a^2 + b^2} \\
 & \downarrow \text{221} \\
 & \frac{4ab}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} - \frac{(a - ib)^2 \arctan\left(\frac{\tan(c + dx)}{\sqrt{a + ib}}\right)}{d\sqrt{a + ib}} + \frac{(a + ib)^2 \arctan\left(\frac{\tan(c + dx)}{\sqrt{a - ib}}\right)}{d\sqrt{a - ib}} \frac{1}{a^2 + b^2}
 \end{aligned}$$

input `Int[(-a + b*Tan[c + d*x])/(a + b*Tan[c + d*x])^(3/2),x]`

output `-(((a + I*b)^2*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d) + ((a - I*b)^2*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d)/(a^2 + b^2) + (4*a*b)/((a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])`

3.371.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]`
- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`
- rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/
(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x]
)^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1
]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

3.371.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2290 vs. $2(112) = 224$.

Time = 0.12 (sec) , antiderivative size = 2291, normalized size of antiderivative = 17.36

method	result	size
derivativedivides	Expression too large to display	2291
default	Expression too large to display	2291
parts	Expression too large to display	3685

input `int((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & 4*a*b/(a^2+b^2)/d/(a+b*\tan(d*x+c))^{(1/2)}+2/d*b^5/(a^2+b^2)^{(5/2)}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan((2*(a+b*\tan(d*x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2))}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})-2/d*b^5/(a^2+b^2)^{(5/2)}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan(((2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-2*(a+b*\tan(d*x+c))^{(1/2))}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})+1/d*b^3/(a^2+b^2)^{(3/2)}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan(((2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-2*(a+b*\tan(d*x+c))^{(1/2))}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})+1/4/d*b^3/(a^2+b^2)^2*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}+(a^2+b^2)^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-1/4/d*b^3/(a^2+b^2)^2*\ln((a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-b*\tan(d*x+c)-a-(a^2+b^2)^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}-1/d*b^3/(a^2+b^2)^{(3/2)}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan((2*(a+b*\tan(d*x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2))}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})+1/d/b/(a^2+b^2)^{(3/2)}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan((2*(a+b*\tan(d*x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2))}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*a^4-1/d*b^3/(a^2+b^2)^{(5/2)}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan((2*(a+b*\tan(d*x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2))}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*a^2-1/d/b/(a^2+b^2)^{(5/2)}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)}*\arctan((2*(a+b*\tan(d*x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2))}/(2*(a^2+b^2)^{(1/2)}-2*a)^{(1/2)})*a^6-1/4/d/b/(a^2+b^2)^2*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*a)^{(1/2)}+(a^2+b^2)^{(1/2)}...
 \end{aligned}$$

3.371.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2320 vs. 2(106) = 212.

Time = 0.29 (sec) , antiderivative size = 2320, normalized size of antiderivative = 17.58

$$\int \frac{-a + b \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="fricas")`

output

```
1/2*(8*sqrt(b*tan(d*x + c) + a)*a*b - ((a^2*b + b^3)*d*tan(d*x + c) + (a^3
+ a*b^2)*d)*sqrt(-(a^5 - 10*a^3*b^2 + 5*a*b^4 + (a^6 + 3*a^4*b^2 + 3*a^2*
b^4 + b^6)*d^2*sqrt(-(25*a^8*b^2 - 100*a^6*b^4 + 110*a^4*b^6 - 20*a^2*b^8
+ b^10)/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2
*b^10 + b^12)*d^4)))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2))*log((5*a^6
*b - 5*a^4*b^3 - 9*a^2*b^5 + b^7)*sqrt(b*tan(d*x + c) + a) + ((a^9 - 6*a^5
*b^4 - 8*a^3*b^6 - 3*a*b^8)*d^3*sqrt(-(25*a^8*b^2 - 100*a^6*b^4 + 110*a^4*
b^6 - 20*a^2*b^8 + b^10)/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 1
5*a^4*b^8 + 6*a^2*b^10 + b^12)*d^4)) + (15*a^6*b^2 - 35*a^4*b^4 + 13*a^2*b
^6 - b^8)*d)*sqrt(-(a^5 - 10*a^3*b^2 + 5*a*b^4 + (a^6 + 3*a^4*b^2 + 3*a^2*
b^4 + b^6)*d^2*sqrt(-(25*a^8*b^2 - 100*a^6*b^4 + 110*a^4*b^6 - 20*a^2*b^8
+ b^10)/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20*a^6*b^6 + 15*a^4*b^8 + 6*a^2
*b^10 + b^12)*d^4)))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2))) + ((a^2*b
+ b^3)*d*tan(d*x + c) + (a^3 + a*b^2)*d)*sqrt(-(a^5 - 10*a^3*b^2 + 5*a*b^
4 + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*sqrt(-(25*a^8*b^2 - 100*a^6*b^
4 + 110*a^4*b^6 - 20*a^2*b^8 + b^10)/((a^12 + 6*a^10*b^2 + 15*a^8*b^4 + 20
*a^6*b^6 + 15*a^4*b^8 + 6*a^2*b^10 + b^12)*d^4)))/((a^6 + 3*a^4*b^2 + 3*a^
2*b^4 + b^6)*d^2))*log((5*a^6*b - 5*a^4*b^3 - 9*a^2*b^5 + b^7)*sqrt(b*tan(
d*x + c) + a) - ((a^9 - 6*a^5*b^4 - 8*a^3*b^6 - 3*a*b^8)*d^3*sqrt(-(25*a^8
*b^2 - 100*a^6*b^4 + 110*a^4*b^6 - 20*a^2*b^8 + b^10)/((a^12 + 6*a^10*b...
```

3.371.6 Sympy [F]

$$\int \frac{-a + b \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx =$$

$$- \int \frac{a}{a \sqrt{a + b \tan(c + dx)} + b \sqrt{a + b \tan(c + dx)} \tan(c + dx)} dx$$

$$- \int \left(- \frac{b \tan(c + dx)}{a \sqrt{a + b \tan(c + dx)} + b \sqrt{a + b \tan(c + dx)} \tan(c + dx)} \right) dx$$

3.371. $\int \frac{-a + b \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx$

input `integrate((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)`

output `-Integral(a/(a*sqrt(a + b*tan(c + d*x)) + b*sqrt(a + b*tan(c + d*x))*tan(c + d*x)), x) - Integral(-b*tan(c + d*x)/(a*sqrt(a + b*tan(c + d*x)) + b*sqrt(a + b*tan(c + d*x))*tan(c + d*x)), x)`

3.371.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{-a + b \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is`

3.371.8 Giac [F(-1)]

Timed out.

$$\int \frac{-a + b \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `Timed out`

3.371.9 Mupad [B] (verification not implemented)

Time = 13.95 (sec) , antiderivative size = 5475, normalized size of antiderivative = 41.48

$$\int \frac{-a + b \tan(c + dx)}{(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

```
input int(-(a - b*tan(c + d*x))/(a + b*tan(c + d*x))^(3/2),x)
```

```
output log(- (((((24*a*b^4*d^2 - 8*a^3*b^2*d^2)^2/4 - b^4*(16*a^6*d^4 + 16*b^6*d^4 + 48*a^2*b^4*d^4 + 48*a^4*b^2*d^4))^(1/2) - 12*a*b^4*d^2 + 4*a^3*b^2*d^2)/(16*(a^6*d^4 + b^6*d^4 + 3*a^2*b^4*d^4 + 3*a^4*b^2*d^4)))^(1/2)*(32*b^13*d^4 + 96*a^2*b^11*d^4 + 64*a^4*b^9*d^4 - 64*a^6*b^7*d^4 - 96*a^8*b^5*d^4 - 32*a^10*b^3*d^4 + (a + b*tan(c + d*x))^(1/2)*(((24*a*b^4*d^2 - 8*a^3*b^2*d^2)^2/4 - b^4*(16*a^6*d^4 + 16*b^6*d^4 + 48*a^2*b^4*d^4 + 48*a^4*b^2*d^4))^(1/2) - 12*a*b^4*d^2 + 4*a^3*b^2*d^2)/(16*(a^6*d^4 + b^6*d^4 + 3*a^2*b^4*d^4 + 3*a^4*b^2*d^4)))^(1/2)*(64*a*b^12*d^5 + 320*a^3*b^10*d^5 + 640*a^5*b^8*d^5 + 640*a^7*b^6*d^5 + 320*a^9*b^4*d^5 + 64*a^11*b^2*d^5)) + (a + b*tan(c + d*x))^(1/2)*(16*b^12*d^3 + 32*a^2*b^10*d^3 - 32*a^6*b^6*d^3 - 16*a^8*b^4*d^3))*(((24*a*b^4*d^2 - 8*a^3*b^2*d^2)^2/4 - b^4*(16*a^6*d^4 + 16*b^6*d^4 + 48*a^2*b^4*d^4 + 48*a^4*b^2*d^4))^(1/2) - 12*a*b^4*d^2 + 4*a^3*b^2*d^2)/(16*(a^6*d^4 + b^6*d^4 + 3*a^2*b^4*d^4 + 3*a^4*b^2*d^4)))^(1/2) - 8*a*b^11*d^2 - 24*a^3*b^9*d^2 - 24*a^5*b^7*d^2 - 8*a^7*b^5*d^2)*(((24*a*b^4*d^2 - 8*a^3*b^2*d^2)^2/4 - b^4*(16*a^6*d^4 + 16*b^6*d^4 + 48*a^2*b^4*d^4 + 48*a^4*b^2*d^4))^(1/2) - 12*a*b^4*d^2 + 4*a^3*b^2*d^2)/(16*(a^6*d^4 + b^6*d^4 + 3*a^2*b^4*d^4 + 3*a^4*b^2*d^4)))^(1/2) + log(- ((-(((24*a*b^4*d^2 - 8*a^3*b^2*d^2)^2/4 - b^4*(16*a^6*d^4 + 16*b^6*d^4 + 48*a^2*b^4*d^4 + 48*a^4*b^2*d^4))^(1/2) + 12*a*b^4*d^2 - 4*a^3*b^2*d^2)/(16*(a^6*d^4 + b^6*d^4 + 3*a^2*b^4*d^4 + 3*a^4*b^2*d^4)))^(1/2)*(32*b^13*d^4 + 96*a^2*b^11...
```

3.372
$$\int \frac{-a+b \tan(c+dx)}{(a+b \tan(c+dx))^{5/2}} dx$$

3.372.1 Optimal result 3536
 3.372.2 Mathematica [C] (verified) 3536
 3.372.3 Rubi [A] (warning: unable to verify) 3537
 3.372.4 Maple [B] (verified) 3540
 3.372.5 Fricas [B] (verification not implemented) 3541
 3.372.6 Sympy [F] 3542
 3.372.7 Maxima [F(-2)] 3543
 3.372.8 Giac [F(-1)] 3543
 3.372.9 Mupad [B] (verification not implemented) 3543

3.372.1 Optimal result

Integrand size = 27, antiderivative size = 174

$$\int \frac{-a+b \tan(c+dx)}{(a+b \tan(c+dx))^{5/2}} dx = \frac{(ia-b)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{(a-ib)^{5/2}d} - \frac{(ia+b)\operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{(a+ib)^{5/2}d} + \frac{4ab}{3(a^2+b^2)d(a+b \tan(c+dx))^{3/2}} + \frac{2b(3a^2-b^2)}{(a^2+b^2)^2 d \sqrt{a+b \tan(c+dx)}}$$

```
output (I*a-b)*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/(a-I*b)^(5/2)/d-(I*a+b)*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/(a+I*b)^(5/2)/d+2*b*(3*a^2-b^2)/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^(1/2)+4/3*a*b/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)
```

3.372.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.31 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.90

$$\int \frac{-a+b \tan(c+dx)}{(a+b \tan(c+dx))^{5/2}} dx = \frac{i \cos(c+dx) \left((a+ib)^2 \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{a+b \tan(c+dx)}{a-ib}\right) - (a-ib)^2 \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{a+b \tan(c+dx)}{a+ib}\right) \right)}{3(a-ib)(a+ib)d(a \cos(c+dx) - b \sin(c+dx))(a+b \tan(c+dx))^{3/2}}$$

3.372.
$$\int \frac{-a+b \tan(c+dx)}{(a+b \tan(c+dx))^{5/2}} dx$$

input `Integrate[(-a + b*Tan[c + d*x])/(a + b*Tan[c + d*x])^(5/2),x]`

output `((-1/3*I)*Cos[c + d*x]*((a + I*b)^2*Hypergeometric2F1[-3/2, 1, -1/2, (a + b*Tan[c + d*x])/(a - I*b)] - (a - I*b)^2*Hypergeometric2F1[-3/2, 1, -1/2, (a + b*Tan[c + d*x])/(a + I*b)])*(a - b*Tan[c + d*x]))/((a - I*b)*(a + I*b)*d*(a*Cos[c + d*x] - b*Sin[c + d*x])*(a + b*Tan[c + d*x])^(3/2))`

3.372.3 Rubi [A] (warning: unable to verify)

Time = 0.86 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.05, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 4012, 25, 3042, 4012, 3042, 4022, 3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{b \tan(c + dx) - a}{(a + b \tan(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{b \tan(c + dx) - a}{(a + b \tan(c + dx))^{5/2}} dx \\
 & \quad \downarrow \text{4012} \\
 & \frac{\int -\frac{a^2 - 2b \tan(c + dx)a - b^2}{(a + b \tan(c + dx))^{3/2}} dx}{a^2 + b^2} + \frac{4ab}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{4ab}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{\int \frac{a^2 - 2b \tan(c + dx)a - b^2}{(a + b \tan(c + dx))^{3/2}} dx}{a^2 + b^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{4ab}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{\int \frac{a^2 - 2b \tan(c + dx)a - b^2}{(a + b \tan(c + dx))^{3/2}} dx}{a^2 + b^2} \\
 & \quad \downarrow \text{4012} \\
 & \frac{4ab}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{\int \frac{a(a^2 - 3b^2) - b(3a^2 - b^2) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx}{a^2 + b^2} - \frac{2b(3a^2 - b^2)}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}}
 \end{aligned}$$

3.372. $\int \frac{-a + b \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \frac{4ab}{3d(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \frac{\int \frac{a(a^2-3b^2)-b(3a^2-b^2)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx}{a^2+b^2} - \frac{2b(3a^2-b^2)}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \\
 & \downarrow 4022 \\
 & \frac{4ab}{3d(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \\
 & - \frac{2b(3a^2-b^2)}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{\frac{1}{2}(a-ib)^3 \int \frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}(a+ib)^3 \int \frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}} dx}{a^2+b^2} \\
 & \downarrow 3042 \\
 & \frac{4ab}{3d(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \\
 & - \frac{2b(3a^2-b^2)}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{\frac{1}{2}(a-ib)^3 \int \frac{1-i\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}(a+ib)^3 \int \frac{i\tan(c+dx)+1}{\sqrt{a+b\tan(c+dx)}} dx}{a^2+b^2} \\
 & \downarrow 4020 \\
 & \frac{4ab}{3d(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \\
 & - \frac{2b(3a^2-b^2)}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{i(a+ib)^3 \int -\frac{1}{(1-i\tan(c+dx))\sqrt{a+b\tan(c+dx)}} d(i\tan(c+dx)) - i(a-ib)^3 \int -\frac{1}{(i\tan(c+dx)+1)\sqrt{a+b\tan(c+dx)}} d(-i\tan(c+dx))}{a^2+b^2} \\
 & \downarrow 25 \\
 & \frac{4ab}{3d(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \\
 & - \frac{2b(3a^2-b^2)}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{i(a-ib)^3 \int \frac{1}{(i\tan(c+dx)+1)\sqrt{a+b\tan(c+dx)}} d(-i\tan(c+dx)) - i(a+ib)^3 \int \frac{1}{(1-i\tan(c+dx))\sqrt{a+b\tan(c+dx)}} d(i\tan(c+dx))}{a^2+b^2} \\
 & \downarrow 73 \\
 & \frac{4ab}{3d(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \\
 & - \frac{2b(3a^2-b^2)}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{(a-ib)^3 \int \frac{1}{-i\tan^2(c+dx) - \frac{ia}{b} + 1} d\sqrt{a+b\tan(c+dx)} + (a+ib)^3 \int \frac{1}{i\tan^2(c+dx) + \frac{ia}{b} + 1} d\sqrt{a+b\tan(c+dx)}}{a^2+b^2} \\
 & \downarrow 221
 \end{aligned}$$

3.372. $\int \frac{-a+b\tan(c+dx)}{(a+b\tan(c+dx))^{5/2}} dx$

$$\frac{4ab}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2b(3a^2 - b^2)}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{(a - ib)^3 \arctan\left(\frac{\tan(c + dx)}{\sqrt{a + ib}}\right)}{d\sqrt{a + ib}} + \frac{(a + ib)^3 \arctan\left(\frac{\tan(c + dx)}{\sqrt{a - ib}}\right)}{d\sqrt{a - ib}}$$

$$a^2 + b^2$$

input `Int[(-a + b*Tan[c + d*x])/(a + b*Tan[c + d*x])^(5/2),x]`

output `(4*a*b)/(3*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) - (((a + I*b)^3*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d) + ((a - I*b)^3*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d))/(a^2 + b^2) - (2*b*(3*a^2 - b^2))/((a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])/(a^2 + b^2)`

3.372.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*
(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

3.372.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3054 vs. $2(150) = 300$.

Time = 0.10 (sec) , antiderivative size = 3055, normalized size of antiderivative = 17.56

method	result	size
derivativedivides	Expression too large to display	3055
default	Expression too large to display	3055
parts	Expression too large to display	4474

```
input int((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x,method=_RETURNVERBOSE)
```

output

```

-1/2*d*b/(a^2+b^2)^3*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3+2/d*b^3/(a^2+b^2)^3/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^2+1/d/b/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^7-5/d*b^3/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^3+1/4/d/b/(a^2+b^2)^(7/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^6-5/4/d*b^3/(a^2+b^2)^(7/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2+7/d*b^5/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a-1/d/b/(a^2+b^2)^(7/2)/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))*a^7+4/3*a*b/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)-2/d*b^3/(a^2+b^2)^2/(a+b*tan(d*x+c))^(1/2)+1/4/d*b^5/(a^2+b^2)^(7/2)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+1/d*b^5/(a^2+b^2)^3/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arcta...

```

3.372.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3695 vs. 2(144) = 288.

Time = 0.34 (sec) , antiderivative size = 3695, normalized size of antiderivative = 21.24

$$\int \frac{-a + b \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="fricas")`

```
output 1/6*(3*((a^4*b^2 + 2*a^2*b^4 + b^6)*d*tan(d*x + c)^2 + 2*(a^5*b + 2*a^3*b^3 + a*b^5)*d*tan(d*x + c) + (a^6 + 2*a^4*b^2 + a^2*b^4)*d)*sqrt(-(a^7 - 21*a^5*b^2 + 35*a^3*b^4 - 7*a*b^6 + (a^10 + 5*a^8*b^2 + 10*a^6*b^4 + 10*a^4*b^6 + 5*a^2*b^8 + b^10)*d^2*sqrt(-(49*a^12*b^2 - 490*a^10*b^4 + 1519*a^8*b^6 - 1484*a^6*b^8 + 511*a^4*b^10 - 42*a^2*b^12 + b^14)/((a^20 + 10*a^18*b^2 + 45*a^16*b^4 + 120*a^14*b^6 + 210*a^12*b^8 + 252*a^10*b^10 + 210*a^8*b^12 + 120*a^6*b^14 + 45*a^4*b^16 + 10*a^2*b^18 + b^20)*d^4)))/((a^10 + 5*a^8*b^2 + 10*a^6*b^4 + 10*a^4*b^6 + 5*a^2*b^8 + b^10)*d^2))*log(-(7*a^8*b - 28*a^6*b^3 - 14*a^4*b^5 + 20*a^2*b^7 - b^9)*sqrt(b*tan(d*x + c) + a) + ((a^14 - a^12*b^2 - 19*a^10*b^4 - 45*a^8*b^6 - 45*a^6*b^8 - 19*a^4*b^10 - a^2*b^12 + b^14)*d^3*sqrt(-(49*a^12*b^2 - 490*a^10*b^4 + 1519*a^8*b^6 - 1484*a^6*b^8 + 511*a^4*b^10 - 42*a^2*b^12 + b^14)/((a^20 + 10*a^18*b^2 + 45*a^16*b^4 + 120*a^14*b^6 + 210*a^12*b^8 + 252*a^10*b^10 + 210*a^8*b^12 + 120*a^6*b^14 + 45*a^4*b^16 + 10*a^2*b^18 + b^20)*d^4)) + 4*(7*a^9*b^2 - 42*a^7*b^4 + 56*a^5*b^6 - 22*a^3*b^8 + a*b^10)*d)*sqrt(-(a^7 - 21*a^5*b^2 + 35*a^3*b^4 - 7*a*b^6 + (a^10 + 5*a^8*b^2 + 10*a^6*b^4 + 10*a^4*b^6 + 5*a^2*b^8 + b^10)*d^2*sqrt(-(49*a^12*b^2 - 490*a^10*b^4 + 1519*a^8*b^6 - 1484*a^6*b^8 + 511*a^4*b^10 - 42*a^2*b^12 + b^14)/((a^20 + 10*a^18*b^2 + 45*a^16*b^4 + 120*a^14*b^6 + 210*a^12*b^8 + 252*a^10*b^10 + 210*a^8*b^12 + 120*a^6*b^14 + 45*a^4*b^16 + 10*a^2*b^18 + b^20)*d^4)))/((a^10 + 5*a^8*b^2 + 10*a^...
```

3.372.6 Sympy [F]

$$\int \frac{-a + b \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx =$$

$$- \int \frac{a}{a^2 \sqrt{a + b \tan(c + dx)} + 2ab \sqrt{a + b \tan(c + dx)} \tan(c + dx) + b^2 \sqrt{a + b \tan(c + dx)} \tan^2(c + dx)} dx$$

$$- \int \left(- \frac{b \tan(c + dx)}{a^2 \sqrt{a + b \tan(c + dx)} + 2ab \sqrt{a + b \tan(c + dx)} \tan(c + dx) + b^2 \sqrt{a + b \tan(c + dx)} \tan^2(c + dx)} \right) dx$$

```
input integrate((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2), x)
```

```
output -Integral(a/(a**2*sqrt(a + b*tan(c + d*x)) + 2*a*b*sqrt(a + b*tan(c + d*x))*tan(c + d*x) + b**2*sqrt(a + b*tan(c + d*x))*tan(c + d*x)**2), x) - Integral(-b*tan(c + d*x)/(a**2*sqrt(a + b*tan(c + d*x)) + 2*a*b*sqrt(a + b*tan(c + d*x))*tan(c + d*x) + b**2*sqrt(a + b*tan(c + d*x))*tan(c + d*x)**2), x)
```

3.372. $\int \frac{-a+b \tan(c+dx)}{(a+b \tan(c+dx))^{5/2}} dx$

3.372.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{-a + b \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \text{Exception raised: ValueError}$$

input `integrate((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is`

3.372.8 Giac [F(-1)]

Timed out.

$$\int \frac{-a + b \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((-a+b*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Timed out`

3.372.9 Mupad [B] (verification not implemented)

Time = 24.86 (sec) , antiderivative size = 8437, normalized size of antiderivative = 48.49

$$\int \frac{-a + b \tan(c + dx)}{(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `int(-(a - b*tan(c + d*x))/(a + b*tan(c + d*x))^(5/2),x)`

output

$$\begin{aligned} & (\log(\left(\left(\left(-4a^7d^2 + (320a^6b^8d^4 - 16a^4b^{10}d^4 - 1760a^8b^6d^4 + 1600a^{10}b^4d^4 - 400a^{12}b^2d^4)\right)^{1/2} + 20a^3b^4d^2 - 40a^5b^2d^2\right)/(a^{10}d^4 + b^{10}d^4 + 5a^2b^8d^4 + 10a^4b^6d^4 + 10a^6b^4d^4 + 5a^8b^2d^4)\right)^{1/2} * (896a^7b^{15}d^4 - 32ab^{21}d^4 - 160a^3b^{19}d^4 - 128a^5b^{17}d^4 - ((a + b\tan(c + dx))^{1/2} * (-4a^7d^2 + (320a^6b^8d^4 - 16a^4b^{10}d^4 - 1760a^8b^6d^4 + 1600a^{10}b^4d^4 - 400a^{12}b^2d^4)^{1/2} + 20a^3b^4d^2 - 40a^5b^2d^2)/(a^{10}d^4 + b^{10}d^4 + 5a^2b^8d^4 + 10a^4b^6d^4 + 10a^6b^4d^4 + 5a^8b^2d^4))^{1/2} * (64ab^{22}d^5 + 640a^3b^{20}d^5 + 2880a^5b^{18}d^5 + 7680a^7b^{16}d^5 + 13440a^9b^{14}d^5 + 16128a^{11}b^{12}d^5 + 13440a^{13}b^{10}d^5 + 7680a^{15}b^8d^5 + 2880a^{17}b^6d^5 + 640a^{19}b^4d^5 + 64a^{21}b^2d^5)) / 4 + 3136a^9b^{13}d^4 + 4928a^{11}b^{11}d^4 + 4480a^{13}b^9d^4 + 2432a^{15}b^7d^4 + 736a^{17}b^5d^4 + 96a^{19}b^3d^4)) / 4 + (a + b\tan(c + dx))^{1/2} * (320a^6b^{14}d^3 - 16a^2b^{18}d^3 + 1024a^8b^{12}d^3 + 1440a^{10}b^{10}d^3 + 1024a^{12}b^8d^3 + 320a^{14}b^6d^3 - 16a^{18}b^2d^3)) * (-4a^7d^2 + (320a^6b^8d^4 - 16a^4b^{10}d^4 - 1760a^8b^6d^4 + 1600a^{10}b^4d^4 - 400a^{12}b^2d^4)^{1/2} + 20a^3b^4d^2 - 40a^5b^2d^2) / (a^{10}d^4 + b^{10}d^4 + 5a^2b^8d^4 + 10a^4b^6d^4 + 10a^6b^4d^4 + 5a^8b^2d^4)^{1/2} / 4 - 16a^4b^{15}d^2 - 96a^6b^{13}d^2 - 240a^8b^{11}d^2 - 320a^{10}b^9d^2 - 240a^{12}b^7d^2 - 96a^{14}b^5d^2 - 16a^{16}b^3d^2 - 320a^{18}b^3d^2 - 16a^{20}b^1d^2 - 16a^{22}b^{-1}d^2) \end{aligned}$$

3.373 $\int \frac{1+i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$

3.373.1 Optimal result 3545
 3.373.2 Mathematica [A] (verified) 3545
 3.373.3 Rubi [A] (warning: unable to verify) 3546
 3.373.4 Maple [B] (verified) 3547
 3.373.5 Fricas [B] (verification not implemented) 3549
 3.373.6 Sympy [F] 3549
 3.373.7 Maxima [B] (verification not implemented) 3550
 3.373.8 Giac [B] (verification not implemented) 3550
 3.373.9 Mupad [B] (verification not implemented) 3551

3.373.1 Optimal result

Integrand size = 27, antiderivative size = 45

$$\int \frac{1+i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx = -\frac{2i \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib}d}$$

output `-2*I*arctanh((a+b*tan(d*x+c))^(1/2)/(a-I*b)^(1/2))/d/(a-I*b)^(1/2)`

3.373.2 Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00

$$\int \frac{1+i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx = -\frac{2i \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a-ib}}\right)}{\sqrt{a-ib}d}$$

input `Integrate[(1 + I*Tan[c + d*x])/Sqrt[a + b*Tan[c + d*x]],x]`

output `((-2*I)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d)`

3.373.3 Rubi [A] (warning: unable to verify)

Time = 0.24 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.78, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$, Rules used = {3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1 + i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1 + i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4020} \\
 & \frac{i \int -\frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{d} \\
 & \quad \downarrow \text{25} \\
 & -\frac{i \int \frac{1}{(1 - i \tan(c + dx))\sqrt{a + b \tan(c + dx)}} d(i \tan(c + dx))}{d} \\
 & \quad \downarrow \text{73} \\
 & \frac{2 \int \frac{1}{\frac{i \tan^2(c + dx)}{b} + \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)}}{bd} \\
 & \quad \downarrow \text{221} \\
 & \frac{2 \arctan\left(\frac{\tan(c + dx)}{\sqrt{a - ib}}\right)}{d\sqrt{a - ib}}
 \end{aligned}$$

input `Int[(1 + I*Tan[c + d*x])/Sqrt[a + b*Tan[c + d*x]],x]`

output `(2*ArcTan[Tan[c + d*x]/Sqrt[a - I*b]])/(Sqrt[a - I*b]*d)`

3.373.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

3.373.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 728 vs. $2(36) = 72$.

Time = 0.10 (sec) , antiderivative size = 729, normalized size of antiderivative = 16.20

method	result
derivativedivides	$\frac{(-i\sqrt{a^2+b^2}-ia+b) \ln\left(\frac{b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}}{2}\right) + \frac{2\left(-i\sqrt{2\sqrt{a^2+b^2}+2a} a + \sqrt{2\sqrt{a^2+b^2}+2a} b\right)}{\sqrt{2\sqrt{a^2+b^2}+2a} \sqrt{a^2+b^2}}}{\sqrt{2\sqrt{a^2+b^2}+2a} \sqrt{a^2+b^2}}$
default	$\frac{(-i\sqrt{a^2+b^2}-ia+b) \ln\left(\frac{b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}}{2}\right) + \frac{2\left(-i\sqrt{2\sqrt{a^2+b^2}+2a} a + \sqrt{2\sqrt{a^2+b^2}+2a} b\right)}{\sqrt{2\sqrt{a^2+b^2}+2a} \sqrt{a^2+b^2}}}{\sqrt{2\sqrt{a^2+b^2}+2a} \sqrt{a^2+b^2}}$
parts	Expression too large to display

input `int((1+I*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2), x, method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/d*(1/(2*(a^2+b^2)^(1/2)+2*a)^(1/2)/(a^2+b^2)^(1/2)*(1/2*(-I*(a^2+b^2)^(1/2) \\ & -I*a+b)*\ln(b*\tan(d*x+c)+a+(a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a) \\ &)^(1/2)+(a^2+b^2)^(1/2))+2*(-I*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a+(2*(a^2+b^2) \\ &)^(1/2)+2*a)^(1/2)*b-1/2*(-I*(a^2+b^2)^(1/2)-I*a+b)*(2*(a^2+b^2)^(1/2)+2*a) \\ &)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*\arctan((2*(a+b*\tan(d*x+c))^(1/2)+(2 \\ & *(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)))+1/(2*(a^2+b^2) \\ &)^(1/2)+2*a)^(1/2)/(a^2+b^2)^(1/2)/((a^2+b^2)^(1/2)*a+a^2+b^2)*(-1/2*(-2*I \\ & *(a^2+b^2)^(1/2)*a^2-I*(a^2+b^2)^(1/2)*b^2-2*I*a^3-2*I*a*b^2+(a^2+b^2)^(1/2) \\ &)*(a*b+a^2*b+b^3)*\ln((a+b*\tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)- \\ & b*\tan(d*x+c)-a-(a^2+b^2)^(1/2))+2*(I*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2) \\ &)^(1/2)*a^2+I*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3+I*(2*(a^2+b^2)^(1/2)+2*a) \\ &)^(1/2)*a*b^2-(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*a*b-(2*(a^2+b^2) \\ &)^(1/2)+2*a)^(1/2)*a^2*b-(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*b^3+1/2*(-2*I*(a^2+ \\ & b^2)^(1/2)*a^2-I*(a^2+b^2)^(1/2)*b^2-2*I*a^3-2*I*a*b^2+(a^2+b^2)^(1/2)*a*b \\ & +a^2*b+b^3)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*a \\ & \arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*\tan(d*x+c))^(1/2))/(2*(a^2+b^2) \\ &)^(1/2)-2*a)^(1/2)))) \end{aligned}$$

3.373.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 249 vs. $2(33) = 66$.

Time = 0.24 (sec) , antiderivative size = 249, normalized size of antiderivative = 5.53

$$\int \frac{1 + i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx$$

$$= \frac{1}{4} \sqrt{-\frac{4i}{(ia + b)d^2}} \log \left(\left((ia + b)de^{(2i dx + 2i c)} + (ia + b)d \right) \sqrt{\frac{(a - ib)e^{(2i dx + 2i c)} + a + ib}{e^{(2i dx + 2i c)} + 1}} \sqrt{-\frac{4i}{(ia + b)d^2}} \right)$$

$$- \frac{1}{4} \sqrt{-\frac{4i}{(ia + b)d^2}} \log \left(\left((-ia - b)de^{(2i dx + 2i c)} + (-ia - b)d \right) \sqrt{\frac{(a - ib)e^{(2i dx + 2i c)} + a + ib}{e^{(2i dx + 2i c)} + 1}} \sqrt{-\frac{4i}{(ia + b)d^2}} \right)$$

input `integrate((1+I*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="fricas")`

output `1/4*sqrt(-4*I/((I*a + b)*d^2))*log(((I*a + b)*d*e^(2*I*d*x + 2*I*c) + (I*a + b)*d)*sqrt(((a - I*b)*e^(2*I*d*x + 2*I*c) + a + I*b)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-4*I/((I*a + b)*d^2)) + 2*(a - I*b)*e^(2*I*d*x + 2*I*c) + 2*a)*e^(-2*I*d*x - 2*I*c)) - 1/4*sqrt(-4*I/((I*a + b)*d^2))*log(((I*a + b)*d*e^(2*I*d*x + 2*I*c) + (-I*a - b)*d)*sqrt(((a - I*b)*e^(2*I*d*x + 2*I*c) + a + I*b)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(-4*I/((I*a + b)*d^2)) + 2*(a - I*b)*e^(2*I*d*x + 2*I*c) + 2*a)*e^(-2*I*d*x - 2*I*c))`

3.373.6 Sympy [F]

$$\int \frac{1 + i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx$$

$$= i \left(\int \left(-\frac{i}{\sqrt{a + b \tan(c + dx)}} \right) dx + \int \frac{\tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx \right)$$

input `integrate((1+I*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)`

output `I*(Integral(-I/sqrt(a + b*tan(c + d*x)), x) + Integral(tan(c + d*x)/sqrt(a + b*tan(c + d*x)), x))`

3.373. $\int \frac{1+i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$

3.373.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 6480 vs. $2(33) = 66$.

Time = 0.64 (sec) , antiderivative size = 6480, normalized size of antiderivative = 144.00

$$\int \frac{1 + i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

```
input integrate((1+I*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="maxima")
```

```
output -1/4*(sqrt(2*a^2 + 2*b^2)*sqrt(a + sqrt(a^2 + b^2))*(2*arctan2((b^2*cos(2*
d*x + 2*c) - a*b*sin(2*d*x + 2*c) + ((a^4 + 2*a^2*b^2 + b^4)*cos(2*d*x + 2
*c)^4 + (a^4 + 2*a^2*b^2 + b^4)*sin(2*d*x + 2*c)^4 + a^4 + 2*a^2*b^2 + b^4
+ 4*(a^4 + a^2*b^2)*cos(2*d*x + 2*c)^3 + 4*(a^3*b + a*b^3)*sin(2*d*x + 2*
c)^3 + 2*(3*a^4 + 2*a^2*b^2 - b^4)*cos(2*d*x + 2*c)^2 + 2*(a^4 + 2*a^2*b^2
+ b^4 + (a^4 + 2*a^2*b^2 + b^4)*cos(2*d*x + 2*c)^2 + 2*(a^4 + a^2*b^2)*co
s(2*d*x + 2*c))*sin(2*d*x + 2*c)^2 + 4*(a^4 + a^2*b^2)*cos(2*d*x + 2*c) +
4*(a^3*b + a*b^3 + (a^3*b + a*b^3)*cos(2*d*x + 2*c)^2 + 2*(a^3*b + a*b^3)*
cos(2*d*x + 2*c))*sin(2*d*x + 2*c))^(1/4)*b*sin(1/2*arctan2(-2*(a*b*cos(2*
d*x + 2*c)^2 - a*b*sin(2*d*x + 2*c)^2 + a*b*cos(2*d*x + 2*c) - (a^2 + (a^2
- b^2)*cos(2*d*x + 2*c))*sin(2*d*x + 2*c))/b^2, (2*a^2*cos(2*d*x + 2*c) +
(a^2 - b^2)*cos(2*d*x + 2*c)^2 - (a^2 - b^2)*sin(2*d*x + 2*c)^2 + a^2 + b
^2 + 2*(2*a*b*cos(2*d*x + 2*c) + a*b)*sin(2*d*x + 2*c))/b^2)), -(a*co
s(2*d*x + 2*c) + b*sin(2*d*x + 2*c) - ((a^4 + 2*a^2*b^2 + b^4)*cos(2*d*x +
2*c)^4 + (a^4 + 2*a^2*b^2 + b^4)*sin(2*d*x + 2*c)^4 + a^4 + 2*a^2*b^2 + b
^4 + 4*(a^4 + a^2*b^2)*cos(2*d*x + 2*c)^3 + 4*(a^3*b + a*b^3)*sin(2*d*x +
2*c)^3 + 2*(3*a^4 + 2*a^2*b^2 - b^4)*cos(2*d*x + 2*c)^2 + 2*(a^4 + 2*a^2*b
^2 + b^4 + (a^4 + 2*a^2*b^2 + b^4)*cos(2*d*x + 2*c)^2 + 2*(a^4 + a^2*b^2)*
cos(2*d*x + 2*c))*sin(2*d*x + 2*c)^2 + 4*(a^4 + a^2*b^2)*cos(2*d*x + 2*c)
+ 4*(a^3*b + a*b^3 + (a^3*b + a*b^3)*cos(2*d*x + 2*c)^2 + 2*(a^3*b + a*...
```

3.373.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 155 vs. $2(33) = 66$.

Time = 0.41 (sec) , antiderivative size = 155, normalized size of antiderivative = 3.44

$$\int \frac{1 + i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx$$

$$= \frac{4i \arctan\left(\frac{2(\sqrt{b \tan(dx+c)+a} - \sqrt{a^2+b^2})\sqrt{b \tan(dx+c)+a}}{a\sqrt{-2a+2\sqrt{a^2+b^2}} - i\sqrt{-2a+2\sqrt{a^2+b^2}}b - \sqrt{a^2+b^2}\sqrt{-2a+2\sqrt{a^2+b^2}}}\right)}{\sqrt{-2a+2\sqrt{a^2+b^2}}d\left(-\frac{ib}{a-\sqrt{a^2+b^2}} + 1\right)}$$

input `integrate((1+I*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `4*I*arctan(2*(sqrt(b*tan(d*x + c) + a)*a - sqrt(a^2 + b^2)*sqrt(b*tan(d*x + c) + a))/(a*sqrt(-2*a + 2*sqrt(a^2 + b^2)) - I*sqrt(-2*a + 2*sqrt(a^2 + b^2))*b - sqrt(a^2 + b^2)*sqrt(-2*a + 2*sqrt(a^2 + b^2)))/(sqrt(-2*a + 2*sqrt(a^2 + b^2))*d*(-I*b/(a - sqrt(a^2 + b^2)) + 1))`

3.373.9 Mupad [B] (verification not implemented)

Time = 10.39 (sec) , antiderivative size = 1410, normalized size of antiderivative = 31.33

$$\int \frac{1 + i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `int((tan(c + d*x)*i + 1)/(a + b*tan(c + d*x))^(1/2),x)`

output

$$\begin{aligned}
& 2*\operatorname{atanh}\left(\frac{32*b^2*((b*1i)/(4*a^2*d^2 + 4*b^2*d^2) - a/(4*a^2*d^2 + 4*b^2*d^2))^{1/2}}{(a + b*\tan(c + d*x))^{1/2}}\right) / \left(\frac{(a^2*b^2*d^2*64i)/(4*a^2*d^3 + 4*b^2*d^3) - (b^2*16i)/d + (64*a*b^3*d^2)/(4*a^2*d^3 + 4*b^2*d^3) - (128*a^2*b^2*((b*1i)/(4*a^2*d^2 + 4*b^2*d^2) - a/(4*a^2*d^2 + 4*b^2*d^2))^{1/2})}{(a + b*\tan(c + d*x))^{1/2}}\right) / \left(\frac{(a^2*b^4*d^2*256i)/(4*a^2*d^3 + 4*b^2*d^3) - (a^2*b^2*64i)/d - (b^4*64i)/d + (256*a^3*b^3*d^2)/(4*a^2*d^3 + 4*b^2*d^3) + (a^4*b^2*d^2*256i)/(4*a^2*d^3 + 4*b^2*d^3) + (256*a*b^5*d^2)/(4*a^2*d^3 + 4*b^2*d^3)}{(a + b*\tan(c + d*x))^{1/2}}\right) \\
& + (a*b^3*((b*1i)/(4*a^2*d^2 + 4*b^2*d^2) - a/(4*a^2*d^2 + 4*b^2*d^2))^{1/2}) * (a + b*\tan(c + d*x))^{1/2} * 128i / \left(\frac{(a^2*b^4*d^2*256i)/(4*a^2*d^3 + 4*b^2*d^3) - (a^2*b^2*64i)/d - (b^4*64i)/d + (256*a^3*b^3*d^2)/(4*a^2*d^3 + 4*b^2*d^3) + (a^4*b^2*d^2*256i)/(4*a^2*d^3 + 4*b^2*d^3) + (256*a*b^5*d^2)/(4*a^2*d^3 + 4*b^2*d^3)}{(a + b*\tan(c + d*x))^{1/2}}\right) \\
& - (a^2*b^2*64i)/d - (b^4*64i)/d + (256*a^3*b^3*d^2)/(4*a^2*d^3 + 4*b^2*d^3) + (a^4*b^2*d^2*256i)/(4*a^2*d^3 + 4*b^2*d^3) + (256*a*b^5*d^2)/(4*a^2*d^3 + 4*b^2*d^3) \\
&) * (- (a - b*1i)/(4*a^2*d^2 + 4*b^2*d^2))^{1/2} + (\log(d*(-1/(d^2*(a - b*1i))))^{1/2} * (a + b*\tan(c + d*x))^{1/2} * 1i + 1) * (-1/(a*d^2 - b*d^2*1i))^{1/2}) / 2 - \log(d*(-1/(d^2*(a - b*1i))))^{1/2} * (a + b*\tan(c + d*x))^{1/2} + 1i) * (-1/(4*(a*d^2 - b*d^2*1i)))^{1/2} + (\log(16*b^2*(a + b*\tan(c + d*x))^{1/2} + 16*b^3*d*(-1/(d^2*(a - b*1i))))^{1/2} - (16*a*b^2*(a + b*\tan(c + d*x))^{1/2}) / (a - b*1i)) * (-1/(a*d^2 - b*d^2*1i))^{1/2}) / 2 - \log(16*b^3*d*(-1/(d^2*(a - b*1i))))^{1/2} - 16*b^2*(a + b*\tan(c + d*x))^{1/2} + (16*a*b^2*(a + b*\tan(c + d*x))^{1/2}) / (a - b*1i)) * (-1/(4*(a*d^2 - b*d^2*1i)))^{1/2} + 2*\operatorname{atanh}\left(\frac{32*b^2*((b*1i)/(4*a^2*d^2 + 4*b^2*d^2) - a/(4*a^2*d^2 + 4*b^2*d^2))^{1/2}}{(a + b*\tan(c + d*x))^{1/2}}\right) \dots
\end{aligned}$$

3.374 $\int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$

3.374.1 Optimal result 3553
 3.374.2 Mathematica [A] (verified) 3553
 3.374.3 Rubi [A] (warning: unable to verify) 3554
 3.374.4 Maple [B] (verified) 3555
 3.374.5 Fracas [B] (verification not implemented) 3557
 3.374.6 Sympy [F] 3557
 3.374.7 Maxima [F(-2)] 3558
 3.374.8 Giac [B] (verification not implemented) 3558
 3.374.9 Mupad [B] (verification not implemented) 3559

3.374.1 Optimal result

Integrand size = 27, antiderivative size = 45

$$\int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx = \frac{2i \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ibd}}$$

output `2*I*arctanh((a+b*tan(d*x+c))^(1/2)/(a+I*b)^(1/2))/d/(a+I*b)^(1/2)`

3.374.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00

$$\int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx = \frac{2i \operatorname{arctanh}\left(\frac{\sqrt{a+b \tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{a+ibd}}$$

input `Integrate[(1 - I*Tan[c + d*x])/Sqrt[a + b*Tan[c + d*x]],x]`

output `((2*I)*ArcTanh[Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d)`

3.374.3 Rubi [A] (warning: unable to verify)

Time = 0.24 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.78, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$, Rules used = {3042, 4020, 25, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4020} \\
 & - \frac{i \int - \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c + dx))}{d} \\
 & \quad \downarrow \text{25} \\
 & \frac{i \int \frac{1}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d(-i \tan(c + dx))}{d} \\
 & \quad \downarrow \text{73} \\
 & \frac{2 \int \frac{1}{-\frac{i \tan^2(c+dx)}{b} - \frac{ia}{b} + 1} d\sqrt{a + b \tan(c + dx)}}{bd} \\
 & \quad \downarrow \text{221} \\
 & \frac{2 \arctan\left(\frac{\tan(c+dx)}{\sqrt{a+ib}}\right)}{d\sqrt{a+ib}}
 \end{aligned}$$

input `Int[(1 - I*Tan[c + d*x])/Sqrt[a + b*Tan[c + d*x]],x]`

output `(2*ArcTan[Tan[c + d*x]/Sqrt[a + I*b]])/(Sqrt[a + I*b]*d)`

3.374.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

3.374.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 740 vs. $2(36) = 72$.

Time = 0.09 (sec) , antiderivative size = 741, normalized size of antiderivative = 16.47

method	result
derivatividedives	$\frac{(-i\sqrt{a^2+b^2}-ia-b) \ln\left(\frac{b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}}{2}\right) + 2\left(-i\sqrt{2\sqrt{a^2+b^2}+2a} a - \sqrt{2\sqrt{a^2+b^2}+2a}\right)}{\sqrt{2\sqrt{a^2+b^2}+2a} \sqrt{a^2+b^2}}$
default	$\frac{(-i\sqrt{a^2+b^2}-ia-b) \ln\left(\frac{b \tan(dx+c)+a+\sqrt{a+b \tan(dx+c)} \sqrt{2\sqrt{a^2+b^2}+2a+\sqrt{a^2+b^2}}}{2}\right) + 2\left(-i\sqrt{2\sqrt{a^2+b^2}+2a} a - \sqrt{2\sqrt{a^2+b^2}+2a}\right)}{\sqrt{2\sqrt{a^2+b^2}+2a} \sqrt{a^2+b^2}}$
parts	Expression too large to display

```
input int((1-I*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2), x, method=_RETURNVERBOSE)
```

```
output 1/d*(-1/(2*(a^2+b^2)^(1/2)+2*a)^(1/2)/(a^2+b^2)^(1/2)*(1/2*(-I*(a^2+b^2)^(1/2)-I*a-b)*ln(b*tan(d*x+c)+a+(a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)+(a^2+b^2)^(1/2))+2*(-I*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a-(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*b-1/2*(-I*(a^2+b^2)^(1/2)-I*a-b)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan((2*(a+b*tan(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)))-1/(2*(a^2+b^2)^(1/2)+2*a)^(1/2)/(a^2+b^2)^(1/2)/((a^2+b^2)^(1/2)*a+a^2+b^2)*(-1/2*(-2*I*(a^2+b^2)^(1/2)*a^2-I*(a^2+b^2)^(1/2)*b^2-2*I*a^3-2*I*a*b^2-(a^2+b^2)^(1/2)*a*b-a^2*b-b^3)*ln((a+b*tan(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)-b*tan(d*x+c)-a-(a^2+b^2)^(1/2))+2*(I*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*a^2+I*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^3+I*(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a*b^2+(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*(a^2+b^2)^(1/2)*a*b+(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*a^2*b+(2*(a^2+b^2)^(1/2)+2*a)^(1/2)*b^3+1/2*(-2*I*(a^2+b^2)^(1/2)*a^2-I*(a^2+b^2)^(1/2)*b^2-2*I*a^3-2*I*a*b^2-(a^2+b^2)^(1/2)*a*b-a^2*b-b^3)*(2*(a^2+b^2)^(1/2)+2*a)^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2)*arctan(((2*(a^2+b^2)^(1/2)+2*a)^(1/2)-2*(a+b*tan(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*a)^(1/2))))
```

3.374. $\int \frac{1-i \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$

3.374.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 267 vs. $2(33) = 66$.

Time = 0.24 (sec) , antiderivative size = 267, normalized size of antiderivative = 5.93

$$\int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx =$$

$$-\frac{1}{4} \sqrt{\frac{4i}{(-ia + b)d^2}} \log \left(\frac{\left((ia - b)de^{(2idx+2ic)} + (ia - b)d \right) \sqrt{\frac{(a-ib)e^{(2idx+2ic)} + a + ib}{e^{(2idx+2ic)} + 1}} \sqrt{\frac{4i}{(-ia+b)d^2}} + 2ae^{(2idx+2ic)}}{(-ia + b)d} \right)$$

$$+\frac{1}{4} \sqrt{\frac{4i}{(-ia + b)d^2}} \log \left(\frac{\left((-ia + b)de^{(2idx+2ic)} + (-ia + b)d \right) \sqrt{\frac{(a-ib)e^{(2idx+2ic)} + a + ib}{e^{(2idx+2ic)} + 1}} \sqrt{\frac{4i}{(-ia+b)d^2}} + 2ae^{(2idx+2ic)}}{(-ia + b)d} \right)$$

input `integrate((1-I*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="fricas")`

output `-1/4*sqrt(4*I/((-I*a + b)*d^2))*log((((I*a - b)*d*e^(2*I*d*x + 2*I*c) + (I*a - b)*d)*sqrt(((a - I*b)*e^(2*I*d*x + 2*I*c) + a + I*b)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(4*I/((-I*a + b)*d^2)) + 2*a*e^(2*I*d*x + 2*I*c) + 2*a + 2*I*b)*e^(-2*I*d*x - 2*I*c)/((-I*a + b)*d) + 1/4*sqrt(4*I/((-I*a + b)*d^2))*log(((((-I*a + b)*d*e^(2*I*d*x + 2*I*c) + (-I*a + b)*d)*sqrt(((a - I*b)*e^(2*I*d*x + 2*I*c) + a + I*b)/(e^(2*I*d*x + 2*I*c) + 1))*sqrt(4*I/((-I*a + b)*d^2)) + 2*a*e^(2*I*d*x + 2*I*c) + 2*a + 2*I*b)*e^(-2*I*d*x - 2*I*c)/((-I*a + b)*d))`

3.374.6 Sympy [F]

$$\int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = -i \left(\int \frac{i}{\sqrt{a + b \tan(c + dx)}} dx + \int \frac{\tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx \right)$$

input `integrate((1-I*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)`

output `-I*(Integral(I/sqrt(a + b*tan(c + d*x)), x) + Integral(tan(c + d*x)/sqrt(a + b*tan(c + d*x)), x))`

3.374. $\int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx$

3.374.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Exception raised: ValueError}$$

input `integrate((1-I*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see `assume?` for more details)Is

3.374.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 155 vs. 2(33) = 66.

Time = 0.40 (sec) , antiderivative size = 155, normalized size of antiderivative = 3.44

$$\int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx$$

$$= -\frac{4i \arctan\left(\frac{2(\sqrt{b \tan(dx+c)+a} - \sqrt{a^2+b^2} \sqrt{b \tan(dx+c)+a})}{a\sqrt{-2a+2\sqrt{a^2+b^2}}+i\sqrt{-2a+2\sqrt{a^2+b^2}}b-\sqrt{a^2+b^2}\sqrt{-2a+2\sqrt{a^2+b^2}}}\right)}{\sqrt{-2a+2\sqrt{a^2+b^2}}d\left(\frac{ib}{a-\sqrt{a^2+b^2}}+1\right)}$$

input `integrate((1-I*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `-4*I*arctan(2*(sqrt(b*tan(d*x + c) + a)*a - sqrt(a^2 + b^2)*sqrt(b*tan(d*x + c) + a))/(a*sqrt(-2*a + 2*sqrt(a^2 + b^2)) + I*sqrt(-2*a + 2*sqrt(a^2 + b^2))*b - sqrt(a^2 + b^2)*sqrt(-2*a + 2*sqrt(a^2 + b^2)))/(sqrt(-2*a + 2*sqrt(a^2 + b^2))*d*(I*b/(a - sqrt(a^2 + b^2)) + 1))`

3.374.9 Mupad [B] (verification not implemented)

Time = 9.22 (sec) , antiderivative size = 1410, normalized size of antiderivative = 31.33

$$\int \frac{1 - i \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

```
input int(-(tan(c + d*x)*1i - 1)/(a + b*tan(c + d*x))^(1/2),x)
```

```
output (log(d*(-1/(d^2*(a - b*1i)))^(1/2)*(a + b*tan(c + d*x))^(1/2) + 1i)*(-1/(a
*d^2 - b*d^2*1i))^(1/2))/2 - 2*atanh((32*b^2*((b*1i)/(4*a^2*d^2 + 4*b^2*d^
2) - a/(4*a^2*d^2 + 4*b^2*d^2))^(1/2)*(a + b*tan(c + d*x))^(1/2))/((a^2*b^
2*d^2*64i)/(4*a^2*d^3 + 4*b^2*d^3) - (b^2*16i)/d + (64*a*b^3*d^2)/(4*a^2*d
^3 + 4*b^2*d^3)) - (128*a^2*b^2*((b*1i)/(4*a^2*d^2 + 4*b^2*d^2) - a/(4*a^2
*d^2 + 4*b^2*d^2))^(1/2)*(a + b*tan(c + d*x))^(1/2))/((a^2*b^4*d^2*256i)/(
4*a^2*d^3 + 4*b^2*d^3) - (a^2*b^2*64i)/d - (b^4*64i)/d + (256*a^3*b^3*d^2)
/(4*a^2*d^3 + 4*b^2*d^3) + (a^4*b^2*d^2*256i)/(4*a^2*d^3 + 4*b^2*d^3) + (2
56*a*b^5*d^2)/(4*a^2*d^3 + 4*b^2*d^3)) + (a*b^3*((b*1i)/(4*a^2*d^2 + 4*b^2
*d^2) - a/(4*a^2*d^2 + 4*b^2*d^2))^(1/2)*(a + b*tan(c + d*x))^(1/2)*128i)/
((a^2*b^4*d^2*256i)/(4*a^2*d^3 + 4*b^2*d^3) - (a^2*b^2*64i)/d - (b^4*64i)/
d + (256*a^3*b^3*d^2)/(4*a^2*d^3 + 4*b^2*d^3) + (a^4*b^2*d^2*256i)/(4*a^2
*d^3 + 4*b^2*d^3) + (256*a*b^5*d^2)/(4*a^2*d^3 + 4*b^2*d^3)))*(-(a - b*1i)/
(4*a^2*d^2 + 4*b^2*d^2))^(1/2) - log(d*(-1/(d^2*(a - b*1i)))^(1/2)*(a + b*
tan(c + d*x))^(1/2)*1i + 1)*(-1/(4*(a*d^2 - b*d^2*1i)))^(1/2) + (log(16*b^
2*(a + b*tan(c + d*x))^(1/2) + 16*b^3*d*(-1/(d^2*(a - b*1i)))^(1/2) - (16*
a*b^2*(a + b*tan(c + d*x))^(1/2))/(a - b*1i))*(-1/(a*d^2 - b*d^2*1i))^(1/2
))/2 - log(16*b^3*d*(-1/(d^2*(a - b*1i)))^(1/2) - 16*b^2*(a + b*tan(c + d*
x))^(1/2) + (16*a*b^2*(a + b*tan(c + d*x))^(1/2))/(a - b*1i))*(-1/(4*(a*d^
2 - b*d^2*1i)))^(1/2) + 2*atanh((32*b^2*((b*1i)/(4*a^2*d^2 + 4*b^2*d^2)...

```

3.375 $\int \frac{3+\tan(x)}{\sqrt{4+3\tan(x)}} dx$

3.375.1 Optimal result	3560
3.375.2 Mathematica [C] (verified)	3560
3.375.3 Rubi [A] (verified)	3561
3.375.4 Maple [B] (verified)	3562
3.375.5 Fricas [A] (verification not implemented)	3562
3.375.6 Sympy [F]	3563
3.375.7 Maxima [F]	3563
3.375.8 Giac [B] (verification not implemented)	3563
3.375.9 Mupad [B] (verification not implemented)	3564

3.375.1 Optimal result

Integrand size = 15, antiderivative size = 30

$$\int \frac{3 + \tan(x)}{\sqrt{4 + 3 \tan(x)}} dx = -\sqrt{2} \arctan \left(\frac{1 - 3 \tan(x)}{\sqrt{2}\sqrt{4 + 3 \tan(x)}} \right)$$

output `-arctan(1/2*(1-3*tan(x))*2^(1/2)/(4+3*tan(x))^(1/2))*2^(1/2)`

3.375.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.77 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.03

$$\int \frac{3 + \tan(x)}{\sqrt{4 + 3 \tan(x)}} dx = -\frac{(1 + 3i)\operatorname{arctanh}\left(\frac{\sqrt{4+3\tan(x)}}{\sqrt{4-3i}}\right)}{\sqrt{4 - 3i}} - \frac{(1 - 3i)\operatorname{arctanh}\left(\frac{\sqrt{4+3\tan(x)}}{\sqrt{4+3i}}\right)}{\sqrt{4 + 3i}}$$

input `Integrate[(3 + Tan[x])/Sqrt[4 + 3*Tan[x]],x]`

output `((-1 - 3*I)*ArcTanh[Sqrt[4 + 3*Tan[x]]/Sqrt[4 - 3*I]])/Sqrt[4 - 3*I] - ((1 - 3*I)*ArcTanh[Sqrt[4 + 3*Tan[x]]/Sqrt[4 + 3*I]])/Sqrt[4 + 3*I]`

3.375.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 4018, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan(x) + 3}{\sqrt{3 \tan(x) + 4}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(x) + 3}{\sqrt{3 \tan(x) + 4}} dx \\ & \quad \downarrow \text{4018} \\ & -2 \int \frac{1}{\frac{(1-3 \tan(x))^2}{3 \tan(x)+4} + 2} d \frac{1-3 \tan(x)}{\sqrt{3 \tan(x) + 4}} \\ & \quad \downarrow \text{216} \\ & -\sqrt{2} \arctan \left(\frac{1-3 \tan(x)}{\sqrt{2} \sqrt{3 \tan(x) + 4}} \right) \end{aligned}$$

input `Int[(3 + Tan[x])/Sqrt[4 + 3*Tan[x]],x]`

output `-(Sqrt[2]*ArcTan[(1 - 3*Tan[x])/(Sqrt[2]*Sqrt[4 + 3*Tan[x]])])`

3.375.3.1 Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`


```
rule 4018 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(a_) + (b_.)*tan[(e_.) + (
f_.)*(x_)]], x_Symbol] :> Simp[-2*(d^2/f) Subst[Int[1/(2*b*c*d - 4*a*d^2
+ x^2), x], x, (b*c - 2*a*d - b*d*Tan[e + f*x])/Sqrt[a + b*Tan[e + f*x]]],
x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0
] && NeQ[c^2 + d^2, 0] && EqQ[2*a*c*d - b*(c^2 - d^2), 0]
```

3.375.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. $2(25) = 50$.

Time = 0.22 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.80

method	result	size
derivativedivides	$\sqrt{2} \arctan\left(\frac{(2\sqrt{4+3\tan(x)}-3\sqrt{2})\sqrt{2}}{2}\right) + \sqrt{2} \arctan\left(\frac{(2\sqrt{4+3\tan(x)}+3\sqrt{2})\sqrt{2}}{2}\right)$	54
default	$\sqrt{2} \arctan\left(\frac{(2\sqrt{4+3\tan(x)}-3\sqrt{2})\sqrt{2}}{2}\right) + \sqrt{2} \arctan\left(\frac{(2\sqrt{4+3\tan(x)}+3\sqrt{2})\sqrt{2}}{2}\right)$	54
parts	$\sqrt{2} \arctan\left(\frac{(2\sqrt{4+3\tan(x)}-3\sqrt{2})\sqrt{2}}{2}\right) + \sqrt{2} \arctan\left(\frac{(2\sqrt{4+3\tan(x)}+3\sqrt{2})\sqrt{2}}{2}\right)$	54

```
input int((3+tan(x))/(4+3*tan(x))^(1/2),x,method=_RETURNVERBOSE)
```

```
output 2^(1/2)*arctan(1/2*(2*(4+3*tan(x))^(1/2)-3*2^(1/2))*2^(1/2))+2^(1/2)*arctan(1/2*(2*(4+3*tan(x))^(1/2)+3*2^(1/2))*2^(1/2))
```

3.375.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{3 + \tan(x)}{\sqrt{4 + 3 \tan(x)}} dx = \sqrt{2} \arctan\left(\frac{3\sqrt{2} \tan(x) - \sqrt{2}}{2\sqrt{3} \tan(x) + 4}\right)$$

```
input integrate((3+tan(x))/(4+3*tan(x))^(1/2),x, algorithm="fricas")
```

```
output sqrt(2)*arctan(1/2*(3*sqrt(2)*tan(x) - sqrt(2))/sqrt(3*tan(x) + 4))
```

3.375.6 Sympy [F]

$$\int \frac{3 + \tan(x)}{\sqrt{4 + 3 \tan(x)}} dx = \int \frac{\tan(x) + 3}{\sqrt{3 \tan(x) + 4}} dx$$

input `integrate((3+tan(x))/(4+3*tan(x))**(1/2),x)`

output `Integral((tan(x) + 3)/sqrt(3*tan(x) + 4), x)`

3.375.7 Maxima [F]

$$\int \frac{3 + \tan(x)}{\sqrt{4 + 3 \tan(x)}} dx = \int \frac{\tan(x) + 3}{\sqrt{3 \tan(x) + 4}} dx$$

input `integrate((3+tan(x))/(4+3*tan(x))^(1/2),x, algorithm="maxima")`

output `integrate((tan(x) + 3)/sqrt(3*tan(x) + 4), x)`

3.375.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. $2(25) = 50$.

Time = 0.33 (sec) , antiderivative size = 65, normalized size of antiderivative = 2.17

$$\int \frac{3 + \tan(x)}{\sqrt{4 + 3 \tan(x)}} dx = \sqrt{2} \arctan \left(\frac{1}{250} \cdot 25^{\frac{3}{4}} \sqrt{10} \left(3 \cdot 25^{\frac{1}{4}} \sqrt{10} + 10 \sqrt{3 \tan(x) + 4} \right) \right) \\ + \sqrt{2} \arctan \left(-\frac{1}{250} \cdot 25^{\frac{3}{4}} \sqrt{10} \left(3 \cdot 25^{\frac{1}{4}} \sqrt{10} - 10 \sqrt{3 \tan(x) + 4} \right) \right)$$

input `integrate((3+tan(x))/(4+3*tan(x))^(1/2),x, algorithm="giac")`

output `sqrt(2)*arctan(1/250*25^(3/4)*sqrt(10)*(3*25^(1/4)*sqrt(10) + 10*sqrt(3*tan(x) + 4))) + sqrt(2)*arctan(-1/250*25^(3/4)*sqrt(10)*(3*25^(1/4)*sqrt(10) - 10*sqrt(3*tan(x) + 4)))`

3.375.9 Mupad [B] (verification not implemented)

Time = 0.86 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.03

$$\int \frac{3 + \tan(x)}{\sqrt{4 + 3 \tan(x)}} dx = \sqrt{2} \left(\operatorname{atan} \left(\sqrt{6 \tan(x) + 8} \left(\frac{1}{10} - \frac{3}{10}i \right) \right) \right. \\ \left. + \operatorname{atan} \left(\sqrt{6 \tan(x) + 8} \left(\frac{1}{10} + \frac{3}{10}i \right) \right) \right)$$

input `int((tan(x) + 3)/(3*tan(x) + 4)^(1/2),x)`output `2^(1/2)*(atan((6*tan(x) + 8)^(1/2)*(1/10 - 3i/10)) + atan((6*tan(x) + 8)^(1/2)*(1/10 + 3i/10)))`

3.376 $\int \frac{1-3 \tan(x)}{\sqrt{4+3 \tan(x)}} dx$

3.376.1 Optimal result 3565
 3.376.2 Mathematica [C] (verified) 3565
 3.376.3 Rubi [A] (verified) 3566
 3.376.4 Maple [B] (verified) 3567
 3.376.5 Fricas [B] (verification not implemented) 3567
 3.376.6 Sympy [F] 3568
 3.376.7 Maxima [F] 3568
 3.376.8 Giac [B] (verification not implemented) 3568
 3.376.9 Mupad [B] (verification not implemented) 3569

3.376.1 Optimal result

Integrand size = 17, antiderivative size = 27

$$\int \frac{1-3 \tan(x)}{\sqrt{4+3 \tan(x)}} dx = \sqrt{2} \operatorname{arctanh}\left(\frac{3+\tan(x)}{\sqrt{2}\sqrt{4+3 \tan(x)}}\right)$$

output `arctanh(1/2*(3+tan(x))*2^(1/2)/(4+3*tan(x))^(1/2))*2^(1/2)`

3.376.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.93 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.26

$$\int \frac{1-3 \tan(x)}{\sqrt{4+3 \tan(x)}} dx = \frac{(3-i) \operatorname{arctanh}\left(\frac{\sqrt{4+3 \tan(x)}}{\sqrt{4-3i}}\right)}{\sqrt{4-3i}} + \frac{(3+i) \operatorname{arctanh}\left(\frac{\sqrt{4+3 \tan(x)}}{\sqrt{4+3i}}\right)}{\sqrt{4+3i}}$$

input `Integrate[(1 - 3*Tan[x])/Sqrt[4 + 3*Tan[x]],x]`

output `((3 - I)*ArcTanh[Sqrt[4 + 3*Tan[x]]/Sqrt[4 - 3*I]]/Sqrt[4 - 3*I] + ((3 + I)*ArcTanh[Sqrt[4 + 3*Tan[x]]/Sqrt[4 + 3*I]]/Sqrt[4 + 3*I])`

3.376.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {3042, 4018, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1 - 3 \tan(x)}{\sqrt{3 \tan(x) + 4}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{1 - 3 \tan(x)}{\sqrt{3 \tan(x) + 4}} dx \\
 & \quad \downarrow \text{4018} \\
 & -18 \int \frac{1}{\frac{81(\tan(x)+3)^2}{3 \tan(x)+4} - 162} d \frac{9(\tan(x) + 3)}{\sqrt{3 \tan(x) + 4}} \\
 & \quad \downarrow \text{220} \\
 & \sqrt{2} \operatorname{arctanh} \left(\frac{\tan(x) + 3}{\sqrt{2} \sqrt{3 \tan(x) + 4}} \right)
 \end{aligned}$$

input `Int[(1 - 3*Tan[x])/Sqrt[4 + 3*Tan[x]],x]`

output `Sqrt[2]*ArcTanh[(3 + Tan[x])/(Sqrt[2]*Sqrt[4 + 3*Tan[x]])]`

3.376.3.1 Defintions of rubi rules used

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4018 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(a_.) + (b_.)*tan[(e_.) + (
f_.)*(x_)]], x_Symbol] :> Simp[-2*(d^2/f) Subst[Int[1/(2*b*c*d - 4*a*d^2
+ x^2), x], x, (b*c - 2*a*d - b*d*Tan[e + f*x])/Sqrt[a + b*Tan[e + f*x]]],
x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0
] && NeQ[c^2 + d^2, 0] && EqQ[2*a*c*d - b*(c^2 - d^2), 0]
```

3.376.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 51 vs. $2(22) = 44$.

Time = 0.05 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.93

method	result	size
derivativedivides	$-\frac{\sqrt{2} \ln(9+3 \tan(x)-3\sqrt{4+3 \tan(x)} \sqrt{2})}{2} + \frac{\sqrt{2} \ln(9+3 \tan(x)+3\sqrt{4+3 \tan(x)} \sqrt{2})}{2}$	52
default	$-\frac{\sqrt{2} \ln(9+3 \tan(x)-3\sqrt{4+3 \tan(x)} \sqrt{2})}{2} + \frac{\sqrt{2} \ln(9+3 \tan(x)+3\sqrt{4+3 \tan(x)} \sqrt{2})}{2}$	52
parts	$-\frac{\sqrt{2} \ln(9+3 \tan(x)-3\sqrt{4+3 \tan(x)} \sqrt{2})}{2} + \frac{\sqrt{2} \ln(9+3 \tan(x)+3\sqrt{4+3 \tan(x)} \sqrt{2})}{2}$	52

```
input int((1-3*tan(x))/(4+3*tan(x))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/2*2^(1/2)*ln(9+3*tan(x)-3*(4+3*tan(x))^(1/2)*2^(1/2))+1/2*2^(1/2)*ln(9+
3*tan(x)+3*(4+3*tan(x))^(1/2)*2^(1/2))
```

3.376.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 47 vs. $2(22) = 44$.

Time = 0.25 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.74

$$\int \frac{1-3 \tan(x)}{\sqrt{4+3 \tan(x)}} dx$$

$$= \frac{1}{2} \sqrt{2} \log \left(\frac{\tan(x)^2 + 2(\sqrt{2} \tan(x) + 3\sqrt{2})\sqrt{3 \tan(x) + 4} + 12 \tan(x) + 17}{\tan(x)^2 + 1} \right)$$

```
input integrate((1-3*tan(x))/(4+3*tan(x))^(1/2),x, algorithm="fricas")
```

output $1/2*\sqrt{2}*\log((\tan(x)^2 + 2*(\sqrt{2})*\tan(x) + 3*\sqrt{2})*\sqrt{3*\tan(x) + 4} + 12*\tan(x) + 17)/(\tan(x)^2 + 1))$

3.376.6 Sympy [F]

$$\int \frac{1 - 3 \tan(x)}{\sqrt{4 + 3 \tan(x)}} dx = - \int \frac{3 \tan(x)}{\sqrt{3 \tan(x) + 4}} dx - \int \left(-\frac{1}{\sqrt{3 \tan(x) + 4}} \right) dx$$

input `integrate((1-3*tan(x))/(4+3*tan(x))**(1/2),x)`

output `-Integral(3*tan(x)/sqrt(3*tan(x) + 4), x) - Integral(-1/sqrt(3*tan(x) + 4), x)`

3.376.7 Maxima [F]

$$\int \frac{1 - 3 \tan(x)}{\sqrt{4 + 3 \tan(x)}} dx = \int -\frac{3 \tan(x) - 1}{\sqrt{3 \tan(x) + 4}} dx$$

input `integrate((1-3*tan(x))/(4+3*tan(x))^(1/2),x, algorithm="maxima")`

output `-integrate((3*tan(x) - 1)/sqrt(3*tan(x) + 4), x)`

3.376.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 57 vs. $2(22) = 44$.

Time = 0.30 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.11

$$\int \frac{1 - 3 \tan(x)}{\sqrt{4 + 3 \tan(x)}} dx = \frac{1}{2} \sqrt{2} \log \left(\frac{3}{5} \cdot 25^{\frac{1}{4}} \sqrt{10} \sqrt{3 \tan(x) + 4} + 3 \tan(x) + 9 \right) - \frac{1}{2} \sqrt{2} \log \left(-\frac{3}{5} \cdot 25^{\frac{1}{4}} \sqrt{10} \sqrt{3 \tan(x) + 4} + 3 \tan(x) + 9 \right)$$

input `integrate((1-3*tan(x))/(4+3*tan(x))^(1/2),x, algorithm="giac")`

output `1/2*sqrt(2)*log(3/5*25^(1/4)*sqrt(10)*sqrt(3*tan(x) + 4) + 3*tan(x) + 9) -
1/2*sqrt(2)*log(-3/5*25^(1/4)*sqrt(10)*sqrt(3*tan(x) + 4) + 3*tan(x) + 9)`

3.376.9 Mupad [B] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.30

$$\int \frac{1 - 3 \tan(x)}{\sqrt{4 + 3 \tan(x)}} dx = \sqrt{2} \left(\operatorname{atan} \left(\sqrt{6 \tan(x) + 8} \left(\frac{1}{10} - \frac{3}{10}i \right) \right) - \operatorname{atan} \left(\sqrt{6 \tan(x) + 8} \left(\frac{1}{10} + \frac{3}{10}i \right) \right) \right) i$$

input `int(-(3*tan(x) - 1)/(3*tan(x) + 4)^(1/2),x)`

output `2^(1/2)*(atan((6*tan(x) + 8)^(1/2)*(1/10 - 3i/10)) - atan((6*tan(x) + 8)^(1/2)*(1/10 + 3i/10)))*1i`

3.377 $\int \frac{4-3 \tan(a+bx)}{\sqrt{4+3 \tan(a+bx)}} dx$

3.377.1 Optimal result 3570
 3.377.2 Mathematica [C] (verified) 3570
 3.377.3 Rubi [A] (verified) 3571
 3.377.4 Maple [A] (verified) 3573
 3.377.5 Fricas [B] (verification not implemented) 3574
 3.377.6 Sympy [F] 3575
 3.377.7 Maxima [F] 3575
 3.377.8 Giac [F(-1)] 3576
 3.377.9 Mupad [B] (verification not implemented) 3576

3.377.1 Optimal result

Integrand size = 25, antiderivative size = 85

$$\int \frac{4-3 \tan(a+bx)}{\sqrt{4+3 \tan(a+bx)}} dx = -\frac{9 \arctan\left(\frac{1-3 \tan(a+bx)}{\sqrt{2}\sqrt{4+3 \tan(a+bx)}}\right)}{5\sqrt{2}b} + \frac{13 \operatorname{arctanh}\left(\frac{3+\tan(a+bx)}{\sqrt{2}\sqrt{4+3 \tan(a+bx)}}\right)}{5\sqrt{2}b}$$

output `-9/10*arctan(1/2*(1-3*tan(b*x+a))*2^(1/2)/(4+3*tan(b*x+a))^(1/2))/b*2^(1/2)+13/10*arctanh(1/2*(3+tan(b*x+a))*2^(1/2)/(4+3*tan(b*x+a))^(1/2))/b*2^(1/2)`

3.377.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.88

$$\int \frac{4-3 \tan(a+bx)}{\sqrt{4+3 \tan(a+bx)}} dx = \frac{(3-4i) \operatorname{arctanh}\left(\frac{\sqrt{4+3 \tan(a+bx)}}{\sqrt{4-3i}}\right)}{\sqrt{4-3ib}} + \frac{(3+4i) \operatorname{arctanh}\left(\frac{\sqrt{4+3 \tan(a+bx)}}{\sqrt{4+3i}}\right)}{\sqrt{4+3ib}}$$

input `Integrate[(4 - 3*Tan[a + b*x])/Sqrt[4 + 3*Tan[a + b*x]], x]`

output $((3 - 4*I)*\text{ArcTanh}[\text{Sqrt}[4 + 3*\text{Tan}[a + b*x]]/\text{Sqrt}[4 - 3*I]])/(\text{Sqrt}[4 - 3*I]*b) + ((3 + 4*I)*\text{ArcTanh}[\text{Sqrt}[4 + 3*\text{Tan}[a + b*x]]/\text{Sqrt}[4 + 3*I]])/(\text{Sqrt}[4 + 3*I]*b)$

3.377.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$, Rules used = {3042, 4019, 27, 3042, 4018, 216, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{4 - 3 \tan(a + bx)}{\sqrt{3 \tan(a + bx) + 4}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{4 - 3 \tan(a + bx)}{\sqrt{3 \tan(a + bx) + 4}} dx \\ & \quad \downarrow \text{4019} \\ & \frac{1}{10} \int \frac{9(\tan(a + bx) + 3)}{\sqrt{3 \tan(a + bx) + 4}} dx - \frac{1}{10} \int -\frac{13(1 - 3 \tan(a + bx))}{\sqrt{3 \tan(a + bx) + 4}} dx \\ & \quad \downarrow \text{27} \\ & \frac{13}{10} \int \frac{1 - 3 \tan(a + bx)}{\sqrt{3 \tan(a + bx) + 4}} dx + \frac{9}{10} \int \frac{\tan(a + bx) + 3}{\sqrt{3 \tan(a + bx) + 4}} dx \\ & \quad \downarrow \text{3042} \\ & \frac{13}{10} \int \frac{1 - 3 \tan(a + bx)}{\sqrt{3 \tan(a + bx) + 4}} dx + \frac{9}{10} \int \frac{\tan(a + bx) + 3}{\sqrt{3 \tan(a + bx) + 4}} dx \\ & \quad \downarrow \text{4018} \\ & \frac{9 \int \frac{1}{\frac{(1 - 3 \tan(a + bx))^2}{3 \tan(a + bx) + 4} + 2} d \frac{1 - 3 \tan(a + bx)}{\sqrt{3 \tan(a + bx) + 4}}}{5b} - \frac{117 \int \frac{1}{\frac{81(\tan(a + bx) + 3)^2}{3 \tan(a + bx) + 4} - 162} d \frac{9(\tan(a + bx) + 3)}{\sqrt{3 \tan(a + bx) + 4}}}{5b} \\ & \quad \downarrow \text{216} \\ & \frac{117 \int \frac{1}{\frac{81(\tan(a + bx) + 3)^2}{3 \tan(a + bx) + 4} - 162} d \frac{9(\tan(a + bx) + 3)}{\sqrt{3 \tan(a + bx) + 4}}}{5b} - \frac{9 \arctan\left(\frac{1 - 3 \tan(a + bx)}{\sqrt{2}\sqrt{3 \tan(a + bx) + 4}}\right)}{5\sqrt{2}b} \\ & \quad \downarrow \text{220} \end{aligned}$$

3.377. $\int \frac{4 - 3 \tan(a + bx)}{\sqrt{4 + 3 \tan(a + bx)}} dx$

$$\frac{13 \operatorname{arctanh}\left(\frac{\tan(ax)+3}{\sqrt{2}\sqrt{3\tan(ax)+4}}\right)}{5\sqrt{2}b} - \frac{9 \arctan\left(\frac{1-3\tan(ax)}{\sqrt{2}\sqrt{3\tan(ax)+4}}\right)}{5\sqrt{2}b}$$

input `Int[(4 - 3*Tan[a + b*x])/Sqrt[4 + 3*Tan[a + b*x]],x]`

output `(-9*ArcTan[(1 - 3*Tan[a + b*x])/(Sqrt[2]*Sqrt[4 + 3*Tan[a + b*x]])])/(5*Sqrt[2]*b) + (13*ArcTanh[(3 + Tan[a + b*x])/(Sqrt[2]*Sqrt[4 + 3*Tan[a + b*x]])])/(5*Sqrt[2]*b)`

3.377.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4018 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2*(d^2/f) Subst[Int[1/(2*b*c*d - 4*a*d^2 + x^2), x], x, (b*c - 2*a*d - b*d*Tan[e + f*x])/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && EqQ[2*a*c*d - b*(c^2 - d^2), 0]`

```
rule 4019 Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := With[{q = Rt[a^2 + b^2, 2]}, Simp[1/(2*q) Int[(a*c + b*d + c*q + (b*c - a*d + d*q)*Tan[e + f*x])/Sqrt[a + b*Tan[e + f*x]], x], x] - Simp[1/(2*q) Int[(a*c + b*d - c*q + (b*c - a*d - d*q)*Tan[e + f*x])/Sqrt[a + b*Tan[e + f*x]], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && NeQ[2*a*c*d - b*(c^2 - d^2), 0] && NiceSqrtQ[a^2 + b^2]
```

3.377.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.58

method	result
derivativedivides	$\frac{13\sqrt{2} \ln(9+3 \tan(bx+a)+3\sqrt{4+3 \tan(bx+a)} \sqrt{2})}{20} + \frac{9\sqrt{2} \arctan\left(\frac{(2\sqrt{4+3 \tan(bx+a)+3\sqrt{2}})\sqrt{2}}{2}\right)}{10} - \frac{13\sqrt{2} \ln(9+3 \tan(bx+a)-3\sqrt{4+3 \tan(bx+a)} \sqrt{2})}{20}$
default	$\frac{13\sqrt{2} \ln(9+3 \tan(bx+a)+3\sqrt{4+3 \tan(bx+a)} \sqrt{2})}{20} + \frac{9\sqrt{2} \arctan\left(\frac{(2\sqrt{4+3 \tan(bx+a)+3\sqrt{2}})\sqrt{2}}{2}\right)}{10} - \frac{13\sqrt{2} \ln(9+3 \tan(bx+a)-3\sqrt{4+3 \tan(bx+a)} \sqrt{2})}{20}$
parts	$-\frac{\sqrt{2} \ln(9+3 \tan(bx+a)-3\sqrt{4+3 \tan(bx+a)} \sqrt{2})}{5b} + \frac{6\sqrt{2} \arctan\left(\frac{(2\sqrt{4+3 \tan(bx+a)-3\sqrt{2}})\sqrt{2}}{2}\right)}{5b} + \frac{\sqrt{2} \ln(9+3 \tan(bx+a)+3\sqrt{4+3 \tan(bx+a)} \sqrt{2})}{5b}$

```
input int((4-3*tan(b*x+a))/(4+3*tan(b*x+a))^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/b*(13/20*2^(1/2)*ln(9+3*tan(b*x+a)+3*(4+3*tan(b*x+a))^(1/2)*2^(1/2))+9/10*2^(1/2)*arctan(1/2*(2*(4+3*tan(b*x+a))^(1/2)+3*2^(1/2))*2^(1/2))-13/20*2^(1/2)*ln(9+3*tan(b*x+a)-3*(4+3*tan(b*x+a))^(1/2)*2^(1/2))+9/10*2^(1/2)*arctan(1/2*(2*(4+3*tan(b*x+a))^(1/2)-3*2^(1/2))*2^(1/2)))
```

3.377.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 309 vs. 2(71) = 142.

Time = 0.25 (sec) , antiderivative size = 309, normalized size of antiderivative = 3.64

$$\begin{aligned}
 & \int \frac{4 - 3 \tan(a + bx)}{\sqrt{4 + 3 \tan(a + bx)}} dx \\
 &= -\frac{1}{10} \sqrt{\frac{117 b^2 \sqrt{-\frac{1}{b^4}} + 44}{b^2}} \log \left(\frac{1}{5} \left(7 b^3 \sqrt{-\frac{1}{b^4}} - 24 b \right) \sqrt{\frac{117 b^2 \sqrt{-\frac{1}{b^4}} + 44}{b^2}} \right. \\
 & \qquad \qquad \qquad \left. + 25 \sqrt{3 \tan(bx + a) + 4} \right) \\
 &+ \frac{1}{10} \sqrt{\frac{117 b^2 \sqrt{-\frac{1}{b^4}} + 44}{b^2}} \log \left(-\frac{1}{5} \left(7 b^3 \sqrt{-\frac{1}{b^4}} - 24 b \right) \sqrt{\frac{117 b^2 \sqrt{-\frac{1}{b^4}} + 44}{b^2}} \right. \\
 & \qquad \qquad \qquad \left. + 25 \sqrt{3 \tan(bx + a) + 4} \right) \\
 &+ \frac{1}{10} \sqrt{-\frac{117 b^2 \sqrt{-\frac{1}{b^4}} - 44}{b^2}} \log \left(\frac{1}{5} \left(7 b^3 \sqrt{-\frac{1}{b^4}} + 24 b \right) \sqrt{-\frac{117 b^2 \sqrt{-\frac{1}{b^4}} - 44}{b^2}} \right. \\
 & \qquad \qquad \qquad \left. + 25 \sqrt{3 \tan(bx + a) + 4} \right) \\
 &- \frac{1}{10} \sqrt{-\frac{117 b^2 \sqrt{-\frac{1}{b^4}} - 44}{b^2}} \log \left(-\frac{1}{5} \left(7 b^3 \sqrt{-\frac{1}{b^4}} + 24 b \right) \sqrt{-\frac{117 b^2 \sqrt{-\frac{1}{b^4}} - 44}{b^2}} \right. \\
 & \qquad \qquad \qquad \left. + 25 \sqrt{3 \tan(bx + a) + 4} \right)
 \end{aligned}$$

input `integrate((4-3*tan(b*x+a))/(4+3*tan(b*x+a))^(1/2),x, algorithm="fracas")`

3.377. $\int \frac{4-3 \tan(a+bx)}{\sqrt{4+3 \tan(a+bx)}} dx$

output $-1/10*\sqrt{((117*b^2*\sqrt{-1/b^4} + 44)/b^2)*\log(1/5*(7*b^3*\sqrt{-1/b^4} - 24*b)*\sqrt{((117*b^2*\sqrt{-1/b^4} + 44)/b^2) + 25*\sqrt{3*\tan(b*x + a) + 4})} + 1/10*\sqrt{((117*b^2*\sqrt{-1/b^4} + 44)/b^2)*\log(-1/5*(7*b^3*\sqrt{-1/b^4} - 24*b)*\sqrt{((117*b^2*\sqrt{-1/b^4} + 44)/b^2) + 25*\sqrt{3*\tan(b*x + a) + 4})} + 1/10*\sqrt{-((117*b^2*\sqrt{-1/b^4} - 44)/b^2)*\log(1/5*(7*b^3*\sqrt{-1/b^4} + 24*b)*\sqrt{-((117*b^2*\sqrt{-1/b^4} - 44)/b^2) + 25*\sqrt{3*\tan(b*x + a) + 4})} - 1/10*\sqrt{-((117*b^2*\sqrt{-1/b^4} - 44)/b^2)*\log(-1/5*(7*b^3*\sqrt{-1/b^4} + 24*b)*\sqrt{-((117*b^2*\sqrt{-1/b^4} - 44)/b^2) + 25*\sqrt{3*\tan(b*x + a) + 4})}$

3.377.6 Sympy [F]

$$\int \frac{4 - 3 \tan(a + bx)}{\sqrt{4 + 3 \tan(a + bx)}} dx = - \int \frac{3 \tan(a + bx)}{\sqrt{3 \tan(a + bx) + 4}} dx - \int \left(-\frac{4}{\sqrt{3 \tan(a + bx) + 4}} \right) dx$$

input `integrate((4-3*tan(b*x+a))/(4+3*tan(b*x+a))**(1/2), x)`

output `-Integral(3*tan(a + b*x)/sqrt(3*tan(a + b*x) + 4), x) - Integral(-4/sqrt(3*tan(a + b*x) + 4), x)`

3.377.7 Maxima [F]

$$\int \frac{4 - 3 \tan(a + bx)}{\sqrt{4 + 3 \tan(a + bx)}} dx = \int -\frac{3 \tan(bx + a) - 4}{\sqrt{3 \tan(bx + a) + 4}} dx$$

input `integrate((4-3*tan(b*x+a))/(4+3*tan(b*x+a))^(1/2), x, algorithm="maxima")`

output `-integrate((3*tan(b*x + a) - 4)/sqrt(3*tan(b*x + a) + 4), x)`

3.377.8 Giac [F(-1)]

Timed out.

$$\int \frac{4 - 3 \tan(a + bx)}{\sqrt{4 + 3 \tan(a + bx)}} dx = \text{Timed out}$$

input `integrate((4-3*tan(b*x+a))/(4+3*tan(b*x+a))^(1/2),x, algorithm="giac")`

output `Timed out`

3.377.9 Mupad [B] (verification not implemented)

Time = 8.94 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.73

$$\begin{aligned} \int \frac{4 - 3 \tan(a + bx)}{\sqrt{4 + 3 \tan(a + bx)}} dx &= \operatorname{atan} \left(\frac{b \sqrt{\frac{-16 - 12i}{25} \frac{12i}{25}} \sqrt{3 \tan(a + bx) + 4}}{2} \right) \sqrt{\frac{-16 - 12i}{25} \frac{12i}{25}} 2i \\ &\quad - \operatorname{atan} \left(\frac{b \sqrt{\frac{-16 + 12i}{25} \frac{12i}{25}} \sqrt{3 \tan(a + bx) + 4}}{2} \right) \sqrt{\frac{-16 + 12i}{25} \frac{12i}{25}} 2i \\ &\quad + 2 \operatorname{atanh} \left(\frac{2b \sqrt{\frac{9 - 27i}{25} \frac{27i}{100}} \sqrt{3 \tan(a + bx) + 4}}{3} \right) \sqrt{\frac{9 - 27i}{25} \frac{27i}{100}} \\ &\quad + 2 \operatorname{atanh} \left(\frac{2b \sqrt{\frac{9 + 27i}{25} \frac{27i}{100}} \sqrt{3 \tan(a + bx) + 4}}{3} \right) \sqrt{\frac{9 + 27i}{25} \frac{27i}{100}} \end{aligned}$$

input `int(-(3*tan(a + b*x) - 4)/(3*tan(a + b*x) + 4)^(1/2),x)`

output `atan((b*((- 16/25 - 12i/25)/b^2)^(1/2)*(3*tan(a + b*x) + 4)^(1/2))/2)*((- 16/25 - 12i/25)/b^2)^(1/2)*2i - atan((b*((- 16/25 + 12i/25)/b^2)^(1/2)*(3*tan(a + b*x) + 4)^(1/2))/2)*((- 16/25 + 12i/25)/b^2)^(1/2)*2i + 2*atanh((2*b*((9/25 - 27i/100)/b^2)^(1/2)*(3*tan(a + b*x) + 4)^(1/2))/3)*((9/25 - 27i/100)/b^2)^(1/2) + 2*atanh((2*b*((9/25 + 27i/100)/b^2)^(1/2)*(3*tan(a + b*x) + 4)^(1/2))/3)*((9/25 + 27i/100)/b^2)^(1/2)`

3.378 $\int \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

3.378.1 Optimal result	3577
3.378.2 Mathematica [C] (verified)	3578
3.378.3 Rubi [A] (verified)	3578
3.378.4 Maple [A] (verified)	3584
3.378.5 Fricas [B] (verification not implemented)	3585
3.378.6 Sympy [F]	3585
3.378.7 Maxima [A] (verification not implemented)	3586
3.378.8 Giac [F(-1)]	3586
3.378.9 Mupad [B] (verification not implemented)	3587

3.378.1 Optimal result

Integrand size = 31, antiderivative size = 278

$$\begin{aligned} & \int \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx \\ &= \frac{(a(A-B) - b(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d} \\ & \quad - \frac{(a(A-B) - b(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d} \\ & \quad - \frac{(b(A-B) + a(A+B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}d} \\ & \quad + \frac{(b(A-B) + a(A+B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}d} \\ & \quad - \frac{2(Ab + aB)\sqrt{\tan(c+dx)}}{d} + \frac{2(aA - bB)\tan^{\frac{3}{2}}(c+dx)}{3d} \\ & \quad + \frac{2(Ab + aB)\tan^{\frac{5}{2}}(c+dx)}{5d} + \frac{2bB\tan^{\frac{7}{2}}(c+dx)}{7d} \end{aligned}$$

output
$$-1/2*(a*(A-B)-b*(A+B))*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/d*2^{(1/2)}-1/2*(a*(A-B)-b*(A+B))*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/d*2^{(1/2)}-1/4*(b*(A-B)+a*(A+B))*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/d*2^{(1/2)}+1/4*(b*(A-B)+a*(A+B))*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/d*2^{(1/2)}-2*(A*b+B*a)*\tan(d*x+c)^{(1/2)}/d+2/3*(A*a-B*b)*\tan(d*x+c)^{(3/2)}/d+2/5*(A*b+B*a)*\tan(d*x+c)^{(5/2)}/d+2/7*b*B*\tan(d*x+c)^{(7/2)}/d$$

3.378.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.74 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.54

$$\int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{-105\sqrt[4]{-1}(ia + b)(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right) + 105(-1)^{3/4}(a + ib)(A + iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{d}$$

input `Integrate[Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output
$$\frac{(-105*(-1)^{(1/4)}*(I*a + b)*(A - I*B)*\operatorname{ArcTan}[(-1)^{(3/4)}*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]] + 105*(-1)^{(3/4)}*(a + I*b)*(A + I*B)*\operatorname{ArcTanh}[(-1)^{(3/4)}*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]] + 2*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]*(-105*(A*b + a*B) + 35*(a*A - b*B)*\operatorname{Tan}[c + d*x] + 21*(A*b + a*B)*\operatorname{Tan}[c + d*x]^2 + 15*b*B*\operatorname{Tan}[c + d*x]^3))/(105*d)}$$

3.378.3 Rubi [A] (verified)

Time = 0.91 (sec) , antiderivative size = 255, normalized size of antiderivative = 0.92, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.581$, Rules used = {3042, 4075, 3042, 4011, 3042, 4011, 3042, 4011, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

↓ 3042

3.378. $\int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \int \tan(c+dx)^{5/2}(a+b \tan(c+dx))(A+B \tan(c+dx))dx \\
& \quad \downarrow 4075 \\
& \int \tan^{5/2}(c+dx)(aA-bB+(Ab+aB) \tan(c+dx))dx + \frac{2bB \tan^{7/2}(c+dx)}{7d} \\
& \quad \downarrow 3042 \\
& \int \tan(c+dx)^{5/2}(aA-bB+(Ab+aB) \tan(c+dx))dx + \frac{2bB \tan^{7/2}(c+dx)}{7d} \\
& \quad \downarrow 4011 \\
& \int \tan^{3/2}(c+dx)(-Ab-aB+(aA-bB) \tan(c+dx))dx + \frac{2(aB+Ab) \tan^{5/2}(c+dx)}{5d} + \frac{2bB \tan^{7/2}(c+dx)}{7d} \\
& \quad \downarrow 3042 \\
& \int \tan(c+dx)^{3/2}(-Ab-aB+(aA-bB) \tan(c+dx))dx + \frac{2(aB+Ab) \tan^{5/2}(c+dx)}{5d} + \\
& \quad \quad \quad \frac{2bB \tan^{7/2}(c+dx)}{7d} \\
& \quad \downarrow 4011 \\
& \int \sqrt{\tan(c+dx)}(-aA+bB-(Ab+aB) \tan(c+dx))dx + \frac{2(aB+Ab) \tan^{5/2}(c+dx)}{5d} + \\
& \quad \quad \quad \frac{2(aA-bB) \tan^{3/2}(c+dx)}{3d} + \frac{2bB \tan^{7/2}(c+dx)}{7d} \\
& \quad \downarrow 3042 \\
& \int \sqrt{\tan(c+dx)}(-aA+bB-(Ab+aB) \tan(c+dx))dx + \frac{2(aB+Ab) \tan^{5/2}(c+dx)}{5d} + \\
& \quad \quad \quad \frac{2(aA-bB) \tan^{3/2}(c+dx)}{3d} + \frac{2bB \tan^{7/2}(c+dx)}{7d} \\
& \quad \downarrow 4011 \\
& \int \frac{Ab+aB-(aA-bB) \tan(c+dx)}{\sqrt{\tan(c+dx)}}dx + \frac{2(aB+Ab) \tan^{5/2}(c+dx)}{5d} + \\
& \quad \frac{2(aA-bB) \tan^{3/2}(c+dx)}{3d} - \frac{2(aB+Ab) \sqrt{\tan(c+dx)}}{d} + \frac{2bB \tan^{7/2}(c+dx)}{7d} \\
& \quad \downarrow 3042
\end{aligned}$$

3.378. $\int \tan^{5/2}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

$$\int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2(aB + Ab) \tan^{\frac{5}{2}}(c + dx)}{5d} + \frac{2(aA - bB) \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2(aB + Ab) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{7}{2}}(c + dx)}{7d}$$

↓ 4017

$$2 \int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} + \frac{2(aB + Ab) \tan^{\frac{5}{2}}(c + dx)}{5d} + \frac{2(aA - bB) \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2(aB + Ab) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{7}{2}}(c + dx)}{7d}$$

↓ 1482

$$2 \left(\frac{1}{2} (a(A + B) + b(A - B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} - \frac{1}{2} (a(A - B) - b(A + B)) \int \frac{\tan(c + dx) + 1}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} \right)$$

$$\frac{2(aB + Ab) \tan^{\frac{5}{2}}(c + dx)}{5d} + \frac{2(aA - bB) \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2(aB + Ab) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{7}{2}}(c + dx)}{7d}$$

↓ 1476

$$2 \left(\frac{1}{2} (a(A + B) + b(A - B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} - \frac{1}{2} (a(A - B) - b(A + B)) \left(\frac{1}{2} \int \frac{1}{\tan(c + dx) - \sqrt{2} \sqrt{\tan(c + dx)}} \right) \right)$$

$$\frac{2(aB + Ab) \tan^{\frac{5}{2}}(c + dx)}{5d} + \frac{2(aA - bB) \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2(aB + Ab) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{7}{2}}(c + dx)}{7d}$$

↓ 1082

$$2 \left(\frac{1}{2} (a(A + B) + b(A - B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} - \frac{1}{2} (a(A - B) - b(A + B)) \left(\int \frac{1}{-\tan(c + dx) - 1} d \left(\frac{1 - \sqrt{2} \sqrt{\tan(c + dx)}}{\sqrt{2}} \right) \right) \right)$$

$$\frac{2(aB + Ab) \tan^{\frac{5}{2}}(c + dx)}{5d} + \frac{2(aA - bB) \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2(aB + Ab) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{7}{2}}(c + dx)}{7d}$$

↓ 217

3.378. $\int \tan^{\frac{5}{2}}(c + dx) (a + b \tan(c + dx)) (A + B \tan(c + dx)) dx$

$$2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)$$

$$\frac{2(aB + Ab) \tan^{\frac{5}{2}}(c+dx)}{5d} + \frac{2(aA - bB) \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2(aB + Ab) \sqrt{\tan(c+dx)}}{d} + \frac{2bB \tan^{\frac{7}{2}}(c+dx)}{7d}$$

↓ 1479

$$2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)$$

$$\frac{2(aB + Ab) \tan^{\frac{5}{2}}(c+dx)}{5d} + \frac{2(aA - bB) \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2(aB + Ab) \sqrt{\tan(c+dx)}}{d} + \frac{2bB \tan^{\frac{7}{2}}(c+dx)}{7d}$$

↓ 25

$$2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right) - \frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right)$$

$$\frac{2(aB + Ab) \tan^{\frac{5}{2}}(c+dx)}{5d} + \frac{2(aA - bB) \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2(aB + Ab) \sqrt{\tan(c+dx)}}{d} + \frac{2bB \tan^{\frac{7}{2}}(c+dx)}{7d}$$

↓ 27

$$2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)$$

$$\frac{2(aB + Ab) \tan^{\frac{5}{2}}(c+dx)}{5d} + \frac{2(aA - bB) \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2(aB + Ab) \sqrt{\tan(c+dx)}}{d} + \frac{2bB \tan^{\frac{7}{2}}(c+dx)}{7d}$$

↓ 1103

3.378. $\int \tan^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))(A+B\tan(c+dx)) dx$

$$2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(a(A-B) - b(A+B)) \right) \\ + \frac{2(aB + Ab) \tan^{\frac{5}{2}}(c+dx)}{5d} + \frac{2(aA - bB) \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2(aB + Ab) \sqrt{\tan(c+dx)}}{d} + \frac{2bB \tan^{\frac{7}{2}}(c+dx)}{7d}$$

input `Int[Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(2*(-1/2*((a*(A - B) - b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]/Sqrt[2]] + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]/Sqrt[2]])) + ((b*(A - B) + a*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d - (2*(A*b + a*B)*Sqrt[Tan[c + d*x]])/d + (2*(a*A - b*B)*Tan[c + d*x]^(3/2))/(3*d) + (2*(A*b + a*B)*Tan[c + d*x]^(5/2))/(5*d) + (2*b*B*Tan[c + d*x]^(7/2))/(7*d)`

3.378.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x], Sqrt[b*Tan[e + f*x]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

```
rule 4075 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

3.378.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 275, normalized size of antiderivative = 0.99

method	result
derivativedivides	$\frac{2Bb \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} + \frac{2Ab \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{2Ba \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{2aA \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} - \frac{2Bb \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} - 2Ab(\sqrt{\tan(dx+c)})$
default	$\frac{2Bb \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} + \frac{2Ab \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{2Ba \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{2aA \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} - \frac{2Bb \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} - 2Ab(\sqrt{\tan(dx+c)})$
parts	$(Ab+Ba) \left(\frac{2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} - 2(\sqrt{\tan(dx+c)}) + \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})}{1-\sqrt{2}(\sqrt{\tan(dx+c)})} \right) \right)}{4} \right)$

```
input int(tan(d*x+c)^(5/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

```
output 1/d*(2/7*B*b*tan(d*x+c)^(7/2)+2/5*A*b*tan(d*x+c)^(5/2)+2/5*B*a*tan(d*x+c)^(
(5/2)+2/3*a*A*tan(d*x+c)^(3/2)-2/3*B*b*tan(d*x+c)^(3/2)-2*A*b*tan(d*x+c)^(
(1/2)-2*B*a*tan(d*x+c)^(1/2)+1/4*(A*b+B*a)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)
)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(
1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(-A*a+B
*b)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x
+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(
1/2)*tan(d*x+c)^(1/2))))
```

$$3.378. \int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

3.378.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2266 vs. $2(236) = 472$.

Time = 0.39 (sec) , antiderivative size = 2266, normalized size of antiderivative = 8.15

$$\int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^(5/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm=
"fricas")
```

```
output -1/210*(105*d*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-
((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B
^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)
/d^4))/d^2)*log(((A*a - B*b)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A
^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^
3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) - ((A^2*B - B^3)*a^3 + (A^3 -
5*A*B^2)*a^2*b - (5*A^2*B - B^3)*a*b^2 - (A^3 - A*B^2)*b^3)*d)*sqrt((2*A*
B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)
*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A
^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2) + ((A^4 - B^
4)*a^4 - 4*(A^3*B + A*B^3)*a^3*b - 4*(A^3*B + A*B^3)*a*b^3 - (A^4 - B^4)*b
^4)*sqrt(tan(d*x + c))) - 105*d*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)
)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b -
2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A
^2*B^2 + B^4)*b^4)/d^4))/d^2)*log(-((A*a - B*b)*d^3*sqrt(-((A^4 - 2*A^2*B^
2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^
2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) - ((A^2*B
- B^3)*a^3 + (A^3 - 5*A*B^2)*a^2*b - (5*A^2*B - B^3)*a*b^2 - (A^3 - A*B^2)
*b^3)*d)*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4
- 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2...
```

3.378.6 Sympy [F]

$$\begin{aligned} & \int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx \\ &= \int (A + B \tan(c + dx))(a + b \tan(c + dx)) \tan^{\frac{5}{2}}(c + dx) dx \end{aligned}$$

input `integrate(tan(d*x+c)**(5/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))*tan(c + d*x)**(5/2), x)`

3.378.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 227, normalized size of antiderivative = 0.82

$$\int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{120 B b \tan(dx + c)^{\frac{7}{2}} + 168 (Ba + Ab) \tan(dx + c)^{\frac{5}{2}} - 210 \sqrt{2}((A - B)a - (A + B)b) \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2} \tan(dx + c) + 1)\right) + 105 \sqrt{2}((A + B)a + (A - B)b) \log(\sqrt{2} \sqrt{\tan(dx + c)} + \tan(dx + c) + 1) - 105 \sqrt{2}((A + B)a + (A - B)b) \log(-\sqrt{2} \sqrt{\tan(dx + c)} + \tan(dx + c) + 1) + 280(Aa - Bb) \tan(dx + c)^{\frac{3}{2}} - 840(Ba + Ab) \sqrt{\tan(dx + c)}}{d}$$

input `integrate(tan(d*x+c)^(5/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/420*(120*B*b*tan(d*x + c)^(7/2) + 168*(B*a + A*b)*tan(d*x + c)^(5/2) - 210*sqrt(2)*((A - B)*a - (A + B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) - 210*sqrt(2)*((A - B)*a - (A + B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + 105*sqrt(2)*((A + B)*a + (A - B)*b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - 105*sqrt(2)*((A + B)*a + (A - B)*b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 280*(A*a - B*b)*tan(d*x + c)^(3/2) - 840*(B*a + A*b)*sqrt(tan(d*x + c)))/d`

3.378.8 Giac [F(-1)]

Timed out.

$$\int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(5/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.378. $\int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

3.378.9 Mupad [B] (verification not implemented)

Time = 18.21 (sec) , antiderivative size = 1522, normalized size of antiderivative = 5.47

$$\int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(tan(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)
```

```
output atan((A^2*a^2*tan(c + d*x)^(1/2)*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4) + (A^2*a*b)/(2*d^2))^(1/2)*32i)/((16*A*b*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2))/d^3 - (16*A^3*a^3)/d + (16*A^3*a*b^2)/d) - (A^2*b^2*tan(c + d*x)^(1/2)*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4) + (A^2*a*b)/(2*d^2))^(1/2)*32i)/((16*A*b*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2))/d^3 - (16*A^3*a^3)/d + (16*A^3*a*b^2)/d))*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4) + (A^2*a*b)/(2*d^2))^(1/2)*2i - atan((A^2*a^2*tan(c + d*x)^(1/2)*((A^2*a*b)/(2*d^2) - (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4))^(1/2)*32i)/((16*A^3*a^3)/d + (16*A*b*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2))/d^3 - (16*A^3*a*b^2)/d) - (A^2*b^2*tan(c + d*x)^(1/2)*((A^2*a*b)/(2*d^2) - (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4))^(1/2)*32i)/((16*A^3*a^3)/d + (16*A*b*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2))/d^3 - (16*A^3*a*b^2)/d))*((A^2*a*b)/(2*d^2) - (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4))^(1/2)*2i + atan((B^2*a^2*tan(c + d*x)^(1/2)*(- (2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^(1/2)/(4*d^4) - (B^2*a*b)/(2*d^2))^(1/2)*32i)/((16*B*a*(2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^(1/2))/d^3 - (16*B^3*b^3)/d + (16*B^3*a^2*b)/d) - (B^2*b^2*tan(c + d*x)^(1/2)*(- (2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^(1/2)/(4*d^4) - (B^2*a*b)/(2*d^2))^(1/2)*32...
```

3.379 $\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

3.379.1 Optimal result	3588
3.379.2 Mathematica [C] (verified)	3589
3.379.3 Rubi [A] (verified)	3589
3.379.4 Maple [A] (verified)	3594
3.379.5 Fricas [B] (verification not implemented)	3595
3.379.6 Sympy [F]	3596
3.379.7 Maxima [A] (verification not implemented)	3596
3.379.8 Giac [F(-1)]	3597
3.379.9 Mupad [B] (verification not implemented)	3597

3.379.1 Optimal result

Integrand size = 31, antiderivative size = 254

$$\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{(b(A-B)+a(A+B)) \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(b(A-B)+a(A+B)) \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(a(A-B)-b(A+B)) \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(a(A-B)-b(A+B)) \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{2(aA-bB)\sqrt{\tan(c+dx)}}{d} + \frac{2(Ab+aB)\tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB\tan^{\frac{5}{2}}(c+dx)}{5d}$$

output

```
-1/2*(b*(A-B)+a*(A+B))*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)-1/2*(
b*(A-B)+a*(A+B))*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)+1/4*(a*(A-B)
-b*(A+B))*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)-1/4*(a*(A-B)
-b*(A+B))*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)+2*(A*a-B*b)*
tan(d*x+c)^(1/2)/d+2/3*(A*b+B*a)*tan(d*x+c)^(3/2)/d+2/5*b*B*tan(d*x+c)^(5/
2)/d
```

3.379.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.10 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.53

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{15\sqrt[4]{-1}(a - ib)(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right) + 15\sqrt[4]{-1}(a + ib)(A + iB) \operatorname{arctanh}\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{15d}$$

input `Integrate[Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(15*(-1)^(1/4)*(a - I*b)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + 15*(-1)^(1/4)*(a + I*b)*(A + I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + 2*Sqrt[Tan[c + d*x]]*(15*(a*A - b*B) + 5*(A*b + a*B)*Tan[c + d*x] + 3*b*B*Tan[c + d*x]^2))/(15*d)`

3.379.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.91, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.548$, Rules used = {3042, 4075, 3042, 4011, 3042, 4011, 3042, 4017, 25, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^{3/2}(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4075}$$

$$\int \tan^{\frac{3}{2}}(c + dx)(aA - bB + (Ab + aB) \tan(c + dx)) dx + \frac{2bB \tan^{\frac{5}{2}}(c + dx)}{5d}$$

$$\downarrow \text{3042}$$

3.379. $\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \int \tan(c+dx)^{3/2}(aA-bB+(Ab+aB)\tan(c+dx))dx + \frac{2bB\tan^{5/2}(c+dx)}{5d} \\
& \quad \downarrow 4011 \\
& \int \sqrt{\tan(c+dx)}(-Ab-aB+(aA-bB)\tan(c+dx))dx + \frac{2(aB+Ab)\tan^{3/2}(c+dx)}{3d} + \\
& \quad \frac{2bB\tan^{5/2}(c+dx)}{5d} \\
& \quad \downarrow 3042 \\
& \int \sqrt{\tan(c+dx)}(-Ab-aB+(aA-bB)\tan(c+dx))dx + \frac{2(aB+Ab)\tan^{3/2}(c+dx)}{3d} + \\
& \quad \frac{2bB\tan^{5/2}(c+dx)}{5d} \\
& \quad \downarrow 4011 \\
& \int \frac{-aA+bB-(Ab+aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}}dx + \frac{2(aB+Ab)\tan^{3/2}(c+dx)}{3d} + \\
& \quad \frac{2(aA-bB)\sqrt{\tan(c+dx)}}{d} + \frac{2bB\tan^{5/2}(c+dx)}{5d} \\
& \quad \downarrow 3042 \\
& \int \frac{-aA+bB-(Ab+aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}}dx + \frac{2(aB+Ab)\tan^{3/2}(c+dx)}{3d} + \\
& \quad \frac{2(aA-bB)\sqrt{\tan(c+dx)}}{d} + \frac{2bB\tan^{5/2}(c+dx)}{5d} \\
& \quad \downarrow 4017 \\
& \frac{2\int \frac{aA-bB+(Ab+aB)\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}}{d} + \frac{2(aB+Ab)\tan^{3/2}(c+dx)}{3d} + \\
& \quad \frac{2(aA-bB)\sqrt{\tan(c+dx)}}{d} + \frac{2bB\tan^{5/2}(c+dx)}{5d} \\
& \quad \downarrow 25 \\
& -\frac{2\int \frac{aA-bB+(Ab+aB)\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}}{d} + \frac{2(aB+Ab)\tan^{3/2}(c+dx)}{3d} + \\
& \quad \frac{2(aA-bB)\sqrt{\tan(c+dx)}}{d} + \frac{2bB\tan^{5/2}(c+dx)}{5d} \\
& \quad \downarrow 1482
\end{aligned}$$

$$\frac{2\left(-\frac{1}{2}(a(A-B) - b(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a(A+B) + b(A-B)) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right)}{\frac{2(aB + Ab) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2(aA - bB) \sqrt{\tan(c+dx)}}{d} + \frac{2bB \tan^{\frac{5}{2}}(c+dx)}{5d}}$$

↓ 1476

$$\frac{2\left(-\frac{1}{2}(a(A-B) - b(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)}} d\right)\right)}{\frac{2(aB + Ab) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2(aA - bB) \sqrt{\tan(c+dx)}}{d} + \frac{2bB \tan^{\frac{5}{2}}(c+dx)}{5d}}$$

↓ 1082

$$\frac{2\left(-\frac{1}{2}(a(A-B) - b(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a(A+B) + b(A-B)) \left(\int \frac{1}{-\tan(c+dx)-1} \frac{d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}}\right)\right)}{\frac{2(aB + Ab) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2(aA - bB) \sqrt{\tan(c+dx)}}{d} + \frac{2bB \tan^{\frac{5}{2}}(c+dx)}{5d}}$$

↓ 217

$$\frac{2\left(-\frac{1}{2}(a(A-B) - b(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}}\right)\right)}{\frac{2(aB + Ab) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2(aA - bB) \sqrt{\tan(c+dx)}}{d} + \frac{2bB \tan^{\frac{5}{2}}(c+dx)}{5d}}$$

↓ 1479

$$\frac{2\left(-\frac{1}{2}(a(A-B) - b(A+B)) \left(-\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}\right)\right)}{\frac{2(aB + Ab) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2(aA - bB) \sqrt{\tan(c+dx)}}{d} + \frac{2bB \tan^{\frac{5}{2}}(c+dx)}{5d}}$$

↓ 25

$$\begin{aligned}
 & \frac{2 \left(-\frac{1}{2}(a(A - B) - b(A + B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2}(a(A + B) + b(A - B)) \left(\int \frac{\sqrt{2}+2\sqrt{\tan(c+dx)}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d} \\
 & \quad \frac{2(aB + Ab) \tan^{\frac{3}{2}}(c + dx)}{3d} + \frac{2(aA - bB)\sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{5}{2}}(c + dx)}{5d} \\
 & \quad \downarrow \text{27} \\
 & \frac{2 \left(-\frac{1}{2}(a(A - B) - b(A + B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c + dx)} \right) - \frac{1}{2}(a(A + B) + b(A - B)) \left(\int \frac{\sqrt{2}+2\sqrt{\tan(c+dx)}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d} \\
 & \quad \frac{2(aB + Ab) \tan^{\frac{3}{2}}(c + dx)}{3d} + \frac{2(aA - bB)\sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{5}{2}}(c + dx)}{5d} \\
 & \quad \downarrow \text{1103} \\
 & \frac{2 \left(-\frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\arctan(1+\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)-1})}{\sqrt{2}} \right) \right)}{d} \\
 & \quad \frac{2(aB + Ab) \tan^{\frac{3}{2}}(c + dx)}{3d} + \frac{2(aA - bB)\sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{5}{2}}(c + dx)}{5d}
 \end{aligned}$$

```
input Int[Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]
```

```
output (2*(-1/2*((b*(A - B) + a*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])) - ((a*(A - B) - b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d + (2*(a*A - b*B)*Sqrt[Tan[c + d*x]])/d + (2*(A*b + a*B)*Tan[c + d*x]^(3/2))/(3*d) + (2*b*B*Tan[c + d*x]^(5/2))/(5*d)
```

3.379.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

3.379. $\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*(a + b*Tan[e + f*x])^m/(f*m), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /;` `FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /;` `FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /;` `FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.379.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{2Bb \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} + \frac{2Ab \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + \frac{2Ba \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + 2aA(\sqrt{\tan(dx+c)}) - 2(\sqrt{\tan(dx+c)})Bb + \frac{(-aA+Bb)\sqrt{2} \left(\ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right)}{4}$
default	$\frac{2Bb \left(\tan^{\frac{5}{2}}(dx+c) \right)}{5} + \frac{2Ab \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + \frac{2Ba \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + 2aA(\sqrt{\tan(dx+c)}) - 2(\sqrt{\tan(dx+c)})Bb + \frac{(-aA+Bb)\sqrt{2} \left(\ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right)}{4}$
parts	$\frac{(Ab+Ba) \left(\frac{2 \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} - \frac{\sqrt{2} \left(\ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right)}{4} \right)}{d}$

input `int(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

$$3.379. \quad \int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

```
output 1/d*(2/5*B*b*tan(d*x+c)^(5/2)+2/3*A*b*tan(d*x+c)^(3/2)+2/3*B*a*tan(d*x+c)^(3/2)+2*a*A*tan(d*x+c)^(1/2)-2*tan(d*x+c)^(1/2)*B*b+1/4*(-A*a+B*b)*2^(1/2)*
*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2))+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(-A*b-B*a)*2^(1/2)*
*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2))+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))))
```

3.379.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2252 vs. 2(216) = 432.

Time = 0.38 (sec) , antiderivative size = 2252, normalized size of antiderivative = 8.87

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output -1/30*(15*d*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-(A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2)*log(((B*a + A*b)*d^3*sqrt(-(A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) + ((A^3 - A*B^2)*a^3 - (5*A^2*B - B^3)*a^2*b - (A^3 - 5*A*B^2)*a*b^2 + (A^2*B - B^3)*b^3)*d)*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-(A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2) + ((A^4 - B^4)*a^4 - 4*(A^3*B + A*B^3)*a^3*b - 4*(A^3*B + A*B^3)*a*b^3 - (A^4 - B^4)*b^4)*sqrt(tan(d*x + c))) - 15*d*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-(A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2)*log(-(B*a + A*b)*d^3*sqrt(-(A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) + ((A^3 - A*B^2)*a^3 - (5*A^2*B - B^3)*a^2*b - (A^3 - 5*A*B^2)*a*b^2 + (A^2*B - B^3)*b^3)*d)*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-(A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2) + ((A^4 - B^4)*a^4 - 4*(A^3*B + A*B^3)*a^3*b - 4*(A^3*B + A*B^3)*a*b^3 - (A^4 - B^4)*b^4)*sqrt(tan(d*x + c)))
```

3.379.6 Sympy [F]

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx))(a + b \tan(c + dx)) \tan^{\frac{3}{2}}(c + dx) dx$$

input `integrate(tan(d*x+c)**(3/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))*tan(c + d*x)**(3/2), x)`

3.379.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 210, normalized size of antiderivative = 0.83

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{24 B b \tan(dx + c)^{\frac{5}{2}} - 30 \sqrt{2}((A + B)a + (A - B)b) \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2} + 2 \sqrt{\tan(dx + c)})\right) - 30 \sqrt{2}((A - B)a + (A + B)b) \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2} - 2 \sqrt{\tan(dx + c)})\right) - 15 \sqrt{2}((A - B)a - (A + B)b) \log(\sqrt{2} \sqrt{\tan(dx + c)} + \tan(dx + c) + 1) + 15 \sqrt{2}((A - B)a - (A + B)b) \log(-\sqrt{2} \sqrt{\tan(dx + c)} + \tan(dx + c) + 1) + 40(Ba + Ab) \tan(dx + c)^{\frac{3}{2}} + 120(Aa - Bb) \sqrt{\tan(dx + c)}}{d}$$

input `integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/60*(24*B*b*tan(d*x + c)^(5/2) - 30*sqrt(2)*((A + B)*a + (A - B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) - 30*sqrt(2)*((A + B)*a + (A - B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - 15*sqrt(2)*((A - B)*a - (A + B)*b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 15*sqrt(2)*((A - B)*a - (A + B)*b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 40*(B*a + A*b)*tan(d*x + c)^(3/2) + 120*(A*a - B*b)*sqrt(tan(d*x + c)))/d`

3.379.8 Giac [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \text{Timed out}$$

```
input integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm=
"giac")
```

```
output Timed out
```

3.379.9 Mupad [B] (verification not implemented)

Time = 14.41 (sec) , antiderivative size = 1492, normalized size of antiderivative = 5.87

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)
```

```
output atan((A^2*a^2*tan(c + d*x)^(1/2)*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4) - (A^2*a*b)/(2*d^2))^(1/2)*32i)/((16*A^3*b^3)/d + (16*A*a*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2))/d^3 - (16*A^3*a^2*b)/d) - (A^2*b^2*tan(c + d*x)^(1/2)*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4) - (A^2*a*b)/(2*d^2))^(1/2)*32i)/((16*A^3*b^3)/d + (16*A*a*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2))/d^3 - (16*A^3*a^2*b)/d)*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4) - (A^2*a*b)/(2*d^2))^(1/2)*2i - atan((A^2*a^2*tan(c + d*x)^(1/2)*(- (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4) - (A^2*a*b)/(2*d^2))^(1/2)*32i)/((16*A*a*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2))/d^3 - (16*A^3*b^3)/d + (16*A^3*a^2*b)/d) - (A^2*b^2*tan(c + d*x)^(1/2)*(- (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4) - (A^2*a*b)/(2*d^2))^(1/2)*32i)/((16*A*a*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2))/d^3 - (16*A^3*b^3)/d + (16*A^3*a^2*b)/d))*(- (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4) - (A^2*a*b)/(2*d^2))^(1/2)*2i - atan((B^2*a^2*tan(c + d*x)^(1/2)*((B^2*a*b)/(2*d^2) - (2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^(1/2)/(4*d^4))^(1/2)*32i)/((16*B^3*a^3)/d + (16*B*b*(2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^(1/2))/d^3 - (16*B^3*a*b^2)/d) - (B^2*b^2*tan(c + d*x)^(1/2)*((B^2*a*b)/(2*d^2) - (2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^(1/2)/(4*d^4))^(1/2)*...
```

3.380 $\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

3.380.1 Optimal result	3598
3.380.2 Mathematica [C] (verified)	3599
3.380.3 Rubi [A] (verified)	3599
3.380.4 Maple [A] (verified)	3604
3.380.5 Fricas [B] (verification not implemented)	3604
3.380.6 Sympy [F]	3605
3.380.7 Maxima [A] (verification not implemented)	3606
3.380.8 Giac [F(-1)]	3606
3.380.9 Mupad [B] (verification not implemented)	3607

3.380.1 Optimal result

Integrand size = 31, antiderivative size = 229

$$\begin{aligned} & \int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))(A + B \tan(c + dx)) dx \\ &= -\frac{(a(A - B) - b(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d} \\ & \quad + \frac{(a(A - B) - b(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d} \\ & \quad + \frac{(b(A - B) + a(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d} \\ & \quad - \frac{(b(A - B) + a(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d} \\ & \quad + \frac{2(Ab + aB)\sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)}{3d} \end{aligned}$$

```
output 1/2*(a*(A-B)-b*(A+B))*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)+1/2*(a
*(A-B)-b*(A+B))*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)+1/4*(b*(A-B)+
a*(A+B))*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)-1/4*(b*(A-B)+
a*(A+B))*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)+2*(A*b+B*a)*t
an(d*x+c)^(1/2)/d+2/3*b*B*tan(d*x+c)^(3/2)/d
```

3.380.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.45 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.50

$$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{3\sqrt[4]{-1}(ia+b)(A-iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right) - 3(-1)^{3/4}(a+ib)(A+iB) \operatorname{arctanh}\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{3d}$$

input `Integrate[Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(3*(-1)^(1/4)*(I*a + b)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] - 3*(-1)^(3/4)*(a + I*b)*(A + I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + 2*Sqrt[Tan[c + d*x]]*(3*A*b + 3*a*B + b*B*Tan[c + d*x]))/(3*d)`

3.380.3 Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 206, normalized size of antiderivative = 0.90, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 4075, 3042, 4011, 3042, 4017, 25, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4075}$$

$$\int \sqrt{\tan(c+dx)}(aA - bB + (Ab + aB) \tan(c+dx)) dx + \frac{2bB \tan^{\frac{3}{2}}(c+dx)}{3d}$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\tan(c+dx)}(aA - bB + (Ab + aB) \tan(c+dx)) dx + \frac{2bB \tan^{\frac{3}{2}}(c+dx)}{3d}$$

$$\downarrow \text{4011}$$

3.380. $\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \int \frac{-Ab - aB + (aA - bB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2(aB + Ab) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)}{3d} \\
& \quad \downarrow \text{3042} \\
& \int \frac{-Ab - aB + (aA - bB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2(aB + Ab) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)}{3d} \\
& \quad \downarrow \text{4017} \\
& \frac{2 \int -\frac{Ab+aB-(aA-bB)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} + \frac{2(aB + Ab) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)}{3d} \\
& \quad \downarrow \text{25} \\
& -\frac{2 \int \frac{Ab+aB-(aA-bB)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} + \frac{2(aB + Ab) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)}{3d} \\
& \quad \downarrow \text{1482} \\
& \frac{2\left(\frac{1}{2}(a(A - B) - b(A + B)) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(a(A + B) + b(A - B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c + dx)}\right)}{d} \\
& \quad + \frac{2(aB + Ab) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)}{3d} \\
& \quad \downarrow \text{1476} \\
& \frac{2\left(\frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c + dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c + dx)}\right)\right)}{d} \\
& \quad + \frac{2(aB + Ab) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)}{3d} \\
& \quad \downarrow \text{1082} \\
& \frac{2\left(\frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}}\right)\right)}{d} - \frac{1}{2}(a(A + B))}{d} \\
& \quad + \frac{2(aB + Ab) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)}{3d} \\
& \quad \downarrow \text{217}
\end{aligned}$$

$$2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a(A + B) + b(A - B)) \int \frac{1}{\tan} \right)$$

$$\frac{2(aB + Ab)\sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)}{3d}$$

↓ 1479

$$2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{1}{\tan} \right) \right)$$

$$\frac{2(aB + Ab)\sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)}{3d}$$

↓ 25

$$2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{1}{\tan} \right) \right)$$

$$\frac{2(aB + Ab)\sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)}{3d}$$

↓ 27

$$2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{1}{\tan} \right) \right)$$

$$\frac{2(aB + Ab)\sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)}{3d}$$

↓ 1103

$$2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{\log}{\tan} \right) \right)$$

$$\frac{2(aB + Ab)\sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)}{3d}$$

input `Int[Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`


```
output (2*(((a*(A - B) - b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 - ((b*(A - B) + a*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d + (2*(A*b + a*B)*Sqrt[Tan[c + d*x]]/d + (2*b*B*Tan[c + d*x]^(3/2))/(3*d))
```

3.380.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 217 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]
```

```
rule 1103 Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

```
rule 1476 Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.380.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{2Bb \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + 2Ab(\sqrt{\tan(dx+c)}) + 2Ba(\sqrt{\tan(dx+c)}) + \frac{(-Ab-Ba)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)} \right) + 2 \arctan \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)} \right) \right)}{4}$
default	$\frac{2Bb \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + 2Ab(\sqrt{\tan(dx+c)}) + 2Ba(\sqrt{\tan(dx+c)}) + \frac{(-Ab-Ba)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)} \right) + 2 \arctan \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)} \right) \right)}{4}$
parts	$\frac{(Ab+Ba) \left(2(\sqrt{\tan(dx+c)}) - \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4} \right)}{d}$

input `int(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*(2/3*B*b*tan(d*x+c)^(3/2)+2*A*b*tan(d*x+c)^(1/2)+2*B*a*tan(d*x+c)^(1/2)+1/4*(-A*b-B*a)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(A*a-B*b)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))`

3.380.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2230 vs. 2(195) = 390.

Time = 0.38 (sec) , antiderivative size = 2230, normalized size of antiderivative = 9.74

$$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))(A+B \tan(c+dx)) dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output

```

1/6*(3*d*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4
- 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 +
B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4)
)/d^2)*log(((A*a - B*b)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B
- A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a
b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) - ((A^2*B - B^3)*a^3 + (A^3 - 5*A*
B^2)*a^2*b - (5*A^2*B - B^3)*a*b^2 - (A^3 - A*B^2)*b^3)*d)*sqrt((2*A*B*a^2
- 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4
- 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B
- A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2) + ((A^4 - B^4)*a^
4 - 4*(A^3*B + A*B^3)*a^3*b - 4*(A^3*B + A*B^3)*a*b^3 - (A^4 - B^4)*b^4)*s
qrt(tan(d*x + c))) - 3*d*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b +
d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4
- 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2
+ B^4)*b^4)/d^4))/d^2)*log(-((A*a - B*b)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4
)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(
A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) - ((A^2*B - B^3)*
a^3 + (A^3 - 5*A*B^2)*a^2*b - (5*A^2*B - B^3)*a*b^2 - (A^3 - A*B^2)*b^3)*d
)*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^
2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)...

```

3.380.6 Sympy [F]

$$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \int (A+B \tan(c+dx))(a+b \tan(c+dx)) \sqrt{\tan(c+dx)} dx$$

input `integrate(tan(d*x+c)**(1/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))*sqrt(tan(c + d*x)), x)`

3.380.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.84

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))(A+B\tan(c+dx)) dx$$

$$= \frac{8Bb\tan(dx+c)^{\frac{3}{2}} + 6\sqrt{2}((A-B)a - (A+B)b)\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(dx+c)}\right)\right) + 6\sqrt{2}((A-B)a - (A+B)b)\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - 2\sqrt{\tan(dx+c)}\right)\right) + 3\sqrt{2}\left((A+B)a + (A-B)b\right)\log\left(\sqrt{2}\sqrt{\tan(dx+c)} + \tan(dx+c) + 1\right) + 3\sqrt{2}\left((A+B)a + (A-B)b\right)\log\left(-\sqrt{2}\sqrt{\tan(dx+c)} + \tan(dx+c) + 1\right) + 24(Ba + Ab)\sqrt{\tan(dx+c)}}{d}$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/12*(8*B*b*tan(d*x + c)^(3/2) + 6*sqrt(2)*((A - B)*a - (A + B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 6*sqrt(2)*((A - B)*a - (A + B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - 3*sqrt(2)*((A + B)*a + (A - B)*b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 3*sqrt(2)*((A + B)*a + (A - B)*b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 24*(B*a + A*b)*sqrt(tan(d*x + c)))/d`

3.380.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))(A+B\tan(c+dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.380.9 Mupad [B] (verification not implemented)

Time = 12.02 (sec) , antiderivative size = 1456, normalized size of antiderivative = 6.36

$$\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)
```

```
output 2*atanh((32*A^2*a^2*tan(c + d*x)^(1/2)*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 -
A^4*a^4*d^4)^(1/2)/(4*d^4) + (A^2*a*b)/(2*d^2))^(1/2))/((16*A*b*(2*A^4*a^
2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2))/d^3 - (16*A^3*a^3)/d + (16*A
^3*a*b^2)/d) - (32*A^2*b^2*tan(c + d*x)^(1/2)*((2*A^4*a^2*b^2*d^4 - A^4*b^
4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4) + (A^2*a*b)/(2*d^2))^(1/2))/((16*A*b*(2
*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2))/d^3 - (16*A^3*a^3)/d
+ (16*A^3*a*b^2)/d))*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2
)/(4*d^4) + (A^2*a*b)/(2*d^2))^(1/2) - 2*atanh((32*A^2*a^2*tan(c + d*x)^(1
/2)*((A^2*a*b)/(2*d^2) - (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(
1/2)/(4*d^4))^(1/2))/((16*A^3*a^3)/d + (16*A*b*(2*A^4*a^2*b^2*d^4 - A^4*b^
4*d^4 - A^4*a^4*d^4)^(1/2))/d^3 - (16*A^3*a*b^2)/d) - (32*A^2*b^2*tan(c +
d*x)^(1/2)*((A^2*a*b)/(2*d^2) - (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4
*d^4)^(1/2)/(4*d^4))^(1/2))/((16*A^3*a^3)/d + (16*A*b*(2*A^4*a^2*b^2*d^4 -
A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2))/d^3 - (16*A^3*a*b^2)/d))*((A^2*a*b)/(2*
d^2) - (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4))^(1/2
) - atan((B^2*a^2*tan(c + d*x)^(1/2)*(- (2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 -
B^4*a^4*d^4)^(1/2)/(4*d^4) - (B^2*a*b)/(2*d^2))^(1/2)*32i)/((16*B*a*(2*B^
4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^(1/2))/d^3 - (16*B^3*b^3)/d + (
16*B^3*a^2*b)/d) - (B^2*b^2*tan(c + d*x)^(1/2)*(- (2*B^4*a^2*b^2*d^4 - B^4
*b^4*d^4 - B^4*a^4*d^4)^(1/2)/(4*d^4) - (B^2*a*b)/(2*d^2))^(1/2)*32i)/(...
```

3.381
$$\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.381.1 Optimal result 3608
 3.381.2 Mathematica [C] (verified) 3609
 3.381.3 Rubi [A] (verified) 3609
 3.381.4 Maple [A] (verified) 3613
 3.381.5 Fricas [B] (verification not implemented) 3614
 3.381.6 Sympy [F] 3614
 3.381.7 Maxima [A] (verification not implemented) 3615
 3.381.8 Giac [F(-1)] 3615
 3.381.9 Mupad [B] (verification not implemented) 3616

3.381.1 Optimal result

Integrand size = 31, antiderivative size = 205

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= -\frac{(b(A - B) + a(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(b(A - B) + a(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(a(A - B) - b(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{(a(A - B) - b(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{2bB\sqrt{\tan(c + dx)}}{d}$$

output

```
1/2*(b*(A-B)+a*(A+B))*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)+1/2*(b
*(A-B)+a*(A+B))*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)-1/4*(a*(A-B)-
b*(A+B))*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)+1/4*(a*(A-B)-
b*(A+B))*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)+2*b*B*tan(d*x
+c)^(1/2)/d
```

3.381.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.46

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \frac{\sqrt[4]{-1}(a - ib)(A - iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right) + \sqrt[4]{-1}(a + ib)(A + iB) \operatorname{arctanh}\left((-1)^{3/4} \sqrt{\tan(c + dx)}\right)}{d}$$

input `Integrate[((a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `-(((-1)^(1/4)*(a - I*b)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + (-1)^(1/4)*(a + I*b)*(A + I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] - 2*b*B*Sqrt[Tan[c + d*x]])/d`

3.381.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.89, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 4075, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\ & \quad \downarrow \text{4075} \\ & \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2bB \sqrt{\tan(c + dx)}}{d} \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2bB \sqrt{\tan(c + dx)}}{d}$$

↓ 4017

$$\frac{2 \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)}}{d} + \frac{2bB \sqrt{\tan(c + dx)}}{d}$$

↓ 1482

$$\frac{2 \left(\frac{1}{2} (a(A - B) - b(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} + \frac{1}{2} (a(A + B) + b(A - B)) \int \frac{\tan(c + dx) + 1}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} \right)}{d} + \frac{2bB \sqrt{\tan(c + dx)}}{d}$$

↓ 1476

$$\frac{2 \left(\frac{1}{2} (a(A - B) - b(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} + \frac{1}{2} (a(A + B) + b(A - B)) \left(\frac{1}{2} \int \frac{1}{\tan(c + dx) - \sqrt{2} \sqrt{\tan(c + dx)}} \right) \right)}{d} + \frac{2bB \sqrt{\tan(c + dx)}}{d}$$

↓ 1082

$$\frac{2 \left(\frac{1}{2} (a(A - B) - b(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} + \frac{1}{2} (a(A + B) + b(A - B)) \left(\frac{\int \frac{1}{-\tan(c + dx) - 1} d(1 - \sqrt{2} \sqrt{\tan(c + dx)})}{\sqrt{2}} \right) \right)}{d} + \frac{2bB \sqrt{\tan(c + dx)}}{d}$$

↓ 217

$$\frac{2 \left(\frac{1}{2} (a(A - B) - b(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} + \frac{1}{2} (a(A + B) + b(A - B)) \left(\frac{\arctan(\sqrt{2} \sqrt{\tan(c + dx)} + 1)}{\sqrt{2}} \right) \right)}{d} + \frac{2bB \sqrt{\tan(c + dx)}}{d}$$

↓ 1479

3.381. $\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$

$$\frac{2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d}$$

$$\frac{2bB\sqrt{\tan(c+dx)}}{d}$$

↓ 25

$$\frac{2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d}$$

$$\frac{2bB\sqrt{\tan(c+dx)}}{d}$$

↓ 27

$$\frac{2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d}$$

$$\frac{2bB\sqrt{\tan(c+dx)}}{d}$$

↓ 1103

$$\frac{2 \left(\frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\log\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)+1}}{1+\sqrt{2}\sqrt{\tan(c+dx)+1}}\right)}{\sqrt{2}} + \frac{\log\left(\frac{1+\sqrt{2}\sqrt{\tan(c+dx)}}{1-\sqrt{2}\sqrt{\tan(c+dx)}}\right)}{\sqrt{2}} \right) \right)}{d}$$

$$\frac{2bB\sqrt{\tan(c+dx)}}{d}$$

input `Int[((a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `(2*(((b*(A - B) + a*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 + ((a*(A - B) - b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d + (2*b*B*Sqrt[Tan[c + d*x]])/d`

3.381.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)], x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.381.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.99

method	result
derivativedivides	$\frac{2(\sqrt{\tan(dx+c)}Bb + \frac{(aA-Bb)\sqrt{2}}{4} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2\arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2\arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4}$
default	$\frac{2(\sqrt{\tan(dx+c)}Bb + \frac{(aA-Bb)\sqrt{2}}{4} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2\arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2\arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4}$
parts	$\frac{(Ab+Ba)\sqrt{2} \left(\ln\left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2\arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2\arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4d}$

input `int((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x,method=_RETURNVERBOSE)`

output `1/d*(2*tan(d*x+c)^(1/2)*B*b+1/4*(A*a-B*b)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(A*b+B*a)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))`

$$3.381. \int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.381.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2216 vs. $2(175) = 350$.

Time = 0.38 (sec) , antiderivative size = 2216, normalized size of antiderivative = 10.81

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm=
"fricas")
```

```
output 1/2*(d*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-(A^4
- 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B
^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))
/d^2)*log((B*a + A*b)*d^3*sqrt(-(A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B -
A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b
^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) + ((A^3 - A*B^2)*a^3 - (5*A^2*B - B
^3)*a^2*b - (A^3 - 5*A*B^2)*a*b^2 + (A^2*B - B^3)*b^3)*d)*sqrt(-(2*A*B*a^2
- 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-(A^4 - 2*A^2*B^2 + B^4)*a^4
- 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B
- A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2) + ((A^4 - B^4)*a^
4 - 4*(A^3*B + A*B^3)*a^3*b - 4*(A^3*B + A*B^3)*a*b^3 - (A^4 - B^4)*b^4)*s
qrt(tan(d*x + c))) - d*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b +
d^2*sqrt(-(A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4
- 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 +
B^4)*b^4)/d^4))/d^2)*log(-(B*a + A*b)*d^3*sqrt(-(A^4 - 2*A^2*B^2 + B^4)
*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A
^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) + ((A^3 - A*B^2)*a
^3 - (5*A^2*B - B^3)*a^2*b - (A^3 - 5*A*B^2)*a*b^2 + (A^2*B - B^3)*b^3)*d)
*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-(A^4 - 2*A^
2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)...
```

3.381.6 Sympy [F]

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\ &= \int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))/sqrt(tan(c + d*x)), x)`

3.381.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.85

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \frac{2\sqrt{2}((A + B)a + (A - B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(dx + c)}\right)\right) + 2\sqrt{2}((A + B)a + (A - B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - 2\sqrt{\tan(dx + c)}\right)\right) + \sqrt{2}((A - B)a - (A + B)b) \log\left(\sqrt{2}\sqrt{\tan(dx + c)} + \tan(dx + c) + 1\right) - \sqrt{2}((A - B)a - (A + B)b) \log\left(-\sqrt{2}\sqrt{\tan(dx + c)} + \tan(dx + c) + 1\right) + 8Bb\sqrt{\tan(dx + c)}}{d}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="maxima")`

output `1/4*(2*sqrt(2)*((A + B)*a + (A - B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A + B)*a + (A - B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((A - B)*a - (A + B)*b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((A - B)*a - (A + B)*b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 8*B*b*sqrt(tan(d*x + c)))/d`

3.381.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="giac")`

output `Timed out`

3.381. $\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

3.381.9 Mupad [B] (verification not implemented)

Time = 10.38 (sec) , antiderivative size = 1420, normalized size of antiderivative = 6.93

$$\begin{aligned}
& \int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
&= 2 \operatorname{atanh} \left(\frac{32 A^2 a^2 \sqrt{\tan(c + dx)} \sqrt{\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4}} - \frac{A^2 a b}{2 d^2}}{\frac{16 A^3 b^3}{d} + \frac{16 A a \sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{d^3} - \frac{16 A^3 a^2 b}{d}} \right) \\
&\quad - \frac{32 A^2 b^2 \sqrt{\tan(c + dx)} \sqrt{\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4}} - \frac{A^2 a b}{2 d^2}}{\frac{16 A^3 b^3}{d} + \frac{16 A a \sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{d^3} - \frac{16 A^3 a^2 b}{d}} \sqrt{\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4}} \\
&\quad - 2 \operatorname{atanh} \left(\frac{32 A^2 a^2 \sqrt{\tan(c + dx)} \sqrt{-\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4}} - \frac{A^2 a b}{2 d^2}}{\frac{16 A a \sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{d^3} - \frac{16 A^3 b^3}{d} + \frac{16 A^3 a^2 b}{d}} \right) \\
&\quad - \frac{32 A^2 b^2 \sqrt{\tan(c + dx)} \sqrt{-\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4}} - \frac{A^2 a b}{2 d^2}}{\frac{16 A a \sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{d^3} - \frac{16 A^3 b^3}{d} + \frac{16 A^3 a^2 b}{d}} \sqrt{-\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4}} \\
&\quad - 2 \operatorname{atanh} \left(\frac{32 B^2 a^2 \sqrt{\tan(c + dx)} \sqrt{\frac{B^2 a b}{2 d^2} - \frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4}}}{\frac{16 B^3 a^3}{d} + \frac{16 B b \sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{d^3} - \frac{16 B^3 a b^2}{d}} \right) \\
&\quad - \frac{32 B^2 b^2 \sqrt{\tan(c + dx)} \sqrt{\frac{B^2 a b}{2 d^2} - \frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4}}}{\frac{16 B^3 a^3}{d} + \frac{16 B b \sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{d^3} - \frac{16 B^3 a b^2}{d}} \sqrt{\frac{B^2 a b}{2 d^2} - \frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4}} \\
&\quad + 2 \operatorname{atanh} \left(\frac{32 B^2 a^2 \sqrt{\tan(c + dx)} \sqrt{\frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4}} + \frac{B^2 a b}{2 d^2}}{\frac{16 B b \sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{d^3} - \frac{16 B^3 a^3}{d} + \frac{16 B^3 a b^2}{d}} \right) \\
&\quad - \frac{32 B^2 b^2 \sqrt{\tan(c + dx)} \sqrt{\frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4}} + \frac{B^2 a b}{2 d^2}}{\frac{16 B b \sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{d^3} - \frac{16 B^3 a^3}{d} + \frac{16 B^3 a b^2}{d}} \sqrt{\frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4}} \\
&\quad + \frac{2 B b \sqrt{\tan(c + dx)}}{d}
\end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x)))/tan(c + d*x)^(1/2),x)`

output

$$\begin{aligned}
& 2*\operatorname{atanh}((32*A^2*a^2*\tan(c + d*x))^{1/2}*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - \\
& A^4*a^4*d^4)^{1/2}/(4*d^4) - (A^2*a*b)/(2*d^2))^{1/2})/((16*A^3*b^3)/d + \\
& (16*A*a*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{1/2})/d^3 - (16*A \\
& ^3*a^2*b)/d) - (32*A^2*b^2*\tan(c + d*x))^{1/2}*((2*A^4*a^2*b^2*d^4 - A^4*b^ \\
& 4*d^4 - A^4*a^4*d^4)^{1/2}/(4*d^4) - (A^2*a*b)/(2*d^2))^{1/2})/((16*A^3*b^ \\
& 3)/d + (16*A*a*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{1/2})/d^3 \\
& - (16*A^3*a^2*b)/d))*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{1/2} \\
&)/(4*d^4) - (A^2*a*b)/(2*d^2))^{1/2} - 2*\operatorname{atanh}((32*A^2*a^2*\tan(c + d*x))^{1} \\
& /2)*(- (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{1/2}/(4*d^4) - (A^ \\
& 2*a*b)/(2*d^2))^{1/2})/((16*A*a*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4 \\
& *d^4)^{1/2})/d^3 - (16*A^3*b^3)/d + (16*A^3*a^2*b)/d) - (32*A^2*b^2*\tan(c \\
& + d*x))^{1/2}*(- (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{1/2}/(4*d \\
& ^4) - (A^2*a*b)/(2*d^2))^{1/2})/((16*A*a*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 \\
& - A^4*a^4*d^4)^{1/2})/d^3 - (16*A^3*b^3)/d + (16*A^3*a^2*b)/d))*(- (2*A^4* \\
& a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{1/2}/(4*d^4) - (A^2*a*b)/(2*d^2) \\
&)^{1/2} - 2*\operatorname{atanh}((32*B^2*a^2*\tan(c + d*x))^{1/2}*((B^2*a*b)/(2*d^2) - (2*B \\
& ^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^{1/2}/(4*d^4))^{1/2})/((16*B^3 \\
& *a^3)/d + (16*B*b*(2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^{1/2})/d \\
& ^3 - (16*B^3*a*b^2)/d) - (32*B^2*b^2*\tan(c + d*x))^{1/2}*((B^2*a*b)/(2*d^2) \\
& - (2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^{1/2}/(4*d^4))^{1/2})...
\end{aligned}$$

3.382
$$\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.382.1 Optimal result 3618
 3.382.2 Mathematica [A] (verified) 3619
 3.382.3 Rubi [A] (verified) 3619
 3.382.4 Maple [A] (verified) 3623
 3.382.5 Fricas [B] (verification not implemented) 3624
 3.382.6 Sympy [F] 3625
 3.382.7 Maxima [A] (verification not implemented) 3625
 3.382.8 Giac [F(-1)] 3626
 3.382.9 Mupad [B] (verification not implemented) 3627

3.382.1 Optimal result

Integrand size = 31, antiderivative size = 205

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{(a(A - B) - b(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(a(A - B) - b(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(b(A - B) + a(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{(b(A - B) + a(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d} - \frac{2aA}{d\sqrt{\tan(c + dx)}}$$

output

```
-1/2*(a*(A-B)-b*(A+B))*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)-1/2*(
a*(A-B)-b*(A+B))*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)-1/4*(b*(A-B)
+a*(A+B))*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)+1/4*(b*(A-B)
+a*(A+B))*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)-2*a*A/d/tan(
d*x+c)^(1/2)
```

3.382.2 Mathematica [A] (verified)

Time = 0.68 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.77

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{-2\sqrt{2}(a(A - B) - b(A + B)) \left(\arctan \left(1 - \sqrt{2}\sqrt{\tan(c + dx)} \right) - \arctan \left(1 + \sqrt{2}\sqrt{\tan(c + dx)} \right) \right) + \sqrt{2}(b(A - B) + a(A + B)) \left(\log \left(1 - \sqrt{2}\sqrt{\tan(c + dx)} \right) + \tan(c + dx) - \log \left(1 + \sqrt{2}\sqrt{\tan(c + dx)} \right) + \tan(c + dx) \right) + (8aA)/\sqrt{\tan(c + dx)}}{d}$$

input `Integrate[((a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `-1/4*(-2*Sqrt[2]*(a*(A - B) - b*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]) + Sqrt[2]*(b*(A - B) + a*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]) + (8*a*A)/Sqrt[Tan[c + d*x]]/d`

3.382.3 Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.89, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.387$, Rules used = {3042, 4074, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx$$

$$\downarrow \text{4074}$$

$$\int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2aA}{d\sqrt{\tan(c + dx)}}$$

$$\downarrow \text{3042}$$

$$\int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2aA}{d\sqrt{\tan(c + dx)}}$$

3.382. $\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

$$\frac{2 \int \frac{Ab+aB-(aA-bB) \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} - \frac{2aA}{d\sqrt{\tan(c+dx)}}$$

↓ 4017

$$\frac{2\left(\frac{1}{2}(a(A+B)+b(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a(A-B)-b(A+B)) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right)}{d} - \frac{2aA}{d\sqrt{\tan(c+dx)}}$$

↓ 1482

$$\frac{2\left(\frac{1}{2}(a(A+B)+b(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a(A-B)-b(A+B)) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}}\right)\right)}{d} - \frac{2aA}{d\sqrt{\tan(c+dx)}}$$

↓ 1476

$$\frac{2\left(\frac{1}{2}(a(A+B)+b(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a(A-B)-b(A+B)) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}}\right)\right)}{d} - \frac{2aA}{d\sqrt{\tan(c+dx)}}$$

↓ 1082

$$\frac{2\left(\frac{1}{2}(a(A+B)+b(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a(A-B)-b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}}\right)\right)}{d} - \frac{2aA}{d\sqrt{\tan(c+dx)}}$$

↓ 217

$$\frac{2\left(\frac{1}{2}(a(A+B)+b(A-B)) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}}\right)\right)}{d} - \frac{2aA}{d\sqrt{\tan(c+dx)}}$$

3.382. $\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

↓ 25

$$\frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2}(a(A-B) - b(A+B)) \right)}{d} - \frac{2aA}{d\sqrt{\tan(c+dx)}}$$

↓ 27

$$\frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2}(a(A-B) - b(A+B)) \right)}{d} - \frac{2aA}{d\sqrt{\tan(c+dx)}}$$

↓ 1103

$$\frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(a(A-B) - b(A+B)) \right)}{d} - \frac{2aA}{d\sqrt{\tan(c+dx)}}$$

input `Int[((a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `(2*(-1/2*((a*(A - B) - b*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])) + ((b*(A - B) + a*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]))))/2))/d - (2*a*A)/(d*Sqrt[Tan[c + d*x]])`

3.382.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4074 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.382.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.99

method	result
derivativedivides	$-\frac{2aA}{\sqrt{\tan(dx+c)}} + \frac{(Ab+Ba)\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4}$
default	$-\frac{2aA}{\sqrt{\tan(dx+c)}} + \frac{(Ab+Ba)\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4}$
parts	$\frac{(Ab+Ba)\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4d}$

input `int((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x,method=_RETURNVERBOSE)`

$$3.382. \int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

output $1/d*(-2*a*A/\tan(dx+c)^{(1/2)}+1/4*(A*b+B*a)*2^{(1/2)}*(\ln((1+2^{(1/2)}*\tan(dx+c)^{(1/2)}+\tan(dx+c))/(-1-2^{(1/2)}*\tan(dx+c)^{(1/2)}+\tan(dx+c))))+2*\arctan(1+2^{(1/2)}*\tan(dx+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(dx+c)^{(1/2)}))+1/4*(-A*a+B*b)*2^{(1/2)}*(\ln((1-2^{(1/2)}*\tan(dx+c)^{(1/2)}+\tan(dx+c))/(1+2^{(1/2)}*\tan(dx+c)^{(1/2)}+\tan(dx+c))))+2*\arctan(1+2^{(1/2)}*\tan(dx+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(dx+c)^{(1/2)}))$

3.382.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2244 vs. 2(175) = 350.

Time = 0.39 (sec), antiderivative size = 2244, normalized size of antiderivative = 10.95

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="fracas")`

output $-1/2*(d*\sqrt{(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*\sqrt{-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4})/d^2)*\log(((A*a - B*b)*d^3*\sqrt{-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4}) - ((A^2*B - B^3)*a^3 + (A^3 - 5*A*B^2)*a^2*b - (5*A^2*B - B^3)*a*b^2 - (A^3 - A*B^2)*b^3)*d)*\sqrt{(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*\sqrt{-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4})/d^2) + ((A^4 - B^4)*a^4 - 4*(A^3*B + A*B^3)*a^3*b - 4*(A^3*B + A*B^3)*a*b^3 - (A^4 - B^4)*b^4)*\sqrt{\tan(dx + c)} - d*\sqrt{(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*\sqrt{-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4})/d^2)*\log(-((A*a - B*b)*d^3*\sqrt{-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4}) - ((A^2*B - B^3)*a^3 + (A^3 - 5*A*B^2)*a^2*b - (5*A^2*B - B^3)*a*b^2 - (A^3 - A*B^2)*b^3)*d)*\sqrt{(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*\sqrt{-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4})/d^2)$

3.382.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))/tan(c + d*x)**(3/2), x)`

3.382.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.85

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{2\sqrt{2}((A - B)a - (A + B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(dx + c)}\right)\right) + 2\sqrt{2}((A - B)a - (A + B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - 2\sqrt{\tan(dx + c)}\right)\right) - \sqrt{2}((A + B)a + (A - B)b) \log(\sqrt{2}\sqrt{\tan(dx + c)} + \tan(dx + c) + 1) + \sqrt{2}((A + B)a + (A - B)b) \log(-\sqrt{2}\sqrt{\tan(dx + c)} + \tan(dx + c) + 1) + 8Aa/\sqrt{\tan(dx + c)}}{d}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="maxima")`

output `-1/4*(2*sqrt(2)*((A - B)*a - (A + B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a - (A + B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*((A + B)*a + (A - B)*b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*((A + B)*a + (A - B)*b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 8*A*a/sqrt(tan(d*x + c)))/d`

3.382.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="giac")`

output `Timed out`

3.382.9 Mupad [B] (verification not implemented)

Time = 10.79 (sec) , antiderivative size = 1420, normalized size of antiderivative = 6.93

$$\begin{aligned}
& \int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\
&= 2 \operatorname{atanh} \left(\frac{32 A^2 a^2 d^3 \sqrt{\tan(c + dx)} \sqrt{\frac{A^2 a b}{2 d^2} - \frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4}}}{16 A b \sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4} + 16 A^3 a^3 d^2 - 16 A^3 a b^2 d^2} \right. \\
&\quad \left. - \frac{32 A^2 b^2 d^3 \sqrt{\tan(c + dx)} \sqrt{\frac{A^2 a b}{2 d^2} - \frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4}}}{16 A b \sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4} + 16 A^3 a^3 d^2 - 16 A^3 a b^2 d^2} \right) \sqrt{\frac{A^2 a b}{2 d^2} - \frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4}} \\
&\quad - 2 \operatorname{atanh} \left(\frac{32 A^2 a^2 d^3 \sqrt{\tan(c + dx)} \sqrt{\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4} + \frac{A^2 a b}{2 d^2}}}{16 A b \sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4} - 16 A^3 a^3 d^2 + 16 A^3 a b^2 d^2} \right. \\
&\quad \left. - \frac{32 A^2 b^2 d^3 \sqrt{\tan(c + dx)} \sqrt{\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4} + \frac{A^2 a b}{2 d^2}}}{16 A b \sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4} - 16 A^3 a^3 d^2 + 16 A^3 a b^2 d^2} \right) \sqrt{\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4}} \\
&\quad - 2 \operatorname{atanh} \left(\frac{32 B^2 a^2 \sqrt{\tan(c + dx)} \sqrt{-\frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4} - \frac{B^2 a b}{2 d^2}}}{\frac{16 B a \sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{d^3} - \frac{16 B^3 b^3}{d} + \frac{16 B^3 a^2 b}{d}} \right. \\
&\quad \left. - \frac{32 B^2 b^2 \sqrt{\tan(c + dx)} \sqrt{-\frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4} - \frac{B^2 a b}{2 d^2}}}{\frac{16 B a \sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{d^3} - \frac{16 B^3 b^3}{d} + \frac{16 B^3 a^2 b}{d}} \right) \sqrt{-\frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4}} \\
&\quad + 2 \operatorname{atanh} \left(\frac{32 B^2 a^2 \sqrt{\tan(c + dx)} \sqrt{\frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4} - \frac{B^2 a b}{2 d^2}}}{\frac{16 B^3 b^3}{d} + \frac{16 B a \sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{d^3} - \frac{16 B^3 a^2 b}{d}} \right. \\
&\quad \left. - \frac{32 B^2 b^2 \sqrt{\tan(c + dx)} \sqrt{\frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4} - \frac{B^2 a b}{2 d^2}}}{\frac{16 B^3 b^3}{d} + \frac{16 B a \sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{d^3} - \frac{16 B^3 a^2 b}{d}} \right) \sqrt{\frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4}} \\
&\quad - \frac{2 A a}{d \sqrt{\tan(c + dx)}}
\end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x)))/tan(c + d*x)^(3/2),x)`

output

$$\begin{aligned}
& 2*\operatorname{atanh}((32*A^2*a^2*d^3*\tan(c + d*x))^{(1/2)}*((A^2*a*b)/(2*d^2) - (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)})/(4*d^4))^{(1/2)})/(16*A*b*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)} + 16*A^3*a^3*d^2 - 16*A^3*a*b^2*d^2) - (32*A^2*b^2*d^3*\tan(c + d*x))^{(1/2)}*((A^2*a*b)/(2*d^2) - (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)})/(4*d^4))^{(1/2)})/(16*A*b*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)} + 16*A^3*a^3*d^2 - 16*A^3*a*b^2*d^2))*((A^2*a*b)/(2*d^2) - (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)})/(4*d^4))^{(1/2)} - 2*\operatorname{atanh}((32*A^2*a^2*d^3*\tan(c + d*x))^{(1/2)}*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)})/(4*d^4) + (A^2*a*b)/(2*d^2))^{(1/2)})/(16*A*b*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)} - 16*A^3*a^3*d^2 + 16*A^3*a*b^2*d^2) - (32*A^2*b^2*d^3*\tan(c + d*x))^{(1/2)}*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)})/(4*d^4) + (A^2*a*b)/(2*d^2))^{(1/2)})/(16*A*b*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)} - 16*A^3*a^3*d^2 + 16*A^3*a*b^2*d^2))*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)})/(4*d^4) + (A^2*a*b)/(2*d^2))^{(1/2)} - 2*\operatorname{atanh}((32*B^2*a^2*\tan(c + d*x))^{(1/2)}*(- (2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^{(1/2)})/(4*d^4) - (B^2*a*b)/(2*d^2))^{(1/2)})/((16*B*a*(2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^{(1/2)})/d^3 - (16*B^3*b^3)/d + (16*B^3*a^2*b)/d) - (32*B^2*b^2*\tan(c + d*x))^{(1/2)}*(- (2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^{(1/2)})/(4*d^4) - (B^2*a*b)/(2*d^2))^{(1/2)})/((16*B*a...
\end{aligned}$$

3.383
$$\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.383.1 Optimal result 3629
 3.383.2 Mathematica [A] (verified) 3630
 3.383.3 Rubi [A] (verified) 3630
 3.383.4 Maple [A] (verified) 3635
 3.383.5 Fricas [B] (verification not implemented) 3635
 3.383.6 Sympy [F] 3636
 3.383.7 Maxima [A] (verification not implemented) 3637
 3.383.8 Giac [F(-1)] 3637
 3.383.9 Mupad [B] (verification not implemented) 3638

3.383.1 Optimal result

Integrand size = 31, antiderivative size = 229

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{(b(A - B) + a(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(b(A - B) + a(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(a(A - B) - b(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(a(A - B) - b(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$- \frac{2aA}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2(Ab + aB)}{d\sqrt{\tan(c + dx)}}$$

output

```
-1/2*(b*(A-B)+a*(A+B))*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)-1/2*(
b*(A-B)+a*(A+B))*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)+1/4*(a*(A-B)
-b*(A+B))*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)-1/4*(a*(A-B)
-b*(A+B))*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)-2*(A*b+B*a)/
d/tan(d*x+c)^(1/2)-2/3*a*A/d/tan(d*x+c)^(3/2)
```

3.383.2 Mathematica [A] (verified)

Time = 0.83 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.78

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{6\sqrt{2}(b(A - B) + a(A + B)) \left(\arctan \left(1 - \sqrt{2}\sqrt{\tan(c + dx)} \right) - \arctan \left(1 + \sqrt{2}\sqrt{\tan(c + dx)} \right) \right) + 3\sqrt{2}}$$

input `Integrate[((a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(6*Sqrt[2]*(b*(A - B) + a*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]) + 3*Sqrt[2]*(a*(A - B) - b*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]) - (8*a*A)/Tan[c + d*x]^(3/2) - (24*(A*b + a*B))/Sqrt[Tan[c + d*x]])/(12*d)`

3.383.3 Rubi [A] (verified)Time = 0.65 (sec) , antiderivative size = 206, normalized size of antiderivative = 0.90, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 4074, 3042, 4012, 25, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx$$

$$\downarrow \text{4074}$$

$$\int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)} dx - \frac{2aA}{3d \tan^{\frac{3}{2}}(c + dx)}$$

$$\downarrow \text{3042}$$

3.383. $\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\tan(c + dx)^{3/2}} dx - \frac{2aA}{3d \tan^{3/2}(c + dx)} \\
& \quad \downarrow 4012 \\
& \int -\frac{aA - bB + (Ab + aB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2(aB + Ab)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{3d \tan^{3/2}(c + dx)} \\
& \quad \downarrow 25 \\
& - \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2(aB + Ab)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{3d \tan^{3/2}(c + dx)} \\
& \quad \downarrow 3042 \\
& - \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2(aB + Ab)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{3d \tan^{3/2}(c + dx)} \\
& \quad \downarrow 4017 \\
& - \frac{2 \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)}}{d} - \frac{2(aB + Ab)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{3d \tan^{3/2}(c + dx)} \\
& \quad \downarrow 1482 \\
& - \frac{2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} + \frac{1}{2}(a(A + B) + b(A - B)) \int \frac{\tan(c + dx) + 1}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} \right)}{d} \\
& \quad - \frac{2(aB + Ab)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{3d \tan^{3/2}(c + dx)} \\
& \quad \downarrow 1476 \\
& - \frac{2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} + \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{1}{2} \int \frac{1}{\tan(c + dx) - \sqrt{2} \sqrt{\tan(c + dx)}} dx \right) \right)}{d} \\
& \quad - \frac{2(aB + Ab)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{3d \tan^{3/2}(c + dx)} \\
& \quad \downarrow 1082 \\
& - \frac{2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} + \frac{1}{2}(a(A + B) + b(A - B)) \left(\int \frac{1}{-\tan(c + dx) - 1} d \left(\frac{1 - \sqrt{2} \sqrt{\tan(c + dx)}}{\sqrt{2}} \right) \right) \right)}{d} \\
& \quad - \frac{2(aB + Ab)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{3d \tan^{3/2}(c + dx)} \\
& \quad \downarrow 217 \\
& \frac{2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} + \frac{1}{2}(a(A + B) + b(A - B)) \left(\int \frac{1}{-\tan(c + dx) - 1} d \left(\frac{1 - \sqrt{2} \sqrt{\tan(c + dx)}}{\sqrt{2}} \right) \right) \right)}{d} \\
& \quad - \frac{2(aB + Ab)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{3d \tan^{3/2}(c + dx)} \\
& \quad \downarrow 217
\end{aligned}$$

3.383. $\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx$

$$2 \left(\frac{1}{2}(a(A-B) - b(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)$$

$$\frac{2(aB + Ab)}{d\sqrt{\tan(c+dx)}} - \frac{2aA}{3d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 1479

$$2 \left(\frac{1}{2}(a(A-B) - b(A+B)) \left(- \frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \right)$$

$$\frac{2(aB + Ab)}{d\sqrt{\tan(c+dx)}} - \frac{2aA}{3d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 25

$$2 \left(\frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(a \right)$$

$$\frac{2(aB + Ab)}{d\sqrt{\tan(c+dx)}} - \frac{2aA}{3d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 27

$$2 \left(\frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)$$

$$\frac{2(aB + Ab)}{d\sqrt{\tan(c+dx)}} - \frac{2aA}{3d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 1103

$$2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{1}{\sqrt{2}} \right) \right)$$

$$\frac{2(aB + Ab)}{d\sqrt{\tan(c+dx)}} - \frac{2aA}{3d \tan^{\frac{3}{2}}(c+dx)}$$

input `Int[((a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

```
output (-2*((b*(A - B) + a*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 + ((a*(A - B) - b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d - (2*a*A)/(3*d*Tan[c + d*x]^(3/2)) - (2*(A*b + a*B))/(d*Sqrt[Tan[c + d*x]])
```

3.383.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]
```

```
rule 1103 Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

```
rule 1476 Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```


rule 1479 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2(d/e), 2]\}, \text{Simp}[e/(2cq) \text{Int}[(q - 2x)/\text{Simp}[d/e + qx - x^2, x], x], x] + \text{Simp}[e/(2cq) \text{Int}[(q + 2x)/\text{Simp}[d/e - qx - x^2, x], x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[cd^2 - ae^2, 0] \&\& \text{NegQ}[d^2e]$

rule 1482 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[ac, 2]\}, \text{Simp}[(dq + ae)/(2ac) \text{Int}[(q + cx^2)/(a + cx^4), x], x] + \text{Simp}[(dq - ae)/(2ac) \text{Int}[(q - cx^2)/(a + cx^4), x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{NeQ}[cd^2 + ae^2, 0] \&\& \text{NeQ}[cd^2 - ae^2, 0] \&\& \text{NegQ}[-a]c]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4012 $\text{Int}[\frac{(a_.) + (b_.)\tan[(e_.) + (f_.)x]^m}{(c_.) + (d_.)\tan[(e_.) + (f_.)x]}, x_Symbol] \rightarrow \text{Simp}[(bc - ad)(a + b\tan[e + fx])^{m+1}/(f(m+1)(a^2 + b^2)), x] + \text{Simp}[1/(a^2 + b^2) \text{Int}[(a + b\tan[e + fx])^{m+1} \text{Simp}[ac + bd - (bc - ad)\tan[e + fx], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[bc - ad, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{LtQ}[m, -1]$

rule 4017 $\text{Int}[\frac{(c_.) + (d_.)\tan[(e_.) + (f_.)x]}{\sqrt{(b_.)\tan[(e_.) + (f_.)x]}}, x_Symbol] \rightarrow \text{Simp}[2/f \text{Subst}[\text{Int}[(bc + dx^2)/(b^2 + x^4), x], x, \text{Sqrt}[b\tan[e + fx]]], x] /; \text{FreeQ}[\{b, c, d, e, f\}, x] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0]$

rule 4074 $\text{Int}[\frac{(a_.) + (b_.)\tan[(e_.) + (f_.)x]^m}{(A_.) + (B_.)\tan[(e_.) + (f_.)x]}, x_Symbol] \rightarrow \text{Simp}[(bc - ad)(A^2 - B^2)(a + b\tan[e + fx])^{m+1}/(b^2f(m+1)(a^2 + b^2)), x] + \text{Simp}[1/(a^2 + b^2) \text{Int}[(a + b\tan[e + fx])^{m+1} \text{Simp}[aAc + bBc + Abd - aBd - (Abc - aBc - aAd - bBd)\tan[e + fx], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \&\& \text{NeQ}[bc - ad, 0] \&\& \text{LtQ}[m, -1] \&\& \text{NeQ}[a^2 + b^2, 0]$

3.383.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 222, normalized size of antiderivative = 0.97

method	result
derivativedivides	$-\frac{2(Ab+Ba)}{\sqrt{\tan(dx+c)}} - \frac{2aA}{3 \tan(dx+c)^{\frac{3}{2}}} + \frac{(-aA+Bb)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(1-\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4}$
default	$-\frac{2(Ab+Ba)}{\sqrt{\tan(dx+c)}} - \frac{2aA}{3 \tan(dx+c)^{\frac{3}{2}}} + \frac{(-aA+Bb)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(1-\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4}$
parts	$\frac{(Ab+Ba) \left(-\frac{\sqrt{2} \left(\ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4} - \frac{2aA}{3 \tan(dx+c)^{\frac{3}{2}}} \right)}{d}$

input `int((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x,method=_RETURNVERBOSE)`

output `1/d*(-2*(A*b+B*a)/tan(d*x+c)^(1/2)-2/3*a*A/tan(d*x+c)^(3/2)+1/4*(-A*a+B*b)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(-A*b-B*a)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))`

3.383.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2275 vs. 2(195) = 390.

Time = 0.41 (sec) , antiderivative size = 2275, normalized size of antiderivative = 9.93

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="fracas")`

output

```
-1/6*(3*d*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2)*log(((B*a + A*b)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) + ((A^3 - A*B^2)*a^3 - (5*A^2*B - B^3)*a^2*b - (A^3 - 5*A*B^2)*a*b^2 + (A^2*B - B^3)*b^3)*d)*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2) + ((A^4 - B^4)*a^4 - 4*(A^3*B + A*B^3)*a^3*b - 4*(A^3*B + A*B^3)*a*b^3 - (A^4 - B^4)*b^4)*sqrt(tan(d*x + c))*tan(d*x + c)^2 - 3*d*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2)*log(-((B*a + A*b)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) + ((A^3 - A*B^2)*a^3 - (5*A^2*B - B^3)*a^2*b - (A^3 - 5*A*B^2)*a*b^2 + (A^2*B - B^3)*b^3)*d)*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4...
```

3.383.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)**(5/2), x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))/tan(c + d*x)**(5/2), x)`

3.383.7 Maxima [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.84

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx =$$

$$6\sqrt{2}((A + B)a + (A - B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(dx + c)}\right)\right) + 6\sqrt{2}((A + B)a + (A - B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - 2\sqrt{\tan(dx + c)}\right)\right) + 3\sqrt{2}((A - B)a - (A + B)b) \log(\sqrt{2}\sqrt{\tan(dx + c)} + \tan(dx + c) + 1) - 3\sqrt{2}((A - B)a - (A + B)b) \log(-\sqrt{2}\sqrt{\tan(dx + c)} + \tan(dx + c) + 1) + 8(Aa + 3(Ba + Ab)\tan(dx + c))/\tan(dx + c)^{\frac{3}{2}}/d$$

```
input integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="maxima")
```

```
output -1/12*(6*sqrt(2)*((A + B)*a + (A - B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 6*sqrt(2)*((A + B)*a + (A - B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + 3*sqrt(2)*((A - B)*a - (A + B)*b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - 3*sqrt(2)*((A - B)*a - (A + B)*b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 8*(A*a + 3*(B*a + A*b)*tan(d*x + c))/tan(d*x + c)^(3/2))/d
```

3.383.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="giac")
```

```
output Timed out
```

3.383.9 Mupad [B] (verification not implemented)

Time = 11.75 (sec) , antiderivative size = 1448, normalized size of antiderivative = 6.32

$$\begin{aligned}
& \int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \\
&= 2 \operatorname{atanh} \left(\frac{32 A^2 a^2 d^3 \sqrt{\tan(c + dx)} \sqrt{-\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4} - \frac{A^2 a b}{2 d^2}}}{16 A a \sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4} - 16 A^3 b^3 d^2 + 16 A^3 a^2 b d^2} \right. \\
&\quad \left. - \frac{32 A^2 b^2 d^3 \sqrt{\tan(c + dx)} \sqrt{-\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4} - \frac{A^2 a b}{2 d^2}}}{16 A a \sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4} - 16 A^3 b^3 d^2 + 16 A^3 a^2 b d^2} \right) \sqrt{-\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4}} \\
&\quad - 2 \operatorname{atanh} \left(\frac{32 A^2 a^2 d^3 \sqrt{\tan(c + dx)} \sqrt{\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4} - \frac{A^2 a b}{2 d^2}}}{16 A a \sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4} + 16 A^3 b^3 d^2 - 16 A^3 a^2 b d^2} \right. \\
&\quad \left. - \frac{32 A^2 b^2 d^3 \sqrt{\tan(c + dx)} \sqrt{\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4} - \frac{A^2 a b}{2 d^2}}}{16 A a \sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4} + 16 A^3 b^3 d^2 - 16 A^3 a^2 b d^2} \right) \sqrt{\frac{\sqrt{-A^4 a^4 d^4 + 2 A^4 a^2 b^2 d^4 - A^4 b^4 d^4}}{4 d^4}} \\
&\quad + 2 \operatorname{atanh} \left(\frac{32 B^2 a^2 d^3 \sqrt{\tan(c + dx)} \sqrt{\frac{B^2 a b}{2 d^2} - \frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4}}}{16 B b \sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4} + 16 B^3 a^3 d^2 - 16 B^3 a b^2 d^2} \right. \\
&\quad \left. - \frac{32 B^2 b^2 d^3 \sqrt{\tan(c + dx)} \sqrt{\frac{B^2 a b}{2 d^2} - \frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4}}}{16 B b \sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4} + 16 B^3 a^3 d^2 - 16 B^3 a b^2 d^2} \right) \sqrt{\frac{B^2 a b}{2 d^2} - \frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4}} \\
&\quad - 2 \operatorname{atanh} \left(\frac{32 B^2 a^2 d^3 \sqrt{\tan(c + dx)} \sqrt{\frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4} + \frac{B^2 a b}{2 d^2}}}{16 B b \sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4} - 16 B^3 a^3 d^2 + 16 B^3 a b^2 d^2} \right. \\
&\quad \left. - \frac{32 B^2 b^2 d^3 \sqrt{\tan(c + dx)} \sqrt{\frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4} + \frac{B^2 a b}{2 d^2}}}{16 B b \sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4} - 16 B^3 a^3 d^2 + 16 B^3 a b^2 d^2} \right) \sqrt{\frac{\sqrt{-B^4 a^4 d^4 + 2 B^4 a^2 b^2 d^4 - B^4 b^4 d^4}}{4 d^4}} \\
&\quad - \frac{\frac{2 A a}{3} + 2 A b \tan(c + dx)}{d \tan(c + dx)^{3/2}} - \frac{2 B a}{d \sqrt{\tan(c + dx)}}
\end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x)))/tan(c + d*x)^(5/2),x)`

output

$$\begin{aligned}
& 2*\operatorname{atanh}((32*A^2*a^2*d^3*\tan(c + d*x))^{(1/2)}*(-(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)})/(4*d^4) - (A^2*a*b)/(2*d^2))^{(1/2)})/(16*A*a*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)} - 16*A^3*b^3*d^2 + 16*A^3*a^2*b*d^2) - \\
& (32*A^2*b^2*d^3*\tan(c + d*x))^{(1/2)}*(-(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)})/(4*d^4) - (A^2*a*b)/(2*d^2))^{(1/2)})/(16*A*a*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)} - 16*A^3*b^3*d^2 + 16*A^3*a^2*b*d^2) - \\
& (32*A^2*b^2*d^3*\tan(c + d*x))^{(1/2)}*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)})/(4*d^4) - (A^2*a*b)/(2*d^2))^{(1/2)})/(16*A*a*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)} + 16*A^3*b^3*d^2 - 16*A^3*a^2*b*d^2) - \\
& (32*A^2*b^2*d^3*\tan(c + d*x))^{(1/2)}*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)})/(4*d^4) - (A^2*a*b)/(2*d^2))^{(1/2)})/(16*A*a*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^{(1/2)} + 16*A^3*b^3*d^2 - 16*A^3*a^2*b*d^2) - \\
& (32*B^2*a^2*d^3*\tan(c + d*x))^{(1/2)}*((B^2*a*b)/(2*d^2) - (2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^{(1/2)})/(4*d^4))^{(1/2)})/(16*B*b*(2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^{(1/2)} + 16*B^3*a^3*d^2 - 16*B^3*a*b^2*d^2) - \\
& (32*B^2*b^2*d^3*\tan(c + d*x))^{(1/2)}*((B^2*a*b)/(2*d^2) - (2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^{(1/2)})/(4*d^4))^{(1/2)})/(16*...
\end{aligned}$$

3.383. $\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

3.384
$$\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

3.384.1 Optimal result 3640
 3.384.2 Mathematica [A] (verified) 3641
 3.384.3 Rubi [A] (verified) 3641
 3.384.4 Maple [A] (verified) 3646
 3.384.5 Fricas [B] (verification not implemented) 3647
 3.384.6 Sympy [F] 3647
 3.384.7 Maxima [A] (verification not implemented) 3648
 3.384.8 Giac [F(-1)] 3648
 3.384.9 Mupad [B] (verification not implemented) 3649

3.384.1 Optimal result

Integrand size = 31, antiderivative size = 254

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx \\ &= -\frac{(a(A - B) - b(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d} \\ &+ \frac{(a(A - B) - b(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d} \\ &+ \frac{(b(A - B) + a(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d} \\ &- \frac{(b(A - B) + a(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d} \\ &- \frac{2aA}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{2(Ab + aB)}{3d \tan^{\frac{3}{2}}(c + dx)} + \frac{2(aA - bB)}{d \sqrt{\tan(c + dx)}} \end{aligned}$$

output

```
1/2*(a*(A-B)-b*(A+B))*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)+1/2*(a
*(A-B)-b*(A+B))*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)+1/4*(b*(A-B)+
a*(A+B))*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)-1/4*(b*(A-B)+
a*(A+B))*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)+2*(A*a-B*b)/d
/tan(d*x+c)^(1/2)-2/5*a*A/d/tan(d*x+c)^(5/2)-2/3*(A*b+B*a)/d/tan(d*x+c)^(3
/2)
```

3.384.2 Mathematica [A] (verified)

Time = 1.32 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.78

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx =$$

$$\frac{30\sqrt{2}(a(A - B) - b(A + B)) \left(\arctan \left(1 - \sqrt{2}\sqrt{\tan(c + dx)} \right) - \arctan \left(1 + \sqrt{2}\sqrt{\tan(c + dx)} \right) \right) - 1}{d}$$

input `Integrate[((a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `-1/60*(30*Sqrt[2]*(a*(A - B) - b*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]) - 15*Sqrt[2]*(b*(A - B) + a*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]) + (24*a*A)/Tan[c + d*x]^(5/2) + (40*(A*b + a*B))/Tan[c + d*x]^(3/2) - (120*(a*A - b*B))/Sqrt[Tan[c + d*x]]/d`

3.384.3 Rubi [A] (verified)Time = 0.81 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.91, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.548$, Rules used = {3042, 4074, 3042, 4012, 25, 3042, 4012, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan(c + dx)^{7/2}} dx$$

$$\downarrow \text{4074}$$

$$\int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)} dx - \frac{2aA}{5d \tan^{\frac{5}{2}}(c + dx)}$$

$$\begin{aligned}
& \int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\tan(c + dx)^{5/2}} dx - \frac{2aA}{5d \tan^{5/2}(c + dx)} \\
& \quad \downarrow 3042 \\
& \int -\frac{aA - bB + (Ab + aB) \tan(c + dx)}{\tan^{3/2}(c + dx)} dx - \frac{2(aB + Ab)}{3d \tan^{3/2}(c + dx)} - \frac{2aA}{5d \tan^{5/2}(c + dx)} \\
& \quad \downarrow 4012 \\
& - \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\tan^{3/2}(c + dx)} dx - \frac{2(aB + Ab)}{3d \tan^{3/2}(c + dx)} - \frac{2aA}{5d \tan^{5/2}(c + dx)} \\
& \quad \downarrow 25 \\
& - \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\tan^{3/2}(c + dx)} dx - \frac{2(aB + Ab)}{3d \tan^{3/2}(c + dx)} - \frac{2aA}{5d \tan^{5/2}(c + dx)} \\
& \quad \downarrow 3042 \\
& - \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\tan(c + dx)^{3/2}} dx - \frac{2(aB + Ab)}{3d \tan^{3/2}(c + dx)} - \frac{2aA}{5d \tan^{5/2}(c + dx)} \\
& \quad \downarrow 4012 \\
& - \int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2(aB + Ab)}{3d \tan^{3/2}(c + dx)} + \frac{2(aA - bB)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{5d \tan^{5/2}(c + dx)} \\
& \quad \downarrow 3042 \\
& - \int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2(aB + Ab)}{3d \tan^{3/2}(c + dx)} + \frac{2(aA - bB)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{5d \tan^{5/2}(c + dx)} \\
& \quad \downarrow 4017 \\
& \frac{2 \int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)}}{d} - \frac{2(aB + Ab)}{3d \tan^{3/2}(c + dx)} + \frac{2(aA - bB)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{5d \tan^{5/2}(c + dx)} \\
& \quad \downarrow 1482 \\
& \frac{2 \left(\frac{1}{2}(a(A + B) + b(A - B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} - \frac{1}{2}(a(A - B) - b(A + B)) \int \frac{\tan(c + dx) + 1}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} \right)}{d} \\
& \quad - \frac{2(aB + Ab)}{3d \tan^{3/2}(c + dx)} + \frac{2(aA - bB)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{5d \tan^{5/2}(c + dx)} \\
& \quad \downarrow 1476 \\
& \frac{2 \left(\frac{1}{2}(a(A + B) + b(A - B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} - \frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{1}{2} \int \frac{1}{\tan(c + dx) - \sqrt{2} \sqrt{\tan(c + dx)}} dx \right) \right)}{d} \\
& \quad - \frac{2(aB + Ab)}{3d \tan^{3/2}(c + dx)} + \frac{2(aA - bB)}{d \sqrt{\tan(c + dx)}} - \frac{2aA}{5d \tan^{5/2}(c + dx)}
\end{aligned}$$

3.384. $\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan^2(c+dx)} dx$

↓ 1082

$$\frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{\int \frac{1}{\tan(c+dx)-1} d\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}} \right) \right)}{d} \\ \frac{2(aB + Ab)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{2(aA - bB)}{d\sqrt{\tan(c+dx)}} - \frac{2aA}{5d \tan^{\frac{5}{2}}(c+dx)}$$

↓ 217

$$\frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \\ \frac{2(aB + Ab)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{2(aA - bB)}{d\sqrt{\tan(c+dx)}} - \frac{2aA}{5d \tan^{\frac{5}{2}}(c+dx)}$$

↓ 1479

$$\frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(- \frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d} \\ \frac{2(aB + Ab)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{2(aA - bB)}{d\sqrt{\tan(c+dx)}} - \frac{2aA}{5d \tan^{\frac{5}{2}}(c+dx)}$$

↓ 25

$$\frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(a \right)}{d} \\ \frac{2(aB + Ab)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{2(aA - bB)}{d\sqrt{\tan(c+dx)}} - \frac{2aA}{5d \tan^{\frac{5}{2}}(c+dx)}$$

↓ 27

$$\frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d} \\ \frac{2(aB + Ab)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{2(aA - bB)}{d\sqrt{\tan(c+dx)}} - \frac{2aA}{5d \tan^{\frac{5}{2}}(c+dx)}$$

↓ 1103

3.384. $\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right) \frac{d}{3d \tan^{\frac{3}{2}}(c+dx) + \frac{2(aA - bB)}{d\sqrt{\tan(c+dx)}} - \frac{2aA}{5d \tan^{\frac{5}{2}}(c+dx)}}$$

input `Int[((a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2), x]`

output `(-2*(-1/2*((a*(A - B) - b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])) + ((b*(A - B) + a*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d - (2*a*A)/(5*d*Tan[c + d*x]^(5/2)) - (2*(A*b + a*B))/(3*d*Tan[c + d*x]^(3/2)) + (2*(a*A - b*B))/(d*Sqrt[Tan[c + d*x]])`

3.384.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

3.384.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 240, normalized size of antiderivative = 0.94

method	result
derivativedivides	$-\frac{2(Ab+Ba)}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(-aA+Bb)}{\sqrt{\tan(dx+c)}} - \frac{2aA}{5 \tan(dx+c)^{\frac{5}{2}}} + \frac{(-Ab-Ba)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan \left(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c)) \right) \right)}{4}$
default	$-\frac{2(Ab+Ba)}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2(-aA+Bb)}{\sqrt{\tan(dx+c)}} - \frac{2aA}{5 \tan(dx+c)^{\frac{5}{2}}} + \frac{(-Ab-Ba)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan \left(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c)) \right) \right)}{4}$
parts	$\frac{(Ab+Ba) \left(-\frac{2}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan \left(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c)) \right) \right)}{4} + 2 \arctan \left(-1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c)) \right) \right)}{d}$

```
input int((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x,method=_RETURNVER
BOSE)
```

```
output 1/d*(-2/3*(A*b+B*a)/tan(d*x+c)^(3/2)-2*(-A*a+B*b)/tan(d*x+c)^(1/2)-2/5*a*A
/tan(d*x+c)^(5/2)+1/4*(-A*b-B*a)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+t
an(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan
(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(A*a-B*b)*2^(1/2
)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+
tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(
d*x+c)^(1/2)))
```

$$3.384. \int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\tan^2(c+dx)} dx$$

3.384.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2290 vs. $2(216) = 432$.

Time = 0.40 (sec) , antiderivative size = 2290, normalized size of antiderivative = 9.02

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm=
"fricas")
```

```
output 1/30*(15*d*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A
^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2
+ B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d
^4))/d^2)*log(((A*a - B*b)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*
B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*
a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) - ((A^2*B - B^3)*a^3 + (A^3 - 5*
A*B^2)*a^2*b - (5*A^2*B - B^3)*a*b^2 - (A^3 - A*B^2)*b^3)*d)*sqrt((2*A*B*a
^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^
4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*
B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2) + ((A^4 - B^4)*
a^4 - 4*(A^3*B + A*B^3)*a^3*b - 4*(A^3*B + A*B^3)*a*b^3 - (A^4 - B^4)*b^4)
*sqrt(tan(d*x + c))*tan(d*x + c)^3 - 15*d*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2
*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B
^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 +
(A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2)*log(-((A*a - B*b)*d^3*sqrt(-((A^4
- 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 +
B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4)
- ((A^2*B - B^3)*a^3 + (A^3 - 5*A*B^2)*a^2*b - (5*A^2*B - B^3)*a*b^2 - (A
^3 - A*B^2)*b^3)*d)*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*
sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - ...
```

3.384.6 Sympy [F]

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx \\ &= \int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)**(7/2), x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))/tan(c + d*x)**(7/2), x)`

3.384.7 Maxima [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.83

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{30 \sqrt{2}((A - B)a - (A + B)b) \arctan\left(\frac{1}{2} \sqrt{2}\left(\sqrt{2} + 2 \sqrt{\tan(dx + c)}\right)\right) + 30 \sqrt{2}((A - B)a - (A + B)b) \arctan\left(\frac{1}{2} \sqrt{2}\left(\sqrt{2} - 2 \sqrt{\tan(dx + c)}\right)\right) + 15 \sqrt{2}((A + B)a + (A - B)b) \log(\sqrt{2} \sqrt{\tan(dx + c)} + \tan(dx + c) + 1) + 15 \sqrt{2}((A + B)a + (A - B)b) \log(-\sqrt{2} \sqrt{\tan(dx + c)} + \tan(dx + c) + 1) + 8(15(Aa - Bb) \tan(dx + c)^2 - 3Aa - 5(Ba + Ab) \tan(dx + c)) / \tan(dx + c)^{\frac{5}{2}}}{d}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2), x, algorithm="maxima")`

output `1/60*(30*sqrt(2)*((A - B)*a - (A + B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 30*sqrt(2)*((A - B)*a - (A + B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - 15*sqrt(2)*((A + B)*a + (A - B)*b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 15*sqrt(2)*((A + B)*a + (A - B)*b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 8*(15*(A*a - B*b)*tan(d*x + c)^2 - 3*A*a - 5*(B*a + A*b)*tan(d*x + c))/tan(d*x + c)^(5/2))/d`

3.384.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2), x, algorithm="giac")`

output `Timed out`

3.384.9 Mupad [B] (verification not implemented)

Time = 14.25 (sec) , antiderivative size = 1473, normalized size of antiderivative = 5.80

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \text{Too large to display}$$

```
input int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x)))/tan(c + d*x)^(7/2),x)
```

```
output 2*atanh((32*A^2*a^2*d^3*tan(c + d*x)^(1/2)*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4) + (A^2*a*b)/(2*d^2))^(1/2))/(16*A*b*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2) - 16*A^3*a^3*d^2 + 16*A^3*a*b^2*d^2) - (32*A^2*b^2*d^3*tan(c + d*x)^(1/2)*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4) + (A^2*a*b)/(2*d^2))^(1/2))/(16*A*b*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2) - 16*A^3*a^3*d^2 + 16*A^3*a*b^2*d^2))*((2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4) + (A^2*a*b)/(2*d^2))^(1/2) - 2*atanh((32*A^2*a^2*d^3*tan(c + d*x)^(1/2)*((A^2*a*b)/(2*d^2) - (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4))^(1/2))/(16*A*b*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2) + 16*A^3*a^3*d^2 - 16*A^3*a*b^2*d^2) - (32*A^2*b^2*d^3*tan(c + d*x)^(1/2)*((A^2*a*b)/(2*d^2) - (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4))^(1/2))/(16*A*b*(2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2) + 16*A^3*a^3*d^2 - 16*A^3*a*b^2*d^2))*((A^2*a*b)/(2*d^2) - (2*A^4*a^2*b^2*d^4 - A^4*b^4*d^4 - A^4*a^4*d^4)^(1/2)/(4*d^4))^(1/2) + 2*atanh((32*B^2*a^2*d^3*tan(c + d*x)^(1/2)*(- (2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^(1/2)/(4*d^4) - (B^2*a*b)/(2*d^2))^(1/2))/(16*B*a*(2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^(1/2) - 16*B^3*b^3*d^2 + 16*B^3*a^2*b*d^2) - (32*B^2*b^2*d^3*tan(c + d*x)^(1/2)*(- (2*B^4*a^2*b^2*d^4 - B^4*b^4*d^4 - B^4*a^4*d^4)^(1/2)/(4*d^4) - (B^2*a*b)/(2*d^2))^(1/2))/(16*B*...
```


3.385 $\int \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.385.1 Optimal result	3650
3.385.2 Mathematica [C] (verified)	3651
3.385.3 Rubi [A] (verified)	3652
3.385.4 Maple [A] (verified)	3659
3.385.5 Fricas [B] (verification not implemented)	3659
3.385.6 Sympy [F]	3660
3.385.7 Maxima [A] (verification not implemented)	3661
3.385.8 Giac [F(-1)]	3661
3.385.9 Mupad [B] (verification not implemented)	3662

3.385.1 Optimal result

Integrand size = 33, antiderivative size = 394

$$\int \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{(a^2(A-B) - b^2(A-B) - 2ab(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(a^2(A-B) - b^2(A-B) - 2ab(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(2ab(A-B) + a^2(A+B) - b^2(A+B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{(2ab(A-B) + a^2(A+B) - b^2(A+B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}d}$$

$$- \frac{2(2aAb + a^2B - b^2B) \sqrt{\tan(c+dx)}}{d} + \frac{2(a^2A - Ab^2 - 2abB) \tan^{\frac{3}{2}}(c+dx)}{3d}$$

$$+ \frac{2(2aAb + a^2B - b^2B) \tan^{\frac{5}{2}}(c+dx)}{5d} + \frac{2b(9Ab + 11aB) \tan^{\frac{7}{2}}(c+dx)}{63d}$$

$$+ \frac{2bB \tan^{\frac{7}{2}}(c+dx)(a + b \tan(c+dx))}{9d}$$

output
$$\begin{aligned} & -1/2*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}) \\ & /d*2^{(1/2)}-1/2*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*\arctan(1+2^{(1/2)}*\tan(d*x+ \\ & c)^{(1/2)})/d*2^{(1/2)}-1/4*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*\ln(1-2^{(1/2)}*\tan \\ & (d*x+c)^{(1/2)}+\tan(d*x+c))/d*2^{(1/2)}+1/4*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))* \\ & \ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/d*2^{(1/2)}-2*(2*A*a*b+B*a^2-B*b^2 \\ &)*\tan(d*x+c)^{(1/2)}/d+2/3*(A*a^2-A*b^2-2*B*a*b)*\tan(d*x+c)^{(3/2)}/d+2/5*(2*A \\ & *a*b+B*a^2-B*b^2)*\tan(d*x+c)^{(5/2)}/d+2/63*b*(9*A*b+11*B*a)*\tan(d*x+c)^{(7/2) \\ &)/d+2/9*b*B*\tan(d*x+c)^{(7/2)}*(a+b*\tan(d*x+c))/d \end{aligned}$$

3.385.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.15 (sec) , antiderivative size = 304, normalized size of antiderivative = 0.77

$$\begin{aligned} & \int \tan^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx \\ & = \frac{2bB\tan^{\frac{7}{2}}(c+dx)(a+b\tan(c+dx))}{9d} + \frac{2}{9} \left(\frac{b(9Ab+11aB)\tan^{\frac{7}{2}}(c+dx)}{7d} \right. \\ & \quad + \frac{i(\frac{9}{2}(a^2A-Ab^2-2abB) - \frac{9}{2}i(2aAb+a^2B-b^2B)) \left(-2\sqrt[4]{-1} \arctan\left((-1)^{3/4} \sqrt{\tan(c+dx)} \right) - 2\sqrt{\tan(c+dx)} \right)}{2d} \\ & \quad \left. - \frac{i(\frac{9}{2}(a^2A-Ab^2-2abB) + \frac{9}{2}i(2aAb+a^2B-b^2B)) \left(-2\sqrt[4]{-1} \operatorname{arctanh}\left((-1)^{3/4} \sqrt{\tan(c+dx)} \right) - 2\sqrt{\tan(c+dx)} \right)}{2d} \right) \end{aligned}$$

input `Integrate[Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output
$$\begin{aligned} & (2*b*B*\tan[c + d*x]^{(7/2)}*(a + b*\tan[c + d*x]))/(9*d) + (2*((b*(9*A*b + 11 \\ & *a*B)*\tan[c + d*x]^{(7/2)})/(7*d) + ((I/2)*((9*(a^2*A - A*b^2 - 2*a*b*B))/2 \\ & - ((9*I)/2)*(2*a*A*b + a^2*B - b^2*B))*(-2*(-1)^{(1/4)}*ArcTan[(-1)^{(3/4)}*Sq \\ & rt[\tan[c + d*x]]] - 2*sqrt[\tan[c + d*x]] - ((2*I)/3)*\tan[c + d*x]^{(3/2)} + \\ & (2*\tan[c + d*x]^{(5/2)})/5))/d - ((I/2)*((9*(a^2*A - A*b^2 - 2*a*b*B))/2 + (\\ & (9*I)/2)*(2*a*A*b + a^2*B - b^2*B))*(-2*(-1)^{(1/4)}*ArcTanh[(-1)^{(3/4)}*sqrt \\ & [\tan[c + d*x]]] - 2*sqrt[\tan[c + d*x]] + ((2*I)/3)*\tan[c + d*x]^{(3/2)} + (2 \\ & *\tan[c + d*x]^{(5/2)})/5))/d)/9 \end{aligned}$$

3.385.3 Rubi [A] (verified)

Time = 1.41 (sec) , antiderivative size = 349, normalized size of antiderivative = 0.89, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4090, 27, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4011, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c+dx)^{5/2}(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4090} \\
 & \frac{2}{9} \int \frac{1}{2} \tan^{\frac{5}{2}}(c+ \\
 & dx) (b(9Ab+11aB) \tan^2(c+dx) + 9(Ba^2+2Aba-b^2B) \tan(c+dx) + a(9aA-7bB)) dx + \\
 & \quad \frac{2bB \tan^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))}{9d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{9} \int \tan^{\frac{5}{2}}(c+ \\
 & dx) (b(9Ab+11aB) \tan^2(c+dx) + 9(Ba^2+2Aba-b^2B) \tan(c+dx) + a(9aA-7bB)) dx + \\
 & \quad \frac{2bB \tan^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))}{9d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{9} \int \tan(c+ \\
 & dx)^{5/2} (b(9Ab+11aB) \tan(c+dx)^2 + 9(Ba^2+2Aba-b^2B) \tan(c+dx) + a(9aA-7bB)) dx + \\
 & \quad \frac{2bB \tan^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))}{9d} \\
 & \quad \downarrow \text{4113} \\
 & \frac{1}{9} \left(\int \tan^{\frac{5}{2}}(c+dx) (9(Aa^2-2bBa-Ab^2) + 9(Ba^2+2Aba-b^2B) \tan(c+dx)) dx + \frac{2b(11aB+9Ab) \tan^{\frac{7}{2}}(c+}{7d} \right. \\
 & \quad \left. \frac{2bB \tan^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))}{9d} \right)
 \end{aligned}$$

3.385. $\int \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

$$\downarrow 3042$$

$$\frac{1}{9} \left(\int \tan(c+dx)^{5/2} (9(Aa^2 - 2bBa - Ab^2) + 9(Ba^2 + 2Aba - b^2B) \tan(c+dx)) dx + \frac{2b(11aB + 9Ab) \tan^{7/2}(c+dx)}{7d} \right. \\ \left. + \frac{2bB \tan^{7/2}(c+dx)(a + b \tan(c+dx))}{9d} \right)$$

$$\downarrow 4011$$

$$\frac{1}{9} \left(\int \tan^{3/2}(c+dx) (9(Aa^2 - 2bBa - Ab^2) \tan(c+dx) - 9(Ba^2 + 2Aba - b^2B)) dx + \frac{18(a^2B + 2aAb - b^2B) \tan^{5/2}(c+dx)}{5d} \right. \\ \left. + \frac{2bB \tan^{7/2}(c+dx)(a + b \tan(c+dx))}{9d} \right)$$

$$\downarrow 3042$$

$$\frac{1}{9} \left(\int \tan(c+dx)^{3/2} (9(Aa^2 - 2bBa - Ab^2) \tan(c+dx) - 9(Ba^2 + 2Aba - b^2B)) dx + \frac{18(a^2B + 2aAb - b^2B) \tan^{5/2}(c+dx)}{5d} \right. \\ \left. + \frac{2bB \tan^{7/2}(c+dx)(a + b \tan(c+dx))}{9d} \right)$$

$$\downarrow 4011$$

$$\frac{1}{9} \left(\int \sqrt{\tan(c+dx)} (-9(Aa^2 - 2bBa - Ab^2) - 9(Ba^2 + 2Aba - b^2B) \tan(c+dx)) dx + \frac{18(a^2B + 2aAb - b^2B) \tan^{3/2}(c+dx)}{5d} \right. \\ \left. + \frac{2bB \tan^{7/2}(c+dx)(a + b \tan(c+dx))}{9d} \right)$$

$$\downarrow 3042$$

$$\frac{1}{9} \left(\int \sqrt{\tan(c+dx)} (-9(Aa^2 - 2bBa - Ab^2) - 9(Ba^2 + 2Aba - b^2B) \tan(c+dx)) dx + \frac{18(a^2B + 2aAb - b^2B) \tan^{3/2}(c+dx)}{5d} \right. \\ \left. + \frac{2bB \tan^{7/2}(c+dx)(a + b \tan(c+dx))}{9d} \right)$$

$$\downarrow 4011$$

$$\frac{1}{9} \left(\int \frac{9(Ba^2 + 2Aba - b^2B) - 9(Aa^2 - 2bBa - Ab^2) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx + \frac{18(a^2B + 2aAb - b^2B) \tan^{5/2}(c+dx)}{5d} + \frac{2bB \tan^{7/2}(c+dx)(a + b \tan(c+dx))}{9d} \right)$$

$$\downarrow 3042$$

3.385. $\int \tan^{5/2}(c+dx)(a + b \tan(c+dx))^2(A + B \tan(c+dx)) dx$

$$\frac{1}{9} \left(\int \frac{9(Ba^2 + 2Aba - b^2B) - 9(Aa^2 - 2bBa - Ab^2) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{18(a^2B + 2aAb - b^2B) \tan^{\frac{5}{2}}(c + dx)}{5d} + \frac{6(a^2A - 2aAb + b^2B) \tan^{\frac{7}{2}}(c + dx)}{9d} \right)$$

↓ 4017

$$\frac{1}{9} \left(\frac{2 \int \frac{9(Ba^2 + 2Aba - b^2B) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{d} + \frac{18(a^2B + 2aAb - b^2B) \tan^{\frac{5}{2}}(c + dx)}{5d} + \frac{6(a^2A - 2aAb + b^2B) \tan^{\frac{7}{2}}(c + dx)}{9d} \right)$$

↓ 27

$$\frac{1}{9} \left(\frac{18 \int \frac{Ba^2 + 2Aba - b^2B - (Aa^2 - 2bBa - Ab^2) \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{d} + \frac{18(a^2B + 2aAb - b^2B) \tan^{\frac{5}{2}}(c + dx)}{5d} + \frac{6(a^2A - 2aAb + b^2B) \tan^{\frac{7}{2}}(c + dx)}{9d} \right)$$

↓ 1482

$$\frac{1}{9} \left(\frac{18 \left(\frac{1}{2}(a^2(A + B) + 2ab(A - B) - b^2(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(a^2(A - B) - 2ab(A + B) - b^2(A + B)) \int \frac{1 + \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} \right)}{d} + \frac{18(a^2B + 2aAb - b^2B) \tan^{\frac{5}{2}}(c + dx)}{5d} + \frac{6(a^2A - 2aAb + b^2B) \tan^{\frac{7}{2}}(c + dx)}{9d} \right)$$

↓ 1476

$$\frac{1}{9} \left(\frac{18 \left(\frac{1}{2}(a^2(A + B) + 2ab(A - B) - b^2(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(a^2(A - B) - 2ab(A + B) - b^2(A + B)) \int \frac{1 + \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} \right)}{d} + \frac{18(a^2B + 2aAb - b^2B) \tan^{\frac{5}{2}}(c + dx)}{5d} + \frac{6(a^2A - 2aAb + b^2B) \tan^{\frac{7}{2}}(c + dx)}{9d} \right)$$

↓ 1082

3.385. $\int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\frac{1}{9} \left(\frac{18 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A+B)) \int \frac{1+\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} \right)$$

$$\frac{2bB \tan^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))}{9d}$$

↓ 217

$$\frac{1}{9} \left(\frac{18 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A+B)) \int \frac{1+\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} \right)$$

$$\frac{2bB \tan^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))}{9d}$$

↓ 1479

$$\frac{1}{9} \left(\frac{18 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{d} \right)}{d} \right)$$

$$\frac{2bB \tan^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))}{9d}$$

↓ 25

$$\frac{1}{9} \left(\frac{18 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{d} \right)}{d} \right)$$

$$\frac{2bB \tan^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))}{9d}$$

↓ 27

3.385. $\int \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

$$\frac{1}{9} \left(\frac{18 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}} \right) \right)}{2bB \tan^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))} \right)$$

\downarrow 1103

$$\frac{1}{9} \left(\frac{18 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{2bB \tan^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))} \right) d$$

input `Int[Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(2*b*B*Tan[c + d*x]^(7/2)*(a + b*Tan[c + d*x]))/(9*d) + ((18*(-1/2*((a^2*(A - B) - b^2*(A - B) - 2*a*b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])) + ((2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d - (18*(2*a*A*b + a^2*B - b^2*B)*Sqrt[Tan[c + d*x]])/d + (6*(a^2*A - A*b^2 - 2*a*b*B)*Tan[c + d*x]^(3/2))/d + (18*(2*a*A*b + a^2*B - b^2*B)*Tan[c + d*x]^(5/2))/(5*d) + (2*b*(9*A*b + 11*a*B)*Tan[c + d*x]^(7/2))/(7*d))/9`

3.385.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 $\text{Int}[(a + b \tan(e + f x))^m (c + d \tan(e + f x) + (f x))], x_{\text{Symbol}}] \rightarrow \text{Simp}[d (a + b \tan[e + f x])^m / (f m), x] + \text{Int}[(a + b \tan[e + f x])^{m-1} \text{Simp}[a c - b d + (b c + a d) \tan[e + f x], x], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]

rule 4017 $\text{Int}[(c + d \tan(e + f x)) / \text{Sqrt}[b \tan(e + f x) + (f x)], x_{\text{Symbol}}] \rightarrow \text{Simp}[2/f \text{ Subst}[\text{Int}[(b c + d x^2) / (b^2 + x^4), x], x, \text{Sqrt}[b \tan[e + f x]]], x] /;$ FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]

rule 4090 $\text{Int}[(a + b \tan(e + f x))^m (A + B \tan(e + f x) + (f x))^n (c + d \tan(e + f x))], x_{\text{Symbol}}] \rightarrow \text{Simp}[b B (a + b \tan[e + f x])^{m-1} (c + d \tan[e + f x])^{n+1} / (d f (m + n)), x] + \text{Simp}[1 / (d (m + n)) \text{Int}[(a + b \tan[e + f x])^{m-2} (c + d \tan[e + f x])^n \text{Simp}[a^2 A d (m + n) - b B (b c (m - 1) + a d (n + 1)) + d (m + n) (2 a A b + B (a^2 - b^2)) \tan[e + f x] - (b B (b c - a d) (m - 1) - b (A b + a B) d (m + n)) \tan[e + f x]^2, x], x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

rule 4113 $\text{Int}[(a + b \tan(e + f x))^m (A + B \tan(e + f x) + (f x)) + (C \tan(e + f x) + (f x))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[C (a + b \tan[e + f x])^{m+1} / (b f (m + 1)), x] + \text{Int}[(a + b \tan[e + f x])^m \text{Simp}[A - C + B \tan[e + f x], x], x] /;$ FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]

3.385.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 374, normalized size of antiderivative = 0.95

method	result
derivativedivides	$\frac{2Bb^2 \left(\tan^{\frac{9}{2}}(dx+c)\right)}{9} + \frac{2Ab^2 \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} + \frac{4Bab \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} + \frac{4Aab \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{2Ba^2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} - \frac{2Bb^2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5}$
default	$\frac{2Bb^2 \left(\tan^{\frac{9}{2}}(dx+c)\right)}{9} + \frac{2Ab^2 \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} + \frac{4Bab \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} + \frac{4Aab \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{2Ba^2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} - \frac{2Bb^2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5}$
parts	$\frac{(Ab^2+2Bab) \left(\frac{2 \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} - \frac{2 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} + \frac{\sqrt{2} \left(\ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})}{4} \right) \right)}{d}$

```
input int(tan(d*x+c)^(5/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 1/d*(2/9*B*b^2*tan(d*x+c)^(9/2)+2/7*A*b^2*tan(d*x+c)^(7/2)+4/7*B*a*b*tan(d
*x+c)^(7/2)+4/5*A*a*b*tan(d*x+c)^(5/2)+2/5*B*a^2*tan(d*x+c)^(5/2)-2/5*B*b^
2*tan(d*x+c)^(5/2)+2/3*A*a^2*tan(d*x+c)^(3/2)-2/3*A*b^2*tan(d*x+c)^(3/2)-4
/3*B*a*b*tan(d*x+c)^(3/2)-4*A*a*b*tan(d*x+c)^(1/2)-2*B*a^2*tan(d*x+c)^(1/2
)+2*tan(d*x+c)^(1/2)*B*b^2+1/4*(2*A*a*b+B*a^2-B*b^2)*2^(1/2)*(ln((1+2^(1/2
))*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*
arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+
1/4*(-A*a^2+A*b^2+2*B*a*b)*2^(1/2)*(ln((1-2^(1/2))*tan(d*x+c)^(1/2)+tan(d*x
+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c
)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))
```

3.385.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4337 vs. 2(348) = 696.

Time = 0.75 (sec) , antiderivative size = 4337, normalized size of antiderivative = 11.01

$$\int \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^(5/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorith
m="fracas")
```

3.385. $\int \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

output `1/630*(315*d*sqrt((2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2)*log(((A*a^2 - 2*B*a*b - A*b^2)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4) - ((A^2*B - B^3)*a^6 + 2*(A^3 - 5*A*B^2)*a^5*b - (23*A^2*B - 7*B^3)*a^4*b^2 - 4*(3*A^3 - 7*A*B^2)*a^3*b^3 + (23*A^2*B - 7*B^3)*a^2*b^4 + 2*(A^3 - 5*A*B^2)*a*b^5 - (A^2*B - B^3)*b^6)*d)*sqrt((2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2) - ((A^4 - B^4)*a^8 - 8*(A^3*B + A*B^3)*a^7*b - 4*(A^4 - B^4)*a^6*b^2 - 8*(A^3*B + A*B^3)*a^5*b^3 - 10*(A^4 - ...`

3.385.6 Sympy [F]

$$\int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^2 \tan^{\frac{5}{2}}(c + dx) dx$$

input `integrate(tan(d*x+c)**(5/2)*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**2*tan(c + d*x)**(5/2), x)`

3.385.7 Maxima [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 329, normalized size of antiderivative = 0.84

$$\int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{280 B b^2 \tan(dx + c)^{\frac{9}{2}} + 360 (2 B a b + A b^2) \tan(dx + c)^{\frac{7}{2}} + 504 (B a^2 + 2 A a b - B b^2) \tan(dx + c)^{\frac{5}{2}} - 630 \sqrt{2} \left((A - B) a^2 - 2(A + B) a b - (A - B) b^2 \right) \arctan\left(\frac{1}{2} \sqrt{2} (\sqrt{2} + 2 \sqrt{\tan(dx + c)})\right) - 630 \sqrt{2} \left((A - B) a^2 - 2(A + B) a b - (A - B) b^2 \right) \arctan\left(-\frac{1}{2} \sqrt{2} (\sqrt{2} - 2 \sqrt{\tan(dx + c)})\right) + 315 \sqrt{2} \left((A + B) a^2 + 2(A - B) a b - (A + B) b^2 \right) \log(\sqrt{2} \sqrt{\tan(dx + c)} + \tan(dx + c) + 1) - 315 \sqrt{2} \left((A + B) a^2 + 2(A - B) a b - (A + B) b^2 \right) \log(-\sqrt{2} \sqrt{\tan(dx + c)} + \tan(dx + c) + 1) + 840 (A a^2 - 2 B a b - A b^2) \tan(dx + c)^{\frac{3}{2}} - 2520 (B a^2 + 2 A a b - B b^2) \sqrt{\tan(dx + c)}}{d}$$

```
input integrate(tan(d*x+c)^(5/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")
```

```
output 1/1260*(280*B*b^2*tan(d*x + c)^(9/2) + 360*(2*B*a*b + A*b^2)*tan(d*x + c)^(7/2) + 504*(B*a^2 + 2*A*a*b - B*b^2)*tan(d*x + c)^(5/2) - 630*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) - 630*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + 315*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - 315*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 840*(A*a^2 - 2*B*a*b - A*b^2)*tan(d*x + c)^(3/2) - 2520*(B*a^2 + 2*A*a*b - B*b^2)*sqrt(tan(d*x + c)))/d
```

3.385.8 Giac [F(-1)]

Timed out.

$$\int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \text{Timed out}$$

```
input integrate(tan(d*x+c)^(5/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")
```

```
output Timed out
```

3.385.9 Mupad [B] (verification not implemented)

Time = 29.53 (sec) , antiderivative size = 3914, normalized size of antiderivative = 9.93

$$\int \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(tan(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)
```

```
output atan((B^2*a^4*tan(c + d*x)^(1/2)*((B^2*a*b^3)/d^2 - (12*B^4*a^2*b^6*d^4 -
B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)
/(4*d^4) - (B^2*a^3*b)/d^2)^(1/2)*32i)/((16*B*a^2*(12*B^4*a^2*b^6*d^4 - B^
4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2))/
d^3 - (192*B^3*a^3*b^3)/d - (16*B*b^2*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 -
B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2))/d^3 + (32*B^
3*a*b^5)/d + (32*B^3*a^5*b)/d) + (B^2*b^4*tan(c + d*x)^(1/2)*((B^2*a*b^3)/
d^2 - (12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4
+ 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4) - (B^2*a^3*b)/d^2)^(1/2)*32i)/((16*B*
a^2*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 +
12*B^4*a^6*b^2*d^4)^(1/2))/d^3 - (192*B^3*a^3*b^3)/d - (16*B*b^2*(12*B^4*
a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*
b^2*d^4)^(1/2))/d^3 + (32*B^3*a*b^5)/d + (32*B^3*a^5*b)/d) - (B^2*a^2*b^2*
tan(c + d*x)^(1/2)*((B^2*a*b^3)/d^2 - (12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 -
B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4) - (B^
2*a^3*b)/d^2)^(1/2)*192i)/((16*B*a^2*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B
^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2))/d^3 - (192*B^
3*a^3*b^3)/d - (16*B*b^2*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 -
38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2))/d^3 + (32*B^3*a*b^5)/d +
(32*B^3*a^5*b)/d))*((B^2*a*b^3)/d^2 - (12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4...
```

$$\mathbf{3.386} \quad \int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

3.386.1 Optimal result	3663
3.386.2 Mathematica [C] (verified)	3664
3.386.3 Rubi [A] (verified)	3664
3.386.4 Maple [A] (verified)	3671
3.386.5 Fricas [B] (verification not implemented)	3671
3.386.6 Sympy [F]	3672
3.386.7 Maxima [A] (verification not implemented)	3673
3.386.8 Giac [F(-1)]	3673
3.386.9 Mupad [B] (verification not implemented)	3674

3.386.1 Optimal result

Integrand size = 33, antiderivative size = 360

$$\begin{aligned} & \int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx \\ &= \frac{(2ab(A-B) + a^2(A+B) - b^2(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d} \\ & \quad - \frac{(2ab(A-B) + a^2(A+B) - b^2(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d} \\ & \quad + \frac{(a^2(A-B) - b^2(A-B) - 2ab(A+B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}d} \\ & \quad - \frac{(a^2(A-B) - b^2(A-B) - 2ab(A+B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}d} \\ & \quad + \frac{2(a^2A - Ab^2 - 2abB) \sqrt{\tan(c+dx)}}{d} + \frac{2(2aAb + a^2B - b^2B) \tan^{\frac{3}{2}}(c+dx)}{3d} \\ & \quad + \frac{2b(7Ab + 9aB) \tan^{\frac{5}{2}}(c+dx)}{35d} + \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))}{7d} \end{aligned}$$

$$3.386. \quad \int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

output
$$\begin{aligned} & -1/2*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}) \\ & /d*2^{(1/2)}-1/2*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*\arctan(1+2^{(1/2)}*\tan(d*x+ \\ & c)^{(1/2)})/d*2^{(1/2)}+1/4*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*\ln(1-2^{(1/2)}*\tan \\ & (d*x+c)^{(1/2)}+\tan(d*x+c))/d*2^{(1/2)}-1/4*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))* \\ & \ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/d*2^{(1/2)}+2*(A*a^2-A*b^2-2*B*a*b \\ &)*\tan(d*x+c)^{(1/2)}/d+2/3*(2*A*a*b+B*a^2-B*b^2)*\tan(d*x+c)^{(3/2)}/d+2/35*b*(\\ & 7*A*b+9*B*a)*\tan(d*x+c)^{(5/2)}/d+2/7*b*B*\tan(d*x+c)^{(5/2)}*(a+b*\tan(d*x+c))/ \\ & d \end{aligned}$$

3.386.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.37 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.49

$$\int \tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

$$= \frac{105\sqrt[4]{-1}(a-ib)^2(A-iB)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)+105\sqrt[4]{-1}(a+ib)^2(A+iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{d}$$

input `Integrate[Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output
$$\begin{aligned} & (105*(-1)^{(1/4)}*(a - I*b)^2*(A - I*B)*\operatorname{ArcTan}[(-1)^{(3/4)}*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]] \\ &] + 105*(-1)^{(1/4)}*(a + I*b)^2*(A + I*B)*\operatorname{ArcTanh}[(-1)^{(3/4)}*\operatorname{Sqrt}[\operatorname{Tan}[c + d \\ & *x]]] + 2*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]*(105*(a^2*A - A*b^2 - 2*a*b*B) + 35*(2*a*A*b \\ & + a^2*B - b^2*B)*\operatorname{Tan}[c + d*x] + 21*b*(A*b + 2*a*B)*\operatorname{Tan}[c + d*x]^2 + 15*b^2 \\ & *B*\operatorname{Tan}[c + d*x]^3))/(105*d) \end{aligned}$$

3.386.3 Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 317, normalized size of antiderivative = 0.88, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 4090, 27, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.386. $\int \tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$

$$\begin{aligned}
& \int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx \\
& \quad \downarrow \text{3042} \\
& \int \tan(c+dx)^{3/2}(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx \\
& \quad \downarrow \text{4090} \\
& \frac{2}{7} \int \frac{1}{2} \tan^{\frac{3}{2}}(c+dx) (b(7Ab+9aB) \tan^2(c+dx) + 7(Ba^2+2Aba-b^2B) \tan(c+dx) + a(7aA-5bB)) dx + \\
& \quad \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))}{7d} \\
& \quad \downarrow \text{27} \\
& \frac{1}{7} \int \tan^{\frac{3}{2}}(c+dx) (b(7Ab+9aB) \tan^2(c+dx) + 7(Ba^2+2Aba-b^2B) \tan(c+dx) + a(7aA-5bB)) dx + \\
& \quad \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))}{7d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \int \tan(c+dx)^{3/2} (b(7Ab+9aB) \tan(c+dx)^2 + 7(Ba^2+2Aba-b^2B) \tan(c+dx) + a(7aA-5bB)) dx + \\
& \quad \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))}{7d} \\
& \quad \downarrow \text{4113} \\
& \frac{1}{7} \left(\int \tan^{\frac{3}{2}}(c+dx) (7(Aa^2-2bBa-Ab^2) + 7(Ba^2+2Aba-b^2B) \tan(c+dx)) dx + \frac{2b(9aB+7Ab) \tan^{\frac{5}{2}}(c+dx)}{5d} \right. \\
& \quad \left. \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))}{7d} \right) \\
& \quad \downarrow \text{3042} \\
& \frac{1}{7} \left(\int \tan(c+dx)^{3/2} (7(Aa^2-2bBa-Ab^2) + 7(Ba^2+2Aba-b^2B) \tan(c+dx)) dx + \frac{2b(9aB+7Ab) \tan^{\frac{5}{2}}(c+dx)}{5d} \right. \\
& \quad \left. \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))}{7d} \right) \\
& \quad \downarrow \text{4011}
\end{aligned}$$

3.386. $\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

$$\frac{1}{7} \left(\int \sqrt{\tan(c+dx)} (7(Aa^2 - 2bBa - Ab^2) \tan(c+dx) - 7(Ba^2 + 2Aba - b^2B)) dx + \frac{14(a^2B + 2aAb - b^2B) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))}{7d} \right)$$

↓ 3042

$$\frac{1}{7} \left(\int \sqrt{\tan(c+dx)} (7(Aa^2 - 2bBa - Ab^2) \tan(c+dx) - 7(Ba^2 + 2Aba - b^2B)) dx + \frac{14(a^2B + 2aAb - b^2B) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))}{7d} \right)$$

↓ 4011

$$\frac{1}{7} \left(\int \frac{-7(Aa^2 - 2bBa - Ab^2) - 7(Ba^2 + 2Aba - b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx + \frac{14(a^2B + 2aAb - b^2B) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))}{7d} \right)$$

↓ 3042

$$\frac{1}{7} \left(\int \frac{-7(Aa^2 - 2bBa - Ab^2) - 7(Ba^2 + 2Aba - b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx + \frac{14(a^2B + 2aAb - b^2B) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))}{7d} \right)$$

↓ 4017

$$\frac{1}{7} \left(\frac{2 \int -\frac{7(Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c+dx))}{\tan^2(c+dx) + 1} d\sqrt{\tan(c+dx)}}{d} + \frac{14(a^2B + 2aAb - b^2B) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{14(a^2B + 2aAb - b^2B) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))}{7d} \right)$$

↓ 27

$$\frac{1}{7} \left(-\frac{14 \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c+dx)}{\tan^2(c+dx) + 1} d\sqrt{\tan(c+dx)}}{d} + \frac{14(a^2B + 2aAb - b^2B) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{14(a^2B + 2aAb - b^2B) \tan^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))}{7d} \right)$$

↓ 1482

3.386. $\int \tan^{\frac{3}{2}}(c+dx)(a + b \tan(c+dx))^2(A + B \tan(c+dx)) dx$

$$\frac{1}{7} \left(\frac{14 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^2(A + B) + 2ab(A - B)) \right)}{d} - \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))}{7d} \right) \downarrow 1476$$

$$\frac{1}{7} \left(\frac{14 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^2(A + B) + 2ab(A - B)) \right)}{d} - \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))}{7d} \right) \downarrow 1082$$

$$\frac{1}{7} \left(\frac{14 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^2(A + B) + 2ab(A - B)) \right)}{d} - \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))}{7d} \right) \downarrow 217$$

$$\frac{1}{7} \left(\frac{14 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^2(A + B) + 2ab(A - B)) \right)}{d} - \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))}{7d} \right) \downarrow 1479$$

$$\frac{1}{7} \left(\frac{14 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \left(- \frac{\int - \frac{\sqrt{2} - 2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int - \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)})}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{d} - \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))}{7d} \right)$$

3.386. $\int \tan^{\frac{3}{2}}(c+dx)(a + b \tan(c+dx))^2(A + B \tan(c+dx)) dx$

$$\begin{array}{c}
\downarrow 25 \\
\frac{1}{7} \left(\frac{14 \left(\frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}} \frac{d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{\frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))}{7d}} \right. \\
\downarrow 27 \\
\frac{1}{7} \left(\frac{14 \left(\frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}} \frac{d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{\frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))}{7d}} \right. \\
\downarrow 1103 \\
\frac{1}{7} \left(\frac{14 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \right)}{\frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))}{7d}} \right)
\end{array}$$

input `Int[Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(2*b*B*Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x]))/(7*d) + ((-14*(((2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 + ((a^2*(A - B) - b^2*(A - B) - 2*a*b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d + (14*(a^2*A - A*b^2 - 2*a*b*B)*Sqrt[Tan[c + d*x]])/d + (14*(2*a*A*b + a^2*B - b^2*B)*Tan[c + d*x]^(3/2))/(3*d) + (2*b*(7*A*b + 9*a*B)*Tan[c + d*x]^(5/2))/(5*d))/7`

3.386.3.1 Defintions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, x], x]$
- rule 27 $\text{Int}[(a_)*(\text{Fx}_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[\text{Fx}, (b_)*(\text{Gx}_) /; \text{FreeQ}[b, x]]$
- rule 217 $\text{Int}[(a_) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 1082 $\text{Int}[(a_) + (b_.)*(x_) + (c_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*c*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \quad \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_) + (e_.)*(x_)]/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$
- rule 1476 $\text{Int}[(d_) + (e_.)*(x_)^2]/((a_) + (c_.)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Simp}[e/(2*c) \quad \text{Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Simp}[e/(2*c) \quad \text{Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{PosQ}[d*e]$
- rule 1479 $\text{Int}[(d_) + (e_.)*(x_)^2]/((a_) + (c_.)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2*(d/e), 2]\}, \text{Simp}[e/(2*c*q) \quad \text{Int}[(q - 2*x)/\text{Simp}[d/e + q*x - x^2, x], x], x] + \text{Simp}[e/(2*c*q) \quad \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{NegQ}[d*e]$
- rule 1482 $\text{Int}[(d_) + (e_.)*(x_)^2]/((a_) + (c_.)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a*c, 2]\}, \text{Simp}[(d*q + a*e)/(2*a*c) \quad \text{Int}[(q + c*x^2)/(a + c*x^4), x], x] + \text{Simp}[(d*q - a*e)/(2*a*c) \quad \text{Int}[(q - c*x^2)/(a + c*x^4), x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{NeQ}[c*d^2 + a*e^2, 0] \ \&\& \ \text{NeQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{NegQ}[(a)*c]$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

3.386.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 333, normalized size of antiderivative = 0.92

method	result
derivativedivides	$\frac{2Bb^2 \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} + \frac{2Ab^2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{4Bab \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{4Aab \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} + \frac{2Ba^2 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} - \frac{2Bb^2 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3}$
default	$\frac{2Bb^2 \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} + \frac{2Ab^2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{4Bab \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{4Aab \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} + \frac{2Ba^2 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} - \frac{2Bb^2 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3}$
parts	$(Ab^2+2Bab) \left(\frac{2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} - 2(\sqrt{\tan(dx+c)}) + \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4} \right)$

```
input int(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 1/d*(2/7*B*b^2*tan(d*x+c)^(7/2)+2/5*A*b^2*tan(d*x+c)^(5/2)+4/5*B*a*b*tan(d
*x+c)^(5/2)+4/3*A*a*b*tan(d*x+c)^(3/2)+2/3*B*a^2*tan(d*x+c)^(3/2)-2/3*B*b^
2*tan(d*x+c)^(3/2)+2*A*a^2*tan(d*x+c)^(1/2)-2*tan(d*x+c)^(1/2)*A*b^2-4*tan
(d*x+c)^(1/2)*B*a*b+1/4*(-A*a^2+A*b^2+2*B*a*b)*2^(1/2)*(ln((1+2^(1/2)*tan(
d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan
(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(-
2*A*a*b-B*a^2+B*b^2)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(
1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2
))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))
```

3.386.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4326 vs. 2(318) = 636.

Time = 0.71 (sec) , antiderivative size = 4326, normalized size of antiderivative = 12.02

$$\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm
m="fricas")
```

```

output 1/210*(105*d*sqrt(-(2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)
*a^3*b - 4*(A^2 - B^2)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16
*(A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3
*B - A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3
*B - A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B -
A*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2)*log(((B*a^2 + 2*A*
a*b - B*b^2)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a
^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^
3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^
5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A
^4 - 2*A^2*B^2 + B^4)*b^8)/d^4) + ((A^3 - A*B^2)*a^6 - 2*(5*A^2*B - B^3)*a
^5*b - (7*A^3 - 23*A*B^2)*a^4*b^2 + 4*(7*A^2*B - 3*B^3)*a^3*b^3 + (7*A^3 -
23*A*B^2)*a^2*b^4 - 2*(5*A^2*B - B^3)*a*b^5 - (A^3 - A*B^2)*b^6)*d)*sqrt(
-(2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 -
B^2)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a
^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^
3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^
5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A
^4 - 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2) - ((A^4 - B^4)*a^8 - 8*(A^3*B + A*B^
3)*a^7*b - 4*(A^4 - B^4)*a^6*b^2 - 8*(A^3*B + A*B^3)*a^5*b^3 - 10*(A^4 ...

```

3.386.6 Sympy [F]

$$\begin{aligned}
 & \int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 &= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^2 \tan^{\frac{3}{2}}(c + dx) dx
 \end{aligned}$$

```

input integrate(tan(d*x+c)**(3/2)*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)

```

```

output Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**2*tan(c + d*x)**(3/2),
x)

```

3.386.7 Maxima [A] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 302, normalized size of antiderivative = 0.84

$$\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{120 B b^2 \tan(dx+c)^{\frac{7}{2}} + 168 (2 B a b + A b^2) \tan(dx+c)^{\frac{5}{2}} - 210 \sqrt{2}((A+B)a^2 + 2(A-B)ab - (A+B))}{d}$$

```
input integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")
```

```
output 1/420*(120*B*b^2*tan(d*x + c)^(7/2) + 168*(2*B*a*b + A*b^2)*tan(d*x + c)^(5/2) - 210*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) - 210*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - 105*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 105*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 280*(B*a^2 + 2*A*a*b - B*b^2)*tan(d*x + c)^(3/2) + 840*(A*a^2 - 2*B*a*b - A*b^2)*sqrt(tan(d*x + c)))/d
```

3.386.8 Giac [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx = \text{Timed out}$$

```
input integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")
```

```
output Timed out
```


3.386.9 Mupad [B] (verification not implemented)

Time = 21.13 (sec) , antiderivative size = 3869, normalized size of antiderivative = 10.75

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)
```

```
output atan((B^2*a^4*tan(c + d*x)^(1/2)*((12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4) - (B^2*a*b^3)/d^2 + (B^2*a^3*b)/d^2)^(1/2)*32i)/((16*B^3*b^6)/d - (16*B^3*a^6)/d - (112*B^3*a^2*b^4)/d + (112*B^3*a^4*b^2)/d + (32*B*a*b*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2))/d^3) + (B^2*b^4*tan(c + d*x)^(1/2)*((12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4) - (B^2*a*b^3)/d^2 + (B^2*a^3*b)/d^2)^(1/2)*32i)/((16*B^3*b^6)/d - (16*B^3*a^6)/d - (112*B^3*a^2*b^4)/d + (112*B^3*a^4*b^2)/d + (32*B*a*b*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2))/d^3) - (B^2*a^2*b^2*tan(c + d*x)^(1/2)*((12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4) - (B^2*a*b^3)/d^2 + (B^2*a^3*b)/d^2)^(1/2)*192i)/((16*B^3*b^6)/d - (16*B^3*a^6)/d - (112*B^3*a^2*b^4)/d + (112*B^3*a^4*b^2)/d + (32*B*a*b*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2))/d^3))*((12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4) - (B^2*a*b^3)/d^2 + (B^2*a^3*b)/d^2)^(1/2)*2i - atan((B^2*a^4*tan(c + d*x)^(1/2)*((B^2*a^3*b)/d^2 - (B^2*a*b^3)/d^2 - (12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2)*32i)...
```

3.387 $\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.387.1 Optimal result	3675
3.387.2 Mathematica [C] (verified)	3676
3.387.3 Rubi [A] (verified)	3676
3.387.4 Maple [A] (verified)	3682
3.387.5 Fricas [B] (verification not implemented)	3683
3.387.6 Sympy [F]	3684
3.387.7 Maxima [A] (verification not implemented)	3684
3.387.8 Giac [F(-1)]	3685
3.387.9 Mupad [B] (verification not implemented)	3685

3.387.1 Optimal result

Integrand size = 33, antiderivative size = 326

$$\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= -\frac{(a^2(A - B) - b^2(A - B) - 2ab(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(a^2(A - B) - b^2(A - B) - 2ab(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(2ab(A - B) + a^2(A + B) - b^2(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(2ab(A - B) + a^2(A + B) - b^2(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{2(2aAb + a^2B - b^2B) \sqrt{\tan(c + dx)}}{d} + \frac{2b(5Ab + 7aB) \tan^{\frac{3}{2}}(c + dx)}{15d}$$

$$+ \frac{2bB \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}{5d}$$

output $\frac{1}{2}(a^2(A-B)-b^2(A-B)-2ab(A+B))\arctan(-1+2^{1/2}\tan(dx+c)^{1/2})/d+2^{1/2}+1/2(a^2(A-B)-b^2(A-B)-2ab(A+B))\arctan(1+2^{1/2}\tan(dx+c)^{1/2})/d+1/4(2ab(A-B)+a^2(A+B)-b^2(A+B))\ln(1-2^{1/2}\tan(dx+c)^{1/2}+\tan(dx+c))/d-1/4(2ab(A-B)+a^2(A+B)-b^2(A+B))\ln(1+2^{1/2}\tan(dx+c)^{1/2}+\tan(dx+c))/d+2(2Aab+B^2a^2-Bb^2)\tan(dx+c)^{1/2}/d+2/15b(5Ab+7Ba)\tan(dx+c)^{3/2}/d+2/5bB\tan(dx+c)^{3/2}(a+b\tan(dx+c))/d$

3.387.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.31 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.46

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2(A+B\tan(c+dx)) dx$$

$$= \frac{15\sqrt[4]{-1}(a-ib)^2(iA+B)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right) - 15(-1)^{3/4}(a+ib)^2(A+iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{d}$$

input `Integrate[Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output $(15(-1)^{1/4}(a - I*b)^2(I*A + B)*\operatorname{ArcTan}[(-1)^{3/4}\sqrt{\tan[c + d*x]}] - 15(-1)^{3/4}(a + I*b)^2(A + I*B)*\operatorname{ArcTanh}[(-1)^{3/4}\sqrt{\tan[c + d*x]}] + 2\sqrt{\tan[c + d*x]}*(15(2aA*b + a^2*B - b^2*B) + 5b*(A*b + 2a*B)*\tan[c + d*x] + 3b^2*B*\tan[c + d*x]^2))/(15*d)$

3.387.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 283, normalized size of antiderivative = 0.87, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 4090, 27, 3042, 4113, 3042, 4011, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2(A+B\tan(c+dx)) dx$$

3.387. $\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2(A+B\tan(c+dx)) dx$

$$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^2(A+B \tan(c+dx))dx$$

↓ 3042

$$\frac{2}{5} \int \frac{1}{2} \sqrt{\tan(c+dx)}(b(5Ab+7aB) \tan^2(c+dx) + 5(Ba^2+2Aba-b^2B) \tan(c+dx) + a(5aA-3bB)) dx + \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))}{5d}$$

↓ 4090

$$\frac{1}{5} \int \sqrt{\tan(c+dx)}(b(5Ab+7aB) \tan^2(c+dx) + 5(Ba^2+2Aba-b^2B) \tan(c+dx) + a(5aA-3bB)) dx + \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))}{5d}$$

↓ 27

$$\frac{1}{5} \int \sqrt{\tan(c+dx)}(b(5Ab+7aB) \tan^2(c+dx) + 5(Ba^2+2Aba-b^2B) \tan(c+dx) + a(5aA-3bB)) dx + \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))}{5d}$$

↓ 3042

$$\frac{1}{5} \int \sqrt{\tan(c+dx)}(b(5Ab+7aB) \tan(c+dx)^2 + 5(Ba^2+2Aba-b^2B) \tan(c+dx) + a(5aA-3bB)) dx + \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))}{5d}$$

↓ 4113

$$\frac{1}{5} \left(\int \sqrt{\tan(c+dx)}(5(Aa^2-2bBa-Ab^2) + 5(Ba^2+2Aba-b^2B) \tan(c+dx)) dx + \frac{2b(7aB+5Ab) \tan^{\frac{3}{2}}(c+dx)}{3d} \right) + \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))}{5d}$$

↓ 3042

$$\frac{1}{5} \left(\int \sqrt{\tan(c+dx)}(5(Aa^2-2bBa-Ab^2) + 5(Ba^2+2Aba-b^2B) \tan(c+dx)) dx + \frac{2b(7aB+5Ab) \tan^{\frac{3}{2}}(c+dx)}{3d} \right) + \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))}{5d}$$

↓ 4011

$$\frac{1}{5} \left(\int \frac{5(Aa^2-2bBa-Ab^2) \tan(c+dx) - 5(Ba^2+2Aba-b^2B)}{\sqrt{\tan(c+dx)}} dx + \frac{10(a^2B+2aAb-b^2B)}{d} \sqrt{\tan(c+dx)} \right) + \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))}{5d}$$

3.387. $\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

↓ 3042

$$\frac{1}{5} \left(\int \frac{5(Aa^2 - 2bBa - Ab^2) \tan(c + dx) - 5(Ba^2 + 2Aba - b^2B)}{\sqrt{\tan(c + dx)}} dx + \frac{10(a^2B + 2aAb - b^2B) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}{5d} \right)$$

↓ 4017

$$\frac{1}{5} \left(\frac{2 \int -\frac{5(Ba^2 + 2Aba - b^2B - (Aa^2 - 2bBa - Ab^2) \tan(c + dx))}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)}}{d} + \frac{10(a^2B + 2aAb - b^2B) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}{5d} \right)$$

↓ 27

$$\frac{1}{5} \left(-\frac{10 \int \frac{Ba^2 + 2Aba - b^2B - (Aa^2 - 2bBa - Ab^2) \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)}}{d} + \frac{10(a^2B + 2aAb - b^2B) \sqrt{\tan(c + dx)}}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}{5d} \right)$$

↓ 1482

$$\frac{1}{5} \left(-\frac{10 \left(\frac{1}{2}(a^2(A + B) + 2ab(A - B) - b^2(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} - \frac{1}{2}(a^2(A - B) - 2ab(A + B)) \right)}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}{5d} \right)$$

↓ 1476

$$\frac{1}{5} \left(-\frac{10 \left(\frac{1}{2}(a^2(A + B) + 2ab(A - B) - b^2(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} - \frac{1}{2}(a^2(A - B) - 2ab(A + B)) \right)}{d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}{5d} \right)$$

↓ 1082

3.387. $\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^2(A-B) - 2ab(A+B)) \int \frac{1+\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} \right. \\ \left. \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))}{5d} \right. \\ \left. \downarrow 217 \right.$$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^2(A-B) - 2ab(A+B)) \int \frac{1+\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} \right. \\ \left. \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))}{5d} \right. \\ \left. \downarrow 1479 \right.$$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{d} \right)}{d} \right. \\ \left. \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))}{5d} \right. \\ \left. \downarrow 25 \right.$$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{d} \right)}{d} \right. \\ \left. \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))}{5d} \right. \\ \left. \downarrow 27 \right.$$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}} \right) \right)}{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} \right) \frac{5d}{5d} \downarrow 1103$$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} \right) \frac{5d}{5d}$$

input `Int[Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(2*b*B*Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x]))/(5*d) + ((-10*(-1/2*((a^2*(A - B) - b^2*(A - B) - 2*a*b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])) + ((2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d + (10*(2*a*A*b + a^2*B - b^2*B)*Sqrt[Tan[c + d*x]])/d + (2*b*(5*A*b + 7*a*B)*Tan[c + d*x]^(3/2))/(3*d))/5`

3.387.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

3.387. $\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`


```
rule 4017 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_
)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sq
rt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] &
& NeQ[c^2 + d^2, 0]
```

```
rule 4090 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4113 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_)
+ (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

3.387.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 292, normalized size of antiderivative = 0.90

method	result
derivativedivides	$\frac{2B b^2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{2A b^2 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} + \frac{4Bab \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} + 4Aab(\sqrt{\tan(dx+c)}) + 2B a^2(\sqrt{\tan(dx+c)}) - 2(\sqrt{\tan(dx+c)})$
default	$\frac{2B b^2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{2A b^2 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} + \frac{4Bab \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} + 4Aab(\sqrt{\tan(dx+c)}) + 2B a^2(\sqrt{\tan(dx+c)}) - 2(\sqrt{\tan(dx+c)})$
parts	$\frac{(A b^2 + 2Bab) \left(\frac{2 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} - \frac{\sqrt{2} \left(\ln \left(\frac{1 - \sqrt{2} (\sqrt{\tan(dx+c)}) + \tan(dx+c)}{1 + \sqrt{2} (\sqrt{\tan(dx+c)}) + \tan(dx+c)} \right) + 2 \arctan(1 + \sqrt{2} (\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1 + \sqrt{2} (\sqrt{\tan(dx+c)}) \right)}{4} \right)}{d}$

```
input int(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)), x, method=_RETURNV
ERBOSE)
```

$$3.387. \quad \int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

output $1/d*(2/5*B*b^2*\tan(d*x+c)^(5/2)+2/3*A*b^2*\tan(d*x+c)^(3/2)+4/3*B*a*b*\tan(d*x+c)^(3/2)+4*A*a*b*\tan(d*x+c)^(1/2)+2*B*a^2*\tan(d*x+c)^(1/2)-2*\tan(d*x+c)^(1/2)*B*b^2+1/4*(-2*A*a*b-B*a^2+B*b^2)*2^(1/2)*(ln((1+2^(1/2)*\tan(d*x+c)^(1/2)+\tan(d*x+c))/(1-2^(1/2)*\tan(d*x+c)^(1/2)+\tan(d*x+c)))+2*\arctan(1+2^(1/2)*\tan(d*x+c)^(1/2))+2*\arctan(-1+2^(1/2)*\tan(d*x+c)^(1/2)))+1/4*(A*a^2-A*b^2-2*B*a*b)*2^(1/2)*(ln((1-2^(1/2)*\tan(d*x+c)^(1/2)+\tan(d*x+c))/(1+2^(1/2)*\tan(d*x+c)^(1/2)+\tan(d*x+c)))+2*\arctan(1+2^(1/2)*\tan(d*x+c)^(1/2))+2*\arctan(-1+2^(1/2)*\tan(d*x+c)^(1/2)))$

3.387.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4283 vs. $2(288) = 576$.

Time = 0.73 (sec) , antiderivative size = 4283, normalized size of antiderivative = 13.14

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output $-1/30*(15*d*\sqrt{(2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2)*a*b^3 + d^2*\sqrt{-(A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8}/d^4)/d^2)*\log(((A*a^2 - 2*B*a*b - A*b^2)*d^3*\sqrt{-(A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8}/d^4) - ((A^2*B - B^3)*a^6 + 2*(A^3 - 5*A*B^2)*a^5*b - (23*A^2*B - 7*B^3)*a^4*b^2 - 4*(3*A^3 - 7*A*B^2)*a^3*b^3 + (23*A^2*B - 7*B^3)*a^2*b^4 + 2*(A^3 - 5*A*B^2)*a*b^5 - (A^2*B - B^3)*b^6)*d)*\sqrt{(2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2)*a*b^3 + d^2*\sqrt{-(A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8}/d^4)/d^2) - ((A^4 - B^4)*a^8 - 8*(A^3*B + A*B^3)*a^7*b - 4*(A^4 - B^4)*a^6*b^2 - 8*(A^3*B + A*B^3)*a^5*b^3 - 10*(A^4 - B...$

3.387.6 Sympy [F]

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

$$= \int (A+B\tan(c+dx))(a+b\tan(c+dx))^2\sqrt{\tan(c+dx)}dx$$

input `integrate(tan(d*x+c)**(1/2)*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**2*sqrt(tan(c + d*x)), x)`

3.387.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 275, normalized size of antiderivative = 0.84

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

$$= \frac{24Bb^2\tan(dx+c)^{\frac{5}{2}} + 30\sqrt{2}((A-B)a^2 - 2(A+B)ab - (A-B)b^2)\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(dx+c)}\right)\right)}{d}$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/60*(24*B*b^2*tan(d*x + c)^(5/2) + 30*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 30*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - 15*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 15*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 40*(2*B*a*b + A*b^2)*tan(d*x + c)^(3/2) + 120*(B*a^2 + 2*A*a*b - B*b^2)*sqrt(tan(d*x + c)))/d`

3.387.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \text{Timed out}$$

```
input integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")
```

output Timed out

3.387.9 Mupad [B] (verification not implemented)

Time = 15.10 (sec) , antiderivative size = 3825, normalized size of antiderivative = 11.73

$$\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)
```

```
output atan((A^2*a^4*tan(c + d*x)^(1/2)*((A^2*a^3*b)/d^2 - (A^2*a*b^3)/d^2 - (12*A^4*a^2*b^6*d^4 - A^4*b^8*d^4 - A^4*a^8*d^4 - 38*A^4*a^4*b^4*d^4 + 12*A^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2)*32i)/((16*A^3*a^6)/d - (16*A^3*b^6)/d + (112*A^3*a^2*b^4)/d - (112*A^3*a^4*b^2)/d + (32*A*a*b*(12*A^4*a^2*b^6*d^4 - A^4*b^8*d^4 - A^4*a^8*d^4 - 38*A^4*a^4*b^4*d^4 + 12*A^4*a^6*b^2*d^4)^(1/2))/d^3) + (A^2*b^4*tan(c + d*x)^(1/2)*((A^2*a^3*b)/d^2 - (A^2*a*b^3)/d^2 - (12*A^4*a^2*b^6*d^4 - A^4*b^8*d^4 - A^4*a^8*d^4 - 38*A^4*a^4*b^4*d^4 + 12*A^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2)*32i)/((16*A^3*a^6)/d - (16*A^3*b^6)/d + (112*A^3*a^2*b^4)/d - (112*A^3*a^4*b^2)/d + (32*A*a*b*(12*A^4*a^2*b^6*d^4 - A^4*b^8*d^4 - A^4*a^8*d^4 - 38*A^4*a^4*b^4*d^4 + 12*A^4*a^6*b^2*d^4)^(1/2))/d^3) - (A^2*a^2*b^2*tan(c + d*x)^(1/2)*((A^2*a^3*b)/d^2 - (A^2*a*b^3)/d^2 - (12*A^4*a^2*b^6*d^4 - A^4*b^8*d^4 - A^4*a^8*d^4 - 38*A^4*a^4*b^4*d^4 + 12*A^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2)*192i)/((16*A^3*a^6)/d - (16*A^3*b^6)/d + (112*A^3*a^2*b^4)/d - (112*A^3*a^4*b^2)/d + (32*A*a*b*(12*A^4*a^2*b^6*d^4 - A^4*b^8*d^4 - A^4*a^8*d^4 - 38*A^4*a^4*b^4*d^4 + 12*A^4*a^6*b^2*d^4)^(1/2))/d^3))*((A^2*a^3*b)/d^2 - (A^2*a*b^3)/d^2 - (12*A^4*a^2*b^6*d^4 - A^4*b^8*d^4 - A^4*a^8*d^4 - 38*A^4*a^4*b^4*d^4 + 12*A^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2)*2i - atan((B^2*a^4*tan(c + d*x)^(1/2)*((12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4) + (B^2*a*b^3)/d^2 - (B^2*a^3*b)/d^2)^(1/2)*32i)...
```

3.388
$$\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.388.1 Optimal result 3686
 3.388.2 Mathematica [C] (verified) 3687
 3.388.3 Rubi [A] (verified) 3687
 3.388.4 Maple [A] (verified) 3692
 3.388.5 Fricas [B] (verification not implemented) 3693
 3.388.6 Sympy [F] 3694
 3.388.7 Maxima [A] (verification not implemented) 3695
 3.388.8 Giac [F(-1)] 3695
 3.388.9 Mupad [B] (verification not implemented) 3696

3.388.1 Optimal result

Integrand size = 33, antiderivative size = 294

$$\int \frac{(a + b \tan(c + dx))^2(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= -\frac{(2ab(A - B) + a^2(A + B) - b^2(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(2ab(A - B) + a^2(A + B) - b^2(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(a^2(A - B) - b^2(A - B) - 2ab(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{(a^2(A - B) - b^2(A - B) - 2ab(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{2b(3Ab + 5aB)\sqrt{\tan(c + dx)}}{3d} + \frac{2bB\sqrt{\tan(c + dx)}(a + b \tan(c + dx))}{3d}$$

output

```
1/2*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/
d*2^(1/2)+1/2*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*arctan(1+2^(1/2)*tan(d*x+c
)^(1/2))/d*2^(1/2)-1/4*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*ln(1-2^(1/2)*tan(
d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)+1/4*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*l
n(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)+2/3*b*(3*A*b+5*B*a)*tan
(d*x+c)^(1/2)/d+2/3*b*B*tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))/d
```

3.388.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.58 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.40

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \frac{-3\sqrt[4]{-1}(a - ib)^2(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right) - 3\sqrt[4]{-1}(a + ib)^2(A + iB) \operatorname{arctanh}\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)}{3d}$$

input `Integrate[((a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `(-3*(-1)^(1/4)*(a - I*b)^2*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] - 3*(-1)^(1/4)*(a + I*b)^2*(A + I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + 2*b*Sqrt[Tan[c + d*x]]*(3*A*b + 6*a*B + b*B*Tan[c + d*x]))/(3*d)`

3.388.3 Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 249, normalized size of antiderivative = 0.85, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.485$, Rules used = {3042, 4090, 27, 3042, 4113, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$\downarrow 4090$$

$$\frac{2}{3} \int \frac{b(3Ab + 5aB) \tan^2(c + dx) + 3(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(3aA - bB)}{2\sqrt{\tan(c + dx)}} dx + \frac{2bB\sqrt{\tan(c + dx)}(a + b \tan(c + dx))}{3d}$$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{1}{3} \int \frac{b(3Ab + 5aB) \tan^2(c + dx) + 3(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(3aA - bB)}{\sqrt{\tan(c + dx)} \frac{2bB \sqrt{\tan(c + dx)}(a + b \tan(c + dx))}{3d}} dx + \\
& \downarrow 3042 \\
& \frac{1}{3} \int \frac{b(3Ab + 5aB) \tan(c + dx)^2 + 3(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(3aA - bB)}{\sqrt{\tan(c + dx)} \frac{2bB \sqrt{\tan(c + dx)}(a + b \tan(c + dx))}{3d}} dx + \\
& \downarrow 4113 \\
& \frac{1}{3} \left(\int \frac{3(Aa^2 - 2bBa - Ab^2) + 3(Ba^2 + 2Aba - b^2B) \tan(c + dx)}{\sqrt{\tan(c + dx)} \frac{2bB \sqrt{\tan(c + dx)}(a + b \tan(c + dx))}{3d}} dx + \frac{2b(5aB + 3Ab) \sqrt{\tan(c + dx)}}{d} \right) + \\
& \downarrow 3042 \\
& \frac{1}{3} \left(\int \frac{3(Aa^2 - 2bBa - Ab^2) + 3(Ba^2 + 2Aba - b^2B) \tan(c + dx)}{\sqrt{\tan(c + dx)} \frac{2bB \sqrt{\tan(c + dx)}(a + b \tan(c + dx))}{3d}} dx + \frac{2b(5aB + 3Ab) \sqrt{\tan(c + dx)}}{d} \right) + \\
& \downarrow 4017 \\
& \frac{1}{3} \left(\frac{2 \int \frac{3(Aa^2 - 2bBa - Ab^2) + (Ba^2 + 2Aba - b^2B) \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)}}{d} + \frac{2b(5aB + 3Ab) \sqrt{\tan(c + dx)}}{d} \right) + \\
& \frac{2bB \sqrt{\tan(c + dx)}(a + b \tan(c + dx))}{3d} \\
& \downarrow 27 \\
& \frac{1}{3} \left(\frac{6 \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)}}{d} + \frac{2b(5aB + 3Ab) \sqrt{\tan(c + dx)}}{d} \right) + \\
& \frac{2bB \sqrt{\tan(c + dx)}(a + b \tan(c + dx))}{3d} \\
& \downarrow 1482
\end{aligned}$$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^2(A + B) + 2ab(A - B) - b^2(A - B)) \int \frac{1 + \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} \right. \\ \left. \frac{2bB \sqrt{\tan(c+dx)}(a + b \tan(c+dx))}{3d} \right) \\ \downarrow 1476$$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^2(A + B) + 2ab(A - B) - b^2(A - B)) \int \frac{1 + \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} \right. \\ \left. \frac{2bB \sqrt{\tan(c+dx)}(a + b \tan(c+dx))}{3d} \right) \\ \downarrow 1082$$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^2(A + B) + 2ab(A - B) - b^2(A - B)) \int \frac{1 + \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} \right. \\ \left. \frac{2bB \sqrt{\tan(c+dx)}(a + b \tan(c+dx))}{3d} \right) \\ \downarrow 217$$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^2(A + B) + 2ab(A - B) - b^2(A - B)) \int \frac{1 + \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} \right. \\ \left. \frac{2bB \sqrt{\tan(c+dx)}(a + b \tan(c+dx))}{3d} \right) \\ \downarrow 1479$$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \left(- \frac{\int - \frac{\sqrt{2} - 2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int - \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1} + \sqrt{\tan(c+dx)})}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1}}{2\sqrt{2}} \right)}{d} \right. \\ \left. \frac{2bB \sqrt{\tan(c+dx)}(a + b \tan(c+dx))}{3d} \right)$$

$$\begin{aligned}
& \downarrow 25 \\
& \frac{1}{3} \left(\frac{6 \left(\frac{1}{2} (a^2(A-B) - 2ab(A+B) - b^2(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right)}{2bB\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} \right)}{3d} \\
& \downarrow 27 \\
& \frac{1}{3} \left(\frac{6 \left(\frac{1}{2} (a^2(A-B) - 2ab(A+B) - b^2(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right)}{2bB\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} \right)}{3d} \\
& \downarrow 1103 \\
& \frac{1}{3} \left(\frac{6 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} (a^2(A-B) - 2ab(A+B) - b^2(A-B)) \right)}{2bB\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} \right)}{3d}
\end{aligned}$$

input `Int[((a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `((6*(((2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])))/2 + ((a^2*(A - B) - b^2*(A - B) - 2*a*b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d + (2*b*(3*A*b + 5*a*B)*Sqrt[Tan[c + d*x]]/d)/3 + (2*b*B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x]))/(3*d)`

3.388.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4090 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

3.388.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 251, normalized size of antiderivative = 0.85

method	result
derivativedivides	$\frac{2Bb^2 \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + 2(\sqrt{\tan(dx+c)}Ab^2 + 4(\sqrt{\tan(dx+c)}Bab + \frac{(Aa^2 - Ab^2 - 2Bab)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))} \right) \right)}{4}$
default	$\frac{2Bb^2 \left(\tan^{\frac{3}{2}}(dx+c) \right)}{3} + 2(\sqrt{\tan(dx+c)}Ab^2 + 4(\sqrt{\tan(dx+c)}Bab + \frac{(Aa^2 - Ab^2 - 2Bab)\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))} \right) \right)}{4}$
parts	$\frac{(Ab^2 + 2Bab) \left(2(\sqrt{\tan(dx+c)}) - \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)))} + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)))} \right)}{4} \right)}{d}$

3.388. $\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

```
input int((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)
```

```
output 1/d*(2/3*B*b^2*tan(d*x+c)^(3/2)+2*tan(d*x+c)^(1/2)*A*b^2+4*tan(d*x+c)^(1/2)
)*B*a*b+1/4*(A*a^2-A*b^2-2*B*a*b)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+
tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*ta
n(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(2*A*a*b+B*a^2-
B*b^2)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(
d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1
+2^(1/2)*tan(d*x+c)^(1/2)))
```

3.388.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4271 vs. 2(258) = 516.

Time = 0.72 (sec) , antiderivative size = 4271, normalized size of antiderivative = 14.53

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorith
m="fricas")
```

```

output -1/6*(3*d*sqrt(-(2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^
3*b - 4*(A^2 - B^2)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A
^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B
- A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B
- A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*
B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2)*log(((B*a^2 + 2*A*a*b
- B*b^2)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*
b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 +
2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 -
4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4
- 2*A^2*B^2 + B^4)*b^8)/d^4) + ((A^3 - A*B^2)*a^6 - 2*(5*A^2*B - B^3)*a^5*
b - (7*A^3 - 23*A*B^2)*a^4*b^2 + 4*(7*A^2*B - 3*B^3)*a^3*b^3 + (7*A^3 - 23
*A*B^2)*a^2*b^4 - 2*(5*A^2*B - B^3)*a*b^5 - (A^3 - A*B^2)*b^6)*d)*sqrt(-(2
*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2
)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*
b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 +
2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 -
4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4
- 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2) - ((A^4 - B^4)*a^8 - 8*(A^3*B + A*B^3)*
a^7*b - 4*(A^4 - B^4)*a^6*b^2 - 8*(A^3*B + A*B^3)*a^5*b^3 - 10*(A^4 - B...

```

3.388.6 Sympy [F]

$$\begin{aligned}
 & \int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
 &= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^2}{\sqrt{\tan(c + dx)}} dx
 \end{aligned}$$

```

input integrate((a+b*tan(d*x+c))**2*(A+B*tan(d*x+c))/tan(d*x+c)**(1/2),x)

```

```

output Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**2/sqrt(tan(c + d*x)),
x)

```

3.388.7 Maxima [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 248, normalized size of antiderivative = 0.84

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \frac{8 B b^2 \tan(dx + c)^{\frac{3}{2}} + 6 \sqrt{2} ((A + B) a^2 + 2 (A - B) a b - (A + B) b^2) \arctan\left(\frac{1}{2} \sqrt{2} (\sqrt{2} + 2 \sqrt{\tan(dx + c)})\right)}{d}$$

input `integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="maxima")`

output `1/12*(8*B*b^2*tan(d*x + c)^(3/2) + 6*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 6*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + 3*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - 3*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 24*(2*B*a*b + A*b^2)*sqrt(tan(d*x + c))/d`

3.388.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="giac")`

output `Timed out`

3.388.9 Mupad [B] (verification not implemented)

Time = 11.19 (sec) , antiderivative size = 3773, normalized size of antiderivative = 12.83

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Too large to display}$$

```
input int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2)/tan(c + d*x)^(1/2),x)
```

```
output atan((B^2*a^4*tan(c + d*x)^(1/2)*((B^2*a^3*b)/d^2 - (B^2*a*b^3)/d^2 - (12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2)*32i)/((16*B^3*a^6)/d - (16*B^3*b^6)/d + (112*B^3*a^2*b^4)/d - (112*B^3*a^4*b^2)/d + (32*B*a*b*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2))/d^3) + (B^2*b^4*tan(c + d*x)^(1/2)*((B^2*a^3*b)/d^2 - (B^2*a*b^3)/d^2 - (12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2)*32i)/((16*B^3*a^6)/d - (16*B^3*b^6)/d + (112*B^3*a^2*b^4)/d - (112*B^3*a^4*b^2)/d + (32*B*a*b*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2))/d^3) - (B^2*a^2*b^2*tan(c + d*x)^(1/2)*((B^2*a^3*b)/d^2 - (B^2*a*b^3)/d^2 - (12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2)*192i)/((16*B^3*a^6)/d - (16*B^3*b^6)/d + (112*B^3*a^2*b^4)/d - (112*B^3*a^4*b^2)/d + (32*B*a*b*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2))/d^3))*((B^2*a^3*b)/d^2 - (B^2*a*b^3)/d^2 - (12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2)*2i - atan((B^2*a^4*tan(c + d*x)^(1/2)*((12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4) - (B^2*a*b^3)/d^2 + (B^2*a^3*b)/d^2)^(1/2)*32i)...
```

3.389
$$\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.389.1 Optimal result 3697
 3.389.2 Mathematica [C] (verified) 3698
 3.389.3 Rubi [A] (verified) 3698
 3.389.4 Maple [A] (verified) 3703
 3.389.5 Fricas [B] (verification not implemented) 3703
 3.389.6 Sympy [F] 3704
 3.389.7 Maxima [A] (verification not implemented) 3705
 3.389.8 Giac [F(-1)] 3705
 3.389.9 Mupad [B] (verification not implemented) 3706

3.389.1 Optimal result

Integrand size = 33, antiderivative size = 276

$$\int \frac{(a + b \tan(c + dx))^2(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{(a^2(A - B) - b^2(A - B) - 2ab(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(a^2(A - B) - b^2(A - B) - 2ab(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(2ab(A - B) + a^2(A + B) - b^2(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{(2ab(A - B) + a^2(A + B) - b^2(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$- \frac{2a^2A}{d\sqrt{\tan(c + dx)}} + \frac{2b^2B\sqrt{\tan(c + dx)}}{d}$$

output

```
-1/2*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))
/d*2^(1/2)-1/2*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*arctan(1+2^(1/2)*tan(d*x+
c)^(1/2))/d*2^(1/2)-1/4*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*ln(1-2^(1/2)*tan
(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)+1/4*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*
ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)-2*a^2*A/d/tan(d*x+c)^(
1/2)+2*b^2*B*tan(d*x+c)^(1/2)/d
```


3.389.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.99 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.76

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{-8b(Ab + 3aB) - 8(a^2A - Ab^2 - 2abB) \operatorname{Hypergeometric2F1}\left(-\frac{1}{4}, 1, \frac{3}{4}, -\tan^2(c + dx)\right) - \sqrt{2}(2aAb + a^2)}{d}$$

input `Integrate[((a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `(-8*b*(A*b + 3*a*B) - 8*(a^2*A - A*b^2 - 2*a*b*B)*Hypergeometric2F1[-1/4, 1, 3/4, -Tan[c + d*x]^2] - Sqrt[2]*(2*a*A*b + a^2*B - b^2*B)*(2*ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - 2*ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]) + Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]])*Sqrt[Tan[c + d*x]] + 8*b*B*(a + b*Tan[c + d*x]))/(4*d*Sqrt[Tan[c + d*x]])`

3.389.3 Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.83, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3042, 4087, 3042, 4113, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx$$

$$\downarrow \text{4087}$$

$$\int \frac{b^2 B \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)}{\sqrt{\tan(c + dx)}} dx - \frac{2a^2 A}{d \sqrt{\tan(c + dx)}}$$

3.389. $\int \frac{(a+b \tan(c+dx))^2 (A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \int \frac{b^2 B \tan(c+dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2Ab + aB)}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2 A}{d\sqrt{\tan(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{Ba^2 + 2Aba - b^2 B - (Aa^2 - 2bBa - Ab^2) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2 A}{d\sqrt{\tan(c+dx)}} + \\
 & \quad \frac{2b^2 B \sqrt{\tan(c+dx)}}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{Ba^2 + 2Aba - b^2 B - (Aa^2 - 2bBa - Ab^2) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2 A}{d\sqrt{\tan(c+dx)}} + \\
 & \quad \frac{2b^2 B \sqrt{\tan(c+dx)}}{d} \\
 & \quad \downarrow \text{4017} \\
 & \frac{2 \int \frac{Ba^2 + 2Aba - b^2 B - (Aa^2 - 2bBa - Ab^2) \tan(c+dx)}{\tan^2(c+dx) + 1} d\sqrt{\tan(c+dx)}}{d} - \frac{2a^2 A}{d\sqrt{\tan(c+dx)}} + \frac{2b^2 B \sqrt{\tan(c+dx)}}{d} \\
 & \quad \downarrow \text{1482} \\
 & \frac{2 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx) + 1} d\sqrt{\tan(c+dx)} - \frac{1}{2} (a^2(A-B) - 2ab(A+B) - b^2(A+B)) \right)}{d} \\
 & \quad + \frac{2a^2 A}{d\sqrt{\tan(c+dx)}} + \frac{2b^2 B \sqrt{\tan(c+dx)}}{d} \\
 & \quad \downarrow \text{1476} \\
 & \frac{2 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx) + 1} d\sqrt{\tan(c+dx)} - \frac{1}{2} (a^2(A-B) - 2ab(A+B) - b^2(A+B)) \right)}{d} \\
 & \quad + \frac{2a^2 A}{d\sqrt{\tan(c+dx)}} + \frac{2b^2 B \sqrt{\tan(c+dx)}}{d} \\
 & \quad \downarrow \text{1082} \\
 & \frac{2 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx) + 1} d\sqrt{\tan(c+dx)} - \frac{1}{2} (a^2(A-B) - 2ab(A+B) - b^2(A+B)) \right)}{d} \\
 & \quad + \frac{2a^2 A}{d\sqrt{\tan(c+dx)}} + \frac{2b^2 B \sqrt{\tan(c+dx)}}{d}
 \end{aligned}$$

3.389. $\int \frac{(a+b \tan(c+dx))^2 (A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

↓ 217

$$2 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2} (a^2(A-B) - 2ab(A+B) - b^2(A+B)) \int \frac{1 + \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) - \frac{2a^2A}{d\sqrt{\tan(c+dx)}} + \frac{2b^2B\sqrt{\tan(c+dx)}}{d}$$

↓ 1479

$$2 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(- \int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2} (a^2(A-B) - 2ab(A+B) - b^2(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right) - \frac{2a^2A}{d\sqrt{\tan(c+dx)}} + \frac{2b^2B\sqrt{\tan(c+dx)}}{d}$$

↓ 25

$$2 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2} (a^2(A-B) - 2ab(A+B) - b^2(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right) - \frac{2a^2A}{d\sqrt{\tan(c+dx)}} + \frac{2b^2B\sqrt{\tan(c+dx)}}{d}$$

↓ 27

$$2 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2} (a^2(A-B) - 2ab(A+B) - b^2(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right) - \frac{2a^2A}{d\sqrt{\tan(c+dx)}} + \frac{2b^2B\sqrt{\tan(c+dx)}}{d}$$

↓ 1103

$$2 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2} (a^2(A-B) - 2ab(A+B) - b^2(A+B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} + \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right) - \frac{2a^2A}{d\sqrt{\tan(c+dx)}} + \frac{2b^2B\sqrt{\tan(c+dx)}}{d}$$

input `Int[((a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `(2*(-1/2*((a^2*(A - B) - b^2*(A - B) - 2*a*b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])) + ((2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d - (2*a^2*A)/(d*Sqrt[Tan[c + d*x]]) + (2*b^2*B*Sqrt[Tan[c + d*x]])/d`

3.389.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4087 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^2*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (B*c - A*d))*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`

rule 4113 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]) + (C_)*tan[(e_) + (f_)*(x_)^2], x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

3.389.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 238, normalized size of antiderivative = 0.86

method	result
derivativedivides	$\frac{2(\sqrt{\tan(dx+c)})Bb^2 - \frac{2Aa^2}{\sqrt{\tan(dx+c)}} + \frac{(2Aab+B a^2 - B b^2)\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right)}{4}}$
default	$\frac{2(\sqrt{\tan(dx+c)})Bb^2 - \frac{2Aa^2}{\sqrt{\tan(dx+c)}} + \frac{(2Aab+B a^2 - B b^2)\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right)}{4}}$
parts	$\frac{(Ab^2+2Bab)\sqrt{2} \left(\ln\left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c)))}{4d}}$

```
input int((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x,method=_RETURNV
ERBOSE)
```

```
output 1/d*(2*tan(d*x+c)^(1/2)*B*b^2-2*A*a^2/tan(d*x+c)^(1/2)+1/4*(2*A*a*b+B*a^2-
B*b^2)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(
d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-
1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(-A*a^2+A*b^2+2*B*a*b)*2^(1/2)*(ln((1-2^(
1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+
2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)
))
```

3.389.5 Fracas [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 4280 vs. $2(244) = 488$.

Time = 0.74 (sec) , antiderivative size = 4280, normalized size of antiderivative = 15.51

$$\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorith
m="fricas")
```

output $\frac{1}{2}*(d*\sqrt{(2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2)*a*b^3 + d^2*\sqrt{-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2)*\log(((A*a^2 - 2*B*a*b - A*b^2)*d^3*\sqrt{-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4} - ((A^2*B - B^3)*a^6 + 2*(A^3 - 5*A*B^2)*a^5*b - (23*A^2*B - 7*B^3)*a^4*b^2 - 4*(3*A^3 - 7*A*B^2)*a^3*b^3 + (23*A^2*B - 7*B^3)*a^2*b^4 + 2*(A^3 - 5*A*B^2)*a*b^5 - (A^2*B - B^3)*b^6)*d)*\sqrt{(2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2)*a*b^3 + d^2*\sqrt{-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2) - ((A^4 - B^4)*a^8 - 8*(A^3*B + A*B^3)*a^7*b - 4*(A^4 - B^4)*a^6*b^2 - 8*(A^3*B + A*B^3)*a^5*b^3 - 10*(A^4 - B^4)*a...$

3.389.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^2}{\tan^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**2*(A+B*tan(d*x+c))/tan(d*x+c)**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**2/tan(c + d*x)**(3/2), x)`

3.389.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 240, normalized size of antiderivative = 0.87

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{8 B b^2 \sqrt{\tan(dx + c)} - 2 \sqrt{2} ((A - B) a^2 - 2 (A + B) a b - (A - B) b^2) \arctan\left(\frac{1}{2} \sqrt{2} (\sqrt{2} + 2 \sqrt{\tan(dx + c)})\right)}{d}$$

```
input integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm
m="maxima")
```

```
output 1/4*(8*B*b^2*sqrt(tan(d*x + c)) - 2*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b -
(A - B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) - 2*sqrt
(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt
(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A +
B)*b^2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((A
+ B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*log(-sqrt(2)*sqrt(tan(d*x + c)) +
tan(d*x + c) + 1) - 8*A*a^2/sqrt(tan(d*x + c)))/d
```

3.389.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm
m="giac")
```

```
output Timed out
```


3.389.9 Mupad [B] (verification not implemented)

Time = 10.69 (sec) , antiderivative size = 3749, normalized size of antiderivative = 13.58

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Too large to display}$$

```
input int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2)/tan(c + d*x)^(3/2),x)
```

```
output 2*atanh((32*A^2*a^4*d^3*tan(c + d*x)^(1/2)*((A^2*a^3*b)/d^2 - (A^2*a*b^3)/
d^2 - (12*A^4*a^2*b^6*d^4 - A^4*b^8*d^4 - A^4*a^8*d^4 - 38*A^4*a^4*b^4*d^4
+ 12*A^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2))/(16*A^3*a^6*d^2 - 16*A^3*b^6*
d^2 + 112*A^3*a^2*b^4*d^2 - 112*A^3*a^4*b^2*d^2 + 32*A*a*b*(12*A^4*a^2*b^6
*d^4 - A^4*b^8*d^4 - A^4*a^8*d^4 - 38*A^4*a^4*b^4*d^4 + 12*A^4*a^6*b^2*d^4
)^(1/2)) + (32*A^2*b^4*d^3*tan(c + d*x)^(1/2)*((A^2*a^3*b)/d^2 - (A^2*a*b^
3)/d^2 - (12*A^4*a^2*b^6*d^4 - A^4*b^8*d^4 - A^4*a^8*d^4 - 38*A^4*a^4*b^4*
d^4 + 12*A^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2))/(16*A^3*a^6*d^2 - 16*A^3*b
^6*d^2 + 112*A^3*a^2*b^4*d^2 - 112*A^3*a^4*b^2*d^2 + 32*A*a*b*(12*A^4*a^2*
b^6*d^4 - A^4*b^8*d^4 - A^4*a^8*d^4 - 38*A^4*a^4*b^4*d^4 + 12*A^4*a^6*b^2*
d^4)^(1/2)) - (192*A^2*a^2*b^2*d^3*tan(c + d*x)^(1/2)*((A^2*a^3*b)/d^2 - (
A^2*a*b^3)/d^2 - (12*A^4*a^2*b^6*d^4 - A^4*b^8*d^4 - A^4*a^8*d^4 - 38*A^4*
a^4*b^4*d^4 + 12*A^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2))/(16*A^3*a^6*d^2 -
16*A^3*b^6*d^2 + 112*A^3*a^2*b^4*d^2 - 112*A^3*a^4*b^2*d^2 + 32*A*a*b*(12*
A^4*a^2*b^6*d^4 - A^4*b^8*d^4 - A^4*a^8*d^4 - 38*A^4*a^4*b^4*d^4 + 12*A^4*
a^6*b^2*d^4)^(1/2)))*((A^2*a^3*b)/d^2 - (A^2*a*b^3)/d^2 - (12*A^4*a^2*b^6*
d^4 - A^4*b^8*d^4 - A^4*a^8*d^4 - 38*A^4*a^4*b^4*d^4 + 12*A^4*a^6*b^2*d^4)
^(1/2)/(4*d^4))^(1/2) - 2*atanh((32*B^2*a^4*tan(c + d*x)^(1/2)*((12*B^4*a^
2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^
2*d^4)^(1/2)/(4*d^4) + (B^2*a*b^3)/d^2 - (B^2*a^3*b)/d^2)^(1/2))/((16*B...
```

3.390 $\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

3.390.1 Optimal result 3707
 3.390.2 Mathematica [C] (verified) 3708
 3.390.3 Rubi [A] (verified) 3708
 3.390.4 Maple [A] (verified) 3713
 3.390.5 Fricas [B] (verification not implemented) 3714
 3.390.6 Sympy [F] 3714
 3.390.7 Maxima [A] (verification not implemented) 3715
 3.390.8 Giac [F(-1)] 3715
 3.390.9 Mupad [B] (verification not implemented) 3716

3.390.1 Optimal result

Integrand size = 33, antiderivative size = 283

$$\int \frac{(a + b \tan(c + dx))^2(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{(2ab(A - B) + a^2(A + B) - b^2(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(2ab(A - B) + a^2(A + B) - b^2(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(a^2(A - B) - b^2(A - B) - 2ab(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(a^2(A - B) - b^2(A - B) - 2ab(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$- \frac{2a^2A}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2a(2Ab + aB)}{d\sqrt{\tan(c + dx)}}$$

output

```
-1/2*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))
/d*2^(1/2)-1/2*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*arctan(1+2^(1/2)*tan(d*x+
c)^(1/2))/d*2^(1/2)+1/4*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*ln(1-2^(1/2)*tan
(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)-1/4*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*
ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)-2*a*(2*A*b+B*a)/d/tan(
d*x+c)^(1/2)-2/3*a^2*A/d/tan(d*x+c)^(3/2)
```

3.390.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.74 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.42

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2(-a^2 A + Ab^2 + 2abB) \operatorname{Hypergeometric2F1}\left(-\frac{3}{4}, 1, \frac{1}{4}, -\tan^2(c + dx)\right) - 6(2aAb + a^2 B - b^2 B) \operatorname{Hypergeometric2F1}\left(-\frac{1}{4}, 1, \frac{3}{4}, -\tan^2(c + dx)\right)}{3d \tan^{\frac{3}{2}}(c + dx)}$$

input `Integrate[((a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(2*(-(a^2*A) + A*b^2 + 2*a*b*B)*Hypergeometric2F1[-3/4, 1, 1/4, -Tan[c + d*x]^2] - 6*(2*a*A*b + a^2*B - b^2*B)*Hypergeometric2F1[-1/4, 1, 3/4, -Tan[c + d*x]^2]*Tan[c + d*x] - 2*b*(A*b + 2*a*B + 3*b*B*Tan[c + d*x]))/(3*d*Tan[c + d*x]^(3/2))`

3.390.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 235, normalized size of antiderivative = 0.83, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 4087, 3042, 4111, 25, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx$$

$$\downarrow \text{4087}$$

$$\int \frac{b^2 B \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)}{\tan^{\frac{3}{2}}(c + dx)} dx - \frac{2a^2 A}{3d \tan^{\frac{3}{2}}(c + dx)}$$

$$\downarrow \text{3042}$$

3.390. $\int \frac{(a+b \tan(c+dx))^2 (A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{b^2 B \tan(c+dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2Ab + aB)}{\tan(c+dx)^{3/2}} dx - \frac{2a^2 A}{3d \tan^{3/2}(c+dx)} \\
& \quad \downarrow \text{4111} \\
& \int -\frac{Aa^2 - 2bBa - Ab^2 - (b^2 B - a(2Ab + aB)) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2 A}{3d \tan^{3/2}(c+dx)} - \\
& \quad \frac{2a(aB + 2Ab)}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow \text{25} \\
& - \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2 B) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2 A}{3d \tan^{3/2}(c+dx)} - \frac{2a(aB + 2Ab)}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& - \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2 B) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2 A}{3d \tan^{3/2}(c+dx)} - \frac{2a(aB + 2Ab)}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow \text{4017} \\
& - \frac{2 \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2 B) \tan(c+dx)}{\tan^2(c+dx) + 1} d\sqrt{\tan(c+dx)}}{d} - \frac{2a^2 A}{3d \tan^{3/2}(c+dx)} - \frac{2a(aB + 2Ab)}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow \text{1482} \\
& - \frac{2\left(\frac{1}{2}(a^2(A - B) - 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx) + 1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a^2(A + B) + 2ab(A - B) - b^2(A - B))\right)}{d} \\
& \quad - \frac{\frac{2a^2 A}{3d \tan^{3/2}(c+dx)} - \frac{2a(aB + 2Ab)}{d\sqrt{\tan(c+dx)}}}{d} \\
& \quad \downarrow \text{1476} \\
& - \frac{2\left(\frac{1}{2}(a^2(A - B) - 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx) + 1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a^2(A + B) + 2ab(A - B) - b^2(A - B))\right)}{d} \\
& \quad - \frac{\frac{2a^2 A}{3d \tan^{3/2}(c+dx)} - \frac{2a(aB + 2Ab)}{d\sqrt{\tan(c+dx)}}}{d} \\
& \quad \downarrow \text{1082}
\end{aligned}$$

3.390. $\int \frac{(a+b \tan(c+dx))^2 (A+B \tan(c+dx))}{\tan^{5/2}(c+dx)} dx$

$$2 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^2(A + B) + 2ab(A - B) - b^2(A - B)) \int \frac{1 + \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)$$

$$\frac{2a^2A}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(aB + 2Ab)}{d\sqrt{\tan(c+dx)}}$$

↓ 217

$$2 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^2(A + B) + 2ab(A - B) - b^2(A - B)) \int \frac{1 + \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)$$

$$\frac{2a^2A}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(aB + 2Ab)}{d\sqrt{\tan(c+dx)}}$$

↓ 1479

$$2 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \left(- \int \frac{\sqrt{2} - 2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} \right) \right)$$

$$\frac{2a^2A}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(aB + 2Ab)}{d\sqrt{\tan(c+dx)}}$$

↓ 25

$$2 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \left(\int \frac{\sqrt{2} - 2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} \right) \right)$$

$$\frac{2a^2A}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(aB + 2Ab)}{d\sqrt{\tan(c+dx)}}$$

↓ 27

$$2 \left(\frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \left(\int \frac{\sqrt{2} - 2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)} + 1}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} \right) \right)$$

$$\frac{2a^2A}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(aB + 2Ab)}{d\sqrt{\tan(c+dx)}}$$

↓ 1103

3.390. $\int \frac{(a+b \tan(c+dx))^2 (A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

$$\frac{2 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} (a^2(A-B) - b^2(A+B)) \right)}{d} - \frac{2a^2A}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2a(aB+2Ab)}{d\sqrt{\tan(c+dx)}}$$

input `Int[((a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(-2*(((2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 + ((a^2*(A - B) - b^2*(A - B) - 2*a*b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d - (2*a^2*A)/(3*d*Tan[c + d*x]^(3/2)) - (2*a*(2*A*b + a*B))/(d*Sqrt[Tan[c + d*x]])`

3.390.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4087 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^2*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (B*c - A*d))*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`

```
rule 4111 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]
```

3.390.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 243, normalized size of antiderivative = 0.86

method	result
derivativedivides	$-\frac{2Aa^2}{3\tan(dx+c)^{\frac{3}{2}}}-\frac{2a(2Ab+Ba)}{\sqrt{\tan(dx+c)}}+\frac{(-Aa^2+Ab^2+2Bab)\sqrt{2}\left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right)+2\arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}))\right)}{4}$
default	$-\frac{2Aa^2}{3\tan(dx+c)^{\frac{3}{2}}}-\frac{2a(2Ab+Ba)}{\sqrt{\tan(dx+c)}}+\frac{(-Aa^2+Ab^2+2Bab)\sqrt{2}\left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right)+2\arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}))\right)}{4}$
parts	$\frac{(Ab^2+2Bab)\sqrt{2}\left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right)+2\arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}))\right)+2\arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}))}{4d}$

```
input int((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x,method=_RETURNV
ERBOSE)
```

```
output 1/d*(-2/3*A*a^2/tan(d*x+c)^(3/2)-2*a*(2*A*b+B*a)/tan(d*x+c)^(1/2)+1/4*(-A*
a^2+A*b^2+2*B*a*b)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-
2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))
+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(-2*A*a*b-B*a^2+B*b^2)*2^(1/2)
*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+t
an(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d
*x+c)^(1/2)))
```

3.390.
$$\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.390.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4313 vs. $2(249) = 498$.

Time = 0.72 (sec) , antiderivative size = 4313, normalized size of antiderivative = 15.24

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm
m="fricas")
```

```
output 1/6*(3*d*sqrt(-(2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3
*b - 4*(A^2 - B^2)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^
3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B -
A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B -
A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B
^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2)*log(((B*a^2 + 2*A*a*b
- B*b^2)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b
- 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 +
2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 -
4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 -
2*A^2*B^2 + B^4)*b^8)/d^4) + ((A^3 - A*B^2)*a^6 - 2*(5*A^2*B - B^3)*a^5*b
- (7*A^3 - 23*A*B^2)*a^4*b^2 + 4*(7*A^2*B - 3*B^3)*a^3*b^3 + (7*A^3 - 23*
A*B^2)*a^2*b^4 - 2*(5*A^2*B - B^3)*a*b^5 - (A^3 - A*B^2)*b^6)*d)*sqrt(-(2*
A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2)
*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b
- 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 +
2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 -
4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 -
2*A^2*B^2 + B^4)*b^8)/d^4))/d^2) - ((A^4 - B^4)*a^8 - 8*(A^3*B + A*B^3)*a
^7*b - 4*(A^4 - B^4)*a^6*b^2 - 8*(A^3*B + A*B^3)*a^5*b^3 - 10*(A^4 - B...
```

3.390.6 Sympy [F]

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^2}{\tan^{\frac{5}{2}}(c + dx)} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))**2*(A+B*tan(d*x+c))/tan(d*x+c)**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**2/tan(c + d*x)**(5/2), x)`

3.390.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 248, normalized size of antiderivative = 0.88

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx =$$

$$6\sqrt{2}((A + B)a^2 + 2(A - B)ab - (A + B)b^2) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(dx + c)}\right)\right) + 6\sqrt{2}((A + B)$$

input `integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="maxima")`

output `-1/12*(6*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 6*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + 3*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - 3*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 8*(A*a^2 + 3*(B*a^2 + 2*A*a*b)*tan(d*x + c))/tan(d*x + c)^(3/2)/d`

3.390.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="giac")`

output `Timed out`

3.390. $\int \frac{(a+b \tan(c+dx))^2 (A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

3.390.9 Mupad [B] (verification not implemented)

Time = 11.81 (sec) , antiderivative size = 3745, normalized size of antiderivative = 13.23

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = \text{Too large to display}$$

```
input int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2)/tan(c + d*x)^(5/2),x)
```

```
output 2*atanh((32*B^2*a^4*d^3*tan(c + d*x)^(1/2)*((B^2*a^3*b)/d^2 - (B^2*a*b^3)/
d^2 - (12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4
+ 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2))/(16*B^3*a^6*d^2 - 16*B^3*b^6*
d^2 + 32*B*a*b*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^
4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2) + 112*B^3*a^2*b^4*d^2 - 112*B^3*a^4*
b^2*d^2) + (32*B^2*b^4*d^3*tan(c + d*x)^(1/2)*((B^2*a^3*b)/d^2 - (B^2*a*b^
3)/d^2 - (12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*
d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2))/(16*B^3*a^6*d^2 - 16*B^3*b
^6*d^2 + 32*B*a*b*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4
*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2) + 112*B^3*a^2*b^4*d^2 - 112*B^3*a
^4*b^2*d^2) - (192*B^2*a^2*b^2*d^3*tan(c + d*x)^(1/2)*((B^2*a^3*b)/d^2 - (
B^2*a*b^3)/d^2 - (12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*
a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2)/(4*d^4))^(1/2))/(16*B^3*a^6*d^2 -
16*B^3*b^6*d^2 + 32*B*a*b*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4
- 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2) + 112*B^3*a^2*b^4*d^2 - 1
12*B^3*a^4*b^2*d^2))*((B^2*a^3*b)/d^2 - (B^2*a*b^3)/d^2 - (12*B^4*a^2*b^6*
d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)
^(1/2)/(4*d^4))^(1/2) - 2*atanh((32*B^2*a^4*d^3*tan(c + d*x)^(1/2)*((12*B^
4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^
6*b^2*d^4)^(1/2)/(4*d^4) - (B^2*a*b^3)/d^2 + (B^2*a^3*b)/d^2)^(1/2))/(1...
```

3.391
$$\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

3.391.1 Optimal result	3717
3.391.2 Mathematica [C] (verified)	3718
3.391.3 Rubi [A] (verified)	3718
3.391.4 Maple [A] (verified)	3724
3.391.5 Fricas [B] (verification not implemented)	3724
3.391.6 Sympy [F]	3725
3.391.7 Maxima [A] (verification not implemented)	3726
3.391.8 Giac [F(-1)]	3726
3.391.9 Mupad [B] (verification not implemented)	3727

3.391.1 Optimal result

Integrand size = 33, antiderivative size = 317

$$\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

$$= -\frac{(a^2(A-B) - b^2(A-B) - 2ab(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(a^2(A-B) - b^2(A-B) - 2ab(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(2ab(A-B) + a^2(A+B) - b^2(A+B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(2ab(A-B) + a^2(A+B) - b^2(A+B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}d}$$

$$- \frac{2a^2A}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2a(2Ab + aB)}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{2\sqrt{2}d}{d\sqrt{\tan(c+dx)}} \frac{2(a^2A - Ab^2 - 2abB)}{d\sqrt{\tan(c+dx)}}$$

output

```
1/2*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/
d*2^(1/2)+1/2*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*arctan(1+2^(1/2)*tan(d*x+c)
)^(1/2))/d*2^(1/2)+1/4*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*ln(1-2^(1/2)*tan(
d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)-1/4*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*l
n(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)+2*(A*a^2-A*b^2-2*B*a*b)
/d/tan(d*x+c)^(1/2)-2/5*a^2*A/d/tan(d*x+c)^(5/2)-2/3*a*(2*A*b+B*a)/d/tan(d
*x+c)^(3/2)
```

3.391.
$$\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

3.391.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.66 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.38

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{2((-3a^2A + 3Ab^2 + 6abB) \operatorname{Hypergeometric2F1}\left(-\frac{5}{4}, 1, -\frac{1}{4}, -\tan^2(c + dx)\right) - 5(2aAb + a^2B - b^2B) \operatorname{Hypergeometric2F1}\left(-\frac{3}{4}, 1, \frac{1}{4}, -\tan^2(c + dx)\right)}{15d \tan^{\frac{5}{2}}(c + dx)}$$

input `Integrate[((a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(2*((-3*a^2*A + 3*A*b^2 + 6*a*b*B)*Hypergeometric2F1[-5/4, 1, -1/4, -Tan[c + d*x]^2] - 5*(2*a*A*b + a^2*B - b^2*B)*Hypergeometric2F1[-3/4, 1, 1/4, -Tan[c + d*x]^2]*Tan[c + d*x] - b*(3*A*b + 6*a*B + 5*b*B*Tan[c + d*x]))/(15*d*Tan[c + d*x]^(5/2))`

3.391.3 Rubi [A] (verified)

Time = 0.92 (sec) , antiderivative size = 269, normalized size of antiderivative = 0.85, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 4087, 3042, 4111, 25, 3042, 4012, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$\downarrow 3042$$

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan(c + dx)^{7/2}} dx$$

$$\downarrow 4087$$

$$\int \frac{b^2 B \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)}{\tan^{\frac{5}{2}}(c + dx)} dx - \frac{2a^2 A}{5d \tan^{\frac{5}{2}}(c + dx)}$$

$$\downarrow 3042$$

3.391. $\int \frac{(a+b \tan(c+dx))^2 (A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{b^2 B \tan(c+dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2Ab + aB)}{\tan(c+dx)^{5/2}} dx - \frac{2a^2 A}{5d \tan^{5/2}(c+dx)} \\
& \quad \downarrow \text{4111} \\
& \int -\frac{Aa^2 - 2bBa - Ab^2 - (b^2 B - a(2Ab + aB)) \tan(c+dx)}{\tan^{3/2}(c+dx)} dx - \frac{2a^2 A}{5d \tan^{5/2}(c+dx)} - \\
& \quad \frac{2a(aB + 2Ab)}{3d \tan^{3/2}(c+dx)} \\
& \quad \downarrow \text{25} \\
& - \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2 B) \tan(c+dx)}{\tan^{3/2}(c+dx)} dx - \frac{2a^2 A}{5d \tan^{5/2}(c+dx)} - \\
& \quad \frac{2a(aB + 2Ab)}{3d \tan^{3/2}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& - \int \frac{Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2 B) \tan(c+dx)}{\tan(c+dx)^{3/2}} dx - \frac{2a^2 A}{5d \tan^{5/2}(c+dx)} - \\
& \quad \frac{2a(aB + 2Ab)}{3d \tan^{3/2}(c+dx)} \\
& \quad \downarrow \text{4012} \\
& - \int \frac{Ba^2 + 2Aba - b^2 B - (Aa^2 - 2bBa - Ab^2) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx + \frac{2(a^2 A - 2abB - Ab^2)}{d\sqrt{\tan(c+dx)}} - \\
& \quad \frac{2a^2 A}{5d \tan^{5/2}(c+dx)} - \frac{2a(aB + 2Ab)}{3d \tan^{3/2}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& - \int \frac{Ba^2 + 2Aba - b^2 B - (Aa^2 - 2bBa - Ab^2) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx + \frac{2(a^2 A - 2abB - Ab^2)}{d\sqrt{\tan(c+dx)}} - \\
& \quad \frac{2a^2 A}{5d \tan^{5/2}(c+dx)} - \frac{2a(aB + 2Ab)}{3d \tan^{3/2}(c+dx)} \\
& \quad \downarrow \text{4017} \\
& - \frac{2 \int \frac{Ba^2 + 2Aba - b^2 B - (Aa^2 - 2bBa - Ab^2) \tan(c+dx)}{\tan^2(c+dx) + 1} d\sqrt{\tan(c+dx)}}{d} + \frac{2(a^2 A - 2abB - Ab^2)}{d\sqrt{\tan(c+dx)}} - \\
& \quad \frac{2a^2 A}{5d \tan^{5/2}(c+dx)} - \frac{2a(aB + 2Ab)}{3d \tan^{3/2}(c+dx)} \\
& \quad \downarrow \text{1482}
\end{aligned}$$

3.391. $\int \frac{(a+b \tan(c+dx))^2 (A+B \tan(c+dx))}{\tan^{7/2}(c+dx)} dx$

$$\frac{2\left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A+B)) \int \frac{1+\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right)}{d}$$

$$\frac{2(a^2A - 2abB - Ab^2)}{d\sqrt{\tan(c+dx)}} - \frac{2a^2A}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2a(aB + 2Ab)}{3d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 1476

$$\frac{2\left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A+B)) \int \frac{1+\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right)}{d}$$

$$\frac{2(a^2A - 2abB - Ab^2)}{d\sqrt{\tan(c+dx)}} - \frac{2a^2A}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2a(aB + 2Ab)}{3d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 1082

$$\frac{2\left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A+B)) \int \frac{1+\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right)}{d}$$

$$\frac{2(a^2A - 2abB - Ab^2)}{d\sqrt{\tan(c+dx)}} - \frac{2a^2A}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2a(aB + 2Ab)}{3d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 217

$$\frac{2\left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A+B)) \int \frac{1+\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right)}{d}$$

$$\frac{2(a^2A - 2abB - Ab^2)}{d\sqrt{\tan(c+dx)}} - \frac{2a^2A}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2a(aB + 2Ab)}{3d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 1479

$$\frac{2\left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}}}{2\sqrt{2}} \right) - \frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A+B)) \int \frac{1+\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right)}{d}$$

$$\frac{2(a^2A - 2abB - Ab^2)}{d\sqrt{\tan(c+dx)}} - \frac{2a^2A}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2a(aB + 2Ab)}{3d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 25

3.391. $\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
& \frac{2 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d} \\
& \frac{2(a^2A - 2abB - Ab^2)}{d\sqrt{\tan(c+dx)}} - \frac{2a^2A}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2a(aB + 2Ab)}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{27} \\
& \frac{2 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}} d\sqrt{\tan(c+dx)} \right) \right)}{d} \\
& \frac{2(a^2A - 2abB - Ab^2)}{d\sqrt{\tan(c+dx)}} - \frac{2a^2A}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2a(aB + 2Ab)}{3d \tan^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{1103} \\
& \frac{2 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d} \\
& \frac{2(a^2A - 2abB - Ab^2)}{d\sqrt{\tan(c+dx)}} - \frac{2a^2A}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2a(aB + 2Ab)}{3d \tan^{\frac{3}{2}}(c+dx)}
\end{aligned}$$

input `Int[((a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(-2*(-1/2*((a^2*(A - B) - b^2*(A - B) - 2*a*b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])) + ((2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d - (2*a^2*A)/(5*d*Tan[c + d*x]^(5/2)) - (2*a*(2*A*b + a*B))/(3*d*Tan[c + d*x]^(3/2)) + (2*(a^2*A - A*b^2 - 2*a*b*B))/(d*Sqrt[Tan[c + d*x]])`

3.391.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4017 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4087 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-(B*c - A*d)*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`

rule 4111 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.391.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 270, normalized size of antiderivative = 0.85

method	result
derivativedivides	$-\frac{2(-Aa^2+Ab^2+2Bab)}{\sqrt{\tan(dx+c)}} - \frac{2Aa^2}{5\tan(dx+c)^{\frac{5}{2}}} - \frac{2a(2Ab+Ba)}{3\tan(dx+c)^{\frac{3}{2}}} + \frac{(-2Aab-Ba^2+Bb^2)\sqrt{2}\left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right)+2\arctan\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right)\right)}{4}$
default	$-\frac{2(-Aa^2+Ab^2+2Bab)}{\sqrt{\tan(dx+c)}} - \frac{2Aa^2}{5\tan(dx+c)^{\frac{5}{2}}} - \frac{2a(2Ab+Ba)}{3\tan(dx+c)^{\frac{3}{2}}} + \frac{(-2Aab-Ba^2+Bb^2)\sqrt{2}\left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right)+2\arctan\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right)\right)}{4}$
parts	$(Ab^2+2Bab)\left(-\frac{\sqrt{2}\left(\ln\left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right)+2\arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}))\right)+2\arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}))\right)}{4}\right)$ d

input `int((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x,method=_RETURNV
ERBOSE)`

output `1/d*(-2*(-A*a^2+A*b^2+2*B*a*b)/tan(d*x+c)^(1/2)-2/5*A*a^2/tan(d*x+c)^(5/2)
-2/3*a*(2*A*b+B*a)/tan(d*x+c)^(3/2)+1/4*(-2*A*a*b-B*a^2+B*b^2)*2^(1/2)*(ln
((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d
*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c
)^(1/2)))+1/4*(A*a^2-A*b^2-2*B*a*b)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2
)
+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*
tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))`

3.391.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4325 vs. 2(279) = 558.

Time = 0.73 (sec) , antiderivative size = 4325, normalized size of antiderivative = 13.64

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm
m="fracas")`

output

```
-1/30*(15*d*sqrt((2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2)*log(((A*a^2 - 2*B*a*b - A*b^2)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4) - ((A^2*B - B^3)*a^6 + 2*(A^3 - 5*A*B^2)*a^5*b - (23*A^2*B - 7*B^3)*a^4*b^2 - 4*(3*A^3 - 7*A*B^2)*a^3*b^3 + (23*A^2*B - 7*B^3)*a^2*b^4 + 2*(A^3 - 5*A*B^2)*a*b^5 - (A^2*B - B^3)*b^6)*d)*sqrt((2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2) - ((A^4 - B^4)*a^8 - 8*(A^3*B + A*B^3)*a^7*b - 4*(A^4 - B^4)*a^6*b^2 - 8*(A^3*B + A*B^3)*a^5*b^3 - 10*(A^4 - B...
```

3.391.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^2}{\tan^{\frac{7}{2}}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**2*(A+B*tan(d*x+c))/tan(d*x+c)**(7/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**2/tan(c + d*x)**(7/2), x)`

3.391.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 276, normalized size of antiderivative = 0.87

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{30 \sqrt{2}((A - B)a^2 - 2(A + B)ab - (A - B)b^2) \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2} + 2 \sqrt{\tan(dx + c)})\right) + 30 \sqrt{2}((A - B)$$

input `integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="maxima")`

output `1/60*(30*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 30*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - 15*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 15*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - 8*(3*A*a^2 - 15*(A*a^2 - 2*B*a*b - A*b^2)*tan(d*x + c)^2 + 5*(B*a^2 + 2*A*a*b)*tan(d*x + c))/tan(d*x + c)^(5/2))/d`

3.391.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="giac")`

output `Timed out`

3.391.9 Mupad [B] (verification not implemented)

Time = 15.11 (sec) , antiderivative size = 3782, normalized size of antiderivative = 11.93

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \text{Too large to display}$$

```
input int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2)/tan(c + d*x)^(7/2),x)
```

```
output 2*atanh((32*B^2*a^4*d^3*tan(c + d*x)^(1/2)*((B^2*a*b^3)/d^2 - (12*B^4*a^2*
b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*
d^4)^(1/2))/(4*d^4) - (B^2*a^3*b)/d^2)^(1/2))/(16*B*a^2*(12*B^4*a^2*b^6*d^4
- B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1
/2) - 16*B*b^2*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^
4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2) - 192*B^3*a^3*b^3*d^2 + 32*B^3*a*b^5
*d^2 + 32*B^3*a^5*b*d^2) + (32*B^2*b^4*d^3*tan(c + d*x)^(1/2)*((B^2*a*b^3)
/d^2 - (12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^
4 + 12*B^4*a^6*b^2*d^4)^(1/2))/(4*d^4) - (B^2*a^3*b)/d^2)^(1/2))/(16*B*a^2*
(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*
B^4*a^6*b^2*d^4)^(1/2) - 16*B*b^2*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*
a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2) - 192*B^3*a^3*b^3
*d^2 + 32*B^3*a*b^5*d^2 + 32*B^3*a^5*b*d^2) - (192*B^2*a^2*b^2*d^3*tan(c +
d*x)^(1/2)*((B^2*a*b^3)/d^2 - (12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8
*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2))/(4*d^4) - (B^2*a^3*b
)/d^2)^(1/2))/(16*B*a^2*(12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 -
38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)^(1/2) - 16*B*b^2*(12*B^4*a^2*b^6*
d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*a^4*b^4*d^4 + 12*B^4*a^6*b^2*d^4)
^(1/2) - 192*B^3*a^3*b^3*d^2 + 32*B^3*a*b^5*d^2 + 32*B^3*a^5*b*d^2))*((B^2
*a*b^3)/d^2 - (12*B^4*a^2*b^6*d^4 - B^4*b^8*d^4 - B^4*a^8*d^4 - 38*B^4*...
```

3.392 $\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.392.1 Optimal result	3728
3.392.2 Mathematica [C] (verified)	3729
3.392.3 Rubi [A] (verified)	3730
3.392.4 Maple [A] (verified)	3737
3.392.5 Fricas [B] (verification not implemented)	3737
3.392.6 Sympy [F]	3738
3.392.7 Maxima [A] (verification not implemented)	3738
3.392.8 Giac [F(-1)]	3739
3.392.9 Mupad [B] (verification not implemented)	3739

3.392.1 Optimal result

Integrand size = 33, antiderivative size = 463

$$\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{(3a^2b(A-B) - b^3(A-B) + a^3(A+B) - 3ab^2(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(3a^2b(A-B) - b^3(A-B) + a^3(A+B) - 3ab^2(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(a^3(A-B) - 3ab^2(A-B) - 3a^2b(A+B) + b^3(A+B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(a^3(A-B) - 3ab^2(A-B) - 3a^2b(A+B) + b^3(A+B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{2(a^3A - 3aAb^2 - 3a^2bB + b^3B) \sqrt{\tan(c+dx)}}{d}$$

$$+ \frac{2(3a^2Ab - Ab^3 + a^3B - 3ab^2B) \tan^{\frac{3}{2}}(c+dx)}{3d}$$

$$+ \frac{2b(27aAb + 22a^2B - 9b^2B) \tan^{\frac{5}{2}}(c+dx)}{45d}$$

$$+ \frac{2b^2(9Ab + 13aB) \tan^{\frac{7}{2}}(c+dx)}{63d} + \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2}{9d}$$

3.392. $\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

output
$$\begin{aligned} & -1/2*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*\arctan(-1+2^{(1/2)}*\tan(dx+c)^{(1/2)})/d*2^{(1/2)}-1/2*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*\arctan(1+2^{(1/2)}*\tan(dx+c)^{(1/2)})/d*2^{(1/2)}+1/4*(a^3*(A-B)-3*a*b^2*(A-B)-3*a^2*b*(A+B)+b^3*(A+B))*\ln(1-2^{(1/2)}*\tan(dx+c)^{(1/2)}+\tan(dx+c))/d*2^{(1/2)}-1/4*(a^3*(A-B)-3*a*b^2*(A-B)-3*a^2*b*(A+B)+b^3*(A+B))*\ln(1+2^{(1/2)}*\tan(dx+c)^{(1/2)}+\tan(dx+c))/d*2^{(1/2)}+2*(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*\tan(dx+c)^{(1/2)}/d+2/3*(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*\tan(dx+c)^{(3/2)}/d+2/45*b*(27*A*a*b+22*B*a^2-9*B*b^2)*\tan(dx+c)^{(5/2)}/d+2/63*b^2*(9*A*b+13*B*a)*\tan(dx+c)^{(7/2)}/d+2/9*b*B*\tan(dx+c)^{(5/2)}*(a+b*\tan(dx+c))^2/d \end{aligned}$$

3.392.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.72 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.48

$$\int \tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx$$

$$= \frac{2(7b(27aAb+22a^2B-9b^2B)\tan^{\frac{5}{2}}(c+dx)+5b^2(9Ab+13aB)\tan^{\frac{7}{2}}(c+dx)+35bB\tan^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx)))}{(315*d)}$$

input `Integrate[Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output
$$\begin{aligned} & (2*(7*b*(27*a*A*b+22*a^2*B-9*b^2*B)*\tan[c+d*x]^{(5/2)}+5*b^2*(9*A*b+13*a*B)*\tan[c+d*x]^{(7/2)}+35*b*B*\tan[c+d*x]^{(5/2)}*(a+b*\tan[c+d*x])^2+(105*(a-I*b)^3*(I*A+B)*(-3*(-1)^{(3/4)}*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\tan[c+d*x]])+\text{Sqrt}[\tan[c+d*x]]*(-3*I+\tan[c+d*x])))/2+(105*(a+I*b)^3*((-I)*A+B)*(3*(-1)^{(3/4)}*\text{ArcTanh}[(-1)^{(3/4)}*\text{Sqrt}[\tan[c+d*x]])+\text{Sqrt}[\tan[c+d*x]]*(3*I+\tan[c+d*x])))/2))/(315*d) \end{aligned}$$

3.392.3 Rubi [A] (verified)

Time = 1.65 (sec) , antiderivative size = 395, normalized size of antiderivative = 0.85, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.697$, Rules used = {3042, 4090, 27, 3042, 4120, 27, 3042, 4113, 3042, 4011, 3042, 4011, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c+dx)^{3/2}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4090} \\
 & \frac{2}{9} \int \frac{1}{2} \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx)) (b(9Ab+13aB) \tan^2(c+dx) + 9(Ba^2+2Aba-b^2B) \tan(c+dx) + a(9aA-5bB)) dx + \\
 & \quad \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2}{9d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{9} \int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx)) (b(9Ab+13aB) \tan^2(c+dx) + 9(Ba^2+2Aba-b^2B) \tan(c+dx) + a(9aA-5bB)) dx + \\
 & \quad \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2}{9d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{9} \int \tan(c+dx)^{3/2}(a+b \tan(c+dx)) (b(9Ab+13aB) \tan(c+dx)^2 + 9(Ba^2+2Aba-b^2B) \tan(c+dx) + a(9aA-5bB)) dx + \\
 & \quad \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2}{9d} \\
 & \quad \downarrow \text{4120} \\
 & \frac{1}{9} \left(\frac{2b^2(13aB+9Ab) \tan^{\frac{7}{2}}(c+dx)}{7d} - \frac{2}{7} \int -\frac{7}{2} \tan^{\frac{3}{2}}(c+dx) ((9aA-5bB)a^2 + b(22Ba^2+27Aba-9b^2B) \tan^2(c+dx) \right. \\
 & \quad \left. + \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2}{9d} \right)
 \end{aligned}$$

3.392. $\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

↓ 27

$$\frac{1}{9} \left(\int \tan^{\frac{3}{2}}(c+dx) ((9aA - 5bB)a^2 + b(22Ba^2 + 27Aba - 9b^2B) \tan^2(c+dx) + 9(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)) \right. \\ \left. \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))^2}{9d} \right)$$

↓ 3042

$$\frac{1}{9} \left(\int \tan(c+dx)^{3/2} ((9aA - 5bB)a^2 + b(22Ba^2 + 27Aba - 9b^2B) \tan(c+dx)^2 + 9(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)) \right. \\ \left. \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))^2}{9d} \right)$$

↓ 4113

$$\frac{1}{9} \left(\int \tan^{\frac{3}{2}}(c+dx) (9(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + 9(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)) dx + \frac{2b(22Ba^2 + 27Aba - 9b^2B)}{9d} \right. \\ \left. \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))^2}{9d} \right)$$

↓ 3042

$$\frac{1}{9} \left(\int \tan(c+dx)^{3/2} (9(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + 9(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)) dx + \frac{2b(22Ba^2 + 27Aba - 9b^2B)}{9d} \right. \\ \left. \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))^2}{9d} \right)$$

↓ 4011

$$\frac{1}{9} \left(\int \sqrt{\tan(c+dx)} (9(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) - 9(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)) dx + \frac{2b(22Ba^2 + 27Aba - 9b^2B)}{9d} \right. \\ \left. \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))^2}{9d} \right)$$

↓ 3042

$$\frac{1}{9} \left(\int \sqrt{\tan(c+dx)} (9(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) - 9(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)) dx + \frac{2b(22Ba^2 + 27Aba - 9b^2B)}{9d} \right. \\ \left. \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a + b \tan(c+dx))^2}{9d} \right)$$

↓ 4011

3.392. $\int \tan^{\frac{3}{2}}(c+dx)(a + b \tan(c+dx))^3(A + B \tan(c+dx)) dx$

$$\frac{1}{9} \left(\int \frac{-9(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) - 9(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2b(22a^2B + 27aAb)}{2bB \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} \right) dx + \frac{2b(22a^2B + 27aAb)}{9d}$$

↓ 3042

$$\frac{1}{9} \left(\int \frac{-9(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) - 9(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2b(22a^2B + 27aAb)}{2bB \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} \right) dx + \frac{2b(22a^2B + 27aAb)}{9d}$$

↓ 4017

$$\frac{1}{9} \left(2 \int \frac{-9(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \frac{2b(22a^2B + 27aAb - 9b^2B) \tan(c + dx)}{5d} \right) dx + \frac{2b(22a^2B + 27aAb)}{9d}$$

↓ 27

$$\frac{1}{9} \left(-18 \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \frac{2b(22a^2B + 27aAb - 9b^2B) \tan(c + dx)}{5d} \right) dx + \frac{2b(22a^2B + 27aAb)}{9d}$$

↓ 1482

$$\frac{1}{9} \left(-18 \left(\frac{1}{2}(a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \frac{1}{2}(a^3(A + B) - 3a^2b(A - B) - 3ab^2(A + B) + b^3(A - B)) \int \frac{1 + \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} \right) + \frac{2b(22a^2B + 27aAb - 9b^2B) \tan(c + dx)}{5d} \right) dx + \frac{2b(22a^2B + 27aAb)}{9d}$$

↓ 1476

$$\frac{1}{9} \left(-18 \left(\frac{1}{2}(a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \frac{1}{2}(a^3(A + B) - 3a^2b(A - B) - 3ab^2(A + B) + b^3(A - B)) \int \frac{1 + \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} \right) + \frac{2b(22a^2B + 27aAb - 9b^2B) \tan(c + dx)}{5d} \right) dx + \frac{2b(22a^2B + 27aAb)}{9d}$$

3.392. $\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

↓ 1082

$$\frac{1}{9} \left(\frac{18 \left(\frac{1}{2}(a^3(A-B) - 3a^2b(A+B) - 3ab^2(A-B) + b^3(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a^3(A+B) \right)}{\right.$$

$$\left. \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2}{9d} \right)$$

↓ 217

$$\frac{1}{9} \left(\frac{18 \left(\frac{1}{2}(a^3(A-B) - 3a^2b(A+B) - 3ab^2(A-B) + b^3(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a^3(A+B) \right)}{\right.$$

$$\left. \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2}{9d} \right)$$

↓ 1479

$$\frac{1}{9} \left(\frac{18 \left(\frac{1}{2}(a^3(A-B) - 3a^2b(A+B) - 3ab^2(A-B) + b^3(A+B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}}}{\right)}{\right.$$

$$\left. \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2}{9d} \right)$$

↓ 25

$$\frac{1}{9} \left(\frac{18 \left(\frac{1}{2}(a^3(A-B) - 3a^2b(A+B) - 3ab^2(A-B) + b^3(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}}}{\right)}{\right.$$

$$\left. \frac{2bB \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2}{9d} \right)$$

↓ 27

$$\frac{1}{9} \left(\frac{18 \left(\frac{1}{2}(a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \right) \left(\int \frac{\sqrt{2} - 2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)} dx \right)}{\frac{2bB \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2}{9d}} \right)$$

↓ 1103

$$\frac{1}{9} \left(\frac{2b(22a^2B + 27aAb - 9b^2B) \tan^{\frac{5}{2}}(c + dx)}{5d} - \frac{18 \left(\frac{1}{2}(a^3(A + B) + 3a^2b(A - B) - 3ab^2(A + B) - b^3(A - B)) \right) \left(\int \frac{\sqrt{2} - 2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)} dx \right)}{\frac{2bB \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2}{9d}} \right)$$

input `Int[Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(2*b*B*Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^2)/(9*d) + ((-18*((3*a^2*b*(A - B) - b^3*(A - B) + a^3*(A + B) - 3*a*b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 + ((a^3*(A - B) - 3*a*b^2*(A - B) - 3*a^2*b*(A + B) + b^3*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d + (18*(a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*Sqrt[Tan[c + d*x]]/d + (6*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*Tan[c + d*x]^(3/2))/d + (2*b*(27*a*A*b + 22*a^2*B - 9*b^2*B)*Tan[c + d*x]^(5/2))/(5*d) + (2*b^2*(9*A*b + 13*a*B)*Tan[c + d*x]^(7/2))/(7*d))/9`

3.392.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

3.392. $\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

- rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*(a + b*Tan[e + f*x])^m/(f*m), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4120 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]`

3.392.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 432, normalized size of antiderivative = 0.93

method	result
derivativedivides	$\frac{2B b^3 \left(\tan^{\frac{9}{2}}(dx+c)\right)}{9} + \frac{2A b^3 \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} + \frac{6B a b^2 \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} + \frac{6A a b^2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{6B a^2 b \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} - \frac{2B b^3 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3}$
default	$\frac{2B b^3 \left(\tan^{\frac{9}{2}}(dx+c)\right)}{9} + \frac{2A b^3 \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} + \frac{6B a b^2 \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} + \frac{6A a b^2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{6B a^2 b \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} - \frac{2B b^3 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3}$
parts	$(A b^3 + 3B a b^2) \left(\frac{2 \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} - \frac{2 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} + \frac{\sqrt{2} \left(\ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right)}{4} \right)$

input `int(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)`

output `1/d*(2/9*B*b^3*tan(d*x+c)^(9/2)+2/7*A*b^3*tan(d*x+c)^(7/2)+6/7*B*a*b^2*tan
(d*x+c)^(7/2)+6/5*A*a*b^2*tan(d*x+c)^(5/2)+6/5*B*a^2*b*tan(d*x+c)^(5/2)-2/
5*B*b^3*tan(d*x+c)^(5/2)+2*A*a^2*b*tan(d*x+c)^(3/2)-2/3*A*b^3*tan(d*x+c)^(
3/2)+2/3*B*a^3*tan(d*x+c)^(3/2)-2*B*a*b^2*tan(d*x+c)^(3/2)+2*A*a^3*tan(d*x
+c)^(1/2)-6*tan(d*x+c)^(1/2)*A*a*b^2-6*tan(d*x+c)^(1/2)*B*a^2*b+2*tan(d*x+
c)^(1/2)*B*b^3+1/4*(-A*a^3+3*A*a*b^2+3*B*a^2*b-B*b^3)*2^(1/2)*(ln(((1+2^(1/
2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))))+2
*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))
+1/4*(-3*A*a^2*b+A*b^3-B*a^3+3*B*a*b^2)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(
1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1
/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))`

3.392.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6209 vs. 2(417) = 834.

Time = 1.40 (sec) , antiderivative size = 6209, normalized size of antiderivative = 13.41

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm
m="fricas")`

3.392. $\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

output Too large to include

3.392.6 Sympy [F]

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^3 \tan^{\frac{3}{2}}(c + dx) dx$$

input `integrate(tan(d*x+c)**(3/2)*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**3*tan(c + d*x)**(3/2), x)`

3.392.7 Maxima [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 398, normalized size of antiderivative = 0.86

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{280 B b^3 \tan(dx + c)^{\frac{9}{2}} + 360 (3 B a b^2 + A b^3) \tan(dx + c)^{\frac{7}{2}} + 504 (3 B a^2 b + 3 A a b^2 - B b^3) \tan(dx + c)^{\frac{5}{2}}}{d}$$

input `integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/1260*(280*B*b^3*tan(d*x + c)^(9/2) + 360*(3*B*a*b^2 + A*b^3)*tan(d*x + c)^(7/2) + 504*(3*B*a^2*b + 3*A*a*b^2 - B*b^3)*tan(d*x + c)^(5/2) - 630*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) - 630*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - 315*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 315*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 840*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*tan(d*x + c)^(3/2) + 2520*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 + B*b^3)*sqrt(tan(d*x + c)))/d`

3.392. $\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.392.8 Giac [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \text{Timed out}$$

```
input integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")
```

output Timed out

3.392.9 Mupad [B] (verification not implemented)

Time = 38.61 (sec) , antiderivative size = 6774, normalized size of antiderivative = 14.63

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)
```

```
output tan(c + d*x)^(1/2)*((2*A*a^3)/d - (6*A*a*b^2)/d) - tan(c + d*x)^(3/2)*((2*A*b^3)/(3*d) - (2*A*a^2*b)/d) + tan(c + d*x)^(3/2)*((2*B*a^3)/(3*d) - (2*B*a*b^2)/d) + tan(c + d*x)^(1/2)*((2*B*b^3)/d - (6*B*a^2*b)/d) - tan(c + d*x)^(5/2)*((2*B*b^3)/(5*d) - (6*B*a^2*b)/(5*d)) - atan((((8*(4*A*a^3*d^2 - 12*A*a*b^2*d^2)*((5*A^2*a^3*b^3)/d^2 - (30*A^4*a^2*b^10*d^4 - A^4*b^12*d^4 - A^4*a^12*d^4 - 255*A^4*a^4*b^8*d^4 + 452*A^4*a^6*b^6*d^4 - 255*A^4*a^8*b^4*d^4 + 30*A^4*a^10*b^2*d^4)^(1/2)/(4*d^4) - (3*A^2*a*b^5)/(2*d^2) - (3*A^2*a^5*b)/(2*d^2))^(1/2))/d^3 - (16*tan(c + d*x)^(1/2)*(A^2*a^6 - A^2*b^6 + 15*A^2*a^2*b^4 - 15*A^2*a^4*b^2))/d^2)*((5*A^2*a^3*b^3)/d^2 - (30*A^4*a^2*b^10*d^4 - A^4*b^12*d^4 - A^4*a^12*d^4 - 255*A^4*a^4*b^8*d^4 + 452*A^4*a^6*b^6*d^4 - 255*A^4*a^8*b^4*d^4 + 30*A^4*a^10*b^2*d^4)^(1/2)/(4*d^4) - (3*A^2*a*b^5)/(2*d^2) - (3*A^2*a^5*b)/(2*d^2))^(1/2)*1i - ((8*(4*A*a^3*d^2 - 12*A*a*b^2*d^2)*((5*A^2*a^3*b^3)/d^2 - (30*A^4*a^2*b^10*d^4 - A^4*b^12*d^4 - A^4*a^12*d^4 - 255*A^4*a^4*b^8*d^4 + 452*A^4*a^6*b^6*d^4 - 255*A^4*a^8*b^4*d^4 + 30*A^4*a^10*b^2*d^4)^(1/2)/(4*d^4) - (3*A^2*a*b^5)/(2*d^2) - (3*A^2*a^5*b)/(2*d^2))^(1/2)*1i)/((16*(3*A^3*a...
```

3.393 $\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.393.1 Optimal result	3740
3.393.2 Mathematica [C] (verified)	3741
3.393.3 Rubi [A] (verified)	3742
3.393.4 Maple [A] (verified)	3748
3.393.5 Fricas [B] (verification not implemented)	3749
3.393.6 Sympy [F]	3749
3.393.7 Maxima [A] (verification not implemented)	3750
3.393.8 Giac [F(-1)]	3750
3.393.9 Mupad [B] (verification not implemented)	3751

3.393.1 Optimal result

Integrand size = 33, antiderivative size = 421

$$\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx =$$

$$\frac{(a^3(A - B) - 3ab^2(A - B) - 3a^2b(A + B) + b^3(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(a^3(A - B) - 3ab^2(A - B) - 3a^2b(A + B) + b^3(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(3a^2b(A - B) - b^3(A - B) + a^3(A + B) - 3ab^2(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(3a^2b(A - B) - b^3(A - B) + a^3(A + B) - 3ab^2(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{2(3a^2Ab - Ab^3 + a^3B - 3ab^2B) \sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{2b(21aAb + 18a^2B - 7b^2B) \tan^{\frac{3}{2}}(c + dx)}{21d}$$

$$+ \frac{2b^2(7Ab + 11aB) \tan^{\frac{5}{2}}(c + dx)}{35d} + \frac{2bB \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2}{7d}$$

output $\frac{1}{2}(a^3(A-B)-3ab^2(A-B)-3a^2b(A+B)+b^3(A+B))\arctan(-1+2^{1/2}\tan(dx+c)^{1/2})/d+1/2(a^3(A-B)-3ab^2(A-B)-3a^2b(A+B)+b^3(A+B))\arctan(1+2^{1/2}\tan(dx+c)^{1/2})/d+1/4(3a^2b(A-B)-b^3(A-B)+a^3(A+B)-3ab^2(A+B))\ln(1-2^{1/2}\tan(dx+c)^{1/2}+\tan(dx+c))/d+1/4(3a^2b(A-B)-b^3(A-B)+a^3(A+B)-3ab^2(A+B))\ln(1+2^{1/2}\tan(dx+c)^{1/2}+\tan(dx+c))/d+2(3Aa^2b-Ab^3+B^3-3Bab^2)\tan(dx+c)^{3/2}/d+2/35b^2(7Ab+11Ba)\tan(dx+c)^{5/2}/d+2/7bB\tan(dx+c)^{3/2}(a+b\tan(dx+c))^2/d$

3.393.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.18 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.47

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx$$

$$= \frac{2\left(\frac{105}{2}(a-ib)^3(iA+B)\left(\sqrt[4]{-1}\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)+\sqrt{\tan(c+dx)}\right)+\frac{105}{2}(a+ib)^3(-iA+B)\left(\sqrt[4]{-1}\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)+\sqrt{\tan(c+dx)}\right)\right)}{d}$$

input `Integrate[Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output $(2*((105*(a - I*b)^3*(I*A + B)*((-1)^(1/4)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]) + Sqrt[Tan[c + d*x]]))/2 + (105*(a + I*b)^3*((-I)*A + B)*((-1)^(1/4)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]) + Sqrt[Tan[c + d*x]]))/2 + 5*b*(21*a*A*b + 18*a^2*B - 7*b^2*B)*Tan[c + d*x]^(3/2) + 3*b^2*(7*A*b + 11*a*B)*Tan[c + d*x]^(5/2) + 15*b*B*Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^2)/(105*d)$

3.393.3 Rubi [A] (verified)

Time = 1.38 (sec) , antiderivative size = 355, normalized size of antiderivative = 0.84, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 4090, 27, 3042, 4120, 27, 3042, 4113, 3042, 4011, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4090} \\
 & \frac{2}{7} \int \frac{1}{2} \sqrt{\tan(c+dx)}(a+b \tan(c+dx)) (b(7Ab+11aB) \tan^2(c+dx) + 7(Ba^2+2Aba-b^2B) \tan(c+dx) + a(7aA-3bB)) dx + \\
 & \quad \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{7} \int \sqrt{\tan(c+dx)}(a+b \tan(c+dx)) (b(7Ab+11aB) \tan^2(c+dx) + 7(Ba^2+2Aba-b^2B) \tan(c+dx) + a(7aA-3bB)) dx + \\
 & \quad \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{7} \int \sqrt{\tan(c+dx)}(a+b \tan(c+dx)) (b(7Ab+11aB) \tan(c+dx)^2 + 7(Ba^2+2Aba-b^2B) \tan(c+dx) + a(7aA-3bB)) dx + \\
 & \quad \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \\
 & \quad \downarrow \text{4120} \\
 & \frac{1}{7} \left(\frac{2b^2(11aB+7Ab) \tan^{\frac{5}{2}}(c+dx)}{5d} - \frac{2}{5} \int -\frac{5}{2} \sqrt{\tan(c+dx)}((7aA-3bB)a^2 + b(18Ba^2+21Aba-7b^2B) \tan^2(c+dx) + \right. \\
 & \quad \left. \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \right)
 \end{aligned}$$

3.393. $\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\downarrow 27$$

$$\frac{1}{7} \left(\int \sqrt{\tan(c+dx)} ((7aA - 3bB)a^2 + b(18Ba^2 + 21Aba - 7b^2B) \tan^2(c+dx) + 7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)) dx + \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \right)$$

$$\downarrow 3042$$

$$\frac{1}{7} \left(\int \sqrt{\tan(c+dx)} ((7aA - 3bB)a^2 + b(18Ba^2 + 21Aba - 7b^2B) \tan(c+dx)^2 + 7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)) dx + \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \right)$$

$$\downarrow 4113$$

$$\frac{1}{7} \left(\int \sqrt{\tan(c+dx)} (7(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + 7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)) dx + \frac{2b(18a^2B + 21aAb - 7b^2B)}{7d} \right)$$

$$\downarrow 3042$$

$$\frac{1}{7} \left(\int \sqrt{\tan(c+dx)} (7(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + 7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)) dx + \frac{2b(18a^2B + 21aAb - 7b^2B)}{7d} \right)$$

$$\downarrow 4011$$

$$\frac{1}{7} \left(\int \frac{7(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) - 7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)}{\sqrt{\tan(c+dx)}} dx + \frac{2b(18a^2B + 21aAb - 7b^2B)}{7d} \right)$$

$$\downarrow 3042$$

$$\frac{1}{7} \left(\int \frac{7(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) - 7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)}{\sqrt{\tan(c+dx)}} dx + \frac{2b(18a^2B + 21aAb - 7b^2B)}{7d} \right)$$

$$\downarrow 4017$$

3.393. $\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\frac{1}{7} \left(\frac{2 \int -\frac{7(Ba^3+3Aba^2-3b^2Ba-Ab^3-(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} + \frac{2b(18a^2B+21aAb-7b^2B)\tan(c+dx)}{3d} \right. \\ \left. \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \right. \\ \left. \downarrow 27 \right.$$

$$\frac{1}{7} \left(-\frac{14 \int \frac{Ba^3+3Aba^2-3b^2Ba-Ab^3-(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} + \frac{2b(18a^2B+21aAb-7b^2B)\tan(c+dx)}{3d} \right. \\ \left. \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \right. \\ \left. \downarrow 1482 \right.$$

$$\frac{1}{7} \left(-\frac{14 \left(\frac{1}{2}(a^3(A+B)+3a^2b(A-B)-3ab^2(A+B)-b^3(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^3(A-B)) \right)}{d} \right. \\ \left. \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \right. \\ \left. \downarrow 1476 \right.$$

$$\frac{1}{7} \left(-\frac{14 \left(\frac{1}{2}(a^3(A+B)+3a^2b(A-B)-3ab^2(A+B)-b^3(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^3(A-B)) \right)}{d} \right. \\ \left. \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \right. \\ \left. \downarrow 1082 \right.$$

$$\frac{1}{7} \left(-\frac{14 \left(\frac{1}{2}(a^3(A+B)+3a^2b(A-B)-3ab^2(A+B)-b^3(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^3(A-B)) \right)}{d} \right. \\ \left. \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \right. \\ \left. \downarrow 217 \right.$$

3.393. $\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\frac{1}{7} \left(\frac{14 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^3(A-B) - B) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} \right. \\ \left. \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \right. \\ \left. \downarrow 1479 \right)$$

$$\frac{1}{7} \left(\frac{14 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(- \int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} - \int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d} \right. \\ \left. \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \right. \\ \left. \downarrow 25 \right)$$

$$\frac{1}{7} \left(\frac{14 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d} \right. \\ \left. \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \right. \\ \left. \downarrow 27 \right)$$

$$\frac{1}{7} \left(\frac{14 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d} \right. \\ \left. \frac{2bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2}{7d} \right. \\ \left. \downarrow 1103 \right)$$

$$\frac{1}{7} \left(\frac{2b(18a^2B + 21aAb - 7b^2B) \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{14 \left(\frac{1}{2}(a^3(A + B) + 3a^2b(A - B) - 3ab^2(A + B) - b^3(A - B)) \right)}{7d} \right) \frac{2bB \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2}{7d}$$

input `Int[Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(2*b*B*Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^2)/(7*d) + ((-14*(-1/2*((a^3*(A - B) - 3*a*b^2*(A - B) - 3*a^2*b*(A + B) + b^3*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])) + ((3*a^2*b*(A - B) - b^3*(A - B) + a^3*(A + B) - 3*a*b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d + (14*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*Sqrt[Tan[c + d*x]])/d + (2*b*(21*a*A*b + 18*a^2*B - 7*b^2*B)*Tan[c + d*x]^(3/2))/(3*d) + (2*b^2*(7*A*b + 11*a*B)*Tan[c + d*x]^(5/2))/(5*d))/7`

3.393.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x], Sqrt[b*Tan[e + f*x]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4120 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]`

3.393.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 372, normalized size of antiderivative = 0.88

method	result
derivativedivides	$\frac{2Bb^3 \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} + \frac{2Ab^3 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{6Bab^2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + 2Aab^2 \left(\tan^{\frac{3}{2}}(dx+c)\right) + 2Ba^2b \left(\tan^{\frac{3}{2}}(dx+c)\right) - \frac{2Bb^3}{7}$
default	$\frac{2Bb^3 \left(\tan^{\frac{7}{2}}(dx+c)\right)}{7} + \frac{2Ab^3 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{6Bab^2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + 2Aab^2 \left(\tan^{\frac{3}{2}}(dx+c)\right) + 2Ba^2b \left(\tan^{\frac{3}{2}}(dx+c)\right) - \frac{2Bb^3}{7}$
parts	$(Ab^3 + 3Bab^2) \left(\frac{2 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} - 2\sqrt{\tan(dx+c)} + \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))} \right) + 2 \arctan(1 + \sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))) \right)}{4} \right)$

3.393. $\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

input `int(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)`

output `1/d*(2/7*B*b^3*tan(d*x+c)^(7/2)+2/5*A*b^3*tan(d*x+c)^(5/2)+6/5*B*a*b^2*tan
(d*x+c)^(5/2)+2*A*a*b^2*tan(d*x+c)^(3/2)+2*B*a^2*b*tan(d*x+c)^(3/2)-2/3*B*
b^3*tan(d*x+c)^(3/2)+6*A*a^2*b*tan(d*x+c)^(1/2)-2*tan(d*x+c)^(1/2)*A*b^3+2
*B*a^3*tan(d*x+c)^(1/2)-6*tan(d*x+c)^(1/2)*B*a*b^2+1/4*(-3*A*a^2*b+A*b^3-B
*a^3+3*B*a*b^2)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(
1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*
arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)
2^(1/2)(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)
^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/
2)*tan(d*x+c)^(1/2))))`

3.393.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6157 vs. $2(379) = 758$.

Time = 1.44 (sec) , antiderivative size = 6157, normalized size of antiderivative = 14.62

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm
m="fricas")`

output Too large to include

3.393.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx \\ &= \int (A+B\tan(c+dx))(a+b\tan(c+dx))^3\sqrt{\tan(c+dx)}dx \end{aligned}$$

input `integrate(tan(d*x+c)**(1/2)*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3*sqrt(tan(c + d*x)), x)`

3.393.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 363, normalized size of antiderivative = 0.86

$$\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{120 B b^3 \tan(dx + c)^{\frac{7}{2}} + 168 (3 B a b^2 + A b^3) \tan(dx + c)^{\frac{5}{2}} + 210 \sqrt{2}((A - B)a^3 - 3(A + B)a^2 b - 3(A - B)a b^2 + (A + B)b^3) \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2} + 2\sqrt{\tan(dx + c)})\right) + 210 \sqrt{2}((A - B)a^3 - 3(A + B)a^2 b - 3(A - B)a b^2 + (A + B)b^3) \arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2} - 2\sqrt{\tan(dx + c)})\right) - 105 \sqrt{2}((A + B)a^3 + 3(A - B)a^2 b - 3(A + B)a b^2 - (A - B)b^3) \log(\sqrt{2}\sqrt{\tan(dx + c)} + \tan(dx + c) + 1) + 105 \sqrt{2}((A + B)a^3 + 3(A - B)a^2 b - 3(A + B)a b^2 - (A - B)b^3) \log(-\sqrt{2}\sqrt{\tan(dx + c)} + \tan(dx + c) + 1) + 280(3 B a^2 b + 3 A a b^2 - B b^3) \tan(dx + c)^{\frac{3}{2}} + 840(B a^3 + 3 A a^2 b - 3 B a b^2 - A b^3) \sqrt{\tan(dx + c)}}{d}$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/420*(120*B*b^3*tan(d*x + c)^(7/2) + 168*(3*B*a*b^2 + A*b^3)*tan(d*x + c)^(5/2) + 210*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 210*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - 105*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 105*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 280*(3*B*a^2*b + 3*A*a*b^2 - B*b^3)*tan(d*x + c)^(3/2) + 840*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)*sqrt(tan(d*x + c)))/d`

3.393.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.393. $\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.393.9 Mupad [B] (verification not implemented)

Time = 25.34 (sec) , antiderivative size = 6716, normalized size of antiderivative = 15.95

$$\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)
```

```
output tan(c + d*x)^(1/2)*((2*B*a^3)/d - (6*B*a*b^2)/d) - tan(c + d*x)^(1/2)*((2*
A*b^3)/d - (6*A*a^2*b)/d) - tan(c + d*x)^(3/2)*((2*B*b^3)/(3*d) - (2*B*a^2
*b)/d) - atan((((8*(4*A*b^3*d^2 - 12*A*a^2*b*d^2)*((3*A^2*a*b^5)/(2*d^2) -
(5*A^2*a^3*b^3)/d^2 - (30*A^4*a^2*b^10*d^4 - A^4*b^12*d^4 - A^4*a^12*d^4
- 255*A^4*a^4*b^8*d^4 + 452*A^4*a^6*b^6*d^4 - 255*A^4*a^8*b^4*d^4 + 30*A^4
*a^10*b^2*d^4)^(1/2)/(4*d^4) + (3*A^2*a^5*b)/(2*d^2))^(1/2))/d^3 - (16*tan
(c + d*x)^(1/2)*(A^2*a^6 - A^2*b^6 + 15*A^2*a^2*b^4 - 15*A^2*a^4*b^2))/d^2
)*((3*A^2*a*b^5)/(2*d^2) - (5*A^2*a^3*b^3)/d^2 - (30*A^4*a^2*b^10*d^4 - A^
4*b^12*d^4 - A^4*a^12*d^4 - 255*A^4*a^4*b^8*d^4 + 452*A^4*a^6*b^6*d^4 - 25
5*A^4*a^8*b^4*d^4 + 30*A^4*a^10*b^2*d^4)^(1/2)/(4*d^4) + (3*A^2*a^5*b)/(2*
d^2))^(1/2)*1i - (((8*(4*A*b^3*d^2 - 12*A*a^2*b*d^2)*((3*A^2*a*b^5)/(2*d^2)
- (5*A^2*a^3*b^3)/d^2 - (30*A^4*a^2*b^10*d^4 - A^4*b^12*d^4 - A^4*a^12*d^
4 - 255*A^4*a^4*b^8*d^4 + 452*A^4*a^6*b^6*d^4 - 255*A^4*a^8*b^4*d^4 + 30*A
^4*a^10*b^2*d^4)^(1/2)/(4*d^4) + (3*A^2*a^5*b)/(2*d^2))^(1/2))/d^3 + (16*t
an(c + d*x)^(1/2)*(A^2*a^6 - A^2*b^6 + 15*A^2*a^2*b^4 - 15*A^2*a^4*b^2))/d
^2)*((3*A^2*a*b^5)/(2*d^2) - (5*A^2*a^3*b^3)/d^2 - (30*A^4*a^2*b^10*d^4 -
A^4*b^12*d^4 - A^4*a^12*d^4 - 255*A^4*a^4*b^8*d^4 + 452*A^4*a^6*b^6*d^4 -
255*A^4*a^8*b^4*d^4 + 30*A^4*a^10*b^2*d^4)^(1/2)/(4*d^4) + (3*A^2*a^5*b)/(
2*d^2))^(1/2)*1i)/(((8*(4*A*b^3*d^2 - 12*A*a^2*b*d^2)*((3*A^2*a*b^5)/(2*d^
2) - (5*A^2*a^3*b^3)/d^2 - (30*A^4*a^2*b^10*d^4 - A^4*b^12*d^4 - A^4*a^...
```

3.394 $\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

3.394.1 Optimal result	3752
3.394.2 Mathematica [C] (verified)	3753
3.394.3 Rubi [A] (verified)	3753
3.394.4 Maple [A] (verified)	3760
3.394.5 Fricas [B] (verification not implemented)	3760
3.394.6 Sympy [F]	3761
3.394.7 Maxima [A] (verification not implemented)	3761
3.394.8 Giac [F(-1)]	3762
3.394.9 Mupad [B] (verification not implemented)	3762

3.394.1 Optimal result

Integrand size = 33, antiderivative size = 380

$$\int \frac{(a + b \tan(c + dx))^3(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx =$$

$$\frac{(3a^2b(A - B) - b^3(A - B) + a^3(A + B) - 3ab^2(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(3a^2b(A - B) - b^3(A - B) + a^3(A + B) - 3ab^2(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(a^3(A - B) - 3ab^2(A - B) - 3a^2b(A + B) + b^3(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{(a^3(A - B) - 3ab^2(A - B) - 3a^2b(A + B) + b^3(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{2b(15aAb + 14a^2B - 5b^2B) \sqrt{\tan(c + dx)}}{5d}$$

$$+ \frac{2b^2(5Ab + 9aB) \tan^{\frac{3}{2}}(c + dx)}{15d} + \frac{2bB\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}{5d}$$

output $\frac{1}{2}(3a^2b(A-B) - b^3(A-B) + a^3(A+B) - 3ab^2(A+B)) \arctan\left(\frac{-1 + 2^{1/2} \tan(dx+c)^{1/2}}{d \cdot 2^{1/2}}\right) + \frac{1}{2}(3a^2b(A-B) - b^3(A-B) + a^3(A+B) - 3ab^2(A+B)) \arctan\left(\frac{1 + 2^{1/2} \tan(dx+c)^{1/2}}{d \cdot 2^{1/2}}\right) - \frac{1}{4}(a^3(A-B) - 3ab^2(A-B) - 3a^2b(A+B) + b^3(A+B)) \ln\left(\frac{1 - 2^{1/2} \tan(dx+c)^{1/2} + \tan(dx+c)}{d \cdot 2^{1/2}}\right) + \frac{1}{4}(a^3(A-B) - 3ab^2(A-B) - 3a^2b(A+B) + b^3(A+B)) \ln\left(\frac{1 + 2^{1/2} \tan(dx+c)^{1/2} + \tan(dx+c)}{d \cdot 2^{1/2}}\right) + \frac{2}{5}b(15Aa^2b + 14B^2a^2 - 5B^2b^2) \tan(dx+c)^{1/2} + \frac{2}{5}b^2(5A^2b + 9B^2a) \tan(dx+c)^{3/2} + \frac{2}{5}b^2B \tan(dx+c)^{1/2} (a + b \tan(dx+c))^2/d$

3.394.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.59 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.40

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \frac{-15\sqrt[4]{-1}(a - ib)^3(A - iB) \arctan\left(\frac{(-1)^{3/4} \sqrt{\tan(c + dx)}}{1}\right) - 15\sqrt[4]{-1}(a + ib)^3(A + iB) \operatorname{arctanh}\left(\frac{(-1)^{3/4}}{1}\right)}{1}$$

input `Integrate[((a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output $(-15(-1)^{1/4}(a - I*b)^3(A - I*B) \operatorname{ArcTan}[-1^{3/4} \operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]] - 15(-1)^{1/4}(a + I*b)^3(A + I*B) \operatorname{ArcTanh}[-1^{3/4} \operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]] + 2*b \operatorname{Sqrt}[\operatorname{Tan}[c + d*x]] * (15*(3*a*A*b + 3*a^2*B - b^2*B) + 5*b*(A*b + 3*a*B) \operatorname{Tan}[c + d*x] + 3*b^2*B \operatorname{Tan}[c + d*x]^2)) / (15*d)$

3.394.3 Rubi [A] (verified)

Time = 1.15 (sec) , antiderivative size = 312, normalized size of antiderivative = 0.82, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 4090, 27, 3042, 4120, 27, 3042, 4113, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.394. $\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

↓ 4090

$$\frac{2}{5} \int \frac{(a + b \tan(c + dx)) (b(5Ab + 9aB) \tan^2(c + dx) + 5(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(5aA - bB))}{2\sqrt{\tan(c + dx)}} dx + \frac{2bB\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}{5d}$$

↓ 27

$$\frac{1}{5} \int \frac{(a + b \tan(c + dx)) (b(5Ab + 9aB) \tan^2(c + dx) + 5(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(5aA - bB))}{\sqrt{\tan(c + dx)}} dx + \frac{2bB\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}{5d}$$

↓ 3042

$$\frac{1}{5} \int \frac{(a + b \tan(c + dx)) (b(5Ab + 9aB) \tan(c + dx)^2 + 5(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(5aA - bB))}{\sqrt{\tan(c + dx)}} dx + \frac{2bB\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}{5d}$$

↓ 4120

$$\frac{1}{5} \left(\frac{2b^2(9aB + 5Ab) \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2}{3} \int - \frac{3((5aA - bB)a^2 + b(14Ba^2 + 15Aba - 5b^2B) \tan^2(c + dx) + 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx))}{2\sqrt{\tan(c + dx)}} dx + \frac{2bB\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}{5d} \right)$$

↓ 27

$$\frac{1}{5} \left(\int \frac{(5aA - bB)a^2 + b(14Ba^2 + 15Aba - 5b^2B) \tan^2(c + dx) + 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2bB\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(\int \frac{(5aA - bB)a^2 + b(14Ba^2 + 15Aba - 5b^2B) \tan(c + dx)^2 + 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2bB \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}{5d} \right)$$

↓ 4113

$$\frac{1}{5} \left(\int \frac{5(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2b(14a^2B + 15aAb - 5b^2B) \sqrt{\tan(c + dx)}}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(\int \frac{5(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2b(14a^2B + 15aAb - 5b^2B) \sqrt{\tan(c + dx)}}{5d} \right)$$

↓ 4017

$$\frac{1}{5} \left(2 \int \frac{5(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} + \frac{2b(14a^2B + 15aAb - 5b^2B) \sqrt{\tan(c + dx)}}{d} \right)$$

↓ 27

$$\frac{1}{5} \left(10 \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} + \frac{2b(14a^2B + 15aAb - 5b^2B) \sqrt{\tan(c + dx)}}{d} \right)$$

↓ 1482

$$\frac{1}{5} \left(10 \left(\frac{1}{2}(a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} + \frac{1}{2}(a^3(A + B) - 3a^2b(A - B) - 3ab^2(A + B) + b^3(A - B)) \int \frac{1 + \tan(c + dx)}{\tan^2(c + dx) + 1} d \sqrt{\tan(c + dx)} \right) + \frac{2b(14a^2B + 15aAb - 5b^2B) \sqrt{\tan(c + dx)}}{d} \right)$$

↓ 1476

3.394. $\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^3(A + B) - \dots \right)}{\frac{2bB\sqrt{\tan(c+dx)}(a + b \tan(c+dx))^2}{5d}} \right) \downarrow 1082$$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^3(A + B) - \dots \right)}{\frac{2bB\sqrt{\tan(c+dx)}(a + b \tan(c+dx))^2}{5d}} \right) \downarrow 217$$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^3(A + B) - \dots \right)}{d \frac{2bB\sqrt{\tan(c+dx)}(a + b \tan(c+dx))^2}{5d}} \right) \downarrow 1479$$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \dots \right) \right)}{\frac{2bB\sqrt{\tan(c+dx)}(a + b \tan(c+dx))^2}{5d}} \right) \downarrow 25$$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^3(A-B) - 3a^2b(A+B) - 3ab^2(A-B) + b^3(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{2bB\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2} + \frac{1}{2} \int \frac{\sqrt{2}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right)$$

\downarrow 27

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^3(A-B) - 3a^2b(A+B) - 3ab^2(A-B) + b^3(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{2bB\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2} + \frac{1}{2} \int \frac{\sqrt{2}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right)$$

\downarrow 1103

$$\frac{1}{5} \left(\frac{2b(14a^2B + 15aAb - 5b^2B) \sqrt{\tan(c+dx)}}{d} + \frac{10 \left(\frac{1}{2} (a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \right)}{2bB\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2} \right)$$

input `Int[((a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `(2*b*B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^2)/(5*d) + ((10*(((3*a^2*b*(A - B) - b^3*(A - B) + a^3*(A + B) - 3*a*b^2*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])))/2 + ((a^3*(A - B) - 3*a*b^2*(A - B) - 3*a^2*b*(A + B) + b^3*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d + (2*b*(15*a*A*b + 14*a^2*B - 5*b^2*B)*Sqrt[Tan[c + d*x]]/d + (2*b^2*(5*A*b + 9*a*B)*Tan[c + d*x]^(3/2))/(3*d))/5`

3.394.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4090 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4120 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]`

3.394.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 314, normalized size of antiderivative = 0.83

method	result
derivativedivides	$\frac{2B b^3 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{2A b^3 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} + 2Ba b^2 \left(\tan^{\frac{3}{2}}(dx+c)\right) + 6(\sqrt{\tan(dx+c)}) A a b^2 + 6(\sqrt{\tan(dx+c)}) B a^2 b - 2(\sqrt{\tan(dx+c)}) A a^2 b^2$
default	$\frac{2B b^3 \left(\tan^{\frac{5}{2}}(dx+c)\right)}{5} + \frac{2A b^3 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} + 2Ba b^2 \left(\tan^{\frac{3}{2}}(dx+c)\right) + 6(\sqrt{\tan(dx+c)}) A a b^2 + 6(\sqrt{\tan(dx+c)}) B a^2 b - 2(\sqrt{\tan(dx+c)}) A a^2 b^2$
parts	$(A b^3 + 3B a b^2) \left(\frac{2 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} - \frac{\sqrt{2} \left(\ln \left(\frac{1 - \sqrt{2} (\sqrt{\tan(dx+c)} + \tan(dx+c))}{1 + \sqrt{2} (\sqrt{\tan(dx+c)} + \tan(dx+c))} \right) + 2 \arctan(1 + \sqrt{2} (\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1 + \sqrt{2} (\sqrt{\tan(dx+c)}) \right)}{4} \right)$

input `int((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x,method=_RETURNV
ERBOSE)`

output `1/d*(2/5*B*b^3*tan(d*x+c)^(5/2)+2/3*A*b^3*tan(d*x+c)^(3/2)+2*B*a*b^2*tan(d
*x+c)^(3/2)+6*tan(d*x+c)^(1/2)*A*a*b^2+6*tan(d*x+c)^(1/2)*B*a^2*b-2*tan(d*
x+c)^(1/2)*B*b^3+1/4*(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*2^(1/2)*(ln((1+2^(1
/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+
2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))
)+1/4*(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(
1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1
/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))`

3.394.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6136 vs. 2(340) = 680.

Time = 1.49 (sec) , antiderivative size = 6136, normalized size of antiderivative = 16.15

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorith
m="fricas")`

output Too large to include

3.394.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^3}{\sqrt{\tan(c + dx)}} dx$$

input `integrate((a+b*tan(d*x+c))**3*(A+B*tan(d*x+c))/tan(d*x+c)**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**3/sqrt(tan(c + d*x)), x)`

3.394.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 327, normalized size of antiderivative = 0.86

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \frac{24 B b^3 \tan(dx + c)^{\frac{5}{2}} + 30 \sqrt{2} ((A + B) a^3 + 3(A - B) a^2 b - 3(A + B) a b^2 - (A - B) b^3) \arctan\left(\frac{1}{2} \sqrt{2} (\sqrt{\tan(dx + c)} + \tan(dx + c))\right) + 30 \sqrt{2} ((A + B) a^3 + 3(A - B) a^2 b - 3(A + B) a b^2 - (A - B) b^3) \arctan\left(\frac{1}{2} \sqrt{2} (\sqrt{\tan(dx + c)} - \tan(dx + c))\right) + 15 \sqrt{2} ((A - B) a^3 - 3(A + B) a^2 b - 3(A - B) a b^2 + (A + B) b^3) \log(\sqrt{2} \sqrt{\tan(dx + c)} + \tan(dx + c) + 1) - 15 \sqrt{2} ((A - B) a^3 - 3(A + B) a^2 b - 3(A - B) a b^2 + (A + B) b^3) \log(-\sqrt{2} \sqrt{\tan(dx + c)} + \tan(dx + c) + 1) + 40(3 B a^2 b + A b^3) \tan(dx + c)^{\frac{3}{2}} + 120(3 B a^2 b + 3 A a b^2 - B b^3) \sqrt{\tan(dx + c)}}{60}$$

input `integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="maxima")`

output `1/60*(24*B*b^3*tan(d*x + c)^(5/2) + 30*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 30*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + 15*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - 15*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 40*(3*B*a^2*b + A*b^3)*tan(d*x + c)^(3/2) + 120*(3*B*a^2*b + 3*A*a*b^2 - B*b^3)*sqrt(tan(d*x + c)))/d`

3.394.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Timed out}$$

```
input integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="giac")
```

output Timed out

3.394.9 Mupad [B] (verification not implemented)

Time = 14.53 (sec) , antiderivative size = 6657, normalized size of antiderivative = 17.52

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Too large to display}$$

```
input int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3)/tan(c + d*x)^(1/2),x)
```

```
output atan((((8*(4*A*a^3*d^2 - 12*A*a*b^2*d^2)*((5*A^2*a^3*b^3)/d^2 - (30*A^4*a^2*b^10*d^4 - A^4*b^12*d^4 - A^4*a^12*d^4 - 255*A^4*a^4*b^8*d^4 + 452*A^4*a^6*b^6*d^4 - 255*A^4*a^8*b^4*d^4 + 30*A^4*a^10*b^2*d^4)^(1/2)/(4*d^4) - (3*A^2*a*b^5)/(2*d^2) - (3*A^2*a^5*b)/(2*d^2))^(1/2))/d^3 - (16*tan(c + d*x)^(1/2)*(A^2*a^6 - A^2*b^6 + 15*A^2*a^2*b^4 - 15*A^2*a^4*b^2))/d^2)*((5*A^2*a^3*b^3)/d^2 - (30*A^4*a^2*b^10*d^4 - A^4*b^12*d^4 - A^4*a^12*d^4 - 255*A^4*a^4*b^8*d^4 + 452*A^4*a^6*b^6*d^4 - 255*A^4*a^8*b^4*d^4 + 30*A^4*a^10*b^2*d^4)^(1/2)/(4*d^4) - (3*A^2*a*b^5)/(2*d^2) - (3*A^2*a^5*b)/(2*d^2))^(1/2)*1i - ((8*(4*A*a^3*d^2 - 12*A*a*b^2*d^2)*((5*A^2*a^3*b^3)/d^2 - (30*A^4*a^2*b^10*d^4 - A^4*b^12*d^4 - A^4*a^12*d^4 - 255*A^4*a^4*b^8*d^4 + 452*A^4*a^6*b^6*d^4 - 255*A^4*a^8*b^4*d^4 + 30*A^4*a^10*b^2*d^4)^(1/2)/(4*d^4) - (3*A^2*a*b^5)/(2*d^2) - (3*A^2*a^5*b)/(2*d^2))^(1/2))/d^3 + (16*tan(c + d*x)^(1/2)*(A^2*a^6 - A^2*b^6 + 15*A^2*a^2*b^4 - 15*A^2*a^4*b^2))/d^2)*((5*A^2*a^3*b^3)/d^2 - (30*A^4*a^2*b^10*d^4 - A^4*b^12*d^4 - A^4*a^12*d^4 - 255*A^4*a^4*b^8*d^4 + 452*A^4*a^6*b^6*d^4 - 255*A^4*a^8*b^4*d^4 + 30*A^4*a^10*b^2*d^4)^(1/2)/(4*d^4) - (3*A^2*a*b^5)/(2*d^2) - (3*A^2*a^5*b)/(2*d^2))^(1/2)*1i)/((16*(3*A^3*a^8*b - A^3*b^9 + 6*A^3*a^4*b^5 + 8*A^3*a^6*b^3))/d^3 + ((8*(4*A*a^3*d^2 - 12*A*a*b^2*d^2)*((5*A^2*a^3*b^3)/d^2 - (30*A^4*a^2*b^10*d^4 - A^4*b^12*d^4 - A^4*a^12*d^4 - 255*A^4*a^4*b^8*d^4 + 452*A^4*a^6*b^6*d^4 - 255*A^4*a^8*b^4*d^4 + 30*A^4*a^10*b^2*d^4)^(1/2)/(4*d^4) - (3...
```

3.395
$$\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.395.1 Optimal result 3763
 3.395.2 Mathematica [C] (verified) 3764
 3.395.3 Rubi [A] (verified) 3765
 3.395.4 Maple [A] (verified) 3771
 3.395.5 Fricas [B] (verification not implemented) 3771
 3.395.6 Sympy [F] 3772
 3.395.7 Maxima [A] (verification not implemented) 3772
 3.395.8 Giac [F(-1)] 3773
 3.395.9 Mupad [B] (verification not implemented) 3773

3.395.1 Optimal result

Integrand size = 33, antiderivative size = 374

$$\int \frac{(a + b \tan(c + dx))^3(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{(a^3(A - B) - 3ab^2(A - B) - 3a^2b(A + B) + b^3(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(a^3(A - B) - 3ab^2(A - B) - 3a^2b(A + B) + b^3(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(3a^2b(A - B) - b^3(A - B) + a^3(A + B) - 3ab^2(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{(3a^2b(A - B) - b^3(A - B) + a^3(A + B) - 3ab^2(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{2b(2a^2A + Ab^2 + 3abB) \sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2aA(a + b \tan(c + dx))^2}{d\sqrt{\tan(c + dx)}}$$

output
$$-1/2*(a^3*(A-B)-3*a*b^2*(A-B)-3*a^2*b*(A+B)+b^3*(A+B))*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/d*2^{(1/2)}-1/2*(a^3*(A-B)-3*a*b^2*(A-B)-3*a^2*b*(A+B)+b^3*(A+B))*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/d*2^{(1/2)}-1/4*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/d*2^{(1/2)}+1/4*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/d*2^{(1/2)}+2*b*(2*A*a^2+A*b^2+3*B*a*b)*\tan(d*x+c)^{(1/2)}/d+2/3*b^2*(3*A*a+B*b)*\tan(d*x+c)^{(3/2)}/d-2*a*A*(a+b*\tan(d*x+c))^2/d/\tan(d*x+c)^{(1/2)}$$

3.395.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 2.86 (sec) , antiderivative size = 264, normalized size of antiderivative = 0.71

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{8b(-12aAb - 17a^2B + 3b^2B) - 3\left(8(a^3A - 3aAb^2 - 3a^2bB + b^3B) \operatorname{Hypergeometric2F1}\left(-\frac{1}{4}, 1, \frac{3}{4}, -\tan^2\right)\right)}{\tan^{\frac{3}{2}}(c + dx)}$$

input `Integrate[((a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output
$$(8*b*(-12*a*A*b - 17*a^2*B + 3*b^2*B) - 3*(8*(a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*\operatorname{Hypergeometric2F1}[-1/4, 1, 3/4, -\operatorname{Tan}[c + d*x]^2] + \operatorname{Sqrt}[2]*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*(2*\operatorname{ArcTan}[1 - \operatorname{Sqrt}[2]*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]] - 2*\operatorname{ArcTan}[1 + \operatorname{Sqrt}[2]*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]] + \operatorname{Log}[1 - \operatorname{Sqrt}[2]*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]] + \operatorname{Tan}[c + d*x]] - \operatorname{Log}[1 + \operatorname{Sqrt}[2]*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]] + \operatorname{Tan}[c + d*x]])*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]] + 8*b*(3*A*b + 7*a*B)*(a + b*\operatorname{Tan}[c + d*x]) + 8*b*B*(a + b*\operatorname{Tan}[c + d*x])^2)/(12*d*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]])$$

3.395.3 Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 303, normalized size of antiderivative = 0.81, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 4088, 27, 3042, 4120, 27, 3042, 4113, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx$$

$$\downarrow \text{4088}$$

$$2 \int \frac{(a + b \tan(c + dx)) (b(3aA + bB) \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(5Ab + aB))}{\frac{2\sqrt{\tan(c + dx)}}{2aA(a + b \tan(c + dx))^2} d\sqrt{\tan(c + dx)}} dx -$$

$$\downarrow \text{27}$$

$$\int \frac{(a + b \tan(c + dx)) (b(3aA + bB) \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(5Ab + aB))}{\frac{\sqrt{\tan(c + dx)}}{2aA(a + b \tan(c + dx))^2} d\sqrt{\tan(c + dx)}} dx -$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx)) (b(3aA + bB) \tan(c + dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(5Ab + aB))}{\frac{\sqrt{\tan(c + dx)}}{2aA(a + b \tan(c + dx))^2} d\sqrt{\tan(c + dx)}} dx -$$

$$\downarrow \text{4120}$$

$$-\frac{2}{3} \int -\frac{3((5Ab + aB)a^2 + b(2Aa^2 + 3bBa + Ab^2) \tan^2(c + dx) - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx))}{2\sqrt{\tan(c + dx)}} dx +$$

$$\frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2aA(a + b \tan(c + dx))^2}{d\sqrt{\tan(c + dx)}}$$

↓ 27

$$\int \frac{(5Ab + aB)a^2 + b(2Aa^2 + 3bBa + Ab^2) \tan^2(c + dx) - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx +$$

$$\frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2aA(a + b \tan(c + dx))^2}{d\sqrt{\tan(c + dx)}}$$

↓ 3042

$$\int \frac{(5Ab + aB)a^2 + b(2Aa^2 + 3bBa + Ab^2) \tan(c + dx)^2 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx +$$

$$\frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2aA(a + b \tan(c + dx))^2}{d\sqrt{\tan(c + dx)}}$$

↓ 4113

$$\int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx +$$

$$\frac{2b(2a^2A + 3abB + Ab^2) \sqrt{\tan(c + dx)}}{d} + \frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2aA(a + b \tan(c + dx))^2}{d\sqrt{\tan(c + dx)}}$$

↓ 3042

$$\int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx +$$

$$\frac{2b(2a^2A + 3abB + Ab^2) \sqrt{\tan(c + dx)}}{d} + \frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2aA(a + b \tan(c + dx))^2}{d\sqrt{\tan(c + dx)}}$$

↓ 4017

$$2 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} +$$

$$\frac{d}{d} + \frac{2b(2a^2A + 3abB + Ab^2) \sqrt{\tan(c + dx)}}{d} + \frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2aA(a + b \tan(c + dx))^2}{d\sqrt{\tan(c + dx)}}$$

↓ 1482

3.395. $\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$

$$\frac{2\left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^3(A-B) - 3a^2b(A+B) + 3ab^2(A-B) + b^3(A+B)) \int \frac{1+\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right)}{d} + \frac{2b(2a^2A + 3abB + Ab^2) \sqrt{\tan(c+dx)}}{d} + \frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2aA(a + b \tan(c+dx))^2}{d\sqrt{\tan(c+dx)}}$$

↓ 1476

$$\frac{2\left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^3(A-B) - 3a^2b(A+B) + 3ab^2(A-B) + b^3(A+B)) \int \frac{1+\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right)}{d} + \frac{2b(2a^2A + 3abB + Ab^2) \sqrt{\tan(c+dx)}}{d} + \frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2aA(a + b \tan(c+dx))^2}{d\sqrt{\tan(c+dx)}}$$

↓ 1082

$$\frac{2\left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^3(A-B) - 3a^2b(A+B) + 3ab^2(A-B) + b^3(A+B)) \int \frac{1+\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right)}{d} + \frac{2b(2a^2A + 3abB + Ab^2) \sqrt{\tan(c+dx)}}{d} + \frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2aA(a + b \tan(c+dx))^2}{d\sqrt{\tan(c+dx)}}$$

↓ 217

$$\frac{2\left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a^3(A-B) - 3a^2b(A+B) + 3ab^2(A-B) + b^3(A+B)) \int \frac{1+\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right)}{d} + \frac{2b(2a^2A + 3abB + Ab^2) \sqrt{\tan(c+dx)}}{d} + \frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2aA(a + b \tan(c+dx))^2}{d\sqrt{\tan(c+dx)}}$$

↓ 1479

$$\frac{2\left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}}\right) - \frac{1}{2}(a^3(A-B) - 3a^2b(A+B) + 3ab^2(A-B) + b^3(A+B)) \int \frac{1+\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right)}{d} + \frac{2b(2a^2A + 3abB + Ab^2) \sqrt{\tan(c+dx)}}{d} + \frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2aA(a + b \tan(c+dx))^2}{d\sqrt{\tan(c+dx)}}$$

↓ 25

3.395. $\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& 2 \left(\frac{1}{2} (a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right. \\
& \left. \frac{2b(2a^2A + 3abB + Ab^2) \sqrt{\tan(c+dx)}}{d} + \frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2aA(a + b \tan(c+dx))^2}{d\sqrt{\tan(c+dx)}} \right) \\
& \quad \downarrow \text{27} \\
& 2 \left(\frac{1}{2} (a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right. \\
& \left. \frac{2b(2a^2A + 3abB + Ab^2) \sqrt{\tan(c+dx)}}{d} + \frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2aA(a + b \tan(c+dx))^2}{d\sqrt{\tan(c+dx)}} \right) \\
& \quad \downarrow \text{1103} \\
& \frac{2b(2a^2A + 3abB + Ab^2) \sqrt{\tan(c+dx)}}{d} + \\
& 2 \left(\frac{1}{2} (a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right. \\
& \left. \frac{2b^2(3aA + bB) \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2aA(a + b \tan(c+dx))^2}{d\sqrt{\tan(c+dx)}} \right)
\end{aligned}$$

input `Int[((a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `(2*(-1/2*((a^3*(A - B) - 3*a*b^2*(A - B) - 3*a^2*b*(A + B) + b^3*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]/Sqrt[2])) + ((3*a^2*b*(A - B) - b^3*(A - B) + a^3*(A + B) - 3*a*b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/d + (2*b*(2*a^2*A + A*b^2 + 3*a*b*B)*Sqrt[Tan[c + d*x]])/d + (2*b^2*(3*a*A + b*B)*Tan[c + d*x]^(3/2))/(3*d) - (2*a*A*(a + b*Tan[c + d*x])^2)/(d*Sqrt[Tan[c + d*x]])`

3.395. $\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

3.395.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4113 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4120 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^((c_) + (d_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]`

3.395.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 286, normalized size of antiderivative = 0.76

method	result
derivativedivides	$\frac{2B b^3 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} + 2(\sqrt{\tan(dx+c)}) A b^3 + 6(\sqrt{\tan(dx+c)}) B a b^2 - \frac{2A a^3}{\sqrt{\tan(dx+c)}} + \frac{(3A a^2 b - A b^3 + B a^3 - 3B a b^2) \sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}\right)\right)}{4}$
default	$\frac{2B b^3 \left(\tan^{\frac{3}{2}}(dx+c)\right)}{3} + 2(\sqrt{\tan(dx+c)}) A b^3 + 6(\sqrt{\tan(dx+c)}) B a b^2 - \frac{2A a^3}{\sqrt{\tan(dx+c)}} + \frac{(3A a^2 b - A b^3 + B a^3 - 3B a b^2) \sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}\right)\right)}{4}$
parts	$\frac{(A b^3 + 3B a b^2) \left(2(\sqrt{\tan(dx+c)}) - \frac{\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}\right)\right)}{4} + 2 \arctan\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})}{1-\sqrt{2}(\sqrt{\tan(dx+c)})}\right) + 2 \arctan\left(\frac{-1+\sqrt{2}(\sqrt{\tan(dx+c)})}{1-\sqrt{2}(\sqrt{\tan(dx+c)})}\right)\right)}{d}$

input `int((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x,method=_RETURNV ERBOSE)`

output `1/d*(2/3*B*b^3*tan(d*x+c)^(3/2)+2*tan(d*x+c)^(1/2)*A*b^3+6*tan(d*x+c)^(1/2)*B*a*b^2-2*A*a^3/tan(d*x+c)^(1/2)+1/4*(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(-A*a^3+3*A*a*b^2+3*B*a^2*b-B*b^3)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))`

3.395.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6137 vs. 2(338) = 676.

Time = 1.47 (sec) , antiderivative size = 6137, normalized size of antiderivative = 16.41

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="fracas")`

output Too large to include

3.395.
$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

3.395.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^3}{\tan^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**3*(A+B*tan(d*x+c))/tan(d*x+c)**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**3/tan(c + d*x)**(3/2), x)`

3.395.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 310, normalized size of antiderivative = 0.83

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{8 B b^3 \tan(dx + c)^{\frac{3}{2}} - \frac{24 A a^3}{\sqrt{\tan(dx+c)}} - 6 \sqrt{2}((A - B)a^3 - 3(A + B)a^2 b - 3(A - B)ab^2 + (A + B)b^3) \arctan}{1}$$

input `integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="maxima")`

output `1/12*(8*B*b^3*tan(d*x + c)^(3/2) - 24*A*a^3/sqrt(tan(d*x + c)) - 6*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) - 6*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + 3*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - 3*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 24*(3*B*a*b^2 + A*b^3)*sqrt(tan(d*x + c))/d`

3.395.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="giac")
```

```
output Timed out
```

3.395.9 Mupad [B] (verification not implemented)

Time = 12.19 (sec) , antiderivative size = 7108, normalized size of antiderivative = 19.01

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Too large to display}$$

```
input int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3)/tan(c + d*x)^(3/2),x)
```

```
output atan((((8*(4*B*a^3*d^2 - 12*B*a*b^2*d^2)*((5*B^2*a^3*b^3)/d^2 - (30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2)/(4*d^4) - (3*B^2*a*b^5)/(2*d^2) - (3*B^2*a^5*b)/(2*d^2))^(1/2))/d^3 - (16*tan(c + d*x)^(1/2)*(B^2*a^6 - B^2*b^6 + 15*B^2*a^2*b^4 - 15*B^2*a^4*b^2))/d^2)*((5*B^2*a^3*b^3)/d^2 - (30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2)/(4*d^4) - (3*B^2*a*b^5)/(2*d^2) - (3*B^2*a^5*b)/(2*d^2))^(1/2)*1i - (((8*(4*B*a^3*d^2 - 12*B*a*b^2*d^2)*((5*B^2*a^3*b^3)/d^2 - (30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2)/(4*d^4) - (3*B^2*a*b^5)/(2*d^2) - (3*B^2*a^5*b)/(2*d^2))^(1/2))/d^3 + (16*tan(c + d*x)^(1/2)*(B^2*a^6 - B^2*b^6 + 15*B^2*a^2*b^4 - 15*B^2*a^4*b^2))/d^2)*((5*B^2*a^3*b^3)/d^2 - (30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2)/(4*d^4) - (3*B^2*a*b^5)/(2*d^2) - (3*B^2*a^5*b)/(2*d^2))^(1/2)*1i)/(((8*(4*B*a^3*d^2 - 12*B*a*b^2*d^2)*((5*B^2*a^3*b^3)/d^2 - (30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2)/(4*d^4) - (3*B^2*a*b^5)/(2*d^2) - (3*B^2*a^5*b)/(2*d^2))^(1/2))/d^3 - (16*tan(c...
```

3.396
$$\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.396.1 Optimal result	3774
3.396.2 Mathematica [C] (verified)	3775
3.396.3 Rubi [A] (verified)	3775
3.396.4 Maple [A] (verified)	3782
3.396.5 Fricas [B] (verification not implemented)	3782
3.396.6 Sympy [F]	3783
3.396.7 Maxima [A] (verification not implemented)	3783
3.396.8 Giac [F(-1)]	3784
3.396.9 Mupad [B] (verification not implemented)	3784

3.396.1 Optimal result

Integrand size = 33, antiderivative size = 372

$$\int \frac{(a + b \tan(c + dx))^3(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{(3a^2b(A - B) - b^3(A - B) + a^3(A + B) - 3ab^2(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(3a^2b(A - B) - b^3(A - B) + a^3(A + B) - 3ab^2(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(a^3(A - B) - 3ab^2(A - B) - 3a^2b(A + B) + b^3(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(a^3(A - B) - 3ab^2(A - B) - 3a^2b(A + B) + b^3(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$- \frac{2a^2(7Ab + 3aB)}{3d\sqrt{\tan(c + dx)}} + \frac{2b^2(aA + 3bB)\sqrt{\tan(c + dx)}}{3d} - \frac{2aA(a + b \tan(c + dx))^2}{3d \tan^{\frac{3}{2}}(c + dx)}$$

output
$$-1/2*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/d*2^{(1/2)}-1/2*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/d*2^{(1/2)}+1/4*(a^3*(A-B)-3*a*b^2*(A-B)-3*a^2*b*(A+B)+b^3*(A+B))*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/d*2^{(1/2)}-1/4*(a^3*(A-B)-3*a*b^2*(A-B)-3*a^2*b*(A+B)+b^3*(A+B))*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/d*2^{(1/2)}-2/3*a^2*(7*A*b+3*B*a)/d/\tan(d*x+c)^{(1/2)}+2/3*b^2*(A*a+3*B*b)*\tan(d*x+c)^{(1/2)}/d-2/3*a*A*(a+b*\tan(d*x+c))^2/d/\tan(d*x+c)^{(3/2)}$$

3.396.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.33 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.44

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx =$$

$$\frac{2((a^3 A - 3aAb^2 - 3a^2bB + b^3 B) \text{Hypergeometric2F1}(-\frac{3}{4}, 1, \frac{1}{4}, -\tan^2(c + dx)) + 3(3a^2 Ab - Ab^3 + a^3 B) \text{Hypergeometric2F1}(-\frac{3}{4}, 1, \frac{1}{4}, -\tan^2(c + dx)))}{3d \tan^{\frac{3}{2}}(c + dx)}$$

input `Integrate[((a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output
$$(-2*((a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*\text{Hypergeometric2F1}[-3/4, 1, 1/4, -\tan[c + d*x]^2] + 3*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*\text{Hypergeometric2F1}[-1/4, 1, 3/4, -\tan[c + d*x]^2]*\tan[c + d*x] + b*(3*a*A*b + 3*a^2*B - b^2*B + 3*b*(A*b + 3*a*B)*\tan[c + d*x] - 3*b^2*B*\tan[c + d*x]^2))/(3*d*\tan[c + d*x]^{(3/2)})$$

3.396.3 Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 302, normalized size of antiderivative = 0.81, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 4088, 27, 3042, 4118, 25, 3042, 4113, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.396.
$$\int \frac{(a+b \tan(c+dx))^3 (A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx$$

↓ 4088

$$\frac{2}{3} \int \frac{(a + b \tan(c + dx)) (b(aA + 3bB) \tan^2(c + dx) - 3(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(7Ab + 3aB))}{\frac{2 \tan^{\frac{3}{2}}(c + dx)}{2aA(a + b \tan(c + dx))^2} \cdot 3d \tan^{\frac{3}{2}}(c + dx)} dx -$$

↓ 27

$$\frac{1}{3} \int \frac{(a + b \tan(c + dx)) (b(aA + 3bB) \tan^2(c + dx) - 3(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(7Ab + 3aB))}{\frac{\tan^{\frac{3}{2}}(c + dx)}{2aA(a + b \tan(c + dx))^2} \cdot 3d \tan^{\frac{3}{2}}(c + dx)} dx -$$

↓ 3042

$$\frac{1}{3} \int \frac{(a + b \tan(c + dx)) (b(aA + 3bB) \tan(c + dx)^2 - 3(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(7Ab + 3aB))}{\frac{\tan(c + dx)^{3/2}}{2aA(a + b \tan(c + dx))^2} \cdot 3d \tan^{\frac{3}{2}}(c + dx)} dx -$$

↓ 4118

$$\frac{1}{3} \left(\int - \frac{b^2(aA + 3bB) \tan^2(c + dx) + 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(3Aa^2 - 9bBa - 10Ab^2)}{\frac{\sqrt{\tan(c + dx)}}{2aA(a + b \tan(c + dx))^2} \cdot 3d \tan^{\frac{3}{2}}(c + dx)} dx - \right.$$

↓ 25

$$\frac{1}{3} \left(- \int \frac{b^2(aA + 3bB) \tan^2(c + dx) + 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(3Aa^2 - 9bBa - 10Ab^2)}{\frac{\sqrt{\tan(c + dx)}}{2aA(a + b \tan(c + dx))^2} \cdot 3d \tan^{\frac{3}{2}}(c + dx)} dx - \right.$$

↓ 3042

3.396. $\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$

$$\frac{1}{3} \left(- \int \frac{-b^2(aA + 3bB) \tan(c + dx)^2 + 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(3Aa^2 - 9bBa - 10Ab^2)}{\sqrt{\tan(c + dx)}} \frac{2aA(a + b \tan(c + dx))^2}{3d \tan^{\frac{3}{2}}(c + dx)} dx \right)$$

↓ 4113

$$\frac{1}{3} \left(- \int \frac{3(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2a^2(3aB + 7Ab)}{d\sqrt{\tan(c + dx)}} + \frac{2aA(a + b \tan(c + dx))^2}{3d \tan^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{3} \left(- \int \frac{3(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx - \frac{2a^2(3aB + 7Ab)}{d\sqrt{\tan(c + dx)}} + \frac{2aA(a + b \tan(c + dx))^2}{3d \tan^{\frac{3}{2}}(c + dx)} \right)$$

↓ 4017

$$\frac{1}{3} \left(- \frac{2 \int \frac{3(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{d} - \frac{2a^2(3aB + 7Ab)}{d\sqrt{\tan(c + dx)}} + \frac{2b^2(aA + 3bB)}{d} + \frac{2aA(a + b \tan(c + dx))^2}{3d \tan^{\frac{3}{2}}(c + dx)} \right)$$

↓ 27

$$\frac{1}{3} \left(- \frac{6 \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{d} - \frac{2a^2(3aB + 7Ab)}{d\sqrt{\tan(c + dx)}} + \frac{2b^2(aA + 3bB)}{d} + \frac{2aA(a + b \tan(c + dx))^2}{3d \tan^{\frac{3}{2}}(c + dx)} \right)$$

↓ 1482

$$\frac{1}{3} \left(- \frac{6 \left(\frac{1}{2}(a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} + \frac{1}{2}(a^3(A + B) + b^3(B + A)) \int \frac{1 + \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} \right)}{d} + \frac{2aA(a + b \tan(c + dx))^2}{3d \tan^{\frac{3}{2}}(c + dx)} \right)$$

3.396. $\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$

$$\downarrow 1476$$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2} (a^3(A-B) - 3a^2b(A+B) - 3ab^2(A-B) + b^3(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^3(A+B)) \right)}{\frac{2aA(a+b\tan(c+dx))^2}{3d\tan^{\frac{3}{2}}(c+dx)}} \right)$$

$$\downarrow 1082$$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2} (a^3(A-B) - 3a^2b(A+B) - 3ab^2(A-B) + b^3(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^3(A+B)) \right)}{\frac{2aA(a+b\tan(c+dx))^2}{3d\tan^{\frac{3}{2}}(c+dx)}} \right)$$

$$\downarrow 217$$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2} (a^3(A-B) - 3a^2b(A+B) - 3ab^2(A-B) + b^3(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^3(A+B)) \right)}{\frac{2aA(a+b\tan(c+dx))^2}{3d\tan^{\frac{3}{2}}(c+dx)}} \right) d$$

$$\downarrow 1479$$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2} (a^3(A-B) - 3a^2b(A+B) - 3ab^2(A-B) + b^3(A+B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \int \right)}{\frac{2aA(a+b\tan(c+dx))^2}{3d\tan^{\frac{3}{2}}(c+dx)}} \right)$$

$$\downarrow 25$$

3.396. $\int \frac{(a+b\tan(c+dx))^3(A+B\tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^3(A-B) - 3a^2b(A+B) - 3ab^2(A-B) + b^3(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right)}{\frac{2aA(a+b\tan(c+dx))^2}{3d\tan^{\frac{3}{2}}(c+dx)}} \right)$$

$$\frac{2aA(a+b\tan(c+dx))^2}{3d\tan^{\frac{3}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^3(A-B) - 3a^2b(A+B) - 3ab^2(A-B) + b^3(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right)}{\frac{2aA(a+b\tan(c+dx))^2}{3d\tan^{\frac{3}{2}}(c+dx)}} \right)$$

$$\frac{2aA(a+b\tan(c+dx))^2}{3d\tan^{\frac{3}{2}}(c+dx)}$$

↓ 1103

$$\frac{1}{3} \left(\frac{2a^2(3aB+7Ab)}{d\sqrt{\tan(c+dx)}} - \frac{6 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right)}{\frac{2aA(a+b\tan(c+dx))^2}{3d\tan^{\frac{3}{2}}(c+dx)}} \right)$$

$$\frac{2aA(a+b\tan(c+dx))^2}{3d\tan^{\frac{3}{2}}(c+dx)}$$

input `Int[((a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `((-6*(((3*a^2*b*(A - B) - b^3*(A - B) + a^3*(A + B) - 3*a*b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]/Sqrt[2]] + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]/Sqrt[2]]))/2 + ((a^3*(A - B) - 3*a*b^2*(A - B) - 3*a^2*b*(A + B) + b^3*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]/Sqrt[2]] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]/(2*Sqrt[2])]))/2)/d - (2*a^2*(7*A*b + 3*a*B))/(d*Sqrt[Tan[c + d*x]] + (2*b^2*(a*A + 3*b*B)*Sqrt[Tan[c + d*x]])/d)/3 - (2*a*A*(a + b*Tan[c + d*x])^2)/(3*d*Tan[c + d*x]^(3/2))`

3.396.3.1 Defintions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, x], x]$
- rule 27 $\text{Int}[(a_)*(\text{Fx}_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[\text{Fx}, (b_)*(\text{Gx}_) /; \text{FreeQ}[b, x]]$
- rule 217 $\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 1082 $\text{Int}[(a_) + (b_)*(x_) + (c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*c*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \quad \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_) + (e_)*(x_)/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$
- rule 1476 $\text{Int}[(d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Simp}[e/(2*c) \quad \text{Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Simp}[e/(2*c) \quad \text{Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{PosQ}[d*e]$
- rule 1479 $\text{Int}[(d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2*(d/e), 2]\}, \text{Simp}[e/(2*c*q) \quad \text{Int}[(q - 2*x)/\text{Simp}[d/e + q*x - x^2, x], x], x] + \text{Simp}[e/(2*c*q) \quad \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{NegQ}[d*e]$
- rule 1482 $\text{Int}[(d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a*c, 2]\}, \text{Simp}[(d*q + a*e)/(2*a*c) \quad \text{Int}[(q + c*x^2)/(a + c*x^4), x], x] + \text{Simp}[(d*q - a*e)/(2*a*c) \quad \text{Int}[(q - c*x^2)/(a + c*x^4), x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{NeQ}[c*d^2 + a*e^2, 0] \ \&\& \ \text{NeQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{NegQ}[-(a)*c]$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^((c_) + (d_)*tan[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4113 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4118 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^((c_) + (d_)*tan[(e_) + (f_)*(x_)]^(n_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c + d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`

3.396.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 278, normalized size of antiderivative = 0.75

method	result
derivativedivides	$2(\sqrt{\tan(dx+c)} B b^3 - \frac{2A a^3}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2a^2(3Ab+Ba)}{\sqrt{\tan(dx+c)}} + \frac{(-A a^3 + 3Aa b^2 + 3B a^2 b - B b^3) \sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))} \right) \right)}{4}}$
default	$2(\sqrt{\tan(dx+c)} B b^3 - \frac{2A a^3}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{2a^2(3Ab+Ba)}{\sqrt{\tan(dx+c)}} + \frac{(-A a^3 + 3Aa b^2 + 3B a^2 b - B b^3) \sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))} \right) \right)}{4}}$
parts	$\frac{(A b^3 + 3B a b^2) \sqrt{2} \left(\ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4d}$

input `int((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x,method=_RETURNV
ERBOSE)`

output `1/d*(2*tan(d*x+c)^(1/2)*B*b^3-2/3*A*a^3/tan(d*x+c)^(3/2)-2*a^2*(3*A*b+B*a)
/tan(d*x+c)^(1/2)+1/4*(-A*a^3+3*A*a*b^2+3*B*a^2*b-B*b^3)*2^(1/2)*(ln((1+2^
(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))
)))+1/4*(-3*A*a^2*b+A*b^3-B*a^3+3*B*a*b^2)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+
c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2
^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))`

3.396.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6162 vs. 2(332) = 664.

Time = 1.44 (sec) , antiderivative size = 6162, normalized size of antiderivative = 16.56

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm
m="fracas")`

output Too large to include

3.396. $\int \frac{(a+b \tan(c+dx))^3 (A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

3.396.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^3}{\tan^{\frac{5}{2}}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**3*(A+B*tan(d*x+c))/tan(d*x+c)**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**3/tan(c + d*x)**(5/2), x)`

3.396.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 310, normalized size of antiderivative = 0.83

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{24 B b^3 \sqrt{\tan(dx + c)} - 6 \sqrt{2}((A + B)a^3 + 3(A - B)a^2 b - 3(A + B)ab^2 - (A - B)b^3) \arctan\left(\frac{1}{2} \sqrt{2}\left(\sqrt{2}\right)\right)}{d}$$

input `integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="maxima")`

output `1/12*(24*B*b^3*sqrt(tan(d*x + c)) - 6*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) - 6*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - 3*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 3*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - 8*(A*a^3 + 3*(B*a^3 + 3*A*a^2*b)*tan(d*x + c))/tan(d*x + c)^(3/2))/d`

3.397 $\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

3.397.1 Optimal result 3785
 3.397.2 Mathematica [C] (verified) 3786
 3.397.3 Rubi [A] (verified) 3786
 3.397.4 Maple [A] (verified) 3793
 3.397.5 Fricas [B] (verification not implemented) 3793
 3.397.6 Sympy [F] 3794
 3.397.7 Maxima [A] (verification not implemented) 3794
 3.397.8 Giac [F(-1)] 3795
 3.397.9 Mupad [B] (verification not implemented) 3795

3.397.1 Optimal result

Integrand size = 33, antiderivative size = 380

$$\int \frac{(a + b \tan(c + dx))^3(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx =$$

$$\frac{(a^3(A - B) - 3ab^2(A - B) - 3a^2b(A + B) + b^3(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(a^3(A - B) - 3ab^2(A - B) - 3a^2b(A + B) + b^3(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(3a^2b(A - B) - b^3(A - B) + a^3(A + B) - 3ab^2(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(3a^2b(A - B) - b^3(A - B) + a^3(A + B) - 3ab^2(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

$$- \frac{2a^2(9Ab + 5aB)}{15d \tan^{\frac{3}{2}}(c + dx)} + \frac{2a(5a^2A - 14Ab^2 - 15abB)}{5d\sqrt{\tan(c + dx)}} - \frac{2aA(a + b \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)}$$

output $\frac{1}{2}(a^3(A-B) - 3ab^2(A-B) - 3a^2b(A+B) + b^3(A+B)) \arctan(-1 + 2^{1/2} \tan(dx+c)^{1/2}) / d \cdot 2^{1/2} + \frac{1}{2}(a^3(A-B) - 3ab^2(A-B) - 3a^2b(A+B) + b^3(A+B)) \arctan(1 + 2^{1/2} \tan(dx+c)^{1/2}) / d \cdot 2^{1/2} + \frac{1}{4}(3a^2b(A-B) - b^3(A-B) + a^3(A+B) - 3ab^2(A+B)) \ln(1 - 2^{1/2} \tan(dx+c)^{1/2} + \tan(dx+c)) / d \cdot 2^{1/2} - \frac{1}{4}(3a^2b(A-B) - b^3(A-B) + a^3(A+B) - 3ab^2(A+B)) \ln(1 + 2^{1/2} \tan(dx+c)^{1/2} + \tan(dx+c)) / d \cdot 2^{1/2} + \frac{2}{5} a^5 (5A^2 - 14Ab^2 - 15B^2) / d \cdot \tan(dx+c)^{1/2} - \frac{2}{15} a^2 (9Ab + 5B^2) / d \cdot \tan(dx+c)^{3/2} - \frac{2}{5} a^2 (a+b \tan(dx+c))^2 / d \cdot \tan(dx+c)^{5/2}$

3.397.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.44 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.44

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \frac{2(3(a^3 A - 3aAb^2 - 3a^2bB + b^3 B) \text{Hypergeometric2F1}(-\frac{5}{4}, 1, -\frac{1}{4}, -\tan^2(c + dx)) + 5(3a^2 Ab - Ab^3 -$$

input `Integrate[((a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output $(-2*(3*(a^3 A - 3a^2 A b^2 - 3a^2 b B + b^3 B) \text{Hypergeometric2F1}[-5/4, 1, -1/4, -\text{Tan}[c + d*x]^2] + 5*(3a^2 A b - A b^3 + a^3 B - 3a^2 b B) \text{Hypergeometric2F1}[-3/4, 1, 1/4, -\text{Tan}[c + d*x]^2] * \text{Tan}[c + d*x] + b*(9a^2 A b + 9a^2 B - 3b^2 B + 5b(A b + 3a B) * \text{Tan}[c + d*x] + 15b^2 B * \text{Tan}[c + d*x]^2)) / (15*d * \text{Tan}[c + d*x]^{5/2})$

3.397.3 Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 312, normalized size of antiderivative = 0.82, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 4088, 27, 3042, 4118, 25, 3042, 4111, 27, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.397. $\int \frac{(a+b \tan(c+dx))^3 (A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan(c + dx)^{7/2}} dx$$

↓ 4088

$$\frac{2}{5} \int \frac{(a + b \tan(c + dx)) (-b(aA - 5bB) \tan^2(c + dx) - 5(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(9Ab + 5aB))}{2 \tan^{\frac{5}{2}}(c + dx) \frac{2aA(a + b \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)}} dx$$

↓ 27

$$\frac{1}{5} \int \frac{(a + b \tan(c + dx)) (-b(aA - 5bB) \tan^2(c + dx) - 5(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(9Ab + 5aB))}{\tan^{\frac{5}{2}}(c + dx) \frac{2aA(a + b \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)}} dx$$

↓ 3042

$$\frac{1}{5} \int \frac{(a + b \tan(c + dx)) (-b(aA - 5bB) \tan(c + dx)^2 - 5(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(9Ab + 5aB))}{\tan(c + dx)^{5/2} \frac{2aA(a + b \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)}} dx$$

↓ 4118

$$\frac{1}{5} \left(\int - \frac{b^2(aA - 5bB) \tan^2(c + dx) + 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(5Aa^2 - 15bBa - 14Ab^2)}{\tan^{\frac{3}{2}}(c + dx) \frac{2aA(a + b \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)}} dx \right)$$

↓ 25

$$\frac{1}{5} \left(- \int \frac{b^2(aA - 5bB) \tan^2(c + dx) + 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(5Aa^2 - 15bBa - 14Ab^2)}{\tan^{\frac{3}{2}}(c + dx) \frac{2aA(a + b \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)}} dx \right)$$

↓ 3042

3.397. $\int \frac{(a+b \tan(c+dx))^3 (A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$\frac{1}{5} \left(- \int \frac{b^2(aA - 5bB) \tan(c + dx)^2 + 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(5Aa^2 - 15bBa - 14Ab^2)}{\tan(c + dx)^{3/2}} \right. \\ \left. \frac{2aA(a + b \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)} \right. \\ \left. \downarrow 4111 \right.$$

$$\frac{1}{5} \left(- \int \frac{5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx + \frac{2a(5a^2A - 15abB - 14Ab^2)}{d\sqrt{\tan(c + dx)}} \right. \\ \left. \frac{2aA(a + b \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)} \right. \\ \left. \downarrow 27 \right.$$

$$\frac{1}{5} \left(-5 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2a(5a^2A - 15abB - 14Ab^2)}{d\sqrt{\tan(c + dx)}} \right. \\ \left. \frac{2aA(a + b \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)} \right. \\ \left. \downarrow 3042 \right.$$

$$\frac{1}{5} \left(-5 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx + \frac{2a(5a^2A - 15abB - 14Ab^2)}{d\sqrt{\tan(c + dx)}} \right. \\ \left. \frac{2aA(a + b \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)} \right. \\ \left. \downarrow 4017 \right.$$

$$\frac{1}{5} \left(- \frac{10 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{d} + \frac{2a(5a^2A - 15abB - 14Ab^2)}{d\sqrt{\tan(c + dx)}} \right. \\ \left. \frac{2aA(a + b \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)} \right. \\ \left. \downarrow 1482 \right.$$

$$\frac{1}{5} \left(- \frac{10 \left(\frac{1}{2}(a^3(A + B) + 3a^2b(A - B) - 3ab^2(A + B) - b^3(A - B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(a^3(A - B) - 3ab^2(A + B) - b^3(A - B)) \right)}{d} \right. \\ \left. \frac{2aA(a + b \tan(c + dx))^2}{5d \tan^{\frac{5}{2}}(c + dx)} \right.$$

3.397. $\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$

↓ 1476

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2} (a^3(A-B) \right)}{\frac{2aA(a+b\tan(c+dx))^2}{5d\tan^{\frac{5}{2}}(c+dx)}} \right)$$

↓ 1082

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2} (a^3(A-B) \right)}{\frac{2aA(a+b\tan(c+dx))^2}{5d\tan^{\frac{5}{2}}(c+dx)}} \right)$$

↓ 217

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2} (a^3(A-B) \right)}{d} \right)$$

↓ 1479

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \int \right)}{\frac{2aA(a+b\tan(c+dx))^2}{5d\tan^{\frac{5}{2}}(c+dx)}} \right)$$

↓ 25

3.397. $\int \frac{(a+b\tan(c+dx))^3(A+B\tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}+2\sqrt{\tan(c+dx)}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{\frac{2aA(a+b\tan(c+dx))^2}{5d\tan^{\frac{5}{2}}(c+dx)}} \right)$$

$$\frac{2aA(a+b\tan(c+dx))^2}{5d\tan^{\frac{5}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}+2\sqrt{\tan(c+dx)}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{\frac{2aA(a+b\tan(c+dx))^2}{5d\tan^{\frac{5}{2}}(c+dx)}} \right)$$

$$\frac{2aA(a+b\tan(c+dx))^2}{5d\tan^{\frac{5}{2}}(c+dx)}$$

↓ 1103

$$\frac{1}{5} \left(\frac{2a(5a^2A - 15abB - 14Ab^2)}{d\sqrt{\tan(c+dx)}} - \frac{2a^2(5aB + 9Ab)}{3d\tan^{\frac{3}{2}}(c+dx)} - \frac{10 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}+2\sqrt{\tan(c+dx)}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{\frac{2aA(a+b\tan(c+dx))^2}{5d\tan^{\frac{5}{2}}(c+dx)}} \right)$$

$$\frac{2aA(a+b\tan(c+dx))^2}{5d\tan^{\frac{5}{2}}(c+dx)}$$

input `Int[((a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `((-10*(-1/2*((a^3*(A - B) - 3*a*b^2*(A - B) - 3*a^2*b*(A + B) + b^3*(A + B)))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])) + ((3*a^2*b*(A - B) - b^3*(A - B) + a^3*(A + B) - 3*a*b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/d - (2*a^2*(9*A*b + 5*a*B))/(3*d*Tan[c + d*x]^(3/2)) + (2*a*(5*a^2*A - 14*A*b^2 - 15*a*b*B))/(d*Sqrt[Tan[c + d*x]]/5 - (2*a*A*(a + b*Tan[c + d*x])^2)/(5*d*Tan[c + d*x]^(5/2)))`

3.397.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^((c_) + (d_)*tan[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4111 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

rule 4118 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^((c_) + (d_)*tan[(e_) + (f_)*(x_)]^(n_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(-b*c - a*d)*(c^2*C - B*c*d + A*d^2)*((c + d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`

3.397.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 290, normalized size of antiderivative = 0.76

method	result
derivativedivides	$-\frac{2Aa^3}{5\tan(dx+c)^{\frac{5}{2}}}-\frac{2a^2(3Ab+Ba)}{3\tan(dx+c)^{\frac{3}{2}}}+\frac{2a(Aa^2-3Ab^2-3Bab)}{\sqrt{\tan(dx+c)}}+\frac{(-3Aa^2b+Ab^3-Ba^3+3Bab^2)\sqrt{2}\left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}\right)\right)}{4d}$
default	$-\frac{2Aa^3}{5\tan(dx+c)^{\frac{5}{2}}}-\frac{2a^2(3Ab+Ba)}{3\tan(dx+c)^{\frac{3}{2}}}+\frac{2a(Aa^2-3Ab^2-3Bab)}{\sqrt{\tan(dx+c)}}+\frac{(-3Aa^2b+Ab^3-Ba^3+3Bab^2)\sqrt{2}\left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}\right)\right)}{4d}$
parts	$\frac{(Ab^3+3Bab^2)\sqrt{2}\left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}\right)\right)+2\arctan\left(1+\sqrt{2}(\sqrt{\tan(dx+c)})\right)+2\arctan\left(-1+\sqrt{2}(\sqrt{\tan(dx+c)})\right)}{4d}$

input `int((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x,method=_RETURNV ERBOSE)`

output `1/d*(-2/5*A*a^3/tan(d*x+c)^(5/2)-2/3*a^2*(3*A*b+B*a)/tan(d*x+c)^(3/2)+2*a*(A*a^2-3*A*b^2-3*B*a*b)/tan(d*x+c)^(1/2)+1/4*(-3*A*a^2*b+A*b^3-B*a^3+3*B*a*b^2)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/4*(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))`

3.397.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6162 vs. 2(340) = 680.

Time = 1.45 (sec) , antiderivative size = 6162, normalized size of antiderivative = 16.22

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="fracas")`

output Too large to include

3.397.
$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

3.397.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^3}{\tan^{\frac{7}{2}}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**3*(A+B*tan(d*x+c))/tan(d*x+c)**(7/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**3/tan(c + d*x)**(7/2), x)`

3.397.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 327, normalized size of antiderivative = 0.86

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{30 \sqrt{2} ((A - B)a^3 - 3(A + B)a^2b - 3(A - B)ab^2 + (A + B)b^3) \arctan\left(\frac{1}{2} \sqrt{2} (\sqrt{2} + 2 \sqrt{\tan(dx + c)})\right)}{1}$$

input `integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="maxima")`

output `1/60*(30*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 30*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - 15*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 15*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - 8*(3*A*a^3 - 15*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2)*tan(d*x + c)^2 + 5*(B*a^3 + 3*A*a^2*b)*tan(d*x + c))/tan(d*x + c)^(5/2))/d`

3.397.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="giac")`

output Timed out

3.397.9 Mupad [B] (verification not implemented)

Time = 14.60 (sec) , antiderivative size = 7591, normalized size of antiderivative = 19.98

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \text{Too large to display}$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3)/tan(c + d*x)^(7/2),x)`

output `2*atanh((32*B^2*a^6*d^3*tan(c + d*x)^(1/2)*((5*B^2*a^3*b^3)/d^2 - (30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2))/(4*d^4) - (3*B^2*a*b^5)/(2*d^2) - (3*B^2*a^5*b)/(2*d^2))^(1/2))/(16*B^3*b^9*d^2 + 16*B*a^3*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) - 288*B^3*a^2*b^7*d^2 + 960*B^3*a^4*b^5*d^2 - 736*B^3*a^6*b^3*d^2 + 48*B^3*a^8*b*d^2 - 48*B*a*b^2*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) - (32*B^2*b^6*d^3*tan(c + d*x)^(1/2)*((5*B^2*a^3*b^3)/d^2 - (30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2))/(4*d^4) - (3*B^2*a*b^5)/(2*d^2) - (3*B^2*a^5*b)/(2*d^2))^(1/2))/(16*B^3*b^9*d^2 + 16*B*a^3*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) - 288*B^3*a^2*b^7*d^2 + 960*B^3*a^4*b^5*d^2 - 736*B^3*a^6*b^3*d^2 + 48*B^3*a^8*b*d^2 - 48*B*a*b^2*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) - 288*B^3*a^2*b^7*d^2 + 960*B^3*a^4*b^5*d^2 - 736*B^3*a^6*b^3*d^2 + 48*B^3*a^8*b*d^2 - 48*B*a*b^2*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) - 288*B^3*a^2*b^7*d^2 + 960*B^3*a^4*b^5*d^2 - 736*B^3*a^6*b^3*d^2 + 48*B^3*a^8*b*d^2 - 48*B*a*b^2*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) - 288*B^3*a^2*b^7*d^2 + 960*B^3*a^4*b^5*d^2 - 736*B^3*a^6*b^3*d^2 + 48*B^3*a^8*b*d^2 - 48*B*a*b^2*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) + (480*B^2*a^2*b^4*d^3*tan(c + d*x)^(1/2)*((5*B^2*a^3*b^3)/d^2 - (30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2))/(4*d^4) - (3*B^2*a*b^5)/(2*d^2) - (3*B^2*a^5*b)/(2*d^2))^(1/2))/(16*B^3*b^9*d^2 + 16*B*a^3*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) - 288*B^3*a^2*b^7*d^2 + 960*B^3*a^4*b^5*d^2 - 736*B^3*a^6*b^3*d^2 + 48*B^3*a^8*b*d^2 - 48*B*a*b^2*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) - 288*B^3*a^2*b^7*d^2 + 960*B^3*a^4*b^5*d^2 - 736*B^3*a^6*b^3*d^2 + 48*B^3*a^8*b*d^2 - 48*B*a*b^2*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) - 288*B^3*a^2*b^7*d^2 + 960*B^3*a^4*b^5*d^2 - 736*B^3*a^6*b^3*d^2 + 48*B^3*a^8*b*d^2 - 48*B*a*b^2*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) + (480*B^2*a^2*b^4*d^3*tan(c + d*x)^(1/2)*((5*B^2*a^3*b^3)/d^2 - (30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2))/(4*d^4) - (3*B^2*a*b^5)/(2*d^2) - (3*B^2*a^5*b)/(2*d^2))^(1/2))/(16*B^3*b^9*d^2 + 16*B*a^3*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) - 288*B^3*a^2*b^7*d^2 + 960*B^3*a^4*b^5*d^2 - 736*B^3*a^6*b^3*d^2 + 48*B^3*a^8*b*d^2 - 48*B*a*b^2*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) - 288*B^3*a^2*b^7*d^2 + 960*B^3*a^4*b^5*d^2 - 736*B^3*a^6*b^3*d^2 + 48*B^3*a^8*b*d^2 - 48*B*a*b^2*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) - 288*B^3*a^2*b^7*d^2 + 960*B^3*a^4*b^5*d^2 - 736*B^3*a^6*b^3*d^2 + 48*B^3*a^8*b*d^2 - 48*B*a*b^2*(30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2) + (480*B^2*a^2*b^4*d^3*tan(c + d*x)^(1/2)*((5*B^2*a^3*b^3)/d^2 - (30*B^4*a^2*b^10*d^4 - B^4*b^12*d^4 - B^4*a^12*d^4 - 255*B^4*a^4*b^8*d^4 + 452*B^4*a^6*b^6*d^4 - 255*B^4*a^8*b^4*d^4 + 30*B^4*a^10*b^2*d^4)^(1/2))/(4*d^4) - (3*B^2*a*b^5)/(2*d^2) - (3*B^2*a^5*b)/(2*d^2))^(1/2))`

3.397. $\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

3.398
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.398.1 Optimal result	3796
3.398.2 Mathematica [C] (verified)	3797
3.398.3 Rubi [A] (verified)	3797
3.398.4 Maple [A] (verified)	3804
3.398.5 Fricas [B] (verification not implemented)	3805
3.398.6 Sympy [F(-1)]	3805
3.398.7 Maxima [A] (verification not implemented)	3806
3.398.8 Giac [F(-1)]	3806
3.398.9 Mupad [B] (verification not implemented)	3807

3.398.1 Optimal result

Integrand size = 33, antiderivative size = 325

$$\begin{aligned} & \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx \\ &= \frac{(a(A-B)+b(A+B)) \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d} \\ & \quad - \frac{(a(A-B)+b(A+B)) \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d} \\ & \quad - \frac{2a^{5/2}(Ab-aB) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{b^{5/2}(a^2+b^2)d} \\ & \quad + \frac{(b(A-B)-a(A+B)) \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d} \\ & \quad - \frac{(b(A-B)-a(A+B)) \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d} \\ & \quad + \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{b^2d} + \frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} \end{aligned}$$

output
$$-2a^{5/2}(A^2b - B^2a) \arctan(b^{1/2} \tan(dx+c)^{1/2}/a^{1/2})/b^{5/2}/(a^2 + b^2)/d - 1/2(a(A-B) + b(A+B)) \arctan(-1 + 2^{1/2} \tan(dx+c)^{1/2})/(a^2 + b^2)/d * 2^{1/2} - 1/2(a(A-B) + b(A+B)) \arctan(1 + 2^{1/2} \tan(dx+c)^{1/2})/(a^2 + b^2)/d * 2^{1/2} + 1/4(b(A-B) - a(A+B)) \ln(1 - 2^{1/2} \tan(dx+c)^{1/2}) + \tan(dx+c)/(a^2 + b^2)/d * 2^{1/2} - 1/4(b(A-B) - a(A+B)) \ln(1 + 2^{1/2} \tan(dx+c)^{1/2}) + \tan(dx+c)/(a^2 + b^2)/d * 2^{1/2} + 2(A^2b - B^2a) \tan(dx+c)^{1/2}/b^2/d + 2/3 * B \tan(dx+c)^{3/2}/b/d$$

3.398.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.23 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.58

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$= \frac{-3\sqrt[4]{-1}(a+ib)b^{5/2}(iA+B) \arctan\left((-1)^{3/4} \sqrt{\tan(c+dx)}\right) + 6a^{5/2}(-Ab+aB) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{d}$$

input `Integrate[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output
$$\frac{(-3(-1)^{1/4}(a+Ib)b^{5/2}(IA+B) \text{ArcTan}[(-1)^{3/4} \text{Sqrt}[\text{Tan}[c+d*x]]] + 6a^{5/2}(-Ab+aB) \text{ArcTan}[\text{Sqrt}[b] \text{Sqrt}[\text{Tan}[c+d*x]]]/\text{Sqrt}[a] + 3(-1)^{1/4}b^{5/2}(Ia+b)(A+IB) \text{ArcTanh}[(-1)^{3/4} \text{Sqrt}[\text{Tan}[c+d*x]]] + 2\text{Sqrt}[b](a^2+b^2) \text{Sqrt}[\text{Tan}[c+d*x]](3A^2b-3a^2B+b^2B \text{Tan}[c+d*x]))}{(3b^{5/2}(a^2+b^2)d)}$$

3.398.3 Rubi [A] (verified)

Time = 1.45 (sec) , antiderivative size = 289, normalized size of antiderivative = 0.89, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4090, 27, 3042, 4130, 27, 3042, 4136, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.398.
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$\begin{aligned}
& \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx \\
& \quad \downarrow 3042 \\
& \int \frac{\tan(c+dx)^{5/2}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx \\
& \quad \downarrow 4090 \\
& \frac{2 \int -\frac{3\sqrt{\tan(c+dx)}(-((Ab-aB) \tan^2(c+dx))+bB \tan(c+dx)+aB)}{2(a+b \tan(c+dx))} dx}{3b} + \frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} \\
& \quad \downarrow 27 \\
& \frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} - \frac{\int \frac{\sqrt{\tan(c+dx)}(-((Ab-aB) \tan^2(c+dx))+bB \tan(c+dx)+aB)}{a+b \tan(c+dx)} dx}{b} \\
& \quad \downarrow 3042 \\
& \frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} - \frac{\int \frac{\sqrt{\tan(c+dx)}(-((Ab-aB) \tan(c+dx)^2)+bB \tan(c+dx)+aB)}{a+b \tan(c+dx)} dx}{b} \\
& \quad \downarrow 4130 \\
& \frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} - \frac{2 \int \frac{A \tan(c+dx)b^2 + (-Ba^2 + Aba + b^2B) \tan^2(c+dx) + a(Ab-aB)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{bd} \\
& \quad \downarrow 27 \\
& \frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} - \frac{\int \frac{A \tan(c+dx)b^2 + (-Ba^2 + Aba + b^2B) \tan^2(c+dx) + a(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{bd} \\
& \quad \downarrow 3042 \\
& \frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} - \frac{\int \frac{A \tan(c+dx)b^2 + (-Ba^2 + Aba + b^2B) \tan(c+dx)^2 + a(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{bd} \\
& \quad \downarrow 4136 \\
& \frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} - \frac{\int \frac{(Ab-aB)b^2 + (aA+bB) \tan(c+dx)b^2}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} + \frac{a^3(Ab-aB) \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{bd} \\
& \quad \downarrow 3042
\end{aligned}$$

3.398. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\frac{\int \frac{(Ab-aB)b^2+(aA+bB)\tan(c+dx)b^2}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} + \frac{a^3(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{bd}$$

\downarrow 4017

$$\frac{2 \int \frac{b^2(Ab-aB+(aA+bB)\tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} + \frac{a^3(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{bd}$$

\downarrow 27

$$\frac{2b^2 \int \frac{Ab-aB+(aA+bB)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} + \frac{a^3(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{bd}$$

\downarrow 1482

$$\frac{2b^2 \left(\frac{1}{2}(b(A-B)-a(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a(A-B)+b(A+B)) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) + \frac{a^3(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2}}{b} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{bd}$$

\downarrow 1476

$$\frac{2b^2 \left(\frac{1}{2}(b(A-B)-a(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}} d\sqrt{\tan(c+dx)} \right) \right)}{d(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{bd}$$

\downarrow 1082

$$\frac{2b^2 \left(\frac{1}{2}(b(A-B)-a(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{bd}$$

\downarrow 217

3.398. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} - \frac{2b^2 \left(\frac{1}{2}(b(A-B)-a(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)} + \frac{a^3(Ab-aB) \int \dots}{b}$$

1479

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} - \frac{2b^2 \left(\frac{1}{2}(b(A-B)-a(A+B)) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(a(A-B)+b(A+B)) \left(\dots \right) \right)}{d(a^2+b^2)} + \frac{a^3(Ab-aB) \int \dots}{b}$$

25

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} - \frac{2b^2 \left(\frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(a(A-B)+b(A+B)) \left(\dots \right) \right)}{d(a^2+b^2)} + \frac{a^3(Ab-aB) \int \dots}{b}$$

27

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} - \frac{2b^2 \left(\frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2}(a(A-B)+b(A+B)) \left(\dots \right) \right)}{d(a^2+b^2)} + \frac{a^3(Ab-aB) \int \dots}{b}$$

1103

$$\frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} - \frac{a^3(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2} + \frac{2b^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(b(A-B)-a(A+B)) \left(\dots \right) \right)}{d(a^2+b^2)} + \frac{a^3(Ab-aB) \int \dots}{b}$$

4117

3.398. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\frac{a^3(Ab-aB) \int \frac{1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} d \tan(c+dx)}{d(a^2+b^2)} + \frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} - \frac{2b^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(b(A-B)-a(A+B)) \right)}{d(a^2+b^2)}$$

73

$$\frac{2a^3(Ab-aB) \int \frac{1}{a+b \tan(c+dx)} d \sqrt{\tan(c+dx)}}{d(a^2+b^2)} + \frac{2B \tan^{\frac{3}{2}}(c+dx)}{3bd} - \frac{2b^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(b(A-B)-a(A+B)) \right)}{d(a^2+b^2)}$$

218

$$\frac{2b^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d(a^2+b^2)}$$

```
input Int[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]
```

```
output -((((2*a^(5/2)*(A*b - a*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[b]*(a^2 + b^2)*d) + (2*b^2*(((a*(A - B) + b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]))/2 + ((b*(A - B) - a*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2))/(a^2 + b^2)*d)/b - (2*(A*b - a*B)*Sqrt[Tan[c + d*x]]/(b*d))/b + (2*B*Tan[c + d*x]^(3/2))/(3*b*d)
```

3.398.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

3.398. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
 -1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
 & (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
 t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
 implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/R
 t[a/b, 2])], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
 eQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
 imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
 2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
 e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
 && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
 -2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
 x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
 reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4090 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

```
rule 4130 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2], x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

```
rule 4136 Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2])/(a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.398.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 292, normalized size of antiderivative = 0.90

method	result
derivativedivides	$\frac{2Bb \left(\frac{\tan^{\frac{3}{2}}(dx+c)}{3}\right) + 2Ab(\sqrt{\tan(dx+c)} - 2Ba(\sqrt{\tan(dx+c)}) - \frac{2a^3(Ab-Ba) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{b^2(a^2+b^2)\sqrt{ab}} + \frac{(-Ab+Ba)\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}}{1-\sqrt{2}}\right)\right)}{b^2(a^2+b^2)\sqrt{ab}}}{b^2}$
default	$\frac{2Bb \left(\frac{\tan^{\frac{3}{2}}(dx+c)}{3}\right) + 2Ab(\sqrt{\tan(dx+c)} - 2Ba(\sqrt{\tan(dx+c)}) - \frac{2a^3(Ab-Ba) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{b^2(a^2+b^2)\sqrt{ab}} + \frac{(-Ab+Ba)\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}}{1-\sqrt{2}}\right)\right)}{b^2(a^2+b^2)\sqrt{ab}}}{b^2}$

```
input int(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

$$3.398. \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

output $1/d*(2/b^2*(1/3*B*b*\tan(dx+c)^{(3/2)}+A*b*\tan(dx+c)^{(1/2)}-B*a*\tan(dx+c)^{(1/2)})-2/b^2*a^3*(A*b-B*a)/(a^2+b^2)/(a*b)^{(1/2)}*\arctan(b*\tan(dx+c)^{(1/2)}/(a*b)^{(1/2)})+2/(a^2+b^2)*(1/8*(-A*b+B*a)*2^{(1/2)}*(\ln((1+2^{(1/2)}*\tan(dx+c)^{(1/2)}+\tan(dx+c)))/(1-2^{(1/2)}*\tan(dx+c)^{(1/2)}+\tan(dx+c))))+2*\arctan(1+2^{(1/2)}*\tan(dx+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(dx+c)^{(1/2)}))+1/8*(-A*a-B*b)*2^{(1/2)}*(\ln((1-2^{(1/2)}*\tan(dx+c)^{(1/2)}+\tan(dx+c)))/(1+2^{(1/2)}*\tan(dx+c)^{(1/2)}+\tan(dx+c))))+2*\arctan(1+2^{(1/2)}*\tan(dx+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(dx+c)^{(1/2)})))))$

3.398.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3067 vs. $2(281) = 562$.

Time = 21.87 (sec) , antiderivative size = 6160, normalized size of antiderivative = 18.95

$$\int \frac{\tan^{5/2}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

input `integrate(tan(dx+c)^(5/2)*(A+B*tan(dx+c))/(a+b*tan(dx+c)),x, algorithm="fricas")`

output Too large to include

3.398.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\tan^{5/2}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Timed out}$$

input `integrate(tan(dx+c)**(5/2)*(A+B*tan(dx+c))/(a+b*tan(dx+c)),x)`

output Timed out

3.398.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 258, normalized size of antiderivative = 0.79

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{24(Ba^4 - Aa^3b) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^2b^2+b^4)\sqrt{ab}} - \frac{3\left(2\sqrt{2}((A-B)a+(A+B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}+2\sqrt{\tan(dx+c)}\right)\right) + 2\sqrt{2}((A-B)a+(A+B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}-2\sqrt{\tan(dx+c)}\right)\right)\right)}{(a^2+b^2)\sqrt{ab}}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/12*(24*(B*a^4 - A*a^3*b)*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/((a^2*b^2 + b^4)*sqrt(a*b)) - 3*(2*sqrt(2)*((A - B)*a + (A + B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a + (A + B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*((A + B)*a - (A - B)*b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*((A + B)*a - (A - B)*b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))/(a^2 + b^2) + 8*(B*b*tan(d*x + c)^(3/2) - 3*(B*a - A*b)*sqrt(tan(d*x + c)))/b^2)/d`

3.398.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output Timed out

3.398.9 Mupad [B] (verification not implemented)

Time = 14.94 (sec) , antiderivative size = 16441, normalized size of antiderivative = 50.59

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

```
input int((tan(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)),x)
```

```
output atan(((((((32*(4*A*a*b^8*d^4 + 8*A*a^3*b^6*d^4 + 4*A*a^5*b^4*d^4))/(b*d^5)
- (32*tan(c + d*x)^(1/2)*(((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4
*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*
a^2*b^2*d^4)))^(1/2)*(16*b^10*d^4 + 16*a^2*b^8*d^4 - 16*a^4*b^6*d^4 - 16*a
^6*b^4*d^4))/(b*d^4)*(((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4
+ 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2
*b^2*d^4)))^(1/2) + (32*tan(c + d*x)^(1/2)*(4*A^2*a^3*b^5*d^2 + 2*A^2*a^5*b
^3*d^2 - 14*A^2*a*b^7*d^2 + 16*A^2*a^7*b*d^2))/(b*d^4))*(((64*A^4*a^2*b^2
*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d
^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) - (32*(A^3*a^2*b^5*d^2
- 15*A^3*a^4*b^3*d^2 + 12*A^3*a^6*b*d^2))/(b*d^5))*(((64*A^4*a^2*b^2*d^4 -
A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(1
6*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) + (32*tan(c + d*x)^(1/2)*(2*
A^4*a^6 - A^4*b^6))/(b*d^4))*(((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*
b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 +
2*a^2*b^2*d^4)))^(1/2)*1i - (((((32*(4*A*a*b^8*d^4 + 8*A*a^3*b^6*d^4 + 4*
A*a^5*b^4*d^4))/(b*d^5) + (32*tan(c + d*x)^(1/2)*(((64*A^4*a^2*b^2*d^4 - A
^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*
(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*(16*b^10*d^4 + 16*a^2*b^8*d^4
- 16*a^4*b^6*d^4 - 16*a^6*b^4*d^4))/(b*d^4))*(((64*A^4*a^2*b^2*d^4 - A...
```

3.399
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.399.1 Optimal result	3808
3.399.2 Mathematica [C] (verified)	3809
3.399.3 Rubi [A] (verified)	3809
3.399.4 Maple [A] (verified)	3815
3.399.5 Fricas [B] (verification not implemented)	3816
3.399.6 Sympy [F]	3816
3.399.7 Maxima [A] (verification not implemented)	3817
3.399.8 Giac [F(-1)]	3817
3.399.9 Mupad [B] (verification not implemented)	3818

3.399.1 Optimal result

Integrand size = 33, antiderivative size = 297

$$\begin{aligned} & \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx \\ &= -\frac{(b(A-B)-a(A+B)) \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d} \\ & \quad + \frac{(b(A-B)-a(A+B)) \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d} \\ & \quad + \frac{2a^{3/2}(Ab-aB) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{b^{3/2}(a^2+b^2)d} \\ & \quad + \frac{(a(A-B)+b(A+B)) \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d} \\ & \quad - \frac{(a(A-B)+b(A+B)) \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d} \\ & \quad + \frac{2B\sqrt{\tan(c+dx)}}{bd} \end{aligned}$$

output $2a^{3/2}(A*b-B*a)*\arctan(b^{1/2}*\tan(d*x+c)^{1/2}/a^{1/2})/b^{3/2}/(a^2+b^2)/d+1/2*(b*(A-B)-a*(A+B))*\arctan(-1+2^{1/2}*\tan(d*x+c)^{1/2})/(a^2+b^2)/d*2^{1/2}+1/2*(b*(A-B)-a*(A+B))*\arctan(1+2^{1/2}*\tan(d*x+c)^{1/2})/(a^2+b^2)/d*2^{1/2}+1/4*(a*(A-B)+b*(A+B))*\ln(1-2^{1/2}*\tan(d*x+c)^{1/2}+\tan(d*x+c))/(a^2+b^2)/d*2^{1/2}-1/4*(a*(A-B)+b*(A+B))*\ln(1+2^{1/2}*\tan(d*x+c)^{1/2}+\tan(d*x+c))/(a^2+b^2)/d*2^{1/2}+2*B*\tan(d*x+c)^{1/2}/b/d$

3.399.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.56

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{\sqrt[4]{-1}(a+ib)b^{3/2}(A-iB)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right) - 2a^{3/2}(-Ab+aB)\arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right) + \sqrt[4]{-1}(a-ib)b^{3/2}(A+iB)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right) + 2a^{3/2}(Ab-aB)\arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{b^{3/2}(a^2+b^2)d}$$

input `Integrate[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output $((-1)^{1/4}*(a + I*b)*b^{3/2}*(A - I*B)*\text{ArcTan}[(-1)^{3/4}*\text{Sqrt}[\text{Tan}[c + d*x]]] - 2*a^{3/2}*(-(A*b) + a*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]] + (-1)^{1/4}*(a - I*b)*b^{3/2}*(A + I*B)*\text{ArcTan}h[(-1)^{3/4}*\text{Sqrt}[\text{Tan}[c + d*x]]] + 2*\text{Sqrt}[b]*(a^2 + b^2)*B*\text{Sqrt}[\text{Tan}[c + d*x]])/(b^{3/2}*(a^2 + b^2)*d)$

3.399.3 Rubi [A] (verified)

Time = 1.09 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.86, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 4090, 27, 3042, 4136, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.399. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$

$$\begin{aligned}
& \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx \\
& \quad \downarrow 3042 \\
& \int \frac{\tan(c+dx)^{3/2}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx \\
& \quad \downarrow 4090 \\
& \frac{2 \int -\frac{((Ab-aB) \tan^2(c+dx)+bB \tan(c+dx)+aB)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b} + \frac{2B\sqrt{\tan(c+dx)}}{bd} \\
& \quad \downarrow 27 \\
& \frac{2B\sqrt{\tan(c+dx)}}{bd} - \frac{\int -\frac{((Ab-aB) \tan^2(c+dx)+bB \tan(c+dx)+aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b} \\
& \quad \downarrow 3042 \\
& \frac{2B\sqrt{\tan(c+dx)}}{bd} - \frac{\int -\frac{((Ab-aB) \tan(c+dx)^2+bB \tan(c+dx)+aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b} \\
& \quad \downarrow 4136 \\
& \frac{2B\sqrt{\tan(c+dx)}}{bd} - \frac{\int \frac{b(aA+bB)-b(Ab-aB) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{a^2(Ab-aB) \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} \\
& \quad \downarrow 3042 \\
& \frac{2B\sqrt{\tan(c+dx)}}{bd} - \frac{\int \frac{b(aA+bB)-b(Ab-aB) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{a^2(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} \\
& \quad \downarrow 4017 \\
& \frac{2B\sqrt{\tan(c+dx)}}{bd} - \frac{2 \int \frac{b(aA+bB-(Ab-aB) \tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} - \frac{a^2(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} \\
& \quad \downarrow 27 \\
& \frac{2B\sqrt{\tan(c+dx)}}{bd} - \frac{2b \int \frac{aA+bB-(Ab-aB) \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} - \frac{a^2(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} \\
& \quad \downarrow 1482
\end{aligned}$$

3.399. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\frac{2B\sqrt{\tan(c+dx)}}{bd} - \frac{2b\left(\frac{1}{2}(a(A-B)+b(A+B))\int\frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}-\frac{1}{2}(b(A-B)-a(A+B))\int\frac{\tan(c+dx)+1}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}\right)-a^2(Ab-aB)\int\frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}(a^2+b^2)}}{d(a^2+b^2)}$$

1476

$$\frac{2B\sqrt{\tan(c+dx)}}{bd} - \frac{2b\left(\frac{1}{2}(a(A-B)+b(A+B))\int\frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}-\frac{1}{2}(b(A-B)-a(A+B))\left(\frac{1}{2}\int\frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}+\frac{1}{2}\int\frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}\right)\right)-a^2(Ab-aB)\int\frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}(a^2+b^2)}}{d(a^2+b^2)}$$

1082

$$\frac{2B\sqrt{\tan(c+dx)}}{bd} - \frac{2b\left(\frac{1}{2}(a(A-B)+b(A+B))\int\frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}-\frac{1}{2}(b(A-B)-a(A+B))\left(\frac{\int\frac{1}{-\tan(c+dx)-1}d\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}}-\frac{\int\frac{1}{-\tan(c+dx)-1}d\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}}\right)\right)-a^2(Ab-aB)\int\frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}(a^2+b^2)}}{d(a^2+b^2)}$$

217

$$\frac{2B\sqrt{\tan(c+dx)}}{bd} - \frac{2b\left(\frac{1}{2}(a(A-B)+b(A+B))\int\frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}-\frac{1}{2}(b(A-B)-a(A+B))\left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}}-\frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}}\right)\right)-a^2(Ab-aB)\int\frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}(a^2+b^2)}}{d(a^2+b^2)}$$

1479

$$\frac{2B\sqrt{\tan(c+dx)}}{bd} - \frac{2b\left(\frac{1}{2}(a(A-B)+b(A+B))\left(\frac{\int\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}}{2\sqrt{2}}-\frac{\int\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}}{2\sqrt{2}}\right)-\frac{1}{2}(b(A-B)-a(A+B))\int\frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}(a^2+b^2)}\right)-a^2(Ab-aB)\int\frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}(a^2+b^2)}}{d(a^2+b^2)}$$

25

$$\frac{2B\sqrt{\tan(c+dx)}}{bd} - \frac{2b\left(\frac{1}{2}(a(A-B)+b(A+B))\left(\frac{\int\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}}{2\sqrt{2}}+\frac{\int\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}}{2\sqrt{2}}\right)-\frac{1}{2}(b(A-B)-a(A+B))\int\frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}(a^2+b^2)}\right)-a^2(Ab-aB)\int\frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}(a^2+b^2)}}{d(a^2+b^2)}$$

27

3.399. $\int\frac{\tan^3(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)}dx$

$$\frac{\frac{2B\sqrt{\tan(c+dx)}}{bd} - 2b\left(\frac{\frac{1}{2}(a(A-B)+b(A+B))\left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)} + \frac{1}{2}\int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}\right) - \frac{1}{2}(b(A-B)-a(A+B))\right)}{d(a^2+b^2)}}{b}$$

1103

$$\frac{\frac{2B\sqrt{\tan(c+dx)}}{bd} - 2b\left(\frac{\frac{1}{2}(a(A-B)+b(A+B))\left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}}\right) - \frac{1}{2}(b(A-B)-a(A+B))\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}}\right)\right)}{d(a^2+b^2)}}{b}$$

4117

$$\frac{\frac{2B\sqrt{\tan(c+dx)}}{bd} - 2b\left(\frac{\frac{1}{2}(a(A-B)+b(A+B))\left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}}\right) - \frac{1}{2}(b(A-B)-a(A+B))\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}}\right)\right)}{d(a^2+b^2)}}{b}$$

73

$$\frac{\frac{2B\sqrt{\tan(c+dx)}}{bd} - 2b\left(\frac{\frac{1}{2}(a(A-B)+b(A+B))\left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}}\right) - \frac{1}{2}(b(A-B)-a(A+B))\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}}\right)\right)}{d(a^2+b^2)}}{b}$$

218

$$\frac{\frac{2B\sqrt{\tan(c+dx)}}{bd} - 2b\left(\frac{\frac{1}{2}(a(A-B)+b(A+B))\left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}}\right) - \frac{1}{2}(b(A-B)-a(A+B))\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}}\right)\right)}{d(a^2+b^2)}}{b}$$

```
input Int[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]
```

output
$$-\left(\frac{-2a^{3/2}(Ab - aB)\text{ArcTan}[\sqrt{b}\sqrt{\tan[c + dx]}/\sqrt{a}]}{\sqrt{b}(a^2 + b^2)d} + \frac{2b(-1/2((b(A - B) - a(A + B))(-\text{ArcTan}[1 - \sqrt{2}\sqrt{\tan[c + dx]})/\sqrt{2}] + \text{ArcTan}[1 + \sqrt{2}\sqrt{\tan[c + dx]})/\sqrt{2}))}{\sqrt{2}} + ((a(A - B) + b(A + B))(-1/2\text{Log}[1 - \sqrt{2}\sqrt{\tan[c + dx]} + \tan[c + dx]]/\sqrt{2} + \text{Log}[1 + \sqrt{2}\sqrt{\tan[c + dx]} + \tan[c + dx]]/(2\sqrt{2}))) / 2)}{(a^2 + b^2)d}\right) / b + \frac{2B\sqrt{\tan[c + dx]}}{bd}$$

3.399.3.1 Defintions of rubi rules used

rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{Int}[Fx, x], x]$

rule 27 $\text{Int}[(a_)*(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$

rule 73 $\text{Int}[(a_ + (b_)*(x_))^m * (c_ + (d_)*(x_))^n, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{Subst}[\text{Int}[x^{p(m+1)-1}(c - a(d/b) + d(x^p/b))^n, x], x, (a + b*x)^{1/p}], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 217 $\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1} * \text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 218 $\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

rule 1082 $\text{Int}[(a_ + (b_)*(x_ + (c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

```
rule 4090 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.399.4 Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.89

method	result
derivativedivides	$\frac{2(\sqrt{\tan(dx+c)}B}{b} + \frac{2a^2(Ab-Ba) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{b(a^2+b^2)\sqrt{ab}} + \frac{(-aA-Bb)\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c)))\right)}{4}$
default	$\frac{2(\sqrt{\tan(dx+c)}B}{b} + \frac{2a^2(Ab-Ba) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{b(a^2+b^2)\sqrt{ab}} + \frac{(-aA-Bb)\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c)))\right)}{4}$

3.399.
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

input `int(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*(2*tan(d*x+c)^(1/2)*B/b+2/b*a^2*(A*b-B*a)/(a^2+b^2)/(a*b)^(1/2)*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))+2/(a^2+b^2)*(1/8*(-A*a-B*b)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))+1/8*(A*b-B*a)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))`

3.399.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3033 vs. $2(257) = 514$.

Time = 9.28 (sec) , antiderivative size = 6092, normalized size of antiderivative = 20.51

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.399.6 Sympy [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \int \frac{(A+B\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)}{a+b\tan(c+dx)} dx$$

input `integrate(tan(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**(3/2)/(a + b*tan(c + d*x)), x)`

3.399.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 234, normalized size of antiderivative = 0.79

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx =$$

$$\frac{8(Ba^3 - Aa^2b) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^2b+b^3)\sqrt{ab}} - \frac{8B\sqrt{\tan(dx+c)}}{b} + \frac{2\sqrt{2}((A+B)a - (A-B)b) \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}+2\sqrt{\tan(dx+c)})\right) + 2\sqrt{2}((A+B)a - (A-B)b) \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}-2\sqrt{\tan(dx+c)})\right)}{(a^2+b^2)\sqrt{2}}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `-1/4*(8*(B*a^3 - A*a^2*b)*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/((a^2*b + b^3)*sqrt(a*b)) - 8*B*sqrt(tan(d*x + c))/b + (2*sqrt(2)*((A + B)*a - (A - B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A + B)*a - (A - B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((A - B)*a + (A + B)*b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((A - B)*a + (A + B)*b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))/(a^2 + b^2))/d`

3.399.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.399.9 Mupad [B] (verification not implemented)

Time = 15.09 (sec) , antiderivative size = 15701, normalized size of antiderivative = 52.87

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

```
input int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)),x)
```

```
output atan(((((((32*(4*B*a*b^8*d^4 + 8*B*a^3*b^6*d^4 + 4*B*a^5*b^4*d^4))/(b*d^5)
- (32*tan(c + d*x)^(1/2)*(((64*B^4*a^2*b^2*d^4 - B^4*(16*a^4*d^4 + 16*b^4
*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*
a^2*b^2*d^4))))^(1/2)*(16*b^10*d^4 + 16*a^2*b^8*d^4 - 16*a^4*b^6*d^4 - 16*a
^6*b^4*d^4))/(b*d^4)*(((64*B^4*a^2*b^2*d^4 - B^4*(16*a^4*d^4 + 16*b^4*d^4
+ 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2
*b^2*d^4))))^(1/2) + (32*tan(c + d*x)^(1/2)*(4*B^2*a^3*b^5*d^2 + 2*B^2*a^5*b
^3*d^2 - 14*B^2*a*b^7*d^2 + 16*B^2*a^7*b*d^2))/(b*d^4))*(((64*B^4*a^2*b^2
*d^4 - B^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b*d
^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4))))^(1/2) - (32*(B^3*a^2*b^5*d^2
- 15*B^3*a^4*b^3*d^2 + 12*B^3*a^6*b*d^2))/(b*d^5))*(((64*B^4*a^2*b^2*d^4 -
B^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b*d^2)/(1
6*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4))))^(1/2) + (32*tan(c + d*x)^(1/2)*(2*
B^4*a^6 - B^4*b^6))/(b*d^4))*(((64*B^4*a^2*b^2*d^4 - B^4*(16*a^4*d^4 + 16*
b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 +
2*a^2*b^2*d^4))))^(1/2)*1i - (((((32*(4*B*a*b^8*d^4 + 8*B*a^3*b^6*d^4 + 4*
B*a^5*b^4*d^4))/(b*d^5) + (32*tan(c + d*x)^(1/2)*(((64*B^4*a^2*b^2*d^4 - B
^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b*d^2)/(16*
(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4))))^(1/2)*(16*b^10*d^4 + 16*a^2*b^8*d^4
- 16*a^4*b^6*d^4 - 16*a^6*b^4*d^4))/(b*d^4))*(((64*B^4*a^2*b^2*d^4 - B...
```

3.400 $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.400.1 Optimal result 3819
 3.400.2 Mathematica [A] (verified) 3820
 3.400.3 Rubi [A] (verified) 3820
 3.400.4 Maple [A] (verified) 3825
 3.400.5 Fricas [B] (verification not implemented) 3826
 3.400.6 Sympy [F] 3826
 3.400.7 Maxima [A] (verification not implemented) 3827
 3.400.8 Giac [F(-1)] 3827
 3.400.9 Mupad [B] (verification not implemented) 3828

3.400.1 Optimal result

Integrand size = 33, antiderivative size = 278

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$= -\frac{(a(A-B)+b(A+B)) \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d}$$

$$+ \frac{(a(A-B)+b(A+B)) \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d}$$

$$- \frac{2\sqrt{a}(Ab-aB) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{b}(a^2+b^2)d}$$

$$- \frac{(b(A-B)-a(A+B)) \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d}$$

$$+ \frac{(b(A-B)-a(A+B)) \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d}$$

output

```
1/2*(a*(A-B)+b*(A+B))*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)+1/2*(a*(A-B)+b*(A+B))*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/4*(b*(A-B)-a*(A+B))*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)+1/4*(b*(A-B)-a*(A+B))*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)-2*(A*b-B*a)*arctan(b^(1/2)*tan(d*x+c)^(1/2)/a^(1/2))*a^(1/2)/(a^2+b^2)/d/b^(1/2)
```

3.400. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.400.2 Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 195, normalized size of antiderivative = 0.70

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{2\sqrt{2}(a(A-B)+b(A+B)) \left(\arctan \left(1 - \sqrt{2}\sqrt{\tan(c+dx)} \right) - \arctan \left(1 + \sqrt{2}\sqrt{\tan(c+dx)} \right) \right) + \dots}{\dots}$$

input `Integrate[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `-1/4*(2*Sqrt[2]*(a*(A - B) + b*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]) + (8*Sqrt[a]*(A*b - a*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/Sqrt[b] - Sqrt[2]*(b*(-A + B) + a*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]])/(a^2 + b^2)*d`

3.400.3 Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.82, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 4095, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx \\ & \quad \downarrow \text{4095} \\ & \frac{\int \frac{Ab-aB+(aA+bB) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2 + b^2} - \frac{a(Ab - aB) \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2 + b^2} \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.400. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\frac{\int \frac{Ab-aB+(aA+bB)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{a(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}$$

↓ 4017

$$\frac{2 \int \frac{Ab-aB+(aA+bB)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} - \frac{a(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}$$

↓ 1482

$$\frac{2\left(\frac{1}{2}(b(A-B)-a(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a(A-B)+b(A+B)) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right)}{d(a^2+b^2)} - \frac{a(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}$$

↓ 1476

$$\frac{2\left(\frac{1}{2}(b(A-B)-a(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}}\right)\right)}{d(a^2+b^2)} - \frac{a(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}$$

↓ 1082

$$\frac{2\left(\frac{1}{2}(b(A-B)-a(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}}\right)\right)}{d(a^2+b^2)} - \frac{a(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}$$

↓ 217

$$\frac{2\left(\frac{1}{2}(b(A-B)-a(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}}\right)\right)}{d(a^2+b^2)} - \frac{a(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}$$

↓ 1479

3.400. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$

$$\frac{2 \left(\frac{1}{2}(b(A - B) - a(A + B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \right)}{d(a^2 + b^2)} \\ \frac{a(Ab - aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2 + b^2} \\ \downarrow \text{25}$$

$$\frac{2 \left(\frac{1}{2}(b(A - B) - a(A + B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \right)}{d(a^2 + b^2)} \\ \frac{a(Ab - aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2 + b^2} \\ \downarrow \text{27}$$

$$\frac{2 \left(\frac{1}{2}(b(A - B) - a(A + B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d(a^2 + b^2)} \\ \frac{a(Ab - aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2 + b^2} \\ \downarrow \text{1103}$$

$$\frac{2 \left(\frac{1}{2}(a(A - B) + b(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(b(A - B) - a(A + B)) \left(\frac{\log}{\sqrt{2}} \right) \right)}{d(a^2 + b^2)} \\ \frac{a(Ab - aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2 + b^2} \\ \downarrow \text{4117}$$

$$\frac{2 \left(\frac{1}{2}(a(A - B) + b(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(b(A - B) - a(A + B)) \left(\frac{\log}{\sqrt{2}} \right) \right)}{d(a^2 + b^2)} \\ \frac{a(Ab - aB) \int \frac{1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} d\tan(c+dx)}{d(a^2 + b^2)} \\ \downarrow \text{73}$$

3.400. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$

$$\begin{aligned}
& \frac{2 \left(\frac{1}{2}(a(A-B) + b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(b(A-B) - a(A+B)) \left(\frac{\log}{\sqrt{2}} \right) \right)}{d(a^2 + b^2)} \\
& \quad \frac{2a(Ab - aB) \int \frac{1}{a+b \tan(c+dx)} d\sqrt{\tan(c+dx)}}{d(a^2 + b^2)} \\
& \quad \quad \quad \downarrow \text{218} \\
& \frac{2 \left(\frac{1}{2}(a(A-B) + b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(b(A-B) - a(A+B)) \left(\frac{\log}{\sqrt{2}} \right) \right)}{d(a^2 + b^2)} \\
& \quad \frac{2\sqrt{a}(Ab - aB) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{bd}(a^2 + b^2)}
\end{aligned}$$

input `Int[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(-2*Sqrt[a]*(A*b - a*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[b]*(a^2 + b^2)*d) + (2*(((a*(A - B) + b*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])))/2 + ((b*(A - B) - a*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)`

3.400.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n_), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

- rule 217 $\text{Int}[(a_ + (b_ \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{-1} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /;$ $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&$
 $\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 218 $\text{Int}[(a_ + (b_ \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a) \cdot \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /;$ $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b]$
- rule 1082 $\text{Int}[(a_ + (b_ \cdot x) + (c_ \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{With}\{q = 1 - 4 \cdot \text{Simplify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /;$ $\text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4 \cdot a \cdot c]) /;$ $\text{FreeQ}\{a, b, c, x\}$
- rule 1103 $\text{Int}[(d_ + (e_ \cdot x))/(a_ + (b_ \cdot x) + (c_ \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$
- rule 1476 $\text{Int}[(d_ + (e_ \cdot x)^2)/(a_ + (c_ \cdot x)^4), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c) \ \text{Int}[1/\text{Simp}[d/e + q \cdot x + x^2, x], x], x] + \text{Simp}[e/(2 \cdot c) \ \text{Int}[1/\text{Simp}[d/e - q \cdot x + x^2, x], x], x] /;$ $\text{FreeQ}\{a, c, d, e, x\} \ \&\& \ \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{PosQ}[d \cdot e]$
- rule 1479 $\text{Int}[(d_ + (e_ \cdot x)^2)/(a_ + (c_ \cdot x)^4), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[-2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c \cdot q) \ \text{Int}[(q - 2 \cdot x)/\text{Simp}[d/e + q \cdot x - x^2, x], x], x] + \text{Simp}[e/(2 \cdot c \cdot q) \ \text{Int}[(q + 2 \cdot x)/\text{Simp}[d/e - q \cdot x - x^2, x], x], x] /;$ $\text{FreeQ}\{a, c, d, e, x\} \ \&\& \ \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{NegQ}[d \cdot e]$
- rule 1482 $\text{Int}[(d_ + (e_ \cdot x)^2)/(a_ + (c_ \cdot x)^4), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[a \cdot c, 2]\}, \text{Simp}[(d \cdot q + a \cdot e)/(2 \cdot a \cdot c) \ \text{Int}[(q + c \cdot x^2)/(a + c \cdot x^4), x], x] + \text{Simp}[(d \cdot q - a \cdot e)/(2 \cdot a \cdot c) \ \text{Int}[(q - c \cdot x^2)/(a + c \cdot x^4), x], x] /;$ $\text{FreeQ}\{a, c, d, e, x\} \ \&\& \ \text{NeQ}[c \cdot d^2 + a \cdot e^2, 0] \ \&\& \ \text{NeQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{NegQ}[(-a) \cdot c]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$ $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4095 `Int((((A_) + (B_)*tan[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[Simp[A*(a*c + b*d) + B*(b*c - a*d) - (A*(b*c - a*d) - B*(a*c + b*d))*Tan[e + f*x], x]/Sqrt[c + d*Tan[e + f*x]], x] - Simp[(b*c - a*d)*((B*a - A*b)/(a^2 + b^2)) Int[(1 + Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*Sqrt[c + d*Tan[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4117 `Int(((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

3.400.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 244, normalized size of antiderivative = 0.88

method	result
derivativedivides	$-\frac{2a(Ab - Ba) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{(a^2+b^2)\sqrt{ab}} + \frac{(Ab - Ba)\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan\left(1+\sqrt{2}(\sqrt{\tan(dx+c)})\right) \right)}{4}$
default	$-\frac{2a(Ab - Ba) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{(a^2+b^2)\sqrt{ab}} + \frac{(Ab - Ba)\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan\left(1+\sqrt{2}(\sqrt{\tan(dx+c)})\right) \right)}{4}$

input `int(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

3.400.
$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

output $1/d*(-2*a*(A*b-B*a)/(a^2+b^2)/(a*b)^{(1/2)}*\arctan(b*\tan(d*x+c)^{(1/2)}/(a*b)^{(1/2)})+2/(a^2+b^2)*(1/8*(A*b-B*a)*2^{(1/2)}*(\ln((1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))/(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}))+1/8*(A*a+B*b)*2^{(1/2)}*(\ln((1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))/(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}))$

3.400.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2983 vs. $2(240) = 480$.

Time = 3.21 (sec) , antiderivative size = 5992, normalized size of antiderivative = 21.55

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.400.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \int \frac{(A+B\tan(c+dx))\sqrt{\tan(c+dx)}}{a+b\tan(c+dx)} dx$$

input `integrate(tan(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(tan(c + d*x))/(a + b*tan(c + d*x)), x)`

3.400.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.78

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{8(Ba^2 - Aab) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^2+b^2)\sqrt{ab}} + \frac{2\sqrt{2}((A-B)a+(A+B)b) \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}+2\sqrt{\tan(dx+c)})\right) + 2\sqrt{2}((A-B)a+(A+B)b) \arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}-2\sqrt{\tan(dx+c)})\right)}{(a^2+b^2)\sqrt{ab}}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/4*(8*(B*a^2 - A*a*b)*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/((a^2 + b^2)*sqrt(a*b)) + (2*sqrt(2)*((A - B)*a + (A + B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a + (A + B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*((A + B)*a - (A - B)*b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*((A + B)*a - (A - B)*b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))/(a^2 + b^2))/d`

3.400.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.400.9 Mupad [B] (verification not implemented)

Time = 13.35 (sec) , antiderivative size = 15090, normalized size of antiderivative = 54.28

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

```
input int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)),x)
```

```
output atan((((32*(13*A^3*a^2*b^4*d^2 + A^3*a^4*b^2*d^2))/d^5 + (((32*(12*A*a*b^7*d^4 + 24*A*a^3*b^5*d^4 + 12*A*a^5*b^3*d^4))/d^5 - (32*tan(c + d*x)^(1/2) * (((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*(16*b^9*d^4 + 16*a^2*b^7*d^4 - 16*a^4*b^5*d^4 - 16*a^6*b^3*d^4))/d^4)*(((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) + (32*tan(c + d*x)^(1/2)*(20*A^2*a^3*b^4*d^2 + 2*A^2*a^5*b^2*d^2 - 14*A^2*a*b^6*d^2))/d^4)*(((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2))* (((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) - (32*tan(c + d*x)^(1/2)*(A^4*b^5 - 2*A^4*a^2*b^3))/d^4)*(((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*1i - (((32*(13*A^3*a^2*b^4*d^2 + A^3*a^4*b^2*d^2))/d^5 + (((32*(12*A*a*b^7*d^4 + 24*A*a^3*b^5*d^4 + 12*A*a^5*b^3*d^4))/d^5 + (32*tan(c + d*x)^(1/2)*(((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*(16*b^9*d^4 + 16*a^2*b^7*d^4 - 16*a^4*b^5*d^4 - 16*a^6*b^3*d^4))/d^4)*(((64*A^4*a^2*b^2*d^4 - A^4*(16*a...
```

3.401
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx$$

3.401.1 Optimal result	3829
3.401.2 Mathematica [A] (verified)	3830
3.401.3 Rubi [A] (verified)	3830
3.401.4 Maple [A] (verified)	3835
3.401.5 Fricas [B] (verification not implemented)	3836
3.401.6 Sympy [F]	3836
3.401.7 Maxima [A] (verification not implemented)	3837
3.401.8 Giac [F(-1)]	3837
3.401.9 Mupad [B] (verification not implemented)	3838

3.401.1 Optimal result

Integrand size = 33, antiderivative size = 278

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx \\ &= \frac{(b(A - B) - a(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2) d} \\ & \quad - \frac{(b(A - B) - a(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2) d} \\ & \quad + \frac{2\sqrt{b}(Ab - aB) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{a}(a^2 + b^2) d} \\ & \quad - \frac{(a(A - B) + b(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2) d} \\ & \quad + \frac{(a(A - B) + b(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2) d} \end{aligned}$$

```
output -1/2*(b*(A-B)-a*(A+B))*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/2*(b*(A-B)-a*(A+B))*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/4*(a*(A-B)+b*(A+B))*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)+1/4*(a*(A-B)+b*(A+B))*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)+2*(A*b-B*a)*arctan(b^(1/2)*tan(d*x+c)^(1/2)/a^(1/2))*b^(1/2)/(a^2+b^2)/d/a^(1/2)
```

3.401.
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx$$

3.401.2 Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.70

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx =$$

$$\frac{2\sqrt{2}(b(-A + B) + a(A + B)) \left(\arctan \left(1 - \sqrt{2}\sqrt{\tan(c + dx)} \right) - \arctan \left(1 + \sqrt{2}\sqrt{\tan(c + dx)} \right) \right) + \dots}{\dots}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])),x]`

output `-1/4*(2*Sqrt[2]*(b*(-A + B) + a*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]) + (8*Sqrt[b]*(-(A*b) + a*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/Sqrt[a] + Sqrt[2]*(a*(A - B) + b*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]))/((a^2 + b^2)*d)`

3.401.3 Rubi [A] (verified)Time = 0.81 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.82, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$, Rules used = {3042, 4096, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx$$

$$\downarrow \text{4096}$$

$$\frac{b(Ab - aB) \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2 + b^2} + \frac{\int \frac{aA+bB-(Ab-aB) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2 + b^2}$$

$$\downarrow \text{3042}$$

3.401. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx$

$$\frac{\int \frac{aA+bB-(Ab-aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} + \frac{b(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}$$

↓ 4017

$$\frac{2 \int \frac{aA+bB-(Ab-aB)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} + \frac{b(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}$$

↓ 1482

$$\frac{2\left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(b(A-B)-a(A+B)) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}\right)}{d(a^2+b^2)} + \frac{b(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}$$

↓ 1476

$$\frac{2\left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}}\right)\right)}{d(a^2+b^2)} + \frac{b(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}$$

↓ 1082

$$\frac{2\left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}}\right)\right)}{d(a^2+b^2)} + \frac{b(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}$$

↓ 217

$$\frac{2\left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}}\right)\right)}{d(a^2+b^2)} + \frac{b(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}$$

↓ 1479

3.401. $\int \frac{A+B\tan(c+dx)}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx$

$$2 \left(\frac{1}{2}(a(A - B) + b(A + B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(b(A - B) - a(A + B)) \right) \frac{dx}{d(a^2 + b^2)}$$

$$\frac{b(Ab - aB) \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2 + b^2}$$

↓ 25

$$2 \left(\frac{1}{2}(a(A - B) + b(A + B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(b(A - B) - a(A + B)) \right) \frac{dx}{d(a^2 + b^2)}$$

$$\frac{b(Ab - aB) \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2 + b^2}$$

↓ 27

$$2 \left(\frac{1}{2}(a(A - B) + b(A + B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2}(b(A - B) - a(A + B)) \right) \frac{dx}{d(a^2 + b^2)}$$

$$\frac{b(Ab - aB) \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2 + b^2}$$

↓ 1103

$$\frac{b(Ab - aB) \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2 + b^2} +$$

$$2 \left(\frac{1}{2}(a(A - B) + b(A + B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(b(A - B) - a(A + B)) \right) \frac{dx}{d(a^2 + b^2)}$$

↓ 4117

$$\frac{b(Ab - aB) \int \frac{1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} d\tan(c + dx)}{d(a^2 + b^2)} +$$

$$2 \left(\frac{1}{2}(a(A - B) + b(A + B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(b(A - B) - a(A + B)) \right) \frac{dx}{d(a^2 + b^2)}$$

↓ 73

3.401. $\int \frac{A+B\tan(c+dx)}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx$

$$\frac{2b(Ab - aB) \int \frac{1}{a+b \tan(c+dx)} d\sqrt{\tan(c+dx)}}{d(a^2 + b^2)} + \frac{2 \left(\frac{1}{2}(a(A - B) + b(A + B)) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(b(A - B) - a(A + B)) \right)}{d(a^2 + b^2)}$$

↓ 218

$$\frac{2\sqrt{b}(Ab - aB) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{ad}(a^2 + b^2)} + \frac{2 \left(\frac{1}{2}(a(A - B) + b(A + B)) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(b(A - B) - a(A + B)) \right)}{d(a^2 + b^2)}$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])),x]`

output `(2*Sqrt[b]*(A*b - a*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[a]*(a^2 + b^2)*d) + (2*(-1/2*((b*(A - B) - a*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])) + ((a*(A - B) + b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)`

3.401.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

- rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4096 `Int((((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_))/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[a*A + b*B - (A*b - a*B)*Tan[e + f*x], x], x] + Simp[b*((A*b - a*B)/(a^2 + b^2)) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4117 `Int(((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

3.401.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 244, normalized size of antiderivative = 0.88

method	result
derivativedivides	$\frac{2(Ab - Ba)b \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{(a^2 + b^2)\sqrt{ab}} + \frac{(aA + Bb)\sqrt{2} \left(\ln\left(\frac{1 + \sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}{1 - \sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}\right) + 2 \arctan(1 + \sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))) \right)}{4}$
default	$\frac{2(Ab - Ba)b \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{(a^2 + b^2)\sqrt{ab}} + \frac{(aA + Bb)\sqrt{2} \left(\ln\left(\frac{1 + \sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}{1 - \sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}\right) + 2 \arctan(1 + \sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))) \right)}{4}$

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

3.401.
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx$$

output $1/d*(2*(A*b-B*a)*b/(a^2+b^2)/(a*b)^{(1/2)*\arctan(b*\tan(d*x+c)^{(1/2))/(a*b)^{(1/2))+2/(a^2+b^2)*(1/8*(A*a+B*b)*2^{(1/2)*(\ln((1+2^{(1/2)*\tan(d*x+c)^{(1/2)+\tan(d*x+c)})/(1-2^{(1/2)*\tan(d*x+c)^{(1/2)+\tan(d*x+c)})))+2*\arctan(1+2^{(1/2)*\tan(d*x+c)^{(1/2)+2*\arctan(-1+2^{(1/2)*\tan(d*x+c)^{(1/2)+1/8*(-A*b+B*a)*2^{(1/2)*(\ln((1-2^{(1/2)*\tan(d*x+c)^{(1/2)+\tan(d*x+c)})/(1+2^{(1/2)*\tan(d*x+c)^{(1/2)+\tan(d*x+c)})))+2*\arctan(1+2^{(1/2)*\tan(d*x+c)^{(1/2)+2*\arctan(-1+2^{(1/2)*\tan(d*x+c)^{(1/2)}}$

3.401.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2999 vs. $2(240) = 480$.

Time = 4.61 (sec) , antiderivative size = 6024, normalized size of antiderivative = 21.67

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.401.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx)) \sqrt{\tan(c + dx)}} dx$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+b*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))*sqrt(tan(c + d*x))), x)`

3.401.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 217, normalized size of antiderivative = 0.78

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx =$$

$$\frac{8 (Bab - Ab^2) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^2+b^2)\sqrt{ab}} - \frac{2\sqrt{2}((A+B)a - (A-B)b) \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2} + 2\sqrt{\tan(dx+c)})\right) + 2\sqrt{2}((A+B)a - (A-B)b) \arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2} - 2\sqrt{\tan(dx+c)})\right)}{(a^2+b^2)\sqrt{ab}}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `-1/4*(8*(B*a*b - A*b^2)*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/((a^2 + b^2)*sqrt(a*b)) - (2*sqrt(2)*((A + B)*a - (A - B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A + B)*a - (A - B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((A - B)*a + (A + B)*b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((A - B)*a + (A + B)*b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))/(a^2 + b^2))/d`

3.401.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.401.9 Mupad [B] (verification not implemented)

Time = 13.41 (sec) , antiderivative size = 14816, normalized size of antiderivative = 53.29

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx = \text{Too large to display}$$

```
input int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + b*tan(c + d*x))),x)
```

```
output atan((((32*(13*B^3*a^2*b^4*d^2 + B^3*a^4*b^2*d^2))/d^5 + (((32*(12*B*a*b^7*d^4 + 24*B*a^3*b^5*d^4 + 12*B*a^5*b^3*d^4))/d^5 - (32*tan(c + d*x)^(1/2) * (((64*B^4*a^2*b^2*d^4 - B^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*(16*b^9*d^4 + 16*a^2*b^7*d^4 - 16*a^4*b^5*d^4 - 16*a^6*b^3*d^4))/d^4)*(((64*B^4*a^2*b^2*d^4 - B^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) + (32*tan(c + d*x)^(1/2)*(20*B^2*a^3*b^4*d^2 + 2*B^2*a^5*b^2*d^2 - 14*B^2*a*b^6*d^2))/d^4)*(((64*B^4*a^2*b^2*d^4 - B^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2))* (((64*B^4*a^2*b^2*d^4 - B^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) - (32*tan(c + d*x)^(1/2)*(B^4*b^5 - 2*B^4*a^2*b^3))/d^4)*(((64*B^4*a^2*b^2*d^4 - B^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*1i - (((32*(13*B^3*a^2*b^4*d^2 + B^3*a^4*b^2*d^2))/d^5 + (((32*(12*B*a*b^7*d^4 + 24*B*a^3*b^5*d^4 + 12*B*a^5*b^3*d^4))/d^5 + (32*tan(c + d*x)^(1/2)*(((64*B^4*a^2*b^2*d^4 - B^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*(16*b^9*d^4 + 16*a^2*b^7*d^4 - 16*a^4*b^5*d^4 - 16*a^6*b^3*d^4))/d^4)*(((64*B^4*a^2*b^2*d^4 - B^4*(16*a...
```

3.402
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$$

3.402.1 Optimal result 3839
 3.402.2 Mathematica [C] (verified) 3840
 3.402.3 Rubi [A] (verified) 3840
 3.402.4 Maple [A] (verified) 3846
 3.402.5 Fricas [B] (verification not implemented) 3847
 3.402.6 Sympy [F] 3847
 3.402.7 Maxima [A] (verification not implemented) 3848
 3.402.8 Giac [F(-1)] 3848
 3.402.9 Mupad [B] (verification not implemented) 3849

3.402.1 Optimal result

Integrand size = 33, antiderivative size = 297

$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$$

$$= \frac{(a(A-B)+b(A+B)) \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d}$$

$$- \frac{(a(A-B)+b(A+B)) \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d}$$

$$- \frac{2b^{3/2}(Ab-aB) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{a^{3/2}(a^2+b^2)d}$$

$$+ \frac{(b(A-B)-a(A+B)) \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d}$$

$$- \frac{(b(A-B)-a(A+B)) \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d} - \frac{2A}{ad\sqrt{\tan(c+dx)}}$$

output

```
-2*b^(3/2)*(A*b-B*a)*arctan(b^(1/2)*tan(d*x+c)^(1/2)/a^(1/2))/a^(3/2)/(a^2+b^2)/d-1/2*(a*(A-B)+b*(A+B))*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/2*(a*(A-B)+b*(A+B))*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)+1/4*(b*(A-B)-a*(A+B))*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)-1/4*(b*(A-B)-a*(A+B))*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)-2*A/a/d/tan(d*x+c)^(1/2)
```

3.402.
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$$

3.402.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.63 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.52

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$= \frac{2b^{3/2}(-Ab+aB) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{a}(a^2+b^2)} + \frac{\sqrt[4]{-1}a((-ia+b)(A-iB) \arctan((-1)^{3/4}\sqrt{\tan(c+dx)})+(ia+b)(A+iB) \operatorname{arctanh}((-1)^{3/4}\sqrt{\tan(c+dx)})}{a^2+b^2}$$

ad

```
input Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])),x
]
```

```
output ((2*b^(3/2)*(-(A*b) + a*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[a]*(a^2 + b^2)) + ((-1)^(1/4)*a*(((I)*a + b)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + (I*a + b)*(A + I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]))/(a^2 + b^2) - (2*A)/Sqrt[Tan[c + d*x]]/(a*d)
```

3.402.3 Rubi [A] (verified)

Time = 1.08 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.86, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 4092, 27, 3042, 4136, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2}(a + b \tan(c + dx))} dx$$

↓ 4092

$$-\frac{2 \int \frac{Ab \tan^2(c+dx)+aA \tan(c+dx)+Ab-aB}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a} - \frac{2A}{ad\sqrt{\tan(c + dx)}}$$

$$\begin{aligned}
 & \int \frac{Ab \tan^2(c+dx) + aA \tan(c+dx) + Ab - aB}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{Ab \tan^2(c+dx) + aA \tan(c+dx) + Ab - aB}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a} - \frac{2A}{ad\sqrt{\tan(c+dx)}} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{Ab \tan(c+dx)^2 + aA \tan(c+dx) + Ab - aB}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a} - \frac{2A}{ad\sqrt{\tan(c+dx)}} \\
 & \quad \downarrow 4136 \\
 & \frac{b^2(Ab-aB) \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} + \frac{\int \frac{a(Ab-aB)+a(aA+bB) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{2A}{ad\sqrt{\tan(c+dx)}} \\
 & \quad \downarrow 3042 \\
 & \frac{b^2(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} + \frac{\int \frac{a(Ab-aB)+a(aA+bB) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{2A}{ad\sqrt{\tan(c+dx)}} \\
 & \quad \downarrow 4017 \\
 & \frac{2 \int \frac{a(Ab-aB+(aA+bB) \tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} + \frac{b^2(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} - \frac{2A}{ad\sqrt{\tan(c+dx)}} \\
 & \quad \downarrow 27 \\
 & \frac{2a \int \frac{Ab-aB+(aA+bB) \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} + \frac{b^2(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} - \frac{2A}{ad\sqrt{\tan(c+dx)}} \\
 & \quad \downarrow 1482 \\
 & \frac{2a \left(\frac{1}{2}(b(A-B)-a(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a(A-B)+b(A+B)) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d(a^2+b^2)} + \frac{b^2(Ab-aB) \int \frac{\tan(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} \\
 & \quad \downarrow 1476 \\
 & \frac{2A}{ad\sqrt{\tan(c+dx)}}
 \end{aligned}$$

3.402. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$\frac{2a \left(\frac{1}{2}(b(A-B) - a(A+B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a(A-B) + b(A+B)) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d(a^2+b^2)}$$

$$\frac{2A}{ad\sqrt{\tan(c+dx)}}$$

↓ 1082

$$\frac{2a \left(\frac{1}{2}(b(A-B) - a(A+B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a(A-B) + b(A+B)) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)}$$

$$\frac{2A}{ad\sqrt{\tan(c+dx)}}$$

↓ 217

$$\frac{2a \left(\frac{1}{2}(b(A-B) - a(A+B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a(A-B) + b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)} + \frac{b}{a}$$

$$\frac{2A}{ad\sqrt{\tan(c+dx)}}$$

↓ 1479

$$\frac{2a \left(\frac{1}{2}(b(A-B) - a(A+B)) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(a(A-B) + b(A+B)) \right)}{d(a^2+b^2)}$$

$$\frac{2A}{ad\sqrt{\tan(c+dx)}}$$

↓ 25

$$\frac{2a \left(\frac{1}{2}(b(A-B) - a(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(a(A-B) + b(A+B)) \right)}{d(a^2+b^2)}$$

$$\frac{2A}{ad\sqrt{\tan(c+dx)}}$$

↓ 27

3.402. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$\begin{aligned}
 & \frac{2a \left(\frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2}(a(A-B)+b(A+B)) \right)}{d(a^2+b^2)} \\
 & \frac{2A}{ad\sqrt{\tan(c+dx)}} \\
 & \downarrow 1103 \\
 & \frac{b^2(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx + 2a \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(b(A-B)-a(A+B)) \right)}{d(a^2+b^2)}}{a} \\
 & \frac{2A}{ad\sqrt{\tan(c+dx)}} \\
 & \downarrow 4117 \\
 & \frac{b^2(Ab-aB) \int \frac{1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} d\tan(c+dx) + 2a \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(b(A-B)-a(A+B)) \right)}{d(a^2+b^2)}}{a} \\
 & \frac{2A}{ad\sqrt{\tan(c+dx)}} \\
 & \downarrow 73 \\
 & \frac{2b^2(Ab-aB) \int \frac{1}{a+b\tan(c+dx)} d\sqrt{\tan(c+dx)} + 2a \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(b(A-B)-a(A+B)) \right)}{d(a^2+b^2)}}{a} \\
 & \frac{2A}{ad\sqrt{\tan(c+dx)}} \\
 & \downarrow 218 \\
 & \frac{2a \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d(a^2+b^2)}}{a} \\
 & \frac{2A}{ad\sqrt{\tan(c+dx)}}
 \end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])),x]`

3.402. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$

```
output -(((2*b^(3/2)*(A*b - a*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(S
qrt[a]*(a^2 + b^2)*d) + (2*a*(((a*(A - B) + b*(A + B))*(-(ArcTan[1 - Sqrt[
2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/S
qrt[2])))/2 + ((b*(A - B) - a*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d
*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c
+ d*x]]/(2*Sqrt[2])))/2))/((a^2 + b^2)*d)/a - (2*A)/(a*d*Sqrt[Tan[c + d*
x]])
```

3.402.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

```
rule 4092 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)
/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^
2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*
B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n +
2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n
+ 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1]
&& (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m]
|| (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

```
rule 4136 Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.402.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.89

method	result
derivativedivides	$-\frac{2(Ab - Ba)b^2 \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{a(a^2 + b^2)\sqrt{ab}} - \frac{2A}{a\sqrt{\tan(dx+c)}} + \frac{(-Ab + Ba)\sqrt{2} \left(\ln\left(\frac{1 + \sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}{1 - \sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}\right) + 2 \arctan\left(1 + \sqrt{2}\right) \right)}{4}$
default	$-\frac{2(Ab - Ba)b^2 \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{a(a^2 + b^2)\sqrt{ab}} - \frac{2A}{a\sqrt{\tan(dx+c)}} + \frac{(-Ab + Ba)\sqrt{2} \left(\ln\left(\frac{1 + \sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}{1 - \sqrt{2}(\sqrt{\tan(dx+c)} + \tan(dx+c))}\right) + 2 \arctan\left(1 + \sqrt{2}\right) \right)}{4}$

3.402. $\int \frac{A+B \tan(c+dx)}{\tan^3(c+dx)(a+b \tan(c+dx))} dx$

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*(-2/a*(A*b-B*a)*b^2/(a^2+b^2)/(a*b)^(1/2)*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))-2/a*A/tan(d*x+c)^(1/2)+2/(a^2+b^2)*(1/8*(-A*b+B*a)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/8*(-A*a-B*b)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))`

3.402.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3055 vs. $2(257) = 514$.

Time = 12.06 (sec) , antiderivative size = 6136, normalized size of antiderivative = 20.66

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.402.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx)) \tan^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(3/2)/(a+b*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))*tan(c + d*x)**(3/2)),x)`

3.402.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 235, normalized size of antiderivative = 0.79

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$= \frac{8 (Bab^2 - Ab^3) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^3 + ab^2)\sqrt{ab}} - \frac{2\sqrt{2}((A-B)a + (A+B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2+2\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}((A-B)a + (A+B)b) \arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2+2\sqrt{\tan(dx+c)}}\right)\right)}{(a^3 + ab^2)\sqrt{ab}}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/4*(8*(B*a*b^2 - A*b^3)*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/((a^3 + a*b^2)*sqrt(a*b)) - (2*sqrt(2)*((A - B)*a + (A + B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a + (A + B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*((A + B)*a - (A - B)*b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*((A + B)*a - (A - B)*b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))/(a^2 + b^2) - 8*A/(a*sqrt(tan(d*x + c))))/d`

3.402.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.402.9 Mupad [B] (verification not implemented)

Time = 13.62 (sec) , antiderivative size = 15318, normalized size of antiderivative = 51.58

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = \text{Too large to display}$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + b*tan(c + d*x))),x)`

output `atan(((tan(c + d*x)^(1/2)*(64*A^4*a^7*b^7*d^5 - 32*A^4*a^9*b^5*d^5) + (((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*((tan(c + d*x)^(1/2)*(512*A^2*a^8*b^8*d^7 - 448*A^2*a^10*b^6*d^7 + 128*A^2*a^12*b^4*d^7 + 64*A^2*a^14*b^2*d^7) - (((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*(tan(c + d*x)^(1/2)*(((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*(512*a^9*b^9*d^9 + 512*a^11*b^7*d^9 - 512*a^13*b^5*d^9 - 512*a^15*b^3*d^9) - 512*A*a^8*b^9*d^8 - 640*A*a^10*b^7*d^8 + 256*A*a^12*b^5*d^8 + 384*A*a^14*b^3*d^8))*(((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) + 128*A^3*a^7*b^8*d^6 - 32*A^3*a^11*b^4*d^6 - 32*A^3*a^13*b^2*d^6))*(((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*i + (tan(c + d*x)^(1/2)*(64*A^4*a^7*b^7*d^5 - 32*A^4*a^9*b^5*d^5) + (((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*((tan(c + d*x)^(1/2)*(512*A^2*a^8*b^8*d^7 - 448*A^2*a^10*b^6*d^7 + 128*A^2*a^12*b^4*d^7 + 64*A^2*a^14*b^2*d^7) - (((64*...`

3.403
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$$

3.403.1 Optimal result 3850
 3.403.2 Mathematica [C] (verified) 3851
 3.403.3 Rubi [A] (verified) 3851
 3.403.4 Maple [A] (verified) 3859
 3.403.5 Fricas [B] (verification not implemented) 3859
 3.403.6 Sympy [F(-1)] 3860
 3.403.7 Maxima [A] (verification not implemented) 3860
 3.403.8 Giac [F(-1)] 3861
 3.403.9 Mupad [B] (verification not implemented) 3861

3.403.1 Optimal result

Integrand size = 33, antiderivative size = 325

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx \\ &= -\frac{(b(A - B) - a(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)d} \\ &+ \frac{(b(A - B) - a(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)d} \\ &+ \frac{2b^{5/2}(Ab - aB) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{a^{5/2}(a^2 + b^2)d} \\ &+ \frac{(a(A - B) + b(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)d} \\ &- \frac{(a(A - B) + b(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)d} \\ &- \frac{2A}{3ad \tan^{\frac{3}{2}}(c + dx)} + \frac{2(Ab - aB)}{a^2d\sqrt{\tan(c + dx)}} \end{aligned}$$

output $2*b^{(5/2)}*(A*b-B*a)*\arctan(b^{(1/2)}*\tan(d*x+c)^{(1/2)}/a^{(1/2)})/a^{(5/2)}/(a^2+b^2)/d+1/2*(b*(A-B)-a*(A+B))*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/(a^2+b^2)/d*2^{(1/2)}+1/2*(b*(A-B)-a*(A+B))*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/(a^2+b^2)/d*2^{(1/2)}+1/4*(a*(A-B)+b*(A+B))*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(a^2+b^2)/d*2^{(1/2)}-1/4*(a*(A-B)+b*(A+B))*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(a^2+b^2)/d*2^{(1/2)}+2*(A*b-B*a)/a^2/d/\tan(d*x+c)^{(1/2)}-2/3*A/a/d/\tan(d*x+c)^{(3/2)}$

3.403.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.63 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.54

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$= \frac{3\sqrt[4]{-1}(A-iB) \arctan\left(\frac{(-1)^{3/4}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{a-ib} + \frac{6b^{5/2}(Ab-aB) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{a^{5/2}(a^2+b^2)} + \frac{3\sqrt[4]{-1}(A+iB) \operatorname{arctanh}\left(\frac{(-1)^{3/4}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{a+ib}$$

$3d$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])),x]`

output $((3*(-1)^{(1/4)}*(A - I*B)*\operatorname{ArcTan}[(-1)^{(3/4)}*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]])/(a - I*b) + (6*b^{(5/2)}*(A*b - a*B)*\operatorname{ArcTan}[\operatorname{Sqrt}[b]*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]]/\operatorname{Sqrt}[a])/(a^{(5/2)}*(a^2 + b^2)) + (3*(-1)^{(1/4)}*(A + I*B)*\operatorname{ArcTanh}[(-1)^{(3/4)}*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]])/(a + I*b) - (2*(a*A + (-3*A*b + 3*a*B)*\operatorname{Tan}[c + d*x]))/(a^2*\operatorname{Tan}[c + d*x]^{(3/2)}))/(3*d)$

3.403.3 Rubi [A] (verified)

Time = 1.45 (sec) , antiderivative size = 289, normalized size of antiderivative = 0.89, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4136, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.403. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$\begin{aligned}
& \int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx \\
& \quad \downarrow 3042 \\
& \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2}(a + b \tan(c + dx))} dx \\
& \quad \downarrow 4092 \\
& \frac{2 \int \frac{3(Ab \tan^2(c + dx) + aA \tan(c + dx) + Ab - aB)}{2 \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx}{3a} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{Ab \tan^2(c + dx) + aA \tan(c + dx) + Ab - aB}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx}{a} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{Ab \tan(c + dx)^2 + aA \tan(c + dx) + Ab - aB}{\tan(c + dx)^{3/2}(a + b \tan(c + dx))} dx}{a} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 4132 \\
& \frac{2 \int -\frac{Aa^2 + B \tan(c + dx)a^2 + bBa - Ab^2 - b(Ab - aB) \tan^2(c + dx)}{2\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{a} - \frac{2(Ab - aB)}{ad\sqrt{\tan(c + dx)}} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{Aa^2 + B \tan(c + dx)a^2 + bBa - Ab^2 - b(Ab - aB) \tan^2(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{a} - \frac{2(Ab - aB)}{ad\sqrt{\tan(c + dx)}} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{Aa^2 + B \tan(c + dx)a^2 + bBa - Ab^2 - b(Ab - aB) \tan(c + dx)^2}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{a} - \frac{2(Ab - aB)}{ad\sqrt{\tan(c + dx)}} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 4136 \\
& \frac{\int \frac{a^2(aA + bB) - a^2(Ab - aB) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx}{a^2 + b^2} - \frac{b^3(Ab - aB) \int \frac{\tan^2(c + dx) + 1}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{a^2 + b^2} - \frac{2(Ab - aB)}{ad\sqrt{\tan(c + dx)}} \\
& \quad \frac{a}{2A} \\
& \quad \frac{2A}{3ad \tan^{\frac{3}{2}}(c + dx)}
\end{aligned}$$

3.403. $\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx$

$$\begin{array}{c}
 \downarrow \text{3042} \\
 \frac{\int \frac{a^2(aA+bB)-a^2(Ab-aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{b^3(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2}}{a} - \frac{2(Ab-aB)}{ad\sqrt{\tan(c+dx)}} \\
 \hline
 \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
 \downarrow \text{4017} \\
 \frac{2 \int \frac{a^2(aA+bB-(Ab-aB)\tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} - \frac{b^3(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2}}{a} - \frac{2(Ab-aB)}{ad\sqrt{\tan(c+dx)}} \\
 \hline
 \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
 \downarrow \text{27} \\
 \frac{2a^2 \int \frac{aA+bB-(Ab-aB)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} - \frac{b^3(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2}}{a} - \frac{2(Ab-aB)}{ad\sqrt{\tan(c+dx)}} \\
 \hline
 \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
 \downarrow \text{1482} \\
 \frac{2a^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(b(A-B)-a(A+B)) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) - b^3(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{d(a^2+b^2)}}{a} - \frac{2(Ab-aB)}{ad\sqrt{\tan(c+dx)}} \\
 \hline
 \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
 \downarrow \text{1476} \\
 \frac{2a^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right) - b^3(Ab-aB) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{d(a^2+b^2)}}{a} - \frac{2(Ab-aB)}{ad\sqrt{\tan(c+dx)}} \\
 \hline
 \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
 \downarrow \text{1082}
 \end{array}$$

3.403. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$2a^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\int \frac{1}{\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)$$

$$d(a^2+b^2)$$

 a a

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)}$$

↓ 217

$$2a^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}})}{\sqrt{2}} - \frac{\arctan(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}})}{\sqrt{2}} \right) \right)$$

$$d(a^2+b^2)$$

 a a

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)}$$

↓ 1479

$$2a^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}})}{\sqrt{2}} - \frac{\arctan(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}})}{\sqrt{2}} \right) \right)$$

$$d(a^2+b^2)$$

 a a

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)}$$

↓ 25

$$2a^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}})}{\sqrt{2}} - \frac{\arctan(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}})}{\sqrt{2}} \right) \right)$$

$$d(a^2+b^2)$$

 a a

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)}$$

↓ 27

3.403. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$2a^2 \left(\frac{\frac{1}{2}(a(A-B)+b(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} \right)}{d(a^2+b^2)} \right)$$

a

a

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)}$$

↓ 1103

$$2a^2 \left(\frac{\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} \right)}{d(a^2+b^2)} \right)$$

a

a

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)}$$

↓ 4117

$$2a^2 \left(\frac{\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} \right)}{d(a^2+b^2)} \right)$$

a

a

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)}$$

↓ 73

$$2a^2 \left(\frac{\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} \right)}{d(a^2+b^2)} \right)$$

a

a

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)}$$

↓ 218

$$2a^2 \left(\frac{\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} \right)}{d(a^2+b^2)} \right)$$

a

a

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)}$$

3.403. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])),x]`

output `-((((-2*b^(5/2)*(A*b - a*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[a]*(a^2 + b^2)*d) + (2*a^2*(-1/2*((b*(A - B) - a*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])) + ((a*(A - B) + b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2))/((a^2 + b^2)*d))/a - (2*(A*b - a*B))/(a*d*Sqrt[Tan[c + d*x]])/a - (2*A)/(3*a*d*Tan[c + d*x]^(3/2))`

3.403.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

3.403. $\int \frac{A+B \tan (c+d x)}{\tan ^2(c+d x)(a+b \tan (c+d x))} d x$

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.403.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 286, normalized size of antiderivative = 0.88

method	result
derivativedivides	$\frac{2b^3(Ab-Ba) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{a^2(a^2+b^2)\sqrt{ab}} - \frac{2A}{3a \tan(dx+c)^{\frac{3}{2}}} - \frac{2(-Ab+Ba)}{a^2\sqrt{\tan(dx+c)}} + \frac{(-aA-Bb)\sqrt{2}}{a^2\sqrt{\tan(dx+c)}} \ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}\right)$
default	$\frac{2b^3(Ab-Ba) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{a^2(a^2+b^2)\sqrt{ab}} - \frac{2A}{3a \tan(dx+c)^{\frac{3}{2}}} - \frac{2(-Ab+Ba)}{a^2\sqrt{\tan(dx+c)}} + \frac{(-aA-Bb)\sqrt{2}}{a^2\sqrt{\tan(dx+c)}} \ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}\right)$

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*(2/a^2*b^3*(A*b-B*a)/(a^2+b^2)/(a*b)^(1/2)*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))-2/3/a*A/tan(d*x+c)^(3/2)-2/a^2*(-A*b+B*a)/tan(d*x+c)^(1/2)+2/(a^2+b^2)*(1/8*(-A*a-B*b)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/8*(A*b-B*a)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))`

3.403.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3129 vs. 2(281) = 562.

Time = 27.01 (sec) , antiderivative size = 6284, normalized size of antiderivative = 19.34

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x, algorithm="fracas")`

output Too large to include

3.403.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(5/2)/(a+b*tan(d*x+c)),x)`

output `Timed out`

3.403.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 256, normalized size of antiderivative = 0.79

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx =$$

$$\frac{24 (Bab^3 - Ab^4) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^4 + a^2b^2)\sqrt{ab}} + \frac{3 \left(2\sqrt{2}((A+B)a - (A-B)b) \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2} + 2\sqrt{\tan(dx+c)})\right) + 2\sqrt{2}((A+B)a - (A-B)b) \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2} - 2\sqrt{\tan(dx+c)})\right)\right)}{(a^4 + a^2b^2)\sqrt{ab}}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `-1/12*(24*(B*a*b^3 - A*b^4)*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/((a^4 + a^2*b^2)*sqrt(a*b)) + 3*(2*sqrt(2)*((A + B)*a - (A - B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A + B)*a - (A - B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((A - B)*a + (A + B)*b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((A - B)*a + (A + B)*b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))/(a^2 + b^2) + 8*(A*a + 3*(B*a - A*b)*tan(d*x + c))/(a^2*tan(d*x + c)^(3/2))/d`

3.403.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = \text{Timed out}$$

```
input integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x, algorithm="giac")
```

```
output Timed out
```

3.403.9 Mupad [B] (verification not implemented)

Time = 15.15 (sec) , antiderivative size = 16111, normalized size of antiderivative = 49.57

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = \text{Too large to display}$$

```
input int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + b*tan(c + d*x))),x)
```

```
output atan((((tan(c + d*x)^(1/2)*(64*A^4*a^14*b^9*d^5 + 32*A^4*a^18*b^5*d^5) + (-
((64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*(((-(
64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*(tan(c +
d*x)^(1/2)*(-(64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2
*b^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4))
)^(1/2)*(512*a^18*b^9*d^9 + 512*a^20*b^7*d^9 - 512*a^22*b^5*d^9 - 512*a^24
*b^3*d^9) - 512*A*a^16*b^10*d^8 - 512*A*a^18*b^8*d^8 + 384*A*a^20*b^6*d^8
+ 256*A*a^22*b^4*d^8 - 128*A*a^24*b^2*d^8) - tan(c + d*x)^(1/2)*(512*A^2*a
^15*b^10*d^7 + 448*A^2*a^19*b^6*d^7 - 128*A^2*a^21*b^4*d^7 - 64*A^2*a^23*b
^2*d^7))*(-(64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b
^2*d^4))^(1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(
1/2) + 384*A^3*a^15*b^9*d^6 - 32*A^3*a^19*b^5*d^6 - 32*A^3*a^21*b^3*d^6))*
(-(64*A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(
1/2) - 8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*1i +
(tan(c + d*x)^(1/2)*(64*A^4*a^14*b^9*d^5 + 32*A^4*a^18*b^5*d^5) + (-((64
A^4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) -
8*A^2*a*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*(((-(64*A
4*a^2*b^2*d^4 - A^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - ...
```

3.404
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.404.1 Optimal result	3862
3.404.2 Mathematica [C] (verified)	3863
3.404.3 Rubi [A] (verified)	3864
3.404.4 Maple [A] (verified)	3871
3.404.5 Fricas [B] (verification not implemented)	3872
3.404.6 Sympy [F]	3872
3.404.7 Maxima [A] (verification not implemented)	3873
3.404.8 Giac [F(-1)]	3873
3.404.9 Mupad [B] (verification not implemented)	3874

3.404.1 Optimal result

Integrand size = 33, antiderivative size = 436

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{(a^2(A-B) - b^2(A-B) + 2ab(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^2 d}$$

$$- \frac{(a^2(A-B) - b^2(A-B) + 2ab(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^2 d}$$

$$+ \frac{a^{3/2}(a^2Ab + 5Ab^3 - 3a^3B - 7ab^2B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{b^{5/2}(a^2+b^2)^2 d}$$

$$+ \frac{(2ab(A-B) - a^2(A+B) + b^2(A+B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^2 d}$$

$$- \frac{(2ab(A-B) - a^2(A+B) + b^2(A+B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^2 d}$$

$$- \frac{(aAb - 3a^2B - 2b^2B) \sqrt{\tan(c+dx)}}{b^2(a^2+b^2)d} + \frac{a(Ab - aB) \tan^{\frac{3}{2}}(c+dx)}{b(a^2+b^2)d(a+b \tan(c+dx))}$$

output $a^{3/2}*(A*a^2*b+5*A*b^3-3*B*a^3-7*B*a*b^2)*\arctan(b^{1/2}*\tan(d*x+c)^{1/2})/a^{1/2})/b^{5/2}/(a^2+b^2)^2/d-1/2*(a^2*(A-B)-b^2*(A-B)+2*a*b*(A+B))*\arctan(-1+2^{1/2}*\tan(d*x+c)^{1/2})/(a^2+b^2)^2/d*2^{1/2}-1/2*(a^2*(A-B)-b^2*(A-B)+2*a*b*(A+B))*\arctan(1+2^{1/2}*\tan(d*x+c)^{1/2})/(a^2+b^2)^2/d*2^{1/2})+1/4*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\ln(1-2^{1/2}*\tan(d*x+c)^{1/2}+\tan(d*x+c))/(a^2+b^2)^2/d*2^{1/2}-1/4*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\ln(1+2^{1/2}*\tan(d*x+c)^{1/2}+\tan(d*x+c))/(a^2+b^2)^2/d*2^{1/2}-(A*a*b-3*B*a^2-2*B*b^2)*\tan(d*x+c)^{1/2}/b^2/(a^2+b^2)/d+a*(A*b-B*a)*\tan(d*x+c)^{3/2}/b/(a^2+b^2)/d/(a+b*\tan(d*x+c))$

3.404.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.49 (sec) , antiderivative size = 275, normalized size of antiderivative = 0.63

$$\int \frac{\tan^{5/2}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= 2 \left((-Ab + 3aB) \sqrt{\tan(c+dx)} + \frac{(a^2Ab + 2Ab^3 - 3a^3B - 4ab^2B) \sqrt{\tan(c+dx)}}{2(a^2+b^2)} + bB \tan^{3/2}(c+dx) - \frac{(\sqrt[4]{-1}(a+ib)^2 b^{5/2}(i))}{\dots} \right)$$

input `Integrate[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2, x]`

output $(2*((-(A*b) + 3*a*B)*\text{Sqrt}[\text{Tan}[c + d*x]] + ((a^2*A*b + 2*A*b^3 - 3*a^3*B - 4*a*b^2*B)*\text{Sqrt}[\text{Tan}[c + d*x]]))/(2*(a^2 + b^2)) + b*B*\text{Tan}[c + d*x]^{3/2} - (((-1)^{1/4}*(a + I*b)^2*b^{5/2}*(I*A + B)*\text{ArcTan}[(-1)^{3/4}*\text{Sqrt}[\text{Tan}[c + d*x]]]) + a^{3/2}*(-(a^2*A*b) - 5*A*b^3 + 3*a^3*B + 7*a*b^2*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]] + (-1)^{3/4}*b^{5/2}*(I*a + b)^2*(I*B)*\text{ArcTanh}[(-1)^{3/4}*\text{Sqrt}[\text{Tan}[c + d*x]]])*(a + b*\text{Tan}[c + d*x])/(2*\text{Sqrt}[b]*(a^2 + b^2)^2))/(b^2*d*(a + b*\text{Tan}[c + d*x]))$

3.404.3 Rubi [A] (verified)

Time = 1.83 (sec) , antiderivative size = 379, normalized size of antiderivative = 0.87, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.697$, Rules used = {3042, 4088, 27, 3042, 4130, 27, 3042, 4136, 27, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^{5/2}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{4088} \\
 & \frac{\int -\frac{\sqrt{\tan(c+dx)}((-3Ba^2+Aba-2b^2B) \tan^2(c+dx)-2b(Ab-aB) \tan(c+dx)+3a(Ab-aB))}{2(a+b \tan(c+dx))} dx}{b(a^2+b^2)} + \\
 & \quad \frac{a(Ab-aB) \tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{27} \\
 & \frac{a(Ab-aB) \tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b \tan(c+dx))} - \\
 & \frac{\int \frac{\sqrt{\tan(c+dx)}((-3Ba^2+Aba-2b^2B) \tan^2(c+dx)-2b(Ab-aB) \tan(c+dx)+3a(Ab-aB))}{a+b \tan(c+dx)} dx}{2b(a^2+b^2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a(Ab-aB) \tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b \tan(c+dx))} - \\
 & \frac{\int \frac{\sqrt{\tan(c+dx)}((-3Ba^2+Aba-2b^2B) \tan(c+dx)^2-2b(Ab-aB) \tan(c+dx)+3a(Ab-aB))}{a+b \tan(c+dx)} dx}{2b(a^2+b^2)} \\
 & \quad \downarrow \text{4130} \\
 & \frac{a(Ab-aB) \tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b \tan(c+dx))} - \\
 & \frac{2 \int -\frac{-2(aA+bB) \tan(c+dx)b^2+(-3Ba^3+Aba^2-4b^2Ba+2Ab^3) \tan^2(c+dx)+a(-3Ba^2+Aba-2b^2B)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b} + \frac{2(-3a^2B+aAb-2b^2B) \sqrt{\tan(c+dx)}}{bd} \\
 & \quad \frac{2b(a^2+b^2)}{2b(a^2+b^2)}
 \end{aligned}$$

3.404. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \int \frac{-2(aA + bB) \tan(c + dx)b^2 + (-3Ba^3 + Aba^2 - 4b^2Ba + 2Ab^3) \tan^2(c + dx) + a(-3Ba^2 + Aba - 2b^2B)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{\frac{2(-3a^2B + aAb - 2b^2B) \sqrt{\tan(c + dx)}}{bd}} \\
 & \frac{2b(a^2 + b^2)}{\downarrow 3042} \\
 & \frac{\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \int \frac{-2(aA + bB) \tan(c + dx)b^2 + (-3Ba^3 + Aba^2 - 4b^2Ba + 2Ab^3) \tan(c + dx)^2 + a(-3Ba^2 + Aba - 2b^2B)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{\frac{2(-3a^2B + aAb - 2b^2B) \sqrt{\tan(c + dx)}}{bd}} \\
 & \frac{2b(a^2 + b^2)}{\downarrow 4136} \\
 & \frac{\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \int \frac{2((-Ba^2 + 2Aba + b^2B)b^2 + (Aa^2 + 2bBa - Ab^2) \tan(c + dx)b^2)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx + \frac{a^2(-3a^3B + a^2Ab - 7ab^2B + 5Ab^3) \int \frac{\tan^2(c + dx) + 1}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{a^2 + b^2}}{\frac{2(-3a^2B + aAb - 2b^2B) \sqrt{\tan(c + dx)}}{bd}} \\
 & \frac{2b(a^2 + b^2)}{\downarrow 27} \\
 & \frac{\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \frac{a^2(-3a^3B + a^2Ab - 7ab^2B + 5Ab^3) \int \frac{\tan^2(c + dx) + 1}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{a^2 + b^2} - \frac{2 \int \frac{(-Ba^2 + 2Aba + b^2B)b^2 + (Aa^2 + 2bBa - Ab^2) \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{a^2 + b^2}}{\frac{2(-3a^2B + aAb - 2b^2B) \sqrt{\tan(c + dx)}}{bd}} \\
 & \frac{2b(a^2 + b^2)}{\downarrow 3042} \\
 & \frac{\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{bd(a^2 + b^2)(a + b \tan(c + dx))} - \frac{a^2(-3a^3B + a^2Ab - 7ab^2B + 5Ab^3) \int \frac{\tan(c + dx)^2 + 1}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{a^2 + b^2} - \frac{2 \int \frac{(-Ba^2 + 2Aba + b^2B)b^2 + (Aa^2 + 2bBa - Ab^2) \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{a^2 + b^2}}{\frac{2(-3a^2B + aAb - 2b^2B) \sqrt{\tan(c + dx)}}{bd}} \\
 & \frac{2b(a^2 + b^2)}{\downarrow 4017}
 \end{aligned}$$

3.404. $\int \frac{\tan^{\frac{5}{2}}(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx$

$$\frac{\frac{2(-3a^2B+aAb-2b^2B)\sqrt{\tan(c+dx)}}{bd} - \frac{a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b\tan(c+dx))} - \frac{a^2(-3a^3B+a^2Ab-7ab^2B+5Ab^3)\int\frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}dx}{a^2+b^2} - 4\int\frac{b^2(-Ba^2+2Aba+b^2B+(Aa^2+2bBa-Ab^2)\tan^2(c+dx)+1)}{d(a^2+b^2)}}{2b(a^2+b^2)}$$

↓ 27

$$\frac{\frac{2(-3a^2B+aAb-2b^2B)\sqrt{\tan(c+dx)}}{bd} - \frac{a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b\tan(c+dx))} - \frac{a^2(-3a^3B+a^2Ab-7ab^2B+5Ab^3)\int\frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}dx}{a^2+b^2} - 4b^2\int\frac{-Ba^2+2Aba+b^2B+(Aa^2+2bBa-Ab^2)\tan^2(c+dx)+1}{d(a^2+b^2)}}{2b(a^2+b^2)}$$

↓ 1482

$$\frac{\frac{2(-3a^2B+aAb-2b^2B)\sqrt{\tan(c+dx)}}{bd} - \frac{a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b\tan(c+dx))} - \frac{a^2(-3a^3B+a^2Ab-7ab^2B+5Ab^3)\int\frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}dx}{a^2+b^2} - 4b^2\left(\frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B)\right)}{2b(a^2+b^2)}$$

↓ 1476

$$\frac{\frac{2(-3a^2B+aAb-2b^2B)\sqrt{\tan(c+dx)}}{bd} - \frac{a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b\tan(c+dx))} - \frac{a^2(-3a^3B+a^2Ab-7ab^2B+5Ab^3)\int\frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}dx}{a^2+b^2} - 4b^2\left(\frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B)\right)}{2b(a^2+b^2)}$$

↓ 1082

$$\frac{\frac{2(-3a^2B+aAb-2b^2B)\sqrt{\tan(c+dx)}}{bd} - \frac{a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b\tan(c+dx))} - \frac{a^2(-3a^3B+a^2Ab-7ab^2B+5Ab^3)\int\frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}dx}{a^2+b^2} - 4b^2\left(\frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B)\right)}{2b(a^2+b^2)}$$

↓ 217

3.404. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

$$\frac{2(-3a^2B+aAb-2b^2B)\sqrt{\tan(c+dx)}}{bd} - \frac{a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b\tan(c+dx))} - \frac{a^2(-3a^3B+a^2Ab-7ab^2B+5Ab^3)\int\frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}dx}{a^2+b^2} - \frac{4b^2\left(\frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B)\right)}{2b(a^2+b^2)}$$

↓ 1479

$$\frac{2(-3a^2B+aAb-2b^2B)\sqrt{\tan(c+dx)}}{bd} - \frac{a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b\tan(c+dx))} - \frac{a^2(-3a^3B+a^2Ab-7ab^2B+5Ab^3)\int\frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}dx}{a^2+b^2} - \frac{4b^2\left(\frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B)\right)}{2b(a^2+b^2)}$$

↓ 25

$$\frac{2(-3a^2B+aAb-2b^2B)\sqrt{\tan(c+dx)}}{bd} - \frac{a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b\tan(c+dx))} - \frac{a^2(-3a^3B+a^2Ab-7ab^2B+5Ab^3)\int\frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}dx}{a^2+b^2} - \frac{4b^2\left(\frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B)\right)}{2b(a^2+b^2)}$$

↓ 27

$$\frac{2(-3a^2B+aAb-2b^2B)\sqrt{\tan(c+dx)}}{bd} - \frac{a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b\tan(c+dx))} - \frac{a^2(-3a^3B+a^2Ab-7ab^2B+5Ab^3)\int\frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}dx}{a^2+b^2} - \frac{4b^2\left(\frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B)\right)}{2b(a^2+b^2)}$$

↓ 1103

$$\frac{2(-3a^2B+aAb-2b^2B)\sqrt{\tan(c+dx)}}{bd} - \frac{a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b\tan(c+dx))} - \frac{a^2(-3a^3B+a^2Ab-7ab^2B+5Ab^3)\int\frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}dx}{a^2+b^2} - \frac{4b^2\left(\frac{1}{2}(a^2(A-B))+2ab(A+B)-b^2(A-B)\right)}{2b(a^2+b^2)}$$

↓ 4117

3.404. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

$$\frac{2(-3a^2B+aAb-2b^2B)\sqrt{\tan(c+dx)}}{bd} - \frac{a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b\tan(c+dx))} - \frac{a^2(-3a^3B+a^2Ab-7ab^2B+5Ab^3)\int\frac{1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}d\tan(c+dx)}{d(a^2+b^2)} - 4b^2\left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B))\right)$$

↓ 73

$$\frac{2(-3a^2B+aAb-2b^2B)\sqrt{\tan(c+dx)}}{bd} - \frac{a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b\tan(c+dx))} - \frac{2a^2(-3a^3B+a^2Ab-7ab^2B+5Ab^3)\int\frac{1}{a+b\tan(c+dx)}d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} - 4b^2\left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B))\right)$$

↓ 218

$$\frac{2(-3a^2B+aAb-2b^2B)\sqrt{\tan(c+dx)}}{bd} - \frac{a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{bd(a^2+b^2)(a+b\tan(c+dx))} - \frac{2a^{3/2}(-3a^3B+a^2Ab-7ab^2B+5Ab^3)\arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{bd}(a^2+b^2)} - 4b^2\left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B))\right)$$

```
input Int[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]
```

```
output -1/2*(-(((2*a^(3/2)*(a^2*A*b + 5*A*b^3 - 3*a^3*B - 7*a*b^2*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[b]*(a^2 + b^2)*d) - (4*b^2*((a^2*(A - B) - b^2*(A - B) + 2*a*b*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])))/2 + ((2*a*b*(A - B) - a^2*(A + B) + b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2))/((a^2 + b^2)*d)/b + (2*(a*A*b - 3*a^2*B - 2*b^2*B)*Sqrt[Tan[c + d*x]])/(b*d))/(b*(a^2 + b^2)) + (a*(A*b - a*B)*Tan[c + d*x]^(3/2))/(b*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))
```

3.404. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

3.404.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

3.404.
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

```
rule 4130 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

```
rule 4136 Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
!GtQ[n, 0] && !LeQ[n, -1]
```

3.404.4 Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 351, normalized size of antiderivative = 0.81

method	result
derivativedivides	$\frac{2a^2 \left(\frac{-\frac{1}{2}Aa^2b - \frac{1}{2}Ab^3 + \frac{1}{2}Ba^3 + \frac{1}{2}Bab^2}{a+b \tan(dx+c)} (\sqrt{\tan(dx+c)}) + \frac{(Aa^2b+5Ab^3-3Ba^3-7Bab^2) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{2\sqrt{ab}} \right)}{\frac{2(\sqrt{\tan(dx+c)})B}{b^2} + \frac{b^2(a^2+b^2)^2}{b^2(a^2+b^2)^2}}$
default	$\frac{2a^2 \left(\frac{-\frac{1}{2}Aa^2b - \frac{1}{2}Ab^3 + \frac{1}{2}Ba^3 + \frac{1}{2}Bab^2}{a+b \tan(dx+c)} (\sqrt{\tan(dx+c)}) + \frac{(Aa^2b+5Ab^3-3Ba^3-7Bab^2) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{2\sqrt{ab}} \right)}{\frac{2(\sqrt{\tan(dx+c)})B}{b^2} + \frac{b^2(a^2+b^2)^2}{b^2(a^2+b^2)^2}}$

```
input int(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNV
ERBOSE)
```

$$3.404. \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

output $1/d*(2*\tan(d*x+c)^{(1/2)}*B/b^2+2*a^2/b^2/(a^2+b^2)^2*((-1/2*A*a^2*b-1/2*A*b^3+1/2*B*a^3+1/2*B*a*b^2)*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))+1/2*(A*a^2*b+5*A*b^3-3*B*a^3-7*B*a*b^2)/(a*b)^{(1/2)}*\arctan(b*\tan(d*x+c)^{(1/2)}/(a*b)^{(1/2)})))+2/(a^2+b^2)^2*(1/8*(-2*A*a*b+B*a^2-B*b^2)*2^{(1/2)}*(\ln((1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})))+1/8*(-A*a^2+A*b^2-2*B*a*b)*2^{(1/2)}*(\ln((1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})))))$

3.404.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5942 vs. 2(397) = 794.

Time = 36.20 (sec) , antiderivative size = 11910, normalized size of antiderivative = 27.32

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

output Too large to include

3.404.6 Sympy [F]

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \int \frac{(A+B\tan(c+dx))\tan^{\frac{5}{2}}(c+dx)}{(a+b\tan(c+dx))^2} dx$$

input `integrate(tan(d*x+c)**(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**(5/2)/(a + b*tan(c + d*x))**2, x)`

3.404.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 377, normalized size of antiderivative = 0.86

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx =$$

$$\frac{4(3Ba^5 - Aa^4b + 7Ba^3b^2 - 5Aa^2b^3) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right) - \frac{4(Ba^3 - Aa^2b)\sqrt{\tan(dx+c)}}{a^3b^2 + ab^4 + (a^2b^3 + b^5)\tan(dx+c)} + \frac{2\sqrt{2}((A-B)a^2 + 2(A+B)ab - (A-B)b^2)}{(a^4b^2 + 2a^2b^4 + b^6)\sqrt{ab}}}{(a^4b^2 + 2a^2b^4 + b^6)\sqrt{ab}}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `-1/4*(4*(3*B*a^5 - A*a^4*b + 7*B*a^3*b^2 - 5*A*a^2*b^3)*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/((a^4*b^2 + 2*a^2*b^4 + b^6)*sqrt(a*b)) - 4*(B*a^3 - A*a^2*b)*sqrt(tan(d*x + c))/(a^3*b^2 + a*b^4 + (a^2*b^3 + b^5)*tan(d*x + c)) + (2*sqrt(2)*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))/(a^4 + 2*a^2*b^2 + b^4) - 8*B*sqrt(tan(d*x + c))/b^2)/d`

3.404.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output `Timed out`

3.404. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

3.404.9 Mupad [B] (verification not implemented)

Time = 41.99 (sec) , antiderivative size = 18313, normalized size of antiderivative = 42.00

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \text{Too large to display}$$

```
input int((tan(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)
```

```
output (log((((((((((128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b^2)^2*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2) - (128*B*a*b^2*(7*a^4 + b^4 + 8*a^2*b^2))/d)*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (64*B^2*a*tan(c + d*x)^(1/2)*(18*a^10 - 15*b^10 + 17*a^2*b^8 - a^4*b^6 + 97*a^6*b^4 + 84*a^8*b^2))/(b^2*d^2*(a^2 + b^2)^2))*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (32*B^3*a^4*(127*a^2*b^6 - 112*b^8 - 9*a^8 + 173*a^4*b^4 + 21*a^6*b^2))/(b^3*d^3*(a^2 + b^2)^3))*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 - (16*B^4*tan(c + d*x)^(1/2)*(9*a^12 + 2*b^12 + 4*a^2*b^10 + 2*a^4*b^8 - 49*a^6*b^6 + 7*a^8*b^4 + 33*a^10*b^2))/(b^3*d^4*(a^2 + b^2)^4))*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 - (16*B^5*a^3*b^2*(3*a^2 + 7*b^2))/(d^5*(a^2 + b^2)^4))*(((192*B^4*a^2*b^6*d^4 - 16*B^4*b^8*d^4 - 16*B^4*a^8*d^4 - 608*B^4*a^4*b^4*d^4 + 192*B^4*a^6*b^2*d^4)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6*d^4 + 6*a^4*b^4*d^4 + 4*a^6*b^2*d^4))^(1/2))/4 + (log((((((((((128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b^2)^2*(-4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16*B^2*a*b^3*d^2 - 16*B^2*a^...
```

$$3.405 \quad \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.405.1 Optimal result	3875
3.405.2 Mathematica [C] (verified)	3876
3.405.3 Rubi [A] (verified)	3876
3.405.4 Maple [A] (verified)	3883
3.405.5 Fricas [B] (verification not implemented)	3883
3.405.6 Sympy [F]	3884
3.405.7 Maxima [A] (verification not implemented)	3884
3.405.8 Giac [F(-1)]	3885
3.405.9 Mupad [B] (verification not implemented)	3885

3.405.1 Optimal result

Integrand size = 33, antiderivative size = 391

$$\begin{aligned} & \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\ &= -\frac{(2ab(A-B) - a^2(A+B) + b^2(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^2 d} \\ & \quad + \frac{(2ab(A-B) - a^2(A+B) + b^2(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^2 d} \\ & \quad + \frac{\sqrt{a}(a^2Ab - 3Ab^3 + a^3B + 5ab^2B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{b^{3/2}(a^2+b^2)^2 d} \\ & \quad + \frac{(a^2(A-B) - b^2(A-B) + 2ab(A+B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^2 d} \\ & \quad - \frac{(a^2(A-B) - b^2(A-B) + 2ab(A+B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^2 d} \\ & \quad + \frac{a(Ab - aB)\sqrt{\tan(c+dx)}}{b(a^2+b^2)d(a+b \tan(c+dx))} \end{aligned}$$

$$3.405. \quad \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

output $\frac{1}{2}*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/(a^2+b^2)^{2/d*2^{(1/2)}}+1/2*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/(a^2+b^2)^{2/d*2^{(1/2)}}+1/4*(a^2*(A-B)-b^2*(A-B)+2*a*b*(A+B))*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(a^2+b^2)^{2/d*2^{(1/2)}}-1/4*(a^2*(A-B)-b^2*(A-B)+2*a*b*(A+B))*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(a^2+b^2)^{2/d*2^{(1/2)}}+(A*a^2*b-3*A*b^3+B*a^3+5*B*a*b^2)*\arctan(b^{(1/2)}*\tan(d*x+c)^{(1/2)}/a^{(1/2)})*a^{(1/2)}/b^{(3/2)}/(a^2+b^2)^{2/d}+a*(A*b-B*a)*\tan(d*x+c)^{(1/2)}/b/(a^2+b^2)/d/(a+b*\tan(d*x+c))$

3.405.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.01 (sec) , antiderivative size = 230, normalized size of antiderivative = 0.59

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{\sqrt{a}(a^2Ab-3Ab^3+a^3B+5ab^2B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right) + \sqrt[4]{-1}b^{3/2}((a+ib)^2(A-iB) \arctan((-1)^{3/4}\sqrt{\tan(c+dx)}) + (a-ib)^2(A+iB) \arctan((-1)^{3/4}\sqrt{\tan(c+dx)})}}{\sqrt{b}(a^2+b^2)^2} + \frac{bd}{bd}$$

input `Integrate[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2, x]`

output $((\text{Sqrt}[a]*(a^2*A*b - 3*A*b^3 + a^3*B + 5*a*b^2*B))*\text{ArcTan}[\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a] + (-1)^{(1/4)}*b^{(3/2)}*((a + I*b)^2*(A - I*B)*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]] + (a - I*b)^2*(A + I*B)*\text{ArcTanh}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]]))/(\text{Sqrt}[b]*(a^2 + b^2)^2) - (2*B*\text{Sqrt}[\text{Tan}[c + d*x]])/(a + b*\text{Tan}[c + d*x]) + ((a*A*b + a^2*B + 2*b^2*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/((a^2 + b^2)*(a + b*\text{Tan}[c + d*x]))/(b*d)$

3.405.3 Rubi [A] (verified)

Time = 1.38 (sec) , antiderivative size = 335, normalized size of antiderivative = 0.86, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 4088, 27, 3042, 4136, 27, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.405. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\begin{aligned}
& \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{\tan(c+dx)^{3/2}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\
& \quad \downarrow \text{4088} \\
& \frac{\int -\frac{((Ba^2+Aba+2b^2B) \tan^2(c+dx))-2b(Ab-aB) \tan(c+dx)+a(Ab-aB)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b(a^2+b^2)} + \frac{a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)(a+b \tan(c+dx))} \\
& \quad \downarrow \text{27} \\
& \frac{a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)(a+b \tan(c+dx))} - \frac{\int -\frac{((Ba^2+Aba+2b^2B) \tan^2(c+dx))-2b(Ab-aB) \tan(c+dx)+a(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{2b(a^2+b^2)} \\
& \quad \downarrow \text{3042} \\
& \frac{a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)(a+b \tan(c+dx))} - \frac{\int -\frac{((Ba^2+Aba+2b^2B) \tan(c+dx)^2)-2b(Ab-aB) \tan(c+dx)+a(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{2b(a^2+b^2)} \\
& \quad \downarrow \text{4136} \\
& \frac{a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)(a+b \tan(c+dx))} - \frac{\int \frac{2(b(Aa^2+2bBa-Ab^2))-b(-Ba^2+2Aba+b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2}}{2b(a^2+b^2)} \\
& \quad \downarrow \text{27} \\
& \frac{a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)(a+b \tan(c+dx))} - \frac{2 \int \frac{b(Aa^2+2bBa-Ab^2))-b(-Ba^2+2Aba+b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2}}{2b(a^2+b^2)} \\
& \quad \downarrow \text{3042} \\
& \frac{a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)(a+b \tan(c+dx))} - \frac{2 \int \frac{b(Aa^2+2bBa-Ab^2))-b(-Ba^2+2Aba+b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2}}{2b(a^2+b^2)} \\
& \quad \downarrow \text{4017}
\end{aligned}$$

3.405. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\frac{\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)(a + b \tan(c + dx))} - 4 \int \frac{b(Aa^2 + 2bBa - Ab^2 - (-Ba^2 + 2Aba + b^2B)\tan(c + dx))}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{a(a^3B + a^2Ab + 5ab^2B - 3Ab^3) \int \frac{\tan(c + dx)^2 + 1}{\sqrt{\tan(c + dx)(a + b \tan(c + dx))}} dx}{a^2 + b^2}}{d(a^2 + b^2)} = \frac{2b(a^2 + b^2)}{27}$$

$$\frac{\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)(a + b \tan(c + dx))} - 4b \int \frac{Aa^2 + 2bBa - Ab^2 - (-Ba^2 + 2Aba + b^2B)\tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{a(a^3B + a^2Ab + 5ab^2B - 3Ab^3) \int \frac{\tan(c + dx)^2 + 1}{\sqrt{\tan(c + dx)(a + b \tan(c + dx))}} dx}{a^2 + b^2}}{d(a^2 + b^2)} = \frac{2b(a^2 + b^2)}{1482}$$

$$\frac{\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)(a + b \tan(c + dx))} - 4b \left(\frac{1}{2}(a^2(A - B) + 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(-a^2(A + B) + 2ab(A - B) + b^2(A + B)) \int \frac{\tan(c + dx) + 1}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} \right)}{d(a^2 + b^2)} = \frac{2b(a^2 + b^2)}{1476}$$

$$\frac{\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)(a + b \tan(c + dx))} - 4b \left(\frac{1}{2}(a^2(A - B) + 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(-a^2(A + B) + 2ab(A - B) + b^2(A + B)) \left(\frac{1}{2} \int \frac{1}{\tan(c + dx) - \sqrt{2}\sqrt{\tan(c + dx)}} \right) \right)}{d(a^2 + b^2)} = \frac{2b(a^2 + b^2)}{1082}$$

$$\frac{\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)(a + b \tan(c + dx))} - 4b \left(\frac{1}{2}(a^2(A - B) + 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(-a^2(A + B) + 2ab(A - B) + b^2(A + B)) \left(\frac{\int \frac{1}{-\tan(c + dx) - 1} d(1 - \sqrt{2}\sqrt{\tan(c + dx)})}{\sqrt{2}} \right) \right)}{d(a^2 + b^2)} = \frac{2b(a^2 + b^2)}{217}$$

$$\frac{\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)(a + b \tan(c + dx))} - 4b \left(\frac{1}{2}(a^2(A - B) + 2ab(A + B) - b^2(A - B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(-a^2(A + B) + 2ab(A - B) + b^2(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c + dx)} + 1)}{\sqrt{2}} \right) \right)}{d(a^2 + b^2)} = \frac{2b(a^2 + b^2)}{1479}$$

3.405. $\int \frac{\tan^{\frac{3}{2}}(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx$

$$\frac{\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)(a + b \tan(c + dx))} - 4b \left(\frac{\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B) + 2ab(A-B) + b^2) \right)}{d(a^2 + b^2)}}{2b(a^2 + b^2)}$$

25

$$\frac{\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)(a + b \tan(c + dx))} - 4b \left(\frac{\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B) + 2ab(A-B) + b^2) \right)}{d(a^2 + b^2)}}{2b(a^2 + b^2)}$$

27

$$\frac{\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)(a + b \tan(c + dx))} - 4b \left(\frac{\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B) + 2ab(A-B) + b^2) \right)}{d(a^2 + b^2)}}{2b(a^2 + b^2)}$$

1103

$$\frac{\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)(a + b \tan(c + dx))} - 4b \left(\frac{\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B) + 2ab(A-B) + b^2) \right)}{d(a^2 + b^2)}}{2b(a^2 + b^2)}$$

4117

$$\frac{\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)(a + b \tan(c + dx))} - 4b \left(\frac{\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B) + 2ab(A-B) + b^2) \right)}{d(a^2 + b^2)}}{2b(a^2 + b^2)}$$

73

3.405. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\frac{4b \left(\frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2} (-a^2(A+B) + 2ab(A-B) + b^2(A-B)) \right)}{bd(a^2 + b^2)(a + b \tan(c + dx)) \sqrt{\tan(c + dx)}} - \frac{1}{2b(a^2 + b^2)}$$

↓ 218

$$\frac{4b \left(\frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2} (-a^2(A+B) + 2ab(A-B) + b^2(A-B)) \right)}{bd(a^2 + b^2)(a + b \tan(c + dx)) \sqrt{\tan(c + dx)}} - \frac{1}{2b(a^2 + b^2)}$$

```
input Int[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]
```

```
output -1/2*((-2*Sqrt[a]*(a^2*A*b - 3*A*b^3 + a^3*B + 5*a*b^2*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[b]*(a^2 + b^2)*d) + (4*b*(-1/2*((2*a*b*(A - B) - a^2*(A + B) + b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]] + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])) + ((a^2*(A - B) - b^2*(A - B) + 2*a*b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)/(b*(a^2 + b^2)) + (a*(A*b - a*B)*Sqrt[Tan[c + d*x]])/(b*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))
```

3.405.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

3.405. $\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

- rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.405.
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4136 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]) + (C_)*tan[(e_) + (f_)*(x_)])^2)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.405.
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.405.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 336, normalized size of antiderivative = 0.86

method	result
derivativedivides	$2a \left(\frac{(A a^2 b + A b^3 - B a^3 - B a b^2) (\sqrt{\tan(dx+c)})}{2b(a+b \tan(dx+c))} + \frac{(A a^2 b - 3A b^3 + B a^3 + 5B a b^2) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{2b\sqrt{ab}} \right) \frac{(-A a^2 + A b^2 - 2B a b)}{(a^2 + b^2)^2} + \dots$
default	$2a \left(\frac{(A a^2 b + A b^3 - B a^3 - B a b^2) (\sqrt{\tan(dx+c)})}{2b(a+b \tan(dx+c))} + \frac{(A a^2 b - 3A b^3 + B a^3 + 5B a b^2) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{2b\sqrt{ab}} \right) \frac{(-A a^2 + A b^2 - 2B a b)}{(a^2 + b^2)^2} + \dots$

input `int(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNV
ERBOSE)`

output `1/d*(2*a/(a^2+b^2)^2*(1/2*(A*a^2*b+A*b^3-B*a^3-B*a*b^2)/b*tan(d*x+c)^(1/2)
/(a+b*tan(d*x+c))+1/2*(A*a^2*b-3*A*b^3+B*a^3+5*B*a*b^2)/b/(a*b)^(1/2)*arct
an(b*tan(d*x+c)^(1/2)/(a*b)^(1/2)))+2/(a^2+b^2)^2*(1/8*(-A*a^2+A*b^2-2*B*a
*b)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x
+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^
(1/2)*tan(d*x+c)^(1/2)))+1/8*(2*A*a*b-B*a^2+B*b^2)*2^(1/2)*(ln((1-2^(1/2)*
tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*ar
ctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))`

3.405.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5899 vs. 2(352) = 704.

Time = 21.05 (sec) , antiderivative size = 11824, normalized size of antiderivative = 30.24

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorith
m="fricas")`

output Too large to include

3.405. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.405.6 Sympy [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \int \frac{(A+B\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)}{(a+b\tan(c+dx))^2} dx$$

input `integrate(tan(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**(3/2)/(a + b*tan(c + d*x))**2, x)`

3.405.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 354, normalized size of antiderivative = 0.91

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{4(Ba^4 + Aa^3b + 5Ba^2b^2 - 3Aab^3) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^4b + 2a^2b^3 + b^5)\sqrt{ab}} - \frac{4(Ba^2 - Aab)\sqrt{\tan(dx+c)}}{a^3b + ab^3 + (a^2b^2 + b^4)\tan(dx+c)} - \frac{2\sqrt{2}((A+B)a^2 - 2(A-B)ab - (A+B)b^2) \arctan\left(\frac{1}{\sqrt{2}}\sqrt{\tan(dx+c)}\right)}{(a^4b + 2a^2b^3 + b^5)\sqrt{ab}}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `1/4*(4*(B*a^4 + A*a^3*b + 5*B*a^2*b^2 - 3*A*a*b^3)*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b)))/((a^4*b + 2*a^2*b^3 + b^5)*sqrt(a*b)) - 4*(B*a^2 - A*a*b)*sqrt(tan(d*x + c))/(a^3*b + a*b^3 + (a^2*b^2 + b^4)*tan(d*x + c)) - (2*sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))/(a^4 + 2*a^2*b^2 + b^4)/d`

3.405. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

3.405.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Timed out}$$

```
input integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm
m="giac")
```

```
output Timed out
```

3.405.9 Mupad [B] (verification not implemented)

Time = 38.27 (sec) , antiderivative size = 17579, normalized size of antiderivative = 44.96

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Too large to display}$$

```
input int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)
```

```
output (log((((((((((128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b^2)^2*((4*(-A^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*A^2*a*b^3*d^2 + 16*A^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2) - (128*A*a*b^2*(5*b^4 - a^4 + 4*a^2*b^2))/d)*((4*(-A^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*A^2*a*b^3*d^2 + 16*A^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (64*A^2*a*b^2*tan(c + d*x)^(1/2)*(a^6 - 15*b^6 + 35*a^2*b^4 - 13*a^4*b^2))/(d^2*(a^2 + b^2)^2))*((4*(-A^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*A^2*a*b^3*d^2 + 16*A^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (32*A^3*a^2*b*(a^6 - 39*b^6 + 43*a^2*b^4 - 13*a^4*b^2))/(d^3*(a^2 + b^2)^3))*((4*(-A^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*A^2*a*b^3*d^2 + 16*A^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 - (16*A^4*b*tan(c + d*x)^(1/2)*(a^8 + 2*b^8 - 5*a^2*b^6 + 17*a^4*b^4 - 7*a^6*b^2))/(d^4*(a^2 + b^2)^4))*((4*(-A^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*A^2*a*b^3*d^2 + 16*A^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (16*A^5*a*b^4*(a^2 - 3*b^2))/(d^5*(a^2 + b^2)^4))*((((192*A^4*a^2*b^6*d^4 - 16*A^4*b^8*d^4 - 16*A^4*a^8*d^4 - 608*A^4*a^4*b^4*d^4 + 192*A^4*a^6*b^2*d^4)^(1/2) - 16*A^2*a*b^3*d^2 + 16*A^2*a^3*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6*d^4 + 6*a^4*b^4*d^4 + 4*a^6*b^2*d^4))^(1/2))/4 + (log((((((((((128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b^2)^2*(-(4*(-A^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16*A^2*a*b^3*d^2 - 16*A^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2) - (128*A*a*b^2*(5*b^4 - a^4 + 4*a^2*b^2))/d)*(-(4...
```

3.405. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

3.406
$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.406.1 Optimal result	3886
3.406.2 Mathematica [C] (verified)	3887
3.406.3 Rubi [A] (verified)	3888
3.406.4 Maple [A] (verified)	3894
3.406.5 Fricas [B] (verification not implemented)	3895
3.406.6 Sympy [F]	3895
3.406.7 Maxima [A] (verification not implemented)	3896
3.406.8 Giac [F(-1)]	3896
3.406.9 Mupad [B] (verification not implemented)	3897

3.406.1 Optimal result

Integrand size = 33, antiderivative size = 391

$$\begin{aligned} & \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\ &= -\frac{(a^2(A-B) - b^2(A-B) + 2ab(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2 + b^2)^2 d} \\ &+ \frac{(a^2(A-B) - b^2(A-B) + 2ab(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2 + b^2)^2 d} \\ &- \frac{(3a^2Ab - Ab^3 - a^3B + 3ab^2B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}(a^2 + b^2)^2 d} \\ &- \frac{(2ab(A-B) - a^2(A+B) + b^2(A+B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}(a^2 + b^2)^2 d} \\ &+ \frac{(2ab(A-B) - a^2(A+B) + b^2(A+B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}(a^2 + b^2)^2 d} \\ &- \frac{(Ab - aB)\sqrt{\tan(c+dx)}}{(a^2 + b^2)d(a + b \tan(c+dx))} \end{aligned}$$

output $\frac{1}{2}(a^2(A-B)-b^2(A-B)+2ab(A+B))\arctan(-1+2^{1/2}\tan(dx+c)^{1/2})/(a^2+b^2)^2/d^{1/2}+1/2(a^2(A-B)-b^2(A-B)+2ab(A+B))\arctan(1+2^{1/2}\tan(dx+c)^{1/2})/(a^2+b^2)^2/d^{1/2}-1/4(2ab(A-B)-a^2(A+B)+b^2(A+B))\ln(1-2^{1/2}\tan(dx+c)^{1/2}+\tan(dx+c))/(a^2+b^2)^2/d^{1/2}+1/4(2ab(A-B)-a^2(A+B)+b^2(A+B))\ln(1+2^{1/2}\tan(dx+c)^{1/2}+\tan(dx+c))/(a^2+b^2)^2/d^{1/2}-(3Aa^2b-Ab^3-Ba^3+3Ba^2b^2)\arctan(b^{1/2}\tan(dx+c)^{1/2}/a^{1/2})/(a^2+b^2)^2/d/a^{1/2}/b^{1/2}-(Ab-Ba)\tan(dx+c)^{1/2}/(a^2+b^2)/d/(a+b\tan(dx+c))$

3.406.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.52 (sec) , antiderivative size = 220, normalized size of antiderivative = 0.56

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{\sqrt{a}(-3a^2Ab+Ab^3+a^3B-3ab^2B)\arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{b}(a^2+b^2)} + \frac{\sqrt[4]{-1}a((a+ib)^2(iA+B)\arctan((-1)^{3/4}\sqrt{\tan(c+dx)})+(a-ib)^2(-iA+B)\arctan((-1)^{1/4}\sqrt{\tan(c+dx)})}{a^2+b^2}$$

input `Integrate[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2, x]`

output $((\sqrt{a}(-3a^2Ab+Ab^3+a^3B-3ab^2B)\text{ArcTan}[\sqrt{b}\sqrt{\text{Tan}[c+d*x]}/\sqrt{a}])/\sqrt{b}(a^2+b^2))+((-1)^{1/4}a((a+I*b)^2(I*A+B)\text{ArcTan}[(-1)^{3/4}\sqrt{\text{Tan}[c+d*x]}]+(a-I*b)^2((-I)*A+B)\text{ArcTan}[(-1)^{3/4}\sqrt{\text{Tan}[c+d*x]}]))/(a^2+b^2)+(-(A*b)+a*B)\text{Sqrt}[\text{Tan}[c+d*x]]+(b*(A*b-a*B)\text{Tan}[c+d*x]^{3/2})/(a+b*\text{Tan}[c+d*x])/(a*(a^2+b^2)*d)$

3.406.3 Rubi [A] (verified)

Time = 1.33 (sec) , antiderivative size = 334, normalized size of antiderivative = 0.85, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 4091, 27, 3042, 4136, 27, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

↓ 3042

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

↓ 4091

$$\frac{\int -\frac{-b(Ab-aB) \tan^2(c+dx)+2b(aA+bB) \tan(c+dx)+b(Ab-aB)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b(a^2+b^2)} - \frac{(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)(a+b \tan(c+dx))}$$

↓ 27

$$\frac{\int \frac{-b(Ab-aB) \tan^2(c+dx)+2b(aA+bB) \tan(c+dx)+b(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{2b(a^2+b^2)} - \frac{(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)(a+b \tan(c+dx))}$$

↓ 3042

$$\frac{\int \frac{-b(Ab-aB) \tan(c+dx)^2+2b(aA+bB) \tan(c+dx)+b(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{2b(a^2+b^2)} - \frac{(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)(a+b \tan(c+dx))}$$

↓ 4136

$$\frac{\int \frac{2(b(-Ba^2+2Aba+b^2B)+b(Aa^2+2bBa-Ab^2) \tan(c+dx))}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} - \frac{b(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2}}{2b(a^2+b^2)}$$

↓ 27

$$\frac{(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)(a+b \tan(c+dx))}$$

3.406. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\frac{2 \int \frac{b(-Ba^2+2Aba+b^2B)+b(Aa^2+2bBa-Ab^2) \tan(c+dx)}{\sqrt{\tan(c+dx)} a^2+b^2} dx - \frac{b(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx}{a^2+b^2}}{2b(a^2+b^2)}$$

$$\frac{(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)(a+b \tan(c+dx))}$$

↓ 3042

$$\frac{2 \int \frac{b(-Ba^2+2Aba+b^2B)+b(Aa^2+2bBa-Ab^2) \tan(c+dx)}{\sqrt{\tan(c+dx)} a^2+b^2} dx - \frac{b(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx}{a^2+b^2}}{2b(a^2+b^2)}$$

$$\frac{(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)(a+b \tan(c+dx))}$$

↓ 4017

$$\frac{4 \int \frac{b(-Ba^2+2Aba+b^2B+(Aa^2+2bBa-Ab^2) \tan(c+dx)) d\sqrt{\tan(c+dx)}}{\tan^2(c+dx)+1} - \frac{b(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx}{a^2+b^2}}{d(a^2+b^2)}$$

$$\frac{2b(a^2+b^2)}{d(a^2+b^2)(a+b \tan(c+dx))}$$

↓ 27

$$\frac{4b \int \frac{-Ba^2+2Aba+b^2B+(Aa^2+2bBa-Ab^2) \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{b(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx}{a^2+b^2}}{d(a^2+b^2)}$$

$$\frac{2b(a^2+b^2)}{d(a^2+b^2)(a+b \tan(c+dx))}$$

↓ 1482

$$\frac{4b \left(\frac{1}{2} (-a^2(A+B)) + 2ab(A-B) + b^2(A+B) \right) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)}$$

$$\frac{(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)(a+b \tan(c+dx))}$$

↓ 1476

3.406. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\frac{4b \left(\frac{1}{2}(-a^2(A+B)) + 2ab(A-B) + b^2(A+B) \right) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)}} \right)}{d(a^2+b^2)} \quad 2b(a^2 + b^2)$$

$$\frac{(Ab - aB)\sqrt{\tan(c + dx)}}{d(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 1082

$$\frac{4b \left(\frac{1}{2}(-a^2(A+B)) + 2ab(A-B) + b^2(A+B) \right) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right)}{d(a^2+b^2)} \quad 2b(a^2 + b^2)$$

$$\frac{(Ab - aB)\sqrt{\tan(c + dx)}}{d(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 217

$$\frac{4b \left(\frac{1}{2}(-a^2(A+B)) + 2ab(A-B) + b^2(A+B) \right) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\sqrt{2}} \right)}{d(a^2+b^2)} \quad 2b(a^2 + b^2)$$

$$\frac{(Ab - aB)\sqrt{\tan(c + dx)}}{d(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 1479

$$4b \left(\frac{1}{2}(-a^2(A+B)) + 2ab(A-B) + b^2(A+B) \right) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\sqrt{2}} \right)$$

$$\frac{(Ab - aB)\sqrt{\tan(c + dx)}}{d(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 25

$$4b \left(\frac{1}{2}(-a^2(A+B)) + 2ab(A-B) + b^2(A+B) \right) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\sqrt{2}} \right)$$

$$\frac{(Ab - aB)\sqrt{\tan(c + dx)}}{d(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 27

3.406. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\frac{4b \left(\frac{1}{2} (-a^2(A+B)) + 2ab(A-B) + b^2(A+B) \right) \left(\int \frac{\sqrt{2-2\sqrt{\tan(c+dx)}}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)} + 1}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right)}{d(a^2 + b^2)}$$

$$\frac{(Ab - aB)\sqrt{\tan(c+dx)}}{d(a^2 + b^2)(a + b\tan(c+dx))}$$

↓ 1103

$$\frac{4b \left(\frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \right) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} (-a^2(A+B) + 2ab(A-B) + b^2(A+B)) \left(\frac{\log(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\log(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right)}{d(a^2 + b^2)}$$

$$\frac{(Ab - aB)\sqrt{\tan(c+dx)}}{d(a^2 + b^2)(a + b\tan(c+dx))}$$

↓ 4117

$$\frac{4b \left(\frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \right) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} (-a^2(A+B) + 2ab(A-B) + b^2(A+B)) \left(\frac{\log(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\log(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right)}{d(a^2 + b^2)}$$

$$\frac{(Ab - aB)\sqrt{\tan(c+dx)}}{d(a^2 + b^2)(a + b\tan(c+dx))}$$

↓ 73

$$\frac{4b \left(\frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \right) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} (-a^2(A+B) + 2ab(A-B) + b^2(A+B)) \left(\frac{\log(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\log(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right)}{d(a^2 + b^2)}$$

$$\frac{(Ab - aB)\sqrt{\tan(c+dx)}}{d(a^2 + b^2)(a + b\tan(c+dx))}$$

↓ 218

$$\frac{4b \left(\frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \right) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} (-a^2(A+B) + 2ab(A-B) + b^2(A+B)) \left(\frac{\log(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\log(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right)}{d(a^2 + b^2)}$$

$$\frac{(Ab - aB)\sqrt{\tan(c+dx)}}{d(a^2 + b^2)(a + b\tan(c+dx))}$$

input `Int[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

$$3.406. \quad \int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

```
output ((-2*Sqrt[b]*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B)*ArcTan[(Sqrt[b]*Sqrt[
Tan[c + d*x]]/Sqrt[a])]/(Sqrt[a]*(a^2 + b^2)*d) + (4*b*(((a^2*(A - B) - b
^2*(A - B) + 2*a*b*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt
[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])))/2 + ((2*a*b*(A - B
) - a^2*(A + B) + b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] +
Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]
/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)/(2*b*(a^2 + b^2)) - ((A*b - a*B)*Sqrt[
Tan[c + d*x]]/((a^2 + b^2)*d*(a + b*Tan[c + d*x])))
```

3.406.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 73 Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

```
rule 4091 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m +
1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n)
+ A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan
[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x] /; FreeQ
[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || Integers
Q[2*m, 2*n])
```

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

```
rule 4136 Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.406.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 332, normalized size of antiderivative = 0.85

method	result
derivativedivides	$2 \left(\frac{\left(\frac{1}{2} A a^2 b + \frac{1}{2} A b^3 - \frac{1}{2} B a^3 - \frac{1}{2} B a b^2\right) (\sqrt{\tan(dx+c)}}{a+b \tan(dx+c)} + \frac{(3 A a^2 b - A b^3 - B a^3 + 3 B a b^2) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{2\sqrt{ab}} \right)}{(a^2+b^2)^2} + \frac{(2 A a b - B a^2)}{(a^2+b^2)^2}$
default	$2 \left(\frac{\left(\frac{1}{2} A a^2 b + \frac{1}{2} A b^3 - \frac{1}{2} B a^3 - \frac{1}{2} B a b^2\right) (\sqrt{\tan(dx+c)}}{a+b \tan(dx+c)} + \frac{(3 A a^2 b - A b^3 - B a^3 + 3 B a b^2) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{2\sqrt{ab}} \right)}{(a^2+b^2)^2} + \frac{(2 A a b - B a^2)}{(a^2+b^2)^2}$

3.406. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

input `int(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNV
ERBOSE)`

output `1/d*(-2/(a^2+b^2)^2*((1/2*A*a^2*b+1/2*A*b^3-1/2*B*a^3-1/2*B*a*b^2)*tan(d*x
+c)^(1/2)/(a+b*tan(d*x+c))+1/2*(3*A*a^2*b-A*b^3-B*a^3+3*B*a*b^2)/(a*b)^(1/
2)*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2)))+2/(a^2+b^2)^2*(1/8*(2*A*a*b-B*a
^2+B*b^2)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*t
an(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan
(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/8*(A*a^2-A*b^2+2*B*a*b)*2^(1/2)*(ln((1-2^
(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))
)+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2
))))))`

3.406.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5901 vs. 2(347) = 694.

Time = 15.76 (sec) , antiderivative size = 11827, normalized size of antiderivative = 30.25

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm
m="fricas")`

output Too large to include

3.406.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \int \frac{(A+B\tan(c+dx))\sqrt{\tan(c+dx)}}{(a+b\tan(c+dx))^2} dx$$

input `integrate(tan(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Integral((A + B*tan(c + d*x))*sqrt(tan(c + d*x))/(a + b*tan(c + d*x))**2,
x)`

3.406. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

3.406.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 341, normalized size of antiderivative = 0.87

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{4(Ba^3-3Aa^2b-3Bab^2+Ab^3)\arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^4+2a^2b^2+b^4)\sqrt{ab}} + \frac{4(Ba-Ab)\sqrt{\tan(dx+c)}}{a^3+ab^2+(a^2b+b^3)\tan(dx+c)} + \frac{2\sqrt{2}((A-B)a^2+2(A+B)ab-(A-B)b^2)\arctan\left(\frac{1}{2}\sqrt{2}\tan(dx+c)\right)}{(a^4+2a^2b^2+b^4)\sqrt{ab}}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `1/4*(4*(B*a^3 - 3*A*a^2*b - 3*B*a*b^2 + A*b^3)*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b)))/((a^4 + 2*a^2*b^2 + b^4)*sqrt(a*b)) + 4*(B*a - A*b)*sqrt(tan(d*x + c))/(a^3 + a*b^2 + (a^2*b + b^3)*tan(d*x + c)) + (2*sqrt(2)*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1)/(a^4 + 2*a^2*b^2 + b^4)/d`

3.406.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output `Timed out`

3.406.9 Mupad [B] (verification not implemented)

Time = 38.28 (sec) , antiderivative size = 17089, normalized size of antiderivative = 43.71

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Too large to display}$$

input `int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`

output `(log((16*B^5*a*b^4*(a^2 - 3*b^2))/(d^5*(a^2 + b^2)^4) - ((((((((((128*B*a*b^2*(5*b^4 - a^4 + 4*a^2*b^2))/d - 128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b^2)^2*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2))^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2))^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 - (64*B^2*a*b^2*tan(c + d*x)^(1/2)*(a^6 - 15*b^6 + 35*a^2*b^4 - 13*a^4*b^2))/(d^2*(a^2 + b^2)^2))*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2))^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 - (32*B^3*a^2*b*(a^6 - 39*b^6 + 43*a^2*b^4 - 13*a^4*b^2))/(d^3*(a^2 + b^2)^3))*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2))^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (16*B^4*b*tan(c + d*x)^(1/2)*(a^8 + 2*b^8 - 5*a^2*b^6 + 17*a^4*b^4 - 7*a^6*b^2))/(d^4*(a^2 + b^2)^4))*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2))^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4*(((192*B^4*a^2*b^6*d^4 - 16*B^4*b^8*d^4 - 16*B^4*a^8*d^4 - 608*B^4*a^4*b^4*d^4 + 192*B^4*a^6*b^2*d^4)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6*d^4 + 6*a^4*b^4*d^4 + 4*a^6*b^2*d^4))^(1/2))/4 + (log((16*B^5*a*b^4*(a^2 - 3*b^2))/(d^5*(a^2 + b^2)^4) - ((((((((((128*B*a*b^2*(5*b^4 - a^4 + 4*a^2*b^2))/d - 128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b^2)^2*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2))^2)^(1/2) + 16*B^2*a*b^3*d^2 - ...`

$$3.407 \quad \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^2} dx$$

3.407.1 Optimal result	3898
3.407.2 Mathematica [C] (verified)	3899
3.407.3 Rubi [A] (verified)	3899
3.407.4 Maple [A] (verified)	3906
3.407.5 Fricas [B] (verification not implemented)	3907
3.407.6 Sympy [F]	3907
3.407.7 Maxima [A] (verification not implemented)	3907
3.407.8 Giac [F(-1)]	3908
3.407.9 Mupad [B] (verification not implemented)	3908

3.407.1 Optimal result

Integrand size = 33, antiderivative size = 391

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx \\ &= \frac{(2ab(A - B) - a^2(A + B) + b^2(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^2 d} \\ & \quad - \frac{(2ab(A - B) - a^2(A + B) + b^2(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^2 d} \\ & \quad + \frac{\sqrt{b}(5a^2Ab + Ab^3 - 3a^3B + ab^2B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{a^{3/2}(a^2 + b^2)^2 d} \\ & \quad - \frac{(a^2(A - B) - b^2(A - B) + 2ab(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^2 d} \\ & \quad + \frac{(a^2(A - B) - b^2(A - B) + 2ab(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^2 d} \\ & \quad + \frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{a(a^2 + b^2)d(a + b \tan(c + dx))} \end{aligned}$$

output
$$\begin{aligned} & -1/2*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}) \\ & / (a^2+b^2)^2/d*2^{(1/2)}-1/2*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\arctan(1+2^{(1/2)} \\ & *\tan(d*x+c)^{(1/2)})/(a^2+b^2)^2/d*2^{(1/2)}-1/4*(a^2*(A-B)-b^2*(A-B)+2*a*b \\ & *(A+B))*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(a^2+b^2)^2/d*2^{(1/2)}+1/ \\ & 4*(a^2*(A-B)-b^2*(A-B)+2*a*b*(A+B))*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+ \\ & c))/(a^2+b^2)^2/d*2^{(1/2)}+(5*A*a^2*b+A*b^3-3*B*a^3+B*a*b^2)*\arctan(b^{(1/2)} \\ & *\tan(d*x+c)^{(1/2)}/a^{(1/2)})*b^{(1/2)}/a^{(3/2)}/(a^2+b^2)^2/d+b*(A*b-B*a)*\tan(d \\ & *x+c)^{(1/2)}/a/(a^2+b^2)/d/(a+b*\tan(d*x+c)) \end{aligned}$$

3.407.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.22 (sec) , antiderivative size = 204, normalized size of antiderivative = 0.52

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx$$

$$= \frac{\sqrt{b}(5a^2Ab + Ab^3 - 3a^3B + ab^2B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{a}(a^2+b^2)} + \frac{\sqrt[4]{-1}(-a(a+ib)^2(A-ib) \arctan((-1)^{3/4}\sqrt{\tan(c+dx)}) - a(a-ib)^2(A+ib) \arctan((-1)^{3/4}\sqrt{\tan(c+dx)})}{a^2+b^2}}{a(a^2 + b^2)d}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^2), x]`

output
$$\begin{aligned} & ((\text{Sqrt}[b]*(5*a^2*A*b + A*b^3 - 3*a^3*B + a*b^2*B))*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan} \\ & [c + d*x]])/\text{Sqrt}[a]])/(\text{Sqrt}[a]*(a^2 + b^2)) + ((-1)^{(1/4)}*(-(a*(a + I*b)^2 \\ & *(A - I*B))*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]]) - a*(a - I*b)^2*(A + I*B) \\ &)*\text{ArcTanh}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]])/(a^2 + b^2) + (b*(A*b - a*B)*\text{Sqrt} \\ & [\text{Tan}[c + d*x]])/(a + b*\text{Tan}[c + d*x])/(a*(a^2 + b^2)*d) \end{aligned}$$

3.407.3 Rubi [A] (verified)

Time = 1.40 (sec) , antiderivative size = 335, normalized size of antiderivative = 0.86, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 4092, 27, 3042, 4136, 27, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.407. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^2} dx$

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx$$

↓ 4092

$$\frac{\int \frac{2Aa^2 + bBa - 2(Ab - aB) \tan(c + dx)a + Ab^2 + b(Ab - aB) \tan^2(c + dx)}{2\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{a(a^2 + b^2)} + \frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 27

$$\frac{\int \frac{2Aa^2 + bBa - 2(Ab - aB) \tan(c + dx)a + Ab^2 + b(Ab - aB) \tan^2(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{2a(a^2 + b^2)} + \frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 3042

$$\frac{\int \frac{2Aa^2 + bBa - 2(Ab - aB) \tan(c + dx)a + Ab^2 + b(Ab - aB) \tan(c + dx)^2}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{2a(a^2 + b^2)} + \frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 4136

$$\frac{\int \frac{2(a(Aa^2 + 2bBa - Ab^2) - a(-Ba^2 + 2Aba + b^2B) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx}{a^2 + b^2} + \frac{b(-3a^3B + 5a^2Ab + ab^2B + Ab^3) \int \frac{\tan^2(c + dx) + 1}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{a^2 + b^2}}{2a(a^2 + b^2)} + \frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 27

$$\frac{2 \int \frac{a(Aa^2 + 2bBa - Ab^2) - a(-Ba^2 + 2Aba + b^2B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx}{a^2 + b^2} + \frac{b(-3a^3B + 5a^2Ab + ab^2B + Ab^3) \int \frac{\tan^2(c + dx) + 1}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{a^2 + b^2}}{2a(a^2 + b^2)} + \frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 3042

$$\frac{2 \int \frac{a(Aa^2 + 2bBa - Ab^2) - a(-Ba^2 + 2Aba + b^2B) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx}{a^2 + b^2} + \frac{b(-3a^3B + 5a^2Ab + ab^2B + Ab^3) \int \frac{\tan(c + dx)^2 + 1}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{a^2 + b^2}}{2a(a^2 + b^2)} + \frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 4017

3.407. $\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx$

$$\frac{4 \int \frac{a(Aa^2+2bBa-Ab^2 - (-Ba^2+2Aba+b^2B) \tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{b(-3a^3B+5a^2Ab+ab^2B+Ab^3)}{a^2+b^2} \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx}{d(a^2+b^2)} + \frac{2a(a^2+b^2)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

↓ 27

$$\frac{4a \int \frac{Aa^2+2bBa-Ab^2 - (-Ba^2+2Aba+b^2B) \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{b(-3a^3B+5a^2Ab+ab^2B+Ab^3)}{a^2+b^2} \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx}{d(a^2+b^2)} + \frac{2a(a^2+b^2)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

↓ 1482

$$\frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d(a^2+b^2)} + \frac{2a(a^2+b^2)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

↓ 1476

$$\frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}} \right) \right)}{d(a^2+b^2)} + \frac{2a(a^2+b^2)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

↓ 1082

$$\frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)} + \frac{2a(a^2+b^2)}{ad(a^2+b^2)(a+b \tan(c+dx))} + \frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

↓ 217

3.407. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))^2}} dx$

$$4a \left(\frac{\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}) \right) \right)}{d(a^2+b^2)}$$

$2a(a^2+b^2)$

$$\frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

↓ 1479

$$4a \left(\frac{\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}) \right)}{d(a^2+b^2)} \right)$$

$$\frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

↓ 25

$$4a \left(\frac{\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}) \right)}{d(a^2+b^2)} \right)$$

$$\frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

↓ 27

$$4a \left(\frac{\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}) \right)}{d(a^2+b^2)} \right)$$

$$\frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

↓ 1103

$$\frac{b(-3a^3B+5a^2Ab+ab^2B+Ab^3) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2} + \frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} + \arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}) \right) - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}) \right) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

↓ 4117

3.407. $\int \frac{A+B\tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2} dx$

$$\frac{b(-3a^3B+5a^2Ab+ab^2B+Ab^3) \int \frac{1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} d \tan(c+dx)}{d(a^2+b^2)} + \frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}{2\sqrt{2}} \right) \right)}{d(a^2+b^2)}$$

2a(a^2

$$\frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)(a + b \tan(c + dx))}$$

73

$$\frac{2b(-3a^3B+5a^2Ab+ab^2B+Ab^3) \int \frac{1}{a+b \tan(c+dx)} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} + \frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}{2\sqrt{2}} \right) \right)}{d(a^2+b^2)}$$

2a(a^2 + b

$$\frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)(a + b \tan(c + dx))}$$

218

$$\frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)(a + b \tan(c + dx))} + \frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}}{2\sqrt{2}} \right) \right) - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A-B))}{d(a^2+b^2)}$$

2a(a^2 + b^2)

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^2),x]`

output `((2*Sqrt[b]*(5*a^2*A*b + A*b^3 - 3*a^3*B + a*b^2*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[a]*(a^2 + b^2)*d) + (4*a*(-1/2*((2*a*b*(A - B) - a^2*(A + B) + b^2*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])) + ((a^2*(A - B) - b^2*(A - B) + 2*a*b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]))) / ((a^2 + b^2)*d)) / (2*a*(a^2 + b^2)) + (b*(A*b - a*B)*Sqrt[Tan[c + d*x]]) / (a*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))`

3.407. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))^2}} dx$

3.407.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`


```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.407.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 336, normalized size of antiderivative = 0.86

method	result
derivativedivides	$\frac{2b \left(\frac{(A a^2 b + A b^3 - B a^3 - B a b^2) (\sqrt{\tan(dx+c)})}{2a(a+b \tan(dx+c))} + \frac{(5A a^2 b + A b^3 - 3B a^3 + B a b^2) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{2a\sqrt{ab}} \right)}{(a^2+b^2)^2} + \frac{(A a^2 - A b^2 + 2B a b)}{(a^2+b^2)^2}$
default	$\frac{2b \left(\frac{(A a^2 b + A b^3 - B a^3 - B a b^2) (\sqrt{\tan(dx+c)})}{2a(a+b \tan(dx+c))} + \frac{(5A a^2 b + A b^3 - 3B a^3 + B a b^2) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{2a\sqrt{ab}} \right)}{(a^2+b^2)^2} + \frac{(A a^2 - A b^2 + 2B a b)}{(a^2+b^2)^2}$

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^2,x,method=_RETURNV
ERBOSE)
```

```
output 1/d*(2*b/(a^2+b^2)^2*(1/2*(A*a^2*b+A*b^3-B*a^3-B*a*b^2)/a*tan(d*x+c)^(1/2)
/(a+b*tan(d*x+c))+1/2*(5*A*a^2*b+A*b^3-3*B*a^3+B*a*b^2)/a/(a*b)^(1/2)*arct
an(b*tan(d*x+c)^(1/2)/(a*b)^(1/2)))+2/(a^2+b^2)^2*(1/8*(A*a^2-A*b^2+2*B*a*
b)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+
c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(
1/2)*tan(d*x+c)^(1/2)))+1/8*(-2*A*a*b+B*a^2-B*b^2)*2^(1/2)*(ln((1-2^(1/2)*
tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*ar
ctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))
```

$$3.407. \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^2} dx$$

3.407.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5903 vs. 2(355) = 710.

Time = 25.29 (sec) , antiderivative size = 11831, normalized size of antiderivative = 30.26

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

output Too large to include

3.407.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^2 \sqrt{\tan(c + dx)}} dx$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+b*tan(d*x+c))**2,x)`

output `Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))**2*sqrt(tan(c + d*x))), x)`

3.407.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 356, normalized size of antiderivative = 0.91

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx = \frac{4(3Ba^3b - 5Aa^2b^2 - Bab^3 - Ab^4) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^5 + 2a^3b^2 + ab^4)\sqrt{ab}} + \frac{4(Bab - Ab^2)\sqrt{\tan(dx+c)}}{a^4 + a^2b^2 + (a^3b + ab^3)\tan(dx+c)} - \frac{2\sqrt{2}((A+B)a^2 - 2(A-B)ab - (A+B)b^2)}{(a^5 + 2a^3b^2 + ab^4)\sqrt{ab}}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

3.407. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^2} dx$

output
$$\frac{-1/4*(4*(3*B*a^3*b - 5*A*a^2*b^2 - B*a*b^3 - A*b^4)*\arctan(b*\sqrt{\tan(dx + c)})/\sqrt{a*b})/((a^5 + 2*a^3*b^2 + a*b^4)*\sqrt{a*b}) + 4*(B*a*b - A*b^2)*\sqrt{\tan(dx + c)}/(a^4 + a^2*b^2 + (a^3*b + a*b^3)*\tan(dx + c)) - (2*\sqrt{2}*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2*\sqrt{\tan(dx + c)}))) + 2*\sqrt{2}*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2*\sqrt{\tan(dx + c)}))) + \sqrt{2}*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*\log(\sqrt{2}*\sqrt{\tan(dx + c)} + \tan(dx + c) + 1) - \sqrt{2}*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*\log(-\sqrt{2}*\sqrt{\tan(dx + c)} + \tan(dx + c) + 1))/(a^4 + 2*a^2*b^2 + b^4))/d$$

3.407.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output Timed out

3.407.9 Mupad [B] (verification not implemented)

Time = 39.00 (sec) , antiderivative size = 17494, normalized size of antiderivative = 44.74

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + b*tan(c + d*x))^2),x)`

output

$$\begin{aligned}
& (\log(- (((((((((256*B*b^3*(2*a^4 - b^4 + a^2*b^2))/d - 128*b^3*\tan(c + d*x) \\
&)^{(1/2)}*(a^2 - b^2)*(a^2 + b^2)^2*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2) \\
&)^{(1/2)} + 16*B^2*a*b^3*d^2 - 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^{(1/2)})* \\
& (((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^{1/2} + 16*B^2*a*b^3*d^2 - 16*B^2 \\
& *a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^{(1/2)})/4 - (64*B^2*a*b^2*\tan(c + d*x)^{(1/2)} \\
& *(a^6 + 17*b^6 - 29*a^2*b^4 + 19*a^4*b^2))/(d^2*(a^2 + b^2)^2))*((4*(-B^4 \\
& *d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^{1/2} + 16*B^2*a*b^3*d^2 - 16*B^2*a^3*b*d \\
& ^2)/(d^4*(a^2 + b^2)^4))^{(1/2)})/4 + (32*B^3*a*b^2*(a^6 + 13*b^6 - 45*a^2*b \\
& ^4 + 39*a^4*b^2))/(d^3*(a^2 + b^2)^3))*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b \\
& ^2)^2)^{1/2} + 16*B^2*a*b^3*d^2 - 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^{(1 \\
& /2)}/4 - (16*B^4*b^3*\tan(c + d*x)^{(1/2)}*(9*a^6 - 3*b^6 + 3*a^2*b^4 - 17*a^ \\
& 4*b^2))/(d^4*(a^2 + b^2)^4))*(((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^{1/2} \\
&) + 16*B^2*a*b^3*d^2 - 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^{(1/2)})/4 - (\\
& 8*B^5*b^3*(9*a^4 - b^4))/(d^5*(a^2 + b^2)^4))*(((192*B^4*a^2*b^6*d^4 - 16* \\
& B^4*b^8*d^4 - 16*B^4*a^8*d^4 - 608*B^4*a^4*b^4*d^4 + 192*B^4*a^6*b^2*d^4)^{ \\
& (1/2)} + 16*B^2*a*b^3*d^2 - 16*B^2*a^3*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^ \\
& 6*d^4 + 6*a^4*b^4*d^4 + 4*a^6*b^2*d^4))^{(1/2)})/4 + (\log(- (((((((((256*B*b \\
& ^3*(2*a^4 - b^4 + a^2*b^2))/d - 128*b^3*\tan(c + d*x)^{(1/2)}*(a^2 - b^2)*(a^ \\
& 2 + b^2)^2*(-(4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^{1/2} - 16*B^2*a*b^3* \\
& d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^{(1/2)})*(-(4*(-B^4*d^4*(a^4...
\end{aligned}$$

3.408
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$$

3.408.1 Optimal result 3910
 3.408.2 Mathematica [C] (verified) 3911
 3.408.3 Rubi [A] (verified) 3912
 3.408.4 Maple [A] (verified) 3920
 3.408.5 Fricas [B] (verification not implemented) 3920
 3.408.6 Sympy [F(-1)] 3921
 3.408.7 Maxima [A] (verification not implemented) 3921
 3.408.8 Giac [F(-1)] 3922
 3.408.9 Mupad [B] (verification not implemented) 3922

3.408.1 Optimal result

Integrand size = 33, antiderivative size = 439

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx \\ &= \frac{(a^2(A - B) - b^2(A - B) + 2ab(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^2 d} \\ & \quad - \frac{(a^2(A - B) - b^2(A - B) + 2ab(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^2 d} \\ & \quad - \frac{b^{3/2}(7a^2 Ab + 3Ab^3 - 5a^3 B - ab^2 B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{a^{5/2}(a^2 + b^2)^2 d} \\ & \quad + \frac{(2ab(A - B) - a^2(A + B) + b^2(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^2 d} \\ & \quad - \frac{(2ab(A - B) - a^2(A + B) + b^2(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^2 d} \\ & \quad - \frac{2a^2 A + 3Ab^2 - abB}{a^2(a^2 + b^2)d\sqrt{\tan(c + dx)}} + \frac{b(Ab - aB)}{a(a^2 + b^2)d\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} \end{aligned}$$

output $-b^{3/2}*(7*A*a^2*b+3*A*b^3-5*B*a^3-B*a*b^2)*\arctan(b^{1/2}*\tan(dx+c)^{1/2})/a^{5/2}/(a^2+b^2)^2/d-1/2*(a^2*(A-B)-b^2*(A-B)+2*a*b*(A+B))*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2})/(a^2+b^2)^2/d*2^{1/2}-1/2*(a^2*(A-B)-b^2*(A-B)+2*a*b*(A+B))*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2})/(a^2+b^2)^2/d*2^{1/2}+1/4*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\ln(1-2^{1/2}*\tan(dx+c)^{1/2})+\tan(dx+c)/(a^2+b^2)^2/d*2^{1/2}-1/4*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\ln(1+2^{1/2}*\tan(dx+c)^{1/2})+\tan(dx+c)/(a^2+b^2)^2/d*2^{1/2}+(-2*A*a^2-3*A*b^2+B*a*b)/a^2/(a^2+b^2)/d/\tan(dx+c)^{1/2}+b*(A*b-B*a)/a/(a^2+b^2)/d/\tan(dx+c)^{1/2}/(a+b*\tan(dx+c))$

3.408.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.51 (sec) , antiderivative size = 239, normalized size of antiderivative = 0.54

$$\int \frac{A + B \tan(c + dx)}{\tan^{3/2}(c + dx)(a + b \tan(c + dx))^2} dx$$

$$= \frac{b^{3/2}(-7a^2Ab-3Ab^3+5a^3B+ab^2B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right) + \sqrt[4]{-1}a(-i(a+ib)^2(A-iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right) + i(a-ib)^2(A+iB)\delta)}{a^{3/2}(a^2+b^2)} + \frac{\sqrt[4]{-1}a(-i(a+ib)^2(A-iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right) + i(a-ib)^2(A+iB)\delta)}{a^2+b^2}$$

$$a(a^2 + b^2) d$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^2),x]`

output $((b^{3/2}*(-7*a^2*A*b - 3*A*b^3 + 5*a^3*B + a*b^2*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(a^{3/2}*(a^2 + b^2)) + ((-1)^{1/4}*a*((-I)*(a + I*b)^2*(A - I*B)*\text{ArcTan}[(-1)^{3/4}*\text{Sqrt}[\text{Tan}[c + d*x]]] + I*(a - I*b)^2*(A + I*B)*\text{ArcTanh}[(-1)^{3/4}*\text{Sqrt}[\text{Tan}[c + d*x]]]))/(a^2 + b^2) + (-2*a^2*A - 3*A*b^2 + a*b*B)/(a*\text{Sqrt}[\text{Tan}[c + d*x]]) + (b*(A*b - a*B))/(\text{Sqrt}[\text{Tan}[c + d*x]])*(a + b*\text{Tan}[c + d*x]))/(a*(a^2 + b^2)*d)$

3.408.3 Rubi [A] (verified)

Time = 1.91 (sec) , antiderivative size = 381, normalized size of antiderivative = 0.87, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.697$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2}(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{4092} \\
 & \frac{\int \frac{2Aa^2 - bBa - 2(Ab - aB) \tan(c + dx)a + 3Ab^2 + 3b(Ab - aB) \tan^2(c + dx)}{2 \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx}{\frac{a(a^2 + b^2)}{b(Ab - aB)}} + \\
 & \quad \frac{ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))}{ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{2Aa^2 - bBa - 2(Ab - aB) \tan(c + dx)a + 3Ab^2 + 3b(Ab - aB) \tan^2(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx}{\frac{2a(a^2 + b^2)}{b(Ab - aB)}} + \\
 & \quad \frac{ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))}{ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{2Aa^2 - bBa - 2(Ab - aB) \tan(c + dx)a + 3Ab^2 + 3b(Ab - aB) \tan(c + dx)^2}{\tan(c + dx)^{3/2}(a + b \tan(c + dx))} dx}{\frac{2a(a^2 + b^2)}{b(Ab - aB)}} + \\
 & \quad \frac{ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))}{ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))} \\
 & \quad \downarrow \text{4132} \\
 & \frac{2 \int \frac{-2Ba^3 + 4Aba^2 + 2(aA + bB) \tan(c + dx)a^2 - b^2Ba + 3Ab^3 + b(2Aa^2 - bBa + 3Ab^2) \tan^2(c + dx)}{2 \sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{\frac{2a(a^2 + b^2)}{b(Ab - aB)}} - \frac{2(2a^2A - abB + 3Ab^2)}{ad \sqrt{\tan(c + dx)}} + \\
 & \quad \frac{ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))}{ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))}
 \end{aligned}$$

3.408. $\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$

$$\begin{aligned}
 & \int \frac{-2Ba^3+4Aba^2+2(aA+bB)\tan(c+dx)a^2-b^2Ba+3Ab^3+b(2Aa^2-bBa+3Ab^2)\tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx - \frac{2(2a^2A-abB+3Ab^2)}{ad\sqrt{\tan(c+dx)}} + \\
 & \frac{2a(a^2+b^2)}{b(Ab-aB)} \\
 & \frac{ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}{\phantom{ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}} \\
 & \downarrow 27 \\
 & \int \frac{-2Ba^3+4Aba^2+2(aA+bB)\tan(c+dx)a^2-b^2Ba+3Ab^3+b(2Aa^2-bBa+3Ab^2)\tan(c+dx)^2}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx - \frac{2(2a^2A-abB+3Ab^2)}{ad\sqrt{\tan(c+dx)}} + \\
 & \frac{2a(a^2+b^2)}{b(Ab-aB)} \\
 & \frac{ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}{\phantom{ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}} \\
 & \downarrow 3042 \\
 & \int \frac{2\left(\frac{(-Ba^2+2Aba+b^2B)a^2+(Aa^2+2bBa-Ab^2)\tan(c+dx)a^2}{\sqrt{\tan(c+dx)}}\right) dx}{a^2+b^2} + \frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3)\int\frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2} - \frac{2(2a^2A-abB+3Ab^2)}{ad\sqrt{\tan(c+dx)}} + \\
 & \frac{2a(a^2+b^2)}{b(Ab-aB)} \\
 & \frac{ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}{\phantom{ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}} \\
 & \downarrow 4136 \\
 & \int \frac{(-Ba^2+2Aba+b^2B)a^2+(Aa^2+2bBa-Ab^2)\tan(c+dx)a^2}{\sqrt{\tan(c+dx)}} dx + \frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3)\int\frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2} - \frac{2(2a^2A-abB+3Ab^2)}{ad\sqrt{\tan(c+dx)}} + \\
 & \frac{2a(a^2+b^2)}{b(Ab-aB)} \\
 & \frac{ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}{\phantom{ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}} \\
 & \downarrow 27 \\
 & 2\int\frac{(-Ba^2+2Aba+b^2B)a^2+(Aa^2+2bBa-Ab^2)\tan(c+dx)a^2}{\sqrt{\tan(c+dx)}} dx + \frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3)\int\frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2} - \frac{2(2a^2A-abB+3Ab^2)}{ad\sqrt{\tan(c+dx)}} + \\
 & \frac{2a(a^2+b^2)}{b(Ab-aB)} \\
 & \frac{ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}{\phantom{ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}} \\
 & \downarrow 3042 \\
 & 2\int\frac{(-Ba^2+2Aba+b^2B)a^2+(Aa^2+2bBa-Ab^2)\tan(c+dx)a^2}{\sqrt{\tan(c+dx)}} dx + \frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3)\int\frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2} - \frac{2(2a^2A-abB+3Ab^2)}{ad\sqrt{\tan(c+dx)}} + \\
 & \frac{2a(a^2+b^2)}{b(Ab-aB)} \\
 & \frac{ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}{\phantom{ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}} \\
 & \downarrow 4017
 \end{aligned}$$

3.408. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^2} dx$

$$\frac{4 \int \frac{a^2(-Ba^2+2Aba+b^2B+(Aa^2+2bBa-Ab^2)\tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3)}{a^2+b^2} \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a} - \frac{2(2a^2A-ab^2)}{ad\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}$$

$$\frac{b(Ab-aB)}{ad(a^2+b^2)\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}$$

↓ 27

$$\frac{4a^2 \int \frac{-Ba^2+2Aba+b^2B+(Aa^2+2bBa-Ab^2)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3)}{a^2+b^2} \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a} - \frac{2(2a^2A-ab^2)}{ad\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}$$

$$\frac{b(Ab-aB)}{ad(a^2+b^2)\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}$$

↓ 1482

$$\frac{4a^2\left(\frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B)\right) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{a} + \frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3)}{a^2+b^2} \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx - \frac{2(2a^2A-ab^2)}{ad\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}$$

$$\frac{b(Ab-aB)}{ad(a^2+b^2)\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}$$

↓ 1476

$$\frac{4a^2\left(\frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B)\right) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}\right)}{a} + \frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3)}{a^2+b^2} \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx - \frac{2(2a^2A-ab^2)}{ad\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}$$

$$\frac{b(Ab-aB)}{ad(a^2+b^2)\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}$$

↓ 1082

$$\frac{4a^2\left(\frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B)\right) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}}\right)}{a} + \frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3)}{a^2+b^2} \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx - \frac{2(2a^2A-ab^2)}{ad\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}$$

$$\frac{b(Ab-aB)}{ad(a^2+b^2)\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}$$

↓ 217

3.408. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^2} dx$

$$4a^2 \left(\frac{1}{2} (-a^2(A+B) + 2ab(A-B) + b^2(A+B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx) + 1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \arctan(1) \right) \right) \frac{1}{d(a^2 + b^2)}$$

a

$$2a(a^2 + b^2)$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \sqrt{\tan(c+dx)}(a + b \tan(c+dx))}$$

↓ 1479

$$4a^2 \left(\frac{1}{2} (-a^2(A+B) + 2ab(A-B) + b^2(A+B)) \left(- \int \frac{\sqrt{2} - 2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \arctan(1) \right) \right) \frac{1}{d(a^2 + b^2)}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \sqrt{\tan(c+dx)}(a + b \tan(c+dx))}$$

↓ 25

$$4a^2 \left(\frac{1}{2} (-a^2(A+B) + 2ab(A-B) + b^2(A+B)) \left(\int \frac{\sqrt{2} - 2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \arctan(1) \right) \right) \frac{1}{d(a^2 + b^2)}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \sqrt{\tan(c+dx)}(a + b \tan(c+dx))}$$

↓ 27

$$4a^2 \left(\frac{1}{2} (-a^2(A+B) + 2ab(A-B) + b^2(A+B)) \left(\int \frac{\sqrt{2} - 2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)} + 1}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \arctan(1) \right) \right) \frac{1}{d(a^2 + b^2)}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \sqrt{\tan(c+dx)}(a + b \tan(c+dx))}$$

↓ 1103

3.408. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

$$\frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3)}{a^2+b^2} \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx + \frac{4a^2\left(\frac{1}{2}(a^2(A-B)+2ab(A+B))-b^2(A-B)\right)\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}}\right)}{a}$$

$$\frac{b(Ab-aB)}{ad(a^2+b^2)\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}$$

↓ 4117

$$\frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3)}{d(a^2+b^2)} \int \frac{1}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} d\tan(c+dx) + \frac{4a^2\left(\frac{1}{2}(a^2(A-B)+2ab(A+B))-b^2(A-B)\right)\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}}\right)}{a}$$

$$\frac{b(Ab-aB)}{ad(a^2+b^2)\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}$$

↓ 73

$$\frac{2b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3)}{d(a^2+b^2)} \int \frac{1}{a+b\tan(c+dx)} d\sqrt{\tan(c+dx)} + \frac{4a^2\left(\frac{1}{2}(a^2(A-B)+2ab(A+B))-b^2(A-B)\right)\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}}\right)}{a}$$

$$\frac{b(Ab-aB)}{ad(a^2+b^2)\sqrt{\tan(c+dx)(a+b\tan(c+dx))}}$$

↓ 218

$$\frac{b(Ab-aB)}{ad(a^2+b^2)\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} + \frac{4a^2\left(\frac{1}{2}(a^2(A-B)+2ab(A+B))-b^2(A-B)\right)\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}}\right) + \frac{1}{2}(-a^2(A+B)+2ab(A-B))}{d(a^2+b^2)}$$

$$\frac{2(2a^2A-abB+3Ab^2)}{ad\sqrt{\tan(c+dx)}}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^2),x]`

3.408. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^2} dx$

```
output (-(((2*b^(3/2)*(7*a^2*A*b + 3*A*b^3 - 5*a^3*B - a*b^2*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]/(Sqrt[a]*(a^2 + b^2)*d) + (4*a^2*((a^2*(A - B) - b^2*(A - B) + 2*a*b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]))/2 + ((2*a*b*(A - B) - a^2*(A + B) + b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)/a) - (2*(2*a^2*A + 3*A*b^2 - a*b*B)/(a*d*Sqrt[Tan[c + d*x]]))/(2*a*(a^2 + b^2)) + (b*(A*b - a*B))/(a*(a^2 + b^2)*d*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x]))
```

3.408.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]
```

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.408.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 352, normalized size of antiderivative = 0.80

method	result
derivativedivides	$2b^2 \left(\frac{\left(\frac{1}{2}Aa^2b + \frac{1}{2}Ab^3 - \frac{1}{2}Ba^3 - \frac{1}{2}Bab^2\right)(\sqrt{\tan(dx+c)}}{a+b\tan(dx+c)} + \frac{(7Aa^2b+3Ab^3-5Ba^3-Bab^2)\arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{2\sqrt{ab}} \right)}{a^2(a^2+b^2)^2} - \frac{2A}{a^2\sqrt{\tan(dx+c)}}$
default	$2b^2 \left(\frac{\left(\frac{1}{2}Aa^2b + \frac{1}{2}Ab^3 - \frac{1}{2}Ba^3 - \frac{1}{2}Bab^2\right)(\sqrt{\tan(dx+c)}}{a+b\tan(dx+c)} + \frac{(7Aa^2b+3Ab^3-5Ba^3-Bab^2)\arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{2\sqrt{ab}} \right)}{a^2(a^2+b^2)^2} - \frac{2A}{a^2\sqrt{\tan(dx+c)}}$

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^2,x,method=_RETURNV
ERBOSE)`

output `1/d*(-2*b^2/a^2/(a^2+b^2)^2*((1/2*A*a^2*b+1/2*A*b^3-1/2*B*a^3-1/2*B*a*b^2)
*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))+1/2*(7*A*a^2*b+3*A*b^3-5*B*a^3-B*a*b^2)
/(a*b)^(1/2)*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2)))-2/a^2*A/tan(d*x+c)^(1
/2)+2/(a^2+b^2)^2*(1/8*(-2*A*a*b+B*a^2-B*b^2)*2^(1/2)*(ln((1+2^(1/2))*tan(d
*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(
1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/8*(-A
*a^2+A*b^2-2*B*a*b)*2^(1/2)*(ln((1-2^(1/2))*tan(d*x+c)^(1/2)+tan(d*x+c))/(1
+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2)
)+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))`

3.408.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5991 vs. 2(396) = 792.

Time = 39.86 (sec) , antiderivative size = 12008, normalized size of antiderivative = 27.35

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^2,x, algorith
m="fracas")`

output `Too large to include`

3.408.
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$$

3.408.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Timed out}$$

```
input integrate((A+B*tan(d*x+c))/tan(d*x+c)**(3/2)/(a+b*tan(d*x+c))**2,x)
```

```
output Timed out
```

3.408.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 396, normalized size of antiderivative = 0.90

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$$

$$\frac{4(5Ba^3b^2 - 7Aa^2b^3 + Bab^4 - 3Ab^5) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right) - 2\sqrt{2}((A-B)a^2 + 2(A+B)ab - (A-B)b^2) \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2} + 2\sqrt{\tan(dx+c)})\right) + 2\sqrt{2}((A-B)a^2 + 2(A+B)ab - (A-B)b^2) \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2} - 2\sqrt{\tan(dx+c)})\right)}{(a^6 + 2a^4b^2 + a^2b^4)\sqrt{ab}}$$

```
input integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^2,x, algorithm
m="maxima")
```

```
output 1/4*(4*(5*B*a^3*b^2 - 7*A*a^2*b^3 + B*a*b^4 - 3*A*b^5)*arctan(b*sqrt(tan(d
*x + c))/sqrt(a*b))/((a^6 + 2*a^4*b^2 + a^2*b^4)*sqrt(a*b)) - (2*sqrt(2)*
(A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2
*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b
^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*((A +
B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan
(d*x + c) + 1) + sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*log(-
sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))/(a^4 + 2*a^2*b^2 + b^4) -
4*(2*A*a^3 + 2*A*a*b^2 + (2*A*a^2*b - B*a*b^2 + 3*A*b^3)*tan(d*x + c))/((a
^4*b + a^2*b^3)*tan(d*x + c)^(3/2) + (a^5 + a^3*b^2)*sqrt(tan(d*x + c)))/
d
```


3.408.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output `Timed out`

3.408.9 Mupad [B] (verification not implemented)

Time = 29.80 (sec) , antiderivative size = 22667, normalized size of antiderivative = 51.63

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + b*tan(c + d*x))^2),x)`

output `(log((((((((((128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b^2)^2*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2) + (128*B*b^2*(2*b^6 - a^6 + 9*a^2*b^4 + 6*a^4*b^2))/(a*d))*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (64*B^2*b^2*tan(c + d*x)^(1/2)*(2*b^8 - a^8 + 5*a^2*b^6 + 67*a^4*b^4 - a^6*b^2))/(a*d^2*(a^2 + b^2)^2))*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 - (32*B^3*b^5*(25*a^6 + b^6 - 13*a^2*b^4 - 85*a^4*b^2))/(a^2*d^3*(a^2 + b^2)^3))*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (16*B^4*b^5*tan(c + d*x)^(1/2)*(b^6 - 27*a^6 + 7*a^2*b^4 + 11*a^4*b^2))/(a^2*d^4*(a^2 + b^2)^4))*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (16*B^5*b^6*(5*a^2 + b^2))/(a*d^5*(a^2 + b^2)^4))*((((192*B^4*a^2*b^6*d^4 - 16*B^4*b^8*d^4 - 16*B^4*a^8*d^4 - 608*B^4*a^4*b^4*d^4 + 192*B^4*a^6*b^2*d^4)^(1/2) - 16*B^2*a*b^3*d^2 + 16*B^2*a^3*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6*d^4 + 6*a^4*b^4*d^4 + 4*a^6*b^2*d^4))^(1/2))/4 + (log((((((((((128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b^2)^2*((4*(-B^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16*B^2*a*b^3*d^2 - 16*B^2*a^3*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2) + (128*B*b^2*(2*b^6 - a^6 + 9*...`

3.409 $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

3.409.1 Optimal result 3923
 3.409.2 Mathematica [C] (verified) 3924
 3.409.3 Rubi [A] (verified) 3925
 3.409.4 Maple [A] (verified) 3934
 3.409.5 Fricas [B] (verification not implemented) 3934
 3.409.6 Sympy [F(-1)] 3935
 3.409.7 Maxima [A] (verification not implemented) 3935
 3.409.8 Giac [F(-1)] 3936
 3.409.9 Mupad [B] (verification not implemented) 3936

3.409.1 Optimal result

Integrand size = 33, antiderivative size = 493

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$$

$$= -\frac{(2ab(A - B) - a^2(A + B) + b^2(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^2 d}$$

$$+ \frac{(2ab(A - B) - a^2(A + B) + b^2(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^2 d}$$

$$+ \frac{b^{5/2}(9a^2 Ab + 5Ab^3 - 7a^3 B - 3ab^2 B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{a^{7/2}(a^2 + b^2)^2 d}$$

$$+ \frac{(a^2(A - B) - b^2(A - B) + 2ab(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^2 d}$$

$$- \frac{(a^2(A - B) - b^2(A - B) + 2ab(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^2 d}$$

$$- \frac{2a^2 A + 5Ab^2 - 3abB}{3a^2(a^2 + b^2)d \tan^{\frac{3}{2}}(c + dx)} + \frac{4a^2 Ab + 5Ab^3 - 2a^3 B - 3ab^2 B}{a^3(a^2 + b^2)d \sqrt{\tan(c + dx)}}$$

$$+ \frac{b(Ab - aB)}{a(a^2 + b^2)d \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}$$

output $b^{5/2} \cdot (9Aa^2b + 5Ab^3 - 7B^2a^3 - 3B^2ab^2) \cdot \arctan(b^{1/2} \tan(dx+c)^{1/2} / a^{1/2}) / a^{7/2} / (a^2+b^2)^{2/d} + 1/2 \cdot (2ab(A-B) - a^2(A+B) + b^2(A+B)) \cdot \arctan(-1+2^{1/2} \tan(dx+c)^{1/2}) / (a^2+b^2)^{2/d} + 1/2 \cdot (2ab(A-B) - a^2(A+B) + b^2(A+B)) \cdot \arctan(1+2^{1/2} \tan(dx+c)^{1/2}) / (a^2+b^2)^{2/d} + 1/4 \cdot (a^2(A-B) - b^2(A-B) + 2ab(A+B)) \cdot \ln(1-2^{1/2} \tan(dx+c)^{1/2} + \tan(dx+c)) / (a^2+b^2)^{2/d} - 1/4 \cdot (a^2(A-B) - b^2(A-B) + 2ab(A+B)) \cdot \ln(1+2^{1/2} \tan(dx+c)^{1/2} + \tan(dx+c)) / (a^2+b^2)^{2/d} + (4Aa^2b + 5Ab^3 - 2B^2a^3 - 3B^2ab^2) / a^3 / (a^2+b^2) / d / \tan(dx+c)^{1/2} + 1/3 \cdot (-2Aa^2 - 5Ab^2 + 3B^2ab) / a^2 / (a^2+b^2) / d / \tan(dx+c)^{3/2} + b \cdot (Ab - Ba) / a / (a^2+b^2) / d / \tan(dx+c)^{3/2} / (a+b \tan(dx+c))$

3.409.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.19 (sec) , antiderivative size = 287, normalized size of antiderivative = 0.58

$$\int \frac{A + B \tan(c + dx)}{\tan^{5/2}(c + dx)(a + b \tan(c + dx))^2} dx$$

$$= \frac{3 \left(\sqrt[4]{-1} a^{7/2} (a+ib)^2 (A-iB) \arctan\left((-1)^{3/4} \sqrt{\tan(c+dx)}\right) + b^{5/2} (9a^2 Ab + 5Ab^3 - 7a^3 B - 3ab^2 B) \arctan\left(\frac{\sqrt{b} \sqrt{\tan(c+dx)}}{\sqrt{a}}\right) + \sqrt[4]{-1} a^{7/2} (a-ib)^2 \right)}{a^{5/2} (a^2 + b^2)} + 3a(a^2 + b^2)$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^2),x]`

output $((3 \cdot (-1)^{1/4} \cdot a^{7/2} \cdot (a + I \cdot b)^2 \cdot (A - I \cdot B) \cdot \text{ArcTan}[(-1)^{3/4} \cdot \text{Sqrt}[\text{Tan}[c + d \cdot x]]] + b^{5/2} \cdot (9 \cdot a^2 \cdot A \cdot b + 5 \cdot A \cdot b^3 - 7 \cdot a^3 \cdot B - 3 \cdot a \cdot b^2 \cdot B) \cdot \text{ArcTan}[(\text{Sqrt}[b] \cdot \text{Sqrt}[\text{Tan}[c + d \cdot x]]) / \text{Sqrt}[a]] + (-1)^{1/4} \cdot a^{7/2} \cdot (a - I \cdot b)^2 \cdot (A + I \cdot B) \cdot \text{ArcTanh}[(-1)^{3/4} \cdot \text{Sqrt}[\text{Tan}[c + d \cdot x]]]) / (a^{5/2} \cdot (a^2 + b^2)) + (-2 \cdot a^2 \cdot A - 5 \cdot A \cdot b^2 + 3 \cdot a \cdot b \cdot B) / (a \cdot \text{Tan}[c + d \cdot x]^{3/2}) + (3 \cdot (4 \cdot a^2 \cdot A \cdot b + 5 \cdot A \cdot b^3 - 2 \cdot a^3 \cdot B - 3 \cdot a \cdot b^2 \cdot B)) / (a^2 \cdot \text{Sqrt}[\text{Tan}[c + d \cdot x]]) + (3 \cdot b \cdot (A \cdot b - a \cdot B)) / (\text{Tan}[c + d \cdot x]^{3/2} \cdot (a + b \cdot \text{Tan}[c + d \cdot x])) / (3 \cdot a \cdot (a^2 + b^2) \cdot d)$

3.409.3 Rubi [A] (verified)

Time = 2.50 (sec) , antiderivative size = 433, normalized size of antiderivative = 0.88, number of steps used = 27, number of rules used = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.788$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2}(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{4092} \\
 & \frac{\int \frac{2Aa^2 - 3bBa - 2(Ab - aB) \tan(c + dx)a + 5Ab^2 + 5b(Ab - aB) \tan^2(c + dx)}{2 \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx}{\frac{a(a^2 + b^2)}{b(Ab - aB)}} + \\
 & \quad \frac{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{2Aa^2 - 3bBa - 2(Ab - aB) \tan(c + dx)a + 5Ab^2 + 5b(Ab - aB) \tan^2(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx}{\frac{2a(a^2 + b^2)}{b(Ab - aB)}} + \\
 & \quad \frac{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{2Aa^2 - 3bBa - 2(Ab - aB) \tan(c + dx)a + 5Ab^2 + 5b(Ab - aB) \tan(c + dx)^2}{\tan(c + dx)^{5/2}(a + b \tan(c + dx))} dx}{\frac{2a(a^2 + b^2)}{b(Ab - aB)}} + \\
 & \quad \frac{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} \\
 & \quad \downarrow \text{4132} \\
 & \frac{2 \int \frac{(-2Ba^3 + 4Aba^2 + 2(aA + bB) \tan(c + dx)a^2 - 3b^2Ba + 5Ab^3 + b(2Aa^2 - 3bBa + 5Ab^2) \tan^2(c + dx))}{2 \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx}{\frac{2a(a^2 + b^2)}{b(Ab - aB)}} - \frac{2(2a^2A - 3abB + 5Ab^2)}{3ad \tan^{\frac{3}{2}}(c + dx)} + \\
 & \quad \frac{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}
 \end{aligned}$$

3.409. $\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$

$$\begin{aligned}
 & \int \frac{-2Ba^3+4Aba^2+2(aA+bB)\tan(c+dx)a^2-3b^2Ba+5Ab^3+b(2Aa^2-3bBa+5Ab^2)\tan^2(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))} dx \\
 & \frac{2(2a^2A-3abB+5Ab^2)}{3ad\tan^{\frac{3}{2}}(c+dx)} + \\
 & \frac{2a(a^2+b^2)}{b(Ab-aB)} \\
 & \frac{ad(a^2+b^2)\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))}{\phantom{ad(a^2+b^2)\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))}} \\
 & \downarrow 3042 \\
 & \int \frac{-2Ba^3+4Aba^2+2(aA+bB)\tan(c+dx)a^2-3b^2Ba+5Ab^3+b(2Aa^2-3bBa+5Ab^2)\tan(c+dx)^2}{\tan(c+dx)^{3/2}(a+b\tan(c+dx))} dx \\
 & \frac{2(2a^2A-3abB+5Ab^2)}{3ad\tan^{\frac{3}{2}}(c+dx)} + \\
 & \frac{2a(a^2+b^2)}{b(Ab-aB)} \\
 & \frac{ad(a^2+b^2)\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))}{\phantom{ad(a^2+b^2)\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))}} \\
 & \downarrow 4132 \\
 & 2 \int \frac{2Aa^4+4bBa^3-2(Ab-aB)\tan(c+dx)a^3-4Ab^2a^2+3b^3Ba-5Ab^4-b(-2Ba^3+4Aba^2-3b^2Ba+5Ab^3)\tan^2(c+dx)}{2\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx \\
 & \frac{2(-2a^3B+4a^2Ab-3ab^2B+5Ab^3)}{ad\sqrt{\tan(c+dx)}} \\
 & \frac{2a(a^2+b^2)}{b(Ab-aB)} \\
 & \frac{ad(a^2+b^2)\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))}{\phantom{ad(a^2+b^2)\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))}} \\
 & \downarrow 27 \\
 & \int \frac{2Aa^4+4bBa^3-2(Ab-aB)\tan(c+dx)a^3-4Ab^2a^2+3b^3Ba-5Ab^4-b(-2Ba^3+4Aba^2-3b^2Ba+5Ab^3)\tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx \\
 & \frac{2(-2a^3B+4a^2Ab-3ab^2B+5Ab^3)}{ad\sqrt{\tan(c+dx)}} \\
 & \frac{2a(a^2+b^2)}{b(Ab-aB)} \\
 & \frac{ad(a^2+b^2)\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))}{\phantom{ad(a^2+b^2)\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))}} \\
 & \downarrow 3042 \\
 & \int \frac{2Aa^4+4bBa^3-2(Ab-aB)\tan(c+dx)a^3-4Ab^2a^2+3b^3Ba-5Ab^4-b(-2Ba^3+4Aba^2-3b^2Ba+5Ab^3)\tan(c+dx)^2}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx \\
 & \frac{2(-2a^3B+4a^2Ab-3ab^2B+5Ab^3)}{ad\sqrt{\tan(c+dx)}} \\
 & \frac{2a(a^2+b^2)}{b(Ab-aB)} \\
 & \frac{ad(a^2+b^2)\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))}{\phantom{ad(a^2+b^2)\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))}} \\
 & \downarrow 4136
 \end{aligned}$$

3.409. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^2} dx$

$$\frac{\int \frac{2(a^3(Aa^2+2bBa-Ab^2)-a^3(-Ba^2+2Aba+b^2B)\tan(c+dx))}{\sqrt{\tan(c+dx)}(a^2+b^2)} dx - \frac{b^3(-7a^3B+9a^2Ab-3ab^2B+5Ab^3) \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2}}{a} - \frac{2(-2a^3B+4a^2Ab-ab^3)}{ad\sqrt{\tan(c+dx)}}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} \quad \frac{2a(a^2 + b^2)}{a}$$

↓ 27

$$\frac{2 \int \frac{a^3(Aa^2+2bBa-Ab^2)-a^3(-Ba^2+2Aba+b^2B)\tan(c+dx)}{\sqrt{\tan(c+dx)}(a^2+b^2)} dx - \frac{b^3(-7a^3B+9a^2Ab-3ab^2B+5Ab^3) \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2}}{a} - \frac{2(-2a^3B+4a^2Ab-ab^3)}{ad\sqrt{\tan(c+dx)}}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} \quad \frac{2a(a^2 + b^2)}{a}$$

↓ 3042

$$\frac{2 \int \frac{a^3(Aa^2+2bBa-Ab^2)-a^3(-Ba^2+2Aba+b^2B)\tan(c+dx)}{\sqrt{\tan(c+dx)}(a^2+b^2)} dx - \frac{b^3(-7a^3B+9a^2Ab-3ab^2B+5Ab^3) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2}}{a} - \frac{2(-2a^3B+4a^2Ab-ab^3)}{ad\sqrt{\tan(c+dx)}}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} \quad \frac{2a(a^2 + b^2)}{a}$$

↓ 4017

$$\frac{4 \int \frac{a^3(Aa^2+2bBa-Ab^2)-(-Ba^2+2Aba+b^2B)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{b^3(-7a^3B+9a^2Ab-3ab^2B+5Ab^3) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2}}{a} - \frac{2(-2a^3B+4a^2Ab-ab^3)}{ad\sqrt{\tan(c+dx)}}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} \quad \frac{2a(a^2 + b^2)}{a}$$

↓ 27

3.409. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

$$\frac{4a^3 \int \frac{Aa^2+2bBa-Ab^2-(-Ba^2+2Aba+b^2B) \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - b^3(-7a^3B+9a^2Ab-3ab^2B+5Ab^3) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))} dx}{d(a^2+b^2)}}{a} - \frac{2(-2a^3B+4ad)}{a}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}$$

↓ 1482

$$\frac{4a^3 \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) - b^3(-7a^3B+9a^2Ab-3ab^2B+5Ab^3)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}$$

↓ 1476

$$\frac{4a^3 \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right) - b^3(-7a^3B+9a^2Ab-3ab^2B+5Ab^3)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}$$

↓ 1082

$$\frac{4a^3 \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) - b^3(-7a^3B+9a^2Ab-3ab^2B+5Ab^3)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}$$

↓ 217

3.409. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

$$4a^3 \left(\frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx) + 1} d\sqrt{\tan(c+dx)} - \frac{1}{2} (-(a^2(A+B)) + 2ab(A-B) + b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) \frac{1}{d(a^2 + b^2)}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}$$

↓ 1479

$$4a^3 \left(\frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(- \int \frac{\sqrt{2} - 2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2} (-(a^2(A+B)) + 2ab(A-B) + b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) \frac{1}{d(a^2 + b^2)}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}$$

↓ 25

$$4a^3 \left(\frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\int \frac{\sqrt{2} - 2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2} (-(a^2(A+B)) + 2ab(A-B) + b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) \frac{1}{d(a^2 + b^2)}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}$$

↓ 27

$$4a^3 \left(\frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\int \frac{\sqrt{2} - 2\sqrt{\tan(c+dx)}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)} + 1}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2} (-(a^2(A+B)) + 2ab(A-B) + b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) \frac{1}{d(a^2 + b^2)}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}$$

↓ 1103

3.409. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

$$\frac{4a^3 \left(\frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2} (-a^2(A+B) + 2ab(A-B) + b^2(A-B)) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}$$

↓ 4117

$$\frac{4a^3 \left(\frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2} (-a^2(A+B) + 2ab(A-B) + b^2(A-B)) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}$$

↓ 73

$$\frac{4a^3 \left(\frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2} (-a^2(A+B) + 2ab(A-B) + b^2(A-B)) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))}$$

↓ 218

$$\frac{b(Ab - aB)}{ad(a^2 + b^2) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} + \frac{4a^3 \left(\frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2} (-a^2(A+B) + 2ab(A-B) + b^2(A-B)) \right)}{d(a^2+b^2)}$$

$$\frac{2(2a^2A - 3abB + 5Ab^2)}{3ad \tan^{\frac{3}{2}}(c+dx)}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^2),x]`

3.409. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

```
output (-(((((-2*b^(5/2)*(9*a^2*A*b + 5*A*b^3 - 7*a^3*B - 3*a*b^2*B)*ArcTan[(Sqrt[
b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[a]*(a^2 + b^2)*d) + (4*a^3*(-1/2*((
2*a*b*(A - B) - a^2*(A + B) + b^2*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[
c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])) + (
(a^2*(A - B) - b^2*(A - B) + 2*a*b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan
[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] +
Tan[c + d*x]]/(2*Sqrt[2])))/2))/((a^2 + b^2)*d))/a - (2*(4*a^2*A*b + 5*A*b
^3 - 2*a^3*B - 3*a*b^2*B))/(a*d*Sqrt[Tan[c + d*x]]))/a - (2*(2*a^2*A + 5*
A*b^2 - 3*a*b*B))/(3*a*d*Tan[c + d*x]^(3/2)))/(2*a*(a^2 + b^2)) + (b*(A*b
- a*B))/(a*(a^2 + b^2)*d*Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x]))
```

3.409.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.409.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 373, normalized size of antiderivative = 0.76

method	result
derivativedivides	$2b^3 \left(\frac{\left(\frac{1}{2} A a^2 b + \frac{1}{2} A b^3 - \frac{1}{2} B a^3 - \frac{1}{2} B a b^2\right) (\sqrt{\tan(dx+c)}}}{a+b \tan(dx+c)} + \frac{(9 A a^2 b + 5 A b^3 - 7 B a^3 - 3 B a b^2) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{2\sqrt{ab}} \right) - \frac{2 A}{3 a^2 \tan(dx+c)} \frac{1}{a^3 (a^2+b^2)^2}$
default	$2b^3 \left(\frac{\left(\frac{1}{2} A a^2 b + \frac{1}{2} A b^3 - \frac{1}{2} B a^3 - \frac{1}{2} B a b^2\right) (\sqrt{\tan(dx+c)}}}{a+b \tan(dx+c)} + \frac{(9 A a^2 b + 5 A b^3 - 7 B a^3 - 3 B a b^2) \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{2\sqrt{ab}} \right) - \frac{2 A}{3 a^2 \tan(dx+c)} \frac{1}{a^3 (a^2+b^2)^2}$

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c))^2,x,method=_RETURNV ERBOSE)`

output `1/d*(2*b^3/a^3/(a^2+b^2)^2*((1/2*A*a^2*b+1/2*A*b^3-1/2*B*a^3-1/2*B*a*b^2)* tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))+1/2*(9*A*a^2*b+5*A*b^3-7*B*a^3-3*B*a*b^2)/(a*b)^(1/2)*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2)))-2/3/a^2*A/tan(d*x+c)^(3/2)-2*(-2*A*b+B*a)/a^3/tan(d*x+c)^(1/2)+2/(a^2+b^2)^2*(1/8*(-A*a^2+A*b^2-2*B*a*b)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)* tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/8*(2*A*a*b-B*a^2+B*b^2)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))`

3.409.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6095 vs. 2(450) = 900.

Time = 75.10 (sec) , antiderivative size = 12216, normalized size of antiderivative = 24.78

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c))^2,x, algorithm="fracas")`

output Too large to include

3.409.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(5/2)/(a+b*tan(d*x+c))**2,x)`

output Timed out

3.409.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 449, normalized size of antiderivative = 0.91

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \frac{12(7Ba^3b^3 - 9Aa^2b^4 + 3Bab^5 - 5Ab^6) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^7 + 2a^5b^2 + a^3b^4)\sqrt{ab}} + \frac{4(2Aa^4 + 2Aa^2b^2 + 3(2Ba^3b - 4Aa^2b^2 + 3Bab^3 - 5Ab^4) \tan(dx+c)^2 + 2(3a^5b + a^3b^3) \tan(dx+c)^{\frac{5}{2}} + (a^6 + a^4b^2) \tan(dx+c))}{(a^7 + 2a^5b^2 + a^3b^4)\sqrt{ab}}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `-1/12*(12*(7*B*a^3*b^3 - 9*A*a^2*b^4 + 3*B*a*b^5 - 5*A*b^6)*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/(a^7 + 2*a^5*b^2 + a^3*b^4)*sqrt(a*b)) + 4*(2*A*a^4 + 2*A*a^2*b^2 + 3*(2*B*a^3*b - 4*A*a^2*b^2 + 3*B*a*b^3 - 5*A*b^4)*tan(d*x + c)^2 + 2*(3*B*a^4 - 5*A*a^3*b + 3*B*a^2*b^2 - 5*A*a*b^3)*tan(d*x + c))/((a^5*b + a^3*b^3)*tan(d*x + c)^(5/2) + (a^6 + a^4*b^2)*tan(d*x + c)^(3/2)) + 3*(2*sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))/(a^4 + 2*a^2*b^2 + b^4))/d`

$$3.409. \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$$

3.409.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Timed out}$$

```
input integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c))^2,x, algorithm
m="giac")
```

```
output Timed out
```

3.409.9 Mupad [B] (verification not implemented)

Time = 27.53 (sec) , antiderivative size = 24620, normalized size of antiderivative = 49.94

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

```
input int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + b*tan(c + d*x))^2),x)
```

```
output (log(80*A^5*a^24*b^20*d^4 - (((192*A^4*a^2*b^6*d^4 - 16*A^4*b^8*d^4 - 16*
A^4*a^8*d^4 - 608*A^4*a^4*b^4*d^4 + 192*A^4*a^6*b^2*d^4)^(1/2) - 16*A^2*a*
b^3*d^2 + 16*A^2*a^3*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6*d^4 + 6*a^4*b^4
*d^4 + 4*a^6*b^2*d^4))^(1/2)*(tan(c + d*x)^(1/2)*(9472*A^4*a^31*b^15*d^5 -
3040*A^4*a^23*b^23*d^5 - 9056*A^4*a^25*b^21*d^5 - 12352*A^4*a^27*b^19*d^5
- 4256*A^4*a^29*b^17*d^5 - 400*A^4*a^21*b^25*d^5 + 13760*A^4*a^33*b^13*d^
5 + 7744*A^4*a^35*b^11*d^5 + 1968*A^4*a^37*b^9*d^5 + 224*A^4*a^39*b^7*d^5
+ 32*A^4*a^41*b^5*d^5) + (((192*A^4*a^2*b^6*d^4 - 16*A^4*b^8*d^4 - 16*A^4
*a^8*d^4 - 608*A^4*a^4*b^4*d^4 + 192*A^4*a^6*b^2*d^4)^(1/2) - 16*A^2*a*b^3
*d^2 + 16*A^2*a^3*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6*d^4 + 6*a^4*b^4*d^
4 + 4*a^6*b^2*d^4))^(1/2)*(12928*A^3*a^25*b^23*d^6 - 800*A^3*a^21*b^27*d^6
- 2080*A^3*a^23*b^25*d^6 - ((tan(c + d*x)^(1/2)*(3200*A^2*a^22*b^28*d^7 +
33920*A^2*a^24*b^26*d^7 + 158208*A^2*a^26*b^24*d^7 + 425536*A^2*a^28*b^22
*d^7 + 727296*A^2*a^30*b^20*d^7 + 820672*A^2*a^32*b^18*d^7 + 615936*A^2*a^
34*b^16*d^7 + 304256*A^2*a^36*b^14*d^7 + 98432*A^2*a^38*b^12*d^7 + 22016*A
^2*a^40*b^10*d^7 + 3072*A^2*a^42*b^8*d^7 - 704*A^2*a^44*b^6*d^7 - 512*A^2*
a^46*b^4*d^7 - 64*A^2*a^48*b^2*d^7) + (((192*A^4*a^2*b^6*d^4 - 16*A^4*b^8
*d^4 - 16*A^4*a^8*d^4 - 608*A^4*a^4*b^4*d^4 + 192*A^4*a^6*b^2*d^4)^(1/2) -
16*A^2*a*b^3*d^2 + 16*A^2*a^3*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6*d^4 +
6*a^4*b^4*d^4 + 4*a^6*b^2*d^4))^(1/2)*(1280*A*a^24*b^28*d^8 - (tan(c + ...
```

3.410
$$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

3.410.1 Optimal result	3937
3.410.2 Mathematica [B] (verified)	3938
3.410.3 Rubi [A] (verified)	3939
3.410.4 Maple [A] (verified)	3948
3.410.5 Fricas [B] (verification not implemented)	3949
3.410.6 Sympy [F(-1)]	3949
3.410.7 Maxima [A] (verification not implemented)	3949
3.410.8 Giac [F(-1)]	3950
3.410.9 Mupad [B] (verification not implemented)	3950

3.410.1 Optimal result

Integrand size = 33, antiderivative size = 600

$$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d}$$

$$- \frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d}$$

$$+ \frac{a^{3/2}(3a^4Ab + 6a^2Ab^3 + 35Ab^5 - 15a^5B - 46a^3b^2B - 63ab^4B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{4b^{7/2}(a^2+b^2)^3 d}$$

$$- \frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d}$$

$$+ \frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d}$$

$$- \frac{(3a^3Ab + 11aAb^3 - 15a^4B - 31a^2b^2B - 8b^4B) \sqrt{\tan(c+dx)}}{4b^3(a^2+b^2)^2 d}$$

$$+ \frac{a(Ab - aB) \tan^{\frac{5}{2}}(c+dx)}{2b(a^2+b^2)d(a+b \tan(c+dx))^2} + \frac{a(a^2Ab + 9Ab^3 - 5a^3B - 13ab^2B) \tan^{\frac{3}{2}}(c+dx)}{4b^2(a^2+b^2)^2 d(a+b \tan(c+dx))}$$

3.410.
$$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

output $\frac{1}{4}a^{3/2}(3Aa^4b+6Aa^2b^3+35Ab^5-15Ba^5-46Ba^3b^2-63Bab^4)\arctan(b^{1/2}\tan(dx+c)^{1/2}/a^{1/2})/b^{7/2}/(a^2+b^2)^3/d-1/2(3a^2b(A-B)-b^3(A-B)-a^3(A+B)+3ab^2(A+B))\arctan(-1+2^{1/2}\tan(dx+c)^{1/2})/(a^2+b^2)^3/d^2-1/2(3a^2b(A-B)-b^3(A-B)-a^3(A+B)+3ab^2(A+B))\arctan(1+2^{1/2}\tan(dx+c)^{1/2})/(a^2+b^2)^3/d^2-1/4(a^3(A-B)-3ab^2(A-B)+3a^2b(A+B)-b^3(A+B))\ln(1-2^{1/2}\tan(dx+c)^{1/2}+\tan(dx+c))/(a^2+b^2)^3/d^2+1/4(a^3(A-B)-3ab^2(A-B)+3a^2b(A+B)-b^3(A+B))\ln(1+2^{1/2}\tan(dx+c)^{1/2}+\tan(dx+c))/(a^2+b^2)^3/d^2-1/4(3Aa^3b+11Aa^2b^3-15Ba^4-31Ba^2b^2-8Bb^4)\tan(dx+c)^{1/2}/b^3/(a^2+b^2)^2/d+1/2a(Ab-Ba)\tan(dx+c)^{5/2}/b/(a^2+b^2)/d/(a+b\tan(dx+c))^2+1/4a(Aa^2b+9Ab^3-5Ba^3-13Bab^2)\tan(dx+c)^{3/2}/b^2/(a^2+b^2)^2/d/(a+b\tan(dx+c))$

3.410.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1351 vs. $2(600) = 1200$.

Time = 6.45 (sec) , antiderivative size = 1351, normalized size of antiderivative = 2.25

$$\int \frac{\tan^{7/2}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Too large to display}$$

input `Integrate[(Tan[c + d*x]^(7/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3, x]`

output $(2*B*\text{Tan}[c + d*x]^{(5/2)})/(b*d*(a + b*\text{Tan}[c + d*x])^2) + (2*(-(((A*b - 5*a*B)*\text{Tan}[c + d*x]^{(3/2)})/(b*d*(a + b*\text{Tan}[c + d*x])^2)) - (2*(-1/6*((-3*a*A*b + 15*a^2*B + b^2*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/(b*d*(a + b*\text{Tan}[c + d*x])^2) - (2*((((a*b^2*(3*a*A*b - 15*a^2*B - b^2*B))/8 - a*((3*b^4*B)/8 - (a*(3*a^2*A*b - 3*A*b^3 - 15*a^3*B - a*b^2*B))/8))*\text{Sqrt}[\text{Tan}[c + d*x]])/(2*a*(a^2 + b^2))*d*(a + b*\text{Tan}[c + d*x])^2) + (((2*(a^2*b*((3*a^2*b^3*(A*b - a*B))/4 + (3*a^2*b*(3*a^2*A*b + 4*A*b^3 - 15*a^3*B - 16*a*b^2*B))/16 - (3*a*b*(3*a^3*A*b - 15*a^4*B - 16*a^2*b^2*B - 4*b^4*B))/16) + (a^2*((3*a^2*b^2*(3*a^2*A*b + 4*A*b^3 - 15*a^3*B - 16*a*b^2*B))/16 - a*((-3*a*b^4*(A*b - a*B))/4 - (3*a^2*(3*a^3*A*b - 15*a^4*B - 16*a^2*b^2*B - 4*b^4*B))/16))))/2 + b^2*((3*a^2*(a^2 + b^2/2)*(3*a^2*A*b + 4*A*b^3 - 15*a^3*B - 16*a*b^2*B))/16 + (a*((-3*a*b^4*(A*b - a*B))/4 - (3*a^2*(3*a^3*A*b - 15*a^4*B - 16*a^2*b^2*B - 4*b^4*B))/16))/2))*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(\text{Sqrt}[a]*\text{Sqrt}[b]*(a^2 + b^2)*d) + (((-3*(-1)^{(1/4)}*a^5*A*b^3*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]])]/(4*d) - (9*(-1)^{(3/4)}*a^4*A*b^4*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]])]/(4*d) + (9*(-1)^{(1/4)}*a^3*A*b^5*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]])]/(4*d) + (3*(-1)^{(3/4)}*a^2*A*b^6*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]])]/(4*d) + (3*(-1)^{(3/4)}*a^5*b^3*B*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]])]/(4*d) - (9*(-1)^{(1/4)}*a^4*b^4*B*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]])]/(4*d) - (9*(-1)^{(3/4)}*a^3*b^5*B*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]])]/(4*...$

3.410.3 Rubi [A] (verified)

Time = 2.75 (sec) , antiderivative size = 528, normalized size of antiderivative = 0.88, number of steps used = 27, number of rules used = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.788$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4130, 27, 3042, 4136, 27, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{7}{2}}(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$$

↓ 3042

$$\int \frac{\tan(c + dx)^{7/2}(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$$

↓ 4088

3.410. $\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\begin{aligned}
 & \int \frac{-\tan^{\frac{3}{2}}(c+dx)((-5Ba^2+Aba-4b^2B)\tan^2(c+dx)-4b(Ab-aB)\tan(c+dx)+5a(Ab-aB))}{2(a+b\tan(c+dx))^2} dx \\
 & \qquad \qquad \qquad \frac{2b(a^2+b^2)}{a(Ab-aB)\tan^{\frac{5}{2}}(c+dx)} + \\
 & \qquad \qquad \qquad \frac{2bd(a^2+b^2)(a+b\tan(c+dx))^2}{27} \\
 & \qquad \qquad \qquad \frac{a(Ab-aB)\tan^{\frac{5}{2}}(c+dx)}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} - \\
 & \int \frac{\tan^{\frac{3}{2}}(c+dx)((-5Ba^2+Aba-4b^2B)\tan^2(c+dx)-4b(Ab-aB)\tan(c+dx)+5a(Ab-aB))}{(a+b\tan(c+dx))^2} dx \\
 & \qquad \qquad \qquad \frac{4b(a^2+b^2)}{3042} \\
 & \qquad \qquad \qquad \frac{a(Ab-aB)\tan^{\frac{5}{2}}(c+dx)}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} - \\
 & \int \frac{\tan(c+dx)^{3/2}((-5Ba^2+Aba-4b^2B)\tan(c+dx)^2-4b(Ab-aB)\tan(c+dx)+5a(Ab-aB))}{(a+b\tan(c+dx))^2} dx \\
 & \qquad \qquad \qquad \frac{4b(a^2+b^2)}{4128} \\
 & \qquad \qquad \qquad \frac{a(Ab-aB)\tan^{\frac{5}{2}}(c+dx)}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} - \\
 & \int \frac{\sqrt{\tan(c+dx)}(8(Aa^2+2bBa-Ab^2)\tan(c+dx)b^2+(-15Ba^4+3Aba^3-31b^2Ba^2+11Ab^3a-8b^4B)\tan^2(c+dx)+3a(-5Ba^3+Ab^2-13b^2Ba+9Ab^3))}{2(a+b\tan(c+dx))} dx - \frac{a(-5Ba^3+Ab^2-13b^2Ba+9Ab^3)}{b(a^2+b^2)} \\
 & \qquad \qquad \qquad \frac{4b(a^2+b^2)}{27} \\
 & \qquad \qquad \qquad \frac{a(Ab-aB)\tan^{\frac{5}{2}}(c+dx)}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} - \\
 & \int \frac{\sqrt{\tan(c+dx)}(8(Aa^2+2bBa-Ab^2)\tan(c+dx)b^2+(-15Ba^4+3Aba^3-31b^2Ba^2+11Ab^3a-8b^4B)\tan^2(c+dx)+3a(-5Ba^3+Ab^2-13b^2Ba+9Ab^3))}{a+b\tan(c+dx)} dx - \frac{a(-5Ba^3+Ab^2-13b^2Ba+9Ab^3)}{2b(a^2+b^2)} \\
 & \qquad \qquad \qquad \frac{4b(a^2+b^2)}{3042} \\
 & \qquad \qquad \qquad \frac{a(Ab-aB)\tan^{\frac{5}{2}}(c+dx)}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} - \\
 & \int \frac{\sqrt{\tan(c+dx)}(8(Aa^2+2bBa-Ab^2)\tan(c+dx)b^2+(-15Ba^4+3Aba^3-31b^2Ba^2+11Ab^3a-8b^4B)\tan^2(c+dx)+3a(-5Ba^3+Ab^2-13b^2Ba+9Ab^3))}{a+b\tan(c+dx)} dx - \frac{a(-5Ba^3+Ab^2-13b^2Ba+9Ab^3)}{2b(a^2+b^2)} \\
 & \qquad \qquad \qquad \frac{4b(a^2+b^2)}{4130}
 \end{aligned}$$

3.410. $\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{-8(-Ba^2 + 2Aba + b^2B) \tan(c + dx)b^3 + (-15Ba^5 + 3Aba^4 - 31b^2Ba^3 + 3Ab^3a^2 - 24b^4Ba + 8Ab^5) \tan^2(c + dx) + a(-15Ba^4 + 3Aba^3 - 31b^2Ba^2 + 11Ab^3a - 8b^4B)}{2\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} \frac{1}{b}$$

$$\frac{2b(a^2 + b^2)}{4b(a^2 + b^2)}$$

↓ 27

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c + dx)}}{bd} - \frac{-8(-Ba^2 + 2Aba + b^2B) \tan(c + dx)b^3 + (-15Ba^5 + 3Aba^4 - 31b^2Ba^3 + 3Ab^3a^2 - 24b^4Ba + 8Ab^5) \tan^2(c + dx) + a(-15Ba^4 + 3Aba^3 - 31b^2Ba^2 + 11Ab^3a - 8b^4B)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} \frac{1}{b}$$

$$\frac{2b(a^2 + b^2)}{4b(a^2 + b^2)}$$

↓ 3042

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c + dx)}}{bd} - \frac{-8(-Ba^2 + 2Aba + b^2B) \tan(c + dx)b^3 + (-15Ba^5 + 3Aba^4 - 31b^2Ba^3 + 3Ab^3a^2 - 24b^4Ba + 8Ab^5) \tan^2(c + dx) + a(-15Ba^4 + 3Aba^3 - 31b^2Ba^2 + 11Ab^3a - 8b^4B)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} \frac{1}{b}$$

$$\frac{2b(a^2 + b^2)}{4b(a^2 + b^2)}$$

↓ 4136

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c + dx)}}{bd} - \frac{8(b^3(Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) - b^3(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \tan(c + dx)) dx}{\sqrt{\tan(c + dx)}(a^2 + b^2)} + \frac{a^2(-15a^5B + 15a^4Ab - 15a^3Ab^2 + 15a^2Ab^3 - 15Ab^4)}{b}$$

$$\frac{2b(a^2 + b^2)}{4b(a^2 + b^2)}$$

↓ 27

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c + dx)}}{bd} - \frac{8(b^3(Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) - b^3(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \tan(c + dx)) dx}{\sqrt{\tan(c + dx)}(a^2 + b^2)} + \frac{a^2(-15a^5B + 15a^4Ab - 15a^3Ab^2 + 15a^2Ab^3 - 15Ab^4)}{b}$$

$$\frac{2b(a^2 + b^2)}{4b(a^2 + b^2)}$$

↓ 3042

3.410. $\int \frac{\tan^{\frac{7}{2}}(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c + dx)}}{bd} - \frac{8 \int \frac{b^3(Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) - b^3(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx}{a^2 + b^2} + \frac{a^2(-15a^5B + 15a^4Ab - 15a^3A^2b + 15a^2Ab^2 - 15aAb^3 + 15a^2b^2B - 15aAb^3 - 8b^4B)}{b}$$

$$\frac{4b(a^2 + b^2)}{2b(a^2 + b^2)}$$

↓ 4017

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c + dx)}}{bd} - \frac{16 \int \frac{b^3(Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) - (-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{d(a^2 + b^2)} + \frac{a^2(-15a^5B + 15a^4Ab - 15a^3A^2b + 15a^2Ab^2 - 15aAb^3 + 15a^2b^2B - 15aAb^3 - 8b^4B)}{b}$$

$$\frac{4b(a^2 + b^2)}{2b(a^2 + b^2)}$$

↓ 27

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c + dx)}}{bd} - \frac{16b^3 \int \frac{Aa^3 + 3bBa^2 - 3Ab^2a - b^3B - (-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{d(a^2 + b^2)} + \frac{a^2(-15a^5B + 15a^4Ab - 15a^3A^2b + 15a^2Ab^2 - 15aAb^3 + 15a^2b^2B - 15aAb^3 - 8b^4B)}{b}$$

$$\frac{4b(a^2 + b^2)}{2b(a^2 + b^2)}$$

↓ 1482

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c + dx)}}{bd} - \frac{16b^3 \left(\frac{1}{2}(a^3(A - B) + 3a^2b(A + B) - 3ab^2(A - B) - b^3(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(-a^3 + 3a^2b - 3ab^2 + b^3) \right)}{d(a^2 + b^2)} + \frac{a^2(-15a^5B + 15a^4Ab - 15a^3A^2b + 15a^2Ab^2 - 15aAb^3 + 15a^2b^2B - 15aAb^3 - 8b^4B)}{b}$$

↓ 1476

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c + dx)}}{bd} - \frac{16b^3 \left(\frac{1}{2}(a^3(A - B) + 3a^2b(A + B) - 3ab^2(A - B) - b^3(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(-a^3 + 3a^2b - 3ab^2 + b^3) \right)}{d(a^2 + b^2)} + \frac{a^2(-15a^5B + 15a^4Ab - 15a^3A^2b + 15a^2Ab^2 - 15aAb^3 + 15a^2b^2B - 15aAb^3 - 8b^4B)}{b}$$

↓ 1082

3.410. $\int \frac{\tan^{\frac{7}{2}}(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{16b^3 \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2} \left(-\left(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B) \right) \sqrt{\tan(c+dx)} \right) \right)}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c+dx)}} - \frac{16b^3 \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2} \left(-\left(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B) \right) \sqrt{\tan(c+dx)} \right) \right)}{bd}$$

↓ 217

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{16b^3 \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2} \left(-\left(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B) \right) \sqrt{\tan(c+dx)} \right) \right)}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c+dx)}} - \frac{16b^3 \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2} \left(-\left(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B) \right) \sqrt{\tan(c+dx)} \right) \right)}{bd}$$

↓ 1479

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{16b^3 \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c+dx)}} - \frac{16b^3 \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{bd}$$

↓ 25

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{16b^3 \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c+dx)}} - \frac{16b^3 \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{bd}$$

↓ 27

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{16b^3 \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c+dx)}} - \frac{16b^3 \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{bd}$$

3.410. $\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

↓ 1103

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c+dx)}} - \frac{16b^3}{a^2 + b^2}$$

↓ 4117

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} d \tan(c+dx)}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c+dx)}} - \frac{d \tan(c+dx)}{d(a^2 + b^2)}$$

↓ 73

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{2a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{1}{a+b \tan(c+dx)} d \sqrt{\tan(c+dx)}}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c+dx)}} + \frac{16b^3}{d(a^2 + b^2)}$$

↓ 218

$$\frac{a(Ab - aB) \tan^{\frac{5}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{16b^3 \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)} + 1)}{2\sqrt{2}} \right) \right)}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B) \sqrt{\tan(c+dx)}} - \frac{1}{d(a^2 + b^2)}$$

input `Int[(Tan[c + d*x]^(7/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

3.410. $\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

```
output (a*(A*b - a*B)*Tan[c + d*x]^(5/2))/(2*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])
^2) - (((2*a^(3/2)*(3*a^4*A*b + 6*a^2*A*b^3 + 35*A*b^5 - 15*a^5*B - 46*
a^3*b^2*B - 63*a*b^4*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqr
t[b]*(a^2 + b^2)*d) + (16*b^3*(-1/2*((3*a^2*b*(A - B) - b^3*(A - B) - a^3*
(A + B) + 3*a*b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[
2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])) + ((a^3*(A - B) - 3
*a*b^2*(A - B) + 3*a^2*b*(A + B) - b^3*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt
[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]
] + Tan[c + d*x]]/(2*Sqrt[2])))/2))/((a^2 + b^2)*d)/b) + (2*(3*a^3*A*b +
11*a*A*b^3 - 15*a^4*B - 31*a^2*b^2*B - 8*b^4*B)*Sqrt[Tan[c + d*x]])/(b*d)
/(2*b*(a^2 + b^2)) - (a*(a^2*A*b + 9*A*b^3 - 5*a^3*B - 13*a*b^2*B)*Tan[c +
d*x]^(3/2))/(b*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(4*b*(a^2 + b^2))
```

3.410.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 73 Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

$$3.410. \quad \int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

$$3.410. \int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
!GtQ[n, 0] && !LeQ[n, -1]
```

3.410.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 466, normalized size of antiderivative = 0.78

method	result
derivativedivides	$\frac{2(\sqrt{\tan(dx+c)}B}{b^3} + \frac{2a^2 \left(\frac{-\frac{5}{8}Aa^4b^2 - \frac{9}{4}Aa^2b^4 - \frac{13}{8}Ab^6 + \frac{9}{8}Ba^5b + \frac{13}{4}Ba^3b^3 + \frac{17}{8}Bab^5 \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) - \frac{a(3Aa^4b + 14Aa^2b^3 + 11Ab^5 - 7Bb^3)}{(a+b \tan(dx+c))^2}}{(a+b \tan(dx+c))^2}$
default	$\frac{2(\sqrt{\tan(dx+c)}B}{b^3} + \frac{2a^2 \left(\frac{-\frac{5}{8}Aa^4b^2 - \frac{9}{4}Aa^2b^4 - \frac{13}{8}Ab^6 + \frac{9}{8}Ba^5b + \frac{13}{4}Ba^3b^3 + \frac{17}{8}Bab^5 \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) - \frac{a(3Aa^4b + 14Aa^2b^3 + 11Ab^5 - 7Bb^3)}{(a+b \tan(dx+c))^2}}{(a+b \tan(dx+c))^2}$

```
input int(tan(d*x+c)^(7/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x,method=_RETURNV
ERBOSE)
```

```
output 1/d*(2*tan(d*x+c)^(1/2)*B/b^3+2*a^2/b^3/(a^2+b^2)^3*((-5/8*A*a^4*b^2-9/4*
A*a^2*b^4-13/8*A*b^6+9/8*B*a^5*b+13/4*B*a^3*b^3+17/8*B*a*b^5)*tan(d*x+c)^(
3/2)-1/8*a*(3*A*a^4*b+14*A*a^2*b^3+11*A*b^5-7*B*b^3-22*B*a^3*b^2-15*B*a*b^
4)*tan(d*x+c)^(1/2))/(a+b*tan(d*x+c))^2+1/8*(3*A*a^4*b+6*A*a^2*b^3+35*A*b^
5-15*B*a^5-46*B*a^3*b^2-63*B*a*b^4)/(a*b)^(1/2)*arctan(b*tan(d*x+c)^(1/2)/
(a*b)^(1/2)))+2/(a^2+b^2)^3*(1/8*(A*a^3-3*A*a*b^2+3*B*a^2*b-B*b^3)*2^(1/2)
*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+t
an(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d
*x+c)^(1/2)))+1/8*(-3*A*a^2*b+A*b^3+B*a^3-3*B*a*b^2)*2^(1/2)*(ln((1-2^(1/2)
)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*
arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))
)
```

$$3.410. \int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

3.410.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8982 vs. $2(545) = 1090$.

Time = 166.93 (sec) , antiderivative size = 17991, normalized size of antiderivative = 29.98

$$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(7/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm m="fricas")`

output Too large to include

3.410.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(7/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)`

output Timed out

3.410.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 571, normalized size of antiderivative = 0.95

$$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \frac{(15Ba^7 - 3Aa^6b + 46Ba^5b^2 - 6Aa^4b^3 + 63Ba^3b^4 - 35Aa^2b^5) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right) - 2\sqrt{2}((A+B)a^3 - 3(A-B)a^2b - 3(A+B)ab^2 + (A-B)a^2b^3 + 3a^4b^5 + 3a^2b^7 + b^9)\sqrt{ab}}{(a^6b^3 + 3a^4b^5 + 3a^2b^7 + b^9)\sqrt{ab}}$$

input `integrate(tan(d*x+c)^(7/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm m="maxima")`

3.410. $\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

output
$$-1/4*((15*B*a^7 - 3*A*a^6*b + 46*B*a^5*b^2 - 6*A*a^4*b^3 + 63*B*a^3*b^4 - 35*A*a^2*b^5)*\arctan(b*\sqrt{\tan(dx + c)}/\sqrt{a*b})/((a^6*b^3 + 3*a^4*b^5 + 3*a^2*b^7 + b^9)*\sqrt{a*b}) - (2*\sqrt{2}*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2*\sqrt{\tan(dx + c)})) + 2*\sqrt{2}*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2*\sqrt{\tan(dx + c)})) + \sqrt{2}*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*\log(\sqrt{2}*\sqrt{\tan(dx + c)} + \tan(dx + c) + 1) - \sqrt{2}*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*\log(-\sqrt{2}*\sqrt{\tan(dx + c)} + \tan(dx + c) + 1))/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - ((9*B*a^5*b - 5*A*a^4*b^2 + 17*B*a^3*b^3 - 13*A*a^2*b^4)*\tan(dx + c)^{(3/2)} + (7*B*a^6 - 3*A*a^5*b + 15*B*a^4*b^2 - 11*A*a^3*b^3)*\sqrt{\tan(dx + c)})/(a^6*b^3 + 2*a^4*b^5 + a^2*b^7 + (a^4*b^5 + 2*a^2*b^7 + b^9)*\tan(dx + c)^2 + 2*(a^5*b^4 + 2*a^3*b^6 + a*b^8)*\tan(dx + c)) - 8*B*\sqrt{\tan(dx + c)}/b^3)/d$$

3.410.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{7}{2}}(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate(tan(dx+c)^(7/2)*(A+B*tan(dx+c))/(a+b*tan(dx+c))^3,x, algorithm="giac")`

output Timed out

3.410.9 Mupad [B] (verification not implemented)

Time = 71.27 (sec) , antiderivative size = 27429, normalized size of antiderivative = 45.72

$$\int \frac{\tan^{\frac{7}{2}}(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

input `int((tan(c + d*x)^(7/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3,x)`

3.410.
$$\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

output $(\log(\frac{(((((128b^3 \tan(c+dx)^{1/2}(a^2-b^2)(a^2+b^2)^2(4(-B^4d^4(a^6-b^6+15a^2b^4-15a^4b^2)^2)^{1/2} + 80B^2a^3b^3d^2 - 24B^2a^5bd^2 - 24B^2a^5bd^2)/(d^4(a^2+b^2)^6))^{1/2} - (64Ba^2b(15a^4+2b^4+41a^2b^2))/d)((4(-B^4d^4(a^6-b^6+15a^2b^4-15a^4b^2)^2)^{1/2} + 80B^2a^3b^3d^2 - 24B^2a^5bd^2 - 24B^2a^5bd^2)/(d^4(a^2+b^2)^6))^{1/2})/4 + (8B^2a \tan(c+dx)^{1/2}(225a^{14} - 184b^{14} + 608a^2b^{12} - 272a^4b^{10} + 3937a^6b^8 + 5804a^8b^6 + 4006a^{10}b^4 + 1380a^{12}b^2))/(b^4d^2(a^2+b^2)^4))((4(-B^4d^4(a^6-b^6+15a^2b^4-15a^4b^2)^2)^{1/2} + 80B^2a^3b^3d^2 - 24B^2a^5bd^2 - 24B^2a^5bd^2)/(d^4(a^2+b^2)^6))^{1/2})/4 - (2B^3a^2(1125a^{14} + 16b^{14} + 6112a^2b^{12} - 17727a^4b^{10} - 23239a^6b^8 - 11174a^8b^6 + 2930a^{10}b^4 + 3525a^{12}b^2))/(b^4d^3(a^2+b^2)^6))((4(-B^4d^4(a^6-b^6+15a^2b^4-15a^4b^2)^2)^{1/2} + 80B^2a^3b^3d^2 - 24B^2a^5bd^2 - 24B^2a^5bd^2)/(d^4(a^2+b^2)^6))^{1/2})/4 - (B^4 \tan(c+dx)^{1/2}(32b^{18} - 225a^{18} + 128a^2b^{16} + 192a^4b^{14} - 3841a^6b^{12} + 18050a^8b^{10} + 26801a^{10}b^8 + 16860a^{12}b^6 + 4049a^{14}b^4 - 30a^{16}b^2))/(b^5d^4(a^2+b^2)^8))((4(-B^4d^4(a^6-b^6+15a^2b^4-15a^4b^2)^2)^{1/2} + 80B^2a^3b^3d^2 - 24B^2a^5bd^2 - 24B^2a^5bd^2)/(d^4(a^2+b^2)^6))^{1/2})/4 - (B^5a^3(225a^{12} + 504b^{12} + 872a^2b^{10} + 4457a^4b^8 + 5916a^6b^6 + 4006a...$

3.410. $\int \frac{\tan^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$3.411 \quad \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

3.411.1 Optimal result	3952
3.411.2 Mathematica [C] (verified)	3953
3.411.3 Rubi [A] (verified)	3954
3.411.4 Maple [A] (verified)	3962
3.411.5 Fricas [B] (verification not implemented)	3962
3.411.6 Sympy [F(-1)]	3963
3.411.7 Maxima [A] (verification not implemented)	3963
3.411.8 Giac [F(-1)]	3964
3.411.9 Mupad [B] (verification not implemented)	3964

3.411.1 Optimal result

Integrand size = 33, antiderivative size = 534

$$\begin{aligned} & \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\ &= \frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d} \\ & \quad - \frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d} \\ & \quad + \frac{\sqrt{a}(a^4Ab + 18a^2Ab^3 - 15Ab^5 + 3a^5B + 6a^3b^2B + 35ab^4B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{4b^{5/2}(a^2+b^2)^3 d} \\ & \quad + \frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d} \\ & \quad - \frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d} \\ & \quad + \frac{a(Ab - aB) \tan^{\frac{3}{2}}(c+dx)}{2b(a^2+b^2)d(a+b \tan(c+dx))^2} - \frac{a(a^2Ab - 7Ab^3 + 3a^3B + 11ab^2B) \sqrt{\tan(c+dx)}}{4b^2(a^2+b^2)^2 d(a+b \tan(c+dx))} \end{aligned}$$

$$3.411. \quad \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

output
$$-1/2*(a^3*(A-B)-3*a*b^2*(A-B)+3*a^2*b*(A+B)-b^3*(A+B))*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/(a^2+b^2)^3/d*2^{(1/2)}-1/2*(a^3*(A-B)-3*a*b^2*(A-B)+3*a^2*b*(A+B)-b^3*(A+B))*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/(a^2+b^2)^3/d*2^{(1/2)}+1/4*(3*a^2*b*(A-B)-b^3*(A-B)-a^3*(A+B)+3*a*b^2*(A+B))*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(a^2+b^2)^3/d*2^{(1/2)}-1/4*(3*a^2*b*(A-B)-b^3*(A-B)-a^3*(A+B)+3*a*b^2*(A+B))*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(a^2+b^2)^3/d*2^{(1/2)}+1/4*(A*a^4*b+18*A*a^2*b^3-15*A*b^5+3*B*a^5+6*B*a^3*b^2+35*B*a*b^4)*\arctan(b^{(1/2)}*\tan(d*x+c)^{(1/2)}/a^{(1/2)})*a^{(1/2)}/b^{(5/2)}/(a^2+b^2)^3/d+1/2*a*(A*b-B*a)*\tan(d*x+c)^{(3/2)}/b/(a^2+b^2)/d/(a+b*\tan(d*x+c))^2-1/4*a*(A*a^2*b-7*A*b^3+3*B*a^3+11*B*a*b^2)*\tan(d*x+c)^{(1/2)}/b^2/(a^2+b^2)^2/d/(a+b*\tan(d*x+c))$$

3.411.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.42 (sec) , antiderivative size = 690, normalized size of antiderivative = 1.29

$$\int \frac{\tan^{5/2}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx = -\frac{2B \tan^{3/2}(c+dx)}{bd(a+b \tan(c+dx))^2}$$

$$2 \left(-\frac{(-Ab-3aB)\sqrt{\tan(c+dx)}}{3bd(a+b \tan(c+dx))^2} - \frac{\left(\frac{1}{4}ab^2(Ab+3aB) - a \left(-\frac{3Ab^3}{4} - \frac{1}{4}a(aAb+3a^2B-3b^2B) \right) \right) \sqrt{\tan(c+dx)}}{2a(a^2+b^2)d(a+b \tan(c+dx))^2} + \frac{2\left(\frac{3}{2}a^3b^3(a^2A-Ab^2+2abB)\right) + \frac{3}{16}a^3b^3}{\dots} \right)$$

input `Integrate[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3, x]`

3.411.
$$\int \frac{\tan^{5/2}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

output $(-2*B*\text{Tan}[c + d*x]^{(3/2)})/(b*d*(a + b*\text{Tan}[c + d*x])^2) - (2*(-1/3*((-(A*b) - 3*a*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/(b*d*(a + b*\text{Tan}[c + d*x])^2) - (2*(((a*b^2*(A*b + 3*a*B))/4 - a*((-3*A*b^3)/4 - (a*(a*A*b + 3*a^2*B - 3*b^2*B))/4))*\text{Sqrt}[\text{Tan}[c + d*x]])/(2*a*(a^2 + b^2)*d*(a + b*\text{Tan}[c + d*x])^2) + (((2*((3*a^3*b^3*(a^2*A - A*b^2 + 2*a*b*B))/2 + (3*a^3*b^2*(a^2*A*b - 7*A*b^3 + 3*a^3*B + 11*a*b^2*B))/16 + (3*a^4*(a^3*A*b + 9*a*A*b^3 + 3*a^4*B + 3*a^2*b^2*B + 8*b^4*B))/16)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(\text{Sqrt}[a]*\text{Sqrt}[b]*(a^2 + b^2)*d) + (-(((-1)^{(1/4)}*((-3*a^2*b^2*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B))/2 + ((3*I)/2)*a^2*b^2*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B))*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]])/d) - ((-1)^{(1/4)}*((-3*a^2*b^2*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B))/2 - ((3*I)/2)*a^2*b^2*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B))*\text{ArcTanh}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]])/d)/(a^2 + b^2))/(a*(a^2 + b^2)) + (((3*a^2*b^2*(a*A*b + 3*a^2*B + 4*b^2*B))/8 - a*((-3*a^2*(a^2*A*b + 4*A*b^3 + 3*a^3*B))/8 - (3*a*b^3*(a*A + b*B))/2))*\text{Sqrt}[\text{Tan}[c + d*x]])/(a*(a^2 + b^2)*d*(a + b*\text{Tan}[c + d*x]))/(2*a*(a^2 + b^2)))/(3*b))/b$

3.411.3 Rubi [A] (verified)

Time = 2.06 (sec) , antiderivative size = 466, normalized size of antiderivative = 0.87, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.697$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4136, 27, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

↓ 3042

$$\int \frac{\tan(c+dx)^{5/2}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

↓ 4088

$$\int -\frac{\sqrt{\tan(c+dx)}(-((3Ba^2+Aba+4b^2B) \tan^2(c+dx))-4b(Ab-aB) \tan(c+dx)+3a(Ab-aB))}{2(a+b \tan(c+dx))^2} dx +$$

$$\frac{a(Ab-aB) \tan^{\frac{3}{2}}(c+dx)}{2bd(a^2+b^2)(a+b \tan(c+dx))^2}$$

↓ 27

3.411. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \int \frac{\sqrt{\tan(c+dx)}(-((3Ba^2 + Aba + 4b^2B) \tan^2(c+dx)) - 4b(Ab - aB) \tan(c+dx) + 3a(Ab - aB))}{(a + b \tan(c+dx))^2} dx}{4b(a^2 + b^2)}$$

↓ 3042

$$\frac{\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \int \frac{\sqrt{\tan(c+dx)}(-((3Ba^2 + Aba + 4b^2B) \tan(c+dx)^2) - 4b(Ab - aB) \tan(c+dx) + 3a(Ab - aB))}{(a + b \tan(c+dx))^2} dx}{4b(a^2 + b^2)}$$

↓ 4128

$$\frac{\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \int -\frac{8(Aa^2 + 2bBa - Ab^2) \tan(c+dx)b^2 + (3Ba^4 + Aba^3 + 3b^2Ba^2 + 9Ab^3a + 8b^4B) \tan^2(c+dx) + a(3Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3)}{2\sqrt{\tan(c+dx)}(a + b \tan(c+dx))} dx}{b(a^2 + b^2)} + \frac{a(3a^3B + a^2Ab + 11ab^2B)}{bd(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 27

$$\frac{\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^3B + a^2Ab + 11ab^2B - 7Ab^3) \sqrt{\tan(c+dx)}}{bd(a^2 + b^2)(a + b \tan(c+dx))} - \int \frac{-8(Aa^2 + 2bBa - Ab^2) \tan(c+dx)b^2 + (3Ba^4 + Aba^3 + 3b^2Ba^2 + 9Ab^3a + 8b^4B) \tan^2(c+dx) + a(3Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3)}{\sqrt{\tan(c+dx)}(a + b \tan(c+dx))}}{2b(a^2 + b^2)}}{4b(a^2 + b^2)}$$

↓ 3042

$$\frac{\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^3B + a^2Ab + 11ab^2B - 7Ab^3) \sqrt{\tan(c+dx)}}{bd(a^2 + b^2)(a + b \tan(c+dx))} - \int \frac{-8(Aa^2 + 2bBa - Ab^2) \tan(c+dx)b^2 + (3Ba^4 + Aba^3 + 3b^2Ba^2 + 9Ab^3a + 8b^4B) \tan(c+dx)^2 + a(3Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3)}{\sqrt{\tan(c+dx)}(a + b \tan(c+dx))}}{2b(a^2 + b^2)}}{4b(a^2 + b^2)}$$

↓ 4136

$$\frac{\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^3B + a^2Ab + 11ab^2B - 7Ab^3) \sqrt{\tan(c+dx)}}{bd(a^2 + b^2)(a + b \tan(c+dx))} - \frac{\int -\frac{8((-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3)b^2 + (Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) \tan(c+dx)b^2)}{\sqrt{\tan(c+dx)}} dx}{a^2 + b^2}}{a^2 + b^2}}{2b(a^2 + b^2)}}{4b(a^2 + b^2)}$$

↓ 27

3.411. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx - 8 \int \frac{(-Ba^3+3A}{bd(a^2+b^2)(a+b \tan(c+dx)) \sqrt{\tan(c+dx)}}}{a^2+b^2} dx}{4b(a^2 + b^2)}$$

↓ 3042

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx - 8 \int \frac{(-Ba^3+3A}{bd(a^2+b^2)(a+b \tan(c+dx)) \sqrt{\tan(c+dx)}}}{a^2+b^2} dx}{4b(a^2 + b^2)}$$

↓ 4017

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx - 16 \int \frac{b^2(-Ba^3+3A}{bd(a^2+b^2)(a+b \tan(c+dx)) \sqrt{\tan(c+dx)}}}{a^2+b^2} dx}{4b(a^2 + b^2)}$$

↓ 27

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx - 16b^2 \int \frac{-Ba^3+3A}{bd(a^2+b^2)(a+b \tan(c+dx)) \sqrt{\tan(c+dx)}}}{4b(a^2 + b^2)}$$

↓ 1482

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx - 16b^2 \left(\frac{1}{2} (-a^3 + 3A) \int \frac{1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx \right)}{4b(a^2 + b^2)}$$

↓ 1476

3.411. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx}{a^2 + b^2} - \frac{16b^2 \left(\frac{1}{2} (-a^3) \right)}{bd(a^2 + b^2)(a + b \tan(c + dx)) \sqrt{\tan(c + dx)}}$$

↓ 1082

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx}{a^2 + b^2} - \frac{16b^2 \left(\frac{1}{2} (-a^3) \right)}{bd(a^2 + b^2)(a + b \tan(c + dx)) \sqrt{\tan(c + dx)}}$$

↓ 217

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx}{a^2 + b^2} - \frac{16b^2 \left(\frac{1}{2} (-a^3) \right)}{bd(a^2 + b^2)(a + b \tan(c + dx)) \sqrt{\tan(c + dx)}}$$

↓ 1479

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx}{a^2 + b^2} - \frac{16b^2 \left(\frac{1}{2} (-a^3) \right)}{bd(a^2 + b^2)(a + b \tan(c + dx)) \sqrt{\tan(c + dx)}}$$

↓ 25

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx}{a^2 + b^2} - \frac{16b^2 \left(\frac{1}{2} (-a^3) \right)}{bd(a^2 + b^2)(a + b \tan(c + dx)) \sqrt{\tan(c + dx)}}$$

↓ 27

3.411. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))} dx}{a^2 + b^2} - \frac{16b^2 \left(\frac{1}{2}(-a^3(A - B) + \dots) \right)}{bd(a^2 + b^2)(a + b \tan(c + dx)) \sqrt{\tan(c + dx)}}$$

1103

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))} dx}{a^2 + b^2} - \frac{16b^2 \left(\frac{1}{2}(a^3(A - B) + \dots) \right)}{bd(a^2 + b^2)(a + b \tan(c + dx)) \sqrt{\tan(c + dx)}}$$

4117

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{1}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))} d \tan(c+dx)}{d(a^2 + b^2)} - \frac{16b^2 \left(\frac{1}{2}(a^3(A - B) + \dots) \right)}{bd(a^2 + b^2)(a + b \tan(c + dx)) \sqrt{\tan(c + dx)}}$$

73

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{2a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{1}{a + b \tan(c + dx)} d \sqrt{\tan(c + dx)}}{d(a^2 + b^2)} - \frac{16b^2 \left(\frac{1}{2}(a^3(A - B) + \dots) \right)}{bd(a^2 + b^2)(a + b \tan(c + dx)) \sqrt{\tan(c + dx)}}$$

218

$$\frac{a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \frac{2\sqrt{a}(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{b}d(a^2 + b^2)} - \frac{16b^2 \left(\frac{1}{2}(a^3(A - B) + \dots) \right)}{bd(a^2 + b^2)(a + b \tan(c + dx)) \sqrt{\tan(c + dx)}}$$

```
input Int[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]
```

3.411. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

```
output (a*(A*b - a*B)*Tan[c + d*x]^(3/2))/(2*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])
^2) - (-1/2*((2*Sqrt[a]*(a^4*A*b + 18*a^2*A*b^3 - 15*A*b^5 + 3*a^5*B + 6*a
^3*b^2*B + 35*a*b^4*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt
[b]*(a^2 + b^2)*d) - (16*b^2*((a^3*(A - B) - 3*a*b^2*(A - B) + 3*a^2*b*(A
+ B) - b^3*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) +
ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]))/2 + ((3*a^2*b*(A - B) - b
^3*(A - B) - a^3*(A + B) + 3*a*b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan
[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] +
Tan[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)/(b*(a^2 + b^2)) + (a*(a^2
*A*b - 7*A*b^3 + 3*a^3*B + 11*a*b^2*B)*Sqrt[Tan[c + d*x]])/(b*(a^2 + b^2)*
d*(a + b*Tan[c + d*x]))/(4*b*(a^2 + b^2))
```

3.411.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 217 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 1082 Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

$$3.411. \quad \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4136 `Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.411.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 451, normalized size of antiderivative = 0.84

method	result
derivativedivides	$2a \frac{\left(\frac{A a^4 b + 10 A a^2 b^3 + 9 A b^5 - 5 B a^5 - 18 B a^3 b^2 - 13 B a b^4}{8b} \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) - a \frac{A a^4 b - 6 A a^2 b^3 - 7 A b^5 + 3 B a^5 + 14 B a^3 b^2 + 11 B a b^4}{8b^2}}{(a+b \tan(dx+c))^2} \frac{1}{(a^2+b^2)^3}$
default	$2a \frac{\left(\frac{A a^4 b + 10 A a^2 b^3 + 9 A b^5 - 5 B a^5 - 18 B a^3 b^2 - 13 B a b^4}{8b} \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) - a \frac{A a^4 b - 6 A a^2 b^3 - 7 A b^5 + 3 B a^5 + 14 B a^3 b^2 + 11 B a b^4}{8b^2}}{(a+b \tan(dx+c))^2} \frac{1}{(a^2+b^2)^3}$

```
input int(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x,method=_RETURNV
ERBOSE)
```

```
output 1/d*(2*a/(a^2+b^2)^3*((1/8*(A*a^4*b+10*A*a^2*b^3+9*A*b^5-5*B*a^5-18*B*a^3*
b^2-13*B*a*b^4)/b*tan(d*x+c)^(3/2)-1/8*a*(A*a^4*b-6*A*a^2*b^3-7*A*b^5+3*B*
a^5+14*B*a^3*b^2+11*B*a*b^4)/b^2*tan(d*x+c)^(1/2))/(a+b*tan(d*x+c))^2+1/8*
(A*a^4*b+18*A*a^2*b^3-15*A*b^5+3*B*a^5+6*B*a^3*b^2+35*B*a*b^4)/b^2/(a*b)^(
1/2)*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2)))+2/(a^2+b^2)^3*(1/8*(-3*A*a^2*
b+A*b^3+B*a^3-3*B*a*b^2)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c
)))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(
1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/8*(-A*a^3+3*A*a*b^2-3*B*a^
2*b+B*b^3)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*
tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arcta
n(-1+2^(1/2)*tan(d*x+c)^(1/2))))
```

3.411.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8913 vs. 2(482) = 964.

Time = 108.21 (sec) , antiderivative size = 17852, normalized size of antiderivative = 33.43

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx = \text{Too large to display}$$

3.411. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm m="fricas")`

output Too large to include

3.411.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)`

output Timed out

3.411.7 Maxima [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 551, normalized size of antiderivative = 1.03

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{(3Ba^6 + Aa^5b + 6Ba^4b^2 + 18Aa^3b^3 + 35Ba^2b^4 - 15Aab^5) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right) - 2\sqrt{2}((A-B)a^3 + 3(A+B)a^2b - 3(A-B)ab^2 - (A+B)b^3) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^6b^2 + 3a^4b^4 + 3a^2b^6 + b^8)\sqrt{ab}}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm m="maxima")`

output $\frac{1}{4} \left((3B^2a^6 + A^2a^5b + 6B^2a^4b^2 + 18A^2a^3b^3 + 35B^2a^2b^4 - 15A^2ab^5) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right) / \left((a^6b^2 + 3a^4b^4 + 3a^2b^6 + b^8) \sqrt{ab} \right) - (2\sqrt{2}((A-B)a^3 + 3(A+B)a^2b - 3(A-B)ab^2 - (A+B)b^3) \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2} + 2\sqrt{\tan(dx+c)})\right) + 2\sqrt{2}((A-B)a^3 + 3(A+B)a^2b - 3(A-B)ab^2 - (A+B)b^3) \arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2} - 2\sqrt{\tan(dx+c)})\right) - \sqrt{2}((A+B)a^3 - 3(A-B)a^2b - 3(A+B)ab^2 + (A-B)b^3) \log(\sqrt{2}\sqrt{\tan(dx+c)} + \tan(dx+c) + 1) + \sqrt{2}((A+B)a^3 - 3(A-B)a^2b - 3(A+B)ab^2 + (A-B)b^3) \log(-\sqrt{2}\sqrt{\tan(dx+c)} + \tan(dx+c) + 1) \right) / (a^6 + 3a^4b^2 + 3a^2b^4 + b^6) - ((5B^2a^4b - A^2a^3b^2 + 13B^2a^2b^3 - 9A^2ab^4) \tan(dx+c)^{3/2} + (3B^2a^5 + A^2a^4b + 11B^2a^3b^2 - 7A^2a^2b^3) \sqrt{\tan(dx+c)}) / (a^6b^2 + 2a^4b^4 + a^2b^6 + (a^4b^4 + 2a^2b^6 + b^8) \tan(dx+c)^2 + 2(a^5b^3 + 2a^3b^5 + ab^7) \tan(dx+c)) \right) / d$

3.411.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{5/2}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="giac")`

output Timed out

3.411.9 Mupad [B] (verification not implemented)

Time = 61.37 (sec) , antiderivative size = 26614, normalized size of antiderivative = 49.84

$$\int \frac{\tan^{5/2}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Too large to display}$$

input `int((tan(c+d*x)^(5/2)*(A+B*tan(c+d*x)))/(a+b*tan(c+d*x))^3,x)`

3.411. $\int \frac{\tan^{5/2}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

output $(\log(\frac{(64Ab^3(11a^2 - 13b^2))/d + 128b^3 \tan(c + dx)^{1/2}}{(a^2 - b^2)(a^2 + b^2)^2((4(-A^4d^4(a^6 - b^6 + 15a^2b^4 - 15a^4b^2))^2)^{1/2} + 80A^2a^3b^3d^2 - 24A^2ab^5d^2 - 24A^2a^5bd^2)/(d^4(a^2 + b^2)^6))^{1/2} + (4(-A^4d^4(a^6 - b^6 + 15a^2b^4 - 15a^4b^2))^2)^{1/2} + 80A^2a^3b^3d^2 - 24A^2ab^5d^2 - 24A^2a^5bd^2)/(d^4(a^2 + b^2)^6))^{1/2})/4 + (8A^2a \tan(c + dx)^{1/2}(a^{10} - 184b^{10} + 833a^2b^8 - 812a^4b^6 + 262a^6b^4 + 44a^8b^2))/(d^2(a^2 + b^2)^4))((4(-A^4d^4(a^6 - b^6 + 15a^2b^4 - 15a^4b^2))^2)^{1/2} + 80A^2a^3b^3d^2 - 24A^2ab^5d^2 - 24A^2a^5bd^2)/(d^4(a^2 + b^2)^6))^{1/2})/4 + (2A^3a^2(5a^{10} - 1199b^{10} + 5017a^2b^8 - 5142a^4b^6 + 1106a^6b^4 + 181a^8b^2))/(d^3(a^2 + b^2)^6))((4(-A^4d^4(a^6 - b^6 + 15a^2b^4 - 15a^4b^2))^2)^{1/2} + 80A^2a^3b^3d^2 - 24A^2ab^5d^2 - 24A^2a^5bd^2)/(d^4(a^2 + b^2)^6))^{1/2})/4 + (A^4 \tan(c + dx)^{1/2}(a^{14} - 32b^{14} + 97a^2b^{12} - 2082a^4b^{10} + 3631a^6b^8 - 2300a^8b^6 + 79a^{10}b^4 + 30a^{12}b^2))/(b^4d^4(a^2 + b^2)^8))((4(-A^4d^4(a^6 - b^6 + 15a^2b^4 - 15a^4b^2))^2)^{1/2} + 80A^2a^3b^3d^2 - 24A^2ab^5d^2 - 24A^2a^5bd^2)/(d^4(a^2 + b^2)^6))^{1/2})/4 + (A^5a(a^{10} - 120b^{10} + 249a^2b^8 - 388a^4b^6 + 302a^6b^4 + 36a^8b^2))/(2b^5d^5(a^2 + b^2)^8))(((480A^4a^2b^{10}d^4 - 16A^4b^{12}d^4 - 16A^4a^{12}d^4 - 4080A^4a^4b^8d^4 + 7232A^4a^6b^6d^4 - 4080A^4a^8...$

3.411. $\int \frac{\tan^5(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$3.412 \quad \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

3.412.1 Optimal result	3966
3.412.2 Mathematica [C] (verified)	3967
3.412.3 Rubi [A] (verified)	3968
3.412.4 Maple [A] (verified)	3975
3.412.5 Fricas [B] (verification not implemented)	3976
3.412.6 Sympy [F]	3976
3.412.7 Maxima [A] (verification not implemented)	3977
3.412.8 Giac [F(-1)]	3977
3.412.9 Mupad [B] (verification not implemented)	3978

3.412.1 Optimal result

Integrand size = 33, antiderivative size = 533

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx =$$

$$\frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d}$$

$$+ \frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d}$$

$$+ \frac{(3a^4Ab - 26a^2Ab^3 + 3Ab^5 + a^5B + 18a^3b^2B - 15ab^4B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{4\sqrt{ab^3/2}(a^2+b^2)^3 d}$$

$$+ \frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d}$$

$$+ \frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d}$$

$$+ \frac{a(Ab - aB)\sqrt{\tan(c+dx)}}{2b(a^2+b^2)d(a+b \tan(c+dx))^2} + \frac{(3a^2Ab - 5Ab^3 + a^3B + 9ab^2B)\sqrt{\tan(c+dx)}}{4b(a^2+b^2)^2d(a+b \tan(c+dx))}$$

$$3.412. \quad \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

output $\frac{1}{2}(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \arctan(-1 + 2^{1/2} \tan(dx+c)^{1/2}) / (a^2+b^2)^{3/2} d^{1/2} + \frac{1}{2}(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \arctan(1 + 2^{1/2} \tan(dx+c)^{1/2}) / (a^2+b^2)^{3/2} d^{1/2} + \frac{1}{4}(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \ln(1 - 2^{1/2} \tan(dx+c)^{1/2} + \tan(dx+c)) / (a^2+b^2)^{3/2} d^{1/2} - \frac{1}{4}(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \ln(1 + 2^{1/2} \tan(dx+c)^{1/2} + \tan(dx+c)) / (a^2+b^2)^{3/2} d^{1/2} + \frac{1}{4}(3Aa^4b - 26Aa^2b^3 + 3Ab^5 + Ba^5 + 18Ba^3b^2 - 15Bab^4) \arctan(b^{1/2} \tan(dx+c)^{1/2} / a^{1/2}) / b^{3/2} / (a^2+b^2)^{3/2} d / a^{1/2} + \frac{1}{2}a(Ab - Ba) \tan(dx+c)^{1/2} / b / (a^2+b^2) / d / (a+b \tan(dx+c))^2 + \frac{1}{4}(3Aa^2b - 5Ab^3 + Ba^3 + 9Bab^2) \tan(dx+c)^{1/2} / b / (a^2+b^2)^2 / d / (a+b \tan(dx+c))$

3.412.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.31 (sec) , antiderivative size = 333, normalized size of antiderivative = 0.62

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{-4B \sqrt{\tan(c+dx)} + \frac{(3aAb+a^2B+4b^2B) \sqrt{\tan(c+dx)}}{a^2+b^2}}{2(a+b \tan(c+dx))} \left(-\frac{3}{4} a^{5/2} \sqrt{b(a^2+b^2)} (3a^2Ab-5Ab^3+a^3B+9ab^2B) \sqrt{\tan(c+dx)} \right)$$

input `Integrate[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3, x]`

output $(-4B \sqrt{\tan[c+dx]} + ((3a^2Ab + a^2B + 4b^2B) \sqrt{\tan[c+dx]})) / (a^2 + b^2) - (2(a + b \tan[c+dx]) * ((-3a^{5/2} \sqrt{b} * (a^2 + b^2) * (3a^2Ab - 5Ab^3 + a^3B + 9a^2b^2B) \sqrt{\tan[c+dx]}) / 4 + ((-3a^2 * (3a^4Ab - 26a^2Ab^3 + 3Ab^5 + a^5B + 18a^3b^2B - 15ab^4B) \operatorname{ArcTan}[(\sqrt{b} \sqrt{\tan[c+dx]}) / \sqrt{a}]) / 4 - 3(-1)^{1/4} * a^{5/2} * b^{3/2} * ((a + I*b)^3 * (A - I*B) * \operatorname{ArcTan}[(-1)^{3/4} \sqrt{\tan[c+dx]}) + (a - I*b)^3 * (A + I*B) * \operatorname{ArcTanh}[(-1)^{3/4} \sqrt{\tan[c+dx]})]) * (a + b \tan[c+dx]))) / (a^{5/2} \sqrt{b} * (a^2 + b^2)^3)) / (6*b*d*(a + b \tan[c+dx])^2)$

3.412. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

3.412.3 Rubi [A] (verified)

Time = 2.04 (sec) , antiderivative size = 462, normalized size of antiderivative = 0.87, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.697$, Rules used = {3042, 4088, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^{3/2}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{4088} \\
 & \frac{\int -\frac{((Ba^2+3Aba+4b^2B) \tan^2(c+dx))-4b(Ab-aB) \tan(c+dx)+a(Ab-aB)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^2} dx}{2b(a^2+b^2)} + \\
 & \quad \frac{a(Ab-aB)\sqrt{\tan(c+dx)}}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{a(Ab-aB)\sqrt{\tan(c+dx)}}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \frac{\int -\frac{((Ba^2+3Aba+4b^2B) \tan^2(c+dx))-4b(Ab-aB) \tan(c+dx)+a(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^2} dx}{4b(a^2+b^2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{a(Ab-aB)\sqrt{\tan(c+dx)}}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \frac{\int -\frac{((Ba^2+3Aba+4b^2B) \tan(c+dx)^2)-4b(Ab-aB) \tan(c+dx)+a(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^2} dx}{4b(a^2+b^2)} \\
 & \quad \downarrow \text{4132} \\
 & \frac{a(Ab-aB)\sqrt{\tan(c+dx)}}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \\
 & \frac{\int -\frac{(Ba^3+3Aba^2+9b^2Ba-5Ab^3) \tan^2(c+dx)-8ab(-Ba^2+2Aba+b^2B) \tan(c+dx)+a(-Ba^3+5Aba^2+7b^2Ba-3Ab^3)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a(a^2+b^2)} - \frac{(a^3B+3a^2Ab+9ab^2B-5Ab^3)}{d(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \downarrow \text{27} \\
 & \frac{a(Ab-aB)\sqrt{\tan(c+dx)}}{2bd(a^2+b^2)(a+b \tan(c+dx))^2} - \frac{(a^3B+3a^2Ab+9ab^2B-5Ab^3)}{d(a^2+b^2)(a+b \tan(c+dx))}
 \end{aligned}$$

3.412. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{\int \frac{a(Ab - aB)\sqrt{\tan(c+dx)}}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} - \frac{-a(Ba^3+3Aba^2+9b^2Ba-5Ab^3)\tan^2(c+dx)-8ab(-Ba^2+2Aba+b^2B)\tan(c+dx)+a(-Ba^3+5Aba^2+7b^2Ba-3Ab^3)}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{2a(a^2+b^2)} - \frac{(a^3B+3a^2Ab+9ab^2B-5Ab^3)\sqrt{\tan(c+dx)}}{d(a^2+b^2)(a+b\tan(c+dx))}}{4b(a^2+b^2)}$$

↓ 3042

$$\frac{\int \frac{a(Ab - aB)\sqrt{\tan(c+dx)}}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} - \frac{-a(Ba^3+3Aba^2+9b^2Ba-5Ab^3)\tan(c+dx)^2-8ab(-Ba^2+2Aba+b^2B)\tan(c+dx)+a(-Ba^3+5Aba^2+7b^2Ba-3Ab^3)}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{2a(a^2+b^2)} - \frac{(a^3B+3a^2Ab+9ab^2B-5Ab^3)\sqrt{\tan(c+dx)}}{d(a^2+b^2)(a+b\tan(c+dx))}}{4b(a^2+b^2)}$$

↓ 4136

$$\frac{\int \frac{a(Ab - aB)\sqrt{\tan(c+dx)}}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} - \frac{8(ab(Aa^3+3bBa^2-3Ab^2a-b^3B)-ab(-Ba^3+3Aba^2+3b^2Ba-Ab^3)\tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{a(a^5B+3a^4Ab+18a^3b^2B-26a^2Ab^3-15ab^4B+3Ab^5)\int \frac{\tan^2(c+dx)}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}}{2a(a^2+b^2)} \frac{4b(a^2+b^2)}{4b(a^2+b^2)}$$

↓ 27

$$\frac{\int \frac{a(Ab - aB)\sqrt{\tan(c+dx)}}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} - \frac{8(ab(Aa^3+3bBa^2-3Ab^2a-b^3B)-ab(-Ba^3+3Aba^2+3b^2Ba-Ab^3)\tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{a(a^5B+3a^4Ab+18a^3b^2B-26a^2Ab^3-15ab^4B+3Ab^5)\int \frac{\tan^2(c+dx)}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}}{2a(a^2+b^2)} \frac{4b(a^2+b^2)}{4b(a^2+b^2)}$$

↓ 3042

$$\frac{\int \frac{a(Ab - aB)\sqrt{\tan(c+dx)}}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} - \frac{8(ab(Aa^3+3bBa^2-3Ab^2a-b^3B)-ab(-Ba^3+3Aba^2+3b^2Ba-Ab^3)\tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{a(a^5B+3a^4Ab+18a^3b^2B-26a^2Ab^3-15ab^4B+3Ab^5)\int \frac{\tan(c+dx)^2}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}}{2a(a^2+b^2)} \frac{4b(a^2+b^2)}{4b(a^2+b^2)}$$

↓ 4017

$$\frac{\int \frac{a(Ab - aB)\sqrt{\tan(c+dx)}}{2bd(a^2+b^2)(a+b\tan(c+dx))^2} - \frac{16\int \frac{ab(Aa^3+3bBa^2-3Ab^2a-b^3B-(-Ba^3+3Aba^2+3b^2Ba-Ab^3)\tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} - \frac{a(a^5B+3a^4Ab+18a^3b^2B-26a^2Ab^3-15ab^4B+3Ab^5)\int \frac{\tan(c+dx)}{\sqrt{\tan(c+dx)(a+b\tan(c+dx))}} dx}{a^2+b^2}}{2a(a^2+b^2)} \frac{4b(a^2+b^2)}{4b(a^2+b^2)}$$

3.412. $\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

$$\begin{aligned} & \downarrow 27 \\ & \frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \\ & \frac{16ab \int \frac{Aa^3 + 3bBa^2 - 3Ab^2a - b^3B - (-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)}}{d(a^2 + b^2)} - \frac{a(a^5B + 3a^4Ab + 18a^3b^2B - 26a^2Ab^3 - 15ab^4B + 3Ab^5) \int \frac{\tan(c + dx)}{\sqrt{\tan(c + dx)}}}{a^2 + b^2} \\ & \hline & \frac{4b(a^2 + b^2)}{2a(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 1482 \\ & \frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \\ & \frac{16ab \left(\frac{1}{2}(a^3(A - B) + 3a^2b(A + B) - 3ab^2(A - B) - b^3(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(-a^3(A + B) + 3a^2b(A - B) + 3ab^2(A + B) - b^3(A - B)) \int \frac{\tan(c + dx)}{\tan^2(c + dx)} \right)}{d(a^2 + b^2)} \\ & \hline & \frac{4b(a^2 + b^2)}{2a(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 1476 \\ & \frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \\ & \frac{16ab \left(\frac{1}{2}(a^3(A - B) + 3a^2b(A + B) - 3ab^2(A - B) - b^3(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(-a^3(A + B) + 3a^2b(A - B) + 3ab^2(A + B) - b^3(A - B)) \left(\frac{1}{2} \int \frac{d \tan(c + dx)}{\tan(c + dx)} \right) \right)}{d(a^2 + b^2)} \\ & \hline & \frac{4b(a^2 + b^2)}{2a(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 1082 \\ & \frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \\ & \frac{16ab \left(\frac{1}{2}(a^3(A - B) + 3a^2b(A + B) - 3ab^2(A - B) - b^3(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(-a^3(A + B) + 3a^2b(A - B) + 3ab^2(A + B) - b^3(A - B)) \left(\int \frac{1}{\tan(c + dx)} \right) \right)}{d(a^2 + b^2)} \\ & \hline & \frac{4b(a^2 + b^2)}{2a(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 217 \\ & \frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \\ & \frac{16ab \left(\frac{1}{2}(a^3(A - B) + 3a^2b(A + B) - 3ab^2(A - B) - b^3(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(-a^3(A + B) + 3a^2b(A - B) + 3ab^2(A + B) - b^3(A - B)) \left(\frac{\arctan(\sqrt{2} \tan(c + dx))}{\sqrt{2}} \right) \right)}{d(a^2 + b^2)} \\ & \hline & \frac{4b(a^2 + b^2)}{2a(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 1479 \\ & \frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{2bd(a^2 + b^2)(a + b \tan(c + dx))^2} - \\ & \frac{16ab \left(\frac{1}{2}(a^3(A - B) + 3a^2b(A + B) - 3ab^2(A - B) - b^3(A + B)) \int \frac{1 - \tan(c + dx)}{\tan^2(c + dx) + 1} d\sqrt{\tan(c + dx)} - \frac{1}{2}(-a^3(A + B) + 3a^2b(A - B) + 3ab^2(A + B) - b^3(A - B)) \left(\frac{\arctan(\sqrt{2} \tan(c + dx))}{\sqrt{2}} \right) \right)}{d(a^2 + b^2)} \\ & \hline & \frac{4b(a^2 + b^2)}{2a(a^2 + b^2)} \end{aligned}$$

3.412. $\int \frac{\tan^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$

$$\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{2bd(a^2 + b^2)(a + b\tan(c + dx))^2} -$$

$$16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)$$

$$d(a^2 + b^2)$$

↓ 25

$$\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{2bd(a^2 + b^2)(a + b\tan(c + dx))^2} -$$

$$16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right) - \frac{1}{2}$$

$$d(a^2 + b^2)$$

↓ 27

$$\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{2bd(a^2 + b^2)(a + b\tan(c + dx))^2} -$$

$$16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right) -$$

$$d(a^2 + b^2)$$

↓ 1103

$$\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{2bd(a^2 + b^2)(a + b\tan(c + dx))^2} -$$

$$16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right) - \frac{1}{2} (- (a^3(A+B)) + 3a^2b$$

$$d(a^2 + b^2)$$

↓ 4117

$$\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{2bd(a^2 + b^2)(a + b\tan(c + dx))^2} -$$

$$16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right) - \frac{1}{2} (- (a^3(A+B)) + 3a^2b$$

$$d(a^2 + b^2)$$

↓ 73

3.412. $\int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

$$\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{2bd(a^2 + b^2)(a + b\tan(c + dx))^2} - \frac{16ab \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right) - \frac{1}{2}(-a^3(A+B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B))}{d(a^2 + b^2)}$$

↓ 218

$$\frac{a(Ab - aB)\sqrt{\tan(c + dx)}}{2bd(a^2 + b^2)(a + b\tan(c + dx))^2} - \frac{16ab \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\log(\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right) - \frac{1}{2}(-a^3(A+B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B))}{d(a^2 + b^2)}$$

```
input Int[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]
```

```
output (a*(A*b - a*B)*Sqrt[Tan[c + d*x]]/(2*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) - (((-2*Sqrt[a]*(3*a^4*A*b - 26*a^2*A*b^3 + 3*A*b^5 + a^5*B + 18*a^3*b^2*B - 15*a*b^4*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[b]*(a^2 + b^2)*d) + (16*a*b*(-1/2*((3*a^2*b*(A - B) - b^3*(A - B) - a^3*(A + B) + 3*a*b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])) + ((a^3*(A - B) - 3*a*b^2*(A - B) + 3*a^2*b*(A + B) - b^3*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)/(2*a*(a^2 + b^2)) - ((3*a^2*A*b - 5*A*b^3 + a^3*B + 9*a*b^2*B)*Sqrt[Tan[c + d*x]])/((a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(4*b*(a^2 + b^2))
```

3.412.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

3.412. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
 -1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
 & (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
 t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
 implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/R
 t[a/b, 2])], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
 eQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
 imp[d*(Log[RemoveContent[a + b*x + c*x^2, x])/b], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
 2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
 e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
 && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
 -2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
 x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
 reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.412.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 450, normalized size of antiderivative = 0.84

method	result
derivativedivides	$\frac{2 \left(\left(\frac{3}{8} A a^4 b - \frac{1}{4} A a^2 b^3 - \frac{5}{8} A b^5 + \frac{1}{8} B a^5 + \frac{5}{4} B a^3 b^2 + \frac{9}{8} B a b^4 \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) + \frac{a \left(5 A a^4 b + 2 A a^2 b^3 - 3 A b^5 - B a^5 + 6 B a^3 b^2 + 7 B a b^4 \right)}{8 b}}{(a+b \tan(dx+c))^2} \frac{1}{(a^2+b^2)^3}$
default	$\frac{2 \left(\left(\frac{3}{8} A a^4 b - \frac{1}{4} A a^2 b^3 - \frac{5}{8} A b^5 + \frac{1}{8} B a^5 + \frac{5}{4} B a^3 b^2 + \frac{9}{8} B a b^4 \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) + \frac{a \left(5 A a^4 b + 2 A a^2 b^3 - 3 A b^5 - B a^5 + 6 B a^3 b^2 + 7 B a b^4 \right)}{8 b}}{(a+b \tan(dx+c))^2} \frac{1}{(a^2+b^2)^3}$

```
input int(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x,method=_RETURNV
ERBOSE)
```

$$3.412. \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

```
output 1/d*(2/(a^2+b^2)^3*((3/8*A*a^4*b-1/4*A*a^2*b^3-5/8*A*b^5+1/8*B*a^5+5/4*B*
a^3*b^2+9/8*B*a*b^4)*tan(d*x+c)^(3/2)+1/8*a*(5*A*a^4*b+2*A*a^2*b^3-3*A*b^5
-B*a^5+6*B*a^3*b^2+7*B*a*b^4)/b*tan(d*x+c)^(1/2))/(a+b*tan(d*x+c))^2+1/8*(
3*A*a^4*b-26*A*a^2*b^3+3*A*b^5+B*a^5+18*B*a^3*b^2-15*B*a*b^4)/b/(a*b)^(1/2
)*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))+2/(a^2+b^2)^3*(1/8*(-A*a^3+3*A*a
*b^2-3*B*a^2*b+B*b^3)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/
(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/
2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/8*(3*A*a^2*b-A*b^3-B*a^3+3*B*
a*b^2)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(
d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1
+2^(1/2)*tan(d*x+c)^(1/2))))
```

3.412.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8932 vs. 2(481) = 962.

Time = 75.16 (sec) , antiderivative size = 17891, normalized size of antiderivative = 33.57

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorith
m="fricas")
```

```
output Too large to include
```

3.412.6 Sympy [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \int \frac{(A+B\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)}{(a+b\tan(c+dx))^3} dx$$

```
input integrate(tan(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)
```

```
output Integral((A + B*tan(c + d*x))*tan(c + d*x)**(3/2)/(a + b*tan(c + d*x))**3,
x)
```

3.412. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

3.412.7 Maxima [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 538, normalized size of antiderivative = 1.01

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{(Ba^5+3Aa^4b+18Ba^3b^2-26Aa^2b^3-15Bab^4+3Ab^5) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right) - 2\sqrt{2}((A+B)a^3-3(A-B)a^2b-3(A+B)ab^2+(A-B)b^3) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right) - \frac{2\sqrt{2}((A+B)a^3-3(A-B)a^2b-3(A+B)ab^2+(A-B)b^3)}{(a^6b+3a^4b^3+3a^2b^5+b^7)\sqrt{ab}}}{(a^6b+3a^4b^3+3a^2b^5+b^7)\sqrt{ab}}$$

```
input integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm
m="maxima")
```

```
output 1/4*((B*a^5 + 3*A*a^4*b + 18*B*a^3*b^2 - 26*A*a^2*b^3 - 15*B*a*b^4 + 3*A*b
^5)*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/((a^6*b + 3*a^4*b^3 + 3*a^2*b^5
+ b^7)*sqrt(a*b)) - (2*sqrt(2)*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)
*a*b^2 + (A - B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c))))
+ 2*sqrt(2)*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b
^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + sqrt(2)*((A - B)
)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*log(sqrt(2)*sqrt(
tan(d*x + c)) + tan(d*x + c) + 1) - sqrt(2)*((A - B)*a^3 + 3*(A + B)*a^2*b
- 3*(A - B)*a*b^2 - (A + B)*b^3)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*
x + c) + 1))/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + ((B*a^3*b + 3*A*a^2*b^2
+ 9*B*a*b^3 - 5*A*b^4)*tan(d*x + c)^(3/2) - (B*a^4 - 5*A*a^3*b - 7*B*a^2*
b^2 + 3*A*a*b^3)*sqrt(tan(d*x + c)))/(a^6*b + 2*a^4*b^3 + a^2*b^5 + (a^4*b
^3 + 2*a^2*b^5 + b^7)*tan(d*x + c)^2 + 2*(a^5*b^2 + 2*a^3*b^4 + a*b^6)*tan
(d*x + c))/d
```

3.412.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Timed out}$$

```
input integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm
m="giac")
```

```
output Timed out
```

3.412. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

3.412.9 Mupad [B] (verification not implemented)

Time = 62.64 (sec) , antiderivative size = 25944, normalized size of antiderivative = 48.68

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Too large to display}$$

```
input int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3,x)
```

```
output (log((((((((((64*B*a*b^3*(11*a^2 - 13*b^2))/d + 128*b^3*tan(c + d*x)^(1/2)
*(a^2 - b^2)*(a^2 + b^2)^2*((4*(-B^4*d^4*(a^6 - b^6 + 15*a^2*b^4 - 15*a^4*
b^2)^2)^(1/2) + 80*B^2*a^3*b^3*d^2 - 24*B^2*a*b^5*d^2 - 24*B^2*a^5*b*d^2)/
(d^4*(a^2 + b^2)^6))^(1/2))*((4*(-B^4*d^4*(a^6 - b^6 + 15*a^2*b^4 - 15*a^4*
*b^2)^2)^(1/2) + 80*B^2*a^3*b^3*d^2 - 24*B^2*a*b^5*d^2 - 24*B^2*a^5*b*d^2)
/(d^4*(a^2 + b^2)^6))^(1/2))/4 + (8*B^2*a*tan(c + d*x)^(1/2)*(a^10 - 184*b
^10 + 833*a^2*b^8 - 812*a^4*b^6 + 262*a^6*b^4 + 44*a^8*b^2))/(d^2*(a^2 + b
^2)^4))*((4*(-B^4*d^4*(a^6 - b^6 + 15*a^2*b^4 - 15*a^4*b^2)^2)^(1/2) + 80*
B^2*a^3*b^3*d^2 - 24*B^2*a*b^5*d^2 - 24*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^6)
)^(1/2))/4 + (2*B^3*a^2*(5*a^10 - 1199*b^10 + 5017*a^2*b^8 - 5142*a^4*b^6
+ 1106*a^6*b^4 + 181*a^8*b^2))/(d^3*(a^2 + b^2)^6))*((4*(-B^4*d^4*(a^6 - b
^6 + 15*a^2*b^4 - 15*a^4*b^2)^2)^(1/2) + 80*B^2*a^3*b^3*d^2 - 24*B^2*a*b^5
*d^2 - 24*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^6))^(1/2))/4 + (B^4*tan(c + d*x)
^(1/2)*(a^14 - 32*b^14 + 97*a^2*b^12 - 2082*a^4*b^10 + 3631*a^6*b^8 - 2300
*a^8*b^6 + 79*a^10*b^4 + 30*a^12*b^2))/(b*d^4*(a^2 + b^2)^8))*((4*(-B^4*d^
4*(a^6 - b^6 + 15*a^2*b^4 - 15*a^4*b^2)^2)^(1/2) + 80*B^2*a^3*b^3*d^2 - 24
*B^2*a*b^5*d^2 - 24*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^6))^(1/2))/4 + (B^5*a*
(a^10 - 120*b^10 + 249*a^2*b^8 - 388*a^4*b^6 + 302*a^6*b^4 + 36*a^8*b^2))/
(2*b*d^5*(a^2 + b^2)^8))*(((480*B^4*a^2*b^10*d^4 - 16*B^4*b^12*d^4 - 16*B^
4*a^12*d^4 - 4080*B^4*a^4*b^8*d^4 + 7232*B^4*a^6*b^6*d^4 - 4080*B^4*a^8...
```

3.413
$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

3.413.1 Optimal result	3979
3.413.2 Mathematica [C] (verified)	3980
3.413.3 Rubi [A] (verified)	3981
3.413.4 Maple [A] (verified)	3989
3.413.5 Fricas [B] (verification not implemented)	3989
3.413.6 Sympy [F]	3990
3.413.7 Maxima [A] (verification not implemented)	3990
3.413.8 Giac [F(-1)]	3991
3.413.9 Mupad [B] (verification not implemented)	3991

3.413.1 Optimal result

Integrand size = 33, antiderivative size = 531

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx =$$

$$\frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d}$$

$$+ \frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d}$$

$$- \frac{(15a^4Ab - 18a^2Ab^3 - Ab^5 - 3a^5B + 26a^3b^2B - 3ab^4B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{4a^{3/2}\sqrt{b}(a^2+b^2)^3 d}$$

$$- \frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d}$$

$$+ \frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d}$$

$$- \frac{(Ab - aB)\sqrt{\tan(c+dx)}}{2(a^2+b^2)d(a+b \tan(c+dx))^2} - \frac{(7a^2Ab - Ab^3 - 3a^3B + 5ab^2B)\sqrt{\tan(c+dx)}}{4a(a^2+b^2)^2 d(a+b \tan(c+dx))}$$

output $\frac{1}{2}(a^3(A-B)-3ab^2(A-B)+3a^2b(A+B)-b^3(A+B))\arctan(-1+2^{1/2}\tan(dx+c)^{1/2})/(a^2+b^2)^{3/2}d^{1/2}+1/2(a^3(A-B)-3ab^2(A-B)+3a^2b(A+B)-b^3(A+B))\arctan(1+2^{1/2}\tan(dx+c)^{1/2})/(a^2+b^2)^{3/2}d^{1/2}-1/4(3a^2b(A-B)-b^3(A-B)-a^3(A+B)+3ab^2(A+B))\ln(1-2^{1/2}\tan(dx+c)^{1/2}+\tan(dx+c))/(a^2+b^2)^{3/2}d^{1/2}+1/4(3a^2b(A-B)-b^3(A-B)-a^3(A+B)+3ab^2(A+B))\ln(1+2^{1/2}\tan(dx+c)^{1/2}+\tan(dx+c))/(a^2+b^2)^{3/2}d^{1/2}-1/4(15Aa^4b-18Aa^2b^3-A^2b^5-3Ba^5+26Ba^3b^2-3B^2a^2b^4)\arctan(b^{1/2}\tan(dx+c)^{1/2}/a^{1/2})/a^{3/2}/(a^2+b^2)^{3/2}d/b^{1/2}-1/2(Ab-Ba)\tan(dx+c)^{1/2}/(a^2+b^2)/d/(a+b\tan(dx+c))^2-1/4(7Aa^2b-A^2b^3-3Ba^3+5B^2a^2b)\tan(dx+c)^{1/2}/a/(a^2+b^2)^{2/2}d/(a+b\tan(dx+c))$

3.413.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.33 (sec) , antiderivative size = 552, normalized size of antiderivative = 1.04

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \frac{b(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{2a(a^2+b^2)d(a+b\tan(c+dx))^2} + \frac{\left(\frac{2(a^3b^2(a^2A-Ab^2+2abB)-\frac{1}{8}ab^3(9a^2Ab+Ab^3-5a^3B+3ab^2B))+\frac{1}{8}a^3b(7a^2Ab-Ab^3-3a^3B+5ab^2B)}{\sqrt{a}\sqrt{b}(a^2+b^2)d} \right) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right) - \frac{(Ab-aB)\sqrt{\tan(c+dx)}}{d(a+b\tan(c+dx))}}{2}$$

input `Integrate[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3, x]`

output

```
(b*(A*b - a*B)*Tan[c + d*x]^(3/2))/(2*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])
^2) + (-(((A*b - a*B)*Sqrt[Tan[c + d*x]])/(d*(a + b*Tan[c + d*x]))) - (2*(
((2*(a^3*b^2*(a^2*A - A*b^2 + 2*a*b*B) - (a*b^3*(9*a^2*A*b + A*b^3 - 5*a^3
*B + 3*a*b^2*B))/8 + (a^3*b*(7*a^2*A*b - A*b^3 - 3*a^3*B + 5*a*b^2*B))/8)*
ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[a]*Sqrt[b]*(a^2 + b^2)
*d) + (-((( -1)^(1/4)*(-a^2*b*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B)) + I
*a^2*b*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B))*ArcTan[(-1)^(3/4)*Sqrt[Tan
[c + d*x]]])/d) - ((( -1)^(1/4)*(-a^2*b*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^
2*B)) - I*a^2*b*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B))*ArcTanh[(-1)^(3/4)
)*Sqrt[Tan[c + d*x]]])/d)/(a^2 + b^2))/(a*(a^2 + b^2)) + ((-1/4*(a*b^3*(A*
b - a*B)) - a*(-3*a^2*b*(A*b - a*B))/4 - a*b^2*(a*A + b*B))*Sqrt[Tan[c +
d*x]])/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])))/b)/(2*a*(a^2 + b^2))
```

3.413.3 Rubi [A] (verified)

Time = 2.08 (sec) , antiderivative size = 464, normalized size of antiderivative = 0.87, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.697$, Rules used = {3042, 4091, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{4091} \\
 & -\frac{\int -\frac{3b(Ab-aB) \tan^2(c+dx)+4b(aA+bB) \tan(c+dx)+b(Ab-aB)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^2} dx}{2b(a^2+b^2)} - \frac{(Ab-aB)\sqrt{\tan(c+dx)}}{2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int -\frac{3b(Ab-aB) \tan^2(c+dx)+4b(aA+bB) \tan(c+dx)+b(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^2} dx}{4b(a^2+b^2)} - \frac{(Ab-aB)\sqrt{\tan(c+dx)}}{2d(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.413. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{\int \frac{-3b(Ab-aB)\tan(c+dx)^2+4b(aA+bB)\tan(c+dx)+b(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2} dx}{4b(a^2+b^2)} - \frac{(Ab-aB)\sqrt{\tan(c+dx)}}{2d(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 4132

$$\frac{\int \frac{-b(-3Ba^3+7Aba^2+5b^2Ba-Ab^3)\tan^2(c+dx)+8ab(Aa^2+2bBa-Ab^2)\tan(c+dx)+b(-5Ba^3+9Aba^2+3b^2Ba+Ab^3)}{2\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a(a^2+b^2)} - \frac{b(-3a^3B+7a^2Ab+5ab^2B-Ab^3)}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

$$\frac{4b(a^2+b^2)}{2d(a^2+b^2)(a+b\tan(c+dx))^2} \frac{(Ab-aB)\sqrt{\tan(c+dx)}}{2d(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 27

$$\frac{\int \frac{-b(-3Ba^3+7Aba^2+5b^2Ba-Ab^3)\tan^2(c+dx)+8ab(Aa^2+2bBa-Ab^2)\tan(c+dx)+b(-5Ba^3+9Aba^2+3b^2Ba+Ab^3)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{2a(a^2+b^2)} - \frac{b(-3a^3B+7a^2Ab+5ab^2B-Ab^3)}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

$$\frac{4b(a^2+b^2)}{2d(a^2+b^2)(a+b\tan(c+dx))^2} \frac{(Ab-aB)\sqrt{\tan(c+dx)}}{2d(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 3042

$$\frac{\int \frac{-b(-3Ba^3+7Aba^2+5b^2Ba-Ab^3)\tan(c+dx)^2+8ab(Aa^2+2bBa-Ab^2)\tan(c+dx)+b(-5Ba^3+9Aba^2+3b^2Ba+Ab^3)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{2a(a^2+b^2)} - \frac{b(-3a^3B+7a^2Ab+5ab^2B-Ab^3)}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

$$\frac{4b(a^2+b^2)}{2d(a^2+b^2)(a+b\tan(c+dx))^2} \frac{(Ab-aB)\sqrt{\tan(c+dx)}}{2d(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 4136

$$\frac{\int \frac{8(ab(-Ba^3+3Aba^2+3b^2Ba-Ab^3)+ab(Aa^3+3bBa^2-3Ab^2a-b^3B)\tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{b(-3a^5B+15a^4Ab+26a^3b^2B-18a^2Ab^3-3ab^4B-Ab^5)}{a^2+b^2} \int \frac{\tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{2a(a^2+b^2)}$$

$$\frac{4b(a^2+b^2)}{2d(a^2+b^2)(a+b\tan(c+dx))^2} \frac{(Ab-aB)\sqrt{\tan(c+dx)}}{2d(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 27

$$\frac{8 \int \frac{ab(-Ba^3+3Aba^2+3b^2Ba-Ab^3)+ab(Aa^3+3bBa^2-3Ab^2a-b^3B)\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{b(-3a^5B+15a^4Ab+26a^3b^2B-18a^2Ab^3-3ab^4B-Ab^5)}{a^2+b^2} \int \frac{\tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{2a(a^2+b^2)}$$

$$\frac{4b(a^2+b^2)}{2d(a^2+b^2)(a+b\tan(c+dx))^2} \frac{(Ab-aB)\sqrt{\tan(c+dx)}}{2d(a^2+b^2)(a+b\tan(c+dx))^2}$$

3.413. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

↓ 3042

$$8 \int \frac{ab(-Ba^3+3Aba^2+3b^2Ba-Ab^3)+ab(Aa^3+3bBa^2-3Ab^2a-b^3B)\tan(c+dx)}{\sqrt{\tan(c+dx)}(a^2+b^2)} dx - \frac{b(-3a^5B+15a^4Ab+26a^3b^2B-18a^2Ab^3-3ab^4B-Ab^5)}{a^2+b^2} \int \frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b)}$$

$$\frac{(Ab - aB)\sqrt{\tan(c + dx)}}{2d(a^2 + b^2)(a + b \tan(c + dx))^2}$$

4b(a² + b²)

↓ 4017

$$16 \int \frac{ab(-Ba^3+3Aba^2+3b^2Ba-Ab^3+(Aa^3+3bBa^2-3Ab^2a-b^3B)\tan(c+dx))d\sqrt{\tan(c+dx)}}{\tan^2(c+dx)+1} - \frac{b(-3a^5B+15a^4Ab+26a^3b^2B-18a^2Ab^3-3ab^4B-Ab^5)}{a^2+b^2} \int \frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}}$$

$$\frac{(Ab - aB)\sqrt{\tan(c + dx)}}{2d(a^2 + b^2)(a + b \tan(c + dx))^2}$$

4b(a² + b²)

↓ 27

$$16ab \int \frac{-Ba^3+3Aba^2+3b^2Ba-Ab^3+(Aa^3+3bBa^2-3Ab^2a-b^3B)\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{b(-3a^5B+15a^4Ab+26a^3b^2B-18a^2Ab^3-3ab^4B-Ab^5)}{a^2+b^2} \int \frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}}$$

$$\frac{(Ab - aB)\sqrt{\tan(c + dx)}}{2d(a^2 + b^2)(a + b \tan(c + dx))^2}$$

4b(a² + b²)

↓ 1482

$$16ab \left(\frac{1}{2}(-a^3(A+B)+3a^2b(A-B)+3ab^2(A+B)-b^3(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)} \right)$$

$$\frac{(Ab - aB)\sqrt{\tan(c + dx)}}{2d(a^2 + b^2)(a + b \tan(c + dx))^2}$$

4b(a² + b²)

↓ 1476

$$16ab \left(\frac{1}{2}(-a^3(A+B)+3a^2b(A-B)+3ab^2(A+B)-b^3(A-B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \left(\frac{1}{2} \int \frac{\tan(c+dx)+1}{\tan(c+dx)} \right) \right)$$

$$\frac{(Ab - aB)\sqrt{\tan(c + dx)}}{2d(a^2 + b^2)(a + b \tan(c + dx))^2}$$

3.413. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

↓ 1082

$$\frac{16ab \left(\frac{1}{2} (- (a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\int \frac{1}{-\tan(c+dx)} \right) \right)}{d(a^2+b^2)}$$

$$\frac{(Ab - aB)\sqrt{\tan(c + dx)}}{2d(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 217

$$\frac{16ab \left(\frac{1}{2} (- (a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)}$$

$$\frac{(Ab - aB)\sqrt{\tan(c + dx)}}{2d(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 1479

$$16ab \left(\frac{1}{2} (- (a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \left(- \frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)$$

$$\frac{(Ab - aB)\sqrt{\tan(c + dx)}}{2d(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 25

$$16ab \left(\frac{1}{2} (- (a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)$$

$$\frac{(Ab - aB)\sqrt{\tan(c + dx)}}{2d(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 27

3.413. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$16ab \left(\frac{1}{2} (-a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B) \right) \left(\int \frac{\sqrt{2-2\sqrt{\tan(c+dx)}}}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)} + 1}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)} + 1} d\sqrt{\tan(c+dx)} \right)$$

$$d(a^2 + b^2)$$

$$\frac{(Ab - aB)\sqrt{\tan(c+dx)}}{2d(a^2 + b^2)(a + b\tan(c+dx))^2}$$

↓ 1103

$$16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) + \frac{1}{2} (-a^3(A+B) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B))$$

$$d(a^2 + b^2)$$

$$\frac{(Ab - aB)\sqrt{\tan(c+dx)}}{2d(a^2 + b^2)(a + b\tan(c+dx))^2}$$

↓ 4117

$$16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) + \frac{1}{2} (-a^3(A+B) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B))$$

$$d(a^2 + b^2)$$

$$\frac{(Ab - aB)\sqrt{\tan(c+dx)}}{2d(a^2 + b^2)(a + b\tan(c+dx))^2}$$

↓ 73

$$16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) + \frac{1}{2} (-a^3(A+B) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B))$$

$$d(a^2 + b^2)$$

$$\frac{(Ab - aB)\sqrt{\tan(c+dx)}}{2d(a^2 + b^2)(a + b\tan(c+dx))^2}$$

↓ 218

$$16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) + \frac{1}{2} (-a^3(A+B) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B))$$

$$d(a^2 + b^2)$$

$$\frac{(Ab - aB)\sqrt{\tan(c+dx)}}{2d(a^2 + b^2)(a + b\tan(c+dx))^2}$$

3.413. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

input `Int[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

output `-1/2*((A*b - a*B)*Sqrt[Tan[c + d*x]])/((a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + (((-2*Sqrt[b]*(15*a^4*A*b - 18*a^2*A*b^3 - A*b^5 - 3*a^5*B + 26*a^3*b^2*B - 3*a*b^4*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[a]*(a^2 + b^2)*d) + (16*a*b*((a^3*(A - B) - 3*a*b^2*(A - B) + 3*a^2*b*(A + B) - b^3*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]))/2 + ((3*a^2*b*(A - B) - b^3*(A - B) - a^3*(A + B) + 3*a*b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)/(2*a*(a^2 + b^2)) - (b*(7*a^2*A*b - A*b^3 - 3*a^3*B + 5*a*b^2*B)*Sqrt[Tan[c + d*x]])/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))/(4*b*(a^2 + b^2))`

3.413.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])]; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4091 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.413.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 451, normalized size of antiderivative = 0.85

method	result
derivativedivides	$\frac{2 \left(\frac{b(7Aa^4b+6Aa^2b^3-Ab^5-3Ba^5+2Ba^3b^2+5Ba^4)}{8a} \left(\tan^{\frac{3}{2}}(dx+c) \right) + \left(\frac{9}{8}Aa^4b + \frac{5}{4}Aa^2b^3 - \frac{5}{8}Ba^5 + \frac{3}{8}Ba^4 + \frac{1}{8}Ab^5 - \frac{1}{4}Ba^3b^2 \right) \right)}{(a+b \tan(dx+c))^2} \frac{1}{(a^2+b^2)^3}$
default	$\frac{2 \left(\frac{b(7Aa^4b+6Aa^2b^3-Ab^5-3Ba^5+2Ba^3b^2+5Ba^4)}{8a} \left(\tan^{\frac{3}{2}}(dx+c) \right) + \left(\frac{9}{8}Aa^4b + \frac{5}{4}Aa^2b^3 - \frac{5}{8}Ba^5 + \frac{3}{8}Ba^4 + \frac{1}{8}Ab^5 - \frac{1}{4}Ba^3b^2 \right) \right)}{(a+b \tan(dx+c))^2} \frac{1}{(a^2+b^2)^3}$

input `int(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x,method=_RETURNV
ERBOSE)`

output `1/d*(-2/(a^2+b^2)^3*((1/8*b*(7*A*a^4*b+6*A*a^2*b^3-A*b^5-3*B*a^5+2*B*a^3*b
^2+5*B*a*b^4)/a*tan(d*x+c)^(3/2)+(9/8*A*a^4*b+5/4*A*a^2*b^3-5/8*B*a^5+3/8*
B*a*b^4+1/8*A*b^5-1/4*B*a^3*b^2)*tan(d*x+c)^(1/2))/(a+b*tan(d*x+c))^2+1/8*
(15*A*a^4*b-18*A*a^2*b^3-A*b^5-3*B*a^5+26*B*a^3*b^2-3*B*a*b^4)/a/(a*b)^(1/
2)*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2)))+2/(a^2+b^2)^3*(1/8*(3*A*a^2*b-A
*b^3-B*a^3+3*B*a*b^2)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/
(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/
2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/8*(A*a^3-3*A*a*b^2+3*B*a^2*b-
B*b^3)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(
d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1
+2^(1/2)*tan(d*x+c)^(1/2))))`

3.413.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8933 vs. 2(477) = 954.

Time = 81.21 (sec) , antiderivative size = 17893, normalized size of antiderivative = 33.70

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm m="fricas")`

output Too large to include

3.413.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \int \frac{(A+B\tan(c+dx))\sqrt{\tan(c+dx)}}{(a+b\tan(c+dx))^3} dx$$

input `integrate(tan(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)`

output `Integral((A + B*tan(c + d*x))*sqrt(tan(c + d*x))/(a + b*tan(c + d*x))**3, x)`

3.413.7 Maxima [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 537, normalized size of antiderivative = 1.01

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{(3Ba^5 - 15Aa^4b - 26Ba^3b^2 + 18Aa^2b^3 + 3Bab^4 + Ab^5) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^7 + 3a^5b^2 + 3a^3b^4 + ab^6)\sqrt{ab}} + \frac{2\sqrt{2}((A-B)a^3 + 3(A+B)a^2b - 3(A-B)ab^2 - (A+B)b^3) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^7 + 3a^5b^2 + 3a^3b^4 + ab^6)\sqrt{ab}}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm m="maxima")`

output $\frac{1}{4}((3B*a^5 - 15A*a^4*b - 26B*a^3*b^2 + 18A*a^2*b^3 + 3B*a*b^4 + A*b^5)*\arctan(b*\sqrt{\tan(dx + c)}/\sqrt{a*b})/((a^7 + 3a^5*b^2 + 3a^3*b^4 + a*b^6)*\sqrt{a*b}) + (2*\sqrt{2}*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2*\sqrt{\tan(dx + c)})) + 2*\sqrt{2}*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2*\sqrt{\tan(dx + c)})) - \sqrt{2}*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*\log(\sqrt{2}*\sqrt{\tan(dx + c)} + \tan(dx + c) + 1) + \sqrt{2}*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*\log(-\sqrt{2}*\sqrt{\tan(dx + c)} + \tan(dx + c) + 1))/(a^6 + 3a^4*b^2 + 3a^2*b^4 + b^6) + ((3B*a^3*b - 7A*a^2*b^2 - 5B*a*b^3 + A*b^4)*\tan(dx + c)^{(3/2)} + (5B*a^4 - 9A*a^3*b - 3B*a^2*b^2 - A*a*b^3)*\sqrt{\tan(dx + c)})/(a^7 + 2a^5*b^2 + a^3*b^4 + (a^5*b^2 + 2a^3*b^4 + a*b^6)*\tan(dx + c)^2 + 2*(a^6*b + 2a^4*b^3 + a^2*b^5)*\tan(dx + c))/d$

3.413.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{\tan(c + dx)}(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate(tan(dx+c)^(1/2)*(A+B*tan(dx+c))/(a+b*tan(dx+c))^3,x, algorithm="giac")`

output Timed out

3.413.9 Mupad [B] (verification not implemented)

Time = 61.16 (sec) , antiderivative size = 26133, normalized size of antiderivative = 49.21

$$\int \frac{\sqrt{\tan(c + dx)}(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

input `int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3,x)`

output $(\log(\frac{(((((64Ab^3(b^4 - 10a^4 + 15a^2b^2)))/(ad) + 128b^3 \tan(c + dx))^{1/2} (a^2 - b^2)(a^2 + b^2)^2 ((4(-A^4d^4(a^6 - b^6 + 15a^2b^4 - 15a^4b^2))^2)^{1/2} + 80A^2a^3b^3d^2 - 24A^2ab^5d^2 - 24A^2a^5bd^2)/(d^4(a^2 + b^2)^6))^{1/2} + 80A^2a^3b^3d^2 - 24A^2ab^5d^2 - 24A^2a^5bd^2)/(d^4(a^2 + b^2)^6))^{1/2} + 80A^2a^3b^3d^2 - 24A^2ab^5d^2 - 24A^2a^5bd^2)/(d^4(a^2 + b^2)^6))^{1/2})/4 + (8A^2b^2 \tan(c + dx))^{1/2} * (8a^{10} + b^{10} - 148a^2b^8 + 902a^4b^6 - 812a^6b^4 + 193a^8b^2))/(a^2d^2(a^2 + b^2)^4) * ((4(-A^4d^4(a^6 - b^6 + 15a^2b^4 - 15a^4b^2))^2)^{1/2} + 80A^2a^3b^3d^2 - 24A^2ab^5d^2 - 24A^2a^5bd^2)/(d^4(a^2 + b^2)^6))^{1/2})/4 - (2A^3b^2(16a^{12} + b^{12} - 71a^2b^{10} - 1382a^4b^8 + 5266a^6b^6 - 4539a^8b^4 + 1189a^{10}b^2))/(a^2d^3(a^2 + b^2)^6) * ((4(-A^4d^4(a^6 - b^6 + 15a^2b^4 - 15a^4b^2))^2)^{1/2} + 80A^2a^3b^3d^2 - 24A^2ab^5d^2 - 24A^2a^5bd^2)/(d^4(a^2 + b^2)^6))^{1/2})/4 - (A^4b^3 \tan(c + dx))^{1/2} * (2a^2b^{10} - b^{12} - 225a^{12} + 49a^4b^8 + 2460a^6b^6 - 3631a^8b^4 + 1922a^{10}b^2))/(a^2d^4(a^2 + b^2)^8) * ((4(-A^4d^4(a^6 - b^6 + 15a^2b^4 - 15a^4b^2))^2)^{1/2} + 80A^2a^3b^3d^2 - 24A^2ab^5d^2 - 24A^2a^5bd^2)/(d^4(a^2 + b^2)^6))^{1/2})/4 + (A^5b^3(7b^8 - 225a^8 + 116a^2b^6 - 270a^4b^4 + 420a^6b^2))/(2a^5d^5(a^2 + b^2)^8) * (((480A^4a^2b^{10}d^4 - 16A^4b^{12}d^4 - 16A^4a^{12}d^4 - 4080A^4a^4b^8d^4 + 7232A^4a^6b^6d^4 - 4...$

3.413. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$3.414 \quad \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^3} dx$$

3.414.1 Optimal result	3993
3.414.2 Mathematica [C] (verified)	3994
3.414.3 Rubi [A] (verified)	3995
3.414.4 Maple [A] (verified)	4003
3.414.5 Fricas [B] (verification not implemented)	4003
3.414.6 Sympy [F(-1)]	4004
3.414.7 Maxima [A] (verification not implemented)	4004
3.414.8 Giac [F(-1)]	4005
3.414.9 Mupad [B] (verification not implemented)	4005

3.414.1 Optimal result

Integrand size = 33, antiderivative size = 534

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3} dx \\ &= \frac{(3a^2b(A - B) - b^3(A - B) - a^3(A + B) + 3ab^2(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^3 d} \\ & \quad - \frac{(3a^2b(A - B) - b^3(A - B) - a^3(A + B) + 3ab^2(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^3 d} \\ & \quad + \frac{\sqrt{b}(35a^4 Ab + 6a^2 Ab^3 + 3Ab^5 - 15a^5 B + 18a^3 b^2 B + ab^4 B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{4a^{5/2}(a^2 + b^2)^3 d} \\ & \quad - \frac{(a^3(A - B) - 3ab^2(A - B) + 3a^2b(A + B) - b^3(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^3 d} \\ & \quad + \frac{(a^3(A - B) - 3ab^2(A - B) + 3a^2b(A + B) - b^3(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^3 d} \\ & \quad + \frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{2a(a^2 + b^2)d(a + b \tan(c + dx))^2} + \frac{b(11a^2 Ab + 3Ab^3 - 7a^3 B + ab^2 B)\sqrt{\tan(c + dx)}}{4a^2(a^2 + b^2)^2 d(a + b \tan(c + dx))} \end{aligned}$$

output
$$-1/2*(3*a^2*b*(A-B)-b^3*(A-B)-a^3*(A+B)+3*a*b^2*(A+B))*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/(a^2+b^2)^3/d*2^{(1/2)}-1/2*(3*a^2*b*(A-B)-b^3*(A-B)-a^3*(A+B)+3*a*b^2*(A+B))*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/(a^2+b^2)^3/d*2^{(1/2)}-1/4*(a^3*(A-B)-3*a*b^2*(A-B)+3*a^2*b*(A+B)-b^3*(A+B))*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(a^2+b^2)^3/d*2^{(1/2)}+1/4*(a^3*(A-B)-3*a*b^2*(A-B)+3*a^2*b*(A+B)-b^3*(A+B))*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(a^2+b^2)^3/d*2^{(1/2)}+1/4*(35*A*a^4*b+6*A*a^2*b^3+3*A*b^5-15*B*a^5+18*B*a^3*b^2+B*a*b^4)*\arctan(b^{(1/2)}*\tan(d*x+c)^{(1/2)}/a^{(1/2)})*b^{(1/2)}/a^{(5/2)}/(a^2+b^2)^3/d+1/2*b*(A*b-B*a)*\tan(d*x+c)^{(1/2)}/a/(a^2+b^2)/d/(a+b*\tan(d*x+c))^2+1/4*b*(11*A*a^2*b+3*A*b^3-7*B*a^3+B*a*b^2)*\tan(d*x+c)^{(1/2)}/a^2/(a^2+b^2)^2/d/(a+b*\tan(d*x+c))$$

3.414.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.94 (sec) , antiderivative size = 288, normalized size of antiderivative = 0.54

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3} dx$$

$$= \frac{2 \left(\frac{1}{2} \sqrt{b} (35a^4 Ab + 6a^2 Ab^3 + 3Ab^5 - 15a^5 B + 18a^3 b^2 B + ab^4 B) \arctan\left(\frac{\sqrt{b} \sqrt{\tan(c+dx)}}{\sqrt{a}}\right) - 2 \sqrt{-1} a^{5/2} ((a+ib)^3 (A-iB) \arctan((-1)^{3/4} \sqrt{\tan(c+dx)}) + (a-ib)^3 (A+iB) \operatorname{ArcTanh}((-1)^{3/4} \sqrt{\tan(c+dx)})) \right)}{a^{3/2} (a^2 + b^2)^2} + \frac{b (11 a^2 A b + 3 A b^3 - 7 a^3 B + a b^2 B) \sqrt{\tan(c+dx)}}{4 a^2 (a^2 + b^2)^2} + \frac{b (A b - B a) \tan(c+dx)}{2 a (a^2 + b^2)}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^3),x]`

output
$$((2*((\text{Sqrt}[b]*(35*a^4*A*b + 6*a^2*A*b^3 + 3*A*b^5 - 15*a^5*B + 18*a^3*b^2*B + a*b^4*B))*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/2 - 2*(-1)^{(1/4)}*a^{(5/2)}*((a + I*b)^3*(A - I*B)*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]] + (a - I*b)^3*(A + I*B)*\text{ArcTanh}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]])))/(a^{(3/2)}*(a^2 + b^2)^2) + (2*b*(A*b - a*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/(a + b*\text{Tan}[c + d*x])^2 + (b*(11*a^2*A*b + 3*A*b^3 - 7*a^3*B + a*b^2*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/(a*(a^2 + b^2)*(a + b*\text{Tan}[c + d*x]))/(4*a*(a^2 + b^2)*d)$$

3.414.3 Rubi [A] (verified)

Time = 2.21 (sec) , antiderivative size = 466, normalized size of antiderivative = 0.87, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.697$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3} dx \\
 & \quad \downarrow \text{4092} \\
 & \frac{\int \frac{4Aa^2 + bBa - 4(Ab - aB) \tan(c + dx)a + 3Ab^2 + 3b(Ab - aB) \tan^2(c + dx)}{2\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx}{2a(a^2 + b^2)} + \frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{4Aa^2 + bBa - 4(Ab - aB) \tan(c + dx)a + 3Ab^2 + 3b(Ab - aB) \tan^2(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx}{4a(a^2 + b^2)} + \frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{4Aa^2 + bBa - 4(Ab - aB) \tan(c + dx)a + 3Ab^2 + 3b(Ab - aB) \tan(c + dx)^2}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx}{4a(a^2 + b^2)} + \frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{4132} \\
 & \frac{\int \frac{8Aa^4 + 9bBa^3 + 3Ab^2a^2 - 8(-Ba^2 + 2Aba + b^2B) \tan(c + dx)a^2 + b^3Ba + 3Ab^4 + b(-7Ba^3 + 11Aba^2 + b^2Ba + 3Ab^3) \tan^2(c + dx)}{2\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx}{a(a^2 + b^2)} + \frac{b(-7a^3B + 11a^2Ab + ab^2B)}{ad(a^2 + b^2)(a + b \tan(c + dx))^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{4a(a^2 + b^2)}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2} + \frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2}
 \end{aligned}$$

3.414. $\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3} dx$

$$\int \frac{8Aa^4 + 9bBa^3 + 3Ab^2a^2 - 8(-Ba^2 + 2Aba + b^2B) \tan(c+dx)a^2 + b^3Ba + 3Ab^4 + b(-7Ba^3 + 11Aba^2 + b^2Ba + 3Ab^3) \tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx + \frac{b(-7a^3B + 11a^2Ab + ab^2B)}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{2ad(a^2+b^2)(a+b \tan(c+dx))^2}$$

↓ 3042

$$\int \frac{8Aa^4 + 9bBa^3 + 3Ab^2a^2 - 8(-Ba^2 + 2Aba + b^2B) \tan(c+dx)a^2 + b^3Ba + 3Ab^4 + b(-7Ba^3 + 11Aba^2 + b^2Ba + 3Ab^3) \tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx + \frac{b(-7a^3B + 11a^2Ab + ab^2B)}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{2ad(a^2+b^2)(a+b \tan(c+dx))^2}$$

↓ 4136

$$\int \frac{8(a^2(Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) - a^2(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \tan(c+dx))}{\sqrt{\tan(c+dx)}(a^2+b^2)} dx + \frac{b(-15a^5B + 35a^4Ab + 18a^3b^2B + 6a^2Ab^3 + ab^4B + 3Ab^5)}{a^2+b^2} \int \frac{\tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{2ad(a^2+b^2)(a+b \tan(c+dx))^2}$$

↓ 27

$$8 \int \frac{a^2(Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) - a^2(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \tan(c+dx)}{\sqrt{\tan(c+dx)}(a^2+b^2)} dx + \frac{b(-15a^5B + 35a^4Ab + 18a^3b^2B + 6a^2Ab^3 + ab^4B + 3Ab^5)}{a^2+b^2} \int \frac{\tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{2ad(a^2+b^2)(a+b \tan(c+dx))^2}$$

↓ 3042

$$8 \int \frac{a^2(Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) - a^2(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \tan(c+dx)}{\sqrt{\tan(c+dx)}(a^2+b^2)} dx + \frac{b(-15a^5B + 35a^4Ab + 18a^3b^2B + 6a^2Ab^3 + ab^4B + 3Ab^5)}{a^2+b^2} \int \frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \frac{b(Ab-aB)\sqrt{\tan(c+dx)}}{2ad(a^2+b^2)(a+b \tan(c+dx))^2}$$

↓ 4017

3.414. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^3} dx$

$$\frac{16 \int \frac{a^2(Aa^3+3bBa^2-3Ab^2a-b^3B - (-Ba^3+3Aba^2+3b^2Ba-Ab^3) \tan(c+dx)) d\sqrt{\tan(c+dx)}}{\tan^2(c+dx)+1} + \frac{b(-15a^5B+35a^4Ab+18a^3b^2B+6a^2Ab^3+ab^4B+3Ab^5) \int \frac{1}{\sqrt{\tan(c+dx)}}}{a^2+b^2}}{d(a^2+b^2)} \quad \frac{4a(a^2+b^2)}{2a(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 27

$$\frac{16a^2 \int \frac{Aa^3+3bBa^2-3Ab^2a-b^3B - (-Ba^3+3Aba^2+3b^2Ba-Ab^3) \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{b(-15a^5B+35a^4Ab+18a^3b^2B+6a^2Ab^3+ab^4B+3Ab^5) \int \frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}}}{a^2+b^2}}{d(a^2+b^2)} \quad \frac{4a(a^2+b^2)}{2a(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 1482

$$\frac{16a^2 \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(-a^3(A+B)+3a^2b(A-B)+3ab^2(A+B)-b^3(A-B)) \int \frac{\tan(c+dx)}{\tan^2(c+dx)} \right)}{d(a^2+b^2)} \quad \frac{4a(a^2+b^2)}{2a(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 1476

$$\frac{16a^2 \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(-a^3(A+B)+3a^2b(A-B)+3ab^2(A+B)-b^3(A-B)) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)} \right) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 1082

$$\frac{16a^2 \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(-a^3(A+B)+3a^2b(A-B)+3ab^2(A+B)-b^3(A-B)) \left(\frac{1}{2} \int \frac{1}{-\tan(c+dx)} \right) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\tan(c + dx)}}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 217

3.414. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^3} dx$

$$16a^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \int \frac{1 - \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2} (- (a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}}) \right) \right) \frac{1}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\tan(c+dx)}}{2ad(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 1479

$$16a^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(- \frac{\int - \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int - \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right) \frac{1}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\tan(c+dx)}}{2ad(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 25

$$16a^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right) \frac{1}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\tan(c+dx)}}{2ad(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 27

$$16a^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right) \frac{1}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\tan(c+dx)}}{2ad(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 1103

3.414. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^3} dx$

$$\frac{b(-15a^5B+35a^4Ab+18a^3b^2B+6a^2Ab^3+ab^4B+3Ab^5) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2} + \frac{16a^2 \left(\frac{1}{2} (a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right) \left(\frac{\log(\tan(c+dx))}{2\sqrt{2}} \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\tan(c+dx)}}{2ad(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 4117

$$\frac{b(-15a^5B+35a^4Ab+18a^3b^2B+6a^2Ab^3+ab^4B+3Ab^5) \int \frac{1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} d\tan(c+dx)}{d(a^2+b^2)} + \frac{16a^2 \left(\frac{1}{2} (a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right) \left(\frac{\log(\tan(c+dx))}{2\sqrt{2}} \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\tan(c+dx)}}{2ad(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 73

$$\frac{2b(-15a^5B+35a^4Ab+18a^3b^2B+6a^2Ab^3+ab^4B+3Ab^5) \int \frac{1}{a+b\tan(c+dx)} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} + \frac{16a^2 \left(\frac{1}{2} (a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right) \left(\frac{\log(\tan(c+dx))}{2\sqrt{2}} \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\tan(c+dx)}}{2ad(a^2+b^2)(a+b\tan(c+dx))^2}$$

↓ 218

$$\frac{b(Ab - aB)\sqrt{\tan(c+dx)}}{2ad(a^2+b^2)(a+b\tan(c+dx))^2} +$$

$$\frac{16a^2 \left(\frac{1}{2} (a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx))}{2\sqrt{2}} \right)}{ad(a^2+b^2)(a+b\tan(c+dx))} + \frac{b(-7a^3B+11a^2Ab+ab^2B+3Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^3),x]`

$$3.414. \quad \int \frac{A+B\tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^3} dx$$

```
output (b*(A*b - a*B)*Sqrt[Tan[c + d*x]]/(2*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])
^2) + (((2*Sqrt[b]*(35*a^4*A*b + 6*a^2*A*b^3 + 3*A*b^5 - 15*a^5*B + 18*a^3
*b^2*B + a*b^4*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]/(Sqrt[a]*(
a^2 + b^2)*d) + (16*a^2*(-1/2*((3*a^2*b*(A - B) - b^3*(A - B) - a^3*(A + B
) + 3*a*b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) +
ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])) + ((a^3*(A - B) - 3*a*b^2
*(A - B) + 3*a^2*b*(A + B) - b^3*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c
+ d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Ta
n[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)/(2*a*(a^2 + b^2)) + (b*(11*
a^2*A*b + 3*A*b^3 - 7*a^3*B + a*b^2*B)*Sqrt[Tan[c + d*x]])/(a*(a^2 + b^2)*
d*(a + b*Tan[c + d*x]))/(4*a*(a^2 + b^2))
```

3.414.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

$$3.414. \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^3} dx$$

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.414.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 452, normalized size of antiderivative = 0.85

method	result
derivativedivides	$2b \frac{\left(\frac{b(11Aa^4b+14Aa^2b^3+3Ab^5-7Ba^5-6Ba^3b^2+Ba^4)}{8a^2} \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) + \frac{(13Aa^4b+18Aa^2b^3+5Ab^5-9Ba^5-10Ba^3b^2-Ba^4)}{8a}}{(a+b \tan(dx+c))^2} + \frac{1}{(a^2+b^2)^3}$
default	$2b \frac{\left(\frac{b(11Aa^4b+14Aa^2b^3+3Ab^5-7Ba^5-6Ba^3b^2+Ba^4)}{8a^2} \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) + \frac{(13Aa^4b+18Aa^2b^3+5Ab^5-9Ba^5-10Ba^3b^2-Ba^4)}{8a}}{(a+b \tan(dx+c))^2} + \frac{1}{(a^2+b^2)^3}$

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^3,x,method=_RETURNV
ERBOSE)`

output `1/d*(2*b/(a^2+b^2)^3*((1/8*b*(11*A*a^4*b+14*A*a^2*b^3+3*A*b^5-7*B*a^5-6*B*
a^3*b^2+B*a*b^4)/a^2*tan(d*x+c)^(3/2)+1/8*(13*A*a^4*b+18*A*a^2*b^3+5*A*b^5
-9*B*a^5-10*B*a^3*b^2-B*a*b^4)/a*tan(d*x+c)^(1/2))/(a+b*tan(d*x+c))^2+1/8*
(35*A*a^4*b+6*A*a^2*b^3+3*A*b^5-15*B*a^5+18*B*a^3*b^2+B*a*b^4)/a^2/(a*b)^(
1/2)*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2)))+2/(a^2+b^2)^3*(1/8*(A*a^3-3*A
*a*b^2+3*B*a^2*b-B*b^3)*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)
)/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(
1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))+1/8*(-3*A*a^2*b+A*b^3+B*a^3-3
*B*a*b^2)*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*
tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan
(-1+2^(1/2)*tan(d*x+c)^(1/2))))`

3.414.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8916 vs. 2(484) = 968.

Time = 118.65 (sec) , antiderivative size = 17859, normalized size of antiderivative = 33.44

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^3,x, algorithm m="fricas")`

output Too large to include

3.414.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+b*tan(d*x+c))**3,x)`

output Timed out

3.414.7 Maxima [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 551, normalized size of antiderivative = 1.03

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3} dx = \frac{(15 Ba^5 b - 35 Aa^4 b^2 - 18 Ba^3 b^3 - 6 Aa^2 b^4 - Bab^5 - 3 Ab^6) \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right) - 2\sqrt{2}((A+B)a^3 - 3(A-B)a^2 b - 3(A+B)ab^2 + (A-B)b^3)}{(a^8 + 3a^6 b^2 + 3a^4 b^4 + a^2 b^6)\sqrt{ab}}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^3,x, algorithm m="maxima")`

output
$$\begin{aligned} & -1/4*((15*B*a^5*b - 35*A*a^4*b^2 - 18*B*a^3*b^3 - 6*A*a^2*b^4 - B*a*b^5 - \\ & 3*A*b^6)*\arctan(b*\sqrt{\tan(dx + c)}/\sqrt{a*b})/((a^8 + 3*a^6*b^2 + 3*a^4* \\ & b^4 + a^2*b^6)*\sqrt{a*b}) - (2*\sqrt{2}*((A + B)*a^3 - 3*(A - B)*a^2*b - 3* \\ & (A + B)*a*b^2 + (A - B)*b^3)*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2*\sqrt{\tan(dx \\ & + c)))) + 2*\sqrt{2}*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A \\ & - B)*b^3)*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2*\sqrt{\tan(dx + c)))) + \sqrt{2}* \\ & ((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*\log(\sqrt{2} \\ &)*\sqrt{\tan(dx + c)} + \tan(dx + c) + 1) - \sqrt{2}*((A - B)*a^3 + 3*(A + B) \\ &)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*\log(-\sqrt{2}*\sqrt{\tan(dx + c)} + \\ & \tan(dx + c) + 1))/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + ((7*B*a^3*b^2 - \\ & 11*A*a^2*b^3 - B*a*b^4 - 3*A*b^5)*\tan(dx + c)^{(3/2)} + (9*B*a^4*b - 13*A*a \\ & ^3*b^2 + B*a^2*b^3 - 5*A*a*b^4)*\sqrt{\tan(dx + c)})/(a^8 + 2*a^6*b^2 + a^4 \\ & *b^4 + (a^6*b^2 + 2*a^4*b^4 + a^2*b^6)*\tan(dx + c)^2 + 2*(a^7*b + 2*a^5*b \\ & ^3 + a^3*b^5)*\tan(dx + c))/d \end{aligned}$$

3.414.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^3,x, algorithm="giac")`

output Timed out

3.414.9 Mupad [B] (verification not implemented)

Time = 63.37 (sec) , antiderivative size = 26707, normalized size of antiderivative = 50.01

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + b*tan(c + d*x))^3),x)`

output

$$\begin{aligned}
& ((A*\tan(c + d*x)^{(1/2)}*(5*b^4 + 13*a^2*b^2))/(4*a*(a^4 + b^4 + 2*a^2*b^2)) \\
& + (A*b*\tan(c + d*x)^{(3/2)}*(3*b^4 + 11*a^2*b^2))/(4*a^2*(a^4 + b^4 + 2*a^2 \\
& *b^2)))/(a^2*d + b^2*d*\tan(c + d*x)^2 + 2*a*b*d*\tan(c + d*x)) - ((\tan(c + \\
& d*x)^{(1/2)}*(B*b^3 + 9*B*a^2*b))/(4*(a^4 + b^4 + 2*a^2*b^2)) - (\tan(c + d*x) \\
&)^{(3/2)}*(B*b^4 - 7*B*a^2*b^2))/(4*a*(a^4 + b^4 + 2*a^2*b^2)))/(a^2*d + b^2 \\
& *d*\tan(c + d*x)^2 + 2*a*b*d*\tan(c + d*x)) + (\log((((((((((64*A*b^2*(3*b^6 \\
& - 2*a^6 + 3*a^2*b^4 + 22*a^4*b^2))/(a^2*d) + 128*b^3*\tan(c + d*x)^{(1/2)}*(a \\
& ^2 - b^2)*(a^2 + b^2)^2*((4*(-A^4*d^4*(a^6 - b^6 + 15*a^2*b^4 - 15*a^4*b^2) \\
&)^2)^{(1/2)} - 80*A^2*a^3*b^3*d^2 + 24*A^2*a*b^5*d^2 + 24*A^2*a^5*b*d^2)/(d^ \\
& 4*(a^2 + b^2)^6))^{(1/2)}))*((4*(-A^4*d^4*(a^6 - b^6 + 15*a^2*b^4 - 15*a^4*b^ \\
& 2)^2)^{(1/2)} - 80*A^2*a^3*b^3*d^2 + 24*A^2*a*b^5*d^2 + 24*A^2*a^5*b*d^2)/(d \\
& ^4*(a^2 + b^2)^6))^{(1/2)})/4 + (8*A^2*b^2*\tan(c + d*x)^{(1/2)}*(9*b^12 - 8*a^ \\
& 12 + 36*a^2*b^10 + 430*a^4*b^8 - 188*a^6*b^6 + 1497*a^8*b^4 + 32*a^10*b^2) \\
&)/(a^3*d^2*(a^2 + b^2)^4))*((4*(-A^4*d^4*(a^6 - b^6 + 15*a^2*b^4 - 15*a^4* \\
& b^2)^2)^{(1/2)} - 80*A^2*a^3*b^3*d^2 + 24*A^2*a*b^5*d^2 + 24*A^2*a^5*b*d^2)/ \\
& (d^4*(a^2 + b^2)^6))^{(1/2)})/4 - (2*A^3*b^3*(45*b^12 - 16*a^12 + 333*a^2*b^ \\
& 10 + 146*a^4*b^8 + 1178*a^6*b^6 - 9791*a^8*b^4 + 1161*a^10*b^2))/(a^3*d^3* \\
& (a^2 + b^2)^6))*((4*(-A^4*d^4*(a^6 - b^6 + 15*a^2*b^4 - 15*a^4*b^2)^2)^{(1/ \\
& 2)} - 80*A^2*a^3*b^3*d^2 + 24*A^2*a*b^5*d^2 + 24*A^2*a^5*b*d^2)/(d^4*(a^2 + \\
& b^2)^6))^{(1/2)})/4 + (A^4*b^5*\tan(c + d*x)^{(1/2)}*(18*a^2*b^10 - 9*b^12 ...
\end{aligned}$$

$$3.415 \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$$

3.415.1 Optimal result	4007
3.415.2 Mathematica [C] (verified)	4008
3.415.3 Rubi [A] (verified)	4009
3.415.4 Maple [A] (verified)	4018
3.415.5 Fricas [B] (verification not implemented)	4019
3.415.6 Sympy [F(-1)]	4019
3.415.7 Maxima [A] (verification not implemented)	4020
3.415.8 Giac [F(-1)]	4020
3.415.9 Mupad [B] (verification not implemented)	4021

3.415.1 Optimal result

Integrand size = 33, antiderivative size = 601

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx \\ &= \frac{(a^3(A - B) - 3ab^2(A - B) + 3a^2b(A + B) - b^3(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^3 d} \\ & \quad - \frac{(a^3(A - B) - 3ab^2(A - B) + 3a^2b(A + B) - b^3(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^3 d} \\ & \quad - \frac{b^{3/2}(63a^4Ab + 46a^2Ab^3 + 15Ab^5 - 35a^5B - 6a^3b^2B - 3ab^4B) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{4a^{7/2}(a^2 + b^2)^3 d} \\ & \quad + \frac{(3a^2b(A - B) - b^3(A - B) - a^3(A + B) + 3ab^2(A + B)) \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^3 d} \\ & \quad - \frac{(3a^2b(A - B) - b^3(A - B) - a^3(A + B) + 3ab^2(A + B)) \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^3 d} \\ & \quad - \frac{8a^4A + 31a^2Ab^2 + 15Ab^4 - 11a^3bB - 3ab^3B}{4a^3(a^2 + b^2)^2 d \sqrt{\tan(c + dx)}} \\ & \quad + \frac{b(Ab - aB)}{2a(a^2 + b^2) d \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} \\ & \quad + \frac{b(13a^2Ab + 5Ab^3 - 9a^3B - ab^2B)}{4a^2(a^2 + b^2)^2 d \sqrt{\tan(c + dx)}(a + b \tan(c + dx))} \end{aligned}$$

$$3.415. \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$$

output
$$-1/4*b^{(3/2)}*(63*A*a^4*b+46*A*a^2*b^3+15*A*b^5-35*B*a^5-6*B*a^3*b^2-3*B*a*b^4)*\arctan(b^{(1/2)}*\tan(d*x+c)^{(1/2)}/a^{(1/2)})/a^{(7/2)}/(a^2+b^2)^3/d-1/2*(a^3*(A-B)-3*a*b^2*(A-B)+3*a^2*b*(A+B)-b^3*(A+B))*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/(a^2+b^2)^3/d*2^{(1/2)}-1/2*(a^3*(A-B)-3*a*b^2*(A-B)+3*a^2*b*(A+B)-b^3*(A+B))*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})/(a^2+b^2)^3/d*2^{(1/2)}+1/4*(3*a^2*b*(A-B)-b^3*(A-B)-a^3*(A+B)+3*a*b^2*(A+B))*\ln(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(a^2+b^2)^3/d*2^{(1/2)}-1/4*(3*a^2*b*(A-B)-b^3*(A-B)-a^3*(A+B)+3*a*b^2*(A+B))*\ln(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(a^2+b^2)^3/d*2^{(1/2)}+1/4*(-8*A*a^4-31*A*a^2*b^2-15*A*b^4+11*B*a^3*b+3*B*a*b^3)/a^3/(a^2+b^2)^2/d/\tan(d*x+c)^{(1/2)}+1/2*b*(A*b-B*a)/a/(a^2+b^2)/d/\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^2+1/4*b*(13*A*a^2*b+5*A*b^3-9*B*a^3-B*a*b^2)/a^2/(a^2+b^2)^2/d/\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))$$

3.415.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.35 (sec) , antiderivative size = 585, normalized size of antiderivative = 0.97

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \frac{b(Ab - aB)}{2a(a^2 + b^2) d \sqrt{\tan(c + dx)(a + b \tan(c + dx))^2}} - \frac{2(-a^4b(a^2A - Ab^2 + 2abB) + \frac{1}{8}a^2b(8a^4A + 31a^2Ab^2 + 15Ab^4 - 11a^3bB - 3ab^3B) + \frac{1}{8}b^2(24a^4Ab + 31a^2Ab^3 + 15Ab^5 - 8a^5B - 3a^3b^2B - 3ab^4B)) \arctan\left(\frac{\sqrt{b}}{\sqrt{a}\sqrt{b}(a^2+b^2)}\right)}{\sqrt{a}\sqrt{b}(a^2+b^2)d} + \dots$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^3),x]`

output $(b*(A*b - a*B))/(2*a*(a^2 + b^2)*d*\text{Sqrt}[\text{Tan}[c + d*x]]*(a + b*\text{Tan}[c + d*x])^2) + (((-2*((2*(-(a^4*b*(a^2*A - A*b^2 + 2*a*b*B)) + (a^2*b*(8*a^4*A + 31*a^2*A*b^2 + 15*A*b^4 - 11*a^3*b*B - 3*a*b^3*B)))/8 + (b^2*(24*a^4*A*b + 31*a^2*A*b^3 + 15*A*b^5 - 8*a^5*B - 3*a^3*b^2*B - 3*a*b^4*B))/8)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(\text{Sqrt}[a]*\text{Sqrt}[b]*(a^2 + b^2)*d) + (-(((-1)^(1/4)*(a^3*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B) - I*a^3*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B))*\text{ArcTan}[(-1)^(3/4)*\text{Sqrt}[\text{Tan}[c + d*x]])/d) - (((-1)^(1/4)*(a^3*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B) + I*a^3*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B))*\text{ArcTanh}[(-1)^(3/4)*\text{Sqrt}[\text{Tan}[c + d*x]])/d)/(a^2 + b^2)))/a - (8*a^4*A + 31*a^2*A*b^2 + 15*A*b^4 - 11*a^3*b*B - 3*a*b^3*B)/(2*a*d*\text{Sqrt}[\text{Tan}[c + d*x]])/(a*(a^2 + b^2)) + ((9*a^2*b*(A*b - a*B))/2 + (b^2*(4*a^2*A + 5*A*b^2 - a*b*B))/2)/(a*(a^2 + b^2)*d*\text{Sqrt}[\text{Tan}[c + d*x]]*(a + b*\text{Tan}[c + d*x]))/(2*a*(a^2 + b^2))$

3.415.3 Rubi [A] (verified)

Time = 3.01 (sec) , antiderivative size = 528, normalized size of antiderivative = 0.88, number of steps used = 27, number of rules used = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.788$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2}(a + b \tan(c + dx))^3} dx$$

↓ 4092

$$\frac{\int \frac{4Aa^2 - bBa - 4(Ab - aB) \tan(c + dx)a + 5Ab^2 + 5b(Ab - aB) \tan^2(c + dx)}{2 \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx}{\frac{2a(a^2 + b^2)}{b(Ab - aB)}} +$$

$$\frac{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}{b(Ab - aB)}$$

↓ 27

3.415. $\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx$

$$\begin{aligned}
& \frac{\int \frac{4Aa^2 - bBa - 4(Ab - aB) \tan(c+dx)a + 5Ab^2 + 5b(Ab - aB) \tan^2(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx}{\frac{4a(a^2 + b^2)}{b(Ab - aB)}} + \\
& \frac{2ad(a^2 + b^2) \sqrt{\tan(c+dx)}(a + b \tan(c+dx))^2}{\downarrow 3042} \\
& \frac{\int \frac{4Aa^2 - bBa - 4(Ab - aB) \tan(c+dx)a + 5Ab^2 + 5b(Ab - aB) \tan(c+dx)^2}{\tan(c+dx)^{3/2}(a+b \tan(c+dx))^2} dx}{\frac{4a(a^2 + b^2)}{b(Ab - aB)}} + \\
& \frac{2ad(a^2 + b^2) \sqrt{\tan(c+dx)}(a + b \tan(c+dx))^2}{\downarrow 4132} \\
& \frac{\int \frac{8Aa^4 - 11bBa^3 + 31Ab^2a^2 - 8(-Ba^2 + 2Aba + b^2B) \tan(c+dx)a^2 - 3b^3Ba + 15Ab^4 + 3b(-9Ba^3 + 13Aba^2 - b^2Ba + 5Ab^3) \tan^2(c+dx)}{2 \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx}{\frac{4a(a^2 + b^2)}{b(Ab - aB)}} + \frac{b(-9a^3B + 13a^2A)}{ad(a^2 + b^2) \sqrt{\tan(c+dx)}} \\
& \frac{2ad(a^2 + b^2) \sqrt{\tan(c+dx)}(a + b \tan(c+dx))^2}{\downarrow 27} \\
& \frac{\int \frac{8Aa^4 - 11bBa^3 + 31Ab^2a^2 - 8(-Ba^2 + 2Aba + b^2B) \tan(c+dx)a^2 - 3b^3Ba + 15Ab^4 + 3b(-9Ba^3 + 13Aba^2 - b^2Ba + 5Ab^3) \tan^2(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx}{\frac{4a(a^2 + b^2)}{2a(a^2 + b^2)}} + \frac{b(-9a^3B + 13a^2A)}{ad(a^2 + b^2) \sqrt{\tan(c+dx)}} \\
& \frac{2ad(a^2 + b^2) \sqrt{\tan(c+dx)}(a + b \tan(c+dx))^2}{\downarrow 3042} \\
& \frac{\int \frac{8Aa^4 - 11bBa^3 + 31Ab^2a^2 - 8(-Ba^2 + 2Aba + b^2B) \tan(c+dx)a^2 - 3b^3Ba + 15Ab^4 + 3b(-9Ba^3 + 13Aba^2 - b^2Ba + 5Ab^3) \tan(c+dx)^2}{\tan(c+dx)^{3/2}(a+b \tan(c+dx))} dx}{\frac{4a(a^2 + b^2)}{2a(a^2 + b^2)}} + \frac{b(-9a^3B + 13a^2A)}{ad(a^2 + b^2) \sqrt{\tan(c+dx)}} \\
& \frac{2ad(a^2 + b^2) \sqrt{\tan(c+dx)}(a + b \tan(c+dx))^2}{\downarrow 4132}
\end{aligned}$$

3.415. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

$$\begin{aligned}
 & \int \frac{-8Ba^5+24Aba^4-3b^2Ba^3+8(Aa^2+2bBa-Ab^2)\tan(c+dx)a^3+31Ab^3a^2-3b^4Ba+15Ab^5+b(8Aa^4-11bBa^3+31Ab^2a^2-3b^3Ba+15Ab^4)\tan^2(c+dx)}{2\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx - 2(8a^5-8Ba^4+4Aba^3-4b^2Ba^2+4Ab^3a-4b^4Ba+4Ab^5) \int \frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx \\
 & \frac{2a(a^2+b^2)}{4a(a^2+b^2)} \\
 & \frac{b(Ab-aB)}{2ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2} \\
 & \downarrow 27 \\
 & \int \frac{-8Ba^5+24Aba^4-3b^2Ba^3+8(Aa^2+2bBa-Ab^2)\tan(c+dx)a^3+31Ab^3a^2-3b^4Ba+15Ab^5+b(8Aa^4-11bBa^3+31Ab^2a^2-3b^3Ba+15Ab^4)\tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx - 2(8a^5-8Ba^4+4Aba^3-4b^2Ba^2+4Ab^3a-4b^4Ba+4Ab^5) \int \frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx \\
 & \frac{2a(a^2+b^2)}{4a(a^2+b^2)} \\
 & \frac{b(Ab-aB)}{2ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2} \\
 & \downarrow 3042 \\
 & \int \frac{-8Ba^5+24Aba^4-3b^2Ba^3+8(Aa^2+2bBa-Ab^2)\tan(c+dx)a^3+31Ab^3a^2-3b^4Ba+15Ab^5+b(8Aa^4-11bBa^3+31Ab^2a^2-3b^3Ba+15Ab^4)\tan(c+dx)^2}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx - 2(8a^5-8Ba^4+4Aba^3-4b^2Ba^2+4Ab^3a-4b^4Ba+4Ab^5) \int \frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx \\
 & \frac{2a(a^2+b^2)}{4a(a^2+b^2)} \\
 & \frac{b(Ab-aB)}{2ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2} \\
 & \downarrow 4136 \\
 & \int \frac{8\left(\frac{-Ba^3+3Aba^2+3b^2Ba-Ab^3}{a^2+b^2}\right)a^3+\left(\frac{Aa^3+3bBa^2-3Ab^2a-b^3B}{a^2+b^2}\right)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx + \frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5)}{a^2+b^2} \int \frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx \\
 & \frac{2a(a^2+b^2)}{4a(a^2+b^2)} \\
 & \frac{b(Ab-aB)}{2ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2} \\
 & \downarrow 27 \\
 & 8 \int \frac{\left(\frac{-Ba^3+3Aba^2+3b^2Ba-Ab^3}{a^2+b^2}\right)a^3+\left(\frac{Aa^3+3bBa^2-3Ab^2a-b^3B}{a^2+b^2}\right)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}} dx + \frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5)}{a^2+b^2} \int \frac{\tan^2(c+dx)}{\sqrt{\tan(c+dx)}} dx \\
 & \frac{2a(a^2+b^2)}{4a(a^2+b^2)} \\
 & \frac{b(Ab-aB)}{2ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2} \\
 & \downarrow 3042
 \end{aligned}$$

3.415. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^3} dx$

$$\frac{8 \int \frac{(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3)a^3 + (Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)} \frac{a^2+b^2}}{a^2+b^2} dx + \frac{b^2(-35a^5B + 63a^4Ab - 6a^3b^2B + 46a^2Ab^3 - 3ab^4B + 15Ab^5) \int \frac{\tan(c+dx)}{\sqrt{\tan(c+dx)}}$$

$$\frac{2a(a^2+b^2)}{a^2+b^2}$$

$$\frac{b(Ab - aB)}{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}$$

↓ 4017

$$\frac{16 \int \frac{a^3(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) + (Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{b^2(-35a^5B + 63a^4Ab - 6a^3b^2B + 46a^2Ab^3 - 3ab^4B + 15Ab^5) \int \sqrt{\tan(c+dx)}}{d(a^2+b^2)}}{a^2+b^2}$$

$$\frac{2a(a^2+b^2)}{a^2+b^2}$$

$$\frac{b(Ab - aB)}{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}$$

↓ 27

$$\frac{16a^3 \int \frac{-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3 + (Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{b^2(-35a^5B + 63a^4Ab - 6a^3b^2B + 46a^2Ab^3 - 3ab^4B + 15Ab^5) \int \sqrt{\tan(c+dx)}}{d(a^2+b^2)}}{a^2+b^2}$$

$$\frac{2a(a^2+b^2)}{a^2+b^2}$$

$$\frac{b(Ab - aB)}{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}$$

↓ 1482

$$\frac{16a^3 \left(\frac{1}{2}(-a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B) \right) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \int \frac{\tan(c+dx)}{\tan^2(c+dx)}}{d(a^2+b^2)}}$$

$$\frac{2a(a^2+b^2)}{a^2+b^2}$$

$$\frac{b(Ab - aB)}{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}$$

↓ 1476

3.415. $\int \frac{A+B \tan(c+dx)}{\tan^3(c+dx)(a+b \tan(c+dx))^3} dx$

$$\frac{16a^3 \left(\frac{1}{2} \left(- (a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B) \right) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \left(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B) \right) \left(\frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}$$

↓ 1082

$$\frac{16a^3 \left(\frac{1}{2} \left(- (a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B) \right) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \left(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B) \right) \left(\frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}$$

↓ 217

$$\frac{16a^3 \left(\frac{1}{2} \left(- (a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B) \right) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \left(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B) \right) \left(\frac{\arctan(\sqrt{\tan(c+dx)})}{a} \right) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}$$

↓ 1479

$$\frac{16a^3 \left(\frac{1}{2} \left(- (a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B) \right) \left(- \frac{\int - \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int - \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}$$

↓ 25

3.415. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

$$16a^3 \left(\frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right) \frac{1}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}$$

↓ 27

$$16a^3 \left(\frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right) \frac{1}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}$$

↓ 1103

$$\frac{b^2(-35a^5B + 63a^4Ab - 6a^3b^2B + 46a^2Ab^3 - 3ab^4B + 15Ab^5) \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} + \frac{16a^3 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arcsin(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}\sqrt{\tan(c+dx)+1}} \right) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}$$

↓ 4117

$$\frac{b^2(-35a^5B + 63a^4Ab - 6a^3b^2B + 46a^2Ab^3 - 3ab^4B + 15Ab^5) \int \frac{1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} d \tan(c+dx)}{d(a^2+b^2)} + \frac{16a^3 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arcsin(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}\sqrt{\tan(c+dx)+1}} \right) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}$$

↓ 73

3.415. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

$$\frac{2b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5)}{d(a^2+b^2)} \int \frac{1}{a+b \tan(c+dx)} d\sqrt{\tan(c+dx)} + \frac{16a^3 \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right) \left(\frac{\arctan(\sqrt{\tan(c+dx)})}{\sqrt{\tan(c+dx)}} \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)}{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2}$$

↓ 218

$$\frac{b(Ab - aB)}{2ad(a^2 + b^2) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} + \frac{16a^3 \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right) \left(\frac{\arctan(\sqrt{\tan(c+dx)})}{\sqrt{\tan(c+dx)}} \right)}{d(a^2+b^2)}$$

$$\frac{b(-9a^3B+13a^2Ab-ab^2B+5Ab^3)}{ad(a^2+b^2)\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} + \frac{2(8a^4A-11a^3bB+31a^2Ab^2-3ab^3B+15Ab^4)}{ad\sqrt{\tan(c+dx)}} - \frac{16a^3 \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right) \left(\frac{\arctan(\sqrt{\tan(c+dx)})}{\sqrt{\tan(c+dx)}} \right)}{d(a^2+b^2)}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^3),x]`

output `(b*(A*b - a*B))/(2*a*(a^2 + b^2)*d*sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^2) + (((-(((2*b^(3/2)*(63*a^4*A*b + 46*a^2*A*b^3 + 15*A*b^5 - 35*a^5*B - 6*a^3*b^2*B - 3*a*b^4*B)*ArcTan[(sqrt[b]*sqrt[Tan[c + d*x]])/sqrt[a]])/(sqrt[a]*(a^2 + b^2)*d) + (16*a^3*((a^3*(A - B) - 3*a*b^2*(A - B) + 3*a^2*b*(A + B) - b^3*(A + B))*(-ArcTan[1 - sqrt[2]*sqrt[Tan[c + d*x]])/sqrt[2]] + ArcTan[1 + sqrt[2]*sqrt[Tan[c + d*x]])/sqrt[2]]))/2 + ((3*a^2*b*(A - B) - b^3*(A - B) - a^3*(A + B) + 3*a*b^2*(A + B))*(-1/2*Log[1 - sqrt[2]*sqrt[Tan[c + d*x]] + Tan[c + d*x]]/sqrt[2] + Log[1 + sqrt[2]*sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*sqrt[2])))/2))/((a^2 + b^2)*d))/a - (2*(8*a^4*A + 31*a^2*A*b^2 + 15*A*b^4 - 11*a^3*b*B - 3*a*b^3*B))/(a*d*sqrt[Tan[c + d*x]]))/(2*a*(a^2 + b^2)) + (b*(13*a^2*A*b + 5*A*b^3 - 9*a^3*B - a*b^2*B))/(a*(a^2 + b^2)*d*sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x]))/(4*a*(a^2 + b^2))`

3.415. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

3.415.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`


```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*
Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.415.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 466, normalized size of antiderivative = 0.78

method	result
derivativedivides	$2b^2 \frac{\left(\frac{15}{8} A a^4 b^2 + \frac{11}{4} A a^2 b^4 + \frac{7}{8} A b^6 - \frac{11}{8} B a^5 b - \frac{7}{4} B a^3 b^3 - \frac{3}{8} B a b^5 \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) + \frac{a(17A a^4 b + 26A a^2 b^3 + 9A b^5 - 13B a^5 - 18B a^3 b^3 - 9B a b^5)}{8}}{(a+b \tan(dx+c))^2} - \frac{a^3(a^2+b^2)^3}{a^3(a^2+b^2)^3}$
default	$2b^2 \frac{\left(\frac{15}{8} A a^4 b^2 + \frac{11}{4} A a^2 b^4 + \frac{7}{8} A b^6 - \frac{11}{8} B a^5 b - \frac{7}{4} B a^3 b^3 - \frac{3}{8} B a b^5 \right) \left(\tan^{\frac{3}{2}}(dx+c) \right) + \frac{a(17A a^4 b + 26A a^2 b^3 + 9A b^5 - 13B a^5 - 18B a^3 b^3 - 9B a b^5)}{8}}{(a+b \tan(dx+c))^2} - \frac{a^3(a^2+b^2)^3}{a^3(a^2+b^2)^3}$

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^3,x,method=_RETURNV
ERBOSE)
```

$$3.415. \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$$

output $1/d*(-2*b^2/a^3/(a^2+b^2)^3*((15/8*A*a^4*b^2+11/4*A*a^2*b^4+7/8*A*b^6-11/8*B*a^5*b-7/4*B*a^3*b^3-3/8*B*a*b^5)*\tan(dx+c)^{(3/2)}+1/8*a*(17*A*a^4*b+26*A*a^2*b^3+9*A*b^5-13*B*a^5-18*B*a^3*b^2-5*B*a*b^4)*\tan(dx+c)^{(1/2)})/(a+b*\tan(dx+c))^2+1/8*(63*A*a^4*b+46*A*a^2*b^3+15*A*b^5-35*B*a^5-6*B*a^3*b^2-3*B*a*b^4)/(a*b)^{(1/2)}*\arctan(b*\tan(dx+c)^{(1/2)}/(a*b)^{(1/2)}))-2/a^3*A/\tan(dx+c)^{(1/2)}+2/(a^2+b^2)^3*(1/8*(-3*A*a^2*b+A*b^3+B*a^3-3*B*a*b^2)*2^{(1/2)}*(\ln((1+2^{(1/2)}*\tan(dx+c)^{(1/2)}+\tan(dx+c))/(1-2^{(1/2)}*\tan(dx+c)^{(1/2)}+\tan(dx+c))))+2*\arctan(1+2^{(1/2)}*\tan(dx+c)^{(1/2)}))+2*\arctan(-1+2^{(1/2)}*\tan(dx+c)^{(1/2)}))+1/8*(-A*a^3+3*A*a*b^2-3*B*a^2*b+B*b^3)*2^{(1/2)}*(\ln((1-2^{(1/2)}*\tan(dx+c)^{(1/2)}+\tan(dx+c))/(1+2^{(1/2)}*\tan(dx+c)^{(1/2)}+\tan(dx+c))))+2*\arctan(1+2^{(1/2)}*\tan(dx+c)^{(1/2)}))+2*\arctan(-1+2^{(1/2)}*\tan(dx+c)^{(1/2)}))$

3.415.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 9032 vs. $2(544) = 1088$.

Time = 189.65 (sec) , antiderivative size = 18091, normalized size of antiderivative = 30.10

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^3,x, algorithm="fricas")`

output Too large to include

3.415.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(3/2)/(a+b*tan(d*x+c))**3,x)`

output Timed out

3.415. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

3.415.7 Maxima [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 607, normalized size of antiderivative = 1.01

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx$$

$$= \frac{(35 B a^5 b^2 - 63 A a^4 b^3 + 6 B a^3 b^4 - 46 A a^2 b^5 + 3 B a b^6 - 15 A b^7) \arctan\left(\frac{b \sqrt{\tan(dx+c)}}{\sqrt{ab}}\right) - \frac{8 A a^6 + 16 A a^4 b^2 + 8 A a^2 b^4 + (8 A a^4 b^2 - 11 B a^3 b^3 + 31 A a^2 b^4 - 15 A a b^5 + 5 A b^6) \tan(dx+c)}{(a^9 + 3 a^7 b^2 + 3 a^5 b^4 + a^3 b^6) \sqrt{ab}}}{(a^7 b^2 + 2 a^5 b^4 + a^3 b^6) \tan(dx+c)}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^3,x, algorithm="maxima")`

output `1/4*((35*B*a^5*b^2 - 63*A*a^4*b^3 + 6*B*a^3*b^4 - 46*A*a^2*b^5 + 3*B*a*b^6 - 15*A*b^7)*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/((a^9 + 3*a^7*b^2 + 3*a^5*b^4 + a^3*b^6)*sqrt(a*b)) - (8*A*a^6 + 16*A*a^4*b^2 + 8*A*a^2*b^4 + (8*A*a^4*b^2 - 11*B*a^3*b^3 + 31*A*a^2*b^4 - 3*B*a*b^5 + 15*A*b^6)*tan(d*x + c)^2 + (16*A*a^5*b - 13*B*a^4*b^2 + 49*A*a^3*b^3 - 5*B*a^2*b^4 + 25*A*a*b^5)*tan(d*x + c))/((a^7*b^2 + 2*a^5*b^4 + a^3*b^6)*tan(d*x + c)^(5/2) + 2*(a^8*b + 2*a^6*b^3 + a^4*b^5)*tan(d*x + c)^(3/2) + (a^9 + 2*a^7*b^2 + a^5*b^4)*sqrt(tan(d*x + c))) - (2*sqrt(2))*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6))/d`

3.415.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^3,x, algorithm="giac")`

output `Timed out`

3.415. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

3.415.9 Mupad [B] (verification not implemented)

Time = 57.04 (sec) , antiderivative size = 35300, normalized size of antiderivative = 58.74

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

```
input int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + b*tan(c + d*x))^3),x)
```

```
output ((B*tan(c + d*x)^(1/2)*(5*b^4 + 13*a^2*b^2))/(4*a*(a^4 + b^4 + 2*a^2*b^2))
+ (B*b*tan(c + d*x)^(3/2)*(3*b^4 + 11*a^2*b^2))/(4*a^2*(a^4 + b^4 + 2*a^2
*b^2)))/(a^2*d + b^2*d*tan(c + d*x)^2 + 2*a*b*d*tan(c + d*x)) - ((2*A)/a +
(A*tan(c + d*x)^2*(15*b^6 + 31*a^2*b^4 + 8*a^4*b^2))/(4*a^3*(a^4 + b^4 +
2*a^2*b^2)) + (A*tan(c + d*x)*(16*a^4*b + 25*b^5 + 49*a^2*b^3))/(4*a^2*(a^
4 + b^4 + 2*a^2*b^2)))/(a^2*d*tan(c + d*x)^(1/2) + b^2*d*tan(c + d*x)^(5/2
) + 2*a*b*d*tan(c + d*x)^(3/2)) + (log(29491200*A^5*a^22*b^35*d^4 - ((tan(
c + d*x)^(1/2)*(7610564608*A^4*a^27*b^33*d^5 - 597688320*A^4*a^23*b^37*d^5
- 1671430144*A^4*a^25*b^35*d^5 - 58982400*A^4*a^21*b^39*d^5 + 85774565376
*A^4*a^29*b^31*d^5 + 385487994880*A^4*a^31*b^29*d^5 + 1104303620096*A^4*a^
33*b^27*d^5 + 2240523796480*A^4*a^35*b^25*d^5 + 3345249468416*A^4*a^37*b^2
3*d^5 + 3717287903232*A^4*a^39*b^21*d^5 + 3053967114240*A^4*a^41*b^19*d^5
+ 1807474491392*A^4*a^43*b^17*d^5 + 726513221632*A^4*a^45*b^15*d^5 + 17076
8990208*A^4*a^47*b^13*d^5 + 10492051456*A^4*a^49*b^11*d^5 - 4917821440*A^4
*a^51*b^9*d^5 - 923009024*A^4*a^53*b^7*d^5 + 8388608*A^4*a^55*b^5*d^5) + (
(((480*A^4*a^2*b^10*d^4 - 16*A^4*b^12*d^4 - 16*A^4*a^12*d^4 - 4080*A^4*a^4
*b^8*d^4 + 7232*A^4*a^6*b^6*d^4 - 4080*A^4*a^8*b^4*d^4 + 480*A^4*a^10*b^2*
d^4)^(1/2) + 80*A^2*a^3*b^3*d^2 - 24*A^2*a*b^5*d^2 - 24*A^2*a^5*b*d^2)/(a^
12*d^4 + b^12*d^4 + 6*a^2*b^10*d^4 + 15*a^4*b^8*d^4 + 20*a^6*b^6*d^4 + 15*
a^8*b^4*d^4 + 6*a^10*b^2*d^4))^(1/2)*(((((((480*A^4*a^2*b^10*d^4 - 16*A...
```

3.416
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.416.1 Optimal result 4022
 3.416.2 Mathematica [A] (verified) 4023
 3.416.3 Rubi [A] (verified) 4023
 3.416.4 Maple [A] (verified) 4027
 3.416.5 Fricas [C] (verification not implemented) 4028
 3.416.6 Sympy [F] 4028
 3.416.7 Maxima [A] (verification not implemented) 4029
 3.416.8 Giac [F(-1)] 4029
 3.416.9 Mupad [B] (verification not implemented) 4029

3.416.1 Optimal result

Integrand size = 36, antiderivative size = 156

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{B \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d} - \frac{B \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d} - \frac{B \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}d} + \frac{B \log\left(1 + \sqrt{2}\sqrt{\tan(c+dx)} + \tan(c+dx)\right)}{2\sqrt{2}d} + \frac{2B \tan^{\frac{3}{2}}(c+dx)}{3d}$$

output

```
-1/2*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)-1/2*B*arctan(1+2^(1/2)
)*tan(d*x+c)^(1/2))/d*2^(1/2)-1/4*B*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+
c))/d*2^(1/2)+1/4*B*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)+2/
3*B*tan(d*x+c)^(3/2)/d
```

3.416.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.57

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{B\left(-3\arctan\left(\sqrt[4]{-\tan^2(c+dx)}\right)\sqrt[4]{-\tan(c+dx)}+3\operatorname{arctanh}\left(\sqrt[4]{-\tan^2(c+dx)}\right)\sqrt[4]{-\tan(c+dx)}+2\tan^{\frac{7}{4}}(c+dx)\right)}{3d\sqrt[4]{\tan(c+dx)}}$$

input `Integrate[(Tan[c + d*x]^(5/2)*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(B*(-3*ArcTan[(-Tan[c + d*x]^2)^(1/4)]*(-Tan[c + d*x])^(1/4) + 3*ArcTanh[(-Tan[c + d*x]^2)^(1/4)]*(-Tan[c + d*x])^(1/4) + 2*Tan[c + d*x]^(7/4)))/(3*d*Tan[c + d*x]^(1/4))`

3.416.3 Rubi [A] (verified)Time = 0.41 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {2011, 3042, 3954, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \tan^{\frac{5}{2}}(c+dx) dx$$

$$\downarrow \text{3042}$$

$$B \int \tan(c+dx)^{5/2} dx$$

$$\downarrow \text{3954}$$

$$B \left(\frac{2 \tan^{\frac{3}{2}}(c+dx)}{3d} - \int \sqrt{\tan(c+dx)} dx \right)$$

3.416. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& B \left(\frac{2 \tan^{\frac{3}{2}}(c+dx)}{3d} - \int \sqrt{\tan(c+dx)} dx \right) \\
& \downarrow 3957 \\
& B \left(\frac{2 \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{\int \frac{\sqrt{\tan(c+dx)}}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} \right) \\
& \downarrow 266 \\
& B \left(\frac{2 \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2 \int \frac{\tan(c+dx)}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)}}{d} \right) \\
& \downarrow 826 \\
& B \left(\frac{2 \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2 \left(\frac{1}{2} \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)} - \frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)} \right)}{d} \right) \\
& \downarrow 1476 \\
& B \left(\frac{2 \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2 \left(\frac{1}{2} \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d \sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d \sqrt{\tan(c+dx)} \right) \right)}{d} \right) \\
& \downarrow 1082 \\
& B \left(\frac{2 \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2 \left(\frac{1}{2} \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right) - \frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}}{d} \right) \\
& \downarrow 217 \\
& B \left(\frac{2 \tan^{\frac{3}{2}}(c+dx)}{3d} - \frac{2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)}}{d} \right) \\
& \downarrow 1479
\end{aligned}$$

3.416. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$B \left(\frac{2 \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2 \left(\frac{1}{2} \left(\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d} + \frac{1}{2} \left(\arctan \left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} \right) - \arctan \left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}} \right) \right) \right)$$

↓ 25

$$B \left(\frac{2 \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2 \left(\frac{1}{2} \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right) \right) + \frac{1}{2} \left(\arctan \left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} \right) - \arctan \left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}} \right) \right)$$

↓ 27

$$B \left(\frac{2 \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2 \left(\frac{1}{2} \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right) \right)$$

↓ 1103

$$B \left(\frac{2 \tan^{\frac{3}{2}}(c + dx)}{3d} - \frac{2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) + \frac{1}{2} \left(\frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d}$$

input `Int[(Tan[c + d*x]^(5/2)*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `B*((-2*((-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]/Sqrt[2]])/2 + (Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]))/2)/d + (2*Tan[c + d*x]^(3/2))/(3*d))`

3.416.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

3.416.
$$\int \frac{\tan^5(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

3.416.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.65

method	result
derivativedivides	$B \frac{\left(\frac{2 \tan^{\frac{3}{2}}(dx+c)}{3} - \sqrt{2} \ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right)}{d}$
default	$B \frac{\left(\frac{2 \tan^{\frac{3}{2}}(dx+c)}{3} - \sqrt{2} \ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right)}{d}$

input `int(tan(d*x+c)^(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

$$3.416. \int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

output $1/d*B*(2/3*\tan(d*x+c)^{(3/2)}-1/4*2^{(1/2)}*(\ln((1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}))$

3.416.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.18

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{4B\tan(dx+c)^{\frac{3}{2}} - 3d\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log\left(d^3\left(-\frac{B^4}{d^4}\right)^{\frac{3}{4}} + B^3\sqrt{\tan(dx+c)}\right) + 3id\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log\left(id^3\left(-\frac{B^4}{d^4}\right)^{\frac{3}{4}} + B^3\sqrt{\tan(dx+c)}\right)}{d}$$

input `integrate(tan(d*x+c)^(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output $1/6*(4*B*\tan(d*x+c)^{(3/2)} - 3*d*(-B^4/d^4)^{(1/4)}*\log(d^3*(-B^4/d^4)^{(3/4)} + B^3*\sqrt{\tan(d*x+c)}) + 3*I*d*(-B^4/d^4)^{(1/4)}*\log(I*d^3*(-B^4/d^4)^{(3/4)} + B^3*\sqrt{\tan(d*x+c)}) - 3*I*d*(-B^4/d^4)^{(1/4)}*\log(-I*d^3*(-B^4/d^4)^{(3/4)} + B^3*\sqrt{\tan(d*x+c)}) + 3*d*(-B^4/d^4)^{(1/4)}*\log(-d^3*(-B^4/d^4)^{(3/4)} + B^3*\sqrt{\tan(d*x+c)})))/d$

3.416.6 Sympy [F]

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = B \int \tan^{\frac{5}{2}}(c+dx) dx$$

input `integrate(tan(d*x+c)**(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `B*Integral(tan(c+d*x)**(5/2),x)`

3.416.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.79

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{8B\tan(dx+c)^{\frac{3}{2}} - 3\left(2\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}+2\sqrt{\tan(dx+c)}\right)\right) + 2\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2}-2\sqrt{\tan(dx+c)}\right)\right)\right)}{d}$$

input `integrate(tan(d*x+c)^(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/12*(8*B*tan(d*x + c)^(3/2) - 3*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*B)/d`

3.416.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.416.9 Mupad [B] (verification not implemented)

Time = 13.96 (sec) , antiderivative size = 16727, normalized size of antiderivative = 107.22

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

input `int((tan(c + d*x)^(5/2)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `atan((((((((32*(4*B*a^2*b^8*d^4 + 8*B*a^4*b^6*d^4 + 4*B*a^6*b^4*d^4))/(b*d^5) - (32*tan(c + d*x)^(1/2)*(((64*B^4*a^6*b^2*d^4 - B^4*a^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a^3*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4))))^(1/2)*(16*b^10*d^4 + 16*a^2*b^8*d^4 - 16*a^4*b^6*d^4 - 16*a^6*b^4*d^4))/(b*d^4))*(((64*B^4*a^6*b^2*d^4 - B^4*a^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a^3*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4))))^(1/2) + (32*tan(c + d*x)^(1/2)*(4*B^2*a^5*b^5*d^2 - 14*B^2*a^3*b^7*d^2 + 2*B^2*a^7*b^3*d^2 + 16*B^2*a^9*b*d^2))/(b*d^4))*(((64*B^4*a^6*b^2*d^4 - B^4*a^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a^3*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4))))^(1/2) - (32*(B^3*a^5*b^5*d^2 - 15*B^3*a^7*b^3*d^2 + 12*B^3*a^9*b*d^2))/(b*d^5))*(((64*B^4*a^6*b^2*d^4 - B^4*a^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a^3*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4))))^(1/2) + (32*tan(c + d*x)^(1/2)*(2*B^4*a^10 - B^4*a^4*b^6))/(b*d^4))*(((64*B^4*a^6*b^2*d^4 - B^4*a^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a^3*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4))))^(1/2) - 8*B^2*a^3*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4))))^(1/2)*1i - (((((32*(4*B*a^2*b^8*d^4 + 8*B*a^4*b^6*d^4 + 4*B*a^6*b^4*d^4))/(b*d^5) + (32*tan(c + d*x)^(1/2)*(((64*B^4*a^6*b^2*d^4 - B^4*a^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a^3*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4))))^(1/2)*(16*b^10*d^4 + 16*a^2*b^8*d^4 - 16*a^4*b^6*d^4 - 16*a^...`

3.416.
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.417
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.417.1 Optimal result 4031
 3.417.2 Mathematica [A] (verified) 4032
 3.417.3 Rubi [A] (verified) 4032
 3.417.4 Maple [A] (verified) 4036
 3.417.5 Fricas [C] (verification not implemented) 4037
 3.417.6 Sympy [F] 4037
 3.417.7 Maxima [A] (verification not implemented) 4038
 3.417.8 Giac [F(-1)] 4038
 3.417.9 Mupad [B] (verification not implemented) 4038

3.417.1 Optimal result

Integrand size = 36, antiderivative size = 154

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{B \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d} - \frac{B \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d} + \frac{B \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}d} - \frac{B \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}d} + \frac{2B\sqrt{\tan(c+dx)}}{d}$$

output

```
-1/2*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)-1/2*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)+1/4*B*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)-1/4*B*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)+2*B*tan(d*x+c)^(1/2)/d
```

3.417.2 Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.90

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{B \left(\frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1+\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} + \frac{\log\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)}{2\sqrt{2}}\right)}{2\sqrt{2}} - \frac{\log\left(\frac{1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)}{2\sqrt{2}}\right)}{2\sqrt{2}} \right)}{d}$$

input `Integrate[(Tan[c + d*x]^(3/2)*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(B*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2] + Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]) + 2*Sqrt[Tan[c + d*x]]))/d`

3.417.3 Rubi [A] (verified)Time = 0.41 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {2011, 3042, 3954, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \tan^{\frac{3}{2}}(c+dx) dx$$

$$\downarrow \text{3042}$$

$$B \int \tan(c+dx)^{3/2} dx$$

$$\downarrow \text{3954}$$

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{d} - \int \frac{1}{\sqrt{\tan(c+dx)}} dx \right)$$

3.417. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& B \left(\frac{2\sqrt{\tan(c+dx)}}{d} - \int \frac{1}{\sqrt{\tan(c+dx)}} dx \right) \\
& \downarrow 3957 \\
& B \left(\frac{2\sqrt{\tan(c+dx)}}{d} - \frac{\int \frac{1}{\sqrt{\tan(c+dx)}(\tan^2(c+dx)+1)} d \tan(c+dx)}{d} \right) \\
& \downarrow 266 \\
& B \left(\frac{2\sqrt{\tan(c+dx)}}{d} - \frac{2 \int \frac{1}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)}}{d} \right) \\
& \downarrow 755 \\
& B \left(\frac{2\sqrt{\tan(c+dx)}}{d} - \frac{2 \left(\frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)} \right)}{d} \right) \\
& \downarrow 1476 \\
& B \left(\frac{2\sqrt{\tan(c+dx)}}{d} - \frac{2 \left(\frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)} + \frac{1}{2} \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d \sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d \sqrt{\tan(c+dx)} \right) \right)}{d} \right) \\
& \downarrow 1082 \\
& B \left(\frac{2\sqrt{\tan(c+dx)}}{d} - \frac{2 \left(\frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)} + \frac{1}{2} \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \right) \\
& \downarrow 217 \\
& B \left(\frac{2\sqrt{\tan(c+dx)}}{d} - \frac{2 \left(\frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)} + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d} \right) \\
& \downarrow 1479
\end{aligned}$$

3.417. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{d} - \frac{2 \left(\frac{1}{2} \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d} \right) + \frac{1}{2} \left(\arctan \left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} \right) - \arctan \left(\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} \right) \right)$$

↓ 25

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{d} - \frac{2 \left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d} \right) + \frac{1}{2} \left(\arctan \left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} \right) - \arctan \left(\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} \right) \right)$$

↓ 27

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{d} - \frac{2 \left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d} \right) + \frac{1}{2} \left(\arctan \left(\frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} \right) - \arctan \left(\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} \right) \right)$$

↓ 1103

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{d} - \frac{2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d} \right) + \frac{1}{2} \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right)$$

input `Int[(Tan[c + d*x]^(3/2)*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `B*((-2*((-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 + (-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]))/2)/d + (2*Sqrt[Tan[c + d*x]])/d)`

3.417.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

3.417.
$$\int \frac{\tan^3(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

3.417.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.66

method	result
derivativedivides	$B \left(2(\sqrt{\tan(dx+c)} - \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right)}{4} \right)$
default	$B \left(2(\sqrt{\tan(dx+c)} - \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right)}{4} \right)$

input `int(tan(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

$$3.417. \int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

output $1/d*B*(2*\tan(d*x+c)^{(1/2)}-1/4*2^{(1/2)}*(\ln((1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}))$

3.417.5 Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.08

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx =$$

$$d\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log\left(B\sqrt{\tan(dx+c)}+d\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}}\right) + i d\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log\left(B\sqrt{\tan(dx+c)}+i d\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}}\right) - i$$

input `integrate(tan(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output $-1/2*(d*(-B^4/d^4)^{(1/4)}*\log(B*\sqrt{\tan(d*x+c)})+d*(-B^4/d^4)^{(1/4)})+I*d*(-B^4/d^4)^{(1/4)}*\log(B*\sqrt{\tan(d*x+c)})+I*d*(-B^4/d^4)^{(1/4)}-I*d*(-B^4/d^4)^{(1/4)}*\log(B*\sqrt{\tan(d*x+c)})-I*d*(-B^4/d^4)^{(1/4)}-d*(-B^4/d^4)^{(1/4)}*\log(B*\sqrt{\tan(d*x+c)})-d*(-B^4/d^4)^{(1/4)}-4*B*\sqrt{\tan(d*x+c)})/d$

3.417.6 Sympy [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = B \int \tan^{\frac{3}{2}}(c+dx) dx$$

input `integrate(tan(d*x+c)**(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `B*Integral(tan(c+d*x)**(3/2),x)`

3.417. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$

3.417.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.80

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx =$$

$$\frac{2\sqrt{2}B \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}+2\sqrt{\tan(dx+c)}\right)\right) + 2\sqrt{2}B \arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2}-2\sqrt{\tan(dx+c)}\right)\right) + \sqrt{2}B \log(\sqrt{2}\sqrt{\tan(dx+c)} + \tan(dx+c) + 1) - \sqrt{2}B \log(-\sqrt{2}\sqrt{\tan(dx+c)} + \tan(dx+c) + 1) - 8B\sqrt{\tan(dx+c)}}{d}$$

input `integrate(tan(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `-1/4*(2*sqrt(2)*B*arctan(1/2*sqrt(2)*(sqrt(2)+2*sqrt(tan(d*x+c))))+2*sqrt(2)*B*arctan(-1/2*sqrt(2)*(sqrt(2)-2*sqrt(tan(d*x+c))))+sqrt(2)*B*log(sqrt(2)*sqrt(tan(d*x+c))+tan(d*x+c)+1)-sqrt(2)*B*log(-sqrt(2)*sqrt(tan(d*x+c))+tan(d*x+c)+1)-8*B*sqrt(tan(d*x+c)))/d`

3.417.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.417.9 Mupad [B] (verification not implemented)

Time = 13.83 (sec) , antiderivative size = 16060, normalized size of antiderivative = 104.29

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

input `int((tan(c + d*x)^(3/2)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `atan((((32*(B^3*a^2*b^7*d^2 - 15*B^3*a^4*b^5*d^2 + 12*B^3*a^6*b^3*d^2))/d^5 - (((32*(4*B*a*b^8*d^4 + 8*B*a^3*b^6*d^4 + 4*B*a^5*b^4*d^4))/d^5 - (32*tan(c + d*x)^(1/2)*(((64*B^4*a^2*b^6*d^4 - B^4*b^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b^3*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4))))^(1/2)*(16*b^9*d^4 + 16*a^2*b^7*d^4 - 16*a^4*b^5*d^4 - 16*a^6*b^3*d^4))/d^4)*(((64*B^4*a^2*b^6*d^4 - B^4*b^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b^3*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) + (32*tan(c + d*x)^(1/2)*(4*B^2*a^3*b^6*d^2 + 2*B^2*a^5*b^4*d^2 + 16*B^2*a^7*b^2*d^2 - 14*B^2*a*b^8*d^2))/d^4)*(((64*B^4*a^2*b^6*d^4 - B^4*b^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b^3*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2))*(((64*B^4*a^2*b^6*d^4 - B^4*b^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b^3*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) + (32*tan(c + d*x)^(1/2)*(B^4*b^9 - 2*B^4*a^6*b^3))/d^4)*(((64*B^4*a^2*b^6*d^4 - B^4*b^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b^3*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*1i - (((32*(B^3*a^2*b^7*d^2 - 15*B^3*a^4*b^5*d^2 + 12*B^3*a^6*b^3*d^2))/d^5 - (((32*(4*B*a*b^8*d^4 + 8*B*a^3*b^6*d^4 + 4*B*a^5*b^4*d^4))/d^5 + (32*tan(c + d*x)^(1/2)*(((64*B^4*a^2*b^6*d^4 - B^4*b^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b^3*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*(16*b^9*d^4...`

3.417. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.418
$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.418.1 Optimal result 4040
 3.418.2 Mathematica [A] (verified) 4041
 3.418.3 Rubi [A] (verified) 4041
 3.418.4 Maple [A] (verified) 4045
 3.418.5 Fricas [C] (verification not implemented) 4045
 3.418.6 Sympy [F] 4046
 3.418.7 Maxima [A] (verification not implemented) 4046
 3.418.8 Giac [F(-1)] 4047
 3.418.9 Mupad [B] (verification not implemented) 4047

3.418.1 Optimal result

Integrand size = 36, antiderivative size = 138

$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = -\frac{B \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d} + \frac{B \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}d} + \frac{B \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}d} - \frac{B \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}d}$$

output `1/2*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)+1/2*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)+1/4*B*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)-1/4*B*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)`

3.418.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.43

$$\int \frac{\sqrt{\tan(c+dx)}(aB + bB \tan(c+dx))}{a + b \tan(c+dx)} dx$$

$$= \frac{B \left(\arctan \left(\sqrt[4]{-\tan^2(c+dx)} \right) - \operatorname{arctanh} \left(\sqrt[4]{-\tan^2(c+dx)} \right) \right) \sqrt[4]{-\tan(c+dx)}}{d \sqrt[4]{\tan(c+dx)}}$$

input `Integrate[(Sqrt[Tan[c + d*x]]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(B*(ArcTan[(-Tan[c + d*x]^2)^(1/4)] - ArcTanh[(-Tan[c + d*x]^2)^(1/4)])*(-Tan[c + d*x])^(1/4))/(d*Tan[c + d*x]^(1/4))`

3.418.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2011, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\tan(c+dx)}(aB + bB \tan(c+dx))}{a + b \tan(c+dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \sqrt{\tan(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$B \int \sqrt{\tan(c+dx)} dx$$

$$\downarrow \text{3957}$$

$$B \int \frac{\sqrt{\tan(c+dx)}}{\tan^2(c+dx)+1} d \tan(c+dx)$$

$$\downarrow \text{266}$$

3.418. $\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\frac{2B \int \frac{\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d}$$

↓ 826

$$\frac{2B \left(\frac{1}{2} \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d}$$

↓ 1476

$$\frac{2B \left(\frac{1}{2} \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d}$$

↓ 1082

$$\frac{2B \left(\frac{1}{2} \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d}$$

↓ 217

$$\frac{2B \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d}$$

↓ 1479

$$\frac{2B \left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{d}$$

↓ 25

$$\frac{2B \left(\frac{1}{2} \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{d}$$

↓ 27

$$\frac{2B \left(\frac{1}{2} \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{d}$$

↓ 1103

3.418. $\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$2B \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)-1})}{2\sqrt{2}} \right) \right) / d$$

input `Int[(Sqrt[Tan[c + d*x]]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(2*B*((-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])/2 + (Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]))/2)/d`

3.418.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2)]^p, x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntegerBinomialQ[a, b, c, 2, m, p, x]`

rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

3.418.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.65

method	result
derivativedivides	$\frac{B\sqrt{2} \left(\ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right)}{4d}$
default	$\frac{B\sqrt{2} \left(\ln \left(\frac{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right)}{4d}$

input `int(tan(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/4/d*B*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))`

3.418.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.19

$$\begin{aligned} & \int \frac{\sqrt{\tan(c+dx)}(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx \\ &= \frac{1}{2} \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \log \left(d^3 \left(-\frac{B^4}{d^4} \right)^{\frac{3}{4}} + B^3 \sqrt{\tan(dx+c)} \right) \\ & \quad - \frac{1}{2} i \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \log \left(i d^3 \left(-\frac{B^4}{d^4} \right)^{\frac{3}{4}} + B^3 \sqrt{\tan(dx+c)} \right) \\ & \quad + \frac{1}{2} i \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \log \left(-i d^3 \left(-\frac{B^4}{d^4} \right)^{\frac{3}{4}} + B^3 \sqrt{\tan(dx+c)} \right) \\ & \quad - \frac{1}{2} \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \log \left(-d^3 \left(-\frac{B^4}{d^4} \right)^{\frac{3}{4}} + B^3 \sqrt{\tan(dx+c)} \right) \end{aligned}$$

input `integrate(tan(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,algorithm="fracas")`

```
output 1/2*(-B^4/d^4)^(1/4)*log(d^3*(-B^4/d^4)^(3/4) + B^3*sqrt(tan(d*x + c))) -
1/2*I*(-B^4/d^4)^(1/4)*log(I*d^3*(-B^4/d^4)^(3/4) + B^3*sqrt(tan(d*x + c))
) + 1/2*I*(-B^4/d^4)^(1/4)*log(-I*d^3*(-B^4/d^4)^(3/4) + B^3*sqrt(tan(d*x
+ c))) - 1/2*(-B^4/d^4)^(1/4)*log(-d^3*(-B^4/d^4)^(3/4) + B^3*sqrt(tan(d*x
+ c)))
```

3.418.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = B \int \sqrt{\tan(c+dx)} dx$$

```
input integrate(tan(d*x+c)**(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x)
```

```
output B*Integral(sqrt(tan(c + d*x)), x)
```

3.418.7 Maxima [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.79

$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{\left(2\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}+2\sqrt{\tan(dx+c)}\right)\right)\right) + 2\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2}-2\sqrt{\tan(dx+c)}\right)\right) - \sqrt{2}\log\left(\frac{\sqrt{2}\sqrt{\tan(dx+c)} + \tan(dx+c) + 1}{\sqrt{2}\sqrt{\tan(dx+c)} - \tan(dx+c) + 1}\right)}{4d}$$

```
input integrate(tan(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")
```

```
output 1/4*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*B/d
```

3.418.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output Timed out

3.418.9 Mupad [B] (verification not implemented)

Time = 13.38 (sec) , antiderivative size = 15753, normalized size of antiderivative = 114.15

$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

input `int((tan(c+d*x)^(1/2)*(B*a+B*b*tan(c+d*x)))/(a+b*tan(c+d*x)),x)`

output `atan((((32*(13*B^3*a^5*b^4*d^2+B^3*a^7*b^2*d^2))/d^5+((32*(12*B*a^2*b^7*d^4+24*B*a^4*b^5*d^4+12*B*a^6*b^3*d^4))/d^5-(32*tan(c+d*x)^(1/2)*(((64*B^4*a^6*b^2*d^4-B^4*a^4*(16*a^4*d^4+16*b^4*d^4+32*a^2*b^2*d^4))^(1/2)-8*B^2*a^3*b*d^2)/(16*(a^4*d^4+b^4*d^4+2*a^2*b^2*d^4)))^(1/2)*(16*b^9*d^4+16*a^2*b^7*d^4-16*a^4*b^5*d^4-16*a^6*b^3*d^4))/d^4)*(((64*B^4*a^6*b^2*d^4-B^4*a^4*(16*a^4*d^4+16*b^4*d^4+32*a^2*b^2*d^4))^(1/2)-8*B^2*a^3*b*d^2)/(16*(a^4*d^4+b^4*d^4+2*a^2*b^2*d^4)))^(1/2)+((32*tan(c+d*x)^(1/2)*(20*B^2*a^5*b^4*d^2-14*B^2*a^3*b^6*d^2+2*B^2*a^7*b^2*d^2))/d^4)*(((64*B^4*a^6*b^2*d^4-B^4*a^4*(16*a^4*d^4+16*b^4*d^4+32*a^2*b^2*d^4))^(1/2)-8*B^2*a^3*b*d^2)/(16*(a^4*d^4+b^4*d^4+2*a^2*b^2*d^4)))^(1/2)*(((64*B^4*a^6*b^2*d^4-B^4*a^4*(16*a^4*d^4+16*b^4*d^4+32*a^2*b^2*d^4))^(1/2)-8*B^2*a^3*b*d^2)/(16*(a^4*d^4+b^4*d^4+2*a^2*b^2*d^4)))^(1/2)-(32*tan(c+d*x)^(1/2)*(B^4*a^4*b^5-2*B^4*a^6*b^3))/d^4)*(((64*B^4*a^6*b^2*d^4-B^4*a^4*(16*a^4*d^4+16*b^4*d^4+32*a^2*b^2*d^4))^(1/2)-8*B^2*a^3*b*d^2)/(16*(a^4*d^4+b^4*d^4+2*a^2*b^2*d^4)))^(1/2)*i1-((((32*(13*B^3*a^5*b^4*d^2+B^3*a^7*b^2*d^2))/d^5+((32*(12*B*a^2*b^7*d^4+24*B*a^4*b^5*d^4+12*B*a^6*b^3*d^4))/d^5+(32*tan(c+d*x)^(1/2)*(((64*B^4*a^6*b^2*d^4-B^4*a^4*(16*a^4*d^4+16*b^4*d^4+32*a^2*b^2*d^4))^(1/2)-8*B^2*a^3*b*d^2)/(16*(a^4*d^4+b^4*d^4+2*a^2*b^2*d^4)))^(1/2)*(16*b^9*d^4+16*a^2*b^7*d^4-16*a^4*b^5*d^4-16*a^6*b^...`

$$3.419 \quad \int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx$$

3.419.1 Optimal result	4048
3.419.2 Mathematica [A] (verified)	4049
3.419.3 Rubi [A] (verified)	4049
3.419.4 Maple [A] (verified)	4053
3.419.5 Fricas [C] (verification not implemented)	4053
3.419.6 Sympy [F]	4054
3.419.7 Maxima [A] (verification not implemented)	4054
3.419.8 Giac [F(-1)]	4055
3.419.9 Mupad [B] (verification not implemented)	4055

3.419.1 Optimal result

Integrand size = 36, antiderivative size = 138

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx = -\frac{B \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d} + \frac{B \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}d} - \frac{B \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d} + \frac{B \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}d}$$

output `1/2*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)+1/2*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)-1/4*B*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)+1/4*B*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)`

3.419.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.80

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx$$

$$= \frac{B \left(-2 \arctan \left(1 - \sqrt{2} \sqrt{\tan(c + dx)} \right) + 2 \arctan \left(1 + \sqrt{2} \sqrt{\tan(c + dx)} \right) - \log \left(1 - \sqrt{2} \sqrt{\tan(c + dx)} \right) + \log \left(1 + \sqrt{2} \sqrt{\tan(c + dx)} \right) \right)}{2\sqrt{2}d}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])),x]`

output `(B*(-2*ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] + 2*ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]] - Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]))/(2*Sqrt[2]*d)`

3.419.3 Rubi [A] (verified)Time = 0.34 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2011, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx$$

$$\downarrow \text{2011}$$

$$B \int \frac{1}{\sqrt{\tan(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$B \int \frac{1}{\sqrt{\tan(c + dx)}} dx$$

$$\downarrow \text{3957}$$

$$\frac{B \int \frac{1}{\sqrt{\tan(c+dx)}(\tan^2(c+dx)+1)} d \tan(c + dx)}{d}$$

3.419. $\int \frac{aB+bB \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx$

$$\begin{aligned}
& \downarrow 266 \\
& \frac{2B \int \frac{1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} \\
& \downarrow 755 \\
& \frac{2B \left(\frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d} \\
& \downarrow 1476 \\
& \frac{2B \left(\frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d} \\
& \downarrow 1082 \\
& \frac{2B \left(\frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \\
& \downarrow 217 \\
& \frac{2B \left(\frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d} \\
& \downarrow 1479 \\
& \frac{2B \left(\frac{1}{2} \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \\
& \downarrow 25 \\
& \frac{2B \left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \\
& \downarrow 27 \\
& \frac{2B \left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) \right)}{d}
\end{aligned}$$

3.419. $\int \frac{aB+bB \tan(c+dx)}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))}} dx$

↓ 1103

$$2B \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)})}{2\sqrt{2}} \right) \right) / d$$

input `Int[(a*B + b*B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])),x]`

output `(2*B*((-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2])/2 + (-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]))/2)/d`

3.419.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplrQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

3.419.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.65

method	result
derivativedivides	$\frac{B\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4d}$
default	$\frac{B\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})) \right)}{4d}$

input `int((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output $1/4/d*B*2^{(1/2)}*(\ln((1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}))$

3.419.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.07

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx = \frac{1}{2} \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \log \left(B\sqrt{\tan(dx + c)} + d \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \right) + \frac{1}{2} i \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \log \left(B\sqrt{\tan(dx + c)} + i d \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \right) - \frac{1}{2} i \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \log \left(B\sqrt{\tan(dx + c)} - i d \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \right) - \frac{1}{2} \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \log \left(B\sqrt{\tan(dx + c)} - d \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \right)$$

3.419. $\int \frac{aB+bB \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output $\frac{1}{2}*(-B^4/d^4)^{(1/4)}*\log(B*\sqrt{\tan(dx + c)} + d*(-B^4/d^4)^{(1/4)}) + \frac{1}{2}*I*(-B^4/d^4)^{(1/4)}*\log(B*\sqrt{\tan(dx + c)} + I*d*(-B^4/d^4)^{(1/4)}) - \frac{1}{2}*I*(-B^4/d^4)^{(1/4)}*\log(B*\sqrt{\tan(dx + c)} - I*d*(-B^4/d^4)^{(1/4)}) - \frac{1}{2}*(-B^4/d^4)^{(1/4)}*\log(B*\sqrt{\tan(dx + c)} - d*(-B^4/d^4)^{(1/4)})$

3.419.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx = B \int \frac{1}{\sqrt{\tan(c + dx)}} dx$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+b*tan(d*x+c)),x)`

output `B*Integral(1/sqrt(tan(c + d*x)), x)`

3.419.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.81

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx = \frac{2\sqrt{2}B \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(dx + c)}\right)\right) + 2\sqrt{2}B \arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2} - 2\sqrt{\tan(dx + c)}\right)\right) + \sqrt{2}}{4d}$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output $\frac{1}{4}*(2*\sqrt{2}*B*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2*\sqrt{\tan(dx + c)}))) + 2*\sqrt{2}*B*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2*\sqrt{\tan(dx + c)}))) + \sqrt{2}*B*\log(\sqrt{2}*\sqrt{\tan(dx + c)} + \tan(dx + c) + 1) - \sqrt{2}*B*\log(-\sqrt{2}*\sqrt{\tan(dx + c)} + \tan(dx + c) + 1))/d$

3.419. $\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx$

3.419.8 Giac [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx = \text{Timed out}$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x, algorithm="giac")`

output Timed out

3.419.9 Mupad [B] (verification not implemented)

Time = 13.81 (sec) , antiderivative size = 15437, normalized size of antiderivative = 111.86

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx = \text{Too large to display}$$

input `int((B*a + B*b*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + b*tan(c + d*x))),x)`

output `atan((((32*(13*B^3*a^2*b^7*d^2 + B^3*a^4*b^5*d^2))/d^5 + (((32*(12*B*a*b^8*d^4 + 24*B*a^3*b^6*d^4 + 12*B*a^5*b^4*d^4))/d^5 - (32*tan(c + d*x)^(1/2) * (((64*B^4*a^2*b^6*d^4 - B^4*b^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b^3*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) * (16*b^9*d^4 + 16*a^2*b^7*d^4 - 16*a^4*b^5*d^4 - 16*a^6*b^3*d^4))/d^4) * (((64*B^4*a^2*b^6*d^4 - B^4*b^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b^3*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) + (32*tan(c + d*x)^(1/2)*(20*B^2*a^3*b^6*d^2 + 2*B^2*a^5*b^4*d^2 - 14*B^2*a*b^8*d^2))/d^4) * (((64*B^4*a^2*b^6*d^4 - B^4*b^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b^3*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) * (((64*B^4*a^2*b^6*d^4 - B^4*b^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b^3*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) - (32*tan(c + d*x)^(1/2)*(B^4*b^9 - 2*B^4*a^2*b^7))/d^4) * (((64*B^4*a^2*b^6*d^4 - B^4*b^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b^3*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) * (((64*B^4*a^2*b^6*d^4 - B^4*b^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a*b^3*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) * (16*b^9*d^4 + 16*a^2*b^7*d^4 - 16*a^4*b^5*d^4 - 16*a^6*b^3*d^4))/d^...`

3.419. $\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx$

3.420
$$\int \frac{aB+bB \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$$

3.420.1 Optimal result 4056
 3.420.2 Mathematica [A] (verified) 4057
 3.420.3 Rubi [A] (verified) 4057
 3.420.4 Maple [A] (verified) 4061
 3.420.5 Fricas [C] (verification not implemented) 4062
 3.420.6 Sympy [F] 4062
 3.420.7 Maxima [A] (verification not implemented) 4063
 3.420.8 Giac [F(-1)] 4063
 3.420.9 Mupad [B] (verification not implemented) 4063

3.420.1 Optimal result

Integrand size = 36, antiderivative size = 154

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = \frac{B \arctan \left(1 - \sqrt{2} \sqrt{\tan(c + dx)} \right)}{\sqrt{2}d} - \frac{B \arctan \left(1 + \sqrt{2} \sqrt{\tan(c + dx)} \right)}{\sqrt{2}d} - \frac{B \log \left(1 - \sqrt{2} \sqrt{\tan(c + dx)} + \tan(c + dx) \right)}{2\sqrt{2}d} + \frac{B \log \left(1 + \sqrt{2} \sqrt{\tan(c + dx)} + \tan(c + dx) \right)}{2\sqrt{2}d} - \frac{2B}{d\sqrt{\tan(c + dx)}}$$

output

```
-1/2*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)-1/2*B*arctan(1+2^(1/2)
)*tan(d*x+c)^(1/2))/d*2^(1/2)-1/4*B*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+
c))/d*2^(1/2)+1/4*B*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)-2*
B/d/tan(d*x+c)^(1/2)
```

3.420.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.51

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$= \frac{B \left(-2 - \arctan \left(\sqrt[4]{-\tan^2(c + dx)} \right) \sqrt[4]{-\tan^2(c + dx)} + \operatorname{arctanh} \left(\sqrt[4]{-\tan^2(c + dx)} \right) \sqrt[4]{-\tan^2(c + dx)} \right)}{d \sqrt{\tan(c + dx)}}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])),x]`

output `(B*(-2 - ArcTan[(-Tan[c + d*x]^2)^(1/4)]*(-Tan[c + d*x]^2)^(1/4) + ArcTanh[(-Tan[c + d*x]^2)^(1/4)]*(-Tan[c + d*x]^2)^(1/4)))/(d*Sqrt[Tan[c + d*x]])`

3.420.3 Rubi [A] (verified)Time = 0.41 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {2011, 3042, 3955, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$\downarrow \text{2011}$$

$$B \int \frac{1}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$B \int \frac{1}{\tan(c + dx)^{3/2}} dx$$

$$\downarrow \text{3955}$$

$$B \left(- \int \sqrt{\tan(c + dx)} dx - \frac{2}{d \sqrt{\tan(c + dx)}} \right)$$

$$\downarrow \text{3042}$$

3.420. $\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$

$$\begin{aligned}
& B\left(-\int \sqrt{\tan(c+dx)}dx - \frac{2}{d\sqrt{\tan(c+dx)}}\right) \\
& \quad \downarrow 3957 \\
& B\left(-\frac{\int \frac{\sqrt{\tan(c+dx)}}{\tan^2(c+dx)+1}d\tan(c+dx)}{d} - \frac{2}{d\sqrt{\tan(c+dx)}}\right) \\
& \quad \downarrow 266 \\
& B\left(-\frac{2\int \frac{\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}}{d} - \frac{2}{d\sqrt{\tan(c+dx)}}\right) \\
& \quad \downarrow 826 \\
& B\left(-\frac{2\left(\frac{1}{2}\int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)} - \frac{1}{2}\int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}\right)}{d} - \frac{2}{d\sqrt{\tan(c+dx)}}\right) \\
& \quad \downarrow 1476 \\
& B\left(-\frac{2\left(\frac{1}{2}\left(\frac{1}{2}\int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)} + \frac{1}{2}\int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}d\sqrt{\tan(c+dx)}\right) - \frac{1}{2}\int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}\right)}{d}\right) \\
& \quad \downarrow 1082 \\
& B\left(-\frac{2\left(\frac{1}{2}\left(\frac{\int \frac{1}{-\tan(c+dx)-1}d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1}d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}}\right) - \frac{1}{2}\int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}\right)}{d}\right) \\
& \quad \downarrow 217 \\
& B\left(-\frac{2\left(\frac{1}{2}\left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}}\right) - \frac{1}{2}\int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}\right)}{d} - \frac{2}{d\sqrt{\tan(c+dx)}}\right) \\
& \quad \downarrow 1479
\end{aligned}$$

3.420. $\int \frac{aB+bB \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right)}{d} \right)$$

↓ 25

$$B \left(\frac{2 \left(\frac{1}{2} \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right)}{d} \right)$$

↓ 27

$$B \left(\frac{2 \left(\frac{1}{2} \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right)}{d} \right)$$

↓ 1103

$$B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d} \right)$$

input `Int[(a*B + b*B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])),x]`

output `B*((-2*((-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])/2 + (Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]))/2)/d - 2/(d*Sqrt[Tan[c + d*x]]))`

3.420.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplrQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3955 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x])^(n + 1)/(b*d*(n + 1)), x] - Simp[1/b^2 Int[(b*Tan[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

3.420.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.66

method	result
derivativedivides	$B \left(\frac{\sqrt{2} \left(\ln \left(\frac{1 - \sqrt{2} (\sqrt{\tan(dx+c)} + \tan(dx+c))}{1 + \sqrt{2} (\sqrt{\tan(dx+c)} + \tan(dx+c))} \right) + 2 \arctan(1 + \sqrt{2} (\sqrt{\tan(dx+c)}) + 2 \arctan(-1 + \sqrt{2} (\sqrt{\tan(dx+c)})) \right)}{4} - \frac{2}{\sqrt{\tan(dx+c)}} \right) / d$
default	$B \left(\frac{\sqrt{2} \left(\ln \left(\frac{1 - \sqrt{2} (\sqrt{\tan(dx+c)} + \tan(dx+c))}{1 + \sqrt{2} (\sqrt{\tan(dx+c)} + \tan(dx+c))} \right) + 2 \arctan(1 + \sqrt{2} (\sqrt{\tan(dx+c)}) + 2 \arctan(-1 + \sqrt{2} (\sqrt{\tan(dx+c)})) \right)}{4} - \frac{2}{\sqrt{\tan(dx+c)}} \right) / d$

input `int((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

$$3.420. \quad \int \frac{aB+bB \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$$

output $1/d*B*(-1/4*2^{(1/2)}*(\ln((1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}))-2/\tan(d*x+c)^{(1/2)})$

3.420.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.40

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx =$$

$$\frac{d\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log\left(d^3\left(-\frac{B^4}{d^4}\right)^{\frac{3}{4}} + B^3\sqrt{\tan(dx + c)}\right) \tan(dx + c) - i d\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log\left(i d^3\left(-\frac{B^4}{d^4}\right)^{\frac{3}{4}} + B^3\sqrt{\tan(dx + c)}\right)}{d^2}$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output $-1/2*(d*(-B^4/d^4)^{(1/4)}*\log(d^3*(-B^4/d^4)^{(3/4)} + B^3*\sqrt{\tan(d*x + c)})*\tan(d*x + c) - I*d*(-B^4/d^4)^{(1/4)}*\log(I*d^3*(-B^4/d^4)^{(3/4)} + B^3*\sqrt{\tan(d*x + c)}))*\tan(d*x + c) + I*d*(-B^4/d^4)^{(1/4)}*\log(-I*d^3*(-B^4/d^4)^{(3/4)} + B^3*\sqrt{\tan(d*x + c)})*\tan(d*x + c) - d*(-B^4/d^4)^{(1/4)}*\log(-d^3*(-B^4/d^4)^{(3/4)} + B^3*\sqrt{\tan(d*x + c)})*\tan(d*x + c) + 4*B*\sqrt{\tan(d*x + c)})/(d*\tan(d*x + c))$

3.420.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = B \int \frac{1}{\tan^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)**(3/2)/(a+b*tan(d*x+c)),x)`

output `B*Integral(tan(c + d*x)**(-3/2), x)`

3.420. $\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$

3.420.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.79

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx =$$

$$\frac{\left(2\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(dx + c)}\right)\right) + 2\sqrt{2} \arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2} - 2\sqrt{\tan(dx + c)}\right)\right) - \sqrt{2}\right)}{d}$$

```
input integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x, algorithm="maxima")
```

```
output -1/4*((2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*B + 8*B/sqrt(tan(d*x + c)))/d
```

3.420.8 Giac [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = \text{Timed out}$$

```
input integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x, algorithm="giac")
```

```
output Timed out
```

3.420.9 Mupad [B] (verification not implemented)

Time = 13.34 (sec) , antiderivative size = 15569, normalized size of antiderivative = 101.10

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = \text{Too large to display}$$

```
input int((B*a + B*b*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + b*tan(c + d*x))),x)
```

3.420. $\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$

output

$$\begin{aligned} & \operatorname{atan}\left(\left(\tan(c+dx)\right)^{1/2}\left(64B^4a^2b^7d^5-32B^4a^4b^5d^5\right)+\left(\left(64B^4a^6b^2d^4-B^4a^4\left(16a^4d^4+16b^4d^4+32a^2b^2d^4\right)\right)^{1/2}-8B^2a^3bd^2\right)/\left(16\left(a^4d^4+b^4d^4+2a^2b^2d^4\right)\right)\right)^{1/2}\left(\left(\tan(c+dx)\right)^{1/2}\left(128B^2a^5b^4d^7-448B^2a^3b^6d^7+64B^2a^7b^2d^7+512B^2ab^8d^7\right)-\left(\left(64B^4a^6b^2d^4-B^4a^4\left(16a^4d^4+16b^4d^4+32a^2b^2d^4\right)\right)^{1/2}-8B^2a^3bd^2\right)/\left(16\left(a^4d^4+b^4d^4+2a^2b^2d^4\right)\right)\right)^{1/2}\left(\tan(c+dx)\right)^{1/2}\left(\left(64B^4a^6b^2d^4-B^4a^4\left(16a^4d^4+16b^4d^4+32a^2b^2d^4\right)\right)^{1/2}-8B^2a^3bd^2\right)/\left(16\left(a^4d^4+b^4d^4+2a^2b^2d^4\right)\right)\right)^{1/2}\left(512b^9d^9+512a^2b^7d^9-512a^4b^5d^9-512a^6b^3d^9\right)-512Bb^9d^8-640Ba^2b^7d^8+256Ba^4b^5d^8+384Ba^6b^3d^8\right)\left(\left(64B^4a^6b^2d^4-B^4a^4\left(16a^4d^4+16b^4d^4+32a^2b^2d^4\right)\right)^{1/2}-8B^2a^3bd^2\right)/\left(16\left(a^4d^4+b^4d^4+2a^2b^2d^4\right)\right)\right)^{1/2}-32B^3a^5b^4d^6-32B^3a^7b^2d^6+128B^3ab^8d^6\right)\left(\left(64B^4a^6b^2d^4-B^4a^4\left(16a^4d^4+16b^4d^4+32a^2b^2d^4\right)\right)^{1/2}-8B^2a^3bd^2\right)/\left(16\left(a^4d^4+b^4d^4+2a^2b^2d^4\right)\right)\right)^{1/2}\left(\tan(c+dx)\right)^{1/2}\left(64B^4a^2b^7d^5-32B^4a^4b^5d^5\right)+\left(\left(64B^4a^6b^2d^4-B^4a^4\left(16a^4d^4+16b^4d^4+32a^2b^2d^4\right)\right)^{1/2}-8B^2a^3bd^2\right)/\left(16\left(a^4d^4+b^4d^4+2a^2b^2d^4\right)\right)\right)^{1/2}\left(\tan(c+dx)\right)^{1/2}\left(128B^2a^5b^4d^7-448B^2a^3b^6d^7+64B^2a^7b^2d^7+512B^2ab^8d^7\right) \dots \end{aligned}$$

3.421
$$\int \frac{aB+bB \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$$

3.421.1 Optimal result 4065
 3.421.2 Mathematica [A] (verified) 4066
 3.421.3 Rubi [A] (verified) 4066
 3.421.4 Maple [A] (verified) 4070
 3.421.5 Fricas [C] (verification not implemented) 4071
 3.421.6 Sympy [F] 4071
 3.421.7 Maxima [A] (verification not implemented) 4072
 3.421.8 Giac [F(-1)] 4072
 3.421.9 Mupad [B] (verification not implemented) 4072

3.421.1 Optimal result

Integrand size = 36, antiderivative size = 156

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = \frac{B \arctan \left(1 - \sqrt{2} \sqrt{\tan(c + dx)} \right)}{\sqrt{2}d} - \frac{B \arctan \left(1 + \sqrt{2} \sqrt{\tan(c + dx)} \right)}{\sqrt{2}d} + \frac{B \log \left(1 - \sqrt{2} \sqrt{\tan(c + dx)} + \tan(c + dx) \right)}{2\sqrt{2}d} - \frac{B \log \left(1 + \sqrt{2} \sqrt{\tan(c + dx)} + \tan(c + dx) \right)}{2\sqrt{2}d} - \frac{2B}{3d \tan^{\frac{3}{2}}(c + dx)}$$

output

```
-1/2*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/d*2^(1/2)-1/2*B*arctan(1+2^(1/2)
)*tan(d*x+c)^(1/2))/d*2^(1/2)+1/4*B*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+
c))/d*2^(1/2)-1/4*B*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/d*2^(1/2)-2/
3*B/d/tan(d*x+c)^(3/2)
```


3.421.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.53

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$= \frac{B \left(-2 + 3 \arctan \left(\sqrt[4]{-\tan^2(c + dx)} \right) (-\tan^2(c + dx))^{\frac{3}{4}} + 3 \operatorname{arctanh} \left(\sqrt[4]{-\tan^2(c + dx)} \right) (-\tan^2(c + dx))^{\frac{3}{4}} \right)}{3d \tan^{\frac{3}{2}}(c + dx)}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])),x]`

output `(B*(-2 + 3*ArcTan[(-Tan[c + d*x]^2)^(1/4)]*(-Tan[c + d*x]^2)^(3/4) + 3*ArcTanh[(-Tan[c + d*x]^2)^(1/4)]*(-Tan[c + d*x]^2)^(3/4)))/(3*d*Tan[c + d*x]^(3/2))`

3.421.3 Rubi [A] (verified)Time = 0.41 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {2011, 3042, 3955, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$\downarrow \text{2011}$$

$$B \int \frac{1}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$B \int \frac{1}{\tan(c + dx)^{5/2}} dx$$

$$\downarrow \text{3955}$$

$$B \left(- \int \frac{1}{\sqrt{\tan(c + dx)}} dx - \frac{2}{3d \tan^{\frac{3}{2}}(c + dx)} \right)$$

$$\begin{aligned}
& \downarrow 3042 \\
& B\left(-\int \frac{1}{\sqrt{\tan(c+dx)}} dx - \frac{2}{3d \tan^{\frac{3}{2}}(c+dx)}\right) \\
& \downarrow 3957 \\
& B\left(-\frac{\int \frac{1}{\sqrt{\tan(c+dx)(\tan^2(c+dx)+1)}} d \tan(c+dx)}{d} - \frac{2}{3d \tan^{\frac{3}{2}}(c+dx)}\right) \\
& \downarrow 266 \\
& B\left(-\frac{2 \int \frac{1}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)}}{d} - \frac{2}{3d \tan^{\frac{3}{2}}(c+dx)}\right) \\
& \downarrow 755 \\
& B\left(-\frac{2\left(\frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)}\right)}{d} - \frac{2}{3d \tan^{\frac{3}{2}}(c+dx)}\right) \\
& \downarrow 1476 \\
& B\left(-\frac{2\left(\frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)} + \frac{1}{2} \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d \sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d \sqrt{\tan(c+dx)}\right)\right)}{d} - \frac{2}{3d \tan^{\frac{3}{2}}(c+dx)}\right) \\
& \downarrow 1082 \\
& B\left(-\frac{2\left(\frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)} + \frac{1}{2} \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}}\right)\right)}{d} - \frac{2}{3d \tan^{\frac{3}{2}}(c+dx)}\right) \\
& \downarrow 217 \\
& B\left(-\frac{2\left(\frac{1}{2} \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d \sqrt{\tan(c+dx)} + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}}\right)\right)}{d} - \frac{2}{3d \tan^{\frac{3}{2}}(c+dx)}\right) \\
& \downarrow 1479
\end{aligned}$$

3.421. $\int \frac{aB+bB \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$B \left(\frac{2 \left(\frac{1}{2} \left(- \frac{\int - \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int - \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right)}{d} \right)$$

↓ 25

$$B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right)}{d} \right)$$

↓ 27

$$B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right)}{d} \right)$$

↓ 1103

$$B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d} \right)$$

input `Int[(a*B + b*B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])),x]`

output `B*((-2*((-ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]/Sqrt[2]))/2 + (-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2]))/2)/d - 2/(3*d*Tan[c + d*x]^(3/2))`

3.421.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3955 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x])^(n + 1)/(b*d*(n + 1)), x] - Simp[1/b^2 Int[(b*Tan[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

3.421.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.65

method	result
derivativedivides	$B \left(\frac{-\frac{2}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})}{4}}{d} \right.$
default	$B \left(\frac{-\frac{2}{3 \tan(dx+c)^{\frac{3}{2}}} - \frac{\sqrt{2} \left(\ln \left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))} \right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}) \right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})}{4}}{d} \right.$

input `int((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

$$3.421. \int \frac{aB+bB \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$$

output $1/d*B*(-2/3/\tan(d*x+c)^{(3/2)}-1/4*2^{(1/2)}*(\ln((1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(1-2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))+2*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}))$

3.421.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.33

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx =$$

$$3d\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log\left(B\sqrt{\tan(dx + c)} + d\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}}\right) \tan(dx + c)^2 + 3id\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log\left(B\sqrt{\tan(dx + c)} + d\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}}\right)$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x, algorithm="fracas")`

output $-1/6*(3*d*(-B^4/d^4)^{(1/4)}*\log(B*\sqrt{\tan(d*x + c)} + d*(-B^4/d^4)^{(1/4)})*\tan(d*x + c)^2 + 3*I*d*(-B^4/d^4)^{(1/4)}*\log(B*\sqrt{\tan(d*x + c)} + d*(-B^4/d^4)^{(1/4)})*\tan(d*x + c)^2 - 3*I*d*(-B^4/d^4)^{(1/4)}*\log(B*\sqrt{\tan(d*x + c)} - d*(-B^4/d^4)^{(1/4)})*\tan(d*x + c)^2 - 3*d*(-B^4/d^4)^{(1/4)}*\log(B*\sqrt{\tan(d*x + c)} - d*(-B^4/d^4)^{(1/4)})*\tan(d*x + c)^2 + 4*B*\sqrt{\tan(d*x + c)}))/(d*\tan(d*x + c)^2)$

3.421.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = B \int \frac{1}{\tan^{\frac{5}{2}}(c + dx)} dx$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)**(5/2)/(a+b*tan(d*x+c)),x)`

output `B*Integral(tan(c + d*x)**(-5/2), x)`

3.421. $\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx$

3.421.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.79

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx =$$

$$\frac{6\sqrt{2}B \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + 2\sqrt{\tan(dx + c)}\right)\right) + 6\sqrt{2}B \arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2} - 2\sqrt{\tan(dx + c)}\right)\right) + 3\sqrt{2}B \log\left(\frac{\sqrt{2}\sqrt{\tan(dx + c)} + \tan(dx + c) + 1}{-\sqrt{2}\sqrt{\tan(dx + c)} + \tan(dx + c) + 1}\right) + 8B/\tan(dx + c)^{\frac{3}{2}}}{d}$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `-1/12*(6*sqrt(2)*B*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 6*sqrt(2)*B*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) + 3*sqrt(2)*B*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - 3*sqrt(2)*B*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + 8*B/tan(d*x + c)^(3/2)) /d`

3.421.8 Giac [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = \text{Timed out}$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.421.9 Mupad [B] (verification not implemented)

Time = 13.57 (sec) , antiderivative size = 16545, normalized size of antiderivative = 106.06

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = \text{Too large to display}$$

input `int((B*a + B*b*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + b*tan(c + d*x))),x)`

output `atan(((tan(c + d*x)^(1/2)*(64*B^4*a^9*b^9*d^5 + 32*B^4*a^13*b^5*d^5) - ((64*B^4*a^6*b^2*d^4 - B^4*a^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a^3*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*((tan(c + d*x)^(1/2)*(512*B^2*a^8*b^10*d^7 + 448*B^2*a^12*b^6*d^7 - 128*B^2*a^14*b^4*d^7 - 64*B^2*a^16*b^2*d^7) - (((64*B^4*a^6*b^2*d^4 - B^4*a^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a^3*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*(tan(c + d*x)^(1/2)*(-((64*B^4*a^6*b^2*d^4 - B^4*a^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a^3*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*(512*a^9*b^9*d^9 + 512*a^11*b^7*d^9 - 512*a^13*b^5*d^9 - 512*a^15*b^3*d^9) - 512*B*a^8*b^10*d^8 - 512*B*a^10*b^8*d^8 + 384*B*a^12*b^6*d^8 + 256*B*a^14*b^4*d^8 - 128*B*a^16*b^2*d^8))*(-((64*B^4*a^6*b^2*d^4 - B^4*a^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a^3*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2) - 384*B^3*a^9*b^9*d^6 + 32*B^3*a^13*b^5*d^6 + 32*B^3*a^15*b^3*d^6))*(-((64*B^4*a^6*b^2*d^4 - B^4*a^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a^3*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*i + (tan(c + d*x)^(1/2)*(64*B^4*a^9*b^9*d^5 + 32*B^4*a^13*b^5*d^5) - (((64*B^4*a^6*b^2*d^4 - B^4*a^4*(16*a^4*d^4 + 16*b^4*d^4 + 32*a^2*b^2*d^4))^(1/2) - 8*B^2*a^3*b*d^2)/(16*(a^4*d^4 + b^4*d^4 + 2*a^2*b^2*d^4)))^(1/2)*((tan(c + d*x)^(1/2)*(512*B^2*a^8*b^10*d^7 + 448*B^2...`

3.421. $\int \frac{aB+bB \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$

3.422
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.422.1 Optimal result 4074
 3.422.2 Mathematica [C] (verified) 4075
 3.422.3 Rubi [A] (verified) 4075
 3.422.4 Maple [A] (verified) 4081
 3.422.5 Fricas [B] (verification not implemented) 4082
 3.422.6 Sympy [F] 4083
 3.422.7 Maxima [A] (verification not implemented) 4084
 3.422.8 Giac [F(-1)] 4084
 3.422.9 Mupad [B] (verification not implemented) 4085

3.422.1 Optimal result

Integrand size = 36, antiderivative size = 256

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{(a+b)B \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d} - \frac{(a+b)B \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d}$$

$$- \frac{2a^{5/2}B \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{b^{3/2}(a^2+b^2)d} - \frac{(a-b)B \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d}$$

$$+ \frac{(a-b)B \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d} + \frac{2B\sqrt{\tan(c+dx)}}{bd}$$

output

```
-2*a^(5/2)*B*arctan(b^(1/2)*tan(d*x+c)^(1/2)/a^(1/2))/b^(3/2)/(a^2+b^2)/d-
1/2*(a+b)*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/2*(a
+b)*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/4*(a-b)*B*1
n(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)+1/4*(a-b)*B*1
n(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)+2*B*tan(d*x+c
)^(1/2)/b/d
```

3.422.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.17 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.61

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{B\left(\sqrt[4]{-1}b^{3/2}(-ia+b)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right) - 2a^{5/2}\arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right) + \sqrt[4]{-1}b^{3/2}(ia+b)a\right)}{b^{3/2}(a^2+b^2)d}$$

input `Integrate[(Tan[c + d*x]^(5/2)*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `(B*((-1)^(1/4)*b^(3/2)*((-I)*a + b)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] - 2*a^(5/2)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]] + (-1)^(1/4)*b^(3/2)*(I*a + b)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + 2*a^2*Sqrt[b]*Sqrt[Tan[c + d*x]] + 2*b^(5/2)*Sqrt[Tan[c + d*x]])/(b^(3/2)*(a^2 + b^2)*d)`

3.422.3 Rubi [A] (verified)

Time = 1.02 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.89, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {2011, 3042, 4049, 27, 3042, 4136, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$\downarrow \text{2011}$$

$$B \int \frac{\tan^{\frac{5}{2}}(c+dx)}{a+b\tan(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$B \int \frac{\tan(c+dx)^{5/2}}{a+b\tan(c+dx)} dx$$

$$\downarrow \text{4049}$$

3.422. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

$$\begin{aligned}
& B \left(\frac{2 \int \frac{-a \tan^2(c+dx) + b \tan(c+dx) + a}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b} + \frac{2\sqrt{\tan(c+dx)}}{bd} \right) \\
& \quad \downarrow 27 \\
& B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{\int \frac{a \tan^2(c+dx) + b \tan(c+dx) + a}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b} \right) \\
& \quad \downarrow 3042 \\
& B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{\int \frac{a \tan(c+dx)^2 + b \tan(c+dx) + a}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b} \right) \\
& \quad \downarrow 4136 \\
& B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{\int \frac{b^2 + a \tan(c+dx)b}{\sqrt{\tan(c+dx)}(a^2+b^2)} dx + \frac{a^3 \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b}}{b} \right) \\
& \quad \downarrow 3042 \\
& B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{\int \frac{b^2 + a \tan(c+dx)b}{\sqrt{\tan(c+dx)}(a^2+b^2)} dx + \frac{a^3 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b}}{b} \right) \\
& \quad \downarrow 4017 \\
& B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{2 \int \frac{b(b+a \tan(c+dx))}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} + \frac{a^3 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b}}{b} \right) \\
& \quad \downarrow 27 \\
& B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{2b \int \frac{b+a \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} + \frac{a^3 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b}}{b} \right) \\
& \quad \downarrow 1482 \\
& B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{2b \left(\frac{1}{2}(a+b) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a-b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d(a^2+b^2)} + \frac{a^3 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{b}}{b} \right)
\end{aligned}$$

3.422. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

↓ 1476

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{2b \left(\frac{1}{2}(a+b) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx) - \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx) + \sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2}(a-b) \int \frac{1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} \right)}{b}$$

↓ 1082

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{2b \left(\frac{1}{2}(a+b) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} \right)}{b}$$

↓ 217

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{2b \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right) + \frac{a^3}{3}}{d(a^2+b^2)}$$

↓ 1479

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{2b \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d(a^2+b^2)}$$

↓ 25

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{2b \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) \right)}{d(a^2+b^2)}$$

↓ 27

3.422. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{2b \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right)}{b}$$

↓ 1103

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{a^3 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} + \frac{2b \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right)}{b}$$

↓ 4117

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{a^3 \int \frac{1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} d \tan(c+dx)}{d(a^2+b^2)} + \frac{2b \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right)}{b}$$

↓ 73

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{2a^3 \int \frac{1}{a+b \tan(c+dx)} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} + \frac{2b \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right)}{b}$$

↓ 218

$$B \left(\frac{2\sqrt{\tan(c+dx)}}{bd} - \frac{2b \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right)}{b}$$

```
input Int[(Tan[c + d*x]^(5/2)*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x
]
```

```
output B*(-(((2*a^(5/2)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[b]*(a
^2 + b^2)*d) + (2*b*(((a + b)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sq
rt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2])))/2 - ((a - b)*(-1
/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sq
rt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2))/((a^2 + b^2)*d)
)/b) + (2*Sqrt[Tan[c + d*x]])/(b*d))
```

3.422.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 217 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 1082 Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

```
rule 1103 Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

$$3.422. \int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

```
rule 4049 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m - 2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d*f*(m + n - 1))), x] + Simp[1/(d*(m + n - 1))
Int[(a + b*Tan[e + f*x])^(m - 3)*(c + d*Tan[e + f*x])^n*Simp[a^3*d*(m + n
- 1) - b^2*(b*c*(m - 2) + a*d*(1 + n)) + b*d*(m + n - 1)*(3*a^2 - b^2)*Tan[
e + f*x] - b^2*(b*c*(m - 2) - a*d*(3*m + 2*n - 4))*Tan[e + f*x]^2, x], x],
x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2
, 0] && NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && GtQ[m, 2] && (GeQ[n, -1] || I
ntegerQ[m]) && !(IGtQ[n, 2] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0]))
)
```

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

```
rule 4136 Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &&
!GtQ[n, 0] && !LeQ[n, -1]
```

3.422.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 242, normalized size of antiderivative = 0.95

method	result
derivativedivides	$B \left(\frac{2(\sqrt{\tan(dx+c)})}{b} - \frac{2a^3 \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{b(a^2+b^2)\sqrt{ab}} \right) + \frac{b\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right)}{4}$
default	$B \left(\frac{2(\sqrt{\tan(dx+c)})}{b} - \frac{2a^3 \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{b(a^2+b^2)\sqrt{ab}} \right) + \frac{b\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))) \right)}{4}$

$$3.422. \int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

input `int(tan(d*x+c)^(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/d*B*(2*tan(d*x+c)^(1/2)/b-2/b*a^3/(a^2+b^2)/(a*b)^(1/2)*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))+2/(a^2+b^2)*(-1/8*b*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))-1/8*a*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))`

3.422.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1606 vs. $2(216) = 432$.

Time = 0.34 (sec) , antiderivative size = 3238, normalized size of antiderivative = 12.65

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fricas")`

output `[1/2*(2*B*a^2*sqrt(-a/b)*log(-(2*b*sqrt(-a/b)*sqrt(tan(d*x + c)) - b*tan(d*x + c) + a)/(b*tan(d*x + c) + a)) - (a^2*b + b^3)*d*sqrt(-(2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2))*log(((a^5 + 2*a^3*b^2 + a*b^4)*d^3*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)) - (B^2*a^2*b - B^2*b^3)*d)*sqrt(-(2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2)) - (B^3*a^2 - B^3*b^2)*sqrt(tan(d*x + c)))] + (a^2*b + b^3)*d*sqrt(-(2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2))*log(-((a^5 + 2*a^3*b^2 + a*b^4)*d^3*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)) - (B^2*a^2*b - B^2*b^3)*d)*sqrt(-(2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2)) - (B^3*a^2 - B^3*b^2)*sqrt(tan(d*x + c)))] + (a^2*b + b^3)*d*sqrt(-(2*B^2*a*b - (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2))*log(((a^5 + 2*a^3*b^2 + a*b^4)*...`

3.422.6 Sympy [F]

$$\int \frac{\tan^{\frac{5}{2}}(c + dx)(aB + bB \tan(c + dx))}{(a + b \tan(c + dx))^2} dx = B \int \frac{\tan^{\frac{5}{2}}(c + dx)}{a + b \tan(c + dx)} dx$$

input `integrate(tan(d*x+c)**(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `B*Integral(tan(c + d*x)**(5/2)/(a + b*tan(c + d*x)), x)`

3.422.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 188, normalized size of antiderivative = 0.73

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx =$$

$$\frac{8Ba^3 \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^2b+b^3)\sqrt{ab}} + \frac{\left(2\sqrt{2}(a+b)\arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}+2\sqrt{\tan(dx+c)})\right)\right) + 2\sqrt{2}(a+b)\arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}-2\sqrt{\tan(dx+c)})\right) - \sqrt{2}(a^2+b^2)}{4d}$$

input `integrate(tan(d*x+c)^(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorith="maxima")`

output `-1/4*(8*B*a^3*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/((a^2*b + b^3)*sqrt(a*b)) + (2*sqrt(2)*(a + b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))) + 2*sqrt(2)*(a + b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))) - sqrt(2)*(a - b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) + sqrt(2)*(a - b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*B/(a^2 + b^2) - 8*B*sqrt(tan(d*x + c))/b)/d`

3.422.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorith="giac")`

output `Timed out`

3.422.9 Mupad [B] (verification not implemented)

Time = 42.44 (sec) , antiderivative size = 18514, normalized size of antiderivative = 72.32

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Too large to display}$$

```
input int((tan(c + d*x)^(5/2)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x
)
```

```
output (log((((((((((128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b^2)^2*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16*B^2*a^3*b^3*d^2 - 16*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2) + (768*B*a^3*b^3*(a^2 + b^2))/d)*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16*B^2*a^3*b^3*d^2 - 16*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (64*B^2*a^3*tan(c + d*x)^(1/2)*(2*a^8 + 15*b^8 - 17*a^2*b^6 + 51*a^4*b^4 + 21*a^6*b^2))/(d^2*(a^2 + b^2)^2))*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16*B^2*a^3*b^3*d^2 - 16*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (32*B^3*a^4*(4*a^8 + b^8 - 77*a^2*b^6 + 47*a^4*b^4 + 33*a^6*b^2))/(d^3*(a^2 + b^2)^3))*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16*B^2*a^3*b^3*d^2 - 16*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (16*B^4*a^4*tan(c + d*x)^(1/2)*(a^10 - 2*b^10 - 4*a^2*b^8 - 27*a^4*b^6 + 15*a^6*b^4 + 9*a^8*b^2))/(b*d^4*(a^2 + b^2)^4))*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16*B^2*a^3*b^3*d^2 - 16*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (8*B^5*a^7*(a^6 + 10*b^6 + 27*a^2*b^4 + 10*a^4*b^2))/(b*d^5*(a^2 + b^2)^4))*(((192*B^4*a^6*b^6*d^4 - 16*B^4*a^4*b^8*d^4 - 16*B^4*a^12*d^4 - 608*B^4*a^8*b^4*d^4 + 192*B^4*a^10*b^2*d^4)^(1/2) + 16*B^2*a^3*b^3*d^2 - 16*B^2*a^5*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6*d^4 + 6*a^4*b^4*d^4 + 4*a^6*b^2*d^4))^(1/2))/4 + (log((((((((((128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b^2)^2*(-(4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a^3*b^3*...
```

3.423
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.423.1 Optimal result 4086
 3.423.2 Mathematica [C] (verified) 4087
 3.423.3 Rubi [A] (verified) 4087
 3.423.4 Maple [A] (verified) 4093
 3.423.5 Fracas [B] (verification not implemented) 4093
 3.423.6 Sympy [F] 4094
 3.423.7 Maxima [A] (verification not implemented) 4095
 3.423.8 Giac [F(-1)] 4095
 3.423.9 Mupad [B] (verification not implemented) 4096

3.423.1 Optimal result

Integrand size = 36, antiderivative size = 237

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{(a-b)B \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d} - \frac{(a-b)B \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d}$$

$$+ \frac{2a^{3/2}B \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{b}(a^2+b^2)d} + \frac{(a+b)B \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d}$$

$$- \frac{(a+b)B \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d}$$

output

```
-1/2*(a-b)*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/2*(a-b)*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)+1/4*(a+b)*B*ln(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)-1/4*(a+b)*B*ln(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)+2*a^(3/2)*B*arctan(b^(1/2)*tan(d*x+c)^(1/2)/a^(1/2))/(a^2+b^2)/d/b^(1/2)
```

3.423.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.18 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.96

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{B\left(3a\left(2\sqrt{2}\sqrt{b}\arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)\right)-2\sqrt{2}\sqrt{b}\arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)\right)+8\sqrt{a}\arctan\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{1+\sqrt{2}\sqrt{\tan(c+dx)}}\right)}{12\sqrt{b}(a^2+b^2)d}$$

input `Integrate[(Tan[c + d*x]^(3/2)*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `(B*(3*a*(2*Sqrt[2]*Sqrt[b]*ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - 2*Sqrt[2]*Sqrt[b]*ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]] + 8*Sqrt[a]*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]] + Sqrt[2]*Sqrt[b]*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Sqrt[2]*Sqrt[b]*Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]) + 8*b^(3/2)*Hypergeometric2F1[3/4, 1, 7/4, -Tan[c + d*x]^2*Tan[c + d*x]^(3/2)))/(12*Sqrt[b]*(a^2 + b^2)*d)`

3.423.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.86, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.472$, Rules used = {2011, 3042, 4056, 25, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$\downarrow \text{2011}$$

$$B \int \frac{\tan^{\frac{3}{2}}(c+dx)}{a+b\tan(c+dx)} dx$$

$$\downarrow \text{3042}$$

3.423. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

$$\begin{aligned}
& B \int \frac{\tan(c+dx)^{3/2}}{a+b \tan(c+dx)} dx \\
& \quad \downarrow 4056 \\
& B \left(\frac{a^2 \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} + \frac{\int -\frac{a-b \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} \right) \\
& \quad \downarrow 25 \\
& B \left(\frac{a^2 \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} - \frac{\int \frac{a-b \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} \right) \\
& \quad \downarrow 3042 \\
& B \left(\frac{a^2 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} - \frac{\int \frac{a-b \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} \right) \\
& \quad \downarrow 4017 \\
& B \left(\frac{a^2 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} - \frac{2 \int \frac{a-b \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} \right) \\
& \quad \downarrow 1482 \\
& B \left(\frac{a^2 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} - \frac{2 \left(\frac{1}{2}(a+b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a-b) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d(a^2+b^2)} \right) \\
& \quad \downarrow 1476 \\
& B \left(\frac{a^2 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} - \frac{2 \left(\frac{1}{2}(a+b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a-b) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}} dx \right) \right)}{d(a^2+b^2)} \right) \\
& \quad \downarrow 1082 \\
& B \left(\frac{a^2 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} - \frac{2 \left(\frac{1}{2}(a+b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a-b) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)} \right) \\
& \quad \downarrow 217
\end{aligned}$$

3.423. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$B \left(\frac{a^2 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2 + b^2} - \frac{2 \left(\frac{1}{2}(a+b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a-b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2 + b^2)} \right)$$

↓ 1479

$$B \left(\frac{a^2 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2 + b^2} - \frac{2 \left(\frac{1}{2}(a+b) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}}}{2\sqrt{2}} \right) \right)}{d(a^2 + b^2)} \right)$$

↓ 25

$$B \left(\frac{a^2 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2 + b^2} - \frac{2 \left(\frac{1}{2}(a+b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}}}{2\sqrt{2}} \right) \right)}{d(a^2 + b^2)} \right)$$

↓ 27

$$B \left(\frac{a^2 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2 + b^2} - \frac{2 \left(\frac{1}{2}(a+b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)}+1} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)}+1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)}} \right) \right)}{d(a^2 + b^2)} \right)$$

↓ 1103

$$B \left(\frac{a^2 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2 + b^2} - \frac{2 \left(\frac{1}{2}(a-b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) + \frac{1}{2}(a+b)}{d(a^2 + b^2)} \right)$$

↓ 4117

3.423. $\int \frac{\tan^3(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$B \left(\frac{a^2 \int \frac{1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} d \tan(c+dx)}{d(a^2+b^2)} - \frac{2 \left(\frac{1}{2}(a-b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)} \right)$$

↓ 73

$$B \left(\frac{2a^2 \int \frac{1}{a+b \tan(c+dx)} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} - \frac{2 \left(\frac{1}{2}(a-b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) + \frac{1}{2}(a+b)}{d(a^2+b^2)} \right)$$

↓ 218

$$B \left(\frac{2a^{3/2} \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{bd}(a^2+b^2)} - \frac{2 \left(\frac{1}{2}(a-b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) + \frac{1}{2}(a+b)}{d(a^2+b^2)} \right)$$

input `Int[(Tan[c + d*x]^(3/2)*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `B*((2*a^(3/2)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[b]*(a^2 + b^2)*d) - (2*(((a - b)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]))/2 + ((a + b)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d))`

3.423.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

3.423. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
 -1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
 & (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
 t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
 implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/R
 t[a/b, 2])], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
 eQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
 imp[d*(Log[RemoveContent[a + b*x + c*x^2, x])/b], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
 2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
 e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
 && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
 -2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
 x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
 reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4056 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(3/2)/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[1/(c^2 + d^2) Int[Simp[a^2*c - b^2*c + 2*a*b*d + (2*a*b*c - a^2*d + b^2*d)*Tan[e + f*x], x]/Sqrt[a + b*Tan[e + f*x]], x], x] + Simp[(b*c - a*d)^2/(c^2 + d^2) Int[(1 + Tan[e + f*x]^2)/(Sqrt[a + b*Tan[e + f*x]]*(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

3.423.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 226, normalized size of antiderivative = 0.95

method	result
derivativedivides	$B \left(\frac{2a^2 \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{(a^2+b^2)\sqrt{ab}} \right) + \frac{a\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})\right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})\right)}{4}$
default	$B \left(\frac{2a^2 \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{(a^2+b^2)\sqrt{ab}} \right) + \frac{a\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})\right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})\right)}{4}$

input `int(tan(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output
$$\frac{1}{d} B \left(\frac{2a^2}{(a^2+b^2)} \frac{1}{(ab)^{1/2}} \arctan\left(\frac{b \tan(dx+c)^{1/2}}{(ab)^{1/2}}\right) + \frac{2}{(a^2+b^2)} \left(-\frac{1}{8} a^{2^{1/2}} \left(\ln\left(\frac{(1+2^{1/2}) \tan(dx+c)^{1/2} + \tan(dx+c)}{(1-2^{1/2}) \tan(dx+c)^{1/2} + \tan(dx+c)}\right) + 2 \arctan(1+2^{1/2} \tan(dx+c)^{1/2}) + 2 \arctan(-1+2^{1/2} \tan(dx+c)^{1/2}) \right) + \frac{1}{8} b^{2^{1/2}} \left(\ln\left(\frac{(1-2^{1/2}) \tan(dx+c)^{1/2} + \tan(dx+c)}{(1+2^{1/2}) \tan(dx+c)^{1/2} + \tan(dx+c)}\right) + 2 \arctan(1+2^{1/2} \tan(dx+c)^{1/2}) + 2 \arctan(-1+2^{1/2} \tan(dx+c)^{1/2}) \right) \right) \right)$$

3.423.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1565 vs. 2(199) = 398.

Time = 0.32 (sec) , antiderivative size = 3156, normalized size of antiderivative = 13.32

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorith="fricas")`

output `[1/2*((a^2 + b^2)*d*sqrt((2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2))*log(((a^4*b + 2*a^2*b^3 + b^5)*d^3*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)) + (B^2*a^3 - B^2*a*b^2)*d)*sqrt((2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2)) - (B^3*a^2 - B^3*b^2)*sqrt(tan(d*x + c))) - (a^2 + b^2)*d*sqrt((2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2))*log(-((a^4*b + 2*a^2*b^3 + b^5)*d^3*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)) + (B^2*a^3 - B^2*a*b^2)*d)*sqrt((2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2)) - (B^3*a^2 - B^3*b^2)*sqrt(tan(d*x + c))) - (a^2 + b^2)*d*sqrt((2*B^2*a*b - (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2))*log(((a^4*b + 2*a^2*b^3 + b^5)*d^3*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)) - (B^2*a^3 - B^...`

3.423.6 Sympy [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = B \int \frac{\tan^{\frac{3}{2}}(c+dx)}{a+b \tan(c+dx)} dx$$

input `integrate(tan(d*x+c)**(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `B*Integral(tan(c + d*x)**(3/2)/(a + b*tan(c + d*x)), x)`

3.423.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.73

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{8Ba^2 \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^2+b^2)\sqrt{ab}} - \frac{\left(2\sqrt{2}(a-b) \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}+2\sqrt{\tan(dx+c)})\right)\right) + 2\sqrt{2}(a-b) \arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}-2\sqrt{\tan(dx+c)})\right) + \sqrt{2}(a+b)}{a^2+b^2}}{4d}$$

```
input integrate(tan(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algo
rithm="maxima")
```

```
output 1/4*(8*B*a^2*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/((a^2 + b^2)*sqrt(a*b)
) - (2*sqrt(2)*(a - b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))
) + 2*sqrt(2)*(a - b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))
) + sqrt(2)*(a + b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - s
qrt(2)*(a + b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*B/(a^2
+ b^2))/d
```

3.423.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Timed out}$$

```
input integrate(tan(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algo
rithm="giac")
```

```
output Timed out
```

3.423.9 Mupad [B] (verification not implemented)

Time = 39.45 (sec) , antiderivative size = 16878, normalized size of antiderivative = 71.22

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Too large to display}$$

```
input int((tan(c + d*x)^(3/2)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x
)
```

```
output (log((((((((((128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b^2)^2*((4*(-B^
4*b^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a^3*b^3*d^2 + 16*B^2*a
*b^5*d^2)/(d^4*(a^2 + b^2)^4))^(1/2) + (768*B*a^2*b^4*(a^2 + b^2))/d)*((4*
(-B^4*b^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a^3*b^3*d^2 + 16*B
^2*a*b^5*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (64*B^2*a*b^2*tan(c + d*x)^(
1/2)*(2*a^8 + 15*b^8 - 17*a^2*b^6 + 51*a^4*b^4 + 21*a^6*b^2))/(d^2*(a^2 +
b^2)^2))*((4*(-B^4*b^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a^3*b
^3*d^2 + 16*B^2*a*b^5*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (32*B^3*a*b^3*(
4*a^8 + b^8 - 77*a^2*b^6 + 47*a^4*b^4 + 33*a^6*b^2))/(d^3*(a^2 + b^2)^3))*
((4*(-B^4*b^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a^3*b^3*d^2 +
16*B^2*a*b^5*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (16*B^4*b^3*tan(c + d*x)
^(1/2)*(a^10 - 2*b^10 - 4*a^2*b^8 - 27*a^4*b^6 + 15*a^6*b^4 + 9*a^8*b^2))/
(d^4*(a^2 + b^2)^4))*((4*(-B^4*b^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) -
16*B^2*a^3*b^3*d^2 + 16*B^2*a*b^5*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (8*
B^5*a^2*b^4*(a^6 + 10*b^6 + 27*a^2*b^4 + 10*a^4*b^2))/(d^5*(a^2 + b^2)^4))
*(((192*B^4*a^2*b^10*d^4 - 16*B^4*b^12*d^4 - 608*B^4*a^4*b^8*d^4 + 192*B^4
*a^6*b^6*d^4 - 16*B^4*a^8*b^4*d^4)^(1/2) - 16*B^2*a^3*b^3*d^2 + 16*B^2*a*b
^5*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6*d^4 + 6*a^4*b^4*d^4 + 4*a^6*b^2*d^4
))^(1/2))/4 + (log((((((((((128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b
^2)^2*(-(4*(-B^4*b^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16*B^2*a^3*...
```

3.424
$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.424.1 Optimal result 4097
 3.424.2 Mathematica [C] (verified) 4098
 3.424.3 Rubi [A] (verified) 4098
 3.424.4 Maple [A] (verified) 4103
 3.424.5 Fricas [B] (verification not implemented) 4104
 3.424.6 Sympy [F] 4105
 3.424.7 Maxima [A] (verification not implemented) 4106
 3.424.8 Giac [F(-1)] 4106
 3.424.9 Mupad [B] (verification not implemented) 4107

3.424.1 Optimal result

Integrand size = 36, antiderivative size = 237

$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= -\frac{(a+b)B \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d} + \frac{(a+b)B \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d}$$

$$- \frac{2\sqrt{a}\sqrt{b}B \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{(a^2+b^2)d} + \frac{(a-b)B \log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d}$$

$$- \frac{(a-b)B \log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}+\tan(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d}$$

```
output 1/2*(a+b)*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)+1/2*(a
+b)*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)+1/4*(a-b)*B*1
n(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)-1/4*(a-b)*B*1
n(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)-2*B*arctan(b^(
1/2)*tan(d*x+c)^(1/2)/a^(1/2))*a^(1/2)*b^(1/2)/(a^2+b^2)/d
```


3.424.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.10 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.86

$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{B\left(-6\sqrt{2}b \arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right) + 6\sqrt{2}b \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right) - 24\sqrt{a}\sqrt{b} \arctan\left(\frac{\sqrt{b}}{\sqrt{a}}\sqrt{\tan(c+dx)}\right)\right)}{12(a^2+b^2)d}$$

input `Integrate[(Sqrt[Tan[c + d*x]]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `(B*(-6*Sqrt[2]*b*ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] + 6*Sqrt[2]*b*ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]] - 24*Sqrt[a]*Sqrt[b]*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]] - 3*Sqrt[2]*b*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] + 3*Sqrt[2]*b*Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] + 8*a*Hypergeometric2F1[3/4, 1, 7/4, -Tan[c + d*x]^2]*Tan[c + d*x]^(3/2)))/(12*(a^2 + b^2)*d)`

3.424.3 Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.86, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {2011, 3042, 4055, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$\downarrow \text{2011}$$

$$B \int \frac{\sqrt{\tan(c+dx)}}{a+b\tan(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
& B \int \frac{\sqrt{\tan(c+dx)}}{a+b \tan(c+dx)} dx \\
& \quad \downarrow \text{4055} \\
& B \left(\frac{\int \frac{b+a \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{ab \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} \right) \\
& \quad \downarrow \text{3042} \\
& B \left(\frac{\int \frac{b+a \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} - \frac{ab \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} \right) \\
& \quad \downarrow \text{4017} \\
& B \left(\frac{2 \int \frac{b+a \tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} - \frac{ab \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} \right) \\
& \quad \downarrow \text{1482} \\
& B \left(\frac{2 \left(\frac{1}{2}(a+b) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a-b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d(a^2+b^2)} - \frac{ab \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} \right) \\
& \quad \downarrow \text{1476} \\
& B \left(\frac{2 \left(\frac{1}{2}(a+b) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2} \right)}{d(a^2+b^2)} \right) \\
& \quad \downarrow \text{1082} \\
& B \left(\frac{2 \left(\frac{1}{2}(a+b) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d(a^2+b^2)} \right) \\
& \quad \downarrow \text{217} \\
& B \left(\frac{2 \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d(a^2+b^2)} \right) -
\end{aligned}$$

3.424. $\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

↓ 1479

$$B \left(\frac{2 \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right.$$

↓ 25

$$B \left(\frac{2 \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right.$$

↓ 27

$$B \left(\frac{2 \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right.$$

↓ 1103

$$B \left(\frac{2 \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right.$$

↓ 4117

$$B \left(\frac{2 \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right.$$

↓ 73

$$B \left(\frac{2 \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d(a^2+b^2)} \right)$$

↓ 218

$$B \left(\frac{2 \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d(a^2+b^2)} \right)$$

input `Int[(Sqrt[Tan[c + d*x]]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `B*((-2*Sqrt[a]*Sqrt[b]*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/((a^2 + b^2)*d) + (2*((a + b)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]))/2 - ((a - b)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)`

3.424.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

- rule 217 $\text{Int}[(a_ + (b_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{-1} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /;$ $\text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&$
 $\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 218 $\text{Int}[(a_ + (b_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a) \cdot \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /;$ $\text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$
- rule 1082 $\text{Int}[(a_ + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4 \cdot \text{Simplify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /;$ $\text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4 \cdot a \cdot c]) /;$ $\text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_ + (e_ \cdot)(x_)) / ((a_ + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /;$ $\text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$
- rule 1476 $\text{Int}[(d_ + (e_ \cdot)(x_)^2) / ((a_ + (c_ \cdot)(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c) \ \text{Int}[1/\text{Simp}[d/e + q \cdot x + x^2, x], x], x] + \text{Simp}[e/(2 \cdot c) \ \text{Int}[1/\text{Simp}[d/e - q \cdot x + x^2, x], x], x]] /;$ $\text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{PosQ}[d \cdot e]$
- rule 1479 $\text{Int}[(d_ + (e_ \cdot)(x_)^2) / ((a_ + (c_ \cdot)(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c \cdot q) \ \text{Int}[(q - 2 \cdot x)/\text{Simp}[d/e + q \cdot x - x^2, x], x], x] + \text{Simp}[e/(2 \cdot c \cdot q) \ \text{Int}[(q + 2 \cdot x)/\text{Simp}[d/e - q \cdot x - x^2, x], x], x]] /;$ $\text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{NegQ}[d \cdot e]$
- rule 1482 $\text{Int}[(d_ + (e_ \cdot)(x_)^2) / ((a_ + (c_ \cdot)(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a \cdot c, 2]\}, \text{Simp}[(d \cdot q + a \cdot e)/(2 \cdot a \cdot c) \ \text{Int}[(q + c \cdot x^2)/(a + c \cdot x^4), x], x] + \text{Simp}[(d \cdot q - a \cdot e)/(2 \cdot a \cdot c) \ \text{Int}[(q - c \cdot x^2)/(a + c \cdot x^4), x], x]] /;$ $\text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{NeQ}[c \cdot d^2 + a \cdot e^2, 0] \ \&\& \ \text{NeQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{NegQ}[(-a) \cdot c]$

```
rule 2011 Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :=
  Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x],
  a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4017 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_
)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sq
rt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] &
& NeQ[c^2 + d^2, 0]
```

```
rule 4055 Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[1/(c^2 + d^2) Int[Simp[a*c + b*d + (b*c -
a*d)*Tan[e + f*x], x]/Sqrt[a + b*Tan[e + f*x]], x], x] - Simp[d*((b*c - a*
d)/(c^2 + d^2)) Int[(1 + Tan[e + f*x]^2)/(Sqrt[a + b*Tan[e + f*x]]*(c + d
*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && N
eQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

```
rule 4117 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_)
+ (f_)*(x_)]^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] :=
  Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
  FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

3.424.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 225, normalized size of antiderivative = 0.95

$$3.424. \quad \int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

method	result
derivativedivides	$B \left(-\frac{2ab \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{(a^2+b^2)\sqrt{ab}} + \frac{b\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})\right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})\right)}{4} \right)$
default	$B \left(-\frac{2ab \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{(a^2+b^2)\sqrt{ab}} + \frac{b\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})\right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})\right)}{4} \right)$

```
input int(tan(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNVERBOSE)
```

```
output 1/d*B*(-2*a*b/(a^2+b^2)/(a*b)^(1/2)*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))+2/(a^2+b^2)*(1/8*b*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))+1/8*a*2^(1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))
```

3.424.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1567 vs. 2(199) = 398.

Time = 0.32 (sec) , antiderivative size = 3160, normalized size of antiderivative = 13.33

$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algo rithm="fracas")
```

output `[1/2*((a^2 + b^2)*d*sqrt(-(2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2))*log(((a^5 + 2*a^3*b^2 + a*b^4)*d^3*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)) - (B^2*a^2*b - B^2*b^3)*d)*sqrt(-(2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2)) - (B^3*a^2 - B^3*b^2)*sqrt(tan(d*x + c))) - (a^2 + b^2)*d*sqrt(-(2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2))*log(-((a^5 + 2*a^3*b^2 + a*b^4)*d^3*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)) - (B^2*a^2*b - B^2*b^3)*d)*sqrt(-(2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2)) - (B^3*a^2 - B^3*b^2)*sqrt(tan(d*x + c))) - (a^2 + b^2)*d*sqrt(-(2*B^2*a*b - (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2))*log(((a^5 + 2*a^3*b^2 + a*b^4)*d^3*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)) + (B^2*a^2...`

3.424.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = B \int \frac{\sqrt{\tan(c+dx)}}{a+b\tan(c+dx)} dx$$

input `integrate(tan(d*x+c)**(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `B*Integral(sqrt(tan(c + d*x))/(a + b*tan(c + d*x)), x)`

3.424.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.73

$$\int \frac{\sqrt{\tan(c+dx)}(aB + bB \tan(c+dx))}{(a + b \tan(c+dx))^2} dx =$$

$$\frac{8 Bab \arctan\left(\frac{b\sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^2+b^2)\sqrt{ab}} - \frac{(2\sqrt{2}(a+b) \arctan\left(\frac{1}{2}\sqrt{2}(\sqrt{2}+2\sqrt{\tan(dx+c)})\right)) + 2\sqrt{2}(a+b) \arctan\left(-\frac{1}{2}\sqrt{2}(\sqrt{2}-2\sqrt{\tan(dx+c)})\right) - \sqrt{2}(a-b) \log\left(\frac{\sqrt{2}(\sqrt{2}+2\sqrt{\tan(dx+c)}) + \tan(dx+c) + 1}{\sqrt{2}(\sqrt{2}-2\sqrt{\tan(dx+c)}) + \tan(dx+c) + 1}\right)}{a^2+b^2}}{4d}$$

input `integrate(tan(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algo
rithm="maxima")`

output `-1/4*(8*B*a*b*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/((a^2 + b^2)*sqrt(a*b
) - (2*sqrt(2)*(a + b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c))
) + 2*sqrt(2)*(a + b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c))
) - sqrt(2)*(a - b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) +
sqrt(2)*(a - b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*B/(a^
2 + b^2))/d`

3.424.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{\tan(c+dx)}(aB + bB \tan(c+dx))}{(a + b \tan(c+dx))^2} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algo
rithm="giac")`

output `Timed out`

3.424.9 Mupad [B] (verification not implemented)

Time = 37.53 (sec) , antiderivative size = 17323, normalized size of antiderivative = 73.09

$$\int \frac{\sqrt{\tan(c+dx)}(aB + bB \tan(c+dx))}{(a + b \tan(c+dx))^2} dx = \text{Too large to display}$$

input `int((tan(c + d*x)^(1/2)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`

output `(log(- (((((((((256*B*a*b^3*(2*a^4 - b^4 + a^2*b^2))/d - 128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b^2)^2*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16*B^2*a^3*b^3*d^2 - 16*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16*B^2*a^3*b^3*d^2 - 16*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 - (64*B^2*a^3*b^2*tan(c + d*x)^(1/2)*(a^6 + 17*b^6 - 29*a^2*b^4 + 19*a^4*b^2))/(d^2*(a^2 + b^2)^2))*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16*B^2*a^3*b^3*d^2 - 16*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (32*B^3*a^4*b^2*(a^6 + 13*b^6 - 45*a^2*b^4 + 39*a^4*b^2))/(d^3*(a^2 + b^2)^3))*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16*B^2*a^3*b^3*d^2 - 16*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 - (16*B^4*a^4*b^3*tan(c + d*x)^(1/2)*(9*a^6 - 3*b^6 + 3*a^2*b^4 - 17*a^4*b^2))/(d^4*(a^2 + b^2)^4))*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16*B^2*a^3*b^3*d^2 - 16*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 - (8*B^5*a^5*b^3*(9*a^4 - b^4))/(d^5*(a^2 + b^2)^4))*(((192*B^4*a^6*b^6*d^4 - 16*B^4*a^4*b^8*d^4 - 16*B^4*a^12*d^4 - 608*B^4*a^8*b^4*d^4 + 192*B^4*a^10*b^2*d^4)^(1/2) + 16*B^2*a^3*b^3*d^2 - 16*B^2*a^5*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6*d^4 + 6*a^4*b^4*d^4 + 4*a^6*b^2*d^4))^(1/2))/4 + (log(- (((((((((256*B*a*b^3*(2*a^4 - b^4 + a^2*b^2))/d - 128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b^2)^2*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a^3*b^3*d^2 + 16*B^...`

3.425
$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)(a + b \tan(c + dx))^2}} dx$$

3.425.1 Optimal result 4108
 3.425.2 Mathematica [C] (verified) 4109
 3.425.3 Rubi [A] (verified) 4109
 3.425.4 Maple [A] (verified) 4115
 3.425.5 Fricas [B] (verification not implemented) 4115
 3.425.6 Sympy [F] 4116
 3.425.7 Maxima [A] (verification not implemented) 4117
 3.425.8 Giac [F(-1)] 4117
 3.425.9 Mupad [B] (verification not implemented) 4118

3.425.1 Optimal result

Integrand size = 36, antiderivative size = 237

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)(a + b \tan(c + dx))^2}} dx$$

$$= -\frac{(a - b)B \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)d} + \frac{(a - b)B \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)d}$$

$$+ \frac{2b^{3/2}B \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c + dx)}}{\sqrt{a}}\right)}{\sqrt{a}(a^2 + b^2)d} - \frac{(a + b)B \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)d}$$

$$+ \frac{(a + b)B \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)d}$$

output

```
1/2*(a-b)*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)+1/2*(a
-b)*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/4*(a+b)*B*ln
(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)+1/4*(a+b)*B*ln
(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)+2*b^(3/2)*B*a
rctan(b^(1/2)*tan(d*x+c)^(1/2)/a^(1/2))/(a^2+b^2)/d/a^(1/2)
```

3.425.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.12 (sec) , antiderivative size = 229, normalized size of antiderivative = 0.97

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx = B \left(\frac{2b^{3/2} \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{a}(a^2 + b^2)d} \right. \\ \left. - \frac{a\left(2\sqrt{2} \arctan\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right) - 2\sqrt{2} \arctan\left(1 + \sqrt{2}\sqrt{\tan(c+dx)}\right) + \sqrt{2} \log\left(1 - \sqrt{2}\sqrt{\tan(c+dx)}\right)\right)}{4(a^2 + b^2)d} \right. \\ \left. - \frac{2b \operatorname{Hypergeometric2F1}\left(\frac{3}{4}, 1, \frac{7}{4}, -\tan^2(c+dx)\right) \tan^{3/2}(c+dx)}{3(a^2 + b^2)d} \right)$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^2), x]`

output `B*((2*b^(3/2)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[a]*(a^2 + b^2)*d) - (a*(2*Sqrt[2]*ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - 2*Sqrt[2]*ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]] + Sqrt[2]*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Sqrt[2]*Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]))/(4*(a^2 + b^2)*d) - (2*b*Hypergeometric2F1[3/4, 1, 7/4, -Tan[c + d*x]^2]*Tan[c + d*x]^(3/2))/(3*(a^2 + b^2)*d))`

3.425.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.86, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {2011, 3042, 4057, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx \\ \downarrow \text{2011} \\ B \int \frac{1}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))} dx$$

3.425. $\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx$

$$\begin{aligned}
& \downarrow \text{3042} \\
& B \int \frac{1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx \\
& \downarrow \text{4057} \\
& B \left(\frac{b^2 \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2} + \frac{\int \frac{a-b\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} \right) \\
& \downarrow \text{3042} \\
& B \left(\frac{b^2 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2} + \frac{\int \frac{a-b\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx}{a^2+b^2} \right) \\
& \downarrow \text{4017} \\
& B \left(\frac{2 \int \frac{a-b\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} + \frac{b^2 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2} \right) \\
& \downarrow \text{1482} \\
& B \left(\frac{2 \left(\frac{1}{2}(a+b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a-b) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d(a^2+b^2)} + \frac{b^2 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2} \right) \\
& \downarrow \text{1476} \\
& B \left(\frac{2 \left(\frac{1}{2}(a+b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a-b) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) \right)}{d(a^2+b^2)} \right) \\
& \downarrow \text{1082} \\
& B \left(\frac{2 \left(\frac{1}{2}(a+b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a-b) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d\left(\frac{1-\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d\left(\frac{\sqrt{2}\sqrt{\tan(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) \right)}{d(a^2+b^2)} \right) \\
& \downarrow \text{217}
\end{aligned}$$

3.425. $\int \frac{aB+bB\tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2} dx$

$$B \left(\frac{2 \left(\frac{1}{2}(a+b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} + \frac{1}{2}(a-b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)} \right) +$$

↓ 1479

$$B \left(\frac{2 \left(\frac{1}{2}(a+b) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(a-b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)} \right) +$$

↓ 25

$$B \left(\frac{2 \left(\frac{1}{2}(a+b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(a-b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)} \right) +$$

↓ 27

$$B \left(\frac{2 \left(\frac{1}{2}(a+b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) + \frac{1}{2}(a-b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)} \right) +$$

↓ 1103

$$B \left(\frac{b^2 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} dx}{a^2+b^2} + \frac{2 \left(\frac{1}{2}(a-b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(a+b) \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right)}{d(a^2+b^2)} \right) +$$

↓ 4117

3.425. $\int \frac{aB+bB\tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^2} dx$

$$B \left(\frac{b^2 \int \frac{1}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))} d\tan(c+dx)}{d(a^2+b^2)} + \frac{2 \left(\frac{1}{2}(a-b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)} \right)$$

↓ 73

$$B \left(\frac{2b^2 \int \frac{1}{a+b\tan(c+dx)} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} + \frac{2 \left(\frac{1}{2}(a-b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) + \frac{1}{2}(a+b)}{d(a^2+b^2)} \right)$$

↓ 218

$$B \left(\frac{2 \left(\frac{1}{2}(a-b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) + \frac{1}{2}(a+b) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)-1})}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right)$$

input `Int[(a*B + b*B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^2),x]`

output `B*((2*b^(3/2)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[a]*(a^2 + b^2)*d) + (2*(((a - b)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]))/2 + ((a + b)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d))`

3.425.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
 -1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
 & (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
 t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
 implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
 eQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
 imp[d*(Log[RemoveContent[a + b*x + c*x^2, x])/b], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
 2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
 e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
 && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
 -2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
 x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
 reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4057 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)/((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[1/(c^2 + d^2) Int[(a + b*Tan[e + f*x])^m *(c - d*Tan[e + f*x]), x], x] + Simp[d^2/(c^2 + d^2) Int[(a + b*Tan[e + f*x])^m *((1 + Tan[e + f*x]^2)/(c + d*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

3.425.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 226, normalized size of antiderivative = 0.95

method	result
derivativedivides	$B \left(\frac{2b^2 \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{(a^2+b^2)\sqrt{ab}} + \frac{a\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})\right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})\right)}{4} \right)$
default	$B \left(\frac{2b^2 \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{(a^2+b^2)\sqrt{ab}} + \frac{a\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}{1-\sqrt{2}(\sqrt{\tan(dx+c)}+\tan(dx+c))}\right) + 2 \arctan(1+\sqrt{2}(\sqrt{\tan(dx+c)})\right) + 2 \arctan(-1+\sqrt{2}(\sqrt{\tan(dx+c)})\right)}{4} \right)$

```
input int((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^2,x,method=_RET
URNVERBOSE)
```

```
output 1/d*B*(2*b^2/(a^2+b^2)/(a*b)^(1/2)*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))+
2/(a^2+b^2)*(1/8*a*2^(1/2)*(ln((1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1-
2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))
+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))-1/8*b*2^(1/2)*(ln((1-2^(1/2)*tan(d
*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(
1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))))
```

3.425.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1565 vs. 2(199) = 398.

Time = 0.34 (sec) , antiderivative size = 3156, normalized size of antiderivative = 13.32

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

```
input integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^2,x, algo
rithm="fricas")
```

output `[-1/2*((a^2 + b^2)*d*sqrt((2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2))*log(((a^4*b + 2*a^2*b^3 + b^5)*d^3*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)) + (B^2*a^3 - B^2*a*b^2)*d)*sqrt((2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2)) - (B^3*a^2 - B^3*b^2)*sqrt(tan(d*x + c))) - (a^2 + b^2)*d*sqrt((2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2))*log(-((a^4*b + 2*a^2*b^3 + b^5)*d^3*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)) + (B^2*a^3 - B^2*a*b^2)*d)*sqrt((2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2)) - (B^3*a^2 - B^3*b^2)*sqrt(tan(d*x + c))) - (a^2 + b^2)*d*sqrt((2*B^2*a*b - (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2))*log(((a^4*b + 2*a^2*b^3 + b^5)*d^3*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)) - (B^2*a^3 - B...`

3.425.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx = B \int \frac{1}{a \sqrt{\tan(c + dx)} + b \tan^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+b*tan(d*x+c))**2,x)`

output `B*Integral(1/(a*sqrt(tan(c + d*x)) + b*tan(c + d*x)**(3/2)), x)`

3.425.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.73

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx$$

$$= \frac{8 B b^2 \arctan\left(\frac{b \sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^2+b^2)\sqrt{ab}} + \frac{\left(2 \sqrt{2}(a-b) \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2}+2 \sqrt{\tan(dx+c)})\right)\right) + 2 \sqrt{2}(a-b) \arctan\left(-\frac{1}{2} \sqrt{2}(\sqrt{2}-2 \sqrt{\tan(dx+c)})\right) + \sqrt{2}(a+b)}{a^2+b^2}$$

$$= \frac{4d}{4d}$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^2,x, algo
rithm="maxima")`

output `1/4*(8*B*b^2*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/((a^2 + b^2)*sqrt(a*b)
) + (2*sqrt(2)*(a - b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c)))
) + 2*sqrt(2)*(a - b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c)))
) + sqrt(2)*(a + b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1) - s
qrt(2)*(a + b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*B/(a^2
+ b^2))/d`

3.425.8 Giac [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^2} dx = \text{Timed out}$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^2,x, algo
rithm="giac")`

output `Timed out`

3.425.9 Mupad [B] (verification not implemented)

Time = 37.13 (sec) , antiderivative size = 16598, normalized size of antiderivative = 70.03

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)(a + b \tan(c + dx))^2}} dx = \text{Too large to display}$$

```
input int((B*a + B*b*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + b*tan(c + d*x))^2),x
)
```

```
output (log((((((((((128*B*b^2*(2*b^6 - a^6 + 9*a^2*b^4 + 6*a^4*b^2))/d + 128*b^3
*tan(c + d*x)^(1/2)*(a^2 - b^2)*(a^2 + b^2)^2*((4*(-B^4*a^4*d^4*(a^4 + b^4
- 6*a^2*b^2)^2)^(1/2) - 16*B^2*a^3*b^3*d^2 + 16*B^2*a^5*b*d^2)/(d^4*(a^2
+ b^2)^4))^(1/2))*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*
B^2*a^3*b^3*d^2 + 16*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (64*B^
2*a*b^2*tan(c + d*x)^(1/2)*(2*b^8 - a^8 + 5*a^2*b^6 + 67*a^4*b^4 - a^6*b^2
))/(d^2*(a^2 + b^2)^2))*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2)
- 16*B^2*a^3*b^3*d^2 + 16*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 -
(32*B^3*a*b^5*(25*a^6 + b^6 - 13*a^2*b^4 - 85*a^4*b^2))/(d^3*(a^2 + b^2)^3
))*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a^3*b^3*d^2
+ 16*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (16*B^4*a^2*b^5*tan(c
+ d*x)^(1/2)*(b^6 - 27*a^6 + 7*a^2*b^4 + 11*a^4*b^2))/(d^4*(a^2 + b^2)^4)
)*((4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) - 16*B^2*a^3*b^3*d^2
+ 16*B^2*a^5*b*d^2)/(d^4*(a^2 + b^2)^4))^(1/2))/4 + (16*B^5*a^4*b^6*(5*a^2
+ b^2))/(d^5*(a^2 + b^2)^4))*(((192*B^4*a^6*b^6*d^4 - 16*B^4*a^4*b^8*d^4
- 16*B^4*a^12*d^4 - 608*B^4*a^8*b^4*d^4 + 192*B^4*a^10*b^2*d^4)^(1/2) - 16
*B^2*a^3*b^3*d^2 + 16*B^2*a^5*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6*d^4 +
6*a^4*b^4*d^4 + 4*a^6*b^2*d^4))^(1/2))/4 + (log((((((((((128*B*b^2*(2*b^6
- a^6 + 9*a^2*b^4 + 6*a^4*b^2))/d + 128*b^3*tan(c + d*x)^(1/2)*(a^2 - b^2)
*(a^2 + b^2)^2*(-(4*(-B^4*a^4*d^4*(a^4 + b^4 - 6*a^2*b^2)^2)^(1/2) + 16...
```

3.426
$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$$

3.426.1 Optimal result 4119
 3.426.2 Mathematica [C] (verified) 4120
 3.426.3 Rubi [A] (verified) 4120
 3.426.4 Maple [A] (verified) 4127
 3.426.5 Fracas [B] (verification not implemented) 4127
 3.426.6 Sympy [F] 4128
 3.426.7 Maxima [A] (verification not implemented) 4129
 3.426.8 Giac [F(-1)] 4129
 3.426.9 Mupad [B] (verification not implemented) 4130

3.426.1 Optimal result

Integrand size = 36, antiderivative size = 256

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$$

$$= \frac{(a + b)B \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)d} - \frac{(a + b)B \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)d}$$

$$- \frac{2b^{5/2}B \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c + dx)}}{\sqrt{a}}\right)}{a^{3/2}(a^2 + b^2)d} - \frac{(a - b)B \log\left(1 - \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)d}$$

$$+ \frac{(a - b)B \log\left(1 + \sqrt{2}\sqrt{\tan(c + dx)} + \tan(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)d} - \frac{2B}{ad\sqrt{\tan(c + dx)}}$$

output

```
-2*b^(5/2)*B*arctan(b^(1/2)*tan(d*x+c)^(1/2)/a^(1/2))/a^(3/2)/(a^2+b^2)/d-
1/2*(a+b)*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/2*(a
+b)*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/4*(a-b)*B*1
n(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)+1/4*(a-b)*B*1
n(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(a^2+b^2)/d*2^(1/2)-2*B/a/d/tan(d
*x+c)^(1/2)
```

3.426.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.52

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$$

$$= \frac{B \left(-(-1)^{3/4}(a + ib) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right) - \frac{2b^{5/2} \arctan \left(\frac{\sqrt{b} \sqrt{\tan(c + dx)}}{\sqrt{a}} \right)}{a^{3/2}} + \sqrt[4]{-1}(ia + b) \operatorname{arctanh} \left(\frac{\sqrt{b} \sqrt{\tan(c + dx)}}{\sqrt{a}} \right) \right)}{(a^2 + b^2) d}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^2),x]`

output `(B*(-((-1)^(3/4)*(a + I*b)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]) - (2*b^(5/2)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/a^(3/2) + (-1)^(1/4)*(I*a + b)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] - (2*(a^2 + b^2))/(a*Sqrt[Tan[c + d*x]])))/((a^2 + b^2)*d)`

3.426.3 Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.89, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {2011, 3042, 4052, 27, 3042, 4136, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$$

$$\downarrow \text{2011}$$

$$B \int \frac{1}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$\downarrow \text{3042}$$

$$B \int \frac{1}{\tan(c + dx)^{3/2}(a + b \tan(c + dx))} dx$$

$$\begin{aligned}
& \downarrow 4052 \\
& B \left(-\frac{2 \int \frac{b \tan^2(c+dx) + a \tan(c+dx) + b}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a} - \frac{2}{ad\sqrt{\tan(c+dx)}} \right) \\
& \downarrow 27 \\
& B \left(-\frac{\int \frac{b \tan^2(c+dx) + a \tan(c+dx) + b}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a} - \frac{2}{ad\sqrt{\tan(c+dx)}} \right) \\
& \downarrow 3042 \\
& B \left(-\frac{\int \frac{b \tan(c+dx)^2 + a \tan(c+dx) + b}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a} - \frac{2}{ad\sqrt{\tan(c+dx)}} \right) \\
& \downarrow 4136 \\
& B \left(-\frac{\frac{\int \frac{\tan(c+dx)a^2 + ba}{\sqrt{\tan(c+dx)}} dx}{a^2 + b^2} + \frac{b^3 \int \frac{\tan^2(c+dx) + 1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2 + b^2}}{a} - \frac{2}{ad\sqrt{\tan(c+dx)}} \right) \\
& \downarrow 3042 \\
& B \left(-\frac{\frac{\int \frac{\tan(c+dx)a^2 + ba}{\sqrt{\tan(c+dx)}} dx}{a^2 + b^2} + \frac{b^3 \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2 + b^2}}{a} - \frac{2}{ad\sqrt{\tan(c+dx)}} \right) \\
& \downarrow 4017 \\
& B \left(-\frac{\frac{2 \int \frac{a(b+a \tan(c+dx))}{\tan^2(c+dx) + 1} d\sqrt{\tan(c+dx)}}{d(a^2 + b^2)} + \frac{b^3 \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2 + b^2}}{a} - \frac{2}{ad\sqrt{\tan(c+dx)}} \right) \\
& \downarrow 27 \\
& B \left(-\frac{\frac{2a \int \frac{b+a \tan(c+dx)}{\tan^2(c+dx) + 1} d\sqrt{\tan(c+dx)}}{d(a^2 + b^2)} + \frac{b^3 \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2 + b^2}}{a} - \frac{2}{ad\sqrt{\tan(c+dx)}} \right) \\
& \downarrow 1482
\end{aligned}$$

3.426. $\int \frac{aB + bB \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

$$B \left(\frac{2a \left(\frac{1}{2}(a+b) \int \frac{\tan(c+dx)+1}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} - \frac{1}{2}(a-b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d(a^2+b^2)} + \frac{b^3 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} \right) - \frac{2}{ad\sqrt{\tan(c+dx)}}$$

↓ 1476

$$B \left(\frac{2a \left(\frac{1}{2}(a+b) \left(\frac{1}{2} \int \frac{1}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} + \frac{1}{2} \int \frac{1}{\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)} \right) - \frac{1}{2}(a-b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d(a^2+b^2)} \right) - \frac{2}{ad\sqrt{\tan(c+dx)}}$$

↓ 1082

$$B \left(\frac{2a \left(\frac{1}{2}(a+b) \left(\frac{\int \frac{1}{-\tan(c+dx)-1} d(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\tan(c+dx)-1} d(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d(a^2+b^2)} \right) + \frac{b^3 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2}$$

↓ 217

$$B \left(\frac{2a \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \int \frac{1-\tan(c+dx)}{\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)} \right)}{d(a^2+b^2)} \right) + \frac{b^3 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2}$$

↓ 1479

$$B \left(\frac{2a \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}}{\tan(c+dx)}}{\tan(c+dx)} \right) \right)}{d(a^2+b^2)} \right) - \frac{2}{ad\sqrt{\tan(c+dx)}}$$

↓ 25

3.426. $\int \frac{aB+bB \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

$$B \left(\frac{2a \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2}(a-b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right) a$$

↓ 27

$$B \left(\frac{2a \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2}(a-b) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\tan(c+dx)}}{\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)+1}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\tan(c+dx)+1}}{\tan(c+dx)+\sqrt{2}} d\sqrt{\tan(c+dx)}}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right) a$$

↓ 1103

$$B \left(\frac{b^3 \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} dx}{a^2+b^2} + \frac{2a \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2}(a-b) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right)}{a}$$

↓ 4117

$$B \left(\frac{b^3 \int \frac{1}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))} d \tan(c+dx)}{d(a^2+b^2)} + \frac{2a \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2}(a-b) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right)}{a}$$

↓ 73

$$B \left(\frac{2b^3 \int \frac{1}{a+b \tan(c+dx)} d\sqrt{\tan(c+dx)}}{d(a^2+b^2)} + \frac{2a \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2}(a-b) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}{2\sqrt{2}} \right)}{d(a^2+b^2)} \right)}{a}$$

↓ 218

3.426. $\int \frac{aB+bB \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

$$B \left(\frac{2a \left(\frac{1}{2}(a+b) \left(\frac{\arctan(\sqrt{2}\sqrt{\tan(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a-b) \left(\frac{\log(\tan(c+dx)+\sqrt{2}\sqrt{\tan(c+dx)+1}}{2\sqrt{2}} - \frac{\log(\tan(c+dx)-\sqrt{2}\sqrt{\tan(c+dx)})}{2\sqrt{2}} \right) \right)}{d(a^2+b^2)} \right) a$$

input `Int[(a*B + b*B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^2),x]`

output `B*(-(((2*b^(5/2)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(Sqrt[a]*(a^2 + b^2)*d) + (2*a*(((a + b)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])/Sqrt[2]))/2 - ((a - b)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]/(2*Sqrt[2])))/2))/((a^2 + b^2)*d)/a) - 2/(a*d*Sqrt[Tan[c + d*x]]))`

3.426.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

3.426. $\int \frac{aB+bB \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4052 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(a^2 + b^2)*(b*c - a*d))), x] + Simp[1/((m + 1)*(a^2 + b^2)*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2) - b*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b^2*d*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && LtQ[m, -1] && (LtQ[n, 0] || IntegerQ[m]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4136 `Int((((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2))/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.426.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 242, normalized size of antiderivative = 0.95

method	result
derivativedivides	$B \left(-\frac{2b^3 \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{a(a^2+b^2)\sqrt{ab}} - \frac{2}{a\sqrt{\tan(dx+c)}} + \frac{b\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}\right) + 2 \arctan\left(1+\sqrt{2}(\sqrt{\tan(dx+c)})\right) \right)}{4} \right)$
default	$B \left(-\frac{2b^3 \arctan\left(\frac{b(\sqrt{\tan(dx+c)})}{\sqrt{ab}}\right)}{a(a^2+b^2)\sqrt{ab}} - \frac{2}{a\sqrt{\tan(dx+c)}} + \frac{b\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}{1-\sqrt{2}(\sqrt{\tan(dx+c)})+\tan(dx+c)}\right) + 2 \arctan\left(1+\sqrt{2}(\sqrt{\tan(dx+c)})\right) \right)}{4} \right)$

```
input int((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^2,x,method=_RET
URNVERBOSE)
```

```
output 1/d*B*(-2/a*b^3/(a^2+b^2)/(a*b)^(1/2)*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))
)-2/a/tan(d*x+c)^(1/2)+2/(a^2+b^2)*(-1/8*b*2^(1/2)*(ln((1+2^(1/2)*tan(d*x
+c)^(1/2)+tan(d*x+c))/(1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))+2*arctan(1+
2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2)))-1/8*a*2^(
1/2)*(ln((1-2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(1+2^(1/2)*tan(d*x+c)^(1/
2)+tan(d*x+c)))+2*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*t
an(d*x+c)^(1/2))))
```

3.426.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1644 vs. 2(216) = 432.

Time = 0.34 (sec) , antiderivative size = 3314, normalized size of antiderivative = 12.95

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

```
input integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^2,x, algo
rithm="fracas")
```

3.426. $\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$

output `[1/2*(2*B*b^2*sqrt(-b/a)*log(-(2*a*sqrt(-b/a)*sqrt(tan(d*x + c)) - b*tan(d*x + c) + a)/(b*tan(d*x + c) + a))*tan(d*x + c) - (a^3 + a*b^2)*d*sqrt(-(2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2))*log(((a^5 + 2*a^3*b^2 + a*b^4)*d^3*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)) - (B^2*a^2*b - B^2*b^3)*d)*sqrt(-(2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2)) - (B^3*a^2 - B^3*b^2)*sqrt(tan(d*x + c)))*tan(d*x + c) + (a^3 + a*b^2)*d*sqrt(-(2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2))*log(-((a^5 + 2*a^3*b^2 + a*b^4)*d^3*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)) - (B^2*a^2*b - B^2*b^3)*d)*sqrt(-(2*B^2*a*b + (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2)) - (B^3*a^2 - B^3*b^2)*sqrt(tan(d*x + c)))*tan(d*x + c) + (a^3 + a*b^2)*d*sqrt(-(2*B^2*a*b - (a^4 + 2*a^2*b^2 + b^4)*d^2*sqrt(-(B^4*a^4 - 2*B^4*a^2*b^2 + B^4*b^4)/((a^8 + 4*a^6*b^2 + 6*a^4*b^4 + 4*a^2*b^6 + b^8)*d^4)))/((a^4 + 2*a^2*b^2 + b^4)*d^2)))]`

3.426.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = B \int \frac{1}{a \tan^{\frac{3}{2}}(c + dx) + b \tan^{\frac{5}{2}}(c + dx)} dx$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)**(3/2)/(a+b*tan(d*x+c))**2,x)`

output `B*Integral(1/(a*tan(c + d*x)**(3/2) + b*tan(c + d*x)**(5/2)), x)`

3.426.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 188, normalized size of antiderivative = 0.73

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx =$$

$$\frac{8 B b^3 \arctan\left(\frac{b \sqrt{\tan(dx+c)}}{\sqrt{ab}}\right)}{(a^3+ab^2)\sqrt{ab}} + \frac{\left(2 \sqrt{2}(a+b) \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2}+2 \sqrt{\tan(dx+c)})\right)\right) + 2 \sqrt{2}(a+b) \arctan\left(-\frac{1}{2} \sqrt{2}(\sqrt{2}-2 \sqrt{\tan(dx+c)})\right) - \sqrt{2}(a+b)}{a^2+b^2}$$

$4d$

```
input integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^2,x, algo
rithm="maxima")
```

```
output -1/4*(8*B*b^3*arctan(b*sqrt(tan(d*x + c))/sqrt(a*b))/((a^3 + a*b^2)*sqrt(a
*b)) + (2*sqrt(2)*(a + b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2*sqrt(tan(d*x + c
)))) + 2*sqrt(2)*(a + b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2*sqrt(tan(d*x + c
)))) - sqrt(2)*(a - b)*log(sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1)
+ sqrt(2)*(a - b)*log(-sqrt(2)*sqrt(tan(d*x + c)) + tan(d*x + c) + 1))*B/(
a^2 + b^2) + 8*B/(a*sqrt(tan(d*x + c))))/d
```

3.426.8 Giac [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Timed out}$$

```
input integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^2,x, algo
rithm="giac")
```

```
output Timed out
```


3.426.9 Mupad [B] (verification not implemented)

Time = 28.98 (sec) , antiderivative size = 22906, normalized size of antiderivative = 89.48

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

```
input int((B*a + B*b*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + b*tan(c + d*x))^2),x
)
```

```
output (log((((192*B^4*a^6*b^6*d^4 - 16*B^4*a^4*b^8*d^4 - 16*B^4*a^12*d^4 - 608*
B^4*a^8*b^4*d^4 + 192*B^4*a^10*b^2*d^4)^(1/2) + 16*B^2*a^3*b^3*d^2 - 16*B^
2*a^5*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6*d^4 + 6*a^4*b^4*d^4 + 4*a^6*b^
2*d^4))^(1/2)*((((192*B^4*a^6*b^6*d^4 - 16*B^4*a^4*b^8*d^4 - 16*B^4*a^12*
d^4 - 608*B^4*a^8*b^4*d^4 + 192*B^4*a^10*b^2*d^4)^(1/2) + 16*B^2*a^3*b^3*d
^2 - 16*B^2*a^5*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6*d^4 + 6*a^4*b^4*d^4
+ 4*a^6*b^2*d^4))^(1/2)*((tan(c + d*x)^(1/2)*(1152*B^2*a^8*b^26*d^7 + 134
40*B^2*a^10*b^24*d^7 + 69056*B^2*a^12*b^22*d^7 + 202752*B^2*a^14*b^20*d^7
+ 372800*B^2*a^16*b^18*d^7 + 443136*B^2*a^18*b^16*d^7 + 337792*B^2*a^20*b^
14*d^7 + 156160*B^2*a^22*b^12*d^7 + 37632*B^2*a^24*b^10*d^7 + 3200*B^2*a^2
6*b^8*d^7 + 704*B^2*a^28*b^6*d^7 + 512*B^2*a^30*b^4*d^7 + 64*B^2*a^32*b^2*
d^7) - (((192*B^4*a^6*b^6*d^4 - 16*B^4*a^4*b^8*d^4 - 16*B^4*a^12*d^4 - 60
8*B^4*a^8*b^4*d^4 + 192*B^4*a^10*b^2*d^4)^(1/2) + 16*B^2*a^3*b^3*d^2 - 16*
B^2*a^5*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6*d^4 + 6*a^4*b^4*d^4 + 4*a^6*
b^2*d^4))^(1/2)*((tan(c + d*x)^(1/2)*(((192*B^4*a^6*b^6*d^4 - 16*B^4*a^4*b
^8*d^4 - 16*B^4*a^12*d^4 - 608*B^4*a^8*b^4*d^4 + 192*B^4*a^10*b^2*d^4)^(1/
2) + 16*B^2*a^3*b^3*d^2 - 16*B^2*a^5*b*d^2)/(a^8*d^4 + b^8*d^4 + 4*a^2*b^6
*d^4 + 6*a^4*b^4*d^4 + 4*a^6*b^2*d^4))^(1/2)*(512*a^9*b^27*d^9 + 5120*a^11
*b^25*d^9 + 22528*a^13*b^23*d^9 + 56320*a^15*b^21*d^9 + 84480*a^17*b^19*d^
9 + 67584*a^19*b^17*d^9 - 67584*a^23*b^13*d^9 - 84480*a^25*b^11*d^9 - 5...
```

3.427 $\int \tan^{\frac{3}{2}}(c+dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.427.1 Optimal result	4131
3.427.2 Mathematica [A] (verified)	4132
3.427.3 Rubi [A] (verified)	4132
3.427.4 Maple [B] (warning: unable to verify)	4136
3.427.5 Fricas [B] (verification not implemented)	4136
3.427.6 Sympy [F]	4137
3.427.7 Maxima [F]	4137
3.427.8 Giac [F(-1)]	4137
3.427.9 Mupad [F(-1)]	4138

3.427.1 Optimal result

Integrand size = 35, antiderivative size = 264

$$\int \tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\sqrt{ia - b}(iA - B) \arctan\left(\frac{\sqrt{ia - b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{d}$$

$$+ \frac{(4aAb - a^2B - 8b^2B) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{4b^{3/2}d}$$

$$+ \frac{\sqrt{ia + b}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia + b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{d}$$

$$+ \frac{(4Ab - aB) \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}{4bd} + \frac{B \sqrt{\tan(c + dx)} (a + b \tan(c + dx))^{3/2}}{2bd}$$

```
output 1/4*(4*A*a*b-B*a^2-8*B*b^2)*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/b^(3/2)/d+(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a-b)^(1/2)/d+(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a+b)^(1/2)/d+1/4*(4*A*b-B*a)*tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/b/d+1/2*B*tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(3/2)/b/d
```

3.427.2 Mathematica [A] (verified)

Time = 3.73 (sec) , antiderivative size = 304, normalized size of antiderivative = 1.15

$$\int \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$= \frac{-\sqrt{a}(-4aAb+a^2B+8b^2B)\operatorname{arcsinh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)\sqrt{1+\frac{b\tan(c+dx)}{a}}+\sqrt{b}\left(-4\sqrt[4]{-1}\sqrt{-a+ibb}(iA+B)\right)}{\dots}$$

input `Integrate[Tan[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(-(Sqrt[a]*(-4*a*A*b + a^2*B + 8*b^2*B)*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a]) + Sqrt[b]*(-4*(-1)^(1/4)*Sqrt[-a + I*b]*b*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])*Sqrt[a + b*Tan[c + d*x]] + 4*(-1)^(3/4)*Sqrt[a + I*b]*b*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])*Sqrt[a + b*Tan[c + d*x]] + Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])*(4*A*b + a*B + 2*b*B*Tan[c + d*x]))/(4*b^(3/2)*d*Sqrt[a + b*Tan[c + d*x]])`

3.427.3 Rubi [A] (verified)

Time = 1.42 (sec) , antiderivative size = 265, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 4090, 27, 3042, 4130, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c+dx)^{3/2}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow \text{4090}$$

$$\begin{aligned}
& \frac{\int -\frac{\sqrt{a+b \tan(c+dx)}(-((4Ab-aB) \tan^2(c+dx))+4bB \tan(c+dx)+aB)}{2\sqrt{\tan(c+dx)}} dx}{\frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{2bd}} + \\
& \quad \downarrow 27 \\
& \frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{2bd} - \frac{\int \frac{\sqrt{a+b \tan(c+dx)}(-((4Ab-aB) \tan^2(c+dx))+4bB \tan(c+dx)+aB)}{\sqrt{\tan(c+dx)}} dx}{4b} \\
& \quad \downarrow 3042 \\
& \frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{2bd} - \frac{\int \frac{\sqrt{a+b \tan(c+dx)}(-((4Ab-aB) \tan(c+dx)^2)+4bB \tan(c+dx)+aB)}{\sqrt{\tan(c+dx)}} dx}{4b} \\
& \quad \downarrow 4130 \\
& \frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{2bd} - \frac{\int \frac{-((-Ba^2+4Aba-8b^2B) \tan^2(c+dx))+8b(Ab+aB) \tan(c+dx)+a(4Ab+aB)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{(4Ab-aB)\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}}{d}}{4b} \\
& \quad \downarrow 27 \\
& \frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{2bd} - \frac{\frac{1}{2} \int \frac{-((-Ba^2+4Aba-8b^2B) \tan^2(c+dx))+8b(Ab+aB) \tan(c+dx)+a(4Ab+aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{(4Ab-aB)\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}}{d}}{4b} \\
& \quad \downarrow 3042 \\
& \frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{2bd} - \frac{\frac{1}{2} \int \frac{-((-Ba^2+4Aba-8b^2B) \tan(c+dx)^2)+8b(Ab+aB) \tan(c+dx)+a(4Ab+aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{(4Ab-aB)\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}}{d}}{4b} \\
& \quad \downarrow 4138 \\
& \frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{2bd} - \frac{\int \frac{-((-Ba^2+4Aba-8b^2B) \tan^2(c+dx))+8b(Ab+aB) \tan(c+dx)+a(4Ab+aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)}}{2d} d \tan(c+dx) - \frac{(4Ab-aB)\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}}{d}}{4b} \\
& \quad \downarrow 2035
\end{aligned}$$

3.427. $\int \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$

$$\begin{aligned}
 & \frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2bd} - \frac{\int \frac{-((-Ba^2+4Aba-8b^2B)\tan^2(c+dx)+8b(Ab+aB)\tan(c+dx)+a(4Ab+aB))d\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}}{d} - \frac{(4Ab-aB)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} \\
 & \qquad \qquad \qquad \downarrow \text{2257} \\
 & \frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2bd} - \frac{\int \left(\frac{Ba^2-4Aba+8b^2B}{\sqrt{a+b\tan(c+dx)}} + \frac{8(b(aA-bB)+b(Ab+aB)\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} \right) d\sqrt{\tan(c+dx)}}{d} - \frac{(4Ab-aB)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} \\
 & \qquad \qquad \qquad \downarrow \text{2009} \\
 & \frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2bd} - \frac{(a^2(-B)+4aAb-8b^2B)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{b}} - \frac{(4Ab-aB)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} + \frac{-4b\sqrt{-b+ia}(-B+ia)\operatorname{arctan}\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d}
 \end{aligned}$$

```
input Int[Tan[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]
```

```
output (B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2))/(2*b*d) - ((-4*Sqrt[I*a - b]*b*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - ((4*a*A*b - a^2*B - 8*b^2*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[b] - 4*b*Sqrt[I*a + b]*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/d - ((4*A*b - a*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d)/(4*b)
```

3.427.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2035 `Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]`

rule 2257 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol
] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a
, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.
) + (f_.)*(x_)])^2, x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))`

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

3.427.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.76 (sec) , antiderivative size = 2183234, normalized size of antiderivative = 8269.83

output too large to display

```
input int((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^(3/2)*(A+B*tan(d*x+c)),x)
```

```
output result too large to display
```

3.427.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8056 vs. $2(212) = 424$.

Time = 3.48 (sec) , antiderivative size = 16118, normalized size of antiderivative = 61.05

$$\int \tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^(3/2)*(A+B*tan(d*x+c)),x, algo
rithm="fracas")
```

```
output Too large to include
```

3.427.6 Sympy [F]

$$\int \tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} \tan^{\frac{3}{2}}(c + dx) dx$$

input `integrate((a+b*tan(d*x+c))**(1/2)*tan(d*x+c)**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))*tan(c + d*x)**(3/2), x)`

3.427.7 Maxima [F]

$$\int \tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \tan(dx + c)^{\frac{3}{2}} dx$$

input `integrate((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*tan(d*x + c)^(3/2), x)`

3.427.8 Giac [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.427.9 Mupad [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^{3/2} (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx$$

input `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`output `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2), x)`

3.428 $\int \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.428.1 Optimal result	4139
3.428.2 Mathematica [A] (verified)	4140
3.428.3 Rubi [A] (verified)	4140
3.428.4 Maple [B] (warning: unable to verify)	4143
3.428.5 Fricas [B] (verification not implemented)	4143
3.428.6 Sympy [F]	4143
3.428.7 Maxima [F]	4144
3.428.8 Giac [F]	4144
3.428.9 Mupad [B] (verification not implemented)	4144

3.428.1 Optimal result

Integrand size = 35, antiderivative size = 201

$$\int \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\sqrt{ia - b}(A + iB) \arctan\left(\frac{\sqrt{ia - b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{d} + \frac{(2Ab + aB) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{\sqrt{bd}}$$

$$- \frac{\sqrt{ia + b}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia + b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{d} + \frac{B \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}{d}$$

```
output (A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a
-b)^(1/2)/d+(2*A*b+B*a)*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(
1/2))/d/b^(1/2)-(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d
*x+c))^(1/2))*(I*a+b)^(1/2)/d+B*tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/d
```

3.428.2 Mathematica [A] (verified)

Time = 2.97 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.19

$$\int \sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$= \frac{\sqrt[4]{-1}\sqrt{-a+ib}(A-iB)\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) + \sqrt[4]{-1}\sqrt{a+ib}(A+iB)\arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d}$$

input `Integrate[Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `((-1)^(1/4)*Sqrt[-a + I*b]*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + (-1)^(1/4)*Sqrt[a + I*b]*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + B*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]] + ((2*A*b + a*B)*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[a + b*Tan[c + d*x]])/(Sqrt[a]*Sqrt[b]*Sqrt[1 + (b*Tan[c + d*x])/a]))/d`

3.428.3 Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 199, normalized size of antiderivative = 0.99, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 4093, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow \text{4093}$$

$$\int \frac{-((2Ab+aB)\tan^2(c+dx)) - 2(aA-bB)\tan(c+dx) + aB}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx + \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d}$$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{1}{2} \int \frac{-((2Ab+aB)\tan^2(c+dx)) - 2(aA-bB)\tan(c+dx) + aB}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \\
& \downarrow 3042 \\
& \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{1}{2} \int \frac{-((2Ab+aB)\tan(c+dx)^2) - 2(aA-bB)\tan(c+dx) + aB}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \\
& \downarrow 4138 \\
& \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{\int \frac{-((2Ab+aB)\tan^2(c+dx)) - 2(aA-bB)\tan(c+dx) + aB}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} d\tan(c+dx)}{2d} \\
& \downarrow 2035 \\
& \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{\int \frac{-((2Ab+aB)\tan^2(c+dx)) - 2(aA-bB)\tan(c+dx) + aB}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} d\sqrt{\tan(c+dx)}}{d} \\
& \downarrow 2257 \\
& \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{\int \left(\frac{-2Ab-aB}{\sqrt{a+b\tan(c+dx)}} + \frac{2(Ab+aB-(aA-bB)\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} \right) d\sqrt{\tan(c+dx)}}{d} \\
& \downarrow 2009 \\
& \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{-\sqrt{-b+ia}(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) - \frac{(aB+2Ab)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{b}} + \sqrt{b+ia}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d}
\end{aligned}$$

input `Int[Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `-((- (Sqrt[I*a - b]*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) - ((2*A*b + a*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[b] + Sqrt[I*a + b]*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/d) + (B*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d`

3.428. $\int \sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx)) dx$

3.428.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2035 `Int[(F_x_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[F_x, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[F_x, x]`
- rule 2257 `Int[(P_x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[P_x*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[P_x, x] && IntegerQ[p]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4093 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(p_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(m + n) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (A*b*c + a*B*c + a*A*d - b*B*d)*(m + n)*Tan[e + f*x] + (A*b*d*(m + n) + B*(a*d*m + b*c*n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[0, m, 1] && LtQ[0, n, 1]`
- rule 4138 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]) + (C_)*tan[(e_) + (f_)*(x_)^2], x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.428.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.43 (sec) , antiderivative size = 2180698, normalized size of antiderivative = 10849.24

output too large to display

input `int(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x)`

output `result too large to display`

3.428.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7955 vs. $2(161) = 322$.

Time = 3.37 (sec) , antiderivative size = 15912, normalized size of antiderivative = 79.16

$$\int \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `Too large to include`

3.428.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx \\ &= \int (A+B \tan(c+dx)) \sqrt{a+b \tan(c+dx)} \sqrt{\tan(c+dx)} dx \end{aligned}$$

input `integrate(tan(d*x+c)**(1/2)*(a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))*sqrt(tan(c + d*x)), x)`

3.428. $\int \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$

3.428.7 Maxima [F]

$$\int \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$= \int (B \tan(dx+c) + A) \sqrt{b \tan(dx+c) + a} \sqrt{\tan(dx+c)} dx$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*sqrt(tan(d*x + c))
, x)`

3.428.8 Giac [F]

$$\int \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$= \int (B \tan(dx+c) + A) \sqrt{b \tan(dx+c) + a} \sqrt{\tan(dx+c)} dx$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algo
rithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*sqrt(tan(d*x + c))
, x)`

3.428.9 Mupad [B] (verification not implemented)

Time = 132.15 (sec) , antiderivative size = 61200, normalized size of antiderivative = 304.48

$$\int \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx = \text{Too large to display}$$

input `int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`

output

$$\begin{aligned}
& ((2*B*a*\tan(c + d*x)^{(1/2)})/((a + b*\tan(c + d*x))^{(1/2)} - a^{(1/2)}) + (2*B* \\
& a*b*\tan(c + d*x)^{(3/2)})/((a + b*\tan(c + d*x))^{(1/2)} - a^{(1/2)})^3)/(d + (b^ \\
& 2*d*\tan(c + d*x)^2)/((a + b*\tan(c + d*x))^{(1/2)} - a^{(1/2)})^4 - (2*b*d*\tan(\\
& c + d*x))/((a + b*\tan(c + d*x))^{(1/2)} - a^{(1/2)})^2 - \operatorname{atan}(\frac{(A^2*b - A^2 \\
& *a^1i + B^2*a^1i - B^2*b + 2*A*B*a + A*B*b*2i)}{(4*d^2)})^{(1/2)} * (\frac{(A^2*b - \\
& A^2*a^1i + B^2*a^1i - B^2*b + 2*A*B*a + A*B*b*2i)}{(4*d^2)})^{(1/2)} * (\frac{(A^2*b \\
& - A^2*a^1i + B^2*a^1i - B^2*b + 2*A*B*a + A*B*b*2i)}{(4*d^2)})^{(1/2)} * (\frac{(27 \\
& 4877906944*(1600*a^{12}*b^{34}*d^8 - 16640*a^{14}*b^{32}*d^8 + 22784*a^{16}*b^{30}*d^8 \\
& + 106496*a^{18}*b^{28}*d^8 + 65536*a^{20}*b^{26}*d^8))}{d^8} - (274877906944*\tan(c \\
& + d*x)*(1600*a^{12}*b^{35}*d^8 - 48000*a^{14}*b^{33}*d^8 + 155136*a^{16}*b^{31}*d^8 + \\
& 466944*a^{18}*b^{29}*d^8 + 262144*a^{20}*b^{27}*d^8))}{(d^8*((a + b*\tan(c + d*x))^{(\\
& 1/2)} - a^{(1/2)})^2)} * (\frac{(A^2*b - A^2*a^1i + B^2*a^1i - B^2*b + 2*A*B*a + A*B* \\
& b*2i)}{(4*d^2)})^{(1/2)} - (219902325552*\tan(c + d*x)^{(1/2)} * (2048*A*a^{20}*b^{27} \\
& *d^6 - 12536*A*a^{16}*b^{31}*d^6 - 3328*A*a^{18}*b^{29}*d^6 - 7160*A*a^{14}*b^{33}*d^6 \\
& + 240*B*a^{13}*b^{34}*d^6 - 720*B*a^{15}*b^{32}*d^6 + 5696*B*a^{17}*b^{30}*d^6 + 1484 \\
& 8*B*a^{19}*b^{28}*d^6 + 8192*B*a^{21}*b^{26}*d^6))}{(d^7*((a + b*\tan(c + d*x))^{(1/2)} \\
&) - a^{(1/2)})} * (\frac{(A^2*b - A^2*a^1i + B^2*a^1i - B^2*b + 2*A*B*a + A*B*b*2i)} \\
& / (4*d^2))^{(1/2)} - (274877906944*(1200*A^2*a^{12}*b^{35}*d^6 - 1600*A^2*a^{14}*b^{ \\
& 33}*d^6 + 272464*A^2*a^{16}*b^{31}*d^6 + 573952*A^2*a^{18}*b^{29}*d^6 + 299008*A^2* \\
& a^{20}*b^{27}*d^6 - 1440*B^2*a^{12}*b^{35}*d^6 + 8352*B^2*a^{14}*b^{33}*d^6 - 320*B...
\end{aligned}$$

$$3.429 \quad \int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.429.1 Optimal result	4146
3.429.2 Mathematica [A] (verified)	4146
3.429.3 Rubi [A] (verified)	4147
3.429.4 Maple [B] (warning: unable to verify)	4151
3.429.5 Fricas [B] (verification not implemented)	4151
3.429.6 Sympy [F]	4151
3.429.7 Maxima [F]	4152
3.429.8 Giac [F(-1)]	4152
3.429.9 Mupad [B] (verification not implemented)	4152

3.429.1 Optimal result

Integrand size = 35, antiderivative size = 169

$$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

$$= -\frac{\sqrt{ia-b}(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} + \frac{2\sqrt{b}B \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$- \frac{\sqrt{ia+b}(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

output `-(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a-b)^(1/2)/d+2*B*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*b^(1/2)/d-(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a+b)^(1/2)/d`

3.429.2 Mathematica [A] (verified)

Time = 1.10 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.21

$$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

$$= \frac{\sqrt[4]{-1}\left(\sqrt{-a+ib}(iA+B) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) + \sqrt{a+ib}(-iA+B) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)\right)}{d}$$

3.429. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

input `Integrate[(Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `((-1)^(1/4)*(Sqrt[-a + I*b]*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) + Sqrt[a + I*b]*((-I)*A + B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) + (2*Sqrt[a]*Sqrt[b]*B*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/Sqrt[a + b*Tan[c + d*x]]/d`

3.429.3 Rubi [A] (verified)

Time = 1.18 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.06, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 4097, 3042, 4099, 3042, 4098, 104, 216, 219, 4117, 65, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
 & \quad \downarrow \text{4097} \\
 & \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx + bB \int \frac{\tan^2(c + dx) + 1}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx + bB \int \frac{\tan(c + dx)^2 + 1}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4099} \\
 & \frac{1}{2}(a + ib)(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(a - ib)(A - \\
 & iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx + bB \int \frac{\tan(c + dx)^2 + 1}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{2}(a+ib)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}(a-ib)(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx + bB \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx \\
& \quad \downarrow 4098 \\
& \frac{(a-ib)(A-iB) \int \frac{1}{(1-i \tan(c+dx)) \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{2d} + \\
& \frac{(a+ib)(A+iB) \int \frac{1}{(i \tan(c+dx)+1) \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{2d} + \\
& bB \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx \\
& \quad \downarrow 104 \\
& \frac{(a+ib)(A+iB) \int \frac{1}{\frac{(ia-b) \tan(c+dx)}{a+b \tan(c+dx)}+1} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} + \\
& \frac{(a-ib)(A-iB) \int \frac{1}{1-\frac{(ia+b) \tan(c+dx)}{a+b \tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} + bB \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx \\
& \quad \downarrow 216 \\
& \frac{(a-ib)(A-iB) \int \frac{1}{1-\frac{(ia+b) \tan(c+dx)}{a+b \tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} + bB \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx + \\
& \frac{(a+ib)(A+iB) \arctan\left(\frac{\sqrt{-b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d \sqrt{-b+ia}} \\
& \quad \downarrow 219 \\
& bB \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx + \frac{(a+ib)(A+iB) \arctan\left(\frac{\sqrt{-b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d \sqrt{-b+ia}} + \\
& \frac{(a-ib)(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d \sqrt{b+ia}} \\
& \quad \downarrow 4117 \\
& \frac{bB \int \frac{1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{d} + \frac{(a+ib)(A+iB) \arctan\left(\frac{\sqrt{-b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d \sqrt{-b+ia}} + \\
& \frac{(a-ib)(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d \sqrt{b+ia}} \\
& \quad \downarrow 65
\end{aligned}$$

3.429. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

$$\begin{aligned}
& \frac{2bB \int \frac{1}{1 - \frac{b \tan(c+dx)}{a+b \tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} + \frac{(a+ib)(A+iB) \arctan\left(\frac{\sqrt{-b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d\sqrt{-b+ia}} + \\
& \frac{(a-ib)(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d\sqrt{b+ia}} \\
& \quad \downarrow \text{219} \\
& \frac{(a+ib)(A+iB) \arctan\left(\frac{\sqrt{-b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d\sqrt{-b+ia}} + \frac{(a-ib)(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d\sqrt{b+ia}} + \\
& \frac{2\sqrt{b}B \operatorname{arctanh}\left(\frac{\sqrt{b} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}
\end{aligned}$$

input `Int[(Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `((a + I*b)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a - b]*d) + (2*Sqrt[b]*B*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/d + ((a - I*b)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d)`

3.429.3.1 Defintions of rubi rules used

rule 65 `Int[1/(Sqrt[(b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 104 `Int[(((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4097 `Int[(Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)])*((A_) + (B_)*tan[(e_) + (f_)*(x_)])]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)], x_Symbol] := Int[Simp[a*A - b*B + (A*b + a*B)*Tan[e + f*x], x]/(Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]), x] + Simp[b*B Int[(1 + Tan[e + f*x]^2)/(Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4098 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

3.429.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 2.28 (sec) , antiderivative size = 2178123, normalized size of antiderivative = 12888.30

output too large to display

input `int((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x)`

output `result too large to display`

3.429.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7969 vs. $2(130) = 260$.

Time = 2.38 (sec) , antiderivative size = 15940, normalized size of antiderivative = 94.32

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algo rithm="fricas")`

output `Too large to include`

3.429.6 Sympy [F]

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \int \frac{(A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)}}{\sqrt{\tan(c + dx)}} dx$$

input `integrate((a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))/sqrt(tan(c + d*x)), x)`

3.429.7 Maxima [F]

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)\sqrt{b \tan(dx + c) + a}}{\sqrt{\tan(dx + c)}} dx$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)/sqrt(tan(d*x + c)), x)`

3.429.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm="giac")`

output `Timed out`

3.429.9 Mupad [B] (verification not implemented)

Time = 20.29 (sec) , antiderivative size = 1141, normalized size of antiderivative = 6.75

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Too large to display}$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2))/tan(c + d*x)^(1/2),x)`

output

$$\begin{aligned} & \operatorname{atanh}\left(\frac{a^{3/2}d \tan(c+dx)^{1/2} \left((-A^4 a^2 d^4)^{1/2} - A^2 b d^2 \right) / d^4}{(-A^4 a^2 d^4)^{1/2} - a d \tan(c+dx)^{1/2} \left((-A^4 a^2 d^4)^{1/2} - A^2 b d^2 \right) / d^4} \right)^{1/2} \\ & + A^2 a^{3/2} b d^3 \tan(c+dx)^{1/2} \left((-A^4 a^2 d^4)^{1/2} - A^2 b d^2 \right) / d^4 \\ & - A^2 a b d^3 \tan(c+dx)^{1/2} \left((-A^4 a^2 d^4)^{1/2} - A^2 b d^2 \right) / d^4 \\ & + (A^3 a^3 d^2 - A a b (-A^4 a^2 d^4)^{1/2} - A b^2 \tan(c+dx) (-A^4 a^2 d^4)^{1/2} - A^3 a^{5/2} d^2 (a + b \tan(c+dx))^{1/2} \\ & + A^3 a^2 b d^2 \tan(c+dx) + A a^{1/2} b (a + b \tan(c+dx))^{1/2} (-A^4 a^2 d^4)^{1/2}) \left((-A^4 a^2 d^4)^{1/2} - A^2 b d^2 \right) / d^4 \\ & - \operatorname{atanh}\left(\frac{a^{3/2}d \tan(c+dx)^{1/2} \left((-A^4 a^2 d^4)^{1/2} + A^2 b d^2 \right) / d^4}{(-A^4 a^2 d^4)^{1/2} + a d \tan(c+dx)^{1/2} \left((-A^4 a^2 d^4)^{1/2} + A^2 b d^2 \right) / d^4} \right)^{1/2} \\ & + A^2 a^{3/2} b d^3 \tan(c+dx)^{1/2} \left((-A^4 a^2 d^4)^{1/2} + A^2 b d^2 \right) / d^4 \\ & + A^2 a b d^3 \tan(c+dx)^{1/2} \left((-A^4 a^2 d^4)^{1/2} + A^2 b d^2 \right) / d^4 \\ & + (A^3 a^3 d^2 + A a b (-A^4 a^2 d^4)^{1/2} + A b^2 \tan(c+dx) (-A^4 a^2 d^4)^{1/2} - A^3 a^{5/2} d^2 (a + b \tan(c+dx))^{1/2} \\ & + A^3 a^2 b d^2 \tan(c+dx) - A a^{1/2} b (a + b \tan(c+dx))^{1/2} (-A^4 a^2 d^4)^{1/2}) \left((-A^4 a^2 d^4)^{1/2} + A^2 b d^2 \right) / d^4 \\ & + \operatorname{atanh}\left(\frac{2 \left(a^{1/2} d \tan(c+dx)^{1/2} \left((-B^4 a^2 d^4)^{1/2} + B^2 b d^2 \right) / d^4 \right)}{(-B^4 a^2 d^4)^{1/2} + B^2 b d^2} \right) / 2 - (d \dots \end{aligned}$$

$$3.430 \quad \int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.430.1 Optimal result	4154
3.430.2 Mathematica [A] (verified)	4154
3.430.3 Rubi [A] (verified)	4155
3.430.4 Maple [B] (warning: unable to verify)	4158
3.430.5 Fricas [B] (verification not implemented)	4159
3.430.6 Sympy [F]	4159
3.430.7 Maxima [F(-1)]	4159
3.430.8 Giac [F(-1)]	4160
3.430.9 Mupad [F(-1)]	4160

3.430.1 Optimal result

Integrand size = 35, antiderivative size = 154

$$\begin{aligned} & \int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx \\ &= -\frac{\sqrt{ia-b}(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} \\ & \quad + \frac{\sqrt{ia+b}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} - \frac{2A\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \end{aligned}$$

output $-(A+I*B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})*(I*a-b)^{(1/2)}/d+(A-I*B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})*(I*a+b)^{(1/2)}/d-2*A*(a+b*\tan(d*x+c))^{(1/2)}/d/\tan(d*x+c)^{(1/2)}$

3.430.2 Mathematica [A] (verified)

Time = 1.12 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.10

$$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx = \frac{\sqrt[4]{-1}\sqrt{-a+ib}(A-iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) + \sqrt[4]{-1}\sqrt{a+ib}(A+iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

3.430. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

input `Integrate[(Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `-(((-1)^(1/4)*Sqrt[-a + I*b]*(A - I*B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]] + (-1)^(1/4)*Sqrt[a + I*b]*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]] + (2*A*Sqrt[a + b*Tan[c + d*x]])/Sqrt[Tan[c + d*x]]/d)`

3.430.3 Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.10, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4091, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx \\
 & \quad \downarrow \text{4091} \\
 & -2 \int -\frac{Ab + aB - (aA - bB) \tan(c + dx)}{2\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx - \frac{2A\sqrt{a + b \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx - \frac{2A\sqrt{a + b \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{Ab + aB - (aA - bB) \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx - \frac{2A\sqrt{a + b \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{4099}
 \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{2}(a+ib)(-B+iA) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}(a-ib)(B+iA) \\
& \quad iA \int \frac{i \tan(c+dx)+1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx - \frac{2A \sqrt{a+b \tan(c+dx)}}{d \sqrt{\tan(c+dx)}} \\
& \quad \downarrow 3042 \\
& -\frac{1}{2}(a+ib)(-B+iA) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}(a-ib)(B+iA) \\
& \quad iA \int \frac{i \tan(c+dx)+1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx - \frac{2A \sqrt{a+b \tan(c+dx)}}{d \sqrt{\tan(c+dx)}} \\
& \quad \downarrow 4098 \\
& \frac{(a-ib)(B+iA) \int \frac{1}{(1-i \tan(c+dx)) \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{2d} - \\
& \frac{(a+ib)(-B+iA) \int \frac{1}{(i \tan(c+dx)+1) \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{2d} - \frac{2A \sqrt{a+b \tan(c+dx)}}{d \sqrt{\tan(c+dx)}} \\
& \quad \downarrow 104 \\
& -\frac{(a+ib)(-B+iA) \int \frac{1}{\frac{(ia-b) \tan(c+dx)}{a+b \tan(c+dx)}+1} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} + \\
& \frac{(a-ib)(B+iA) \int \frac{1}{1-\frac{(ia+b) \tan(c+dx)}{a+b \tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} - \frac{2A \sqrt{a+b \tan(c+dx)}}{d \sqrt{\tan(c+dx)}} \\
& \quad \downarrow 216 \\
& \frac{(a-ib)(B+iA) \int \frac{1}{1-\frac{(ia+b) \tan(c+dx)}{a+b \tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} - \\
& \frac{(a+ib)(-B+iA) \arctan\left(\frac{\sqrt{-b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d \sqrt{-b+ia}} - \frac{2A \sqrt{a+b \tan(c+dx)}}{d \sqrt{\tan(c+dx)}} \\
& \quad \downarrow 219 \\
& -\frac{(a+ib)(-B+iA) \arctan\left(\frac{\sqrt{-b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d \sqrt{-b+ia}} + \frac{(a-ib)(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d \sqrt{b+ia}} - \\
& \quad \frac{2A \sqrt{a+b \tan(c+dx)}}{d \sqrt{\tan(c+dx)}}
\end{aligned}$$

input `Int[(Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

```
output -(((a + I*b)*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a +
b*Tan[c + d*x]])/(Sqrt[I*a - b]*d) + ((a - I*b)*(I*A + B)*ArcTanh[(Sqrt[
I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d)
- (2*A*Sqrt[a + b*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]])
```

3.430.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 104 Int[(((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)))/((e_.) + (f_.)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4091 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m +
1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n)
+ A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan
[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x] /; FreeQ
[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || Integers
Q[2*m, 2*n])
```

```
rule 4098 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*
x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]
```

```
rule 4099 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2,
0]
```

3.430.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.06 (sec) , antiderivative size = 2176213, normalized size of antiderivative = 14131.25

output too large to display

```
input int((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x)
```

```
output result too large to display
```

$$3.430. \quad \int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.430.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7865 vs. $2(122) = 244$.

Time = 1.41 (sec) , antiderivative size = 7865, normalized size of antiderivative = 51.07

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algo
rithm="fricas")`

output Too large to include

3.430.6 Sympy [F]

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)}}{\tan^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))/tan(c + d*x)**(3/2)
, x)`

3.430.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algo
rithm="maxima")`

output Timed out

3.430. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

3.430.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorith="giac")`

output `Timed out`

3.430.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)}}{\tan(c + dx)^{3/2}} dx \end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2))/tan(c + d*x)^(3/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2))/tan(c + d*x)^(3/2),x)`

$$3.431 \quad \int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.431.1 Optimal result 4161
 3.431.2 Mathematica [A] (verified) 4161
 3.431.3 Rubi [A] (verified) 4162
 3.431.4 Maple [B] (warning: unable to verify) 4166
 3.431.5 Fricas [B] (verification not implemented) 4167
 3.431.6 Sympy [F] 4167
 3.431.7 Maxima [F] 4167
 3.431.8 Giac [F(-1)] 4168
 3.431.9 Mupad [F(-1)] 4168

3.431.1 Optimal result

Integrand size = 35, antiderivative size = 199

$$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

$$= \frac{\sqrt{ia-b}(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} + \frac{\sqrt{ia+b}(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$- \frac{2A\sqrt{a+b \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2(Ab+3aB)\sqrt{a+b \tan(c+dx)}}{3ad\sqrt{\tan(c+dx)}}$$

```
output (I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a
-b)^(1/2)/d+(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)
)^(1/2))*(I*a+b)^(1/2)/d-2/3*(A*b+3*B*a)*(a+b*tan(d*x+c))^(1/2)/a/d/tan(d*
x+c)^(1/2)-2/3*A*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(3/2)
```

3.431.2 Mathematica [A] (verified)

Time = 1.75 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.97

$$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

$$= \frac{-3\sqrt[4]{-1}\sqrt{-a+ib}(iA+B) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) + 3(-1)^{3/4}\sqrt{a+ib}(A+iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{3d}$$

3.431. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

input `Integrate[(Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(-3*(-1)^(1/4)*Sqrt[-a + I*b]*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + 3*(-1)^(3/4)*Sqrt[a + I*b]*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - (2*Sqrt[a + b*Tan[c + d*x]]*(a*A + (A*b + 3*a*B)*Tan[c + d*x]))/(a*Tan[c + d*x]^(3/2)))/(3*d)`

3.431.3 Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.12, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 4091, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx \\ & \quad \downarrow \text{4091} \\ & -\frac{2}{3} \int -\frac{-2Ab \tan^2(c + dx) - 3(aA - bB) \tan(c + dx) + Ab + 3aB}{2 \tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx - \frac{2A \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \\ & \quad \downarrow \text{27} \\ & \frac{1}{3} \int \frac{-2Ab \tan^2(c + dx) - 3(aA - bB) \tan(c + dx) + Ab + 3aB}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx - \frac{2A \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \\ & \quad \downarrow \text{3042} \\ & \frac{1}{3} \int \frac{-2Ab \tan(c + dx)^2 - 3(aA - bB) \tan(c + dx) + Ab + 3aB}{\tan(c + dx)^{3/2} \sqrt{a + b \tan(c + dx)}} dx - \frac{2A \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \\ & \quad \downarrow \text{4132} \end{aligned}$$

3.431. $\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$

$$\begin{aligned}
& \frac{1}{3} \left(-\frac{2 \int \frac{3(a(aA-bB)+a(Ab+aB)\tan(c+dx))}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a} - \frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right) - \\
& \quad \frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow 27 \\
& \frac{1}{3} \left(-\frac{3 \int \frac{a(aA-bB)+a(Ab+aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a} - \frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right) - \\
& \quad \frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{3} \left(-\frac{3 \int \frac{a(aA-bB)+a(Ab+aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a} - \frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right) - \\
& \quad \frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow 4099 \\
& -\frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \\
& \frac{1}{3} \left(-\frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} - \frac{3\left(\frac{1}{2}a(a+ib)(A+iB)\int \frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}a(a-ib)(A-iB)\right)}{a} \right) \\
& \quad \downarrow 3042 \\
& -\frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \\
& \frac{1}{3} \left(-\frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} - \frac{3\left(\frac{1}{2}a(a+ib)(A+iB)\int \frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}a(a-ib)(A-iB)\right)}{a} \right) \\
& \quad \downarrow 4098 \\
& -\frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \\
& \frac{1}{3} \left(-\frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} - \frac{3\left(\frac{a(a-ib)(A-iB)\int \frac{1}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} d\tan(c+dx)}{2d} + \frac{a(a+ib)(A+iB)}{a}\right)}{a} \right)
\end{aligned}$$

3.431. $\int \frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \downarrow 104 \\
 & -\frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \\
 & \frac{1}{3} \left(\frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} - \frac{3 \left(\frac{a(a+ib)(A+iB) \int \frac{1}{\frac{(ia-b)\tan(c+dx)}{a+b\tan(c+dx)}+1} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{a} + \frac{a(a-ib)(A-iB) \int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{a} \right)}{a} \right)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 216 \\
 & -\frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \\
 & \frac{1}{3} \left(\frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} - \frac{3 \left(\frac{a(a-ib)(A-iB) \int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{a} + \frac{a(a+ib)(A+iB) \arctan\left(\frac{\sqrt{-b+ia}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} \right)}{a} \right)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 219 \\
 & -\frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \\
 & \frac{1}{3} \left(\frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} - \frac{3 \left(\frac{a(a+ib)(A+iB) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} + \frac{a(a-ib)(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}} \right)}{a} \right)
 \end{aligned}$$

input `Int[(Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(-2*A*Sqrt[a + b*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) + ((-3*((a*(a + I*b)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d) + (a*(a - I*b)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d)))/a - (2*(A*b + 3*a*B)*Sqrt[a + b*Tan[c + d*x]]/(a*d*Sqrt[Tan[c + d*x]]))/3`

3.431. $\int \frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

3.431.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4091 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegerQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.431.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.83 (sec) , antiderivative size = 2181119, normalized size of antiderivative = 10960.40

output too large to display

input `int((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x)`

output `result too large to display`

$$3.431. \quad \int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.431.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7946 vs. $2(159) = 318$.

Time = 1.45 (sec) , antiderivative size = 7946, normalized size of antiderivative = 39.93

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algo
rithm="fricas")`

output Too large to include

3.431.6 Sympy [F]

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)}}{\tan^{\frac{5}{2}}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))/tan(c + d*x)**(5/2)
, x)`

3.431.7 Maxima [F]

$$\begin{aligned} & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a}}{\tan(dx + c)^{\frac{5}{2}}} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)/tan(d*x + c)^(5/2)
, x)`

3.431. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

3.431.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorith="giac")`

output `Timed out`

3.431.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)}}{\tan(c + dx)^{5/2}} dx \end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2))/tan(c + d*x)^(5/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2))/tan(c + d*x)^(5/2),x)`

3.432
$$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

3.432.1 Optimal result 4169
 3.432.2 Mathematica [A] (verified) 4170
 3.432.3 Rubi [A] (verified) 4170
 3.432.4 Maple [B] (warning: unable to verify) 4176
 3.432.5 Fricas [B] (verification not implemented) 4176
 3.432.6 Sympy [F] 4176
 3.432.7 Maxima [F(-1)] 4177
 3.432.8 Giac [F(-1)] 4177
 3.432.9 Mupad [F(-1)] 4177

3.432.1 Optimal result

Integrand size = 35, antiderivative size = 250

$$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

$$= \frac{\sqrt{ia-b}(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} - \frac{\sqrt{ia+b}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$- \frac{2A\sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2(Ab+5aB)\sqrt{a+b \tan(c+dx)}}{15ad \tan^{\frac{3}{2}}(c+dx)}$$

$$+ \frac{2(15a^2A+2Ab^2-5abB)\sqrt{a+b \tan(c+dx)}}{15a^2d\sqrt{\tan(c+dx)}}$$

output

```
(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a-b)^(1/2)/d-(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a+b)^(1/2)/d+2/15*(15*A*a^2+2*A*b^2-5*B*a*b)*(a+b*tan(d*x+c))^(1/2)/a^2/d/tan(d*x+c)^(1/2)-2/5*A*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(5/2)-2/15*(A*b+5*B*a)*(a+b*tan(d*x+c))^(1/2)/a/d/tan(d*x+c)^(3/2)
```


3.432.2 Mathematica [A] (verified)

Time = 3.43 (sec) , antiderivative size = 226, normalized size of antiderivative = 0.90

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$= \frac{15\sqrt[4]{-1}\sqrt{-a + ib}(A - iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a + ib}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) + 15\sqrt[4]{-1}\sqrt{a + ib}(A + iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{a + ib}}{\sqrt{a + b \tan(c + dx)}}\right)}{15d}$$

input `Integrate[(Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2), x]`

output `(15*(-1)^(1/4)*Sqrt[-a + I*b]*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + 15*(-1)^(1/4)*Sqrt[a + I*b]*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + (2*Sqrt[a + b*Tan[c + d*x]]*(-3*a^2*A - a*(A*b + 5*a*B)*Tan[c + d*x] + (15*a^2*A + 2*A*b^2 - 5*a*b*B)*Tan[c + d*x]^2))/(a^2*Tan[c + d*x]^(5/2)))/(15*d)`

3.432.3 Rubi [A] (verified)

Time = 1.76 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.16, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 4091, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan(c + dx)^{7/2}} dx$$

$$\downarrow \text{4091}$$

$$-\frac{2}{5} \int -\frac{-4Ab \tan^2(c + dx) - 5(aA - bB) \tan(c + dx) + Ab + 5aB}{2 \tan^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx - \frac{2A \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)}$$

3.432. $\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{1}{5} \int \frac{-4Ab \tan^2(c+dx) - 5(aA - bB) \tan(c+dx) + Ab + 5aB}{\tan^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx - \frac{2A \sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \downarrow 3042 \\
& \frac{1}{5} \int \frac{-4Ab \tan(c+dx)^2 - 5(aA - bB) \tan(c+dx) + Ab + 5aB}{\tan(c+dx)^{5/2} \sqrt{a+b \tan(c+dx)}} dx - \frac{2A \sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \downarrow 4132 \\
& \frac{1}{5} \left(\frac{2 \int \frac{15Aa^2 - 5bBa + 15(Ab+aB) \tan(c+dx)a + 2Ab^2 + 2b(Ab+5aB) \tan^2(c+dx)}{2 \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(5aB + Ab) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) - \\
& \quad \frac{2A \sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \downarrow 27 \\
& \frac{1}{5} \left(\frac{\int \frac{15Aa^2 - 5bBa + 15(Ab+aB) \tan(c+dx)a + 2Ab^2 + 2b(Ab+5aB) \tan^2(c+dx)}{\tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(5aB + Ab) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) - \\
& \quad \frac{2A \sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \downarrow 3042 \\
& \frac{1}{5} \left(\frac{\int \frac{15Aa^2 - 5bBa + 15(Ab+aB) \tan(c+dx)a + 2Ab^2 + 2b(Ab+5aB) \tan(c+dx)^2}{\tan(c+dx)^{3/2} \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(5aB + Ab) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) - \\
& \quad \frac{2A \sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \downarrow 4132 \\
& \frac{1}{5} \left(\frac{-\frac{2 \int -\frac{15(a^2(Ab+aB) - a^2(aA-bB) \tan(c+dx))}{2 \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(15a^2A - 5abB + 2Ab^2) \sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}}}{3a} - \frac{2(5aB + Ab) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) - \\
& \quad \frac{2A \sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)}
\end{aligned}$$

3.432. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{1}{5} \left(\frac{15 \int \frac{a^2(Ab+aB)-a^2(aA-bB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a} - \frac{2(15a^2A-5abB+2Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} - \frac{2(5aB+Ab)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} \right) - \\
 & \frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} \\
 & \downarrow 3042 \\
 & \frac{1}{5} \left(\frac{15 \int \frac{a^2(Ab+aB)-a^2(aA-bB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a} - \frac{2(15a^2A-5abB+2Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} - \frac{2(5aB+Ab)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} \right) - \\
 & \frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} \\
 & \downarrow 4099 \\
 & -\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} + \\
 & \frac{1}{5} \left(\frac{2(5aB+Ab)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-5abB+2Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{15\left(\frac{1}{2}a^2(a-ib)(B+iA)\int\frac{i\tan(c+dx)+}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}\right)}{3a} \right) \\
 & \downarrow 3042 \\
 & -\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} + \\
 & \frac{1}{5} \left(\frac{2(5aB+Ab)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-5abB+2Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{15\left(\frac{1}{2}a^2(a-ib)(B+iA)\int\frac{i\tan(c+dx)+}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}\right)}{3a} \right) \\
 & \downarrow 4098
 \end{aligned}$$

3.432. $\int \frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$\frac{1}{5} \left(\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} + \frac{2(5aB+Ab)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-5abB+2Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{15 \left(\frac{a^2(a-ib)(B+iA) \int \frac{1}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}} dx \right)}{2d} \right)$$

↓ 104

$$\frac{1}{5} \left(\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} + \frac{2(5aB+Ab)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-5abB+2Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{15 \left(\frac{a^2(a-ib)(B+iA) \int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} dx}{d} \right)}{3a} \right)$$

↓ 216

$$\frac{1}{5} \left(\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} + \frac{2(5aB+Ab)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-5abB+2Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{15 \left(\frac{a^2(a-ib)(B+iA) \int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} dx}{d} \right)}{3a} \right)$$

↓ 219

$$\frac{1}{5} \left(\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} + \frac{2(5aB+Ab)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-5abB+2Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{15 \left(\frac{a^2(a-ib)(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}} \right)}{3a} \right)$$

3.432. $\int \frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

input `Int[(Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(-2*A*Sqrt[a + b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) + ((-2*(A*b + 5*a*B)*Sqrt[a + b*Tan[c + d*x]])/(3*a*d*Tan[c + d*x]^(3/2)) - ((15*(-((a^2*(a + I*b)*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a - b]*d)) + (a^2*(a - I*b)*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d)))/a - (2*(15*a^2*A + 2*A*b^2 - 5*a*b*B)*Sqrt[a + b*Tan[c + d*x]]/(a*d*Sqrt[Tan[c + d*x]]))/(3*a))/5`

3.432.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 104 `Int[(((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4091 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.432.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.40 (sec) , antiderivative size = 2183172, normalized size of antiderivative = 8732.69

output too large to display

input `int((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x)`

output `result too large to display`

3.432.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7964 vs. 2(204) = 408.

Time = 1.42 (sec) , antiderivative size = 7964, normalized size of antiderivative = 31.86

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algo rithm="fricas")`

output `Too large to include`

3.432.6 Sympy [F]

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)}}{\tan^{\frac{7}{2}}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(7/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))/tan(c + d*x)**(7/2), x)`

3.432. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

3.432.7 Maxima [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algo
rithm="maxima")
```

output Timed out

3.432.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algo
rithm="giac")
```

output Timed out

3.432.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx \\ &= \int \frac{(A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)}}{\tan(c + dx)^{7/2}} dx \end{aligned}$$

```
input int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2))/tan(c + d*x)^(7/2),x
)
```

```
output int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2))/tan(c + d*x)^(7/2),
x)
```

3.432. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

3.433
$$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

3.433.1 Optimal result	4178
3.433.2 Mathematica [A] (verified)	4179
3.433.3 Rubi [A] (verified)	4179
3.433.4 Maple [B] (warning: unable to verify)	4186
3.433.5 Fricas [B] (verification not implemented)	4186
3.433.6 Sympy [F(-1)]	4187
3.433.7 Maxima [F]	4187
3.433.8 Giac [F(-1)]	4187
3.433.9 Mupad [F(-1)]	4188

3.433.1 Optimal result

Integrand size = 35, antiderivative size = 314

$$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

$$= -\frac{\sqrt{ia-b}(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$-\frac{\sqrt{ia+b}(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} - \frac{2A\sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

$$-\frac{2(Ab+7aB)\sqrt{a+b \tan(c+dx)}}{35ad \tan^{\frac{5}{2}}(c+dx)} + \frac{2(35a^2A+4Ab^2-7abB)\sqrt{a+b \tan(c+dx)}}{105a^2d \tan^{\frac{3}{2}}(c+dx)}$$

$$+ \frac{2(35a^2Ab-8Ab^3+105a^3B+14ab^2B)\sqrt{a+b \tan(c+dx)}}{105a^3d\sqrt{\tan(c+dx)}}$$

output

```
-(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a-b)^(1/2)/d-(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a+b)^(1/2)/d+2/105*(35*A*a^2*b-8*A*b^3+105*B*a^3+14*B*a*b^2)*(a+b*tan(d*x+c))^(1/2)/a^3/d/tan(d*x+c)^(1/2)-2/7*A*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(7/2)-2/35*(A*b+7*B*a)*(a+b*tan(d*x+c))^(1/2)/a/d/tan(d*x+c)^(5/2)+2/105*(35*A*a^2+4*A*b^2-7*B*a*b)*(a+b*tan(d*x+c))^(1/2)/a^2/d/tan(d*x+c)^(3/2)
```

3.433.2 Mathematica [A] (verified)

Time = 4.42 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.84

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$= \frac{105\sqrt[4]{-1}\sqrt{-a + ib}(iA + B) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a + ib}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) - 105(-1)^{3/4}\sqrt{a + ib}(A + iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a + ib}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{105d}$$

input `Integrate[(Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2),x]`

output `(105*(-1)^(1/4)*Sqrt[-a + I*b]*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - 105*(-1)^(3/4)*Sqrt[a + I*b]*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + (2*Sqrt[a + b*Tan[c + d*x]]*(-15*a^3*A - 3*a^2*(A*b + 7*a*B)*Tan[c + d*x] + a*(35*a^2*A + 4*A*b^2 - 7*a*b*B)*Tan[c + d*x]^2 + (35*a^2*A*b - 8*A*b^3 + 105*a^3*B + 14*a*b^2*B)*Tan[c + d*x]^3))/(a^3*Tan[c + d*x]^(7/2)))/(105*d)`

3.433.3 Rubi [A] (verified)

Time = 2.27 (sec) , antiderivative size = 355, normalized size of antiderivative = 1.13, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.543$, Rules used = {3042, 4091, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan(c + dx)^{9/2}} dx$$

↓ 4091

3.433. $\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$

$$\begin{aligned}
& -\frac{2}{7} \int \frac{-6Ab \tan^2(c+dx) - 7(aA - bB) \tan(c+dx) + Ab + 7aB}{2 \tan^{\frac{7}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx - \frac{2A \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow 27 \\
& \frac{1}{7} \int \frac{-6Ab \tan^2(c+dx) - 7(aA - bB) \tan(c+dx) + Ab + 7aB}{\tan^{\frac{7}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx - \frac{2A \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{7} \int \frac{-6Ab \tan(c+dx)^2 - 7(aA - bB) \tan(c+dx) + Ab + 7aB}{\tan(c+dx)^{7/2} \sqrt{a+b \tan(c+dx)}} dx - \frac{2A \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow 4132 \\
& \frac{1}{7} \left(-\frac{2 \int \frac{35Aa^2 - 7bBa + 35(Ab+aB) \tan(c+dx)a + 4Ab^2 + 4b(Ab+7aB) \tan^2(c+dx)}{2 \tan^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2(7aB + Ab) \sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} \right) - \\
& \quad \frac{2A \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow 27 \\
& \frac{1}{7} \left(-\frac{\int \frac{35Aa^2 - 7bBa + 35(Ab+aB) \tan(c+dx)a + 4Ab^2 + 4b(Ab+7aB) \tan^2(c+dx)}{\tan^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2(7aB + Ab) \sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} \right) - \\
& \quad \frac{2A \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow 3042 \\
& \frac{1}{7} \left(-\frac{\int \frac{35Aa^2 - 7bBa + 35(Ab+aB) \tan(c+dx)a + 4Ab^2 + 4b(Ab+7aB) \tan(c+dx)^2}{\tan(c+dx)^{5/2} \sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2(7aB + Ab) \sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} \right) - \\
& \quad \frac{2A \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} \\
& \quad \downarrow 4132
\end{aligned}$$

3.433. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

$$\frac{1}{7} \left(\frac{2 \int -\frac{105Ba^3+35Aba^2-105(aA-bB)\tan(c+dx)a^2+14b^2Ba-8Ab^3-2b(35Aa^2-7bBa+4Ab^2)\tan^2(c+dx)}{2 \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(35a^2A-7abB+4Ab^2)\sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) - 5a$$

$$\frac{2A\sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{7} \left(\frac{\int \frac{105Ba^3+35Aba^2-105(aA-bB)\tan(c+dx)a^2+14b^2Ba-8Ab^3-2b(35Aa^2-7bBa+4Ab^2)\tan^2(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(35a^2A-7abB+4Ab^2)\sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) - 5a$$

$$\frac{2A\sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{\int \frac{105Ba^3+35Aba^2-105(aA-bB)\tan(c+dx)a^2+14b^2Ba-8Ab^3-2b(35Aa^2-7bBa+4Ab^2)\tan(c+dx)^2}{\tan(c+dx)^{3/2}\sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(35a^2A-7abB+4Ab^2)\sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) - 5a$$

$$\frac{2A\sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 4132

$$\frac{1}{7} \left(\frac{2 \int \frac{105((aA-bB)a^3+(Ab+aB)\tan(c+dx)a^3)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(105a^3B+35a^2Ab+14ab^2B-8Ab^3)\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}{3a} - \frac{2(35a^2A-7abB+4Ab^2)\sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) - 5a$$

$$\frac{2A\sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 27

3.433. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

$$\frac{1}{7} \left(\frac{-\frac{105 \int \frac{(aA-bB)a^3+(Ab+aB)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a} - \frac{2(105a^3B+35a^2Ab+14ab^2B-8Ab^3)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}{3a} - \frac{2(35a^2A-7abB+4Ab^2)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} \right) - \frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{-\frac{105 \int \frac{(aA-bB)a^3+(Ab+aB)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a} - \frac{2(105a^3B+35a^2Ab+14ab^2B-8Ab^3)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}{3a} - \frac{2(35a^2A-7abB+4Ab^2)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} \right) - \frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)}$$

↓ 4099

$$-\frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} +$$

$$\frac{1}{7} \left(-\frac{2(7aB+Ab)\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-7abB+4Ab^2)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} + \frac{2(105a^3B+35a^2Ab+14ab^2B-8Ab^3)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right) - \frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$-\frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} +$$

$$\frac{1}{7} \left(-\frac{2(7aB+Ab)\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-7abB+4Ab^2)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} + \frac{2(105a^3B+35a^2Ab+14ab^2B-8Ab^3)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right) - \frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)}$$

↓ 4098

3.433. $\int \frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

$$\begin{aligned}
 & -\frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \\
 \frac{1}{7} \left(& -\frac{2(7aB+Ab)\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-7abB+4Ab^2)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} + \frac{2(105a^3B+35a^2Ab+14ab^2B-8Ab^3)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right)
 \end{aligned}$$

↓ 104

$$\begin{aligned}
 & -\frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \\
 \frac{1}{7} \left(& -\frac{2(7aB+Ab)\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-7abB+4Ab^2)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} + \frac{2(105a^3B+35a^2Ab+14ab^2B-8Ab^3)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right)
 \end{aligned}$$

↓ 216

$$\begin{aligned}
 & -\frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \\
 \frac{1}{7} \left(& -\frac{2(7aB+Ab)\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-7abB+4Ab^2)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} + \frac{2(105a^3B+35a^2Ab+14ab^2B-8Ab^3)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right)
 \end{aligned}$$

↓ 219

3.433. $\int \frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

$$-\frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \left(\frac{1}{7} \left[-\frac{2(7aB+Ab)\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-7abB+4Ab^2)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} + \frac{2(105a^3B+35a^2Ab+14ab^2B-8Ab^3)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right] \right)$$

input `Int[(Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2),x]`

output `(-2*A*Sqrt[a + b*Tan[c + d*x]])/(7*d*Tan[c + d*x]^(7/2)) + ((-2*(A*b + 7*a*B)*Sqrt[a + b*Tan[c + d*x]])/(5*a*d*Tan[c + d*x]^(5/2)) - ((-2*(35*a^2*A + 4*A*b^2 - 7*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(3*a*d*Tan[c + d*x]^(3/2)) + ((-105*((a^3*(a + I*b)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d) + (a^3*(a - I*b)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d)))/a - (2*(35*a^2*A*b - 8*A*b^3 + 105*a^3*B + 14*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]]/(a*d*Sqrt[Tan[c + d*x]]))/(3*a))/(5*a))/7`

3.433.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && IntegerQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[(((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4091 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`
- rule 4098 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`
- rule 4099 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`


```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

3.433.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.62 (sec) , antiderivative size = 2185304, normalized size of antiderivative = 6959.57

output too large to display

```
input int((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x)
```

```
output result too large to display
```

3.433.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8039 vs. $2(259) = 518$.

Time = 1.42 (sec) , antiderivative size = 8039, normalized size of antiderivative = 25.60

$$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algo
rithm="fricas")
```

```
output Too large to include
```

3.433. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

3.433.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(9/2),x)`

output `Timed out`

3.433.7 Maxima [F]

$$\begin{aligned} & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx \\ &= \int \frac{(B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a}}{\tan(dx + c)^{\frac{9}{2}}} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)/tan(d*x + c)^(9/2), x)`

3.433.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algorithm="giac")`

output `Timed out`

3.433. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

3.433.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)}}{\tan(c + dx)^{9/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2))/tan(c + d*x)^(9/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2))/tan(c + d*x)^(9/2),x)`

3.434 $\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.434.1 Optimal result	4189
3.434.2 Mathematica [A] (verified)	4190
3.434.3 Rubi [A] (verified)	4191
3.434.4 Maple [B] (warning: unable to verify)	4195
3.434.5 Fricas [B] (verification not implemented)	4196
3.434.6 Sympy [F(-1)]	4196
3.434.7 Maxima [F]	4196
3.434.8 Giac [F(-1)]	4197
3.434.9 Mupad [F(-1)]	4197

3.434.1 Optimal result

Integrand size = 35, antiderivative size = 323

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{(ia - b)^{3/2}(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) + \frac{(6a^2Ab - 16Ab^3 - a^3B - 24ab^2B) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{8b^{3/2}d} + \frac{(ia + b)^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} + \frac{(6aAb - a^2B - 8b^2B) \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}{8bd} + \frac{(6Ab - aB) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}}{12bd} + \frac{B \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}}{3bd}$$

output $(I*a-b)^{(3/2)}*(A+I*B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})/d+1/8*(6*A*a^2*b-16*A*b^3-B*a^3-24*B*a*b^2)*\operatorname{arctanh}(b^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})/b^{(3/2)}/d+(I*a+b)^{(3/2)}*(A-I*B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})/d+1/8*(6*A*a*b-B*a^2-8*B*b^2)*\tan(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(1/2)}/b/d+1/12*(6*A*b-B*a)*\tan(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(3/2)}/b/d+1/3*B*\tan(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(5/2)}/b/d$

3.434.2 Mathematica [A] (verified)

Time = 4.65 (sec) , antiderivative size = 347, normalized size of antiderivative = 1.07

$$\int \tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx = \frac{24\sqrt[4]{-1}(-a+ib)^{3/2}b(iA+B)\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)+24(-1)^{3/4}(a+ib)^{3/2}}{\dots}$$

input `Integrate[Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]), x]`

output $(24*(-1)^{(1/4)}*(-a + I*b)^{(3/2)}*b*(I*A + B)*\operatorname{ArcTan}(((-1)^{(1/4)}*\sqrt{-a + I*b})*\sqrt{\tan[c + d*x]})/\sqrt{a + b*\tan[c + d*x]}) + 24*(-1)^{(3/4)}*(a + I*b)^{(3/2)}*b*(A + I*B)*\operatorname{ArcTan}(((-1)^{(1/4)}*\sqrt{a + I*b})*\sqrt{\tan[c + d*x]})/\sqrt{a + b*\tan[c + d*x]}) - 3*(-6*a*A*b + a^2*B + 8*b^2*B)*\sqrt{\tan[c + d*x]})*\sqrt{a + b*\tan[c + d*x]}) + 2*(6*A*b - a*B)*\sqrt{\tan[c + d*x]})*(a + b*\tan[c + d*x])^{(3/2)} + 8*B*\sqrt{\tan[c + d*x]})*(a + b*\tan[c + d*x])^{(5/2)} - (3*\sqrt{a}*(-6*a^2*A*b + 16*A*b^3 + a^3*B + 24*a*b^2*B)*\operatorname{ArcSinh}((\sqrt{b})*\sqrt{\tan[c + d*x]})/\sqrt{a})*\sqrt{1 + (b*\tan[c + d*x])/a})/(\sqrt{b})*\sqrt{a + b*\tan[c + d*x]})/(24*b*d)$

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd} - \frac{\frac{3}{4} \int \frac{\sqrt{a+b\tan(c+dx)}(-((-Ba^2+6Aba-8b^2B)\tan^2(c+dx))+8b(Ab+aB)\tan(c+dx)+a(2Ab+aB))}{\sqrt{\tan(c+dx)}} dx - \frac{(6Ab-aB)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{2d}}{6b}$$

↓ 3042

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd} - \frac{\frac{3}{4} \int \frac{\sqrt{a+b\tan(c+dx)}(-((-Ba^2+6Aba-8b^2B)\tan(c+dx)^2)+8b(Ab+aB)\tan(c+dx)+a(2Ab+aB))}{\sqrt{\tan(c+dx)}} dx - \frac{(6Ab-aB)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{2d}}{6b}$$

↓ 4130

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd} - \frac{\frac{3}{4} \left(\int \frac{-((-Ba^3+6Aba^2-24b^2Ba-16Ab^3)\tan^2(c+dx))+16b(Ba^2+2Aba-b^2B)\tan(c+dx)+a(Ba^2+10Aba-8b^2B)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{(a^2(-B)+6aAb-8b^2B)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{2d} \right)}{6b}$$

↓ 27

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd} - \frac{\frac{3}{4} \left(\frac{1}{2} \int \frac{-((-Ba^3+6Aba^2-24b^2Ba-16Ab^3)\tan^2(c+dx))+16b(Ba^2+2Aba-b^2B)\tan(c+dx)+a(Ba^2+10Aba-8b^2B)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{(a^2(-B)+6aAb-8b^2B)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{2d} \right)}{6b}$$

↓ 3042

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd} - \frac{\frac{3}{4} \left(\frac{1}{2} \int \frac{-((-Ba^3+6Aba^2-24b^2Ba-16Ab^3)\tan(c+dx)^2)+16b(Ba^2+2Aba-b^2B)\tan(c+dx)+a(Ba^2+10Aba-8b^2B)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{(a^2(-B)+6aAb-8b^2B)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{2d} \right)}{6b}$$

↓ 4138

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd} - \frac{\frac{3}{4} \left(\frac{\int \frac{-((-Ba^3+6Aba^2-24b^2Ba-16Ab^3)\tan^2(c+dx))+16b(Ba^2+2Aba-b^2B)\tan(c+dx)+a(Ba^2+10Aba-8b^2B)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} d\tan(c+dx)}{2d} - \frac{(a^2(-B)+6aAb-8b^2B)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{2d} \right)}{6b}$$

↓ 2035

3.434. $\int \tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx$

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd} - \frac{\int \frac{-((-Ba^3+6Aba^2-24b^2Ba-16Ab^3)\tan^2(c+dx)+16b(Ba^2+2Aba-b^2B)\tan(c+dx)+a(Ba^2+10Aba-8b^2B))d\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}}{d} - \frac{(a^2(-B)+6aAb-8b^2)}{d}$$

6b

2257

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd} - \frac{\int \left(\frac{Ba^3-6Aba^2+24b^2Ba+16Ab^3}{\sqrt{a+b\tan(c+dx)}} + \frac{16(b(Aa^2-2bBa-Ab^2)+b(Ba^2+2Aba-b^2B)\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} \right) d\sqrt{\tan(c+dx)}}{d} - \frac{(a^2(-B)+6aAb-8b^2B)\sqrt{\tan(c+dx)}}{d}$$

6b

2009

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd} - \frac{(6Ab-aB)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d} + \frac{3}{4} \left(-\frac{(a^2(-B)+6aAb-8b^2B)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} + \frac{(a^3(-B)+6a^2Ab-24ab^2B-16b^3)}{d} \right)$$

6b

input `Int[Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(5/2))/(3*b*d) - (-1/2*((6*A*b - a*B)*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2))/d + (3*((-8*(I*a - b)^(3/2)*b*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - ((6*a^2*A*b - 16*A*b^3 - a^3*B - 24*a*b^2*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/Sqrt[b] - 8*b*(I*a + b)^(3/2)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/d - ((6*a*A*b - a^2*B - 8*b^2*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d)/4)/(6*b)`

3.434.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2035 `Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[Fx, x]`
- rule 2257 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4130 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
) + (f_.)*(x_)^2), x_Symbol] :> Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && (!IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

3.434.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.90 (sec) , antiderivative size = 2403184, normalized size of antiderivative = 7440.20

output too large to display

```
input int(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x)
```

```
output result too large to display
```

3.434.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12212 vs. $2(264) = 528$.

Time = 5.20 (sec) , antiderivative size = 24426, normalized size of antiderivative = 75.62

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output Too large to include

3.434.6 Sympy [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(3/2)*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output Timed out

3.434.7 Maxima [F]

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{3}{2}} \tan(dx + c)^{\frac{3}{2}} dx$$

input `integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)*tan(d*x + c)^(3/2), x)`

3.434. $\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.434.8 Giac [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.434.9 Mupad [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int \tan(c + dx)^{3/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2} dx$$

input `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2),x)`

output `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2), x)`

3.435 $\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.435.1 Optimal result	4198
3.435.2 Mathematica [A] (verified)	4199
3.435.3 Rubi [A] (verified)	4199
3.435.4 Maple [B] (warning: unable to verify)	4203
3.435.5 Fricas [B] (verification not implemented)	4204
3.435.6 Sympy [F]	4204
3.435.7 Maxima [F]	4204
3.435.8 Giac [F(-1)]	4205
3.435.9 Mupad [F(-1)]	4205

3.435.1 Optimal result

Integrand size = 35, antiderivative size = 268

$$\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{(a + ib)^2(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia - bd}}$$

$$+ \frac{(12aAb + 3a^2B - 8b^2B) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{4\sqrt{bd}}$$

$$+ \frac{(ia + b)^{3/2}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$+ \frac{(4Ab + 5aB)\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}}{4d} + \frac{bB \tan^{3/2}(c + dx)\sqrt{a + b \tan(c + dx)}}{2d}$$

output

```
(I*a+b)^(3/2)*(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d+(a+I*b)^2*(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d/(I*a-b)^(1/2)+1/4*(12*A*a*b+3*B*a^2-8*B*b^2)*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d/b^(1/2)+1/4*(4*A*b+5*B*a)*tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/d+1/2*b*B*(a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^(3/2)/d
```

3.435.2 Mathematica [A] (verified)

Time = 2.70 (sec) , antiderivative size = 290, normalized size of antiderivative = 1.08

$$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{-4\sqrt[4]{-1}(-a+ib)^{3/2}(A-iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) + 4\sqrt[4]{-1}(a+ib)^{3/2}(A+B \tan(c+dx))}{1}$$

input `Integrate[Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(-4*(-1)^(1/4)*(-a + I*b)^(3/2)*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + 4*(-1)^(1/4)*(a + I*b)^(3/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + (4*A*b + 5*a*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]] + 2*b*B*Tan[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]] + (Sqrt[a]*(12*a*A*b + 3*a^2*B - 8*b^2*B)*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/(Sqrt[b]*Sqrt[a + b*Tan[c + d*x]])/(4*d)`

3.435.3 Rubi [A] (verified)

Time = 1.55 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.98, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 4090, 27, 3042, 4130, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$$

↓ 3042

$$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$$

↓ 4090

$$\frac{1}{2} \int \frac{\sqrt{\tan(c+dx)}(b(4Ab+5aB)\tan^2(c+dx)+4(Ba^2+2Aba-b^2B)\tan(c+dx)+a(4aA-3bB))}{2\sqrt{a+b\tan(c+dx)}} dx + \frac{bB \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}{2d} \downarrow 27$$

$$\frac{1}{4} \int \frac{\sqrt{\tan(c+dx)}(b(4Ab+5aB)\tan^2(c+dx)+4(Ba^2+2Aba-b^2B)\tan(c+dx)+a(4aA-3bB))}{\sqrt{a+b\tan(c+dx)}} dx + \frac{bB \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}{2d} \downarrow 3042$$

$$\frac{1}{4} \int \frac{\sqrt{\tan(c+dx)}(b(4Ab+5aB)\tan(c+dx)^2+4(Ba^2+2Aba-b^2B)\tan(c+dx)+a(4aA-3bB))}{\sqrt{a+b\tan(c+dx)}} dx + \frac{bB \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}{2d} \downarrow 4130$$

$$\frac{1}{4} \left(\frac{\int -\frac{b(3Ba^2+12Aba-8b^2B)\tan^2(c+dx)-8b(Aa^2-2bBa-Ab^2)\tan(c+dx)+ab(4Ab+5aB)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{b} + \frac{(5aB+4Ab)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} \right) + \frac{bB \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}{2d} \downarrow 27$$

$$\frac{1}{4} \left(\frac{(5aB+4Ab)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{\int -\frac{b(3Ba^2+12Aba-8b^2B)\tan^2(c+dx)-8b(Aa^2-2bBa-Ab^2)\tan(c+dx)+ab(4Ab+5aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}}{2b} \right) + \frac{bB \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}{2d} \downarrow 3042$$

$$\frac{1}{4} \left(\frac{(5aB+4Ab)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{\int -\frac{b(3Ba^2+12Aba-8b^2B)\tan(c+dx)^2-8b(Aa^2-2bBa-Ab^2)\tan(c+dx)+ab(4Ab+5aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}}{2b} \right) + \frac{bB \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}{2d}$$

3.435. $\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx$

$$\begin{aligned} & \downarrow 4138 \\ & \frac{1}{4} \left(\frac{(5aB + 4Ab)\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}}{d} - \frac{\int \frac{-b(3Ba^2 + 12Aba - 8b^2B)\tan^2(c + dx) - 8b(Aa^2 - 2bBa - Ab^2)\tan(c + dx) + bB \tan^{\frac{3}{2}}(c + dx)\sqrt{a + b \tan(c + dx)}}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}(\tan^2(c + dx) + 1)} dx}{2bd} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 2035 \\ & \frac{1}{4} \left(\frac{(5aB + 4Ab)\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}}{d} - \frac{\int \frac{-b(3Ba^2 + 12Aba - 8b^2B)\tan^2(c + dx) - 8b(Aa^2 - 2bBa - Ab^2)\tan(c + dx) + bB \tan^{\frac{3}{2}}(c + dx)\sqrt{a + b \tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}(\tan^2(c + dx) + 1)} dx}{bd} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 2257 \\ & \frac{1}{4} \left(\frac{(5aB + 4Ab)\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}}{d} - \frac{\int \left(\frac{8(b(Ba^2 + 2Aba - b^2B) - b(Aa^2 - 2bBa - Ab^2)\tan(c + dx))}{\sqrt{a + b \tan(c + dx)}(\tan^2(c + dx) + 1)} - \frac{b(3Ba^2 + 12Aba - 8b^2B)\tan^{\frac{3}{2}}(c + dx)\sqrt{a + b \tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}} \right) dx}{bd} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 2009 \\ & \frac{1}{4} \left(\frac{(5aB + 4Ab)\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}}{d} - \frac{bB \tan^{\frac{3}{2}}(c + dx)\sqrt{a + b \tan(c + dx)}}{2d} - \frac{-\sqrt{b}(3a^2B + 12aAb - 8b^2B) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) + 4b \operatorname{arctanh}\left(\frac{\sqrt{a + b \tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{bd} \right) \end{aligned}$$

input `Int[Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(b*B*Tan[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]])/(2*d) + (-((4*(I*a - b)^(3/2)*b*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - Sqrt[b]*(12*a*A*b + 3*a^2*B - 8*b^2*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - 4*b*(I*a + b)^(3/2)*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(b*d)) + ((4*A*b + 5*a*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d)/4`

3.435.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2035 `Int[(F_x_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[Fx, x]`
- rule 2257 `Int[(P_x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4090 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4130 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
) + (f_.)*(x_)^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && (!GtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

3.435.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.90 (sec) , antiderivative size = 2400957, normalized size of antiderivative = 8958.79

output too large to display

```
input int(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x)
```

```
output result too large to display
```

3.435.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12134 vs. $2(216) = 432$.

Time = 4.74 (sec) , antiderivative size = 24274, normalized size of antiderivative = 90.57

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorith="fricas")`

output Too large to include

3.435.6 Sympy [F]

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx = \int (A + B\tan(c+dx))(a+b\tan(c+dx))^{\frac{3}{2}}\sqrt{\tan(c+dx)}dx$$

input `integrate(tan(d*x+c)**(1/2)*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)*sqrt(tan(c + d*x)), x)`

3.435.7 Maxima [F]

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}(A + B\tan(c+dx))dx = \int (B\tan(dx+c) + A)(b\tan(dx+c) + a)^{\frac{3}{2}}\sqrt{\tan(dx+c)}dx$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorith="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)*sqrt(tan(d*x + c)), x)`

3.435.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.435.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx = \int \sqrt{\tan(c+dx)}(A+B\tan(c+dx))(a+b\tan(c+dx))^{3/2} dx$$

input `int(tan(c+d*x)^(1/2)*(A+B*tan(c+d*x))*(a+b*tan(c+d*x))^(3/2),x)`

output `int(tan(c+d*x)^(1/2)*(A+B*tan(c+d*x))*(a+b*tan(c+d*x))^(3/2),x)`

3.436
$$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.436.1 Optimal result 4206
 3.436.2 Mathematica [A] (verified) 4207
 3.436.3 Rubi [A] (verified) 4207
 3.436.4 Maple [B] (warning: unable to verify) 4210
 3.436.5 Fricas [B] (verification not implemented) 4210
 3.436.6 Sympy [F] 4211
 3.436.7 Maxima [F] 4211
 3.436.8 Giac [F(-1)] 4211
 3.436.9 Mupad [F(-1)] 4212

3.436.1 Optimal result

Integrand size = 35, antiderivative size = 204

$$\int \frac{(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx =$$

$$\frac{(ia - b)^{3/2}(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$+ \frac{\sqrt{b}(2Ab + 3aB) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$- \frac{(ia + b)^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$+ \frac{bB \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}{d}$$

output

```
-(I*a-b)^(3/2)*(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d-(I*a+b)^(3/2)*(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d+(2*A*b+3*B*a)*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*b^(1/2)/d+b*B*tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/d
```

3.436.2 Mathematica [A] (verified)

Time = 1.11 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.19

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \frac{-\sqrt[4]{-1}(-a + ib)^{3/2}(iA + B) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{\tan(c + dx)}}$$

input `Integrate[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `(-((-1)^(1/4)*(-a + I*b)^(3/2)*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]) - (-1)^(3/4)*(a + I*b)^(3/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]] + b*B*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]] + (Sqrt[a]*Sqrt[b]*(2*A*b + 3*a*B)*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/Sqrt[a + b*Tan[c + d*x]]/d`

3.436.3 Rubi [A] (verified)Time = 1.10 (sec) , antiderivative size = 200, normalized size of antiderivative = 0.98, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 4090, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\ & \quad \downarrow \text{4090} \\ & \int \frac{b(2Ab + 3aB) \tan^2(c + dx) + 2(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(2aA - bB)}{2\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)} + bB\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx + \end{aligned}$$

3.436. $\int \frac{(a+b \tan(c+dx))^{3/2} (A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{1}{2} \int \frac{b(2Ab + 3aB) \tan^2(c + dx) + 2(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(2aA - bB)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx + \\
 & \quad \frac{bB \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}{d} \\
 & \downarrow 3042 \\
 & \frac{1}{2} \int \frac{b(2Ab + 3aB) \tan(c + dx)^2 + 2(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(2aA - bB)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx + \\
 & \quad \frac{bB \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}{d} \\
 & \downarrow 4138 \\
 & \int \frac{b(2Ab + 3aB) \tan^2(c + dx) + 2(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(2aA - bB)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)} (\tan^2(c + dx) + 1)} d \tan(c + dx) + \\
 & \quad \frac{2d}{bB \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} \\
 & \downarrow 2035 \\
 & \int \frac{b(2Ab + 3aB) \tan^2(c + dx) + 2(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(2aA - bB)}{\sqrt{a + b \tan(c + dx)} (\tan^2(c + dx) + 1)} d \sqrt{\tan(c + dx)} + \\
 & \quad \frac{d}{bB \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} \\
 & \downarrow 2257 \\
 & \int \left(\frac{b(2Ab + 3aB)}{\sqrt{a + b \tan(c + dx)}} + \frac{2(Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx))}{\sqrt{a + b \tan(c + dx)} (\tan^2(c + dx) + 1)} \right) d \sqrt{\tan(c + dx)} + \\
 & \quad \frac{d}{bB \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} \\
 & \downarrow 2009 \\
 & \frac{bB \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}{d} + \\
 & \frac{-((-b + ia)^{3/2} (A + iB) \arctan\left(\frac{\sqrt{-b + ia} \sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) + \sqrt{b} (3aB + 2Ab) \operatorname{arctanh}\left(\frac{\sqrt{b} \sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) - (b + ia)^{3/2} (A + iB) \arctan\left(\frac{\sqrt{-b + ia} \sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right))}{d}
 \end{aligned}$$

input `Int[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

3.436. $\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$

```
output (-((I*a - b)^(3/2)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) + Sqrt[b]*(2*A*b + 3*a*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) - (I*a + b)^(3/2)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/d + (b*B*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d
```

3.436.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2035 Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[Fx, x]
```

```
rule 2257 Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4090 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```



```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

3.436.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.95 (sec) , antiderivative size = 2396039, normalized size of antiderivative = 11745.29

output too large to display

```
input int((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x)
```

```
output result too large to display
```

3.436.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12109 vs. $2(164) = 328$.

Time = 4.77 (sec) , antiderivative size = 24220, normalized size of antiderivative = 118.73

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algo
rithm="fracas")
```

```
output Too large to include
```

3.436.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2}}{\sqrt{\tan(c + dx)}} dx$$

input `integrate((a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)/sqrt(tan(c + d*x)), x)`

3.436.7 Maxima [F]

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^{3/2}}{\sqrt{\tan(dx + c)}} dx$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorith="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)/sqrt(tan(d*x + c)), x)`

3.436.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorith="giac")`

output `Timed out`

3.436.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2}}{\sqrt{\tan(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/tan(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/tan(c + d*x)^(1/2),x)`

3.437
$$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.437.1 Optimal result 4213
 3.437.2 Mathematica [A] (verified) 4214
 3.437.3 Rubi [A] (verified) 4214
 3.437.4 Maple [B] (warning: unable to verify) 4217
 3.437.5 Fricas [B] (verification not implemented) 4217
 3.437.6 Sympy [F] 4218
 3.437.7 Maxima [F(-1)] 4218
 3.437.8 Giac [F(-1)] 4218
 3.437.9 Mupad [F(-1)] 4219

3.437.1 Optimal result

Integrand size = 35, antiderivative size = 209

$$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx =$$

$$-\frac{(a+ib)^2(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia-bd}} + \frac{2b^{3/2}B \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$-\frac{(ia+b)^{3/2}(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} - \frac{2aA\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}$$

```
output 2*b^(3/2)*B*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d-(I*
a+b)^(3/2)*(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))
^(1/2))/d-(a+I*b)^2*(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan
(d*x+c))^(1/2))/d/(I*a-b)^(1/2)-2*a*A*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(
1/2)
```

3.437.2 Mathematica [A] (verified)

Time = 4.46 (sec) , antiderivative size = 416, normalized size of antiderivative = 1.99

$$\int \frac{(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \frac{\sqrt[4]{-1}(-a + ib)^{3/2}(A - iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\tan^{3/2}(c + dx)}$$

input `Integrate[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `((-1)^(1/4)*(-a + I*b)^(3/2)*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]] - (-1)^(1/4)*a*A*Sqrt[a + I*b]*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]] - (-1)^(3/4)*A*Sqrt[a + I*b]*b*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]] - (-1)^(3/4)*a*Sqrt[a + I*b]*B*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]] + (-1)^(1/4)*Sqrt[a + I*b]*b*B*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]] - (2*a*A*Sqrt[a + b*Tan[c + d*x]])/Sqrt[Tan[c + d*x]] + (2*Sqrt[a]*b^(3/2)*B*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/Sqrt[a + b*Tan[c + d*x]]]/d`

3.437.3 Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 200, normalized size of antiderivative = 0.96, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 4088, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx$$

↓ 4088

3.437. $\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{3/2}(c+dx)} dx$

$$\begin{aligned}
& 2 \int \frac{b^2 B \tan^2(c+dx) - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2Ab + aB)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \\
& \quad \frac{2aA\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 27 \\
& \int \frac{b^2 B \tan^2(c+dx) - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2Ab + aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \\
& \quad \frac{2aA\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 3042 \\
& \int \frac{b^2 B \tan(c+dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2Ab + aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \\
& \quad \frac{2aA\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 4138 \\
& \frac{\int \frac{b^2 B \tan^2(c+dx) - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2Ab + aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} d\tan(c+dx)}{d} - \frac{2aA\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 2035 \\
& \frac{2 \int \frac{b^2 B \tan^2(c+dx) - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2Ab + aB)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} d\sqrt{\tan(c+dx)}}{d} - \frac{2aA\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 2257 \\
& \frac{2 \int \left(\frac{Bb^2}{\sqrt{a+b\tan(c+dx)}} + \frac{Ba^2 + 2Aba - b^2B - (Aa^2 - 2bBa - Ab^2) \tan(c+dx)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} \right) d\sqrt{\tan(c+dx)}}{d} - \\
& \quad \frac{2aA\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow 2009 \\
& - \frac{2aA\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \\
& \frac{2 \left(\frac{1}{2}(-b+ia)^{3/2}(-B+ia) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) - \frac{1}{2}(b+ia)^{3/2}(B+ia) \operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) + b^{3/2}E \right)}{d}
\end{aligned}$$

input `Int[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

$$3.437. \quad \int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\tan^{3/2}(c+dx)} dx$$

```
output (2*(((I*a - b)^(3/2)*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/S
qrt[a + b*Tan[c + d*x]]])/2 + b^(3/2)*B*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]
])/Sqrt[a + b*Tan[c + d*x]]] - ((I*a + b)^(3/2)*(I*A + B)*ArcTanh[(Sqrt[I*
a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/2))/d - (2*a*A*Sqrt[
a + b*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]])
```

3.437.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2035 Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]
```

```
rule 2257 Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol
] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a
, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4088 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x
])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2))
Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*
(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n
+ 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[
e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n
+ 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] &
& LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])
```

3.437.
$$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{3/2}(c+dx)} dx$$

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

3.437.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.91 (sec) , antiderivative size = 2394883, normalized size of antiderivative = 11458.77

output too large to display

```
input int((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x)
```

```
output result too large to display
```

3.437.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12148 vs. $2(167) = 334$.

Time = 4.52 (sec) , antiderivative size = 24297, normalized size of antiderivative = 116.25

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algo
rithm="fracas")
```

```
output Too large to include
```

3.437. $\int \frac{(a+b \tan(c+dx))^{3/2} (A+B \tan(c+dx))}{\tan^{3/2}(c+dx)} dx$

3.437.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2}}{\tan^{3/2}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)/tan(c + d*x)**(3/2), x)`

3.437.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algo
rithm="maxima")`

output `Timed out`

3.437.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algo
rithm="giac")`

output `Timed out`

3.437.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2}}{\tan(c + dx)^{3/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/tan(c + d*x)^(3/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/tan(c + d*x)^(3/2),x)`

3.438
$$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.438.1 Optimal result 4220
 3.438.2 Mathematica [A] (verified) 4221
 3.438.3 Rubi [A] (verified) 4221
 3.438.4 Maple [B] (warning: unable to verify) 4226
 3.438.5 Fricas [B] (verification not implemented) 4226
 3.438.6 Sympy [F] 4227
 3.438.7 Maxima [F] 4227
 3.438.8 Giac [F(-1)] 4227
 3.438.9 Mupad [F(-1)] 4228

3.438.1 Optimal result

Integrand size = 35, antiderivative size = 196

$$\int \frac{(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx = \frac{(ia - b)^{3/2}(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} + \frac{(ia + b)^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} - \frac{2aA\sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2(4Ab + 3aB)\sqrt{a + b \tan(c + dx)}}{3d\sqrt{\tan(c + dx)}}$$

output $(I*a-b)^{(3/2)}*(A+I*B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})/d+(I*a+b)^{(3/2)}*(A-I*B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})/d-2/3*(4*A*b+3*B*a)*(a+b*\tan(d*x+c))^{(1/2)}/d/\tan(d*x+c)^{(1/2)}-2/3*a*A*(a+b*\tan(d*x+c))^{(1/2)}/d/\tan(d*x+c)^{(3/2)}$

3.438.2 Mathematica [A] (verified)

Time = 1.00 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.21

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \frac{3\sqrt[4]{-1} \left((-a + ib)^{3/2} (iA + B) \arctan \left(\frac{\sqrt[4]{-1} \sqrt{-a + ib} \sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}} \right) \right)}{\tan^{5/2}(c + dx)}$$

input `Integrate[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(3*(-1)^(1/4)*((-a + I*b)^(3/2)*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) + I*(a + I*b)^(3/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(3/2) - 3*b*B*Sqrt[a + b*Tan[c + d*x]] + (-2*a*A + 3*b*B)*Sqrt[a + b*Tan[c + d*x]] - 2*(4*A*b + 3*a*B)*Tan[c + d*x]*Sqrt[a + b*Tan[c + d*x]]/(3*d*Tan[c + d*x]^(3/2))`

3.438.3 Rubi [A] (verified)Time = 1.34 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.15, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 4088, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx$$

$$\downarrow \text{4088}$$

$$\frac{2}{3} \int \frac{-b(2aA - 3bB) \tan^2(c + dx) - 3(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(4Ab + 3aB)}{2 \tan^{3/2}(c + dx) \sqrt{a + b \tan(c + dx)}} dx - \frac{2aA \sqrt{a + b \tan(c + dx)}}{3d \tan^{3/2}(c + dx)}$$

3.438. $\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{1}{3} \int \frac{-b(2aA - 3bB) \tan^2(c + dx) - 3(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(4Ab + 3aB)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx - \\
& \quad \frac{2aA \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \\
& \downarrow 3042 \\
& \frac{1}{3} \int \frac{-b(2aA - 3bB) \tan(c + dx)^2 - 3(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(4Ab + 3aB)}{\tan(c + dx)^{\frac{3}{2}} \sqrt{a + b \tan(c + dx)}} dx - \\
& \quad \frac{2aA \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \\
& \downarrow 4132 \\
& \frac{1}{3} \left(- \frac{2 \int \frac{3(a(Aa^2 - 2bBa - Ab^2) + a(Ba^2 + 2Aba - b^2B) \tan(c + dx))}{2\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx}{a} - \frac{2(3aB + 4Ab) \sqrt{a + b \tan(c + dx)}}{d \sqrt{\tan(c + dx)}} \right) - \\
& \quad \frac{2aA \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \\
& \downarrow 27 \\
& \frac{1}{3} \left(- \frac{3 \int \frac{a(Aa^2 - 2bBa - Ab^2) + a(Ba^2 + 2Aba - b^2B) \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx}{a} - \frac{2(3aB + 4Ab) \sqrt{a + b \tan(c + dx)}}{d \sqrt{\tan(c + dx)}} \right) - \\
& \quad \frac{2aA \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \\
& \downarrow 3042 \\
& \frac{1}{3} \left(- \frac{3 \int \frac{a(Aa^2 - 2bBa - Ab^2) + a(Ba^2 + 2Aba - b^2B) \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx}{a} - \frac{2(3aB + 4Ab) \sqrt{a + b \tan(c + dx)}}{d \sqrt{\tan(c + dx)}} \right) - \\
& \quad \frac{2aA \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \\
& \downarrow 4099 \\
& - \frac{2aA \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} + \\
& \frac{1}{3} \left(- \frac{2(3aB + 4Ab) \sqrt{a + b \tan(c + dx)}}{d \sqrt{\tan(c + dx)}} - \frac{3 \left(\frac{1}{2} a(a - ib)^2 (A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2} a(a + ib)^2 (A + iB) \int \frac{i \tan(c + dx) - 1}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \right)}{a} \right)
\end{aligned}$$

3.438. $\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$

$$\begin{aligned} & \downarrow 3042 \\ & -\frac{2aA\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \\ \frac{1}{3} & \left(-\frac{2(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{3\left(\frac{1}{2}a(a-ib)^2(A-ib)\int\frac{i\tan(c+dx)+1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx + \frac{1}{2}a(a+ib)^2(A-ib)\int\frac{1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx\right)}{a} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 4098 \\ & -\frac{2aA\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \\ \frac{1}{3} & \left(-\frac{2(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{3\left(\frac{a(a-ib)^2(A-ib)\int\frac{1}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}d\tan(c+dx)}{2d} + \frac{a(a+ib)^2(A+ib)\int\frac{1}{(1+i\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}d\tan(c+dx)}{2d}\right)}{a} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 104 \\ & -\frac{2aA\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \\ \frac{1}{3} & \left(-\frac{2(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{3\left(\frac{a(a-ib)^2(A-ib)\int\frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} + \frac{a(a+ib)^2(A+ib)\int\frac{1}{1+\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d}\right)}{a} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 216 \\ & -\frac{2aA\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \\ \frac{1}{3} & \left(-\frac{2(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{3\left(\frac{a(a-ib)^2(A-ib)\int\frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} + \frac{a(a+ib)^2(A+ib)\arctan\left(\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+\sqrt{a+b\tan(c+dx)}}}\right)}{a} \right) \end{aligned}$$

$$\downarrow 219$$

3.438. $\int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

$$-\frac{2aA\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \frac{1}{3} \left(\frac{2(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{3 \left(\frac{a(a+ib)^2(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} + \frac{a(a-ib)^2(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{b+ia}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}} \right)}{a} \right)$$

input `Int[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2), x]`

output `(-2*a*A*Sqrt[a + b*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) + ((-3*((a + I*b)^2*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d) + (a*(a - I*b)^2*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d))/a - (2*(4*A*b + 3*a*B)*Sqrt[a + b*Tan[c + d*x]]/(d*Sqrt[Tan[c + d*x]]))/3`

3.438.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 104 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q], x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

3.438. $\int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`


```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

3.438.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.91 (sec) , antiderivative size = 2398858, normalized size of antiderivative = 12239.07

output too large to display

```
input int((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x)
```

```
output result too large to display
```

3.438.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12024 vs. 2(156) = 312.

Time = 2.46 (sec) , antiderivative size = 12024, normalized size of antiderivative = 61.35

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algo
rithm="fracas")
```

```
output Too large to include
```

3.438. $\int \frac{(a+b \tan(c+dx))^{3/2} (A+B \tan(c+dx))}{\tan^{5/2}(c+dx)} dx$

3.438.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2}}{\tan^{5/2}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)/tan(c + d*x)**(5/2), x)`

3.438.7 Maxima [F]

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^{3/2}}{\tan(dx + c)^{5/2}} dx$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algo rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)/tan(d*x + c)^(5/2), x)`

3.438.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algo rithm="giac")`

output `Timed out`

3.438.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2}}{\tan(c + dx)^{5/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/tan(c + d*x)^(5/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/tan(c + d*x)^(5/2),x)`

$$3.439 \quad \int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{7/2}(c+dx)} dx$$

3.439.1 Optimal result 4229
 3.439.2 Mathematica [A] (verified) 4230
 3.439.3 Rubi [A] (verified) 4230
 3.439.4 Maple [B] (warning: unable to verify) 4236
 3.439.5 Fricas [B] (verification not implemented) 4236
 3.439.6 Sympy [F] 4237
 3.439.7 Maxima [F(-1)] 4237
 3.439.8 Giac [F(-1)] 4237
 3.439.9 Mupad [F(-1)] 4238

3.439.1 Optimal result

Integrand size = 35, antiderivative size = 259

$$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{7/2}(c+dx)} dx = \frac{(a+ib)^2(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia-b}d}$$

$$+ \frac{(ia+b)^{3/2}(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} - \frac{2aA\sqrt{a+b \tan(c+dx)}}{5d \tan^{5/2}(c+dx)}$$

$$- \frac{2(6Ab+5aB)\sqrt{a+b \tan(c+dx)}}{15d \tan^{3/2}(c+dx)} + \frac{2(15a^2A-3Ab^2-20abB)\sqrt{a+b \tan(c+dx)}}{15ad\sqrt{\tan(c+dx)}}$$

```
output (I*a+b)^(3/2)*(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d+(a+I*b)^2*(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d/(I*a-b)^(1/2)+2/15*(15*A*a^2-3*A*b^2-20*B*a*b)*(a+b*tan(d*x+c))^(1/2)/a/d/tan(d*x+c)^(1/2)-2/5*a*A*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(5/2)-2/15*(6*A*b+5*B*a)*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(3/2)
```

3.439.2 Mathematica [A] (verified)

Time = 3.34 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \frac{-30\sqrt[4]{-1}a \left((-a + ib)^{3/2} (A - iB) \arctan \left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{a+b\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}} \right) \right)}{\tan^{7/2}(c + dx)}$$

input `Integrate[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(-30*(-1)^(1/4)*a*((-a + I*b)^(3/2)*(A - I*B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]) - (a + I*b)^(3/2)*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(5/2) - 15*a*b*B*Sqrt[a + b*Tan[c + d*x]] - 3*a*(4*a*A - 5*b*B)*Sqrt[a + b*Tan[c + d*x]] - 4*a*(6*A*b + 5*a*B)*Tan[c + d*x]*Sqrt[a + b*Tan[c + d*x]] + 4*(15*a^2*A - 3*A*b^2 - 20*a*b*B)*Tan[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]])/(30*a*d*Tan[c + d*x]^(5/2))`

3.439.3 Rubi [A] (verified)

Time = 1.84 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.12, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 4088, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan(c + dx)^{7/2}} dx$$

↓ 4088

$$\frac{2}{5} \int \frac{-b(4aA - 5bB) \tan^2(c + dx) - 5(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(6Ab + 5aB)}{2 \tan^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx -$$

$$\frac{2aA \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{5} \int \frac{-b(4aA - 5bB) \tan^2(c + dx) - 5(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(6Ab + 5aB)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx -$$

$$\frac{2aA \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{5} \int \frac{-b(4aA - 5bB) \tan(c + dx)^2 - 5(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(6Ab + 5aB)}{\tan(c + dx)^{5/2} \sqrt{a + b \tan(c + dx)}} dx -$$

$$\frac{2aA \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)}$$

↓ 4132

$$\frac{1}{5} \left(\frac{2 \int \frac{2ab(6Ab + 5aB) \tan^2(c + dx) + 15a(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(15Aa^2 - 20bBa - 3Ab^2)}{2 \tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx}{3a} - \frac{2(5aB + 6Ab) \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right)$$

$$\frac{2aA \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{5} \left(\frac{\int \frac{2ab(6Ab + 5aB) \tan^2(c + dx) + 15a(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(15Aa^2 - 20bBa - 3Ab^2)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx}{3a} - \frac{2(5aB + 6Ab) \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right)$$

$$\frac{2aA \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{5} \left(\frac{\int \frac{2ab(6Ab + 5aB) \tan(c + dx)^2 + 15a(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(15Aa^2 - 20bBa - 3Ab^2)}{\tan(c + dx)^{3/2} \sqrt{a + b \tan(c + dx)}} dx}{3a} - \frac{2(5aB + 6Ab) \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right)$$

$$\frac{2aA \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)}$$

↓ 4132

3.439. $\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx$

$$\frac{1}{5} \left(-\frac{2 \int \frac{15(a^2(Ba^2+2Aba-b^2B)-a^2(Aa^2-2bBa-Ab^2)\tan(c+dx)) dx}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}}{a} - \frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} \right)$$

$$\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{5} \left(\frac{15 \int \frac{a^2(Ba^2+2Aba-b^2B)-a^2(Aa^2-2bBa-Ab^2)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a} - \frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} \right)$$

$$\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{5} \left(\frac{15 \int \frac{a^2(Ba^2+2Aba-b^2B)-a^2(Aa^2-2bBa-Ab^2)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a} - \frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} \right)$$

$$\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}$$

↓ 4099

$$-\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} +$$

$$\frac{1}{5} \left(-\frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{15\left(\frac{1}{2}a^2(a-ib)^2(B+iA)\int \frac{i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx\right)}{3a} \right)$$

↓ 3042

$$-\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} +$$

$$\frac{1}{5} \left(-\frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{15\left(\frac{1}{2}a^2(a-ib)^2(B+iA)\int \frac{i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx\right)}{3a} \right)$$

3.439. $\int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \downarrow 4098 \\
 & -\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} + \\
 \frac{1}{5} \left(\right. & -\frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{15 \left(\frac{a^2(a-ib)^2(B+iA) \int \frac{\sqrt{\tan(c+dx)}}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}} dx \right)}{2d} \\
 & \left. \right)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 104 \\
 & -\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} + \\
 \frac{1}{5} \left(\right. & -\frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{15 \left(\frac{a^2(a-ib)^2(B+iA) \int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} dx}{d} \right)}{3a} \\
 & \left. \right)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 216 \\
 & -\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} + \\
 \frac{1}{5} \left(\right. & -\frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{15 \left(\frac{a^2(a-ib)^2(B+iA) \int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} dx}{d} \right)}{3a} \\
 & \left. \right)
 \end{aligned}$$

\downarrow 219

3.439. $\int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$-\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} + \frac{1}{5} \left(\frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{15 \left(\frac{a^2(a-ib)^2(B+ia)\operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{a+b\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}} \right)}{3a} \right)$$

input `Int[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2), x]`

output `(-2*a*A*Sqrt[a + b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) + ((-2*(6*A*b + 5*A*B)*Sqrt[a + b*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - ((15*(-((a^2*(a + I*b)^2*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a - b]*d)) + (a^2*(a - I*b)^2*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]/(Sqrt[I*a + b]*d))))/a - (2*(15*a^2*A - 3*A*b^2 - 20*a*b*B)*Sqrt[a + b*Tan[c + d*x]]/(d*Sqrt[Tan[c + d*x]]))/(3*a))/5`

3.439.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 104 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q], x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

3.439.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.88 (sec) , antiderivative size = 2400946, normalized size of antiderivative = 9270.06

output too large to display

```
input int((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x)
```

```
output result too large to display
```

3.439.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12093 vs. $2(211) = 422$.

Time = 2.32 (sec) , antiderivative size = 12093, normalized size of antiderivative = 46.69

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algo
rithm="fracas")
```

```
output Too large to include
```

3.439. $\int \frac{(a+b \tan(c+dx))^{3/2} (A+B \tan(c+dx))}{\tan^{7/2}(c+dx)} dx$

3.439.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2}}{\tan^{7/2}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(7/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)/tan(c + d*x)**(7/2), x)`

3.439.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="maxima")`

output `Timed out`

3.439.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="giac")`

output `Timed out`

3.439.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2}}{\tan(c + dx)^{7/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/tan(c + d*x)^(7/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/tan(c + d*x)^(7/2),x)`

$$3.440 \quad \int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

3.440.1 Optimal result	4239
3.440.2 Mathematica [A] (verified)	4240
3.440.3 Rubi [A] (verified)	4240
3.440.4 Maple [B] (warning: unable to verify)	4247
3.440.5 Fricas [B] (verification not implemented)	4247
3.440.6 Sympy [F(-1)]	4248
3.440.7 Maxima [F]	4248
3.440.8 Giac [F(-1)]	4248
3.440.9 Mupad [F(-1)]	4249

3.440.1 Optimal result

Integrand size = 35, antiderivative size = 311

$$\int \frac{(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx =$$

$$\frac{(ia - b)^{3/2}(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$- \frac{(ia + b)^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$- \frac{2aA\sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{2(8Ab + 7aB)\sqrt{a + b \tan(c + dx)}}{35d \tan^{\frac{5}{2}}(c + dx)}$$

$$+ \frac{2(35a^2A - 3Ab^2 - 42abB)\sqrt{a + b \tan(c + dx)}}{105ad \tan^{\frac{3}{2}}(c + dx)}$$

$$+ \frac{2(140a^2Ab + 6Ab^3 + 105a^3B - 21ab^2B)\sqrt{a + b \tan(c + dx)}}{105a^2d\sqrt{\tan(c + dx)}}$$

```
output -(I*a-b)^(3/2)*(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d-(I*a+b)^(3/2)*(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d+2/105*(140*A*a^2*b+6*A*b^3+105*B*a^3-21*B*a*b^2)*(a+b*tan(d*x+c))^(1/2)/a^2/d/tan(d*x+c)^(1/2)-2/7*a*A*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(7/2)-2/35*(8*A*b+7*B*a)*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(5/2)+2/105*(35*A*a^2-3*A*b^2-42*B*a*b)*(a+b*tan(d*x+c))^(1/2)/a/d/tan(d*x+c)^(3/2)
```

$$3.440. \quad \int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

3.440.2 Mathematica [A] (verified)

Time = 5.66 (sec) , antiderivative size = 346, normalized size of antiderivative = 1.11

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \frac{-35a^3bB\sqrt{a + b \tan(c + dx)} - 5a^3(6aA - 7bB)\sqrt{a + b \tan(c + dx)}}{\tan^{9/2}(c + dx)}$$

input `Integrate[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2),x]`

output `(-35*a^3*b*B*Sqrt[a + b*Tan[c + d*x]] - 5*a^3*(6*a*A - 7*b*B)*Sqrt[a + b*Tan[c + d*x]] - 6*a^3*(8*A*b + 7*a*B)*Tan[c + d*x]*Sqrt[a + b*Tan[c + d*x]] + a*Tan[c + d*x]^2*(-105*(-1)^(3/4)*a^2*((-a + I*b)^(3/2)*(A - I*B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]) + (a + I*b)^(3/2)*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(3/2) + 2*a*(35*a^2*A - 3*A*b^2 - 42*a*b*B)*Sqrt[a + b*Tan[c + d*x]] + 2*(140*a^2*A*b + 6*A*b^3 + 10*5*a^3*B - 21*a*b^2*B)*Tan[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/(105*a^3*d*Tan[c + d*x]^(7/2))`

3.440.3 Rubi [A] (verified)

Time = 2.37 (sec) , antiderivative size = 352, normalized size of antiderivative = 1.13, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.543$, Rules used = {3042, 4088, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan(c + dx)^{9/2}} dx$$

↓ 4088

3.440. $\int \frac{(a+b \tan(c+dx))^{3/2} (A+B \tan(c+dx))}{\tan^{9/2}(c+dx)} dx$

$$\frac{2}{7} \int \frac{-b(6aA - 7bB) \tan^2(c + dx) - 7(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(8Ab + 7aB)}{2 \tan^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx - \frac{2aA \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{7} \int \frac{-b(6aA - 7bB) \tan^2(c + dx) - 7(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(8Ab + 7aB)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx - \frac{2aA \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \int \frac{-b(6aA - 7bB) \tan(c + dx)^2 - 7(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(8Ab + 7aB)}{\tan(c + dx)^{7/2} \sqrt{a + b \tan(c + dx)}} dx - \frac{2aA \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 4132

$$\frac{1}{7} \left(\frac{2 \int \frac{4ab(8Ab+7aB) \tan^2(c+dx)+35a(Ba^2+2Aba-b^2B) \tan(c+dx)+a(35Aa^2-42bBa-3Ab^2)}{2 \tan^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2(7aB+8Ab) \sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right) - \frac{2aA \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{7} \left(\frac{\int \frac{4ab(8Ab+7aB) \tan^2(c+dx)+35a(Ba^2+2Aba-b^2B) \tan(c+dx)+a(35Aa^2-42bBa-3Ab^2)}{\tan^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2(7aB+8Ab) \sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right) - \frac{2aA \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{\int \frac{4ab(8Ab+7aB) \tan(c+dx)^2+35a(Ba^2+2Aba-b^2B) \tan(c+dx)+a(35Aa^2-42bBa-3Ab^2)}{\tan(c+dx)^{5/2} \sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2(7aB+8Ab) \sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right) - \frac{2aA \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 4132

3.440. $\int \frac{(a+b \tan(c+dx))^{3/2} (A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

$$\frac{1}{7} \left(\frac{2 \int \frac{-105(Aa^2 - 2bBa - Ab^2) \tan(c+dx)a^2 - 2b(35Aa^2 - 42bBa - 3Ab^2) \tan^2(c+dx)a + (105Ba^3 + 140Aba^2 - 21b^2Ba + 6Ab^3)a}{2 \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx - \frac{2(35a^2A - 42abB - 3Aa^2)}{3d} \right)$$

$$\frac{2aA \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{7} \left(\int \frac{-105(Aa^2 - 2bBa - Ab^2) \tan(c+dx)a^2 - 2b(35Aa^2 - 42bBa - 3Ab^2) \tan^2(c+dx)a + (105Ba^3 + 140Aba^2 - 21b^2Ba + 6Ab^3)a}{\tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx - \frac{2(35a^2A - 42abB - 3Aa^2)}{3d \tan^{\frac{3}{2}}(c+dx)} \right)$$

$$\frac{2aA \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \left(\int \frac{-105(Aa^2 - 2bBa - Ab^2) \tan(c+dx)a^2 - 2b(35Aa^2 - 42bBa - 3Ab^2) \tan(c+dx)^2 a + (105Ba^3 + 140Aba^2 - 21b^2Ba + 6Ab^3)a}{\tan(c+dx)^{\frac{3}{2}} \sqrt{a+b \tan(c+dx)}} dx - \frac{2(35a^2A - 42abB - 3Aa^2)}{3d \tan^{\frac{3}{2}}(c+dx)} \right)$$

$$\frac{2aA \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 4132

$$\frac{1}{7} \left(\int \frac{2 \int \frac{105((Aa^2 - 2bBa - Ab^2)a^3 + (Ba^2 + 2Aba - b^2B) \tan(c+dx)a^3)}{2 \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx - \frac{2(105a^3B + 140a^2Ab - 21ab^2B + 6Ab^3) \sqrt{a+b \tan(c+dx)}}{d \sqrt{\tan(c+dx)}}}{3a} - \frac{2(35a^2A - 42abB - 3Aa^2)}{3d \tan^{\frac{3}{2}}(c+dx)} \right)$$

$$\frac{2aA \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 27

3.440. $\int \frac{(a+b \tan(c+dx))^{3/2} (A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

$$\frac{1}{7} \left(\frac{105 \int \frac{(Aa^2 - 2bBa - Ab^2)a^3 + (Ba^2 + 2Aba - b^2B) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{2(105a^3B + 140a^2Ab - 21ab^2B + 6Ab^3)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2(35a^2A - 42abB - 3Ab^2)\sqrt{a+b \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)$$

$$\frac{2aA\sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{105 \int \frac{(Aa^2 - 2bBa - Ab^2)a^3 + (Ba^2 + 2Aba - b^2B) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{2(105a^3B + 140a^2Ab - 21ab^2B + 6Ab^3)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2(35a^2A - 42abB - 3Ab^2)\sqrt{a+b \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)$$

$$\frac{2aA\sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 4099

$$-\frac{2aA\sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} +$$

$$\frac{1}{7} \left(\frac{2(7aB + 8Ab)\sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A - 42abB - 3Ab^2)\sqrt{a+b \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{2(105a^3B + 140a^2Ab - 21ab^2B + 6Ab^3)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)$$

↓ 3042

$$-\frac{2aA\sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} +$$

$$\frac{1}{7} \left(\frac{2(7aB + 8Ab)\sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A - 42abB - 3Ab^2)\sqrt{a+b \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{2(105a^3B + 140a^2Ab - 21ab^2B + 6Ab^3)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)$$

↓ 4098

3.440. $\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

$$\begin{aligned}
 & -\frac{2aA\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \\
 \frac{1}{7} \left(& -\frac{2(7aB+8Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-42abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \frac{2(105a^3B+140a^2Ab-21ab^2B+6Ab^3)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)
 \end{aligned}$$

↓ 104

$$\begin{aligned}
 & -\frac{2aA\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \\
 \frac{1}{7} \left(& -\frac{2(7aB+8Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-42abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \frac{2(105a^3B+140a^2Ab-21ab^2B+6Ab^3)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)
 \end{aligned}$$

↓ 216

$$\begin{aligned}
 & -\frac{2aA\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \\
 \frac{1}{7} \left(& -\frac{2(7aB+8Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-42abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \frac{2(105a^3B+140a^2Ab-21ab^2B+6Ab^3)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)
 \end{aligned}$$

↓ 219

3.440. $\int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

$$-\frac{2aA\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \left(\frac{1}{7} \left[-\frac{2(7aB+8Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-42abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} + \frac{2(105a^3B+140a^2Ab-21ab^2B+6Ab^3)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right] \right)$$

```
input Int[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2),x
]
```

```
output (-2*a*A*Sqrt[a + b*Tan[c + d*x]]/(7*d*Tan[c + d*x]^(7/2)) + ((-2*(8*A*b +
7*a*B)*Sqrt[a + b*Tan[c + d*x]]/(5*d*Tan[c + d*x]^(5/2)) - ((-2*(35*a^2*
A - 3*A*b^2 - 42*a*b*B)*Sqrt[a + b*Tan[c + d*x]]/(3*d*Tan[c + d*x]^(3/2))
+ ((-105*((a^3*(a + I*b)^2*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d
*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a - b]*d) + (a^3*(a - I*b)^2*(A -
I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])
]/(Sqrt[I*a + b]*d)))/a - (2*(140*a^2*A*b + 6*A*b^3 + 105*a^3*B - 21*a*b^2
*B)*Sqrt[a + b*Tan[c + d*x]]/(d*Sqrt[Tan[c + d*x]]))/(3*a))/(5*a))/7
```

3.440.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_)] /; FreeQ[b, x]
```

```
rule 104 Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))/((e_) + (f_)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

3.440. $\int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

3.440.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.98 (sec) , antiderivative size = 2403086, normalized size of antiderivative = 7726.96

output too large to display

```
input int((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x)
```

```
output result too large to display
```

3.440.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12128 vs. $2(259) = 518$.

Time = 2.46 (sec) , antiderivative size = 12128, normalized size of antiderivative = 39.00

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algo
rithm="fracas")
```

```
output Too large to include
```

3.440. $\int \frac{(a+b \tan(c+dx))^{3/2} (A+B \tan(c+dx))}{\tan^{9/2}(c+dx)} dx$

3.440.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(9/2),x)`

output `Timed out`

3.440.7 Maxima [F]

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^{3/2}}{\tan^2(dx + c)} dx$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)/tan(d*x + c)^(9/
2), x)`

3.440.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algo
rithm="giac")`

output `Timed out`

3.440. $\int \frac{(a+b \tan(c+dx))^{3/2} (A+B \tan(c+dx))}{\tan^{9/2}(c+dx)} dx$

3.440.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2}}{\tan(c + dx)^{9/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/tan(c + d*x)^(9/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/tan(c + d*x)^(9/2),x)`

3.441
$$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$$

3.441.1 Optimal result 4250
 3.441.2 Mathematica [A] (verified) 4251
 3.441.3 Rubi [A] (verified) 4252
 3.441.4 Maple [B] (warning: unable to verify) 4262
 3.441.5 Fricas [B] (verification not implemented) 4262
 3.441.6 Sympy [F(-1)] 4262
 3.441.7 Maxima [F(-1)] 4263
 3.441.8 Giac [F(-1)] 4263
 3.441.9 Mupad [F(-1)] 4263

3.441.1 Optimal result

Integrand size = 35, antiderivative size = 382

$$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx = \frac{(ia-b)^{3/2}(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} - \frac{(ia+b)^{3/2}(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} - \frac{2aA\sqrt{a+b \tan(c+dx)}}{9d \tan^{\frac{9}{2}}(c+dx)} - \frac{2(10Ab+9aB)\sqrt{a+b \tan(c+dx)}}{63d \tan^{\frac{7}{2}}(c+dx)} + \frac{2(21a^2A-Ab^2-24abB)\sqrt{a+b \tan(c+dx)}}{105ad \tan^{\frac{5}{2}}(c+dx)} + \frac{2(126a^2Ab+4Ab^3+105a^3B-9ab^2B)\sqrt{a+b \tan(c+dx)}}{315a^2d \tan^{\frac{3}{2}}(c+dx)} - \frac{2(315a^4A-63a^2Ab^2+8Ab^4-420a^3bB-18ab^3B)\sqrt{a+b \tan(c+dx)}}{315a^3d\sqrt{\tan(c+dx)}}$$

```
output (I*a-b)^(3/2)*(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d-(I*a+b)^(3/2)*(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d-2/315*(315*A*a^4-63*A*a^2*b^2+8*A*b^4-420*B*a^3*b-18*B*a*b^3)*(a+b*tan(d*x+c))^(1/2)/a^3/d/tan(d*x+c)^(1/2)-2/9*a*A*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(9/2)-2/63*(10*A*b+9*B*a)*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(7/2)+2/105*(21*A*a^2-A*b^2-24*B*a*b)*(a+b*tan(d*x+c))^(1/2)/a/d/tan(d*x+c)^(5/2)+2/315*(126*A*a^2*b+4*A*b^3+105*B*a^3-9*B*a*b^2)*(a+b*tan(d*x+c))^(1/2)/a^2/d/tan(d*x+c)^(3/2)
```

3.441.
$$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$$

3.441.2 Mathematica [A] (verified)

Time = 6.78 (sec) , antiderivative size = 474, normalized size of antiderivative = 1.24

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx =$$

$$-\frac{bB\sqrt{a + b \tan(c + dx)}}{4d \tan^{\frac{9}{2}}(c + dx)} + \frac{1}{4} - \frac{(8aA - 9bB)\sqrt{a + b \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

3.441. $\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx$

input `Integrate[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(11/2),x]`

output `-1/4*(b*B*Sqrt[a + b*Tan[c + d*x]]/(d*Tan[c + d*x]^(9/2)) + (-1/9*((8*a*A - 9*b*B)*Sqrt[a + b*Tan[c + d*x]]/(d*Tan[c + d*x]^(9/2)) + (2*((-4*a*(10*A*b + 9*a*B)*Sqrt[a + b*Tan[c + d*x]]/(7*d*Tan[c + d*x]^(7/2)) - (2*((-6*a*(21*a^2*A - A*b^2 - 24*a*b*B)*Sqrt[a + b*Tan[c + d*x]]/(5*d*Tan[c + d*x]^(5/2)) - (2*((a*(126*a^2*A*b + 4*A*b^3 + 105*a^3*B - 9*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]]/(d*Tan[c + d*x]^(3/2)) - (2*((-945*a^4*((-1)^(1/4))*(-a + I*b)^(3/2)*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - (-1)^(1/4)*(a + I*b)^(3/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]))/(4*d) + (3*a*(315*a^4*A - 63*a^2*A*b^2 + 8*A*b^4 - 420*a^3*b*B - 18*a*b^3*B)*Sqrt[a + b*Tan[c + d*x]]/(2*d*Sqrt[Tan[c + d*x]])))/(3*a))/(5*a))/(7*a))/(9*a))/4`

3.441.3 Rubi [A] (verified)

Time = 2.98 (sec) , antiderivative size = 430, normalized size of antiderivative = 1.13, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.629$, Rules used = {3042, 4088, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan(c + dx)^{11/2}} dx$$

↓ 4088

$$\frac{2}{9} \int \frac{-b(8aA - 9bB) \tan^2(c + dx) - 9(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(10Ab + 9aB)}{2 \tan^{\frac{9}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} + \frac{2aA \sqrt{a + b \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)}} dx -$$

↓ 27

3.441. $\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx$

$$\begin{aligned}
& \frac{1}{9} \int \frac{-b(8aA - 9bB) \tan^2(c + dx) - 9(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(10Ab + 9aB)}{\tan^{\frac{9}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx - \\
& \quad \frac{2aA \sqrt{a + b \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{9} \int \frac{-b(8aA - 9bB) \tan(c + dx)^2 - 9(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(10Ab + 9aB)}{\tan(c + dx)^{9/2} \sqrt{a + b \tan(c + dx)}} dx - \\
& \quad \frac{2aA \sqrt{a + b \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} \\
& \quad \downarrow \text{4132} \\
& \frac{1}{9} \left(\frac{2 \int \frac{3(2ab(10Ab + 9aB) \tan^2(c + dx) + 21a(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(21Aa^2 - 24bBa - Ab^2))}{2 \tan^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx}{7a} - \frac{2(9aB + 10Ab) \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} \right) \\
& \quad \frac{2aA \sqrt{a + b \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{9} \left(\frac{3 \int \frac{2ab(10Ab + 9aB) \tan^2(c + dx) + 21a(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(21Aa^2 - 24bBa - Ab^2)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx}{7a} - \frac{2(9aB + 10Ab) \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} \right) \\
& \quad \frac{2aA \sqrt{a + b \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{9} \left(\frac{3 \int \frac{2ab(10Ab + 9aB) \tan(c + dx)^2 + 21a(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(21Aa^2 - 24bBa - Ab^2)}{\tan(c + dx)^{7/2} \sqrt{a + b \tan(c + dx)}} dx}{7a} - \frac{2(9aB + 10Ab) \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} \right) \\
& \quad \frac{2aA \sqrt{a + b \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} \\
& \quad \downarrow \text{4132}
\end{aligned}$$

3.441. $\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx$

$$\frac{1}{9} \left(3 \left(\frac{2 \int \frac{-105(Aa^2 - 2bBa - Ab^2) \tan(c+dx)a^2 - 4b(21Aa^2 - 24bBa - Ab^2) \tan^2(c+dx)a + (105Ba^3 + 126Aba^2 - 9b^2Ba + 4Ab^3)a}{2 \tan^{\frac{5}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2(21a^2A - 24abB - Ab^2)}{5d \tan^{\frac{5}{2}}(c+dx)} \right) - 7a \right)$$

$$\frac{2aA\sqrt{a+b \tan(c+dx)}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{9} \left(3 \left(\frac{\int \frac{-105(Aa^2 - 2bBa - Ab^2) \tan(c+dx)a^2 - 4b(21Aa^2 - 24bBa - Ab^2) \tan^2(c+dx)a + (105Ba^3 + 126Aba^2 - 9b^2Ba + 4Ab^3)a}{\tan^{\frac{5}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2(21a^2A - 24abB - Ab^2)}{5d \tan^{\frac{5}{2}}(c+dx)} \right) - 7a \right)$$

$$\frac{2aA\sqrt{a+b \tan(c+dx)}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{9} \left(3 \left(\frac{\int \frac{-105(Aa^2 - 2bBa - Ab^2) \tan(c+dx)a^2 - 4b(21Aa^2 - 24bBa - Ab^2) \tan(c+dx)^2a + (105Ba^3 + 126Aba^2 - 9b^2Ba + 4Ab^3)a}{\tan(c+dx)^{5/2}\sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2(21a^2A - 24abB - Ab^2)}{5d \tan^{\frac{5}{2}}(c+dx)} \right) - 7a \right)$$

$$\frac{2aA\sqrt{a+b \tan(c+dx)}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

↓ 4132

3.441. $\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$

$$\left(\frac{1}{9} \right) \left(\frac{3}{3} \right) \left(\frac{\int \frac{315(Ba^2+2Aba-b^2B)\tan(c+dx)a^3+2b(105Ba^3+126Aba^2-9b^2Ba+4Ab^3)\tan^2(c+dx)a+(315Aa^4-420bBa^3-63Ab^2a^2-18b^3Ba+8Ab^4)a}{2\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}} dx}{5a} \right)$$

7a

$$\frac{2aA\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)}$$

↓ 27

$$\left(\frac{1}{9} \right) \left(\frac{3}{3} \right) \left(\frac{\int \frac{315(Ba^2+2Aba-b^2B)\tan(c+dx)a^3+2b(105Ba^3+126Aba^2-9b^2Ba+4Ab^3)\tan^2(c+dx)a+(315Aa^4-420bBa^3-63Ab^2a^2-18b^3Ba+8Ab^4)a}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}} dx}{5a} \right)$$

7a

$$\frac{2aA\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)}$$

↓ 3042

$$\left(\frac{1}{9} \right) \left(\frac{3}{3} \right) \left(\frac{\int \frac{315(Ba^2+2Aba-b^2B)\tan(c+dx)a^3+2b(105Ba^3+126Aba^2-9b^2Ba+4Ab^3)\tan(c+dx)^2a+(315Aa^4-420bBa^3-63Ab^2a^2-18b^3Ba+8Ab^4)a}{\tan(c+dx)^{3/2}\sqrt{a+b\tan(c+dx)}} dx}{5a} \right)$$

7a

$$\frac{2aA\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)}$$

3.441. $\int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$

↓ 4132

$$\left(\frac{1}{9} \right) \left(3 \left(\frac{2 \int \frac{315(a^4(Ba^2+2Aba-b^2B)-a^4(Aa^2-2bBa-Ab^2)\tan(c+dx))}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{2(315a^4A-420a^3bB-63a^2Ab^2-18ab^3B+8Ab^4)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2(105a^3B}{5a} \right) \right) - 7a$$

$$\frac{2aA\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)}$$

↓ 27

$$\left(\frac{1}{9} \right) \left(3 \left(\frac{315 \int \frac{a^4(Ba^2+2Aba-b^2B)-a^4(Aa^2-2bBa-Ab^2)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{2(315a^4A-420a^3bB-63a^2Ab^2-18ab^3B+8Ab^4)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2(105a^3B}{5a} \right) \right) - 7a$$

$$\frac{2aA\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)}$$

↓ 3042

3.441. $\int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$

$$\frac{1}{9} \left(\frac{3 \left(\frac{315 \int \frac{a^4 (Ba^2 + 2Aba - b^2 B) - a^4 (Aa^2 - 2bBa - Ab^2) \tan(c+dx)}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx - \frac{2(315a^4 A - 420a^3 bB - 63a^2 Ab^2 - 18ab^3 B + 8Ab^4) \sqrt{a+b \tan(c+dx)}}{d \sqrt{\tan(c+dx)}} - \frac{2(105a^3 B}{7a} \right)}{7a} \right)$$

$$\frac{2aA \sqrt{a + b \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 4099

$$- \frac{2aA \sqrt{a + b \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} +$$

$$\frac{1}{9} \left(\frac{2(9aB + 10Ab) \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{3 \left(- \frac{2(21a^2 A - 24abB - Ab^2) \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} + \frac{2(105a^3 B + 126a^2 Ab - 9ab^2 B + 4Ab^3) \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right)}{7d \tan^{\frac{7}{2}}(c + dx)} \right)$$

↓ 3042

$$- \frac{2aA \sqrt{a + b \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} +$$

$$\frac{1}{9} \left(\frac{2(9aB + 10Ab) \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{3 \left(- \frac{2(21a^2 A - 24abB - Ab^2) \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} + \frac{2(105a^3 B + 126a^2 Ab - 9ab^2 B + 4Ab^3) \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right)}{7d \tan^{\frac{7}{2}}(c + dx)} \right)$$

3.441. $\int \frac{(a+b \tan(c+dx))^{3/2} (A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$

$$\frac{1}{9} \left(\frac{2(9aB + 10Ab)\sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{2aA\sqrt{a + b \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} + 3 \left(-\frac{2(21a^2A - 24abB - Ab^2)\sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} + \frac{2(105a^3B + 126a^2Ab - 9ab^2B + 4Ab^3)\sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right) \right)$$

$$\frac{1}{9} \left(\frac{2(9aB + 10Ab)\sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{2aA\sqrt{a + b \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} + 3 \left(-\frac{2(21a^2A - 24abB - Ab^2)\sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} + \frac{2(105a^3B + 126a^2Ab - 9ab^2B + 4Ab^3)\sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right) \right)$$

↓ 216

3.441. $\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx$

$$\frac{1}{9} \left(\frac{2(9aB + 10Ab)\sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{\frac{2aA\sqrt{a + b \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} + 3 \left(-\frac{2(21a^2A - 24abB - Ab^2)\sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} + \frac{2(105a^3B + 126a^2Ab - 9ab^2B + 4Ab^3)\sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right)}{\right)}$$

↓ 219

$$\frac{1}{9} \left(\frac{2(9aB + 10Ab)\sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{\frac{2aA\sqrt{a + b \tan(c + dx)}}{9d \tan^{\frac{9}{2}}(c + dx)} + 3 \left(-\frac{2(21a^2A - 24abB - Ab^2)\sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} + \frac{2(105a^3B + 126a^2Ab - 9ab^2B + 4Ab^3)\sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right)}{\right)}$$

```
input Int[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(11/2), x]
```

3.441. $\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx$

```
output (-2*a*A*Sqrt[a + b*Tan[c + d*x]]/(9*d*Tan[c + d*x]^(9/2)) + ((-2*(10*A*b
+ 9*a*B)*Sqrt[a + b*Tan[c + d*x]]/(7*d*Tan[c + d*x]^(7/2)) - (3*((-2*(21*
a^2*A - A*b^2 - 24*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2
)) + ((-2*(126*a^2*A*b + 4*A*b^3 + 105*a^3*B - 9*a*b^2*B)*Sqrt[a + b*Tan[c
+ d*x]])/(3*d*Tan[c + d*x]^(3/2)) - ((315*(-((a^4*(a + I*b)^2*(I*A - B)*A
rcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]))/(Sqrt[
I*a - b]*d)) + (a^4*(a - I*b)^2*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[
c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]))/(Sqrt[I*a + b]*d)))/a - (2*(315*a^4*
A - 63*a^2*A*b^2 + 8*A*b^4 - 420*a^3*b*B - 18*a*b^3*B)*Sqrt[a + b*Tan[c +
d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a))/(5*a))/(7*a))/9
```

3.441.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 104 Int[(((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)))/((e_.) + (f_.)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.441.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.90 (sec) , antiderivative size = 2403057, normalized size of antiderivative = 6290.73

output too large to display

input `int((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x)`

output `result too large to display`

3.441.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12202 vs. $2(324) = 648$.

Time = 2.43 (sec) , antiderivative size = 12202, normalized size of antiderivative = 31.94

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x, algorithm="fricas")`

output `Too large to include`

3.441.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(11/2),x)`

output `Timed out`

3.441.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x, algorithm="maxima")`

output `Timed out`

3.441.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x, algorithm="giac")`

output `Timed out`

3.441.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2}}{\tan(c + dx)^{11/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/tan(c + d*x)^(11/2), x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/tan(c + d*x)^(11/2), x)`

3.441. $\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{11/2}(c+dx)} dx$

3.442 $\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.442.1 Optimal result	4264
3.442.2 Mathematica [A] (verified)	4265
3.442.3 Rubi [A] (verified)	4266
3.442.4 Maple [B] (warning: unable to verify)	4270
3.442.5 Fricas [B] (verification not implemented)	4271
3.442.6 Sympy [F(-1)]	4271
3.442.7 Maxima [F]	4271
3.442.8 Giac [F(-1)]	4272
3.442.9 Mupad [F(-1)]	4272

3.442.1 Optimal result

Integrand size = 35, antiderivative size = 397

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx =$$

$$\begin{aligned} & - \frac{(ia - b)^{5/2}(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} \\ & + \frac{(40a^3Ab - 320aAb^3 - 5a^4B - 240a^2b^2B + 128b^4B) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{64b^{3/2}d} \\ & - \frac{(ia + b)^{5/2}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} \\ & + \frac{(40a^2Ab - 64Ab^3 - 5a^3B - 112ab^2B) \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}{64bd} \\ & + \frac{(40aAb - 5a^2B - 48b^2B) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}}{96bd} \\ & + \frac{(8Ab - aB) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}}{24bd} \\ & + \frac{B \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{7/2}}{4bd} \end{aligned}$$

output
$$-(I*a-b)^{(5/2)}*(I*A-B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})/d+1/64*(40*A*a^3*b-320*A*a*b^3-5*B*a^4-240*B*a^2*b^2+128*B*b^4)*\operatorname{arctanh}(b^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})/b^{(3/2)}/d-(I*a+b)^{(5/2)}*(I*A+B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})/d+1/64*(40*A*a^2*b-64*A*b^3-5*B*a^3-112*B*a*b^2)*\tan(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(1/2)}/b/d+1/96*(40*A*a*b-5*B*a^2-48*B*b^2)*\tan(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(3/2)}/b/d+1/24*(8*A*b-B*a)*\tan(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(5/2)}/b/d+1/4*B*\tan(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(7/2)}/b/d$$

3.442.2 Mathematica [A] (verified)

Time = 4.98 (sec) , antiderivative size = 411, normalized size of antiderivative = 1.04

$$\int \tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx = \frac{-192\sqrt[4]{-1}(-a+ib)^{5/2}b(iA+B)\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)+192(-1)^{3/4}(a+ib)^{5/2}b(iA+B)\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{1}$$

input `Integrate[Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]), x]`

output
$$(-192*(-1)^{(1/4)}*(-a + I*b)^{(5/2)}*b*(I*A + B)*\operatorname{ArcTan}(((-1)^{(1/4)}*\sqrt{-a + I*b})*\sqrt{\tan[c + d*x]})/\sqrt{a + b*\tan[c + d*x]}) + 192*(-1)^{(3/4)}*(a + I*b)^{(5/2)}*b*(A + I*B)*\operatorname{ArcTan}(((-1)^{(1/4)}*\sqrt{a + I*b})*\sqrt{\tan[c + d*x]})/\sqrt{a + b*\tan[c + d*x]}) - 3*(-40*a^2*A*b + 64*A*b^3 + 5*a^3*B + 112*a*b^2*B)*\sqrt{\tan[c + d*x]}*\sqrt{a + b*\tan[c + d*x]} - 2*(-40*a*A*b + 5*a^2*B + 48*b^2*B)*\sqrt{\tan[c + d*x]}*(a + b*\tan[c + d*x])^{(3/2)} + 8*(8*A*b - a*B)*\sqrt{\tan[c + d*x]}*(a + b*\tan[c + d*x])^{(5/2)} + 48*B*\sqrt{\tan[c + d*x]}*(a + b*\tan[c + d*x])^{(7/2)} - (3*\sqrt{a}*(-40*a^3*A*b + 320*a*A*b^3 + 5*a^4*B + 240*a^2*b^2*B - 128*b^4*B)*\operatorname{ArcSinh}(\sqrt{b}*\sqrt{\tan[c + d*x]})/\sqrt{a})*\sqrt{1 + (b*\tan[c + d*x])/a})/(\sqrt{b}*\sqrt{a + b*\tan[c + d*x]})/(192*b*d)$$

3.442.3 Rubi [A] (verified)

Time = 2.56 (sec) , antiderivative size = 400, normalized size of antiderivative = 1.01, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.486$, Rules used = {3042, 4090, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c+dx)^{3/2}(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4090} \\
 & \int \frac{(a+b \tan(c+dx))^{5/2}(-((8Ab-aB) \tan^2(c+dx))+8bB \tan(c+dx)+aB)}{2\sqrt{\tan(c+dx)}} dx + \\
 & \quad \frac{4b}{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{7/2}} \\
 & \quad \quad \quad \downarrow \text{27} \\
 & \quad \quad \quad \frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{7/2}}{4bd} - \\
 & \int \frac{(a+b \tan(c+dx))^{5/2}(-((8Ab-aB) \tan^2(c+dx))+8bB \tan(c+dx)+aB)}{\sqrt{\tan(c+dx)}} dx \\
 & \quad \quad \quad \frac{8b}{8b} \\
 & \quad \quad \quad \downarrow \text{3042} \\
 & \quad \quad \quad \frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{7/2}}{4bd} - \\
 & \int \frac{(a+b \tan(c+dx))^{5/2}(-((8Ab-aB) \tan(c+dx)^2)+8bB \tan(c+dx)+aB)}{\sqrt{\tan(c+dx)}} dx \\
 & \quad \quad \quad \frac{8b}{8b} \\
 & \quad \quad \quad \downarrow \text{4130} \\
 & \quad \quad \quad \frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{7/2}}{4bd} - \\
 & \frac{1}{3} \int \frac{(a+b \tan(c+dx))^{3/2}(-((-5Ba^2+40Aba-48b^2B) \tan^2(c+dx))+48b(Ab+aB) \tan(c+dx)+a(8Ab+5aB))}{2\sqrt{\tan(c+dx)}} dx - \frac{(8Ab-aB)\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{7/2}}{3d} \\
 & \quad \quad \quad \downarrow \text{27}
 \end{aligned}$$

3.442. $\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd} - \frac{\frac{1}{6} \int \frac{(a+b\tan(c+dx))^{3/2} (-((-5Ba^2+40Aba-48b^2B)\tan^2(c+dx)+48b(Ab+aB)\tan(c+dx)+a(8Ab+5aB)))}{\sqrt{\tan(c+dx)}} dx - \frac{(8Ab-aB)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3d}}{8b}$$

↓ 3042

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd} - \frac{\frac{1}{6} \int \frac{(a+b\tan(c+dx))^{3/2} (-((-5Ba^2+40Aba-48b^2B)\tan^2(c+dx)+48b(Ab+aB)\tan(c+dx)+a(8Ab+5aB)))}{\sqrt{\tan(c+dx)}} dx - \frac{(8Ab-aB)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3d}}{8b}$$

↓ 4130

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd} - \frac{\frac{1}{6} \left(\frac{3}{2} \int \frac{3\sqrt{a+b\tan(c+dx)} (-((-5Ba^3+40Aba^2-112b^2Ba-64Ab^3)\tan^2(c+dx)+64b(Ba^2+2Aba-b^2B)\tan(c+dx)+a(5Ba^2+24Aba-16b^2B)))}{2\sqrt{\tan(c+dx)}} dx - \frac{(8Ab-aB)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3d} \right)}{8b}$$

↓ 27

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd} - \frac{\frac{1}{6} \left(\frac{3}{4} \int \frac{\sqrt{a+b\tan(c+dx)} (-((-5Ba^3+40Aba^2-112b^2Ba-64Ab^3)\tan^2(c+dx)+64b(Ba^2+2Aba-b^2B)\tan(c+dx)+a(5Ba^2+24Aba-16b^2B)))}{\sqrt{\tan(c+dx)}} dx - \frac{(8Ab-aB)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3d} \right)}{8b}$$

↓ 3042

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd} - \frac{\frac{1}{6} \left(\frac{3}{4} \int \frac{\sqrt{a+b\tan(c+dx)} (-((-5Ba^3+40Aba^2-112b^2Ba-64Ab^3)\tan^2(c+dx)+64b(Ba^2+2Aba-b^2B)\tan(c+dx)+a(5Ba^2+24Aba-16b^2B)))}{\sqrt{\tan(c+dx)}} dx - \frac{(8Ab-aB)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3d} \right)}{8b}$$

↓ 4130

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd} - \frac{\frac{1}{6} \left(\frac{3}{4} \left(\int \frac{-((-5Ba^4+40Aba^3-240b^2Ba^2-320Ab^3a+128b^4B)\tan^2(c+dx)+128b(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx)+a(5Ba^3+88Aba^2-16b^4B))}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{(8Ab-aB)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3d} \right) \right)}{8b}$$

↓ 27

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd} - \frac{\frac{1}{6} \left(\frac{3}{4} \left(\frac{1}{2} \int \frac{-((-5Ba^4+40Aba^3-240b^2Ba^2-320Ab^3a+128b^4B)\tan^2(c+dx)+128b(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx)+a(5Ba^3+88Aba^2-16b^4B))}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{(8Ab-aB)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3d} \right) \right)}{8b}$$

3.442. $\int \tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$

↓ 3042

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd} - \frac{\frac{1}{6}\left(\frac{3}{4}\left(\frac{1}{2}\int\frac{-((-5Ba^4+40Aba^3-240b^2Ba^2-320Ab^3a+128b^4B)\tan(c+dx)^2)+128b(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx)+a(5Ba^3+88Aba^2-144b^2Ba-64Ab^3)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx\right)\right)}{1}$$

↓ 4138

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd} - \frac{\frac{1}{6}\left(\frac{3}{4}\left(\frac{\int\frac{-((-5Ba^4+40Aba^3-240b^2Ba^2-320Ab^3a+128b^4B)\tan^2(c+dx))+128b(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx)+a(5Ba^3+88Aba^2-144b^2Ba-64Ab^3)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}dx}{2d}\right)\right)}{1}$$

↓ 2035

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd} - \frac{\frac{1}{6}\left(\frac{3}{4}\left(\frac{\int\frac{-((-5Ba^4+40Aba^3-240b^2Ba^2-320Ab^3a+128b^4B)\tan^2(c+dx))+128b(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx)+a(5Ba^3+88Aba^2-144b^2Ba-64Ab^3)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}dx}{d}\right)\right)}{1}$$

↓ 2257

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd} - \frac{\frac{1}{6}\left(\frac{3}{4}\left(\frac{\int\left(\frac{5Ba^4-40Aba^3+240b^2Ba^2+320Ab^3a-128b^4B}{\sqrt{a+b\tan(c+dx)}}+\frac{128\left(b\left(Aa^3-3bBa^2-3Ab^2a+b^3B\right)+b\left(Ba^3+3Aba^2-3b^2Ba-Ab^3\right)\tan(c+dx)\right)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}\right)dx}{d}\right)\right)}{1}d\sqrt{\tan(c+dx)}$$

↓ 2009

$$\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd} - \frac{(8Ab-aB)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3d} + \frac{1}{6}\left(-\frac{(-5a^2B+40aAb-48b^2B)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d} + \frac{3}{4}\left(-\frac{(-5a^3B+40a^2Ab-48ab^2)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{1/2}}{2d}\right)\right)$$

input `Int[Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

```
output (B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(7/2))/(4*b*d) - (-1/3*((8*A*b
- a*B)*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(5/2))/d + (-1/2*((40*A*A*b
- 5*a^2*B - 48*b^2*B)*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2))/d +
(3*((64*(I*a - b)^(5/2)*b*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x
]])/Sqrt[a + b*Tan[c + d*x]]] - ((40*a^3*A*b - 320*A*A*b^3 - 5*a^4*B - 240
*a^2*b^2*B + 128*b^4*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Ta
n[c + d*x]]])/Sqrt[b] + 64*b*(I*a + b)^(5/2)*(I*A + B)*ArcTanh[(Sqrt[I*a +
b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/d - ((40*a^2*A*b - 64*A
*b^3 - 5*a^3*B - 112*a*b^2*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])
/d)/4)/6)/(8*b)
```

3.442.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2035 Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]
```

```
rule 2257 Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol
] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a
, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4090 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4130 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

3.442.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.06 (sec) , antiderivative size = 2659561, normalized size of antiderivative = 6699.15

output too large to display

```
input int(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x)
```

$$3.442. \quad \int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{\frac{5}{2}}(A + B \tan(c + dx)) dx$$

output result too large to display

3.442.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17728 vs. 2(334) = 668.

Time = 9.41 (sec) , antiderivative size = 35458, normalized size of antiderivative = 89.31

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.442.6 Sympy [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(3/2)*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output Timed out

3.442.7 Maxima [F]

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{5}{2}} \tan(dx + c)^{\frac{3}{2}} dx$$

input `integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)*tan(d*x + c)^(3/2), x)`

3.442.8 Giac [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.442.9 Mupad [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int \tan(c + dx)^{3/2}(A + B \tan(c + dx))(a + b \tan(c + dx))^{5/2} dx$$

input `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)`

output `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2), x)`

3.443 $\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.443.1 Optimal result	4273
3.443.2 Mathematica [A] (verified)	4274
3.443.3 Rubi [A] (verified)	4275
3.443.4 Maple [B] (warning: unable to verify)	4279
3.443.5 Fricas [B] (verification not implemented)	4280
3.443.6 Sympy [F(-1)]	4280
3.443.7 Maxima [F]	4280
3.443.8 Giac [F(-1)]	4281
3.443.9 Mupad [F(-1)]	4281

3.443.1 Optimal result

Integrand size = 35, antiderivative size = 316

$$\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx =$$

$$\frac{(ia - b)^{5/2}(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$+ \frac{(30a^2Ab - 16Ab^3 + 5a^3B - 40ab^2B) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{8\sqrt{bd}}$$

$$+ \frac{(ia + b)^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$+ \frac{(14aAb + 5a^2B - 8b^2B) \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}{8d}$$

$$+ \frac{(2Ab + 3aB) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}}{4d}$$

$$+ \frac{bB \tan^{3/2}(c + dx)(a + b \tan(c + dx))^{3/2}}{3d}$$

output $-(I*a-b)^{(5/2)}*(A+I*B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})/d+(I*a+b)^{(5/2)}*(A-I*B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})/d+1/8*(30*A*a^2*b-16*A*b^3+5*B*a^3-40*B*a*b^2)*\operatorname{arctanh}(b^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})/d/b^{(1/2)}+1/8*(14*A*a*b+5*B*a^2-8*B*b^2)*\tan(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(1/2)}/d+1/4*(2*A*b+3*B*a)*\tan(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(3/2)}/d+1/3*b*B*\tan(d*x+c)^{(3/2)}*(a+b*\tan(d*x+c))^{(3/2)}/d$

3.443.2 Mathematica [A] (verified)

Time = 4.83 (sec) , antiderivative size = 345, normalized size of antiderivative = 1.09

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx = \frac{24\sqrt[4]{-1}(-a+ib)^{5/2}(A-iB)\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)+24\sqrt[4]{-1}(a+ib)^{5/2}(A+B\tan(c+dx))}{d}$$

input `Integrate[Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output $(24*(-1)^{(1/4)}*(-a+I*b)^{(5/2)}*(A-I*B)*\operatorname{ArcTan}(((1/4)*\sqrt{-a+I*b})*\sqrt{\tan[c+d*x]})/\sqrt{a+b*\tan[c+d*x]})+24*(-1)^{(1/4)}*(a+I*b)^{(5/2)}*(A+I*B)*\operatorname{ArcTan}(((1/4)*\sqrt{a+I*b})*\sqrt{\tan[c+d*x]})/\sqrt{a+b*\tan[c+d*x]})+3*(14*a*A*b+5*a^2*B-8*b^2*B)*\sqrt{\tan[c+d*x]}\sqrt{a+b*\tan[c+d*x]}+6*(2*A*b+3*a*B)*\sqrt{\tan[c+d*x]}*(a+b*\tan[c+d*x])^{(3/2)}+8*b*B*\tan[c+d*x]^{(3/2)}*(a+b*\tan[c+d*x])^{(3/2)}+(3*\sqrt{a}*(30*a^2*A*b-16*A*b^3+5*a^3*B-40*a*b^2*B)*\operatorname{ArcSinh}((\sqrt{b})*\sqrt{\tan[c+d*x]})/\sqrt{a})*\sqrt{1+(b*\tan[c+d*x])/a})/(\sqrt{b})*\sqrt{a+b*\tan[c+d*x]})/(24*d)$

3.443.3 Rubi [A] (verified)

Time = 2.04 (sec) , antiderivative size = 325, normalized size of antiderivative = 1.03, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 4090, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4090}$$

$$\frac{1}{3} \int \frac{3}{2} \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)} (b(2Ab+3aB) \tan^2(c+dx) + 2(Ba^2+2Aba-b^2B) \tan(c+dx) + a(2aA + bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2})) dx$$

$$\downarrow \text{27}$$

$$\frac{1}{2} \int \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)} (b(2Ab+3aB) \tan^2(c+dx) + 2(Ba^2+2Aba-b^2B) \tan(c+dx) + a(2aA + bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2})) dx$$

$$\downarrow \text{3042}$$

$$\frac{1}{2} \int \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)} (b(2Ab+3aB) \tan(c+dx)^2 + 2(Ba^2+2Aba-b^2B) \tan(c+dx) + a(2aA + bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2})) dx$$

$$\downarrow \text{4130}$$

$$\frac{1}{2} \left(\frac{\int -\frac{\sqrt{a+b \tan(c+dx)}(-b(5Ba^2+14Aba-8b^2B) \tan^2(c+dx) - 8b(Aa^2-2bBa-Ab^2) \tan(c+dx) + ab(2Ab+3aB)) dx}{2\sqrt{\tan(c+dx)}}}{2b} + \frac{(3aB+2Ab)\sqrt{\tan(c+dx)}}{2} \right) + \frac{bB \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}}{3d}$$

$$\downarrow \text{27}$$

3.443. $\int \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\frac{1}{2} \left(\frac{(3aB + 2Ab)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d} - \frac{\int \frac{\sqrt{a+b\tan(c+dx)}(-b(5Ba^2+14Aba-8b^2B)\tan^2(c+dx)-8b(Aa^2-2bBa^2-2b^2Aa^2))}{\sqrt{\tan(c+dx)}} dx}{4b} \right)$$

$$\frac{bB \tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}}{3d}$$

↓ 3042

$$\frac{1}{2} \left(\frac{(3aB + 2Ab)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d} - \frac{\int \frac{\sqrt{a+b\tan(c+dx)}(-b(5Ba^2+14Aba-8b^2B)\tan(c+dx)^2-8b(Aa^2-2bBa^2-2b^2Aa^2))}{\sqrt{\tan(c+dx)}} dx}{4b} \right)$$

$$\frac{bB \tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}}{3d}$$

↓ 4130

$$\frac{1}{2} \left(\frac{(3aB + 2Ab)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d} - \frac{\int \frac{-b(5Ba^3+30Aba^2-40b^2Ba-16Ab^3)\tan^2(c+dx)-16b(Aa^3-3bBa^2-2b^2Aa^2)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} \right)$$

$$\frac{bB \tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}}{3d}$$

↓ 27

$$\frac{1}{2} \left(\frac{(3aB + 2Ab)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d} - \frac{\frac{1}{2} \int \frac{-b(5Ba^3+30Aba^2-40b^2Ba-16Ab^3)\tan^2(c+dx)-16b(Aa^3-3bBa^2-2b^2Aa^2)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} \right)$$

$$\frac{bB \tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}}{3d}$$

↓ 3042

$$\frac{1}{2} \left(\frac{(3aB + 2Ab)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d} - \frac{\frac{1}{2} \int \frac{-b(5Ba^3+30Aba^2-40b^2Ba-16Ab^3)\tan(c+dx)^2-16b(Aa^3-3bBa^2-2b^2Aa^2)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} \right)$$

$$\frac{bB \tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}}{3d}$$

↓ 4138

3.443. $\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$

$$\frac{1}{2} \left(\frac{(3aB + 2Ab)\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}}{2d} - \frac{\int \frac{-b(5Ba^3 + 30Aba^2 - 40b^2Ba - 16Ab^3) \tan^2(c + dx) - 16b(Aa^3 - 3bBa^2 - 3Ab^2a + b^3)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}(\tan^2(c + dx) + 1)} dx}{2d} \right) - \frac{bB \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}}{3d}$$

↓ 2035

$$\frac{1}{2} \left(\frac{(3aB + 2Ab)\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}}{2d} - \frac{\int \frac{-b(5Ba^3 + 30Aba^2 - 40b^2Ba - 16Ab^3) \tan^2(c + dx) - 16b(Aa^3 - 3bBa^2 - 3Ab^2a + b^3)}{\sqrt{a + b \tan(c + dx)}(\tan^2(c + dx) + 1)} dx}{d} \right) - \frac{bB \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}}{3d}$$

↓ 2257

$$\frac{1}{2} \left(\frac{(3aB + 2Ab)\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}}{2d} - \frac{\int \left(\frac{16(b(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) - b(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)}(\tan^2(c + dx) + 1)} \right) dx}{d} \right) - \frac{bB \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}}{3d}$$

↓ 2009

$$\frac{bB \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}}{3d} + \frac{1}{2} \left(\frac{(3aB + 2Ab)\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}}{2d} - \frac{b(5a^2B + 14aAb - 8b^2B)\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}}{d} + \frac{-\sqrt{b}(5a^3B + b^3)}{d} \right)$$

input `Int[Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

```
output (b*B*Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2))/(3*d) + (((2*A*b + 3*a
*B)*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2))/(2*d) - ((8*(I*a - b)^(
5/2)*b*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[
c + d*x]]) - Sqrt[b]*(30*a^2*A*b - 16*A*b^3 + 5*a^3*B - 40*a*b^2*B)*ArcTan
h[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) - 8*b*(I*a + b)^(
5/2)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c
+ d*x]])]/d - (b*(14*a*A*b + 5*a^2*B - 8*b^2*B)*Sqrt[Tan[c + d*x]]*Sqrt[a
+ b*Tan[c + d*x]])/d)/(4*b))/2
```

3.443.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2035 Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]
```

```
rule 2257 Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol
] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a
, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4090 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4130 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

3.443.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.04 (sec) , antiderivative size = 2657119, normalized size of antiderivative = 8408.60

output too large to display

```
input int(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x)
```

$$3.443. \quad \int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

output result too large to display

3.443.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17621 vs. 2(260) = 520.

Time = 7.58 (sec) , antiderivative size = 35244, normalized size of antiderivative = 111.53

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorith="fricas")`

output Too large to include

3.443.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(1/2)*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output Timed out

3.443.7 Maxima [F]

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))dx = \int (B\tan(dx+c)+A)(b\tan(dx+c)+a)^{\frac{5}{2}}\sqrt{\tan(dx+c)}dx$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)*sqrt(tan(d*x + c)), x)`

3.443.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.443.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int \sqrt{\tan(c + dx)}(A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2} dx$$

input `int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)`

output `int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2), x)`

$$3.444 \quad \int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.444.1 Optimal result	4282
3.444.2 Mathematica [A] (verified)	4283
3.444.3 Rubi [A] (verified)	4283
3.444.4 Maple [B] (warning: unable to verify)	4287
3.444.5 Fricas [B] (verification not implemented)	4288
3.444.6 Sympy [F]	4288
3.444.7 Maxima [F]	4288
3.444.8 Giac [F(-1)]	4289
3.444.9 Mupad [F(-1)]	4289

3.444.1 Optimal result

Integrand size = 35, antiderivative size = 260

$$\int \frac{(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \frac{(ia - b)^{5/2}(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$+ \frac{\sqrt{b}(20aAb + 15a^2B - 8b^2B) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{4d}$$

$$+ \frac{(ia + b)^{5/2}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$+ \frac{b(4Ab + 7aB)\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}}{4d} + \frac{bB\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}}{2d}$$

output $(I*a-b)^{(5/2)}*(I*A-B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})/d+(I*a+b)^{(5/2)}*(I*A+B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})/d+1/4*(20*A*a*b+15*B*a^2-8*B*b^2)*\operatorname{arctanh}(b^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})*b^{(1/2)}/d+1/4*b*(4*A*b+7*B*a)*\tan(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(1/2)}/d+1/2*b*B*\tan(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(3/2)}/d$

3.444.2 Mathematica [A] (verified)

Time = 2.86 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.12

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \frac{4\sqrt[4]{-1}(-a + ib)^{5/2}(iA + B) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{\tan(c + dx)}}$$

input `Integrate[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `(4*(-1)^(1/4)*(-a + I*b)^(5/2)*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - 4*(-1)^(3/4)*(a + I*b)^(5/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + b*(4*A*b + 7*a*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]] + 2*b*B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2) + (Sqrt[a]*Sqrt[b]*(20*a*A*b + 15*a^2*B - 8*b^2*B)*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/Sqrt[a + b*Tan[c + d*x]]/(4*d)`

3.444.3 Rubi [A] (verified)

Time = 1.54 (sec) , antiderivative size = 257, normalized size of antiderivative = 0.99, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 4090, 27, 3042, 4130, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

↓ 4090

3.444. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

$$\frac{1}{2} \int \frac{\sqrt{a+b \tan(c+dx)}(b(4Ab+7aB) \tan^2(c+dx) + 4(Ba^2+2Aba-b^2B) \tan(c+dx) + a(4aA-bB))}{2\sqrt{\tan(c+dx)} \frac{bB\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{2d}} dx +$$

↓ 27

$$\frac{1}{4} \int \frac{\sqrt{a+b \tan(c+dx)}(b(4Ab+7aB) \tan^2(c+dx) + 4(Ba^2+2Aba-b^2B) \tan(c+dx) + a(4aA-bB))}{\sqrt{\tan(c+dx)} \frac{bB\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{2d}} dx +$$

↓ 3042

$$\frac{1}{4} \int \frac{\sqrt{a+b \tan(c+dx)}(b(4Ab+7aB) \tan(c+dx)^2 + 4(Ba^2+2Aba-b^2B) \tan(c+dx) + a(4aA-bB))}{\sqrt{\tan(c+dx)} \frac{bB\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{2d}} dx +$$

↓ 4130

$$\frac{1}{4} \left(\int \frac{b(15Ba^2+20Aba-8b^2B) \tan^2(c+dx) + 8(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx) + a(8Aa^2-9bBa)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)} \frac{bB\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{2d}} \right)$$

↓ 27

$$\frac{1}{4} \left(\frac{1}{2} \int \frac{b(15Ba^2+20Aba-8b^2B) \tan^2(c+dx) + 8(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx) + a(8Aa^2-9bBa)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)} \frac{bB\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{2d}} \right)$$

↓ 3042

$$\frac{1}{4} \left(\frac{1}{2} \int \frac{b(15Ba^2+20Aba-8b^2B) \tan(c+dx)^2 + 8(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx) + a(8Aa^2-9bBa)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)} \frac{bB\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{2d}} \right)$$

↓ 4138

$$\frac{1}{4} \left(\frac{\int \frac{b(15Ba^2+20Aba-8b^2B) \tan^2(c+dx)+8(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx)+a(8Aa^2-9bBa-4Ab^2)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} d \tan(c+dx)}{2d} + \frac{b(7aB + \dots)}{\dots} \right)$$

\downarrow 2035

$$\frac{1}{4} \left(\frac{\int \frac{b(15Ba^2+20Aba-8b^2B) \tan^2(c+dx)+8(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx)+a(8Aa^2-9bBa-4Ab^2)}{\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} d \sqrt{\tan(c+dx)}}{d} + \frac{b(7aB + \dots)}{\dots} \right)$$

\downarrow 2257

$$\frac{1}{4} \left(\frac{\int \left(\frac{b(15Ba^2+20Aba-8b^2B)}{\sqrt{a+b \tan(c+dx)}} + \frac{8(Aa^3-3bBa^2-3Ab^2a+b^3B+(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} \right) d \sqrt{\tan(c+dx)}}{d} + \frac{b(7aB + \dots)}{\dots} \right)$$

\downarrow 2009

$$\frac{1}{4} \left(\frac{bB \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{2d} + \frac{b(7aB + 4Ab) \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}}{d} + \frac{\sqrt{b}(15a^2B + 20aAb - 8b^2B) \operatorname{arctanh}\left(\frac{\sqrt{b} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) + 4 \dots}{\dots} \right)$$

input `Int[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `(b*B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2))/(2*d) + ((4*(I*a - b)^(5/2)*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])] + Sqrt[b]*(20*a*A*b + 15*a^2*B - 8*b^2*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])] + 4*(I*a + b)^(5/2)*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/d + (b*(4*A*b + 7*a*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d/4`

3.444.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2035 `Int[(F_x_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[Fx, x]`
- rule 2257 `Int[(P_x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4130 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
) + (f_.)*(x_)^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

3.444.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.03 (sec) , antiderivative size = 2654895, normalized size of antiderivative = 10211.13

output too large to display

```
input int((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x)
```

```
output result too large to display
```

3.444.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17603 vs. $2(210) = 420$.

Time = 7.66 (sec) , antiderivative size = 35212, normalized size of antiderivative = 135.43

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algo
rithm="fricas")`

output Too large to include

3.444.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2}}{\sqrt{\tan(c + dx)}} dx$$

input `integrate((a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(5/2)/sqrt(tan(c + d*x
)), x)`

3.444.7 Maxima [F]

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^{5/2}}{\sqrt{\tan(dx + c)}} dx$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)/sqrt(tan(d*x + c
)), x)`

3.444. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$

3.444.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algo
rithm="giac")`

output `Timed out`

3.444.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2}}{\sqrt{\tan(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(1/2),x
)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(1/2),
x)`

3.445
$$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.445.1 Optimal result 4290
 3.445.2 Mathematica [A] (verified) 4291
 3.445.3 Rubi [A] (verified) 4291
 3.445.4 Maple [B] (warning: unable to verify) 4295
 3.445.5 Fricas [B] (verification not implemented) 4296
 3.445.6 Sympy [F] 4296
 3.445.7 Maxima [F(-1)] 4296
 3.445.8 Giac [F(-1)] 4297
 3.445.9 Mupad [F(-1)] 4297

3.445.1 Optimal result

Integrand size = 35, antiderivative size = 241

$$\int \frac{(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \frac{(ia - b)^{5/2}(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$+ \frac{b^{3/2}(2Ab + 5aB) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} - \frac{(ia + b)^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$+ \frac{b(2aA + bB) \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}{d} - \frac{2aA(a + b \tan(c + dx))^{3/2}}{d \sqrt{\tan(c + dx)}}$$

output

```
(I*a-b)^(5/2)*(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)^(1/2))/d+b^(3/2)*(2*A*b+5*B*a)*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)^(1/2))/d-(I*a+b)^(5/2)*(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)^(1/2))/d+b*(2*A*a+B*b)*tan(d*x+c)^(1/2)*(a+b*tan(d*x+c)^(1/2))/d-2*a*A*(a+b*tan(d*x+c))^(3/2)/d/tan(d*x+c)^(1/2)
```

3.445.2 Mathematica [A] (verified)

Time = 2.71 (sec) , antiderivative size = 358, normalized size of antiderivative = 1.49

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \frac{(-1)^{3/4} (-a + ib)^{5/2} (iA + B) \arctan\left(\frac{\sqrt[4]{-1} \sqrt{-a + ib} \sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{\tan^{3/2}(c + dx)}$$

input `Integrate[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `((-1)^(3/4)*(-a + I*b)^(5/2)*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]] - (-1)^(1/4)*(a + I*b)^(5/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]] - ((a - I*b)^2*(A - I*B)*Sqrt[a + b*Tan[c + d*x]]/Sqrt[Tan[c + d*x]] - ((a + I*b)^2*(A + I*B)*Sqrt[a + b*Tan[c + d*x]]/Sqrt[Tan[c + d*x]] + (b*B*(a + b*Tan[c + d*x])^(3/2))/Sqrt[Tan[c + d*x]] + (b*(2*A*b + 5*a*B)*Sqrt[a + b*Tan[c + d*x]]*((Sqrt[b]*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/Sqrt[a]))/Sqrt[a] - Sqrt[1 + (b*Tan[c + d*x])/a]/Sqrt[Tan[c + d*x]])/Sqrt[1 + (b*Tan[c + d*x])/a])/d`

3.445.3 Rubi [A] (verified)

Time = 1.51 (sec) , antiderivative size = 237, normalized size of antiderivative = 0.98, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 4088, 27, 3042, 4130, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx$$

↓ 4088

3.445. $\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx$

$$2 \int \frac{\sqrt{a + b \tan(c + dx)} (b(2aA + bB) \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(4Ab + aB))}{2\sqrt{\tan(c + dx)} \frac{2aA(a + b \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}}} dx -$$

↓ 27

$$\int \frac{\sqrt{a + b \tan(c + dx)} (b(2aA + bB) \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(4Ab + aB))}{\sqrt{\tan(c + dx)} \frac{2aA(a + b \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}}} dx -$$

↓ 3042

$$\int \frac{\sqrt{a + b \tan(c + dx)} (b(2aA + bB) \tan(c + dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(4Ab + aB))}{\sqrt{\tan(c + dx)} \frac{2aA(a + b \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}}} dx -$$

↓ 4130

$$\int \frac{b^2(2Ab + 5aB) \tan^2(c + dx) - 2(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx) + a(2Ba^2 + 6Aba - b^2B)}{2\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)} \frac{b(2aA + bB) \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}{d} - \frac{2aA(a + b \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}}} dx +$$

↓ 27

$$\frac{1}{2} \int \frac{b^2(2Ab + 5aB) \tan^2(c + dx) - 2(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx) + a(2Ba^2 + 6Aba - b^2B)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)} \frac{b(2aA + bB) \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}{d} - \frac{2aA(a + b \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}}} dx +$$

↓ 3042

$$\frac{1}{2} \int \frac{b^2(2Ab + 5aB) \tan(c + dx)^2 - 2(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx) + a(2Ba^2 + 6Aba - b^2B)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)} \frac{b(2aA + bB) \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}{d} - \frac{2aA(a + b \tan(c + dx))^{3/2}}{d\sqrt{\tan(c + dx)}}} dx +$$

↓ 4138

3.445. $\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx$

$$\begin{aligned}
& \int \frac{b^2(2Ab+5aB)\tan^2(c+dx)-2(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx)+a(2Ba^2+6Aba-b^2B)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} d\tan(c+dx) \\
& \quad + \frac{2d}{b(2aA+bB)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} - \frac{2aA(a+b\tan(c+dx))^{3/2}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow \text{2035} \\
& \int \frac{b^2(2Ab+5aB)\tan^2(c+dx)-2(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx)+a(2Ba^2+6Aba-b^2B)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} d\sqrt{\tan(c+dx)} \\
& \quad + \frac{d}{b(2aA+bB)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} - \frac{2aA(a+b\tan(c+dx))^{3/2}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow \text{2257} \\
& \int \left(\frac{(2Ab+5aB)b^2}{\sqrt{a+b\tan(c+dx)}} + \frac{2(Ba^3+3Aba^2-3b^2Ba-Ab^3-(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} \right) d\sqrt{\tan(c+dx)} \\
& \quad + \frac{d}{b(2aA+bB)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} - \frac{2aA(a+b\tan(c+dx))^{3/2}}{d\sqrt{\tan(c+dx)}} \\
& \quad \downarrow \text{2009} \\
& \frac{(-b+ia)^{5/2}(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) + b^{3/2}(5aB+2Ab)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) - (b+ia)^{5/2}(A-iB)}{b(2aA+bB)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} - \frac{2aA(a+b\tan(c+dx))^{3/2}}{d\sqrt{\tan(c+dx)}}
\end{aligned}$$

input `Int[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2), x]`

output `((I*a - b)^(5/2)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + b^(3/2)*(2*A*b + 5*a*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - (I*a + b)^(5/2)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]/d + (b*(2*a*A + b*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d - (2*a*A*(a + b*Tan[c + d*x])^(3/2))/(d*Sqrt[Tan[c + d*x]])`

3.445. $\int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\tan^{3/2}(c+dx)} dx$

3.445.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2035 `Int[(F_x_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[Fx, x]`
- rule 2257 `Int[(P_x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4138 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.445.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.08 (sec) , antiderivative size = 2653772, normalized size of antiderivative = 11011.50

output too large to display

input `int((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x)`

output `result too large to display`

3.445.
$$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.445.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17614 vs. $2(197) = 394$.

Time = 7.59 (sec) , antiderivative size = 35230, normalized size of antiderivative = 146.18

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algo
rithm="fricas")`

output Too large to include

3.445.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2}}{\tan^{3/2}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(5/2)/tan(c + d*x)**(3
/2), x)`

3.445.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algo
rithm="maxima")`

output Timed out

3.445. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{3/2}(c+dx)} dx$

3.445.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algo
rithm="giac")`

output `Timed out`

3.445.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{3/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2}}{\tan(c + dx)^{3/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(3/2),x
)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(3/2),
x)`

3.446
$$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.446.1 Optimal result	4298
3.446.2 Mathematica [C] (verified)	4299
3.446.3 Rubi [A] (verified)	4299
3.446.4 Maple [B] (warning: unable to verify)	4303
3.446.5 Fricas [B] (verification not implemented)	4304
3.446.6 Sympy [F]	4304
3.446.7 Maxima [F]	4304
3.446.8 Giac [F(-1)]	4305
3.446.9 Mupad [F(-1)]	4305

3.446.1 Optimal result

Integrand size = 35, antiderivative size = 240

$$\int \frac{(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx =$$

$$-\frac{(ia - b)^{5/2}(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} + \frac{2b^{5/2}B \operatorname{Arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$-\frac{(ia + b)^{5/2}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$-\frac{2a(2Ab + aB)\sqrt{a + b \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} - \frac{2aA(a + b \tan(c + dx))^{3/2}}{3d \tan^{\frac{3}{2}}(c + dx)}$$

```
output -(I*a-b)^(5/2)*(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d+2*b^(5/2)*B*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d-(I*a+b)^(5/2)*(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d-2*a*(2*A*b+B*a)*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(1/2)-2/3*a*A*(a+b*tan(d*x+c))^(3/2)/d/tan(d*x+c)^(3/2)
```

3.446.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 4.47 (sec) , antiderivative size = 417, normalized size of antiderivative = 1.74

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \frac{iab(A + iB) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, -\frac{3}{2}, -\frac{1}{2}, -\frac{b \tan(c + dx)}{a}\right)}{\tan^{5/2}(c + dx)}$$

input `Integrate[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(I*a*b*(A + I*B)*Hypergeometric2F1[-3/2, -3/2, -1/2, -((b*Tan[c + d*x])/a)]*Sqrt[a + b*Tan[c + d*x]] - a*b*(I*A + B)*Hypergeometric2F1[-3/2, -3/2, -1/2, -((b*Tan[c + d*x])/a)]*Sqrt[a + b*Tan[c + d*x]] + (I*a + b)*(A - I*B)*Sqrt[1 + (b*Tan[c + d*x])/a]*(3*(-1)^(1/4)*(-a + I*b)^(3/2)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]*Tan[c + d*x]^(3/2) + (I*a + (-3*a + (4*I)*b)*Tan[c + d*x])*Sqrt[a + b*Tan[c + d*x]]) + (I*a - b)*(A + I*B)*Sqrt[1 + (b*Tan[c + d*x])/a]*(3*(-1)^(1/4)*(a + I*b)^(3/2)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(3/2) + (I*a + (3*a + (4*I)*b)*Tan[c + d*x])*Sqrt[a + b*Tan[c + d*x]]))/(3*d*Tan[c + d*x]^(3/2)*Sqrt[1 + (b*Tan[c + d*x])/a])`

3.446.3 Rubi [A] (verified)

Time = 1.43 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx$$

3.446. $\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx$

↓ 4088

$$\frac{2}{3} \int \frac{3\sqrt{a+b \tan(c+dx)}(b^2 B \tan^2(c+dx) - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2Ab + aB))}{\frac{2 \tan^{\frac{3}{2}}(c+dx)}{2aA(a+b \tan(c+dx))^{3/2}} - \frac{3d \tan^{\frac{3}{2}}(c+dx)}{3d \tan^{\frac{3}{2}}(c+dx)}} dx -$$

↓ 27

$$\int \frac{\sqrt{a+b \tan(c+dx)}(b^2 B \tan^2(c+dx) - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2Ab + aB))}{\frac{\tan^{\frac{3}{2}}(c+dx)}{2aA(a+b \tan(c+dx))^{3/2}} - \frac{3d \tan^{\frac{3}{2}}(c+dx)}{3d \tan^{\frac{3}{2}}(c+dx)}} dx -$$

↓ 3042

$$\int \frac{\sqrt{a+b \tan(c+dx)}(b^2 B \tan(c+dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2Ab + aB))}{\frac{\tan(c+dx)^{3/2}}{2aA(a+b \tan(c+dx))^{3/2}} - \frac{3d \tan^{\frac{3}{2}}(c+dx)}{3d \tan^{\frac{3}{2}}(c+dx)}} dx -$$

↓ 4128

$$2 \int -\frac{-B \tan^2(c+dx)b^3 + a(Aa^2 - 3bBa - 3Ab^2) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)}{\frac{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}}{2a(aB + 2Ab)\sqrt{a+b \tan(c+dx)}} - \frac{2aA(a+b \tan(c+dx))^{3/2}}{3d \tan^{\frac{3}{2}}(c+dx)}} dx -$$

↓ 27

$$- \int \frac{-B \tan^2(c+dx)b^3 + a(Aa^2 - 3bBa - 3Ab^2) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)}{\frac{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}}{2a(aB + 2Ab)\sqrt{a+b \tan(c+dx)}} - \frac{2aA(a+b \tan(c+dx))^{3/2}}{3d \tan^{\frac{3}{2}}(c+dx)}} dx -$$

↓ 3042

$$- \int \frac{-B \tan(c+dx)^2 b^3 + a(Aa^2 - 3bBa - 3Ab^2) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)}{\frac{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}}{2a(aB + 2Ab)\sqrt{a+b \tan(c+dx)}} - \frac{2aA(a+b \tan(c+dx))^{3/2}}{3d \tan^{\frac{3}{2}}(c+dx)}} dx -$$

↓ 4138

3.446. $\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$

$$\begin{aligned}
& \frac{\int \frac{-B \tan^2(c+dx)b^3+a(Aa^2-3bBa-3Ab^2)+(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} d \tan(c+dx)}{\frac{2a(aB+2Ab)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2aA(a+b \tan(c+dx))^{3/2}}{3d \tan^{3/2}(c+dx)}} \\
& \quad \downarrow \text{2035} \\
& \frac{2 \int \frac{-B \tan^2(c+dx)b^3+a(Aa^2-3bBa-3Ab^2)+(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} d \sqrt{\tan(c+dx)}}{\frac{2a(aB+2Ab)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2aA(a+b \tan(c+dx))^{3/2}}{3d \tan^{3/2}(c+dx)}} \\
& \quad \downarrow \text{2257} \\
& \frac{2 \int \left(\frac{Aa^3-3bBa^2-3Ab^2a+b^3B+(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} - \frac{b^3B}{\sqrt{a+b \tan(c+dx)}} \right) d \sqrt{\tan(c+dx)}}{\frac{2a(aB+2Ab)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2aA(a+b \tan(c+dx))^{3/2}}{3d \tan^{3/2}(c+dx)}} \\
& \quad \downarrow \text{2009} \\
& \frac{2 \left(\frac{1}{2}(-b+ia)^{5/2}(-B+iA) \arctan \left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right) + \frac{1}{2}(b+ia)^{5/2}(B+iA) \operatorname{arctanh} \left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right) - b^{5/2} \right)}{\frac{2a(aB+2Ab)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2aA(a+b \tan(c+dx))^{3/2}}{3d \tan^{3/2}(c+dx)}}
\end{aligned}$$

input `Int[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2), x]`

output `(-2*(((I*a - b)^(5/2)*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/2 - b^(5/2)*B*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) + ((I*a + b)^(5/2)*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/2))/d - (2*a*(2*A*b + a*B)*Sqrt[a + b*Tan[c + d*x]]/(d*Sqrt[Tan[c + d*x]]) - (2*a*A*(a + b*Tan[c + d*x])^(3/2))/(3*d*Tan[c + d*x]^(3/2))`

3.446. $\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{5/2}(c+dx)} dx$

3.446.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2035 `Int[(F_x_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[Fx, x]`
- rule 2257 `Int[(P_x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

```
rule 4128 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

3.446.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.02 (sec) , antiderivative size = 2654078, normalized size of antiderivative = 11058.66

output too large to display

```
input int((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x)
```

```
output result too large to display
```

3.446.
$$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{5/2}(c+dx)} dx$$

3.446.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17675 vs. $2(194) = 388$.

Time = 7.33 (sec) , antiderivative size = 35349, normalized size of antiderivative = 147.29

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorith="fricas")`

output Too large to include

3.446.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2}}{\tan^{5/2}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(5/2)/tan(c + d*x)**(5/2), x)`

3.446.7 Maxima [F]

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^{5/2}}{\tan(dx + c)^{5/2}} dx$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorith="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)/tan(d*x + c)^(5/2), x)`

3.446. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{5/2}(c+dx)} dx$

3.446.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algo
rithm="giac")`

output `Timed out`

3.446.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{5/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2}}{\tan(c + dx)^{5/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(5/2),x
)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(5/2),
x)`

$$3.447 \quad \int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

3.447.1 Optimal result	4306
3.447.2 Mathematica [A] (verified)	4307
3.447.3 Rubi [A] (verified)	4307
3.447.4 Maple [B] (warning: unable to verify)	4313
3.447.5 Fricas [B] (verification not implemented)	4313
3.447.6 Sympy [F(-1)]	4314
3.447.7 Maxima [F(-1)]	4314
3.447.8 Giac [F(-1)]	4314
3.447.9 Mupad [F(-1)]	4315

3.447.1 Optimal result

Integrand size = 35, antiderivative size = 247

$$\int \frac{(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{7}{2}}(c + dx)} dx =$$

$$-\frac{(ia - b)^{5/2}(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$+ \frac{(ia + b)^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} - \frac{2a(8Ab + 5aB)\sqrt{a + b \tan(c + dx)}}{15d \tan^{\frac{3}{2}}(c + dx)}$$

$$+ \frac{2(15a^2A - 23Ab^2 - 35abB)\sqrt{a + b \tan(c + dx)}}{15d\sqrt{\tan(c + dx)}} - \frac{2aA(a + b \tan(c + dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c + dx)}$$

output

```
-(I*a-b)^(5/2)*(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d+(I*a+b)^(5/2)*(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d+2/15*(15*A*a^2-23*A*b^2-35*B*a*b)*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(1/2)-2/15*a*(8*A*b+5*B*a)*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(3/2)-2/5*a*A*(a+b*tan(d*x+c))^(3/2)/d/tan(d*x+c)^(5/2)
```

3.447.2 Mathematica [A] (verified)

Time = 2.29 (sec) , antiderivative size = 321, normalized size of antiderivative = 1.30

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \frac{60\sqrt[4]{-1} \left((-a + ib)^{5/2} (A - iB) \arctan \left(\frac{\sqrt[4]{-1} \sqrt{-a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right) \right)}{\tan^{7/2}(c + dx)}$$

input `Integrate[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

output `(60*(-1)^(1/4)*((-a + I*b)^(5/2)*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]] + (a + I*b)^(5/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(5/2) + 15*b*(-2*A*b + a*B)*Sqrt[a + b*Tan[c + d*x]] - 3*(8*a^2*A - 10*A*b^2 - 15*a*b*B)*Sqrt[a + b*Tan[c + d*x]] - 4*(22*a*A*b + 10*a^2*B - 15*b^2*B)*Tan[c + d*x]*Sqrt[a + b*Tan[c + d*x]] + 8*(15*a^2*A - 23*A*b^2 - 35*a*b*B)*Tan[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]] - 60*b*B*(a + b*Tan[c + d*x])^(3/2))/(60*d*Tan[c + d*x]^(5/2))`

3.447.3 Rubi [A] (verified)

Time = 1.89 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.15, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan(c + dx)^{7/2}} dx$$

↓ 4088

3.447. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{7/2}(c+dx)} dx$

$$\frac{2}{5} \int \frac{\sqrt{a+b \tan(c+dx)}(-b(2aA-5bB) \tan^2(c+dx)-5(Aa^2-2bBa-Ab^2) \tan(c+dx)+a(8Ab+5aB))}{2 \tan^{\frac{5}{2}}(c+dx)} dx$$

$$\frac{2aA(a+b \tan(c+dx))^{\frac{3}{2}}}{5d \tan^{\frac{5}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{5} \int \frac{\sqrt{a+b \tan(c+dx)}(-b(2aA-5bB) \tan^2(c+dx)-5(Aa^2-2bBa-Ab^2) \tan(c+dx)+a(8Ab+5aB))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

$$\frac{2aA(a+b \tan(c+dx))^{\frac{3}{2}}}{5d \tan^{\frac{5}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{5} \int \frac{\sqrt{a+b \tan(c+dx)}(-b(2aA-5bB) \tan(c+dx)^2-5(Aa^2-2bBa-Ab^2) \tan(c+dx)+a(8Ab+5aB))}{\tan(c+dx)^{\frac{5}{2}}} dx$$

$$\frac{2aA(a+b \tan(c+dx))^{\frac{3}{2}}}{5d \tan^{\frac{5}{2}}(c+dx)}$$

↓ 4128

$$\frac{1}{5} \left(\frac{2}{3} \int -\frac{b(10Ba^2+22Aba-15b^2B) \tan^2(c+dx)+15(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx)+a(15Aa^2-15Ab^2)}{2 \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx \right)$$

$$\frac{2aA(a+b \tan(c+dx))^{\frac{3}{2}}}{5d \tan^{\frac{5}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{5} \left(-\frac{1}{3} \int \frac{b(10Ba^2+22Aba-15b^2B) \tan^2(c+dx)+15(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx)+a(15Aa^2-15Ab^2)}{\tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx \right)$$

$$\frac{2aA(a+b \tan(c+dx))^{\frac{3}{2}}}{5d \tan^{\frac{5}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{5} \left(-\frac{1}{3} \int \frac{b(10Ba^2+22Aba-15b^2B) \tan(c+dx)^2+15(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx)+a(15Aa^2-15Ab^2)}{\tan(c+dx)^{\frac{3}{2}} \sqrt{a+b \tan(c+dx)}} dx \right)$$

$$\frac{2aA(a+b \tan(c+dx))^{\frac{3}{2}}}{5d \tan^{\frac{5}{2}}(c+dx)}$$

↓ 4132

3.447. $\int \frac{(a+b \tan(c+dx))^{\frac{5}{2}}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
& \frac{1}{5} \left(\frac{1}{3} \left(\frac{2 \int -\frac{15(a(Ba^3+3Aba^2-3b^2Ba-Ab^3))-a(Aa^3-3bBa^2-3Ab^2a+b^3B) \tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} + \frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) \right. \\
& \qquad \qquad \qquad \frac{2aA(a+b \tan(c+dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \qquad \qquad \qquad \downarrow \text{27} \\
& \frac{1}{5} \left(\frac{1}{3} \left(\frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{15 \int \frac{a(Ba^3+3Aba^2-3b^2Ba-Ab^3))-a(Aa^3-3bBa^2-3Ab^2a+b^3B) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}}}{a} \right) \right. \\
& \qquad \qquad \qquad \frac{2aA(a+b \tan(c+dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \qquad \qquad \qquad \downarrow \text{3042} \\
& \frac{1}{5} \left(\frac{1}{3} \left(\frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{15 \int \frac{a(Ba^3+3Aba^2-3b^2Ba-Ab^3))-a(Aa^3-3bBa^2-3Ab^2a+b^3B) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}}}{a} \right) \right. \\
& \qquad \qquad \qquad \frac{2aA(a+b \tan(c+dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c+dx)} \\
& \qquad \qquad \qquad \downarrow \text{4099} \\
& \qquad \qquad \qquad -\frac{2aA(a+b \tan(c+dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c+dx)} + \\
& \frac{1}{5} \left(-\frac{2a(5aB+8Ab)\sqrt{a+b \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{1}{3} \left(\frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{15(\frac{1}{2}a(a-ib))}{a} \right) \right. \\
& \qquad \qquad \qquad \downarrow \text{3042} \\
& \qquad \qquad \qquad -\frac{2aA(a+b \tan(c+dx))^{3/2}}{5d \tan^{\frac{5}{2}}(c+dx)} + \\
& \frac{1}{5} \left(-\frac{2a(5aB+8Ab)\sqrt{a+b \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} + \frac{1}{3} \left(\frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{15(\frac{1}{2}a(a-ib))}{a} \right) \right. \\
& \qquad \qquad \qquad \downarrow \text{4098}
\end{aligned}$$

3.447. $\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$

$$\begin{aligned}
 & -\frac{2aA(a+b\tan(c+dx))^{3/2}}{5d\tan^{5/2}(c+dx)} + \\
 \frac{1}{5} & \left(-\frac{2a(5aB+8Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)} + \frac{1}{3} \left(\frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - 15 \left(\frac{a(a-ib)^3(B-ib)}{d^2} \right) \right) \right) \\
 & \quad \downarrow 104 \\
 & -\frac{2aA(a+b\tan(c+dx))^{3/2}}{5d\tan^{5/2}(c+dx)} + \\
 \frac{1}{5} & \left(-\frac{2a(5aB+8Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)} + \frac{1}{3} \left(\frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - 15 \left(\frac{a(a-ib)^3(B-ib)}{d^2} \right) \right) \right) \\
 & \quad \downarrow 216 \\
 & -\frac{2aA(a+b\tan(c+dx))^{3/2}}{5d\tan^{5/2}(c+dx)} + \\
 \frac{1}{5} & \left(-\frac{2a(5aB+8Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)} + \frac{1}{3} \left(\frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - 15 \left(\frac{a(a-ib)^3(B-ib)}{d^2} \right) \right) \right) \\
 & \quad \downarrow 219 \\
 & -\frac{2aA(a+b\tan(c+dx))^{3/2}}{5d\tan^{5/2}(c+dx)} + \\
 \frac{1}{5} & \left(-\frac{2a(5aB+8Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)} + \frac{1}{3} \left(\frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - 15 \left(\frac{a(a-ib)^3(B-ib)}{d^2} \right) \right) \right)
 \end{aligned}$$

input `Int[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(7/2),x]`

3.447. $\int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\tan^{7/2}(c+dx)} dx$

```
output (-2*a*A*(a + b*Tan[c + d*x])^(3/2))/(5*d*Tan[c + d*x]^(5/2)) + ((-2*a*(8*A
*b + 5*a*B)*Sqrt[a + b*Tan[c + d*x]]/(3*d*Tan[c + d*x]^(3/2)) + ((-15*(-(
(a*(a + I*b)^3*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a
+ b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d)) + (a*(a - I*b)^3*(I*A + B)*ArcTanh[
(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a +
b]*d)))/a + (2*(15*a^2*A - 23*A*b^2 - 35*a*b*B)*Sqrt[a + b*Tan[c + d*x]]/
(d*Sqrt[Tan[c + d*x]]))/3)/5
```

3.447.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 104 Int[(((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)))/((e_.) + (f_.)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

3.447.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.02 (sec) , antiderivative size = 2652302, normalized size of antiderivative = 10738.06

output too large to display

```
input int((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x)
```

```
output result too large to display
```

3.447.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18802 vs. $2(201) = 402$.

Time = 5.06 (sec) , antiderivative size = 18802, normalized size of antiderivative = 76.12

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algo
rithm="fracas")
```

```
output Too large to include
```

3.447. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{7/2}(c+dx)} dx$

3.447.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(7/2),x)`

output `Timed out`

3.447.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="maxima")`

output `Timed out`

3.447.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(7/2),x, algorithm="giac")`

output `Timed out`

3.447.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{7/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2}}{\tan(c + dx)^{7/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(7/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(7/2),x)`

3.448
$$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

3.448.1 Optimal result 4316
 3.448.2 Mathematica [A] (verified) 4317
 3.448.3 Rubi [A] (verified) 4317
 3.448.4 Maple [B] (warning: unable to verify) 4324
 3.448.5 Fricas [B] (verification not implemented) 4324
 3.448.6 Sympy [F(-1)] 4325
 3.448.7 Maxima [F] 4325
 3.448.8 Giac [F(-1)] 4325
 3.448.9 Mupad [F(-1)] 4326

3.448.1 Optimal result

Integrand size = 35, antiderivative size = 309

$$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx = \frac{(ia-b)^{5/2}(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$+ \frac{(ia+b)^{5/2}(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} - \frac{2a(10Ab+7aB)\sqrt{a+b \tan(c+dx)}}{35d \tan^{\frac{5}{2}}(c+dx)}$$

$$+ \frac{2(35a^2A-45Ab^2-77abB)\sqrt{a+b \tan(c+dx)}}{105d \tan^{\frac{3}{2}}(c+dx)}$$

$$+ \frac{2(245a^2Ab-15Ab^3+105a^3B-161ab^2B)\sqrt{a+b \tan(c+dx)}}{105ad\sqrt{\tan(c+dx)}}$$

$$- \frac{2aA(a+b \tan(c+dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

output

```
(I*a-b)^(5/2)*(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d+(I*a+b)^(5/2)*(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d+2/105*(245*A*a^2*b-15*A*b^3+105*B*a^3-161*B*a*b^2)*(a+b*tan(d*x+c))^(1/2)/a/d/tan(d*x+c)^(1/2)-2/35*a*(10*A*b+7*B*a)*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(5/2)+2/105*(35*A*a^2-45*A*b^2-77*B*a*b)*(a+b*tan(d*x+c))^(1/2)/d/tan(d*x+c)^(3/2)-2/7*a*A*(a+b*tan(d*x+c))^(3/2)/d/tan(d*x+c)^(7/2)
```

3.448.2 Mathematica [A] (verified)

Time = 6.80 (sec) , antiderivative size = 381, normalized size of antiderivative = 1.23

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \frac{420\sqrt[4]{-1}a \left((-a + ib)^{5/2} (iA + B) \arctan \left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{a+b\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}} \right) \right)}{\tan^{9/2}(c + dx)}$$

input `Integrate[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2),x]`

output `(420*(-1)^(1/4)*a*((-a + I*b)^(5/2)*(I*A + B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]) + (a + I*b)^(5/2)*((-I)*A + B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(7/2) - 35*a*b*(4*A*b + a*B)*Sqrt[a + b*Tan[c + d*x]] - 5*a*(24*a^2*A - 28*A*b^2 - 49*a*b*B)*Sqrt[a + b*Tan[c + d*x]] - 6*a*(60*a*A*b + 28*a^2*B - 35*b^2*B)*Tan[c + d*x]*Sqrt[a + b*Tan[c + d*x]] + 8*a*(35*a^2*A - 45*A*b^2 - 77*a*b*B)*Tan[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]] + 8*(245*a^2*A*b - 15*A*b^3 + 105*a^3*B - 161*a*b^2*B)*Tan[c + d*x]^3*Sqrt[a + b*Tan[c + d*x]] - 210*a*b*B*(a + b*Tan[c + d*x])^(3/2))/(420*a*d*Tan[c + d*x]^(7/2))`

3.448.3 Rubi [A] (verified)Time = 2.39 (sec) , antiderivative size = 350, normalized size of antiderivative = 1.13, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.543$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan(c + dx)^{9/2}} dx$$

↓ 4088

3.448. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{9/2}(c+dx)} dx$

$$\frac{2}{7} \int \frac{\sqrt{a+b \tan(c+dx)}(-b(4aA-7bB) \tan^2(c+dx)-7(Aa^2-2bBa-Ab^2) \tan(c+dx)+a(10Ab+7aB))}{2 \tan^{\frac{7}{2}}(c+dx)} dx$$

$$\frac{2aA(a+b \tan(c+dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{7} \int \frac{\sqrt{a+b \tan(c+dx)}(-b(4aA-7bB) \tan^2(c+dx)-7(Aa^2-2bBa-Ab^2) \tan(c+dx)+a(10Ab+7aB))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

$$\frac{2aA(a+b \tan(c+dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \int \frac{\sqrt{a+b \tan(c+dx)}(-b(4aA-7bB) \tan(c+dx)^2-7(Aa^2-2bBa-Ab^2) \tan(c+dx)+a(10Ab+7aB))}{\tan(c+dx)^{7/2}} dx$$

$$\frac{2aA(a+b \tan(c+dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 4128

$$\frac{1}{7} \left(\frac{2}{5} \int -\frac{b(28Ba^2+60Aba-35b^2B) \tan^2(c+dx)+35(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx)+a(35Aa^2-35Ab^2)}{2 \tan^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx \right)$$

$$\frac{2aA(a+b \tan(c+dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{7} \left(-\frac{1}{5} \int \frac{b(28Ba^2+60Aba-35b^2B) \tan^2(c+dx)+35(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx)+a(35Aa^2-35Ab^2)}{\tan^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx \right)$$

$$\frac{2aA(a+b \tan(c+dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(-\frac{1}{5} \int \frac{b(28Ba^2+60Aba-35b^2B) \tan(c+dx)^2+35(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx)+a(35Aa^2-35Ab^2)}{\tan(c+dx)^{5/2} \sqrt{a+b \tan(c+dx)}} dx \right)$$

$$\frac{2aA(a+b \tan(c+dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c+dx)}$$

↓ 4132

3.448. $\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2 \int -\frac{-2ab(35Aa^2 - 77bBa - 45Ab^2) \tan^2(c+dx) - 105a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) + a(105Ba^3 + 245Aba^2 - 161b^2Ba - 15Ab^3)}{2 \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}}{3a} \right) \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2(35a^2A - 77abB - 45Ab^2) \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{\int -\frac{-2ab(35Aa^2 - 77bBa - 45Ab^2) \tan^2(c+dx) - 105a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) + a(105Ba^3 + 245Aba^2 - 161b^2Ba - 15Ab^3)}{\tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}}{3a} \right) \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2(35a^2A - 77abB - 45Ab^2) \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{\int -\frac{-2ab(35Aa^2 - 77bBa - 45Ab^2) \tan(c+dx)^2 - 105a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) + a(105Ba^3 + 245Aba^2 - 161b^2Ba - 15Ab^3)}{\tan(c+dx)^{3/2} \sqrt{a+b \tan(c+dx)}}}{3a} \right) \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 4132

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2(35a^2A - 77abB - 45Ab^2) \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2 \int \frac{105((Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)a^2 + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx) + a(105Ba^3 + 245Aba^2 - 161b^2Ba - 15Ab^3))}{2 \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}}}{a} \right) \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{7} \left(\frac{1}{5} \left(\frac{2(35a^2A - 77abB - 45Ab^2) \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{105 \int \frac{(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)a^2 + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx) + a(105Ba^3 + 245Aba^2 - 161b^2Ba - 15Ab^3)}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}}}{a} \right) \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c + dx)}$$

↓ 3042

3.448. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \frac{1}{7} \left(\frac{1}{5} \left(\frac{2(35a^2A - 77abB - 45Ab^2) \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{105 \int \frac{(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)a^2 + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx}{a} \right) \right. \\
 & \qquad \qquad \qquad \frac{2aA(a + b \tan(c + dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c + dx)} \\
 & \qquad \qquad \qquad \downarrow 4099 \\
 & \qquad \qquad \qquad - \frac{2aA(a + b \tan(c + dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c + dx)} + \\
 & \frac{1}{7} \left(- \frac{2a(7aB + 10Ab) \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} + \frac{1}{5} \left(\frac{2(35a^2A - 77abB - 45Ab^2) \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2(105a^3B + 24a^2A)}{a} \right) \right. \\
 & \qquad \qquad \qquad \downarrow 3042 \\
 & \qquad \qquad \qquad - \frac{2aA(a + b \tan(c + dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c + dx)} + \\
 & \frac{1}{7} \left(- \frac{2a(7aB + 10Ab) \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} + \frac{1}{5} \left(\frac{2(35a^2A - 77abB - 45Ab^2) \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2(105a^3B + 24a^2A)}{a} \right) \right. \\
 & \qquad \qquad \qquad \downarrow 4098 \\
 & \qquad \qquad \qquad - \frac{2aA(a + b \tan(c + dx))^{3/2}}{7d \tan^{\frac{7}{2}}(c + dx)} + \\
 & \frac{1}{7} \left(- \frac{2a(7aB + 10Ab) \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} + \frac{1}{5} \left(\frac{2(35a^2A - 77abB - 45Ab^2) \sqrt{a + b \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} - \frac{2(105a^3B + 24a^2A)}{a} \right) \right. \\
 & \qquad \qquad \qquad \downarrow 104
 \end{aligned}$$

3.448. $\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{9}{2}}(c + dx)} dx$

$$\begin{aligned}
& -\frac{2aA(a+b\tan(c+dx))^{3/2}}{7d\tan^{7/2}(c+dx)} + \\
\frac{1}{7} & \left(-\frac{2a(7aB+10Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{5/2}(c+dx)} + \frac{1}{5} \left(\frac{2(35a^2A-77abB-45Ab^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)} - \frac{2(105a^3B+24a^2A)}{3d\tan^{3/2}(c+dx)} \right) \right) \\
& \quad \downarrow \text{216} \\
& -\frac{2aA(a+b\tan(c+dx))^{3/2}}{7d\tan^{7/2}(c+dx)} + \\
\frac{1}{7} & \left(-\frac{2a(7aB+10Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{5/2}(c+dx)} + \frac{1}{5} \left(\frac{2(35a^2A-77abB-45Ab^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)} - \frac{2(105a^3B+24a^2A)}{3d\tan^{3/2}(c+dx)} \right) \right) \\
& \quad \downarrow \text{219} \\
& -\frac{2aA(a+b\tan(c+dx))^{3/2}}{7d\tan^{7/2}(c+dx)} + \\
\frac{1}{7} & \left(-\frac{2a(7aB+10Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{5/2}(c+dx)} + \frac{1}{5} \left(\frac{2(35a^2A-77abB-45Ab^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)} - \frac{2(105a^3B+24a^2A)}{3d\tan^{3/2}(c+dx)} \right) \right)
\end{aligned}$$

input `Int[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(9/2),x]`


```
output (-2*a*A*(a + b*Tan[c + d*x])^(3/2))/(7*d*Tan[c + d*x]^(7/2)) + ((-2*a*(10*
A*b + 7*a*B)*Sqrt[a + b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) + ((2*(35*
a^2*A - 45*A*b^2 - 77*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(
3/2)) - ((-105*((a^2*(a + I*b)^3*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[
c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d) + (a^2*(a - I*b)^3
*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d
*x]]])/(Sqrt[I*a + b]*d)))/a - (2*(245*a^2*A*b - 15*A*b^3 + 105*a^3*B - 16
1*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a)/5)/7
```

3.448.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 104 Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))/((e_) + (f_)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

```
rule 219 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

3.448.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.06 (sec) , antiderivative size = 2654465, normalized size of antiderivative = 8590.50

output too large to display

```
input int((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x)
```

```
output result too large to display
```

3.448.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18856 vs. $2(257) = 514$.

Time = 4.85 (sec) , antiderivative size = 18856, normalized size of antiderivative = 61.02

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algo
rithm="fracas")
```

```
output Too large to include
```

3.448. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{9/2}(c+dx)} dx$

3.448.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(9/2),x)`

output `Timed out`

3.448.7 Maxima [F]

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^{5/2}}{\tan^2(dx + c)^{9/2}} dx$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)/tan(d*x + c)^(9/
2), x)`

3.448.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(9/2),x, algo
rithm="giac")`

output `Timed out`

3.448.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{9/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2}}{\tan(c + dx)^{9/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(9/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(9/2),x)`

$$3.449 \quad \int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$$

3.449.1 Optimal result	4327
3.449.2 Mathematica [A] (verified)	4329
3.449.3 Rubi [A] (verified)	4330
3.449.4 Maple [B] (warning: unable to verify)	4338
3.449.5 Fricas [B] (verification not implemented)	4338
3.449.6 Sympy [F(-1)]	4338
3.449.7 Maxima [F(-1)]	4339
3.449.8 Giac [F(-1)]	4339
3.449.9 Mupad [F(-1)]	4339

3.449.1 Optimal result

Integrand size = 35, antiderivative size = 378

$$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx = \frac{(ia-b)^{5/2}(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$- \frac{(ia+b)^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} - \frac{2a(4Ab+3aB)\sqrt{a+b \tan(c+dx)}}{21d \tan^{\frac{7}{2}}(c+dx)}$$

$$+ \frac{2(21a^2A-25Ab^2-45abB)\sqrt{a+b \tan(c+dx)}}{105d \tan^{\frac{5}{2}}(c+dx)}$$

$$+ \frac{2(231a^2Ab-5Ab^3+105a^3B-135ab^2B)\sqrt{a+b \tan(c+dx)}}{315ad \tan^{\frac{3}{2}}(c+dx)}$$

$$- \frac{2(315a^4A-483a^2Ab^2-10Ab^4-735a^3bB+45ab^3B)\sqrt{a+b \tan(c+dx)}}{315a^2d\sqrt{\tan(c+dx)}}$$

$$- \frac{2aA(a+b \tan(c+dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c+dx)}$$

output $(I*a-b)^{(5/2)}*(A+I*B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})/d-(I*a+b)^{(5/2)}*(A-I*B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})/d-2/315*(315*A*a^4-483*A*a^2*b^2-10*A*b^4-735*B*a^3*b+45*B*a*b^3)*(a+b*\tan(d*x+c))^{(1/2)}/a^2/d/\tan(d*x+c)^{(1/2)}-2/21*a*(4*A*b+3*B*a)*(a+b*\tan(d*x+c))^{(1/2)}/d/\tan(d*x+c)^{(7/2)}+2/105*(21*A*a^2-25*A*b^2-45*B*a*b)*(a+b*\tan(d*x+c))^{(1/2)}/d/\tan(d*x+c)^{(5/2)}+2/315*(231*A*a^2*b-5*A*b^3+105*B*a^3-135*B*a*b^2)*(a+b*\tan(d*x+c))^{(1/2)}/a/d/\tan(d*x+c)^{(3/2)}-2/9*a*A*(a+b*\tan(d*x+c))^{(3/2)}/d/\tan(d*x+c)^{(9/2)}$

3.449. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{1/2}(c+dx)} dx$

3.449.2 Mathematica [A] (verified)

Time = 7.15 (sec) , antiderivative size = 543, normalized size of antiderivative = 1.44

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx = -\frac{bB(a + b \tan(c + dx))^{3/2}}{3d \tan^{\frac{9}{2}}(c + dx)}$$

$$+\frac{1}{3} - \frac{3b(2Ab + aB)\sqrt{a + b \tan(c + dx)}}{8d \tan^{\frac{9}{2}}(c + dx)} + \frac{1}{4} - \frac{(16a^2A - 18Ab^2 - 33abB)\sqrt{a + b \tan(c + dx)}}{6d \tan^{\frac{9}{2}}(c + dx)} - \frac{6a(38a}{2}$$

3.449. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$

input `Integrate[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(11/2),x]`

output `-1/3*(b*B*(a + b*Tan[c + d*x])^(3/2))/(d*Tan[c + d*x]^(9/2)) + ((-3*b*(2*A*b + a*B)*Sqrt[a + b*Tan[c + d*x]])/(8*d*Tan[c + d*x]^(9/2)) + (-1/6*((16*a^2*A - 18*A*b^2 - 33*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Tan[c + d*x]^(9/2)) - (2*((6*a*(38*a*A*b + 18*a^2*B - 21*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(7*d*Tan[c + d*x]^(7/2)) - (2*((18*a^2*(21*a^2*A - 25*A*b^2 - 45*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) - (2*((-3*a^2*(231*a^2*A*b - 5*A*b^3 + 105*a^3*B - 135*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Tan[c + d*x]^(3/2)) - (2*((-2835*a^4*(-1)^(1/4)*(-a + I*b)^(5/2)*(A - I*B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) + (-1)^(1/4)*(a + I*b)^(5/2)*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])))/(4*d) - (9*a^2*(315*a^4*A - 483*a^2*A*b^2 - 10*A*b^4 - 735*a^3*b*B + 45*a*b^3*B)*Sqrt[a + b*Tan[c + d*x]])/(2*d*Sqrt[Tan[c + d*x]])))/(3*a))/(5*a))/(7*a))/(9*a))/4)/3`

3.449.3 Rubi [A] (verified)

Time = 3.00 (sec) , antiderivative size = 428, normalized size of antiderivative = 1.13, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.629$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan(c + dx)^{11/2}} dx$$

↓ 4088

$$\frac{2}{9} \int \frac{3\sqrt{a + b \tan(c + dx)} (-b(2aA - 3bB) \tan^2(c + dx) - 3(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(4Ab + 3aB))}{2 \tan^{9/2}(c + dx) + \frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{9/2}(c + dx)}} dx$$

3.449. $\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx$

↓ 27

$$\frac{1}{3} \int \frac{\sqrt{a + b \tan(c + dx)} (-b(2aA - 3bB) \tan^2(c + dx) - 3(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(4Ab + 3aB))}{\tan^{\frac{9}{2}}(c + dx)} dx$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \int \frac{\sqrt{a + b \tan(c + dx)} (-b(2aA - 3bB) \tan(c + dx)^2 - 3(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(4Ab + 3aB))}{\tan(c + dx)^{9/2}} dx$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 4128

$$\frac{1}{3} \left(\frac{2}{7} \int -\frac{b(18Ba^2 + 38Aba - 21b^2B) \tan^2(c + dx) + 21(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(21Aa^2 - 21Ab^2)}{2 \tan^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{3} \left(-\frac{1}{7} \int \frac{b(18Ba^2 + 38Aba - 21b^2B) \tan^2(c + dx) + 21(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(21Aa^2 - 21Ab^2)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \left(-\frac{1}{7} \int \frac{b(18Ba^2 + 38Aba - 21b^2B) \tan(c + dx)^2 + 21(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx) + a(21Aa^2 - 21Ab^2)}{\tan(c + dx)^{7/2} \sqrt{a + b \tan(c + dx)}} dx \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 4132

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2 \int -\frac{-4ab(21Aa^2 - 45bBa - 25Ab^2) \tan^2(c+dx) - 105a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) + a(105Ba^3 + 231Aba^2 - 135b^2Ba - 5Ab^3)}{2 \tan^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}}{5a} \right) \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2) \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{\int -\frac{4ab(21Aa^2 - 45bBa - 25Ab^2) \tan^2(c+dx) - 105a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) + a(105Ba^3 + 231Aba^2 - 135b^2Ba - 5Ab^3)}{\tan^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}}{5a} \right) \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2) \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{\int -\frac{4ab(21Aa^2 - 45bBa - 25Ab^2) \tan(c+dx)^2 - 105a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) + a(105Ba^3 + 231Aba^2 - 135b^2Ba - 5Ab^3)}{\tan(c+dx)^{5/2} \sqrt{a+b \tan(c+dx)}}}{5a} \right) \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 4132

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2) \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{2 \int \frac{315(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)a^2 + 2b(105Ba^3 + 231Aba^2 - 135b^2Ba - 5Ab^3)}{2 \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}}{5a} \right) \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{3} \left(\frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2) \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{\int \frac{315(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)a^2 + 2b(105Ba^3 + 231Aba^2 - 135b^2Ba - 5Ab^3)}{\tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}}{5a} \right) \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)}$$

3.449. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{\frac{11}{2}}(c+dx)} dx$

$$\begin{aligned} & \downarrow 3042 \\ \frac{1}{3} \left(\frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2) \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{\int \frac{315(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)a^2 + 2b(105Ba^3 + 231Aba^2 - 135Ab^2a + b^3B) \tan^2(c + dx)}{\tan(c + dx)^5} dx}{\tan(c + dx)^5} \right) \right. \\ & \left. \frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 4132 \\ \frac{1}{3} \left(\frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2) \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{2 \int \frac{315(a^3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) - a^3(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx))}{2\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx}{a} \right) \right. \\ & \left. \frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ \frac{1}{3} \left(\frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2) \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{315 \int \frac{a^3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) - a^3(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx}{a} \right) \right. \\ & \left. \frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ \frac{1}{3} \left(\frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2) \sqrt{a + b \tan(c + dx)}}{5d \tan^{\frac{5}{2}}(c + dx)} - \frac{315 \int \frac{a^3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) - a^3(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx}{a} \right) \right. \\ & \left. \frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c + dx)} \right) \end{aligned}$$

$$\begin{aligned} & \downarrow 4099 \\ 3.449. \quad \int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx \end{aligned}$$

$$\begin{aligned}
& -\frac{2aA(a+b\tan(c+dx))^{3/2}}{9d\tan^{9/2}(c+dx)} + \\
\frac{1}{3} & \left(-\frac{2a(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{7d\tan^{7/2}(c+dx)} + \frac{1}{7} \left(\frac{2(21a^2A-45abB-25Ab^2)\sqrt{a+b\tan(c+dx)}}{5d\tan^{5/2}(c+dx)} - \frac{2(105a^3B+231a^2A)}{9d\tan^{9/2}(c+dx)} \right) \right) \\
& \quad \downarrow 3042 \\
& -\frac{2aA(a+b\tan(c+dx))^{3/2}}{9d\tan^{9/2}(c+dx)} + \\
\frac{1}{3} & \left(-\frac{2a(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{7d\tan^{7/2}(c+dx)} + \frac{1}{7} \left(\frac{2(21a^2A-45abB-25Ab^2)\sqrt{a+b\tan(c+dx)}}{5d\tan^{5/2}(c+dx)} - \frac{2(105a^3B+231a^2A)}{9d\tan^{9/2}(c+dx)} \right) \right) \\
& \quad \downarrow 4098 \\
& -\frac{2aA(a+b\tan(c+dx))^{3/2}}{9d\tan^{9/2}(c+dx)} + \\
\frac{1}{3} & \left(-\frac{2a(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{7d\tan^{7/2}(c+dx)} + \frac{1}{7} \left(\frac{2(21a^2A-45abB-25Ab^2)\sqrt{a+b\tan(c+dx)}}{5d\tan^{5/2}(c+dx)} - \frac{2(105a^3B+231a^2A)}{9d\tan^{9/2}(c+dx)} \right) \right) \\
& \quad \downarrow 104 \\
& -\frac{2aA(a+b\tan(c+dx))^{3/2}}{9d\tan^{9/2}(c+dx)} + \\
\frac{1}{3} & \left(-\frac{2a(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{7d\tan^{7/2}(c+dx)} + \frac{1}{7} \left(\frac{2(21a^2A-45abB-25Ab^2)\sqrt{a+b\tan(c+dx)}}{5d\tan^{5/2}(c+dx)} - \frac{2(105a^3B+231a^2A)}{9d\tan^{9/2}(c+dx)} \right) \right) \\
& \quad \downarrow 216
\end{aligned}$$

3.449. $\int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\tan^{11/2}(c+dx)} dx$

$$\begin{aligned}
 & -\frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{9/2}(c + dx)} + \\
 & \left(\frac{1}{3} \left(-\frac{2a(3aB + 4Ab)\sqrt{a + b \tan(c + dx)}}{7d \tan^{7/2}(c + dx)} + \frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2)\sqrt{a + b \tan(c + dx)}}{5d \tan^{5/2}(c + dx)} - \frac{2(105a^3B + 231a^2A - 45abB - 25Ab^2)}{3d \tan^{3/2}(c + dx)} \right) \right) \right) \\
 & \quad \downarrow \text{219} \\
 & -\frac{2aA(a + b \tan(c + dx))^{3/2}}{9d \tan^{9/2}(c + dx)} + \\
 & \left(\frac{1}{3} \left(-\frac{2a(3aB + 4Ab)\sqrt{a + b \tan(c + dx)}}{7d \tan^{7/2}(c + dx)} + \frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2)\sqrt{a + b \tan(c + dx)}}{5d \tan^{5/2}(c + dx)} - \frac{2(105a^3B + 231a^2A - 45abB - 25Ab^2)}{3d \tan^{3/2}(c + dx)} \right) \right) \right)
 \end{aligned}$$

input `Int[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(11/2), x]`

output `(-2*a*A*(a + b*Tan[c + d*x])^(3/2))/(9*d*Tan[c + d*x]^(9/2)) + ((-2*a*(4*A*b + 3*a*B)*Sqrt[a + b*Tan[c + d*x]])/(7*d*Tan[c + d*x]^(7/2)) + ((2*(21*a^2*A - 25*A*b^2 - 45*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) - ((-2*(231*a^2*A*b - 5*A*b^3 + 105*a^3*B - 135*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - ((315*(-((a^3*(a + I*b)^3*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d)) + (a^3*(a - I*b)^3*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d)))/a - (2*(315*a^4*A - 483*a^2*A*b^2 - 10*A*b^4 - 735*a^3*b*B + 45*a*b^3*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a)/(5*a))/7)/3`

3.449. $\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{11/2}(c+dx)} dx$

3.449.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && ! (ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.449.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.05 (sec) , antiderivative size = 2659448, normalized size of antiderivative = 7035.58

output too large to display

input `int((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x)`

output `result too large to display`

3.449.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18922 vs. $2(320) = 640$.

Time = 4.90 (sec) , antiderivative size = 18922, normalized size of antiderivative = 50.06

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x, algorithm="fracas")`

output `Too large to include`

3.449.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(11/2),x)`

output `Timed out`

3.449.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x, algorithm="maxima")`

output `Timed out`

3.449.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(11/2),x, algorithm="giac")`

output `Timed out`

3.449.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{11/2}(c + dx)} dx = \text{Hanged}$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(11/2), x)`

output `\text{Hanged}`

$$3.450 \quad \int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$$

3.450.1 Optimal result	4340
3.450.2 Mathematica [A] (verified)	4342
3.450.3 Rubi [A] (verified)	4343
3.450.4 Maple [B] (warning: unable to verify)	4354
3.450.5 Fricas [B] (verification not implemented)	4354
3.450.6 Sympy [F(-1)]	4354
3.450.7 Maxima [F(-1)]	4355
3.450.8 Giac [F(-1)]	4355
3.450.9 Mupad [F(-1)]	4355

3.450.1 Optimal result

Integrand size = 35, antiderivative size = 460

$$\int \frac{(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{13}{2}}(c + dx)} dx =$$

$$\frac{(ia - b)^{5/2}(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d}$$

$$- \frac{(ia + b)^{5/2}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} - \frac{2a(14Ab + 11aB)\sqrt{a + b \tan(c + dx)}}{99d \tan^{\frac{9}{2}}(c + dx)}$$

$$+ \frac{2(99a^2A - 113Ab^2 - 209abB)\sqrt{a + b \tan(c + dx)}}{693d \tan^{\frac{7}{2}}(c + dx)}$$

$$+ \frac{2(495a^2Ab - 5Ab^3 + 231a^3B - 275ab^2B)\sqrt{a + b \tan(c + dx)}}{1155ad \tan^{\frac{5}{2}}(c + dx)}$$

$$- \frac{2(1155a^4A - 1485a^2Ab^2 - 20Ab^4 - 2541a^3bB + 55ab^3B)\sqrt{a + b \tan(c + dx)}}{3465a^2d \tan^{\frac{3}{2}}(c + dx)}$$

$$- \frac{2(8085a^4Ab - 495a^2Ab^3 + 40Ab^5 + 3465a^5B - 5313a^3b^2B - 110ab^4B)\sqrt{a + b \tan(c + dx)}}{3465a^3d \sqrt{\tan(c + dx)}}$$

$$- \frac{2aA(a + b \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

$$3.450. \quad \int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$$

output

$$\begin{aligned}
& -(I*a-b)^{(5/2)}*(I*A-B)*\arctan((I*a-b)^{(1/2)}*\tan(dx+c)^{(1/2)}/(a+b*\tan(dx+c))^{(1/2)})/d-(I*a+b)^{(5/2)}*(I*A+B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(dx+c)^{(1/2)}/(a+b*\tan(dx+c))^{(1/2)})/d-2/3465*(8085*A*a^4*b-495*A*a^2*b^3+40*A*b^5+3465*B*a^5-5313*B*a^3*b^2-110*B*a*b^4)*(a+b*\tan(dx+c))^{(1/2)}/a^3/d/\tan(dx+c)^{(1/2)}-2/99*a*(14*A*b+11*B*a)*(a+b*\tan(dx+c))^{(1/2)}/d/\tan(dx+c)^{(9/2)}+2/693*(99*A*a^2-113*A*b^2-209*B*a*b)*(a+b*\tan(dx+c))^{(1/2)}/d/\tan(dx+c)^{(7/2)}+2/1155*(495*A*a^2*b-5*A*b^3+231*B*a^3-275*B*a*b^2)*(a+b*\tan(dx+c))^{(1/2)}/a/d/\tan(dx+c)^{(5/2)}-2/3465*(1155*A*a^4-1485*A*a^2*b^2-20*A*b^4-2541*B*a^3*b+55*B*a*b^3)*(a+b*\tan(dx+c))^{(1/2)}/a^2/d/\tan(dx+c)^{(3/2)}-2/11*a*A*(a+b*\tan(dx+c))^{(3/2)}/d/\tan(dx+c)^{(11/2)}
\end{aligned}$$

3.450.
$$\int \frac{(a+b \tan(cx+d))^{5/2} (A+B \tan(cx+d))}{\tan^{13/2}(cx+d)} dx$$

3.450.2 Mathematica [A] (verified)

Time = 7.31 (sec) , antiderivative size = 632, normalized size of antiderivative = 1.37

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{13}{2}}(c + dx)} dx = -\frac{bB(a + b \tan(c + dx))^{3/2}}{4d \tan^{\frac{11}{2}}(c + dx)}$$

3.450. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$

input `Integrate[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(13/2),x]`

output `-1/4*(b*B*(a + b*Tan[c + d*x])^(3/2))/(d*Tan[c + d*x]^(11/2)) + (-1/10*(b*(8*A*b + 5*a*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Tan[c + d*x]^(11/2)) + (-1/22*((80*a^2*A - 88*A*b^2 - 165*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Tan[c + d*x]^(11/2)) - (2*((5*a*(184*a*A*b + 88*a^2*B - 99*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(18*d*Tan[c + d*x]^(9/2)) - (2*((10*a^2*(99*a^2*A - 113*A*b^2 - 209*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(7*d*Tan[c + d*x]^(7/2)) - (2*((-3*a^2*(495*a^2*A*b - 5*A*b^3 + 231*a^3*B - 275*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Tan[c + d*x]^(5/2)) - (2*((-5*a^2*(1155*a^4*A - 1485*a^2*A*b^2 - 20*A*b^4 - 2541*a^3*b*B + 55*a*b^3*B)*Sqrt[a + b*Tan[c + d*x]])/(2*d*Tan[c + d*x]^(3/2)) - (2*((51975*a^5*((-1)^(3/4)*(-a + I*b)^(5/2)*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - (-1)^(3/4)*(a + I*b)^(5/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]))/(8*d) + (15*a^2*(8085*a^4*A*b - 495*a^2*A*b^3 + 40*A*b^5 + 3465*a^5*B - 5313*a^3*b^2*B - 110*a*b^4*B)*Sqrt[a + b*Tan[c + d*x]]/(4*d*Sqrt[Tan[c + d*x]])))/(3*a)))/(5*a)))/(7*a)))/(9*a)))/(11*a))/5)/4`

3.450.3 Rubi [A] (verified)

Time = 3.68 (sec) , antiderivative size = 509, normalized size of antiderivative = 1.11, number of steps used = 26, number of rules used = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{13}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan(c + dx)^{13/2}} dx$$

↓ 4088

3.450. $\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{13}{2}}(c + dx)} dx$

$$\frac{2}{11} \int \frac{\sqrt{a+b \tan(c+dx)}(-b(8aA-11bB) \tan^2(c+dx)-11(Aa^2-2bBa-Ab^2) \tan(c+dx)+a(14Ab+11aB)}{2 \tan^{\frac{11}{2}}(c+dx)} dx$$

$$\frac{2aA(a+b \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{11} \int \frac{\sqrt{a+b \tan(c+dx)}(-b(8aA-11bB) \tan^2(c+dx)-11(Aa^2-2bBa-Ab^2) \tan(c+dx)+a(14Ab+11aB)}{\tan^{\frac{11}{2}}(c+dx)} dx$$

$$\frac{2aA(a+b \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{11} \int \frac{\sqrt{a+b \tan(c+dx)}(-b(8aA-11bB) \tan(c+dx)^2-11(Aa^2-2bBa-Ab^2) \tan(c+dx)+a(14Ab+11aB)}{\tan(c+dx)^{11/2}} dx$$

$$\frac{2aA(a+b \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)}$$

↓ 4128

$$\frac{1}{11} \left(\frac{2}{9} \int \frac{b(88Ba^2+184Aba-99b^2B) \tan^2(c+dx)+99(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx)+a(99Aa^2+11aB)}{2 \tan^{\frac{9}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx \right)$$

$$\frac{2aA(a+b \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{11} \left(-\frac{1}{9} \int \frac{b(88Ba^2+184Aba-99b^2B) \tan^2(c+dx)+99(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx)+a(99Aa^2+11aB)}{\tan^{\frac{9}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx \right)$$

$$\frac{2aA(a+b \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{11} \left(-\frac{1}{9} \int \frac{b(88Ba^2+184Aba-99b^2B) \tan(c+dx)^2+99(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx)+a(99Aa^2+11aB)}{\tan(c+dx)^{9/2} \sqrt{a+b \tan(c+dx)}} dx \right)$$

$$\frac{2aA(a+b \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)}$$

↓ 4132

3.450. $\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$

$$\frac{1}{11} \left(\frac{1}{9} \left(2 \int \frac{3(-2ab(99Aa^2 - 209bBa - 113Ab^2) \tan^2(c+dx) - 231a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) + a(231Ba^3 + 495Aba^2 - 275b^2Ba - 113Ab^2))}{2 \tan^{\frac{7}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx - \frac{3 \int \frac{-2ab(99Aa^2 - 209bBa - 113Ab^2) \tan^2(c+dx) - 231a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) + a(231Ba^3 + 495Aba^2 - 275b^2Ba - 113Ab^2)}{\tan^{\frac{7}{2}}(c+dx)} dx \right) - \frac{2aA(a+b \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)} \right)$$

↓ 27

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} - \frac{3 \int \frac{-2ab(99Aa^2 - 209bBa - 113Ab^2) \tan^2(c+dx) - 231a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) + a(231Ba^3 + 495Aba^2 - 275b^2Ba - 113Ab^2)}{\tan^{\frac{7}{2}}(c+dx)} dx \right) - \frac{2aA(a+b \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} - \frac{3 \int \frac{-2ab(99Aa^2 - 209bBa - 113Ab^2) \tan(c+dx)^2 - 231a(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx) + a(231Ba^3 + 495Aba^2 - 275b^2Ba - 113Ab^2)}{\tan(c+dx)} dx \right) - \frac{2aA(a+b \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)} \right)$$

↓ 4132

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} - \frac{3 \left(\frac{2 \int \frac{1155(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c+dx)a^2 + 4b(231Ba^3 + 495Aba^2 - 275b^2Ba - 113Ab^2)}{\tan^{\frac{7}{2}}(c+dx)} dx \right)}{\tan^{\frac{11}{2}}(c+dx)} \right) - \frac{2aA(a+b \tan(c+dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c+dx)} \right)$$

↓ 27

3.450. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{\int \frac{1155(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)a^2 + 4b(231Ba^3 + 44Aa^2b - 33Ab^2 - 3b^3A)}{7d \tan^{\frac{7}{2}}(c + dx)} dx}{3} \right) \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{\int \frac{1155(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx)a^2 + 4b(231Ba^3 + 44Aa^2b - 33Ab^2 - 3b^3A)}{7d \tan^{\frac{7}{2}}(c + dx)} dx}{3} \right) \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 4132

$$\frac{1}{11} \left(\frac{1}{9} \left(\frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{\int \frac{-3465(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx)a^3 - 2b(1155Aa^2 - 33Ab^2 - 3b^3A)}{7d \tan^{\frac{7}{2}}(c + dx)} dx}{3} \right) \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 27

3.450. $\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{13}{2}}(c + dx)} dx$

$$\frac{1}{11} \left(\frac{1}{9} \frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \int \frac{-3465(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx)a^3 - 2b(1155Aa^4}{\dots} \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)} \downarrow 3042$$

$$\frac{1}{11} \left(\frac{1}{9} \frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \int \frac{-3465(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c+dx)a^3 - 2b(1155Aa^4}{\dots} \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)} \downarrow 4132$$

$$\frac{1}{11} \left(\frac{1}{9} \frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \int \frac{3465((Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)a^4 + (Ba^3 + 3Aba^2 - 3b^2B) \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}}{2 \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)} a} \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

3.450. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$

↓ 27

$$\frac{1}{11} \left(\frac{1}{9} \frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{3465 \int \frac{(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)a^4 + (Ba^3 + 3Aba^2 - 3b^2Ba^2)}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}}}{a} \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{11} \left(\frac{1}{9} \frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a + b \tan(c + dx)}}{7d \tan^{\frac{7}{2}}(c + dx)} - \frac{3465 \int \frac{(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)a^4 + (Ba^3 + 3Aba^2 - 3b^2Ba^2)}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}}}{a} \right)$$

$$\frac{2aA(a + b \tan(c + dx))^{3/2}}{11d \tan^{\frac{11}{2}}(c + dx)}$$

↓ 4099

3.450. $\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$

$$\begin{aligned}
 & -\frac{2aA(a+b\tan(c+dx))^{3/2}}{11d\tan^{\frac{11}{2}}(c+dx)} + \\
 & \left(\frac{1}{11} \left(-\frac{2a(11aB+14Ab)\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} \frac{2(99a^2A-209abB-113Ab^2)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} - \frac{2(2a^2+3ab+2b^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{5}{2}}(c+dx)} \right) \right)
 \end{aligned}$$

↓ 3042

$$\begin{aligned}
 & -\frac{2aA(a+b\tan(c+dx))^{3/2}}{11d\tan^{\frac{11}{2}}(c+dx)} + \\
 & \left(\frac{1}{11} \left(-\frac{2a(11aB+14Ab)\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} \frac{2(99a^2A-209abB-113Ab^2)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} - \frac{2(2a^2+3ab+2b^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{5}{2}}(c+dx)} \right) \right)
 \end{aligned}$$

↓ 4098

$$\begin{aligned}
 & -\frac{2aA(a+b\tan(c+dx))^{3/2}}{11d\tan^{\frac{11}{2}}(c+dx)} + \\
 & \left(\frac{1}{11} \left(-\frac{2a(11aB+14Ab)\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} \frac{2(99a^2A-209abB-113Ab^2)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} - \frac{2(2a^2+3ab+2b^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{5}{2}}(c+dx)} \right) \right)
 \end{aligned}$$

3.450. $\int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\tan^{\frac{13}{2}}(c+dx)} dx$

$$\begin{aligned}
 & \downarrow 104 \\
 & -\frac{2aA(a+b\tan(c+dx))^{3/2}}{11d\tan^{\frac{11}{2}}(c+dx)} + \\
 & \left(\frac{1}{11} - \frac{2a(11aB+14Ab)\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} \frac{2(99a^2A-209abB-113Ab^2)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} - \frac{2(2a^2+3ab+2b^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{5}{2}}(c+dx)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 216 \\
 & -\frac{2aA(a+b\tan(c+dx))^{3/2}}{11d\tan^{\frac{11}{2}}(c+dx)} + \\
 & \left(\frac{1}{11} - \frac{2a(11aB+14Ab)\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} \frac{2(99a^2A-209abB-113Ab^2)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} - \frac{2(2a^2+3ab+2b^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{5}{2}}(c+dx)} \right)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 219 \\
 & -\frac{2aA(a + b \tan(c + dx))^{3/2}}{11d \tan^{11/2}(c + dx)} + \\
 & \left(\frac{1}{11} - \frac{2a(11aB + 14Ab)\sqrt{a + b \tan(c + dx)}}{9d \tan^{9/2}(c + dx)} + \frac{1}{9} \right) \frac{2(99a^2A - 209abB - 113Ab^2)\sqrt{a + b \tan(c + dx)}}{7d \tan^{7/2}(c + dx)} - \dots
 \end{aligned}$$

```
input Int[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(13/2),
x]
```

```
output (-2*a*A*(a + b*Tan[c + d*x])^(3/2))/(11*d*Tan[c + d*x]^(11/2)) + ((-2*a*(1
4*A*b + 11*a*B)*Sqrt[a + b*Tan[c + d*x]])/(9*d*Tan[c + d*x]^(9/2)) + ((2*(
99*a^2*A - 113*A*b^2 - 209*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(7*d*Tan[c + d
*x]^(7/2)) - (3*((-2*(495*a^2*A*b - 5*A*b^3 + 231*a^3*B - 275*a*b^2*B)*Sqr
t[a + b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) - ((-2*(1155*a^4*A - 1485*
a^2*A*b^2 - 20*A*b^4 - 2541*a^3*b*B + 55*a*b^3*B)*Sqrt[a + b*Tan[c + d*x]]
)/(3*d*Tan[c + d*x]^(3/2)) + ((-3465*((a^4*(a + I*b)^3*(A + I*B)*ArcTan[(S
qrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]
*d) + (a^4*(a - I*b)^3*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]]
)/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d)))/a - (2*(8085*a^4*A*b - 49
5*a^2*A*b^3 + 40*A*b^5 + 3465*a^5*B - 5313*a^3*b^2*B - 110*a*b^4*B)*Sqrt[a
+ b*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a)/(5*a))/(7*a))/9)/11
```

3.450. $\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{13/2}(c+dx)} dx$

3.450.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && ! (ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.450.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 5.88 (sec) , antiderivative size = 2660696, normalized size of antiderivative = 5784.12

output too large to display

input `int((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(13/2),x)`

output `result too large to display`

3.450.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18981 vs. $2(396) = 792$.

Time = 4.91 (sec) , antiderivative size = 18981, normalized size of antiderivative = 41.26

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{13}{2}}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(13/2),x, algorithm="fricas")`

output `Too large to include`

3.450.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{\frac{13}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)**(13/2),x)`

output `Timed out`

3.450.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{13/2}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(13/2),x, algorithm="maxima")
```

```
output Timed out
```

3.450.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{13/2}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/tan(d*x+c)^(13/2),x, algorithm="giac")
```

```
output Timed out
```

3.450.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\tan^{13/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2}}{\tan(c + dx)^{13/2}} dx$$

```
input int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(13/2), x)
```

```
output int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(13/2), x)
```

3.450. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\tan^{13/2}(c+dx)} dx$

3.451
$$\int \frac{(a+b \tan(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c+dx)\right)}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.451.1 Optimal result 4356
 3.451.2 Mathematica [A] (verified) 4357
 3.451.3 Rubi [A] (verified) 4357
 3.451.4 Maple [B] (warning: unable to verify) 4361
 3.451.5 Fricas [B] (verification not implemented) 4362
 3.451.6 Sympy [F] 4362
 3.451.7 Maxima [F] 4363
 3.451.8 Giac [F(-1)] 4363
 3.451.9 Mupad [F(-1)] 4363

3.451.1 Optimal result

Integrand size = 43, antiderivative size = 253

$$\int \frac{(a + b \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx)\right)}{\tan^{\frac{5}{2}}(c + dx)} dx = \frac{(ia - b)^{5/2}(2a - 3ib)B \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{2ad}$$

$$+ \frac{2b^{5/2}B \operatorname{Arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{d} - \frac{(2a + 3ib)(ia + b)^{5/2}B \operatorname{Arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{2ad}$$

$$- \frac{2(a^2 + 3b^2)B \sqrt{a + b \tan(c + dx)}}{d \sqrt{\tan(c + dx)}} - \frac{bB(a + b \tan(c + dx))^{3/2}}{d \tan^{\frac{3}{2}}(c + dx)}$$

output

```
1/2*(I*a-b)^(5/2)*(2*a-3*I*b)*B*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b
*tan(d*x+c))^(1/2))/a/d+2*b^(5/2)*B*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*
tan(d*x+c))^(1/2))/d-1/2*(2*a+3*I*b)*(I*a+b)^(5/2)*B*arctanh((I*a+b)^(1/2)
*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/a/d-2*(a^2+3*b^2)*B*(a+b*tan(d*x
+c))^(1/2)/d/tan(d*x+c)^(1/2)-b*B*(a+b*tan(d*x+c))^(3/2)/d/tan(d*x+c)^(3/2)
)
```

3.451.
$$\int \frac{(a+b \tan(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c+dx)\right)}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.451.2 Mathematica [A] (verified)

Time = 4.86 (sec) , antiderivative size = 356, normalized size of antiderivative = 1.41

$$\int \frac{(a + b \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{\frac{5}{2}}(c + dx)} dx = \frac{B \cos(c + dx)(3b + 2a \tan(c + dx)) \left(4\sqrt{ab}^{5/2} \operatorname{arcsinh} \right)}{\tan^{\frac{5}{2}}(c + dx)}$$

input `Integrate[((a + b*Tan[c + d*x])^(5/2)*((3*b*B)/(2*a) + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2),x]`

output `(B*Cos[c + d*x]*(3*b + 2*a*Tan[c + d*x])*(4*Sqrt[a]*b^(5/2)*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Tan[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]] - Sqrt[1 + (b*Tan[c + d*x])/a]*((-1)^(1/4)*(-a + I*b)^(5/2)*(2*a + (3*I)*b)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]]*Tan[c + d*x]^(3/2) + (-1)^(1/4)*(a + I*b)^(5/2)*(2*a - (3*I)*b)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]]*Tan[c + d*x]^(3/2) + 2*a*Sqrt[a + b*Tan[c + d*x]]*(a*b + (2*a^2 + 7*b^2)*Tan[c + d*x])))/(2*a*d*(3*b*Cos[c + d*x] + 2*a*Sin[c + d*x])*Tan[c + d*x]^(3/2)*Sqrt[1 + (b*Tan[c + d*x])/a])`

3.451.3 Rubi [A] (verified)Time = 1.57 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.01, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.256$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{\frac{5}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan(c + dx)^{5/2}} dx$$

↓ 4088

3.451. $\int \frac{(a + b \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{\frac{5}{2}}(c + dx)} dx$

$$\frac{2}{3} \int \frac{3\sqrt{a+b \tan(c+dx)} \left(2b^2 B \tan^2(c+dx) + b \left(\frac{3b^2}{a} + a \right) B \tan(c+dx) + 2(a^2 + 3b^2) B \right)}{4 \tan^{\frac{3}{2}}(c+dx) \frac{bB(a+b \tan(c+dx))^{3/2}}{d \tan^{\frac{3}{2}}(c+dx)}} dx -$$

↓ 27

$$\frac{1}{2} \int \frac{\sqrt{a+b \tan(c+dx)} \left(2b^2 B \tan^2(c+dx) + b \left(\frac{3b^2}{a} + a \right) B \tan(c+dx) + 2(a^2 + 3b^2) B \right)}{\tan^{\frac{3}{2}}(c+dx) \frac{bB(a+b \tan(c+dx))^{3/2}}{d \tan^{\frac{3}{2}}(c+dx)}} dx -$$

↓ 3042

$$\frac{1}{2} \int \frac{\sqrt{a+b \tan(c+dx)} \left(2b^2 B \tan(c+dx)^2 + b \left(\frac{3b^2}{a} + a \right) B \tan(c+dx) + 2(a^2 + 3b^2) B \right)}{\tan(c+dx)^{3/2} \frac{bB(a+b \tan(c+dx))^{3/2}}{d \tan^{\frac{3}{2}}(c+dx)}} dx -$$

↓ 4128

$$\frac{1}{2} \left(2 \int \frac{2B \tan^2(c+dx)b^3 + 3(a^2 + 3b^2) Bb - \frac{(2a^4 + 3b^2 a^2 - 3b^4) B \tan(c+dx)}{a}}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{4B(a^2 + 3b^2) \sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) \frac{bB(a+b \tan(c+dx))^{3/2}}{d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{2} \left(\int \frac{2B \tan^2(c+dx)b^3 + 3(a^2 + 3b^2) Bb - \frac{(2a^4 + 3b^2 a^2 - 3b^4) B \tan(c+dx)}{a}}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{4B(a^2 + 3b^2) \sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) \frac{bB(a+b \tan(c+dx))^{3/2}}{d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{2} \left(\int \frac{2B \tan(c+dx)^2 b^3 + 3(a^2 + 3b^2) Bb - \frac{(2a^4 + 3b^2 a^2 - 3b^4) B \tan(c+dx)}{a}}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{4B(a^2 + 3b^2) \sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) \frac{bB(a+b \tan(c+dx))^{3/2}}{d \tan^{\frac{3}{2}}(c+dx)}$$

↓ 4138

3.451. $\int \frac{(a+b \tan(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c+dx) \right)}{\tan^{\frac{5}{2}}(c+dx)} dx$

$$\frac{1}{2} \left(\frac{\int \frac{2B \tan^2(c+dx)b^3 + 3(a^2 + 3b^2)Bb - \frac{(2a^4 + 3b^2a^2 - 3b^4)B \tan(c+dx)}{a}}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} d \tan(c+dx)}{d} - \frac{4B(a^2 + 3b^2) \sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) -$$

$$\frac{bB(a+b \tan(c+dx))^{3/2}}{d \tan^{3/2}(c+dx)}$$

↓ 2035

$$\frac{1}{2} \left(\frac{2 \int \frac{2B \tan^2(c+dx)b^3 + 3(a^2 + 3b^2)Bb - \frac{(2a^4 + 3b^2a^2 - 3b^4)B \tan(c+dx)}{a}}{\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} d \sqrt{\tan(c+dx)}}{d} - \frac{4B(a^2 + 3b^2) \sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) -$$

$$\frac{bB(a+b \tan(c+dx))^{3/2}}{d \tan^{3/2}(c+dx)}$$

↓ 2257

$$\frac{1}{2} \left(\frac{2 \int \left(\frac{2Bb^3}{\sqrt{a+b \tan(c+dx)}} + \frac{ab(3a^2 + 7b^2)B - (2a^4 + 3b^2a^2 - 3b^4)B \tan(c+dx)}{a\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} \right) d \sqrt{\tan(c+dx)}}{d} - \frac{4B(a^2 + 3b^2) \sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) -$$

$$\frac{bB(a+b \tan(c+dx))^{3/2}}{d \tan^{3/2}(c+dx)}$$

↓ 2009

$$- \frac{bB(a+b \tan(c+dx))^{3/2}}{d \tan^{3/2}(c+dx)} +$$

$$\frac{1}{2} \left(- \frac{4B(a^2 + 3b^2) \sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{2 \left(\frac{B(2a-3ib)(-b+ia)^{5/2} \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{2a} + 2b^{5/2} B \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \right)}{d} \right)$$

input `Int[((a + b*Tan[c + d*x])^(5/2)*((3*b*B)/(2*a) + B*Tan[c + d*x]))/Tan[c + d*x]^(5/2), x]`

3.451. $\int \frac{(a+b \tan(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c+dx) \right)}{\tan^{5/2}(c+dx)} dx$

```
output -((b*B*(a + b*Tan[c + d*x])^(3/2))/(d*Tan[c + d*x]^(3/2))) + ((2*(((I*a -
b)^(5/2)*(2*a - (3*I)*b)*B*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[
a + b*Tan[c + d*x]]])/(2*a) + 2*b^(5/2)*B*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*
x]])/Sqrt[a + b*Tan[c + d*x]]] - ((2*a + (3*I)*b)*(I*a + b)^(5/2)*B*ArcTan
h[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(2*a))/d
- (4*(a^2 + 3*b^2)*B*Sqrt[a + b*Tan[c + d*x]]/(d*Sqrt[Tan[c + d*x]]))/2
```

3.451.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2035 Int[(F_x_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]
```

```
rule 2257 Int[(P_x_)*((d_) + (e_)*(x_)^(2))^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol
] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a
, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4088 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x
])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2))
Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*
(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n
+ 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[
e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n
+ 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] &
& LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])
```

$$3.451. \int \frac{(a+b \tan(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c+dx) \right)}{\tan^{\frac{5}{2}}(c+dx)} dx$$

```
rule 4128 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

3.451.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 5.86 (sec) , antiderivative size = 1491744, normalized size of antiderivative = 5896.22

output too large to display

```
input int((a+b*tan(d*x+c))^(5/2)*(3/2*b*B/a+B*tan(d*x+c))/tan(d*x+c)^(5/2),x)
```

```
output result too large to display
```

$$3.451. \quad \int \frac{(a+b \tan(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c+dx) \right)}{\tan^{\frac{5}{2}}(c+dx)} dx$$

3.451.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8171 vs. $2(205) = 410$.

Time = 2.54 (sec) , antiderivative size = 16341, normalized size of antiderivative = 64.59

$$\int \frac{(a + b \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(3/2*b*B/a+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="fricas")`

output Too large to include

3.451.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx = \frac{B \left(\int \frac{2a^3 \sqrt{a+b \tan(c+dx)}}{\tan^{3/2}(c+dx)} dx + \int \frac{3b^3 \sqrt{a+b \tan(c+dx)}}{\sqrt{\tan(c+dx)}} dx + \int \right)}{\tan^{5/2}(c + dx)}$$

input `integrate((a+b*tan(d*x+c))**(5/2)*(3/2*b*B/a+B*tan(d*x+c))/tan(d*x+c)**(5/2),x)`

output `B*(Integral(2*a**3*sqrt(a + b*tan(c + d*x))/tan(c + d*x)**(3/2), x) + Integral(3*b**3*sqrt(a + b*tan(c + d*x))/sqrt(tan(c + d*x)), x) + Integral(6*a*b**2*sqrt(a + b*tan(c + d*x))/tan(c + d*x)**(3/2), x) + Integral(2*a*b**2*sqrt(a + b*tan(c + d*x))*sqrt(tan(c + d*x)), x) + Integral(3*a**2*b*sqrt(a + b*tan(c + d*x))/tan(c + d*x)**(5/2), x) + Integral(4*a**2*b*sqrt(a + b*tan(c + d*x))/sqrt(tan(c + d*x)), x))/(2*a)`

3.451. $\int \frac{(a+b \tan(c+dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c+dx) \right)}{\tan^{5/2}(c+dx)} dx$

3.451.7 Maxima [F]

$$\int \frac{(a + b \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx = \int \frac{(2B \tan(dx + c) + \frac{3Bb}{a})(b \tan(dx + c) + a)^{5/2}}{2 \tan(dx + c)^{5/2}} dx$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(3/2*b*B/a+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="maxima")`

output `1/2*integrate((2*B*tan(d*x + c) + 3*B*b/a)*(b*tan(d*x + c) + a)^(5/2)/tan(d*x + c)^(5/2), x)`

3.451.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(3/2*b*B/a+B*tan(d*x+c))/tan(d*x+c)^(5/2),x, algorithm="giac")`

output `Timed out`

3.451.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx = \int \frac{(B \tan(c + dx) + \frac{3Bb}{2a})(a + b \tan(c + dx))^{5/2}}{\tan(c + dx)^{5/2}} dx$$

input `int(((B*tan(c + d*x) + (3*B*b)/(2*a))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(5/2), x)`

output `int(((B*tan(c + d*x) + (3*B*b)/(2*a))*(a + b*tan(c + d*x))^(5/2))/tan(c + d*x)^(5/2), x)`

3.451. $\int \frac{(a + b \tan(c + dx))^{5/2} \left(\frac{3bB}{2a} + B \tan(c + dx) \right)}{\tan^{5/2}(c + dx)} dx$

3.452
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

3.452.1 Optimal result 4364
 3.452.2 Mathematica [A] (verified) 4365
 3.452.3 Rubi [A] (verified) 4365
 3.452.4 Maple [B] (warning: unable to verify) 4368
 3.452.5 Fricas [B] (verification not implemented) 4368
 3.452.6 Sympy [F] 4368
 3.452.7 Maxima [F] 4369
 3.452.8 Giac [F(-1)] 4369
 3.452.9 Mupad [F(-1)] 4369

3.452.1 Optimal result

Integrand size = 35, antiderivative size = 206

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = -\frac{(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia-bd}} + \frac{(2Ab-aB) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{b^{3/2}d} - \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia+bd}} + \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}}{bd}$$

output

```
(2*A*b-B*a)*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/b^(3/2)/d-(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d/(I*a-b)^(1/2)-(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d/(I*a+b)^(1/2)+B*tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/b/d
```

3.452.2 Mathematica [A] (verified)

Time = 2.01 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.19

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

$$= \frac{\sqrt[4]{-1}b(iA+B)\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{-a+ib}} + \frac{(-1)^{3/4}b(A+iB)\arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{a+ib}} + \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd}$$

input `Integrate[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `(((-1)^(1/4)*b*(I*A + B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[-a + I*b] + ((-1)^(3/4)*b*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[a + I*b] + B*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]] + (Sqrt[a]*(2*A*b - a*B)*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/(Sqrt[b]*Sqrt[a + b*Tan[c + d*x]])/(b*d)`

3.452.3 Rubi [A] (verified)Time = 0.95 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 4090, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)^{3/2}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

$$\downarrow \text{4090}$$

$$\frac{\int -\frac{((2Ab-aB)\tan^2(c+dx))+2bB\tan(c+dx)+aB}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{b} + \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd}$$

3.452. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

$$\begin{aligned}
 & \int \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd} - \frac{\int \frac{-((2Ab-aB)\tan^2(c+dx))+2bB\tan(c+dx)+aB}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{2b} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd} - \frac{\int \frac{-((2Ab-aB)\tan(c+dx)^2)+2bB\tan(c+dx)+aB}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{2b} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd} - \frac{\int \frac{-((2Ab-aB)\tan^2(c+dx))+2bB\tan(c+dx)+aB}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} d\tan(c+dx)}{2bd} \\
 & \quad \downarrow \text{4138} \\
 & \int \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd} - \frac{\int \frac{-((2Ab-aB)\tan^2(c+dx))+2bB\tan(c+dx)+aB}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} d\sqrt{\tan(c+dx)}}{bd} \\
 & \quad \downarrow \text{2035} \\
 & \int \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd} - \frac{\int \left(\frac{aB-2Ab}{\sqrt{a+b\tan(c+dx)}} + \frac{2(Ab+B\tan(c+dx)b)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} \right) d\sqrt{\tan(c+dx)}}{bd} \\
 & \quad \downarrow \text{2257} \\
 & \frac{b(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{-b+ia}} - \frac{(2Ab-aB)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{b}} + \frac{b(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{b+ia}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd} - \frac{b(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{-b+ia}} - \frac{(2Ab-aB)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{b}} + \frac{b(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{b+ia}}
 \end{aligned}$$

input `Int[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `-((b*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[I*a - b] - ((2*A*b - a*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[b] + (b*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[I*a + b]/(b*d) + (B*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/(b*d)`

3.452. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

3.452.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2035 `Int[(F_x_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[Fx, x]`
- rule 2257 `Int[(P_x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`
- rule 4138 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

$$3.452. \int \frac{\tan^3(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

3.452.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.95 (sec) , antiderivative size = 1889462, normalized size of antiderivative = 9172.15

output too large to display

input `int(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x)`

output `result too large to display`

3.452.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10163 vs. $2(166) = 332$.

Time = 4.85 (sec) , antiderivative size = 20328, normalized size of antiderivative = 98.68

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="fricas")`

output `Too large to include`

3.452.6 Sympy [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(A+B\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx$$

input `integrate(tan(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**(3/2)/sqrt(a + b*tan(c + d*x))
, x)`

3.452. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

3.452.7 Maxima [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^{\frac{3}{2}}}{\sqrt{b\tan(dx+c)+a}} dx$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^(3/2)/sqrt(b*tan(d*x + c) + a), x)`

3.452.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.452.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{\tan(c+dx)^{3/2}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

input `int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x)`

output `int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x)`

3.452. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

3.453 $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

3.453.1 Optimal result 4370
 3.453.2 Mathematica [A] (verified) 4371
 3.453.3 Rubi [A] (verified) 4371
 3.453.4 Maple [B] (warning: unable to verify) 4375
 3.453.5 Fricas [B] (verification not implemented) 4375
 3.453.6 Sympy [F] 4376
 3.453.7 Maxima [F] 4376
 3.453.8 Giac [F(-1)] 4376
 3.453.9 Mupad [B] (verification not implemented) 4377

3.453.1 Optimal result

Integrand size = 35, antiderivative size = 168

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = \frac{(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia - bd}} + \frac{2B \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{bd}} - \frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia + bd}}$$

```
output (I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d/(I
*a-b)^(1/2)+2*B*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d
/b^(1/2)-(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(
1/2))/d/(I*a+b)^(1/2)
```

3.453.2 Mathematica [A] (verified)

Time = 1.40 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.22

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

$$= \frac{\sqrt[4]{-1} \left(-\frac{(A-iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} + \frac{(A+iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{a+ib}} \right) + \frac{2\sqrt{a} \operatorname{Barcsinh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)}{\sqrt{b}\sqrt{a+b \tan(c+dx)}}}{d}$$

```
input Integrate[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]
```

```
output ((-1)^(1/4)*(-(((A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[-a + I*b]) + ((A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[a + I*b]) + (2*Sqrt[a]*B*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/(Sqrt[b]*Sqrt[a + b*Tan[c + d*x]]))/d
```

3.453.3 Rubi [A] (verified)

Time = 1.09 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 4097, 3042, 4099, 3042, 4098, 104, 216, 219, 4117, 65, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

↓ 3042

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

↓ 4097

$$\int \frac{A \tan(c+dx) - B}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx + B \int \frac{\tan^2(c+dx) + 1}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx$$

$$\begin{aligned}
& \downarrow 3042 \\
& \int \frac{A \tan(c+dx) - B}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx + B \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx \\
& \downarrow 4099 \\
& \frac{1}{2}(-B+iA) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2}(B+iA) \int \frac{i \tan(c+dx) + 1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx + B \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx \\
& \downarrow 3042 \\
& \frac{1}{2}(-B+iA) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2}(B+iA) \int \frac{i \tan(c+dx) + 1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx + B \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx \\
& \downarrow 4098 \\
& \frac{(B+iA) \int \frac{1}{(1-i \tan(c+dx)) \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{2d} + \\
& \frac{(-B+iA) \int \frac{1}{(i \tan(c+dx)+1) \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{2d} + \\
& B \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx \\
& \downarrow 104 \\
& \frac{(-B+iA) \int \frac{1}{\frac{(ia-b) \tan(c+dx)}{a+b \tan(c+dx)} + 1} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} - \frac{(B+iA) \int \frac{1}{1 - \frac{(ia+b) \tan(c+dx)}{a+b \tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} + \\
& B \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx \\
& \downarrow 216 \\
& - \frac{(B+iA) \int \frac{1}{1 - \frac{(ia+b) \tan(c+dx)}{a+b \tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} + B \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx + \\
& \frac{(-B+iA) \arctan \left(\frac{\sqrt{-b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right)}{d \sqrt{-b+ia}} \\
& \downarrow 219
\end{aligned}$$

3.453. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\begin{aligned}
& B \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{(-B+iA) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} - \\
& \quad \frac{(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}} \\
& \quad \downarrow 4117 \\
& B \int \frac{1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} d\tan(c+dx) + \frac{(-B+iA) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} - \\
& \quad \frac{(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}} \\
& \quad \downarrow 65 \\
& 2B \int \frac{1}{1-\frac{b\tan(c+dx)}{a+b\tan(c+dx)}} d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}} + \frac{(-B+iA) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} - \\
& \quad \frac{(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}} \\
& \quad \downarrow 219 \\
& \frac{(-B+iA) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} - \frac{(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}} + \\
& \quad \frac{2B \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{bd}}
\end{aligned}$$

input `Int[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `((I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a - b]*d) + (2*B*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[b]*d) - ((I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d)`

3.453.3.1 Defintions of rubi rules used

- rule 65 `Int[1/(Sqrt[(b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`
- rule 104 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4097 `Int[(Sqrt[(a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]))/Sqrt[(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Int[Simp[a*A - b*B + (A*b + a*B)*Tan[e + f*x], x]/(Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]), x] + Simp[b*B Int[(1 + Tan[e + f*x]^2)/(Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

```
rule 4099 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2,
0]
```

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

3.453.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.95 (sec) , antiderivative size = 1885950, normalized size of antiderivative = 11225.89

output too large to display

```
input int(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x)
```

```
output result too large to display
```

3.453.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10116 vs. $2(132) = 264$.

Time = 4.15 (sec) , antiderivative size = 20234, normalized size of antiderivative = 120.44

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="fricas")
```

```
output Too large to include
```

3.453. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

3.453.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(A+B\tan(c+dx))\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}} dx$$

input `integrate(tan(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(tan(c + d*x))/sqrt(a + b*tan(c + d*x)), x)`

3.453.7 Maxima [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\sqrt{\tan(dx+c)}}{\sqrt{b\tan(dx+c)+a}} dx$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(tan(d*x + c))/sqrt(b*tan(d*x + c) + a), x)`

3.453.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Timed out`

3.453.9 Mupad [B] (verification not implemented)

Time = 103.96 (sec) , antiderivative size = 30600, normalized size of antiderivative = 182.14

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Too large to display}$$

```
input int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x
)
```

```
output atan((((B^2 - A^2 + A*B*2i)/(4*(a*d^2*i + b*d^2)))^(1/2)*(((B^2 - A^2 +
A*B*2i)/(4*(a*d^2*i + b*d^2)))^(1/2)*(((B^2 - A^2 + A*B*2i)/(4*(a*d^2*i
i + b*d^2)))^(1/2)*(((274877906944*(1600*a^12*b^34*d^8 - 16640*a^14*b^32*
d^8 + 22784*a^16*b^30*d^8 + 106496*a^18*b^28*d^8 + 65536*a^20*b^26*d^8))/d
^8 - (274877906944*tan(c + d*x)*(1600*a^12*b^35*d^8 - 48000*a^14*b^33*d^8
+ 155136*a^16*b^31*d^8 + 466944*a^18*b^29*d^8 + 262144*a^20*b^27*d^8))/(d^
8*((a + b*tan(c + d*x))^(1/2) - a^(1/2))^2))*((B^2 - A^2 + A*B*2i)/(4*(a*d
^2*i + b*d^2)))^(1/2) - (219902325552*tan(c + d*x)^(1/2)*(240*A*a^13*b^3
3*d^6 + 3064*A*a^15*b^31*d^6 + 8960*A*a^17*b^29*d^6 + 6144*A*a^19*b^27*d^6
+ 6920*B*a^14*b^32*d^6 + 9472*B*a^16*b^30*d^6 - 5632*B*a^18*b^28*d^6 - 81
92*B*a^20*b^26*d^6))/(d^7*((a + b*tan(c + d*x))^(1/2) - a^(1/2))))*((B^2 -
A^2 + A*B*2i)/(4*(a*d^2*i + b*d^2)))^(1/2) - (274877906944*(19216*A^2*a^
14*b^31*d^6 - 1440*A^2*a^12*b^33*d^6 - 22016*A^2*a^16*b^29*d^6 - 45056*A^2
*a^18*b^27*d^6 + 1200*B^2*a^12*b^33*d^6 - 16640*B^2*a^14*b^31*d^6 + 279040
*B^2*a^16*b^29*d^6 + 561152*B^2*a^18*b^27*d^6 + 262144*B^2*a^20*b^25*d^6 +
16480*A*B*a^13*b^32*d^6 - 25792*A*B*a^15*b^30*d^6 + 34816*A*B*a^17*b^28*d
^6 + 81920*A*B*a^19*b^26*d^6))/d^8 + (274877906944*tan(c + d*x)*(46704*A^2
*a^14*b^32*d^6 - 1440*A^2*a^12*b^34*d^6 - 137216*A^2*a^16*b^30*d^6 - 12288
0*A^2*a^18*b^28*d^6 + 65536*A^2*a^20*b^26*d^6 + 1200*B^2*a^12*b^34*d^6 - 5
2320*B^2*a^14*b^32*d^6 + 1200640*B^2*a^16*b^30*d^6 + 2306048*B^2*a^18*b...
```


3.454
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx$$

3.454.1 Optimal result 4378
 3.454.2 Mathematica [A] (verified) 4378
 3.454.3 Rubi [A] (verified) 4379
 3.454.4 Maple [B] (warning: unable to verify) 4381
 3.454.5 Fricas [B] (verification not implemented) 4381
 3.454.6 Sympy [F] 4382
 3.454.7 Maxima [F] 4382
 3.454.8 Giac [F(-1)] 4383
 3.454.9 Mupad [B] (verification not implemented) 4383

3.454.1 Optimal result

Integrand size = 35, antiderivative size = 123

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx = \frac{(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia - bd}} + \frac{(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia + bd}}$$

```
output (A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d/(I
*a-b)^(1/2)+(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)
)^(1/2))/d/(I*a+b)^(1/2)
```

3.454.2 Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.11

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx = \frac{\sqrt[4]{-1} \left(-\frac{(iA+B) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} + \frac{(-iA+B) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{a+ib}} \right)}{d}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]]),x]`

output `((-1)^(1/4)*(-(((I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[-a + I*b]) + (((-I)*A + B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[a + I*b]))/d`

3.454.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4099} \\
 & \frac{1}{2}(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2}(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4098} \\
 & \frac{(A - iB) \int \frac{1}{(1 - i \tan(c + dx)) \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} d \tan(c + dx)}{2d} + \frac{(A + iB) \int \frac{1}{(i \tan(c + dx) + 1) \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} d \tan(c + dx)}{2d}
 \end{aligned}$$

3.454. $\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx$

$$\begin{aligned}
 & \downarrow 104 \\
 & \frac{(A + iB) \int \frac{1}{\frac{(ia-b)\tan(c+dx)}{a+b\tan(c+dx)} + 1} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} + \frac{(A - iB) \int \frac{1}{1 - \frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} \\
 & \downarrow 216 \\
 & \frac{(A - iB) \int \frac{1}{1 - \frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} + \frac{(A + iB) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} \\
 & \downarrow 219 \\
 & \frac{(A + iB) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} + \frac{(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}}
 \end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]]),x]`

output `((A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a - b]*d) + ((A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d)`

3.454.3.1 Defintions of rubi rules used

rule 104 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

3.454.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.92 (sec) , antiderivative size = 1878820, normalized size of antiderivative = 15274.96

output too large to display

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2),x)`

output `result too large to display`

3.454.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 9913 vs. 2(95) = 190.

Time = 2.98 (sec) , antiderivative size = 9913, normalized size of antiderivative = 80.59

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

3.454. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="fricas")`

output Too large to include

3.454.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)} \sqrt{\tan(c + dx)}} dx$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+b*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))/(sqrt(a + b*tan(c + d*x))*sqrt(tan(c + d*x))
, x)`

3.454.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx = \int \frac{B \tan(dx + c) + A}{\sqrt{b \tan(dx + c) + a} \sqrt{\tan(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)/(sqrt(b*tan(d*x + c) + a)*sqrt(tan(d*x + c)
)), x)`

3.454.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="giac")`

output `Timed out`

3.454.9 Mupad [B] (verification not implemented)

Time = 59.96 (sec) , antiderivative size = 8223, normalized size of antiderivative = 66.85

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(1/2)),x
)`

output

$$\begin{aligned} & \operatorname{atan}\left(\frac{(B^5 b^9 d^7 \tan(c+dx))^{1/2} \left(-((-16B^4 a^2 d^4)^{1/2} + 4B^2 b d^2) / (16a^2 d^4 + 16b^2 d^4)\right)^{1/2} 1280i}{(a + b \tan(c+dx))^{1/2} - a^{1/2}}\right) \\ & + \frac{(B^3 b^{10} d^9 \tan(c+dx))^{1/2} \left(-((-16B^4 a^2 d^4)^{1/2} + 4B^2 b d^2) / (16a^2 d^4 + 16b^2 d^4)\right)^{3/2} 10240i}{(a + b \tan(c+dx))^{1/2} - a^{1/2}} \\ & + \frac{(B b^{11} d^{11} \tan(c+dx))^{1/2} \left(-((-16B^4 a^2 d^4)^{1/2} + 4B^2 b d^2) / (16a^2 d^4 + 16b^2 d^4)\right)^{5/2} 20480i}{(a + b \tan(c+dx))^{1/2} - a^{1/2}} \\ & + \frac{(B a^{10} b d^{11} \tan(c+dx))^{1/2} \left(-((-16B^4 a^2 d^4)^{1/2} + 4B^2 b d^2) / (16a^2 d^4 + 16b^2 d^4)\right)^{5/2} 196608i}{(a + b \tan(c+dx))^{1/2} - a^{1/2}} \\ & + \frac{(B^5 a^8 b d^7 \tan(c+dx))^{1/2} \left(-((-16B^4 a^2 d^4)^{1/2} + 4B^2 b d^2) / (16a^2 d^4 + 16b^2 d^4)\right)^{1/2} 12288i}{(a + b \tan(c+dx))^{1/2} - a^{1/2}} \\ & + \frac{(B a^2 b^9 d^{11} \tan(c+dx))^{1/2} \left(-((-16B^4 a^2 d^4)^{1/2} + 4B^2 b d^2) / (16a^2 d^4 + 16b^2 d^4)\right)^{5/2} 274432i}{(a + b \tan(c+dx))^{1/2} - a^{1/2}} \\ & + \frac{(B a^4 b^7 d^{11} \tan(c+dx))^{1/2} \left(-((-16B^4 a^2 d^4)^{1/2} + 4B^2 b d^2) / (16a^2 d^4 + 16b^2 d^4)\right)^{5/2} 897024i}{(a + b \tan(c+dx))^{1/2} - a^{1/2}} \\ & + \frac{(B a^6 b^5 d^{11} \tan(c+dx))^{1/2} \left(-((-16B^4 a^2 d^4)^{1/2} + 4B^2 b d^2) / (16a^2 d^4 + 16b^2 d^4)\right)^{5/2} 1249280i}{(a + b \tan(c+dx))^{1/2} - a^{1/2}} \\ & + \frac{(B a^8 b^3 d^{11} \tan(c+dx))^{1/2} \left(-((-16B^4 a^2 d^4)^{1/2} + 4B^2 b d^2) / (16a^2 d^4 + 16b^2 d^4)\right)^{5/2} 802816i}{(a + b \tan(c+dx))^{1/2} - a^{1/2}} \\ & + \frac{(B^5 a^2 b^7 d^7 \tan(c+dx))^{1/2} \left(-((-16B^4 a^2 d^4)^{1/2} + 4B^2 b d^2) / (16a^2 d^4 + 16b^2 d^4)\right)^{1/2} 1280i}{(a + b \tan(c+dx))^{1/2} - a^{1/2}} \end{aligned}$$

$$3.455 \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx$$

3.455.1 Optimal result	4385
3.455.2 Mathematica [A] (verified)	4385
3.455.3 Rubi [A] (verified)	4386
3.455.4 Maple [B] (warning: unable to verify)	4389
3.455.5 Fricas [B] (verification not implemented)	4390
3.455.6 Sympy [F]	4390
3.455.7 Maxima [F]	4390
3.455.8 Giac [F(-1)]	4391
3.455.9 Mupad [F(-1)]	4391

3.455.1 Optimal result

Integrand size = 35, antiderivative size = 159

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)\sqrt{a + b \tan(c + dx)}} dx = -\frac{(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia - bd}} + \frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia + bd}} - \frac{2A\sqrt{a + b \tan(c + dx)}}{ad\sqrt{\tan(c + dx)}}$$

output $-(I*A-B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})/d/(I*a-b)^{(1/2)}+(I*A+B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})/d/(I*a+b)^{(1/2)}-2*A*(a+b*\tan(d*x+c))^{(1/2)}/a/d/\tan(d*x+c)^{(1/2)}$

3.455.2 Mathematica [A] (verified)

Time = 0.55 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.08

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)\sqrt{a + b \tan(c + dx)}} dx = \frac{\sqrt[4]{-1}(A-iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} - \frac{\sqrt[4]{-1}(A+iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{a+ib}} - \frac{2A\sqrt{a+b \tan(c+dx)}}{a\sqrt{\tan(c+dx)}}$$

3.455. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]]),x]`

output `(((-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[-a + I*b] - ((-1)^(1/4)*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[a + I*b] - (2*A*Sqrt[a + b*Tan[c + d*x]])/(a*Sqrt[Tan[c + d*x]]))/d`

3.455.3 Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.04, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4092, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2} \sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4092} \\
 & \frac{2 \int -\frac{aB - aA \tan(c + dx)}{2\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx}{a} - \frac{2A \sqrt{a + b \tan(c + dx)}}{ad \sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{aB - aA \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx}{a} - \frac{2A \sqrt{a + b \tan(c + dx)}}{ad \sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{aB - aA \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx}{a} - \frac{2A \sqrt{a + b \tan(c + dx)}}{ad \sqrt{\tan(c + dx)}} \\
 & \quad \downarrow \text{4099}
 \end{aligned}$$

3.455. $\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx$

$$\begin{aligned}
& \frac{2A\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \\
& \frac{\frac{1}{2}a(B+iA)\int\frac{i\tan(c+dx)+1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx - \frac{1}{2}a(-B+iA)\int\frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx}{a} \\
& \quad \downarrow 3042 \\
& \frac{2A\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \\
& \frac{\frac{1}{2}a(B+iA)\int\frac{i\tan(c+dx)+1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx - \frac{1}{2}a(-B+iA)\int\frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx}{a} \\
& \quad \downarrow 4098 \\
& \frac{2A\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \\
& \frac{a(B+iA)\int\frac{1}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}d\tan(c+dx) - a(-B+iA)\int\frac{1}{(i\tan(c+dx)+1)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}d\tan(c+dx)}{2d} \\
& \quad \downarrow 104 \\
& \frac{2A\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \\
& \frac{a(B+iA)\int\frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}} - a(-B+iA)\int\frac{1}{\frac{(ia-b)\tan(c+dx)}{a+b\tan(c+dx)}+1}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} \\
& \quad \downarrow 216 \\
& \frac{2A\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \\
& \frac{a(B+iA)\int\frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}} - a(-B+iA)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d} \\
& \quad \downarrow 219 \\
& -\frac{2A\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{a(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}} - \frac{a(-B+iA)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}}
\end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]]),x]`

3.455. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}} dx$

```
output 
$$\frac{-((a*(I*A - B)*\text{ArcTan}[(\text{Sqrt}[I*a - b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a + b*\text{Tan}[c + d*x]]))/(\text{Sqrt}[I*a - b]*d) + (a*(I*A + B)*\text{ArcTanh}[(\text{Sqrt}[I*a + b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a + b*\text{Tan}[c + d*x]]))/(\text{Sqrt}[I*a + b]*d))/a - (2*A*\text{Sqrt}[a + b*\text{Tan}[c + d*x]])/(a*d*\text{Sqrt}[\text{Tan}[c + d*x]])}$$

```

3.455.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 104 Int[(((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4092 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)
/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^
2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*
B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n +
2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n
+ 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1]
&& (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m]
|| (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4098 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*
x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]
```

```
rule 4099 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2,
0]
```

3.455.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.92 (sec) , antiderivative size = 1886236, normalized size of antiderivative = 11863.12

output too large to display

```
input int((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2)/tan(d*x+c)^(3/2),x)
```

```
output result too large to display
```

$$3.455. \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx$$

3.455.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10036 vs. $2(124) = 248$.

Time = 2.98 (sec) , antiderivative size = 10036, normalized size of antiderivative = 63.12

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2)/tan(d*x+c)^(3/2),x, algo
rithm="fricas")`

output Too large to include

3.455.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)} \tan^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2)/tan(d*x+c)**(3/2),x)`

output `Integral((A + B*tan(c + d*x))/(sqrt(a + b*tan(c + d*x))*tan(c + d*x)**(3/2)), x)`

3.455.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \int \frac{B \tan(dx + c) + A}{\sqrt{b \tan(dx + c) + a} \tan(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2)/tan(d*x+c)^(3/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)/(sqrt(b*tan(d*x + c) + a)*tan(d*x + c)^(3/2)), x)`

3.455.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2)/tan(d*x+c)^(3/2),x, algorith="giac")`

output `Timed out`

3.455.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2} \sqrt{a + b \tan(c + dx)}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(1/2)),x)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(1/2)),x)`

3.456
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx$$

3.456.1 Optimal result	4392
3.456.2 Mathematica [A] (verified)	4393
3.456.3 Rubi [A] (verified)	4393
3.456.4 Maple [B] (warning: unable to verify)	4397
3.456.5 Fricas [B] (verification not implemented)	4398
3.456.6 Sympy [F]	4398
3.456.7 Maxima [F]	4398
3.456.8 Giac [F(-1)]	4399
3.456.9 Mupad [F(-1)]	4399

3.456.1 Optimal result

Integrand size = 35, antiderivative size = 203

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)\sqrt{a + b \tan(c + dx)}} dx = -\frac{(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia - bd}} - \frac{(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia + bd}} - \frac{2A\sqrt{a + b \tan(c + dx)}}{3ad \tan^{\frac{3}{2}}(c + dx)} + \frac{2(2Ab - 3aB)\sqrt{a + b \tan(c + dx)}}{3a^2d\sqrt{\tan(c + dx)}}$$

output
$$-(A+I*B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})/d/(I*a-b)^{(1/2)}-(A-I*B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})/d/(I*a+b)^{(1/2)}+2/3*(2*A*b-3*B*a)*(a+b*\tan(d*x+c))^{(1/2)}/a^2/d/\tan(d*x+c)^{(1/2)}-2/3*A*(a+b*\tan(d*x+c))^{(1/2)}/a/d/\tan(d*x+c)^{(3/2)}$$

3.456.2 Mathematica [A] (verified)

Time = 2.32 (sec) , antiderivative size = 195, normalized size of antiderivative = 0.96

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx$$

$$= \frac{3 \sqrt[4]{-1} (iA+B) \arctan\left(\frac{\sqrt[4]{-1} \sqrt{-a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} + \frac{3(-1)^{3/4} (A+iB) \arctan\left(\frac{\sqrt[4]{-1} \sqrt{a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{a+ib}} - \frac{2\sqrt{a+b \tan(c+dx)}(aA+(-2A^2+3aB)\tan(c+dx))}{a^2 \tan^{\frac{3}{2}}(c+dx)}$$

$$\frac{\hspace{10em}}{3d}$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*Sqrt[a + b*Tan[c + d*x]]),x]`

output `((3*(-1)^(1/4)*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[-a + I*b] + (3*(-1)^(3/4)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[a + I*b] - (2*Sqrt[a + b*Tan[c + d*x]]*(a*A + (-2*A*b + 3*a*B)*Tan[c + d*x]))/(a^2*Tan[c + d*x]^(3/2))/(3*d)`

3.456.3 Rubi [A] (verified)

Time = 1.15 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.08, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2} \sqrt{a + b \tan(c + dx)}} dx$$

↓ 4092

$$-\frac{2 \int \frac{2Ab \tan^2(c+dx)+3aA \tan(c+dx)+2Ab-3aB}{2 \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2A \sqrt{a + b \tan(c + dx)}}{3ad \tan^{\frac{3}{2}}(c + dx)}$$

↓ 27

3.456. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx$

$$\begin{aligned}
 & \frac{\int \frac{2Ab \tan^2(c+dx) + 3aA \tan(c+dx) + 2Ab - 3aB}{\tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2A \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{2Ab \tan(c+dx)^2 + 3aA \tan(c+dx) + 2Ab - 3aB}{\tan(c+dx)^{3/2} \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2A \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow \text{4132} \\
 & \frac{2 \int -\frac{3(Aa^2 + B \tan(c+dx)a^2)}{2\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(2Ab - 3aB) \sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} - \frac{2A \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow \text{27} \\
 & \frac{3 \int \frac{Aa^2 + B \tan(c+dx)a^2}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(2Ab - 3aB) \sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} - \frac{2A \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3 \int \frac{Aa^2 + B \tan(c+dx)a^2}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(2Ab - 3aB) \sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} - \frac{2A \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow \text{4099} \\
 & \frac{2A \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} - \frac{2(2Ab - 3aB) \sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} + \frac{3 \left(\frac{1}{2} a^2 (A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2} a^2 (A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx \right)}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2A \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} - \frac{2(2Ab - 3aB) \sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} + \frac{3 \left(\frac{1}{2} a^2 (A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2} a^2 (A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx \right)}{a} \\
 & \quad \downarrow \text{4098}
 \end{aligned}$$

3.456. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx$

$$\frac{-\frac{2(2Ab-3aB)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{2A\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)}}{3a} + \frac{3\left(\frac{a^2(A-iB)\int\frac{1}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}d\tan(c+dx)}{2d} + \frac{a^2(A+iB)\int\frac{1}{(i\tan(c+dx)+1)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}d\tan(c+dx)}{2d}\right)}{a}$$

104

$$\frac{-\frac{2(2Ab-3aB)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{2A\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)}}{3a} + \frac{3\left(\frac{a^2(A+iB)\int\frac{1}{\frac{(ia-b)\tan(c+dx)}{a+b\tan(c+dx)}+1}d-\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} + \frac{a^2(A-iB)\int\frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}}d-\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d}\right)}{a}$$

216

$$\frac{-\frac{2(2Ab-3aB)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{2A\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)}}{3a} + \frac{3\left(\frac{a^2(A-iB)\int\frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}}d-\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} + \frac{a^2(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}}\right)}{a}$$

219

$$\frac{-\frac{2(2Ab-3aB)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{2A\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)}}{3a} + \frac{3\left(\frac{a^2(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} + \frac{a^2(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}}\right)}{a}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*Sqrt[a + b*Tan[c + d*x]]),x]`

output `(-2*A*Sqrt[a + b*Tan[c + d*x]])/(3*a*d*Tan[c + d*x]^(3/2)) - ((3*((a^2*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a - b]*d) + (a^2*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d)))/a - (2*(2*A*b - 3*a*B)*Sqrt[a + b*Tan[c + d*x]]/(a*d*Sqrt[Tan[c + d*x]]))/(3*a)`

3.456.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && ! (ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.456.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.89 (sec) , antiderivative size = 1888895, normalized size of antiderivative = 9304.90

output too large to display

input `int((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2)/tan(d*x+c)^(5/2),x)`

output `result too large to display`

$$3.456. \quad \int \frac{A+B \tan(c+dx)}{\tan^2(c+dx) \sqrt{a+b \tan(c+dx)}} dx$$

3.456.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10067 vs. $2(163) = 326$.

Time = 3.63 (sec) , antiderivative size = 10067, normalized size of antiderivative = 49.59

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2)/tan(d*x+c)^(5/2),x, algo
rithm="fricas")`

output Too large to include

3.456.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)} \tan^{\frac{5}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2)/tan(d*x+c)**(5/2),x)`

output `Integral((A + B*tan(c + d*x))/(sqrt(a + b*tan(c + d*x))*tan(c + d*x)**(5/2
)), x)`

3.456.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \int \frac{B \tan(dx + c) + A}{\sqrt{b \tan(dx + c) + a} \tan(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2)/tan(d*x+c)^(5/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)/(sqrt(b*tan(d*x + c) + a)*tan(d*x + c)^(5/2
)), x)`

3.456.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2)/tan(d*x+c)^(5/2),x, algorith="giac")`

output `Timed out`

3.456.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2} \sqrt{a + b \tan(c + dx)}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + b*tan(c + d*x))^(1/2)),x)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + b*tan(c + d*x))^(1/2)),x)`

3.457
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{7}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx$$

3.457.1 Optimal result 4400
 3.457.2 Mathematica [A] (verified) 4401
 3.457.3 Rubi [A] (verified) 4401
 3.457.4 Maple [B] (warning: unable to verify) 4406
 3.457.5 Fricas [B] (verification not implemented) 4407
 3.457.6 Sympy [F] 4407
 3.457.7 Maxima [F] 4407
 3.457.8 Giac [F(-1)] 4408
 3.457.9 Mupad [F(-1)] 4408

3.457.1 Optimal result

Integrand size = 35, antiderivative size = 256

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx)\sqrt{a + b \tan(c + dx)}} dx = \frac{(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia - bd}} - \frac{(iA + B)\operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia + bd}} - \frac{2A\sqrt{a + b \tan(c + dx)}}{5ad \tan^{\frac{5}{2}}(c + dx)} + \frac{2(4Ab - 5aB)\sqrt{a + b \tan(c + dx)}}{15a^2d \tan^{\frac{3}{2}}(c + dx)} + \frac{2(15a^2A - 8Ab^2 + 10abB)\sqrt{a + b \tan(c + dx)}}{15a^3d\sqrt{\tan(c + dx)}}$$

output $(I*A-B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})/d/(I*a-b)^{(1/2)}-(I*A+B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})/d/(I*a+b)^{(1/2)}+2/15*(15*A*a^2-8*A*b^2+10*B*a*b)*(a+b*\tan(d*x+c))^{(1/2)}/a^3/d/\tan(d*x+c)^{(1/2)}-2/5*A*(a+b*\tan(d*x+c))^{(1/2)}/a/d/\tan(d*x+c)^{(5/2)}+2/15*(4*A*b-5*B*a)*(a+b*\tan(d*x+c))^{(1/2)}/a^2/d/\tan(d*x+c)^{(3/2)}$

3.457.2 Mathematica [A] (verified)

Time = 5.79 (sec) , antiderivative size = 227, normalized size of antiderivative = 0.89

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx$$

$$= \frac{15 \sqrt[4]{-1} (A - iB) \arctan\left(\frac{\sqrt[4]{-1} \sqrt{-a + ib} \sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{\sqrt{-a + ib}} + \frac{15 \sqrt[4]{-1} (A + iB) \arctan\left(\frac{\sqrt[4]{-1} \sqrt{a + ib} \sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{\sqrt{a + ib}} + \frac{2 \sqrt{a + b \tan(c + dx)} (-3a + b \tan(c + dx))}{15d}$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(7/2)*Sqrt[a + b*Tan[c + d*x]]),x]`

output `((-15*(-1)^(1/4)*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[-a + I*b] + (15*(-1)^(1/4)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[a + I*b] + (2*Sqrt[a + b*Tan[c + d*x]]*(-3*a^2*A - a*(-4*A*b + 5*a*B)*Tan[c + d*x] + (15*a^2*A - 8*A*b^2 + 10*a*b*B)*Tan[c + d*x]^2))/(a^3*Tan[c + d*x]^(5/2)))/(15*d)`

3.457.3 Rubi [A] (verified)

Time = 1.67 (sec) , antiderivative size = 282, normalized size of antiderivative = 1.10, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{7/2} \sqrt{a + b \tan(c + dx)}} dx$$

↓ 4092

3.457. $\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx$

$$\begin{array}{c}
\frac{2 \int \frac{4Ab \tan^2(c+dx) + 5aA \tan(c+dx) + 4Ab - 5aB}{2 \tan^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2A \sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} \\
\downarrow 27 \\
\frac{\int \frac{4Ab \tan^2(c+dx) + 5aA \tan(c+dx) + 4Ab - 5aB}{\tan^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2A \sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} \\
\downarrow 3042 \\
\frac{\int \frac{4Ab \tan(c+dx)^2 + 5aA \tan(c+dx) + 4Ab - 5aB}{\tan(c+dx)^{5/2} \sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2A \sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} \\
\downarrow 4132 \\
\frac{2 \int -\frac{15Aa^2 + 15B \tan(c+dx)a^2 + 10bBa - 8Ab^2 - 2b(4Ab - 5aB) \tan^2(c+dx)}{2 \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(4Ab - 5aB) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
\hline
\frac{5a}{2A \sqrt{a+b \tan(c+dx)}} \\
\frac{2A \sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} \\
\downarrow 27 \\
\frac{\int \frac{15Aa^2 + 15B \tan(c+dx)a^2 + 10bBa - 8Ab^2 - 2b(4Ab - 5aB) \tan^2(c+dx)}{\tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(4Ab - 5aB) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
\hline
\frac{5a}{2A \sqrt{a+b \tan(c+dx)}} \\
\frac{2A \sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} \\
\downarrow 3042 \\
\frac{\int \frac{15Aa^2 + 15B \tan(c+dx)a^2 + 10bBa - 8Ab^2 - 2b(4Ab - 5aB) \tan(c+dx)^2}{\tan(c+dx)^{3/2} \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(4Ab - 5aB) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
\hline
\frac{5a}{2A \sqrt{a+b \tan(c+dx)}} \\
\frac{2A \sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} \\
\downarrow 4132 \\
\frac{2 \int -\frac{15(a^3 B - a^3 A \tan(c+dx))}{2 \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(15a^2 A + 10abB - 8Ab^2) \sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} - \frac{2(4Ab - 5aB) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
\hline
\frac{5a}{2A \sqrt{a+b \tan(c+dx)}} \\
\frac{2A \sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)}
\end{array}$$

3.457. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{7}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{15 \int \frac{a^3 B - a^3 A \tan(c+dx)}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(15a^2 A + 10abB - 8Ab^2) \sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} - \frac{2(4Ab - 5aB) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
 & \frac{5a}{2A \sqrt{a+b \tan(c+dx)}} \\
 & \frac{5ad \tan^{\frac{5}{2}}(c+dx)}{\downarrow 3042} \\
 & \frac{15 \int \frac{a^3 B - a^3 A \tan(c+dx)}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(15a^2 A + 10abB - 8Ab^2) \sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} - \frac{2(4Ab - 5aB) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \\
 & \frac{5a}{2A \sqrt{a+b \tan(c+dx)}} \\
 & \frac{5ad \tan^{\frac{5}{2}}(c+dx)}{\downarrow 4099} \\
 & \frac{2A \sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab - 5aB) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} + \frac{2(15a^2 A + 10abB - 8Ab^2) \sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} + \frac{15 \left(\frac{1}{2} a^3 (B+iA) \int \frac{i \tan(c+dx)+1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2} a^3 (-B+iA) \int \frac{1}{\sqrt{\tan(c+dx)}} dx \right)}{3a} \\
 & \frac{5a}{\downarrow 3042} \\
 & \frac{2A \sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab - 5aB) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} + \frac{2(15a^2 A + 10abB - 8Ab^2) \sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} + \frac{15 \left(\frac{1}{2} a^3 (B+iA) \int \frac{i \tan(c+dx)+1}{\sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2} a^3 (-B+iA) \int \frac{1}{\sqrt{\tan(c+dx)}} dx \right)}{3a} \\
 & \frac{5a}{\downarrow 4098} \\
 & \frac{2A \sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab - 5aB) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} + \frac{2(15a^2 A + 10abB - 8Ab^2) \sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} + \frac{15 \left(\frac{a^3 (B+iA) \int \frac{1}{(1-i \tan(c+dx)) \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{2d} \right)}{3a} \\
 & \frac{5a}{\downarrow 104}
 \end{aligned}$$

3.457. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{7}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx$

$$\frac{-\frac{2A\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} + \frac{-\frac{2(15a^2A+10abB-8Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}{3a} + \frac{\left(\frac{a^{3(B+iA)}\int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}} - \frac{a^{3(-B+iA)}}{a}\right)}{5a}$$

216

$$\frac{-\frac{2A\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} + \frac{-\frac{2(15a^2A+10abB-8Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}{3a} + \frac{\left(\frac{a^{3(B+iA)}\int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}} - \frac{a^{3(-B+iA)}}{a}\right)}{5a}$$

219

$$\frac{-\frac{2A\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} + \frac{-\frac{2(15a^2A+10abB-8Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}{3a} + \frac{\left(\frac{a^{3(B+iA)}\operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) - \frac{a^{3(-B+iA)}\operatorname{arctan}\left(\frac{\sqrt{a-b}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}}}{5a}$$

```
input Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(7/2)*Sqrt[a + b*Tan[c + d*x]]),x]
```

```
output (-2*A*Sqrt[a + b*Tan[c + d*x]]/(5*a*d*Tan[c + d*x]^(5/2)) - ((-2*(4*A*b - 5*a*B)*Sqrt[a + b*Tan[c + d*x]]/(3*a*d*Tan[c + d*x]^(3/2)) + ((15*(-((a^3*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]))/(Sqrt[I*a - b]*d)) + (a^3*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]/(Sqrt[I*a + b]*d)))/a - (2*(15*a^2*A - 8*A*b^2 + 10*a*b*B)*Sqrt[a + b*Tan[c + d*x]]/(a*d*Sqrt[Tan[c + d*x]]))/(3*a))/(5*a)
```

3.457. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{7}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}} dx$

3.457.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && ! (ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.457.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.94 (sec) , antiderivative size = 1890924, normalized size of antiderivative = 7386.42

output too large to display

input `int((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2)/tan(d*x+c)^(7/2),x)`

output `result too large to display`

3.457.
$$\int \frac{A+B \tan(c+dx)}{\tan^2(c+dx) \sqrt{a+b \tan(c+dx)}} dx$$

3.457.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10125 vs. $2(210) = 420$.

Time = 3.06 (sec) , antiderivative size = 10125, normalized size of antiderivative = 39.55

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2)/tan(d*x+c)^(7/2),x, algo
rithm="fricas")`

output Too large to include

3.457.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)} \tan^{\frac{7}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2)/tan(d*x+c)**(7/2),x)`

output `Integral((A + B*tan(c + d*x))/(sqrt(a + b*tan(c + d*x))*tan(c + d*x)**(7/2
)), x)`

3.457.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \int \frac{B \tan(dx + c) + A}{\sqrt{b \tan(dx + c) + a} \tan(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2)/tan(d*x+c)^(7/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)/(sqrt(b*tan(d*x + c) + a)*tan(d*x + c)^(7/2
)), x)`

3.457.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2)/tan(d*x+c)^(7/2),x, algorith="giac")`

output `Timed out`

3.457.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{7/2} \sqrt{a + b \tan(c + dx)}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(7/2)*(a + b*tan(c + d*x))^(1/2)),x)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(7/2)*(a + b*tan(c + d*x))^(1/2)),x)`

3.458
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.458.1 Optimal result 4409
 3.458.2 Mathematica [A] (verified) 4410
 3.458.3 Rubi [A] (verified) 4410
 3.458.4 Maple [B] (warning: unable to verify) 4413
 3.458.5 Fricas [B] (verification not implemented) 4413
 3.458.6 Sympy [F] 4414
 3.458.7 Maxima [F] 4414
 3.458.8 Giac [F(-1)] 4414
 3.458.9 Mupad [F(-1)] 4415

3.458.1 Optimal result

Integrand size = 35, antiderivative size = 219

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx =$$

$$-\frac{(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia-b)^{3/2}d} + \frac{2B \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{b^{3/2}d}$$

$$-\frac{(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia+b)^{3/2}d} + \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{b(a^2+b^2)d\sqrt{a+b \tan(c+dx)}}$$

output

```
-(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a-b)^(3/2)/d+2*B*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/b^(3/2)/d-(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a+b)^(3/2)/d+2*a*(A*b-B*a)*tan(d*x+c)^(1/2)/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^(1/2)
```


3.458.2 Mathematica [A] (verified)

Time = 2.03 (sec) , antiderivative size = 341, normalized size of antiderivative = 1.56

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{3}{2}}} dx =$$

$$2\sqrt{a}\sqrt{-a+ib}\sqrt{a+ib}(a^2+b^2) \operatorname{Barcsinh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right) \sqrt{1+\frac{b \tan(c+dx)}{a}} + \sqrt{b}\left(\sqrt[4]{-1}(a+ib)^{\frac{3}{2}}b(iA+B)\right)$$

input `Integrate[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `-((2*Sqrt[a]*Sqrt[-a + I*b]*Sqrt[a + I*b]*(a^2 + b^2)*B*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a] + Sqrt[b]*((-1)^(1/4)*(a + I*b)^(3/2)*b*(I*A + B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]] + Sqrt[-a + I*b]*(2*a*Sqrt[a + I*b]*(A*b - a*B)*Sqrt[Tan[c + d*x]] + (-1)^(1/4)*b*(I*a + b)*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]]))/((-a + I*b)^(3/2)*(a + I*b)^(3/2)*b^(3/2)*d*Sqrt[a + b*Tan[c + d*x]])`

3.458.3 Rubi [A] (verified)Time = 1.14 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.16, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3042, 4088, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{3}{2}}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)^{\frac{3}{2}}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{3}{2}}} dx$$

$$\downarrow \text{4088}$$

3.458. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{3}{2}}} dx$

$$\begin{aligned}
& \frac{2 \int -\frac{((a^2+b^2)B \tan^2(c+dx)) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{b(a^2+b^2)} + \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
& \quad \downarrow 27 \\
& \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b \tan(c+dx)}} - \frac{\int -\frac{((a^2+b^2)B \tan^2(c+dx)) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{b(a^2+b^2)} \\
& \quad \downarrow 3042 \\
& \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b \tan(c+dx)}} - \frac{\int -\frac{((a^2+b^2)B \tan(c+dx)^2) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{b(a^2+b^2)} \\
& \quad \downarrow 4138 \\
& \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b \tan(c+dx)}} - \frac{\int -\frac{((a^2+b^2)B \tan^2(c+dx)) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} d \tan(c+dx)}{bd(a^2+b^2)} \\
& \quad \downarrow 2035 \\
& \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b \tan(c+dx)}} - \frac{2 \int -\frac{((a^2+b^2)B \tan^2(c+dx)) - b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} d \sqrt{\tan(c+dx)}}{bd(a^2+b^2)} \\
& \quad \downarrow 2257 \\
& \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b \tan(c+dx)}} - \frac{2 \int \left(\frac{b(aA+bB) - b(Ab-aB) \tan(c+dx)}{\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} - \frac{(a^2+b^2)B}{\sqrt{a+b \tan(c+dx)}} \right) d \sqrt{\tan(c+dx)}}{bd(a^2+b^2)} \\
& \quad \downarrow 2009 \\
& \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b \tan(c+dx)}} - \frac{2 \left(-\frac{B(a^2+b^2) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{b}} + \frac{b(a-ib)(A+ib) \operatorname{arctan}\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{2\sqrt{-b+ia}} + \frac{b(a+ib)(A-ib) \operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{2\sqrt{b+ia}} \right)}{bd(a^2+b^2)}
\end{aligned}$$

input `Int[(Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x])]/(a + b*Tan[c + d*x])^(3/2),x]`

$$3.458. \quad \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

```
output (-2*((a - I*b)*b*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt
[a + b*Tan[c + d*x]])/(2*Sqrt[I*a - b]) - ((a^2 + b^2)*B*ArcTanh[(Sqrt[b]
*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[b] + ((a + I*b)*b*(A
- I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]
])/(2*Sqrt[I*a + b]))/(b*(a^2 + b^2)*d) + (2*a*(A*b - a*B)*Sqrt[Tan[c + d
*x]])/(b*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])
```

3.458.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2035 Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]
```

```
rule 2257 Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol
] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a
, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4088 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x
])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2))
Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*
(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n
+ 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[
e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n
+ 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] &
& LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])
```

3.458.
$$\int \frac{\tan^3(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

3.458.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.55 (sec) , antiderivative size = 1560668, normalized size of antiderivative = 7126.34

output too large to display

```
input int(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x)
```

```
output result too large to display
```

3.458.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18781 vs. 2(176) = 352.

Time = 10.95 (sec) , antiderivative size = 37564, normalized size of antiderivative = 171.53

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algo
rithm="fracas")
```

```
output Too large to include
```

3.458. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$

3.458.6 Sympy [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(A+B\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx$$

input `integrate(tan(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**(3/2)/(a + b*tan(c + d*x))**(3/2), x)`

3.458.7 Maxima [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^{\frac{3}{2}}}{(b\tan(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorith="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^(3/2)/(b*tan(d*x + c) + a)^(3/2), x)`

3.458.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorith="giac")`

output `Timed out`

3.458.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx = \int \frac{\tan(c+dx)^{\frac{3}{2}}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx$$

input `int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x)`

output `int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x)`

$$3.459 \quad \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.459.1 Optimal result	4416
3.459.2 Mathematica [A] (verified)	4416
3.459.3 Rubi [A] (verified)	4417
3.459.4 Maple [B] (warning: unable to verify)	4420
3.459.5 Fricas [B] (verification not implemented)	4421
3.459.6 Sympy [F]	4421
3.459.7 Maxima [F]	4421
3.459.8 Giac [F(-1)]	4422
3.459.9 Mupad [F(-1)]	4422

3.459.1 Optimal result

Integrand size = 35, antiderivative size = 170

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = -\frac{(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia-b)^{3/2}d} + \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia+b)^{3/2}d} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{(a^2+b^2)d\sqrt{a+b \tan(c+dx)}}$$

output `-(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a-b)^(3/2)/d+(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a+b)^(3/2)/d-2*(A*b-B*a)*tan(d*x+c)^(1/2)/(a^2+b^2)/d/(a+b*tan(d*x+c))^(1/2)`

3.459.2 Mathematica [A] (verified)

Time = 1.70 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.41

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = -\frac{\sqrt[4]{-1}a(a+ib)(A-iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} + \frac{\sqrt[4]{-1}a(a-ib)(A+iB) \operatorname{arctanh}\left(\frac{\sqrt[4]{-1}\sqrt{-a-ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a-ib}}$$

input `Integrate[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2), x]`

3.459. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

output $(-(((-1)^{1/4} * a * (a + I * b) * (A - I * B) * \text{ArcTan} [((-1)^{1/4} * \text{Sqrt} [-a + I * b] * \text{Sqrt} [\text{Tan} [c + d * x]]) / \text{Sqrt} [a + b * \text{Tan} [c + d * x]]) / \text{Sqrt} [-a + I * b]) + ((-1)^{1/4} * a * (a - I * b) * (A + I * B) * \text{ArcTan} [((-1)^{1/4} * \text{Sqrt} [a + I * b] * \text{Sqrt} [\text{Tan} [c + d * x]]) / \text{Sqrt} [a + b * \text{Tan} [c + d * x]]) / \text{Sqrt} [a + I * b] + (2 * b * (A * b - a * B) * \text{Tan} [c + d * x]^{3/2}) / \text{Sqrt} [a + b * \text{Tan} [c + d * x]] + 2 * (- (A * b) + a * B) * \text{Sqrt} [\text{Tan} [c + d * x]] * \text{Sqrt} [a + b * \text{Tan} [c + d * x]]) / (a * (a^2 + b^2) * d)$

3.459.3 Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.19, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4091, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

↓ 3042

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

↓ 4091

$$\frac{2 \int -\frac{b(Ab-aB)+b(aA+bB) \tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{b(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b \tan(c+dx)}}$$

↓ 27

$$\frac{\int \frac{b(Ab-aB)+b(aA+bB) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{b(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b \tan(c+dx)}}$$

↓ 3042

$$\frac{\int \frac{b(Ab-aB)+b(aA+bB) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{b(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b \tan(c+dx)}}$$

↓ 4099

3.459. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \frac{-\frac{2(Ab - aB)\sqrt{\tan(c + dx)}}{d(a^2 + b^2)\sqrt{a + b\tan(c + dx)}} + \frac{\frac{1}{2}b(b + ia)(A + iB) \int \frac{1 - i\tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b\tan(c + dx)}} dx - \frac{1}{2}b(-b + ia)(A - iB) \int \frac{i\tan(c + dx) + 1}{\sqrt{\tan(c + dx)}\sqrt{a + b\tan(c + dx)}} dx}{b(a^2 + b^2)}}{b(a^2 + b^2)} \\
& \quad \downarrow \text{3042} \\
& \frac{-\frac{2(Ab - aB)\sqrt{\tan(c + dx)}}{d(a^2 + b^2)\sqrt{a + b\tan(c + dx)}} + \frac{\frac{1}{2}b(b + ia)(A + iB) \int \frac{1 - i\tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b\tan(c + dx)}} dx - \frac{1}{2}b(-b + ia)(A - iB) \int \frac{i\tan(c + dx) + 1}{\sqrt{\tan(c + dx)}\sqrt{a + b\tan(c + dx)}} dx}{b(a^2 + b^2)}}{b(a^2 + b^2)} \\
& \quad \downarrow \text{4098} \\
& \frac{b(b + ia)(A + iB) \int \frac{1}{(i\tan(c + dx) + 1)\sqrt{\tan(c + dx)}\sqrt{a + b\tan(c + dx)}} d\tan(c + dx) - \frac{b(-b + ia)(A - iB) \int \frac{1}{(1 - i\tan(c + dx))\sqrt{\tan(c + dx)}\sqrt{a + b\tan(c + dx)}} d\tan(c + dx)}{2d}}{b(a^2 + b^2)} \\
& \quad \downarrow \text{104} \\
& \frac{b(b + ia)(A + iB) \int \frac{1}{\frac{(ia - b)\tan(c + dx)}{a + b\tan(c + dx)} + 1} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b\tan(c + dx)}} - \frac{b(-b + ia)(A - iB) \int \frac{1}{1 - \frac{(ia + b)\tan(c + dx)}{a + b\tan(c + dx)}} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b\tan(c + dx)}}}{d}}{b(a^2 + b^2)} \\
& \quad \downarrow \text{216} \\
& \frac{b(b + ia)(A + iB) \arctan\left(\frac{\sqrt{-b + ia}\sqrt{\tan(c + dx)}}{\sqrt{a + b\tan(c + dx)}}\right) - \frac{b(-b + ia)(A - iB) \int \frac{1}{1 - \frac{(ia + b)\tan(c + dx)}{a + b\tan(c + dx)}} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b\tan(c + dx)}}}{d}}{d\sqrt{-b + ia}}}{b(a^2 + b^2)} \\
& \quad \downarrow \text{219} \\
& \frac{b(b + ia)(A + iB) \arctan\left(\frac{\sqrt{-b + ia}\sqrt{\tan(c + dx)}}{\sqrt{a + b\tan(c + dx)}}\right) - \frac{b(-b + ia)(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{b + ia}\sqrt{\tan(c + dx)}}{\sqrt{a + b\tan(c + dx)}}\right)}{d\sqrt{b + ia}}}{d\sqrt{-b + ia}}}{b(a^2 + b^2)}
\end{aligned}$$

3.459. $\int \frac{\sqrt{\tan(c + dx)}(A + B\tan(c + dx))}{(a + b\tan(c + dx))^{3/2}} dx$

```
input Int[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x
]
```

```
output ((b*(I*a + b)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a +
b*Tan[c + d*x]])/(Sqrt[I*a - b]*d) - ((I*a - b)*b*(A - I*B)*ArcTanh[(Sqr
t[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d
))/(b*(a^2 + b^2)) - (2*(A*b - a*B)*Sqrt[Tan[c + d*x]])/((a^2 + b^2)*d*Sqr
t[a + b*Tan[c + d*x]])
```

3.459.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 104 Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))/((e_) + (f_)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x
] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

```
rule 219 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4091 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m +
1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n)
+ A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan
[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x] /; FreeQ
[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || Integers
Q[2*m, 2*n])
```

```
rule 4098 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*
x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]
```

```
rule 4099 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2,
0]
```

3.459.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.43 (sec) , antiderivative size = 1559497, normalized size of antiderivative = 9173.51

output too large to display

```
input int(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x)
```

```
output result too large to display
```

$$3.459. \quad \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.459.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18532 vs. $2(138) = 276$.

Time = 7.01 (sec) , antiderivative size = 18532, normalized size of antiderivative = 109.01

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algo
rithm="fricas")`

output Too large to include

3.459.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(A+B\tan(c+dx))\sqrt{\tan(c+dx)}}{(a+b\tan(c+dx))^{3/2}} dx$$

input `integrate(tan(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(tan(c + d*x))/(a + b*tan(c + d*x))**(3/
2), x)`

3.459.7 Maxima [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(B\tan(dx+c)+A)\sqrt{\tan(dx+c)}}{(b\tan(dx+c)+a)^{3/2}} dx$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(tan(d*x + c))/(b*tan(d*x + c) + a)^(3/
2), x)`

3.459. $\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$

3.459.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algo
rithm="giac")`

output `Timed out`

3.459.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$$

input `int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x
)`

output `int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),
x)`

3.460
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx$$

3.460.1 Optimal result 4423
 3.460.2 Mathematica [A] (verified) 4423
 3.460.3 Rubi [A] (verified) 4424
 3.460.4 Maple [B] (warning: unable to verify) 4427
 3.460.5 Fracas [B] (verification not implemented) 4428
 3.460.6 Sympy [F] 4429
 3.460.7 Maxima [F] 4429
 3.460.8 Giac [F(-1)] 4429
 3.460.9 Mupad [F(-1)] 4430

3.460.1 Optimal result

Integrand size = 35, antiderivative size = 175

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \frac{(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia - b)^{3/2}d} + \frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia + b)^{3/2}d} + \frac{2b(Ab - aB) \sqrt{\tan(c + dx)}}{a(a^2 + b^2) d \sqrt{a + b \tan(c + dx)}}$$

output `(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a-b)^(3/2)/d+(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a+b)^(3/2)/d+2*b*(A*b-B*a)*tan(d*x+c)^(1/2)/a/(a^2+b^2)/d/(a+b*tan(d*x+c))^(1/2)`

3.460.2 Mathematica [A] (verified)

Time = 0.89 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.15

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \frac{\sqrt[4]{-1}(-ia+b)(A-iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} + \frac{\sqrt[4]{-1}(a-ib)(-iA+B)}{(a^2 + b^2) d}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)), x]`

output $((-1)^{(1/4)} * ((-I) * a + b) * (A - I * B) * \text{ArcTan} [((-1)^{(1/4)} * \text{Sqrt} [-a + I * b] * \text{Sqrt} [\text{Tan} [c + d * x]]) / \text{Sqrt} [a + b * \text{Tan} [c + d * x]]) / \text{Sqrt} [-a + I * b] + ((-1)^{(1/4)} * (a - I * b) * ((-I) * A + B) * \text{ArcTan} [((-1)^{(1/4)} * \text{Sqrt} [a + I * b] * \text{Sqrt} [\text{Tan} [c + d * x]]) / \text{Sqrt} [a + b * \text{Tan} [c + d * x]]) / \text{Sqrt} [a + I * b] + (2 * b * (A * b - a * B) * \text{Sqrt} [\text{Tan} [c + d * x]]) / (a * \text{Sqrt} [a + b * \text{Tan} [c + d * x]]) / ((a^2 + b^2) * d)$

3.460.3 Rubi [A] (verified)

Time = 0.95 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.16, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4092, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} (a + b \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} (a + b \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{4092} \\
 & \frac{2 \int \frac{a(A + bB) - a(Ab - aB) \tan(c + dx)}{2 \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx}{a(a^2 + b^2)} + \frac{2b(Ab - aB) \sqrt{\tan(c + dx)}}{ad(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{a(A + bB) - a(Ab - aB) \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx}{a(a^2 + b^2)} + \frac{2b(Ab - aB) \sqrt{\tan(c + dx)}}{ad(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a(A + bB) - a(Ab - aB) \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx}{a(a^2 + b^2)} + \frac{2b(Ab - aB) \sqrt{\tan(c + dx)}}{ad(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} \\
 & \quad \downarrow \text{4099} \\
 & \frac{2b(Ab - aB) \sqrt{\tan(c + dx)}}{ad(a^2 + b^2) \sqrt{a + b \tan(c + dx)}} + \\
 & \frac{\frac{1}{2} a(a - ib)(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2} a(a + ib)(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx}{a(a^2 + b^2)}
 \end{aligned}$$

3.460. $\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)} (a + b \tan(c + dx))^{3/2}} dx$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{2b(Ab - aB)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b\tan(c + dx)}} + \\ & \frac{\frac{1}{2}a(a - ib)(A + iB) \int \frac{1 - i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}a(a + ib)(A - iB) \int \frac{i\tan(c+dx)+1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 4098 \\ & \frac{2b(Ab - aB)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b\tan(c + dx)}} + \\ & \frac{a(a+ib)(A-iB) \int \frac{1}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} d\tan(c+dx) + \frac{a(a-ib)(A+iB) \int \frac{1}{(i\tan(c+dx)+1)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} d\tan(c+dx)}{a(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 104 \\ & \frac{2b(Ab - aB)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b\tan(c + dx)}} + \\ & \frac{a(a-ib)(A+iB) \int \frac{1}{\frac{(ia-b)\tan(c+dx)}{a+b\tan(c+dx)}+1} d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}} + \frac{a(a+ib)(A-iB) \int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{a(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 216 \\ & \frac{2b(Ab - aB)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b\tan(c + dx)}} + \\ & \frac{a(a+ib)(A-iB) \int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}} + \frac{a(a-ib)(A+iB) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{a(a^2 + b^2) d\sqrt{-b+ia}} \end{aligned}$$

$$\begin{aligned} & \downarrow 219 \\ & \frac{2b(Ab - aB)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b\tan(c + dx)}} + \\ & \frac{a(a-ib)(A+iB) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) + \frac{a(a+ib)(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}}}{a(a^2 + b^2)} \end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)),x]`


```
output ((a*(a - I*b)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a +
b*Tan[c + d*x]])/(Sqrt[I*a - b]*d) + (a*(a + I*b)*(A - I*B)*ArcTanh[(Sqr
t[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d
))/(a*(a^2 + b^2)) + (2*b*(A*b - a*B)*Sqrt[Tan[c + d*x]])/(a*(a^2 + b^2)*d
*Sqrt[a + b*Tan[c + d*x]])
```

3.460.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 104 Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

```
rule 219 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4092 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)
/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^
2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*
B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n +
2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n
+ 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1]
&& (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m]
|| (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4098 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*
x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]
```

```
rule 4099 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2,
0]
```

3.460.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 4926 vs. $2(149) = 298$.

Time = 29.70 (sec) , antiderivative size = 4927, normalized size of antiderivative = 28.15

method	result	size
default	Expression too large to display	4927
parts	Expression too large to display	1559557

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x,method=_RET
URNVERBOSE)
```

$$3.460. \quad \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))^{3/2}}} dx$$

output `1/4/d*2^(1/2)/a/(b+(a^2+b^2)^(1/2))^(1/2)/(a^2+b^2)^(3/2)*(A*ln(1/(1-cos(d*x+c)))*(a*(1-cos(d*x+c))^2*csc(d*x+c)+2*(a^2+b^2)^(1/2)*(1-cos(d*x+c))-2*(1-cos(d*x+c))*(a*(1-cos(d*x+c))^2*csc(d*x+c)^2-2*b*(csc(d*x+c)-cot(d*x+c))-a)*csc(d*x+c))^(1/2)*(-b+(a^2+b^2)^(1/2))^(1/2)*sin(d*x+c)-2*b*(1-cos(d*x+c))-a*sin(d*x+c)))*a^2*(-b+(a^2+b^2)^(1/2))^(1/2)*((1-cos(d*x+c))*(a*(1-cos(d*x+c))^2*csc(d*x+c)^2-2*b*(csc(d*x+c)-cot(d*x+c))-a)*csc(d*x+c))^(1/2)*(b+(a^2+b^2)^(1/2))^(1/2)-A*ln(1/(1-cos(d*x+c)))*(a*(1-cos(d*x+c))^2*csc(d*x+c)+2*(a^2+b^2)^(1/2)*(1-cos(d*x+c))-2*((1-cos(d*x+c))*(a*(1-cos(d*x+c))^2*csc(d*x+c)^2-2*b*(csc(d*x+c)-cot(d*x+c))-a)*csc(d*x+c))^(1/2)*(-b+(a^2+b^2)^(1/2))^(1/2)*sin(d*x+c)-2*b*(1-cos(d*x+c))-a*sin(d*x+c)))*b^2*(-b+(a^2+b^2)^(1/2))^(1/2)*((1-cos(d*x+c))*(a*(1-cos(d*x+c))^2*csc(d*x+c)^2-2*b*(csc(d*x+c)-cot(d*x+c))-a)*csc(d*x+c))^(1/2)*(b+(a^2+b^2)^(1/2))^(1/2)-A*(a^2+b^2)^(1/2)*ln(1/(1-cos(d*x+c)))*(a*(1-cos(d*x+c))^2*csc(d*x+c)+2*(a^2+b^2)^(1/2)*(1-cos(d*x+c))-2*((1-cos(d*x+c))*(a*(1-cos(d*x+c))^2*csc(d*x+c)^2-2*b*(csc(d*x+c)-cot(d*x+c))-a)*csc(d*x+c))^(1/2)*(-b+(a^2+b^2)^(1/2))^(1/2)*sin(d*x+c)-2*b*(1-cos(d*x+c))-a*sin(d*x+c)))*((1-cos(d*x+c))*(a*(1-cos(d*x+c))^2*csc(d*x+c)^2-2*b*(csc(d*x+c)-cot(d*x+c))-a)*csc(d*x+c))^(1/2)*(-b+(a^2+b^2)^(1/2))^(1/2)*(b+(a^2+b^2)^(1/2))^(1/2)*b-A*ln(1/(1-cos(d*x+c)))*(a*(1-cos(d*x+c))^2*csc(d*x+c)+2*(a^2+b^2)^(1/2)*(1-cos(d*x+c))+2*((1-cos(d*x+c))*(a*(1-cos(d*x+c))^2*csc(d*x+c)^2-2*b*(csc(d*x+c)-cot(d*x+c))-...`

3.460.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18563 vs. $2(143) = 286$.

Time = 6.95 (sec) , antiderivative size = 18563, normalized size of antiderivative = 106.07

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x, algo rithm="fracas")`

output `Too large to include`

3.460.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{\frac{3}{2}} \sqrt{\tan(c + dx)}} dx$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+b*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))**(3/2)*sqrt(tan(c + d*x))), x)`

3.460.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a)^{\frac{3}{2}} \sqrt{\tan(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x, algorith="maxima")`

output `integrate((B*tan(d*x + c) + A)/((b*tan(d*x + c) + a)^(3/2)*sqrt(tan(d*x + c))), x)`

3.460.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x, algorith="giac")`

output `Timed out`

3.460.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(3/2)),x)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(3/2)),x)`

3.461
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$$

3.461.1 Optimal result 4431
 3.461.2 Mathematica [A] (verified) 4432
 3.461.3 Rubi [A] (verified) 4432
 3.461.4 Maple [B] (warning: unable to verify) 4436
 3.461.5 Fracas [B] (verification not implemented) 4437
 3.461.6 Sympy [F] 4437
 3.461.7 Maxima [F(-1)] 4437
 3.461.8 Giac [F(-1)] 4438
 3.461.9 Mupad [F(-1)] 4438

3.461.1 Optimal result

Integrand size = 35, antiderivative size = 216

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \frac{(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia - b)^{3/2}d} - \frac{(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia + b)^{3/2}d} - \frac{2A}{ad\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} - \frac{2b(a^2A + 2Ab^2 - abB) \sqrt{\tan(c + dx)}}{a^2(a^2 + b^2)d\sqrt{a + b \tan(c + dx)}}$$

output `(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a-b)^(3/2)/d-(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a+b)^(3/2)/d-2*A/a/d/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2)-2*b*(A*a^2+2*A*b^2-B*a*b)*tan(d*x+c)^(1/2)/a^2/(a^2+b^2)/d/(a+b*tan(d*x+c))^(1/2)`

3.461.2 Mathematica [A] (verified)

Time = 2.33 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.15

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \frac{\sqrt[4]{-1} \left(\frac{(a+ib)(A-iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} - \frac{(a-ib)(A+iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} \right)}{a^2+b^2}$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2)),x]`

output `(((-1)^(1/4)*(((a + I*b)*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[-a + I*b] - ((a - I*b)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[a + I*b]))/(a^2 + b^2) - (2*A)/(a*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]]) - (2*b*(a^2*A + 2*A*b^2 - a*b*B)*Sqrt[Tan[c + d*x]])/(a^2*(a^2 + b^2)*Sqrt[a + b*Tan[c + d*x]]))/d`

3.461.3 Rubi [A] (verified)

Time = 1.32 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.19, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2}(a + b \tan(c + dx))^{3/2}} dx$$

↓ 4092

$$\frac{2 \int \frac{2Ab \tan^2(c+dx)+aA \tan(c+dx)+2Ab-aB}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{a} - \frac{2A}{ad\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}}$$

3.461. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \int \frac{2Ab \tan^2(c+dx) + aA \tan(c+dx) + 2Ab - aB}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx \quad \downarrow 27 \\
 & \frac{2A}{a \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} \\
 & \int \frac{2Ab \tan(c+dx)^2 + aA \tan(c+dx) + 2Ab - aB}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx \quad \downarrow 3042 \\
 & \frac{2A}{a \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} \\
 & \frac{2 \int \frac{(Ab-aB)a^2 + (aA+bB) \tan(c+dx)a^2}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \quad \downarrow 4132 \\
 & \frac{2A}{a \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} \\
 & \int \frac{(Ab-aB)a^2 + (aA+bB) \tan(c+dx)a^2}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx + \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \quad \downarrow 27 \\
 & \frac{2A}{a \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} \\
 & \int \frac{(Ab-aB)a^2 + (aA+bB) \tan(c+dx)a^2}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx + \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \quad \downarrow 3042 \\
 & \frac{2A}{a \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} \\
 & \int \frac{(Ab-aB)a^2 + (aA+bB) \tan(c+dx)a^2}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx + \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \quad \downarrow 4099 \\
 & \frac{2A}{a \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} \\
 & \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{1}{2} \frac{a^2(b+ia)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2} a^2(-b+ia)(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} \\
 & \frac{2A}{a \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}} \\
 & \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{1}{2} \frac{a^2(b+ia)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2} a^2(-b+ia)(A-iB) \int \frac{i \tan(c+dx)+1}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} \\
 & \frac{2A}{a \sqrt{\tan(c+dx)} \sqrt{a+b \tan(c+dx)}}
 \end{aligned}$$

3.461. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$

$$\begin{aligned} & \downarrow 4098 \\ & \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{2A}{\frac{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{a^2(b+ia)(A+iB)\int \frac{1}{(i\tan(c+dx)+1)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}d\tan(c+dx)} - \frac{a^2(-b+ia)(A-iB)\int \frac{1}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}d\tan(c+dx)}}{a(a^2+b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 104 \\ & \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{2A}{\frac{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{a^2(b+ia)(A+iB)\int \frac{1}{\frac{(ia-b)\tan(c+dx)}{a+b\tan(c+dx)}+1}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}} - \frac{a^2(-b+ia)(A-iB)\int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}} \end{aligned}$$

$$\begin{aligned} & \downarrow 216 \\ & \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{2A}{\frac{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{a^2(b+ia)(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)} - \frac{a^2(-b+ia)(A-iB)\int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}} \end{aligned}$$

$$\begin{aligned} & \downarrow 219 \\ & \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{2A}{\frac{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{a^2(b+ia)(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)} - \frac{a^2(-b+ia)(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}} \end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2)),x]`

output `(-2*A)/(a*d*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]]) - (((a^2*(I*a + b)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d) - (a^2*(I*a - b)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d))/(a*(a^2 + b^2)) + (2*b*(a^2*A + 2*A*b^2 - a*b*B)*Sqrt[Tan[c + d*x]])/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])/a`

3.461.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4098 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*
x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]
```

```
rule 4099 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2,
0]
```

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
(!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

3.461.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 5197 vs. $2(186) = 372$.

Time = 3.93 (sec) , antiderivative size = 5198, normalized size of antiderivative = 24.06

method	result	size
default	Expression too large to display	5198
parts	Expression too large to display	1560457

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x,method=_RET
URNVERBOSE)
```

$$3.461. \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$$

output result too large to display

3.461.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18689 vs. 2(180) = 360.
Time = 7.04 (sec) , antiderivative size = 18689, normalized size of antiderivative = 86.52

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x, algorithm="fricas")`

output Too large to include

3.461.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{\frac{3}{2}} \tan^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(3/2)/(a+b*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))**(3/2)*tan(c + d*x)**(3/2)), x)`

3.461.7 Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output Timed out

3.461. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$

3.461.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x, algorith="giac")`

output `Timed out`

3.461.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Hanged}$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(3/2)),x)`

output `\text{Hanged}`

3.462
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$$

3.462.1 Optimal result 4439
 3.462.2 Mathematica [A] (verified) 4440
 3.462.3 Rubi [A] (verified) 4440
 3.462.4 Maple [C] (warning: unable to verify) 4445
 3.462.5 Fracas [B] (verification not implemented) 4446
 3.462.6 Sympy [F] 4446
 3.462.7 Maxima [F(-1)] 4446
 3.462.8 Giac [F(-1)] 4447
 3.462.9 Mupad [F(-1)] 4447

3.462.1 Optimal result

Integrand size = 35, antiderivative size = 276

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx =$$

$$-\frac{(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia - b)^{3/2}d} - \frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia + b)^{3/2}d}$$

$$-\frac{2A}{3ad \tan^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} + \frac{2(4Ab - 3aB)}{3a^2 d \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}$$

$$+ \frac{2b(5a^2 Ab + 8Ab^3 - 3a^3 B - 6ab^2 B) \sqrt{\tan(c + dx)}}{3a^3 (a^2 + b^2) d \sqrt{a + b \tan(c + dx)}}$$

```
output -(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*
a-b)^(3/2)/d-(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c
))^(1/2))/(I*a+b)^(3/2)/d+2/3*(4*A*b-3*B*a)/a^2/d/tan(d*x+c)^(1/2)/(a+b*ta
n(d*x+c))^(1/2)+2/3*b*(5*A*a^2*b+8*A*b^3-3*B*a^3-6*B*a*b^2)*tan(d*x+c)^(1/
2)/a^3/(a^2+b^2)/d/(a+b*tan(d*x+c))^(1/2)-2/3*A/a/d/(a+b*tan(d*x+c))^(1/2)
/tan(d*x+c)^(3/2)
```

3.462.2 Mathematica [A] (verified)

Time = 3.26 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.08

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \frac{3 \sqrt[4]{-1} a \left(\frac{(a+ib)(iA+B) \arctan\left(\frac{\sqrt[4]{-1} \sqrt{-a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} \right) + \frac{(ia+b)(A+iB) \arctan\left(\frac{\sqrt[4]{-1} \sqrt{-a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}}}{a^2 + b^2}$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^(3/2)),x]`

output `((3*(-1)^(1/4)*a*(((a + I*b)*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[-a + I*b] + ((I*a + b)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[a + I*b]))/(a^2 + b^2) - (2*A)/(Tan[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]]) + (8*A*b - 6*a*B)/(a*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]]) + (2*b*(5*a^2*A*b + 8*A*b^3 - 3*a^3*B - 6*a*b^2*B)*Sqrt[Tan[c + d*x]])/(a^2*(a^2 + b^2)*Sqrt[a + b*Tan[c + d*x]]))/(3*a*d)`

3.462.3 Rubi [A] (verified)Time = 1.80 (sec) , antiderivative size = 316, normalized size of antiderivative = 1.14, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2}(a + b \tan(c + dx))^{3/2}} dx$$

↓ 4092

3.462. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{2 \int \frac{4Ab \tan^2(c+dx) + 3aA \tan(c+dx) + 4Ab - 3aB}{2 \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx}{3a} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{4Ab \tan^2(c+dx) + 3aA \tan(c+dx) + 4Ab - 3aB}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx}{3a} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{4Ab \tan(c+dx)^2 + 3aA \tan(c+dx) + 4Ab - 3aB}{\tan(c+dx)^{3/2}(a+b \tan(c+dx))^{3/2}} dx}{3a} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 4132 \\
 & \frac{2 \int -\frac{3Aa^2 + 3B \tan(c+dx)a^2 + 6bBa - 8Ab^2 - 2b(4Ab - 3aB) \tan^2(c+dx)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{a} - \frac{2(4Ab - 3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \\
 & \quad \frac{3a}{2A} \\
 & \quad \frac{3ad \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{3Aa^2 + 3B \tan(c+dx)a^2 + 6bBa - 8Ab^2 - 2b(4Ab - 3aB) \tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{a} - \frac{2(4Ab - 3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \\
 & \quad \frac{3a}{2A} \\
 & \quad \frac{3ad \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{3Aa^2 + 3B \tan(c+dx)a^2 + 6bBa - 8Ab^2 - 2b(4Ab - 3aB) \tan(c+dx)^2}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{a} - \frac{2(4Ab - 3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \\
 & \quad \frac{3a}{2A} \\
 & \quad \frac{3ad \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 4132 \\
 & \frac{2 \int \frac{3(a^3(aA+bB) - a^3(Ab-aB) \tan(c+dx))}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-3a^3B + 5a^2Ab - 6ab^2B + 8Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} - \frac{2(4Ab - 3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \\
 & \quad \frac{3a}{2A} \\
 & \quad \frac{3ad \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 27
 \end{aligned}$$

3.462. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{3 \int \frac{a^3(aA+bB)-a^3(Ab-aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{2b(-3a^3B+5a^2Ab-6ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{a} - \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} \\
 & \quad \frac{3a}{2A} \\
 & \quad \frac{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}{\phantom{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}} \\
 & \quad \downarrow 3042 \\
 & \frac{3 \int \frac{a^3(aA+bB)-a^3(Ab-aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{2b(-3a^3B+5a^2Ab-6ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{a} - \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} \\
 & \quad \frac{3a}{2A} \\
 & \quad \frac{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}{\phantom{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}} \\
 & \quad \downarrow 4099 \\
 & \quad \frac{2A}{} \\
 & \quad \frac{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}{\phantom{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}} - \frac{2b(-3a^3B+5a^2Ab-6ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{3\left(\frac{1}{2}a^3(a-ib)(A+iB) \int \frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}a^3(a+ib)(A-iB) \int \frac{1+i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx\right)}{a(a^2+b^2)} \\
 & \quad \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} + \frac{3a}{} \\
 & \quad \downarrow 3042 \\
 & \quad \frac{2A}{} \\
 & \quad \frac{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}{\phantom{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}} - \frac{2b(-3a^3B+5a^2Ab-6ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{3\left(\frac{1}{2}a^3(a-ib)(A+iB) \int \frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}a^3(a+ib)(A-iB) \int \frac{1+i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx\right)}{a(a^2+b^2)} \\
 & \quad \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} + \frac{3a}{} \\
 & \quad \downarrow 4098 \\
 & \quad \frac{2A}{} \\
 & \quad \frac{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}{\phantom{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}} - \frac{2b(-3a^3B+5a^2Ab-6ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{3\left(\frac{a^3(a+ib)(A-iB) \int \frac{1}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{(1+i\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx\right)}{2d(a^2+b^2)} \\
 & \quad \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} + \frac{3a}{} \\
 & \quad \downarrow 104
 \end{aligned}$$

3.462. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$

$$\frac{-\frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} + \frac{-\frac{2b(-3a^3B+5a^2Ab-6ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{2A}{3ad\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}}{3a} + \frac{\left(\frac{a^3(a-ib)(A+iB) \int \frac{1}{\frac{(ia-b)\tan(c+dx)}{a+b\tan(c+dx)} + 1} d - \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}} \right)}{a(a^2+b^2)}$$

216

$$\frac{-\frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} + \frac{-\frac{2b(-3a^3B+5a^2Ab-6ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{2A}{3ad\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}}{3a} + \frac{\left(\frac{a^3(a+ib)(A-iB) \int \frac{1}{1 - \frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d - \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}} \right)}{a(a^2+b^2)}$$

219

$$\frac{-\frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} + \frac{-\frac{2b(-3a^3B+5a^2Ab-6ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{2A}{3ad\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}}{3a} + \frac{\left(\frac{a^3(a-ib)(A+iB) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) + a^3(a+ib)(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} \right)}{a(a^2+b^2)}$$

```
input Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^(3/2)),x]
```

```
output (-2*A)/(3*a*d*Tan[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]]) - ((-2*(4*A*b - 3*a*B))/(a*d*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]]) + ((3*((a^3*(a - I*b)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d) + (a^3*(a + I*b)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d)))/(a*(a^2 + b^2)) - (2*b*(5*a^2*A*b + 8*A*b^3 - 3*a^3*B - 6*a*b^2*B)*Sqrt[Tan[c + d*x]])/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]))/a)/(3*a)
```

3.462. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$

3.462.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4098 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*
x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]
```

```
rule 4099 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2,
0]
```

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

3.462.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 3.99 (sec) , antiderivative size = 9712, normalized size of antiderivative = 35.19

method	result	size
default	Expression too large to display	9712
parts	Expression too large to display	1562265

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c))^(3/2),x,method=_RET
URNVERBOSE)
```

$$3.462. \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$$

output result too large to display

3.462.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18754 vs. 2(227) = 454.
Time = 7.05 (sec) , antiderivative size = 18754, normalized size of antiderivative = 67.95

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c))^(3/2),x, algorithm="fricas")`

output Too large to include

3.462.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{\frac{3}{2}} \tan^{\frac{5}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(5/2)/(a+b*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))**(3/2)*tan(c + d*x)**(5/2)), x)`

3.462.7 Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output Timed out

3.462. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$

3.462.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c))^(3/2),x, algorith="giac")`

output `Timed out`

3.462.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2} (a + b \tan(c + dx))^{3/2}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + b*tan(c + d*x))^(3/2)),x)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + b*tan(c + d*x))^(3/2)),x)`

3.463
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

3.463.1 Optimal result 4448
 3.463.2 Mathematica [B] (verified) 4449
 3.463.3 Rubi [A] (verified) 4450
 3.463.4 Maple [B] (warning: unable to verify) 4453
 3.463.5 Fricas [B] (verification not implemented) 4454
 3.463.6 Sympy [F(-1)] 4454
 3.463.7 Maxima [F] 4455
 3.463.8 Giac [F(-1)] 4455
 3.463.9 Mupad [F(-1)] 4455

3.463.1 Optimal result

Integrand size = 35, antiderivative size = 282

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia-b)^{5/2}d}$$

$$+ \frac{2B \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{b^{5/2}d} - \frac{(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia+b)^{5/2}d}$$

$$+ \frac{2a(Ab-aB) \tan^{\frac{3}{2}}(c+dx)}{3b(a^2+b^2)d(a+b \tan(c+dx))^{3/2}} + \frac{2a(2Ab^3-a(a^2+3b^2)B) \sqrt{\tan(c+dx)}}{b^2(a^2+b^2)^2 d \sqrt{a+b \tan(c+dx)}}$$

output `(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a-b)^(5/2)/d+2*B*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/b^(5/2)/d-(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a+b)^(5/2)/d+2*a*(2*A*b^3-a*(a^2+3*b^2)*B)*tan(d*x+c)^(1/2)/b^2/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^(1/2)+2/3*a*(A*b-B*a)*tan(d*x+c)^(3/2)/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)`

3.463.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 596 vs. $2(282) = 564$.

Time = 6.46 (sec) , antiderivative size = 596, normalized size of antiderivative = 2.11

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \frac{(A-iB) \left(\frac{3\sqrt[4]{-1} \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{(-a+ib)^{5/2}} + \frac{\tan^{\frac{3}{2}}(c+dx)}{(a-ib)(a+b\tan(c+dx))} \right)}{3d} + \frac{(A+iB) \left(\frac{3\sqrt[4]{-1} \arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{(a+ib)^{5/2}} - \frac{\tan^{\frac{3}{2}}(c+dx)}{(a+ib)(a+b\tan(c+dx))^{3/2}} - \frac{3i\sqrt{\tan(c+dx)}}{(a+ib)^2\sqrt{a+b\tan(c+dx)}} \right)}{3d} + \frac{(iA-B)\sqrt{a+b\tan(c+dx)} \left(\frac{b^2\tan^2(c+dx)}{(a+b\tan(c+dx))^2} + \frac{3b\tan(c+dx)}{a+b\tan(c+dx)} - \frac{3\sqrt{b}\operatorname{arcsinh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)\sqrt{\tan(c+dx)}}{\sqrt{a}\sqrt{1+\frac{b\tan(c+dx)}{a}}}\right)}{3b^3d\sqrt{\tan(c+dx)}} + \frac{(iA+B)\sqrt{a+b\tan(c+dx)} \left(\frac{b^2\tan^2(c+dx)}{(a+b\tan(c+dx))^2} + \frac{3b\tan(c+dx)}{a+b\tan(c+dx)} - \frac{3\sqrt{b}\operatorname{arcsinh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a}}\right)\sqrt{\tan(c+dx)}}{\sqrt{a}\sqrt{1+\frac{b\tan(c+dx)}{a}}}\right)}{3b^3d\sqrt{\tan(c+dx)}}$$

input `Integrate[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]`

output `((A - I*B)*((3*(-1)^(1/4)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]])/(-a + I*b)^(5/2) + Tan[c + d*x]^(3/2)/((a - I*b)*(a + b*Tan[c + d*x])^(3/2)) - ((3*I)*Sqrt[Tan[c + d*x]]/((a - I*b)^2*Sqrt[a + b*Tan[c + d*x]])))/(3*d) - ((A + I*B)*((3*(-1)^(1/4)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]])/(a + I*b)^(5/2) - Tan[c + d*x]^(3/2)/((a + I*b)*(a + b*Tan[c + d*x])^(3/2)) - ((3*I)*Sqrt[Tan[c + d*x]]/((a + I*b)^2*Sqrt[a + b*Tan[c + d*x]])))/(3*d) + ((I*A - B)*Sqrt[a + b*Tan[c + d*x]]*((b^2*Tan[c + d*x]^2)/(a + b*Tan[c + d*x]^2) + (3*b*Tan[c + d*x])/(a + b*Tan[c + d*x]) - (3*Sqrt[b]*ArcSinh[Sqrt[b]*Sqrt[Tan[c + d*x]]]/Sqrt[a]*Sqrt[Tan[c + d*x]]/(Sqrt[a]*Sqrt[1 + (b*Tan[c + d*x])/a])))/(3*b^3*d*Sqrt[Tan[c + d*x]]) - ((I*A + B)*Sqrt[a + b*Tan[c + d*x]]*((b^2*Tan[c + d*x]^2)/(a + b*Tan[c + d*x]^2) + (3*b*Tan[c + d*x])/(a + b*Tan[c + d*x]) - (3*Sqrt[b]*ArcSinh[Sqrt[b]*Sqrt[Tan[c + d*x]]]/Sqrt[a]*Sqrt[Tan[c + d*x]]/(Sqrt[a]*Sqrt[1 + (b*Tan[c + d*x])/a])))/(3*b^3*d*Sqrt[Tan[c + d*x]]))`

$$3.463. \quad \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$$

3.463.3 Rubi [A] (verified)

Time = 1.73 (sec) , antiderivative size = 344, normalized size of antiderivative = 1.22, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.314$, Rules used = {3042, 4088, 27, 3042, 4128, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{5}{2}}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^{\frac{5}{2}}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{5}{2}}} dx \\
 & \quad \downarrow \text{4088} \\
 & \frac{2 \int -\frac{3\sqrt{\tan(c+dx)}(-((a^2+b^2)B \tan^2(c+dx))-b(Ab-aB) \tan(c+dx)+a(Ab-aB))}{2(a+b \tan(c+dx))^{\frac{3}{2}}} dx}{3b(a^2+b^2)} + \\
 & \quad \frac{2a(Ab-aB) \tan^{\frac{3}{2}}(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^{\frac{3}{2}}} \\
 & \quad \downarrow \text{27} \\
 & \frac{2a(Ab-aB) \tan^{\frac{3}{2}}(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^{\frac{3}{2}}} - \\
 & \frac{\int \frac{\sqrt{\tan(c+dx)}(-((a^2+b^2)B \tan^2(c+dx))-b(Ab-aB) \tan(c+dx)+a(Ab-aB))}{(a+b \tan(c+dx))^{\frac{3}{2}}} dx}{b(a^2+b^2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{2a(Ab-aB) \tan^{\frac{3}{2}}(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^{\frac{3}{2}}} - \\
 & \frac{\int \frac{\sqrt{\tan(c+dx)}(-((a^2+b^2)B \tan(c+dx)^2)-b(Ab-aB) \tan(c+dx)+a(Ab-aB))}{(a+b \tan(c+dx))^{\frac{3}{2}}} dx}{b(a^2+b^2)} \\
 & \quad \downarrow \text{4128} \\
 & \frac{2a(Ab-aB) \tan^{\frac{3}{2}}(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^{\frac{3}{2}}} - \\
 & \frac{2 \int \frac{(Aa^2+2bBa-Ab^2) \tan(c+dx)b^2-(a^2+b^2)^2 B \tan^2(c+dx)+a(2Ab^3-a(a^2+3b^2)B)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{b(a^2+b^2)} - \frac{2a(2Ab^3-aB(a^2+3b^2))\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{2a(Ab-aB) \tan^{\frac{3}{2}}(c+dx)}{3bd(a^2+b^2)(a+b \tan(c+dx))^{\frac{3}{2}}} - \frac{2a(2Ab^3-aB(a^2+3b^2))\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b \tan(c+dx)}}
 \end{aligned}$$

3.463. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{5}{2}}} dx$

$$\begin{aligned}
& \frac{\int \frac{2a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{(Aa^2 + 2bBa - Ab^2) \tan(c + dx)b^2 - (a^2 + b^2)^2 B \tan^2(c + dx) + a(2Ab^3 - a(a^2 + 3b^2)B)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx}{b(a^2 + b^2)} - \frac{2a(2Ab^3 - aB(a^2 + 3b^2))\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{2a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{(Aa^2 + 2bBa - Ab^2) \tan(c + dx)b^2 - (a^2 + b^2)^2 B \tan^2(c + dx) + a(2Ab^3 - a(a^2 + 3b^2)B)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx}{b(a^2 + b^2)} - \frac{2a(2Ab^3 - aB(a^2 + 3b^2))\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} \\
& \quad \downarrow \text{4138} \\
& \frac{\int \frac{2a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{(Aa^2 + 2bBa - Ab^2) \tan(c + dx)b^2 - (a^2 + b^2)^2 B \tan^2(c + dx) + a(2Ab^3 - a(a^2 + 3b^2)B)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}(\tan^2(c + dx) + 1)} d \tan(c + dx)}{bd(a^2 + b^2)} - \frac{2a(2Ab^3 - aB(a^2 + 3b^2))\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} \\
& \quad \downarrow \text{2035} \\
& \frac{2 \int \frac{2a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{(Aa^2 + 2bBa - Ab^2) \tan(c + dx)b^2 - (a^2 + b^2)^2 B \tan^2(c + dx) + a(2Ab^3 - a(a^2 + 3b^2)B)}{\sqrt{a + b \tan(c + dx)}(\tan^2(c + dx) + 1)} d \sqrt{\tan(c + dx)}}{bd(a^2 + b^2)} - \frac{2a(2Ab^3 - aB(a^2 + 3b^2))\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} \\
& \quad \downarrow \text{2257} \\
& \frac{2 \int \left(\frac{(-Ba^2 + 2Aba + b^2B)b^2 + (Aa^2 + 2bBa - Ab^2) \tan(c + dx)b^2}{\sqrt{a + b \tan(c + dx)}(\tan^2(c + dx) + 1)} - \frac{(a^2 + b^2)^2 B}{\sqrt{a + b \tan(c + dx)}} \right) d \sqrt{\tan(c + dx)}}{bd(a^2 + b^2)} - \frac{2a(2Ab^3 - aB(a^2 + 3b^2))\sqrt{\tan(c + dx)}}{bd(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} \\
& \quad \downarrow \text{2009}
\end{aligned}$$

3.463. $\int \frac{\tan^{\frac{5}{2}}(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx$

$$\frac{2a(Ab - aB) \tan^{\frac{3}{2}}(c + dx)}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{\frac{3}{2}}} - \frac{2 \left(-\frac{B(a^2 + b^2)^2 \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{b}} + \frac{b^2(a-ib)^2(-B+iA) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{2\sqrt{-b+ia}} \right)}{bd(a^2 + b^2)} - \frac{2a(2Ab^3 - aB(a^2 + 3b^2))\sqrt{\tan(c+dx)}}{bd(a^2 + b^2)\sqrt{a+b \tan(c+dx)}} + \frac{b^2(a+ib)^2(-B+iA) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{2\sqrt{-b+ia}}$$

$$b(a^2 + b^2)$$

input `Int[(Tan[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]`

output `(2*a*(A*b - a*B)*Tan[c + d*x]^(3/2))/(3*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) - ((2*((a - I*b)^2*b^2*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(2*Sqrt[I*a - b]) - ((a^2 + b^2)^2*B*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/Sqrt[b] - ((a + I*b)^2*b^2*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(2*Sqrt[I*a + b])))/(b*(a^2 + b^2)*d) - (2*a*(2*A*b^3 - a*(a^2 + 3*b^2)*B)*Sqrt[Tan[c + d*x]])/(b*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])/(b*(a^2 + b^2))`

3.463.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2035 `Int[(F_x_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[F_x, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[F_x, x]`

rule 2257 `Int[(P_x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[P_x*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[P_x, x] && IntegerQ[p]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

$$3.463. \quad \int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{5}{2}}} dx$$

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4138 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.463.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 2.18 (sec) , antiderivative size = 2978162, normalized size of antiderivative = 10560.86

output too large to display

3.463.
$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{5}{2}}} dx$$

input `int(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x)`

output `result too large to display`

3.463.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27883 vs. $2(236) = 472$.

Time = 22.13 (sec) , antiderivative size = 55768, normalized size of antiderivative = 197.76

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="fricas")`

output `Too large to include`

3.463.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output `Timed out`

3.463.7 Maxima [F]

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^{\frac{5}{2}}}{(b\tan(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^(5/2)/(b*tan(d*x + c) + a)^(5/2), x)`

3.463.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Timed out`

3.463.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{\tan(c+dx)^{5/2}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$$

input `int((tan(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)`

output `int((tan(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)`

3.463. $\int \frac{\tan^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

3.464
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

3.464.1 Optimal result 4456
 3.464.2 Mathematica [A] (verified) 4457
 3.464.3 Rubi [A] (verified) 4457
 3.464.4 Maple [B] (warning: unable to verify) 4462
 3.464.5 Fricas [B] (verification not implemented) 4462
 3.464.6 Sympy [F] 4463
 3.464.7 Maxima [F] 4463
 3.464.8 Giac [F(-1)] 4463
 3.464.9 Mupad [F(-1)] 4464

3.464.1 Optimal result

Integrand size = 35, antiderivative size = 244

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia-b)^{5/2}d} + \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia+b)^{5/2}d} + \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3b(a^2+b^2)d(a+b \tan(c+dx))^{3/2}} + \frac{2(2a^2Ab-4Ab^3+a^3B+7ab^2B)\sqrt{\tan(c+dx)}}{3b(a^2+b^2)^2d\sqrt{a+b \tan(c+dx)}}$$

output

```
(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a-b)^(5/2)/d+(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a+b)^(5/2)/d+2/3*(2*A*a^2*b-4*A*b^3+B*a^3+7*B*a*b^2)*tan(d*x+c)^(1/2)/b/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^(1/2)+2/3*a*(A*b-B*a)*tan(d*x+c)^(1/2)/b/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)
```

3.464.2 Mathematica [A] (verified)

Time = 3.35 (sec) , antiderivative size = 308, normalized size of antiderivative = 1.26

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = -\frac{3B\sqrt{\tan(c+dx)}}{(a+b \tan(c+dx))^{3/2}} + \frac{(2aAb+a^2B+3b^2B)\sqrt{\tan(c+dx)}}{(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \frac{3\sqrt[4]{-1}b}{(a+ib)^2(iA+...)} + \dots$$

```
input Integrate[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]
```

```
output ((-3*B*Sqrt[Tan[c + d*x]])/(a + b*Tan[c + d*x])^(3/2) + ((2*a*A*b + a^2*B + 3*b^2*B)*Sqrt[Tan[c + d*x]])/((a^2 + b^2)*(a + b*Tan[c + d*x])^(3/2)) + (3*(-1)^(1/4)*b*(((a + I*b)^2*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[-a + I*b] + (I*(a - I*b)^2*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[a + I*b] + (2*(2*a^2*A*b - 4*A*b^3 + a^3*B + 7*a*b^2*B)*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]/(a^2 + b^2)^2)/(3*b*d)
```

3.464.3 Rubi [A] (verified)

Time = 1.51 (sec) , antiderivative size = 293, normalized size of antiderivative = 1.20, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 4088, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx \xrightarrow{3042} \int \frac{\tan(c+dx)^{3/2}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx \xrightarrow{4088} \dots$$

$$\begin{aligned}
& \frac{2 \int -\frac{((Ba^2+2Aba+3b^2B) \tan^2(c+dx)) - 3b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3b(a^2+b^2)} + \\
& \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
& \quad \downarrow \text{27} \\
& \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b \tan(c+dx))^{3/2}} - \\
& \frac{\int -\frac{((Ba^2+2Aba+3b^2B) \tan^2(c+dx)) - 3b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3b(a^2+b^2)} \\
& \quad \downarrow \text{3042} \\
& \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b \tan(c+dx))^{3/2}} - \\
& \frac{\int -\frac{((Ba^2+2Aba+3b^2B) \tan(c+dx)^2) - 3b(Ab-aB) \tan(c+dx) + a(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3b(a^2+b^2)} \\
& \quad \downarrow \text{4132} \\
& \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b \tan(c+dx))^{3/2}} - \\
& \frac{2 \int \frac{3(ab(Aa^2+2bBa-Ab^2)) - ab(-Ba^2+2Aba+b^2B) \tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2(a^3B+2a^2Ab+7ab^2B-4Ab^3)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{3b(a^2+b^2)} \\
& \quad \downarrow \text{27} \\
& \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b \tan(c+dx))^{3/2}} - \\
& \frac{3 \int \frac{ab(Aa^2+2bBa-Ab^2)) - ab(-Ba^2+2Aba+b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2(a^3B+2a^2Ab+7ab^2B-4Ab^3)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{3b(a^2+b^2)} \\
& \quad \downarrow \text{3042} \\
& \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b \tan(c+dx))^{3/2}} - \\
& \frac{3 \int \frac{ab(Aa^2+2bBa-Ab^2)) - ab(-Ba^2+2Aba+b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2(a^3B+2a^2Ab+7ab^2B-4Ab^3)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{3b(a^2+b^2)} \\
& \quad \downarrow \text{4099}
\end{aligned}$$

3.464. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{5}{2}}} dx$

$$\frac{2a(Ab - aB)\sqrt{\tan(c + dx)}}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2(a^3B + 2a^2Ab + 7ab^2B - 4Ab^3)\sqrt{\tan(c + dx)}}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{3\left(\frac{1}{2}ab(a - ib)^2(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}ab(a + ib)^2(A - iB) \int \frac{i \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx\right)}{a(a^2 + b^2)}$$

$3b(a^2 + b^2)$

↓ 3042

$$\frac{2a(Ab - aB)\sqrt{\tan(c + dx)}}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2(a^3B + 2a^2Ab + 7ab^2B - 4Ab^3)\sqrt{\tan(c + dx)}}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{3\left(\frac{1}{2}ab(a - ib)^2(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}ab(a + ib)^2(A - iB) \int \frac{i \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx\right)}{a(a^2 + b^2)}$$

$3b(a^2 + b^2)$

↓ 4098

$$\frac{2a(Ab - aB)\sqrt{\tan(c + dx)}}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2(a^3B + 2a^2Ab + 7ab^2B - 4Ab^3)\sqrt{\tan(c + dx)}}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{3\left(\frac{ab(a - ib)^2(A + iB) \int \frac{1}{(i \tan(c + dx) + 1)\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} d \tan(c + dx)}{2d} + \frac{ab(a + ib)^2(A - iB) \int \frac{1}{(1 - i \tan(c + dx))\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} d \tan(c + dx)}{2d}\right)}{a(a^2 + b^2)}$$

$3b(a^2 + b^2)$

↓ 104

$$\frac{2a(Ab - aB)\sqrt{\tan(c + dx)}}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2(a^3B + 2a^2Ab + 7ab^2B - 4Ab^3)\sqrt{\tan(c + dx)}}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{3\left(\frac{ab(a - ib)^2(A + iB) \int \frac{1}{\frac{(ia - b) \tan(c + dx)}{a + b \tan(c + dx)} + 1} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}}{d} + \frac{ab(a + ib)^2(A - iB) \int \frac{1}{1 - \frac{(ia + b) \tan(c + dx)}{a + b \tan(c + dx)}} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}}{d}\right)}{a(a^2 + b^2)}$$

$3b(a^2 + b^2)$

↓ 216

$$\frac{2a(Ab - aB)\sqrt{\tan(c + dx)}}{3bd(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} - \frac{2(a^3B + 2a^2Ab + 7ab^2B - 4Ab^3)\sqrt{\tan(c + dx)}}{d(a^2 + b^2)\sqrt{a + b \tan(c + dx)}} + \frac{3\left(\frac{ab(a + ib)^2(A - iB) \int \frac{1}{1 - \frac{(ia + b) \tan(c + dx)}{a + b \tan(c + dx)}} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}}{d} + \frac{ab(a - ib)^2(A + iB) \arctan\left(\frac{\sqrt{-b + ia}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{d\sqrt{-b + ia}}\right)}{a(a^2 + b^2)}$$

$3b(a^2 + b^2)$

↓ 219

3.464. $\int \frac{\tan^3(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx$

$$\frac{\frac{2a(Ab - aB)\sqrt{\tan(c + dx)}}{3bd(a^2 + b^2)(a + b\tan(c + dx))^{3/2}} - \frac{2(a^3B + 2a^2Ab + 7ab^2B - 4Ab^3)\sqrt{\tan(c + dx)}}{d(a^2 + b^2)\sqrt{a + b\tan(c + dx)}} + \frac{3\left(\frac{ab(a - ib)^2(A + iB)\arctan\left(\frac{\sqrt{-b + ia}\sqrt{\tan(c + dx)}}{\sqrt{a + b\tan(c + dx)}}\right)}{d\sqrt{-b + ia}} + \frac{ab(a + ib)^2(A - iB)\operatorname{arctanh}\left(\frac{\sqrt{b + ia}\sqrt{\tan(c + dx)}}{\sqrt{a + b\tan(c + dx)}}\right)}{d\sqrt{b + ia}}\right)}{a(a^2 + b^2)}}{3b(a^2 + b^2)}$$

input `Int[(Tan[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2), x]`

output `(2*a*(A*b - a*B)*Sqrt[Tan[c + d*x]]/(3*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) - ((3*((a - I*b)^2*b*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a - b]*d) + (a*(a + I*b)^2*b*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d)))/(a*(a^2 + b^2)) - (2*(2*a^2*A*b - 4*A*b^3 + a^3*B + 7*a*b^2*B)*Sqrt[Tan[c + d*x]]/((a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]))/(3*b*(a^2 + b^2))`

3.464.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 104 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q], x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

3.464. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

3.464.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 5.39 (sec) , antiderivative size = 2978130, normalized size of antiderivative = 12205.45

output too large to display

```
input int(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x)
```

```
output result too large to display
```

3.464.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27576 vs. $2(204) = 408$.

Time = 14.84 (sec) , antiderivative size = 27576, normalized size of antiderivative = 113.02

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="fracas")
```

```
output Too large to include
```

3.464. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

3.464.6 Sympy [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx$$

input `integrate(tan(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**(3/2)/(a + b*tan(c + d*x))**(5/2), x)`

3.464.7 Maxima [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^{\frac{3}{2}}}{(b\tan(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorith="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^(3/2)/(b*tan(d*x + c) + a)^(5/2), x)`

3.464.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorith="giac")`

output `Timed out`

3.464.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx = \int \frac{\tan(c+dx)^{\frac{3}{2}}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx$$

input `int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)`

output `int((tan(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)`

3.465
$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

3.465.1 Optimal result 4465
 3.465.2 Mathematica [A] (verified) 4466
 3.465.3 Rubi [A] (verified) 4466
 3.465.4 Maple [B] (warning: unable to verify) 4471
 3.465.5 Fricas [B] (verification not implemented) 4471
 3.465.6 Sympy [F] 4471
 3.465.7 Maxima [F] 4472
 3.465.8 Giac [F(-1)] 4472
 3.465.9 Mupad [F(-1)] 4472

3.465.1 Optimal result

Integrand size = 35, antiderivative size = 244

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = -\frac{(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia-b)^{5/2}d}$$

$$+ \frac{(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia+b)^{5/2}d} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{3(a^2+b^2)d(a+b \tan(c+dx))^{3/2}}$$

$$- \frac{2(5a^2Ab-Ab^3-2a^3B+4ab^2B)\sqrt{\tan(c+dx)}}{3a(a^2+b^2)^2d\sqrt{a+b \tan(c+dx)}}$$

```
output - (I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*
a-b)^(5/2)/d+(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)
)^(1/2))/(I*a+b)^(5/2)/d-2/3*(5*A*a^2*b-A*b^3-2*B*a^3+4*B*a*b^2)*tan(d*x+
c)^(1/2)/a/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^(1/2)-2/3*(A*b-B*a)*tan(d*x+c)^(
1/2)/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)
```


3.465.2 Mathematica [A] (verified)

Time = 3.74 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.31

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{2b(Ab-aB) \tan^{\frac{3}{2}}(c+dx)}{(a+b \tan(c+dx))^{3/2}} + \frac{6b(2aAb-a^2B+b^2B) \tan^{\frac{3}{2}}(c+dx)}{(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{\sqrt[4]{-1}a^{a+ib}}{3}$$

input `Integrate[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]`

output `((2*b*(A*b - a*B)*Tan[c + d*x]^(3/2))/(a + b*Tan[c + d*x])^(3/2) + (6*b*(2*a*A*b - a^2*B + b^2*B)*Tan[c + d*x]^(3/2))/((a^2 + b^2)*Sqrt[a + b*Tan[c + d*x]]) + (3*(-(((-1)^(1/4)*a*(a + I*b)^2*(A - I*B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[-a + I*b]) + ((-1)^(1/4)*a*(a - I*b)^2*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]])/Sqrt[a + I*b] + 2*(-2*a*A*b + a^2*B - b^2*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/(a^2 + b^2))/(3*a*(a^2 + b^2)*d)`

3.465.3 Rubi [A] (verified)

Time = 1.48 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.22, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 4091, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

↓ 4091

$$\begin{aligned}
 & \frac{2 \int -\frac{2b(Ab-aB) \tan^2(c+dx)+3b(aA+bB) \tan(c+dx)+b(Ab-aB)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3b(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{3d(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 27 \\
 & \frac{\int -\frac{2b(Ab-aB) \tan^2(c+dx)+3b(aA+bB) \tan(c+dx)+b(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3b(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{3d(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 3042 \\
 & \frac{\int -\frac{2b(Ab-aB) \tan(c+dx)^2+3b(aA+bB) \tan(c+dx)+b(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3b(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{3d(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 4132 \\
 & \frac{2 \int \frac{3(ab(-Ba^2+2Aba+b^2B)+ab(Aa^2+2bBa-Ab^2)) \tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-2a^3B+5a^2Ab+4ab^2B-Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 27 \\
 & \frac{3 \int \frac{ab(-Ba^2+2Aba+b^2B)+ab(Aa^2+2bBa-Ab^2)) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-2a^3B+5a^2Ab+4ab^2B-Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 3042 \\
 & \frac{3 \int \frac{ab(-Ba^2+2Aba+b^2B)+ab(Aa^2+2bBa-Ab^2)) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-2a^3B+5a^2Ab+4ab^2B-Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow 4099 \\
 & \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{3d(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \\
 & \frac{2b(-2a^3B+5a^2Ab+4ab^2B-Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{3\left(\frac{1}{2}ab(a-ib)^2(-B+iA) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2}ab(a+ib)^2(B+iA) \int \frac{i \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx\right)}{a(a^2+b^2)} \\
 & \quad \downarrow 3042 \\
 & \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{3d(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \frac{2b(-2a^3B+5a^2Ab+4ab^2B-Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{3\left(\frac{1}{2}ab(a-ib)^2(-B+iA) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2}ab(a+ib)^2(B+iA) \int \frac{i \tan(c+dx)}{\sqrt{\tan(c+dx)}} dx\right)}{a(a^2+b^2)}
 \end{aligned}$$

3.465. $\int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\frac{-\frac{2(Ab - aB)\sqrt{\tan(c + dx)}}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-2a^3B + 5a^2Ab + 4ab^2B - Ab^3)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}}}{3b(a^2 + b^2)} + \frac{3\left(\frac{1}{2}ab(a - ib)^2(-B + iA) \int \frac{1 - i \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx - \frac{1}{2}ab(a + ib)^2(B + iA) \int \frac{i \tan(c + dx)}{\sqrt{\tan(c + dx)}} dx\right)}{a(a^2 + b^2)}$$

↓ 4098

$$\frac{-\frac{2(Ab - aB)\sqrt{\tan(c + dx)}}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-2a^3B + 5a^2Ab + 4ab^2B - Ab^3)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}}}{3b(a^2 + b^2)} + \frac{3\left(\frac{ab(a - ib)^2(-B + iA) \int \frac{1}{(i \tan(c + dx) + 1)\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} d \tan(c + dx) - \frac{ab(a + ib)^2(B + iA)}{a(a^2 + b^2)}\right)}{a(a^2 + b^2)}$$

↓ 104

$$\frac{-\frac{2(Ab - aB)\sqrt{\tan(c + dx)}}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-2a^3B + 5a^2Ab + 4ab^2B - Ab^3)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}}}{3b(a^2 + b^2)} + \frac{3\left(\frac{ab(a - ib)^2(-B + iA) \int \frac{1}{\frac{(ia - b) \tan(c + dx)}{a + b \tan(c + dx)} + 1} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}} - \frac{ab(a + ib)^2(B + iA) \int \frac{1}{1 - \frac{(ia + b) \tan(c + dx)}{a + b \tan(c + dx)}} d \tan(c + dx)}{a(a^2 + b^2)}\right)}{a(a^2 + b^2)}$$

↓ 216

$$\frac{-\frac{2(Ab - aB)\sqrt{\tan(c + dx)}}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-2a^3B + 5a^2Ab + 4ab^2B - Ab^3)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}}}{3b(a^2 + b^2)} + \frac{3\left(\frac{ab(a - ib)^2(-B + iA) \arctan\left(\frac{\sqrt{-b + ia}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{d\sqrt{-b + ia}} - \frac{ab(a + ib)^2(B + iA) \int \frac{1}{1 - \frac{(ia + b) \tan(c + dx)}{a + b \tan(c + dx)}} d \tan(c + dx)}{d}\right)}{a(a^2 + b^2)}$$

↓ 219

$$\frac{-\frac{2(Ab - aB)\sqrt{\tan(c + dx)}}{3d(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-2a^3B + 5a^2Ab + 4ab^2B - Ab^3)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}}}{3b(a^2 + b^2)} + \frac{3\left(\frac{ab(a - ib)^2(-B + iA) \arctan\left(\frac{\sqrt{-b + ia}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{d\sqrt{-b + ia}} - \frac{ab(a + ib)^2(B + iA) \operatorname{arctanh}\left(\frac{\sqrt{b + ia}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{d\sqrt{b + ia}}\right)}{a(a^2 + b^2)}$$

input `Int[(Sqrt[Tan[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2), x]`

3.465. $\int \frac{\sqrt{\tan(c + dx)}(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx$

```
output (-2*(A*b - a*B)*Sqrt[Tan[c + d*x]]/(3*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) + ((3*((a*(a - I*b)^2*b*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a - b]*d) - (a*(a + I*b)^2*b*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a + b]*d)))/(a*(a^2 + b^2)) - (2*b*(5*a^2*A*b - A*b^3 - 2*a^3*B + 4*a*b^2*B)*Sqrt[Tan[c + d*x]]/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]))/(3*b*(a^2 + b^2))
```

3.465.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 104 Int[(((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

rule 4091 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.465.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 2.54 (sec) , antiderivative size = 2978176, normalized size of antiderivative = 12205.64

output too large to display

input `int(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x)`

output `result too large to display`

3.465.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27593 vs. 2(200) = 400.

Time = 14.52 (sec) , antiderivative size = 27593, normalized size of antiderivative = 113.09

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorith="fricas")`

output `Too large to include`

3.465.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\sqrt{\tan(c+dx)}}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx$$

input `integrate(tan(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(tan(c + d*x))/(a + b*tan(c + d*x))**(5/2), x)`

3.465.7 Maxima [F]

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(B\tan(dx+c)+A)\sqrt{\tan(dx+c)}}{(b\tan(dx+c)+a)^{5/2}} dx$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(tan(d*x + c))/(b*tan(d*x + c) + a)^(5/2), x)`

3.465.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Timed out`

3.465.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$$

input `int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)`

output `int((tan(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)`

3.466 $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx$

3.466.1 Optimal result 4473
 3.466.2 Mathematica [A] (verified) 4474
 3.466.3 Rubi [A] (verified) 4474
 3.466.4 Maple [B] (warning: unable to verify) 4479
 3.466.5 Fricas [B] (verification not implemented) 4479
 3.466.6 Sympy [F(-1)] 4479
 3.466.7 Maxima [F] 4480
 3.466.8 Giac [F(-1)] 4480
 3.466.9 Mupad [F(-1)] 4480

3.466.1 Optimal result

Integrand size = 35, antiderivative size = 247

$$\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx = -\frac{(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia-b)^{5/2}d}$$

$$-\frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia+b)^{5/2}d} + \frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{3a(a^2+b^2)d(a+b \tan(c+dx))^{3/2}}$$

$$+ \frac{2b(8a^2Ab+2Ab^3-5a^3B+ab^2B)\sqrt{\tan(c+dx)}}{3a^2(a^2+b^2)^2d\sqrt{a+b \tan(c+dx)}}$$

```
output -(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*
a-b)^(5/2)/d-(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)
)^(1/2))/(I*a+b)^(5/2)/d+2/3*b*(8*A*a^2*b+2*A*b^3-5*B*a^3+B*a*b^2)*tan(d*
x+c)^(1/2)/a^2/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^(1/2)+2/3*b*(A*b-B*a)*tan(d*
x+c)^(1/2)/a/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)
```


3.466.2 Mathematica [A] (verified)

Time = 2.75 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.11

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}} dx = \frac{-3\sqrt[4]{-1} \left(\frac{(a+ib)^2(iA+B) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} \right) + \frac{i(a-ib)^2(A+...}{...}}{...}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(5/2)),x]`

output `(-3*(-1)^(1/4)*(((a + I*b)^2*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]])/Sqrt[-a + I*b] + (I*(a - I*b)^2*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]])/Sqrt[a + I*b]) + (2*b*(a^2 + b^2)*(A*b - a*B)*Sqrt[Tan[c + d*x]])/(a*(a + b*Tan[c + d*x])^(3/2)) + (2*b*(8*a^2*A*b + 2*A*b^3 - 5*a^3*B + a*b^2*B)*Sqrt[Tan[c + d*x]])/(a^2*Sqrt[a + b*Tan[c + d*x]]))/(3*(a^2 + b^2)^2*d)`

3.466.3 Rubi [A] (verified)

Time = 1.57 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.21, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}} dx$$

↓ 4092

$$\frac{2 \int \frac{3Aa^2 + bBa - 3(Ab - aB) \tan(c + dx) a + 2Ab^2 + 2b(Ab - aB) \tan^2(c + dx)}{2\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx}{3a(a^2 + b^2)} + \frac{2b(Ab - aB)\sqrt{\tan(c + dx)}}{3ad(a^2 + b^2)(a + b \tan(c + dx))^{3/2}}$$

3.466. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \int \frac{3Aa^2 + bBa - 3(Ab - aB) \tan(c+dx)a + 2Ab^2 + 2b(Ab - aB) \tan^2(c+dx)}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))^{3/2}}} dx \\
& \quad \frac{2b(Ab - aB) \sqrt{\tan(c+dx)}}{3ad(a^2 + b^2)(a + b \tan(c+dx))^{3/2}} \\
& \quad \frac{3Aa^2 + bBa - 3(Ab - aB) \tan(c+dx)a + 2Ab^2 + 2b(Ab - aB) \tan(c+dx)^2}{3a(a^2 + b^2)} dx \\
& \quad \frac{2b(Ab - aB) \sqrt{\tan(c+dx)}}{3ad(a^2 + b^2)(a + b \tan(c+dx))^{3/2}} \\
& \quad 2 \int \frac{3(a^2(Aa^2 + 2bBa - Ab^2) - a^2(-Ba^2 + 2Aba + b^2B) \tan(c+dx))}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx \\
& \quad \frac{2b(-5a^3B + 8a^2Ab + ab^2B + 2Ab^3) \sqrt{\tan(c+dx)}}{ad(a^2 + b^2)\sqrt{a+b \tan(c+dx)}} + \\
& \quad \frac{3a(a^2 + b^2)}{3ad(a^2 + b^2)(a + b \tan(c+dx))^{3/2}} \\
& \quad \frac{2b(Ab - aB) \sqrt{\tan(c+dx)}}{3ad(a^2 + b^2)(a + b \tan(c+dx))^{3/2}} \\
& \quad 3 \int \frac{a^2(Aa^2 + 2bBa - Ab^2) - a^2(-Ba^2 + 2Aba + b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx \\
& \quad \frac{2b(-5a^3B + 8a^2Ab + ab^2B + 2Ab^3) \sqrt{\tan(c+dx)}}{ad(a^2 + b^2)\sqrt{a+b \tan(c+dx)}} + \\
& \quad \frac{3a(a^2 + b^2)}{3ad(a^2 + b^2)(a + b \tan(c+dx))^{3/2}} \\
& \quad \frac{2b(Ab - aB) \sqrt{\tan(c+dx)}}{3ad(a^2 + b^2)(a + b \tan(c+dx))^{3/2}} \\
& \quad 3 \int \frac{a^2(Aa^2 + 2bBa - Ab^2) - a^2(-Ba^2 + 2Aba + b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx \\
& \quad \frac{2b(-5a^3B + 8a^2Ab + ab^2B + 2Ab^3) \sqrt{\tan(c+dx)}}{ad(a^2 + b^2)\sqrt{a+b \tan(c+dx)}} + \\
& \quad \frac{3a(a^2 + b^2)}{3ad(a^2 + b^2)(a + b \tan(c+dx))^{3/2}} \\
& \quad \frac{2b(Ab - aB) \sqrt{\tan(c+dx)}}{3ad(a^2 + b^2)(a + b \tan(c+dx))^{3/2}} \\
& \quad \frac{2b(Ab - aB) \sqrt{\tan(c+dx)}}{3ad(a^2 + b^2)(a + b \tan(c+dx))^{3/2}} + \\
& \quad \frac{2b(-5a^3B + 8a^2Ab + ab^2B + 2Ab^3) \sqrt{\tan(c+dx)}}{ad(a^2 + b^2)\sqrt{a+b \tan(c+dx)}} + \frac{3\left(\frac{1}{2}a^2(a-ib)^2(A+ib) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}a^2(a+ib)^2(A-ib) \int \frac{i \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx\right)}{a(a^2 + b^2)} \\
& \quad \frac{3a(a^2 + b^2)}{3a(a^2 + b^2)} \\
& \quad \frac{2b(Ab - aB) \sqrt{\tan(c+dx)}}{3ad(a^2 + b^2)(a + b \tan(c+dx))^{3/2}}
\end{aligned}$$

3.466. $\int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))^{5/2}}} dx$

$$\frac{\frac{2b(Ab - aB)\sqrt{\tan(c + dx)}}{3ad(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-5a^3B + 8a^2Ab + ab^2B + 2Ab^3)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}}}{3a(a^2 + b^2)} + \frac{3\left(\frac{1}{2}a^2(a - ib)^2(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx + \frac{1}{2}a^2(a + ib)^2(A - iB) \int \frac{i \tan(c + dx)}{\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} dx\right)}{a(a^2 + b^2)}$$

4098

$$\frac{\frac{2b(Ab - aB)\sqrt{\tan(c + dx)}}{3ad(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-5a^3B + 8a^2Ab + ab^2B + 2Ab^3)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}}}{3a(a^2 + b^2)} + \frac{3\left(\frac{a^2(a - ib)^2(A + iB) \int \frac{1}{(i \tan(c + dx) + 1)\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} d \tan(c + dx)}{2d} + \frac{a^2(a + ib)^2(A - iB) \int \frac{1}{(1 - i \tan(c + dx))\sqrt{\tan(c + dx)}\sqrt{a + b \tan(c + dx)}} d \tan(c + dx)}{2d}\right)}{a(a^2 + b^2)}$$

104

$$\frac{\frac{2b(Ab - aB)\sqrt{\tan(c + dx)}}{3ad(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-5a^3B + 8a^2Ab + ab^2B + 2Ab^3)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}}}{3a(a^2 + b^2)} + \frac{3\left(\frac{a^2(a - ib)^2(A + iB) \int \frac{1}{\frac{(ia - b) \tan(c + dx)}{a + b \tan(c + dx)} + 1} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}}{d} + \frac{a^2(a + ib)^2(A - iB) \int \frac{1}{1 - \frac{(ia + b) \tan(c + dx)}{a + b \tan(c + dx)}} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}}{d}\right)}{a(a^2 + b^2)}$$

216

$$\frac{\frac{2b(Ab - aB)\sqrt{\tan(c + dx)}}{3ad(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-5a^3B + 8a^2Ab + ab^2B + 2Ab^3)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}}}{3a(a^2 + b^2)} + \frac{3\left(\frac{a^2(a + ib)^2(A - iB) \int \frac{1}{1 - \frac{(ia + b) \tan(c + dx)}{a + b \tan(c + dx)}} d \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}}{d} + \frac{a^2(a - ib)^2(A + iB) \arctan\left(\frac{\sqrt{-b + ia}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{d\sqrt{-b + ia}}\right)}{a(a^2 + b^2)}$$

219

$$\frac{\frac{2b(Ab - aB)\sqrt{\tan(c + dx)}}{3ad(a^2 + b^2)(a + b \tan(c + dx))^{3/2}} + \frac{2b(-5a^3B + 8a^2Ab + ab^2B + 2Ab^3)\sqrt{\tan(c + dx)}}{ad(a^2 + b^2)\sqrt{a + b \tan(c + dx)}}}{3a(a^2 + b^2)} + \frac{3\left(\frac{a^2(a - ib)^2(A + iB) \arctan\left(\frac{\sqrt{-b + ia}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{d\sqrt{-b + ia}} + \frac{a^2(a + ib)^2(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{b + ia}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{d\sqrt{b + ia}}\right)}{a(a^2 + b^2)}$$

```
input Int[(A + B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(5/2)),x
]
```

3.466. $\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}} dx$

```
output (2*b*(A*b - a*B)*Sqrt[Tan[c + d*x]]/(3*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x
])^(3/2)) + ((3*((a^2*(a - I*b)^2*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan
[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a - b]*d) + (a^2*(a + I*b)^
2*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c +
d*x]])]/(Sqrt[I*a + b]*d)))/(a*(a^2 + b^2)) + (2*b*(8*a^2*A*b + 2*A*b^3 -
5*a^3*B + a*b^2*B)*Sqrt[Tan[c + d*x]]/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c +
d*x]]))/(3*a*(a^2 + b^2))
```

3.466.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 104 Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))/((e_.) + (f_.)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x
] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.466.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 3.07 (sec) , antiderivative size = 2975233, normalized size of antiderivative = 12045.48

output too large to display

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(5/2),x)`

output `result too large to display`

3.466.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27639 vs. 2(208) = 416.

Time = 14.52 (sec) , antiderivative size = 27639, normalized size of antiderivative = 111.90

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(5/2),x, algorith="fracas")`

output `Too large to include`

3.466.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+b*tan(d*x+c))**(5/2),x)`

output `Timed out`

3.466.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a)^{5/2} \sqrt{\tan(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)/((b*tan(d*x + c) + a)^(5/2)*sqrt(tan(d*x +
c))), x)`

3.466.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="giac")`

output `Timed out`

3.466.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(5/2)),x
)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(5/2)),
x)`

$$3.467 \quad \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$$

3.467.1 Optimal result	4481
3.467.2 Mathematica [A] (verified)	4482
3.467.3 Rubi [A] (verified)	4482
3.467.4 Maple [B] (warning: unable to verify)	4487
3.467.5 Fracas [B] (verification not implemented)	4488
3.467.6 Sympy [F(-1)]	4488
3.467.7 Maxima [F(-1)]	4488
3.467.8 Giac [F(-1)]	4489
3.467.9 Mupad [F(-1)]	4489

3.467.1 Optimal result

Integrand size = 35, antiderivative size = 301

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \frac{(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia - b)^{5/2}d} - \frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia + b)^{5/2}d} - \frac{2A}{ad\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} - \frac{2b(3a^2A + 4Ab^2 - abB) \sqrt{\tan(c + dx)}}{3a^2(a^2 + b^2)d(a + b \tan(c + dx))^{3/2}} - \frac{2b(3a^4A + 17a^2Ab^2 + 8Ab^4 - 8a^3bB - 2ab^3B) \sqrt{\tan(c + dx)}}{3a^3(a^2 + b^2)^2d\sqrt{a + b \tan(c + dx)}}$$

```
output (I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a
-b)^(5/2)/d-(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)
)^(1/2))/(I*a+b)^(5/2)/d-2/3*b*(3*A*a^4+17*A*a^2*b^2+8*A*b^4-8*B*a^3*b-2*B
*a*b^3)*tan(d*x+c)^(1/2)/a^3/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^(1/2)-2*A/a/d/
tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2)-2/3*b*(3*A*a^2+4*A*b^2-B*a*b)*tan(
d*x+c)^(1/2)/a^2/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)
```

3.467. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$

3.467.2 Mathematica [A] (verified)

Time = 4.03 (sec) , antiderivative size = 326, normalized size of antiderivative = 1.08

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = -\frac{6aA}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} - \frac{2b(3a^2A+4Ab^2-abB)\sqrt{\tan(c+dx)}}{(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \frac{3\sqrt[4]{-a^2+b^2} \operatorname{arctan}\left(\frac{\sqrt{\tan(c+dx)}}{\sqrt[4]{-a^2+b^2}}\right)}{\sqrt[4]{-a^2+b^2}}$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(5/2)),x]`

output `((-6*a*A)/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)) - (2*b*(3*a^2*A + 4*A*b^2 - a*b*B)*Sqrt[Tan[c + d*x]])/((a^2 + b^2)*(a + b*Tan[c + d*x])^(3/2)) + (3*(-1)^(1/4)*a^3*((a + I*b)^2*(A - I*B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[-a + I*b] - ((a - I*b)^2*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[a + I*b] - (2*b*(3*a^4*A + 17*a^2*A*b^2 + 8*A*b^4 - 8*a^3*b*B - 2*a*b^3*B)*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(a*(a^2 + b^2)^2))/(3*a^2*d)`

3.467.3 Rubi [A] (verified)

Time = 1.99 (sec) , antiderivative size = 361, normalized size of antiderivative = 1.20, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2}(a + b \tan(c + dx))^{5/2}} dx$$

↓ 4092

3.467. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{2 \int \frac{4Ab \tan^2(c+dx) + aA \tan(c+dx) + 4Ab - aB}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2A}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{4Ab \tan^2(c+dx) + aA \tan(c+dx) + 4Ab - aB}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2A}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{4Ab \tan(c+dx)^2 + aA \tan(c+dx) + 4Ab - aB}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2A}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 4132 \\
 & \frac{2 \int \frac{-3Ba^3 + 9Aba^2 + 3(aA + bB) \tan(c+dx)a^2 - 2b^2Ba + 8Ab^3 + 2b(3Aa^2 - bBa + 4Ab^2) \tan^2(c+dx)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3a(a^2+b^2)} + \frac{2b(3a^2A - abB + 4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \frac{2A^a}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{-3Ba^3 + 9Aba^2 + 3(aA + bB) \tan(c+dx)a^2 - 2b^2Ba + 8Ab^3 + 2b(3Aa^2 - bBa + 4Ab^2) \tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3a(a^2+b^2)} + \frac{2b(3a^2A - abB + 4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \frac{2A^a}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{-3Ba^3 + 9Aba^2 + 3(aA + bB) \tan(c+dx)a^2 - 2b^2Ba + 8Ab^3 + 2b(3Aa^2 - bBa + 4Ab^2) \tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3a(a^2+b^2)} + \frac{2b(3a^2A - abB + 4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \frac{2A^a}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 4132 \\
 & \frac{2 \int \frac{3\left((-Ba^2 + 2Aba + b^2B)a^3 + (Aa^2 + 2bBa - Ab^2) \tan(c+dx)a^3\right)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(3a^4A - 8a^3bB + 17a^2Ab^2 - 2ab^3B + 8Ab^4)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{2b(3a^2A - abB + 4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \frac{2A^a}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}
 \end{aligned}$$

3.467. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$

↓ 27

$$\frac{\int \frac{(-Ba^2+2Aba+b^2B)a^3+(Aa^2+2bBa-Ab^2)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{2b(3a^4A-8a^3bB+17a^2Ab^2-2ab^3B+8Ab^4)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{3a(a^2+b^2)} + \frac{2b(3a^2A-abB+4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))} - \frac{2A}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}$$

↓ 3042

$$\frac{\int \frac{(-Ba^2+2Aba+b^2B)a^3+(Aa^2+2bBa-Ab^2)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{2b(3a^4A-8a^3bB+17a^2Ab^2-2ab^3B+8Ab^4)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{3a(a^2+b^2)} + \frac{2b(3a^2A-abB+4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))} - \frac{2A}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}$$

↓ 4099

$$\frac{2b(3a^2A-abB+4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{2A}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} - \frac{2b(3a^4A-8a^3bB+17a^2Ab^2-2ab^3B+8Ab^4)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{3\left(\frac{1}{2}a^3(a-ib)^2(-B+iA)\int \frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx\right)}{3a(a^2+b^2)}$$

↓ 3042

$$\frac{2b(3a^2A-abB+4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{2A}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} - \frac{2b(3a^4A-8a^3bB+17a^2Ab^2-2ab^3B+8Ab^4)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{3\left(\frac{1}{2}a^3(a-ib)^2(-B+iA)\int \frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx\right)}{3a(a^2+b^2)}$$

↓ 4098

$$\frac{2b(3a^2A-abB+4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{2A}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} - \frac{2b(3a^4A-8a^3bB+17a^2Ab^2-2ab^3B+8Ab^4)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{3\left(\frac{a^3(a-ib)^2(-B+iA)\int \frac{1}{(i\tan(c+dx)+1)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx\right)}{3a(a^2+b^2)}$$

↓ 104

3.467. $\int \frac{A+B\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}} dx$

$$\frac{2A}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} + \frac{2b(3a^4A-8a^3bB+17a^2Ab^2-2ab^3B+8Ab^4)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{3 \left(\frac{a^3(a-ib)^2(-B+iA) \int \frac{1}{\frac{(ia-b)\tan(c+dx)}{a+b\tan(c+dx)}+1} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}} \right)}{3a(a^2+b^2)}$$

216

$$\frac{2A}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} + \frac{2b(3a^4A-8a^3bB+17a^2Ab^2-2ab^3B+8Ab^4)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{3 \left(\frac{a^3(a-ib)^2(-B+iA) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} \right)}{3a(a^2+b^2)}$$

219

$$\frac{2A}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} + \frac{2b(3a^4A-8a^3bB+17a^2Ab^2-2ab^3B+8Ab^4)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{3 \left(\frac{a^3(a-ib)^2(-B+iA) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} \right)}{3a(a^2+b^2)}$$

```
input Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(5/2)),x]
```

```
output (-2*A)/(a*d*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)) - ((2*b*(3*a^2*A + 4*A*b^2 - a*b*B)*Sqrt[Tan[c + d*x]])/(3*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) + ((3*((a^3*(a - I*b)^2*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a - b]*d) - (a^3*(a + I*b)^2*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a + b]*d)))/(a*(a^2 + b^2)) + (2*b*(3*a^4*A + 17*a^2*A*b^2 + 8*A*b^4 - 8*a^3*b*B - 2*a*b^3*B)*Sqrt[Tan[c + d*x]])/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]])/(3*a*(a^2 + b^2))/a
```

3.467. $\int \frac{A+B \tan(c+dx)}{\tan^3(c+dx)(a+b \tan(c+dx))^{5/2}} dx$

3.467.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.467.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 2.10 (sec) , antiderivative size = 2976756, normalized size of antiderivative = 9889.55

output too large to display

input `int((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(5/2),x)`

output `result too large to display`

3.467.
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$$

3.467.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27749 vs. $2(257) = 514$.

Time = 14.45 (sec) , antiderivative size = 27749, normalized size of antiderivative = 92.19

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="fracas")`

output Too large to include

3.467.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(3/2)/(a+b*tan(d*x+c))**(5/2),x)`

output Timed out

3.467.7 Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="maxima")`

output Timed out

3.467. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$

3.467.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="giac")`

output `Timed out`

3.467.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{3/2} (a + b \tan(c + dx))^{5/2}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(5/2)),x
)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(5/2)),
x)`

3.468
$$\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$$

3.468.1 Optimal result	4490
3.468.2 Mathematica [A] (verified)	4491
3.468.3 Rubi [A] (verified)	4491
3.468.4 Maple [B] (warning: unable to verify)	4497
3.468.5 Fracas [B] (verification not implemented)	4498
3.468.6 Sympy [F(-1)]	4498
3.468.7 Maxima [F(-1)]	4498
3.468.8 Giac [F(-1)]	4499
3.468.9 Mupad [F(-1)]	4499

3.468.1 Optimal result

Integrand size = 35, antiderivative size = 359

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \frac{(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia - b)^{5/2}d} + \frac{(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(ia + b)^{5/2}d} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} + \frac{2(2Ab - aB)}{a^2 d \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} + \frac{2b(7a^2 Ab + 8Ab^3 - 3a^3 B - 4ab^2 B) \sqrt{\tan(c + dx)}}{3a^3 (a^2 + b^2) d (a + b \tan(c + dx))^{3/2}} + \frac{2b(8a^4 Ab + 30a^2 Ab^3 + 16Ab^5 - 3a^5 B - 17a^3 b^2 B - 8ab^4 B) \sqrt{\tan(c + dx)}}{3a^4 (a^2 + b^2)^2 d \sqrt{a + b \tan(c + dx)}}$$

```
output (A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a-b)^(5/2)/d+(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/(I*a+b)^(5/2)/d+2/3*b*(8*A*a^4*b+30*A*a^2*b^3+16*A*b^5-3*B*a^5-17*B*a^3*b^2-8*B*a*b^4)*tan(d*x+c)^(1/2)/a^4/(a^2+b^2)^2/d/(a+b*tan(d*x+c))^(1/2)+2*(2*A*b-B*a)/a^2/d/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2)+2/3*b*(7*A*a^2*b+8*A*b^3-3*B*a^3-4*B*a*b^2)*tan(d*x+c)^(1/2)/a^3/(a^2+b^2)/d/(a+b*tan(d*x+c))^(3/2)-2/3*A/a/d/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2)
```

3.468.2 Mathematica [A] (verified)

Time = 3.89 (sec) , antiderivative size = 383, normalized size of antiderivative = 1.07

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = -\frac{6A}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} + \frac{6(6Ab-3aB)}{a\sqrt{\tan(c+dx)(a+b \tan(c+dx))^{3/2}}} + \frac{6b(7a^2A}{a$$

input `Integrate[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^(5/2)),x]`

output `((-6*A)/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2)) + (6*(6*A*b - 3*a*B))/(a*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)) + (6*b*(7*a^2*A*b + 8*A*b^3 - 3*a^3*B - 4*a*b^2*B)*Sqrt[Tan[c + d*x]])/(a^2*(a^2 + b^2)*(a + b*Tan[c + d*x])^(3/2)) + (9*(-1)^(3/4)*a^4*((a + I*b)^2*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[-a + I*b] + ((a - I*b)^2*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[a + I*b]) + (6*b*(8*a^4*A*b + 30*a^2*A*b^3 + 16*A*b^5 - 3*a^5*B - 17*a^3*b^2*B - 8*a*b^4*B)*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]])/(a^3*(a^2 + b^2)^2))/(9*a*d)`

3.468.3 Rubi [A] (verified)

Time = 2.60 (sec) , antiderivative size = 424, normalized size of antiderivative = 1.18, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.543$, Rules used = {3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2}(a + b \tan(c + dx))^{5/2}} dx$$

↓ 4092

3.468. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{2 \int \frac{3(2Ab \tan^2(c+dx) + aA \tan(c+dx) + 2Ab - aB)}{2 \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx}{3a} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{2Ab \tan^2(c+dx) + aA \tan(c+dx) + 2Ab - aB}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{2Ab \tan(c+dx)^2 + aA \tan(c+dx) + 2Ab - aB}{\tan(c+dx)^{3/2}(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 4132 \\
 & \frac{2 \int -\frac{Aa^2 + B \tan(c+dx)a^2 + 4bBa - 8Ab^2 - 4b(2Ab - aB) \tan^2(c+dx)}{2 \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2(2Ab - aB)}{ad \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \\
 & \quad \frac{a}{2A} \\
 & \quad \frac{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}}{27} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{Aa^2 + B \tan(c+dx)a^2 + 4bBa - 8Ab^2 - 4b(2Ab - aB) \tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2(2Ab - aB)}{ad \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \\
 & \quad \frac{a}{2A} \\
 & \quad \frac{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}}{3042} \\
 & \quad \downarrow 3042 \\
 & \frac{\int \frac{Aa^2 + B \tan(c+dx)a^2 + 4bBa - 8Ab^2 - 4b(2Ab - aB) \tan(c+dx)^2}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2(2Ab - aB)}{ad \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \\
 & \quad \frac{a}{2A} \\
 & \quad \frac{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}}{4132} \\
 & \quad \downarrow 4132 \\
 & \frac{2 \int \frac{3Aa^4 + 9bBa^3 - 3(Ab - aB) \tan(c+dx)a^3 - 14Ab^2a^2 + 8b^3Ba - 16Ab^4 - 2b(-3Ba^3 + 7Aba^2 - 4b^2Ba + 8Ab^3) \tan^2(c+dx)}{2 \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3a(a^2+b^2)} - \frac{2b(-3a^3B + 7a^2Ab - 4ab^2B + 8Ab^3)}{3ad(a^2+b^2)(a+b \tan(c+dx))} \\
 & \quad \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} \\
 & \quad \downarrow 27
 \end{aligned}$$

3.468. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$

$$\int \frac{3Aa^4 + 9bBa^3 - 3(Ab - aB) \tan(c+dx)a^3 - 14Ab^2a^2 + 8b^3Ba - 16Ab^4 - 2b(-3Ba^3 + 7Aba^2 - 4b^2Ba + 8Ab^3) \tan^2(c+dx)}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))^{3/2}} \cdot 3a(a^2+b^2)} dx - \frac{2b(-3a^3B + 7a^2Ab - 4ab^2B + 8Ab^3) \sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))}$$

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}}$$

3042

$$\int \frac{3Aa^4 + 9bBa^3 - 3(Ab - aB) \tan(c+dx)a^3 - 14Ab^2a^2 + 8b^3Ba - 16Ab^4 - 2b(-3Ba^3 + 7Aba^2 - 4b^2Ba + 8Ab^3) \tan(c+dx)^2}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))^{3/2}} \cdot 3a(a^2+b^2)} dx - \frac{2b(-3a^3B + 7a^2Ab - 4ab^2B + 8Ab^3) \sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))}$$

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}}$$

4132

$$2 \int \frac{3(a^4(Aa^2 + 2bBa - Ab^2) - a^4(-Ba^2 + 2Aba + b^2B) \tan(c+dx))}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{2b(-3a^5B + 8a^4Ab - 17a^3b^2B + 30a^2Ab^3 - 8ab^4B + 16Ab^5) \sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} - \frac{2b(-3a^3B + 7a^2Ab - 4ab^2B + 8Ab^3) \sqrt{\tan(c+dx)}}{3ad(a^2+b^2)}$$

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}}$$

27

$$3 \int \frac{a^4(Aa^2 + 2bBa - Ab^2) - a^4(-Ba^2 + 2Aba + b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{2b(-3a^5B + 8a^4Ab - 17a^3b^2B + 30a^2Ab^3 - 8ab^4B + 16Ab^5) \sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} - \frac{2b(-3a^3B + 7a^2Ab - 4ab^2B + 8Ab^3) \sqrt{\tan(c+dx)}}{3ad(a^2+b^2)}$$

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}}$$

3042

$$3 \int \frac{a^4(Aa^2 + 2bBa - Ab^2) - a^4(-Ba^2 + 2Aba + b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{2b(-3a^5B + 8a^4Ab - 17a^3b^2B + 30a^2Ab^3 - 8ab^4B + 16Ab^5) \sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} - \frac{2b(-3a^3B + 7a^2Ab - 4ab^2B + 8Ab^3) \sqrt{\tan(c+dx)}}{3ad(a^2+b^2)}$$

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}}$$

3.468. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$

$$\begin{array}{c}
 \downarrow 4099 \\
 2A \\
 \hline
 3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2} \\
 \hline
 \frac{2(2Ab-aB)}{ad\sqrt{\tan(c+dx)(a+b \tan(c+dx))^{3/2}}} + \frac{-\frac{2b(-3a^3B+7a^2Ab-4ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{-\frac{2b(-3a^5B+8a^4Ab-17a^3b^2B+30a^2Ab^3-8ab^4B+16Ab^5)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}
 \end{array}$$

a

$$\begin{array}{c}
 \downarrow 3042 \\
 2A \\
 \hline
 3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2} \\
 \hline
 \frac{2(2Ab-aB)}{ad\sqrt{\tan(c+dx)(a+b \tan(c+dx))^{3/2}}} + \frac{-\frac{2b(-3a^3B+7a^2Ab-4ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{-\frac{2b(-3a^5B+8a^4Ab-17a^3b^2B+30a^2Ab^3-8ab^4B+16Ab^5)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}
 \end{array}$$

a

$$\begin{array}{c}
 \downarrow 4098 \\
 2A \\
 \hline
 3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2} \\
 \hline
 \frac{2(2Ab-aB)}{ad\sqrt{\tan(c+dx)(a+b \tan(c+dx))^{3/2}}} + \frac{-\frac{2b(-3a^3B+7a^2Ab-4ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{-\frac{2b(-3a^5B+8a^4Ab-17a^3b^2B+30a^2Ab^3-8ab^4B+16Ab^5)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}
 \end{array}$$

$$\begin{array}{c}
 \downarrow 104 \\
 2A \\
 \hline
 3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2} \\
 \hline
 \frac{2(2Ab-aB)}{ad\sqrt{\tan(c+dx)(a+b \tan(c+dx))^{3/2}}} + \frac{-\frac{2b(-3a^3B+7a^2Ab-4ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{-\frac{2b(-3a^5B+8a^4Ab-17a^3b^2B+30a^2Ab^3-8ab^4B+16Ab^5)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}
 \end{array}$$

a

$$\downarrow 216$$

3.468. $\int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} - \frac{2(2Ab-aB)}{ad \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} + \frac{-2b(-3a^3B+7a^2Ab-4ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \frac{-2b(-3a^5B+8a^4Ab-17a^3b^2B+30a^2Ab^3-8ab^4B+16Ab^5)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}$$

a

↓ 219

$$\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} - \frac{2(2Ab-aB)}{ad \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} + \frac{-2b(-3a^3B+7a^2Ab-4ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \frac{-2b(-3a^5B+8a^4Ab-17a^3b^2B+30a^2Ab^3-8ab^4B+16Ab^5)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}$$

a

input `Int[(A + B*Tan[c + d*x])/(Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^(5/2)),x]`

output `(-2*A)/(3*a*d*Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2)) - ((-2*(2*A*b - a*B))/(a*d*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)) + ((-2*b*(7*a^2*A*b + 8*A*b^3 - 3*a^3*B - 4*a*b^2*B)*Sqrt[Tan[c + d*x]])/(3*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) + ((3*((a^4*(a - I*b)^2*(A + I*B)*ArcTan[Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]/(Sqrt[I*a - b]*d) + (a^4*(a + I*b)^2*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]/(Sqrt[I*a + b]*d)))/(a*(a^2 + b^2)) - (2*b*(8*a^4*A*b + 30*a^2*A*b^3 + 16*A*b^5 - 3*a^5*B - 17*a^3*b^2*B - 8*a*b^4*B)*Sqrt[Tan[c + d*x]])/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]))/(3*a*(a^2 + b^2))/a)/a`

3.468.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4098 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*
x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]
```

```
rule 4099 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2,
0]
```

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
(!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

3.468.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 5.47 (sec) , antiderivative size = 2982515, normalized size of antiderivative = 8307.84

output too large to display

```
input int((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c))^(5/2),x)
```

```
output result too large to display
```

$$3.468. \quad \int \frac{A+B \tan(c+dx)}{\tan^2(c+dx)(a+b \tan(c+dx))^{5/2}} dx$$

3.468.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27865 vs. $2(308) = 616$.

Time = 14.52 (sec) , antiderivative size = 27865, normalized size of antiderivative = 77.62

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="fracas")`

output Too large to include

3.468.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)**(5/2)/(a+b*tan(d*x+c))**(5/2),x)`

output Timed out

3.468.7 Maxima [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="maxima")`

output Timed out

3.468.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/tan(d*x+c)^(5/2)/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="giac")`

output `Timed out`

3.468.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{\tan(c + dx)^{5/2} (a + b \tan(c + dx))^{5/2}} dx$$

input `int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + b*tan(c + d*x))^(5/2)),x
)`

output `int((A + B*tan(c + d*x))/(tan(c + d*x)^(5/2)*(a + b*tan(c + d*x))^(5/2)),
x)`

3.469
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.469.1 Optimal result	4500
3.469.2 Mathematica [A] (verified)	4500
3.469.3 Rubi [A] (verified)	4501
3.469.4 Maple [B] (warning: unable to verify)	4503
3.469.5 Fricas [B] (verification not implemented)	4503
3.469.6 Sympy [F]	4503
3.469.7 Maxima [F]	4504
3.469.8 Giac [F(-1)]	4504
3.469.9 Mupad [F(-1)]	4504

3.469.1 Optimal result

Integrand size = 38, antiderivative size = 155

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = -\frac{B \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia-bd}} + \frac{2B \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{bd}} - \frac{B \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia+bd}}$$

output
$$-B*\arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(dx+c)}}{\sqrt{a+b*\tan(dx+c)}}\right)/\sqrt{ia-bd} + 2*B*\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(dx+c)}}{\sqrt{a+b*\tan(dx+c)}}\right)/\sqrt{bd} - B*\operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(dx+c)}}{\sqrt{a+b*\tan(dx+c)}}\right)/\sqrt{ia+bd}$$

3.469.2 Mathematica [A] (verified)

Time = 0.84 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.23

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{B \left((-1)^{3/4} \left(\frac{\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} \right) + \frac{\arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{a+ib}} \right)}{d}$$

3.469.
$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

input `Integrate[(Tan[c + d*x]^(3/2)*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `(B*((-1)^(3/4)*(ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[-a + I*b] + ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[a + I*b] + (2*Sqrt[a]*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/Sqrt[b]*Sqrt[a + b*Tan[c + d*x]))/d`

3.469.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.95, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.132$, Rules used = {2011, 3042, 4058, 614, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{3}{2}}} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{\tan^{\frac{3}{2}}(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{\tan(c+dx)^{\frac{3}{2}}}{\sqrt{a+b \tan(c+dx)}} dx \\
 & \quad \downarrow \text{4058} \\
 & \frac{B \int \frac{\tan^{\frac{3}{2}}(c+dx)}{\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} d \tan(c+dx)}{d} \\
 & \quad \downarrow \text{614} \\
 & \frac{B \int \left(\frac{1}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} - \frac{1}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} \right) d \tan(c+dx)}{d} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

3.469. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{3}{2}}} dx$

$$\frac{B \left(-\frac{\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b}\tan(c+dx)}\right)}{\sqrt{-b+ia}} + \frac{2\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b}\tan(c+dx)}\right)}{\sqrt{b}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b}\tan(c+dx)}\right)}{\sqrt{b+ia}} \right)}{d}$$

input `Int[(Tan[c + d*x]^(3/2)*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `(B*(-(ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[I*a - b]) + (2*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[b] - ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[I*a + b]))/d`

3.469.3.1 Defintions of rubi rules used

rule 614 `Int[((e_.)*(x_))^(m_)/(Sqrt[(c_) + (d_.)*(x_)]*((a_) + (b_.)*(x_)^2)), x_Symbol] := Simp[e^(m + 1/2) Int[ExpandIntegrand[1/(Sqrt[e*x]*Sqrt[c + d*x]), x^(m + 1/2)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[m - 1/2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4058 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*((c + d*ff*x)^n/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.469. $\int \frac{\tan^3(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$

3.469.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.31 (sec) , antiderivative size = 943434, normalized size of antiderivative = 6086.67

output too large to display

input `int(tan(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x)`

output `result too large to display`

3.469.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4592 vs. $2(123) = 246$.

Time = 1.05 (sec) , antiderivative size = 9186, normalized size of antiderivative = 59.26

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate(tan(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,
algorithm="fricas")`

output `Too large to include`

3.469.6 Sympy [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = B \int \frac{\tan^{\frac{3}{2}}(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx$$

input `integrate(tan(d*x+c)**(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x
)`

output `B*Integral(tan(c + d*x)**(3/2)/sqrt(a + b*tan(c + d*x)), x)`

3.469.7 Maxima [F]

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx = \int \frac{(Bb\tan(dx+c)+Ba)\tan(dx+c)^{\frac{3}{2}}}{(b\tan(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(tan(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,
algorithm="maxima")`

output `integrate((B*b*tan(d*x + c) + B*a)*tan(d*x + c)^(3/2)/(b*tan(d*x + c) + a)
^(3/2), x)`

3.469.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,
algorithm="giac")`

output `Timed out`

3.469.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx = \int \frac{\tan(c+dx)^{\frac{3}{2}}(Ba+Bb\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx$$

input `int((tan(c + d*x)^(3/2)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/
2),x)`

output `int((tan(c + d*x)^(3/2)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/
2), x)`

3.469. $\int \frac{\tan^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx$

$$3.470 \quad \int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.470.1 Optimal result	4505
3.470.2 Mathematica [A] (verified)	4505
3.470.3 Rubi [A] (verified)	4506
3.470.4 Maple [B] (warning: unable to verify)	4508
3.470.5 Fricas [B] (verification not implemented)	4508
3.470.6 Sympy [F]	4509
3.470.7 Maxima [F]	4510
3.470.8 Giac [F(-1)]	4510
3.470.9 Mupad [F(-1)]	4510

3.470.1 Optimal result

Integrand size = 38, antiderivative size = 117

$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{iB \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia-bd}} - \frac{iB \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia+bd}}$$

output `I*B*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d/(I*a-b)^(1/2)-I*B*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d/(I*a+b)^(1/2)`

3.470.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.06

$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{\sqrt[4]{-1}B \left(-\frac{\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} + \frac{\arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{a+ib}} \right)}{d}$$

input `Integrate[(Sqrt[Tan[c + d*x]]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

3.470. $\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

output $((-1)^{(1/4)}*B*(-(\text{ArcTan}[((-1)^{(1/4)}*\text{Sqrt}[-a + I*b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a + b*\text{Tan}[c + d*x]])/\text{Sqrt}[-a + I*b]) + \text{ArcTan}[((-1)^{(1/4)}*\text{Sqrt}[a + I*b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a + b*\text{Tan}[c + d*x]])/\text{Sqrt}[a + I*b]))/d$

3.470.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.97, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.184$, Rules used = {2011, 3042, 4058, 613, 104, 218, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} dx \\
 & \quad \downarrow \text{4058} \\
 & B \int \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} d \tan(c+dx) \\
 & \quad \downarrow \text{613} \\
 & \frac{B \left(\frac{1}{2} \int \frac{1}{\sqrt{\tan(c+dx)}(\tan(c+dx)+i)\sqrt{a+b \tan(c+dx)}} d \tan(c+dx) - \frac{1}{2} \int \frac{1}{(i-\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} d \tan(c+dx) \right)}{d} \\
 & \quad \downarrow \text{104} \\
 & \frac{B \left(\int \frac{1}{\frac{(a-ib) \tan(c+dx)}{a+b \tan(c+dx)} + i} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} - \int \frac{1}{i - \frac{(a+ib) \tan(c+dx)}{a+b \tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right)}{d} \\
 & \quad \downarrow \text{218}
 \end{aligned}$$

3.470. $\int \frac{\sqrt{\tan(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

$$\begin{array}{c}
 B \left(\int \frac{1}{\frac{(a-ib)\tan(c+dx)}{a+b\tan(c+dx)} + i} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}} + \frac{i \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{-b+ia}} \right) \\
 \hline
 d \\
 \downarrow \text{221} \\
 B \left(\frac{i \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{-b+ia}} - \frac{i \operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{b+ia}} \right) \\
 \hline
 d
 \end{array}$$

input `Int[(Sqrt[Tan[c + d*x]]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `(B*((I*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[I*a - b] - (I*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[I*a + b]))/d`

3.470.3.1 Defintions of rubi rules used

rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 613 `Int[Sqrt[(e_)*(x_)]/(Sqrt[(c_) + (d_)*(x_)]*((a_) + (b_)*(x_)^2)), x_Symbol] := Simp[e/(2*b) Int[1/(Sqrt[e*x]*Sqrt[c + d*x]*(Rt[-a/b, 2] + x)), x], x] - Simp[e/(2*b) Int[1/(Sqrt[e*x]*Sqrt[c + d*x]*(Rt[-a/b, 2] - x)), x], x] /; FreeQ[{a, b, c, d, e}, x]`

```
rule 2011 Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] :=
  Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x,
  a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4058 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)])^(n_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*((c + d*ff*x)^n/(1 + ff^2*x^2)), x], x,
Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

3.470.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.57 (sec) , antiderivative size = 940499, normalized size of antiderivative = 8038.45

output too large to display

```
input int(tan(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x)
```

```
output result too large to display
```

3.470.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4509 vs. 2(89) = 178.

Time = 0.75 (sec) , antiderivative size = 4509, normalized size of antiderivative = 38.54

$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

```
input integrate(tan(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,
algorithm="fricas")
```

3.470. $\int \frac{\sqrt{\tan(c+dx)}(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$

output

```
-1/8*sqrt(((a^2 + b^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^2 -
B^2*b)/((a^2 + b^2)*d^2))*log((2*(2*B^3*a^4*b + 4*B^3*a^2*b^3 + (B^3*a^5 +
3*B^3*a^3*b^2 + 4*B^3*a*b^4)*tan(d*x + c) + (2*(B*a^5*b + 3*B*a^3*b^3 + 2
*B*a*b^5)*d^2*tan(d*x + c) - (B*a^6 + 4*B*a^4*b^2 + 7*B*a^2*b^4 + 4*B*b^6)
*d^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)))*sqrt(b*tan(d*x + c) +
a)*sqrt(tan(d*x + c)) + ((B^2*a^6 + 7*B^2*a^4*b^2 + 12*B^2*a^2*b^4)*d*tan(
d*x + c)^2 + 2*(B^2*a^5*b + B^2*a^3*b^3 - 4*B^2*a*b^5)*d*tan(d*x + c) - (B
^2*a^6 + 3*B^2*a^4*b^2 + 4*B^2*a^2*b^4)*d - 2*((a^4*b^3 + 5*a^2*b^5 + 4*b^
7)*d^3*tan(d*x + c)^2 + (a^7 + 6*a^5*b^2 + 13*a^3*b^4 + 8*a*b^6)*d^3*tan(d
*x + c) + (a^6*b + 3*a^4*b^3 + 2*a^2*b^5)*d^3)*sqrt(-B^4*a^2/((a^4 + 2*a^2
*b^2 + b^4)*d^4)))*sqrt(((a^2 + b^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4
)*d^4))*d^2 - B^2*b)/((a^2 + b^2)*d^2)))/(tan(d*x + c)^2 + 1)) - 1/8*sqrt(
((a^2 + b^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^2 - B^2*b)/((a
^2 + b^2)*d^2))*log(-2*(2*B^3*a^4*b + 4*B^3*a^2*b^3 + (B^3*a^5 + 3*B^3*a^
3*b^2 + 4*B^3*a*b^4)*tan(d*x + c) + (2*(B*a^5*b + 3*B*a^3*b^3 + 2*B*a*b^5)
*d^2*tan(d*x + c) - (B*a^6 + 4*B*a^4*b^2 + 7*B*a^2*b^4 + 4*B*b^6)*d^2)*sqr
t(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)))*sqrt(b*tan(d*x + c) + a)*sqrt(t
an(d*x + c)) + ((B^2*a^6 + 7*B^2*a^4*b^2 + 12*B^2*a^2*b^4)*d*tan(d*x + c)^
2 + 2*(B^2*a^5*b + B^2*a^3*b^3 - 4*B^2*a*b^5)*d*tan(d*x + c) - (B^2*a^6 +
3*B^2*a^4*b^2 + 4*B^2*a^2*b^4)*d - 2*((a^4*b^3 + 5*a^2*b^5 + 4*b^7)*d^3...
```

3.470.6 Sympy [F]

$$\int \frac{\sqrt{\tan(c+dx)}(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = B \int \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}} dx$$

input

```
integrate(tan(d*x+c)**(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x
)
```

output

```
B*Integral(sqrt(tan(c + d*x))/sqrt(a + b*tan(c + d*x)), x)
```

3.470.7 Maxima [F]

$$\int \frac{\sqrt{\tan(c+dx)}(aB + bB \tan(c+dx))}{(a + b \tan(c+dx))^{3/2}} dx = \int \frac{(Bb \tan(dx+c) + Ba)\sqrt{\tan(dx+c)}}{(b \tan(dx+c) + a)^{3/2}} dx$$

input `integrate(tan(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,
algorithm="maxima")`

output `integrate((B*b*tan(d*x + c) + B*a)*sqrt(tan(d*x + c))/(b*tan(d*x + c) + a)
^(3/2), x)`

3.470.8 Giac [F(-1)]

Timed out.

$$\int \frac{\sqrt{\tan(c+dx)}(aB + bB \tan(c+dx))}{(a + b \tan(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,
algorithm="giac")`

output `Timed out`

3.470.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\tan(c+dx)}(aB + bB \tan(c+dx))}{(a + b \tan(c+dx))^{3/2}} dx = \int \frac{\sqrt{\tan(c+dx)}(Ba + Bb \tan(c+dx))}{(a + b \tan(c+dx))^{3/2}} dx$$

input `int((tan(c + d*x)^(1/2)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/
2),x)`

output `int((tan(c + d*x)^(1/2)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/
2), x)`

3.471
$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx$$

3.471.1 Optimal result 4511
 3.471.2 Mathematica [A] (verified) 4511
 3.471.3 Rubi [A] (verified) 4512
 3.471.4 Maple [B] (warning: unable to verify) 4513
 3.471.5 Fricas [B] (verification not implemented) 4514
 3.471.6 Sympy [F] 4514
 3.471.7 Maxima [F] 4515
 3.471.8 Giac [F(-1)] 4515
 3.471.9 Mupad [F(-1)] 4515

3.471.1 Optimal result

Integrand size = 38, antiderivative size = 111

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \frac{B \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia-bd}} + \frac{B \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia+bd}}$$

output `B*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d/(I*a-b)^(1/2)+B*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d/(I*a+b)^(1/2)`

3.471.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.13

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \frac{(-1)^{3/4} B \left(-\frac{\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} - \frac{\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{a+ib}} \right)}{d}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)),x]`

3.471.
$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx$$

output $((-1)^{3/4} * B * (-\text{ArcTan}[((-1)^{1/4} * \text{Sqrt}[-a + I * b] * \text{Sqrt}[\text{Tan}[c + d * x]]] / \text{Sqrt}[a + b * \text{Tan}[c + d * x]]] / \text{Sqrt}[-a + I * b]) - \text{ArcTan}[((-1)^{1/4} * \text{Sqrt}[a + I * b] * \text{Sqrt}[\text{Tan}[c + d * x]]] / \text{Sqrt}[a + b * \text{Tan}[c + d * x]]] / \text{Sqrt}[a + I * b]) / d$

3.471.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.97, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.132$, Rules used = {2011, 3042, 4058, 615, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{1}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4058} \\
 & \frac{B \int \frac{1}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)} (\tan^2(c + dx) + 1)} d \tan(c + dx)}{d} \\
 & \quad \downarrow \text{615} \\
 & \frac{B \int \left(\frac{i}{2(i - \tan(c + dx)) \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} + \frac{i}{2 \sqrt{\tan(c + dx)} (\tan(c + dx) + i) \sqrt{a + b \tan(c + dx)}} \right) d \tan(c + dx)}{d} \\
 & \quad \downarrow \text{2009} \\
 & \frac{B \left(\frac{\arctan\left(\frac{\sqrt{-b + ia} \sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{\sqrt{-b + ia}} + \frac{\text{arctanh}\left(\frac{\sqrt{b + ia} \sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{\sqrt{b + ia}} \right)}{d}
 \end{aligned}$$

input $\text{Int}[(a * B + b * B * \text{Tan}[c + d * x]) / (\text{Sqrt}[\text{Tan}[c + d * x]] * (a + b * \text{Tan}[c + d * x])^{3/2}), x]$

3.471. $\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx$

```
output (B*(ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[I*a - b] + ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[I*a + b]))/d
```

3.471.3.1 Defintions of rubi rules used

```
rule 615 Int[((e_.)*(x_)^(m_.)*((c_) + (d_.)*(x_)^(n_.))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && ILtQ[p, 0]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2011 Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4058 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*((c + d*ff*x)^n/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

3.471.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.54 (sec) , antiderivative size = 940264, normalized size of antiderivative = 8470.85

output too large to display

```
input int((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x)
```

```
output result too large to display
```

$$3.471. \quad \int \frac{aB+bB \tan(c+dx)}{\sqrt{\tan(c+dx)(a+b \tan(c+dx))^{3/2}} dx$$

3.471.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4469 vs. $2(87) = 174$.

Time = 0.77 (sec) , antiderivative size = 4469, normalized size of antiderivative = 40.26

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

```
input integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x,
algorithm="fricas")
```

```
output 1/8*sqrt(-((a^2 + b^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^2 -
B^2*b)/((a^2 + b^2)*d^2))*log(1/2*(2*(2*B^3*a^4*b + 4*B^3*a^2*b^3 + (B^3*a^5 + 3*B^3*a^3*b^2 + 4*B^3*a*b^4)*tan(d*x + c) + (2*(B*a^5*b + 3*B*a^3*b^3 + 2*B*a*b^5)*d^2*tan(d*x + c) - (B*a^6 + 4*B*a^4*b^2 + 7*B*a^2*b^4 + 4*B*b^6)*d^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)))*sqrt(b*tan(d*x + c) + a)*sqrt(tan(d*x + c)) + (2*(B^2*a^3*b^3 + 4*B^2*a*b^5)*d*tan(d*x + c)^2 + 2*(B^2*a^6 + 5*B^2*a^4*b^2 + 8*B^2*a^2*b^4)*d*tan(d*x + c) + 2*(B^2*a^5*b + 2*B^2*a^3*b^3)*d + ((a^7 + 8*a^5*b^2 + 19*a^3*b^4 + 12*a*b^6)*d^3*tan(d*x + c)^2 + 2*(a^6*b + 2*a^4*b^3 - 3*a^2*b^5 - 4*b^7)*d^3*tan(d*x + c) - (a^7 + 4*a^5*b^2 + 7*a^3*b^4 + 4*a*b^6)*d^3)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)))*sqrt(-((a^2 + b^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^2 - B^2*b)/((a^2 + b^2)*d^2)))/(tan(d*x + c)^2 + 1)) + 1/8*sqrt(-((a^2 + b^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^2 - B^2*b)/((a^2 + b^2)*d^2))*log(-1/2*(2*(2*B^3*a^4*b + 4*B^3*a^2*b^3 + (B^3*a^5 + 3*B^3*a^3*b^2 + 4*B^3*a*b^4)*tan(d*x + c) + (2*(B*a^5*b + 3*B*a^3*b^3 + 2*B*a*b^5)*d^2*tan(d*x + c) - (B*a^6 + 4*B*a^4*b^2 + 7*B*a^2*b^4 + 4*B*b^6)*d^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)))*sqrt(b*tan(d*x + c) + a)*sqrt(tan(d*x + c)) + (2*(B^2*a^3*b^3 + 4*B^2*a*b^5)*d*tan(d*x + c)^2 + 2*(B^2*a^6 + 5*B^2*a^4*b^2 + 8*B^2*a^2*b^4)*d*tan(d*x + c) + 2*(B^2*a^5*b + 2*B^2*a^3*b^3)*d + ((a^7 + 8*a^5*b^2 + 19*a^3*b^4 + 12*a*b^6)*d^3*tan(d*...
```

3.471.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = B \int \frac{1}{\sqrt{a + b \tan(c + dx)} \sqrt{\tan(c + dx)}} dx$$

```
input integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)**(1/2)/(a+b*tan(d*x+c))**(3/2),x
)
```

3.471. $\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx$

output `B*Integral(1/(sqrt(a + b*tan(c + d*x))*sqrt(tan(c + d*x))), x)`

3.471.7 Maxima [F]

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \int \frac{Bb \tan(dx + c) + Ba}{(b \tan(dx + c) + a)^{\frac{3}{2}} \sqrt{\tan(dx + c)}} dx$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x,
algorithm="maxima")`

output `integrate((B*b*tan(d*x + c) + B*a)/((b*tan(d*x + c) + a)^(3/2)*sqrt(tan(d*
x + c))), x)`

3.471.8 Giac [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x,
algorithm="giac")`

output `Timed out`

3.471.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \int \frac{Ba + Bb \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx$$

input `int((B*a + B*b*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(3/2
)),x)`

output `int((B*a + B*b*tan(c + d*x))/(tan(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(3/2
)), x)`

3.471. $\int \frac{aB + bB \tan(c + dx)}{\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{3/2}} dx$

3.472
$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx$$

3.472.1 Optimal result 4516
 3.472.2 Mathematica [A] (verified) 4516
 3.472.3 Rubi [A] (verified) 4517
 3.472.4 Maple [B] (warning: unable to verify) 4520
 3.472.5 Fricas [B] (verification not implemented) 4520
 3.472.6 Sympy [F] 4521
 3.472.7 Maxima [F(-1)] 4522
 3.472.8 Giac [F(-1)] 4522
 3.472.9 Mupad [F(-1)] 4522

3.472.1 Optimal result

Integrand size = 38, antiderivative size = 150

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = -\frac{iB \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia-bd}} + \frac{iB \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{ia+bd}} - \frac{2B\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}$$

output `-I*B*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d/(I*a-b)^(1/2)+I*B*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))/d/(I*a+b)^(1/2)-2*B*(a+b*tan(d*x+c))^(1/2)/a/d/tan(d*x+c)^(1/2)`

3.472.2 Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.05

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \frac{B \left(\frac{\sqrt[4]{-1} \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} - \frac{\sqrt[4]{-1} \arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{a+ib}} \right)}{d}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2)),x]`

3.472.
$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx$$

output $(B*((-1)^{(1/4)}*\text{ArcTan}[((-1)^{(1/4)}*\text{Sqrt}[-a + I*b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a + b*\text{Tan}[c + d*x]])/\text{Sqrt}[-a + I*b] - ((-1)^{(1/4)}*\text{ArcTan}[((-1)^{(1/4)}*\text{Sqrt}[a + I*b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a + b*\text{Tan}[c + d*x]])/\text{Sqrt}[a + I*b] - (2*\text{Sqrt}[a + b*\text{Tan}[c + d*x]])/(a*\text{Sqrt}[\text{Tan}[c + d*x]]))/d$

3.472.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.99, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2011, 3042, 4052, 27, 3042, 4058, 613, 104, 218, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx$$

↓ 2011

$$B \int \frac{1}{\tan^{\frac{3}{2}}(c + dx)\sqrt{a + b \tan(c + dx)}} dx$$

↓ 3042

$$B \int \frac{1}{\tan(c + dx)^{3/2}\sqrt{a + b \tan(c + dx)}} dx$$

↓ 4052

$$B \left(-\frac{2 \int \frac{a\sqrt{\tan(c+dx)}}{2\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2\sqrt{a + b \tan(c + dx)}}{ad\sqrt{\tan(c + dx)}} \right)$$

↓ 27

$$B \left(-\int \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}} dx - \frac{2\sqrt{a + b \tan(c + dx)}}{ad\sqrt{\tan(c + dx)}} \right)$$

↓ 3042

$$B \left(-\int \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}} dx - \frac{2\sqrt{a + b \tan(c + dx)}}{ad\sqrt{\tan(c + dx)}} \right)$$

↓ 4058

$$B \left(-\frac{\int \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} d \tan(c+dx)}{d} - \frac{2\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right)$$

↓ 613

$$B \left(-\frac{2\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} - \frac{\frac{1}{2} \int \frac{1}{\sqrt{\tan(c+dx)}(\tan(c+dx)+i)\sqrt{a+b \tan(c+dx)}} d \tan(c+dx) - \frac{1}{2} \int \frac{1}{(i-\tan(c+dx))\sqrt{\tan(c+dx)}} d \tan(c+dx)}{d} \right)$$

↓ 104

$$B \left(-\frac{2\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} - \frac{\int \frac{1}{\frac{(a-ib)\tan(c+dx)}{a+b \tan(c+dx)}+i} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} - \int \frac{1}{i-\frac{(a+ib)\tan(c+dx)}{a+b \tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} \right)$$

↓ 218

$$B \left(-\frac{2\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} - \frac{\int \frac{1}{\frac{(a-ib)\tan(c+dx)}{a+b \tan(c+dx)}+i} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} + \frac{i \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-b+ia}}}{d} \right)$$

↓ 221

$$B \left(-\frac{2\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} - \frac{\frac{i \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-b+ia}} - \frac{i \operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{b+ia}}}{d} \right)$$

input `Int[(a*B + b*B*Tan[c + d*x])/(Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2)),x]`

output `B*(-(((I*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[I*a - b] - (I*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[I*a + b])/d - (2*Sqrt[a + b*Tan[c + d*x]])/(a*d*Sqrt[Tan[c + d*x]]))`

3.472.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 613 `Int[Sqrt[(e_)*(x_)]/(Sqrt[(c_) + (d_)*(x_)]*((a_) + (b_)*(x_)^2)), x_Symbol] := Simp[e/(2*b) Int[1/(Sqrt[e*x]*Sqrt[c + d*x]*(Rt[-a/b, 2] + x)), x], x] - Simp[e/(2*b) Int[1/(Sqrt[e*x]*Sqrt[c + d*x]*(Rt[-a/b, 2] - x)), x], x] /; FreeQ[{a, b, c, d, e}, x]`
- rule 2011 `Int[(u_)*((a_) + (b_)*(v_)^(m_))*((c_) + (d_)*(v_)^(n_)), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4052 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_), x_Symbol] := Simp[b^2*(a + b*Tan[e + f*x])^(m + 1)*((c
+ d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(a^2 + b^2)*(b*c - a*d))), x] + Simp[1
/((m + 1)*(a^2 + b^2)*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c +
d*Tan[e + f*x])^n*Simp[a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2) - b*(b*c -
a*d)*(m + 1)*Tan[e + f*x] - b^2*d*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /
; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0]
&& NeQ[c^2 + d^2, 0] && IntegerQ[2*m] && LtQ[m, -1] && (LtQ[n, 0] || Integ
erQ[m]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4058 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*((c + d*ff*x)^n/(1 + ff^2*x^2)), x], x,
Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

3.472.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.58 (sec) , antiderivative size = 943929, normalized size of antiderivative = 6292.86

output too large to display

```
input int((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x)
```

```
output result too large to display
```

3.472.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4608 vs. $2(118) = 236$.

Time = 0.76 (sec) , antiderivative size = 4608, normalized size of antiderivative = 30.72

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

```
input integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x,
algorithm="fricas")
```

3.472. $\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx$

output `1/8*(a*d*sqrt(((a^2 + b^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^2 - B^2*b)/((a^2 + b^2)*d^2))*log((2*(2*B^3*a^4*b + 4*B^3*a^2*b^3 + (B^3*a^5 + 3*B^3*a^3*b^2 + 4*B^3*a*b^4)*tan(d*x + c) + (2*(B*a^5*b + 3*B*a^3*b^3 + 2*B*a*b^5)*d^2*tan(d*x + c) - (B*a^6 + 4*B*a^4*b^2 + 7*B*a^2*b^4 + 4*B*b^6)*d^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)))*sqrt(b*tan(d*x + c) + a)*sqrt(tan(d*x + c)) + ((B^2*a^6 + 7*B^2*a^4*b^2 + 12*B^2*a^2*b^4)*d*tan(d*x + c)^2 + 2*(B^2*a^5*b + B^2*a^3*b^3 - 4*B^2*a*b^5)*d*tan(d*x + c) - (B^2*a^6 + 3*B^2*a^4*b^2 + 4*B^2*a^2*b^4)*d - 2*((a^4*b^3 + 5*a^2*b^5 + 4*b^7)*d^3*tan(d*x + c)^2 + (a^7 + 6*a^5*b^2 + 13*a^3*b^4 + 8*a*b^6)*d^3*tan(d*x + c) + (a^6*b + 3*a^4*b^3 + 2*a^2*b^5)*d^3)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)))*sqrt(((a^2 + b^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^2 - B^2*b)/((a^2 + b^2)*d^2)))/(tan(d*x + c)^2 + 1))*tan(d*x + c) + a*d*sqrt(((a^2 + b^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^2 - B^2*b)/((a^2 + b^2)*d^2))*log(-2*(2*B^3*a^4*b + 4*B^3*a^2*b^3 + (B^3*a^5 + 3*B^3*a^3*b^2 + 4*B^3*a*b^4)*tan(d*x + c) + (2*(B*a^5*b + 3*B*a^3*b^3 + 2*B*a*b^5)*d^2*tan(d*x + c) - (B*a^6 + 4*B*a^4*b^2 + 7*B*a^2*b^4 + 4*B*b^6)*d^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)))*sqrt(b*tan(d*x + c) + a)*sqrt(tan(d*x + c)) + ((B^2*a^6 + 7*B^2*a^4*b^2 + 12*B^2*a^2*b^4)*d*tan(d*x + c)^2 + 2*(B^2*a^5*b + B^2*a^3*b^3 - 4*B^2*a*b^5)*d*tan(d*x + c) - (B^2*a^6 + 3*B^2*a^4*b^2 + 4*B^2*a^2*b^4)*d - 2*((a^4*b^3 + 5*a^2...`

3.472.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{\frac{3}{2}}} dx = B \int \frac{1}{\sqrt{a + b \tan(c + dx)} \tan^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)**(3/2)/(a+b*tan(d*x+c))**(3/2), x)`

output `B*Integral(1/(sqrt(a + b*tan(c + d*x))*tan(c + d*x)**(3/2)), x)`

3.472.7 Maxima [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

```
input integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x,
algorithm="maxima")
```

output Timed out

3.472.8 Giac [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

```
input integrate((B*a+b*B*tan(d*x+c))/tan(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x,
algorithm="giac")
```

output Timed out

3.472.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \int \frac{B a + B b \tan(c + dx)}{\tan(c + dx)^{3/2} (a + b \tan(c + dx))^{3/2}} dx$$

```
input int((B*a + B*b*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(3/2)),x)
```

```
output int((B*a + B*b*tan(c + d*x))/(tan(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(3/2)), x)
```

3.473 $\int (a + b \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx$

3.473.1 Optimal result	4523
3.473.2 Mathematica [A] (verified)	4524
3.473.3 Rubi [A] (warning: unable to verify)	4524
3.473.4 Maple [C] (verified)	4528
3.473.5 Fricas [B] (verification not implemented)	4529
3.473.6 Sympy [F]	4529
3.473.7 Maxima [F]	4530
3.473.8 Giac [F]	4530
3.473.9 Mupad [B] (verification not implemented)	4530

3.473.1 Optimal result

Integrand size = 25, antiderivative size = 379

$$\int (a + b \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx = -\frac{1}{4}(a - ib)^{2/3}(A - iB)x + \frac{\sqrt{3}(a - ib)^{2/3}(iA + B) \arctan\left(\frac{1 + \sqrt[3]{a + b \tan(c + dx)}}{\sqrt[3]{a - ib}}\right) - \frac{1}{4}(a + ib)^{2/3}(A + iB)x - \frac{\sqrt{3}(a + ib)^{2/3}(iA - B) \arctan\left(\frac{1 + \sqrt[3]{a + b \tan(c + dx)}}{\sqrt[3]{a - ib}}\right)}{2d}$$

output

```
-1/4*(a-I*b)^(2/3)*(A-I*B)*x-1/4*(a+I*b)^(2/3)*(A+I*B)*x-1/4*(a+I*b)^(2/3)
*(I*A-B)*ln(cos(d*x+c))/d+1/4*(a-I*b)^(2/3)*(I*A+B)*ln(cos(d*x+c))/d+3/4*(
a-I*b)^(2/3)*(I*A+B)*ln((a-I*b)^(1/3)-(a+b*tan(d*x+c))^(1/3))/d-3/4*(a+I*b)
)^(2/3)*(I*A-B)*ln((a+I*b)^(1/3)-(a+b*tan(d*x+c))^(1/3))/d+1/2*(a-I*b)^(2/
3)*(I*A+B)*arctan(1/3*(1+2*(a+b*tan(d*x+c))^(1/3)/(a-I*b)^(1/3))*3^(1/2))*
3^(1/2)/d-1/2*(a+I*b)^(2/3)*(I*A-B)*arctan(1/3*(1+2*(a+b*tan(d*x+c))^(1/3)
/(a+I*b)^(1/3))*3^(1/2))*3^(1/2)/d+3/2*B*(a+b*tan(d*x+c))^(2/3)/d
```

3.473.2 Mathematica [A] (verified)

Time = 1.07 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.69

$$\int (a + b \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx = \frac{i \left((A - iB) \left((a - ib)^{2/3} \left(2\sqrt{3} \arctan \left(\frac{1 + \sqrt[3]{a + b \tan(c + dx)}}{\sqrt[3]{a - ib}} \right) \right) - \log(i + \tan(c + dx)) \right) \right)}{d}$$

input `Integrate[(a + b*Tan[c + d*x])^(2/3)*(A + B*Tan[c + d*x]),x]`

output `((I/4)*((A - I*B)*((a - I*b)^(2/3)*(2*sqrt[3]*ArcTan[(1 + (2*(a + b*Tan[c + d*x])^(1/3)))/(a - I*b)^(1/3)]/sqrt[3]] - Log[I + Tan[c + d*x]] + 3*Log[(a - I*b)^(1/3) - (a + b*Tan[c + d*x])^(1/3)]) + 3*(a + b*Tan[c + d*x])^(2/3)) - (A + I*B)*((a + I*b)^(2/3)*(2*sqrt[3]*ArcTan[(1 + (2*(a + b*Tan[c + d*x])^(1/3)))/(a + I*b)^(1/3)]/sqrt[3]] - Log[I - Tan[c + d*x]] + 3*Log[(a + I*b)^(1/3) - (a + b*Tan[c + d*x])^(1/3)]) + 3*(a + b*Tan[c + d*x])^(2/3)))/d`

3.473.3 Rubi [A] (warning: unable to verify)Time = 0.70 (sec) , antiderivative size = 269, normalized size of antiderivative = 0.71, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.440$, Rules used = {3042, 4011, 3042, 4022, 3042, 4020, 25, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + b \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + b \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4011} \\ & \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx + \frac{3B(a + b \tan(c + dx))^{2/3}}{2d} \end{aligned}$$

$$\begin{aligned}
 & \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx + \frac{3B(a + b \tan(c + dx))^{2/3}}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2}(a + ib)(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx + \frac{1}{2}(a - ib)(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt[3]{a + b \tan(c + dx)}} dx + \\
 & \quad \frac{3B(a + b \tan(c + dx))^{2/3}}{2d} \\
 & \quad \downarrow \text{4022} \\
 & \frac{1}{2}(a + ib)(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx + \frac{1}{2}(a - ib)(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt[3]{a + b \tan(c + dx)}} dx + \\
 & \quad \frac{3B(a + b \tan(c + dx))^{2/3}}{2d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{i(a - ib)(A - iB) \int -\frac{1}{(1 - i \tan(c + dx)) \sqrt[3]{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} - \\
 & \frac{i(a + ib)(A + iB) \int -\frac{1}{(i \tan(c + dx) + 1) \sqrt[3]{a + b \tan(c + dx)}} d(-i \tan(c + dx))}{2d} + \\
 & \quad \frac{3B(a + b \tan(c + dx))^{2/3}}{2d} \\
 & \quad \downarrow \text{4020} \\
 & \frac{i(a - ib)(A - iB) \int \frac{1}{(1 - i \tan(c + dx)) \sqrt[3]{a + b \tan(c + dx)}} d(i \tan(c + dx))}{2d} + \\
 & \frac{i(a + ib)(A + iB) \int \frac{1}{(i \tan(c + dx) + 1) \sqrt[3]{a + b \tan(c + dx)}} d(-i \tan(c + dx))}{2d} + \frac{3B(a + b \tan(c + dx))^{2/3}}{2d} \\
 & \quad \downarrow \text{25} \\
 & \frac{i(a - ib)(A - iB) \left(\frac{3}{2} \int \frac{1}{-\tan^2(c + dx) + (a - ib)^{2/3} + \sqrt[3]{a - ib} \sqrt[3]{a + b \tan(c + dx)}} d \sqrt[3]{a + b \tan(c + dx)} - \frac{3 \int \frac{1}{\sqrt[3]{a - ib} - i \tan(c + dx)}}{2d} \right)}{2d} \\
 & \frac{i(a + ib)(A + iB) \left(\frac{3}{2} \int \frac{1}{-\tan^2(c + dx) + (a + ib)^{2/3} + \sqrt[3]{a + ib} \sqrt[3]{a + b \tan(c + dx)}} d \sqrt[3]{a + b \tan(c + dx)} - \frac{3 \int \frac{1}{i \tan(c + dx) + \sqrt[3]{a + b \tan(c + dx)}}}{2d} \right)}{2d} \\
 & \quad \downarrow \text{67} \\
 & \frac{3B(a + b \tan(c + dx))^{2/3}}{2d}
 \end{aligned}$$

$$\begin{array}{c}
\downarrow 16 \\
\frac{i(a-ib)(A-ib) \left(\frac{3}{2} \int \frac{1}{-\tan^2(c+dx) + (a-ib)^{2/3} + \sqrt[3]{a-ib} \sqrt[3]{a+b \tan(c+dx)}} d\sqrt[3]{a+b \tan(c+dx)} - \frac{\log(1-i \tan(c+dx))}{2\sqrt[3]{a-ib}} \right)}{2d} \\
\frac{i(a+ib)(A+ib) \left(\frac{3}{2} \int \frac{1}{-\tan^2(c+dx) + (a+ib)^{2/3} + \sqrt[3]{a+ib} \sqrt[3]{a+b \tan(c+dx)}} d\sqrt[3]{a+b \tan(c+dx)} - \frac{\log(1+i \tan(c+dx))}{2\sqrt[3]{a+ib}} \right)}{2d} \\
\frac{3B(a+b \tan(c+dx))^{2/3}}{2d} \\
\downarrow 1082 \\
\frac{i(a-ib)(A-ib) \left(-\frac{3 \int \frac{1}{\tan^2(c+dx)-3} d \left(\frac{2i \tan(c+dx)}{\sqrt[3]{a-ib}} + 1 \right)}{\sqrt[3]{a-ib}} - \frac{\log(1-i \tan(c+dx))}{2\sqrt[3]{a-ib}} + \frac{3 \log(\sqrt[3]{a-ib} - i \tan(c+dx))}{2\sqrt[3]{a-ib}} \right)}{2d} \\
\frac{i(a+ib)(A+ib) \left(-\frac{3 \int \frac{1}{\tan^2(c+dx)-3} d \left(1 - \frac{2i \tan(c+dx)}{\sqrt[3]{a+ib}} \right)}{\sqrt[3]{a+ib}} - \frac{\log(1+i \tan(c+dx))}{2\sqrt[3]{a+ib}} + \frac{3 \log(\sqrt[3]{a+ib} + i \tan(c+dx))}{2\sqrt[3]{a+ib}} \right)}{2d} \\
+ \\
\frac{3B(a+b \tan(c+dx))^{2/3}}{2d} \\
\downarrow 217 \\
\frac{i(a-ib)(A-ib) \left(\frac{i\sqrt{3} \arctanh\left(\frac{\tan(c+dx)}{\sqrt{3}}\right)}{\sqrt[3]{a-ib}} - \frac{\log(1-i \tan(c+dx))}{2\sqrt[3]{a-ib}} + \frac{3 \log(\sqrt[3]{a-ib} - i \tan(c+dx))}{2\sqrt[3]{a-ib}} \right)}{2d} \\
\frac{i(a+ib)(A+ib) \left(-\frac{i\sqrt{3} \arctanh\left(\frac{\tan(c+dx)}{\sqrt{3}}\right)}{\sqrt[3]{a+ib}} - \frac{\log(1+i \tan(c+dx))}{2\sqrt[3]{a+ib}} + \frac{3 \log(\sqrt[3]{a+ib} + i \tan(c+dx))}{2\sqrt[3]{a+ib}} \right)}{2d} \\
+ \\
\frac{3B(a+b \tan(c+dx))^{2/3}}{2d}
\end{array}$$

input `Int[(a + b*Tan[c + d*x])^(2/3)*(A + B*Tan[c + d*x]),x]`

```
output ((I/2)*(a - I*b)*(A - I*B)*((I*Sqrt[3]*ArcTanh[Tan[c + d*x]/Sqrt[3]])/(a -
I*b)^(1/3) - Log[1 - I*Tan[c + d*x]]/(2*(a - I*b)^(1/3)) + (3*Log[(a - I*
b)^(1/3) - I*Tan[c + d*x]]/(2*(a - I*b)^(1/3))))/d - ((I/2)*(a + I*b)*(A
+ I*B)*((-I)*Sqrt[3]*ArcTanh[Tan[c + d*x]/Sqrt[3]])/(a + I*b)^(1/3) - Log
[1 + I*Tan[c + d*x]]/(2*(a + I*b)^(1/3)) + (3*Log[(a + I*b)^(1/3) + I*Tan[
c + d*x]]/(2*(a + I*b)^(1/3))))/d + (3*B*(a + b*Tan[c + d*x])^(2/3))/(2*d
)
```

3.473.3.1 Defintions of rubi rules used

```
rule 16 Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a +
b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 67 Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(1/3)), x_Symbol] := With[
{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q), x
] + (Simp[3/(2*b) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)],
x] - Simp[3/(2*b*q) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x])] /
; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4011 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int
[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2,
0] && GtQ[m, 0]
```

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

3.473.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.56 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.26

method	result
derivativedivides	$\frac{\frac{3(a+b \tan(dx+c))^{\frac{2}{3}} B}{2} + \frac{\left(\frac{(Ab+Ba)R^4 + B(-a^2-b^2)R}{R^5 - R^2 a} \ln \left((a+b \tan(dx+c))^{\frac{1}{3}} - R \right) \right)}{\sum_{R=\text{RootOf}(_Z^6 - 2a_Z^3 + a^2 + b^2)}}}{d}$
default	$\frac{\frac{3(a+b \tan(dx+c))^{\frac{2}{3}} B}{2} + \frac{\left(\frac{(Ab+Ba)R^4 + B(-a^2-b^2)R}{R^5 - R^2 a} \ln \left((a+b \tan(dx+c))^{\frac{1}{3}} - R \right) \right)}{\sum_{R=\text{RootOf}(_Z^6 - 2a_Z^3 + a^2 + b^2)}}}{d}$
parts	$\frac{Ab \left(\frac{\sum_{R=\text{RootOf}(_Z^6 - 2a_Z^3 + a^2 + b^2)} \frac{-R^2 \ln \left((a+b \tan(dx+c))^{\frac{1}{3}} - R \right)}{-R^3 - a}}{2d} \right)}{2d} + B \left(\frac{3(a+b \tan(dx+c))^{\frac{2}{3}}}{2} + \frac{\sum_{R=\text{RootOf}(_Z^6 - 2a_Z^3 + a^2 + b^2)} \frac{-R^2 \ln \left((a+b \tan(dx+c))^{\frac{1}{3}} - R \right)}{-R^3 - a}}{2d} \right)$

3.473. $\int (a + b \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx$

input `int((a+b*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*(3/2*(a+b*tan(d*x+c))^(2/3)*B+1/2*sum(((A*b+B*a)*_R^4+B*(-a^2-b^2)*_R)/(_R^5-_R^2*a)*ln((a+b*tan(d*x+c))^(1/3)-_R),_R=RootOf(_Z^6-2*_Z^3*a+a^2+b^2)))`

3.473.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5405 vs. $2(281) = 562$.

Time = 1.89 (sec) , antiderivative size = 5405, normalized size of antiderivative = 14.26

$$\int (a + b \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.473.6 Sympy [F]

$$\int (a + b \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx)) (a + b \tan(c + dx))^{2/3} dx$$

input `integrate((a+b*tan(d*x+c))**(2/3)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(2/3), x)`

3.473.7 Maxima [F]

$$\int (a + b \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{2/3} dx$$

input `integrate((a+b*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(2/3), x)`

3.473.8 Giac [F]

$$\int (a + b \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{2/3} dx$$

input `integrate((a+b*tan(d*x+c))^(2/3)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `undef`

3.473.9 Mupad [B] (verification not implemented)

Time = 25.43 (sec) , antiderivative size = 3945, normalized size of antiderivative = 10.41

$$\int (a + b \tan(c + dx))^{2/3} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(2/3),x)`

output $\log\left(\left(\left(2*(-B^6*a^2*b^2*d^6)^{1/2} + B^3*a^2*d^3 - B^3*b^2*d^3\right)/d^6\right)^{2/3} * \left(\left(\left(2*(-B^6*a^2*b^2*d^6)^{1/2} + B^3*a^2*d^3 - B^3*b^2*d^3\right)/d^6\right)^{1/3} * (1944*a*b^4 * \left(\left(2*(-B^6*a^2*b^2*d^6)^{1/2} + B^3*a^2*d^3 - B^3*b^2*d^3\right)/d^6\right)^{2/3} * (a^2 + b^2) - (1944*B^2*b^4 * (a^2 + b^2)^2 * (a + b*\tan(c + d*x))^{1/3})/d^2\right)/2 + (972*B^3*a*b^4 * (3*b^4 - a^4 + 2*a^2*b^2)/d^3)/4 + (243*B^5*b^4 * (a^2 - b^2) * (a^2 + b^2)^2 * (a + b*\tan(c + d*x))^{1/3})/d^5 * \left(\left(\left(-4*B^6*a^2*b^2*d^6\right)^{1/2} + B^3*a^2*d^3 - B^3*b^2*d^3\right)/(8*d^6)\right)^{1/3} + \log\left(\left(\left(-2*(-B^6*a^2*b^2*d^6)^{1/2} - B^3*a^2*d^3 + B^3*b^2*d^3\right)/d^6\right)^{2/3} * \left(\left(\left(-2*(-B^6*a^2*b^2*d^6)^{1/2} - B^3*a^2*d^3 + B^3*b^2*d^3\right)/d^6\right)^{1/3} * (1944*a*b^4 * \left(\left(2*(-B^6*a^2*b^2*d^6)^{1/2} - B^3*a^2*d^3 + B^3*b^2*d^3\right)/d^6\right)^{2/3} * (a^2 + b^2) - (1944*B^2*b^4 * (a^2 + b^2)^2 * (a + b*\tan(c + d*x))^{1/3})/d^2\right)/2 + (972*B^3*a*b^4 * (3*b^4 - a^4 + 2*a^2*b^2)/d^3)/4 + (243*B^5*b^4 * (a^2 - b^2) * (a^2 + b^2)^2 * (a + b*\tan(c + d*x))^{1/3})/d^5 * \left(\left(\left(-4*B^6*a^2*b^2*d^6\right)^{1/2} - B^3*a^2*d^3 + B^3*b^2*d^3\right)/(8*d^6)\right)^{1/3} + \log\left(\left(\left(-A^6*d^6 * (a^2 - b^2)^2\right)^{1/2} - 2*A^3*a*b*d^3\right)/d^6\right)^{2/3} * \left(\left(\left(1944*a*b^4 * \left(\left(-A^6*d^6 * (a^2 - b^2)^2\right)^{1/2} - 2*A^3*a*b*d^3\right)/d^6\right)^{2/3} * (a^2 + b^2) + (1944*A^2*b^4 * (a^2 + b^2)^2 * (a + b*\tan(c + d*x))^{1/3})/d^2\right) * \left(\left(\left(-A^6*d^6 * (a^2 - b^2)^2\right)^{1/2} - 2*A^3*a*b*d^3\right)/d^6\right)^{1/3}\right)/2 + (972*A^3*b^5 * (3*a^4 - b^4 + 2*a^2*b^2)/d^3)/4 + (486*A^5*a*b^5 * (a^2 + b^2)^2 * (a + b*\tan(c + d*x))^{1/3})/d^5 * \left(\left(2*A^6*a^2*b^2*d^6 - A^6*b^4*d^6 - A^6*a^4*d^6\right)^{1/2}/(8*d^6) - (A^3...$

3.474 $\int \sqrt[3]{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx$

3.474.1 Optimal result	4532
3.474.2 Mathematica [A] (verified)	4533
3.474.3 Rubi [A] (warning: unable to verify)	4534
3.474.4 Maple [C] (verified)	4537
3.474.5 Fricas [B] (verification not implemented)	4538
3.474.6 Sympy [F]	4539
3.474.7 Maxima [F]	4540
3.474.8 Giac [F]	4540
3.474.9 Mupad [B] (verification not implemented)	4540

3.474.1 Optimal result

Integrand size = 25, antiderivative size = 377

$$\begin{aligned} & \int \sqrt[3]{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx \\ &= -\frac{1}{4}\sqrt[3]{a - ib}(A - iB)x - \frac{1}{4}\sqrt[3]{a + ib}(A + iB)x \\ &\quad - \frac{\sqrt{3}\sqrt[3]{a - ib}(iA + B) \arctan\left(\frac{1 + \sqrt[3]{a + b \tan(c + dx)}}{\sqrt[3]{a - ib}}\right)}{2d} \\ &\quad + \frac{\sqrt{3}\sqrt[3]{a + ib}(iA - B) \arctan\left(\frac{1 + \sqrt[3]{a + b \tan(c + dx)}}{\sqrt[3]{a + ib}}\right)}{2d} \\ &\quad - \frac{\sqrt[3]{a + ib}(iA - B) \log(\cos(c + dx))}{4d} + \frac{\sqrt[3]{a - ib}(iA + B) \log(\cos(c + dx))}{4d} \\ &\quad + \frac{3\sqrt[3]{a - ib}(iA + B) \log\left(\sqrt[3]{a - ib} - \sqrt[3]{a + b \tan(c + dx)}\right)}{4d} \\ &\quad - \frac{3\sqrt[3]{a + ib}(iA - B) \log\left(\sqrt[3]{a + ib} - \sqrt[3]{a + b \tan(c + dx)}\right)}{4d} + \frac{3B\sqrt[3]{a + b \tan(c + dx)}}{d} \end{aligned}$$

output
$$\begin{aligned} & -1/4*(a-I*b)^{(1/3)}*(A-I*B)*x-1/4*(a+I*b)^{(1/3)}*(A+I*B)*x-1/4*(a+I*b)^{(1/3)} \\ & *(I*A-B)*\ln(\cos(d*x+c))/d+1/4*(a-I*b)^{(1/3)}*(I*A+B)*\ln(\cos(d*x+c))/d+3/4*(\\ & a-I*b)^{(1/3)}*(I*A+B)*\ln((a-I*b)^{(1/3)}-(a+b*\tan(d*x+c))^{(1/3)})/d-3/4*(a+I*b \\ &)^{(1/3)}*(I*A-B)*\ln((a+I*b)^{(1/3)}-(a+b*\tan(d*x+c))^{(1/3)})/d-1/2*(a-I*b)^{(1/ \\ & 3)}*(I*A+B)*\arctan(1/3*(1+2*(a+b*\tan(d*x+c))^{(1/3)})/(a-I*b)^{(1/3}))*3^{(1/2)}* \\ & 3^{(1/2)}/d+1/2*(a+I*b)^{(1/3)}*(I*A-B)*\arctan(1/3*(1+2*(a+b*\tan(d*x+c))^{(1/3)} \\ & / (a+I*b)^{(1/3}))*3^{(1/2)})*3^{(1/2)}/d+3*B*(a+b*\tan(d*x+c))^{(1/3)}/d \end{aligned}$$

3.474.2 Mathematica [A] (verified)

Time = 0.99 (sec) , antiderivative size = 347, normalized size of antiderivative = 0.92

$$\int \sqrt[3]{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= i \left((A - iB) \left(-\frac{1}{2} \sqrt[3]{a - ib} \left(2\sqrt{3} \arctan \left(\frac{1 + \sqrt[3]{a + b \tan(c + dx)}}{\sqrt[3]{a - ib}} \right) \right) - 2 \log \left(\sqrt[3]{a - ib} - \sqrt[3]{a + b \tan(c + dx)} \right) \right) \right)$$

input `Integrate[(a + b*Tan[c + d*x])^(1/3)*(A + B*Tan[c + d*x]),x]`

output
$$\begin{aligned} & ((I/2)*((A - I*B)*(-1/2*((a - I*b)^{(1/3)}*(2*Sqrt[3]*ArcTan[(1 + (2*(a + b* \\ & Tan[c + d*x])^{(1/3)}))/(a - I*b)^{(1/3)})/Sqrt[3]] - 2*Log[(a - I*b)^{(1/3)} - (\\ & a + b*Tan[c + d*x])^{(1/3)}] + Log[(a - I*b)^{(2/3)} + (a - I*b)^{(1/3)}*(a + b* \\ & Tan[c + d*x])^{(1/3)} + (a + b*Tan[c + d*x])^{(2/3)}])) + 3*(a + b*Tan[c + d*x \\ &])^{(1/3)} - (A + I*B)*(-1/2*((a + I*b)^{(1/3)}*(2*Sqrt[3]*ArcTan[(1 + (2*(a \\ & + b*Tan[c + d*x])^{(1/3)}))/(a + I*b)^{(1/3)})/Sqrt[3]] - 2*Log[(a + I*b)^{(1/3)} \\ & - (a + b*Tan[c + d*x])^{(1/3)}] + Log[(a + I*b)^{(2/3)} + (a + I*b)^{(1/3)}*(a \\ & + b*Tan[c + d*x])^{(1/3)} + (a + b*Tan[c + d*x])^{(2/3)}])) + 3*(a + b*Tan[c + \\ & d*x])^{(1/3)}))/d \end{aligned}$$

3.474.3 Rubi [A] (warning: unable to verify)

Time = 0.73 (sec) , antiderivative size = 267, normalized size of antiderivative = 0.71, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.440$, Rules used = {3042, 4011, 3042, 4022, 3042, 4020, 25, 69, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt[3]{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt[3]{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4011} \\
 & \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx + \frac{3B \sqrt[3]{a + b \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{aA - bB + (Ab + aB) \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx + \frac{3B \sqrt[3]{a + b \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{4022} \\
 & \frac{1}{2}(a + ib)(A + iB) \int \frac{1 - i \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx + \frac{1}{2}(a - ib)(A - iB) \int \frac{i \tan(c + dx) + 1}{(a + b \tan(c + dx))^{2/3}} dx + \\
 & \quad \frac{3B \sqrt[3]{a + b \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2}(a + ib)(A + iB) \int \frac{1 - i \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx + \frac{1}{2}(a - ib)(A - iB) \int \frac{i \tan(c + dx) + 1}{(a + b \tan(c + dx))^{2/3}} dx + \\
 & \quad \frac{3B \sqrt[3]{a + b \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{4020} \\
 & \frac{i(a - ib)(A - iB) \int -\frac{1}{(1 - i \tan(c + dx))(a + b \tan(c + dx))^{2/3}} d(i \tan(c + dx))}{2d} - \\
 & \frac{i(a + ib)(A + iB) \int -\frac{1}{(i \tan(c + dx) + 1)(a + b \tan(c + dx))^{2/3}} d(-i \tan(c + dx))}{2d} + \frac{3B \sqrt[3]{a + b \tan(c + dx)}}{d} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{i(a-ib)(A-iB) \int \frac{1}{(1-i \tan(c+dx))(a+b \tan(c+dx))^{2/3}} d(i \tan(c+dx))}{2d} + \\
 & \frac{i(a+ib)(A+iB) \int \frac{1}{(i \tan(c+dx)+1)(a+b \tan(c+dx))^{2/3}} d(-i \tan(c+dx))}{2d} + \frac{3B \sqrt[3]{a+b \tan(c+dx)}}{d} \\
 & \qquad \qquad \qquad \downarrow 69 \\
 & \frac{i(a-ib)(A-iB) \left(-\frac{3 \int \frac{1}{-\tan^2(c+dx)+(a-ib)^{2/3}+\sqrt[3]{a-ib} \sqrt[3]{a+b \tan(c+dx)}} d \sqrt[3]{a+b \tan(c+dx)}}{2 \sqrt[3]{a-ib}} - \frac{3 \int \frac{1}{\sqrt[3]{a-ib}-i \tan(c+dx)}}{d} \right)}{2d} \\
 & \frac{i(a+ib)(A+iB) \left(-\frac{3 \int \frac{1}{-\tan^2(c+dx)+(a+ib)^{2/3}+\sqrt[3]{a+ib} \sqrt[3]{a+b \tan(c+dx)}} d \sqrt[3]{a+b \tan(c+dx)}}{2 \sqrt[3]{a+ib}} - \frac{3 \int \frac{1}{i \tan(c+dx)+\sqrt[3]{a+ib}}}{d} \right)}{2d} \\
 & \qquad \qquad \qquad \downarrow 16 \\
 & \frac{i(a-ib)(A-iB) \left(-\frac{3 \int \frac{1}{-\tan^2(c+dx)+(a-ib)^{2/3}+\sqrt[3]{a-ib} \sqrt[3]{a+b \tan(c+dx)}} d \sqrt[3]{a+b \tan(c+dx)}}{2 \sqrt[3]{a-ib}} - \frac{\log(1-i \tan(c+dx))}{2(a-ib)^{2/3}} \right)}{2d} \\
 & \frac{i(a+ib)(A+iB) \left(-\frac{3 \int \frac{1}{-\tan^2(c+dx)+(a+ib)^{2/3}+\sqrt[3]{a+ib} \sqrt[3]{a+b \tan(c+dx)}} d \sqrt[3]{a+b \tan(c+dx)}}{2 \sqrt[3]{a+ib}} - \frac{\log(1+i \tan(c+dx))}{2(a+ib)^{2/3}} \right)}{2d} \\
 & \qquad \qquad \qquad \downarrow 1082 \\
 & \frac{i(a-ib)(A-iB) \left(\frac{3 \int \frac{1}{\tan^2(c+dx)-3} d \left(\frac{2i \tan(c+dx)}{\sqrt[3]{a-ib}} + 1 \right)}{(a-ib)^{2/3}} - \frac{\log(1-i \tan(c+dx))}{2(a-ib)^{2/3}} + \frac{3 \log \left(\sqrt[3]{a-ib}-i \tan(c+dx) \right)}{2(a-ib)^{2/3}} \right)}{2d} \\
 & \frac{i(a+ib)(A+iB) \left(\frac{3 \int \frac{1}{\tan^2(c+dx)-3} d \left(1 - \frac{2i \tan(c+dx)}{\sqrt[3]{a+ib}} \right)}{(a+ib)^{2/3}} - \frac{\log(1+i \tan(c+dx))}{2(a+ib)^{2/3}} + \frac{3 \log \left(\sqrt[3]{a+ib}+i \tan(c+dx) \right)}{2(a+ib)^{2/3}} \right)}{2d} + \\
 & \frac{3B \sqrt[3]{a+b \tan(c+dx)}}{d}
 \end{aligned}$$

3.474. $\int \sqrt[3]{a+b \tan(c+dx)}(A+B \tan(c+dx)) dx$

$$\begin{array}{c}
 \downarrow 217 \\
 \frac{i(a-ib)(A-ib) \left(-\frac{i\sqrt{3}\operatorname{arctanh}\left(\frac{\tan(c+dx)}{\sqrt{3}}\right)}{(a-ib)^{2/3}} - \frac{\log(1-i\tan(c+dx))}{2(a-ib)^{2/3}} + \frac{3\log\left(\sqrt[3]{a-ib-i\tan(c+dx)}\right)}{2(a-ib)^{2/3}} \right)}{2d} - \\
 \frac{i(a+ib)(A+ib) \left(\frac{i\sqrt{3}\operatorname{arctanh}\left(\frac{\tan(c+dx)}{\sqrt{3}}\right)}{(a+ib)^{2/3}} - \frac{\log(1+i\tan(c+dx))}{2(a+ib)^{2/3}} + \frac{3\log\left(\sqrt[3]{a+ib+i\tan(c+dx)}\right)}{2(a+ib)^{2/3}} \right)}{2d} + \\
 \frac{3B\sqrt[3]{a+b\tan(c+dx)}}{d}
 \end{array}$$

input `Int[(a + b*Tan[c + d*x])^(1/3)*(A + B*Tan[c + d*x]),x]`

output `((I/2)*(a - I*b)*(A - I*B)*((-I)*Sqrt[3]*ArcTanh[Tan[c + d*x]/Sqrt[3]])/(a - I*b)^(2/3) - Log[1 - I*Tan[c + d*x]]/(2*(a - I*b)^(2/3)) + (3*Log[(a - I*b)^(1/3) - I*Tan[c + d*x]])/(2*(a - I*b)^(2/3)))/d - ((I/2)*(a + I*b)*(A + I*B)*((I*Sqrt[3]*ArcTanh[Tan[c + d*x]/Sqrt[3]])/(a + I*b)^(2/3) - Log[1 + I*Tan[c + d*x]]/(2*(a + I*b)^(2/3)) + (3*Log[(a + I*b)^(1/3) + I*Tan[c + d*x]])/(2*(a + I*b)^(2/3)))/d + (3*B*(a + b*Tan[c + d*x])^(1/3))/d`

3.474.3.1 Defintions of rubi rules used

rule 16 `Int[(c_)/((a_) + (b_)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 69 `Int[1/(((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(2/3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q^2), x] + (-Simp[3/(2*b*q) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Simp[3/(2*b*q^2) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x]) /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`

rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

```
rule 1082 Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4011 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]
```

```
rule 4020 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

3.474.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.61 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.26

method	result
derivativedivides	$\frac{3B(a+b \tan(dx+c))^{\frac{1}{3}} + \frac{\left(\frac{(Ab+Ba)R^3 - Ba^2 - Bb^2}{\sum_{R=\text{RootOf}(_Z^6 - 2a_Z^3 + a^2 + b^2)} \ln((a+b \tan(dx+c))^{\frac{1}{3}} - R)} \right)}{R^5 - R^2 a}}{d^2}$
default	$\frac{3B(a+b \tan(dx+c))^{\frac{1}{3}} + \frac{\left(\frac{(Ab+Ba)R^3 - Ba^2 - Bb^2}{\sum_{R=\text{RootOf}(_Z^6 - 2a_Z^3 + a^2 + b^2)} \ln((a+b \tan(dx+c))^{\frac{1}{3}} - R)} \right)}{R^5 - R^2 a}}{d^2}$
parts	$\frac{Ab \left(\frac{\sum_{R=\text{RootOf}(_Z^6 - 2a_Z^3 + a^2 + b^2)} \frac{R \ln((a+b \tan(dx+c))^{\frac{1}{3}} - R)}{R^3 - a}}{2d} \right) + B \left(\frac{3(a+b \tan(dx+c))^{\frac{1}{3}} + \frac{\left(\frac{(Ab+Ba)R^3 - Ba^2 - Bb^2}{\sum_{R=\text{RootOf}(_Z^6 - 2a_Z^3 + a^2 + b^2)} \ln((a+b \tan(dx+c))^{\frac{1}{3}} - R)} \right)}{R^5 - R^2 a}}{d^2} \right)}{d^2}$

```
input int((a+b*tan(d*x+c))^(1/3)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output 1/d*(3*B*(a+b*tan(d*x+c))^(1/3)+1/2*sum(((A*b+B*a)*_R^3-B*a^2-B*b^2)/(_R^5
-_R^2*a)*ln((a+b*tan(d*x+c))^(1/3)-_R),_R=RootOf(_Z^6-2*_Z^3*a+a^2+b^2)))
```

3.474.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2785 vs. 2(281) = 562.

Time = 0.37 (sec) , antiderivative size = 2785, normalized size of antiderivative = 7.39

$$\int \sqrt[3]{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^(1/3)*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

output `1/4*(2*d*(-(d^3*sqrt(-((A^6 - 6*A^4*B^2 + 9*A^2*B^4)*a^2 - 2*(3*A^5*B - 10*A^3*B^3 + 3*A*B^5)*a*b + (9*A^4*B^2 - 6*A^2*B^4 + B^6)*b^2)/d^6) + (3*A^2*B - B^3)*a + (A^3 - 3*A*B^2)*b)/d^3)^(1/3)*log(((A^5 - 2*A^3*B^2 - 3*A*B^4)*a - (3*A^4*B + 2*A^2*B^3 - B^5)*b)*(b*tan(d*x + c) + a)^(1/3) + (A*d^4*sqrt(-((A^6 - 6*A^4*B^2 + 9*A^2*B^4)*a^2 - 2*(3*A^5*B - 10*A^3*B^3 + 3*A*B^5)*a*b + (9*A^4*B^2 - 6*A^2*B^4 + B^6)*b^2)/d^6) - ((A^3*B - 3*A*B^3)*a - (3*A^2*B^2 - B^4)*b)*d)*(-(d^3*sqrt(-((A^6 - 6*A^4*B^2 + 9*A^2*B^4)*a^2 - 2*(3*A^5*B - 10*A^3*B^3 + 3*A*B^5)*a*b + (9*A^4*B^2 - 6*A^2*B^4 + B^6)*b^2)/d^6) + (3*A^2*B - B^3)*a + (A^3 - 3*A*B^2)*b)/d^3)^(1/3) - (sqrt(-3)*d + d)*(-(d^3*sqrt(-((A^6 - 6*A^4*B^2 + 9*A^2*B^4)*a^2 - 2*(3*A^5*B - 10*A^3*B^3 + 3*A*B^5)*a*b + (9*A^4*B^2 - 6*A^2*B^4 + B^6)*b^2)/d^6) + (3*A^2*B - B^3)*a + (A^3 - 3*A*B^2)*b)/d^3)^(1/3)*log(((A^5 - 2*A^3*B^2 - 3*A*B^4)*a - (3*A^4*B + 2*A^2*B^3 - B^5)*b)*(b*tan(d*x + c) + a)^(1/3) + 1/2*(sqrt(-3)*((A^3*B - 3*A*B^3)*a - (3*A^2*B^2 - B^4)*b)*d + ((A^3*B - 3*A*B^3)*a - (3*A^2*B^2 - B^4)*b)*d - (sqrt(-3)*A*d^4 + A*d^4)*sqrt(-((A^6 - 6*A^4*B^2 + 9*A^2*B^4)*a^2 - 2*(3*A^5*B - 10*A^3*B^3 + 3*A*B^5)*a*b + (9*A^4*B^2 - 6*A^2*B^4 + B^6)*b^2)/d^6))*(-(d^3*sqrt(-((A^6 - 6*A^4*B^2 + 9*A^2*B^4)*a^2 - 2*(3*A^5*B - 10*A^3*B^3 + 3*A*B^5)*a*b + (9*A^4*B^2 - 6*A^2*B^4 + B^6)*b^2)/d^6) + (3*A^2*B - B^3)*a + (A^3 - 3*A*B^2)*b)/d^3)^(1/3)) + (sqrt(-3)*d - d)*(-(d^3*sqrt(-((A^6 - 6*A^4*B^2 + 9*A^2*B^4)*a^2 - 2*(3*A^5*B - 10*A^3*B^3 + 3*A*B^5)*a*b + (9*A^4*B^2 - 6*A^2*B^4 + B^6)*b^2)/d^6) + (3*A^2*B - B^3)*a + (A^3 - 3*A*B^2)*b)/d^3)^(1/3))`

3.474.6 Sympy [F]

$$\int \sqrt[3]{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx)) \sqrt[3]{a + b \tan(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**(1/3)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(1/3), x)`

3.474.7 Maxima [F]

$$\begin{aligned} & \int \sqrt[3]{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{1}{3}} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))^(1/3)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(1/3), x)`

3.474.8 Giac [F]

$$\begin{aligned} & \int \sqrt[3]{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{1}{3}} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))^(1/3)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `undef`

3.474.9 Mupad [B] (verification not implemented)

Time = 20.45 (sec) , antiderivative size = 2537, normalized size of antiderivative = 6.73

$$\int \sqrt[3]{a + b \tan(c + dx)}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `int((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/3),x)`

output

$$\begin{aligned} & \log(a*d^7*((-A^6*a^2*d^6)^{(1/2)} - A^3*b*d^3)/d^6)^{(4/3)} + A*b*(a + b*\tan(c + d*x))^{(1/3)} * (-A^6*a^2*d^6)^{(1/2)} - A^4*a^2*d^3*(a + b*\tan(c + d*x))^{(1/3)} \\ & + 2*A^3*a*b*d^4*((-A^6*a^2*d^6)^{(1/2)} - A^3*b*d^3)/d^6)^{(1/3)} * (((-A^6*a^2*d^6)^{(1/2)} - A^3*b*d^3)/(8*d^6))^{(1/3)} + \log(A*b*(a + b*\tan(c + d*x))^{(1/3)} * (-A^6*a^2*d^6)^{(1/2)} + A^4*a^2*d^3*(a + b*\tan(c + d*x))^{(1/3)} + a*d * (-((-A^6*a^2*d^6)^{(1/2)} + A^3*b*d^3)/d^6)^{(1/3)} * (-A^6*a^2*d^6)^{(1/2)} - A^3*a*b*d^4 * (-((-A^6*a^2*d^6)^{(1/2)} + A^3*b*d^3)/d^6)^{(1/3)} * (-((-A^6*a^2*d^6)^{(1/2)} + A^3*b*d^3)/(8*d^6))^{(1/3)} + \log(d * (((-B^6*b^2*d^6)^{(1/2)} + B^3*a*d^3)/d^6)^{(1/3)} - B*(a + b*\tan(c + d*x))^{(1/3)} * (((-B^6*b^2*d^6)^{(1/2)} + B^3*a*d^3)/(8*d^6))^{(1/3)} + \log(d * (-((-B^6*b^2*d^6)^{(1/2)} - B^3*a*d^3)/d^6)^{(1/3)} - B*(a + b*\tan(c + d*x))^{(1/3)} * (-((-B^6*b^2*d^6)^{(1/2)} - B^3*a*d^3)/(8*d^6))^{(1/3)} + (\log(- ((3^{(1/2)}*1i - 1) * (((3^{(1/2)}*1i - 1)^2 * (1944*a*b^4*(3^{(1/2)}*1i - 1) * (((-B^6*b^2*d^6)^{(1/2)} + B^3*a*d^3)/d^6)^{(1/3)} * (a^2 + b^2) - (3888*B*a*b^4*(a^2 + b^2)*(a + b*\tan(c + d*x))^{(1/3)}))/d * (((-B^6*b^2*d^6)^{(1/2)} + B^3*a*d^3)/d^6)^{(2/3)})/16 - (972*B^3*b^4*(a^4 - b^4))/d^3 * (((-B^6*b^2*d^6)^{(1/2)} + B^3*a*d^3)/d^6)^{(1/3)})/4 - (486*B^4*b^4*(a^4 - b^4)*(a + b*\tan(c + d*x))^{(1/3)})/d^4 * (3^{(1/2)}*1i - 1) * ((-B^6*b^2*d^6)^{(1/2)}/(8*d^6) + (B^3*a)/(8*d^3))^{(1/3)})/2 - (\log(- ((3^{(1/2)}*1i + 1) * (((3^{(1/2)}*1i + 1)^2 * (1944*a*b^4*(3^{(1/2)}*1i + 1) * (((-B^6*b^2*d^6)^{(1/2)} + B^3*a*d^3)/d^6)^{(1/3)} * (a^2 + b^2) + (3888*B*a*b^4*(a^2 + b^2)*(a + b*\tan(c + ... \end{aligned}$$

$$3.475 \quad \int \frac{A+B \tan(c+dx)}{\sqrt[3]{a+b \tan(c+dx)}} dx$$

3.475.1 Optimal result	4542
3.475.2 Mathematica [A] (verified)	4543
3.475.3 Rubi [A] (warning: unable to verify)	4544
3.475.4 Maple [C] (verified)	4547
3.475.5 Fricas [B] (verification not implemented)	4548
3.475.6 Sympy [F]	4548
3.475.7 Maxima [F]	4549
3.475.8 Giac [A] (verification not implemented)	4549
3.475.9 Mupad [B] (verification not implemented)	4549

3.475.1 Optimal result

Integrand size = 25, antiderivative size = 357

$$\int \frac{A+B \tan(c+dx)}{\sqrt[3]{a+b \tan(c+dx)}} dx = -\frac{(A-iB)x}{4\sqrt[3]{a-ib}} - \frac{(A+iB)x}{4\sqrt[3]{a+ib}}$$

$$+ \frac{\sqrt{3}(iA+B) \arctan\left(\frac{1+\sqrt[3]{a+b \tan(c+dx)}}{\frac{\sqrt[3]{a-ib}}{\sqrt{3}}}\right)}{2\sqrt[3]{a-ibd}}$$

$$- \frac{\sqrt{3}(iA-B) \arctan\left(\frac{1+\sqrt[3]{a+b \tan(c+dx)}}{\frac{\sqrt[3]{a+ib}}{\sqrt{3}}}\right)}{2\sqrt[3]{a+ibd}}$$

$$- \frac{(iA-B) \log(\cos(c+dx))}{4\sqrt[3]{a+ibd}} + \frac{(iA+B) \log(\cos(c+dx))}{4\sqrt[3]{a-ibd}}$$

$$+ \frac{3(iA+B) \log\left(\sqrt[3]{a-ib} - \sqrt[3]{a+b \tan(c+dx)}\right)}{4\sqrt[3]{a-ibd}}$$

$$- \frac{3(iA-B) \log\left(\sqrt[3]{a+ib} - \sqrt[3]{a+b \tan(c+dx)}\right)}{4\sqrt[3]{a+ibd}}$$

output
$$-1/4*(A-I*B)*x/(a-I*b)^{(1/3)}-1/4*(A+I*B)*x/(a+I*b)^{(1/3)}-1/4*(I*A-B)*\ln(\cos(dx+c))/(a-I*b)^{(1/3)}/d+1/4*(I*A+B)*\ln(\cos(dx+c))/(a-I*b)^{(1/3)}/d+3/4*(I*A+B)*\ln((a-I*b)^{(1/3)}-(a+b*\tan(dx+c))^{(1/3)})/(a-I*b)^{(1/3)}/d-3/4*(I*A-B)*\ln((a+I*b)^{(1/3)}-(a+b*\tan(dx+c))^{(1/3)})/(a+I*b)^{(1/3)}/d+1/2*(I*A+B)*\arctan(1/3*(1+2*(a+b*\tan(dx+c))^{(1/3)})/(a-I*b)^{(1/3)})*3^{(1/2)}*3^{(1/2)}/(a-I*b)^{(1/3)}/d-1/2*(I*A-B)*\arctan(1/3*(1+2*(a+b*\tan(dx+c))^{(1/3)})/(a+I*b)^{(1/3)})*3^{(1/2)}*3^{(1/2)}/(a+I*b)^{(1/3)}/d$$

3.475.2 Mathematica [A] (verified)

Time = 0.46 (sec) , antiderivative size = 227, normalized size of antiderivative = 0.64

$$\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx$$

$$= \frac{i \left((A-iB) \left(2\sqrt{3} \arctan \left(\frac{1 + \sqrt[3]{a + b \tan(c + dx)}}{\frac{\sqrt[3]{a - ib}}{\sqrt{3}}} \right) - \log(i + \tan(c + dx)) + 3 \log \left(\sqrt[3]{a - ib} - \sqrt[3]{a + b \tan(c + dx)} \right) \right)}{\sqrt[3]{a - ib}} \right)}{4d}$$

input `Integrate[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^(1/3),x]`

output
$$\left(\frac{(I/4)*((A - I*B)*(2*sqrt[3]*ArcTan[(1 + (2*(a + b*Tan[c + d*x])^{(1/3)})/(a - I*b)^{(1/3)})/sqrt[3]] - Log[I + Tan[c + d*x]] + 3*Log[(a - I*b)^{(1/3)} - (a + b*Tan[c + d*x])^{(1/3)}])/(a - I*b)^{(1/3)} - ((A + I*B)*(2*sqrt[3]*ArcTan[(1 + (2*(a + b*Tan[c + d*x])^{(1/3)})/(a + I*b)^{(1/3)})/sqrt[3]] - Log[I - Tan[c + d*x]] + 3*Log[(a + I*b)^{(1/3)} - (a + b*Tan[c + d*x])^{(1/3)}])/(a + I*b)^{(1/3)))/d$$

3.475.3 Rubi [A] (warning: unable to verify)

Time = 0.54 (sec) , antiderivative size = 233, normalized size of antiderivative = 0.65, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$, Rules used = {3042, 4022, 3042, 4020, 25, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4022} \\
 & \frac{1}{2}(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx + \frac{1}{2}(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt[3]{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2}(A + iB) \int \frac{1 - i \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx + \frac{1}{2}(A - iB) \int \frac{i \tan(c + dx) + 1}{\sqrt[3]{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4020} \\
 & \frac{i(A - iB) \int -\frac{1}{(1 - i \tan(c + dx)) \sqrt[3]{a + b \tan(c + dx)}} d(i \tan(c + dx))}{i(A + iB) \int -\frac{2d}{(i \tan(c + dx) + 1) \sqrt[3]{a + b \tan(c + dx)}} d(-i \tan(c + dx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{i(A + iB) \int \frac{1}{(i \tan(c + dx) + 1) \sqrt[3]{a + b \tan(c + dx)}} d(-i \tan(c + dx))}{i(A - iB) \int \frac{2d}{(1 - i \tan(c + dx)) \sqrt[3]{a + b \tan(c + dx)}} d(i \tan(c + dx))} \\
 & \quad \downarrow \text{67}
 \end{aligned}$$

3.475. $\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx$

$$\frac{i(A - iB) \left(\frac{3}{2} \int \frac{1}{-\tan^2(c+dx) + (a-ib)^{2/3} + \sqrt[3]{a-ib} \sqrt[3]{a+b \tan(c+dx)}} d \sqrt[3]{a+b \tan(c+dx)} - \frac{3 \int \frac{1}{\sqrt[3]{a-ib-i \tan(c+dx)}} d \sqrt[3]{a-ib-i \tan(c+dx)}}{2 \sqrt[3]{a-ib}} \right)}{2d} - \frac{i(A + iB) \left(\frac{3}{2} \int \frac{1}{-\tan^2(c+dx) + (a+ib)^{2/3} + \sqrt[3]{a+ib} \sqrt[3]{a+b \tan(c+dx)}} d \sqrt[3]{a+b \tan(c+dx)} - \frac{3 \int \frac{1}{i \tan(c+dx) + \sqrt[3]{a+ib}} d \sqrt[3]{a+ib}}{2 \sqrt[3]{a+ib}} \right)}{2d}$$

↓ 16

$$\frac{i(A - iB) \left(\frac{3}{2} \int \frac{1}{-\tan^2(c+dx) + (a-ib)^{2/3} + \sqrt[3]{a-ib} \sqrt[3]{a+b \tan(c+dx)}} d \sqrt[3]{a+b \tan(c+dx)} - \frac{\log(1-i \tan(c+dx))}{2 \sqrt[3]{a-ib}} + \frac{3 \log(\sqrt[3]{a-ib-i \tan(c+dx)})}{2 \sqrt[3]{a-ib}} \right)}{2d} - \frac{i(A + iB) \left(\frac{3}{2} \int \frac{1}{-\tan^2(c+dx) + (a+ib)^{2/3} + \sqrt[3]{a+ib} \sqrt[3]{a+b \tan(c+dx)}} d \sqrt[3]{a+b \tan(c+dx)} - \frac{\log(1+i \tan(c+dx))}{2 \sqrt[3]{a+ib}} + \frac{3 \log(\sqrt[3]{a+ib+i \tan(c+dx)})}{2 \sqrt[3]{a+ib}} \right)}{2d}$$

↓ 1082

$$\frac{i(A - iB) \left(-\frac{3 \int \frac{1}{\tan^2(c+dx)-3} d \left(\frac{2i \tan(c+dx)}{\sqrt[3]{a-ib}} + 1 \right)}{\sqrt[3]{a-ib}} - \frac{\log(1-i \tan(c+dx))}{2 \sqrt[3]{a-ib}} + \frac{3 \log(\sqrt[3]{a-ib-i \tan(c+dx)})}{2 \sqrt[3]{a-ib}} \right)}{2d} - \frac{i(A + iB) \left(-\frac{3 \int \frac{1}{\tan^2(c+dx)-3} d \left(1 - \frac{2i \tan(c+dx)}{\sqrt[3]{a+ib}} \right)}{\sqrt[3]{a+ib}} - \frac{\log(1+i \tan(c+dx))}{2 \sqrt[3]{a+ib}} + \frac{3 \log(\sqrt[3]{a+ib+i \tan(c+dx)})}{2 \sqrt[3]{a+ib}} \right)}{2d}$$

↓ 217

$$\frac{i(A - iB) \left(\frac{i\sqrt{3} \arctanh\left(\frac{\tan(c+dx)}{\sqrt{3}}\right)}{\sqrt[3]{a-ib}} - \frac{\log(1-i \tan(c+dx))}{2 \sqrt[3]{a-ib}} + \frac{3 \log(\sqrt[3]{a-ib-i \tan(c+dx)})}{2 \sqrt[3]{a-ib}} \right)}{2d} - \frac{i(A + iB) \left(-\frac{i\sqrt{3} \arctanh\left(\frac{\tan(c+dx)}{\sqrt{3}}\right)}{\sqrt[3]{a+ib}} - \frac{\log(1+i \tan(c+dx))}{2 \sqrt[3]{a+ib}} + \frac{3 \log(\sqrt[3]{a+ib+i \tan(c+dx)})}{2 \sqrt[3]{a+ib}} \right)}{2d}$$

input `Int[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^(1/3),x]`

3.475. $\int \frac{A+B \tan(c+dx)}{\sqrt[3]{a+b \tan(c+dx)}} dx$


```
output ((I/2)*(A - I*B)*((I*Sqrt[3]*ArcTanh[Tan[c + d*x]/Sqrt[3]])/(a - I*b)^(1/3)
) - Log[1 - I*Tan[c + d*x]]/(2*(a - I*b)^(1/3)) + (3*Log[(a - I*b)^(1/3) -
I*Tan[c + d*x]]/(2*(a - I*b)^(1/3))))/d - ((I/2)*(A + I*B)*((-I)*Sqrt[3
]*ArcTanh[Tan[c + d*x]/Sqrt[3]])/(a + I*b)^(1/3) - Log[1 + I*Tan[c + d*x]]
/(2*(a + I*b)^(1/3)) + (3*Log[(a + I*b)^(1/3) + I*Tan[c + d*x]]/(2*(a + I
*b)^(1/3))))/d
```

3.475.3.1 Defintions of rubi rules used

```
rule 16 Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a +
b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 67 Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(1/3)), x_Symbol] := With[
{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q), x
] + (Simp[3/(2*b) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)],
x] - Simp[3/(2*b*q) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x])] /
; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

3.475.
$$\int \frac{A+B \tan(c+dx)}{\sqrt[3]{a+b \tan(c+dx)}} dx$$

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

3.475.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.43 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.20

method	result
derivativedivides	$\frac{\sum_{R=\text{RootOf}(_Z^6-2a_Z^3+a^2+b^2)} \frac{(B_R^4+(Ab-Ba)_R) \ln\left(\frac{(a+b \tan(dx+c))^{\frac{1}{3}}-_R}{_R^5-_R^2 a}\right)}{2d}}{\sum_{R=\text{RootOf}(_Z^6-2a_Z^3+a^2+b^2)} \frac{(B_R^4+(Ab-Ba)_R) \ln\left(\frac{(a+b \tan(dx+c))^{\frac{1}{3}}-_R}{_R^5-_R^2 a}\right)}{2d}}$
default	$\frac{\sum_{R=\text{RootOf}(_Z^6-2a_Z^3+a^2+b^2)} \frac{(B_R^4+(Ab-Ba)_R) \ln\left(\frac{(a+b \tan(dx+c))^{\frac{1}{3}}-_R}{_R^5-_R^2 a}\right)}{2d}}{\sum_{R=\text{RootOf}(_Z^6-2a_Z^3+a^2+b^2)} \frac{(B_R^4+(Ab-Ba)_R) \ln\left(\frac{(a+b \tan(dx+c))^{\frac{1}{3}}-_R}{_R^5-_R^2 a}\right)}{2d}}$
parts	$\frac{Ab \left(\sum_{R=\text{RootOf}(_Z^6-2a_Z^3+a^2+b^2)} \frac{\ln\left(\frac{(a+b \tan(dx+c))^{\frac{1}{3}}-_R}{_R^5-_R^2 a}\right)}{2d} \right)}{\sum_{R=\text{RootOf}(_Z^6-2a_Z^3+a^2+b^2)} \frac{\ln\left(\frac{(a+b \tan(dx+c))^{\frac{1}{3}}-_R}{_R^5-_R^2 a}\right)}{2d}} + \frac{B \left(\sum_{R=\text{RootOf}(_Z^6-2a_Z^3+a^2+b^2)} \frac{\ln\left(\frac{(a+b \tan(dx+c))^{\frac{1}{3}}-_R}{_R^5-_R^2 a}\right)}{2d} \right)}{\sum_{R=\text{RootOf}(_Z^6-2a_Z^3+a^2+b^2)} \frac{\ln\left(\frac{(a+b \tan(dx+c))^{\frac{1}{3}}-_R}{_R^5-_R^2 a}\right)}{2d}}$

```
input int((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/3),x,method=_RETURNVERBOSE)
```

```
output 1/2/d*sum((B*_R^4+(A*b-B*a)*_R)/(_R^5-_R^2*a)*ln((a+b*tan(d*x+c))^(1/3)-_R
),_R=RootOf(_Z^6-2*_Z^3*a+a^2+b^2))
```

$$3.475. \int \frac{A+B \tan(c+dx)}{\sqrt[3]{a+b \tan(c+dx)}} dx$$

3.475.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4121 vs. $2(263) = 526$.

Time = 0.44 (sec) , antiderivative size = 4121, normalized size of antiderivative = 11.54

$$\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/3),x, algorithm="fricas")`

output `1/4*(sqrt(-3) - 1)*(-((a^2 + b^2)*d^3*sqrt(-((A^6 - 6*A^4*B^2 + 9*A^2*B^4)*a^2 + 2*(3*A^5*B - 10*A^3*B^3 + 3*A*B^5)*a*b + (9*A^4*B^2 - 6*A^2*B^4 + B^6)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^6)) + (3*A^2*B - B^3)*a - (A^3 - 3*A*B^2)*b)/((a^2 + b^2)*d^3))^(1/3)*log(1/2*(sqrt(-3)*((A^5 - 4*A^3*B^2 + 3*A*B^4)*a^2 + (5*A^4*B - 10*A^2*B^3 + B^5)*a*b + 2*(3*A^3*B^2 - A*B^4)*b^2)*d^2 + ((A^5 - 4*A^3*B^2 + 3*A*B^4)*a^2 + (5*A^4*B - 10*A^2*B^3 + B^5)*a*b + 2*(3*A^3*B^2 - A*B^4)*b^2)*d^2 + (sqrt(-3)*(2*A*B*a^3 + 2*A*B*a*b^2 - (A^2 - B^2)*a^2*b - (A^2 - B^2)*b^3)*d^5 + (2*A*B*a^3 + 2*A*B*a*b^2 - (A^2 - B^2)*a^2*b - (A^2 - B^2)*b^3)*d^5)*sqrt(-((A^6 - 6*A^4*B^2 + 9*A^2*B^4)*a^2 + 2*(3*A^5*B - 10*A^3*B^3 + 3*A*B^5)*a*b + (9*A^4*B^2 - 6*A^2*B^4 + B^6)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^6)))*(-((a^2 + b^2)*d^3*sqrt(-((A^6 - 6*A^4*B^2 + 9*A^2*B^4)*a^2 + 2*(3*A^5*B - 10*A^3*B^3 + 3*A*B^5)*a*b + (9*A^4*B^2 - 6*A^2*B^4 + B^6)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^6)) + (3*A^2*B - B^3)*a - (A^3 - 3*A*B^2)*b)/((a^2 + b^2)*d^3))^(2/3) - ((A^7 - A^5*B^2 - 5*A^3*B^4 - 3*A*B^6)*a + (3*A^6*B + 5*A^4*B^3 + A^2*B^5 - B^7)*b)*(b*tan(d*x + c) + a)^(1/3) - 1/4*(sqrt(-3) + 1)*(-((a^2 + b^2)*d^3*sqrt(-((A^6 - 6*A^4*B^2 + 9*A^2*B^4)*a^2 + 2*(3*A^5*B - 10*A^3*B^3 + 3*A*B^5)*a*b + (9*A^4*B^2 - 6*A^2*B^4 + B^6)*b^2)/((a^4 + 2*a^2*b^2 + b^4)*d^6)) + (3*A^2*B - B^3)*a - (A^3 - 3*A*B^2)*b)/((a^2 + b^2)*d^3))^(1/3)*log(-1/2*(sqrt(-3)*((A^5 - 4*A^3*B^2 + 3*A*B^4)*a^2 + (5*A^4*B - 10*A^2*B^3 + B^5)*a*b + 2*(3*...`

3.475.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/3),x)`

output `Integral((A + B*tan(c + d*x))/(a + b*tan(c + d*x))**(1/3), x)`

3.475. $\int \frac{A+B \tan(c+dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx$

3.475.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a)^{\frac{1}{3}}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/3),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)/(b*tan(d*x + c) + a)^(1/3), x)`

3.475.8 Giac [A] (verification not implemented)

Time = 3.43 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.41

$$\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx$$

$$= \frac{\left(\frac{A^3 - 3iA^2B - 3AB^2 + iB^3}{8ia + 8b}\right)^{\frac{1}{3}} \log\left(-a + ib - (-a^2 + 2iab + b^2)^{\frac{1}{3}}(b \tan(dx + c) + a)^{\frac{1}{3}}\right) + \left(-\frac{A^3 + 3iA^2B - 3AB^2 - iB^3}{8ia - 8b}\right)^{\frac{1}{3}} \log\left(-a - ib + (-a^2 + 2iab + b^2)^{\frac{1}{3}}(b \tan(dx + c) + a)^{\frac{1}{3}}\right)}{d}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/3),x, algorithm="giac")`

output `((((A^3 - 3*I*A^2*B - 3*A*B^2 + I*B^3)/(8*I*a + 8*b))^(1/3)*log(-a + I*b - (-a^2 + 2*I*a*b + b^2)^(1/3)*(b*tan(d*x + c) + a)^(1/3)) + (-A^3 + 3*I*A^2*B - 3*A*B^2 - I*B^3)/(8*I*a - 8*b))^(1/3)*log(-a - I*b + (a^2 + 2*I*a*b - b^2)^(1/3)*(b*tan(d*x + c) + a)^(1/3)))/d`

3.475.9 Mupad [B] (verification not implemented)

Time = 20.96 (sec) , antiderivative size = 3228, normalized size of antiderivative = 9.04

$$\int \frac{A + B \tan(c + dx)}{\sqrt[3]{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `int((A + B*tan(c + d*x))/(a + b*tan(c + d*x))^(1/3),x)`

3.476 $\int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^{2/3}} dx$

3.476.1 Optimal result 4551
 3.476.2 Mathematica [A] (verified) 4552
 3.476.3 Rubi [A] (warning: unable to verify) 4553
 3.476.4 Maple [C] (verified) 4556
 3.476.5 Fricas [B] (verification not implemented) 4557
 3.476.6 Sympy [F] 4557
 3.476.7 Maxima [F] 4557
 3.476.8 Giac [F] 4558
 3.476.9 Mupad [B] (verification not implemented) 4558

3.476.1 Optimal result

Integrand size = 25, antiderivative size = 357

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx = -\frac{(A - iB)x}{4(a - ib)^{2/3}} - \frac{(A + iB)x}{4(a + ib)^{2/3}}$$

$$- \frac{\sqrt{3}(iA + B) \arctan\left(\frac{1 + \sqrt[3]{a + b \tan(c + dx)}}{\frac{\sqrt[3]{a - ib}}{\sqrt{3}}}\right)}{2(a - ib)^{2/3}d}$$

$$+ \frac{\sqrt{3}(iA - B) \arctan\left(\frac{1 + \sqrt[3]{a + b \tan(c + dx)}}{\frac{\sqrt[3]{a + ib}}{\sqrt{3}}}\right)}{2(a + ib)^{2/3}d} - \frac{(iA - B) \log(\cos(c + dx))}{4(a + ib)^{2/3}d}$$

$$+ \frac{(iA + B) \log(\cos(c + dx))}{4(a - ib)^{2/3}d} + \frac{3(iA + B) \log\left(\sqrt[3]{a - ib} - \sqrt[3]{a + b \tan(c + dx)}\right)}{4(a - ib)^{2/3}d}$$

$$- \frac{3(iA - B) \log\left(\sqrt[3]{a + ib} - \sqrt[3]{a + b \tan(c + dx)}\right)}{4(a + ib)^{2/3}d}$$

output $-1/4*(A-I*B)*x/(a-I*b)^(2/3)-1/4*(A+I*B)*x/(a+I*b)^(2/3)-1/4*(I*A-B)*\ln(\cos(dx+c))/(a-I*b)^(2/3)/d+1/4*(I*A+B)*\ln(\cos(dx+c))/(a-I*b)^(2/3)/d+3/4*(I*A+B)*\ln((a-I*b)^(1/3)-(a+b*\tan(dx+c))^(1/3))/(a-I*b)^(2/3)/d-3/4*(I*A-B)*\ln((a+I*b)^(1/3)-(a+b*\tan(dx+c))^(1/3))/(a+I*b)^(2/3)/d-1/2*(I*A+B)*\arctan(1/3*(1+2*(a+b*\tan(dx+c))^(1/3))/(a-I*b)^(1/3))*3^(1/2)/3^(1/2)/(a-I*b)^(2/3)/d+1/2*(I*A-B)*\arctan(1/3*(1+2*(a+b*\tan(dx+c))^(1/3))/(a+I*b)^(1/3))*3^(1/2)/3^(1/2)/(a+I*b)^(2/3)/d$

3.476.2 Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 305, normalized size of antiderivative = 0.85

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx = \frac{i \left((A - iB) \left(2\sqrt{3} \arctan \left(\frac{1 + \sqrt[3]{a + b \tan(c + dx)}}{\sqrt[3]{a - ib}} \right) - 2 \log \left(\sqrt[3]{a - ib} - \sqrt[3]{a + b \tan(c + dx)} \right) \right)}{(a - ib)^2}$$

input `Integrate[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^(2/3),x]`

output $((I/4)*(-(((A - I*B)*(2*sqrt[3]*ArcTan[(1 + (2*(a + b*Tan[c + d*x])^(1/3)))/(a - I*b)^(1/3))/sqrt[3]] - 2*Log[(a - I*b)^(1/3) - (a + b*Tan[c + d*x])^(1/3)] + Log[(a - I*b)^(2/3) + (a - I*b)^(1/3)*(a + b*Tan[c + d*x])^(1/3) + (a + b*Tan[c + d*x])^(2/3)]))/(a - I*b)^(2/3)) + ((A + I*B)*(2*sqrt[3]*ArcTan[(1 + (2*(a + b*Tan[c + d*x])^(1/3)))/(a + I*b)^(1/3))/sqrt[3]] - 2*Log[(a + I*b)^(1/3) - (a + b*Tan[c + d*x])^(1/3)] + Log[(a + I*b)^(2/3) + (a + I*b)^(1/3)*(a + b*Tan[c + d*x])^(1/3) + (a + b*Tan[c + d*x])^(2/3)]))/(a + I*b)^(2/3))/d$

3.476.3 Rubi [A] (warning: unable to verify)

Time = 0.57 (sec) , antiderivative size = 233, normalized size of antiderivative = 0.65, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$, Rules used = {3042, 4022, 3042, 4020, 25, 69, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx \\
 & \quad \downarrow \text{4022} \\
 & \frac{1}{2}(A + iB) \int \frac{1 - i \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx + \frac{1}{2}(A - iB) \int \frac{i \tan(c + dx) + 1}{(a + b \tan(c + dx))^{2/3}} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2}(A + iB) \int \frac{1 - i \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx + \frac{1}{2}(A - iB) \int \frac{i \tan(c + dx) + 1}{(a + b \tan(c + dx))^{2/3}} dx \\
 & \quad \downarrow \text{4020} \\
 & \frac{i(A - iB) \int -\frac{1}{(1 - i \tan(c + dx))(a + b \tan(c + dx))^{2/3}} d(i \tan(c + dx))}{2d} - \\
 & \frac{i(A + iB) \int -\frac{1}{(i \tan(c + dx) + 1)(a + b \tan(c + dx))^{2/3}} d(-i \tan(c + dx))}{2d} \\
 & \quad \downarrow \text{25} \\
 & \frac{i(A + iB) \int \frac{1}{(i \tan(c + dx) + 1)(a + b \tan(c + dx))^{2/3}} d(-i \tan(c + dx))}{2d} - \\
 & \frac{i(A - iB) \int \frac{1}{(1 - i \tan(c + dx))(a + b \tan(c + dx))^{2/3}} d(i \tan(c + dx))}{2d} \\
 & \quad \downarrow \text{69}
 \end{aligned}$$

$$i(A - iB) \left(\frac{3 \int \frac{1}{-\tan^2(c+dx)+(a-ib)^{2/3} + \sqrt[3]{a-ib} \sqrt[3]{a+b \tan(c+dx)}} d \sqrt[3]{a+b \tan(c+dx)}}{2 \sqrt[3]{a-ib}} - \frac{3 \int \frac{1}{\sqrt[3]{a-ib} - i \tan(c+dx)}}{2(a-ib)^{2/3}} d \sqrt[3]{a+b \tan(c+dx)} \right)$$

$$i(A + iB) \left(\frac{3 \int \frac{1}{-\tan^2(c+dx)+(a+ib)^{2/3} + \sqrt[3]{a+ib} \sqrt[3]{a+b \tan(c+dx)}} d \sqrt[3]{a+b \tan(c+dx)}}{2 \sqrt[3]{a+ib}} - \frac{3 \int \frac{1}{i \tan(c+dx) + \sqrt[3]{a+ib}} d \sqrt[3]{a+b \tan(c+dx)}}{2(a+ib)^{2/3}} \right)$$

2d

↓ 16

$$i(A - iB) \left(\frac{3 \int \frac{1}{-\tan^2(c+dx)+(a-ib)^{2/3} + \sqrt[3]{a-ib} \sqrt[3]{a+b \tan(c+dx)}} d \sqrt[3]{a+b \tan(c+dx)}}{2 \sqrt[3]{a-ib}} - \frac{\log(1-i \tan(c+dx))}{2(a-ib)^{2/3}} + \frac{3 \log(\sqrt[3]{a-ib} - i \tan(c+dx))}{2(a-ib)^{2/3}} \right)$$

$$i(A + iB) \left(\frac{3 \int \frac{1}{-\tan^2(c+dx)+(a+ib)^{2/3} + \sqrt[3]{a+ib} \sqrt[3]{a+b \tan(c+dx)}} d \sqrt[3]{a+b \tan(c+dx)}}{2 \sqrt[3]{a+ib}} - \frac{\log(1+i \tan(c+dx))}{2(a+ib)^{2/3}} + \frac{3 \log(\sqrt[3]{a+ib} + i \tan(c+dx))}{2(a+ib)^{2/3}} \right)$$

2d

↓ 1082

$$i(A - iB) \left(\frac{3 \int \frac{1}{\tan^2(c+dx)-3} d \left(\frac{2i \tan(c+dx)}{\sqrt[3]{a-ib}} + 1 \right)}{(a-ib)^{2/3}} - \frac{\log(1-i \tan(c+dx))}{2(a-ib)^{2/3}} + \frac{3 \log(\sqrt[3]{a-ib} - i \tan(c+dx))}{2(a-ib)^{2/3}} \right)$$

2d

$$i(A + iB) \left(\frac{3 \int \frac{1}{\tan^2(c+dx)-3} d \left(1 - \frac{2i \tan(c+dx)}{\sqrt[3]{a+ib}} \right)}{(a+ib)^{2/3}} - \frac{\log(1+i \tan(c+dx))}{2(a+ib)^{2/3}} + \frac{3 \log(\sqrt[3]{a+ib} + i \tan(c+dx))}{2(a+ib)^{2/3}} \right)$$

2d

↓ 217

$$i(A - iB) \left(-\frac{i\sqrt{3} \operatorname{arctanh}\left(\frac{\tan(c+dx)}{\sqrt{3}}\right)}{(a-ib)^{2/3}} - \frac{\log(1-i \tan(c+dx))}{2(a-ib)^{2/3}} + \frac{3 \log(\sqrt[3]{a-ib} - i \tan(c+dx))}{2(a-ib)^{2/3}} \right)$$

2d

$$i(A + iB) \left(\frac{i\sqrt{3} \operatorname{arctanh}\left(\frac{\tan(c+dx)}{\sqrt{3}}\right)}{(a+ib)^{2/3}} - \frac{\log(1+i \tan(c+dx))}{2(a+ib)^{2/3}} + \frac{3 \log(\sqrt[3]{a+ib} + i \tan(c+dx))}{2(a+ib)^{2/3}} \right)$$

2d

input `Int[(A + B*Tan[c + d*x])/(a + b*Tan[c + d*x])^(2/3),x]`

```
output ((I/2)*(A - I*B)*((-I)*Sqrt[3]*ArcTanh[Tan[c + d*x]/Sqrt[3]]/(a - I*b)^(
2/3) - Log[1 - I*Tan[c + d*x]]/(2*(a - I*b)^(2/3)) + (3*Log[(a - I*b)^(1/3
) - I*Tan[c + d*x]]/(2*(a - I*b)^(2/3))))/d - ((I/2)*(A + I*B)*((I*Sqrt[3
]*ArcTanh[Tan[c + d*x]/Sqrt[3]]/(a + I*b)^(2/3) - Log[1 + I*Tan[c + d*x]]
/(2*(a + I*b)^(2/3)) + (3*Log[(a + I*b)^(1/3) + I*Tan[c + d*x]]/(2*(a + I
*b)^(2/3)))))/d
```

3.476.3.1 Defintions of rubi rules used

```
rule 16 Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a +
b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]
```

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 69 Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(2/3)), x_Symbol] := With[
{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q^2),
x] + (-Simp[3/(2*b*q) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1
/3)], x] - Simp[3/(2*b*q^2) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)],
x])] /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4020 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```

3.476.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.40 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.19

method	result
derivativedivides	$\frac{\sum_{R=\text{RootOf}(_Z^6-2a_Z^3+a^2+b^2)} \frac{(B_R^3+Ab-Ba) \ln\left(\frac{(a+b \tan(dx+c))^{\frac{1}{3}}-R}{a}\right)}{_R^5-_R^2 a}}{2d}$
default	$\frac{\sum_{R=\text{RootOf}(_Z^6-2a_Z^3+a^2+b^2)} \frac{(B_R^3+Ab-Ba) \ln\left(\frac{(a+b \tan(dx+c))^{\frac{1}{3}}-R}{a}\right)}{_R^5-_R^2 a}}{2d}$
parts	$\frac{Ab \left(\sum_{R=\text{RootOf}(_Z^6-2a_Z^3+a^2+b^2)} \frac{\ln\left(\frac{(a+b \tan(dx+c))^{\frac{1}{3}}-R}{a}\right)}{_R^2(_R^3-a)} \right)}{2d} + \frac{B \left(\sum_{R=\text{RootOf}(_Z^6-2a_Z^3+a^2+b^2)} \frac{\ln\left(\frac{(a+b \tan(dx+c))^{\frac{1}{3}}-R}{a}\right)}{_R^2(_R^3-a)} \right)}{2d}$

```
input int((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(2/3),x,method=_RETURNVERBOSE)
```

```
output 1/2/d*sum((B*_R^3+A*b-B*a)/(_R^5-_R^2*a)*ln((a+b*tan(d*x+c))^(1/3)-_R),_R=
RootOf(_Z^6-2*_Z^3*a+a^2+b^2))
```

$$3.476. \int \frac{A+B \tan(c+dx)}{(a+b \tan(c+dx))^{2/3}} dx$$

3.476.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6073 vs. $2(263) = 526$.

Time = 1.35 (sec) , antiderivative size = 6073, normalized size of antiderivative = 17.01

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(2/3),x, algorithm="fricas")`

output Too large to include

3.476.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(2/3),x)`

output `Integral((A + B*tan(c + d*x))/(a + b*tan(c + d*x))**(2/3), x)`

3.476.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a)^{2/3}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(2/3),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)/(b*tan(d*x + c) + a)^(2/3), x)`

3.476.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a)^{2/3}} dx$$

input `integrate((A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(2/3),x, algorithm="giac")`

output `undef`

3.476.9 Mupad [B] (verification not implemented)

Time = 23.04 (sec) , antiderivative size = 4562, normalized size of antiderivative = 12.78

$$\int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{2/3}} dx = \text{Too large to display}$$

input `int((A + B*tan(c + d*x))/(a + b*tan(c + d*x))^(2/3),x)`

output `log((((16*(-B^6*a^2*b^2*d^6)^(1/2) + 8*B^3*a^2*d^3 - 8*B^3*b^2*d^3)/(d^6*(a^2 + b^2)^2))^(1/3)*(1944*a*b^4*(-B^6*a^2*b^2*d^6)^(1/2) - 1944*B^3*a*b^6*d^3 + 243*B*b^8*d^5*(a + b*tan(c + d*x))^(1/3)*((16*(-B^6*a^2*b^2*d^6)^(1/2) + 8*B^3*a^2*d^3 - 8*B^3*b^2*d^3)/(d^6*(a^2 + b^2)^2))^(2/3) - 243*B*a^4*b^4*d^5*(a + b*tan(c + d*x))^(1/3)*((16*(-B^6*a^2*b^2*d^6)^(1/2) + 8*B^3*a^2*d^3 - 8*B^3*b^2*d^3)/(d^6*(a^2 + b^2)^2))^(2/3)))/(4*d^6*(a^2 + b^2)) + (486*B^4*b^4*(a + b*tan(c + d*x))^(1/3))/d^4)*((((16*B^3*a^2*d^3 - 16*B^3*b^2*d^3)^2/4 - B^6*(64*a^4*d^6 + 64*b^4*d^6 + 128*a^2*b^2*d^6))^(1/2) + 8*B^3*a^2*d^3 - 8*B^3*b^2*d^3)/(64*(a^4*d^6 + b^4*d^6 + 2*a^2*b^2*d^6)))^(1/3) + log((486*B^4*b^4*(a + b*tan(c + d*x))^(1/3))/d^4 - (((-16*(-B^6*a^2*b^2*d^6)^(1/2) - 8*B^3*a^2*d^3 + 8*B^3*b^2*d^3)/(d^6*(a^2 + b^2)^2))^(1/3)*(1944*a*b^4*(-B^6*a^2*b^2*d^6)^(1/2) + 1944*B^3*a*b^6*d^3 - 243*B*b^8*d^5*(a + b*tan(c + d*x))^(1/3)*(-16*(-B^6*a^2*b^2*d^6)^(1/2) - 8*B^3*a^2*d^3 + 8*B^3*b^2*d^3)/(d^6*(a^2 + b^2)^2))^(2/3) + 243*B*a^4*b^4*d^5*(a + b*tan(c + d*x))^(1/3)*(-16*(-B^6*a^2*b^2*d^6)^(1/2) - 8*B^3*a^2*d^3 + 8*B^3*b^2*d^3)/(d^6*(a^2 + b^2)^2)))/(4*d^6*(a^2 + b^2))*(-((((16*B^3*a^2*d^3 - 16*B^3*b^2*d^3)^2/4 - B^6*(64*a^4*d^6 + 64*b^4*d^6 + 128*a^2*b^2*d^6))^(1/2) - 8*B^3*a^2*d^3 + 8*B^3*b^2*d^3)/(64*(a^4*d^6 + b^4*d^6 + 2*a^2*b^2*d^6)))^(1/3) + log((((1944*a*b^4*(a^2 + b^2)*((8*(-A^6*d^6*(a^2 - b^2)^2)^(1/2) + 16*A^3*a*b*d^3)/(d^6*(a^2 + b^2)^2))^(1/3) + (7776*A*a*b^...`

$$3.477 \quad \int \frac{i - \tan(e + fx)}{\sqrt[3]{c + d \tan(e + fx)}} dx$$

3.477.1 Optimal result	4559
3.477.2 Mathematica [A] (verified)	4559
3.477.3 Rubi [A] (warning: unable to verify)	4560
3.477.4 Maple [C] (verified)	4562
3.477.5 Fricas [B] (verification not implemented)	4563
3.477.6 Sympy [F]	4564
3.477.7 Maxima [F]	4564
3.477.8 Giac [B] (verification not implemented)	4564
3.477.9 Mupad [B] (verification not implemented)	4565

3.477.1 Optimal result

Integrand size = 27, antiderivative size = 148

$$\int \frac{i - \tan(e + fx)}{\sqrt[3]{c + d \tan(e + fx)}} dx = -\frac{ix}{2\sqrt[3]{c - id}} - \frac{\sqrt{3} \arctan\left(\frac{1 + \sqrt[3]{c + d \tan(e + fx)}}{\frac{\sqrt[3]{c - id}}{\sqrt{3}}}\right)}{\sqrt[3]{c - id}f} - \frac{\log(\cos(e + fx))}{2\sqrt[3]{c - id}f} - \frac{3 \log\left(\sqrt[3]{c - id} - \sqrt[3]{c + d \tan(e + fx)}\right)}{2\sqrt[3]{c - id}f}$$

output
$$-1/2*I*x/(c-I*d)^{(1/3)}-1/2*\ln(\cos(f*x+e))/(c-I*d)^{(1/3)}/f-3/2*\ln((c-I*d)^{(1/3)}-(c+d*\tan(f*x+e))^{(1/3)})/(c-I*d)^{(1/3)}/f-\arctan(1/3*(1+2*(c+d*\tan(f*x+e))^{(1/3)})/(c-I*d)^{(1/3}))*3^{(1/2)}/(c-I*d)^{(1/3)}/f$$

3.477.2 Mathematica [A] (verified)

Time = 0.67 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.70

$$\int \frac{i - \tan(e + fx)}{\sqrt[3]{c + d \tan(e + fx)}} dx$$

$$= \frac{-2\sqrt{3} \arctan\left(\frac{1 + \sqrt[3]{c + d \tan(e + fx)}}{\frac{\sqrt[3]{c - id}}{\sqrt{3}}}\right) + \log(i + \tan(e + fx)) - 3 \log\left(\sqrt[3]{c - id} - \sqrt[3]{c + d \tan(e + fx)}\right)}{2\sqrt[3]{c - id}f}$$

3.477. $\int \frac{i - \tan(e + fx)}{\sqrt[3]{c + d \tan(e + fx)}} dx$

input `Integrate[(I - Tan[e + f*x])/(c + d*Tan[e + f*x])^(1/3),x]`

output `(-2*Sqrt[3]*ArcTan[(1 + (2*(c + d*Tan[e + f*x])^(1/3))/(c - I*d)^(1/3)]/Sqrt[3]] + Log[I + Tan[e + f*x]] - 3*Log[(c - I*d)^(1/3) - (c + d*Tan[e + f*x])^(1/3)])/(2*(c - I*d)^(1/3)*f)`

3.477.3 Rubi [A] (warning: unable to verify)

Time = 0.30 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.71, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3042, 4020, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{-\tan(e+fx) + i}{\sqrt[3]{c + d \tan(e+fx)}} dx$$

↓ 3042

$$\int \frac{-\tan(e+fx) + i}{\sqrt[3]{c + d \tan(e+fx)}} dx$$

↓ 4020

$$-\frac{i \int \frac{1}{(1-i \tan(e+fx)) \sqrt[3]{c + d \tan(e+fx)}} d(-\tan(e+fx))}{f}$$

↓ 67

$$i \left(-\frac{3}{2} i \int \frac{1}{\tan^2(e+fx) + (c-id)^{2/3} + \sqrt[3]{c-id} \sqrt[3]{c + d \tan(e+fx)}} d \sqrt[3]{c + d \tan(e+fx)} + \frac{3i \int \frac{1}{\tan(e+fx) + \sqrt[3]{c-id}} d \sqrt[3]{c + d \tan(e+fx)}}{2 \sqrt[3]{c-id}} \right)$$

↓ 16

$$i \left(-\frac{3}{2} i \int \frac{1}{\tan^2(e+fx) + (c-id)^{2/3} + \sqrt[3]{c-id} \sqrt[3]{c + d \tan(e+fx)}} d \sqrt[3]{c + d \tan(e+fx)} + \frac{i \log(\tan(e+fx) + i)}{2 \sqrt[3]{c-id}} - \frac{3i \log(\tan(e+fx) + i)}{2 \sqrt[3]{c-id}} \right)$$

↓ 1082

3.477. $\int \frac{i - \tan(e+fx)}{\sqrt[3]{c + d \tan(e+fx)}} dx$

$$\frac{i \left(\frac{3i f \frac{1}{-\tan^2(e+fx)-3} d \left(1 - \frac{2 \tan(e+fx)}{\sqrt[3]{c-id}} \right)}{\sqrt[3]{c-id}} + \frac{i \log(\tan(e+fx)+i)}{2\sqrt[3]{c-id}} - \frac{3i \log(\tan(e+fx)+\sqrt[3]{c-id})}{2\sqrt[3]{c-id}} \right)}{f}$$

↓ 217

$$\frac{i \left(\frac{i\sqrt{3} \arctan\left(\frac{\tan(e+fx)}{\sqrt{3}}\right)}{\sqrt[3]{c-id}} + \frac{i \log(\tan(e+fx)+i)}{2\sqrt[3]{c-id}} - \frac{3i \log(\tan(e+fx)+\sqrt[3]{c-id})}{2\sqrt[3]{c-id}} \right)}{f}$$

input `Int[(I - Tan[e + f*x])/(c + d*Tan[e + f*x])^(1/3),x]`

output `((-I)*((I*Sqrt[3]*ArcTan[Tan[e + f*x]/Sqrt[3]])/(c - I*d)^(1/3) + ((I/2)*Log[I + Tan[e + f*x]]/(c - I*d)^(1/3) - (((3*I)/2)*Log[(c - I*d)^(1/3) + Tan[e + f*x]]/(c - I*d)^(1/3)))/f`

3.477.3.1 Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 67 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(1/3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q), x] + (Simp[3/(2*b) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Simp[3/(2*b*q) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x])] /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

3.477. $\int \frac{i - \tan(e+fx)}{\sqrt[3]{c + d \tan(e+fx)}} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

3.477.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.36 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.28

method	result
derivativedivides	$-\frac{\sum_{R=\text{RootOf}(_Z^3+id-c)} \frac{\ln\left((c+d\tan(fx+e))^{\frac{1}{3}}-R\right)}{R}}{f}$
default	$-\frac{\sum_{R=\text{RootOf}(_Z^3+id-c)} \frac{\ln\left((c+d\tan(fx+e))^{\frac{1}{3}}-R\right)}{R}}{f}$
parts	$\frac{id\left(\sum_{R=\text{RootOf}(_Z^3-2c_Z^3+c^2+d^2)} \frac{\ln\left((c+d\tan(fx+e))^{\frac{1}{3}}-R\right)}{R(R^3-c)}\right)}{2f} - \frac{\sum_{R=\text{RootOf}(_Z^3-2c_Z^3+c^2+d^2)} \frac{\ln\left((c+d\tan(fx+e))^{\frac{1}{3}}-R\right)}{R}}{2f}$

input `int((1-tan(f*x+e))/(c+d*tan(f*x+e))^(1/3),x,method=_RETURNVERBOSE)`

output `-1/f*sum(1/_R*ln((c+d*tan(f*x+e))^(1/3)-_R),_R=RootOf(_Z^3+I*d-c))`

3.477.
$$\int \frac{i - \tan(e+fx)}{\sqrt[3]{c+d \tan(e+fx)}} dx$$

3.477.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 278 vs. $2(109) = 218$.

Time = 0.25 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.88

$$\int \frac{i - \tan(e + fx)}{\sqrt[3]{c + d \tan(e + fx)}} dx$$

$$= \frac{1}{2} (i\sqrt{3} - 1) \left(-\frac{i}{(ic + d)f^3} \right)^{\frac{1}{3}} \log \left(\frac{1}{2} (\sqrt{3}(ic + d)f^2 + (c - id)f^2) \left(-\frac{i}{(ic + d)f^3} \right)^{\frac{2}{3}} \right. \\ \left. + \left(\frac{(c - id)e^{(2ifx + 2ie)} + c + id}{e^{(2ifx + 2ie)} + 1} \right)^{\frac{1}{3}} \right) \\ + \frac{1}{2} (-i\sqrt{3} - 1) \left(-\frac{i}{(ic + d)f^3} \right)^{\frac{1}{3}} \log \left(\frac{1}{2} (\sqrt{3}(-ic - d)f^2 + (c - id)f^2) \left(-\frac{i}{(ic + d)f^3} \right)^{\frac{2}{3}} \right. \\ \left. + \left(\frac{(c - id)e^{(2ifx + 2ie)} + c + id}{e^{(2ifx + 2ie)} + 1} \right)^{\frac{1}{3}} \right) \\ + \left(-\frac{i}{(ic + d)f^3} \right)^{\frac{1}{3}} \log \left(-(c - id)f^2 \left(-\frac{i}{(ic + d)f^3} \right)^{\frac{2}{3}} \right. \\ \left. + \left(\frac{(c - id)e^{(2ifx + 2ie)} + c + id}{e^{(2ifx + 2ie)} + 1} \right)^{\frac{1}{3}} \right)$$

input `integrate((I-tan(f*x+e))/(c+d*tan(f*x+e))^(1/3),x, algorithm="fracas")`

output `1/2*(I*sqrt(3) - 1)*(-I/((I*c + d)*f^3))^(1/3)*log(1/2*(sqrt(3)*(I*c + d)*f^2 + (c - I*d)*f^2)*(-I/((I*c + d)*f^3))^(2/3) + (((c - I*d)*e^(2*I*f*x + 2*I*e) + c + I*d)/(e^(2*I*f*x + 2*I*e) + 1))^(1/3)) + 1/2*(-I*sqrt(3) - 1)*(-I/((I*c + d)*f^3))^(1/3)*log(1/2*(sqrt(3)*(-I*c - d)*f^2 + (c - I*d)*f^2)*(-I/((I*c + d)*f^3))^(2/3) + (((c - I*d)*e^(2*I*f*x + 2*I*e) + c + I*d)/(e^(2*I*f*x + 2*I*e) + 1))^(1/3)) + (-I/((I*c + d)*f^3))^(1/3)*log(-(c - I*d)*f^2*(-I/((I*c + d)*f^3))^(2/3) + (((c - I*d)*e^(2*I*f*x + 2*I*e) + c + I*d)/(e^(2*I*f*x + 2*I*e) + 1))^(1/3))`

3.477.6 Sympy [F]

$$\int \frac{i - \tan(e + fx)}{\sqrt[3]{c + d \tan(e + fx)}} dx = - \int \left(-\frac{i}{\sqrt[3]{c + d \tan(e + fx)}} \right) dx - \int \frac{\tan(e + fx)}{\sqrt[3]{c + d \tan(e + fx)}} dx$$

input `integrate((I-tan(f*x+e))/(c+d*tan(f*x+e))**(1/3),x)`

output `-Integral(-I/(c + d*tan(e + f*x))**(1/3), x) - Integral(tan(e + f*x)/(c + d*tan(e + f*x))**(1/3), x)`

3.477.7 Maxima [F]

$$\int \frac{i - \tan(e + fx)}{\sqrt[3]{c + d \tan(e + fx)}} dx = \int -\frac{\tan(fx + e) - i}{(d \tan(fx + e) + c)^{\frac{1}{3}}} dx$$

input `integrate((I-tan(f*x+e))/(c+d*tan(f*x+e))^(1/3),x, algorithm="maxima")`

output `-integrate((tan(f*x + e) - I)/(d*tan(f*x + e) + c)^(1/3), x)`

3.477.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 795 vs. $2(109) = 218$.

Time = 0.66 (sec) , antiderivative size = 795, normalized size of antiderivative = 5.37

$$\int \frac{i - \tan(e + fx)}{\sqrt[3]{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input `integrate((I-tan(f*x+e))/(c+d*tan(f*x+e))^(1/3),x, algorithm="giac")`

output

```

-(c - I*d)^(2/3)*log((d*tan(f*x + e) + c)^(1/3) - (c - I*d)^(1/3))/(c*f -
I*d*f) - (sqrt(3)*c*cos(1/6*pi*sgn(c)*sgn(d) - 1/6*pi*sgn(d) - 1/3*arctan(
d/c))^2 - sqrt(3)*c*sin(1/6*pi*sgn(c)*sgn(d) - 1/6*pi*sgn(d) - 1/3*arctan(
d/c))^2 + 2*c*cos(1/6*pi*sgn(c)*sgn(d) - 1/6*pi*sgn(d) - 1/3*arctan(d/c))*
sin(1/6*pi*sgn(c)*sgn(d) - 1/6*pi*sgn(d) - 1/3*arctan(d/c))*arctan(1/3*sq
rt(3)*(2*(d*tan(f*x + e) + c)^(1/3) + (c - I*d)^(1/3))/(c - I*d)^(1/3))/((
c^2 + d^2)^(2/3)*f) - I*(sqrt(3)*d*cos(1/6*pi*sgn(c)*sgn(d) - 1/6*pi*sgn(d
) - 1/3*arctan(d/c))^2 - sqrt(3)*d*sin(1/6*pi*sgn(c)*sgn(d) - 1/6*pi*sgn(d
) - 1/3*arctan(d/c))^2 + 2*d*cos(1/6*pi*sgn(c)*sgn(d) - 1/6*pi*sgn(d) - 1/
3*arctan(d/c))*sin(1/6*pi*sgn(c)*sgn(d) - 1/6*pi*sgn(d) - 1/3*arctan(d/c))
)*arctan(1/3*sqrt(3)*(2*(d*tan(f*x + e) + c)^(1/3) + (c - I*d)^(1/3))/(c -
I*d)^(1/3))/((c^2 + d^2)^(2/3)*f) - 1/2*(2*sqrt(3)*c*cos(1/6*pi*sgn(c)*sg
n(d) - 1/6*pi*sgn(d) - 1/3*arctan(d/c))*sin(1/6*pi*sgn(c)*sgn(d) - 1/6*pi*
sgn(d) - 1/3*arctan(d/c)) - c*cos(1/6*pi*sgn(c)*sgn(d) - 1/6*pi*sgn(d) - 1
/3*arctan(d/c))^2 + c*sin(1/6*pi*sgn(c)*sgn(d) - 1/6*pi*sgn(d) - 1/3*arcta
n(d/c))^2)*log((c^2 + d^2)^(1/3)*cos(1/6*pi*sgn(c)*sgn(d) - 1/6*pi*sgn(d)
- 1/3*arctan(d/c))^2 + (c^2 + d^2)^(1/3)*sin(1/6*pi*sgn(c)*sgn(d) - 1/6*pi
*sgn(d) - 1/3*arctan(d/c))^2 + (d*tan(f*x + e) + c)^(2/3) + (d*tan(f*x + e
) + c)^(1/3)*(c - I*d)^(1/3))/((c^2 + d^2)^(2/3)*f) - 1/2*I*(2*sqrt(3)*d*c
os(1/6*pi*sgn(c)*sgn(d) - 1/6*pi*sgn(d) - 1/3*arctan(d/c))*sin(1/6*pi*s...

```

3.477.9 Mupad [B] (verification not implemented)

Time = 22.94 (sec) , antiderivative size = 2982, normalized size of antiderivative = 20.15

$$\int \frac{i - \tan(e + fx)}{\sqrt[3]{c + d \tan(e + fx)}} dx = \text{Too large to display}$$

input `int(-(tan(e + f*x) - 1i)/(c + d*tan(e + f*x))^(1/3),x)`

output

```

log(d^5*f*(c + d*tan(e + f*x))^(1/3)*243i + ((1944*d^4*f^4*(c^2 - d^2)*(c
+ d*tan(e + f*x))^(1/3) - 3*2^(1/3)*c*d^4*f^6*(c^2 + d^2)*(-(11664*f^3*((c
^2*d^12*(c^2 + d^2)^2)/f^6)^(1/2) + d^9*11664i + c^2*d^7*11664i)/(d^6*f^3*
(c^2 + d^2)^2))^(2/3))*((11664*f^3*((c^2*d^12*(c^2 + d^2)^2)/f^6)^(1/2) + d
^9*11664i + c^2*d^7*11664i))/(93312*d^6*f^3*(c^2 + d^2)^2))*(-(f^3*((4*(72
9*d^8 + 729*c^2*d^6)*(46656*d^10 + 93312*c^2*d^8 + 46656*c^4*d^6))/f^6 - (
11664*d^9 + 11664*c^2*d^7)^2/f^6)^(1/2) + d^9*11664i + c^2*d^7*11664i)/(93
312*f^3*(d^10 + 2*c^2*d^8 + c^4*d^6)))^(1/3) + log(d^5*f*(c + d*tan(e + f*
x))^(1/3)*243i + ((1944*d^4*f^4*(c^2 - d^2)*(c + d*tan(e + f*x))^(1/3) - 3
*2^(1/3)*c*d^4*f^6*(c^2 + d^2)*(-(d^9*11664i - 11664*f^3*((c^2*d^12*(c^2 +
d^2)^2)/f^6)^(1/2) + c^2*d^7*11664i)/(d^6*f^3*(c^2 + d^2)^2))^(2/3))*((d^9
*11664i - 11664*f^3*((c^2*d^12*(c^2 + d^2)^2)/f^6)^(1/2) + c^2*d^7*11664i)
)/(93312*d^6*f^3*(c^2 + d^2)^2))*(-(d^9*11664i - f^3*((4*(729*d^8 + 729*c^
2*d^6)*(46656*d^10 + 93312*c^2*d^8 + 46656*c^4*d^6))/f^6 - (11664*d^9 + 11
664*c^2*d^7)^2/f^6)^(1/2) + c^2*d^7*11664i)/(93312*f^3*(d^10 + 2*c^2*d^8 +
c^4*d^6)))^(1/3) + (log(-((-1/(f^3*(c - d*i))))^(2/3)*(((1/(f^3*(c - d*
i))))^(1/3)*((1944*d^4*(c^2 - d^2)*(c + d*tan(e + f*x))^(1/3))/f^2 - 1944*
c*d^4*(-1/(f^3*(c - d*i))))^(2/3)*(c^2 + d^2)))/2 - (972*d^4*(c^2 + d^2))/
f^3)/4 - (243*c*d^4*(c + d*tan(e + f*x))^(1/3))/f^5*(-1/(c*f^3 - d*f^3*i)
i))^(1/3))/2 + log(((7776*c*d^4*(c^2 + d^2)*(-1i/(8*f^3*(c*i - d))))^(2...

```

3.477.
$$\int \frac{i - \tan(e + fx)}{\sqrt[3]{c + d \tan(e + fx)}} dx$$

3.478 $\int \frac{d - c \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx$

3.478.1 Optimal result 4567
 3.478.2 Mathematica [A] (verified) 4568
 3.478.3 Rubi [A] (warning: unable to verify) 4568
 3.478.4 Maple [C] (verified) 4572
 3.478.5 Fricas [B] (verification not implemented) 4572
 3.478.6 Sympy [F] 4574
 3.478.7 Maxima [F] 4574
 3.478.8 Giac [F] 4575
 3.478.9 Mupad [B] (verification not implemented) 4575

3.478.1 Optimal result

Integrand size = 26, antiderivative size = 299

$$\int \frac{d - c \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx = -\frac{1}{4}i\sqrt[3]{c - id}x + \frac{1}{4}i\sqrt[3]{c + id}x$$

$$+ \frac{\sqrt{3}\sqrt[3]{c - id} \arctan\left(\frac{1 + \sqrt[3]{c + d \tan(e + fx)}}{\frac{\sqrt[3]{c - id}}{\sqrt{3}}}\right)}{2f}$$

$$+ \frac{\sqrt{3}\sqrt[3]{c + id} \arctan\left(\frac{1 + \sqrt[3]{c + d \tan(e + fx)}}{\frac{\sqrt[3]{c + id}}{\sqrt{3}}}\right)}{2f} - \frac{\sqrt[3]{c - id} \log(\cos(e + fx))}{4f}$$

$$- \frac{\sqrt[3]{c + id} \log(\cos(e + fx))}{4f} - \frac{3\sqrt[3]{c - id} \log\left(\sqrt[3]{c - id} - \sqrt[3]{c + d \tan(e + fx)}\right)}{4f}$$

$$- \frac{3\sqrt[3]{c + id} \log\left(\sqrt[3]{c + id} - \sqrt[3]{c + d \tan(e + fx)}\right)}{4f}$$

output
$$-1/4*I*(c-I*d)^{(1/3)*x}+1/4*I*(c+I*d)^{(1/3)*x}-1/4*(c-I*d)^{(1/3)*\ln(\cos(f*x+e))}/f-1/4*(c+I*d)^{(1/3)*\ln(\cos(f*x+e))}/f-3/4*(c-I*d)^{(1/3)*\ln((c-I*d)^{(1/3)}-(c+d*\tan(f*x+e))^{(1/3)})}/f-3/4*(c+I*d)^{(1/3)*\ln((c+I*d)^{(1/3)}-(c+d*\tan(f*x+e))^{(1/3)})}/f+1/2*(c-I*d)^{(1/3)*\arctan(1/3*(1+2*(c+d*\tan(f*x+e))^{(1/3)})/(c-I*d)^{(1/3)})}*3^{(1/2)}/f+1/2*(c+I*d)^{(1/3)*\arctan(1/3*(1+2*(c+d*\tan(f*x+e))^{(1/3)})/(c+I*d)^{(1/3)})}*3^{(1/2)}/f$$

3.478.2 Mathematica [A] (verified)

Time = 0.40 (sec) , antiderivative size = 330, normalized size of antiderivative = 1.10

$$\int \frac{d - c \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx = \frac{2\sqrt{3}\sqrt[3]{c - id} \arctan\left(\frac{1 + \sqrt[3]{c + d \tan(e + fx)}}{\sqrt[3]{c - id}}\right) + 2\sqrt{3}\sqrt[3]{c + id} \arctan\left(\frac{1 + \sqrt[3]{c + d \tan(e + fx)}}{\sqrt[3]{c - id}}\right)}{2}$$

input `Integrate[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x])^(2/3),x]`

output
$$(2*\text{Sqrt}[3]*(c - I*d)^{(1/3)*\text{ArcTan}[(1 + (2*(c + d*\text{Tan}[e + f*x])^{(1/3)})/(c - I*d)^{(1/3)})]/\text{Sqrt}[3]} + 2*\text{Sqrt}[3]*(c + I*d)^{(1/3)*\text{ArcTan}[(1 + (2*(c + d*\text{Tan}[e + f*x])^{(1/3)})/(c + I*d)^{(1/3)})]/\text{Sqrt}[3]} - 2*(c - I*d)^{(1/3)*\text{Log}[(c - I*d)^{(1/3)} - (c + d*\text{Tan}[e + f*x])^{(1/3)}]} - 2*(c + I*d)^{(1/3)*\text{Log}[(c + I*d)^{(1/3)} - (c + d*\text{Tan}[e + f*x])^{(1/3)}]} + (c - I*d)^{(1/3)*\text{Log}[(c - I*d)^{(2/3)} + (c - I*d)^{(1/3)*(c + d*\text{Tan}[e + f*x])^{(1/3)}]} + (c + d*\text{Tan}[e + f*x])^{(2/3)}] + (c + I*d)^{(1/3)*\text{Log}[(c + I*d)^{(2/3)} + (c + I*d)^{(1/3)*(c + d*\text{Tan}[e + f*x])^{(1/3)}]} + (c + d*\text{Tan}[e + f*x])^{(2/3)}})/(4*f)$$

3.478.3 Rubi [A] (warning: unable to verify)

Time = 0.59 (sec) , antiderivative size = 235, normalized size of antiderivative = 0.79, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.346$, Rules used = {3042, 4022, 3042, 4020, 25, 69, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{d - c \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx$$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & \int \frac{d - c \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx \\
 & \downarrow \text{4022} \\
 & \frac{1}{2}(d + ic) \int \frac{i \tan(e + fx) + 1}{(c + d \tan(e + fx))^{2/3}} dx - \frac{1}{2}(-d + ic) \int \frac{1 - i \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx \\
 & \downarrow \text{3042} \\
 & \frac{1}{2}(d + ic) \int \frac{i \tan(e + fx) + 1}{(c + d \tan(e + fx))^{2/3}} dx - \frac{1}{2}(-d + ic) \int \frac{1 - i \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx \\
 & \downarrow \text{4020} \\
 & \frac{i(d + ic) \int -\frac{1}{(1 - i \tan(e + fx))(c + d \tan(e + fx))^{2/3}} d(i \tan(e + fx))}{2f} + \\
 & \frac{i(-d + ic) \int -\frac{1}{(i \tan(e + fx) + 1)(c + d \tan(e + fx))^{2/3}} d(-i \tan(e + fx))}{2f} \\
 & \downarrow \text{25} \\
 & \frac{i(d + ic) \int \frac{1}{(1 - i \tan(e + fx))(c + d \tan(e + fx))^{2/3}} d(i \tan(e + fx))}{2f} - \\
 & \frac{i(-d + ic) \int \frac{1}{(i \tan(e + fx) + 1)(c + d \tan(e + fx))^{2/3}} d(-i \tan(e + fx))}{2f} \\
 & \downarrow \text{69} \\
 & i(d + ic) \left(-\frac{3 \int \frac{1}{-\tan^2(e + fx) + (c - id)^{2/3} + \sqrt[3]{c - id} \sqrt[3]{c + d \tan(e + fx)}} d^3 \sqrt[3]{c + d \tan(e + fx)}}{2 \sqrt[3]{c - id}} - \frac{3 \int \frac{1}{\sqrt[3]{c - id} - i \tan(e + fx)} d^3 \sqrt[3]{c + d \tan(e + fx)}}{2(c - id)^{2/3}} \right) \\
 & \hline
 & i(-d + ic) \left(-\frac{3 \int \frac{1}{-\tan^2(e + fx) + (c + id)^{2/3} + \sqrt[3]{c + id} \sqrt[3]{c + d \tan(e + fx)}} d^3 \sqrt[3]{c + d \tan(e + fx)}}{2 \sqrt[3]{c + id}} - \frac{3 \int \frac{1}{i \tan(e + fx) + \sqrt[3]{c + id}} d^3 \sqrt[3]{c + d \tan(e + fx)}}{2(c + id)^{2/3}} \right) \\
 & \hline
 & \qquad \qquad \qquad 2f \\
 & \downarrow \text{16}
 \end{aligned}$$

3.478. $\int \frac{d - c \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx$

$$i(d + ic) \left(-\frac{3 \int \frac{1}{-\tan^2(e+fx) + (c-id)^{2/3} + \sqrt[3]{c-id} \sqrt[3]{c+d \tan(e+fx)}} d \sqrt[3]{c+d \tan(e+fx)}}{2 \sqrt[3]{c-id}} - \frac{\log(1-i \tan(e+fx))}{2(c-id)^{2/3}} + \frac{3 \log(\sqrt[3]{c-id})}{2(c-id)^{2/3}} \right)$$

$$i(-d + ic) \left(-\frac{3 \int \frac{1}{-\tan^2(e+fx) + (c+id)^{2/3} + \sqrt[3]{c+id} \sqrt[3]{c+d \tan(e+fx)}} d \sqrt[3]{c+d \tan(e+fx)}}{2 \sqrt[3]{c+id}} - \frac{\log(1+i \tan(e+fx))}{2(c+id)^{2/3}} + \frac{3 \log(\sqrt[3]{c+id})}{2(c+id)^{2/3}} \right)$$

$2f$

↓ 1082

$$i(d + ic) \left(\frac{3 \int \frac{1}{\tan^2(e+fx) - 3} d \left(\frac{2i \tan(e+fx)}{\sqrt[3]{c-id}} + 1 \right)}{(c-id)^{2/3}} - \frac{\log(1-i \tan(e+fx))}{2(c-id)^{2/3}} + \frac{3 \log(\sqrt[3]{c-id} - i \tan(e+fx))}{2(c-id)^{2/3}} \right)$$

$2f$

$$i(-d + ic) \left(\frac{3 \int \frac{1}{\tan^2(e+fx) - 3} d \left(1 - \frac{2i \tan(e+fx)}{\sqrt[3]{c+id}} \right)}{(c+id)^{2/3}} - \frac{\log(1+i \tan(e+fx))}{2(c+id)^{2/3}} + \frac{3 \log(\sqrt[3]{c+id} + i \tan(e+fx))}{2(c+id)^{2/3}} \right)$$

$2f$

↓ 217

$$i(d + ic) \left(-\frac{i\sqrt{3} \operatorname{arctanh}\left(\frac{\tan(e+fx)}{\sqrt{3}}\right)}{(c-id)^{2/3}} - \frac{\log(1-i \tan(e+fx))}{2(c-id)^{2/3}} + \frac{3 \log(\sqrt[3]{c-id} - i \tan(e+fx))}{2(c-id)^{2/3}} \right)$$

$2f$

$$i(-d + ic) \left(\frac{i\sqrt{3} \operatorname{arctanh}\left(\frac{\tan(e+fx)}{\sqrt{3}}\right)}{(c+id)^{2/3}} - \frac{\log(1+i \tan(e+fx))}{2(c+id)^{2/3}} + \frac{3 \log(\sqrt[3]{c+id} + i \tan(e+fx))}{2(c+id)^{2/3}} \right)$$

$2f$

input `Int[(d - c*Tan[e + f*x])/(c + d*Tan[e + f*x])^(2/3),x]`

output `((I/2)*(I*c + d)*(((-I)*Sqrt[3]*ArcTanh[Tan[e + f*x]/Sqrt[3]])/(c - I*d)^(2/3) - Log[1 - I*Tan[e + f*x]]/(2*(c - I*d)^(2/3)) + (3*Log[(c - I*d)^(1/3) - I*Tan[e + f*x]])/(2*(c - I*d)^(2/3)))/f + ((I/2)*(I*c - d)*((I*Sqrt[3]*ArcTanh[Tan[e + f*x]/Sqrt[3]])/(c + I*d)^(2/3) - Log[1 + I*Tan[e + f*x]]/(2*(c + I*d)^(2/3)) + (3*Log[(c + I*d)^(1/3) + I*Tan[e + f*x]])/(2*(c + I*d)^(2/3)))/f`

3.478.3.1 Defintions of rubi rules used

- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 69 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(2/3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q^2), x] + (-Simp[3/(2*b*q) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Simp[3/(2*b*q^2) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x])] /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

3.478.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.46 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.24

method	result
derivativedivides	$-\frac{\sum_{R=\text{RootOf}(-Z^6-2cZ^3+c^2+d^2)} \frac{(-R^3 c - c^2 - d^2) \ln\left((c+d \tan(fx+e))^{\frac{1}{3}} - R\right)}{-R^5 - R^2 c}}{2f}$
default	$-\frac{\sum_{R=\text{RootOf}(-Z^6-2cZ^3+c^2+d^2)} \frac{(-R^3 c - c^2 - d^2) \ln\left((c+d \tan(fx+e))^{\frac{1}{3}} - R\right)}{-R^5 - R^2 c}}{2f}$
parts	$\frac{d^2 \left(\sum_{R=\text{RootOf}(-Z^6-2cZ^3+c^2+d^2)} \frac{\ln\left((c+d \tan(fx+e))^{\frac{1}{3}} - R\right)}{-R^2 (-R^3 - c)} \right)}{2f} - \frac{c \left(\sum_{R=\text{RootOf}(-Z^6-2cZ^3+c^2+d^2)} \frac{\ln\left((c+d \tan(fx+e))^{\frac{1}{3}} - R\right)}{-R^2 (-R^3 - c)} \right)}{2f}$

input `int((d-c*tan(f*x+e))/(c+d*tan(f*x+e))^(2/3),x,method=_RETURNVERBOSE)`

output `-1/2/f*sum((_R^3*c-c^2-d^2)/(_R^5-_R^2*c)*ln((c+d*tan(f*x+e))^(1/3)-_R),_R=RootOf(_Z^6-2*_Z^3+c^2+d^2))`

3.478.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 443 vs. 2(217) = 434.

3.478. $\int \frac{d-c \tan(e+fx)}{(c+d \tan(e+fx))^{2/3}} dx$

Time = 0.25 (sec) , antiderivative size = 443, normalized size of antiderivative = 1.48

$$\begin{aligned}
 & \int \frac{d - c \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx = \\
 & -\frac{1}{4} (\sqrt{-3} + 1) \left(-\frac{f^3 \sqrt{-\frac{d^2}{f^6}} + c}{f^3} \right)^{\frac{1}{3}} \log \left(-\frac{1}{2} (\sqrt{-3}f + f) \left(-\frac{f^3 \sqrt{-\frac{d^2}{f^6}} + c}{f^3} \right)^{\frac{1}{3}} \right. \\
 & \left. + (d \tan(fx + e) + c)^{\frac{1}{3}} \right) \\
 & + \frac{1}{4} (\sqrt{-3} - 1) \left(-\frac{f^3 \sqrt{-\frac{d^2}{f^6}} + c}{f^3} \right)^{\frac{1}{3}} \log \left(\frac{1}{2} (\sqrt{-3}f - f) \left(-\frac{f^3 \sqrt{-\frac{d^2}{f^6}} + c}{f^3} \right)^{\frac{1}{3}} \right. \\
 & \left. + (d \tan(fx + e) + c)^{\frac{1}{3}} \right) \\
 & -\frac{1}{4} (\sqrt{-3} + 1) \left(\frac{f^3 \sqrt{-\frac{d^2}{f^6}} - c}{f^3} \right)^{\frac{1}{3}} \log \left(-\frac{1}{2} (\sqrt{-3}f + f) \left(\frac{f^3 \sqrt{-\frac{d^2}{f^6}} - c}{f^3} \right)^{\frac{1}{3}} \right. \\
 & \left. + (d \tan(fx + e) + c)^{\frac{1}{3}} \right) \\
 & + \frac{1}{4} (\sqrt{-3} - 1) \left(\frac{f^3 \sqrt{-\frac{d^2}{f^6}} - c}{f^3} \right)^{\frac{1}{3}} \log \left(\frac{1}{2} (\sqrt{-3}f - f) \left(\frac{f^3 \sqrt{-\frac{d^2}{f^6}} - c}{f^3} \right)^{\frac{1}{3}} \right. \\
 & \left. + (d \tan(fx + e) + c)^{\frac{1}{3}} \right) \\
 & + \frac{1}{2} \left(-\frac{f^3 \sqrt{-\frac{d^2}{f^6}} + c}{f^3} \right)^{\frac{1}{3}} \log \left(f \left(-\frac{f^3 \sqrt{-\frac{d^2}{f^6}} + c}{f^3} \right)^{\frac{1}{3}} + (d \tan(fx + e) + c)^{\frac{1}{3}} \right) \\
 & + \frac{1}{2} \left(\frac{f^3 \sqrt{-\frac{d^2}{f^6}} - c}{f^3} \right)^{\frac{1}{3}} \log \left(f \left(\frac{f^3 \sqrt{-\frac{d^2}{f^6}} - c}{f^3} \right)^{\frac{1}{3}} + (d \tan(fx + e) + c)^{\frac{1}{3}} \right)
 \end{aligned}$$

3.478. $\int \frac{d - c \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx$

input `integrate((d-c*tan(f*x+e))/(c+d*tan(f*x+e))^(2/3),x, algorithm="fricas")`

output `-1/4*(sqrt(-3) + 1)*(-(f^3*sqrt(-d^2/f^6) + c)/f^3)^(1/3)*log(-1/2*(sqrt(-3)*f + f)*(-(f^3*sqrt(-d^2/f^6) + c)/f^3)^(1/3) + (d*tan(f*x + e) + c)^(1/3)) + 1/4*(sqrt(-3) - 1)*(-(f^3*sqrt(-d^2/f^6) + c)/f^3)^(1/3)*log(1/2*(sqrt(-3)*f - f)*(-(f^3*sqrt(-d^2/f^6) + c)/f^3)^(1/3) + (d*tan(f*x + e) + c)^(1/3)) - 1/4*(sqrt(-3) + 1)*((f^3*sqrt(-d^2/f^6) - c)/f^3)^(1/3)*log(-1/2*(sqrt(-3)*f + f)*((f^3*sqrt(-d^2/f^6) - c)/f^3)^(1/3) + (d*tan(f*x + e) + c)^(1/3)) + 1/4*(sqrt(-3) - 1)*((f^3*sqrt(-d^2/f^6) - c)/f^3)^(1/3)*log(1/2*(sqrt(-3)*f - f)*((f^3*sqrt(-d^2/f^6) - c)/f^3)^(1/3) + (d*tan(f*x + e) + c)^(1/3)) + 1/2*(-(f^3*sqrt(-d^2/f^6) + c)/f^3)^(1/3)*log(f*(-(f^3*sqrt(-d^2/f^6) + c)/f^3)^(1/3) + (d*tan(f*x + e) + c)^(1/3)) + 1/2*((f^3*sqrt(-d^2/f^6) - c)/f^3)^(1/3)*log(f*((f^3*sqrt(-d^2/f^6) - c)/f^3)^(1/3) + (d*tan(f*x + e) + c)^(1/3))`

3.478.6 Sympy [F]

$$\int \frac{d - c \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx = - \int \left(- \frac{d}{(c + d \tan(e + fx))^{2/3}} \right) dx - \int \frac{c \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx$$

input `integrate((d-c*tan(f*x+e))/(c+d*tan(f*x+e))**(2/3),x)`

output `-Integral(-d/(c + d*tan(e + f*x))**(2/3), x) - Integral(c*tan(e + f*x)/(c + d*tan(e + f*x))**(2/3), x)`

3.478.7 Maxima [F]

$$\int \frac{d - c \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx = \int - \frac{c \tan(fx + e) - d}{(d \tan(fx + e) + c)^{2/3}} dx$$

input `integrate((d-c*tan(f*x+e))/(c+d*tan(f*x+e))^(2/3),x, algorithm="maxima")`

output `-integrate((c*tan(f*x + e) - d)/(d*tan(f*x + e) + c)^(2/3), x)`

3.478.8 Giac [F]

$$\int \frac{d - c \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx = \int -\frac{c \tan(fx + e) - d}{(d \tan(fx + e) + c)^{2/3}} dx$$

input `integrate((d-c*tan(f*x+e))/(c+d*tan(f*x+e))^(2/3),x, algorithm="giac")`

output `undef`

3.478.9 Mupad [B] (verification not implemented)

Time = 23.06 (sec) , antiderivative size = 4308, normalized size of antiderivative = 14.41

$$\int \frac{d - c \tan(e + fx)}{(c + d \tan(e + fx))^{2/3}} dx = \text{Too large to display}$$

input `int((d - c*tan(e + f*x))/(c + d*tan(e + f*x))^(2/3),x)`

output `log((((1944*c*d^4*(c^2 + d^2)*((8*(-d^6*f^6*(c^2 - d^2)^2)^(1/2) + 16*c*d^4*f^3)/(f^6*(c^2 + d^2)^2))^(1/3) + (7776*c*d^6*(c + d*tan(e + f*x))^(1/3))/f)*((8*(-d^6*f^6*(c^2 - d^2)^2)^(1/2) + 16*c*d^4*f^3)/(f^6*(c^2 + d^2)^2))^(2/3))/16 - (972*d^8)/f^3)*((8*(-d^6*f^6*(c^2 - d^2)^2)^(1/2) + 16*c*d^4*f^3)/(f^6*(c^2 + d^2)^2))^(1/3))/4 - (486*d^8*(c + d*tan(e + f*x))^(1/3))/f^4)*(((256*c^2*d^8*f^6 - d^6*(64*c^4*f^6 + 64*d^4*f^6 + 128*c^2*d^2*f^6))^(1/2) + 16*c*d^4*f^3)/(64*(c^4*f^6 + d^4*f^6 + 2*c^2*d^2*f^6)))^(1/3) + log((((1944*c*d^4*(c^2 + d^2)*(-8*(-d^6*f^6*(c^2 - d^2)^2)^(1/2) - 16*c*d^4*f^3)/(f^6*(c^2 + d^2)^2))^(1/3) + (7776*c*d^6*(c + d*tan(e + f*x))^(1/3))/f)*(-8*(-d^6*f^6*(c^2 - d^2)^2)^(1/2) - 16*c*d^4*f^3)/(f^6*(c^2 + d^2)^2))^(2/3))/16 - (972*d^8)/f^3)*(-8*(-d^6*f^6*(c^2 - d^2)^2)^(1/2) - 16*c*d^4*f^3)/(f^6*(c^2 + d^2)^2))^(1/3))/4 - (486*d^8*(c + d*tan(e + f*x))^(1/3))/f^4)*(-((256*c^2*d^8*f^6 - d^6*(64*c^4*f^6 + 64*d^4*f^6 + 128*c^2*d^2*f^6))^(1/2) - 16*c*d^4*f^3)/(64*(c^4*f^6 + d^4*f^6 + 2*c^2*d^2*f^6)))^(1/3) + log(- (486*c^4*d^4*(c + d*tan(e + f*x))^(1/3))/f^4 - (((16*(-c^8*d^2*f^6)^(1/2) - 8*c^5*f^3 + 8*c^3*d^2*f^3)/(f^6*(c^2 + d^2)^2))^(1/3)*(1944*c*d^4*(-c^8*d^2*f^6)^(1/2) + 1944*c^4*d^6*f^3 + 243*c^5*d^4*f^5*(c + d*tan(e + f*x))^(1/3))*((16*(-c^8*d^2*f^6)^(1/2) - 8*c^5*f^3 + 8*c^3*d^2*f^3)/(f^6*(c^2 + d^2)^2))^(2/3) - 243*c*d^8*f^5*(c + d*tan(e + f*x))^(1/3))*((16*(-c^8*d^2*f^6)^(1/2) - 8*c^5*f^3 + 8*c^3*d^2*f^3)/(f^6*(c^2 + d^2)^2))...`

3.479 $\int \tan^m(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

3.479.1 Optimal result	4576
3.479.2 Mathematica [A] (verified)	4577
3.479.3 Rubi [A] (verified)	4578
3.479.4 Maple [F]	4582
3.479.5 Fricas [F]	4582
3.479.6 Sympy [F]	4583
3.479.7 Maxima [F]	4583
3.479.8 Giac [F]	4584
3.479.9 Mupad [F(-1)]	4584

3.479.1 Optimal result

Integrand size = 31, antiderivative size = 403

$$\int \tan^m(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx =$$

$$-\frac{b(Ab^3(12+7m+m^2)+4ab^2B(12+7m+m^2)-2a^3B(19+8m+m^2)-a^2Ab(68+37m+5m^2)) \tan^{1+m}(c+dx)}{d(1+m)(3+m)(4+m)}$$

$$+\frac{(a^4A-6a^2Ab^2+Ab^4-4a^3bB+4ab^3B) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right) \tan^{1+m}(c+dx)}{d(1+m)}$$

$$+\frac{b^2(2aAb(4+m)^2-b^2B(12+7m+m^2)+a^2B(26+9m+m^2)) \tan^{2+m}(c+dx)}{d(2+m)(3+m)(4+m)}$$

$$+\frac{(4a^3Ab-4aAb^3+a^4B-6a^2b^2B+b^4B) \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, -\tan^2(c+dx)\right) \tan^{2+m}(c+dx)}{d(2+m)}$$

$$+\frac{b(Ab(4+m)+aB(7+m)) \tan^{1+m}(c+dx)(a+b \tan(c+dx))^2}{d(3+m)(4+m)}$$

$$+\frac{bB \tan^{1+m}(c+dx)(a+b \tan(c+dx))^3}{d(4+m)}$$

output
$$-b*(A*b^3*(m^2+7*m+12)+4*a*b^2*B*(m^2+7*m+12)-2*a^3*B*(m^2+8*m+19)-a^2*A*b*(5*m^2+37*m+68))*\tan(d*x+c)^{(1+m)}/d/(4+m)/(m^2+4*m+3)+(A*a^4-6*A*a^2*b^2+A*b^4-4*B*a^3*b+4*B*a*b^3)*\text{hypergeom}([1, 1/2+1/2*m], [3/2+1/2*m], -\tan(d*x+c)^2)*\tan(d*x+c)^{(1+m)}/d/(1+m)+b^2*(2*a*A*b*(4+m)^2-b^2*B*(m^2+7*m+12)+a^2*B*(m^2+9*m+26))*\tan(d*x+c)^{(2+m)}/d/(2+m)/(3+m)/(4+m)+(4*A*a^3*b-4*A*a*b^3+B*a^4-6*B*a^2*b^2+B*b^4)*\text{hypergeom}([1, 1+1/2*m], [2+1/2*m], -\tan(d*x+c)^2)*\tan(d*x+c)^{(2+m)}/d/(2+m)+b*(A*b*(4+m)+a*B*(7+m))*\tan(d*x+c)^{(1+m)*(a+b*\tan(d*x+c))^2}/d/(3+m)/(4+m)+b*B*\tan(d*x+c)^{(1+m)*(a+b*\tan(d*x+c))^3}/d/(4+m)$$

3.479.2 Mathematica [A] (verified)

Time = 4.83 (sec) , antiderivative size = 355, normalized size of antiderivative = 0.88

$$\int \tan^m(c+dx)(a+b\tan(c+dx))^4(A+B\tan(c+dx)) dx$$

$$= \frac{\tan^{1+m}(c+dx) \left(-b(2+m) \left(Ab^3(12+7m+m^2) + 4ab^2B(12+7m+m^2) - 2a^3B(19+8m+m^2) - a^2A \right) \right)}{d(1+m)(2+m)(3+m)(4+m)}$$

input `Integrate[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output
$$\begin{aligned} & (\tan[c + d*x]^{(1+m)} * (-b*(2+m)*(A*b^3*(12+7*m+m^2) + 4*a*b^2*B*(12+7*m+m^2) - 2*a^3*B*(19+8*m+m^2) - a^2*A*b*(68+37*m+5*m^2))) + \\ & (a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*a^3*b*B + 4*a*b^3*B)*(2+m)*(3+m)*(4+m)*\text{Hypergeometric2F1}[1, (1+m)/2, (3+m)/2, -\tan[c + d*x]^2] + b^2*(1+m)* \\ & (2*a*A*b*(4+m)^2 - b^2*B*(12+7*m+m^2) + a^2*B*(26+9*m+m^2))*\tan[c + d*x] + \\ & (4*a^3*A*b - 4*a*A*b^3 + a^4*B - 6*a^2*b^2*B + b^4*B)*(1+m)*(3+m)*(4+m)*\text{Hypergeometric2F1}[1, (2+m)/2, (4+m)/2, -\tan[c + d*x]^2]* \\ & \tan[c + d*x] + b*(1+m)*(2+m)*(A*b*(4+m) + a*B*(7+m))*(a + b*\tan[c + d*x])^2 + \\ & b*B*(1+m)*(2+m)*(3+m)*(a + b*\tan[c + d*x])^3) / (d*(1+m)*(2+m)*(3+m)*(4+m)) \end{aligned}$$

3.479.3 Rubi [A] (verified)

Time = 2.44 (sec) , antiderivative size = 416, normalized size of antiderivative = 1.03, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 4090, 25, 3042, 4130, 3042, 4120, 25, 3042, 4113, 3042, 4021, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^m(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c+dx)^m(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4090} \\
 & \frac{\int -\tan^m(c+dx)(a+b \tan(c+dx))^2(-b(Ab(m+4)+aB(m+7)) \tan^2(c+dx) - (Ba^2+2Aba-b^2B)(m+4) \tan(c+dx) + bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^3) dx}{m+4} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \tan^m(c+dx)(a+b \tan(c+dx))^2(-b(Ab(m+4)+aB(m+7)) \tan^2(c+dx) - (Ba^2+2Aba-b^2B)(m+4) \tan(c+dx) + bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^3) dx}{m+4} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \tan(c+dx)^m(a+b \tan(c+dx))^2(-b(Ab(m+4)+aB(m+7)) \tan(c+dx)^2 - (Ba^2+2Aba-b^2B)(m+4) \tan(c+dx) + bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^3) dx}{m+4} \\
 & \quad \downarrow \text{4130} \\
 & \frac{\int \tan^m(c+dx)(a+b \tan(c+dx))(-b(B(m^2+9m+26)a^2+2Ab(m+4)^2a-b^2B(m^2+7m+12)) \tan^2(c+dx) - (Ba^3+3Aba^2-3b^2Ba-Ab^3)(m+3)(m+4) \tan(c+dx) + bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^3) dx}{m+3} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.479. $\int \tan^m(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

$$\frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^3}{d(m+4)} - \frac{\int \tan(c+dx)^m (a+b \tan(c+dx)) (-b(B(m^2+9m+26)a^2+2Ab(m+4)^2a-b^2B(m^2+7m+12)) \tan(c+dx)^2 - (Ba^3+3Aba^2-3b^2Ba-Ab^3)(m+3)(m+4))}{m+3}$$

4120

$$\frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^3}{d(m+4)} - \frac{\int -\tan^m(c+dx) ((m+2)(a(m+3)(bB(m+1)-aA(m+4))+b(m+1)(Ab(m+4)+aB(m+7)))a^2+b(m+2)(-2B(m^2+8m+19)a^3-Ab(5m^2+37m+68)a^2+4b^2B(m^2+7m+6))}{m+2}$$

25

$$\frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^3}{d(m+4)} - \frac{\int \tan^m(c+dx) ((m+2)(a(m+3)(bB(m+1)-aA(m+4))+b(m+1)(Ab(m+4)+aB(m+7)))a^2+b(m+2)(-2B(m^2+8m+19)a^3-Ab(5m^2+37m+68)a^2+4b^2B(m^2+7m+6))}{m+2}$$

3042

$$\frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^3}{d(m+4)} - \frac{\int \tan(c+dx)^m ((m+2)(a(m+3)(bB(m+1)-aA(m+4))+b(m+1)(Ab(m+4)+aB(m+7)))a^2+b(m+2)(-2B(m^2+8m+19)a^3-Ab(5m^2+37m+68)a^2+4b^2B(m^2+7m+6))}{m+2}$$

4113

$$\frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^3}{d(m+4)} - \frac{\int \tan^m(c+dx) (-((Aa^4-4bBa^3-6Ab^2a^2+4b^3Ba+Ab^4)(m+2)(m+3)(m+4)) - (Ba^4+4Aba^3-6b^2Ba^2-4Ab^3a+b^4B)(m+2)(m+3) \tan(c+dx)(m+4)) dx + \frac{b(m+2)}{m+2}}$$

3042

$$\frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^3}{d(m+4)} - \frac{\int \tan(c+dx)^m (-((Aa^4-4bBa^3-6Ab^2a^2+4b^3Ba+Ab^4)(m+2)(m+3)(m+4)) - (Ba^4+4Aba^3-6b^2Ba^2-4Ab^3a+b^4B)(m+2)(m+3) \tan(c+dx)(m+4)) dx + \frac{b(m+2)}{m+2}}$$

4021

3.479. $\int \tan^m(c+dx)(a+b \tan(c+dx))^4(A+B \tan(c+dx)) dx$

$$\frac{bB \tan^{m+1}(c + dx)(a + b \tan(c + dx))^3}{d(m + 4)} - \frac{-(m+2)(m+3)(m+4)(a^4 B + 4a^3 b B - 6a^2 b^2 B - 4a b^3 + b^4 B) \int \tan^{m+1}(c+dx) dx - (m+2)(m+3)(m+4)(a^4 A - 4a^3 b B - 6a^2 b^2 B + 4a b^3 B + Ab^4) \int \tan^m(c+dx) dx + \frac{b}{m+2}}$$

↓ 3042

$$\frac{bB \tan^{m+1}(c + dx)(a + b \tan(c + dx))^3}{d(m + 4)} - \frac{-(m+2)(m+3)(m+4)(a^4 A - 4a^3 b B - 6a^2 b^2 B + 4a b^3 B + Ab^4) \int \tan(c+dx)^m dx - (m+2)(m+3)(m+4)(a^4 B + 4a^3 Ab - 6a^2 b^2 B - 4a Ab^3 + b^4 B) \int \tan(c+dx)^{m+1} dx + \frac{b}{m+2}}$$

↓ 3957

$$\frac{bB \tan^{m+1}(c + dx)(a + b \tan(c + dx))^3}{d(m + 4)} - \frac{(m+2)(m+3)(m+4)(a^4 B + 4a^3 Ab - 6a^2 b^2 B - 4a Ab^3 + b^4 B) \int \frac{\tan^{m+1}(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx) - (m+2)(m+3)(m+4)(a^4 A - 4a^3 b B - 6a^2 b^2 B + 4a b^3 B + Ab^4) \int \frac{\tan^m(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} - \frac{b}{m+2}$$

↓ 278

$$\frac{bB \tan^{m+1}(c + dx)(a + b \tan(c + dx))^3}{d(m + 4)} - \frac{b(m+2)(-2a^3 B(m^2 + 8m + 19) - a^2 Ab(5m^2 + 37m + 68) + 4ab^2 B(m^2 + 7m + 12) + Ab^3(m^2 + 7m + 12)) \tan^{m+1}(c+dx) - (m+2)(m+3)(m+4)(a^4 A - 4a^3 b B - 6a^2 b^2 B + 4a b^3 B + Ab^4) \int \tan^m(c+dx) dx}{d(m+1)}$$

input `Int[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^4*(A + B*Tan[c + d*x]),x]`

output `(b*B*Tan[c + d*x]^(1 + m)*(a + b*Tan[c + d*x])^3)/(d*(4 + m)) - ((b*(A*b*(4 + m) + a*B*(7 + m))*Tan[c + d*x]^(1 + m)*(a + b*Tan[c + d*x])^2)/(d*(3 + m))) + ((b^2*(2*a*A*b*(4 + m)^2 - b^2*B*(12 + 7*m + m^2) + a^2*B*(26 + 9*m + m^2))*Tan[c + d*x]^(2 + m))/(d*(2 + m))) + ((b*(2 + m)*(A*b^3*(12 + 7*m + m^2) + 4*a*b^2*B*(12 + 7*m + m^2) - 2*a^3*B*(19 + 8*m + m^2) - a^2*A*b*(68 + 37*m + 5*m^2))*Tan[c + d*x]^(1 + m))/(d*(1 + m)) - ((a^4*A - 6*a^2*A*b^2 + A*b^4 - 4*a^3*b*B + 4*a*b^3*B)*(2 + m)*(3 + m)*(4 + m)*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(1 + m))/(d*(1 + m)) - (((a^3*A*b - 4*a*A*b^3 + a^4*B - 6*a^2*b^2*B + b^4*B)*(3 + m)*(4 + m)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(2 + m))/d)/(2 + m))/(3 + m))/(4 + m)`

3.479. $\int \tan^m(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx$

3.479.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`
- rule 4021 `Int[((b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Tan[e + f*x])^m, x], x] + Simp[d/b Int[(b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x] && NeQ[c^2 + d^2, 0] && !IntegerQ[2*m]`
- rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`
- rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4120 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f_)*(x_)])^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]`

rule 4130 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f_)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.479.4 Maple [F]

$$\int \tan(dx + c)^m (a + b \tan(dx + c))^4 (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^m*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x)`

3.479.5 Fracas [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^4 \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*b^4*tan(d*x + c)^5 + A*a^4 + (4*B*a*b^3 + A*b^4)*tan(d*x + c)^4 + 2*(3*B*a^2*b^2 + 2*A*a*b^3)*tan(d*x + c)^3 + 2*(2*B*a^3*b + 3*A*a^2*b^2)*tan(d*x + c)^2 + (B*a^4 + 4*A*a^3*b)*tan(d*x + c))*tan(d*x + c)^m, x)`

3.479.6 Sympy [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx \\ &= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^4 \tan^m(c + dx) dx \end{aligned}$$

input `integrate(tan(d*x+c)**m*(a+b*tan(d*x+c))**4*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**4*tan(c + d*x)**m, x)`

3.479.7 Maxima [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^4(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^4 \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^4*tan(d*x + c)^m, x)`

3.479.8 Giac [F]

$$\begin{aligned} & \int \tan^m(c+dx)(a+b\tan(c+dx))^4(A+B\tan(c+dx)) dx \\ &= \int (B\tan(dx+c)+A)(b\tan(dx+c)+a)^4 \tan(dx+c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^4*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^4*tan(d*x + c)^m, x)`

3.479.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \tan^m(c+dx)(a+b\tan(c+dx))^4(A+B\tan(c+dx)) dx \\ &= \int \tan(c+dx)^m (A+B\tan(c+dx)) (a+b\tan(c+dx))^4 dx \end{aligned}$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^4,x)`

output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^4, x)`

3.480 $\int \tan^m(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.480.1 Optimal result	4585
3.480.2 Mathematica [A] (verified)	4586
3.480.3 Rubi [A] (verified)	4586
3.480.4 Maple [F]	4590
3.480.5 Fricas [F]	4590
3.480.6 Sympy [F]	4591
3.480.7 Maxima [F]	4591
3.480.8 Giac [F]	4591
3.480.9 Mupad [F(-1)]	4592

3.480.1 Optimal result

Integrand size = 31, antiderivative size = 267

$$\int \tan^m(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{b(3aAb(3+m) - b^2B(3+m) + 2a^2B(4+m)) \tan^{1+m}(c+dx)}{d(1+m)(3+m)}$$

$$+ \frac{(a^3A - 3aAb^2 - 3a^2bB + b^3B) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right) \tan^{1+m}(c+dx)}{d(1+m)}$$

$$+ \frac{b^2(Ab(3+m) + aB(5+m)) \tan^{2+m}(c+dx)}{d(2+m)(3+m)}$$

$$+ \frac{(3a^2Ab - Ab^3 + a^3B - 3ab^2B) \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, -\tan^2(c+dx)\right) \tan^{2+m}(c+dx)}{d(2+m)}$$

$$+ \frac{bB \tan^{1+m}(c+dx)(a+b \tan(c+dx))^2}{d(3+m)}$$

output

```

b*(3*a*A*b*(3+m)-b^2*B*(3+m)+2*a^2*B*(4+m))*tan(d*x+c)^(1+m)/d/(1+m)/(3+m)
+(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*hypergeom([1, 1/2+1/2*m],[3/2+1/2*m],-tan(d*x+c)^2)*tan(d*x+c)^(1+m)/d/(1+m)+b^2*(A*b*(3+m)+a*B*(5+m))*tan(d*x+c)^(2+m)/d/(2+m)/(3+m)+(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)*hypergeom([1, 1+1/2*m],[2+1/2*m],-tan(d*x+c)^2)*tan(d*x+c)^(2+m)/d/(2+m)+b*B*tan(d*x+c)^(1+m)*(a+b*tan(d*x+c))^2/d/(3+m)

```


3.480.2 Mathematica [A] (verified)

Time = 2.52 (sec) , antiderivative size = 232, normalized size of antiderivative = 0.87

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{\tan^{1+m}(c + dx) (b(2 + m) (3aAb(3 + m) - b^2B(3 + m) + 2a^2B(4 + m)) + (a^3A - 3aAb^2 - 3a^2bB + b^3B))}{d(1 + m)(2 + m)(3 + m)}$$

input `Integrate[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`output `(Tan[c + d*x]^(1 + m)*(b*(2 + m)*(3*a*A*b*(3 + m) - b^2*B*(3 + m) + 2*a^2*B*(4 + m)) + (a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*(2 + m)*(3 + m)*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2] + b^2*(1 + m)*(A*b*(3 + m) + a*B*(5 + m))*Tan[c + d*x] + (3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*(1 + m)*(3 + m)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x] + b*B*(1 + m)*(2 + m)*(a + b*Tan[c + d*x])^2)/(d*(1 + m)*(2 + m)*(3 + m))`**3.480.3 Rubi [A] (verified)**Time = 1.43 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.06, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3042, 4090, 25, 3042, 4120, 25, 3042, 4113, 3042, 4021, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^m(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4090}$$

$$\frac{\int -\tan^m(c + dx)(a + b \tan(c + dx)) (-b(Ab(m + 3) + aB(m + 5)) \tan^2(c + dx) - (Ba^2 + 2Aba - b^2B) (m + 3))}{d(m + 3)} + \frac{bB \tan^{m+1}(c + dx)(a + b \tan(c + dx))^2}{m + 3}$$

$$3.480. \quad \int \tan^m(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$\begin{aligned} & \downarrow 25 \\ & \frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^2}{d(m+3)} - \\ & \frac{\int \tan^m(c+dx)(a+b \tan(c+dx))(-b(Ab(m+3)+aB(m+5)) \tan^2(c+dx) - (Ba^2+2Aba-b^2B)(m+3) \tan(c+dx))}{m+3} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^2}{d(m+3)} - \\ & \frac{\int \tan(c+dx)^m(a+b \tan(c+dx))(-b(Ab(m+3)+aB(m+5)) \tan(c+dx)^2 - (Ba^2+2Aba-b^2B)(m+3) \tan(c+dx))}{m+3} \end{aligned}$$

$$\begin{aligned} & \downarrow 4120 \\ & \frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^2}{d(m+3)} - \\ & \frac{\int -\tan^m(c+dx)((m+2)(bB(m+1)-aA(m+3))a^2-b(m+2)(2B(m+4)a^2+3Ab(m+3)a-b^2B(m+3)) \tan^2(c+dx) - (Ba^3+3Aba^2-3b^2Ba-Ab^3)(m+2))}{m+2} \\ & \frac{}{m+3} \end{aligned}$$

$$\begin{aligned} & \downarrow 25 \\ & \frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^2}{d(m+3)} - \\ & \frac{\int \tan^m(c+dx)((m+2)(bB(m+1)-aA(m+3))a^2-b(m+2)(2B(m+4)a^2+3Ab(m+3)a-b^2B(m+3)) \tan^2(c+dx) - (Ba^3+3Aba^2-3b^2Ba-Ab^3)(m+2))}{m+2} \\ & \frac{}{m+3} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^2}{d(m+3)} - \\ & \frac{\int \tan(c+dx)^m((m+2)(bB(m+1)-aA(m+3))a^2-b(m+2)(2B(m+4)a^2+3Ab(m+3)a-b^2B(m+3)) \tan(c+dx)^2 - (Ba^3+3Aba^2-3b^2Ba-Ab^3)(m+2))}{m+2} \\ & \frac{}{m+3} \end{aligned}$$

$$\begin{aligned} & \downarrow 4113 \\ & \frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^2}{d(m+3)} - \\ & \frac{\int \tan^m(c+dx)(-((Aa^3-3bBa^2-3Ab^2a+b^3B)(m+2)(m+3)) - (Ba^3+3Aba^2-3b^2Ba-Ab^3)(m+2) \tan(c+dx)(m+3)) dx - \frac{b(m+2)(2a^2B(m+4)+3a^3B(m+3))}{m+2}}{m+2} \\ & \frac{}{m+3} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^2}{d(m+3)} - \\ & \frac{\int \tan(c+dx)^m(-((Aa^3-3bBa^2-3Ab^2a+b^3B)(m+2)(m+3)) - (Ba^3+3Aba^2-3b^2Ba-Ab^3)(m+2) \tan(c+dx)(m+3)) dx - \frac{b(m+2)(2a^2B(m+4)+3a^3B(m+3))}{m+2}}{m+2} \\ & \frac{}{m+3} \end{aligned}$$

3.480. $\int \tan^m(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\begin{aligned} & \downarrow 4021 \\ & \frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^2}{d(m+3)} - \\ & \frac{-(m+2)(m+3)(a^3B+3a^2Ab-3ab^2B-Ab^3) \int \tan^{m+1}(c+dx)dx - (m+2)(m+3)(a^3A-3a^2bB-3aAb^2+b^3B) \int \tan^m(c+dx)dx - \frac{b(m+2)(2a^2B(m+3))}{m+2}}{m+2} \\ & \qquad \qquad \qquad m+3 \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^2}{d(m+3)} - \\ & \frac{-(m+2)(m+3)(a^3A-3a^2bB-3aAb^2+b^3B) \int \tan(c+dx)^m dx - (m+2)(m+3)(a^3B+3a^2Ab-3ab^2B-Ab^3) \int \tan(c+dx)^{m+1} dx - \frac{b(m+2)(2a^2B(m+3))}{m+2}}{m+2} \\ & \qquad \qquad \qquad m+3 \end{aligned}$$

$$\begin{aligned} & \downarrow 3957 \\ & \frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^2}{d(m+3)} - \\ & \frac{\frac{(m+2)(m+3)(a^3B+3a^2Ab-3ab^2B-Ab^3) \int \frac{\tan^{m+1}(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} - \frac{(m+2)(m+3)(a^3A-3a^2bB-3aAb^2+b^3B) \int \frac{\tan^m(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} - \frac{b(m+2)(2a^2B(m+3))}{m+2}}{m+2} \\ & \qquad \qquad \qquad m+3 \end{aligned}$$

$$\begin{aligned} & \downarrow 278 \\ & \frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))^2}{d(m+3)} - \\ & \frac{\frac{b(m+2)(2a^2B(m+4)+3aAb(m+3)-b^2B(m+3)) \tan^{m+1}(c+dx)}{d(m+1)} - \frac{(m+2)(m+3)(a^3A-3a^2bB-3aAb^2+b^3B) \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+1}{2}, \frac{m+3}{2}, -\tan^2(c+dx)\right)}{d(m+1)}}{m+2} \end{aligned}$$

```
input Int[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]
```

```
output (b*B*Tan[c + d*x]^(1 + m)*(a + b*Tan[c + d*x])^2)/(d*(3 + m)) - (-((b^2*(A *b*(3 + m) + a*B*(5 + m))*Tan[c + d*x]^(2 + m))/(d*(2 + m))) + (-((b*(2 + m)*(3*a*A*b*(3 + m) - b^2*B*(3 + m) + 2*a^2*B*(4 + m))*Tan[c + d*x]^(1 + m))/(d*(1 + m))) - ((a^3*A - 3*a*A*b^2 - 3*a^2*b*B + b^3*B)*(2 + m)*(3 + m)*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(1 + m))/(d*(1 + m)) - ((3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*(3 + m)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(2 + m))/d)/(2 + m)/(3 + m)
```

3.480. $\int \tan^m(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.480.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`
- rule 4021 `Int[((b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Tan[e + f*x])^m, x], x] + Simp[d/b Int[(b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x] && NeQ[c^2 + d^2, 0] && !IntegerQ[2*m]`
- rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`
- rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

```
rule 4120 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*
*(x_)])^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp
[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

3.480.4 Maple [F]

$$\int \tan(dx + c)^m (a + b \tan(dx + c))^3 (A + B \tan(dx + c)) dx$$

```
input int(tan(d*x+c)^m*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x)
```

```
output int(tan(d*x+c)^m*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x)
```

3.480.5 Fracas [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^3 \tan(dx + c)^m dx \end{aligned}$$

```
input integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="f
ricas")
```

```
output integral((B*b^3*tan(d*x + c)^4 + A*a^3 + (3*B*a*b^2 + A*b^3)*tan(d*x + c)^
3 + 3*(B*a^2*b + A*a*b^2)*tan(d*x + c)^2 + (B*a^3 + 3*A*a^2*b)*tan(d*x + c
))*tan(d*x + c)^m, x)
```

3.480.6 Sympy [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ &= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^3 \tan^m(c + dx) dx \end{aligned}$$

input `integrate(tan(d*x+c)**m*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)), x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**3*tan(c + d*x)**m, x)`

3.480.7 Maxima [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^3 \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)), x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^3*tan(d*x + c)^m, x)`

3.480.8 Giac [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^3 \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)), x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^3*tan(d*x + c)^m, x)`

3.480.9 Mupad [F(-1)]

Timed out.

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^m (A + B \tan(c + dx)) (a + b \tan(c + dx))^3 dx$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3, x)`

3.481 $\int \tan^m(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.481.1 Optimal result	4593
3.481.2 Mathematica [A] (verified)	4594
3.481.3 Rubi [A] (verified)	4594
3.481.4 Maple [F]	4597
3.481.5 Fricas [F]	4597
3.481.6 Sympy [F]	4598
3.481.7 Maxima [F]	4598
3.481.8 Giac [F]	4598
3.481.9 Mupad [F(-1)]	4599

3.481.1 Optimal result

Integrand size = 31, antiderivative size = 194

$$\int \tan^m(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{b(Ab(2+m) + aB(3+m)) \tan^{1+m}(c+dx)}{d(1+m)(2+m)}$$

$$+ \frac{(a^2A - Ab^2 - 2abB) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right) \tan^{1+m}(c+dx)}{d(1+m)}$$

$$+ \frac{(2aAb + a^2B - b^2B) \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, -\tan^2(c+dx)\right) \tan^{2+m}(c+dx)}{d(2+m)}$$

$$+ \frac{bB \tan^{1+m}(c+dx)(a+b \tan(c+dx))}{d(2+m)}$$

```
output b*(A*b*(2+m)+a*B*(3+m))*tan(d*x+c)^(1+m)/d/(1+m)/(2+m)+(A*a^2-A*b^2-2*B*a*
b)*hypergeom([1, 1/2+1/2*m],[3/2+1/2*m],-tan(d*x+c)^2)*tan(d*x+c)^(1+m)/d/
(1+m)+(2*A*a*b+B*a^2-B*b^2)*hypergeom([1, 1+1/2*m],[2+1/2*m],-tan(d*x+c)^2
)*tan(d*x+c)^(2+m)/d/(2+m)+b*B*tan(d*x+c)^(1+m)*(a+b*tan(d*x+c))/d/(2+m)
```


3.481.2 Mathematica [A] (verified)

Time = 0.74 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.80

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{\tan^{1+m}(c + dx) \left(\frac{b(Ab(2+m) + aB(3+m))}{1+m} + \frac{(a^2A - Ab^2 - 2abB)(2+m) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right)}{1+m} \right) + (2aAb + \dots)}{d(2 + \dots)}$$

```
input Integrate[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]
```

```
output (Tan[c + d*x]^(1 + m)*((b*(A*b*(2 + m) + a*B*(3 + m)))/(1 + m) + ((a^2*A - A*b^2 - 2*a*b*B)*(2 + m)*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2])/(1 + m) + (2*a*A*b + a^2*B - b^2*B)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x] + b*B*(a + b*Tan[c + d*x]))/(d*(2 + m))
```

3.481.3 Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.02, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3042, 4090, 25, 3042, 4113, 3042, 4021, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

↓ 3042

$$\int \tan(c + dx)^m(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

↓ 4090

$$\int \frac{-\tan^m(c + dx) (-b(Ab(m + 2) + aB(m + 3)) \tan^2(c + dx) - (Ba^2 + 2Aba - b^2B) (m + 2) \tan(c + dx) + a(bB \tan^{m+1}(c + dx)(a + b \tan(c + dx)))}{d(m + 2)} dx$$

↓ 25

3.481. $\int \tan^m(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\frac{\frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))}{d(m+2)} - \int \tan^m(c+dx) (-b(Ab(m+2) + aB(m+3)) \tan^2(c+dx) - (Ba^2 + 2Aba - b^2B)(m+2) \tan(c+dx) + a(bB(m+2) - (Aa^2 - 2bBa - Ab^2)(m+2))) dx - \frac{b(aB(m+3) + Ab(m+2))}{d(m+1)}}{m+2}$$

↓ 3042

$$\frac{\frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))}{d(m+2)} - \int \tan(c+dx)^m (-b(Ab(m+2) + aB(m+3)) \tan(c+dx)^2 - (Ba^2 + 2Aba - b^2B)(m+2) \tan(c+dx) + a(bB(m+2) - (Aa^2 - 2bBa - Ab^2)(m+2))) dx - \frac{b(aB(m+3) + Ab(m+2))}{d(m+1)}}{m+2}$$

↓ 4113

$$\frac{\frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))}{d(m+2)} - \int \tan^m(c+dx) (-((Aa^2 - 2bBa - Ab^2)(m+2)) - (Ba^2 + 2Aba - b^2B) \tan(c+dx)(m+2)) dx - \frac{b(aB(m+3) + Ab(m+2))}{d(m+1)}}{m+2}$$

↓ 3042

$$\frac{\frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))}{d(m+2)} - \int \tan(c+dx)^m (-((Aa^2 - 2bBa - Ab^2)(m+2)) - (Ba^2 + 2Aba - b^2B) \tan(c+dx)(m+2)) dx - \frac{b(aB(m+3) + Ab(m+2))}{d(m+1)}}{m+2}$$

↓ 4021

$$\frac{\frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))}{d(m+2)} - (m+2)(a^2B + 2aAb - b^2B) \int \tan^{m+1}(c+dx) dx - (m+2)(a^2A - 2abB - Ab^2) \int \tan^m(c+dx) dx - \frac{b(aB(m+3) + Ab(m+2))}{d(m+1)}}{m+2}$$

↓ 3042

$$\frac{\frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))}{d(m+2)} - (m+2)(a^2A - 2abB - Ab^2) \int \tan(c+dx)^m dx - (m+2)(a^2B + 2aAb - b^2B) \int \tan(c+dx)^{m+1} dx - \frac{b(aB(m+3) + Ab(m+2))}{d(m+1)}}{m+2}$$

↓ 3957

$$\frac{\frac{bB \tan^{m+1}(c+dx)(a+b \tan(c+dx))}{d(m+2)} - \frac{(m+2)(a^2B + 2aAb - b^2B) \int \frac{\tan^{m+1}(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} - \frac{(m+2)(a^2A - 2abB - Ab^2) \int \frac{\tan^m(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} - \frac{b(aB(m+3) + Ab(m+2))}{d(m+1)}}{m+2}$$

↓ 278

3.481. $\int \tan^m(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

$$\frac{bB \tan^{m+1}(c + dx)(a + b \tan(c + dx))}{d(m+2)} - \frac{(m+2)(a^2A - 2abB - Ab^2) \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+1}{2}, \frac{m+3}{2}, -\tan^2(c+dx)\right)}{d(m+1)} - \frac{(a^2B + 2aAb - b^2B) \tan^{m+2}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+4}{2}, -\tan^2(c+dx)\right)}{d(m+2)}$$

```
input Int[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]
```

```
output (b*B*Tan[c + d*x]^(1 + m)*(a + b*Tan[c + d*x]))/(d*(2 + m)) - (-(b*(A*b*(2 + m) + a*B*(3 + m))*Tan[c + d*x]^(1 + m))/(d*(1 + m))) - ((a^2*A - A*b^2 - 2*a*b*B)*(2 + m)*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(1 + m))/(d*(1 + m)) - ((2*a*A*b + a^2*B - b^2*B)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(2 + m))/d)/(2 + m)
```

3.481.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 278 Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(p_)), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3957 Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

```
rule 4021 Int[((b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Tan[e + f*x])^m, x], x] + Simp[d/b Int[(b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x] && NeQ[c^2 + d^2, 0] && !IntegerQ[2*m]
```

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

3.481.4 Maple [F]

$$\int \tan(dx + c)^m (a + b \tan(dx + c))^2 (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^m*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x)`

3.481.5 Fracas [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^2 \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*b^2*tan(d*x + c)^3 + A*a^2 + (2*B*a*b + A*b^2)*tan(d*x + c)^2 + (B*a^2 + 2*A*a*b)*tan(d*x + c))*tan(d*x + c)^m, x)`

3.481.6 Sympy [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx \\ &= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^2 \tan^m(c + dx) dx \end{aligned}$$

input `integrate(tan(d*x+c)**m*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**2*tan(c + d*x)**m, x)`

3.481.7 Maxima [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^2 \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^2*tan(d*x + c)^m, x)`

3.481.8 Giac [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^2 \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^2*tan(d*x + c)^m, x)`

3.481.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx \\ &= \int \tan(c + dx)^m (A + B \tan(c + dx)) (a + b \tan(c + dx))^2 dx \end{aligned}$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)`

output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2, x)`

3.482 $\int \tan^m(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

3.482.1 Optimal result	4600
3.482.2 Mathematica [A] (verified)	4601
3.482.3 Rubi [A] (verified)	4601
3.482.4 Maple [F]	4603
3.482.5 Fracas [F]	4603
3.482.6 Sympy [F]	4604
3.482.7 Maxima [F]	4604
3.482.8 Giac [F]	4604
3.482.9 Mupad [F(-1)]	4605

3.482.1 Optimal result

Integrand size = 29, antiderivative size = 127

$$\int \tan^m(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{bB \tan^{1+m}(c+dx)}{d(1+m)}$$

$$+ \frac{(aA - bB) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right) \tan^{1+m}(c+dx)}{d(1+m)}$$

$$+ \frac{(Ab + aB) \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, -\tan^2(c+dx)\right) \tan^{2+m}(c+dx)}{d(2+m)}$$

```
output b*B*tan(d*x+c)^(1+m)/d/(1+m)+(A*a-B*b)*hypergeom([1, 1/2+1/2*m], [3/2+1/2*m], -tan(d*x+c)^2)*tan(d*x+c)^(1+m)/d/(1+m)+(A*b+B*a)*hypergeom([1, 1+1/2*m], [2+1/2*m], -tan(d*x+c)^2)*tan(d*x+c)^(2+m)/d/(2+m)
```

3.482.2 Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.85

$$\int \tan^m(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{\tan^{1+m}(c+dx) \left(\frac{bB}{1+m} + \frac{(aA-bB) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right)}{1+m} + \frac{(Ab+aB) \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, -\tan^2(c+dx)\right)}{2+m} \right)}{d}$$

input `Integrate[Tan[c + d*x]^m*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(Tan[c + d*x]^(1 + m)*((b*B)/(1 + m) + ((a*A - b*B)*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2])/(1 + m) + ((A*b + a*B)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x])/(2 + m))/d`

3.482.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.241$, Rules used = {3042, 4075, 3042, 4021, 3042, 3957, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^m(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$\downarrow 3042$$

$$\int \tan(c+dx)^m(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$\downarrow 4075$$

$$\int \tan^m(c+dx)(aA-bB+(Ab+aB) \tan(c+dx)) dx + \frac{bB \tan^{m+1}(c+dx)}{d(m+1)}$$

$$\downarrow 3042$$

$$\int \tan(c+dx)^m(aA-bB+(Ab+aB) \tan(c+dx)) dx + \frac{bB \tan^{m+1}(c+dx)}{d(m+1)}$$

$$\downarrow 4021$$

$$\begin{aligned}
& (aB + Ab) \int \tan^{m+1}(c + dx) dx + (aA - bB) \int \tan^m(c + dx) dx + \frac{bB \tan^{m+1}(c + dx)}{d(m+1)} \\
& \quad \downarrow \text{3042} \\
& (aA - bB) \int \tan(c + dx)^m dx + (aB + Ab) \int \tan(c + dx)^{m+1} dx + \frac{bB \tan^{m+1}(c + dx)}{d(m+1)} \\
& \quad \downarrow \text{3957} \\
& \frac{(aB + Ab) \int \frac{\tan^{m+1}(c+dx)}{\tan^2(c+dx)+1} d \tan(c + dx)}{d} + \frac{(aA - bB) \int \frac{\tan^m(c+dx)}{\tan^2(c+dx)+1} d \tan(c + dx)}{d} + \\
& \quad \frac{bB \tan^{m+1}(c + dx)}{d(m+1)} \\
& \quad \downarrow \text{278} \\
& \frac{(aA - bB) \tan^{m+1}(c + dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+1}{2}, \frac{m+3}{2}, -\tan^2(c + dx)\right)}{d(m+1)} + \\
& \frac{(aB + Ab) \tan^{m+2}(c + dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+4}{2}, -\tan^2(c + dx)\right)}{d(m+2)} + \\
& \quad \frac{bB \tan^{m+1}(c + dx)}{d(m+1)}
\end{aligned}$$

input `Int[Tan[c + d*x]^m*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(b*B*Tan[c + d*x]^(1 + m))/(d*(1 + m)) + ((a*A - b*B)*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(1 + m))/(d*(1 + m)) + ((A*b + a*B)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(2 + m))/(d*(2 + m))`

3.482.3.1 Defintions of rubi rules used

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[b/d Subst[Int [x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4021 `Int[((b_.)*tan[(e_.) + (f_.)*(x_)]^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Tan[e + f*x])^m, x], x] + Simp[d/b Int [(b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x] && NeQ[c^2 + d^2, 0] && !IntegerQ[2*m]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.482.4 Maple [F]

$$\int \tan(dx + c)^m (a + b \tan(dx + c)) (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^m*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

3.482.5 Fracas [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a) \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*b*tan(d*x + c)^2 + A*a + (B*a + A*b)*tan(d*x + c))*tan(d*x + c)^m, x)`

3.482.6 Sympy [F]

$$\int \tan^m(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx))(a + b \tan(c + dx)) \tan^m(c + dx) dx$$

input `integrate(tan(d*x+c)**m*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))*tan(c + d*x)**m, x)`

3.482.7 Maxima [F]

$$\int \tan^m(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a) \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)*tan(d*x + c)^m, x)`

3.482.8 Giac [F]

$$\int \tan^m(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a) \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)*tan(d*x + c)^m, x)`

3.482.9 Mupad [F(-1)]

Timed out.

$$\int \tan^m(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^m (A + B \tan(c + dx)) (a + b \tan(c + dx)) dx$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)`output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)), x)`

3.483
$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.483.1 Optimal result 4606
 3.483.2 Mathematica [A] (verified) 4607
 3.483.3 Rubi [A] (verified) 4607
 3.483.4 Maple [F] 4610
 3.483.5 Fricas [F] 4610
 3.483.6 Sympy [F] 4611
 3.483.7 Maxima [F] 4611
 3.483.8 Giac [F] 4611
 3.483.9 Mupad [F(-1)] 4612

3.483.1 Optimal result

Integrand size = 31, antiderivative size = 185

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$= \frac{(aA + bB) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right) \tan^{1+m}(c+dx)}{(a^2 + b^2) d(1+m)}$$

$$+ \frac{b(Ab - aB) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, -\frac{b \tan(c+dx)}{a}\right) \tan^{1+m}(c+dx)}{a(a^2 + b^2) d(1+m)}$$

$$- \frac{(Ab - aB) \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, -\tan^2(c+dx)\right) \tan^{2+m}(c+dx)}{(a^2 + b^2) d(2+m)}$$

```
output (A*a+B*b)*hypergeom([1, 1/2+1/2*m], [3/2+1/2*m], -tan(d*x+c)^2)*tan(d*x+c)^(
1+m)/(a^2+b^2)/d/(1+m)+b*(A*b-B*a)*hypergeom([1, 1+m], [2+m], -b*tan(d*x+c)/
a)*tan(d*x+c)^(1+m)/a/(a^2+b^2)/d/(1+m)-(A*b-B*a)*hypergeom([1, 1+1/2*m], [
2+1/2*m], -tan(d*x+c)^2)*tan(d*x+c)^(2+m)/(a^2+b^2)/d/(2+m)
```

3.483.2 Mathematica [A] (verified)

Time = 1.01 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.78

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$= \frac{\tan^{1+m}(c+dx) \left((aA+bB) \operatorname{Hypergeometric2F1} \left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx) \right) + \frac{(Ab-aB)(b(2+m) \operatorname{Hypergeometric2F1} \left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx) \right))}{(a^2+b^2)d(1+m)} \right)}{(a^2+b^2)d(1+m)}$$

input `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`output `(Tan[c + d*x]^(1 + m)*((a*A + b*B)*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2] + ((A*b - a*B)*(b*(2 + m)*Hypergeometric2F1[1, 1 + m, 2 + m, -(b*Tan[c + d*x])/a]) - a*(1 + m)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]))/(a*(2 + m)))/((a^2 + b^2)*d*(1 + m))`**3.483.3 Rubi [A] (verified)**Time = 0.83 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.96, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3042, 4096, 3042, 4021, 3042, 3957, 278, 4117, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\tan(c+dx)^m(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$\downarrow \text{4096}$$

$$\frac{\int \tan^m(c+dx)(aA+bB - (Ab-aB) \tan(c+dx)) dx}{a^2+b^2} + \frac{b(Ab-aB) \int \frac{\tan^m(c+dx)(\tan^2(c+dx)+1)}{a+b \tan(c+dx)} dx}{a^2+b^2}$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
& \frac{\int \tan(c+dx)^m (aA + bB - (Ab - aB) \tan(c+dx)) dx}{a^2 + b^2} + \frac{b(Ab - aB) \int \frac{\tan(c+dx)^m (\tan(c+dx)^2 + 1)}{a + b \tan(c+dx)} dx}{a^2 + b^2} \\
& \quad \downarrow \text{4021} \\
& \frac{(aA + bB) \int \tan^m(c+dx) dx - (Ab - aB) \int \tan^{m+1}(c+dx) dx}{a^2 + b^2} + \\
& \quad \frac{b(Ab - aB) \int \frac{\tan(c+dx)^m (\tan(c+dx)^2 + 1)}{a + b \tan(c+dx)} dx}{a^2 + b^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(aA + bB) \int \tan(c+dx)^m dx - (Ab - aB) \int \tan(c+dx)^{m+1} dx}{a^2 + b^2} + \\
& \quad \frac{b(Ab - aB) \int \frac{\tan(c+dx)^m (\tan(c+dx)^2 + 1)}{a + b \tan(c+dx)} dx}{a^2 + b^2} \\
& \quad \downarrow \text{3957} \\
& \frac{(aA + bB) \int \frac{\tan^m(c+dx)}{\tan^2(c+dx) + 1} d \tan(c+dx)}{d} - \frac{(Ab - aB) \int \frac{\tan^{m+1}(c+dx)}{\tan^2(c+dx) + 1} d \tan(c+dx)}{d} + \\
& \quad \frac{b(Ab - aB) \int \frac{\tan(c+dx)^m (\tan(c+dx)^2 + 1)}{a + b \tan(c+dx)} dx}{a^2 + b^2} \\
& \quad \downarrow \text{278} \\
& \frac{b(Ab - aB) \int \frac{\tan(c+dx)^m (\tan(c+dx)^2 + 1)}{a + b \tan(c+dx)} dx}{a^2 + b^2} + \\
& \frac{(aA + bB) \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+1}{2}, \frac{m+3}{2}, -\tan^2(c+dx)\right)}{d(m+1)} - \frac{(Ab - aB) \tan^{m+2}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+4}{2}, -\tan^2(c+dx)\right)}{d(m+2)} \\
& \quad \downarrow \text{4117} \\
& \frac{b(Ab - aB) \int \frac{\tan^m(c+dx)}{a + b \tan(c+dx)} d \tan(c+dx)}{d(a^2 + b^2)} + \\
& \frac{(aA + bB) \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+1}{2}, \frac{m+3}{2}, -\tan^2(c+dx)\right)}{d(m+1)} - \frac{(Ab - aB) \tan^{m+2}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+4}{2}, -\tan^2(c+dx)\right)}{d(m+2)} \\
& \quad \downarrow \text{74} \\
& \frac{b(Ab - aB) \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, m+1, m+2, -\frac{b \tan(c+dx)}{a}\right)}{ad(m+1)(a^2 + b^2)} + \\
& \frac{(aA + bB) \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+1}{2}, \frac{m+3}{2}, -\tan^2(c+dx)\right)}{d(m+1)} - \frac{(Ab - aB) \tan^{m+2}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+4}{2}, -\tan^2(c+dx)\right)}{d(m+2)} \\
& \quad \downarrow \\
& \frac{b(Ab - aB) \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, m+1, m+2, -\frac{b \tan(c+dx)}{a}\right)}{a^2 + b^2}
\end{aligned}$$

3.483. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

input `Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(b*(A*b - a*B)*Hypergeometric2F1[1, 1 + m, 2 + m, -((b*Tan[c + d*x])/a)]*Tan[c + d*x]^(1 + m))/(a*(a^2 + b^2)*d*(1 + m)) + (((a*A + b*B)*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(1 + m))/(d*(1 + m)) - ((A*b - a*B)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(2 + m))/(d*(2 + m)))/(a^2 + b^2)`

3.483.3.1 Defintions of rubi rules used

rule 74 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))`

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

rule 4021 `Int[((b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Tan[e + f*x])^m, x], x] + Simp[d/b Int[(b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x] && NeQ[c^2 + d^2, 0] && !IntegerQ[2*m]`

rule 4096 `Int[(((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[a*A + b*B - (A*b - a*B)*Tan[e + f*x], x], x] + Simp[b*(A*b - a*B)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

3.483.4 Maple [F]

$$\int \frac{\tan(dx + c)^m (A + B \tan(dx + c))}{a + b \tan(dx + c)} dx$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

3.483.5 Fracas [F]

$$\int \frac{\tan^m(c + dx)(A + B \tan(c + dx))}{a + b \tan(c + dx)} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)^m}{b \tan(dx + c) + a} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*tan(d*x + c) + A)*tan(d*x + c)^m/(b*tan(d*x + c) + a), x)`

3.483.6 Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \int \frac{(A+B\tan(c+dx))\tan^m(c+dx)}{a+b\tan(c+dx)} dx$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**m/(a + b*tan(c + d*x)), x)`

3.483.7 Maxima [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{b\tan(dx+c)+a} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(b*tan(d*x + c) + a), x)`

3.483.8 Giac [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{b\tan(dx+c)+a} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(b*tan(d*x + c) + a), x)`

3.483.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \int \frac{\tan(c+dx)^m (A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

input `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`output `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)), x)`

3.484
$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.484.1 Optimal result 4613
 3.484.2 Mathematica [A] (verified) 4614
 3.484.3 Rubi [A] (verified) 4614
 3.484.4 Maple [F] 4618
 3.484.5 Fricas [F] 4618
 3.484.6 Sympy [F] 4619
 3.484.7 Maxima [F] 4619
 3.484.8 Giac [F] 4619
 3.484.9 Mupad [F(-1)] 4620

3.484.1 Optimal result

Integrand size = 31, antiderivative size = 282

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{(a^2A - Ab^2 + 2abB) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right) \tan^{1+m}(c+dx)}{(a^2 + b^2)^2 d(1+m)}$$

$$+ \frac{b(a^2Ab(2-m) - Ab^3m + ab^2B(1+m) - a^3(B - Bm)) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, -\frac{b \tan(c+dx)}{a}\right)}{a^2(a^2 + b^2)^2 d(1+m)}$$

$$- \frac{(2aAb - a^2B + b^2B) \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, -\tan^2(c+dx)\right) \tan^{2+m}(c+dx)}{(a^2 + b^2)^2 d(2+m)}$$

$$+ \frac{b(Ab - aB) \tan^{1+m}(c+dx)}{a(a^2 + b^2) d(a + b \tan(c+dx))}$$

output

```
(A*a^2-A*b^2+2*B*a*b)*hypergeom([1, 1/2+1/2*m], [3/2+1/2*m], -tan(d*x+c)^2)*
tan(d*x+c)^(1+m)/(a^2+b^2)^2/d/(1+m)+b*(a^2*A*b*(2-m)-A*b^3*m+a*b^2*B*(1+m)
)-a^3*(-B*m+B))*hypergeom([1, 1+m], [2+m], -b*tan(d*x+c)/a)*tan(d*x+c)^(1+m)
/a^2/(a^2+b^2)^2/d/(1+m)-(2*A*a*b-B*a^2+B*b^2)*hypergeom([1, 1+1/2*m], [2+1
/2*m], -tan(d*x+c)^2)*tan(d*x+c)^(2+m)/(a^2+b^2)^2/d/(2+m)+b*(A*b-B*a)*tan(
d*x+c)^(1+m)/a/(a^2+b^2)/d/(a+b*tan(d*x+c))
```

3.484.2 Mathematica [A] (verified)

Time = 3.33 (sec) , antiderivative size = 239, normalized size of antiderivative = 0.85

$$\int \frac{\tan^m(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx$$

$$= \frac{\tan^{1+m}(c + dx) \left(\frac{b(-a^2Ab(-2+m)+a^3B(-1+m)-Ab^3m+ab^2B(1+m)) \operatorname{Hypergeometric2F1}\left(1, 1+m, 2+m, -\frac{b \tan(c+dx)}{a}\right)}{a(a^2+b^2)(1+m)} + \frac{b(Ab-c)}{a+b \tan(c+dx)} \right)}{a^2+b^2}$$

input `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `(Tan[c + d*x]^(1 + m)*((b*(-(a^2*A*b*(-2 + m)) + a^3*B*(-1 + m) - A*b^3*m + a*b^2*B*(1 + m))*Hypergeometric2F1[1, 1 + m, 2 + m, -(b*Tan[c + d*x])/a]))/(a*(a^2 + b^2)*(1 + m)) + (b*(A*b - a*B))/(a + b*Tan[c + d*x]) + (a*((a^2*A - A*b^2 + 2*a*b*B)*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2])/(1 + m) + ((-2*a*A*b + a^2*B - b^2*B)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x])/(2 + m))/(a^2 + b^2))/d`

3.484.3 Rubi [A] (verified)

Time = 1.42 (sec) , antiderivative size = 292, normalized size of antiderivative = 1.04, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4092, 3042, 4136, 3042, 4021, 3042, 3957, 278, 4117, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^m(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx$$

↓ 3042

$$\int \frac{\tan(c + dx)^m(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx$$

↓ 4092

$$\frac{\int \frac{\tan^m(c+dx)(Aa^2+bB(m+1)a-(Ab-aB)\tan(c+dx)a-b(Ab-aB)m\tan^2(c+dx)-Ab^2m)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} + \frac{b(Ab-aB)\tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

↓ 3042

$$\frac{\int \frac{\tan(c+dx)^m(Aa^2+bB(m+1)a-(Ab-aB)\tan(c+dx)a-b(Ab-aB)m\tan(c+dx)^2-Ab^2m)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} + \frac{b(Ab-aB)\tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

↓ 4136

$$\frac{\int \tan^m(c+dx)(a(Aa^2+2bBa-Ab^2)-a(-Ba^2+2Aba+b^2B)\tan(c+dx)) dx}{a^2+b^2} - \frac{b(a^3B(1-m)-a^2Ab(2-m)-ab^2B(m+1)+Ab^3m)}{a^2+b^2} \int \frac{\tan^m(c+dx)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} + \frac{b(Ab-aB)\tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

↓ 3042

$$\frac{\int \tan(c+dx)^m(a(Aa^2+2bBa-Ab^2)-a(-Ba^2+2Aba+b^2B)\tan(c+dx)) dx}{a^2+b^2} - \frac{b(a^3B(1-m)-a^2Ab(2-m)-ab^2B(m+1)+Ab^3m)}{a^2+b^2} \int \frac{\tan(c+dx)^m}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} + \frac{b(Ab-aB)\tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

↓ 4021

$$\frac{a(a^2A+2abB-Ab^2) \int \tan^m(c+dx) dx - a(a^2(-B)+2aAb+b^2B) \int \tan^{m+1}(c+dx) dx}{a^2+b^2} - \frac{b(a^3B(1-m)-a^2Ab(2-m)-ab^2B(m+1)+Ab^3m)}{a^2+b^2} \int \frac{\tan(c+dx)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} + \frac{b(Ab-aB)\tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

↓ 3042

$$\frac{a(a^2A+2abB-Ab^2) \int \tan(c+dx)^m dx - a(a^2(-B)+2aAb+b^2B) \int \tan(c+dx)^{m+1} dx}{a^2+b^2} - \frac{b(a^3B(1-m)-a^2Ab(2-m)-ab^2B(m+1)+Ab^3m)}{a^2+b^2} \int \frac{\tan(c+dx)}{a+b\tan(c+dx)} dx}{a(a^2+b^2)} + \frac{b(Ab-aB)\tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b\tan(c+dx))}$$

↓ 3957

3.484. $\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

$$\frac{\frac{a(a^2A+2abB-Ab^2) \int \frac{\tan^m(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d} - \frac{a(a^2(-B)+2aAb+b^2B) \int \frac{\tan^{m+1}(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx)}{d}}{a^2+b^2} - \frac{b(a^3B(1-m)-a^2Ab(2-m)-ab^2B(m+1)+A)}{a^2+b^2}}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{ad(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 278

$$\frac{\frac{a(a^2A+2abB-Ab^2) \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+1}{2}, \frac{m+3}{2}, -\tan^2(c+dx)\right)}{d(m+1)} - \frac{a(a^2(-B)+2aAb+b^2B) \tan^{m+2}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+2}{2}, -\tan^2(c+dx)\right)}{d(m+2)}}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{ad(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 4117

$$\frac{\frac{a(a^2A+2abB-Ab^2) \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+1}{2}, \frac{m+3}{2}, -\tan^2(c+dx)\right)}{d(m+1)} - \frac{a(a^2(-B)+2aAb+b^2B) \tan^{m+2}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+2}{2}, -\tan^2(c+dx)\right)}{d(m+2)}}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{ad(a^2 + b^2)(a + b \tan(c + dx))}$$

↓ 74

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{ad(a^2 + b^2)(a + b \tan(c + dx))} + \frac{\frac{a(a^2A+2abB-Ab^2) \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+1}{2}, \frac{m+3}{2}, -\tan^2(c+dx)\right)}{d(m+1)} - \frac{a(a^2(-B)+2aAb+b^2B) \tan^{m+2}(c+dx) \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+2}{2}, -\tan^2(c+dx)\right)}{d(m+2)}}{a^2+b^2}$$

$$a(a^2 + b^2)$$

input `Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `(b*(A*b - a*B)*Tan[c + d*x]^(1 + m))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x]) + (-(b*(a^3*B*(1 - m) - a^2*A*b*(2 - m) + A*b^3*m - a*b^2*B*(1 + m))*Hypergeometric2F1[1, 1 + m, 2 + m, -(b*Tan[c + d*x])/a]*Tan[c + d*x]^(1 + m))/(a*(a^2 + b^2)*d*(1 + m))) + ((a*(a^2*A - A*b^2 + 2*a*b*B)*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(1 + m))/(d*(1 + m)) - (a*(2*a*A*b - a^2*B + b^2*B)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(2 + m))/(d*(2 + m)))/(a^2 + b^2))/(a*(a^2 + b^2))`

3.484. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.484.3.1 Defintions of rubi rules used

- rule 74 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))`
- rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`
- rule 4021 `Int[((b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Tan[e + f*x])^m, x], x] + Simp[d/b Int[(b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x] && NeQ[c^2 + d^2, 0] && !IntegerQ[2*m]`
- rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4136 `Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.484.4 Maple [F]

$$\int \frac{\tan(dx + c)^m (A + B \tan(dx + c))}{(a + b \tan(dx + c))^2} dx$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x)`

output `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x)`

3.484.5 Fracas [F]

$$\int \frac{\tan^m(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)^m}{(b \tan(dx + c) + a)^2} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fracas")`

output `integral((B*tan(d*x + c) + A)*tan(d*x + c)^m/(b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2), x)`

3.484.6 Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \int \frac{(A+B\tan(c+dx))\tan^m(c+dx)}{(a+b\tan(c+dx))^2} dx$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**m/(a + b*tan(c + d*x))**2, x)`

3.484.7 Maxima [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{(b\tan(dx+c)+a)^2} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(b*tan(d*x + c) + a)^2, x)`

3.484.8 Giac [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{(b\tan(dx+c)+a)^2} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(b*tan(d*x + c) + a)^2, x)`

3.484.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx = \int \frac{\tan(c+dx)^m (A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

input `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`output `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^2, x)`

3.485
$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

3.485.1 Optimal result 4621
 3.485.2 Mathematica [A] (verified) 4622
 3.485.3 Rubi [A] (verified) 4622
 3.485.4 Maple [F] 4628
 3.485.5 Fricas [F] 4628
 3.485.6 Sympy [F] 4629
 3.485.7 Maxima [F] 4629
 3.485.8 Giac [F] 4629
 3.485.9 Mupad [F(-1)] 4630

3.485.1 Optimal result

Integrand size = 31, antiderivative size = 438

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$= \frac{(a^3 A - 3aAb^2 + 3a^2bB - b^3 B) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right) \tan^{1+m}(c+dx)}{(a^2 + b^2)^3 d(1+m)}$$

$$- \frac{b(Ab^5(1-m)m + ab^4Bm(1+m) - 2a^3b^2B(3+m-m^2) + 2a^2Ab^3(1+3m-m^2) - a^4Ab(6-5m+m^2))}{2a^3(a^2 + b^2)^3 d}$$

$$- \frac{(3a^2Ab - Ab^3 - a^3B + 3ab^2B) \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, -\tan^2(c+dx)\right) \tan^{2+m}(c+dx)}{(a^2 + b^2)^3 d(2+m)}$$

$$+ \frac{b(Ab - aB) \tan^{1+m}(c+dx)}{2a(a^2 + b^2) d(a + b \tan(c+dx))^2}$$

$$+ \frac{b(Ab^3(1-m) - a^3B(3-m) + a^2Ab(5-m) + ab^2B(1+m)) \tan^{1+m}(c+dx)}{2a^2(a^2 + b^2)^2 d(a + b \tan(c+dx))}$$

```
output (A*a^3-3*A*a*b^2+3*B*a^2*b-B*b^3)*hypergeom([1, 1/2+1/2*m], [3/2+1/2*m], -tan(d*x+c)^2)*tan(d*x+c)^(1+m)/(a^2+b^2)^3/d/(1+m)-1/2*b*(A*b^5*(1-m)*m+a*b^4*B*m*(1+m)-2*a^3*b^2*B*(-m^2+m+3)+2*a^2*A*b^3*(-m^2+3*m+1)-a^4*A*b*(m^2-5*m+6)+a^5*B*(m^2-3*m+2))*hypergeom([1, 1+m], [2+m], -b*tan(d*x+c)/a)*tan(d*x+c)^(1+m)/a^3/(a^2+b^2)^3/d/(1+m)-(3*A*a^2*b-A*b^3-B*a^3+3*B*a*b^2)*hypergeom([1, 1+1/2*m], [2+1/2*m], -tan(d*x+c)^2)*tan(d*x+c)^(2+m)/(a^2+b^2)^3/d/(2+m)+1/2*b*(A*b-B*a)*tan(d*x+c)^(1+m)/a/(a^2+b^2)/d/(a+b*tan(d*x+c))^2+1/2*b*(A*b^3*(1-m)-a^3*B*(3-m)+a^2*A*b*(5-m)+a*b^2*B*(1+m))*tan(d*x+c)^(1+m)/a^2/(a^2+b^2)^2/d/(a+b*tan(d*x+c))
```

3.485.
$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

3.485.2 Mathematica [A] (verified)

Time = 6.41 (sec) , antiderivative size = 530, normalized size of antiderivative = 1.21

$$\int \frac{\tan^m(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx = \frac{b(Ab - aB) \tan^{1+m}(c + dx)}{2a(a^2 + b^2)d(a + b \tan(c + dx))^2} + \frac{(-a(-2ab(Ab - aB) - ab(Ab - aB)(1 - m)) + b^2(2a^2A + Ab^2(1 - m) + abB(1 + m))) \tan^{1+m}(c + dx)}{a(a^2 + b^2)d(a + b \tan(c + dx))} + \frac{(2a^3b(2aAb - a^2B + b^2B) - a^2bm(Ab^3(1 - m) - a^3b^3)) \tan^{1+m}(c + dx)}{a^2(a^2 + b^2)d(a + b \tan(c + dx))}$$

input `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

output `(b*(A*b - a*B)*Tan[c + d*x]^(1 + m))/(2*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) + (((-a*(-2*a*b*(A*b - a*B) - a*b*(A*b - a*B)*(1 - m))) + b^2*(2*a^2*A + A*b^2*(1 - m) + a*b*B*(1 + m)))*Tan[c + d*x]^(1 + m))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])) + (((2*a^3*b*(2*a*A*b - a^2*B + b^2*B) - a^2*b*m*(A*b^3*(1 - m) - a^3*B*(3 - m) + a^2*A*b*(5 - m) + a*b^2*B*(1 + m)) + b^2*(-(a^2*b*(A*b - a*B)*(3 - m)*(1 + m)) + (a^2 - b^2*m)*(2*a^2*A + A*b^2*(1 - m) + a*b*B*(1 + m))))*Hypergeometric2F1[1, 1 + m, 2 + m, -(b*Tan[c + d*x])/a])*Tan[c + d*x]^(1 + m))/(a*(a^2 + b^2)*d*(1 + m)) + ((2*a^2*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B)*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(1 + m))/(d*(1 + m)) - (2*a^2*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(2 + m))/(d*(2 + m)))/(a^2 + b^2))/(2*a*(a^2 + b^2))`

3.485.3 Rubi [A] (verified)

Time = 2.46 (sec) , antiderivative size = 464, normalized size of antiderivative = 1.06, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 4092, 3042, 4132, 25, 3042, 4136, 27, 3042, 4021, 3042, 3957, 278, 4117, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^m(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$$

↓ 3042

$$\begin{aligned}
 & \int \frac{\tan(c+dx)^m (A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{4092} \\
 & \int \frac{\tan^m(c+dx) (2Aa^2+bB(m+1)a-2(Ab-aB) \tan(c+dx)a+b(Ab-aB)(1-m) \tan^2(c+dx)+Ab^2(1-m))}{(a+b \tan(c+dx))^2} dx \\
 & \quad + \frac{2a(a^2+b^2)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} + \frac{b(Ab-aB) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^m (2Aa^2+bB(m+1)a-2(Ab-aB) \tan(c+dx)a+b(Ab-aB)(1-m) \tan(c+dx)^2+Ab^2(1-m))}{(a+b \tan(c+dx))^2} dx \\
 & \quad + \frac{2a(a^2+b^2)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} + \frac{b(Ab-aB) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow \text{4132} \\
 & \int - \frac{\tan^m(c+dx) (b(Ab-aB)(3-m)(m+1)a^2+2(-Ba^2+2Aba+b^2B) \tan(c+dx)a^2+bm(-B(3-m)a^3+Ab(5-m)a^2+b^2B(m+1)a+Ab^3(1-m)) \tan^2(c+dx)-(a^2-b^2))}{a(a^2+b^2)} dx \\
 & \quad + \frac{2a(a^2+b^2)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} + \frac{b(Ab-aB) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{b(a^3(-B)(3-m)+a^2Ab(5-m)+ab^2B(m+1)+Ab^3(1-m)) \tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b \tan(c+dx))} - \int \frac{\tan^m(c+dx) (b(Ab-aB)(3-m)(m+1)a^2+2(-Ba^2+2Aba+b^2B) \tan(c+dx)a^2+bm(-B(3-m)a^3+Ab(5-m)a^2+b^2B(m+1)a+Ab^3(1-m)) \tan^2(c+dx)-(a^2-b^2))}{a(a^2+b^2)} dx \\
 & \quad + \frac{2a(a^2+b^2)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} + \frac{b(Ab-aB) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{b(a^3(-B)(3-m)+a^2Ab(5-m)+ab^2B(m+1)+Ab^3(1-m)) \tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b \tan(c+dx))} - \int \frac{\tan(c+dx)^m (b(Ab-aB)(3-m)(m+1)a^2+2(-Ba^2+2Aba+b^2B) \tan(c+dx)a^2+bm(-B(3-m)a^3+Ab(5-m)a^2+b^2B(m+1)a+Ab^3(1-m)) \tan^2(c+dx)-(a^2-b^2))}{a(a^2+b^2)} dx \\
 & \quad + \frac{2a(a^2+b^2)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} + \frac{b(Ab-aB) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} \\
 & \quad \downarrow \text{4136}
 \end{aligned}$$

3.485. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{b(a^3(-B)(3-m)+a^2Ab(5-m)+ab^2B(m+1)+Ab^3(1-m)) \tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b \tan(c+dx))} - \frac{\int -2 \tan^m(c+dx) (a^2(Aa^3+3bBa^2-3Ab^2a-b^3B)-a^2(-Ba^3+3Aba^2+3b^3B))}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{2ad (a^2 + b^2) (a + b \tan(c + dx))^2}$$

↓ 27

$$\frac{b(a^3(-B)(3-m)+a^2Ab(5-m)+ab^2B(m+1)+Ab^3(1-m)) \tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b \tan(c+dx))} - \frac{b(a^5B(m^2-3m+2)-a^4Ab(m^2-5m+6)-2a^3b^2B(-m^2+m+3)+2a^2Ab^3)}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{2ad (a^2 + b^2) (a + b \tan(c + dx))^2}$$

↓ 3042

$$\frac{b(a^3(-B)(3-m)+a^2Ab(5-m)+ab^2B(m+1)+Ab^3(1-m)) \tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b \tan(c+dx))} - \frac{b(a^5B(m^2-3m+2)-a^4Ab(m^2-5m+6)-2a^3b^2B(-m^2+m+3)+2a^2Ab^3)}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{2ad (a^2 + b^2) (a + b \tan(c + dx))^2}$$

↓ 4021

$$\frac{b(a^3(-B)(3-m)+a^2Ab(5-m)+ab^2B(m+1)+Ab^3(1-m)) \tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b \tan(c+dx))} - \frac{b(a^5B(m^2-3m+2)-a^4Ab(m^2-5m+6)-2a^3b^2B(-m^2+m+3)+2a^2Ab^3)}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{2ad (a^2 + b^2) (a + b \tan(c + dx))^2}$$

↓ 3042

$$\frac{b(a^3(-B)(3-m)+a^2Ab(5-m)+ab^2B(m+1)+Ab^3(1-m)) \tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b \tan(c+dx))} - \frac{b(a^5B(m^2-3m+2)-a^4Ab(m^2-5m+6)-2a^3b^2B(-m^2+m+3)+2a^2Ab^3)}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{2ad (a^2 + b^2) (a + b \tan(c + dx))^2}$$

↓ 3957

3.485. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{b(a^3(-B)(3-m)+a^2Ab(5-m)+ab^2B(m+1)+Ab^3(1-m)) \tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b \tan(c+dx))} - \frac{b(a^5B(m^2-3m+2)-a^4Ab(m^2-5m+6)-2a^3b^2B(-m^2+m+3)+2a^2Ab^3)}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 278

$$\frac{b(a^3(-B)(3-m)+a^2Ab(5-m)+ab^2B(m+1)+Ab^3(1-m)) \tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b \tan(c+dx))} - \frac{b(a^5B(m^2-3m+2)-a^4Ab(m^2-5m+6)-2a^3b^2B(-m^2+m+3)+2a^2Ab^3)}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 4117

$$\frac{b(a^3(-B)(3-m)+a^2Ab(5-m)+ab^2B(m+1)+Ab^3(1-m)) \tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b \tan(c+dx))} - \frac{b(a^5B(m^2-3m+2)-a^4Ab(m^2-5m+6)-2a^3b^2B(-m^2+m+3)+2a^2Ab^3)}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2}$$

↓ 74

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{2ad(a^2 + b^2)(a + b \tan(c + dx))^2} +$$

$$\frac{b(a^3(-B)(3-m)+a^2Ab(5-m)+ab^2B(m+1)+Ab^3(1-m)) \tan^{m+1}(c+dx)}{ad(a^2+b^2)(a+b \tan(c+dx))} - \frac{b(a^5B(m^2-3m+2)-a^4Ab(m^2-5m+6)-2a^3b^2B(-m^2+m+3)+2a^2Ab^3)}{ad(a^2+b^2)(a+b \tan(c+dx))}$$

input `Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

3.485. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$


```
output (b*(A*b - a*B)*Tan[c + d*x]^(1 + m))/(2*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x
])^2) + ((b*(A*b^3*(1 - m) - a^3*B*(3 - m) + a^2*A*b*(5 - m) + a*b^2*B*(1
+ m))*Tan[c + d*x]^(1 + m))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])) - ((b*(
A*b^5*(1 - m)*m + a*b^4*B*m*(1 + m) - 2*a^3*b^2*B*(3 + m - m^2) + 2*a^2*A*
b^3*(1 + 3*m - m^2) - a^4*A*b*(6 - 5*m + m^2) + a^5*B*(2 - 3*m + m^2))*Hyp
ergeometric2F1[1, 1 + m, 2 + m, -(b*Tan[c + d*x])/a]*Tan[c + d*x]^(1 + m
))/(a*(a^2 + b^2)*d*(1 + m)) - (2*((a^2*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b
^3*B)*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2]*Tan[c +
d*x]^(1 + m))/(d*(1 + m)) - (a^2*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B)*H
ypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(2
+ m))/(d*(2 + m))))/(a^2 + b^2))/(a*(a^2 + b^2))/(2*a*(a^2 + b^2))
```

3.485.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 74 Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x
)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x]
/; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0]
&& !(EqQ[n, -2^(-1)]) && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0]))
```

```
rule 278 Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((
c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (
-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0]
|| GtQ[a, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3957 Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int
[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] &&
!IntegerQ[n]
```

rule 4021 `Int[((b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Tan[e + f*x])^m, x], x] + Simp[d/b Int[(b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{b, c, d, e, f, m}, x] && NeQ[c^2 + d^2, 0] && !IntegerQ[2*m]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x]))], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] & !GtQ[n, 0] && !LeQ[n, -1]`

3.485.4 Maple [F]

$$\int \frac{\tan(dx + c)^m (A + B \tan(dx + c))}{(a + b \tan(dx + c))^3} dx$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x)`

output `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x)`

3.485.5 Fricas [F]

$$\int \frac{\tan^m(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)^m}{(b \tan(dx + c) + a)^3} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="fricas")`

output `integral((B*tan(d*x + c) + A)*tan(d*x + c)^m/(b^3*tan(d*x + c)^3 + 3*a*b^2*tan(d*x + c)^2 + 3*a^2*b*tan(d*x + c) + a^3), x)`

3.485.6 Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \int \frac{(A+B\tan(c+dx))\tan^m(c+dx)}{(a+b\tan(c+dx))^3} dx$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**m/(a + b*tan(c + d*x))**3, x)`

3.485.7 Maxima [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{(b\tan(dx+c)+a)^3} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(b*tan(d*x + c) + a)^3, x)`

3.485.8 Giac [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{(b\tan(dx+c)+a)^3} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(b*tan(d*x + c) + a)^3, x)`

3.485.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \int \frac{\tan(c+dx)^m (A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

input `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3,x)`output `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3, x)`

3.486 $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

3.486.1 Optimal result	4631
3.486.2 Mathematica [B] (verified)	4632
3.486.3 Rubi [A] (verified)	4633
3.486.4 Maple [F]	4640
3.486.5 Fricas [F]	4640
3.486.6 Sympy [F]	4640
3.486.7 Maxima [F]	4641
3.486.8 Giac [F]	4641
3.486.9 Mupad [F(-1)]	4641

3.486.1 Optimal result

Integrand size = 31, antiderivative size = 659

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

$$= \frac{(a^4 A - 6a^2 Ab^2 + Ab^4 + 4a^3 bB - 4ab^3 B) \operatorname{Hypergeometric2F1}\left(1, \frac{1+m}{2}, \frac{3+m}{2}, -\tan^2(c+dx)\right) \tan^{1+m}(c+dx)}{(a^2 + b^2)^4 d(1+m)}$$

$$- \frac{b(ab^6 Bm(1 - m^2) + 3a^2 Ab^5 m(2 - 5m + m^2) + Ab^7 m(2 - 3m + m^2) + 3a^3 b^4 B(2 + 5m + 2m^2 - m^3))}{(a^2 + b^2)^4 d(1+m)}$$

$$- \frac{(4a^3 Ab - 4aAb^3 - a^4 B + 6a^2 b^2 B - b^4 B) \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, -\tan^2(c+dx)\right) \tan^{2+m}(c+dx)}{(a^2 + b^2)^4 d(2+m)}$$

$$+ \frac{b(Ab - aB) \tan^{1+m}(c+dx)}{3a(a^2 + b^2) d(a + b \tan(c+dx))^3}$$

$$+ \frac{b(Ab^3(2 - m) - a^3 B(5 - m) + a^2 Ab(8 - m) + ab^2 B(1 + m)) \tan^{1+m}(c+dx)}{6a^2(a^2 + b^2)^2 d(a + b \tan(c+dx))^2}$$

$$+ \frac{b(ab^4 B(1 - m^2) + 2a^3 b^2 B(7 + 3m - m^2) + a^4 Ab(26 - 9m + m^2) + 2a^2 Ab^3(2 - 6m + m^2) - a^5 B(11 - m^2))}{6a^3(a^2 + b^2)^3 d(a + b \tan(c+dx))}$$

output $(Aa^4 - 6Aa^2b^2 + Ab^4 + 4Bab^3 - 4Bab^3) \operatorname{hypergeom}([1, 1/2 + 1/2m], [3/2 + 1/2m], -\tan(dx+c)^2) \tan(dx+c)^{(1+m)} / (a^2+b^2)^4 / d / (1+m) - 1/6b(a^6b^6 Bm^2(-m^2+1) + 3a^2Ab^5m(m^2-5m+2) + Ab^7m(m^2-3m+2) + 3a^3b^4B(-m^3+2m^2+5m+2) + a^7B(-m^3+6m^2-11m+6) - a^6Ab(-m^3+9m^2-26m+24) + 3a^4Ab^3(m^3-7m^2+10m+8) - 3a^5b^2B(m^3-4m^2-m+12)) \operatorname{hypergeom}([1, 1+m], [2+m], -b \tan(dx+c)/a) \tan(dx+c)^{(1+m)} / a^4 / (a^2+b^2)^4 / d / (1+m) - (4Aa^3b - 4Aa^2b^3 - Ba^4 + 6Bab^2 - Bb^4) \operatorname{hypergeom}([1, 1+1/2m], [2+1/2m], -\tan(dx+c)^2) \tan(dx+c)^{(2+m)} / (a^2+b^2)^4 / d / (2+m) + 1/3b(Ab-Ba) \tan(dx+c)^{(1+m)} / a / (a^2+b^2) / d / (a+b \tan(dx+c))^3 + 1/6b(Ab^3(2-m) - a^3B(5-m) + a^2Ab(8-m) + ab^2B(1+m)) \tan(dx+c)^{(1+m)} / a^2 / (a^2+b^2)^2 / d / (a+b \tan(dx+c))^2 + 1/6b(a^4b^4B(-m^2+1) + 2a^3b^2B(-m^2+3m+7) + a^4Ab(m^2-9m+26) + 2a^2Ab^3(m^2-6m+2) - a^5B(m^2-6m+11) + Ab^5(m^2-3m+2)) \tan(dx+c)^{(1+m)} / a^3 / (a^2+b^2)^3 / d / (a+b \tan(dx+c))$

3.486.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1425 vs. $2(659) = 1318$.

Time = 6.47 (sec) , antiderivative size = 1425, normalized size of antiderivative = 2.16

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx = \text{Too large to display}$$

input `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]`

output

```
(b*(A*b - a*B)*Tan[c + d*x]^(1 + m))/(3*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x
])^3) + (((-a*(-3*a*b*(A*b - a*B) - a*b*(A*b - a*B)*(2 - m))) + b^2*(3*a^
2*A + A*b^2*(2 - m) + a*b*B*(1 + m))) * Tan[c + d*x]^(1 + m))/(2*a*(a^2 + b^
2)*d*(a + b*Tan[c + d*x])^2) + (((b^2*(-(a^2*b*(A*b - a*B)*(5 - m)*(1 + m)
) + (2*a^2 + b^2*(1 - m))*(3*a^2*A + A*b^2*(2 - m) + a*b*B*(1 + m))) - a*(
-6*a^2*b*(2*a*A*b - a^2*B + b^2*B) - a*b*(1 - m)*(A*b^3*(2 - m) - a^3*B*(5
- m) + a^2*A*b*(8 - m) + a*b^2*B*(1 + m)))) * Tan[c + d*x]^(1 + m))/(a*(a^2
+ b^2)*d*(a + b*Tan[c + d*x])) + (((a^2*b*(6*a^3*(2*a*A*b - a^2*B + b^2*B
) - b^2*(1 - m)*(A*b^3*(2 - m) - a^3*B*(5 - m) + a^2*A*b*(8 - m) + a*b^2*B
*(1 + m)) + b*(-(a^2*b*(A*b - a*B)*(5 - m)*(1 + m)) + (2*a^2 + b^2*(1 - m)
)*(3*a^2*A + A*b^2*(2 - m) + a*b*B*(1 + m)))) - a^2*m*(b^2*(-(a^2*b*(A*b -
a*B)*(5 - m)*(1 + m)) + (2*a^2 + b^2*(1 - m))*(3*a^2*A + A*b^2*(2 - m) +
a*b*B*(1 + m))) - a*(-6*a^2*b*(2*a*A*b - a^2*B + b^2*B) - a*b*(1 - m)*(A*b
^3*(2 - m) - a^3*B*(5 - m) + a^2*A*b*(8 - m) + a*b^2*B*(1 + m)))) + b^2*((
a^2 - b^2*m)*(-(a^2*b*(A*b - a*B)*(5 - m)*(1 + m)) + (2*a^2 + b^2*(1 - m)
)*(3*a^2*A + A*b^2*(2 - m) + a*b*B*(1 + m))) + a*(1 + m)*(-6*a^2*b*(2*a*A*b
- a^2*B + b^2*B) - a*b*(1 - m)*(A*b^3*(2 - m) - a^3*B*(5 - m) + a^2*A*b*(
8 - m) + a*b^2*B*(1 + m)))))*Hypergeometric2F1[1, 1 + m, 2 + m, -(b*Tan[c
+ d*x])/a]*Tan[c + d*x]^(1 + m))/(a*(a^2 + b^2)*d*(1 + m)) + ((6*a^7*A*H
ypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]...
```

3.486.3 Rubi [A] (verified)

Time = 4.06 (sec) , antiderivative size = 704, normalized size of antiderivative = 1.07, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.581$, Rules used = {3042, 4092, 3042, 4132, 25, 3042, 4132, 25, 3042, 4136, 27, 3042, 4021, 3042, 3957, 278, 4117, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

↓ 3042

$$\int \frac{\tan(c+dx)^m(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$$

↓ 4092

$$\begin{aligned}
 & \int \frac{\tan^m(c+dx)(3Aa^2+bB(m+1)a-3(Ab-aB)\tan(c+dx)a+b(Ab-aB)(2-m)\tan^2(c+dx)+Ab^2(2-m))}{(a+b\tan(c+dx))^3} dx \\
 & \qquad \qquad \qquad \frac{3a(a^2+b^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} + \\
 & \qquad \qquad \qquad \frac{b(Ab-aB)\tan^{m+1}(c+dx)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \int \frac{\tan^m(c+dx)^m(3Aa^2+bB(m+1)a-3(Ab-aB)\tan(c+dx)a+b(Ab-aB)(2-m)\tan(c+dx)^2+Ab^2(2-m))}{(a+b\tan(c+dx))^3} dx \\
 & \qquad \qquad \qquad \frac{3a(a^2+b^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} + \\
 & \qquad \qquad \qquad \frac{b(Ab-aB)\tan^{m+1}(c+dx)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \qquad \qquad \qquad \downarrow \text{4132} \\
 & \int -\frac{\tan^m(c+dx)(b(Ab-aB)(5-m)(m+1)a^2+6(-Ba^2+2Aba+b^2B)\tan(c+dx)a^2-b(1-m)(-B(5-m)a^3+Ab(8-m)a^2+b^2B(m+1)a+Ab^3(2-m))\tan^2(c+dx)-)}{(a+b\tan(c+dx))^2} \\
 & \qquad \qquad \qquad \frac{3a(a^2+b^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \qquad \qquad \qquad \frac{b(Ab-aB)\tan^{m+1}(c+dx)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \qquad \qquad \qquad \downarrow \text{25} \\
 & \frac{b(a^3(-B)(5-m)+a^2Ab(8-m)+ab^2B(m+1)+Ab^3(2-m))\tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2} - \int \frac{\tan^m(c+dx)(b(Ab-aB)(5-m)(m+1)a^2+6(-Ba^2+2Aba+b^2B)\tan(c+dx)-)}{(a+b\tan(c+dx))^2} \\
 & \qquad \qquad \qquad \frac{3a(a^2+b^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \qquad \qquad \qquad \frac{b(Ab-aB)\tan^{m+1}(c+dx)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \frac{b(a^3(-B)(5-m)+a^2Ab(8-m)+ab^2B(m+1)+Ab^3(2-m))\tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2} - \int \frac{\tan^m(c+dx)^m(b(Ab-aB)(5-m)(m+1)a^2+6(-Ba^2+2Aba+b^2B)\tan(c+dx)-)}{(a+b\tan(c+dx))^2} \\
 & \qquad \qquad \qquad \frac{3a(a^2+b^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \qquad \qquad \qquad \frac{b(Ab-aB)\tan^{m+1}(c+dx)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \qquad \qquad \qquad \downarrow \text{4132} \\
 & \frac{b(a^3(-B)(5-m)+a^2Ab(8-m)+ab^2B(m+1)+Ab^3(2-m))\tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b\tan(c+dx))^2} - \int \frac{\tan^m(c+dx)(6Aa^6+bB(m^3-6m^2+11m+18)a^5-Ab^2(m^3-9m^2+26m))}{(a+b\tan(c+dx))^2} \\
 & \qquad \qquad \qquad \frac{3a(a^2+b^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3} \\
 & \qquad \qquad \qquad \frac{b(Ab-aB)\tan^{m+1}(c+dx)}{3ad(a^2+b^2)(a+b\tan(c+dx))^3}
 \end{aligned}$$

3.486. $\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$

↓ 25

$$\frac{b(a^3(-B)(5-m)+a^2Ab(8-m)+ab^2B(m+1)+Ab^3(2-m)) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} - \frac{\int \frac{\tan^m(c+dx)(6Aa^6+bB(m^3-6m^2+11m+18)a^5-Ab^2(m^3-9m^2+26m-12))}{(a+b \tan(c+dx))^3} dx}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{3ad (a^2 + b^2) (a + b \tan(c + dx))^3}$$

↓ 3042

$$\frac{b(a^3(-B)(5-m)+a^2Ab(8-m)+ab^2B(m+1)+Ab^3(2-m)) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} - \frac{\int \frac{\tan(c+dx)^m (6Aa^6+bB(m^3-6m^2+11m+18)a^5-Ab^2(m^3-9m^2+26m-12))}{(a+b \tan(c+dx))^3} dx}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{3ad (a^2 + b^2) (a + b \tan(c + dx))^3}$$

↓ 4136

$$\frac{b(a^3(-B)(5-m)+a^2Ab(8-m)+ab^2B(m+1)+Ab^3(2-m)) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} - \frac{\int 6 \tan^m(c+dx) (a^3(Aa^4+4bBa^3-6Ab^2a^2-4b^3Ba+Ab^4)-a^3(-Ba^4+4bBa^3-6Ab^2a^2-4b^3Ba+Ab^4))}{(a+b \tan(c+dx))^3} dx}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{3ad (a^2 + b^2) (a + b \tan(c + dx))^3}$$

↓ 27

$$\frac{b(a^3(-B)(5-m)+a^2Ab(8-m)+ab^2B(m+1)+Ab^3(2-m)) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} - \frac{6 \int \tan^m(c+dx) (a^3(Aa^4+4bBa^3-6Ab^2a^2-4b^3Ba+Ab^4)-a^3(-Ba^4+4bBa^3-6Ab^2a^2-4b^3Ba+Ab^4))}{(a+b \tan(c+dx))^3} dx}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{3ad (a^2 + b^2) (a + b \tan(c + dx))^3}$$

↓ 3042

$$\frac{b(a^3(-B)(5-m)+a^2Ab(8-m)+ab^2B(m+1)+Ab^3(2-m)) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} - \frac{6 \int \tan(c+dx)^m (a^3(Aa^4+4bBa^3-6Ab^2a^2-4b^3Ba+Ab^4)-a^3(-Ba^4+4bBa^3-6Ab^2a^2-4b^3Ba+Ab^4))}{(a+b \tan(c+dx))^3} dx}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{3ad (a^2 + b^2) (a + b \tan(c + dx))^3}$$

3.486. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

↓ 4021

$$\frac{b(a^3(-B)(5-m)+a^2Ab(8-m)+ab^2B(m+1)+Ab^3(2-m)) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} - \frac{6(a^3(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4)) \int \tan^m(c+dx) dx - a^3(a^4(-1))}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{3ad (a^2 + b^2) (a + b \tan(c + dx))^3}$$

↓ 3042

$$\frac{b(a^3(-B)(5-m)+a^2Ab(8-m)+ab^2B(m+1)+Ab^3(2-m)) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} - \frac{6(a^3(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4)) \int \tan(c+dx)^m dx - a^3(a^4(-1))}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{3ad (a^2 + b^2) (a + b \tan(c + dx))^3}$$

↓ 3957

$$\frac{b(a^3(-B)(5-m)+a^2Ab(8-m)+ab^2B(m+1)+Ab^3(2-m)) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} - \frac{6 \left(\frac{a^3(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4)}{d} \int \frac{\tan^m(c+dx)}{\tan^2(c+dx)+1} d \tan(c+dx) \right)}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{3ad (a^2 + b^2) (a + b \tan(c + dx))^3}$$

↓ 278

$$\frac{b(a^3(-B)(5-m)+a^2Ab(8-m)+ab^2B(m+1)+Ab^3(2-m)) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} - \frac{6 \left(\frac{a^3(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4)}{d(m+1)} \tan^{m+1}(c+dx) \text{ Hypergeometric} \right)}{a^2+b^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{3ad (a^2 + b^2) (a + b \tan(c + dx))^3}$$

↓ 4117

3.486. $\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx$

$$\frac{b(a^3(-B)(5-m)+a^2Ab(8-m)+ab^2B(m+1)+Ab^3(2-m)) \tan^{m+1}(c+dx)}{2ad(a^2+b^2)(a+b \tan(c+dx))^2} - \frac{6 \left(\frac{a^3(a^4A+4a^3bB-6a^2Ab^2-4ab^3B+Ab^4) \tan^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left[1, 1+m, 2+m, -\frac{(b \tan(c+dx))}{a}\right]}{d(m+1)} \right)}{ad(a^2+b^2)(a+b \tan(c+dx))^2}$$

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{3ad(a^2 + b^2)(a + b \tan(c + dx))^3}$$

↓ 74

$$\frac{b(Ab - aB) \tan^{m+1}(c + dx)}{3ad(a^2 + b^2)(a + b \tan(c + dx))^3} + \frac{b(a^5(-B)(m^2-6m+11)+a^4Ab(m^2-9m+26)+2a^3b^2B(-m^2+3m+7))}{ad(a^2+b^2)(a+b \tan(c+dx))^2}$$

input `Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^4,x]`

output `(b*(A*b - a*B)*Tan[c + d*x]^(1 + m))/(3*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^3) + ((b*(A*b^3*(2 - m) - a^3*B*(5 - m) + a^2*A*b*(8 - m) + a*b^2*B*(1 + m))*Tan[c + d*x]^(1 + m))/(2*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^2) - (-((b*(a*b^4*B*(1 - m^2) + 2*a^3*b^2*B*(7 + 3*m - m^2) + a^4*A*b*(26 - 9*m + m^2) + 2*a^2*A*b^3*(2 - 6*m + m^2) - a^5*B*(11 - 6*m + m^2) + A*b^5*(2 - 3*m + m^2))*Tan[c + d*x]^(1 + m))/(a*(a^2 + b^2)*d*(a + b*Tan[c + d*x]))) - (-((b*(a*b^6*B*m*(1 - m^2) + 3*a^2*A*b^5*m*(2 - 5*m + m^2) + A*b^7*m*(2 - 3*m + m^2) + 3*a^3*b^4*B*(2 + 5*m + 2*m^2 - m^3) + a^7*B*(6 - 11*m + 6*m^2 - m^3) - a^6*A*b*(24 - 26*m + 9*m^2 - m^3) + 3*a^4*A*b^3*(8 + 10*m - 7*m^2 + m^3) - 3*a^5*b^2*B*(12 - m - 4*m^2 + m^3))*Hypergeometric2F1[1, 1 + m, 2 + m, -(b*Tan[c + d*x])/a]*Tan[c + d*x]^(1 + m))/(a*(a^2 + b^2)*d*(1 + m))) + (6*((a^3*(a^4*A - 6*a^2*A*b^2 + A*b^4 + 4*a^3*b*B - 4*a*b^3*B)*Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(1 + m))/(d*(1 + m)) - (a^3*(4*a^3*A*b - 4*a*A*b^3 - a^4*B + 6*a^2*b^2*B - b^4*B)*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, -Tan[c + d*x]^2]*Tan[c + d*x]^(2 + m))/(d*(2 + m)))/(a^2 + b^2))/(a*(a^2 + b^2)))/(2*a*(a^2 + b^2)))/(3*a*(a^2 + b^2))`

3.486.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 74 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m+1)/(b*(m+1)))*Hypergeometric2F1[-n, m+1, m+2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))`
- rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m+1)/(c*(m+1)))*Hypergeometric2F1[-p, (m+1)/2, (m+1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3957 `Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`
- rule 4021 `Int[((b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c Int[(b*Tan[e + f*x])^m, x], x] + Simp[d/b Int[(b*Tan[e + f*x])^(m+1), x], x] /; FreeQ[{b, c, d, e, f, m}, x] && NeQ[c^2 + d^2, 0] && !IntegerQ[2*m]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.486.4 Maple [F]

$$\int \frac{\tan(dx+c)^m (A+B \tan(dx+c))}{(a+b \tan(dx+c))^4} dx$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x)`

output `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x)`

3.486.5 Fricas [F]

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx = \int \frac{(B \tan(dx+c) + A) \tan(dx+c)^m}{(b \tan(dx+c) + a)^4} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="fricas")`

output `integral((B*tan(d*x + c) + A)*tan(d*x + c)^m/(b^4*tan(d*x + c)^4 + 4*a*b^3*tan(d*x + c)^3 + 6*a^2*b^2*tan(d*x + c)^2 + 4*a^3*b*tan(d*x + c) + a^4), x)`

3.486.6 Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^4} dx = \int \frac{(A+B \tan(c+dx)) \tan^m(c+dx)}{(a+b \tan(c+dx))^4} dx$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**4,x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**m/(a + b*tan(c + d*x))**4, x)`

3.486.7 Maxima [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{(b\tan(dx+c)+a)^4} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(b*tan(d*x + c) + a)^4, x)`

3.486.8 Giac [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{(b\tan(dx+c)+a)^4} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^4,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(b*tan(d*x + c) + a)^4, x)`

3.486.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx = \int \frac{\tan(c+dx)^m(A+B\tan(c+dx))}{(a+b\tan(c+dx))^4} dx$$

input `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^4,x)`

output `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^4, x)`

3.487 $\int \tan^m(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.487.1 Optimal result	4642
3.487.2 Mathematica [F]	4643
3.487.3 Rubi [A] (verified)	4643
3.487.4 Maple [F]	4645
3.487.5 Fricas [F]	4646
3.487.6 Sympy [F(-1)]	4646
3.487.7 Maxima [F]	4646
3.487.8 Giac [F(-1)]	4647
3.487.9 Mupad [F(-1)]	4647

3.487.1 Optimal result

Integrand size = 33, antiderivative size = 193

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{a^2(A + iB) \operatorname{AppellF1}\left(1 + m, -\frac{5}{2}, 1, 2 + m, -\frac{b \tan(c + dx)}{a}, -i \tan(c + dx)\right) \tan^{1+m}(c + dx) + B \tan(c + dx)}{2d(1 + m) \sqrt{1 + \frac{b \tan(c + dx)}{a}}} + \frac{a^2(A - iB) \operatorname{AppellF1}\left(1 + m, -\frac{5}{2}, 1, 2 + m, -\frac{b \tan(c + dx)}{a}, i \tan(c + dx)\right) \tan^{1+m}(c + dx) \sqrt{a + b \tan(c + dx)}}{2d(1 + m) \sqrt{1 + \frac{b \tan(c + dx)}{a}}}$$

output

```
1/2*a^2*(A+I*B)*AppellF1(1+m,1,-5/2,2+m,-I*tan(d*x+c),-b*tan(d*x+c)/a)*(a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^(1+m)/d/(1+m)/(1+b*tan(d*x+c)/a)^(1/2)+1/2*a^2*(A-I*B)*AppellF1(1+m,1,-5/2,2+m,I*tan(d*x+c),-b*tan(d*x+c)/a)*(a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^(1+m)/d/(1+m)/(1+b*tan(d*x+c)/a)^(1/2)
```

3.487.2 Mathematica [F]

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int \tan^m(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

input `Integrate[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Integrate[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

3.487.3 Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3042, 4086, 3042, 4085, 152, 150}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(c + dx)^m(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4086} \\ & \frac{1}{2}(A + iB) \int (1 - i \tan(c + dx)) \tan^m(c + dx)(a + b \tan(c + dx))^{5/2} dx + \frac{1}{2}(A - iB) \int (i \tan(c + dx) + 1) \tan^m(c + dx)(a + b \tan(c + dx))^{5/2} dx \\ & \quad \downarrow \text{3042} \\ & \frac{1}{2}(A + iB) \int (1 - i \tan(c + dx)) \tan(c + dx)^m(a + b \tan(c + dx))^{5/2} dx + \frac{1}{2}(A - iB) \int (i \tan(c + dx) + 1) \tan(c + dx)^m(a + b \tan(c + dx))^{5/2} dx \\ & \quad \downarrow \text{4085} \end{aligned}$$

$$\begin{aligned}
& \frac{(A - iB) \int \frac{\tan^m(c+dx)(a+b \tan(c+dx))^{5/2}}{1-i \tan(c+dx)} d \tan(c+dx)}{2d} + \\
& \frac{(A + iB) \int \frac{\tan^m(c+dx)(a+b \tan(c+dx))^{5/2}}{i \tan(c+dx)+1} d \tan(c+dx)}{2d} \\
& \quad \downarrow \text{152} \\
& \frac{a^2(A - iB) \sqrt{a + b \tan(c+dx)} \int \frac{\tan^m(c+dx) \left(\frac{b \tan(c+dx)}{a} + 1\right)^{5/2}}{1-i \tan(c+dx)} d \tan(c+dx)}{2d \sqrt{\frac{b \tan(c+dx)}{a} + 1}} + \\
& \frac{a^2(A + iB) \sqrt{a + b \tan(c+dx)} \int \frac{\tan^m(c+dx) \left(\frac{b \tan(c+dx)}{a} + 1\right)^{5/2}}{i \tan(c+dx)+1} d \tan(c+dx)}{2d \sqrt{\frac{b \tan(c+dx)}{a} + 1}} \\
& \quad \downarrow \text{150} \\
& \frac{a^2(A + iB) \tan^{m+1}(c+dx) \sqrt{a + b \tan(c+dx)} \operatorname{AppellF1}\left(m+1, -\frac{5}{2}, 1, m+2, -\frac{b \tan(c+dx)}{a}, -i \tan(c+dx)\right)}{2d(m+1) \sqrt{\frac{b \tan(c+dx)}{a} + 1}} + \\
& \frac{a^2(A - iB) \tan^{m+1}(c+dx) \sqrt{a + b \tan(c+dx)} \operatorname{AppellF1}\left(m+1, -\frac{5}{2}, 1, m+2, -\frac{b \tan(c+dx)}{a}, i \tan(c+dx)\right)}{2d(m+1) \sqrt{\frac{b \tan(c+dx)}{a} + 1}}
\end{aligned}$$

input `Int[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(a^2*(A + I*B)*AppellF1[1 + m, -5/2, 1, 2 + m, -((b*Tan[c + d*x])/a), (-I)*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*Sqrt[a + b*Tan[c + d*x]]/(2*d*(1 + m)*Sqrt[1 + (b*Tan[c + d*x])/a]) + (a^2*(A - I*B)*AppellF1[1 + m, -5/2, 1, 2 + m, -((b*Tan[c + d*x])/a), I*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*Sqrt[a + b*Tan[c + d*x]]/(2*d*(1 + m)*Sqrt[1 + (b*Tan[c + d*x])/a])`

3.487.3.1 Defintions of rubi rules used

rule 150 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_] :> Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])`

- rule 152 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_))*((e_) + (f_.)*(x_)^(p_)), x_] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4085 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && EqQ[A^2 + B^2, 0]`
- rule 4086 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`

3.487.4 Maple [F]

$$\int \tan(dx + c)^m (a + b \tan(dx + c))^{5/2} (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^m*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x)`

3.487.5 Fracas [F]

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{5/2} \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm m="fricas")`

output `integral((B*b^2*tan(d*x + c)^3 + A*a^2 + (2*B*a*b + A*b^2)*tan(d*x + c)^2 + (B*a^2 + 2*A*a*b)*tan(d*x + c))*sqrt(b*tan(d*x + c) + a)*tan(d*x + c)^m, x)`

3.487.6 Sympy [F(-1)]

Timed out.

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**m*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.487.7 Maxima [F]

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{5/2} \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm m="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)*tan(d*x + c)^m, x)`

3.487.8 Giac [F(-1)]

Timed out.

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.487.9 Mupad [F(-1)]

Timed out.

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int \tan(c + dx)^m (A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2} dx$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)`

output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2), x)`

3.488 $\int \tan^m(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.488.1 Optimal result	4648
3.488.2 Mathematica [F]	4649
3.488.3 Rubi [A] (verified)	4649
3.488.4 Maple [F]	4651
3.488.5 Fricas [F]	4652
3.488.6 Sympy [F]	4652
3.488.7 Maxima [F]	4652
3.488.8 Giac [F(-1)]	4653
3.488.9 Mupad [F(-1)]	4653

3.488.1 Optimal result

Integrand size = 33, antiderivative size = 189

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{a(A + iB) \operatorname{AppellF1}\left(1 + m, -\frac{3}{2}, 1, 2 + m, -\frac{b \tan(c + dx)}{a}, -i \tan(c + dx)\right) \tan^{1+m}(c + dx) + B \tan(c + dx)}{2d(1 + m)\sqrt{1 + \frac{b \tan(c + dx)}{a}}} + \frac{a(A - iB) \operatorname{AppellF1}\left(1 + m, -\frac{3}{2}, 1, 2 + m, -\frac{b \tan(c + dx)}{a}, i \tan(c + dx)\right) \tan^{1+m}(c + dx) \sqrt{a + b \tan(c + dx)}}{2d(1 + m)\sqrt{1 + \frac{b \tan(c + dx)}{a}}}$$

```
output 1/2*a*(A+I*B)*AppellF1(1+m,1,-3/2,2+m,-I*tan(d*x+c),-b*tan(d*x+c)/a)*(a+b*
tan(d*x+c))^(1/2)*tan(d*x+c)^(1+m)/d/(1+m)/(1+b*tan(d*x+c)/a)^(1/2)+1/2*a*
(A-I*B)*AppellF1(1+m,1,-3/2,2+m,I*tan(d*x+c),-b*tan(d*x+c)/a)*(a+b*tan(d*x
+c))^(1/2)*tan(d*x+c)^(1+m)/d/(1+m)/(1+b*tan(d*x+c)/a)^(1/2)
```

3.488.2 Mathematica [F]

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int \tan^m(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$$

input `Integrate[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `Integrate[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

3.488.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3042, 4086, 3042, 4085, 152, 150}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \tan(c + dx)^m(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4086} \\ & \frac{1}{2}(A + iB) \int (1 - i \tan(c + dx)) \tan^m(c + dx)(a + b \tan(c + dx))^{3/2} dx + \frac{1}{2}(A - iB) \int (i \tan(c + dx) + 1) \tan^m(c + dx)(a + b \tan(c + dx))^{3/2} dx \\ & \quad \downarrow \text{3042} \\ & \frac{1}{2}(A + iB) \int (1 - i \tan(c + dx)) \tan(c + dx)^m(a + b \tan(c + dx))^{3/2} dx + \frac{1}{2}(A - iB) \int (i \tan(c + dx) + 1) \tan(c + dx)^m(a + b \tan(c + dx))^{3/2} dx \\ & \quad \downarrow \text{4085} \end{aligned}$$

$$\begin{aligned}
& \frac{(A - iB) \int \frac{\tan^m(c+dx)(a+b \tan(c+dx))^{3/2}}{1-i \tan(c+dx)} d \tan(c+dx)}{2d} + \\
& \frac{(A + iB) \int \frac{\tan^m(c+dx)(a+b \tan(c+dx))^{3/2}}{i \tan(c+dx)+1} d \tan(c+dx)}{2d} \\
& \quad \downarrow \text{152} \\
& \frac{a(A - iB) \sqrt{a + b \tan(c+dx)} \int \frac{\tan^m(c+dx) \left(\frac{b \tan(c+dx)}{a} + 1\right)^{3/2}}{1-i \tan(c+dx)} d \tan(c+dx)}{2d \sqrt{\frac{b \tan(c+dx)}{a} + 1}} + \\
& \frac{a(A + iB) \sqrt{a + b \tan(c+dx)} \int \frac{\tan^m(c+dx) \left(\frac{b \tan(c+dx)}{a} + 1\right)^{3/2}}{i \tan(c+dx)+1} d \tan(c+dx)}{2d \sqrt{\frac{b \tan(c+dx)}{a} + 1}} \\
& \quad \downarrow \text{150} \\
& \frac{a(A + iB) \tan^{m+1}(c+dx) \sqrt{a + b \tan(c+dx)} \operatorname{AppellF1}\left(m+1, -\frac{3}{2}, 1, m+2, -\frac{b \tan(c+dx)}{a}, -i \tan(c+dx)\right)}{2d(m+1) \sqrt{\frac{b \tan(c+dx)}{a} + 1}} + \\
& \frac{a(A - iB) \tan^{m+1}(c+dx) \sqrt{a + b \tan(c+dx)} \operatorname{AppellF1}\left(m+1, -\frac{3}{2}, 1, m+2, -\frac{b \tan(c+dx)}{a}, i \tan(c+dx)\right)}{2d(m+1) \sqrt{\frac{b \tan(c+dx)}{a} + 1}}
\end{aligned}$$

input `Int[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(a*(A + I*B)*AppellF1[1 + m, -3/2, 1, 2 + m, -((b*Tan[c + d*x])/a), (-I)*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*Sqrt[a + b*Tan[c + d*x]]/(2*d*(1 + m)*Sqrt[1 + (b*Tan[c + d*x])/a]) + (a*(A - I*B)*AppellF1[1 + m, -3/2, 1, 2 + m, -((b*Tan[c + d*x])/a), I*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*Sqrt[a + b*Tan[c + d*x]]/(2*d*(1 + m)*Sqrt[1 + (b*Tan[c + d*x])/a])`

3.488.3.1 Defintions of rubi rules used

rule 150 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_] :> Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])`

- rule 152 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_))*((e_) + (f_.)*(x_)^(p_)), x_] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4085 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && EqQ[A^2 + B^2, 0]`
- rule 4086 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`

3.488.4 Maple [F]

$$\int \tan(dx + c)^m (a + b \tan(dx + c))^{3/2} (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^m*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x)`

3.488.5 Fracas [F]

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{3}{2}} \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*b*tan(d*x + c)^2 + A*a + (B*a + A*b)*tan(d*x + c))*sqrt(b*tan(d*x + c) + a)*tan(d*x + c)^m, x)`

3.488.6 Sympy [F]

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (A + B \tan(c + dx))(a + b \tan(c + dx))^{\frac{3}{2}} \tan^m(c + dx) dx$$

input `integrate(tan(d*x+c)**m*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)*tan(c + d*x)**m, x)`

3.488.7 Maxima [F]

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{3}{2}} \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)*tan(d*x + c)^m, x)`

3.488.8 Giac [F(-1)]

Timed out.

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.488.9 Mupad [F(-1)]

Timed out.

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int \tan(c + dx)^m (A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2} dx$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2),x)`

output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2), x)`

3.489 $\int \tan^m(c+dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.489.1 Optimal result	4654
3.489.2 Mathematica [F]	4655
3.489.3 Rubi [A] (verified)	4655
3.489.4 Maple [F]	4657
3.489.5 Fracas [F]	4658
3.489.6 Sympy [F]	4658
3.489.7 Maxima [F]	4658
3.489.8 Giac [F]	4659
3.489.9 Mupad [F(-1)]	4659

3.489.1 Optimal result

Integrand size = 33, antiderivative size = 187

$$\int \tan^m(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{(A + iB) \operatorname{AppellF1}\left(1 + m, -\frac{1}{2}, 1, 2 + m, -\frac{b \tan(c + dx)}{a}, -i \tan(c + dx)\right) \tan^{1+m}(c + dx) \sqrt{a + b \tan(c + dx)}}{2d(1 + m) \sqrt{1 + \frac{b \tan(c + dx)}{a}}} + \frac{(A - iB) \operatorname{AppellF1}\left(1 + m, -\frac{1}{2}, 1, 2 + m, -\frac{b \tan(c + dx)}{a}, i \tan(c + dx)\right) \tan^{1+m}(c + dx) \sqrt{a + b \tan(c + dx)}}{2d(1 + m) \sqrt{1 + \frac{b \tan(c + dx)}{a}}}$$

output `1/2*(A+I*B)*AppellF1(1+m,1,-1/2,2+m,-I*tan(d*x+c),-b*tan(d*x+c)/a)*(a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^(1+m)/d/(1+m)/(1+b*tan(d*x+c)/a)^(1/2)+1/2*(A-I*B)*AppellF1(1+m,1,-1/2,2+m,I*tan(d*x+c),-b*tan(d*x+c)/a)*(a+b*tan(d*x+c))^(1/2)*tan(d*x+c)^(1+m)/d/(1+m)/(1+b*tan(d*x+c)/a)^(1/2)`

3.489.2 Mathematica [F]

$$\int \tan^m(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \tan^m(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

input `Integrate[Tan[c + d*x]^m*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `Integrate[Tan[c + d*x]^m*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]), x]`

3.489.3 Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3042, 4086, 3042, 4085, 152, 150}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^m(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^m \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$\downarrow \text{4086}$$

$$\frac{1}{2}(A + iB) \int (1 - i \tan(c + dx)) \tan^m(c + dx) \sqrt{a + b \tan(c + dx)} dx + \frac{1}{2}(A - iB) \int (i \tan(c + dx) + 1) \tan^m(c + dx) \sqrt{a + b \tan(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\frac{1}{2}(A + iB) \int (1 - i \tan(c + dx)) \tan(c + dx)^m \sqrt{a + b \tan(c + dx)} dx + \frac{1}{2}(A - iB) \int (i \tan(c + dx) + 1) \tan(c + dx)^m \sqrt{a + b \tan(c + dx)} dx$$

$$\downarrow \text{4085}$$

$$\begin{aligned}
& \frac{(A - iB) \int \frac{\tan^m(c+dx) \sqrt{a+b \tan(c+dx)}}{1-i \tan(c+dx)} d \tan(c+dx)}{2d} + \\
& \frac{(A + iB) \int \frac{\tan^m(c+dx) \sqrt{a+b \tan(c+dx)}}{i \tan(c+dx)+1} d \tan(c+dx)}{2d} \\
& \quad \downarrow \text{152} \\
& \frac{(A - iB) \sqrt{a+b \tan(c+dx)} \int \frac{\tan^m(c+dx) \sqrt{\frac{b \tan(c+dx)}{a}+1}}{1-i \tan(c+dx)} d \tan(c+dx)}{2d \sqrt{\frac{b \tan(c+dx)}{a}+1}} + \\
& \frac{(A + iB) \sqrt{a+b \tan(c+dx)} \int \frac{\tan^m(c+dx) \sqrt{\frac{b \tan(c+dx)}{a}+1}}{i \tan(c+dx)+1} d \tan(c+dx)}{2d \sqrt{\frac{b \tan(c+dx)}{a}+1}} \\
& \quad \downarrow \text{150} \\
& \frac{(A + iB) \tan^{m+1}(c+dx) \sqrt{a+b \tan(c+dx)} \operatorname{AppellF1}\left(m+1, -\frac{1}{2}, 1, m+2, -\frac{b \tan(c+dx)}{a}, -i \tan(c+dx)\right)}{2d(m+1) \sqrt{\frac{b \tan(c+dx)}{a}+1}} + \\
& \frac{(A - iB) \tan^{m+1}(c+dx) \sqrt{a+b \tan(c+dx)} \operatorname{AppellF1}\left(m+1, -\frac{1}{2}, 1, m+2, -\frac{b \tan(c+dx)}{a}, i \tan(c+dx)\right)}{2d(m+1) \sqrt{\frac{b \tan(c+dx)}{a}+1}}
\end{aligned}$$

input `Int[Tan[c + d*x]^m*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `((A + I*B)*AppellF1[1 + m, -1/2, 1, 2 + m, -((b*Tan[c + d*x])/a), (-I)*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*Sqrt[a + b*Tan[c + d*x]])/(2*d*(1 + m)*Sqrt[1 + (b*Tan[c + d*x])/a]) + ((A - I*B)*AppellF1[1 + m, -1/2, 1, 2 + m, -((b*Tan[c + d*x])/a), I*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*Sqrt[a + b*Tan[c + d*x]])/(2*d*(1 + m)*Sqrt[1 + (b*Tan[c + d*x])/a])`

3.489.3.1 Defintions of rubi rules used

rule 150 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_))*((e_) + (f_.)*(x_)^(p_), x_] := Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])`

rule 152 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_))*((e_) + (f_.)*(x_)^(p_)), x_] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4085 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && EqQ[A^2 + B^2, 0]`

rule 4086 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`

3.489.4 Maple [F]

$$\int \tan(dx + c)^m \sqrt{a + b \tan(dx + c)} (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^m*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x)`

3.489.5 Fracas [F]

$$\int \tan^m(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm m="fricas")`

output `integral((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*tan(d*x + c)^m, x)`

3.489.6 Sympy [F]

$$\int \tan^m(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} \tan^m(c + dx) dx$$

input `integrate(tan(d*x+c)**m*(a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))*tan(c + d*x)**m, x)`

3.489.7 Maxima [F]

$$\int \tan^m(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm m="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*tan(d*x + c)^m, x)`

3.489.8 Giac [F]

$$\int \tan^m(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm
m="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*tan(d*x + c)^m, x)`

3.489.9 Mupad [F(-1)]

Timed out.

$$\int \tan^m(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^m (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`

output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2), x)`

3.490
$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

3.490.1 Optimal result 4660
 3.490.2 Mathematica [F] 4661
 3.490.3 Rubi [A] (verified) 4661
 3.490.4 Maple [F] 4664
 3.490.5 Fricas [F] 4664
 3.490.6 Sympy [F] 4664
 3.490.7 Maxima [F] 4665
 3.490.8 Giac [F(-1)] 4665
 3.490.9 Mupad [F(-1)] 4665

3.490.1 Optimal result

Integrand size = 33, antiderivative size = 187

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

$$= \frac{(A+iB) \operatorname{AppellF1}\left(1+m, \frac{1}{2}, 1, 2+m, -\frac{b \tan(c+dx)}{a}, -i \tan(c+dx)\right) \tan^{1+m}(c+dx) \sqrt{1+\frac{b \tan(c+dx)}{a}}}{2d(1+m) \sqrt{a+b \tan(c+dx)}} + \frac{(A-iB) \operatorname{AppellF1}\left(1+m, \frac{1}{2}, 1, 2+m, -\frac{b \tan(c+dx)}{a}, i \tan(c+dx)\right) \tan^{1+m}(c+dx) \sqrt{1+\frac{b \tan(c+dx)}{a}}}{2d(1+m) \sqrt{a+b \tan(c+dx)}}$$

output

```
1/2*(A+I*B)*AppellF1(1+m,1,1/2,2+m,-I*tan(d*x+c),-b*tan(d*x+c)/a)*(1+b*tan(d*x+c)/a)^(1/2)*tan(d*x+c)^(1+m)/d/(1+m)/(a+b*tan(d*x+c))^(1/2)+1/2*(A-I*B)*AppellF1(1+m,1,1/2,2+m,I*tan(d*x+c),-b*tan(d*x+c)/a)*(1+b*tan(d*x+c)/a)^(1/2)*tan(d*x+c)^(1+m)/d/(1+m)/(a+b*tan(d*x+c))^(1/2)
```

3.490.2 Mathematica [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

input `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]], x]`

output `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]], x]`

3.490.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3042, 4086, 3042, 4085, 152, 150}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\tan(c+dx)^m(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx \\ & \quad \downarrow \text{4086} \\ & \frac{1}{2}(A+iB) \int \frac{(1-i\tan(c+dx))\tan^m(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}(A-iB) \int \frac{(i\tan(c+dx)+1)\tan^m(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \frac{1}{2}(A+iB) \int \frac{(1-i\tan(c+dx))\tan(c+dx)^m}{\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}(A-iB) \int \frac{(i\tan(c+dx)+1)\tan(c+dx)^m}{\sqrt{a+b\tan(c+dx)}} dx \\ & \quad \downarrow \text{4085} \end{aligned}$$

$$\frac{(A - iB) \int \frac{\tan^m(c+dx)}{(1-i \tan(c+dx))\sqrt{a+b \tan(c+dx)}} d \tan(c + dx)}{2d} + \frac{(A + iB) \int \frac{\tan^m(c+dx)}{(i \tan(c+dx)+1)\sqrt{a+b \tan(c+dx)}} d \tan(c + dx)}{2d}$$

↓ 152

$$\frac{(A - iB) \sqrt{\frac{b \tan(c+dx)}{a} + 1} \int \frac{\tan^m(c+dx)}{(1-i \tan(c+dx))\sqrt{\frac{b \tan(c+dx)}{a} + 1}} d \tan(c + dx)}{2d\sqrt{a + b \tan(c + dx)}} + \frac{(A + iB) \sqrt{\frac{b \tan(c+dx)}{a} + 1} \int \frac{\tan^m(c+dx)}{(i \tan(c+dx)+1)\sqrt{\frac{b \tan(c+dx)}{a} + 1}} d \tan(c + dx)}{2d\sqrt{a + b \tan(c + dx)}}$$

↓ 150

$$\frac{(A + iB) \tan^{m+1}(c + dx) \sqrt{\frac{b \tan(c+dx)}{a} + 1} \operatorname{AppellF1}\left(m + 1, \frac{1}{2}, 1, m + 2, -\frac{b \tan(c+dx)}{a}, -i \tan(c + dx)\right)}{2d(m + 1)\sqrt{a + b \tan(c + dx)}} + \frac{(A - iB) \tan^{m+1}(c + dx) \sqrt{\frac{b \tan(c+dx)}{a} + 1} \operatorname{AppellF1}\left(m + 1, \frac{1}{2}, 1, m + 2, -\frac{b \tan(c+dx)}{a}, i \tan(c + dx)\right)}{2d(m + 1)\sqrt{a + b \tan(c + dx)}}$$

input `Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `((A + I*B)*AppellF1[1 + m, 1/2, 1, 2 + m, -((b*Tan[c + d*x])/a), (-I)*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*Sqrt[1 + (b*Tan[c + d*x])/a])/(2*d*(1 + m)*Sqrt[a + b*Tan[c + d*x]]) + ((A - I*B)*AppellF1[1 + m, 1/2, 1, 2 + m, -((b*Tan[c + d*x])/a), I*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*Sqrt[1 + (b*Tan[c + d*x])/a])/(2*d*(1 + m)*Sqrt[a + b*Tan[c + d*x]])`

3.490.3.1 Defintions of rubi rules used

rule 150 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_] := Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])`

rule 152 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4085 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && EqQ[A^2 + B^2, 0]`

rule 4086 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`

3.490.4 Maple [F]

$$\int \frac{\tan(dx+c)^m (A+B \tan(dx+c))}{\sqrt{a+b \tan(dx+c)}} dx$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x)`

output `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x)`

3.490.5 Fricas [F]

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = \int \frac{(B \tan(dx+c) + A) \tan(dx+c)^m}{\sqrt{b \tan(dx+c) + a}} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="fricas")`

output `integral((B*tan(d*x + c) + A)*tan(d*x + c)^m/sqrt(b*tan(d*x + c) + a), x)`

3.490.6 Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = \int \frac{(A+B \tan(c+dx)) \tan^m(c+dx)}{\sqrt{a+b \tan(c+dx)}} dx$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**m/sqrt(a + b*tan(c + d*x)), x)`

3.490.7 Maxima [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{\sqrt{b\tan(dx+c)+a}} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm
m="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/sqrt(b*tan(d*x + c) + a), x)`

3.490.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm
m="giac")`

output `Timed out`

3.490.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{\tan(c+dx)^m(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

input `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x)`

output `int((tan(c + d*x)^m*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2), x)`

3.491
$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.491.1 Optimal result 4666
 3.491.2 Mathematica [F] 4666
 3.491.3 Rubi [A] (verified) 4667
 3.491.4 Maple [F] 4669
 3.491.5 Fricas [F] 4669
 3.491.6 Sympy [F] 4670
 3.491.7 Maxima [F] 4670
 3.491.8 Giac [F(-1)] 4670
 3.491.9 Mupad [F(-1)] 4671

3.491.1 Optimal result

Integrand size = 33, antiderivative size = 193

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{(A+iB) \operatorname{AppellF1}\left(1+m, \frac{3}{2}, 1, 2+m, -\frac{b \tan(c+dx)}{a}, -i \tan(c+dx)\right)}{2ad(1+m)\sqrt{a+b \tan(c+dx)}} + \frac{(A-iB) \operatorname{AppellF1}\left(1+m, \frac{3}{2}, 1, 2+m, -\frac{b \tan(c+dx)}{a}, i \tan(c+dx)\right) \tan^{1+m}(c+dx) \sqrt{1+\frac{b \tan(c+dx)}{a}}}{2ad(1+m)\sqrt{a+b \tan(c+dx)}}$$

```
output 1/2*(A+I*B)*AppellF1(1+m,1,3/2,2+m,-I*tan(d*x+c),-b*tan(d*x+c)/a)*(1+b*tan(d*x+c)/a)^(1/2)*tan(d*x+c)^(1+m)/a/d/(1+m)/(a+b*tan(d*x+c))^(1/2)+1/2*(A-I*B)*AppellF1(1+m,1,3/2,2+m,I*tan(d*x+c),-b*tan(d*x+c)/a)*(1+b*tan(d*x+c)/a)^(1/2)*tan(d*x+c)^(1+m)/a/d/(1+m)/(a+b*tan(d*x+c))^(1/2)
```

3.491.2 Mathematica [F]

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

```
input Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2), x]
```

```
output Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2), x]
```

3.491.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3042, 4086, 3042, 4085, 152, 150}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\tan(c+dx)^m(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{4086} \\
 & \frac{1}{2}(A+iB) \int \frac{(1-i \tan(c+dx)) \tan^m(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx + \frac{1}{2}(A-iB) \int \frac{(i \tan(c+dx)+1) \tan^m(c+dx)}{(a+b \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2}(A+iB) \int \frac{(1-i \tan(c+dx)) \tan(c+dx)^m}{(a+b \tan(c+dx))^{3/2}} dx + \frac{1}{2}(A-iB) \int \frac{(i \tan(c+dx)+1) \tan(c+dx)^m}{(a+b \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{4085} \\
 & \frac{(A-iB) \int \frac{\tan^m(c+dx)}{(1-i \tan(c+dx))(a+b \tan(c+dx))^{3/2}} d \tan(c+dx)}{2d} + \\
 & \frac{(A+iB) \int \frac{\tan^m(c+dx)}{(i \tan(c+dx)+1)(a+b \tan(c+dx))^{3/2}} d \tan(c+dx)}{2d} \\
 & \quad \downarrow \text{152} \\
 & \frac{(A-iB) \sqrt{\frac{b \tan(c+dx)}{a} + 1} \int \frac{\tan^m(c+dx)}{(1-i \tan(c+dx)) \left(\frac{b \tan(c+dx)}{a} + 1\right)^{3/2}} d \tan(c+dx)}{2ad \sqrt{a+b \tan(c+dx)}} + \\
 & \frac{(A+iB) \sqrt{\frac{b \tan(c+dx)}{a} + 1} \int \frac{\tan^m(c+dx)}{(i \tan(c+dx)+1) \left(\frac{b \tan(c+dx)}{a} + 1\right)^{3/2}} d \tan(c+dx)}{2ad \sqrt{a+b \tan(c+dx)}} \\
 & \quad \downarrow \text{150}
 \end{aligned}$$

$$\frac{(A + iB) \tan^{m+1}(c + dx) \sqrt{\frac{b \tan(c+dx)}{a} + 1} \operatorname{AppellF1}\left(m + 1, \frac{3}{2}, 1, m + 2, -\frac{b \tan(c+dx)}{a}, -i \tan(c + dx)\right)}{2ad(m + 1) \sqrt{a + b \tan(c + dx)}} +$$

$$\frac{(A - iB) \tan^{m+1}(c + dx) \sqrt{\frac{b \tan(c+dx)}{a} + 1} \operatorname{AppellF1}\left(m + 1, \frac{3}{2}, 1, m + 2, -\frac{b \tan(c+dx)}{a}, i \tan(c + dx)\right)}{2ad(m + 1) \sqrt{a + b \tan(c + dx)}}$$

input `Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `((A + I*B)*AppellF1[1 + m, 3/2, 1, 2 + m, -((b*Tan[c + d*x])/a), (-I)*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*Sqrt[1 + (b*Tan[c + d*x])/a]/(2*a*d*(1 + m)*Sqrt[a + b*Tan[c + d*x]]) + ((A - I*B)*AppellF1[1 + m, 3/2, 1, 2 + m, -(b*Tan[c + d*x])/a, I*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*Sqrt[1 + (b*Tan[c + d*x])/a]/(2*a*d*(1 + m)*Sqrt[a + b*Tan[c + d*x]])`

3.491.3.1 Defintions of rubi rules used

rule 150 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_] := Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])`

rule 152 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n])Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4085 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && EqQ[A^2 + B^2, 0]`

rule 4086 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`

3.491.4 Maple [F]

$$\int \frac{\tan(dx + c)^m (A + B \tan(dx + c))}{(a + b \tan(dx + c))^{\frac{3}{2}}} dx$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x)`

output `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x)`

3.491.5 Fracas [F]

$$\int \frac{\tan^m(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)^m}{(b \tan(dx + c) + a)^{\frac{3}{2}}} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm m="fricas")`

output `integral((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*tan(d*x + c)^m/(b^2*tan(d*x + c)^2 + 2*a*b*tan(d*x + c) + a^2), x)`

3.491.6 Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(A+B\tan(c+dx))\tan^m(c+dx)}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**m/(a + b*tan(c + d*x))**(3/2), x)`

3.491.7 Maxima [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{(b\tan(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(b*tan(d*x + c) + a)^(3/2), x)`

3.491.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `Timed out`

3.491.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{\tan(c+dx)^m (A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$$

input `int((tan(c + d*x))^m*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x)`

output `int((tan(c + d*x))^m*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2), x)`

3.492
$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

3.492.1 Optimal result 4672
 3.492.2 Mathematica [F] 4672
 3.492.3 Rubi [A] (verified) 4673
 3.492.4 Maple [F] 4675
 3.492.5 Fracas [F] 4675
 3.492.6 Sympy [F] 4676
 3.492.7 Maxima [F] 4676
 3.492.8 Giac [F(-1)] 4676
 3.492.9 Mupad [F(-1)] 4677

3.492.1 Optimal result

Integrand size = 33, antiderivative size = 193

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{(A+iB) \operatorname{AppellF1}\left(1+m, \frac{5}{2}, 1, 2+m, -\frac{b \tan(c+dx)}{a}, -i \tan(c+dx)\right)}{2a^2 d(1+m) \sqrt{a+b \tan(c+dx)}} + \frac{(A-iB) \operatorname{AppellF1}\left(1+m, \frac{5}{2}, 1, 2+m, -\frac{b \tan(c+dx)}{a}, i \tan(c+dx)\right) \tan^{1+m}(c+dx) \sqrt{1+\frac{b \tan(c+dx)}{a}}}{2a^2 d(1+m) \sqrt{a+b \tan(c+dx)}}$$

output `1/2*(A+I*B)*AppellF1(1+m,1,5/2,2+m,-I*tan(d*x+c),-b*tan(d*x+c)/a)*(1+b*tan(d*x+c)/a)^(1/2)*tan(d*x+c)^(1+m)/a^2/d/(1+m)/(a+b*tan(d*x+c))^(1/2)+1/2*(A-I*B)*AppellF1(1+m,1,5/2,2+m,I*tan(d*x+c),-b*tan(d*x+c)/a)*(1+b*tan(d*x+c)/a)^(1/2)*tan(d*x+c)^(1+m)/a^2/d/(1+m)/(a+b*tan(d*x+c))^(1/2)`

3.492.2 Mathematica [F]

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

input `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2), x]`

output `Integrate[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2), x]`

3.492.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3042, 4086, 3042, 4085, 152, 150}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\tan^m(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{\tan(c+dx)^m(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

↓ 4086

$$\frac{1}{2}(A+iB) \int \frac{(1-i \tan(c+dx)) \tan^m(c+dx)}{(a+b \tan(c+dx))^{5/2}} dx + \frac{1}{2}(A-iB) \int \frac{(i \tan(c+dx)+1) \tan^m(c+dx)}{(a+b \tan(c+dx))^{5/2}} dx$$

↓ 3042

$$\frac{1}{2}(A+iB) \int \frac{(1-i \tan(c+dx)) \tan(c+dx)^m}{(a+b \tan(c+dx))^{5/2}} dx + \frac{1}{2}(A-iB) \int \frac{(i \tan(c+dx)+1) \tan(c+dx)^m}{(a+b \tan(c+dx))^{5/2}} dx$$

↓ 4085

$$\frac{(A-iB) \int \frac{\tan^m(c+dx)}{(1-i \tan(c+dx))(a+b \tan(c+dx))^{5/2}} d \tan(c+dx)}{2d} + \frac{(A+iB) \int \frac{\tan^m(c+dx)}{(i \tan(c+dx)+1)(a+b \tan(c+dx))^{5/2}} d \tan(c+dx)}{2d}$$

↓ 152

$$\frac{(A-iB) \sqrt{\frac{b \tan(c+dx)}{a} + 1} \int \frac{\tan^m(c+dx)}{(1-i \tan(c+dx)) \left(\frac{b \tan(c+dx)}{a} + 1\right)^{5/2}} d \tan(c+dx)}{2a^2 d \sqrt{a+b \tan(c+dx)}} + \frac{(A+iB) \sqrt{\frac{b \tan(c+dx)}{a} + 1} \int \frac{\tan^m(c+dx)}{(i \tan(c+dx)+1) \left(\frac{b \tan(c+dx)}{a} + 1\right)^{5/2}} d \tan(c+dx)}{2a^2 d \sqrt{a+b \tan(c+dx)}}$$

↓ 150

$$\frac{(A + iB) \tan^{m+1}(c + dx) \sqrt{\frac{b \tan(c+dx)}{a} + 1} \operatorname{AppellF1}\left(m + 1, \frac{5}{2}, 1, m + 2, -\frac{b \tan(c+dx)}{a}, -i \tan(c + dx)\right)}{2a^2 d(m + 1) \sqrt{a + b \tan(c + dx)}} +$$

$$\frac{(A - iB) \tan^{m+1}(c + dx) \sqrt{\frac{b \tan(c+dx)}{a} + 1} \operatorname{AppellF1}\left(m + 1, \frac{5}{2}, 1, m + 2, -\frac{b \tan(c+dx)}{a}, i \tan(c + dx)\right)}{2a^2 d(m + 1) \sqrt{a + b \tan(c + dx)}}$$

input `Int[(Tan[c + d*x]^m*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x]`

output `((A + I*B)*AppellF1[1 + m, 5/2, 1, 2 + m, -((b*Tan[c + d*x])/a), (-I)*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*Sqrt[1 + (b*Tan[c + d*x])/a])/(2*a^2*d*(1 + m)*Sqrt[a + b*Tan[c + d*x]]) + ((A - I*B)*AppellF1[1 + m, 5/2, 1, 2 + m, -((b*Tan[c + d*x])/a), I*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*Sqrt[1 + (b*Tan[c + d*x])/a])/(2*a^2*d*(1 + m)*Sqrt[a + b*Tan[c + d*x]])`

3.492.3.1 Defintions of rubi rules used

rule 150 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_] := Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])`

rule 152 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n])Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4085 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && EqQ[A^2 + B^2, 0]`

rule 4086 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`

3.492.4 Maple [F]

$$\int \frac{\tan(dx + c)^m (A + B \tan(dx + c))}{(a + b \tan(dx + c))^{\frac{5}{2}}} dx$$

input `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x)`

output `int(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x)`

3.492.5 Fracas [F]

$$\int \frac{\tan^m(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{5/2}} dx = \int \frac{(B \tan(dx + c) + A) \tan(dx + c)^m}{(b \tan(dx + c) + a)^{\frac{5}{2}}} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm m="fricas")`

output `integral((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*tan(d*x + c)^m/(b^3*tan(d*x + c)^3 + 3*a*b^2*tan(d*x + c)^2 + 3*a^2*b*tan(d*x + c) + a^3), x)`

3.492.6 Sympy [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\tan^m(c+dx)}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx$$

input `integrate(tan(d*x+c)**m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*tan(c + d*x)**m/(a + b*tan(c + d*x))**(5/2), x)`

3.492.7 Maxima [F]

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(B\tan(dx+c)+A)\tan(dx+c)^m}{(b\tan(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*tan(d*x + c)^m/(b*tan(d*x + c) + a)^(5/2), x)`

3.492.8 Giac [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(tan(d*x+c)^m*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `Timed out`

3.492.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^m(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{\tan(c+dx)^m (A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$$

input `int((tan(c + d*x))^m*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)`output `int((tan(c + d*x))^m*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2), x)`

3.493 $\int \tan^m(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.493.1 Optimal result	4678
3.493.2 Mathematica [F]	4679
3.493.3 Rubi [A] (verified)	4679
3.493.4 Maple [F]	4681
3.493.5 Fricas [F]	4681
3.493.6 Sympy [F]	4682
3.493.7 Maxima [F]	4682
3.493.8 Giac [F]	4682
3.493.9 Mupad [F(-1)]	4683

3.493.1 Optimal result

Integrand size = 31, antiderivative size = 183

$$\int \tan^m(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \frac{(A+iB) \operatorname{AppellF1}\left(1+m, -n, 1, 2+m, -\frac{b \tan(c+dx)}{a}, -i \tan(c+dx)\right) \tan^{1+m}(c+dx)(a+b \tan(c+dx))}{2d(1+m)} + \frac{(A-iB) \operatorname{AppellF1}\left(1+m, -n, 1, 2+m, -\frac{b \tan(c+dx)}{a}, i \tan(c+dx)\right) \tan^{1+m}(c+dx)(a+b \tan(c+dx))}{2d(1+m)}$$

output

```
1/2*(A+I*B)*AppellF1(1+m,1,-n,2+m,-I*tan(d*x+c),-b*tan(d*x+c)/a)*tan(d*x+c)^(1+m)*(a+b*tan(d*x+c))^n/d/(1+m)/((1+b*tan(d*x+c)/a)^n)+1/2*(A-I*B)*AppellF1(1+m,1,-n,2+m,I*tan(d*x+c),-b*tan(d*x+c)/a)*tan(d*x+c)^(1+m)*(a+b*tan(d*x+c))^n/d/(1+m)/((1+b*tan(d*x+c)/a)^n)
```

3.493.2 Mathematica [F]

$$\int \tan^m(c+dx)(a+b\tan(c+dx))^n(A+B\tan(c+dx)) dx$$

$$= \int \tan^m(c+dx)(a+b\tan(c+dx))^n(A+B\tan(c+dx)) dx$$

input `Integrate[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

output `Integrate[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

3.493.3 Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.194$, Rules used = {3042, 4086, 3042, 4085, 152, 150}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^m(c+dx)(A+B\tan(c+dx))(a+b\tan(c+dx))^n dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c+dx)^m(A+B\tan(c+dx))(a+b\tan(c+dx))^n dx$$

$$\downarrow \text{4086}$$

$$\frac{1}{2}(A+iB) \int (1-i\tan(c+dx)) \tan^m(c+dx)(a+b\tan(c+dx))^n dx + \frac{1}{2}(A-iB) \int (i\tan(c+dx) + 1) \tan^m(c+dx)(a+b\tan(c+dx))^n dx$$

$$\downarrow \text{3042}$$

$$\frac{1}{2}(A+iB) \int (1-i\tan(c+dx)) \tan(c+dx)^m(a+b\tan(c+dx))^n dx + \frac{1}{2}(A-iB) \int (i\tan(c+dx) + 1) \tan(c+dx)^m(a+b\tan(c+dx))^n dx$$

$$\downarrow \text{4085}$$

$$\begin{aligned}
& \frac{(A - iB) \int \frac{\tan^m(c+dx)(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d \tan(c+dx)}{2d} + \\
& \frac{(A + iB) \int \frac{\tan^m(c+dx)(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d \tan(c+dx)}{2d} \\
& \quad \downarrow \text{152} \\
& \frac{(A - iB)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1 \right)^{-n} \int \frac{\tan^m(c+dx) \left(\frac{b \tan(c+dx)}{a} + 1 \right)^n}{1-i \tan(c+dx)} d \tan(c+dx)}{2d} + \\
& \frac{(A + iB)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1 \right)^{-n} \int \frac{\tan^m(c+dx) \left(\frac{b \tan(c+dx)}{a} + 1 \right)^n}{i \tan(c+dx)+1} d \tan(c+dx)}{2d} \\
& \quad \downarrow \text{150} \\
& \frac{(A + iB) \tan^{m+1}(c + dx)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1 \right)^{-n} \text{AppellF1} \left(m + 1, -n, 1, m + 2, -\frac{b \tan(c+dx)}{a}, -i \right)}{2d(m + 1)} + \\
& \frac{(A - iB) \tan^{m+1}(c + dx)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1 \right)^{-n} \text{AppellF1} \left(m + 1, -n, 1, m + 2, -\frac{b \tan(c+dx)}{a}, i \right)}{2d(m + 1)}
\end{aligned}$$

input `Int[Tan[c + d*x]^m*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `((A + I*B)*AppellF1[1 + m, -n, 1, 2 + m, -((b*Tan[c + d*x])/a), (-I)*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*(a + b*Tan[c + d*x])^n)/(2*d*(1 + m)*(1 + (b*Tan[c + d*x])/a)^n) + ((A - I*B)*AppellF1[1 + m, -n, 1, 2 + m, -((b*Tan[c + d*x])/a), I*Tan[c + d*x]]*Tan[c + d*x]^(1 + m)*(a + b*Tan[c + d*x])^n)/(2*d*(1 + m)*(1 + (b*Tan[c + d*x])/a)^n)`

3.493.3.1 Defintions of rubi rules used

rule 150 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_] := Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])`

rule 152 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4085 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && EqQ[A^2 + B^2, 0]`

rule 4086 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`

3.493.4 Maple [F]

$$\int \tan(dx + c)^m (a + b \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^m*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^m*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.493.5 Fracas [F]

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^m dx \end{aligned}$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output `integral((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*tan(d*x + c)^m, x)`

3.493.6 Sympy [F]

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^n \tan^m(c + dx) dx$$

input `integrate(tan(d*x+c)**m*(a+b*tan(d*x+c))**n*(A+B*tan(d*x+c)), x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**n*tan(c + d*x)**m, x)`

3.493.7 Maxima [F]

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)), x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*tan(d*x + c)^m, x)`

3.493.8 Giac [F]

$$\int \tan^m(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^m dx$$

input `integrate(tan(d*x+c)^m*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*tan(d*x + c)^m, x)`

3.493.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \tan^m(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int \tan(c + dx)^m (A + B \tan(c + dx)) (a + b \tan(c + dx))^n dx \end{aligned}$$

input `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n,x)`

output `int(tan(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n, x)`

3.494 $\int \tan^4(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.494.1 Optimal result	4684
3.494.2 Mathematica [A] (verified)	4685
3.494.3 Rubi [A] (verified)	4686
3.494.4 Maple [F]	4690
3.494.5 Fricas [F]	4691
3.494.6 Sympy [F]	4691
3.494.7 Maxima [F]	4691
3.494.8 Giac [F]	4692
3.494.9 Mupad [F(-1)]	4692

3.494.1 Optimal result

Integrand size = 31, antiderivative size = 387

$$\int \tan^4(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx =$$

$$\frac{(Ab^3(2+n)(3+n)(4+n) - a(b^2B(3+n)(4+n) - 2a(3aB - Ab(4+n))))(a+b \tan(c+dx))^{1+n}}{b^4d(1+n)(2+n)(3+n)(4+n)}$$

$$+ \frac{(A - iB) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a-ib}\right)(a+b \tan(c+dx))^{1+n}}{2(ia+b)d(1+n)}$$

$$- \frac{(A + iB) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a+ib}\right)(a+b \tan(c+dx))^{1+n}}{2(ia-b)d(1+n)}$$

$$- \frac{(b^2B(3+n)(4+n) - 2a(3aB - Ab(4+n))) \tan(c+dx)(a+b \tan(c+dx))^{1+n}}{b^3d(2+n)(3+n)(4+n)}$$

$$- \frac{(3aB - Ab(4+n)) \tan^2(c+dx)(a+b \tan(c+dx))^{1+n}}{b^2d(3+n)(4+n)}$$

$$+ \frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{1+n}}{bd(4+n)}$$

output $-(A*b^3*(2+n)*(3+n)*(4+n)-a*(b^2*B*(3+n)*(4+n)-2*a*(3*B*a-A*b*(4+n)))*(a+b*\tan(dx+c))^{(1+n)}/b^4/d/(1+n)/(2+n)/(3+n)/(4+n)+1/2*(A-I*B)*\text{hypergeom}([1, 1+n], [2+n], (a+b*\tan(dx+c))/(a-I*b))*(a+b*\tan(dx+c))^{(1+n)}/(I*a+b)/d/(1+n)-1/2*(A+I*B)*\text{hypergeom}([1, 1+n], [2+n], (a+b*\tan(dx+c))/(a+I*b))*(a+b*\tan(dx+c))^{(1+n)}/(I*a-b)/d/(1+n)-(b^2*B*(3+n)*(4+n)-2*a*(3*B*a-A*b*(4+n)))*\tan(dx+c)*(a+b*\tan(dx+c))^{(1+n)}/b^3/d/(2+n)/(3+n)/(4+n)-(3*B*a-A*b*(4+n))*\tan(dx+c)^2*(a+b*\tan(dx+c))^{(1+n)}/b^2/d/(3+n)/(4+n)+B*\tan(dx+c)^3*(a+b*\tan(dx+c))^{(1+n)}/b/d/(4+n)$

3.494.2 Mathematica [A] (verified)

Time = 6.10 (sec) , antiderivative size = 384, normalized size of antiderivative = 0.99

$$\int \tan^4(c+dx)(a+b\tan(c+dx))^n(A+B\tan(c+dx)) dx$$

$$= \frac{(a+b\tan(c+dx))^{1+n} \left(i \left(2i(a-ib)(a+ib)(6a^3B-2a^2Ab(4+n)-ab^2B(3+n)(4+n)+Ab^3(2+n) \right) \right)}{}$$

input `Integrate[Tan[c + d*x]^4*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output $((a + b*\text{Tan}[c + d*x])^{(1 + n)}*(I*((2*I)*(a - I*b)*(a + I*b)*(6*a^3*B - 2*a^2*A*b*(4 + n) - a*b^2*B*(3 + n)*(4 + n) + A*b^3*(2 + n)*(3 + n)*(4 + n)) - (a + I*b)*b^4*(A - I*B)*(24 + 26*n + 9*n^2 + n^3)*\text{Hypergeometric2F1}[1, 1 + n, 2 + n, (a + b*\text{Tan}[c + d*x])/(a - I*b)] + (a - I*b)*b^4*(A + I*B)*(24 + 26*n + 9*n^2 + n^3)*\text{Hypergeometric2F1}[1, 1 + n, 2 + n, (a + b*\text{Tan}[c + d*x])/(a + I*b)]) + 2*(a - I*b)*(a + I*b)*b*(1 + n)*(6*a^2*B - 2*a*A*b*(4 + n) - b^2*B*(3 + n)*(4 + n))*\text{Tan}[c + d*x] - 2*(a - I*b)*(a + I*b)*b^2*(1 + n)*(2 + n)*(3*a*B - A*b*(4 + n))*\text{Tan}[c + d*x]^2 + 2*(a - I*b)*(a + I*b)*b^3*B*(1 + n)*(2 + n)*(3 + n)*\text{Tan}[c + d*x]^3))/(2*(a - I*b)*(a + I*b)*b^4*d*(1 + n)*(2 + n)*(3 + n)*(4 + n))$

3.494.3 Rubi [A] (verified)

Time = 1.97 (sec) , antiderivative size = 412, normalized size of antiderivative = 1.06, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.548$, Rules used = {3042, 4090, 25, 3042, 4130, 25, 3042, 4130, 25, 3042, 4113, 3042, 4022, 3042, 4020, 25, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \tan^4(c+dx)(A+B \tan(c+dx))(a+b \tan(c+dx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan(c+dx)^4(A+B \tan(c+dx))(a+b \tan(c+dx))^n dx \\
 & \quad \downarrow \text{4090} \\
 & \frac{\int -\tan^2(c+dx)(a+b \tan(c+dx))^n ((3aB - Ab(n+4)) \tan^2(c+dx) + bB(n+4) \tan(c+dx) + 3aB) dx}{b(n+4)} + \\
 & \quad \frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} \\
 & \quad \downarrow \text{25} \\
 & \frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \\
 & \frac{\int \tan^2(c+dx)(a+b \tan(c+dx))^n ((3aB - Ab(n+4)) \tan^2(c+dx) + bB(n+4) \tan(c+dx) + 3aB) dx}{b(n+4)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \\
 & \frac{\int \tan(c+dx)^2(a+b \tan(c+dx))^n ((3aB - Ab(n+4)) \tan(c+dx)^2 + bB(n+4) \tan(c+dx) + 3aB) dx}{b(n+4)} \\
 & \quad \downarrow \text{4130} \\
 & \frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \\
 & \frac{\int -\tan(c+dx)(a+b \tan(c+dx))^n (-A(n+3)(n+4) \tan(c+dx)b^2 + (6Ba^2 - 2Ab(n+4)a - b^2B(n+3)(n+4)) \tan^2(c+dx) + 2a(3aB - Ab(n+4))) dx}{b(n+3)} + \\
 & \quad \frac{ }{b(n+4)} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

3.494. $\int \tan^4(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

$$\frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \frac{\tan^2(c+dx)(3aB-Ab(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \frac{\int \tan(c+dx)(a+b \tan(c+dx))^n (-A(n+3)(n+4) \tan(c+dx)b^2 + (6Ba^2-2Ab(n+4)a-b^2B(n+3)(n+4)) \tan(c+dx)b^3 + (6Ba^3-2Ab(n+4)a^2-b^2B(n+3)(n+4)a-b^3B(n+2)(n+3)(n+4)) \tan(c+dx)b^4}{b(n+3)}}{b(n+4)}$$

↓ 3042

$$\frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \frac{\tan^2(c+dx)(3aB-Ab(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \frac{\int \tan(c+dx)(a+b \tan(c+dx))^n (-A(n+3)(n+4) \tan(c+dx)b^2 + (6Ba^2-2Ab(n+4)a-b^2B(n+3)(n+4)) \tan(c+dx)b^3 + (6Ba^3-2Ab(n+4)a^2-b^2B(n+3)(n+4)a-b^3B(n+2)(n+3)(n+4)) \tan(c+dx)b^4}{b(n+3)}}{b(n+4)}$$

↓ 4130

$$\frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \frac{\tan^2(c+dx)(3aB-Ab(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \frac{\int -(a+b \tan(c+dx))^n (-B(n+2)(n+3)(n+4) \tan(c+dx)b^3 + (6Ba^3-2Ab(n+4)a^2-b^2B(n+3)(n+4)a-b^3B(n+2)(n+3)(n+4)) \tan(c+dx)b^4)}{b(n+2)}}{b(n+4)}$$

↓ 25

$$\frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \frac{\tan^2(c+dx)(3aB-Ab(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \frac{\tan(c+dx)(6a^2B-2aAb(n+4)-b^2B(n+3)(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{\int (a+b \tan(c+dx))^n (-B(n+2)(n+3)(n+4) \tan(c+dx)b^3 + (6Ba^3-2Ab(n+4)a^2-b^2B(n+3)(n+4)a-b^3B(n+2)(n+3)(n+4)) \tan(c+dx)b^4)}{b(n+2)}}{b(n+4)}$$

↓ 3042

$$\frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \frac{\tan^2(c+dx)(3aB-Ab(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \frac{\tan(c+dx)(6a^2B-2aAb(n+4)-b^2B(n+3)(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{\int (a+b \tan(c+dx))^n (-B(n+2)(n+3)(n+4) \tan(c+dx)b^3 + (6Ba^3-2Ab(n+4)a^2-b^2B(n+3)(n+4)a-b^3B(n+2)(n+3)(n+4)) \tan(c+dx)b^4)}{b(n+2)}}{b(n+4)}$$

↓ 4113

$$\frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \frac{\tan^2(c+dx)(3aB-Ab(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \frac{\tan(c+dx)(6a^2B-2aAb(n+4)-b^2B(n+3)(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{\int (a+b \tan(c+dx))^n (-A(n+3)(n+4) \tan(c+dx)b^2 + (6Ba^2-2Ab(n+4)a-b^2B(n+3)(n+4)) \tan(c+dx)b^3 + (6Ba^3-2Ab(n+4)a^2-b^2B(n+3)(n+4)a-b^3B(n+2)(n+3)(n+4)) \tan(c+dx)b^4)}{b(n+2)}}{b(n+4)}$$

↓ 3042

3.494. $\int \tan^4(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

$$\frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \frac{\tan^2(c+dx)(3aB-Ab(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \frac{\tan(c+dx)(6a^2B-2aAb(n+4)-b^2B(n+3)(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{\int(a+b \tan(c+dx))^n(-A(n+1)+B)dx}{b(n+4)}$$

↓ 4022

$$\frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \frac{\tan^2(c+dx)(3aB-Ab(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \frac{\tan(c+dx)(6a^2B-2aAb(n+4)-b^2B(n+3)(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{1}{2}b^3(n+2)(n+3)(n+4)(A+B)$$

↓ 3042

$$\frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \frac{\tan^2(c+dx)(3aB-Ab(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \frac{\tan(c+dx)(6a^2B-2aAb(n+4)-b^2B(n+3)(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{1}{2}b^3(n+2)(n+3)(n+4)(A+B)$$

↓ 4020

$$\frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \frac{\tan^2(c+dx)(3aB-Ab(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \frac{\tan(c+dx)(6a^2B-2aAb(n+4)-b^2B(n+3)(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{ib^3(n+2)(n+3)(n+4)(A-iB)}{b(n+4)}$$

↓ 25

$$\frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \frac{\tan^2(c+dx)(3aB-Ab(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \frac{\tan(c+dx)(6a^2B-2aAb(n+4)-b^2B(n+3)(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{ib^3(n+2)(n+3)(n+4)(A-iB)}{b(n+4)}$$

↓ 78

$$\frac{B \tan^3(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+4)} - \frac{\tan^2(c+dx)(3aB-Ab(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \frac{\tan(c+dx)(6a^2B-2aAb(n+4)-b^2B(n+3)(n+4))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{(6a^3B-2a^2Ab(n+4)-ab^2B)}{b(n+4)}$$

3.494. $\int \tan^4(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

input `Int[Tan[c + d*x]^4*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `(B*Tan[c + d*x]^3*(a + b*Tan[c + d*x])^(1 + n))/(b*d*(4 + n)) - (((3*a*B - A*b*(4 + n))*Tan[c + d*x]^2*(a + b*Tan[c + d*x])^(1 + n))/(b*d*(3 + n)) - (((6*a^2*B - 2*a*A*b*(4 + n) - b^2*B*(3 + n)*(4 + n))*Tan[c + d*x]*(a + b*Tan[c + d*x])^(1 + n))/(b*d*(2 + n)) - (((6*a^3*B - 2*a^2*A*b*(4 + n) - a*b^2*B*(3 + n)*(4 + n) + A*b^3*(2 + n)*(3 + n)*(4 + n))*(a + b*Tan[c + d*x])^(1 + n))/(b*d*(1 + n)) + ((I/2)*b^3*(A - I*B)*(2 + n)*(3 + n)*(4 + n)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a - I*b)]*(a + b*Tan[c + d*x])^(1 + n))/((a - I*b)*d*(1 + n)) - ((I/2)*b^3*(A + I*B)*(2 + n)*(3 + n)*(4 + n)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a + I*b)]*(a + b*Tan[c + d*x])^(1 + n))/((a + I*b)*d*(1 + n)))/(b*(2 + n))/(b*(3 + n))/(b*(4 + n))`

3.494.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.494.4 Maple [F]

$$\int \tan(dx + c)^4 (a + b \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^4*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^4*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.494.5 Fricas [F]

$$\int \tan^4(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^4 dx$$

input `integrate(tan(d*x+c)^4*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*tan(d*x + c)^5 + A*tan(d*x + c)^4)*(b*tan(d*x + c) + a)^n, x)`

3.494.6 Sympy [F]

$$\int \tan^4(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^n \tan^4(c + dx) dx$$

input `integrate(tan(d*x+c)**4*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n*tan(c + d*x)**4, x)`

3.494.7 Maxima [F]

$$\int \tan^4(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^4 dx$$

input `integrate(tan(d*x+c)^4*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*tan(d*x + c)^4, x)`

3.494.8 Giac [F]

$$\int \tan^4(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^4 dx$$

input `integrate(tan(d*x+c)^4*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*tan(d*x + c)^4, x)`

3.494.9 Mupad [F(-1)]

Timed out.

$$\int \tan^4(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^4 (A + B \tan(c + dx)) (a + b \tan(c + dx))^n dx$$

input `int(tan(c + d*x)^4*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n,x)`

output `int(tan(c + d*x)^4*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n, x)`

3.495 $\int \tan^3(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.495.1 Optimal result	4693
3.495.2 Mathematica [A] (verified)	4694
3.495.3 Rubi [A] (verified)	4694
3.495.4 Maple [F]	4698
3.495.5 Fricas [F]	4699
3.495.6 Sympy [F]	4699
3.495.7 Maxima [F]	4699
3.495.8 Giac [F]	4700
3.495.9 Mupad [F(-1)]	4700

3.495.1 Optimal result

Integrand size = 31, antiderivative size = 291

$$\int \tan^3(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \frac{(2a^2B - aAb(3+n) - b^2B(6+5n+n^2))(a+b \tan(c+dx))^{1+n}}{b^3d(1+n)(2+n)(3+n)}$$

$$+ \frac{(iA+B) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a-ib}\right)(a+b \tan(c+dx))^{1+n}}{2(ia+b)d(1+n)}$$

$$+ \frac{(A+iB) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a+ib}\right)(a+b \tan(c+dx))^{1+n}}{2(a+ib)d(1+n)}$$

$$- \frac{(2aB - Ab(3+n)) \tan(c+dx)(a+b \tan(c+dx))^{1+n}}{b^2d(2+n)(3+n)}$$

$$+ \frac{B \tan^2(c+dx)(a+b \tan(c+dx))^{1+n}}{bd(3+n)}$$

output

```
(2*B*a^2-a*A*b*(3+n)-b^2*B*(n^2+5*n+6))*(a+b*tan(d*x+c))^(1+n)/b^3/d/(1+n)
/(2+n)/(3+n)+1/2*(I*A+B)*hypergeom([1, 1+n], [2+n], (a+b*tan(d*x+c))/(a-I*b))
*(a+b*tan(d*x+c))^(1+n)/(I*a+b)/d/(1+n)+1/2*(A+I*B)*hypergeom([1, 1+n], [2
+n], (a+b*tan(d*x+c))/(a+I*b))*(a+b*tan(d*x+c))^(1+n)/(a+I*b)/d/(1+n)-(2*B*
a-A*b*(3+n))*tan(d*x+c)*(a+b*tan(d*x+c))^(1+n)/b^2/d/(2+n)/(3+n)+B*tan(d*x
+c)^2*(a+b*tan(d*x+c))^(1+n)/b/d/(3+n)
```

3.495.2 Mathematica [A] (verified)

Time = 2.61 (sec) , antiderivative size = 281, normalized size of antiderivative = 0.97

$$\int \tan^3(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \frac{(a + b \tan(c + dx))^{1+n} \left(2(a - ib)(a + ib) (2a^2B - aAb(3 + n) - b^2B(2 + n)(3 + n)) + (a + ib)b^3(A - iB) \right)}{...}$$

input `Integrate[Tan[c + d*x]^3*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `((a + b*Tan[c + d*x])^(1 + n)*(2*(a - I*b)*(a + I*b)*(2*a^2*B - a*A*b*(3 + n) - b^2*B*(2 + n)*(3 + n)) + (a + I*b)*b^3*(A - I*B)*(6 + 5*n + n^2)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a - I*b)] + (a - I*b)*b^3*(A + I*B)*(6 + 5*n + n^2)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a + I*b)] - 2*(a - I*b)*(a + I*b)*b*(1 + n)*(2*a*B - A*b*(3 + n))*Tan[c + d*x] + 2*(a - I*b)*(a + I*b)*b^2*B*(1 + n)*(2 + n)*Tan[c + d*x]^2)/(2*(a - I*b)*(a + I*b)*b^3*d*(1 + n)*(2 + n)*(3 + n))`

3.495.3 Rubi [A] (verified)

Time = 1.29 (sec) , antiderivative size = 319, normalized size of antiderivative = 1.10, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 4090, 25, 3042, 4130, 25, 3042, 4113, 3042, 4022, 3042, 4020, 25, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^3(c + dx)(A + B \tan(c + dx))(a + b \tan(c + dx))^n dx$$

↓ 3042

$$\int \tan(c + dx)^3(A + B \tan(c + dx))(a + b \tan(c + dx))^n dx$$

↓ 4090

$$\frac{\int -\tan(c + dx)(a + b \tan(c + dx))^n ((2aB - Ab(n + 3)) \tan^2(c + dx) + bB(n + 3) \tan(c + dx) + 2aB) dx}{b(n + 3)} + \frac{B \tan^2(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 3)}$$

$$\begin{aligned}
 & \downarrow 25 \\
 & \frac{B \tan^2(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \\
 & \frac{\int \tan(c+dx)(a+b \tan(c+dx))^n ((2aB - Ab(n+3)) \tan^2(c+dx) + bB(n+3) \tan(c+dx) + 2aB) dx}{b(n+3)} \\
 & \downarrow 3042 \\
 & \frac{B \tan^2(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \\
 & \frac{\int \tan(c+dx)(a+b \tan(c+dx))^n ((2aB - Ab(n+3)) \tan(c+dx)^2 + bB(n+3) \tan(c+dx) + 2aB) dx}{b(n+3)} \\
 & \downarrow 4130 \\
 & \frac{B \tan^2(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \\
 & \frac{\int -(a+b \tan(c+dx))^n (-A(n+2)(n+3) \tan(c+dx)b^2 + (2Ba^2 - Ab(n+3)a - b^2B(n+2)(n+3)) \tan^2(c+dx) + a(2aB - Ab(n+3))) dx}{b(n+2)} + \frac{\tan(c+dx)(2aB - Ab(n+3))}{b(n+3)} \\
 & \downarrow 25 \\
 & \frac{B \tan^2(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \\
 & \frac{\frac{\tan(c+dx)(2aB - Ab(n+3))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{\int (a+b \tan(c+dx))^n (-A(n+2)(n+3) \tan(c+dx)b^2 + (2Ba^2 - Ab(n+3)a - b^2B(n+2)(n+3)) \tan^2(c+dx) + a(2aB - Ab(n+3))) dx}{b(n+2)}}{b(n+3)} \\
 & \downarrow 3042 \\
 & \frac{B \tan^2(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \\
 & \frac{\frac{\tan(c+dx)(2aB - Ab(n+3))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{\int (a+b \tan(c+dx))^n (-A(n+2)(n+3) \tan(c+dx)b^2 + (2Ba^2 - Ab(n+3)a - b^2B(n+2)(n+3)) \tan^2(c+dx) + a(2aB - Ab(n+3))) dx}{b(n+2)}}{b(n+3)} \\
 & \downarrow 4113 \\
 & \frac{B \tan^2(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+3)} - \\
 & \frac{\frac{\tan(c+dx)(2aB - Ab(n+3))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{\int (a+b \tan(c+dx))^n (b^2B(n+2)(n+3) - Ab^2(n+2)(n+3) \tan(c+dx)) dx + \frac{(2a^2B - aAb(n+3) - b^2)}{b(n+2)}}{b(n+2)}}{b(n+3)} \\
 & \downarrow 3042
 \end{aligned}$$

$$\frac{B \tan^2(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 3)} - \frac{\frac{\tan(c+dx)(2aB - Ab(n+3))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{\int (a+b \tan(c+dx))^n (b^2 B(n+2)(n+3) - Ab^2(n+2)(n+3) \tan(c+dx)) dx + \frac{(2a^2 B - aAb(n+3) - b^2)}{b(n+2)}}{b(n+3)}}{b(n+3)}$$

↓ 4022

$$\frac{B \tan^2(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 3)} - \frac{\frac{\tan(c+dx)(2aB - Ab(n+3))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{-\frac{1}{2}b^2(n+2)(n+3)(-B+iA) \int (1-i \tan(c+dx))(a+b \tan(c+dx))^n dx + \frac{1}{2}b^2(n+2)(n+3)(B+iA)}{b(n+3)}}{b(n+3)}$$

↓ 3042

$$\frac{B \tan^2(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 3)} - \frac{\frac{\tan(c+dx)(2aB - Ab(n+3))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{-\frac{1}{2}b^2(n+2)(n+3)(-B+iA) \int (1-i \tan(c+dx))(a+b \tan(c+dx))^n dx + \frac{1}{2}b^2(n+2)(n+3)(B+iA)}{b(n+3)}}{b(n+3)}$$

↓ 4020

$$\frac{B \tan^2(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 3)} - \frac{\frac{\tan(c+dx)(2aB - Ab(n+3))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{ib^2(n+2)(n+3)(B+iA) \int -\frac{(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d(i \tan(c+dx))}{2d} + \frac{ib^2(n+2)(n+3)(-B+iA) \int -\frac{(a+b \tan(c+dx))}{i \tan(c+dx)}}{2d}}{b(n+3)}}{b(n+3)}$$

↓ 25

$$\frac{B \tan^2(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 3)} - \frac{\frac{\tan(c+dx)(2aB - Ab(n+3))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{ib^2(n+2)(n+3)(B+iA) \int \frac{(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d(i \tan(c+dx))}{2d} - \frac{ib^2(n+2)(n+3)(-B+iA) \int \frac{(a+b \tan(c+dx))}{i \tan(c+dx)}}{2d}}{b(n+3)}}{b(n+3)}$$

↓ 78

$$\frac{B \tan^2(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 3)} - \frac{\frac{\tan(c+dx)(2aB - Ab(n+3))(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{(2a^2 B - aAb(n+3) - b^2 B(n+2)(n+3))(a+b \tan(c+dx))^{n+1}}{bd(n+1)} - \frac{ib^2(n+2)(n+3)(B+iA)(a+b \tan(c+dx))^n}{2a}}{b(n+3)}}{b(n+3)}$$

input `Int [Tan [c + d*x]^3*(a + b*Tan [c + d*x])^n*(A + B*Tan [c + d*x]), x]`

3.495. $\int \tan^3(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$

```
output (B*Tan[c + d*x]^2*(a + b*Tan[c + d*x])^(1 + n))/(b*d*(3 + n)) - (((2*a*B -
A*b*(3 + n))*Tan[c + d*x]*(a + b*Tan[c + d*x])^(1 + n))/(b*d*(2 + n)) - (
((2*a^2*B - a*A*b*(3 + n) - b^2*B*(2 + n)*(3 + n))*(a + b*Tan[c + d*x])^(1
+ n))/(b*d*(1 + n)) - ((I/2)*b^2*(I*A + B)*(2 + n)*(3 + n)*Hypergeometric
2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a - I*b)]*(a + b*Tan[c + d*x])^
(1 + n))/((a - I*b)*d*(1 + n)) - ((I/2)*b^2*(I*A - B)*(2 + n)*(3 + n)*Hype
rgeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a + I*b)]*(a + b*Tan[
c + d*x])^(1 + n))/((a + I*b)*d*(1 + n)))/(b*(2 + n))/(b*(3 + n))
```

3.495.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 78 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b
*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m
+ 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x]
&& !IntegerQ[m] && IntegerQ[n]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4020 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 +
c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[
b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]
```

```
rule 4022 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(
1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m
*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c
- a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]
```


rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.495.4 Maple [F]

$$\int \tan(dx + c)^3 (a + b \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^3*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^3*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.495.5 Fricas [F]

$$\int \tan^3(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^3 dx$$

input `integrate(tan(d*x+c)^3*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*tan(d*x + c)^4 + A*tan(d*x + c)^3)*(b*tan(d*x + c) + a)^n, x)`

3.495.6 Sympy [F]

$$\int \tan^3(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^n \tan^3(c + dx) dx$$

input `integrate(tan(d*x+c)**3*(a+b*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**n*tan(c + d*x)**3, x)`

3.495.7 Maxima [F]

$$\int \tan^3(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^3 dx$$

input `integrate(tan(d*x+c)^3*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*tan(d*x + c)^3, x)`

3.495.8 Giac [F]

$$\int \tan^3(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^3 dx$$

input `integrate(tan(d*x+c)^3*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*tan(d*x + c)^3, x)`

3.495.9 Mupad [F(-1)]

Timed out.

$$\int \tan^3(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^3 (A + B \tan(c + dx)) (a + b \tan(c + dx))^n dx$$

input `int(tan(c + d*x)^3*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n,x)`

output `int(tan(c + d*x)^3*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n, x)`

3.496 $\int \tan^2(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.496.1 Optimal result	4701
3.496.2 Mathematica [A] (verified)	4702
3.496.3 Rubi [A] (verified)	4702
3.496.4 Maple [F]	4705
3.496.5 Fricas [F]	4706
3.496.6 Sympy [F]	4706
3.496.7 Maxima [F]	4706
3.496.8 Giac [F]	4707
3.496.9 Mupad [F(-1)]	4707

3.496.1 Optimal result

Integrand size = 31, antiderivative size = 219

$$\begin{aligned} & \int \tan^2(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx \\ &= -\frac{(aB - Ab(2+n))(a+b \tan(c+dx))^{1+n}}{b^2d(1+n)(2+n)} \\ & \quad + \frac{(iA + B) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a-ib}\right) (a+b \tan(c+dx))^{1+n}}{2(a-ib)d(1+n)} \\ & \quad + \frac{(A + iB) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a+ib}\right) (a+b \tan(c+dx))^{1+n}}{2(ia-b)d(1+n)} \\ & \quad + \frac{B \tan(c+dx)(a+b \tan(c+dx))^{1+n}}{bd(2+n)} \end{aligned}$$

```
output -(B*a-A*b*(2+n))*(a+b*tan(d*x+c))^(1+n)/b^2/d/(1+n)/(2+n)+1/2*(I*A+B)*hypergeom([1, 1+n],[2+n],(a+b*tan(d*x+c))/(a-I*b))*(a+b*tan(d*x+c))^(1+n)/(a-I*b)/d/(1+n)+1/2*(A+I*B)*hypergeom([1, 1+n],[2+n],(a+b*tan(d*x+c))/(a+I*b))*(a+b*tan(d*x+c))^(1+n)/(I*a-b)/d/(1+n)+B*tan(d*x+c)*(a+b*tan(d*x+c))^(1+n)/b/d/(2+n)
```

3.496.2 Mathematica [A] (verified)

Time = 1.50 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.77

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \frac{(a + b \tan(c + dx))^{1+n} \left(\frac{4Ab - 2aB + 2Abn}{b + bn} + \frac{b(iA+B)(2+n) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a-ib}\right)}{(a-ib)(1+n)} \right) + \frac{b(-iA+B)(2+n) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a+ib}\right)}{(a+ib)(1+n)}}{2bd(2+n)}$$

input `Integrate[Tan[c + d*x]^2*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `((a + b*Tan[c + d*x])^(1 + n)*((4*A*b - 2*a*B + 2*A*b*n)/(b + b*n) + (b*(I *A + B)*(2 + n)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a - I*b)])/((a - I*b)*(1 + n)) + (b*(-I)*A + B)*(2 + n)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a + I*b)])/((a + I*b)*(1 + n)) + 2*B*Tan[c + d*x]))/(2*b*d*(2 + n))`

3.496.3 Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.07, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.355$, Rules used = {3042, 4090, 25, 3042, 4113, 3042, 4022, 3042, 4020, 25, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^2(c + dx)(A + B \tan(c + dx))(a + b \tan(c + dx))^n dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c + dx)^2(A + B \tan(c + dx))(a + b \tan(c + dx))^n dx$$

$$\downarrow \text{4090}$$

$$\frac{\int -(a + b \tan(c + dx))^n ((aB - Ab(n + 2)) \tan^2(c + dx) + bB(n + 2) \tan(c + dx) + aB) dx}{b(n + 2)} + \frac{B \tan(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 2)}$$

$$\downarrow \text{25}$$

3.496. $\int \tan^2(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$

$$\begin{aligned}
 & \frac{B \tan(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 2)} - \frac{\int (a + b \tan(c + dx))^n ((aB - Ab(n + 2)) \tan^2(c + dx) + bB(n + 2) \tan(c + dx) + aB) dx}{b(n + 2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{B \tan(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 2)} - \frac{\int (a + b \tan(c + dx))^n ((aB - Ab(n + 2)) \tan(c + dx)^2 + bB(n + 2) \tan(c + dx) + aB) dx}{b(n + 2)} \\
 & \quad \downarrow \text{4113} \\
 & \frac{B \tan(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 2)} - \frac{\int (a + b \tan(c + dx))^n (Ab(n + 2) + bB \tan(c + dx)(n + 2)) dx + \frac{(aB - Ab(n + 2))(a + b \tan(c + dx))^{n+1}}{bd(n + 1)}}{b(n + 2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{B \tan(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 2)} - \frac{\int (a + b \tan(c + dx))^n (Ab(n + 2) + bB \tan(c + dx)(n + 2)) dx + \frac{(aB - Ab(n + 2))(a + b \tan(c + dx))^{n+1}}{bd(n + 1)}}{b(n + 2)} \\
 & \quad \downarrow \text{4022} \\
 & \frac{B \tan(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 2)} - \frac{\frac{1}{2}b(n + 2)(A + iB) \int (1 - i \tan(c + dx))(a + b \tan(c + dx))^n dx + \frac{1}{2}b(n + 2)(A - iB) \int (i \tan(c + dx) + 1)(a + b \tan(c + dx))^n dx}{b(n + 2)}}{b(n + 2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{B \tan(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 2)} - \frac{\frac{1}{2}b(n + 2)(A + iB) \int (1 - i \tan(c + dx))(a + b \tan(c + dx))^n dx + \frac{1}{2}b(n + 2)(A - iB) \int (i \tan(c + dx) + 1)(a + b \tan(c + dx))^n dx}{b(n + 2)}}{b(n + 2)} \\
 & \quad \downarrow \text{4020} \\
 & \frac{B \tan(c + dx)(a + b \tan(c + dx))^{n+1}}{bd(n + 2)} - \frac{\frac{ib(n + 2)(A - iB) \int -\frac{(a + b \tan(c + dx))^n}{1 - i \tan(c + dx)} d(i \tan(c + dx))}{2d} - \frac{ib(n + 2)(A + iB) \int -\frac{(a + b \tan(c + dx))^n}{i \tan(c + dx) + 1} d(-i \tan(c + dx))}{2d} + \frac{(aB - Ab(n + 2))(a + b \tan(c + dx))^{n+1}}{bd(n + 1)}}{b(n + 2)}}{b(n + 2)} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

$$\frac{\frac{B \tan(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{ib(n+2)(A-iB) \int \frac{(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d(i \tan(c+dx))}{2d} + \frac{ib(n+2)(A+iB) \int \frac{(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d(-i \tan(c+dx))}{2d} + \frac{(aB-Ab(n+2))(a+b \tan(c+dx))}{bd(n+1)}}{b(n+2)}$$

↓ 78

$$\frac{\frac{B \tan(c+dx)(a+b \tan(c+dx))^{n+1}}{bd(n+2)} - \frac{ib(n+2)(A-iB)(a+b \tan(c+dx))^{n+1} \text{Hypergeometric2F1}\left(1, n+1, n+2, \frac{a+b \tan(c+dx)}{a-ib}\right)}{2d(n+1)(a-ib)} + \frac{ib(n+2)(A+iB)(a+b \tan(c+dx))^{n+1} \text{Hypergeometric2F1}\left(1, n+1, n+2, \frac{a+b \tan(c+dx)}{a+ib}\right)}{2d(n+1)(a+ib)}}{b(n+2)}$$

input `Int[Tan[c + d*x]^2*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

output `(B*Tan[c + d*x]*(a + b*Tan[c + d*x])^(1 + n))/(b*d*(2 + n)) - (((a*B - A*b*(2 + n))*(a + b*Tan[c + d*x])^(1 + n))/(b*d*(1 + n)) - ((I/2)*b*(A - I*B)*(2 + n)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a - I*b)]*(a + b*Tan[c + d*x])^(1 + n))/((a - I*b)*d*(1 + n)) + ((I/2)*b*(A + I*B)*(2 + n)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a + I*b)]*(a + b*Tan[c + d*x])^(1 + n))/((a + I*b)*d*(1 + n)))/(b*(2 + n))`

3.496.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 78 `Int[((a_) + (b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

3.496.4 Maple [F]

$$\int \tan(dx + c)^2 (a + b \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^2*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^2*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.496.5 Fricas [F]

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^2 dx$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*tan(d*x + c)^3 + A*tan(d*x + c)^2)*(b*tan(d*x + c) + a)^n, x)`

3.496.6 Sympy [F]

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^n \tan^2(c + dx) dx$$

input `integrate(tan(d*x+c)**2*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n*tan(c + d*x)**2, x)`

3.496.7 Maxima [F]

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^2 dx$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*tan(d*x + c)^2, x)`

3.496.8 Giac [F]

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^2 dx$$

input `integrate(tan(d*x+c)^2*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*tan(d*x + c)^2, x)`

3.496.9 Mupad [F(-1)]

Timed out.

$$\int \tan^2(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^2 (A + B \tan(c + dx)) (a + b \tan(c + dx))^n dx$$

input `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n,x)`

output `int(tan(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n, x)`

3.497 $\int \tan(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.497.1 Optimal result	4708
3.497.2 Mathematica [A] (verified)	4709
3.497.3 Rubi [A] (verified)	4709
3.497.4 Maple [F]	4712
3.497.5 Fricas [F]	4712
3.497.6 Sympy [F]	4712
3.497.7 Maxima [F]	4713
3.497.8 Giac [F]	4713
3.497.9 Mupad [F(-1)]	4713

3.497.1 Optimal result

Integrand size = 29, antiderivative size = 168

$$\int \tan(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \frac{B(a+b \tan(c+dx))^{1+n}}{bd(1+n)}$$

$$- \frac{(A-iB) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a-ib}\right) (a+b \tan(c+dx))^{1+n}}{2(a-ib)d(1+n)}$$

$$- \frac{(A+iB) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a+ib}\right) (a+b \tan(c+dx))^{1+n}}{2(a+ib)d(1+n)}$$

output

```
B*(a+b*tan(d*x+c))^(1+n)/b/d/(1+n)-1/2*(A-I*B)*hypergeom([1, 1+n], [2+n], (a+b*tan(d*x+c))/(a-I*b))*(a+b*tan(d*x+c))^(1+n)/(a-I*b)/d/(1+n)-1/2*(A+I*B)*hypergeom([1, 1+n], [2+n], (a+b*tan(d*x+c))/(a+I*b))*(a+b*tan(d*x+c))^(1+n)/(a+I*b)/d/(1+n)
```

3.497.2 Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.74

$$\int \tan(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \frac{\left(\frac{2B}{b} - \frac{(A - iB) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a-ib}\right)}{a-ib} - \frac{(A+iB) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a+ib}\right)}{a+ib} \right) (a + b \tan(c + dx))}{2d(1+n)}$$

input `Integrate[Tan[c + d*x]*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`output `((2*B)/b - ((A - I*B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a - I*b)])/(a - I*b) - ((A + I*B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a + I*b)])/(a + I*b))*(a + b*Tan[c + d*x])^(1 + n)/(2*d*(1 + n))`**3.497.3 Rubi [A] (verified)**Time = 0.53 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.04, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {3042, 4075, 3042, 4022, 3042, 4020, 25, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan(c + dx)(A + B \tan(c + dx))(a + b \tan(c + dx))^n dx$$

$$\downarrow 3042$$

$$\int \tan(c + dx)(A + B \tan(c + dx))(a + b \tan(c + dx))^n dx$$

$$\downarrow 4075$$

$$\int (A \tan(c + dx) - B)(a + b \tan(c + dx))^n dx + \frac{B(a + b \tan(c + dx))^{n+1}}{bd(n+1)}$$

$$\downarrow 3042$$

$$\int (A \tan(c + dx) - B)(a + b \tan(c + dx))^n dx + \frac{B(a + b \tan(c + dx))^{n+1}}{bd(n+1)}$$

$$\downarrow 4022$$

 3.497. $\int \tan(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \frac{1}{2}(-B + iA) \int (1 - i \tan(c + dx))(a + b \tan(c + dx))^n dx - \frac{1}{2}(B + iA) \int (i \tan(c + dx) + 1)(a + \\
& \quad b \tan(c + dx))^n dx + \frac{B(a + b \tan(c + dx))^{n+1}}{bd(n + 1)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{2}(-B + iA) \int (1 - i \tan(c + dx))(a + b \tan(c + dx))^n dx - \frac{1}{2}(B + iA) \int (i \tan(c + dx) + 1)(a + \\
& \quad b \tan(c + dx))^n dx + \frac{B(a + b \tan(c + dx))^{n+1}}{bd(n + 1)} \\
& \quad \downarrow \text{4020} \\
& \frac{i(B + iA) \int -\frac{(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d(i \tan(c + dx))}{2d} - \\
& \frac{i(-B + iA) \int -\frac{(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d(-i \tan(c + dx))}{2d} + \frac{B(a + b \tan(c + dx))^{n+1}}{bd(n + 1)} \\
& \quad \downarrow \text{25} \\
& \frac{i(B + iA) \int \frac{(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d(i \tan(c + dx))}{2d} + \frac{i(-B + iA) \int \frac{(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d(-i \tan(c + dx))}{2d} + \\
& \quad \frac{B(a + b \tan(c + dx))^{n+1}}{bd(n + 1)} \\
& \quad \downarrow \text{78} \\
& \frac{i(B + iA)(a + b \tan(c + dx))^{n+1} \text{Hypergeometric2F1}\left(1, n + 1, n + 2, \frac{a+b \tan(c+dx)}{a-ib}\right)}{2d(n + 1)(a - ib)} + \\
& \frac{i(-B + iA)(a + b \tan(c + dx))^{n+1} \text{Hypergeometric2F1}\left(1, n + 1, n + 2, \frac{a+b \tan(c+dx)}{a+ib}\right)}{2d(n + 1)(a + ib)} + \\
& \quad \frac{B(a + b \tan(c + dx))^{n+1}}{bd(n + 1)}
\end{aligned}$$

input `Int[Tan[c + d*x]*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `(B*(a + b*Tan[c + d*x])^(1 + n))/(b*d*(1 + n)) + ((I/2)*(I*A + B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a - I*b)]*(a + b*Tan[c + d*x])^(1 + n))/((a - I*b)*d*(1 + n)) + ((I/2)*(I*A - B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a + I*b)]*(a + b*Tan[c + d*x])^(1 + n))/((a + I*b)*d*(1 + n))`

3.497.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 78 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4020 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`
- rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.497.4 Maple [F]

$$\int \tan(dx + c) (a + b \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.497.5 Fricas [F]

$$\begin{aligned} & \int \tan(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c) dx \end{aligned}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*tan(d*x + c)^2 + A*tan(d*x + c))*(b*tan(d*x + c) + a)^n, x)`

3.497.6 Sympy [F]

$$\begin{aligned} & \int \tan(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^n \tan(c + dx) dx \end{aligned}$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**n*tan(c + d*x), x)`

3.497.7 Maxima [F]

$$\int \tan(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c) dx$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*tan(d*x + c), x)`

3.497.8 Giac [F]

$$\int \tan(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c) dx$$

input `integrate(tan(d*x+c)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*tan(d*x + c), x)`

3.497.9 Mupad [F(-1)]

Timed out.

$$\int \tan(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx) (A + B \tan(c + dx)) (a + b \tan(c + dx))^n dx$$

input `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n,x)`

output `int(tan(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n, x)`

3.498 $\int (a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$

3.498.1 Optimal result	4714
3.498.2 Mathematica [A] (verified)	4714
3.498.3 Rubi [A] (verified)	4715
3.498.4 Maple [F]	4717
3.498.5 Fracas [F]	4717
3.498.6 Sympy [F]	4717
3.498.7 Maxima [F]	4718
3.498.8 Giac [F]	4718
3.498.9 Mupad [F(-1)]	4718

3.498.1 Optimal result

Integrand size = 23, antiderivative size = 143

$$\int (a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \frac{(A - iB) \operatorname{Hypergeometric2F1}\left(1, 1 + n, 2 + n, \frac{a + b \tan(c + dx)}{a - ib}\right) (a + b \tan(c + dx))^{1+n}}{2(ia + b)d(1 + n)}$$

$$+ \frac{(iA - B) \operatorname{Hypergeometric2F1}\left(1, 1 + n, 2 + n, \frac{a + b \tan(c + dx)}{a + ib}\right) (a + b \tan(c + dx))^{1+n}}{2(a + ib)d(1 + n)}$$

output `1/2*(A-I*B)*hypergeom([1, 1+n], [2+n], (a+b*tan(d*x+c))/(a-I*b))*(a+b*tan(d*x+c))^(1+n)/(I*a+b)/d/(1+n)+1/2*(I*A-B)*hypergeom([1, 1+n], [2+n], (a+b*tan(d*x+c))/(a+I*b))*(a+b*tan(d*x+c))^(1+n)/(a+I*b)/d/(1+n)`

3.498.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.84

$$\int (a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \frac{i \left(-\frac{(A - iB) \operatorname{Hypergeometric2F1}\left(1, 1 + n, 2 + n, \frac{a + b \tan(c + dx)}{a - ib}\right)}{a - ib} + \frac{(A + iB) \operatorname{Hypergeometric2F1}\left(1, 1 + n, 2 + n, \frac{a + b \tan(c + dx)}{a + ib}\right)}{a + ib} \right) (a + b \tan(c + dx))^{1+n}}{2d(1 + n)}$$

input `Integrate[(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `((I/2)*(-((A - I*B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a - I*b)])/(a - I*b)) + ((A + I*B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a + I*b)])/(a + I*b))*(a + b*Tan[c + d*x])^(1 + n))/(d*(1 + n))`

3.498.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.01, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$, Rules used = {3042, 4022, 3042, 4020, 25, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (A + B \tan(c + dx))(a + b \tan(c + dx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (A + B \tan(c + dx))(a + b \tan(c + dx))^n dx \\
 & \quad \downarrow \text{4022} \\
 & \frac{1}{2}(A + iB) \int (1 - i \tan(c + dx))(a + b \tan(c + dx))^n dx + \frac{1}{2}(A - iB) \int (i \tan(c + dx) + 1)(a + b \tan(c + dx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2}(A + iB) \int (1 - i \tan(c + dx))(a + b \tan(c + dx))^n dx + \frac{1}{2}(A - iB) \int (i \tan(c + dx) + 1)(a + b \tan(c + dx))^n dx \\
 & \quad \downarrow \text{4020} \\
 & \frac{i(A - iB) \int -\frac{(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d(i \tan(c + dx))}{2d} - \frac{i(A + iB) \int -\frac{(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d(-i \tan(c + dx))}{2d} \\
 & \quad \downarrow \text{25} \\
 & \frac{i(A + iB) \int \frac{(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d(-i \tan(c + dx))}{2d} - \frac{i(A - iB) \int \frac{(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d(i \tan(c + dx))}{2d} \\
 & \quad \downarrow \text{78}
 \end{aligned}$$

3.498. $\int (a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$

$$\frac{i(A + iB)(a + b \tan(c + dx))^{n+1} \operatorname{Hypergeometric2F1}\left(1, n + 1, n + 2, \frac{a + b \tan(c + dx)}{a + ib}\right)}{2d(n + 1)(a + ib)} - \frac{i(A - iB)(a + b \tan(c + dx))^{n+1} \operatorname{Hypergeometric2F1}\left(1, n + 1, n + 2, \frac{a + b \tan(c + dx)}{a - ib}\right)}{2d(n + 1)(a - ib)}$$

input `Int[(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `((-1/2*I)*(A - I*B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a - I*b)]*(a + b*Tan[c + d*x])^(1 + n))/((a - I*b)*d*(1 + n)) + ((I/2)*(A + I*B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a + I*b)]*(a + b*Tan[c + d*x])^(1 + n))/((a + I*b)*d*(1 + n))`

3.498.3.1 Defintions of rubi rules used

rule 25 `Int[-(F*x_), x_Symbol] := Simp[Identity[-1] Int[F*x, x], x]`

rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

3.498.4 Maple [F]

$$\int (a + b \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.498.5 Fracas [F]

$$\int (a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n dx$$

input `integrate((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n, x)`

3.498.6 Sympy [F]

$$\begin{aligned} & \int (a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ & = \int (A + B \tan(c + dx)) (a + b \tan(c + dx))^n dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**n, x)`

3.498.7 Maxima [F]

$$\int (a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n dx$$

input `integrate((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n, x)`

3.498.8 Giac [F]

$$\int (a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n dx$$

input `integrate((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n, x)`

3.498.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int (a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (A + B \tan(c + dx)) (a + b \tan(c + dx))^n dx \end{aligned}$$

input `int((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n,x)`

output `int((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n, x)`

3.499 $\int \cot(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.499.1 Optimal result	4719
3.499.2 Mathematica [A] (verified)	4720
3.499.3 Rubi [A] (verified)	4720
3.499.4 Maple [F]	4723
3.499.5 Fricas [F]	4724
3.499.6 Sympy [F]	4724
3.499.7 Maxima [F]	4724
3.499.8 Giac [F]	4725
3.499.9 Mupad [F(-1)]	4725

3.499.1 Optimal result

Integrand size = 29, antiderivative size = 190

$$\int \cot(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \frac{(iA+B) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a-ib}\right) (a+b \tan(c+dx))^{1+n}}{2(ia+b)d(1+n)}$$

$$+ \frac{(A+iB) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a+ib}\right) (a+b \tan(c+dx))^{1+n}}{2(a+ib)d(1+n)}$$

$$- \frac{A \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, 1+\frac{b \tan(c+dx)}{a}\right) (a+b \tan(c+dx))^{1+n}}{ad(1+n)}$$

```
output 1/2*(I*A+B)*hypergeom([1, 1+n],[2+n],(a+b*tan(d*x+c))/(a-I*b))*(a+b*tan(d*x+c))^(1+n)/(I*a+b)/d/(1+n)+1/2*(A+I*B)*hypergeom([1, 1+n],[2+n],(a+b*tan(d*x+c))/(a+I*b))*(a+b*tan(d*x+c))^(1+n)/(a+I*b)/d/(1+n)-A*hypergeom([1, 1+n],[2+n],1+b*tan(d*x+c)/a)*(a+b*tan(d*x+c))^(1+n)/a/d/(1+n)
```

3.499.2 Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.89

$$\int \cot(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \frac{\left(a(a + ib)(A - iB) \operatorname{Hypergeometric2F1} \left(1, 1 + n, 2 + n, \frac{a + b \tan(c + dx)}{a - ib} \right) + (a - ib) \left(a(A + iB) \operatorname{Hypergeometric2F1} \left(1, 1 + n, 2 + n, \frac{a + b \tan(c + dx)}{a + ib} \right) \right)}{2}$$

input `Integrate[Cot[c + d*x]*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`output `((a*(a + I*b)*(A - I*B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a - I*b)] + (a - I*b)*(a*(A + I*B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a + I*b)] - 2*A*(a + I*b)*Hypergeometric2F1[1, 1 + n, 2 + n, 1 + (b*Tan[c + d*x])/a]))*(a + b*Tan[c + d*x])^(1 + n)/(2*a*(a - I*b)*(a + I*b)*d*(1 + n))`**3.499.3 Rubi [A] (verified)**Time = 0.78 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$, Rules used = {3042, 4096, 3042, 4022, 3042, 4020, 25, 78, 4117, 75}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot(c + dx)(A + B \tan(c + dx))(a + b \tan(c + dx))^n dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))^n}{\tan(c + dx)} dx$$

$$\downarrow \text{4096}$$

$$\int (B - A \tan(c + dx))(a + b \tan(c + dx))^n dx + A \int \cot(c + dx)(a + b \tan(c + dx))^n (\tan^2(c + dx) + 1) dx$$

$$\downarrow \text{3042}$$

$$\int (B - A \tan(c + dx))(a + b \tan(c + dx))^n dx + A \int \frac{(a + b \tan(c + dx))^n (\tan(c + dx)^2 + 1)}{\tan(c + dx)} dx$$

$$\begin{aligned}
& \downarrow 4022 \\
& -\frac{1}{2}(-B + iA) \int (1 - i \tan(c + dx))(a + b \tan(c + dx))^n dx + \frac{1}{2}(B + iA) \int (i \tan(c + dx) + 1)(a + \\
& \quad b \tan(c + dx))^n dx + A \int \frac{(a + b \tan(c + dx))^n (\tan(c + dx)^2 + 1)}{\tan(c + dx)} dx \\
& \downarrow 3042 \\
& -\frac{1}{2}(-B + iA) \int (1 - i \tan(c + dx))(a + b \tan(c + dx))^n dx + \frac{1}{2}(B + iA) \int (i \tan(c + dx) + 1)(a + \\
& \quad b \tan(c + dx))^n dx + A \int \frac{(a + b \tan(c + dx))^n (\tan(c + dx)^2 + 1)}{\tan(c + dx)} dx \\
& \downarrow 4020 \\
& \frac{i(B + iA) \int -\frac{(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d(i \tan(c + dx))}{2d} + \frac{i(-B + iA) \int -\frac{(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d(-i \tan(c + dx))}{2d} + \\
& \quad A \int \frac{(a + b \tan(c + dx))^n (\tan(c + dx)^2 + 1)}{\tan(c + dx)} dx \\
& \downarrow 25 \\
& -\frac{i(B + iA) \int \frac{(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d(i \tan(c + dx))}{2d} - \frac{i(-B + iA) \int \frac{(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d(-i \tan(c + dx))}{2d} + \\
& \quad A \int \frac{(a + b \tan(c + dx))^n (\tan(c + dx)^2 + 1)}{\tan(c + dx)} dx \\
& \downarrow 78 \\
& A \int \frac{(a + b \tan(c + dx))^n (\tan(c + dx)^2 + 1)}{\tan(c + dx)} dx - \\
& \frac{i(B + iA)(a + b \tan(c + dx))^{n+1} \operatorname{Hypergeometric2F1}\left(1, n + 1, n + 2, \frac{a+b \tan(c+dx)}{a-ib}\right)}{2d(n + 1)(a - ib)} - \\
& \frac{i(-B + iA)(a + b \tan(c + dx))^{n+1} \operatorname{Hypergeometric2F1}\left(1, n + 1, n + 2, \frac{a+b \tan(c+dx)}{a+ib}\right)}{2d(n + 1)(a + ib)} \\
& \downarrow 4117 \\
& \frac{A \int \cot(c + dx)(a + b \tan(c + dx))^n d \tan(c + dx)}{d} - \\
& \frac{i(B + iA)(a + b \tan(c + dx))^{n+1} \operatorname{Hypergeometric2F1}\left(1, n + 1, n + 2, \frac{a+b \tan(c+dx)}{a-ib}\right)}{2d(n + 1)(a - ib)} - \\
& \frac{i(-B + iA)(a + b \tan(c + dx))^{n+1} \operatorname{Hypergeometric2F1}\left(1, n + 1, n + 2, \frac{a+b \tan(c+dx)}{a+ib}\right)}{2d(n + 1)(a + ib)} \\
& \downarrow 75
\end{aligned}$$

$$\frac{i(B + iA)(a + b \tan(c + dx))^{n+1} \operatorname{Hypergeometric2F1}\left(1, n + 1, n + 2, \frac{a + b \tan(c + dx)}{a - ib}\right)}{2d(n + 1)(a - ib)} - \frac{i(-B + iA)(a + b \tan(c + dx))^{n+1} \operatorname{Hypergeometric2F1}\left(1, n + 1, n + 2, \frac{a + b \tan(c + dx)}{a + ib}\right)}{2d(n + 1)(a + ib)} - \frac{A(a + b \tan(c + dx))^{n+1} \operatorname{Hypergeometric2F1}\left(1, n + 1, n + 2, \frac{b \tan(c + dx)}{a} + 1\right)}{ad(n + 1)}$$

input `Int[Cot[c + d*x]*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `((-1/2*I)*(I*A + B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a - I*b)]*(a + b*Tan[c + d*x])^(1 + n))/((a - I*b)*d*(1 + n)) - ((I/2)*(I*A - B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a + I*b)]*(a + b*Tan[c + d*x])^(1 + n))/((a + I*b)*d*(1 + n)) - (A*Hypergeometric2F1[1, 1 + n, 2 + n, 1 + (b*Tan[c + d*x])/a]*(a + b*Tan[c + d*x])^(1 + n))/(a*d*(1 + n))`

3.499.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 75 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((c + d*x)^(n + 1)/(d*(n + 1)*(-d/(b*c))^m)*Hypergeometric2F1[-m, n + 1, n + 2, 1 + d*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[n] && (IntegerQ[m] || GtQ[-d/(b*c), 0])`

rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`

rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

rule 4096 `Int((((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[a*A + b*B - (A*b - a*B)*Tan[e + f*x], x], x] + Simp[b*((A*b - a*B)/(a^2 + b^2)) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

3.499.4 Maple [F]

$$\int \cot(dx + c)(a + b \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(cot(d*x+c)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(cot(d*x+c)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.499.5 Fricas [F]

$$\int \cot(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c) dx$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*cot(d*x + c)*tan(d*x + c) + A*cot(d*x + c))*(b*tan(d*x + c) + a)^n, x)`

3.499.6 Sympy [F]

$$\int \cot(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^n \cot(c + dx) dx$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**n*cot(c + d*x), x)`

3.499.7 Maxima [F]

$$\int \cot(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c) dx$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*cot(d*x + c), x)`

3.499.8 Giac [F]

$$\int \cot(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c) dx$$

input `integrate(cot(d*x+c)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*cot(d*x + c), x)`

3.499.9 Mupad [F(-1)]

Timed out.

$$\int \cot(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx) (A + B \tan(c + dx)) (a + b \tan(c + dx))^n dx$$

input `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n,x)`

output `int(cot(c + d*x)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n, x)`

3.500 $\int \cot^2(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.500.1 Optimal result	4726
3.500.2 Mathematica [A] (verified)	4727
3.500.3 Rubi [A] (verified)	4727
3.500.4 Maple [F]	4731
3.500.5 Fricas [F]	4732
3.500.6 Sympy [F]	4732
3.500.7 Maxima [F]	4732
3.500.8 Giac [F]	4733
3.500.9 Mupad [F(-1)]	4733

3.500.1 Optimal result

Integrand size = 31, antiderivative size = 228

$$\int \cot^2(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= -\frac{A \cot(c+dx)(a+b \tan(c+dx))^{1+n}}{ad}$$

$$- \frac{(A-iB) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a-ib}\right) (a+b \tan(c+dx))^{1+n}}{2(ia+b)d(1+n)}$$

$$+ \frac{(A+iB) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a+ib}\right) (a+b \tan(c+dx))^{1+n}}{2(ia-b)d(1+n)}$$

$$- \frac{(aB+Abn) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, 1+\frac{b \tan(c+dx)}{a}\right) (a+b \tan(c+dx))^{1+n}}{a^2d(1+n)}$$

```
output -A*cot(d*x+c)*(a+b*tan(d*x+c))^(1+n)/a/d-1/2*(A-I*B)*hypergeom([1, 1+n], [2+n], (a+b*tan(d*x+c))/(a-I*b))*(a+b*tan(d*x+c))^(1+n)/(I*a+b)/d/(1+n)+1/2*(A+I*B)*hypergeom([1, 1+n], [2+n], (a+b*tan(d*x+c))/(a+I*b))*(a+b*tan(d*x+c))^(1+n)/(I*a-b)/d/(1+n)-(A*b*n+B*a)*hypergeom([1, 1+n], [2+n], 1+b*tan(d*x+c)/a)*(a+b*tan(d*x+c))^(1+n)/a^2/d/(1+n)
```

3.500.2 Mathematica [A] (verified)

Time = 0.46 (sec) , antiderivative size = 202, normalized size of antiderivative = 0.89

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \frac{\left(a^2(a + ib)(A - iB) \operatorname{Hypergeometric2F1}\left(1, 1 + n, 2 + n, \frac{a + b \tan(c + dx)}{a - ib}\right) - (a - ib) \left(a^2(A + iB) \operatorname{Hypergeometric2F1}\left(1, 1 + n, 2 + n, \frac{a + b \tan(c + dx)}{a - ib}\right) \right) \right)}{2a^2(a - ib)((-I)a + b)d(1 + n)}$$

input `Integrate[Cot[c + d*x]^2*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`output `((a^2*(a + I*b)*(A - I*B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a - I*b)] - (a - I*b)*(a^2*(A + I*B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a + I*b)] + 2*((-I)*a + b)*(a*B*Hypergeometric2F1[1, 1 + n, 2 + n, 1 + (b*Tan[c + d*x])/a] - A*b*Hypergeometric2F1[2, 1 + n, 2 + n, 1 + (b*Tan[c + d*x])/a]))*(a + b*Tan[c + d*x])^(1 + n))/(2*a^2*(a - I*b)*((-I)*a + b)*d*(1 + n))`**3.500.3 Rubi [A] (verified)**Time = 1.07 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 4092, 25, 3042, 4136, 25, 3042, 4022, 3042, 4020, 25, 78, 4117, 75}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^2(c + dx)(A + B \tan(c + dx))(a + b \tan(c + dx))^n dx$$

$$\downarrow 3042$$

$$\int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))^n}{\tan(c + dx)^2} dx$$

$$\downarrow 4092$$

$$\int -\cot(c + dx)(a + b \tan(c + dx))^n (Abn \tan^2(c + dx) - aA \tan(c + dx) + aB + Abn) dx$$

$$\frac{A \cot(c + dx)(a + b \tan(c + dx))^{n+1}}{ad}$$

$$\downarrow 25$$

3.500. $\int \cot^2(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \frac{\int \cot(c+dx)(a+b \tan(c+dx))^n (Abn \tan^2(c+dx) - aA \tan(c+dx) + aB + Abn) dx}{\frac{A \cot(c+dx)(a+b \tan(c+dx))^{n+1}}{ad}} \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{(a+b \tan(c+dx))^n (Abn \tan(c+dx)^2 - aA \tan(c+dx) + aB + Abn)}{\tan(c+dx)} dx}{a} - \frac{A \cot(c+dx)(a+b \tan(c+dx))^{n+1}}{ad} \\
& \quad \downarrow \text{4136} \\
& \frac{\int -(a+b \tan(c+dx))^n (aA + aB \tan(c+dx)) dx + (aB + Abn) \int \cot(c+dx)(a+b \tan(c+dx))^n (\tan^2(c+dx) + 1) dx}{\frac{A \cot(c+dx)(a+b \tan(c+dx))^{n+1}}{ad}} \\
& \quad \downarrow \text{25} \\
& \frac{(aB + Abn) \int \cot(c+dx)(a+b \tan(c+dx))^n (\tan^2(c+dx) + 1) dx - \int (a+b \tan(c+dx))^n (aA + aB \tan(c+dx)) dx}{\frac{A \cot(c+dx)(a+b \tan(c+dx))^{n+1}}{ad}} \\
& \quad \downarrow \text{3042} \\
& \frac{(aB + Abn) \int \frac{(a+b \tan(c+dx))^n (\tan(c+dx)^2 + 1)}{\tan(c+dx)} dx - \int (a+b \tan(c+dx))^n (aA + aB \tan(c+dx)) dx}{\frac{A \cot(c+dx)(a+b \tan(c+dx))^{n+1}}{ad}} \\
& \quad \downarrow \text{4022} \\
& \frac{(aB + Abn) \int \frac{(a+b \tan(c+dx))^n (\tan(c+dx)^2 + 1)}{\tan(c+dx)} dx - \frac{1}{2} a(A + iB) \int (1 - i \tan(c+dx))(a+b \tan(c+dx))^n dx - \frac{1}{2} a(A - iB) \int (1 + i \tan(c+dx))(a+b \tan(c+dx))^n dx}{\frac{A \cot(c+dx)(a+b \tan(c+dx))^{n+1}}{ad} + a} \\
& \quad \downarrow \text{3042} \\
& \frac{(aB + Abn) \int \frac{(a+b \tan(c+dx))^n (\tan(c+dx)^2 + 1)}{\tan(c+dx)} dx - \frac{1}{2} a(A + iB) \int (1 - i \tan(c+dx))(a+b \tan(c+dx))^n dx - \frac{1}{2} a(A - iB) \int (1 + i \tan(c+dx))(a+b \tan(c+dx))^n dx}{\frac{A \cot(c+dx)(a+b \tan(c+dx))^{n+1}}{ad} + a} \\
& \quad \downarrow \text{4020}
\end{aligned}$$

3.500. $\int \cot^2(c+dx)(a+b \tan(c+dx))^n (A+B \tan(c+dx)) dx$

3.500.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 75 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((c + d*x)^(n + 1)/(d*(n + 1)*(-d/(b*c))^m)*Hypergeometric2F1[-m, n + 1, n + 2, 1 + d*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[n] && (IntegerQ[m] || GtQ[-d/(b*c), 0])`
- rule 78 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*(a + b*x)^(m + 1)/(b^(n + 1)*(m + 1))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4020 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`

```
rule 4092 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)
/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^
2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*
B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2
)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n
+ 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1]
&& (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m]
|| (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

```
rule 4136 Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.500.4 Maple [F]

$$\int \cot(dx + c)^2 (a + b \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

```
input int(cot(d*x+c)^2*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)
```

```
output int(cot(d*x+c)^2*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)
```

3.500.5 Fricas [F]

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c)^2 dx$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*cot(d*x + c)^2*tan(d*x + c) + A*cot(d*x + c)^2)*(b*tan(d*x + c) + a)^n, x)`

3.500.6 Sympy [F]

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^n \cot^2(c + dx) dx$$

input `integrate(cot(d*x+c)**2*(a+b*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**n*cot(c + d*x)**2, x)`

3.500.7 Maxima [F]

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c)^2 dx$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*cot(d*x + c)^2, x)`

3.500.8 Giac [F]

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c)^2 dx$$

input `integrate(cot(d*x+c)^2*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*cot(d*x + c)^2, x)`

3.500.9 Mupad [F(-1)]

Timed out.

$$\int \cot^2(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^2 (A + B \tan(c + dx)) (a + b \tan(c + dx))^n dx$$

input `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n,x)`

output `int(cot(c + d*x)^2*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n, x)`

3.501 $\int \cot^3(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.501.1 Optimal result	4734
3.501.2 Mathematica [A] (verified)	4735
3.501.3 Rubi [A] (verified)	4735
3.501.4 Maple [F]	4740
3.501.5 Fricas [F]	4740
3.501.6 Sympy [F]	4741
3.501.7 Maxima [F]	4741
3.501.8 Giac [F]	4742
3.501.9 Mupad [F(-1)]	4742

3.501.1 Optimal result

Integrand size = 31, antiderivative size = 292

$$\int \cot^3(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= -\frac{(2aB - Ab(1 - n)) \cot(c+dx)(a+b \tan(c+dx))^{1+n}}{2a^2d}$$

$$- \frac{A \cot^2(c+dx)(a+b \tan(c+dx))^{1+n}}{2ad}$$

$$- \frac{(iA + B) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a-ib}\right) (a+b \tan(c+dx))^{1+n}}{2(ia+b)d(1+n)}$$

$$- \frac{(A + iB) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, \frac{a+b \tan(c+dx)}{a+ib}\right) (a+b \tan(c+dx))^{1+n}}{2(a+ib)d(1+n)}$$

$$+ \frac{(2a^2A - 2abBn + Ab^2(1 - n)n) \operatorname{Hypergeometric2F1}\left(1, 1+n, 2+n, 1 + \frac{b \tan(c+dx)}{a}\right) (a+b \tan(c+dx))^{1+n}}{2a^3d(1+n)}$$

output

```
-1/2*(2*B*a-A*b*(1-n))*cot(d*x+c)*(a+b*tan(d*x+c))^(1+n)/a^2/d-1/2*A*cot(d
*x+c)^2*(a+b*tan(d*x+c))^(1+n)/a/d-1/2*(I*A+B)*hypergeom([1, 1+n], [2+n], (a
+b*tan(d*x+c))/(a-I*b))*(a+b*tan(d*x+c))^(1+n)/(I*a+b)/d/(1+n)-1/2*(A+I*B)
*hypergeom([1, 1+n], [2+n], (a+b*tan(d*x+c))/(a+I*b))*(a+b*tan(d*x+c))^(1+n)
/(a+I*b)/d/(1+n)+1/2*(2*A*a^2-2*a*b*B*n+A*b^2*(1-n)*n)*hypergeom([1, 1+n],
[2+n], 1+b*tan(d*x+c)/a)*(a+b*tan(d*x+c))^(1+n)/a^3/d/(1+n)
```

3.501.2 Mathematica [A] (verified)

Time = 0.59 (sec) , antiderivative size = 230, normalized size of antiderivative = 0.79

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx =$$

$$\left(a^3(a + ib)(A - iB) \operatorname{Hypergeometric2F1} \left(1, 1 + n, 2 + n, \frac{a + b \tan(c + dx)}{a - ib} \right) + (a - ib) \left(a^3(A + iB) \operatorname{Hypergeometric2F1} \left(1, 1 + n, 2 + n, \frac{a + b \tan(c + dx)}{a - ib} \right) \right) \right) dx$$

input `Integrate[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `-1/2*((a^3*(a + I*b)*(A - I*B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a - I*b)] + (a - I*b)*(a^3*(A + I*B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a + I*b)] - 2*(a + I*b)*(a^2*A*Hypergeometric2F1[1, 1 + n, 2 + n, 1 + (b*Tan[c + d*x])/a] + b*(a*B*Hypergeometric2F1[2, 1 + n, 2 + n, 1 + (b*Tan[c + d*x])/a] - A*b*Hypergeometric2F1[3, 1 + n, 2 + n, 1 + (b*Tan[c + d*x])/a]))*(a + b*Tan[c + d*x])^(1 + n))/(a^3*(a - I*b)*(a + I*b)*d*(1 + n))`

3.501.3 Rubi [A] (verified)

Time = 1.65 (sec) , antiderivative size = 317, normalized size of antiderivative = 1.09, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.516$, Rules used = {3042, 4092, 25, 3042, 4132, 3042, 4136, 27, 3042, 4022, 3042, 4020, 25, 78, 4117, 75}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^3(c + dx)(A + B \tan(c + dx))(a + b \tan(c + dx))^n dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))^n}{\tan(c + dx)^3} dx$$

$$\downarrow \text{4092}$$

$$\frac{\int -\cot^2(c+dx)(a+b\tan(c+dx))^n(-Ab(1-n)\tan^2(c+dx)-2aA\tan(c+dx)+2aB-A(b-bn))dx}{\frac{A\cot^2(c+dx)(a+b\tan(c+dx))^{n+1}}{2ad}} \quad \downarrow \quad 25$$

$$\frac{\int \cot^2(c+dx)(a+b\tan(c+dx))^n(-Ab(1-n)\tan^2(c+dx)-2aA\tan(c+dx)+2aB-Ab(1-n))dx}{\frac{A\cot^2(c+dx)(a+b\tan(c+dx))^{n+1}}{2ad}} \quad \downarrow \quad 3042$$

$$\frac{\int \frac{(a+b\tan(c+dx))^n(-Ab(1-n)\tan^2(c+dx)-2aA\tan(c+dx)+2aB-Ab(1-n))}{\tan^2(c+dx)}dx}{\frac{A\cot^2(c+dx)(a+b\tan(c+dx))^{n+1}}{2ad}} \quad \downarrow \quad 4132$$

$$\frac{\int \cot(c+dx)(a+b\tan(c+dx))^n(2Aa^2+2B\tan(c+dx)a^2-2bBna-b(2aB-Ab(1-n))n\tan^2(c+dx)+Ab^2(1-n)n)dx}{a} - \frac{\cot(c+dx)(2aB-Ab(1-n))}{a} \quad \downarrow \quad 3042$$

$$\frac{\int \frac{(a+b\tan(c+dx))^n(2Aa^2+2B\tan(c+dx)a^2-2bBna-b(2aB-Ab(1-n))n\tan^2(c+dx)+Ab^2(1-n)n)}{\tan(c+dx)}dx}{a} - \frac{\cot(c+dx)(2aB-Ab(1-n))(a+b\tan(c+dx))^n}{ad} \quad \downarrow \quad 4136$$

$$\frac{(2a^2A-2abBn+Ab^2(1-n))\int \cot(c+dx)(a+b\tan(c+dx))^n(\tan^2(c+dx)+1)dx + \int 2(a^2B-a^2A\tan(c+dx))(a+b\tan(c+dx))^n dx}{a} - \frac{\cot(c+dx)}{a} \quad \downarrow \quad 27$$

$$\frac{(2a^2A-2abBn+Ab^2(1-n))\int \cot(c+dx)(a+b\tan(c+dx))^n(\tan^2(c+dx)+1)dx + 2\int (a^2B-a^2A\tan(c+dx))(a+b\tan(c+dx))^n dx}{a} - \frac{\cot(c+dx)}{a} \quad \downarrow \quad 2a$$

$$\frac{A\cot^2(c+dx)(a+b\tan(c+dx))^{n+1}}{2ad} \quad \downarrow \quad 2a$$

3.501. $\int \cot^3(c+dx)(a+b\tan(c+dx))^n(A+B\tan(c+dx))dx$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \frac{(2a^2A - 2abBn + Ab^2(1-n)n) \int \frac{(a+b \tan(c+dx))^n (\tan(c+dx)^2+1)}{\tan(c+dx)} dx + 2 \int (a^2B - a^2A \tan(c+dx)) (a+b \tan(c+dx))^n dx}{a} - \frac{\cot(c+dx)(2aB - Ab(1-n))}{a}}{\frac{A \cot^2(c+dx)(a+b \tan(c+dx))^{n+1}}{2ad}} \\
 & \downarrow 4022 \\
 & \frac{A \cot^2(c+dx)(a+b \tan(c+dx))^{n+1}}{2ad} + \frac{\cot(c+dx)(2aB - Ab(1-n))(a+b \tan(c+dx))^{n+1}}{ad} - \frac{(2a^2A - 2abBn + Ab^2(1-n)n) \int \frac{(a+b \tan(c+dx))^n (\tan(c+dx)^2+1)}{\tan(c+dx)} dx + 2 \left(\frac{1}{2}a^2(B+iA) \int (i \tan(c+dx))^{n+1} dx\right)}{2a} \\
 & \downarrow 3042 \\
 & \frac{A \cot^2(c+dx)(a+b \tan(c+dx))^{n+1}}{2ad} + \frac{\cot(c+dx)(2aB - Ab(1-n))(a+b \tan(c+dx))^{n+1}}{ad} - \frac{(2a^2A - 2abBn + Ab^2(1-n)n) \int \frac{(a+b \tan(c+dx))^n (\tan(c+dx)^2+1)}{\tan(c+dx)} dx + 2 \left(\frac{1}{2}a^2(B+iA) \int (i \tan(c+dx))^{n+1} dx\right)}{2a} \\
 & \downarrow 4020 \\
 & \frac{A \cot^2(c+dx)(a+b \tan(c+dx))^{n+1}}{2ad} + \frac{\cot(c+dx)(2aB - Ab(1-n))(a+b \tan(c+dx))^{n+1}}{ad} - \frac{(2a^2A - 2abBn + Ab^2(1-n)n) \int \frac{(a+b \tan(c+dx))^n (\tan(c+dx)^2+1)}{\tan(c+dx)} dx + 2 \left(\frac{ia^2(B+iA) \int \frac{(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} dx\right)}{2a} \\
 & \downarrow 25 \\
 & \frac{A \cot^2(c+dx)(a+b \tan(c+dx))^{n+1}}{2ad} + \frac{\cot(c+dx)(2aB - Ab(1-n))(a+b \tan(c+dx))^{n+1}}{ad} - \frac{(2a^2A - 2abBn + Ab^2(1-n)n) \int \frac{(a+b \tan(c+dx))^n (\tan(c+dx)^2+1)}{\tan(c+dx)} dx + 2 \left(-\frac{ia^2(B+iA) \int \frac{(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} dx\right)}{2a} \\
 & \downarrow 78 \\
 & \frac{A \cot^2(c+dx)(a+b \tan(c+dx))^{n+1}}{2ad} + \frac{\cot(c+dx)(2aB - Ab(1-n))(a+b \tan(c+dx))^{n+1}}{ad} - \frac{(2a^2A - 2abBn + Ab^2(1-n)n) \int \frac{(a+b \tan(c+dx))^n (\tan(c+dx)^2+1)}{\tan(c+dx)} dx + 2 \left(\frac{ia^2(B+iA)(a+b \tan(c+dx))^{n+1}}{a}\right)}{2a} \\
 & \downarrow 4117
 \end{aligned}$$

3.501. $\int \cot^3(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

$$\begin{aligned}
 & \frac{-\frac{A \cot^2(c+dx)(a+b \tan(c+dx))^{n+1}}{2ad} + \frac{\cot(c+dx)(2aB-Ab(1-n))(a+b \tan(c+dx))^{n+1}}{ad} - \frac{(2a^2A-2abBn+Ab^2(1-n)) \int \cot(c+dx)(a+b \tan(c+dx))^n d \tan(c+dx)}{d}}{2a} + 2 \left(-\frac{ia^2(B+iA)(a+b \tan(c+dx))^{n+1}}{ad} \right) \\
 & \quad \downarrow 75 \\
 & \frac{-\frac{A \cot^2(c+dx)(a+b \tan(c+dx))^{n+1}}{2ad} + \frac{\cot(c+dx)(2aB-Ab(1-n))(a+b \tan(c+dx))^{n+1}}{ad} - \frac{(2a^2A-2abBn+Ab^2(1-n))(a+b \tan(c+dx))^{n+1} \text{Hypergeometric2F1}\left(1, n+1, n+2, \frac{b \tan(c+dx)}{a}\right) + 1}{ad(n+1)}}{2a}
 \end{aligned}$$

input `Int[Cot[c + d*x]^3*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `-1/2*(A*Cot[c + d*x]^2*(a + b*Tan[c + d*x])^(1 + n))/(a*d) + (-(((2*a*B - A*b*(1 - n))*Cot[c + d*x]*(a + b*Tan[c + d*x])^(1 + n))/(a*d)) - (((2*a^2*A - 2*a*b*B*n + A*b^2*(1 - n)*n)*Hypergeometric2F1[1, 1 + n, 2 + n, 1 + (b*Tan[c + d*x])/a]*(a + b*Tan[c + d*x])^(1 + n))/(a*d*(1 + n)))) + 2*(((-1/2*I)*a^2*(I*A + B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a - I*b)]*(a + b*Tan[c + d*x])^(1 + n))/((a - I*b)*d*(1 + n)) - ((I/2)*a^2*(I*A - B)*Hypergeometric2F1[1, 1 + n, 2 + n, (a + b*Tan[c + d*x])/(a + I*b)]*(a + b*Tan[c + d*x])^(1 + n))/((a + I*b)*d*(1 + n))))/a)/(2*a)`

3.501.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 75 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((c + d*x)^(n + 1)/(d*(n + 1)*(-d/(b*c))^m)*Hypergeometric2F1[-m, n + 1, n + 2, 1 + d*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[n] && (IntegerQ[m] || GtQ[-d/(b*c), 0])`

- rule 78 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b*c - a*d)^n*(a + b*x)^(m + 1)/(b^(n + 1)*(m + 1))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4020 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[c*(d/f) Subst[Int[(a + (b/d)*x)^m/(d^2 + c*x), x], x, d*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[c^2 + d^2, 0]`
- rule 4022 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c + I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 - I*Tan[e + f*x]), x], x] + Simp[(c - I*d)/2 Int[(a + b*Tan[e + f*x])^m*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !IntegerQ[m]`
- rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`
- rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.501.4 Maple [F]

$$\int \cot(dx + c)^3 (a + b \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

```
input int(cot(d*x+c)^3*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)
```

```
output int(cot(d*x+c)^3*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)
```

3.501.5 Fricas [F]

$$\begin{aligned} & \int \cot^3(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c)^3 dx \end{aligned}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*cot(d*x + c)^3*tan(d*x + c) + A*cot(d*x + c)^3)*(b*tan(d*x + c) + a)^n, x)`

3.501.6 Sympy [F]

$$\begin{aligned} & \int \cot^3(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^n \cot^3(c + dx) dx \end{aligned}$$

input `integrate(cot(d*x+c)**3*(a+b*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**n*cot(c + d*x)**3, x)`

3.501.7 Maxima [F]

$$\begin{aligned} & \int \cot^3(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c)^3 dx \end{aligned}$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*cot(d*x + c)^3, x)`

3.501.8 Giac [F]

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c)^3 dx$$

input `integrate(cot(d*x+c)^3*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*cot(d*x + c)^3, x)`

3.501.9 Mupad [F(-1)]

Timed out.

$$\int \cot^3(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^3 (A + B \tan(c + dx)) (a + b \tan(c + dx))^n dx$$

input `int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n,x)`

output `int(cot(c + d*x)^3*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n, x)`

3.502 $\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.502.1 Optimal result	4743
3.502.2 Mathematica [C] (verified)	4743
3.502.3 Rubi [A] (verified)	4744
3.502.4 Maple [B] (verified)	4747
3.502.5 Fricas [B] (verification not implemented)	4748
3.502.6 Sympy [F(-1)]	4748
3.502.7 Maxima [B] (verification not implemented)	4749
3.502.8 Giac [F]	4749
3.502.9 Mupad [F(-1)]	4750

3.502.1 Optimal result

Integrand size = 34, antiderivative size = 103

$$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{2\sqrt[4]{-1}a(A-iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d}$$

$$+ \frac{2a(A-iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a(iA+B)\cot^{\frac{3}{2}}(c+dx)}{3d} - \frac{2aA\cot^{\frac{5}{2}}(c+dx)}{5d}$$

```
output 2*(-1)^(1/4)*a*(A-I*B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d-2/3*a*(I*A+B)
)*cot(d*x+c)^(3/2)/d-2/5*a*A*cot(d*x+c)^(5/2)/d+2*a*(A-I*B)*cot(d*x+c)^(1/
2)/d
```

3.502.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.76 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.53

$$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx =$$

$$-\frac{2a\cot^{\frac{3}{2}}(c+dx)(3A\cot(c+dx)+5(iA+B)\operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, i \tan(c+dx)\right))}{15d}$$

input `Integrate[Cot[c + d*x]^(7/2)*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(-2*a*Cot[c + d*x]^(3/2)*(3*A*Cot[c + d*x] + 5*(I*A + B)*Hypergeometric2F1[-3/2, 1, -1/2, I*Tan[c + d*x]]))/(15*d)`

3.502.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.324$, Rules used = {3042, 4064, 3042, 4075, 3042, 4011, 3042, 4011, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \cot(c + dx)^{7/2}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4064} \\
 & \int \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + ia)(A \cot(c + dx) + B) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \left(-\tan\left(c + dx + \frac{\pi}{2}\right)\right)^{3/2} \left(-a \tan\left(c + dx + \frac{\pi}{2}\right) + ia\right) \left(B - A \tan\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{4075} \\
 & -\frac{2aA \cot^{\frac{5}{2}}(c + dx)}{5d} + \int \cot^{\frac{3}{2}}(c + dx)(a(iA + B) \cot(c + dx) - a(A - iB)) dx \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2aA \cot^{\frac{5}{2}}(c + dx)}{5d} + \\
 & \int \left(-\tan\left(c + dx + \frac{\pi}{2}\right)\right)^{3/2} \left(-a(A - iB) - a(iA + B) \tan\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{4011}
 \end{aligned}$$

3.502. $\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \int \sqrt{\cot(c+dx)}(-a(iA+B) - a(A-iB)\cot(c+dx))dx - \frac{2a(B+iA)\cot^{\frac{3}{2}}(c+dx)}{3d} - \\
& \quad \frac{2aA\cot^{\frac{5}{2}}(c+dx)}{5d} \\
& \quad \downarrow \text{3042} \\
& \int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(a(A-iB)\tan\left(c+dx+\frac{\pi}{2}\right) - a(iA+B)\right)dx - \\
& \quad \frac{2a(B+iA)\cot^{\frac{3}{2}}(c+dx)}{3d} - \frac{2aA\cot^{\frac{5}{2}}(c+dx)}{5d} \\
& \quad \downarrow \text{4011} \\
& \int \frac{a(A-iB) - a(iA+B)\cot(c+dx)}{\sqrt{\cot(c+dx)}}dx - \frac{2a(B+iA)\cot^{\frac{3}{2}}(c+dx)}{3d} + \\
& \quad \frac{2a(A-iB)\sqrt{\cot(c+dx)}}{d} - \frac{2aA\cot^{\frac{5}{2}}(c+dx)}{5d} \\
& \quad \downarrow \text{3042} \\
& \int \frac{a(A-iB) + a(iA+B)\tan\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}}dx - \frac{2a(B+iA)\cot^{\frac{3}{2}}(c+dx)}{3d} + \\
& \quad \frac{2a(A-iB)\sqrt{\cot(c+dx)}}{d} - \frac{2aA\cot^{\frac{5}{2}}(c+dx)}{5d} \\
& \quad \downarrow \text{4016} \\
& \frac{2a^2(A-iB)^2 \int \frac{1}{-a(A-iB)-a(iA+B)\cot(c+dx)}d\sqrt{\cot(c+dx)}}{d} - \frac{2a(B+iA)\cot^{\frac{3}{2}}(c+dx)}{3d} + \\
& \quad \frac{2a(A-iB)\sqrt{\cot(c+dx)}}{d} - \frac{2aA\cot^{\frac{5}{2}}(c+dx)}{5d} \\
& \quad \downarrow \text{221} \\
& \frac{2\sqrt[4]{-1}a(A-iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} - \frac{2a(B+iA)\cot^{\frac{3}{2}}(c+dx)}{3d} + \\
& \quad \frac{2a(A-iB)\sqrt{\cot(c+dx)}}{d} - \frac{2aA\cot^{\frac{5}{2}}(c+dx)}{5d}
\end{aligned}$$

input `Int[Cot[c + d*x]^(7/2)*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(2*(-1)^(1/4)*a*(A - I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]])/d + (2*a*(A - I*B)*Sqrt[Cot[c + d*x]])/d - (2*a*(I*A + B)*Cot[c + d*x]^(3/2))/(3*d) - (2*a*A*Cot[c + d*x]^(5/2))/(5*d)`

3.502. $\int \cot^{\frac{7}{2}}(c+dx)(a+ia\tan(c+dx))(A+B\tan(c+dx))dx$

3.502.3.1 Defintions of rubi rules used

- rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`
- rule 4016 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`
- rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)])*(g_.)^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.502.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 550 vs. $2(84) = 168$.

Time = 0.61 (sec) , antiderivative size = 551, normalized size of antiderivative = 5.35

method	result
derivativedivides	$-\frac{a\left(\frac{1}{\tan(dx+c)}\right)^{\frac{7}{2}}\tan(dx+c)\left(30iA\sqrt{2}\arctan\left(-1+\sqrt{2}\sqrt{\tan(dx+c)}\right)\tan(dx+c)^{\frac{5}{2}}+30iB\sqrt{2}\arctan\left(-1+\sqrt{2}\sqrt{\tan(dx+c)}\right)\tan(dx+c)^{\frac{5}{2}}\right)}{\dots}$
default	$-\frac{a\left(\frac{1}{\tan(dx+c)}\right)^{\frac{7}{2}}\tan(dx+c)\left(30iA\sqrt{2}\arctan\left(-1+\sqrt{2}\sqrt{\tan(dx+c)}\right)\tan(dx+c)^{\frac{5}{2}}+30iB\sqrt{2}\arctan\left(-1+\sqrt{2}\sqrt{\tan(dx+c)}\right)\tan(dx+c)^{\frac{5}{2}}\right)}{\dots}$

```
input int(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output -1/60*a/d*(1/tan(d*x+c))^(7/2)*tan(d*x+c)*(30*I*A*2^(1/2)*arctan(-1+2^(1/2)
)*tan(d*x+c)^(1/2))*tan(d*x+c)^(5/2)+30*I*B*2^(1/2)*arctan(-1+2^(1/2)*tan(
d*x+c)^(1/2))*tan(d*x+c)^(5/2)+30*I*B*2^(1/2)*arctan(1+2^(1/2)*tan(d*x+c)^(
1/2))*tan(d*x+c)^(5/2)+15*I*B*2^(1/2)*ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d
*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))*tan(d*x+c)^(5/2)+15*I*A*
2^(1/2)*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*tan(d*x+c)^(1
/2)-tan(d*x+c)-1))*tan(d*x+c)^(5/2)+40*I*A*tan(d*x+c)-30*A*2^(1/2)*arctan(
1+2^(1/2)*tan(d*x+c)^(1/2))*tan(d*x+c)^(5/2)-30*A*2^(1/2)*arctan(-1+2^(1/2)
)*tan(d*x+c)^(1/2))*tan(d*x+c)^(5/2)-15*A*2^(1/2)*ln(-(2^(1/2)*tan(d*x+c)^(
1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))*tan(d*x+c)^(5
/2)+15*B*2^(1/2)*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*tan(
d*x+c)^(1/2)-tan(d*x+c)-1))*tan(d*x+c)^(5/2)+30*B*2^(1/2)*arctan(1+2^(1/2)
)*tan(d*x+c)^(1/2))*tan(d*x+c)^(5/2)+30*B*2^(1/2)*arctan(-1+2^(1/2)*tan(d*x
+c)^(1/2))*tan(d*x+c)^(5/2)+30*I*A*2^(1/2)*arctan(1+2^(1/2)*tan(d*x+c)^(1/
2))*tan(d*x+c)^(5/2)-120*A*tan(d*x+c)^2+120*I*B*tan(d*x+c)^2+40*B*tan(d*x+
c)+24*A)
```

3.502.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 434 vs. $2(81) = 162$.

Time = 0.27 (sec) , antiderivative size = 434, normalized size of antiderivative = 4.21

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx =$$

$$15 \left(de^{(4i dx + 4i c)} - 2 de^{(2i dx + 2i c)} + d \right) \sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^2}{d^2}} \log \left(-\frac{2 \left((A - iB)ae^{(2i dx + 2i c)} - (i de^{(2i dx + 2i c)} - i d) \sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^2}{d^2}} \right)}{(i \dots)} \right)$$

input `integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-1/30*(15*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*log(-2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) - (I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*sqrt(((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a) - 15*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*log(-2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) - (-I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*sqrt(((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a) - 4*((23*A - 20*I*B)*a*e^(4*I*d*x + 4*I*c) - 6*(4*A - 5*I*B)*a*e^(2*I*d*x + 2*I*c) + (13*A - 10*I*B)*a)*sqrt(((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))/(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)`

3.502.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(7/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.502. $\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

3.502.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 190 vs. $2(81) = 162$.

Time = 0.37 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.84

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{15 \left(2 \sqrt{2}((i - 1) A + (i + 1) B) \arctan \left(\frac{1}{2} \sqrt{2} \left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}} \right) \right) + 2 \sqrt{2}((i - 1) A + (i + 1) B) \arctan \right.}{}$$

input `integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/60*(15*(2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a + 120*(A - I*B)*a/sqrt(tan(d*x + c)) + 40*(-I*A - B)*a/tan(d*x + c)^(3/2) - 24*A*a/tan(d*x + c)^(5/2))/d`

3.502.8 Giac [F]

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a) \cot(dx + c)^{\frac{7}{2}} dx$$

input `integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)*cot(d*x + c)^(7/2), x)`

3.502.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{7/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) \text{ li}) dx$$

input `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`output `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i), x)`

3.503 $\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.503.1 Optimal result	4751
3.503.2 Mathematica [C] (verified)	4751
3.503.3 Rubi [A] (verified)	4752
3.503.4 Maple [B] (verified)	4754
3.503.5 Fricas [B] (verification not implemented)	4755
3.503.6 Sympy [F(-1)]	4756
3.503.7 Maxima [B] (verification not implemented)	4756
3.503.8 Giac [F]	4757
3.503.9 Mupad [F(-1)]	4757

3.503.1 Optimal result

Integrand size = 34, antiderivative size = 78

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= -\frac{2\sqrt[4]{-1}a(iA+B)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} - \frac{2a(iA+B)\sqrt{\cot(c+dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)}{3d}$$

```
output -2*(-1)^(1/4)*a*(I*A+B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d-2/3*a*A*cot
(d*x+c)^(3/2)/d-2*a*(I*A+B)*cot(d*x+c)^(1/2)/d
```

3.503.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.53 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.69

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx =$$

$$\frac{2a\sqrt{\cot(c+dx)}(A \cot(c+dx) + 3(iA+B) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, i \tan(c+dx)\right))}{3d}$$

input `Integrate[Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(-2*a*Sqrt[Cot[c + d*x]]*(A*Cot[c + d*x] + 3*(I*A + B)*Hypergeometric2F1[-1/2, 1, 1/2, I*Tan[c + d*x]]))/(3*d)`

3.503.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4064, 3042, 4075, 3042, 4011, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \cot(c + dx)^{5/2}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4064} \\
 & \int \sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)(A \cot(c + dx) + B) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{-\tan\left(c + dx + \frac{\pi}{2}\right)}\left(-a \tan\left(c + dx + \frac{\pi}{2}\right) + ia\right)\left(B - A \tan\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{4075} \\
 & -\frac{2aA \cot^{\frac{3}{2}}(c + dx)}{3d} + \int \sqrt{\cot(c + dx)}(a(iA + B) \cot(c + dx) - a(A - iB)) dx \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2aA \cot^{\frac{3}{2}}(c + dx)}{3d} + \int \sqrt{-\tan\left(c + dx + \frac{\pi}{2}\right)}\left(-a(A - iB) - a(iA + B) \tan\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{4011} \\
 & \int \frac{-a(iA + B) - a(A - iB) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx - \frac{2a(B + iA) \sqrt{\cot(c + dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c + dx)}{3d}
 \end{aligned}$$

3.503. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\begin{aligned}
 & \int \frac{a(A - iB) \tan(c + dx + \frac{\pi}{2}) - a(iA + B)}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx - \frac{2a(B + iA)\sqrt{\cot(c + dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c + dx)}{3d} \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \frac{2a^2(B + iA)^2 \int \frac{1}{a(iA+B) - a(A-iB)\cot(c+dx)} d\sqrt{\cot(c + dx)}}{d} - \frac{2a(B + iA)\sqrt{\cot(c + dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c + dx)}{3d} \\
 & \qquad \qquad \qquad \downarrow \text{4016} \\
 & \frac{2\sqrt[4]{-1}a(B + iA)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c + dx)}\right)}{d} - \frac{2a(B + iA)\sqrt{\cot(c + dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c + dx)}{3d} \\
 & \qquad \qquad \qquad \downarrow \text{221}
 \end{aligned}$$

```
input Int[Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]
```

```
output (-2*(-1)^(1/4)*a*(I*A + B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]]]/d - (2*a*(I*A + B)*Sqrt[Cot[c + d*x]]/d - (2*a*A*Cot[c + d*x]^(3/2))/(3*d))
```

3.503.3.1 Defintions of rubi rules used

```
rule 221 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4011 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]
```

3.503. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.503.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 527 vs. 2(64) = 128.

Time = 0.45 (sec) , antiderivative size = 528, normalized size of antiderivative = 6.77

method	result
derivativedivides	$-\frac{a\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}}\tan(dx+c)\left(6iA\arctan\left(1+\sqrt{2}\sqrt{\tan(dx+c)}\right)\sqrt{2}\tan(dx+c)^{\frac{3}{2}}+6iA\arctan\left(-1+\sqrt{2}\sqrt{\tan(dx+c)}\right)\sqrt{2}\tan(dx+c)^{\frac{3}{2}}\right)}{\dots}$
default	$-\frac{a\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}}\tan(dx+c)\left(6iA\arctan\left(1+\sqrt{2}\sqrt{\tan(dx+c)}\right)\sqrt{2}\tan(dx+c)^{\frac{3}{2}}+6iA\arctan\left(-1+\sqrt{2}\sqrt{\tan(dx+c)}\right)\sqrt{2}\tan(dx+c)^{\frac{3}{2}}\right)}{\dots}$

input `int(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNV ERBOSE)`

$$3.503. \int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

output

```
-1/12*a/d*(1/tan(d*x+c))^(5/2)*tan(d*x+c)*(6*I*A*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(3/2)+6*I*A*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(3/2)+3*I*A*ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))*2^(1/2)*tan(d*x+c)^(3/2)-3*I*B*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1))*2^(1/2)*tan(d*x+c)^(3/2)-6*I*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(3/2)-6*I*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(3/2)+3*A*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1))*2^(1/2)*tan(d*x+c)^(3/2)+6*A*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(3/2)+6*A*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(3/2)+6*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(3/2)+6*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(3/2)+3*B*ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))*2^(1/2)*tan(d*x+c)^(3/2)+24*I*A*tan(d*x+c)+24*B*tan(d*x+c)+8*A)
```

3.503.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 382 vs. $2(62) = 124$.

Time = 0.25 (sec) , antiderivative size = 382, normalized size of antiderivative = 4.90

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx =$$

$$3 \left(de^{(2i dx + 2i c)} - d \right) \sqrt{-\frac{(i A^2 + 2 AB - i B^2) a^2}{d^2}} \log \left(-\frac{2 \left((A - i B) a e^{(2i dx + 2i c)} + (de^{(2i dx + 2i c)} - d) \sqrt{-\frac{(i A^2 + 2 AB - i B^2) a^2}{d^2}} \sqrt{\frac{i e^{(2i dx + 2i c)}}{e^{(2i dx + 2i c)}}}}{(i A + B) a} \right)}{e^{(2i dx + 2i c)}} \right)$$

input `integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output

```
-1/6*(3*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)
*log(-2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) + (d*e^(2*I*d*x + 2*I*c) - d)*sqrt
(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2
*I*d*x + 2*I*c) - 1)))e^(-2*I*d*x - 2*I*c)/((I*A + B)*a) - 3*(d*e^(2*I*d
*x + 2*I*c) - d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*log(-2*((A - I*B)*
a*e^(2*I*d*x + 2*I*c) - (d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(I*A^2 + 2*A*B -
I*B^2)*a^2/d^2)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1
))))e^(-2*I*d*x - 2*I*c)/((I*A + B)*a) + 4*((4*I*A + 3*B)*a*e^(2*I*d*x +
2*I*c) + (-2*I*A - 3*B)*a)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x +
2*I*c) - 1)))/(d*e^(2*I*d*x + 2*I*c) - d)
```

3.503.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(5/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.503.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 174 vs. $2(62) = 124$.

Time = 0.40 (sec) , antiderivative size = 174, normalized size of antiderivative = 2.23

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx =$$

$$\frac{3 \left(2\sqrt{2}(-i+1)A + (i-1)B \right) \arctan \left(\frac{1}{2}\sqrt{2} \left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}} \right) \right) + 2\sqrt{2}(-i+1)A + (i-1)B}{\dots}$$

input `integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

3.503. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

output
$$\begin{aligned} & -1/12*(3*(2*\sqrt{2})*(-(I + 1)*A + (I - 1)*B)*\arctan(1/2*\sqrt{2}*(\sqrt{2} + \\ & 2/\sqrt{\tan(dx + c)})) + 2*\sqrt{2})*(-(I + 1)*A + (I - 1)*B)*\arctan(-1/2*\sqrt{2} \\ & *(\sqrt{2} - 2/\sqrt{\tan(dx + c)})) - \sqrt{2}*((I - 1)*A + (I + 1)*B) \\ & * \log(\sqrt{2}/\sqrt{\tan(dx + c)} + 1/\tan(dx + c) + 1) + \sqrt{2}*((I - 1)*A \\ & + (I + 1)*B)* \log(-\sqrt{2}/\sqrt{\tan(dx + c)} + 1/\tan(dx + c) + 1)) * a - 2 \\ & 4*(-I*A - B)*a/\sqrt{\tan(dx + c)} + 8*A*a/\tan(dx + c)^{(3/2)}/d \end{aligned}$$

3.503.8 Giac [F]

$$\begin{aligned} & \int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\ & = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a) \cot(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)*cot(d*x + c)^(5/2), x)`

3.503.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\ & = \int \cot(c + dx)^{5/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) li) dx \end{aligned}$$

input `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li),x)`

output `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li), x)`

3.504 $\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.504.1 Optimal result	4758
3.504.2 Mathematica [A] (verified)	4758
3.504.3 Rubi [A] (verified)	4759
3.504.4 Maple [B] (verified)	4761
3.504.5 Fricas [B] (verification not implemented)	4762
3.504.6 Sympy [F]	4762
3.504.7 Maxima [B] (verification not implemented)	4763
3.504.8 Giac [F]	4763
3.504.9 Mupad [F(-1)]	4764

3.504.1 Optimal result

Integrand size = 34, antiderivative size = 53

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= -\frac{2\sqrt[4]{-1}a(A-iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} - \frac{2aA\sqrt{\cot(c+dx)}}{d}$$

output `-2*(-1)^(1/4)*a*(A-I*B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d-2*a*A*cot(d*x+c)^(1/2)/d`

3.504.2 Mathematica [A] (verified)

Time = 0.77 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.21

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= -\frac{2ia\sqrt{\cot(c+dx)}\left(-iA+\sqrt[4]{-1}(A-iB)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)\sqrt{\tan(c+dx)}\right)}{d}$$

input `Integrate[Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output $((-2*I)*a*\text{Sqrt}[\text{Cot}[c + d*x]]*((-I)*A + (-1)^{(1/4)}*(A - I*B)*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]]*\text{Sqrt}[\text{Tan}[c + d*x]]))/d$

3.504.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.206$, Rules used = {3042, 4064, 3042, 4075, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \cot(c + dx)^{3/2}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4064} \\ & \int \frac{(a \cot(c + dx) + ia)(A \cot(c + dx) + B)}{\sqrt{\cot(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(-a \tan(c + dx + \frac{\pi}{2}) + ia)(B - A \tan(c + dx + \frac{\pi}{2}))}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx \\ & \quad \downarrow \text{4075} \\ & -\frac{2aA\sqrt{\cot(c + dx)}}{d} + \int \frac{a(iA + B)\cot(c + dx) - a(A - iB)}{\sqrt{\cot(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & -\frac{2aA\sqrt{\cot(c + dx)}}{d} + \int \frac{-a(A - iB) - a(iA + B)\tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx \\ & \quad \downarrow \text{4016} \\ & -\frac{2aA\sqrt{\cot(c + dx)}}{d} + \frac{2a^2(A - iB)^2}{d} \int \frac{1}{a(A - iB) + a(iA + B)\cot(c + dx)} d\sqrt{\cot(c + dx)} \\ & \quad \downarrow \text{221} \end{aligned}$$

3.504. $\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

$$-\frac{2aA\sqrt{\cot(c+dx)}}{d} - \frac{2\sqrt[4]{-1}a(A-iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d}$$

input `Int[Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(-2*(-1)^(1/4)*a*(A - I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]])/d - (2*a*A*Sqrt[Cot[c + d*x]])/d`

3.504.3.1 Defintions of rubi rules used

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m+n) Int[(g*Cot[e + f*x])^(p-m-n)*(b+a*Cot[e + f*x])^m*(d+c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m+1)/(b*f*(m+1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.504.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 507 vs. $2(44) = 88$.

Time = 0.36 (sec) , antiderivative size = 508, normalized size of antiderivative = 9.58

method	result
derivativedivides	$\frac{a \left(\frac{1}{\tan(dx+c)} \right)^{\frac{3}{2}} \tan(dx+c) \left(2iA \arctan \left(1 + \sqrt{2} \sqrt{\tan(dx+c)} \right) \sqrt{2} \sqrt{\tan(dx+c)} + 2iA \arctan \left(-1 + \sqrt{2} \sqrt{\tan(dx+c)} \right) \sqrt{2} \right)}{\dots}$
default	$\frac{a \left(\frac{1}{\tan(dx+c)} \right)^{\frac{3}{2}} \tan(dx+c) \left(2iA \arctan \left(1 + \sqrt{2} \sqrt{\tan(dx+c)} \right) \sqrt{2} \sqrt{\tan(dx+c)} + 2iA \arctan \left(-1 + \sqrt{2} \sqrt{\tan(dx+c)} \right) \sqrt{2} \right)}{\dots}$

```
input int(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 1/4*a/d*(1/tan(d*x+c))^(3/2)*tan(d*x+c)*(2*I*A*arctan(1+2^(1/2)*tan(d*x+c)
^(1/2))*2^(1/2)*tan(d*x+c)^(1/2)+2*I*A*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))
*2^(1/2)*tan(d*x+c)^(1/2)+I*A*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/
(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1))*2^(1/2)*tan(d*x+c)^(1/2)+2*I*B*ar
ctan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(1/2)+2*I*B*arctan(-1+
2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(1/2)+I*B*ln(-(2^(1/2)*tan(d*
x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))*2^(1/2)*
tan(d*x+c)^(1/2)-2*A*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)
^(1/2)-2*A*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(1/2)-A*
ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+ta
n(d*x+c)))*2^(1/2)*tan(d*x+c)^(1/2)+2*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))
*2^(1/2)*tan(d*x+c)^(1/2)+2*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*
tan(d*x+c)^(1/2)+B*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*ta
n(d*x+c)^(1/2)-tan(d*x+c)-1))*2^(1/2)*tan(d*x+c)^(1/2)-8*A)
```


3.504.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 316 vs. $2(43) = 86$.

Time = 0.24 (sec) , antiderivative size = 316, normalized size of antiderivative = 5.96

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx =$$

$$4Aa \sqrt{\frac{ie^{(2i dx+2i c)+i}}{e^{(2i dx+2i c)}-1}} - \sqrt{\frac{(-iA^2-2AB+iB^2)a^2}{d^2}} d \log \left(\frac{2 \left((A-iB)ae^{(2i dx+2i c)} - (ide^{(2i dx+2i c)}-id) \sqrt{-\frac{(-iA^2-2AB+iB^2)a^2}{d^2}} \right)}{(iA+B)a} \right)$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-1/2*(4*A*a*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)) - sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*d*log(-2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) - (I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a)) + sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*d*log(-2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) - (-I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a))/d`

3.504.6 Sympy [F]

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= ia \left(\int \left(-iA \cot^{\frac{3}{2}}(c+dx) \right) dx + \int A \tan(c+dx) \cot^{\frac{3}{2}}(c+dx) dx \right.$$

$$\left. + \int B \tan^2(c+dx) \cot^{\frac{3}{2}}(c+dx) dx + \int \left(-iB \tan(c+dx) \cot^{\frac{3}{2}}(c+dx) \right) dx \right)$$

input `integrate(cot(d*x+c)**(3/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `I*a*(Integral(-I*A*cot(c + d*x)**(3/2), x) + Integral(A*tan(c + d*x)*cot(c + d*x)**(3/2), x) + Integral(B*tan(c + d*x)**2*cot(c + d*x)**(3/2), x) + Integral(-I*B*tan(c + d*x)*cot(c + d*x)**(3/2), x))`

3.504. $\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

3.504.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 155 vs. $2(43) = 86$.

Time = 0.37 (sec) , antiderivative size = 155, normalized size of antiderivative = 2.92

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx =$$

$$\frac{\left(2\sqrt{2}((i-1)A + (i+1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}((i-1)A + (i+1)B) \arctan\right.}{-}$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/4*((2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a + 8*A*a/sqrt(tan(d*x + c)))/d`

3.504.8 Giac [F]

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a) \cot(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)*cot(d*x + c)^(3/2), x)`

3.504.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{3/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) \text{ li}) dx$$

input `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`output `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i), x)`

3.505 $\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$

3.505.1 Optimal result	4765
3.505.2 Mathematica [A] (verified)	4765
3.505.3 Rubi [A] (verified)	4766
3.505.4 Maple [B] (verified)	4768
3.505.5 Fricas [B] (verification not implemented)	4768
3.505.6 Sympy [F]	4769
3.505.7 Maxima [B] (verification not implemented)	4770
3.505.8 Giac [F]	4770
3.505.9 Mupad [F(-1)]	4771

3.505.1 Optimal result

Integrand size = 34, antiderivative size = 55

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2\sqrt[4]{-1}a(iA + B)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c + dx)}\right)}{d} + \frac{2iaB}{d\sqrt{\cot(c + dx)}}$$

output `2*(-1)^(1/4)*a*(I*A+B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d+2*I*a*B/d/cot(d*x+c)^(1/2)`

3.505.2 Mathematica [A] (verified)

Time = 0.73 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.15

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2a\left(iB - \frac{\sqrt[4]{-1}(A-iB)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{\sqrt{\tan(c+dx)}}\right)}{d\sqrt{\cot(c + dx)}}$$

input `Integrate[Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output $(2*a*(I*B - ((-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]])/Sqrt[Tan[c + d*x]]/(d*Sqrt[Cot[c + d*x]])$

3.505.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.206$, Rules used = {3042, 4064, 3042, 4074, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{(a \cot(c+dx)+ia)(A \cot(c+dx)+B)}{\cot^{\frac{3}{2}}(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-a \tan(c+dx+\frac{\pi}{2})+ia)(B-A \tan(c+dx+\frac{\pi}{2}))}{(-\tan(c+dx+\frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{4074} \\
 & \int \frac{a(iA+B)+a(A-iB) \cot(c+dx)}{\sqrt{\cot(c+dx)}} dx + \frac{2iaB}{d\sqrt{\cot(c+dx)}} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{a(iA+B)-a(A-iB) \tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx + \frac{2iaB}{d\sqrt{\cot(c+dx)}} \\
 & \quad \downarrow \text{4016} \\
 & \frac{2a^2(B+iA)^2}{d} \int \frac{1}{a(A-iB) \cot(c+dx)-a(iA+B)} d\sqrt{\cot(c+dx)} + \frac{2iaB}{d\sqrt{\cot(c+dx)}} \\
 & \quad \downarrow \text{221}
 \end{aligned}$$

3.505. $\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

$$\frac{2\sqrt[4]{-1}a(B + iA)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c + dx)}\right)}{d} + \frac{2iaB}{d\sqrt{\cot(c + dx)}}$$

input `Int[Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(2*(-1)^(1/4)*a*(I*A + B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]]]/d + ((2*I)*a*B)/(d*Sqrt[Cot[c + d*x]])`

3.505.3.1 Defintions of rubi rules used

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4016 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)])*(g_.)^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4074 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.505.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 421 vs. 2(45) = 90.

Time = 0.36 (sec) , antiderivative size = 422, normalized size of antiderivative = 7.67

method	result
derivativedivides	$a\sqrt{\frac{1}{\tan(dx+c)}}\sqrt{\tan(dx+c)}\left(2iA\sqrt{2}\arctan\left(1+\sqrt{2}\sqrt{\tan(dx+c)}\right)+2iA\sqrt{2}\arctan\left(-1+\sqrt{2}\sqrt{\tan(dx+c)}\right)+iA\sqrt{2}\ln\left(-\frac{1+\sqrt{2}\sqrt{\tan(dx+c)}}{1-\sqrt{2}\sqrt{\tan(dx+c)}}\right)\right)$
default	$a\sqrt{\frac{1}{\tan(dx+c)}}\sqrt{\tan(dx+c)}\left(2iA\sqrt{2}\arctan\left(1+\sqrt{2}\sqrt{\tan(dx+c)}\right)+2iA\sqrt{2}\arctan\left(-1+\sqrt{2}\sqrt{\tan(dx+c)}\right)+iA\sqrt{2}\ln\left(-\frac{1+\sqrt{2}\sqrt{\tan(dx+c)}}{1-\sqrt{2}\sqrt{\tan(dx+c)}}\right)\right)$

```
input int(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 1/4*a/d*(1/tan(d*x+c))^(1/2)*tan(d*x+c)^(1/2)*(2*I*A*2^(1/2)*arctan(1+2^(1
/2)*tan(d*x+c)^(1/2))+2*I*A*2^(1/2)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))+I*
A*2^(1/2)*ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c
)^(1/2)+tan(d*x+c))) -I*B*2^(1/2)*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c
)))/(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)) -2*I*B*2^(1/2)*arctan(1+2^(1/2)
*tan(d*x+c)^(1/2)) -2*I*B*2^(1/2)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))+A*2^(
1/2)*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*tan(d*x+c)^(1/2)
-tan(d*x+c)-1))+2*A*2^(1/2)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*A*2^(1/2)
*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))+8*I*B*tan(d*x+c)^(1/2)+2*B*2^(1/2)*ar
ctan(1+2^(1/2)*tan(d*x+c)^(1/2))+2*B*2^(1/2)*arctan(-1+2^(1/2)*tan(d*x+c)^(
1/2))+B*2^(1/2)*ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*ta
n(d*x+c)^(1/2)+tan(d*x+c)))
```

3.505.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 364 vs. 2(43) = 86.

Time = 0.25 (sec) , antiderivative size = 364, normalized size of antiderivative = 6.62

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{(de^{(2i dx+2i c)} + d) \sqrt{-\frac{(i A^2+2 AB-i B^2)a^2}{d^2}} \log\left(-\frac{2\left((A-i B)ae^{(2i dx+2i c)}+(de^{(2i dx+2i c)}-d)\sqrt{-\frac{(i A^2+2 AB-i B^2)a^2}{d^2}}\sqrt{\frac{ie^{(2i dx+2i c)}}{e^{(2i dx+2i c)}}}\right)}{(i A+B)a}}\right)}{1}$$

3.505. $\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/2*((d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*log(-2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) + (d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a)) - (d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*log(-2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) - (d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a)) + 4*(B*a*e^(2*I*d*x + 2*I*c) - B*a)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))/(d*e^(2*I*d*x + 2*I*c) + d)`

3.505.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx \\ &= ia \left(\int (-iA \sqrt{\cot(c+dx)}) dx + \int A \tan(c+dx) \sqrt{\cot(c+dx)} dx \right. \\ & \quad \left. + \int B \tan^2(c+dx) \sqrt{\cot(c+dx)} dx + \int (-iB \tan(c+dx) \sqrt{\cot(c+dx)}) dx \right) \end{aligned}$$

input `integrate(cot(d*x+c)**(1/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `I*a*(Integral(-I*A*sqrt(cot(c + d*x)), x) + Integral(A*tan(c + d*x)*sqrt(cot(c + d*x)), x) + Integral(B*tan(c + d*x)**2*sqrt(cot(c + d*x)), x) + Integral(-I*B*tan(c + d*x)*sqrt(cot(c + d*x)), x))`

3.505.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 155 vs. $2(43) = 86$.

Time = 0.76 (sec) , antiderivative size = 155, normalized size of antiderivative = 2.82

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{8iBa\sqrt{\tan(dx+c)} + \left(2\sqrt{2}(-(i+1)A+(i-1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}(-(i+1)A+(i-1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) - \sqrt{2}((i-1)A+(i+1)B) \log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + \frac{1}{\tan(dx+c)} + 1\right) + \sqrt{2}((i-1)A+(i+1)B) \log\left(-\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + \frac{1}{\tan(dx+c)} + 1\right)\right)a}{d}$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/4*(8*I*B*a*sqrt(tan(d*x + c)) + (2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a)/d`

3.505.8 Giac [F]

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \int (B \tan(dx+c) + A)(ia \tan(dx+c) + a)\sqrt{\cot(dx+c)} dx$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)*sqrt(cot(d*x + c)), x)`

3.505.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \int \sqrt{\cot(c+dx)}(A+B \tan(c+dx))(a+a \tan(c+dx) li) dx$$

input `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i),x)`output `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i), x)`

3.506
$$\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.506.1 Optimal result 4772
 3.506.2 Mathematica [A] (verified) 4772
 3.506.3 Rubi [A] (verified) 4773
 3.506.4 Maple [B] (verified) 4775
 3.506.5 Fricas [B] (verification not implemented) 4776
 3.506.6 Sympy [F] 4776
 3.506.7 Maxima [B] (verification not implemented) 4777
 3.506.8 Giac [F] 4777
 3.506.9 Mupad [F(-1)] 4778

3.506.1 Optimal result

Integrand size = 34, antiderivative size = 80

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \frac{2\sqrt[4]{-1}a(A - iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c + dx)}\right)}{d} + \frac{2iaB}{3d \cot^{3/2}(c + dx)} + \frac{2a(iA + B)}{d\sqrt{\cot(c + dx)}}$$

output `2*(-1)^(1/4)*a*(A-I*B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d+2/3*I*a*B/d/cot(d*x+c)^(3/2)+2*a*(I*A+B)/d/cot(d*x+c)^(1/2)`

3.506.2 Mathematica [A] (verified)

Time = 1.04 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.08

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \frac{2ia\sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}\left(3(A - iB)\left(\sqrt[4]{-1}\arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right) + \sqrt{\tan(c + dx)}\right) + B\right)}{3d}$$

input `Integrate[((a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output $((2I/3)*a*\text{Sqrt}[\text{Cot}[c + dx]]*\text{Sqrt}[\text{Tan}[c + dx]]*(3*(A - I*B)*((-1)^{1/4})*\text{ArcTan}[(-1)^{3/4}*\text{Sqrt}[\text{Tan}[c + dx]]] + \text{Sqrt}[\text{Tan}[c + dx]]) + B*\text{Tan}[c + dx]^{3/2})/d$

3.506.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.265$, Rules used = {3042, 4064, 3042, 4074, 3042, 4012, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\ & \quad \downarrow 3042 \\ & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\ & \quad \downarrow 4064 \\ & \int \frac{(a \cot(c + dx) + ia)(A \cot(c + dx) + B)}{\cot^{5/2}(c + dx)} dx \\ & \quad \downarrow 3042 \\ & \int \frac{(-a \tan(c + dx + \frac{\pi}{2}) + ia)(B - A \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{5/2}} dx \\ & \quad \downarrow 4074 \\ & \int \frac{a(iA + B) + a(A - iB) \cot(c + dx)}{\cot^{3/2}(c + dx)} dx + \frac{2iaB}{3d \cot^{3/2}(c + dx)} \\ & \quad \downarrow 3042 \\ & \int \frac{a(iA + B) - a(A - iB) \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}} dx + \frac{2iaB}{3d \cot^{3/2}(c + dx)} \\ & \quad \downarrow 4012 \\ & \int \frac{a(A - iB) - a(iA + B) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx + \frac{2a(B + iA)}{d \sqrt{\cot(c + dx)}} + \frac{2iaB}{3d \cot^{3/2}(c + dx)} \\ & \quad \downarrow 3042 \end{aligned}$$

$$\int \frac{a(A - iB) + a(iA + B) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx + \frac{2a(B + iA)}{d\sqrt{\cot(c + dx)}} + \frac{2iaB}{3d \cot^{\frac{3}{2}}(c + dx)}$$

↓ 4016

$$\frac{2a^2(A - iB)^2 \int \frac{1}{-a(A - iB) - a(iA + B) \cot(c + dx)} d\sqrt{\cot(c + dx)}}{d} + \frac{2a(B + iA)}{d\sqrt{\cot(c + dx)}} + \frac{2iaB}{3d \cot^{\frac{3}{2}}(c + dx)}$$

↓ 221

$$\frac{2\sqrt[4]{-1}a(A - iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c + dx)}\right)}{d} + \frac{2a(B + iA)}{d\sqrt{\cot(c + dx)}} + \frac{2iaB}{3d \cot^{\frac{3}{2}}(c + dx)}$$

input `Int[((a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `(2*(-1)^(1/4)*a*(A - I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]])/d + (((2*I)/3)*a*B)/(d*Cot[c + d*x]^(3/2)) + (2*a*(I*A + B))/(d*Sqrt[Cot[c + d*x]])`

3.506.3.1 Defintions of rubi rules used

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

```
rule 4064 Int[(cot[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(m_.))*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp [g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c *Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !Integer Q[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(m_))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b *c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]
```

3.506.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 219 vs. 2(65) = 130.

Time = 0.36 (sec) , antiderivative size = 220, normalized size of antiderivative = 2.75

method	result
derivativedivides	$a \left(\frac{(-iB+A)\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan(1+\sqrt{2}\sqrt{\cot(dx+c)}) + 2 \arctan(-1+\sqrt{2}\sqrt{\cot(dx+c)}) \right)}{4} \right) + \dots$
default	$a \left(\frac{(-iB+A)\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan(1+\sqrt{2}\sqrt{\cot(dx+c)}) + 2 \arctan(-1+\sqrt{2}\sqrt{\cot(dx+c)}) \right)}{4} \right) + \dots$

```
input int((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x,method=_RETURNV ERBOSE)
```

```
output -a/d*(1/4*(A-I*B)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+c ot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+ 2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/4*(-I*A-B)*2^(1/2)*(ln((1+cot(d*x +c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*a rctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))-2 *(I*A+B)/cot(d*x+c)^(1/2)-2/3*I*B/cot(d*x+c)^(3/2))
```

$$3.506. \int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.506.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 429 vs. $2(62) = 124$.

Time = 0.25 (sec) , antiderivative size = 429, normalized size of antiderivative = 5.36

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx =$$

$$3 \left(de^{(4i dx + 4i c)} + 2 de^{(2i dx + 2i c)} + d \right) \sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^2}{d^2}} \log \left(-\frac{2 \left((A - iB)ae^{(2i dx + 2i c)} - (i de^{(2i dx + 2i c)} - i d) \sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^2}{d^2}} \right)}{(iA + B)a} \right)$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="fricas")`

output `-1/6*(3*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*log(-2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) - (I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c))/(I*A + B)*a) - 3*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*log(-2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) - (I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^2/d^2)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c))/(I*A + B)*a) - 4*((3*A - 4*I*B)*a*e^(4*I*d*x + 4*I*c) + 2*I*B*a*e^(2*I*d*x + 2*I*c) - (3*A - 2*I*B)*a)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))/(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)`

3.506.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= ia \left(\int \left(-\frac{iA}{\sqrt{\cot(c + dx)}} \right) dx + \int \frac{A \tan(c + dx)}{\sqrt{\cot(c + dx)}} dx + \int \frac{B \tan^2(c + dx)}{\sqrt{\cot(c + dx)}} dx + \int \left(-\frac{iB \tan(c + dx)}{\sqrt{\cot(c + dx)}} \right) dx \right)$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)**(1/2),x)`

output `I*a*(Integral(-I*A/sqrt(cot(c + d*x)), x) + Integral(A*tan(c + d*x)/sqrt(cot(c + d*x)), x) + Integral(B*tan(c + d*x)**2/sqrt(cot(c + d*x)), x) + Integral(-I*B*tan(c + d*x)/sqrt(cot(c + d*x)), x))`

3.506.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 177 vs. $2(62) = 124$.

Time = 0.29 (sec) , antiderivative size = 177, normalized size of antiderivative = 2.21

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \frac{8 \left(i B a - \frac{3(-i A - B)a}{\tan(dx+c)} \right) \tan(dx + c)^{\frac{3}{2}} + 3 \left(2\sqrt{2}((i-1)A + (i+1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) \right)}{d}$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="maxima")`

output `1/12*(8*(I*B*a - 3*(-I*A - B)*a/tan(d*x + c))*tan(d*x + c)^(3/2) + 3*(2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a)/d`

3.506.8 Giac [F]

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)/sqrt(cot(d*x + c)), x)`

3.506.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) 1i)}{\sqrt{\cot(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i))/cot(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i))/cot(c + d*x)^(1/2), x)`

3.507 $\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

3.507.1 Optimal result 4779
 3.507.2 Mathematica [A] (verified) 4779
 3.507.3 Rubi [A] (verified) 4780
 3.507.4 Maple [B] (verified) 4783
 3.507.5 Fricas [B] (verification not implemented) 4784
 3.507.6 Sympy [F] 4784
 3.507.7 Maxima [B] (verification not implemented) 4785
 3.507.8 Giac [F] 4785
 3.507.9 Mupad [F(-1)] 4786

3.507.1 Optimal result

Integrand size = 34, antiderivative size = 105

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= -\frac{2\sqrt[4]{-1}a(iA + B)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c + dx)}\right)}{d} + \frac{2iaB}{5d \cot^{\frac{5}{2}}(c + dx)} + \frac{2a(iA + B)}{3d \cot^{\frac{3}{2}}(c + dx)} + \frac{2a(A - iB)}{d\sqrt{\cot(c + dx)}}$$

output `-2*(-1)^(1/4)*a*(I*A+B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d+2/5*I*a*B/d/cot(d*x+c)^(5/2)+2/3*a*(I*A+B)/d/cot(d*x+c)^(3/2)+2*a*(A-I*B)/d/cot(d*x+c)^(1/2)`

3.507.2 Mathematica [A] (verified)

Time = 1.99 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.94

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{2ia\sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}\left(3B \tan^{\frac{5}{2}}(c + dx) + 5(A - iB)\right)\left(-3(-1)^{3/4} \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right)\right)}{15d}$$

input `Integrate[((a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(3/2),x]`

output `((2*I)/15)*a*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(3*B*Tan[c + d*x]^(5/2) + 5*(A - I*B)*(-3*(-1)^(3/4)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + Sqrt[Tan[c + d*x]]*(-3*I + Tan[c + d*x])))/d`

3.507.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3042, 4064, 3042, 4074, 3042, 4012, 3042, 4012, 25, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\cot(c + dx)^{3/2}} dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{(a \cot(c + dx) + ia)(A \cot(c + dx) + B)}{\cot^{\frac{7}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-a \tan(c + dx + \frac{\pi}{2}) + ia)(B - A \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{7/2}} dx \\
 & \quad \downarrow \text{4074} \\
 & \int \frac{a(iA + B) + a(A - iB) \cot(c + dx)}{\cot^{\frac{5}{2}}(c + dx)} dx + \frac{2iaB}{5d \cot^{\frac{5}{2}}(c + dx)} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{a(iA + B) - a(A - iB) \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{5/2}} dx + \frac{2iaB}{5d \cot^{\frac{5}{2}}(c + dx)} \\
 & \quad \downarrow \text{4012}
 \end{aligned}$$

3.507. $\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \int \frac{a(A - iB) - a(iA + B) \cot(c + dx)}{\cot^{\frac{3}{2}}(c + dx)} dx + \frac{2a(B + iA)}{3d \cot^{\frac{3}{2}}(c + dx)} + \frac{2iaB}{5d \cot^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \int \frac{a(A - iB) + a(iA + B) \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}} dx + \frac{2a(B + iA)}{3d \cot^{\frac{3}{2}}(c + dx)} + \frac{2iaB}{5d \cot^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{4012} \\
& \int -\frac{a(iA + B) + a(A - iB) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx + \frac{2a(B + iA)}{3d \cot^{\frac{3}{2}}(c + dx)} + \frac{2a(A - iB)}{d\sqrt{\cot(c + dx)}} + \frac{2iaB}{5d \cot^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{25} \\
& -\int \frac{a(iA + B) + a(A - iB) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx + \frac{2a(B + iA)}{3d \cot^{\frac{3}{2}}(c + dx)} + \frac{2a(A - iB)}{d\sqrt{\cot(c + dx)}} + \frac{2iaB}{5d \cot^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& -\int \frac{a(iA + B) - a(A - iB) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx + \frac{2a(B + iA)}{3d \cot^{\frac{3}{2}}(c + dx)} + \frac{2a(A - iB)}{d\sqrt{\cot(c + dx)}} + \\
& \quad \frac{2iaB}{5d \cot^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{4016} \\
& -\frac{2a^2(B + iA)^2 \int \frac{1}{a(A - iB) \cot(c + dx) - a(iA + B)} d\sqrt{\cot(c + dx)}}{d} + \frac{2a(B + iA)}{3d \cot^{\frac{3}{2}}(c + dx)} + \frac{2a(A - iB)}{d\sqrt{\cot(c + dx)}} + \\
& \quad \frac{2iaB}{5d \cot^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{221} \\
& -\frac{2\sqrt[4]{-1}a(B + iA) \operatorname{arctanh}\left((-1)^{3/4} \sqrt{\cot(c + dx)}\right)}{d} + \frac{2a(B + iA)}{3d \cot^{\frac{3}{2}}(c + dx)} + \frac{2a(A - iB)}{d\sqrt{\cot(c + dx)}} + \\
& \quad \frac{2iaB}{5d \cot^{\frac{5}{2}}(c + dx)}
\end{aligned}$$

input `Int[((a + I*a*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(3/2),x]`

output `(-2*(-1)^(1/4)*a*(I*A + B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]])/d + (((2*I)/5)*a*B)/(d*Cot[c + d*x]^(5/2)) + (2*a*(I*A + B))/(3*d*Cot[c + d*x]^(3/2)) + (2*a*(A - I*B))/(d*Sqrt[Cot[c + d*x]])`

3.507. $\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

3.507.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`
- rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`
- rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4074 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.507.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 469 vs. $2(85) = 170$.

Time = 0.40 (sec) , antiderivative size = 470, normalized size of antiderivative = 4.48

method	result
derivativedivides	$-\frac{a \left(120iB \sqrt{\tan(dx+c)} + 30iA\sqrt{2} \arctan\left(1+\sqrt{2} \sqrt{\tan(dx+c)}\right) - 40iA \tan(dx+c)^{\frac{3}{2}} - 30iB\sqrt{2} \arctan\left(-1+\sqrt{2} \sqrt{\tan(dx+c)}\right) \right)}{\dots}$
default	$-\frac{a \left(120iB \sqrt{\tan(dx+c)} + 30iA\sqrt{2} \arctan\left(1+\sqrt{2} \sqrt{\tan(dx+c)}\right) - 40iA \tan(dx+c)^{\frac{3}{2}} - 30iB\sqrt{2} \arctan\left(-1+\sqrt{2} \sqrt{\tan(dx+c)}\right) \right)}{\dots}$

input `int((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x,method=_RETURNV
ERBOSE)`

output
$$\begin{aligned} & -1/60*a/d*(120*I*B*\tan(d*x+c)^{(1/2)}+30*I*A*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}) \\ & *2^{(1/2)}-40*I*A*\tan(d*x+c)^{(3/2)}-30*I*B*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}) \\ & *2^{(1/2)}-30*I*B*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})*2^{(1/2)}+15*I*A*\ln(- \\ & (2^{(1/2)}*\tan(d*x+c)^{(1/2)}-\tan(d*x+c)-1)/(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d* \\ & x+c)))*2^{(1/2)}-15*I*B*\ln(-(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(2^{(1/2)} \\ & * \tan(d*x+c)^{(1/2)}-\tan(d*x+c)-1))*2^{(1/2)}+30*I*A*\arctan(-1+2^{(1/2)}*\tan(d*x+ \\ & c)^{(1/2)})*2^{(1/2)}-40*B*\tan(d*x+c)^{(3/2)}+15*A*2^{(1/2)}*\ln(-(1+2^{(1/2)}*\tan(d* \\ & x+c)^{(1/2)}+\tan(d*x+c))/(2^{(1/2)}*\tan(d*x+c)^{(1/2)}-\tan(d*x+c)-1))+30*A*2^{(1/2)} \\ & *\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+30*A*2^{(1/2)}*\arctan(-1+2^{(1/2)}*\tan(d \\ & *x+c)^{(1/2)})-24*I*B*\tan(d*x+c)^{(5/2)}+30*B*2^{(1/2)}*\arctan(1+2^{(1/2)}*\tan(d*x \\ & +c)^{(1/2)})+30*B*2^{(1/2)}*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})+15*B*2^{(1/2)}* \\ & \ln(-(2^{(1/2)}*\tan(d*x+c)^{(1/2)}-\tan(d*x+c)-1)/(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan \\ & (d*x+c)))-120*A*\tan(d*x+c)^{(1/2)}/(1/\tan(d*x+c))^{(3/2)}/\tan(d*x+c)^{(3/2)} \end{aligned}$$

3.507.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 482 vs. $2(81) = 162$.

Time = 0.27 (sec) , antiderivative size = 482, normalized size of antiderivative = 4.59

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx =$$

$$15 \left(de^{(6i dx + 6i c)} + 3 de^{(4i dx + 4i c)} + 3 de^{(2i dx + 2i c)} + d \right) \sqrt{-\frac{(i A^2 + 2 AB - i B^2) a^2}{d^2}} \log \left(-\frac{2 \left((A - i B) a e^{(2i dx + 2i c)} + (de^{(6i dx + 6i c)} + 3 de^{(4i dx + 4i c)} + 3 de^{(2i dx + 2i c)} + d) \right)}{\dots} \right)$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algorithm="fricas")`

output `-1/30*(15*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*log(-2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) + (d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a) - 15*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*log(-2*((A - I*B)*a*e^(2*I*d*x + 2*I*c) - (d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^2/d^2)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/((I*A + B)*a) + 4*((20*I*A + 23*B)*a*e^(6*I*d*x + 6*I*c) + (10*I*A + B)*a*e^(4*I*d*x + 4*I*c) + (-20*I*A - 11*B)*a*e^(2*I*d*x + 2*I*c) + (-10*I*A - 13*B)*a)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))/(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)`

3.507.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= ia \left(\int \left(-\frac{iA}{\cot^{\frac{3}{2}}(c + dx)} \right) dx + \int \frac{A \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)} dx + \int \frac{B \tan^2(c + dx)}{\cot^{\frac{3}{2}}(c + dx)} dx + \int \left(-\frac{iB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)} \right) dx \right)$$

3.507. $\int \frac{(a+ia \tan(c+dx))(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)**(3/2),x)`

output `I*a*(Integral(-I*A/cot(c + d*x)**(3/2), x) + Integral(A*tan(c + d*x)/cot(c + d*x)**(3/2), x) + Integral(B*tan(c + d*x)**2/cot(c + d*x)**(3/2), x) + Integral(-I*B*tan(c + d*x)/cot(c + d*x)**(3/2), x))`

3.507.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 191 vs. $2(81) = 162$.

Time = 0.31 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.82

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx =$$

$$8 \left(-3i Ba - \frac{5(iA+B)a}{\tan(dx+c)} - \frac{15(A-iB)a}{\tan(dx+c)^2} \right) \tan(dx+c)^{\frac{5}{2}} + 15 \left(2\sqrt{2}(-i+1)A + (i-1)B \right) \arctan\left(\frac{1}{2}\sqrt{2}\left(\frac{A+B}{\tan(dx+c)} + 1\right)\right)$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algorithm="maxima")`

output `-1/60*(8*(-3*I*B*a - 5*(I*A + B)*a/tan(d*x + c) - 15*(A - I*B)*a/tan(d*x + c)^2)*tan(d*x + c)^(5/2) + 15*(2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a/d`

3.507.8 Giac [F]

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)}{\cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+I*a*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)/cot(d*x + c)^(3/2), x)`

3.507.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) 1i)}{\cot(c + dx)^{3/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i))/cot(c + d*x)^(3/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i))/cot(c + d*x)^(3/2), x)`

3.508 $\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.508.1 Optimal result	4787
3.508.2 Mathematica [C] (verified)	4787
3.508.3 Rubi [A] (verified)	4788
3.508.4 Maple [B] (verified)	4791
3.508.5 Fricas [B] (verification not implemented)	4792
3.508.6 Sympy [F(-1)]	4793
3.508.7 Maxima [A] (verification not implemented)	4793
3.508.8 Giac [F]	4794
3.508.9 Mupad [F(-1)]	4794

3.508.1 Optimal result

Integrand size = 36, antiderivative size = 128

$$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{4\sqrt{-1}a^2(A-iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} + \frac{4a^2(A-iB)\sqrt{\cot(c+dx)}}{d}$$

$$- \frac{2a^2(7iA+5B)\cot^{\frac{3}{2}}(c+dx)}{15d} - \frac{2A\cot^{\frac{3}{2}}(c+dx)(ia^2+a^2\cot(c+dx))}{5d}$$

```
output 4*(-1)^(1/4)*a^2*(A-I*B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d-2/15*a^2*(
7*I*A+5*B)*cot(d*x+c)^(3/2)/d-2/5*A*cot(d*x+c)^(3/2)*(I*a^2+a^2*cot(d*x+c)
)/d+4*a^2*(A-I*B)*cot(d*x+c)^(1/2)/d
```

3.508.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.58 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.58

$$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx =$$

$$\frac{2a^2\sqrt{\cot(c+dx)}(\cot(c+dx)(5(2iA+B)+3A\cot(c+dx))-30(A-iB)\operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1\right))}{15d}$$

3.508. $\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

input `Integrate[Cot[c + d*x]^(7/2)*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(-2*a^2*Sqrt[Cot[c + d*x]]*(Cot[c + d*x]*(5*((2*I)*A + B) + 3*A*Cot[c + d*x]) - 30*(A - I*B)*Hypergeometric2F1[-1/2, 1, 1/2, I*Tan[c + d*x]]))/(15*d)`

3.508.3 Rubi [A] (verified)

Time = 0.84 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.04, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3042, 4064, 3042, 4077, 27, 3042, 4075, 3042, 4011, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \cot(c + dx)^{7/2}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4064} \\
 & \int \sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2(A \cot(c + dx) + B) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{-\tan\left(c + dx + \frac{\pi}{2}\right)} \left(-a \tan\left(c + dx + \frac{\pi}{2}\right) + ia\right)^2 \left(B - A \tan\left(c + dx + \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{4077} \\
 & -\frac{2}{5} \int \frac{1}{2} \sqrt{\cot(c + dx)} (\cot(c + dx)a + ia)(a(3A - 5iB) - a(7iA + 5B) \cot(c + dx)) dx - \\
 & \quad \frac{2A \cot^{\frac{3}{2}}(c + dx) (a^2 \cot(c + dx) + ia^2)}{5d} \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{5} \int \sqrt{\cot(c + dx)} (\cot(c + dx)a + ia)(a(3A - 5iB) - a(7iA + 5B) \cot(c + dx)) dx - \\
 & \quad \frac{2A \cot^{\frac{3}{2}}(c + dx) (a^2 \cot(c + dx) + ia^2)}{5d}
 \end{aligned}$$

3.508. $\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

↓ 3042

$$-\frac{1}{5} \int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)} \left(ia - a \tan\left(c+dx+\frac{\pi}{2}\right) \right) \left(a(3A-5iB) + a(7iA+5B) \tan\left(c+dx+\frac{\pi}{2}\right) \right) dx - \frac{2A \cot^{\frac{3}{2}}(c+dx) (a^2 \cot(c+dx) + ia^2)}{5d}$$

↓ 4075

$$\frac{1}{5} \left(- \int \sqrt{\cot(c+dx)} (10(iA+B)a^2 + 10(A-iB) \cot(c+dx)a^2) dx - \frac{2a^2(5B+7iA) \cot^{\frac{3}{2}}(c+dx)}{3d} \right) - \frac{2A \cot^{\frac{3}{2}}(c+dx) (a^2 \cot(c+dx) + ia^2)}{5d}$$

↓ 3042

$$\frac{1}{5} \left(- \int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)} \left(10a^2(iA+B) - 10a^2(A-iB) \tan\left(c+dx+\frac{\pi}{2}\right) \right) dx - \frac{2a^2(5B+7iA) \cot^{\frac{3}{2}}(c+dx)}{3d} \right) - \frac{2A \cot^{\frac{3}{2}}(c+dx) (a^2 \cot(c+dx) + ia^2)}{5d}$$

↓ 4011

$$\frac{1}{5} \left(- \int \frac{10a^2(iA+B) \cot(c+dx) - 10a^2(A-iB)}{\sqrt{\cot(c+dx)}} dx - \frac{2a^2(5B+7iA) \cot^{\frac{3}{2}}(c+dx)}{3d} + \frac{20a^2(A-iB) \sqrt{\cot(c+dx)}}{d} \right) - \frac{2A \cot^{\frac{3}{2}}(c+dx) (a^2 \cot(c+dx) + ia^2)}{5d}$$

↓ 3042

$$\frac{1}{5} \left(- \int \frac{-10(A-iB)a^2 - 10(iA+B) \tan\left(c+dx+\frac{\pi}{2}\right) a^2}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}} dx - \frac{2a^2(5B+7iA) \cot^{\frac{3}{2}}(c+dx)}{3d} + \frac{20a^2(A-iB) \sqrt{\cot(c+dx)}}{d} \right) - \frac{2A \cot^{\frac{3}{2}}(c+dx) (a^2 \cot(c+dx) + ia^2)}{5d}$$

↓ 4016

$$\frac{1}{5} \left(- \frac{200a^4(A-iB)^2 \int \frac{1}{10(A-iB)a^2+10(iA+B) \cot(c+dx)a^2} d\sqrt{\cot(c+dx)}}{d} - \frac{2a^2(5B+7iA) \cot^{\frac{3}{2}}(c+dx)}{3d} + \frac{20a^2(A-iB) \sqrt{\cot(c+dx)}}{d} \right) - \frac{2A \cot^{\frac{3}{2}}(c+dx) (a^2 \cot(c+dx) + ia^2)}{5d}$$

3.508. $\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

↓ 221

$$\frac{1}{5} \left(\frac{20\sqrt[4]{-1}a^2(A - iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c + dx)}\right)}{d} - \frac{2a^2(5B + 7iA)\cot^{3/2}(c + dx)}{3d} + \frac{20a^2(A - iB)\sqrt{\cot(c + dx)}}{d} - \frac{2A\cot^{3/2}(c + dx)(a^2\cot(c + dx) + ia^2)}{5d} \right)$$

input `Int[Cot[c + d*x]^(7/2)*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(-2*A*Cot[c + d*x]^(3/2)*(I*a^2 + a^2*Cot[c + d*x]))/(5*d) + ((20*(-1)^(1/4)*a^2*(A - I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]])/d + (20*a^2*(A - I*B)*Sqrt[Cot[c + d*x]])/d - (2*a^2*((7*I)*A + 5*B)*Cot[c + d*x]^(3/2))/(3*d))/5`

3.508.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(m_.))*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp [g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(m_.))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4077 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(m_.))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.508.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 252 vs. 2(108) = 216.

Time = 0.72 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.98

method	result
derivativedivides	$a^2 \left(\frac{2A \cot(dx+c)^{\frac{5}{2}}}{5} + \frac{4iA \cot(dx+c)^{\frac{3}{2}}}{3} + \frac{2B \cot(dx+c)^{\frac{3}{2}}}{3} + 4iB \sqrt{\cot(dx+c)} - 4A \sqrt{\cot(dx+c)} + \frac{(-2iB+2A)\sqrt{2}}{\ln\left(\frac{1+\cot(dx+c)}{1-\cot(dx+c)}\right)} \right)$
default	$a^2 \left(\frac{2A \cot(dx+c)^{\frac{5}{2}}}{5} + \frac{4iA \cot(dx+c)^{\frac{3}{2}}}{3} + \frac{2B \cot(dx+c)^{\frac{3}{2}}}{3} + 4iB \sqrt{\cot(dx+c)} - 4A \sqrt{\cot(dx+c)} + \frac{(-2iB+2A)\sqrt{2}}{\ln\left(\frac{1+\cot(dx+c)}{1-\cot(dx+c)}\right)} \right)$

input `int(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

$$3.508. \int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

output
$$-a^2/d*(2/5*A*\cot(d*x+c)^(5/2)+4/3*I*A*\cot(d*x+c)^(3/2)+2/3*B*\cot(d*x+c)^(3/2)+4*I*B*\cot(d*x+c)^(1/2)-4*A*\cot(d*x+c)^(1/2)+1/4*(2*A-2*I*B)*2^(1/2)*(\ln((1+\cot(d*x+c)+2^(1/2)*\cot(d*x+c)^(1/2))/(1+\cot(d*x+c)-2^(1/2)*\cot(d*x+c)^(1/2))))+2*\arctan(1+2^(1/2)*\cot(d*x+c)^(1/2))+2*\arctan(-1+2^(1/2)*\cot(d*x+c)^(1/2)))+1/4*(-2*I*A-2*B)*2^(1/2)*(\ln((1+\cot(d*x+c)-2^(1/2)*\cot(d*x+c)^(1/2))/(1+\cot(d*x+c)+2^(1/2)*\cot(d*x+c)^(1/2))))+2*\arctan(1+2^(1/2)*\cot(d*x+c)^(1/2))+2*\arctan(-1+2^(1/2)*\cot(d*x+c)^(1/2))$$

3.508.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 448 vs. $2(104) = 208$.

Time = 0.26 (sec) , antiderivative size = 448, normalized size of antiderivative = 3.50

$$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx =$$

$$15 \sqrt{-\frac{(-iA^2-2AB+iB^2)a^4}{d^2}} (de^{(4i dx+4i c)} - 2de^{(2i dx+2i c)} + d) \log \left(\frac{2 \left((A-iB)a^2 e^{(2i dx+2i c)} - \sqrt{-\frac{(-iA^2-2AB+iB^2)a^4}{d^2}} \right)}{(-iA} \right.$$

input `integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output
$$-1/15*(15*\sqrt{-(-I*A^2-2*A*B+I*B^2)*a^4/d^2}*(d*e^{(4*I*d*x+4*I*c)}-2*d*e^{(2*I*d*x+2*I*c)}+d)*\log(2*((A-I*B)*a^2*e^{(2*I*d*x+2*I*c)}-\sqrt{-(-I*A^2-2*A*B+I*B^2)*a^4/d^2}*(I*d*e^{(2*I*d*x+2*I*c)}-I*d)*\sqrt{((I*e^{(2*I*d*x+2*I*c)}+I)/(e^{(2*I*d*x+2*I*c)}-1))}*e^{(-2*I*d*x-2*I*c)}/((-I*A-B)*a^2))-15*\sqrt{-(-I*A^2-2*A*B+I*B^2)*a^4/d^2}*(d*e^{(4*I*d*x+4*I*c)}-2*d*e^{(2*I*d*x+2*I*c)}+d)*\log(2*((A-I*B)*a^2*e^{(2*I*d*x+2*I*c)}-\sqrt{-(-I*A^2-2*A*B+I*B^2)*a^4/d^2}*(-I*d*e^{(2*I*d*x+2*I*c)}+I*d)*\sqrt{((I*e^{(2*I*d*x+2*I*c)}+I)/(e^{(2*I*d*x+2*I*c)}-1))}*e^{(-2*I*d*x-2*I*c)}/((-I*A-B)*a^2))-2*((43*A-35*I*B)*a^2*e^{(4*I*d*x+4*I*c)}-6*(9*A-10*I*B)*a^2*e^{(2*I*d*x+2*I*c)}+(23*A-25*I*B)*a^2)*\sqrt{((I*e^{(2*I*d*x+2*I*c)}+I)/(e^{(2*I*d*x+2*I*c)}-1))}/(d*e^{(4*I*d*x+4*I*c)}-2*d*e^{(2*I*d*x+2*I*c)}+d)$$

3.508. $\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.508.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(7/2)*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.508.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.55

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$\frac{15 \left(2\sqrt{2}(-i-1)A - (i+1)B \right) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}(-i-1)A - (i+1)B}{1}$$

input `integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/30*(15*(2*sqrt(2)*(-(I - 1)*A - (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I - 1)*A - (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a^2 - 120*(A - I*B)*a^2/sqrt(tan(d*x + c)) - 20*(-2*I*A - B)*a^2/tan(d*x + c)^(3/2) + 12*A*a^2/tan(d*x + c)^(5/2))/d`

3.508.8 Giac [F]

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^2 \cot(dx + c)^{\frac{7}{2}} dx$$

input `integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^2*cot(d*x + c)^(7/2), x)`

3.508.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{7/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) 1i)^2 dx$$

input `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2,x)`

output `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2, x)`

3.509 $\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.509.1 Optimal result	4795
3.509.2 Mathematica [A] (verified)	4795
3.509.3 Rubi [A] (verified)	4796
3.509.4 Maple [B] (verified)	4799
3.509.5 Fricas [B] (verification not implemented)	4800
3.509.6 Sympy [F(-1)]	4800
3.509.7 Maxima [B] (verification not implemented)	4801
3.509.8 Giac [F]	4801
3.509.9 Mupad [F(-1)]	4802

3.509.1 Optimal result

Integrand size = 36, antiderivative size = 103

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= -\frac{4\sqrt[4]{-1}a^2(iA+B)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} - \frac{2a^2(5iA+3B)\sqrt{\cot(c+dx)}}{3d} - \frac{2A\sqrt{\cot(c+dx)}(ia^2+a^2 \cot(c+dx))}{3d}$$

output `-4*(-1)^(1/4)*a^2*(I*A+B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d-2/3*a^2*(5*I*A+3*B)*cot(d*x+c)^(1/2)/d-2/3*A*(I*a^2+a^2*cot(d*x+c))*cot(d*x+c)^(1/2)/d`

3.509.2 Mathematica [A] (verified)

Time = 3.07 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.76

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx =$$

$$\frac{2a^2 \cot^{\frac{3}{2}}(c+dx) \left(A + 3(2iA + B) \tan(c+dx) - 6\sqrt[4]{-1}(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right) \right) \tan^{\frac{3}{2}}}{3d}$$

3.509. $\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

input `Integrate[Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(-2*a^2*Cot[c + d*x]^(3/2)*(A + 3*((2*I)*A + B)*Tan[c + d*x] - 6*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Tan[c + d*x]^(3/2))/(3*d)`

3.509.3 Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {3042, 4064, 3042, 4077, 27, 3042, 4075, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \cot(c + dx)^{5/2}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{(a \cot(c + dx) + ia)^2(A \cot(c + dx) + B)}{\sqrt{\cot(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-a \tan(c + dx + \frac{\pi}{2}) + ia)^2(B - A \tan(c + dx + \frac{\pi}{2}))}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4077} \\
 & -\frac{2}{3} \int \frac{(\cot(c + dx)a + ia)(a(A - 3iB) - a(5iA + 3B) \cot(c + dx))}{2\sqrt{\cot(c + dx)}} dx - \\
 & \quad \frac{2A\sqrt{\cot(c + dx)}(a^2 \cot(c + dx) + ia^2)}{3d} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.509. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& -\frac{1}{3} \int \frac{(\cot(c+dx)a+ia)(a(A-3iB)-a(5iA+3B)\cot(c+dx))}{\sqrt{\cot(c+dx)}} dx - \\
& \quad \frac{2A\sqrt{\cot(c+dx)}(a^2\cot(c+dx)+ia^2)}{3d} \\
& \quad \downarrow \text{3042} \\
& -\frac{1}{3} \int \frac{(ia-a\tan(c+dx+\frac{\pi}{2}))(a(A-3iB)+a(5iA+3B)\tan(c+dx+\frac{\pi}{2}))}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx - \\
& \quad \frac{2A\sqrt{\cot(c+dx)}(a^2\cot(c+dx)+ia^2)}{3d} \\
& \quad \downarrow \text{4075} \\
& \frac{1}{3} \left(-\int \frac{6(iA+B)a^2+6(A-iB)\cot(c+dx)a^2}{\sqrt{\cot(c+dx)}} dx - \frac{2a^2(3B+5iA)\sqrt{\cot(c+dx)}}{d} \right) - \\
& \quad \frac{2A\sqrt{\cot(c+dx)}(a^2\cot(c+dx)+ia^2)}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(-\int \frac{6a^2(iA+B)-6a^2(A-iB)\tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx - \frac{2a^2(3B+5iA)\sqrt{\cot(c+dx)}}{d} \right) - \\
& \quad \frac{2A\sqrt{\cot(c+dx)}(a^2\cot(c+dx)+ia^2)}{3d} \\
& \quad \downarrow \text{4016} \\
& \frac{1}{3} \left(-\frac{72a^4(B+iA)^2 \int \frac{1}{6a^2(A-iB)\cot(c+dx)-6a^2(iA+B)}}{d} d\sqrt{\cot(c+dx)} - \frac{2a^2(3B+5iA)\sqrt{\cot(c+dx)}}{d} \right) - \\
& \quad \frac{2A\sqrt{\cot(c+dx)}(a^2\cot(c+dx)+ia^2)}{3d} \\
& \quad \downarrow \text{221} \\
& \frac{1}{3} \left(-\frac{12\sqrt[4]{-1}a^2(B+iA)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} - \frac{2a^2(3B+5iA)\sqrt{\cot(c+dx)}}{d} \right) - \\
& \quad \frac{2A\sqrt{\cot(c+dx)}(a^2\cot(c+dx)+ia^2)}{3d}
\end{aligned}$$

input `Int[Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

```
output ((-12*(-1)^(1/4)*a^2*(I*A + B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]]]/d -
(2*a^2*((5*I)*A + 3*B)*Sqrt[Cot[c + d*x]]/d)/3 - (2*A*Sqrt[Cot[c + d*x]]
*(I*a^2 + a^2*Cot[c + d*x]))/(3*d)
```

3.509.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 221 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4016 Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_
)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b
*Tan[e + f*x]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]
```

```
rule 4064 Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(
x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp
[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c
*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !Integer
Q[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4075 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_)
+ (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B
*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*
x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c,
d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]
```

```
rule 4077 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Sim
p[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan
[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c -
a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] &&
GtQ[m, 1] && !LtQ[n, -1]
```

3.509.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 229 vs. 2(86) = 172.

Time = 0.70 (sec) , antiderivative size = 230, normalized size of antiderivative = 2.23

method	result
derivativedivides	$a^2 \left(\frac{2A \cot(dx+c)^{\frac{3}{2}}}{3} + 4iA \sqrt{\cot(dx+c)} + 2B \sqrt{\cot(dx+c)} + \frac{(-2iA-2B)\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)} \right) \right)}{4} \right)$
default	$a^2 \left(\frac{2A \cot(dx+c)^{\frac{3}{2}}}{3} + 4iA \sqrt{\cot(dx+c)} + 2B \sqrt{\cot(dx+c)} + \frac{(-2iA-2B)\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)} \right) \right)}{4} \right)$

```
input int(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETUR
NVERBOSE)
```

```
output -a^2/d*(2/3*A*cot(d*x+c)^(3/2)+4*I*A*cot(d*x+c)^(1/2)+2*B*cot(d*x+c)^(1/2)
+1/4*(-2*I*A-2*B)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+c
ot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+
2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))+1/4*(-2*A+2*I*B)*2^(1/2)*(ln((1+cot
(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))
+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)
)))
```

$$3.509. \int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

3.509.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 394 vs. $2(83) = 166$.

Time = 0.25 (sec) , antiderivative size = 394, normalized size of antiderivative = 3.83

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx =$$

$$3 \sqrt{-\frac{(iA^2+2AB-iB^2)a^4}{d^2}} (de^{(2i dx+2i c)} - d) \log \left(\frac{2 \left((A-iB)a^2 e^{(2i dx+2i c)} + \sqrt{-\frac{(iA^2+2AB-iB^2)a^4}{d^2}} (de^{(2i dx+2i c)} - d) \sqrt{\frac{ie^{(2i dx+2i c)}}{e^{(2i dx+2i c)}}}}{(iA-B)a^2} \right)}{(-iA-B)a^2} \right)$$

```
input integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output -1/3*(3*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)
*log(2*((A - I*B)*a^2*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*
a^4/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(
2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^2)) - 3*sqrt(-(
I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log(2*((A - I*
B)*a^2*e^(2*I*d*x + 2*I*c) - sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(d*e^(
2*I*d*x + 2*I*c) - d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c)
- 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^2)) + 2*((7*I*A + 3*B)*a^2*e^(
2*I*d*x + 2*I*c) + (-5*I*A - 3*B)*a^2)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e
^(2*I*d*x + 2*I*c) - 1)))/(d*e^(2*I*d*x + 2*I*c) - d)
```

3.509.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx = \text{Timed out}$$

```
input integrate(cot(d*x+c)**(5/2)*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)
```

```
output Timed out
```

3.509. $\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.509.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 180 vs. $2(83) = 166$.

Time = 0.30 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.75

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$\frac{3 \left(2\sqrt{2}(-(i+1)A + (i-1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}(-(i+1)A + (i-1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) - \sqrt{2}((i-1)A + (i+1)B) \log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + \frac{1}{\tan(dx+c)} + 1\right) + \sqrt{2}((i-1)A + (i+1)B) \log\left(-\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + \frac{1}{\tan(dx+c)} + 1\right) \right) a^2 - 12(-2iA - B)a^2/\sqrt{\tan(dx+c)} + 4Aa^2/\tan(dx+c)^{\frac{3}{2}}}{d}$$

input `integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/6*(3*(2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a^2 - 12*(-2*I*A - B)*a^2/sqrt(tan(d*x + c)) + 4*A*a^2/tan(d*x + c)^(3/2))/d`

3.509.8 Giac [F]

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^2 \cot(dx + c)^{\frac{5}{2}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^2*cot(d*x + c)^(5/2), x)`

3.509.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{5/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) i)^2 dx$$

input `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2,x)`output `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2, x)`

3.510 $\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.510.1 Optimal result	4803
3.510.2 Mathematica [A] (verified)	4803
3.510.3 Rubi [A] (verified)	4804
3.510.4 Maple [B] (verified)	4807
3.510.5 Fricas [B] (verification not implemented)	4807
3.510.6 Sympy [F]	4808
3.510.7 Maxima [B] (verification not implemented)	4809
3.510.8 Giac [F]	4809
3.510.9 Mupad [F(-1)]	4810

3.510.1 Optimal result

Integrand size = 36, antiderivative size = 99

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= -\frac{4\sqrt[4]{-1}a^2(A-iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} - \frac{2a^2(A+iB)\sqrt{\cot(c+dx)}}{d} + \frac{2iB(ia^2+a^2\cot(c+dx))}{d\sqrt{\cot(c+dx)}}$$

```
output -4*(-1)^(1/4)*a^2*(A-I*B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d+2*I*B*(I*a^2+a^2*cot(d*x+c))/d/cot(d*x+c)^(1/2)-2*a^2*(A+I*B)*cot(d*x+c)^(1/2)/d
```

3.510.2 Mathematica [A] (verified)

Time = 2.57 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.70

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx =$$

$$-\frac{2a^2\sqrt{\cot(c+dx)}\left(A+2\sqrt[4]{-1}(iA+B)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)\sqrt{\tan(c+dx)}+B \tan(c+dx)\right)}{d}$$

input `Integrate[Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(-2*a^2*Sqrt[Cot[c + d*x]]*(A + 2*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sqrt[Tan[c + d*x]] + B*Tan[c + d*x])/d`

3.510.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {3042, 4064, 3042, 4076, 27, 3042, 4075, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \cot(c + dx)^{3/2}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{(a \cot(c + dx) + ia)^2(A \cot(c + dx) + B)}{\cot^{\frac{3}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-a \tan(c + dx + \frac{\pi}{2}) + ia)^2(B - A \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{4076} \\
 & 2 \int \frac{(\cot(c + dx)a + ia)(a(iA + 3B) + a(A + iB) \cot(c + dx))}{2\sqrt{\cot(c + dx)}} dx + \frac{2iB(a^2 \cot(c + dx) + ia^2)}{d\sqrt{\cot(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(\cot(c + dx)a + ia)(a(iA + 3B) + a(A + iB) \cot(c + dx))}{\sqrt{\cot(c + dx)}} dx + \frac{2iB(a^2 \cot(c + dx) + ia^2)}{d\sqrt{\cot(c + dx)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.510. $\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \int \frac{(ia - a \tan(c + dx + \frac{\pi}{2})) (a(iA + 3B) - a(A + iB) \tan(c + dx + \frac{\pi}{2}))}{\sqrt{-\tan(c + dx + \frac{\pi}{2})} \frac{2iB(a^2 \cot(c + dx) + ia^2)}{d\sqrt{\cot(c + dx)}}} dx + \\
& \quad \downarrow \text{4075} \\
& \int \frac{2a^2(iA + B) \cot(c + dx) - 2a^2(A - iB)}{\sqrt{\cot(c + dx)}} dx - \frac{2a^2(A + iB)\sqrt{\cot(c + dx)}}{d} + \\
& \quad \frac{2iB(a^2 \cot(c + dx) + ia^2)}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow \text{3042} \\
& \int \frac{-2(A - iB)a^2 - 2(iA + B) \tan(c + dx + \frac{\pi}{2}) a^2}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx - \frac{2a^2(A + iB)\sqrt{\cot(c + dx)}}{d} + \\
& \quad \frac{2iB(a^2 \cot(c + dx) + ia^2)}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow \text{4016} \\
& \frac{8a^4(A - iB)^2}{d} \int \frac{1}{2(A - iB)a^2 + 2(iA + B) \cot(c + dx)a^2} d\sqrt{\cot(c + dx)} - \frac{2a^2(A + iB)\sqrt{\cot(c + dx)}}{d} + \\
& \quad \frac{2iB(a^2 \cot(c + dx) + ia^2)}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow \text{221} \\
& - \frac{4\sqrt[4]{-1}a^2(A - iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c + dx)}\right)}{d} - \frac{2a^2(A + iB)\sqrt{\cot(c + dx)}}{d} + \\
& \quad \frac{2iB(a^2 \cot(c + dx) + ia^2)}{d\sqrt{\cot(c + dx)}}
\end{aligned}$$

input `Int[Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(-4*(-1)^(1/4)*a^2*(A - I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]])/d - (2*a^2*(A + I*B)*Sqrt[Cot[c + d*x])/d + ((2*I)*B*(I*a^2 + a^2*Cot[c + d*x]))/(d*Sqrt[Cot[c + d*x]])`

3.510.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`
- rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])], x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`
- rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.510.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 217 vs. $2(85) = 170$.

Time = 0.39 (sec) , antiderivative size = 218, normalized size of antiderivative = 2.20

method	result
derivativedivides	$a^2 \left(2A\sqrt{\cot(dx+c)} + \frac{(2iB-2A)\sqrt{2} \left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right) + 2\arctan(1+\sqrt{2}\sqrt{\cot(dx+c)}) + 2\arctan(-1+\sqrt{2}\sqrt{\cot(dx+c)}) \right)}{4} \right)$
default	$a^2 \left(2A\sqrt{\cot(dx+c)} + \frac{(2iB-2A)\sqrt{2} \left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right) + 2\arctan(1+\sqrt{2}\sqrt{\cot(dx+c)}) + 2\arctan(-1+\sqrt{2}\sqrt{\cot(dx+c)}) \right)}{4} \right)$

input `int(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `-a^2/d*(2*A*cot(d*x+c)^(1/2)+1/4*(-2*A+2*I*B)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/4*(2*I*A+2*B)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+2*B/cot(d*x+c)^(1/2))`

3.510.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 385 vs. $2(81) = 162$.

Time = 0.25 (sec) , antiderivative size = 385, normalized size of antiderivative = 3.89

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{\sqrt{-\frac{(-iA^2-2AB+iB^2)a^4}{d^2}}(de^{(2i dx+2i c)}+d) \log\left(\frac{2\left((A-iB)a^2e^{(2i dx+2i c)}-\sqrt{-\frac{(-iA^2-2AB+iB^2)a^4}{d^2}}(ide^{(2i dx+2i c)}-i)d\right)\sqrt{\frac{ie^{(2i dx+2i c)}}{e^{(2i dx+2i c)}}}}{(-iA-B)a^2}\right)}{(-iA-B)a^2}$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,algorithm="fracas")`

3.510. $\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

output $(\sqrt{-(-IA^2 - 2AB + IB^2)}a^4/d^2)(de^{(2Id*x + 2I*c)} + d)\log(2 * ((A - IB)a^2e^{(2Id*x + 2I*c)} - \sqrt{-(-IA^2 - 2AB + IB^2)}a^4/d^2)(Id*e^{(2Id*x + 2I*c)} - Id)\sqrt{(Ie^{(2Id*x + 2I*c)} + I)/(e^{(2Id*x + 2I*c)} - 1)})e^{(-2Id*x - 2I*c)/((-IA - B)a^2)} - \sqrt{-(-IA^2 - 2AB + IB^2)}a^4/d^2)(de^{(2Id*x + 2I*c)} + d)\log(2 * ((A - IB)a^2e^{(2Id*x + 2I*c)} - \sqrt{-(-IA^2 - 2AB + IB^2)}a^4/d^2)(-Id*e^{(2Id*x + 2I*c)} + Id)\sqrt{(Ie^{(2Id*x + 2I*c)} + I)/(e^{(2Id*x + 2I*c)} - 1)})e^{(-2Id*x - 2I*c)/((-IA - B)a^2)} - 2 * ((A - IB)a^2e^{(2Id*x + 2I*c)} + (A + IB)a^2)\sqrt{(Ie^{(2Id*x + 2I*c)} + I)/(e^{(2Id*x + 2I*c)} - 1)})/(de^{(2Id*x + 2I*c)} + d)$

3.510.6 Sympy [F]

$$\begin{aligned} & \int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx \\ &= -a^2 \left(\int (-A \cot^{\frac{3}{2}}(c + dx)) dx + \int A \tan^2(c + dx) \cot^{\frac{3}{2}}(c + dx) dx \right. \\ & \quad + \int (-B \tan(c + dx) \cot^{\frac{3}{2}}(c + dx)) dx + \int B \tan^3(c + dx) \cot^{\frac{3}{2}}(c + dx) dx \\ & \quad \quad \quad + \int (-2iA \tan(c + dx) \cot^{\frac{3}{2}}(c + dx)) dx \\ & \quad \quad \quad \left. + \int (-2iB \tan^2(c + dx) \cot^{\frac{3}{2}}(c + dx)) dx \right) \end{aligned}$$

input `integrate(cot(d*x+c)**(3/2)*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `-a**2*(Integral(-A*cot(c + d*x)**(3/2), x) + Integral(A*tan(c + d*x)**2*cot(c + d*x)**(3/2), x) + Integral(-B*tan(c + d*x)*cot(c + d*x)**(3/2), x) + Integral(B*tan(c + d*x)**3*cot(c + d*x)**(3/2), x) + Integral(-2*I*A*tan(c + d*x)*cot(c + d*x)**(3/2), x) + Integral(-2*I*B*tan(c + d*x)**2*cot(c + d*x)**(3/2), x))`

3.510.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 174 vs. $2(81) = 162$.

Time = 0.32 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.76

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$\frac{4Ba^2\sqrt{\tan(dx+c)} - \left(2\sqrt{2}(-(i-1)A - (i+1)B)\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}(-(i-1)A - (i+1)B)\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) - \sqrt{2}(-(i-1)A - (i+1)B)\log\left(\frac{\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}}{\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}}\right) + 1/\tan(dx+c) + 1\right)a^2 + 4Aa^2/\sqrt{\tan(dx+c)}}{d}$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/2*(4*B*a^2*sqrt(tan(d*x + c)) - (2*sqrt(2)*(-(I - 1)*A - (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I - 1)*A - (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a^2 + 4*A*a^2/sqrt(tan(d*x + c)))/d`

3.510.8 Giac [F]

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^2 \cot(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^2*cot(d*x + c)^(3/2), x)`

3.510.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{3/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) i)^2 dx$$

input `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2,x)`output `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2, x)`

3.511 $\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.511.1 Optimal result	4811
3.511.2 Mathematica [A] (verified)	4811
3.511.3 Rubi [A] (verified)	4812
3.511.4 Maple [B] (verified)	4815
3.511.5 Fricas [B] (verification not implemented)	4815
3.511.6 Sympy [F]	4816
3.511.7 Maxima [B] (verification not implemented)	4817
3.511.8 Giac [F]	4817
3.511.9 Mupad [F(-1)]	4818

3.511.1 Optimal result

Integrand size = 36, antiderivative size = 105

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{4\sqrt[4]{-1}a^2(iA + B)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c + dx)}\right)}{d} - \frac{2a^2(3A - 5iB)}{3d\sqrt{\cot(c + dx)}} + \frac{2iB(ia^2 + a^2 \cot(c + dx))}{3d \cot^{3/2}(c + dx)}$$

```
output 4*(-1)^(1/4)*a^2*(I*A+B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d+2/3*I*B*(I
*a^2+a^2*cot(d*x+c))/d/cot(d*x+c)^(3/2)-2/3*a^2*(3*A-5*I*B)/d/cot(d*x+c)^(
1/2)
```

3.511.2 Mathematica [A] (verified)

Time = 4.51 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.86

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx =$$

$$\frac{2a^2\sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}\left(6\sqrt[4]{-1}(A - iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right) + \sqrt{\tan(c + dx)}(3A - 5iB)\right)}{3d}$$

input `Integrate[Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(-2*a^2*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(6*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + Sqrt[Tan[c + d*x]]*(3*A - (6*I)*B + B*Tan[c + d*x]))/(3*d)`

3.511.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.306$, Rules used = {3042, 4064, 3042, 4076, 27, 3042, 4074, 27, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{(a \cot(c+dx)+ia)^2(A \cot(c+dx)+B)}{\cot^{\frac{5}{2}}(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-a \tan(c+dx+\frac{\pi}{2})+ia)^2(B-A \tan(c+dx+\frac{\pi}{2}))}{(-\tan(c+dx+\frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{4076} \\
 & \frac{2}{3} \int \frac{(\cot(c+dx)a+ia)(a(3iA+5B)+a(3A-iB) \cot(c+dx))}{2 \cot^{\frac{3}{2}}(c+dx)} dx + \frac{2iB(a^2 \cot(c+dx)+ia^2)}{3d \cot^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{3} \int \frac{(\cot(c+dx)a+ia)(a(3iA+5B)+a(3A-iB) \cot(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx + \frac{2iB(a^2 \cot(c+dx)+ia^2)}{3d \cot^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.511. $\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \frac{1}{3} \int \frac{(ia - a \tan(c + dx + \frac{\pi}{2})) (a(3iA + 5B) - a(3A - iB) \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2} \frac{2iB(a^2 \cot(c + dx) + ia^2)}{3d \cot^{\frac{3}{2}}(c + dx)}} dx + \\
& \quad \downarrow \text{4074} \\
& \frac{1}{3} \left(\int \frac{6((iA + B)a^2 + (A - iB) \cot(c + dx)a^2)}{\sqrt{\cot(c + dx)}} dx - \frac{2a^2(3A - 5iB)}{d\sqrt{\cot(c + dx)}} \right) + \frac{2iB(a^2 \cot(c + dx) + ia^2)}{3d \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{27} \\
& \frac{1}{3} \left(6 \int \frac{(iA + B)a^2 + (A - iB) \cot(c + dx)a^2}{\sqrt{\cot(c + dx)}} dx - \frac{2a^2(3A - 5iB)}{d\sqrt{\cot(c + dx)}} \right) + \frac{2iB(a^2 \cot(c + dx) + ia^2)}{3d \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(6 \int \frac{a^2(iA + B) - a^2(A - iB) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx - \frac{2a^2(3A - 5iB)}{d\sqrt{\cot(c + dx)}} \right) + \\
& \quad \frac{2iB(a^2 \cot(c + dx) + ia^2)}{3d \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{4016} \\
& \frac{1}{3} \left(\frac{12a^4(B + iA)^2 \int \frac{1}{a^2(A - iB) \cot(c + dx) - a^2(iA + B)} d\sqrt{\cot(c + dx)}}{d} - \frac{2a^2(3A - 5iB)}{d\sqrt{\cot(c + dx)}} \right) + \\
& \quad \frac{2iB(a^2 \cot(c + dx) + ia^2)}{3d \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{221} \\
& \frac{1}{3} \left(\frac{12\sqrt[4]{-1}a^2(B + iA) \operatorname{arctanh}((-1)^{3/4} \sqrt{\cot(c + dx)})}{d} - \frac{2a^2(3A - 5iB)}{d\sqrt{\cot(c + dx)}} \right) + \\
& \quad \frac{2iB(a^2 \cot(c + dx) + ia^2)}{3d \cot^{\frac{3}{2}}(c + dx)}
\end{aligned}$$

input `Int[Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `((12*(-1)^(1/4)*a^2*(I*A + B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]])/d - (2*a^2*(3*A - (5*I)*B))/(d*Sqrt[Cot[c + d*x]])/3 + (((2*I)/3)*B*(I*a^2 + a^2*Cot[c + d*x]))/(d*Cot[c + d*x]^(3/2))`

3.511.3.1 Defintions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 221 $\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) * \text{ArcTanh}[x / \text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4016 $\text{Int}[(c_*) + (d_*) * \tan[(e_*) + (f_*)(x_)] / \text{Sqrt}[(b_*) * \tan[(e_*) + (f_*)(x_)]], x_Symbol] \rightarrow \text{Simp}[2 * (c^2/f) \text{ Subst}[\text{Int}[1/(b*c - d*x^2), x], x, \text{Sqrt}[b * \tan[e + f*x]]], x] /; \text{FreeQ}[\{b, c, d, e, f\}, x] \ \&\& \ \text{EqQ}[c^2 + d^2, 0]$
- rule 4064 $\text{Int}[(\cot[(e_*) + (f_*)(x_)] * (g_*)^p) * ((a_*) + (b_*) * \tan[(e_*) + (f_*)(x_)])^m * ((c_*) + (d_*) * \tan[(e_*) + (f_*)(x_)])^n, x_Symbol] \rightarrow \text{Simp}[g^{m+n} \text{ Int}[(g * \cot[e + f*x])^{p-m-n} * (b + a * \cot[e + f*x])^m * (d + c * \cot[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \ \&\& \ !\text{IntegerQ}[p] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{IntegerQ}[n]$
- rule 4074 $\text{Int}[(a_*) + (b_*) * \tan[(e_*) + (f_*)(x_)]^m * ((A_*) + (B_*) * \tan[(e_*) + (f_*)(x_)] * ((c_*) + (d_*) * \tan[(e_*) + (f_*)(x_)]), x_Symbol] \rightarrow \text{Simp}[(b * c - a * d) * (A * b - a * B) * (a + b * \tan[e + f*x])^{m+1} / (b * f * (m+1) * (a^2 + b^2)), x] + \text{Simp}[1 / (a^2 + b^2) \text{ Int}[(a + b * \tan[e + f*x])^{m+1} * \text{Simp}[a * A * c + b * B * c + A * b * d - a * B * d - (A * b * c - a * B * c - a * A * d - b * B * d) * \tan[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[b * c - a * d, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{NeQ}[a^2 + b^2, 0]$
- rule 4076 $\text{Int}[(a_*) + (b_*) * \tan[(e_*) + (f_*)(x_)]^m * ((A_*) + (B_*) * \tan[(e_*) + (f_*)(x_)] * ((c_*) + (d_*) * \tan[(e_*) + (f_*)(x_)])^n, x_Symbol] \rightarrow \text{Simp}[(-a^2) * (B * c - A * d) * (a + b * \tan[e + f*x])^{m-1} * ((c + d * \tan[e + f*x])^{n+1} / (d * f * (b * c + a * d) * (n+1))), x] - \text{Simp}[a / (d * (b * c + a * d) * (n+1)) \text{ Int}[(a + b * \tan[e + f*x])^{m-1} * (c + d * \tan[e + f*x])^{n+1} * \text{Simp}[A * b * d * (m - n - 2) - B * (b * c * (m - 1) + a * d * (n + 1)) + (a * A * d * (m + n) - B * (a * c * (m - 1) + b * d * (n + 1))] * \tan[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[b * c - a * d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{LtQ}[n, -1]$

3.511.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 224 vs. $2(87) = 174$.

Time = 0.37 (sec) , antiderivative size = 225, normalized size of antiderivative = 2.14

method	result
derivativedivides	$a^2 \left(-\frac{2(2iB-A)}{\sqrt{\cot(dx+c)}} + \frac{2B}{3 \cot(dx+c)^{\frac{3}{2}}} + \frac{(2iA+2B)\sqrt{2}}{4} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)} \right) + 2 \arctan \left(\frac{1-\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)} \right) \right) \right)$
default	$a^2 \left(-\frac{2(2iB-A)}{\sqrt{\cot(dx+c)}} + \frac{2B}{3 \cot(dx+c)^{\frac{3}{2}}} + \frac{(2iA+2B)\sqrt{2}}{4} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)} \right) + 2 \arctan \left(\frac{1-\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)} \right) \right) \right)$

input `int(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `-a^2/d*(-2*(2*I*B-A)/cot(d*x+c)^(1/2)+2/3*B/cot(d*x+c)^(3/2)+1/4*(2*I*A+2*B)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/4*(2*A-2*I*B)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))`

3.511.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 439 vs. $2(83) = 166$.

Time = 0.25 (sec) , antiderivative size = 439, normalized size of antiderivative = 4.18

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{3 \sqrt{-\frac{(iA^2+2AB-iB^2)a^4}{d^2}} (de^{(4i dx+4i c)} + 2 de^{(2i dx+2i c)} + d) \log \left(\frac{2 \left((A-iB)a^2 e^{(2i dx+2i c)} + \sqrt{-\frac{(iA^2+2AB-iB^2)a^4}{d^2}} (de^{(2i dx+2i c)} + d) \right)}{(-iA-B)a^2} \right)}{1}$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fracas")`

3.511. $\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$

output `1/3*(3*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log(2*((A - I*B)*a^2*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^2)) - 3*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log(2*((A - I*B)*a^2*e^(2*I*d*x + 2*I*c) - sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^4/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^2)) - 2*((-3*I*A - 7*B)*a^2*e^(4*I*d*x + 4*I*c) + 2*B*a^2*e^(2*I*d*x + 2*I*c) + (3*I*A + 5*B)*a^2)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))/(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)`

3.511.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx \\ &= -a^2 \left(\int \left(-A\sqrt{\cot(c+dx)} \right) dx + \int A \tan^2(c+dx)\sqrt{\cot(c+dx)} dx \right. \\ & \quad \left. + \int \left(-B \tan(c+dx)\sqrt{\cot(c+dx)} \right) dx + \int B \tan^3(c+dx)\sqrt{\cot(c+dx)} dx \right. \\ & \quad \left. + \int \left(-2iA \tan(c+dx)\sqrt{\cot(c+dx)} \right) dx \right. \\ & \quad \left. + \int \left(-2iB \tan^2(c+dx)\sqrt{\cot(c+dx)} \right) dx \right) \end{aligned}$$

input `integrate(cot(d*x+c)**(1/2)*(a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `-a**2*(Integral(-A*sqrt(cot(c + d*x)), x) + Integral(A*tan(c + d*x)**2*sqrt(cot(c + d*x)), x) + Integral(-B*tan(c + d*x)*sqrt(cot(c + d*x)), x) + Integral(B*tan(c + d*x)**3*sqrt(cot(c + d*x)), x) + Integral(-2*I*A*tan(c + d*x)*sqrt(cot(c + d*x)), x) + Integral(-2*I*B*tan(c + d*x)**2*sqrt(cot(c + d*x)), x))`

3.511.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 180 vs. $2(83) = 166$.

Time = 0.30 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.71

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{3 \left(2\sqrt{2}(-(i+1)A+(i-1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}+\frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}(-(i+1)A+(i-1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}-\frac{2}{\sqrt{\tan(dx+c)}}\right)\right) - \sqrt{2}((i-1)A+(i+1)B) \log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}}+1\right) + \sqrt{2}((i-1)A+(i+1)B) \log\left(-\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}}+1\right) a^2 - 4(Ba^2+3(A-2iB)a^2/\tan(dx+c))\tan(dx+c)^{3/2} \right)}{d}$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/6*(3*(2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a^2 - 4*(B*a^2 + 3*(A - 2*I*B)*a^2/tan(d*x + c))*tan(d*x + c)^(3/2)/d`

3.511.8 Giac [F]

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \int (B \tan(dx+c) + A)(ia \tan(dx+c) + a)^2 \sqrt{\cot(dx+c)} dx$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^2*sqrt(cot(d*x + c)), x)`

3.511.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int \sqrt{\cot(c + dx)}(A + B \tan(c + dx)) (a + a \tan(c + dx) i)^2 dx$$

input `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*i)^2,x)`output `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*i)^2, x)`

$$3.512 \quad \int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.512.1 Optimal result	4819
3.512.2 Mathematica [A] (verified)	4819
3.512.3 Rubi [A] (verified)	4820
3.512.4 Maple [B] (verified)	4823
3.512.5 Fricas [B] (verification not implemented)	4824
3.512.6 Sympy [F]	4825
3.512.7 Maxima [A] (verification not implemented)	4826
3.512.8 Giac [F]	4826
3.512.9 Mupad [F(-1)]	4827

3.512.1 Optimal result

Integrand size = 36, antiderivative size = 130

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\ &= \frac{4\sqrt{-1}a^2(A - iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c + dx)}\right)}{d} \\ & \quad - \frac{2a^2(5A - 7iB)}{15d \cot^{3/2}(c + dx)} + \frac{4a^2(iA + B)}{d\sqrt{\cot(c + dx)}} + \frac{2iB(ia^2 + a^2 \cot(c + dx))}{5d \cot^{5/2}(c + dx)} \end{aligned}$$

output `4*(-1)^(1/4)*a^2*(A-I*B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d-2/15*a^2*(5*A-7*I*B)/d/cot(d*x+c)^(3/2)+2/5*I*B*(I*a^2+a^2*cot(d*x+c))/d/cot(d*x+c)^(5/2)+4*a^2*(I*A+B)/d/cot(d*x+c)^(1/2)`

3.512.2 Mathematica [A] (verified)

Time = 4.87 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.75

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\ &= \frac{2a^2 \left(30(iA + B) + \frac{30\sqrt{-1}(iA+B)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)}{\sqrt{\tan(c+dx)}} - 5(A - 2iB)\tan(c + dx) - 3B \tan^2(c + dx) \right)}{15d\sqrt{\cot(c + dx)}} \end{aligned}$$

input `Integrate[((a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `(2*a^2*(30*(I*A + B) + (30*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]])/Sqrt[Tan[c + d*x]] - 5*(A - (2*I)*B)*Tan[c + d*x] - 3*B*Tan[c + d*x]^2)/(15*d*Sqrt[Cot[c + d*x]])`

3.512.3 Rubi [A] (verified)

Time = 0.90 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.06, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.361$, Rules used = {3042, 4064, 3042, 4076, 27, 3042, 4074, 27, 3042, 4012, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{(a \cot(c + dx) + ia)^2 (A \cot(c + dx) + B)}{\cot^{7/2}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-a \tan(c + dx + \frac{\pi}{2}) + ia)^2 (B - A \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{7/2}} dx \\
 & \quad \downarrow \text{4076} \\
 & \frac{2}{5} \int \frac{(\cot(c + dx)a + ia)(a(5iA + 7B) + a(5A - 3iB) \cot(c + dx))}{2 \cot^{5/2}(c + dx)} dx + \frac{2iB(a^2 \cot(c + dx) + ia^2)}{5d \cot^{5/2}(c + dx)} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{5} \int \frac{(\cot(c + dx)a + ia)(a(5iA + 7B) + a(5A - 3iB) \cot(c + dx))}{\cot^{5/2}(c + dx)} dx + \frac{2iB(a^2 \cot(c + dx) + ia^2)}{5d \cot^{5/2}(c + dx)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.512. $\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$

$$\begin{aligned}
& \frac{1}{5} \int \frac{(ia - a \tan(c + dx + \frac{\pi}{2})) (a(5iA + 7B) - a(5A - 3iB) \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{5/2} \frac{2iB(a^2 \cot(c + dx) + ia^2)}{5d \cot^{\frac{5}{2}}(c + dx)}} dx + \\
& \quad \downarrow \text{4074} \\
& \frac{1}{5} \left(\int \frac{10((iA + B)a^2 + (A - iB) \cot(c + dx)a^2)}{\cot^{\frac{3}{2}}(c + dx) \frac{2iB(a^2 \cot(c + dx) + ia^2)}{5d \cot^{\frac{5}{2}}(c + dx)}} dx - \frac{2a^2(5A - 7iB)}{3d \cot^{\frac{3}{2}}(c + dx)} \right) + \\
& \quad \downarrow \text{27} \\
& \frac{1}{5} \left(10 \int \frac{(iA + B)a^2 + (A - iB) \cot(c + dx)a^2}{\cot^{\frac{3}{2}}(c + dx)} dx - \frac{2a^2(5A - 7iB)}{3d \cot^{\frac{3}{2}}(c + dx)} \right) + \frac{2iB(a^2 \cot(c + dx) + ia^2)}{5d \cot^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \left(10 \int \frac{a^2(iA + B) - a^2(A - iB) \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2} \frac{2iB(a^2 \cot(c + dx) + ia^2)}{5d \cot^{\frac{5}{2}}(c + dx)}} dx - \frac{2a^2(5A - 7iB)}{3d \cot^{\frac{3}{2}}(c + dx)} \right) + \\
& \quad \downarrow \text{4012} \\
& \frac{1}{5} \left(10 \left(\int \frac{a^2(A - iB) - a^2(iA + B) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx + \frac{2a^2(B + iA)}{d\sqrt{\cot(c + dx)}} \right) - \frac{2a^2(5A - 7iB)}{3d \cot^{\frac{3}{2}}(c + dx)} \right) + \\
& \quad \frac{2iB(a^2 \cot(c + dx) + ia^2)}{5d \cot^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{5} \left(10 \left(\int \frac{(A - iB)a^2 + (iA + B) \tan(c + dx + \frac{\pi}{2}) a^2}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx + \frac{2a^2(B + iA)}{d\sqrt{\cot(c + dx)}} \right) - \frac{2a^2(5A - 7iB)}{3d \cot^{\frac{3}{2}}(c + dx)} \right) + \\
& \quad \frac{2iB(a^2 \cot(c + dx) + ia^2)}{5d \cot^{\frac{5}{2}}(c + dx)} \\
& \quad \downarrow \text{4016} \\
& \frac{1}{5} \left(10 \left(\frac{2a^4(A - iB)^2 \int \frac{1}{-((A - iB)a^2 - (iA + B) \cot(c + dx)a^2)} d\sqrt{\cot(c + dx)}}{d} + \frac{2a^2(B + iA)}{d\sqrt{\cot(c + dx)}} \right) - \frac{2a^2(5A - 7iB)}{3d \cot^{\frac{3}{2}}(c + dx)} \right) + \\
& \quad \frac{2iB(a^2 \cot(c + dx) + ia^2)}{5d \cot^{\frac{5}{2}}(c + dx)}
\end{aligned}$$

3.512. $\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$

↓ 221

$$\frac{1}{5} \left(10 \left(\frac{2\sqrt[4]{-1}a^2(A - iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c + dx)}\right)}{d} + \frac{2a^2(B + iA)}{d\sqrt{\cot(c + dx)}} \right) - \frac{2a^2(5A - 7iB)}{3d\cot^{3/2}(c + dx)} \right) + \frac{2iB(a^2\cot(c + dx) + ia^2)}{5d\cot^{5/2}(c + dx)}$$

input `Int[((a + I*a*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `(10*((2*(-1)^(1/4)*a^2*(A - I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]])/d + (2*a^2*(I*A + B))/(d*Sqrt[Cot[c + d*x]])) - (2*a^2*(5*A - (7*I)*B))/(3*d*Cot[c + d*x]^(3/2)))/5 + (((2*I)/5)*B*(I*a^2 + a^2*Cot[c + d*x]))/(d*Cot[c + d*x]^(5/2))`

3.512.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

3.512. $\int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

```
rule 4064 Int[(cot[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp [g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]
```

```
rule 4076 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

3.512.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 242 vs. 2(109) = 218.

Time = 0.38 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.87

method	result
derivativedivides	$a^2 \left(-\frac{2(2iA+2B)}{\sqrt{\cot(dx+c)}} - \frac{2(2iB-A)}{3 \cot(dx+c)^{\frac{3}{2}}} + \frac{2B}{5 \cot(dx+c)^{\frac{5}{2}}} + \frac{(-2iB+2A)\sqrt{2}}{4} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)} \right) \right) \right)$
default	$a^2 \left(-\frac{2(2iA+2B)}{\sqrt{\cot(dx+c)}} - \frac{2(2iB-A)}{3 \cot(dx+c)^{\frac{3}{2}}} + \frac{2B}{5 \cot(dx+c)^{\frac{5}{2}}} + \frac{(-2iB+2A)\sqrt{2}}{4} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)} \right) \right) \right)$

$$3.512. \int \frac{(a+ia \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

input `int((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x,method=_RETURNVERBOSE)`

output `-a^2/d*(-2*(2*I*A+2*B)/cot(d*x+c)^(1/2)-2/3*(2*I*B-A)/cot(d*x+c)^(3/2)+2/5*B/cot(d*x+c)^(5/2)+1/4*(2*A-2*I*B)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/4*(-2*I*A-2*B)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))`

3.512.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 505 vs. $2(104) = 208$.

Time = 0.26 (sec) , antiderivative size = 505, normalized size of antiderivative = 3.88

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx =$$

$$15 \sqrt{-\frac{(-i A^2 - 2AB + i B^2)a^4}{d^2}} (de^{(6i dx + 6i c)} + 3de^{(4i dx + 4i c)} + 3de^{(2i dx + 2i c)} + d) \log \left(\frac{2 \left((A - i B)a^2 e^{(2i dx + 2i c)} - \sqrt{-\frac{(-i A^2 - 2AB + i B^2)a^4}{d^2}} \right)}{\dots} \right)$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="fracas")`

output
$$\begin{aligned} & -1/15*(15*\sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2}*(d*e^{(6*I*d*x + 6*I*c)} + \\ & 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} + d)*\log(2*((A - I*B)*a \\ & ^2*e^{(2*I*d*x + 2*I*c)} - \sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2}*(I*d*e^{(2 \\ & *I*d*x + 2*I*c)} - I*d)*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I \\ & c)} - 1))}*e^{(-2*I*d*x - 2*I*c)}/((-I*A - B)*a^2)) - 15*\sqrt{-(-I*A^2 - 2*A* \\ & B + I*B^2)*a^4/d^2}*(d*e^{(6*I*d*x + 6*I*c)} + 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d \\ & *e^{(2*I*d*x + 2*I*c)} + d)*\log(2*((A - I*B)*a^2*e^{(2*I*d*x + 2*I*c)} - \sqrt{ \\ & -(-I*A^2 - 2*A*B + I*B^2)*a^4/d^2}*(-I*d*e^{(2*I*d*x + 2*I*c)} + I*d)*\sqrt{((\\ & I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))}*e^{(-2*I*d*x - 2*I*c} \\ &))/((-I*A - B)*a^2)) - 2*((35*A - 43*I*B)*a^2*e^{(6*I*d*x + 6*I*c)} + (25*A - \\ & 11*I*B)*a^2*e^{(4*I*d*x + 4*I*c)} - (35*A - 31*I*B)*a^2*e^{(2*I*d*x + 2*I*c)} \\ & - (25*A - 23*I*B)*a^2)*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I \\ & *c)} - 1))}/(d*e^{(6*I*d*x + 6*I*c)} + 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d \\ & *x + 2*I*c)} + d) \end{aligned}$$

3.512.6 Sympy [F]

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^2(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\ & = -a^2 \left(\int \left(-\frac{A}{\sqrt{\cot(c + dx)}} \right) dx + \int \frac{A \tan^2(c + dx)}{\sqrt{\cot(c + dx)}} dx + \int \left(-\frac{B \tan(c + dx)}{\sqrt{\cot(c + dx)}} \right) dx \right. \\ & \quad \left. + \int \frac{B \tan^3(c + dx)}{\sqrt{\cot(c + dx)}} dx + \int \left(-\frac{2iA \tan(c + dx)}{\sqrt{\cot(c + dx)}} \right) dx + \int \left(-\frac{2iB \tan^2(c + dx)}{\sqrt{\cot(c + dx)}} \right) dx \right) \end{aligned}$$

input `integrate((a+I*a*tan(d*x+c))**2*(A+B*tan(d*x+c))/cot(d*x+c)**(1/2),x)`

output
$$\begin{aligned} & -a**2*(Integral(-A/\sqrt{\cot(c + d*x)}, x) + Integral(A*tan(c + d*x)**2/\sqrt{ \\ & \cot(c + d*x)}, x) + Integral(-B*tan(c + d*x)/\sqrt{\cot(c + d*x)}, x) + In \\ & tegral(B*tan(c + d*x)**3/\sqrt{\cot(c + d*x)}, x) + Integral(-2*I*A*tan(c + \\ & d*x)/\sqrt{\cot(c + d*x)}, x) + Integral(-2*I*B*tan(c + d*x)**2/\sqrt{\cot(c + \\ & d*x)}, x)) \end{aligned}$$

3.512.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.53

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx =$$

$$\frac{4 \left(3 B a^2 + \frac{5(A - 2iB)a^2}{\tan(dx+c)} - \frac{30(iA+B)a^2}{\tan(dx+c)^2} \right) \tan(dx+c)^{\frac{5}{2}} + 15 \left(2\sqrt{2}(-(i-1)A - (i+1)B) \arctan\left(\frac{1}{2}\sqrt{2}\right) \right)}{d}$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="maxima")`

output `-1/30*(4*(3*B*a^2 + 5*(A - 2*I*B)*a^2/tan(d*x + c) - 30*(I*A + B)*a^2/tan(d*x + c)^2)*tan(d*x + c)^(5/2) + 15*(2*sqrt(2)*(-(I - 1)*A - (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I - 1)*A - (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a^2)/d`

3.512.8 Giac [F]

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^2}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^2*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^2/sqrt(cot(d*x + c)), x)`

3.512.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) 1i)^2}{\sqrt{\cot(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2)/cot(c + d*x)^(1/2),x)`output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^2)/cot(c + d*x)^(1/2), x)`

3.513 $\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.513.1 Optimal result	4828
3.513.2 Mathematica [C] (verified)	4829
3.513.3 Rubi [A] (verified)	4829
3.513.4 Maple [A] (verified)	4833
3.513.5 Fricas [B] (verification not implemented)	4834
3.513.6 Sympy [F(-1)]	4834
3.513.7 Maxima [A] (verification not implemented)	4835
3.513.8 Giac [F]	4835
3.513.9 Mupad [F(-1)]	4836

3.513.1 Optimal result

Integrand size = 36, antiderivative size = 171

$$\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{8\sqrt[4]{-1}a^3(iA+B)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} + \frac{8a^3(iA+B)\sqrt{\cot(c+dx)}}{d}$$

$$+ \frac{8a^3(23A-21iB)\cot^{\frac{3}{2}}(c+dx)}{105d} - \frac{2aA\cot^{\frac{3}{2}}(c+dx)(ia+a\cot(c+dx))^2}{7d}$$

$$- \frac{2(11iA+7B)\cot^{\frac{3}{2}}(c+dx)(ia^3+a^3\cot(c+dx))}{35d}$$

```
output 8*(-1)^(1/4)*a^3*(I*A+B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d+8/105*a^3*
(23*A-21*I*B)*cot(d*x+c)^(3/2)/d-2/7*a*A*cot(d*x+c)^(3/2)*(I*a+a*cot(d*x+c)
)^2/d-2/35*(11*I*A+7*B)*cot(d*x+c)^(3/2)*(I*a^3+a^3*cot(d*x+c))/d+8*a^3*(
I*A+B)*cot(d*x+c)^(1/2)/d
```

3.513.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.93 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.66

$$\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{2a^3 \cot^{\frac{3}{2}}(c+dx) (35iB + 21iA \cot(c+dx) + 63B \cot(c+dx) + 45A \cot^2(c+dx) - 60A \cot^2(c+dx) \operatorname{Hy}}{}$$

input `Integrate[Cot[c + d*x]^(9/2)*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(2*a^3*Cot[c + d*x]^(3/2)*((35*I)*B + (21*I)*A*Cot[c + d*x] + 63*B*Cot[c + d*x] + 45*A*Cot[c + d*x]^2 - 60*A*Cot[c + d*x]^2*Hypergeometric2F1[-7/2, 1, -5/2, I*Tan[c + d*x]] - 84*B*Cot[c + d*x]*Hypergeometric2F1[-5/2, 1, -3/2, I*Tan[c + d*x]]))/(105*d)`

3.513.3 Rubi [A] (verified)

Time = 1.15 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.06, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3042, 4064, 3042, 4077, 27, 3042, 4077, 27, 3042, 4075, 3042, 4011, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{9/2}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4064}$$

$$\int \sqrt{\cot(c+dx)}(a \cot(c+dx) + ia)^3(A \cot(c+dx) + B) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(-a \tan\left(c+dx+\frac{\pi}{2}\right) + ia\right)^3\left(B - A \tan\left(c+dx+\frac{\pi}{2}\right)\right) dx$$

3.513. $\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \downarrow 4077 \\
& -\frac{2}{7} \int \frac{1}{2} \sqrt{\cot(c+dx)} (\cot(c+dx)a+ia)^2 (a(3A-7iB) - a(11iA+7B)\cot(c+dx)) dx - \\
& \quad \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + ia)^2}{7d} \\
& \quad \downarrow 27 \\
& -\frac{1}{7} \int \sqrt{\cot(c+dx)} (\cot(c+dx)a+ia)^2 (a(3A-7iB) - a(11iA+7B)\cot(c+dx)) dx - \\
& \quad \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + ia)^2}{7d} \\
& \quad \downarrow 3042 \\
& -\frac{1}{7} \int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)} \left(ia - a \tan\left(c+dx+\frac{\pi}{2}\right)\right)^2 \left(a(3A-7iB) + a(11iA+7B)\tan\left(c+dx+\frac{\pi}{2}\right)\right) dx - \\
& \quad \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + ia)^2}{7d} \\
& \quad \downarrow 4077 \\
& \frac{1}{7} \left(\frac{2}{5} \int -2\sqrt{\cot(c+dx)} (\cot(c+dx)a+ia) (2(6iA+7B)a^2 + (23A-21iB)\cot(c+dx)a^2) dx - \frac{2(7B+11iA)}{5} \cot(c+dx) \right) \\
& \quad \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + ia)^2}{7d} \\
& \quad \downarrow 27 \\
& \frac{1}{7} \left(-\frac{4}{5} \int \sqrt{\cot(c+dx)} (\cot(c+dx)a+ia) (2(6iA+7B)a^2 + (23A-21iB)\cot(c+dx)a^2) dx - \frac{2(7B+11iA)}{5} \cot(c+dx) \right) \\
& \quad \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + ia)^2}{7d} \\
& \quad \downarrow 3042 \\
& \frac{1}{7} \left(-\frac{4}{5} \int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)} \left(ia - a \tan\left(c+dx+\frac{\pi}{2}\right)\right) \left(2a^2(6iA+7B) - a^2(23A-21iB)\tan\left(c+dx+\frac{\pi}{2}\right)\right) dx - \right. \\
& \quad \left. \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + ia)^2}{7d} \right) \\
& \quad \downarrow 4075
\end{aligned}$$

3.513. $\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\frac{1}{7} \left(-\frac{4}{5} \left(\int \sqrt{\cot(c+dx)} (35a^3(iA+B) \cot(c+dx) - 35a^3(A-iB)) dx - \frac{2a^3(23A-21iB) \cot^{\frac{3}{2}}(c+dx)}{3d} \right) - \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + ia)^2}{7d} \right)$$

↓ 3042

$$\frac{1}{7} \left(-\frac{4}{5} \left(\int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)} \left(-35(A-iB)a^3 - 35(iA+B) \tan\left(c+dx+\frac{\pi}{2}\right) a^3 \right) dx - \frac{2a^3(23A-21iB) \cot^{\frac{3}{2}}(c+dx)}{3d} \right) - \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + ia)^2}{7d} \right)$$

↓ 4011

$$\frac{1}{7} \left(-\frac{4}{5} \left(\int \frac{-35(iA+B)a^3 - 35(A-iB) \cot(c+dx)a^3}{\sqrt{\cot(c+dx)}} dx - \frac{2a^3(23A-21iB) \cot^{\frac{3}{2}}(c+dx)}{3d} - \frac{70a^3(B+iA)\sqrt{\cot(c+dx)}}{d} \right) - \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + ia)^2}{7d} \right)$$

↓ 3042

$$\frac{1}{7} \left(-\frac{4}{5} \left(\int \frac{35a^3(A-iB) \tan\left(c+dx+\frac{\pi}{2}\right) - 35a^3(iA+B)}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}} dx - \frac{2a^3(23A-21iB) \cot^{\frac{3}{2}}(c+dx)}{3d} - \frac{70a^3(B+iA)\sqrt{\cot(c+dx)}}{d} \right) - \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + ia)^2}{7d} \right)$$

↓ 4016

$$\frac{1}{7} \left(-\frac{4}{5} \left(\frac{2450a^6(B+iA)^2 \int \frac{1}{35a^3(iA+B) - 35a^3(A-iB) \cot(c+dx)} d\sqrt{\cot(c+dx)}}{d} - \frac{2a^3(23A-21iB) \cot^{\frac{3}{2}}(c+dx)}{3d} - \frac{70a^3(B+iA)\sqrt{\cot(c+dx)}}{d} \right) - \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + ia)^2}{7d} \right)$$

↓ 221

$$\frac{1}{7} \left(-\frac{4}{5} \left(-\frac{70\sqrt[4]{-1}a^3(B+iA) \operatorname{arctanh}\left((-1)^{3/4} \sqrt{\cot(c+dx)}\right)}{d} - \frac{2a^3(23A-21iB) \cot^{\frac{3}{2}}(c+dx)}{3d} - \frac{70a^3(B+iA)\sqrt{\cot(c+dx)}}{d} \right) - \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + ia)^2}{7d} \right)$$

3.513. $\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

input `Int[Cot[c + d*x]^(9/2)*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(-2*a*A*Cot[c + d*x]^(3/2)*(I*a + a*Cot[c + d*x])^2)/(7*d) + ((-2*((11*I)*A + 7*B)*Cot[c + d*x]^(3/2)*(I*a^3 + a^3*Cot[c + d*x]))/(5*d) - (4*((-70*(-1)^(1/4)*a^3*(I*A + B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]]])/d - (70*a^3*(I*A + B)*Sqrt[Cot[c + d*x]])/d - (2*a^3*(23*A - (21*I)*B)*Cot[c + d*x]^(3/2))/(3*d))/5)/7`

3.513.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4016 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)])*(g_.)^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4077 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.513.4 Maple [A] (verified)

Time = 1.31 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.61

method	result
derivativedivides	$a^3 \left(-\frac{2A \cot(dx+c)^{\frac{7}{2}}}{7} - \frac{6iA \cot(dx+c)^{\frac{5}{2}}}{5} - \frac{2B \cot(dx+c)^{\frac{5}{2}}}{5} - 2iB \cot(dx+c)^{\frac{3}{2}} + \frac{8A \cot(dx+c)^{\frac{3}{2}}}{3} + 8iA \sqrt{\cot(dx+c)} + 8B \sqrt{\cot(dx+c)} \right)$
default	$a^3 \left(-\frac{2A \cot(dx+c)^{\frac{7}{2}}}{7} - \frac{6iA \cot(dx+c)^{\frac{5}{2}}}{5} - \frac{2B \cot(dx+c)^{\frac{5}{2}}}{5} - 2iB \cot(dx+c)^{\frac{3}{2}} + \frac{8A \cot(dx+c)^{\frac{3}{2}}}{3} + 8iA \sqrt{\cot(dx+c)} + 8B \sqrt{\cot(dx+c)} \right)$

input `int(cot(d*x+c)^(9/2)*(a+i*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `a^3/d*(-2/7*A*cot(d*x+c)^(7/2)-6/5*I*A*cot(d*x+c)^(5/2)-2/5*B*cot(d*x+c)^(5/2)-2*I*B*cot(d*x+c)^(3/2)+8/3*A*cot(d*x+c)^(3/2)+8*I*A*cot(d*x+c)^(1/2)+8*B*cot(d*x+c)^(1/2)-1/4*(4*I*A+4*B)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))-1/4*(4*A-4*I*B)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))`

$$3.513. \int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

3.513.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 508 vs. $2(139) = 278$.

Time = 0.28 (sec) , antiderivative size = 508, normalized size of antiderivative = 2.97

$$\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{2 \left(105 \sqrt{-\frac{(iA^2+2AB-iB^2)a^6}{d^2}} (de^{(6i dx+6i c)} - 3de^{(4i dx+4i c)} + 3de^{(2i dx+2i c)} - d) \log \left(\frac{2 \left((A-iB)a^3 e^{(2i dx+2i c)} + \sqrt{\dots} \right)}{\dots} \right) \right)}{\dots}$$

input `integrate(cot(d*x+c)^(9/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `2/105*(105*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*log(2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3)) - 105*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*log(2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) - sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3)) - 2*((-319*I*A - 273*B)*a^3*e^(6*I*d*x + 6*I*c) + 2*(323*I*A + 336*B)*a^3*e^(4*I*d*x + 4*I*c) + (-551*I*A - 567*B)*a^3*e^(2*I*d*x + 2*I*c) + 4*(41*I*A + 42*B)*a^3)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))/(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)`

3.513.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(9/2)*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

3.513. $\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

output Timed out

3.513.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.27

$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{105 \left(2\sqrt{2}(-(i+1)A + (i-1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}(-(i+1)A + (i-1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) - \sqrt{2}((i-1)A + (i+1)B) \log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + \frac{1}{\tan(dx+c)} + 1\right) + \sqrt{2}((i-1)A + (i+1)B) \log\left(-\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + \frac{1}{\tan(dx+c)} + 1\right) \right) a^3 + 840(IA + B)a^3/\sqrt{\tan(dx+c)} + 70(4A - 3IB)a^3/\tan(dx+c)^{\frac{3}{2}} + 42(-3IA - B)a^3/\tan(dx+c)^{\frac{5}{2}} - 30Aa^3/\tan(dx+c)^{\frac{7}{2}}}{d}$$

input `integrate(cot(d*x+c)^(9/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/105*(105*(2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a^3 + 840*(I*A + B)*a^3/sqrt(tan(d*x + c)) + 70*(4*A - 3*I*B)*a^3/tan(d*x + c)^(3/2) + 42*(-3*I*A - B)*a^3/tan(d*x + c)^(5/2) - 30*A*a^3/tan(d*x + c)^(7/2))/d`

3.513.8 Giac [F]

$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^3 \cot(dx + c)^{\frac{9}{2}} dx$$

input `integrate(cot(d*x+c)^(9/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^3*cot(d*x + c)^(9/2), x)`

3.513. $\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.513.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{9/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) i)^3 dx$$

input `int(cot(c + d*x)^(9/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3,x)`output `int(cot(c + d*x)^(9/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3, x)`

3.514 $\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.514.1 Optimal result	4837
3.514.2 Mathematica [C] (verified)	4838
3.514.3 Rubi [A] (verified)	4838
3.514.4 Maple [B] (verified)	4842
3.514.5 Fricas [B] (verification not implemented)	4842
3.514.6 Sympy [F(-1)]	4843
3.514.7 Maxima [A] (verification not implemented)	4843
3.514.8 Giac [F]	4844
3.514.9 Mupad [F(-1)]	4844

3.514.1 Optimal result

Integrand size = 36, antiderivative size = 146

$$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{8\sqrt{-1}a^3(A-iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} + \frac{16a^3(6A-5iB)\sqrt{\cot(c+dx)}}{15d} - \frac{2aA\sqrt{\cot(c+dx)}(ia+a \cot(c+dx))^2}{5d} - \frac{2(9iA+5B)\sqrt{\cot(c+dx)}(ia^3+a^3 \cot(c+dx))}{15d}$$

```
output 8*(-1)^(1/4)*a^3*(A-I*B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d+16/15*a^3*(6*A-5*I*B)*cot(d*x+c)^(1/2)/d-2/5*a*A*(I*a+a*cot(d*x+c))^2*cot(d*x+c)^(1/2)/d-2/15*(9*I*A+5*B)*(I*a^3+a^3*cot(d*x+c))*cot(d*x+c)^(1/2)/d
```

3.514.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 3.58 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.77

$$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx =$$

$$\frac{2a^3 \sqrt{\cot(c+dx)}(-15iB - 5iA \cot(c+dx) - 15B \cot(c+dx) - 9A \cot^2(c+dx) + 12A \cot^2(c+dx))}{15d}$$

input `Integrate[Cot[c + d*x]^(7/2)*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(-2*a^3*Sqrt[Cot[c + d*x]]*((-15*I)*B - (5*I)*A*Cot[c + d*x] - 15*B*Cot[c + d*x] - 9*A*Cot[c + d*x]^2 + 12*A*Cot[c + d*x]^2*Hypergeometric2F1[-5/2, 1, -3/2, I*Tan[c + d*x]] + 20*B*Cot[c + d*x]*Hypergeometric2F1[-3/2, 1, -1/2, I*Tan[c + d*x]]))/(15*d)`

3.514.3 Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.05, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.361$, Rules used = {3042, 4064, 3042, 4077, 27, 3042, 4077, 27, 3042, 4075, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{7/2}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4064}$$

$$\int \frac{(a \cot(c+dx) + ia)^3(A \cot(c+dx) + B)}{\sqrt{\cot(c+dx)}} dx$$

$$\downarrow \text{3042}$$

3.514. $\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \int \frac{(-a \tan(c + dx + \frac{\pi}{2}) + ia)^3 (B - A \tan(c + dx + \frac{\pi}{2}))}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow 4077 \\
& -\frac{2}{5} \int \frac{(\cot(c + dx)a + ia)^2 (a(A - 5iB) - a(9iA + 5B) \cot(c + dx))}{2\sqrt{\cot(c + dx)}} dx - \\
& \quad \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2}{5d} \\
& \quad \downarrow 27 \\
& -\frac{1}{5} \int \frac{(\cot(c + dx)a + ia)^2 (a(A - 5iB) - a(9iA + 5B) \cot(c + dx))}{\sqrt{\cot(c + dx)}} dx - \\
& \quad \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2}{5d} \\
& \quad \downarrow 3042 \\
& -\frac{1}{5} \int \frac{(ia - a \tan(c + dx + \frac{\pi}{2}))^2 (a(A - 5iB) + a(9iA + 5B) \tan(c + dx + \frac{\pi}{2}))}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx - \\
& \quad \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2}{5d} \\
& \quad \downarrow 4077 \\
& \frac{1}{5} \left(\frac{2}{3} \int -\frac{2(\cot(c + dx)a + ia) ((3iA + 5B)a^2 + 2(6A - 5iB) \cot(c + dx)a^2)}{\sqrt{\cot(c + dx)}} dx - \frac{2(5B + 9iA)\sqrt{\cot(c + dx)}(a^3 \cot(c + dx) + ia^2)}{3d} \right. \\
& \quad \left. \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2}{5d} \right) \\
& \quad \downarrow 27 \\
& \frac{1}{5} \left(-\frac{4}{3} \int \frac{(\cot(c + dx)a + ia) ((3iA + 5B)a^2 + 2(6A - 5iB) \cot(c + dx)a^2)}{\sqrt{\cot(c + dx)}} dx - \frac{2(5B + 9iA)\sqrt{\cot(c + dx)}(a^3 \cot(c + dx) + ia^2)}{3d} \right. \\
& \quad \left. \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2}{5d} \right) \\
& \quad \downarrow 3042 \\
& \frac{1}{5} \left(-\frac{4}{3} \int \frac{(ia - a \tan(c + dx + \frac{\pi}{2})) (a^2(3iA + 5B) - 2a^2(6A - 5iB) \tan(c + dx + \frac{\pi}{2}))}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx - \frac{2(5B + 9iA)\sqrt{\cot(c + dx)}(a^3 \cot(c + dx) + ia^2)}{3d} \right. \\
& \quad \left. \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2}{5d} \right)
\end{aligned}$$

3.514. $\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

↓ 4075

$$\frac{1}{5} \left(-\frac{4}{3} \left(\int \frac{15a^3(iA+B)\cot(c+dx) - 15a^3(A-iB)}{\sqrt{\cot(c+dx)}} dx - \frac{4a^3(6A-5iB)\sqrt{\cot(c+dx)}}{d} \right) - \frac{2(5B+9iA)\sqrt{\cot(c+dx)}}{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx)+ia)^2} \right)$$

↓ 3042

$$\frac{1}{5} \left(-\frac{4}{3} \left(\int \frac{-15(A-iB)a^3 - 15(iA+B)\tan(c+dx+\frac{\pi}{2})a^3}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx - \frac{4a^3(6A-5iB)\sqrt{\cot(c+dx)}}{d} \right) - \frac{2(5B+9iA)\sqrt{\cot(c+dx)}}{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx)+ia)^2} \right)$$

↓ 4016

$$\frac{1}{5} \left(-\frac{4}{3} \left(\frac{450a^6(A-iB)^2 \int \frac{1}{15(A-iB)a^3+15(iA+B)\cot(c+dx)a^3} d\sqrt{\cot(c+dx)}}{d} - \frac{4a^3(6A-5iB)\sqrt{\cot(c+dx)}}{d} \right) - \frac{2(5B+9iA)\sqrt{\cot(c+dx)}}{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx)+ia)^2} \right)$$

↓ 221

$$\frac{1}{5} \left(-\frac{4}{3} \left(-\frac{30\sqrt[4]{-1}a^3(A-iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} - \frac{4a^3(6A-5iB)\sqrt{\cot(c+dx)}}{d} \right) - \frac{2(5B+9iA)\sqrt{\cot(c+dx)}}{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx)+ia)^2} \right)$$

input `Int[Cot[c + d*x]^(7/2)*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(-2*a*A*Sqrt[Cot[c + d*x]]*(I*a + a*Cot[c + d*x])^2)/(5*d) + ((-4*((-30*(-1)^(1/4)*a^3*(A - I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]]])/d - (4*a^3*(6*A - (5*I)*B)*Sqrt[Cot[c + d*x]]/d))/3 - (2*((9*I)*A + 5*B)*Sqrt[Cot[c + d*x]]*(I*a^3 + a^3*Cot[c + d*x]))/(3*d))/5`

3.514.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`
- rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`
- rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])], x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`
- rule 4077 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.514.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 251 vs. 2(123) = 246.

Time = 1.24 (sec) , antiderivative size = 252, normalized size of antiderivative = 1.73

method	result
derivativedivides	$a^3 \left(-\frac{2A \cot(dx+c)^{\frac{5}{2}}}{5} - 2iA \cot(dx+c)^{\frac{3}{2}} - \frac{2B \cot(dx+c)^{\frac{3}{2}}}{3} - 6iB \sqrt{\cot(dx+c)} + 8A \sqrt{\cot(dx+c)} - \frac{(-4iB+4A)\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)}{1+\cot(dx+c)} \right) \right)}{\dots} \right)$
default	$a^3 \left(-\frac{2A \cot(dx+c)^{\frac{5}{2}}}{5} - 2iA \cot(dx+c)^{\frac{3}{2}} - \frac{2B \cot(dx+c)^{\frac{3}{2}}}{3} - 6iB \sqrt{\cot(dx+c)} + 8A \sqrt{\cot(dx+c)} - \frac{(-4iB+4A)\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)}{1+\cot(dx+c)} \right) \right)}{\dots} \right)$

input `int(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output
$$a^3/d * (-2/5 * A * \cot(dx+c)^{(5/2)} - 2 * I * A * \cot(dx+c)^{(3/2)} - 2/3 * B * \cot(dx+c)^{(3/2)} - 6 * I * B * \cot(dx+c)^{(1/2)} + 8 * A * \cot(dx+c)^{(1/2)} - 1/4 * (4 * A - 4 * I * B) * 2^{(1/2)} * (\ln((1 + \cot(dx+c) + 2^{(1/2)} * \cot(dx+c)^{(1/2)}) / (1 + \cot(dx+c) - 2^{(1/2)} * \cot(dx+c)^{(1/2)}))) + 2 * \arctan(1 + 2^{(1/2)} * \cot(dx+c)^{(1/2)}) + 2 * \arctan(-1 + 2^{(1/2)} * \cot(dx+c)^{(1/2)})) - 1/4 * (-4 * I * A - 4 * B) * 2^{(1/2)} * (\ln((1 + \cot(dx+c) - 2^{(1/2)} * \cot(dx+c)^{(1/2)}) / (1 + \cot(dx+c) + 2^{(1/2)} * \cot(dx+c)^{(1/2)}))) + 2 * \arctan(1 + 2^{(1/2)} * \cot(dx+c)^{(1/2)}) + 2 * \arctan(-1 + 2^{(1/2)} * \cot(dx+c)^{(1/2)}))$$

3.514.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 449 vs. 2(118) = 236.

Time = 0.27 (sec) , antiderivative size = 449, normalized size of antiderivative = 3.08

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx =$$

$$2 \left(15 \sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^6}{d^2}} (de^{(4i dx + 4i c)} - 2 de^{(2i dx + 2i c)} + d) \log \left(\frac{2 \left((A - iB)a^3 e^{(2i dx + 2i c)} - \sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^6}{d^2}} \right)}{\dots} \right) \right)$$

input `integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output
$$\begin{aligned} & -2/15*(15*\sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2}*(d*e^{(4*I*d*x + 4*I*c)} - \\ & 2*d*e^{(2*I*d*x + 2*I*c)} + d)*\log(2*((A - I*B)*a^3*e^{(2*I*d*x + 2*I*c)} - \sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2}*(I*d*e^{(2*I*d*x + 2*I*c)} - I*d)*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))})*e^{(-2*I*d*x - 2*I*c)}/((-I*A - B)*a^3)) - 15*\sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2}*(d*e^{(4*I*d*x + 4*I*c)} - 2*d*e^{(2*I*d*x + 2*I*c)} + d)*\log(2*((A - I*B)*a^3*e^{(2*I*d*x + 2*I*c)} - \sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2}*(-I*d*e^{(2*I*d*x + 2*I*c)} + I*d)*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))})*e^{(-2*I*d*x - 2*I*c)}/((-I*A - B)*a^3)) - 2*((39*A - 25*I*B)*a^3*e^{(4*I*d*x + 4*I*c)} - 3*(19*A - 15*I*B)*a^3*e^{(2*I*d*x + 2*I*c)} + 4*(6*A - 5*I*B)*a^3)*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))}/(d*e^{(4*I*d*x + 4*I*c)} - 2*d*e^{(2*I*d*x + 2*I*c)} + d) \end{aligned}$$

3.514.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(7/2)*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output Timed out

3.514.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.37

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \frac{15 \left(2\sqrt{2}((i-1)A + (i+1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}((i-1)A + (i+1)B) \arctan\right)}{\dots}$$

input `integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

$$3.514. \quad \int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

output $1/15*(15*(2*\sqrt{2})*((I - 1)*A + (I + 1)*B)*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2/\sqrt{\tan(dx + c)})) + 2*\sqrt{2}*((I - 1)*A + (I + 1)*B)*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2/\sqrt{\tan(dx + c)})) + \sqrt{2}*(-(I + 1)*A + (I - 1)*B)*\log(\sqrt{2}/\sqrt{\tan(dx + c)} + 1/\tan(dx + c) + 1) - \sqrt{2}*(-(I + 1)*A + (I - 1)*B)*\log(-\sqrt{2}/\sqrt{\tan(dx + c)} + 1/\tan(dx + c) + 1))*a^3 + 30*(4*A - 3*I*B)*a^3/\sqrt{\tan(dx + c)} + 10*(-3*I*A - B)*a^3/\tan(dx + c)^{(3/2)} - 6*A*a^3/\tan(dx + c)^{(5/2)}/d$

3.514.8 Giac [F]

$$\int \cot^{7/2}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^3 \cot(dx + c)^{7/2} dx$$

input `integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^3*cot(d*x + c)^(7/2), x)`

3.514.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{7/2}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{7/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^3 dx$$

input `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^3,x)`

output `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^3, x)`

3.515 $\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.515.1 Optimal result	4845
3.515.2 Mathematica [C] (verified)	4845
3.515.3 Rubi [A] (verified)	4846
3.515.4 Maple [B] (verified)	4849
3.515.5 Fricas [B] (verification not implemented)	4850
3.515.6 Sympy [F(-1)]	4851
3.515.7 Maxima [A] (verification not implemented)	4851
3.515.8 Giac [F]	4852
3.515.9 Mupad [F(-1)]	4852

3.515.1 Optimal result

Integrand size = 36, antiderivative size = 138

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= -\frac{8\sqrt{-1}a^3(iA+B)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} - \frac{16ia^3A\sqrt{\cot(c+dx)}}{3d}$$

$$+ \frac{2iaB(ia+a \cot(c+dx))^2}{d\sqrt{\cot(c+dx)}} - \frac{2(A+3iB)\sqrt{\cot(c+dx)}(ia^3+a^3 \cot(c+dx))}{3d}$$

output

```
-8*(-1)^(1/4)*a^3*(I*A+B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d+2*I*a*B*(I*a+a*cot(d*x+c))^2/d/cot(d*x+c)^(1/2)-16/3*I*a^3*A*cot(d*x+c)^(1/2)/d-2/3*(A+3*I*B)*(I*a^3+a^3*cot(d*x+c))*cot(d*x+c)^(1/2)/d
```

3.515.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 3.00 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.82

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx =$$

$$-\frac{2a^3(3iB-3iA \cot(c+dx)-9B \cot(c+dx)-3A \cot^2(c+dx)+4A \cot^2(c+dx)) \operatorname{Hypergeometric2F1}}{3d\sqrt{\cot(c+dx)}}$$

3.515. $\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

input `Integrate[Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(-2*a^3*((3*I)*B - (3*I)*A*Cot[c + d*x] - 9*B*Cot[c + d*x] - 3*A*Cot[c + d*x]^2 + 4*A*Cot[c + d*x]^2*Hypergeometric2F1[-3/2, 1, -1/2, I*Tan[c + d*x]] + 12*B*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, I*Tan[c + d*x]])/(3*d*Sqrt[Cot[c + d*x]])`

3.515.3 Rubi [A] (verified)

Time = 0.94 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.02, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.361$, Rules used = {3042, 4064, 3042, 4076, 27, 3042, 4077, 27, 3042, 4075, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \cot(c + dx)^{5/2}(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{(a \cot(c + dx) + ia)^3(A \cot(c + dx) + B)}{\cot^{\frac{3}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-a \tan(c + dx + \frac{\pi}{2}) + ia)^3(B - A \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}} dx \\
 & \quad \downarrow \text{4076} \\
 & 2 \int \frac{(\cot(c + dx)a + ia)^2(a(iA + 5B) + a(A + 3iB) \cot(c + dx))}{2\sqrt{\cot(c + dx)}} dx + \frac{2iaB(a \cot(c + dx) + ia)^2}{d\sqrt{\cot(c + dx)}} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{(\cot(c + dx)a + ia)^2(a(iA + 5B) + a(A + 3iB) \cot(c + dx))}{\sqrt{\cot(c + dx)}} dx + \frac{2iaB(a \cot(c + dx) + ia)^2}{d\sqrt{\cot(c + dx)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.515. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \int \frac{(ia - a \tan(c + dx + \frac{\pi}{2}))^2 (a(iA + 5B) - a(A + 3iB) \tan(c + dx + \frac{\pi}{2}))}{\sqrt{-\tan(c + dx + \frac{\pi}{2})} \frac{2iaB(a \cot(c + dx) + ia)^2}{d\sqrt{\cot(c + dx)}}} dx + \\
& \quad \downarrow 4077 \\
& -\frac{2}{3} \int \frac{2(\cot(c + dx)a + ia) (a^2(A - 3iB) - 2ia^2 A \cot(c + dx))}{\sqrt{\cot(c + dx)}} dx - \\
& \frac{2(A + 3iB)\sqrt{\cot(c + dx)}(a^3 \cot(c + dx) + ia^3)}{3d} + \frac{2iaB(a \cot(c + dx) + ia)^2}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow 27 \\
& -\frac{4}{3} \int \frac{(\cot(c + dx)a + ia) (a^2(A - 3iB) - 2ia^2 A \cot(c + dx))}{\sqrt{\cot(c + dx)}} dx - \\
& \frac{2(A + 3iB)\sqrt{\cot(c + dx)}(a^3 \cot(c + dx) + ia^3)}{3d} + \frac{2iaB(a \cot(c + dx) + ia)^2}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow 3042 \\
& -\frac{4}{3} \int \frac{(ia - a \tan(c + dx + \frac{\pi}{2})) ((A - 3iB)a^2 + 2iA \tan(c + dx + \frac{\pi}{2}) a^2)}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx - \\
& \frac{2(A + 3iB)\sqrt{\cot(c + dx)}(a^3 \cot(c + dx) + ia^3)}{3d} + \frac{2iaB(a \cot(c + dx) + ia)^2}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow 4075 \\
& -\frac{4}{3} \left(\int \frac{3(iA + B)a^3 + 3(A - iB) \cot(c + dx)a^3}{\sqrt{\cot(c + dx)}} dx + \frac{4ia^3 A \sqrt{\cot(c + dx)}}{d} \right) - \\
& \frac{2(A + 3iB)\sqrt{\cot(c + dx)}(a^3 \cot(c + dx) + ia^3)}{3d} + \frac{2iaB(a \cot(c + dx) + ia)^2}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow 3042 \\
& -\frac{4}{3} \left(\int \frac{3a^3(iA + B) - 3a^3(A - iB) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx + \frac{4ia^3 A \sqrt{\cot(c + dx)}}{d} \right) - \\
& \frac{2(A + 3iB)\sqrt{\cot(c + dx)}(a^3 \cot(c + dx) + ia^3)}{3d} + \frac{2iaB(a \cot(c + dx) + ia)^2}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow 4016
\end{aligned}$$

3.515. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$-\frac{4}{3} \left(\frac{18a^6(B+iA)^2 \int \frac{1}{3a^3(A-iB)\cot(c+dx)-3a^3(iA+B)} d\sqrt{\cot(c+dx)} + \frac{4ia^3A\sqrt{\cot(c+dx)}}{d} \right) - \frac{2(A+3iB)\sqrt{\cot(c+dx)}(a^3\cot(c+dx)+ia^3)}{3d} + \frac{2iaB(a\cot(c+dx)+ia)^2}{d\sqrt{\cot(c+dx)}}$$

↓ 221

$$-\frac{4}{3} \left(\frac{6\sqrt[4]{-1}a^3(B+iA)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} + \frac{4ia^3A\sqrt{\cot(c+dx)}}{d} \right) - \frac{2(A+3iB)\sqrt{\cot(c+dx)}(a^3\cot(c+dx)+ia^3)}{3d} + \frac{2iaB(a\cot(c+dx)+ia)^2}{d\sqrt{\cot(c+dx)}}$$

input `Int[Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(-4*((6*(-1)^(1/4)*a^3*(I*A + B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]]])/d + ((4*I)*a^3*A*Sqrt[Cot[c + d*x]]/d))/3 + ((2*I)*a*B*(I*a + a*Cot[c + d*x])^2)/(d*Sqrt[Cot[c + d*x]]) - (2*(A + (3*I)*B)*Sqrt[Cot[c + d*x]]*(I*a^3 + a^3*Cot[c + d*x]))/(3*d)`

3.515.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4016 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp [g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4076 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

rule 4077 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

3.515.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 240 vs. $2(116) = 232$.

Time = 1.21 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.75

$$3.515. \quad \int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

method	result
derivativedivides	$a^3 \left(-\frac{2A \cot(dx+c)^{\frac{3}{2}}}{3} - 6iA \sqrt{\cot(dx+c)} - 2B \sqrt{\cot(dx+c)} - \frac{(-4iA-4B)\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(1 + \frac{\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) \right)}{4} \right)$
default	$a^3 \left(-\frac{2A \cot(dx+c)^{\frac{3}{2}}}{3} - 6iA \sqrt{\cot(dx+c)} - 2B \sqrt{\cot(dx+c)} - \frac{(-4iA-4B)\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(1 + \frac{\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) \right)}{4} \right)$

```
input int(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output a^3/d*(-2/3*A*cot(d*x+c)^(3/2)-6*I*A*cot(d*x+c)^(1/2)-2*B*cot(d*x+c)^(1/2)-1/4*(-4*I*A-4*B)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))-1/4*(4*I*B-4*A)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))-2*I*B/cot(d*x+c)^(1/2))
```

3.515.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 409 vs. 2(110) = 220.

Time = 0.26 (sec) , antiderivative size = 409, normalized size of antiderivative = 2.96

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx =$$

$$2 \left(3 \sqrt{-\frac{(iA^2+2AB-iB^2)a^6}{d^2}} (de^{(4i dx+4i c)} - d) \log \left(\frac{2 \left((A-iB)a^3 e^{(2i dx+2i c)} + \sqrt{-\frac{(iA^2+2AB-iB^2)a^6}{d^2}} (de^{(2i dx+2i c)} - d) \right) \sqrt{-\frac{(iA^2+2AB-iB^2)a^6}{d^2}}}{(-iA-B)a^3} \right) \right)$$

```
input integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fracas")
```

3.515. $\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

output
$$\begin{aligned} & -2/3*(3*\sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2}*(d*e^{(4*I*d*x + 4*I*c)} - d) \\ & * \log(2*((A - I*B)*a^3*e^{(2*I*d*x + 2*I*c)} + \sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2} \\ & *(d*e^{(2*I*d*x + 2*I*c)} - d)*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))) \\ & *e^{(-2*I*d*x - 2*I*c)}/((-I*A - B)*a^3)) - 3*\sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2} \\ & *(d*e^{(4*I*d*x + 4*I*c)} - d)* \log(2*((A - I*B)*a^3*e^{(2*I*d*x + 2*I*c)} - \sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2} \\ & *(d*e^{(2*I*d*x + 2*I*c)} - d)*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))) \\ & *e^{(-2*I*d*x - 2*I*c)}/((-I*A - B)*a^3)) + 2*((5*I*A + 3*B)*a^3*e^{(4*I*d*x + 4*I*c)} \\ & + (I*A - 3*B)*a^3*e^{(2*I*d*x + 2*I*c)} - 4*I*A*a^3)*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))} \\ & / (d*e^{(4*I*d*x + 4*I*c)} - d) \end{aligned}$$

3.515.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(5/2)*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output Timed out

3.515.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.41

$$\begin{aligned} & \int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ & = \frac{-6i Ba^3 \sqrt{\tan(dx + c)} - 3 \left(2 \sqrt{2} (-(i + 1) A + (i - 1) B) \arctan \left(\frac{1}{2} \sqrt{2} \left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx + c)}} \right) \right) + 2 \sqrt{2} (-(i + 1) A + (i - 1) B) \right)}{1} \end{aligned}$$

input `integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

3.515. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

output `1/3*(-6*I*B*a^3*sqrt(tan(d*x + c)) - 3*(2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a^3 + 6*(-3*I*A - B)*a^3/sqrt(tan(d*x + c)) - 2*A*a^3/tan(d*x + c)^(3/2))/d`

3.515.8 Giac [F]

$$\begin{aligned} & \int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^3 \cot(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

input `integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^3*cot(d*x + c)^(5/2), x)`

3.515.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ &= \int \cot(c + dx)^{5/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^3 dx \end{aligned}$$

input `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^3,x)`

output `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^3, x)`

3.516 $\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.516.1 Optimal result	4853
3.516.2 Mathematica [C] (verified)	4853
3.516.3 Rubi [A] (verified)	4854
3.516.4 Maple [A] (verified)	4857
3.516.5 Fricas [B] (verification not implemented)	4858
3.516.6 Sympy [F]	4859
3.516.7 Maxima [A] (verification not implemented)	4859
3.516.8 Giac [F]	4860
3.516.9 Mupad [F(-1)]	4860

3.516.1 Optimal result

Integrand size = 36, antiderivative size = 142

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= -\frac{8\sqrt{-1}a^3(A-iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} - \frac{16ia^3B\sqrt{\cot(c+dx)}}{3d}$$

$$+ \frac{2iaB(ia+a \cot(c+dx))^2}{3d \cot^{\frac{3}{2}}(c+dx)} - \frac{2(3A-7iB)(ia^3+a^3 \cot(c+dx))}{3d\sqrt{\cot(c+dx)}}$$

```
output -8*(-1)^(1/4)*a^3*(A-I*B)*arctanh((-1)^(3/4)*cot(dx+c)^(1/2))/d+2/3*I*a*B
*(I*a+a*cot(dx+c))^2/d/cot(dx+c)^(3/2)-2/3*(3*A-7*I*B)*(I*a^3+a^3*cot(dx+c))/d/cot(dx+c)^(1/2)-16/3*I*a^3*B*cot(dx+c)^(1/2)/d
```

3.516.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 2.95 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.80

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx =$$

$$\frac{2a^3\sqrt{\cot(c+dx)}\left(-9A+12A \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, i \tan(c+dx)\right)+12\sqrt{-1}B \operatorname{arctan}\left((-1)^3\right)\right)}{3d}$$

3.516. $\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

input `Integrate[Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(-2*a^3*Sqrt[Cot[c + d*x]]*(-9*A + 12*A*Hypergeometric2F1[-1/2, 1, 1/2, I*Tan[c + d*x]] + 12*(-1)^(1/4)*B*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sqrt[Tan[c + d*x]] + (3*I)*A*Tan[c + d*x] + 9*B*Tan[c + d*x] + I*B*Tan[c + d*x]^2))/(3*d)`

3.516.3 Rubi [A] (verified)

Time = 0.96 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.03, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.361$, Rules used = {3042, 4064, 3042, 4076, 27, 3042, 4076, 27, 3042, 4075, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \cot(c + dx)^{3/2}(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{(a \cot(c + dx) + ia)^3(A \cot(c + dx) + B)}{\cot^{\frac{5}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-a \tan(c + dx + \frac{\pi}{2}) + ia)^3(B - A \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{5/2}} dx \\
 & \quad \downarrow \text{4076} \\
 & \frac{2}{3} \int \frac{(\cot(c + dx)a + ia)^2(a(3iA + 7B) + a(3A + iB) \cot(c + dx))}{2 \cot^{\frac{3}{2}}(c + dx)} dx + \frac{2iaB(a \cot(c + dx) + ia)^2}{3d \cot^{\frac{3}{2}}(c + dx)} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{3} \int \frac{(\cot(c + dx)a + ia)^2(a(3iA + 7B) + a(3A + iB) \cot(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx + \frac{2iaB(a \cot(c + dx) + ia)^2}{3d \cot^{\frac{3}{2}}(c + dx)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.516. $\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\frac{1}{3} \int \frac{(ia - a \tan(c + dx + \frac{\pi}{2}))^2 (a(3iA + 7B) - a(3A + iB) \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2} \frac{2iaB(a \cot(c + dx) + ia)^2}{3d \cot^{\frac{3}{2}}(c + dx)}} dx +$$

↓ 4076

$$\frac{1}{3} \left(2 \int \frac{2(\cot(c + dx)a + ia) ((3iA + 5B)a^2 + 2iB \cot(c + dx)a^2)}{\sqrt{\cot(c + dx)} \frac{2iaB(a \cot(c + dx) + ia)^2}{3d \cot^{\frac{3}{2}}(c + dx)}} dx - \frac{2(3A - 7iB) (a^3 \cot(c + dx) + ia^3)}{d\sqrt{\cot(c + dx)}} \right) +$$

↓ 27

$$\frac{1}{3} \left(4 \int \frac{(\cot(c + dx)a + ia) ((3iA + 5B)a^2 + 2iB \cot(c + dx)a^2)}{\sqrt{\cot(c + dx)} \frac{2iaB(a \cot(c + dx) + ia)^2}{3d \cot^{\frac{3}{2}}(c + dx)}} dx - \frac{2(3A - 7iB) (a^3 \cot(c + dx) + ia^3)}{d\sqrt{\cot(c + dx)}} \right) +$$

↓ 3042

$$\frac{1}{3} \left(4 \int \frac{(ia - a \tan(c + dx + \frac{\pi}{2})) (a^2(3iA + 5B) - 2ia^2B \tan(c + dx + \frac{\pi}{2}))}{\sqrt{-\tan(c + dx + \frac{\pi}{2})} \frac{2iaB(a \cot(c + dx) + ia)^2}{3d \cot^{\frac{3}{2}}(c + dx)}} dx - \frac{2(3A - 7iB) (a^3 \cot(c + dx) + ia^3)}{d\sqrt{\cot(c + dx)}} \right) +$$

↓ 4075

$$\frac{1}{3} \left(4 \left(\int \frac{3a^3(iA + B) \cot(c + dx) - 3a^3(A - iB)}{\sqrt{\cot(c + dx)}} dx - \frac{4ia^3B\sqrt{\cot(c + dx)}}{d} \right) - \frac{2(3A - 7iB) (a^3 \cot(c + dx) + ia^3)}{d\sqrt{\cot(c + dx)}} \right) +$$

↓ 3042

$$\frac{1}{3} \left(4 \left(\int \frac{-3(A - iB)a^3 - 3(iA + B) \tan(c + dx + \frac{\pi}{2}) a^3}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx - \frac{4ia^3B\sqrt{\cot(c + dx)}}{d} \right) - \frac{2(3A - 7iB) (a^3 \cot(c + dx) + ia^3)}{d\sqrt{\cot(c + dx)}} \right) +$$

↓ 4016

3.516. $\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\frac{1}{3} \left(4 \left(\frac{18a^6(A-iB)^2 \int \frac{1}{3(A-iB)a^3+3(iA+B)\cot(c+dx)a^3} d\sqrt{\cot(c+dx)} - \frac{4ia^3B\sqrt{\cot(c+dx)}}{d} \right) - \frac{2(3A-7iB)(a^3 \cot(c+dx))}{d\sqrt{\cot(c+dx)}} \right) - \frac{2iaB(a \cot(c+dx) + ia)^2}{3d \cot^{\frac{3}{2}}(c+dx)}$$

↓ 221

$$\frac{1}{3} \left(4 \left(-\frac{6\sqrt[4]{-1}a^3(A-iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} - \frac{4ia^3B\sqrt{\cot(c+dx)}}{d} \right) - \frac{2(3A-7iB)(a^3 \cot(c+dx))}{d\sqrt{\cot(c+dx)}} \right) - \frac{2iaB(a \cot(c+dx) + ia)^2}{3d \cot^{\frac{3}{2}}(c+dx)}$$

input `Int[Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `((2*I)/3)*a*B*(I*a + a*Cot[c + d*x])^2/(d*Cot[c + d*x]^(3/2)) + 4*((-6*(-1)^(1/4)*a^3*(A - I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]])/d - ((4*I)*a^3*B*Sqrt[Cot[c + d*x]])/d - (2*(3*A - (7*I)*B)*(I*a^3 + a^3*Cot[c + d*x]))/(d*Sqrt[Cot[c + d*x]])/3`

3.516.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4016 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp [g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

rule 4076 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.516.4 Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.66

method	result
derivativedivides	$a^3 \left(-\frac{(4iB-4A)\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan(1+\sqrt{2}\sqrt{\cot(dx+c)}) + 2 \arctan(-1+\sqrt{2}\sqrt{\cot(dx+c)}) \right)}{4} \right) - \frac{(4iA+4B)\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan(1+\sqrt{2}\sqrt{\cot(dx+c)}) + 2 \arctan(-1+\sqrt{2}\sqrt{\cot(dx+c)}) \right)}{4}$
default	$a^3 \left(-\frac{(4iB-4A)\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan(1+\sqrt{2}\sqrt{\cot(dx+c)}) + 2 \arctan(-1+\sqrt{2}\sqrt{\cot(dx+c)}) \right)}{4} \right) - \frac{(4iA+4B)\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan(1+\sqrt{2}\sqrt{\cot(dx+c)}) + 2 \arctan(-1+\sqrt{2}\sqrt{\cot(dx+c)}) \right)}{4}$

input `int(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

$$3.516. \int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

output $a^3/d*(-1/4*(4*I*B-4*A)*2^{(1/2)}*(\ln((1+\cot(dx+c)+2^{(1/2)}*\cot(dx+c)^{(1/2)})/(1+\cot(dx+c)-2^{(1/2)}*\cot(dx+c)^{(1/2)}))+2*\arctan(1+2^{(1/2)}*\cot(dx+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\cot(dx+c)^{(1/2)}))-1/4*(4*I*A+4*B)*2^{(1/2)}*(\ln((1+\cot(dx+c)-2^{(1/2)}*\cot(dx+c)^{(1/2)})/(1+\cot(dx+c)+2^{(1/2)}*\cot(dx+c)^{(1/2)}))+2*\arctan(1+2^{(1/2)}*\cot(dx+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\cot(dx+c)^{(1/2)}))-2*A*\cot(dx+c)^{(1/2)}+2*(-I*A-3*B)/\cot(dx+c)^{(1/2)}-2/3*I*B/\cot(dx+c)^{(3/2)})$

3.516.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 442 vs. $2(112) = 224$.

Time = 0.25 (sec) , antiderivative size = 442, normalized size of antiderivative = 3.11

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{2 \left(3 \sqrt{-\frac{(-i A^2 - 2 AB + i B^2) a^6}{d^2}} (d e^{(4i dx + 4i c)} + 2 d e^{(2i dx + 2i c)} + d) \log \left(\frac{2 \left((A - i B) a^3 e^{(2i dx + 2i c)} - \sqrt{-\frac{(-i A^2 - 2 AB + i B^2) a^6}{d^2}} \right)}{(-i A - B) a^3} \right) \right)}{(-i A - B) a^3}$$

input `integrate(cot(dx+c)^(3/2)*(a+I*a*tan(dx+c))^3*(A+B*tan(dx+c)),x, algorithm="fracas")`

output $2/3*(3*\sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2}*(d*e^{(4*I*d*x + 4*I*c)} + 2*d*e^{(2*I*d*x + 2*I*c)} + d)*\log(2*((A - I*B)*a^3*e^{(2*I*d*x + 2*I*c)} - \sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2}*(I*d*e^{(2*I*d*x + 2*I*c)} - I*d)*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))}*e^{(-2*I*d*x - 2*I*c)})/((-I*A - B)*a^3)) - 3*\sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2}*(d*e^{(4*I*d*x + 4*I*c)} + 2*d*e^{(2*I*d*x + 2*I*c)} + d)*\log(2*((A - I*B)*a^3*e^{(2*I*d*x + 2*I*c)} - \sqrt{-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2}*(-I*d*e^{(2*I*d*x + 2*I*c)} + I*d)*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))}*e^{(-2*I*d*x - 2*I*c)})/((-I*A - B)*a^3)) - 2*((3*A - 5*I*B)*a^3*e^{(4*I*d*x + 4*I*c)} + (3*A + I*B)*a^3*e^{(2*I*d*x + 2*I*c)} + 4*I*B*a^3)*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))}/(d*e^{(4*I*d*x + 4*I*c)} + 2*d*e^{(2*I*d*x + 2*I*c)} + d)$

3.516.6 Sympy [F]

$$\begin{aligned} & \int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx \\ &= -ia^3 \left(\int iA \cot^{\frac{3}{2}}(c+dx) dx + \int (-3A \tan(c+dx) \cot^{\frac{3}{2}}(c+dx)) dx \right. \\ & \quad + \int A \tan^3(c+dx) \cot^{\frac{3}{2}}(c+dx) dx + \int (-3B \tan^2(c+dx) \cot^{\frac{3}{2}}(c+dx)) dx \\ & \quad + \int B \tan^4(c+dx) \cot^{\frac{3}{2}}(c+dx) dx + \int (-3iA \tan^2(c+dx) \cot^{\frac{3}{2}}(c+dx)) dx \\ & \quad \left. + \int iB \tan(c+dx) \cot^{\frac{3}{2}}(c+dx) dx + \int (-3iB \tan^3(c+dx) \cot^{\frac{3}{2}}(c+dx)) dx \right) \end{aligned}$$

input `integrate(cot(d*x+c)**(3/2)*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `-I*a**3*(Integral(I*A*cot(c + d*x)**(3/2), x) + Integral(-3*A*tan(c + d*x)*cot(c + d*x)**(3/2), x) + Integral(A*tan(c + d*x)**3*cot(c + d*x)**(3/2), x) + Integral(-3*B*tan(c + d*x)**2*cot(c + d*x)**(3/2), x) + Integral(B*tan(c + d*x)**4*cot(c + d*x)**(3/2), x) + Integral(-3*I*A*tan(c + d*x)**2*cot(c + d*x)**(3/2), x) + Integral(I*B*tan(c + d*x)*cot(c + d*x)**(3/2), x) + Integral(-3*I*B*tan(c + d*x)**3*cot(c + d*x)**(3/2), x))`

3.516.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.39

$$\begin{aligned} & \int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx = \\ & \quad 3 \left(2\sqrt{2}((i-1)A + (i+1)B) \arctan \left(\frac{1}{2}\sqrt{2} \left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}} \right) \right) + 2\sqrt{2}((i-1)A + (i+1)B) \arctan \right. \end{aligned}$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/3*(3*(2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a^3 + 6*A*a^3/sqrt(tan(d*x + c)) - 2*(-I*B*a^3 - 3*(I*A + 3*B)*a^3/tan(d*x + c))*tan(d*x + c)^(3/2))/d`

3.516.8 Giac [F]

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^3 \cot(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^3*cot(d*x + c)^(3/2), x)`

3.516.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{3/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^3 dx$$

input `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^3,x)`

output `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^3, x)`

3.517 $\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.517.1 Optimal result	4861
3.517.2 Mathematica [A] (verified)	4861
3.517.3 Rubi [A] (verified)	4862
3.517.4 Maple [A] (verified)	4866
3.517.5 Fricas [B] (verification not implemented)	4866
3.517.6 Sympy [F]	4867
3.517.7 Maxima [A] (verification not implemented)	4868
3.517.8 Giac [F]	4868
3.517.9 Mupad [F(-1)]	4869

3.517.1 Optimal result

Integrand size = 36, antiderivative size = 148

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{8\sqrt{-1}a^3(iA + B)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c + dx)}\right)}{d} - \frac{16a^3(5A - 6iB)}{15d\sqrt{\cot(c + dx)}} + \frac{2iaB(ia + a \cot(c + dx))^2}{5d \cot^{5/2}(c + dx)} - \frac{2(5A - 9iB)(ia^3 + a^3 \cot(c + dx))}{15d \cot^{3/2}(c + dx)}$$

output

```
8*(-1)^(1/4)*a^3*(I*A+B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d+2/5*I*a*B*(I*a+a*cot(d*x+c))^2/d/cot(d*x+c)^(5/2)-2/15*(5*A-9*I*B)*(I*a^3+a^3*cot(d*x+c))/d/cot(d*x+c)^(3/2)-16/15*a^3*(5*A-6*I*B)/d/cot(d*x+c)^(1/2)
```

3.517.2 Mathematica [A] (verified)

Time = 5.94 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.74

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx =$$

$$\frac{2ia^3 \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(-60\sqrt[4]{-1}(iA + B) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right) + \sqrt{\tan(c + dx)} \right)}{15d}$$

input `Integrate[Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(((-2*I)/15)*a^3*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-60*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + Sqrt[Tan[c + d*x]]*((-45*I)*A - 60*B + 5*(A - (3*I)*B)*Tan[c + d*x] + 3*B*Tan[c + d*x]^2))/d`

3.517.3 Rubi [A] (verified)

Time = 1.07 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.05, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3042, 4064, 3042, 4076, 27, 3042, 4076, 27, 3042, 4074, 27, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx \\ & \quad \downarrow \text{4064} \\ & \int \frac{(a \cot(c+dx)+ia)^3(A \cot(c+dx)+B)}{\cot^{\frac{7}{2}}(c+dx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(-a \tan(c+dx+\frac{\pi}{2})+ia)^3(B-A \tan(c+dx+\frac{\pi}{2}))}{(-\tan(c+dx+\frac{\pi}{2}))^{7/2}} dx \\ & \quad \downarrow \text{4076} \\ & \frac{2}{5} \int \frac{(\cot(c+dx)a+ia)^2(a(5iA+9B)+a(5A-iB) \cot(c+dx))}{2 \cot^{\frac{5}{2}}(c+dx)} dx + \frac{2iaB(a \cot(c+dx)+ia)^2}{5d \cot^{\frac{5}{2}}(c+dx)} \\ & \quad \downarrow \text{27} \\ & \frac{1}{5} \int \frac{(\cot(c+dx)a+ia)^2(a(5iA+9B)+a(5A-iB) \cot(c+dx))}{\cot^{\frac{5}{2}}(c+dx)} dx + \frac{2iaB(a \cot(c+dx)+ia)^2}{5d \cot^{\frac{5}{2}}(c+dx)} \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.517. $\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\frac{1}{5} \int \frac{(ia - a \tan(c + dx + \frac{\pi}{2}))^2 (a(5iA + 9B) - a(5A - iB) \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{5/2} \frac{2iaB(a \cot(c + dx) + ia)^2}{5d \cot^{\frac{5}{2}}(c + dx)}} dx +$$

↓ 4076

$$\frac{1}{5} \left(\frac{2}{3} \int \frac{2(\cot(c + dx)a + ia) (2(5iA + 6B)a^2 + (5A - 3iB) \cot(c + dx)a^2)}{\cot^{\frac{3}{2}}(c + dx) \frac{2iaB(a \cot(c + dx) + ia)^2}{5d \cot^{\frac{5}{2}}(c + dx)}} dx - \frac{2(5A - 9iB) (a^3 \cot(c + dx) + ia^3)}{3d \cot^{\frac{3}{2}}(c + dx)} \right)$$

↓ 27

$$\frac{1}{5} \left(\frac{4}{3} \int \frac{(\cot(c + dx)a + ia) (2(5iA + 6B)a^2 + (5A - 3iB) \cot(c + dx)a^2)}{\cot^{\frac{3}{2}}(c + dx) \frac{2iaB(a \cot(c + dx) + ia)^2}{5d \cot^{\frac{5}{2}}(c + dx)}} dx - \frac{2(5A - 9iB) (a^3 \cot(c + dx) + ia^3)}{3d \cot^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{5} \left(\frac{4}{3} \int \frac{(ia - a \tan(c + dx + \frac{\pi}{2})) (2a^2(5iA + 6B) - a^2(5A - 3iB) \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2} \frac{2iaB(a \cot(c + dx) + ia)^2}{5d \cot^{\frac{5}{2}}(c + dx)}} dx - \frac{2(5A - 9iB) (a^3 \cot(c + dx) + ia^3)}{3d \cot^{\frac{3}{2}}(c + dx)} \right)$$

↓ 4074

$$\frac{1}{5} \left(\frac{4}{3} \left(\int \frac{15((iA + B)a^3 + (A - iB) \cot(c + dx)a^3)}{\sqrt{\cot(c + dx)}} dx - \frac{4a^3(5A - 6iB)}{d\sqrt{\cot(c + dx)}} \right) - \frac{2(5A - 9iB) (a^3 \cot(c + dx) + ia^3)}{3d \cot^{\frac{3}{2}}(c + dx)} \right) \frac{2iaB(a \cot(c + dx) + ia)^2}{5d \cot^{\frac{5}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{5} \left(\frac{4}{3} \left(15 \int \frac{(iA + B)a^3 + (A - iB) \cot(c + dx)a^3}{\sqrt{\cot(c + dx)}} dx - \frac{4a^3(5A - 6iB)}{d\sqrt{\cot(c + dx)}} \right) - \frac{2(5A - 9iB) (a^3 \cot(c + dx) + ia^3)}{3d \cot^{\frac{3}{2}}(c + dx)} \right) \frac{2iaB(a \cot(c + dx) + ia)^2}{5d \cot^{\frac{5}{2}}(c + dx)}$$

↓ 3042

3.517. $\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\frac{1}{5} \left(\frac{4}{3} \left(15 \int \frac{a^3(iA+B) - a^3(A-iB) \tan(c+dx + \frac{\pi}{2})}{\sqrt{-\tan(c+dx + \frac{\pi}{2})}} dx - \frac{4a^3(5A-6iB)}{d\sqrt{\cot(c+dx)}} \right) - \frac{2(5A-9iB)(a^3 \cot(c+dx) + ia)}{3d \cot^{\frac{3}{2}}(c+dx)} \right) - \frac{2iaB(a \cot(c+dx) + ia)^2}{5d \cot^{\frac{5}{2}}(c+dx)}$$

↓ 4016

$$\frac{1}{5} \left(\frac{4}{3} \left(\frac{30a^6(B+ia)^2 \int \frac{1}{a^3(A-iB) \cot(c+dx) - a^3(iA+B)} d\sqrt{\cot(c+dx)}}{d} - \frac{4a^3(5A-6iB)}{d\sqrt{\cot(c+dx)}} \right) - \frac{2(5A-9iB)(a^3 \cot(c+dx) + ia)}{3d \cot^{\frac{3}{2}}(c+dx)} \right) - \frac{2iaB(a \cot(c+dx) + ia)^2}{5d \cot^{\frac{5}{2}}(c+dx)}$$

↓ 221

$$\frac{1}{5} \left(\frac{4}{3} \left(\frac{30\sqrt[4]{-1}a^3(B+ia) \operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} - \frac{4a^3(5A-6iB)}{d\sqrt{\cot(c+dx)}} \right) - \frac{2(5A-9iB)(a^3 \cot(c+dx) + ia)}{3d \cot^{\frac{3}{2}}(c+dx)} \right) - \frac{2iaB(a \cot(c+dx) + ia)^2}{5d \cot^{\frac{5}{2}}(c+dx)}$$

input `Int[Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `((2*I)/5)*a*B*(I*a + a*Cot[c + d*x])^2/(d*Cot[c + d*x]^(5/2)) + ((4*((30*(-1)^(1/4)*a^3*(I*A + B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]]])/d - (4*a^3*(5*A - (6*I)*B))/(d*Sqrt[Cot[c + d*x]])))/3 - (2*(5*A - (9*I)*B)*(I*a^3 + a^3*Cot[c + d*x]))/(3*d*Cot[c + d*x]^(3/2))/5`

3.517.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4016 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4074 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])], x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

3.517.4 Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.64

method	result
derivativedivides	$a^3 \left(\frac{(4iA+4B)\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan(1+\sqrt{2}\sqrt{\cot(dx+c)}) + 2 \arctan(-1+\sqrt{2}\sqrt{\cot(dx+c)}) \right)}{4} - \frac{(-4iB)}{4} \right)$
default	$a^3 \left(\frac{(4iA+4B)\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan(1+\sqrt{2}\sqrt{\cot(dx+c)}) + 2 \arctan(-1+\sqrt{2}\sqrt{\cot(dx+c)}) \right)}{4} - \frac{(-4iB)}{4} \right)$

```
input int(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)
```

```
output a^3/d*(-1/4*(4*I*A+4*B)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))-1/4*(4*A-4*I*B)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+2/3*(-I*A-3*B)/cot(d*x+c)^(3/2)+2*(4*I*B-3*A)/cot(d*x+c)^(1/2)-2/5*I*B/cot(d*x+c)^(5/2))
```

3.517.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 502 vs. 2(118) = 236.

Time = 0.27 (sec) , antiderivative size = 502, normalized size of antiderivative = 3.39

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{2 \left(15 \sqrt{-\frac{(iA^2+2AB-iB^2)a^6}{d^2}} (de^{(6i dx+6i c)} + 3 de^{(4i dx+4i c)} + 3 de^{(2i dx+2i c)} + d) \log \left(\frac{2 \left((A-iB)a^3 e^{(2i dx+2i c)} + \sqrt{-\dots}} \right)}{\dots} \right) \right)}{\dots}$$

```
input integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fracas")
```

3.517. $\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$

output

```

2/15*(15*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3
*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*log(2*((A - I*B)*a^3
*e^(2*I*d*x + 2*I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^6/d^2)*(d*e^(2*I*d*
x + 2*I*c) - d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1
)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3)) - 15*sqrt(-(I*A^2 + 2*A*B - I*B^
2)*a^6/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*
d*x + 2*I*c) + d)*log(2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) - sqrt(-(I*A^2
+ 2*A*B - I*B^2)*a^6/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt((I*e^(2*I*d*x +
2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*
a^3)) - 2*((-25*I*A - 39*B)*a^3*e^(6*I*d*x + 6*I*c) + 2*(-10*I*A - 9*B)*a^
3*e^(4*I*d*x + 4*I*c) + (25*I*A + 33*B)*a^3*e^(2*I*d*x + 2*I*c) + 4*(5*I*A
+ 6*B)*a^3)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))/
(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c)
+ d)

```

3.517.6 Sympy [F]

$$\begin{aligned}
& \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
&= -ia^3 \left(\int ia \sqrt{\cot(c+dx)} dx + \int (-3A \tan(c+dx) \sqrt{\cot(c+dx)}) dx \right. \\
&\quad + \int A \tan^3(c+dx) \sqrt{\cot(c+dx)} dx + \int (-3B \tan^2(c+dx) \sqrt{\cot(c+dx)}) dx \\
&\quad + \int B \tan^4(c+dx) \sqrt{\cot(c+dx)} dx + \int (-3iA \tan^2(c+dx) \sqrt{\cot(c+dx)}) dx \\
&\quad \left. + \int iB \tan(c+dx) \sqrt{\cot(c+dx)} dx + \int (-3iB \tan^3(c+dx) \sqrt{\cot(c+dx)}) dx \right)
\end{aligned}$$

input `integrate(cot(d*x+c)**(1/2)*(a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output

```

-I*a**3*(Integral(I*A*sqrt(cot(c + d*x)), x) + Integral(-3*A*tan(c + d*x)*
sqrt(cot(c + d*x)), x) + Integral(A*tan(c + d*x)**3*sqrt(cot(c + d*x)), x)
+ Integral(-3*B*tan(c + d*x)**2*sqrt(cot(c + d*x)), x) + Integral(B*tan(c
+ d*x)**4*sqrt(cot(c + d*x)), x) + Integral(-3*I*A*tan(c + d*x)**2*sqrt(c
ot(c + d*x)), x) + Integral(I*B*tan(c + d*x)*sqrt(cot(c + d*x)), x) + Inte
gral(-3*I*B*tan(c + d*x)**3*sqrt(cot(c + d*x)), x))

```

3.517.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.37

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{15 \left(2\sqrt{2}(-(i+1)A+(i-1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}+\frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}(-(i+1)A+(i-1)B) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}-\frac{2}{\sqrt{\tan(dx+c)}}\right)\right) - \sqrt{2}((i-1)A+(i+1)B) \log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}}+1\right) + \sqrt{2}((i-1)A+(i+1)B) \log\left(-\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}}+1\right) a^3 - 2(3iB a^3 - 5(-iA - 3B) a^3/\tan(dx+c) + 15(3A - 4iB) a^3/\tan(dx+c)^2) \tan(dx+c)^{5/2} \right)}{d}$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/15*(15*(2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*(-(I + 1)*A + (I - 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*((I - 1)*A + (I + 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*((I - 1)*A + (I + 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a^3 - 2*(3*I*B*a^3 - 5*(-I*A - 3*B)*a^3/tan(d*x + c) + 15*(3*A - 4*I*B)*a^3/tan(d*x + c)^2)*tan(d*x + c)^(5/2))/d`

3.517.8 Giac [F]

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \int (B \tan(dx+c) + A)(ia \tan(dx+c) + a)^3 \sqrt{\cot(dx+c)} dx$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^3*sqrt(cot(d*x + c)), x)`

3.517.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int \sqrt{\cot(c + dx)}(A + B \tan(c + dx)) (a + a \tan(c + dx) i)^3 dx$$

input `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*i)^3,x)`output `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*i)^3, x)`

3.518
$$\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.518.1 Optimal result 4870
 3.518.2 Mathematica [A] (verified) 4870
 3.518.3 Rubi [A] (verified) 4871
 3.518.4 Maple [A] (verified) 4875
 3.518.5 Fricas [B] (verification not implemented) 4876
 3.518.6 Sympy [F] 4877
 3.518.7 Maxima [A] (verification not implemented) 4878
 3.518.8 Giac [F] 4878
 3.518.9 Mupad [F(-1)] 4879

3.518.1 Optimal result

Integrand size = 36, antiderivative size = 173

$$\int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \frac{8\sqrt[4]{-1}a^3(A - iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c + dx)}\right)}{d} - \frac{8a^3(21A - 23iB)}{105d \cot^{3/2}(c + dx)}$$

$$+ \frac{8a^3(iA + B)}{d\sqrt{\cot(c + dx)}} + \frac{2iaB(ia + a \cot(c + dx))^2}{7d \cot^{7/2}(c + dx)} - \frac{2(7A - 11iB)(ia^3 + a^3 \cot(c + dx))}{35d \cot^{5/2}(c + dx)}$$

output

```
8*(-1)^(1/4)*a^3*(A-I*B)*arctanh((-1)^(3/4)*cot(d*x+c)^(1/2))/d-8/105*a^3*(21*A-23*I*B)/d/cot(d*x+c)^(3/2)+2/7*I*a*B*(I*a+a*cot(d*x+c))^2/d/cot(d*x+c)^(7/2)-2/35*(7*A-11*I*B)*(I*a^3+a^3*cot(d*x+c))/d/cot(d*x+c)^(5/2)+8*a^3*(I*A+B)/d/cot(d*x+c)^(1/2)
```

3.518.2 Mathematica [A] (verified)

Time = 6.34 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.76

$$\int \frac{(a + ia \tan(c + dx))^3(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \frac{2a^3 \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(420\sqrt[4]{-1}(iA + B) \arctan\left((-1)^{3/4}\sqrt{\tan(c + dx)}\right) + \sqrt{\tan(c + dx)}(420\right)}{105d}$$

input `Integrate[((a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `(2*a^3*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(420*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]] + Sqrt[Tan[c + d*x]]*(420*(I*A + B) - 35*(3*A - (4*I)*B)*Tan[c + d*x] - (21*I)*(A - (3*I)*B)*Tan[c + d*x]^2 - (15*I)*B*Tan[c + d*x]^3))/(105*d)`

3.518.3 Rubi [A] (verified)

Time = 1.25 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.08, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {3042, 4064, 3042, 4076, 27, 3042, 4076, 27, 3042, 4074, 27, 3042, 4012, 3042, 4016, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{(a \cot(c + dx) + ia)^3 (A \cot(c + dx) + B)}{\cot^{\frac{9}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-a \tan(c + dx + \frac{\pi}{2}) + ia)^3 (B - A \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{9/2}} dx \\
 & \quad \downarrow \text{4076} \\
 & \frac{2}{7} \int \frac{(\cot(c + dx)a + ia)^2 (a(7iA + 11B) + a(7A - 3iB) \cot(c + dx))}{2 \cot^{\frac{7}{2}}(c + dx)} dx + \\
 & \quad \frac{2iaB(a \cot(c + dx) + ia)^2}{7d \cot^{\frac{7}{2}}(c + dx)} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.518. $\int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

$$\frac{1}{7} \int \frac{(\cot(c+dx)a+ia)^2(a(7iA+11B)+a(7A-3iB)\cot(c+dx))}{\cot^{\frac{7}{2}}(c+dx)} dx + \frac{2iaB(a\cot(c+dx)+ia)^2}{7d\cot^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \int \frac{(ia-a\tan(c+dx+\frac{\pi}{2}))^2(a(7iA+11B)-a(7A-3iB)\tan(c+dx+\frac{\pi}{2}))}{(-\tan(c+dx+\frac{\pi}{2}))^{7/2}} dx + \frac{2iaB(a\cot(c+dx)+ia)^2}{7d\cot^{\frac{7}{2}}(c+dx)}$$

↓ 4076

$$\frac{1}{7} \left(\frac{2}{5} \int \frac{2(\cot(c+dx)a+ia)((21iA+23B)a^2+2(7A-6iB)\cot(c+dx)a^2)}{\cot^{\frac{5}{2}}(c+dx)} dx - \frac{2(7A-11iB)(a^3\cot(c+dx))}{5d\cot^{\frac{5}{2}}(c+dx)} \right) + \frac{2iaB(a\cot(c+dx)+ia)^2}{7d\cot^{\frac{7}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{7} \left(\frac{4}{5} \int \frac{(\cot(c+dx)a+ia)((21iA+23B)a^2+2(7A-6iB)\cot(c+dx)a^2)}{\cot^{\frac{5}{2}}(c+dx)} dx - \frac{2(7A-11iB)(a^3\cot(c+dx))}{5d\cot^{\frac{5}{2}}(c+dx)} \right) + \frac{2iaB(a\cot(c+dx)+ia)^2}{7d\cot^{\frac{7}{2}}(c+dx)}$$

↓ 3042

$$\frac{1}{7} \left(\frac{4}{5} \int \frac{(ia-a\tan(c+dx+\frac{\pi}{2}))(a^2(21iA+23B)-2a^2(7A-6iB)\tan(c+dx+\frac{\pi}{2}))}{(-\tan(c+dx+\frac{\pi}{2}))^{5/2}} dx - \frac{2(7A-11iB)(a^3\cot(c+dx))}{5d\cot^{\frac{5}{2}}(c+dx)} \right) + \frac{2iaB(a\cot(c+dx)+ia)^2}{7d\cot^{\frac{7}{2}}(c+dx)}$$

↓ 4074

$$\frac{1}{7} \left(\frac{4}{5} \left(\int \frac{35((iA+B)a^3+(A-iB)\cot(c+dx)a^3)}{\cot^{\frac{3}{2}}(c+dx)} dx - \frac{2a^3(21A-23iB)}{3d\cot^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7A-11iB)(a^3\cot(c+dx))}{5d\cot^{\frac{5}{2}}(c+dx)} \right) + \frac{2iaB(a\cot(c+dx)+ia)^2}{7d\cot^{\frac{7}{2}}(c+dx)}$$

↓ 27

3.518. $\int \frac{(a+ia\tan(c+dx))^3(A+B\tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

$$\frac{1}{7} \left(\frac{4}{5} \left(35 \int \frac{(iA+B)a^3 + (A-iB)\cot(c+dx)a^3}{\cot^{\frac{3}{2}}(c+dx)} dx - \frac{2a^3(21A-23iB)}{3d\cot^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7A-11iB)(a^3\cot(c+dx) + ia^3)}{5d\cot^{\frac{5}{2}}(c+dx)} \right. \\ \left. \frac{2iaB(a\cot(c+dx) + ia)^2}{7d\cot^{\frac{7}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{4}{5} \left(35 \int \frac{a^3(iA+B) - a^3(A-iB)\tan(c+dx + \frac{\pi}{2})}{(-\tan(c+dx + \frac{\pi}{2}))^{3/2}} dx - \frac{2a^3(21A-23iB)}{3d\cot^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7A-11iB)(a^3\cot(c+dx) + ia^3)}{5d\cot^{\frac{5}{2}}(c+dx)} \right. \\ \left. \frac{2iaB(a\cot(c+dx) + ia)^2}{7d\cot^{\frac{7}{2}}(c+dx)} \right)$$

↓ 4012

$$\frac{1}{7} \left(\frac{4}{5} \left(35 \left(\int \frac{a^3(A-iB) - a^3(iA+B)\cot(c+dx)}{\sqrt{\cot(c+dx)}} dx + \frac{2a^3(B+iA)}{d\sqrt{\cot(c+dx)}} \right) - \frac{2a^3(21A-23iB)}{3d\cot^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7A-11iB)(a^3\cot(c+dx) + ia^3)}{5d\cot^{\frac{5}{2}}(c+dx)} \right. \\ \left. \frac{2iaB(a\cot(c+dx) + ia)^2}{7d\cot^{\frac{7}{2}}(c+dx)} \right)$$

↓ 3042

$$\frac{1}{7} \left(\frac{4}{5} \left(35 \left(\int \frac{(A-iB)a^3 + (iA+B)\tan(c+dx + \frac{\pi}{2})a^3}{\sqrt{-\tan(c+dx + \frac{\pi}{2})}} dx + \frac{2a^3(B+iA)}{d\sqrt{\cot(c+dx)}} \right) - \frac{2a^3(21A-23iB)}{3d\cot^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7A-11iB)(a^3\cot(c+dx) + ia^3)}{5d\cot^{\frac{5}{2}}(c+dx)} \right. \\ \left. \frac{2iaB(a\cot(c+dx) + ia)^2}{7d\cot^{\frac{7}{2}}(c+dx)} \right)$$

↓ 4016

$$\frac{1}{7} \left(\frac{4}{5} \left(35 \left(\frac{2a^6(A-iB)^2 \int \frac{1}{-((A-iB)a^3 - (iA+B)\cot(c+dx))a^3} d\sqrt{\cot(c+dx)}}{d} + \frac{2a^3(B+iA)}{d\sqrt{\cot(c+dx)}} \right) - \frac{2a^3(21A-23iB)}{3d\cot^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7A-11iB)(a^3\cot(c+dx) + ia^3)}{5d\cot^{\frac{5}{2}}(c+dx)} \right. \\ \left. \frac{2iaB(a\cot(c+dx) + ia)^2}{7d\cot^{\frac{7}{2}}(c+dx)} \right)$$

↓ 221

$$\frac{1}{7} \left(\frac{4}{5} \left(35 \left(\frac{2\sqrt[4]{-1}a^3(A-iB)\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\cot(c+dx)}\right)}{d} + \frac{2a^3(B+iA)}{d\sqrt{\cot(c+dx)}} \right) - \frac{2a^3(21A-23iB)}{3d\cot^{\frac{3}{2}}(c+dx)} \right) - \frac{2(7A-11iB)(a^3\cot(c+dx) + ia^3)}{5d\cot^{\frac{5}{2}}(c+dx)} \right. \\ \left. \frac{2iaB(a\cot(c+dx) + ia)^2}{7d\cot^{\frac{7}{2}}(c+dx)} \right)$$

3.518. $\int \frac{(a+ia\tan(c+dx))^3(A+B\tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

input `Int[((a + I*a*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `((2*I/7)*a*B*(I*a + a*Cot[c + d*x])^2)/(d*Cot[c + d*x]^(7/2)) + ((4*(35*((-1)^(1/4)*a^3*(A - I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Cot[c + d*x]]])/d + (2*a^3*(I*A + B))/(d*Sqrt[Cot[c + d*x]]) - (2*a^3*(21*A - (23*I)*B))/(3*d*Cot[c + d*x]^(3/2)))/5 - (2*(7*A - (11*I)*B)*(I*a^3 + a^3*Cot[c + d*x]))/(5*d*Cot[c + d*x]^(5/2)))/7`

3.518.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4016 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2*(c^2/f) Subst[Int[1/(b*c - d*x^2), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && EqQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)])*(g_.)^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

```
rule 4076 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m -
n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) +
b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

3.518.4 Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.51

method	result
derivativedivides	$a^3 \left(\frac{-\frac{2iA}{5} - \frac{6B}{5}}{\cot(dx+c)^{\frac{5}{2}}} + \frac{8iA+8B}{\sqrt{\cot(dx+c)}} + \frac{\frac{8iB}{3} - 2A}{\cot(dx+c)^{\frac{3}{2}}} - \frac{2iB}{7 \cot(dx+c)^{\frac{7}{2}}} - \frac{(-4iB+4A)\sqrt{2}}{4} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) \right) \right)$
default	$a^3 \left(\frac{-\frac{2iA}{5} - \frac{6B}{5}}{\cot(dx+c)^{\frac{5}{2}}} + \frac{8iA+8B}{\sqrt{\cot(dx+c)}} + \frac{\frac{8iB}{3} - 2A}{\cot(dx+c)^{\frac{3}{2}}} - \frac{2iB}{7 \cot(dx+c)^{\frac{7}{2}}} - \frac{(-4iB+4A)\sqrt{2}}{4} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) \right) \right)$

```
input int((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x,method=_RETUR
NVERBOSE)
```

$$3.518. \int \frac{(a+ia \tan(c+dx))^3(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

```
output a^3/d*(2/5*(-I*A-3*B)/cot(d*x+c)^(5/2)+2*(4*I*A+4*B)/cot(d*x+c)^(1/2)+2/3*
(4*I*B-3*A)/cot(d*x+c)^(3/2)-2/7*I*B/cot(d*x+c)^(7/2)-1/4*(4*A-4*I*B)*2^(1
/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(
d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*co
t(d*x+c)^(1/2)))-1/4*(-4*I*A-4*B)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*
x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*co
t(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))
```

3.518.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 563 vs. $2(139) = 278$.

Time = 0.28 (sec) , antiderivative size = 563, normalized size of antiderivative = 3.25

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx =$$

$$2 \left(105 \sqrt{-\frac{(-i A^2 - 2AB + i B^2)a^6}{d^2}} (de^{(8i dx + 8i c)} + 4 de^{(6i dx + 6i c)} + 6 de^{(4i dx + 4i c)} + 4 de^{(2i dx + 2i c)} + d) \log \left(\frac{2 \left(\right)}{\right)} \right)$$

```
input integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algori
thm="fracas")
```

```

output -2/105*(105*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2)*(d*e^(8*I*d*x + 8*I*c)
+ 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*x + 2*
I*c) + d)*log(2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) - sqrt(-(-I*A^2 - 2*A*B
+ I*B^2)*a^6/d^2)*(I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt((I*e^(2*I*d*x + 2*
I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3
)) - 105*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^6/d^2)*(d*e^(8*I*d*x + 8*I*c) +
4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*x + 2*I*c)
) + d)*log(2*((A - I*B)*a^3*e^(2*I*d*x + 2*I*c) - sqrt(-(-I*A^2 - 2*A*B +
I*B^2)*a^6/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt((I*e^(2*I*d*x + 2*I*
c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/((-I*A - B)*a^3))
- 2*((273*A - 319*I*B)*a^3*e^(8*I*d*x + 8*I*c) + 3*(133*A - 109*I*B)*a^3*
e^(6*I*d*x + 6*I*c) - 5*(21*A - 19*I*B)*a^3*e^(4*I*d*x + 4*I*c) - 3*(133*A
- 129*I*B)*a^3*e^(2*I*d*x + 2*I*c) - 4*(42*A - 41*I*B)*a^3)*sqrt((I*e^(2*
I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))/(d*e^(8*I*d*x + 8*I*c) + 4
*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) + 4*d*e^(2*I*d*x + 2*I*c)
+ d)

```

3.518.6 Sympy [F]

$$\begin{aligned}
& \int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\
&= -ia^3 \left(\int \frac{iA}{\sqrt{\cot(c + dx)}} dx + \int \left(-\frac{3A \tan(c + dx)}{\sqrt{\cot(c + dx)}} \right) dx + \int \frac{A \tan^3(c + dx)}{\sqrt{\cot(c + dx)}} dx \right. \\
&\quad \left. + \int \left(-\frac{3B \tan^2(c + dx)}{\sqrt{\cot(c + dx)}} \right) dx + \int \frac{B \tan^4(c + dx)}{\sqrt{\cot(c + dx)}} dx + \int \left(-\frac{3iA \tan^2(c + dx)}{\sqrt{\cot(c + dx)}} \right) dx \right. \\
&\quad \left. + \int \frac{iB \tan(c + dx)}{\sqrt{\cot(c + dx)}} dx + \int \left(-\frac{3iB \tan^3(c + dx)}{\sqrt{\cot(c + dx)}} \right) dx \right)
\end{aligned}$$

```

input integrate((a+I*a*tan(d*x+c))**3*(A+B*tan(d*x+c))/cot(d*x+c)**(1/2),x)

```

```

output -I*a**3*(Integral(I*A/sqrt(cot(c + d*x)), x) + Integral(-3*A*tan(c + d*x)/
sqrt(cot(c + d*x)), x) + Integral(A*tan(c + d*x)**3/sqrt(cot(c + d*x)), x)
+ Integral(-3*B*tan(c + d*x)**2/sqrt(cot(c + d*x)), x) + Integral(B*tan(c
+ d*x)**4/sqrt(cot(c + d*x)), x) + Integral(-3*I*A*tan(c + d*x)**2/sqrt(c
ot(c + d*x)), x) + Integral(I*B*tan(c + d*x)/sqrt(cot(c + d*x)), x) + Inte
gral(-3*I*B*tan(c + d*x)**3/sqrt(cot(c + d*x)), x))

```

3.518. $\int \frac{(a+ia \tan(c+dx))^3 (A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

3.518.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.28

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx =$$

$$\frac{2 \left(15i B a^3 - \frac{21(-iA - 3B)a^3}{\tan(dx+c)} + \frac{35(3A - 4iB)a^3}{\tan(dx+c)^2} - \frac{420(iA+B)a^3}{\tan(dx+c)^3} \right) \tan(dx+c)^{\frac{7}{2}} - 105 \left(2\sqrt{2}((i-1)A + (i+1)B) \right)}{1}$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="maxima")`

output `-1/105*(2*(15*I*B*a^3 - 21*(-I*A - 3*B)*a^3/tan(d*x + c) + 35*(3*A - 4*I*B)*a^3/tan(d*x + c)^2 - 420*(I*A + B)*a^3/tan(d*x + c)^3)*tan(d*x + c)^(7/2) - 105*(2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((I - 1)*A + (I + 1)*B)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*(-(I + 1)*A + (I - 1)*B)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*a^3)/d`

3.518.8 Giac [F]

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^3}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^3*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^3/sqrt(cot(d*x + c)), x)`

3.518.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) i)^3}{\sqrt{\cot(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3)/cot(c + d*x)^(1/2),x)`output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^3)/cot(c + d*x)^(1/2), x)`

3.519
$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

3.519.1 Optimal result 4880
 3.519.2 Mathematica [A] (verified) 4881
 3.519.3 Rubi [A] (verified) 4881
 3.519.4 Maple [B] (verified) 4886
 3.519.5 Fricas [B] (verification not implemented) 4887
 3.519.6 Sympy [F(-1)] 4888
 3.519.7 Maxima [F(-2)] 4889
 3.519.8 Giac [F] 4889
 3.519.9 Mupad [F(-1)] 4889

3.519.1 Optimal result

Integrand size = 36, antiderivative size = 297

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= -\frac{\left(\frac{1}{4}-\frac{i}{4}\right)\left((6+i)A+(1+4i)B\right) \arctan\left(1-\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}ad}$$

$$+\frac{\left((7-5i)A+(5+3i)B\right) \arctan\left(1+\sqrt{2}\sqrt{\cot(c+dx)}\right)}{4\sqrt{2}ad}$$

$$+\frac{5(iA-B)\sqrt{\cot(c+dx)}}{2ad}-\frac{(7A+3iB)\cot^{\frac{3}{2}}(c+dx)}{6ad}+\frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{2d(ia+a \cot(c+dx))}$$

$$+\frac{\left((7+5i)A-(5-3i)B\right) \log\left(1-\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{8\sqrt{2}ad}$$

$$+\frac{\left((-7-5i)A+(5-3i)B\right) \log\left(1+\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{8\sqrt{2}ad}$$

output

```
-1/6*(7*A+3*I*B)*cot(d*x+c)^(3/2)/a/d+1/2*(A+I*B)*cot(d*x+c)^(5/2)/d/(I*a+
a*cot(d*x+c))+1/8-1/8*I*((6+I)*A+(1+4*I)*B)*arctan(-1+2^(1/2)*cot(d*x+c)
^(1/2))/a/d*2^(1/2)+1/8*((7-5*I)*A+(5+3*I)*B)*arctan(1+2^(1/2)*cot(d*x+c)
^(1/2))/a/d*2^(1/2)+1/16*((7+5*I)*A+(-5+3*I)*B)*ln(1+cot(d*x+c)-2^(1/2)*cot
(d*x+c)^(1/2))/a/d*2^(1/2)+1/16*((-7-5*I)*A+(5-3*I)*B)*ln(1+cot(d*x+c)+2^(
1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/2)+5/2*(I*A-B)*cot(d*x+c)^(1/2)/a/d
```

3.519.
$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

3.519.2 Mathematica [A] (verified)

Time = 3.47 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.63

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= \frac{\cot^{\frac{3}{2}}(c+dx) \left(4iA + 8A \tan(c+dx) + 12iB \tan(c+dx) + 15iA \tan^2(c+dx) - 15B \tan^2(c+dx) + 3\sqrt{-1} \right)}{6a \tan(c+dx)}$$

input `Integrate[(Cot[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `(Cot[c + d*x]^(3/2)*((4*I)*A + 8*A*Tan[c + d*x] + (12*I)*B*Tan[c + d*x] + (15*I)*A*Tan[c + d*x]^2 - 15*B*Tan[c + d*x]^2 + 3*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Tan[c + d*x]^(3/2)*(-I + Tan[c + d*x]) + 6*(-1)^(1/4)*(3*A + (2*I)*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Tan[c + d*x]^(3/2)*(-I + Tan[c + d*x])))/(6*a*d*(-I + Tan[c + d*x]))`

3.519.3 Rubi [A] (verified)Time = 1.10 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.89, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 4064, 3042, 4078, 27, 3042, 4011, 3042, 4011, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\cot(c+dx)^{5/2}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A \cot(c+dx) + B)}{a \cot(c+dx) + ia} dx$$

$$\downarrow \text{3042}$$

3.519. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

$$\begin{aligned}
& \int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{5/2} (B - A \tan(c+dx+\frac{\pi}{2}))}{-a \tan(c+dx+\frac{\pi}{2}) + ia} dx \\
& \quad \downarrow 4078 \\
& \int \frac{-\frac{1}{2} \cot^{\frac{3}{2}}(c+dx)(5a(iA-B) - a(7A+3iB) \cot(c+dx)) dx}{2a^2} + \frac{(A+iB) \cot^{\frac{5}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} \\
& \quad \downarrow 27 \\
& \frac{(A+iB) \cot^{\frac{5}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} - \frac{\int \cot^{\frac{3}{2}}(c+dx)(5a(iA-B) - a(7A+3iB) \cot(c+dx)) dx}{4a^2} \\
& \quad \downarrow 3042 \\
& \frac{(A+iB) \cot^{\frac{5}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} - \\
& \frac{\int (-\tan(c+dx+\frac{\pi}{2}))^{3/2} (5a(iA-B) + a(7A+3iB) \tan(c+dx+\frac{\pi}{2})) dx}{4a^2} \\
& \quad \downarrow 4011 \\
& \frac{(A+iB) \cot^{\frac{5}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} - \\
& \frac{\int \sqrt{\cot(c+dx)}(a(7A+3iB) + 5a(iA-B) \cot(c+dx)) dx + \frac{2a(7A+3iB) \cot^{\frac{3}{2}}(c+dx)}{3d}}{4a^2} \\
& \quad \downarrow 3042 \\
& \frac{(A+iB) \cot^{\frac{5}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} - \\
& \frac{\int \sqrt{-\tan(c+dx+\frac{\pi}{2})} (a(7A+3iB) - 5a(iA-B) \tan(c+dx+\frac{\pi}{2})) dx + \frac{2a(7A+3iB) \cot^{\frac{3}{2}}(c+dx)}{3d}}{4a^2} \\
& \quad \downarrow 4011 \\
& \frac{(A+iB) \cot^{\frac{5}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} - \\
& \frac{\int \frac{a(7A+3iB) \cot(c+dx) - 5a(iA-B)}{\sqrt{\cot(c+dx)}} dx + \frac{2a(7A+3iB) \cot^{\frac{3}{2}}(c+dx)}{3d} - \frac{10a(-B+iA) \sqrt{\cot(c+dx)}}{d}}{4a^2} \\
& \quad \downarrow 3042 \\
& \frac{(A+iB) \cot^{\frac{5}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} - \\
& \frac{\int \frac{-5a(iA-B) - a(7A+3iB) \tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx + \frac{2a(7A+3iB) \cot^{\frac{3}{2}}(c+dx)}{3d} - \frac{10a(-B+iA) \sqrt{\cot(c+dx)}}{d}}{4a^2}
\end{aligned}$$

3.519. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

$$\begin{aligned}
 & \downarrow 4017 \\
 & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \\
 & \frac{2 \int \frac{a(5(iA - B) - (7A + 3iB) \cot(c + dx))}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} + \frac{2a(7A + 3iB) \cot^{\frac{3}{2}}(c + dx)}{3d} - \frac{10a(-B + iA) \sqrt{\cot(c + dx)}}{d} \\
 & \frac{4a^2}{4a^2} \\
 & \downarrow 27 \\
 & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \\
 & \frac{2a \int \frac{5(iA - B) - (7A + 3iB) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} + \frac{2a(7A + 3iB) \cot^{\frac{3}{2}}(c + dx)}{3d} - \frac{10a(-B + iA) \sqrt{\cot(c + dx)}}{d} \\
 & \frac{4a^2}{4a^2} \\
 & \downarrow 1482 \\
 & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \\
 & \frac{2a \left(\frac{1}{2}((7 + 5i)A - (5 - 3i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \left(\frac{1}{2} - \frac{i}{2}\right)((6 + i)A + (1 + 4i)B) \int \frac{\cot(c + dx) + 1}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} \right)}{d} + \frac{2a(7A + 3iB) \cot^{\frac{3}{2}}(c + dx)}{3d} \\
 & \frac{4a^2}{4a^2} \\
 & \downarrow 1476 \\
 & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \\
 & \frac{2a \left(\frac{1}{2}((7 + 5i)A - (5 - 3i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \left(\frac{1}{2} - \frac{i}{2}\right)((6 + i)A + (1 + 4i)B) \left(\frac{1}{2} \int \frac{1}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)} + \frac{1}{2} \int \frac{1}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)} \right) \right)}{d} \\
 & \frac{4a^2}{4a^2} \\
 & \downarrow 1082 \\
 & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \\
 & \frac{2a \left(\frac{1}{2}((7 + 5i)A - (5 - 3i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \left(\frac{1}{2} - \frac{i}{2}\right)((6 + i)A + (1 + 4i)B) \left(\frac{\int \frac{1}{-\cot(c + dx) - 1} d(1 - \sqrt{2}\sqrt{\cot(c + dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c + dx) - 1} d(\sqrt{2}\sqrt{\cot(c + dx)})}{\sqrt{2}} \right) \right)}{d} \\
 & \frac{4a^2}{4a^2} \\
 & \downarrow 217 \\
 & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \\
 & \frac{2a \left(\frac{1}{2}((7 + 5i)A - (5 - 3i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \left(\frac{1}{2} - \frac{i}{2}\right)((6 + i)A + (1 + 4i)B) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\cot(c + dx)} + 1}{\sqrt{2}})}{\sqrt{2}} - \frac{\arctan(\frac{1 - \sqrt{2}\sqrt{\cot(c + dx)}}{\sqrt{2}})}{\sqrt{2}} \right) \right)}{d} \\
 & \frac{4a^2}{4a^2} \\
 & \downarrow 1479
 \end{aligned}$$

3.519. $\int \frac{\cot^{\frac{5}{2}}(c + dx)(A + B \tan(c + dx))}{a + ia \tan(c + dx)} dx$

$$\frac{2a \left(\frac{1}{2}((7+5i)A - (5-3i)B) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) - \left(\frac{1}{2}-\frac{i}{2}\right)((6+i)A+(1+4i)B) \right)}{d} \quad 4a^2$$

25

$$\frac{2a \left(\frac{1}{2}((7+5i)A - (5-3i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) - \left(\frac{1}{2}-\frac{i}{2}\right)((6+i)A+(1+4i)B) \right)}{d} \quad 4a^2$$

27

$$\frac{2a \left(\frac{1}{2}((7+5i)A - (5-3i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) - \left(\frac{1}{2}-\frac{i}{2}\right)((6+i)A+(1+4i)B) \right)}{d} \quad 4a^2$$

1103

$$\frac{2a \left(\frac{1}{2}((7+5i)A - (5-3i)B) \left(\frac{\log(\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right) - \left(\frac{1}{2}-\frac{i}{2}\right)((6+i)A+(1+4i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \quad 4a^2$$

input `Int[(Cot[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `((A + I*B)*Cot[c + d*x]^(5/2))/(2*d*(I*a + a*Cot[c + d*x])) - ((-10*a*(I*A - B)*Sqrt[Cot[c + d*x]])/d + (2*a*(7*A + (3*I)*B)*Cot[c + d*x]^(3/2))/(3*d) + (2*a*((-1/2 + I/2)*((6 + I)*A + (1 + 4*I)*B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + (((7 + 5*I)*A - (5 - 3*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d)/(4*a^2)`

3.519. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

3.519.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*(a + b*Tan[e + f*x])^m/(f*m), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)])*(g_.)^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4078 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.519.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 542 vs. $2(246) = 492$.

Time = 0.41 (sec) , antiderivative size = 543, normalized size of antiderivative = 1.83

3.519.
$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

method	result
derivativedivides	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \left(-12iB \tan(dx+c)^{\frac{3}{2}} \arctan\left(\left(\frac{1}{2}+\frac{i}{2}\right)\sqrt{\tan(dx+c)}\sqrt{2}\right)+12iB \tan(dx+c)^{\frac{5}{2}} \arctan\left(\left(\frac{1}{2}+\frac{i}{2}\right)\sqrt{\tan(dx+c)}\sqrt{2}\right)\right)$
default	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \left(-12iB \tan(dx+c)^{\frac{3}{2}} \arctan\left(\left(\frac{1}{2}+\frac{i}{2}\right)\sqrt{\tan(dx+c)}\sqrt{2}\right)+12iB \tan(dx+c)^{\frac{5}{2}} \arctan\left(\left(\frac{1}{2}+\frac{i}{2}\right)\sqrt{\tan(dx+c)}\sqrt{2}\right)\right)$

```
input int(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 1/12/a/d*(1/tan(d*x+c))^(5/2)*tan(d*x+c)*(-12*I*B*tan(d*x+c)^(3/2)*arctan(
(1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))+12*I*B*tan(d*x+c)^(5/2)*arctan((1/2+
1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-3*I*B*tan(d*x+c)^(5/2)*arctan((1/2-1/2*I)
*tan(d*x+c)^(1/2)*2^(1/2))-18*I*A*tan(d*x+c)^(5/2)*arctan((1/2+1/2*I)*tan(
d*x+c)^(1/2)*2^(1/2))+3*A*tan(d*x+c)^(5/2)*arctan((1/2-1/2*I)*tan(d*x+c)^(
1/2)*2^(1/2))+18*A*tan(d*x+c)^(5/2)*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(
1/2))+3*B*tan(d*x+c)^(5/2)*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))+1
2*B*tan(d*x+c)^(5/2)*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-3*I*B*ta
n(d*x+c)^(3/2)*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-18*I*A*tan(d*x
+c)^(3/2)*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-3*I*A*tan(d*x+c)^(3
/2)*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))+3*I*A*tan(d*x+c)^(5/2)*ar
ctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-12*I*B*tan(d*x+c)*2^(1/2)+3*A*t
an(d*x+c)^(3/2)*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-18*A*tan(d*x+
c)^(3/2)*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-3*B*tan(d*x+c)^(3/2)
*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))+12*B*tan(d*x+c)^(3/2)*arctan
((1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-4*I*A*2^(1/2)+15*B*tan(d*x+c)^2*2^(
1/2)-15*I*A*tan(d*x+c)^2*2^(1/2)-8*A*tan(d*x+c)*2^(1/2)*2^(1/2)/(-tan(d*x
+c)+I)
```

3.519.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 716 vs. 2(222) = 444.

Time = 0.27 (sec) , antiderivative size = 716, normalized size of antiderivative = 2.41

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx =$$

$$3 \left(a d e^{(4i dx+4i c)} - a d e^{(2i dx+2i c)} \right) \sqrt{\frac{-i A^2-2 AB+i B^2}{a^2 d^2}} \log \left(-\frac{2 \left((a d e^{(2i dx+2i c)}-a d) \sqrt{\frac{i e^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}-1}} \sqrt{\frac{-i A^2-2 AB+i B^2}{a^2 d^2}} + i A+B \right)}{i A+B} \right)$$

3.519. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="fricas")`

output `-1/24*(3*(a*d*e^(4*I*d*x + 4*I*c) - a*d*e^(2*I*d*x + 2*I*c))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2))*log(-2*((a*d*e^(2*I*d*x + 2*I*c) - a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B) - 3*(a*d*e^(4*I*d*x + 4*I*c) - a*d*e^(2*I*d*x + 2*I*c))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2))*log(2*((a*d*e^(2*I*d*x + 2*I*c) - a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B) - 6*(a*d*e^(4*I*d*x + 4*I*c) - a*d*e^(2*I*d*x + 2*I*c))*sqrt((9*I*A^2 - 12*A*B - 4*I*B^2)/(a^2*d^2))*log(((a*d*e^(2*I*d*x + 2*I*c) - a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((9*I*A^2 - 12*A*B - 4*I*B^2)/(a^2*d^2)) + 3*A + 2*I*B)*e^(-2*I*d*x - 2*I*c)/(a*d)) + 6*(a*d*e^(4*I*d*x + 4*I*c) - a*d*e^(2*I*d*x + 2*I*c))*sqrt((9*I*A^2 - 12*A*B - 4*I*B^2)/(a^2*d^2))*log(-((a*d*e^(2*I*d*x + 2*I*c) - a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((9*I*A^2 - 12*A*B - 4*I*B^2)/(a^2*d^2)) - 3*A - 2*I*B)*e^(-2*I*d*x - 2*I*c)/(a*d)) - 2*((19*I*A - 27*B)*e^(4*I*d*x + 4*I*c) - 2*(19*I*A - 15*B)*e^(2*I*d*x + 2*I*c) + 3*I*A - 3*B)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))/(a*d*e^(4*I*d*x + 4*I*c) - a*d*e^(2*I*d*x + 2*I*c))`

3.519.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)`

output `Timed out`

3.519. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

3.519.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.519.8 Giac [F]

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{5}{2}}}{ia\tan(dx+c)+a} dx$$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x+c)+A)*cot(d*x+c)^(5/2)/(I*a*tan(d*x+c)+a),x)`

3.519.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \int \frac{\cot(c+dx)^{5/2}(A+B\tan(c+dx))}{a+a\tan(c+dx)\text{li}} dx$$

input `int((cot(c+d*x)^(5/2)*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*1i),x)`

output `int((cot(c+d*x)^(5/2)*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*1i),x)`

3.519. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$

3.520
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

3.520.1 Optimal result 4890
 3.520.2 Mathematica [A] (verified) 4891
 3.520.3 Rubi [A] (verified) 4891
 3.520.4 Maple [B] (verified) 4896
 3.520.5 Fricas [B] (verification not implemented) 4897
 3.520.6 Sympy [F] 4898
 3.520.7 Maxima [F(-2)] 4898
 3.520.8 Giac [F] 4899
 3.520.9 Mupad [F(-1)] 4899

3.520.1 Optimal result

Integrand size = 36, antiderivative size = 268

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= \frac{((-5-3i)A+(3-i)B) \arctan\left(1-\sqrt{2}\sqrt{\cot(c+dx)}\right)}{4\sqrt{2}ad}$$

$$+ \frac{((5+3i)A-(3-i)B) \arctan\left(1+\sqrt{2}\sqrt{\cot(c+dx)}\right)}{4\sqrt{2}ad}$$

$$- \frac{(5A+iB)\sqrt{\cot(c+dx)}}{2ad} + \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{2d(ia+a \cot(c+dx))}$$

$$- \frac{\left(\frac{1}{8}-\frac{i}{8}\right) \left((4+i)A+(1+2i)B\right) \log\left(1-\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{\sqrt{2}ad}$$

$$+ \frac{\left((5-3i)A+(3+i)B\right) \log\left(1+\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{8\sqrt{2}ad}$$

output

```
1/2*(A+I*B)*cot(d*x+c)^(3/2)/d/(I*a+a*cot(d*x+c))-1/8*((-5-3*I)*A+(3-I)*B)
*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/2)+1/8*((5+3*I)*A+(-3+I)*B)*
arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/2)+(-1/16+1/16*I)*((4+I)*A+(1+
2*I)*B)*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/2)+1/16*((5-3*I)
)*A+(3+I)*B)*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/2)-1/2*(5*
A+I*B)*cot(d*x+c)^(1/2)/a/d
```

3.520.
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

3.520.2 Mathematica [A] (verified)

Time = 2.38 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.59

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx =$$

$$\frac{\sqrt{\cot(c+dx)} \left(-4iA + 5A \tan(c+dx) + iB \tan(c+dx) + \sqrt[4]{-1}(iA+B) \arctan \left((-1)^{3/4} \sqrt{\tan(c+dx)} \right) \right)}{a+ia \tan(c+dx)}$$

input `Integrate[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `-1/2*(Sqrt[Cot[c + d*x]]*((-4*I)*A + 5*A*Tan[c + d*x] + I*B*Tan[c + d*x] + (-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sqrt[Tan[c + d*x]]*(-I + Tan[c + d*x]) + 2*(-1)^(1/4)*((-2*I)*A + B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sqrt[Tan[c + d*x]]*(-I + Tan[c + d*x])))/(a*d*(-I + Tan[c + d*x]))`

3.520.3 Rubi [A] (verified)Time = 0.87 (sec) , antiderivative size = 234, normalized size of antiderivative = 0.87, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.528$, Rules used = {3042, 4064, 3042, 4078, 27, 3042, 4011, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\cot(c+dx)^{3/2}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A \cot(c+dx) + B)}{a \cot(c+dx) + ia} dx$$

$$\downarrow \text{3042}$$

3.520. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

$$\begin{aligned}
& \int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{3/2} (B - A \tan(c+dx+\frac{\pi}{2}))}{-a \tan(c+dx+\frac{\pi}{2}) + ia} dx \\
& \quad \downarrow 4078 \\
& \int \frac{-\frac{1}{2} \sqrt{\cot(c+dx)} (3a(iA-B) - a(5A+iB) \cot(c+dx)) dx}{2a^2} + \frac{(A+iB) \cot^{\frac{3}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} \\
& \quad \downarrow 27 \\
& \frac{(A+iB) \cot^{\frac{3}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} - \frac{\int \sqrt{\cot(c+dx)} (3a(iA-B) - a(5A+iB) \cot(c+dx)) dx}{4a^2} \\
& \quad \downarrow 3042 \\
& \frac{(A+iB) \cot^{\frac{3}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} - \frac{\int \sqrt{-\tan(c+dx+\frac{\pi}{2})} (3a(iA-B) + a(5A+iB) \tan(c+dx+\frac{\pi}{2})) dx}{4a^2} \\
& \quad \downarrow 4011 \\
& \frac{(A+iB) \cot^{\frac{3}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} - \frac{\int \frac{a(5A+iB)+3a(iA-B) \cot(c+dx)}{\sqrt{\cot(c+dx)}} dx + \frac{2a(5A+iB) \sqrt{\cot(c+dx)}}{d}}{4a^2} \\
& \quad \downarrow 3042 \\
& \frac{(A+iB) \cot^{\frac{3}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} - \frac{\int \frac{a(5A+iB)-3a(iA-B) \tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx + \frac{2a(5A+iB) \sqrt{\cot(c+dx)}}{d}}{4a^2} \\
& \quad \downarrow 4017 \\
& \frac{(A+iB) \cot^{\frac{3}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} - \frac{2 \int -\frac{a(5A+iB)+3(iA-B) \cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)} + \frac{2a(5A+iB) \sqrt{\cot(c+dx)}}{d}}{4a^2} \\
& \quad \downarrow 25 \\
& \frac{(A+iB) \cot^{\frac{3}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} - \frac{\frac{2a(5A+iB) \sqrt{\cot(c+dx)}}{d} - \frac{2 \int \frac{a(5A+iB)+3(iA-B) \cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)}}{d}}{4a^2} \\
& \quad \downarrow 27 \\
& \frac{(A+iB) \cot^{\frac{3}{2}}(c+dx)}{2d(a \cot(c+dx) + ia)} - \frac{\frac{2a(5A+iB) \sqrt{\cot(c+dx)}}{d} - \frac{2a \int \frac{5A+iB+3(iA-B) \cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)}}{d}}{4a^2} \\
& \quad \downarrow 1482
\end{aligned}$$

3.520. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

$$\frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \frac{2a(5A + iB)\sqrt{\cot(c + dx)}}{d}}{4a^2} \xrightarrow{1476} \frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \frac{2a\left(\frac{1}{2}((5-3i)A + (3+i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}((5+3i)A - (3-i)B) \int \frac{\cot(c + dx) + 1}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}\right)}{d}}{4a^2}$$

$$\xrightarrow{1082} \frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \frac{2a\left(\frac{1}{2}((5-3i)A + (3+i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}((5+3i)A - (3-i)B) \left(\frac{1}{2} \int \frac{1}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}\right)\right)}{d}}{4a^2}$$

$$\xrightarrow{217} \frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \frac{2a\left(\frac{1}{2}((5-3i)A + (3+i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}((5+3i)A - (3-i)B) \left(\frac{\int \frac{1}{-\cot(c + dx) - 1} d(1 - \sqrt{2}\sqrt{\cot(c + dx)})}{\sqrt{2}}\right)\right)}{d}}{4a^2}$$

$$\xrightarrow{1479} \frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \frac{2a\left(\frac{1}{2}((5-3i)A + (3+i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}} - \arctan(1 - \sqrt{2}\sqrt{\cot(c + dx)})\right)\right)}{d}}{4a^2}$$

$$\xrightarrow{25} \frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \frac{2a\left(\frac{1}{2}((5-3i)A + (3+i)B) \left(\frac{\int \frac{\sqrt{2} - 2\sqrt{\cot(c + dx)}}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}}\right)\right)}{d}}{4a^2}$$

$$\xrightarrow{27} \frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \frac{2a\left(\frac{1}{2}((5-3i)A + (3+i)B) \left(\frac{\int \frac{\sqrt{2} - 2\sqrt{\cot(c + dx)}}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}}\right)\right)}{d}}{4a^2}$$

3.520. $\int \frac{\cot^{\frac{3}{2}}(c + dx)(A + B \tan(c + dx))}{a + ia \tan(c + dx)} dx$

$$\frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \frac{2a(5A + iB)\sqrt{\cot(c + dx)}}{d}}{4a^2} - \frac{2a \left(\frac{1}{2}((5-3i)A + (3+i)B) \left(\int \frac{\sqrt{2-2\sqrt{\cot(c+dx)}}}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{d}}{4a^2}$$

↓ 1103

$$\frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{2d(a \cot(c + dx) + ia)} - \frac{2a(5A + iB)\sqrt{\cot(c + dx)}}{d}}{4a^2} - \frac{2a \left(\frac{1}{2}((5+3i)A - (3-i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}((5-3i)A + (3+i)B) \left(\frac{\log(\cot(c+dx))}{d} \right) \right)}{d}}{4a^2}$$

input `Int[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `((A + I*B)*Cot[c + d*x]^(3/2))/(2*d*(I*a + a*Cot[c + d*x])) - ((2*a*(5*A + I*B)*Sqrt[Cot[c + d*x]])/d - (2*a*(((5 + 3*I)*A - (3 - I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]))/2 + (((5 - 3*I)*A + (3 + I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d)/(4*a^2)`

3.520.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.520.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 512 vs. 2(222) = 444.

Time = 0.36 (sec) , antiderivative size = 513, normalized size of antiderivative = 1.91

method	result
derivativedivides	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{3}{2}} \tan(dx+c) \left(2iB \tan(dx+c)^{\frac{3}{2}} \arctan\left(\left(\frac{1}{2} + \frac{i}{2}\right) \sqrt{\tan(dx+c)} \sqrt{2}\right) + iB \tan(dx+c) \sqrt{2} + iB \sqrt{\tan(dx+c)} \arctan\left(\left(\frac{1}{2} + \frac{i}{2}\right) \sqrt{\tan(dx+c)} \sqrt{2}\right)\right)$
default	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{3}{2}} \tan(dx+c) \left(2iB \tan(dx+c)^{\frac{3}{2}} \arctan\left(\left(\frac{1}{2} + \frac{i}{2}\right) \sqrt{\tan(dx+c)} \sqrt{2}\right) + iB \tan(dx+c) \sqrt{2} + iB \sqrt{\tan(dx+c)} \arctan\left(\left(\frac{1}{2} + \frac{i}{2}\right) \sqrt{\tan(dx+c)} \sqrt{2}\right)\right)$

input `int(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,method=_RETURNV ERBOSE)`

$$3.520. \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

```
output 1/4/a/d*(1/tan(d*x+c))^(3/2)*tan(d*x+c)*(2*I*B*tan(d*x+c)^(3/2)*arctan((1/
2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))+I*B*tan(d*x+c)*2^(1/2)+I*B*tan(d*x+c)^(
1/2)*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-I*A*tan(d*x+c)^(1/2)*arc
tan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))+4*A*tan(d*x+c)^(3/2)*arctan((1/2
+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))+A*tan(d*x+c)^(3/2)*arctan((1/2-1/2*I)*ta
n(d*x+c)^(1/2)*2^(1/2))-2*B*tan(d*x+c)^(3/2)*arctan((1/2+1/2*I)*tan(d*x+c)
^(1/2)*2^(1/2))-B*tan(d*x+c)^(3/2)*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(
1/2))-I*B*tan(d*x+c)^(3/2)*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-4*
I*A*tan(d*x+c)^(1/2)*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))+4*I*A*ta
n(d*x+c)^(3/2)*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))+2*I*B*tan(d*x+
c)^(1/2)*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-I*A*tan(d*x+c)^(3/2)
*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-4*I*A*2^(1/2)+4*A*tan(d*x+c)
^(1/2)*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-A*tan(d*x+c)^(1/2)*arc
tan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))+5*A*tan(d*x+c)*2^(1/2)+2*B*tan(d
*x+c)^(1/2)*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-B*tan(d*x+c)^(1/2)
)*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2)))2^(1/2)/(-tan(d*x+c)+I)
```

3.520.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 623 vs. $2(201) = 402$.

Time = 0.26 (sec) , antiderivative size = 623, normalized size of antiderivative = 2.32

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx =$$

$$\frac{\left(ad \sqrt{\frac{iA^2+2AB-iB^2}{a^2d^2}} e^{(2i dx+2i c)} \log \left(-\frac{2 \left((iade^{(2i dx+2i c)}-iad) \sqrt{\frac{ie^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}-1}} \sqrt{\frac{iA^2+2AB-iB^2}{a^2d^2}} + (A-iB)e^{(2i dx+2i c)} \right) e^{(2i dx+2i c)}}{iA+B} \right)}{1} \right)}{1}$$

```
input integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm
m="fracas")
```

3.520. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$


```
output -1/8*(a*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(
-2*((I*a*d*e^(2*I*d*x + 2*I*c) - I*a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(
e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2)) + (A - I
*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B) - a*d*sqrt((I*A^2
+ 2*A*B - I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(-2*((-I*a*d*e^(2*I*d*
x + 2*I*c) + I*a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c)
- 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I
*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B) - 2*a*d*sqrt((-4*I*A^2 + 4*A*B + I*B^
2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(((a*d*e^(2*I*d*x + 2*I*c) - a*d)*sq
rt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-4*I*A^2 +
4*A*B + I*B^2)/(a^2*d^2)) + 2*I*A - B)*e^(-2*I*d*x - 2*I*c)/(a*d)) + 2*a*d
*sqrt((-4*I*A^2 + 4*A*B + I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(-((a*d
*e^(2*I*d*x + 2*I*c) - a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x +
2*I*c) - 1))*sqrt((-4*I*A^2 + 4*A*B + I*B^2)/(a^2*d^2)) - 2*I*A + B)*e^(-
2*I*d*x - 2*I*c)/(a*d)) + 2*((9*A + I*B)*e^(2*I*d*x + 2*I*c) - A - I*B)*sq
rt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-2*I*d*x - 2
*I*c)/(a*d)
```

3.520.6 Sympy [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = -\frac{i \left(\int \frac{A \cot^{\frac{3}{2}}(c+dx)}{\tan(c+dx)-i} dx + \int \frac{B \tan(c+dx) \cot^{\frac{3}{2}}(c+dx)}{\tan(c+dx)-i} dx \right)}{a}$$

```
input integrate(cot(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)
```

```
output -I*(Integral(A*cot(c + d*x)**(3/2)/(tan(c + d*x) - I), x) + Integral(B*tan
(c + d*x)*cot(c + d*x)**(3/2)/(tan(c + d*x) - I), x))/a
```

3.520.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorith
m="maxima")
```

3.520. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

3.520.8 Giac [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{ia\tan(dx+c)+a} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(3/2)/(I*a*tan(d*x + c) + a), x)`

3.520.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx = \int \frac{\cot(c+dx)^{3/2}(A+B\tan(c+dx))}{a+a\tan(c+dx)1i} dx$$

input `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i),x)`

output `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i), x)`

3.521
$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

3.521.1 Optimal result 4900
 3.521.2 Mathematica [A] (verified) 4901
 3.521.3 Rubi [A] (verified) 4901
 3.521.4 Maple [A] (verified) 4906
 3.521.5 Fricas [B] (verification not implemented) 4907
 3.521.6 Sympy [F] 4907
 3.521.7 Maxima [F(-2)] 4908
 3.521.8 Giac [F(-2)] 4908
 3.521.9 Mupad [F(-1)] 4909

3.521.1 Optimal result

Integrand size = 36, antiderivative size = 235

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= \frac{\left(\frac{1}{4} - \frac{i}{4}\right) \left((2+i)A+B\right) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}ad}$$

$$- \frac{\left(\frac{1}{4} - \frac{i}{4}\right) \left((2+i)A+B\right) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}ad} + \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(ia+a \cot(c+dx))}$$

$$- \frac{\left((3+i)A - (1+i)B\right) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{8\sqrt{2}ad}$$

$$+ \frac{\left((3+i)A - (1+i)B\right) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{8\sqrt{2}ad}$$

output

```
(-1/8+1/8*I)*((2+I)*A+B)*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/2)+(
-1/8+1/8*I)*((2+I)*A+B)*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/2)-1/1
6*((3+I)*A-(1+I)*B)*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/2)+
1/16*((3+I)*A-(1+I)*B)*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/
2)+1/2*(A+I*B)*cot(d*x+c)^(1/2)/d/(I*a+a*cot(d*x+c))
```

3.521.2 Mathematica [A] (verified)

Time = 3.15 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.58

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= \frac{\sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}\left((-iA+B)\sqrt{\tan(c+dx)} - \sqrt[4]{-1}(A-iB) \arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)\right)}{2ad(-i+\tan(c+dx))}$$

input `Integrate[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((-I)*A + B)*Sqrt[Tan[c + d*x]] - (-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*(-I + Tan[c + d*x]) - 2*(-1)^(1/4)*A*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*(-I + Tan[c + d*x])))/(2*a*d*(-I + Tan[c + d*x]))`

3.521.3 Rubi [A] (verified)Time = 0.71 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.87, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.472$, Rules used = {3042, 4064, 3042, 4078, 27, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{\sqrt{\cot(c+dx)}(A \cot(c+dx) + B)}{a \cot(c+dx) + ia} dx$$

$$\downarrow \text{3042}$$

3.521. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

$$\begin{aligned}
 & \int \frac{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(B-A\tan(c+dx+\frac{\pi}{2}))}{-a\tan(c+dx+\frac{\pi}{2})+ia} dx \\
 & \quad \downarrow 4078 \\
 & \frac{\int -\frac{a(iA-B)-a(3A-iB)\cot(c+dx)}{2\sqrt{\cot(c+dx)}} dx}{2a^2} + \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(a\cot(c+dx)+ia)} \\
 & \quad \downarrow 27 \\
 & \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(a\cot(c+dx)+ia)} - \frac{\int \frac{a(iA-B)-a(3A-iB)\cot(c+dx)}{\sqrt{\cot(c+dx)}} dx}{4a^2} \\
 & \quad \downarrow 3042 \\
 & \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(a\cot(c+dx)+ia)} - \frac{\int \frac{a(iA-B)+a(3A-iB)\tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{4a^2} \\
 & \quad \downarrow 4017 \\
 & \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(a\cot(c+dx)+ia)} - \frac{\int -\frac{a(iA-B)-(3A-iB)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{2a^2d} \\
 & \quad \downarrow 25 \\
 & \frac{\int \frac{a(iA-B)-(3A-iB)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{2a^2d} + \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(a\cot(c+dx)+ia)} \\
 & \quad \downarrow 27 \\
 & \frac{\int \frac{iA-B-(3A-iB)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{2ad} + \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(a\cot(c+dx)+ia)} \\
 & \quad \downarrow 1482 \\
 & \frac{\frac{1}{2}((3+i)A-(1+i)B)\int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - (\frac{1}{2}-\frac{i}{2})(B+(2+i)A)\int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{2ad} + \\
 & \quad \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(a\cot(c+dx)+ia)} \\
 & \quad \downarrow 1476 \\
 & \frac{\frac{1}{2}((3+i)A-(1+i)B)\int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - (\frac{1}{2}-\frac{i}{2})(B+(2+i)A)\left(\frac{1}{2}\int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}\right)}{2ad} \\
 & \quad \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(a\cot(c+dx)+ia)}
 \end{aligned}$$

3.521. $\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{a+ia\tan(c+dx)} dx$

↓ 1082

$$\frac{\frac{1}{2}((3+i)A - (1+i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \left(\frac{1}{2} - \frac{i}{2}\right) (B + (2+i)A) \left(\frac{\int \frac{1}{\cot(c+dx)-1} d\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right)}{2ad} \\ \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(a\cot(c+dx)+ia)}$$

↓ 217

$$\frac{\frac{1}{2}((3+i)A - (1+i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \left(\frac{1}{2} - \frac{i}{2}\right) (B + (2+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \arctan(1) \right)}{2ad} \\ \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(a\cot(c+dx)+ia)}$$

↓ 1479

$$\frac{\frac{1}{2}((3+i)A - (1+i)B) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right) (B + (2+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \arctan(1) \right)}{2ad} \\ \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(a\cot(c+dx)+ia)}$$

↓ 25

$$\frac{\frac{1}{2}((3+i)A - (1+i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right) (B + (2+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \arctan(1) \right)}{2ad} \\ \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(a\cot(c+dx)+ia)}$$

↓ 27

$$\frac{\frac{1}{2}((3+i)A - (1+i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) - \left(\frac{1}{2} - \frac{i}{2}\right) (B + (2+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \arctan(1) \right)}{2ad} \\ \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(a\cot(c+dx)+ia)}$$

↓ 1103

3.521. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

$$\frac{\frac{1}{2}((3+i)A - (1+i)B) \left(\frac{\log(\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right) - \left(\frac{1}{2} - \frac{i}{2}\right)(B + (2+i)A)}{2ad} \\ \frac{(A+iB)\sqrt{\cot(c+dx)}}{2d(a\cot(c+dx) + ia)}$$

input `Int[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x]),x]`

output `((A + I*B)*Sqrt[Cot[c + d*x]]/(2*d*(I*a + a*Cot[c + d*x])) + ((-1/2 + I/2)*((2 + I)*A + B)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + (((3 + I)*A - (1 + I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/(2*a*d)`

3.521.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2(d/e), 2]\}, \text{Simp}[e/(2c) \text{Int}[1/\text{Simp}[d/e + qx + x^2, x], x], x] + \text{Simp}[e/(2c) \text{Int}[1/\text{Simp}[d/e - qx + x^2, x], x], x]] /; \text{FreeQ}\{a, c, d, e\}, x] \&\& \text{EqQ}[c^2d^2 - a^2e^2, 0] \&\& \text{PosQ}[d^2e]$

rule 1479 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2(d/e), 2]\}, \text{Simp}[e/(2cq) \text{Int}[(q - 2x)/\text{Simp}[d/e + qx - x^2, x], x], x] + \text{Simp}[e/(2cq) \text{Int}[(q + 2x)/\text{Simp}[d/e - qx - x^2, x], x], x]] /; \text{FreeQ}\{a, c, d, e\}, x] \&\& \text{EqQ}[c^2d^2 - a^2e^2, 0] \&\& \text{NegQ}[d^2e]$

rule 1482 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a^2c, 2]\}, \text{Simp}[(dq + ae)/(2ac) \text{Int}[(q + cx^2)/(a + cx^4), x], x] + \text{Simp}[(dq - ae)/(2ac) \text{Int}[(q - cx^2)/(a + cx^4), x], x]] /; \text{FreeQ}\{a, c, d, e\}, x] \&\& \text{NeQ}[c^2d^2 + a^2e^2, 0] \&\& \text{NeQ}[c^2d^2 - a^2e^2, 0] \&\& \text{NegQ}[(-a)^2c]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4017 $\text{Int}[\frac{(c_.) + (d_.)\tan[(e_.) + (f_.)x]}{\text{Sqrt}[(b_.)\tan[(e_.) + (f_.)x] + (f_.)x]}], x_Symbol] \rightarrow \text{Simp}[2/f \text{Subst}[\text{Int}[(b^2c + dx^2)/(b^2 + x^4), x], x, \text{Sqrt}[b^2\tan[e + fx]]], x] /; \text{FreeQ}\{b, c, d, e, f\}, x] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0]$

rule 4064 $\text{Int}[(\cot[(e_.) + (f_.)x])^m (g_.)^p ((a_.) + (b_.)\tan[(e_.) + (f_.)x])^n], x_Symbol] \rightarrow \text{Simp}[g^{m+n} \text{Int}[(g^m \cot[e + fx])^{p-m-n} (b + a^m \cot[e + fx])^m (d + c^m \cot[e + fx])^n, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, p\}, x] \&\& \text{!IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4078 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.521.4 Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 342, normalized size of antiderivative = 1.46

method	result
derivativedivides	$\frac{\sqrt{\frac{1}{\tan(dx+c)}} \sqrt{\tan(dx+c)} \left(iA\sqrt{\tan(dx+c)}\sqrt{2}+2iA\arctan\left(\left(\frac{1}{2}+\frac{i}{2}\right)\sqrt{\tan(dx+c)}\sqrt{2}\right)\tan(dx+c)-iA\arctan\left(\left(\frac{1}{2}-\frac{i}{2}\right)\sqrt{\tan(dx+c)}\sqrt{2}\right)\tan(dx+c) \right)}{\sqrt{\frac{1}{\tan(dx+c)}} \sqrt{\tan(dx+c)}}$
default	$\frac{\sqrt{\frac{1}{\tan(dx+c)}} \sqrt{\tan(dx+c)} \left(iA\sqrt{\tan(dx+c)}\sqrt{2}+2iA\arctan\left(\left(\frac{1}{2}+\frac{i}{2}\right)\sqrt{\tan(dx+c)}\sqrt{2}\right)\tan(dx+c)-iA\arctan\left(\left(\frac{1}{2}-\frac{i}{2}\right)\sqrt{\tan(dx+c)}\sqrt{2}\right)\tan(dx+c) \right)}{\sqrt{\frac{1}{\tan(dx+c)}} \sqrt{\tan(dx+c)}}$

input `int(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x,method=_RETURNV
ERBOSE)`

output $\frac{1}{4} \frac{a}{d} \frac{1}{\tan(d*x+c)^{1/2}} \tan(d*x+c)^{1/2} (I*A*\tan(d*x+c)^{1/2} * 2^{1/2} + 2*I*A*\arctan((1/2+1/2*I)*\tan(d*x+c)^{1/2} * 2^{1/2}) * \tan(d*x+c) - I*A*\arctan((1/2-1/2*I)*\tan(d*x+c)^{1/2} * 2^{1/2}) * \tan(d*x+c) + I*B*\arctan((1/2-1/2*I)*\tan(d*x+c)^{1/2} * 2^{1/2}) * \tan(d*x+c) + 2*I*A*\arctan((1/2+1/2*I)*\tan(d*x+c)^{1/2} * 2^{1/2}) + I*A*\arctan((1/2-1/2*I)*\tan(d*x+c)^{1/2} * 2^{1/2}) - 2*A*\arctan((1/2+1/2*I)*\tan(d*x+c)^{1/2} * 2^{1/2}) * \tan(d*x+c) - A*\arctan((1/2-1/2*I)*\tan(d*x+c)^{1/2} * 2^{1/2}) * \tan(d*x+c) - B*\tan(d*x+c)^{1/2} * 2^{1/2} + I*B*\arctan((1/2-1/2*I)*\tan(d*x+c)^{1/2} * 2^{1/2}) - B*\arctan((1/2-1/2*I)*\tan(d*x+c)^{1/2} * 2^{1/2}) * \tan(d*x+c) + 2*A*\arctan((1/2+1/2*I)*\tan(d*x+c)^{1/2} * 2^{1/2}) - A*\arctan((1/2-1/2*I)*\tan(d*x+c)^{1/2} * 2^{1/2}) + B*\arctan((1/2-1/2*I)*\tan(d*x+c)^{1/2} * 2^{1/2})) * 2^{1/2} / (-\tan(d*x+c) + I)$

3.521.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 570 vs. $2(178) = 356$.

Time = 0.26 (sec) , antiderivative size = 570, normalized size of antiderivative = 2.43

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$$

$$= \left(ad \sqrt{\frac{-iA^2-2AB+iB^2}{a^2d^2}} e^{(2i dx+2i c)} \log \left(-\frac{2 \left((ade^{(2i dx+2i c)}-ad) \sqrt{\frac{ie^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}-1}} \sqrt{\frac{-iA^2-2AB+iB^2}{a^2d^2}} + (A-iB)e^{(2i dx+2i c)} \right) e^{(-2i dx-2i c)}}{iA+B} \right) \right)$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm m="fracas")`

output `1/8*(a*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(-2*((a*d*e^(2*I*d*x + 2*I*c) - a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - a*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(2*((a*d*e^(2*I*d*x + 2*I*c) - a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 2*a*d*sqrt(I*A^2/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(-((a*d*e^(2*I*d*x + 2*I*c) - a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt(I*A^2/(a^2*d^2)) + A)*e^(-2*I*d*x - 2*I*c)/(a*d)) + 2*a*d*sqrt(I*A^2/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(((a*d*e^(2*I*d*x + 2*I*c) - a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt(I*A^2/(a^2*d^2)) - A)*e^(-2*I*d*x - 2*I*c)/(a*d)) + 2*((-I*A + B)*e^(2*I*d*x + 2*I*c) + I*A - B)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*e^(-2*I*d*x - 2*I*c)/(a*d)`

3.521.6 Sympy [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = -\frac{i \left(\int \frac{A \sqrt{\cot(c+dx)}}{\tan(c+dx)-i} dx + \int \frac{B \tan(c+dx) \sqrt{\cot(c+dx)}}{\tan(c+dx)-i} dx \right)}{a}$$

input `integrate(cot(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x)`

3.521. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx$

output `-I*(Integral(A*sqrt(cot(c + d*x))/(tan(c + d*x) - I), x) + Integral(B*tan(c + d*x)*sqrt(cot(c + d*x))/(tan(c + d*x) - I), x))/a`

3.521.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\cot(c + dx)}(A + B \tan(c + dx))}{a + ia \tan(c + dx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.521.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\cot(c + dx)}(A + B \tan(c + dx))}{a + ia \tan(c + dx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c)),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command :INPUT:sage2OUTPUT:Not invertible Error: Bad Argument Value`

3.521.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+ia \tan(c+dx)} dx = \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+a \tan(c+dx) \text{ li}} dx$$

input `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i),x)`output `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i), x)`

$$3.522 \quad \int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+ia \tan(c+dx))}} dx$$

3.522.1 Optimal result 4910
 3.522.2 Mathematica [A] (verified) 4911
 3.522.3 Rubi [A] (verified) 4911
 3.522.4 Maple [A] (verified) 4916
 3.522.5 Fricas [B] (verification not implemented) 4916
 3.522.6 Sympy [F] 4917
 3.522.7 Maxima [F(-2)] 4918
 3.522.8 Giac [F] 4918
 3.522.9 Mupad [F(-1)] 4918

3.522.1 Optimal result

Integrand size = 36, antiderivative size = 237

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))}} dx$$

$$= \frac{\left(\frac{1}{4} - \frac{i}{4}\right) (A + (2 - i)B) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}ad}$$

$$- \frac{\left(\frac{1}{4} - \frac{i}{4}\right) (A + (2 - i)B) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}ad} + \frac{(iA - B)\sqrt{\cot(c + dx)}}{2d(ia + a \cot(c + dx))}$$

$$+ \frac{\left(\frac{1}{8} + \frac{i}{8}\right) (A - (2 + i)B) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{\sqrt{2}ad}$$

$$- \frac{\left(\frac{1}{8} + \frac{i}{8}\right) (A - (2 + i)B) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{\sqrt{2}ad}$$

```
output (-1/8+1/8*I)*(A+(2-I)*B)*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/2)+(
-1/8+1/8*I)*(A+(2-I)*B)*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/2)+(1/
16+1/16*I)*(A-(2+I)*B)*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/
2)-(1/16+1/16*I)*(A-(2+I)*B)*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/a/d
*2^(1/2)+1/2*(I*A-B)*cot(d*x+c)^(1/2)/d/(I*a+a*cot(d*x+c))
```

3.522.2 Mathematica [A] (verified)

Time = 3.38 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.57

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))} dx$$

$$= \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left((A + iB) \sqrt{\tan(c + dx)} + \sqrt[4]{-1} (iA + B) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right) \right)}{2ad(-i + \tan(c + dx))}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((A + I*B)*Sqrt[Tan[c + d*x]] + (-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]*(-I + Tan[c + d*x])] - 2*(-1)^(1/4)*B*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]*(-I + Tan[c + d*x])])/(2*a*d*(-I + Tan[c + d*x]))`

3.522.3 Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 207, normalized size of antiderivative = 0.87, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.472$, Rules used = {3042, 4064, 3042, 4079, 27, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{A \cot(c + dx) + B}{\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} dx$$

$$\downarrow \text{3042}$$

$$\begin{aligned}
 & \int \frac{B - A \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})} (-a \tan(c + dx + \frac{\pi}{2}) + ia)} dx \\
 & \quad \downarrow \text{4079} \\
 & \frac{\int \frac{a(A-3iB) - a(iA-B) \cot(c+dx)}{2\sqrt{\cot(c+dx)}} dx}{2a^2} + \frac{(-B + iA) \sqrt{\cot(c + dx)}}{2d(a \cot(c + dx) + ia)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{a(A-3iB) - a(iA-B) \cot(c+dx)}{\sqrt{\cot(c+dx)}} dx}{4a^2} + \frac{(-B + iA) \sqrt{\cot(c + dx)}}{2d(a \cot(c + dx) + ia)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{a(A-3iB) + a(iA-B) \tan(c+dx + \frac{\pi}{2})}{\sqrt{-\tan(c+dx + \frac{\pi}{2})}} dx}{4a^2} + \frac{(-B + iA) \sqrt{\cot(c + dx)}}{2d(a \cot(c + dx) + ia)} \\
 & \quad \downarrow \text{4017} \\
 & \frac{\int -\frac{a(A-3iB) - (iA-B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c + dx)}}{2a^2d} + \frac{(-B + iA) \sqrt{\cot(c + dx)}}{2d(a \cot(c + dx) + ia)} \\
 & \quad \downarrow \text{25} \\
 & \frac{(-B + iA) \sqrt{\cot(c + dx)}}{2d(a \cot(c + dx) + ia)} - \frac{\int \frac{a(A-3iB) - (iA-B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c + dx)}}{2a^2d} \\
 & \quad \downarrow \text{27} \\
 & \frac{(-B + iA) \sqrt{\cot(c + dx)}}{2d(a \cot(c + dx) + ia)} - \frac{\int \frac{A-3iB - (iA-B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c + dx)}}{2ad} \\
 & \quad \downarrow \text{1482} \\
 & \frac{(-B + iA) \sqrt{\cot(c + dx)}}{2d(a \cot(c + dx) + ia)} - \\
 & \frac{(\frac{1}{2} + \frac{i}{2}) (A - (2 + i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c + dx)} + (\frac{1}{2} - \frac{i}{2}) (A + (2 - i)B) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c + dx)}}{2ad} \\
 & \quad \downarrow \text{1476} \\
 & \frac{(-B + iA) \sqrt{\cot(c + dx)}}{2d(a \cot(c + dx) + ia)} - \\
 & \frac{(\frac{1}{2} + \frac{i}{2}) (A - (2 + i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c + dx)} + (\frac{1}{2} - \frac{i}{2}) (A + (2 - i)B) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c + dx)} \right)}{2ad} \\
 & \quad \downarrow \text{1082}
 \end{aligned}$$

3.522. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+ia \tan(c+dx))}} dx$

$$\frac{\frac{(-B + iA)\sqrt{\cot(c + dx)}}{2d(a \cot(c + dx) + ia)} - \left(\frac{1}{2} + \frac{i}{2}\right) (A - (2 + i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \left(\frac{1}{2} - \frac{i}{2}\right) (A + (2 - i)B) \left(\frac{\int \frac{1}{\cot(c + dx) - 1} d(1 - \sqrt{2}\sqrt{\cot(c + dx)})}{\sqrt{2}}\right)}{2ad}$$

$$\downarrow \text{217}$$

$$\frac{\frac{(-B + iA)\sqrt{\cot(c + dx)}}{2d(a \cot(c + dx) + ia)} - \left(\frac{1}{2} + \frac{i}{2}\right) (A - (2 + i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \left(\frac{1}{2} - \frac{i}{2}\right) (A + (2 - i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}} - \frac{\arctan(1)}{\sqrt{2}}\right)}{2ad}$$

$$\downarrow \text{1479}$$

$$\frac{\frac{(-B + iA)\sqrt{\cot(c + dx)}}{2d(a \cot(c + dx) + ia)} - \left(\frac{1}{2} + \frac{i}{2}\right) (A - (2 + i)B) \left(-\frac{\int -\frac{\sqrt{2} - 2\sqrt{\cot(c + dx)}}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}}\right) + \left(\frac{1}{2} - \frac{i}{2}\right) (A + (2 - i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}} - \frac{\arctan(1)}{\sqrt{2}}\right)}{2ad}$$

$$\downarrow \text{25}$$

$$\frac{\frac{(-B + iA)\sqrt{\cot(c + dx)}}{2d(a \cot(c + dx) + ia)} - \left(\frac{1}{2} + \frac{i}{2}\right) (A - (2 + i)B) \left(\frac{\int \frac{\sqrt{2} - 2\sqrt{\cot(c + dx)}}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}}\right) + \left(\frac{1}{2} - \frac{i}{2}\right) (A + (2 - i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}} - \frac{\arctan(1)}{\sqrt{2}}\right)}{2ad}$$

$$\downarrow \text{27}$$

$$\frac{\frac{(-B + iA)\sqrt{\cot(c + dx)}}{2d(a \cot(c + dx) + ia)} - \left(\frac{1}{2} + \frac{i}{2}\right) (A - (2 + i)B) \left(\frac{\int \frac{\sqrt{2} - 2\sqrt{\cot(c + dx)}}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c + dx) + 1}}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}\right) + \left(\frac{1}{2} - \frac{i}{2}\right) (A + (2 - i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}} - \frac{\arctan(1)}{\sqrt{2}}\right)}{2ad}$$

$$\downarrow \text{1103}$$

$$\frac{\frac{(-B + iA)\sqrt{\cot(c + dx)}}{2d(a \cot(c + dx) + ia)} - \left(\frac{1}{2} - \frac{i}{2}\right) (A + (2 - i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\cot(c + dx)})}{\sqrt{2}}\right) + \left(\frac{1}{2} + \frac{i}{2}\right) (A - (2 + i)B) \left(\frac{\log(\cot(c + dx))}{\sqrt{2}}\right)}{2ad}$$

3.522. $\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))}} dx$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])),x]`

output `((I*A - B)*Sqrt[Cot[c + d*x]]/(2*d*(I*a + a*Cot[c + d*x])) - ((1/2 - I/2)*(A + (2 - I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]/Sqrt[2]) + (1/2 + I/2)*(A - (2 + I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/(2*a*d)`

3.522.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)])*(g_.)^p)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^m)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^n, x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^m)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^n, x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.522.4 Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.54

method	result	size
derivativedivides	$\frac{4\left(\frac{iA}{4} + \frac{B}{4}\right) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} + \frac{i\left(-\frac{i(iA-B)\sqrt{\cot(dx+c)}}{i+\cot(dx+c)} + \frac{4iB \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}\right)}{2}$	127
default	$\frac{4\left(\frac{iA}{4} + \frac{B}{4}\right) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} + \frac{i\left(-\frac{i(iA-B)\sqrt{\cot(dx+c)}}{i+\cot(dx+c)} + \frac{4iB \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}\right)}{2}$	127

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)),x,method=_RETURNV
ERBOSE)`

output `1/a/d*(4*(1/4*I*A+1/4*B)/(2^(1/2)-I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))+1/2*I*(-I*(I*A-B)*cot(d*x+c)^(1/2)/(I+cot(d*x+c))+4*I*B/(2^(1/2)+I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2))))`

3.522.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 571 vs. 2(178) = 356.

Time = 0.27 (sec) , antiderivative size = 571, normalized size of antiderivative = 2.41

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))}} dx$$

$$\left(ad \sqrt{\frac{iA^2 + 2AB - iB^2}{a^2 d^2}} e^{(2i dx + 2i c)} \log \left(-\frac{2 \left((i a d e^{(2i dx + 2i c)} - i a d) \sqrt{\frac{i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} - 1}} \sqrt{\frac{iA^2 + 2AB - iB^2}{a^2 d^2}} + (A - iB) e^{(2i dx + 2i c)} \right)}{iA + B} \right) e^{(-2i dx - 2i c)} \right)$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)),x, algorithm
m="fracas")`

```
output 1/8*(a*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(-
2*((I*a*d*e^(2*I*d*x + 2*I*c) - I*a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e
^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2)) + (A - I*
B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - a*d*sqrt((I*A^2
+ 2*A*B - I*B^2)/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(-2*((-I*a*d*e^(2*I*d*x
+ 2*I*c) + I*a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) -
1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I*
c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 2*a*d*sqrt(I*B^2/(a^2*d^2))*e^(2*I*d
*x + 2*I*c)*log(-((a*d*e^(2*I*d*x + 2*I*c) - a*d)*sqrt((I*e^(2*I*d*x + 2*I
*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt(I*B^2/(a^2*d^2)) + B)*e^(-2*I*d*x
- 2*I*c)/(a*d)) + 2*a*d*sqrt(I*B^2/(a^2*d^2))*e^(2*I*d*x + 2*I*c)*log(((a
*d*e^(2*I*d*x + 2*I*c) - a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x
+ 2*I*c) - 1))*sqrt(I*B^2/(a^2*d^2)) - B)*e^(-2*I*d*x - 2*I*c)/(a*d)) + 2
*((A + I*B)*e^(2*I*d*x + 2*I*c) - A - I*B)*sqrt((I*e^(2*I*d*x + 2*I*c) + I
)/(e^(2*I*d*x + 2*I*c) - 1))*e^(-2*I*d*x - 2*I*c)/(a*d)
```

3.522.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))} dx$$

$$= \frac{i \left(\int \frac{A}{\tan(c+dx)\sqrt{\cot(c+dx)} - i\sqrt{\cot(c+dx)}} dx + \int \frac{B \tan(c+dx)}{\tan(c+dx)\sqrt{\cot(c+dx)} - i\sqrt{\cot(c+dx)}} dx \right)}{a}$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)**(1/2)/(a+I*a*tan(d*x+c)),x)
```

```
output -I*(Integral(A/(tan(c + d*x)*sqrt(cot(c + d*x)) - I*sqrt(cot(c + d*x))), x
) + Integral(B*tan(c + d*x)/(tan(c + d*x)*sqrt(cot(c + d*x)) - I*sqrt(cot(
c + d*x))), x))/a
```

3.522.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)),x, algorithm m="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.522.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))}} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)\sqrt{\cot(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)),x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)*sqrt(cot(d*x + c))), x)`

3.522.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + a \tan(c + dx) 1i)}} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)), x)`

3.523
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} dx$$

3.523.1 Optimal result 4919
 3.523.2 Mathematica [A] (verified) 4920
 3.523.3 Rubi [A] (verified) 4920
 3.523.4 Maple [A] (verified) 4925
 3.523.5 Fricas [B] (verification not implemented) 4926
 3.523.6 Sympy [F] 4927
 3.523.7 Maxima [F(-2)] 4927
 3.523.8 Giac [F] 4927
 3.523.9 Mupad [F(-1)] 4928

3.523.1 Optimal result

Integrand size = 36, antiderivative size = 276

$$\begin{aligned} & \int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} dx \\ &= \frac{((1-3i)A+(3+5i)B) \arctan\left(1-\sqrt{2}\sqrt{\cot(c+dx)}\right)}{4\sqrt{2}ad} \\ & \quad + \frac{\left(\frac{1}{4}+\frac{i}{4}\right) ((1+2i)A-(4+i)B) \arctan\left(1+\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}ad} \\ & \quad - \frac{A+5iB}{2ad\sqrt{\cot(c+dx)}} + \frac{iA-B}{2d\sqrt{\cot(c+dx)}(ia+a \cot(c+dx))} \\ & \quad - \frac{\left(\frac{1}{8}+\frac{i}{8}\right) ((2+i)A+(1+4i)B) \log\left(1-\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{\sqrt{2}ad} \\ & \quad + \frac{\left(\frac{1}{8}+\frac{i}{8}\right) ((2+i)A+(1+4i)B) \log\left(1+\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{\sqrt{2}ad} \end{aligned}$$

```
output -1/8*((1-3*I)*A+(3+5*I)*B)*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/2)
+(1/8+1/8*I)*((1+2*I)*A-(4+I)*B)*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/2)
-(1/16+1/16*I)*((2+I)*A+(1+4*I)*B)*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/2)
+(1/16+1/16*I)*((2+I)*A+(1+4*I)*B)*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/a/d*2^(1/2)
+1/2*(-A-5*I*B)/a/d/cot(d*x+c)^(1/2)+1/2*(I*A-B)/d/(I*a+a*cot(d*x+c))/cot(d*x+c)^(1/2)
```

3.523.2 Mathematica [A] (verified)

Time = 4.02 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.56

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$= \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(\sqrt[4]{-1} (A - iB) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right) (-i + \tan(c + dx)) - 2 \sqrt[4]{-1} \right)}{2ad(-i + \dots)}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*(-I + Tan[c + d*x]) - 2*(-1)^(1/4)*(A + (2*I)*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*(-I + Tan[c + d*x]) + I*Sqrt[Tan[c + d*x]]*(A + (5*I)*B - 4*B*Tan[c + d*x])))/(2*a*d*(-I + Tan[c + d*x]))`

3.523.3 Rubi [A] (verified)

Time = 0.87 (sec) , antiderivative size = 242, normalized size of antiderivative = 0.88, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.528$, Rules used = {3042, 4064, 3042, 4079, 27, 3042, 4012, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2}(a + ia \tan(c + dx))} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{A \cot(c + dx) + B}{\cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + ia)} dx$$

$$\downarrow \text{3042}$$

3.523. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

$$\begin{aligned}
 & \int \frac{B - A \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2} (-a \tan(c + dx + \frac{\pi}{2}) + ia)} dx \\
 & \quad \downarrow 4079 \\
 & \frac{\int -\frac{a(A+5iB)+3a(iA-B)\cot(c+dx)}{2\cot^{\frac{3}{2}}(c+dx)} dx}{2a^2} + \frac{-B + iA}{2d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} \\
 & \quad \downarrow 27 \\
 & \frac{-B + iA}{2d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{\int \frac{a(A+5iB)+3a(iA-B)\cot(c+dx)}{\cot^{\frac{3}{2}}(c+dx)} dx}{4a^2} \\
 & \quad \downarrow 3042 \\
 & \frac{-B + iA}{2d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{\int \frac{a(A+5iB)-3a(iA-B)\tan(c+dx+\frac{\pi}{2})}{(-\tan(c+dx+\frac{\pi}{2}))^{3/2}} dx}{4a^2} \\
 & \quad \downarrow 4012 \\
 & \frac{-B + iA}{2d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{\int \frac{3a(iA-B)-a(A+5iB)\cot(c+dx)}{\sqrt{\cot(c+dx)}} dx + \frac{2a(A+5iB)}{d\sqrt{\cot(c+dx)}}}{4a^2} \\
 & \quad \downarrow 3042 \\
 & \frac{-B + iA}{2d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{\int \frac{3a(iA-B)+a(A+5iB)\tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx + \frac{2a(A+5iB)}{d\sqrt{\cot(c+dx)}}}{4a^2} \\
 & \quad \downarrow 4017 \\
 & \frac{-B + iA}{2d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{2 \int -\frac{a(3(iA-B)-(A+5iB)\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{2a(A+5iB)}{d\sqrt{\cot(c+dx)}}}{4a^2} \\
 & \quad \downarrow 25 \\
 & \frac{-B + iA}{2d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{\frac{2a(A+5iB)}{d\sqrt{\cot(c+dx)}} - \frac{2 \int \frac{a(3(iA-B)-(A+5iB)\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d}}{4a^2} \\
 & \quad \downarrow 27 \\
 & \frac{-B + iA}{2d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{\frac{2a(A+5iB)}{d\sqrt{\cot(c+dx)}} - \frac{2a \int \frac{3(iA-B)-(A+5iB)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d}}{4a^2} \\
 & \quad \downarrow 1482
 \end{aligned}$$

3.523. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

$$\begin{aligned}
 & \frac{2a(A+5iB)}{d\sqrt{\cot(c+dx)}} - \frac{-B+iA}{2d\sqrt{\cot(c+dx)}(a\cot(c+dx)+ia)} \\
 & \frac{2a\left(\left(\frac{1}{2}+\frac{i}{2}\right)((2+i)A+(1+4i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \left(\frac{1}{2}+\frac{i}{2}\right)((1+2i)A-(4+i)B) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{d} \\
 & \frac{4a^2}{1476} \\
 & \frac{2a(A+5iB)}{d\sqrt{\cot(c+dx)}} - \frac{-B+iA}{2d\sqrt{\cot(c+dx)}(a\cot(c+dx)+ia)} \\
 & \frac{2a\left(\left(\frac{1}{2}+\frac{i}{2}\right)((2+i)A+(1+4i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \left(\frac{1}{2}+\frac{i}{2}\right)((1+2i)A-(4+i)B) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}\right)\right)}{d} \\
 & \frac{4a^2}{1082} \\
 & \frac{2a(A+5iB)}{d\sqrt{\cot(c+dx)}} - \frac{-B+iA}{2d\sqrt{\cot(c+dx)}(a\cot(c+dx)+ia)} \\
 & \frac{2a\left(\left(\frac{1}{2}+\frac{i}{2}\right)((2+i)A+(1+4i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \left(\frac{1}{2}+\frac{i}{2}\right)((1+2i)A-(4+i)B) \left(\int \frac{1}{-\cot(c+dx)-1} \frac{d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}}\right)\right)}{d} \\
 & \frac{4a^2}{217} \\
 & \frac{2a(A+5iB)}{d\sqrt{\cot(c+dx)}} - \frac{-B+iA}{2d\sqrt{\cot(c+dx)}(a\cot(c+dx)+ia)} \\
 & \frac{2a\left(\left(\frac{1}{2}+\frac{i}{2}\right)((2+i)A+(1+4i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \left(\frac{1}{2}+\frac{i}{2}\right)((1+2i)A-(4+i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}}\right)\right)}{d} \\
 & \frac{4a^2}{1479} \\
 & \frac{2a(A+5iB)}{d\sqrt{\cot(c+dx)}} - \frac{-B+iA}{2d\sqrt{\cot(c+dx)}(a\cot(c+dx)+ia)} \\
 & \frac{2a\left(\left(\frac{1}{2}+\frac{i}{2}\right)((2+i)A+(1+4i)B) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}}\right)\right)}{d} \\
 & \frac{4a^2}{25} \\
 & \frac{2a(A+5iB)}{d\sqrt{\cot(c+dx)}} - \frac{-B+iA}{2d\sqrt{\cot(c+dx)}(a\cot(c+dx)+ia)} \\
 & \frac{2a\left(\left(\frac{1}{2}+\frac{i}{2}\right)((2+i)A+(1+4i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}}\right)\right)}{d} + \left(\frac{1}{2}+\frac{i}{2}\right) \\
 & \frac{4a^2}{25}
 \end{aligned}$$

3.523. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

$$\begin{array}{c}
\downarrow 27 \\
\frac{-B + iA}{2d\sqrt{\cot(c + dx)}(a\cot(c + dx) + ia)} - \\
\frac{2a\left(\left(\frac{1}{2} + \frac{i}{2}\right)((2+i)A + (1+4i)B)\left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}\right) + \frac{2a(A+5iB)}{d\sqrt{\cot(c+dx)}}}{4a^2} \\
\downarrow 1103 \\
\frac{-B + iA}{2d\sqrt{\cot(c + dx)}(a\cot(c + dx) + ia)} - \\
\frac{2a\left(\left(\frac{1}{2} + \frac{i}{2}\right)((1+2i)A - (4+i)B)\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}}\right) + \left(\frac{1}{2} + \frac{i}{2}\right)((2+i)A + (1+4i)B)\left(\frac{\log(\cot(c+dx))}{d}\right) + \frac{2a(A+5iB)}{d\sqrt{\cot(c+dx)}}\right)}{4a^2}
\end{array}$$

input `Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])),x]`

output `(I*A - B)/(2*d*Sqrt[Cot[c + d*x]]*(I*a + a*Cot[c + d*x])) - ((2*a*(A + (5*I)*B))/(d*Sqrt[Cot[c + d*x]]) - (2*a*((1/2 + I/2)*((1 + 2*I)*A - (4 + I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + (1/2 + I/2)*((2 + I)*A + (1 + 4*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/d)/(4*a^2)`

3.523.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.523.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.52

method	result	size
derivativedivides	$\frac{-\frac{2iB}{\sqrt{\cot(dx+c)}} + \frac{4\left(\frac{A}{4} - \frac{iB}{4}\right) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} + \frac{i\left(\frac{i(B+A)\sqrt{\cot(dx+c)}}{i+\cot(dx+c)} + \frac{4(iA-2B) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}\right)}{2}}{ad}$	143
default	$\frac{-\frac{2iB}{\sqrt{\cot(dx+c)}} + \frac{4\left(\frac{A}{4} - \frac{iB}{4}\right) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} + \frac{i\left(\frac{i(B+A)\sqrt{\cot(dx+c)}}{i+\cot(dx+c)} + \frac{4(iA-2B) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}\right)}{2}}{ad}$	143

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c)),x,method=_RETURNV ERBOSE)`

$$3.523. \int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} dx$$

output $1/a/d*(-2*I*B/\cot(d*x+c)^{(1/2)}+4*(1/4*A-1/4*I*B)/(2^{(1/2)}-I*2^{(1/2)})*\arctan(2*\cot(d*x+c)^{(1/2)}/(2^{(1/2)}-I*2^{(1/2)}))+1/2*I*(I*(A+I*B)*\cot(d*x+c)^{(1/2)})/(I+\cot(d*x+c))+4*(I*A-2*B)/(2^{(1/2)}+I*2^{(1/2)})*\arctan(2*\cot(d*x+c)^{(1/2)}/(2^{(1/2)}+I*2^{(1/2)}))$

3.523.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 700 vs. $2(199) = 398$.

Time = 0.27 (sec) , antiderivative size = 700, normalized size of antiderivative = 2.54

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx =$$

$$\frac{(ade^{(4i dx + 4i c)} + ade^{(2i dx + 2i c)}) \sqrt{\frac{-i A^2 - 2 AB + i B^2}{a^2 d^2}} \log \left(- \frac{2 \left((ade^{(2i dx + 2i c)} - ad) \sqrt{\frac{i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} - 1}} \sqrt{\frac{-i A^2 - 2 AB + i B^2}{a^2 d^2}} + (A + B) \right)}{i A + B} \right)}{}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c)),x, algorithm="fracas")`

output $-1/8*((a*d*e^{(4*I*d*x + 4*I*c)} + a*d*e^{(2*I*d*x + 2*I*c)})*\sqrt{((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2))}*\log(-2*((a*d*e^{(2*I*d*x + 2*I*c)} - a*d)*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))*\sqrt{((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2))} + (A - I*B)*e^{(2*I*d*x + 2*I*c)}*e^{(-2*I*d*x - 2*I*c)/(I*A + B)} - (a*d*e^{(4*I*d*x + 4*I*c)} + a*d*e^{(2*I*d*x + 2*I*c)})*\sqrt{((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2))}*\log(2*((a*d*e^{(2*I*d*x + 2*I*c)} - a*d)*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))*\sqrt{((-I*A^2 - 2*A*B + I*B^2)/(a^2*d^2))} - (A - I*B)*e^{(2*I*d*x + 2*I*c)}*e^{(-2*I*d*x - 2*I*c)/(I*A + B)} + 2*(a*d*e^{(4*I*d*x + 4*I*c)} + a*d*e^{(2*I*d*x + 2*I*c)})*\sqrt{((I*A^2 - 4*A*B - 4*I*B^2)/(a^2*d^2))}*\log(-((a*d*e^{(2*I*d*x + 2*I*c)} - a*d)*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))*\sqrt{((I*A^2 - 4*A*B - 4*I*B^2)/(a^2*d^2))} + A + 2*I*B)*e^{(-2*I*d*x - 2*I*c)/(a*d)} - 2*(a*d*e^{(4*I*d*x + 4*I*c)} + a*d*e^{(2*I*d*x + 2*I*c)})*\sqrt{((I*A^2 - 4*A*B - 4*I*B^2)/(a^2*d^2))}*\log(((a*d*e^{(2*I*d*x + 2*I*c)} - a*d)*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))*\sqrt{((I*A^2 - 4*A*B - 4*I*B^2)/(a^2*d^2))} - A - 2*I*B)*e^{(-2*I*d*x - 2*I*c)/(a*d)} - 2*((I*A - 9*B)*e^{(4*I*d*x + 4*I*c)} + 8*B*e^{(2*I*d*x + 2*I*c)} - I*A + B)*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)))/(a*d*e^{(4*I*d*x + 4*I*c)} + a*d*e^{(2*I*d*x + 2*I*c)}))$

3.523. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

3.523.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$= -\frac{i \left(\int \frac{A}{\tan(c+dx) \cot^{\frac{3}{2}}(c+dx) - i \cot^{\frac{3}{2}}(c+dx)} dx + \int \frac{B \tan(c+dx)}{\tan(c+dx) \cot^{\frac{3}{2}}(c+dx) - i \cot^{\frac{3}{2}}(c+dx)} dx \right)}{a}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(3/2)/(a+I*a*tan(d*x+c)),x)`

output `-I*(Integral(A/(tan(c + d*x)*cot(c + d*x)**(3/2) - I*cot(c + d*x)**(3/2)), x) + Integral(B*tan(c + d*x)/(tan(c + d*x)*cot(c + d*x)**(3/2) - I*cot(c + d*x)**(3/2)), x))/a`

3.523.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.523.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a) \cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)*cot(d*x + c)^(3/2)), x)`

3.523.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{\frac{3}{2}} (a + a \tan(c + dx) \text{ li})} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + a*tan(c + d*x)*li)),x)`output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + a*tan(c + d*x)*li)), x)`

$$3.524 \quad \int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$$

3.524.1 Optimal result	4929
3.524.2 Mathematica [A] (verified)	4930
3.524.3 Rubi [A] (verified)	4930
3.524.4 Maple [A] (verified)	4936
3.524.5 Fricas [B] (verification not implemented)	4937
3.524.6 Sympy [F]	4938
3.524.7 Maxima [F(-2)]	4938
3.524.8 Giac [F]	4938
3.524.9 Mupad [F(-1)]	4939

3.524.1 Optimal result

Integrand size = 36, antiderivative size = 307

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx \\ &= \frac{\left(\frac{1}{4} + \frac{i}{4}\right) \left((4 + i)A + (1 + 6i)B\right) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}ad} \\ & \quad - \frac{\left(\frac{1}{4} + \frac{i}{4}\right) \left((4 + i)A + (1 + 6i)B\right) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}ad} \\ & \quad - \frac{3A + 7iB}{6ad \cot^{\frac{3}{2}}(c + dx)} - \frac{5(iA - B)}{2ad \sqrt{\cot(c + dx)}} + \frac{iA - B}{2d \cot^{\frac{3}{2}}(c + dx)(ia + a \cot(c + dx))} \\ & \quad + \frac{\left((3 - 5i)A + (5 + 7i)B\right) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{8\sqrt{2}ad} \\ & \quad + \frac{\left(\frac{1}{8} + \frac{i}{8}\right) \left((1 + 4i)A - (6 + i)B\right) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{\sqrt{2}ad} \end{aligned}$$

output $1/6*(-3*A-7*I*B)/a/d/\cot(d*x+c)^{(3/2)}+1/2*(I*A-B)/d/\cot(d*x+c)^{(3/2)}/(I*a+a*\cot(d*x+c))- (1/8+1/8*I)*((4+I)*A+(1+6*I)*B)*\arctan(-1+2^{(1/2)*\cot(d*x+c)^{(1/2)})}/a/d*2^{(1/2)}-(1/8+1/8*I)*((4+I)*A+(1+6*I)*B)*\arctan(1+2^{(1/2)*\cot(d*x+c)^{(1/2)})}/a/d*2^{(1/2)}+1/16*((3-5*I)*A+(5+7*I)*B)*\ln(1+\cot(d*x+c)-2^{(1/2)*\cot(d*x+c)^{(1/2)})}/a/d*2^{(1/2)}+(1/16+1/16*I)*((1+4*I)*A-(6+I)*B)*\ln(1+\cot(d*x+c)+2^{(1/2)*\cot(d*x+c)^{(1/2)})}/a/d*2^{(1/2)}-5/2*(I*A-B)/a/d/\cot(d*x+c)^{(1/2)}$

$$3.524. \quad \int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$$

3.524.2 Mathematica [A] (verified)

Time = 5.57 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.58

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$= \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(-3(-1)^{3/4}(A - iB) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right) (-i + \tan(c + dx)) + 6(-1)^{1/4}((-2i)A + 3B) \operatorname{ArcTanh} \left[(-1)^{3/4} \sqrt{\tan(c + dx)} \right] (-i + \tan(c + dx)) + \sqrt{\tan(c + dx)} (-15(A + iB) + 4((-3i)A + 2B) \tan(c + dx) - (4i)B \tan^2(c + dx)) \right)}{6a d (-i + \tan(c + dx))}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-3*(-1)^(3/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*(-I + Tan[c + d*x]) + 6*(-1)^(1/4)*((-2*I)*A + 3*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*(-I + Tan[c + d*x]) + Sqrt[Tan[c + d*x]]*(-15*(A + I*B) + 4*((-3*I)*A + 2*B)*Tan[c + d*x] - (4*I)*B*Tan[c + d*x]^2)))/(6*a*d*(-I + Tan[c + d*x]))`

3.524.3 Rubi [A] (verified)Time = 1.04 (sec) , antiderivative size = 267, normalized size of antiderivative = 0.87, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.611$, Rules used = {3042, 4064, 3042, 4079, 27, 3042, 4012, 3042, 4012, 25, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{5/2}(a + ia \tan(c + dx))} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{A \cot(c + dx) + B}{\cot^{\frac{5}{2}}(c + dx)(a \cot(c + dx) + ia)} dx$$

$$\downarrow \text{3042}$$

3.524. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

$$\begin{aligned}
& \int \frac{B - A \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{5/2} (-a \tan(c + dx + \frac{\pi}{2}) + ia)} dx \\
& \quad \downarrow 4079 \\
& \frac{\int -\frac{a(3A+7iB)+5a(iA-B)\cot(c+dx)}{2\cot^{\frac{5}{2}}(c+dx)} dx}{2a^2} + \frac{-B + iA}{2d\cot^{\frac{3}{2}}(c + dx)(a\cot(c + dx) + ia)} \\
& \quad \downarrow 27 \\
& \frac{-B + iA}{2d\cot^{\frac{3}{2}}(c + dx)(a\cot(c + dx) + ia)} - \frac{\int \frac{a(3A+7iB)+5a(iA-B)\cot(c+dx)}{\cot^{\frac{5}{2}}(c+dx)} dx}{4a^2} \\
& \quad \downarrow 3042 \\
& \frac{-B + iA}{2d\cot^{\frac{3}{2}}(c + dx)(a\cot(c + dx) + ia)} - \frac{\int \frac{a(3A+7iB)-5a(iA-B)\tan(c+dx+\frac{\pi}{2})}{(-\tan(c+dx+\frac{\pi}{2}))^{5/2}} dx}{4a^2} \\
& \quad \downarrow 4012 \\
& \frac{-B + iA}{2d\cot^{\frac{3}{2}}(c + dx)(a\cot(c + dx) + ia)} - \frac{\int \frac{5a(iA-B)-a(3A+7iB)\cot(c+dx)}{\cot^{\frac{3}{2}}(c+dx)} dx + \frac{2a(3A+7iB)}{3d\cot^{\frac{3}{2}}(c+dx)}}{4a^2} \\
& \quad \downarrow 3042 \\
& \frac{-B + iA}{2d\cot^{\frac{3}{2}}(c + dx)(a\cot(c + dx) + ia)} - \frac{\int \frac{5a(iA-B)+a(3A+7iB)\tan(c+dx+\frac{\pi}{2})}{(-\tan(c+dx+\frac{\pi}{2}))^{3/2}} dx + \frac{2a(3A+7iB)}{3d\cot^{\frac{3}{2}}(c+dx)}}{4a^2} \\
& \quad \downarrow 4012 \\
& \frac{-B + iA}{2d\cot^{\frac{3}{2}}(c + dx)(a\cot(c + dx) + ia)} - \frac{\int -\frac{a(3A+7iB)+5a(iA-B)\cot(c+dx)}{\sqrt{\cot(c+dx)}} dx + \frac{2a(3A+7iB)}{3d\cot^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\cot(c+dx)}}}{4a^2} \\
& \quad \downarrow 25 \\
& \frac{-B + iA}{2d\cot^{\frac{3}{2}}(c + dx)(a\cot(c + dx) + ia)} - \frac{\int \frac{a(3A+7iB)+5a(iA-B)\cot(c+dx)}{\sqrt{\cot(c+dx)}} dx + \frac{2a(3A+7iB)}{3d\cot^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\cot(c+dx)}}}{4a^2} \\
& \quad \downarrow 3042
\end{aligned}$$

3.524. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

$$\begin{array}{c}
 \frac{-B + iA}{2d \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + ia)} \\
 - \int \frac{a(3A+7iB) - 5a(iA-B) \tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx + \frac{2a(3A+7iB)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\cot(c+dx)}} \\
 \hline
 4a^2 \\
 \downarrow 4017 \\
 \frac{-B + iA}{2d \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + ia)} \\
 - \frac{2 \int -\frac{a(3A+7iB+5(iA-B) \cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} + \frac{2a(3A+7iB)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\cot(c+dx)}} \\
 \hline
 4a^2 \\
 \downarrow 25 \\
 \frac{-B + iA}{2d \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + ia)} \\
 2 \int \frac{a(3A+7iB+5(iA-B) \cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{2a(3A+7iB)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\cot(c+dx)}} \\
 \hline
 4a^2 \\
 \downarrow 27 \\
 \frac{-B + iA}{2a \int \frac{3A+7iB+5(iA-B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}} + \frac{2a(3A+7iB)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\cot(c+dx)}} \\
 \hline
 4a^2 \\
 \downarrow 1482 \\
 \frac{-B + iA}{2a \left(\frac{1}{2}((3-5i)A+(5+7i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \left(\frac{1}{2}+\frac{i}{2}\right)((4+i)A+(1+6i)B) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)} + \frac{2a(3A+7iB)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{10a(-B+iA)}{d\sqrt{\cot(c+dx)}} \\
 \hline
 4a^2 \\
 \downarrow 1476 \\
 \frac{-B + iA}{2a \left(\frac{1}{2}((3-5i)A+(5+7i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \left(\frac{1}{2}+\frac{i}{2}\right)((4+i)A+(1+6i)B) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)} \\
 \hline
 4a^2 \\
 \downarrow 1082
 \end{array}$$

3.524. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

$$\frac{-B + iA}{2d \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + ia)} - \frac{2a \left(\frac{1}{2}((3-5i)A + (5+7i)B) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \left(\frac{1}{2} + \frac{i}{2}\right)((4+i)A + (1+6i)B) \left(\int \frac{1}{\cot(c+dx)-1} \frac{d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \int \frac{1}{\cot(c+dx)-1} \frac{d(\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{d}}{4a^2}$$

↓ 217

$$\frac{-B + iA}{2d \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + ia)} - \frac{2a \left(\frac{1}{2}((3-5i)A + (5+7i)B) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \left(\frac{1}{2} + \frac{i}{2}\right)((4+i)A + (1+6i)B) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\cot(c+dx)}+1}{\sqrt{2}})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{d}}{4a^2} +$$

↓ 1479

$$\frac{-B + iA}{2d \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + ia)} - \frac{2a \left(\frac{1}{2}((3-5i)A + (5+7i)B) \left(-\int \frac{\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)}+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \int \frac{\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)}+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \left(\frac{1}{2} + \frac{i}{2}\right)((4+i)A + (1+6i)B) \right)}{d}}{4a^2}$$

↓ 25

$$\frac{-B + iA}{2d \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + ia)} - \frac{2a \left(\frac{1}{2}((3-5i)A + (5+7i)B) \left(\int \frac{\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)}+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \int \frac{\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)}+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \left(\frac{1}{2} + \frac{i}{2}\right)((4+i)A + (1+6i)B) \right)}{d}}{4a^2}$$

↓ 27

$$\frac{-B + iA}{2d \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + ia)} - \frac{2a \left(\frac{1}{2}((3-5i)A + (5+7i)B) \left(\int \frac{\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)}+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\frac{\sqrt{2}\sqrt{\cot(c+dx)}+1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)}+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \left(\frac{1}{2} + \frac{i}{2}\right)((4+i)A + (1+6i)B) \right)}{d}}{4a^2}$$

↓ 1103

3.524. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

$$\frac{-B + iA}{2d \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + ia)} - \frac{2a \left(\frac{1}{2} + \frac{i}{2} \right) \left((4+i)A + (1+6i)B \right) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} \left((3-5i)A + (5+7i)B \right) \left(\frac{\log(\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right)}{4a^2}$$

input `Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])),x]`

output `(I*A - B)/(2*d*Cot[c + d*x]^(3/2)*(I*a + a*Cot[c + d*x])) - ((2*a*(3*A + (7*I)*B))/(3*d*Cot[c + d*x]^(3/2)) + (10*a*(I*A - B))/(d*Sqrt[Cot[c + d*x]]) + (2*a*((1/2 + I/2)*((4 + I)*A + (1 + 6*I)*B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + (((3 - 5*I)*A + (5 + 7*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d)/(4*a^2)`

3.524.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

3.524. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

```
rule 4079 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

3.524.4 Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.52

method	result
derivativedivides	$\frac{4\left(-\frac{iA}{4}-\frac{B}{4}\right)\arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} + \frac{-2iA+2B}{\sqrt{\cot(dx+c)}} - \frac{2iB}{3\cot(dx+c)^{\frac{3}{2}}} - \frac{i\left(-\frac{i(iA-B)\sqrt{\cot(dx+c)}}{i+\cot(dx+c)} + \frac{4(3iB+2A)\arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}\right)}{2}$
default	$\frac{4\left(-\frac{iA}{4}-\frac{B}{4}\right)\arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} + \frac{-2iA+2B}{\sqrt{\cot(dx+c)}} - \frac{2iB}{3\cot(dx+c)^{\frac{3}{2}}} - \frac{i\left(-\frac{i(iA-B)\sqrt{\cot(dx+c)}}{i+\cot(dx+c)} + \frac{4(3iB+2A)\arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}\right)}{2}$

```
input int((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output 1/a/d*(4*(-1/4*I*A-1/4*B)/(2^(1/2)-I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2
^(1/2)-I*2^(1/2)))+2*(-I*A+B)/cot(d*x+c)^(1/2)-2/3*I*B/cot(d*x+c)^(3/2)-1/
2*I*(-I*(I*A-B)*cot(d*x+c)^(1/2)/(I+cot(d*x+c))+4*(2*A+3*I*B)/(2^(1/2)+I*2
^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2))))
```

3.524.
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$$

3.524.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 794 vs. $2(222) = 444$.

Time = 0.27 (sec) , antiderivative size = 794, normalized size of antiderivative = 2.59

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx =$$

$$3 \left(ade^{(6i dx + 6i c)} + 2 ade^{(4i dx + 4i c)} + ade^{(2i dx + 2i c)} \right) \sqrt{\frac{i A^2 + 2 AB - i B^2}{a^2 d^2}} \log \left(- \frac{2 \left((i ade^{(2i dx + 2i c)} - i ad) \sqrt{\frac{i e^{(2i dx + 2i c)} + \dots}{e^{(2i dx + 2i c)} - \dots}} \right)}{\dots} \right)$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c)),x, algorithm="fricas")
```

```
output -1/24*(3*(a*d*e^(6*I*d*x + 6*I*c) + 2*a*d*e^(4*I*d*x + 4*I*c) + a*d*e^(2*I*d*x + 2*I*c))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2))*log(-2*((I*a*d*e^(2*I*d*x + 2*I*c) - I*a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I*c))e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*(a*d*e^(6*I*d*x + 6*I*c) + 2*a*d*e^(4*I*d*x + 4*I*c) + a*d*e^(2*I*d*x + 2*I*c))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2))*log(-2*((-I*a*d*e^(2*I*d*x + 2*I*c) + I*a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^2*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I*c))e^(-2*I*d*x - 2*I*c)/(I*A + B)) + 6*(a*d*e^(6*I*d*x + 6*I*c) + 2*a*d*e^(4*I*d*x + 4*I*c) + a*d*e^(2*I*d*x + 2*I*c))*sqrt((-4*I*A^2 + 12*A*B + 9*I*B^2)/(a^2*d^2))*log(-((a*d*e^(2*I*d*x + 2*I*c) - a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-4*I*A^2 + 12*A*B + 9*I*B^2)/(a^2*d^2)) + 2*I*A - 3*B)*e^(-2*I*d*x - 2*I*c)/(a*d)) - 6*(a*d*e^(6*I*d*x + 6*I*c) + 2*a*d*e^(4*I*d*x + 4*I*c) + a*d*e^(2*I*d*x + 2*I*c))*sqrt((-4*I*A^2 + 12*A*B + 9*I*B^2)/(a^2*d^2))*log(((a*d*e^(2*I*d*x + 2*I*c) - a*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-4*I*A^2 + 12*A*B + 9*I*B^2)/(a^2*d^2)) - 2*I*A + 3*B)*e^(-2*I*d*x - 2*I*c)/(a*d)) + 2*((27*A + 19*I*B)*e^(6*I*d*x + 6*I*c) + (3*A + 19*I*B)*e^(4*I*d*x + 4*I*c) - (27*A + 35*I*B)*e^(2*I*d*x + 2*I*c) - 3*A - 3*I*B)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*...
```


3.524.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx$$

$$= - \frac{i \left(\int \frac{A}{\tan(c+dx) \cot^{\frac{5}{2}}(c+dx) - i \cot^{\frac{5}{2}}(c+dx)} dx + \int \frac{B \tan(c+dx)}{\tan(c+dx) \cot^{\frac{5}{2}}(c+dx) - i \cot^{\frac{5}{2}}(c+dx)} dx \right)}{a}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(5/2)/(a+I*a*tan(d*x+c)),x)`

output `-I*(Integral(A/(tan(c + d*x)*cot(c + d*x)**(5/2) - I*cot(c + d*x)**(5/2)), x) + Integral(B*tan(c + d*x)/(tan(c + d*x)*cot(c + d*x)**(5/2) - I*cot(c + d*x)**(5/2)), x))/a`

3.524.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.524.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a) \cot(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)*cot(d*x + c)^(5/2)), x)`

3.524. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))} dx$

3.524.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{5/2} (a + a \tan(c + dx) li)} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + a*tan(c + d*x)*li)),x)`output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + a*tan(c + d*x)*li)), x)`

3.525
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.525.1 Optimal result 4940
 3.525.2 Mathematica [A] (verified) 4941
 3.525.3 Rubi [A] (verified) 4941
 3.525.4 Maple [B] (verified) 4947
 3.525.5 Fricas [B] (verification not implemented) 4948
 3.525.6 Sympy [F] 4949
 3.525.7 Maxima [F(-2)] 4950
 3.525.8 Giac [F] 4950
 3.525.9 Mupad [F(-1)] 4950

3.525.1 Optimal result

Integrand size = 36, antiderivative size = 317

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$= -\frac{\left(\frac{1}{16} - \frac{i}{16}\right) \left((2+23i)A - (7+2i)B\right) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}a^2d}$$

$$+ \frac{\left((25+21i)A - (9-5i)B\right) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{16\sqrt{2}a^2d}$$

$$- \frac{5(5A+iB)\sqrt{\cot(c+dx)}}{8a^2d} + \frac{(7A+3iB)\cot^{\frac{3}{2}}(c+dx)}{8a^2d(i+\cot(c+dx))} + \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{4d(ia+a\cot(c+dx))^2}$$

$$- \frac{\left(\frac{1}{32} - \frac{i}{32}\right) \left((23+2i)A + (2+7i)B\right) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{\sqrt{2}a^2d}$$

$$+ \frac{\left(\frac{1}{32} - \frac{i}{32}\right) \left((23+2i)A + (2+7i)B\right) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{\sqrt{2}a^2d}$$

output

```
1/8*(7*A+3*I*B)*cot(d*x+c)^(3/2)/a^2/d/(I+cot(d*x+c))+1/4*(A+I*B)*cot(d*x+c)^(5/2)/d/(I*a+a*cot(d*x+c))^2+(1/32-1/32*I)*((2+23*I)*A-(7+2*I)*B)*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/a^2/d*2^(1/2)+1/32*((25+21*I)*A+(-9+5*I)*B)*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/a^2/d*2^(1/2)+(-1/64+1/64*I)*((23+2*I)*A+(2+7*I)*B)*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/a^2/d*2^(1/2)+(1/64-1/64*I)*((23+2*I)*A+(2+7*I)*B)*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/a^2/d*2^(1/2)-5/8*(5*A+I*B)*cot(d*x+c)^(1/2)/a^2/d
```

3.525.
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.525.2 Mathematica [A] (verified)

Time = 5.42 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.70

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx =$$

$$\frac{\sqrt{\cot(c+dx)} \left(-16A + (-1)^{3/4} (23A + 7iB) \operatorname{arctanh} \left((-1)^{3/4} \sqrt{\tan(c+dx)} \right) \sec^2(c+dx) (\cos(2(c+dx))) \right)}{a^2 d (-1 + \tan(c+dx))^2}$$

input `Integrate[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `-1/8*(Sqrt[Cot[c + d*x]]*(-16*A + (-1)^(3/4)*(23*A + (7*I)*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)])*Sqrt[Tan[c + d*x]] + 2*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*((-I)*Cos[2*(c + d*x)] + Sin[2*(c + d*x)])*Sqrt[Tan[c + d*x]] - (43*I)*A*Tan[c + d*x] + 7*B*Tan[c + d*x] + 25*A*Tan[c + d*x]^2 + (5*I)*B*Tan[c + d*x]^2)/(a^2*d*(-1 + Tan[c + d*x])^2)`

3.525.3 Rubi [A] (verified)

Time = 1.24 (sec) , antiderivative size = 286, normalized size of antiderivative = 0.90, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3042, 4064, 3042, 4078, 27, 3042, 4078, 3042, 4011, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\cot(c+dx)^{3/2}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A \cot(c+dx) + B)}{(a \cot(c+dx) + ia)^2} dx$$

3.525. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
& \int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{5/2} (B - A \tan(c+dx+\frac{\pi}{2}))}{(-a \tan(c+dx+\frac{\pi}{2}) + ia)^2} dx \\
& \quad \downarrow 3042 \\
& \int -\frac{\cot^{\frac{3}{2}}(c+dx)(5a(iA-B)-a(9A+iB)\cot(c+dx))}{2(\cot(c+dx)a+ia)} dx + \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} \\
& \quad \downarrow 4078 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \int \frac{\cot^{\frac{3}{2}}(c+dx)(5a(iA-B)-a(9A+iB)\cot(c+dx))}{\cot(c+dx)a+ia} dx \\
& \quad \downarrow 27 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{3/2} (5a(iA-B)+a(9A+iB)\tan(c+dx+\frac{\pi}{2}))}{ia-a\tan(c+dx+\frac{\pi}{2})} dx \\
& \quad \downarrow 3042 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \frac{\int \sqrt{\cot(c+dx)}(3a^2(7iA-3B)-5a^2(5A+iB)\cot(c+dx)) dx}{2a^2} - \frac{(7A+3iB)\cot^{\frac{3}{2}}(c+dx)}{d(\cot(c+dx)+i)} \\
& \quad \downarrow 4078 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \frac{\int \sqrt{-\tan(c+dx+\frac{\pi}{2})}(3(7iA-3B)a^2+5(5A+iB)\tan(c+dx+\frac{\pi}{2})a^2) dx}{2a^2} - \frac{(7A+3iB)\cot^{\frac{3}{2}}(c+dx)}{d(\cot(c+dx)+i)} \\
& \quad \downarrow 3042 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \frac{\int \frac{5(5A+iB)a^2+3(7iA-3B)\cot(c+dx)a^2}{\sqrt{\cot(c+dx)}} dx + \frac{10a^2(5A+iB)\sqrt{\cot(c+dx)}}{d}}{2a^2} - \frac{(7A+3iB)\cot^{\frac{3}{2}}(c+dx)}{d(\cot(c+dx)+i)} \\
& \quad \downarrow 4011 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \frac{\int \frac{5a^2(5A+iB)-3a^2(7iA-3B)\tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx + \frac{10a^2(5A+iB)\sqrt{\cot(c+dx)}}{d}}{2a^2} - \frac{(7A+3iB)\cot^{\frac{3}{2}}(c+dx)}{d(\cot(c+dx)+i)} \\
& \quad \downarrow 3042 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \frac{\int \frac{5a^2(5A+iB)-3a^2(7iA-3B)\tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx + \frac{10a^2(5A+iB)\sqrt{\cot(c+dx)}}{d}}{2a^2} - \frac{(7A+3iB)\cot^{\frac{3}{2}}(c+dx)}{d(\cot(c+dx)+i)} \\
& \quad \downarrow 4017
\end{aligned}$$

3.525. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$

$$\begin{aligned}
 & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \\
 & \frac{2 \int \frac{a^2(5(5A+iB)+3(7iA-3B) \cot(c+dx)) d\sqrt{\cot(c+dx)}}{\cot^2(c+dx)+1} + 10a^2(5A+iB)\sqrt{\cot(c+dx)}}{2a^2} - \frac{(7A+3iB) \cot^{\frac{3}{2}}(c+dx)}{d(\cot(c+dx)+i)} \\
 & \frac{8a^2}{25} \\
 & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \\
 & \frac{10a^2(5A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2 \int \frac{a^2(5(5A+iB)+3(7iA-3B) \cot(c+dx)) d\sqrt{\cot(c+dx)}}{\cot^2(c+dx)+1}}{2a^2} - \frac{(7A+3iB) \cot^{\frac{3}{2}}(c+dx)}{d(\cot(c+dx)+i)} \\
 & \frac{8a^2}{27} \\
 & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \\
 & \frac{10a^2(5A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a^2 \int \frac{5(5A+iB)+3(7iA-3B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{2a^2} - \frac{(7A+3iB) \cot^{\frac{3}{2}}(c+dx)}{d(\cot(c+dx)+i)} \\
 & \frac{8a^2}{1482} \\
 & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \\
 & \frac{10a^2(5A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a^2 \left(\left(\frac{1}{2} - \frac{i}{2} \right) ((23+2i)A + (2+7i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} ((25+21i)A - (9-5i)B) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{2a^2} - \frac{(7A+3iB) \cot^{\frac{3}{2}}(c+dx)}{d(\cot(c+dx)+i)} \\
 & \frac{8a^2}{1476} \\
 & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \\
 & \frac{10a^2(5A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a^2 \left(\left(\frac{1}{2} - \frac{i}{2} \right) ((23+2i)A + (2+7i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} ((25+21i)A - (9-5i)B) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{2a^2} - \frac{(7A+3iB) \cot^{\frac{3}{2}}(c+dx)}{d(\cot(c+dx)+i)} \\
 & \frac{8a^2}{1082} \\
 & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \\
 & \frac{10a^2(5A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a^2 \left(\left(\frac{1}{2} - \frac{i}{2} \right) ((23+2i)A + (2+7i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} ((25+21i)A - (9-5i)B) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{2a^2} - \frac{(7A+3iB) \cot^{\frac{3}{2}}(c+dx)}{d(\cot(c+dx)+i)} \\
 & \frac{8a^2}{217}
 \end{aligned}$$

3.525. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \frac{10a^2(5A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a^2 \left(\left(\frac{1}{2} - \frac{i}{2}\right) ((23+2i)A + (2+7i)B) \int \frac{1 - \cot^{\frac{5}{2}}(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} ((25+21i)A - (9-5i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{2a^2 d}$$

$$8a^2$$

↓ 1479

$$\frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \frac{10a^2(5A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a^2 \left(\left(\frac{1}{2} - \frac{i}{2}\right) ((23+2i)A + (2+7i)B) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) \right)}{2a^2 d}$$

$$8a^2$$

↓ 25

$$\frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \frac{10a^2(5A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a^2 \left(\left(\frac{1}{2} - \frac{i}{2}\right) ((23+2i)A + (2+7i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) \right)}{2a^2 d}$$

$$8a^2$$

↓ 27

$$\frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \frac{10a^2(5A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a^2 \left(\left(\frac{1}{2} - \frac{i}{2}\right) ((23+2i)A + (2+7i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{2a^2 d}$$

$$8a^2$$

↓ 1103

$$\frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \frac{10a^2(5A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a^2 \left(\frac{1}{2} ((25+21i)A - (9-5i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \left(\frac{1}{2} - \frac{i}{2}\right) ((23+2i)A + (2+7i)B) \left(\frac{\log(\cot(c+dx))}{\sqrt{2}} \right) \right)}{2a^2 d}$$

$$8a^2$$

input `Int[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

3.525. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

```
output ((A + I*B)*Cot[c + d*x]^(5/2))/(4*d*(I*a + a*Cot[c + d*x])^2) - (-(((7*A +
(3*I)*B)*Cot[c + d*x]^(3/2))/(d*(I + Cot[c + d*x]))) + ((10*a^2*(5*A + I*
B)*Sqrt[Cot[c + d*x]])/d - (2*a^2*(((25 + 21*I)*A - (9 - 5*I)*B)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]))/2 + (1/2 - I/2)*((23 + 2*I)*A + (2 + 7*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2]))))/d)/(2*a^2))/(8*a^2)
```

3.525.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]
```

```
rule 1103 Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

```
rule 1476 Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```


rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^m*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^p*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^m*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^n, x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

```
rule 4078 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.525.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 761 vs. $2(261) = 522$.

Time = 0.41 (sec) , antiderivative size = 762, normalized size of antiderivative = 2.40

method	result	size
derivativedivides	Expression too large to display	762
default	Expression too large to display	762

```
input int(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x,method=_RETUR
NVERBOSE)
```

output
$$-1/16/a^2/d*(1/\tan(dx+c))^{3/2}*\tan(dx+c)*(23*I*A*\tan(dx+c)^{5/2}*\arctan((1/2+1/2*I)*\tan(dx+c)^{1/2}*2^{1/2})-2*I*A*\tan(dx+c)^{5/2}*\arctan((1/2-1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))+7*I*B*\tan(dx+c)^{5/2}*\arctan((1/2+1/2*I)*\tan(dx+c)^{1/2}*2^{1/2})-2*I*B*\tan(dx+c)^{5/2}*\arctan((1/2-1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))-46*I*A*\tan(dx+c)^{3/2}*\arctan((1/2+1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))-4*I*A*\tan(dx+c)^{3/2}*\arctan((1/2-1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))+14*I*B*\tan(dx+c)^{3/2}*\arctan((1/2+1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))+4*I*B*\tan(dx+c)^{3/2}*\arctan((1/2-1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))-23*I*A*\tan(dx+c)^{1/2}*\arctan((1/2+1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))+2*I*A*\tan(dx+c)^{1/2}*\arctan((1/2-1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))-7*I*B*\tan(dx+c)^{1/2}*\arctan((1/2+1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))+2*I*B*\tan(dx+c)^{1/2}*\arctan((1/2-1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))+5*I*B*\tan(dx+c)^2*2^{1/2}-43*I*A*\tan(dx+c)*2^{1/2}+7*B*\tan(dx+c)*2^{1/2}+25*A*\tan(dx+c)^2*2^{1/2}-16*A*2^{1/2}+2*A*\tan(dx+c)^{5/2}*\arctan((1/2-1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))+23*A*\tan(dx+c)^{5/2}*\arctan((1/2+1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))-2*B*\tan(dx+c)^{5/2}*\arctan((1/2-1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))-7*B*\tan(dx+c)^{5/2}*\arctan((1/2+1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))-4*A*\tan(dx+c)^{3/2}*\arctan((1/2-1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))+46*A*\tan(dx+c)^{3/2}*\arctan((1/2+1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))-4*B*\tan(dx+c)^{3/2}*\arctan((1/2-1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))+14*B*\tan(dx+c)^{3/2}*\arctan((1/2+1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))-14*B*\tan(dx+c)^{3/2}*\arctan((1/2-1/2*I)*\tan(dx+c)^{1/2}*2^{1/2}))$$

3.525.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 669 vs. $2(234) = 468$.

Time = 0.27 (sec) , antiderivative size = 669, normalized size of antiderivative = 2.11

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \frac{\left(2a^2d\sqrt{\frac{iA^2+2AB-iB^2}{a^4d^2}}e^{(4idx+4ic)}\log\left(-\frac{2\left((ia^2de^{(2idx+2ic)}-ia^2d)\sqrt{\frac{ie^{(2idx+2ic)}+i}{e^{(2idx+2ic)}-1}}\sqrt{\frac{iA^2+2AB-iB^2}{a^4d^2}}+(A-iB)e^{(2idx+2ic)}\right)}{iA+B}\right)}{iA+B}\right)}{iA+B}$$

input `integrate(cot(dx+c)^(3/2)*(A+B*tan(dx+c))/(a+I*a*tan(dx+c))^2,x, algorithm="fracas")`

```

output -1/32*(2*a^2*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)
*log(-2*((I*a^2*d*e^(2*I*d*x + 2*I*c) - I*a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c)
+ I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))
+ (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 2*a^2*
d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(-2*((-I*
a^2*d*e^(2*I*d*x + 2*I*c) + I*a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(
2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2)) + (A - I*B)
*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - a^2*d*sqrt((-529*I
*A^2 + 322*A*B + 49*I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(1/8*((a^2*d*
e^(2*I*d*x + 2*I*c) - a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x
+ 2*I*c) - 1))*sqrt((-529*I*A^2 + 322*A*B + 49*I*B^2)/(a^4*d^2)) + 23*I*A
- 7*B)*e^(-2*I*d*x - 2*I*c)/(a^2*d)) + a^2*d*sqrt((-529*I*A^2 + 322*A*B +
49*I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(-1/8*((a^2*d*e^(2*I*d*x + 2*I
*c) - a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*s
qrt((-529*I*A^2 + 322*A*B + 49*I*B^2)/(a^4*d^2)) - 23*I*A + 7*B)*e^(-2*I*d
*x - 2*I*c)/(a^2*d)) + 2*(6*(7*A + I*B)*e^(4*I*d*x + 4*I*c) - (9*A + 5*I*B
)*e^(2*I*d*x + 2*I*c) - A - I*B)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*
d*x + 2*I*c) - 1))*e^(-4*I*d*x - 4*I*c)/(a^2*d)

```

3.525.6 Sympy [F]

$$\begin{aligned}
 & \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx \\
 &= -\frac{\int \frac{A \cot^{\frac{3}{2}}(c+dx)}{\tan^2(c+dx)-2i \tan(c+dx)-1} dx + \int \frac{B \tan(c+dx) \cot^{\frac{3}{2}}(c+dx)}{\tan^2(c+dx)-2i \tan(c+dx)-1} dx}{a^2}
 \end{aligned}$$

```

input integrate(cot(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**2,x)

```

```

output -(Integral(A*cot(c + d*x)**(3/2)/(tan(c + d*x)**2 - 2*I*tan(c + d*x) - 1),
x) + Integral(B*tan(c + d*x)*cot(c + d*x)**(3/2)/(tan(c + d*x)**2 - 2*I*t
an(c + d*x) - 1), x))/a**2

```

3.525.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.525.8 Giac [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{(ia\tan(dx+c)+a)^2} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")`

output `integrate((B*tan(d*x+c)+A)*cot(d*x+c)^(3/2)/(I*a*tan(d*x+c)+a)^2, x)`

3.525.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \int \frac{\cot(c+dx)^{\frac{3}{2}}(A+B\tan(c+dx))}{(a+a\tan(c+dx)1i)^2} dx$$

input `int((cot(c+d*x)^(3/2)*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*1i)^2,x)`

output `int((cot(c+d*x)^(3/2)*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*1i)^2, x)`

3.525. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$

3.526
$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

3.526.1 Optimal result 4951
 3.526.2 Mathematica [A] (verified) 4952
 3.526.3 Rubi [A] (verified) 4952
 3.526.4 Maple [B] (verified) 4957
 3.526.5 Fricas [B] (verification not implemented) 4958
 3.526.6 Sympy [F] 4959
 3.526.7 Maxima [F(-2)] 4960
 3.526.8 Giac [F] 4960
 3.526.9 Mupad [F(-1)] 4960

3.526.1 Optimal result

Integrand size = 36, antiderivative size = 284

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$= \frac{((9-5i)A+(1-3i)B) \arctan\left(1-\sqrt{2}\sqrt{\cot(c+dx)}\right)}{16\sqrt{2}a^2d}$$

$$+ \frac{\left(\frac{1}{16}+\frac{i}{16}\right)((-2+7i)A+(1+2i)B) \arctan\left(1+\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}a^2d}$$

$$+ \frac{(5A+iB)\sqrt{\cot(c+dx)}}{8a^2d(i+\cot(c+dx))} + \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{4d(ia+a \cot(c+dx))^2}$$

$$+ \frac{\left(\frac{1}{32}+\frac{i}{32}\right)((-7+2i)A+(2+i)B) \log\left(1-\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{\sqrt{2}a^2d}$$

$$+ \frac{((9+5i)A-(1+3i)B) \log\left(1+\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{32\sqrt{2}a^2d}$$

```
output 1/4*(A+I*B)*cot(d*x+c)^(3/2)/d/(I*a+a*cot(d*x+c))^2-1/32*((9-5*I)*A+(1-3*I)
)*B)*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/a^2/d*2^(1/2)+(1/32+1/32*I)*((-2+
7*I)*A+(1+2*I)*B)*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/a^2/d*2^(1/2)+(1/64+1
/64*I)*((-7+2*I)*A+(2+I)*B)*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/a^2/
d*2^(1/2)+1/64*((9+5*I)*A-(1+3*I)*B)*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1
/2))/a^2/d*2^(1/2)+1/8*(5*A+I*B)*cot(d*x+c)^(1/2)/a^2/d/(I+cot(d*x+c))
```

3.526.2 Mathematica [A] (verified)

Time = 4.20 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.70

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= \frac{\sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}\left(2\sqrt[4]{-1}(A-iB)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)\sec^2(c+dx)(\cos(2(c+dx)))\right)}{}$$

input `Integrate[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])2,x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(2*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)]) + (-1)^(1/4)*(7*A - I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)]) + Sqrt[Tan[c + d*x]]*(-7*A - (3*I)*B + ((-5*I)*A + B)*Tan[c + d*x]))/(8*a^2*d*(-I + Tan[c + d*x])^2)`

3.526.3 Rubi [A] (verified)Time = 1.00 (sec) , antiderivative size = 248, normalized size of antiderivative = 0.87, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.528$, Rules used = {3042, 4064, 3042, 4078, 27, 3042, 4078, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A\cot(c+dx)+B)}{(a\cot(c+dx)+ia)^2} dx$$

$$\begin{aligned}
& \int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{3/2} (B - A \tan(c+dx+\frac{\pi}{2}))}{(-a \tan(c+dx+\frac{\pi}{2}) + ia)^2} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{-\sqrt{\cot(c+dx)}(3a(iA-B)-a(7A-iB)\cot(c+dx))}{2(\cot(c+dx)a+ia)} dx + \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} \\
& \quad \downarrow \text{4078} \\
& \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \frac{\int \frac{\sqrt{\cot(c+dx)}(3a(iA-B)-a(7A-iB)\cot(c+dx))}{\cot(c+dx)a+ia} dx}{8a^2} \\
& \quad \downarrow \text{27} \\
& \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \frac{\int \frac{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(3a(iA-B)+a(7A-iB)\tan(c+dx+\frac{\pi}{2}))}{ia-a\tan(c+dx+\frac{\pi}{2})} dx}{8a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \frac{\int \frac{a^2(5iA-B)-3a^2(3A-iB)\cot(c+dx)}{\sqrt{\cot(c+dx)}} dx}{2a^2} - \frac{(5A+iB)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} \\
& \quad \downarrow \text{4078} \\
& \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \frac{\int \frac{(5iA-B)a^2+3(3A-iB)\tan(c+dx+\frac{\pi}{2})a^2}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{2a^2} - \frac{(5A+iB)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} \\
& \quad \downarrow \text{3042} \\
& \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \frac{\int \frac{a^2(5iA-B-3(3A-iB)\cot(c+dx))d\sqrt{\cot(c+dx)}}{\cot^2(c+dx)+1}}{a^2d} - \frac{(5A+iB)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} \\
& \quad \downarrow \text{4017} \\
& \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \frac{\int \frac{a^2(5iA-B-3(3A-iB)\cot(c+dx))d\sqrt{\cot(c+dx)}}{\cot^2(c+dx)+1}}{a^2d} - \frac{(5A+iB)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} \\
& \quad \downarrow \text{25} \\
& \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \frac{\int \frac{5iA-B-3(3A-iB)\cot(c+dx)}{\cot^2(c+dx)+1}d\sqrt{\cot(c+dx)}}{d} - \frac{(5A+iB)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} \\
& \quad \downarrow \text{27} \\
& \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{4d(a\cot(c+dx)+ia)^2} - \frac{\int \frac{5iA-B-3(3A-iB)\cot(c+dx)}{\cot^2(c+dx)+1}d\sqrt{\cot(c+dx)}}{d} - \frac{(5A+iB)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)}
\end{aligned}$$

3.526. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((9+5i)A - (1+3i)B) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx) + (\frac{1}{2} + \frac{i}{2})} + ((1+2i)B - (2-7i)A) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} - \frac{(5A+iB)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)}}{8a^2}$$

$$\frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((9+5i)A - (1+3i)B) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx) + (\frac{1}{2} + \frac{i}{2})} + ((1+2i)B - (2-7i)A) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right)}{d}}{8a^2}$$

$$\frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((9+5i)A - (1+3i)B) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx) + (\frac{1}{2} + \frac{i}{2})} + ((1+2i)B - (2-7i)A) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)-1} d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right)}{d}}{8a^2}$$

$$\frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((9+5i)A - (1+3i)B) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx) + (\frac{1}{2} + \frac{i}{2})} + ((1+2i)B - (2-7i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right)}{d}}{8a^2}$$

$$\frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((9+5i)A - (1+3i)B) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + (\frac{1}{2} + \frac{i}{2})((1+2i)B - (2-7i)A)}{d}}{8a^2}$$

$$\frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((9+5i)A - (1+3i)B) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx) + (\frac{1}{2} + \frac{i}{2})} + ((1+2i)B - (2-7i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right)}{d}}{8a^2}$$

3.526. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$

$$\frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((9+5i)A - (1+3i)B) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + (\frac{1}{2} + \frac{i}{2})((1+2i)B - (2-7i)A)}{d}}{8a^2}$$

↓ 27

$$\frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((9+5i)A - (1+3i)B) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + (\frac{1}{2} + \frac{i}{2})((1+2i)B - (2-7i)A)}{d}}{8a^2}$$

↓ 1103

$$\frac{\frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{4d(a \cot(c + dx) + ia)^2} - \frac{(\frac{1}{2} + \frac{i}{2})((1+2i)B - (2-7i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}((9+5i)A - (1+3i)B) \left(\frac{\log(\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right)}{d}}{8a^2}$$

input `Int[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^2,x]`

output `((A + I*B)*Cot[c + d*x]^(3/2))/(4*d*(I*a + a*Cot[c + d*x])^2) - (-(((5*A + I*B)*Sqrt[Cot[c + d*x]])/(d*(I + Cot[c + d*x]))) - ((1/2 + I/2)*((-2 + 7*I)*A + (1 + 2*I)*B)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + (((9 + 5*I)*A - (1 + 3*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d)/(8*a^2)`

3.526.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] & & NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.526.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 680 vs. $2(235) = 470$.

Time = 0.42 (sec) , antiderivative size = 681, normalized size of antiderivative = 2.40

method	result	size
derivativedivides	Expression too large to display	681
default	Expression too large to display	681

input `int(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output

```

-1/16/a^2/d*(1/tan(d*x+c))^(1/2)*tan(d*x+c)^(1/2)*(-B*arctan((1/2+1/2*I)*t
an(d*x+c)^(1/2)*2^(1/2))+2*A*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))+
2*B*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))+7*A*arctan((1/2+1/2*I)*ta
n(d*x+c)^(1/2)*2^(1/2))-7*I*A*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))
+2*I*A*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))-2*I*B*arctan((1/2-1/2*
I)*tan(d*x+c)^(1/2)*2^(1/2))+7*A*tan(d*x+c)^(1/2)*2^(1/2)-I*B*arctan((1/2+
1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))+2*B*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2
^(1/2))*tan(d*x+c)+B*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))*tan(d*x+
c)^2-B*tan(d*x+c)^(3/2)*2^(1/2)-7*A*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2
^(1/2))*tan(d*x+c)^2-2*A*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))*tan(d
*x+c)^2-2*B*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))*tan(d*x+c)^2+14*A
*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))*tan(d*x+c)-4*A*arctan((1/2-1
/2*I)*tan(d*x+c)^(1/2)*2^(1/2))*tan(d*x+c)+4*B*arctan((1/2-1/2*I)*tan(d*x+
c)^(1/2)*2^(1/2))*tan(d*x+c)+5*I*A*tan(d*x+c)^(3/2)*2^(1/2)+7*I*A*arctan((
1/2+1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))*tan(d*x+c)^2-2*I*A*arctan((1/2-1/2*I)
*tan(d*x+c)^(1/2)*2^(1/2))*tan(d*x+c)^2+2*I*B*arctan((1/2-1/2*I)*tan(d*x+c
)^(1/2)*2^(1/2))*tan(d*x+c)^2+14*I*A*arctan((1/2+1/2*I)*tan(d*x+c)^(1/2)*2
^(1/2))*tan(d*x+c)+4*I*A*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(1/2))*tan(
d*x+c)+3*I*B*tan(d*x+c)^(1/2)*2^(1/2)-2*I*B*arctan((1/2+1/2*I)*tan(d*x+c)
^(1/2)*2^(1/2))*tan(d*x+c)+4*I*B*arctan((1/2-1/2*I)*tan(d*x+c)^(1/2)*2^(...

```

3.526.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 666 vs. $2(211) = 422$.

Time = 0.26 (sec) , antiderivative size = 666, normalized size of antiderivative = 2.35

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$$

$$= \left(2a^2d\sqrt{\frac{-iA^2-2AB+iB^2}{a^4d^2}}e^{(4idx+4ic)} \log \left(-\frac{2\left((a^2de^{(2idx+2ic)}-a^2d)\sqrt{\frac{ie^{(2idx+2ic)}+i}{e^{(2idx+2ic)}-1}}\sqrt{\frac{-iA^2-2AB+iB^2}{a^4d^2}}+(A-iB)e^{(2idx+2ic)}\right)}{iA+B} \right) \right)$$

input

```

integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algo
rithm="fricas")

```

3.526. $\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx$

output `1/32*(2*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)
log(-2((a^2*d*e^(2*I*d*x + 2*I*c) - a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c) +
I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2)) +
(A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 2*a^2*d*s
qrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(2*((a^2*d*
e^(2*I*d*x + 2*I*c) - a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x
+ 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2)) - (A - I*B)*e^(2*I
*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - a^2*d*sqrt((49*I*A^2 + 14
*A*B - I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(-1/8*((a^2*d*e^(2*I*d*x +
2*I*c) - a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1
))*sqrt((49*I*A^2 + 14*A*B - I*B^2)/(a^4*d^2)) + 7*A - I*B)*e^(-2*I*d*x -
2*I*c)/(a^2*d)) + a^2*d*sqrt((49*I*A^2 + 14*A*B - I*B^2)/(a^4*d^2))*e^(4*I
*d*x + 4*I*c)*log(1/8*((a^2*d*e^(2*I*d*x + 2*I*c) - a^2*d)*sqrt((I*e^(2*I*
d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((49*I*A^2 + 14*A*B - I*B
^2)/(a^4*d^2)) - 7*A + I*B)*e^(-2*I*d*x - 2*I*c)/(a^2*d)) - 2*(2*(3*I*A -
B)*e^(4*I*d*x + 4*I*c) - (5*I*A - B)*e^(2*I*d*x + 2*I*c) - I*A + B)*sqrt((
I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*e^(-4*I*d*x - 4*I*c
)/(a^2*d)`

3.526.6 Sympy [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^2} dx$$

$$= -\frac{\int \frac{A\sqrt{\cot(c+dx)}}{\tan^2(c+dx)-2i \tan(c+dx)-1} dx + \int \frac{B \tan(c+dx)\sqrt{\cot(c+dx)}}{\tan^2(c+dx)-2i \tan(c+dx)-1} dx}{a^2}$$

input `integrate(cot(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**2,x)`

output `-(Integral(A*sqrt(cot(c + d*x))/(tan(c + d*x)**2 - 2*I*tan(c + d*x) - 1),
x) + Integral(B*tan(c + d*x)*sqrt(cot(c + d*x))/(tan(c + d*x)**2 - 2*I*tan
(c + d*x) - 1), x))/a**2`

3.526.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.526.8 Giac [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \int \frac{(B\tan(dx+c)+A)\sqrt{\cot(dx+c)}}{(ia\tan(dx+c)+a)^2} dx$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")`

output `integrate((B*tan(d*x+c)+A)*sqrt(cot(d*x+c))/(I*a*tan(d*x+c)+a)^2, x)`

3.526.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^2} dx = \int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+a\tan(c+dx)li)^2} dx$$

input `int((cot(c+d*x)^(1/2)*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*li)^2,x)`

output `int((cot(c+d*x)^(1/2)*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*li)^2, x)`

3.527 $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+ia \tan(c+dx))^2}} dx$

3.527.1 Optimal result 4961
 3.527.2 Mathematica [A] (verified) 4962
 3.527.3 Rubi [A] (verified) 4962
 3.527.4 Maple [A] (verified) 4968
 3.527.5 Fricas [B] (verification not implemented) 4968
 3.527.6 Sympy [F] 4969
 3.527.7 Maxima [F(-2)] 4970
 3.527.8 Giac [F] 4970
 3.527.9 Mupad [F(-1)] 4970

3.527.1 Optimal result

Integrand size = 36, antiderivative size = 274

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))^2}} dx$$

$$= -\frac{((-1 + 3i)A + (1 + 3i)B) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{16\sqrt{2}a^2d}$$

$$+ \frac{((-1 + 3i)A + (1 + 3i)B) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{16\sqrt{2}a^2d}$$

$$+ \frac{(3iA + B)\sqrt{\cot(c + dx)}}{8a^2d(i + \cot(c + dx))} + \frac{(A + iB)\sqrt{\cot(c + dx)}}{4d(ia + a \cot(c + dx))^2}$$

$$+ \frac{((1 + 3i)A + (1 - 3i)B) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{32\sqrt{2}a^2d}$$

$$- \frac{((1 + 3i)A + (1 - 3i)B) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{32\sqrt{2}a^2d}$$

output

```
1/32*((-1+3*I)*A+(1+3*I)*B)*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/a^2/d*2^(1/2)+1/32*((-1+3*I)*A+(1+3*I)*B)*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/a^2/d*2^(1/2)+1/64*((1+3*I)*A+(1-3*I)*B)*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/a^2/d*2^(1/2)-1/64*((1+3*I)*A+(1-3*I)*B)*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/a^2/d*2^(1/2)+1/8*(3*I*A+B)*cot(d*x+c)^(1/2)/a^2/d/(I+cot(d*x+c))+1/4*(A+I*B)*cot(d*x+c)^(1/2)/d/(I*a+a*cot(d*x+c))^2
```

3.527. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+ia \tan(c+dx))^2}} dx$

3.527.2 Mathematica [A] (verified)

Time = 3.16 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.72

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^2} dx$$

$$= \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(-2\sqrt[4]{-1} (iA + B) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right) \sec^2(c + dx) (\cos(2(c + dx))) \right)}{\dots}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^2),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-2*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)]) + (-1)^(1/4)*((-I)*A + B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)]) + Sqrt[Tan[c + d*x]]*((-3*I)*A - B + (A - (3*I)*B)*Tan[c + d*x]))/(8*a^2*d*(-I + Tan[c + d*x])^2)`

3.527.3 Rubi [A] (verified)

Time = 0.98 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.88, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 4064, 3042, 4078, 27, 3042, 4079, 25, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^2} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{\sqrt{\cot(c + dx)}(A \cot(c + dx) + B)}{(a \cot(c + dx) + ia)^2} dx$$

3.527. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
& \int \frac{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(B-A \tan\left(c+dx+\frac{\pi}{2}\right)\right)}{\left(-a \tan\left(c+dx+\frac{\pi}{2}\right)+ia\right)^2} dx && \downarrow \text{3042} \\
& \frac{\int -\frac{a(iA-B)-a(5A-3iB) \cot(c+dx)}{2\sqrt{\cot(c+dx)}(\cot(c+dx)a+ia)} dx}{4a^2} + \frac{(A+iB)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx)+ia)^2} && \downarrow \text{4078} \\
& \frac{(A+iB)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx)+ia)^2} - \frac{\int \frac{a(iA-B)-a(5A-3iB) \cot(c+dx)}{\sqrt{\cot(c+dx)}(\cot(c+dx)a+ia)} dx}{8a^2} && \downarrow \text{27} \\
& \frac{(A+iB)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx)+ia)^2} - \frac{\int \frac{a(iA-B)+a(5A-3iB) \tan\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}(ia-a \tan\left(c+dx+\frac{\pi}{2}\right))} dx}{8a^2} && \downarrow \text{3042} \\
& \frac{(A+iB)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx)+ia)^2} - \frac{\int -\frac{a^2(A-3iB)-a^2(3iA+B) \cot(c+dx)}{\sqrt{\cot(c+dx)}} dx}{2a^2} - \frac{(B+3iA)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} && \downarrow \text{4079} \\
& \frac{(A+iB)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx)+ia)^2} - \frac{\int \frac{a^2(A-3iB)-a^2(3iA+B) \cot(c+dx)}{\sqrt{\cot(c+dx)}} dx}{2a^2} - \frac{(B+3iA)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} && \downarrow \text{25} \\
& \frac{(A+iB)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx)+ia)^2} - \frac{\int \frac{(A-3iB)a^2+(3iA+B) \tan\left(c+dx+\frac{\pi}{2}\right)a^2}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}} dx}{2a^2} - \frac{(B+3iA)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} && \downarrow \text{3042} \\
& \frac{(A+iB)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx)+ia)^2} - \frac{\int -\frac{a^2(A-3iB)-(3iA+B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{a^2d} - \frac{(B+3iA)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} && \downarrow \text{4017} \\
& \frac{(A+iB)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx)+ia)^2} - \frac{\int \frac{a^2(A-3iB)-(3iA+B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{a^2d} - \frac{(B+3iA)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} && \downarrow \text{25} \\
& \frac{(A+iB)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx)+ia)^2} - \frac{\int \frac{a^2(A-3iB)-(3iA+B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{a^2d} - \frac{(B+3iA)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} && \downarrow \text{25}
\end{aligned}$$

3.527. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
 & \downarrow 27 \\
 & \frac{(A + iB)\sqrt{\cot(c + dx)}}{4d(a \cot(c + dx) + ia)^2} - \frac{\int \frac{A - 3iB - (3iA + B)\cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} - \frac{(B + 3iA)\sqrt{\cot(c + dx)}}{d(\cot(c + dx) + i)} \\
 & \downarrow 1482 \\
 & \frac{(A + iB)\sqrt{\cot(c + dx)}}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((1 + 3i)A + (1 - 3i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}((1 - 3i)A - (1 + 3i)B) \int \frac{\cot(c + dx) + 1}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} - \frac{(B + 3iA)\sqrt{\cot(c + dx)}}{d(\cot(c + dx) + i)} \\
 & \downarrow 1476 \\
 & \frac{(A + iB)\sqrt{\cot(c + dx)}}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((1 + 3i)A + (1 - 3i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}((1 - 3i)A - (1 + 3i)B) \left(\frac{1}{2} \int \frac{1}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)} + \frac{1}{2} \int \frac{1}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)} \right)}{d} \\
 & \downarrow 1082 \\
 & \frac{(A + iB)\sqrt{\cot(c + dx)}}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((1 + 3i)A + (1 - 3i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}((1 - 3i)A - (1 + 3i)B) \left(\int \frac{1}{-\cot(c + dx) - 1} \frac{d(1 - \sqrt{2}\sqrt{\cot(c + dx)})}{\sqrt{2}} - \int \frac{1}{-\cot(c + dx) - 1} \frac{d(\sqrt{2}\sqrt{\cot(c + dx)})}{\sqrt{2}} \right)}{d} \\
 & \downarrow 217 \\
 & \frac{(A + iB)\sqrt{\cot(c + dx)}}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((1 + 3i)A + (1 - 3i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}((1 - 3i)A - (1 + 3i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\cot(c + dx)})}{\sqrt{2}} \right)}{d} - \frac{(B + 3iA)\sqrt{\cot(c + dx)}}{d(\cot(c + dx) + i)} \\
 & \downarrow 1479 \\
 & \frac{(A + iB)\sqrt{\cot(c + dx)}}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((1 + 3i)A + (1 - 3i)B) \left(-\frac{\int \frac{\sqrt{2} - 2\sqrt{\cot(c + dx)}}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} \right) + \frac{1}{2}((1 - 3i)A - (1 + 3i)B) \left(\arctan(\sqrt{2}\sqrt{\cot(c + dx) + 1}) - \arctan(1 - \sqrt{2}\sqrt{\cot(c + dx)}) \right)}{d} \\
 & \downarrow 25
 \end{aligned}$$

3.527. $\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))^2}} dx$

$$\frac{\frac{(A + iB)\sqrt{\cot(c + dx)}}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((1+3i)A+(1-3i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}((1-3i)A-(1+3i)B) \left(\frac{\arctan(\dots)}{\dots} \right)}{d}}{8a^2}$$

↓ 27

$$\frac{\frac{(A + iB)\sqrt{\cot(c + dx)}}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((1+3i)A+(1-3i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2}((1-3i)A-(1+3i)B) \left(\frac{\arctan(\dots)}{\dots} \right)}{d}}{8a^2}$$

↓ 1103

$$\frac{\frac{(A + iB)\sqrt{\cot(c + dx)}}{4d(a \cot(c + dx) + ia)^2} - \frac{\frac{1}{2}((1-3i)A-(1+3i)B) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}})}{\sqrt{2}} - \frac{\arctan(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}})}{\sqrt{2}} \right) + \frac{1}{2}((1+3i)A+(1-3i)B) \left(\frac{\log(\frac{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}}{2\sqrt{2}})}{\dots} - \frac{\log(\cot(\dots))}{\dots} \right)}{d}}{8a^2}$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^2), x]`

output `((A + I*B)*Sqrt[Cot[c + d*x]])/(4*d*(I*a + a*Cot[c + d*x])^2) - (-((((3*I)*A + B)*Sqrt[Cot[c + d*x]])/(d*(I + Cot[c + d*x]))) + (((((1 - 3*I)*A - (1 + 3*I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]] + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]))/2 + (((((1 + 3*I)*A + (1 - 3*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2]] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d)/(8*a^2)`

3.527.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

$$3.527. \int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+ia \tan(c+dx))^2}} dx$$

- rule 217 $\text{Int}[(a_ + (b_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{(-1)} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /;$ $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 1082 $\text{Int}[(a_ + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}\{q = 1 - 4 \cdot \text{Simplify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /;$ $\text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4 \cdot a \cdot c]) /;$ $\text{FreeQ}\{a, b, c, x\}$
- rule 1103 $\text{Int}[(d_ + (e_ \cdot)(x_)) / ((a_ + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$
- rule 1476 $\text{Int}[(d_ + (e_ \cdot)(x_)^2) / ((a_ + (c_ \cdot)(x_)^4), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c) \ \text{Int}[1/\text{Simp}[d/e + q \cdot x + x^2, x], x], x] + \text{Simp}[e/(2 \cdot c) \ \text{Int}[1/\text{Simp}[d/e - q \cdot x + x^2, x], x], x] /;$ $\text{FreeQ}\{a, c, d, e, x\} \ \&\& \ \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{PosQ}[d \cdot e]$
- rule 1479 $\text{Int}[(d_ + (e_ \cdot)(x_)^2) / ((a_ + (c_ \cdot)(x_)^4), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[-2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c \cdot q) \ \text{Int}[(q - 2 \cdot x)/\text{Simp}[d/e + q \cdot x - x^2, x], x], x] + \text{Simp}[e/(2 \cdot c \cdot q) \ \text{Int}[(q + 2 \cdot x)/\text{Simp}[d/e - q \cdot x - x^2, x], x], x] /;$ $\text{FreeQ}\{a, c, d, e, x\} \ \&\& \ \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{NegQ}[d \cdot e]$
- rule 1482 $\text{Int}[(d_ + (e_ \cdot)(x_)^2) / ((a_ + (c_ \cdot)(x_)^4), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[a \cdot c, 2]\}, \text{Simp}[(d \cdot q + a \cdot e)/(2 \cdot a \cdot c) \ \text{Int}[(q + c \cdot x^2)/(a + c \cdot x^4), x], x] + \text{Simp}[(d \cdot q - a \cdot e)/(2 \cdot a \cdot c) \ \text{Int}[(q - c \cdot x^2)/(a + c \cdot x^4), x], x] /;$ $\text{FreeQ}\{a, c, d, e, x\} \ \&\& \ \text{NeQ}[c \cdot d^2 + a \cdot e^2, 0] \ \&\& \ \text{NeQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{NegQ}[(-a) \cdot c]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$ $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.527.4 Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.53

method	result
derivativedivides	$\frac{i(-iB+A) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{2\sqrt{2}-2i\sqrt{2}} + \frac{i\left(\frac{\left(-\frac{iB}{2} + \frac{3A}{2}\right) \cot(dx+c)^{\frac{3}{2}} + \left(\frac{3B}{2} + \frac{iA}{2}\right) \sqrt{\cot(dx+c)}}{(i+\cot(dx+c))^2} + \frac{(iB+A) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}\right)}{4}}{a^2 d}$
default	$\frac{i(-iB+A) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{2\sqrt{2}-2i\sqrt{2}} + \frac{i\left(\frac{\left(-\frac{iB}{2} + \frac{3A}{2}\right) \cot(dx+c)^{\frac{3}{2}} + \left(\frac{3B}{2} + \frac{iA}{2}\right) \sqrt{\cot(dx+c)}}{(i+\cot(dx+c))^2} + \frac{(iB+A) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}\right)}{4}}{a^2 d}$

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

output `1/a^2/d*(1/2*I*(A-I*B)/(2^(1/2)-I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))+1/4*I*((-1/2*I*B+3/2*A)*cot(d*x+c)^(3/2)+(3/2*B+1/2*I*A)*cot(d*x+c)^(1/2)/(I+cot(d*x+c))^2+(A+I*B)/(2^(1/2)+I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2))))`

3.527.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 663 vs. 2(211) = 422.

Time = 0.27 (sec) , antiderivative size = 663, normalized size of antiderivative = 2.42

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))^2}} dx$$

$$= \left(2 a^2 d \sqrt{\frac{i A^2 + 2 A B - i B^2}{a^4 d^2}} e^{(4i dx + 4i c)} \log \left(- \frac{2 \left((i a^2 d e^{(2i dx + 2i c)} - i a^2 d) \sqrt{\frac{i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} - 1}} \sqrt{\frac{i A^2 + 2 A B - i B^2}{a^4 d^2}} + (A - i B) e^{(2i dx + 2i c)} \right)}{i A + B} \right) \right)$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="fricas")`

output `1/32*(2*a^2*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(-2*((I*a^2*d*e^(2*I*d*x + 2*I*c) - I*a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 2*a^2*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(-2*((-I*a^2*d*e^(2*I*d*x + 2*I*c) + I*a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) + a^2*d*sqrt((-I*A^2 + 2*A*B + I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(1/8*((a^2*d*e^(2*I*d*x + 2*I*c) - a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 + 2*A*B + I*B^2)/(a^4*d^2)) + I*A - B)*e^(-2*I*d*x - 2*I*c)/(a^2*d)) - a^2*d*sqrt((-I*A^2 + 2*A*B + I*B^2)/(a^4*d^2))*e^(4*I*d*x + 4*I*c)*log(-1/8*((a^2*d*e^(2*I*d*x + 2*I*c) - a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 + 2*A*B + I*B^2)/(a^4*d^2)) - I*A + B)*e^(-2*I*d*x - 2*I*c)/(a^2*d)) + 2*(2*(A - I*B)*e^(4*I*d*x + 4*I*c) - (A - 3*I*B)*e^(2*I*d*x + 2*I*c) - A - I*B)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*e^(-4*I*d*x - 4*I*c)/(a^2*d)`

3.527.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))^2}} dx = \frac{\int \frac{A}{\tan^2(c+dx)\sqrt{\cot(c+dx)-2i \tan(c+dx)\sqrt{\cot(c+dx)}-\sqrt{\cot(c+dx)}}} dx + \int \frac{B \tan(c+dx)}{\tan^2(c+dx)\sqrt{\cot(c+dx)-2i \tan(c+dx)\sqrt{\cot(c+dx)}-\sqrt{\cot(c+dx)}}} dx}{a^2}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(1/2)/(a+I*a*tan(d*x+c))**2,x)`

output `-(Integral(A/(tan(c + d*x)**2*sqrt(cot(c + d*x)) - 2*I*tan(c + d*x)*sqrt(cot(c + d*x)) - sqrt(cot(c + d*x))), x) + Integral(B*tan(c + d*x)/(tan(c + d*x)**2*sqrt(cot(c + d*x)) - 2*I*tan(c + d*x)*sqrt(cot(c + d*x)) - sqrt(cot(c + d*x))), x))/a**2`

3.527.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.527.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^2} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^2 \sqrt{\cot(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)^2*sqrt(cot(d*x + c))), x)`

3.527.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^2} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + a \tan(c + dx) 1i)^2} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^2),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^2), x)`

3.527. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^2} dx$

3.528 $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

3.528.1 Optimal result 4971
 3.528.2 Mathematica [A] (verified) 4972
 3.528.3 Rubi [A] (verified) 4972
 3.528.4 Maple [A] (verified) 4977
 3.528.5 Fracas [B] (verification not implemented) 4978
 3.528.6 Sympy [F] 4979
 3.528.7 Maxima [F(-2)] 4979
 3.528.8 Giac [F] 4979
 3.528.9 Mupad [F(-1)] 4980

3.528.1 Optimal result

Integrand size = 36, antiderivative size = 284

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx$$

$$= -\frac{\left(\frac{1}{16} + \frac{i}{16}\right) \left((2 + i)A + (7 - 2i)B\right) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}a^2d}$$

$$+ \frac{\left((1 + 3i)A + (9 + 5i)B\right) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{16\sqrt{2}a^2d}$$

$$+ \frac{(A + 5iB)\sqrt{\cot(c + dx)}}{8a^2d(i + \cot(c + dx))} + \frac{(iA - B)\sqrt{\cot(c + dx)}}{4d(ia + a \cot(c + dx))^2}$$

$$+ \frac{\left((1 - 3i)A - (9 - 5i)B\right) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{32\sqrt{2}a^2d}$$

$$+ \frac{\left(\frac{1}{32} + \frac{i}{32}\right) \left((1 + 2i)A + (2 - 7i)B\right) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{\sqrt{2}a^2d}$$

output

```
(1/32+1/32*I)*((2+I)*A+(7-2*I)*B)*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/a^2/d*2^(1/2)+1/32*((1+3*I)*A+(9+5*I)*B)*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/a^2/d*2^(1/2)+1/64*((1-3*I)*A+(-9+5*I)*B)*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/a^2/d*2^(1/2)+(1/64+1/64*I)*((1+2*I)*A+(2-7*I)*B)*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/a^2/d*2^(1/2)+1/8*(A+5*I*B)*cot(d*x+c)^(1/2)/a^2/d/(I+cot(d*x+c))+1/4*(I*A-B)*cot(d*x+c)^(1/2)/d/(I*a+a*cot(d*x+c))^2
```

3.528. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

3.528.2 Mathematica [A] (verified)

Time = 4.38 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.69

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx$$

$$= \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(-2\sqrt[4]{-1} (A - iB) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right) \sec^2(c + dx) (\cos(2(c + dx))) \right)}{\dots}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^2),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-2*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)]) + (-1)^(1/4)*(A - (7*I)*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)]) - Sqrt[Tan[c + d*x]]*(A + (5*I)*B + ((3*I)*A - 7*B)*Tan[c + d*x]))/(8*a^2*d*(-I + Tan[c + d*x])^2)`

3.528.3 Rubi [A] (verified)Time = 1.00 (sec) , antiderivative size = 246, normalized size of antiderivative = 0.87, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$, Rules used = {3042, 4064, 3042, 4079, 27, 3042, 4079, 25, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2}(a + ia \tan(c + dx))^2} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{A \cot(c + dx) + B}{\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} dx$$

3.528. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
& \int \frac{B - A \tan\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{-\tan\left(c + dx + \frac{\pi}{2}\right)} \left(-a \tan\left(c + dx + \frac{\pi}{2}\right) + ia\right)^2} dx && \downarrow \text{3042} \\
& \frac{\int \frac{a(A-7iB) - 3a(iA-B) \cot(c+dx)}{2\sqrt{\cot(c+dx)}(\cot(c+dx)a+ia)} dx}{4a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx) + ia)^2} && \downarrow \text{4079} \\
& \frac{\int \frac{a(A-7iB) - 3a(iA-B) \cot(c+dx)}{\sqrt{\cot(c+dx)}(\cot(c+dx)a+ia)} dx}{8a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx) + ia)^2} && \downarrow \text{27} \\
& \frac{\int \frac{a(A-7iB) + 3a(iA-B) \tan\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}(ia-a \tan\left(c+dx+\frac{\pi}{2}\right))} dx}{8a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx) + ia)^2} && \downarrow \text{3042} \\
& \frac{\int -\frac{3(iA+3B)a^2 + (A+5iB) \cot(c+dx)a^2}{\sqrt{\cot(c+dx)}} dx}{8a^2} + \frac{(A+5iB)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx) + ia)^2} && \downarrow \text{4079} \\
& \frac{(A+5iB)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} - \frac{\int \frac{3(iA+3B)a^2 + (A+5iB) \cot(c+dx)a^2}{\sqrt{\cot(c+dx)}} dx}{8a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx) + ia)^2} && \downarrow \text{25} \\
& \frac{(A+5iB)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} - \frac{\int \frac{3a^2(iA+3B) - a^2(A+5iB) \tan\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}} dx}{8a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx) + ia)^2} && \downarrow \text{3042} \\
& \frac{(A+5iB)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} - \frac{\int -\frac{a^2(3(iA+3B) + (A+5iB) \cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{8a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx) + ia)^2} && \downarrow \text{4017} \\
& \frac{\int \frac{a^2(3(iA+3B) + (A+5iB) \cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{8a^2} + \frac{(A+5iB)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{4d(a \cot(c+dx) + ia)^2} && \downarrow \text{25}
\end{aligned}$$

3.528. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{\int \frac{3(iA+3B)+(A+5iB)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{8a^2} + \frac{(A+5iB)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{4d(a\cot(c+dx)+ia)^2} \\
& \downarrow 1482 \\
& \frac{\left(\frac{1}{2}+\frac{i}{2}\right)((1+2i)A+(2-7i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}((1+3i)A+(9+5i)B) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{(A+5iB)\sqrt{\cot(c+dx)}}{d(\cot(c+dx)+i)}}{8a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{4d(a\cot(c+dx)+ia)^2} \\
& \downarrow 1476 \\
& \frac{\left(\frac{1}{2}+\frac{i}{2}\right)((1+2i)A+(2-7i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}((1+3i)A+(9+5i)B) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}\right)}{8a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{4d(a\cot(c+dx)+ia)^2} \\
& \downarrow 1082 \\
& \frac{\left(\frac{1}{2}+\frac{i}{2}\right)((1+2i)A+(2-7i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}((1+3i)A+(9+5i)B) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right) - \int \frac{1}{-\cot(c+dx)-1} d\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{d}\right)}{8a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{4d(a\cot(c+dx)+ia)^2} \\
& \downarrow 217 \\
& \frac{\left(\frac{1}{2}+\frac{i}{2}\right)((1+2i)A+(2-7i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}((1+3i)A+(9+5i)B) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{d}\right)}{8a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{4d(a\cot(c+dx)+ia)^2} \\
& \downarrow 1479
\end{aligned}$$

3.528. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

$$\frac{\left(\frac{1}{2} + \frac{i}{2}\right)((1+2i)A + (2-7i)B) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}((1+3i)A + (9+5i)B)}{d} + \frac{8a^2}{8a^2}$$

$$\frac{(-B + iA)\sqrt{\cot(c + dx)}}{4d(a \cot(c + dx) + ia)^2}$$

↓ 25

$$\frac{\left(\frac{1}{2} + \frac{i}{2}\right)((1+2i)A + (2-7i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}((1+3i)A + (9+5i)B)}{d} + \frac{8a^2}{8a^2}$$

$$\frac{(-B + iA)\sqrt{\cot(c + dx)}}{4d(a \cot(c + dx) + ia)^2}$$

↓ 27

$$\frac{\left(\frac{1}{2} + \frac{i}{2}\right)((1+2i)A + (2-7i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2}((1+3i)A + (9+5i)B)}{d} + \frac{8a^2}{8a^2}$$

$$\frac{(-B + iA)\sqrt{\cot(c + dx)}}{4d(a \cot(c + dx) + ia)^2}$$

↓ 1103

$$\frac{\frac{1}{2}((1+3i)A + (9+5i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \left(\frac{1}{2} + \frac{i}{2}\right)((1+2i)A + (2-7i)B) \left(\frac{\log(\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} - \log[1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)]/(2\sqrt{2}) \right)}{d} + \frac{8a^2}{8a^2}$$

$$\frac{(-B + iA)\sqrt{\cot(c + dx)}}{4d(a \cot(c + dx) + ia)^2}$$

input `Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^2),x]`

output `((I*A - B)*Sqrt[Cot[c + d*x]])/(4*d*(I*a + a*Cot[c + d*x])^2) + (((A + (5*I)*B)*Sqrt[Cot[c + d*x]])/(d*(I + Cot[c + d*x])) + (((((1 + 3*I)*A + (9 + 5*I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]))/2 + (1/2 + I/2)*((1 + 2*I)*A + (2 - 7*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/d)/(8*a^2)`

3.528. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

3.528.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.528.4 Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.51

method	result
derivativedivides	$\frac{-\left(\frac{5iB}{2} - \frac{A}{2}\right) \cot(dx+c)^{\frac{3}{2}} + \left(\frac{7B}{2} - \frac{3iA}{2}\right) \sqrt{\cot(dx+c)}}{4(i+\cot(dx+c))^2} - \frac{(-7iB+A) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{4(\sqrt{2+i\sqrt{2}})} - \frac{i(iA+B) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{2(\sqrt{2-i\sqrt{2}})}$
default	$\frac{-\left(\frac{5iB}{2} - \frac{A}{2}\right) \cot(dx+c)^{\frac{3}{2}} + \left(\frac{7B}{2} - \frac{3iA}{2}\right) \sqrt{\cot(dx+c)}}{4(i+\cot(dx+c))^2} - \frac{(-7iB+A) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{4(\sqrt{2+i\sqrt{2}})} - \frac{i(iA+B) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{2(\sqrt{2-i\sqrt{2}})}$

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

$$3.528. \int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$$

output $1/a^2/d*(-1/4*((-5/2*I*B-1/2*A)*\cot(dx+c)^{(3/2)}+(7/2*B-3/2*I*A)*\cot(dx+c)^{(1/2)})/(I+\cot(dx+c))^2-1/4*(-7*I*B+A)/(2^{(1/2)}+I*2^{(1/2)})*\arctan(2*\cot(dx+c)^{(1/2)}/(2^{(1/2)}+I*2^{(1/2)}))-1/2*I*(I*A+B)/(2^{(1/2)}-I*2^{(1/2)})*\arctan(2*\cot(dx+c)^{(1/2)}/(2^{(1/2)}-I*2^{(1/2)}))$

3.528.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 666 vs. $2(209) = 418$.

Time = 0.26 (sec) , antiderivative size = 666, normalized size of antiderivative = 2.35

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx =$$

$$\left(2 a^2 d \sqrt{\frac{-i A^2 - 2 A B + i B^2}{a^4 d^2}} e^{(4i dx + 4i c)} \log \left(-\frac{2 \left((a^2 d e^{(2i dx + 2i c)} - a^2 d) \sqrt{\frac{i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} - 1}} \sqrt{\frac{-i A^2 - 2 A B + i B^2}{a^4 d^2}} + (A - i B) e^{(2i dx + 2i c)} \right)}{i A + B} \right) \right)$$

input `integrate((A+B*tan(dx+c))/cot(dx+c)^(3/2)/(a+I*a*tan(dx+c))^2,x, algorithm="fracas")`

output $-1/32*(2*a^2*d*\sqrt{((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2))}*e^{(4*I*d*x + 4*I*c)}*\log(-2*((a^2*d*e^{(2*I*d*x + 2*I*c)} - a^2*d)*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}*\sqrt{((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2))} + (A - I*B)*e^{(2*I*d*x + 2*I*c)}*e^{(-2*I*d*x - 2*I*c)/(I*A + B)}) - 2*a^2*d*\sqrt{((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2))}*e^{(4*I*d*x + 4*I*c)}*\log(2*((a^2*d*e^{(2*I*d*x + 2*I*c)} - a^2*d)*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}*\sqrt{((-I*A^2 - 2*A*B + I*B^2)/(a^4*d^2))} - (A - I*B)*e^{(2*I*d*x + 2*I*c)}*e^{(-2*I*d*x - 2*I*c)/(I*A + B)}) + a^2*d*\sqrt{(I*A^2 + 14*A*B - 49*I*B^2)/(a^4*d^2)}*e^{(4*I*d*x + 4*I*c)}*\log(-1/8*((a^2*d*e^{(2*I*d*x + 2*I*c)} - a^2*d)*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}*\sqrt{(I*A^2 + 14*A*B - 49*I*B^2)/(a^4*d^2)} + A - 7*I*B)*e^{(-2*I*d*x - 2*I*c)/(a^2*d)} - a^2*d*\sqrt{(I*A^2 + 14*A*B - 49*I*B^2)/(a^4*d^2)}*e^{(4*I*d*x + 4*I*c)}*\log(1/8*((a^2*d*e^{(2*I*d*x + 2*I*c)} - a^2*d)*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}*\sqrt{(I*A^2 + 14*A*B - 49*I*B^2)/(a^4*d^2)} - A + 7*I*B)*e^{(-2*I*d*x - 2*I*c)/(a^2*d)} + 2*(2*(I*A - 3*B)*e^{(4*I*d*x + 4*I*c)} - (3*I*A - 7*B)*e^{(2*I*d*x + 2*I*c)} + I*A - B)*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)})*e^{(-4*I*d*x - 4*I*c)/(a^2*d)}$

3.528. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

3.528.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx = \frac{\int \frac{A}{\tan^2(c + dx) \cot^{\frac{3}{2}}(c + dx) - 2i \tan(c + dx) \cot^{\frac{3}{2}}(c + dx) - \cot^{\frac{3}{2}}(c + dx)} dx + \int \frac{B \tan(c + dx)}{\tan^2(c + dx) \cot^{\frac{3}{2}}(c + dx) - 2i \tan(c + dx) \cot^{\frac{3}{2}}(c + dx) - \cot^{\frac{3}{2}}(c + dx)} dx}{a^2}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(3/2)/(a+I*a*tan(d*x+c))**2,x)`

output `-(Integral(A/(tan(c + d*x)**2*cot(c + d*x)**(3/2) - 2*I*tan(c + d*x)*cot(c + d*x)**(3/2) - cot(c + d*x)**(3/2)), x) + Integral(B*tan(c + d*x)/(tan(c + d*x)**2*cot(c + d*x)**(3/2) - 2*I*tan(c + d*x)*cot(c + d*x)**(3/2) - cot(c + d*x)**(3/2)), x))/a**2`

3.528.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.528.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^2 \cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)^2*cot(d*x + c)^(3/2)), x)`

3.528. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

3.528.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2} (a + a \tan(c + dx) li)^2} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + a*tan(c + d*x)*li)^2),x)`output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + a*tan(c + d*x)*li)^2), x)`

$$3.529 \quad \int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$$

3.529.1 Optimal result	4981
3.529.2 Mathematica [A] (verified)	4982
3.529.3 Rubi [A] (verified)	4982
3.529.4 Maple [A] (verified)	4988
3.529.5 Fricas [B] (verification not implemented)	4989
3.529.6 Sympy [F(-1)]	4990
3.529.7 Maxima [F(-2)]	4991
3.529.8 Giac [F]	4991
3.529.9 Mupad [F(-1)]	4991

3.529.1 Optimal result

Integrand size = 36, antiderivative size = 319

$$\begin{aligned} & \int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx \\ &= -\frac{\left(\frac{1}{16}-\frac{i}{16}\right)\left((2+7i)A-(23+2i)B\right) \arctan\left(1-\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}a^2d} \\ & \quad + \frac{\left((9+5i)A-(25-21i)B\right) \arctan\left(1+\sqrt{2}\sqrt{\cot(c+dx)}\right)}{16\sqrt{2}a^2d} + \frac{5(iA-5B)}{8a^2d\sqrt{\cot(c+dx)}} \\ & \quad + \frac{3A+7iB}{8a^2d\sqrt{\cot(c+dx)}(i+\cot(c+dx))} + \frac{iA-B}{4d\sqrt{\cot(c+dx)}(ia+a \cot(c+dx))^2} \\ & \quad - \frac{\left(\frac{1}{32}-\frac{i}{32}\right)\left((7+2i)A+(2+23i)B\right) \log\left(1-\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{\sqrt{2}a^2d} \\ & \quad + \frac{\left(\frac{1}{32}-\frac{i}{32}\right)\left((7+2i)A+(2+23i)B\right) \log\left(1+\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{\sqrt{2}a^2d} \end{aligned}$$

output $(1/32-1/32*I)*((2+7*I)*A-(23+2*I)*B)*\arctan(-1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/a^{2/d}*2^{(1/2)}+1/32*((9+5*I)*A+(-25+21*I)*B)*\arctan(1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/a^{2/d}*2^{(1/2)}+(-1/64+1/64*I)*((7+2*I)*A+(2+23*I)*B)*\ln(1+\cot(d*x+c)-2^{(1/2)}*\cot(d*x+c)^{(1/2)})/a^{2/d}*2^{(1/2)}+(1/64-1/64*I)*((7+2*I)*A+(2+23*I)*B)*\ln(1+\cot(d*x+c)+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/a^{2/d}*2^{(1/2)}+5/8*(I*A-5*B)/a^{2/d}/\cot(d*x+c)^{(1/2)}+1/8*(3*A+7*I*B)/a^{2/d}/(I+\cot(d*x+c))/\cot(d*x+c)^{(1/2)}+1/4*(I*A-B)/d/(I*a+a*\cot(d*x+c))^2/\cot(d*x+c)^{(1/2)}$

$$3.529. \quad \int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$$

3.529.2 Mathematica [A] (verified)

Time = 4.94 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.67

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx = \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(-2\sqrt[4]{-1}(iA + B) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right) \sec^2(c + dx) (\cos(2(c + dx))) \right)}{\dots}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^2),x]`

output `-1/8*(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-2*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)]) + (-1)^(3/4)*(7*A + (23*I)*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^2*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)]) + Sqrt[Tan[c + d*x]]*((5*I)*(A + (5*I)*B) - (7*A + (43*I)*B)*Tan[c + d*x] + 16*B*Tan[c + d*x]^2)))/(a^2*d*(-I + Tan[c + d*x])^2)`

3.529.3 Rubi [A] (verified)Time = 1.27 (sec) , antiderivative size = 288, normalized size of antiderivative = 0.90, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.639$, Rules used = {3042, 4064, 3042, 4079, 27, 3042, 4079, 25, 3042, 4012, 25, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{5/2}(a + ia \tan(c + dx))^2} dx \\ & \quad \downarrow \text{4064} \\ & \int \frac{A \cot(c + dx) + B}{\cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + ia)^2} dx \end{aligned}$$

3.529. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
 &\int \frac{B - A \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2} (-a \tan(c + dx + \frac{\pi}{2}) + ia)^2} dx \\
 &\quad \downarrow \text{3042} \\
 &\int \frac{-\frac{a(A+9iB)+5a(iA-B) \cot(c+dx)}{2 \cot^{\frac{3}{2}}(c+dx)(\cot(c+dx)a+ia)} dx}{4a^2} + \frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} \\
 &\quad \downarrow \text{4079} \\
 &\frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} - \frac{\int \frac{a(A+9iB)+5a(iA-B) \cot(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(\cot(c+dx)a+ia)} dx}{8a^2} \\
 &\quad \downarrow \text{27} \\
 &\frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} - \frac{\int \frac{a(A+9iB)-5a(iA-B) \tan(c+dx+\frac{\pi}{2})}{(-\tan(c+dx+\frac{\pi}{2}))^{3/2}(ia-a \tan(c+dx+\frac{\pi}{2}))} dx}{8a^2} \\
 &\quad \downarrow \text{3042} \\
 &\frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} - \frac{\int \frac{5a^2(iA-5B)-3a^2(3A+7iB) \cot(c+dx)}{\cot^{\frac{3}{2}}(c+dx)} dx}{8a^2} - \frac{3A+7iB}{d\sqrt{\cot(c+dx)}(\cot(c+dx)+i)} \\
 &\quad \downarrow \text{4079} \\
 &\frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} - \frac{\int \frac{5a^2(iA-5B)-3a^2(3A+7iB) \cot(c+dx)}{\cot^{\frac{3}{2}}(c+dx)} dx}{8a^2} - \frac{3A+7iB}{d\sqrt{\cot(c+dx)}(\cot(c+dx)+i)} \\
 &\quad \downarrow \text{25} \\
 &\frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} - \frac{\int \frac{5(iA-5B)a^2+3(3A+7iB) \tan(c+dx+\frac{\pi}{2})a^2}{(-\tan(c+dx+\frac{\pi}{2}))^{3/2}} dx}{8a^2} - \frac{3A+7iB}{d\sqrt{\cot(c+dx)}(\cot(c+dx)+i)} \\
 &\quad \downarrow \text{3042} \\
 &\frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} - \frac{\int \frac{5(iA-5B)a^2+3(3A+7iB) \tan(c+dx+\frac{\pi}{2})a^2}{(-\tan(c+dx+\frac{\pi}{2}))^{3/2}} dx}{8a^2} - \frac{3A+7iB}{d\sqrt{\cot(c+dx)}(\cot(c+dx)+i)} \\
 &\quad \downarrow \text{4012} \\
 &\frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} - \frac{\int \frac{3(3A+7iB)a^2+5(iA-5B) \cot(c+dx)a^2}{\sqrt{\cot(c+dx)}} dx + \frac{10a^2(-5B+iA)}{d\sqrt{\cot(c+dx)}}}{8a^2} - \frac{3A+7iB}{d\sqrt{\cot(c+dx)}(\cot(c+dx)+i)} \\
 &\quad \downarrow \\
 &\frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} - \frac{\int \frac{3(3A+7iB)a^2+5(iA-5B) \cot(c+dx)a^2}{\sqrt{\cot(c+dx)}} dx + \frac{10a^2(-5B+iA)}{d\sqrt{\cot(c+dx)}}}{8a^2} - \frac{3A+7iB}{d\sqrt{\cot(c+dx)}(\cot(c+dx)+i)}
 \end{aligned}$$

3.529. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
 & \downarrow 25 \\
 & \frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} - \\
 & \frac{\frac{10a^2(-5B+iA)}{d\sqrt{\cot(c+dx)}} - \int \frac{3(3A+7iB)a^2+5(iA-5B)\cot(c+dx)a^2}{\sqrt{\cot(c+dx)}} dx}{2a^2} - \frac{3A+7iB}{d\sqrt{\cot(c+dx)}(\cot(c+dx)+i)} \\
 & \frac{8a^2}{\downarrow 3042} \\
 & \frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} - \\
 & \frac{\frac{10a^2(-5B+iA)}{d\sqrt{\cot(c+dx)}} - \int \frac{3a^2(3A+7iB)-5a^2(iA-5B)\tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{2a^2} - \frac{3A+7iB}{d\sqrt{\cot(c+dx)}(\cot(c+dx)+i)} \\
 & \frac{8a^2}{\downarrow 4017} \\
 & \frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} - \\
 & \frac{\frac{10a^2(-5B+iA)}{d\sqrt{\cot(c+dx)}} - 2 \int \frac{a^2(3(3A+7iB)+5(iA-5B)\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{2a^2} - \frac{3A+7iB}{d\sqrt{\cot(c+dx)}(\cot(c+dx)+i)} \\
 & \frac{8a^2}{\downarrow 25} \\
 & \frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} - \\
 & \frac{2 \int \frac{a^2(3(3A+7iB)+5(iA-5B)\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{10a^2(-5B+iA)}{d\sqrt{\cot(c+dx)}}}{2a^2} - \frac{3A+7iB}{d\sqrt{\cot(c+dx)}(\cot(c+dx)+i)} \\
 & \frac{8a^2}{\downarrow 27} \\
 & \frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} - \\
 & \frac{\frac{2a^2 \int \frac{3(3A+7iB)+5(iA-5B)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{2a^2} + \frac{10a^2(-5B+iA)}{d\sqrt{\cot(c+dx)}}}{2a^2} - \frac{3A+7iB}{d\sqrt{\cot(c+dx)}(\cot(c+dx)+i)} \\
 & \frac{8a^2}{\downarrow 1482} \\
 & \frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} - \\
 & \frac{2a^2 \left(\left(\frac{1}{2} - \frac{i}{2} \right) ((7+2i)A + (2+23i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} ((9+5i)A - (25-21i)B) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right) + \frac{10a^2(-5B+iA)}{d\sqrt{\cot(c+dx)}}}{2a^2} - \frac{3A+7iB}{d\sqrt{\cot(c+dx)}(\cot(c+dx)+i)} \\
 & \frac{8a^2}{\downarrow}
 \end{aligned}$$

3.529. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned} & \downarrow 1476 \\ & \frac{-B + iA}{4d\sqrt{\cot(c+dx)}(a\cot(c+dx) + ia)^2} \\ & \frac{2a^2 \left(\left(\frac{1}{2} - \frac{i}{2}\right) ((7+2i)A + (2+23i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} ((9+5i)A - (25-21i)B) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{2a^2} \\ & \frac{8a^2}{2a^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 1082 \\ & \frac{-B + iA}{4d\sqrt{\cot(c+dx)}(a\cot(c+dx) + ia)^2} \\ & \frac{2a^2 \left(\left(\frac{1}{2} - \frac{i}{2}\right) ((7+2i)A + (2+23i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} ((9+5i)A - (25-21i)B) \left(\int \frac{1}{\cot(c+dx) - \sqrt{2}} d\sqrt{\cot(c+dx)} - \int \frac{1}{\cot(c+dx) + \sqrt{2}} d\sqrt{\cot(c+dx)} \right) \right)}{2a^2} \\ & \frac{8a^2}{2a^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 217 \\ & \frac{-B + iA}{4d\sqrt{\cot(c+dx)}(a\cot(c+dx) + ia)^2} \\ & \frac{2a^2 \left(\left(\frac{1}{2} - \frac{i}{2}\right) ((7+2i)A + (2+23i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} ((9+5i)A - (25-21i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{2a^2} + \frac{10}{d} \\ & \frac{8a^2}{2a^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 1479 \\ & \frac{-B + iA}{4d\sqrt{\cot(c+dx)}(a\cot(c+dx) + ia)^2} \\ & \frac{2a^2 \left(\left(\frac{1}{2} - \frac{i}{2}\right) ((7+2i)A + (2+23i)B) \left(- \int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} ((9+5i)A - (25-21i)B) \right)}{2a^2} \\ & \frac{8a^2}{2a^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 25 \\ & \frac{-B + iA}{4d\sqrt{\cot(c+dx)}(a\cot(c+dx) + ia)^2} \\ & \frac{2a^2 \left(\left(\frac{1}{2} - \frac{i}{2}\right) ((7+2i)A + (2+23i)B) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} ((9+5i)A - (25-21i)B) \right)}{2a^2} \\ & \frac{8a^2}{2a^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \end{aligned}$$

3.529. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

$$\begin{aligned}
 & \frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} \\
 & 2a^2 \left(\left(\frac{1}{2} - \frac{i}{2} \right) ((7+2i)A + (2+23i)B) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} ((9+5i)A - (25-21i)B) \right) \\
 & \frac{2a^2}{8a^2} \\
 & \downarrow 1103 \\
 & \frac{-B + iA}{4d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} \\
 & 2a^2 \left(\frac{1}{2} ((9+5i)A - (25-21i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \left(\frac{1}{2} - \frac{i}{2} \right) ((7+2i)A + (2+23i)B) \left(\frac{\log(\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right) \right) \\
 & \frac{2a^2}{8a^2}
 \end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^2),x]`

output `(I*A - B)/(4*d*Sqrt[Cot[c + d*x]]*(I*a + a*Cot[c + d*x])^2) - (-((3*A + (7*I)*B)/(d*Sqrt[Cot[c + d*x]]*(I + Cot[c + d*x]))) - ((10*a^2*(I*A - 5*B))/(d*Sqrt[Cot[c + d*x]]) + (2*a^2*(((9 + 5*I)*A - (25 - 21*I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]))/2 + (1/2 - I/2)*((7 + 2*I)*A + (2 + 23*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/d)/(2*a^2)/(8*a^2)`

3.529.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

3.529. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.529.4 Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.49

method	result
derivativedivides	$\frac{-\frac{i(-iB+A) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{2(\sqrt{2-i\sqrt{2}})} - \frac{2B}{\sqrt{\cot(dx+c)}} + \frac{\left(\frac{5iA}{2} - \frac{9B}{2}\right) \cot(dx+c)^{\frac{3}{2}} + \left(-\frac{7A}{2} - \frac{11iB}{2}\right) \sqrt{\cot(dx+c)}}{4(i+\cot(dx+c))^2} + \frac{(7iA-23B) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{4\sqrt{2+4i\sqrt{2}}}}{a^2 d}$
default	$\frac{-\frac{i(-iB+A) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{2(\sqrt{2-i\sqrt{2}})} - \frac{2B}{\sqrt{\cot(dx+c)}} + \frac{\left(\frac{5iA}{2} - \frac{9B}{2}\right) \cot(dx+c)^{\frac{3}{2}} + \left(-\frac{7A}{2} - \frac{11iB}{2}\right) \sqrt{\cot(dx+c)}}{4(i+\cot(dx+c))^2} + \frac{(7iA-23B) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{4\sqrt{2+4i\sqrt{2}}}}{a^2 d}$

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^2,x,method=_RETURNVERBOSE)`

$$3.529. \int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$$

output $1/a^2/d*(-1/2*I*(A-I*B)/(2^(1/2)-I*2^(1/2))*\arctan(2*\cot(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))-2*B/\cot(d*x+c)^(1/2)+1/4*((5/2*I*A-9/2*B)*\cot(d*x+c)^(3/2)+(-7/2*A-11/2*I*B)*\cot(d*x+c)^(1/2))/(I+\cot(d*x+c))^2+1/4*(7*I*A-23*B)/(2^(1/2)+I*2^(1/2))*\arctan(2*\cot(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))$

3.529.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 761 vs. $2(234) = 468$.

Time = 0.27 (sec) , antiderivative size = 761, normalized size of antiderivative = 2.39

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx =$$

$$2 \left(a^2 de^{(6i dx + 6i c)} + a^2 de^{(4i dx + 4i c)} \right) \sqrt{\frac{i A^2 + 2 AB - i B^2}{a^4 d^2}} \log \left(- \frac{2 \left((i a^2 de^{(2i dx + 2i c)} - i a^2 d) \sqrt{\frac{i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} - 1}} \sqrt{\frac{i A^2 + 2 AB - i B^2}{a^4 d^2}} \right)}{i A + B} \right)$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="fracas")`

output

```
-1/32*(2*(a^2*d*e^(6*I*d*x + 6*I*c) + a^2*d*e^(4*I*d*x + 4*I*c))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))*log(-2*((I*a^2*d*e^(2*I*d*x + 2*I*c) - I*a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B) - 2*(a^2*d*e^(6*I*d*x + 6*I*c) + a^2*d*e^(4*I*d*x + 4*I*c))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2))*log(-2*((-I*a^2*d*e^(2*I*d*x + 2*I*c) + I*a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^4*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B) - (a^2*d*e^(6*I*d*x + 6*I*c) + a^2*d*e^(4*I*d*x + 4*I*c))*sqrt((-49*I*A^2 + 322*A*B + 529*I*B^2)/(a^4*d^2))*log(1/8*((a^2*d*e^(2*I*d*x + 2*I*c) - a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-49*I*A^2 + 322*A*B + 529*I*B^2)/(a^4*d^2)) + 7*I*A - 23*B)*e^(-2*I*d*x - 2*I*c)/(a^2*d) + (a^2*d*e^(6*I*d*x + 6*I*c) + a^2*d*e^(4*I*d*x + 4*I*c))*sqrt((-49*I*A^2 + 322*A*B + 529*I*B^2)/(a^4*d^2))*log(-1/8*((a^2*d*e^(2*I*d*x + 2*I*c) - a^2*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-49*I*A^2 + 322*A*B + 529*I*B^2)/(a^4*d^2)) - 7*I*A + 23*B)*e^(-2*I*d*x - 2*I*c)/(a^2*d) - 2*(6*(A + 7*I*B)*e^(6*I*d*x + 6*I*c) - (A + 33*I*B)*e^(4*I*d*x + 4*I*c) - 2*(3*A + 5*I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))/(a^2*d*e^(6*I*d*x + 6*I*c) + a^2*d*e^(4*I*...
```

3.529.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(5/2)/(a+I*a*tan(d*x+c))**2,x)`

output `Timed out`

3.529.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: exp: undefined: 0 to a negative exponent.`

3.529.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^2 \cot(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^2,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)^2*cot(d*x + c)^(5/2)), x)`

3.529.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^2} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{\frac{5}{2}}(a + a \tan(c + dx) li)^2} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + a*tan(c + d*x)*li)^2),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + a*tan(c + d*x)*li)^2), x)`

3.529. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^2} dx$

3.530
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.530.1 Optimal result	4992
3.530.2 Mathematica [A] (verified)	4993
3.530.3 Rubi [A] (verified)	4993
3.530.4 Maple [A] (verified)	5000
3.530.5 Fricas [B] (verification not implemented)	5001
3.530.6 Sympy [F]	5002
3.530.7 Maxima [F(-2)]	5003
3.530.8 Giac [F]	5003
3.530.9 Mupad [F(-1)]	5003

3.530.1 Optimal result

Integrand size = 36, antiderivative size = 367

$$\begin{aligned} & \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\ &= -\frac{\left(\frac{1}{16} - \frac{i}{16}\right) \left((1+29i)A - (6+i)B \right) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}a^3d} \\ &+ \frac{\left((30+28i)A - (7-5i)B \right) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{16\sqrt{2}a^3d} - \frac{5(6A+iB)\sqrt{\cot(c+dx)}}{8a^3d} \\ &+ \frac{(A+iB)\cot^{\frac{7}{2}}(c+dx)}{6d(ia+a \cot(c+dx))^3} + \frac{(5A+2iB)\cot^{\frac{5}{2}}(c+dx)}{12ad(ia+a \cot(c+dx))^2} + \frac{7(4A+iB)\cot^{\frac{3}{2}}(c+dx)}{24d(ia^3+a^3 \cot(c+dx))} \\ &- \frac{\left(\frac{1}{32} - \frac{i}{32}\right) \left((29+i)A + (1+6i)B \right) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{\sqrt{2}a^3d} \\ &+ \frac{\left(\frac{1}{32} - \frac{i}{32}\right) \left((29+i)A + (1+6i)B \right) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{\sqrt{2}a^3d} \end{aligned}$$

output $\frac{1}{6}(A+I*B)*\cot(d*x+c)^{(7/2)}/d/(I*a+a*\cot(d*x+c))^{3}+1/12*(5*A+2*I*B)*\cot(d*x+c)^{(5/2)}/a/d/(I*a+a*\cot(d*x+c))^{2}+7/24*(4*A+I*B)*\cot(d*x+c)^{(3/2)}/d/(I*a^{3}+a^{3}*\cot(d*x+c))+(1/32-1/32*I)*((1+29*I)*A-(6+I)*B)*\arctan(-1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/a^{3}/d*2^{(1/2)}+1/32*((30+28*I)*A+(-7+5*I)*B)*\arctan(1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/a^{3}/d*2^{(1/2)}+(-1/64+1/64*I)*((29+I)*A+(1+6*I)*B)*\ln(1+\cot(d*x+c)-2^{(1/2)}*\cot(d*x+c)^{(1/2)})/a^{3}/d*2^{(1/2)}+(1/64-1/64*I)*((29+I)*A+(1+6*I)*B)*\ln(1+\cot(d*x+c)+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/a^{3}/d*2^{(1/2)}-5/8*(6*A+I*B)*\cot(d*x+c)^{(1/2)}/a^{3}/d$

3.530.2 Mathematica [A] (verified)

Time = 6.00 (sec) , antiderivative size = 248, normalized size of antiderivative = 0.68

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{\sqrt{\cot(c+dx)} \left(-48iA + 3\sqrt[4]{-1}(A-iB) \arctan \left((-1)^{3/4} \sqrt{\tan(c+dx)} \right) \sec^3(c+dx)(\cos(3(c+dx)) + i \right)}{\dots}$$

input `Integrate[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output $(\text{Sqrt}[\text{Cot}[c + d*x]]*((-48*I)*A + 3*(-1)^{(1/4)}*(A - I*B)*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]]*\text{Sec}[c + d*x]^3*(\text{Cos}[3*(c + d*x)] + I*\text{Sin}[3*(c + d*x)])*\text{Sqrt}[\text{Tan}[c + d*x]] - 3*(-1)^{(1/4)}*(29*A + (6*I)*B)*\text{ArcTanh}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]]*\text{Sec}[c + d*x]^3*(\text{Cos}[3*(c + d*x)] + I*\text{Sin}[3*(c + d*x)])*\text{Sqrt}[\text{Tan}[c + d*x]] + 204*A*\text{Tan}[c + d*x] + (27*I)*B*\text{Tan}[c + d*x] + (242*I)*A*\text{Tan}[c + d*x]^2 - 38*B*\text{Tan}[c + d*x]^2 - 90*A*\text{Tan}[c + d*x]^3 - (15*I)*B*\text{Tan}[c + d*x]^3))/(24*a^3*d*(-I + \text{Tan}[c + d*x])^3)$

3.530.3 Rubi [A] (verified)

Time = 1.64 (sec) , antiderivative size = 342, normalized size of antiderivative = 0.93, number of steps used = 26, number of rules used = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.694$, Rules used = {3042, 4064, 3042, 4078, 27, 3042, 4078, 27, 3042, 4078, 27, 3042, 4078, 27, 3042, 4011, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.530. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\
& \quad \downarrow 3042 \\
& \int \frac{\cot(c+dx)^{3/2}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx \\
& \quad \downarrow 4064 \\
& \int \frac{\cot^{\frac{7}{2}}(c+dx)(A \cot(c+dx)+B)}{(a \cot(c+dx)+ia)^3} dx \\
& \quad \downarrow 3042 \\
& \int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{7/2}(B-A \tan(c+dx+\frac{\pi}{2}))}{(-a \tan(c+dx+\frac{\pi}{2})+ia)^3} dx \\
& \quad \downarrow 4078 \\
& \frac{\int -\frac{\cot^{\frac{5}{2}}(c+dx)(7a(iA-B)-a(13A+iB) \cot(c+dx))}{2(\cot(c+dx)a+ia)^2} dx}{6a^2} + \frac{(A+iB) \cot^{\frac{7}{2}}(c+dx)}{6d(a \cot(c+dx)+ia)^3} \\
& \quad \downarrow 27 \\
& \frac{(A+iB) \cot^{\frac{7}{2}}(c+dx)}{6d(a \cot(c+dx)+ia)^3} - \frac{\int \frac{\cot^{\frac{5}{2}}(c+dx)(7a(iA-B)-a(13A+iB) \cot(c+dx))}{(\cot(c+dx)a+ia)^2} dx}{12a^2} \\
& \quad \downarrow 3042 \\
& \frac{(A+iB) \cot^{\frac{7}{2}}(c+dx)}{6d(a \cot(c+dx)+ia)^3} - \frac{\int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{5/2}(7a(iA-B)+a(13A+iB) \tan(c+dx+\frac{\pi}{2}))}{(ia-a \tan(c+dx+\frac{\pi}{2}))^2} dx}{12a^2} \\
& \quad \downarrow 4078 \\
& \frac{(A+iB) \cot^{\frac{7}{2}}(c+dx)}{6d(a \cot(c+dx)+ia)^3} - \frac{\int \frac{2 \cot^{\frac{3}{2}}(c+dx)(5a^2(5iA-2B)-a^2(31A+4iB) \cot(c+dx))}{\cot(c+dx)a+ia} dx}{4a^2} - \frac{a(5A+2iB) \cot^{\frac{5}{2}}(c+dx)}{d(a \cot(c+dx)+ia)^2} \\
& \quad \downarrow 27 \\
& \frac{(A+iB) \cot^{\frac{7}{2}}(c+dx)}{6d(a \cot(c+dx)+ia)^3} - \frac{\int \frac{\cot^{\frac{3}{2}}(c+dx)(5a^2(5iA-2B)-a^2(31A+4iB) \cot(c+dx))}{2a^2} dx}{12a^2} - \frac{a(5A+2iB) \cot^{\frac{5}{2}}(c+dx)}{d(a \cot(c+dx)+ia)^2} \\
& \quad \downarrow 3042
\end{aligned}$$

3.530. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{\frac{(A+iB)\cot\frac{7}{2}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{3/2}(5(5iA-2B)a^2+(31A+4iB)\tan(c+dx+\frac{\pi}{2})a^2)}{ia-a\tan(c+dx+\frac{\pi}{2})} dx}{2a^2}}{12a^2} - \frac{a(5A+2iB)\cot\frac{5}{2}(c+dx)}{d(a\cot(c+dx)+ia)^2} \\
 & \quad \downarrow 4078 \\
 & \frac{\frac{(A+iB)\cot\frac{7}{2}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \int \frac{3\sqrt{\cot(c+dx)}(7a^3(4iA-B)-5a^3(6A+iB)\cot(c+dx))}{2a^2} dx - \frac{7a^2(4A+iB)\cot\frac{3}{2}(c+dx)}{d(a\cot(c+dx)+ia)}}{2a^2}}{12a^2} - \frac{a(5A+2iB)\cot\frac{5}{2}(c+dx)}{d(a\cot(c+dx)+ia)^2} \\
 & \quad \downarrow 27 \\
 & \frac{\frac{(A+iB)\cot\frac{7}{2}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{3\int\sqrt{\cot(c+dx)}(7a^3(4iA-B)-5a^3(6A+iB)\cot(c+dx))}{2a^2} dx - \frac{7a^2(4A+iB)\cot\frac{3}{2}(c+dx)}{d(a\cot(c+dx)+ia)}}{2a^2}}{12a^2} - \frac{a(5A+2iB)\cot\frac{5}{2}(c+dx)}{d(a\cot(c+dx)+ia)^2} \\
 & \quad \downarrow 3042 \\
 & \frac{\frac{(A+iB)\cot\frac{7}{2}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{3\int\sqrt{-\tan(c+dx+\frac{\pi}{2})}(7(4iA-B)a^3+5(6A+iB)\tan(c+dx+\frac{\pi}{2})a^3) dx}{2a^2} - \frac{7a^2(4A+iB)\cot\frac{3}{2}(c+dx)}{d(a\cot(c+dx)+ia)}}{2a^2}}{12a^2} - \frac{a(5A+2iB)\cot\frac{5}{2}(c+dx)}{d(a\cot(c+dx)+ia)^2} \\
 & \quad \downarrow 4011 \\
 & \frac{\frac{(A+iB)\cot\frac{7}{2}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{3\left(\int\frac{5(6A+iB)a^3+7(4iA-B)\cot(c+dx)a^3}{\sqrt{\cot(c+dx)}} dx + \frac{10a^3(6A+iB)\sqrt{\cot(c+dx)}}{d}\right)}{2a^2} - \frac{7a^2(4A+iB)\cot\frac{3}{2}(c+dx)}{d(a\cot(c+dx)+ia)}}{2a^2}}{12a^2} - \frac{a(5A+2iB)\cot\frac{5}{2}(c+dx)}{d(a\cot(c+dx)+ia)^2} \\
 & \quad \downarrow 3042 \\
 & \frac{\frac{(A+iB)\cot\frac{7}{2}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{3\left(\int\frac{5a^3(6A+iB)-7a^3(4iA-B)\tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx + \frac{10a^3(6A+iB)\sqrt{\cot(c+dx)}}{d}\right)}{2a^2} - \frac{7a^2(4A+iB)\cot\frac{3}{2}(c+dx)}{d(a\cot(c+dx)+ia)}}{2a^2}}{12a^2} - \frac{a(5A+2iB)\cot\frac{5}{2}(c+dx)}{d(a\cot(c+dx)+ia)^2} \\
 & \quad \downarrow 4017
 \end{aligned}$$

3.530. $\int \frac{\cot\frac{3}{2}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{(A + iB) \cot^{\frac{7}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\
 & \frac{3 \left(\frac{2 \int -\frac{a^3(5(6A+iB)+7(4iA-B) \cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} + \frac{10a^3(6A+iB)\sqrt{\cot(c+dx)}}{d} \right)}{2a^2} - \frac{7a^2(4A+iB) \cot^{\frac{3}{2}}(c+dx)}{d(a \cot(c+dx)+ia)} - \frac{a(5A+2iB) \cot^{\frac{5}{2}}(c+dx)}{d(a \cot(c+dx)+ia)^2} \\
 & \frac{12a^2}{2a^2} \\
 & \downarrow 25 \\
 & \frac{(A + iB) \cot^{\frac{7}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\
 & \frac{3 \left(\frac{10a^3(6A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2 \int \frac{a^3(5(6A+iB)+7(4iA-B) \cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} \right)}{2a^2} - \frac{7a^2(4A+iB) \cot^{\frac{3}{2}}(c+dx)}{d(a \cot(c+dx)+ia)} - \frac{a(5A+2iB) \cot^{\frac{5}{2}}(c+dx)}{d(a \cot(c+dx)+ia)^2} \\
 & \frac{12a^2}{2a^2} \\
 & \downarrow 27 \\
 & \frac{(A + iB) \cot^{\frac{7}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\
 & \frac{3 \left(\frac{10a^3(6A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a^3 \int \frac{5(6A+iB)+7(4iA-B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} \right)}{2a^2} - \frac{7a^2(4A+iB) \cot^{\frac{3}{2}}(c+dx)}{d(a \cot(c+dx)+ia)} - \frac{a(5A+2iB) \cot^{\frac{5}{2}}(c+dx)}{d(a \cot(c+dx)+ia)^2} \\
 & \frac{12a^2}{2a^2} \\
 & \downarrow 1482 \\
 & \frac{(A + iB) \cot^{\frac{7}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\
 & \frac{3 \left(\frac{10a^3(6A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a^3 \left(\left(\frac{1}{2} - \frac{i}{2} \right) ((29+i)A + (1+6i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} ((30+28i)A - (7-5i)B) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d} \right)}{2a^2} - \frac{7a^2(4A+iB) \cot^{\frac{3}{2}}(c+dx)}{d(a \cot(c+dx)+ia)} - \frac{a(5A+2iB) \cot^{\frac{5}{2}}(c+dx)}{d(a \cot(c+dx)+ia)^2} \\
 & \frac{12a^2}{2a^2} \\
 & \downarrow 1476 \\
 & \frac{(A + iB) \cot^{\frac{7}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\
 & \frac{3 \left(\frac{10a^3(6A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a^3 \left(\left(\frac{1}{2} - \frac{i}{2} \right) ((29+i)A + (1+6i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} ((30+28i)A - (7-5i)B) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{d} \right)}{2a^2} - \frac{7a^2(4A+iB) \cot^{\frac{3}{2}}(c+dx)}{d(a \cot(c+dx)+ia)} - \frac{a(5A+2iB) \cot^{\frac{5}{2}}(c+dx)}{d(a \cot(c+dx)+ia)^2} \\
 & \frac{12a^2}{2a^2} \\
 & \downarrow 1082
 \end{aligned}$$

3.530. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\frac{(A + iB) \cot^{\frac{7}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \frac{10a^3(6A + iB)\sqrt{\cot(c + dx)}}{d} - \frac{2a^3 \left(\left(\frac{1}{2} - \frac{i}{2} \right) ((29 + i)A + (1 + 6i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2} ((30 + 28i)A - (7 - 5i)B) \left(\int \frac{1}{-\cot(c + dx) - 1} d(1 - \sqrt{2}\sqrt{\cot(c + dx)}) \right) \right)}{d}$$

2a² 2a²

12a²

↓ 217

$$\frac{(A + iB) \cot^{\frac{7}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \frac{10a^3(6A + iB)\sqrt{\cot(c + dx)}}{d} - \frac{2a^3 \left(\left(\frac{1}{2} - \frac{i}{2} \right) ((29 + i)A + (1 + 6i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2} ((30 + 28i)A - (7 - 5i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx)} + 1)}{\sqrt{2}} - \arctan(1) \right) \right)}{d}$$

2a² 2a²

12a²

↓ 1479

$$\frac{(A + iB) \cot^{\frac{7}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \frac{10a^3(6A + iB)\sqrt{\cot(c + dx)}}{d} - \frac{2a^3 \left(\left(\frac{1}{2} - \frac{i}{2} \right) ((29 + i)A + (1 + 6i)B) \left(- \frac{\int \frac{\sqrt{2} - 2\sqrt{\cot(c + dx)}}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx)} + 1} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c + dx)} + 1)}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx)} + 1} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} \right) \right)}{d}$$

2a² 2a²

12a²

↓ 25

$$\frac{(A + iB) \cot^{\frac{7}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \frac{10a^3(6A + iB)\sqrt{\cot(c + dx)}}{d} - \frac{2a^3 \left(\left(\frac{1}{2} - \frac{i}{2} \right) ((29 + i)A + (1 + 6i)B) \left(\frac{\int \frac{\sqrt{2} - 2\sqrt{\cot(c + dx)}}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx)} + 1} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c + dx)} + 1)}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx)} + 1} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} \right) \right)}{d}$$

2a² 2a²

12a²

↓ 27

3.530. $\int \frac{\cot^{\frac{3}{2}}(c + dx)(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^3} dx$

$$\frac{(A + iB) \cot^{\frac{7}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \frac{10a^3(6A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a^3 \left(\left(\frac{1}{2} - \frac{i}{2}\right) ((29+i)A + (1+6i)B) \left(\int \frac{\sqrt{2-2\sqrt{\cot(c+dx)}}}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)} + 1} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)} + 1}{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)} + 1} d\sqrt{\cot(c+dx)} \right) \right)}{2a^2}$$

$$\frac{(A + iB) \cot^{\frac{7}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \frac{10a^3(6A+iB)\sqrt{\cot(c+dx)}}{d} - \frac{2a^3 \left(\frac{1}{2} ((30+28i)A - (7-5i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \left(\frac{1}{2} - \frac{i}{2}\right) ((29+i)A + (1+6i)B) \left(\frac{\log(\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)} + 1)}{d} \right) \right)}{2a^2}$$

12a²

↓ 1103

```
input Int[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]
```

```
output ((A + I*B)*Cot[c + d*x]^(7/2))/(6*d*(I*a + a*Cot[c + d*x])^3) - (-((a*(5*A + (2*I)*B)*Cot[c + d*x]^(5/2))/(d*(I*a + a*Cot[c + d*x])^2)) + ((-7*a^2*(4*A + I*B)*Cot[c + d*x]^(3/2))/(d*(I*a + a*Cot[c + d*x])) + (3*((10*a^3*(6*A + I*B)*Sqrt[Cot[c + d*x]])/d - (2*a^3*(((30 + 28*I)*A - (7 - 5*I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2])))/2 + (1/2 - I/2)*((29 + I)*A + (1 + 6*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/d)/(2*a^2)/(2*a^2))/(12*a^2)
```

3.530.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

3.530. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

rule 217 $\text{Int}[(a_ + (b_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{(-1)} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /;$ $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&$
 $\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 1082 $\text{Int}[(a_ + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}\{q = 1 - 4 \cdot \text{Simplify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /;$ $\text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4 \cdot a \cdot c]) /;$ $\text{FreeQ}\{a, b, c, x\}$

rule 1103 $\text{Int}[(d_ + (e_ \cdot)(x_)) / ((a_ + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$

rule 1476 $\text{Int}[(d_ + (e_ \cdot)(x_)^2) / ((a_ + (c_ \cdot)(x_)^4), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c) \ \text{Int}[1/\text{Simp}[d/e + q \cdot x + x^2, x], x], x] + \text{Simp}[e/(2 \cdot c) \ \text{Int}[1/\text{Simp}[d/e - q \cdot x + x^2, x], x], x] /;$ $\text{FreeQ}\{a, c, d, e, x\} \ \&\& \ \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{PosQ}[d \cdot e]$

rule 1479 $\text{Int}[(d_ + (e_ \cdot)(x_)^2) / ((a_ + (c_ \cdot)(x_)^4), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[-2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c \cdot q) \ \text{Int}[(q - 2 \cdot x)/\text{Simp}[d/e + q \cdot x - x^2, x], x], x] + \text{Simp}[e/(2 \cdot c \cdot q) \ \text{Int}[(q + 2 \cdot x)/\text{Simp}[d/e - q \cdot x - x^2, x], x], x] /;$ $\text{FreeQ}\{a, c, d, e, x\} \ \&\& \ \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{NegQ}[d \cdot e]$

rule 1482 $\text{Int}[(d_ + (e_ \cdot)(x_)^2) / ((a_ + (c_ \cdot)(x_)^4), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[a \cdot c, 2]\}, \text{Simp}[(d \cdot q + a \cdot e)/(2 \cdot a \cdot c) \ \text{Int}[(q + c \cdot x^2)/(a + c \cdot x^4), x], x] + \text{Simp}[(d \cdot q - a \cdot e)/(2 \cdot a \cdot c) \ \text{Int}[(q - c \cdot x^2)/(a + c \cdot x^4), x], x] /;$ $\text{FreeQ}\{a, c, d, e, x\} \ \&\& \ \text{NeQ}[c \cdot d^2 + a \cdot e^2, 0] \ \&\& \ \text{NeQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{NegQ}[(-a) \cdot c]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$ $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*(a + b*Tan[e + f*x])^m/(f*m), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4078 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

3.530.4 Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.49

method	result
derivativedivides	$\frac{-2A\sqrt{\cot(dx+c)} + \frac{4\left(-\frac{iA}{16} - \frac{B}{16}\right) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} + i\left(\frac{i(20iA-9B)\cot(dx+c)^{\frac{5}{2}} + \left(-\frac{98iA}{3} + \frac{38B}{3}\right)\cot(dx+c)^{\frac{3}{2}} + (5iB+14A)}{(i+\cot(dx+c))^3}\right)}{a^3d}$
default	$\frac{-2A\sqrt{\cot(dx+c)} + \frac{4\left(-\frac{iA}{16} - \frac{B}{16}\right) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} + i\left(\frac{i(20iA-9B)\cot(dx+c)^{\frac{5}{2}} + \left(-\frac{98iA}{3} + \frac{38B}{3}\right)\cot(dx+c)^{\frac{3}{2}} + (5iB+14A)}{(i+\cot(dx+c))^3}\right)}{a^3d}$

3.530. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

input `int(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/a^3/d*(-2*A*cot(d*x+c)^(1/2)+4*(-1/16*I*A-1/16*B)/(2^(1/2)-I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))+1/8*I*((I*(20*I*A-9*B)*cot(d*x+c)^(5/2)+(-98/3*I*A+38/3*B)*cot(d*x+c)^(3/2)+(14*A+5*I*B)*cot(d*x+c)^(1/2))/(I+cot(d*x+c))^3+2*(29*A+6*I*B)/(2^(1/2)+I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2))))`

3.530.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 688 vs. $2(276) = 552$.

Time = 0.28 (sec) , antiderivative size = 688, normalized size of antiderivative = 1.87

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx =$$

$$\left(3 a^3 d \sqrt{\frac{i A^2+2 A B-i B^2}{a^6 d^2}} e^{(6i dx+6i c)} \log \left(-\frac{2 \left((i a^3 d e^{(2i dx+2i c)}-i a^3 d) \sqrt{\frac{i e^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}-1}} \sqrt{\frac{i A^2+2 A B-i B^2}{a^6 d^2}}+(A-i B) e^{(2i dx+2i c)} \right)}{i A+B} \right) \right)$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="fracas")`

output

```
-1/96*(3*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)
*log(-2*((I*a^3*d*e^(2*I*d*x + 2*I*c) - I*a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c)
+ I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2))
+ (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*a^3*
d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(-2*((-I*
a^3*d*e^(2*I*d*x + 2*I*c) + I*a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(
2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2)) + (A - I*B)
*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*a^3*d*sqrt((-841
*I*A^2 + 348*A*B + 36*I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(1/8*((a^3*
d*e^(2*I*d*x + 2*I*c) - a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*
x + 2*I*c) - 1))*sqrt((-841*I*A^2 + 348*A*B + 36*I*B^2)/(a^6*d^2)) + 29*I*
A - 6*B)*e^(-2*I*d*x - 2*I*c)/(a^3*d)) + 3*a^3*d*sqrt((-841*I*A^2 + 348*A*
B + 36*I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(-1/8*((a^3*d*e^(2*I*d*x +
2*I*c) - a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1
))*sqrt((-841*I*A^2 + 348*A*B + 36*I*B^2)/(a^6*d^2)) - 29*I*A + 6*B)*e^(-2
*I*d*x - 2*I*c)/(a^3*d)) + 2*(2*(73*A + 10*I*B)*e^(6*I*d*x + 6*I*c) - (41*
A + 14*I*B)*e^(4*I*d*x + 4*I*c) - (8*A + 5*I*B)*e^(2*I*d*x + 2*I*c) - A -
I*B)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*e^(-6*I*
d*x - 6*I*c)/(a^3*d)
```

3.530.6 Sympy [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{i \left(\int \frac{A \cot^{\frac{3}{2}}(c+dx)}{\tan^3(c+dx) - 3i \tan^2(c+dx) - 3 \tan(c+dx) + i} dx + \int \frac{B \tan(c+dx) \cot^{\frac{3}{2}}(c+dx)}{\tan^3(c+dx) - 3i \tan^2(c+dx) - 3 \tan(c+dx) + i} dx \right)}{a^3}$$

input `integrate(cot(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)`

output `I*(Integral(A*cot(c + d*x)**(3/2)/(tan(c + d*x)**3 - 3*I*tan(c + d*x)**2 - 3*tan(c + d*x) + I), x) + Integral(B*tan(c + d*x)*cot(c + d*x)**(3/2)/(tan(c + d*x)**3 - 3*I*tan(c + d*x)**2 - 3*tan(c + d*x) + I), x))/a**3`

3.530.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.530.8 Giac [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{(ia\tan(dx+c)+a)^3} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output `integrate((B*tan(d*x+c)+A)*cot(d*x+c)^(3/2)/(I*a*tan(d*x+c)+a)^3, x)`

3.530.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx = \int \frac{\cot(c+dx)^{3/2}(A+B\tan(c+dx))}{(a+a\tan(c+dx)1i)^3} dx$$

input `int((cot(c+d*x)^(3/2)*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*1i)^3,x)`

output `int((cot(c+d*x)^(3/2)*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*1i)^3, x)`

3.530. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

3.531
$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.531.1 Optimal result 5004
 3.531.2 Mathematica [A] (verified) 5005
 3.531.3 Rubi [A] (verified) 5005
 3.531.4 Maple [A] (verified) 5011
 3.531.5 Fricas [B] (verification not implemented) 5012
 3.531.6 Sympy [F] 5013
 3.531.7 Maxima [F(-2)] 5013
 3.531.8 Giac [F(-2)] 5013
 3.531.9 Mupad [F(-1)] 5014

3.531.1 Optimal result

Integrand size = 36, antiderivative size = 318

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= -\frac{((-7+5i)A+2iB) \arctan\left(1-\sqrt{2}\sqrt{\cot(c+dx)}\right)}{16\sqrt{2}a^3d}$$

$$+\frac{((-7+5i)A+2iB) \arctan\left(1+\sqrt{2}\sqrt{\cot(c+dx)}\right)}{16\sqrt{2}a^3d} + \frac{(A+iB) \cot^{\frac{5}{2}}(c+dx)}{6d(ia+a \cot(c+dx))^3}$$

$$+\frac{(4A+iB) \cot^{\frac{3}{2}}(c+dx)}{12ad(ia+a \cot(c+dx))^2} + \frac{5A\sqrt{\cot(c+dx)}}{8d(ia^3+a^3 \cot(c+dx))}$$

$$-\frac{((7+5i)A-2iB) \log\left(1-\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{32\sqrt{2}a^3d}$$

$$+\frac{((7+5i)A-2iB) \log\left(1+\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{32\sqrt{2}a^3d}$$

```
output 1/6*(A+I*B)*cot(d*x+c)^(5/2)/d/(I*a+a*cot(d*x+c))^3+1/12*(4*A+I*B)*cot(d*x+c)^(3/2)/a/d/(I*a+a*cot(d*x+c))^2+1/32*((-7+5*I)*A+2*I*B)*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)+1/32*((-7+5*I)*A+2*I*B)*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)-1/64*((7+5*I)*A-2*I*B)*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)+1/64*((7+5*I)*A-2*I*B)*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)+5/8*A*cot(d*x+c)^(1/2)/d/(I*a^3+a^3*cot(d*x+c))
```

3.531.
$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.531.2 Mathematica [A] (verified)

Time = 4.59 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.70

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx =$$

$$\frac{i\sqrt{\cot(c+dx)}\sec^3(c+dx)\left(12\sqrt[4]{-1}(A-iB)\arctan\left((-1)^{3/4}\sqrt{\tan(c+dx)}\right)(\cos(3(c+dx))+i\sin(3(c+dx)))\right)}{(a+ia\tan(c+dx))^3}$$

input `Integrate[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])
^3,x]`

output `((-1/96*I)*Sqrt[Cot[c + d*x]]*Sec[c + d*x]^3*(12*(-1)^(1/4)*(A - I*B)*ArcT
an[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)])
+ 12*(-1)^(1/4)*(6*A - I*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*(Cos[3*
(c + d*x)] + I*Sin[3*(c + d*x)]) - 4*Cos[c + d*x]*(6*A + (3*I)*B + 3*(7*A
+ I*B)*Cos[2*(c + d*x)] + ((19*I)*A - B)*Sin[2*(c + d*x)])*Sqrt[Tan[c + d*
x]]*Sqrt[Tan[c + d*x]])/(a^3*d*(-I + Tan[c + d*x])^3)`

3.531.3 Rubi [A] (verified)Time = 1.33 (sec) , antiderivative size = 293, normalized size of antiderivative = 0.92, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.611$, Rules used = {3042, 4064, 3042, 4078, 27, 3042, 4078, 27, 3042, 4078, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A\cot(c+dx)+B)}{(a\cot(c+dx)+ia)^3} dx$$

$$\begin{aligned}
& \int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{5/2} (B - A \tan(c+dx+\frac{\pi}{2}))}{(-a \tan(c+dx+\frac{\pi}{2}) + ia)^3} dx \\
& \quad \downarrow 3042 \\
& \int \frac{-\frac{\cot^{\frac{3}{2}}(c+dx)(5a(iA-B)-a(11A-iB)\cot(c+dx))}{2(\cot(c+dx)a+ia)^2} dx}{6a^2} + \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} \\
& \quad \downarrow 4078 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{\int \frac{\cot^{\frac{3}{2}}(c+dx)(5a(iA-B)-a(11A-iB)\cot(c+dx))}{(\cot(c+dx)a+ia)^2} dx}{12a^2} \\
& \quad \downarrow 27 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{\int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{3/2} (5a(iA-B)+a(11A-iB)\tan(c+dx+\frac{\pi}{2}))}{(ia-a\tan(c+dx+\frac{\pi}{2}))^2} dx}{12a^2} \\
& \quad \downarrow 3042 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{\int \frac{6\sqrt{\cot(c+dx)}(a^2(4iA-B)-a^2(6A-iB)\cot(c+dx))}{\cot(c+dx)a+ia} dx}{4a^2} - \frac{a(4A+iB)\cot^{\frac{3}{2}}(c+dx)}{d(a\cot(c+dx)+ia)^2} \\
& \quad \downarrow 4078 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{3 \int \frac{\sqrt{\cot(c+dx)}(a^2(4iA-B)-a^2(6A-iB)\cot(c+dx))}{\cot(c+dx)a+ia} dx}{2a^2} - \frac{a(4A+iB)\cot^{\frac{3}{2}}(c+dx)}{d(a\cot(c+dx)+ia)^2} \\
& \quad \downarrow 27 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{3 \int \frac{\sqrt{-\tan(c+dx+\frac{\pi}{2})}((4iA-B)a^2+(6A-iB)\tan(c+dx+\frac{\pi}{2})a^2)}{ia-a\tan(c+dx+\frac{\pi}{2})} dx}{2a^2} - \frac{a(4A+iB)\cot^{\frac{3}{2}}(c+dx)}{d(a\cot(c+dx)+ia)^2} \\
& \quad \downarrow 3042 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{3 \left(\frac{\int \frac{5ia^3A-a^3(7A-2iB)\cot(c+dx)}{\sqrt{\cot(c+dx)}} dx}{2a^2} - \frac{5a^2A\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} \right)}{2a^2} - \frac{a(4A+iB)\cot^{\frac{3}{2}}(c+dx)}{d(a\cot(c+dx)+ia)^2} \\
& \quad \downarrow 4078 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{3 \left(\frac{\int \frac{5ia^3A-a^3(7A-2iB)\cot(c+dx)}{\sqrt{\cot(c+dx)}} dx}{2a^2} - \frac{5a^2A\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} \right)}{2a^2} - \frac{a(4A+iB)\cot^{\frac{3}{2}}(c+dx)}{d(a\cot(c+dx)+ia)^2} \\
& \quad \downarrow 3042 \\
& \frac{(A+iB)\cot^{\frac{5}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{3 \left(\frac{\int \frac{5ia^3A-a^3(7A-2iB)\cot(c+dx)}{\sqrt{\cot(c+dx)}} dx}{2a^2} - \frac{5a^2A\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} \right)}{2a^2} - \frac{a(4A+iB)\cot^{\frac{3}{2}}(c+dx)}{d(a\cot(c+dx)+ia)^2}
\end{aligned}$$

3.531. $\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^3} dx$

$$\frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \frac{\left(\frac{\int \frac{5iAa^3 + (7A - 2iB) \tan(c + dx + \frac{\pi}{2}) a^3}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx}{2a^2} - \frac{5a^2 A \sqrt{\cot(c + dx)}}{d(a \cot(c + dx) + ia)} \right)}{12a^2}$$

↓ 4017

$$\frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \frac{\left(\frac{\int \frac{a^3(5iA - (7A - 2iB) \cot(c + dx))}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{a^2 d} - \frac{5a^2 A \sqrt{\cot(c + dx)}}{d(a \cot(c + dx) + ia)} \right)}{12a^2} - \frac{a(4A + iB) \cot^{\frac{3}{2}}(c + dx)}{d(a \cot(c + dx) + ia)^2}$$

↓ 25

$$\frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \frac{\left(-\frac{\int \frac{a^3(5iA - (7A - 2iB) \cot(c + dx))}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{a^2 d} - \frac{5a^2 A \sqrt{\cot(c + dx)}}{d(a \cot(c + dx) + ia)} \right)}{12a^2} - \frac{a(4A + iB) \cot^{\frac{3}{2}}(c + dx)}{d(a \cot(c + dx) + ia)^2}$$

↓ 27

$$\frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \frac{\left(-\frac{a \int \frac{5iA - (7A - 2iB) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} - \frac{5a^2 A \sqrt{\cot(c + dx)}}{d(a \cot(c + dx) + ia)} \right)}{12a^2} - \frac{a(4A + iB) \cot^{\frac{3}{2}}(c + dx)}{d(a \cot(c + dx) + ia)^2}$$

↓ 1482

$$\frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \frac{\left(-\frac{a \left(\frac{1}{2}((7 + 5i)A - 2iB) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}(2iB - (7 - 5i)A) \int \frac{\cot(c + dx) + 1}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} \right)}{d} - \frac{5a^2 A \sqrt{\cot(c + dx)}}{d(a \cot(c + dx) + ia)} \right)}{12a^2} - \frac{a(4A + iB) \cot^{\frac{3}{2}}(c + dx)}{d(a \cot(c + dx) + ia)^2}$$

↓ 1476

$$\frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \frac{\left(-\frac{a \left(\frac{1}{2}((7 + 5i)A - 2iB) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}(2iB - (7 - 5i)A) \left(\frac{1}{2} \int \frac{1}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)} + \frac{1}{2} \int \frac{1}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)} \right) \right)}{d} - \frac{5a^2 A \sqrt{\cot(c + dx)}}{d(a \cot(c + dx) + ia)} \right)}{12a^2}$$

3.531. $\int \frac{\sqrt{\cot(c + dx)}(A + B \tan(c + dx))}{(a + ia \tan(c + dx))^3} dx$

$$\begin{aligned} & \downarrow 1082 \\ & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\ & 3 \left(\frac{a \left(\frac{1}{2}((7+5i)A - 2iB) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)+1} + \frac{1}{2}(2iB - (7-5i)A) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1 - \sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)-1} d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \right) \\ & \hline & 2a^2 \qquad \qquad \qquad 12a^2 \end{aligned}$$

$$\begin{aligned} & \downarrow 217 \\ & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\ & 3 \left(\frac{a \left(\frac{1}{2}((7+5i)A - 2iB) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)+1} + \frac{1}{2}(2iB - (7-5i)A) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}})}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{d} - \frac{5a^2 A \sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)} \right) \\ & \hline & 2a^2 \qquad \qquad \qquad 12a^2 \end{aligned}$$

$$\begin{aligned} & \downarrow 1479 \\ & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\ & 3 \left(\frac{a \left(\frac{1}{2}((7+5i)A - 2iB) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(2iB - (7-5i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}) \right) \right)}{d} \right) \\ & \hline & 2a^2 \qquad \qquad \qquad 12a^2 \end{aligned}$$

$$\begin{aligned} & \downarrow 25 \\ & \frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\ & 3 \left(\frac{a \left(\frac{1}{2}((7+5i)A - 2iB) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(2iB - (7-5i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}) \right) \right)}{d} \right) \\ & \hline & 2a^2 \qquad \qquad \qquad 12a^2 \end{aligned}$$

$$\downarrow 27$$

3.531. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$

$$\begin{array}{c}
\frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\
3 \frac{\left(a \left(\frac{1}{2}((7+5i)A - 2iB) \left(\int \frac{\sqrt{2} - 2\sqrt{\cot(c+dx)}}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)} + 1} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)} + 1}{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)} + 1} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2}(2iB - (7-5i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{d} \\
\hline
2a^2 \qquad \qquad \qquad 12a^2 \\
\downarrow 1103 \\
\frac{(A + iB) \cot^{\frac{5}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\
3 \frac{\left(-\frac{5a^2 A \sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)} - a \left(\frac{1}{2}(2iB - (7-5i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}((7+5i)A - 2iB) \left(\frac{\log(\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)})}{2\sqrt{2}} \right) \right) \right)}{d} \\
\hline
2a^2 \qquad \qquad \qquad 12a^2
\end{array}$$

input `Int[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^3,x]`

output `((A + I*B)*Cot[c + d*x]^(5/2))/(6*d*(I*a + a*Cot[c + d*x])^3) - ((a*(4*A + I*B)*Cot[c + d*x]^(3/2))/(d*(I*a + a*Cot[c + d*x])) + (3*((-5*a^2*A*Sqrt[Cot[c + d*x]])/(d*(I*a + a*Cot[c + d*x])) - (a*(((7 + 5*I)*A + (2*I)*B)*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]))/2 + (((7 + 5*I)*A - (2*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2))/d)/(2*a^2)/(12*a^2)`

3.531.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

$$3.531. \quad \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

```
rule 4064 Int[(cot[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(m_))*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp [g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4078 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(m_))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp [(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp [1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

3.531.4 Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.51

method	result
derivativedivides	$\frac{i \left(\frac{-i(2iB+9A) \cot(dx+c)^{\frac{5}{2}} + \left(\frac{2iB}{3} + \frac{38A}{3}\right) \cot(dx+c)^{\frac{3}{2}} + 5iA\sqrt{\cot(dx+c)}}{(i+\cot(dx+c))^3} + \frac{2(6iA+B) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}} \right)}{8} + \frac{4\left(-\frac{A}{16} + \frac{iB}{16}\right) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}$
default	$\frac{i \left(\frac{-i(2iB+9A) \cot(dx+c)^{\frac{5}{2}} + \left(\frac{2iB}{3} + \frac{38A}{3}\right) \cot(dx+c)^{\frac{3}{2}} + 5iA\sqrt{\cot(dx+c)}}{(i+\cot(dx+c))^3} + \frac{2(6iA+B) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}} \right)}{8} + \frac{4\left(-\frac{A}{16} + \frac{iB}{16}\right) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}$

```
input int(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)
```

```
output 1/a^3/d*(1/8*I*((-I*(2*I*B+9*A)*cot(d*x+c)^(5/2)+(2/3*I*B+38/3*A)*cot(d*x+c)^(3/2)+5*I*A*cot(d*x+c)^(1/2))/(I+cot(d*x+c))^3+2*(6*I*A+B)/(2^(1/2)+I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2))))+4*(-1/16*A+1/16*I*B)/(2^(1/2)-I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))
```

$$3.531. \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

3.531.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 683 vs. $2(247) = 494$.

Time = 0.27 (sec) , antiderivative size = 683, normalized size of antiderivative = 2.15

$$\int \frac{\sqrt{\cot(c+dx)(A+B \tan(c+dx))}}{(a+ia \tan(c+dx))^3} dx$$

$$= \left(3 a^3 d \sqrt{\frac{-i A^2 - 2 A B + i B^2}{a^6 d^2}} e^{(6i dx + 6i c)} \log \left(-\frac{2 \left((a^3 d e^{(2i dx + 2i c)} - a^3 d) \sqrt{\frac{i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} - 1}} \sqrt{\frac{-i A^2 - 2 A B + i B^2}{a^6 d^2}} + (A - i B) e^{(2i dx + 2i c)} \right)}{i A + B} \right) \right)$$

```
input integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorith
thm="fricas")
```

```
output 1/96*(3*a^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)
*log(-2*((a^3*d*e^(2*I*d*x + 2*I*c) - a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c) +
I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2)) +
(A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*a^3*d*s
qrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(2*((a^3*d*
e^(2*I*d*x + 2*I*c) - a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x
+ 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2)) - (A - I*B)*e^(2*I
*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*a^3*d*sqrt((36*I*A^2 +
12*A*B - I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(-1/8*((a^3*d*e^(2*I*d*x
+ 2*I*c) - a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) -
1))*sqrt((36*I*A^2 + 12*A*B - I*B^2)/(a^6*d^2)) + 6*A - I*B)*e^(-2*I*d*x
- 2*I*c)/(a^3*d)) + 3*a^3*d*sqrt((36*I*A^2 + 12*A*B - I*B^2)/(a^6*d^2))*e^
(6*I*d*x + 6*I*c)*log(1/8*((a^3*d*e^(2*I*d*x + 2*I*c) - a^3*d)*sqrt((I*e^
(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((36*I*A^2 + 12*A*B -
I*B^2)/(a^6*d^2)) - 6*A + I*B)*e^(-2*I*d*x - 2*I*c)/(a^3*d)) - 2*(2*(10*I
*A - B)*e^(6*I*d*x + 6*I*c) - (14*I*A + B)*e^(4*I*d*x + 4*I*c) - (5*I*A -
2*B)*e^(2*I*d*x + 2*I*c) - I*A + B)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2
*I*d*x + 2*I*c) - 1)))e^(-6*I*d*x - 6*I*c)/(a^3*d)
```

3.531.6 Sympy [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx$$

$$= \frac{i \left(\int \frac{A\sqrt{\cot(c+dx)}}{\tan^3(c+dx)-3i \tan^2(c+dx)-3 \tan(c+dx)+i} dx + \int \frac{B \tan(c+dx)\sqrt{\cot(c+dx)}}{\tan^3(c+dx)-3i \tan^2(c+dx)-3 \tan(c+dx)+i} dx \right)}{a^3}$$

input `integrate(cot(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**3,x)`

output `I*(Integral(A*sqrt(cot(c + d*x))/(tan(c + d*x)**3 - 3*I*tan(c + d*x)**2 - 3*tan(c + d*x) + I), x) + Integral(B*tan(c + d*x)*sqrt(cot(c + d*x))/(tan(c + d*x)**3 - 3*I*tan(c + d*x)**2 - 3*tan(c + d*x) + I), x))/a**3`

3.531.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.531.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \text{Exception raised: TypeError}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:Not invertible Error: Bad Argument
Value

3.531.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^3} dx = \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+a \tan(c+dx) 1i)^3} dx$$

input `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3,x)`

output `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^3, x)`
)

3.532 $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+ia \tan(c+dx))^3}} dx$

3.532.1 Optimal result 5015
 3.532.2 Mathematica [A] (verified) 5016
 3.532.3 Rubi [A] (verified) 5016
 3.532.4 Maple [A] (verified) 5022
 3.532.5 Fricas [B] (verification not implemented) 5023
 3.532.6 Sympy [F] 5024
 3.532.7 Maxima [F(-2)] 5024
 3.532.8 Giac [F] 5024
 3.532.9 Mupad [F(-1)] 5025

3.532.1 Optimal result

Integrand size = 36, antiderivative size = 316

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))^3}} dx$$

$$= -\frac{\left(\frac{1}{16} + \frac{i}{16}\right) \left((1 + i)A + B\right) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}a^3d}$$

$$+ \frac{\left(\frac{1}{16} + \frac{i}{16}\right) \left((1 + i)A + B\right) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}a^3d}$$

$$+ \frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{6d(ia + a \cot(c + dx))^3} + \frac{A\sqrt{\cot(c + dx)}}{4ad(ia + a \cot(c + dx))^2} + \frac{(2iA + B)\sqrt{\cot(c + dx)}}{8d(ia^3 + a^3 \cot(c + dx))}$$

$$+ \frac{(2iA + (1 - i)B) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{32\sqrt{2}a^3d}$$

$$- \frac{(2iA + (1 - i)B) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{32\sqrt{2}a^3d}$$

output

```
1/6*(A+I*B)*cot(d*x+c)^(3/2)/d/(I*a+a*cot(d*x+c))^3+(1/32+1/32*I)*((1+I)*A
+B)*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)+(1/32+1/32*I)*((1+I)
*A+B)*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)+1/64*(2*I*A+(1-I)*B
)*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)-1/64*(2*I*A+(1-I)
)*B)*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)+1/4*A*cot(d*x
+c)^(1/2)/a/d/(I*a+a*cot(d*x+c))^2+1/8*(2*I*A+B)*cot(d*x+c)^(1/2)/d/(I*a^3
+a^3*cot(d*x+c))
```

3.532. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+ia \tan(c+dx))^3}} dx$

3.532.2 Mathematica [A] (verified)

Time = 5.79 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.62

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^3} dx =$$

$$\frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(3\sqrt[4]{-1} (A - iB) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right) \sec^3(c + dx) (\cos(3(c + dx))) \right)}{a^3 d (-I + \tan(c + dx))^3}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^3),x]`

output `-1/24*(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(3*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^3*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)]) + 3*(-1)^(1/4)*A*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^3*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)]) + Sqrt[Tan[c + d*x]]*(-3*I + Tan[c + d*x])*((2*I)*A + B + (3*I)*B*Tan[c + d*x]))/(a^3*d*(-I + Tan[c + d*x])^3)`

3.532.3 Rubi [A] (verified)Time = 1.29 (sec) , antiderivative size = 291, normalized size of antiderivative = 0.92, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.611$, Rules used = {3042, 4064, 3042, 4078, 27, 3042, 4078, 27, 3042, 4079, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^3} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{\cot^{\frac{3}{2}}(c + dx)(A \cot(c + dx) + B)}{(a \cot(c + dx) + ia)^3} dx$$

3.532. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{3/2} (B-A \tan(c+dx+\frac{\pi}{2}))}{(-a \tan(c+dx+\frac{\pi}{2})+ia)^3} dx \\
& \downarrow 4078 \\
& \frac{\int -\frac{3\sqrt{\cot(c+dx)}(a(iA-B)-a(3A-iB)\cot(c+dx))}{2(\cot(c+dx)a+ia)^2} dx}{6a^2} + \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{6d(a \cot(c+dx)+ia)^3} \\
& \downarrow 27 \\
& \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{6d(a \cot(c+dx)+ia)^3} - \frac{\int \frac{\sqrt{\cot(c+dx)}(a(iA-B)-a(3A-iB)\cot(c+dx))}{(\cot(c+dx)a+ia)^2} dx}{4a^2} \\
& \downarrow 3042 \\
& \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{6d(a \cot(c+dx)+ia)^3} - \frac{\int \frac{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(a(iA-B)+a(3A-iB)\tan(c+dx+\frac{\pi}{2}))}{(ia-a \tan(c+dx+\frac{\pi}{2}))^2} dx}{4a^2} \\
& \downarrow 4078 \\
& \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{6d(a \cot(c+dx)+ia)^3} - \frac{\int \frac{2(ia^2A-a^2(3A-2iB)\cot(c+dx))}{\sqrt{\cot(c+dx)}(\cot(c+dx)a+ia)} dx}{4a^2} - \frac{aA\sqrt{\cot(c+dx)}}{d(a \cot(c+dx)+ia)^2} \\
& \downarrow 27 \\
& \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{6d(a \cot(c+dx)+ia)^3} - \frac{\int \frac{ia^2A-a^2(3A-2iB)\cot(c+dx)}{\sqrt{\cot(c+dx)}(\cot(c+dx)a+ia)} dx}{2a^2} - \frac{aA\sqrt{\cot(c+dx)}}{d(a \cot(c+dx)+ia)^2} \\
& \downarrow 3042 \\
& \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{6d(a \cot(c+dx)+ia)^3} - \frac{\int \frac{ia^2+(3A-2iB)\tan(c+dx+\frac{\pi}{2})a^2}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(ia-a \tan(c+dx+\frac{\pi}{2}))} dx}{2a^2} - \frac{aA\sqrt{\cot(c+dx)}}{d(a \cot(c+dx)+ia)^2} \\
& \downarrow 4079 \\
& \frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{6d(a \cot(c+dx)+ia)^3} - \frac{\int \frac{iBa^3+(2iA+B)\cot(c+dx)a^3}{\sqrt{\cot(c+dx)}} dx}{2a^2} - \frac{a^2(B+2iA)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx)+ia)} - \frac{aA\sqrt{\cot(c+dx)}}{d(a \cot(c+dx)+ia)^2} \\
& \downarrow 3042
\end{aligned}$$

3.532. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^3} dx$

$$\frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{\int \frac{ia^3B - a^3(2iA+B)\tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{2a^2} - \frac{a^2(B+2iA)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} - \frac{aA\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2}$$

↓ 4017

$$\frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{\int -\frac{a^3(iB+(2iA+B)\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{a^2d} - \frac{a^2(B+2iA)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} - \frac{aA\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2}$$

↓ 25

$$\frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{\int \frac{a^3(iB+(2iA+B)\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{a^2d} - \frac{a^2(B+2iA)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} - \frac{aA\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2}$$

↓ 27

$$\frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{a \int \frac{iB+(2iA+B)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} - \frac{a^2(B+2iA)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} - \frac{aA\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2}$$

↓ 1482

$$\frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{a\left(\left(\frac{1}{2}+\frac{i}{2}\right)(B+(1+i)A) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(2iA+(1-i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{2a^2} - \frac{a^2(B+2iA)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} - \frac{aA\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2}$$

↓ 1476

$$\frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{a\left(\left(\frac{1}{2}+\frac{i}{2}\right)(B+(1+i)A)\left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}\right) - \frac{1}{2}(2iA+(1-i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{2a^2} - \frac{a^2(B+2iA)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} - \frac{aA\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2}$$

↓ 1082

$$\frac{(A+iB)\cot^{\frac{3}{2}}(c+dx)}{6d(a\cot(c+dx)+ia)^3} - \frac{a\left(\left(\frac{1}{2}+\frac{i}{2}\right)(B+(1+i)A)\left(\frac{\int \frac{1}{\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{\cot(c+dx)-1} d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}}\right) - \frac{1}{2}(2iA+(1-i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{2a^2} - \frac{a^2(B+2iA)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} - \frac{aA\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2}$$

3.532. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+ia \tan(c+dx))^3}} dx$

$$\begin{aligned} & \downarrow 217 \\ & \frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\ & \frac{a \left(\left(\frac{1}{2} + \frac{i}{2} \right) (B + (1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} (2iA + (1-i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d} - \frac{a^2 (B+2iA) \sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)} \\ & \frac{\hspace{10em}}{2a^2} \\ & \frac{\hspace{10em}}{4a^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 1479 \\ & \frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\ & \frac{a \left(\left(\frac{1}{2} + \frac{i}{2} \right) (B + (1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} (2iA + (1-i)B) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} - \int \frac{\sqrt{2}+2\sqrt{\cot(c+dx)}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{d} \\ & \frac{\hspace{10em}}{2a^2} \\ & \frac{\hspace{10em}}{4a^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 25 \\ & \frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\ & \frac{a \left(\left(\frac{1}{2} + \frac{i}{2} \right) (B + (1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} (2iA + (1-i)B) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \int \frac{\sqrt{2}+2\sqrt{\cot(c+dx)}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{d} \\ & \frac{\hspace{10em}}{2a^2} \\ & \frac{\hspace{10em}}{4a^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\ & \frac{a \left(\left(\frac{1}{2} + \frac{i}{2} \right) (B + (1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} (2iA + (1-i)B) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}+2\sqrt{\cot(c+dx)}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{d} \\ & \frac{\hspace{10em}}{2a^2} \\ & \frac{\hspace{10em}}{4a^2} \end{aligned}$$

$$\begin{aligned} & \downarrow 1103 \\ & \frac{(A + iB) \cot^{\frac{3}{2}}(c + dx)}{6d(a \cot(c + dx) + ia)^3} - \\ & \frac{a \left(\left(\frac{1}{2} + \frac{i}{2} \right) (B + (1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} (2iA + (1-i)B) \left(\frac{\log(\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d} - \frac{a^2 (B+2iA) \sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)} \\ & \frac{\hspace{10em}}{2a^2} \\ & \frac{\hspace{10em}}{4a^2} \end{aligned}$$

3.532. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^3} dx$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^3),x]`

output `((A + I*B)*Cot[c + d*x]^(3/2))/(6*d*(I*a + a*Cot[c + d*x])^3) - (-((a*A*Sqrt[Cot[c + d*x]])/(d*(I*a + a*Cot[c + d*x])^2)) + (-((a^2*((2*I)*A + B)*Sqrt[Cot[c + d*x]])/(d*(I*a + a*Cot[c + d*x]))) - (a*((1/2 + I/2)*((1 + I)*A + B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) - (((2*I)*A + (1 - I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d)/(2*a^2))/(4*a^2)`

3.532.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

3.532.
$$\int \frac{A+B \tan (c+d x)}{\sqrt{\cot (c+d x)\left(a+i a \tan (c+d x)\right)^3}} d x$$

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.532.4 Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.49

method	result
derivativedivides	$\frac{4\left(\frac{iA}{16} + \frac{B}{16}\right) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} + \frac{i\left(\frac{-i(2iA+B)\cot(dx+c)^{\frac{5}{2}} + \left(\frac{2iA}{3} + \frac{10B}{3}\right)\cot(dx+c)^{\frac{3}{2}} + iB\sqrt{\cot(dx+c)}}{(i+\cot(dx+c))^3} + \frac{2A \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}\right)}{8a^3d}$
default	$\frac{4\left(\frac{iA}{16} + \frac{B}{16}\right) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}} + \frac{i\left(\frac{-i(2iA+B)\cot(dx+c)^{\frac{5}{2}} + \left(\frac{2iA}{3} + \frac{10B}{3}\right)\cot(dx+c)^{\frac{3}{2}} + iB\sqrt{\cot(dx+c)}}{(i+\cot(dx+c))^3} + \frac{2A \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{\sqrt{2+i\sqrt{2}}}\right)}{8a^3d}$

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/a^3/d*(4*(1/16*I*A+1/16*B)/(2^(1/2)-I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))+1/8*I*((-I*(2*I*A+B)*cot(d*x+c)^(5/2)+(2/3*I*A+10/3*B)*cot(d*x+c)^(3/2)+I*B*cot(d*x+c)^(1/2))/(I+cot(d*x+c))^3+2*A/(2^(1/2)+I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2))))`

3.532.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 635 vs. $2(247) = 494$.

Time = 0.28 (sec) , antiderivative size = 635, normalized size of antiderivative = 2.01

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))^3}} dx$$

$$\left(3a^3 d \sqrt{\frac{iA^2 + 2AB - iB^2}{a^6 d^2}} e^{(6i dx + 6i c)} \log \left(-\frac{2 \left((i a^3 d e^{(2i dx + 2i c)} - i a^3 d) \sqrt{\frac{i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} - 1}} \sqrt{\frac{iA^2 + 2AB - iB^2}{a^6 d^2}} + (A - iB) e^{(2i dx + 2i c)} \right)}{iA + B} \right) \right)$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^3,x, algorith
thm="fricas")
```

```
output 1/96*(3*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*
log(-2*((I*a^3*d*e^(2*I*d*x + 2*I*c) - I*a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c)
) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2))
+ (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*a^3*d
*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(-2*((-I*a
^3*d*e^(2*I*d*x + 2*I*c) + I*a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2
*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2)) + (A - I*B)*
e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) + 24*a^3*d*sqrt(-1/64
*I*A^2/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(1/8*(8*(a^3*d*e^(2*I*d*x + 2*I*c)
) - a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sq
rt(-1/64*I*A^2/(a^6*d^2)) + I*A)*e^(-2*I*d*x - 2*I*c)/(a^3*d)) - 24*a^3*d*s
qrt(-1/64*I*A^2/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(-1/8*(8*(a^3*d*e^(2*I*d
*x + 2*I*c) - a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c)
- 1))*sqrt(-1/64*I*A^2/(a^6*d^2)) - I*A)*e^(-2*I*d*x - 2*I*c)/(a^3*d)) +
2*(2*(A - 2*I*B)*e^(6*I*d*x + 6*I*c) + (A + 4*I*B)*e^(4*I*d*x + 4*I*c) - (
2*A - I*B)*e^(2*I*d*x + 2*I*c) - A - I*B)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)
/(e^(2*I*d*x + 2*I*c) - 1))*e^(-6*I*d*x - 6*I*c)/(a^3*d)
```

3.532.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^3} dx$$

$$= \frac{i \left(\int \frac{A}{\tan^3(c+dx)\sqrt{\cot(c+dx)} - 3i \tan^2(c+dx)\sqrt{\cot(c+dx)} - 3 \tan(c+dx)\sqrt{\cot(c+dx)} + i\sqrt{\cot(c+dx)}} dx + \int \frac{B}{\tan^3(c+dx)\sqrt{\cot(c+dx)} - 3i \tan^2(c+dx)\sqrt{\cot(c+dx)} - 3 \tan(c+dx)\sqrt{\cot(c+dx)} + i\sqrt{\cot(c+dx)}} dx \right)}{a^3}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(1/2)/(a+I*a*tan(d*x+c))**3,x)`

output `I*(Integral(A/(tan(c + d*x)**3*sqrt(cot(c + d*x)) - 3*I*tan(c + d*x)**2*sqrt(cot(c + d*x)) - 3*tan(c + d*x)*sqrt(cot(c + d*x)) + I*sqrt(cot(c + d*x))), x) + Integral(B*tan(c + d*x)/(tan(c + d*x)**3*sqrt(cot(c + d*x)) - 3*I*tan(c + d*x)**2*sqrt(cot(c + d*x)) - 3*tan(c + d*x)*sqrt(cot(c + d*x)) + I*sqrt(cot(c + d*x))), x))/a**3`

3.532.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.532.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^3} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^3 \sqrt{\cot(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)^3*sqrt(cot(d*x + c))), x)`

3.532.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^3} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + a \tan(c + dx) i)^3} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + a*tan(c + d*x)*i)^3),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + a*tan(c + d*x)*i)^3), x)`

3.533
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$$

3.533.1 Optimal result	5026
3.533.2 Mathematica [A] (verified)	5027
3.533.3 Rubi [A] (verified)	5027
3.533.4 Maple [A] (verified)	5033
3.533.5 Fricas [B] (verification not implemented)	5034
3.533.6 Sympy [F(-1)]	5035
3.533.7 Maxima [F(-2)]	5035
3.533.8 Giac [F]	5035
3.533.9 Mupad [F(-1)]	5036

3.533.1 Optimal result

Integrand size = 36, antiderivative size = 308

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx \\ &= -\frac{((1 + i)A + 2B) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{16\sqrt{2}a^3d} \\ & \quad + \frac{((1 + i)A + 2B) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{16\sqrt{2}a^3d} + \frac{(A + iB)\sqrt{\cot(c + dx)}}{6d(ia + a \cot(c + dx))^3} \\ & \quad + \frac{(2iA + B)\sqrt{\cot(c + dx)}}{12ad(ia + a \cot(c + dx))^2} + \frac{A\sqrt{\cot(c + dx)}}{8d(ia^3 + a^3 \cot(c + dx))} \\ & \quad - \frac{((-1 + i)A + 2B) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{32\sqrt{2}a^3d} \\ & \quad + \frac{((-1 + i)A + 2B) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{32\sqrt{2}a^3d} \end{aligned}$$

output

```
1/32*((1+I)*A+2*B)*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)+1/32*
((1+I)*A+2*B)*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)-1/64*((-1+I)
)*A+2*B)*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)+1/64*((-1
+I)*A+2*B)*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)+1/6*(A+
I*B)*cot(d*x+c)^(1/2)/d/(I*a+a*cot(d*x+c))^3+1/12*(2*I*A+B)*cot(d*x+c)^(1/
2)/a/d/(I*a+a*cot(d*x+c))^2+1/8*A*cot(d*x+c)^(1/2)/d/(I*a^3+a^3*cot(d*x+c)
)
```

3.533.
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$$

3.533.2 Mathematica [A] (verified)

Time = 4.33 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.66

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx$$

$$= \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(3\sqrt[4]{-1} (iA + B) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right) \sec^3(c + dx) (\cos(3(c + dx))) \right)}{24a^3 d (-I + \tan(c + dx))^3}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^3),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(3*(-1)^(1/4)*(I*A + B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^3*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)]) - 3*(-1)^(1/4)*B*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^3*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)]) - I*Sqrt[Tan[c + d*x]]*(-I + 3*Tan[c + d*x])*((-3*I)*A + (A - (2*I)*B)*Tan[c + d*x]))/(24*a^3*d*(-I + Tan[c + d*x])^3)`

3.533.3 Rubi [A] (verified)Time = 1.28 (sec) , antiderivative size = 287, normalized size of antiderivative = 0.93, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.611$, Rules used = {3042, 4064, 3042, 4078, 27, 3042, 4079, 27, 3042, 4079, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2}(a + ia \tan(c + dx))^3} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{\sqrt{\cot(c + dx)}(A \cot(c + dx) + B)}{(a \cot(c + dx) + ia)^3} dx$$

3.533. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(B-A \tan\left(c+dx+\frac{\pi}{2}\right)\right)}{\left(-a \tan\left(c+dx+\frac{\pi}{2}\right)+ia\right)^3} dx \\
& \int -\frac{a(iA-B)-a(7A-5iB) \cot(c+dx)}{2\sqrt{\cot(c+dx)}(\cot(c+dx)a+ia)^2} dx + \frac{(A+iB)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx)+ia)^3} \\
& \frac{(A+iB)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx)+ia)^3} - \frac{\int \frac{a(iA-B)-a(7A-5iB) \cot(c+dx)}{\sqrt{\cot(c+dx)}(\cot(c+dx)a+ia)^2} dx}{12a^2} \\
& \frac{(A+iB)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx)+ia)^3} - \frac{\int \frac{a(iA-B)+a(7A-5iB) \tan\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(ia-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)^2} dx}{12a^2} \\
& \frac{(A+iB)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx)+ia)^3} - \frac{\int \frac{6\left(iBa^2+(2iA+B) \cot(c+dx)a^2\right)}{\sqrt{\cot(c+dx)}(\cot(c+dx)a+ia)} dx}{4a^2} - \frac{a(B+2iA)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx)+ia)^2} \\
& \frac{(A+iB)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx)+ia)^3} - \frac{3 \int \frac{iBa^2+(2iA+B) \cot(c+dx)a^2}{\sqrt{\cot(c+dx)}(\cot(c+dx)a+ia)} dx}{2a^2} - \frac{a(B+2iA)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx)+ia)^2} \\
& \frac{(A+iB)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx)+ia)^3} - \frac{3 \int \frac{ia^2B-a^2(2iA+B) \tan\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(ia-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)} dx}{2a^2} - \frac{a(B+2iA)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx)+ia)^2} \\
& \frac{(A+iB)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx)+ia)^3} - \frac{3\left(\frac{\int \frac{(iA+2B)a^3+A \cot(c+dx)a^3}{\sqrt{\cot(c+dx)}} dx}{2a^2} - \frac{a^2 A \sqrt{\cot(c+dx)}}{d(a \cot(c+dx)+ia)}\right)}{2a^2} - \frac{a(B+2iA)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx)+ia)^2}
\end{aligned}$$

3.533. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
 & \frac{(A + iB)\sqrt{\cot(c + dx)}}{6d(a \cot(c + dx) + ia)^3} - \frac{3 \left(\frac{\int \frac{a^3(iA+2B) - a^3 A \tan\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}} dx}{2a^2} - \frac{a^2 A \sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)} \right)}{12a^2} - \frac{a(B+2iA)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)^2} \\
 & \quad \downarrow 4017 \\
 & \frac{(A + iB)\sqrt{\cot(c + dx)}}{6d(a \cot(c + dx) + ia)^3} - \frac{3 \left(\frac{\int -\frac{a^3(\cot(c+dx)A+iA+2B)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{a^2 d} - \frac{a^2 A \sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)} \right)}{12a^2} - \frac{a(B+2iA)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)^2} \\
 & \quad \downarrow 25 \\
 & \frac{(A + iB)\sqrt{\cot(c + dx)}}{6d(a \cot(c + dx) + ia)^3} - \frac{3 \left(-\frac{\int \frac{a^3(\cot(c+dx)A+iA+2B)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{a^2 d} - \frac{a^2 A \sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)} \right)}{12a^2} - \frac{a(B+2iA)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)^2} \\
 & \quad \downarrow 27 \\
 & \frac{(A + iB)\sqrt{\cot(c + dx)}}{6d(a \cot(c + dx) + ia)^3} - \frac{3 \left(-\frac{a \int \frac{\cot(c+dx)A+iA+2B}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} - \frac{a^2 A \sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)} \right)}{12a^2} - \frac{a(B+2iA)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)^2} \\
 & \quad \downarrow 1482 \\
 & \frac{(A + iB)\sqrt{\cot(c + dx)}}{6d(a \cot(c + dx) + ia)^3} - \frac{3 \left(-\frac{a \left(\frac{1}{2}(2B - (1-i)A) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(2B + (1+i)A) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d} - \frac{a^2 A \sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)} \right)}{12a^2} - \frac{a(B+2iA)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)^2} \\
 & \quad \downarrow 1476 \\
 & \frac{(A + iB)\sqrt{\cot(c + dx)}}{6d(a \cot(c + dx) + ia)^3} - \frac{3 \left(-\frac{a \left(\frac{1}{2}(2B - (1-i)A) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(2B + (1+i)A) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{d} - \frac{a^2 A \sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)} \right)}{12a^2} - \frac{a(B+2iA)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)^2} \\
 & \quad \downarrow 1082
 \end{aligned}$$

3.533. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\frac{(A + iB)\sqrt{\cot(c + dx)}}{6d(a \cot(c + dx) + ia)^3} - \frac{a \left(\frac{1}{2}(2B - (1-i)A) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(2B + (1+i)A) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1 - \sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)-1} d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \frac{12a^2}{2a^2}$$

217

$$\frac{(A + iB)\sqrt{\cot(c + dx)}}{6d(a \cot(c + dx) + ia)^3} - \frac{a \left(\frac{1}{2}(2B - (1-i)A) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(2B + (1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{d} - \frac{a^2 A \sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)} \frac{12a^2}{2a^2}$$

1479

$$\frac{(A + iB)\sqrt{\cot(c + dx)}}{6d(a \cot(c + dx) + ia)^3} - \frac{a \left(\frac{1}{2}(2B - (1-i)A) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(2B + (1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \frac{12a^2}{2a^2}$$

25

$$\frac{(A + iB)\sqrt{\cot(c + dx)}}{6d(a \cot(c + dx) + ia)^3} - \frac{a \left(\frac{1}{2}(2B - (1-i)A) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(2B + (1+i)A) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \frac{12a^2}{2a^2}$$

27

3.533. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\frac{(A + iB)\sqrt{\cot(c + dx)}}{6d(a \cot(c + dx) + ia)^3} - \frac{a \left(\frac{1}{2}(2B - (1-i)A) \left(\int \frac{\sqrt{2} - 2\sqrt{\cot(c+dx)}}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)} + 1} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)} + 1}{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)} + 1} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2}(2B + (1+i)A) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right)}{2a^2} \right)}{12a^2}$$

↓ 1103

$$\frac{(A + iB)\sqrt{\cot(c + dx)}}{6d(a \cot(c + dx) + ia)^3} - \frac{a \left(\frac{1}{2}(2B + (1+i)A) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)} + 1}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1 - \sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) + \frac{1}{2}(2B - (1-i)A) \left(\frac{\log\left(\frac{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)} + 1}{2\sqrt{2}}\right)}{\sqrt{2}} \right) \right) - \frac{a^2 A \sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)}}{2a^2} \right)}{12a^2}$$

```
input Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^3),x]
```

```
output ((A + I*B)*Sqrt[Cot[c + d*x]])/(6*d*(I*a + a*Cot[c + d*x])^3) - (-((a*((2*I)*A + B)*Sqrt[Cot[c + d*x]])/(d*(I*a + a*Cot[c + d*x])^2)) + (3*(-((a^2*A*Sqrt[Cot[c + d*x]])/(d*(I*a + a*Cot[c + d*x]))) - (a*(((1 + I)*A + 2*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2])))/2 + (((-1 + I)*A + 2*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/d)/(2*a^2))/(12*a^2)
```

3.533.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

3.533. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])]; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(m_))*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp [g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^(m*(d + c *Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !Integer Q[p] && IntegerQ[m] && IntegerQ[n]`

rule 4078 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(m_))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp [(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a *A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(m_))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Simp [(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.533.4 Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.49

method	result
derivativedivides	$\frac{-A \cot(dx+c)^{\frac{5}{2}} + \left(-\frac{10iA}{3} - \frac{2B}{3}\right) \cot(dx+c)^{\frac{3}{2}} + (-2iB+A)\sqrt{\cot(dx+c)}}{8(i+\cot(dx+c))^3} + \frac{iB \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{4\sqrt{2+4i\sqrt{2}}} + \frac{4\left(\frac{A}{16} - \frac{iB}{16}\right) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}}$
default	$\frac{-A \cot(dx+c)^{\frac{5}{2}} + \left(-\frac{10iA}{3} - \frac{2B}{3}\right) \cot(dx+c)^{\frac{3}{2}} + (-2iB+A)\sqrt{\cot(dx+c)}}{8(i+\cot(dx+c))^3} + \frac{iB \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{4\sqrt{2+4i\sqrt{2}}} + \frac{4\left(\frac{A}{16} - \frac{iB}{16}\right) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}}$

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

$$3.533. \int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$$

output $1/a^3/d*(-1/8*(-A*\cot(d*x+c)^(5/2)+(-10/3*I*A-2/3*B)*\cot(d*x+c)^(3/2)+(A-2*I*B)*\cot(d*x+c)^(1/2))/(I+\cot(d*x+c))^3+1/4*I*B/(2^(1/2)+I*2^(1/2))*\arctan(2*\cot(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))+4*(1/16*A-1/16*I*B)/(2^(1/2)-I*2^(1/2))*\arctan(2*\cot(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))$

3.533.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 637 vs. $2(247) = 494$.

Time = 0.26 (sec) , antiderivative size = 637, normalized size of antiderivative = 2.07

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx =$$

$$\left(3 a^3 d \sqrt{\frac{-i A^2 - 2 A B + i B^2}{a^6 d^2}} e^{(6i dx + 6i c)} \log \left(-\frac{2 \left((a^3 d e^{(2i dx + 2i c)} - a^3 d) \sqrt{\frac{i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} - 1}} \sqrt{\frac{-i A^2 - 2 A B + i B^2}{a^6 d^2}} + (A - i B) e^{(2i dx + 2i c)} \right)}{i A + B} \right) \right)$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="fracas")`

output $-1/96*(3*a^3*d*\sqrt{((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2))}*e^{(6*I*d*x + 6*I*c)}*\log(-2*((a^3*d*e^{(2*I*d*x + 2*I*c)} - a^3*d)*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}*\sqrt{((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2))} + (A - I*B)*e^{(2*I*d*x + 2*I*c)}*e^{(-2*I*d*x - 2*I*c)/(I*A + B)}) - 3*a^3*d*\sqrt{((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2))}*e^{(6*I*d*x + 6*I*c)}*\log(2*((a^3*d*e^{(2*I*d*x + 2*I*c)} - a^3*d)*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}*\sqrt{((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2))} - (A - I*B)*e^{(2*I*d*x + 2*I*c)}*e^{(-2*I*d*x - 2*I*c)/(I*A + B)}) - 24*a^3*d*\sqrt{(-1/64*I*B^2/(a^6*d^2))}*e^{(6*I*d*x + 6*I*c)}*\log(1/8*(8*(a^3*d*e^{(2*I*d*x + 2*I*c)} - a^3*d)*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}*\sqrt{(-1/64*I*B^2/(a^6*d^2))} + I*B)*e^{(-2*I*d*x - 2*I*c)/(a^3*d)} + 24*a^3*d*\sqrt{(-1/64*I*B^2/(a^6*d^2))}*e^{(6*I*d*x + 6*I*c)}*\log(-1/8*(8*(a^3*d*e^{(2*I*d*x + 2*I*c)} - a^3*d)*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}*\sqrt{(-1/64*I*B^2/(a^6*d^2))} - I*B)*e^{(-2*I*d*x - 2*I*c)/(a^3*d)} + 2*(2*(2*I*A + B)*e^{(6*I*d*x + 6*I*c)} - (4*I*A + 5*B)*e^{(4*I*d*x + 4*I*c)} - (I*A - 4*B)*e^{(2*I*d*x + 2*I*c)} + I*A - B)*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)))*e^{(-6*I*d*x - 6*I*c)/(a^3*d)}$

3.533.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(3/2)/(a+I*a*tan(d*x+c))**3,x)`

output `Timed out`

3.533.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.533.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^3 \cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)^3*cot(d*x + c)^(3/2)), x)`

3.533.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2} (a + a \tan(c + dx) li)^3} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + a*tan(c + d*x)*li)^3),x)`output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + a*tan(c + d*x)*li)^3), x)`

3.534 $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

3.534.1 Optimal result 5037
 3.534.2 Mathematica [A] (verified) 5038
 3.534.3 Rubi [A] (verified) 5038
 3.534.4 Maple [A] (verified) 5044
 3.534.5 Fricas [B] (verification not implemented) 5045
 3.534.6 Sympy [F(-1)] 5046
 3.534.7 Maxima [F(-2)] 5046
 3.534.8 Giac [F] 5046
 3.534.9 Mupad [F(-1)] 5047

3.534.1 Optimal result

Integrand size = 36, antiderivative size = 310

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx$$

$$= -\frac{(2A + (5 - 7i)B) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{16\sqrt{2}a^3d}$$

$$+ \frac{(2A + (5 - 7i)B) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{16\sqrt{2}a^3d} + \frac{(iA - B)\sqrt{\cot(c + dx)}}{6d(ia + a \cot(c + dx))^3}$$

$$+ \frac{(A + 4iB)\sqrt{\cot(c + dx)}}{12ad(ia + a \cot(c + dx))^2} + \frac{5B\sqrt{\cot(c + dx)}}{8d(ia^3 + a^3 \cot(c + dx))}$$

$$- \frac{(2A - (5 + 7i)B) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{32\sqrt{2}a^3d}$$

$$+ \frac{(2A - (5 + 7i)B) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{32\sqrt{2}a^3d}$$

output

```
1/32*(2*A+(5-7*I)*B)*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)+1/3
2*(2*A+(5-7*I)*B)*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)-1/64*(2
*A-(5+7*I)*B)*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)+1/64
*(2*A-(5+7*I)*B)*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/a^3/d*2^(1/2)+1
/6*(I*A-B)*cot(d*x+c)^(1/2)/d/(I*a+a*cot(d*x+c))^3+1/12*(A+4*I*B)*cot(d*x+
c)^(1/2)/a/d/(I*a+a*cot(d*x+c))^2+5/8*B*cot(d*x+c)^(1/2)/d/(I*a^3+a^3*cot(
d*x+c))
```

3.534. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

3.534.2 Mathematica [A] (verified)

Time = 5.76 (sec) , antiderivative size = 217, normalized size of antiderivative = 0.70

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx =$$

$$\frac{\sqrt{\cot(c + dx)} \sec^3(c + dx) \left(-3\sqrt[4]{-1}(A - iB) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right) (\cos(3(c + dx)) + i \sin(3(c + dx))) \right)}{a^3 d (-I + \tan(c + dx))^3}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^3),x]`

output `-1/24*(Sqrt[Cot[c + d*x]]*Sec[c + d*x]^3*(-3*(-1)^(1/4)*(A - I*B)*ArcTan[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)]) + 3*(-1)^(1/4)*(A - (6*I)*B)*ArcTanh[(-1)^(3/4)*Sqrt[Tan[c + d*x]]]*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)]) + Cos[c + d*x]*(3*(A + (2*I)*B) - 3*(A + (7*I)*B)*Cos[2*(c + d*x)] + ((-I)*A + 19*B)*Sin[2*(c + d*x)])*Sqrt[Tan[c + d*x]])*Sqrt[Tan[c + d*x]])/(a^3*d*(-I + Tan[c + d*x])^3)`

3.534.3 Rubi [A] (verified)Time = 1.33 (sec) , antiderivative size = 288, normalized size of antiderivative = 0.93, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.611$, Rules used = {3042, 4064, 3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{5/2}(a + ia \tan(c + dx))^3} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{A \cot(c + dx) + B}{\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} dx$$

3.534. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{B - A \tan\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{-\tan\left(c + dx + \frac{\pi}{2}\right)} \left(-a \tan\left(c + dx + \frac{\pi}{2}\right) + ia\right)^3} dx && \downarrow \text{3042} \\
& \frac{\int \frac{a(A-11iB) - 5a(iA-B) \cot(c+dx)}{2\sqrt{\cot(c+dx)}(\cot(c+dx)a+ia)^2} dx}{6a^2} + \frac{(-B + iA)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx) + ia)^3} && \downarrow \text{4079} \\
& \frac{\int \frac{a(A-11iB) - 5a(iA-B) \cot(c+dx)}{\sqrt{\cot(c+dx)}(\cot(c+dx)a+ia)^2} dx}{12a^2} + \frac{(-B + iA)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx) + ia)^3} && \downarrow \text{27} \\
& \frac{\int \frac{a(A-11iB) + 5a(iA-B) \tan\left(c+dx + \frac{\pi}{2}\right)}{\sqrt{-\tan\left(c+dx + \frac{\pi}{2}\right)} \left(ia - a \tan\left(c+dx + \frac{\pi}{2}\right)\right)^2} dx}{12a^2} + \frac{(-B + iA)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx) + ia)^3} && \downarrow \text{3042} \\
& \frac{\int -\frac{6\left((iA+6B)a^2 + (A+4iB) \cot(c+dx)a^2\right)}{\sqrt{\cot(c+dx)}(\cot(c+dx)a+ia)} dx}{12a^2} + \frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)^2} + \frac{(-B + iA)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx) + ia)^3} && \downarrow \text{4079} \\
& \frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)^2} - \frac{3 \int \frac{(iA+6B)a^2 + (A+4iB) \cot(c+dx)a^2}{\sqrt{\cot(c+dx)}(\cot(c+dx)a+ia)} dx}{12a^2} + \frac{(-B + iA)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx) + ia)^3} && \downarrow \text{27} \\
& \frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)^2} - \frac{3 \int \frac{a^2(iA+6B) - a^2(A+4iB) \tan\left(c+dx + \frac{\pi}{2}\right)}{\sqrt{-\tan\left(c+dx + \frac{\pi}{2}\right)} \left(ia - a \tan\left(c+dx + \frac{\pi}{2}\right)\right)} dx}{12a^2} + \frac{(-B + iA)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx) + ia)^3} && \downarrow \text{3042} \\
& \frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)^2} - \frac{3 \left(\frac{\int \frac{(2A-7iB)a^3 + 5B \cot(c+dx)a^3}{\sqrt{\cot(c+dx)}} dx}{2a^2} - \frac{5a^2 B \sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)} \right)}{12a^2} + \frac{(-B + iA)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx) + ia)^3} && \downarrow \text{4079} \\
& \frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)^2} - \frac{3 \left(\frac{\int \frac{(2A-7iB)a^3 + 5B \cot(c+dx)a^3}{\sqrt{\cot(c+dx)}} dx}{2a^2} - \frac{5a^2 B \sqrt{\cot(c+dx)}}{d(a \cot(c+dx) + ia)} \right)}{12a^2} + \frac{(-B + iA)\sqrt{\cot(c+dx)}}{6d(a \cot(c+dx) + ia)^3} && \downarrow \text{3042}
\end{aligned}$$

3.534. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\frac{\frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2} - \frac{3 \left(\frac{\int \frac{a^3(2A-7iB)-5a^3B \tan\left(c+dx+\frac{\pi}{2}\right) dx}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}}{2a^2} - \frac{5a^2B\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} \right)}{12a^2}}{12a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{6d(a\cot(c+dx)+ia)^3}$$

↓ 4017

$$\frac{\frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2} - \frac{3 \left(\frac{\int -\frac{a^3(2A-7iB+5B\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{a^2d} - \frac{5a^2B\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} \right)}{2a^2}}{12a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{6d(a\cot(c+dx)+ia)^3}$$

↓ 25

$$\frac{\frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2} - \frac{3 \left(\frac{\int \frac{a^3(2A-7iB+5B\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{a^2d} - \frac{5a^2B\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} \right)}{2a^2}}{12a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{6d(a\cot(c+dx)+ia)^3}$$

↓ 27

$$\frac{\frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2} - \frac{3 \left(-\frac{a \int \frac{2A-7iB+5B\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} - \frac{5a^2B\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} \right)}{2a^2}}{12a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{6d(a\cot(c+dx)+ia)^3}$$

↓ 1482

$$\frac{\frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2} - \frac{3 \left(-\frac{a \left(\frac{1}{2}(2A-(5+7i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(2A+(5-7i)B) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d} - \frac{5a^2B\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} \right)}{2a^2}}{12a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{6d(a\cot(c+dx)+ia)^3}$$

↓ 1476

$$\frac{\frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2} - \frac{3 \left(-\frac{a \left(\frac{1}{2}(2A-(5+7i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(2A+(5-7i)B) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{d}}{2a^2}}{12a^2} + \frac{(-B+iA)\sqrt{\cot(c+dx)}}{6d(a\cot(c+dx)+ia)^3}$$

3.534. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

↓ 1082

$$\frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2} - \frac{3 \left(a \left(\frac{1}{2}(2A-(5+7i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(2A+(5-7i)B) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \int \frac{1}{-\cot(c+dx)+1} d(1+\sqrt{2}\sqrt{\cot(c+dx)}) \right) \right)}{d} \right)}{2a^2}$$

$$\frac{(-B+iA)\sqrt{\cot(c+dx)}}{6d(a\cot(c+dx)+ia)^3}$$

12a²

↓ 217

$$\frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2} - \frac{3 \left(a \left(\frac{1}{2}(2A-(5+7i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(2A+(5-7i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{d} \right)}{2a^2}$$

$$\frac{(-B+iA)\sqrt{\cot(c+dx)}}{6d(a\cot(c+dx)+ia)^3}$$

12a²

↓ 1479

$$\frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2} - \frac{3 \left(a \left(\frac{1}{2}(2A-(5+7i)B) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) \right)}{d} \right)}{2a^2}$$

$$\frac{(-B+iA)\sqrt{\cot(c+dx)}}{6d(a\cot(c+dx)+ia)^3}$$

12a²

↓ 25

$$\frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2} - \frac{3 \left(a \left(\frac{1}{2}(2A-(5+7i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) \right) + \frac{1}{2}(2A+(5-7i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{d} \right)}{2a^2}$$

$$\frac{(-B+iA)\sqrt{\cot(c+dx)}}{6d(a\cot(c+dx)+ia)^3}$$

12a²

↓ 27

3.534. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\frac{\frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2} - \frac{3 \left(\frac{a \left(\frac{1}{2}(2A-(5+7i)B) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2}(2A-(5+7i)B) \left(\frac{\log}{d} \right) \right)}{12a^2}}{\frac{(-B+iA)\sqrt{\cot(c+dx)}}{6d(a\cot(c+dx)+ia)^3}}}{\frac{a(A+4iB)\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)^2} - \frac{3 \left(-\frac{5a^2B\sqrt{\cot(c+dx)}}{d(a\cot(c+dx)+ia)} - \frac{a \left(\frac{1}{2}(2A+(5-7i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(2A-(5+7i)B) \left(\frac{\log}{d} \right) \right)}{12a^2}}{\frac{(-B+iA)\sqrt{\cot(c+dx)}}{6d(a\cot(c+dx)+ia)^3}}}$$

↓ 1103

```
input Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^3),x]
```

```
output ((I*A - B)*Sqrt[Cot[c + d*x]]/(6*d*(I*a + a*Cot[c + d*x])^3) + ((a*(A + (4*I)*B)*Sqrt[Cot[c + d*x]]/(d*(I*a + a*Cot[c + d*x])^2) - (3*((-5*a^2*B*Sqrt[Cot[c + d*x]]/(d*(I*a + a*Cot[c + d*x])) - (a*((2*A + (5 - 7*I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]/Sqrt[2]))/2 + ((2*A - (5 + 7*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2))/d))/(2*a^2))/(12*a^2)
```

3.534.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

3.534. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

- rule 217 $\text{Int}[(a_+ + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{-1} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /;$ $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 1082 $\text{Int}[(a_+ + (b_+)(x_+) + (c_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{With}\{q = 1 - 4 \cdot \text{Simplify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /;$ $\text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4 \cdot a \cdot c]) /;$ $\text{FreeQ}\{a, b, c, x\}$
- rule 1103 $\text{Int}[(d_+ + (e_+)(x_+))/(a_+ + (b_+)(x_+) + (c_+)(x_+)^2), x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /;$ $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$
- rule 1476 $\text{Int}[(d_+ + (e_+)(x_+)^2)/(a_+ + (c_+)(x_+)^4), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c) \ \text{Int}[1/\text{Simp}[d/e + q \cdot x + x^2, x], x], x] + \text{Simp}[e/(2 \cdot c) \ \text{Int}[1/\text{Simp}[d/e - q \cdot x + x^2, x], x], x] /;$ $\text{FreeQ}\{a, c, d, e, x\} \ \&\& \ \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{PosQ}[d \cdot e]$
- rule 1479 $\text{Int}[(d_+ + (e_+)(x_+)^2)/(a_+ + (c_+)(x_+)^4), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[-2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c \cdot q) \ \text{Int}[(q - 2 \cdot x)/\text{Simp}[d/e + q \cdot x - x^2, x], x], x] + \text{Simp}[e/(2 \cdot c \cdot q) \ \text{Int}[(q + 2 \cdot x)/\text{Simp}[d/e - q \cdot x - x^2, x], x], x] /;$ $\text{FreeQ}\{a, c, d, e, x\} \ \&\& \ \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{NegQ}[d \cdot e]$
- rule 1482 $\text{Int}[(d_+ + (e_+)(x_+)^2)/(a_+ + (c_+)(x_+)^4), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[a \cdot c, 2]\}, \text{Simp}[(d \cdot q + a \cdot e)/(2 \cdot a \cdot c) \ \text{Int}[(q + c \cdot x^2)/(a + c \cdot x^4), x], x] + \text{Simp}[(d \cdot q - a \cdot e)/(2 \cdot a \cdot c) \ \text{Int}[(q - c \cdot x^2)/(a + c \cdot x^4), x], x] /;$ $\text{FreeQ}\{a, c, d, e, x\} \ \&\& \ \text{NeQ}[c \cdot d^2 + a \cdot e^2, 0] \ \&\& \ \text{NeQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{NegQ}[(-a) \cdot c]$
- rule 3042 $\text{Int}[u_+, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$ $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.534.4 Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.51

method	result
derivativedivides	$\frac{5B \cot(dx+c)^{\frac{5}{2}} + \left(\frac{38iB}{3} + \frac{2A}{3}\right) \cot(dx+c)^{\frac{3}{2}} + (2iA-9B)\sqrt{\cot(dx+c)}}{8(i+\cot(dx+c))^3} + \frac{(iA+6B) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{4\sqrt{2+4i\sqrt{2}}} + \frac{4\left(-\frac{iA}{16} - \frac{B}{16}\right) \arctan\left(\frac{2\sqrt{c}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}}$
default	$\frac{5B \cot(dx+c)^{\frac{5}{2}} + \left(\frac{38iB}{3} + \frac{2A}{3}\right) \cot(dx+c)^{\frac{3}{2}} + (2iA-9B)\sqrt{\cot(dx+c)}}{8(i+\cot(dx+c))^3} + \frac{(iA+6B) \arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{4\sqrt{2+4i\sqrt{2}}} + \frac{4\left(-\frac{iA}{16} - \frac{B}{16}\right) \arctan\left(\frac{2\sqrt{c}}{\sqrt{2-i\sqrt{2}}}\right)}{\sqrt{2-i\sqrt{2}}}$

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)`

output `1/a^3/d*(1/8*(5*B*cot(d*x+c)^(5/2)+(38/3*I*B+2/3*A)*cot(d*x+c)^(3/2)+(2*I*A-9*B)*cot(d*x+c)^(1/2))/(I+cot(d*x+c))^3+1/4*(I*A+6*B)/(2^(1/2)+I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))+4*(-1/16*I*A-1/16*B)/(2^(1/2)-I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))`

$$3.534. \int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$$

3.534.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 685 vs. $2(245) = 490$.

Time = 0.26 (sec) , antiderivative size = 685, normalized size of antiderivative = 2.21

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx =$$

$$\left(3a^3 d \sqrt{\frac{iA^2 + 2AB - iB^2}{a^6 d^2}} e^{(6i dx + 6i c)} \log \left(-\frac{2 \left((i a^3 d e^{(2i dx + 2i c)} - i a^3 d) \sqrt{\frac{i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} - 1}} \sqrt{\frac{iA^2 + 2AB - iB^2}{a^6 d^2}} + (A - iB) e^{(2i dx + 2i c)} \right)}{iA + B} \right) \right)$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^3,x, algorith
thm="fracas")
```

```
output -1/96*(3*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)
*log(-2*((I*a^3*d*e^(2*I*d*x + 2*I*c) - I*a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*
c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2))
+ (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*a^3*
d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(-2*((-I*
a^3*d*e^(2*I*d*x + 2*I*c) + I*a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(
2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^6*d^2)) + (A - I*B)
*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x - 2*I*c)/(I*A + B)) - 3*a^3*d*sqrt((-I*A
^2 - 12*A*B + 36*I*B^2)/(a^6*d^2))*e^(6*I*d*x + 6*I*c)*log(1/8*((a^3*d*e^(
2*I*d*x + 2*I*c) - a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2
*I*c) - 1))*sqrt((-I*A^2 - 12*A*B + 36*I*B^2)/(a^6*d^2)) + I*A + 6*B)*e^(-
2*I*d*x - 2*I*c)/(a^3*d)) + 3*a^3*d*sqrt((-I*A^2 - 12*A*B + 36*I*B^2)/(a^6
*d^2))*e^(6*I*d*x + 6*I*c)*log(-1/8*((a^3*d*e^(2*I*d*x + 2*I*c) - a^3*d)*s
qrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 -
12*A*B + 36*I*B^2)/(a^6*d^2)) - I*A - 6*B)*e^(-2*I*d*x - 2*I*c)/(a^3*d)) +
2*(2*(A + 10*I*B)*e^(6*I*d*x + 6*I*c) - (5*A + 26*I*B)*e^(4*I*d*x + 4*I*c
) + (4*A + 7*I*B)*e^(2*I*d*x + 2*I*c) - A - I*B)*sqrt((I*e^(2*I*d*x + 2*I*
c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*e^(-6*I*d*x - 6*I*c)/(a^3*d)
```

3.534.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(5/2)/(a+I*a*tan(d*x+c))**3,x)`

output `Timed out`

3.534.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.534.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^3 \cot(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)^3*cot(d*x + c)^(5/2)), x)`

3.534.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{5/2} (a + a \tan(c + dx) li)^3} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + a*tan(c + d*x)*li)^3),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + a*tan(c + d*x)*li)^3), x)`

3.535
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$$

3.535.1 Optimal result	5048
3.535.2 Mathematica [A] (verified)	5049
3.535.3 Rubi [A] (verified)	5049
3.535.4 Maple [A] (verified)	5057
3.535.5 Fricas [B] (verification not implemented)	5057
3.535.6 Sympy [F(-1)]	5058
3.535.7 Maxima [F(-2)]	5059
3.535.8 Giac [F]	5059
3.535.9 Mupad [F(-1)]	5059

3.535.1 Optimal result

Integrand size = 36, antiderivative size = 367

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx \\ &= \frac{\left(\frac{1}{16} + \frac{i}{16}\right) \left((1 + 6i)A - (29 + i)B\right) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}a^3d} \\ &+ \frac{\left((5 - 7i)A + (28 + 30i)B\right) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{16\sqrt{2}a^3d} \\ &+ \frac{5(A + 6iB)}{8a^3d\sqrt{\cot(c + dx)}} + \frac{iA - B}{6d\sqrt{\cot(c + dx)}(ia + a \cot(c + dx))^3} \\ &+ \frac{2A + 5iB}{12ad\sqrt{\cot(c + dx)}(ia + a \cot(c + dx))^2} - \frac{7(iA - 4B)}{24d\sqrt{\cot(c + dx)}(ia^3 + a^3 \cot(c + dx))} \\ &+ \frac{\left(\frac{1}{32} + \frac{i}{32}\right) \left((6 + i)A + (1 + 29i)B\right) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{\sqrt{2}a^3d} \\ &- \frac{\left(\frac{1}{32} + \frac{i}{32}\right) \left((6 + i)A + (1 + 29i)B\right) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{\sqrt{2}a^3d} \end{aligned}$$

3.535.
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$$

output $(-1/32-1/32*I)*((1+6*I)*A-(29+I)*B)*\arctan(-1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/a^3/d*2^{(1/2)}+1/32*((5-7*I)*A+(28+30*I)*B)*\arctan(1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/a^3/d*2^{(1/2)}+(1/64+1/64*I)*((6+I)*A+(1+29*I)*B)*\ln(1+\cot(d*x+c)-2^{(1/2)}*\cot(d*x+c)^{(1/2)})/a^3/d*2^{(1/2)}-(1/64+1/64*I)*((6+I)*A+(1+29*I)*B)*\ln(1+\cot(d*x+c)+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/a^3/d*2^{(1/2)}+5/8*(A+6*I*B)/a^3/d/\cot(d*x+c)^{(1/2)}+1/6*(I*A-B)/d/(I*a+a*\cot(d*x+c))^3/\cot(d*x+c)^{(1/2)}+1/12*(2*A+5*I*B)/a/d/(I*a+a*\cot(d*x+c))^2/\cot(d*x+c)^{(1/2)}-7/24*(I*A-4*B)/d/(I*a^3+a^3*\cot(d*x+c))/\cot(d*x+c)^{(1/2)}$

3.535.2 Mathematica [A] (verified)

Time = 7.09 (sec) , antiderivative size = 242, normalized size of antiderivative = 0.66

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx$$

$$= \frac{\sqrt{\cot(c + dx)} \sec^3(c + dx) \left(-12\sqrt[4]{-1}(iA + B) \arctan \left((-1)^{3/4} \sqrt{\tan(c + dx)} \right) (\cos(3(c + dx))) + i \sin(3(c + dx)) \right)}{\cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(7/2)*(a + I*a*Tan[c + d*x])^3),x]`

output $(\text{Sqrt}[\text{Cot}[c + d*x]]*\text{Sec}[c + d*x]^3*(-12*(-1)^{(1/4)}*(I*A + B)*\text{ArcTan}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]]*(\text{Cos}[3*(c + d*x)] + I*\text{Sin}[3*(c + d*x)]) + 12*(-1)^{(3/4)}*(6*A + (29*I)*B)*\text{ArcTanh}[(-1)^{(3/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]*(\text{Cos}[3*(c + d*x)] + I*\text{Sin}[3*(c + d*x)]) + (2*I)*((9*A + (33*I)*B)*\text{Cos}[c + d*x] + 21*(A + (7*I)*B)*\text{Cos}[3*(c + d*x)] + (2*I)*(19*A + (97*I)*B + (19*A + (145*I)*B)*\text{Cos}[2*(c + d*x)])*\text{Sin}[c + d*x])*\text{Sqrt}[\text{Tan}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]])/(96*a^3*d*(-I + \text{Tan}[c + d*x])^3)$

3.535.3 Rubi [A] (verified)

Time = 1.62 (sec) , antiderivative size = 346, normalized size of antiderivative = 0.94, number of steps used = 26, number of rules used = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.694$, Rules used = {3042, 4064, 3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4012, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.535. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\begin{aligned}
& \int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{7/2}(a + ia \tan(c + dx))^3} dx \\
& \quad \downarrow \text{4064} \\
& \int \frac{A \cot(c + dx) + B}{\cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + ia)^3} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{B - A \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}(-a \tan(c + dx + \frac{\pi}{2}) + ia)^3} dx \\
& \quad \downarrow \text{4079} \\
& \frac{\int -\frac{a(A+13iB)+7a(iA-B)\cot(c+dx)}{2\cot^{\frac{3}{2}}(c+dx)(\cot(c+dx)a+ia)^2} dx}{6a^2} + \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} \\
& \quad \downarrow \text{27} \\
& \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} - \frac{\int \frac{a(A+13iB)+7a(iA-B)\cot(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(\cot(c+dx)a+ia)^2} dx}{12a^2} \\
& \quad \downarrow \text{3042} \\
& \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} - \frac{\int \frac{a(A+13iB)-7a(iA-B)\tan(c+dx+\frac{\pi}{2})}{(-\tan(c+dx+\frac{\pi}{2}))^{3/2}(ia-a\tan(c+dx+\frac{\pi}{2}))^2} dx}{12a^2} \\
& \quad \downarrow \text{4079} \\
& \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} - \frac{\int -\frac{2(a^2(4iA-31B)-5a^2(2A+5iB)\cot(c+dx))}{\cot^{\frac{3}{2}}(c+dx)(\cot(c+dx)a+ia)} dx}{12a^2} - \frac{a(2A+5iB)}{d\sqrt{\cot(c+dx)}(a \cot(c+dx)+ia)^2} \\
& \quad \downarrow \text{27} \\
& \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} - \frac{\int \frac{a^2(4iA-31B)-5a^2(2A+5iB)\cot(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(\cot(c+dx)a+ia)} dx}{2a^2} - \frac{a(2A+5iB)}{d\sqrt{\cot(c+dx)}(a \cot(c+dx)+ia)^2} \\
& \quad \downarrow \\
& \frac{-B + iA}{12a^2}
\end{aligned}$$

3.535. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\begin{array}{c}
 \downarrow 3042 \\
 \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} - \\
 \frac{\int \frac{(4iA - 31B)a^2 + 5(2A + 5iB) \tan(c + dx + \frac{\pi}{2})a^2}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}(ia - a \tan(c + dx + \frac{\pi}{2}))} dx}{2a^2}}{2a^2} - \frac{a(2A + 5iB)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} \\
 \frac{12a^2}{\downarrow 4079} \\
 \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} - \\
 \frac{\int \frac{3(5(A + 6iB)a^3 + 7(iA - 4B) \cot(c + dx)a^3)}{\cot^{\frac{3}{2}}(c + dx)} dx}{2a^2}}{2a^2} - \frac{7a^2(-4B + iA)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{a(2A + 5iB)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} \\
 \frac{12a^2}{\downarrow 27} \\
 \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} - \\
 \frac{3 \int \frac{5(A + 6iB)a^3 + 7(iA - 4B) \cot(c + dx)a^3}{\cot^{\frac{3}{2}}(c + dx)} dx}{2a^2}}{2a^2} - \frac{7a^2(-4B + iA)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{a(2A + 5iB)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} \\
 \frac{12a^2}{\downarrow 3042} \\
 \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} - \\
 \frac{3 \int \frac{5a^3(A + 6iB) - 7a^3(iA - 4B) \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}} dx}{2a^2}}{2a^2} - \frac{7a^2(-4B + iA)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{a(2A + 5iB)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} \\
 \frac{12a^2}{\downarrow 4012} \\
 \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} - \\
 \frac{3 \left(\int \frac{7a^3(iA - 4B) - 5a^3(A + 6iB) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx + \frac{10a^3(A + 6iB)}{d\sqrt{\cot(c + dx)}} \right)}{2a^2}}{2a^2} - \frac{7a^2(-4B + iA)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{a(2A + 5iB)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} \\
 \frac{12a^2}{\downarrow 3042}
 \end{array}$$

3.535. $\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx$

$$\begin{aligned}
 & \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} \\
 & \frac{3 \left(\int \frac{7(iA - 4B)a^3 + 5(A + 6iB) \tan\left(c + dx + \frac{\pi}{2}\right)a^3}{\sqrt{-\tan\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{10a^3(A + 6iB)}{d\sqrt{\cot(c + dx)}} \right)}{2a^2} \\
 & - \frac{7a^2(-4B + iA)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{a(2A + 5iB)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} \\
 & \frac{12a^2}{2a^2} \\
 & \quad \downarrow \text{4017} \\
 & \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} \\
 & \frac{3 \left(\frac{2 \int -\frac{a^3(7(iA - 4B) - 5(A + 6iB) \cot(c + dx))}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{2a^2} + \frac{10a^3(A + 6iB)}{d\sqrt{\cot(c + dx)}} \right)}{2a^2} \\
 & - \frac{7a^2(-4B + iA)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{a(2A + 5iB)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} \\
 & \frac{12a^2}{2a^2} \\
 & \quad \downarrow \text{25} \\
 & \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} \\
 & \frac{3 \left(\frac{10a^3(A + 6iB)}{d\sqrt{\cot(c + dx)}} - \frac{2 \int \frac{a^3(7(iA - 4B) - 5(A + 6iB) \cot(c + dx))}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{2a^2} \right)}{2a^2} \\
 & - \frac{7a^2(-4B + iA)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{a(2A + 5iB)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} \\
 & \frac{12a^2}{2a^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} \\
 & \frac{3 \left(\frac{10a^3(A + 6iB)}{d\sqrt{\cot(c + dx)}} - \frac{2a^3 \int \frac{7(iA - 4B) - 5(A + 6iB) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{2a^2} \right)}{2a^2} \\
 & - \frac{7a^2(-4B + iA)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{a(2A + 5iB)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} \\
 & \frac{12a^2}{2a^2} \\
 & \quad \downarrow \text{1482} \\
 & \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} \\
 & \frac{3 \left(\frac{10a^3(A + 6iB)}{d\sqrt{\cot(c + dx)}} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((6 + i)A + (1 + 29i)B) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \left(\frac{1}{2} + \frac{i}{2} \right) ((1 + 6i)A - (29 + i)B) \int \frac{\cot(c + dx) + 1}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} \right)}{2a^2} \right)}{2a^2} \\
 & - \frac{7a^2(-4B + iA)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)} - \frac{a(2A + 5iB)}{d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^2} \\
 & \frac{12a^2}{2a^2} \\
 & \quad \downarrow \text{1476}
 \end{aligned}$$

3.535. $\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx$

$$\begin{array}{c}
 \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} \\
 \frac{\left(\frac{10a^3(A+6iB)}{d\sqrt{\cot(c+dx)}} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((6+i)A + (1+29i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \left(\frac{1}{2} + \frac{i}{2} \right) ((1+6i)A - (29+i)B) \left(\frac{1}{d} \int \frac{1}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{2a^2}}{2a^2} \\
 \hline
 \frac{1082}{\downarrow} \\
 \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} \\
 \frac{\left(\frac{10a^3(A+6iB)}{d\sqrt{\cot(c+dx)}} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((6+i)A + (1+29i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \left(\frac{1}{2} + \frac{i}{2} \right) ((1+6i)A - (29+i)B) \left(\frac{1}{d} \int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)}) - \frac{1}{\sqrt{2}} \int \frac{1}{\cot(c+dx)+1} d(1+\sqrt{2}\sqrt{\cot(c+dx)}) \right) \right)}{2a^2}}{2a^2} \\
 \hline
 \frac{217}{\downarrow} \\
 \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} \\
 \frac{\left(\frac{10a^3(A+6iB)}{d\sqrt{\cot(c+dx)}} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((6+i)A + (1+29i)B) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \left(\frac{1}{2} + \frac{i}{2} \right) ((1+6i)A - (29+i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{2a^2}}{2a^2} \\
 \hline
 \frac{1479}{\downarrow} \\
 \frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} \\
 \frac{\left(\frac{10a^3(A+6iB)}{d\sqrt{\cot(c+dx)}} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2} \right) ((6+i)A + (1+29i)B) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) \right)}{2a^2}}{2a^2} \right)}{2a^2} \\
 \hline
 \frac{25}{\downarrow}
 \end{array}$$

3.535. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

$$\frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2}\right) ((6+i)A + (1+29i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \left(\frac{1}{2} + \frac{i}{2}\right) \frac{10a^3(A+6iB)}{d\sqrt{\cot(c+dx)}} \right)}{2a^2}$$

27

$$\frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2}\right) ((6+i)A + (1+29i)B) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{d} \right) + \left(\frac{1}{2} + \frac{i}{2}\right) \frac{10a^3(A+6iB)}{d\sqrt{\cot(c+dx)}} \right)}{2a^2}$$

1103

$$\frac{-B + iA}{6d\sqrt{\cot(c + dx)}(a \cot(c + dx) + ia)^3} - \frac{2a^3 \left(\left(\frac{1}{2} + \frac{i}{2}\right) ((1+6i)A - (29+i)B) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \left(\frac{1}{2} + \frac{i}{2}\right) ((6+i)A + (1+29i)B) \left(\frac{\log(\cot(c+dx))}{d} \right) \right)}{2a^2}$$

```
input Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(7/2)*(a + I*a*Tan[c + d*x])^3),x]
```

```
output (I*A - B)/(6*d*Sqrt[Cot[c + d*x]]*(I*a + a*Cot[c + d*x])^3) - (-((a*(2*A + (5*I)*B))/(d*Sqrt[Cot[c + d*x]]*(I*a + a*Cot[c + d*x])^2)) - ((-7*a^2*(I*A - 4*B))/(d*Sqrt[Cot[c + d*x]]*(I*a + a*Cot[c + d*x]))) + (3*((10*a^3*(A + (6*I)*B))/(d*Sqrt[Cot[c + d*x]]) - (2*a^3*((1/2 + I/2)*((1 + 6*I)*A - (29 + I)*B)*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + (1/2 + I/2)*((6 + I)*A + (1 + 29*I)*B)*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/d)/(2*a^2))/(2*a^2))/(12*a^2)
```

3.535. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

3.535.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4017 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)])*(g_.)^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4079 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.535.4 Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.49

method	result
derivativedivides	$\frac{-i(5iA-14B)\cot(dx+c)^{\frac{5}{2}} + \left(\frac{38iA}{3} - \frac{98B}{3}\right)\cot(dx+c)^{\frac{3}{2}} + (-20iB-9A)\sqrt{\cot(dx+c)}}{8(i+\cot(dx+c))^3} + \frac{(29iB+6A)\arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{4\sqrt{2}+4i\sqrt{2}} + \frac{2iB}{\sqrt{\cot(dx+c)}}$
default	$\frac{-i(5iA-14B)\cot(dx+c)^{\frac{5}{2}} + \left(\frac{38iA}{3} - \frac{98B}{3}\right)\cot(dx+c)^{\frac{3}{2}} + (-20iB-9A)\sqrt{\cot(dx+c)}}{8(i+\cot(dx+c))^3} + \frac{(29iB+6A)\arctan\left(\frac{2\sqrt{\cot(dx+c)}}{\sqrt{2+i\sqrt{2}}}\right)}{4\sqrt{2}+4i\sqrt{2}} + \frac{2iB}{\sqrt{\cot(dx+c)}}$

```
input int((A+B*tan(d*x+c))/cot(d*x+c)^(7/2)/(a+I*a*tan(d*x+c))^3,x,method=_RETURNVERBOSE)
```

```
output 1/a^3/d*(1/8*(-I*(5*I*A-14*B)*cot(d*x+c)^(5/2)+(38/3*I*A-98/3*B)*cot(d*x+c)^(3/2)+(-9*A-20*I*B)*cot(d*x+c)^(1/2))/(I+cot(d*x+c))^3+1/4*(6*A+29*I*B)/(2^(1/2)+I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)+I*2^(1/2)))+2*I*B/cot(d*x+c)^(1/2)+4*(-1/16*A+1/16*I*B)/(2^(1/2)-I*2^(1/2))*arctan(2*cot(d*x+c)^(1/2)/(2^(1/2)-I*2^(1/2)))
```

3.535.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 783 vs. 2(274) = 548.

Time = 0.28 (sec) , antiderivative size = 783, normalized size of antiderivative = 2.13

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx$$

$$= \frac{3 \left(a^3 de^{(8i dx+8i c)} + a^3 de^{(6i dx+6i c)} \right) \sqrt{\frac{-iA^2-2AB+iB^2}{a^6d^2}} \log \left(-\frac{2 \left((a^3 de^{(2i dx+2i c)} - a^3 d) \sqrt{\frac{ie^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}-1}} \sqrt{\frac{-iA^2-2AB+iB^2}{a^6d^2}} \right)}{iA+B} \right)}{1}$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(7/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="fracas")
```


output

```

1/96*(3*(a^3*d*e^(8*I*d*x + 8*I*c) + a^3*d*e^(6*I*d*x + 6*I*c))*sqrt((-I*A
^2 - 2*A*B + I*B^2)/(a^6*d^2))*log(-2*((a^3*d*e^(2*I*d*x + 2*I*c) - a^3*d)
*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2
- 2*A*B + I*B^2)/(a^6*d^2)) + (A - I*B)*e^(2*I*d*x + 2*I*c))*e^(-2*I*d*x -
2*I*c)/(I*A + B) - 3*(a^3*d*e^(8*I*d*x + 8*I*c) + a^3*d*e^(6*I*d*x + 6*I
*c))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2))*log(2*((a^3*d*e^(2*I*d*x + 2
*I*c) - a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))
*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^6*d^2)) - (A - I*B)*e^(2*I*d*x + 2*I*c))
*e^(-2*I*d*x - 2*I*c)/(I*A + B) + 3*(a^3*d*e^(8*I*d*x + 8*I*c) + a^3*d*e^
(6*I*d*x + 6*I*c))*sqrt((36*I*A^2 - 348*A*B - 841*I*B^2)/(a^6*d^2))*log(1/
8*((a^3*d*e^(2*I*d*x + 2*I*c) - a^3*d)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e
^(2*I*d*x + 2*I*c) - 1))*sqrt((36*I*A^2 - 348*A*B - 841*I*B^2)/(a^6*d^2))
+ 6*A + 29*I*B)*e^(-2*I*d*x - 2*I*c)/(a^3*d) - 3*(a^3*d*e^(8*I*d*x + 8*I*
c) + a^3*d*e^(6*I*d*x + 6*I*c))*sqrt((36*I*A^2 - 348*A*B - 841*I*B^2)/(a^6
*d^2))*log(-1/8*((a^3*d*e^(2*I*d*x + 2*I*c) - a^3*d)*sqrt((I*e^(2*I*d*x +
2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((36*I*A^2 - 348*A*B - 841*I*B^
2)/(a^6*d^2)) - 6*A - 29*I*B)*e^(-2*I*d*x - 2*I*c)/(a^3*d) - 2*(2*(10*I*A
- 73*B)*e^(8*I*d*x + 8*I*c) + 3*(-2*I*A + 35*B)*e^(6*I*d*x + 6*I*c) - (19
*I*A - 49*B)*e^(4*I*d*x + 4*I*c) + 3*(2*I*A - 3*B)*e^(2*I*d*x + 2*I*c) - I
*A + B)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))/(a...

```

3.535.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(7/2)/(a+I*a*tan(d*x+c))**3,x)`

output `Timed out`

3.535.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(7/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expr: undefined: 0 to a negative exponent.`

3.535.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^3 \cot(dx + c)^{\frac{7}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(7/2)/(a+I*a*tan(d*x+c))^3,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)^3*cot(d*x + c)^(7/2)), x)`

3.535.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^3} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{7/2} (a + a \tan(c + dx) li)^3} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(7/2)*(a + a*tan(c + d*x)*li)^3),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(7/2)*(a + a*tan(c + d*x)*li)^3), x)`

3.535. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^3} dx$

3.536 $\int \cot^{\frac{7}{2}}(c+dx) \sqrt{a + ia \tan(c + dx)} (A+B \tan(c+dx)) dx$

3.536.1 Optimal result	5060
3.536.2 Mathematica [A] (verified)	5061
3.536.3 Rubi [A] (verified)	5061
3.536.4 Maple [B] (verified)	5065
3.536.5 Fricas [B] (verification not implemented)	5066
3.536.6 Sympy [F(-1)]	5066
3.536.7 Maxima [B] (verification not implemented)	5067
3.536.8 Giac [F]	5068
3.536.9 Mupad [F(-1)]	5068

3.536.1 Optimal result

Integrand size = 38, antiderivative size = 198

$$\int \cot^{\frac{7}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= - \frac{(1 + i)\sqrt{a}(A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{2(13A - 5iB) \sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)}}{15d}$$

$$- \frac{2(iA + 5B) \cot^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{15d} - \frac{2A \cot^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{5d}$$

```
output (-1-I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*a^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-2/15*(I*A+5*B)*cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(1/2)/d-2/5*A*cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(1/2)/d+2/15*(13*A-5*I*B)*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/d
```

3.536.2 Mathematica [A] (verified)

Time = 5.88 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.78

$$\int \cot^{\frac{7}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx =$$

$$\frac{i\left(\frac{15\sqrt{2}a(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{\sqrt{ia\tan(c+dx)}} + 2\cot^3(c+dx)\sqrt{a+ia\tan(c+dx)}(-3iA+(A-5iB)\tan(c+dx))\right)}{15d\sqrt{\cot(c+dx)}}$$

input `Integrate[Cot[c + d*x]^(7/2)*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `((-1/15*I)*((15*Sqrt[2]*a*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])]/Sqrt[a + I*a*Tan[c + d*x]]])/Sqrt[I*a*Tan[c + d*x]] + 2*Cot[c + d*x]^3*Sqrt[a + I*a*Tan[c + d*x]]*((-3*I)*A + (A - (5*I)*B)*Tan[c + d*x] + ((13*I)*A + 5*B)*Tan[c + d*x]^2)))/(d*Sqrt[Cot[c + d*x]])`

3.536.3 Rubi [A] (verified)Time = 1.29 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.10, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4729, 3042, 4081, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{7}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{7/2}\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\int \frac{\sqrt{ia\tan(c+dx)a+a(A+B\tan(c+dx))}}{\tan^{\frac{7}{2}}(c+dx)}dx$$

$$\downarrow \text{3042}$$

3.536. $\int \cot^{\frac{7}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$

$$\begin{aligned}
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\sqrt{i \tan(c+dx)a+a}(A+B \tan(c+dx))}{\tan(c+dx)^{7/2}} dx \\
& \quad \downarrow 4081 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA+5B)-4aA \tan(c+dx))}{2 \tan^{\frac{5}{2}}(c+dx)} dx}{5a} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right) \\
& \quad \downarrow 27 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA+5B)-4aA \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx}{5a} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right) \\
& \quad \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA+5B)-4aA \tan(c+dx))}{\tan(c+dx)^{5/2}} dx}{5a} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} \right) \\
& \quad \downarrow 4081 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int -\frac{\sqrt{i \tan(c+dx)a+a}((13A-5iB)a^2+2(iA+5B) \tan(c+dx)a^2)}{2 \tan^{\frac{3}{2}}(c+dx)} dx}{5a} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{5d} \right) \\
& \quad \downarrow 27 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((13A-5iB)a^2+2(iA+5B) \tan(c+dx)a^2)}{\tan^{\frac{3}{2}}(c+dx)} dx}{5a} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{5d} \right) \\
& \quad \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{\sqrt{i \tan(c+dx)a+a}((13A-5iB)a^2+2(iA+5B) \tan(c+dx)a^2)}{\tan(c+dx)^{3/2}} dx}{5a} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} - \frac{2A\sqrt{a+ia \tan(c+dx)}}{5d} \right)
\end{aligned}$$

3.536. $\int \cot^{\frac{7}{2}}(c+dx) \sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$

$$\begin{array}{c} \downarrow 4081 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int \frac{15a^3(iA+B)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx}{a} - \frac{2a^2(13A-5iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right) \\ \hline 5a \end{array}$$

$$\begin{array}{c} \downarrow 27 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(- \frac{15a^2(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{3a} - \frac{2a^2(13A-5iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right) \\ \hline 5a \end{array}$$

$$\begin{array}{c} \downarrow 3042 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(- \frac{15a^2(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{3a} - \frac{2a^2(13A-5iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right) \\ \hline 5a \end{array}$$

$$\begin{array}{c} \downarrow 4027 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(- \frac{30ia^4(B+iA) \int \frac{1}{i \tan(c+dx)a^2 - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} - \frac{2a^2(13A-5iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right) \\ \hline 5a \end{array}$$

$$\begin{array}{c} \downarrow 218 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(- \frac{(15-15i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^2(13A-5iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a(5B+iA)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right) \\ \hline 5a \end{array}$$

input `Int[Cot[c + d*x]^(7/2)*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

3.536. $\int \cot^{\frac{7}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx)) dx$

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A*Sqrt[a + I*a*Tan[c + d*x]])/(
5*d*Tan[c + d*x]^(5/2)) + ((-2*a*(I*A + 5*B)*Sqrt[a + I*a*Tan[c + d*x]])/(
3*d*Tan[c + d*x]^(3/2)) - (((15 - 15*I)*a^(5/2)*(I*A + B)*ArcTanh[((1 + I)
*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a^2*(13*A
- (5*I)*B)*Sqrt[a + I*a*Tan[c + d*x]]/(d*Sqrt[Tan[c + d*x]]))/(3*a))/(5*
a))
```

3.536.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4027 Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_)
+ (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_.)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.536.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 637 vs. 2(161) = 322.

Time = 0.68 (sec) , antiderivative size = 638, normalized size of antiderivative = 3.22

method	result
derivativedivides	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{7}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))}}{\left(15iB\sqrt{2} \ln\left(-\frac{-2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+ia-3a \tan(dx+c)}}{\tan(dx+c)+i}\right)\right)}$
default	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{7}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))}}{\left(15iB\sqrt{2} \ln\left(-\frac{-2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+ia-3a \tan(dx+c)}}{\tan(dx+c)+i}\right)\right)}$

```
input int(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

```
output -1/30/d*(1/tan(d*x+c))^(7/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)*(15*I*B
*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2
)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^4+15*I*A*2^(1/2)*ln(-(-
2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d
*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3-15*A*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(
1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c
)+I))*a*tan(d*x+c)^4-20*I*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(
1/2)*tan(d*x+c)^3+15*B*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*
(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3
-56*I*A*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+52
*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3+20*I*B*
(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-40*B*(-I*a)^(
1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+12*I*A*(-I*a)^(1/
2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-16*A*(-I*a)^(1/2)*tan(d*x+c)*(a*t
an(d*x+c)*(1+I*tan(d*x+c)))^(1/2))/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)/(
-tan(d*x+c)+I)/(-I*a)^(1/2)
```

$$3.536. \quad \int \cot^{\frac{7}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

3.536.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 481 vs. $2(151) = 302$.

Time = 0.26 (sec) , antiderivative size = 481, normalized size of antiderivative = 2.43

$$\int \cot^{\frac{7}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx =$$

$$15\sqrt{2}(de^{4i dx+4i c} - 2de^{2i dx+2i c} + d)\sqrt{-\frac{(-iA^2-2AB+iB^2)a}{d^2}}\log\left(-\frac{4\left((A-iB)ae^{i dx+i c}-(i de^{2i dx+2i c}-i d)\right)\sqrt{-\frac{(-iA^2-2AB+iB^2)a}{d^2}}}{\dots}\right)$$

input `integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-1/30*(15*sqrt(2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*log(-4*((A - I*B)*a*e^(I*d*x + I*c) - (I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-I*d*x - I*c)/(I*A + B)) - 15*sqrt(2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*log(-4*((A - I*B)*a*e^(I*d*x + I*c) - (-I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-I*d*x - I*c)/(I*A + B)) - 4*sqrt(2)*((17*A - 10*I*B)*e^(5*I*d*x + 5*I*c) - 10*(2*A - I*B)*e^(3*I*d*x + 3*I*c) + 15*A*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))/(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)`

3.536.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(7/2)*(a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.536. $\int \cot^{\frac{7}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$

3.536.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1409 vs. $2(151) = 302$.

Time = 0.74 (sec) , antiderivative size = 1409, normalized size of antiderivative = 7.12

$$\int \cot^{\frac{7}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")
```

```
output 1/30*(sqrt(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) +
1)*((30*((I + 1)*A - (I - 1)*B)*cos(3*d*x + 3*c) + (-(39*I + 39)*A + (25*I
- 25)*B)*cos(d*x + c) + 30*((I - 1)*A + (I + 1)*B)*sin(3*d*x + 3*c) + (-(
39*I - 39)*A - (25*I + 25)*B)*sin(d*x + c))*cos(3/2*arctan2(sin(2*d*x + 2*
c), cos(2*d*x + 2*c) - 1)) + (30*(-(I - 1)*A - (I + 1)*B)*cos(3*d*x + 3*c)
+ ((39*I - 39)*A + (25*I + 25)*B)*cos(d*x + c) + 30*((I + 1)*A - (I - 1)*
B)*sin(3*d*x + 3*c) + (-(39*I + 39)*A + (25*I - 25)*B)*sin(d*x + c))*sin(3
/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1))*sqrt(a) + 15*(2*((- (I
- 1)*A - (I + 1)*B)*cos(2*d*x + 2*c)^2 + (-(I - 1)*A - (I + 1)*B)*sin(2*d
*x + 2*c)^2 + 2*((I - 1)*A + (I + 1)*B)*cos(2*d*x + 2*c) - (I - 1)*A - (I
+ 1)*B)*arctan2(2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x +
2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1))
+ 2*sin(d*x + c), 2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x
+ 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1)
) + 2*cos(d*x + c)) + ((-(I + 1)*A + (I - 1)*B)*cos(2*d*x + 2*c)^2 + (-(I
+ 1)*A + (I - 1)*B)*sin(2*d*x + 2*c)^2 + 2*((I + 1)*A - (I - 1)*B)*cos(2*d
*x + 2*c) - (I + 1)*A + (I - 1)*B)*log(4*cos(d*x + c)^2 + 4*sin(d*x + c)^2
+ 4*sqrt(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) + 1
)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1))^2 + sin(1/2*ar
ctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1))^2) + 8*(cos(2*d*x + 2*c)...
```

3.536.8 Giac [F]

$$\int \cot^{\frac{7}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{ia \tan(dx + c) + a} \cot(dx + c)^{\frac{7}{2}} dx$$

input `integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)*cot(d*x + c)^(7/2), x)`

3.536.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{7/2} (A + B \tan(c + dx)) \sqrt{a + a \tan(c + dx)} \operatorname{li} dx$$

input `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2), x)`

output `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2), x)`

3.537 $\int \cot^{\frac{5}{2}}(c+dx) \sqrt{a + ia \tan(c + dx)} (A+B \tan(c+dx)) dx$

3.537.1 Optimal result	5069
3.537.2 Mathematica [A] (verified)	5069
3.537.3 Rubi [A] (verified)	5070
3.537.4 Maple [B] (verified)	5073
3.537.5 Fricas [B] (verification not implemented)	5074
3.537.6 Sympy [F(-1)]	5074
3.537.7 Maxima [B] (verification not implemented)	5075
3.537.8 Giac [F]	5076
3.537.9 Mupad [F(-1)]	5076

3.537.1 Optimal result

Integrand size = 38, antiderivative size = 155

$$\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{(1 + i)\sqrt{a}(iA + B)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d}$$

$$- \frac{2(iA + 3B)\sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)}}{3d} - \frac{2A \cot^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)}}{3d}$$

```
output (1+I)*(I*A+B)*arctanh(((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*a^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-2/3*A*cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(1/2)/d-2/3*(I*A+3*B)*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/d
```

3.537.2 Mathematica [A] (verified)

Time = 4.15 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.79

$$\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\sqrt{\cot(c + dx)} \left(3\sqrt{2}(iA + B)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{ia \tan(c + dx)} - 2(iA + 3B + A \cot(c + dx))\sqrt{a + ia \tan(c + dx)} \right)}{3d}$$

3.537. $\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

input `Integrate[Cot[c + d*x]^(5/2)*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*(3*Sqrt[2]*(I*A + B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]]*Sqrt[I*a*Tan[c + d*x]] - 2*(I*A + 3*B + A*Cot[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]]))/(3*d)`

3.537.3 Rubi [A] (verified)

Time = 0.93 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.05, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4729, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \cot(c + dx)^{5/2} \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4729} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\sqrt{i \tan(c + dx) a + a} (A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\sqrt{i \tan(c + dx) a + a} (A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx \\
 & \quad \downarrow \text{4081} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{2 \int \frac{\sqrt{i \tan(c + dx) a + a} (a(iA + 3B) - 2aA \tan(c + dx))}{2 \tan^{\frac{3}{2}}(c + dx)} dx}{3a} - \frac{2A \sqrt{a + ia \tan(c + dx)}}{3d \tan^{\frac{3}{2}}(c + dx)} \right) \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.537. $\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{\int\frac{\sqrt{i\tan(c+dx)a+a(a(iA+3B)-2aA\tan(c+dx))}dx}{\tan^{\frac{3}{2}}(c+dx)}-\frac{2A\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{\int\frac{\sqrt{i\tan(c+dx)a+a(a(iA+3B)-2aA\tan(c+dx))}dx}{\tan(c+dx)^{3/2}}-\frac{2A\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}\right)$$

↓ 4081

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2\int-\frac{3a^2(A-iB)\sqrt{i\tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}}dx-\frac{2a(3B+iA)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a}-\frac{2A\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{-3a(A-iB)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx-\frac{2a(3B+iA)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a}-\frac{2A\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{-3a(A-iB)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx-\frac{2a(3B+iA)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a}-\frac{2A\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}\right)$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{6ia^3(A-iB)\int\frac{1}{-\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a}-ia}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}-\frac{2a(3B+iA)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a}-\frac{2A\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}\right)$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(3-3i)a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}-\frac{2a(3B+iA)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-\frac{2A\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}\right)$$

3.537. $\int \cot^{\frac{5}{2}}(c+dx)\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$

input `Int[Cot[c + d*x]^(5/2)*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) + (((-3 + 3*I)*a^(3/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d - (2*a*(I*A + 3*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a)`

3.537.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.537.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 560 vs. 2(126) = 252.

Time = 0.56 (sec) , antiderivative size = 561, normalized size of antiderivative = 3.62

method	result
derivativedivides	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} \left(3iA\sqrt{2} \ln\left(-\frac{-2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+ia-3a \tan(dx+c)}}{\tan(dx+c)+i}\right)}\right)}{}$
default	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} \left(3iA\sqrt{2} \ln\left(-\frac{-2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+ia-3a \tan(dx+c)}}{\tan(dx+c)+i}\right)}\right)}{}$

```
input int(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

```
output -1/6/d*(1/tan(d*x+c))^(5/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)*(3*I*A*1
n(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*
tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^3-3*I*B*ln(-(-2*2^(1/2)*(-
-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan
(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^2+3*B*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2
)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I
)*a*tan(d*x+c)^3-12*B*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x
+c)))^(1/2)+3*A*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)
))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^2-8*A*(-
I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)-4*I*A*(-I*a)^(
1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2+12*I*B*tan(d*x+c)*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)+4*I*A*(a*tan(d*x+c)*(1+
I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2))/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)/(
-tan(d*x+c)+I)/(-I*a)^(1/2)
```

$$3.537. \quad \int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

3.537.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 430 vs. $2(118) = 236$.

Time = 0.25 (sec) , antiderivative size = 430, normalized size of antiderivative = 2.77

$$\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx =$$

$$3\sqrt{2} (de^{(2i dx + 2i c)} - d) \sqrt{-\frac{(i A^2 + 2 AB - i B^2)a}{d^2}} \log \left(-\frac{4 \left((A - i B) a e^{(i dx + i c)} + (de^{(2i dx + 2i c)} - d) \sqrt{-\frac{(i A^2 + 2 AB - i B^2)a}{d^2}} \sqrt{e^{(2i dx + 2i c)}} \right)}{i A + B} \right)$$

```
input integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output -1/6*(3*sqrt(2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*log(-4*((A - I*B)*a*e^(I*d*x + I*c) + (d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-I*d*x - I*c)/(I*A + B) - 3*sqrt(2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*log(-4*((A - I*B)*a*e^(I*d*x + I*c) - (d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-I*d*x - I*c)/(I*A + B) + 4*sqrt(2)*((2*I*A + 3*B)*e^(3*I*d*x + 3*I*c) - 3*B*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))/(d*e^(2*I*d*x + 2*I*c) - d)
```

3.537.6 SymPy [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

```
input integrate(cot(d*x+c)**(5/2)*(a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)
```

```
output Timed out
```

3.537. $\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.537.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1157 vs. $2(118) = 236$.

Time = 0.50 (sec) , antiderivative size = 1157, normalized size of antiderivative = 7.46

$$\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")
```

```
output 1/6*(2*sqrt(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) +
1)*((3*(-(I - 1)*A - (I + 1)*B)*cos(3*d*x + 3*c) + (-(I - 1)*A + (3*I + 3)
)*B)*cos(d*x + c) + 3*((I + 1)*A - (I - 1)*B)*sin(3*d*x + 3*c) + ((I + 1)*
A + (3*I - 3)*B)*sin(d*x + c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x
+ 2*c) - 1)) + (3*(-(I + 1)*A + (I - 1)*B)*cos(3*d*x + 3*c) + (-(I + 1)*A
- (3*I - 3)*B)*cos(d*x + c) + 3*(-(I - 1)*A - (I + 1)*B)*sin(3*d*x + 3*c)
+ (-(I - 1)*A + (3*I + 3)*B)*sin(d*x + c))*sin(3/2*arctan2(sin(2*d*x + 2*
c), cos(2*d*x + 2*c) - 1)))*sqrt(a) + 3*(2*((-(I + 1)*A + (I - 1)*B)*cos(2
*d*x + 2*c)^2 + (-(I + 1)*A + (I - 1)*B)*sin(2*d*x + 2*c)^2 + 2*((I + 1)*A
- (I - 1)*B)*cos(2*d*x + 2*c) - (I + 1)*A + (I - 1)*B)*arctan2(2*(cos(2*d
*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*a
rctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1)) + 2*sin(d*x + c), 2*(cos(2
*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2
*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1)) + 2*cos(d*x + c)) + (((I
- 1)*A + (I + 1)*B)*cos(2*d*x + 2*c)^2 + ((I - 1)*A + (I + 1)*B)*sin(2*d*
x + 2*c)^2 + 2*(-(I - 1)*A - (I + 1)*B)*cos(2*d*x + 2*c) + (I - 1)*A + (I
+ 1)*B)*log(4*cos(d*x + c)^2 + 4*sin(d*x + c)^2 + 4*sqrt(cos(2*d*x + 2*c)^
2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) + 1)*(cos(1/2*arctan2(sin(2*d*
x + 2*c), cos(2*d*x + 2*c) - 1))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos
(2*d*x + 2*c) - 1))^2) + 8*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2...
```

3.537.8 Giac [F]

$$\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{ia \tan(dx + c) + a} \cot(dx + c)^{\frac{5}{2}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)*cot(d*x + c)^(5/2), x)`

3.537.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{5/2} (A + B \tan(c + dx)) \sqrt{a + a \tan(c + dx)} \operatorname{li} dx$$

input `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2), x)`

output `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2), x)`

3.538 $\int \cot^{\frac{3}{2}}(c+dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.538.1 Optimal result	5077
3.538.2 Mathematica [A] (verified)	5077
3.538.3 Rubi [A] (verified)	5078
3.538.4 Maple [B] (verified)	5080
3.538.5 Fricas [B] (verification not implemented)	5081
3.538.6 Sympy [F(-1)]	5082
3.538.7 Maxima [B] (verification not implemented)	5082
3.538.8 Giac [F]	5083
3.538.9 Mupad [F(-1)]	5084

3.538.1 Optimal result

Integrand size = 38, antiderivative size = 110

$$\int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{(1 + i)\sqrt{a}(A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d} - \frac{2A\sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)}}{d}$$

```
output (1+I)*(A-I*B)*arctanh(((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*a^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-2*A*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/d
```

3.538.2 Mathematica [A] (verified)

Time = 3.04 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.93

$$\int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\sqrt{\cot(c + dx)} \left(\sqrt{2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{ia \tan(c + dx)} - 2A\sqrt{a + ia \tan(c + dx)} \right)}{d}$$

3.538. $\int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

input `Integrate[Cot[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*(Sqrt[2]*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]]*Sqrt[I*a*Tan[c + d*x]] - 2*A*Sqrt[a + I*a*Tan[c + d*x]]))/d`

3.538.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.01, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {3042, 4729, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \cot(c + dx)^{3/2} \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4729} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\sqrt{i \tan(c + dx) a + a} (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\sqrt{i \tan(c + dx) a + a} (A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx \\
 & \quad \downarrow \text{4081} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{2 \int \frac{a(iA+B)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx}{a} - \frac{2A\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} \right) \\
 & \quad \downarrow \text{27} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left((B + iA) \int \frac{\sqrt{i \tan(c + dx) a + a}}{\sqrt{\tan(c + dx)}} dx - \frac{2A\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} \right) \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.538. $\int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

$$\begin{aligned} & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left((B+iA)\int\frac{\sqrt{ia\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx-\frac{2A\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right) \\ & \qquad \qquad \qquad \downarrow 4027 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2ia^2(B+iA)\int\frac{1}{-\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a}-ia}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{ia\tan(c+dx)a+a}}}{d}-\frac{2A\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right) \\ & \qquad \qquad \qquad \downarrow 218 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(1-i)\sqrt{a}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}-\frac{2A\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right) \end{aligned}$$

input `Int[Cot[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((1 - I)*Sqrt[a]*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*A *Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]])`

3.538.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4027 Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2)), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]
```

3.538.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 441 vs. 2(91) = 182.

Time = 0.56 (sec) , antiderivative size = 442, normalized size of antiderivative = 4.02

method	result
derivativedivides	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{3}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} \left(iB \ln \left(-\frac{-2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+ia-3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \sqrt{2} a \right)$
default	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{3}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} \left(iB \ln \left(-\frac{-2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+ia-3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \sqrt{2} a \right)$

```
input int(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_RETURVERBOSE)
```

$$3.538. \quad \int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

output $\frac{1}{2}d*(1/\tan(dx+c))^{3/2}*\tan(dx+c)*(a*(1+I*\tan(dx+c)))^{1/2}*(I*B*2^{1/2}*\ln(-(-2*2^{1/2})*(-I*a)^{1/2}*(a*\tan(dx+c)*(1+I*\tan(dx+c)))^{1/2}+I*a-3*a*\tan(dx+c))/(\tan(dx+c)+I))*a*\tan(dx+c)^2+I*A*2^{1/2}*\ln(-(-2*2^{1/2})*(-I*a)^{1/2}*(a*\tan(dx+c)*(1+I*\tan(dx+c)))^{1/2}+I*a-3*a*\tan(dx+c))/(\tan(dx+c)+I))*a*\tan(dx+c)-A*\ln(-(-2*2^{1/2})*(-I*a)^{1/2}*(a*\tan(dx+c)*(1+I*\tan(dx+c)))^{1/2}+I*a-3*a*\tan(dx+c))/(\tan(dx+c)+I))*2^{1/2}*a*\tan(dx+c)^2+B*2^{1/2}*\ln(-(-2*2^{1/2})*(-I*a)^{1/2}*(a*\tan(dx+c)*(1+I*\tan(dx+c)))^{1/2}+I*a-3*a*\tan(dx+c))/(\tan(dx+c)+I))*a*\tan(dx+c)-4*I*A*(a*\tan(dx+c)*(1+I*\tan(dx+c)))^{1/2}*(-I*a)^{1/2}+4*A*(-I*a)^{1/2}*\tan(dx+c)*(a*\tan(dx+c)*(1+I*\tan(dx+c)))^{1/2})/(a*\tan(dx+c)*(1+I*\tan(dx+c)))^{1/2}/(-I*a)^{1/2}/(-\tan(dx+c)+I)$

3.538.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 370 vs. $2(85) = 170$.

Time = 0.25 (sec) , antiderivative size = 370, normalized size of antiderivative = 3.36

$$\int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx =$$

$$4\sqrt{2}A\sqrt{\frac{a}{e^{2i dx+2i c}+1}}\sqrt{\frac{ie^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}-1}}e^{(i dx+i c)} - \sqrt{2}d\sqrt{-\frac{(-iA^2-2AB+iB^2)a}{d^2}}\log\left(-\frac{4\left((A-iB)ae^{(i dx+i c)}-(ide^{2i c})\right)}{\dots}\right)$$

input `integrate(cot(dx+c)^(3/2)*(a+I*a*tan(dx+c))^(1/2)*(A+B*tan(dx+c)),x, algorithm="fricas")`

output $-1/2*(4*\sqrt{2}*A*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))*e^{(I*d*x + I*c)} - \sqrt{2}*d*\sqrt{(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*\log(-4*((A - I*B)*a*e^{(I*d*x + I*c)} - (I*d*e^{(2*I*d*x + 2*I*c)} - I*d)*\sqrt{(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)))})*e^{(-I*d*x - I*c)/(I*A + B)} + \sqrt{2}*d*\sqrt{(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*\log(-4*((A - I*B)*a*e^{(I*d*x + I*c)} - (-I*d*e^{(2*I*d*x + 2*I*c)} + I*d)*\sqrt{(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)})*\sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)))})*e^{(-I*d*x - I*c)/(I*A + B)))/d$

3.538. $\int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.538.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(3/2)*(a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.538.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 558 vs. $2(85) = 170$.

Time = 0.40 (sec) , antiderivative size = 558, normalized size of antiderivative = 5.07

$$\int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\left(2((i-1)A + (i+1)B) \arctan\left(2(\cos(2dx + 2c))^2 + \sin(2dx + 2c)^2 - 2\cos(2dx + 2c) + 1 \right)^{\frac{1}{4}} \sin\left(\frac{1}{2}\right) \right)}{1}$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output $\frac{1}{2} * ((2 * ((I - 1) * A + (I + 1) * B) * \arctan2(2 * (\cos(2 * d * x + 2 * c)^2 + \sin(2 * d * x + 2 * c)^2 - 2 * \cos(2 * d * x + 2 * c) + 1)^{(1/4)} * \sin(1/2 * \arctan2(\sin(2 * d * x + 2 * c), \cos(2 * d * x + 2 * c) - 1))) + 2 * \sin(d * x + c), 2 * (\cos(2 * d * x + 2 * c)^2 + \sin(2 * d * x + 2 * c)^2 - 2 * \cos(2 * d * x + 2 * c) + 1)^{(1/4)} * \cos(1/2 * \arctan2(\sin(2 * d * x + 2 * c), \cos(2 * d * x + 2 * c) - 1))) + 2 * \cos(d * x + c)) - ((- (I + 1) * A + (I - 1) * B) * \log(4 * \cos(d * x + c)^2 + 4 * \sin(d * x + c)^2 + 4 * \sqrt{\cos(2 * d * x + 2 * c)^2 + \sin(2 * d * x + 2 * c)^2 - 2 * \cos(2 * d * x + 2 * c) + 1}) * (\cos(1/2 * \arctan2(\sin(2 * d * x + 2 * c), \cos(2 * d * x + 2 * c) - 1)))^2 + \sin(1/2 * \arctan2(\sin(2 * d * x + 2 * c), \cos(2 * d * x + 2 * c) - 1)))^2) + 8 * (\cos(2 * d * x + 2 * c)^2 + \sin(2 * d * x + 2 * c)^2 - 2 * \cos(2 * d * x + 2 * c) + 1)^{(1/4)} * (\cos(d * x + c) * \cos(1/2 * \arctan2(\sin(2 * d * x + 2 * c), \cos(2 * d * x + 2 * c) - 1))) + \sin(d * x + c) * \sin(1/2 * \arctan2(\sin(2 * d * x + 2 * c), \cos(2 * d * x + 2 * c) - 1)))) * (\cos(2 * d * x + 2 * c)^2 + \sin(2 * d * x + 2 * c)^2 - 2 * \cos(2 * d * x + 2 * c) + 1)^{(1/4)} * \sqrt{a} + 4 * ((- (I + 1) * A * \cos(d * x + c) - (I - 1) * A * \sin(d * x + c)) * \cos(1/2 * \arctan2(\sin(2 * d * x + 2 * c), \cos(2 * d * x + 2 * c) - 1))) + ((I - 1) * A * \cos(d * x + c) - (I + 1) * A * \sin(d * x + c)) * \sin(1/2 * \arctan2(\sin(2 * d * x + 2 * c), \cos(2 * d * x + 2 * c) - 1))) * \sqrt{a}) / ((\cos(2 * d * x + 2 * c)^2 + \sin(2 * d * x + 2 * c)^2 - 2 * \cos(2 * d * x + 2 * c) + 1)^{(1/4)} * d)$

3.538.8 Giac [F]

$$\int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{ia \tan(dx + c) + a} \cot(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)*cot(d*x + c)^(3/2), x)`

3.538.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{3/2} (A + B \tan(c + dx)) \sqrt{a + a \tan(c + dx)} \operatorname{li} dx$$

input `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2), x)`

output `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2), x)`

3.539 $\int \sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.539.1 Optimal result	5085
3.539.2 Mathematica [A] (verified)	5086
3.539.3 Rubi [A] (verified)	5086
3.539.4 Maple [B] (verified)	5089
3.539.5 Fricas [B] (verification not implemented)	5090
3.539.6 Sympy [F]	5091
3.539.7 Maxima [F]	5092
3.539.8 Giac [F]	5092
3.539.9 Mupad [F(-1)]	5092

3.539.1 Optimal result

Integrand size = 38, antiderivative size = 152

$$\int \sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= -\frac{2(-1)^{3/4} \sqrt{a} B \arctan\left(\frac{(-1)^{3/4} \sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d} + \frac{(1 - i) \sqrt{a} (A - iB) \operatorname{arctanh}\left(\frac{(1+i) \sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d}$$

```
output -2*(-1)^(3/4)*B*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*a^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(1-I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*a^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d
```

3.539.2 Mathematica [A] (verified)

Time = 3.44 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.06

$$\int \sqrt{\cot(c+dx)}\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$= \frac{a\sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}\left(\frac{\sqrt{2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)\sqrt{\tan(c+dx)}}{\sqrt{ia\tan(c+dx)}} + \frac{2\sqrt[4]{-1}\operatorname{Arcsinh}\left(\sqrt[4]{-1}\sqrt{\tan(c+dx)}\right)}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}$$

input `Integrate[Sqrt[Cot[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(a*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((Sqrt[2]*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[Tan[c + d*x]])/Sqrt[I*a*Tan[c + d*x]] + (2*(-1)^(1/4)*B*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]])*Sqrt[1 + I*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]))/d`

3.539.3 Rubi [A] (verified)

Time = 0.81 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.88, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {3042, 4729, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\cot(c+dx)}\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\cot(c+dx)}\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\int \frac{\sqrt{ia\tan(c+dx)a+a}(A+B\tan(c+dx))}{\sqrt{\tan(c+dx)}}dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\int \frac{\sqrt{ia\tan(c+dx)a+a}(A+B\tan(c+dx))}{\sqrt{\tan(c+dx)}}dx$$

3.539. $\int \sqrt{\cot(c+dx)}\sqrt{a+ia\tan(c+dx)}(A+B\tan(c+dx))dx$

$$\begin{aligned} & \downarrow 4084 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left((A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + \frac{iB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{a} \right) \\ & \downarrow 3042 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left((A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + \frac{iB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{a} \right) \\ & \downarrow 4027 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{iB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{a} - \frac{2ia^2(A-iB) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} \right) \\ & \downarrow 218 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{iB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{a} + \frac{(1-i)\sqrt{a}(A-iB) \operatorname{arctanh} \left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}} \right)}{d} \right) \\ & \downarrow 4082 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{iaB \int \frac{1}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{d} + \frac{(1-i)\sqrt{a}(A-iB) \operatorname{arctanh} \left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}} \right)}{d} \right) \\ & \downarrow 65 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2iaB \int \frac{1}{1-\frac{ia \tan(c+dx)}{i \tan(c+dx)a+a}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} + \frac{(1-i)\sqrt{a}(A-iB) \operatorname{arctanh} \left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}} \right)}{d} \right) \\ & \downarrow 216 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(1-i)\sqrt{a}(A-iB) \operatorname{arctanh} \left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}} \right)}{d} - \frac{2(-1)^{3/4}\sqrt{a}B \operatorname{arctan} \left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}} \right)}{d} \right) \end{aligned}$$

input `Int[Sqrt[Cot[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `((-2*(-1)^(3/4)*Sqrt[a]*B*ArcTan[(-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]]]/Sqrt[a + I*a*Tan[c + d*x]])/d + ((1 - I)*Sqrt[a]*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d)*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]`

3.539.3.1 Defintions of rubi rules used

rule 65 `Int[1/(Sqrt[(b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4082 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

```
rule 4084 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x]
- Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[
e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.539.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 505 vs. 2(121) = 242.

Time = 0.56 (sec) , antiderivative size = 506, normalized size of antiderivative = 3.33

method	result
derivativedivides	$\frac{\sqrt{\frac{1}{\tan(dx+c)}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a \left(iA\sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+3a \tan(dx+c)-ia}}{\tan(dx+c)+i} \right) \sqrt{ia} t}{\dots}$
default	$\frac{\sqrt{\frac{1}{\tan(dx+c)}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a \left(iA\sqrt{2} \ln \left(\frac{2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+3a \tan(dx+c)-ia}}{\tan(dx+c)+i} \right) \sqrt{ia} t}{\dots}$

```
input int(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```


input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `1/2*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*log(-4*((A - I*B)*a*e^(I*d*x + I*c) + (d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-I*d*x - I*c)/(I*A + B) - 1/2*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*log(-4*((A - I*B)*a*e^(I*d*x + I*c) - (d*e^(2*I*d*x + 2*I*c) - d)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-I*d*x - I*c)/(I*A + B) - 1/4*sqrt(4*I*B^2*a/d^2)*log(-16*(3*B*a^2*e^(2*I*d*x + 2*I*c) - B*a^2 + sqrt(2)*(I*a*d*e^(3*I*d*x + 3*I*c) - I*a*d*e^(I*d*x + I*c))*sqrt(4*I*B^2*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-2*I*d*x - 2*I*c)/B + 1/4*sqrt(4*I*B^2*a/d^2)*log(-16*(3*B*a^2*e^(2*I*d*x + 2*I*c) - B*a^2 + sqrt(2)*(-I*a*d*e^(3*I*d*x + 3*I*c) + I*a*d*e^(I*d*x + I*c))*sqrt(4*I*B^2*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-2*I*d*x - 2*I*c)/B)`

3.539.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{\cot(c+dx)} \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx \\ &= \int \sqrt{ia(\tan(c+dx)-i)} (A+B \tan(c+dx)) \sqrt{\cot(c+dx)} dx \end{aligned}$$

input `integrate(cot(d*x+c)**(1/2)*(a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))*sqrt(cot(c + d*x)), x)`

3.539.7 Maxima [F]

$$\int \sqrt{\cot(c+dx)} \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$= \int (B \tan(dx+c) + A) \sqrt{ia \tan(dx+c) + a} \sqrt{\cot(dx+c)} dx$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)*sqrt(cot(d*x + c)), x)`

3.539.8 Giac [F]

$$\int \sqrt{\cot(c+dx)} \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$= \int (B \tan(dx+c) + A) \sqrt{ia \tan(dx+c) + a} \sqrt{\cot(dx+c)} dx$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)*sqrt(cot(d*x + c)), x)`

3.539.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cot(c+dx)} \sqrt{a+ia \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$= \int \sqrt{\cot(c+dx)} (A+B \tan(c+dx)) \sqrt{a+a \tan(c+dx)} li dx$$

input `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2),
x)`

output `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2),
x)`

$$3.540 \quad \int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.540.1 Optimal result	5094
3.540.2 Mathematica [A] (verified)	5095
3.540.3 Rubi [A] (verified)	5095
3.540.4 Maple [B] (verified)	5099
3.540.5 Fricas [B] (verification not implemented)	5100
3.540.6 Sympy [F]	5101
3.540.7 Maxima [F]	5102
3.540.8 Giac [F]	5102
3.540.9 Mupad [F(-1)]	5102

3.540.1 Optimal result

Integrand size = 38, antiderivative size = 192

$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

$$= -\frac{(-1)^{3/4} \sqrt{a}(2A-iB) \arctan\left(\frac{(-1)^{3/4} \sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{(1+i) \sqrt{a}(A-iB) \operatorname{arctanh}\left(\frac{(1+i) \sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}} + \frac{B \sqrt{a+ia \tan(c+dx)}}{d \sqrt{\cot(c+dx)}}$$

output
$$-(-1)^{(3/4)}*(2*A-I*B)*\arctan((-1)^{(3/4)}*a^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+I*a*\tan(d*x+c))^{(1/2)})*a^{(1/2)}*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/d-(1+I)*(A-I*B)*\operatorname{arctanh}((1+I)*a^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+I*a*\tan(d*x+c))^{(1/2)})*a^{(1/2)}*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/d+B*(a+I*a*\tan(d*x+c))^{(1/2)}/d/\cot(d*x+c)^{(1/2)}$$

3.540.2 Mathematica [A] (verified)

Time = 3.75 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.21

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(\frac{B \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}}{d} \right.$$

$$- \frac{i\sqrt{2}(\frac{1}{2}a^2(2A - iB) - \frac{1}{2}ia^2B) \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right) \sqrt{\tan(c + dx)}}{d\sqrt{ia \tan(c + dx)}} + \frac{\sqrt[4]{-1}a^2(2A - iB) \operatorname{arcsinh}\left(\sqrt[4]{-1}\sqrt{\tan(c + dx)}\right) \sqrt{1 + ia \tan(c + dx)}}{d\sqrt{a + ia \tan(c + dx)}} \Bigg) + \frac{\dots}{a}$$

```
input Integrate[(Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((B*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/d + (((-I)*Sqrt[2]*((a^2*(2*A - I*B))/2 - (I/2)*a^2*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]]*Sqrt[Tan[c + d*x]])/(d*Sqrt[I*a*Tan[c + d*x]]) + ((-1)^(1/4)*a^2*(2*A - I*B)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]*Sqrt[1 + I*Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]]))/a]
```

3.540.3 Rubi [A] (verified)

Time = 1.11 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.93, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.342$, Rules used = {3042, 4729, 3042, 4080, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

↓ 3042

3.540. $\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$

$$\begin{aligned}
& \int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\
& \quad \downarrow 4729 \\
& \sqrt{\tan(c + dx)}\sqrt{\cot(c + dx)} \int \sqrt{\tan(c + dx)}\sqrt{i \tan(c + dx)a + a}(A + B \tan(c + dx)) dx \\
& \quad \downarrow 3042 \\
& \sqrt{\tan(c + dx)}\sqrt{\cot(c + dx)} \int \sqrt{\tan(c + dx)}\sqrt{i \tan(c + dx)a + a}(A + B \tan(c + dx)) dx \\
& \quad \downarrow 4080 \\
& \sqrt{\tan(c + dx)}\sqrt{\cot(c + dx)} \left(\frac{\int -\frac{\sqrt{i \tan(c+dx)a+a}(aB-a(2A-iB) \tan(c+dx))}{2\sqrt{\tan(c+dx)}} dx}{a} + \frac{B\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} \right) \\
& \quad \downarrow 27 \\
& \sqrt{\tan(c + dx)}\sqrt{\cot(c + dx)} \left(\frac{B\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(aB-a(2A-iB) \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{2a} \right) \\
& \quad \downarrow 3042 \\
& \sqrt{\tan(c + dx)}\sqrt{\cot(c + dx)} \left(\frac{B\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(aB-a(2A-iB) \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{2a} \right) \\
& \quad \downarrow 4084 \\
& \sqrt{\tan(c + dx)}\sqrt{\cot(c + dx)} \left(\frac{B\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2a(B + iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - (B + 2iA)}{2a} \right) \\
& \quad \downarrow 3042 \\
& \sqrt{\tan(c + dx)}\sqrt{\cot(c + dx)} \left(\frac{B\sqrt{\tan(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2a(B + iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - (B + 2iA)}{2a} \right) \\
& \quad \downarrow 4027
\end{aligned}$$

3.540. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{4ia^3(B+iA) \int \frac{1}{\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{d} \right)$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{(2-2i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} \right)$$

↓ 4082

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{(2-2i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} \right)$$

↓ 65

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{(2-2i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d} - \frac{2\sqrt[4]{-1}a^{3/2}(B+2iA)\operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} \right)$$

input `Int[(Sqrt[a + I*a*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output $\text{Sqrt}[\text{Cot}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]*(-1/2*((2*(-1)^(1/4)*a^(3/2))*((2*I)*A + B)*\text{ArcTan}[((-1)^(3/4)*\text{Sqrt}[a]*\text{Sqrt}[\text{Tan}[c + d*x]]]/\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])]/d + ((2 - 2*I)*a^(3/2)*(I*A + B)*\text{ArcTanh}[((1 + I)*\text{Sqrt}[a]*\text{Sqrt}[\text{Tan}[c + d*x]]]/\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])]/d)/a + (B*\text{Sqrt}[\text{Tan}[c + d*x]]*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])/d$

3.540.3.1 Defintions of rubi rules used

rule 27 $\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \&\& \text{!MatchQ}[F_x, (b_)*(G_x_)] /; \text{FreeQ}[b, x]$

rule 65 $\text{Int}[1/(\text{Sqrt}[(b_)*(x_)]*\text{Sqrt}[(c_) + (d_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[2 \text{ Subst}[\text{Int}[1/(b - d*x^2), x], x, \text{Sqrt}[b*x]/\text{Sqrt}[c + d*x]], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{!GtQ}[c, 0]$

rule 216 $\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \&\& (\text{GtQ}[a, 0] \parallel \text{GtQ}[b, 0])$

rule 218 $\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4027 $\text{Int}[\text{Sqrt}[(a_) + (b_)*\text{tan}[(e_) + (f_)*(x_)]]/\text{Sqrt}[(c_) + (d_)*\text{tan}[(e_) + (f_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[-2*a*(b/f) \text{ Subst}[\text{Int}[1/(a*c - b*d - 2*a^2*x^2), x], x, \text{Sqrt}[c + d*\text{Tan}[e + f*x]]/\text{Sqrt}[a + b*\text{Tan}[e + f*x]]], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0]$

```
rule 4080 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[
1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Sim
p[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*T
an[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c
- a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]
```

```
rule 4082 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 + b^2, 0] && EqQ[A*b + a*B, 0]
```

```
rule 4084 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x]
- Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[
e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.540.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 315 vs. $2(155) = 310$.

Time = 0.68 (sec) , antiderivative size = 316, normalized size of antiderivative = 1.65

$$3.540. \quad \int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

method	result
derivativedivides	$-\frac{\sqrt{a(1+i \tan(dx+c))} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \left(B \sqrt{-ia} \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}} \right) \right)}{}$
default	$-\frac{\sqrt{a(1+i \tan(dx+c))} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \left(B \sqrt{-ia} \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}} \right) \right)}{}$

```
input int((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x,method=_R
ETURNVERBOSE)
```

```
output -1/2/d*(a*(1+I*tan(d*x+c)))^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(B
*(-I*a)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(
1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a-2*B*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/
2)*(I*a)^(1/2)*(-I*a)^(1/2)+2*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(
1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*(-I*a)^(1/2)-2^(1/2)*
ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a
*tan(d*x+c))/(tan(d*x+c)+I))*(I*a)^(1/2)*a)/(1/tan(d*x+c))^(1/2)/(1+I*tan(
d*x+c))/tan(d*x+c)/(I*a)^(1/2)/a/(-I*a)^(1/2)
```

3.540.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 793 vs. 2(148) = 296.

Time = 0.26 (sec) , antiderivative size = 793, normalized size of antiderivative = 4.13

$$\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx = \text{Too large to display}$$

```
input integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, al
gorithm="fricas")
```

output

```
-1/4*(2*sqrt(2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)
*a/d^2)*log(-4*((A - I*B)*a*e^(I*d*x + I*c) - (I*d*e^(2*I*d*x + 2*I*c) - I
*d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)
)*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-I*d*x -
I*c)/(I*A + B) - 2*sqrt(2)*(d*e^(2*I*d*x + 2*I*c) + d)*sqrt(-(-I*A^2 - 2
*A*B + I*B^2)*a/d^2)*log(-4*((A - I*B)*a*e^(I*d*x + I*c) - (-I*d*e^(2*I*d*
x + 2*I*c) + I*d)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x
+ 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)
))e^(-I*d*x - I*c)/(I*A + B) + 4*sqrt(2)*(I*B*e^(3*I*d*x + 3*I*c) - I*B*
e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*
I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)) - (d*e^(2*I*d*x + 2*I*c) + d)*sqrt((4
*I*A^2 + 4*A*B - I*B^2)*a/d^2)*log(-16*(3*(2*I*A + B)*a^2*e^(2*I*d*x + 2*I
*c) + (-2*I*A - B)*a^2 + 2*sqrt(2)*(a*d*e^(3*I*d*x + 3*I*c) - a*d*e^(I*d*x
+ I*c))*sqrt((4*I*A^2 + 4*A*B - I*B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c)
+ 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-2*
I*d*x - 2*I*c)/(2*I*A + B) + (d*e^(2*I*d*x + 2*I*c) + d)*sqrt((4*I*A^2 +
4*A*B - I*B^2)*a/d^2)*log(-16*(3*(2*I*A + B)*a^2*e^(2*I*d*x + 2*I*c) + (-2
*I*A - B)*a^2 - 2*sqrt(2)*(a*d*e^(3*I*d*x + 3*I*c) - a*d*e^(I*d*x + I*c))*
sqrt((4*I*A^2 + 4*A*B - I*B^2)*a/d^2)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sq
rt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-2*I*d*x ...
```

3.540.6 Sympy [F]

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{\sqrt{ia (\tan(c + dx) - i)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

input `integrate((a+I*a*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)**(1/2),x)`

output `Integral(sqrt(I*a*(tan(c + d*x) - I))*(A + B*tan(c + d*x))/sqrt(cot(c + d*x)), x)`

3.540.7 Maxima [F]

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)\sqrt{ia \tan(dx + c) + a}}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)/sqrt(cot(d*x + c)), x)`

3.540.8 Giac [F]

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)\sqrt{ia \tan(dx + c) + a}}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(I*a*tan(d*x + c) + a)/sqrt(cot(d*x + c)), x)`

3.540.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + ia \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(A + B \tan(c + dx)) \sqrt{a + a \tan(c + dx)} \operatorname{li}}{\sqrt{\cot(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2))/cot(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(1/2))/cot(c + d*x)^(1/2), x)`

3.540. $\int \frac{\sqrt{a+ia \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

3.541 $\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.541.1 Optimal result	5104
3.541.2 Mathematica [A] (verified)	5105
3.541.3 Rubi [A] (verified)	5105
3.541.4 Maple [B] (verified)	5110
3.541.5 Fricas [B] (verification not implemented)	5111
3.541.6 Sympy [F(-1)]	5112
3.541.7 Maxima [B] (verification not implemented)	5112
3.541.8 Giac [F]	5113
3.541.9 Mupad [F(-1)]	5114

3.541.1 Optimal result

Integrand size = 38, antiderivative size = 245

$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{(2 - 2i)a^{3/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d} + \frac{4a(67iA + 63B)\sqrt{\cot(c + dx)}\sqrt{a + ia \tan(c + dx)}}{105d} + \frac{4a(19A - 21iB)\cot^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{105d} - \frac{2a(8iA + 7B)\cot^{\frac{5}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{35d} - \frac{2aA\cot^{\frac{7}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{7d}$$

output

```
(2-2*I)*a^(3/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+4/105*a*(19*A-21*I*B)*cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(1/2)/d-2/35*a*(8*I*A+7*B)*cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(1/2)/d-2/7*a*A*cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(1/2)/d+4/105*a*(67*I*A+63*B)*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/d
```

3.541.2 Mathematica [A] (verified)

Time = 7.45 (sec) , antiderivative size = 359, normalized size of antiderivative = 1.47

$$\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{2 \cot^{\frac{5}{2}}(c+dx) \left(-15a^2A(i+\cot(c+dx))^2 \tan(c+dx) - 3(3iA+7B)(a+ia \tan(c+dx)) \right)}{\dots}$$

input `Integrate[Cot[c + d*x]^(9/2)*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(2*Cot[c + d*x]^(5/2)*(-15*a^2*A*(I + Cot[c + d*x])^2*Tan[c + d*x] - 3*((3*I)*A + 7*B)*(a + I*a*Tan[c + d*x])^2 + (29*A - (21*I)*B)*Tan[c + d*x]*(a + I*a*Tan[c + d*x])^2 - (105*a*(A - I*B)*Tan[c + d*x]^2*(-((-1)^(1/4))*a*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]]*Sqrt[Tan[c + d*x]]*(-I + Tan[c + d*x])) + Sqrt[a]*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[I*a*Tan[c + d*x]]*(-I + Tan[c + d*x]) + Sqrt[1 + I*Tan[c + d*x]]*(a*(-I + Tan[c + d*x]) + I*Sqrt[2]*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]*Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]))/Sqrt[1 + I*Tan[c + d*x]])/(105*d*Sqrt[a + I*a*Tan[c + d*x]])`

3.541.3 Rubi [A] (verified)Time = 1.59 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.10, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.447$, Rules used = {3042, 4729, 3042, 4076, 27, 3042, 4081, 25, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{9/2}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4729}$$

3.541. $\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a + a)^{3/2}(A + B \tan(c+dx))}{\tan^{9/2}(c+dx)} dx$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a + a)^{3/2}(A + B \tan(c+dx))}{\tan(c+dx)^{9/2}} dx$$

↓ 4076

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2}{7} \int \frac{\sqrt{i \tan(c+dx)a + a}(a(8iA + 7B) - a(6A - 7iB) \tan(c+dx))}{2 \tan^{7/2}(c+dx)} dx - \frac{2aA\sqrt{a}}{7d \tan^{7/2}(c+dx)} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \int \frac{\sqrt{i \tan(c+dx)a + a}(a(8iA + 7B) - a(6A - 7iB) \tan(c+dx))}{\tan^{7/2}(c+dx)} dx - \frac{2aA\sqrt{a}}{7d \tan^{7/2}(c+dx)} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \int \frac{\sqrt{i \tan(c+dx)a + a}(a(8iA + 7B) - a(6A - 7iB) \tan(c+dx))}{\tan(c+dx)^{7/2}} dx - \frac{2aA\sqrt{a}}{7d \tan^{7/2}(c+dx)} \right)$$

↓ 4081

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(\frac{2 \int -\frac{\sqrt{i \tan(c+dx)a + a}((19A - 21iB)a^2 + 2(8iA + 7B) \tan(c+dx)a^2)}{\tan^{5/2}(c+dx)} dx}{5a} - \frac{2a(7B + 8iA)\sqrt{a}}{5d \tan^{5/2}(c+dx)} \right) \right)$$

↓ 25

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(-\frac{2 \int \frac{\sqrt{i \tan(c+dx)a + a}((19A - 21iB)a^2 + 2(8iA + 7B) \tan(c+dx)a^2)}{\tan^{5/2}(c+dx)} dx}{5a} - \frac{2a(7B + 8iA)\sqrt{a}}{5d \tan^{5/2}(c+dx)} \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(-\frac{2 \int \frac{\sqrt{i \tan(c+dx)a + a}((19A - 21iB)a^2 + 2(8iA + 7B) \tan(c+dx)a^2)}{\tan(c+dx)^{5/2}} dx}{5a} - \frac{2a(7B + 8iA)\sqrt{a}}{5d \tan^{5/2}(c+dx)} \right) \right)$$

↓ 4081

3.541. $\int \cot^{9/2}(c+dx)(a + ia \tan(c+dx))^{3/2}(A + B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(\frac{2 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(67iA+63B)-2a^3(19A-21iB) \tan(c+dx))}{2 \tan^{\frac{3}{2}}(c+dx)} dx - \frac{2a^2(19A-21iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(\frac{2 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(67iA+63B)-2a^3(19A-21iB) \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx - \frac{2a^2(19A-21iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(\frac{2 \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(67iA+63B)-2a^3(19A-21iB) \tan(c+dx))}{\tan(c+dx)^{3/2}} dx - \frac{2a^2(19A-21iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} \right) \right)$$

↓ 4081

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(\frac{2 \left(\frac{\left(2 \int \frac{105a^4(A-iB)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx - \frac{2a^3(63B+67iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)}{3a} - \frac{2a^2(19A-21iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(\frac{2 \left(\frac{-105a^3(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(63B+67iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)}{3a} - \frac{2a^2(19A-21iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} \right)$$

3.541. $\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}(A+B \tan(c+dx)) dx$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(- \frac{2 \left(\frac{-105a^3(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(63B+67iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2a^2(19A-21iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} \right) \right)$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(- \frac{2 \left(\frac{210ia^5(A-iB) \int \frac{1}{-2 \tan(c+dx)a^2 - i \tan(c+dx)a+a} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} - \frac{2a^3(63B+67iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2a^2(19A-21iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} \right) \right)$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(- \frac{2 \left(\frac{-(105-105i)a^{7/2}(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^3(63B+67iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{2a^2(19A-21iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)} \right)}{5a} \right) \right)$$

input `Int[Cot[c + d*x]^(9/2)*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*a*A*Sqrt[a + I*a*Tan[c + d*x]])/(7*d*Tan[c + d*x]^(7/2)) + ((-2*a*((8*I)*A + 7*B)*Sqrt[a + I*a*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) - (2*((-2*a^2*(19*A - (21*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) + (((-105 + 105*I)*a^(7/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/d - (2*a^3*((67*I)*A + 63*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]])))/(3*a)))/(5*a))/7)`

3.541.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4076 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`
- rule 4081 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.541.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 803 vs. $2(200) = 400$.

Time = 0.58 (sec) , antiderivative size = 804, normalized size of antiderivative = 3.28

method	result
derivativedivides	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{9}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a \left(504B\sqrt{ia} \sqrt{-ia} \tan(dx+c)^3 \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} + 536iA\sqrt{a}\right)$
default	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{9}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a \left(504B\sqrt{ia} \sqrt{-ia} \tan(dx+c)^3 \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} + 536iA\sqrt{a}\right)$

```
input int(cot(d*x+c)^(9/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

```
output 1/210/d*(1/tan(d*x+c))^(9/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)*a*(504*
B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1
/2)+536*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3*(a*tan(d*x+c)*(1+I*tan(d
*x+c)))^(1/2)+105*I*(I*a)^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)
*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan
(d*x+c)^4+152*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*t
an(d*x+c)))^(1/2)-420*I*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*ta
n(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^4+4
20*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)
)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^4+210*I*ln(1/2*(2*I*a*ta
n(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2
))*(-I*a)^(1/2)*a*tan(d*x+c)^4-105*(I*a)^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2
)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I)
)*2^(1/2)*a*tan(d*x+c)^4-96*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan
(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+210*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+
c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(
d*x+c)^4-168*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*
tan(d*x+c)))^(1/2)-84*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*
(1+I*tan(d*x+c)))^(1/2)-60*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(
1/2)*(-I*a)^(1/2))/(I*a)^(1/2)/(-I*a)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+...
```

3.541.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 568 vs. 2(187) = 374.

Time = 0.26 (sec) , antiderivative size = 568, normalized size of antiderivative = 2.32

$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{105 \sqrt{2} \sqrt{-\frac{(iA^2 + 2AB - iB^2)a^3}{d^2}} (de^{(6i dx + 6i c)} - 3de^{(4i dx + 4i c)} + 3de^{(2i dx + 2i c)} - d) \log \left(\frac{4 \dots}{\dots} \right)}{\dots}$$

```
input integrate(cot(d*x+c)^(9/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, al
gorithm="fracas")
```

output `1/105*(105*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*log(4*((A - I*B)*a^2*e^(I*d*x + I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a)) - 105*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)*log(4*((A - I*B)*a^2*e^(I*d*x + I*c) - sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a)) - 2*sqrt(2)*((-211*I*A - 189*B)*a*e^(7*I*d*x + 7*I*c) + 7*(53*I*A + 57*B)*a*e^(5*I*d*x + 5*I*c) + 35*(-11*I*A - 9*B)*a*e^(3*I*d*x + 3*I*c) + 105*(I*A + B)*a*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))/(d*e^(6*I*d*x + 6*I*c) - 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) - d)`

3.541.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{\frac{3}{2}}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(9/2)*(a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.541.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 3787 vs. $2(187) = 374$.

Time = 3.82 (sec) , antiderivative size = 3787, normalized size of antiderivative = 15.46

$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{\frac{3}{2}}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(9/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

3.541. $\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{\frac{3}{2}}(A + B \tan(c + dx)) dx$

output

```
-1/420*(sqrt(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c)
+ 1)*(3*(280*(-(I - 1)*A - (I + 1)*B)*a*cos(7*d*x + 7*c) + 140*((I - 1)*A
+ (3*I + 3)*B)*a*cos(5*d*x + 5*c) + 7*(-(19*I - 19)*A - (29*I + 29)*B)*a*c
os(3*d*x + 3*c) + (-47*I - 47)*A + (63*I + 63)*B)*a*cos(d*x + c) + 280*((
I + 1)*A - (I - 1)*B)*a*sin(7*d*x + 7*c) + 140*(-(I + 1)*A + (3*I - 3)*B)*
a*sin(5*d*x + 5*c) + 7*((19*I + 19)*A - (29*I - 29)*B)*a*sin(3*d*x + 3*c)
+ ((47*I + 47)*A + (63*I - 63)*B)*a*sin(d*x + c))*cos(7/2*arctan2(sin(2*d*
x + 2*c), cos(2*d*x + 2*c) - 1)) + 4((((141*I - 141)*A + (119*I + 119)*B)
*a*cos(d*x + c) + (-141*I + 141)*A + (119*I - 119)*B)*a*sin(d*x + c))*cos
(2*d*x + 2*c)^2 + ((141*I - 141)*A + (119*I + 119)*B)*a*cos(d*x + c) + (((
141*I - 141)*A + (119*I + 119)*B)*a*cos(d*x + c) + (-141*I + 141)*A + (11
9*I - 119)*B)*a*sin(d*x + c))*sin(2*d*x + 2*c)^2 + (-141*I + 141)*A + (11
9*I - 119)*B)*a*sin(d*x + c) + 210*((-(I - 1)*A - (I + 1)*B)*a*cos(2*d*x +
2*c)^2 + -(I - 1)*A - (I + 1)*B)*a*sin(2*d*x + 2*c)^2 + 2*((I - 1)*A + (
I + 1)*B)*a*cos(2*d*x + 2*c) + -(I - 1)*A - (I + 1)*B)*a*cos(3*d*x + 3*c
) + 2*((-(141*I - 141)*A - (119*I + 119)*B)*a*cos(d*x + c) + ((141*I + 141
)*A - (119*I - 119)*B)*a*sin(d*x + c))*cos(2*d*x + 2*c) + 210*((I + 1)*A
- (I - 1)*B)*a*cos(2*d*x + 2*c)^2 + ((I + 1)*A - (I - 1)*B)*a*sin(2*d*x +
2*c)^2 + 2*(-(I + 1)*A + (I - 1)*B)*a*cos(2*d*x + 2*c) + ((I + 1)*A - (I -
1)*B)*a*sin(3*d*x + 3*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x...
```

3.541.8 Giac [F]

$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{\frac{3}{2}}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{\frac{3}{2}} \cot(dx + c)^{\frac{9}{2}} dx$$

input

```
integrate(cot(d*x+c)^(9/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, al
gorithm="giac")
```

output

```
integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^(
9/2), x)
```


3.541.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{9/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) i)^{3/2} dx$$

input `int(cot(c + d*x)^(9/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2), x)`

output `int(cot(c + d*x)^(9/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2), x)`

3.542 $\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.542.1 Optimal result	5115
3.542.2 Mathematica [A] (verified)	5116
3.542.3 Rubi [A] (verified)	5116
3.542.4 Maple [B] (verified)	5120
3.542.5 Fricas [B] (verification not implemented)	5121
3.542.6 Sympy [F(-1)]	5122
3.542.7 Maxima [B] (verification not implemented)	5122
3.542.8 Giac [F]	5123
3.542.9 Mupad [F(-1)]	5124

3.542.1 Optimal result

Integrand size = 38, antiderivative size = 201

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx =$$

$$\frac{(2 + 2i)a^{3/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{4a(9A - 10iB)\sqrt{\cot(c + dx)}\sqrt{a + ia \tan(c + dx)}}{15d}$$

$$- \frac{2a(6iA + 5B)\cot^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{15d}$$

$$- \frac{2aA\cot^{\frac{5}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{5d}$$

output

```
(-2-2*I)*a^(3/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-2/15*a*(6*I*A+5*B)*cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(1/2)/d-2/5*a*A*cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(1/2)/d+4/15*a*(9*A-10*I*B)*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/d
```

3.542.2 Mathematica [A] (verified)

Time = 6.64 (sec) , antiderivative size = 325, normalized size of antiderivative = 1.62

$$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A + B \tan(c+dx)) dx = \frac{2 \cot^{\frac{3}{2}}(c+dx) \left(-3a^2 A (i + \cot(c+dx))^2 \tan(c+dx) + a^2 (3iA + 5B) (-i + \tan(c+dx)) \right)}{\dots}$$

input `Integrate[Cot[c + d*x]^(7/2)*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(2*Cot[c + d*x]^(3/2)*(-3*a^2*A*(I + Cot[c + d*x])^2*Tan[c + d*x] + a^2*((3*I)*A + 5*B)*(-I + Tan[c + d*x])^2 + (15*a*(A - I*B)*Tan[c + d*x]*(Sqrt[a]*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*(1 + I*Tan[c + d*x])*Sqrt[I*a*Tan[c + d*x]] - (-1)^(3/4)*a*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]]*Sqrt[Tan[c + d*x]]*(-I + Tan[c + d*x]) + Sqrt[1 + I*Tan[c + d*x]]*(a + I*a*Tan[c + d*x] - Sqrt[2]*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]))/Sqrt[1 + I*Tan[c + d*x]])/(15*d*Sqrt[a + I*a*Tan[c + d*x]])`

3.542.3 Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.07, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4729, 3042, 4076, 27, 3042, 4081, 25, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A + B \tan(c+dx)) dx$$

↓ 3042

$$\int \cot(c+dx)^{7/2}(a+ia \tan(c+dx))^{3/2}(A + B \tan(c+dx)) dx$$

↓ 4729

3.542. $\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A + B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a + a)^{3/2}(A + B \tan(c+dx))}{\tan^{7/2}(c+dx)} dx$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a + a)^{3/2}(A + B \tan(c+dx))}{\tan(c+dx)^{7/2}} dx$$

↓ 4076

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2}{5} \int \frac{\sqrt{i \tan(c+dx)a + a}(a(6iA + 5B) - a(4A - 5iB) \tan(c+dx))}{2 \tan^{5/2}(c+dx)} dx - \frac{2aA\sqrt{a+i}}{5d \tan^{5/2}(c+dx)} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \int \frac{\sqrt{i \tan(c+dx)a + a}(a(6iA + 5B) - a(4A - 5iB) \tan(c+dx))}{\tan^{5/2}(c+dx)} dx - \frac{2aA\sqrt{a+i}}{5d \tan^{5/2}(c+dx)} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \int \frac{\sqrt{i \tan(c+dx)a + a}(a(6iA + 5B) - a(4A - 5iB) \tan(c+dx))}{\tan(c+dx)^{5/2}} dx - \frac{2aA\sqrt{a+i}}{5d \tan^{5/2}(c+dx)} \right)$$

↓ 4081

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \left(\frac{2 \int -\frac{\sqrt{i \tan(c+dx)a + a}((9A-10iB)a^2 + (6iA+5B) \tan(c+dx)a^2)}{\tan^{3/2}(c+dx)} dx}{3a} - \frac{2a(5B + 6iA)\sqrt{a+i}}{3d \tan^{3/2}(c+dx)} \right) \right)$$

↓ 25

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \left(-\frac{2 \int \frac{\sqrt{i \tan(c+dx)a + a}((9A-10iB)a^2 + (6iA+5B) \tan(c+dx)a^2)}{\tan^{3/2}(c+dx)} dx}{3a} - \frac{2a(5B + 6iA)\sqrt{a+i}}{3d \tan^{3/2}(c+dx)} \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \left(-\frac{2 \int \frac{\sqrt{i \tan(c+dx)a + a}((9A-10iB)a^2 + (6iA+5B) \tan(c+dx)a^2)}{\tan(c+dx)^{3/2}} dx}{3a} - \frac{2a(5B + 6iA)\sqrt{a+i}}{3d \tan^{3/2}(c+dx)} \right) \right)$$

↓ 4081

3.542. $\int \cot^{7/2}(c+dx)(a + ia \tan(c+dx))^{3/2}(A + B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \left(- \frac{2 \left(\frac{15a^3(iA+B)\sqrt{i\tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx - \frac{2a^2(9A-10iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)}{3a} - \frac{2a(5B+6iA)}{3d} \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \left(- \frac{2 \left(15a^2(B+iA) \int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(9A-10iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)}{3a} - \frac{2a(5B+6iA)}{3d} \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \left(- \frac{2 \left(15a^2(B+iA) \int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(9A-10iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)}{3a} - \frac{2a(5B+6iA)}{3d} \right) \right)$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \left(- \frac{2 \left(\frac{30ia^4(B+iA) \int \frac{1}{-\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}} - \frac{2a^2(9A-10iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)}{3a} - \frac{2a(5B+6iA)}{3d} \right) \right)$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \left(- \frac{2 \left(\frac{(15-15i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} - \frac{2a^2(9A-10iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)}{3a} - \frac{2a(5B+6iA)}{3d} \right) \right)$$

```
input Int[Cot[c + d*x]^(7/2)*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]
]
```

output $\text{Sqrt}[\text{Cot}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]*((-2*a*A*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])/(5*d*\text{Tan}[c + d*x]^{(5/2)}) + ((-2*a*((6*I)*A + 5*B)*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])/(3*d*\text{Tan}[c + d*x]^{(3/2)}) - (2*((15 - 15*I)*a^{(5/2)}*(I*A + B)*\text{ArcTanh}[\frac{((1 + I)*\text{Sqrt}[a]*\text{Sqrt}[\text{Tan}[c + d*x]])}{\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]]])/d - (2*a^2*(9*A - (10*I)*B)*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])/(d*\text{Sqrt}[\text{Tan}[c + d*x]])))/(3*a))/5$

3.542.3.1 Defintions of rubi rules used

rule 25 $\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$

rule 27 $\text{Int}[(a_*)(F_x), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_*)(G_x)] /; \text{FreeQ}[b, x]$

rule 218 $\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4027 $\text{Int}[\text{Sqrt}[(a_*) + (b_*)*\text{tan}[(e_*) + (f_*)(x_)]]/\text{Sqrt}[(c_*) + (d_*)*\text{tan}[(e_*) + (f_*)(x_)]], x_Symbol] \rightarrow \text{Simp}[-2*a*(b/f) \quad \text{Subst}[\text{Int}[1/(a*c - b*d - 2*a^2*x^2), x], x, \text{Sqrt}[c + d*\text{Tan}[e + f*x]]/\text{Sqrt}[a + b*\text{Tan}[e + f*x]]], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0]$

rule 4076 $\text{Int}[(a_*) + (b_*)*\text{tan}[(e_*) + (f_*)(x_)])^{(m_*)}*((A_*) + (B_*)*\text{tan}[(e_*) + (f_*)(x_)])^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(-a^2)*(B*c - A*d)*(a + b*\text{Tan}[e + f*x])^{(m-1)}*((c + d*\text{Tan}[e + f*x])^{(n+1)})/(d*f*(b*c + a*d)*(n+1)), x] - \text{Simp}[a/(d*(b*c + a*d)*(n+1)) \quad \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m-1)}*(c + d*\text{Tan}[e + f*x])^{(n+1)}*\text{Simp}[A*b*d*(m-n-2) - B*(b*c*(m-1) + a*d*(n+1)) + (a*A*d*(m+n) - B*(a*c*(m-1) + b*d*(n+1))]*\text{Tan}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{LtQ}[n, -1]$

```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.542.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 714 vs. 2(164) = 328.

Time = 0.53 (sec) , antiderivative size = 715, normalized size of antiderivative = 3.56

method	result
derivativedivides	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{7}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a \left(-72A\sqrt{ia} \sqrt{-ia} \tan(dx+c)^2 \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+60iA}\right)}{...}$
default	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{7}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a \left(-72A\sqrt{ia} \sqrt{-ia} \tan(dx+c)^2 \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+60iA}\right)}{...}$

```
input int(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

3.542. $\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

output

```
-1/30/d*(1/tan(d*x+c))^(7/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)*a*(-72*
A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1
/2)+60*I*A*(-I*a)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(
d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*tan(d*x+c)^3-15*I^2^(1/2)*(I*
a)^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2
)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3+80*I*B*(-I*a)^(1/2)*(
I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+60*B*(-I*a)^(
1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*
a)^(1/2)+a)/(I*a)^(1/2))*a*tan(d*x+c)^3+30*I*(-I*a)^(1/2)*ln(1/2*(2*I*a*ta
n(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2
))*a*tan(d*x+c)^3-15*2^(1/2)*(I*a)^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*t
an(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*ta
n(d*x+c)^3+24*I*A*(-I*a)^(1/2)*(I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*t
an(d*x+c)))^(1/2)-30*(-I*a)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)
*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*tan(d*x+c)^3+20*B*(
I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+1
2*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)*(-I*a)^(1/2)/(-I*a)
^(1/2)/(I*a)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
```

3.542.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 510 vs. $2(153) = 306$.

Time = 0.27 (sec) , antiderivative size = 510, normalized size of antiderivative = 2.54

$$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx =$$

$$15\sqrt{2}\sqrt{-\frac{(-iA^2-2AB+iB^2)a^3}{d^2}}(de^{(4i dx+4i c)} - 2de^{(2i dx+2i c)} + d) \log \left(\frac{4 \left((A-iB)a^2e^{(i dx+i c)} - \sqrt{-\frac{(-iA^2-2AB+iB^2)a^3}{d^2}} \right)}{\dots} \right)$$

input

```
integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, al
gorithm="fricas")
```

3.542. $\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

output

```
-1/15*(15*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(d*e^(4*I*d*x +
4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((A - I*B)*a^2*e^(I*d*x + I*c)
- sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(I*d*e^(2*I*d*x + 2*I*c) - I*d)
*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*
I*d*x + 2*I*c) - 1)))e^(-I*d*x - I*c)/((-I*A - B)*a)) - 15*sqrt(2)*sqrt(-
(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x
+ 2*I*c) + d)*log(4*((A - I*B)*a^2*e^(I*d*x + I*c) - sqrt(-(-I*A^2 - 2*A*B
+ I*B^2)*a^3/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(a/(e^(2*I*d*x + 2
*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e
^(-I*d*x - I*c)/((-I*A - B)*a)) - 2*sqrt(2)*((27*A - 25*I*B)*a*e^(5*I*d*x
+ 5*I*c) - 10*(3*A - 4*I*B)*a*e^(3*I*d*x + 3*I*c) + 15*(A - I*B)*a*e^(I*d*
x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) +
I)/(e^(2*I*d*x + 2*I*c) - 1)))/(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*x + 2
*I*c) + d)
```

3.542.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(7/2)*(a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.542.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1457 vs. $2(153) = 306$.

Time = 0.74 (sec) , antiderivative size = 1457, normalized size of antiderivative = 7.25

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

3.542. $\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

output

```

1/15*(2*sqrt(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c)
+ 1)*((15*((I + 1)*A - (I - 1)*B)*a*cos(3*d*x + 3*c) + (-16*I + 16)*A + (
15*I - 15)*B)*a*cos(d*x + c) + 15*((I - 1)*A + (I + 1)*B)*a*sin(3*d*x + 3*
c) + (-16*I - 16)*A - (15*I + 15)*B)*a*sin(d*x + c))*cos(3/2*arctan2(sin(
2*d*x + 2*c), cos(2*d*x + 2*c) - 1)) + (15*(-(I - 1)*A - (I + 1)*B)*a*cos(
3*d*x + 3*c) + ((16*I - 16)*A + (15*I + 15)*B)*a*cos(d*x + c) + 15*((I + 1
)*A - (I - 1)*B)*a*sin(3*d*x + 3*c) + (-16*I + 16)*A + (15*I - 15)*B)*a*s
in(d*x + c))*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1)))*sqr
t(a) + 15*(2*((-I - 1)*A - (I + 1)*B)*a*cos(2*d*x + 2*c)^2 + (-I - 1)*A
- (I + 1)*B)*a*sin(2*d*x + 2*c)^2 + 2*((I - 1)*A + (I + 1)*B)*a*cos(2*d*x
+ 2*c) + (-I - 1)*A - (I + 1)*B)*a*arctan2(2*(cos(2*d*x + 2*c)^2 + sin(2
*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*d*x +
2*c), cos(2*d*x + 2*c) - 1)) + 2*sin(d*x + c), 2*(cos(2*d*x + 2*c)^2 + sin
(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x
+ 2*c), cos(2*d*x + 2*c) - 1)) + 2*cos(d*x + c)) + ((-I + 1)*A + (I - 1)*
B)*a*cos(2*d*x + 2*c)^2 + (-I + 1)*A + (I - 1)*B)*a*sin(2*d*x + 2*c)^2 +
2*((I + 1)*A - (I - 1)*B)*a*cos(2*d*x + 2*c) + (-I + 1)*A + (I - 1)*B)*a
*log(4*cos(d*x + c)^2 + 4*sin(d*x + c)^2 + 4*sqrt(cos(2*d*x + 2*c)^2 + sin
(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) + 1)*(cos(1/2*arctan2(sin(2*d*x + 2*c
), cos(2*d*x + 2*c) - 1))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d..

```

3.542.8 Giac [F]

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{\frac{3}{2}}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{\frac{3}{2}} \cot(dx + c)^{\frac{7}{2}} dx$$

input

```

integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, al
gorithm="giac")

```

output

```

integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^(
7/2), x)

```

3.542.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \int \cot(c+dx)^{7/2}(A+B \tan(c+dx))(a+a \tan(c+dx) i)^{3/2} dx$$

input `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2), x)`

output `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2), x)`

3.543 $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.543.1 Optimal result	5125
3.543.2 Mathematica [A] (verified)	5125
3.543.3 Rubi [A] (verified)	5126
3.543.4 Maple [B] (verified)	5129
3.543.5 Fricas [B] (verification not implemented)	5130
3.543.6 Sympy [F(-1)]	5131
3.543.7 Maxima [B] (verification not implemented)	5131
3.543.8 Giac [F]	5132
3.543.9 Mupad [F(-1)]	5133

3.543.1 Optimal result

Integrand size = 38, antiderivative size = 157

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{(2 + 2i)a^{3/2}(iA + B)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d} - \frac{2a(4iA + 3B)\sqrt{\cot(c + dx)}\sqrt{a + ia \tan(c + dx)}}{3d} - \frac{2aA \cot^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{3d}$$

output $(2+2*I)*a^{(3/2)}*(I*A+B)*\operatorname{arctanh}((1+I)*a^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+I*a*\tan(d*x+c))^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/d-2/3*a*A*\cot(d*x+c)^{(3/2)}*(a+I*a*\tan(d*x+c))^{(1/2)}/d-2/3*a*(4*I*A+3*B)*\cot(d*x+c)^{(1/2)}*(a+I*a*\tan(d*x+c))^{(1/2)}/d$

3.543.2 Mathematica [A] (verified)

Time = 5.68 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.85

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{2a\sqrt{\cot(c + dx)}\left(aA \cot(c + dx)(-i + \tan(c + dx))^2 + \frac{3(A-iB)\left(-\sqrt[4]{-1}\operatorname{arcsinh}\left(\sqrt[4]{-1}\right)\right)}{3(A-iB)}\right)}{3d}$$

3.543. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

input `Integrate[Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(2*a*Sqrt[Cot[c + d*x]]*(a*A*Cot[c + d*x]*(-I + Tan[c + d*x])^2 + (3*(A - I*B)*(-(-1)^(1/4)*a*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]])*Sqrt[Tan[c + d*x]]*(-I + Tan[c + d*x])) + Sqrt[a]*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[I*a*Tan[c + d*x]]*(-I + Tan[c + d*x]) + Sqrt[1 + I*Tan[c + d*x]]*(a*(-I + Tan[c + d*x]) + I*Sqrt[2]*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]))/Sqrt[1 + I*Tan[c + d*x]])/(3*d*Sqrt[a + I*a*Tan[c + d*x]])`

3.543.3 Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4729, 3042, 4076, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \cot(c + dx)^{5/2}(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$$

$$\downarrow 4729$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{(i \tan(c + dx)a + a)^{3/2}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow 3042$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{(i \tan(c + dx)a + a)^{3/2}(A + B \tan(c + dx))}{\tan(c + dx)^{5/2}} dx$$

$$\downarrow 4076$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{2}{3} \int \frac{\sqrt{i \tan(c + dx)a + a}(a(4iA + 3B) - a(2A - 3iB) \tan(c + dx))}{2 \tan^{\frac{3}{2}}(c + dx)} dx - \frac{2aA\sqrt{a + ia \tan(c + dx)}}{3d \tan(c + dx)} \right)$$

$$\downarrow 27$$

3.543. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\int\frac{\sqrt{i\tan(c+dx)a+a}(a(4iA+3B)-a(2A-3iB)\tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)}dx-\frac{2aA\sqrt{a+}}{3d\tan}$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\int\frac{\sqrt{i\tan(c+dx)a+a}(a(4iA+3B)-a(2A-3iB)\tan(c+dx))}{\tan(c+dx)^{3/2}}dx-\frac{2aA\sqrt{a+}}{3d\tan}$$

↓ 4081

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{2\int-\frac{3a^2(A-iB)\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx}{a}-\frac{2a(3B+4iA)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right)-\frac{2aA}{3d\tan}$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(-6a(A-iB)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx-\frac{2a(3B+4iA)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(-6a(A-iB)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx-\frac{2a(3B+4iA)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right)\right)$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{12ia^3(A-iB)\int\frac{1}{-\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a}-ia}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{d}-\frac{2a(3B+4iA)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right)\right)$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(-\frac{(6-6i)a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}-\frac{2a(3B+4iA)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right)\right)$$

input `Int[Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*a*A*Sqrt[a + I*a*Tan[c + d*x]])
/(3*d*Tan[c + d*x]^(3/2)) + (((-6 + 6*I)*a^(3/2)*(A - I*B)*ArcTanh[((1 + I)
)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a*((4*I)
*A + 3*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/3)
```

3.543.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4027 Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_)
+ (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4076 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.543.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 625 vs. $2(128) = 256$.

Time = 0.53 (sec) , antiderivative size = 626, normalized size of antiderivative = 3.99

method	result
derivativedivides	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a \left(-12iB\sqrt{-ia} \ln\left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) \sqrt{ia+a}}{2\sqrt{ia}}\right)}{\right)}$
default	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a \left(-12iB\sqrt{-ia} \ln\left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) \sqrt{ia+a}}{2\sqrt{ia}}\right)}{\right)}$

```
input int(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

3.543. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$


```

output -1/6/d*(1/tan(d*x+c))^(5/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)*a*(-12*I
*B*(-I*a)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))
^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*tan(d*x+c)^2+3*I*(I*a)^(1/2)*2^(1/2)*
ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a
*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2+16*I*A*(I*a)^(1/2)*(-I*a)^(1/2
)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+12*A*(-I*a)^(1/2)*ln(1/
2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)
/(I*a)^(1/2))*a*tan(d*x+c)^2+6*I*(-I*a)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(
a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*tan(d*x
+c)^2-3*(I*a)^(1/2)*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+
I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2+12
*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/
2)+6*(-I*a)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)
)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*tan(d*x+c)^2+4*A*(a*tan(d*x+c)*(1+I
*tan(d*x+c)))^(1/2)*(I*a)^(1/2)*(-I*a)^(1/2))/(I*a)^(1/2)/(-I*a)^(1/2)/(a*
tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)

```

3.543.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 460 vs. $2(119) = 238$.

Time = 0.26 (sec) , antiderivative size = 460, normalized size of antiderivative = 2.93

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx =$$

$$3\sqrt{2}\sqrt{-\frac{(iA^2+2AB-iB^2)a^3}{d^2}}(de^{(2i dx+2i c)}-d) \log \left(\frac{4 \left((A-iB)a^2 e^{(i dx+i c)} + \sqrt{-\frac{(iA^2+2AB-iB^2)a^3}{d^2}} (de^{(2i dx+2i c)}-d) \sqrt{\frac{1}{e^{(2i dx+2i c)}}}}{(-iA-B)a} \right)}{(-iA-B)a} \right)$$

```

input integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, al
gorithm="fracas")

```

output
$$\begin{aligned} & -1/3*(3*\sqrt{2}*\sqrt{-(I*A^2 + 2*A*B - I*B^2)}*a^3/d^2)*(d*e^{(2*I*d*x + 2*I*c)} - d)*\log(4*((A - I*B)*a^2*e^{(I*d*x + I*c)} + \sqrt{-(I*A^2 + 2*A*B - I*B^2)}*a^3/d^2)*(d*e^{(2*I*d*x + 2*I*c)} - d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}) \\ & * \sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))} * e^{(-I*d*x - I*c)/((-I*A - B)*a)} - 3*\sqrt{2}*\sqrt{-(I*A^2 + 2*A*B - I*B^2)}*a^3/d^2*(d \\ & * e^{(2*I*d*x + 2*I*c)} - d)*\log(4*((A - I*B)*a^2*e^{(I*d*x + I*c)} - \sqrt{-(I*A^2 + 2*A*B - I*B^2)}*a^3/d^2)*(d*e^{(2*I*d*x + 2*I*c)} - d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}) \\ & * \sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))} * e^{(-I*d*x - I*c)/((-I*A - B)*a)} + 2*\sqrt{2}*((5*I*A + 3*B)*a*e^{(3*I*d*x + 3*I*c)} + 3*(-I*A - B)*a*e^{(I*d*x + I*c)})*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)} \\ & * \sqrt{((I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1))}/(d*e^{(2*I*d*x + 2*I*c)} - d) \end{aligned}$$

3.543.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(5/2)*(a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output Timed out

3.543.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1113 vs. $2(119) = 238$.

Time = 0.51 (sec) , antiderivative size = 1113, normalized size of antiderivative = 7.09

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

3.543. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

output

```

1/3*(2*sqrt(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) +
1)*((3*(-(I - 1)*A - (I + 1)*B)*a*cos(3*d*x + 3*c) + ((I - 1)*A + (3*I +
3)*B)*a*cos(d*x + c) + 3*((I + 1)*A - (I - 1)*B)*a*sin(3*d*x + 3*c) + (-(I
+ 1)*A + (3*I - 3)*B)*a*sin(d*x + c))*cos(3/2*arctan2(sin(2*d*x + 2*c), c
os(2*d*x + 2*c) - 1)) + (3*(-(I + 1)*A + (I - 1)*B)*a*cos(3*d*x + 3*c) + (
(I + 1)*A - (3*I - 3)*B)*a*cos(d*x + c) + 3*(-(I - 1)*A - (I + 1)*B)*a*sin
(3*d*x + 3*c) + ((I - 1)*A + (3*I + 3)*B)*a*sin(d*x + c))*sin(3/2*arctan2(
sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1)))*sqrt(a) + 3*(2*((-(I + 1)*A + (I
- 1)*B)*a*cos(2*d*x + 2*c)^2 + (-(I + 1)*A + (I - 1)*B)*a*sin(2*d*x + 2*c
)^2 + 2*((I + 1)*A - (I - 1)*B)*a*cos(2*d*x + 2*c) + (-(I + 1)*A + (I - 1)
*B)*a)*arctan2(2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x +
2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1)) +
2*sin(d*x + c), 2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x
+ 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1))
+ 2*cos(d*x + c)) + (((I - 1)*A + (I + 1)*B)*a*cos(2*d*x + 2*c)^2 + ((I -
1)*A + (I + 1)*B)*a*sin(2*d*x + 2*c)^2 + 2*(-(I - 1)*A - (I + 1)*B)*a*cos
(2*d*x + 2*c) + ((I - 1)*A + (I + 1)*B)*a)*log(4*cos(d*x + c)^2 + 4*sin(d*
x + c)^2 + 4*sqrt(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x +
2*c) + 1)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1))^2 + si
n(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1))^2) + 8*(cos(2*d*...

```

3.543.8 Giac [F]

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{\frac{3}{2}}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{\frac{3}{2}} \cot(dx + c)^{\frac{5}{2}} dx$$

input

```

integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, al
gorithm="giac")

```

output

```

integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^(
5/2), x)

```

3.543.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{5/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) i)^{3/2} dx$$

input `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2), x)`

output `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2), x)`

3.544 $\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{\frac{3}{2}}(A + B \tan(c + dx)) dx$

3.544.1 Optimal result	5134
3.544.2 Mathematica [A] (verified)	5135
3.544.3 Rubi [A] (verified)	5136
3.544.4 Maple [B] (verified)	5139
3.544.5 Fricas [B] (verification not implemented)	5140
3.544.6 Sympy [F(-1)]	5141
3.544.7 Maxima [F(-1)]	5142
3.544.8 Giac [F]	5142
3.544.9 Mupad [F(-1)]	5142

3.544.1 Optimal result

Integrand size = 38, antiderivative size = 186

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{\frac{3}{2}}(A + B \tan(c + dx)) dx = \frac{2\sqrt[4]{-1}a^{3/2}B \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d} + \frac{(2 + 2i)a^{3/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d} - \frac{2aA\sqrt{\cot(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d}$$

output

```
2*(-1)^(1/4)*a^(3/2)*B*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(2+2*I)*a^(3/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-2*a*A*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c)^(1/2))/d
```

3.544.2 Mathematica [A] (verified)

Time = 7.77 (sec) , antiderivative size = 357, normalized size of antiderivative = 1.92

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A + B \tan(c+dx)) dx = \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)} \left(-\frac{2A(a+ia \tan(c+dx))^{3/2}}{d\sqrt{\tan(c+dx)}} \right. \\ \left. + \frac{i(a^2A + \frac{1}{2}ia^2(3iA+B))}{2} \left(-\frac{2i\sqrt{2}a \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)\sqrt{ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{2ia^{3/2} \operatorname{arcsinh}\left(\frac{\sqrt{ia \tan(c+dx)}}{\sqrt{a}}\right)\sqrt{1+i \tan(c+dx)}\sqrt{ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} \right) \right) + \frac{\dots}{a}$$

input `Integrate[Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]), x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A*(a + I*a*Tan[c + d*x])^(3/2))/(d*Sqrt[Tan[c + d*x]]) + (2*((-I)*(a^2*A + (I/2)*a^2*((3*I)*A + B))*((-2*I)*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) + ((2*I)*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])))/a + (I*a^2*A*(-((-1)^(3/4)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/Sqrt[1 + I*Tan[c + d*x]]) + Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]))/d)/a`

$$3.544. \quad \int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A + B \tan(c+dx)) dx$$

3.544.3 Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.90, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.342$, Rules used = {3042, 4729, 3042, 4076, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{3/2}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^{3/2}(A+B \tan(c+dx))}{\tan(c+dx)^{3/2}} dx$$

$$\downarrow \text{4076}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(2 \int \frac{\sqrt{i \tan(c+dx)a+a}(a(2iA+B)+iaB \tan(c+dx))}{2\sqrt{\tan(c+dx)}} dx - \frac{2aA\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)$$

$$\downarrow \text{27}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{\sqrt{i \tan(c+dx)a+a}(a(2iA+B)+iaB \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx - \frac{2aA\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{\sqrt{i \tan(c+dx)a+a}(a(2iA+B)+iaB \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx - \frac{2aA\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)$$

$$\downarrow \text{4084}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(2a(B+ia) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - B \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a}}{\sqrt{\tan(c+dx)}} dx \right)$$

3.544. $\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(2a(B+iA)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx-B\int\frac{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}\right)$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{4ia^3(B+iA)\int\frac{1}{-\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a}-ia}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{d}-B\int\frac{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}\right)$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-B\int\frac{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx+\frac{(2-2i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}\right)$$

↓ 4082

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{a^2B\int\frac{1}{\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}}d\tan(c+dx)}{d}+\frac{(2-2i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}\right)$$

↓ 65

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2a^2B\int\frac{1}{1-\frac{ia\tan(c+dx)}{i\tan(c+dx)a+a}}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{d}+\frac{(2-2i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}\right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(2-2i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}+\frac{2\sqrt{-1}a^{3/2}B\operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}\right)$$

input `Int[Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`


```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((2*(-1)^(1/4)*a^(3/2)*B*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d + ((2 - 2*I)*a^(3/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a*A*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]])
```

3.544.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 65 Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]
```

```
rule 216 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4027 Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

```
rule 4076 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

```
rule 4082 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 + b^2, 0] && EqQ[A*b + a*B, 0]
```

```
rule 4084 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x]
- Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[
e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.544.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 528 vs. $2(150) = 300$.

Time = 0.55 (sec) , antiderivative size = 529, normalized size of antiderivative = 2.84

$$3.544. \quad \int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$$

method	result
derivativedivides	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{3}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a \left(4iA \ln\left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}}\right) \sqrt{-ia} a \tan\right)$
default	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{3}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a \left(4iA \ln\left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}}\right) \sqrt{-ia} a \tan\right)$

```
input int(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

```
output 1/2/d*(1/tan(d*x+c))^(3/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)*a*(4*I*A*
ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/
2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)-I*ln(-(-2*2^(1/2)*(-I*a)^(1/2
)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I)
)*2^(1/2)*(I*a)^(1/2)*a*tan(d*x+c)+2*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d
*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*t
an(d*x+c)+2*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(
1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)-ln(-(-2*2^(1/2)
*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(t
an(d*x+c)+I))*2^(1/2)*(I*a)^(1/2)*a*tan(d*x+c)-4*A*(a*tan(d*x+c)*(1+I*tan(
d*x+c)))^(1/2)*(I*a)^(1/2))*(-I*a)^(1/2)-2*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*ta
n(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*
a*tan(d*x+c))/(-I*a)^(1/2)/(I*a)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/
2)
```

3.544.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 674 vs. 2(142) = 284.

Time = 0.26 (sec) , antiderivative size = 674, normalized size of antiderivative = 3.62

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}(A+B \tan(c+dx)) dx =$$

$$8\sqrt{2}Aa\sqrt{\frac{a}{e^{(2i dx+2i c)}+1}}\sqrt{\frac{ie^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}-1}}e^{(i dx+i c)} - 4\sqrt{2}\sqrt{\frac{-(-iA^2-2AB+iB^2)a^3}{d^2}}d \log\left(\frac{4\left((A-iB)a^2e^{(i dx+i c)}-\sqrt{-\frac{(-iA^2-2AB+iB^2)a^3}{d^2}}\right)}{\dots}\right)$$

3.544. $\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{\frac{3}{2}}(A+B \tan(c+dx)) dx$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `-1/4*(8*sqrt(2)*A*a*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*e^(I*d*x + I*c) - 4*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*d*log(4*((A - I*B)*a^2*e^(I*d*x + I*c) - sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a) + 4*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*d*log(4*((A - I*B)*a^2*e^(I*d*x + I*c) - sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a) + sqrt(-4*I*B^2*a^3/d^2)*d*log(-16*(3*B*a^2*e^(2*I*d*x + 2*I*c) - B*a^2 + sqrt(2)*sqrt(-4*I*B^2*a^3/d^2)*(d*e^(3*I*d*x + 3*I*c) - d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/B - sqrt(-4*I*B^2*a^3/d^2)*d*log(-16*(3*B*a^2*e^(2*I*d*x + 2*I*c) - B*a^2 - sqrt(2)*sqrt(-4*I*B^2*a^3/d^2)*(d*e^(3*I*d*x + 3*I*c) - d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/B)/d`

3.544.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{\frac{3}{2}}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(3/2)*(a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.544.7 Maxima [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `Timed out`

3.544.8 Giac [F]

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{\frac{3}{2}} \cot(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^(3/2), x)`

3.544.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{3/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) 1i)^{3/2} dx$$

input `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2), x)`

output `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2), x)`

3.544. $\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.545 $\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.545.1 Optimal result	5143
3.545.2 Mathematica [A] (verified)	5144
3.545.3 Rubi [A] (verified)	5144
3.545.4 Maple [B] (verified)	5148
3.545.5 Fricas [B] (verification not implemented)	5149
3.545.6 Sympy [F]	5150
3.545.7 Maxima [F]	5151
3.545.8 Giac [F]	5151
3.545.9 Mupad [F(-1)]	5151

3.545.1 Optimal result

Integrand size = 38, antiderivative size = 196

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx =$$

$$\frac{(-1)^{3/4}a^{3/2}(2iA + 3B) \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{(2 - 2i)a^{3/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{iaB\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\cot(c + dx)}}$$

```
output -(-1)^(3/4)*a^(3/2)*(2*I*A+3*B)*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)
/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(2-2*I)*a^(
3/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/
2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+I*a*B*(a+I*a*tan(d*x+c))^(1/2)/d/c
ot(d*x+c)^(1/2)
```

3.545.2 Mathematica [A] (verified)

Time = 8.71 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.43

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{a\sqrt{\cot(c+dx)}\left(\sqrt[4]{-1}a \operatorname{Barcsinh}\left(\sqrt[4]{-1}\sqrt{\tan(c+dx)}\right)\right)(1+i \tan(c+dx))\sqrt{\tan(c+dx)}}{\dots}$$

input `Integrate[Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]), x]`

output `(a*Sqrt[Cot[c + d*x]]*((-1)^(1/4)*a*B*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]]*(1 + I*Tan[c + d*x])*Sqrt[Tan[c + d*x]] - 2*Sqrt[a]*(A - I*B)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[I*a*Tan[c + d*x]]*(-I + Tan[c + d*x]) - Sqrt[1 + I*Tan[c + d*x]]*(a*B*Tan[c + d*x]*(-I + Tan[c + d*x]) + 2*Sqrt[2]*(I*A + B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]))/(d*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])`

3.545.3 Rubi [A] (verified)

Time = 1.12 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.93, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.342$, Rules used = {3042, 4729, 3042, 4077, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx \\ & \quad \downarrow \text{4729} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^{3/2}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.545. $\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a + a)^{3/2}(A + B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

↓ 4077

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{\sqrt{i \tan(c+dx)a + a}(a(2A - iB) + a(2iA + 3B) \tan(c+dx))}{2\sqrt{\tan(c+dx)}} dx + \frac{iaB\sqrt{\tan(c+dx)}}{\sqrt{\tan(c+dx)}} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \int \frac{\sqrt{i \tan(c+dx)a + a}(a(2A - iB) + a(2iA + 3B) \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx + \frac{iaB\sqrt{\tan(c+dx)}}{\sqrt{\tan(c+dx)}} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \int \frac{\sqrt{i \tan(c+dx)a + a}(a(2A - iB) + a(2iA + 3B) \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx + \frac{iaB\sqrt{\tan(c+dx)}}{\sqrt{\tan(c+dx)}} \right)$$

↓ 4084

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \left(4a(A - iB) \int \frac{\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx - (2A - 3iB) \int \frac{(a - ia \tan(c+dx))\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \left(4a(A - iB) \int \frac{\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx - (2A - 3iB) \int \frac{(a - ia \tan(c+dx))\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx \right) \right)$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \left(-\frac{8ia^3(A - iB) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a + a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a + a}}}{d} - (2A - 3iB) \int \frac{(a - ia \tan(c+dx))\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx \right) \right)$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \left(\frac{(4 - 4i)a^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - (2A - 3iB) \int \frac{(a - ia \tan(c+dx))\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx \right) \right)$$

↓ 4082

3.545. $\int \sqrt{\cot(c+dx)}(a + ia \tan(c+dx))^{3/2}(A + B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{(4-4i)a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}-\frac{a^2(2A-3iB)\int\frac{1}{\sqrt{\tan(c+dx)}}}{d}\right)\right)$$

↓ 65

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{(4-4i)a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}-\frac{2a^2(2A-3iB)\int\frac{1}{1-\frac{ia\tan(c+dx)}{i\tan(c+dx)}}}{d}\right)\right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{2\sqrt[4]{-1}a^{3/2}(2A-3iB)\operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}+\frac{(4-4i)a^{3/2}(A-iB)\operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}\right)\right)$$

input `Int[Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((2*(-1)^(1/4)*a^(3/2)*(2*A - (3*I)*B)*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d + ((4 - 4*I)*a^(3/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d)/2 + (I*a*B*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/d)`

3.545.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 65 `Int[1/(Sqrt[(b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4077 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n)), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

```
rule 4084 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x]
- Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[
e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x]
/; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.545.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 491 vs. 2(158) = 316.

Time = 0.53 (sec) , antiderivative size = 492, normalized size of antiderivative = 2.51

method	result
derivativedivides	$\frac{\sqrt{\frac{1}{\tan(dx+c)}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a \left(-iB \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) \sqrt{ia+a}}{2\sqrt{ia}} \right) \right) \sqrt{-ia} a+2i}{}$
default	$\frac{\sqrt{\frac{1}{\tan(dx+c)}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a \left(-iB \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)}(1+i \tan(dx+c)) \sqrt{ia+a}}{2\sqrt{ia}} \right) \right) \sqrt{-ia} a+2i}{}$

```
input int(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

3.545. $\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

output $\frac{1}{2}d \cdot \left(\frac{1}{\tan(dx+c)}\right)^{1/2} \cdot \tan(dx+c) \cdot \left(a(1+i \tan(dx+c))\right)^{1/2} \cdot a \cdot (-iB \ln\left(\frac{1}{2}(2Ia \tan(dx+c) + 2(a \tan(dx+c)(1+i \tan(dx+c)))^{1/2} \cdot (Ia)^{1/2} + a) / (Ia)^{1/2}\right) \cdot (-Ia)^{1/2} \cdot a + 2I \cdot B \cdot (Ia)^{1/2} \cdot (-Ia)^{1/2} \cdot (a \tan(dx+c) \cdot (1+i \tan(dx+c)))^{1/2} + I \cdot \ln\left((2 \cdot 2^{1/2}) \cdot (-Ia)^{1/2} \cdot (a \tan(dx+c) \cdot (1+i \tan(dx+c)))^{1/2} + 3a \cdot \tan(dx+c) - Ia\right) / (\tan(dx+c) + I) \cdot (Ia)^{1/2} \cdot 2^{1/2} \cdot a + 2A \cdot \ln\left(\frac{1}{2}(2Ia \tan(dx+c) + 2(a \tan(dx+c)(1+i \tan(dx+c)))^{1/2} \cdot (Ia)^{1/2} + a) / (Ia)^{1/2}\right) \cdot (-Ia)^{1/2} \cdot a + 2I \cdot \ln\left(\frac{1}{2}(2Ia \tan(dx+c) + 2(a \tan(dx+c)(1+i \tan(dx+c)))^{1/2} \cdot (Ia)^{1/2} + a) / (Ia)^{1/2}\right) \cdot a \cdot (-Ia)^{1/2} - 2^{1/2} \cdot \ln\left((2 \cdot 2^{1/2}) \cdot (-Ia)^{1/2} \cdot (a \tan(dx+c) \cdot (1+i \tan(dx+c)))^{1/2} + 3a \cdot \tan(dx+c) - Ia\right) / (\tan(dx+c) + I) \cdot (Ia)^{1/2} \cdot a + 2 \cdot \ln\left(\frac{1}{2}(2Ia \tan(dx+c) + 2(a \tan(dx+c)(1+i \tan(dx+c)))^{1/2} \cdot (Ia)^{1/2} + a) / (Ia)^{1/2}\right) \cdot (-Ia)^{1/2} \cdot a) / (Ia)^{1/2} / (-Ia)^{1/2} / (a \tan(dx+c) \cdot (1+i \tan(dx+c)))^{1/2}$

3.545.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 827 vs. $2(148) = 296$.

Time = 0.28 (sec) , antiderivative size = 827, normalized size of antiderivative = 4.22

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \text{Too large to display}$$

input `integrate(cot(dx+c)^(1/2)*(a+I*a*tan(dx+c))^(3/2)*(A+B*tan(dx+c)),x, algorithm="fricas")`

output `1/4*(4*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log(4*((A - I*B)*a^2*e^(I*d*x + I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a) - 4*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log(4*((A - I*B)*a^2*e^(I*d*x + I*c) - sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a) + 4*sqrt(2)*(B*a*e^(3*I*d*x + 3*I*c) - B*a*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)) - sqrt((-4*I*A^2 - 12*A*B + 9*I*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log(-16*(3*(2*I*A + 3*B)*a^2*e^(2*I*d*x + 2*I*c) + (-2*I*A - 3*B)*a^2 + 2*sqrt(2)*sqrt((-4*I*A^2 - 12*A*B + 9*I*B^2)*a^3/d^2)*(I*d*e^(3*I*d*x + 3*I*c) - I*d*e^(I*d*x + I*c)))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/(2*I*A + 3*B) + sqrt((-4*I*A^2 - 12*A*B + 9*I*B^2)*a^3/d^2)*(d*e^(2*I*d*x + 2*I*c) + d)*log(-16*(3*(2*I*A + 3*B)*a^2*e^(2*I*d*x + 2*I*c) + (-2*I*A - 3*B)*a^2 + 2*sqrt(2)*sqrt((-4*I*A^2 - 12*A*B + 9*I*B^2)*a^3/d^2)*(-I*d*e^(3*I*d*x + 3*I*c) + I*d*e^(I*d*x + I*c)))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)...`

3.545.6 Sympy [F]

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (ia(\tan(c + dx) - i))^{3/2} (A + B \tan(c + dx)) \sqrt{\cot(c + dx)} dx$$

input `integrate(cot(d*x+c)**(1/2)*(a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(3/2)*(A + B*tan(c + d*x))*sqrt(cot(c + d*x)), x)`

3.545.7 Maxima [F]

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \int (B \tan(dx+c)+A)(ia \tan(dx+c)+a)^{3/2} \sqrt{\cot(dx+c)} dx$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x+c)+A)*(I*a*tan(d*x+c)+a)^(3/2)*sqrt(cot(d*x+c)),x)`

3.545.8 Giac [F]

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \int (B \tan(dx+c)+A)(ia \tan(dx+c)+a)^{3/2} \sqrt{\cot(dx+c)} dx$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x+c)+A)*(I*a*tan(d*x+c)+a)^(3/2)*sqrt(cot(d*x+c)),x)`

3.545.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \int \sqrt{\cot(c+dx)}(A+B \tan(c+dx))(a+a \tan(c+dx) li)^{3/2} dx$$

input `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2),
x)`

output `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2),
x)`

3.546
$$\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.546.1 Optimal result 5153
 3.546.2 Mathematica [A] (verified) 5154
 3.546.3 Rubi [A] (verified) 5154
 3.546.4 Maple [B] (verified) 5159
 3.546.5 Fricas [B] (verification not implemented) 5160
 3.546.6 Sympy [F] 5161
 3.546.7 Maxima [F] 5162
 3.546.8 Giac [F] 5162
 3.546.9 Mupad [F(-1)] 5162

3.546.1 Optimal result

Integrand size = 38, antiderivative size = 244

$$\int \frac{(a + ia \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx =$$

$$\frac{(-1)^{3/4}a^{3/2}(12A - 11iB) \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{4d}$$

$$\frac{(2 + 2i)a^{3/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{iaB\sqrt{a + ia \tan(c + dx)}}{2d \cot^{3/2}(c + dx)} + \frac{a(4iA + 5B)\sqrt{a + ia \tan(c + dx)}}{4d\sqrt{\cot(c + dx)}}$$

output

```
-1/4*(-1)^(3/4)*a^(3/2)*(12*A-11*I*B)*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)
^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-(2+2*
I)*a^(3/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c
))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+1/2*I*a*B*(a+I*a*tan(d*x+c)
^(1/2)/d/cot(d*x+c)^(3/2)+1/4*a*(4*I*A+5*B)*(a+I*a*tan(d*x+c))^(1/2)/d/cot
(d*x+c)^(1/2)
```


3.546.2 Mathematica [A] (verified)

Time = 10.72 (sec) , antiderivative size = 376, normalized size of antiderivative = 1.54

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(\frac{B \sqrt{\tan(c + dx)} (a + ia \tan(c + dx))}{2d} \right. \\ \left. + \frac{i(\frac{1}{2}a^2(4A - 3iB) - \frac{1}{2}ia^2B)}{a} \left(-\frac{2i\sqrt{2}a \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c + dx)}}{\sqrt{a + ia \tan(c + dx)}}\right) \sqrt{ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)}} + \frac{2ia^{3/2} \operatorname{arcsinh}\left(\frac{\sqrt{ia \tan(c + dx)}}{\sqrt{a}}\right) \sqrt{1 + i \tan(c + dx)} \sqrt{ia \tan(c + dx)}}{d\sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}} \right) \right)$$

input `Integrate[((a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]], x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((B*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(3/2))/(2*d) + (((-1)*((a^2*(4*A - (3*I)*B))/2 - (I/2)*a^2*B))*((-2*I)*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) + ((2*I)*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]))/a + ((I/2)*a^2*(4*A - (3*I)*B)*(-((-1)^(3/4)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]]*Sqrt[a + I*a*Tan[c + d*x]])/Sqrt[1 + I*Tan[c + d*x]]) + Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]))/d)/(2*a))`

3.546.3 Rubi [A] (verified)

Time = 1.48 (sec) , antiderivative size = 233, normalized size of antiderivative = 0.95, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {3042, 4729, 3042, 4077, 27, 3042, 4080, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.546. $\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$

$$\begin{aligned}
& \int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\
& \quad \downarrow \text{4729} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \sqrt{\tan(c + dx)} (i \tan(c + dx) a + a)^{3/2} (A + B \tan(c + dx)) dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \sqrt{\tan(c + dx)} (i \tan(c + dx) a + a)^{3/2} (A + B \tan(c + dx)) dx \\
& \quad \downarrow \text{4077} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{2} \int \frac{1}{2} \sqrt{\tan(c + dx)} \sqrt{i \tan(c + dx) a + a} (a(4A - 3iB) + a(4iA + 5B) \tan(c + dx)) dx \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{4} \int \sqrt{\tan(c + dx)} \sqrt{i \tan(c + dx) a + a} (a(4A - 3iB) + a(4iA + 5B) \tan(c + dx)) dx \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{4} \int \sqrt{\tan(c + dx)} \sqrt{i \tan(c + dx) a + a} (a(4A - 3iB) + a(4iA + 5B) \tan(c + dx)) dx \right) \\
& \quad \downarrow \text{4080} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{4} \left(\frac{\int -\frac{\sqrt{i \tan(c + dx) a + a} (a^2(4iA + 5B) - a^2(12A - 11iB) \tan(c + dx))}{2\sqrt{\tan(c + dx)}} dx}{a} + \frac{a(5B + 4iA) \sqrt{\tan(c + dx)}}{\sqrt{\tan(c + dx)}} \right) \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{4} \left(\frac{a(5B + 4iA) \sqrt{\tan(c + dx)} \sqrt{a + ia \tan(c + dx)}}{d} - \frac{\int \frac{\sqrt{i \tan(c + dx) a + a} (a^2(4iA + 5B) - a^2(12A - 11iB) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx}{2a} \right) \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.546. $\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{a(5B+4iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{\int\frac{\sqrt{i\tan(c+dx)a+a(a^2(4iA+5B)-\sqrt{\tan(c+dx)})}}{\sqrt{\tan(c+dx)}}}{2a}\right)\right)$$

↓ 4084

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{a(5B+4iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{16a^2(B+iA)\int\frac{\sqrt{i\tan(c+dx)a-\sqrt{\tan(c+dx)}}}{\sqrt{\tan(c+dx)}}}{16a^2(B+iA)\int\frac{\sqrt{i\tan(c+dx)a-\sqrt{\tan(c+dx)}}}{\sqrt{\tan(c+dx)}}}\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{a(5B+4iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{16a^2(B+iA)\int\frac{\sqrt{i\tan(c+dx)a-\sqrt{\tan(c+dx)}}}{\sqrt{\tan(c+dx)}}}{16a^2(B+iA)\int\frac{\sqrt{i\tan(c+dx)a-\sqrt{\tan(c+dx)}}}{\sqrt{\tan(c+dx)}}}\right)\right)$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{a(5B+4iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{32ia^4(B+iA)\int\frac{1}{\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a}}}{\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a}}}\right)\right)$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{a(5B+4iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{(16-16i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{1+\sqrt{\tan(c+dx)}}{\sqrt{\tan(c+dx)}}\right)}{d}\right)\right)$$

↓ 4082

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{a(5B+4iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{(16-16i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{1+\sqrt{\tan(c+dx)}}{\sqrt{\tan(c+dx)}}\right)}{d}\right)\right)$$

↓ 65

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{a(5B+4iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{(16-16i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{1+i\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}\right)\right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{a(5B+4iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{2\sqrt[4]{-1}a^{5/2}(11B+12iA)\operatorname{arctan}\left(\frac{c-d\sqrt{\tan(c+dx)}}{d}\right)}{d}\right)\right)$$

```
input Int[((a + I*a*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]], x]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((I/2)*a*B*Tan[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]])/d + (-1/2*((2*(-1)^(1/4))*a^(5/2)*((12*I)*A + 11*B)*ArcTan[(-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]]]/Sqrt[a + I*a*Tan[c + d*x]])/d + ((16 - 16*I)*a^(5/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d)/a + (a*((4*I)*A + 5*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/d)/4
```

3.546.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F x_), x_Symbol] := Simp[a Int[F x, x], x] /; FreeQ[a, x] && !MatchQ[F x, (b_)*(G x_) /; FreeQ[b, x]]
```

```
rule 65 Int[1/(Sqrt[(b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

3.546. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4077 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

rule 4080 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`

rule 4082 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

```
rule 4084 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Sim
p[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x]
- Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[
e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.546.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 454 vs. 2(194) = 388.

Time = 0.66 (sec) , antiderivative size = 455, normalized size of antiderivative = 1.86

method	result
derivativedivides	$-\frac{\sqrt{a(1+i \tan(dx+c))} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \left(-4iB \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia} \sqrt{-ia} \tan(dx+c)+4\right)}{\dots}$
default	$-\frac{\sqrt{a(1+i \tan(dx+c))} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \left(-4iB \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia} \sqrt{-ia} \tan(dx+c)+4\right)}{\dots}$

```
input int((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x,method=_R
ETURNVERBOSE)
```

3.546. $\int \frac{(a+ia \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

output
$$-1/8/d*(a*(1+I*\tan(d*x+c)))^{1/2}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2}*(-4*I*B*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2}*(I*a)^{1/2}*(-I*a)^{1/2}*\tan(d*x+c)+4*I*A*\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2}*(I*a)^{1/2}+a)/(I*a)^{1/2})*a*(-I*a)^{1/2}-8*I*A*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2}*(I*a)^{1/2}*(-I*a)^{1/2}+5*B*(-I*a)^{1/2}*\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2}*(I*a)^{1/2}+a)/(I*a)^{1/2}))*a-10*B*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2}*(I*a)^{1/2}*(-I*a)^{1/2}+16*I*\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2}*(I*a)^{1/2}+a)/(I*a)^{1/2})*a*(-I*a)^{1/2}-8*2^{1/2}*\ln(-(-2*2^{1/2}*(-I*a)^{1/2}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2}+I*a-3*a*\tan(d*x+c))/(tan(d*x+c)+I))*a)^{1/2})*a)/(1/\tan(d*x+c))^{1/2}/(1+I*\tan(d*x+c))/\tan(d*x+c)/(I*a)^{1/2}/(-I*a)^{1/2}$$

3.546.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 913 vs. $2(182) = 364$.

Time = 0.27 (sec) , antiderivative size = 913, normalized size of antiderivative = 3.74

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \text{Too large to display}$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="fracas")`

output

```
-1/16*(16*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(d*e^(4*I*d*x +
4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((A - I*B)*a^2*e^(I*d*x + I*c)
- sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(I*d*e^(2*I*d*x + 2*I*c) - I*d)
*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*
I*d*x + 2*I*c) - 1)))e^(-I*d*x - I*c)/((-I*A - B)*a)) - 16*sqrt(2)*sqrt(-
(-I*A^2 - 2*A*B + I*B^2)*a^3/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x
+ 2*I*c) + d)*log(4*((A - I*B)*a^2*e^(I*d*x + I*c) - sqrt(-(-I*A^2 - 2*A*B
+ I*B^2)*a^3/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(a/(e^(2*I*d*x + 2
*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e
^(-I*d*x - I*c)/((-I*A - B)*a)) - 4*sqrt(2)*((4*A - 7*I*B)*a*e^(5*I*d*x +
5*I*c) + 4*I*B*a*e^(3*I*d*x + 3*I*c) - (4*A - 3*I*B)*a*e^(I*d*x + I*c))*sq
rt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d
*x + 2*I*c) - 1)) - sqrt((144*I*A^2 + 264*A*B - 121*I*B^2)*a^3/d^2)*(d*e^(
4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log(-16*(3*(12*I*A + 11*B)
*a^2*e^(2*I*d*x + 2*I*c) + (-12*I*A - 11*B)*a^2 + 2*sqrt(2)*sqrt((144*I*A^
2 + 264*A*B - 121*I*B^2)*a^3/d^2)*(d*e^(3*I*d*x + 3*I*c) - d*e^(I*d*x + I*
c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^
(2*I*d*x + 2*I*c) - 1)))e^(-2*I*d*x - 2*I*c)/(12*I*A + 11*B)) + sqrt((144
*I*A^2 + 264*A*B - 121*I*B^2)*a^3/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I
*d*x + 2*I*c) + d)*log(-16*(3*(12*I*A + 11*B)*a^2*e^(2*I*d*x + 2*I*c) + ...
```

3.546.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \int \frac{(ia(\tan(c + dx) - i))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

input `integrate((a+I*a*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)**(1/2),x)`

output `Integral((I*a*(tan(c + d*x) - I))**(3/2)*(A + B*tan(c + d*x))/sqrt(cot(c + d*x)), x)`

3.546.7 Maxima [F]

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{3/2}}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)/sqrt(cot(d*x + c)), x)`

3.546.8 Giac [F]

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{3/2}}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(3/2)/sqrt(cot(d*x + c)), x)`

3.546.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) i)^{3/2}}{\sqrt{\cot(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2))/cot(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(3/2))/cot(c + d*x)^(1/2), x)`

3.547 $\int \cot^{\frac{11}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.547.1 Optimal result	5163
3.547.2 Mathematica [A] (verified)	5164
3.547.3 Rubi [A] (verified)	5165
3.547.4 Maple [B] (verified)	5170
3.547.5 Fricas [B] (verification not implemented)	5171
3.547.6 Sympy [F(-1)]	5172
3.547.7 Maxima [B] (verification not implemented)	5172
3.547.8 Giac [F]	5173
3.547.9 Mupad [F(-1)]	5174

3.547.1 Optimal result

Integrand size = 38, antiderivative size = 297

$$\int \cot^{\frac{11}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{(4 + 4i)a^{5/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d} - \frac{8a^2(197A - 195iB)\sqrt{\cot(c + dx)}\sqrt{a + ia \tan(c + dx)}}{315d} + \frac{8a^2(59iA + 60B)\cot^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{315d} + \frac{2a^2(46A - 45iB)\cot^{\frac{5}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{105d} - \frac{2a^2(4iA + 3B)\cot^{\frac{7}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{21d} - \frac{2aA\cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{9d}$$

output

```
(4+4*I)*a^(5/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+8/315*a^2*(59*I*A+60*B)*cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(1/2)/d+2/105*a^2*(46*A-45*I*B)*cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(1/2)/d-2/21*a^2*(4*I*A+3*B)*cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(1/2)/d-8/315*a^2*(197*A-195*I*B)*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/d-2/9*a*A*cot(d*x+c)^(9/2)*(a+I*a*tan(d*x+c))^(3/2)/d
```

3.547. $\int \cot^{\frac{11}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.547.2 Mathematica [A] (verified)

Time = 11.11 (sec) , antiderivative size = 580, normalized size of antiderivative = 1.95

$$\int \cot^{\frac{11}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A + B \tan(c+dx)) dx = \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)} \left(-\frac{2A(a+ia \tan(c+dx))^{5/2}}{9d \tan^{\frac{9}{2}}(c+dx)} \right. \\ \left. + 2 \left(-\frac{a(5iA+9B)(a+ia \tan(c+dx))^{5/2}}{7d \tan^{\frac{7}{2}}(c+dx)} + \frac{a^2(53A-45iB)(a+ia \tan(c+dx))^{5/2} - \frac{63}{4}a^2(iA+B)}{10d \tan^{\frac{5}{2}}(c+dx)} \right) \left(\frac{4i\sqrt{2}a^2 \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) \right)$$

input `Integrate[Cot[c + d*x]^(11/2)*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]), x]`

output $\text{Sqrt}[\text{Cot}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]*((-2*A*(a + I*a*\text{Tan}[c + d*x])^{(5/2)})/(9*d*\text{Tan}[c + d*x]^{(9/2)}) + (2*(-1/7*(a*((5*I)*A + 9*B)*(a + I*a*\text{Tan}[c + d*x])^{(5/2)})/(d*\text{Tan}[c + d*x]^{(7/2)}) + (2*((a^2*(53*A - (45*I)*B)*(a + I*a*\text{Tan}[c + d*x])^{(5/2)})/(10*d*\text{Tan}[c + d*x]^{(5/2)}) - (63*a^2*(I*A + B)*(((4*I)*\text{Sqrt}[2]*a^2*\text{ArcTanh}[(\text{Sqrt}[2]*\text{Sqrt}[I*a*\text{Tan}[c + d*x]])/\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])]*\text{Sqrt}[I*a*\text{Tan}[c + d*x]])/(d*\text{Sqrt}[\text{Tan}[c + d*x]]) - ((4*I)*a^{(5/2)}*\text{ArcSinh}[\text{Sqrt}[I*a*\text{Tan}[c + d*x]]/\text{Sqrt}[a]]*\text{Sqrt}[1 + I*\text{Tan}[c + d*x]]*\text{Sqrt}[I*a*\text{Tan}[c + d*x]])/(d*\text{Sqrt}[\text{Tan}[c + d*x]]*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]) + (5*(-1)^{(3/4)}*a^2*\text{ArcSinh}[(-1)^{(1/4)}*\text{Sqrt}[\text{Tan}[c + d*x]]*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])/(d*\text{Sqrt}[1 + I*\text{Tan}[c + d*x]]) - (2*a^2*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])/(3*d*\text{Tan}[c + d*x]^{(3/2)}) - (((14*I)/3)*a^2*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])/(d*\text{Sqrt}[\text{Tan}[c + d*x]]) - (I*a^{(3/2)}*\text{ArcSinh}[\text{Sqrt}[I*a*\text{Tan}[c + d*x]]/\text{Sqrt}[a]]*\text{Sqrt}[I*a*\text{Tan}[c + d*x]]*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])/(d*\text{Sqrt}[1 + I*\text{Tan}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]))/4)/(7*a)))/(9*a))$

3.547.3 Rubi [A] (verified)

Time = 1.95 (sec) , antiderivative size = 322, normalized size of antiderivative = 1.08, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.526$, Rules used = {3042, 4729, 3042, 4076, 27, 3042, 4076, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot^{\frac{11}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \cot(c + dx)^{11/2}(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx \\ & \quad \downarrow \text{4729} \\ & \sqrt{\tan(c + dx)}\sqrt{\cot(c + dx)} \int \frac{(i \tan(c + dx)a + a)^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\tan(c + dx)}\sqrt{\cot(c + dx)} \int \frac{(i \tan(c + dx)a + a)^{5/2}(A + B \tan(c + dx))}{\tan(c + dx)^{11/2}} dx \\ & \quad \downarrow \text{4076} \end{aligned}$$

3.547. $\int \cot^{\frac{11}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2}{9}\int\frac{3(i\tan(c+dx)a+a)^{3/2}(a(4iA+3B)-a(2A-3iB)\tan(c+dx))}{2\tan^{\frac{9}{2}}(c+dx)}dx-\frac{2aA(a-3B)}{9a}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\int\frac{(i\tan(c+dx)a+a)^{3/2}(a(4iA+3B)-a(2A-3iB)\tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)}dx-\frac{2aA(a-3B)}{9a}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\int\frac{(i\tan(c+dx)a+a)^{3/2}(a(4iA+3B)-a(2A-3iB)\tan(c+dx))}{\tan(c+dx)^{9/2}}dx-\frac{2aA(a-3B)}{9a}\right)$$

↓ 4076

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{2}{7}\int-\frac{\sqrt{i\tan(c+dx)a+a}((46A-45iB)a^2+(38iA+39B)\tan(c+dx)a^2)}{2\tan^{\frac{7}{2}}(c+dx)}dx\right)\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(-\frac{1}{7}\int\frac{\sqrt{i\tan(c+dx)a+a}((46A-45iB)a^2+(38iA+39B)\tan(c+dx)a^2)}{\tan^{\frac{7}{2}}(c+dx)}dx\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(-\frac{1}{7}\int\frac{\sqrt{i\tan(c+dx)a+a}((46A-45iB)a^2+(38iA+39B)\tan(c+dx)a^2)}{\tan(c+dx)^{7/2}}dx\right)\right)$$

↓ 4081

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{1}{7}\left(\frac{2a^2(46A-45iB)\sqrt{a+ia\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}-\frac{2\int\frac{2\sqrt{i\tan(c+dx)a+a}(a^3(59iA+60B)-a^3)}{\tan^{\frac{5}{2}}(c+dx)}}{5a}\right)\right)\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{1}{7}\left(\frac{2a^2(46A-45iB)\sqrt{a+ia\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}-\frac{4\int\frac{\sqrt{i\tan(c+dx)a+a}(a^3(59iA+60B)-a^3)}{\tan^{\frac{5}{2}}(c+dx)}}{5a}\right)\right)\right)$$

↓ 3042

3.547. $\int \cot^{\frac{11}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{1}{7}\left(\frac{2a^2(46A-45iB)\sqrt{a+ia\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}-4\int\frac{\sqrt{i\tan(c+dx)a+a}(a^3(59iA+60B)-a^3)}{\tan(c+dx)^{5/2}}\right)\right)\right)-\frac{4}{5a}$$

↓ 4081

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{1}{7}\left(\frac{2a^2(46A-45iB)\sqrt{a+ia\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}-4\left(\frac{2\int-\frac{\sqrt{i\tan(c+dx)a+a}((197A-195iB)a^4+)}{2\tan^{\frac{3}{2}}(c+dx)}\right)}{3a}\right)\right)\right)-\frac{4}{3a}$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{1}{7}\left(\frac{2a^2(46A-45iB)\sqrt{a+ia\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}-4\left(-\frac{\int\frac{\sqrt{i\tan(c+dx)a+a}((197A-195iB)a^4+2)}{\tan^{\frac{3}{2}}(c+dx)}\right)}{3a}\right)\right)\right)-\frac{4}{3a}$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{1}{7}\left(\frac{2a^2(46A-45iB)\sqrt{a+ia\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}-4\left(-\frac{\int\frac{\sqrt{i\tan(c+dx)a+a}((197A-195iB)a^4+2)}{\tan(c+dx)^{3/2}}\right)}{3a}\right)\right)\right)-\frac{4}{3a}$$

↓ 4081

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{1}{7}\left(\frac{2a^2(46A-45iB)\sqrt{a+ia\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}-4\left(-\frac{2\int\frac{315a^5(iA+B)\sqrt{i\tan(c+dx)a+a}dx}{2\sqrt{\tan(c+dx)}}}{a}\right)}{3a}\right)\right)\right)-\frac{4}{3a}$$

↓ 27

3.547. $\int \cot^{\frac{11}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}(A+B\tan(c+dx))dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A-45iB)\sqrt{a+ia\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} - 4 \left(-\frac{315a^4(B+iA)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx}{3a} \right) \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A-45iB)\sqrt{a+ia\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} - 4 \left(-\frac{315a^4(B+iA)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx}{3a} \right) \right) \right)$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A-45iB)\sqrt{a+ia\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} - 4 \left(-\frac{630ia^6(B+iA)\int\frac{1}{\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a}-ia}}{d} \right) \right) \right)$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \left(\frac{1}{7} \left(\frac{2a^2(46A-45iB)\sqrt{a+ia\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} - 4 \left(-\frac{2a^3(60B+59iA)\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)} \right) \right) \right)$$

input `Int[Cot[c + d*x]^(11/2)*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]), x]`

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*a*A*(a + I*a*Tan[c + d*x])^(3/2))
)/(9*d*Tan[c + d*x]^(9/2)) + ((-2*a^2*((4*I)*A + 3*B)*Sqrt[a + I*a*Tan[c
+ d*x]])/(7*d*Tan[c + d*x]^(7/2)) + ((2*a^2*(46*A - (45*I)*B)*Sqrt[a + I*a
*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) - (4*((-2*a^3*((59*I)*A + 60*B)*S
qrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (((315 - 315*I)*a^(9
/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Ta
n[c + d*x]]])/d - (2*a^4*(197*A - (195*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(
d*Sqrt[Tan[c + d*x]]))/(3*a)))/(5*a))/7)/3
```

3.547.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4027 Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4076 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```



```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.547.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 894 vs. $2(244) = 488$.

Time = 1.60 (sec) , antiderivative size = 895, normalized size of antiderivative = 3.01

method	result
derivativedivides	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{11}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a^2 \left(-1576A \tan(dx+c)^4 \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia} \sqrt{-ia} + 126\right)$
default	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{11}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a^2 \left(-1576A \tan(dx+c)^4 \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia} \sqrt{-ia} + 126\right)$

```
input int(cot(d*x+c)^(11/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_
RETURNVERBOSE)
```

```
output 1/315/d*(1/tan(d*x+c))^(11/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)*a^2*(-
1576*A*tan(d*x+c)^4*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)*(-I*
a)^(1/2)+1260*I*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)
)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^5+472*I*A*t
an(d*x+c)^3*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)*(-I*a)^(1/2)
+480*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3*(a*tan(d*x+c)*(1+I*tan(d*x+c)
)))^(1/2)-190*I*A*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1
/2)*(-I*a)^(1/2)+1260*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(
d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^5-270
*I*B*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)*(-I*a)
^(1/2)-315*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1
/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*(I*a)^(1/2)*2^(1/2)*a*tan(d*x+c)^5
+276*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)
)))^(1/2)+630*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^
(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^5-630*ln(1/2*(
2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I
*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^5-315*I*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*
(I*a)^(1/2)*2^(1/2)*a*tan(d*x+c)^5+1560*I*B*tan(d*x+c)^4*(a*tan(d*x+c)*(1+
I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)*(-I*a)^(1/2)-90*B*(I*a)^(1/2)*(-I*a)^(...
```

3.547.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 629 vs. 2(229) = 458.

Time = 0.27 (sec) , antiderivative size = 629, normalized size of antiderivative = 2.12

$$\int \cot^{\frac{11}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{2 \left(315 \sqrt{2} \sqrt{-\frac{(-iA^2 - 2AB + iB^2)a^5}{d^2}} (de^{(8i dx + 8i c)} - 4 de^{(6i dx + 6i c)} + 6 de^{(4i dx + 4i c)} - 4 de^{(2i dx + 2i c)}) \right)}{\dots}$$

```
input integrate(cot(d*x+c)^(11/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, a
lgorithm="fricas")
```

output `2/315*(315*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((A - I*B)*a^3*e^(I*d*x + I*c) - sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 315*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((A - I*B)*a^3*e^(I*d*x + I*c) - sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 2*sqrt(2)*(2*(323*A - 300*I*B)*a^2*e^(9*I*d*x + 9*I*c) - 27*(61*A - 65*I*B)*a^2*e^(7*I*d*x + 7*I*c) + 63*(37*A - 35*I*B)*a^2*e^(5*I*d*x + 5*I*c) - 1365*(A - I*B)*a^2*e^(3*I*d*x + 3*I*c) + 315*(A - I*B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))/(d*e^(8*I*d*x + 8*I*c) - 4*d*e^(6*I*d*x + 6*I*c) + 6*d*e^(4*I*d*x + 4*I*c) - 4*d*e^(2*I*d*x + 2*I*c) + d)`

3.547.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{11}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(11/2)*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.547.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 4543 vs. $2(229) = 458$.

Time = 4.83 (sec) , antiderivative size = 4543, normalized size of antiderivative = 15.30

$$\int \cot^{\frac{11}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

3.547. $\int \cot^{\frac{11}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

input `integrate(cot(d*x+c)^(11/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/1260*(sqrt(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) + 1)*((5040*((I + 1)*A - (I - 1)*B)*a^2*cos(7*d*x + 7*c) + 16800*(-(I + 1)*A + (I - 1)*B)*a^2*cos(5*d*x + 5*c) + 20496*((I + 1)*A - (I - 1)*B)*a^2*cos(3*d*x + 3*c) + (-9071*I + 9071)*A + (8841*I - 8841)*B)*a^2*cos(d*x + c) + 5040*((I - 1)*A + (I + 1)*B)*a^2*sin(7*d*x + 7*c) + 16800*(-(I - 1)*A - (I + 1)*B)*a^2*sin(5*d*x + 5*c) + 20496*((I - 1)*A + (I + 1)*B)*a^2*sin(3*d*x + 3*c) + (-9071*I - 9071)*A - (8841*I + 8841)*B)*a^2*sin(d*x + c))*cos(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1)) + 8*((-(121*I + 121)*A + (75*I - 75)*B)*a^2*cos(d*x + c) + (-(121*I - 121)*A - (75*I + 75)*B)*a^2*sin(d*x + c) + ((-(121*I + 121)*A + (75*I - 75)*B)*a^2*cos(d*x + c) + (-(121*I - 121)*A - (75*I + 75)*B)*a^2*sin(d*x + c))*cos(2*d*x + 2*c)^2 + ((-(121*I + 121)*A + (75*I - 75)*B)*a^2*cos(d*x + c) + (-(121*I - 121)*A - (75*I + 75)*B)*a^2*sin(d*x + c))*sin(2*d*x + 2*c)^2 + 630*(((I + 1)*A - (I - 1)*B)*a^2*cos(2*d*x + 2*c)^2 + ((I + 1)*A - (I - 1)*B)*a^2*sin(2*d*x + 2*c)^2 + 2*(-(I + 1)*A + (I - 1)*B)*a^2*cos(2*d*x + 2*c) + ((I + 1)*A - (I - 1)*B)*a^2*cos(3*d*x + 3*c) + 2*(((121*I + 121)*A - (75*I - 75)*B)*a^2*cos(d*x + c) + ((121*I - 121)*A + (75*I + 75)*B)*a^2*sin(d*x + c))*cos(2*d*x + 2*c) + 630*(((I - 1)*A + (I + 1)*B)*a^2*cos(2*d*x + 2*c)^2 + ((I - 1)*A + (I + 1)*B)*a^2*sin(2*d*x + 2*c)^2 + 2*(-(I - 1)*A - (I + 1)*B)*a^2*cos(2*d*x + 2*c) + ((I - 1)*A + (I + 1)*B)*a^2*sin(3*d*x + 3*c))*co...`

3.547.8 Giac [F]

$$\int \cot^{\frac{11}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{\frac{5}{2}} \cot(dx + c)^{\frac{11}{2}} dx$$

input `integrate(cot(d*x+c)^(11/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^(11/2), x)`

3.547. $\int \cot^{\frac{11}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.547.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{11}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \int \cot(c+dx)^{11/2} (A+B \tan(c+dx)) (a+a \tan(c+dx) i)^{5/2} dx$$

input `int(cot(c + d*x)^(11/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),x)`

output `int(cot(c + d*x)^(11/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2), x)`

3.548 $\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.548.1 Optimal result	5175
3.548.2 Mathematica [B] (verified)	5176
3.548.3 Rubi [A] (verified)	5177
3.548.4 Maple [B] (verified)	5181
3.548.5 Fricas [B] (verification not implemented)	5182
3.548.6 Sympy [F(-1)]	5183
3.548.7 Maxima [B] (verification not implemented)	5183
3.548.8 Giac [F]	5184
3.548.9 Mupad [F(-1)]	5185

3.548.1 Optimal result

Integrand size = 38, antiderivative size = 251

$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{(4 - 4i)a^{5/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d} + \frac{4a^2(130iA + 133B)\sqrt{\cot(c + dx)}\sqrt{a + ia \tan(c + dx)}}{105d} + \frac{2a^2(80A - 77iB)\cot^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{105d} - \frac{2a^2(10iA + 7B)\cot^{\frac{5}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{35d} - \frac{2aA\cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{7d}$$

output

```
(4-4*I)*a^(5/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+2/105*a^2*(80*A-77*I*B)*cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(1/2)/d-2/35*a^2*(10*I*A+7*B)*cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(1/2)/d+4/105*a^2*(130*I*A+133*B)*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/d-2/7*a*A*cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(3/2)/d
```

3.548.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 524 vs. $2(251) = 502$.

Time = 9.42 (sec) , antiderivative size = 524, normalized size of antiderivative = 2.09

$$\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A + B \tan(c+dx)) dx = \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)} \left(-\frac{2A(a+ia \tan(c+dx))^{5/2}}{7d \tan^{\frac{7}{2}}(c+dx)} \right. \\ \left. + 2 \left(-\frac{a(5iA+7B)(a+ia \tan(c+dx))^{5/2}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{7}{2}a(A-iB) \left(\frac{4i\sqrt{2}a^2 \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)\sqrt{ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{4ia^{5/2} \operatorname{arcsinh}\left(\frac{\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d\sqrt{\tan(c+dx)}} \right) \right) \right)$$

input `Integrate[Cot[c + d*x]^(9/2)*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A*(a + I*a*Tan[c + d*x])^(5/2))/(7*d*Tan[c + d*x]^(7/2)) + (2*(-1/5*(a*((5*I)*A + 7*B)*(a + I*a*Tan[c + d*x])^(5/2))/(d*Tan[c + d*x]^(5/2)) - (7*a*(A - I*B)*((4*I)*Sqrt[2]*a^2*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - ((4*I)*a^(5/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + (5*(-1)^(3/4)*a^2*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]) - (2*a^2*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (((14*I)/3)*a^2*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - (I*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])))/2)/(7*a)`

3.548.3 Rubi [A] (verified)

Time = 1.55 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.07, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {3042, 4729, 3042, 4076, 27, 3042, 4076, 27, 3042, 4081, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{9/2}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^{5/2}(A+B \tan(c+dx))}{\tan(c+dx)^{9/2}} dx$$

$$\downarrow \text{4076}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2}{7} \int \frac{(i \tan(c+dx)a+a)^{3/2}(a(10iA+7B)-a(4A-7iB) \tan(c+dx))}{2 \tan^{\frac{7}{2}}(c+dx)} dx - \frac{2aA(a+ia \tan(c+dx))^{5/2}}{7} \right)$$

$$\downarrow \text{27}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \int \frac{(i \tan(c+dx)a+a)^{3/2}(a(10iA+7B)-a(4A-7iB) \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx - \frac{2aA(a+ia \tan(c+dx))^{5/2}}{7} \right)$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \int \frac{(i \tan(c+dx)a+a)^{3/2}(a(10iA+7B)-a(4A-7iB) \tan(c+dx))}{\tan(c+dx)^{7/2}} dx - \frac{2aA(a+ia \tan(c+dx))^{5/2}}{7} \right)$$

$$\downarrow \text{4076}$$

3.548. $\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{2}{5}\int-\frac{\sqrt{i\tan(c+dx)a+a((80A-77iB)a^2+3(20iA+21B)\tan(c+dx)a^2)}}{2\tan^{\frac{5}{2}}(c+dx)}dx\right.\right.$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(-\frac{1}{5}\int\frac{\sqrt{i\tan(c+dx)a+a((80A-77iB)a^2+3(20iA+21B)\tan(c+dx)a^2)}}{\tan^{\frac{5}{2}}(c+dx)}dx\right.\right.$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(-\frac{1}{5}\int\frac{\sqrt{i\tan(c+dx)a+a((80A-77iB)a^2+3(20iA+21B)\tan(c+dx)a^2)}}{\tan(c+dx)^{5/2}}dx\right.\right.$$

↓ 4081

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{2a^2(80A-77iB)\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{2\int\frac{\sqrt{i\tan(c+dx)a+a(a^3(130iA+133B)-c)}}{\tan^{\frac{3}{2}}(c+dx)}}{3a}\right.\right.\right.$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{2a^2(80A-77iB)\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{2\int\frac{\sqrt{i\tan(c+dx)a+a(a^3(130iA+133B)-c)}}{\tan(c+dx)^{3/2}}}{3a}\right.\right.\right.$$

↓ 4081

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{2a^2(80A-77iB)\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{2\left(\frac{2\int-\frac{105a^4(A-iB)\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx}{a}\right)}{3a}\right.\right.\right.$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{2a^2(80A-77iB)\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{2\left(-210a^3(A-iB)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}\right)}{3a}\right.\right.\right.$$

↓ 3042

3.548. $\int \cot^{\frac{9}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}(A+B\tan(c+dx))dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{2a^2(80A-77iB)\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{2(-210a^3(A-iB)\int\frac{\sqrt{i\tan(c+dx)a-}}{\sqrt{\tan(c+dx)}}}\right)\right)\right)$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{2a^2(80A-77iB)\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{2\left(\frac{420ia^5(A-iB)\int\frac{1}{-2\tan(c+dx)a^2-ia}}{d}\right)}\right)\right)$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{2a^2(80A-77iB)\sqrt{a+ia\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{2\left(-\frac{(210-210i)a^{7/2}(A-iB)\operatorname{arctanh}\left(\frac{1}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}\right)}\right)\right)$$

```
input Int[Cot[c + d*x]^(9/2)*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x
]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*a*A*(a + I*a*Tan[c + d*x])^(3/2
))/((7*d*Tan[c + d*x]^(7/2)) + ((-2*a^2*((10*I)*A + 7*B)*Sqrt[a + I*a*Tan[c
+ d*x]])/(5*d*Tan[c + d*x]^(5/2)) + ((2*a^2*(80*A - (77*I)*B)*Sqrt[a + I*
a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (2*(((-210 + 210*I)*a^(7/2)*(A
- I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c +
d*x]]])/d - (2*a^3*((130*I)*A + 133*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt
[Tan[c + d*x]])))/(3*a))/5)/7)
```

3.548.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`
- rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.548.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 805 vs. $2(206) = 412$.

Time = 1.67 (sec) , antiderivative size = 806, normalized size of antiderivative = 3.21

method	result
derivativedivides	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{9}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a^2 \left(532B\sqrt{ia} \sqrt{-ia} \tan(dx+c)^3 \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - 90iA\sqrt{a}\right)$
default	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{9}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a^2 \left(532B\sqrt{ia} \sqrt{-ia} \tan(dx+c)^3 \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} - 90iA\sqrt{a}\right)$

```
input int(cot(d*x+c)^(9/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

```
output 1/105/d*(1/tan(d*x+c))^(9/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)*a^2*(53
2*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^
(1/2)-90*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*
x+c)))^(1/2)+520*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3*(a*tan(d*x+c)*(
1+I*tan(d*x+c)))^(1/2)+160*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(
d*x+c)*(1+I*tan(d*x+c)))^(1/2)-420*I*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d
*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*t
an(d*x+c)^4+420*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)
)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^4-154*I*B*(
I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
-105*(I*a)^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)
)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^4+105*I
*(I*a)^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^
(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^4+210*ln(1/
2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)
/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)^4+210*I*ln(1/2*(2*I*a*tan(d*x+c)+2
*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(
1/2)*a*tan(d*x+c)^4-42*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)
)*(1+I*tan(d*x+c)))^(1/2)-30*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)
^(1/2)*(-I*a)^(1/2))/(I*a)^(1/2)/(-I*a)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*...
```

3.548.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 577 vs. 2(193) = 386.

Time = 0.27 (sec) , antiderivative size = 577, normalized size of antiderivative = 2.30

$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{2 \left(105 \sqrt{2} \sqrt{-\frac{(iA^2 + 2AB - iB^2)a^5}{d^2}} (de^{(6i dx + 6i c)} - 3 de^{(4i dx + 4i c)} + 3 de^{(2i dx + 2i c)} - d) \log \left(\dots \right) \right)}{\dots}$$

```
input integrate(cot(d*x+c)^(9/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, al
gorithm="fracas")
```

output
$$\begin{aligned} & 2/105*(105*\sqrt{2})*\sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2}*(d*e^{(6*I*d*x + 6*I*c)} - 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} - d)*\log(4*((A - I*B)*a^3*e^{(I*d*x + I*c)} + \sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2}*(d*e^{(2*I*d*x + 2*I*c)} - d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}))e^{(-I*d*x - I*c)}/((-I*A - B)*a^2)) - 105*\sqrt{2})*\sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2}*(d*e^{(6*I*d*x + 6*I*c)} - 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} - d)*\log(4*((A - I*B)*a^3*e^{(I*d*x + I*c)} - \sqrt{-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2}*(d*e^{(2*I*d*x + 2*I*c)} - d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}))e^{(-I*d*x - I*c)}/((-I*A - B)*a^2)) - 2*\sqrt{2})*(2*(-100*I*A - 91*B)*a^2*e^{(7*I*d*x + 7*I*c)} + 7*(55*I*A + 61*B)*a^2*e^{(5*I*d*x + 5*I*c)} + 350*(-I*A - B)*a^2*e^{(3*I*d*x + 3*I*c)} + 105*(I*A + B)*a^2*e^{(I*d*x + I*c)})*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)})/(d*e^{(6*I*d*x + 6*I*c)} - 3*d*e^{(4*I*d*x + 4*I*c)} + 3*d*e^{(2*I*d*x + 2*I*c)} - d) \end{aligned}$$

3.548.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(9/2)*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.548.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 4087 vs. $2(193) = 386$.

Time = 1.69 (sec) , antiderivative size = 4087, normalized size of antiderivative = 16.28

$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(9/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

3.548.
$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

output

```
-1/105*(sqrt(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c)
+ 1)*(3*(140*(-(I - 1)*A - (I + 1)*B)*a^2*cos(7*d*x + 7*c) + 140*((I - 1)*
A + (2*I + 2)*B)*a^2*cos(5*d*x + 5*c) + 21*(-(4*I - 4)*A - (9*I + 9)*B)*a^
2*cos(3*d*x + 3*c) + ((4*I - 4)*A + (49*I + 49)*B)*a^2*cos(d*x + c) + 140*
((I + 1)*A - (I - 1)*B)*a^2*sin(7*d*x + 7*c) + 140*(-(I + 1)*A + (2*I - 2)
*B)*a^2*sin(5*d*x + 5*c) + 21*((4*I + 4)*A - (9*I - 9)*B)*a^2*sin(3*d*x +
3*c) + (-(4*I + 4)*A + (49*I - 49)*B)*a^2*sin(d*x + c))*cos(7/2*arctan2(si
n(2*d*x + 2*c), cos(2*d*x + 2*c) - 1)) + 4*((65*I - 65)*A + (56*I + 56)*B
)*a^2*cos(d*x + c) + (-(65*I + 65)*A + (56*I - 56)*B)*a^2*sin(d*x + c) + (
((65*I - 65)*A + (56*I + 56)*B)*a^2*cos(d*x + c) + (-(65*I + 65)*A + (56*I
- 56)*B)*a^2*sin(d*x + c))*cos(2*d*x + 2*c)^2 + (((65*I - 65)*A + (56*I +
56)*B)*a^2*cos(d*x + c) + (-(65*I + 65)*A + (56*I - 56)*B)*a^2*sin(d*x +
c))*sin(2*d*x + 2*c)^2 + 105*((-(I - 1)*A - (I + 1)*B)*a^2*cos(2*d*x + 2*c)
)^2 + (-(I - 1)*A - (I + 1)*B)*a^2*sin(2*d*x + 2*c)^2 + 2*((I - 1)*A + (I
+ 1)*B)*a^2*cos(2*d*x + 2*c) + (-(I - 1)*A - (I + 1)*B)*a^2*cos(3*d*x + 3
*c) + 2*((-(65*I - 65)*A - (56*I + 56)*B)*a^2*cos(d*x + c) + ((65*I + 65)*
A - (56*I - 56)*B)*a^2*sin(d*x + c))*cos(2*d*x + 2*c) + 105*((I + 1)*A -
(I - 1)*B)*a^2*cos(2*d*x + 2*c)^2 + ((I + 1)*A - (I - 1)*B)*a^2*sin(2*d*x
+ 2*c)^2 + 2*(-(I + 1)*A + (I - 1)*B)*a^2*cos(2*d*x + 2*c) + ((I + 1)*A -
(I - 1)*B)*a^2*sin(3*d*x + 3*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos...
```

3.548.8 Giac [F]

$$\int \cot^{\frac{9}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{\frac{5}{2}} \cot(dx + c)^{\frac{9}{2}} dx$$

input

```
integrate(cot(d*x+c)^(9/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, al
gorithm="giac")
```

output

```
integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^(
9/2), x)
```

3.548.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \int \cot(c+dx)^{9/2}(A+B \tan(c+dx))(a+a \tan(c+dx) i)^{5/2} dx$$

input `int(cot(c + d*x)^(9/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2), x)`

output `int(cot(c + d*x)^(9/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2), x)`

3.549 $\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.549.1 Optimal result	5186
3.549.2 Mathematica [B] (verified)	5187
3.549.3 Rubi [A] (verified)	5187
3.549.4 Maple [B] (verified)	5191
3.549.5 Fricas [B] (verification not implemented)	5192
3.549.6 Sympy [F(-1)]	5193
3.549.7 Maxima [B] (verification not implemented)	5193
3.549.8 Giac [F]	5194
3.549.9 Mupad [F(-1)]	5195

3.549.1 Optimal result

Integrand size = 38, antiderivative size = 205

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx =$$

$$\frac{(4 + 4i)a^{5/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{2a^2(38A - 35iB)\sqrt{\cot(c + dx)}\sqrt{a + ia \tan(c + dx)}}{15d}$$

$$- \frac{2a^2(8iA + 5B)\cot^{\frac{3}{2}}(c + dx)\sqrt{a + ia \tan(c + dx)}}{15d}$$

$$- \frac{2aA\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{5d}$$

output $(-4-4*I)*a^{(5/2)}*(A-I*B)*\operatorname{arctanh}\left(\frac{(1+I)*a^{(1/2)}*\tan(d*x+c)^{(1/2)}}{(a+I*a*\tan(d*x+c))^{(1/2)}}\right)*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/d-2/15*a^2*(8*I*A+5*B)*\cot(d*x+c)^{(3/2)}*(a+I*a*\tan(d*x+c))^{(1/2)}/d+2/15*a^2*(38*A-35*I*B)*\cot(d*x+c)^{(1/2)}*(a+I*a*\tan(d*x+c))^{(1/2)}/d-2/5*a*A*\cot(d*x+c)^{(5/2)}*(a+I*a*\tan(d*x+c))^{(3/2)}/d$

3.549.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 468 vs. $2(205) = 410$.

Time = 8.46 (sec) , antiderivative size = 468, normalized size of antiderivative = 2.28

$$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)} \left(-\frac{2A(a+ia \tan(c+dx))^{5/2}}{5d \tan^{\frac{5}{2}}(c+dx)} \right. \\ \left. + (iA+B) \left(\frac{4i\sqrt{2}a^2 \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{4ia^{5/2} \operatorname{arcsinh}\left(\frac{\sqrt{ia \tan(c+dx)}}{\sqrt{a}}\right) \sqrt{1+i \tan(c+dx)}}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} \right) \right)$$

input `Integrate[Cot[c + d*x]^(7/2)*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A*(a + I*a*Tan[c + d*x])^(5/2))/(5*d*Tan[c + d*x]^(5/2)) + (I*A + B)*(((4*I)*Sqrt[2]*a^2*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - ((4*I)*a^(5/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + (5*(-1)^(3/4)*a^2*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]) - (2*a^2*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (((14*I)/3)*a^2*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - (I*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]]))`

3.549.3 Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4729, 3042, 4076, 27, 3042, 4076, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.549. $\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$$

↓ 3042

$$\int \cot(c+dx)^{7/2}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$$

↓ 4729

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^{5/2}(A+B \tan(c+dx))}{\tan(c+dx)^{7/2}} dx$$

↓ 4076

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2}{5} \int \frac{(i \tan(c+dx)a+a)^{3/2}(a(8iA+5B)-a(2A-5iB) \tan(c+dx))}{2 \tan^{\frac{5}{2}}(c+dx)} dx - \frac{2aA(a-5a^2)}{5a^2} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \int \frac{(i \tan(c+dx)a+a)^{3/2}(a(8iA+5B)-a(2A-5iB) \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx - \frac{2aA(a-5a^2)}{5a^2} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \int \frac{(i \tan(c+dx)a+a)^{3/2}(a(8iA+5B)-a(2A-5iB) \tan(c+dx))}{\tan(c+dx)^{5/2}} dx - \frac{2aA(a-5a^2)}{5a^2} \right)$$

↓ 4076

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \left(\frac{2}{3} \int -\frac{\sqrt{i \tan(c+dx)a+a}((38A-35iB)a^2+(22iA+25B) \tan(c+dx)a^2)}{2 \tan^{\frac{3}{2}}(c+dx)} dx \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \left(-\frac{1}{3} \int \frac{\sqrt{i \tan(c+dx)a+a}((38A-35iB)a^2+(22iA+25B) \tan(c+dx)a^2)}{\tan^{\frac{3}{2}}(c+dx)} dx \right) \right)$$

↓ 3042

3.549. $\int \cot^{\frac{7}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(-\frac{1}{3}\int\frac{\sqrt{i\tan(c+dx)a+a}((38A-35iB)a^2+(22iA+25B)\tan(c+dx)a^2)}{\tan(c+dx)^{3/2}}dx\right.\right.$$

↓ 4081

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\left(\frac{2a^2(38A-35iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-\frac{2\int\frac{30a^3(iA+B)\sqrt{i\tan(c+dx)a+a}dx}{\sqrt{\tan(c+dx)}}}{a}\right)\right.\right.$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\left(\frac{2a^2(38A-35iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-60a^2(B+iA)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}\right.\right.\right.$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\left(\frac{2a^2(38A-35iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-60a^2(B+iA)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}\right.\right.\right.$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\left(\frac{120ia^4(B+iA)\int\frac{1}{-\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a}-ia}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{d}+\frac{2a^2(38A-35iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right)\right.\right.$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\left(\frac{2a^2(38A-35iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-\frac{(60-60i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{1}{\sqrt{\tan(c+dx)}}\right)}{d}\right)\right.\right.$$

input `Int[Cot[c + d*x]^(7/2)*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*a*A*(a + I*a*Tan[c + d*x])^(3/2))
)/(5*d*Tan[c + d*x]^(5/2)) + ((-2*a^2*((8*I)*A + 5*B)*Sqrt[a + I*a*Tan[c
+ d*x]])/(3*d*Tan[c + d*x]^(3/2)) + (((-60 + 60*I)*a^(5/2)*(I*A + B)*ArcTan
h[(((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/d + (
2*a^2*(38*A - (35*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]
)/3)/5)
```

3.549.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4027 Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_)
+ (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4076 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n
+ 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[
(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n
- 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b
*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]
```

```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.549.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 716 vs. $2(168) = 336$.

Time = 1.59 (sec) , antiderivative size = 717, normalized size of antiderivative = 3.50

method	result
derivativedivides	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{7}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a^2 \left(-76A\sqrt{ia} \sqrt{-ia} \tan(dx+c)^2 \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+60iA}\right)}{...}$
default	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{7}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a^2 \left(-76A\sqrt{ia} \sqrt{-ia} \tan(dx+c)^2 \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+60iA}\right)}{...}$

```
input int(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

3.549. $\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

```
output -1/15/d*(1/tan(d*x+c))^(7/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)*a^2*(-7
6*A*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^
(1/2)+60*I*A*(-I*a)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*ta
n(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*tan(d*x+c)^3-15*I^2^(1/2)*(
I*a)^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1
/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3+70*I*B*(-I*a)^(1/2)
*(I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+60*B*(-I*a
)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(
I*a)^(1/2)+a)/(I*a)^(1/2))*a*tan(d*x+c)^3+30*I*(-I*a)^(1/2)*ln(1/2*(2*I*a*
tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1
/2))*a*tan(d*x+c)^3-15*2^(1/2)*(I*a)^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a
*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*
tan(d*x+c)^3+22*I*A*(-I*a)^(1/2)*(I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I
*tan(d*x+c)))^(1/2)-30*(-I*a)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+
c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*tan(d*x+c)^3+10*B
*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
+6*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)*(-I*a)^(1/2))/(-I*a
)^(1/2)/(I*a)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
```

3.549.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 515 vs. 2(157) = 314.

Time = 0.26 (sec) , antiderivative size = 515, normalized size of antiderivative = 2.51

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx =$$

$$2 \left(15 \sqrt{2} \sqrt{-\frac{(-i A^2 - 2 AB + i B^2) a^5}{d^2}} (de^{(4i dx + 4i c)} - 2 de^{(2i dx + 2i c)} + d) \log \left(\frac{4 \left((A - i B) a^3 e^{(i dx + i c)} - \sqrt{-\frac{(-i A^2 - 2 AB + i B^2)}{d^2}} \right)}{\dots} \right) \right)$$

```
input integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, al
gorithm="fricas")
```

output

```
-2/15*(15*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(d*e^(4*I*d*x +
4*I*c) - 2*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((A - I*B)*a^3*e^(I*d*x + I*c)
- sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(I*d*e^(2*I*d*x + 2*I*c) - I*d)
*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*
I*d*x + 2*I*c) - 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 15*sqrt(2)*sqrt
(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I*d*
x + 2*I*c) + d)*log(4*((A - I*B)*a^3*e^(I*d*x + I*c) - sqrt(-(-I*A^2 - 2*A
*B + I*B^2)*a^5/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(a/(e^(2*I*d*x +
2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))
*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 2*sqrt(2)*(2*(13*A - 10*I*B)*a^2*e^(
5*I*d*x + 5*I*c) - 35*(A - I*B)*a^2*e^(3*I*d*x + 3*I*c) + 15*(A - I*B)*a^2
*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2
*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))/(d*e^(4*I*d*x + 4*I*c) - 2*d*e^(2*I
*d*x + 2*I*c) + d)
```

3.549.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(7/2)*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output Timed out

3.549.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1545 vs. $2(157) = 314$.

Time = 0.69 (sec) , antiderivative size = 1545, normalized size of antiderivative = 7.54

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

3.549. $\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

output `2/15*(sqrt(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) + 1)*((30*((I + 1)*A - (I - 1)*B)*a^2*cos(3*d*x + 3*c) + (-31*I + 31)*A + (25*I - 25)*B)*a^2*cos(d*x + c) + 30*((I - 1)*A + (I + 1)*B)*a^2*sin(3*d*x + 3*c) + (-31*I - 31)*A - (25*I + 25)*B)*a^2*sin(d*x + c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1)) + (30*(-(I - 1)*A - (I + 1)*B)*a^2*cos(3*d*x + 3*c) + ((31*I - 31)*A + (25*I + 25)*B)*a^2*cos(d*x + c) + 30*((I + 1)*A - (I - 1)*B)*a^2*sin(3*d*x + 3*c) + (-31*I + 31)*A + (25*I - 25)*B)*a^2*sin(d*x + c))*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1)))*sqrt(a) + 15*(2*((- (I - 1)*A - (I + 1)*B)*a^2*cos(2*d*x + 2*c)^2 + (- (I - 1)*A - (I + 1)*B)*a^2*sin(2*d*x + 2*c)^2 + 2*((I - 1)*A + (I + 1)*B)*a^2*cos(2*d*x + 2*c) + (- (I - 1)*A - (I + 1)*B)*a^2)*arctan2(2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) + 1)^(1/4)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1)) + 2*sin(d*x + c), 2*(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) + 1)^(1/4)*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1)) + 2*cos(d*x + c)) + ((- (I + 1)*A + (I - 1)*B)*a^2*cos(2*d*x + 2*c)^2 + (- (I + 1)*A + (I - 1)*B)*a^2*sin(2*d*x + 2*c)^2 + 2*((I + 1)*A - (I - 1)*B)*a^2*cos(2*d*x + 2*c) + (- (I + 1)*A + (I - 1)*B)*a^2)*log(4*cos(d*x + c)^2 + 4*sin(d*x + c)^2 + 4*sqrt(cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 - 2*cos(2*d*x + 2*c) + 1)*(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c) - 1))^2 + sin(1/2*arc...`

3.549.8 Giac [F]

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{\frac{5}{2}} \cot(dx + c)^{\frac{7}{2}} dx$$

input `integrate(cot(d*x+c)^(7/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^(7/2), x)`

3.549.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{7/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) i)^{5/2} dx$$

input `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2), x)`

output `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2), x)`

3.550 $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.550.1 Optimal result	5196
3.550.2 Mathematica [B] (verified)	5197
3.550.3 Rubi [A] (verified)	5198
3.550.4 Maple [B] (verified)	5202
3.550.5 Fricas [B] (verification not implemented)	5203
3.550.6 Sympy [F(-1)]	5204
3.550.7 Maxima [F]	5205
3.550.8 Giac [F]	5205
3.550.9 Mupad [F(-1)]	5205

3.550.1 Optimal result

Integrand size = 38, antiderivative size = 230

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{2(-1)^{3/4}a^{5/2}B \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d} + \frac{(4 + 4i)a^{5/2}(iA + B)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d} - \frac{2a^2(2iA + B)\sqrt{\cot(c + dx)}\sqrt{a + ia \tan(c + dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}{3d}$$

```
output 2*(-1)^(3/4)*a^(5/2)*B*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(4+4*I)*a^(5/2)*(I*A+B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c)^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-2*a^2*(2*I*A+B)*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c)^(1/2))/d-2/3*a*A*cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c)^(3/2))/d
```

3.550.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 575 vs. $2(230) = 460$.

Time = 8.30 (sec) , antiderivative size = 575, normalized size of antiderivative = 2.50

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A$$

$$+ B \tan(c + dx)) dx = \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)} - \frac{2A(a + ia \tan(c + dx))^{5/2}}{3d \tan^{\frac{3}{2}}(c + dx)}$$

$$+ 2 \left(-\frac{a(5iA+3B)(a+ia \tan(c+dx))^{5/2}}{d\sqrt{\tan(c+dx)}} + \frac{2 \left(ia^4(5iA+3B)\sqrt{a+ia \tan(c+dx)} \left(-\frac{3}{4}(-1)^{3/4} \operatorname{arcsinh} \left(\sqrt[4]{-1} \sqrt{\tan(c+dx)} \right) + \frac{5}{4} \sqrt{1+i \tan(c+dx)} \sqrt{\tan(c+dx)} \right)}{d\sqrt{1+i \tan(c+dx)}} \right) \right)$$

input `Integrate[Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A*(a + I*a*Tan[c + d*x])^(5/2))/(3*d*Tan[c + d*x]^(3/2)) + (2*(-((a*((5*I)*A + 3*B)*(a + I*a*Tan[c + d*x])^(5/2))/(d*Sqrt[Tan[c + d*x]])) + (2*((I*a^4*((5*I)*A + 3*B)*Sqrt[a + I*a*Tan[c + d*x]]*((-3*(-1)^(3/4)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]])/4 + (5*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/4 + (I/2)*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(3/2)))/(d*Sqrt[1 + I*Tan[c + d*x]]) + (a*((-1/4*I)*a^3*(23*A - (15*I)*B) + a^3*((5*I)*A + 3*B))*(((4*I)*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[Tan[c + d*x]])/Sqrt[I*a*Tan[c + d*x]] + ((4*I)*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/(Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + I*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]] + (I*Sqrt[a]*ArcSinh[Sqrt[I*a*Tan[c + d*x]])/Sqrt[a]]*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])))/d)/a)/(3*a)`

3.550.3 Rubi [A] (verified)

Time = 1.39 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.92, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {3042, 4729, 3042, 4076, 27, 3042, 4076, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

↓ 3042

$$\int \cot(c + dx)^{5/2}(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

↓ 4729

$$\sqrt{\tan(c + dx)}\sqrt{\cot(c + dx)} \int \frac{(i \tan(c + dx)a + a)^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{5}{2}}(c + dx)} dx$$

↓ 3042

3.550. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a + a)^{5/2}(A + B \tan(c+dx))}{\tan(c+dx)^{5/2}} dx$$

↓ 4076

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2}{3} \int \frac{3(i \tan(c+dx)a + a)^{3/2}(a(2iA + B) + iaB \tan(c+dx))}{2 \tan^{3/2}(c+dx)} dx - \frac{2aA(a + ia \tan(c+dx))}{3d \tan^{3/2}(c+dx)} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{(i \tan(c+dx)a + a)^{3/2}(a(2iA + B) + iaB \tan(c+dx))}{\tan^{3/2}(c+dx)} dx - \frac{2aA(a + ia \tan(c+dx))}{3d \tan^{3/2}(c+dx)} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{(i \tan(c+dx)a + a)^{3/2}(a(2iA + B) + iaB \tan(c+dx))}{\tan(c+dx)^{3/2}} dx - \frac{2aA(a + ia \tan(c+dx))}{3d \tan^{3/2}(c+dx)} \right)$$

↓ 4076

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(2 \int -\frac{\sqrt{i \tan(c+dx)a + a}((4A - 3iB)a^2 + B \tan(c+dx)a^2)}{2\sqrt{\tan(c+dx)}} dx - \frac{2a^2(B + 2iA)\sqrt{a}}{d\sqrt{\tan(c+dx)}} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(- \int \frac{\sqrt{i \tan(c+dx)a + a}((4A - 3iB)a^2 + B \tan(c+dx)a^2)}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(B + 2iA)\sqrt{a}}{d\sqrt{\tan(c+dx)}} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(- \int \frac{\sqrt{i \tan(c+dx)a + a}((4A - 3iB)a^2 + B \tan(c+dx)a^2)}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(B + 2iA)\sqrt{a}}{d\sqrt{\tan(c+dx)}} \right)$$

↓ 4084

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-4a^2(A - iB) \int \frac{\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx - iaB \int \frac{(a - ia \tan(c+dx))\sqrt{i \tan(c+dx)}}{\sqrt{\tan(c+dx)}} dx \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-4a^2(A - iB) \int \frac{\sqrt{i \tan(c+dx)a + a}}{\sqrt{\tan(c+dx)}} dx - iaB \int \frac{(a - ia \tan(c+dx))\sqrt{i \tan(c+dx)}}{\sqrt{\tan(c+dx)}} dx \right)$$

3.550. $\int \cot^{5/2}(c+dx)(a + ia \tan(c+dx))^{5/2}(A + B \tan(c+dx)) dx$

$$\begin{aligned} & \downarrow 4027 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{8ia^4(A-iB) \int \frac{1}{-\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} - iaB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx \right) \\ & \downarrow 218 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-iaB \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{(4-4i)a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} \right) \\ & \downarrow 4082 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{ia^3B \int \frac{1}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} d \tan(c+dx)}{d} - \frac{(4-4i)a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} \right) \\ & \downarrow 65 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2ia^3B \int \frac{1}{1-\frac{ia \tan(c+dx)}{i \tan(c+dx)a+a}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{d} - \frac{(4-4i)a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} \right) \\ & \downarrow 216 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{(4-4i)a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{2(-1)^{3/4}a^{5/2}B \operatorname{arctan}\left(\frac{(-1)^{3/4}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} \right) \end{aligned}$$

input `Int[Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((2*(-1)^(3/4)*a^(5/2)*B*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/d - ((4 - 4*I)*a^(5/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d - (2*a^2*((2*I)*A + B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - (2*a*A*(a + I*a*Tan[c + d*x])^(3/2))/(3*d*Tan[c + d*x]^(3/2))`

3.550. $\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.550.3.1 Defintions of rubi rules used

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 65 $\text{Int}[1/(\text{Sqrt}[(b_*)(x_)]*\text{Sqrt}[(c_)+(d_*)(x_)]), x_Symbol] \rightarrow \text{Simp}[2 \text{ Subst}[\text{Int}[1/(b-d*x^2), x], x, \text{Sqrt}[b*x]/\text{Sqrt}[c+d*x]], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ !\text{GtQ}[c, 0]$
- rule 216 $\text{Int}[(a_)+(b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$
- rule 218 $\text{Int}[(a_)+(b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4027 $\text{Int}[\text{Sqrt}[(a_)+(b_)*\tan[(e_)+(f_*)(x_)]]/\text{Sqrt}[(c_)+(d_)*\tan[(e_)+(f_*)(x_)]], x_Symbol] \rightarrow \text{Simp}[-2*a*(b/f) \text{ Subst}[\text{Int}[1/(a*c-b*d-2*a^2*x^2), x], x, \text{Sqrt}[c+d*\text{Tan}[e+f*x]]/\text{Sqrt}[a+b*\text{Tan}[e+f*x]]], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c-a*d, 0] \ \&\& \ \text{EqQ}[a^2+b^2, 0] \ \&\& \ \text{NeQ}[c^2+d^2, 0]$
- rule 4076 $\text{Int}[(a_)+(b_)*\tan[(e_)+(f_*)(x_)])^{(m_)*((A_)+(B_)*\tan[(e_)+(f_*)(x_)])^{(n_)}], x_Symbol] \rightarrow \text{Simp}[(-a^2)*(B*c-A*d)*(a+b*\text{Tan}[e+f*x])^{(m-1)}*((c+d*\text{Tan}[e+f*x])^{(n+1)})/(d*f*(b*c+a*d)*(n+1)), x] - \text{Simp}[a/(d*(b*c+a*d)*(n+1)) \text{ Int}[(a+b*\text{Tan}[e+f*x])^{(m-1)}*(c+d*\text{Tan}[e+f*x])^{(n+1)}*\text{Simp}[A*b*d*(m-n-2)-B*(b*c*(m-1)+a*d*(n+1))+(a*A*d*(m+n)-B*(a*c*(m-1)+b*d*(n+1))*\text{Tan}[e+f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[b*c-a*d, 0] \ \&\& \ \text{EqQ}[a^2+b^2, 0] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{LtQ}[n, -1]$

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

rule 4729 `Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.550.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 627 vs. $2(186) = 372$.

Time = 1.66 (sec) , antiderivative size = 628, normalized size of antiderivative = 2.73

method	result
derivativedivides	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a^2 \left(-9iB\sqrt{-ia} \ln\left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}}\right)}{2\sqrt{ia}}$
default	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a^2 \left(-9iB\sqrt{-ia} \ln\left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}}\right)}{2\sqrt{ia}}$

input `int(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNERVERBOSE)`

3.550. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{\frac{5}{2}}(A + B \tan(c + dx)) dx$

```

output -1/3/d*(1/tan(d*x+c))^(5/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)*a^2*(-9*
I*B*(-I*a)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))
)^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*tan(d*x+c)^2+3*I*(I*a)^(1/2)*2^(1/2)
*a*tan(d*x+c))/(tan(d*x+c)+I)*a*tan(d*x+c)^2+14*I*A*(-I*a)^(1/2)*(I*a)^(1/2)
/2*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+12*A*(-I*a)^(1/2)*ln(1
/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a
)/(I*a)^(1/2))*a*tan(d*x+c)^2+6*I*(-I*a)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*tan(d*
x+c)^2-3*(I*a)^(1/2)*2^(1/2)*ln(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1
+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I)*a*tan(d*x+c)^2+6
*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
+6*(-I*a)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)
)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*tan(d*x+c)^2+2*A*(a*tan(d*x+c)*(1+I
*tan(d*x+c)))^(1/2)*(I*a)^(1/2)*(-I*a)^(1/2))/(-I*a)^(1/2)/(I*a)^(1/2)/(a*
tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)

```

3.550.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 780 vs. $2(176) = 352$.

Time = 0.28 (sec) , antiderivative size = 780, normalized size of antiderivative = 3.39

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx =$$

$$24\sqrt{2}\sqrt{-\frac{(iA^2+2AB-iB^2)a^5}{d^2}}(de^{(2i dx+2i c)}-d) \log \left(\frac{4 \left((A-iB)a^3 e^{(i dx+ic)} + \sqrt{-\frac{(iA^2+2AB-iB^2)a^5}{d^2}} (de^{(2i dx+2i c)}-d) \sqrt{e^{(2i dx+2i c)}} \right)}{(-iA-B)a^2} \right)$$

```

input integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, al
gorithm="fracas")

```

output

```
-1/12*(24*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log(4*((A - I*B)*a^3*e^(I*d*x + I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 24*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log(4*((A - I*B)*a^3*e^(I*d*x + I*c) - sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1)))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-I*d*x - I*c)/((-I*A - B)*a^2)) + 8*sqrt(2)*((8*I*A + 3*B)*a^2*e^(3*I*d*x + 3*I*c) + 3*(-2*I*A - B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)) - 3*sqrt(4*I*B^2*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log(-16*(3*B*a^3*e^(2*I*d*x + 2*I*c) - B*a^3 + sqrt(2)*sqrt(4*I*B^2*a^5/d^2)*(I*d*e^(3*I*d*x + 3*I*c) - I*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/(B*a)) + 3*sqrt(4*I*B^2*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*log(-16*(3*B*a^3*e^(2*I*d*x + 2*I*c) - B*a^3 + sqrt(2)*sqrt(4*I*B^2*a^5/d^2))*(-I*d*e^(3*I*d*x + 3*I*c) + I*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-2*I*d*x - 2*I*c)/(B*a)))/(d*e^(2*I*d*x + 2*I*c) - d)
```

3.550.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(5/2)*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output Timed out

3.550.7 Maxima [F]

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \int (B \tan(dx+c)+A)(ia \tan(dx+c)+a)^{\frac{5}{2}} \cot(dx+c)^{\frac{5}{2}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x+c)+A)*(I*a*tan(d*x+c)+a)^(5/2)*cot(d*x+c)^(5/2),x)`

3.550.8 Giac [F]

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \int (B \tan(dx+c)+A)(ia \tan(dx+c)+a)^{\frac{5}{2}} \cot(dx+c)^{\frac{5}{2}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x+c)+A)*(I*a*tan(d*x+c)+a)^(5/2)*cot(d*x+c)^(5/2),x)`

3.550.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \int \cot(c+dx)^{5/2} (A+B \tan(c+dx)) (a+a \tan(c+dx) li)^{5/2} dx$$

input `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),
x)`

output `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2),
x)`

3.551 $\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.551.1 Optimal result	5207
3.551.2 Mathematica [B] (verified)	5208
3.551.3 Rubi [A] (verified)	5209
3.551.4 Maple [B] (verified)	5213
3.551.5 Fricas [B] (verification not implemented)	5214
3.551.6 Sympy [F(-1)]	5215
3.551.7 Maxima [F(-1)]	5216
3.551.8 Giac [F]	5216
3.551.9 Mupad [F(-1)]	5216

3.551.1 Optimal result

Integrand size = 38, antiderivative size = 236

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{(-1)^{3/4}a^{5/2}(2A - 5iB) \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d} + \frac{(4 + 4i)a^{5/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d} + \frac{a^2(2iA - B)\sqrt{a + ia \tan(c + dx)}}{d\sqrt{\cot(c + dx)}} - \frac{2aA\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{3/2}}{d}$$

```
output (-1)^(3/4)*a^(5/2)*(2*A-5*I*B)*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(4+4*I)*a^(5/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+a^2*(2*I*A-B)*(a+I*a*tan(d*x+c))^(1/2)/d/cot(d*x+c)^(1/2)-2*a*A*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(3/2)/d
```

3.551.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 506 vs. $2(236) = 472$.

Time = 8.25 (sec) , antiderivative size = 506, normalized size of antiderivative = 2.14

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A$$

$$+ B \tan(c + dx)) dx = \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(-\frac{2A(a + ia \tan(c + dx))^{5/2}}{d \sqrt{\tan(c + dx)}} \right.$$

$$+ \left(\frac{2ia^3 A \sqrt{a + ia \tan(c + dx)} \left(-\frac{3}{4}(-1)^{3/4} \operatorname{arcsinh} \left(\sqrt[4]{-1} \sqrt{\tan(c + dx)} \right) + \frac{5}{4} \sqrt{1 + i \tan(c + dx)} \sqrt{\tan(c + dx)} + \frac{1}{2} i \sqrt{1 + i \tan(c + dx)} \tan^{\frac{3}{2}}(c + dx) \right)}{d \sqrt{1 + i \tan(c + dx)}} \right)$$

input `Integrate[Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A*(a + I*a*Tan[c + d*x])^(5/2))/(d*Sqrt[Tan[c + d*x]]) + (2*((2*I)*a^3*A*Sqrt[a + I*a*Tan[c + d*x]]*((-3*(-1)^(3/4)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]])/4 + (5*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/4 + (I/2)*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(3/2)))/(d*Sqrt[1 + I*Tan[c + d*x]]) + (a*(2*a^2*A + (I/2)*a^2*((5*I)*A + B))*((-4*I)*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[Tan[c + d*x]])/Sqrt[I*a*Tan[c + d*x]] + ((4*I)*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/(Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + I*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]] + (I*Sqrt[a]*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])))/d)/a)`

3.551.3 Rubi [A] (verified)

Time = 1.45 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.94, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {3042, 4729, 3042, 4076, 27, 3042, 4077, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c + dx)^{3/2}(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{(i \tan(c + dx)a + a)^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{(i \tan(c + dx)a + a)^{5/2}(A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx$$

$$\downarrow \text{4076}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(2 \int \frac{(i \tan(c + dx)a + a)^{3/2}(a(4iA + B) + a(2A + iB) \tan(c + dx))}{2\sqrt{\tan(c + dx)}} dx - \frac{2aA(a + ia \tan(c + dx))}{d\sqrt{\tan(c + dx)}} \right)$$

3.551. $\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\begin{aligned} & \downarrow 27 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\int\frac{(i\tan(c+dx)a+a)^{3/2}(a(4iA+B)+a(2A+iB)\tan(c+dx))}{\sqrt{\tan(c+dx)}}dx-\frac{2aA(a+ia)}{d\sqrt{\tan(c+dx)}}\right) \\ & \downarrow 3042 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\int\frac{(i\tan(c+dx)a+a)^{3/2}(a(4iA+B)+a(2A+iB)\tan(c+dx))}{\sqrt{\tan(c+dx)}}dx-\frac{2aA(a+ia)}{d\sqrt{\tan(c+dx)}}\right) \\ & \downarrow 4077 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\int\frac{\sqrt{i\tan(c+dx)a+a}(3a^2(2iA+B)-a^2(2A-5iB)\tan(c+dx))}{2\sqrt{\tan(c+dx)}}dx+\frac{a^2(-B+ia)}{d\sqrt{\tan(c+dx)}}\right) \\ & \downarrow 27 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\int\frac{\sqrt{i\tan(c+dx)a+a}(3a^2(2iA+B)-a^2(2A-5iB)\tan(c+dx))}{\sqrt{\tan(c+dx)}}dx+\frac{a^2(-B+ia)}{d\sqrt{\tan(c+dx)}}\right) \\ & \downarrow 3042 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\int\frac{\sqrt{i\tan(c+dx)a+a}(3a^2(2iA+B)-a^2(2A-5iB)\tan(c+dx))}{\sqrt{\tan(c+dx)}}dx+\frac{a^2(-B+ia)}{d\sqrt{\tan(c+dx)}}\right) \\ & \downarrow 4084 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(8a^2(B+iA)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx-a(5B+2iA)\int\frac{(a-ia\tan(c+dx))\sqrt{\tan(c+dx)}}{\sqrt{\tan(c+dx)}}dx\right)\right) \\ & \downarrow 3042 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(8a^2(B+iA)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx-a(5B+2iA)\int\frac{(a-ia\tan(c+dx))\sqrt{\tan(c+dx)}}{\sqrt{\tan(c+dx)}}dx\right)\right) \\ & \downarrow 4027 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(-\frac{16ia^4(B+iA)\int\frac{1}{-\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a}-ia}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{d}-a(5B+2iA)\int\frac{(a-ia\tan(c+dx))\sqrt{\tan(c+dx)}}{\sqrt{\tan(c+dx)}}dx\right)\right) \end{aligned}$$

3.551. $\int \cot^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}(A+B\tan(c+dx))dx$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{(8-8i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}-a(5B+2iA)\int\frac{(a-ia\tan(c+dx))^{3/2}}{\sqrt{\tan(c+dx)}}dx\right)\right)$$

↓ 4082

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{(8-8i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}-\frac{a^3(5B+2iA)\int\frac{1}{\sqrt{\tan(c+dx)}}dx}{d}\right)\right)$$

↓ 65

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{(8-8i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}-\frac{2a^3(5B+2iA)\int\frac{1}{1-\frac{ia\tan(c+dx)}{i\tan(c+dx)}}dx}{d}\right)\right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{2\sqrt[4]{-1}a^{5/2}(5B+2iA)\operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}+\frac{(8-8i)a^{5/2}(B+iA)\operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}\right)\right)$$

input `Int[Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((2*(-1)^(1/4)*a^(5/2)*((2*I)*A + 5*B)*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/d + ((8 - 8*I)*a^(5/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/d)/2 + (a^2*((2*I)*A - B)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/d - (2*a*A*(a + I*a*Tan[c + d*x])^(3/2))/(d*Sqrt[Tan[c + d*x]])`

3.551.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4076 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a^2)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(b*c + a*d)*(n + 1))), x] - Simp[a/(d*(b*c + a*d)*(n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*b*d*(m - n - 2) - B*(b*c*(m - 1) + a*d*(n + 1)) + (a*A*d*(m + n) - B*(a*c*(m - 1) + b*d*(n + 1)))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && LtQ[n, -1]`

```
rule 4077 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan
[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c -
a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a
, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] &&
GtQ[m, 1] && !LtQ[n, -1]
```

```
rule 4082 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a
^2 + b^2, 0] && EqQ[A*b + a*B, 0]
```

```
rule 4084 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x]
- Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[
e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.551.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 572 vs. $2(193) = 386$.

Time = 0.54 (sec) , antiderivative size = 573, normalized size of antiderivative = 2.43

$$3.551. \quad \int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

method	result
derivativedivides	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{3}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a^2 \left(6iA \ln\left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}}\right) \sqrt{-ia} a \tan(dx+c)\right)$
default	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{3}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a^2 \left(6iA \ln\left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia+a}}{2\sqrt{ia}}\right) \sqrt{-ia} a \tan(dx+c)\right)$

```
input int(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

```
output 1/2/d*(1/tan(d*x+c))^(3/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)*a^2*(6*I*
A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(
1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)-2*I*ln(-(-2*2^(1/2)*(-I*a)^(
1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c
)+I))*2^(1/2)*(I*a)^(1/2)*a*tan(d*x+c)+3*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*t
an(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)
*a*tan(d*x+c)-2*B*(I*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*t
an(d*x+c)))^(1/2)+4*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*
x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a*tan(d*x+c)-2*ln(-(-
2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(
d*x+c))/(tan(d*x+c)+I))*2^(1/2)*(I*a)^(1/2)*a*tan(d*x+c)-4*A*(a*tan(d*x+c)
*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)*(-I*a)^(1/2)-4*ln(1/2*(2*I*a*tan(d*x+
c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I
*a)^(1/2)*a*tan(d*x+c))/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)/(I*a)^(1/2)/
(-I*a)^(1/2)
```

3.551.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 849 vs. 2(182) = 364.

Time = 0.28 (sec) , antiderivative size = 849, normalized size of antiderivative = 3.60

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, al
gorithm="fricas")
```

3.551. $\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

output $\frac{1}{4}(8\sqrt{2})\sqrt{-(-IA^2 - 2AB + IB^2)a^5/d^2}(de^{(2I dx + 2Ic)} + d)\log(4((A - IB)a^3e^{(I dx + Ic)} - \sqrt{-(-IA^2 - 2AB + IB^2)a^5/d^2})(I de^{(2I dx + 2Ic)} - Id)\sqrt{a/(e^{(2I dx + 2Ic)} + 1)})\sqrt{(Ie^{(2I dx + 2Ic)} + I)/(e^{(2I dx + 2Ic)} - 1))e^{(-I dx - Ic)/((-IA - B)a^2)} - 8\sqrt{2})\sqrt{-(-IA^2 - 2AB + IB^2)a^5/d^2}(de^{(2I dx + 2Ic)} + d)\log(4((A - IB)a^3e^{(I dx + Ic)} - \sqrt{-(-IA^2 - 2AB + IB^2)a^5/d^2})(-I de^{(2I dx + 2Ic)} + Id)\sqrt{a/(e^{(2I dx + 2Ic)} + 1)})\sqrt{(Ie^{(2I dx + 2Ic)} + I)/(e^{(2I dx + 2Ic)} - 1))e^{(-I dx - Ic)/((-IA - B)a^2)} - 4\sqrt{2})((2A - IB)a^2e^{(3I dx + 3Ic)} + (2A + IB)a^2e^{(I dx + Ic)})\sqrt{a/(e^{(2I dx + 2Ic)} + 1)}\sqrt{(Ie^{(2I dx + 2Ic)} + I)/(e^{(2I dx + 2Ic)} - 1)} - \sqrt{((4IA^2 + 20AB - 25IB^2)a^5/d^2)(de^{(2I dx + 2Ic)} + d)\log(-16(3(2IA + 5B)a^3e^{(2I dx + 2Ic)} + (-2IA - 5B)a^3 + 2\sqrt{2})\sqrt{((4IA^2 + 20AB - 25IB^2)a^5/d^2)(de^{(3I dx + 3Ic)} - de^{(I dx + Ic)})}\sqrt{a/(e^{(2I dx + 2Ic)} + 1)})\sqrt{(Ie^{(2I dx + 2Ic)} + I)/(e^{(2I dx + 2Ic)} - 1))e^{(-2I dx - 2Ic)/((2IA + 5B)a)} + \sqrt{((4IA^2 + 20AB - 25IB^2)a^5/d^2)(de^{(2I dx + 2Ic)} + d)\log(-16(3(2IA + 5B)a^3e^{(2I dx + 2Ic)} + (-2IA - 5B)a^3 - 2\sqrt{2})\sqrt{((4IA^2 + 20AB - 25IB^2)a^5/d^2)(de^{(3I dx + 3Ic)} - de^{(I dx + Ic)})}\sqrt{a/(e^{(2I dx + 2Ic)} + ...$

3.551.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(3/2)*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output Timed out

3.551.7 Maxima [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `Timed out`

3.551.8 Giac [F]

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{\frac{5}{2}} \cot(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^(3/2), x)`

3.551.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{3/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) 1i)^{5/2} dx$$

input `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2), x)`

output `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2), x)`

3.551. $\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.552 $\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.552.1 Optimal result	5217
3.552.2 Mathematica [A] (verified)	5218
3.552.3 Rubi [A] (verified)	5219
3.552.4 Maple [B] (verified)	5223
3.552.5 Fricas [B] (verification not implemented)	5224
3.552.6 Sympy [F(-1)]	5225
3.552.7 Maxima [F(-1)]	5225
3.552.8 Giac [F]	5225
3.552.9 Mupad [F(-1)]	5226

3.552.1 Optimal result

Integrand size = 38, antiderivative size = 246

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx =$$

$$\frac{(-1)^{3/4}a^{5/2}(20iA + 23B) \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{4d}$$

$$+ \frac{(4 - 4i)a^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d}$$

$$- \frac{a^2(4A - 7iB)\sqrt{a + ia \tan(c + dx)}}{4d\sqrt{\cot(c + dx)}} + \frac{iaB(a + ia \tan(c + dx))^{3/2}}{2d\sqrt{\cot(c + dx)}}$$

```
output -1/4*(-1)^(3/4)*a^(5/2)*(20*I*A+23*B)*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)
^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(4-4*
I)*a^(5/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c
))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-1/4*a^2*(4*A-7*I*B)*(a+I*a*t
an(d*x+c)^(1/2)/d/cot(d*x+c)^(1/2)+1/2*I*a*B*(a+I*a*tan(d*x+c))^(3/2)/d/c
ot(d*x+c)^(1/2)
```


3.552.2 Mathematica [A] (verified)

Time = 11.23 (sec) , antiderivative size = 454, normalized size of antiderivative = 1.85

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)} \left(\frac{ia^2 B \sqrt{a+ia \tan(c+dx)} \left(-\frac{3}{4}(-1)^{3/4} \operatorname{arcsinh}\left(\sqrt[4]{-1}\sqrt{\tan(c+dx)}\right)\right)}{\dots} \right. \\ \left. + \frac{a(iaA+aB) \left(-\frac{4i\sqrt{2}a \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)\sqrt{\tan(c+dx)}}{\sqrt{ia \tan(c+dx)}} + \frac{4ia^{3/2} \operatorname{arcsinh}\left(\frac{\sqrt{ia \tan(c+dx)}}{\sqrt{a}}\right)\sqrt{1+i \tan(c+dx)}\sqrt{\tan(c+dx)}}{\sqrt{ia \tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}\right)}{d} \right)$$

input `Integrate[Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((I*a^2*B*Sqrt[a + I*a*Tan[c + d*x]]*((-3*(-1)^(3/4)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]])]/4 + (5*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/4 + (I/2)*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(3/2)))/(d*Sqrt[1 + I*Tan[c + d*x]]) + (a*(I*a*A + a*B)*((-4*I)*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])]/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[Tan[c + d*x]]/Sqrt[I*a*Tan[c + d*x]] + ((4*I)*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/(Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + I*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]] + (I*Sqrt[a]*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])))/d`

3.552.3 Rubi [A] (verified)

Time = 1.47 (sec) , antiderivative size = 233, normalized size of antiderivative = 0.95, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {3042, 4729, 3042, 4077, 27, 3042, 4077, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^{5/2}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^{5/2}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

$$\downarrow \text{4077}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \int \frac{(i \tan(c+dx)a+a)^{3/2}(a(4A-iB)+a(4iA+7B) \tan(c+dx))}{2\sqrt{\tan(c+dx)}} dx + \frac{iaB\sqrt{\tan(c+dx)}}{2} \right)$$

$$\downarrow \text{27}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \int \frac{(i \tan(c+dx)a+a)^{3/2}(a(4A-iB)+a(4iA+7B) \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx + \frac{iaB\sqrt{\tan(c+dx)}}{4} \right)$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \int \frac{(i \tan(c+dx)a+a)^{3/2}(a(4A-iB)+a(4iA+7B) \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx + \frac{iaB\sqrt{\tan(c+dx)}}{4} \right)$$

$$\downarrow \text{4077}$$

3.552. $\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\int\frac{\sqrt{i\tan(c+dx)a+a}(3(4A-3iB)a^2+(20iA+23B)\tan(c+dx)a^2)}{2\sqrt{\tan(c+dx)}}dx-a^2\right.\right.$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{1}{2}\int\frac{\sqrt{i\tan(c+dx)a+a}(3(4A-3iB)a^2+(20iA+23B)\tan(c+dx)a^2)}{\sqrt{\tan(c+dx)}}dx-a^2\right.\right.$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{1}{2}\int\frac{\sqrt{i\tan(c+dx)a+a}(3(4A-3iB)a^2+(20iA+23B)\tan(c+dx)a^2)}{\sqrt{\tan(c+dx)}}dx-a^2\right.\right.$$

↓ 4084

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{1}{2}\left(32a^2(A-iB)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx-a(20A-23iB)\int\frac{(a-ia\tan(c+dx))}{\sqrt{\tan(c+dx)}}dx\right.\right.\right.$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{1}{2}\left(32a^2(A-iB)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx-a(20A-23iB)\int\frac{(a-ia\tan(c+dx))}{\sqrt{\tan(c+dx)}}dx\right.\right.\right.$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{1}{2}\left(-\frac{64ia^4(A-iB)\int\frac{1}{-\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a}-ia}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{d}-a(20A-23iB)\int\frac{(a-ia\tan(c+dx))}{\sqrt{\tan(c+dx)}}dx\right.\right.\right.$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{1}{2}\left(\frac{(32-32i)a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}-a(20A-23iB)\int\frac{(a-ia\tan(c+dx))}{\sqrt{\tan(c+dx)}}dx\right.\right.\right.$$

↓ 4082

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{1}{2}\left(\frac{(32-32i)a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}-\frac{a^3(20A-23iB)\int\frac{(a-ia\tan(c+dx))}{\sqrt{\tan(c+dx)}}dx}{d}\right.\right.\right.$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \left(\frac{1}{2} \left(\frac{(32-32i)a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} - \frac{2a^3(20A-23iB)}{d} \right) \right) \right)$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \left(\frac{1}{2} \left(\frac{2\sqrt[4]{-1}a^{5/2}(20A-23iB)\operatorname{arctan}\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} + \frac{(32-32i)a^{5/2}(A-iB)}{d} \right) \right) \right)$$

input `Int[Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((I/2)*a*B*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(3/2))/d + (((2*(-1)^(1/4))*a^(5/2)*(20*A - (23*I)*B)*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]]]/Sqrt[a + I*a*Tan[c + d*x]])/d + ((32 - 32*I)*a^(5/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d)/2 - (a^2*(4*A - (7*I)*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/d)/4)`

3.552.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 65 `Int[1/(Sqrt[(b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4077 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.552.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 571 vs. $2(196) = 392$.

Time = 0.53 (sec) , antiderivative size = 572, normalized size of antiderivative = 2.33

method	result
derivativedivides	$-\frac{\sqrt{\frac{1}{\tan(dx+c)}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a^2 \left(-18iB\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} + 4B\sqrt{ia} \sqrt{-ia} \tan(dx+c) \right)}{\dots}$
default	$-\frac{\sqrt{\frac{1}{\tan(dx+c)}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} a^2 \left(-18iB\sqrt{ia} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} + 4B\sqrt{ia} \sqrt{-ia} \tan(dx+c) \right)}{\dots}$

```
input int(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_R
ETURNVERBOSE)
```

```
output -1/8/d*(1/tan(d*x+c))^(1/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)*a^2*(-18
*I*B*(-I*a)^(1/2)*(I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+4*B*(I
*a)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+9*
I*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)
^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a-8*I*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*ta
n(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*2^(1/
2)*(I*a)^(1/2)*a+8*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)*(-I
*a)^(1/2)-12*A*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(
1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a+8*2^(1/2)*ln((2*2^(1/2)*(-
I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan
(d*x+c)+I)*(I*a)^(1/2)*a-16*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1
+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*(-I*a)^(1/2)-16*ln(1/2
*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)*(I*a)^(1/2)+a)/
(I*a)^(1/2))*(-I*a)^(1/2)*a)/(-I*a)^(1/2)/(I*a)^(1/2)/(a*tan(d*x+c)*(1+I*t
an(d*x+c)))^(1/2)
```

3.552.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 923 vs. $2(184) = 368$.

Time = 0.27 (sec) , antiderivative size = 923, normalized size of antiderivative = 3.75

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")
```

```
output 1/16*(32*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((A - I*B)*a^3*e^(I*d*x + I*c) + sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 32*sqrt(2)*sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((A - I*B)*a^3*e^(I*d*x + I*c) - sqrt(-(I*A^2 + 2*A*B - I*B^2)*a^5/d^2)*(d*e^(2*I*d*x + 2*I*c) - d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-I*d*x - I*c)/((-I*A - B)*a^2)) + 4*sqrt(2)*((4*I*A + 11*B)*a^2*e^(5*I*d*x + 5*I*c) - 4*B*a^2*e^(3*I*d*x + 3*I*c) + (-4*I*A - 7*B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)) - sqrt((-400*I*A^2 - 920*A*B + 529*I*B^2)*a^5/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log(-16*(3*(20*I*A + 23*B)*a^3*e^(2*I*d*x + 2*I*c) + (-20*I*A - 23*B)*a^3 + 2*sqrt(2)*sqrt((-400*I*A^2 - 920*A*B + 529*I*B^2)*a^5/d^2)*(I*d*e^(3*I*d*x + 3*I*c) - I*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-2*I*d*x - 2*I*c)/((20*I*A + 23*B)*a) + sqrt((-400*I*A^2 - 920*A*B + 529*I*B^2)*a^5/d^2)*(d*e^(4*I*d*x + 4*I*c) + 2*d*e^(2*I*d*x + 2*I*c) + d)*log(-16*(3*(20*I*A + 23*B)*a^3*e^(2*I*d*x + 2...))
```

3.552.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(1/2)*(a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.552.7 Maxima [F(-1)]

Timed out.

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `Timed out`

3.552.8 Giac [F]

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{5/2} \sqrt{\cot(dx + c)} dx$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)*sqrt(cot(d*x + c)), x)`

3.552.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \int \sqrt{\cot(c+dx)}(A+B \tan(c+dx))(a+a \tan(c+dx) \text{li})^{5/2} dx$$

input `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2), x)`

output `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2), x)`

3.553
$$\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.553.1 Optimal result 5227
 3.553.2 Mathematica [A] (verified) 5228
 3.553.3 Rubi [A] (verified) 5229
 3.553.4 Maple [B] (verified) 5234
 3.553.5 Fricas [B] (verification not implemented) 5234
 3.553.6 Sympy [F(-1)] 5235
 3.553.7 Maxima [F] 5236
 3.553.8 Giac [F] 5236
 3.553.9 Mupad [F(-1)] 5236

3.553.1 Optimal result

Integrand size = 38, antiderivative size = 292

$$\int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx =$$

$$\frac{(-1)^{3/4}a^{5/2}(46A - 45iB) \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{(4 + 4i)a^{5/2}(A - iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}} -$$

$$\frac{a^2(2A - 3iB)\sqrt{a + ia \tan(c + dx)}}{4d \cot^{3/2}(c + dx)} + \frac{a^2(18iA + 19B)\sqrt{a + ia \tan(c + dx)}}{8d\sqrt{\cot(c + dx)}} + \frac{iaB(a + ia \tan(c + dx))^{3/2}}{3d \cot^{3/2}(c + dx)}$$

```
output -1/8*(-1)^(3/4)*a^(5/2)*(46*A-45*I*B)*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)
^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-(4+4*
I)*a^(5/2)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c
))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-1/4*a^2*(2*A-3*I*B)*(a+I*a*t
an(d*x+c)^(1/2)/d/cot(d*x+c)^(3/2)+1/8*a^2*(18*I*A+19*B)*(a+I*a*tan(d*x+c
))^(1/2)/d/cot(d*x+c)^(1/2)+1/3*I*a*B*(a+I*a*tan(d*x+c)^(3/2)/d/cot(d*x+c
)^(3/2)
```

3.553.2 Mathematica [A] (verified)

Time = 11.27 (sec) , antiderivative size = 524, normalized size of antiderivative = 1.79

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(\frac{B \sqrt{\tan(c + dx)} (a + ia \tan(c + dx))}{3d} \right.$$

$$+ \frac{ia^3 (6A - 5iB) \sqrt{a + ia \tan(c + dx)} \left(-\frac{3}{4} (-1)^{3/4} \operatorname{arcsinh} \left(\sqrt[4]{-1} \sqrt{\tan(c + dx)} \right) + \frac{5}{4} \sqrt{1 + i \tan(c + dx)} \sqrt{\tan(c + dx)} + \frac{1}{2} i \sqrt{1 + i \tan(c + dx)} \tan^{\frac{3}{2}}(c + dx) \right)}{2d \sqrt{1 + i \tan(c + dx)}} \right)$$

input `Integrate[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]], x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((B*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(5/2))/(3*d) + (((I/2)*a^3*(6*A - (5*I)*B)*Sqrt[a + I*a*Tan[c + d*x]]*((-3*(-1)^(3/4)*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]])]/4 + (5*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/4 + (I/2)*Sqrt[1 + I*Tan[c + d*x]]*Tan[c + d*x]^(3/2)))/(d*Sqrt[1 + I*Tan[c + d*x]]) + (a*((a^2*(6*A - (5*I)*B))/2 - (I/2)*a^2*B)*((-4*I)*Sqrt[2]*a*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*Sqrt[Tan[c + d*x]])/Sqrt[I*a*Tan[c + d*x]] + ((4*I)*a^(3/2)*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[1 + I*Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/(Sqrt[I*a*Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + I*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]] + (I*Sqrt[a]*ArcSinh[Sqrt[I*a*Tan[c + d*x]]/Sqrt[a]]*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/(Sqrt[1 + I*Tan[c + d*x]]*Sqrt[I*a*Tan[c + d*x]])))/d)/(3*a)`

3.553.3 Rubi [A] (verified)

Time = 1.85 (sec) , antiderivative size = 286, normalized size of antiderivative = 0.98, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4729, 3042, 4077, 27, 3042, 4077, 27, 3042, 4080, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \sqrt{\tan(c + dx)} (i \tan(c + dx) a + a)^{5/2} (A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \sqrt{\tan(c + dx)} (i \tan(c + dx) a + a)^{5/2} (A + B \tan(c + dx)) dx$$

$$\downarrow \text{4077}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{3} \int \frac{3}{2} \sqrt{\tan(c + dx)} (i \tan(c + dx) a + a)^{3/2} (a(2A - iB) + a(2iA + 3B) \tan(c + dx)) dx \right)$$

$$\downarrow \text{27}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{2} \int \sqrt{\tan(c + dx)} (i \tan(c + dx) a + a)^{3/2} (a(2A - iB) + a(2iA + 3B) \tan(c + dx)) dx \right)$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{2} \int \sqrt{\tan(c + dx)} (i \tan(c + dx) a + a)^{3/2} (a(2A - iB) + a(2iA + 3B) \tan(c + dx)) dx \right)$$

$$\downarrow \text{4077}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{1}{2}\int\frac{1}{2}\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a((14A-13iB)a^2+(18iA+19B)\tan(c+dx))}\right)\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{1}{4}\int\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a((14A-13iB)a^2+(18iA+19B)\tan(c+dx))}\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{1}{4}\int\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a((14A-13iB)a^2+(18iA+19B)\tan(c+dx))}\right)\right)$$

↓ 4080

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{1}{4}\left(\frac{\int-\frac{\sqrt{i\tan(c+dx)a+a(a^3(18iA+19B)-a^3(46A-45iB)\tan(c+dx))}{2\sqrt{\tan(c+dx)}}dx}{a}+a^2(19B+18iA)\right)\right)\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{1}{4}\left(\frac{a^2(19B+18iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{\int\frac{\sqrt{i\tan(c+dx)a+a(a^3(18iA+19B)-a^3(46A-45iB)\tan(c+dx))}}{2\sqrt{\tan(c+dx)}}dx}{a}\right)\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{1}{4}\left(\frac{a^2(19B+18iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{\int\frac{\sqrt{i\tan(c+dx)a+a(a^3(18iA+19B)-a^3(46A-45iB)\tan(c+dx))}}{2\sqrt{\tan(c+dx)}}dx}{a}\right)\right)\right)$$

↓ 4084

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{1}{4}\left(\frac{a^2(19B+18iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{64a^3(B+iA)\int\frac{\sqrt{i\tan(c+dx)a+a(a^3(18iA+19B)-a^3(46A-45iB)\tan(c+dx))}}{2\sqrt{\tan(c+dx)}}dx}{a}\right)\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{1}{4}\left(\frac{a^2(19B+18iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{64a^3(B+iA)\int\frac{\sqrt{i\tan(c+dx)a+a(a^3(18iA+19B)-a^3(46A-45iB)\tan(c+dx))}}{2\sqrt{\tan(c+dx)}}dx}{a}\right)\right)\right)$$

3.553. $\int \frac{(a+ia\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{1}{4}\left(\frac{a^2(19B+18iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{-a^2(45B+46iA)\int\frac{\sqrt{\tan(c+dx)}}{\cot(c+dx)}dx}{a}\right)\right)\right)$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{1}{4}\left(\frac{a^2(19B+18iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{(64-64i)a^{7/2}(B+iA)\arctan\left(\frac{\sqrt{\tan(c+dx)}}{\cot(c+dx)}\right)}{a}\right)\right)\right)$$

↓ 4082

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{1}{4}\left(\frac{a^2(19B+18iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{(64-64i)a^{7/2}(B+iA)\arctan\left(\frac{\sqrt{\tan(c+dx)}}{\cot(c+dx)}\right)}{a}\right)\right)\right)$$

↓ 65

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{1}{4}\left(\frac{a^2(19B+18iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{(64-64i)a^{7/2}(B+iA)\arctan\left(\frac{\sqrt{\tan(c+dx)}}{\cot(c+dx)}\right)}{a}\right)\right)\right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{1}{4}\left(\frac{a^2(19B+18iA)\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}{d}-\frac{2\sqrt[4]{-1}a^{7/2}(45B+46iA)\arctan\left(\frac{\sqrt{\tan(c+dx)}}{\cot(c+dx)}\right)}{a}\right)\right)\right)$$

```
input Int[((a + I*a*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]], x]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((I/3)*a*B*Tan[c + d*x]^(3/2)*(a +
I*a*Tan[c + d*x])^(3/2))/d + (-1/2*(a^2*(2*A - (3*I)*B)*Tan[c + d*x]^(3/2)
*Sqrt[a + I*a*Tan[c + d*x]])/d + (-1/2*((2*(-1)^(1/4)*a^(7/2)*((46*I)*A +
45*B)*ArcTan[(-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c +
d*x]]])/d + ((64 - 64*I)*a^(7/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[T
an[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d)/a + (a^2*((18*I)*A + 19*B)*S
qrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]])/d)/4)/2)
```

3.553.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 65 Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Sub
st[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d
}, x] && !GtQ[c, 0]
```

```
rule 216 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4027 Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_)
+ (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

rule 4077 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n) + B*(a*c*(m - 1) - b*d*(n + 1)) - (B*(b*c - a*d)*(m - 1) - d*(A*b + a*B)*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[m, 1] && !LtQ[n, -1]`

rule 4080 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

rule 4729 `Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.553.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 541 vs. $2(234) = 468$.

Time = 0.58 (sec) , antiderivative size = 542, normalized size of antiderivative = 1.86

method	result
derivativedivides	$-\frac{\sqrt{a(1+i \tan(dx+c))} a \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \left(57B\sqrt{-ia} \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia}}{2\sqrt{ia}} \right) \right)}{\dots}$
default	$-\frac{\sqrt{a(1+i \tan(dx+c))} a \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \left(57B\sqrt{-ia} \ln \left(\frac{2ia \tan(dx+c)+2\sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{ia}}{2\sqrt{ia}} \right) \right)}{\dots}$

```
input int((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x,method=_R
ETURNVERBOSE)
```

```
output -1/48/d*(a*(1+I*tan(d*x+c)))^(1/2)*a*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
*(57*B*(-I*a)^(1/2)*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+
c))))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2)*a-114*B*(a*tan(d*x+c)*(1+I*tan(d*x+
c)))^(1/2)*(I*a)^(1/2)*(-I*a)^(1/2)-52*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan
(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+16*B*(I*a)^(1/2)*(-I*a)^(1/2)*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2+54*I*A*ln(1/2*(2*I*a*tan
(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2)
)*(-I*a)^(1/2)*a-108*I*A*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d
*x+c)))^(1/2)+24*A*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)
))^(1/2)*tan(d*x+c)+192*I*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x+c)*(1+I*tan
(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*a*(-I*a)^(1/2)-96*2^(1/2)*ln(-
(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan
(d*x+c))/(tan(d*x+c)+I))*(I*a)^(1/2)*a)/(1/tan(d*x+c))^(1/2)/(1+I*tan(d*x+
c))/tan(d*x+c)/(I*a)^(1/2)/(-I*a)^(1/2)
```

3.553.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1013 vs. $2(220) = 440$.

Time = 0.29 (sec) , antiderivative size = 1013, normalized size of antiderivative = 3.47

$$\int \frac{(a + ia \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \text{Too large to display}$$

3.553. $\int \frac{(a+ia \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="fricas")`

output `-1/96*(192*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((A - I*B)*a^3*e^(I*d*x + I*c) - sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(I*d*e^(2*I*d*x + 2*I*c) - I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 192*sqrt(2)*sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*log(4*((A - I*B)*a^3*e^(I*d*x + I*c) - sqrt(-(-I*A^2 - 2*A*B + I*B^2)*a^5/d^2)*(-I*d*e^(2*I*d*x + 2*I*c) + I*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-I*d*x - I*c)/((-I*A - B)*a^2)) - 4*sqrt(2)*((66*A - 91*I*B)*a^2*e^(7*I*d*x + 7*I*c) + 7*(66*A - I*B)*a^2*e^(5*I*d*x + 5*I*c) - (66*A - 59*I*B)*a^2*e^(3*I*d*x + 3*I*c) - 3*(14*A - 13*I*B)*a^2*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)) - 3*sqrt((2116*I*A^2 + 4140*A*B - 2025*I*B^2)*a^5/d^2)*(d*e^(6*I*d*x + 6*I*c) + 3*d*e^(4*I*d*x + 4*I*c) + 3*d*e^(2*I*d*x + 2*I*c) + d)*log(-16*(3*(46*I*A + 45*B)*a^3*e^(2*I*d*x + 2*I*c) + (-46*I*A - 45*B)*a^3 + 2*sqrt(2)*sqrt((2116*I*A^2 + 4140*A*B - 2025*I*B^2)*a^5/d^2)*(d*e^(3*I*d*x + 3*I*c) - d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))e^(-2*I*d*x - 2*I*c)/((46*I*A + 45*B)*a)...`

3.553.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)**(1/2),x)`

output `Timed out`

3.553.7 Maxima [F]

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{5/2}}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)/sqrt(cot(d*x + c)), x)`

3.553.8 Giac [F]

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^{5/2}}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^(5/2)/sqrt(cot(d*x + c)), x)`

3.553.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) i)^{5/2}}{\sqrt{\cot(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/cot(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^(5/2))/cot(c + d*x)^(1/2), x)`

3.554
$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

3.554.1 Optimal result 5237
 3.554.2 Mathematica [A] (verified) 5238
 3.554.3 Rubi [A] (verified) 5238
 3.554.4 Maple [B] (verified) 5242
 3.554.5 Fracas [B] (verification not implemented) 5243
 3.554.6 Sympy [F(-1)] 5244
 3.554.7 Maxima [F(-2)] 5244
 3.554.8 Giac [F] 5245
 3.554.9 Mupad [F(-1)] 5245

3.554.1 Optimal result

Integrand size = 38, antiderivative size = 211

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

$$= \frac{\left(\frac{1}{2} + \frac{i}{2}\right) (iA + B) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{\sqrt{ad}}$$

$$+ \frac{(A+iB) \cot^{\frac{3}{2}}(c+dx)}{d\sqrt{a+ia \tan(c+dx)}} + \frac{(7iA-9B)\sqrt{\cot(c+dx)}\sqrt{a+ia \tan(c+dx)}}{3ad}$$

$$- \frac{(5A+3iB) \cot^{\frac{3}{2}}(c+dx)\sqrt{a+ia \tan(c+dx)}}{3ad}$$

```
output (1/2+1/2*I)*(I*A+B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/a^(1/2)+(A+I*B)*cot(d*x+c)^(3/2)/d/(a+I*a*tan(d*x+c))^(1/2)-1/3*(5*A+3*I*B)*cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^(1/2)/a/d+1/3*(7*I*A-9*B)*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/a/d
```

3.554.
$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

3.554.2 Mathematica [A] (verified)

Time = 5.22 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.67

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

$$= \frac{3\sqrt{2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ia \tan(c+dx)}} - \frac{2(7A+9iB+(-2iA+6B) \cot(c+dx)+2A \cot^2(c+dx))}{\sqrt{a+ia \tan(c+dx)}}}{6d\sqrt{\cot(c+dx)}}$$

input `Integrate[(Cot[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `((-3*Sqrt[2]*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/Sqrt[I*a*Tan[c + d*x]] - (2*(7*A + (9*I)*B + ((-2*I)*A + 6*B)*Cot[c + d*x] + 2*A*Cot[c + d*x]^2))/Sqrt[a + I*a*Tan[c + d*x]])/(6*d*Sqrt[Cot[c + d*x]])`

3.554.3 Rubi [A] (verified)Time = 1.20 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.04, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4729, 3042, 4079, 27, 3042, 4081, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\cot(c+dx)^{5/2}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)\sqrt{ia \tan(c+dx)a+a}} dx$$

$$\downarrow \text{3042}$$

3.554. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

$$\begin{aligned}
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^{5/2} \sqrt{ia \tan(c+dx)+a}} dx \\
 & \qquad \qquad \qquad \downarrow \text{4079} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{\sqrt{ia \tan(c+dx)+a}(a(5A+3iB)-4a(iA-B) \tan(c+dx))}{2 \tan^{5/2}(c+dx)} dx}{a^2} + \frac{A+iB}{d \tan^{3/2}(c+dx) \sqrt{ia \tan(c+dx)+a}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{27} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{\sqrt{ia \tan(c+dx)+a}(a(5A+3iB)-4a(iA-B) \tan(c+dx))}{\tan^{5/2}(c+dx)} dx}{2a^2} + \frac{A+iB}{d \tan^{3/2}(c+dx) \sqrt{ia \tan(c+dx)+a}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{\sqrt{ia \tan(c+dx)+a}(a(5A+3iB)-4a(iA-B) \tan(c+dx))}{\tan(c+dx)^{5/2}} dx}{2a^2} + \frac{A+iB}{d \tan^{3/2}(c+dx) \sqrt{ia \tan(c+dx)+a}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{4081} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int -\frac{\sqrt{ia \tan(c+dx)+a}((7iA-9B)a^2+2(5A+3iB) \tan(c+dx)a^2)}{2 \tan^{3/2}(c+dx)} dx}{3a} - \frac{2a(5A+3iB) \sqrt{ia \tan(c+dx)+a}}{3d \tan^{3/2}(c+dx)} + \frac{A+iB}{d \tan^{3/2}(c+dx) \sqrt{ia \tan(c+dx)+a}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{27} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{\sqrt{ia \tan(c+dx)+a}((7iA-9B)a^2+2(5A+3iB) \tan(c+dx)a^2)}{\tan^{3/2}(c+dx)} dx}{3a} - \frac{2a(5A+3iB) \sqrt{ia \tan(c+dx)+a}}{3d \tan^{3/2}(c+dx)} + \frac{A+iB}{d \tan^{3/2}(c+dx) \sqrt{ia \tan(c+dx)+a}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{\sqrt{ia \tan(c+dx)+a}((7iA-9B)a^2+2(5A+3iB) \tan(c+dx)a^2)}{\tan(c+dx)^{3/2}} dx}{3a} - \frac{2a(5A+3iB) \sqrt{ia \tan(c+dx)+a}}{3d \tan^{3/2}(c+dx)} + \frac{A+iB}{d \tan^{3/2}(c+dx) \sqrt{ia \tan(c+dx)+a}} \right)
 \end{aligned}$$

3.554. $\int \frac{\cot^{5/2}(c+dx)(A+B \tan(c+dx))}{\sqrt{ia \tan(c+dx)+a}} dx$

$$\begin{array}{c} \downarrow 4081 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int \frac{3a^3(A-iB)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx - \frac{2a^2(-9B+7iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a(5A+3iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)}}{3a} \right. \\ \left. \frac{2a^2}{2a^2} \right) + \end{array}$$

$$\begin{array}{c} \downarrow 27 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{3a^2(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(-9B+7iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a(5A+3iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)}}{3a} \right. \\ \left. \frac{2a^2}{2a^2} \right) + \end{array}$$

$$\begin{array}{c} \downarrow 3042 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{3a^2(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(-9B+7iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a(5A+3iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)}}{3a} \right. \\ \left. \frac{2a^2}{2a^2} \right) + \end{array}$$

$$\begin{array}{c} \downarrow 4027 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{6ia^4(A-iB) \int \frac{1}{-2 \tan(c+dx)a^2 - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} - \frac{2a^2(-9B+7iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a(5A+3iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)}}{3a} \right. \\ \left. \frac{2a^2}{2a^2} \right) + \end{array}$$

$$\begin{array}{c} \downarrow 218 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(3-3i)a^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) - \frac{2a^2(-9B+7iA)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{2a(5A+3iB)\sqrt{a+ia \tan(c+dx)}}{3d \tan^{\frac{3}{2}}(c+dx)}}{d} \right. \\ \left. \frac{2a^2}{2a^2} \right) + \end{array}$$

3.554. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

input `Int[(Cot[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((A + I*B)/(d*Tan[c + d*x]^(3/2)*Sqrt[a + I*a*Tan[c + d*x]]) + ((-2*a*(5*A + (3*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - (((3 - 3*I)*a^(5/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a^2*((7*I)*A - 9*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a))/(2*a^2))`

3.554.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.554.
$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$


```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.554.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 747 vs. $2(171) = 342$.

Time = 0.62 (sec) , antiderivative size = 748, normalized size of antiderivative = 3.55

method	result
derivativedivides	$\left(\frac{1}{i \tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} \left(28iA \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{-ia} \tan(dx+c)^3 + 3iA \sqrt{2} \ln\left(\frac{2\sqrt{a(1+i \tan(dx+c))}}{1+i \tan(dx+c)}\right)\right)$
default	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} \left(28iA \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{-ia} \tan(dx+c)^3 + 3iA \sqrt{2} \ln\left(\frac{2\sqrt{a(1+i \tan(dx+c))}}{1+i \tan(dx+c)}\right)\right)$

```
input int(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x,method=_R
ETURNVERBOSE)
```

3.554.
$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

```
output 1/12/d*(1/tan(d*x+c))^(5/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)/a*(28*I*
A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3+3*I*A*2^
(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a
*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*a*tan(d*x+c)^4+3*B*2^(1/2)*ln((2*2^(1/2)*
(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(ta
n(d*x+c)+I))*a*tan(d*x+c)^4-36*B*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I
*a)^(1/2)*tan(d*x+c)^3-3*I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x
+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*a*tan(d*x+
c)^2+6*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))
)^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*a*tan(d*x+c)^3+36*A*(a*tan(d*x
+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2-6*I*B*2^(1/2)*ln((2*
2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-
I*a)/(tan(d*x+c)+I))*a*tan(d*x+c)^3-3*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)
*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))
*a*tan(d*x+c)^2+60*I*B*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)*
tan(d*x+c)^2+24*B*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^
(1/2)+8*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)/(a*tan(d*x+c
)*(1+I*tan(d*x+c)))^(1/2)/(-tan(d*x+c)+I)^2/(-I*a)^(1/2)
```

3.554.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 487 vs. 2(161) = 322.

Time = 0.25 (sec) , antiderivative size = 487, normalized size of antiderivative = 2.31

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx =$$

$$3\sqrt{2} \left(ade^{(3i dx+3i c)} - ade^{(i dx+i c)} \right) \sqrt{-\frac{iA^2+2AB-iB^2}{ad^2}} \log \left(-\frac{4 \left((A-iB)ae^{(i dx+i c)} + (ade^{(2i dx+2i c)} - ad) \sqrt{\frac{a}{e^{(2i dx+2i c)} + iA+B}} \right)}{iA+B} \right)$$

```
input integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, al
gorithm="fricas")
```

output
$$\begin{aligned} & -1/12*(3*\sqrt{2}*(a*d*e^{(3*I*d*x + 3*I*c)} - a*d*e^{(I*d*x + I*c)})*\sqrt{-(I*A^2 + 2*A*B - I*B^2)/(a*d^2)}*\log(-4*((A - I*B)*a*e^{(I*d*x + I*c)} + (a*d*e^{(2*I*d*x + 2*I*c)} - a*d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}*\sqrt{-(I*A^2 + 2*A*B - I*B^2)/(a*d^2)}))*e^{(-I*d*x - I*c)/(I*A + B)} - 3*\sqrt{2}*(a*d*e^{(3*I*d*x + 3*I*c)} - a*d*e^{(I*d*x + I*c)})*\sqrt{-(I*A^2 + 2*A*B - I*B^2)/(a*d^2)}*\log(-4*((A - I*B)*a*e^{(I*d*x + I*c)} - (a*d*e^{(2*I*d*x + 2*I*c)} - a*d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}*\sqrt{-(I*A^2 + 2*A*B - I*B^2)/(a*d^2)}))*e^{(-I*d*x - I*c)/(I*A + B)} \\ & + 2*\sqrt{2}*((-7*I*A + 15*B)*e^{(4*I*d*x + 4*I*c)} + 18*(I*A - B)*e^{(2*I*d*x + 2*I*c)} - 3*I*A + 3*B)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)))/(a*d*e^{(3*I*d*x + 3*I*c)} - a*d*e^{(I*d*x + I*c)}) \end{aligned}$$

3.554.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{5}{2}}(c + dx)(A + B \tan(c + dx))}{\sqrt{a + ia \tan(c + dx)}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/2),x)`

output Timed out

3.554.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^{\frac{5}{2}}(c + dx)(A + B \tan(c + dx))}{\sqrt{a + ia \tan(c + dx)}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.

3.554.
$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

3.554.8 Giac [F]

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{5}{2}}}{\sqrt{ia\tan(dx+c)+a}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(5/2)/sqrt(I*a*tan(d*x + c) + a), x)`

3.554.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{\cot(c+dx)^{5/2}(A+B\tan(c+dx))}{\sqrt{a+a\tan(c+dx)} \operatorname{li}} dx$$

input `int((cot(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2),x)`

output `int((cot(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2), x)`

3.555
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

3.555.1 Optimal result	5246
3.555.2 Mathematica [A] (verified)	5246
3.555.3 Rubi [A] (verified)	5247
3.555.4 Maple [B] (verified)	5250
3.555.5 Fricas [B] (verification not implemented)	5251
3.555.6 Sympy [F]	5252
3.555.7 Maxima [F(-2)]	5252
3.555.8 Giac [F]	5253
3.555.9 Mupad [F(-1)]	5253

3.555.1 Optimal result

Integrand size = 38, antiderivative size = 163

$$\begin{aligned} & \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx \\ &= \frac{\left(\frac{1}{2} + \frac{i}{2}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{\sqrt{ad}} \\ & \quad + \frac{(A+iB)\sqrt{\cot(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} - \frac{(3A+iB)\sqrt{\cot(c+dx)}\sqrt{a+ia \tan(c+dx)}}{ad} \end{aligned}$$

output `(1/2+1/2*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/a^(1/2)+(A+I*B)*cot(d*x+c)^(1/2)/d/(a+I*a*tan(d*x+c))^(1/2)-(3*A+I*B)*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/a/d`

3.555.2 Mathematica [A] (verified)

Time = 4.10 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.74

$$\begin{aligned} & \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx \\ &= \frac{\frac{\sqrt{2}(iA+B)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ia \tan(c+dx)}} + \frac{-6iA+2B-4A \cot(c+dx)}{\sqrt{a+ia \tan(c+dx)}}}{2d\sqrt{\cot(c+dx)}} \end{aligned}$$

3.555.
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

input `Integrate[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `((Sqrt[2]*(I*A + B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/Sqrt[I*a*Tan[c + d*x]] + ((-6*I)*A + 2*B - 4*A*Cot[c + d*x])/Sqrt[a + I*a*Tan[c + d*x]])/(2*d*Sqrt[Cot[c + d*x]])`

3.555.3 Rubi [A] (verified)

Time = 0.90 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4729, 3042, 4079, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cot(c+dx)^{3/2}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx \\
 & \quad \downarrow \text{4729} \\
 & \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx) \sqrt{i \tan(c+dx)} a+a} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^{3/2} \sqrt{i \tan(c+dx)} a+a} dx \\
 & \quad \downarrow \text{4079} \\
 & \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \left(\frac{\int \frac{\sqrt{i \tan(c+dx)} a+a(a(3A+iB)-2a(iA-B) \tan(c+dx))}{2 \tan^{\frac{3}{2}}(c+dx)} dx}{a^2} + \frac{A+iB}{d \sqrt{\tan(c+dx)} \sqrt{a+ia \tan(c+dx)}} \right) \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.555. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{\int\frac{\sqrt{i\tan(c+dx)a+a(a(3A+iB)-2a(iA-B)\tan(c+dx))}dx}{\tan^{\frac{3}{2}}(c+dx)}\frac{dx}{2a^2}+\frac{A+iB}{d\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{\int\frac{\sqrt{i\tan(c+dx)a+a(a(3A+iB)-2a(iA-B)\tan(c+dx))}dx}{\tan(c+dx)^{3/2}}\frac{dx}{2a^2}+\frac{A+iB}{d\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}\right)$$

↓ 4081

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2\int\frac{a^2(iA+B)\sqrt{i\tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}}dx-\frac{2a(3A+iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2}+\frac{A+iB}{d\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{a(B+iA)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx-\frac{2a(3A+iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2}+\frac{A+iB}{d\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{a(B+iA)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx-\frac{2a(3A+iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2}+\frac{A+iB}{d\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}\right)$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2ia^3(B+iA)\int\frac{1}{-2\tan(c+dx)a^2-ia}\frac{d\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}-\frac{2a(3A+iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2}+\frac{A+iB}{d\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}\right)$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(1-i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}-\frac{2a(3A+iB)\sqrt{a+ia\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2}+\frac{A+iB}{d\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}\right)$$

3.555. $\int\frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}}dx$

input `Int[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((A + I*B)/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + (((1 - I)*a^(3/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a*(3*A + I*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(2*a^2)`

3.555.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

3.555.
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$


```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.555.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 702 vs. 2(133) = 266.

Time = 0.60 (sec) , antiderivative size = 703, normalized size of antiderivative = 4.31

method	result
derivativedivides	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{3}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))}}{iB\sqrt{2} \ln\left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+3a \tan(dx+c)-ia}}{\tan(dx+c)+i}\right)} a t$
default	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{3}{2}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))}}{iB\sqrt{2} \ln\left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+3a \tan(dx+c)-ia}}{\tan(dx+c)+i}\right)} a t$

```
input int(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x,method=_R
ETURNVERBOSE)
```

3.555.
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

output

```

-1/4/d*(1/tan(d*x+c))^(3/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)/a*(I*B*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^3+2*I*A*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^2-A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*a*tan(d*x+c)^3+12*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2-I*B*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)+2*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*a*tan(d*x+c)^2+4*I*B*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2+A*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)-20*I*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)*tan(d*x+c)+4*B*tan(d*x+c)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)-8*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)/(-I*a)^(1/2)/(-tan(d*x+c)+I)^2

```

3.555.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 426 vs. $2(125) = 250$.

Time = 0.25 (sec) , antiderivative size = 426, normalized size of antiderivative = 2.61

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx =$$

$$\left(\sqrt{2}ad \sqrt{-\frac{-iA^2-2AB+iB^2}{ad^2}} e^{(idx+ic)} \log \left(-\frac{4 \left((A-iB)ae^{(idx+ic)} + (iade^{(2idx+2ic)} - iad) \sqrt{\frac{a}{e^{(2idx+2ic)}+1}} \sqrt{\frac{ie^{(2idx+2ic)}+i}{e^{(2idx+2ic)}-1}}}{iA+B} \right)} \right)$$

input

```

integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="fricas")

```

3.555. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

```
output -1/4*(sqrt(2)*a*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2))*e^(I*d*x + I*c)*
log(-4*((A - I*B)*a*e^(I*d*x + I*c) + (I*a*d*e^(2*I*d*x + 2*I*c) - I*a*d)*
sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I
*d*x + 2*I*c) - 1))*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2)))*e^(-I*d*x - I
*c)/(I*A + B)) - sqrt(2)*a*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2))*e^(I*
d*x + I*c)*log(-4*((A - I*B)*a*e^(I*d*x + I*c) + (-I*a*d*e^(2*I*d*x + 2*I*
c) + I*a*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c)
+ I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2)))*e
^(-I*d*x - I*c)/(I*A + B)) + 2*sqrt(2)*((5*A + I*B)*e^(2*I*d*x + 2*I*c) -
A - I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I
)/(e^(2*I*d*x + 2*I*c) - 1))*e^(-I*d*x - I*c)/(a*d)
```

3.555.6 Sympy [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \int \frac{(A+B \tan(c+dx)) \cot^{\frac{3}{2}}(c+dx)}{\sqrt{ia(\tan(c+dx)-i)}} dx$$

```
input integrate(cot(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/2),x)
```

```
output Integral((A + B*tan(c + d*x))*cot(c + d*x)**(3/2)/sqrt(I*a*(tan(c + d*x) -
I)), x)
```

3.555.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \text{Exception raised: RuntimeError}$$

```
input integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, al
gorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un
defined.
```

3.555. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$

3.555.8 Giac [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{\sqrt{ia\tan(dx+c)+a}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(3/2)/sqrt(I*a*tan(d*x + c) + a), x)`

3.555.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{\cot(c+dx)^{3/2}(A+B\tan(c+dx))}{\sqrt{a+a\tan(c+dx)} \operatorname{li}} dx$$

input `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2),x)`

output `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2), x)`

3.556
$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

3.556.1 Optimal result	5254
3.556.2 Mathematica [A] (verified)	5254
3.556.3 Rubi [A] (verified)	5255
3.556.4 Maple [B] (verified)	5257
3.556.5 Fracas [B] (verification not implemented)	5258
3.556.6 Sympy [F]	5259
3.556.7 Maxima [F(-2)]	5259
3.556.8 Giac [F]	5260
3.556.9 Mupad [F(-1)]	5260

3.556.1 Optimal result

Integrand size = 38, antiderivative size = 119

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

$$= \frac{\left(\frac{1}{2} - \frac{i}{2}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{\sqrt{ad}}$$

$$+ \frac{A + iB}{d\sqrt{\cot(c+dx)}\sqrt{a+ia \tan(c+dx)}}$$

output `(1/2-1/2*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/a^(1/2)+(A+I*B)/d/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2)`

3.556.2 Mathematica [A] (verified)

Time = 2.84 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \frac{\sqrt{2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ia \tan(c+dx)}} + \frac{2(A+iB)}{2d\sqrt{\cot(c+dx)}}$$

input `Integrate[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]], x]`

3.556.
$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

output $((\text{Sqrt}[2]*(A - I*B)*\text{ArcTanh}[(\text{Sqrt}[2]*\text{Sqrt}[I*a*\text{Tan}[c + d*x]])/\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])/\text{Sqrt}[I*a*\text{Tan}[c + d*x]] + (2*(A + I*B))/\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])/(2*d*\text{Sqrt}[\text{Cot}[c + d*x]])$

3.556.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.01, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {3042, 4729, 3042, 4079, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx \\ & \quad \downarrow \text{4729} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} dx \\ & \quad \downarrow \text{4079} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(A-iB)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx}{a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \right) \\ & \quad \downarrow \text{27} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \right) \\ & \quad \downarrow \text{3042} \end{aligned}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A-iB)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx}{2a}+\frac{(A+iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}\right)$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A+iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}-\frac{ia(A-iB)\int\frac{1}{-\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a}-ia}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{d}\right)$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(\frac{1}{2}-\frac{i}{2})(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{\sqrt{ad}}+\frac{(A+iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}\right)$$

input `Int[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[a + I*a*Tan[c + d*x]],x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((1/2 - I/2)*(A - I*B)*ArcTanh[(((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/(Sqrt[a]*d) + ((A + I*B)*Sqrt[Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]]))`

3.556.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4027 Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_)
+ (f_)*(x_)], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4079 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.556.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 640 vs. 2(95) = 190.

Time = 0.69 (sec) , antiderivative size = 641, normalized size of antiderivative = 5.39

method	result
derivativedivides	$-\frac{\sqrt{\frac{1}{\tan(dx+c)}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))}}{iA\sqrt{2} \ln\left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+3a \tan(dx+c)-ia}}{\tan(dx+c)+i}\right)} a \tan(dx+c)$
default	$-\frac{\sqrt{\frac{1}{\tan(dx+c)}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))}}{iA\sqrt{2} \ln\left(\frac{2\sqrt{2}\sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+3a \tan(dx+c)-ia}}{\tan(dx+c)+i}\right)} a \tan(dx+c)$

```
input int(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x,method=_R
ETURNVERBOSE)
```

$$3.556. \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$


```
output -1/4/d*(1/tan(d*x+c))^(1/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)/a*(I*A*2
^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*
a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*a*tan(d*x+c)^2-2*I*B*2^(1/2)*ln((2*2^(1/
2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/
(tan(d*x+c)+I))*a*tan(d*x+c)+B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d
*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*a*tan(d*
x+c)^2-I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c
)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*a+2*A*ln((2*2^(1/2)*(-I*a)^(
1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)
+I))*2^(1/2)*a*tan(d*x+c)+4*I*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*
a)^(1/2)*tan(d*x+c)-B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+
I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*a+4*I*B*(a*tan(d*
x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)-4*B*tan(d*x+c)*(a*tan(d*x+c)*(1+
I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)+4*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2
)*(-I*a)^(1/2)/(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)/(-tan(d*x+c)+I)^2/(-
I*a)^(1/2)
```

3.556.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 423 vs. 2(89) = 178.

Time = 0.25 (sec) , antiderivative size = 423, normalized size of antiderivative = 3.55

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx$$

$$= \left(\sqrt{2}ad \sqrt{-\frac{iA^2+2AB-iB^2}{ad^2}} e^{(i dx+ic)} \log \left(-\frac{4 \left((A-iB)ae^{(i dx+ic)}+(ade^{(2i dx+2i c)}-ad) \sqrt{\frac{a}{e^{(2i dx+2i c)}+1}} \sqrt{\frac{ie^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}-1}} \sqrt{-iA}}{iA+B} \right)} \right) \right)$$

```
input integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, al
gorithm="fricas")
```

output $\frac{1}{4}(\sqrt{2})a*d*\sqrt{-(I*A^2 + 2*A*B - I*B^2)/(a*d^2)}*e^{(I*d*x + I*c)}*\log(-4*((A - I*B)*a*e^{(I*d*x + I*c)} + (a*d*e^{(2*I*d*x + 2*I*c)} - a*d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}*\sqrt{-(I*A^2 + 2*A*B - I*B^2)/(a*d^2)}))e^{(-I*d*x - I*c)}/(I*A + B) - \sqrt{2})a*d*\sqrt{-(I*A^2 + 2*A*B - I*B^2)/(a*d^2)}*e^{(I*d*x + I*c)}*\log(-4*((A - I*B)*a*e^{(I*d*x + I*c)} - (a*d*e^{(2*I*d*x + 2*I*c)} - a*d)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}*\sqrt{-(I*A^2 + 2*A*B - I*B^2)/(a*d^2)}))e^{(-I*d*x - I*c)}/(I*A + B) - 2*\sqrt{2}*((I*A - B)*e^{(2*I*d*x + 2*I*c)} - I*A + B)*\sqrt{a/(e^{(2*I*d*x + 2*I*c)} + 1)}*\sqrt{(I*e^{(2*I*d*x + 2*I*c)} + I)/(e^{(2*I*d*x + 2*I*c)} - 1)}))e^{(-I*d*x - I*c)}/(a*d)$

3.556.6 Sympy [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \int \frac{(A+B \tan(c+dx)) \sqrt{\cot(c+dx)}}{\sqrt{ia}(\tan(c+dx)-i)} dx$$

input `integrate(cot(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(cot(c + d*x))/sqrt(I*a*(tan(c + d*x) - I)), x)`

3.556.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+ia \tan(c+dx)}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.556.8 Giac [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\sqrt{\cot(dx+c)}}{\sqrt{ia\tan(dx+c)+a}} dx$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(cot(d*x + c))/sqrt(I*a*tan(d*x + c) + a), x)`

3.556.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+ia\tan(c+dx)}} dx = \int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+a\tan(c+dx)} \operatorname{li}} dx$$

input `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2),x)`

output `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(1/2), x)`

3.557 $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}\sqrt{a+ia \tan(c+dx)}} dx$

3.557.1 Optimal result 5261
 3.557.2 Mathematica [A] (verified) 5262
 3.557.3 Rubi [A] (verified) 5262
 3.557.4 Maple [B] (verified) 5266
 3.557.5 Fricas [B] (verification not implemented) 5267
 3.557.6 Sympy [F] 5268
 3.557.7 Maxima [F(-2)] 5269
 3.557.8 Giac [F] 5269
 3.557.9 Mupad [F(-1)] 5269

3.557.1 Optimal result

Integrand size = 38, antiderivative size = 196

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}\sqrt{a + ia \tan(c + dx)}} dx$$

$$= -\frac{2\sqrt{-1}B \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{\sqrt{ad}}$$

$$- \frac{\left(\frac{1}{2} + \frac{i}{2}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{\sqrt{ad}}$$

$$+ \frac{iA - B}{d\sqrt{\cot(c + dx)}\sqrt{a + ia \tan(c + dx)}}$$

output

```
-2*(-1)^(1/4)*B*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/a^(1/2)-(1/2+1/2*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/a^(1/2)+(I*A-B)/d/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2)
```

3.557.2 Mathematica [A] (verified)

Time = 2.83 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.97

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx$$

$$= \frac{\left(\frac{1}{2} + \frac{i}{2}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left((-2 + 2i)(-1)^{3/4} \sqrt{a} \operatorname{Barcsinh}\left(\sqrt[4]{-1} \sqrt{\tan(c + dx)}\right)\right) \sqrt{1 + i \tan(c + dx)}}{\sqrt{ad} \sqrt{a + ia \tan(c + dx)}}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]),x]`

output `((1/2 + I/2)*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2 + 2*I)*(-1)^(3/4)*Sqrt[a]*B*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]]*Sqrt[1 + I*Tan[c + d*x]] + (1 + I)*Sqrt[a]*(A + I*B)*Sqrt[Tan[c + d*x]] - (A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]]*Sqrt[a + I*a*Tan[c + d*x]])/(Sqrt[a]*d*Sqrt[a + I*a*Tan[c + d*x]])`

3.557.3 Rubi [A] (verified)

Time = 1.11 (sec) , antiderivative size = 181, normalized size of antiderivative = 0.92, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.342$, Rules used = {3042, 4729, 3042, 4078, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx$$

↓ 4729

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\sqrt{\tan(c + dx)} (A + B \tan(c + dx))}{\sqrt{i \tan(c + dx) a + a}} dx$$

↓ 3042

3.557. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)} \sqrt{a+ia \tan(c+dx)}} dx$

$$\begin{aligned}
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{i \tan(c+dx)a+a}} dx \\
 & \quad \downarrow \text{4078} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA-B)+2iaB \tan(c+dx))}{2\sqrt{\tan(c+dx)}} dx}{a^2} \right) \\
 & \quad \downarrow \text{27} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA-B)+2iaB \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{2a^2} \right) \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} - \frac{\int \frac{\sqrt{i \tan(c+dx)a+a}(a(iA-B)+2iaB \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{2a^2} \right) \\
 & \quad \downarrow \text{4084} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} - \frac{a(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - 2B \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a^2} \right) \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} - \frac{a(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - 2B \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a^2} \right) \\
 & \quad \downarrow \text{4027} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} - \frac{2ia^3(B+iA) \int \frac{1}{\frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} - 2B \int \frac{(a-ia \tan(c+dx))\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{2a^2} \right) \\
 & \quad \downarrow \text{218}
 \end{aligned}$$

3.557. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}\sqrt{a+ia \tan(c+dx)}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}-\frac{(1-i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}-\frac{2B\int\frac{(a-ia\tan(c+dx))^{3/2}}{\sqrt{\tan(c+dx)}}dx}{2a^2}\right)$$

↓ 4082

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}-\frac{(1-i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}-\frac{2a^2B\int\frac{1}{\sqrt{\tan(c+dx)}}dx}{2a^2}\right)$$

↓ 65

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}-\frac{(1-i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}-\frac{4a^2B\int\frac{1}{1-\frac{ia}{i\tan(c+dx)}}dx}{2a^2}\right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}-\frac{(1-i)a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}+\frac{4\sqrt[4]{-1}a^{3/2}B}{2a^2}\right)$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-1/2*((-1)^(1/4)*a^(3/2)*B*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d + ((1 - I)*a^(3/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d)/a^2 + ((I*A - B)*Sqrt[Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]])`

3.557.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

rule 4729 `Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.557.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 606 vs. $2(155) = 310$.

Time = 0.57 (sec) , antiderivative size = 607, normalized size of antiderivative = 3.10

method	result
derivativedivides	$-\frac{\sqrt{a(1+i \tan(dx+c))} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \left(iB\sqrt{ia} \ln \left(-\frac{-2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+ia-3a \tan(dx+c)}}{\tan(dx+c)+i} \right)}{\right)}$
default	$-\frac{\sqrt{a(1+i \tan(dx+c))} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \left(iB\sqrt{ia} \ln \left(-\frac{-2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+ia-3a \tan(dx+c)}}{\tan(dx+c)+i} \right)}{\right)}$

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2),x,method=_R ETURNVERBOSE)`

$$3.557. \int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)} \sqrt{a+ia \tan(c+dx)}} dx$$

```
output -1/4/d*(a*(1+I*tan(d*x+c)))^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I
*B*(I*a)^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))
)^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)+I*A*(I*a)
^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+
I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a-A*(I*a)^(1/2)*ln(-(-2*2^(1/2
)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(
tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)+4*I*B*(I*a)^(1/2)*(-I*a)^(1/2)*(a*tan(
d*x+c)*(1+I*tan(d*x+c)))^(1/2)-4*I*B*ln(1/2*(2*I*a*tan(d*x+c)+2*(a*tan(d*x
+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-I*a)^(1/2)*a+B*(
I*a)^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1
/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a+4*B*ln(1/2*(2*I*a*tan(d*
x+c)+2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(1/2)+a)/(I*a)^(1/2))*(-
I*a)^(1/2)*a*tan(d*x+c)+4*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(I*a)^(
1/2)*(-I*a)^(1/2))/(1/tan(d*x+c))^(1/2)/a^2/(1+I*tan(d*x+c))/tan(d*x+c)/(I
*a)^(1/2)/(-I*a)^(1/2)/(-tan(d*x+c)+I)
```

3.557.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 736 vs. 2(148) = 296.

Time = 0.26 (sec) , antiderivative size = 736, normalized size of antiderivative = 3.76

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx$$

$$= \left(\sqrt{2} ad \sqrt{-\frac{-iA^2 - 2AB + iB^2}{ad^2}} e^{(i dx + i c)} \log \left(-\frac{4 \left((A - iB) a e^{(i dx + i c)} + (i a d e^{(2i dx + 2i c)} - i ad) \sqrt{\frac{a}{e^{(2i dx + 2i c)} + 1}} \sqrt{\frac{i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} - 1}} \right)}{i A + B} \right) \right)$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2),x, al
gorithm="fracas")
```

output

```

1/4*(sqrt(2)*a*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2))*e^(I*d*x + I*c)*log(-4*((A - I*B)*a*e^(I*d*x + I*c) + (I*a*d*e^(2*I*d*x + 2*I*c) - I*a*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2)))*e^(-I*d*x - I*c)/(I*A + B)) - sqrt(2)*a*d*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2))*e^(I*d*x + I*c)*log(-4*((A - I*B)*a*e^(I*d*x + I*c) + (-I*a*d*e^(2*I*d*x + 2*I*c) + I*a*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt(-(-I*A^2 - 2*A*B + I*B^2)/(a*d^2)))*e^(-I*d*x - I*c)/(I*A + B)) + a*d*sqrt(-4*I*B^2/(a*d^2))*e^(I*d*x + I*c)*log(-16*(3*B*a^2*e^(2*I*d*x + 2*I*c) - B*a^2 + sqrt(2)*(a^2*d*e^(3*I*d*x + 3*I*c) - a^2*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt(-4*I*B^2/(a*d^2)))*e^(-2*I*d*x - 2*I*c)/B) - a*d*sqrt(-4*I*B^2/(a*d^2))*e^(I*d*x + I*c)*log(-16*(3*B*a^2*e^(2*I*d*x + 2*I*c) - B*a^2 - sqrt(2)*(a^2*d*e^(3*I*d*x + 3*I*c) - a^2*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt(-4*I*B^2/(a*d^2)))*e^(-2*I*d*x - 2*I*c)/B) + 2*sqrt(2)*((A + I*B)*e^(2*I*d*x + 2*I*c) - A - I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-I*d*x - I*c)/(a*d)

```

3.557.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{ia (\tan(c + dx) - i)} \sqrt{\cot(c + dx)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(1/2)/(a+I*a*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))/(sqrt(I*a*(tan(c + d*x) - I))*sqrt(cot(c + d*x))), x)`

3.557.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.557.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx = \int \frac{B \tan(dx + c) + A}{\sqrt{ia \tan(dx + c) + a} \sqrt{\cot(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/(sqrt(I*a*tan(d*x + c) + a)*sqrt(cot(d*x + c))), x)`

3.557.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + a \tan(c + dx)} li} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(1/2)), x)`

3.557. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)} \sqrt{a+ia \tan(c+dx)}} dx$

3.558 $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

3.558.1 Optimal result 5270
 3.558.2 Mathematica [A] (verified) 5271
 3.558.3 Rubi [A] (verified) 5271
 3.558.4 Maple [B] (verified) 5275
 3.558.5 Fricas [B] (verification not implemented) 5276
 3.558.6 Sympy [F] 5277
 3.558.7 Maxima [F(-2)] 5277
 3.558.8 Giac [F] 5278
 3.558.9 Mupad [F(-1)] 5278

3.558.1 Optimal result

Integrand size = 38, antiderivative size = 214

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{(\frac{1}{4} + \frac{i}{4})(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{a^{3/2}d} + \frac{(A+iB)\sqrt{\cot(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} + \frac{(11A+5iB)\sqrt{\cot(c+dx)}}{6ad\sqrt{a+ia \tan(c+dx)}} - \frac{(25A+7iB)\sqrt{\cot(c+dx)}\sqrt{a+ia \tan(c+dx)}}{6a^2d}$$

output

```
(1/4+1/4*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/a^(3/2)/d+1/6*(11*A+5*I*B)*cot(d*x+c)^(1/2)/a/d/(a+I*a*tan(d*x+c))^(1/2)-1/6*(25*A+7*I*B)*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/a^2/d+1/3*(A+I*B)*cot(d*x+c)^(1/2)/d/(a+I*a*tan(d*x+c))^(3/2)
```

3.558.2 Mathematica [A] (verified)

Time = 7.15 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.92

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{\frac{3}{2}}} dx = \frac{i\left(3\sqrt{2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)(i+\cot(c+dx))\sqrt{a+ia\tan(c+dx)}\right)}{12d\cot^{\frac{5}{2}}(c+dx)(ia\tan(c+dx))^{\frac{3}{2}}(-i-\cot(c+dx))}$$

input `Integrate[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`

output `((I/12)*(3*Sqrt[2]*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*(I + Cot[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]] + (2*a*((-39*I)*A + 9*B - 12*A*Cot[c + d*x] + (25*A + (7*I)*B)*Tan[c + d*x]))/Sqrt[I*a*Tan[c + d*x]])/(d*Cot[c + d*x]^(5/2)*(I*a*Tan[c + d*x])^(3/2)*(-I + Tan[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]])`

3.558.3 Rubi [A] (verified)Time = 1.25 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.03, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4729, 3042, 4079, 27, 3042, 4079, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{\frac{3}{2}}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\cot(c+dx)^{\frac{3}{2}}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{\frac{3}{2}}} dx \\ & \quad \downarrow \text{4729} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(i\tan(c+dx)a+a)^{\frac{3}{2}}} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B\tan(c+dx)}{\tan(c+dx)^{\frac{3}{2}}(i\tan(c+dx)a+a)^{\frac{3}{2}}} dx \end{aligned}$$

3.558. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{\frac{3}{2}}} dx$

$$\begin{array}{c}
\downarrow 4079 \\
\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(7A+iB)-4a(iA-B)\tan(c+dx)}{2\tan^{\frac{3}{2}}(c+dx)\sqrt{i\tan(c+dx)a+a}} dx}{3a^2} + \frac{A+iB}{3d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{3/2}} \right) \\
\downarrow 27 \\
\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(7A+iB)-4a(iA-B)\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} + \frac{A+iB}{3d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{3/2}} \right) \\
\downarrow 3042 \\
\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(7A+iB)-4a(iA-B)\tan(c+dx)}{\tan(c+dx)^{3/2}\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} + \frac{A+iB}{3d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{3/2}} \right) \\
\downarrow 4079 \\
\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{\sqrt{i\tan(c+dx)a+a}(a^2(25A+7iB)-2a^2(11iA-5B)\tan(c+dx)) dx}{2\tan^{\frac{3}{2}}(c+dx)a^2} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}}{6a^2} + \frac{A+iB}{3d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{3/2}} \right) \\
\downarrow 27 \\
\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{\sqrt{i\tan(c+dx)a+a}(a^2(25A+7iB)-2a^2(11iA-5B)\tan(c+dx)) dx}{\tan^{\frac{3}{2}}(c+dx)2a^2} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}}{6a^2} + \frac{A+iB}{3d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{3/2}} \right) \\
\downarrow 3042 \\
\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{\sqrt{i\tan(c+dx)a+a}(a^2(25A+7iB)-2a^2(11iA-5B)\tan(c+dx)) dx}{\tan(c+dx)^{3/2}2a^2} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia\tan(c+dx)}}}{6a^2} + \frac{A+iB}{3d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{3/2}} \right) \\
\downarrow 4081
\end{array}$$

3.558. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int \frac{3a^3(iA+B)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx - \frac{2a^2(25A+7iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} \right) + \dots$$

27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{3a^2(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(25A+7iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} \right) + \dots$$

3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{3a^2(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^2(25A+7iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} \right) + \dots$$

4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{6ia^4(B+iA) \int \frac{1}{i \tan(c+dx)a+a} dx - \frac{d \sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} - \frac{2a^2(25A+7iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} \right) + \dots$$

218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(3-3i)a^{5/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) - \frac{2a^2(25A+7iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{a(11A+5iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} \right) + \dots$$

```
input Int[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2), x]
```


output $\text{Sqrt}[\text{Cot}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]*((A + I*B)/(3*d*\text{Sqrt}[\text{Tan}[c + d*x]]*(a + I*a*\text{Tan}[c + d*x])^{3/2}) + ((a*(11*A + (5*I)*B))/(d*\text{Sqrt}[\text{Tan}[c + d*x]]*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]) + (((3 - 3*I)*a^{5/2}*(I*A + B)*\text{ArcTan}h[((1 + I)*\text{Sqrt}[a]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]])/d - (2*a^2*(2*5*A + (7*I)*B)*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])/(d*\text{Sqrt}[\text{Tan}[c + d*x]]))/(2*a^2)/(6*a^2))$

3.558.3.1 Defintions of rubi rules used

rule 27 $\text{Int}[(a_)*(F_x_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_)] /; \text{FreeQ}[b, x]$

rule 218 $\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4027 $\text{Int}[\text{Sqrt}[(a_) + (b_)*\text{tan}[(e_) + (f_)*(x_)]]/\text{Sqrt}[(c_) + (d_)*\text{tan}[(e_) + (f_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[-2*a*(b/f) \text{ Subst}[\text{Int}[1/(a*c - b*d - 2*a^2*x^2), x], x, \text{Sqrt}[c + d*\text{Tan}[e + f*x]]/\text{Sqrt}[a + b*\text{Tan}[e + f*x]]], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0]$

rule 4079 $\text{Int}[(a_) + (b_)*\text{tan}[(e_) + (f_)*(x_)])^{(m_)}*((A_) + (B_)*\text{tan}[(e_) + (f_)*(x_)])^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a*A + b*B)*(a + b*\text{Tan}[e + f*x])^m*((c + d*\text{Tan}[e + f*x])^{(n+1)})/(2*f*m*(b*c - a*d)), x] + \text{Simp}[1/(2*a*m*(b*c - a*d)) \text{ Int}[(a + b*\text{Tan}[e + f*x])^{(m+1)}*(c + d*\text{Tan}[e + f*x])^n*\text{Simp}[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*\text{Tan}[e + f*x], x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, n\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{LtQ}[m, 0] \ \&\& \ !\text{GtQ}[n, 0]$

```
rule 4081 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n +
1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e +
f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*
m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; Fr
eeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2,
0] && LtQ[n, -1]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.558.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 938 vs. $2(172) = 344$.

Time = 0.61 (sec) , antiderivative size = 939, normalized size of antiderivative = 4.39

method	result	size
derivativedivides	Expression too large to display	939
default	Expression too large to display	939

```
input int(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x,method=_R
ETURNVERBOSE)
```

```

output 1/24/d*(1/tan(d*x+c))^(3/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)*(3*I*B*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^4+9*I*A*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^3-3*A*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^4-9*I*B*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^2+28*I*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3+9*B*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3-3*I*A*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)-256*I*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^2+9*A*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^2+100*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)^3-36*I*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)-3*B*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)+64*B*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+48*I*A*(-I*a)^(...

```

3.558.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 465 vs. $2(162) = 324$.

Time = 0.26 (sec) , antiderivative size = 465, normalized size of antiderivative = 2.17

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx =$$

$$\left(3\sqrt{\frac{1}{2}}a^2d\sqrt{\frac{iA^2+2AB-iB^2}{a^3d^2}}e^{(3i dx+3i c)} \log \left(-\frac{4\left(\sqrt{2}\sqrt{\frac{1}{2}}(ia^2de^{(2i dx+2i c)}-ia^2d)\sqrt{\frac{a}{e^{(2i dx+2i c)}+1}}\sqrt{\frac{ie^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}-1}}\sqrt{\frac{iA^2+2AB-iB^2}{a^3d^2}}\right)}{iA+B} \right) \right)$$

```

input integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="fricas")

```

3.558. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx$

output `-1/12*(3*sqrt(1/2)*a^2*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(I*a^2*d*e^(2*I*d*x + 2*I*c) - I*a^2*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^3*d^2)) + (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 3*sqrt(1/2)*a^2*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(-I*a^2*d*e^(2*I*d*x + 2*I*c) + I*a^2*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^3*d^2)) + (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) + sqrt(2)*(2*(19*A + 4*I*B)*e^(4*I*d*x + 4*I*c) - (13*A + 7*I*B)*e^(2*I*d*x + 2*I*c) - A - I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*e^(-3*I*d*x - 3*I*c)/(a^2*d)`

3.558.6 Sympy [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \int \frac{(A+B \tan(c+dx)) \cot^{\frac{3}{2}}(c+dx)}{(ia(\tan(c+dx)-i))^{\frac{3}{2}}} dx$$

input `integrate(cot(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**(3/2)/(I*a*(tan(c + d*x) - I))**(3/2), x)`

3.558.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is not defined.`

3.558. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

3.558.8 Giac [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{(ia\tan(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(3/2)/(I*a*tan(d*x + c) + a)^(3/2), x)`

3.558.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \int \frac{\cot(c+dx)^{3/2}(A+B\tan(c+dx))}{(a+a\tan(c+dx)li)^{3/2}} dx$$

input `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(3/2),x)`

output `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(3/2), x)`

3.559
$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$$

3.559.1 Optimal result 5279
 3.559.2 Mathematica [A] (verified) 5279
 3.559.3 Rubi [A] (verified) 5280
 3.559.4 Maple [B] (verified) 5283
 3.559.5 Fricas [B] (verification not implemented) 5284
 3.559.6 Sympy [F] 5284
 3.559.7 Maxima [F(-2)] 5285
 3.559.8 Giac [F] 5285
 3.559.9 Mupad [F(-1)] 5285

3.559.1 Optimal result

Integrand size = 38, antiderivative size = 168

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{\left(\frac{1}{4} - \frac{i}{4}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{a^{3/2}d} + \frac{A + iB}{3d\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{3/2}} + \frac{7A + iB}{6ad\sqrt{\cot(c+dx)}\sqrt{a+ia \tan(c+dx)}}$$

output $(1/4-1/4*I)*(A-I*B)*\operatorname{arctanh}((1+I)*a^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+I*a*\tan(d*x+c)))^{(1/2)}*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/a^{(3/2)}/d+1/6*(7*A+I*B)/a/d/\cot(d*x+c)^{(1/2)}/(a+I*a*\tan(d*x+c))^{(1/2)}+1/3*(A+I*B)/d/\cot(d*x+c)^{(1/2)}/(a+I*a*\tan(d*x+c))^{(3/2)}$

3.559.2 Mathematica [A] (verified)

Time = 5.06 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.87

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{3\sqrt{2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ia \tan(c+dx)}} + \frac{6(-3iA+B)+2(7A+iB) \tan(c+dx)}{(-i+\tan(c+dx))\sqrt{a+ia \tan(c+dx)}} \frac{1}{12ad\sqrt{\cot(c+dx)}}$$

input $\operatorname{Integrate}[(\operatorname{Sqrt}[\operatorname{Cot}[c+d*x]])*(A+B*\operatorname{Tan}[c+d*x]))/(a+I*a*\operatorname{Tan}[c+d*x])^{(3/2)},x]$

3.559.
$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$$

output $((3*\text{Sqrt}[2]*(A - I*B)*\text{ArcTanh}[(\text{Sqrt}[2]*\text{Sqrt}[I*a*\text{Tan}[c + d*x]])/\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]])/\text{Sqrt}[I*a*\text{Tan}[c + d*x]] + (6*((-3*I)*A + B) + 2*(7*A + I*B)*\text{Tan}[c + d*x])/((-I + \text{Tan}[c + d*x])*\text{Sqrt}[a + I*a*\text{Tan}[c + d*x]]))/((12*a*d*\text{Sqrt}[\text{Cot}[c + d*x]]))$

3.559.3 Rubi [A] (verified)

Time = 0.89 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.02, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4729, 3042, 4079, 27, 3042, 4079, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$$

↓ 3042

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$$

↓ 4729

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^{3/2}} dx$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^{3/2}} dx$$

↓ 4079

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(5A-iB)-2a(iA-B) \tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} dx}{3a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(5A-iB)-2a(iA-B) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i \tan(c+dx)a+a}} dx}{6a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} \right)$$

↓ 3042

$$\begin{aligned}
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(5A-iB)-2a(iA-B)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} \right) \\
& \quad \downarrow \text{4079} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{3a^2(A-iB)\sqrt{i\tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx}{6a^2} + \frac{a(7A+iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\frac{3}{2}(A-iB)\int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + \frac{a(7A+iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}}{6a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\frac{3}{2}(A-iB)\int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + \frac{a(7A+iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}}{6a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} \right) \\
& \quad \downarrow \text{4027} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\frac{a(7A+iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{3ia^2(A-iB)\int \frac{1}{\frac{-2\tan(c+dx)a^2}{i\tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{6a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} \right) \\
& \quad \downarrow \text{218} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\left(\frac{3}{2}-\frac{3i}{2}\right)\sqrt{a}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} + \frac{a(7A+iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} \right)
\end{aligned}$$

input `Int[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(3/2),x]`


```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((A + I*B)*Sqrt[Tan[c + d*x]])/(3*d
*(a + I*a*Tan[c + d*x])^(3/2)) + (((3/2 - (3*I)/2)*Sqrt[a]*(A - I*B)*ArcTa
nh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d + (
a*(7*A + I*B)*Sqrt[Tan[c + d*x]]/(d*Sqrt[a + I*a*Tan[c + d*x]]))/(6*a^2))
```

3.559.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4027 Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4079 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.559.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 875 vs. $2(134) = 268$.

Time = 0.57 (sec) , antiderivative size = 876, normalized size of antiderivative = 5.21

method	result
derivativedivides	$\sqrt{\frac{1}{\tan(dx+c)}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} \left(3iA\sqrt{2} \ln \left(-\frac{-2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+ia-3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \right) a t$
default	$\sqrt{\frac{1}{\tan(dx+c)}} \tan(dx+c) \sqrt{a(1+i \tan(dx+c))} \left(3iA\sqrt{2} \ln \left(-\frac{-2\sqrt{2} \sqrt{-ia} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))+ia-3a \tan(dx+c)}}{\tan(dx+c)+i} \right) \right) a t$

input `int(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x,method=_R
ETURNVERBOSE)`

output $\frac{1}{24}d*(1/\tan(d*x+c))^{(1/2)}*\tan(d*x+c)*(a*(1+I*\tan(d*x+c)))^{(1/2)}*(3*I*A*1$
 $\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+I*a-3*a*$
 $\tan(d*x+c))/(\tan(d*x+c)+I))*2^{(1/2)}*a*\tan(d*x+c)^3-9*I*B*\ln(-(-2*2^{(1/2)}*(-$
 $-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+I*a-3*a*\tan(d*x+c))/(\tan$
 $(d*x+c)+I))*2^{(1/2)}*a*\tan(d*x+c)^2+3*B*2^{(1/2)}*\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}$
 $)*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+I*a-3*a*\tan(d*x+c))/(\tan(d*x+c)+I)$
 $)*a*\tan(d*x+c)^3-9*I*A*\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan$
 $(d*x+c)))^{(1/2)}+I*a-3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*2^{(1/2)}*a*\tan(d*x+c)+2$
 $8*I*A*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*(-I*a)^{(1/2)}*\tan(d*x+c)^2+9*A*$
 $\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+I*a-3*a*$
 $*\tan(d*x+c))/(\tan(d*x+c)+I))*2^{(1/2)}*a*\tan(d*x+c)^2+3*I*B*\ln(-(-2*2^{(1/2)}*(-$
 $-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+I*a-3*a*\tan(d*x+c))/(\tan$
 $(d*x+c)+I))*2^{(1/2)}*a+16*I*B*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*(-I*a)$
 $^{(1/2)}*\tan(d*x+c)-9*B*2^{(1/2)}*\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*($
 $1+I*\tan(d*x+c)))^{(1/2)}+I*a-3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*a*\tan(d*x+c)-4*$
 $B*(-I*a)^{(1/2)}*\tan(d*x+c)^2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}-36*I*A*($
 $a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*(-I*a)^{(1/2)}-3*A*\ln(-(-2*2^{(1/2)}*(-I*$
 $a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+I*a-3*a*\tan(d*x+c))/(\tan(d*$
 $x+c)+I))*2^{(1/2)}*a+64*A*(-I*a)^{(1/2)}*\tan(d*x+c)*(a*\tan(d*x+c)*(1+I*\tan(d*x$
 $+c)))^{(1/2)}+12*B*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*(-I*a)^{(1/2)}/a^...$

3.559. $\int \frac{\sqrt{\cot(c+dx)(A+B \tan(c+dx))}}{(a+ia \tan(c+dx))^{3/2}} dx$

3.559.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 463 vs. $2(126) = 252$.

Time = 0.26 (sec) , antiderivative size = 463, normalized size of antiderivative = 2.76

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \frac{\left(3 \sqrt{\frac{1}{2}} a^2 d \sqrt{\frac{-iA^2-2AB+iB^2}{a^3 d^2}} e^{(3i dx+3i c)} \log \left(\frac{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (a^2 d e^{(2i dx+2i c)} - a^2 d) \sqrt{\frac{a}{(e^{(2i dx+2i c)} + 1)})} \sqrt{\frac{(I e^{(2i dx+2i c)} + I)}{(e^{(2i dx+2i c)} - 1)})} \sqrt{\frac{(-iA^2-2AB+iB^2)}{(a^3 d^2)}} + (A - IB) a e^{(I dx + I c)} e^{(-I dx - I c)} / (IA + B)}{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (a^2 d e^{(2i dx+2i c)} - a^2 d) \sqrt{\frac{a}{(e^{(2i dx+2i c)} + 1)})} \sqrt{\frac{(I e^{(2i dx+2i c)} + I)}{(e^{(2i dx+2i c)} - 1)})} \sqrt{\frac{(-iA^2-2AB+iB^2)}{(a^3 d^2)}} - (A - IB) a e^{(I dx + I c)} e^{(-I dx - I c)} / (IA + B)} - \sqrt{2} (2(4IA - B) e^{(4I dx + 4I c)} - (7IA - B) e^{(2I dx + 2I c)} - IA + B) \sqrt{\frac{a}{(e^{(2i dx+2i c)} + 1)})} \sqrt{\frac{(I e^{(2i dx+2i c)} + I)}{(e^{(2i dx+2i c)} - 1)})} e^{(-3I dx - 3I c)} / (a^2 d)} \right)}{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (a^2 d e^{(2i dx+2i c)} - a^2 d) \sqrt{\frac{a}{(e^{(2i dx+2i c)} + 1)})} \sqrt{\frac{(I e^{(2i dx+2i c)} + I)}{(e^{(2i dx+2i c)} - 1)})} \sqrt{\frac{(-iA^2-2AB+iB^2)}{(a^3 d^2)}} + (A - IB) a e^{(I dx + I c)} e^{(-I dx - I c)} / (IA + B)} - \sqrt{2} (2(4IA - B) e^{(4I dx + 4I c)} - (7IA - B) e^{(2I dx + 2I c)} - IA + B) \sqrt{\frac{a}{(e^{(2i dx+2i c)} + 1)})} \sqrt{\frac{(I e^{(2i dx+2i c)} + I)}{(e^{(2i dx+2i c)} - 1)})} e^{(-3I dx - 3I c)} / (a^2 d)} \right)}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="fricas")`

output `1/12*(3*sqrt(1/2)*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(a^2*d*e^(2*I*d*x + 2*I*c) - a^2*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2)) + (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B) - 3*sqrt(1/2)*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(4*(sqrt(2)*sqrt(1/2)*(a^2*d*e^(2*I*d*x + 2*I*c) - a^2*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2)) - (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B) - sqrt(2)*(2*(4*I*A - B)*e^(4*I*d*x + 4*I*c) - (7*I*A - B)*e^(2*I*d*x + 2*I*c) - I*A + B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*e^(-3*I*d*x - 3*I*c)/(a^2*d)`

3.559.6 Sympy [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx = \int \frac{(A+B \tan(c+dx)) \sqrt{\cot(c+dx)}}{(ia(\tan(c+dx) - i))^{\frac{3}{2}}} dx$$

input `integrate(cot(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(cot(c + d*x))/(I*a*(tan(c + d*x) - I))**(3/2), x)`

3.559. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{3/2}} dx$

3.559.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.559.8 Giac [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \int \frac{(B\tan(dx+c)+A)\sqrt{\cot(dx+c)}}{(ia\tan(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(cot(d*x + c))/(I*a*tan(d*x + c) + a)^(3/2), x)`

3.559.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx = \int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+a\tan(c+dx)1i)^{3/2}} dx$$

input `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(3/2),x)`

output `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(3/2), x)`

3.559. $\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{3/2}} dx$

$$3.560 \quad \int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+ia \tan(c+dx))}^{3/2}} dx$$

3.560.1 Optimal result	5286
3.560.2 Mathematica [A] (verified)	5286
3.560.3 Rubi [A] (verified)	5287
3.560.4 Maple [B] (verified)	5290
3.560.5 Fricas [B] (verification not implemented)	5291
3.560.6 Sympy [F]	5292
3.560.7 Maxima [F(-2)]	5292
3.560.8 Giac [F]	5293
3.560.9 Mupad [F(-1)]	5293

3.560.1 Optimal result

Integrand size = 38, antiderivative size = 170

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))}^{3/2}} dx =$$

$$\frac{\left(\frac{1}{4} + \frac{i}{4}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{a^{3/2}d}$$

$$+ \frac{iA - B}{3d\sqrt{\cot(c + dx)(a + ia \tan(c + dx))}^{3/2}} + \frac{iA + 5B}{6ad\sqrt{\cot(c + dx)}\sqrt{a + ia \tan(c + dx)}}$$

output `(-1/4-1/4*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/a^(3/2)/d+1/6*(I*A+5*B)/a/d/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2)+1/3*(I*A-B)/d/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(3/2)`

3.560.2 Mathematica [A] (verified)

Time = 5.08 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.87

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))}^{3/2}} dx = -\frac{3i\sqrt{2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{\sqrt{ia \tan(c+dx)}} + \frac{6(A-iB)+2(iA+5B)\tan(c+dx)}{(-i+\tan(c+dx))\sqrt{a+ia \tan(c+dx)}} + \frac{6(A-iB)+2(iA+5B)\tan(c+dx)}{12ad\sqrt{\cot(c + dx)}}$$

3.560. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+ia \tan(c+dx))}^{3/2}} dx$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^(3/2)),x]`

output `(((-3*I)*Sqrt[2]*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])]/Sqrt[I*a*Tan[c + d*x]] + (6*(A - I*B) + 2*(I*A + 5*B)*Tan[c + d*x])/((-I + Tan[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]]))/(12*a*d*Sqrt[Cot[c + d*x]])`

3.560.3 Rubi [A] (verified)

Time = 0.92 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.03, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.289$, Rules used = {3042, 4729, 3042, 4078, 27, 3042, 4079, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{4729} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\sqrt{\tan(c + dx)}(A + B \tan(c + dx))}{(i \tan(c + dx)a + a)^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\sqrt{\tan(c + dx)}(A + B \tan(c + dx))}{(i \tan(c + dx)a + a)^{3/2}} dx \\
 & \quad \downarrow \text{4078} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{(-B + iA) \sqrt{\tan(c + dx)}}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\int \frac{a(iA - B) - 2a(A - 2iB) \tan(c + dx)}{2\sqrt{\tan(c + dx)} \sqrt{i \tan(c + dx)a + a}} dx}{3a^2} \right) \\
 & \quad \downarrow \text{27} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{(-B + iA) \sqrt{\tan(c + dx)}}{3d(a + ia \tan(c + dx))^{3/2}} - \frac{\int \frac{a(iA - B) - 2a(A - 2iB) \tan(c + dx)}{\sqrt{\tan(c + dx)} \sqrt{i \tan(c + dx)a + a}} dx}{6a^2} \right)
 \end{aligned}$$

3.560. $\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} - \frac{\int \frac{a(iA-B)-2a(A-2iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} \right) \\
& \downarrow 4079 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} - \frac{\frac{\int \frac{3a^2(iA+B)\sqrt{i\tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx}{a^2} - \frac{a(5B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}}{6a^2} \right) \\
& \downarrow 27 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} - \frac{\frac{3}{2}(B+iA)\int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{a(5B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}}{6a^2} \right) \\
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} - \frac{\frac{3}{2}(B+iA)\int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{a(5B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}}{6a^2} \right) \\
& \downarrow 4027 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} - \frac{\frac{3ia^2(B+iA)\int \frac{1}{\frac{2\tan(c+dx)a^2}{i\tan(c+dx)a+a} - ia} d - \frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{6a^2} - \frac{a(5B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} \right) \\
& \downarrow 218 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} - \frac{\left(\frac{3}{2}-\frac{3i}{2}\right)\sqrt{a}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} - \frac{a(5B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}}{6a^2} \right)
\end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^(3/2)),x]`

$$3.560. \quad \int \frac{A+B\tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia\tan(c+dx))^{3/2}} dx$$

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((I*A - B)*Sqrt[Tan[c + d*x]])/(3*d
*(a + I*a*Tan[c + d*x])^(3/2)) - (((3/2 - (3*I)/2)*Sqrt[a]*(I*A + B)*ArcTan
h[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (
a*(I*A + 5*B)*Sqrt[Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]]))/(6*a^2))
```

3.560.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4027 Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4078 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[(-(A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```



```
rule 4079 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.560.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 662 vs. $2(136) = 272$.

Time = 0.53 (sec) , antiderivative size = 663, normalized size of antiderivative = 3.90

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(dx+c))} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \left(-12B \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{-ia} - 3iB \ln \left(-\frac{-2\sqrt{2}\sqrt{-a}}{\dots} \right) \right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(dx+c))} \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \left(-12B \sqrt{a \tan(dx+c)(1+i \tan(dx+c))} \sqrt{-ia} - 3iB \ln \left(-\frac{-2\sqrt{2}\sqrt{-a}}{\dots} \right) \right)}{\dots}$

```
input int((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(3/2),x,method=_R
ETURNVERBOSE)
```

output $1/24/d*(a*(1+I*\tan(d*x+c)))^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*(-12*B*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*(-I*a)^{(1/2)}-3*I*B*2^{(1/2)}*\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+I*a-3*a*\tan(d*x+c))/(tan(d*x+c)+I))*a-20*I*B*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)+6*B*2^{(1/2)}*\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+I*a-3*a*\tan(d*x+c))/(tan(d*x+c)+I))*a*\tan(d*x+c)+3*I*B*2^{(1/2)}*\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+I*a-3*a*\tan(d*x+c))/(tan(d*x+c)+I))*a*\tan(d*x+c)^2-12*I*A*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+3*A*\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+I*a-3*a*\tan(d*x+c))/(tan(d*x+c)+I))*2^{(1/2)}*a+4*A*(-I*a)^{(1/2)}*\tan(d*x+c)*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+6*I*A*2^{(1/2)}*\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+I*a-3*a*\tan(d*x+c))/(tan(d*x+c)+I))*a*\tan(d*x+c)-3*A*\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}+I*a-3*a*\tan(d*x+c))/(tan(d*x+c)+I))*2^{(1/2)}*a*\tan(d*x+c)^2/(1/tan(d*x+c))^{(1/2)}/a^3/(1+I*\tan(d*x+c))/tan(d*x+c)/(-I*a)^{(1/2)}/(-tan(d*x+c)+I)^2$

3.560.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 461 vs. 2(126) = 252.

Time = 0.25 (sec) , antiderivative size = 461, normalized size of antiderivative = 2.71

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx = \frac{\left(3 \sqrt{\frac{1}{2}} a^2 d \sqrt{\frac{i A^2 + 2 AB - i B^2}{a^3 d^2}} e^{(3i dx + 3i c)} \log \left(-\frac{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (i a^2 d e^{(2i dx + 2i c)} \right)} \right)}{\dots} \right)}{\dots}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="fricas")`

```
output 1/12*(3*sqrt(1/2)*a^2*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^3*d^2))*e^(3*I*d*x
+ 3*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(I*a^2*d*e^(2*I*d*x + 2*I*c) - I*a^2*d
)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2
*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^3*d^2)) + (A - I*B)*
a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 3*sqrt(1/2)*a^2*d*sqrt((I
*A^2 + 2*A*B - I*B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(-4*(sqrt(2)*sqrt(
1/2)*(-I*a^2*d*e^(2*I*d*x + 2*I*c) + I*a^2*d)*sqrt(a/(e^(2*I*d*x + 2*I*c)
+ 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*
A^2 + 2*A*B - I*B^2)/(a^3*d^2)) + (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x -
I*c)/(I*A + B)) + sqrt(2)*(2*(A - 2*I*B)*e^(4*I*d*x + 4*I*c) - (A - 5*I*B
)*e^(2*I*d*x + 2*I*c) - A - I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I
*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*e^(-3*I*d*x - 3*I*c)
/(a^2*d)
```

3.560.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{(ia (\tan(c + dx) - i))^{3/2} \sqrt{\cot(c + dx)}} dx$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)**(1/2)/(a+I*a*tan(d*x+c))**(3/2),x)
```

```
output Integral((A + B*tan(c + d*x))/((I*a*(tan(c + d*x) - I))**(3/2)*sqrt(cot(c
+ d*x))), x)
```

3.560.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(3/2),x, al
gorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un
defined.
```

3.560. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{3/2}} dx$

3.560.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^{3/2} \sqrt{\cot(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)^(3/2)*sqrt(cot(d*x + c))), x)`

3.560.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + a \tan(c + dx) 1i)^{3/2}} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(3/2)),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(3/2)), x)`

3.561
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$$

3.561.1 Optimal result 5294
 3.561.2 Mathematica [A] (verified) 5295
 3.561.3 Rubi [A] (verified) 5295
 3.561.4 Maple [B] (verified) 5300
 3.561.5 Fricas [B] (verification not implemented) 5301
 3.561.6 Sympy [F] 5302
 3.561.7 Maxima [F(-2)] 5302
 3.561.8 Giac [F] 5302
 3.561.9 Mupad [F(-1)] 5303

3.561.1 Optimal result

Integrand size = 38, antiderivative size = 243

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \frac{2(-1)^{3/4} B \arctan\left(\frac{(-1)^{3/4} \sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{a^{3/2} d}$$

$$+ \frac{\left(\frac{1}{4} + \frac{i}{4}\right) (iA + B) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a} \sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{a^{3/2} d}$$

$$+ \frac{iA - B}{3d \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} + \frac{A + 3iB}{2ad \sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)}}$$

output

```
2*(-1)^(3/4)*B*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/a^(3/2)/d+(1/4+1/4*I)*(I*A+B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/a^(3/2)/d+1/2*(A+3*I*B)/a/d/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2)+1/3*(I*A-B)/d/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(3/2)
```

3.561.2 Mathematica [A] (verified)

Time = 5.87 (sec) , antiderivative size = 232, normalized size of antiderivative = 0.95

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \frac{\left(\frac{1}{12} + \frac{i}{12}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left((12 + 12i)^4 \sqrt{-1} \sqrt{a} \operatorname{ArcSinh}\left[\frac{(-1)^{1/4} \sqrt{\tan(c + dx)}}{(12 + 12i)^4 \sqrt{-1} \sqrt{a}}\right]\right)}{\dots}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(3/2)),x]`

output `((1/12 + I/12)*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((12 + 12*I)*(-1)^(1/4)*Sqrt[a]*B*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]]*(1 + I*Tan[c + d*x])^(3/2) + 3*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*(-I + Tan[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]] + (1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]]*(-3*(A + (3*I)*B) + ((-5*I)*A + 11*B)*Tan[c + d*x]))/(a^(3/2)*d*(-I + Tan[c + d*x])*Sqrt[a + I*a*Tan[c + d*x]])`

3.561.3 Rubi [A] (verified)Time = 1.48 (sec) , antiderivative size = 232, normalized size of antiderivative = 0.95, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {3042, 4729, 3042, 4078, 27, 3042, 4078, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2}(a + ia \tan(c + dx))^{3/2}} dx \\ & \quad \downarrow \text{4729} \\ & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\tan^{\frac{3}{2}}(c + dx)(A + B \tan(c + dx))}{(i \tan(c + dx)a + a)^{3/2}} dx \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.561. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\tan(c+dx)^{3/2}(A+B \tan(c+dx))}{(i \tan(c+dx)a+a)^{3/2}} dx \\
 & \qquad \qquad \qquad \downarrow \text{4078} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{3\sqrt{\tan(c+dx)}(a(iA-B)+2iaB \tan(c+dx))}{2\sqrt{i \tan(c+dx)}a+a} dx}{3a^2} \right) \\
 & \qquad \qquad \qquad \downarrow \text{27} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{\sqrt{\tan(c+dx)}(a(iA-B)+2iaB \tan(c+dx))}{\sqrt{i \tan(c+dx)}a+a} dx}{2a^2} \right) \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{\sqrt{\tan(c+dx)}(a(iA-B)+2iaB \tan(c+dx))}{\sqrt{i \tan(c+dx)}a+a} dx}{2a^2} \right) \\
 & \qquad \qquad \qquad \downarrow \text{4078} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int -\frac{\sqrt{i \tan(c+dx)}a+a((A+3iB)a^2+4B \tan(c+dx)a^2)}{2\sqrt{\tan(c+dx)}} dx}{a^2} - \frac{a(A+3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{27} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{\sqrt{i \tan(c+dx)}a+a((A+3iB)a^2+4B \tan(c+dx)a^2)}{\sqrt{\tan(c+dx)}} dx}{2a^2} - \frac{a(A+3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{3042} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{3/2}} - \frac{\int \frac{\sqrt{i \tan(c+dx)}a+a((A+3iB)a^2+4B \tan(c+dx)a^2)}{\sqrt{\tan(c+dx)}} dx}{2a^2} - \frac{a(A+3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{4084}
 \end{aligned}$$

3.561. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{3/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} - \frac{a^2(A-iB)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx+4iaB\int\frac{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)}}{\sqrt{\tan(c+dx)}}dx}{2a^2}\right) - \frac{\phantom{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}}{2a^2}$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} - \frac{a^2(A-iB)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx+4iaB\int\frac{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)}}{\sqrt{\tan(c+dx)}}dx}{2a^2}\right) - \frac{\phantom{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}}{2a^2}$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} - \frac{4iaB\int\frac{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx - \frac{2ia^4(A-iB)\int\frac{1}{2\tan(c+dx)}dx}{2a^2}}{2a^2}\right) - \frac{\phantom{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}}{2a^2}$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} - \frac{4iaB\int\frac{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx + \frac{(1-i)a^{5/2}(A-iB)\arctan\frac{1}{d}}{2a^2}}{2a^2}\right) - \frac{\phantom{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}}{2a^2}$$

↓ 4082

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} - \frac{4ia^3B\int\frac{1}{\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}}d\tan(c+dx) + \frac{(1-i)a^{5/2}(A-iB)\arctan\frac{1}{d}}{2a^2}}{2a^2}\right) - \frac{\phantom{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}}{2a^2}$$

↓ 65

3.561. $\int \frac{A+B\tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{3/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} - \frac{8ia^3B \int \frac{1}{1-\frac{ia\tan(c+dx)}{a+ia\tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{ia\tan(c+dx)+a}} + \frac{(1-i)a^{5/2}(A-iB)\arctan\left(\frac{\sqrt{\tan(c+dx)}}{\sqrt{ia\tan(c+dx)+a}}\right)}{2a^2}}{2a^2} \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{3d(a+ia\tan(c+dx))^{3/2}} - \frac{(1-i)a^{5/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right) - 8(-1)^{3/4}a^{5/2}B\operatorname{arctan}\left(\frac{\sqrt{\tan(c+dx)}}{\sqrt{ia\tan(c+dx)+a}}\right)}{2a^2}}{2a^2} \right)$$

input `Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(3/2)),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((I*A - B)*Tan[c + d*x]^(3/2))/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) - (((-8*(-1)^(3/4)*a^(5/2)*B*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d + ((1 - I)*a^(5/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d)/(2*a^2) - (a*(A + (3*I)*B)*Sqrt[Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]]))/(2*a^2))`

3.561.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

- rule 218 $\text{Int}[(a_+) + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a) \cdot \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] \text{ ; FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$
- rule 3042 $\text{Int}[u_+, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$
- rule 4027 $\text{Int}[\text{Sqrt}[(a_+) + (b_+) \cdot \tan[(e_+) + (f_+)(x_+)]] / \text{Sqrt}[(c_+) + (d_+) \cdot \tan[(e_+) + (f_+)(x_+)]], x_Symbol] \rightarrow \text{Simp}[-2 \cdot a \cdot (b/f) \text{ Subst}[\text{Int}[1/(a \cdot c - b \cdot d - 2 \cdot a^2 \cdot x^2), x], x, \text{Sqrt}[c + d \cdot \tan[e + f \cdot x]] / \text{Sqrt}[a + b \cdot \tan[e + f \cdot x]]], x] \text{ ; FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0]$
- rule 4078 $\text{Int}[(a_+) + (b_+) \cdot \tan[(e_+) + (f_+)(x_+)])^{(m_+)} \cdot ((A_+) + (B_+) \cdot \tan[(e_+) + (f_+)(x_+)])^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[(-A \cdot b - a \cdot B) \cdot (a + b \cdot \tan[e + f \cdot x])^m \cdot (c + d \cdot \tan[e + f \cdot x])^n / (2 \cdot a \cdot f \cdot m), x] + \text{Simp}[1 / (2 \cdot a^2 \cdot m) \text{ Int}[(a + b \cdot \tan[e + f \cdot x])^{(m+1)} \cdot (c + d \cdot \tan[e + f \cdot x])^{(n-1)} \cdot \text{Simp}[A \cdot (a \cdot c \cdot m + b \cdot d \cdot n) - B \cdot (b \cdot c \cdot m + a \cdot d \cdot n) - d \cdot (b \cdot B \cdot (m - n) - a \cdot A \cdot (m + n)) \cdot \tan[e + f \cdot x], x], x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{LtQ}[m, 0] \ \&\& \ \text{GtQ}[n, 0]$
- rule 4082 $\text{Int}[(a_+) + (b_+) \cdot \tan[(e_+) + (f_+)(x_+)])^{(m_+)} \cdot ((A_+) + (B_+) \cdot \tan[(e_+) + (f_+)(x_+)])^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[b \cdot (B/f) \text{ Subst}[\text{Int}[(a + b \cdot x)^{(m-1)} \cdot (c + d \cdot x)^n, x], x, \tan[e + f \cdot x]], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B, m, n\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{EqQ}[A \cdot b + a \cdot B, 0]$
- rule 4084 $\text{Int}[(a_+) + (b_+) \cdot \tan[(e_+) + (f_+)(x_+)])^{(m_+)} \cdot ((A_+) + (B_+) \cdot \tan[(e_+) + (f_+)(x_+)])^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[(A \cdot b + a \cdot B) / b \text{ Int}[(a + b \cdot \tan[e + f \cdot x])^m \cdot (c + d \cdot \tan[e + f \cdot x])^n, x], x] - \text{Simp}[B / b \text{ Int}[(a + b \cdot \tan[e + f \cdot x])^m \cdot (c + d \cdot \tan[e + f \cdot x])^n \cdot (a - b \cdot \tan[e + f \cdot x]), x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, A, B, m, n\}, x] \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{NeQ}[A \cdot b + a \cdot B, 0]$

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.561.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1232 vs. $2(192) = 384$.

Time = 0.54 (sec) , antiderivative size = 1233, normalized size of antiderivative = 5.07

method	result	size
derivativedivides	Expression too large to display	1233
default	Expression too large to display	1233

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(3/2),x,method=_RETURNERVERBOSE)`

output
$$\frac{1}{24} \frac{I}{d} \frac{1}{(\tan(d*x+c))^{3/2}} \frac{1}{\tan(d*x+c)} \frac{(a*(1+I*\tan(d*x+c)))^{1/2}}{a^{2*(2+4*I*B*(-I*a)^{1/2}*\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}} \frac{(I*a)^{1/2}+a}{(I*a)^{1/2}} \frac{a+20*A*(I*a)^{1/2}}{(I*a)^{1/2}} \frac{(-I*a)^{1/2}*\tan(d*x+c)^2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2}}{44*I*B*(-I*a)^{1/2}} \frac{(I*a)^{1/2}}{(I*a)^{1/2}} \frac{(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2}}{\tan(d*x+c)^2-3*A^2^{1/2}} \frac{(I*a)^{1/2}}{(I*a)^{1/2}} \frac{\ln(-(-2*2^{1/2})*(-I*a)^{1/2}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}}{I*a-3*a*\tan(d*x+c)} \frac{1}{(\tan(d*x+c)+I)} \frac{a*\tan(d*x+c)^3-3*I*A^2^{1/2}}{(I*a)^{1/2}} \frac{(I*a)^{1/2}}{(I*a)^{1/2}} \frac{\ln(-(-2*2^{1/2})*(-I*a)^{1/2}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}}{I*a-3*a*\tan(d*x+c)} \frac{1}{(\tan(d*x+c)+I)} \frac{a-9*I*B*2^{1/2}}{(I*a)^{1/2}} \frac{(I*a)^{1/2}}{(I*a)^{1/2}} \frac{\ln(-(-2*2^{1/2})*(-I*a)^{1/2}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}}{I*a-3*a*\tan(d*x+c)} \frac{1}{(\tan(d*x+c)+I)} \frac{a*\tan(d*x+c)-32*I*A*(-I*a)^{1/2}}{(I*a)^{1/2}} \frac{(I*a)^{1/2}}{(I*a)^{1/2}} \frac{(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{1/2}}{\tan(d*x+c)+9*B*2^{1/2}} \frac{(I*a)^{1/2}}{(I*a)^{1/2}} \frac{\ln(-(-2*2^{1/2})*(-I*a)^{1/2}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}}{I*a-3*a*\tan(d*x+c)} \frac{1}{(\tan(d*x+c)+I)} \frac{a*\tan(d*x+c)^2+24*B*(-I*a)^{1/2}}{(I*a)^{1/2}} \frac{(I*a)^{1/2}}{(I*a)^{1/2}} \frac{\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}}{(I*a)^{1/2}+a} \frac{(I*a)^{1/2}}{(I*a)^{1/2}} \frac{a*\tan(d*x+c)^3-72*I*B*(-I*a)^{1/2}}{(I*a)^{1/2}} \frac{(I*a)^{1/2}}{(I*a)^{1/2}} \frac{\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}}{(I*a)^{1/2}} \frac{(I*a)^{1/2}}{(I*a)^{1/2}} \frac{a*\tan(d*x+c)^2+3*I*B*2^{1/2}}{(I*a)^{1/2}} \frac{(I*a)^{1/2}}{(I*a)^{1/2}} \frac{\ln(-(-2*2^{1/2})*(-I*a)^{1/2}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}}{(I*a)^{1/2}+I*a-3*a*\tan(d*x+c)} \frac{1}{(\tan(d*x+c)+I)} \frac{a*\tan(d*x+c)^3+9*A*(I*a)^{1/2}}{(I*a)^{1/2}} \frac{(I*a)^{1/2}}{(I*a)^{1/2}} \frac{\ln(-(-2*2^{1/2})*(-I*a)^{1/2}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))))^{1/2}}{I*a-3*...}$$

3.561.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 782 vs. $2(181) = 362$.

Time = 0.28 (sec) , antiderivative size = 782, normalized size of antiderivative = 3.22

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx =$$

$$\left(3 \sqrt{\frac{1}{2}} a^2 d \sqrt{\frac{-iA^2 - 2AB + iB^2}{a^3 d^2}} e^{(3i dx + 3i c)} \log \left(-\frac{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (a^2 d e^{(2i dx + 2i c)} - a^2 d) \sqrt{\frac{a}{e^{(2i dx + 2i c)} + 1}} \sqrt{\frac{i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} - 1}} \sqrt{\frac{-iA^2 - 2AB + iB^2}{a^3 d^2}}}{iA + B} \right) \right.$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="fracas")
```

```
output -1/12*(3*sqrt(1/2)*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(a^2*d*e^(2*I*d*x + 2*I*c) - a^2*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2)) + (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 3*sqrt(1/2)*a^2*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(4*(sqrt(2)*sqrt(1/2)*(a^2*d*e^(2*I*d*x + 2*I*c) - a^2*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^3*d^2)) - (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 3*a^2*d*sqrt(4*I*B^2/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(-16*(3*B*a^2*e^(2*I*d*x + 2*I*c) - B*a^2 + sqrt(2)*(I*a^3*d*e^(3*I*d*x + 3*I*c) - I*a^3*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt(4*I*B^2/(a^3*d^2)))*e^(-2*I*d*x - 2*I*c)/B + 3*a^2*d*sqrt(4*I*B^2/(a^3*d^2))*e^(3*I*d*x + 3*I*c)*log(-16*(3*B*a^2*e^(2*I*d*x + 2*I*c) - B*a^2 + sqrt(2)*(-I*a^3*d*e^(3*I*d*x + 3*I*c) + I*a^3*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt(4*I*B^2/(a^3*d^2)))*e^(-2*I*d*x - 2*I*c)/B + sqrt(2)*(2*(2*I*A - 5*B)*e^(4*I*d*x + 4*I*c) - (5*I*A - 11*B)*e^(2*I*d*x + 2*I*c) + I*A - B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c))...
```

3.561.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{(ia(\tan(c + dx) - i))^{\frac{3}{2}} \cot^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(3/2)/(a+I*a*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))/((I*a*(tan(c + d*x) - I))**(3/2)*cot(c + d*x)**(3/2)), x)`

3.561.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.561.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^{\frac{3}{2}} \cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^(3/2)), x)`

3.561.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2} (a + a \tan(c + dx) i)^{3/2}} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + a*tan(c + d*x)*1i)^(3/2)),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + a*tan(c + d*x)*1i)^(3/2)), x)`

3.562
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$$

3.562.1 Optimal result 5304
 3.562.2 Mathematica [A] (verified) 5305
 3.562.3 Rubi [A] (verified) 5305
 3.562.4 Maple [B] (verified) 5310
 3.562.5 Fricas [B] (verification not implemented) 5311
 3.562.6 Sympy [F(-1)] 5311
 3.562.7 Maxima [F(-2)] 5312
 3.562.8 Giac [F] 5312
 3.562.9 Mupad [F(-1)] 5312

3.562.1 Optimal result

Integrand size = 38, antiderivative size = 260

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{(\frac{1}{8} + \frac{i}{8})(A-iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{a^{5/2}d} + \frac{(A+iB)\sqrt{\cot(c+dx)}}{5d(a+ia \tan(c+dx))^{5/2}} + \frac{(17A+7iB)\sqrt{\cot(c+dx)}}{30ad(a+ia \tan(c+dx))^{3/2}} + \frac{(151A+41iB)\sqrt{\cot(c+dx)}}{60a^2d\sqrt{a+ia \tan(c+dx)}} - \frac{(317A+67iB)\sqrt{\cot(c+dx)}\sqrt{a+ia \tan(c+dx)}}{60a^3d}$$

output

```
(1/8+1/8*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/a^(5/2)/d+1/60*(151*A+41*I*B)*cot(d*x+c)^(1/2)/a^2/d/(a+I*a*tan(d*x+c))^(1/2)-1/60*(317*A+67*I*B)*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^(1/2)/a^3/d+1/5*(A+I*B)*cot(d*x+c)^(1/2)/d/(a+I*a*tan(d*x+c))^(5/2)+1/30*(17*A+7*I*B)*cot(d*x+c)^(1/2)/a/d/(a+I*a*tan(d*x+c))^(3/2)
```

3.562.2 Mathematica [A] (verified)

Time = 9.21 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.86

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx = \frac{-15i\sqrt{2}(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{ia \tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)(i+\cot(c+dx))^2\sqrt{a}}{120a^2d \cot^{\frac{5}{2}}(c+dx)\sqrt{ia \tan(c+dx)}}$$

input `Integrate[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `((-15*I)*Sqrt[2]*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*(I + Cot[c + d*x])^2*Sqrt[a + I*a*Tan[c + d*x]] + (2*a*(160*((-5*I)*A + B) - 15*(41*A + (7*I)*B)*Cot[c + d*x] + (120*I)*A*Cot[c + d*x]^2 + (317*A + (67*I)*B)*Tan[c + d*x]))/Sqrt[I*a*Tan[c + d*x]]/(120*a^2*d*Cot[c + d*x]^(5/2)*Sqrt[I*a*Tan[c + d*x]]*(-I + Tan[c + d*x])^2*Sqrt[a + I*a*Tan[c + d*x]])`

3.562.3 Rubi [A] (verified)Time = 1.60 (sec) , antiderivative size = 274, normalized size of antiderivative = 1.05, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.447$, Rules used = {3042, 4729, 3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4081, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\cot(c+dx)^{\frac{3}{2}}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx \\ & \quad \downarrow \text{4729} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(i \tan(c+dx)a+a)^{\frac{5}{2}}} dx \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.562. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{\frac{5}{2}}} dx$

$$\begin{aligned}
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B\tan(c+dx)}{\tan(c+dx)^{3/2}(i\tan(c+dx)a+a)^{5/2}} dx \\
& \quad \downarrow 4079 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(11A+iB)-6a(iA-B)\tan(c+dx)}{2\tan^{\frac{3}{2}}(c+dx)(i\tan(c+dx)a+a)^{3/2}} dx}{5a^2} + \frac{A+iB}{5d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{5/2}} \right) \\
& \quad \downarrow 27 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(11A+iB)-6a(iA-B)\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(i\tan(c+dx)a+a)^{3/2}} dx}{10a^2} + \frac{A+iB}{5d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{5/2}} \right) \\
& \quad \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(11A+iB)-6a(iA-B)\tan(c+dx)}{\tan(c+dx)^{3/2}(i\tan(c+dx)a+a)^{3/2}} dx}{10a^2} + \frac{A+iB}{5d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{5/2}} \right) \\
& \quad \downarrow 4079 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a^2(83A+13iB)-4a^2(17iA-7B)\tan(c+dx)}{2\tan^{\frac{3}{2}}(c+dx)\sqrt{i\tan(c+dx)a+a}} dx}{3a^2} + \frac{a(17A+7iB)}{3d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{3/2}} + \frac{1}{5d\sqrt{\tan(c+dx)}} \right) \\
& \quad \downarrow 27 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a^2(83A+13iB)-4a^2(17iA-7B)\tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} + \frac{a(17A+7iB)}{3d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{3/2}} + \frac{1}{5d\sqrt{\tan(c+dx)}} \right) \\
& \quad \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a^2(83A+13iB)-4a^2(17iA-7B)\tan(c+dx)}{\tan(c+dx)^{3/2}\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} + \frac{a(17A+7iB)}{3d\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^{3/2}} + \frac{1}{5d\sqrt{\tan(c+dx)}} \right) \\
& \quad \downarrow 4079
\end{aligned}$$

3.562. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(317A+67iB)-2a^3(151iA-41B) \tan(c+dx)) dx}{2 \tan^{\frac{3}{2}}(c+dx)} + \frac{a^2(151A+41iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}}{6a^2} + \frac{3d\sqrt{\tan(c+dx)}}{10a^2} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(317A+67iB)-2a^3(151iA-41B) \tan(c+dx)) dx}{\tan^{\frac{3}{2}}(c+dx)} + \frac{a^2(151A+41iB)}{2a^2 d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}}{6a^2} + \frac{3d\sqrt{\tan(c+dx)}}{10a^2} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{\sqrt{i \tan(c+dx)a+a} (a^3(317A+67iB)-2a^3(151iA-41B) \tan(c+dx)) dx}{\tan(c+dx)^{3/2}} + \frac{a^2(151A+41iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}}{6a^2} + \frac{3d\sqrt{\tan(c+dx)}}{10a^2} \right)$$

↓ 4081

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int \frac{15a^4(iA+B)\sqrt{i \tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx - \frac{2a^3(317A+67iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{a^2(151A+41iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}}{2a^2} + \frac{3d\sqrt{\tan(c+dx)}}{10a^2} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{15a^3(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(317A+67iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} + \frac{a^2(151A+41iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}}}{2a^2} + \frac{3d\sqrt{\tan(c+dx)}}{10a^2} \right)$$

↓ 3042

3.562. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{15a^3(B+iA) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - \frac{2a^3(317A+67iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{a^2(151A+41iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} + \frac{3d\sqrt{t}}{10a^2} \right) \\
 & \qquad \qquad \qquad \downarrow 4027 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{30ia^5(B+iA) \int \frac{1}{- \frac{2 \tan(c+dx)a^2}{i \tan(c+dx)a+a} - ia} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}} - \frac{2a^3(317A+67iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{2a^2} + \frac{a^2(151A+41iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} + \frac{3d\sqrt{t}}{10a^2} \right) \\
 & \qquad \qquad \qquad \downarrow 218 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\frac{a^2(151A+41iB)}{d\sqrt{\tan(c+dx)}\sqrt{a+ia \tan(c+dx)}} + \frac{(15-15i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} - \frac{2a^3(317A+67iB)\sqrt{a+ia \tan(c+dx)}}{d\sqrt{\tan(c+dx)}}}{6a^2} + \frac{3d\sqrt{t}}{10a^2} \right)
 \end{aligned}$$

```
input Int[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((A + I*B)/(5*d*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^(5/2)) + ((a*(17*A + (7*I)*B))/(3*d*Sqrt[Tan[c + d*x]])*(a + I*a*Tan[c + d*x])^(3/2)) + ((a^2*(151*A + (41*I)*B))/(d*Sqrt[Tan[c + d*x]]*Sqrt[a + I*a*Tan[c + d*x]]) + (((15 - 15*I)*a^(7/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d - (2*a^3*(317*A + (67*I)*B)*Sqrt[a + I*a*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]])))/(2*a^2))/(6*a^2))/(10*a^2))
```

3.562. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

3.562.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`
- rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.562.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1165 vs. $2(210) = 420$.

Time = 0.57 (sec) , antiderivative size = 1166, normalized size of antiderivative = 4.48

method	result	size
derivativedivides	Expression too large to display	1166
default	Expression too large to display	1166

```
input int(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x,method=_R
ETURNVERBOSE)
```

```
output 1/240/d*(1/tan(d*x+c))^(3/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)*(-15*I*
B*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/
2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I)*a*tan(d*x+c)-15*I*B*2^(1/2)*ln(-(-2
*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*
x+c))/(tan(d*x+c)+I)*a*tan(d*x+c)^5+15*A*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(
1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)
+I)*a*tan(d*x+c)^5+90*I*B*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x
+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I)*a*tan(d*x
+c)^3+60*I*A*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*
x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I)*a*tan(d*x+c)^2-60*B*2^(1/
2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-
3*a*tan(d*x+c))/(tan(d*x+c)+I)*a*tan(d*x+c)^4+4468*I*A*(a*tan(d*x+c)*(1+I
*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3+1060*I*B*(a*tan(d*x+c)*(1+I
tan(d*x+c)))^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2-1268*A*(a*tan(d*x+c)*(1+I*tan
(d*x+c)))^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^4-90*A*2^(1/2)*ln(-(-2*2^(1/2)*(-I
*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d
*x+c)+I)*a*tan(d*x+c)^3-60*I*A*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*ta
n(d*x+c)*(1+I*tan(d*x+c))))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I)*a*tan
(d*x+c)^4-2940*I*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)*tan(
d*x+c)-908*B*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)*tan(d*x...
```

$$3.562. \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$$

3.562.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 482 vs. $2(198) = 396$.

Time = 0.27 (sec) , antiderivative size = 482, normalized size of antiderivative = 1.85

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx =$$

$$\left(15\sqrt{\frac{1}{2}}a^3d\sqrt{\frac{iA^2+2AB-iB^2}{a^5d^2}}e^{(5i dx+5i c)} \log\left(-\frac{4\left(\sqrt{2}\sqrt{\frac{1}{2}}(ia^3de^{(2i dx+2i c)}-ia^3d)\sqrt{\frac{a}{e^{(2i dx+2i c)}+1}}\sqrt{\frac{ie^{(2i dx+2i c)}+i}{e^{(2i dx+2i c)}-1}}\sqrt{\frac{iA^2+2AB-iB^2}{a^5d^2}}\right)}{iA+B}\right) \right)$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="fricas")`

output `-1/120*(15*sqrt(1/2)*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(I*a^3*d*e^(2*I*d*x + 2*I*c) - I*a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2)) + (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 15*sqrt(1/2)*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(-I*a^3*d*e^(2*I*d*x + 2*I*c) + I*a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2)) + (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) + sqrt(2)*((463*A + 83*I*B)*e^(6*I*d*x + 6*I*c) - 2*(97*A + 32*I*B)*e^(4*I*d*x + 4*I*c) - 2*(13*A + 8*I*B)*e^(2*I*d*x + 2*I*c) - 3*A - 3*I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1)))*e^(-5*I*d*x - 5*I*c)/(a^3*d)`

3.562.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(5/2),x)`

output `Timed out`

3.562. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx$

3.562.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.562.8 Giac [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{(ia\tan(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(3/2)/(I*a*tan(d*x + c) + a)^(5/2), x)`

3.562.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx = \int \frac{\cot(c+dx)^{3/2}(A+B\tan(c+dx))}{(a+a\tan(c+dx)1i)^{5/2}} dx$$

input `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2),x)`

output `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + a*tan(c + d*x)*1i)^(5/2), x)`

3.562. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx$

3.563
$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$$

3.563.1 Optimal result 5313
 3.563.2 Mathematica [A] (verified) 5314
 3.563.3 Rubi [A] (verified) 5314
 3.563.4 Maple [B] (verified) 5318
 3.563.5 Fricas [B] (verification not implemented) 5319
 3.563.6 Sympy [F] 5319
 3.563.7 Maxima [F(-2)] 5320
 3.563.8 Giac [F] 5320
 3.563.9 Mupad [F(-1)] 5320

3.563.1 Optimal result

Integrand size = 38, antiderivative size = 214

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{\left(\frac{1}{8} - \frac{i}{8}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{a^{5/2}d} + \frac{A + iB}{5d\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{5/2}} + \frac{13A + 3iB}{30ad\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{3/2}} + \frac{67A - 3iB}{60a^2d\sqrt{\cot(c+dx)}\sqrt{a+ia \tan(c+dx)}}$$

```
output (1/8-1/8*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/a^(5/2)/d+1/60*(67*A-3*I*B)/a^2/d/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2)+1/5*(A+I*B)/d/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(5/2)+1/30*(13*A+3*I*B)/a/d/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(3/2)
```


3.563.2 Mathematica [A] (verified)

Time = 5.72 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.03

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{\cot^{3/2}(c+dx) \sec^2(c+dx)(ia \tan(c+dx))^{3/2}}{(2(19A+9iB+(8$$

input `Integrate[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])
^(5/2),x]`

output `(Cot[c + d*x]^(3/2)*Sec[c + d*x]^2*(I*a*Tan[c + d*x])^(3/2)*(2*(19*A + (9*I)*B + (86*A + (6*I)*B)*Cos[2*(c + d*x)] + (80*I)*A*Sin[2*(c + d*x)])*Sqrt[I*a*Tan[c + d*x]] + 15*Sqrt[2]*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)])*Sqrt[a + I*a*Tan[c + d*x]])/(120*a^4*d*(-I + Tan[c + d*x])^2*Sqrt[a + I*a*Tan[c + d*x]])`

3.563.3 Rubi [A] (verified)

Time = 1.26 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.06, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4729, 3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx \\ & \quad \downarrow \text{4729} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(i \tan(c+dx)a+a)^{5/2}} dx \end{aligned}$$

3.563. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
& \downarrow 4079 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(9A-iB)-4a(iA-B)\tan(c+dx)}{2\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)^{3/2}} dx}{5a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} \right) \\
& \downarrow 27 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(9A-iB)-4a(iA-B)\tan(c+dx)}{\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)^{3/2}} dx}{10a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} \right) \\
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(9A-iB)-4a(iA-B)\tan(c+dx)}{\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)^{3/2}} dx}{10a^2} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} \right) \\
& \downarrow 4079 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a^2(41A-9iB)-2a^2(13iA-3B)\tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}} dx}{3a^2} + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} \right) \\
& \downarrow 27 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a^2(41A-9iB)-2a^2(13iA-3B)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} \right) \\
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a^2(41A-9iB)-2a^2(13iA-3B)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} + \frac{(A+iB)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} \right) \\
& \downarrow 4079 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{15a^3(A-iB)\sqrt{i\tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx}{a^2} + \frac{a^2(67A-3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}} + \frac{(A+iB)}{5d(a+ia\tan(c+dx))^{5/2}} \right)
\end{aligned}$$

3.563. $\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+ia\tan(c+dx))^{5/2}} dx$

$$\begin{array}{c}
\downarrow 27 \\
\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\frac{15}{2}a(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + \frac{a^2(67A-3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} + \frac{(A+iB)}{5d(a+ia)} \right) \\
\downarrow 3042 \\
\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\frac{15}{2}a(A-iB) \int \frac{\sqrt{i \tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx + \frac{a^2(67A-3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} + \frac{(A+iB)}{5d(a+ia)} \right) \\
\downarrow 4027 \\
\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\frac{a^2(67A-3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}} - \frac{15ia^3(A-iB) \int \frac{1}{-2 \tan(c+dx)a^2} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i \tan(c+dx)a+a}}}{6a^2}}{10a^2} + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} \right) \\
\downarrow 218 \\
\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\left(\frac{15}{2} - \frac{15i}{2}\right)a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right)}{d} + \frac{a^2(67A-3iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia \tan(c+dx)}}}{6a^2} + \frac{a(13A+3iB)\sqrt{\tan(c+dx)}}{3d(a+ia \tan(c+dx))^{3/2}} \right)
\end{array}$$

input `Int[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + I*a*Tan[c + d*x])^(5/2),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((A + I*B)*Sqrt[Tan[c + d*x]])/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) + ((a*(13*A + (3*I)*B)*Sqrt[Tan[c + d*x]])/(3*d*(a + I*a*Tan[c + d*x])^(3/2)) + (((15/2 - (15*I)/2)*a^(3/2)*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d + (a^2*(67*A - (3*I)*B)*Sqrt[Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]]))/(6*a^2))/(10*a^2))`

3.563.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]]/Sqrt[(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4079 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`
- rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.563.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1093 vs. $2(172) = 344$.

Time = 0.58 (sec) , antiderivative size = 1094, normalized size of antiderivative = 5.11

method	result	size
derivativedivides	Expression too large to display	1094
default	Expression too large to display	1094

```
input int(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x,method=_R
ETURNVERBOSE)
```

```
output -1/240/d*(1/tan(d*x+c))^(1/2)*tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)/a^3*(1
5*I*A*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(
1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*a-90*I*A*2^(1/2)*ln((2*2^(1/2)*(-
I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(
d*x+c)+I))*a*tan(d*x+c)^2+15*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d
*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*a*tan(d*
x+c)^4+12*B*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^
3+60*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c))
)^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*a*tan(d*x+c)+60*A*2^(1/2)*ln((
2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c
)-I*a)/(tan(d*x+c)+I))*a*tan(d*x+c)^3+908*A*(a*tan(d*x+c)*(1+I*tan(d*x+c))
)^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2+268*I*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(
1/2)*(-I*a)^(1/2)*tan(d*x+c)^3-60*I*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*
a*tan(d*x+c)^3-90*B*2^(1/2)*ln((2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I
tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*x+c)+I))*a*tan(d*x+c)^2-12*I
*B*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^2-60*I*B*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)-60*A*ln((2*2^(1/2)*(-I*
a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+3*a*tan(d*x+c)-I*a)/(tan(d*
x+c)+I))*2^(1/2)*a*tan(d*x+c)-1060*I*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^...
```

3.563.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 481 vs. $2(162) = 324$.

Time = 0.26 (sec) , antiderivative size = 481, normalized size of antiderivative = 2.25

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \frac{\left(15 \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{-iA^2-2AB+iB^2}{a^5 d^2}} e^{(5i dx+5i c)} \log \left(-\frac{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (a^3 d e^{(2i dx+2i c)} - a^3 d) \sqrt{\frac{a}{(e^{(2i dx+2i c)} + 1)})} \sqrt{\frac{(I e^{(2i dx+2i c)} + I)}{(e^{(2i dx+2i c)} - 1)})} \sqrt{\frac{(-iA^2-2AB+iB^2)}{(a^5 d^2)}} + (A - IB) a e^{(I dx + I c)} e^{(-I dx - I c)} / (IA + B)}{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (a^3 d e^{(2i dx+2i c)} - a^3 d) \sqrt{\frac{a}{(e^{(2i dx+2i c)} + 1)})} \sqrt{\frac{(I e^{(2i dx+2i c)} + I)}{(e^{(2i dx+2i c)} - 1)})} \sqrt{\frac{(-iA^2-2AB+iB^2)}{(a^5 d^2)}} - (A - IB) a e^{(I dx + I c)} e^{(-I dx - I c)} / (IA + B)} + \sqrt{2} * ((-83IA + 3B) e^{(6I dx + 6I c)} - 2 * (-32IA - 3B) e^{(4I dx + 4I c)} - 2 * (-8IA + 3B) e^{(2I dx + 2I c)} + 3IA - 3B) \sqrt{\frac{a}{(e^{(2i dx+2i c)} + 1)})} \sqrt{\frac{(I e^{(2i dx+2i c)} + I)}{(e^{(2i dx+2i c)} - 1)})} e^{(-5I dx - 5I c)} / (a^3 d)} \right)}{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (a^3 d e^{(2i dx+2i c)} - a^3 d) \sqrt{\frac{a}{(e^{(2i dx+2i c)} + 1)})} \sqrt{\frac{(I e^{(2i dx+2i c)} + I)}{(e^{(2i dx+2i c)} - 1)})} \sqrt{\frac{(-iA^2-2AB+iB^2)}{(a^5 d^2)}} + (A - IB) a e^{(I dx + I c)} e^{(-I dx - I c)} / (IA + B)}\right)}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="fricas")`

output `1/120*(15*sqrt(1/2)*a^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(a^3*d*e^(2*I*d*x + 2*I*c) - a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2)) + (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 15*sqrt(1/2)*a^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(4*(sqrt(2)*sqrt(1/2)*(a^3*d*e^(2*I*d*x + 2*I*c) - a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2)) - (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) + sqrt(2)*((-83*I*A + 3*B)*e^(6*I*d*x + 6*I*c) - 2*(-32*I*A - 3*B)*e^(4*I*d*x + 4*I*c) - 2*(-8*I*A + 3*B)*e^(2*I*d*x + 2*I*c) + 3*I*A - 3*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*e^(-5*I*d*x - 5*I*c)/(a^3*d)`

3.563.6 Sympy [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \int \frac{(A+B \tan(c+dx)) \sqrt{\cot(c+dx)}}{(ia(\tan(c+dx) - i))^{5/2}} dx$$

input `integrate(cot(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(cot(c + d*x))/(I*a*(tan(c + d*x) - I))* (5/2), x)`

3.563. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

3.563.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.563.8 Giac [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \int \frac{(B \tan(dx+c)+A)\sqrt{\cot(dx+c)}}{(ia \tan(dx+c)+a)^{5/2}} dx$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(d*x+c)+A)*sqrt(cot(d*x+c))/(I*a*tan(d*x+c)+a)^(5/2),x)`

3.563.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx = \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+a \tan(c+dx) 1i)^{5/2}} dx$$

input `int((cot(c+d*x)^(1/2)*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*1i)^(5/2),x)`

output `int((cot(c+d*x)^(1/2)*(A+B*tan(c+d*x)))/(a+a*tan(c+d*x)*1i)^(5/2),x)`

3.563. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+ia \tan(c+dx))^{5/2}} dx$

$$3.564 \quad \int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+ia \tan(c+dx))}^{5/2}} dx$$

3.564.1 Optimal result 5321
 3.564.2 Mathematica [A] (verified) 5322
 3.564.3 Rubi [A] (verified) 5322
 3.564.4 Maple [B] (verified) 5326
 3.564.5 Fricas [B] (verification not implemented) 5327
 3.564.6 Sympy [F] 5328
 3.564.7 Maxima [F(-2)] 5328
 3.564.8 Giac [F] 5329
 3.564.9 Mupad [F(-1)] 5329

3.564.1 Optimal result

Integrand size = 38, antiderivative size = 216

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + ia \tan(c + dx))}^{5/2}} dx =$$

$$\frac{\left(\frac{1}{8} + \frac{i}{8}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{a^{5/2}d}$$

$$+ \frac{iA - B}{5d\sqrt{\cot(c + dx)(a + ia \tan(c + dx))}^{5/2}} + \frac{3iA + 7B}{30ad\sqrt{\cot(c + dx)(a + ia \tan(c + dx))}^{3/2}}$$

$$- \frac{3iA - 13B}{60a^2d\sqrt{\cot(c + dx)}\sqrt{a + ia \tan(c + dx)}}$$

```
output (-1/8-1/8*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/a^(5/2)/d+1/60*(-3*I*A+13*B)/a^2/d/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2)+1/5*(I*A-B)/d/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(5/2)+1/30*(3*I*A+7*B)/a/d/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(3/2)
```


3.564.2 Mathematica [A] (verified)

Time = 5.63 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.03

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx = \frac{\sec^2(c + dx) \left(-2i(9A - iB + 2(3A - 7iB) \cos(2(c + dx))) + 2 \right)}{12} + \dots$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^(5/2)),x]`

output `(Sec[c + d*x]^2*((-2*I)*(9*A - I*B + 2*(3*A - (7*I)*B)*Cos[2*(c + d*x)] + 20*B*Sin[2*(c + d*x)])*Sqrt[I*a*Tan[c + d*x]] + 15*Sqrt[2]*(I*A + B)*ArcTan[((Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)])*Sqrt[a + I*a*Tan[c + d*x]])]/(120*a^2*d*Sqrt[Cot[c + d*x]]*Sqrt[I*a*Tan[c + d*x]]*(-I + Tan[c + d*x])^2*Sqrt[a + I*a*Tan[c + d*x]])`

3.564.3 Rubi [A] (verified)Time = 1.26 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.06, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4729, 3042, 4078, 27, 3042, 4079, 27, 3042, 4079, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{4729} \\ & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\sqrt{\tan(c + dx)}(A + B \tan(c + dx))}{(i \tan(c + dx)a + a)^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\sqrt{\tan(c + dx)}(A + B \tan(c + dx))}{(i \tan(c + dx)a + a)^{5/2}} dx \end{aligned}$$

3.564. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \downarrow 4078 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}}-\frac{\int\frac{a(iA-B)-2a(2A-3iB)\tan(c+dx)}{2\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)^{3/2}}dx}{5a^2}\right) \\
 & \downarrow 27 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}}-\frac{\int\frac{a(iA-B)-2a(2A-3iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)^{3/2}}dx}{10a^2}\right) \\
 & \downarrow 3042 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}}-\frac{\int\frac{a(iA-B)-2a(2A-3iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)^{3/2}}dx}{10a^2}\right) \\
 & \downarrow 4079 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}}-\frac{\int\frac{a^2(9iA+B)-2a^2(3A-7iB)\tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}}dx}{3a^2}-\frac{a(7B+3iA)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}}\right) \\
 & \downarrow 27 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}}-\frac{\int\frac{a^2(9iA+B)-2a^2(3A-7iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}}dx}{6a^2}-\frac{a(7B+3iA)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}}\right) \\
 & \downarrow 3042 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}}-\frac{\int\frac{a^2(9iA+B)-2a^2(3A-7iB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}}dx}{6a^2}-\frac{a(7B+3iA)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}}\right) \\
 & \downarrow 4079 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}}-\frac{\int\frac{15a^3(iA+B)\sqrt{i\tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}}dx}{a^2}+\frac{a^2(-13B+3iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}-\frac{a(7B+3iA)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}}\right)
 \end{aligned}$$

3.564. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+ia \tan(c+dx))^{5/2}}} dx$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{15}{2}a(B+iA)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx + \frac{a^2(-13B+3iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}}{6a^2} - \frac{a(7E)}{3d(a)}$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{15}{2}a(B+iA)\int\frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx + \frac{a^2(-13B+3iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}}{6a^2} - \frac{a(7E)}{3d(a)}$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{15ia^3(B+iA)\int\frac{1}{-2\tan(c+dx)a^2}d\frac{\sqrt{\tan(c+dx)}}{i\tan(c+dx)a+a} - ia}{d\sqrt{a+ia\tan(c+dx)}} - \frac{a^2(-13B+3iA)\sqrt{\tan(c+dx)}}{6a^2}}{10a^2}$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\sqrt{\tan(c+dx)}}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{(\frac{15}{2}-\frac{15i}{2})a^{3/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right) + \frac{a^2(-13B+3iA)}{d\sqrt{a+ia\tan(c+dx)}}}{6a^2}}{10a^2}$$

```
input Int[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^(5/2)),x]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((I*A - B)*Sqrt[Tan[c + d*x]])/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) - (-1/3*(a*((3*I)*A + 7*B)*Sqrt[Tan[c + d*x]])/(d*(a + I*a*Tan[c + d*x])^(3/2)) + (((15/2 - (15*I)/2)*a^(3/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])/d + (a^2*((3*I)*A - 13*B)*Sqrt[Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]])))/(6*a^2)/(10*a^2)
```

3.564. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{5/2}} dx$

3.564.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4078 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)), x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]`
- rule 4079 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m - b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && !GtQ[n, 0]`

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.564.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 855 vs. $2(174) = 348$.

Time = 0.54 (sec) , antiderivative size = 856, normalized size of antiderivative = 3.96

method	result
derivativedivides	$\frac{i\sqrt{a(1+i\tan(dx+c))}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))}\left(-45iB\ln\left(-\frac{-2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))+ia-3a\tan(dx+c)}}{\tan(dx+c)+i}\right)\right)}{\dots}$
default	$\frac{i\sqrt{a(1+i\tan(dx+c))}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))}\left(-45iB\ln\left(-\frac{-2\sqrt{2}\sqrt{-ia}\sqrt{a\tan(dx+c)(1+i\tan(dx+c))+ia-3a\tan(dx+c)}}{\tan(dx+c)+i}\right)\right)}{\dots}$

```
input int((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(5/2),x,method=_R
ETURNVERBOSE)
```

output

```
-1/240*I/d*(a*(1+I*tan(d*x+c)))^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
)*(-45*I*B*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x
+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2-15*A*ln(-(-
2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d
*x+c))/(tan(d*x+c)+I))*2^(1/2)*a+12*I*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*ta
n(d*x+c)))^(1/2)*tan(d*x+c)^2+15*I*A*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*
(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*
a*tan(d*x+c)^3+45*A*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x
+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*2^(1/2)*a*tan(d*x+c)^2+16
0*I*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*tan(d*x+c)+60*B*(
a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)-45*I*A*2^(1/2)*ln(-(-2*2
^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+
c))/(tan(d*x+c)+I))*a*tan(d*x+c)+60*I*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*ta
n(d*x+c)))^(1/2)-45*B*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(
1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)-52
*B*(-I*a)^(1/2)*tan(d*x+c)^2*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+15*I*B*
2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)
+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a+15*B*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)
^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+
c)+I))*a*tan(d*x+c)^3)/(1/tan(d*x+c))^(1/2)/a^4/(1+I*tan(d*x+c))/tan(d*...
```

3.564.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 482 vs. $2(162) = 324$.

Time = 0.25 (sec) , antiderivative size = 482, normalized size of antiderivative = 2.23

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx = \left(15 \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{iA^2 + 2AB - iB^2}{a^5 d^2}} e^{(5i dx + 5i c)} \log \left(-\frac{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (i a^3 d e^{(2i dx + 2i c)} \right)} \right)} \right)$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="fricas")`

output `1/120*(15*sqrt(1/2)*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(I*a^3*d*e^(2*I*d*x + 2*I*c) - I*a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2)) + (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 15*sqrt(1/2)*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(-I*a^3*d*e^(2*I*d*x + 2*I*c) + I*a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2)) + (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) + sqrt(2)*((3*A - 17*I*B)*e^(6*I*d*x + 6*I*c) + 2*(3*A + 8*I*B)*e^(4*I*d*x + 4*I*c) - 2*(3*A - 2*I*B)*e^(2*I*d*x + 2*I*c) - 3*A - 3*I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*e^(-5*I*d*x - 5*I*c)/(a^3*d)`

3.564.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{(ia (\tan(c + dx) - i))^{5/2} \sqrt{\cot(c + dx)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(1/2)/(a+I*a*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))/((I*a*(tan(c + d*x) - I))**(5/2)*sqrt(cot(c + d*x))), x)`

3.564.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is not defined.`

3.564. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^{5/2}} dx$

3.564.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^{5/2} \sqrt{\cot(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)^(5/2)*sqrt(cot(d*x + c))), x)`

3.564.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + a \tan(c + dx) 1i)^{5/2}} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(5/2)),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + a*tan(c + d*x)*1i)^(5/2)), x)`

$$3.565 \quad \int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$$

3.565.1 Optimal result	5330
3.565.2 Mathematica [A] (verified)	5331
3.565.3 Rubi [A] (verified)	5331
3.565.4 Maple [B] (verified)	5335
3.565.5 Fricas [B] (verification not implemented)	5336
3.565.6 Sympy [F(-1)]	5337
3.565.7 Maxima [F(-2)]	5337
3.565.8 Giac [F]	5338
3.565.9 Mupad [F(-1)]	5338

3.565.1 Optimal result

Integrand size = 38, antiderivative size = 214

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \frac{\left(\frac{1}{8} + \frac{i}{8}\right) (iA + B) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)}}{a^{5/2}d} + \frac{iA - B}{5d \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} + \frac{A + 11iB}{30ad \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^{3/2}} + \frac{13A - 37iB}{60a^2d \sqrt{\cot(c + dx)} \sqrt{a + ia \tan(c + dx)}}$$

```
output (1/8+1/8*I)*(I*A+B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/a^(5/2)/d+1/60*(13*A-37*I*B)/a^2/d/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2)+1/5*(I*A-B)/d/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(5/2)+1/30*(A+11*I*B)/a/d/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(3/2)
```

3.565.2 Mathematica [A] (verified)

Time = 7.13 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.05

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \frac{\cot^{\frac{3}{2}}(c + dx) \sec^2(c + dx)(ia \tan(c + dx))^{3/2} \left(2(A + 11iB + 2\right)}{2(A + 11iB + 2)}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(5/2)),x]`

output `(Cot[c + d*x]^(3/2)*Sec[c + d*x]^2*(I*a*Tan[c + d*x])^(3/2)*(2*(A + (11*I)*B + 2*(7*A - (13*I)*B)*Cos[2*(c + d*x)] + 20*(I*A + B)*Sin[2*(c + d*x)])*Sqrt[I*a*Tan[c + d*x]] - 15*Sqrt[2]*(A - I*B)*ArcTanh[(Sqrt[2]*Sqrt[I*a*Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)])*Sqrt[a + I*a*Tan[c + d*x]])/(120*a^4*d*(-I + Tan[c + d*x])^2*Sqrt[a + I*a*Tan[c + d*x]])`

3.565.3 Rubi [A] (verified)Time = 1.28 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.06, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$, Rules used = {3042, 4729, 3042, 4078, 27, 3042, 4078, 27, 3042, 4079, 27, 3042, 4027, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2}(a + ia \tan(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{4729} \\ & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\tan^{\frac{3}{2}}(c + dx)(A + B \tan(c + dx))}{(i \tan(c + dx)a + a)^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\tan(c + dx)^{3/2}(A + B \tan(c + dx))}{(i \tan(c + dx)a + a)^{5/2}} dx \end{aligned}$$

3.565. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$

$$\begin{aligned} & \downarrow 4078 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\int \frac{\sqrt{\tan(c+dx)}(3a(iA-B)-2a(A-4iB)\tan(c+dx))}{2(i\tan(c+dx)a+a)^{3/2}} dx}{5a^2}\right) \\ & \downarrow 27 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\int \frac{\sqrt{\tan(c+dx)}(3a(iA-B)-2a(A-4iB)\tan(c+dx))}{(i\tan(c+dx)a+a)^{3/2}} dx}{10a^2}\right) \\ & \downarrow 3042 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\int \frac{\sqrt{\tan(c+dx)}(3a(iA-B)-2a(A-4iB)\tan(c+dx))}{(i\tan(c+dx)a+a)^{3/2}} dx}{10a^2}\right) \\ & \downarrow 4078 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\int -\frac{(A+11iB)a^2+2(7iA+13B)\tan(c+dx)a^2}{2\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}} dx}{3a^2} - \frac{a(A+11iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))}\right) \\ & \downarrow 27 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\int \frac{(A+11iB)a^2+2(7iA+13B)\tan(c+dx)a^2}{\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} - \frac{a(A+11iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}}\right) \\ & \downarrow 3042 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\int \frac{(A+11iB)a^2+2(7iA+13B)\tan(c+dx)a^2}{\sqrt{\tan(c+dx)}\sqrt{i\tan(c+dx)a+a}} dx}{6a^2} - \frac{a(A+11iB)\sqrt{\tan(c+dx)}}{3d(a+ia\tan(c+dx))^{3/2}}\right) \\ & \downarrow 4079 \end{aligned}$$

3.565. $\int \frac{A+B\tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\int \frac{15a^3(A-iB)\sqrt{i\tan(c+dx)a+a}}{2\sqrt{\tan(c+dx)}} dx}{a^2} - \frac{a^2(13A-37iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{a(A+iB)}{3d(a+ia\tan(c+dx))}\right) - \frac{a^2(13A-37iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{a(A+iB)}{3d(a+ia\tan(c+dx))}$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{15}{2}a(A-iB)\int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{6a^2} - \frac{a^2(13A-37iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{a(A+iB)}{3d(a+ia\tan(c+dx))}\right) - \frac{a^2(13A-37iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{a(A+iB)}{3d(a+ia\tan(c+dx))}$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{15}{2}a(A-iB)\int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx}{6a^2} - \frac{a^2(13A-37iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{a(A+iB)}{3d(a+ia\tan(c+dx))}\right) - \frac{a^2(13A-37iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{a(A+iB)}{3d(a+ia\tan(c+dx))}$$

↓ 4027

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{15ia^3(A-iB)\int \frac{1}{-2\tan(c+dx)a^2-ia}}{d} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{i\tan(c+dx)a+a}}}{6a^2} - \frac{a^2(13A-37iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{a(A+iB)}{3d(a+ia\tan(c+dx))}\right) - \frac{a^2(13A-37iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{a(A+iB)}{3d(a+ia\tan(c+dx))}$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(-B+iA)\tan^{\frac{3}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\left(\frac{15}{2}-\frac{15i}{2}\right)a^{3/2}(A-iB)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d} - \frac{a^2(13A-37iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}}}{6a^2} - \frac{a(A+iB)}{3d(a+ia\tan(c+dx))}\right) - \frac{a^2(13A-37iB)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{a(A+iB)}{3d(a+ia\tan(c+dx))}$$

```
input Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^(5/2)),x]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((I*A - B)*Tan[c + d*x]^(3/2))/(5*d
*(a + I*a*Tan[c + d*x])^(5/2)) - (-1/3*(a*(A + (11*I)*B)*Sqrt[Tan[c + d*x]
])/d*(a + I*a*Tan[c + d*x])^(3/2)) + (((15/2 - (15*I)/2)*a^(3/2)*(A - I*B
)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])
)/d - (a^2*(13*A - (37*I)*B)*Sqrt[Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d
*x]])))/(6*a^2)/(10*a^2))
```

3.565.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4027 Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_)
+ (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*
a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /;
FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && N
eQ[c^2 + d^2, 0]
```

```
rule 4078 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-(A*b - a*B))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

```
rule 4079 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(a*A + b*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(2*f*m*(
b*c - a*d))), x] + Simp[1/(2*a*m*(b*c - a*d)) Int[(a + b*Tan[e + f*x])^(m
+ 1)*(c + d*Tan[e + f*x])^n*Simp[A*(b*c*m - a*d*(2*m + n + 1)) + B*(a*c*m
- b*d*(n + 1)) + d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x], x], x] /; Free
Q[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0]
&& LtQ[m, 0] && !GtQ[n, 0]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.565.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1105 vs. $2(172) = 344$.

Time = 0.54 (sec) , antiderivative size = 1106, normalized size of antiderivative = 5.17

method	result	size
derivativedivides	Expression too large to display	1106
default	Expression too large to display	1106

```
input int((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(5/2),x,method=_R
ETURNVERBOSE)
```

```
output 1/240/d/(1/tan(d*x+c))^(3/2)/tan(d*x+c)*(a*(1+I*tan(d*x+c)))^(1/2)/a^3*(60
*I*B*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^
(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)-148*B*(a*tan(d*x+c)
*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)*tan(d*x+c)^3-60*I*B*2^(1/2)*ln(-(-2*
2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x
+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^3+15*B*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1
/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+
I))*a*tan(d*x+c)^4-212*A*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2
)*tan(d*x+c)^2+15*I*A*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(
1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a-90*I*A*2^(1/2
)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3
*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^2+60*A*2^(1/2)*ln(-(-2*2^(1/2)
*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(t
an(d*x+c)+I))*a*tan(d*x+c)^3-52*I*A*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x
+c)))^(1/2)*tan(d*x+c)^3+308*I*B*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(d*x+
c)))^(1/2)*tan(d*x+c)^2-90*B*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d
*x+c)*(1+I*tan(d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x
+c)^2+15*I*A*2^(1/2)*ln(-(-2*2^(1/2)*(-I*a)^(1/2)*(a*tan(d*x+c)*(1+I*tan(
d*x+c)))^(1/2)+I*a-3*a*tan(d*x+c))/(tan(d*x+c)+I))*a*tan(d*x+c)^4-60*I*B*(
a*tan(d*x+c)*(1+I*tan(d*x+c)))^(1/2)*(-I*a)^(1/2)-60*A*2^(1/2)*ln(-(-2*...
```

3.565.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 482 vs. 2(160) = 320.

Time = 0.25 (sec) , antiderivative size = 482, normalized size of antiderivative = 2.25

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx =$$

$$\left(15 \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{-i A^2 - 2AB + i B^2}{a^5 d^2}} e^{(5i dx + 5i c)} \log \left(-\frac{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (a^3 d e^{(2i dx + 2i c)} - a^3 d) \sqrt{\frac{a}{e^{(2i dx + 2i c)} + 1}} \sqrt{\frac{i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} - 1}} \sqrt{\frac{-i A^2 - 2AB + i B^2}{a^5 d^2}}}{i A + B} \right) \right)$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(5/2),x, al
gorithm="fricas")
```

```
output -1/120*(15*sqrt(1/2)*a^3*d*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2))*e^(5*I
*d*x + 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(a^3*d*e^(2*I*d*x + 2*I*c) - a^3*d
)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2
*I*d*x + 2*I*c) - 1))*sqrt((-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2)) + (A - I*B)
*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 15*sqrt(1/2)*a^3*d*sqrt(
(-I*A^2 - 2*A*B + I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(4*(sqrt(2)*sqr
t(1/2)*(a^3*d*e^(2*I*d*x + 2*I*c) - a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1
))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((-I*A^
2 - 2*A*B + I*B^2)/(a^5*d^2)) - (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I
*c)/(I*A + B)) - sqrt(2)*((-17*I*A - 23*B)*e^(6*I*d*x + 6*I*c) - 2*(-8*I*A
- 17*B)*e^(4*I*d*x + 4*I*c) - 2*(-2*I*A + 7*B)*e^(2*I*d*x + 2*I*c) - 3*I*
A + 3*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I
)/(e^(2*I*d*x + 2*I*c) - 1))*e^(-5*I*d*x - 5*I*c)/(a^3*d)
```

3.565.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)**(3/2)/(a+I*a*tan(d*x+c))**(5/2),x)
```

```
output Timed out
```

3.565.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(5/2),x, al
gorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un
defined.
```

3.565. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$

3.565.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^{\frac{5}{2}} \cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^(3/2)), x)`

3.565.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2} (a + a \tan(c + dx) li)^{5/2}} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + a*tan(c + d*x)*li)^(5/2)),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + a*tan(c + d*x)*li)^(5/2)), x)`

3.566
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$$

3.566.1 Optimal result	5339
3.566.2 Mathematica [A] (verified)	5340
3.566.3 Rubi [A] (verified)	5340
3.566.4 Maple [B] (verified)	5345
3.566.5 Fricas [B] (verification not implemented)	5346
3.566.6 Sympy [F(-1)]	5347
3.566.7 Maxima [F(-2)]	5348
3.566.8 Giac [F]	5348
3.566.9 Mupad [F(-1)]	5348

3.566.1 Optimal result

Integrand size = 38, antiderivative size = 289

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \frac{2\sqrt[4]{-1}B \arctan\left(\frac{(-1)^{3/4}\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{a^{5/2}d}$$

$$+ \frac{\left(\frac{1}{8} + \frac{i}{8}\right) (A - iB) \operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{a^{5/2}d}$$

$$+ \frac{iA - B}{5d \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} + \frac{A + 3iB}{6ad \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^{3/2}}$$

$$- \frac{iA - 7B}{4a^2d\sqrt{\cot(c + dx)}\sqrt{a + ia \tan(c + dx)}}$$

output

```
2*(-1)^(1/4)*B*arctan((-1)^(3/4)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/a^(5/2)/d+(1/8+1/8*I)*(A-I*B)*arctanh((1+I)*a^(1/2)*tan(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/a^(5/2)/d+1/4*(-I*A+7*B)/a^2/d/cot(d*x+c)^(1/2)/(a+I*a*tan(d*x+c))^(1/2)+1/5*(I*A-B)/d/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^(5/2)+1/6*(A+3*I*B)/a/d/cot(d*x+c)^(3/2)/(a+I*a*tan(d*x+c))^(3/2)
```

3.566.
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$$

3.566.2 Mathematica [A] (verified)

Time = 9.97 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.11

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \frac{\left(\frac{1}{120} + \frac{i}{120}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left((-120 - 120i) \sqrt[4]{-1}\right)}{\dots}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^(5/2)),x]`

output `((1/120 + I/120)*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-120 - 120*I)*(-1)^(1/4)*Sqrt[a]*B*ArcSinh[(-1)^(1/4)*Sqrt[Tan[c + d*x]]]*Sec[c + d*x]^3*(Cos[3*(c + d*x)] + I*Sin[3*(c + d*x)]) - Sec[c + d*x]^2*Sqrt[1 + I*Tan[c + d*x]]*((-1 - I)*Sqrt[a]*(-11*A - (21*I)*B + 2*(13*A + (63*I)*B)*Cos[2*(c + d*x)] + (20*I)*(A + (6*I)*B)*Sin[2*(c + d*x)])*Sqrt[Tan[c + d*x]] + 15*(A - I*B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]])*(Cos[2*(c + d*x)] + I*Sin[2*(c + d*x)])*Sqrt[a + I*a*Tan[c + d*x]])/(a^(5/2)*d*Sqrt[1 + I*Tan[c + d*x]]*(-I + Tan[c + d*x])^2*Sqrt[a + I*a*Tan[c + d*x]])`

3.566.3 Rubi [A] (verified)

Time = 1.82 (sec) , antiderivative size = 285, normalized size of antiderivative = 0.99, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4729, 3042, 4078, 27, 3042, 4078, 27, 3042, 4078, 27, 3042, 4084, 3042, 4027, 218, 4082, 65, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{5/2}(a + ia \tan(c + dx))^{5/2}} dx \\ & \quad \downarrow \text{4729} \\ & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\tan^{\frac{5}{2}}(c + dx)(A + B \tan(c + dx))}{(i \tan(c + dx)a + a)^{5/2}} dx \end{aligned}$$

3.566. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\tan(c+dx)^{5/2}(A+B \tan(c+dx))}{(i \tan(c+dx)a+a)^{5/2}} dx \\
 & \downarrow 4078 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA) \tan^{\frac{5}{2}}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{5 \tan^{\frac{3}{2}}(c+dx)(a(iA-B)+2iaB \tan(c+dx))}{2(i \tan(c+dx)a+a)^{3/2}} dx}{5a^2} \right) \\
 & \downarrow 27 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA) \tan^{\frac{5}{2}}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\tan^{\frac{3}{2}}(c+dx)(a(iA-B)+2iaB \tan(c+dx))}{(i \tan(c+dx)a+a)^{3/2}} dx}{2a^2} \right) \\
 & \downarrow 3042 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA) \tan^{\frac{5}{2}}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\tan(c+dx)^{3/2}(a(iA-B)+2iaB \tan(c+dx))}{(i \tan(c+dx)a+a)^{3/2}} dx}{2a^2} \right) \\
 & \downarrow 4078 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA) \tan^{\frac{5}{2}}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int -\frac{3\sqrt{\tan(c+dx)}((A+3iB)a^2+4B \tan(c+dx)a^2)}{2\sqrt{i \tan(c+dx)a+a}} dx}{3a^2} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{5/2}} \right) \\
 & \downarrow 27 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA) \tan^{\frac{5}{2}}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\sqrt{\tan(c+dx)}((A+3iB)a^2+4B \tan(c+dx)a^2)}{\sqrt{i \tan(c+dx)a+a}} dx}{2a^2} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{5/2}} \right) \\
 & \downarrow 3042 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA) \tan^{\frac{5}{2}}(c+dx)}{5d(a+ia \tan(c+dx))^{5/2}} - \frac{\int \frac{\sqrt{\tan(c+dx)}((A+3iB)a^2+4B \tan(c+dx)a^2)}{\sqrt{i \tan(c+dx)a+a}} dx}{2a^2} - \frac{a(A+3iB) \tan^{\frac{3}{2}}(c+dx)}{3d(a+ia \tan(c+dx))^{5/2}} \right) \\
 & \downarrow 4078
 \end{aligned}$$

3.566. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{a^2(-7B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \int \frac{\sqrt{i\tan(c+dx)a+a((iA-7B)a^3+8iB\tan(c+dx))}}{2\sqrt{\tan(c+dx)}}}{2a^2}}{2a^2} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{a^2(-7B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \int \frac{\sqrt{i\tan(c+dx)a+a((iA-7B)a^3+8iB\tan(c+dx))}}{\sqrt{\tan(c+dx)}}}{2a^2}}{2a^2} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{a^2(-7B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \int \frac{\sqrt{i\tan(c+dx)a+a((iA-7B)a^3+8iB\tan(c+dx))}}{\sqrt{\tan(c+dx)}}}{2a^2}}{2a^2} \right)$$

↓ 4084

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{a^2(-7B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{a^3(B+iA)\int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - 8a^2B\int \frac{(a-ia)}{2a^2}}{2a^2}}{2a^2}}{2a^2} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{a^2(-7B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{a^3(B+iA)\int \frac{\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}} dx - 8a^2B\int \frac{(a-ia)}{2a^2}}{2a^2}}{2a^2}}{2a^2} \right)$$

↓ 4027

3.566. $\int \frac{A+B\tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{a^2(-7B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{-8a^2B\int\frac{(a-ia\tan(c+dx))\sqrt{i\tan(c+dx)a+a}}{\sqrt{\tan(c+dx)}}dx}{2a^2}}{2a^2} \right)$$

↓ 218

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{a^2(-7B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{(1-i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}}{2a^2}}{2a^2} \right)$$

↓ 4082

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{a^2(-7B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{(1-i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}}{2a^2}}{2a^2} \right)$$

↓ 65

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{a^2(-7B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{(1-i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}}{2a^2}}{2a^2} \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA)\tan^{\frac{5}{2}}(c+dx)}{5d(a+ia\tan(c+dx))^{5/2}} - \frac{\frac{a^2(-7B+iA)\sqrt{\tan(c+dx)}}{d\sqrt{a+ia\tan(c+dx)}} - \frac{(1-i)a^{7/2}(B+iA)\operatorname{arctanh}\left(\frac{(1+i)\sqrt{a}\sqrt{\tan(c+dx)}}{\sqrt{a+ia\tan(c+dx)}}\right)}{d}}{2a^2}}{2a^2} \right)$$

3.566. $\int \frac{A+B\tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^{5/2}} dx$

input `Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^(5/2)),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((I*A - B)*Tan[c + d*x]^(5/2))/(5*d*(a + I*a*Tan[c + d*x])^(5/2)) - (-1/3*(a*(A + (3*I)*B)*Tan[c + d*x]^(3/2)))/(d*(a + I*a*Tan[c + d*x])^(3/2)) + (-1/2*((16*(-1)^(1/4)*a^(7/2)*B*ArcTan[((-1)^(3/4)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d + ((1 - I)*a^(7/2)*(I*A + B)*ArcTanh[((1 + I)*Sqrt[a]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*a*Tan[c + d*x]]])/d)/a^2 + (a^2*(I*A - 7*B)*Sqrt[Tan[c + d*x]])/(d*Sqrt[a + I*a*Tan[c + d*x]])/(2*a^2))/(2*a^2)`

3.566.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4027 `Int[Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]]/Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[-2*a*(b/f) Subst[Int[1/(a*c - b*d - 2*a^2*x^2), x], x, Sqrt[c + d*Tan[e + f*x]]/Sqrt[a + b*Tan[e + f*x]]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

3.566. $\int \frac{A+B \tan (c+d x)}{\cot ^{\frac{5}{2}}(c+d x)(a+i a \tan (c+d x))^{5 / 2}} d x$

```
rule 4078 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(-A*b - a*B)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(2*a*f*m)),
x] + Simp[1/(2*a^2*m) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*
x])^(n - 1)*Simp[A*(a*c*m + b*d*n) - B*(b*c*m + a*d*n) - d*(b*B*(m - n) - a
*A*(m + n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0] && GtQ[n, 0]
```

```
rule 4082 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]],
x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[
a^2 + b^2, 0] && EqQ[A*b + a*B, 0]
```

```
rule 4084 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x]
- Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[
e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.566.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1551 vs. $2(230) = 460$.

Time = 0.55 (sec) , antiderivative size = 1552, normalized size of antiderivative = 5.37

method	result	size
derivativedivides	Expression too large to display	1552
default	Expression too large to display	1552

3.566.
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$$

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^(5/2),x,method=_R
ETURNVERBOSE)`

output
$$\begin{aligned} & -1/240/d/(1/\tan(d*x+c))^{(5/2)}/\tan(d*x+c)^2*(a*(1+I*\tan(d*x+c)))^{(1/2)}/a^3* \\ & (240*B*(-I*a)^{(1/2)}*\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+ \\ & c)))^{(1/2)}*(I*a)^{(1/2)+a}/(I*a)^{(1/2)}*a-15*A*(I*a)^{(1/2)}*\ln(-(-2*2^{(1/2)}* \\ & (-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)+I*a-3*a*\tan(d*x+c))/(\tan \\ & (d*x+c)+I))*2^{(1/2)}*a+60*I*A*(I*a)^{(1/2)}*\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}*(a* \\ & \tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)+I*a-3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*2^{(\\ & 1/2)}*a*\tan(d*x+c)^3-420*B*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*(I*a)^{(1/2)} \\ &)*(-I*a)^{(1/2)}+240*B*\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x \\ & +c)))^{(1/2)}*(I*a)^{(1/2)+a}/(I*a)^{(1/2)}*(-I*a)^{(1/2)}*a*\tan(d*x+c)^4-1440*B \\ & *\ln(1/2*(2*I*a*\tan(d*x+c)+2*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*(I*a)^{(1 \\ & /2)+a}/(I*a)^{(1/2)}*(-I*a)^{(1/2)}*a*\tan(d*x+c)^2-220*A*(I*a)^{(1/2)}*(-I*a)^{(\\ & 1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)+148*A*(I*a)^{(1/2)}*(- \\ & I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)^3+1548*B*(I*a) \\ & ^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)}*\tan(d*x+c)^2-15* \\ & A*(I*a)^{(1/2)}*\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c))) \\ & ^{(1/2)+I*a-3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*2^{(1/2)}*a*\tan(d*x+c)^4+60*B*(I* \\ & a)^{(1/2)}*\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2 \\ &)+I*a-3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*2^{(1/2)}*a*\tan(d*x+c)^3+90*A*(I*a)^{(1 \\ & /2)}*\ln(-(-2*2^{(1/2)}*(-I*a)^{(1/2)}*(a*\tan(d*x+c)*(1+I*\tan(d*x+c)))^{(1/2)+I*a \\ & -3*a*\tan(d*x+c))/(\tan(d*x+c)+I))*2^{(1/2)}*a*\tan(d*x+c)^2-60*B*(I*a)^{(1/2)... \end{aligned}$$

3.566.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 799 vs. $2(217) = 434$.

Time = 0.27 (sec) , antiderivative size = 799, normalized size of antiderivative = 2.76

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx =$$

$$\left(15 \sqrt{\frac{1}{2}} a^3 d \sqrt{\frac{i A^2 + 2 AB - i B^2}{a^5 d^2}} e^{(5i dx + 5i c)} \log \left(- \frac{4 \left(\sqrt{2} \sqrt{\frac{1}{2}} (i a^3 d e^{(2i dx + 2i c)} - i a^3 d) \sqrt{\frac{a}{e^{(2i dx + 2i c)} + 1}} \sqrt{\frac{i e^{(2i dx + 2i c)} + i}{e^{(2i dx + 2i c)} - 1}} \sqrt{\frac{i A^2 + 2 AB}{a^5 d}}}{i A + B} \right) \right)$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^(5/2),x, al
gorithm="fricas")`

$$3.566. \quad \int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$$

output

```
-1/120*(15*sqrt(1/2)*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(I*a^3*d*e^(2*I*d*x + 2*I*c) - I*a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2)) + (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) - 15*sqrt(1/2)*a^3*d*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-4*(sqrt(2)*sqrt(1/2)*(-I*a^3*d*e^(2*I*d*x + 2*I*c) + I*a^3*d)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt((I*A^2 + 2*A*B - I*B^2)/(a^5*d^2)) + (A - I*B)*a*e^(I*d*x + I*c))*e^(-I*d*x - I*c)/(I*A + B)) + 30*a^3*d*sqrt(-4*I*B^2/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-16*(3*B*a^2*e^(2*I*d*x + 2*I*c) - B*a^2 + sqrt(2)*(a^4*d*e^(3*I*d*x + 3*I*c) - a^4*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt(-4*I*B^2/(a^5*d^2)))*e^(-2*I*d*x - 2*I*c)/B) - 30*a^3*d*sqrt(-4*I*B^2/(a^5*d^2))*e^(5*I*d*x + 5*I*c)*log(-16*(3*B*a^2*e^(2*I*d*x + 2*I*c) - B*a^2 - sqrt(2)*(a^4*d*e^(3*I*d*x + 3*I*c) - a^4*d*e^(I*d*x + I*c))*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))*sqrt(-4*I*B^2/(a^5*d^2)))*e^(-2*I*d*x - 2*I*c)/B) + sqrt(2)*((23*A + 123*I*B)*e^(6*I*d*x + 6*I*c) - 2*(17*A + 72*I*B)*e^(4*I*d*x + 4*I*c) + 2*(7*A + 12*I*B))*e^(2*I*d*x + 2*I*c) - 3*A - 3*I*B)*sqrt(a/(e^(2*I*d*x + 2*I*c) + 1))*...
```

3.566.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(5/2)/(a+I*a*tan(d*x+c))**(5/2),x)`

output `Timed out`

3.566.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.
```

3.566.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{B \tan(dx + c) + A}{(ia \tan(dx + c) + a)^{\frac{5}{2}} \cot(dx + c)^{\frac{5}{2}}} dx$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+I*a*tan(d*x+c))^(5/2),x, algorithm="giac")
```

```
output integrate((B*tan(d*x + c) + A)/((I*a*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^(5/2)), x)
```

3.566.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{\frac{5}{2}}(a + a \tan(c + dx) li)^{\frac{5}{2}}} dx$$

```
input int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + a*tan(c + d*x)*li)^(5/2)),x)
```

```
output int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + a*tan(c + d*x)*li)^(5/2)), x)
```

3.566. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^{5/2}} dx$

3.567 $\int \cot^m(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.567.1 Optimal result	5349
3.567.2 Mathematica [F]	5349
3.567.3 Rubi [A] (verified)	5350
3.567.4 Maple [F]	5353
3.567.5 Fricas [F]	5354
3.567.6 Sympy [F]	5354
3.567.7 Maxima [F]	5354
3.567.8 Giac [F]	5355
3.567.9 Mupad [F(-1)]	5355

3.567.1 Optimal result

Integrand size = 34, antiderivative size = 179

$$\int \cot^m(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \frac{(A-iB) \operatorname{AppellF1}(1-m, 1-n, 1, 2-m, -i \tan(c+dx), i \tan(c+dx)) \cot^{-1+m}(c+dx)(1+i \tan(c+dx))}{d(1-m)} + \frac{iB \cot^{-1+m}(c+dx) \operatorname{Hypergeometric2F1}(1-m, 1-n, 2-m, -i \tan(c+dx))(1+i \tan(c+dx))^{-n}(a+i \tan(c+dx))^n}{d(1-m)}$$

```
output (A-I*B)*AppellF1(1-m,1-n,1,2-m,-I*tan(d*x+c),I*tan(d*x+c))*cot(d*x+c)^(-1+m)*(a+I*a*tan(d*x+c))^n/d/(1-m)/((1+I*tan(d*x+c))^n)+I*B*cot(d*x+c)^(-1+m)*hypergeom([1-n, 1-m],[2-m],[-I*tan(d*x+c)]*(a+I*a*tan(d*x+c))^n/d/(1-m)/((1+I*tan(d*x+c))^n)
```

3.567.2 Mathematica [F]

$$\int \cot^m(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \int \cot^m(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

input `Integrate[Cot[c + d*x]^m*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `Integrate[Cot[c + d*x]^m*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

3.567.3 Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.12, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.382$, Rules used = {3042, 4729, 3042, 4084, 3042, 4047, 25, 27, 152, 150, 4082, 76, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \cot(c + dx)^m(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4729} \\
 & \tan^m(c + dx) \cot^m(c + dx) \int \tan^{-m}(c + dx)(i \tan(c + dx)a + a)^n(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \tan^m(c + dx) \cot^m(c + dx) \int \tan(c + dx)^{-m}(i \tan(c + dx)a + a)^n(A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4084} \\
 & dx \left((A - iB) \int \tan^{-m}(c + dx)(i \tan(c + dx)a + a)^n dx + \frac{iB \int \tan^{-m}(c + dx)(a - ia \tan(c + dx))(i \tan(c + dx))}{a} \right) \\
 & \quad \downarrow \text{3042} \\
 & dx \left((A - iB) \int \tan(c + dx)^{-m}(i \tan(c + dx)a + a)^n dx + \frac{iB \int \tan(c + dx)^{-m}(a - ia \tan(c + dx))(i \tan(c + dx))}{a} \right) \\
 & \quad \downarrow \text{4047} \\
 & dx \left(\frac{\tan^m(c + dx) \cot^m(c + dx) \left(ia^2(A - iB) \int -\frac{\tan^{-m}(c+dx)(i \tan(c+dx)a+a)^{n-1}}{a(a-ia \tan(c+dx))} d(ia \tan(c + dx)) \right)}{d} + \frac{iB \int \tan(c + dx)^{-m}(a - ia \tan(c + dx))(i \tan(c + dx))}{a} \right)
 \end{aligned}$$

3.567. $\int \cot^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \downarrow 25 \\
& \tan^m(c+dx) \cot^m(c+dx) \\
dx) \left(\frac{iB \int \tan(c+dx)^{-m} (a - ia \tan(c+dx)) (i \tan(c+dx)a + a)^n dx}{a} - \frac{ia^2(A - iB) \int \frac{\tan^{-m}(c+dx)(i \tan(c+dx)a + a)^n}{a(ia \tan(c+dx) - a)}}{d} \right) \\
& \downarrow 27 \\
& \tan^m(c+dx) \cot^m(c+dx) \\
dx) \left(\frac{iB \int \tan(c+dx)^{-m} (a - ia \tan(c+dx)) (i \tan(c+dx)a + a)^n dx}{a} - \frac{ia(A - iB) \int \frac{\tan^{-m}(c+dx)(i \tan(c+dx)a + a)^n}{a - ia \tan(c+dx)}}{d} \right) \\
& \downarrow 152 \\
& \tan^m(c+dx) \cot^m(c+dx) \\
dx) \left(\frac{iB \int \tan(c+dx)^{-m} (a - ia \tan(c+dx)) (i \tan(c+dx)a + a)^n dx}{a} - \frac{i(A - iB)(1 + i \tan(c+dx))^{-n} (a + ia \tan(c+dx))^n}{d} \right) \\
& \downarrow 150 \\
& \tan^m(c+dx) \cot^m(c+dx) \\
dx) \left(\frac{iB \int \tan(c+dx)^{-m} (a - ia \tan(c+dx)) (i \tan(c+dx)a + a)^n dx}{a} + \frac{(A - iB) \tan^{1-m}(c+dx) (1 + i \tan(c+dx))^n}{d} \right) \\
& \downarrow 4082 \\
& \tan^m(c+dx) \cot^m(c+dx) \\
dx) \left(\frac{iaB \int \tan^{-m}(c+dx) (i \tan(c+dx)a + a)^{n-1} d \tan(c+dx)}{d} + \frac{(A - iB) \tan^{1-m}(c+dx) (1 + i \tan(c+dx))^{-n}}{d} \right) \\
& \downarrow 76 \\
& \tan^m(c+dx) \cot^m(c+dx) \\
dx) \left(\frac{iB(1 + i \tan(c+dx))^{-n} (a + ia \tan(c+dx))^n \int (i \tan(c+dx) + 1)^{n-1} \tan^{-m}(c+dx) d \tan(c+dx)}{d} + \frac{(A - iB) \tan^{1-m}(c+dx) (1 + i \tan(c+dx))^{-n}}{d} \right) \\
& \downarrow 74 \\
& \tan^m(c+dx) \cot^m(c+dx) \\
dx) \left(\frac{(A - iB) \tan^{1-m}(c+dx) (1 + i \tan(c+dx))^{-n} (a + ia \tan(c+dx))^n \operatorname{AppellF1}(1 - m, 1 - n, 1, 2 - m, -i \tan(c+dx))}{d(1 - m)} \right)
\end{aligned}$$

input `Int[Cot[c + d*x]^m*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

```
output Cot[c + d*x]^m*Tan[c + d*x]^m*((A - I*B)*AppellF1[1 - m, 1 - n, 1, 2 - m,
(-I)*Tan[c + d*x], I*Tan[c + d*x]]*Tan[c + d*x]^(1 - m)*(a + I*a*Tan[c +
d*x])^n)/(d*(1 - m)*(1 + I*Tan[c + d*x])^n) + (I*B*Hypergeometric2F1[1 - m
, 1 - n, 2 - m, (-I)*Tan[c + d*x]]*Tan[c + d*x]^(1 - m)*(a + I*a*Tan[c + d
*x])^n)/(d*(1 - m)*(1 + I*Tan[c + d*x])^n))
```

3.567.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 74 Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x
)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x]
/; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0]
&& !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))
```

```
rule 76 Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[c^IntPart
[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*
(x/c))^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !Integer
Q[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0] && ((RationalQ[m] && !(EqQ[n, -2
^(-1)] && EqQ[c^2 - d^2, 0])) || !RationalQ[n])
```

```
rule 150 Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_
] := Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2
, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !In
tegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])
```

```
rule 152 Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_
] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n])
Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x], x] /; FreeQ[{b, c, d, e, f, m,
n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]
```

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4047 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

rule 4729 `Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.567.4 Maple [F]

$$\int \cot(dx + c)^m (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(cot(d*x+c)^m*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(cot(d*x+c)^m*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.567.5 Fracas [F]

$$\int \cot^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^m dx$$

input `integrate(cot(d*x+c)^m*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(((A - I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*(2*a*e^(2*I*d*x + 2*I*c))/(e^(2*I*d*x + 2*I*c) + 1))^n*((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))^m/(e^(2*I*d*x + 2*I*c) + 1), x)`

3.567.6 Sympy [F]

$$\int \cot^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (ia(\tan(c + dx) - i))^n(A + B \tan(c + dx)) \cot^m(c + dx) dx$$

input `integrate(cot(d*x+c)**m*(a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x))*cot(c + d*x)**m, x)`

3.567.7 Maxima [F]

$$\int \cot^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^m dx$$

input `integrate(cot(d*x+c)^m*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*cot(d*x + c)^m, x)`

3.567.8 Giac [F]

$$\int \cot^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^m dx$$

input `integrate(cot(d*x+c)^m*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*cot(d*x + c)^m, x)`

3.567.9 Mupad [F(-1)]

Timed out.

$$\int \cot^m(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^m (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^n dx$$

input `int(cot(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n,x)`

output `int(cot(c + d*x)^m*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n, x)`

3.568 $\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.568.1 Optimal result	5356
3.568.2 Mathematica [F]	5357
3.568.3 Rubi [A] (warning: unable to verify)	5357
3.568.4 Maple [F]	5363
3.568.5 Fricas [F]	5364
3.568.6 Sympy [F(-1)]	5364
3.568.7 Maxima [F]	5364
3.568.8 Giac [F]	5365
3.568.9 Mupad [F(-1)]	5365

3.568.1 Optimal result

Integrand size = 36, antiderivative size = 247

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= -\frac{2(3B+2iAn)\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^n}{3d} - \frac{2A \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{3d} - \frac{2(A-iB) \operatorname{AppellF1}\left(\frac{1}{2}, 1-n, 1, \frac{3}{2}, -i \tan(c+dx), i \tan(c+dx)\right) (1+i \tan(c+dx))^{-n} (a+ia \tan(c+dx))^n}{d\sqrt{\cot(c+dx)}} - \frac{2(1-2n)(3iB-2An) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1-n, \frac{3}{2}, -i \tan(c+dx)\right) (1+i \tan(c+dx))^{-n} (a+ia \tan(c+dx))^n}{3d\sqrt{\cot(c+dx)}}$$

```
output -2/3*A*cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^n/d-2/3*(3*B+2*I*A*n)*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n/d-2*(A-I*B)*AppellF1(1/2,1-n,1,3/2,-I*tan(d*x+c),I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/cot(d*x+c)^(1/2)/((1+I*tan(d*x+c))^n)-2/3*(1-2*n)*(3*I*B-2*A*n)*hypergeom([1/2, 1-n],[3/2],-I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/cot(d*x+c)^(1/2)/((1+I*tan(d*x+c))^n)
```

3.568.2 Mathematica [F]

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

input `Integrate[Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `Integrate[Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

3.568.3 Rubi [A] (warning: unable to verify)

Time = 1.52 (sec) , antiderivative size = 301, normalized size of antiderivative = 1.22, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3042, 4729, 3042, 4081, 27, 3042, 4081, 27, 3042, 4084, 3042, 4047, 25, 27, 148, 27, 334, 333, 4082, 76, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{5/2}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^n(A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^n(A+B \tan(c+dx))}{\tan(c+dx)^{5/2}} dx$$

$$\downarrow \text{4081}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2\int\frac{(i\tan(c+dx)a+a)^n(a(3B+2iAn)-aA(3-2n)\tan(c+dx))}{2\tan^{\frac{3}{2}}(c+dx)}dx}{3a}-\frac{2A(a+ia\tan(c+dx))^n}{3d\tan^{\frac{3}{2}}(c+dx)}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{\int\frac{(i\tan(c+dx)a+a)^n(a(3B+2iAn)-aA(3-2n)\tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)}dx}{3a}-\frac{2A(a+ia\tan(c+dx))^n}{3d\tan^{\frac{3}{2}}(c+dx)}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{\int\frac{(i\tan(c+dx)a+a)^n(a(3B+2iAn)-aA(3-2n)\tan(c+dx))}{\tan(c+dx)^{3/2}}dx}{3a}-\frac{2A(a+ia\tan(c+dx))^n}{3d\tan^{\frac{3}{2}}(c+dx)}\right)$$

↓ 4081

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2\int\frac{(i\tan(c+dx)a+a)^n\left(a^2\left(6iBn-A(4n^2-2n+3)\right)-a^2(1-2n)(3B+2iAn)\tan(c+dx)\right)}{2\sqrt{\tan(c+dx)}}dx}{3a}-\frac{2a(3B+2iAn)(a+ia\tan(c+dx))}{d\sqrt{\tan(c+dx)}}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{\int\frac{(i\tan(c+dx)a+a)^n\left(a^2\left(6iBn-A(4n^2-2n+3)\right)-a^2(1-2n)(3B+2iAn)\tan(c+dx)\right)}{\sqrt{\tan(c+dx)}}dx}{3a}-\frac{2a(3B+2iAn)(a+ia\tan(c+dx))}{d\sqrt{\tan(c+dx)}}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{\int\frac{(i\tan(c+dx)a+a)^n\left(a^2\left(6iBn-A(4n^2-2n+3)\right)-a^2(1-2n)(3B+2iAn)\tan(c+dx)\right)}{\sqrt{\tan(c+dx)}}dx}{3a}-\frac{2a(3B+2iAn)(a+ia\tan(c+dx))}{d\sqrt{\tan(c+dx)}}\right)$$

↓ 4084

3.568. $\int \cot^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^n(A+B\tan(c+dx))dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{-3a^2(A-iB) \int \frac{(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{a} - \frac{2a(\dots)}{3a} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{-3a^2(A-iB) \int \frac{(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{a} - \frac{2a(\dots)}{3a} \right)$$

↓ 4047

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{3ia^4(A-iB) \int -\frac{(i \tan(c+dx)a+a)^{n-1}}{a \sqrt{\tan(c+dx)}(a-ia \tan(c+dx))} d(ia \tan(c+dx))}{a} - a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{3a} \right)$$

↓ 25

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{3ia^4(A-iB) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{a \sqrt{\tan(c+dx)}(a-ia \tan(c+dx))} d(ia \tan(c+dx))}{a} - a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{3a} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{3ia^3(A-iB) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)}(a-ia \tan(c+dx))} d(ia \tan(c+dx))}{a} - a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{3a} \right)$$

↓ 148

3.568. $\int \cot^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{6a^4(A-iB) \int \frac{(a-ia^3 \tan^2(c+dx))^{n-1}}{a(ia^2 \tan^2(c+dx)+1)} d\sqrt{\tan(c+dx)}}{d} - \frac{a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)}{\sqrt{\tan(c+dx)}}}{a} \right) \frac{3a}{3a}$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{6a^3(A-iB) \int \frac{(a-ia^3 \tan^2(c+dx))^{n-1}}{ia^2 \tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} - \frac{a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)}{\sqrt{\tan(c+dx)}}}{a} \right) \frac{3a}{3a}$$

↓ 334

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{6a^2(A-iB)(a-ia^3 \tan^2(c+dx))^n (1-ia^2 \tan^2(c+dx))^{-n} \int \frac{(1-ia^2 \tan^2(c+dx))^{n-1}}{ia^2 \tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} - \frac{a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}}}{a} \right) \frac{3a}{3a}$$

↓ 333

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{a(1-2n)(-2An+3iB) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx - \frac{6ia^3(A-iB) \tan(c+dx)(a-ia^3 \tan^2(c+dx))^n}{a} \right) \frac{3a}{3a}$$

↓ 4082

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{a^3(1-2n)(-2An+3iB) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)}} d \tan(c+dx)}{d} - \frac{6ia^3(A-iB) \tan(c+dx)(a-ia^3 \tan^2(c+dx))^n (1-ia^2 \tan^2(c+dx))^{-n}}{a} \right) \frac{3a}{3a}$$

↓ 76

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{a^2(1-2n)(-2An+3iB)(1+i\tan(c+dx))^{-n}(a+ia\tan(c+dx))^n \int \frac{(i\tan(c+dx)+1)^{n-1} d\tan(c+dx)}{\sqrt{\tan(c+dx)}} - 6ia^3(A-iB)\tan(c+dx)}{d} \right)$$

↓ 74

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a^2(1-2n)(-2An+3iB)\sqrt{\tan(c+dx)}(1+i\tan(c+dx))^{-n}(a+ia\tan(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1-n, \frac{3}{2}, -i\tan(c+dx)\right)}{d} \right)$$

```
input Int[Cot[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A*(a + I*a*Tan[c + d*x])^n)/(3*d*Tan[c + d*x]^(3/2)) + ((-2*a*(3*B + (2*I)*A*n)*(a + I*a*Tan[c + d*x])^n)/(d*Sqrt[Tan[c + d*x]])) + ((-2*a^2*(1 - 2*n)*((3*I)*B - 2*A*n)*Hypergeometric2F1[1/2, 1 - n, 3/2, (-I)*Tan[c + d*x]]*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + I*Tan[c + d*x])^n) - ((6*I)*a^3*(A - I*B)*AppellF1[1/2, 1, 1 - n, 3/2, (-I)*a^2*Tan[c + d*x]^2, I*a^2*Tan[c + d*x]^2]*Tan[c + d*x]*(a - I*a^3*Tan[c + d*x]^2)^n)/(d*(1 - I*a^2*Tan[c + d*x]^2)^n))/a)/(3*a))
```

3.568.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 74 Int[((b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))
```

3.568. $\int \cot^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^n(A+B\tan(c+dx))dx$

- rule 76 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0] && ((RationalQ[m] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0])) || !RationalQ[n])`
- rule 148 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_.))*((e_) + (f_.)*(x_)^(p_.)), x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c + d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x]] /; FreeQ[{b, c, d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`
- rule 333 `Int[((a_) + (b_.)*(x_)^2)^(p_))*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`
- rule 334 `Int[((a_) + (b_.)*(x_)^2)^(p_))*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4047 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4081 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2)), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

rule 4729 `Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.568.4 Maple [F]

$$\int \cot(dx + c)^{\frac{5}{2}} (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.568. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$

3.568.5 Fricas [F]

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^{\frac{5}{2}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral(-(A - I*B)*e^(4*I*d*x + 4*I*c) + 2*A*e^(2*I*d*x + 2*I*c) + A + I*B)*(2*a*e^(2*I*d*x + 2*I*c)/(e^(2*I*d*x + 2*I*c) + 1))^n*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))/(e^(4*I*d*x + 4*I*c) - 2*e^(2*I*d*x + 2*I*c) + 1), x)`

3.568.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(5/2)*(a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.568.7 Maxima [F]

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^{\frac{5}{2}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*cot(d*x + c)^(5/2), x)`

3.568. $\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$

3.568.8 Giac [F]

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^{\frac{5}{2}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*cot(d*x + c)^(5/2), x)`

3.568.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{\frac{5}{2}} (A + B \tan(c + dx)) (a + a \tan(c + dx) i)^n dx$$

input `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n,x)`

output `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n, x)`

3.569 $\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.569.1 Optimal result	5366
3.569.2 Mathematica [F]	5367
3.569.3 Rubi [A] (warning: unable to verify)	5367
3.569.4 Maple [F]	5372
3.569.5 Fricas [F]	5372
3.569.6 Sympy [F(-1)]	5373
3.569.7 Maxima [F]	5373
3.569.8 Giac [F]	5374
3.569.9 Mupad [F(-1)]	5374

3.569.1 Optimal result

Integrand size = 36, antiderivative size = 194

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= -\frac{2A\sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^n}{d} + \frac{2(iA+B) \operatorname{AppellF1}\left(\frac{1}{2}, 1-n, 1, \frac{3}{2}, -i \tan(c+dx), i \tan(c+dx)\right) (1+i \tan(c+dx))^{-n}(a+ia \tan(c+dx))}{d\sqrt{\cot(c+dx)}} - \frac{2iA(1-2n) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1-n, \frac{3}{2}, -i \tan(c+dx)\right) (1+i \tan(c+dx))^{-n}(a+ia \tan(c+dx))}{d\sqrt{\cot(c+dx)}}$$

```
output -2*A*cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n/d+2*(I*A+B)*AppellF1(1/2,1-n,1,
3/2,-I*tan(d*x+c),I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/cot(d*x+c)^(1/2)/((
1+I*tan(d*x+c))^n)-2*I*A*(1-2*n)*hypergeom([1/2, 1-n],[3/2],-I*tan(d*x+c))
*(a+I*a*tan(d*x+c))^n/d/cot(d*x+c)^(1/2)/((1+I*tan(d*x+c))^n)
```

3.569.2 Mathematica [F]

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

input `Integrate[Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

output `Integrate[Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

3.569.3 Rubi [A] (warning: unable to verify)

Time = 1.11 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.24, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4729, 3042, 4081, 27, 3042, 4084, 3042, 4047, 25, 27, 148, 27, 334, 333, 4082, 76, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{3/2}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^n(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^n(A+B \tan(c+dx))}{\tan(c+dx)^{3/2}} dx$$

$$\downarrow \text{4081}$$

3.569. $\int \cot^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2\int\frac{(i\tan(c+dx)a+a)^n(a(B+2iAn)-aA(1-2n)\tan(c+dx))dx}{2\sqrt{\tan(c+dx)}}}{a}-\frac{2A(a+ia\tan(c+dx))^n}{d\sqrt{\tan(c+dx)}}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{\int\frac{(i\tan(c+dx)a+a)^n(a(B+2iAn)-aA(1-2n)\tan(c+dx))dx}{\sqrt{\tan(c+dx)}}}{a}-\frac{2A(a+ia\tan(c+dx))^n}{d\sqrt{\tan(c+dx)}}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{\int\frac{(i\tan(c+dx)a+a)^n(a(B+2iAn)-aA(1-2n)\tan(c+dx))dx}{\sqrt{\tan(c+dx)}}}{a}-\frac{2A(a+ia\tan(c+dx))^n}{d\sqrt{\tan(c+dx)}}\right)$$

↓ 4084

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{a(B+iA)\int\frac{(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}}dx-iA(1-2n)\int\frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}}dx}{a}-\frac{2A(a+ia\tan(c+dx))^n}{d\sqrt{\tan(c+dx)}}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{a(B+iA)\int\frac{(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}}dx-iA(1-2n)\int\frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}}dx}{a}-\frac{2A(a+ia\tan(c+dx))^n}{d\sqrt{\tan(c+dx)}}\right)$$

↓ 4047

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{ia^3(B+iA)\int-\frac{(i\tan(c+dx)a+a)^{n-1}}{a\sqrt{\tan(c+dx)}(a-ia\tan(c+dx))}d(ia\tan(c+dx))}{d}-iA(1-2n)\int\frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}}dx}{a}-\frac{2A(a+ia\tan(c+dx))^n}{d\sqrt{\tan(c+dx)}}\right)$$

↓ 25

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{-\frac{ia^3(B+iA)\int\frac{(i\tan(c+dx)a+a)^{n-1}}{a\sqrt{\tan(c+dx)}(a-ia\tan(c+dx))}d(ia\tan(c+dx))}{d}-iA(1-2n)\int\frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}}dx}{a}-\frac{2A(a+ia\tan(c+dx))^n}{d\sqrt{\tan(c+dx)}}\right)$$

↓ 27

3.569. $\int \cot^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n(A+B\tan(c+dx))dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{-\frac{ia^2(B+iA) \int \frac{(i \tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)}(a-ia \tan(c+dx))} d(ia \tan(c+dx))}{d} - iA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)}{\sqrt{\tan(c+dx)}}}{a} \right)$$

↓ 148

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a^3(B+iA) \int \frac{(a-ia^3 \tan^2(c+dx))^{n-1}}{a(ia^2 \tan^2(c+dx)+1)} d\sqrt{\tan(c+dx)}}{d} - iA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)}{\sqrt{\tan(c+dx)}}}{a} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a^2(B+iA) \int \frac{(a-ia^3 \tan^2(c+dx))^{n-1}}{ia^2 \tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} - iA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)}{\sqrt{\tan(c+dx)}}}{a} \right)$$

↓ 334

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(B+iA)(1-ia^2 \tan^2(c+dx))^{-n}(a-ia^3 \tan^2(c+dx))^n \int \frac{(1-ia^2 \tan^2(c+dx))^{n-1}}{ia^2 \tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} - iA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)}{\sqrt{\tan(c+dx)}}}{a} \right)$$

↓ 333

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2ia^2(B+iA) \tan(c+dx)(1-ia^2 \tan^2(c+dx))^{-n}(a-ia^3 \tan^2(c+dx))^n \operatorname{AppellF1}(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2 \tan^2(c+dx))}{d} - iA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)}{\sqrt{\tan(c+dx)}}}{a} \right)$$

↓ 4082

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2ia^2(B+iA) \tan(c+dx)(1-ia^2 \tan^2(c+dx))^{-n}(a-ia^3 \tan^2(c+dx))^n \operatorname{AppellF1}(\frac{1}{2}, 1, 1-n, \frac{3}{2}, -ia^2 \tan^2(c+dx))}{d} - iA(1-2n) \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)}{\sqrt{\tan(c+dx)}}}{a} \right)$$

↓ 76

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2ia^2(B+IA)\tan(c+dx)(1-ia^2\tan^2(c+dx))^{-n}(a-ia^3\tan^2(c+dx))^n\text{AppellF1}\left(\frac{1}{2},1,1-n,\frac{3}{2},-ia^2\tan^2(c+dx)\right)}{d}\right)$$

↓ 74

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2ia^2(B+IA)\tan(c+dx)(1-ia^2\tan^2(c+dx))^{-n}(a-ia^3\tan^2(c+dx))^n\text{AppellF1}\left(\frac{1}{2},1,1-n,\frac{3}{2},-ia^2\tan^2(c+dx)\right)}{d}\right)$$

input `Int[Cot[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A*(a + I*a*Tan[c + d*x])^n)/(d*Sqrt[Tan[c + d*x]]) + (((-2*I)*a*A*(1 - 2*n)*Hypergeometric2F1[1/2, 1 - n, 3/2, (-I)*Tan[c + d*x]]*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + I*Tan[c + d*x])^n) + ((2*I)*a^2*(I*A + B)*AppellF1[1/2, 1, 1 - n, 3/2, (-I)*a^2*Tan[c + d*x]^2, I*a^2*Tan[c + d*x]^2]*Tan[c + d*x]*(a - I*a^3*Tan[c + d*x]^2)^n)/(d*(1 - I*a^2*Tan[c + d*x]^2)^n))/a`

3.569.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 74 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0]))`

rule 76 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^(m*(1 + d*(x/c))^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0] && ((RationalQ[m] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0])) || !RationalQ[n])`

3.569. $\int \cot^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n(A+B\tan(c+dx))dx$

- rule 148 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_.))*((e_) + (f_.)*(x_)^(p_.)), x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c + d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x] /; FreeQ[{b, c, d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`
- rule 333 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`
- rule 334 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4047 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4081 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*d - B*c)*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(a*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*(b*d*m - a*c*(n + 1)) - B*(b*c*m + a*d*(n + 1)) - a*(B*c - A*d)*(m + n + 1)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[n, -1]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

rule 4729 `Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.569.4 Maple [F]

$$\int \cot(dx + c)^{\frac{3}{2}} (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.569.5 Fracas [F]

$$\begin{aligned} & \int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fracas")`

3.569. $\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx$

output `integral(((I*A + B)*e^(2*I*d*x + 2*I*c) + I*A - B)*(2*a*e^(2*I*d*x + 2*I*c))/(e^(2*I*d*x + 2*I*c) + 1))^n*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))/(e^(2*I*d*x + 2*I*c) - 1), x)`

3.569.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(3/2)*(a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output Timed out

3.569.7 Maxima [F]

$$\begin{aligned} & \int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*cot(d*x + c)^(3/2), x)`

3.569.8 Giac [F]

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n \cot(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*cot(d*x + c)^(3/2), x)`

3.569.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{3/2} (A + B \tan(c + dx)) (a + a \tan(c + dx) li)^n dx$$

input `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n,x)`

output `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n, x)`

3.570 $\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$

3.570.1 Optimal result	5375
3.570.2 Mathematica [F]	5375
3.570.3 Rubi [A] (warning: unable to verify)	5376
3.570.4 Maple [F]	5380
3.570.5 Fricas [F]	5380
3.570.6 Sympy [F]	5381
3.570.7 Maxima [F]	5381
3.570.8 Giac [F]	5381
3.570.9 Mupad [F(-1)]	5382

3.570.1 Optimal result

Integrand size = 36, antiderivative size = 158

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \frac{2(A - iB) \operatorname{AppellF1}\left(\frac{1}{2}, 1 - n, 1, \frac{3}{2}, -i \tan(c + dx), i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n} (a + ia \tan(c + dx))^n}{d \sqrt{\cot(c + dx)}} + \frac{2iB \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1 - n, \frac{3}{2}, -i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n} (a + ia \tan(c + dx))^n}{d \sqrt{\cot(c + dx)}}$$

```
output 2*(A-I*B)*AppellF1(1/2,1-n,1,3/2,-I*tan(d*x+c),I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/cot(d*x+c)^(1/2)/((1+I*tan(d*x+c))^n)+2*I*B*hypergeom([1/2, 1-n],[3/2],-I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/cot(d*x+c)^(1/2)/((1+I*tan(d*x+c))^n)
```

3.570.2 Mathematica [F]

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

input `Integrate[Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `Integrate[Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

3.570.3 Rubi [A] (warning: unable to verify)

Time = 0.84 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.25, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3042, 4729, 3042, 4084, 3042, 4047, 25, 27, 148, 27, 334, 333, 4082, 76, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4729} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^n(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(i \tan(c+dx)a+a)^n(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx \\
 & \quad \downarrow \text{4084} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left((A-iB) \int \frac{(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx + \frac{iB \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{a} \right) \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left((A-iB) \int \frac{(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx + \frac{iB \int \frac{(a-ia \tan(c+dx))(i \tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{a} \right) \\
 & \quad \downarrow \text{4047}
 \end{aligned}$$

3.570. $\int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{ia^2(A-iB)\int-\frac{(i\tan(c+dx)a+a)^{n-1}}{a\sqrt{\tan(c+dx)}(a-ia\tan(c+dx))}d(ia\tan(c+dx))}{d}+\frac{iB\int\frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}}dx}{a}\right)$$

↓ 25

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{iB\int\frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}}dx}{a}-\frac{ia^2(A-iB)\int\frac{(i\tan(c+dx)a+a)^{n-1}}{a\sqrt{\tan(c+dx)}(a-ia\tan(c+dx))}d(ia\tan(c+dx))}{d}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{iB\int\frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}}dx}{a}-\frac{ia(A-iB)\int\frac{(i\tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)}(a-ia\tan(c+dx))}d(ia\tan(c+dx))}{d}\right)$$

↓ 148

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2a^2(A-iB)\int\frac{(a-ia^3\tan^2(c+dx))^{n-1}}{a(ia^2\tan^2(c+dx)+1)}d\sqrt{\tan(c+dx)}}{d}+\frac{iB\int\frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}}dx}{a}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2a(A-iB)\int\frac{(a-ia^3\tan^2(c+dx))^{n-1}}{ia^2\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}}{d}+\frac{iB\int\frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}}dx}{a}\right)$$

↓ 334

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2(A-iB)(a-ia^3\tan^2(c+dx))^n(1-ia^2\tan^2(c+dx))^{-n}\int\frac{(1-ia^2\tan^2(c+dx))^{n-1}}{ia^2\tan^2(c+dx)+1}d\sqrt{\tan(c+dx)}}{d}\right)$$

↓ 333

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{iB\int\frac{(a-ia\tan(c+dx))(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}}dx}{a}+\frac{2ia(A-iB)\tan(c+dx)(a-ia^3\tan^2(c+dx))^n}{a}\right)$$

↓ 4082

3.570. $\int \sqrt{\cot(c+dx)}(a+ia\tan(c+dx))^n(A+B\tan(c+dx))dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{iaB\int\frac{(i\tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)}}d\tan(c+dx)}{d}+\frac{2ia(A-iB)\tan(c+dx)(a-ia^3\tan^2(c+dx))}{d}\right)$$

↓ 76

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{iB(1+i\tan(c+dx))^{-n}(a+ia\tan(c+dx))^n\int\frac{(i\tan(c+dx)+1)^{n-1}}{\sqrt{\tan(c+dx)}}d\tan(c+dx)}{d}+\frac{2ia(A-iB)\tan(c+dx)(1-ia^2\tan^2(c+dx))^{-n}(a-ia^3\tan^2(c+dx))^n\text{AppellF1}}{d}\right)$$

↓ 74

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2ia(A-iB)\tan(c+dx)(1-ia^2\tan^2(c+dx))^{-n}(a-ia^3\tan^2(c+dx))^n\text{AppellF1}}{d}\right)$$

input `Int[Sqrt[Cot[c + d*x]]*(a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((2*I)*B*Hypergeometric2F1[1/2, 1 - n, 3/2, (-I)*Tan[c + d*x]]*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + I*Tan[c + d*x])^n) + ((2*I)*a*(A - I*B)*AppellF1[1/2, 1, 1 - n, 3/2, (-I)*a^2*Tan[c + d*x]^2, I*a^2*Tan[c + d*x]^2]*Tan[c + d*x]*(a - I*a^3*Tan[c + d*x]^2)^n)/(d*(1 - I*a^2*Tan[c + d*x]^2)^n))`

3.570.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 74 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)]) && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0]))`

- rule 76 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0] && ((RationalQ[m] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0])) || !RationalQ[n])`
- rule 148 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_.))*((e_) + (f_.)*(x_)^(p_.)), x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c + d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x]] /; FreeQ[{b, c, d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`
- rule 333 `Int[((a_) + (b_.)*(x_)^2)^(p_))*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`
- rule 334 `Int[((a_) + (b_.)*(x_)^2)^(p_))*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4047 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4082 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

rule 4729 `Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.570.4 Maple [F]

$$\int \sqrt{\cot(dx + c)} (a + ia \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.570.5 Fracas [F]

$$\begin{aligned} & \int \sqrt{\cot(c + dx)} (a + ia \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A) (ia \tan(dx + c) + a)^n \sqrt{\cot(dx + c)} dx \end{aligned}$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output `integral(((A - I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*(2*a*e^(2*I*d*x + 2*I*c))/(e^(2*I*d*x + 2*I*c) + 1))^n*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))/(e^(2*I*d*x + 2*I*c) + 1), x)`

3.570.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx \\ &= \int (ia(\tan(c+dx)-i))^n(A+B \tan(c+dx)) \sqrt{\cot(c+dx)} dx \end{aligned}$$

input `integrate(cot(d*x+c)**(1/2)*(a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x))*sqrt(cot(c + d*x)), x)`

3.570.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx \\ &= \int (B \tan(dx+c)+A)(ia \tan(dx+c)+a)^n \sqrt{\cot(dx+c)} dx \end{aligned}$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*sqrt(cot(d*x + c)), x)`

3.570.8 Giac [F]

$$\begin{aligned} & \int \sqrt{\cot(c+dx)}(a+ia \tan(c+dx))^n(A+B \tan(c+dx)) dx \\ &= \int (B \tan(dx+c)+A)(ia \tan(dx+c)+a)^n \sqrt{\cot(dx+c)} dx \end{aligned}$$

input `integrate(cot(d*x+c)^(1/2)*(a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n*sqrt(cot(d*x + c)), x)`

3.570.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cot(c + dx)}(a + ia \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \sqrt{\cot(c + dx)}(A + B \tan(c + dx)) (a + a \tan(c + dx) \text{ li})^n dx$$

input `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n,x)`output `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + a*tan(c + d*x)*li)^n, x)`

3.571
$$\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.571.1 Optimal result 5383
 3.571.2 Mathematica [F] 5383
 3.571.3 Rubi [A] (warning: unable to verify) 5384
 3.571.4 Maple [F] 5389
 3.571.5 Fricas [F] 5389
 3.571.6 Sympy [F] 5390
 3.571.7 Maxima [F] 5390
 3.571.8 Giac [F] 5391
 3.571.9 Mupad [F(-1)] 5391

3.571.1 Optimal result

Integrand size = 36, antiderivative size = 215

$$\int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \frac{2B(a + ia \tan(c + dx))^n}{d(1 + 2n)\sqrt{\cot(c + dx)}} - \frac{2(iA + B) \operatorname{AppellF1}\left(\frac{1}{2}, 1 - n, 1, \frac{3}{2}, -i \tan(c + dx), i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n}(a + ia \tan(c + dx))^n}{d\sqrt{\cot(c + dx)}} + \frac{2(2Bn + iA(1 + 2n)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1 - n, \frac{3}{2}, -i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n}(a + ia \tan(c + dx))^n}{d(1 + 2n)\sqrt{\cot(c + dx)}}$$

```
output 2*B*(a+I*a*tan(d*x+c))^n/d/(1+2*n)/cot(d*x+c)^(1/2)-2*(I*A+B)*AppellF1(1/2,1-n,1,3/2,-I*tan(d*x+c),I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/cot(d*x+c)^(1/2)/((1+I*tan(d*x+c))^n)+2*(2*B*n+I*A*(1+2*n))*hypergeom([1/2, 1-n],[3/2,-I*tan(d*x+c)]*(a+I*a*tan(d*x+c))^n/d/(1+2*n)/cot(d*x+c)^(1/2)/((1+I*tan(d*x+c))^n))
```

3.571.2 Mathematica [F]

$$\int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

input `Integrate[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `Integrate[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]], x]`

3.571.3 Rubi [A] (warning: unable to verify)

Time = 1.17 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.25, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3042, 4729, 3042, 4080, 27, 3042, 4084, 3042, 4047, 25, 27, 148, 27, 334, 333, 4082, 76, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

↓ 4729

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \sqrt{\tan(c + dx)} (i \tan(c + dx) a + a)^n (A + B \tan(c + dx)) dx$$

↓ 3042

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \sqrt{\tan(c + dx)} (i \tan(c + dx) a + a)^n (A + B \tan(c + dx)) dx$$

↓ 4080

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{2 \int -\frac{(i \tan(c + dx) a + a)^n (a B - a(2nA + A - 2iBn) \tan(c + dx))}{2\sqrt{\tan(c + dx)}} dx}{a(2n + 1)} + \frac{2B \sqrt{\tan(c + dx)} (a + ia \tan(c + dx))}{d(2n + 1)} \right)$$

↓ 27

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{2B \sqrt{\tan(c + dx)} (a + ia \tan(c + dx))^n}{d(2n + 1)} - \frac{\int \frac{(i \tan(c + dx) a + a)^n (a B - a(2nA + A - 2iBn) \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx}{a(2n + 1)} \right)$$

3.571. $\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$

$$\begin{aligned} & \downarrow 3042 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} & \left(\frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \frac{\int \frac{(i\tan(c+dx)a+a)^n(aB-a(2nA+A-2iBn)\tan(c+dx))}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)} \right) \\ & \downarrow 4084 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} & \left(\frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \frac{a(2n+1)(B+iA)\int \frac{(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)} \right) \\ & \downarrow 3042 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} & \left(\frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \frac{a(2n+1)(B+iA)\int \frac{(i\tan(c+dx)a+a)^n}{\sqrt{\tan(c+dx)}} dx}{a(2n+1)} \right) \\ & \downarrow 4047 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} & \left(\frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \frac{ia^3(2n+1)(B+iA)\int -\frac{(i\tan(c+dx)a+a)^{n-1}}{a\sqrt{\tan(c+dx)}(a-ia\tan(c+dx))} dx}{d} \right) \\ & \downarrow 25 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} & \left(\frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \frac{ia^3(2n+1)(B+iA)\int \frac{(i\tan(c+dx)a+a)^{n-1}}{a\sqrt{\tan(c+dx)}(a-ia\tan(c+dx))} dx}{d} \right) \\ & \downarrow 27 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} & \left(\frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \frac{ia^2(2n+1)(B+iA)\int \frac{(i\tan(c+dx)a+a)^{n-1}}{\sqrt{\tan(c+dx)}(a-ia\tan(c+dx))} dx}{d} \right) \\ & \downarrow 148 \end{aligned}$$

3.571. $\int \frac{(a+ia\tan(c+dx))^n(A+B\tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \frac{2a^3(2n+1)(B+iA) \int \frac{(a-ia^3\tan^2(c+dx))^{n-1}}{a(ia^2\tan^2(c+dx)+1)} d\sqrt{\tan(c+dx)}}{d} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \frac{2a^2(2n+1)(B+iA) \int \frac{(a-ia^3\tan^2(c+dx))^{n-1}}{ia^2\tan^2(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} \right)$$

↓ 334

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \frac{2a(2n+1)(B+iA)(1-ia^2\tan^2(c+dx))^{-n}(a-ia^3\tan^2(c+dx))^{n-1}}{d} \right)$$

↓ 333

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \frac{2ia^2(2n+1)(B+iA)\tan(c+dx)(1-ia^2\tan^2(c+dx))^{-n}}{d} \right)$$

↓ 4082

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \frac{2ia^2(2n+1)(B+iA)\tan(c+dx)(1-ia^2\tan^2(c+dx))^{-n}}{d} \right)$$

↓ 76

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \frac{2ia^2(2n+1)(B+iA)\tan(c+dx)(1-ia^2\tan^2(c+dx))^{-n}}{d} \right)$$

↓ 74

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)} - \frac{2ia^2(2n+1)(B+ia)\tan(c+dx)(1-ia^2\tan^2(c+dx))^{-n}}{d(2n+1)}\right)$$

input `Int[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((2*B*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + 2*n)) - ((-2*a*(2*B*n + I*A*(1 + 2*n))*Hypergeometric2F1[1/2, 1 - n, 3/2, (-I)*Tan[c + d*x]]*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + I*Tan[c + d*x])^n) + ((2*I)*a^2*(I*A + B)*(1 + 2*n)*AppellF1[1/2, 1, 1 - n, 3/2, (-I)*a^2*Tan[c + d*x]^2, I*a^2*Tan[c + d*x]^2]*Tan[c + d*x]*(a - I*a^3*Tan[c + d*x]^2)^n)/(d*(1 - I*a^2*Tan[c + d*x]^2)^n))/(a*(1 + 2*n))`

3.571.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 74 `Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)]) && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0]))`

rule 76 `Int[((b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0] && ((RationalQ[m] && !(EqQ[n, -2^(-1)]) && EqQ[c^2 - d^2, 0])) || !RationalQ[n]`

rule 148 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_.))*((e_) + (f_.)*(x_)^(p_.), x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c + d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x] /; FreeQ[{b, c, d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`

rule 333 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 334 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4047 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4080 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`

rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

rule 4729 `Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.571.4 Maple [F]

$$\int \frac{(a + ia \tan(dx + c))^n (A + B \tan(dx + c))}{\sqrt{\cot(dx + c)}} dx$$

input `int((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x)`

output `int((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x)`

3.571.5 Fracas [F]

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\ &= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n}{\sqrt{\cot(dx + c)}} dx \end{aligned}$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="fracas")`

output `integral(((I*A - B)*e^(4*I*d*x + 4*I*c) + 2*B*e^(2*I*d*x + 2*I*c) + I*A - B)*(2*a*e^(2*I*d*x + 2*I*c)/(e^(2*I*d*x + 2*I*c) + 1))^n*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))/(e^(4*I*d*x + 4*I*c) + 2*e^(2*I*d*x + 2*I*c) + 1), x)`

3.571.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(ia(\tan(c + dx) - i))^n (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)**(1/2),x)`

output `Integral((I*a*(tan(c + d*x) - I))^n*(A + B*tan(c + d*x))/sqrt(cot(c + d*x)), x)`

3.571.7 Maxima [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(i a \tan(dx + c) + a)^n}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n/sqrt(cot(d*x + c)), x)`

3.571.8 Giac [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(i a \tan(dx + c) + a)^n}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n/sqrt(cot(d*x + c)), x)`

3.571.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) 1i)^n}{\sqrt{\cot(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n)/cot(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n)/cot(c + d*x)^(1/2), x)`

$$3.572 \quad \int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$$

3.572.1 Optimal result	5392
3.572.2 Mathematica [F]	5393
3.572.3 Rubi [A] (warning: unable to verify)	5393
3.572.4 Maple [F]	5399
3.572.5 Fricas [F]	5399
3.572.6 Sympy [F]	5400
3.572.7 Maxima [F]	5400
3.572.8 Giac [F]	5400
3.572.9 Mupad [F(-1)]	5401

3.572.1 Optimal result

Integrand size = 36, antiderivative size = 291

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx \\ &= \frac{2B(a + ia \tan(c + dx))^n}{d(3 + 2n) \cot^{\frac{3}{2}}(c + dx)} - \frac{2(2iBn - A(3 + 2n))(a + ia \tan(c + dx))^n}{d(1 + 2n)(3 + 2n)\sqrt{\cot(c + dx)}} \\ & \quad - \frac{2(A - iB) \operatorname{AppellF1}\left(\frac{1}{2}, 1 - n, 1, \frac{3}{2}, -i \tan(c + dx), i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n}(a + ia \tan(c + dx))^n}{d\sqrt{\cot(c + dx)}} \\ & \quad + \frac{2(2An(3 + 2n) - iB(3 + 6n + 4n^2)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1 - n, \frac{3}{2}, -i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n}(a + ia \tan(c + dx))^n}{d(1 + 2n)(3 + 2n)\sqrt{\cot(c + dx)}} \end{aligned}$$

```
output 2*B*(a+I*a*tan(d*x+c))^n/d/(3+2*n)/cot(d*x+c)^(3/2)-2*(2*I*B*n-A*(3+2*n))*
(a+I*a*tan(d*x+c))^n/d/(4*n^2+8*n+3)/cot(d*x+c)^(1/2)-2*(A-I*B)*AppellF1(1
/2,1-n,1,3/2,-I*tan(d*x+c),I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/cot(d*x+c)
^(1/2)/((1+I*tan(d*x+c))^n)+2*(2*A*n*(3+2*n)-I*B*(4*n^2+6*n+3))*hypergeom(
[1/2, 1-n],[3/2],-I*tan(d*x+c))*(a+I*a*tan(d*x+c))^n/d/(4*n^2+8*n+3)/cot(d
*x+c)^(1/2)/((1+I*tan(d*x+c))^n)
```

3.572.2 Mathematica [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

input `Integrate[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(3/2), x]`

output `Integrate[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(3/2), x]`

3.572.3 Rubi [A] (warning: unable to verify)

Time = 1.67 (sec) , antiderivative size = 351, normalized size of antiderivative = 1.21, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3042, 4729, 3042, 4080, 27, 3042, 4080, 27, 3042, 4084, 3042, 4047, 25, 27, 148, 27, 334, 333, 4082, 76, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot(c + dx)^{3/2}} dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \tan^{\frac{3}{2}}(c + dx) (i \tan(c + dx) a + a)^n (A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \tan(c + dx)^{3/2} (i \tan(c + dx) a + a)^n (A + B \tan(c + dx)) dx$$

$$\downarrow \text{4080}$$

3.572. $\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2\int-\frac{1}{2}\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)^n(3aB+a(2iBn-A(2n+3))\tan(c+dx)}{a(2n+3)}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}-\frac{\int\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)^n(3aB}{a(2n-$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}-\frac{\int\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)^n(3aB}{a(2n-$$

↓ 4080

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}-\frac{2\int-\frac{(i\tan(c+dx)a+a)^n(a^2(2iBn-A(2n+3))-a^2(2iAn(2n+3))}{2\sqrt{\tan(c+dx)}}}{a(2n+1)}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}-\frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}-\frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)}\right)$$

↓ 4084

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}-\frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)}\right)$$

↓ 3042

3.572. $\int \frac{(a+ia\tan(c+dx))^n(A+B\tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}-\frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)}\right)$$

↓ 4047

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}-\frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)}\right)$$

↓ 25

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}-\frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}-\frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)}\right)$$

↓ 148

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}-\frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)}\right)$$

↓ 27

3.572. $\int \frac{(a+ia\tan(c+dx))^n(A+B\tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)}\right)$$

↓ 334

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)}\right)$$

↓ 333

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)}\right)$$

↓ 4082

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)}\right)$$

↓ 76

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)}\right)$$

↓ 74

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)} - \frac{2a(-A(2n+3)+2iBn)\sqrt{\tan(c+dx)}(a+ia\tan(c+dx))^n}{d(2n+1)}\right)$$

3.572. $\int \frac{(a+ia\tan(c+dx))^n(A+B\tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

input `Int[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(3/2),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((2*B*Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^n)/(d*(3 + 2*n)) - ((2*a*((2*I)*B*n - A*(3 + 2*n))*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + 2*n)) - ((2*a^2*(2*A*n*(3 + 2*n) - I*B*(3 + 6*n + 4*n^2))*Hypergeometric2F1[1/2, 1 - n, 3/2, (-I)*Tan[c + d*x]]*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + I*Tan[c + d*x])^n) - ((2*I)*a^3*(A - I*B)*(3 + 8*n + 4*n^2)*AppellF1[1/2, 1, 1 - n, 3/2, (-I)*a^2*Tan[c + d*x]^2, I*a^2*Tan[c + d*x]^2]*Tan[c + d*x]*(a - I*a^3*Tan[c + d*x]^2)^n)/(d*(1 - I*a^2*Tan[c + d*x]^2)^n))/(a*(1 + 2*n))/(a*(3 + 2*n)))`

3.572.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 74 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))`

rule 76 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n, x], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0] && ((RationalQ[m] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0])) || !RationalQ[n])`

rule 148 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_.)*((e_) + (f_.)*(x_))^(p_.), x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c + d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x] /; FreeQ[{b, c, d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`

- rule 333 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`
- rule 334 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4047 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4080 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`
- rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

```
rule 4084 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Sim
p[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x]
- Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[
e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c -
a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]
```

```
rule 4729 Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.572.4 Maple [F]

$$\int \frac{(a + ia \tan(dx + c))^n (A + B \tan(dx + c))}{\cot(dx + c)^{\frac{3}{2}}} dx$$

```
input int((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x)
```

```
output int((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x)
```

3.572.5 Fracas [F]

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n}{\cot(dx + c)^{\frac{3}{2}}} dx \end{aligned}$$

```
input integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algori
thm="fracas")
```

```
output integral(-((A - I*B)*e^(6*I*d*x + 6*I*c) - (A - 3*I*B)*e^(4*I*d*x + 4*I*c)
- (A + 3*I*B)*e^(2*I*d*x + 2*I*c) + A + I*B)*(2*a*e^(2*I*d*x + 2*I*c)/(e^
(2*I*d*x + 2*I*c) + 1))^n*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2
*I*c) - 1))/(e^(6*I*d*x + 6*I*c) + 3*e^(4*I*d*x + 4*I*c) + 3*e^(2*I*d*x +
2*I*c) + 1), x)
```

3.572. $\int \frac{(a+ia \tan(c+dx))^n (A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

3.572.6 Sympy [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(ia(\tan(c + dx) - i))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c))/cot(d*x+c)**(3/2),x)`

output `Integral((I*a*(tan(c + d*x) - I))**n*(A + B*tan(c + d*x))/cot(c + d*x)**(3/2), x)`

3.572.7 Maxima [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n}{\cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n/cot(d*x + c)^(3/2), x)`

3.572.8 Giac [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n}{\cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n/cot(d*x + c)^(3/2), x)`

3.572.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) i)^n}{\cot(c + dx)^{3/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n)/cot(c + d*x)^(3/2),x)`

output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n)/cot(c + d*x)^(3/2), x)`

3.573
$$\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\cot^{\frac{5}{2}}(c+dx)} dx$$

3.573.1 Optimal result 5402
 3.573.2 Mathematica [F] 5403
 3.573.3 Rubi [A] (warning: unable to verify) 5403
 3.573.4 Maple [F] 5410
 3.573.5 Fricas [F] 5410
 3.573.6 Sympy [F(-1)] 5411
 3.573.7 Maxima [F] 5411
 3.573.8 Giac [F] 5411
 3.573.9 Mupad [F(-1)] 5412

3.573.1 Optimal result

Integrand size = 36, antiderivative size = 383

$$\int \frac{(a + ia \tan(c + dx))^n(A + B \tan(c + dx))}{\cot^{\frac{5}{2}}(c + dx)} dx$$

$$= \frac{2B(a + ia \tan(c + dx))^n}{d(5 + 2n) \cot^{\frac{5}{2}}(c + dx)} - \frac{2(2iBn - A(5 + 2n))(a + ia \tan(c + dx))^n}{d(3 + 2n)(5 + 2n) \cot^{\frac{3}{2}}(c + dx)}$$

$$- \frac{2(2iAn(5 + 2n) + B(15 + 10n + 4n^2))(a + ia \tan(c + dx))^n}{d(1 + 2n)(3 + 2n)(5 + 2n)\sqrt{\cot(c + dx)}}$$

$$+ \frac{2(iA + B) \operatorname{AppellF1}\left(\frac{1}{2}, 1 - n, 1, \frac{3}{2}, -i \tan(c + dx), i \tan(c + dx)\right) (1 + i \tan(c + dx))^{-n}(a + ia \tan(c + dx))}{d\sqrt{\cot(c + dx)}}$$

$$- \frac{2(4Bn(9 + 8n + 2n^2) + iA(15 + 36n + 32n^2 + 8n^3)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 1 - n, \frac{3}{2}, -i \tan(c + dx)\right)}{d(1 + 2n)(3 + 2n)(5 + 2n)\sqrt{\cot(c + dx)}}$$

```
output 2*B*(a+I*a*tan(d*x+c))^n/d/(5+2*n)/cot(d*x+c)^(5/2)-2*(2*I*B*n-A*(5+2*n))*
(a+I*a*tan(d*x+c))^n/d/(3+2*n)/(5+2*n)/cot(d*x+c)^(3/2)-2*(2*I*A*n*(5+2*n)
+B*(4*n^2+10*n+15))*(a+I*a*tan(d*x+c))^n/d/(5+2*n)/(4*n^2+8*n+3)/cot(d*x+c
)^(1/2)+2*(I*A+B)*AppellF1(1/2,1-n,1,3/2,-I*tan(d*x+c),I*tan(d*x+c))*(a+I*
a*tan(d*x+c))^n/d/cot(d*x+c)^(1/2)/((1+I*tan(d*x+c))^n)-2*(4*B*n*(2*n^2+8*
n+9)+I*A*(8*n^3+32*n^2+36*n+15))*hypergeom([1/2, 1-n],[3/2],-I*tan(d*x+c))
*(a+I*a*tan(d*x+c))^n/d/(5+2*n)/(4*n^2+8*n+3)/cot(d*x+c)^(1/2)/((1+I*tan(d
*x+c))^n)
```

3.573.2 Mathematica [F]

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{5}{2}}(c + dx)} dx$$

input `Integrate[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(5/2), x]`

output `Integrate[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(5/2), x]`

3.573.3 Rubi [A] (warning: unable to verify)

Time = 2.25 (sec) , antiderivative size = 442, normalized size of antiderivative = 1.15, number of steps used = 25, number of rules used = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4729, 3042, 4080, 27, 3042, 4080, 27, 3042, 4080, 27, 3042, 4084, 3042, 4047, 25, 27, 148, 27, 334, 333, 4082, 76, 74}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot(c + dx)^{5/2}} dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \tan^{\frac{5}{2}}(c + dx) (i \tan(c + dx) a + a)^n (A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \tan(c + dx)^{5/2} (i \tan(c + dx) a + a)^n (A + B \tan(c + dx)) dx$$

$$\downarrow \text{4080}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2\int-\frac{1}{2}\tan^{\frac{3}{2}}(c+dx)(i\tan(c+dx)a+a)^n(5aB+a(2iBn-A(2n+5))\tan(c+dx))}{a(2n+5)}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+5)}-\frac{\int\tan^{\frac{3}{2}}(c+dx)(i\tan(c+dx)a+a)^n(5aB+a(2iBn-A(2n+5))\tan(c+dx))}{a(2n+5)}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+5)}-\frac{\int\tan(c+dx)^{3/2}(i\tan(c+dx)a+a)^n(5aB+a(2iBn-A(2n+5))\tan(c+dx))}{a(2n+5)}\right)$$

↓ 4080

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+5)}-\frac{2\int-\frac{1}{2}\sqrt{\tan(c+dx)}(i\tan(c+dx)a+a)^n(3a^2(2iBn-A(2n+5))\tan(c+dx))}{a(2n+5)}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+5)}-\frac{2a(-A(2n+5)+2iBn)\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+5)}-\frac{2a(-A(2n+5)+2iBn)\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}\right)$$

↓ 4080

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+5)}-\frac{2a(-A(2n+5)+2iBn)\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}\right)$$

↓ 27

3.573. $\int \frac{(a+ia\tan(c+dx))^n(A+B\tan(c+dx))}{\cot^{\frac{5}{2}}(c+dx)} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn)\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn)\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}\right)$$

↓ 4084

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn)\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn)\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}\right)$$

↓ 4047

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2B\tan^{\frac{5}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn)\tan^{\frac{3}{2}}(c+dx)(a+ia\tan(c+dx))^n}{d(2n+3)}\right)$$

↓ 25

3.573. $\int \frac{(a+ia\tan(c+dx))^n(A+B\tan(c+dx))}{\cot^{\frac{5}{2}}(c+dx)} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} \right)$$

↓ 148

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} \right)$$

↓ 334

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} \right)$$

3.573. $\int \frac{(a+ia \tan(c+dx))^n(A+B \tan(c+dx))}{\cot^{\frac{5}{2}}(c+dx)} dx$

↓ 333

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} \right)$$

↓ 4082

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} \right)$$

↓ 76

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} \right)$$

↓ 74

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B \tan^{\frac{5}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+5)} - \frac{2a(-A(2n+5)+2iBn) \tan^{\frac{3}{2}}(c+dx)(a+ia \tan(c+dx))^n}{d(2n+3)} \right)$$

input `Int[((a + I*a*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(5/2), x]`

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((2*B*Tan[c + d*x]^(5/2)*(a + I*a*Tan[c + d*x])^n)/(d*(5 + 2*n)) - ((2*a*((2*I)*B*n - A*(5 + 2*n))*Tan[c + d*x]^(3/2)*(a + I*a*Tan[c + d*x])^n)/(d*(3 + 2*n)) - ((-2*a^2*((2*I)*A*n*(5 + 2*n) + B*(15 + 10*n + 4*n^2))*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + 2*n)) + ((-2*a^3*(4*B*n*(9 + 8*n + 2*n^2) + I*A*(15 + 36*n + 32*n^2 + 8*n^3))*Hypergeometric2F1[1/2, 1 - n, 3/2, (-I)*Tan[c + d*x]]*Sqrt[Tan[c + d*x]]*(a + I*a*Tan[c + d*x])^n)/(d*(1 + I*Tan[c + d*x])^n) + ((2*I)*a^4*(I*A + B)*(15 + 46*n + 36*n^2 + 8*n^3)*AppellF1[1/2, 1, 1 - n, 3/2, (-I)*a^2*Tan[c + d*x]^2, I*a^2*Tan[c + d*x]^2]*Tan[c + d*x]*(a - I*a^3*Tan[c + d*x]^2)^n)/(d*(1 - I*a^2*Tan[c + d*x]^2)^n)/(a*(1 + 2*n)))/(a*(3 + 2*n)))/(a*(5 + 2*n))
```

3.573.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 74 Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m + 1)/(b*(m + 1)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))
```

```
rule 76 Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n, x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0] && !GtQ[-d/(b*c), 0] && ((RationalQ[m] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0])) || !RationalQ[n])
```

```
rule 148 Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_.)*((e_) + (f_.)*(x_))^(p_.), x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c + d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x] /; FreeQ[{b, c, d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]
```

- rule 333 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`
- rule 334 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4047 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(b/f) Subst[Int[(a + x)^(m - 1)*((c + (d/b)*x)^n/(b^2 + a*x)), x], x, b*Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4080 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp[1/(a*(m + n)) Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^(n - 1)*Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (a*A*d*(m + n) - B*(b*d*m - a*c*n))*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && GtQ[n, 0]`
- rule 4082 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(B/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && EqQ[A*b + a*B, 0]`

rule 4084 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b + a*B)/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n, x], x] - Simp[B/b Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(a - b*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && NeQ[A*b + a*B, 0]`

rule 4729 `Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.573.4 Maple [F]

$$\int \frac{(a + ia \tan(dx + c))^n (A + B \tan(dx + c))}{\cot(dx + c)^{\frac{5}{2}}} dx$$

input `int((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(5/2),x)`

output `int((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(5/2),x)`

3.573.5 Fracas [F]

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(B \tan(dx + c) + A)(ia \tan(dx + c) + a)^n}{\cot(dx + c)^{\frac{5}{2}}} dx \end{aligned}$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(5/2),x, algorithm="fracas")`

output `integral(((I*A + B)*e^(8*I*d*x + 8*I*c) - 2*(I*A + 2*B)*e^(6*I*d*x + 6*I*c) + 6*B*e^(4*I*d*x + 4*I*c) - 2*(-I*A + 2*B)*e^(2*I*d*x + 2*I*c) - I*A + B)*(2*a*e^(2*I*d*x + 2*I*c)/(e^(2*I*d*x + 2*I*c) + 1))^n*sqrt((I*e^(2*I*d*x + 2*I*c) + I)/(e^(2*I*d*x + 2*I*c) - 1))/(e^(8*I*d*x + 8*I*c) + 4*e^(6*I*d*x + 6*I*c) + 6*e^(4*I*d*x + 4*I*c) + 4*e^(2*I*d*x + 2*I*c) + 1), x)`

3.573. $\int \frac{(a+ia \tan(c+dx))^n (A+B \tan(c+dx))}{\cot^{\frac{5}{2}}(c+dx)} dx$

3.573.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{5}{2}}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(d*x+c))**n*(A+B*tan(d*x+c))/cot(d*x+c)**(5/2),x)`

output `Timed out`

3.573.7 Maxima [F]

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(B \tan(dx + c) + A)(i a \tan(dx + c) + a)^n}{\cot(dx + c)^{\frac{5}{2}}} dx \end{aligned}$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(5/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n/cot(d*x + c)^(5/2), x)`

3.573.8 Giac [F]

$$\begin{aligned} & \int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{5}{2}}(c + dx)} dx \\ &= \int \frac{(B \tan(dx + c) + A)(i a \tan(dx + c) + a)^n}{\cot(dx + c)^{\frac{5}{2}}} dx \end{aligned}$$

input `integrate((a+I*a*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(5/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(I*a*tan(d*x + c) + a)^n/cot(d*x + c)^(5/2), x)`

3.573.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{5}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + a \tan(c + dx) 1i)^n}{\cot(c + dx)^{5/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n)/cot(c + d*x)^(5/2),x)`output `int(((A + B*tan(c + d*x))*(a + a*tan(c + d*x)*1i)^n)/cot(c + d*x)^(5/2), x)`

3.574 $\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

3.574.1 Optimal result	5413
3.574.2 Mathematica [A] (verified)	5414
3.574.3 Rubi [A] (verified)	5414
3.574.4 Maple [B] (verified)	5419
3.574.5 Fricas [B] (verification not implemented)	5420
3.574.6 Sympy [F(-1)]	5420
3.574.7 Maxima [A] (verification not implemented)	5421
3.574.8 Giac [F]	5421
3.574.9 Mupad [F(-1)]	5422

3.574.1 Optimal result

Integrand size = 31, antiderivative size = 229

$$\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= -\frac{(b(A-B)+a(A+B)) \arctan\left(1-\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(b(A-B)+a(A+B)) \arctan\left(1+\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d} - \frac{2(Ab+aB)\sqrt{\cot(c+dx)}}{d}$$

$$- \frac{2aA \cot^{\frac{3}{2}}(c+dx)}{3d} + \frac{(a(A-B)-b(A+B)) \log\left(1-\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(a(A-B)-b(A+B)) \log\left(1+\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{2\sqrt{2}d}$$

output

```
-2/3*a*A*cot(d*x+c)^(3/2)/d+1/2*(b*(A-B)+a*(A+B))*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+1/2*(b*(A-B)+a*(A+B))*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+1/4*(a*(A-B)-b*(A+B))*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/4*(a*(A-B)-b*(A+B))*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-2*(A*b+B*a)*cot(d*x+c)^(1/2)/d
```

3.574.2 Mathematica [A] (verified)

Time = 1.03 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.86

$$\int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))(A+B\tan(c+dx))dx$$

$$= \frac{\sqrt{\cot(c+dx)}\left(6\sqrt{2}(b(A-B)+a(A+B))\left(\arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)\right)-\arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)\right)}{12d}$$

input `Integrate[Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*(6*Sqrt[2]*(b*(A - B) + a*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]) + 3*Sqrt[2]*(a*(A - B) - b*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]) - (8*a*A)/Tan[c + d*x]^(3/2) - (24*(A*b + a*B))/Sqrt[Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/(12*d)`

3.574.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 206, normalized size of antiderivative = 0.90, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.516$, Rules used = {3042, 4064, 3042, 4075, 3042, 4011, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))(A+B\tan(c+dx))dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{5/2}(a+b\tan(c+dx))(A+B\tan(c+dx))dx$$

$$\downarrow \text{4064}$$

$$\int \sqrt{\cot(c+dx)}(a\cot(c+dx)+b)(A\cot(c+dx)+B)dx$$

$$\downarrow \text{3042}$$

3.574. $\int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))(A+B\tan(c+dx))dx$

$$\begin{aligned}
& \int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(b-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)\left(B-A \tan\left(c+dx+\frac{\pi}{2}\right)\right) dx \\
& \quad \downarrow 4075 \\
& \int \sqrt{\cot(c+dx)}(-aA+bB+(Ab+aB) \cot(c+dx)) dx - \frac{2aA \cot^{\frac{3}{2}}(c+dx)}{3d} \\
& \quad \downarrow 3042 \\
& \int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(-aA+bB-(Ab+aB) \tan\left(c+dx+\frac{\pi}{2}\right)\right) dx - \frac{2aA \cot^{\frac{3}{2}}(c+dx)}{3d} \\
& \quad \downarrow 4011 \\
& \int \frac{-Ab-aB-(aA-bB) \cot(c+dx)}{\sqrt{\cot(c+dx)}} dx - \frac{2(aB+Ab) \sqrt{\cot(c+dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)}{3d} \\
& \quad \downarrow 3042 \\
& \int \frac{-Ab-aB-(bB-aA) \tan\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}} dx - \frac{2(aB+Ab) \sqrt{\cot(c+dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)}{3d} \\
& \quad \downarrow 4017 \\
& \frac{2 \int \frac{Ab+aB+(aA-bB) \cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)}}{d} - \frac{2(aB+Ab) \sqrt{\cot(c+dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)}{3d} \\
& \quad \downarrow 1482 \\
& \frac{2\left(\frac{1}{2}(a(A+B)+b(A-B)) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)} - \frac{1}{2}(a(A-B)-b(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)}\right)}{d} \\
& \quad \frac{2(aB+Ab) \sqrt{\cot(c+dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)}{3d} \\
& \quad \downarrow 1476 \\
& \frac{2\left(\frac{1}{2}(a(A+B)+b(A-B))\left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2} \sqrt{\cot(c+dx)+1}} d \sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2} \sqrt{\cot(c+dx)+1}} d \sqrt{\cot(c+dx)}\right)\right)}{d} \\
& \quad \frac{2(aB+Ab) \sqrt{\cot(c+dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)}{3d} \\
& \quad \downarrow 1082
\end{aligned}$$

3.574. $\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

$$\frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)-1} d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) - \frac{1}{2}(a(A-B) - b(A+B)) \int \frac{1}{\cot(c+dx)} dx \right)}{d} - \frac{2(aB + Ab)\sqrt{\cot(c+dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)}{3d}$$

↓ 217

$$\frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a(A-B) - b(A+B)) \int \frac{1}{\cot(c+dx)} dx \right)}{d} - \frac{2(aB + Ab)\sqrt{\cot(c+dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)}{3d}$$

↓ 1479

$$\frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a(A-B) - b(A+B)) \left(-\int \frac{1}{\cot(c+dx)} dx \right) \right)}{d} - \frac{2(aB + Ab)\sqrt{\cot(c+dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)}{3d}$$

↓ 25

$$\frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a(A-B) - b(A+B)) \left(\int \frac{1}{\cot(c+dx)} dx \right) \right)}{d} - \frac{2(aB + Ab)\sqrt{\cot(c+dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)}{3d}$$

↓ 27

$$\frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a(A-B) - b(A+B)) \left(\int \frac{1}{\cot(c+dx)} dx \right) \right)}{d} - \frac{2(aB + Ab)\sqrt{\cot(c+dx)}}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)}{3d}$$

↓ 1103

3.574. $\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

$$\frac{2\left(\frac{1}{2}(a(A+B) + b(A-B))\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}}\right) - \frac{1}{2}(a(A-B) - b(A+B))\left(\frac{\log\left(\frac{2(aB+Ab)\sqrt{\cot(c+dx)}}{d} - \frac{2aA\cot^{\frac{3}{2}}(c+dx)}{3d}\right)}{d}\right)}{d}\right)}{d}$$

input `Int[Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(-2*(A*b + a*B)*Sqrt[Cot[c + d*x]]/d - (2*a*A*Cot[c + d*x]^(3/2))/(3*d) + (2*((b*(A - B) + a*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]/Sqrt[2]]))/2 - ((a*(A - B) - b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d`

3.574.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2(d/e), 2]\}, \text{Simp}[e/(2c) \text{Int}[1/\text{Simp}[d/e + qx + x^2, x], x], x] + \text{Simp}[e/(2c) \text{Int}[1/\text{Simp}[d/e - qx + x^2, x], x], x]] /; \text{FreeQ}\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[d*e]$

rule 1479 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2(d/e), 2]\}, \text{Simp}[e/(2c*q) \text{Int}[(q - 2x)/\text{Simp}[d/e + qx - x^2, x], x], x] + \text{Simp}[e/(2c*q) \text{Int}[(q + 2x)/\text{Simp}[d/e - qx - x^2, x], x], x]] /; \text{FreeQ}\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{NegQ}[d*e]$

rule 1482 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a*c, 2]\}, \text{Simp}[(d*q + a*e)/(2*a*c) \text{Int}[(q + c*x^2)/(a + c*x^4), x], x] + \text{Simp}[(d*q - a*e)/(2*a*c) \text{Int}[(q - c*x^2)/(a + c*x^4), x], x]] /; \text{FreeQ}\{a, c, d, e\}, x] \&\& \text{NeQ}[c*d^2 + a*e^2, 0] \&\& \text{NeQ}[c*d^2 - a*e^2, 0] \&\& \text{NegQ}[(-a)*c]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4011 $\text{Int}[\frac{(a_.) + (b_.)\tan[(e_.) + (f_.)x]}{(c_.) + (d_.)\tan[(e_.) + (f_.)x]}]^m, x_Symbol] \rightarrow \text{Simp}[d*((a + b*\text{Tan}[e + f*x])^m/(f*m)), x] + \text{Int}[(a + b*\text{Tan}[e + f*x])^{m-1}*\text{Simp}[a*c - b*d + (b*c + a*d)*\text{Tan}[e + f*x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{GtQ}[m, 0]$

rule 4017 $\text{Int}[\frac{(c_.) + (d_.)\tan[(e_.) + (f_.)x]}{\sqrt{(b_.)\tan[(e_.) + (f_.)x] + (c_.) + (d_.)\tan[(e_.) + (f_.)x]}}, x_Symbol] \rightarrow \text{Simp}[2/f \text{Subst}[\text{Int}[(b*c + d*x^2)/(b^2 + x^4), x], x, \text{Sqrt}[b*\text{Tan}[e + f*x]]], x] /; \text{FreeQ}\{b, c, d, e, f\}, x] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0]$

rule 4064 $\text{Int}[\frac{(\cot[(e_.) + (f_.)x])^{p_.*} * ((a_.) + (b_.)\tan[(e_.) + (f_.)x])^{m_.*} * ((c_.) + (d_.)\tan[(e_.) + (f_.)x])^{n_.*}}{(a_.) + (b_.)\tan[(e_.) + (f_.)x]}], x_Symbol] \rightarrow \text{Simp}[g^{m+n} \text{Int}[(g*\text{Cot}[e + f*x])^{p-m-n}*(b + a*\text{Cot}[e + f*x])^m*(d + c*\text{Cot}[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, p\}, x] \&\& \text{!IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4075 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.574.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 534 vs. $2(195) = 390$.

Time = 0.47 (sec) , antiderivative size = 535, normalized size of antiderivative = 2.34

method	result
derivativedivides	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \left(3A\sqrt{2} \ln\left(-\frac{1+\sqrt{2}\sqrt{\tan(dx+c)+\tan(dx+c)}}{\sqrt{2}\sqrt{\tan(dx+c)-\tan(dx+c)-1}}\right) \tan(dx+c)^{\frac{3}{2}} a+6A\sqrt{2} \arctan\left(1+\sqrt{2}\sqrt{\tan(dx+c)-\tan(dx+c)-1}\right)}{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \left(3A\sqrt{2} \ln\left(-\frac{1+\sqrt{2}\sqrt{\tan(dx+c)+\tan(dx+c)}}{\sqrt{2}\sqrt{\tan(dx+c)-\tan(dx+c)-1}}\right) \tan(dx+c)^{\frac{3}{2}} a+6A\sqrt{2} \arctan\left(1+\sqrt{2}\sqrt{\tan(dx+c)-\tan(dx+c)-1}\right)}\right)}$
default	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \left(3A\sqrt{2} \ln\left(-\frac{1+\sqrt{2}\sqrt{\tan(dx+c)+\tan(dx+c)}}{\sqrt{2}\sqrt{\tan(dx+c)-\tan(dx+c)-1}}\right) \tan(dx+c)^{\frac{3}{2}} a+6A\sqrt{2} \arctan\left(1+\sqrt{2}\sqrt{\tan(dx+c)-\tan(dx+c)-1}\right)}{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \left(3A\sqrt{2} \ln\left(-\frac{1+\sqrt{2}\sqrt{\tan(dx+c)+\tan(dx+c)}}{\sqrt{2}\sqrt{\tan(dx+c)-\tan(dx+c)-1}}\right) \tan(dx+c)^{\frac{3}{2}} a+6A\sqrt{2} \arctan\left(1+\sqrt{2}\sqrt{\tan(dx+c)-\tan(dx+c)-1}\right)}\right)}$

input `int(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/12/d*(1/\tan(d*x+c))^{5/2}*\tan(d*x+c)*(3*A*2^{1/2}*\ln(-1+2^{1/2}*\tan(d*x+c)^{1/2}+\tan(d*x+c))/(2^{1/2}*\tan(d*x+c)^{1/2}-\tan(d*x+c)-1))*\tan(d*x+c)^{3/2} \\ & +6*A*2^{1/2}*\arctan(1+2^{1/2}*\tan(d*x+c)^{1/2})*\tan(d*x+c)^{3/2} \\ & +6*A*2^{1/2}*\arctan(1+2^{1/2}*\tan(d*x+c)^{1/2})*\tan(d*x+c)^{3/2} \\ & +6*A*2^{1/2}*\arctan(-1+2^{1/2}*\tan(d*x+c)^{1/2})*\tan(d*x+c)^{3/2} \\ & +6*A*2^{1/2}*\arctan(-1+2^{1/2}*\tan(d*x+c)^{1/2})*\tan(d*x+c)^{3/2} \\ & +3*A*2^{1/2}*\ln(-2^{1/2}*\tan(d*x+c)^{1/2}-\tan(d*x+c)-1)/(1+2^{1/2}*\tan(d*x+c)^{1/2}+\tan(d*x+c)) \\ &)*\tan(d*x+c)^{3/2} \\ & +3*B*2^{1/2}*\ln(-1+2^{1/2}*\tan(d*x+c)^{1/2}+\tan(d*x+c))/(2^{1/2}*\tan(d*x+c)^{1/2}-\tan(d*x+c)-1))*\tan(d*x+c)^{3/2} \\ & +6*B*2^{1/2}*\arctan(1+2^{1/2}*\tan(d*x+c)^{1/2})*\tan(d*x+c)^{3/2} \\ & +6*B*2^{1/2}*\arctan(-1+2^{1/2}*\tan(d*x+c)^{1/2})*\tan(d*x+c)^{3/2} \\ & +6*B*2^{1/2}*\arctan(-1+2^{1/2}*\tan(d*x+c)^{1/2})*\tan(d*x+c)^{3/2} \\ & +3*B*2^{1/2}*\ln(-2^{1/2}*\tan(d*x+c)^{1/2}-\tan(d*x+c)-1)/(1+2^{1/2}*\tan(d*x+c)^{1/2}+\tan(d*x+c)) \\ &)*\tan(d*x+c)^{3/2} \\ & +24*A*\tan(d*x+c)*b+24*B*\tan(d*x+c)*a+8*A*a \end{aligned}$$

3.574.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2267 vs. 2(195) = 390.

Time = 0.70 (sec) , antiderivative size = 2267, normalized size of antiderivative = 9.90

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm=
"fricas")
```

```
output -1/6*(3*d*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-(A
^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2
+ B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d
^4))/d^2)*log(((B*a + A*b)*d^3*sqrt(-(A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*
B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*
a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) + ((A^3 - A*B^2)*a^3 - (5*A^2*B
- B^3)*a^2*b - (A^3 - 5*A*B^2)*a*b^2 + (A^2*B - B^3)*b^3)*d)*sqrt(-(2*A*B*
a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-(A^4 - 2*A^2*B^2 + B^4)*a
^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3
*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2) + ((A^4 - B^4)
*a^4 - 4*(A^3*B + A*B^3)*a^3*b - 4*(A^3*B + A*B^3)*a*b^3 - (A^4 - B^4)*b^4
)*sqrt(tan(d*x + c))*tan(d*x + c) - 3*d*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*
(A^2 - B^2)*a*b + d^2*sqrt(-(A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^
3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 +
(A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2)*log(-((B*a + A*b)*d^3*sqrt(-(A^4
- 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B
^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4)
+ ((A^3 - A*B^2)*a^3 - (5*A^2*B - B^3)*a^2*b - (A^3 - 5*A*B^2)*a*b^2 + (A^
2*B - B^3)*b^3)*d)*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*
sqrt(-(A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - ...
```

3.574.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \text{Timed out}$$

```
input integrate(cot(d*x+c)**(5/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)
```

3.574. $\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

output Timed out

3.574.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.86

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{6\sqrt{2}((A+B)a + (A-B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 6\sqrt{2}((A+B)a + (A-B)b) \arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) - 3\sqrt{2}((A-B)a - (A+B)b) \log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + \frac{1}{\tan(dx+c)} + 1\right) + 3\sqrt{2}((A-B)a - (A+B)b) \log\left(-\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + \frac{1}{\tan(dx+c)} + 1\right) - 8Aa \tan^{\frac{3}{2}}(dx+c) - 24(Ba + Ab) \sqrt{\tan(dx+c)}}{d}$$

input `integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/12*(6*sqrt(2)*((A + B)*a + (A - B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 6*sqrt(2)*((A + B)*a + (A - B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - 3*sqrt(2)*((A - B)*a - (A + B)*b)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + 3*sqrt(2)*((A - B)*a - (A + B)*b)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - 8*A*a/tan(d*x + c)^(3/2) - 24*(B*a + A*b)/sqrt(tan(d*x + c)))/d`

3.574.8 Giac [F]

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a) \cot(dx + c)^{\frac{5}{2}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)*cot(d*x + c)^(5/2), x)`

3.574. $\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

3.574.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{5/2} (A + B \tan(c + dx)) (a + b \tan(c + dx)) dx$$

input `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)`output `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)), x)`

3.575 $\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

3.575.1 Optimal result	5423
3.575.2 Mathematica [A] (verified)	5424
3.575.3 Rubi [A] (verified)	5424
3.575.4 Maple [B] (verified)	5428
3.575.5 Fricas [B] (verification not implemented)	5429
3.575.6 Sympy [F]	5430
3.575.7 Maxima [A] (verification not implemented)	5431
3.575.8 Giac [F]	5431
3.575.9 Mupad [F(-1)]	5432

3.575.1 Optimal result

Integrand size = 31, antiderivative size = 205

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= -\frac{(a(A-B)-b(A+B)) \arctan\left(1-\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(a(A-B)-b(A+B)) \arctan\left(1+\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d} - \frac{2aA\sqrt{\cot(c+dx)}}{d}$$

$$- \frac{(b(A-B)+a(A+B)) \log\left(1-\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{(b(A-B)+a(A+B)) \log\left(1+\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{2\sqrt{2}d}$$

output

```
1/2*(a*(A-B)-b*(A+B))*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+1/2*(a
*(A-B)-b*(A+B))*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/4*(b*(A-B)+
a*(A+B))*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+1/4*(b*(A-B)+
a*(A+B))*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-2*a*A*cot(d*x
+c)^(1/2)/d
```

3.575.2 Mathematica [A] (verified)

Time = 0.53 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.87

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \frac{\sqrt{\cot(c+dx)} \left(2\sqrt{2}(a(A-B) - b(A+B)) \left(\arctan \left(1 - \sqrt{2}\sqrt{\tan(c+dx)} \right) - \arctan \left(1 + \sqrt{2}\sqrt{\tan(c+dx)} \right) \right) \right)}{4d}$$

input `Integrate[Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*(2*Sqrt[2]*(a*(A - B) - b*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]) - Sqrt[2]*(b*(A - B) + a*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]) - (8*a*A)/Sqrt[Tan[c + d*x]])*Sqrt[Tan[c + d*x]])/(4*d)`

3.575.3 Rubi [A] (verified)

Time = 0.61 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.89, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.452$, Rules used = {3042, 4064, 3042, 4075, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{3/2}(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4064}$$

$$\int \frac{(a \cot(c+dx) + b)(A \cot(c+dx) + B)}{\sqrt{\cot(c+dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(b - a \tan(c+dx + \frac{\pi}{2}))(B - A \tan(c+dx + \frac{\pi}{2}))}{\sqrt{-\tan(c+dx + \frac{\pi}{2})}} dx$$

3.575. $\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \int \frac{-aA + bB + (Ab + aB) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx - \frac{2aA\sqrt{\cot(c + dx)}}{d} \\
& \quad \downarrow \text{4075} \\
& \int \frac{-aA + bB - (Ab + aB) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx - \frac{2aA\sqrt{\cot(c + dx)}}{d} \\
& \quad \downarrow \text{3042} \\
& \frac{2 \int \frac{aA - bB - (Ab + aB) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} - \frac{2aA\sqrt{\cot(c + dx)}}{d} \\
& \quad \downarrow \text{4017} \\
& \quad \downarrow \text{1482} \\
& \frac{2\left(\frac{1}{2}(a(A + B) + b(A - B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}(a(A - B) - b(A + B)) \int \frac{\cot(c + dx) + 1}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}\right)}{d} \\
& \quad \downarrow \text{1476} \\
& \frac{2\left(\frac{1}{2}(a(A + B) + b(A - B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{1}{2} \int \frac{1}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}}\right)\right)}{d} \\
& \quad \downarrow \text{1082} \\
& \frac{2\left(\frac{1}{2}(a(A + B) + b(A - B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\int \frac{1}{-\cot(c + dx) - 1} d(1 - \sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}}\right)\right)}{d} \\
& \quad \downarrow \text{217} \\
& \frac{2\left(\frac{1}{2}(a(A + B) + b(A - B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}}\right)\right)}{d} \\
& \quad \downarrow \text{1479}
\end{aligned}$$

3.575. $\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

$$\begin{aligned}
& \frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) + \frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\log\left[1 - \sqrt{2}\sqrt{\cot(c+dx)}\right]}{\sqrt{2}} + \frac{\log\left[1 + \sqrt{2}\sqrt{\cot(c+dx)}\right]}{\sqrt{2}} + \frac{\log\left[1 + \sqrt{2}\sqrt{\cot(c+dx)+1}\right]}{\sqrt{2}} + \frac{\log\left[1 - \sqrt{2}\sqrt{\cot(c+dx)+1}\right]}{\sqrt{2}} \right)}{d} \right. \\
& \qquad \qquad \qquad \frac{2aA\sqrt{\cot(c+dx)}}{d} \\
& \qquad \qquad \qquad \downarrow \text{25} \\
& \frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) + \frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\log\left[1 - \sqrt{2}\sqrt{\cot(c+dx)}\right]}{\sqrt{2}} + \frac{\log\left[1 + \sqrt{2}\sqrt{\cot(c+dx)}\right]}{\sqrt{2}} + \frac{\log\left[1 + \sqrt{2}\sqrt{\cot(c+dx)+1}\right]}{\sqrt{2}} + \frac{\log\left[1 - \sqrt{2}\sqrt{\cot(c+dx)+1}\right]}{\sqrt{2}} \right)}{d} \right. \\
& \qquad \qquad \qquad \frac{2aA\sqrt{\cot(c+dx)}}{d} \\
& \qquad \qquad \qquad \downarrow \text{27} \\
& \frac{2 \left(\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) + \frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\log\left[1 - \sqrt{2}\sqrt{\cot(c+dx)}\right]}{\sqrt{2}} + \frac{\log\left[1 + \sqrt{2}\sqrt{\cot(c+dx)}\right]}{\sqrt{2}} + \frac{\log\left[1 + \sqrt{2}\sqrt{\cot(c+dx)+1}\right]}{\sqrt{2}} + \frac{\log\left[1 - \sqrt{2}\sqrt{\cot(c+dx)+1}\right]}{\sqrt{2}} \right)}{d} \right. \\
& \qquad \qquad \qquad \frac{2aA\sqrt{\cot(c+dx)}}{d} \\
& \qquad \qquad \qquad \downarrow \text{1103} \\
& \frac{2 \left(\frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) + \frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\log\left[1 - \sqrt{2}\sqrt{\cot(c+dx)}\right]}{\sqrt{2}} + \frac{\log\left[1 + \sqrt{2}\sqrt{\cot(c+dx)}\right]}{\sqrt{2}} + \frac{\log\left[1 + \sqrt{2}\sqrt{\cot(c+dx)+1}\right]}{\sqrt{2}} + \frac{\log\left[1 - \sqrt{2}\sqrt{\cot(c+dx)+1}\right]}{\sqrt{2}} \right)}{d} \right. \\
& \qquad \qquad \qquad \frac{2aA\sqrt{\cot(c+dx)}}{d}
\end{aligned}$$

input `Int[Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `(-2*a*A*Sqrt[Cot[c + d*x]])/d + (2*(((a*(A - B) - b*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]))/2 + ((b*(A - B) + a*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d`

3.575.3.1 Defintions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, x], x]$
- rule 27 $\text{Int}[(a_)*(\text{Fx}_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[\text{Fx}, (b_)*(\text{Gx}_)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[(a_) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 1082 $\text{Int}[(a_) + (b_.)*(x_) + (c_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*c*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \quad \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_) + (e_.)*(x_)]/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$
- rule 1476 $\text{Int}[(d_) + (e_.)*(x_)^2]/((a_) + (c_.)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Simp}[e/(2*c) \quad \text{Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Simp}[e/(2*c) \quad \text{Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{PosQ}[d*e]$
- rule 1479 $\text{Int}[(d_) + (e_.)*(x_)^2]/((a_) + (c_.)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2*(d/e), 2]\}, \text{Simp}[e/(2*c*q) \quad \text{Int}[(q - 2*x)/\text{Simp}[d/e + q*x - x^2, x], x], x] + \text{Simp}[e/(2*c*q) \quad \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{NegQ}[d*e]$
- rule 1482 $\text{Int}[(d_) + (e_.)*(x_)^2]/((a_) + (c_.)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a*c, 2]\}, \text{Simp}[(d*q + a*e)/(2*a*c) \quad \text{Int}[(q + c*x^2)/(a + c*x^4), x], x] + \text{Simp}[(d*q - a*e)/(2*a*c) \quad \text{Int}[(q - c*x^2)/(a + c*x^4), x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{NeQ}[c*d^2 + a*e^2, 0] \ \&\& \ \text{NeQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{NegQ}[-(a)*c]$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4075 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[B*d*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A*c - B*d + (B*c + A*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m}, x] && NeQ[b*c - a*d, 0] && !LeQ[m, -1]`

3.575.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 511 vs. $2(175) = 350$.

Time = 0.37 (sec) , antiderivative size = 512, normalized size of antiderivative = 2.50

method	result
derivativedivides	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{3}{2}} \tan(dx+c) \left(A\sqrt{2} \ln\left(-\frac{1+\sqrt{2}\sqrt{\tan(dx+c)+\tan(dx+c)}}{\sqrt{2}\sqrt{\tan(dx+c)-\tan(dx+c)-1}}\right) \sqrt{\tan(dx+c)} b - 2A\sqrt{2} \arctan\left(1+\sqrt{2}\sqrt{\tan(dx+c)}\right) \right)$
default	$\left(\frac{1}{\tan(dx+c)}\right)^{\frac{3}{2}} \tan(dx+c) \left(A\sqrt{2} \ln\left(-\frac{1+\sqrt{2}\sqrt{\tan(dx+c)+\tan(dx+c)}}{\sqrt{2}\sqrt{\tan(dx+c)-\tan(dx+c)-1}}\right) \sqrt{\tan(dx+c)} b - 2A\sqrt{2} \arctan\left(1+\sqrt{2}\sqrt{\tan(dx+c)}\right) \right)$

input `int(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)), x, method=_RETURNVERBOSE)`

3.575. $\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))(A+B\tan(c+dx)) dx$

output $\frac{1}{4}d*(1/\tan(dx+c))^{3/2}*\tan(dx+c)*(A^{1/2}*\ln(-(1+2^{1/2}*\tan(dx+c))^{1/2}+\tan(dx+c)))/(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1))*\tan(dx+c)^{1/2}*b-2*A^{1/2}*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2}))*\tan(dx+c)^{1/2}*a+2*A^{1/2}*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2}))*\tan(dx+c)^{1/2}*b-2*A^{1/2}*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2}))*\tan(dx+c)^{1/2}*a+2*A^{1/2}*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2}))*\tan(dx+c)^{1/2}*b-A^{1/2}*\ln(-(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1)/(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c)))*\tan(dx+c)^{1/2}*a+B^{1/2}*\ln(-(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c)))/(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1))*\tan(dx+c)^{1/2}*a+2*B^{1/2}*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2}))*\tan(dx+c)^{1/2}*a+2*B^{1/2}*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2}))*\tan(dx+c)^{1/2}*b+2*B^{1/2}*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2}))*\tan(dx+c)^{1/2}*a+2*B^{1/2}*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2}))*\tan(dx+c)^{1/2}*b+B^{1/2}*\ln(-(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1)/(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c)))*\tan(dx+c)^{1/2}*b-8*A*a)$

3.575.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2212 vs. $2(175) = 350$.

Time = 0.70 (sec) , antiderivative size = 2212, normalized size of antiderivative = 10.79

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))(A+B\tan(c+dx))dx = \text{Too large to display}$$

input `integrate(cot(dx+c)^(3/2)*(a+b*tan(dx+c))*(A+B*tan(dx+c)),x, algorithm="fracas")`

output

```
-1/2*(d*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4
- 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B
^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))
/d^2)*log(((A*a - B*b)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B -
A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b
^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) - ((A^2*B - B^3)*a^3 + (A^3 - 5*A*B
^2)*a^2*b - (5*A^2*B - B^3)*a*b^2 - (A^3 - A*B^2)*b^3)*d)*sqrt((2*A*B*a^2
- 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 -
8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B -
A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2) + ((A^4 - B^4)*a^4
- 4*(A^3*B + A*B^3)*a^3*b - 4*(A^3*B + A*B^3)*a*b^3 - (A^4 - B^4)*b^4)*sq
rt(tan(d*x + c))) - d*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^
2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 -
10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B
^4)*b^4)/d^4))/d^2)*log(-((A*a - B*b)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a
^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3
*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) - ((A^2*B - B^3)*a^3
+ (A^3 - 5*A*B^2)*a^2*b - (5*A^2*B - B^3)*a*b^2 - (A^3 - A*B^2)*b^3)*d)*s
qrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B
^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^...
```

3.575.6 Sympy [F]

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx))(a + b \tan(c + dx)) \cot^{\frac{3}{2}}(c + dx) dx$$

input `integrate(cot(d*x+c)**(3/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))*cot(c + d*x)**(3/2), x)`

3.575.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.87

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{2\sqrt{2}((A - B)a - (A + B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}((A - B)a - (A + B)b) \arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + \sqrt{2}((A + B)a + (A - B)b) \log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + \frac{1}{\tan(dx+c)} + 1\right) - \sqrt{2}((A + B)a + (A - B)b) \log\left(-\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + \frac{1}{\tan(dx+c)} + 1\right) - 8Aa/\sqrt{\tan(dx+c)}}{d}$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/4*(2*sqrt(2)*((A - B)*a - (A + B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a - (A + B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)*((A + B)*a + (A - B)*b)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*((A + B)*a + (A - B)*b)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - 8*A*a/sqrt(tan(d*x + c)))/d`

3.575.8 Giac [F]

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a) \cot(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)*cot(d*x + c)^(3/2), x)`

3.575.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))(A + B \tan(c + dx)) dx \\ &= \int \cot(c + dx)^{3/2} (A + B \tan(c + dx)) (a + b \tan(c + dx)) dx \end{aligned}$$

input `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)`output `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)), x)`

3.576 $\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

3.576.1 Optimal result	5433
3.576.2 Mathematica [A] (verified)	5434
3.576.3 Rubi [A] (verified)	5434
3.576.4 Maple [B] (verified)	5439
3.576.5 Fricas [B] (verification not implemented)	5439
3.576.6 Sympy [F]	5440
3.576.7 Maxima [A] (verification not implemented)	5441
3.576.8 Giac [F]	5441
3.576.9 Mupad [F(-1)]	5442

3.576.1 Optimal result

Integrand size = 31, antiderivative size = 205

$$\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$$

$$= \frac{(b(A - B) + a(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(b(A - B) + a(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d} + \frac{2bB}{d\sqrt{\cot(c + dx)}}$$

$$- \frac{(a(A - B) - b(A + B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{(a(A - B) - b(A + B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

output

```
-1/2*(b*(A-B)+a*(A+B))*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/2*(
b*(A-B)+a*(A+B))*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/4*(a*(A-B)
-b*(A+B))*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+1/4*(a*(A-B)
-b*(A+B))*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+2*b*B/d/cot(
d*x+c)^(1/2)
```


3.576.2 Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.87

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))(A+B\tan(c+dx))dx = \frac{\sqrt{\cot(c+dx)}\left(2\sqrt{2}(b(A-B)+a(A+B))\left(\arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)\right)-\arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)\right)}{d}$$

input `Integrate[Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

output `-1/4*(Sqrt[Cot[c + d*x]]*(2*Sqrt[2]*(b*(A - B) + a*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]) + Sqrt[2]*(a*(A - B) - b*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]) - 8*b*B*Sqrt[Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/d`

3.576.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.89, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.484$, Rules used = {3042, 4064, 3042, 4074, 3042, 4017, 25, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))(A+B\tan(c+dx))dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))(A+B\tan(c+dx))dx \\ & \quad \downarrow \text{4064} \\ & \int \frac{(a\cot(c+dx)+b)(A\cot(c+dx)+B)}{\cot^{\frac{3}{2}}(c+dx)}dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(b-a\tan(c+dx+\frac{\pi}{2}))(B-A\tan(c+dx+\frac{\pi}{2}))}{(-\tan(c+dx+\frac{\pi}{2}))^{3/2}}dx \end{aligned}$$

3.576. $\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))(A+B\tan(c+dx))dx$

$$\begin{aligned}
& \int \frac{Ab + aB + (aA - bB) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx + \frac{2bB}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow 4074 \\
& \int \frac{Ab + aB - (aA - bB) \tan\left(c + dx + \frac{\pi}{2}\right)}{\sqrt{-\tan\left(c + dx + \frac{\pi}{2}\right)}} dx + \frac{2bB}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow 3042 \\
& \frac{2 \int -\frac{Ab + aB + (aA - bB) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} + \frac{2bB}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow 4017 \\
& \frac{2bB}{d\sqrt{\cot(c + dx)}} - \frac{2 \int \frac{Ab + aB + (aA - bB) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} \\
& \quad \downarrow 25 \\
& \frac{2bB}{d\sqrt{\cot(c + dx)}} - \frac{2 \int \frac{Ab + aB + (aA - bB) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} \\
& \quad \downarrow 1482 \\
& \frac{2\left(\frac{1}{2}(a(A - B) - b(A + B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \frac{1}{2}(a(A + B) + b(A - B)) \int \frac{\cot(c + dx) + 1}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}\right)}{d} \\
& \quad \downarrow 1476 \\
& \frac{2\left(\frac{1}{2}(a(A - B) - b(A + B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{1}{2} \int \frac{1}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx)}}\right)\right)}{d} \\
& \quad \downarrow 1082 \\
& \frac{2\left(\frac{1}{2}(a(A - B) - b(A + B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{\int \frac{1}{-\cot(c + dx) - 1} d(1 - \sqrt{2}\sqrt{\cot(c + dx)})}{\sqrt{2}}\right)\right)}{d} \\
& \quad \downarrow 217 \\
& \frac{2bB}{d\sqrt{\cot(c + dx)}}
\end{aligned}$$

3.576. $\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

$$2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}} - \right) \right)$$

$$\frac{2bB}{d\sqrt{\cot(c + dx)}}$$

↓ 1479

$$2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \left(- \frac{\int - \frac{\sqrt{2} - 2\sqrt{\cot(c + dx)}}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} - \frac{\int - \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}} - \right) \right)$$

$$\frac{2bB}{d\sqrt{\cot(c + dx)}}$$

↓ 25

$$2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\int \frac{\sqrt{2} - 2\sqrt{\cot(c + dx)}}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}} - \right) \right)$$

$$\frac{2bB}{d\sqrt{\cot(c + dx)}}$$

↓ 27

$$2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\int \frac{\sqrt{2} - 2\sqrt{\cot(c + dx)}}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c + dx) + 1}}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)} \right) - \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}} - \right) \right)$$

$$\frac{2bB}{d\sqrt{\cot(c + dx)}}$$

↓ 1103

$$2 \left(\frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{\log(\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx) + 1})}{2\sqrt{2}} - \frac{\log(\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1})}{2\sqrt{2}} \right) - \frac{1}{2}(a(A + B) + b(A - B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}} - \right) \right)$$

$$\frac{2bB}{d\sqrt{\cot(c + dx)}}$$

input `Int[Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]),x]`

```
output (2*b*B)/(d*Sqrt[Cot[c + d*x]]) + (2*(-1/2*((b*(A - B) + a*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2])) + ((a*(A - B) - b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d
```

3.576.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]
```

```
rule 1103 Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

```
rule 1476 Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

```
rule 1479 Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4074 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c + b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.576.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 425 vs. 2(175) = 350.

Time = 0.36 (sec) , antiderivative size = 426, normalized size of antiderivative = 2.08

method	result
derivativedivides	$\sqrt{\frac{1}{\tan(dx+c)}} \sqrt{\tan(dx+c)} \left(A \ln \left(-\frac{1+\sqrt{2} \sqrt{\tan(dx+c)} + \tan(dx+c)}{\sqrt{2} \sqrt{\tan(dx+c)} - \tan(dx+c) - 1} \right) \sqrt{2} a + 2A \arctan \left(1 + \sqrt{2} \sqrt{\tan(dx+c)} \right) \sqrt{2} a + 2A \arctan \left(-1 + \sqrt{2} \sqrt{\tan(dx+c)} \right) \sqrt{2} a + 2A \arctan \left(-\frac{1-\sqrt{2} \sqrt{\tan(dx+c)} + \tan(dx+c)}{\sqrt{2} \sqrt{\tan(dx+c)} - \tan(dx+c) - 1} \right) \sqrt{2} a \right)$
default	$\sqrt{\frac{1}{\tan(dx+c)}} \sqrt{\tan(dx+c)} \left(A \ln \left(-\frac{1+\sqrt{2} \sqrt{\tan(dx+c)} + \tan(dx+c)}{\sqrt{2} \sqrt{\tan(dx+c)} - \tan(dx+c) - 1} \right) \sqrt{2} a + 2A \arctan \left(1 + \sqrt{2} \sqrt{\tan(dx+c)} \right) \sqrt{2} a + 2A \arctan \left(-1 + \sqrt{2} \sqrt{\tan(dx+c)} \right) \sqrt{2} a + 2A \arctan \left(-\frac{1-\sqrt{2} \sqrt{\tan(dx+c)} + \tan(dx+c)}{\sqrt{2} \sqrt{\tan(dx+c)} - \tan(dx+c) - 1} \right) \sqrt{2} a \right)$

input `int(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/4/d*(1/tan(d*x+c))^(1/2)*tan(d*x+c)^(1/2)*(A*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1))*2^(1/2)*a+2*A*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*a+2*A*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*b+2*A*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*a+2*A*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*b+A*ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))*2^(1/2)*b-B*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1))*2^(1/2)*b+2*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*a-2*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*b+2*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*a-2*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*b+B*ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))*2^(1/2)*a+8*B*b*tan(d*x+c)^(1/2)`

3.576.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2216 vs. 2(175) = 350.

Time = 0.71 (sec) , antiderivative size = 2216, normalized size of antiderivative = 10.81

$$\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="fricas")`

3.576. $\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))(A + B \tan(c + dx)) dx$

output

```

1/2*(d*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4
- 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B
^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))
/d^2)*log(((B*a + A*b)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B -
A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b
^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) + ((A^3 - A*B^2)*a^3 - (5*A^2*B - B
^3)*a^2*b - (A^3 - 5*A*B^2)*a*b^2 + (A^2*B - B^3)*b^3)*d)*sqrt(-(2*A*B*a^2
- 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4
- 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B
- A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2) + ((A^4 - B^4)*a^
4 - 4*(A^3*B + A*B^3)*a^3*b - 4*(A^3*B - A*B^3)*a*b^3 - (A^4 - B^4)*b^4)*s
qrt(tan(d*x + c))) - d*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b +
d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4
- 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 +
B^4)*b^4)/d^4))/d^2)*log(-((B*a + A*b)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)
*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A
^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) + ((A^3 - A*B^2)*a
^3 - (5*A^2*B - B^3)*a^2*b - (A^3 - 5*A*B^2)*a*b^2 + (A^2*B - B^3)*b^3)*d)
*sqrt(-(2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^
2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)...

```

3.576.6 Sympy [F]

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))(A+B\tan(c+dx)) dx$$

$$= \int (A+B\tan(c+dx))(a+b\tan(c+dx))\sqrt{\cot(c+dx)} dx$$

input `integrate(cot(d*x+c)**(1/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))*sqrt(cot(c + d*x)), x)`

3.576.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.87

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))(A+B\tan(c+dx)) dx =$$

$$2\sqrt{2}((A+B)a+(A-B)b)\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}+\frac{2}{\sqrt{\tan(dx+c)}}\right)\right)+2\sqrt{2}((A+B)a+(A-B)b)\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}-\frac{2}{\sqrt{\tan(dx+c)}}\right)\right)-\sqrt{2}((A-B)a-(A+B)b)\log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}}+1\right)+\sqrt{2}((A-B)a-(A+B)b)\log\left(-\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}}+1\right)-8Bb\sqrt{\tan(dx+c)}/d$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/4*(2*sqrt(2)*((A + B)*a + (A - B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A + B)*a + (A - B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*((A - B)*a - (A + B)*b)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*((A - B)*a - (A + B)*b)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - 8*B*b*sqrt(tan(d*x + c))/d`

3.576.8 Giac [F]

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))(A+B\tan(c+dx)) dx$$

$$= \int (B\tan(dx+c)+A)(b\tan(dx+c)+a)\sqrt{\cot(dx+c)} dx$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)*sqrt(cot(d*x + c)), x)`

3.576.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))(A+B \tan(c+dx)) dx$$

$$= \int \sqrt{\cot(c+dx)}(A+B \tan(c+dx))(a+b \tan(c+dx)) dx$$

input `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)),x)`output `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x)), x)`

3.577 $\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

3.577.1 Optimal result 5443
 3.577.2 Mathematica [A] (verified) 5444
 3.577.3 Rubi [A] (verified) 5444
 3.577.4 Maple [A] (verified) 5449
 3.577.5 Fricas [B] (verification not implemented) 5449
 3.577.6 Sympy [F] 5450
 3.577.7 Maxima [A] (verification not implemented) 5451
 3.577.8 Giac [F] 5451
 3.577.9 Mupad [F(-1)] 5452

3.577.1 Optimal result

Integrand size = 31, antiderivative size = 229

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \frac{(a(A - B) - b(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(a(A - B) - b(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d} + \frac{2bB}{3d \cot^{\frac{3}{2}}(c + dx)}$$

$$+ \frac{2(Ab + aB)}{d\sqrt{\cot(c + dx)}} + \frac{(b(A - B) + a(A + B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(b(A - B) + a(A + B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

```
output 2/3*b*B/d/cot(d*x+c)^(3/2)-1/2*(a*(A-B)-b*(A+B))*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/2*(a*(A-B)-b*(A+B))*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+1/4*(b*(A-B)+a*(A+B))*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/4*(b*(A-B)+a*(A+B))*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+2*(A*b+B*a)/d/cot(d*x+c)^(1/2)
```

3.577.2 Mathematica [A] (verified)

Time = 0.56 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.86

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx =$$

$$\frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(6\sqrt{2}(a(A - B) - b(A + B)) \left(\arctan \left(1 - \sqrt{2} \sqrt{\tan(c + dx)} \right) \right) - \arctan \left(1 + \sqrt{2} \sqrt{\tan(c + dx)} \right) \right)}{\sqrt{\cot(c + dx)}}$$

input `Integrate[((a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `-1/12*(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(6*Sqrt[2]*(a*(A - B) - b*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])] - 3*Sqrt[2]*(b*(A - B) + a*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]) - 24*(A*b + a*B)*Sqrt[Tan[c + d*x]] - 8*b*B*Tan[c + d*x]^(3/2))/d`

3.577.3 Rubi [A] (verified)Time = 0.80 (sec) , antiderivative size = 206, normalized size of antiderivative = 0.90, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.548$, Rules used = {3042, 4064, 3042, 4074, 3042, 4012, 3042, 4017, 25, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{(a \cot(c + dx) + b)(A \cot(c + dx) + B)}{\cot^{\frac{5}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

3.577. $\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

$$\begin{aligned}
& \int \frac{(b - a \tan(c + dx + \frac{\pi}{2})) (B - A \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{5/2}} dx \\
& \quad \downarrow 4074 \\
& \int \frac{Ab + aB + (aA - bB) \cot(c + dx)}{\cot^{3/2}(c + dx)} dx + \frac{2bB}{3d \cot^{3/2}(c + dx)} \\
& \quad \downarrow 3042 \\
& \int \frac{Ab + aB - (aA - bB) \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}} dx + \frac{2bB}{3d \cot^{3/2}(c + dx)} \\
& \quad \downarrow 4012 \\
& \int \frac{aA - bB - (Ab + aB) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx + \frac{2(aB + Ab)}{d\sqrt{\cot(c + dx)}} + \frac{2bB}{3d \cot^{3/2}(c + dx)} \\
& \quad \downarrow 3042 \\
& \int \frac{aA - bB - (-Ab - aB) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx + \frac{2(aB + Ab)}{d\sqrt{\cot(c + dx)}} + \frac{2bB}{3d \cot^{3/2}(c + dx)} \\
& \quad \downarrow 4017 \\
& \frac{2 \int -\frac{aA - bB - (Ab + aB) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} + \frac{2(aB + Ab)}{d\sqrt{\cot(c + dx)}} + \frac{2bB}{3d \cot^{3/2}(c + dx)} \\
& \quad \downarrow 25 \\
& -\frac{2 \int \frac{aA - bB - (Ab + aB) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} + \frac{2(aB + Ab)}{d\sqrt{\cot(c + dx)}} + \frac{2bB}{3d \cot^{3/2}(c + dx)} \\
& \quad \downarrow 1482 \\
& \frac{2\left(-\frac{1}{2}(a(A + B) + b(A - B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \frac{1}{2}(a(A - B) - b(A + B)) \int \frac{\cot(c + dx) + 1}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}\right)}{d} \\
& \quad + \frac{2(aB + Ab)}{d\sqrt{\cot(c + dx)}} + \frac{2bB}{3d \cot^{3/2}(c + dx)} \\
& \quad \downarrow 1476 \\
& \frac{2\left(-\frac{1}{2}(a(A + B) + b(A - B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \frac{1}{2}(a(A - B) - b(A + B)) \left(\frac{1}{2} \int \frac{1}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx)}} dx\right)\right)}{d} \\
& \quad + \frac{2(aB + Ab)}{d\sqrt{\cot(c + dx)}} + \frac{2bB}{3d \cot^{3/2}(c + dx)}
\end{aligned}$$

3.577. $\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$

↓ 1082

$$\frac{2 \left(-\frac{1}{2}(a(A+B) + b(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d\frac{(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{d} + \frac{2(aB + Ab)}{d\sqrt{\cot(c+dx)}} + \frac{2bB}{3d \cot^{\frac{3}{2}}(c+dx)}$$

↓ 217

$$\frac{2 \left(-\frac{1}{2}(a(A+B) + b(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} \right) \right)}{d} + \frac{2(aB + Ab)}{d\sqrt{\cot(c+dx)}} + \frac{2bB}{3d \cot^{\frac{3}{2}}(c+dx)}$$

↓ 1479

$$\frac{2 \left(-\frac{1}{2}(a(A+B) + b(A-B)) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) \right)}{d} + \frac{2(aB + Ab)}{d\sqrt{\cot(c+dx)}} + \frac{2bB}{3d \cot^{\frac{3}{2}}(c+dx)}$$

↓ 25

$$\frac{2 \left(-\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) \right) - \frac{1}{2}(a(A-B) - b(A+B)) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} \right)}{d} + \frac{2(aB + Ab)}{d\sqrt{\cot(c+dx)}} + \frac{2bB}{3d \cot^{\frac{3}{2}}(c+dx)}$$

↓ 27

$$\frac{2 \left(-\frac{1}{2}(a(A+B) + b(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{d} + \frac{2(aB + Ab)}{d\sqrt{\cot(c+dx)}} + \frac{2bB}{3d \cot^{\frac{3}{2}}(c+dx)}$$

↓ 1103

3.577. $\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

$$\frac{2\left(-\frac{1}{2}(a(A-B) - b(A+B))\left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right) - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}}\right) - \frac{1}{2}(a(A+B) + b(A-B))\left(\frac{1}{\sqrt{2}}\right)\right)}{d} + \frac{2(aB + Ab)}{d\sqrt{\cot(c+dx)}} + \frac{2bB}{3d\cot^{\frac{3}{2}}(c+dx)}$$

input `Int[((a + b*Tan[c + d*x])*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `(2*b*B)/(3*d*Cot[c + d*x]^(3/2)) + (2*(A*b + a*B))/(d*Sqrt[Cot[c + d*x]]) + (2*(-1/2*((a*(A - B) - b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2])) - ((b*(A - B) + a*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d`

3.577.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

3.577. $\int \frac{(a+b \tan(c+dx))(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

rule 1476 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2(d/e), 2]\}, \text{Simp}[e/(2c) \text{Int}[1/\text{Simp}[d/e + qx + x^2, x], x], x] + \text{Simp}[e/(2c) \text{Int}[1/\text{Simp}[d/e - qx + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c d^2 - a e^2, 0] \&\& \text{PosQ}[d e]$

rule 1479 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2(d/e), 2]\}, \text{Simp}[e/(2c q) \text{Int}[(q - 2x)/\text{Simp}[d/e + qx - x^2, x], x], x] + \text{Simp}[e/(2c q) \text{Int}[(q + 2x)/\text{Simp}[d/e - qx - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c d^2 - a e^2, 0] \&\& \text{NegQ}[d e]$

rule 1482 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a c, 2]\}, \text{Simp}[(d q + a e)/(2 a c) \text{Int}[(q + c x^2)/(a + c x^4), x], x] + \text{Simp}[(d q - a e)/(2 a c) \text{Int}[(q - c x^2)/(a + c x^4), x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{NeQ}[c d^2 + a e^2, 0] \&\& \text{NeQ}[c d^2 - a e^2, 0] \&\& \text{NegQ}[(-a) c]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4012 $\text{Int}[\frac{(a_.) + (b_.)\tan[(e_.) + (f_.)x]}{(c_.) + (d_.)\tan[(e_.) + (f_.)x]}]^m, x_Symbol] \rightarrow \text{Simp}[(b c - a d) \frac{(a + b \tan[e + f x])^{m+1}}{(f(m+1)(a^2 + b^2))}, x] + \text{Simp}[1/(a^2 + b^2) \text{Int}[(a + b \tan[e + f x])^{m+1} \text{Simp}[a c + b d - (b c - a d) \tan[e + f x], x], x]] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{NeQ}[b c - a d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{LtQ}[m, -1]$

rule 4017 $\text{Int}[\frac{(c_.) + (d_.)\tan[(e_.) + (f_.)x]}{\sqrt{(b_.)\tan[(e_.) + (f_.)x]}}, x_Symbol] \rightarrow \text{Simp}[2/f \text{Subst}[\text{Int}[(b c + d x^2)/(b^2 + x^4), x], x, \text{Sqrt}[b \tan[e + f x]]], x] /; \text{FreeQ}[\{b, c, d, e, f\}, x] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0]$

rule 4064 $\text{Int}[(\cot[(e_.) + (f_.)x] (g_.)^p)^m \frac{(a_.) + (b_.)\tan[(e_.) + (f_.)x]}{(c_.) + (d_.)\tan[(e_.) + (f_.)x]}]^n, x_Symbol] \rightarrow \text{Simp}[g^{m+n} \text{Int}[(g \cot[e + f x])^{p-m-n} (b + a \cot[e + f x])^m (d + c \cot[e + f x])^n, x]] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \&\& !\text{IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

```
rule 4074 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b
*c - a*d)*(A*b - a*B)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2
))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*A*c
+ b*B*c + A*b*d - a*B*d - (A*b*c - a*B*c - a*A*d - b*B*d)*Tan[e + f*x], x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m
, -1] && NeQ[a^2 + b^2, 0]
```

3.577.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.97

method	result
derivativedivides	$-\frac{2(Ab+Ba)}{\sqrt{\cot(dx+c)}} - \frac{2Bb}{3 \cot(dx+c)^{\frac{3}{2}}} + \frac{(Aa-Bb)\sqrt{2} \left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right) + 2 \arctan\left(1+\sqrt{2}\sqrt{\cot(dx+c)}\right) + 2 \arctan(-1+\sqrt{2}\sqrt{\cot(dx+c)}) \right)}{4}$
default	$-\frac{2(Ab+Ba)}{\sqrt{\cot(dx+c)}} - \frac{2Bb}{3 \cot(dx+c)^{\frac{3}{2}}} + \frac{(Aa-Bb)\sqrt{2} \left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right) + 2 \arctan\left(1+\sqrt{2}\sqrt{\cot(dx+c)}\right) + 2 \arctan(-1+\sqrt{2}\sqrt{\cot(dx+c)}) \right)}{4}$

```
input int((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2), x, method=_RETURNVER
BOSE)
```

```
output -1/d*(-2*(A*b+B*a)/cot(d*x+c)^(1/2)-2/3*B*b/cot(d*x+c)^(3/2)+1/4*(A*a-B*b)
*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)
*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/
2)*cot(d*x+c)^(1/2)))+1/4*(-A*b-B*a)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot
(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)
*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))))
```

3.577.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2239 vs. 2(195) = 390.

Time = 0.70 (sec) , antiderivative size = 2239, normalized size of antiderivative = 9.78

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="fricas")`

output `1/6*(3*d*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4)))/d^2*log(((A*a - B*b)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) - ((A^2*B - B^3)*a^3 + (A^3 - 5*A*B^2)*a^2*b - (5*A^2*B - B^3)*a*b^2 - (A^3 - A*B^2)*b^3)*d)*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2) + ((A^4 - B^4)*a^4 - 4*(A^3*B + A*B^3)*a^3*b - 4*(A^3*B + A*B^3)*a*b^3 - (A^4 - B^4)*b^4)*sqrt(tan(d*x + c))) - 3*d*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4))/d^2*log(-((A*a - B*b)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4) - ((A^2*B - B^3)*a^3 + (A^3 - 5*A*B^2)*a^2*b - (5*A^2*B - B^3)*a*b^2 - (A^3 - A*B^2)*b^3)*d)*sqrt((2*A*B*a^2 - 2*A*B*b^2 + 2*(A^2 - B^2)*a*b + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^4 - 8*(A^3*B - A*B^3)*a^3*b - 2*(A^4 - 10*A^2*B^2 + B^4)*a^2*b^2 + 8*(A^3*B - A*B^3)*a*b^3 + (A^4 - 2*A^2*B^2 + B^4)*b^4)/d^4)))/d^2)`

3.577.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))/sqrt(cot(c + d*x)), x)`

3.577.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.86

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx =$$

$$\frac{6\sqrt{2}((A - B)a - (A + B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 6\sqrt{2}((A - B)a - (A + B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 3\sqrt{2}((A + B)a + (A - B)b) \log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + \frac{1}{\tan(dx+c)} + 1\right) - 3\sqrt{2}((A + B)a + (A - B)b) \log\left(-\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + \frac{1}{\tan(dx+c)} + 1\right) - 8(Bb + 3(Ba + Ab)/\tan(dx+c))\tan(dx+c)^{3/2}}{d}$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="maxima")`

output `-1/12*(6*sqrt(2)*((A - B)*a - (A + B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 6*sqrt(2)*((A - B)*a - (A + B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + 3*sqrt(2)*((A + B)*a + (A - B)*b)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - 3*sqrt(2)*((A + B)*a + (A - B)*b)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - 8*(B*b + 3*(B*a + A*b)/tan(d*x + c))*tan(d*x + c)^(3/2))/d`

3.577.8 Giac [F]

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+b*tan(d*x+c))*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)/sqrt(cot(d*x + c)), x)`

3.577.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x)))/cot(c + d*x)^(1/2),x)`output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x)))/cot(c + d*x)^(1/2), x)`

3.578 $\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.578.1 Optimal result	5453
3.578.2 Mathematica [A] (verified)	5454
3.578.3 Rubi [A] (verified)	5454
3.578.4 Maple [A] (verified)	5461
3.578.5 Fricas [B] (verification not implemented)	5461
3.578.6 Sympy [F(-1)]	5462
3.578.7 Maxima [A] (verification not implemented)	5463
3.578.8 Giac [F]	5463
3.578.9 Mupad [F(-1)]	5464

3.578.1 Optimal result

Integrand size = 33, antiderivative size = 326

$$\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \frac{(a^2(A-B) - b^2(A-B) - 2ab(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(a^2(A-B) - b^2(A-B) - 2ab(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{2(a^2A - Ab^2 - 2abB) \sqrt{\cot(c+dx)}}{d}$$

$$- \frac{2a(7Ab + 5aB) \cot^{\frac{3}{2}}(c+dx)}{15d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)(b + a \cot(c+dx))}{5d}$$

$$+ \frac{(2ab(A-B) + a^2(A+B) - b^2(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(2ab(A-B) + a^2(A+B) - b^2(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

output
$$\begin{aligned} & -2/15*a*(7*A*b+5*B*a)*\cot(d*x+c)^{(3/2)}/d-2/5*a*A*\cot(d*x+c)^{(3/2)}*(b+a*\cot \\ & (d*x+c))/d-1/2*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*\arctan(-1+2^{(1/2)}*\cot(d*x \\ & +c)^{(1/2)})/d*2^{(1/2)}-1/2*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*\arctan(1+2^{(1/2)} \\ &)*\cot(d*x+c)^{(1/2)})/d*2^{(1/2)}+1/4*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*\ln(1+c \\ & \ot(d*x+c)-2^{(1/2)}*\cot(d*x+c)^{(1/2)})/d*2^{(1/2)}-1/4*(2*a*b*(A-B)+a^2*(A+B)-b \\ & ^2*(A+B))*\ln(1+\cot(d*x+c)+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/d*2^{(1/2)}+2*(A*a^2-A*b \\ & ^2-2*B*a*b)*\cot(d*x+c)^{(1/2)}/d \end{aligned}$$

3.578.2 Mathematica [A] (verified)

Time = 2.02 (sec) , antiderivative size = 255, normalized size of antiderivative = 0.78

$$\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx =$$

$$\frac{\sqrt{\cot(c+dx)} \left(30\sqrt{2}(a^2(A-B) + b^2(-A+B) - 2ab(A+B)) \left(\arctan \left(1 - \sqrt{2}\sqrt{\tan(c+dx)} \right) \right) - \arctan \left(1 + \sqrt{2}\sqrt{\tan(c+dx)} \right) \right)}{d}$$

input `Integrate[Cot[c + d*x]^(7/2)*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output
$$\begin{aligned} & -1/60*(\text{Sqrt}[\text{Cot}[c + d*x]]*(30*\text{Sqrt}[2]*(a^2*(A - B) + b^2*(-A + B) - 2*a*b* \\ & (A + B))*(\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]] - \text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt} \\ & [\text{Tan}[c + d*x]]) - 15*\text{Sqrt}[2]*(2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))* \\ & (\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]] - \text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt} \\ & [\text{Tan}[c + d*x]] + \text{Tan}[c + d*x])) + (24*a^2*A)/\text{Tan}[c + d*x]^{(5/2)} + (40*a*(2 \\ & *A*b + a*B))/\text{Tan}[c + d*x]^{(3/2)} - (120*(a^2*A - A*b^2 - 2*a*b*B))/\text{Sqrt}[\text{Tan} \\ & [c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]/d \end{aligned}$$

3.578.3 Rubi [A] (verified)

Time = 1.12 (sec) , antiderivative size = 283, normalized size of antiderivative = 0.87, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 4064, 3042, 4090, 27, 3042, 4113, 3042, 4011, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.578. $\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx \\
& \quad \downarrow \text{3042} \\
& \int \cot(c+dx)^{7/2}(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx \\
& \quad \downarrow \text{4064} \\
& \int \sqrt{\cot(c+dx)}(a \cot(c+dx)+b)^2(A \cot(c+dx)+B) dx \\
& \quad \downarrow \text{3042} \\
& \int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(b-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)^2\left(B-A \tan\left(c+dx+\frac{\pi}{2}\right)\right) dx \\
& \quad \downarrow \text{4090} \\
& -\frac{2}{5} \int \frac{1}{2} \sqrt{\cot(c+dx)}\left(-a(7Ab+5aB) \cot^2(c+dx)+5\left(Aa^2-2bBa-Ab^2\right) \cot(c+dx)+b(3aA-5bB)\right) dx - \\
& \quad \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx)+b)}{5d} \\
& \quad \downarrow \text{27} \\
& -\frac{1}{5} \int \sqrt{\cot(c+dx)}\left(-a(7Ab+5aB) \cot^2(c+dx)+5\left(Aa^2-2bBa-Ab^2\right) \cot(c+dx)+b(3aA-5bB)\right) dx - \\
& \quad \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx)+b)}{5d} \\
& \quad \downarrow \text{3042} \\
& -\frac{1}{5} \int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(-a(7Ab+5aB) \tan\left(c+dx+\frac{\pi}{2}\right)^2-5\left(Aa^2-2bBa-Ab^2\right) \tan\left(c+dx+\frac{\pi}{2}\right)+b\right) dx - \\
& \quad \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx)+b)}{5d} \\
& \quad \downarrow \text{4113} \\
& \frac{1}{5} \left(- \int \sqrt{\cot(c+dx)}\left(5\left(Ba^2+2Aba-b^2B\right)+5\left(Aa^2-2bBa-Ab^2\right) \cot(c+dx)\right) dx - \frac{2a(5aB+7Ab) \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx)+b)}{3d} \right) \\
& \quad \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx)+b)}{5d} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.578. $\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

$$\frac{1}{5} \left(- \int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)} \left(5(Ba^2+2Aba-b^2B) - 5(Aa^2-2bBa-Ab^2) \tan\left(c+dx+\frac{\pi}{2}\right) \right) dx - \frac{2a(5aB}{5d} \right. \\ \left. \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx)+b)}{5d} \right) \\ \downarrow \text{4011}$$

$$\frac{1}{5} \left(- \int \frac{5(Ba^2+2Aba-b^2B) \cot(c+dx) - 5(Aa^2-2bBa-Ab^2)}{\sqrt{\cot(c+dx)}} dx + \frac{10(a^2A-2abB-Ab^2) \sqrt{\cot(c+dx)}}{d} \right. \\ \left. \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx)+b)}{5d} \right) \\ \downarrow \text{3042}$$

$$\frac{1}{5} \left(- \int \frac{-5(Aa^2-2bBa-Ab^2) - 5(Ba^2+2Aba-b^2B) \tan\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}} dx + \frac{10(a^2A-2abB-Ab^2) \sqrt{\cot(c+dx)}}{d} \right. \\ \left. \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx)+b)}{5d} \right) \\ \downarrow \text{4017}$$

$$\frac{1}{5} \left(- \frac{2 \int \frac{5(Aa^2-2bBa-Ab^2-(Ba^2+2Aba-b^2B) \cot(c+dx))}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)}}{d} + \frac{10(a^2A-2abB-Ab^2) \sqrt{\cot(c+dx)}}{d} - \frac{2a(5aB}{5d} \right. \\ \left. \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx)+b)}{5d} \right) \\ \downarrow \text{27}$$

$$\frac{1}{5} \left(- \frac{10 \int \frac{Aa^2-2bBa-Ab^2-(Ba^2+2Aba-b^2B) \cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)}}{d} + \frac{10(a^2A-2abB-Ab^2) \sqrt{\cot(c+dx)}}{d} - \frac{2a(5aB}{5d} \right. \\ \left. \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx)+b)}{5d} \right) \\ \downarrow \text{1482}$$

$$\frac{1}{5} \left(- \frac{10 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)} + \frac{1}{2}(a^2(A-B) - 2ab(A+B) - \right. \right. \\ \left. \left. \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx)+b)}{5d} \right)}{d} \right)$$

3.578. $\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

↓ 1476

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} (a^2(A-B) - 2ab(A+B)) \right)}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)}{5d} \right)$$

↓ 1082

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} (a^2(A-B) - 2ab(A+B)) \right)}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)}{5d} \right)$$

↓ 217

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} (a^2(A-B) - 2ab(A+B)) \right)}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)}{5d} \right)$$

↓ 1479

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(- \frac{\int - \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int - \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right)}{d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)}{5d} \right)$$

↓ 25

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \right) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} \frac{1}{2\sqrt{2}} \right)}{\frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)}{5d}} \right)$$

↓ 27

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \right) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} \right)}{\frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)}{5d}} \right)$$

↓ 1103

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A-B)) \right) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B))}{d}}{\frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)}{5d}} \right)$$

input `Int[Cot[c + d*x]^(7/2)*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(-2*a*A*Cot[c + d*x]^(3/2)*(b + a*Cot[c + d*x]))/(5*d) + ((10*(a^2*A - A*b^2 - 2*a*b*B)*Sqrt[Cot[c + d*x]])/d - (2*a*(7*A*b + 5*a*B)*Cot[c + d*x]^(3/2))/(3*d) - (10*(((a^2*(A - B) - b^2*(A - B) - 2*a*b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]))/2 + ((2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d)/5`

3.578.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Int[(a + b*Tan[e + f*x])^(m - 1)*Simp[a*c - b*d + (b*c + a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]`

rule 4017 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)])*(g_.)^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

3.578.4 Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 293, normalized size of antiderivative = 0.90

method	result
derivativedivides	$-\frac{2Aa^2 \cot(dx+c)^{\frac{5}{2}}}{5} + \frac{4Aab \cot(dx+c)^{\frac{3}{2}}}{3} + \frac{2Ba^2 \cot(dx+c)^{\frac{3}{2}}}{3} - 2Aa^2 \sqrt{\cot(dx+c)} + 2Ab^2 \sqrt{\cot(dx+c)} + 4Bab \sqrt{\cot(dx+c)} +$
default	$-\frac{2Aa^2 \cot(dx+c)^{\frac{5}{2}}}{5} + \frac{4Aab \cot(dx+c)^{\frac{3}{2}}}{3} + \frac{2Ba^2 \cot(dx+c)^{\frac{3}{2}}}{3} - 2Aa^2 \sqrt{\cot(dx+c)} + 2Ab^2 \sqrt{\cot(dx+c)} + 4Bab \sqrt{\cot(dx+c)} +$

```
input int(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output -1/d*(2/5*A*a^2*cot(d*x+c)^(5/2)+4/3*A*a*b*cot(d*x+c)^(3/2)+2/3*B*a^2*cot(
d*x+c)^(3/2)-2*A*a^2*cot(d*x+c)^(1/2)+2*A*b^2*cot(d*x+c)^(1/2)+4*B*a*b*cot
(d*x+c)^(1/2)+1/4*(A*a^2-A*b^2-2*B*a*b)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*
cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1
/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/4*(-2*A*a*b
-B*a^2+B*b^2)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d
*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*ar
ctan(-1+2^(1/2)*cot(d*x+c)^(1/2))))
```

3.578.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4325 vs. 2(288) = 576.

Time = 2.27 (sec) , antiderivative size = 4325, normalized size of antiderivative = 13.27

$$\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorith
m="fricas")
```

```

output -1/30*(15*d*sqrt((2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a
^3*b - 4*(A^2 - B^2)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(
A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B
- A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B
- A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A
*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2)*log(((A*a^2 - 2*B*a*
b - A*b^2)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7
*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3
+ 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5
- 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4
- 2*A^2*B^2 + B^4)*b^8)/d^4) - ((A^2*B - B^3)*a^6 + 2*(A^3 - 5*A*B^2)*a^5
*b - (23*A^2*B - 7*B^3)*a^4*b^2 - 4*(3*A^3 - 7*A*B^2)*a^3*b^3 + (23*A^2*B
- 7*B^3)*a^2*b^4 + 2*(A^3 - 5*A*B^2)*a*b^5 - (A^2*B - B^3)*b^6)*d)*sqrt((2
*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2
)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*
b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 +
2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 -
4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4
- 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2) - ((A^4 - B^4)*a^8 - 8*(A^3*B + A*B^3)*
a^7*b - 4*(A^4 - B^4)*a^6*b^2 - 8*(A^3*B + A*B^3)*a^5*b^3 - 10*(A^4 - B...

```

3.578.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \text{Timed out}$$

```
input integrate(cot(d*x+c)**(7/2)*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)
```

```
output Timed out
```

3.578.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 279, normalized size of antiderivative = 0.86

$$\int \cot^{\frac{7}{2}}(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx =$$

$$30\sqrt{2}((A-B)a^2 - 2(A+B)ab - (A-B)b^2) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 30\sqrt{2}((A-B)a^2 -$$

input `integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/60*(30*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 30*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + 15*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - 15*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + 24*A*a^2/tan(d*x + c)^(5/2) - 120*(A*a^2 - 2*B*a*b - A*b^2)/sqrt(tan(d*x + c)) + 40*(B*a^2 + 2*A*a*b)/tan(d*x + c)^(3/2))/d`

3.578.8 Giac [F]

$$\int \cot^{\frac{7}{2}}(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

$$= \int (B\tan(dx+c) + A)(b\tan(dx+c) + a)^2 \cot(dx+c)^{\frac{7}{2}} dx$$

input `integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^2*cot(d*x + c)^(7/2), x)`

3.578.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{7/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^2 dx$$

input `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)`output `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2, x)`

3.579 $\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.579.1 Optimal result	5465
3.579.2 Mathematica [A] (verified)	5466
3.579.3 Rubi [A] (verified)	5466
3.579.4 Maple [A] (verified)	5472
3.579.5 Fricas [B] (verification not implemented)	5473
3.579.6 Sympy [F(-1)]	5473
3.579.7 Maxima [A] (verification not implemented)	5474
3.579.8 Giac [F]	5474
3.579.9 Mupad [F(-1)]	5475

3.579.1 Optimal result

Integrand size = 33, antiderivative size = 294

$$\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= -\frac{(2ab(A-B) + a^2(A+B) - b^2(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(2ab(A-B) + a^2(A+B) - b^2(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$- \frac{2a(5Ab + 3aB)\sqrt{\cot(c+dx)}}{3d} - \frac{2aA\sqrt{\cot(c+dx)}(b + a \cot(c+dx))}{3d}$$

$$+ \frac{(a^2(A-B) - b^2(A-B) - 2ab(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(a^2(A-B) - b^2(A-B) - 2ab(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

output

```
1/2*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/
d*2^(1/2)+1/2*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*arctan(1+2^(1/2)*cot(d*x+c
)^(1/2))/d*2^(1/2)+1/4*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*ln(1+cot(d*x+c)-2
^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/4*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*l
n(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-2/3*a*(5*A*b+3*B*a)*cot
(d*x+c)^(1/2)/d-2/3*a*A*(b+a*cot(d*x+c))*cot(d*x+c)^(1/2)/d
```

3.579. $\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.579.2 Mathematica [A] (verified)

Time = 1.29 (sec) , antiderivative size = 226, normalized size of antiderivative = 0.77

$$\int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

$$= \frac{\sqrt{\cot(c+dx)}\left(6\sqrt{2}(2ab(A-B)+a^2(A+B)-b^2(A+B))\left(\arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)\right)-\arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)\right)+3\sqrt{2}(a^2(A-B)+b^2(-A+B)-2ab(A+B))\left(\log\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)\right)+\log\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)-\frac{8a^2A}{\tan(c+dx)^{\frac{3}{2}}}-\frac{(24a(2Ab+a^2))\sqrt{\tan(c+dx)}}{12d}}{12d}$$

input `Integrate[Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*(6*Sqrt[2]*(2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]) + 3*Sqrt[2]*(a^2*(A - B) + b^2*(-A + B) - 2*a*b*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]) - (8*a^2*A)/Tan[c + d*x]^(3/2) - (24*a*(2*A*b + a^2))/Sqrt[Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/(12*d)`

3.579.3 Rubi [A] (verified)

Time = 0.96 (sec) , antiderivative size = 249, normalized size of antiderivative = 0.85, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 4064, 3042, 4090, 27, 3042, 4113, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{5/2}(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

$$\downarrow \text{4064}$$

$$\int \frac{(a\cot(c+dx)+b)^2(A\cot(c+dx)+B)}{\sqrt{\cot(c+dx)}}dx$$

$$\downarrow \text{3042}$$

3.579. $\int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$

$$\begin{aligned}
& \int \frac{(b - a \tan(c + dx + \frac{\pi}{2}))^2 (B - A \tan(c + dx + \frac{\pi}{2}))}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx \\
& \quad \downarrow \text{4090} \\
& -\frac{2}{3} \int \frac{-a(5Ab + 3aB) \cot^2(c + dx) + 3(Aa^2 - 2bBa - Ab^2) \cot(c + dx) + b(aA - 3bB)}{2\sqrt{\cot(c + dx)}} dx - \\
& \quad \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)}{3d} \\
& \quad \downarrow \text{27} \\
& -\frac{1}{3} \int \frac{-a(5Ab + 3aB) \cot^2(c + dx) + 3(Aa^2 - 2bBa - Ab^2) \cot(c + dx) + b(aA - 3bB)}{\sqrt{\cot(c + dx)}} dx - \\
& \quad \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)}{3d} \\
& \quad \downarrow \text{3042} \\
& -\frac{1}{3} \int \frac{-a(5Ab + 3aB) \tan(c + dx + \frac{\pi}{2})^2 - 3(Aa^2 - 2bBa - Ab^2) \tan(c + dx + \frac{\pi}{2}) + b(aA - 3bB)}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx - \\
& \quad \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)}{3d} \\
& \quad \downarrow \text{4113} \\
& \frac{1}{3} \left(- \int \frac{3(Ba^2 + 2Aba - b^2B) + 3(Aa^2 - 2bBa - Ab^2) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx - \frac{2a(3aB + 5Ab)\sqrt{\cot(c + dx)}}{d} \right) - \\
& \quad \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)}{3d} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \left(- \int \frac{3(Ba^2 + 2Aba - b^2B) - 3(Aa^2 - 2bBa - Ab^2) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx - \frac{2a(3aB + 5Ab)\sqrt{\cot(c + dx)}}{d} \right) - \\
& \quad \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)}{3d} \\
& \quad \downarrow \text{4017}
\end{aligned}$$

3.579. $\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\frac{1}{3} \left(\frac{2 \int -\frac{3(Ba^2+2Aba-b^2B+(Aa^2-2bBa-Ab^2)\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} - \frac{2a(3aB+5Ab)\sqrt{\cot(c+dx)}}{d} \right) -$$

$$\frac{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx)+b)}{3d}$$

↓ 27

$$\frac{1}{3} \left(\frac{6 \int \frac{Ba^2+2Aba-b^2B+(Aa^2-2bBa-Ab^2)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} - \frac{2a(3aB+5Ab)\sqrt{\cot(c+dx)}}{d} \right) -$$

$$\frac{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx)+b)}{3d}$$

↓ 1482

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^2(A+B)+2ab(A-B)-b^2(A+B)) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a^2(A-B)-2ab(A+B)-b^2) \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right)}{d} \right) -$$

$$\frac{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx)+b)}{3d}$$

↓ 1476

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^2(A+B)+2ab(A-B)-b^2(A+B)) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{d} \right) -$$

$$\frac{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx)+b)}{3d}$$

↓ 1082

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^2(A+B)+2ab(A-B)-b^2(A+B)) \left(\int \frac{1}{-\cot(c+dx)-1} \frac{d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \int \frac{1}{-\cot(c+dx)-1} \frac{d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \right) -$$

$$\frac{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx)+b)}{3d}$$

↓ 217

3.579. $\int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx)) dx$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a^2(A-B) - b^2(A+B)) \right)}{d} - \frac{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx) + b)}{3d} \right)$$

↓ 1479

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a^2(A-B) - b^2(A+B)) \right)}{d} - \frac{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx) + b)}{3d} \right)$$

↓ 25

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a^2(A-B) - b^2(A+B)) \right)}{d} - \frac{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx) + b)}{3d} \right)$$

↓ 27

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a^2(A-B) - b^2(A+B)) \right)}{d} - \frac{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx) + b)}{3d} \right)$$

↓ 1103

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A+B)) \right)}{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx)+b)} - \frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A+B)) \right)}{3d}$$

input `Int[Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(-2*a*A*Sqrt[Cot[c + d*x]]*(b + a*Cot[c + d*x]))/(3*d) + ((-2*a*(5*A*b + 3*a*B)*Sqrt[Cot[c + d*x]])/d + (6*(((2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]))/2 - ((a^2*(A - B) - b^2*(A - B) - 2*a*b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2))/d)/3`

3.579.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

3.579. $\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

rule 1476 $\text{Int}[\{(d_)+(e_)(x_)^2\}/\{(a_)+(c_)(x_)^4\}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Simp}[e/(2*c) \text{Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Simp}[e/(2*c) \text{Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[d*e]$

rule 1479 $\text{Int}[\{(d_)+(e_)(x_)^2\}/\{(a_)+(c_)(x_)^4\}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2*(d/e), 2]\}, \text{Simp}[e/(2*c*q) \text{Int}[(q - 2*x)/\text{Simp}[d/e + q*x - x^2, x], x], x] + \text{Simp}[e/(2*c*q) \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{NegQ}[d*e]$

rule 1482 $\text{Int}[\{(d_)+(e_)(x_)^2\}/\{(a_)+(c_)(x_)^4\}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a*c, 2]\}, \text{Simp}[(d*q + a*e)/(2*a*c) \text{Int}[(q + c*x^2)/(a + c*x^4), x], x] + \text{Simp}[(d*q - a*e)/(2*a*c) \text{Int}[(q - c*x^2)/(a + c*x^4), x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{NeQ}[c*d^2 + a*e^2, 0] \&\& \text{NeQ}[c*d^2 - a*e^2, 0] \&\& \text{NegQ}[(-a)*c]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4017 $\text{Int}[\{(c_)+(d_)*\tan[(e_)+(f_)(x_)]\}/\text{Sqrt}[(b_)*\tan[(e_)+(f_)(x_)]], x_Symbol] \rightarrow \text{Simp}[2/f \text{Subst}[\text{Int}[(b*c + d*x^2)/(b^2 + x^4), x], x, \text{Sqrt}[b*\tan[e + f*x]]], x] /; \text{FreeQ}[\{b, c, d, e, f\}, x] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0]$

rule 4064 $\text{Int}[(\cot[(e_)+(f_)(x_)]*(g_))^p*\{(a_)+(b_)*\tan[(e_)+(f_)(x_)]\}^m*\{(c_)+(d_)*\tan[(e_)+(f_)(x_)]\}^n, x_Symbol] \rightarrow \text{Simp}[g^{m+n} \text{Int}[(g*\cot[e + f*x])^{p-m-n}*(b + a*\cot[e + f*x])^m*(d + c*\cot[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \&\& !\text{IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

3.579.4 Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 252, normalized size of antiderivative = 0.86

method	result
derivativedivides	$-\frac{2Aa^2 \cot(dx+c)^{\frac{3}{2}}}{3} + 4abA\sqrt{\cot(dx+c)} + 2Ba^2\sqrt{\cot(dx+c)} + \frac{(-2abA - Ba^2 + Bb^2)\sqrt{2} \left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right) \right)}{2}$
default	$-\frac{2Aa^2 \cot(dx+c)^{\frac{3}{2}}}{3} + 4abA\sqrt{\cot(dx+c)} + 2Ba^2\sqrt{\cot(dx+c)} + \frac{(-2abA - Ba^2 + Bb^2)\sqrt{2} \left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right) \right)}{2}$

input `int(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNV ERBOSE)`

output `-1/d*(2/3*A*a^2*cot(d*x+c)^(3/2)+4*a*b*A*cot(d*x+c)^(1/2)+2*B*a^2*cot(d*x+c)^(1/2)+1/4*(-2*A*a*b-B*a^2+B*b^2)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/4*(-A*a^2+A*b^2+2*B*a*b)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))`

3.579. $\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.579.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4305 vs. $2(258) = 516$.

Time = 2.31 (sec) , antiderivative size = 4305, normalized size of antiderivative = 14.64

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm
m="fricas")
```

```
output 1/6*(3*d*sqrt(-(2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3
*b - 4*(A^2 - B^2)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^
3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B -
A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B -
A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B
^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2)*log(((B*a^2 + 2*A*a*b
- B*b^2)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b
- 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 +
2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 -
4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 -
2*A^2*B^2 + B^4)*b^8)/d^4) + ((A^3 - A*B^2)*a^6 - 2*(5*A^2*B - B^3)*a^5*b
- (7*A^3 - 23*A*B^2)*a^4*b^2 + 4*(7*A^2*B - 3*B^3)*a^3*b^3 + (7*A^3 - 23*
A*B^2)*a^2*b^4 - 2*(5*A^2*B - B^3)*a*b^5 - (A^3 - A*B^2)*b^6)*d)*sqrt(-(2*
A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2)
*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b
- 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 +
2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 -
4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 -
2*A^2*B^2 + B^4)*b^8)/d^4))/d^2) - ((A^4 - B^4)*a^8 - 8*(A^3*B + A*B^3)*a
^7*b - 4*(A^4 - B^4)*a^6*b^2 - 8*(A^3*B + A*B^3)*a^5*b^3 - 10*(A^4 - B^...
```

3.579.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \text{Timed out}$$

```
input integrate(cot(d*x+c)**(5/2)*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)
```

3.579. $\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

output Timed out

3.579.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 252, normalized size of antiderivative = 0.86

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{6\sqrt{2}((A + B)a^2 + 2(A - B)ab - (A + B)b^2) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 6\sqrt{2}((A + B)a^2 + 2(A - B)ab - (A + B)b^2) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 3\sqrt{2}((A - B)a^2 - 2(A + B)ab + (A + B)b^2) \log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + 1\right) + 3\sqrt{2}((A - B)a^2 - 2(A + B)ab + (A + B)b^2) \log\left(-\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + 1\right) - 8Aa^2/\tan(dx+c)^{\frac{3}{2}} - 24(Ba^2 + 2Aab)/\sqrt{\tan(dx+c)}}{d}$$

input `integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/12*(6*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 6*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - 3*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + 3*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - 8*A*a^2/tan(d*x + c)^(3/2) - 24*(B*a^2 + 2*A*a*b)/sqrt(tan(d*x + c))/d`

3.579.8 Giac [F]

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^2 \cot(dx + c)^{\frac{5}{2}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^2*cot(d*x + c)^(5/2), x)`

3.579. $\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.579.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{5/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^2 dx$$

input `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)`output `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2, x)`

3.580 $\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.580.1 Optimal result	5476
3.580.2 Mathematica [A] (verified)	5477
3.580.3 Rubi [A] (verified)	5477
3.580.4 Maple [A] (verified)	5482
3.580.5 Fricas [B] (verification not implemented)	5483
3.580.6 Sympy [F]	5484
3.580.7 Maxima [A] (verification not implemented)	5484
3.580.8 Giac [F]	5485
3.580.9 Mupad [F(-1)]	5485

3.580.1 Optimal result

Integrand size = 33, antiderivative size = 276

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= -\frac{(a^2(A-B) - b^2(A-B) - 2ab(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(a^2(A-B) - b^2(A-B) - 2ab(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{2b^2B}{d\sqrt{\cot(c+dx)}} - \frac{2a^2A\sqrt{\cot(c+dx)}}{d}$$

$$- \frac{(2ab(A-B) + a^2(A+B) - b^2(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{(2ab(A-B) + a^2(A+B) - b^2(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

output

```
1/2*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/
d*2^(1/2)+1/2*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*arctan(1+2^(1/2)*cot(d*x+c
)^(1/2))/d*2^(1/2)-1/4*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*ln(1+cot(d*x+c)-2
^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+1/4*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*l
n(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+2*b^2*B/d/cot(d*x+c)^(1
/2)-2*a^2*A*cot(d*x+c)^(1/2)/d
```

3.580. $\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

3.580.2 Mathematica [A] (verified)

Time = 0.90 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.80

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

$$= \frac{\sqrt{\cot(c+dx)}\left(2\sqrt{2}(a^2(A-B)+b^2(-A+B)-2ab(A+B))\left(\arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)\right)-\arctan\right)}{4d}$$

input `Integrate[Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*(2*Sqrt[2]*(a^2*(A - B) + b^2*(-A + B) - 2*a*b*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]) - Sqrt[2]*(2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]) - (8*a^2*A)/Sqrt[Tan[c + d*x]] + 8*b^2*B*Sqrt[Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/(4*d)`

3.580.3 Rubi [A] (verified)

Time = 0.90 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.83, number of steps used = 18, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.515$, Rules used = {3042, 4064, 3042, 4087, 25, 3042, 4113, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

$$\downarrow 3042$$

$$\int \cot(c+dx)^{3/2}(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

$$\downarrow 4064$$

$$\int \frac{(a\cot(c+dx)+b)^2(A\cot(c+dx)+B)}{\cot^{\frac{3}{2}}(c+dx)}dx$$

$$\downarrow 3042$$

3.580. $\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$

$$\begin{aligned}
& \int \frac{(b - a \tan(c + dx + \frac{\pi}{2}))^2 (B - A \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}} dx \\
& \quad \downarrow 4087 \\
& \frac{2b^2 B}{d\sqrt{\cot(c + dx)}} - \int -\frac{a^2 A \cot^2(c + dx) + (Ba^2 + 2Aba - b^2 B) \cot(c + dx) + b(Ab + 2aB)}{\sqrt{\cot(c + dx)}} dx \\
& \quad \downarrow 25 \\
& \int \frac{a^2 A \cot^2(c + dx) + (Ba^2 + 2Aba - b^2 B) \cot(c + dx) + b(Ab + 2aB)}{\sqrt{\cot(c + dx)}} dx + \frac{2b^2 B}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow 3042 \\
& \int \frac{a^2 A \tan(c + dx + \frac{\pi}{2})^2 - (Ba^2 + 2Aba - b^2 B) \tan(c + dx + \frac{\pi}{2}) + b(Ab + 2aB)}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx + \\
& \quad \frac{2b^2 B}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow 4113 \\
& \int \frac{-Aa^2 + 2bBa + Ab^2 + (Ba^2 + 2Aba - b^2 B) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx - \frac{2a^2 A \sqrt{\cot(c + dx)}}{d} + \\
& \quad \frac{2b^2 B}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow 3042 \\
& \int \frac{-Aa^2 + 2bBa + Ab^2 - (Ba^2 + 2Aba - b^2 B) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx - \frac{2a^2 A \sqrt{\cot(c + dx)}}{d} + \\
& \quad \frac{2b^2 B}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow 4017 \\
& \frac{2 \int \frac{Aa^2 - 2bBa - Ab^2 - (Ba^2 + 2Aba - b^2 B) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} - \frac{2a^2 A \sqrt{\cot(c + dx)}}{d} + \frac{2b^2 B}{d\sqrt{\cot(c + dx)}} \\
& \quad \downarrow 1482 \\
& \frac{2 \left(\frac{1}{2} (a^2(A + B) + 2ab(A - B) - b^2(A + B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2} (a^2(A - B) - 2ab(A + B) - b^2(A - B)) \int \frac{1 + \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} \right)}{d} \\
& \quad + \frac{2a^2 A \sqrt{\cot(c + dx)}}{d} + \frac{2b^2 B}{d\sqrt{\cot(c + dx)}}
\end{aligned}$$

3.580. $\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2 (A + B \tan(c + dx)) dx$

$$\downarrow 1476$$

$$\frac{2\left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A+B)) \int \frac{1+\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{d} + \frac{2a^2A\sqrt{\cot(c+dx)}}{d} + \frac{2b^2B}{d\sqrt{\cot(c+dx)}}$$

$$\downarrow 1082$$

$$\frac{2\left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A+B)) \int \frac{1+\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{d} + \frac{2a^2A\sqrt{\cot(c+dx)}}{d} + \frac{2b^2B}{d\sqrt{\cot(c+dx)}}$$

$$\downarrow 217$$

$$\frac{2\left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A+B)) \int \frac{1+\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{d} + \frac{2a^2A\sqrt{\cot(c+dx)}}{d} + \frac{2b^2B}{d\sqrt{\cot(c+dx)}}$$

$$\downarrow 1479$$

$$\frac{2\left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(-\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}\right) + \frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}\right)\right)}{d} + \frac{2a^2A\sqrt{\cot(c+dx)}}{d} + \frac{2b^2B}{d\sqrt{\cot(c+dx)}}$$

$$\downarrow 25$$

$$\frac{2\left(\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}\right) + \frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}\right)\right)}{d} + \frac{2a^2A\sqrt{\cot(c+dx)}}{d} + \frac{2b^2B}{d\sqrt{\cot(c+dx)}}$$

$$\downarrow 27$$

3.580. $\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^2(A+B\tan(c+dx)) dx$

$$2 \left(\frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\int \frac{\sqrt{2-2\sqrt{\cot(c+dx)}}}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1}} dx \right) \right.$$

$$\frac{2a^2 A \sqrt{\cot(c+dx)}}{d} + \frac{2b^2 B}{d \sqrt{\cot(c+dx)}}$$

↓ 1103

$$2 \left(\frac{1}{2} (a^2(A-B) - 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} (a^2(A+B) + 2ab(A-B) - b^2(A+B)) \right)$$

$$\frac{2a^2 A \sqrt{\cot(c+dx)}}{d} + \frac{2b^2 B}{d \sqrt{\cot(c+dx)}}$$

input `Int[Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(2*b^2*B)/(d*Sqrt[Cot[c + d*x]]) - (2*a^2*A*Sqrt[Cot[c + d*x]])/d + (2*(((a^2*(A - B) - b^2*(A - B) - 2*a*b*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]))/2 + ((2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d`

3.580.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])]; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`


```
rule 4064 Int[(cot[(e_.) + (f_.)*(x_)]*(g_.))^(p_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(m_.))*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp [g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]
```

```
rule 4087 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Simp[(- (B*c - A*d))*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

```
rule 4113 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(m_.))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

3.580.4 Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 239, normalized size of antiderivative = 0.87

method	result
derivativedivides	$\frac{2Aa^2\sqrt{\cot(dx+c)} + \frac{(-Aa^2 + Ab^2 + 2Bab)\sqrt{2}\left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right) + 2\arctan\left(1+\sqrt{2}\sqrt{\cot(dx+c)}\right) + 2\arctan(-\dots)}{4}}{4}$
default	$\frac{2Aa^2\sqrt{\cot(dx+c)} + \frac{(-Aa^2 + Ab^2 + 2Bab)\sqrt{2}\left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right) + 2\arctan\left(1+\sqrt{2}\sqrt{\cot(dx+c)}\right) + 2\arctan(-\dots)}{4}}{4}$

```
input int(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)), x, method=_RETURNV ERBOSE)
```

$$3.580. \int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

output
$$-1/d*(2*A*a^2*cot(d*x+c)^(1/2)+1/4*(-A*a^2+A*b^2+2*B*a*b)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/4*(2*A*a*b+B*a^2-B*b^2)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))-2*B*b^2/cot(d*x+c)^(1/2))$$

3.580.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4248 vs. 2(244) = 488.

Time = 2.28 (sec) , antiderivative size = 4248, normalized size of antiderivative = 15.39

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output
$$\begin{aligned} & 1/2*(d*\sqrt{(2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b} \\ & - 4*(A^2 - B^2)*a*b^3 + d^2*\sqrt{-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B \\ & - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A* \\ & B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A* \\ & B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3) \\ & *a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4)/d^2)*\log(((A*a^2 - 2*B*a*b - A \\ & *b^2)*d^3*\sqrt{-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b - \\ & 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 + 2*(\\ & 19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 - 4*(\\ & 3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 - 2* \\ & A^2*B^2 + B^4)*b^8)/d^4) - ((A^2*B - B^3)*a^6 + 2*(A^3 - 5*A*B^2)*a^5*b - \\ & (23*A^2*B - 7*B^3)*a^4*b^2 - 4*(3*A^3 - 7*A*B^2)*a^3*b^3 + (23*A^2*B - 7*B \\ & ^3)*a^2*b^4 + 2*(A^3 - 5*A*B^2)*a*b^5 - (A^2*B - B^3)*b^6)*d)*\sqrt{(2*A*B* \\ & a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2)*a*b \\ & ^3 + d^2*\sqrt{-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*b - 4 \\ & *(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 + 2*(1 \\ & 9*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 - 4*(3 \\ & *A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4 - 2*A \\ & ^2*B^2 + B^4)*b^8)/d^4)/d^2) - ((A^4 - B^4)*a^8 - 8*(A^3*B + A*B^3)*a^7*b \\ & - 4*(A^4 - B^4)*a^6*b^2 - 8*(A^3*B + A*B^3)*a^5*b^3 - 10*(A^4 - B^4)*a \dots \end{aligned}$$

3.580.
$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

3.580.6 Sympy [F]

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^2 \cot^{\frac{3}{2}}(c + dx) dx$$

input `integrate(cot(d*x+c)**(3/2)*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**2*cot(c + d*x)**(3/2), x)`

3.580.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 244, normalized size of antiderivative = 0.88

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{8 B b^2 \sqrt{\tan(dx + c)} + 2 \sqrt{2}((A - B)a^2 - 2(A + B)ab - (A - B)b^2) \arctan\left(\frac{1}{2} \sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + \dots}{d}$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/4*(8*B*b^2*sqrt(tan(d*x + c)) + 2*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - 8*A*a^2/sqrt(tan(d*x + c)))/d`

3.580.8 Giac [F]

$$\begin{aligned} & \int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^2 \cot(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^2*cot(d*x + c)^(3/2), x)`

3.580.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx \\ &= \int \cot(c + dx)^{3/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^2 dx \end{aligned}$$

input `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)`

output `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2, x)`

3.581 $\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

3.581.1 Optimal result	5486
3.581.2 Mathematica [A] (verified)	5487
3.581.3 Rubi [A] (verified)	5487
3.581.4 Maple [A] (verified)	5493
3.581.5 Fricas [B] (verification not implemented)	5493
3.581.6 Sympy [F]	5494
3.581.7 Maxima [A] (verification not implemented)	5495
3.581.8 Giac [F]	5495
3.581.9 Mupad [F(-1)]	5496

3.581.1 Optimal result

Integrand size = 33, antiderivative size = 283

$$\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \frac{(2ab(A - B) + a^2(A + B) - b^2(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(2ab(A - B) + a^2(A + B) - b^2(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{2b^2B}{3d \cot^{\frac{3}{2}}(c + dx)} + \frac{2b(Ab + 2aB)}{d\sqrt{\cot(c + dx)}}$$

$$- \frac{(a^2(A - B) - b^2(A - B) - 2ab(A + B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{(a^2(A - B) - b^2(A - B) - 2ab(A + B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

output

```
2/3*b^2*B/d/cot(d*x+c)^(3/2)-1/2*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*arctan(
-1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/2*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B
))*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/4*(a^2*(A-B)-b^2*(A-B)-2
*a*b*(A+B))*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+1/4*(a^2*(
A-B)-b^2*(A-B)-2*a*b*(A+B))*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(
1/2)+2*b*(A*b+2*B*a)/d/cot(d*x+c)^(1/2)
```

3.581.2 Mathematica [A] (verified)

Time = 0.61 (sec) , antiderivative size = 226, normalized size of antiderivative = 0.80

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx =$$

$$\sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}\left(6\sqrt{2}(2ab(A-B)+a^2(A+B)-b^2(A+B))\left(\arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)\right)\right)$$

input `Integrate[Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `-1/12*(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(6*Sqrt[2]*(2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]) + 3*Sqrt[2]*(a^2*(A - B) + b^2*(-A + B) - 2*a*b*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]) - 24*b*(A*b + 2*a*B)*Sqrt[Tan[c + d*x]] - 8*b^2*B*Tan[c + d*x]^(3/2)))/d`

3.581.3 Rubi [A] (verified)

Time = 0.96 (sec) , antiderivative size = 235, normalized size of antiderivative = 0.83, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 4064, 3042, 4087, 25, 3042, 4111, 3042, 4017, 25, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

↓ 3042

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$$

↓ 4064

$$\int \frac{(a\cot(c+dx)+b)^2(A\cot(c+dx)+B)}{\cot^{\frac{5}{2}}(c+dx)}dx$$

↓ 3042

3.581. $\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx$

$$\begin{aligned}
& \int \frac{(b - a \tan(c + dx + \frac{\pi}{2}))^2 (B - A \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{5/2}} dx \\
& \quad \downarrow \text{4087} \\
& \frac{2b^2 B}{3d \cot^{\frac{3}{2}}(c + dx)} - \int -\frac{a^2 A \cot^2(c + dx) + (Ba^2 + 2Aba - b^2 B) \cot(c + dx) + b(Ab + 2aB)}{\cot^{\frac{3}{2}}(c + dx)} dx \\
& \quad \downarrow \text{25} \\
& \int \frac{a^2 A \cot^2(c + dx) + (Ba^2 + 2Aba - b^2 B) \cot(c + dx) + b(Ab + 2aB)}{\cot^{\frac{3}{2}}(c + dx)} dx + \frac{2b^2 B}{3d \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \int \frac{a^2 A \tan(c + dx + \frac{\pi}{2})^2 - (Ba^2 + 2Aba - b^2 B) \tan(c + dx + \frac{\pi}{2}) + b(Ab + 2aB)}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}} dx + \\
& \quad \frac{2b^2 B}{3d \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{4111} \\
& \int \frac{Ba^2 + 2Aba - b^2 B + (a^2 A - b(Ab + 2aB)) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx + \frac{2b(2aB + Ab)}{d\sqrt{\cot(c + dx)}} + \frac{2b^2 B}{3d \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{3042} \\
& \int \frac{Ba^2 + 2Aba - b^2 B - (a^2 A - b(Ab + 2aB)) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx + \frac{2b(2aB + Ab)}{d\sqrt{\cot(c + dx)}} + \\
& \quad \frac{2b^2 B}{3d \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{4017} \\
& \frac{2 \int -\frac{Ba^2 + 2Aba - b^2 B + (Aa^2 - 2bBa - Ab^2) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} + \frac{2b(2aB + Ab)}{d\sqrt{\cot(c + dx)}} + \frac{2b^2 B}{3d \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{25} \\
& \frac{2 \int \frac{Ba^2 + 2Aba - b^2 B + (Aa^2 - 2bBa - Ab^2) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} + \frac{2b(2aB + Ab)}{d\sqrt{\cot(c + dx)}} + \frac{2b^2 B}{3d \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow \text{1482}
\end{aligned}$$

3.581. $\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$

$$\frac{2\left(\frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A-B)) \int \frac{1+\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{d} \\ + \frac{2b(2aB + Ab)}{d\sqrt{\cot(c+dx)}} + \frac{2b^2B}{3d \cot^{\frac{3}{2}}(c+dx)} \\ \downarrow \text{1476}$$

$$\frac{2\left(\frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A-B)) \int \frac{1+\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{d} \\ + \frac{2b(2aB + Ab)}{d\sqrt{\cot(c+dx)}} + \frac{2b^2B}{3d \cot^{\frac{3}{2}}(c+dx)} \\ \downarrow \text{1082}$$

$$\frac{2\left(\frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A-B)) \int \frac{1+\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{d} \\ + \frac{2b(2aB + Ab)}{d\sqrt{\cot(c+dx)}} + \frac{2b^2B}{3d \cot^{\frac{3}{2}}(c+dx)} \\ \downarrow \text{217}$$

$$\frac{2\left(\frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A-B)) \int \frac{1+\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{d} \\ + \frac{2b(2aB + Ab)}{d\sqrt{\cot(c+dx)}} + \frac{2b^2B}{3d \cot^{\frac{3}{2}}(c+dx)} \\ \downarrow \text{1479}$$

$$\frac{2\left(\frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A-B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A-B)) \int \frac{1+\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{d} \\ + \frac{2b(2aB + Ab)}{d\sqrt{\cot(c+dx)}} + \frac{2b^2B}{3d \cot^{\frac{3}{2}}(c+dx)} \\ \downarrow \text{25}$$

3.581. $\int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$

$$2 \left(\frac{1}{2}(a^2(A - B) - 2ab(A + B) - b^2(A - B)) \right) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) d$$

$$\frac{2b(2aB + Ab)}{d\sqrt{\cot(c + dx)}} + \frac{2b^2B}{3d \cot^{\frac{3}{2}}(c + dx)}$$

27

$$2 \left(\frac{1}{2}(a^2(A - B) - 2ab(A + B) - b^2(A - B)) \right) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) d$$

$$\frac{2b(2aB + Ab)}{d\sqrt{\cot(c + dx)}} + \frac{2b^2B}{3d \cot^{\frac{3}{2}}(c + dx)}$$

1103

$$2 \left(\frac{1}{2}(a^2(A - B) - 2ab(A + B) - b^2(A - B)) \right) \left(\frac{\log(\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right) d$$

$$\frac{2b(2aB + Ab)}{d\sqrt{\cot(c + dx)}} + \frac{2b^2B}{3d \cot^{\frac{3}{2}}(c + dx)}$$

input `Int[Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]),x]`

output `(2*b^2*B)/(3*d*Cot[c + d*x]^(3/2)) + (2*b*(A*b + 2*a*B))/(d*Sqrt[Cot[c + d*x]]) + (2*(-1/2*((2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2])) + ((a^2*(A - B) - b^2*(A - B) - 2*a*b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d`

3.581.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4087 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^2*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(- (B*c - A*d))*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]`

rule 4111 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

3.581.4 Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 244, normalized size of antiderivative = 0.86

method	result
derivativedivides	$-\frac{2Bb^2}{3\cot(dx+c)^{\frac{3}{2}}}-\frac{2b(Ab+2Ba)}{\sqrt{\cot(dx+c)}}+\frac{(2abA+Ba^2-Bb^2)\sqrt{2}\left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right)+2\arctan\left(1+\sqrt{2}\sqrt{\cot(dx+c)}\right)+2\right)}{4}$
default	$-\frac{2Bb^2}{3\cot(dx+c)^{\frac{3}{2}}}-\frac{2b(Ab+2Ba)}{\sqrt{\cot(dx+c)}}+\frac{(2abA+Ba^2-Bb^2)\sqrt{2}\left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right)+2\arctan\left(1+\sqrt{2}\sqrt{\cot(dx+c)}\right)+2\right)}{4}$

```
input int(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output -1/d*(-2/3*B*b^2/cot(d*x+c)^(3/2)-2*b*(A*b+2*B*a)/cot(d*x+c)^(1/2)+1/4*(2*
A*a*b+B*a^2-B*b^2)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+
cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))
+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/4*(A*a^2-A*b^2-2*B*a*b)*2^(1/2)*
(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+
c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*
x+c)^(1/2))))
```

3.581.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4281 vs. 2(249) = 498.

Time = 2.33 (sec) , antiderivative size = 4281, normalized size of antiderivative = 15.13

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^2(A+B\tan(c+dx))dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x,algorith
m="fricas")
```

output

```

-1/6*(3*d*sqrt(-(2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^
3*b - 4*(A^2 - B^2)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A
^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B
- A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B
- A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*
B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2)*log(((B*a^2 + 2*A*a*b
- B*b^2)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*
b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 +
2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 -
4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4
- 2*A^2*B^2 + B^4)*b^8)/d^4) + ((A^3 - A*B^2)*a^6 - 2*(5*A^2*B - B^3)*a^5*
b - (7*A^3 - 23*A*B^2)*a^4*b^2 + 4*(7*A^2*B - 3*B^3)*a^3*b^3 + (7*A^3 - 23
*A*B^2)*a^2*b^4 - 2*(5*A^2*B - B^3)*a*b^5 - (A^3 - A*B^2)*b^6)*d)*sqrt(-(2
*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2
)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7*
b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 +
2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 -
4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4
- 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2) - ((A^4 - B^4)*a^8 - 8*(A^3*B + A*B^3)*
a^7*b - 4*(A^4 - B^4)*a^6*b^2 - 8*(A^3*B + A*B^3)*a^5*b^3 - 10*(A^4 - B...

```

3.581.6 Sympy [F]

$$\int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \int (A+B \tan(c+dx))(a+b \tan(c+dx))^2 \sqrt{\cot(c+dx)} dx$$

input `integrate(cot(d*x+c)**(1/2)*(a+b*tan(d*x+c))**2*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**2*sqrt(cot(c + d*x)), x)`

3.581.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.90

$$\int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx =$$

$$6 \sqrt{2}((A+B)a^2+2(A-B)ab-(A+B)b^2) \arctan\left(\frac{1}{2} \sqrt{2}\left(\sqrt{2}+\frac{2}{\sqrt{\tan(dx+c)}}\right)\right)+6 \sqrt{2}((A+B)a^2+2$$

```
input integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="maxima")
```

```
output -1/12*(6*sqrt(2)*((A+B)*a^2+2*(A-B)*a*b-(A+B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2)+2/sqrt(tan(d*x+c))))+6*sqrt(2)*((A+B)*a^2+2*(A-B)*a*b-(A+B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2)-2/sqrt(tan(d*x+c))))-3*sqrt(2)*((A-B)*a^2-2*(A+B)*a*b-(A-B)*b^2)*log(sqrt(2)/sqrt(tan(d*x+c))+1/tan(d*x+c)+1)+3*sqrt(2)*((A-B)*a^2-2*(A+B)*a*b-(A-B)*b^2)*log(-sqrt(2)/sqrt(tan(d*x+c))+1/tan(d*x+c)+1)-8*(B*b^2+3*(2*B*a*b+A*b^2)/tan(d*x+c))*tan(d*x+c)^(3/2)/d
```

3.581.8 Giac [F]

$$\int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^2(A+B \tan(c+dx)) dx$$

$$= \int (B \tan(dx+c)+A)(b \tan(dx+c)+a)^2 \sqrt{\cot(dx+c)} dx$$

```
input integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^2*(A+B*tan(d*x+c)),x, algorithm="giac")
```

```
output integrate((B*tan(d*x+c)+A)*(b*tan(d*x+c)+a)^2*sqrt(cot(d*x+c)),x)
```

3.581.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^2(A + B \tan(c + dx)) dx$$

$$= \int \sqrt{\cot(c + dx)}(A + B \tan(c + dx))(a + b \tan(c + dx))^2 dx$$

input `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2,x)`output `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2, x)`

3.582
$$\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.582.1 Optimal result 5497
 3.582.2 Mathematica [A] (verified) 5498
 3.582.3 Rubi [A] (verified) 5498
 3.582.4 Maple [A] (verified) 5504
 3.582.5 Fricas [B] (verification not implemented) 5505
 3.582.6 Sympy [F] 5505
 3.582.7 Maxima [A] (verification not implemented) 5506
 3.582.8 Giac [F] 5506
 3.582.9 Mupad [F(-1)] 5507

3.582.1 Optimal result

Integrand size = 33, antiderivative size = 317

$$\int \frac{(a + b \tan(c + dx))^2(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \frac{(a^2(A - B) - b^2(A - B) - 2ab(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(a^2(A - B) - b^2(A - B) - 2ab(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{2b^2B}{5d \cot^{5/2}(c + dx)} + \frac{2b(Ab + 2aB)}{3d \cot^{3/2}(c + dx)} + \frac{2(2aAb + a^2B - b^2B)}{d\sqrt{\cot(c + dx)}}$$

$$+ \frac{(2ab(A - B) + a^2(A + B) - b^2(A + B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(2ab(A - B) + a^2(A + B) - b^2(A + B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

output

```
2/5*b^2*B/d/cot(d*x+c)^(5/2)+2/3*b*(A*b+2*B*a)/d/cot(d*x+c)^(3/2)-1/2*(a^2
*(A-B)-b^2*(A-B)-2*a*b*(A+B))*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)
)-1/2*(a^2*(A-B)-b^2*(A-B)-2*a*b*(A+B))*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))
/d*2^(1/2)+1/4*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*ln(1+cot(d*x+c)-2^(1/2)*c
ot(d*x+c)^(1/2))/d*2^(1/2)-1/4*(2*a*b*(A-B)+a^2*(A+B)-b^2*(A+B))*ln(1+cot(
d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+2*(2*A*a*b+B*a^2-B*b^2)/d/cot(d
*x+c)^(1/2)
```


3.582.2 Mathematica [A] (verified)

Time = 1.23 (sec) , antiderivative size = 255, normalized size of antiderivative = 0.80

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx =$$

$$\frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(30\sqrt{2}(a^2(A - B) + b^2(-A + B) - 2ab(A + B)) \left(\arctan \left(1 - \sqrt{2} \sqrt{\tan(c + dx)} \right) \right) \right)}{d}$$

input `Integrate[((a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `-1/60*(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(30*Sqrt[2]*(a^2*(A - B) + b^2*(-A + B) - 2*a*b*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]) - 15*Sqrt[2]*(2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]) - 120*(2*a*A*b + a^2*B - b^2*B)*Sqrt[Tan[c + d*x]] - 40*b*(A*b + 2*a*B)*Tan[c + d*x]^(3/2) - 24*b^2*B*Tan[c + d*x]^(5/2))/d`

3.582.3 Rubi [A] (verified)

Time = 1.11 (sec) , antiderivative size = 269, normalized size of antiderivative = 0.85, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 4064, 3042, 4087, 25, 3042, 4111, 3042, 4012, 3042, 4017, 25, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{(a \cot(c + dx) + b)^2 (A \cot(c + dx) + B)}{\cot^{\frac{7}{2}}(c + dx)} dx$$

$$\begin{aligned}
& \int \frac{(b - a \tan(c + dx + \frac{\pi}{2}))^2 (B - A \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{7/2}} dx && \downarrow \text{3042} \\
& \frac{2b^2 B}{5d \cot^{\frac{5}{2}}(c + dx)} - \int \frac{a^2 A \cot^2(c + dx) + (Ba^2 + 2Aba - b^2 B) \cot(c + dx) + b(Ab + 2aB)}{\cot^{\frac{5}{2}}(c + dx)} dx && \downarrow \text{4087} \\
& \int \frac{a^2 A \cot^2(c + dx) + (Ba^2 + 2Aba - b^2 B) \cot(c + dx) + b(Ab + 2aB)}{\cot^{\frac{5}{2}}(c + dx)} dx + \frac{2b^2 B}{5d \cot^{\frac{5}{2}}(c + dx)} && \downarrow \text{25} \\
& \int \frac{a^2 A \tan(c + dx + \frac{\pi}{2})^2 - (Ba^2 + 2Aba - b^2 B) \tan(c + dx + \frac{\pi}{2}) + b(Ab + 2aB)}{(-\tan(c + dx + \frac{\pi}{2}))^{5/2}} dx + \frac{2b^2 B}{5d \cot^{\frac{5}{2}}(c + dx)} && \downarrow \text{3042} \\
& \int \frac{Ba^2 + 2Aba - b^2 B + (a^2 A - b(Ab + 2aB)) \cot(c + dx)}{\cot^{\frac{3}{2}}(c + dx)} dx + \frac{2b(2aB + Ab)}{3d \cot^{\frac{3}{2}}(c + dx)} + \frac{2b^2 B}{5d \cot^{\frac{5}{2}}(c + dx)} && \downarrow \text{4111} \\
& \int \frac{Ba^2 + 2Aba - b^2 B - (a^2 A - b(Ab + 2aB)) \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}} dx + \frac{2b(2aB + Ab)}{3d \cot^{\frac{3}{2}}(c + dx)} + \frac{2b^2 B}{5d \cot^{\frac{5}{2}}(c + dx)} && \downarrow \text{3042} \\
& \int \frac{Aa^2 - b(Ab + 2aB) - (Ba^2 + 2Aba - b^2 B) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx + \frac{2(a^2 B + 2aAb - b^2 B)}{d \sqrt{\cot(c + dx)}} + \frac{2b(2aB + Ab)}{3d \cot^{\frac{3}{2}}(c + dx)} + \frac{2b^2 B}{5d \cot^{\frac{5}{2}}(c + dx)} && \downarrow \text{4012} \\
& \int \frac{Aa^2 - b(Ab + 2aB) - (-Ba^2 - 2Aba + b^2 B) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx + \frac{2(a^2 B + 2aAb - b^2 B)}{d \sqrt{\cot(c + dx)}} + \frac{2b(2aB + Ab)}{3d \cot^{\frac{3}{2}}(c + dx)} + \frac{2b^2 B}{5d \cot^{\frac{5}{2}}(c + dx)} && \downarrow \text{3042}
\end{aligned}$$

3.582. $\int \frac{(a+b \tan(c+dx))^2 (A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

$$\begin{aligned}
& \downarrow 4017 \\
& \frac{2 \int -\frac{Aa^2-2bBa-Ab^2-(Ba^2+2Aba-b^2B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} + \frac{2(a^2B+2aAb-b^2B)}{d\sqrt{\cot(c+dx)}} + \\
& \quad \frac{2b(2aB+Ab)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{2b^2B}{5d \cot^{\frac{5}{2}}(c+dx)} \\
& \downarrow 25 \\
& \frac{2 \int \frac{Aa^2-2bBa-Ab^2-(Ba^2+2Aba-b^2B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} + \frac{2(a^2B+2aAb-b^2B)}{d\sqrt{\cot(c+dx)}} + \\
& \quad \frac{2b(2aB+Ab)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{2b^2B}{5d \cot^{\frac{5}{2}}(c+dx)} \\
& \downarrow 1482 \\
& \frac{2\left(-\frac{1}{2}(a^2(A+B)+2ab(A-B)-b^2(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a^2(A-B)-2ab(A+B)-b^2(A+B))\right)}{d} \\
& \quad + \frac{2(a^2B+2aAb-b^2B)}{d\sqrt{\cot(c+dx)}} + \frac{2b(2aB+Ab)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{2b^2B}{5d \cot^{\frac{5}{2}}(c+dx)} \\
& \downarrow 1476 \\
& \frac{2\left(-\frac{1}{2}(a^2(A+B)+2ab(A-B)-b^2(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a^2(A-B)-2ab(A+B)-b^2(A+B))\right)}{d} \\
& \quad + \frac{2(a^2B+2aAb-b^2B)}{d\sqrt{\cot(c+dx)}} + \frac{2b(2aB+Ab)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{2b^2B}{5d \cot^{\frac{5}{2}}(c+dx)} \\
& \downarrow 1082 \\
& \frac{2\left(-\frac{1}{2}(a^2(A+B)+2ab(A-B)-b^2(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a^2(A-B)-2ab(A+B)-b^2(A+B))\right)}{d} \\
& \quad + \frac{2(a^2B+2aAb-b^2B)}{d\sqrt{\cot(c+dx)}} + \frac{2b(2aB+Ab)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{2b^2B}{5d \cot^{\frac{5}{2}}(c+dx)} \\
& \downarrow 217 \\
& \frac{2\left(-\frac{1}{2}(a^2(A+B)+2ab(A-B)-b^2(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a^2(A-B)-2ab(A+B)-b^2(A+B))\right)}{d} \\
& \quad + \frac{2(a^2B+2aAb-b^2B)}{d\sqrt{\cot(c+dx)}} + \frac{2b(2aB+Ab)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{2b^2B}{5d \cot^{\frac{5}{2}}(c+dx)}
\end{aligned}$$

3.582. $\int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

$$\begin{aligned}
& \downarrow 1479 \\
& 2 \left(-\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}}{2\sqrt{2}} \right) \right. \\
& \hline
& \frac{2(a^2B + 2aAb - b^2B)}{d\sqrt{\cot(c+dx)}} + \frac{2b(2aB + Ab)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{2b^2B}{5d \cot^{\frac{5}{2}}(c+dx)} \\
& \downarrow 25 \\
& 2 \left(-\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right. \\
& \hline
& \frac{2(a^2B + 2aAb - b^2B)}{d\sqrt{\cot(c+dx)}} + \frac{2b(2aB + Ab)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{2b^2B}{5d \cot^{\frac{5}{2}}(c+dx)} \\
& \downarrow 27 \\
& 2 \left(-\frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} \right) \right. \\
& \hline
& \frac{2(a^2B + 2aAb - b^2B)}{d\sqrt{\cot(c+dx)}} + \frac{2b(2aB + Ab)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{2b^2B}{5d \cot^{\frac{5}{2}}(c+dx)} \\
& \downarrow 1103 \\
& 2 \left(-\frac{1}{2}(a^2(A-B) - 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a^2(A+B) + 2ab(A-B) - b^2(A+B)) \right. \\
& \hline
& \frac{2(a^2B + 2aAb - b^2B)}{d\sqrt{\cot(c+dx)}} + \frac{2b(2aB + Ab)}{3d \cot^{\frac{3}{2}}(c+dx)} + \frac{2b^2B}{5d \cot^{\frac{5}{2}}(c+dx)}
\end{aligned}$$

input `Int[((a + b*Tan[c + d*x])^2*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

```
output (2*b^2*B)/(5*d*Cot[c + d*x]^(5/2)) + (2*b*(A*b + 2*a*B))/(3*d*Cot[c + d*x]
^(3/2)) + (2*(2*a*A*b + a^2*B - b^2*B))/(d*Sqrt[Cot[c + d*x]]) + (2*(-1/2*
((a^2*(A - B) - b^2*(A - B) - 2*a*b*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Co
t[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2])) -
((2*a*b*(A - B) + a^2*(A + B) - b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[C
ot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]
+ Cot[c + d*x]]/(2*Sqrt[2])))/2))/d
```

3.582.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

```
rule 1103 Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

```
rule 1476 Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

```
rule 4087 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(- (B*c - A*d))*(b*c - a*d)^2*((c + d*Tan[e + f*x])^(n + 1)/(f*d^2*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[B*(b*c - a*d)^2 + A*d*(a^2*c - b^2*c + 2*a*b*d) + d*(B*(a^2*c - b^2*c + 2*a*b*d) + A*(2*a*b*c - a^2*d + b^2*d))*Tan[e + f*x] + b^2*B*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

```
rule 4111 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]
```

3.582.4 Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 271, normalized size of antiderivative = 0.85

method	result
derivativedivides	$\frac{(A a^2 - A b^2 - 2 B a b) \sqrt{2} \left(\ln \left(\frac{1 + \cot(dx+c) + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c) - \sqrt{2} \sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1 + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \sqrt{2} \sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{-1 + \sqrt{2} \sqrt{\cot(dx+c)}}{-1 + \sqrt{2} \sqrt{\cot(dx+c)}} \right) \right)}{4} + \dots$
default	$\frac{(A a^2 - A b^2 - 2 B a b) \sqrt{2} \left(\ln \left(\frac{1 + \cot(dx+c) + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c) - \sqrt{2} \sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1 + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \sqrt{2} \sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{-1 + \sqrt{2} \sqrt{\cot(dx+c)}}{-1 + \sqrt{2} \sqrt{\cot(dx+c)}} \right) \right)}{4} + \dots$

```
input int((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x,method=_RETURNV ERBOSE)
```

```
output -1/d*(1/4*(A*a^2-A*b^2-2*B*a*b)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/4*(-2*A*a*b-B*a^2+B*b^2)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))-2*(2*A*a*b+B*a^2-B*b^2)/cot(d*x+c)^(1/2)-2/5*B*b^2/cot(d*x+c)^(5/2)-2/3*b*(A*b+2*B*a)/cot(d*x+c)^(3/2))
```

$$3.582. \int \frac{(a+b \tan(c+dx))^2(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.582.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4293 vs. $2(279) = 558$.

Time = 2.37 (sec) , antiderivative size = 4293, normalized size of antiderivative = 13.54

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm
m="fricas")
```

```
output -1/30*(15*d*sqrt((2*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a
^3*b - 4*(A^2 - B^2)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(
A^3*B - A*B^3)*a^7*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B
- A*B^3)*a^5*b^3 + 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B
- A*B^3)*a^3*b^5 - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A
*B^3)*a*b^7 + (A^4 - 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2)*log(((A*a^2 - 2*B*a
b - A*b^2)*d^3*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7
*b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3
+ 2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5
- 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4
- 2*A^2*B^2 + B^4)*b^8)/d^4) - ((A^2*B - B^3)*a^6 + 2*(A^3 - 5*A*B^2)*a^5
*b - (23*A^2*B - 7*B^3)*a^4*b^2 - 4*(3*A^3 - 7*A*B^2)*a^3*b^3 + (23*A^2*B
- 7*B^3)*a^2*b^4 + 2*(A^3 - 5*A*B^2)*a*b^5 - (A^2*B - B^3)*b^6)*d)*sqrt((2
*A*B*a^4 - 12*A*B*a^2*b^2 + 2*A*B*b^4 + 4*(A^2 - B^2)*a^3*b - 4*(A^2 - B^2
)*a*b^3 + d^2*sqrt(-((A^4 - 2*A^2*B^2 + B^4)*a^8 - 16*(A^3*B - A*B^3)*a^7
b - 4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^6*b^2 + 112*(A^3*B - A*B^3)*a^5*b^3 +
2*(19*A^4 - 102*A^2*B^2 + 19*B^4)*a^4*b^4 - 112*(A^3*B - A*B^3)*a^3*b^5 -
4*(3*A^4 - 22*A^2*B^2 + 3*B^4)*a^2*b^6 + 16*(A^3*B - A*B^3)*a*b^7 + (A^4
- 2*A^2*B^2 + B^4)*b^8)/d^4))/d^2) - ((A^4 - B^4)*a^8 - 8*(A^3*B + A*B^3)*
a^7*b - 4*(A^4 - B^4)*a^6*b^2 - 8*(A^3*B + A*B^3)*a^5*b^3 - 10*(A^4 - B...
```

3.582.6 Sympy [F]

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\ &= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^2}{\sqrt{\cot(c + dx)}} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))**2*(A+B*tan(d*x+c))/cot(d*x+c)**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**2/sqrt(cot(c + d*x)),
x)`

3.582.7 Maxima [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 282, normalized size of antiderivative = 0.89

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \frac{8 \left(3 B b^2 + \frac{5 (2 B a b + A b^2)}{\tan(dx+c)} + \frac{15 (B a^2 + 2 A a b - B b^2)}{\tan(dx+c)^2} \right) \tan(dx+c)^{\frac{5}{2}} - 30 \sqrt{2} ((A - B) a^2 - 2 (A + B) a b - (A - B))}{1}$$

input `integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="maxima")`

output `1/60*(8*(3*B*b^2 + 5*(2*B*a*b + A*b^2)/tan(d*x + c) + 15*(B*a^2 + 2*A*a*b - B*b^2)/tan(d*x + c)^2)*tan(d*x + c)^(5/2) - 30*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) - 30*sqrt(2)*((A - B)*a^2 - 2*(A + B)*a*b - (A - B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - 15*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + 15*sqrt(2)*((A + B)*a^2 + 2*(A - B)*a*b - (A + B)*b^2)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))/d`

3.582.8 Giac [F]

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^2}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+b*tan(d*x+c))^2*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^2/sqrt(cot(d*x + c)), x)`

3.582.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^2 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^2}{\sqrt{\cot(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2)/cot(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^2)/cot(c + d*x)^(1/2), x)`

3.583 $\int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.583.1 Optimal result	5508
3.583.2 Mathematica [A] (verified)	5509
3.583.3 Rubi [A] (verified)	5510
3.583.4 Maple [B] (verified)	5517
3.583.5 Fricas [B] (verification not implemented)	5518
3.583.6 Sympy [F(-1)]	5519
3.583.7 Maxima [A] (verification not implemented)	5519
3.583.8 Giac [F]	5520
3.583.9 Mupad [F(-1)]	5520

3.583.1 Optimal result

Integrand size = 33, antiderivative size = 421

$$\int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{(3a^2b(A-B) - b^3(A-B) + a^3(A+B) - 3ab^2(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(3a^2b(A-B) - b^3(A-B) + a^3(A+B) - 3ab^2(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{2(3a^2Ab - Ab^3 + a^3B - 3ab^2B) \sqrt{\cot(c+dx)}}{d}$$

$$+ \frac{2a(7a^2A - 18Ab^2 - 21abB) \cot^{\frac{3}{2}}(c+dx)}{21d}$$

$$- \frac{2a^2(11Ab + 7aB) \cot^{\frac{5}{2}}(c+dx)}{35d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx)(b + a \cot(c+dx))^2}{7d}$$

$$- \frac{(a^3(A-B) - 3ab^2(A-B) - 3a^2b(A+B) + b^3(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{(a^3(A-B) - 3ab^2(A-B) - 3a^2b(A+B) + b^3(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

output $\frac{2}{21}a(7Aa^2-18Ab^2-21Bab)\cot(dx+c)^{3/2}/d-2/35a^2(11Ab+7Ba)\cot(dx+c)^{5/2}/d-2/7aA\cot(dx+c)^{3/2}(b+a\cot(dx+c))^2/d-1/2(3a^2b(A-B)-b^3(A-B)+a^3(A+B)-3ab^2(A+B))\arctan(-1+2^{1/2}\cot(dx+c)^{1/2})/d2^{1/2}-1/2(3a^2b(A-B)-b^3(A-B)+a^3(A+B)-3ab^2(A+B))\arctan(1+2^{1/2}\cot(dx+c)^{1/2})/d2^{1/2}-1/4(a^3(A-B)-3ab^2(A-B)-3a^2b(A+B)+b^3(A+B))\ln(1+\cot(dx+c)-2^{1/2}\cot(dx+c)^{1/2})/d2^{1/2}+1/4(a^3(A-B)-3ab^2(A-B)-3a^2b(A+B)+b^3(A+B))\ln(1+\cot(dx+c)+2^{1/2}\cot(dx+c)^{1/2})/d2^{1/2}+2(3Aa^2b-Ab^3+Ba^3-3Bab^2)\cot(dx+c)^{1/2}/d$

3.583.2 Mathematica [A] (verified)

Time = 3.91 (sec) , antiderivative size = 326, normalized size of antiderivative = 0.77

$$\int \cot^{\frac{9}{2}}(c+dx)(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx$$

$$= \frac{2\sqrt{\cot(c+dx)}\left(-\frac{(3a^2b(A-B)+b^3(-A+B)+a^3(A+B)-3ab^2(A+B))\left(\arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)-\arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)\right)}{2\sqrt{2}}\right)}{1}$$

input `Integrate[Cot[c + d*x]^(9/2)*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output $(2\sqrt{\cot[c+dx]}*(-1/2*((3a^2b(A-B)+b^3(-A+B)+a^3(A+B)-3ab^2(A+B))*(\text{ArcTan}[1-\sqrt{2}\sqrt{\tan[c+dx]})-\text{ArcTan}[1+\sqrt{2}\sqrt{\tan[c+dx]})])/ \sqrt{2} - ((a^3(A-B)+3ab^2(-A+B)-3a^2b(A+B)+b^3(A+B))*(\text{Log}[1-\sqrt{2}\sqrt{\tan[c+dx]}+\tan[c+dx]]-\text{Log}[1+\sqrt{2}\sqrt{\tan[c+dx]}+\tan[c+dx]]))/(4\sqrt{2}) - (a^3A)/(7\tan[c+dx]^{7/2}) - (a^2(3Aab+aB))/(5\tan[c+dx]^{5/2}) + (a(a^2A-3Ab^2-3abB))/(3\tan[c+dx]^{3/2}) + (3a^2Ab-Ab^3+a^3B-3ab^2B)/\sqrt{\tan[c+dx]})\sqrt{\tan[c+dx]})/d$

3.583.3 Rubi [A] (verified)

Time = 1.59 (sec) , antiderivative size = 355, normalized size of antiderivative = 0.84, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.697$, Rules used = {3042, 4064, 3042, 4090, 27, 3042, 4120, 27, 3042, 4113, 3042, 4011, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \cot(c+dx)^{9/2}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4064} \\
 & \int \sqrt{\cot(c+dx)}(a \cot(c+dx)+b)^3(A \cot(c+dx)+B) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(b-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)^3\left(B-A \tan\left(c+dx+\frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{4090} \\
 & -\frac{2}{7} \int \frac{1}{2} \sqrt{\cot(c+dx)}(b+a \cot(c+dx))(-a(11Ab+7aB) \cot^2(c+dx)+7(Aa^2-2bBa-Ab^2) \cot(c+dx)+b(3aA-7bB)) dx - \\
 & \quad \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx)+b)^2}{7d} \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{7} \int \sqrt{\cot(c+dx)}(b+a \cot(c+dx))(-a(11Ab+7aB) \cot^2(c+dx)+7(Aa^2-2bBa-Ab^2) \cot(c+dx)+b(3aA-7bB)) dx - \\
 & \quad \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx)+b)^2}{7d} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{1}{7} \int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(b-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)\left(-a(11Ab+7aB) \tan\left(c+dx+\frac{\pi}{2}\right)^2-7(Aa^2-2bBa- \right. \\
 & \quad \left. \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx)+b)^2}{7d}\right) dx
 \end{aligned}$$

3.583. $\int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\downarrow 4120$$

$$\frac{1}{7} \left(-\frac{2}{5} \int \frac{5}{2} \sqrt{\cot(c+dx)} ((3aA - 7bB)b^2 + a(7Aa^2 - 21bBa - 18Ab^2) \cot^2(c+dx) + 7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)) \cot(c+dx) - 7(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \right) dx + \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)^2}{7d}$$

$$\downarrow 27$$

$$\frac{1}{7} \left(- \int \sqrt{\cot(c+dx)} ((3aA - 7bB)b^2 + a(7Aa^2 - 21bBa - 18Ab^2) \cot^2(c+dx) + 7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)) \cot(c+dx) - 7(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \right) dx + \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)^2}{7d}$$

$$\downarrow 3042$$

$$\frac{1}{7} \left(- \int \sqrt{-\tan\left(c+dx + \frac{\pi}{2}\right)} \left((3aA - 7bB)b^2 + a(7Aa^2 - 21bBa - 18Ab^2) \tan\left(c+dx + \frac{\pi}{2}\right)^2 - 7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \right) \tan\left(c+dx + \frac{\pi}{2}\right) - 7(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \right) dx + \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)^2}{7d}$$

$$\downarrow 4113$$

$$\frac{1}{7} \left(- \int \sqrt{\cot(c+dx)} (7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c+dx) - 7(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)) dx + \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)^2}{7d}$$

$$\downarrow 3042$$

$$\frac{1}{7} \left(- \int \sqrt{-\tan\left(c+dx + \frac{\pi}{2}\right)} \left(-7(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) - 7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan\left(c+dx + \frac{\pi}{2}\right) \right) \tan\left(c+dx + \frac{\pi}{2}\right) - 7(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \right) dx + \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)^2}{7d}$$

$$\downarrow 4011$$

$$\frac{1}{7} \left(- \int \frac{-7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) - 7(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \cot(c+dx)}{\sqrt{\cot(c+dx)}} dx + \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)^2}{7d}$$

$$\downarrow 3042$$

3.583. $\int \cot^{\frac{9}{2}}(c+dx)(a + b \tan(c+dx))^3(A + B \tan(c+dx)) dx$

$$\frac{1}{7} \left(- \int \frac{7(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx + \frac{\pi}{2}) - 7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx + \frac{2a(7a^2A - 21abB - 18Ab^2) \cot(c + dx)}{7d} \right)$$

$$\frac{2aA \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + b)^2}{7d}$$

↓ 4017

$$\frac{1}{7} \left(- \frac{2 \int \frac{7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 + (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \cot(c + dx))}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} + \frac{2a(7a^2A - 21abB - 18Ab^2) \cot(c + dx)}{3d} \right)$$

$$\frac{2aA \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + b)^2}{7d}$$

↓ 27

$$\frac{1}{7} \left(- \frac{14 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 + (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} + \frac{2a(7a^2A - 21abB - 18Ab^2) \cot(c + dx)}{3d} \right)$$

$$\frac{2aA \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + b)^2}{7d}$$

↓ 1482

$$\frac{1}{7} \left(- \frac{14 \left(\frac{1}{2}(a^3(A + B) + 3a^2b(A - B) - 3ab^2(A + B) - b^3(A - B)) \int \frac{\cot(c + dx) + 1}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \frac{1}{2}(a^3(A - B)) \right)}{d} \right)$$

$$\frac{2aA \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + b)^2}{7d}$$

↓ 1476

$$\frac{1}{7} \left(- \frac{14 \left(\frac{1}{2}(a^3(A + B) + 3a^2b(A - B) - 3ab^2(A + B) - b^3(A - B)) \left(\frac{1}{2} \int \frac{1}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)} \right) \right)}{d} \right)$$

$$\frac{2aA \cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + b)^2}{7d}$$

↓ 1082

$$\frac{1}{7} \left(\frac{14 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \right) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)+1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right)}{\frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)^2}{7d}} \right)$$

↓ 217

$$\frac{1}{7} \left(\frac{14 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \right) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} \right)}{d}$$

↓ 1479

$$\frac{1}{7} \left(\frac{14 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \right) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} \right)}{\frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)^2}{7d}} \right)$$

↓ 25

$$\frac{1}{7} \left(\frac{14 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \right) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} \right)}{\frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)^2}{7d}} \right)$$

↓ 27

$$\frac{1}{7} \left(\frac{14 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{\frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)^2}{7d}} \right)$$

↓ 1103

$$\frac{1}{7} \left(\frac{2a(7a^2A - 21abB - 18Ab^2) \cot^{\frac{3}{2}}(c+dx)}{3d} - \frac{2a^2(7aB + 11Ab) \cot^{\frac{5}{2}}(c+dx)}{5d} - \frac{14 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{\frac{2aA \cot^{\frac{3}{2}}(c+dx)(a \cot(c+dx) + b)^2}{7d}} \right)$$

input `Int[Cot[c + d*x]^(9/2)*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(-2*a*A*Cot[c + d*x]^(3/2)*(b + a*Cot[c + d*x])^2)/(7*d) + ((14*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*Sqrt[Cot[c + d*x]])/d + (2*a*(7*a^2*A - 18*A*b^2 - 21*a*b*B)*Cot[c + d*x]^(3/2))/(3*d) - (2*a^2*(11*A*b + 7*a*B)*Cot[c + d*x]^(5/2))/(5*d) - (14*((3*a^2*b*(A - B) - b^3*(A - B) + a^3*(A + B) - 3*a*b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]))/2 - ((a^3*(A - B) - 3*a*b^2*(A - B) - 3*a^2*b*(A + B) + b^3*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d)/7`

3.583.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4011 $\text{Int}[(a + b \tan(e + f x))^m (c + d \tan(e + f x) + (f x))], x_{\text{Symbol}}] \rightarrow \text{Simp}[d (a + b \tan[e + f x])^m / (f m), x] + \text{Int}[(a + b \tan[e + f x])^{m-1} \text{Simp}[a c - b d + (b c + a d) \tan[e + f x], x], x] /;$ FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && GtQ[m, 0]

rule 4017 $\text{Int}[(c + d \tan(e + f x)) / \text{Sqrt}[b \tan(e + f x) + (f x)], x_{\text{Symbol}}] \rightarrow \text{Simp}[2/f \text{ Subst}[\text{Int}[(b c + d x^2) / (b^2 + x^4), x], x, \text{Sqrt}[b \tan[e + f x]]], x] /;$ FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]

rule 4064 $\text{Int}[(\cot(e + f x) + (f x))^p (a + b \tan(e + f x) + (f x))^m (c + d \tan(e + f x) + (f x))^n], x_{\text{Symbol}}] \rightarrow \text{Simp}[g^{m+n} \text{Int}[(g \cot[e + f x])^{p-m-n} (b + a \cot[e + f x])^m (d + c \cot[e + f x])^n], x] /;$ FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]

rule 4090 $\text{Int}[(a + b \tan(e + f x))^m (A + B \tan(e + f x) + (f x)) (c + d \tan(e + f x) + (f x))^n], x_{\text{Symbol}}] \rightarrow \text{Simp}[b B (a + b \tan[e + f x])^{m-1} (c + d \tan[e + f x])^{n+1} / (d f (m + n)), x] + \text{Simp}[1 / (d (m + n)) \text{Int}[(a + b \tan[e + f x])^{m-2} (c + d \tan[e + f x])^n \text{Simp}[a^2 A d (m + n) - b B (b c (m - 1) + a d (n + 1)) + d (m + n) (2 a A b + B (a^2 - b^2)) \tan[e + f x] - (b B (b c - a d) (m - 1) - b (A b + a B) d (m + n)) \tan[e + f x]^2, x], x], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))

rule 4113 $\text{Int}[(a + b \tan(e + f x))^m (A + B \tan(e + f x) + (f x)) + (C \tan(e + f x) + (f x))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[C (a + b \tan[e + f x])^{m+1} / (b f (m + 1)), x] + \text{Int}[(a + b \tan[e + f x])^m \text{Simp}[A - C + B \tan[e + f x], x], x] /;$ FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]

```
rule 4120 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_) + (C_)*tan[(e_) + (f
_)*(x_)])^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

3.583.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1191 vs. $2(379) = 758$.

Time = 1.18 (sec) , antiderivative size = 1192, normalized size of antiderivative = 2.83

method	result	size
derivativedivides	Expression too large to display	1192
default	Expression too large to display	1192

```
input int(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

output $1/420/d*(1/\tan(dx+c))^{9/2}*\tan(dx+c)*(-840*A*\tan(dx+c)^3*b^3-168*B*\tan(dx+c)*a^3-630*B*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*\tan(dx+c)^{7/2})*a*b^2-630*B*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*\tan(dx+c)^{7/2})*a^2*b-630*B*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*\tan(dx+c)^{7/2})*a*b^2-315*B*\ln(-(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1)/(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c)))*2^{1/2}*\tan(dx+c)^{7/2})*a*b^2-315*A*\ln(-1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c))/(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1))*2^{1/2}*\tan(dx+c)^{7/2})*a*b^2+630*A*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*\tan(dx+c)^{7/2})*a^2*b-630*A*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*\tan(dx+c)^{7/2})*a*b^2+630*A*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*\tan(dx+c)^{7/2})*a^2*b-630*A*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*\tan(dx+c)^{7/2})*a*b^2+315*A*\ln(-(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1)/(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c)))*2^{1/2}*\tan(dx+c)^{7/2})*a^2*b-315*B*\ln(-1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c))/(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1))*2^{1/2}*\tan(dx+c)^{7/2})*a^2*b-630*B*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*\tan(dx+c)^{7/2})*a^2*b+2520*A*\tan(dx+c)^3*a^2*b-120*A*a^3-2520*B*\tan(dx+c)^3*a*b^2-840*A*\tan(dx+c)^2*a*b^2-840*B*\tan(dx+c)^2*a^2*b-504*A*\tan(dx+c)*a^2*b+840*B*\tan(dx+c)^3*a^3+280*A*\tan(dx+c)^2*a^3+210*A*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*\tan(dx+c)^{7/2})*a^3-210*A*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*\tan(dx+c)^{7/2})*b^3-...$

3.583.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6214 vs. $2(379) = 758$.

Time = 6.61 (sec) , antiderivative size = 6214, normalized size of antiderivative = 14.76

$$\int \cot^{\frac{9}{2}}(c+dx)(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx = \text{Too large to display}$$

input `integrate(cot(dx+c)^(9/2)*(a+b*tan(dx+c))^3*(A+B*tan(dx+c)),x, algorithm="fricas")`

output Too large to include

3.583. $\int \cot^{\frac{9}{2}}(c+dx)(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx$

3.583.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(9/2)*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.583.7 Maxima [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 366, normalized size of antiderivative = 0.87

$$\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx =$$

$$\frac{210 \sqrt{2}((A + B)a^3 + 3(A - B)a^2b - 3(A + B)ab^2 - (A - B)b^3) \arctan\left(\frac{1}{2} \sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + \dots}{\dots}$$

input `integrate(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/420*(210*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 210*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - 105*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + 105*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + 120*A*a^3/tan(d*x + c)^(7/2) - 840*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)/sqrt(tan(d*x + c)) - 280*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2)/tan(d*x + c)^(3/2) + 168*(B*a^3 + 3*A*a^2*b)/tan(d*x + c)^(5/2))/d`

3.583.8 Giac [F]

$$\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^3 \cot(dx + c)^{\frac{9}{2}} dx$$

input `integrate(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^3*cot(d*x + c)^(9/2), x)`

3.583.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{9/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^3 dx$$

input `int(cot(c + d*x)^(9/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`

output `int(cot(c + d*x)^(9/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3, x)`

3.584 $\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.584.1 Optimal result	5521
3.584.2 Mathematica [A] (verified)	5522
3.584.3 Rubi [A] (verified)	5523
3.584.4 Maple [B] (verified)	5529
3.584.5 Fricas [B] (verification not implemented)	5530
3.584.6 Sympy [F(-1)]	5531
3.584.7 Maxima [A] (verification not implemented)	5531
3.584.8 Giac [F]	5532
3.584.9 Mupad [F(-1)]	5532

3.584.1 Optimal result

Integrand size = 33, antiderivative size = 380

$$\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{(a^3(A-B) - 3ab^2(A-B) - 3a^2b(A+B) + b^3(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(a^3(A-B) - 3ab^2(A-B) - 3a^2b(A+B) + b^3(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{2a(5a^2A - 14Ab^2 - 15abB) \sqrt{\cot(c+dx)}}{5d}$$

$$- \frac{2a^2(9Ab + 5aB) \cot^{\frac{3}{2}}(c+dx)}{15d} - \frac{2aA \sqrt{\cot(c+dx)}(b + a \cot(c+dx))^2}{5d}$$

$$+ \frac{(3a^2b(A-B) - b^3(A-B) + a^3(A+B) - 3ab^2(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(3a^2b(A-B) - b^3(A-B) + a^3(A+B) - 3ab^2(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

output
$$\begin{aligned} & -2/15*a^2*(9*A*b+5*B*a)*cot(d*x+c)^(3/2)/d-1/2*(a^3*(A-B)-3*a*b^2*(A-B)-3* \\ & a^2*b*(A+B)+b^3*(A+B))*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/2*(\\ & a^3*(A-B)-3*a*b^2*(A-B)-3*a^2*b*(A+B)+b^3*(A+B))*arctan(1+2^(1/2)*cot(d*x+ \\ & c)^(1/2))/d*2^(1/2)+1/4*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))* \\ & ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/4*(3*a^2*b*(A-B)-b^3* \\ & *(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/ \\ & d*2^(1/2)+2/5*a*(5*A*a^2-14*A*b^2-15*B*a*b)*cot(d*x+c)^(1/2)/d-2/5*a*A*(b+ \\ & a*cot(d*x+c))^2*cot(d*x+c)^(1/2)/d \end{aligned}$$

3.584.2 Mathematica [A] (verified)

Time = 2.47 (sec) , antiderivative size = 286, normalized size of antiderivative = 0.75

$$\int \cot^{7/2}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{2\sqrt{\cot(c+dx)} \left(-\frac{(a^3(A-B)+3ab^2(-A+B)-3a^2b(A+B)+b^3(A+B))(\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})-\arctan(1+\sqrt{2}\sqrt{\tan(c+dx)})}{2\sqrt{2}} \right)}{1}$$

input `Integrate[Cot[c + d*x]^(7/2)*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output
$$\begin{aligned} & (2*\text{Sqrt}[\text{Cot}[c + d*x]]*(-1/2*((a^3*(A - B) + 3*a*b^2*(-A + B) - 3*a^2*b*(A \\ & + B) + b^3*(A + B))*(\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]] - \text{ArcTan}[1 + \text{S} \\ & \text{qrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]]))/\text{Sqrt}[2] + ((3*a^2*b*(A - B) + b^3*(-A + B) + \\ & a^3*(A + B) - 3*a*b^2*(A + B))*(\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c \\ & + d*x]] - \text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]]))/(4*\text{Sqrt}[2] \\ &) - (a^3*A)/(5*\text{Tan}[c + d*x]^(5/2)) - (a^2*(3*A*b + a*B))/(3*\text{Tan}[c + d*x]^(\\ & 3/2)) + (a*(a^2*A - 3*A*b^2 - 3*a*b*B))/\text{Sqrt}[\text{Tan}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d \\ & *x]])/d \end{aligned}$$

3.584.3 Rubi [A] (verified)

Time = 1.32 (sec) , antiderivative size = 312, normalized size of antiderivative = 0.82, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 4064, 3042, 4090, 27, 3042, 4120, 27, 3042, 4113, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \cot(c+dx)^{7/2}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{(a \cot(c+dx)+b)^3(A \cot(c+dx)+B)}{\sqrt{\cot(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b-a \tan(c+dx+\frac{\pi}{2}))^3(B-A \tan(c+dx+\frac{\pi}{2}))}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx \\
 & \quad \downarrow \text{4090} \\
 & -\frac{2}{5} \int \frac{(b+a \cot(c+dx))(-a(9Ab+5aB) \cot^2(c+dx)+5(Aa^2-2bBa-Ab^2) \cot(c+dx)+b(aA-5bB))}{2\sqrt{\cot(c+dx)}} dx \\
 & \quad \frac{2aA\sqrt{\cot(c+dx)}(a \cot(c+dx)+b)^2}{5d} \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{5} \int \frac{(b+a \cot(c+dx))(-a(9Ab+5aB) \cot^2(c+dx)+5(Aa^2-2bBa-Ab^2) \cot(c+dx)+b(aA-5bB))}{\sqrt{\cot(c+dx)}} dx \\
 & \quad \frac{2aA\sqrt{\cot(c+dx)}(a \cot(c+dx)+b)^2}{5d} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$3.584. \quad \int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$-\frac{1}{5} \int \frac{(b - a \tan(c + dx + \frac{\pi}{2})) \left(-a(9Ab + 5aB) \tan(c + dx + \frac{\pi}{2})^2 - 5(Aa^2 - 2bBa - Ab^2) \tan(c + dx + \frac{\pi}{2}) + b \right)}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx + \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2}{5d}$$

↓ 4120

$$\frac{1}{5} \left(-\frac{2}{3} \int \frac{3((aA - 5bB)b^2 + a(5Aa^2 - 15bBa - 14Ab^2) \cot^2(c + dx) + 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c + dx))}{2\sqrt{\cot(c + dx)}} dx + \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2}{5d} \right)$$

↓ 27

$$\frac{1}{5} \left(- \int \frac{(aA - 5bB)b^2 + a(5Aa^2 - 15bBa - 14Ab^2) \cot^2(c + dx) + 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx + \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(- \int \frac{(aA - 5bB)b^2 + a(5Aa^2 - 15bBa - 14Ab^2) \tan(c + dx + \frac{\pi}{2})^2 - 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx + \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2}{5d} \right)$$

↓ 4113

$$\frac{1}{5} \left(- \int \frac{5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c + dx) - 5(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)}{\sqrt{\cot(c + dx)}} dx + \frac{2a(5a^2A - 15abB - 5b^2B)}{5d} + \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2}{5d} \right)$$

↓ 3042

$$\frac{1}{5} \left(- \int \frac{-5(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) - 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx + \frac{2a(5a^2A - 15abB - 5b^2B)}{5d} + \frac{2aA\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2}{5d} \right)$$

↓ 4017

3.584. $\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\frac{1}{5} \left(-\frac{2 \int \frac{5(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B - (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} + \frac{2a(5a^2A - 15abB - 14Ab^2) \sqrt{\cot(c+dx)}}{d} \right) - \frac{2aA\sqrt{\cot(c+dx)}(a \cot(c+dx) + b)^2}{5d}$$

↓ 27

$$\frac{1}{5} \left(-\frac{10 \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B - (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} + \frac{2a(5a^2A - 15abB - 14Ab^2) \sqrt{\cot(c+dx)}}{d} \right) - \frac{2aA\sqrt{\cot(c+dx)}(a \cot(c+dx) + b)^2}{5d}$$

↓ 1482

$$\frac{1}{5} \left(-\frac{10 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(a^3(A-B) \right)}{d} \right) - \frac{2aA\sqrt{\cot(c+dx)}(a \cot(c+dx) + b)^2}{5d}$$

↓ 1476

$$\frac{1}{5} \left(-\frac{10 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(a^3(A-B) \right)}{d} \right) - \frac{2aA\sqrt{\cot(c+dx)}(a \cot(c+dx) + b)^2}{5d}$$

↓ 1082

$$\frac{1}{5} \left(-\frac{10 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(a^3(A-B) \right)}{d} \right) - \frac{2aA\sqrt{\cot(c+dx)}(a \cot(c+dx) + b)^2}{5d}$$

↓ 217

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(a^3(A-B) \right)}{d} \right. \\ \left. \frac{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx)+b)^2}{5d} \right. \\ \left. \downarrow 1479 \right)$$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \int \frac{\sqrt{2}+2\sqrt{\cot(c+dx)}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) \right)}{d} \right. \\ \left. \frac{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx)+b)^2}{5d} \right. \\ \left. \downarrow 25 \right)$$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \int \frac{\sqrt{2}+2\sqrt{\cot(c+dx)}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) \right)}{d} \right. \\ \left. \frac{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx)+b)^2}{5d} \right. \\ \left. \downarrow 27 \right)$$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}+2\sqrt{\cot(c+dx)}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) \right)}{d} \right. \\ \left. \frac{2aA\sqrt{\cot(c+dx)}(a\cot(c+dx)+b)^2}{5d} \right. \\ \left. \downarrow 1103 \right)$$

$$\frac{1}{5} \left(\frac{2a(5a^2A - 15abB - 14Ab^2) \sqrt{\cot(c + dx)}}{d} - \frac{2a^2(5aB + 9Ab) \cot^{\frac{3}{2}}(c + dx)}{3d} - \frac{10 \left(\frac{1}{2}(a^3(A - B) - 3a^2b(A + B) + b^3(A + B)) \right)}{5d} \right)$$

input `Int[Cot[c + d*x]^(7/2)*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(-2*a*A*Sqrt[Cot[c + d*x]]*(b + a*Cot[c + d*x])^2)/(5*d) + ((2*a*(5*a^2*A - 14*A*b^2 - 15*a*b*B)*Sqrt[Cot[c + d*x]])/d - (2*a^2*(9*A*b + 5*a*B)*Cot[c + d*x]^(3/2))/(3*d) - (10*((a^3*(A - B) - 3*a*b^2*(A - B) - 3*a^2*b*(A + B) + b^3*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]))/2 + ((3*a^2*b*(A - B) - b^3*(A - B) + a^3*(A + B) - 3*a*b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d)/5`

3.584.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)])*(g_.)^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

```
rule 4090 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4113 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Si
mp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] &&
NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

```
rule 4120 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n +
1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Si
mp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*
d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] &&
!LtQ[n, -1]
```

3.584.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1133 vs. $2(340) = 680$.

Time = 1.22 (sec) , antiderivative size = 1134, normalized size of antiderivative = 2.98

method	result	size
derivativedivides	Expression too large to display	1134
default	Expression too large to display	1134

```
input int(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

$$3.584. \quad \int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

output
$$-1/60/d*(1/\tan(d*x+c))^{(7/2)}*\tan(d*x+c)*(40*B*\tan(d*x+c)*a^3-30*A*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*b^3-45*B*\ln(-(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(2^{(1/2)}*\tan(d*x+c)^{(1/2)}-\tan(d*x+c)-1))*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*a*b^2+90*A*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*a^2*b+90*A*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*a*b^2+90*A*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*a^2*b+90*A*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*a*b^2+45*A*\ln(-(2^{(1/2)}*\tan(d*x+c)^{(1/2)}-\tan(d*x+c)-1)/(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*a*b^2+90*B*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*a^2*b-90*B*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*a*b^2+45*B*\ln(-(2^{(1/2)}*\tan(d*x+c)^{(1/2)}-\tan(d*x+c)-1)/(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c)))*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*a^2*b+90*B*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*a^2*b-90*B*\arctan(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*a*b^2+45*A*\ln(-(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(2^{(1/2)}*\tan(d*x+c)^{(1/2)}-\tan(d*x+c)-1))*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*a^2*b+24*A*a^3+360*A*\tan(d*x+c)^2*a*b^2+360*B*\tan(d*x+c)^2*a^2*b+120*A*\tan(d*x+c)*a^2*b-30*A*\arctan(-1+2^{(1/2)}*\tan(d*x+c)^{(1/2)})*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*a^3-120*A*\tan(d*x+c)^2*a^3-15*A*\ln(-(1+2^{(1/2)}*\tan(d*x+c)^{(1/2)}+\tan(d*x+c))/(2^{(1/2)}*\tan(d*x+c)^{(1/2)}-\tan(d*x+c)-1))*2^{(1/2)}*\tan(d*x+c)^{(5/2)}*...$$

3.584.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6162 vs. $2(340) = 680$.

Time = 6.46 (sec) , antiderivative size = 6162, normalized size of antiderivative = 16.22

$$\int \cot^{\frac{7}{2}}(c+dx)(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output Too large to include

3.584.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(7/2)*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.584.7 Maxima [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 330, normalized size of antiderivative = 0.87

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx =$$

$$\frac{30\sqrt{2}((A - B)a^3 - 3(A + B)a^2b - 3(A - B)ab^2 + (A + B)b^3) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 30\sqrt{2}((A - B)a^3 - 3(A + B)a^2b - 3(A - B)ab^2 + (A + B)b^3) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 15\sqrt{2}((A + B)a^3 + 3(A - B)a^2b - 3(A + B)ab^2 - (A - B)b^3) \log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}}\right) + 15\sqrt{2}((A + B)a^3 + 3(A - B)a^2b - 3(A + B)ab^2 - (A - B)b^3) \log\left(-\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}}\right) + \frac{24Aa^3}{\tan(dx+c)^{5/2}} - \frac{120(Aa^3 - 3Ba^2b - 3Aab^2 - 2Bb^3)}{\sqrt{\tan(dx+c)}} + \frac{40(Ba^3 + 3Aa^2b)}{\tan(dx+c)^{3/2}}}{d}$$

input `integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/60*(30*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 30*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + 15*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - 15*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + 24*A*a^3/tan(d*x + c)^(5/2) - 120*(A*a^3 - 3*B*a^2*b - 3*A*a*b^2 - 2*B*b^3)/sqrt(tan(d*x + c)) + 40*(B*a^3 + 3*A*a^2*b)/tan(d*x + c)^(3/2))/d`

3.584.8 Giac [F]

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^3 \cot(dx + c)^{\frac{7}{2}} dx$$

input `integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^3*cot(d*x + c)^(7/2), x)`

3.584.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{7/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^3 dx$$

input `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`

output `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3, x)`

3.585 $\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.585.1 Optimal result	5533
3.585.2 Mathematica [A] (verified)	5534
3.585.3 Rubi [A] (verified)	5534
3.585.4 Maple [B] (verified)	5541
3.585.5 Fricas [B] (verification not implemented)	5542
3.585.6 Sympy [F(-1)]	5542
3.585.7 Maxima [A] (verification not implemented)	5542
3.585.8 Giac [F]	5543
3.585.9 Mupad [F(-1)]	5543

3.585.1 Optimal result

Integrand size = 33, antiderivative size = 374

$$\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx =$$

$$\frac{(3a^2b(A-B) - b^3(A-B) + a^3(A+B) - 3ab^2(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(3a^2b(A-B) - b^3(A-B) + a^3(A+B) - 3ab^2(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$- \frac{2a(3aAb + a^2B + 2b^2B) \sqrt{\cot(c+dx)}}{d}$$

$$- \frac{2a^2(aA + 3bB) \cot^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB(b + a \cot(c+dx))^2}{d\sqrt{\cot(c+dx)}}$$

$$+ \frac{(a^3(A-B) - 3ab^2(A-B) - 3a^2b(A+B) + b^3(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(a^3(A-B) - 3ab^2(A-B) - 3a^2b(A+B) + b^3(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

3.585. $\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

output
$$\begin{aligned} & -2/3*a^2*(A*a+3*B*b)*\cot(d*x+c)^{(3/2)}/d+1/2*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*\arctan(-1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/d*2^{(1/2)}+1/2*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*\arctan(1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/d*2^{(1/2)}+1/4*(a^3*(A-B)-3*a*b^2*(A+B)-3*a^2*b*(A+B)+b^3*(A+B))*\ln(1+\cot(d*x+c)-2^{(1/2)}*\cot(d*x+c)^{(1/2)})/d*2^{(1/2)}-1/4*(a^3*(A-B)-3*a*b^2*(A+B)-3*a^2*b*(A+B)+b^3*(A+B))*\ln(1+\cot(d*x+c)+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/d*2^{(1/2)}+2*b*B*(b+a*\cot(d*x+c))^2/d/\cot(d*x+c)^{(1/2)}-2*a*(3*A*a*b+B*a^2+2*B*b^2)*\cot(d*x+c)^{(1/2)}/d \end{aligned}$$

3.585.2 Mathematica [A] (verified)

Time = 2.19 (sec) , antiderivative size = 270, normalized size of antiderivative = 0.72

$$\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$= \frac{2\sqrt{\cot(c+dx)} \left(\frac{(3a^2b(A-B)+b^3(-A+B)+a^3(A+B)-3ab^2(A+B)) \left(\arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right) - \arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right) \right)}{2\sqrt{2}} + (a \right)}{\dots}$$

input `Integrate[Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output
$$\begin{aligned} & (2*\text{Sqrt}[\text{Cot}[c + d*x]]*((3*a^2*b*(A - B) + b^3*(-A + B) + a^3*(A + B) - 3*a*b^2*(A + B))*(\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]] - \text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]])/(2*\text{Sqrt}[2]) + ((a^3*(A - B) + 3*a*b^2*(-A + B) - 3*a^2*b*(A + B) + b^3*(A + B))*(\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]] - \text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]]))/(4*\text{Sqrt}[2]) - (a^3*A)/(3*\text{Tan}[c + d*x]^(3/2)) - (a^2*(3*A*b + a*B))/\text{Sqrt}[\text{Tan}[c + d*x]] + b^3*B*\text{Sqrt}[\text{Tan}[c + d*x]])*\text{Sqrt}[\text{Tan}[c + d*x]])/d \end{aligned}$$

3.585.3 Rubi [A] (verified)

Time = 1.29 (sec) , antiderivative size = 303, normalized size of antiderivative = 0.81, number of steps used = 21, number of rules used = 20, $\frac{\text{number of rules}}{\text{integrand size}} = 0.606$, Rules used = {3042, 4064, 3042, 4088, 27, 3042, 4120, 27, 3042, 4113, 3042, 4017, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$3.585. \quad \int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$$

$$\begin{aligned}
& \int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
& \quad \downarrow \text{3042} \\
& \int \cot(c+dx)^{5/2}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
& \quad \downarrow \text{4064} \\
& \int \frac{(a \cot(c+dx)+b)^3(A \cot(c+dx)+B)}{\cot^{\frac{3}{2}}(c+dx)} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(b-a \tan(c+dx+\frac{\pi}{2}))^3(B-A \tan(c+dx+\frac{\pi}{2}))}{(-\tan(c+dx+\frac{\pi}{2}))^{3/2}} dx \\
& \quad \downarrow \text{4088} \\
& \frac{2bB(a \cot(c+dx)+b)^2}{d\sqrt{\cot(c+dx)}} - \\
& 2 \int \frac{(b+a \cot(c+dx))(a(aA+3bB) \cot^2(c+dx)+(Ba^2+2Aba-b^2B) \cot(c+dx)+b(Ab+5aB))}{2\sqrt{\cot(c+dx)}} dx \\
& \quad \downarrow \text{27} \\
& \int \frac{(b+a \cot(c+dx))(a(aA+3bB) \cot^2(c+dx)+(Ba^2+2Aba-b^2B) \cot(c+dx)+b(Ab+5aB))}{\sqrt{\cot(c+dx)}} dx + \\
& \quad \frac{2bB(a \cot(c+dx)+b)^2}{d\sqrt{\cot(c+dx)}} \\
& \quad \downarrow \text{3042} \\
& \int \frac{(b-a \tan(c+dx+\frac{\pi}{2})) \left(a(aA+3bB) \tan(c+dx+\frac{\pi}{2})^2 - (Ba^2+2Aba-b^2B) \tan(c+dx+\frac{\pi}{2}) + b(Ab+5aB) \right)}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx + \\
& \quad \frac{2bB(a \cot(c+dx)+b)^2}{d\sqrt{\cot(c+dx)}} \\
& \quad \downarrow \text{4120} \\
& \frac{2}{3} \int \frac{3((Ab+5aB)b^2+a(Ba^2+3Aba+2b^2B) \cot^2(c+dx)-(Aa^3-3bBa^2-3Ab^2a+b^3B) \cot(c+dx))}{2\sqrt{\cot(c+dx)}} dx - \\
& \quad \frac{2a^2(aA+3bB) \cot^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB(a \cot(c+dx)+b)^2}{d\sqrt{\cot(c+dx)}}
\end{aligned}$$

3.585. $\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

↓ 27

$$\int \frac{(Ab + 5aB)b^2 + a(Ba^2 + 3Aba + 2b^2B) \cot^2(c + dx) - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx -$$

$$\frac{2a^2(aA + 3bB) \cot^{\frac{3}{2}}(c + dx)}{3d} + \frac{2bB(a \cot(c + dx) + b)^2}{d\sqrt{\cot(c + dx)}}$$

↓ 3042

$$\int \frac{(Ab + 5aB)b^2 + a(Ba^2 + 3Aba + 2b^2B) \tan(c + dx + \frac{\pi}{2})^2 - (-Aa^3 + 3bBa^2 + 3Ab^2a - b^3B) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx -$$

$$\frac{2a^2(aA + 3bB) \cot^{\frac{3}{2}}(c + dx)}{3d} + \frac{2bB(a \cot(c + dx) + b)^2}{d\sqrt{\cot(c + dx)}}$$

↓ 4113

$$\int \frac{-Ba^3 - 3Aba^2 + 3b^2Ba + Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx -$$

$$\frac{2a(a^2B + 3aAb + 2b^2B) \sqrt{\cot(c + dx)}}{d} - \frac{2a^2(aA + 3bB) \cot^{\frac{3}{2}}(c + dx)}{3d} + \frac{2bB(a \cot(c + dx) + b)^2}{d\sqrt{\cot(c + dx)}}$$

↓ 3042

$$\int \frac{-Ba^3 - 3Aba^2 + 3b^2Ba + Ab^3 - (-Aa^3 + 3bBa^2 + 3Ab^2a - b^3B) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx -$$

$$\frac{2a(a^2B + 3aAb + 2b^2B) \sqrt{\cot(c + dx)}}{d} - \frac{2a^2(aA + 3bB) \cot^{\frac{3}{2}}(c + dx)}{3d} + \frac{2bB(a \cot(c + dx) + b)^2}{d\sqrt{\cot(c + dx)}}$$

↓ 4017

$$2 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 + (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} -$$

$$\frac{d}{2a(a^2B + 3aAb + 2b^2B) \sqrt{\cot(c + dx)}} - \frac{2a^2(aA + 3bB) \cot^{\frac{3}{2}}(c + dx)}{3d} + \frac{2bB(a \cot(c + dx) + b)^2}{d\sqrt{\cot(c + dx)}}$$

↓ 1482

$$2 \left(\frac{1}{2} (a^3(A + B) + 3a^2b(A - B) - 3ab^2(A + B) - b^3(A - B)) \int \frac{\cot(c + dx) + 1}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \frac{1}{2} (a^3(A - B) - 3a^2b(A + B) + 3ab^2(A - B) + b^3(A + B)) \int \frac{1}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} \right) -$$

$$\frac{d}{2a(a^2B + 3aAb + 2b^2B) \sqrt{\cot(c + dx)}} - \frac{2a^2(aA + 3bB) \cot^{\frac{3}{2}}(c + dx)}{3d} + \frac{2bB(a \cot(c + dx) + b)^2}{d\sqrt{\cot(c + dx)}}$$

↓ 1476

3.585. $\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\frac{2\left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B))\left(\frac{1}{2}\int \frac{1}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)} + 1} d\sqrt{\cot(c+dx)} + \frac{1}{2}\int \frac{1}{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)} + 1} d\sqrt{\cot(c+dx)}\right) + \frac{2a(a^2B + 3aAb + 2b^2B)\sqrt{\cot(c+dx)}}{d} - \frac{2a^2(aA + 3bB)\cot^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB(a\cot(c+dx) + b)^2}{d\sqrt{\cot(c+dx)}}}{d}$$

↓ 1082

$$\frac{2\left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B))\left(\frac{\int \frac{1}{-\cot(c+dx) - 1} d(1 - \sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx) - 1} d(1 + \sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}}\right) + \frac{2a(a^2B + 3aAb + 2b^2B)\sqrt{\cot(c+dx)}}{d} - \frac{2a^2(aA + 3bB)\cot^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB(a\cot(c+dx) + b)^2}{d\sqrt{\cot(c+dx)}}}{d}$$

↓ 217

$$\frac{2\left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B))\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}}\right) + \frac{2a(a^2B + 3aAb + 2b^2B)\sqrt{\cot(c+dx)}}{d} - \frac{2a^2(aA + 3bB)\cot^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB(a\cot(c+dx) + b)^2}{d\sqrt{\cot(c+dx)}}}{d}$$

↓ 1479

$$\frac{2\left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B))\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}}\right) + \frac{2a(a^2B + 3aAb + 2b^2B)\sqrt{\cot(c+dx)}}{d} - \frac{2a^2(aA + 3bB)\cot^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB(a\cot(c+dx) + b)^2}{d\sqrt{\cot(c+dx)}}}{d}$$

↓ 25

$$\frac{2\left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B))\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}}\right) + \frac{2a(a^2B + 3aAb + 2b^2B)\sqrt{\cot(c+dx)}}{d} - \frac{2a^2(aA + 3bB)\cot^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB(a\cot(c+dx) + b)^2}{d\sqrt{\cot(c+dx)}}}{d}$$

↓ 27

3.585. $\int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^3(A+B\tan(c+dx)) dx$

$$\begin{aligned}
& 2 \left(\frac{1}{2} (a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right) \\
& \frac{2a(a^2B + 3aAb + 2b^2B) \sqrt{\cot(c+dx)}}{d} - \frac{2a^2(aA + 3bB) \cot^{\frac{3}{2}}(c+dx)}{3d} + \frac{2bB(a \cot(c+dx) + b)^2}{d\sqrt{\cot(c+dx)}} \\
& \quad \downarrow \text{1103} \\
& - \frac{2a(a^2B + 3aAb + 2b^2B) \sqrt{\cot(c+dx)}}{d} - \frac{2a^2(aA + 3bB) \cot^{\frac{3}{2}}(c+dx)}{3d} + \\
& 2 \left(\frac{1}{2} (a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right) \\
& \frac{2bB(a \cot(c+dx) + b)^2}{d\sqrt{\cot(c+dx)}}
\end{aligned}$$

input `Int[Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(-2*a*(3*a*A*b + a^2*B + 2*b^2*B)*Sqrt[Cot[c + d*x]]/d - (2*a^2*(a*A + 3*b*B)*Cot[c + d*x]^(3/2))/(3*d) + (2*b*B*(b + a*Cot[c + d*x])^2)/(d*Sqrt[Cot[c + d*x]]) + (2*(((3*a^2*b*(A - B) - b^3*(A - B) + a^3*(A + B) - 3*a*b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]))/2 - ((a^3*(A - B) - 3*a*b^2*(A - B) - 3*a^2*b*(A + B) + b^3*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d`

3.585.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

3.585. $\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp [g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4113 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[C*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]`

rule 4120 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[b*C*Tan[e + f*x]*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 2))), x] - Simp[1/(d*(n + 2)) Int[(c + d*Tan[e + f*x])^n*Simp[b*c*C - a*A*d*(n + 2) - (A*b + a*B - b*C)*d*(n + 2)*Tan[e + f*x] - (a*C*d*(n + 2) - b*(c*C - B*d*(n + 2)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && !LtQ[n, -1]`

3.585.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1103 vs. $2(338) = 676$.

Time = 1.22 (sec) , antiderivative size = 1104, normalized size of antiderivative = 2.95

method	result	size
derivativdivides	Expression too large to display	1104
default	Expression too large to display	1104

```
input int(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

```
output -1/12/d*(1/tan(d*x+c))^(5/2)*tan(d*x+c)*(24*B*tan(d*x+c)*a^3+18*A*arctan(1
+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(3/2)*a^2*b-18*A*arctan(1+2^(
1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(3/2)*a*b^2+18*A*arctan(-1+2^(1
/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(3/2)*a^2*b-9*A*ln(-(1+2^(1/2)*ta
n(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1))*2^(1/2
)*tan(d*x+c)^(3/2)*a*b^2-18*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*t
an(d*x+c)^(3/2)*a^2*b-18*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(
d*x+c)^(3/2)*a*b^2-18*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*
x+c)^(3/2)*a^2*b-18*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+
c)^(3/2)*a*b^2-9*B*ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*
tan(d*x+c)^(1/2)+tan(d*x+c)))*2^(1/2)*tan(d*x+c)^(3/2)*a*b^2-18*A*arctan(-
1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(3/2)*a*b^2+9*A*ln(-(2^(1/2
)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))*
2^(1/2)*tan(d*x+c)^(3/2)*a^2*b-9*B*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x
+c))/(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1))*2^(1/2)*tan(d*x+c)^(3/2)*a^2
*b+8*A*a^3+72*A*tan(d*x+c)*a^2*b-24*B*b^3*tan(d*x+c)^2+6*A*arctan(-1+2^(1/
2)*tan(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(3/2)*a^3-6*A*arctan(-1+2^(1/2)*ta
n(d*x+c)^(1/2))*2^(1/2)*tan(d*x+c)^(3/2)*b^3-3*A*ln(-(2^(1/2)*tan(d*x+c)^(
1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))*2^(1/2)*tan(d*
x+c)^(3/2)*b^3+3*B*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)...
```

3.585.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6154 vs. $2(338) = 676$.

Time = 6.38 (sec) , antiderivative size = 6154, normalized size of antiderivative = 16.45

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output Too large to include

3.585.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(5/2)*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output Timed out

3.585.7 Maxima [A] (verification not implemented)

Time = 0.65 (sec) , antiderivative size = 314, normalized size of antiderivative = 0.84

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{24 B b^3 \sqrt{\tan(dx + c)} + 6 \sqrt{2}((A + B)a^3 + 3(A - B)a^2 b - 3(A + B)ab^2 - (A - B)b^3) \arctan\left(\frac{1}{2} \sqrt{2}(\sqrt{2} \tan(dx + c) - 1)\right)}{1}$$

input `integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

3.585. $\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

output $1/12*(24*B*b^3*\sqrt{\tan(dx + c)} + 6*\sqrt{2}*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2/\sqrt{\tan(dx + c)})) + 6*\sqrt{2}*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2/\sqrt{\tan(dx + c)})) - 3*\sqrt{2}*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*\log(\sqrt{2}/\sqrt{\tan(dx + c)} + 1/\tan(dx + c) + 1) + 3*\sqrt{2}*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*\log(-\sqrt{2}/\sqrt{\tan(dx + c)} + 1/\tan(dx + c) + 1) - 8*A*a^3/\tan(dx + c)^{(3/2)} - 24*(B*a^3 + 3*A*a^2*b)/\sqrt{\tan(dx + c)})/d$

3.585.8 Giac [F]

$$\int \cot^{5/2}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^3 \cot(dx + c)^{5/2} dx$$

input `integrate(cot(dx+c)^(5/2)*(a+b*tan(dx+c))^3*(A+B*tan(dx+c)),x, algorithm="giac")`

output `integrate((B*tan(dx + c) + A)*(b*tan(dx + c) + a)^3*cot(dx + c)^(5/2), x)`

3.585.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{5/2}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{5/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^3 dx$$

input `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`

output `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3, x)`

3.585. $\int \cot^{5/2}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.586 $\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

3.586.1 Optimal result	5544
3.586.2 Mathematica [A] (verified)	5545
3.586.3 Rubi [A] (verified)	5545
3.586.4 Maple [B] (verified)	5552
3.586.5 Fricas [B] (verification not implemented)	5553
3.586.6 Sympy [F]	5554
3.586.7 Maxima [A] (verification not implemented)	5554
3.586.8 Giac [F]	5555
3.586.9 Mupad [F(-1)]	5555

3.586.1 Optimal result

Integrand size = 33, antiderivative size = 372

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx =$$

$$\frac{(a^3(A-B) - 3ab^2(A-B) - 3a^2b(A+B) + b^3(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{(a^3(A-B) - 3ab^2(A-B) - 3a^2b(A+B) + b^3(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{2b^2(3Ab + 7aB)}{3d\sqrt{\cot(c+dx)}} - \frac{2a^2(3aA + bB)\sqrt{\cot(c+dx)}}{3d} + \frac{2bB(b + a \cot(c+dx))^2}{3d \cot^{\frac{3}{2}}(c+dx)}$$

$$- \frac{(3a^2b(A-B) - b^3(A-B) + a^3(A+B) - 3ab^2(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{(3a^2b(A-B) - b^3(A-B) + a^3(A+B) - 3ab^2(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

output $\frac{2}{3}bB(b+a\cot(dx+c))^2/d/\cot(dx+c)^{(3/2)}+1/2*(a^3*(A-B)-3*a*b^2*(A-B)-3*a^2*b*(A+B)+b^3*(A+B))*\arctan(-1+2^{(1/2)}*\cot(dx+c)^{(1/2)})/d*2^{(1/2)}+1/2*(a^3*(A-B)-3*a*b^2*(A-B)-3*a^2*b*(A+B)+b^3*(A+B))*\arctan(1+2^{(1/2)}*\cot(dx+c)^{(1/2)})/d*2^{(1/2)}-1/4*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*\ln(1+\cot(dx+c)-2^{(1/2)}*\cot(dx+c)^{(1/2)})/d*2^{(1/2)}+1/4*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*\ln(1+\cot(dx+c)+2^{(1/2)}*\cot(dx+c)^{(1/2)})/d*2^{(1/2)}+2/3*b^2*(3*A*b+7*B*a)/d/\cot(dx+c)^{(1/2)}-2/3*a^2*(3*A*a+B*b)*\cot(dx+c)^{(1/2)}/d$

3.586.2 Mathematica [A] (verified)

Time = 2.24 (sec) , antiderivative size = 270, normalized size of antiderivative = 0.73

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx$$

$$= \frac{2\sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}\left(\frac{(a^3(A-B)+3ab^2(-A+B)-3a^2b(A+B)+b^3(A+B))(\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})-\arctan(1+\sqrt{2}\sqrt{\tan(c+dx)})}{2\sqrt{2}}\right)}{\dots}$$

input `Integrate[Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output $(2*\text{Sqrt}[\text{Cot}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]*(((a^3*(A - B) + 3*a*b^2*(-A + B) - 3*a^2*b*(A + B) + b^3*(A + B))*(\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]] - \text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]])))/(2*\text{Sqrt}[2]) - ((3*a^2*b*(A - B) + b^3*(-A + B) + a^3*(A + B) - 3*a*b^2*(A + B))*(\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]] - \text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]])))/(4*\text{Sqrt}[2]) - (a^3*A)/\text{Sqrt}[\text{Tan}[c + d*x]] + b^2*(A*b + 3*a*B)*\text{Sqrt}[\text{Tan}[c + d*x]] + (b^3*B*\text{Tan}[c + d*x]^(3/2))/3)/d$

3.586.3 Rubi [A] (verified)

Time = 1.32 (sec) , antiderivative size = 302, normalized size of antiderivative = 0.81, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.636$, Rules used = {3042, 4064, 3042, 4088, 27, 3042, 4118, 25, 3042, 4113, 3042, 4017, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$3.586. \quad \int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx$$

$$\begin{aligned}
& \int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
& \quad \downarrow \text{3042} \\
& \int \cot(c+dx)^{3/2}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
& \quad \downarrow \text{4064} \\
& \int \frac{(a \cot(c+dx)+b)^3(A \cot(c+dx)+B)}{\cot^{\frac{5}{2}}(c+dx)} dx \\
& \quad \downarrow \text{3042} \\
& \int \frac{(b-a \tan(c+dx+\frac{\pi}{2}))^3(B-A \tan(c+dx+\frac{\pi}{2}))}{(-\tan(c+dx+\frac{\pi}{2}))^{5/2}} dx \\
& \quad \downarrow \text{4088} \\
& \frac{2bB(a \cot(c+dx)+b)^2}{3d \cot^{\frac{3}{2}}(c+dx)} - \\
& \frac{2}{3} \int -\frac{(b+a \cot(c+dx))(a(3aA+bB) \cot^2(c+dx)+3(Ba^2+2Aba-b^2B) \cot(c+dx)+b(3Ab+7aB))}{2 \cot^{\frac{3}{2}}(c+dx)} dx \\
& \quad \downarrow \text{27} \\
& \frac{1}{3} \int \frac{(b+a \cot(c+dx))(a(3aA+bB) \cot^2(c+dx)+3(Ba^2+2Aba-b^2B) \cot(c+dx)+b(3Ab+7aB))}{\cot^{\frac{3}{2}}(c+dx)} dx + \\
& \quad \frac{2bB(a \cot(c+dx)+b)^2}{3d \cot^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{3042} \\
& \frac{1}{3} \int \frac{(b-a \tan(c+dx+\frac{\pi}{2})) \left(a(3aA+bB) \tan(c+dx+\frac{\pi}{2})^2 - 3(Ba^2+2Aba-b^2B) \tan(c+dx+\frac{\pi}{2}) + b(3Ab+7aB) \right)}{(-\tan(c+dx+\frac{\pi}{2}))^{3/2}} dx + \\
& \quad \frac{2bB(a \cot(c+dx)+b)^2}{3d \cot^{\frac{3}{2}}(c+dx)} \\
& \quad \downarrow \text{4118} \\
& \frac{1}{3} \left(\frac{2b^2(7aB+3Ab)}{d\sqrt{\cot(c+dx)}} - \int -\frac{a^2(3aA+bB) \cot^2(c+dx)+3(Ba^3+3Aba^2-3b^2Ba-Ab^3) \cot(c+dx)+b(10Ba^2+3Ab^2)}{\sqrt{\cot(c+dx)}} dx \right) \\
& \quad \frac{2bB(a \cot(c+dx)+b)^2}{3d \cot^{\frac{3}{2}}(c+dx)}
\end{aligned}$$

3.586. $\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

↓ 25

$$\frac{1}{3} \left(\int \frac{a^2(3aA + bB) \cot^2(c + dx) + 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c + dx) + b(10Ba^2 + 9Aba - 3b^2B)}{\sqrt{\cot(c + dx)}} dx + \frac{2bB(a \cot(c + dx) + b)^2}{3d \cot^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{3} \left(\int \frac{a^2(3aA + bB) \tan(c + dx + \frac{\pi}{2})^2 - 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx + \frac{\pi}{2}) + b(10Ba^2 + 9Aba - 3b^2B)}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx + \frac{2bB(a \cot(c + dx) + b)^2}{3d \cot^{\frac{3}{2}}(c + dx)} \right)$$

↓ 4113

$$\frac{1}{3} \left(\int \frac{3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c + dx) - 3(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)}{\sqrt{\cot(c + dx)}} dx - \frac{2a^2(3aA + bB) \sqrt{\cot(c + dx)}}{d} + \frac{2bB(a \cot(c + dx) + b)^2}{3d \cot^{\frac{3}{2}}(c + dx)} \right)$$

↓ 3042

$$\frac{1}{3} \left(\int \frac{-3(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) - 3(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx - \frac{2a^2(3aA + bB) \sqrt{\cot(c + dx)}}{d} + \frac{2bB(a \cot(c + dx) + b)^2}{3d \cot^{\frac{3}{2}}(c + dx)} \right)$$

↓ 4017

$$\frac{1}{3} \left(2 \int \frac{3(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) - (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c + dx)}{\cot^2(c + dx) + 1} d \sqrt{\cot(c + dx)} - \frac{2a^2(3aA + bB) \sqrt{\cot(c + dx)}}{d} + \frac{2bB(a \cot(c + dx) + b)^2}{3d \cot^{\frac{3}{2}}(c + dx)} \right)$$

↓ 27

$$\frac{1}{3} \left(\frac{6 \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B - (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} - \frac{2a^2(3aA + bB)\sqrt{\cot(c+dx)}}{d} + \frac{2b}{d} \right. \\ \left. \frac{2bB(a \cot(c+dx) + b)^2}{3d \cot^{\frac{3}{2}}(c+dx)} \right. \\ \left. \downarrow 1482 \right.$$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(a^3(A-B) - \dots \right)}{d} \right. \\ \left. \frac{2bB(a \cot(c+dx) + b)^2}{3d \cot^{\frac{3}{2}}(c+dx)} \right. \\ \left. \downarrow 1476 \right.$$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(a^3(A-B) - \dots \right)}{d} \right. \\ \left. \frac{2bB(a \cot(c+dx) + b)^2}{3d \cot^{\frac{3}{2}}(c+dx)} \right. \\ \left. \downarrow 1082 \right.$$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(a^3(A-B) - \dots \right)}{d} \right. \\ \left. \frac{2bB(a \cot(c+dx) + b)^2}{3d \cot^{\frac{3}{2}}(c+dx)} \right. \\ \left. \downarrow 217 \right.$$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(a^3(A-B) - \dots \right)}{d} \right. \\ \left. \frac{2bB(a \cot(c+dx) + b)^2}{3d \cot^{\frac{3}{2}}(c+dx)} \right. \\ \left. \downarrow 1479 \right.$$

3.586. $\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx$

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \right) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \int -\frac{\sqrt{2}}{\cot(c+dx)} dx}{\dots} \right)}{\dots} \right)$$

$$\frac{2bB(a \cot(c+dx) + b)^2}{3d \cot^{\frac{3}{2}}(c+dx)}$$

↓ 25

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \right) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \int \frac{\sqrt{2}}{\cot(c+dx)} dx}{\dots} \right)}{\dots} \right)$$

$$\frac{2bB(a \cot(c+dx) + b)^2}{3d \cot^{\frac{3}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{3} \left(\frac{6 \left(\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \right) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}}{\cot(c+dx)} dx}{\dots} \right)}{\dots} \right)$$

$$\frac{2bB(a \cot(c+dx) + b)^2}{3d \cot^{\frac{3}{2}}(c+dx)}$$

↓ 1103

$$\frac{1}{3} \left(-\frac{2a^2(3aA + bB)\sqrt{\cot(c+dx)}}{d} + \frac{6 \left(\frac{1}{2}(a^3(A-B) - 3a^2b(A+B) - 3ab^2(A-B) + b^3(A+B)) \right) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)}}{\dots}\right)}{\dots} \right)}{\dots} \right)$$

$$\frac{2bB(a \cot(c+dx) + b)^2}{3d \cot^{\frac{3}{2}}(c+dx)}$$

input `Int[Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

```
output (2*b*B*(b + a*Cot[c + d*x])^2)/(3*d*Cot[c + d*x]^(3/2)) + ((2*b^2*(3*A*b +
7*a*B))/(d*Sqrt[Cot[c + d*x]]) - (2*a^2*(3*a*A + b*B)*Sqrt[Cot[c + d*x]])
/d + (6*(((a^3*(A - B) - 3*a*b^2*(A - B) - 3*a^2*b*(A + B) + b^3*(A + B))*
(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sq
rt[Cot[c + d*x]]/Sqrt[2]))/2 + ((3*a^2*b*(A - B) - b^3*(A - B) + a^3*(A +
B) - 3*a*b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c +
d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt
[2])))/2))/d)/3
```

3.586.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 217 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

```
rule 1103 Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

```
rule 1476 Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

```
rule 4113 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*((a +
b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1))), x] + Int[(a + b*Tan[e + f*x])^m*Simp[A - C + B*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B, C, m}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && !LeQ[m, -1]
```

```
rule 4118 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)
*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)
*(x_)]^2), x_Symbol] := Simp[(-(b*c - a*d))*(c^2*C - B*c*d + A*d^2)*((c + d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2 + d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n, -1]
```

3.586.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1103 vs. $2(332) = 664$.

Time = 0.45 (sec) , antiderivative size = 1104, normalized size of antiderivative = 2.97

method	result	size
derivativedivides	Expression too large to display	1104
default	Expression too large to display	1104

```
input int(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

$$3.586. \quad \int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

output $1/12/d*(1/\tan(dx+c))^{3/2}*\tan(dx+c)*(18*B*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2}))*2^{1/2}*\tan(dx+c)^{1/2}*a^2*b-18*B*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2}))*2^{1/2}*\tan(dx+c)^{1/2}*a*b^2+9*B*\ln(-(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1)/(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c)))*2^{1/2}*\tan(dx+c)^{1/2})*a^2*b+9*A*\ln(-(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c))/(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1))*2^{1/2}*\tan(dx+c)^{1/2}*a^2*b+18*A*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2}))*2^{1/2}*\tan(dx+c)^{1/2}*a^2*b+18*A*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2}))*2^{1/2}*\tan(dx+c)^{1/2}*a*b^2+18*A*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2}))*2^{1/2}*\tan(dx+c)^{1/2}*a^2*b+18*A*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2}))*2^{1/2}*\tan(dx+c)^{1/2}*a*b^2+9*A*\ln(-(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1)/(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c)))*2^{1/2}*\tan(dx+c)^{1/2}*a*b^2-9*B*\ln(-(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c))/(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1))*2^{1/2}*\tan(dx+c)^{1/2}*a*b^2+18*B*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2}))*2^{1/2}*\tan(dx+c)^{1/2}*a^2*b-18*B*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2}))*2^{1/2}*\tan(dx+c)^{1/2}*a*b^2+72*B*a*b^2*\tan(dx+c)-24*A*a^3-3*B*\ln(-(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1)/(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c)))*2^{1/2}*\tan(dx+c)^{1/2}*b^3+8*B*b^3*\tan(dx+c)^2+24*A*b^3*\tan(dx+c)-6*B*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2}))*2^{1/2}*\tan(dx+c)^{1/2}*b^3-3*A*\ln(-(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c))/(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1))*2^{1/2}*\tan(dx+c)^{1/2}*b^3-6*A*\arct...$

3.586.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6105 vs. $2(332) = 664$.

Time = 6.20 (sec) , antiderivative size = 6105, normalized size of antiderivative = 16.41

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx = \text{Too large to display}$$

input `integrate(cot(dx+c)^(3/2)*(a+b*tan(dx+c))^3*(A+B*tan(dx+c)),x, algorithm="fracas")`

output Too large to include

3.586.6 Sympy [F]

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \int (A + B \tan(c + dx))(a + b \tan(c + dx))^3 \cot^{\frac{3}{2}}(c + dx) dx$$

input `integrate(cot(d*x+c)**(3/2)*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**3*cot(c + d*x)**(3/2), x)`

3.586.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 316, normalized size of antiderivative = 0.85

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx =$$

$$\frac{24 A a^3}{\sqrt{\tan(dx+c)}} - 6 \sqrt{2}((A - B)a^3 - 3(A + B)a^2b - 3(A - B)ab^2 + (A + B)b^3) \arctan\left(\frac{1}{2} \sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right)$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `-1/12*(24*A*a^3/sqrt(tan(d*x + c)) - 6*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) - 6*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - 3*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + 3*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - 8*(B*b^3 + 3*(3*B*a*b^2 + A*b^3)/tan(d*x + c))*tan(d*x + c)^(3/2)/d`

3.586.8 Giac [F]

$$\begin{aligned} & \int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^3 \cot(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^3*cot(d*x + c)^(3/2), x)`

3.586.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx \\ &= \int \cot(c + dx)^{3/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^3 dx \end{aligned}$$

input `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`

output `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3, x)`

3.587 $\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

3.587.1 Optimal result	5556
3.587.2 Mathematica [A] (verified)	5557
3.587.3 Rubi [A] (verified)	5558
3.587.4 Maple [B] (verified)	5565
3.587.5 Fricas [B] (verification not implemented)	5566
3.587.6 Sympy [F]	5567
3.587.7 Maxima [A] (verification not implemented)	5567
3.587.8 Giac [F]	5568
3.587.9 Mupad [F(-1)]	5568

3.587.1 Optimal result

Integrand size = 33, antiderivative size = 380

$$\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$$

$$= \frac{(3a^2b(A - B) - b^3(A - B) + a^3(A + B) - 3ab^2(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(3a^2b(A - B) - b^3(A - B) + a^3(A + B) - 3ab^2(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{2b^2(5Ab + 9aB)}{15d \cot^{\frac{3}{2}}(c + dx)} + \frac{2b(15aAb + 14a^2B - 5b^2B)}{5d\sqrt{\cot(c + dx)}} + \frac{2bB(b + a \cot(c + dx))^2}{5d \cot^{\frac{5}{2}}(c + dx)}$$

$$- \frac{(a^3(A - B) - 3ab^2(A - B) - 3a^2b(A + B) + b^3(A + B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

$$+ \frac{(a^3(A - B) - 3ab^2(A - B) - 3a^2b(A + B) + b^3(A + B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

3.587. $\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

output $\frac{2}{15}b^2(5A^2b+9B^2a)/d/\cot(dx+c)^{3/2}+2/5*b*B*(b+a*\cot(dx+c))^2/d/\cot(dx+c)^{5/2}-1/2*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*\arctan(-1+2^{1/2}*\cot(dx+c)^{1/2})/d*2^{1/2}-1/2*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*\arctan(1+2^{1/2}*\cot(dx+c)^{1/2})/d*2^{1/2}-1/4*(a^3*(A-B)-3*a*b^2*(A-B)-3*a^2*b*(A+B)+b^3*(A+B))*\ln(1+\cot(dx+c)-2^{1/2}*\cot(dx+c)^{1/2})/d*2^{1/2}+1/4*(a^3*(A-B)-3*a*b^2*(A-B)-3*a^2*b*(A+B)+b^3*(A+B))*\ln(1+\cot(dx+c)+2^{1/2}*\cot(dx+c)^{1/2})/d*2^{1/2}+2/5*b*(15*A*a*b+14*B*a^2-5*B*b^2)/d/\cot(dx+c)^{1/2}$

3.587.2 Mathematica [A] (verified)

Time = 1.32 (sec) , antiderivative size = 287, normalized size of antiderivative = 0.76

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx$$

$$= \frac{2\sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}\left(-\frac{(3a^2b(A-B)+b^3(-A+B)+a^3(A+B)-3ab^2(A+B))\left(\arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)-\arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)\right)}{2\sqrt{2}}\right)}{d}$$

input `Integrate[Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output $(2*\text{Sqrt}[\text{Cot}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]*(-1/2*((3*a^2*b*(A - B) + b^3*(-A + B) + a^3*(A + B) - 3*a*b^2*(A + B))*(\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]] - \text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]]))/\text{Sqrt}[2] - ((a^3*(A - B) + 3*a*b^2*(-A + B) - 3*a^2*b*(A + B) + b^3*(A + B))*(\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]] - \text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]])))/(4*\text{Sqrt}[2]) + b*(3*a*A*b + 3*a^2*B - b^2*B)*\text{Sqrt}[\text{Tan}[c + d*x]] + (b^2*(A*b + 3*a*B)*\text{Tan}[c + d*x]^{3/2})/3 + (b^3*B*\text{Tan}[c + d*x]^{5/2})/5)/d$

3.587.3 Rubi [A] (verified)

Time = 1.39 (sec) , antiderivative size = 312, normalized size of antiderivative = 0.82, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4064, 3042, 4088, 27, 3042, 4118, 25, 3042, 4111, 27, 3042, 4017, 25, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{(a \cot(c+dx)+b)^3(A \cot(c+dx)+B)}{\cot^{\frac{7}{2}}(c+dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b-a \tan(c+dx+\frac{\pi}{2}))^3(B-A \tan(c+dx+\frac{\pi}{2}))}{(-\tan(c+dx+\frac{\pi}{2}))^{7/2}} dx \\
 & \quad \downarrow \text{4088} \\
 & \frac{2bB(a \cot(c+dx)+b)^2}{5d \cot^{\frac{5}{2}}(c+dx)} - \\
 & \frac{2}{5} \int -\frac{(b+a \cot(c+dx))(a(5aA-bB) \cot^2(c+dx)+5(Ba^2+2Aba-b^2B) \cot(c+dx)+b(5Ab+9aB))}{2 \cot^{\frac{5}{2}}(c+dx)} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{5} \int \frac{(b+a \cot(c+dx))(a(5aA-bB) \cot^2(c+dx)+5(Ba^2+2Aba-b^2B) \cot(c+dx)+b(5Ab+9aB))}{\cot^{\frac{5}{2}}(c+dx)} dx + \\
 & \quad \frac{2bB(a \cot(c+dx)+b)^2}{5d \cot^{\frac{5}{2}}(c+dx)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{1}{5} \int \frac{(b - a \tan(c + dx + \frac{\pi}{2})) (a(5aA - bB) \tan(c + dx + \frac{\pi}{2})^2 - 5(Ba^2 + 2Aba - b^2B) \tan(c + dx + \frac{\pi}{2}) + b(5aA - bB))}{(-\tan(c + dx + \frac{\pi}{2}))^{5/2} \frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}}$$

↓ 4118

$$\frac{1}{5} \left(\frac{2b^2(9aB + 5Ab)}{3d \cot^{\frac{3}{2}}(c + dx)} - \int - \frac{a^2(5aA - bB) \cot^2(c + dx) + 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c + dx) + b(14Ba^2 + 15Aba - 5b^2B)}{\cot^{\frac{3}{2}}(c + dx)} \frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)} \right)$$

↓ 25

$$\frac{1}{5} \left(\int \frac{a^2(5aA - bB) \cot^2(c + dx) + 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c + dx) + b(14Ba^2 + 15Aba - 5b^2B)}{\cot^{\frac{3}{2}}(c + dx)} \frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)} dx \right)$$

↓ 3042

$$\frac{1}{5} \left(\int \frac{a^2(5aA - bB) \tan(c + dx + \frac{\pi}{2})^2 - 5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx + \frac{\pi}{2}) + b(14Ba^2 + 15Aba - 5b^2B)}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2} \frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}} \right)$$

↓ 4111

$$\frac{1}{5} \left(\int \frac{5(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 + (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \cot(c + dx))}{\sqrt{\cot(c + dx)} \frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}} dx + \frac{2b(14a^2B + 15aAb - 5b^2B)}{d\sqrt{\cot(c + dx)}} \right)$$

↓ 27

$$\frac{1}{5} \left(5 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 + (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \cot(c + dx)}{\sqrt{\cot(c + dx)} \frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}} dx + \frac{2b(14a^2B + 15aAb - 5b^2B)}{d\sqrt{\cot(c + dx)}} \right)$$

↓ 3042

3.587. $\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\frac{1}{5} \left(5 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx + \frac{2b(14a^2B + 15aAb)}{d\sqrt{\cot(c + dx)}} \right) + \frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}$$

↓ 4017

$$\frac{1}{5} \left(\frac{10 \int -\frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 + (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} + \frac{2b(14a^2B + 15aAb - 5b^2B)}{d\sqrt{\cot(c + dx)}} \right) + \frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}$$

↓ 25

$$\frac{1}{5} \left(-\frac{10 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 + (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} + \frac{2b(14a^2B + 15aAb - 5b^2B)}{d\sqrt{\cot(c + dx)}} \right) + \frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}$$

↓ 1482

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \frac{1}{2}(a^3(A + B) + b^3(A - B)) \int \frac{1 + \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} \right)}{d} \right) + \frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}$$

↓ 1476

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \frac{1}{2}(a^3(A + B) + b^3(A - B)) \int \frac{1 + \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} \right)}{d} \right) + \frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}$$

↓ 1082

3.587. $\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3(A + B \tan(c + dx)) dx$

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a^3(A + B) - \dots \right)}{\dots} \right)$$

$$\frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}$$

↓ 217

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a^3(A + B) - \dots \right)}{\dots} \right)$$

$$\frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}$$

↓ 1479

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \dots \right)}{\dots} \right)}{\dots} \right)$$

$$\frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}$$

↓ 25

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2}(a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right)}{\dots} \right)}{\dots} \right)$$

$$\frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{5} \left(\frac{10 \left(\frac{1}{2} (a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \right) \left(\int \frac{\sqrt{2} - 2\sqrt{\cot(c+dx)}}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)} + 1} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)} dx \right)}{\frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}} \right)$$

↓ 1103

$$\frac{1}{5} \left(\frac{2b(14a^2B + 15aAb - 5b^2B)}{d\sqrt{\cot(c + dx)}} + \frac{10 \left(\frac{1}{2} (a^3(A - B) - 3a^2b(A + B) - 3ab^2(A - B) + b^3(A + B)) \right) \left(\frac{\log(\cot(c+dx) + \sqrt{\cot(c+dx)})}{\cot(c+dx) + \sqrt{\cot(c+dx)}} \right)}{\frac{2bB(a \cot(c + dx) + b)^2}{5d \cot^{\frac{5}{2}}(c + dx)}} \right)$$

input `Int[Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]),x]`

output `(2*b*B*(b + a*Cot[c + d*x])^2)/(5*d*Cot[c + d*x]^(5/2)) + ((2*b^2*(5*A*b + 9*a*B))/(3*d*Cot[c + d*x]^(3/2)) + (2*b*(15*a*A*b + 14*a^2*B - 5*b^2*B))/(d*Sqrt[Cot[c + d*x]]) + (10*(-1/2*((3*a^2*b*(A - B) - b^3*(A - B) + a^3*(A + B) - 3*a*b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2])) + ((a^3*(A - B) - 3*a*b^2*(A - B) - 3*a^2*b*(A + B) + b^3*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d)/5`

3.587.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

- rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4111 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A - C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]`

```
rule 4118 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.
)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)])^2), x_Symbol] := Simp[(-(b*c - a*d))*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2)) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]
```

3.587.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 955 vs. $2(340) = 680$.

Time = 0.45 (sec) , antiderivative size = 956, normalized size of antiderivative = 2.52

method	result	size
derivativedivides	Expression too large to display	956
default	Expression too large to display	956

```
input int(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x,method=_RETURNV
ERBOSE)
```

output $1/60/d*(1/\tan(dx+c))^{1/2}*\tan(dx+c)^{1/2}*(15*A*\ln(-(1+2^{1/2}*\tan(dx+c))^{1/2}+\tan(dx+c)))/(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1))*2^{1/2}*a^3+30*A*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*a^3-30*A*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*b^3+30*A*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*(1/2)*a^3-30*A*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*b^3-15*A*\ln(-(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1)/(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c))))*2^{1/2}*b^3+15*B*\ln(-(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c)))/(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1))*2^{1/2}*b^3+30*B*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*a^3+30*B*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*b^3+30*B*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*a^3+30*B*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*b^3+15*B*\ln(-(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1)/(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c)))*2^{1/2}*a^3+120*B*a*b^2*\tan(dx+c)^{3/2}+360*A*a*b^2*\tan(dx+c)^{1/2}+360*B*a^2*b*\tan(dx+c)^{1/2}+24*B*b^3*\tan(dx+c)^{5/2}+40*A*b^3*\tan(dx+c)^{3/2}-120*B*b^3*\tan(dx+c)^{1/2}-45*A*\ln(-(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c)))/(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1))*2^{1/2}*a*b^2+90*A*\arctan(1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*a^2*b-90*A*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*a^2*b-90*A*\arctan(-1+2^{1/2}*\tan(dx+c)^{1/2})*2^{1/2}*a*b^2+45*A*\ln(-(2^{1/2}*\tan(dx+c)^{1/2}-\tan(dx+c)-1)/(1+2^{1/2}*\tan(dx+c)^{1/2}+\tan(dx+c)))*2^{1/2}*a^2*b-45*B*\ln(-1...$

3.587.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6147 vs. $2(340) = 680$.

Time = 6.19 (sec) , antiderivative size = 6147, normalized size of antiderivative = 16.18

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx = \text{Too large to display}$$

input `integrate(cot(dx+c)^(1/2)*(a+b*tan(dx+c))^3*(A+B*tan(dx+c)),x, algorithm="fracas")`

output Too large to include

3.587.6 Sympy [F]

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx$$

$$= \int (A+B\tan(c+dx))(a+b\tan(c+dx))^3\sqrt{\cot(c+dx)}dx$$

input `integrate(cot(d*x+c)**(1/2)*(a+b*tan(d*x+c))**3*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**3*sqrt(cot(c + d*x)), x)`

3.587.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 334, normalized size of antiderivative = 0.88

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^3(A+B\tan(c+dx))dx$$

$$= \frac{8\left(3Bb^3 + \frac{5(3Bab^2+Ab^3)}{\tan(dx+c)} + \frac{15(3Ba^2b+3Aab^2-Bb^3)}{\tan(dx+c)^2}\right)\tan(dx+c)^{\frac{5}{2}} - 30\sqrt{2}((A+B)a^3 + 3(A-B)a^2b - 3(A$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `1/60*(8*(3*B*b^3 + 5*(3*B*a*b^2 + A*b^3)/tan(d*x + c) + 15*(3*B*a^2*b + 3*A*a*b^2 - B*b^3)/tan(d*x + c)^2)*tan(d*x + c)^(5/2) - 30*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) - 30*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + 15*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - 15*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))/d`

3.587.8 Giac [F]

$$\begin{aligned} & \int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\ &= \int (B \tan(dx+c)+A)(b \tan(dx+c)+a)^3 \sqrt{\cot(dx+c)} dx \end{aligned}$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^3*(A+B*tan(d*x+c)),x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^3*sqrt(cot(d*x + c)), x)`

3.587.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^3(A+B \tan(c+dx)) dx \\ &= \int \sqrt{\cot(c+dx)}(A+B \tan(c+dx))(a+b \tan(c+dx))^3 dx \end{aligned}$$

input `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3,x)`

output `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3, x)`

3.588
$$\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.588.1 Optimal result	5569
3.588.2 Mathematica [A] (verified)	5570
3.588.3 Rubi [A] (verified)	5571
3.588.4 Maple [A] (verified)	5578
3.588.5 Fricas [B] (verification not implemented)	5579
3.588.6 Sympy [F]	5579
3.588.7 Maxima [A] (verification not implemented)	5580
3.588.8 Giac [F]	5580
3.588.9 Mupad [F(-1)]	5581

3.588.1 Optimal result

Integrand size = 33, antiderivative size = 421

$$\int \frac{(a + b \tan(c + dx))^3(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \frac{(a^3(A - B) - 3ab^2(A - B) - 3a^2b(A + B) + b^3(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d}$$

$$- \frac{(a^3(A - B) - 3ab^2(A - B) - 3a^2b(A + B) + b^3(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d}$$

$$+ \frac{2b^2(7Ab + 11aB)}{35d \cot^{\frac{5}{2}}(c + dx)} + \frac{2b(21aAb + 18a^2B - 7b^2B)}{21d \cot^{\frac{3}{2}}(c + dx)}$$

$$+ \frac{2(3a^2Ab - Ab^3 + a^3B - 3ab^2B)}{d\sqrt{\cot(c + dx)}} + \frac{2bB(b + a \cot(c + dx))^2}{7d \cot^{\frac{7}{2}}(c + dx)}$$

$$+ \frac{(3a^2b(A - B) - b^3(A - B) + a^3(A + B) - 3ab^2(A + B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

$$- \frac{(3a^2b(A - B) - b^3(A - B) + a^3(A + B) - 3ab^2(A + B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

output $\frac{2}{35}b^2(7A*b+11B*a)/d/\cot(dx+c)^{(5/2)}+2/21*b*(21A*a*b+18B*a^2-7B*b^2)/d/\cot(dx+c)^{(3/2)}+2/7*b*B*(b+a*\cot(dx+c))^2/d/\cot(dx+c)^{(7/2)}-1/2*(a^3*(A-B)-3*a*b^2*(A-B)-3*a^2*b*(A+B)+b^3*(A+B))*\arctan(-1+2^{(1/2)}*\cot(dx+c)^{(1/2)})/d*2^{(1/2)}-1/2*(a^3*(A-B)-3*a*b^2*(A-B)-3*a^2*b*(A+B)+b^3*(A+B))*\arctan(1+2^{(1/2)}*\cot(dx+c)^{(1/2)})/d*2^{(1/2)}+1/4*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*\ln(1+\cot(dx+c)-2^{(1/2)}*\cot(dx+c)^{(1/2)})/d*2^{(1/2)}-1/4*(3*a^2*b*(A-B)-b^3*(A-B)+a^3*(A+B)-3*a*b^2*(A+B))*\ln(1+\cot(dx+c)+2^{(1/2)}*\cot(dx+c)^{(1/2)})/d*2^{(1/2)}+2*(3A*a^2*b-A*b^3+B*a^3-3B*a*b^2)/d/\cot(dx+c)^{(1/2)}$

3.588.2 Mathematica [A] (verified)

Time = 2.69 (sec) , antiderivative size = 327, normalized size of antiderivative = 0.78

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \frac{2\sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}\left(-\frac{(a^3(A-B)+3ab^2(-A+B)-3a^2b(A+B)+b^3(A+B))(\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})-\arctan(1+\sqrt{2}\sqrt{\tan(c+dx)})}{2\sqrt{2}}\right)}{1}$$

input `Integrate[((a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output $(2*\text{Sqrt}[\text{Cot}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]*(-1/2*((a^3*(A - B) + 3*a*b^2*(-A + B) - 3*a^2*b*(A + B) + b^3*(A + B))*(\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]] - \text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]]))/\text{Sqrt}[2] + ((3*a^2*b*(A - B) + b^3*(-A + B) + a^3*(A + B) - 3*a*b^2*(A + B))*(\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]] - \text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]])))/(4*\text{Sqrt}[2]) + (3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B)*\text{Sqrt}[\text{Tan}[c + d*x]] + (b*(3*a*A*b + 3*a^2*B - b^2*B)*\text{Tan}[c + d*x]^{(3/2)})/3 + (b^2*(A*b + 3*a*B)*\text{Tan}[c + d*x]^{(5/2)})/5 + (b^3*B*\text{Tan}[c + d*x]^{(7/2)})/7)/d$

3.588.3 Rubi [A] (verified)

Time = 1.64 (sec) , antiderivative size = 358, normalized size of antiderivative = 0.85, number of steps used = 25, number of rules used = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$, Rules used = {3042, 4064, 3042, 4088, 27, 3042, 4118, 25, 3042, 4111, 27, 3042, 4012, 3042, 4017, 25, 1482, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{(a \cot(c + dx) + b)^3 (A \cot(c + dx) + B)}{\cot^{\frac{9}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(b - a \tan(c + dx + \frac{\pi}{2}))^3 (B - A \tan(c + dx + \frac{\pi}{2}))}{(-\tan(c + dx + \frac{\pi}{2}))^{9/2}} dx \\
 & \quad \downarrow \text{4088} \\
 & \frac{2bB(a \cot(c + dx) + b)^2}{7d \cot^{\frac{7}{2}}(c + dx)} - \\
 & \frac{2}{7} \int \frac{(b + a \cot(c + dx)) (a(7aA - 3bB) \cot^2(c + dx) + 7(Ba^2 + 2Aba - b^2B) \cot(c + dx) + b(7Ab + 11aB))}{2 \cot^{\frac{7}{2}}(c + dx)} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{7} \int \frac{(b + a \cot(c + dx)) (a(7aA - 3bB) \cot^2(c + dx) + 7(Ba^2 + 2Aba - b^2B) \cot(c + dx) + b(7Ab + 11aB))}{\cot^{\frac{7}{2}}(c + dx)} dx + \\
 & \quad \frac{2bB(a \cot(c + dx) + b)^2}{7d \cot^{\frac{7}{2}}(c + dx)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{1}{7} \int \frac{(b - a \tan(c + dx + \frac{\pi}{2})) (a(7aA - 3bB) \tan(c + dx + \frac{\pi}{2})^2 - 7(Ba^2 + 2Aba - b^2B) \tan(c + dx + \frac{\pi}{2}) + b(7aA - 3bB))}{(-\tan(c + dx + \frac{\pi}{2}))^{7/2} \frac{2bB(a \cot(c + dx) + b)^2}{7d \cot^{\frac{7}{2}}(c + dx)}} dx$$

↓ 4118

$$\frac{1}{7} \left(\frac{2b^2(11aB + 7Ab)}{5d \cot^{\frac{5}{2}}(c + dx)} - \int \frac{a^2(7aA - 3bB) \cot^2(c + dx) + 7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c + dx) + b(18Ba^2 + 21Aba - 7b^2B)}{\cot^{\frac{5}{2}}(c + dx)} dx \right) \frac{2bB(a \cot(c + dx) + b)^2}{7d \cot^{\frac{7}{2}}(c + dx)}$$

↓ 25

$$\frac{1}{7} \left(\int \frac{a^2(7aA - 3bB) \cot^2(c + dx) + 7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c + dx) + b(18Ba^2 + 21Aba - 7b^2B)}{\cot^{\frac{5}{2}}(c + dx)} dx \right) \frac{2bB(a \cot(c + dx) + b)^2}{7d \cot^{\frac{7}{2}}(c + dx)}$$

↓ 3042

$$\frac{1}{7} \left(\int \frac{a^2(7aA - 3bB) \tan(c + dx + \frac{\pi}{2})^2 - 7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \tan(c + dx + \frac{\pi}{2}) + b(18Ba^2 + 21Aba - 7b^2B)}{(-\tan(c + dx + \frac{\pi}{2}))^{5/2} \frac{2bB(a \cot(c + dx) + b)^2}{7d \cot^{\frac{7}{2}}(c + dx)}} dx \right)$$

↓ 4111

$$\frac{1}{7} \left(\int \frac{7(Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 + (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \cot(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx + \frac{2b(18a^2B + 21aAb - 7b^2B)}{3d \cot^{\frac{3}{2}}(c + dx)} \right) \frac{2bB(a \cot(c + dx) + b)^2}{7d \cot^{\frac{7}{2}}(c + dx)}$$

↓ 27

$$\frac{1}{7} \left(7 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 + (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \cot(c + dx)}{\cot^{\frac{3}{2}}(c + dx)} dx + \frac{2b(18a^2B + 21aAb - 7b^2B)}{3d \cot^{\frac{3}{2}}(c + dx)} \right) \frac{2bB(a \cot(c + dx) + b)^2}{7d \cot^{\frac{7}{2}}(c + dx)}$$

↓ 3042

3.588. $\int \frac{(a+b \tan(c+dx))^3 (A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

$$\frac{1}{7} \left(7 \int \frac{Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3 - (Aa^3 - 3bBa^2 - 3Ab^2a + b^3B) \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2} \frac{2bB(a \cot(c + dx) + b)^2}{7d \cot^{\frac{7}{2}}(c + dx)}} dx + \frac{2b(18a^2B + 21aAb - Ab^2)}{3d \cot^{\frac{3}{2}}(c + dx)} \right)$$

↓ 4012

$$\frac{1}{7} \left(7 \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B - (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c + dx)}{\sqrt{\cot(c + dx)} \frac{2bB(a \cot(c + dx) + b)^2}{7d \cot^{\frac{7}{2}}(c + dx)}} dx + \frac{2(a^3B + 3a^2Ab - 3ab^2B - Ab^3)}{d \sqrt{\cot(c + dx)}} \right)$$

↓ 3042

$$\frac{1}{7} \left(7 \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B - (-Ba^3 - 3Aba^2 + 3b^2Ba + Ab^3) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})} \frac{2bB(a \cot(c + dx) + b)^2}{7d \cot^{\frac{7}{2}}(c + dx)}} dx + \frac{2(a^3B + 3a^2Ab - 3ab^2B - Ab^3)}{d \sqrt{\cot(c + dx)}} \right)$$

↓ 4017

$$\frac{1}{7} \left(7 \int \frac{2 \int -\frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B - (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} + \frac{2(a^3B + 3a^2Ab - 3ab^2B - Ab^3)}{d \sqrt{\cot(c + dx)}} \right)$$

↓ 25

$$\frac{1}{7} \left(7 \int \frac{2(a^3B + 3a^2Ab - 3ab^2B - Ab^3)}{d \sqrt{\cot(c + dx)}} - \frac{2 \int \frac{Aa^3 - 3bBa^2 - 3Ab^2a + b^3B - (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d} \right)$$

↓ 1482

$$\frac{1}{7} \left(7 \int \frac{2\left(-\frac{1}{2}(a^3(A + B) + 3a^2b(A - B) - 3ab^2(A + B) - b^3(A - B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \frac{1}{2}(a^3(A - B) + 3a^2b(A + B) - 3ab^2(A - B) - b^3(A + B)) \int \frac{1 + \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}\right)}{d} \right)$$

↓

3.588. $\int \frac{(a+b \tan(c+dx))^3 (A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

↓ 1476

$$\frac{1}{7} \left(7 \left(\frac{2 \left(-\frac{1}{2} (a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2} (a^3(A-B) - b^3(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{2bB(a \cot(c+dx) + b)^2} \right) \right)$$

↓ 1082

$$\frac{1}{7} \left(7 \left(\frac{2 \left(-\frac{1}{2} (a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2} (a^3(A-B) - b^3(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{2bB(a \cot(c+dx) + b)^2} \right) \right)$$

↓ 217

$$\frac{1}{7} \left(7 \left(\frac{2 \left(-\frac{1}{2} (a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2} (a^3(A-B) - b^3(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{2bB(a \cot(c+dx) + b)^2} \right) \right)$$

↓ 1479

$$\frac{1}{7} \left(7 \left(\frac{2 \left(-\frac{1}{2} (a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) - \frac{1}{2} (a^3(A-B) - b^3(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{2bB(a \cot(c+dx) + b)^2} \right) \right)$$

↓ 25

$$\frac{1}{7} \left(\frac{2 \left(-\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \right) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}+2\sqrt{\cot(c+dx)}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right)}{7d \cot^{\frac{7}{2}}(c+dx)} \right)$$

$$\frac{2bB(a \cot(c+dx) + b)^2}{7d \cot^{\frac{7}{2}}(c+dx)}$$

↓ 27

$$\frac{1}{7} \left(\frac{2 \left(-\frac{1}{2}(a^3(A+B) + 3a^2b(A-B) - 3ab^2(A+B) - b^3(A-B)) \right) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}+2\sqrt{\cot(c+dx)}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right)}{7d \cot^{\frac{7}{2}}(c+dx)} \right)$$

$$\frac{2bB(a \cot(c+dx) + b)^2}{7d \cot^{\frac{7}{2}}(c+dx)}$$

↓ 1103

$$\frac{1}{7} \left(\frac{2b(18a^2B + 21aAb - 7b^2B)}{3d \cot^{\frac{3}{2}}(c+dx)} + 7 \left(\frac{2 \left(-\frac{1}{2}(a^3(A-B) - 3a^2b(A+B) - 3ab^2(A-B) + b^3(A+B)) \right) \left(\arctan(\sqrt{2} \frac{\sqrt{\cot(c+dx)-1}}{\sqrt{\cot(c+dx)+1}}) \right)}{7d \cot^{\frac{7}{2}}(c+dx)} \right) \right)$$

$$\frac{2bB(a \cot(c+dx) + b)^2}{7d \cot^{\frac{7}{2}}(c+dx)}$$

input `Int[((a + b*Tan[c + d*x])^3*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `(2*b*B*(b + a*Cot[c + d*x])^2)/(7*d*Cot[c + d*x]^(7/2)) + ((2*b^2*(7*A*b + 11*a*B))/(5*d*Cot[c + d*x]^(5/2)) + (2*b*(21*a*A*b + 18*a^2*B - 7*b^2*B))/(3*d*Cot[c + d*x]^(3/2)) + 7*((2*(3*a^2*A*b - A*b^3 + a^3*B - 3*a*b^2*B))/(d*Sqrt[Cot[c + d*x]]) + (2*(-1/2*((a^3*(A - B) - 3*a*b^2*(A - B) - 3*a^2*b*(A + B) + b^3*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2])) - ((3*a^2*b*(A - B) - b^3*(A - B) + a^3*(A + B) - 3*a*b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/d)/7`

3.588.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4012 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(b*c - a*d)*((a + b*Tan[e + f*x])^(m + 1)/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*c + b*d - (b*c - a*d)*Tan[e + f*x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && LtQ[m, -1]`

rule 4017 `Int[((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((A*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`


```
rule 4111 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 -
a*b*B + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x
] + Simp[1/(a^2 + b^2) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[b*B + a*(A -
C) - (A*b - a*B - b*C)*Tan[e + f*x], x], x] /; FreeQ[{a, b, e, f, A, B
, C}, x] && NeQ[A*b^2 - a*b*B + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0
]
```

```
rule 4118 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.
)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f
_.)*(x_)]^2), x_Symbol] := Simp[(-(b*c - a*d))*(c^2*C - B*c*d + A*d^2)*((c
+ d*Tan[e + f*x])^(n + 1)/(d^2*f*(n + 1)*(c^2 + d^2))), x] + Simp[1/(d*(c^2
+ d^2) Int[(c + d*Tan[e + f*x])^(n + 1)*Simp[a*d*(A*c - c*C + B*d) + b*
(c^2*C - B*c*d + A*d^2) + d*(A*b*c + a*B*c - b*c*C - a*A*d + b*B*d + a*C*d)
*Tan[e + f*x] + b*C*(c^2 + d^2)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b,
c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[c^2 + d^2, 0] && LtQ[n
, -1]
```

3.588.4 Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 328, normalized size of antiderivative = 0.78

method	result
derivativedivides	$-\frac{2(3Aa^2b - Ab^3 + Ba^3 - 3Bab^2)}{\sqrt{\cot(dx+c)}} - \frac{2Bb^3}{7\cot(dx+c)^{\frac{7}{2}}} - \frac{2b(3abA + 3Ba^2 - Bb^2)}{3\cot(dx+c)^{\frac{3}{2}}} - \frac{2b^2(Ab + 3Ba)}{5\cot(dx+c)^{\frac{5}{2}}} + \frac{(Aa^3 - 3Aab^2 - 3Ba^2b + Bb^3)\sqrt{c+d\tan(dx+c)}}{\cot(dx+c)}$
default	$-\frac{2(3Aa^2b - Ab^3 + Ba^3 - 3Bab^2)}{\sqrt{\cot(dx+c)}} - \frac{2Bb^3}{7\cot(dx+c)^{\frac{7}{2}}} - \frac{2b(3abA + 3Ba^2 - Bb^2)}{3\cot(dx+c)^{\frac{3}{2}}} - \frac{2b^2(Ab + 3Ba)}{5\cot(dx+c)^{\frac{5}{2}}} + \frac{(Aa^3 - 3Aab^2 - 3Ba^2b + Bb^3)\sqrt{c+d\tan(dx+c)}}{\cot(dx+c)}$

```
input int((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2), x, method=_RETURNV
ERBOSE)
```

3.588.
$$\int \frac{(a+b \tan(c+dx))^3(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

output
$$\begin{aligned} & -1/d*(-2*(3*A*a^2*b-A*b^3+B*a^3-3*B*a*b^2)/\cot(d*x+c)^{(1/2)}-2/7*B*b^3/\cot(d*x+c)^{(7/2)}-2/3*b*(3*A*a*b+3*B*a^2-B*b^2)/\cot(d*x+c)^{(3/2)}-2/5*b^2*(A*b+3*B*a)/\cot(d*x+c)^{(5/2)}+1/4*(A*a^3-3*A*a*b^2-3*B*a^2*b+B*b^3)*2^{(1/2)}*(\ln((1+\cot(d*x+c)+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(1+\cot(d*x+c)-2^{(1/2)}*\cot(d*x+c)^{(1/2)}))+2*\arctan(1+2^{(1/2)}*\cot(d*x+c)^{(1/2)}))+2*\arctan(-1+2^{(1/2)}*\cot(d*x+c)^{(1/2)}))+1/4*(-3*A*a^2*b+A*b^3-B*a^3+3*B*a*b^2)*2^{(1/2)}*(\ln((1+\cot(d*x+c)-2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(1+\cot(d*x+c)+2^{(1/2)}*\cot(d*x+c)^{(1/2)}))+2*\arctan(1+2^{(1/2)}*\cot(d*x+c)^{(1/2)}))+2*\arctan(-1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})) \end{aligned}$$

3.588.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6167 vs. $2(379) = 758$.

Time = 6.20 (sec) , antiderivative size = 6167, normalized size of antiderivative = 14.65

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="fricas")`

output Too large to include

3.588.6 Sympy [F]

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\ & = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^3}{\sqrt{\cot(c + dx)}} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))**3*(A+B*tan(d*x+c))/cot(d*x+c)**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**3/sqrt(cot(c + d*x)), x)`

3.588.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 370, normalized size of antiderivative = 0.88

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \frac{8 \left(15 B b^3 + \frac{21 (3 B a b^2 + A b^3)}{\tan(dx+c)} + \frac{35 (3 B a^2 b + 3 A a b^2 - B b^3)}{\tan(dx+c)^2} + \frac{105 (B a^3 + 3 A a^2 b - 3 B a b^2 - A b^3)}{\tan(dx+c)^3} \right) \tan(dx+c)^{\frac{7}{2}} - 210 \sqrt{2} ((A + B) a^2 b - 3 (A - B) a b^2 + (A + B) b^3) \arctan\left(\frac{1}{2} \sqrt{2} (\sqrt{2} + 2/\sqrt{\tan(dx+c)})}\right) - 210 \sqrt{2} ((A - B) a^2 b - 3 (A + B) a b^2 + (A + B) b^3) \arctan\left(-\frac{1}{2} \sqrt{2} (\sqrt{2} - 2/\sqrt{\tan(dx+c)})}\right) - 105 \sqrt{2} ((A + B) a^3 + 3 (A - B) a^2 b - 3 (A + B) a b^2 - (A - B) b^3) \log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + \frac{1}{\tan(dx+c)} + 1\right) + 105 \sqrt{2} ((A + B) a^3 + 3 (A - B) a^2 b - 3 (A + B) a b^2 - (A - B) b^3) \log\left(-\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + \frac{1}{\tan(dx+c)} + 1\right)}{d}$$

```
input integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="maxima")
```

```
output 1/420*(8*(15*B*b^3 + 21*(3*B*a*b^2 + A*b^3)/tan(d*x + c) + 35*(3*B*a^2*b + 3*A*a*b^2 - B*b^3)/tan(d*x + c)^2 + 105*(B*a^3 + 3*A*a^2*b - 3*B*a*b^2 - A*b^3)/tan(d*x + c)^3)*tan(d*x + c)^(7/2) - 210*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) - 210*sqrt(2)*((A - B)*a^3 - 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 + (A + B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - 105*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + 105*sqrt(2)*((A + B)*a^3 + 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 - (A - B)*b^3)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))/d
```

3.588.8 Giac [F]

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^3}{\sqrt{\cot(dx + c)}} dx$$

```
input integrate((a+b*tan(d*x+c))^3*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="giac")
```

```
output integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^3/sqrt(cot(d*x + c)), x)
```

3.588.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^3 (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^3}{\sqrt{\cot(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3)/cot(c + d*x)^(1/2),x)`output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^3)/cot(c + d*x)^(1/2), x)`

3.589
$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.589.1 Optimal result	5582
3.589.2 Mathematica [A] (verified)	5583
3.589.3 Rubi [A] (warning: unable to verify)	5583
3.589.4 Maple [B] (verified)	5591
3.589.5 Fricas [B] (verification not implemented)	5592
3.589.6 Sympy [F(-1)]	5593
3.589.7 Maxima [A] (verification not implemented)	5593
3.589.8 Giac [F]	5594
3.589.9 Mupad [F(-1)]	5594

3.589.1 Optimal result

Integrand size = 33, antiderivative size = 325

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$= \frac{(b(A-B) - a(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2 + b^2) d}$$

$$- \frac{(b(A-B) - a(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2 + b^2) d}$$

$$- \frac{2b^{5/2}(Ab - aB) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{a^{5/2}(a^2 + b^2) d} + \frac{2(Ab - aB)\sqrt{\cot(c+dx)}}{a^2 d}$$

$$- \frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad} + \frac{(a(A-B) + b(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2 + b^2) d}$$

$$- \frac{(a(A-B) + b(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2 + b^2) d}$$

output
$$-2b^{5/2}(Ab-Ba)\arctan(a^{1/2}\cot(dx+c)^{1/2}/b^{1/2})/a^{5/2}/(a^2+b^2)/d-2/3A\cot(dx+c)^{3/2}/a/d-1/2*(b(A-B)-a(A+B))\arctan(-1+2^{1/2}\cot(dx+c)^{1/2})/(a^2+b^2)/d*2^{1/2}-1/2*(b(A-B)-a(A+B))\arctan(1+2^{1/2}\cot(dx+c)^{1/2})/(a^2+b^2)/d*2^{1/2}+1/4*(a(A-B)+b(A+B))*\ln(1+\cot(dx+c)-2^{1/2}\cot(dx+c)^{1/2})/(a^2+b^2)/d*2^{1/2}-1/4*(a(A-B)+b(A+B))*\ln(1+\cot(dx+c)+2^{1/2}\cot(dx+c)^{1/2})/(a^2+b^2)/d*2^{1/2}+2*(Ab-Ba)\cot(dx+c)^{1/2}/a^2/d$$

3.589.2 Mathematica [A] (verified)

Time = 1.76 (sec) , antiderivative size = 272, normalized size of antiderivative = 0.84

$$\int \frac{\cot^{5/2}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{\sqrt{\cot(c+dx)} \left(-\frac{6\sqrt{2}(b(-A+B)+a(A+B))(\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})-\arctan(1+\sqrt{2}\sqrt{\tan(c+dx)})}{a^2+b^2} + \frac{24b^{5/2}(-Ab+aB)\arctan(1+\sqrt{2}\sqrt{\tan(c+dx)})}{a^{5/2}(a^2+b^2)} \right)}{a^2+b^2}$$

input `Integrate[(Cot[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output
$$-1/12*(\text{Sqrt}[\text{Cot}[c + d*x]]*((-6*\text{Sqrt}[2]*(b*(-A + B) + a*(A + B))*(\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]) - \text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]])/(a^2 + b^2) + (24*b^{5/2}*(-(A*b) + a*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(a^{5/2}*(a^2 + b^2)) - (3*\text{Sqrt}[2]*(a*(A - B) + b*(A + B))*(\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]] - \text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]])))/(a^2 + b^2) + (8*A)/(a*\text{Tan}[c + d*x]^{3/2}) + (24*(-(A*b) + a*B))/(a^2*\text{Sqrt}[\text{Tan}[c + d*x]]))*\text{Sqrt}[\text{Tan}[c + d*x]]/d$$

3.589.3 Rubi [A] (warning: unable to verify)

Time = 1.73 (sec) , antiderivative size = 286, normalized size of antiderivative = 0.88, number of steps used = 26, number of rules used = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.758$, Rules used = {3042, 4064, 3042, 4090, 27, 3042, 4130, 27, 3042, 4136, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.589.
$$\int \frac{\cot^{5/2}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$\begin{aligned}
& \int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx \\
& \quad \downarrow 3042 \\
& \int \frac{\cot(c+dx)^{5/2}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx \\
& \quad \downarrow 4064 \\
& \int \frac{\cot^{\frac{5}{2}}(c+dx)(A \cot(c+dx)+B)}{a \cot(c+dx)+b} dx \\
& \quad \downarrow 3042 \\
& \int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{5/2}(B-A \tan(c+dx+\frac{\pi}{2}))}{b-a \tan(c+dx+\frac{\pi}{2})} dx \\
& \quad \downarrow 4090 \\
& \frac{2 \int \frac{3\sqrt{\cot(c+dx)}((Ab-aB) \cot^2(c+dx)+aA \cot(c+dx)+Ab)}{2(b+a \cot(c+dx))} dx}{3a} - \frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{\sqrt{\cot(c+dx)}((Ab-aB) \cot^2(c+dx)+aA \cot(c+dx)+Ab)}{b+a \cot(c+dx)} dx}{a} - \frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{\sqrt{-\tan(c+dx+\frac{\pi}{2})}((Ab-aB) \tan(c+dx+\frac{\pi}{2})^2-aA \tan(c+dx+\frac{\pi}{2})+Ab)}{b-a \tan(c+dx+\frac{\pi}{2})} dx}{a} - \frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad} \\
& \quad \downarrow 4130 \\
& \frac{2 \int \frac{-B \cot(c+dx)a^2-(Aa^2+bBa-Ab^2) \cot^2(c+dx)+b(Ab-aB)}{2\sqrt{\cot(c+dx)}(b+a \cot(c+dx))} dx}{a} - \frac{2(Ab-aB)\sqrt{\cot(c+dx)}}{ad} - \frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{-B \cot(c+dx)a^2-(Aa^2+bBa-Ab^2) \cot^2(c+dx)+b(Ab-aB)}{\sqrt{\cot(c+dx)}(b+a \cot(c+dx))} dx}{a} - \frac{2(Ab-aB)\sqrt{\cot(c+dx)}}{ad} - \frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{B \tan(c+dx+\frac{\pi}{2})a^2+(-Aa^2-bBa+Ab^2) \tan(c+dx+\frac{\pi}{2})^2+b(Ab-aB)}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx}{a} - \frac{2(Ab-aB)\sqrt{\cot(c+dx)}}{ad} - \frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad}
\end{aligned}$$

3.589. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\int \frac{a^2(Ab-aB)-a^2(aA+bB)\cot(c+dx)}{\sqrt{\cot(c+dx)}(a^2+b^2)} dx + \frac{b^3(Ab-aB) \int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)}(b+a \cot(c+dx))} dx}{a^2+b^2}$$

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad} - \frac{2(Ab-aB)\sqrt{\cot(c+dx)}}{ad}$$

4136

$$\int \frac{(Ab-aB)a^2+(aA+bB)\tan(c+dx+\frac{\pi}{2})a^2}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(a^2+b^2)} dx + \frac{b^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx}{a^2+b^2}$$

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad} - \frac{2(Ab-aB)\sqrt{\cot(c+dx)}}{ad}$$

3042

$$2 \int \frac{a^2(Ab-aB-(aA+bB)\cot(c+dx)) d\sqrt{\cot(c+dx)}}{\cot^2(c+dx)+1} + \frac{b^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx}{a^2+b^2}$$

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad} - \frac{2(Ab-aB)\sqrt{\cot(c+dx)}}{ad}$$

4017

$$b^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx - 2 \int \frac{a^2(Ab-aB-(aA+bB)\cot(c+dx)) d\sqrt{\cot(c+dx)}}{\cot^2(c+dx)+1}$$

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad} - \frac{2(Ab-aB)\sqrt{\cot(c+dx)}}{ad}$$

25

$$b^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx - 2a^2 \int \frac{Ab-aB-(aA+bB)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}$$

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad} - \frac{2(Ab-aB)\sqrt{\cot(c+dx)}}{ad}$$

27

$$\int \frac{a^2(Ab-aB-(aA+bB)\cot(c+dx)) d\sqrt{\cot(c+dx)}}{\cot^2(c+dx)+1} - \frac{b^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx}{a^2+b^2}$$

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad} - \frac{2(Ab-aB)\sqrt{\cot(c+dx)}}{ad}$$

1482

3.589. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\frac{b^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2a^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(b(A-B)-a(A+B)) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)} dx \right)}{d(a^2+b^2)}$$

a

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad}$$

↓ 1476

$$\frac{b^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2a^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)} dx \right) \right)}{d(a^2+b^2)}$$

a

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad}$$

↓ 1082

$$\frac{b^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2a^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(b(A-B)-a(A+B)) \left(\int \frac{1}{-\cot(c+dx)} dx \right) \right)}{d(a^2+b^2)}$$

a

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad}$$

↓ 217

$$\frac{b^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2a^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)}$$

a

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad}$$

↓ 1479

$$\frac{b^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2a^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)})}{\cot(c+dx)}}{\cot(c+dx)} \right) \right)}{d(a^2+b^2)}$$

a

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad}$$

3.589. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

↓ 25

$$\frac{b^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2a^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)})}{\cot(c+dx)+\sqrt{2}} dx}{2\sqrt{2}} \right) \right)}{a}$$

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad}$$

↓ 27

$$\frac{b^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2a^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)}}{\cot(c+dx)+\sqrt{2}} dx \right) \right)}{a}$$

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad}$$

↓ 1103

$$\frac{b^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2a^2 \left(\frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(a(A+B)+b(A-B)) \right)}{a}$$

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad}$$

↓ 4117

$$\frac{b^3(Ab-aB) \int \frac{1}{\sqrt{\cot(c+dx)(b+a\cot(c+dx))}} d(-\cot(c+dx))}{d(a^2+b^2)} - \frac{2a^2 \left(\frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(a(A+B)+b(A-B)) \right)}{d(a^2+b^2)}$$

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad}$$

↓ 73

$$\frac{2b^3(Ab-aB) \int \frac{1}{a \cot^2(c+dx)+b} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} - \frac{2a^2 \left(\frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(a(A-B)+b(A+B)) \right)}{d(a^2+b^2)}$$

$$\frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad}$$

3.589. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

↓ 218

$$\frac{2b^{5/2}(Ab-aB) \arctan\left(\frac{\sqrt{a} \cot(c+dx)}{\sqrt{b}}\right) - 2a^2 \left(\frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}}\right) + \frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}}\right) + \frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}}\right)\right)}{\sqrt{ad}(a^2+b^2)} - \frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad}$$

input `Int[(Cot[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(-2*A*Cot[c + d*x]^(3/2))/(3*a*d) - ((-2*(A*b - a*B)*Sqrt[Cot[c + d*x]])/(a*d) - ((2*b^(5/2)*(A*b - a*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]])/(Sqrt[a]*(a^2 + b^2)*d) - (2*a^2*((b*(A - B) - a*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]] + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]))/2 + ((a*(A - B) + b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)/a/a`

3.589.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

- rule 218 $\text{Int}[(a_.) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$
- rule 1082 $\text{Int}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_.) + (e_.)*(x_)] / ((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$
- rule 1476 $\text{Int}[(d_.) + (e_.)*(x_)^2] / ((a_.) + (c_.)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Simp}[e/(2*c) \ \text{Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Simp}[e/(2*c) \ \text{Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{PosQ}[d*e]$
- rule 1479 $\text{Int}[(d_.) + (e_.)*(x_)^2] / ((a_.) + (c_.)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2*(d/e), 2]\}, \text{Simp}[e/(2*c*q) \ \text{Int}[(q - 2*x)/\text{Simp}[d/e + q*x - x^2, x], x], x] + \text{Simp}[e/(2*c*q) \ \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{NegQ}[d*e]$
- rule 1482 $\text{Int}[(d_.) + (e_.)*(x_)^2] / ((a_.) + (c_.)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a*c, 2]\}, \text{Simp}[(d*q + a*e)/(2*a*c) \ \text{Int}[(q + c*x^2)/(a + c*x^4), x], x] + \text{Simp}[(d*q - a*e)/(2*a*c) \ \text{Int}[(q - c*x^2)/(a + c*x^4), x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{NeQ}[c*d^2 + a*e^2, 0] \ \&\& \ \text{NeQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{NegQ}[(a)*c]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4017 $\text{Int}[(c_.) + (d_.)*\text{tan}[(e_.) + (f_.)*(x_)]) / \text{Sqrt}[(b_.)*\text{tan}[(e_.) + (f_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[2/f \ \text{Subst}[\text{Int}[(b*c + d*x^2)/(b^2 + x^4), x], x, \text{Sqrt}[b*\text{Tan}[e + f*x]], x] /; \text{FreeQ}[\{b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[c^2 - d^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0]$

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp [g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c *Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !Integer Q[p] && IntegerQ[m] && IntegerQ[n]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b *(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Sim p[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C *(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C* m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.589.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 771 vs. 2(281) = 562.

Time = 0.50 (sec) , antiderivative size = 772, normalized size of antiderivative = 2.38

method	result
derivativedivides	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \left(3A \tan(dx+c)^{\frac{3}{2}} \sqrt{2} \sqrt{ab} \ln\left(-\frac{1+\sqrt{2} \sqrt{\tan(dx+c)+\tan(dx+c)}}{\sqrt{2} \sqrt{\tan(dx+c)-\tan(dx+c)-1}}\right) a^3 + 6A \tan(dx+c)^{\frac{3}{2}} \sqrt{2} \sqrt{ab} a\right)}{\dots}$
default	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{5}{2}} \tan(dx+c) \left(3A \tan(dx+c)^{\frac{3}{2}} \sqrt{2} \sqrt{ab} \ln\left(-\frac{1+\sqrt{2} \sqrt{\tan(dx+c)+\tan(dx+c)}}{\sqrt{2} \sqrt{\tan(dx+c)-\tan(dx+c)-1}}\right) a^3 + 6A \tan(dx+c)^{\frac{3}{2}} \sqrt{2} \sqrt{ab} a\right)}{\dots}$

```
input int(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

$$3.589. \int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

output

```
-1/12/d*(1/tan(d*x+c))^(5/2)*tan(d*x+c)*(3*A*tan(d*x+c)^(3/2)*2^(1/2)*(a*b)^(1/2)*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1))*a^3+6*A*tan(d*x+c)^(3/2)*2^(1/2)*(a*b)^(1/2)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*a^3-6*A*tan(d*x+c)^(3/2)*2^(1/2)*(a*b)^(1/2)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*a^2*b+6*A*tan(d*x+c)^(3/2)*2^(1/2)*(a*b)^(1/2)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*a^3-6*A*tan(d*x+c)^(3/2)*2^(1/2)*(a*b)^(1/2)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*a^2*b-3*A*tan(d*x+c)^(3/2)*2^(1/2)*(a*b)^(1/2)*ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))*a^2*b+3*B*tan(d*x+c)^(3/2)*2^(1/2)*(a*b)^(1/2)*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1))*a^2*b+6*B*tan(d*x+c)^(3/2)*2^(1/2)*(a*b)^(1/2)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*a^3+6*B*tan(d*x+c)^(3/2)*2^(1/2)*(a*b)^(1/2)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*a^2*b+6*B*tan(d*x+c)^(3/2)*2^(1/2)*(a*b)^(1/2)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*a^3+6*B*tan(d*x+c)^(3/2)*2^(1/2)*(a*b)^(1/2)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*a^2*b+3*B*tan(d*x+c)^(3/2)*2^(1/2)*(a*b)^(1/2)*ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))*a^3-24*A*tan(d*x+c)^(3/2)*arctan(b*tan(d*x+c)^(1/2))/(a*b)^(1/2)*b^4+24*B*tan(d*x+c)^(3/2)*arctan(b*tan(d*x+c)^(1/2))/(a*b)^(1/2)*a*b^3-24*A*tan(d*x+c)*(a*b)^(1/2)*a^2*b-24*A*tan(d*x+c)*(a*b)^(1/2)*b^3+24*B*tan(d*x+c)*(a*b)^(1/2)*a^3+24*B*tan(d*x+c)*(a*b)^(1/2)*a*...
```

3.589.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3115 vs. $2(281) = 562$.

Time = 26.27 (sec) , antiderivative size = 6260, normalized size of antiderivative = 19.26

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.589. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$

3.589.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Timed out`

3.589.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 262, normalized size of antiderivative = 0.81

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{24(Bab^3 - Ab^4) \arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right)}{(a^4 + a^2b^2)\sqrt{ab}} + \frac{3\left(2\sqrt{2}((A+B)a - (A-B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}((A+B)a - (A-B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right)\right)}{(a^4 + a^2b^2)\sqrt{ab}}$$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/12*(24*(B*a*b^3 - A*b^4)*arctan(a/(sqrt(a*b)*sqrt(tan(d*x + c))))/((a^4 + a^2*b^2)*sqrt(a*b)) + 3*(2*sqrt(2)*((A + B)*a - (A - B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A + B)*a - (A - B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*((A - B)*a + (A + B)*b)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*((A - B)*a + (A + B)*b)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))/(a^2 + b^2) - 8*(A*a/tan(d*x + c)^(3/2) + 3*(B*a - A*b)/sqrt(tan(d*x + c)))/a^2)/d`

3.589.8 Giac [F]

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{5}{2}}}{b\tan(dx+c)+a} dx$$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(5/2)/(b*tan(d*x + c) + a), x)`

3.589.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \int \frac{\cot(c+dx)^{5/2}(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

input `int((cot(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `int((cot(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)), x)`

3.590
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.590.1 Optimal result 5595
 3.590.2 Mathematica [A] (verified) 5596
 3.590.3 Rubi [A] (warning: unable to verify) 5596
 3.590.4 Maple [B] (verified) 5603
 3.590.5 Fricas [B] (verification not implemented) 5604
 3.590.6 Sympy [F] 5605
 3.590.7 Maxima [A] (verification not implemented) 5605
 3.590.8 Giac [F] 5606
 3.590.9 Mupad [F(-1)] 5606

3.590.1 Optimal result

Integrand size = 33, antiderivative size = 297

$$\begin{aligned} & \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx \\ &= -\frac{(a(A-B)+b(A+B)) \arctan\left(1-\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d} \\ & \quad + \frac{(a(A-B)+b(A+B)) \arctan\left(1+\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d} \\ & \quad + \frac{2b^{3/2}(Ab-aB) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{a^{3/2}(a^2+b^2)d} - \frac{2A\sqrt{\cot(c+dx)}}{ad} \\ & \quad + \frac{(b(A-B)-a(A+B)) \log\left(1-\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d} \\ & \quad - \frac{(b(A-B)-a(A+B)) \log\left(1+\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d} \end{aligned}$$

output

```
2*b^(3/2)*(A*b-B*a)*arctan(a^(1/2)*cot(d*x+c)^(1/2)/b^(1/2))/a^(3/2)/(a^2+b^2)/d+1/2*(a*(A-B)+b*(A+B))*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)+1/2*(a*(A-B)+b*(A+B))*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)+1/4*(b*(A-B)-a*(A+B))*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/4*(b*(A-B)-a*(A+B))*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-2*A*cot(d*x+c)^(1/2)/a/d
```

3.590.
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.590.2 Mathematica [A] (verified)

Time = 0.95 (sec) , antiderivative size = 249, normalized size of antiderivative = 0.84

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$= \frac{\sqrt{\cot(c+dx)} \left(\frac{2\sqrt{2}(a(A-B)+b(A+B))(\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})-\arctan(1+\sqrt{2}\sqrt{\tan(c+dx)})}{a^2+b^2} \right) + \frac{8b^{3/2}(-Ab+aB) \arctan\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{a}\right)}{a^{3/2}(a^2+b^2)}}{1}$$

input `Integrate[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*((2*Sqrt[2]*(a*(A - B) + b*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]))/(a^2 + b^2) + (8*b^(3/2)*(-(A*b) + a*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/(a^(3/2)*(a^2 + b^2)) - (Sqrt[2]*(b*(-A + B) + a*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]))/(a^2 + b^2) - (8*A)/(a*Sqrt[Tan[c + d*x]]))*Sqrt[Tan[c + d*x]]/(4*d)`

3.590.3 Rubi [A] (warning: unable to verify)Time = 1.41 (sec) , antiderivative size = 250, normalized size of antiderivative = 0.84, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4064, 3042, 4090, 27, 3042, 4136, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\cot(c+dx)^{3/2}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$\downarrow \text{4064}$$

3.590. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\begin{aligned}
& \int \frac{\cot^{\frac{3}{2}}(c+dx)(A \cot(c+dx) + B)}{a \cot(c+dx) + b} dx \\
& \quad \downarrow 3042 \\
& \int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{3/2}(B - A \tan(c+dx+\frac{\pi}{2}))}{b - a \tan(c+dx+\frac{\pi}{2})} dx \\
& \quad \downarrow 4090 \\
& - \frac{2 \int \frac{(Ab-aB) \cot^2(c+dx) + aA \cot(c+dx) + Ab}{2\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} dx}{a} - \frac{2A\sqrt{\cot(c+dx)}}{ad} \\
& \quad \downarrow 27 \\
& - \frac{\int \frac{(Ab-aB) \cot^2(c+dx) + aA \cot(c+dx) + Ab}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} dx}{a} - \frac{2A\sqrt{\cot(c+dx)}}{ad} \\
& \quad \downarrow 3042 \\
& - \frac{\int \frac{(Ab-aB) \tan(c+dx+\frac{\pi}{2})^2 - aA \tan(c+dx+\frac{\pi}{2}) + Ab}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a} - \frac{2A\sqrt{\cot(c+dx)}}{ad} \\
& \quad \downarrow 4136 \\
& - \frac{\frac{b^2(Ab-aB) \int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} dx}{a^2+b^2} + \frac{\int \frac{a(aA+bB)+a(Ab-aB) \cot(c+dx)}{\sqrt{\cot(c+dx)}} dx}{a^2+b^2}}{a} - \frac{2A\sqrt{\cot(c+dx)}}{ad} \\
& \quad \downarrow 3042 \\
& - \frac{b^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} + \frac{\int \frac{a(aA+bB)-a(Ab-aB) \tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{a^2+b^2}}{a} - \frac{2A\sqrt{\cot(c+dx)}}{ad} \\
& \quad \downarrow 4017 \\
& - \frac{b^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} + \frac{2 \int -\frac{a(aA+bB+(Ab-aB) \cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)}}{a} - \frac{2A\sqrt{\cot(c+dx)}}{ad} \\
& \quad \downarrow 25
\end{aligned}$$

3.590. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\frac{b^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2 \int \frac{a(aA+bB+(Ab-aB) \cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)}$$

$$\frac{2A\sqrt{\cot(c+dx)}}{ad}$$

↓ 27

$$\frac{b^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2a \int \frac{aA+bB+(Ab-aB) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)}$$

$$\frac{2A\sqrt{\cot(c+dx)}}{ad}$$

↓ 1482

$$\frac{b^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2a\left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(b(A-B)-a(A+B)) \int \frac{1}{\cot(c+dx)-1} d\sqrt{\cot(c+dx)}\right)}{d(a^2+b^2)}$$

$$\frac{2A\sqrt{\cot(c+dx)}}{ad}$$

↓ 1476

$$\frac{b^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2a\left(\frac{1}{2}(a(A-B)+b(A+B))\left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}\right)\right)}{d(a^2+b^2)}$$

$$\frac{2A\sqrt{\cot(c+dx)}}{ad}$$

↓ 1082

$$\frac{b^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2a\left(\frac{1}{2}(a(A-B)+b(A+B))\left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)-1} d(1+\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}}\right)\right)}{d(a^2+b^2)}$$

$$\frac{2A\sqrt{\cot(c+dx)}}{ad}$$

↓ 217

3.590. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\frac{b^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} = \frac{2a \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}) - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2}}{d(a^2+b^2)}$$

$$\frac{2A\sqrt{\cot(c+dx)}}{ad}$$

↓ 1479

$$\frac{b^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} = \frac{2a \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}) - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2}}{a}$$

$$\frac{2A\sqrt{\cot(c+dx)}}{ad}$$

↓ 25

$$\frac{b^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} = \frac{2a \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}) - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2}}{a}$$

$$\frac{2A\sqrt{\cot(c+dx)}}{ad}$$

↓ 27

$$\frac{b^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} = \frac{2a \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}) - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2}}{a}$$

$$\frac{2A\sqrt{\cot(c+dx)}}{ad}$$

↓ 1103

$$\frac{b^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} = \frac{2a \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}) - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2}}{a}$$

$$\frac{2A\sqrt{\cot(c+dx)}}{ad}$$

↓ 4117

3.590. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\frac{b^2(Ab-aB) \int \frac{1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} d(-\cot(c+dx))}{d(a^2+b^2)} - \frac{2a \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{a}$$

$$\frac{2A\sqrt{\cot(c+dx)}}{ad} \quad \downarrow \quad 73$$

$$\frac{2b^2(Ab-aB) \int \frac{1}{a \cot^2(c+dx)+b} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} - \frac{2a \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2}(b(A-B)-a(A+B))}{d(a^2+b^2)}$$

$$\frac{2A\sqrt{\cot(c+dx)}}{ad} \quad \downarrow \quad 218$$

$$\frac{2b^{3/2}(Ab-aB) \arctan\left(\frac{\sqrt{a} \cot(c+dx)}{\sqrt{b}}\right)}{\sqrt{ad}(a^2+b^2)} - \frac{2a \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2}(b(A-B)-a(A+B))}{d(a^2+b^2)}$$

$$\frac{2A\sqrt{\cot(c+dx)}}{ad}$$

```
input Int[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]
```

```
output (-2*A*Sqrt[Cot[c + d*x]]/(a*d) - ((2*b^(3/2)*(A*b - a*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]]/(Sqrt[a]*(a^2 + b^2)*d) - (2*a*(((a*(A - B) + b*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]))/2 - ((b*(A - B) - a*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2))/((a^2 + b^2)*d))/a
```

3.590.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

3.590. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
 -1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
 & (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
 t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
 implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
 eQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
 imp[d*(Log[RemoveContent[a + b*x + c*x^2, x])/b], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
 2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
 e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
 && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
 -2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
 x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
 reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4090 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.590.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 685 vs. 2(257) = 514.

Time = 0.38 (sec) , antiderivative size = 686, normalized size of antiderivative = 2.31

method	result
derivativedivides	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{3}{2}} \tan(dx+c) \left(A\sqrt{ab} \sqrt{2} \ln\left(-\frac{1+\sqrt{2} \sqrt{\tan(dx+c)+\tan(dx+c)}}{\sqrt{2} \sqrt{\tan(dx+c)-\tan(dx+c)-1}}\right) \sqrt{\tan(dx+c)} ab+2A\sqrt{ab} \sqrt{2} \arctan\left(1+\sqrt{2}\right)}{\dots}$
default	$-\frac{\left(\frac{1}{\tan(dx+c)}\right)^{\frac{3}{2}} \tan(dx+c) \left(A\sqrt{ab} \sqrt{2} \ln\left(-\frac{1+\sqrt{2} \sqrt{\tan(dx+c)+\tan(dx+c)}}{\sqrt{2} \sqrt{\tan(dx+c)-\tan(dx+c)-1}}\right) \sqrt{\tan(dx+c)} ab+2A\sqrt{ab} \sqrt{2} \arctan\left(1+\sqrt{2}\right)}{\dots}$

```
input int(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

$$3.590. \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

output

```

-1/4/d*(1/tan(d*x+c))^(3/2)*tan(d*x+c)*(A*(a*b)^(1/2)*2^(1/2)*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1))*tan(d*x+c)^(1/2)*a*b+2*A*(a*b)^(1/2)*2^(1/2)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*tan(d*x+c)^(1/2)*a^2+2*A*(a*b)^(1/2)*2^(1/2)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*tan(d*x+c)^(1/2)*a*b+2*A*(a*b)^(1/2)*2^(1/2)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*tan(d*x+c)^(1/2)*a^2+2*A*(a*b)^(1/2)*2^(1/2)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*tan(d*x+c)^(1/2)*a*b+A*(a*b)^(1/2)*2^(1/2)*ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))) *tan(d*x+c)^(1/2)*a^2-B*(a*b)^(1/2)*2^(1/2)*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1))*tan(d*x+c)^(1/2)*a^2-2*B*(a*b)^(1/2)*2^(1/2)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*tan(d*x+c)^(1/2)*a^2+2*B*(a*b)^(1/2)*2^(1/2)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*tan(d*x+c)^(1/2)*a*b-2*B*(a*b)^(1/2)*2^(1/2)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*tan(d*x+c)^(1/2)*a^2+2*B*(a*b)^(1/2)*2^(1/2)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*tan(d*x+c)^(1/2)*a*b+B*(a*b)^(1/2)*2^(1/2)*ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))) *tan(d*x+c)^(1/2)*a*b+8*A*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))*tan(d*x+c)^(1/2)*b^3-8*B*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))*tan(d*x+c)^(1/2)*a*b^2+8*A*(a*b)^(1/2)*a^2+8*A*(a*b)^(1/2)*b^2)/(a^2+b^2)/a/(a*b)^(1/2)

```

3.590.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3013 vs. $2(257) = 514$.

Time = 11.53 (sec) , antiderivative size = 6056, normalized size of antiderivative = 20.39

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.590. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$

3.590.6 Sympy [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \int \frac{(A+B\tan(c+dx))\cot^{\frac{3}{2}}(c+dx)}{a+b\tan(c+dx)} dx$$

input `integrate(cot(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**(3/2)/(a + b*tan(c + d*x)), x)`

3.590.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 239, normalized size of antiderivative = 0.80

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{8(Bab^2 - Ab^3) \arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right) - 2\sqrt{2}((A-B)a + (A+B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}((A-B)a + (A+B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right)}{(a^3 + ab^2)\sqrt{ab}}$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `-1/4*(8*(B*a*b^2 - A*b^3)*arctan(a/(sqrt(a*b)*sqrt(tan(d*x + c))))/((a^3 + a*b^2)*sqrt(a*b)) - (2*sqrt(2)*((A - B)*a + (A + B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a + (A + B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)*((A + B)*a - (A - B)*b)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*((A + B)*a - (A - B)*b)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))/(a^2 + b^2) + 8*A/(a*sqrt(tan(d*x + c))))/d`

3.590.8 Giac [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{b\tan(dx+c)+a} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(3/2)/(b*tan(d*x + c) + a), x)`

3.590.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \int \frac{\cot(c+dx)^{3/2}(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

input `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)), x)`

3.591
$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.591.1 Optimal result 5607
 3.591.2 Mathematica [A] (verified) 5608
 3.591.3 Rubi [A] (warning: unable to verify) 5608
 3.591.4 Maple [B] (verified) 5614
 3.591.5 Fricas [B] (verification not implemented) 5615
 3.591.6 Sympy [F] 5615
 3.591.7 Maxima [A] (verification not implemented) 5615
 3.591.8 Giac [F] 5616
 3.591.9 Mupad [F(-1)] 5616

3.591.1 Optimal result

Integrand size = 33, antiderivative size = 278

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$= -\frac{(b(A-B) - a(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2 + b^2)d}$$

$$+ \frac{(b(A-B) - a(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2 + b^2)d}$$

$$- \frac{2\sqrt{b}(Ab - aB) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{\sqrt{a}(a^2 + b^2)d}$$

$$- \frac{(a(A-B) + b(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2 + b^2)d}$$

$$+ \frac{(a(A-B) + b(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2 + b^2)d}$$

output

```
1/2*(b*(A-B)-a*(A+B))*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)+1/2*(b*(A-B)-a*(A+B))*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/4*(a*(A-B)+b*(A+B))*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)+1/4*(a*(A-B)+b*(A+B))*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-2*(A*b-B*a)*arctan(a^(1/2)*cot(d*x+c)^(1/2)/b^(1/2))*b^(1/2)/(a^2+b^2)/d/a^(1/2)
```

3.591.2 Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.77

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$= \frac{\sqrt{\cot(c+dx)} \left(-2\sqrt{2}(b(-A+B) + a(A+B)) \left(\arctan \left(1 - \sqrt{2}\sqrt{\tan(c+dx)} \right) - \arctan \left(1 + \sqrt{2}\sqrt{\tan(c+dx)} \right) \right) \right)}{4(a^2 + b^2)d}$$

input `Integrate[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*(-2*Sqrt[2]*(b*(-A + B) + a*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]])] + (8*Sqrt[b]*(A*b - a*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/Sqrt[a] - Sqrt[2]*(a*(A - B) + b*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]))*Sqrt[Tan[c + d*x]]/(4*(a^2 + b^2)*d)`

3.591.3 Rubi [A] (warning: unable to verify)Time = 1.06 (sec) , antiderivative size = 224, normalized size of antiderivative = 0.81, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.576$, Rules used = {3042, 4064, 3042, 4095, 25, 3042, 4017, 25, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{\sqrt{\cot(c+dx)}(A \cot(c+dx) + B)}{a \cot(c+dx) + b} dx$$

3.591. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\begin{aligned}
& \int \frac{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(B-A \tan\left(c+dx+\frac{\pi}{2}\right)\right)}{b-a \tan\left(c+dx+\frac{\pi}{2}\right)} dx \\
& \quad \downarrow \text{3042} \\
& \frac{b(Ab-aB) \int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} dx}{a^2+b^2} + \frac{\int -\frac{Ab-aB-(aA+bB) \cot(c+dx)}{\sqrt{\cot(c+dx)}} dx}{a^2+b^2} \\
& \quad \downarrow \text{4095} \\
& \frac{b(Ab-aB) \int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} dx}{a^2+b^2} - \frac{\int \frac{Ab-aB-(aA+bB) \cot(c+dx)}{\sqrt{\cot(c+dx)}} dx}{a^2+b^2} \\
& \quad \downarrow \text{25} \\
& \frac{b(Ab-aB) \int \frac{\tan\left(c+dx+\frac{\pi}{2}\right)^2+1}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)\left(b-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)}} dx}{a^2+b^2} - \frac{\int \frac{Ab-aB-(-aA-bB) \tan\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}} dx}{a^2+b^2} \\
& \quad \downarrow \text{3042} \\
& \frac{b(Ab-aB) \int \frac{\tan\left(c+dx+\frac{\pi}{2}\right)^2+1}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)\left(b-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)}} dx}{a^2+b^2} - \frac{2 \int -\frac{Ab-aB-(aA+bB) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d\left(a^2+b^2\right)} \\
& \quad \downarrow \text{4017} \\
& \frac{b(Ab-aB) \int \frac{\tan\left(c+dx+\frac{\pi}{2}\right)^2+1}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)\left(b-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)}} dx}{a^2+b^2} + \frac{2 \int \frac{Ab-aB-(aA+bB) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d\left(a^2+b^2\right)} \\
& \quad \downarrow \text{25} \\
& \frac{b(Ab-aB) \int \frac{\tan\left(c+dx+\frac{\pi}{2}\right)^2+1}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)\left(b-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)}} dx}{a^2+b^2} - \frac{2\left(-\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(b(A-B)-a(A+B)) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{d\left(a^2+b^2\right)} \\
& \quad \downarrow \text{1482} \\
& \frac{b(Ab-aB) \int \frac{\tan\left(c+dx+\frac{\pi}{2}\right)^2+1}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)\left(b-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)}} dx}{a^2+b^2} - \frac{2\left(-\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)}}\right)\right)}{d\left(a^2+b^2\right)} \\
& \quad \downarrow \text{1476} \\
& \frac{b(Ab-aB) \int \frac{\tan\left(c+dx+\frac{\pi}{2}\right)^2+1}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)\left(b-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)}} dx}{a^2+b^2} - \frac{2\left(-\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)}}\right)\right)}{d\left(a^2+b^2\right)}
\end{aligned}$$

3.591. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\begin{aligned} & \downarrow 1082 \\ & \frac{b(Ab - aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2 + b^2} - \\ & \frac{2 \left(-\frac{1}{2}(a(A-B) + b(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(b(A-B) - a(A+B)) \left(\int \frac{1}{-\cot(c+dx)-1} d\frac{(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 217 \\ & \frac{b(Ab - aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2 + b^2} - \\ & \frac{2 \left(-\frac{1}{2}(a(A-B) + b(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(b(A-B) - a(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 1479 \\ & \frac{b(Ab - aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2 + b^2} - \\ & \frac{2 \left(-\frac{1}{2}(a(A-B) + b(A+B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) \right)}{d(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 25 \\ & \frac{b(Ab - aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2 + b^2} - \\ & \frac{2 \left(-\frac{1}{2}(a(A-B) + b(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) \right) - \frac{1}{2}(b(A+B))}{d(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{b(Ab - aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2 + b^2} - \\ & \frac{2 \left(-\frac{1}{2}(a(A-B) + b(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{d(a^2 + b^2)} \end{aligned}$$

$$\downarrow 1103$$

3.591. $\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$

$$\frac{b(Ab - aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2 + b^2} -$$

$$\frac{2 \left(-\frac{1}{2}(b(A - B) - a(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} \right) - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a(A - B) + b(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} \right) - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right)}{d(a^2 + b^2)} -$$

↓ 4117

$$\frac{b(Ab - aB) \int \frac{1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} d(-\cot(c + dx))}{d(a^2 + b^2)} -$$

$$\frac{2 \left(-\frac{1}{2}(b(A - B) - a(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} \right) - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a(A - B) + b(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} \right) - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right)}{d(a^2 + b^2)} -$$

↓ 73

$$\frac{2b(Ab - aB) \int \frac{1}{a \cot^2(c+dx)+b} d\sqrt{\cot(c + dx)}}{d(a^2 + b^2)} -$$

$$\frac{2 \left(-\frac{1}{2}(b(A - B) - a(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} \right) - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a(A - B) + b(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} \right) - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right)}{d(a^2 + b^2)} -$$

↓ 218

$$\frac{2\sqrt{b}(Ab - aB) \arctan\left(\frac{\sqrt{a} \cot(c+dx)}{\sqrt{b}}\right)}{\sqrt{ad}(a^2 + b^2)} -$$

$$\frac{2 \left(-\frac{1}{2}(b(A - B) - a(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} \right) - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(a(A - B) + b(A + B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} \right) - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right)}{d(a^2 + b^2)} -$$

input `Int[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(2*Sqrt[b]*(A*b - a*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]])/(Sqrt[a]*(a^2 + b^2)*d) - (2*(-1/2*((b*(A - B) - a*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2])) - ((a*(A - B) + b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)`

3.591.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4095 `Int[(((A_) + (B_)*tan[(e_) + (f_)*(x_)])*Sqrt[(c_) + (d_)*tan[(e_) + (f_)*(x_)]])/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[Simp[A*(a*c + b*d) + B*(b*c - a*d) - (A*(b*c - a*d) - B*(a*c + b*d))*Tan[e + f*x], x]/Sqrt[c + d*Tan[e + f*x]], x] - Simp[(b*c - a*d)*((B*a - A*b)/(a^2 + b^2)) Int[(1 + Tan[e + f*x]^2)/((a + b*Tan[e + f*x])*Sqrt[c + d*Tan[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :=
  Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
  FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

3.591.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 530 vs. $2(240) = 480$.

Time = 0.40 (sec) , antiderivative size = 531, normalized size of antiderivative = 1.91

method	result
derivativedivides	$\frac{1}{\sqrt{\tan(dx+c)}} \sqrt{\tan(dx+c)} \left(A \ln \left(-\frac{1+\sqrt{2} \sqrt{\tan(dx+c)+\tan(dx+c)}}{\sqrt{2} \sqrt{\tan(dx+c)-\tan(dx+c)-1}} \right) \sqrt{2} \sqrt{ab} a + 2A \arctan \left(1+\sqrt{2} \sqrt{\tan(dx+c)} \right) \sqrt{2} \sqrt{ab} a \right)$
default	$\frac{1}{\sqrt{\tan(dx+c)}} \sqrt{\tan(dx+c)} \left(A \ln \left(-\frac{1+\sqrt{2} \sqrt{\tan(dx+c)+\tan(dx+c)}}{\sqrt{2} \sqrt{\tan(dx+c)-\tan(dx+c)-1}} \right) \sqrt{2} \sqrt{ab} a + 2A \arctan \left(1+\sqrt{2} \sqrt{\tan(dx+c)} \right) \sqrt{2} \sqrt{ab} a \right)$

```
input int(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

```
output 1/4/d*(1/tan(d*x+c))^(1/2)*tan(d*x+c)^(1/2)*(A*ln(-(1+2^(1/2)*tan(d*x+c)^(
1/2)+tan(d*x+c))/(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1))*2^(1/2)*(a*b)^(1
/2)*a+2*A*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*(a*b)^(1/2)*a-2*A*arc
tan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*(a*b)^(1/2)*b+2*A*arctan(-1+2^(1/2
)*tan(d*x+c)^(1/2))*2^(1/2)*(a*b)^(1/2)*a-2*A*arctan(-1+2^(1/2)*tan(d*x+c
)^(1/2))*2^(1/2)*(a*b)^(1/2)*b-A*ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1
)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))*2^(1/2)*(a*b)^(1/2)*b+B*ln(-(1+
2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-
1))*2^(1/2)*(a*b)^(1/2)*b+2*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*(
a*b)^(1/2)*a+2*B*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*(a*b)^(1/2)*b+
2*B*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*(a*b)^(1/2)*a+2*B*arctan(-
1+2^(1/2)*tan(d*x+c)^(1/2))*2^(1/2)*(a*b)^(1/2)*b+B*ln(-(2^(1/2)*tan(d*x+c
)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))*2^(1/2)*(a*
b)^(1/2)*a+8*A*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))*b^2-8*B*arctan(b*tan
(d*x+c)^(1/2)/(a*b)^(1/2))*a*b/(a^2+b^2)/(a*b)^(1/2)
```

3.591.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2995 vs. 2(240) = 480.

Time = 4.33 (sec) , antiderivative size = 6020, normalized size of antiderivative = 21.65

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.591.6 Sympy [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx = \int \frac{(A+B\tan(c+dx))\sqrt{\cot(c+dx)}}{a+b\tan(c+dx)} dx$$

input `integrate(cot(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(cot(c + d*x))/(a + b*tan(c + d*x)), x)`

3.591.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.79

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{8(Bab-Ab^2)\arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right)}{(a^2+b^2)\sqrt{ab}} - \frac{2\sqrt{2}((A+B)a-(A-B)b)\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}+\frac{2}{\sqrt{\tan(dx+c)}}\right)\right)}{(a^2+b^2)\sqrt{ab}} + 2\sqrt{2}((A+B)a-(A-B)b)\arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2}+\frac{2}{\sqrt{\tan(dx+c)}}\right)\right)}{(a^2+b^2)\sqrt{ab}}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

3.591. $\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{a+b\tan(c+dx)} dx$

output $\frac{1}{4} \cdot (8 \cdot (B \cdot a \cdot b - A \cdot b^2) \cdot \arctan(a / (\sqrt{a \cdot b} \cdot \sqrt{\tan(dx + c)}))) / ((a^2 + b^2) \cdot \sqrt{a \cdot b}) - (2 \cdot \sqrt{2}) \cdot ((A + B) \cdot a - (A - B) \cdot b) \cdot \arctan(1/2 \cdot \sqrt{2} \cdot (\sqrt{2} + 2/\sqrt{\tan(dx + c)})) + 2 \cdot \sqrt{2} \cdot ((A + B) \cdot a - (A - B) \cdot b) \cdot \arctan(-1/2 \cdot \sqrt{2} \cdot (\sqrt{2} - 2/\sqrt{\tan(dx + c)})) - \sqrt{2} \cdot ((A - B) \cdot a + (A + B) \cdot b) \cdot \log(\sqrt{2}/\sqrt{\tan(dx + c)} + 1/\tan(dx + c) + 1) + \sqrt{2} \cdot ((A - B) \cdot a + (A + B) \cdot b) \cdot \log(-\sqrt{2}/\sqrt{\tan(dx + c)} + 1/\tan(dx + c) + 1) / (a^2 + b^2) / d$

3.591.8 Giac [F]

$$\int \frac{\sqrt{\cot(c + dx)}(A + B \tan(c + dx))}{a + b \tan(c + dx)} dx = \int \frac{(B \tan(dx + c) + A) \sqrt{\cot(dx + c)}}{b \tan(dx + c) + a} dx$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(cot(d*x + c))/(b*tan(d*x + c) + a), x)`

3.591.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cot(c + dx)}(A + B \tan(c + dx))}{a + b \tan(c + dx)} dx = \int \frac{\sqrt{\cot(c + dx)}(A + B \tan(c + dx))}{a + b \tan(c + dx)} dx$$

input `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x)), x)`

3.592
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))}} dx$$

3.592.1 Optimal result 5617
 3.592.2 Mathematica [A] (verified) 5618
 3.592.3 Rubi [A] (warning: unable to verify) 5618
 3.592.4 Maple [A] (verified) 5624
 3.592.5 Fricas [B] (verification not implemented) 5625
 3.592.6 Sympy [F] 5625
 3.592.7 Maxima [A] (verification not implemented) 5625
 3.592.8 Giac [F] 5626
 3.592.9 Mupad [F(-1)] 5626

3.592.1 Optimal result

Integrand size = 33, antiderivative size = 278

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))}} dx \\ &= \frac{(a(A - B) + b(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}(a^2 + b^2) d} \\ & \quad - \frac{(a(A - B) + b(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}(a^2 + b^2) d} \\ & \quad + \frac{2\sqrt{a}(Ab - aB) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{\sqrt{b}(a^2 + b^2) d} \\ & \quad - \frac{(b(A - B) - a(A + B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}(a^2 + b^2) d} \\ & \quad + \frac{(b(A - B) - a(A + B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}(a^2 + b^2) d} \end{aligned}$$

output

```
-1/2*(a*(A-B)+b*(A+B))*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/2*(a*(A-B)+b*(A+B))*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/4*(b*(A-B)-a*(A+B))*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)+1/4*(b*(A-B)-a*(A+B))*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)+2*(A*b-B*a)*arctan(a^(1/2)*cot(d*x+c)^(1/2)/b^(1/2))*a^(1/2)/(a^2+b^2)/d/b^(1/2)
```

3.592.
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))}} dx$$

3.592.2 Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.77

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx =$$

$$\frac{\sqrt{\cot(c + dx)} \left(2\sqrt{2}(a(A - B) + b(A + B)) \left(\arctan \left(1 - \sqrt{2}\sqrt{\tan(c + dx)} \right) - \arctan \left(1 + \sqrt{2}\sqrt{\tan(c + dx)} \right) \right) \right)}{(a^2 + b^2)d}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])),x]`

output `-1/4*(Sqrt[Cot[c + d*x]]*(2*Sqrt[2]*(a*(A - B) + b*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]) + (8*Sqrt[a]*(A*b - a*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])/Sqrt[b] - Sqrt[2]*(b*(-A + B) + a*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] + Tan[c + d*x] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]] + Tan[c + d*x]))*Sqrt[Tan[c + d*x]]/((a^2 + b^2)*d)`

3.592.3 Rubi [A] (warning: unable to verify)

Time = 1.09 (sec) , antiderivative size = 224, normalized size of antiderivative = 0.81, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {3042, 4064, 3042, 4096, 3042, 4017, 25, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx$$

$$\downarrow \text{4064}$$

$$\int \frac{A \cot(c + dx) + B}{\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} dx$$

3.592. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))} dx$

$$\begin{aligned}
& \int \frac{B - A \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})} (b - a \tan(c + dx + \frac{\pi}{2}))} dx \\
& \quad \downarrow \text{3042} \\
& \frac{\int \frac{aA + bB + (Ab - aB) \cot(c + dx)}{\sqrt{\cot(c + dx)}} dx}{a^2 + b^2} - \frac{a(Ab - aB) \int \frac{\cot^2(c + dx) + 1}{\sqrt{\cot(c + dx)} (b + a \cot(c + dx))} dx}{a^2 + b^2} \\
& \quad \downarrow \text{4096} \\
& \frac{\int \frac{aA + bB - (Ab - aB) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx}{a^2 + b^2} - \frac{a(Ab - aB) \int \frac{\tan(c + dx + \frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c + dx + \frac{\pi}{2})} (b - a \tan(c + dx + \frac{\pi}{2}))} dx}{a^2 + b^2} \\
& \quad \downarrow \text{3042} \\
& \frac{2 \int -\frac{aA + bB + (Ab - aB) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d(a^2 + b^2)} - \frac{a(Ab - aB) \int \frac{\tan(c + dx + \frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c + dx + \frac{\pi}{2})} (b - a \tan(c + dx + \frac{\pi}{2}))} dx}{a^2 + b^2} \\
& \quad \downarrow \text{4017} \\
& \frac{a(Ab - aB) \int \frac{\tan(c + dx + \frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c + dx + \frac{\pi}{2})} (b - a \tan(c + dx + \frac{\pi}{2}))} dx}{a^2 + b^2} - \frac{2 \int \frac{aA + bB + (Ab - aB) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d(a^2 + b^2)} \\
& \quad \downarrow \text{25} \\
& \frac{2 \left(\frac{1}{2} (b(A - B) - a(A + B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \frac{1}{2} (a(A - B) + b(A + B)) \int \frac{\cot(c + dx) + 1}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} \right)}{d(a^2 + b^2)} \\
& \quad \downarrow \text{1482} \\
& \frac{a(Ab - aB) \int \frac{\tan(c + dx + \frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c + dx + \frac{\pi}{2})} (b - a \tan(c + dx + \frac{\pi}{2}))} dx}{a^2 + b^2} \\
& \quad \downarrow \text{1476} \\
& \frac{2 \left(\frac{1}{2} (b(A - B) - a(A + B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \frac{1}{2} (a(A - B) + b(A + B)) \left(\frac{1}{2} \int \frac{1}{\cot(c + dx) - \sqrt{2} \sqrt{\cot(c + dx)}} \right) \right)}{d(a^2 + b^2)} \\
& \quad \downarrow \text{1082} \\
& \frac{a(Ab - aB) \int \frac{\tan(c + dx + \frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c + dx + \frac{\pi}{2})} (b - a \tan(c + dx + \frac{\pi}{2}))} dx}{a^2 + b^2}
\end{aligned}$$

3.592. $\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} (a + b \tan(c + dx))} dx$

$$2 \left(\frac{1}{2}(b(A-B) - a(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a(A-B) + b(A+B)) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right) \frac{d(a^2+b^2)}{d(a^2+b^2)}$$

$$\frac{a(Ab - aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2}$$

↓ 217

$$2 \left(\frac{1}{2}(b(A-B) - a(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(a(A-B) + b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right) \frac{d(a^2+b^2)}{d(a^2+b^2)}$$

$$\frac{a(Ab - aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2}$$

↓ 1479

$$2 \left(\frac{1}{2}(b(A-B) - a(A+B)) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(a(A-B) + b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right) \frac{d(a^2+b^2)}{d(a^2+b^2)}$$

$$\frac{a(Ab - aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2}$$

↓ 25

$$2 \left(\frac{1}{2}(b(A-B) - a(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) - \frac{1}{2}(a(A-B) + b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right) \frac{d(a^2+b^2)}{d(a^2+b^2)}$$

$$\frac{a(Ab - aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2}$$

↓ 27

$$2 \left(\frac{1}{2}(b(A-B) - a(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) - \frac{1}{2}(a(A-B) + b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right) \frac{d(a^2+b^2)}{d(a^2+b^2)}$$

$$\frac{a(Ab - aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2}$$

3.592. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))}} dx$

↓ 1103

$$\frac{2 \left(\frac{1}{2}(b(A - B) - a(A + B)) \left(\frac{\log(\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(a(A - B) + b(A + B)) \right)}{d(a^2 + b^2)}$$

$$\frac{a(Ab - aB) \int \frac{\tan(c+dx + \frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c+dx + \frac{\pi}{2})(b - a \tan(c+dx + \frac{\pi}{2}))}} dx}{a^2 + b^2}$$

↓ 4117

$$\frac{2 \left(\frac{1}{2}(b(A - B) - a(A + B)) \left(\frac{\log(\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(a(A - B) + b(A + B)) \right)}{d(a^2 + b^2)}$$

$$\frac{a(Ab - aB) \int \frac{1}{\sqrt{\cot(c+dx)(b + a \cot(c+dx))}} d(-\cot(c + dx))}{d(a^2 + b^2)}$$

↓ 73

$$\frac{2a(Ab - aB) \int \frac{1}{a \cot^2(c+dx) + b} d\sqrt{\cot(c + dx)}}{d(a^2 + b^2)} +$$

$$\frac{2 \left(\frac{1}{2}(b(A - B) - a(A + B)) \left(\frac{\log(\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(a(A - B) + b(A + B)) \right)}{d(a^2 + b^2)}$$

↓ 218

$$\frac{2 \left(\frac{1}{2}(b(A - B) - a(A + B)) \left(\frac{\log(\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right) - \frac{1}{2}(a(A - B) + b(A + B)) \right)}{d(a^2 + b^2)}$$

$$\frac{2\sqrt{a}(Ab - aB) \arctan\left(\frac{\sqrt{a}\cot(c+dx)}{\sqrt{b}}\right)}{\sqrt{bd}(a^2 + b^2)}$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])),x]`

```
output (-2*Sqrt[a]*(A*b - a*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]]/(Sqrt[b]*(
a^2 + b^2)*d) + (2*(-1/2*((a*(A - B) + b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqr
rt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2
])) + ((b*(A - B) - a*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] +
Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]
/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)
```

3.592.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 73 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 217 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 1082 Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

```
rule 1103 Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])/Sqrt[(b_.)*tan[(e_.) + (f_.)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)])*(g_.)^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

```
rule 4096 Int[(((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[a*A + b*B - (A*b - a*B)*Tan[e + f*x], x], x] + Simp[b*((A*b - a*B)/(a^2 + b^2)) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

3.592.4 Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 244, normalized size of antiderivative = 0.88

method	result
derivativedivides	$\frac{2 \left(\frac{(Aa+Bb)\sqrt{2} \left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right) + 2 \arctan\left(1+\sqrt{2}\sqrt{\cot(dx+c)}\right) + 2 \arctan\left(-1+\sqrt{2}\sqrt{\cot(dx+c)}\right) \right)}{8} \right) (Ab-Ba)}{a^2+b^2}$
default	$\frac{2 \left(\frac{(Aa+Bb)\sqrt{2} \left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right) + 2 \arctan\left(1+\sqrt{2}\sqrt{\cot(dx+c)}\right) + 2 \arctan\left(-1+\sqrt{2}\sqrt{\cot(dx+c)}\right) \right)}{8} \right) (Ab-Ba)}{a^2+b^2}$

```
input int((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c)), x, method=_RETURNVERBOSE)
```

```
output 1/d*(-2/(a^2+b^2)*(1/8*(A*a+B*b)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/8*(A*b-B*a)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+2*(A*b-B*a)*a/(a^2+b^2)/(a*b)^(1/2)*arctan(a*cot(d*x+c)^(1/2)/(a*b)^(1/2)))
```

$$3.592. \int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))} dx$$

3.592.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2983 vs. 2(240) = 480.

Time = 3.09 (sec) , antiderivative size = 5992, normalized size of antiderivative = 21.55

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.592.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx)) \sqrt{\cot(c + dx)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(1/2)/(a+b*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))*sqrt(cot(c + d*x))), x)`

3.592.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 220, normalized size of antiderivative = 0.79

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx = \frac{8(Ba^2 - Aab) \arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right)}{(a^2+b^2)\sqrt{ab}} + \frac{2\sqrt{2}((A-B)a+(A+B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}((A-B)a+(A+B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right)}{(a^2+b^2)\sqrt{ab}}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x, algorithm="maxima")`

3.592. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))} dx$

output
$$\frac{-1/4*(8*(B*a^2 - A*a*b)*\arctan(a/(\sqrt{a*b}*\sqrt{\tan(dx + c)})))/((a^2 + b^2)*\sqrt{a*b}) + (2*\sqrt{2}*((A - B)*a + (A + B)*b)*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2/\sqrt{\tan(dx + c)})) + 2*\sqrt{2}*((A - B)*a + (A + B)*b)*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2/\sqrt{\tan(dx + c)})) + \sqrt{2}*((A + B)*a - (A - B)*b)*\log(\sqrt{2}/\sqrt{\tan(dx + c)} + 1/\tan(dx + c) + 1) - \sqrt{2}*((A + B)*a - (A - B)*b)*\log(-\sqrt{2}/\sqrt{\tan(dx + c)} + 1/\tan(dx + c) + 1))/(a^2 + b^2)/d}$$

3.592.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a)\sqrt{\cot(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((b*tan(d*x + c) + a)*sqrt(cot(d*x + c))), x)`

3.592.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))), x)`

3.593
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$$

3.593.1 Optimal result 5627
 3.593.2 Mathematica [A] (verified) 5628
 3.593.3 Rubi [A] (warning: unable to verify) 5628
 3.593.4 Maple [A] (verified) 5635
 3.593.5 Fricas [B] (verification not implemented) 5636
 3.593.6 Sympy [F] 5636
 3.593.7 Maxima [A] (verification not implemented) 5636
 3.593.8 Giac [F] 5637
 3.593.9 Mupad [F(-1)] 5637

3.593.1 Optimal result

Integrand size = 33, antiderivative size = 297

$$\begin{aligned} & \int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx \\ &= \frac{(b(A-B)-a(A+B)) \arctan\left(1-\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d} \\ & \quad - \frac{(b(A-B)-a(A+B)) \arctan\left(1+\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)d} \\ & \quad - \frac{2a^{3/2}(Ab-aB) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{b^{3/2}(a^2+b^2)d} + \frac{2B}{bd\sqrt{\cot(c+dx)}} \\ & \quad + \frac{(a(A-B)+b(A+B)) \log\left(1-\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d} \\ & \quad - \frac{(a(A-B)+b(A+B)) \log\left(1+\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)d} \end{aligned}$$

output

```
-2*a^(3/2)*(A*b-B*a)*arctan(a^(1/2)*cot(d*x+c)^(1/2)/b^(1/2))/b^(3/2)/(a^2+b^2)/d-1/2*(b*(A-B)-a*(A+B))*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/2*(b*(A-B)-a*(A+B))*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)+1/4*(a*(A-B)+b*(A+B))*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)-1/4*(a*(A-B)+b*(A+B))*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(a^2+b^2)/d*2^(1/2)+2*B/b/d/cot(d*x+c)^(1/2)
```

3.593.2 Mathematica [A] (verified)

Time = 0.60 (sec) , antiderivative size = 251, normalized size of antiderivative = 0.85

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = \frac{\sqrt{\cot(c + dx)} \left(2\sqrt{2}b^{3/2}(b(A - B) - a(A + B)) \left(\arctan \left(1 - \sqrt{2}\sqrt{\tan(c + dx)} \right) - \arctan \left(1 + \sqrt{2}\sqrt{\tan(c + dx)} \right) \right) \right)}{\dots}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])),x]`

output `-1/4*(Sqrt[Cot[c + d*x]]*(2*Sqrt[2]*b^(3/2)*(b*(A - B) - a*(A + B))*(ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]] - ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]]) + 8*a^(3/2)*(-(A*b) + a*B)*ArcTan[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]] - Sqrt[2]*b^(3/2)*(a*(A - B) + b*(A + B))*(Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]]) - 8*Sqrt[b]*(a^2 + b^2)*B*Sqrt[Tan[c + d*x]]*Sqrt[Tan[c + d*x]])/(b^(3/2)*(a^2 + b^2)*d)`

3.593.3 Rubi [A] (warning: unable to verify)

Time = 1.45 (sec) , antiderivative size = 249, normalized size of antiderivative = 0.84, number of steps used = 23, number of rules used = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3042, 4064, 3042, 4092, 27, 3042, 4136, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2}(a + b \tan(c + dx))} dx \\ & \quad \downarrow \text{4064} \\ & \int \frac{A \cot(c + dx) + B}{\cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + b)} dx \end{aligned}$$

$$\begin{aligned}
 & \downarrow 3042 \\
 & \int \frac{B - A \tan\left(c + dx + \frac{\pi}{2}\right)}{\left(-\tan\left(c + dx + \frac{\pi}{2}\right)\right)^{3/2} \left(b - a \tan\left(c + dx + \frac{\pi}{2}\right)\right)} dx \\
 & \downarrow 4092 \\
 & \frac{2 \int \frac{-aB \cot^2(c+dx) - bB \cot(c+dx) + Ab - aB}{2\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} dx}{b} + \frac{2B}{bd\sqrt{\cot(c+dx)}} \\
 & \downarrow 27 \\
 & \frac{\int \frac{-aB \cot^2(c+dx) - bB \cot(c+dx) + Ab - aB}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} dx}{b} + \frac{2B}{bd\sqrt{\cot(c+dx)}} \\
 & \downarrow 3042 \\
 & \frac{\int \frac{-aB \tan\left(c+dx+\frac{\pi}{2}\right)^2 + bB \tan\left(c+dx+\frac{\pi}{2}\right) + Ab - aB}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)\left(b-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)}} dx}{b} + \frac{2B}{bd\sqrt{\cot(c+dx)}} \\
 & \downarrow 4136 \\
 & \frac{a^2(Ab-aB) \int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} dx}{a^2+b^2} + \frac{\int \frac{b(Ab-aB)-b(aA+bB) \cot(c+dx)}{\sqrt{\cot(c+dx)}} dx}{a^2+b^2} + \frac{2B}{bd\sqrt{\cot(c+dx)}} \\
 & \downarrow 3042 \\
 & \frac{a^2(Ab-aB) \int \frac{\tan\left(c+dx+\frac{\pi}{2}\right)^2+1}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)\left(b-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)}} dx}{a^2+b^2} + \frac{\int \frac{b(Ab-aB)+b(aA+bB) \tan\left(c+dx+\frac{\pi}{2}\right)}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}} dx}{a^2+b^2} + \frac{2B}{bd\sqrt{\cot(c+dx)}} \\
 & \downarrow 4017 \\
 & \frac{a^2(Ab-aB) \int \frac{\tan\left(c+dx+\frac{\pi}{2}\right)^2+1}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)\left(b-a \tan\left(c+dx+\frac{\pi}{2}\right)\right)}} dx}{a^2+b^2} + \frac{2 \int \frac{-b(Ab-aB)-(aA+bB) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} + \\
 & \frac{b}{2B} \\
 & \frac{2B}{bd\sqrt{\cot(c+dx)}} \\
 & \downarrow 25
 \end{aligned}$$

3.593. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$\begin{aligned}
 & \frac{a^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2 \int \frac{b(Ab-aB-(aA+bB) \cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} + \\
 & \qquad \qquad \qquad \frac{\frac{b}{2B}}{bd\sqrt{\cot(c+dx)}} \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & \frac{a^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2b \int \frac{Ab-aB-(aA+bB) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} + \\
 & \qquad \qquad \qquad \frac{\frac{b}{2B}}{bd\sqrt{\cot(c+dx)}} \\
 & \qquad \qquad \qquad \downarrow 1482 \\
 & \frac{a^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2b \left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(b(A-B)-a(A+B)) \int \frac{\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d(a^2+b^2)} + \\
 & \qquad \qquad \qquad \frac{\frac{2B}{b}}{bd\sqrt{\cot(c+dx)}} \\
 & \qquad \qquad \qquad \downarrow 1476 \\
 & \frac{a^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2b \left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{1}{2} \int \frac{\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right) \right)}{d(a^2+b^2)} + \\
 & \qquad \qquad \qquad \frac{\frac{2B}{b}}{bd\sqrt{\cot(c+dx)}} \\
 & \qquad \qquad \qquad \downarrow 1082 \\
 & \frac{a^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2b \left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{1}{2} \int \frac{\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right) \right)}{d(a^2+b^2)} + \\
 & \qquad \qquad \qquad \frac{\frac{2B}{b}}{bd\sqrt{\cot(c+dx)}} \\
 & \qquad \qquad \qquad \downarrow 217
 \end{aligned}$$

3.593. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$\frac{a^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2b \left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)}$$

$$\frac{2B}{bd\sqrt{\cot(c+dx)}}$$

↓ 1479

$$\frac{a^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2b \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}}}{2\sqrt{2}} \right) \right)}{b}$$

$$\frac{2B}{bd\sqrt{\cot(c+dx)}}$$

↓ 25

$$\frac{a^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2b \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}}}{2\sqrt{2}} \right) \right)}{b}$$

$$\frac{2B}{bd\sqrt{\cot(c+dx)}}$$

↓ 27

$$\frac{a^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2b \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}} \right) \right)}{b}$$

$$\frac{2B}{bd\sqrt{\cot(c+dx)}}$$

↓ 1103

$$\frac{a^2(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2b \left(\frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}})}{\sqrt{2}} - \frac{\arctan(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}})}{\sqrt{2}} \right) \right) + \frac{1}{2}(a(A-B)+b(A+B)) \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}}}{b}$$

$$\frac{2B}{bd\sqrt{\cot(c+dx)}}$$

↓ 4117

3.593. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$\begin{aligned}
 & \frac{a^2(Ab-aB) \int \frac{1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} d(-\cot(c+dx))}{d(a^2+b^2)} - \frac{2b \left(\frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} \right)}{b} \\
 & \qquad \qquad \qquad \frac{2B}{bd\sqrt{\cot(c+dx)}} \\
 & \qquad \qquad \qquad \downarrow \text{73} \\
 & \frac{2a^2(Ab-aB) \int \frac{1}{a \cot^2(c+dx)+b} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} - \frac{2b \left(\frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(a(A-B)+b(A+B)) \right)}{b} \\
 & \qquad \qquad \qquad \frac{2B}{bd\sqrt{\cot(c+dx)}} \\
 & \qquad \qquad \qquad \downarrow \text{218} \\
 & \frac{2a^{3/2}(Ab-aB) \arctan\left(\frac{\sqrt{a}\cot(c+dx)}{\sqrt{b}}\right)}{\sqrt{bd}(a^2+b^2)} - \frac{2b \left(\frac{1}{2}(b(A-B)-a(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2}(a(A-B)+b(A+B)) \right)}{b} \\
 & \qquad \qquad \qquad \frac{2B}{bd\sqrt{\cot(c+dx)}}
 \end{aligned}$$

```
input Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])),x]
```

```
output (2*B)/(b*d*Sqrt[Cot[c + d*x]]) + ((2*a^(3/2)*(A*b - a*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]])/(Sqrt[b]*(a^2 + b^2)*d) - (2*b*((b*(A - B) - a*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]))/2 + ((a*(A - B) + b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)/b
```

3.593.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

3.593. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$

- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
 -1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
 & (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
 t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
 implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
 eQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
 imp[d*(Log[RemoveContent[a + b*x + c*x^2, x])/b], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
 2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
 e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
 && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
 -2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
 x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
 reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.593.4 Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.89

method	result
derivativedivides	$\frac{\frac{2B}{b\sqrt{\cot(dx+c)}}}{2} - \frac{(Ab-Ba)\sqrt{2} \left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right) + 2\arctan\left(\frac{1+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\sqrt{2}\sqrt{\cot(dx+c)}}\right) + 2\arctan\left(\frac{-1+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\sqrt{2}\sqrt{\cot(dx+c)}}\right) \right)}{8}$
default	$\frac{\frac{2B}{b\sqrt{\cot(dx+c)}}}{2} - \frac{(Ab-Ba)\sqrt{2} \left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right) + 2\arctan\left(\frac{1+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\sqrt{2}\sqrt{\cot(dx+c)}}\right) + 2\arctan\left(\frac{-1+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\sqrt{2}\sqrt{\cot(dx+c)}}\right) \right)}{8}$

```
input int((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x,method=_RETURNVER
BOSE)
```

```
output 1/d*(2*B/b/cot(d*x+c)^(1/2)-2/(a^2+b^2)*(1/8*(A*b-B*a)*2^(1/2)*(ln((1+cot(
d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+
2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))
)+1/8*(-A*a-B*b)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1co
t(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2
*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))))-2/b*(A*b-B*a)*a^2/(a^2+b^2)/(a*b)^(
1/2)*arctan(a*cot(d*x+c)^(1/2)/(a*b)^(1/2)))
```

$$3.593. \int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$$

3.593.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3033 vs. 2(257) = 514.

Time = 8.78 (sec) , antiderivative size = 6092, normalized size of antiderivative = 20.51

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.593.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx)) \cot^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(3/2)/(a+b*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))*cot(c + d*x)**(3/2)), x)`

3.593.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 238, normalized size of antiderivative = 0.80

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$= \frac{8(Ba^3 - Aa^2b) \arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right)}{(a^2b + b^3)\sqrt{ab}} + \frac{8B\sqrt{\tan(dx+c)}}{b} + \frac{2\sqrt{2}((A+B)a - (A-B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}((A+B)a - (A-B)b)}{(a^2b + b^3)\sqrt{ab}}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x, algorithm="maxima")`

3.593. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$

output $1/4*(8*(B*a^3 - A*a^2*b)*\arctan(a/(\sqrt{a*b}*\sqrt{\tan(d*x + c)})))/((a^2*b + b^3)*\sqrt{a*b}) + 8*B*\sqrt{\tan(d*x + c)}/b + (2*\sqrt{2}*((A + B)*a - (A - B)*b)*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2/\sqrt{\tan(d*x + c)})) + 2*\sqrt{2}*((A + B)*a - (A - B)*b)*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2/\sqrt{\tan(d*x + c)})) - \sqrt{2}*((A - B)*a + (A + B)*b)*\log(\sqrt{2}/\sqrt{\tan(d*x + c)}) + 1/\tan(d*x + c) + 1 + \sqrt{2}*((A - B)*a + (A + B)*b)*\log(-\sqrt{2}/\sqrt{\tan(d*x + c)}) + 1/\tan(d*x + c) + 1)/(a^2 + b^2))/d$

3.593.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a) \cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((b*tan(d*x + c) + a)*cot(d*x + c)^(3/2)), x)`

3.593.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2} (a + b \tan(c + dx))} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))), x)`

3.594
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$$

3.594.1 Optimal result 5638
 3.594.2 Mathematica [A] (verified) 5639
 3.594.3 Rubi [A] (warning: unable to verify) 5639
 3.594.4 Maple [A] (verified) 5647
 3.594.5 Fricas [B] (verification not implemented) 5648
 3.594.6 Sympy [F] 5648
 3.594.7 Maxima [A] (verification not implemented) 5649
 3.594.8 Giac [F] 5649
 3.594.9 Mupad [F(-1)] 5650

3.594.1 Optimal result

Integrand size = 33, antiderivative size = 325

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx \\ &= -\frac{(a(A - B) + b(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)d} \\ & \quad + \frac{(a(A - B) + b(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)d} \\ & \quad + \frac{2a^{5/2}(Ab - aB) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{b^{5/2}(a^2 + b^2)d} + \frac{2B}{3bd \cot^{\frac{3}{2}}(c + dx)} + \frac{2(Ab - aB)}{b^2d\sqrt{\cot(c + dx)}} \\ & \quad + \frac{(b(A - B) - a(A + B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)d} \\ & \quad - \frac{(b(A - B) - a(A + B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)d} \end{aligned}$$

output $2*a^{(5/2)}*(A*b-B*a)*\arctan(a^{(1/2)}*\cot(d*x+c)^{(1/2)}/b^{(1/2)})/b^{(5/2)}/(a^2+b^2)/d+2/3*B/b/d/\cot(d*x+c)^{(3/2)}+1/2*(a*(A-B)+b*(A+B))*\arctan(-1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)/d*2^{(1/2)}+1/2*(a*(A-B)+b*(A+B))*\arctan(1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)/d*2^{(1/2)}+1/4*(b*(A-B)-a*(A+B))*\ln(1+\cot(d*x+c)-2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)/d*2^{(1/2)}-1/4*(b*(A-B)-a*(A+B))*\ln(1+\cot(d*x+c)+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)/d*2^{(1/2)}+2*(A*b-B*a)/b^2/d/\cot(d*x+c)^{(1/2)}$

3.594.2 Mathematica [A] (verified)

Time = 1.11 (sec) , antiderivative size = 272, normalized size of antiderivative = 0.84

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$= \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(\frac{6\sqrt{2}(a(A-B)+b(A+B))(\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})-\arctan(1+\sqrt{2}\sqrt{\tan(c+dx)})}{a^2+b^2} \right) + \frac{24a^{5/2}(-A+B)}{b^2}}{12d}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])),x]`

output $(\text{Sqrt}[\text{Cot}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]*((6*\text{Sqrt}[2]*(a*(A - B) + b*(A + B)) * (\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]] - \text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]])))/(a^2 + b^2) + (24*a^{(5/2)}*(-(A*b) + a*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(b^{(5/2)}*(a^2 + b^2)) - (3*\text{Sqrt}[2]*(b*(-A + B) + a*(A + B))*(\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]] - \text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]]))/(a^2 + b^2) + (24*(A*b - a*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/b^2 + (8*B*\text{Tan}[c + d*x]^{(3/2)})/b)/(12*d)$

3.594.3 Rubi [A] (warning: unable to verify)

Time = 1.85 (sec) , antiderivative size = 285, normalized size of antiderivative = 0.88, number of steps used = 26, number of rules used = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.758$, Rules used = {3042, 4064, 3042, 4092, 27, 3042, 4132, 27, 3042, 4136, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.594. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$\begin{aligned}
& \int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx \\
& \quad \downarrow 3042 \\
& \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{5/2}(a + b \tan(c + dx))} dx \\
& \quad \downarrow 4064 \\
& \int \frac{A \cot(c + dx) + B}{\cot^{\frac{5}{2}}(c + dx)(a \cot(c + dx) + b)} dx \\
& \quad \downarrow 3042 \\
& \int \frac{B - A \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{5/2}(b - a \tan(c + dx + \frac{\pi}{2}))} dx \\
& \quad \downarrow 4092 \\
& \frac{2 \int \frac{3(-aB \cot^2(c+dx) - bB \cot(c+dx) + Ab - aB)}{2 \cot^{\frac{3}{2}}(c+dx)(b+a \cot(c+dx))} dx}{3b} + \frac{2B}{3bd \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 27 \\
& \frac{\int \frac{-aB \cot^2(c+dx) - bB \cot(c+dx) + Ab - aB}{\cot^{\frac{3}{2}}(c+dx)(b+a \cot(c+dx))} dx}{b} + \frac{2B}{3bd \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 3042 \\
& \frac{\int \frac{-aB \tan(c+dx + \frac{\pi}{2})^2 + bB \tan(c+dx + \frac{\pi}{2}) + Ab - aB}{(-\tan(c+dx + \frac{\pi}{2}))^{3/2}(b - a \tan(c+dx + \frac{\pi}{2}))} dx}{b} + \frac{2B}{3bd \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 4132 \\
& \frac{2 \int -\frac{Ba^2 + (Ab - aB) \cot^2(c+dx)a + Aba + b^2B + Ab^2 \cot(c+dx)}{2\sqrt{\cot(c+dx)}(b+a \cot(c+dx))} dx}{b} + \frac{2(Ab - aB)}{bd\sqrt{\cot(c+dx)}} + \frac{2B}{3bd \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 27 \\
& \frac{\frac{2(Ab - aB)}{bd\sqrt{\cot(c+dx)}} - \frac{\int -\frac{Ba^2 + (Ab - aB) \cot^2(c+dx)a + Aba + b^2B + Ab^2 \cot(c+dx)}{\sqrt{\cot(c+dx)}(b+a \cot(c+dx))} dx}{b}}{b} + \frac{2B}{3bd \cot^{\frac{3}{2}}(c + dx)} \\
& \quad \downarrow 3042
\end{aligned}$$

3.594. $\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx$

$$\begin{aligned}
 & \frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{\int \frac{-Ba^2+(Ab-aB)\tan(c+dx+\frac{\pi}{2})^2 a+Ab a+b^2 B-Ab^2 \tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx}{b} + \frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow 4136 \\
 & \frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{\int \frac{(aA+bB)b^2+(Ab-aB)\cot(c+dx)b^2}{\sqrt{\cot(c+dx)}} dx}{a^2+b^2} + \frac{a^3(Ab-aB) \int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)}(b+a \cot(c+dx))} dx}{b} + \frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow 3042 \\
 & \frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{\int \frac{b^2(aA+bB)-b^2(Ab-aB)\tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{a^2+b^2} + \frac{a^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx}{b} + \frac{b}{2B} \\
 & \quad \quad \quad \frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow 4017 \\
 & \frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{2 \int \frac{b^2(aA+bB+(Ab-aB)\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} + \frac{a^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx}{b} + \frac{b}{2B} \\
 & \quad \quad \quad \frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow 25 \\
 & \frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{a^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx}{a^2+b^2} - \frac{2 \int \frac{b^2(aA+bB+(Ab-aB)\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} + \frac{b}{2B} \\
 & \quad \quad \quad \frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)} \\
 & \quad \downarrow 27 \\
 & \frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{a^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx}{a^2+b^2} - \frac{2b^2 \int \frac{aA+bB+(Ab-aB)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} + \frac{b}{2B} \\
 & \quad \quad \quad \frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)}
 \end{aligned}$$

3.594. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$

↓ 1482

$$\frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{a^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2b^2\left(\frac{1}{2}(a(A-B)+b(A+B)) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(b(A-B)-a(A+B)) \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}\right)}{d(a^2+b^2)}$$

$$\frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)}$$

↓ 1476

$$\frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{a^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2b^2\left(\frac{1}{2}(a(A-B)+b(A+B))\left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}\right)\right)}{b}$$

$$\frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)}$$

↓ 1082

$$\frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{a^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2b^2\left(\frac{1}{2}(a(A-B)+b(A+B))\left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)+1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}}\right)\right)}{b}$$

$$\frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)}$$

↓ 217

$$\frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{a^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2b^2\left(\frac{1}{2}(a(A-B)+b(A+B))\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}}\right)\right)}{d(a^2+b^2)}$$

$$\frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)}$$

↓ 1479

3.594. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$\frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{a^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 2b^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)$$

$$\frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)}$$

↓ 25

$$\frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{a^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 2b^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)$$

$$\frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)}$$

↓ 27

$$\frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{a^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 2b^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)$$

$$\frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)}$$

↓ 1103

$$\frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{a^3(Ab-aB) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 2b^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)$$

$$\frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)}$$

↓ 4117

3.594. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$\frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{a^3(Ab-aB) \int \frac{1}{\sqrt{\cot(c+dx)(b+a\cot(c+dx))}} d(-\cot(c+dx))}{d(a^2+b^2)} - \frac{2b^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{b}$$

$$\frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)}$$

↓ 73

$$\frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{2a^3(Ab-aB) \int \frac{1}{a \cot^2(c+dx)+b} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} - \frac{2b^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{b}$$

$$\frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)}$$

↓ 218

$$\frac{2(Ab-aB)}{bd\sqrt{\cot(c+dx)}} - \frac{2a^{5/2}(Ab-aB) \arctan\left(\frac{\sqrt{a}\cot(c+dx)}{\sqrt{b}}\right)}{\sqrt{bd}(a^2+b^2)} - \frac{2b^2 \left(\frac{1}{2}(a(A-B)+b(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2}(b(A-B))}{d(a^2+b^2)}$$

$$\frac{2B}{3bd \cot^{\frac{3}{2}}(c+dx)}$$

input `Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])),x]`

output `(2*B)/(3*b*d*Cot[c + d*x]^(3/2)) + ((2*(A*b - a*B))/(b*d*Sqrt[Cot[c + d*x]]) - ((2*a^(5/2)*(A*b - a*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]])/(Sqrt[b]*(a^2 + b^2)*d) - (2*b^2*((a*(A - B) + b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]))/2 - ((b*(A - B) - a*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/(a^2 + b^2*d)/b/b`

3.594. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$

3.594.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;`
`FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /;`
`FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!LtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /;`
`FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.594.4 Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 286, normalized size of antiderivative = 0.88

method	result
derivativedivides	$\frac{2a^3(Ab-Ba) \arctan\left(\frac{a\sqrt{\cot(dx+c)}}{\sqrt{ab}}\right)}{b^2(a^2+b^2)\sqrt{ab}} - \frac{2 \left(\frac{(-Aa-Bb)\sqrt{2}}{8} \left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right) + 2 \arctan(1+\sqrt{2}\sqrt{\cot(dx+c)}) + 2 \arctan(1-\sqrt{2}\sqrt{\cot(dx+c)}) \right) \right)}{8}$
default	$\frac{2a^3(Ab-Ba) \arctan\left(\frac{a\sqrt{\cot(dx+c)}}{\sqrt{ab}}\right)}{b^2(a^2+b^2)\sqrt{ab}} - \frac{2 \left(\frac{(-Aa-Bb)\sqrt{2}}{8} \left(\ln\left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}\right) + 2 \arctan(1+\sqrt{2}\sqrt{\cot(dx+c)}) + 2 \arctan(1-\sqrt{2}\sqrt{\cot(dx+c)}) \right) \right)}{8}$

3.594. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/d*(2/b^2*a^3*(A*b-B*a)/(a^2+b^2)/(a*b)^(1/2)*arctan(a*cot(d*x+c)^(1/2)/(a*b)^(1/2))-2/(a^2+b^2)*(1/8*(-A*a-B*b)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/8*(-A*b+B*a)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+2/3*B/b/cot(d*x+c)^(3/2)+2*(A*b-B*a)/b^2/cot(d*x+c)^(1/2)`

3.594.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3075 vs. $2(281) = 562$.

Time = 21.24 (sec) , antiderivative size = 6176, normalized size of antiderivative = 19.00

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.594.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx)) \cot^{\frac{5}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(5/2)/(a+b*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))*cot(c + d*x)**(5/2)),x)`

3.594.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 262, normalized size of antiderivative = 0.81

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx =$$

$$\frac{24(Ba^4 - Aa^3b) \arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right)}{(a^2b^2 + b^4)\sqrt{ab}} - \frac{8\left(Bb - \frac{3(Ba - Ab)}{\tan(dx+c)}\right) \tan(dx+c)^{\frac{3}{2}}}{b^2} - \frac{3\left(2\sqrt{2}((A-B)a + (A+B)b) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right)\right)}{b^2}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `-1/12*(24*(B*a^4 - A*a^3*b)*arctan(a/(sqrt(a*b)*sqrt(tan(d*x + c))))/((a^2 *b^2 + b^4)*sqrt(a*b)) - 8*(B*b - 3*(B*a - A*b)/tan(d*x + c))*tan(d*x + c)^(3/2)/b^2 - 3*(2*sqrt(2)*((A - B)*a + (A + B)*b)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a + (A + B)*b)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)*((A + B)*a - (A - B)*b)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*((A + B)*a - (A - B)*b)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))/(a^2 + b^2))/d`

3.594.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a) \cot(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((b*tan(d*x + c) + a)*cot(d*x + c)^(5/2)), x)`

3.594.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{\frac{5}{2}}(a + b \tan(c + dx))} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + b*tan(c + d*x))),x)`output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + b*tan(c + d*x))), x)`

$$3.595 \quad \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.595.1 Optimal result	5651
3.595.2 Mathematica [A] (verified)	5652
3.595.3 Rubi [A] (warning: unable to verify)	5653
3.595.4 Maple [B] (verified)	5661
3.595.5 Fricas [B] (verification not implemented)	5662
3.595.6 Sympy [F]	5663
3.595.7 Maxima [A] (verification not implemented)	5663
3.595.8 Giac [F]	5664
3.595.9 Mupad [F(-1)]	5664

3.595.1 Optimal result

Integrand size = 33, antiderivative size = 438

$$\begin{aligned} & \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\ &= -\frac{(a^2(A-B) - b^2(A-B) + 2ab(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^2 d} \\ & \quad + \frac{(a^2(A-B) - b^2(A-B) + 2ab(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^2 d} \\ & \quad + \frac{b^{3/2}(7a^2 Ab + 3Ab^3 - 5a^3 B - ab^2 B) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{a^{5/2}(a^2+b^2)^2 d} \\ & \quad - \frac{(2a^2 A + 3Ab^2 - abB) \sqrt{\cot(c+dx)}}{a^2(a^2+b^2) d} + \frac{b(Ab - aB) \cot^{\frac{3}{2}}(c+dx)}{a(a^2+b^2) d(b+a \cot(c+dx))} \\ & \quad + \frac{(2ab(A-B) - a^2(A+B) + b^2(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^2 d} \\ & \quad - \frac{(2ab(A-B) - a^2(A+B) + b^2(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^2 d} \end{aligned}$$

$$3.595. \quad \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

output $b^{3/2}*(7*A*a^2*b+3*A*b^3-5*B*a^3-B*a*b^2)*\arctan(a^{1/2}*\cot(d*x+c)^{1/2})/b^{1/2})/a^{5/2}/(a^2+b^2)^2/d+b*(A*b-B*a)*\cot(d*x+c)^{3/2}/a/(a^2+b^2)/d/(b+a*\cot(d*x+c))+1/2*(a^2*(A-B)-b^2*(A-B)+2*a*b*(A+B))*\arctan(-1+2^{1/2}*\cot(d*x+c)^{1/2})/(a^2+b^2)^2/d*2^{1/2}+1/2*(a^2*(A-B)-b^2*(A-B)+2*a*b*(A+B))*\arctan(1+2^{1/2}*\cot(d*x+c)^{1/2})/(a^2+b^2)^2/d*2^{1/2}+1/4*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\ln(1+\cot(d*x+c)-2^{1/2}*\cot(d*x+c)^{1/2})/(a^2+b^2)^2/d*2^{1/2}-1/4*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\ln(1+\cot(d*x+c)+2^{1/2}*\cot(d*x+c)^{1/2})/(a^2+b^2)^2/d*2^{1/2}-(2*A*a^2+3*A*b^2-B*a*b)*\cot(d*x+c)^{1/2}/a^2/(a^2+b^2)/d$

3.595.2 Mathematica [A] (verified)

Time = 5.88 (sec) , antiderivative size = 383, normalized size of antiderivative = 0.87

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{\sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)} \left(\frac{2\sqrt{2}(a^2(A-B)+b^2(-A+B)+2ab(A+B))(\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})-\arctan(1+\sqrt{2}\sqrt{\tan(c+dx)})}{(a^2+b^2)^2} \right)}{\dots}$$

input `Integrate[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2, x]`

output $(\text{Sqrt}[\text{Cot}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]*((2*\text{Sqrt}[2]*(a^2*(A - B) + b^2*(-A + B) + 2*a*b*(A + B))*(\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]] - \text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]]))/(a^2 + b^2)^2 + (4*b^{3/2}*(-(A*b) + a*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(a^{5/2}*(a^2 + b^2)) - (8*b^{3/2}*(3*a^2*A*b + A*b^3 - 2*a^3*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(a^{5/2}*(a^2 + b^2)^2) - (\text{Sqrt}[2]*(2*a*b*(-A + B) + a^2*(A + B) - b^2*(A + B))*(\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]] - \text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]])))/(a^2 + b^2)^2 - (8*A)/(a^2*\text{Sqrt}[\text{Tan}[c + d*x]]) + (4*b^2*(-(A*b) + a*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/(a^2*(a^2 + b^2)*(a + b*\text{Tan}[c + d*x])))/(4*d)$

3.595. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.595.3 Rubi [A] (warning: unable to verify)

Time = 2.11 (sec) , antiderivative size = 377, normalized size of antiderivative = 0.86, number of steps used = 27, number of rules used = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.788$, Rules used = {3042, 4064, 3042, 4088, 27, 3042, 4130, 27, 3042, 4136, 27, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\cot(c+dx)^{3/2}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{\cot^{\frac{5}{2}}(c+dx)(A \cot(c+dx)+B)}{(a \cot(c+dx)+b)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{5/2}(B-A \tan(c+dx+\frac{\pi}{2}))}{(b-a \tan(c+dx+\frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{4088} \\
 & \frac{b(Ab-aB) \cot^{\frac{3}{2}}(c+dx)}{ad(a^2+b^2)(a \cot(c+dx)+b)} - \\
 & \frac{\int -\frac{\sqrt{\cot(c+dx)}((2Aa^2-bBa+3Ab^2) \cot^2(c+dx)-2a(Ab-aB) \cot(c+dx)+3b(Ab-aB))}{2(b+a \cot(c+dx))} dx}{a(a^2+b^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\sqrt{\cot(c+dx)}((2Aa^2-bBa+3Ab^2) \cot^2(c+dx)-2a(Ab-aB) \cot(c+dx)+3b(Ab-aB))}{b+a \cot(c+dx)} dx}{2a(a^2+b^2)} + \\
 & \frac{b(Ab-aB) \cot^{\frac{3}{2}}(c+dx)}{ad(a^2+b^2)(a \cot(c+dx)+b)} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.595. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\begin{aligned}
 & \frac{\int \frac{\sqrt{-\tan(c+dx+\frac{\pi}{2})} \left((2Aa^2 - bBa + 3Ab^2) \tan(c+dx+\frac{\pi}{2})^2 + 2a(Ab - aB) \tan(c+dx+\frac{\pi}{2}) + 3b(Ab - aB) \right)}{b - a \tan(c+dx+\frac{\pi}{2})} dx}{2a(a^2 + b^2)} + \\
 & \frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)} \\
 & \quad \downarrow \text{4130} \\
 & \frac{2 \int \frac{2(aA + bB) \cot(c+dx)a^2 + (-2Ba^3 + 4Aba^2 - b^2Ba + 3Ab^3) \cot^2(c+dx) + b(2Aa^2 - bBa + 3Ab^2)}{2\sqrt{\cot(c+dx)}(b + a \cot(c+dx))} dx}{a} - \frac{2(2a^2A - abB + 3Ab^2)\sqrt{\cot(c+dx)}}{ad} + \\
 & \frac{2a(a^2 + b^2)}{ad(a^2 + b^2)(a \cot(c + dx) + b)} \frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{2(aA + bB) \cot(c+dx)a^2 + (-2Ba^3 + 4Aba^2 - b^2Ba + 3Ab^3) \cot^2(c+dx) + b(2Aa^2 - bBa + 3Ab^2)}{\sqrt{\cot(c+dx)}(b + a \cot(c+dx))} dx}{a} - \frac{2(2a^2A - abB + 3Ab^2)\sqrt{\cot(c+dx)}}{ad} + \\
 & \frac{2a(a^2 + b^2)}{ad(a^2 + b^2)(a \cot(c + dx) + b)} \frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{-2(aA + bB) \tan(c+dx+\frac{\pi}{2})a^2 + (-2Ba^3 + 4Aba^2 - b^2Ba + 3Ab^3) \tan(c+dx+\frac{\pi}{2})^2 + b(2Aa^2 - bBa + 3Ab^2)}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b - a \tan(c+dx+\frac{\pi}{2}))} dx}{a} - \frac{2(2a^2A - abB + 3Ab^2)\sqrt{\cot(c+dx)}}{ad} + \\
 & \frac{2a(a^2 + b^2)}{ad(a^2 + b^2)(a \cot(c + dx) + b)} \frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)} \\
 & \quad \downarrow \text{4136} \\
 & \frac{\int \frac{2((Aa^2 + 2bBa - Ab^2)a^2 + (-Ba^2 + 2Aba + b^2B) \cot(c+dx)a^2)}{\sqrt{\cot(c+dx)}(a^2 + b^2)} dx}{a} + \frac{b^2(-5a^3B + 7a^2Ab - ab^2B + 3Ab^3) \int \frac{\cot^2(c+dx) + 1}{\sqrt{\cot(c+dx)}(b + a \cot(c+dx))} dx}{a^2 + b^2} - \frac{2(2a^2A - abB + 3Ab^2)\sqrt{\cot(c+dx)}}{ad} + \\
 & \frac{2a(a^2 + b^2)}{ad(a^2 + b^2)(a \cot(c + dx) + b)} \frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

3.595. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\frac{2 \int \frac{(Aa^2+2bBa-Ab^2)a^2+(-Ba^2+2Aba+b^2B) \cot(c+dx)a^2}{\sqrt{\cot(c+dx)}} dx + b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} dx}{a^2+b^2} - \frac{2(2a^2A-abB+3a^2b^2)}{a}$$

$$\frac{2a(a^2+b^2)}{ad(a^2+b^2)(a \cot(c+dx)+b)}$$

↓ 3042

$$\frac{2 \int \frac{a^2(Aa^2+2bBa-Ab^2)-a^2(-Ba^2+2Aba+b^2B) \tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx + b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2(2a^2A-abB+3a^2b^2)}{a}$$

$$\frac{2a(a^2+b^2)}{ad(a^2+b^2)(a \cot(c+dx)+b)}$$

↓ 4017

$$4 \int \frac{a^2(Aa^2+2bBa-Ab^2+(-Ba^2+2Aba+b^2B) \cot(c+dx)) d\sqrt{\cot(c+dx)}}{\cot^2(c+dx)+1} + \frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{2(2a^2A-abB+3a^2b^2)}{a}$$

$$\frac{2a(a^2+b^2)}{ad(a^2+b^2)(a \cot(c+dx)+b)}$$

↓ 25

$$b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx - \frac{4 \int \frac{a^2(Aa^2+2bBa-Ab^2+(-Ba^2+2Aba+b^2B) \cot(c+dx)) d\sqrt{\cot(c+dx)}}{\cot^2(c+dx)+1}}{a^2+b^2} - \frac{2(2a^2A-abB+3a^2b^2)}{a}$$

$$\frac{2a(a^2+b^2)}{ad(a^2+b^2)(a \cot(c+dx)+b)}$$

↓ 27

$$b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx - \frac{4a^2 \int \frac{Aa^2+2bBa-Ab^2+(-Ba^2+2Aba+b^2B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{a^2+b^2} - \frac{2(2a^2A-abB+3a^2b^2)}{a}$$

$$\frac{2a(a^2+b^2)}{ad(a^2+b^2)(a \cot(c+dx)+b)}$$

3.595. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

↓ 1482

$$\frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 4a^2 \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)-1} \right) - \frac{a}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

2a(a² + b²)

↓ 1476

$$\frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 4a^2 \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)}} dx \right) \right) - \frac{a}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 1082

$$\frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 4a^2 \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right) - \frac{a}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 217

$$\frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 4a^2 \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right) - \frac{a}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 1479

3.595. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 4a^2 \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - a \right) \right)$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 25

$$\frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 4a^2 \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - a \right) \right)$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 27

$$\frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 4a^2 \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - a \right) \right)$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 1103

$$\frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 4a^2 \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - a \right) \right)$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 4117

3.595. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\frac{b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} d(-\cot(c+dx))}{d(a^2+b^2)} - 4a^2 \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \arctan(1-\sqrt{\cot(c+dx)}) \right) \right)$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 73

$$\frac{2b^2(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \int \frac{1}{a \cot^2(c+dx)+b} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} - 4a^2 \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \arctan(1-\sqrt{\cot(c+dx)}) \right) \right)$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 218

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{ad(a^2 + b^2)(a \cot(c + dx) + b)} +$$

$$\frac{2b^{3/2}(-5a^3B+7a^2Ab-ab^2B+3Ab^3) \arctan\left(\frac{\sqrt{a} \cot(c+dx)}{\sqrt{b}}\right)}{\sqrt{ad}(a^2+b^2)} - 4a^2 \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \arctan(1-\sqrt{\cot(c+dx)}) \right) \right)$$

input `Int[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `(b*(A*b - a*B)*Cot[c + d*x]^(3/2))/(a*(a^2 + b^2)*d*(b + a*Cot[c + d*x])) + ((-2*(2*a^2*A + 3*A*b^2 - a*b*B)*Sqrt[Cot[c + d*x]])/(a*d) - ((2*b^(3/2)*(7*a^2*A*b + 3*A*b^3 - 5*a^3*B - a*b^2*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]])/(Sqrt[a]*(a^2 + b^2)*d) - (4*a^2*((a^2*(A - B) - b^2*(A - B) + 2*a*b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]))/2 - ((2*a*b*(A - B) - a^2*(A + B) + b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/(a*(a^2 + b^2)*d)/a/(2*a*(a^2 + b^2))`

3.595. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

3.595.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2(d/e), 2]\}, \text{Simp}[e/(2cq) \text{Int}[(q - 2x)/\text{Simp}[d/e + qx - x^2, x], x], x] + \text{Simp}[e/(2cq) \text{Int}[(q + 2x)/\text{Simp}[d/e - qx - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c^2d^2 - a^2e^2, 0] \&\& \text{NegQ}[d^2e]$

rule 1482 $\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a^2c, 2]\}, \text{Simp}[(dq + ae)/(2ac) \text{Int}[(q + cx^2)/(a + cx^4), x], x] + \text{Simp}[(dq - ae)/(2ac) \text{Int}[(q - cx^2)/(a + cx^4), x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{NeQ}[c^2d^2 + a^2e^2, 0] \&\& \text{NeQ}[c^2d^2 - a^2e^2, 0] \&\& \text{NegQ}[(a)c]$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4017 $\text{Int}[\frac{(c_.) + (d_.)\tan[(e_.) + (f_.)x]}{\sqrt{(b_.)\tan[(e_.) + (f_.)x] + (c_.) + (d_.)\tan[(e_.) + (f_.)x]}}], x_Symbol] \rightarrow \text{Simp}[2/f \text{Subst}[\text{Int}[(bc + dx^2)/(b^2 + x^4), x], x, \text{Sqrt}[b\tan[e + fx]]], x] /; \text{FreeQ}[\{b, c, d, e, f\}, x] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0]$

rule 4064 $\text{Int}[(\cot[(e_.) + (f_.)x] + (g_.)^p)((a_.) + (b_.)\tan[(e_.) + (f_.)x])^m((c_.) + (d_.)\tan[(e_.) + (f_.)x])^n], x_Symbol] \rightarrow \text{Simp}[g^{m+n} \text{Int}[(g\cot[e + fx])^{p-m-n}(b + a\cot[e + fx])^m(d + c\cot[e + fx])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \&\& \text{IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4088 $\text{Int}[(a_.) + (b_.)\tan[(e_.) + (f_.)x])^m((A_.) + (B_.)\tan[(e_.) + (f_.)x])^n((c_.) + (d_.)\tan[(e_.) + (f_.)x])^n], x_Symbol] \rightarrow \text{Simp}[(b^2c - a^2d)(B^2c - A^2d)(a + b\tan[e + fx])^{m-1}((c + d\tan[e + fx])^{n+1}/(d^2f(n+1)(c^2 + d^2))), x] - \text{Simp}[1/(d^2f(n+1)(c^2 + d^2)) \text{Int}[(a + b\tan[e + fx])^{m-2}(c + d\tan[e + fx])^{n+1} \text{Simp}[a^2d^2(b^2d^2(m-1) - a^2c^2(n+1)) + (b^2B^2c - (A^2b + a^2B^2)d)(b^2c^2(m-1) + a^2d^2(n+1)) - d^2((a^2A - b^2B)(b^2c - a^2d) + (A^2b + a^2B^2)(a^2c + b^2d))(n+1)\tan[e + fx] - b^2(d^2(A^2b^2c + a^2B^2c - a^2A^2d)(m+n) - b^2B^2(c^2(m-1) - d^2(n+1)))\tan[e + fx]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \&\& \text{NeQ}[b^2c - a^2d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{GtQ}[m, 1] \&\& \text{LtQ}[n, -1] \&\& (\text{IntegerQ}[m] \mid \mid \text{IntegersQ}[2m, 2n])]$

3.595.
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

```
rule 4130 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C
*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(GtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.595.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2216 vs. 2(396) = 792.

Time = 0.39 (sec) , antiderivative size = 2217, normalized size of antiderivative = 5.06

method	result	size
derivativedivides	Expression too large to display	2217
default	Expression too large to display	2217

```
input int(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNV
ERBOSE)
```

$$3.595. \quad \int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

output

```
-1/4/d*(1/tan(d*x+c))^(3/2)*tan(d*x+c)*(12*A*(a*b)^(1/2)*tan(d*x+c)*b^5+12
*A*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))*tan(d*x+c)^(3/2)*b^6+16*A*(a*b)^(
1/2)*a^3*b^2+8*A*(a*b)^(1/2)*a*b^4+8*A*(a*b)^(1/2)*a^5+12*A*arctan(b*tan(
d*x+c)^(1/2)/(a*b)^(1/2))*tan(d*x+c)^(1/2)*a*b^5-4*B*(a*b)^(1/2)*tan(d*x+c
)*a^3*b^2-4*B*(a*b)^(1/2)*tan(d*x+c)*a*b^4-20*B*arctan(b*tan(d*x+c)^(1/2)/
(a*b)^(1/2))*tan(d*x+c)^(1/2)*a^4*b^2-4*B*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(
1/2))*tan(d*x+c)^(1/2)*a^2*b^4+8*A*(a*b)^(1/2)*a^4*b*tan(d*x+c)+28*A*arct
an(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))*tan(d*x+c)^(3/2)*a^2*b^4-20*B*arctan(b*
tan(d*x+c)^(1/2)/(a*b)^(1/2))*tan(d*x+c)^(3/2)*a^3*b^3-4*B*arctan(b*tan(d*
x+c)^(1/2)/(a*b)^(1/2))*tan(d*x+c)^(3/2)*a*b^5+20*A*(a*b)^(1/2)*tan(d*x+c
)*a^2*b^3+28*A*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))*tan(d*x+c)^(1/2)*a^3*
b^3+4*A*(a*b)^(1/2)*2^(1/2)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*tan(d*x+c)^(
3/2)*a^3*b^2-2*A*(a*b)^(1/2)*2^(1/2)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*t
an(d*x+c)^(3/2)*a^2*b^3+2*A*(a*b)^(1/2)*2^(1/2)*arctan(-1+2^(1/2)*tan(d*x+
c)^(1/2))*tan(d*x+c)^(3/2)*a^4*b+4*A*(a*b)^(1/2)*2^(1/2)*arctan(-1+2^(1/2)
*tan(d*x+c)^(1/2))*tan(d*x+c)^(3/2)*a^3*b^2+2*A*(a*b)^(1/2)*2^(1/2)*ln(-(1
+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)
-1))*tan(d*x+c)^(1/2)*a^4*b+4*A*(a*b)^(1/2)*2^(1/2)*arctan(1+2^(1/2)*tan(d
*x+c)^(1/2))*tan(d*x+c)^(1/2)*a^4*b-2*A*(a*b)^(1/2)*2^(1/2)*arctan(1+2^(1/
2)*tan(d*x+c)^(1/2))*tan(d*x+c)^(1/2)*a^3*b^2+4*A*(a*b)^(1/2)*2^(1/2)*a...
```

3.595.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5937 vs. $2(397) = 794$.

Time = 37.59 (sec) , antiderivative size = 11904, normalized size of antiderivative = 27.18

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="fracas")`

output Too large to include

3.595. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

3.595.6 Sympy [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \int \frac{(A+B\tan(c+dx))\cot^{\frac{3}{2}}(c+dx)}{(a+b\tan(c+dx))^2} dx$$

input `integrate(cot(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**(3/2)/(a + b*tan(c + d*x))**2, x)`

3.595.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 383, normalized size of antiderivative = 0.87

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \frac{4(5Ba^3b^2-7Aa^2b^3+Bab^4-3Ab^5)\arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right)}{(a^6+2a^4b^2+a^2b^4)\sqrt{ab}} - \frac{2\sqrt{2}((A-B)a^2+2(A+B)ab-(A-B)b^2)\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}+\frac{2}{\sqrt{\tan(dx+c)}}\right)\right)}{(a^6+2a^4b^2+a^2b^4)\sqrt{ab}}$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `-1/4*(4*(5*B*a^3*b^2 - 7*A*a^2*b^3 + B*a*b^4 - 3*A*b^5)*arctan(a/(sqrt(a*b)*sqrt(tan(d*x + c))))/((a^6 + 2*a^4*b^2 + a^2*b^4)*sqrt(a*b)) - (2*sqrt(2)*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))/(a^4 + 2*a^2*b^2 + b^4) - 4*(B*a*b^2 - A*b^3)/((a^4*b + a^2*b^3 + (a^5 + a^3*b^2)/tan(d*x + c))*sqrt(tan(d*x + c))) + 8*A/(a^2*sqrt(tan(d*x + c))))/d`

3.595. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

3.595.8 Giac [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{(b\tan(dx+c)+a)^2} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(3/2)/(b*tan(d*x + c) + a)^2, x)`

3.595.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \int \frac{\cot(c+dx)^{3/2}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

input `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`

output `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^2, x)`

3.596
$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

3.596.1 Optimal result	5665
3.596.2 Mathematica [A] (verified)	5666
3.596.3 Rubi [A] (warning: unable to verify)	5667
3.596.4 Maple [B] (verified)	5674
3.596.5 Fricas [B] (verification not implemented)	5675
3.596.6 Sympy [F]	5675
3.596.7 Maxima [A] (verification not implemented)	5675
3.596.8 Giac [F]	5676
3.596.9 Mupad [F(-1)]	5676

3.596.1 Optimal result

Integrand size = 33, antiderivative size = 392

$$\begin{aligned} & \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\ &= -\frac{(2ab(A-B) - a^2(A+B) + b^2(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^2 d} \\ &+ \frac{(2ab(A-B) - a^2(A+B) + b^2(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^2 d} \\ &- \frac{\sqrt{b}(5a^2Ab + Ab^3 - 3a^3B + ab^2B) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{a^{3/2}(a^2+b^2)^2 d} \\ &+ \frac{b(Ab - aB)\sqrt{\cot(c+dx)}}{a(a^2+b^2)d(b+a \cot(c+dx))} \\ &- \frac{(a^2(A-B) - b^2(A-B) + 2ab(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^2 d} \\ &+ \frac{(a^2(A-B) - b^2(A-B) + 2ab(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^2 d} \end{aligned}$$

output $\frac{1}{2} \cdot (2ab(A-B) - a^2(A+B) + b^2(A+B)) \arctan(-1 + \sqrt{2} \cot(dx+c)^{1/2}) / (a^2+b^2)^{1/2} + \frac{1}{2} \cdot (2ab(A-B) - a^2(A+B) + b^2(A+B)) \arctan(1 + \sqrt{2} \cot(dx+c)^{1/2}) / (a^2+b^2)^{1/2} - \frac{1}{4} \cdot (a^2(A-B) - b^2(A-B) + 2ab(A+B)) \ln(1 + \cot(dx+c)^{1/2}) / (a^2+b^2)^{1/2} + \frac{1}{4} \cdot (a^2(A-B) - b^2(A-B) + 2ab(A+B)) \ln(1 + \cot(dx+c)^{1/2}) / (a^2+b^2)^{1/2} - (5Aa^2b + Ab^3 - 3Ba^3 + B^2a^2b) \arctan(a^{1/2} \cot(dx+c)^{1/2} / b^{1/2}) \cdot b^{1/2} / a^{3/2} / (a^2+b^2)^{1/2} + b(Ab - Ba) \cot(dx+c)^{1/2} / a / (a^2+b^2)^{1/2} / d / (b + a \cot(dx+c))$

3.596.2 Mathematica [A] (verified)

Time = 2.79 (sec) , antiderivative size = 341, normalized size of antiderivative = 0.87

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$$

$$= \frac{\sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)} \left(2\sqrt{2}(2ab(A-B) - a^2(A+B) + b^2(A+B)) \left(\arctan\left(1 - \sqrt{2} \sqrt{\tan(c+dx)}\right) \right) \right)}{4(a^2+b^2)^2 d}$$

input `Integrate[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2, x]`

output $(\text{Sqrt}[\text{Cot}[c + d*x]] * \text{Sqrt}[\text{Tan}[c + d*x]] * (2 * \text{Sqrt}[2] * (2 * a * b * (A - B) - a^2 * (A + B) + b^2 * (A + B)) * (\text{ArcTan}[1 - \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[c + d*x]]] - \text{ArcTan}[1 + \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[c + d*x]]]) + (4 * \text{Sqrt}[b] * (a^2 + b^2) * (A * b - a * B) * \text{ArcTan}[(\text{Sqrt}[b] * \text{Sqrt}[\text{Tan}[c + d*x]]) / \text{Sqrt}[a]]) / a^{3/2} + (8 * \text{Sqrt}[b] * (2 * a * A * b - a^2 * B + b^2 * B) * \text{ArcTan}[(\text{Sqrt}[b] * \text{Sqrt}[\text{Tan}[c + d*x]]) / \text{Sqrt}[a]]) / \text{Sqrt}[a] - \text{Sqrt}[2] * (a^2 * (A - B) + b^2 * (-A + B) + 2 * a * b * (A + B)) * (\text{Log}[1 - \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]] - \text{Log}[1 + \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]]) + (4 * b * (a^2 + b^2) * (A * b - a * B) * \text{Sqrt}[\text{Tan}[c + d*x]]) / (a * (a + b * \text{Tan}[c + d*x]))) / (4 * (a^2 + b^2)^2 * d)$

3.596.3 Rubi [A] (warning: unable to verify)

Time = 1.60 (sec) , antiderivative size = 331, normalized size of antiderivative = 0.84, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.697$, Rules used = {3042, 4064, 3042, 4088, 27, 3042, 4136, 27, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{\cot^{\frac{3}{2}}(c+dx)(A \cot(c+dx)+B)}{(a \cot(c+dx)+b)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{3/2}(B-A \tan(c+dx+\frac{\pi}{2}))}{(b-a \tan(c+dx+\frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{4088} \\
 & \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a \cot(c+dx)+b)} - \frac{\int -\frac{(2Aa^2+bBa+Ab^2)\cot^2(c+dx)-2a(Ab-aB)\cot(c+dx)+b(Ab-aB)}{2\sqrt{\cot(c+dx)}(b+a \cot(c+dx))} dx}{a(a^2+b^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{(2Aa^2+bBa+Ab^2)\cot^2(c+dx)-2a(Ab-aB)\cot(c+dx)+b(Ab-aB)}{\sqrt{\cot(c+dx)}(b+a \cot(c+dx))} dx}{2a(a^2+b^2)} + \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a \cot(c+dx)+b)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{(2Aa^2+bBa+Ab^2)\tan(c+dx+\frac{\pi}{2})^2+2a(Ab-aB)\tan(c+dx+\frac{\pi}{2})+b(Ab-aB)}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx}{2a(a^2+b^2)} + \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a \cot(c+dx)+b)} \\
 & \quad \downarrow \text{4136}
 \end{aligned}$$

$$\frac{\int -\frac{2(a(-Ba^2+2Aba+b^2B)-a(Aa^2+2bBa-Ab^2)\cot(c+dx))}{\sqrt{\cot(c+dx)}}dx + \frac{b(-3a^3B+5a^2Ab+ab^2B+Ab^3)\int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)(b+a\cot(c+dx))}}dx}{a^2+b^2}}{a^2+b^2} +$$

$$\frac{2a(a^2+b^2)}{ad(a^2+b^2)(a\cot(c+dx)+b)} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a\cot(c+dx)+b)}$$

27

$$\frac{b(-3a^3B+5a^2Ab+ab^2B+Ab^3)\int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)(b+a\cot(c+dx))}}dx}{a^2+b^2} - \frac{2\int \frac{a(-Ba^2+2Aba+b^2B)-a(Aa^2+2bBa-Ab^2)\cot(c+dx)}{\sqrt{\cot(c+dx)}}dx}{a^2+b^2}}{a^2+b^2} +$$

$$\frac{2a(a^2+b^2)}{ad(a^2+b^2)(a\cot(c+dx)+b)} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a\cot(c+dx)+b)}$$

3042

$$\frac{b(-3a^3B+5a^2Ab+ab^2B+Ab^3)\int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}}dx}{a^2+b^2} - \frac{2\int \frac{a(-Ba^2+2Aba+b^2B)+a(Aa^2+2bBa-Ab^2)\tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}}dx}{a^2+b^2}}{a^2+b^2} +$$

$$\frac{2a(a^2+b^2)}{ad(a^2+b^2)(a\cot(c+dx)+b)} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a\cot(c+dx)+b)}$$

4017

$$\frac{b(-3a^3B+5a^2Ab+ab^2B+Ab^3)\int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}}dx}{a^2+b^2} - \frac{4\int \frac{a(-Ba^2+2Aba+b^2B-(Aa^2+2bBa-Ab^2)\cot(c+dx))}{\cot^2(c+dx)+1}d\sqrt{\cot(c+dx)}}{d(a^2+b^2)}}{a^2+b^2} +$$

$$\frac{2a(a^2+b^2)}{ad(a^2+b^2)(a\cot(c+dx)+b)} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a\cot(c+dx)+b)}$$

25

$$\frac{4\int \frac{a(-Ba^2+2Aba+b^2B-(Aa^2+2bBa-Ab^2)\cot(c+dx))}{\cot^2(c+dx)+1}d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} + \frac{b(-3a^3B+5a^2Ab+ab^2B+Ab^3)\int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}}dx}{a^2+b^2}}{a^2+b^2} +$$

$$\frac{2a(a^2+b^2)}{ad(a^2+b^2)(a\cot(c+dx)+b)} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a\cot(c+dx)+b)}$$

27

3.596. $\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

$$\frac{4a \int \frac{-Ba^2 + 2Aba + b^2B - (Aa^2 + 2bBa - Ab^2) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{b(-3a^3B + 5a^2Ab + ab^2B + Ab^3) \int \frac{\tan(c+dx + \frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c+dx + \frac{\pi}{2})(b-a \tan(c+dx + \frac{\pi}{2}))}} dx}{a^2+b^2}}{d(a^2+b^2)} + \frac{2a(a^2+b^2)}{ad(a^2+b^2)(a \cot(c+dx)+b)} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a \cot(c+dx)+b)}$$

↓ 1482

$$\frac{4a \left(\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(-a^2(A+B) + 2ab(A-B) + b^2(A+B)) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d(a^2+b^2)} + \frac{2a(a^2+b^2)}{ad(a^2+b^2)(a \cot(c+dx)+b)} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a \cot(c+dx)+b)}$$

↓ 1476

$$\frac{4a \left(\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(-a^2(A+B) + 2ab(A-B) + b^2(A+B)) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1}} \right) \right)}{d(a^2+b^2)} + \frac{2a(a^2+b^2)}{ad(a^2+b^2)(a \cot(c+dx)+b)} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a \cot(c+dx)+b)}$$

↓ 1082

$$\frac{4a \left(\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(-a^2(A+B) + 2ab(A-B) + b^2(A+B)) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d(a^2+b^2)} + \frac{2a(a^2+b^2)}{ad(a^2+b^2)(a \cot(c+dx)+b)} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a \cot(c+dx)+b)}$$

↓ 217

$$\frac{4a \left(\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2}(-a^2(A+B) + 2ab(A-B) + b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}) \right) \right)}{d(a^2+b^2)} + \frac{2a(a^2+b^2)}{ad(a^2+b^2)(a \cot(c+dx)+b)} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a \cot(c+dx)+b)}$$

↓ 1479

3.596. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$4a \left(\frac{\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \right)}{d(a^2+b^2)} + \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B))$$

$$\frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 25

$$4a \left(\frac{\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \right)}{d(a^2+b^2)} + \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B))$$

$$\frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 27

$$4a \left(\frac{\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \right)}{d(a^2+b^2)} + \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B))$$

$$\frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 1103

$$\frac{b(-3a^3B+5a^2Ab+ab^2B+Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} + \frac{4a \left(\frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B) \right) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right)}{2a(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 4117

$$\frac{b(-3a^3B+5a^2Ab+ab^2B+Ab^3) \int \frac{1}{\sqrt{\cot(c+dx)(b+a\cot(c+dx))}} d(-\cot(c+dx))}{d(a^2+b^2)} + \frac{4a \left(\frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B) \right) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right)}{2a(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{ad(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 73

3.596. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

$$\frac{4a \left(\frac{1}{2} (-a^2(A+B)) + 2ab(A-B) + b^2(A+B) \right) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)}+1}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) + \frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\log\left(\frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a\cot(c+dx)+b)}\right)}{d(a^2+b^2)} \right)}{2a(a^2+b^2)}$$

$\frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a\cot(c+dx)+b)}$
 \downarrow 218
 $\frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a\cot(c+dx)+b)} +$

$$\frac{4a \left(\frac{1}{2} (-a^2(A+B)) + 2ab(A-B) + b^2(A+B) \right) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)}+1}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) + \frac{1}{2} (a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\log\left(\frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a\cot(c+dx)+b)}\right)}{d(a^2+b^2)} \right)}{2a(a^2+b^2)}$$

input `Int[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^2,x]`

output `(b*(A*b - a*B)*Sqrt[Cot[c + d*x]]/(a*(a^2 + b^2)*d*(b + a*Cot[c + d*x])) + ((2*Sqrt[b]*(5*a^2*A*b + A*b^3 - 3*a^3*B + a*b^2*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]])/(Sqrt[a]*(a^2 + b^2)*d) + (4*a*((2*a*b*(A - B) - a^2*(A + B) + b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]/Sqrt[2]] + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]/Sqrt[2]]))/2 + ((a^2*(A - B) - b^2*(A - B) + 2*a*b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/(a^2 + b^2)*d)/(2*a*(a^2 + b^2))`

3.596.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

3.596. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^2} dx$

- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 $\text{Int}[\frac{(c_.) + (d_.)\tan[e_.] + (f_.)x_]}{\sqrt{(b_.)\tan[e_.] + (f_.)x_}}], x_{\text{Symbol}}] \rightarrow \text{Simp}[2/f \text{ Subst}[\text{Int}[(b*c + d*x^2)/(b^2 + x^4), x], x, \sqrt{b*\tan[e + f*x]}], x] /; \text{FreeQ}\{b, c, d, e, f\}, x] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0]$

rule 4064 $\text{Int}[(\cot[e_.] + (f_.)x_]*(g_.)^{(p_.)}*((a_.) + (b_.)\tan[e_.] + (f_.)x_))^{(m_.)}*((c_.) + (d_.)\tan[e_.] + (f_.)x_)]^{(n_.)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[g^{(m+n)} \text{ Int}[(g*\cot[e + f*x])^{(p-m-n)}*(b + a*\cot[e + f*x])^m*(d + c*\cot[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, p\}, x] \&\& \text{!IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4088 $\text{Int}[(a_.) + (b_.)\tan[e_.] + (f_.)x_)]^{(m_.)}*((A_.) + (B_.)\tan[e_.] + (f_.)x_)]^{(n_.)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(b*c - a*d)*(B*c - A*d)*(a + b*\tan[e + f*x])^{(m-1)}*((c + d*\tan[e + f*x])^{(n+1)})/(d*f*(n+1)*(c^2 + d^2)), x] - \text{Simp}[1/(d*(n+1)*(c^2 + d^2)) \text{ Int}[(a + b*\tan[e + f*x])^{(m-2)}*(c + d*\tan[e + f*x])^{(n+1)}*\text{Simp}[a*A*d*(b*d*(m-1) - a*c*(n+1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m-1) + a*d*(n+1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n+1)*\tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m+n) - b*B*(c^2*(m-1) - d^2*(n+1)))*\tan[e + f*x]^2, x], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{GtQ}[m, 1] \&\& \text{LtQ}[n, -1] \&\& (\text{IntegerQ}[m] \mid\mid \text{IntegersQ}[2*m, 2*n])$

rule 4117 $\text{Int}[(a_.) + (b_.)\tan[e_.] + (f_.)x_)]^{(m_.)}*((c_.) + (d_.)\tan[e_.] + (f_.)x_)]^{(n_.)}*((A_.) + (C_.)\tan[e_.] + (f_.)x_)]^2, x_{\text{Symbol}}] \rightarrow \text{Simp}[A/f \text{ Subst}[\text{Int}[(a + b*x)^m*(c + d*x)^n, x], x, \tan[e + f*x]], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, C, m, n\}, x] \&\& \text{EqQ}[A, C]$

rule 4136 $\text{Int}[(c_.) + (d_.)\tan[e_.] + (f_.)x_)]^{(n_.)}*((A_.) + (B_.)\tan[e_.] + (f_.)x_)] + (C_.)\tan[e_.] + (f_.)x_)]^2)/((a_.) + (b_.)\tan[e_.] + (f_.)x_)), x_{\text{Symbol}}] \rightarrow \text{Simp}[1/(a^2 + b^2) \text{ Int}[(c + d*\tan[e + f*x])^n*\text{Simp}[b*B + a*(A - C) + (a*B - b*(A - C))*\tan[e + f*x], x], x], x] + \text{Simp}[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) \text{ Int}[(c + d*\tan[e + f*x])^n*((1 + \tan[e + f*x]^2)/(a + b*\tan[e + f*x])), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, C, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{!GtQ}[n, 0] \&\& \text{!LeQ}[n, -1]$

3.596.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1935 vs. $2(352) = 704$.

Time = 0.42 (sec) , antiderivative size = 1936, normalized size of antiderivative = 4.94

method	result	size
derivativdivides	Expression too large to display	1936
default	Expression too large to display	1936

```
input int(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x,method=_RETURNV
ERBOSE)
```

```
output 1/4/d*(1/tan(d*x+c))^(1/2)*tan(d*x+c)^(1/2)*(4*A*arctan(b*tan(d*x+c)^(1/2)
/(a*b)^(1/2))*b^5*tan(d*x+c)+4*A*(a*b)^(1/2)*tan(d*x+c)^(1/2)*b^4+20*A*arc
tan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))*a^3*b^2+4*A*arctan(b*tan(d*x+c)^(1/2)/
(a*b)^(1/2))*a*b^4-12*B*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))*a^4*b+4*B*a
rctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))*a^2*b^3+2*B*2^(1/2)*(a*b)^(1/2)*arct
an(1+2^(1/2)*tan(d*x+c)^(1/2))*a^4+2*B*2^(1/2)*(a*b)^(1/2)*arctan(-1+2^(1/
2)*tan(d*x+c)^(1/2))*a^4+B*2^(1/2)*(a*b)^(1/2)*ln(-(2^(1/2)*tan(d*x+c)^(1/
2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c)))*a^4-12*B*arctan(
b*tan(d*x+c)^(1/2)/(a*b)^(1/2))*a^3*b^2*tan(d*x+c)+4*B*arctan(b*tan(d*x+c)
^(1/2)/(a*b)^(1/2))*a*b^4*tan(d*x+c)+4*A*(a*b)^(1/2)*tan(d*x+c)^(1/2)*a^2*
b^2-4*B*(a*b)^(1/2)*tan(d*x+c)^(1/2)*a^3*b-4*B*(a*b)^(1/2)*tan(d*x+c)^(1/2
)*a*b^3+A*2^(1/2)*(a*b)^(1/2)*ln(-(1+2^(1/2)*tan(d*x+c)^(1/2)+tan(d*x+c))/
(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1))*a^4+2*A*2^(1/2)*(a*b)^(1/2)*arcta
n(1+2^(1/2)*tan(d*x+c)^(1/2))*a^4+2*A*2^(1/2)*(a*b)^(1/2)*arctan(-1+2^(1/2
)*tan(d*x+c)^(1/2))*a^4+20*A*arctan(b*tan(d*x+c)^(1/2)/(a*b)^(1/2))*a^2*b^
3*tan(d*x+c)-2*B*2^(1/2)*(a*b)^(1/2)*arctan(1+2^(1/2)*tan(d*x+c)^(1/2))*a^
2*b^2+4*B*2^(1/2)*(a*b)^(1/2)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*a^3*b-2*
B*2^(1/2)*(a*b)^(1/2)*arctan(-1+2^(1/2)*tan(d*x+c)^(1/2))*a^2*b^2-B*2^(1/2
)*(a*b)^(1/2)*ln(-(2^(1/2)*tan(d*x+c)^(1/2)-tan(d*x+c)-1)/(1+2^(1/2)*tan(d
*x+c)^(1/2)+tan(d*x+c)))*a^2*b^2-4*A*2^(1/2)*(a*b)^(1/2)*arctan(-1+2^(1...
```

3.596.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5899 vs. $2(354) = 708$.

Time = 23.69 (sec) , antiderivative size = 11827, normalized size of antiderivative = 30.17

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm m="fracas")`

output Too large to include

3.596.6 Sympy [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx = \int \frac{(A+B\tan(c+dx))\sqrt{\cot(c+dx)}}{(a+b\tan(c+dx))^2} dx$$

input `integrate(cot(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**2,x)`

output `Integral((A + B*tan(c + d*x))*sqrt(cot(c + d*x))/(a + b*tan(c + d*x))**2, x)`

3.596.7 Maxima [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 362, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$$

$$= \frac{4(3Ba^3b-5Aa^2b^2-Bab^3-Ab^4)\arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right)}{(a^5+2a^3b^2+ab^4)\sqrt{ab}} - \frac{2\sqrt{2}((A+B)a^2-2(A-B)ab-(A+B)b^2)\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}+\frac{2}{\sqrt{\tan(dx+c)}}\right)\right)}{(a^5+2a^3b^2+ab^4)\sqrt{ab}} + 2$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm m="maxima")`

3.596. $\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^2} dx$

output $\frac{1}{4} \cdot (4 \cdot (3B \cdot a^3 \cdot b - 5A \cdot a^2 \cdot b^2 - B \cdot a \cdot b^3 - A \cdot b^4) \cdot \arctan(a / (\sqrt{a \cdot b}) \cdot \sqrt{\tan(dx + c)})) / ((a^5 + 2a^3 \cdot b^2 + a \cdot b^4) \cdot \sqrt{a \cdot b}) - (2 \cdot \sqrt{2}) \cdot ((A + B) \cdot a^2 - 2 \cdot (A - B) \cdot a \cdot b - (A + B) \cdot b^2) \cdot \arctan(1/2 \cdot \sqrt{2} \cdot (\sqrt{2} + 2 / \sqrt{\tan(dx + c)})) + 2 \cdot \sqrt{2} \cdot ((A + B) \cdot a^2 - 2 \cdot (A - B) \cdot a \cdot b - (A + B) \cdot b^2) \cdot \arctan(-1/2 \cdot \sqrt{2} \cdot (\sqrt{2} - 2 / \sqrt{\tan(dx + c)})) - \sqrt{2} \cdot ((A - B) \cdot a^2 + 2 \cdot (A + B) \cdot a \cdot b - (A - B) \cdot b^2) \cdot \log(\sqrt{2} / \sqrt{\tan(dx + c)}) + 1 / \tan(dx + c) + 1 + \sqrt{2} \cdot ((A - B) \cdot a^2 + 2 \cdot (A + B) \cdot a \cdot b - (A - B) \cdot b^2) \cdot \log(-\sqrt{2} / \sqrt{\tan(dx + c)}) + 1 / \tan(dx + c) + 1) / (a^4 + 2a^2 \cdot b^2 + b^4) - 4 \cdot (B \cdot a \cdot b - A \cdot b^2) / ((a^3 \cdot b + a \cdot b^3 + (a^4 + a^2 \cdot b^2) / \tan(dx + c)) \cdot \sqrt{\tan(dx + c)}) / d$

3.596.8 Giac [F]

$$\int \frac{\sqrt{\cot(c + dx)}(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx = \int \frac{(B \tan(dx + c) + A) \sqrt{\cot(dx + c)}}{(b \tan(dx + c) + a)^2} dx$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(cot(d*x + c))/(b*tan(d*x + c) + a)^2, x)`

3.596.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cot(c + dx)}(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx = \int \frac{\sqrt{\cot(c + dx)}(A + B \tan(c + dx))}{(a + b \tan(c + dx))^2} dx$$

input `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^2,x)`

output `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^2, x)`

$$3.597 \quad \int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))^2}} dx$$

3.597.1 Optimal result	5677
3.597.2 Mathematica [A] (verified)	5678
3.597.3 Rubi [A] (warning: unable to verify)	5679
3.597.4 Maple [A] (verified)	5686
3.597.5 Fricas [B] (verification not implemented)	5686
3.597.6 Sympy [F]	5687
3.597.7 Maxima [A] (verification not implemented)	5687
3.597.8 Giac [F]	5688
3.597.9 Mupad [F(-1)]	5688

3.597.1 Optimal result

Integrand size = 33, antiderivative size = 390

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))^2}} dx \\ &= \frac{(a^2(A - B) - b^2(A - B) + 2ab(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^2 d} \\ & \quad - \frac{(a^2(A - B) - b^2(A - B) + 2ab(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^2 d} \\ & \quad + \frac{(3a^2 Ab - Ab^3 - a^3 B + 3ab^2 B) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{\sqrt{a}\sqrt{b}(a^2 + b^2)^2 d} - \frac{(Ab - aB)\sqrt{\cot(c + dx)}}{(a^2 + b^2) d(b + a \cot(c + dx))} \\ & \quad - \frac{(2ab(A - B) - a^2(A + B) + b^2(A + B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^2 d} \\ & \quad + \frac{(2ab(A - B) - a^2(A + B) + b^2(A + B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^2 d} \end{aligned}$$

3.597. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))^2}} dx$

output
$$\begin{aligned} & -1/2*(a^2*(A-B)-b^2*(A-B)+2*a*b*(A+B))*\arctan(-1+2^{(1/2)}*\cot(d*x+c)^{(1/2)}) \\ & / (a^2+b^2)^{2/d*2^{(1/2)}}-1/2*(a^2*(A-B)-b^2*(A-B)+2*a*b*(A+B))*\arctan(1+2^{(1/2)} \\ & *\cot(d*x+c)^{(1/2)})/(a^2+b^2)^{2/d*2^{(1/2)}}-1/4*(2*a*b*(A-B)-a^2*(A+B)+b^2 \\ & *(A+B))*\ln(1+\cot(d*x+c)-2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)^{2/d*2^{(1/2)}}+1/ \\ & 4*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\ln(1+\cot(d*x+c)+2^{(1/2)}*\cot(d*x+c)^{(1/2)}) \\ & / (a^2+b^2)^{2/d*2^{(1/2)}}+(3*A*a^2*b-A*b^3-B*a^3+3*B*a*b^2)*\arctan(a^{(1/2)} \\ & *\cot(d*x+c)^{(1/2)}/b^{(1/2)})/(a^2+b^2)^{2/d/a^{(1/2)}/b^{(1/2)}}-(A*b-B*a)*\cot(d*x \\ & +c)^{(1/2)}/(a^2+b^2)/d/(b+a*\cot(d*x+c)) \end{aligned}$$

3.597.2 Mathematica [A] (verified)

Time = 3.02 (sec) , antiderivative size = 336, normalized size of antiderivative = 0.86

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))^2}} dx = \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(2\sqrt{2}(a^2(A - B) + b^2(-A + B) + 2ab(A + B)) \left(\arctan \left(1 - \sqrt{2} \sqrt{\tan(c + dx)} \right) \right) \right)}{\dots}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^2),x]`

output
$$\begin{aligned} & -1/4*(\text{Sqrt}[\text{Cot}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]*(2*\text{Sqrt}[2]*(a^2*(A - B) + b^2* \\ & (-A + B) + 2*a*b*(A + B))*(\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]] - \text{ArcTan} \\ & [1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]]) - (4*(a^2 + b^2)*(-(A*b) + a*B)*\text{ArcTan}[(\\ & \text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(\text{Sqrt}[a]*\text{Sqrt}[b]) + (8*\text{Sqrt}[b]*(a^2* \\ & A - A*b^2 + 2*a*b*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/\text{Sqrt}[a] \\ & + \text{Sqrt}[2]*(2*a*b*(A - B) - a^2*(A + B) + b^2*(A + B))*(\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt} \\ & [\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]] - \text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan} \\ & [c + d*x]]) - (4*(a^2 + b^2)*(-(A*b) + a*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/(a + b*\text{Tan} \\ & [c + d*x]))/((a^2 + b^2)^2*d) \end{aligned}$$

3.597.3 Rubi [A] (warning: unable to verify)

Time = 1.64 (sec) , antiderivative size = 330, normalized size of antiderivative = 0.85, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.697$, Rules used = {3042, 4064, 3042, 4091, 27, 3042, 4136, 27, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))^2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))^2}} dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{\sqrt{\cot(c + dx)}(A \cot(c + dx) + B)}{(a \cot(c + dx) + b)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(B - A \tan(c + dx + \frac{\pi}{2}))}{(b - a \tan(c + dx + \frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{4091} \\
 & \frac{\int -\frac{a(Ab - aB) \cot^2(c + dx) - 2a(aA + bB) \cot(c + dx) + a(Ab - aB)}{2\sqrt{\cot(c + dx)}(b + a \cot(c + dx))} dx}{a(a^2 + b^2)} - \frac{(Ab - aB)\sqrt{\cot(c + dx)}}{d(a^2 + b^2)(a \cot(c + dx) + b)} \\
 & \quad \downarrow \text{27} \\
 & -\frac{\int \frac{-a(Ab - aB) \cot^2(c + dx) - 2a(aA + bB) \cot(c + dx) + a(Ab - aB)}{\sqrt{\cot(c + dx)}(b + a \cot(c + dx))} dx}{2a(a^2 + b^2)} - \frac{(Ab - aB)\sqrt{\cot(c + dx)}}{d(a^2 + b^2)(a \cot(c + dx) + b)} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\int \frac{-a(Ab - aB) \tan(c + dx + \frac{\pi}{2})^2 + 2a(aA + bB) \tan(c + dx + \frac{\pi}{2}) + a(Ab - aB)}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(b - a \tan(c + dx + \frac{\pi}{2}))} dx}{2a(a^2 + b^2)} - \frac{(Ab - aB)\sqrt{\cot(c + dx)}}{d(a^2 + b^2)(a \cot(c + dx) + b)} \\
 & \quad \downarrow \text{4136}
 \end{aligned}$$

3.597. $\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))^2}} dx$

$$\frac{\int -\frac{2(a(Aa^2+2bBa-Ab^2)+a(-Ba^2+2Aba+b^2B)\cot(c+dx))}{\sqrt{\cot(c+dx)}(a^2+b^2)}dx + \frac{a(a^3(-B)+3a^2Ab+3ab^2B-Ab^3)}{a^2+b^2} \int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)(b+a\cot(c+dx))}}dx}{\frac{2a(a^2+b^2)(Ab-aB)\sqrt{\cot(c+dx)}}{d(a^2+b^2)(a\cot(c+dx)+b)}} \quad \downarrow \quad 27$$

$$\frac{a(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)(b+a\cot(c+dx))}}dx - 2 \int \frac{a(Aa^2+2bBa-Ab^2)+a(-Ba^2+2Aba+b^2B)\cot(c+dx)}{\sqrt{\cot(c+dx)}(a^2+b^2)}dx}{\frac{2a(a^2+b^2)(Ab-aB)\sqrt{\cot(c+dx)}}{d(a^2+b^2)(a\cot(c+dx)+b)}} \quad \downarrow \quad 3042$$

$$\frac{a(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}}dx - 2 \int \frac{a(Aa^2+2bBa-Ab^2)-a(-Ba^2+2Aba+b^2B)\tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}}dx}{\frac{2a(a^2+b^2)(Ab-aB)\sqrt{\cot(c+dx)}}{d(a^2+b^2)(a\cot(c+dx)+b)}} \quad \downarrow \quad 4017$$

$$\frac{a(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}}dx - 4 \int -\frac{a(Aa^2+2bBa-Ab^2+(-Ba^2+2Aba+b^2B)\cot(c+dx))}{\cot^2(c+dx)+1}d\sqrt{\cot(c+dx)}}{\frac{2a(a^2+b^2)(Ab-aB)\sqrt{\cot(c+dx)}}{d(a^2+b^2)(a\cot(c+dx)+b)}} \quad \downarrow \quad 25$$

$$\frac{4 \int \frac{a(Aa^2+2bBa-Ab^2+(-Ba^2+2Aba+b^2B)\cot(c+dx))}{\cot^2(c+dx)+1}d\sqrt{\cot(c+dx)} + \frac{a(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}}dx}{a^2+b^2}}{\frac{2a(a^2+b^2)(Ab-aB)\sqrt{\cot(c+dx)}}{d(a^2+b^2)(a\cot(c+dx)+b)}} \quad \downarrow \quad 27$$

3.597. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))^2}} dx$

$$\frac{4a \int \frac{Aa^2+2bBa-Ab^2+(-Ba^2+2Aba+b^2B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{a(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))} dx}{a^2+b^2}}{d(a^2+b^2)}$$

$$\frac{2a(a^2+b^2)(Ab-aB)\sqrt{\cot(c+dx)}}{d(a^2+b^2)(a \cot(c+dx)+b)}$$

↓ 1482

$$\frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d(a^2+b^2)}$$

$$\frac{2a(a^2+b^2)(Ab-aB)\sqrt{\cot(c+dx)}}{d(a^2+b^2)(a \cot(c+dx)+b)}$$

↓ 1476

$$\frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) - \frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d(a^2+b^2)}$$

$$\frac{2a(a^2+b^2)(Ab-aB)\sqrt{\cot(c+dx)}}{d(a^2+b^2)(a \cot(c+dx)+b)}$$

↓ 1082

$$\frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)-1} d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d(a^2+b^2)}$$

$$\frac{2a(a^2+b^2)(Ab-aB)\sqrt{\cot(c+dx)}}{d(a^2+b^2)(a \cot(c+dx)+b)}$$

↓ 217

$$\frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d(a^2+b^2)}$$

$$\frac{2a(a^2+b^2)(Ab-aB)\sqrt{\cot(c+dx)}}{d(a^2+b^2)(a \cot(c+dx)+b)}$$

↓ 1479

3.597. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))^2}} dx$

$$\frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \right)}{d(a^2+b^2)}$$

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{d(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 25

$$\frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \right)}{d(a^2+b^2)}$$

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{d(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 27

$$\frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \right)}{d(a^2+b^2)}$$

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{d(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 1103

$$\frac{a(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} + \frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \right)}{d(a^2+b^2)}$$

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{d(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 4117

$$\frac{a(a^3(-B)+3a^2Ab+3ab^2B-Ab^3) \int \frac{1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} d(-\cot(c+dx))}{d(a^2+b^2)} + \frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \right)}{d(a^2+b^2)}$$

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{d(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 73

3.597. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))^2}} dx$

$$\frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \right)}{d(a^2+b^2)}$$

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{d(a^2 + b^2)(a \cot(c + dx) + b)}$$

↓ 218

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{d(a^2 + b^2)(a \cot(c + dx) + b)} - \frac{4a \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(-a^2(A+B)+2ab(A-B)+b^2(A+B)) \right)}{d(a^2+b^2)}$$

$$2a(a^2 + b^2)$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^2),x]`

output `-(((A*b - a*B)*Sqrt[Cot[c + d*x]])/((a^2 + b^2)*d*(b + a*Cot[c + d*x]))) - ((2*Sqrt[a]*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]])/(Sqrt[b]*(a^2 + b^2)*d) + (4*a*((a^2*(A - B) - b^2*(A - B) + 2*a*b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]/Sqrt[2]] + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]/Sqrt[2]]))/2 - ((2*a*b*(A - B) - a^2*(A + B) + b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)/(2*a*(a^2 + b^2))`

3.597.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

3.597. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))^2}} dx$

rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & (LtQ[a, 0] || LtQ[b, 0])`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4091 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4136 `Int((((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]) + (C_)*tan[(e_) + (f_)*(x_)])^2)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.597.4 Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 332, normalized size of antiderivative = 0.85

method	result
derivativedivides	$\frac{2 \left((A a^2 - A b^2 + 2 B a b) \sqrt{2} \left(\ln \left(\frac{1 + \cot(dx+c) + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c) - \sqrt{2} \sqrt{\cot(dx+c)}} \right) + 2 \arctan(1 + \sqrt{2} \sqrt{\cot(dx+c)}) + 2 \arctan(-1 + \sqrt{2} \sqrt{\cot(dx+c)}) \right) \right)}{8}$
default	$\frac{2 \left((A a^2 - A b^2 + 2 B a b) \sqrt{2} \left(\ln \left(\frac{1 + \cot(dx+c) + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c) - \sqrt{2} \sqrt{\cot(dx+c)}} \right) + 2 \arctan(1 + \sqrt{2} \sqrt{\cot(dx+c)}) + 2 \arctan(-1 + \sqrt{2} \sqrt{\cot(dx+c)}) \right) \right)}{8}$

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^2,x,method=_RETURNV
ERBOSE)`

output `1/d*(-2/(a^2+b^2)^2*(1/8*(A*a^2-A*b^2+2*B*a*b)*2^(1/2)*(ln((1+cot(d*x+c)+2
^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan
(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/8*(2
*A*a*b-B*a^2+B*b^2)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1
+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2
) + 2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))))+2/(a^2+b^2)^2*((-1/2*A*a^2*b-1/2
*A*b^3+1/2*B*a^3+1/2*B*a*b^2)*cot(d*x+c)^(1/2)/(b+a*cot(d*x+c))+1/2*(3*A*a
^2*b-A*b^3-B*a^3+3*B*a*b^2)/(a*b)^(1/2)*arctan(a*cot(d*x+c)^(1/2)/(a*b)^(1
/2))))`

3.597.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5901 vs. 2(348) = 696.

Time = 14.68 (sec) , antiderivative size = 11827, normalized size of antiderivative = 30.33

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))^2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^2,x, algorith
m="fracas")`

output Too large to include

3.597.
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))^2}} dx$$

3.597.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^2} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^2 \sqrt{\cot(c + dx)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(1/2)/(a+b*tan(d*x+c))**2,x)`

output `Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))**2*sqrt(cot(c + d*x))), x)`

3.597.7 Maxima [A] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 347, normalized size of antiderivative = 0.89

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^2} dx = \frac{4(Ba^3 - 3Aa^2b - 3Bab^2 + Ab^3) \arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right)}{(a^4 + 2a^2b^2 + b^4)\sqrt{ab}} + \frac{2\sqrt{2}((A-B)a^2 + 2(A+B)ab - (A-B)b^2) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right)}{(a^4 + 2a^2b^2 + b^4)\sqrt{ab}} + \dots$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^2,x, algorithm="maxima")`

output `-1/4*(4*(B*a^3 - 3*A*a^2*b - 3*B*a*b^2 + A*b^3)*arctan(a/(sqrt(a*b)*sqrt(tan(d*x + c))))/((a^4 + 2*a^2*b^2 + b^4)*sqrt(a*b)) + (2*sqrt(2))*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1)/((a^4 + 2*a^2*b^2 + b^4) - 4*(B*a - A*b)/((a^2*b + b^3 + (a^3 + a*b^2)/tan(d*x + c))*sqrt(tan(d*x + c))))/d`

3.597.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^2} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a)^2 \sqrt{\cot(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)/((b*tan(d*x + c) + a)^2*sqrt(cot(d*x + c))), x)`

3.597.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^2} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^2} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))^2),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))^2), x)`

$$3.598 \quad \int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$$

3.598.1 Optimal result	5689
3.598.2 Mathematica [A] (verified)	5690
3.598.3 Rubi [A] (warning: unable to verify)	5691
3.598.4 Maple [A] (verified)	5698
3.598.5 Fricas [B] (verification not implemented)	5698
3.598.6 Sympy [F]	5699
3.598.7 Maxima [A] (verification not implemented)	5699
3.598.8 Giac [F]	5700
3.598.9 Mupad [F(-1)]	5700

3.598.1 Optimal result

Integrand size = 33, antiderivative size = 392

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx \\ &= \frac{(2ab(A - B) - a^2(A + B) + b^2(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^2 d} \\ & \quad - \frac{(2ab(A - B) - a^2(A + B) + b^2(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^2 d} \\ & \quad - \frac{\sqrt{a}(a^2 Ab - 3Ab^3 + a^3 B + 5ab^2 B) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{b^{3/2}(a^2 + b^2)^2 d} \\ & \quad + \frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{b(a^2 + b^2)d(b + a \cot(c + dx))} \\ & \quad + \frac{(a^2(A - B) - b^2(A - B) + 2ab(A + B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^2 d} \\ & \quad - \frac{(a^2(A - B) - b^2(A - B) + 2ab(A + B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^2 d} \end{aligned}$$

output
$$\begin{aligned} & -1/2*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\arctan(-1+2^{(1/2)}*\cot(d*x+c)^{(1/2)}) \\ & / (a^2+b^2)^2/d*2^{(1/2)}-1/2*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\arctan(1+2^{(1/2)} \\ & /2)*\cot(d*x+c)^{(1/2)})/(a^2+b^2)^2/d*2^{(1/2)}+1/4*(a^2*(A-B)-b^2*(A-B)+2*a*b \\ & *(A+B))*\ln(1+\cot(d*x+c)-2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)^2/d*2^{(1/2)}-1/ \\ & 4*(a^2*(A-B)-b^2*(A-B)+2*a*b*(A+B))*\ln(1+\cot(d*x+c)+2^{(1/2)}*\cot(d*x+c)^{(1/2)}) \\ & / (a^2+b^2)^2/d*2^{(1/2)}-(A*a^2*b-3*A*b^3+B*a^3+5*B*a*b^2))*\arctan(a^{(1/2)} \\ & *\cot(d*x+c)^{(1/2)}/b^{(1/2)})*a^{(1/2)}/b^{(3/2)}/(a^2+b^2)^2/d+a*(A*b-B*a)*\cot(d \\ & *x+c)^{(1/2)}/b/(a^2+b^2)/d/(b+a*\cot(d*x+c)) \end{aligned}$$

3.598.2 Mathematica [A] (verified)

Time = 2.68 (sec) , antiderivative size = 342, normalized size of antiderivative = 0.87

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$$

$$= \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(-2\sqrt{2}(2ab(A - B) - a^2(A + B) + b^2(A + B)) \left(\arctan \left(1 - \sqrt{2} \sqrt{\tan(c + dx)} \right) \right) \right)}{\dots}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^2),x]`

output
$$\begin{aligned} & (\text{Sqrt}[\text{Cot}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]*(-2*\text{Sqrt}[2]*(2*a*b*(A - B) - a^2*(A \\ & + B) + b^2*(A + B))*(\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]] - \text{ArcTan}[1 + \\ & \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]]) + (4*\text{Sqrt}[a]*(a^2 + b^2)*(A*b - a*B)*\text{ArcTan}[(\\ & \text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/b^{(3/2)} + (8*\text{Sqrt}[a]*(-2*A*b^3 + a*(\\ & a^2 + 3*b^2)*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/b^{(3/2)} + \text{Sqrt}[2]*(a^2*(A - B) + b^2*(-A + B) + 2*a*b*(A + B))*(\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]] - \text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]]) + (4*a*(a^2 + b^2)*(A*b - a*B)*\text{Sqrt}[\text{Tan}[c + d*x]]/(b*(a + b*\text{Tan}[c + d*x]))))/ (4*(a^2 + b^2)^2*d) \end{aligned}$$

3.598.3 Rubi [A] (warning: unable to verify)

Time = 1.69 (sec) , antiderivative size = 331, normalized size of antiderivative = 0.84, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.697$, Rules used = {3042, 4064, 3042, 4092, 27, 3042, 4136, 27, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2}(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{A \cot(c + dx) + B}{\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B - A \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(b - a \tan(c + dx + \frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{4092} \\
 & \frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{bd(a^2 + b^2)(a \cot(c + dx) + b)} - \frac{\int -\frac{Ba^2 - (Ab - aB)\cot^2(c + dx)a + Aba + 2b^2B + 2b(Ab - aB)\cot(c + dx)}{2\sqrt{\cot(c + dx)}(b + a \cot(c + dx))} dx}{b(a^2 + b^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{Ba^2 - (Ab - aB)\cot^2(c + dx)a + Aba + 2b^2B + 2b(Ab - aB)\cot(c + dx)}{\sqrt{\cot(c + dx)}(b + a \cot(c + dx))} dx}{2b(a^2 + b^2)} + \frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{bd(a^2 + b^2)(a \cot(c + dx) + b)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{Ba^2 - (Ab - aB)\tan(c + dx + \frac{\pi}{2})^2 a + Aba + 2b^2B - 2b(Ab - aB)\tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(b - a \tan(c + dx + \frac{\pi}{2}))} dx}{2b(a^2 + b^2)} + \frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{bd(a^2 + b^2)(a \cot(c + dx) + b)} \\
 & \quad \downarrow \text{4136}
 \end{aligned}$$

$$\frac{\int \frac{2(b(-Ba^2+2Aba+b^2B)-b(Aa^2+2bBa-Ab^2)\cot(c+dx))}{\sqrt{\cot(c+dx)}} dx + \frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)(b+a\cot(c+dx))}} dx}{a^2+b^2}}{2b(a^2+b^2)} + \frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

27

$$\frac{2 \int \frac{b(-Ba^2+2Aba+b^2B)-b(Aa^2+2bBa-Ab^2)\cot(c+dx)}{\sqrt{\cot(c+dx)}} dx + \frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)(b+a\cot(c+dx))}} dx}{a^2+b^2}}{2b(a^2+b^2)} + \frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

3042

$$\frac{2 \int \frac{b(-Ba^2+2Aba+b^2B)+b(Aa^2+2bBa-Ab^2)\tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx + \frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2}}{2b(a^2+b^2)} + \frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

4017

$$\frac{4 \int -\frac{b(-Ba^2+2Aba+b^2B)-(Aa^2+2bBa-Ab^2)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2}}{2b(a^2+b^2)} + \frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

25

$$\frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{4 \int \frac{b(-Ba^2+2Aba+b^2B)-(Aa^2+2bBa-Ab^2)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)}}{2b(a^2+b^2)} + \frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

27

3.598. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

$$\frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{4b \int \frac{-Ba^2+2Aba+b^2B-(Aa^2+2bBa-Ab^2)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} +$$

$$\frac{2b(a^2+b^2)}{bd(a^2+b^2)(a\cot(c+dx)+b)} \frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

↓ 1482

$$\frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{4b\left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{2b(a^2+b^2)}$$

$$\frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

↓ 1476

$$\frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{4b\left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{2b(a^2+b^2)}$$

$$\frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

↓ 1082

$$\frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{4b\left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{2b(a^2+b^2)}$$

$$\frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

↓ 217

$$\frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{4b\left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}\right)}{2b(a^2+b^2)}$$

$$\frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

↓ 1479

3.598. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

$$\frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 4b \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)}} \frac{dx}{2\sqrt{2}} \right) \right)$$

$$\frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

↓ 25

$$\frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 4b \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)}} \frac{dx}{2\sqrt{2}} \right) \right)$$

$$\frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

↓ 27

$$\frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 4b \left(\frac{1}{2}(a^2(A-B)+2ab(A+B)-b^2(A-B)) \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)}} \frac{dx}{2\sqrt{2}} \right) \right)$$

$$\frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

↓ 1103

$$\frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 4b \left(\frac{1}{2}(-(a^2(A+B))+2ab(A-B)+b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)$$

$$\frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

↓ 4117

$$\frac{a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{1}{\sqrt{\cot(c+dx)(b+a\cot(c+dx))}} d(-\cot(c+dx))}{d(a^2+b^2)} - 4b \left(\frac{1}{2}(-(a^2(A+B))+2ab(A-B)+b^2(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)$$

$$\frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a\cot(c+dx)+b)}$$

↓ 73

3.598. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

$$\frac{2a(a^3B+a^2Ab+5ab^2B-3Ab^3) \int \frac{1}{a \cot^2(c+dx)+b} d\sqrt{\cot(c+dx)} - 4b \left(\frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B) \right) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \arctan(1-\sqrt{2}\sqrt{\cot(c+dx)+1}) \right)}{d(a^2+b^2)} = \frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{bd(a^2+b^2)(a \cot(c+dx)+b)} + \frac{2\sqrt{a}(a^3B+a^2Ab+5ab^2B-3Ab^3) \arctan\left(\frac{\sqrt{a} \cot(c+dx)}{\sqrt{b}}\right) - 4b \left(\frac{1}{2}(-a^2(A+B))+2ab(A-B)+b^2(A+B) \right) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}} - \arctan(1-\sqrt{2}\sqrt{\cot(c+dx)+1}) \right)}{\sqrt{bd}(a^2+b^2)}$$

$2b(a^2 + b^2)$

```
input Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^2),x]
```

```
output (a*(A*b - a*B)*Sqrt[Cot[c + d*x]])/(b*(a^2 + b^2)*d*(b + a*Cot[c + d*x]))
+ ((2*Sqrt[a]*(a^2*A*b - 3*A*b^3 + a^3*B + 5*a*b^2*B)*ArcTan[(Sqrt[a]*Cot[
c + d*x])/Sqrt[b]])/(Sqrt[b]*(a^2 + b^2)*d) - (4*b*((2*a*b*(A - B) - a^2*
(A + B) + b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]/Sqrt[2]]
+ ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]/Sqrt[2]]))/2 + ((a^2*(A - B) - b^2
*(A - B) + 2*a*b*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c
+ d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*S
qrt[2])))/2)/((a^2 + b^2)*d)/(2*b*(a^2 + b^2))
```

3.598.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

3.598. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

- rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`
- rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)]*(g_))^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2), x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4136 `Int((((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)]) + (C_)*tan[(e_) + (f_)*(x_)])^2)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.598.4 Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 336, normalized size of antiderivative = 0.86

method	result
derivativedivides	$\frac{\left((2abA - B a^2 + B b^2) \sqrt{2} \left(\ln \left(\frac{1 + \cot(dx+c) + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c) - \sqrt{2} \sqrt{\cot(dx+c)}} \right) + 2 \arctan(1 + \sqrt{2} \sqrt{\cot(dx+c)}) + 2 \arctan(-1 + \sqrt{2} \sqrt{\cot(dx+c)}) \right) \right)}{8}$
default	$\frac{\left((2abA - B a^2 + B b^2) \sqrt{2} \left(\ln \left(\frac{1 + \cot(dx+c) + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c) - \sqrt{2} \sqrt{\cot(dx+c)}} \right) + 2 \arctan(1 + \sqrt{2} \sqrt{\cot(dx+c)}) + 2 \arctan(-1 + \sqrt{2} \sqrt{\cot(dx+c)}) \right) \right)}{8}$

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^2,x,method=_RETURNV
ERBOSE)`

output `1/d*(-2/(a^2+b^2)^2*(1/8*(2*A*a*b-B*a^2+B*b^2)*2^(1/2)*(ln((1+cot(d*x+c)+2
^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan
(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/8*(-
A*a^2+A*b^2-2*B*a*b)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(
1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2
))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))) -2*a/(a^2+b^2)^2*(-1/2*(A*a^2*b+
A*b^3-B*a^3-B*a*b^2)/b*cot(d*x+c)^(1/2)/(b+a*cot(d*x+c))+1/2*(A*a^2*b-3*A*
b^3+B*a^3+5*B*a*b^2)/b/(a*b)^(1/2)*arctan(a*cot(d*x+c)^(1/2)/(a*b)^(1/2)))`

3.598.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5899 vs. 2(353) = 706.

Time = 19.64 (sec) , antiderivative size = 11824, normalized size of antiderivative = 30.16

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^2,x, algorith
m="fracas")`

output `Too large to include`

3.598.
$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$$

3.598.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^2 \cot^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(3/2)/(a+b*tan(d*x+c))**2,x)`

output `Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))**2*cot(c + d*x)**(3/2)), x)`

3.598.7 Maxima [A] (verification not implemented)

Time = 0.73 (sec) , antiderivative size = 360, normalized size of antiderivative = 0.92

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \frac{4(Ba^4 + Aa^3b + 5Ba^2b^2 - 3Aab^3) \arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right)}{(a^4b + 2a^2b^3 + b^5)\sqrt{ab}} - \frac{2\sqrt{2}((A+B)a^2 - 2(A-B)ab - (A+B)b^2) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right)}{(a^4b + 2a^2b^3 + b^5)\sqrt{ab}}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^2,x, algorithm m="maxima")`

output `-1/4*(4*(B*a^4 + A*a^3*b + 5*B*a^2*b^2 - 3*A*a*b^3)*arctan(a/(sqrt(a*b)*sqrt(tan(d*x + c))))/((a^4*b + 2*a^2*b^3 + b^5)*sqrt(a*b)) - (2*sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A + B)*a^2 - 2*(A - B)*a*b - (A + B)*b^2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*((A - B)*a^2 + 2*(A + B)*a*b - (A - B)*b^2)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))/(a^4 + 2*a^2*b^2 + b^4) + 4*(B*a^2 - A*a*b)/((a^2*b^2 + b^4 + (a^3*b + a*b^3)/tan(d*x + c))*sqrt(tan(d*x + c)))/d`

3.598.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a)^2 \cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^2,x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)/((b*tan(d*x + c) + a)^2*cot(d*x + c)^(3/2)), x)`

3.598.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2} (a + b \tan(c + dx))^2} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))^2),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))^2), x)`

3.599 $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

3.599.1 Optimal result 5701
 3.599.2 Mathematica [A] (verified) 5702
 3.599.3 Rubi [A] (warning: unable to verify) 5703
 3.599.4 Maple [A] (verified) 5711
 3.599.5 Fricas [B] (verification not implemented) 5712
 3.599.6 Sympy [F(-1)] 5712
 3.599.7 Maxima [A] (verification not implemented) 5712
 3.599.8 Giac [F(-1)] 5713
 3.599.9 Mupad [F(-1)] 5713

3.599.1 Optimal result

Integrand size = 33, antiderivative size = 437

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$$

$$= -\frac{(a^2(A - B) - b^2(A - B) + 2ab(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^2 d}$$

$$+ \frac{(a^2(A - B) - b^2(A - B) + 2ab(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^2 d}$$

$$- \frac{a^{3/2}(a^2 Ab + 5Ab^3 - 3a^3 B - 7ab^2 B) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{b^{5/2}(a^2 + b^2)^2 d}$$

$$- \frac{aAb - 3a^2 B - 2b^2 B}{b^2(a^2 + b^2)d\sqrt{\cot(c + dx)}} + \frac{a(Ab - aB)}{b(a^2 + b^2)d\sqrt{\cot(c + dx)}(b + a \cot(c + dx))}$$

$$+ \frac{(2ab(A - B) - a^2(A + B) + b^2(A + B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^2 d}$$

$$- \frac{(2ab(A - B) - a^2(A + B) + b^2(A + B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^2 d}$$

output
$$-a^{3/2}*(A*a^2*b+5*A*b^3-3*B*a^3-7*B*a*b^2)*\arctan(a^{1/2}*\cot(d*x+c)^{(1/2)}/b^{(1/2)})/b^{(5/2)}/(a^2+b^2)^2/d+1/2*(a^2*(A-B)-b^2*(A-B)+2*a*b*(A+B))*\arctan(-1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)^2/d*2^{(1/2)}+1/2*(a^2*(A-B)-b^2*(A-B)+2*a*b*(A+B))*\arctan(1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)^2/d*2^{(1/2)}+1/4*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\ln(1+\cot(d*x+c)-2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)^2/d*2^{(1/2)}-1/4*(2*a*b*(A-B)-a^2*(A+B)+b^2*(A+B))*\ln(1+\cot(d*x+c)+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)^2/d*2^{(1/2)}+(-A*a*b+3*B*a^2+2*B*b^2)/b^2/(a^2+b^2)/d/\cot(d*x+c)^{(1/2)}+a*(A*b-B*a)/b/(a^2+b^2)/d/(b+a*\cot(d*x+c))/\cot(d*x+c)^{(1/2)}$$

3.599.2 Mathematica [A] (verified)

Time = 3.63 (sec) , antiderivative size = 390, normalized size of antiderivative = 0.89

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$$

$$= \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(\frac{2\sqrt{2}(a^2(A-B)+b^2(-A+B)+2ab(A+B))(\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})-\arctan(1+\sqrt{2}\sqrt{\tan(c+dx)})}{(a^2+b^2)^2} \right)}{\dots}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^2),x]`

output
$$\left(\text{Sqrt}[\text{Cot}[c + d*x]] * \text{Sqrt}[\text{Tan}[c + d*x]] * \left((2 * \text{Sqrt}[2] * (a^2 * (A - B) + b^2 * (-A + B) + 2 * a * b * (A + B))) * (\text{ArcTan}[1 - \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[c + d*x]]] - \text{ArcTan}[1 + \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[c + d*x]]) \right) / (a^2 + b^2)^2 + (4 * a^{3/2} * (-A * b) + a * B) * \text{ArcTan}[(\text{Sqrt}[b] * \text{Sqrt}[\text{Tan}[c + d*x]]) / \text{Sqrt}[a]] / (b^{5/2} * (a^2 + b^2)) + (8 * a^{3/2} * (a^2 * A * b + 3 * A * b^3 - 2 * a^3 * B - 4 * a * b^2 * B)) * \text{ArcTan}[(\text{Sqrt}[b] * \text{Sqrt}[\text{Tan}[c + d*x]]) / \text{Sqrt}[a]] / (b^{5/2} * (a^2 + b^2)^2) - (\text{Sqrt}[2] * (2 * a * b * (-A + B) + a^2 * (A + B) - b^2 * (A + B))) * (\text{Log}[1 - \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]] - \text{Log}[1 + \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]]) \right) / (a^2 + b^2)^2 + (8 * B * \text{Sqrt}[\text{Tan}[c + d*x]]) / b^2 + (4 * a^2 * (-A * b) + a * B) * \text{Sqrt}[\text{Tan}[c + d*x]] / (b^2 * (a^2 + b^2) * (a + b * \text{Tan}[c + d*x])) \right) / (4 * d)$$

3.599.3 Rubi [A] (warning: unable to verify)

Time = 2.26 (sec) , antiderivative size = 375, normalized size of antiderivative = 0.86, number of steps used = 27, number of rules used = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.788$, Rules used = {3042, 4064, 3042, 4092, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{5/2}(a + b \tan(c + dx))^2} dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{A \cot(c + dx) + B}{\cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + b)^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B - A \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}(b - a \tan(c + dx + \frac{\pi}{2}))^2} dx \\
 & \quad \downarrow \text{4092} \\
 & \frac{a(Ab - aB)}{bd(a^2 + b^2) \sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \int \frac{-3Ba^2 + 3(Ab - aB) \cot^2(c + dx)a + Aba - 2b^2B - 2b(Ab - aB) \cot(c + dx)}{2 \cot^{\frac{3}{2}}(c + dx)(b + a \cot(c + dx))} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{a(Ab - aB)}{bd(a^2 + b^2) \sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \int \frac{-3Ba^2 + 3(Ab - aB) \cot^2(c + dx)a + Aba - 2b^2B - 2b(Ab - aB) \cot(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(b + a \cot(c + dx))} dx \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \int \frac{-3Ba^2 + 3(Ab - aB)\tan(c + dx + \frac{\pi}{2})^2 a + Aba - 2b^2 B + 2b(Ab - aB)\tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}(b - a \tan(c + dx + \frac{\pi}{2}))} dx$$

$$\frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} \downarrow 4132$$

$$\frac{2 \int -\frac{-3Ba^3 + Aba^2 + (-3Ba^2 + Aba - 2b^2 B)\cot^2(c + dx)a - 4b^2 Ba + 2Ab^3 - 2b^2(aA + bB)\cot(c + dx)}{2\sqrt{\cot(c + dx)}(b + a \cot(c + dx))} dx}{2b(a^2 + b^2)} + \frac{2(-3a^2 B + aAb - 2b^2 B)}{bd\sqrt{\cot(c + dx)}}$$

$$\frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} \downarrow 27$$

$$\frac{2(-3a^2 B + aAb - 2b^2 B)}{bd\sqrt{\cot(c + dx)}} - \int \frac{-3Ba^3 + Aba^2 + (-3Ba^2 + Aba - 2b^2 B)\cot^2(c + dx)a - 4b^2 Ba + 2Ab^3 - 2b^2(aA + bB)\cot(c + dx)}{\sqrt{\cot(c + dx)}(b + a \cot(c + dx))} dx}{2b(a^2 + b^2)}$$

$$\frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} \downarrow 3042$$

$$\frac{2(-3a^2 B + aAb - 2b^2 B)}{bd\sqrt{\cot(c + dx)}} - \int \frac{-3Ba^3 + Aba^2 + (-3Ba^2 + Aba - 2b^2 B)\tan(c + dx + \frac{\pi}{2})^2 a - 4b^2 Ba + 2Ab^3 + 2b^2(aA + bB)\tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(b - a \tan(c + dx + \frac{\pi}{2}))} dx}{2b(a^2 + b^2)}$$

$$\frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} \downarrow 4136$$

$$\frac{2(-3a^2 B + aAb - 2b^2 B)}{bd\sqrt{\cot(c + dx)}} - \frac{\int -\frac{2((Aa^2 + 2bBa - Ab^2)b^2 + (-Ba^2 + 2Aba + b^2 B)\cot(c + dx)b^2)}{\sqrt{\cot(c + dx)}} dx}{a^2 + b^2} + \frac{a^2(-3a^3 B + a^2 Ab - 7ab^2 B + 5Ab^3) \int \frac{\cot^2(c + dx) + 1}{\sqrt{\cot(c + dx)}(b + a \cot(c + dx))} dx}{a^2 + b^2}}{2b(a^2 + b^2)}$$

$$\frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} \downarrow 27$$

$$\frac{2(-3a^2 B + aAb - 2b^2 B)}{bd\sqrt{\cot(c + dx)}} - \frac{a^2(-3a^3 B + a^2 Ab - 7ab^2 B + 5Ab^3) \int \frac{\cot^2(c + dx) + 1}{\sqrt{\cot(c + dx)}(b + a \cot(c + dx))} dx}{a^2 + b^2} - \frac{2 \int \frac{(Aa^2 + 2bBa - Ab^2)b^2 + (-Ba^2 + 2Aba + b^2 B)\cot(c + dx)}{\sqrt{\cot(c + dx)}}}{a^2 + b^2}}{2b(a^2 + b^2)}$$

3.599. $\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \\ & \frac{2(-3a^2B + aAb - 2b^2B)}{bd\sqrt{\cot(c + dx)}} - \frac{a^2(-3a^3B + a^2Ab - 7ab^2B + 5Ab^3) \int \frac{\tan(c + dx + \frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(b - a \tan(c + dx + \frac{\pi}{2}))} dx}{a^2 + b^2} - \frac{2 \int \frac{b^2(Aa^2 + 2bBa - Ab^2) - b^2(-Ba^2 + 2Aba + b^2)}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}}}{a^2 + b^2} \\ & \frac{2b(a^2 + b^2)}{b} \end{aligned}$$

$$\begin{aligned} & \downarrow 4017 \\ & \frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \\ & \frac{2(-3a^2B + aAb - 2b^2B)}{bd\sqrt{\cot(c + dx)}} - \frac{a^2(-3a^3B + a^2Ab - 7ab^2B + 5Ab^3) \int \frac{\tan(c + dx + \frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(b - a \tan(c + dx + \frac{\pi}{2}))} dx}{a^2 + b^2} - \frac{4 \int \frac{b^2(Aa^2 + 2bBa - Ab^2 + (-Ba^2 + 2Aba + b^2) \cot^2(c + dx) + 1)}{\cot^2(c + dx) + 1}}{d(a^2 + b^2)} \\ & \frac{2b(a^2 + b^2)}{b} \end{aligned}$$

$$\begin{aligned} & \downarrow 25 \\ & \frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \\ & \frac{2(-3a^2B + aAb - 2b^2B)}{bd\sqrt{\cot(c + dx)}} - \frac{4 \int \frac{b^2(Aa^2 + 2bBa - Ab^2 + (-Ba^2 + 2Aba + b^2) \cot(c + dx))}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d(a^2 + b^2)} + \frac{a^2(-3a^3B + a^2Ab - 7ab^2B + 5Ab^3) \int \frac{\tan(c + dx)}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}}}{a^2 + b^2} \\ & \frac{2b(a^2 + b^2)}{b} \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \\ & \frac{2(-3a^2B + aAb - 2b^2B)}{bd\sqrt{\cot(c + dx)}} - \frac{4b^2 \int \frac{Aa^2 + 2bBa - Ab^2 + (-Ba^2 + 2Aba + b^2) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}}{d(a^2 + b^2)} + \frac{a^2(-3a^3B + a^2Ab - 7ab^2B + 5Ab^3) \int \frac{\tan(c + dx)}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}}}{a^2 + b^2} \\ & \frac{2b(a^2 + b^2)}{b} \end{aligned}$$

$$\begin{aligned} & \downarrow 1482 \\ & \frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \\ & \frac{2(-3a^2B + aAb - 2b^2B)}{bd\sqrt{\cot(c + dx)}} - \frac{4b^2 \left(\frac{1}{2}(a^2(A - B) + 2ab(A + B) - b^2(A - B)) \int \frac{\cot(c + dx) + 1}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} - \frac{1}{2}(-a^2(A + B) + 2ab(A - B) + b^2(A + B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx)} \right)}{d(a^2 + b^2)} \\ & \frac{2b(a^2 + b^2)}{b} \end{aligned}$$

3.599. $\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$

$$\begin{aligned} & \downarrow 1476 \\ & \frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \\ & \frac{2(-3a^2B + aAb - 2b^2B)}{bd\sqrt{\cot(c + dx)}} - \frac{4b^2\left(\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B))\left(\frac{1}{2}\int \frac{1}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)} + \frac{1}{2}\int \frac{1}{\cot(c + dx) + \sqrt{2}\sqrt{\cot(c + dx) + 1}} d\sqrt{\cot(c + dx)}\right)\right)}{d(a^2 + b^2)} \end{aligned}$$

2b

$$\begin{aligned} & \downarrow 1082 \\ & \frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \\ & \frac{2(-3a^2B + aAb - 2b^2B)}{bd\sqrt{\cot(c + dx)}} - \frac{4b^2\left(\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B))\left(\frac{\int \frac{1}{-\cot(c + dx) - 1} d(1 - \sqrt{2}\sqrt{\cot(c + dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c + dx) - 1} d(\sqrt{2}\sqrt{\cot(c + dx) + 1})}{\sqrt{2}}\right)\right)}{d(a^2 + b^2)} \end{aligned}$$

2b(a² +

$$\begin{aligned} & \downarrow 217 \\ & \frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \\ & \frac{2(-3a^2B + aAb - 2b^2B)}{bd\sqrt{\cot(c + dx)}} - \frac{4b^2\left(\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B))\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\cot(c + dx)})}{\sqrt{2}}\right) - \frac{1}{2}(-(a^2(A+B)) + 2ab(A-B))\right)}{d(a^2 + b^2)} \end{aligned}$$

2b(a² + b²)

$$\begin{aligned} & \downarrow 1479 \\ & \frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \\ & \frac{2(-3a^2B + aAb - 2b^2B)}{bd\sqrt{\cot(c + dx)}} - \frac{4b^2\left(\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B))\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c + dx)} + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}\sqrt{\cot(c + dx)})}{\sqrt{2}}\right) - \frac{1}{2}(-(a^2(A+B)) + 2ab(A-B))\right)}{d(a^2 + b^2)} \end{aligned}$$

d(a² + b²)

↓ 25

3.599. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

$$\frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \frac{4b^2 \left(\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(-(a^2(A+B)) + 2ab(A-B)) \right)}{bd\sqrt{\cot(c+dx)}} - \frac{2(-3a^2B + aAb - 2b^2B)}{bd\sqrt{\cot(c+dx)}} - \frac{d(a^2 + b^2)}{d(a^2 + b^2)}$$

27

$$\frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \frac{4b^2 \left(\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(-(a^2(A+B)) + 2ab(A-B)) \right)}{bd\sqrt{\cot(c+dx)}} - \frac{2(-3a^2B + aAb - 2b^2B)}{bd\sqrt{\cot(c+dx)}} - \frac{d(a^2 + b^2)}{d(a^2 + b^2)}$$

1103

$$\frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \frac{a^2(-3a^3B + a^2Ab - 7ab^2B + 5Ab^3) \int \frac{\tan(c+dx+\frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{bd\sqrt{\cot(c+dx)}} + \frac{4b^2 \left(\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(-(a^2(A+B)) + 2ab(A-B)) \right)}{bd\sqrt{\cot(c+dx)}} - \frac{2(-3a^2B + aAb - 2b^2B)}{bd\sqrt{\cot(c+dx)}} - \frac{d(a^2 + b^2)}{d(a^2 + b^2)}$$

4117

$$\frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \frac{a^2(-3a^3B + a^2Ab - 7ab^2B + 5Ab^3) \int \frac{1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} d(-\cot(c+dx))}{bd\sqrt{\cot(c+dx)}} + \frac{4b^2 \left(\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(-(a^2(A+B)) + 2ab(A-B)) \right)}{bd\sqrt{\cot(c+dx)}} - \frac{2(-3a^2B + aAb - 2b^2B)}{bd\sqrt{\cot(c+dx)}} - \frac{d(a^2 + b^2)}{d(a^2 + b^2)}$$

73

$$\frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)} - \frac{4b^2 \left(\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2}(-(a^2(A+B)) + 2ab(A-B)) \right)}{bd\sqrt{\cot(c+dx)}} - \frac{2(-3a^2B + aAb - 2b^2B)}{bd\sqrt{\cot(c+dx)}} - \frac{d(a^2 + b^2)}{d(a^2 + b^2)}$$

218

3.599. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

2b(a

$$\frac{a(Ab - aB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a\cot(c + dx) + b)} - \frac{4b^2\left(\frac{1}{2}(a^2(A-B) + 2ab(A+B) - b^2(A-B))\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}}\right) - \frac{1}{2}(-(a^2(A+B)) + 2ab(A-B))\right)}{d(a^2+b^2)}$$

$$\frac{2(-3a^2B+aAb-2b^2B)}{bd\sqrt{\cot(c+dx)}} - \frac{2b(a^2 + b^2)}{d(a^2+b^2)}$$

input `Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^2),x]`

output `(a*(A*b - a*B))/(b*(a^2 + b^2)*d*Sqrt[Cot[c + d*x]]*(b + a*Cot[c + d*x])) - ((2*(a*A*b - 3*a^2*B - 2*b^2*B))/(b*d*Sqrt[Cot[c + d*x]]) - ((2*a^(3/2)*(a^2*A*b + 5*A*b^3 - 3*a^3*B - 7*a*b^2*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]])/(Sqrt[b]*(a^2 + b^2)*d) + (4*b^2*((a^2*(A - B) - b^2*(A - B) + 2*a*b*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]))/2 - ((2*a*b*(A - B) - a^2*(A + B) + b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/(a^2 + b^2)*d)/b)/(2*b*(a^2 + b^2))`

3.599.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

- rule 218 $\text{Int}[(a_.) + (b_.) \cdot (x_.)^2]^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a) \cdot \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$
- rule 1082 $\text{Int}[(a_.) + (b_.) \cdot (x_.) + (c_.) \cdot (x_.)^2]^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4 \cdot \text{Simplify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4 \cdot a \cdot c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_.) + (e_.) \cdot (x_.)] / [(a_.) + (b_.) \cdot (x_.) + (c_.) \cdot (x_.)^2], x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$
- rule 1476 $\text{Int}[(d_.) + (e_.) \cdot (x_.)^2] / [(a_.) + (c_.) \cdot (x_.)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c) \ \text{Int}[1/\text{Simp}[d/e + q \cdot x + x^2, x], x], x] + \text{Simp}[e/(2 \cdot c) \ \text{Int}[1/\text{Simp}[d/e - q \cdot x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{PosQ}[d \cdot e]$
- rule 1479 $\text{Int}[(d_.) + (e_.) \cdot (x_.)^2] / [(a_.) + (c_.) \cdot (x_.)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c \cdot q) \ \text{Int}[(q - 2 \cdot x)/\text{Simp}[d/e + q \cdot x - x^2, x], x], x] + \text{Simp}[e/(2 \cdot c \cdot q) \ \text{Int}[(q + 2 \cdot x)/\text{Simp}[d/e - q \cdot x - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{NegQ}[d \cdot e]$
- rule 1482 $\text{Int}[(d_.) + (e_.) \cdot (x_.)^2] / [(a_.) + (c_.) \cdot (x_.)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a \cdot c, 2]\}, \text{Simp}[(d \cdot q + a \cdot e)/(2 \cdot a \cdot c) \ \text{Int}[(q + c \cdot x^2)/(a + c \cdot x^4), x], x] + \text{Simp}[(d \cdot q - a \cdot e)/(2 \cdot a \cdot c) \ \text{Int}[(q - c \cdot x^2)/(a + c \cdot x^4), x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{NeQ}[c \cdot d^2 + a \cdot e^2, 0] \ \&\& \ \text{NeQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{NegQ}[-(a) \cdot c]$
- rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4017 $\text{Int}[(c_.) + (d_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)] / \text{Sqrt}[(b_.) \cdot \tan[(e_.) + (f_.) \cdot (x_.)]], x_Symbol] \rightarrow \text{Simp}[2/f \ \text{Subst}[\text{Int}[(b \cdot c + d \cdot x^2)/(b^2 + x^4), x], x, \text{Sqrt}[b \cdot \text{Tan}[e + f \cdot x]], x] /; \text{FreeQ}[\{b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[c^2 - d^2, 0] \ \&\& \ \text{NeQ}[c^2 + d^2, 0]$

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp [g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_))*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2])/(a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.599.4 Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 351, normalized size of antiderivative = 0.80

method	result
derivativedivides	$\frac{2 \left((-A a^2 + A b^2 - 2Bab) \sqrt{2} \left(\ln \left(\frac{1 + \cot(dx+c) + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c) - \sqrt{2} \sqrt{\cot(dx+c)}} \right) + 2 \arctan(1 + \sqrt{2} \sqrt{\cot(dx+c)}) + 2 \arctan(-1 + \sqrt{2} \sqrt{\cot(dx+c)}) \right)}{8} \right)}{a^2}$
default	$\frac{2 \left((-A a^2 + A b^2 - 2Bab) \sqrt{2} \left(\ln \left(\frac{1 + \cot(dx+c) + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c) - \sqrt{2} \sqrt{\cot(dx+c)}} \right) + 2 \arctan(1 + \sqrt{2} \sqrt{\cot(dx+c)}) + 2 \arctan(-1 + \sqrt{2} \sqrt{\cot(dx+c)}) \right)}{8} \right)}{a^2}$

```
input int((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c))^2,x,method=_RETURNV
ERBOSE)
```

```
output 1/d*(-2/(a^2+b^2)^2*(1/8*(-A*a^2+A*b^2-2*B*a*b)*2^(1/2)*(ln((1+cot(d*x+c)+
2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan
(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/8*(
-2*A*a*b+B*a^2-B*b^2)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/
(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/
2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+2*B/b^2/cot(d*x+c)^(1/2)-2*a^2
/b^2/(a^2+b^2)^2*((1/2*A*a^2*b+1/2*A*b^3-1/2*B*a^3-1/2*B*a*b^2)*cot(d*x+c)
^(1/2)/(b+a*cot(d*x+c))+1/2*(A*a^2*b+5*A*b^3-3*B*a^3-7*B*a*b^2)/(a*b)^(1/2)
)*arctan(a*cot(d*x+c)^(1/2)/(a*b)^(1/2)))
```

3.599.
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$$

3.599.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5952 vs. $2(396) = 792$.

Time = 35.18 (sec) , antiderivative size = 11930, normalized size of antiderivative = 27.30

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c))^2,x, algorithm m="fracas")`

output Too large to include

3.599.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(5/2)/(a+b*tan(d*x+c))**2,x)`

output Timed out

3.599.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 402, normalized size of antiderivative = 0.92

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^2} dx$$

$$= \frac{4(3Ba^5 - Aa^4b + 7Ba^3b^2 - 5Aa^2b^3) \arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right)}{(a^4b^2 + 2a^2b^4 + b^6)\sqrt{ab}} + \frac{2\sqrt{2}((A-B)a^2 + 2(A+B)ab - (A-B)b^2) \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right)}{\dots}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c))^2,x, algorithm m="maxima")`

3.599. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^2} dx$

output $\frac{1}{4} \cdot (4 \cdot (3B \cdot a^5 - A \cdot a^4 \cdot b + 7B \cdot a^3 \cdot b^2 - 5A \cdot a^2 \cdot b^3) \cdot \arctan(a / (\sqrt{a \cdot b} \cdot \sqrt{\tan(dx + c)}))) / ((a^4 \cdot b^2 + 2a^2 \cdot b^4 + b^6) \cdot \sqrt{a \cdot b}) + (2 \cdot \sqrt{2} \cdot ((A - B) \cdot a^2 + 2 \cdot (A + B) \cdot a \cdot b - (A - B) \cdot b^2) \cdot \arctan(1/2 \cdot \sqrt{2} \cdot (\sqrt{2} + 2/\sqrt{\tan(dx + c)}))) + 2 \cdot \sqrt{2} \cdot ((A - B) \cdot a^2 + 2 \cdot (A + B) \cdot a \cdot b - (A - B) \cdot b^2) \cdot \arctan(-1/2 \cdot \sqrt{2} \cdot (\sqrt{2} - 2/\sqrt{\tan(dx + c)}))) + \sqrt{2} \cdot ((A + B) \cdot a^2 - 2 \cdot (A - B) \cdot a \cdot b - (A + B) \cdot b^2) \cdot \log(\sqrt{2} / \sqrt{\tan(dx + c)} + 1 / \tan(dx + c) + 1) - \sqrt{2} \cdot ((A + B) \cdot a^2 - 2 \cdot (A - B) \cdot a \cdot b - (A + B) \cdot b^2) \cdot \log(-\sqrt{2} / \sqrt{\tan(dx + c)} + 1 / \tan(dx + c) + 1)) / (a^4 + 2a^2 \cdot b^2 + b^4) + 4 \cdot (2B \cdot a^2 \cdot b + 2B \cdot b^3 + (3B \cdot a^3 - A \cdot a^2 \cdot b + 2B \cdot a \cdot b^2) / \tan(dx + c)) / ((a^2 \cdot b^3 + b^5) / \sqrt{\tan(dx + c)} + (a^3 \cdot b^2 + a \cdot b^4) / \tan(dx + c)^{3/2}) / d$

3.599.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{5/2}(c + dx)(a + b \tan(c + dx))^2} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c))^2,x, algorithm="giac")`

output Timed out

3.599.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{5/2}(c + dx)(a + b \tan(c + dx))^2} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{5/2} (a + b \tan(c + dx))^2} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + b*tan(c + d*x))^2),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + b*tan(c + d*x))^2), x)`

3.600
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

3.600.1 Optimal result	5714
3.600.2 Mathematica [A] (verified)	5715
3.600.3 Rubi [A] (warning: unable to verify)	5716
3.600.4 Maple [B] (verified)	5726
3.600.5 Fricas [B] (verification not implemented)	5727
3.600.6 Sympy [F]	5728
3.600.7 Maxima [A] (verification not implemented)	5728
3.600.8 Giac [F]	5729
3.600.9 Mupad [F(-1)]	5729

3.600.1 Optimal result

Integrand size = 33, antiderivative size = 601

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx =$$

$$\frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d}$$

$$+ \frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d}$$

$$+ \frac{b^{3/2}(63a^4Ab + 46a^2Ab^3 + 15Ab^5 - 35a^5B - 6a^3b^2B - 3ab^4B) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{4a^{7/2}(a^2+b^2)^3 d}$$

$$- \frac{(8a^4A + 31a^2Ab^2 + 15Ab^4 - 11a^3bB - 3ab^3B) \sqrt{\cot(c+dx)}}{4a^3(a^2+b^2)^2 d}$$

$$+ \frac{b(Ab - aB) \cot^{\frac{5}{2}}(c+dx)}{2a(a^2+b^2)d(b+a \cot(c+dx))^2} + \frac{b(13a^2Ab + 5Ab^3 - 9a^3B - ab^2B) \cot^{\frac{3}{2}}(c+dx)}{4a^2(a^2+b^2)^2 d(b+a \cot(c+dx))}$$

$$+ \frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d}$$

$$- \frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d}$$

3.600.
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

output $\frac{1}{4}b^{3/2}(63Aa^4b+46Aa^2b^3+15Ab^5-35Ba^5-6Ba^3b^2-3Bab^4)\arctan(a^{1/2}\cot(dx+c)^{1/2}/b^{1/2})/a^{7/2}/(a^2+b^2)^3/d+1/2b*(A*b-B*a)\cot(dx+c)^{5/2}/a/(a^2+b^2)/d/(b+a\cot(dx+c))^2+1/4b*(13Aa^2*b+5Aa*b^3-9Ba^3-Ba*b^2)\cot(dx+c)^{3/2}/a^2/(a^2+b^2)^2/d/(b+a\cot(dx+c))+1/2*(a^3*(A-B)-3a*b^2*(A-B)+3a^2*b*(A+B)-b^3*(A+B))\arctan(-1+2^{1/2}\cot(dx+c)^{1/2})/(a^2+b^2)^3/d*2^{1/2}+1/2*(a^3*(A-B)-3a*b^2*(A-B)+3a^2*b*(A+B)-b^3*(A+B))\arctan(1+2^{1/2}\cot(dx+c)^{1/2})/(a^2+b^2)^3/d*2^{1/2}+1/4*(3a^2*b*(A-B)-b^3*(A-B)-a^3*(A+B)+3a*b^2*(A+B))*\ln(1+\cot(dx+c))-2^{1/2}\cot(dx+c)^{1/2})/(a^2+b^2)^3/d*2^{1/2}-1/4*(3a^2*b*(A-B)-b^3*(A-B)-a^3*(A+B)+3a*b^2*(A+B))*\ln(1+\cot(dx+c)+2^{1/2}\cot(dx+c)^{1/2})/(a^2+b^2)^3/d*2^{1/2}-1/4*(8Aa^4+31Aa^2b^2+15Ab^4-11Ba^3b-3Bab^3)\cot(dx+c)^{1/2}/a^3/(a^2+b^2)^2/d$

3.600.2 Mathematica [A] (verified)

Time = 6.56 (sec) , antiderivative size = 617, normalized size of antiderivative = 1.03

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

$$= \frac{2\sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}\left(\frac{(a^3(A-B)-3ab^2(A-B)+3a^2b(A+B)-b^3(A+B))\left(\sqrt{2}\arctan\left(1-\sqrt{2}\sqrt{\tan(c+dx)}\right)-\sqrt{2}\arctan\left(1+\sqrt{2}\sqrt{\tan(c+dx)}\right)\right)}{4(a^2+b^2)^3}\right)}{1}$$

input `Integrate[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3, x]`

output $(2\sqrt{\cot[c + dx]}\sqrt{\tan[c + dx]}*((a^3(A - B) - 3ab^2(A - B) + 3a^2b(A + B) - b^3(A + B))*(\sqrt{2}\operatorname{ArcTan}[1 - \sqrt{2}\sqrt{\tan[c + dx]}] - \sqrt{2}\operatorname{ArcTan}[1 + \sqrt{2}\sqrt{\tan[c + dx]}]))/ (4(a^2 + b^2)^3) - (3b^{3/2}(Ab - aB)\operatorname{ArcTan}[(\sqrt{b}\sqrt{\tan[c + dx]})/\sqrt{a}]) / (8a^{7/2}(a^2 + b^2)) - (b^{3/2}(3a^2Ab + Ab^3 - 2a^3B)\operatorname{ArcTan}[(\sqrt{b}\sqrt{\tan[c + dx]})/\sqrt{a}]) / (2a^{7/2}(a^2 + b^2)^2) - (b^{3/2}(6a^4Ab + 3a^2Ab^3 + Ab^5 - 3a^5B + a^3b^2B)\operatorname{ArcTan}[(\sqrt{b}\sqrt{\tan[c + dx]})/\sqrt{a}]) / (a^{7/2}(a^2 + b^2)^3) + ((3a^2b(A - B) - b^3(A - B) - a^3(A + B) + 3ab^2(A + B))*(\sqrt{2}\operatorname{Log}[1 - \sqrt{2}\sqrt{\tan[c + dx]}] + \operatorname{Tan}[c + dx]) - \sqrt{2}\operatorname{Log}[1 + \sqrt{2}\sqrt{\tan[c + dx]}] + \operatorname{Tan}[c + dx])) / (8(a^2 + b^2)^3) - A / (a^3\sqrt{\tan[c + dx]}) - (b^2(Ab - aB)\sqrt{\tan[c + dx]}) / (4a^2(a^2 + b^2)(a + b\tan[c + dx])^2) - (3b^2(Ab - aB)\sqrt{\tan[c + dx]}) / (8a^3(a^2 + b^2)(a + b\tan[c + dx])) - (b^2(3a^2Ab + Ab^3 - 2a^3B)\sqrt{\tan[c + dx]}) / (2a^3(a^2 + b^2)^2(a + b\tan[c + dx])))) / d$

3.600.3 Rubi [A] (warning: unable to verify)

Time = 3.15 (sec) , antiderivative size = 524, normalized size of antiderivative = 0.87, number of steps used = 30, number of rules used = 29, $\frac{\text{number of rules}}{\text{integrand size}} = 0.879$, Rules used = {3042, 4064, 3042, 4088, 27, 3042, 4128, 27, 3042, 4130, 27, 3042, 4136, 27, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^{\frac{3}{2}}(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$$

↓ 3042

$$\int \frac{\cot(c + dx)^{3/2}(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$$

↓ 4064

$$\int \frac{\cot^{\frac{7}{2}}(c + dx)(A \cot(c + dx) + B)}{(a \cot(c + dx) + b)^3} dx$$

↓ 3042

$$\int \frac{(-\tan(c + dx + \frac{\pi}{2}))^{7/2}(B - A \tan(c + dx + \frac{\pi}{2}))}{(b - a \tan(c + dx + \frac{\pi}{2}))^3} dx$$

3.600. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\begin{aligned}
& \downarrow 4088 \\
& \frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2} - \\
& \frac{\int -\frac{\cot^{\frac{3}{2}}(c+dx)((4Aa^2 - bBa + 5Ab^2) \cot^2(c+dx) - 4a(Ab - aB) \cot(c+dx) + 5b(Ab - aB))}{2(b+a \cot(c+dx))^2} dx}{2a(a^2 + b^2)} \\
& \downarrow 27 \\
& \frac{\int \frac{\cot^{\frac{3}{2}}(c+dx)((4Aa^2 - bBa + 5Ab^2) \cot^2(c+dx) - 4a(Ab - aB) \cot(c+dx) + 5b(Ab - aB))}{(b+a \cot(c+dx))^2} dx}{4a(a^2 + b^2)} + \\
& \frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2} \\
& \downarrow 3042 \\
& \frac{\int \frac{(-\tan(c+dx + \frac{\pi}{2}))^{3/2}((4Aa^2 - bBa + 5Ab^2) \tan^2(c+dx + \frac{\pi}{2}) + 4a(Ab - aB) \tan(c+dx + \frac{\pi}{2}) + 5b(Ab - aB))}{(b-a \tan(c+dx + \frac{\pi}{2}))^2} dx}{4a(a^2 + b^2)} + \\
& \frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2} \\
& \downarrow 4128 \\
& \frac{b(-9a^3B + 13a^2Ab - ab^2B + 5Ab^3) \cot^{\frac{3}{2}}(c+dx)}{ad(a^2 + b^2)(a \cot(c+dx) + b)} - \frac{\int -\frac{\sqrt{\cot(c+dx)}(-8(-Ba^2 + 2Aba + b^2B) \cot(c+dx)a^2 + (8Aa^4 - 11bBa^3 + 31Ab^2a^2 - 3b^3Ba + 15Ab^4) \cot^2(c+dx) + 3b(-9Ba^3 + 13Aba^2 - b^2Ba + 5Ab^3))}{2(b+a \cot(c+dx))}}{a(a^2 + b^2)} dx}{4a(a^2 + b^2)} \\
& \frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2} \\
& \downarrow 27 \\
& \frac{\int \frac{\sqrt{\cot(c+dx)}(-8(-Ba^2 + 2Aba + b^2B) \cot(c+dx)a^2 + (8Aa^4 - 11bBa^3 + 31Ab^2a^2 - 3b^3Ba + 15Ab^4) \cot^2(c+dx) + 3b(-9Ba^3 + 13Aba^2 - b^2Ba + 5Ab^3))}{b+a \cot(c+dx)}}{2a(a^2 + b^2)} dx}{4a(a^2 + b^2)} + \frac{b(-9a^3B + 13a^2Ab - ab^2B + 5Ab^3) \cot^{\frac{3}{2}}(c+dx)}{ad(a^2 + b^2)(a \cot(c+dx) + b)} \\
& \frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2} \\
& \downarrow 3042
\end{aligned}$$

3.600. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\int \frac{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(8\left(-Ba^2+2Aba+b^2B\right)\tan\left(c+dx+\frac{\pi}{2}\right)a^2+\left(8Aa^4-11bBa^3+31Ab^2a^2-3b^3Ba+15Ab^4\right)\tan\left(c+dx+\frac{\pi}{2}\right)^2+3b\left(-9Ba^3+13Aba^2-b^2Ba+5Ab^3\right)}{b-a\tan\left(c+dx+\frac{\pi}{2}\right)} dx$$

$$\frac{2a(a^2+b^2)}{2a(a^2+b^2)}$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 4130

$$2 \int \frac{8\left(Aa^2+2bBa-Ab^2\right)\cot(c+dx)a^3+\left(-8Ba^5+24Aba^4-3b^2Ba^3+31Ab^3a^2-3b^4Ba+15Ab^5\right)\cot^2(c+dx)+b\left(8Aa^4-11bBa^3+31Ab^2a^2-3b^3Ba+15Ab^4\right)}{2\sqrt{\cot(c+dx)}(b+a\cot(c+dx))} dx$$

$$\frac{2a(a^2+b^2)}{2a(a^2+b^2)}$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 27

$$2 \int \frac{8\left(Aa^2+2bBa-Ab^2\right)\cot(c+dx)a^3+\left(-8Ba^5+24Aba^4-3b^2Ba^3+31Ab^3a^2-3b^4Ba+15Ab^5\right)\cot^2(c+dx)+b\left(8Aa^4-11bBa^3+31Ab^2a^2-3b^3Ba+15Ab^4\right)}{\sqrt{\cot(c+dx)}(b+a\cot(c+dx))} dx$$

$$\frac{2a(a^2+b^2)}{2a(a^2+b^2)}$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 3042

$$2 \int \frac{-8\left(Aa^2+2bBa-Ab^2\right)\tan\left(c+dx+\frac{\pi}{2}\right)a^3+\left(-8Ba^5+24Aba^4-3b^2Ba^3+31Ab^3a^2-3b^4Ba+15Ab^5\right)\tan\left(c+dx+\frac{\pi}{2}\right)^2+b\left(8Aa^4-11bBa^3+31Ab^2a^2-3b^3Ba+15Ab^4\right)}{\sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)}\left(b-a\tan\left(c+dx+\frac{\pi}{2}\right)\right)} dx$$

$$\frac{2a(a^2+b^2)}{2a(a^2+b^2)}$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 4136

$$2 \int \frac{8\left(\left(Aa^3+3bBa^2-3Ab^2a-b^3B\right)a^3+\left(-Ba^3+3Aba^2+3b^2Ba-Ab^3\right)\cot(c+dx)a^3\right)}{\sqrt{\cot(c+dx)}(a^2+b^2)} dx + \frac{b^2\left(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5\right)}{a^2+b^2} \int \frac{\cot(c+dx)}{\sqrt{\cot(c+dx)}} dx$$

$$\frac{2a(a^2+b^2)}{2a(a^2+b^2)}$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 27

3.600. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

$$\frac{8 \int \frac{(Aa^3+3bBa^2-3Ab^2a-b^3B)a^3+(-Ba^3+3Aba^2+3b^2Ba-Ab^3) \cot(c+dx)a^3}{\sqrt{\cot(c+dx)} a^2+b^2} dx + \frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5) \int \frac{\cot^2(c+dx)}{\sqrt{\cot(c+dx)}} dx}{a^2+b^2}}{a} = \frac{2a(a^2+b^2)}{4a(a^2+b^2)}$$

$$\frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 3042

$$\frac{8 \int \frac{a^3(Aa^3+3bBa^2-3Ab^2a-b^3B)-a^3(-Ba^3+3Aba^2+3b^2Ba-Ab^3) \tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})} a^2+b^2} dx + \frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5) \int \frac{\sqrt{-\tan(c+dx+\frac{\pi}{2})}}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{a^2+b^2}}{a} = \frac{2a(a^2+b^2)}{4a(a^2+b^2)}$$

$$\frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 4017

$$\frac{16 \int -\frac{a^3(Aa^3+3bBa^2-3Ab^2a-b^3B+(-Ba^3+3Aba^2+3b^2Ba-Ab^3) \cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5) \int \frac{\sqrt{\cot(c+dx)}}{\sqrt{\cot(c+dx)}} dx}{a^2+b^2}}{d(a^2+b^2)} = \frac{2a(a^2+b^2)}{4a(a^2+b^2)}$$

$$\frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 25

$$\frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx - 16 \int \frac{a^3(Aa^3+3bBa^2-3Ab^2a-b^3B+(-Ba^3+3Aba^2+3b^2Ba-Ab^3) \cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{a^2+b^2}}{a} = \frac{2a(a^2+b^2)}{4a(a^2+b^2)}$$

$$\frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 27

3.600. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{16a^3 \int \frac{Aa^3+3bBa^2-3Ab^2a-b^3B+(-Ba^3+3Aba^2)}{\cot^2(c+dx)+1} dx}{d(a^2+b^2)}$$

$$\frac{b(Ab-aB) \cot^{\frac{5}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

$$4a(a^2 +$$

↓ 1482

$$\frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{16a^3 \int \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B))\right)}{a}$$

$$\frac{b(Ab-aB) \cot^{\frac{5}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

↓ 1476

$$\frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{16a^3 \int \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B))\right)}{a}$$

$$\frac{b(Ab-aB) \cot^{\frac{5}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

↓ 1082

$$\frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{16a^3 \int \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B))\right)}{a}$$

$$\frac{b(Ab-aB) \cot^{\frac{5}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

↓ 217

3.600. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5)}{a^2+b^2} \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx - 16a^3 \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right)$$

$$\frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 1479

$$\frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5)}{a^2+b^2} \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx - 16a^3 \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right)$$

$$\frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 25

$$\frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5)}{a^2+b^2} \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx - 16a^3 \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right)$$

$$\frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 27

$$\frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5)}{a^2+b^2} \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx - 16a^3 \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right)$$

$$\frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 1103

3.600. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - 16a^3 \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right)$$

$$\frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 4117

$$\frac{b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5) \int \frac{1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} d(-\cot(c+dx))}{d(a^2+b^2)} - 16a^3 \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right)$$

$$\frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 73

$$\frac{2b^2(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5) \int \frac{1}{a \cot^2(c+dx)+b} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} - 16a^3 \left(\frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right)$$

$$\frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 218

$$\frac{b(Ab - aB) \cot^{\frac{5}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2} +$$

$$\frac{b(-9a^3B+13a^2Ab-ab^2B+5Ab^3) \cot^{\frac{3}{2}}(c+dx)}{ad(a^2+b^2)(a \cot(c+dx)+b)} + \frac{2(8a^4A-11a^3bB+31a^2Ab^2-3ab^3B+15Ab^4) \sqrt{\cot(c+dx)}}{ad} - \frac{2b^{3/2}(-35a^5B+63a^4Ab-6a^3b^2B+46a^2Ab^3-3ab^4B+15Ab^5)}{\sqrt{ad}}$$

input `Int[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

3.600. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

```
output (b*(A*b - a*B)*Cot[c + d*x]^(5/2))/(2*a*(a^2 + b^2)*d*(b + a*Cot[c + d*x])
^2) + ((b*(13*a^2*A*b + 5*A*b^3 - 9*a^3*B - a*b^2*B)*Cot[c + d*x]^(3/2))/(
a*(a^2 + b^2)*d*(b + a*Cot[c + d*x])) + ((-2*(8*a^4*A + 31*a^2*A*b^2 + 15*
A*b^4 - 11*a^3*b*B - 3*a*b^3*B)*Sqrt[Cot[c + d*x]])/(a*d) - ((2*b^(3/2)*(6
3*a^4*A*b + 46*a^2*A*b^3 + 15*A*b^5 - 35*a^5*B - 6*a^3*b^2*B - 3*a*b^4*B)*
ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]])/(Sqrt[a]*(a^2 + b^2)*d) - (16*a^3*
((a^3*(A - B) - 3*a*b^2*(A - B) + 3*a^2*b*(A + B) - b^3*(A + B))*(-(ArcTa
n[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c
+ d*x]]]/Sqrt[2]))/2 - ((3*a^2*b*(A - B) - b^3*(A - B) - a^3*(A + B) + 3*
a*b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sq
rt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2
))/((a^2 + b^2)*d)/a)/(2*a*(a^2 + b^2))/(4*a*(a^2 + b^2))
```

3.600.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_.) + (f_.)*(x_)]*(g_.))^(p_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp [g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

3.600.
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

```
rule 4130 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2], x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

```
rule 4136 Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_) + (C_.)*tan[(e_.) + (f_.)*(x_)^2]))/(a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)], x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
!GtQ[n, 0] && !LeQ[n, -1]
```

3.600.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4580 vs. $2(549) = 1098$.

Time = 0.42 (sec) , antiderivative size = 4581, normalized size of antiderivative = 7.62

method	result	size
derivativedivides	Expression too large to display	4581
default	Expression too large to display	4581

```
input int(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x,method=_RETURNV
ERBOSE)
```

3.600.
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

output $-1/4/d*(1/\tan(dx+c))^{3/2}*\tan(dx+c)*(15*A*\arctan(b*\tan(dx+c)^{1/2}/(a*b)^{1/2}))*\tan(dx+c)^{5/2}*b^9+15*A*(a*b)^{1/2}*\tan(dx+c)^2*b^8+24*A*(a*b)^{1/2}*a^6*b^2+24*A*(a*b)^{1/2}*a^4*b^4+8*A*(a*b)^{1/2}*a^2*b^6+8*A*(a*b)^{1/2}*a^8+39*A*(a*b)^{1/2}*\tan(dx+c)^2*a^4*b^4+46*A*(a*b)^{1/2}*\tan(dx+c)^2*a^2*b^6-11*B*(a*b)^{1/2}*\tan(dx+c)^2*a^5*b^3-14*B*(a*b)^{1/2}*\tan(dx+c)^2*a^3*b^5-3*B*(a*b)^{1/2}*\tan(dx+c)^2*a*b^7+65*A*(a*b)^{1/2}*\tan(dx+c)*a^5*b^3+74*A*(a*b)^{1/2}*\tan(dx+c)*a^3*b^5+25*A*(a*b)^{1/2}*\tan(dx+c)*a*b^7+8*A*(a*b)^{1/2}*a^6*b^2*\tan(dx+c)^2+63*A*\arctan(b*\tan(dx+c)^{1/2}/(a*b)^{1/2}))*\tan(dx+c)^{1/2}*a^6*b^3+46*A*\arctan(b*\tan(dx+c)^{1/2}/(a*b)^{1/2}))*\tan(dx+c)^{1/2}*a^4*b^5+15*A*\arctan(b*\tan(dx+c)^{1/2}/(a*b)^{1/2}))*\tan(dx+c)^{1/2}*a^2*b^7-13*B*(a*b)^{1/2}*\tan(dx+c)*a^6*b^2-18*B*(a*b)^{1/2}*\tan(dx+c)*a^4*b^4-5*B*(a*b)^{1/2}*\tan(dx+c)*a^2*b^6-35*B*\arctan(b*\tan(dx+c)^{1/2}/(a*b)^{1/2}))*\tan(dx+c)^{1/2}*a^7*b^2-6*B*\arctan(b*\tan(dx+c)^{1/2}/(a*b)^{1/2}))*\tan(dx+c)^{1/2}*a^5*b^4-3*B*\arctan(b*\tan(dx+c)^{1/2}/(a*b)^{1/2}))*\tan(dx+c)^{1/2}*a^3*b^6+16*A*(a*b)^{1/2}*a^7*b*\tan(dx+c)-70*B*\arctan(b*\tan(dx+c)^{1/2}/(a*b)^{1/2}))*\tan(dx+c)^{3/2}*a^6*b^3-12*B*\arctan(b*\tan(dx+c)^{1/2}/(a*b)^{1/2}))*\tan(dx+c)^{3/2}*a^4*b^5-6*B*\arctan(b*\tan(dx+c)^{1/2}/(a*b)^{1/2}))*\tan(dx+c)^{3/2}*a^2*b^7+63*A*\arctan(b*\tan(dx+c)^{1/2}/(a*b)^{1/2}))*\tan(dx+c)^{5/2}*a^4*b^5+46*A*\arctan(b*\tan(dx+c)^{1/2}/(a*b)^{1/2}))*\tan(dx+c)^{5/2}*a^2*b^7-35*B*\arctan(b*\tan...$

3.600.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8979 vs. $2(544) = 1088$.

Time = 187.83 (sec) , antiderivative size = 17988, normalized size of antiderivative = 29.93

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Too large to display}$$

input `integrate(cot(dx+c)^(3/2)*(A+B*tan(dx+c))/(a+b*tan(dx+c))^3,x, algorithm="fracas")`

output Too large to include

3.600. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

3.600.6 Sympy [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx = \int \frac{(A+B \tan(c+dx)) \cot^{\frac{3}{2}}(c+dx)}{(a+b \tan(c+dx))^3} dx$$

input `integrate(cot(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**(3/2)/(a + b*tan(c + d*x))**3, x)`

3.600.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 577, normalized size of antiderivative = 0.96

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx = \frac{(35Ba^5b^2 - 63Aa^4b^3 + 6Ba^3b^4 - 46Aa^2b^5 + 3Bab^6 - 15Ab^7) \arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right) - \frac{2\sqrt{2}((A-B)a^3 + 3(A+B)a^2b - 3(A-B)ab^2 - (A+B)b^3)}{(a^9 + 3a^7b^2 + 3a^5b^4 + a^3b^6)\sqrt{ab}}}{1}$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="maxima")`

output `-1/4*((35*B*a^5*b^2 - 63*A*a^4*b^3 + 6*B*a^3*b^4 - 46*A*a^2*b^5 + 3*B*a*b^6 - 15*A*b^7)*arctan(a/(sqrt(a*b)*sqrt(tan(d*x + c))))/((a^9 + 3*a^7*b^2 + 3*a^5*b^4 + a^3*b^6)*sqrt(a*b)) - (2*sqrt(2))*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2))*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2))*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2))*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - ((11*B*a^3*b^3 - 15*A*a^2*b^4 + 3*B*a*b^5 - 7*A*b^6)/sqrt(tan(d*x + c)) + (13*B*a^4*b^2 - 17*A*a^3*b^3 + 5*B*a^2*b^4 - 9*A*a*b^5)/tan(d*x + c)^(3/2))/(a^7*b^2 + 2*a^5*b^4 + a^3*b^6 + 2*(a^8*b + 2*a^6*b^3 + a^4*b^5)/tan(d*x + c) + (a^9 + 2*a^7*b^2 + a^5*b^4)/tan(d*x + c)^2) + 8*A/(a^3*sqrt(tan(d*x + c))))/d`

3.600. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

3.600.8 Giac [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{(b\tan(dx+c)+a)^3} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(3/2)/(b*tan(d*x + c) + a)^3, x)`

3.600.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Hanged}$$

input `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3,x)`

output `\text{Hanged}`

$$\mathbf{3.601} \quad \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

3.601.1 Optimal result	5730
3.601.2 Mathematica [A] (verified)	5731
3.601.3 Rubi [A] (warning: unable to verify)	5732
3.601.4 Maple [B] (verified)	5740
3.601.5 Fricas [B] (verification not implemented)	5741
3.601.6 Sympy [F]	5742
3.601.7 Maxima [A] (verification not implemented)	5742
3.601.8 Giac [F]	5743
3.601.9 Mupad [F(-1)]	5743

3.601.1 Optimal result

Integrand size = 33, antiderivative size = 534

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx =$$

$$\frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d}$$

$$+ \frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d}$$

$$- \frac{\sqrt{b}(35a^4Ab + 6a^2Ab^3 + 3Ab^5 - 15a^5B + 18a^3b^2B + ab^4B) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{4a^{5/2}(a^2+b^2)^3 d}$$

$$+ \frac{b(Ab - aB) \cot^{\frac{3}{2}}(c+dx)}{2a(a^2+b^2)d(b+a \cot(c+dx))^2} + \frac{b(11a^2Ab + 3Ab^3 - 7a^3B + ab^2B) \sqrt{\cot(c+dx)}}{4a^2(a^2+b^2)^2 d(b+a \cot(c+dx))}$$

$$- \frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d}$$

$$+ \frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d}$$

$$3.601. \quad \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

output $\frac{1}{2} b (A b - B a) \cot(d x + c)^{3/2} / a (a^2 + b^2) / d (b + a \cot(d x + c))^2 + \frac{1}{2} (3 a^2 b (A - B) - b^3 (A - B) - a^3 (A + B) + 3 a b^2 (A + B)) \arctan(-1 + 2^{1/2} \cot(d x + c)^{1/2}) / (a^2 + b^2)^3 / d^{1/2} + \frac{1}{2} (3 a^2 b (A - B) - b^3 (A - B) - a^3 (A + B) + 3 a b^2 (A + B)) \arctan(1 + 2^{1/2} \cot(d x + c)^{1/2}) / (a^2 + b^2)^3 / d^{1/2} - \frac{1}{4} (a^3 (A - B) - 3 a b^2 (A - B) + 3 a^2 b (A + B) - b^3 (A + B)) \ln(1 + \cot(d x + c) - 2^{1/2} \cot(d x + c)^{1/2}) / (a^2 + b^2)^3 / d^{1/2} + \frac{1}{4} (a^3 (A - B) - 3 a b^2 (A - B) + 3 a^2 b (A + B) - b^3 (A + B)) \ln(1 + \cot(d x + c) + 2^{1/2} \cot(d x + c)^{1/2}) / (a^2 + b^2)^3 / d^{1/2} - \frac{1}{4} (35 A a^4 b + 6 A a^2 b^3 + 3 A b^5 - 15 B a^5 + 18 B a^3 b^2 + B a b^4) \arctan(a^{1/2} \cot(d x + c)^{1/2} / b^{1/2}) * b^{1/2} / a^{5/2} / (a^2 + b^2)^3 / d + \frac{1}{4} b (11 A a^2 b + 3 A b^3 - 7 B a^3 + B a b^2) \cot(d x + c)^{1/2} / a^2 / (a^2 + b^2)^2 / d / (b + a \cot(d x + c))$

3.601.2 Mathematica [A] (verified)

Time = 6.47 (sec) , antiderivative size = 581, normalized size of antiderivative = 1.09

$$\int \frac{\sqrt{\cot(c + dx)}(A + B \tan(c + dx))}{(a + b \tan(c + dx))^3} dx$$

$$= \frac{2 \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(\frac{(3 a^2 b (A - B) - b^3 (A - B) - a^3 (A + B) + 3 a b^2 (A + B)) \left(\sqrt{2} \arctan\left(\frac{1 - \sqrt{2} \sqrt{\tan(c + dx)}}{1 + \sqrt{2} \sqrt{\tan(c + dx)}}\right) - \sqrt{2} \arctan\left(\frac{1 + \sqrt{2} \sqrt{\tan(c + dx)}}{1 - \sqrt{2} \sqrt{\tan(c + dx)}}\right)\right)}{4 (a^2 + b^2)^3} \right)}{(a + b \tan(c + dx))^3}$$

input `Integrate[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3, x]`

output $(2 \sqrt{\text{Cot}[c + d x]} \sqrt{\text{Tan}[c + d x]} * (((3 a^2 b (A - B) - b^3 (A - B) - a^3 (A + B) + 3 a b^2 (A + B)) * (\text{Sqrt}[2] * \text{ArcTan}[1 - \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[c + d x]]] - \text{Sqrt}[2] * \text{ArcTan}[1 + \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[c + d x]]]) / (4 * (a^2 + b^2)^3) + (3 \sqrt{b} * (A b - a B) * \text{ArcTan}[(\text{Sqrt}[b] * \text{Sqrt}[\text{Tan}[c + d x]]) / \text{Sqrt}[a]]) / (8 a^{5/2} * (a^2 + b^2)) + (\text{Sqrt}[b] * (2 a A b - a^2 B + b^2 B) * \text{ArcTan}[(\text{Sqrt}[b] * \text{Sqrt}[\text{Tan}[c + d x]]) / \text{Sqrt}[a]]) / (2 a^{3/2} * (a^2 + b^2)^2) + (\text{Sqrt}[b] * (3 a^2 A b - A b^3 - a^3 B + 3 a b^2 B) * \text{ArcTan}[(\text{Sqrt}[b] * \text{Sqrt}[\text{Tan}[c + d x]]) / \text{Sqrt}[a]]) / (\text{Sqrt}[a] * (a^2 + b^2)^3) - ((a^3 (A - B) - 3 a b^2 (A - B) + 3 a^2 b (A + B) - b^3 (A + B)) * (\text{Sqrt}[2] * \text{Log}[1 - \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[c + d x]] + \text{Tan}[c + d x]] - \text{Sqrt}[2] * \text{Log}[1 + \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[c + d x]] + \text{Tan}[c + d x]])) / (8 * (a^2 + b^2)^3) + (b * (A b - a B) * \text{Sqrt}[\text{Tan}[c + d x]]) / (4 a * (a^2 + b^2) * (a + b \text{Tan}[c + d x])^2) + (3 b * (A b - a B) * \text{Sqrt}[\text{Tan}[c + d x]]) / (8 a^2 * (a^2 + b^2) * (a + b \text{Tan}[c + d x])) + (b * (2 a A b - a^2 B + b^2 B) * \text{Sqrt}[\text{Tan}[c + d x]]) / (2 a * (a^2 + b^2)^2 * (a + b \text{Tan}[c + d x])))) / d$

3.601. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

3.601.3 Rubi [A] (warning: unable to verify)

Time = 2.38 (sec) , antiderivative size = 462, normalized size of antiderivative = 0.87, number of steps used = 27, number of rules used = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.788$, Rules used = {3042, 4064, 3042, 4088, 27, 3042, 4128, 27, 3042, 4136, 27, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{\cot^{\frac{5}{2}}(c+dx)(A \cot(c+dx)+B)}{(a \cot(c+dx)+b)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-\tan(c+dx+\frac{\pi}{2}))^{5/2}(B-A \tan(c+dx+\frac{\pi}{2}))}{(b-a \tan(c+dx+\frac{\pi}{2}))^3} dx \\
 & \quad \downarrow \text{4088} \\
 & \frac{b(Ab-aB) \cot^{\frac{3}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2} - \\
 & \frac{\int -\frac{\sqrt{\cot(c+dx)}((4Aa^2+bBa+3Ab^2) \cot^2(c+dx)-4a(Ab-aB) \cot(c+dx)+3b(Ab-aB))}{2(b+a \cot(c+dx))^2} dx}{2a(a^2+b^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\sqrt{\cot(c+dx)}((4Aa^2+bBa+3Ab^2) \cot^2(c+dx)-4a(Ab-aB) \cot(c+dx)+3b(Ab-aB))}{(b+a \cot(c+dx))^2} dx}{4a(a^2+b^2)} + \\
 & \frac{b(Ab-aB) \cot^{\frac{3}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.601. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{\int \frac{\sqrt{-\tan(c+dx+\frac{\pi}{2})} \left((4Aa^2+bBa+3Ab^2) \tan(c+dx+\frac{\pi}{2})^2 + 4a(Ab-aB) \tan(c+dx+\frac{\pi}{2}) + 3b(Ab-aB) \right) dx}{(b-a \tan(c+dx+\frac{\pi}{2}))^2}}{4a(a^2+b^2)} + \frac{b(Ab-aB) \cot^{\frac{3}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

↓ 4128

$$\frac{\frac{b(-7a^3B+11a^2Ab+ab^2B+3Ab^3) \sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a \cot(c+dx)+b)} - \int \frac{-8(-Ba^2+2Aba+b^2B) \cot(c+dx)a^2 + (8Aa^4+9bBa^3+3Ab^2a^2+b^3Ba+3Ab^4) \cot^2(c+dx) + b(-7Ba^3+11Aba^2+b^2Ba+3Ab^3)}{2\sqrt{\cot(c+dx)}(b+a \cot(c+dx))}}{a(a^2+b^2)}}{4a(a^2+b^2)} + \frac{b(Ab-aB) \cot^{\frac{3}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

↓ 27

$$\frac{\int \frac{-8(-Ba^2+2Aba+b^2B) \cot(c+dx)a^2 + (8Aa^4+9bBa^3+3Ab^2a^2+b^3Ba+3Ab^4) \cot^2(c+dx) + b(-7Ba^3+11Aba^2+b^2Ba+3Ab^3)}{\sqrt{\cot(c+dx)}(b+a \cot(c+dx))} dx}{2a(a^2+b^2)} + \frac{b(-7a^3B+11a^2Ab+ab^2B+3Ab^3)}{ad(a^2+b^2)(a \cot(c+dx)+b)}}{4a(a^2+b^2)} + \frac{b(Ab-aB) \cot^{\frac{3}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

↓ 3042

$$\frac{\int \frac{8(-Ba^2+2Aba+b^2B) \tan(c+dx+\frac{\pi}{2})a^2 + (8Aa^4+9bBa^3+3Ab^2a^2+b^3Ba+3Ab^4) \tan(c+dx+\frac{\pi}{2})^2 + b(-7Ba^3+11Aba^2+b^2Ba+3Ab^3)}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx}{2a(a^2+b^2)} + \frac{b(-7a^3B+11a^2Ab+ab^2B+3Ab^3)}{ad(a^2+b^2)(a \cot(c+dx)+b)}}{4a(a^2+b^2)} + \frac{b(Ab-aB) \cot^{\frac{3}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

↓ 4136

$$\frac{\int \frac{8(a^2(-Ba^3+3Aba^2+3b^2Ba-Ab^3)) - a^2(Aa^3+3bBa^2-3Ab^2a-b^3B) \cot(c+dx)}{\sqrt{\cot(c+dx)}} dx}{a^2+b^2} + \frac{b(-15a^5B+35a^4Ab+18a^3b^2B+6a^2Ab^3+ab^4B+3Ab^5)}{a^2+b^2} \int \frac{\cot^2(c+dx)}{\sqrt{\cot(c+dx)}(b+a \cot(c+dx))}}{2a(a^2+b^2)}}{4a(a^2+b^2)} + \frac{b(Ab-aB) \cot^{\frac{3}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

↓ 27

3.601. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{b(-15a^5B+35a^4Ab+18a^3b^2B+6a^2Ab^3+ab^4B+3Ab^5) \int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} dx}{a^2+b^2} - \frac{8 \int \frac{a^2(-Ba^3+3Aba^2+3b^2Ba-Ab^3)-a^2(Aa^3+3bBa^2-3Ab^2a-b^3B)}{\sqrt{\cot(c+dx)}}}{a^2+b^2} dx}{2a(a^2+b^2)}$$

$$4a(a^2+b^2)$$

$$\frac{b(Ab-aB) \cot^{\frac{3}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

↓ 3042

$$\frac{b(-15a^5B+35a^4Ab+18a^3b^2B+6a^2Ab^3+ab^4B+3Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{8 \int \frac{(-Ba^3+3Aba^2+3b^2Ba-Ab^3)a^2+(Aa^3+3bBa^2-3Ab^2a-b^3B)}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}}}{a^2+b^2} dx}{2a(a^2+b^2)}$$

$$4a(a^2+b^2)$$

$$\frac{b(Ab-aB) \cot^{\frac{3}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

↓ 4017

$$\frac{b(-15a^5B+35a^4Ab+18a^3b^2B+6a^2Ab^3+ab^4B+3Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{16 \int \frac{a^2(-Ba^3+3Aba^2+3b^2Ba-Ab^3)-(Aa^3+3bBa^2-3Ab^2a-b^3B) \cot(c+dx)}{\cot^2(c+dx)+1}}{d(a^2+b^2)} dx}{2a(a^2+b^2)}$$

$$4a(a^2+b^2)$$

$$\frac{b(Ab-aB) \cot^{\frac{3}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

↓ 25

$$\frac{16 \int \frac{a^2(-Ba^3+3Aba^2+3b^2Ba-Ab^3)-(Aa^3+3bBa^2-3Ab^2a-b^3B) \cot(c+dx)}{\cot^2(c+dx)+1}}{d(a^2+b^2)} d\sqrt{\cot(c+dx)} + \frac{b(-15a^5B+35a^4Ab+18a^3b^2B+6a^2Ab^3+ab^4B+3Ab^5) \int \frac{1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}}}{a^2+b^2} dx}{2a(a^2+b^2)}$$

$$4a(a^2+b^2)$$

$$\frac{b(Ab-aB) \cot^{\frac{3}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

↓ 27

$$\frac{16a^2 \int \frac{-Ba^3+3Aba^2+3b^2Ba-Ab^3-(Aa^3+3bBa^2-3Ab^2a-b^3B) \cot(c+dx)}{\cot^2(c+dx)+1}}{d(a^2+b^2)} d\sqrt{\cot(c+dx)} + \frac{b(-15a^5B+35a^4Ab+18a^3b^2B+6a^2Ab^3+ab^4B+3Ab^5) \int \frac{1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}}}{a^2+b^2} dx}{2a(a^2+b^2)}$$

$$4a(a^2+b^2)$$

$$\frac{b(Ab-aB) \cot^{\frac{3}{2}}(c+dx)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

3.601. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

↓ 1482

$$\frac{16a^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \int \frac{\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d(a^2+b^2)}$$

$$2a(a^2+b^2)$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 1476

$$\frac{16a^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)} d\sqrt{\cot(c+dx)} \right) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 1082

$$\frac{16a^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \left(\int \frac{1}{-\cot(c+dx)} d\sqrt{\cot(c+dx)} \right) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 217

$$\frac{16a^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \int \frac{1 - \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{\cot(c+dx)}} \right) \right)}{d(a^2+b^2)}$$

$$2a(a^2+b^2)$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 1479

3.601. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$16a^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(-\frac{\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) \right)$$

$$d(a^2+b^2)$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 25

$$16a^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) \right) + \frac{1}{2} \left(\right)$$

$$d(a^2+b^2)$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 27

$$16a^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right) + \frac{1}{2} \left(\right)$$

$$d(a^2+b^2)$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 1103

$$b(-15a^5B + 35a^4Ab + 18a^3b^2B + 6a^2Ab^3 + ab^4B + 3Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx + 16a^2 \left(\frac{1}{2} (-a^3(A+B) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B) + 3ab^2(A-B) - b^3(A+B)) \right)$$

$$a^2+b^2$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 4117

3.601. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

$$\frac{b(-15a^5B+35a^4Ab+18a^3b^2B+6a^2Ab^3+ab^4B+3Ab^5) \int \frac{1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} d(-\cot(c+dx))}{d(a^2+b^2)} + \frac{16a^2 \left(\frac{1}{2}(-a^3(A+B))+3a^2b(A-B)+3ab^2(A+B)-b^3(A-B) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 73

$$\frac{16a^2 \left(\frac{1}{2}(-a^3(A+B))+3a^2b(A-B)+3ab^2(A+B)-b^3(A-B) \right) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) + \frac{1}{2}(a^3(A-B)+3a^2b(A+B)-3ab^2(A-B))}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 218

$$\frac{b(Ab - aB) \cot^{\frac{3}{2}}(c + dx)}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2} +$$

$$\frac{16a^2 \left(\frac{1}{2}(-a^3(A+B))+3a^2b(A-B)+3ab^2(A+B)-b^3(A-B) \right) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right)}{ad(a^2+b^2)(a \cot(c+dx)+b)} + \frac{b(-7a^3B+11a^2Ab+ab^2B+3Ab^3)\sqrt{\cot(c+dx)}}{ad(a^2+b^2)(a \cot(c+dx)+b)}$$

input `Int[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^3,x]`

output `(b*(A*b - a*B)*Cot[c + d*x]^(3/2))/(2*a*(a^2 + b^2)*d*(b + a*Cot[c + d*x])^2) + ((b*(11*a^2*A*b + 3*A*b^3 - 7*a^3*B + a*b^2*B)*Sqrt[Cot[c + d*x]])/(a*(a^2 + b^2)*d*(b + a*Cot[c + d*x])) + ((2*Sqrt[b]*(35*a^4*A*b + 6*a^2*A*b^3 + 3*A*b^5 - 15*a^5*B + 18*a^3*b^2*B + a*b^4*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]])/(Sqrt[a]*(a^2 + b^2)*d) + (16*a^2*((3*a^2*b*(A - B) - b^3*(A - B) - a^3*(A + B) + 3*a*b^2*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]))/2 + ((a^3*(A - B) - 3*a*b^2*(A - B) + 3*a^2*b*(A + B) - b^3*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/(a^2 + b^2)*d)/(2*a*(a^2 + b^2))/(4*a*(a^2 + b^2))`

3.601. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

3.601.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 $\text{Int}[\frac{(d_+) + (e_+)(x_+)^2}{(a_+) + (c_+)(x_+)^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2*(d/e), 2]\}, \text{Simp}[e/(2*c*q) \text{Int}[(q - 2*x)/\text{Simp}[d/e + q*x - x^2, x], x], x] + \text{Simp}[e/(2*c*q) \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{NegQ}[d*e]$

rule 1482 $\text{Int}[\frac{(d_+) + (e_+)(x_+)^2}{(a_+) + (c_+)(x_+)^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[a*c, 2]\}, \text{Simp}[(d*q + a*e)/(2*a*c) \text{Int}[(q + c*x^2)/(a + c*x^4), x], x] + \text{Simp}[(d*q - a*e)/(2*a*c) \text{Int}[(q - c*x^2)/(a + c*x^4), x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{NeQ}[c*d^2 + a*e^2, 0] \&\& \text{NeQ}[c*d^2 - a*e^2, 0] \&\& \text{NegQ}[(a)*c]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4017 $\text{Int}[\frac{(c_+) + (d_+)*\tan[(e_+) + (f_+)(x_+)]}{\text{Sqrt}[(b_+)*\tan[(e_+) + (f_+)(x_+)]}], x_Symbol] \rightarrow \text{Simp}[2/f \text{Subst}[\text{Int}[(b*c + d*x^2)/(b^2 + x^4), x], x, \text{Sqrt}[b*\text{Tan}[e + f*x]]], x] /; \text{FreeQ}[\{b, c, d, e, f\}, x] \&\& \text{NeQ}[c^2 - d^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0]$

rule 4064 $\text{Int}[(\cot[(e_+) + (f_+)(x_+)]*(g_+))^{(p_+)}*((a_+) + (b_+)*\tan[(e_+) + (f_+)(x_+)]^{(m_+)})^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[g^{(m+n)} \text{Int}[(g*\cot[e + f*x])^{(p-m-n)}*(b + a*\cot[e + f*x])^m*(d + c*\cot[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, p\}, x] \&\& !\text{IntegerQ}[p] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n]$

rule 4088 $\text{Int}[\frac{((a_+) + (b_+)*\tan[(e_+) + (f_+)(x_+)])^{(m_+)}*((A_+) + (B_+)*\tan[(e_+) + (f_+)(x_+)])^{(n_+)}}{(c_+) + (d_+)*\tan[(e_+) + (f_+)(x_+)]^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[(b*c - a*d)*(B*c - A*d)*(a + b*\text{Tan}[e + f*x])^{(m-1)}*((c + d*\text{Tan}[e + f*x])^{(n+1)})^{(n+1)}/(d*f*(n+1)*(c^2 + d^2)), x] - \text{Simp}[1/(d*(n+1)*(c^2 + d^2)) \text{Int}[(a + b*\text{Tan}[e + f*x])^{(m-2)}*(c + d*\text{Tan}[e + f*x])^{(n+1)}*\text{Simp}[a*A*d*(b*d*(m-1) - a*c*(n+1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m-1) + a*d*(n+1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n+1)*\text{Tan}[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m+n) - b*B*(c^2*(m-1) - d^2*(n+1)))*\text{Tan}[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{NeQ}[a^2 + b^2, 0] \&\& \text{NeQ}[c^2 + d^2, 0] \&\& \text{GtQ}[m, 1] \&\& \text{LtQ}[n, -1] \&\& (\text{IntegerQ}[m] || \text{IntegersQ}[2*m, 2*n])$

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)^2], x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

```
rule 4128 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)^2] + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2], x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)^2] + (C_.)*tan[(e_.) + (f_.)*(x_)^2]))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.601.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4190 vs. $2(486) = 972$.

Time = 0.40 (sec) , antiderivative size = 4191, normalized size of antiderivative = 7.85

method	result	size
derivativedivides	Expression too large to display	4191
default	Expression too large to display	4191

```
input int(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x,method=_RETURNV
ERBOSE)
```

$$3.601. \quad \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

output $\frac{1}{4}d*(1/\tan(dx+c))^{(1/2)}*\tan(dx+c)^{(1/2)}*(-15*B*\arctan(b*\tan(dx+c))^{(1/2)}/(a*b)^{(1/2)}*a^5*b^3*\tan(dx+c)^2+18*B*\arctan(b*\tan(dx+c))^{(1/2)}/(a*b)^{(1/2)}*a^3*b^5*\tan(dx+c)^2+B*\arctan(b*\tan(dx+c))^{(1/2)}/(a*b)^{(1/2)}*a*b^7*\tan(dx+c)^2-7*B*(a*b)^{(1/2)}*\tan(dx+c)^{(3/2)}*a^5*b^2-6*B*(a*b)^{(1/2)}*\tan(dx+c)^{(3/2)}*a^3*b^4+B*(a*b)^{(1/2)}*\tan(dx+c)^{(3/2)}*a*b^6+70*A*\arctan(b*\tan(dx+c))^{(1/2)}/(a*b)^{(1/2)}*a^5*b^3*\tan(dx+c)+12*A*\arctan(b*\tan(dx+c))^{(1/2)}/(a*b)^{(1/2)}*a^3*b^5*\tan(dx+c)+6*A*\arctan(b*\tan(dx+c))^{(1/2)}/(a*b)^{(1/2)}*a*b^7*\tan(dx+c)+A*\ln(-(1+2^{(1/2)}*\tan(dx+c))^{(1/2)}+\tan(dx+c))/(2^{(1/2)}*\tan(dx+c)^{(1/2)}-\tan(dx+c)-1)*2^{(1/2)}*(a*b)^{(1/2)}*a^7+2*A*\arctan(1+2^{(1/2)}*\tan(dx+c))^{(1/2)}*2^{(1/2)}*(a*b)^{(1/2)}*a^7+2*A*\arctan(-1+2^{(1/2)}*\tan(dx+c))^{(1/2)}*2^{(1/2)}*(a*b)^{(1/2)}*a^7-30*B*\arctan(b*\tan(dx+c))^{(1/2)}/(a*b)^{(1/2)}*a^6*b^2*\tan(dx+c)+36*B*\arctan(b*\tan(dx+c))^{(1/2)}/(a*b)^{(1/2)}*a^4*b^4*\tan(dx+c)+2*B*\arctan(b*\tan(dx+c))^{(1/2)}/(a*b)^{(1/2)}*a^2*b^6*\tan(dx+c)+2*B*\arctan(1+2^{(1/2)}*\tan(dx+c))^{(1/2)}*2^{(1/2)}*(a*b)^{(1/2)}*a^7+2*B*\arctan(-1+2^{(1/2)}*\tan(dx+c))^{(1/2)}*2^{(1/2)}*(a*b)^{(1/2)}*a^7+B*\ln(-(2^{(1/2)}*\tan(dx+c))^{(1/2)}-\tan(dx+c)-1)/(1+2^{(1/2)}*\tan(dx+c))^{(1/2)}+\tan(dx+c))*2^{(1/2)}*(a*b)^{(1/2)}*a^7+13*A*(a*b)^{(1/2)}*\tan(dx+c)^{(1/2)}*a^5*b^2+18*A*(a*b)^{(1/2)}*\tan(dx+c)^{(1/2)}*a^3*b^4+5*A*(a*b)^{(1/2)}*\tan(dx+c)^{(1/2)}*a*b^6-9*B*(a*b)^{(1/2)}*\tan(dx+c)^{(1/2)}*a^6*b^2*B*\arctan(-1+2^{(1/2)}*\tan(dx+c))^{(1/2)}*2^{(1/2)}*(a*b)^{(1/2)}*a^4*b^3-3*B*\ln(-(2^{(1/2)}*\tan(dx+c))^{(1/2)}-\tan(dx+c)+1)/(1+2^{(1/2)}*\tan(dx+c))^{(1/2)}+\tan(dx+c))*2^{(1/2)}*(a*b)^{(1/2)}*a^7-13*A*(a*b)^{(1/2)}*\tan(dx+c)^{(1/2)}*a^5*b^2-18*A*(a*b)^{(1/2)}*\tan(dx+c)^{(1/2)}*a^3*b^4-5*A*(a*b)^{(1/2)}*\tan(dx+c)^{(1/2)}*a*b^6+9*B*(a*b)^{(1/2)}*\tan(dx+c)^{(1/2)}*a^6*b^2*B*\arctan(1+2^{(1/2)}*\tan(dx+c))^{(1/2)}*2^{(1/2)}*(a*b)^{(1/2)}*a^7-2*B*\arctan(-1+2^{(1/2)}*\tan(dx+c))^{(1/2)}*2^{(1/2)}*(a*b)^{(1/2)}*a^7$

3.601.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8921 vs. $2(484) = 968$.

Time = 124.59 (sec) , antiderivative size = 17874, normalized size of antiderivative = 33.47

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \text{Too large to display}$$

input `integrate(cot(dx+c)^(1/2)*(A+B*tan(dx+c))/(a+b*tan(dx+c))^3,x, algorithm="fricas")`

output Too large to include

3.601. $\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$

3.601.6 Sympy [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx = \int \frac{(A+B \tan(c+dx)) \sqrt{\cot(c+dx)}}{(a+b \tan(c+dx))^3} dx$$

input `integrate(cot(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**3,x)`

output `Integral((A + B*tan(c + d*x))*sqrt(cot(c + d*x))/(a + b*tan(c + d*x))**3, x)`

3.601.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 558, normalized size of antiderivative = 1.04

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$$

$$\frac{(15Ba^5b-35Aa^4b^2-18Ba^3b^3-6Aa^2b^4-Bab^5-3Ab^6) \arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right) - 2\sqrt{2}((A+B)a^3-3(A-B)a^2b-3(A+B)ab^2+(A-B)b^3) a}{(a^8+3a^6b^2+3a^4b^4+a^2b^6)\sqrt{ab}}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm="maxima")`

output `1/4*((15*B*a^5*b - 35*A*a^4*b^2 - 18*B*a^3*b^3 - 6*A*a^2*b^4 - B*a*b^5 - 3*A*b^6)*arctan(a/(sqrt(a*b)*sqrt(tan(d*x + c))))/((a^8 + 3*a^6*b^2 + 3*a^4*b^4 + a^2*b^6)*sqrt(a*b)) - (2*sqrt(2)*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - ((7*B*a^3*b^2 - 11*A*a^2*b^3 - B*a*b^4 - 3*A*b^5)/sqrt(tan(d*x + c)) + (9*B*a^4*b - 13*A*a^3*b^2 + B*a^2*b^3 - 5*A*a*b^4)/tan(d*x + c)^(3/2))/(a^6*b^2 + 2*a^4*b^4 + a^2*b^6 + 2*(a^7*b + 2*a^5*b^3 + a^3*b^5)/tan(d*x + c) + (a^8 + 2*a^6*b^2 + a^4*b^4)/tan(d*x + c)^2))/d`

3.601. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^3} dx$

3.601.8 Giac [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \int \frac{(B\tan(dx+c)+A)\sqrt{\cot(dx+c)}}{(b\tan(dx+c)+a)^3} dx$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^3,x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)*sqrt(cot(d*x + c))/(b*tan(d*x + c) + a)^3, x)`

3.601.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx = \int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^3} dx$$

input `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3,x)`

output `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^3, x)`

$$3.602 \quad \int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))^3}} dx$$

3.602.1 Optimal result	5744
3.602.2 Mathematica [A] (verified)	5745
3.602.3 Rubi [A] (warning: unable to verify)	5746
3.602.4 Maple [A] (verified)	5754
3.602.5 Fricas [B] (verification not implemented)	5755
3.602.6 Sympy [F]	5755
3.602.7 Maxima [A] (verification not implemented)	5756
3.602.8 Giac [F]	5756
3.602.9 Mupad [F(-1)]	5757

3.602.1 Optimal result

Integrand size = 33, antiderivative size = 534

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))^3}} dx \\ &= \frac{(a^3(A - B) - 3ab^2(A - B) + 3a^2b(A + B) - b^3(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^3 d} \\ & \quad - \frac{(a^3(A - B) - 3ab^2(A - B) + 3a^2b(A + B) - b^3(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^3 d} \\ & \quad + \frac{(15a^4Ab - 18a^2Ab^3 - Ab^5 - 3a^5B + 26a^3b^2B - 3ab^4B) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{4a^{3/2}\sqrt{b}(a^2 + b^2)^3 d} \\ & \quad + \frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{2a(a^2 + b^2)d(b + a \cot(c + dx))^2} - \frac{(9a^2Ab + Ab^3 - 5a^3B + 3ab^2B)\sqrt{\cot(c + dx)}}{4a(a^2 + b^2)^2d(b + a \cot(c + dx))} \\ & \quad - \frac{(3a^2b(A - B) - b^3(A - B) - a^3(A + B) + 3ab^2(A + B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^3 d} \\ & \quad + \frac{(3a^2b(A - B) - b^3(A - B) - a^3(A + B) + 3ab^2(A + B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^3 d} \end{aligned}$$

output
$$-1/2*(a^3*(A-B)-3*a*b^2*(A-B)+3*a^2*b*(A+B)-b^3*(A+B))*\arctan(-1+2^{(1/2)*\cot(dx+c)^{(1/2)}}/(a^2+b^2)^{3/d*2^{(1/2)}}-1/2*(a^3*(A-B)-3*a*b^2*(A-B)+3*a^2*b*(A+B)-b^3*(A+B))*\arctan(1+2^{(1/2)*\cot(dx+c)^{(1/2)}}/(a^2+b^2)^{3/d*2^{(1/2)}})-1/4*(3*a^2*b*(A-B)-b^3*(A-B)-a^3*(A+B)+3*a*b^2*(A+B))*\ln(1+\cot(dx+c)-2^{(1/2)*\cot(dx+c)^{(1/2)}}/(a^2+b^2)^{3/d*2^{(1/2)}})+1/4*(3*a^2*b*(A-B)-b^3*(A-B)-a^3*(A+B)+3*a*b^2*(A+B))*\ln(1+\cot(dx+c)+2^{(1/2)*\cot(dx+c)^{(1/2)}}/(a^2+b^2)^{3/d*2^{(1/2)}})+1/4*(15*A*a^4*b-18*A*a^2*b^3-A*b^5-3*B*a^5+26*B*a^3*b^2-3*B*a*b^4)*\arctan(a^{(1/2)*\cot(dx+c)^{(1/2)}}/b^{(1/2)})/a^{(3/2)}/(a^2+b^2)^{3/d/b^{(1/2)}}+1/2*b*(A*b-B*a)*\cot(dx+c)^{(1/2)}/a/(a^2+b^2)/d/(b+a*\cot(dx+c))^{-2}-1/4*(9*A*a^2*b+A*b^3-5*B*a^3+3*B*a*b^2)*\cot(dx+c)^{(1/2)}/a/(a^2+b^2)^2/d/(b+a*\cot(dx+c))$$

3.602.2 Mathematica [A] (verified)

Time = 6.44 (sec) , antiderivative size = 576, normalized size of antiderivative = 1.08

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3} dx$$

$$= 2\sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)} \left(-\frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B))(\sqrt{2}\arctan(1 - \sqrt{2}\sqrt{\tan(c+dx)}) - \sqrt{2}\arctan(1 + \sqrt{2}\sqrt{\tan(c+dx)}))}{4(a^2+b^2)^3} \right)$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^3),x]`

output
$$(2*\text{Sqrt}[\text{Cot}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]*(-1/4*((a^3*(A - B) - 3*a*b^2*(A - B) + 3*a^2*b*(A + B) - b^3*(A + B))*(\text{Sqrt}[2]*\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]] - \text{Sqrt}[2]*\text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]]))/(a^2 + b^2)^3 - (3*(A*b - a*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(8*a^{(3/2)}*\text{Sqrt}[b]*(a^2 + b^2)) - (\text{Sqrt}[b]*(a^2*A - A*b^2 + 2*a*b*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(2*a^{(3/2)}*(a^2 + b^2)^2) - (\text{Sqrt}[b]*(a^3*A - 3*a*A*b^2 + 3*a^2*b*B - b^3*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(\text{Sqrt}[a]*(a^2 + b^2)^3) - ((3*a^2*b*(A - B) - b^3*(A - B) - a^3*(A + B) + 3*a*b^2*(A + B))*(\text{Sqrt}[2]*\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]] - \text{Sqrt}[2]*\text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]])))/(8*(a^2 + b^2)^3) - ((A*b - a*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/(4*(a^2 + b^2)*(a + b*\text{Tan}[c + d*x])^2) - (3*(A*b - a*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/(8*a*(a^2 + b^2)*(a + b*\text{Tan}[c + d*x])) - (b*(a^2*A - A*b^2 + 2*a*b*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/(2*a*(a^2 + b^2)^2*(a + b*\text{Tan}[c + d*x])))/d$$

3.602.
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^3} dx$$

3.602.3 Rubi [A] (warning: unable to verify)

Time = 2.36 (sec) , antiderivative size = 459, normalized size of antiderivative = 0.86, number of steps used = 27, number of rules used = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.788$, Rules used = {3042, 4064, 3042, 4088, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3} dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{\cot^{\frac{3}{2}}(c + dx)(A \cot(c + dx) + B)}{(a \cot(c + dx) + b)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(-\tan(c + dx + \frac{\pi}{2}))^{3/2} (B - A \tan(c + dx + \frac{\pi}{2}))}{(b - a \tan(c + dx + \frac{\pi}{2}))^3} dx \\
 & \quad \downarrow \text{4088} \\
 & \frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2} - \frac{\int -\frac{(4Aa^2 + 3bBa + Ab^2)\cot^2(c + dx) - 4a(Ab - aB)\cot(c + dx) + b(Ab - aB)}{2\sqrt{\cot(c + dx)}(b + a \cot(c + dx))^2} dx}{2a(a^2 + b^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{(4Aa^2 + 3bBa + Ab^2)\cot^2(c + dx) - 4a(Ab - aB)\cot(c + dx) + b(Ab - aB)}{\sqrt{\cot(c + dx)}(b + a \cot(c + dx))^2} dx}{4a(a^2 + b^2)} + \frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{(4Aa^2 + 3bBa + Ab^2)\tan(c + dx + \frac{\pi}{2})^2 + 4a(Ab - aB)\tan(c + dx + \frac{\pi}{2}) + b(Ab - aB)}{\sqrt{-\tan(c + dx + \frac{\pi}{2})(b - a \tan(c + dx + \frac{\pi}{2}))^2}} dx}{4a(a^2 + b^2)} + \\
 & \quad \frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2} \\
 & \quad \downarrow \text{4132}
 \end{aligned}$$

3.602. $\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3} dx$

$$\frac{\int \frac{-b(-5Ba^3+9Aba^2+3b^2Ba+Ab^3) \cot^2(c+dx) - 8ab(Aa^2+2bBa-Ab^2) \cot(c+dx) + b(-3Ba^3+7Aba^2+5b^2Ba-Ab^3)}{2\sqrt{\cot(c+dx)}(b+a \cot(c+dx))} dx}{b(a^2+b^2)} - \frac{(-5a^3B+9a^2Ab+3ab^2B+Ab^3)}{d(a^2+b^2)(a \cot(c+dx))}$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

27

$$\frac{\int \frac{-b(-5Ba^3+9Aba^2+3b^2Ba+Ab^3) \cot^2(c+dx) - 8ab(Aa^2+2bBa-Ab^2) \cot(c+dx) + b(-3Ba^3+7Aba^2+5b^2Ba-Ab^3)}{\sqrt{\cot(c+dx)}(b+a \cot(c+dx))} dx}{2b(a^2+b^2)} - \frac{(-5a^3B+9a^2Ab+3ab^2B+Ab^3)}{d(a^2+b^2)(a \cot(c+dx))}$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

3042

$$\frac{\int \frac{-b(-5Ba^3+9Aba^2+3b^2Ba+Ab^3) \tan(c+dx+\frac{\pi}{2})^2 + 8ab(Aa^2+2bBa-Ab^2) \tan(c+dx+\frac{\pi}{2}) + b(-3Ba^3+7Aba^2+5b^2Ba-Ab^3)}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}(b-a \tan(c+dx+\frac{\pi}{2}))} dx}{2b(a^2+b^2)} - \frac{(-5a^3B+9a^2Ab+3ab^2B+Ab^3)}{d(a^2+b^2)(a \cot(c+dx))}$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

4136

$$\frac{\int \frac{8(ab(Aa^3+3bBa^2-3Ab^2a-b^3B)+ab(-Ba^3+3Aba^2+3b^2Ba-Ab^3) \cot(c+dx))}{\sqrt{\cot(c+dx)}} dx}{a^2+b^2} + \frac{b(-3a^5B+15a^4Ab+26a^3b^2B-18a^2Ab^3-3ab^4B-Ab^5) \int \frac{\cot^2(c+dx)}{\sqrt{\cot(c+dx)}} dx}{a^2+b^2}$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

27

$$\frac{b(-3a^5B+15a^4Ab+26a^3b^2B-18a^2Ab^3-3ab^4B-Ab^5) \int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)}(b+a \cot(c+dx))} dx}{a^2+b^2} - 8 \int \frac{ab(Aa^3+3bBa^2-3Ab^2a-b^3B)+ab(-Ba^3+3Aba^2+3b^2Ba-Ab^3)}{\sqrt{\cot(c+dx)}} dx}{a^2+b^2}$$

$$\frac{4a(a^2+b^2)}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2} \frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

3042

3.602. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^3} dx$

$$\frac{b(-3a^5B+15a^4Ab+26a^3b^2B-18a^2Ab^3-3ab^4B-Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{8 \int \frac{ab(Aa^3+3bBa^2-3Ab^2a-b^3B)-ab(-Ba^3+3Aba^2)}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{a^2+b^2}$$

$$\frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

$4a(a^2+b^2)$

↓ 4017

$$\frac{b(-3a^5B+15a^4Ab+26a^3b^2B-18a^2Ab^3-3ab^4B-Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{16 \int \frac{ab(Aa^3+3bBa^2-3Ab^2a-b^3B+(-Ba^3+3Aba^2+3b^2Ba-Ab^3)\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)}$$

$$\frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

$4a(a^2+b^2)$

↓ 25

$$\frac{16 \int \frac{ab(Aa^3+3bBa^2-3Ab^2a-b^3B+(-Ba^3+3Aba^2+3b^2Ba-Ab^3)\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} + \frac{b(-3a^5B+15a^4Ab+26a^3b^2B-18a^2Ab^3-3ab^4B-Ab^5) \int \frac{1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{a^2+b^2}$$

$$\frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

$4a(a^2+b^2)$

↓ 27

$$\frac{16ab \int \frac{Aa^3+3bBa^2-3Ab^2a-b^3B+(-Ba^3+3Aba^2+3b^2Ba-Ab^3)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} + \frac{b(-3a^5B+15a^4Ab+26a^3b^2B-18a^2Ab^3-3ab^4B-Ab^5) \int \frac{1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{a^2+b^2}$$

$$\frac{b(Ab-aB)\sqrt{\cot(c+dx)}}{2ad(a^2+b^2)(a \cot(c+dx)+b)^2}$$

$4a(a^2+b^2)$

↓ 1482

3.602. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))^3}} dx$

$$\frac{16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d(a^2+b^2)}$$

$$2b(a^2+b^2)$$

$$\frac{b(Ab - aB)\sqrt{\cot(c+dx)}}{2ad(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 1476

$$\frac{16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\cot(c+dx)}}{2ad(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 1082

$$\frac{16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)-1} d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right) - \frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\cot(c+dx)}}{2ad(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 217

$$\frac{16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right) - \frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)}$$

$$2b(a^2+b^2)$$

$$\frac{b(Ab - aB)\sqrt{\cot(c+dx)}}{2ad(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 1479

3.602. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^3} dx$

$$16ab \left(\frac{\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} (-a^3(A+B) + 3a^2b(A-B) + 3ab^2(A-B) - b^3(A+B))}{d(a^2+b^2)} \right)$$

$$\frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 25

$$16ab \left(\frac{\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} (-a^3(A+B) + 3a^2b(A-B) + 3ab^2(A-B) - b^3(A+B))}{d(a^2+b^2)} \right)$$

$$\frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 27

$$16ab \left(\frac{\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} (-a^3(A+B) + 3a^2b(A-B) + 3ab^2(A-B) - b^3(A+B))}{d(a^2+b^2)} \right)$$

$$\frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 1103

$$\frac{b(-3a^5B + 15a^4Ab + 26a^3b^2B - 18a^2Ab^3 - 3ab^4B - Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} + \frac{16ab \left(\frac{\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} (-a^3(A+B) + 3a^2b(A-B) + 3ab^2(A-B) - b^3(A+B))}{d(a^2+b^2)} \right)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 4117

3.602. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))^3}} dx$

$$\frac{b(-3a^5B+15a^4Ab+26a^3b^2B-18a^2Ab^3-3ab^4B-Ab^5)}{d(a^2+b^2)} \int \frac{1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} d(-\cot(c+dx)) + \frac{16ab}{d(a^2+b^2)} \left(\frac{1}{2} (a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right)$$

$$\frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 73

$$\frac{16ab \left(\frac{1}{2} (a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)}+1}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) - \frac{1}{2} (-a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B)}{d(a^2+b^2)}$$

$$\frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 218

$$\frac{b(Ab - aB)\sqrt{\cot(c + dx)}}{2ad(a^2 + b^2)(a \cot(c + dx) + b)^2} +$$

$$\frac{16ab \left(\frac{1}{2} (a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right) \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)}+1}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) - \frac{(-5a^3B+9a^2Ab+3ab^2B+Ab^3)\sqrt{\cot(c+dx)}}{d(a^2+b^2)(a \cot(c+dx)+b)}}{d(a^2+b^2)}$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^3),x]`

output `(b*(A*b - a*B)*Sqrt[Cot[c + d*x]])/(2*a*(a^2 + b^2)*d*(b + a*Cot[c + d*x])^2) + (-(((9*a^2*A*b + A*b^3 - 5*a^3*B + 3*a*b^2*B)*Sqrt[Cot[c + d*x]])/((a^2 + b^2)*d*(b + a*Cot[c + d*x]))) - ((2*Sqrt[b]*(15*a^4*A*b - 18*a^2*A*b^3 - A*b^5 - 3*a^5*B + 26*a^3*b^2*B - 3*a*b^4*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]])/(Sqrt[a]*(a^2 + b^2)*d) + (16*a*b*((a^3*(A - B) - 3*a*b^2*(A - B) + 3*a^2*b*(A + B) - b^3*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]))/2 - ((3*a^2*b*(A - B) - b^3*(A - B) - a^3*(A + B) + 3*a*b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d))/(2*b*(a^2 + b^2)))/(4*a*(a^2 + b^2))`

3.602. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))^3}} dx$

3.602.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`


```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4136 Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^
n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
!GtQ[n, 0] && !LeQ[n, -1]
```

3.602.4 Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 451, normalized size of antiderivative = 0.84

method	result
derivativedivides	$\frac{2 \left(\left(-\frac{9}{8} A a^4 b - \frac{5}{4} A a^2 b^3 - \frac{1}{8} A b^5 + \frac{5}{8} B a^5 + \frac{1}{4} B a^3 b^2 - \frac{3}{8} B a b^4 \right) \cot(dx+c) \frac{3}{2} - \frac{b(7A a^4 b + 6A a^2 b^3 - A b^5 - 3B a^5 + 2B a^3 b^2 + 5B a b^4) \sqrt{a^2 + b^2}}{8a} \right)}{(b+a \cot(dx+c))^2 (a^2+b^2)^3}$
default	$\frac{2 \left(\left(-\frac{9}{8} A a^4 b - \frac{5}{4} A a^2 b^3 - \frac{1}{8} A b^5 + \frac{5}{8} B a^5 + \frac{1}{4} B a^3 b^2 - \frac{3}{8} B a b^4 \right) \cot(dx+c) \frac{3}{2} - \frac{b(7A a^4 b + 6A a^2 b^3 - A b^5 - 3B a^5 + 2B a^3 b^2 + 5B a b^4) \sqrt{a^2 + b^2}}{8a} \right)}{(b+a \cot(dx+c))^2 (a^2+b^2)^3}$

3.602. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))^3}} dx$

```
input int((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^3,x,method=_RETURNV
ERBOSE)
```

```
output 1/d*(2/(a^2+b^2)^3*((( -9/8*A*a^4*b-5/4*A*a^2*b^3-1/8*A*b^5+5/8*B*a^5+1/4*B
*a^3*b^2-3/8*B*a*b^4)*cot(d*x+c)^(3/2)-1/8*b*(7*A*a^4*b+6*A*a^2*b^3-A*b^5-
3*B*a^5+2*B*a^3*b^2+5*B*a*b^4)/a*cot(d*x+c)^(1/2))/(b+a*cot(d*x+c))^2+1/8*
(15*A*a^4*b-18*A*a^2*b^3-A*b^5-3*B*a^5+26*B*a^3*b^2-3*B*a*b^4)/a/(a*b)^(1/
2)*arctan(a*cot(d*x+c)^(1/2)/(a*b)^(1/2)))-2/(a^2+b^2)^3*(1/8*(A*a^3-3*A*a
*b^2+3*B*a^2*b-B*b^3)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/
(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/
2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/8*(3*A*a^2*b-A*b^3-B*a^3+3*B*
a*b^2)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2
^(1/2)*cot(d*x+c)^(1/2))))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1
+2^(1/2)*cot(d*x+c)^(1/2))))))
```

3.602.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8942 vs. $2(482) = 964$.

Time = 85.90 (sec) , antiderivative size = 17912, normalized size of antiderivative = 33.54

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^3,x, algorith
m="fricas")
```

```
output Too large to include
```

3.602.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^3 \sqrt{\cot(c + dx)}} dx$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)**(1/2)/(a+b*tan(d*x+c))**3,x)
```

```
output Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))**3*sqrt(cot(c + d*x)))
, x)
```

3.602. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^3} dx$

3.602.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 544, normalized size of antiderivative = 1.02

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3} dx = \frac{(3Ba^5 - 15Aa^4b - 26Ba^3b^2 + 18Aa^2b^3 + 3Bab^4 + Ab^5) \arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right)}{(a^7 + 3a^5b^2 + 3a^3b^4 + ab^6)\sqrt{ab}} + \frac{2\sqrt{2}((A-B)a^3 + 3(A+B)a^2b - 3(A-B)ab^2 - (A+B)b^3)}{(a^7 + 3a^5b^2 + 3a^3b^4 + ab^6)\sqrt{ab}}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^3,x, algorithm="maxima")`

output `-1/4*((3*B*a^5 - 15*A*a^4*b - 26*B*a^3*b^2 + 18*A*a^2*b^3 + 3*B*a*b^4 + A*b^5)*arctan(a/(sqrt(a*b)*sqrt(tan(d*x + c))))/((a^7 + 3*a^5*b^2 + 3*a^3*b^4 + a*b^6)*sqrt(a*b)) + (2*sqrt(2)*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - ((3*B*a^3*b - 7*A*a^2*b^2 - 5*B*a*b^3 + A*b^4)/sqrt(tan(d*x + c)) + (5*B*a^4 - 9*A*a^3*b - 3*B*a^2*b^2 - A*a*b^3)/tan(d*x + c)^(3/2))/(a^5*b^2 + 2*a^3*b^4 + a*b^6 + 2*(a^6*b + 2*a^4*b^3 + a^2*b^5)/tan(d*x + c) + (a^7 + 2*a^5*b^2 + a^3*b^4)/tan(d*x + c)^2))/d`

3.602.8 Giac [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a)^3 \sqrt{\cot(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^3,x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)/((b*tan(d*x + c) + a)^3*sqrt(cot(d*x + c))), x)`

3.602. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^3} dx$

3.602.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^3} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))^3),x)`output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))^3), x)`

3.603 $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

3.603.1 Optimal result	5758
3.603.2 Mathematica [A] (verified)	5759
3.603.3 Rubi [A] (warning: unable to verify)	5760
3.603.4 Maple [A] (verified)	5768
3.603.5 Fricas [B] (verification not implemented)	5769
3.603.6 Sympy [F(-1)]	5770
3.603.7 Maxima [A] (verification not implemented)	5770
3.603.8 Giac [F(-1)]	5771
3.603.9 Mupad [F(-1)]	5771

3.603.1 Optimal result

Integrand size = 33, antiderivative size = 530

$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$$

$$= \frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d}$$

$$- \frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d}$$

$$- \frac{(3a^4Ab - 26a^2Ab^3 + 3Ab^5 + a^5B + 18a^3b^2B - 15ab^4B) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{4\sqrt{ab}^{3/2}(a^2+b^2)^3 d}$$

$$- \frac{(Ab - aB)\sqrt{\cot(c+dx)}}{2(a^2+b^2)d(b+a \cot(c+dx))^2} + \frac{(5a^2Ab - 3Ab^3 - a^3B + 7ab^2B)\sqrt{\cot(c+dx)}}{4b(a^2+b^2)^2 d(b+a \cot(c+dx))}$$

$$+ \frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d}$$

$$- \frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d}$$

3.603. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

output
$$-1/2*(3*a^2*b*(A-B)-b^3*(A-B)-a^3*(A+B)+3*a*b^2*(A+B))*\arctan(-1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)^3/d*2^{(1/2)}-1/2*(3*a^2*b*(A-B)-b^3*(A-B)-a^3*(A+B)+3*a*b^2*(A+B))*\arctan(1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)^3/d*2^{(1/2)}+1/4*(a^3*(A-B)-3*a*b^2*(A-B)+3*a^2*b*(A+B)-b^3*(A+B))*\ln(1+\cot(d*x+c)-2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)^3/d*2^{(1/2)}-1/4*(a^3*(A-B)-3*a*b^2*(A-B)+3*a^2*b*(A+B)-b^3*(A+B))*\ln(1+\cot(d*x+c)+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)^3/d*2^{(1/2)}-1/4*(3*A*a^4*b-26*A*a^2*b^3+3*A*b^5+B*a^5+18*B*a^3*b^2-15*B*a*b^4)*\arctan(a^{(1/2)}*\cot(d*x+c)^{(1/2)}/b^{(1/2)})/b^{(3/2)}/(a^2+b^2)^3/d/a^{(1/2)}-1/2*(A*b-B*a)*\cot(d*x+c)^{(1/2)}/(a^2+b^2)/d/(b+a*\cot(d*x+c))^2+1/4*(5*A*a^2*b-3*A*b^3-B*a^3+7*B*a*b^2)*\cot(d*x+c)^{(1/2)}/b/(a^2+b^2)^2/d/(b+a*\cot(d*x+c))$$

3.603.2 Mathematica [A] (verified)

Time = 6.46 (sec) , antiderivative size = 586, normalized size of antiderivative = 1.11

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx$$

$$= \frac{2\sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}\left(-\frac{(3a^2b(A-B)-b^3(A-B)-a^3(A+B)+3ab^2(A+B))(\sqrt{2}\arctan(1-\sqrt{2}\sqrt{\tan(c+dx)})-\sqrt{2}\arctan(1+\sqrt{2}\sqrt{\tan(c+dx)}))}{4(a^2+b^2)^3}\right)}{1}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^3),x]`

output
$$(2*\text{Sqrt}[\text{Cot}[c + d*x]]*\text{Sqrt}[\text{Tan}[c + d*x]]*(-1/4*((3*a^2*b*(A - B) - b^3*(A - B) - a^3*(A + B) + 3*a*b^2*(A + B))*(\text{Sqrt}[2]*\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]] - \text{Sqrt}[2]*\text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]]]))/(a^2 + b^2)^3 + (3*(A*b - a*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(8*\text{Sqrt}[a]*b^{(3/2)}*(a^2 + b^2)) - (\text{Sqrt}[b]*(3*a^2*A*b - A*b^3 - a^3*B + 3*a*b^2*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(\text{Sqrt}[a]*(a^2 + b^2)^3) - ((2*A*b^3 - a*(a^2 + 3*b^2)*B)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a]])/(2*\text{Sqrt}[a]*b^{(3/2)}*(a^2 + b^2)^2) + ((a^3*(A - B) - 3*a*b^2*(A - B) + 3*a^2*b*(A + B) - b^3*(A + B))*(\text{Sqrt}[2]*\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]] - \text{Sqrt}[2]*\text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[c + d*x]] + \text{Tan}[c + d*x]])))/(8*(a^2 + b^2)^3) + (a*(A*b - a*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/(4*b*(a^2 + b^2)*(a + b*\text{Tan}[c + d*x])^2) + (3*(A*b - a*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/(8*b*(a^2 + b^2)*(a + b*\text{Tan}[c + d*x])) - ((2*A*b^3 - a*(a^2 + 3*b^2)*B)*\text{Sqrt}[\text{Tan}[c + d*x]])/(2*b*(a^2 + b^2)^2*(a + b*\text{Tan}[c + d*x])))/d$$

3.603.
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$$

3.603.3 Rubi [A] (warning: unable to verify)

Time = 2.38 (sec) , antiderivative size = 459, normalized size of antiderivative = 0.87, number of steps used = 27, number of rules used = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.788$, Rules used = {3042, 4064, 3042, 4091, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2}(a + b \tan(c + dx))^3} dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{\sqrt{\cot(c + dx)}(A \cot(c + dx) + B)}{(a \cot(c + dx) + b)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(B - A \tan(c + dx + \frac{\pi}{2}))}{(b - a \tan(c + dx + \frac{\pi}{2}))^3} dx \\
 & \quad \downarrow \text{4091} \\
 & \frac{\int -\frac{-3a(Ab - aB) \cot^2(c + dx) - 4a(aA + bB) \cot(c + dx) + a(Ab - aB)}{2\sqrt{\cot(c + dx)}(b + a \cot(c + dx))^2} dx}{2a(a^2 + b^2)} - \frac{(Ab - aB)\sqrt{\cot(c + dx)}}{2d(a^2 + b^2)(a \cot(c + dx) + b)^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{-3a(Ab - aB) \cot^2(c + dx) - 4a(aA + bB) \cot(c + dx) + a(Ab - aB)}{\sqrt{\cot(c + dx)}(b + a \cot(c + dx))^2} dx}{4a(a^2 + b^2)} - \frac{(Ab - aB)\sqrt{\cot(c + dx)}}{2d(a^2 + b^2)(a \cot(c + dx) + b)^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{-3a(Ab - aB) \tan(c + dx + \frac{\pi}{2})^2 + 4a(aA + bB) \tan(c + dx + \frac{\pi}{2}) + a(Ab - aB)}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(b - a \tan(c + dx + \frac{\pi}{2}))^2} dx}{4a(a^2 + b^2)} - \frac{(Ab - aB)\sqrt{\cot(c + dx)}}{2d(a^2 + b^2)(a \cot(c + dx) + b)^2} \\
 & \quad \downarrow \text{4132}
 \end{aligned}$$

3.603. $\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx$

$$\begin{aligned}
 & \int \frac{-a(-Ba^3+5Aba^2+7b^2Ba-3Ab^3)\cot^2(c+dx)+8ab(-Ba^2+2Aba+b^2B)\cot(c+dx)+a(Ba^3+3Aba^2+9b^2Ba-5Ab^3)}{2\sqrt{\cot(c+dx)(b+a\cot(c+dx))}} dx - \frac{a(a^3(-B)+5a^2Ab+7ab^2B)}{bd(a^2+b^2)(a\cot(c+dx)+b)} \\
 & \frac{4a(a^2+b^2)}{2d(a^2+b^2)(a\cot(c+dx)+b)^2} \\
 & \quad \downarrow \text{27} \\
 & \int \frac{-a(-Ba^3+5Aba^2+7b^2Ba-3Ab^3)\cot^2(c+dx)+8ab(-Ba^2+2Aba+b^2B)\cot(c+dx)+a(Ba^3+3Aba^2+9b^2Ba-5Ab^3)}{\sqrt{\cot(c+dx)(b+a\cot(c+dx))}} dx - \frac{a(a^3(-B)+5a^2Ab+7ab^2B)}{bd(a^2+b^2)(a\cot(c+dx)+b)} \\
 & \frac{4a(a^2+b^2)}{2d(a^2+b^2)(a\cot(c+dx)+b)^2} \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{-a(-Ba^3+5Aba^2+7b^2Ba-3Ab^3)\tan(c+dx+\frac{\pi}{2})^2-8ab(-Ba^2+2Aba+b^2B)\tan(c+dx+\frac{\pi}{2})+a(Ba^3+3Aba^2+9b^2Ba-5Ab^3)}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx - \frac{a(a^3(-B)+5a^2Ab+7ab^2B)}{bd(a^2+b^2)(a\cot(c+dx)+b)} \\
 & \frac{4a(a^2+b^2)}{2d(a^2+b^2)(a\cot(c+dx)+b)^2} \\
 & \quad \downarrow \text{4136} \\
 & \int \frac{8(ab(-Ba^3+3Aba^2+3b^2Ba-Ab^3)-ab(Aa^3+3bBa^2-3Ab^2a-b^3B)\cot(c+dx))}{\sqrt{\cot(c+dx)}} dx + \frac{a(a^5B+3a^4Ab+18a^3b^2B-26a^2Ab^3-15ab^4B+3Ab^5)}{a^2+b^2} \int \frac{\cot^2(c+dx)}{\sqrt{\cot(c+dx)(b+a\cot(c+dx))}} dx \\
 & \frac{4a(a^2+b^2)}{2b(a^2+b^2)} \\
 & \frac{4a(a^2+b^2)}{2d(a^2+b^2)(a\cot(c+dx)+b)^2} \\
 & \quad \downarrow \text{27} \\
 & 8 \int \frac{ab(-Ba^3+3Aba^2+3b^2Ba-Ab^3)-ab(Aa^3+3bBa^2-3Ab^2a-b^3B)\cot(c+dx)}{\sqrt{\cot(c+dx)}} dx + \frac{a(a^5B+3a^4Ab+18a^3b^2B-26a^2Ab^3-15ab^4B+3Ab^5)}{a^2+b^2} \int \frac{\cot^2(c+dx)}{\sqrt{\cot(c+dx)(b+a\cot(c+dx))}} dx \\
 & \frac{4a(a^2+b^2)}{2b(a^2+b^2)} \\
 & \frac{4a(a^2+b^2)}{2d(a^2+b^2)(a\cot(c+dx)+b)^2} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

3.603. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

$$\frac{8 \int \frac{ab(-Ba^3+3Aba^2+3b^2Ba-Ab^3)+ab(Aa^3+3bBa^2-3Ab^2a-b^3B) \tan(c+dx+\frac{\pi}{2})}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{a^2+b^2} + \frac{a(a^5B+3a^4Ab+18a^3b^2B-26a^2Ab^3-15ab^4B+3Ab^5) \int \frac{1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{a^2+b^2}$$

$$4a(a^2+b^2)$$

$$\frac{(Ab-aB)\sqrt{\cot(c+dx)}}{2d(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 4017

$$\frac{16 \int \frac{ab(-Ba^3+3Aba^2+3b^2Ba-Ab^3)-(Aa^3+3bBa^2-3Ab^2a-b^3B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} + \frac{a(a^5B+3a^4Ab+18a^3b^2B-26a^2Ab^3-15ab^4B+3Ab^5) \int \frac{1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{a^2+b^2}$$

$$2b(a^2+b^2)$$

$$4a(a^2+b^2)$$

$$\frac{(Ab-aB)\sqrt{\cot(c+dx)}}{2d(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 25

$$\frac{a(a^5B+3a^4Ab+18a^3b^2B-26a^2Ab^3-15ab^4B+3Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{16 \int \frac{ab(-Ba^3+3Aba^2+3b^2Ba-Ab^3)-(Aa^3+3bBa^2-3Ab^2a-b^3B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)}$$

$$2b(a^2+b^2)$$

$$4a(a^2+b^2)$$

$$\frac{(Ab-aB)\sqrt{\cot(c+dx)}}{2d(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 27

$$\frac{a(a^5B+3a^4Ab+18a^3b^2B-26a^2Ab^3-15ab^4B+3Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{16ab \int \frac{-Ba^3+3Aba^2+3b^2Ba-Ab^3-(Aa^3+3bBa^2-3Ab^2a-b^3B) \cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)}$$

$$2b(a^2+b^2)$$

$$4a(a^2+b^2)$$

$$\frac{(Ab-aB)\sqrt{\cot(c+dx)}}{2d(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 1482

3.603. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

$$\frac{a(a^5 B + 3a^4 Ab + 18a^3 b^2 B - 26a^2 Ab^3 - 15ab^4 B + 3Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2 + b^2} - 16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2 b(A+B) - 3ab^2(A-B) - b^3(A+B)) \right)$$

$$2b(a^2 + b^2)$$

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{2d(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 1476

$$\frac{a(a^5 B + 3a^4 Ab + 18a^3 b^2 B - 26a^2 Ab^3 - 15ab^4 B + 3Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2 + b^2} - 16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2 b(A+B) - 3ab^2(A-B) - b^3(A+B)) \right)$$

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{2d(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 1082

$$\frac{a(a^5 B + 3a^4 Ab + 18a^3 b^2 B - 26a^2 Ab^3 - 15ab^4 B + 3Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2 + b^2} - 16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2 b(A+B) - 3ab^2(A-B) - b^3(A+B)) \right)$$

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{2d(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 217

$$\frac{a(a^5 B + 3a^4 Ab + 18a^3 b^2 B - 26a^2 Ab^3 - 15ab^4 B + 3Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a \tan(c+dx+\frac{\pi}{2}))}} dx}{a^2 + b^2} - 16ab \left(\frac{1}{2} (a^3(A-B) + 3a^2 b(A+B) - 3ab^2(A-B) - b^3(A+B)) \right)$$

$$2b(a^2 + b^2)$$

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{2d(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 1479

3.603. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

$$\frac{a(a^5B+3a^4Ab+18a^3b^2B-26a^2Ab^3-15ab^4B+3Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} \quad 16ab \left(\frac{1}{2} (a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right)$$

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{2d(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 25

$$\frac{a(a^5B+3a^4Ab+18a^3b^2B-26a^2Ab^3-15ab^4B+3Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} \quad 16ab \left(\frac{1}{2} (a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right)$$

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{2d(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 27

$$\frac{a(a^5B+3a^4Ab+18a^3b^2B-26a^2Ab^3-15ab^4B+3Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} \quad 16ab \left(\frac{1}{2} (a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right)$$

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{2d(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 1103

$$\frac{a(a^5B+3a^4Ab+18a^3b^2B-26a^2Ab^3-15ab^4B+3Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} \quad 16ab \left(\frac{1}{2} (-a^3(A+B)+3a^2b(A-B)+3ab^2(A+B)-b^3(A-B)) \right)$$

$$\frac{(Ab - aB)\sqrt{\cot(c + dx)}}{2d(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 4117

3.603. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

$$\frac{a(a^5 B + 3a^4 Ab + 18a^3 b^2 B - 26a^2 Ab^3 - 15ab^4 B + 3Ab^5) \int \frac{1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} d(-\cot(c+dx))}{d(a^2+b^2)} - \frac{16ab \left(\frac{1}{2} (-a^3(A+B)) + 3a^2 b(A-B) + 3ab^2(A+B) - b^3(A-B) \right)}{d(a^2+b^2)}$$

$$\frac{(Ab - aB) \sqrt{\cot(c+dx)}}{2d(a^2+b^2)(a \cot(c+dx) + b)^2}$$

↓ 73

$$\frac{2a(a^5 B + 3a^4 Ab + 18a^3 b^2 B - 26a^2 Ab^3 - 15ab^4 B + 3Ab^5) \int \frac{1}{a \cot^2(c+dx) + b} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} - \frac{16ab \left(\frac{1}{2} (-a^3(A+B)) + 3a^2 b(A-B) + 3ab^2(A+B) - b^3(A-B) \right)}{d(a^2+b^2)}$$

$$\frac{(Ab - aB) \sqrt{\cot(c+dx)}}{2d(a^2+b^2)(a \cot(c+dx) + b)^2}$$

↓ 218

$$\frac{(Ab - aB) \sqrt{\cot(c+dx)}}{2d(a^2+b^2)(a \cot(c+dx) + b)^2} - \frac{2\sqrt{a}(a^5 B + 3a^4 Ab + 18a^3 b^2 B - 26a^2 Ab^3 - 15ab^4 B + 3Ab^5) \arctan\left(\frac{\sqrt{a} \cot(c+dx)}{\sqrt{b}}\right) - 16ab \left(\frac{1}{2} (-a^3(A+B)) + 3a^2 b(A-B) + 3ab^2(A+B) - b^3(A-B) \right)}{bd(a^2+b^2)(a \cot(c+dx) + b) \sqrt{b} d(a^2+b^2)}$$

input `Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^3),x]`

output `-1/2*((A*b - a*B)*Sqrt[Cot[c + d*x]])/((a^2 + b^2)*d*(b + a*Cot[c + d*x])^2) - ((a*(5*a^2*A*b - 3*A*b^3 - a^3*B + 7*a*b^2*B)*Sqrt[Cot[c + d*x]])/(b*(a^2 + b^2)*d*(b + a*Cot[c + d*x]))) - ((2*Sqrt[a]*(3*a^4*A*b - 26*a^2*A*b^3 + 3*A*b^5 + a^5*B + 18*a^3*b^2*B - 15*a*b^4*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]])/(Sqrt[b]*(a^2 + b^2)*d) - (16*a*b*((3*a^2*b*(A - B) - b^3*(A - B) - a^3*(A + B) + 3*a*b^2*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2])))/2 + ((a^3*(A - B) - 3*a*b^2*(A - B) + 3*a^2*b*(A + B) - b^3*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)/(2*b*(a^2 + b^2))/(4*a*(a^2 + b^2))`

3.603. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

3.603.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4091 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2), x_Symbol] :=
Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] +
Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /;
FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] :=
Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] +
Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /;
FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.603.4 Maple [A] (verified)

Time = 0.57 (sec) , antiderivative size = 450, normalized size of antiderivative = 0.85

method	result
derivativedivides	$\frac{2 \left(\frac{a(5A^4b+2Aa^2b^3-3Ab^5-Ba^5+6Ba^3b^2+7Bab^4) \cot(dx+c)^{\frac{3}{2}}}{8b} + \frac{(-\frac{3}{8}Aa^4b+\frac{5}{8}Ab^5-\frac{5}{4}Ba^3b^2-\frac{9}{8}Bab^4+\frac{1}{4}Aa^2b^3-\frac{1}{8}Ba^5)}{(b+a \cot(dx+c))^2} \right)}{(a^2+b^2)^3}$
default	$\frac{2 \left(\frac{a(5A^4b+2Aa^2b^3-3Ab^5-Ba^5+6Ba^3b^2+7Bab^4) \cot(dx+c)^{\frac{3}{2}}}{8b} + \frac{(-\frac{3}{8}Aa^4b+\frac{5}{8}Ab^5-\frac{5}{4}Ba^3b^2-\frac{9}{8}Bab^4+\frac{1}{4}Aa^2b^3-\frac{1}{8}Ba^5)}{(b+a \cot(dx+c))^2} \right)}{(a^2+b^2)^3}$

```
input int((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^3,x,method=_RETURNV
ERBOSE)
```

```
output 1/d*(-2/(a^2+b^2)^3*((-1/8*a*(5*A*a^4*b+2*A*a^2*b^3-3*A*b^5-B*a^5+6*B*a^3*
b^2+7*B*a*b^4)/b*cot(d*x+c)^(3/2)+(-3/8*A*a^4*b+5/8*A*b^5-5/4*B*a^3*b^2-9/
8*B*a*b^4+1/4*A*a^2*b^3-1/8*B*a^5)*cot(d*x+c)^(1/2))/(b+a*cot(d*x+c))^2+1/
8*(3*A*a^4*b-26*A*a^2*b^3+3*A*b^5+B*a^5+18*B*a^3*b^2-15*B*a*b^4)/b/(a*b)^(
1/2)*arctan(a*cot(d*x+c)^(1/2)/(a*b)^(1/2)))-2/(a^2+b^2)^3*(1/8*(3*A*a^2*b
-A*b^3-B*a^3+3*B*a*b^2)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)
)/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(
1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/8*(-A*a^3+3*A*a*b^2-3*B*a^2
*b+B*b^3)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)
)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan
(-1+2^(1/2)*cot(d*x+c)^(1/2))))
```

3.603.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8942 vs. 2(477) = 954.

Time = 73.40 (sec) , antiderivative size = 17911, normalized size of antiderivative = 33.79

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^3,x, algorith
m="fricas")
```

```
output Too large to include
```

3.603. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

3.603.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(3/2)/(a+b*tan(d*x+c))**3,x)`

output `Timed out`

3.603.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 545, normalized size of antiderivative = 1.03

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \frac{(Ba^5 + 3Aa^4b + 18Ba^3b^2 - 26Aa^2b^3 - 15Bab^4 + 3Ab^5) \arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right) - 2\sqrt{2}((A+B)a^3 - 3(A-B)a^2b - 3(A+B)ab^2 + (A-B)b^3)}{(a^6b + 3a^4b^3 + 3a^2b^5 + b^7)\sqrt{ab}}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^3,x, algorithm="maxima")`

output `-1/4*((B*a^5 + 3*A*a^4*b + 18*B*a^3*b^2 - 26*A*a^2*b^3 - 15*B*a*b^4 + 3*A*b^5)*arctan(a/(sqrt(a*b)*sqrt(tan(d*x + c))))/((a^6*b + 3*a^4*b^3 + 3*a^2*b^5 + b^7)*sqrt(a*b)) - (2*sqrt(2)*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) - ((B*a^3*b + 3*A*a^2*b^2 + 9*B*a*b^3 - 5*A*b^4)/sqrt(tan(d*x + c)) - (B*a^4 - 5*A*a^3*b - 7*B*a^2*b^2 + 3*A*a*b^3)/tan(d*x + c)^(3/2))/(a^4*b^3 + 2*a^2*b^5 + b^7 + 2*(a^5*b^2 + 2*a^3*b^4 + a*b^6)/tan(d*x + c) + (a^6*b + 2*a^4*b^3 + a^2*b^5)/tan(d*x + c)^2))/d`

3.603.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^3,x, algorithm m="giac")`

output `Timed out`

3.603.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2} (a + b \tan(c + dx))^3} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))^3),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))^3), x)`

3.604 $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

3.604.1 Optimal result	5772
3.604.2 Mathematica [A] (verified)	5773
3.604.3 Rubi [A] (warning: unable to verify)	5774
3.604.4 Maple [A] (verified)	5782
3.604.5 Fricas [B] (verification not implemented)	5783
3.604.6 Sympy [F(-1)]	5784
3.604.7 Maxima [A] (verification not implemented)	5784
3.604.8 Giac [F(-1)]	5785
3.604.9 Mupad [F(-1)]	5785

3.604.1 Optimal result

Integrand size = 33, antiderivative size = 534

$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3} dx =$$

$$\frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d}$$

$$+ \frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}(a^2+b^2)^3 d}$$

$$- \frac{\sqrt{a}(a^4Ab + 18a^2Ab^3 - 15Ab^5 + 3a^5B + 6a^3b^2B + 35ab^4B) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{4b^{5/2}(a^2+b^2)^3 d}$$

$$+ \frac{a(Ab - aB)\sqrt{\cot(c+dx)}}{2b(a^2+b^2)d(b+a \cot(c+dx))^2} - \frac{a(a^2Ab - 7Ab^3 + 3a^3B + 11ab^2B)\sqrt{\cot(c+dx)}}{4b^2(a^2+b^2)^2d(b+a \cot(c+dx))}$$

$$+ \frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d}$$

$$- \frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}(a^2+b^2)^3 d}$$

3.604. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

output $\frac{1}{2}(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \arctan\left(\frac{-1 + \sqrt{2} \cot(dx+c)^{1/2}}{\sqrt{a^2+b^2}}\right) + \frac{1}{2}(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \arctan\left(\frac{1 + \sqrt{2} \cot(dx+c)^{1/2}}{\sqrt{a^2+b^2}}\right) + \frac{1}{4}(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \ln\left(\frac{1 + \cot(dx+c)^{1/2}}{\sqrt{a^2+b^2}}\right) - \frac{1}{4}(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \ln\left(\frac{1 + \cot(dx+c)^{1/2}}{\sqrt{a^2+b^2}}\right) - \frac{1}{4}(Aa^4b + 18Aa^2b^3 - 15Ab^5 + 3Bb^5 + 6Ba^3b^2 + 35Bab^4) \arctan\left(\frac{a^{1/2} \cot(dx+c)^{1/2}}{b^{1/2}}\right) + \frac{a^{1/2}}{b^{5/2}} \sqrt{\frac{a^2+b^2}{d}} + \frac{1}{2} \frac{a(b-Ba) \cot(dx+c)^{1/2}}{b \sqrt{a^2+b^2}} + \frac{1}{4} \frac{a^2(Aa^2b - 7Ab^3 + 3Bb^3 + 11Bab^2) \cot(dx+c)^{1/2}}{b^2 \sqrt{a^2+b^2}} + \frac{1}{4} \frac{a^2(Aa^2b - 7Ab^3 + 3Bb^3 + 11Bab^2) \cot(dx+c)^{1/2}}{b^2 \sqrt{a^2+b^2}}$

3.604.2 Mathematica [A] (verified)

Time = 6.53 (sec) , antiderivative size = 610, normalized size of antiderivative = 1.14

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^3} dx$$

$$= \frac{2\sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)} \left(\frac{(a^3(A-B) - 3ab^2(A-B) + 3a^2b(A+B) - b^3(A+B)) \left(\sqrt{2} \arctan\left(\frac{1 - \sqrt{2}\sqrt{\tan(c+dx)}}{1 + \sqrt{2}\sqrt{\tan(c+dx)}}\right) - \sqrt{2} \arctan\left(\frac{1 + \sqrt{2}\sqrt{\tan(c+dx)}}{1 - \sqrt{2}\sqrt{\tan(c+dx)}}\right)\right)}{4(a^2+b^2)^3} \right)}{d}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^3),x]`

output $(2\sqrt{\cot[c + dx]}\sqrt{\tan[c + dx]} * (((a^3(A - B) - 3ab^2(A - B) + 3a^2b(A + B) - b^3(A + B)) * (\sqrt{2} \operatorname{ArcTan}[1 - \sqrt{2}\sqrt{\tan[c + dx]}] - \sqrt{2} \operatorname{ArcTan}[1 + \sqrt{2}\sqrt{\tan[c + dx]}]]) / (4(a^2 + b^2)^3) - (3\sqrt{a}(Ab - aB) \operatorname{ArcTan}[(\sqrt{b}\sqrt{\tan[c + dx]})/\sqrt{a}]) / (8b^{5/2}(a^2 + b^2)) + (\sqrt{a}(a^2Ab + 3Aab^3 - 2a^3B - 4ab^2B) \operatorname{ArcTan}[(\sqrt{b}\sqrt{\tan[c + dx]})/\sqrt{a}]) / (2b^{5/2}(a^2 + b^2)^2) + (\sqrt{a}(a^2Ab^3 - 3Aab^5 + a^5B + 3a^3b^2B + 6ab^4B) \operatorname{ArcTan}[(\sqrt{b}\sqrt{\tan[c + dx]})/\sqrt{a}]) / (b^{5/2}(a^2 + b^2)^3) + ((3a^2b(A - B) - b^3(A - B) - a^3(A + B) + 3ab^2(A + B)) * (\sqrt{2} \operatorname{Log}[1 - \sqrt{2}\sqrt{\tan[c + dx]}] + \operatorname{Tan}[c + dx] - \sqrt{2} \operatorname{Log}[1 + \sqrt{2}\sqrt{\tan[c + dx]}] + \operatorname{Tan}[c + dx])) / (8(a^2 + b^2)^3) - (a^2(Ab - aB) \sqrt{\tan[c + dx]}) / (4b^2(a^2 + b^2)(a + b \tan[c + dx])^2) - (3a(Ab - aB) \sqrt{\tan[c + dx]}) / (8b^2(a^2 + b^2)(a + b \tan[c + dx])) + (a(a^2Ab + 3Aab^3 - 2a^3B - 4ab^2B) \sqrt{\tan[c + dx]}) / (2b^2(a^2 + b^2)^2(a + b \tan[c + dx])))) / d$

$$3.604. \quad \int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$$

3.604.3 Rubi [A] (warning: unable to verify)

Time = 2.40 (sec) , antiderivative size = 463, normalized size of antiderivative = 0.87, number of steps used = 27, number of rules used = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.788$, Rules used = {3042, 4064, 3042, 4092, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{5/2}(a + b \tan(c + dx))^3} dx \\
 & \quad \downarrow \text{4064} \\
 & \int \frac{A \cot(c + dx) + B}{\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{B - A \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(b - a \tan(c + dx + \frac{\pi}{2}))^3} dx \\
 & \quad \downarrow \text{4092} \\
 & \frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2} - \frac{\int -\frac{3Ba^2 - 3(Ab - aB)\cot^2(c + dx)a + Aba + 4b^2B + 4b(Ab - aB)\cot(c + dx)}{2\sqrt{\cot(c + dx)}(b + a \cot(c + dx))^2} dx}{2b(a^2 + b^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{3Ba^2 - 3(Ab - aB)\cot^2(c + dx)a + Aba + 4b^2B + 4b(Ab - aB)\cot(c + dx)}{\sqrt{\cot(c + dx)}(b + a \cot(c + dx))^2} dx}{4b(a^2 + b^2)} + \frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int \frac{3Ba^2 - 3(Ab - aB)\tan(c + dx + \frac{\pi}{2})^2 a + Aba + 4b^2B - 4b(Ab - aB)\tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(b - a \tan(c + dx + \frac{\pi}{2}))^2} dx}{4b(a^2 + b^2)} + \\
 & \quad \frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2} \\
 & \quad \downarrow \text{4132}
 \end{aligned}$$

3.604. $\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^3} dx$

$$\int \frac{3Ba^4 + Aba^3 + 3b^2Ba^2 + 9Ab^3a + (3Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3) \cot^2(c+dx)a + 8b^4B - 8b^2(Aa^2 + 2bBa - Ab^2) \cot(c+dx)}{2\sqrt{\cot(c+dx)(b+a \cot(c+dx))} b(a^2+b^2)} dx - \frac{a(3a^3B + a^2Ab + 11ab^2B - 7Ab^3)}{bd(a^2+b^2)(a \cot(c+dx) + b)}$$

$$\frac{4b(a^2 + b^2)}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2} \frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

27

$$\int \frac{3Ba^4 + Aba^3 + 3b^2Ba^2 + 9Ab^3a + (3Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3) \cot^2(c+dx)a + 8b^4B - 8b^2(Aa^2 + 2bBa - Ab^2) \cot(c+dx)}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))} 2b(a^2+b^2)} dx - \frac{a(3a^3B + a^2Ab + 11ab^2B - 7Ab^3)}{bd(a^2+b^2)(a \cot(c+dx) + b)}$$

$$\frac{4b(a^2 + b^2)}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2} \frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

3042

$$\int \frac{3Ba^4 + Aba^3 + 3b^2Ba^2 + 9Ab^3a + (3Ba^3 + Aba^2 + 11b^2Ba - 7Ab^3) \tan(c+dx + \frac{\pi}{2})^2 a + 8b^4B + 8b^2(Aa^2 + 2bBa - Ab^2) \tan(c+dx + \frac{\pi}{2})}{\sqrt{-\tan(c+dx + \frac{\pi}{2})(b-a \tan(c+dx + \frac{\pi}{2}))} 2b(a^2+b^2)} dx - \frac{a(3a^3B + a^2Ab + 11ab^2B - 7Ab^3)}{bd(a^2+b^2)(a \cot(c+dx) + b)}$$

$$\frac{4b(a^2 + b^2)}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2} \frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

4136

$$\int \frac{8((Aa^3 + 3bBa^2 - 3Ab^2a - b^3B)b^2 + (-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \cot(c+dx)b^2)}{\sqrt{\cot(c+dx)} \frac{a^2+b^2}} dx + \frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\cot^2(c+dx)}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} dx}{a^2+b^2}$$

$$\frac{4b(a^2 + b^2)}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2} \frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

27

$$\frac{a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\cot^2(c+dx)+1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} dx}{a^2+b^2} - \frac{8 \int \frac{(Aa^3 + 3bBa^2 - 3Ab^2a - b^3B)b^2 + (-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) \cot(c+dx)}{\sqrt{\cot(c+dx)}} dx}{a^2+b^2}$$

$$\frac{4b(a^2 + b^2)}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2} \frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

3042

3.604. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

$$\frac{a(3a^5B+a^4Ab+6a^3b^2B+18a^2Ab^3+35ab^4B-15Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{8 \int \frac{b^2(Aa^3+3bBa^2-3Ab^2a-b^3B)-b^2(-Ba^3+3Aba^2+3b^2Ba-Ab^3) \cot(c+dx)}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{a^2+b^2}$$

$$\frac{4b(a^2+b^2)}{2b(a^2+b^2)}$$

$$\frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{2bd(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 4017

$$\frac{a(3a^5B+a^4Ab+6a^3b^2B+18a^2Ab^3+35ab^4B-15Ab^5) \int \frac{\tan(c+dx+\frac{\pi}{2})^2+1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})(b-a\tan(c+dx+\frac{\pi}{2}))}} dx}{a^2+b^2} - \frac{16 \int \frac{b^2(Aa^3+3bBa^2-3Ab^2a-b^3B+(-Ba^3+3Aba^2+3b^2Ba-Ab^3)\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)}$$

$$\frac{4b(a^2+b^2)}{2b(a^2+b^2)}$$

$$\frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{2bd(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 25

$$\frac{16 \int \frac{b^2(Aa^3+3bBa^2-3Ab^2a-b^3B+(-Ba^3+3Aba^2+3b^2Ba-Ab^3)\cot(c+dx))}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} + \frac{a(3a^5B+a^4Ab+6a^3b^2B+18a^2Ab^3+35ab^4B-15Ab^5) \int \frac{1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{a^2+b^2}$$

$$\frac{4b(a^2+b^2)}{2b(a^2+b^2)}$$

$$\frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{2bd(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 27

$$\frac{16b^2 \int \frac{Aa^3+3bBa^2-3Ab^2a-b^3B+(-Ba^3+3Aba^2+3b^2Ba-Ab^3)\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d(a^2+b^2)} + \frac{a(3a^5B+a^4Ab+6a^3b^2B+18a^2Ab^3+35ab^4B-15Ab^5) \int \frac{1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx}{a^2+b^2}$$

$$\frac{4b(a^2+b^2)}{2b(a^2+b^2)}$$

$$\frac{a(Ab-aB)\sqrt{\cot(c+dx)}}{2bd(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 1482

3.604. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

$$\frac{16b^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)-1} - \frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)+1} \right)}{d(a^2+b^2)}$$

$$2b(a^2+b^2)$$

$$\frac{a(Ab - aB)\sqrt{\cot(c+dx)}}{2bd(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 1476

$$\frac{16b^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{1}{2} \int \frac{1}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)+1} + \frac{1}{2} \int \frac{1}{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)+1} \right) - \frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)+1} \right)}{d(a^2+b^2)}$$

$$\frac{a(Ab - aB)\sqrt{\cot(c+dx)}}{2bd(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 1082

$$\frac{16b^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)-1} d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) - \frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)+1} \right)}{d(a^2+b^2)}$$

$$\frac{a(Ab - aB)\sqrt{\cot(c+dx)}}{2bd(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 217

$$\frac{16b^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)) \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)+1} \right)}{d(a^2+b^2)}$$

$$2b(a^2+b^2)$$

$$\frac{a(Ab - aB)\sqrt{\cot(c+dx)}}{2bd(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 1479

3.604. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

$$16b^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B)) \right)$$

$$d(a^2+b^2)$$

$$\frac{a(Ab - aB)\sqrt{\cot(c+dx)}}{2bd(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 25

$$16b^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B)) \right)$$

$$d(a^2+b^2)$$

$$\frac{a(Ab - aB)\sqrt{\cot(c+dx)}}{2bd(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 27

$$16b^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} (-(a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B)) \right)$$

$$d(a^2+b^2)$$

$$\frac{a(Ab - aB)\sqrt{\cot(c+dx)}}{2bd(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 1103

$$a(3a^5B + a^4Ab + 6a^3b^2B + 18a^2Ab^3 + 35ab^4B - 15Ab^5) \int \frac{\tan(c+dx + \frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c+dx + \frac{\pi}{2})(b - a\tan(c+dx + \frac{\pi}{2}))}} dx + \frac{16b^2 \left(\frac{1}{2} (a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \right)}{a^2+b^2}$$

$$\frac{a(Ab - aB)\sqrt{\cot(c+dx)}}{2bd(a^2+b^2)(a\cot(c+dx)+b)^2}$$

↓ 4117

3.604. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

$$\frac{a(3a^5B+a^4Ab+6a^3b^2B+18a^2Ab^3+35ab^4B-15Ab^5) \int \frac{1}{\sqrt{\cot(c+dx)(b+a \cot(c+dx))}} d(-\cot(c+dx))}{d(a^2+b^2)} + \frac{16b^2 \left(\frac{1}{2} (a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \right)}{d(a^2+b^2)}$$

$$\frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 73

$$\frac{16b^2 \left(\frac{1}{2} (a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\cot(c+dx)}+1}{\sqrt{2}})}{\sqrt{2}} - \frac{\arctan(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}})}{\sqrt{2}} \right) \right) - \frac{1}{2} (-a^3(A+B)+3a^2b(A-B)+3ab^2(A+B))}{d(a^2+b^2)}$$

$$\frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2}$$

↓ 218

$$\frac{a(Ab - aB)\sqrt{\cot(c + dx)}}{2bd(a^2 + b^2)(a \cot(c + dx) + b)^2} + \frac{16b^2 \left(\frac{1}{2} (a^3(A-B)+3a^2b(A+B)-3ab^2(A-B)-b^3(A+B)) \left(\frac{\arctan(\frac{\sqrt{2}\sqrt{\cot(c+dx)}+1}{\sqrt{2}})}{\sqrt{2}} - \frac{\arctan(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}})}{\sqrt{2}} \right) \right) - \frac{1}{2} (-a^3(A+B)+3a^2b(A-B)+3ab^2(A+B))}{d(a^2+b^2)}$$

input `Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^3),x]`

output `(a*(A*b - a*B)*Sqrt[Cot[c + d*x]])/(2*b*(a^2 + b^2)*d*(b + a*Cot[c + d*x])^2) + (-((a*(a^2*A*b - 7*A*b^3 + 3*a^3*B + 11*a*b^2*B)*Sqrt[Cot[c + d*x]])/(b*(a^2 + b^2)*d*(b + a*Cot[c + d*x]))) + ((2*Sqrt[a]*(a^4*A*b + 18*a^2*A*b^3 - 15*A*b^5 + 3*a^5*B + 6*a^3*b^2*B + 35*a*b^4*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]])/(Sqrt[b]*(a^2 + b^2)*d) + (16*b^2*((a^3*(A - B) - 3*a*b^2*(A - B) + 3*a^2*b*(A + B) - b^3*(A + B))*(-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]))/2 - (((3*a^2*b*(A - B) - b^3*(A - B) - a^3*(A + B) + 3*a*b^2*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2))/((a^2 + b^2)*d))/(2*b*(a^2 + b^2)))/(4*b*(a^2 + b^2))`

3.604. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

3.604.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2), x_Symbol] :=
Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] +
Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /;
FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4136 `Int((((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)^2])/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] :=
Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x], x] +
Simp[(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /;
FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && !GtQ[n, 0] && !LeQ[n, -1]`

3.604.4 Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 451, normalized size of antiderivative = 0.84

method	result
derivativedivides	$2a \left(\frac{a(Aa^4b - 6Aa^2b^3 - 7Ab^5 + 3Ba^5 + 14Ba^3b^2 + 11Bab^4) \cot(dx+c)^{\frac{3}{2}} - (Aa^4b + 10Aa^2b^3 + 9Ab^5 - 5Ba^5 - 18Ba^3b^2 - 13Bab^4)}{8b^2} \right) \frac{1}{(b+a \cot(dx+c))^2} - \frac{1}{(a^2+b^2)^3}$
default	$2a \left(\frac{a(Aa^4b - 6Aa^2b^3 - 7Ab^5 + 3Ba^5 + 14Ba^3b^2 + 11Bab^4) \cot(dx+c)^{\frac{3}{2}} - (Aa^4b + 10Aa^2b^3 + 9Ab^5 - 5Ba^5 - 18Ba^3b^2 - 13Bab^4)}{8b^2} \right) \frac{1}{(b+a \cot(dx+c))^2} - \frac{1}{(a^2+b^2)^3}$

```
input int((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c))^3,x,method=_RETURNV
ERBOSE)
```

```
output 1/d*(-2*a/(a^2+b^2)^3*((1/8*a*(A*a^4*b-6*A*a^2*b^3-7*A*b^5+3*B*a^5+14*B*a^
3*b^2+11*B*a*b^4)/b^2*cot(d*x+c)^(3/2)-1/8*(A*a^4*b+10*A*a^2*b^3+9*A*b^5-5
*B*a^5-18*B*a^3*b^2-13*B*a*b^4)/b*cot(d*x+c)^(1/2))/(b+a*cot(d*x+c))^2+1/8
*(A*a^4*b+18*A*a^2*b^3-15*A*b^5+3*B*a^5+6*B*a^3*b^2+35*B*a*b^4)/b^2/(a*b)^(
1/2)*arctan(a*cot(d*x+c)^(1/2)/(a*b)^(1/2)))-2/(a^2+b^2)^3*(1/8*(-A*a^3+3
*A*a*b^2-3*B*a^2*b+B*b^3)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/
2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)
^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/8*(-3*A*a^2*b+A*b^3+B*a^3
-3*B*a*b^2)*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x
+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arct
an(-1+2^(1/2)*cot(d*x+c)^(1/2))))
```

3.604.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8923 vs. 2(482) = 964.

Time = 106.83 (sec) , antiderivative size = 17872, normalized size of antiderivative = 33.47

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c))^3,x, algorith
m="fricas")
```

```
output Too large to include
```

3.604. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

3.604.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(5/2)/(a+b*tan(d*x+c))**3,x)`

output `Timed out`

3.604.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 556, normalized size of antiderivative = 1.04

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \frac{(3Ba^6 + Aa^5b + 6Ba^4b^2 + 18Aa^3b^3 + 35Ba^2b^4 - 15Aab^5) \arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right) - 2\sqrt{2}((A-B)a^3 + 3(A+B)a^2b - 3(A-B)ab^2 - (A+B)b^3)}{(a^6b^2 + 3a^4b^4 + 3a^2b^6 + b^8)\sqrt{ab}}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c))^3,x, algorithm="maxima")`

output `-1/4*((3*B*a^6 + A*a^5*b + 6*B*a^4*b^2 + 18*A*a^3*b^3 + 35*B*a^2*b^4 - 15*A*a*b^5)*arctan(a/(sqrt(a*b)*sqrt(tan(d*x + c))))/((a^6*b^2 + 3*a^4*b^4 + 3*a^2*b^6 + b^8)*sqrt(a*b)) - (2*sqrt(2)*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2 + (A - B)*b^3)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + ((5*B*a^4*b - A*a^3*b^2 + 13*B*a^2*b^3 - 9*A*a*b^4)/sqrt(tan(d*x + c)) + (3*B*a^5 + A*a^4*b + 11*B*a^3*b^2 - 7*A*a^2*b^3)/tan(d*x + c)^(3/2))/(a^4*b^4 + 2*a^2*b^6 + b^8 + 2*(a^5*b^3 + 2*a^3*b^5 + a*b^7)/tan(d*x + c) + (a^6*b^2 + 2*a^4*b^4 + a^2*b^6)/tan(d*x + c)^2))/d`

3.604. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

3.604.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c))^3,x, algorithm m="giac")`

output `Timed out`

3.604.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{5/2} (a + b \tan(c + dx))^3} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + b*tan(c + d*x))^3),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + b*tan(c + d*x))^3), x)`

3.605 $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

3.605.1 Optimal result	5786
3.605.2 Mathematica [A] (verified)	5787
3.605.3 Rubi [A] (warning: unable to verify)	5788
3.605.4 Maple [A] (verified)	5797
3.605.5 Fricas [B] (verification not implemented)	5798
3.605.6 Sympy [F(-1)]	5798
3.605.7 Maxima [A] (verification not implemented)	5799
3.605.8 Giac [F(-1)]	5799
3.605.9 Mupad [F(-1)]	5800

3.605.1 Optimal result

Integrand size = 33, antiderivative size = 600

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3} dx =$$

$$\frac{(3a^2b(A - B) - b^3(A - B) - a^3(A + B) + 3ab^2(A + B)) \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^3 d}$$

$$+ \frac{(3a^2b(A - B) - b^3(A - B) - a^3(A + B) + 3ab^2(A + B)) \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}(a^2 + b^2)^3 d}$$

$$- \frac{a^{3/2}(3a^4Ab + 6a^2Ab^3 + 35Ab^5 - 15a^5B - 46a^3b^2B - 63ab^4B) \arctan\left(\frac{\sqrt{a}\sqrt{\cot(c+dx)}}{\sqrt{b}}\right)}{4b^{7/2}(a^2 + b^2)^3 d}$$

$$- \frac{3a^3Ab + 11aAb^3 - 15a^4B - 31a^2b^2B - 8b^4B}{4b^3(a^2 + b^2)^2 d \sqrt{\cot(c + dx)}}$$

$$+ \frac{a(Ab - aB)}{2b(a^2 + b^2) d \sqrt{\cot(c + dx)}(b + a \cot(c + dx))^2}$$

$$+ \frac{a(a^2Ab + 9Ab^3 - 5a^3B - 13ab^2B)}{4b^2(a^2 + b^2)^2 d \sqrt{\cot(c + dx)}(b + a \cot(c + dx))}$$

$$- \frac{(a^3(A - B) - 3ab^2(A - B) + 3a^2b(A + B) - b^3(A + B)) \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^3 d}$$

$$+ \frac{(a^3(A - B) - 3ab^2(A - B) + 3a^2b(A + B) - b^3(A + B)) \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}(a^2 + b^2)^3 d}$$

3.605. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

output

$$\begin{aligned}
& -1/4*a^{(3/2)}*(3*A*a^4*b+6*A*a^2*b^3+35*A*b^5-15*B*a^5-46*B*a^3*b^2-63*B*a*b^4)*\arctan(a^{(1/2)}*\cot(d*x+c)^{(1/2)}/b^{(1/2)})/b^{(7/2)}/(a^2+b^2)^3/d+1/2*(3 \\
& *a^2*b*(A-B)-b^3*(A-B)-a^3*(A+B)+3*a*b^2*(A+B))*\arctan(-1+2^{(1/2)}*\cot(d*x+ \\
& c)^{(1/2)})/(a^2+b^2)^3/d*2^{(1/2)}+1/2*(3*a^2*b*(A-B)-b^3*(A-B)-a^3*(A+B)+3*a \\
& *b^2*(A+B))*\arctan(1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)^3/d*2^{(1/2)}-1/4*(\\
& a^3*(A-B)-3*a*b^2*(A-B)+3*a^2*b*(A+B)-b^3*(A+B))*\ln(1+\cot(d*x+c)-2^{(1/2)}*c \\
& \cot(d*x+c)^{(1/2)})/(a^2+b^2)^3/d*2^{(1/2)}+1/4*(a^3*(A-B)-3*a*b^2*(A-B)+3*a^2* \\
& b*(A+B)-b^3*(A+B))*\ln(1+\cot(d*x+c)+2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(a^2+b^2)^3/d \\
& *2^{(1/2)}+1/4*(-3*A*a^3*b-11*A*a*b^3+15*B*a^4+31*B*a^2*b^2+8*B*b^4)/b^3/(a^ \\
& 2+b^2)^2/d/\cot(d*x+c)^{(1/2)}+1/2*a*(A*b-B*a)/b/(a^2+b^2)/d/(b+a*\cot(d*x+c)) \\
& ^2/\cot(d*x+c)^{(1/2)}+1/4*a*(A*a^2*b+9*A*b^3-5*B*a^3-13*B*a*b^2)/b^2/(a^2+b^ \\
& 2)^2/d/(b+a*\cot(d*x+c))/\cot(d*x+c)^{(1/2)}
\end{aligned}$$

3.605.2 Mathematica [A] (verified)

Time = 6.58 (sec) , antiderivative size = 639, normalized size of antiderivative = 1.06

$$\begin{aligned}
& \int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3} dx \\
& = \frac{2\sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)} \left(\frac{(3a^2b(A-B) - b^3(A-B) - a^3(A+B) + 3ab^2(A+B)) \left(\sqrt{2} \arctan\left(\frac{1 - \sqrt{2}\sqrt{\tan(c+dx)}}{1 + \sqrt{2}\sqrt{\tan(c+dx)}} \right) - \sqrt{2} \arctan\left(\frac{1 + \sqrt{2}\sqrt{\tan(c+dx)}}{1 - \sqrt{2}\sqrt{\tan(c+dx)}} \right) \right)}{4(a^2 + b^2)^3} \right)}{1}
\end{aligned}$$

input

```
Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(7/2)*(a + b*Tan[c + d*x])^3),x]
```

output $(2\sqrt{\cot[c + dx]}\sqrt{\tan[c + dx]}*((3a^2b(A - B) - b^3(A - B) - a^3(A + B) + 3ab^2(A + B))*(\sqrt{2}\operatorname{ArcTan}[1 - \sqrt{2}\sqrt{\tan[c + dx]}] - \sqrt{2}\operatorname{ArcTan}[1 + \sqrt{2}\sqrt{\tan[c + dx]}]))/ (4(a^2 + b^2)^3) + (3a^{3/2}(Ab - aB)\operatorname{ArcTan}[(\sqrt{b}\sqrt{\tan[c + dx]})/\sqrt{a}]) / (8b^{7/2}(a^2 + b^2)) - (a^{3/2}(2a^2Ab + 4Ab^3 - 3a^3B - 5ab^2B)\operatorname{ArcTan}[(\sqrt{b}\sqrt{\tan[c + dx]})/\sqrt{a}]) / (2b^{7/2}(a^2 + b^2)^2) + (a^{3/2}(a^4Ab + 3a^2Ab^3 + 6Ab^5 - 3a^5B - 9a^3b^2B - 10ab^4B)\operatorname{ArcTan}[(\sqrt{b}\sqrt{\tan[c + dx]})/\sqrt{a}]) / (b^{7/2}(a^2 + b^2)^3) - ((a^3(A - B) - 3ab^2(A - B) + 3a^2b(A + B) - b^3(A + B))*(\sqrt{2}\operatorname{Log}[1 - \sqrt{2}\sqrt{\tan[c + dx]} + \tan[c + dx]] - \sqrt{2}\operatorname{Log}[1 + \sqrt{2}\sqrt{\tan[c + dx]} + \tan[c + dx]])) / (8(a^2 + b^2)^3) + (B\sqrt{\tan[c + dx]})/b^3 + (a^3(Ab - aB)\sqrt{\tan[c + dx]}) / (4b^3(a^2 + b^2)(a + b\tan[c + dx])^2) + (3a^2(Ab - aB)\sqrt{\tan[c + dx]}) / (8b^3(a^2 + b^2)(a + b\tan[c + dx])) - (a^2(2a^2Ab + 4Ab^3 - 3a^3B - 5ab^2B)\sqrt{\tan[c + dx]}) / (2b^3(a^2 + b^2)^2(a + b\tan[c + dx])))/d$

3.605.3 Rubi [A] (warning: unable to verify)

Time = 3.31 (sec) , antiderivative size = 524, normalized size of antiderivative = 0.87, number of steps used = 30, number of rules used = 29, $\frac{\text{number of rules}}{\text{integrand size}} = 0.879$, Rules used = {3042, 4064, 3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4136, 27, 3042, 4017, 25, 27, 1482, 1476, 1082, 217, 1479, 25, 27, 1103, 4117, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{7/2}(a + b \tan(c + dx))^3} dx \\ & \quad \downarrow \text{4064} \\ & \int \frac{A \cot(c + dx) + B}{\cot^{\frac{3}{2}}(c + dx)(a \cot(c + dx) + b)^3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{B - A \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}(b - a \tan(c + dx + \frac{\pi}{2}))^3} dx \end{aligned}$$

3.605. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

$$\begin{aligned}
 & \downarrow 4092 \\
 & \frac{a(Ab - aB)}{2bd(a^2 + b^2) \sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \int \frac{-5Ba^2 + 5(Ab - aB) \cot^2(c + dx)a + Aba - 4b^2B - 4b(Ab - aB) \cot(c + dx)}{2 \cot^{\frac{3}{2}}(c + dx)(b + a \cot(c + dx))^2} dx \\
 & \frac{2bd(a^2 + b^2)}{2b(a^2 + b^2)} \\
 & \downarrow 27 \\
 & \frac{a(Ab - aB)}{2bd(a^2 + b^2) \sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \int \frac{-5Ba^2 + 5(Ab - aB) \cot^2(c + dx)a + Aba - 4b^2B - 4b(Ab - aB) \cot(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(b + a \cot(c + dx))^2} dx \\
 & \frac{2bd(a^2 + b^2)}{4b(a^2 + b^2)} \\
 & \downarrow 3042 \\
 & \frac{a(Ab - aB)}{2bd(a^2 + b^2) \sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \int \frac{-5Ba^2 + 5(Ab - aB) \tan(c + dx + \frac{\pi}{2})^2 a + Aba - 4b^2B + 4b(Ab - aB) \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}(b - a \tan(c + dx + \frac{\pi}{2}))^2} dx \\
 & \frac{2bd(a^2 + b^2)}{4b(a^2 + b^2)} \\
 & \downarrow 4132 \\
 & \frac{a(Ab - aB)}{2bd(a^2 + b^2) \sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \int \frac{-15Ba^4 + 3Aba^3 - 31b^2Ba^2 + 11Ab^3a + 3(-5Ba^3 + Aba^2 - 13b^2Ba + 9Ab^3) \cot^2(c + dx)a - 8b^4B + 8b^2(Aa^2 + 2bBa - Ab^2) \cot(c + dx)}{2 \cot^{\frac{3}{2}}(c + dx)(b + a \cot(c + dx))} dx - \frac{a(-5a^3B + a^2Ab - 3a^2bB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}} \\
 & \frac{2bd(a^2 + b^2)}{4b(a^2 + b^2)} \\
 & \downarrow 27 \\
 & \frac{a(Ab - aB)}{2bd(a^2 + b^2) \sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \int \frac{-15Ba^4 + 3Aba^3 - 31b^2Ba^2 + 11Ab^3a + 3(-5Ba^3 + Aba^2 - 13b^2Ba + 9Ab^3) \cot^2(c + dx)a - 8b^4B + 8b^2(Aa^2 + 2bBa - Ab^2) \cot(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(b + a \cot(c + dx))} dx - \frac{a(-5a^3B + a^2Ab - 3a^2bB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}} \\
 & \frac{2bd(a^2 + b^2)}{2b(a^2 + b^2)} \\
 & \frac{2bd(a^2 + b^2)}{4b(a^2 + b^2)} \\
 & \downarrow 3042 \\
 & \frac{a(Ab - aB)}{2bd(a^2 + b^2) \sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \int \frac{-15Ba^4 + 3Aba^3 - 31b^2Ba^2 + 11Ab^3a + 3(-5Ba^3 + Aba^2 - 13b^2Ba + 9Ab^3) \tan(c + dx + \frac{\pi}{2})^2 a - 8b^4B - 8b^2(Aa^2 + 2bBa - Ab^2) \tan(c + dx + \frac{\pi}{2})}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}(b - a \tan(c + dx + \frac{\pi}{2}))} dx - \frac{a(-5a^3B + a^2Ab - 3a^2bB)}{bd(a^2 + b^2)\sqrt{\cot(c + dx)}} \\
 & \frac{2bd(a^2 + b^2)}{2b(a^2 + b^2)} \\
 & \frac{2bd(a^2 + b^2)}{4b(a^2 + b^2)}
 \end{aligned}$$

3.605. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

$$\begin{aligned} & \downarrow 4132 \\ & \frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \\ & \frac{2 \int \frac{-15Ba^5 + 3Aba^4 - 31b^2Ba^3 + 3Ab^3a^2 + (-15Ba^4 + 3Aba^3 - 31b^2Ba^2 + 11Ab^3a - 8b^4B) \cot^2(c + dx)a - 24b^4Ba + 8Ab^5 - 8b^3(-Ba^2 + 2Aba + b^2B) \cot(c + dx)}{2\sqrt{\cot(c + dx)}(b + a \cot(c + dx))} dx}{b} \\ & \frac{4b(a^2 + b^2)}{2b(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \\ & \frac{2 \left(\frac{-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B}{bd\sqrt{\cot(c + dx)}} \int \frac{-15Ba^5 + 3Aba^4 - 31b^2Ba^3 + 3Ab^3a^2 + (-15Ba^4 + 3Aba^3 - 31b^2Ba^2 + 11Ab^3a - 8b^4B) \cot^2(c + dx)a - 24b^4Ba + 8Ab^5 - 8b^3(-Ba^2 + 2Aba + b^2B) \cot(c + dx)}{\sqrt{\cot(c + dx)}(b + a \cot(c + dx))} dx \right)}{2b(a^2 + b^2)} \\ & \frac{4b(a^2 + b^2)}{2b(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \\ & \frac{2 \left(\frac{-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B}{bd\sqrt{\cot(c + dx)}} \int \frac{-15Ba^5 + 3Aba^4 - 31b^2Ba^3 + 3Ab^3a^2 + (-15Ba^4 + 3Aba^3 - 31b^2Ba^2 + 11Ab^3a - 8b^4B) \tan(c + dx + \frac{\pi}{2})^2 a - 24b^4Ba + 8Ab^5 - 8b^3(-Ba^2 + 2Aba + b^2B) \tan(c + dx + \frac{\pi}{2})}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(b - a \tan(c + dx + \frac{\pi}{2}))} dx \right)}{2b(a^2 + b^2)} \\ & \frac{4b(a^2 + b^2)}{2b(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 4136 \\ & \frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \\ & \frac{2 \left(\frac{-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B}{bd\sqrt{\cot(c + dx)}} \int \frac{8(b^3(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3) - b^3(Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) \cot(c + dx))}{\sqrt{\cot(c + dx)}} dx + \frac{a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5)}{a^2 + b^2} \right)}{2b(a^2 + b^2)} \\ & \frac{4b(a^2 + b^2)}{2b(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \\ & \frac{2 \left(\frac{-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B}{bd\sqrt{\cot(c + dx)}} \int \frac{a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5)}{a^2 + b^2} \frac{\cot^2(c + dx) + 1}{\sqrt{\cot(c + dx)}(b + a \cot(c + dx))} dx + \frac{8 \int \frac{b^3(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3)}{b} dx}{b} \right)}{2b(a^2 + b^2)} \\ & \frac{4b(a^2 + b^2)}{2b(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \\ & \frac{2 \left(\frac{-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B}{bd\sqrt{\cot(c + dx)}} \int \frac{a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5)}{a^2 + b^2} \frac{\cot^2(c + dx) + 1}{\sqrt{\cot(c + dx)}(b + a \cot(c + dx))} dx + \frac{8 \int \frac{b^3(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3)}{b} dx}{b} \right)}{2b(a^2 + b^2)} \\ & \frac{4b(a^2 + b^2)}{2b(a^2 + b^2)} \end{aligned}$$

3.605. $\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3} dx$

$$\frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \frac{a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\tan(c + dx + \frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(b - a \tan(c + dx + \frac{\pi}{2}))} dx}{bd\sqrt{\cot(c + dx)}(a^2 + b^2)} - \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)}{bd\sqrt{\cot(c + dx)}} - \frac{a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5)}{a^2 + b^2} - \frac{8 \int \frac{\tan(c + dx + \frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(b - a \tan(c + dx + \frac{\pi}{2}))} dx}{b}$$

$$\frac{4b(a^2 + b^2)}{2b(a^2 + b^2)}$$

↓ 4017

$$\frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \frac{a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\tan(c + dx + \frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(b - a \tan(c + dx + \frac{\pi}{2}))} dx}{bd\sqrt{\cot(c + dx)}(a^2 + b^2)} - \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)}{bd\sqrt{\cot(c + dx)}} - \frac{a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5)}{a^2 + b^2} - \frac{16 \int \frac{\tan(c + dx + \frac{\pi}{2})^2 + 1}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}(b - a \tan(c + dx + \frac{\pi}{2}))} dx}{b}$$

$$\frac{4b(a^2 + b^2)}{2b(a^2 + b^2)}$$

↓ 25

$$\frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \frac{16 \int \frac{b^3(-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3 - (Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) \cot(c + dx))}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5)}{bd\sqrt{\cot(c + dx)}(a^2 + b^2)} - \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)}{bd\sqrt{\cot(c + dx)}} - \frac{a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5)}{a^2 + b^2} - \frac{b}{b}$$

$$\frac{4b(a^2 + b^2)}{2b(a^2 + b^2)}$$

↓ 27

$$\frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \frac{16b^3 \int \frac{-Ba^3 + 3Aba^2 + 3b^2Ba - Ab^3 - (Aa^3 + 3bBa^2 - 3Ab^2a - b^3B) \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5)}{bd\sqrt{\cot(c + dx)}(a^2 + b^2)} - \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)}{bd\sqrt{\cot(c + dx)}} - \frac{a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5)}{a^2 + b^2} - \frac{b}{b}$$

$$\frac{4b(a^2 + b^2)}{2b(a^2 + b^2)}$$

↓ 1482

$$\frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \frac{16b^3 \left(\frac{1}{2}(a^3(A - B) + 3a^2b(A + B) - 3ab^2(A - B) - b^3(A + B)) \int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}(-a^3(A + B) + 3a^2b(A - B) + 3ab^2(A + B) - b^3(A - B)) \int \frac{1 + \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} \right) + a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5)}{bd\sqrt{\cot(c + dx)}(a^2 + b^2)} - \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)}{bd\sqrt{\cot(c + dx)}} - \frac{a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5)}{a^2 + b^2} - \frac{b}{b}$$

3.605. $\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3} dx$

$$\begin{aligned} & \downarrow 1476 \\ & \frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \\ & \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)}{bd\sqrt{\cot(c + dx)}} - \frac{16b^3\left(\frac{1}{2}(a^3(A - B) + 3a^2b(A + B) - 3ab^2(A - B) - b^3(A + B))\int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}(-a^3(A + B) + 3a^2b(A - B) + 3ab^2(A + B) - b^3(A - B))\int \frac{1 + \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}\right)}{bd\sqrt{\cot(c + dx)}} \end{aligned}$$

$$\begin{aligned} & \downarrow 1082 \\ & \frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \\ & \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)}{bd\sqrt{\cot(c + dx)}} - \frac{16b^3\left(\frac{1}{2}(a^3(A - B) + 3a^2b(A + B) - 3ab^2(A - B) - b^3(A + B))\int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}(-a^3(A + B) + 3a^2b(A - B) + 3ab^2(A + B) - b^3(A - B))\int \frac{1 + \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}\right)}{bd\sqrt{\cot(c + dx)}} \end{aligned}$$

$$\begin{aligned} & \downarrow 217 \\ & \frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \\ & \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)}{bd\sqrt{\cot(c + dx)}} - \frac{16b^3\left(\frac{1}{2}(a^3(A - B) + 3a^2b(A + B) - 3ab^2(A - B) - b^3(A + B))\int \frac{1 - \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)} + \frac{1}{2}(-a^3(A + B) + 3a^2b(A - B) + 3ab^2(A + B) - b^3(A - B))\int \frac{1 + \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}\right)}{d(a^2 + b^2)} \end{aligned}$$

$$\begin{aligned} & \downarrow 1479 \\ & \frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \\ & \frac{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)}{bd\sqrt{\cot(c + dx)}} - \frac{16b^3\left(\frac{1}{2}(a^3(A - B) + 3a^2b(A + B) - 3ab^2(A - B) - b^3(A + B))\left(\int \frac{\sqrt{2} - 2\sqrt{\cot(c + dx)}}{\cot(c + dx) - \sqrt{2}\sqrt{\cot(c + dx)} + 1} d\sqrt{\cot(c + dx)}\right) + \frac{1}{2}(-a^3(A + B) + 3a^2b(A - B) + 3ab^2(A + B) - b^3(A - B))\int \frac{1 + \cot(c + dx)}{\cot^2(c + dx) + 1} d\sqrt{\cot(c + dx)}\right)}{bd\sqrt{\cot(c + dx)}} \end{aligned}$$

\(\downarrow\) 25

3.605. $\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3} dx$

$$\frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \frac{16b^3 \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\int \frac{\sqrt{2} - 2\sqrt{\cot(c+dx)}}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)} + 1} d\sqrt{\cot(c+dx)} + \int \frac{\sqrt{2} + 2\sqrt{\cot(c+dx)}}{\cot(c+dx) + \sqrt{2}\sqrt{\cot(c+dx)} + 1} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)} dx \right) \right)}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)bd\sqrt{\cot(c+dx)}}$$

27

$$\frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \frac{16b^3 \left(\frac{1}{2}(a^3(A-B) + 3a^2b(A+B) - 3ab^2(A-B) - b^3(A+B)) \left(\int \frac{\sqrt{2} - 2\sqrt{\cot(c+dx)}}{\cot(c+dx) - \sqrt{2}\sqrt{\cot(c+dx)} + 1} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)} dx \right) \right)}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)bd\sqrt{\cot(c+dx)}}$$

1103

$$\frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \frac{a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{\tan(c+dx + \frac{\pi}{2})^2 + 1}{-\tan(c+dx + \frac{\pi}{2})(b - a \tan(c+dx + \frac{\pi}{2}))} dx}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)bd\sqrt{\cot(c+dx)}(a^2 + b^2)}$$

4117

$$\frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a \cot(c + dx) + b)^2} - \frac{a^2(-15a^5B + 3a^4Ab - 46a^3b^2B + 6a^2Ab^3 - 63ab^4B + 35Ab^5) \int \frac{1}{\sqrt{\cot(c+dx)}(b + a \cot(c+dx))} d(-\cot(c+dx))}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)bd\sqrt{\cot(c+dx)}(a^2 + b^2)}$$

73

3.605. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

$$\frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a\cot(c + dx) + b)^2} - \frac{16b^3\left(\frac{1}{2}(-a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)\right)\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}}\right)}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)} - \frac{bd\sqrt{\cot(c+dx)}}{bd\sqrt{\cot(c+dx)}}$$

↓ 218

$$\frac{a(Ab - aB)}{2bd(a^2 + b^2)\sqrt{\cot(c + dx)}(a\cot(c + dx) + b)^2} - \frac{16b^3\left(\frac{1}{2}(-a^3(A+B)) + 3a^2b(A-B) + 3ab^2(A+B) - b^3(A-B)\right)\left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)}+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}}\right)}{2(-15a^4B + 3a^3Ab - 31a^2b^2B + 11aAb^3 - 8b^4B)} - \frac{bd\sqrt{\cot(c+dx)}}{bd\sqrt{\cot(c+dx)}}$$

input `Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(7/2)*(a + b*Tan[c + d*x])^3),x]`

output `(a*(A*b - a*B))/(2*b*(a^2 + b^2)*d*Sqrt[Cot[c + d*x]]*(b + a*Cot[c + d*x])^2 - ((a*(a^2*A*b + 9*A*b^3 - 5*a^3*B - 13*a*b^2*B))/(b*(a^2 + b^2)*d*Sqrt[Cot[c + d*x]]*(b + a*Cot[c + d*x]))) + ((2*(3*a^3*A*b + 11*a*A*b^3 - 15*a^4*B - 31*a^2*b^2*B - 8*b^4*B))/(b*d*Sqrt[Cot[c + d*x]]) - ((2*a^(3/2)*(3*a^4*A*b + 6*a^2*A*b^3 + 35*A*b^5 - 15*a^5*B - 46*a^3*b^2*B - 63*a*b^4*B)*ArcTan[(Sqrt[a]*Cot[c + d*x])/Sqrt[b]])/(Sqrt[b]*(a^2 + b^2)*d) + (16*b^3*((3*a^2*b*(A - B) - b^3*(A - B) - a^3*(A + B) + 3*a*b^2*(A + B))*(-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]))/2 + ((a^3*(A - B) - 3*a*b^2*(A - B) + 3*a^2*b*(A + B) - b^3*(A + B))*(-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/2)/((a^2 + b^2)*d)/b/(2*b*(a^2 + b^2))/(4*b*(a^2 + b^2))`

3.605.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

$$3.605. \int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$$

- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
 -1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
 & (LtQ[a, 0] || LtQ[b, 0])`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
 t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
 implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
 eQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
 imp[d*(Log[RemoveContent[a + b*x + c*x^2, x])/b], x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
 2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
 e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
 && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`
- rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
 -2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
 x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
 reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 1482 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a*c, 2]}, Simp[(d*q + a*e)/(2*a*c) Int[(q + c*x^2)/(a + c*x^4), x], x] + Simp[(d*q - a*e)/(2*a*c) Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4017 `Int[((c_) + (d_)*tan[(e_) + (f_)*(x_)])/Sqrt[(b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] := Simp[2/f Subst[Int[(b*c + d*x^2)/(b^2 + x^4), x], x, Sqrt[b*Tan[e + f*x]]], x] /; FreeQ[{b, c, d, e, f}, x] && NeQ[c^2 - d^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4064 `Int[(cot[(e_) + (f_)*(x_)])*(g_)^(p_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[g^(m + n) Int[(g*Cot[e + f*x])^(p - m - n)*(b + a*Cot[e + f*x])^m*(d + c*Cot[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && !IntegerQ[p] && IntegerQ[m] && IntegerQ[n]`

rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4117 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (C_)*tan[(e_) + (f_)*(x_)])^2, x_Symbol] := Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4136 Int[(((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.)
+ (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2))/((a_.) + (b_.)*tan[(e_.)
+ (f_.)*(x_)]), x_Symbol] := Simp[1/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*
Simp[b*B + a*(A - C) + (a*B - b*(A - C))*Tan[e + f*x], x], x] + Simp[
(A*b^2 - a*b*B + a^2*C)/(a^2 + b^2) Int[(c + d*Tan[e + f*x])^n*((1 + Tan[
e + f*x]^2)/(a + b*Tan[e + f*x])), x], x] /; FreeQ[{a, b, c, d, e, f, A, B,
C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] &
& !GtQ[n, 0] && !LeQ[n, -1]
```

3.605.4 Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 466, normalized size of antiderivative = 0.78

method	result
derivativedivides	$\frac{2B}{b^3 \sqrt{\cot(dx+c)}} \frac{2 \left((-3A a^2 b + A b^3 + B a^3 - 3B a b^2) \sqrt{2} \left(\ln \left(\frac{1 + \cot(dx+c) + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c) - \sqrt{2} \sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1 + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c)} \right) + 2 \arctan \left(\frac{1 + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c)} \right) \right)}{8}$
default	$\frac{2B}{b^3 \sqrt{\cot(dx+c)}} \frac{2 \left((-3A a^2 b + A b^3 + B a^3 - 3B a b^2) \sqrt{2} \left(\ln \left(\frac{1 + \cot(dx+c) + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c) - \sqrt{2} \sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1 + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c)} \right) + 2 \arctan \left(\frac{1 + \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c)} \right) \right)}{8}$

```
input int((A+B*tan(d*x+c))/cot(d*x+c)^(7/2)/(a+b*tan(d*x+c))^3,x,method=_RETURNV
ERBOSE)
```

$$3.605. \int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$$

```
output 1/d*(2*B/b^3/cot(d*x+c)^(1/2)-2/(a^2+b^2)^3*(1/8*(-3*A*a^2*b+A*b^3+B*a^3-3
*B*a*b^2)*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)
)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan
(-1+2^(1/2)*cot(d*x+c)^(1/2)))+1/8*(A*a^3-3*A*a*b^2+3*B*a^2*b-B*b^3)*2^(1/
2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d
*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot
(d*x+c)^(1/2)))+2*a^2/b^3/(a^2+b^2)^3*((( -3/8*A*a^5*b-7/4*A*b^3*a^3-11/8*
A*a*b^5+7/8*B*a^6+11/4*B*a^4*b^2+15/8*B*a^2*b^4)*cot(d*x+c)^(3/2)-1/8*b*(5
*A*a^4*b+18*A*a^2*b^3+13*A*b^5-9*B*a^5-26*B*a^3*b^2-17*B*a*b^4)*cot(d*x+c)
^(1/2))/(b+a*cot(d*x+c))^2-1/8*(3*A*a^4*b+6*A*a^2*b^3+35*A*b^5-15*B*a^5-46
*B*a^3*b^2-63*B*a*b^4)/(a*b)^(1/2)*arctan(a*cot(d*x+c)^(1/2)/(a*b)^(1/2))
)
```

3.605.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8992 vs. $2(545) = 1090$.

Time = 163.94 (sec) , antiderivative size = 18011, normalized size of antiderivative = 30.02

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \text{Too large to display}$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(7/2)/(a+b*tan(d*x+c))^3,x, algorith
m="fricas")
```

```
output Too large to include
```

3.605.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \text{Timed out}$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)**(7/2)/(a+b*tan(d*x+c))**3,x)
```

```
output Timed out
```

3.605. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

3.605.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 612, normalized size of antiderivative = 1.02

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3} dx$$

$$\frac{(15Ba^7 - 3Aa^6b + 46Ba^5b^2 - 6Aa^4b^3 + 63Ba^3b^4 - 35Aa^2b^5) \arctan\left(\frac{a}{\sqrt{ab}\sqrt{\tan(dx+c)}}\right) - 2\sqrt{2}((A+B)a^3 - 3(A-B)a^2b - 3(A+B)ab^2 + (A-B)b^3)}{(a^6b^3 + 3a^4b^5 + 3a^2b^7 + b^9)\sqrt{ab}}$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(7/2)/(a+b*tan(d*x+c))^3,x, algorithm
m="maxima")
```

```
output 1/4*((15*B*a^7 - 3*A*a^6*b + 46*B*a^5*b^2 - 6*A*a^4*b^3 + 63*B*a^3*b^4 - 3
5*A*a^2*b^5)*arctan(a/(sqrt(a*b)*sqrt(tan(d*x + c))))/((a^6*b^3 + 3*a^4*b^
5 + 3*a^2*b^7 + b^9)*sqrt(a*b)) - (2*sqrt(2)*((A + B)*a^3 - 3*(A - B)*a^2*
b - 3*(A + B)*a*b^2 + (A - B)*b^3)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(ta
n(d*x + c)))) + 2*sqrt(2)*((A + B)*a^3 - 3*(A - B)*a^2*b - 3*(A + B)*a*b^2
+ (A - B)*b^3)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sq
rt(2)*((A - B)*a^3 + 3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*log(
sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*((A - B)*a^3 +
3*(A + B)*a^2*b - 3*(A - B)*a*b^2 - (A + B)*b^3)*log(-sqrt(2)/sqrt(tan(d*x
+ c)) + 1/tan(d*x + c) + 1))/(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6) + (8*B*a
^4*b^2 + 16*B*a^2*b^4 + 8*B*b^6 + (25*B*a^5*b - 5*A*a^4*b^2 + 49*B*a^3*b^3
- 13*A*a^2*b^4 + 16*B*a*b^5)/tan(d*x + c) + (15*B*a^6 - 3*A*a^5*b + 31*B*
a^4*b^2 - 11*A*a^3*b^3 + 8*B*a^2*b^4)/tan(d*x + c)^2)/((a^4*b^5 + 2*a^2*b^
7 + b^9)/sqrt(tan(d*x + c)) + 2*(a^5*b^4 + 2*a^3*b^6 + a*b^8)/tan(d*x + c)
^(3/2) + (a^6*b^3 + 2*a^4*b^5 + a^2*b^7)/tan(d*x + c)^(5/2))/d
```

3.605.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \text{Timed out}$$

```
input integrate((A+B*tan(d*x+c))/cot(d*x+c)^(7/2)/(a+b*tan(d*x+c))^3,x, algorithm
m="giac")
```

output Timed out

3.605. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^3} dx$

3.605.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^3} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{7/2} (a + b \tan(c + dx))^3} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(7/2)*(a + b*tan(c + d*x))^3),x)`output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(7/2)*(a + b*tan(c + d*x))^3), x)`

3.606
$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.606.1 Optimal result 5801
 3.606.2 Mathematica [A] (verified) 5802
 3.606.3 Rubi [A] (verified) 5802
 3.606.4 Maple [A] (verified) 5806
 3.606.5 Fricas [C] (verification not implemented) 5807
 3.606.6 Sympy [F] 5807
 3.606.7 Maxima [A] (verification not implemented) 5808
 3.606.8 Giac [F] 5808
 3.606.9 Mupad [B] (verification not implemented) 5809

3.606.1 Optimal result

Integrand size = 36, antiderivative size = 156

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = -\frac{B \arctan\left(1-\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d} + \frac{B \arctan\left(1+\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d} - \frac{2B \cot^{\frac{3}{2}}(c+dx)}{3d} + \frac{B \log\left(1-\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{2\sqrt{2}d} - \frac{B \log\left(1+\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{2\sqrt{2}d}$$

output

```
-2/3*B*cot(d*x+c)^(3/2)/d+1/2*B*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+1/2*B*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+1/4*B*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/4*B*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)
```


3.606.2 Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.57

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{B\left(-3\arctan\left(\sqrt[4]{-\cot^2(c+dx)}\right)\sqrt[4]{-\cot(c+dx)}+3\operatorname{arctanh}\left(\sqrt[4]{-\cot^2(c+dx)}\right)\sqrt[4]{-\cot(c+dx)}\right)}{3d\sqrt[4]{\cot(c+dx)}}$$

input `Integrate[(Cot[c + d*x]^(5/2)*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `-1/3*(B*(-3*ArcTan[(-Cot[c + d*x]^2)^(1/4)]*(-Cot[c + d*x])^(1/4) + 3*ArcTanh[(-Cot[c + d*x]^2)^(1/4)]*(-Cot[c + d*x])^(1/4) + 2*Cot[c + d*x]^(7/4)))/(d*Cot[c + d*x]^(1/4))`

3.606.3 Rubi [A] (verified)Time = 0.46 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {2011, 3042, 3954, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\cot^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx \\ & \quad \downarrow \text{2011} \\ & B \int \cot^{\frac{5}{2}}(c+dx) dx \\ & \quad \downarrow \text{3042} \\ & B \int \left(-\tan\left(c+dx+\frac{\pi}{2}\right)\right)^{5/2} dx \\ & \quad \downarrow \text{3954} \\ & B \left(-\int \sqrt{\cot(c+dx)} dx - \frac{2\cot^{\frac{3}{2}}(c+dx)}{3d}\right) \end{aligned}$$

3.606. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& B \left(- \int \sqrt{-\tan \left(c + dx + \frac{\pi}{2} \right)} dx - \frac{2 \cot^{\frac{3}{2}}(c + dx)}{3d} \right) \\
& \downarrow 3957 \\
& B \left(\frac{\int \frac{\sqrt{\cot(c+dx)}}{\cot^2(c+dx)+1} d \cot(c + dx)}{d} - \frac{2 \cot^{\frac{3}{2}}(c + dx)}{3d} \right) \\
& \downarrow 266 \\
& B \left(\frac{2 \int \frac{\cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c + dx)}}{d} - \frac{2 \cot^{\frac{3}{2}}(c + dx)}{3d} \right) \\
& \downarrow 826 \\
& B \left(\frac{2 \left(\frac{1}{2} \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d \sqrt{\cot(c + dx)} - \frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c + dx)} \right)}{d} - \frac{2 \cot^{\frac{3}{2}}(c + dx)}{3d} \right) \\
& \downarrow 1476 \\
& B \left(\frac{2 \left(\frac{1}{2} \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d \sqrt{\cot(c + dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d \sqrt{\cot(c + dx)} \right) - \frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c + dx)} \right)}{d} \right) \\
& \downarrow 1082 \\
& B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)-1} d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c + dx)} \right)}{d} \right) \\
& \downarrow 217 \\
& B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c + dx)} \right)}{d} - \frac{2 \cot^{\frac{3}{2}}(c + dx)}{3d} \right) \\
& \downarrow 1479
\end{aligned}$$

3.606. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$B \left(\frac{2 \left(\frac{1}{2} \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right)}{d} \right)$$

↓ 25

$$B \left(\frac{2 \left(\frac{1}{2} \left(- \int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right)}{d} \right)$$

↓ 27

$$B \left(\frac{2 \left(\frac{1}{2} \left(- \int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right)}{d} \right)$$

↓ 1103

$$B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log(\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d} \right)$$

input `Int[(Cot[c + d*x]^(5/2)*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `B*((-2*Cot[c + d*x]^(3/2))/(3*d) + (2*((-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2])/2 + (Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2]))/2))/d`

3.606.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplrQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

3.606.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.65

method	result	size
default	$\frac{B \left(-\frac{2 \cot(dx+c)^{\frac{3}{2}}}{3} + \frac{\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1+\sqrt{2}\sqrt{\cot(dx+c)}}{4} \right) + 2 \arctan \left(\frac{-1+\sqrt{2}\sqrt{\cot(dx+c)}}{4} \right) \right)}{d}$	102

input `int(cot(d*x+c)^(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `B/d*(-2/3*cot(d*x+c)^(3/2)+1/4*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))`

3.606. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.606.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 458, normalized size of antiderivative = 2.94

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx =$$

$$3d\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log\left(\frac{B\sqrt{\frac{\cos(2dx+2c)+1}{\sin(2dx+2c)}} \sin(2dx+2c) + (d\cos(2dx+2c)+d)\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}}}{\cos(2dx+2c)+1}\right) \sin(2dx+2c) - 3d\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log\left(\dots\right)$$

```
input integrate(cot(d*x+c)^(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")
```

```
output -1/6*(3*d*(-B^4/d^4)^(1/4)*log((B*sqrt((cos(2*d*x + 2*c)) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) + (d*cos(2*d*x + 2*c) + d)*(-B^4/d^4)^(1/4))/(cos(2*d*x + 2*c) + 1)*sin(2*d*x + 2*c) - 3*d*(-B^4/d^4)^(1/4)*log((B*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) - (d*cos(2*d*x + 2*c) + d)*(-B^4/d^4)^(1/4))/(cos(2*d*x + 2*c) + 1)*sin(2*d*x + 2*c) - 3*I*d*(-B^4/d^4)^(1/4)*log((B*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) - (I*d*cos(2*d*x + 2*c) + I*d)*(-B^4/d^4)^(1/4))/(cos(2*d*x + 2*c) + 1))*sin(2*d*x + 2*c) + 3*I*d*(-B^4/d^4)^(1/4)*log((B*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) - (-I*d*cos(2*d*x + 2*c) - I*d)*(-B^4/d^4)^(1/4))/(cos(2*d*x + 2*c) + 1))*sin(2*d*x + 2*c) + 4*(B*cos(2*d*x + 2*c) + B)*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c)))/(d*sin(2*d*x + 2*c))
```

3.606.6 Sympy [F]

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = B \int \cot^{\frac{5}{2}}(c+dx) dx$$

```
input integrate(cot(d*x+c)**(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x)
```

```
output B*Integral(cot(c + d*x)**(5/2), x)
```

3.606. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$

3.606.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.81

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{3\left(2\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2}+\frac{2}{\sqrt{\tan(dx+c)}}\right)\right)+2\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2}-\frac{2}{\sqrt{\tan(dx+c)}}\right)\right)-\sqrt{2}\log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}}\right)\right)}{12d}$$

input `integrate(cot(d*x+c)^(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/12*(3*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)+2/sqrt(tan(d*x+c))))+2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)-2/sqrt(tan(d*x+c))))-sqrt(2)*log(sqrt(2)/sqrt(tan(d*x+c))+1/tan(d*x+c)+1)+sqrt(2)*log(-sqrt(2)/sqrt(tan(d*x+c))+1/tan(d*x+c)+1))*B-8*B/tan(d*x+c)^(3/2))/d`

3.606.8 Giac [F]

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \int \frac{(Bb\tan(dx+c)+Ba)\cot(dx+c)^{\frac{5}{2}}}{b\tan(dx+c)+a} dx$$

input `integrate(cot(d*x+c)^(5/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*b*tan(d*x+c)+B*a)*cot(d*x+c)^(5/2)/(b*tan(d*x+c)+a),x)`

3.606.9 Mupad [B] (verification not implemented)

Time = 9.93 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.41

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{(-1)^{1/4} B \operatorname{atan}\left((-1)^{1/4} \sqrt{\frac{1}{\tan(c+dx)}}\right)}{d} - \frac{2B\left(\frac{1}{\tan(c+dx)}\right)^{3/2}}{3d} - \frac{(-1)^{1/4} B \operatorname{atanh}\left((-1)^{1/4} \sqrt{\frac{1}{\tan(c+dx)}}\right)}{d}$$

input `int((cot(c + d*x)^(5/2)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`output `((-1)^(1/4)*B*atan((-1)^(1/4)*(1/tan(c + d*x))^(1/2))/d - (2*B*(1/tan(c + d*x))^(3/2))/(3*d) - ((-1)^(1/4)*B*atanh((-1)^(1/4)*(1/tan(c + d*x))^(1/2)))/d`

3.607
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.607.1 Optimal result 5810
 3.607.2 Mathematica [A] (verified) 5811
 3.607.3 Rubi [A] (verified) 5811
 3.607.4 Maple [A] (verified) 5815
 3.607.5 Fricas [C] (verification not implemented) 5816
 3.607.6 Sympy [F] 5816
 3.607.7 Maxima [A] (verification not implemented) 5817
 3.607.8 Giac [F] 5817
 3.607.9 Mupad [B] (verification not implemented) 5817

3.607.1 Optimal result

Integrand size = 36, antiderivative size = 154

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = -\frac{B \arctan\left(1-\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d} + \frac{B \arctan\left(1+\sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d} - \frac{2B\sqrt{\cot(c+dx)}}{d} - \frac{B \log\left(1-\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{2\sqrt{2}d} + \frac{B \log\left(1+\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)\right)}{2\sqrt{2}d}$$

output `1/2*B*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+1/2*B*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/4*B*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+1/4*B*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-2*B*cot(d*x+c)^(1/2)/d`

3.607.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.91

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx =$$

$$\frac{B\left(\frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1+\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} + 2\sqrt{\cot(c+dx)} + \frac{\log\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)}{2\sqrt{2}}\right)}{2\sqrt{2}} - \frac{\log\left(\frac{1+\sqrt{2}\sqrt{\cot(c+dx)}+\cot(c+dx)}{2\sqrt{2}}\right)}{2\sqrt{2}}\right)}{d}$$

input `Integrate[(Cot[c + d*x]^(3/2)*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `-((B*(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2] - ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2] + 2*Sqrt[Cot[c + d*x]] + Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2])))/d)`

3.607.3 Rubi [A] (verified)Time = 0.41 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {2011, 3042, 3954, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \cot^{\frac{3}{2}}(c+dx) dx$$

$$\downarrow \text{3042}$$

$$B \int \left(-\tan\left(c+dx+\frac{\pi}{2}\right)\right)^{3/2} dx$$

$$\downarrow \text{3954}$$

$$B \left(-\int \frac{1}{\sqrt{\cot(c+dx)}} dx - \frac{2\sqrt{\cot(c+dx)}}{d}\right)$$

3.607. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& B \left(- \int \frac{1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx - \frac{2\sqrt{\cot(c+dx)}}{d} \right) \\
& \downarrow 3957 \\
& B \left(\frac{\int \frac{1}{\sqrt{\cot(c+dx)(\cot^2(c+dx)+1)}} d \cot(c+dx)}{d} - \frac{2\sqrt{\cot(c+dx)}}{d} \right) \\
& \downarrow 266 \\
& B \left(\frac{2 \int \frac{1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} - \frac{2\sqrt{\cot(c+dx)}}{d} \right) \\
& \downarrow 755 \\
& B \left(\frac{2 \left(\frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d} - \frac{2\sqrt{\cot(c+dx)}}{d} \right) \\
& \downarrow 1476 \\
& B \left(\frac{2 \left(\frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{d} \right) \\
& \downarrow 1082 \\
& B \left(\frac{2 \left(\frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)-1} d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \right) \\
& \downarrow 217 \\
& B \left(\frac{2 \left(\frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{d} - \frac{2\sqrt{\cot(c+dx)}}{d} \right) \\
& \downarrow 1479
\end{aligned}$$

3.607. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$B \left(\frac{2 \left(\frac{1}{2} \left(\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} - \int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \right)$$

↓ 25

$$B \left(\frac{2 \left(\frac{1}{2} \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \right)$$

↓ 27

$$B \left(\frac{2 \left(\frac{1}{2} \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \right)$$

↓ 1103

$$B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log(\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d} \right)$$

input `Int[(Cot[c + d*x]^(3/2)*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `B*((-2*Sqrt[Cot[c + d*x]])/d + (2*((-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]))/2 + (-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2]))/2)/d)`

3.607.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3954 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Simp[b^2 Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

3.607.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.66

method	result	size
default	$\frac{B \left(-2\sqrt{\cot(dx+c)} + \frac{\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan(1+\sqrt{2}\sqrt{\cot(dx+c)}) + 2 \arctan(-1+\sqrt{2}\sqrt{\cot(dx+c)}) \right)}{4} \right)}{d}$	102

input `int(cot(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `B/d*(-2*cot(d*x+c)^(1/2)+1/4*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2)))`

3.607. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$

3.607.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 422, normalized size of antiderivative = 2.74

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx =$$

$$d\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log\left(\frac{B^3\sqrt{\frac{\cos(2dx+2c)+1}{\sin(2dx+2c)}}\sin(2dx+2c)+(d^3\cos(2dx+2c)+d^3)\left(-\frac{B^4}{d^4}\right)^{\frac{3}{4}}}{\cos(2dx+2c)+1}\right) - d\left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log\left(\frac{B^3\sqrt{\frac{\cos(2dx+2c)+1}{\sin(2dx+2c)}}\sin(2dx+2c)+(d^3\cos(2dx+2c)+d^3)\left(-\frac{B^4}{d^4}\right)^{\frac{3}{4}}}{\cos(2dx+2c)+1}\right)$$

input `integrate(cot(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="fricas")`

output `-1/2*(d*(-B^4/d^4)^(1/4)*log((B^3*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) + (d^3*cos(2*d*x + 2*c) + d^3)*(-B^4/d^4)^(3/4))/(cos(2*d*x + 2*c) + 1)) - d*(-B^4/d^4)^(1/4)*log((B^3*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) - (d^3*cos(2*d*x + 2*c) + d^3)*(-B^4/d^4)^(3/4))/(cos(2*d*x + 2*c) + 1)) + I*d*(-B^4/d^4)^(1/4)*log((B^3*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) - (I*d^3*cos(2*d*x + 2*c) + I*d^3)*(-B^4/d^4)^(3/4))/(cos(2*d*x + 2*c) + 1)) - I*d*(-B^4/d^4)^(1/4)*log((B^3*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) - (-I*d^3*cos(2*d*x + 2*c) - I*d^3)*(-B^4/d^4)^(3/4))/(cos(2*d*x + 2*c) + 1)) + 4*B*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c)))/d`

3.607.6 Sympy [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = B \int \cot^{\frac{3}{2}}(c+dx) dx$$

input `integrate(cot(d*x+c)**(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `B*Integral(cot(c + d*x)**(3/2), x)`

3.607. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$

3.607.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.82

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{2\sqrt{2}B \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}B \arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + \sqrt{2}B \log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}}\right)}{4d}$$

```
input integrate(cot(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")
```

```
output 1/4*(2*sqrt(2)*B*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*B*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)*B*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*B*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - 8*B/sqrt(tan(d*x + c)))/d
```

3.607.8 Giac [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \int \frac{(Bb\tan(dx+c) + Ba)\cot(dx+c)^{\frac{3}{2}}}{b\tan(dx+c) + a} dx$$

```
input integrate(cot(d*x+c)^(3/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")
```

```
output integrate((B*b*tan(d*x + c) + B*a)*cot(d*x + c)^(3/2)/(b*tan(d*x + c) + a), x)
```

3.607.9 Mupad [B] (verification not implemented)

Time = 9.82 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.42

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = -\frac{2B\sqrt{\frac{1}{\tan(c+dx)}}}{d}$$

$$-\frac{(-1)^{1/4}B\operatorname{atan}\left((-1)^{1/4}\sqrt{\frac{1}{\tan(c+dx)}}\right)}{d} \operatorname{li}$$

$$-\frac{(-1)^{1/4}B\operatorname{atanh}\left((-1)^{1/4}\sqrt{\frac{1}{\tan(c+dx)}}\right)}{d} \operatorname{li}$$

3.607. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$

input `int((cot(c + d*x)^(3/2)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `- (2*B*(1/tan(c + d*x))^(1/2))/d - ((-1)^(1/4)*B*atan((-1)^(1/4)*(1/tan(c + d*x))^(1/2))*1i)/d - ((-1)^(1/4)*B*atanh((-1)^(1/4)*(1/tan(c + d*x))^(1/2))*1i)/d`

3.607.
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.608
$$\int \frac{\sqrt{\cot(c+dx)}(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$$

3.608.1 Optimal result	5819
3.608.2 Mathematica [A] (verified)	5820
3.608.3 Rubi [A] (verified)	5820
3.608.4 Maple [A] (verified)	5824
3.608.5 Fricas [C] (verification not implemented)	5824
3.608.6 Sympy [F]	5825
3.608.7 Maxima [A] (verification not implemented)	5825
3.608.8 Giac [F]	5826
3.608.9 Mupad [B] (verification not implemented)	5826

3.608.1 Optimal result

Integrand size = 36, antiderivative size = 138

$$\int \frac{\sqrt{\cot(c+dx)}(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx = \frac{B \arctan\left(1 - \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d} - \frac{B \arctan\left(1 + \sqrt{2}\sqrt{\cot(c+dx)}\right)}{\sqrt{2}d} - \frac{B \log\left(1 - \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d} + \frac{B \log\left(1 + \sqrt{2}\sqrt{\cot(c+dx)} + \cot(c+dx)\right)}{2\sqrt{2}d}$$

```
output -1/2*B*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/2*B*arctan(1+2^(1/2)
)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/4*B*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/
2))/d*2^(1/2)+1/4*B*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)
```

3.608.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.43

$$\int \frac{\sqrt{\cot(c+dx)}(aB + bB \tan(c+dx))}{a + b \tan(c+dx)} dx$$

$$= \frac{B \left(-\arctan \left(\sqrt[4]{-\cot^2(c+dx)} \right) + \operatorname{arctanh} \left(\sqrt[4]{-\cot^2(c+dx)} \right) \right) \sqrt[4]{-\cot(c+dx)}}{d \sqrt[4]{\cot(c+dx)}}$$

input `Integrate[(Sqrt[Cot[c + d*x]]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(B*(-ArcTan[(-Cot[c + d*x]^2)^(1/4)] + ArcTanh[(-Cot[c + d*x]^2)^(1/4)])*(-Cot[c + d*x])^(1/4))/(d*Cot[c + d*x]^(1/4))`

3.608.3 Rubi [A] (verified)Time = 0.35 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2011, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cot(c+dx)}(aB + bB \tan(c+dx))}{a + b \tan(c+dx)} dx$$

$$\downarrow \text{2011}$$

$$B \int \sqrt{\cot(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$B \int \sqrt{-\tan\left(c+dx + \frac{\pi}{2}\right)} dx$$

$$\downarrow \text{3957}$$

$$B \int \frac{\sqrt{\cot(c+dx)}}{\cot^2(c+dx)+1} d \cot(c+dx)$$

$$\downarrow \text{266}$$

3.608. $\int \frac{\sqrt{\cot(c+dx)}(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\frac{2B \int \frac{\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d}$$

↓ 826

$$\frac{2B \left(\frac{1}{2} \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d}$$

↓ 1476

$$\frac{2B \left(\frac{1}{2} \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) - \frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d}$$

↓ 1082

$$\frac{2B \left(\frac{1}{2} \left(\int \frac{1}{-\cot(c+dx)-1} \frac{d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \int \frac{1}{-\cot(c+dx)-1} \frac{d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d}$$

↓ 217

$$\frac{2B \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d}$$

↓ 1479

$$\frac{2B \left(\frac{1}{2} \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d}$$

↓ 25

$$\frac{2B \left(\frac{1}{2} \left(- \int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} - \int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d}$$

↓ 27

$$\frac{2B \left(\frac{1}{2} \left(- \int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d}$$

↓ 1103

3.608. $\int \frac{\sqrt{\cot(c+dx)}(aB+bB \tan(c+dx))}{a+b \tan(c+dx)} dx$

$$\frac{2B \left(\frac{1}{2} \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right)}{\sqrt{2}} - \frac{\arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log\left(\frac{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}}{2\sqrt{2}}\right)}{2\sqrt{2}} - \frac{\log\left(\frac{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)}}{2\sqrt{2}}\right)}{2\sqrt{2}} \right) \right)}{d}$$

input `Int[(Sqrt[Cot[c + d*x]]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x]),x]`

output `(-2*B*((-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2])/2 + (Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2]))/2)/d`

3.608.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2)]^p, x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntegerBinomialQ[a, b, c, 2, m, p, x]`

rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

3.608.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.65

method	result	size
default	$-\frac{B\sqrt{2}\left(\ln\left(\frac{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}\right)+2\arctan\left(1+\sqrt{2}\sqrt{\cot(dx+c)}\right)+2\arctan\left(-1+\sqrt{2}\sqrt{\cot(dx+c)}\right)\right)}{4d}$	90

input `int(cot(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output
$$-1/4*B/d*2^{(1/2)}*(\ln((1+\cot(d*x+c)-2^{(1/2)}*\cot(d*x+c)^{(1/2)})/(1+\cot(d*x+c)+2^{(1/2)}*\cot(d*x+c)^{(1/2)}))+2*\arctan(1+2^{(1/2)}*\cot(d*x+c)^{(1/2)})+2*\arctan(-1+2^{(1/2)}*\cot(d*x+c)^{(1/2)}))$$

3.608.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 362, normalized size of antiderivative = 2.62

$$\int \frac{\sqrt{\cot(c+dx)}(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

$$= \frac{1}{2} \left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log \left(\frac{B\sqrt{\frac{\cos(2dx+2c)+1}{\sin(2dx+2c)}} \sin(2dx+2c) + (d\cos(2dx+2c)+d) \left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}}}{\cos(2dx+2c)+1} \right)$$

$$- \frac{1}{2} \left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log \left(\frac{B\sqrt{\frac{\cos(2dx+2c)+1}{\sin(2dx+2c)}} \sin(2dx+2c) - (d\cos(2dx+2c)+d) \left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}}}{\cos(2dx+2c)+1} \right)$$

$$- \frac{1}{2} i \left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log \left(\frac{B\sqrt{\frac{\cos(2dx+2c)+1}{\sin(2dx+2c)}} \sin(2dx+2c) - (id\cos(2dx+2c)+id) \left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}}}{\cos(2dx+2c)+1} \right)$$

$$+ \frac{1}{2} i \left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log \left(\frac{B\sqrt{\frac{\cos(2dx+2c)+1}{\sin(2dx+2c)}} \sin(2dx+2c) - (-id\cos(2dx+2c)-id) \left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}}}{\cos(2dx+2c)+1} \right)$$

input `integrate(cot(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x,algorithm="fracas")`

3.608.
$$\int \frac{\sqrt{\cot(c+dx)}(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$$

output $1/2*(-B^4/d^4)^{(1/4)}*\log((B*\sqrt{(\cos(2*d*x + 2*c) + 1)/\sin(2*d*x + 2*c)})*\sin(2*d*x + 2*c) + (d*\cos(2*d*x + 2*c) + d)*(-B^4/d^4)^{(1/4)})/(\cos(2*d*x + 2*c) + 1)) - 1/2*(-B^4/d^4)^{(1/4)}*\log((B*\sqrt{(\cos(2*d*x + 2*c) + 1)/\sin(2*d*x + 2*c)})*\sin(2*d*x + 2*c) - (d*\cos(2*d*x + 2*c) + d)*(-B^4/d^4)^{(1/4)})/(\cos(2*d*x + 2*c) + 1)) - 1/2*I*(-B^4/d^4)^{(1/4)}*\log((B*\sqrt{(\cos(2*d*x + 2*c) + 1)/\sin(2*d*x + 2*c)})*\sin(2*d*x + 2*c) - (I*d*\cos(2*d*x + 2*c) + I*d)*(-B^4/d^4)^{(1/4)})/(\cos(2*d*x + 2*c) + 1)) + 1/2*I*(-B^4/d^4)^{(1/4)}*\log((B*\sqrt{(\cos(2*d*x + 2*c) + 1)/\sin(2*d*x + 2*c)})*\sin(2*d*x + 2*c) - (-I*d*\cos(2*d*x + 2*c) - I*d)*(-B^4/d^4)^{(1/4)})/(\cos(2*d*x + 2*c) + 1))$

3.608.6 Sympy [F]

$$\int \frac{\sqrt{\cot(c+dx)}(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = B \int \sqrt{\cot(c+dx)} dx$$

input `integrate(cot(d*x+c)**(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x)`

output `B*Integral(sqrt(cot(c + d*x)), x)`

3.608.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.82

$$\int \frac{\sqrt{\cot(c+dx)}(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx = \frac{\left(2\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}\arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) - \sqrt{2}\log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}}\right)\right)}{4d}$$

input `integrate(cot(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output $-1/4*(2*\sqrt{2}*\arctan(1/2*\sqrt{2}*(\sqrt{2} + 2/\sqrt{\tan(d*x + c)}))) + 2*\sqrt{2}*\arctan(-1/2*\sqrt{2}*(\sqrt{2} - 2/\sqrt{\tan(d*x + c)})) - \sqrt{2}*\log(\sqrt{2}/\sqrt{\tan(d*x + c)} + 1/\tan(d*x + c) + 1) + \sqrt{2}*\log(-\sqrt{2}/\sqrt{\tan(d*x + c)} + 1/\tan(d*x + c) + 1))*B/d$

3.608. $\int \frac{\sqrt{\cot(c+dx)}(aB+bB\tan(c+dx))}{a+b\tan(c+dx)} dx$

3.608.8 Giac [F]

$$\int \frac{\sqrt{\cot(c+dx)}(aB + bB \tan(c+dx))}{a + b \tan(c+dx)} dx = \int \frac{(Bb \tan(dx+c) + Ba)\sqrt{\cot(dx+c)}}{b \tan(dx+c) + a} dx$$

input `integrate(cot(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*b*tan(d*x + c) + B*a)*sqrt(cot(d*x + c))/(b*tan(d*x + c) + a), x)`

3.608.9 Mupad [B] (verification not implemented)

Time = 9.51 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.30

$$\int \frac{\sqrt{\cot(c+dx)}(aB + bB \tan(c+dx))}{a + b \tan(c+dx)} dx$$

$$= -\frac{(-1)^{1/4} B \left(\operatorname{atan} \left((-1)^{1/4} \sqrt{\frac{1}{\tan(c+dx)}} \right) - \operatorname{atanh} \left((-1)^{1/4} \sqrt{\frac{1}{\tan(c+dx)}} \right) \right)}{d}$$

input `int((cot(c + d*x)^(1/2)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x)),x)`

output `-((-1)^(1/4)*B*(atan((-1)^(1/4)*(1/tan(c + d*x))^(1/2)) - atanh((-1)^(1/4)*(1/tan(c + d*x))^(1/2)))/d`

3.609
$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))}} dx$$

3.609.1 Optimal result 5827
 3.609.2 Mathematica [A] (verified) 5828
 3.609.3 Rubi [A] (verified) 5828
 3.609.4 Maple [A] (verified) 5832
 3.609.5 Fricas [C] (verification not implemented) 5832
 3.609.6 Sympy [F] 5833
 3.609.7 Maxima [A] (verification not implemented) 5833
 3.609.8 Giac [F] 5834
 3.609.9 Mupad [B] (verification not implemented) 5834

3.609.1 Optimal result

Integrand size = 36, antiderivative size = 138

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))}} dx = \frac{B \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d} - \frac{B \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d} + \frac{B \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d} - \frac{B \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

output

```
-1/2*B*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/2*B*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+1/4*B*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)-1/4*B*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)
```

3.609.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.80

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx$$

$$= \frac{B \left(2 \arctan \left(1 - \sqrt{2} \sqrt{\cot(c + dx)} \right) - 2 \arctan \left(1 + \sqrt{2} \sqrt{\cot(c + dx)} \right) + \log \left(1 - \sqrt{2} \sqrt{\cot(c + dx)} + \cot(c + dx) \right) + \log \left(1 + \sqrt{2} \sqrt{\cot(c + dx)} + \cot(c + dx) \right) \right)}{2\sqrt{2}d}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])),x]`

output `(B*(2*ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]] - 2*ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]) + Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]])/(2*Sqrt[2]*d)`

3.609.3 Rubi [A] (verified)Time = 0.36 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2011, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx$$

$$\downarrow \text{2011}$$

$$B \int \frac{1}{\sqrt{\cot(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$B \int \frac{1}{\sqrt{-\tan(c + dx + \frac{\pi}{2})}} dx$$

$$\downarrow \text{3957}$$

$$\frac{B \int \frac{1}{\sqrt{\cot(c+dx)(\cot^2(c+dx)+1)}} d \cot(c + dx)}{d}$$

3.609. $\int \frac{aB+bB \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))} dx$

$$\begin{aligned} & \downarrow 266 \\ & \frac{2B \int \frac{1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} \\ & \downarrow 755 \\ & \frac{2B \left(\frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d} \\ & \downarrow 1476 \\ & \frac{2B \left(\frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) \right)}{d} \\ & \downarrow 1082 \\ & \frac{2B \left(\frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)-1} d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \\ & \downarrow 217 \\ & \frac{2B \left(\frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{d} \\ & \downarrow 1479 \\ & \frac{2B \left(\frac{1}{2} \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \\ & \downarrow 25 \\ & \frac{2B \left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \\ & \downarrow 27 \\ & \frac{2B \left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \end{aligned}$$

3.609. $\int \frac{aB+bB \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))}} dx$

↓ 1103

$$\frac{2B \left(\frac{1}{2} \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\sqrt{2}}\right) - \arctan\left(\frac{1-\sqrt{2}\sqrt{\cot(c+dx)}}{\sqrt{2}}\right)}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log\left(\frac{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}}{2\sqrt{2}}\right) - \log\left(\frac{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)}}{2\sqrt{2}}\right)}{\sqrt{2}} \right) \right)}{d}$$

input `Int[(a*B + b*B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])),x]`

output `(-2*B*((-ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]]]/Sqrt[2])/2 + (-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2]))/2)/d`

3.609.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

3.609. $\int \frac{aB+bB \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))}} dx$

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

3.609.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.65

method	result	size
default	$-\frac{B\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(1+\sqrt{2}\sqrt{\cot(dx+c)} \right) + 2 \arctan \left(-1+\sqrt{2}\sqrt{\cot(dx+c)} \right) \right)}{4d}$	90

```
input int((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x,method=_RETURN
NVERBOSE)
```

```
output -1/4*B/d*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)
-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(
-1+2^(1/2)*cot(d*x+c)^(1/2)))
```

3.609.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 386, normalized size of antiderivative = 2.80

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))}} dx$$

$$= \frac{1}{2} \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \log \left(\frac{B^3 \sqrt{\frac{\cos(2dx+2c)+1}{\sin(2dx+2c)}} \sin(2dx+2c) + (d^3 \cos(2dx+2c) + d^3) \left(-\frac{B^4}{d^4} \right)^{\frac{3}{4}}}{\cos(2dx+2c) + 1} \right)$$

$$- \frac{1}{2} \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \log \left(\frac{B^3 \sqrt{\frac{\cos(2dx+2c)+1}{\sin(2dx+2c)}} \sin(2dx+2c) - (d^3 \cos(2dx+2c) + d^3) \left(-\frac{B^4}{d^4} \right)^{\frac{3}{4}}}{\cos(2dx+2c) + 1} \right)$$

$$+ \frac{1}{2} i \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \log \left(\frac{B^3 \sqrt{\frac{\cos(2dx+2c)+1}{\sin(2dx+2c)}} \sin(2dx+2c) - (i d^3 \cos(2dx+2c) + i d^3) \left(-\frac{B^4}{d^4} \right)^{\frac{3}{4}}}{\cos(2dx+2c) + 1} \right)$$

$$- \frac{1}{2} i \left(-\frac{B^4}{d^4} \right)^{\frac{1}{4}} \log \left(\frac{B^3 \sqrt{\frac{\cos(2dx+2c)+1}{\sin(2dx+2c)}} \sin(2dx+2c) - (-i d^3 \cos(2dx+2c) - i d^3) \left(-\frac{B^4}{d^4} \right)^{\frac{3}{4}}}{\cos(2dx+2c) + 1} \right)$$

```
input integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x,algori
thm="fracas")
```

3.609.
$$\int \frac{aB+bB \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))}} dx$$

```
output 1/2*(-B^4/d^4)^(1/4)*log((B^3*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c)
)*sin(2*d*x + 2*c) + (d^3*cos(2*d*x + 2*c) + d^3)*(-B^4/d^4)^(3/4))/(cos(2
*d*x + 2*c) + 1)) - 1/2*(-B^4/d^4)^(1/4)*log((B^3*sqrt((cos(2*d*x + 2*c) +
1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) - (d^3*cos(2*d*x + 2*c) + d^3)*(-B^
4/d^4)^(3/4))/(cos(2*d*x + 2*c) + 1)) + 1/2*I*(-B^4/d^4)^(1/4)*log((B^3*sq
rt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) - (I*d^3*cos(
2*d*x + 2*c) + I*d^3)*(-B^4/d^4)^(3/4))/(cos(2*d*x + 2*c) + 1)) - 1/2*I*(-
B^4/d^4)^(1/4)*log((B^3*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(
2*d*x + 2*c) - (-I*d^3*cos(2*d*x + 2*c) - I*d^3)*(-B^4/d^4)^(3/4))/(cos(2*
d*x + 2*c) + 1))
```

3.609.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx = B \int \frac{1}{\sqrt{\cot(c + dx)}} dx$$

```
input integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)**(1/2)/(a+b*tan(d*x+c)),x)
```

```
output B*Integral(1/sqrt(cot(c + d*x)), x)
```

3.609.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.84

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx = \frac{2\sqrt{2}B \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2}B \arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + \sqrt{2}B \log\left(\frac{\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}}{\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}}\right)}{4d}$$

```
input integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x, algori
thm="maxima")
```

```
output -1/4*(2*sqrt(2)*B*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2
*sqrt(2)*B*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + sqrt(2)
*B*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - sqrt(2)*B*log(-s
qrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))/d
```

3.609. $\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx$

3.609.8 Giac [F]

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx = \int \frac{Bb \tan(dx + c) + Ba}{(b \tan(dx + c) + a)\sqrt{\cot(dx + c)}} dx$$

input `integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*b*tan(d*x + c) + B*a)/((b*tan(d*x + c) + a)*sqrt(cot(d*x + c))), x)`

3.609.9 Mupad [B] (verification not implemented)

Time = 9.41 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.34

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))} dx = \frac{(-1)^{1/4} B \operatorname{atan}\left((-1)^{1/4} \sqrt{\frac{1}{\tan(c+dx)}}\right) \operatorname{li}}{d} + \frac{(-1)^{1/4} B \operatorname{atanh}\left((-1)^{1/4} \sqrt{\frac{1}{\tan(c+dx)}}\right) \operatorname{li}}{d}$$

input `int((B*a + B*b*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))),x)`

output `((-1)^(1/4)*B*atan((-1)^(1/4)*(1/tan(c + d*x))^(1/2))*1i)/d + ((-1)^(1/4)*B*atanh((-1)^(1/4)*(1/tan(c + d*x))^(1/2))*1i)/d`

3.610
$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

3.610.1 Optimal result 5835
 3.610.2 Mathematica [A] (verified) 5836
 3.610.3 Rubi [A] (verified) 5836
 3.610.4 Maple [A] (verified) 5840
 3.610.5 Fricas [C] (verification not implemented) 5841
 3.610.6 Sympy [F] 5841
 3.610.7 Maxima [A] (verification not implemented) 5842
 3.610.8 Giac [F] 5842
 3.610.9 Mupad [B] (verification not implemented) 5842

3.610.1 Optimal result

Integrand size = 36, antiderivative size = 154

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = -\frac{B \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d} + \frac{B \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d} + \frac{2B}{d\sqrt{\cot(c + dx)}} + \frac{B \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d} - \frac{B \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

output

```
1/2*B*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+1/2*B*arctan(1+2^(1/2)
*cot(d*x+c)^(1/2))/d*2^(1/2)+1/4*B*ln(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2
))/d*2^(1/2)-1/4*B*ln(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/d*2^(1/2)+2*B
/d/cot(d*x+c)^(1/2)
```

3.610.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.51

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$= \frac{B \left(2 + \arctan \left(\sqrt[4]{-\cot^2(c + dx)} \right) \sqrt[4]{-\cot^2(c + dx)} - \operatorname{arctanh} \left(\sqrt[4]{-\cot^2(c + dx)} \right) \sqrt[4]{-\cot^2(c + dx)} \right)}{d \sqrt{\cot(c + dx)}}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])),x]`

output `(B*(2 + ArcTan[(-Cot[c + d*x]^2)^(1/4)]*(-Cot[c + d*x]^2)^(1/4) - ArcTanh[(-Cot[c + d*x]^2)^(1/4)]*(-Cot[c + d*x]^2)^(1/4)))/(d*Sqrt[Cot[c + d*x]])`

3.610.3 Rubi [A] (verified)Time = 0.43 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {2011, 3042, 3955, 3042, 3957, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$\downarrow \text{2011}$$

$$B \int \frac{1}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$B \int \frac{1}{(-\tan(c + dx + \frac{\pi}{2}))^{3/2}} dx$$

$$\downarrow \text{3955}$$

$$B \left(\frac{2}{d \sqrt{\cot(c + dx)}} - \int \sqrt{\cot(c + dx)} dx \right)$$

$$\begin{aligned}
& \downarrow 3042 \\
& B \left(\frac{2}{d\sqrt{\cot(c+dx)}} - \int \sqrt{-\tan\left(c+dx+\frac{\pi}{2}\right)} dx \right) \\
& \downarrow 3957 \\
& B \left(\frac{\int \frac{\sqrt{\cot(c+dx)}}{\cot^2(c+dx)+1} d\cot(c+dx)}{d} + \frac{2}{d\sqrt{\cot(c+dx)}} \right) \\
& \downarrow 266 \\
& B \left(\frac{2 \int \frac{\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)}}{d} + \frac{2}{d\sqrt{\cot(c+dx)}} \right) \\
& \downarrow 826 \\
& B \left(\frac{2 \left(\frac{1}{2} \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} - \frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d} + \frac{2}{d\sqrt{\cot(c+dx)}} \right) \\
& \downarrow 1476 \\
& B \left(\frac{2 \left(\frac{1}{2} \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) - \frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d} \right) \\
& \downarrow 1082 \\
& B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)-1} d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d} \right) \\
& \downarrow 217 \\
& B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) - \frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d\sqrt{\cot(c+dx)} \right)}{d} + \frac{2}{d\sqrt{\cot(c+dx)}} \right) \\
& \downarrow 1479
\end{aligned}$$

3.610. $\int \frac{aB+bB \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\int -\frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} + \frac{\int -\frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \right)$$

↓ 25

$$B \left(\frac{2 \left(\frac{1}{2} \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \right)$$

↓ 27

$$B \left(\frac{2 \left(\frac{1}{2} \left(-\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \right)$$

↓ 1103

$$B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log(\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d} \right)$$

input `Int[(a*B + b*B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])),x]`

output `B*(2/(d*Sqrt[Cot[c + d*x]]) + (2*((-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2])/2 + (Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2]) - Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2]))/2))/d)`

3.610.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplifierQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3955 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x])^(n + 1)/(b*d*(n + 1)), x] - Simp[1/b^2 Int[(b*Tan[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

3.610.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.66

method	result	size
default	$\frac{B \left(\frac{\sqrt{2} \left(\ln \left(\frac{1 + \cot(dx+c) - \sqrt{2} \sqrt{\cot(dx+c)}}{1 + \cot(dx+c) + \sqrt{2} \sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1 + \sqrt{2} \sqrt{\cot(dx+c)}}{2} \right) + 2 \arctan \left(\frac{-1 + \sqrt{2} \sqrt{\cot(dx+c)}}{2} \right) \right)}{4} + \frac{2}{\sqrt{\cot(dx+c)}} \right)}{d}$	102

input `int((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `B/d*(1/4*2^(1/2)*(ln((1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))+2/cot(d*x+c)^(1/2))`

3.610. $\int \frac{aB+bB \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$

3.610.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 471, normalized size of antiderivative = 3.06

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$4 B \sqrt{\frac{\cos(2 dx + 2 c) + 1}{\sin(2 dx + 2 c)}} \sin(2 dx + 2 c) - (d \cos(2 dx + 2 c) + d) \left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log \left(\frac{B \sqrt{\frac{\cos(2 dx + 2 c) + 1}{\sin(2 dx + 2 c)}} \sin(2 dx + 2 c) + (d \cos(2 dx + 2 c) + d)}{\cos(2 dx + 2 c) + 1} \right)$$

```
input integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x, algorithm="fracas")
```

```
output 1/2*(4*B*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) -
(d*cos(2*d*x + 2*c) + d)*(-B^4/d^4)^(1/4)*log((B*sqrt((cos(2*d*x + 2*c) +
1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) + (d*cos(2*d*x + 2*c) + d)*(-B^4/d^4)^(1/4))/(cos(2*d*x + 2*c) + 1)) +
(d*cos(2*d*x + 2*c) + d)*(-B^4/d^4)^(1/4)*log((B*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) -
(d*cos(2*d*x + 2*c) + d)*(-B^4/d^4)^(1/4))/(cos(2*d*x + 2*c) + 1)) + (I*d*cos(2*d*x + 2*c) + I*d)*(-B^4/d^4)^(1/4)*log((B*sqrt((cos(2*d*x + 2*c) +
1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) - (I*d*cos(2*d*x + 2*c) + I*d)*(-B^4/d^4)^(1/4))/(cos(2*d*x + 2*c) + 1)) +
(-I*d*cos(2*d*x + 2*c) - I*d)*(-B^4/d^4)^(1/4)*log((B*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) -
(-I*d*cos(2*d*x + 2*c) - I*d)*(-B^4/d^4)^(1/4))/(cos(2*d*x + 2*c) + 1)))/(d*cos(2*d*x + 2*c) + d)
```

3.610.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = B \int \frac{1}{\cot^{\frac{3}{2}}(c + dx)} dx$$

```
input integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)**(3/2)/(a+b*tan(d*x+c)),x)
```

```
output B*Integral(cot(c + d*x)**(-3/2), x)
```

3.610. $\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$

3.610.7 Maxima [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.82

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$= \frac{\left(2\sqrt{2} \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 2\sqrt{2} \arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) - \sqrt{2} \log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}}\right)\right)}{4d}$$

input `integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/4*((2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) - sqrt(2)*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + sqrt(2)*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1))*B + 8*B*sqrt(tan(d*x + c)))/d`

3.610.8 Giac [F]

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = \int \frac{Bb \tan(dx + c) + Ba}{(b \tan(dx + c) + a) \cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*b*tan(d*x + c) + B*a)/((b*tan(d*x + c) + a)*cot(d*x + c)^(3/2)), x)`

3.610.9 Mupad [B] (verification not implemented)

Time = 9.02 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.42

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx = \frac{2B}{d \sqrt{\frac{1}{\tan(c+dx)}}} + \frac{(-1)^{1/4} B \operatorname{atan}\left((-1)^{1/4} \sqrt{\frac{1}{\tan(c+dx)}}\right)}{d} - \frac{(-1)^{1/4} B \operatorname{atanh}\left((-1)^{1/4} \sqrt{\frac{1}{\tan(c+dx)}}\right)}{d}$$

3.610. $\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))} dx$

input `int((B*a + B*b*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))),x)`

output `(2*B)/(d*(1/tan(c + d*x))^(1/2)) + ((-1)^(1/4)*B*atan((-1)^(1/4)*(1/tan(c + d*x))^(1/2)))/d - ((-1)^(1/4)*B*atanh((-1)^(1/4)*(1/tan(c + d*x))^(1/2)))/d`

3.610.
$$\int \frac{aB + bB \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))} dx$$

3.611
$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

3.611.1 Optimal result 5844
 3.611.2 Mathematica [A] (verified) 5845
 3.611.3 Rubi [A] (verified) 5845
 3.611.4 Maple [A] (verified) 5849
 3.611.5 Fricas [C] (verification not implemented) 5850
 3.611.6 Sympy [F] 5850
 3.611.7 Maxima [A] (verification not implemented) 5851
 3.611.8 Giac [F] 5851
 3.611.9 Mupad [B] (verification not implemented) 5852

3.611.1 Optimal result

Integrand size = 36, antiderivative size = 156

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = -\frac{B \arctan\left(1 - \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d} + \frac{B \arctan\left(1 + \sqrt{2}\sqrt{\cot(c + dx)}\right)}{\sqrt{2}d} + \frac{2B}{3d \cot^{\frac{3}{2}}(c + dx)} - \frac{B \log\left(1 - \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d} + \frac{B \log\left(1 + \sqrt{2}\sqrt{\cot(c + dx)} + \cot(c + dx)\right)}{2\sqrt{2}d}$$

output
$$\frac{2}{3} \frac{B}{d} \frac{1}{\cot(dx+c)^{3/2}} + \frac{1}{2} \frac{B \arctan(-1 + 2^{1/2} \cot(dx+c)^{1/2})}{d 2^{1/2}} + \frac{1}{2} \frac{B \arctan(1 + 2^{1/2} \cot(dx+c)^{1/2})}{d 2^{1/2}} - \frac{1}{4} \frac{B \ln(1 + \cot(dx+c) - 2^{1/2} \cot(dx+c)^{1/2})}{d 2^{1/2}} + \frac{1}{4} \frac{B \ln(1 + \cot(dx+c) + 2^{1/2} \cot(dx+c)^{1/2})}{d 2^{1/2}}$$

3.611.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.53

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = \frac{B \left(-2 + 3 \arctan \left(\sqrt[4]{-\cot^2(c + dx)} \right) (-\cot^2(c + dx))^{\frac{3}{4}} + 3 \operatorname{arctanh} \left(\sqrt[4]{-\cot^2(c + dx)} \right) (-\cot^2(c + dx))^{\frac{3}{4}} \right)}{3d \cot^{\frac{3}{2}}(c + dx)}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])),x]`

output `-1/3*(B*(-2 + 3*ArcTan[(-Cot[c + d*x]^2)^(1/4)]*(-Cot[c + d*x]^2)^(3/4) + 3*ArcTanh[(-Cot[c + d*x]^2)^(1/4)]*(-Cot[c + d*x]^2)^(3/4)))/(d*Cot[c + d*x]^(3/2))`

3.611.3 Rubi [A] (verified)Time = 0.42 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {2011, 3042, 3955, 3042, 3957, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx \\ & \quad \downarrow \text{2011} \\ & B \int \frac{1}{\cot^{\frac{5}{2}}(c + dx)} dx \\ & \quad \downarrow \text{3042} \\ & B \int \frac{1}{(-\tan(c + dx + \frac{\pi}{2}))^{\frac{5}{2}}} dx \\ & \quad \downarrow \text{3955} \\ & B \left(\frac{2}{3d \cot^{\frac{3}{2}}(c + dx)} - \int \frac{1}{\sqrt{\cot(c + dx)}} dx \right) \end{aligned}$$

3.611. $\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& B \left(\frac{2}{3d \cot^{\frac{3}{2}}(c+dx)} - \int \frac{1}{\sqrt{-\tan(c+dx+\frac{\pi}{2})}} dx \right) \\
& \downarrow 3957 \\
& B \left(\frac{\int \frac{1}{\sqrt{\cot(c+dx)(\cot^2(c+dx)+1)}} d \cot(c+dx)}{d} + \frac{2}{3d \cot^{\frac{3}{2}}(c+dx)} \right) \\
& \downarrow 266 \\
& B \left(\frac{2 \int \frac{1}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)}}{d} + \frac{2}{3d \cot^{\frac{3}{2}}(c+dx)} \right) \\
& \downarrow 755 \\
& B \left(\frac{2 \left(\frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{\cot(c+dx)+1}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)} \right)}{d} + \frac{2}{3d \cot^{\frac{3}{2}}(c+dx)} \right) \\
& \downarrow 1476 \\
& B \left(\frac{2 \left(\frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)} + \frac{1}{2} \left(\frac{1}{2} \int \frac{1}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d \sqrt{\cot(c+dx)} + \frac{1}{2} \int \frac{1}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d \sqrt{\cot(c+dx)} \right) \right)}{d} \right) \\
& \downarrow 1082 \\
& B \left(\frac{2 \left(\frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)} + \frac{1}{2} \left(\frac{\int \frac{1}{-\cot(c+dx)-1} d(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} - \frac{\int \frac{1}{-\cot(c+dx)-1} d(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right) \right)}{d} \right) \\
& \downarrow 217 \\
& B \left(\frac{2 \left(\frac{1}{2} \int \frac{1-\cot(c+dx)}{\cot^2(c+dx)+1} d \sqrt{\cot(c+dx)} + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) \right)}{d} + \frac{2}{3d \cot^{\frac{3}{2}}(c+dx)} \right) \\
& \downarrow 1479
\end{aligned}$$

3.611. $\int \frac{aB+bB \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$

$$B \left(\frac{2 \left(\frac{1}{2} \left(- \frac{\int - \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} - \frac{\int - \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right)}{d} \right)$$

↓ 25

$$B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right)}{d} \right)$$

↓ 27

$$B \left(\frac{2 \left(\frac{1}{2} \left(\int \frac{\sqrt{2}-2\sqrt{\cot(c+dx)}}{\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)}}{2\sqrt{2}} + \frac{1}{2} \int \frac{\sqrt{2}\sqrt{\cot(c+dx)+1}}{\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1}} d\sqrt{\cot(c+dx)} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} \right)}{d} \right)$$

↓ 1103

$$B \left(\frac{2 \left(\frac{1}{2} \left(\frac{\arctan(\sqrt{2}\sqrt{\cot(c+dx)+1})}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}\sqrt{\cot(c+dx)})}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log(\cot(c+dx)+\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} - \frac{\log(\cot(c+dx)-\sqrt{2}\sqrt{\cot(c+dx)+1})}{2\sqrt{2}} \right) \right)}{d} \right)$$

input `Int[(a*B + b*B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])),x]`

output `B*(2/(3*d*Cot[c + d*x]^(3/2)) + (2*((-(ArcTan[1 - Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2]) + ArcTan[1 + Sqrt[2]*Sqrt[Cot[c + d*x]])/Sqrt[2])/2 + (-1/2*Log[1 - Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/Sqrt[2] + Log[1 + Sqrt[2]*Sqrt[Cot[c + d*x]] + Cot[c + d*x]]/(2*Sqrt[2]))/2)/d)`

3.611.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 266 `Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`
- rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3955 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(b*Tan[c + d*x])^(n + 1)/(b*d*(n + 1)), x] - Simp[1/b^2 Int[(b*Tan[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1]`

rule 3957 `Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b/d Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]`

3.611.4 Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.65

method	result	size
default	$B \left(\frac{2}{3 \cot(dx+c)^{\frac{3}{2}}} + \frac{\sqrt{2} \left(\ln \left(\frac{1+\cot(dx+c)+\sqrt{2}\sqrt{\cot(dx+c)}}{1+\cot(dx+c)-\sqrt{2}\sqrt{\cot(dx+c)}} \right) + 2 \arctan \left(\frac{1+\sqrt{2}\sqrt{\cot(dx+c)}}{-1+\sqrt{2}\sqrt{\cot(dx+c)}} \right) \right)}{4} \right) / d$	102

input `int((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `B/d*(2/3/cot(d*x+c)^(3/2)+1/4*2^(1/2)*(ln((1+cot(d*x+c)+2^(1/2)*cot(d*x+c)^(1/2))/(1+cot(d*x+c)-2^(1/2)*cot(d*x+c)^(1/2)))+2*arctan(1+2^(1/2)*cot(d*x+c)^(1/2))+2*arctan(-1+2^(1/2)*cot(d*x+c)^(1/2))))`

3.611. $\int \frac{aB+bB \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))} dx$

3.611.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 503, normalized size of antiderivative = 3.22

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx =$$

$$3(d \cos(2dx + 2c) + d) \left(-\frac{B^4}{d^4}\right)^{\frac{1}{4}} \log \left(\frac{B^3 \sqrt{\frac{\cos(2dx+2c)+1}{\sin(2dx+2c)}} \sin(2dx+2c) + (d^3 \cos(2dx+2c) + d^3) \left(-\frac{B^4}{d^4}\right)^{\frac{3}{4}}}{\cos(2dx+2c)+1} \right) - 3(d \cos($$

```
input integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x, algorithm="fracas")
```

```
output -1/6*(3*(d*cos(2*d*x + 2*c) + d)*(-B^4/d^4)^(1/4)*log((B^3*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) + (d^3*cos(2*d*x + 2*c) + d^3)*(-B^4/d^4)^(3/4))/(cos(2*d*x + 2*c) + 1)) - 3*(d*cos(2*d*x + 2*c) + d)*(-B^4/d^4)^(1/4)*log((B^3*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) - (d^3*cos(2*d*x + 2*c) + d^3)*(-B^4/d^4)^(3/4))/(cos(2*d*x + 2*c) + 1)) + 3*(I*d*cos(2*d*x + 2*c) + I*d)*(-B^4/d^4)^(1/4)*log((B^3*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) - (I*d^3*cos(2*d*x + 2*c) + I*d^3)*(-B^4/d^4)^(3/4))/(cos(2*d*x + 2*c) + 1)) + 3*(-I*d*cos(2*d*x + 2*c) - I*d)*(-B^4/d^4)^(1/4)*log((B^3*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c))*sin(2*d*x + 2*c) - (-I*d^3*cos(2*d*x + 2*c) - I*d^3)*(-B^4/d^4)^(3/4))/(cos(2*d*x + 2*c) + 1)) + 4*(B*cos(2*d*x + 2*c) - B)*sqrt((cos(2*d*x + 2*c) + 1)/sin(2*d*x + 2*c)))/(d*cos(2*d*x + 2*c) + d)
```

3.611.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = B \int \frac{1}{\cot^{\frac{5}{2}}(c + dx)} dx$$

```
input integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)**(5/2)/(a+b*tan(d*x+c)),x)
```

```
output B*Integral(cot(c + d*x)**(-5/2), x)
```

3.611. $\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx$

3.611.7 Maxima [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.82

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx$$

$$= \frac{6\sqrt{2}B \arctan\left(\frac{1}{2}\sqrt{2}\left(\sqrt{2} + \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 6\sqrt{2}B \arctan\left(-\frac{1}{2}\sqrt{2}\left(\sqrt{2} - \frac{2}{\sqrt{\tan(dx+c)}}\right)\right) + 3\sqrt{2}B \log\left(\frac{\sqrt{2}}{\sqrt{\tan(dx+c)}} + 1\right) + 8B \tan(dx+c)^{\frac{3}{2}}}{12d}$$

input `integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x, algorithm="maxima")`

output `1/12*(6*sqrt(2)*B*arctan(1/2*sqrt(2)*(sqrt(2) + 2/sqrt(tan(d*x + c)))) + 6*sqrt(2)*B*arctan(-1/2*sqrt(2)*(sqrt(2) - 2/sqrt(tan(d*x + c)))) + 3*sqrt(2)*B*log(sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) - 3*sqrt(2)*B*log(-sqrt(2)/sqrt(tan(d*x + c)) + 1/tan(d*x + c) + 1) + 8*B*tan(d*x + c)^(3/2))/d`

3.611.8 Giac [F]

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = \int \frac{Bb \tan(dx + c) + Ba}{(b \tan(dx + c) + a) \cot(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*b*tan(d*x + c) + B*a)/((b*tan(d*x + c) + a)*cot(d*x + c)^(5/2)), x)`

3.611.9 Mupad [B] (verification not implemented)

Time = 9.89 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.42

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))} dx = \frac{2B}{3d \left(\frac{1}{\tan(c+dx)}\right)^{3/2}} - \frac{(-1)^{1/4} B \operatorname{atan}\left((-1)^{1/4} \sqrt{\frac{1}{\tan(c+dx)}}\right) \operatorname{li}}{d} - \frac{(-1)^{1/4} B \operatorname{atanh}\left((-1)^{1/4} \sqrt{\frac{1}{\tan(c+dx)}}\right) \operatorname{li}}{d}$$

input `int((B*a + B*b*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + b*tan(c + d*x))),x)`output `(2*B)/(3*d*(1/tan(c + d*x))^(3/2)) - ((-1)^(1/4)*B*atan((-1)^(1/4)*(1/tan(c + d*x))^(1/2))*1i)/d - ((-1)^(1/4)*B*atanh((-1)^(1/4)*(1/tan(c + d*x))^(1/2))*1i)/d`

3.612 $\int \cot^{\frac{9}{2}}(c+dx) \sqrt{a + b \tan(c + dx)} (A+B \tan(c+dx)) dx$

3.612.1 Optimal result	5853
3.612.2 Mathematica [A] (verified)	5854
3.612.3 Rubi [A] (verified)	5854
3.612.4 Maple [B] (warning: unable to verify)	5861
3.612.5 Fricas [B] (verification not implemented)	5861
3.612.6 Sympy [F(-1)]	5861
3.612.7 Maxima [F]	5862
3.612.8 Giac [F(-2)]	5862
3.612.9 Mupad [F(-1)]	5862

3.612.1 Optimal result

Integrand size = 35, antiderivative size = 354

$$\int \cot^{\frac{9}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= -\frac{\sqrt{ia - b}(iA - B) \arctan\left(\frac{\sqrt{ia - b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d}$$

$$- \frac{\sqrt{ia + b}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia + b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{2(35a^2 Ab - 8Ab^3 + 105a^3 B + 14ab^2 B) \sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}}{105a^3 d}$$

$$+ \frac{2(35a^2 A + 4Ab^2 - 7abB) \cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{105a^2 d}$$

$$- \frac{2(Ab + 7aB) \cot^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{35ad} - \frac{2A \cot^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{7d}$$

output

```
-(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a-b)^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a+b)^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+2/105*(35*A*a^2+4*A*b^2-7*B*a*b)*cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(1/2)/a^2/d-2/35*(A*b+7*B*a)*cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(1/2)/a/d-2/7*A*cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^(1/2)/d+2/105*(35*A*a^2*b-8*A*b^3+105*B*a^3+14*B*a*b^2)*cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/a^3/d
```

3.612.2 Mathematica [A] (verified)

Time = 4.49 (sec) , antiderivative size = 291, normalized size of antiderivative = 0.82

$$\int \cot^{\frac{9}{2}}(c+dx) \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$= \cot^{\frac{7}{2}}(c+dx) \left(105 \sqrt[4]{-1} a^3 \sqrt{-a+ib} (iA+B) \arctan \left(\frac{\sqrt[4]{-1} \sqrt{-a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right) \tan^{\frac{7}{2}}(c+dx) - 105(-1)^{3/4} \right)$$

input `Integrate[Cot[c + d*x]^(9/2)*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(Cot[c + d*x]^(7/2)*(105*(-1)^(1/4)*a^3*Sqrt[-a + I*b]*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(7/2) - 105*(-1)^(3/4)*a^3*Sqrt[a + I*b]*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(7/2) + 2*Sqrt[a + b*Tan[c + d*x]]*(-15*a^3*A - 3*a^2*(A*b + 7*a*B)*Tan[c + d*x] + a*(35*a^2*A + 4*A*b^2 - 7*a*b*B)*Tan[c + d*x]^2 + (35*a^2*A*b - 8*A*b^3 + 105*a^3*B + 14*a*b^2*B)*Tan[c + d*x]^3))/(105*a^3*d)`

3.612.3 Rubi [A] (verified)

Time = 2.64 (sec) , antiderivative size = 376, normalized size of antiderivative = 1.06, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 4729, 3042, 4091, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{9}{2}}(c+dx) \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{9/2} \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \int \frac{\sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

3.612. $\int \cot^{\frac{9}{2}}(c+dx) \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \int \frac{\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\tan(c+dx)^{9/2}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2}{7} \int \frac{-6Ab\tan^2(c+dx) - 7(aA - bB)\tan(c+dx) + Ab + 7aB}{2\tan^{7/2}(c+dx)\sqrt{a+b\tan(c+dx)}} dx - \frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{7/2}(c+dx)} \right) \\
& \quad \downarrow \text{4091} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2}{7} \int \frac{-6Ab\tan^2(c+dx) - 7(aA - bB)\tan(c+dx) + Ab + 7aB}{2\tan^{7/2}(c+dx)\sqrt{a+b\tan(c+dx)}} dx - \frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{7/2}(c+dx)} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \int \frac{-6Ab\tan^2(c+dx) - 7(aA - bB)\tan(c+dx) + Ab + 7aB}{\tan^{7/2}(c+dx)\sqrt{a+b\tan(c+dx)}} dx - \frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{7/2}(c+dx)} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \int \frac{-6Ab\tan(c+dx)^2 - 7(aA - bB)\tan(c+dx) + Ab + 7aB}{\tan(c+dx)^{7/2}\sqrt{a+b\tan(c+dx)}} dx - \frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{7/2}(c+dx)} \right) \\
& \quad \downarrow \text{4132} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(-\frac{2 \int \frac{35Aa^2 - 7bBa + 35(Ab+aB)\tan(c+dx)a + 4Ab^2 + 4b(Ab+7aB)\tan^2(c+dx)}{2\tan^{5/2}(c+dx)\sqrt{a+b\tan(c+dx)}} dx}{5a} - \frac{2(7aB + Ab)}{5ad} \right) \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(-\frac{\int \frac{35Aa^2 - 7bBa + 35(Ab+aB)\tan(c+dx)a + 4Ab^2 + 4b(Ab+7aB)\tan^2(c+dx)}{\tan^{5/2}(c+dx)\sqrt{a+b\tan(c+dx)}} dx}{5a} - \frac{2(7aB + Ab)}{5ad} \right) \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(-\frac{\int \frac{35Aa^2 - 7bBa + 35(Ab+aB)\tan(c+dx)a + 4Ab^2 + 4b(Ab+7aB)\tan(c+dx)^2}{\tan(c+dx)^{5/2}\sqrt{a+b\tan(c+dx)}} dx}{5a} - \frac{2(7aB + Ab)}{5ad} \right) \right) \\
& \quad \downarrow \text{4132}
\end{aligned}$$

3.612. $\int \cot^{9/2}(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(- \frac{2 \int - \frac{105Ba^3+35Aba^2-105(aA-bB)\tan(c+dx)a^2+14b^2Ba-8Ab^3-2b(35Aa^2-7bBa+4Ab^2)\tan^2(c+dx)}{2 \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{5a}{5a} \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(- \frac{\int \frac{105Ba^3+35Aba^2-105(aA-bB)\tan(c+dx)a^2+14b^2Ba-8Ab^3-2b(35Aa^2-7bBa+4Ab^2)\tan^2(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{5a}{5a} \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(- \frac{\int \frac{105Ba^3+35Aba^2-105(aA-bB)\tan(c+dx)a^2+14b^2Ba-8Ab^3-2b(35Aa^2-7bBa+4Ab^2)\tan(c+dx)^2}{\tan(c+dx)^{3/2}\sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{5a}{5a} \right) \right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(- \frac{2 \int \frac{105((aA-bB)a^3+(Ab+aB)\tan(c+dx)a^3)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(105a^3B+35a^2Ab+14ab^2B-8Ab^3)\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}{3a} - \frac{5a}{5a} \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(- \frac{105 \int \frac{(aA-bB)a^3+(Ab+aB)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(105a^3B+35a^2Ab+14ab^2B-8Ab^3)\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}{3a} - \frac{5a}{5a} \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(-\frac{105 \int \frac{(aA-bB)a^3+(Ab+aB)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a} - \frac{2(105a^3B+35a^2Ab+14ab^2B-8Ab^3)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} - \frac{2(35a^2A-7abB)}{3a} \right) - \frac{2(35a^2A-7abB)}{5a} \right)$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(-\frac{2(7aB+Ab)\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-7abB)}{3a} \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(-\frac{2(7aB+Ab)\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-7abB)}{3a} \right) \right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(-\frac{2(7aB+Ab)\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-7abB)}{3a} \right) \right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(-\frac{2(7aB+Ab)\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-7abB)}{3a} \right) \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(-\frac{2(7aB+Ab)\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-7abB)}{3a} \right) \right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(-\frac{2(7aB+Ab)\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-7abB)}{3a} \right) \right)$$

input `Int[Cot[c + d*x]^(9/2)*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A*Sqrt[a + b*Tan[c + d*x]])/(7*d*Tan[c + d*x]^(7/2)) + ((-2*(A*b + 7*a*B)*Sqrt[a + b*Tan[c + d*x]])/(5*a*d*Tan[c + d*x]^(5/2)) - ((-2*(35*a^2*A + 4*A*b^2 - 7*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(3*a*d*Tan[c + d*x]^(3/2)) + ((-105*((a^3*(a + I*b)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a - b]*d) + (a^3*(a - I*b)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a + b]*d)))/a - (2*(35*a^2*A*b - 8*A*b^3 + 105*a^3*B + 14*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(a*d*Sqrt[Tan[c + d*x]]))/(3*a))/(5*a))/7`

3.612.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4091 `Int[(((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegerQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && ! (ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.612.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.38 (sec) , antiderivative size = 2185615, normalized size of antiderivative = 6174.05

output too large to display

input `int(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x)`

output `result too large to display`

3.612.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8179 vs. $2(291) = 582$.

Time = 1.40 (sec) , antiderivative size = 8179, normalized size of antiderivative = 23.10

$$\int \cot^{\frac{9}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `Too large to include`

3.612.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(9/2)*(a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.612.7 Maxima [F]

$$\int \cot^{\frac{9}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \cot(dx + c)^{\frac{9}{2}} dx$$

input `integrate(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*cot(d*x + c)^(9/2)
, x)`

3.612.8 Giac [F(-2)]

Exception generated.

$$\int \cot^{\frac{9}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Exception raised: TypeError}$$

input `integrate(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algo
rithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(con
st gen &`

3.612.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{9/2} (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx$$

input `int(cot(c + d*x)^(9/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`

output `int(cot(c + d*x)^(9/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2), x)`

3.613 $\int \cot^{\frac{7}{2}}(c+dx) \sqrt{a + b \tan(c + dx)} (A+B \tan(c+dx)) dx$

3.613.1 Optimal result	5864
3.613.2 Mathematica [A] (verified)	5865
3.613.3 Rubi [A] (verified)	5865
3.613.4 Maple [B] (warning: unable to verify)	5871
3.613.5 Fricas [B] (verification not implemented)	5871
3.613.6 Sympy [F(-1)]	5871
3.613.7 Maxima [F]	5872
3.613.8 Giac [F(-2)]	5872
3.613.9 Mupad [F(-1)]	5872

3.613.1 Optimal result

Integrand size = 35, antiderivative size = 290

$$\int \cot^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\sqrt{ia - b}(A + iB) \arctan\left(\frac{\sqrt{ia - b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d} - \frac{\sqrt{ia + b}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia + b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d} + \frac{2(15a^2A + 2Ab^2 - 5abB) \sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}}{15a^2d} - \frac{2(Ab + 5aB) \cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{15ad} - \frac{2A \cot^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{5d}$$

```
output (A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a-b)^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a+b)^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-2/15*(A*b+5*B*a)*cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(1/2)/a/d-2/5*A*cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(1/2)/d+2/15*(15*A*a^2+2*A*b^2-5*B*a*b)*cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/a^2/d
```

3.613.2 Mathematica [A] (verified)

Time = 2.48 (sec) , antiderivative size = 252, normalized size of antiderivative = 0.87

$$\int \cot^{\frac{7}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$= \cot^{\frac{5}{2}}(c+dx) \left(15\sqrt[4]{-1}a^2\sqrt{-a+ib}(A-iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) \tan^{\frac{5}{2}}(c+dx) + 15\sqrt[4]{-1}a^2\sqrt{a+ib}(A+iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) \tan^{\frac{5}{2}}(c+dx) + 2\sqrt{a+b\tan(c+dx)}(-3a^2A - a(Ab+5aB)\tan(c+dx) + (15a^2A + 2Ab^2 - 5abB)\tan^2(c+dx)) \right) / (15a^2d)$$

input `Integrate[Cot[c + d*x]^(7/2)*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(Cot[c + d*x]^(5/2)*(15*(-1)^(1/4)*a^2*Sqrt[-a + I*b]*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(5/2) + 15*(-1)^(1/4)*a^2*Sqrt[a + I*b]*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(5/2) + 2*Sqrt[a + b*Tan[c + d*x]]*(-3*a^2*A - a*(A*b + 5*a*B)*Tan[c + d*x] + (15*a^2*A + 2*A*b^2 - 5*a*b*B)*Tan[c + d*x]^2))/(15*a^2*d)`

3.613.3 Rubi [A] (verified)

Time = 1.94 (sec) , antiderivative size = 310, normalized size of antiderivative = 1.07, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.514$, Rules used = {3042, 4729, 3042, 4091, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{7}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{7/2}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)}dx$$

$$\downarrow \text{3042}$$

3.613. $\int \cot^{\frac{7}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\int\frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\tan(c+dx)^{7/2}}dx$$

↓ 4091

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2}{5}\int-\frac{-4Ab\tan^2(c+dx)-5(aA-bB)\tan(c+dx)+Ab+5aB}{2\tan^{5/2}(c+dx)\sqrt{a+b\tan(c+dx)}}dx-\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{5/2}(c+dx)}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\int\frac{-4Ab\tan^2(c+dx)-5(aA-bB)\tan(c+dx)+Ab+5aB}{\tan^{5/2}(c+dx)\sqrt{a+b\tan(c+dx)}}dx-\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{5/2}(c+dx)}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\int\frac{-4Ab\tan(c+dx)^2-5(aA-bB)\tan(c+dx)+Ab+5aB}{\tan(c+dx)^{5/2}\sqrt{a+b\tan(c+dx)}}dx-\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{5/2}(c+dx)}\right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(-\frac{2\int\frac{15Aa^2-5bBa+15(Ab+aB)\tan(c+dx)a+2Ab^2+2b(Ab+5aB)\tan^2(c+dx)}{2\tan^{3/2}(c+dx)\sqrt{a+b\tan(c+dx)}}dx}{3a}-\frac{2(5aB+Ab)}{3ad}\right)\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(-\frac{\int\frac{15Aa^2-5bBa+15(Ab+aB)\tan(c+dx)a+2Ab^2+2b(Ab+5aB)\tan^2(c+dx)}{\tan^{3/2}(c+dx)\sqrt{a+b\tan(c+dx)}}dx}{3a}-\frac{2(5aB+Ab)}{3ad}\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(-\frac{\int\frac{15Aa^2-5bBa+15(Ab+aB)\tan(c+dx)a+2Ab^2+2b(Ab+5aB)\tan(c+dx)^2}{\tan(c+dx)^{3/2}\sqrt{a+b\tan(c+dx)}}dx}{3a}-\frac{2(5aB+Ab)}{3ad}\right)\right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(-\frac{2\int-\frac{15(a^2(Ab+aB)-a^2(aA-bB)\tan(c+dx))}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx}{3a}-\frac{2(15a^2A-5abB+2Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}{3a}-\frac{2(5aB+Ab)}{3ad}\right)\right)$$

3.613. $\int \cot^{7/2}(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(-\frac{15\int\frac{a^2(Ab+aB)-a^2(aA-bB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx}{a}-\frac{2(15a^2A-5abB+2Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}\right)-\frac{2(5aB)}{3a}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(-\frac{15\int\frac{a^2(Ab+aB)-a^2(aA-bB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx}{a}-\frac{2(15a^2A-5abB+2Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}\right)-\frac{2(5aB)}{3a}\right)$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}+\frac{1}{5}\left(-\frac{2(5aB+Ab)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)}-\frac{2(15a^2A-5abB)}{ad}\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}+\frac{1}{5}\left(-\frac{2(5aB+Ab)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)}-\frac{2(15a^2A-5abB)}{ad}\right)\right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}+\frac{1}{5}\left(-\frac{2(5aB+Ab)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)}-\frac{2(15a^2A-5abB)}{ad}\right)\right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} + \frac{1}{5} \left(-\frac{2(5aB+Ab)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-5abB)}{ad} \right) \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} + \frac{1}{5} \left(-\frac{2(5aB+Ab)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-5abB)}{ad} \right) \right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} + \frac{1}{5} \left(-\frac{2(5aB+Ab)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{2(15a^2A-5abB)}{ad} \right) \right)$$

input `Int[Cot[c + d*x]^(7/2)*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A*Sqrt[a + b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) + ((-2*(A*b + 5*a*B)*Sqrt[a + b*Tan[c + d*x]])/(3*a*d*Tan[c + d*x]^(3/2)) - ((15*((a^2*(a + I*b)*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a - b]*d)) + (a^2*(a - I*b)*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a + b]*d)))/a - (2*(15*a^2*A + 2*A*b^2 - 5*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(a*d*Sqrt[Tan[c + d*x]]))/(3*a))/5`

3.613.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4091 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegerQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0]))`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.613.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.18 (sec) , antiderivative size = 2183483, normalized size of antiderivative = 7529.25

output too large to display

input `int(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x)`

output `result too large to display`

3.613.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8104 vs. $2(236) = 472$.

Time = 1.40 (sec) , antiderivative size = 8104, normalized size of antiderivative = 27.94

$$\int \cot^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `Too large to include`

3.613.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(7/2)*(a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.613.7 Maxima [F]

$$\int \cot^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \cot(dx + c)^{\frac{7}{2}} dx$$

input `integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*cot(d*x + c)^(7/2)
, x)`

3.613.8 Giac [F(-2)]

Exception generated.

$$\int \cot^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Exception raised: TypeError}$$

input `integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algo
rithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(con
st gen &`

3.613.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{7/2} (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx$$

input `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`

output `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2), x)`

3.614 $\int \cot^{\frac{5}{2}}(c+dx) \sqrt{a + b \tan(c + dx)} (A+B \tan(c+dx)) dx$

3.614.1 Optimal result	5874
3.614.2 Mathematica [A] (verified)	5875
3.614.3 Rubi [A] (verified)	5875
3.614.4 Maple [B] (warning: unable to verify)	5880
3.614.5 Fricas [B] (verification not implemented)	5880
3.614.6 Sympy [F(-1)]	5881
3.614.7 Maxima [F]	5881
3.614.8 Giac [F(-2)]	5881
3.614.9 Mupad [F(-1)]	5882

3.614.1 Optimal result

Integrand size = 35, antiderivative size = 239

$$\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \frac{\sqrt{ia - b}(iA - B) \arctan\left(\frac{\sqrt{ia - b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d} + \frac{\sqrt{ia + b}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia + b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d} - \frac{2(Ab + 3aB) \sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}}{3ad} - \frac{2A \cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{3d}$$

output

```
(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a-b)^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a+b)^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-2/3*A*cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(1/2)/d-2/3*(A*b+3*B*a)*cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/a/d
```

3.614.2 Mathematica [A] (verified)

Time = 2.02 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.90

$$\int \cot^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$= \cot^{\frac{3}{2}}(c+dx) \left(-3\sqrt[4]{-1} a \sqrt{-a+ib} (iA+B) \arctan \left(\frac{\sqrt[4]{-1} \sqrt{-a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right) \tan^{\frac{3}{2}}(c+dx) + 3(-1)^{3/4} a \sqrt{a} \right)$$

input `Integrate[Cot[c + d*x]^(5/2)*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(Cot[c + d*x]^(3/2)*(-3*(-1)^(1/4)*a*Sqrt[-a + I*b]*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(3/2) + 3*(-1)^(3/4)*a*Sqrt[a + I*b]*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(3/2) - 2*Sqrt[a + b*Tan[c + d*x]]*(a*A + (A*b + 3*a*B)*Tan[c + d*x]))/(3*a*d)`

3.614.3 Rubi [A] (verified)Time = 1.39 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.02, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 4729, 3042, 4091, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{5/2} \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \int \frac{\sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx$$

$$\downarrow \text{3042}$$

3.614. $\int \cot^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\tan(c+dx)^{5/2}} dx$$

↓ 4091

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2}{3} \int -\frac{-2Ab\tan^2(c+dx) - 3(aA - bB)\tan(c+dx) + Ab + 3aB}{2\tan^{3/2}(c+dx)\sqrt{a+b\tan(c+dx)}} dx - \frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \int \frac{-2Ab\tan^2(c+dx) - 3(aA - bB)\tan(c+dx) + Ab + 3aB}{\tan^{3/2}(c+dx)\sqrt{a+b\tan(c+dx)}} dx - \frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \int \frac{-2Ab\tan(c+dx)^2 - 3(aA - bB)\tan(c+dx) + Ab + 3aB}{\tan(c+dx)^{3/2}\sqrt{a+b\tan(c+dx)}} dx - \frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)} \right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \left(-\frac{2 \int \frac{3(aA-bB)+a(Ab+aB)\tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a} - \frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right) - \frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \left(-\frac{3 \int \frac{a(aA-bB)+a(Ab+aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a} - \frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right) - \frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \left(-\frac{3 \int \frac{a(aA-bB)+a(Ab+aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a} - \frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right) - \frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)} \right)$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)} + \frac{1}{3} \left(-\frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} - \frac{3\left(\frac{1}{2}a+ib\right)}{ad\sqrt{\tan(c+dx)}} \right) \right)$$

↓ 3042

3.614. $\int \cot^{5/2}(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}+\frac{1}{3}\left(-\frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}-\frac{3\left(\frac{1}{2}a+ib\right)}{\dots}\right)\right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}+\frac{1}{3}\left(-\frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}-\frac{3\left(\frac{a(a-ib)(A-ib)}{\dots}\right)}{\dots}\right)\right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}+\frac{1}{3}\left(-\frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}-\frac{3\left(\frac{a(a+ib)(A+ib)}{\dots}\right)}{\dots}\right)\right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}+\frac{1}{3}\left(-\frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}-\frac{3\left(\frac{a(a-ib)(A-ib)}{\dots}\right)}{\dots}\right)\right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}+\frac{1}{3}\left(-\frac{2(3aB+Ab)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}-\frac{3\left(\frac{a(a+ib)(A+ib)}{\dots}\right)}{\dots}\right)\right)$$

input `Int[Cot[c + d*x]^(5/2)*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A*Sqrt[a + b*Tan[c + d*x]])/(3*
d*Tan[c + d*x]^(3/2)) + ((-3*((a*(a + I*b)*(A + I*B)*ArcTan[(Sqrt[I*a - b]
*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a - b]*d) + (a*(a
- I*b)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan
[c + d*x]]]/(Sqrt[I*a + b]*d)))/a - (2*(A*b + 3*a*B)*Sqrt[a + b*Tan[c + d
*x]])/(a*d*Sqrt[Tan[c + d*x]]))/3)
```

3.614.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 104 Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))/((e_.) + (f_.)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4091 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.614.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.37 (sec) , antiderivative size = 2181430, normalized size of antiderivative = 9127.32

output too large to display

```
input int(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x)
```

```
output result too large to display
```

3.614.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8070 vs. $2(191) = 382$.

Time = 1.39 (sec) , antiderivative size = 8070, normalized size of antiderivative = 33.77

$$\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algo
rithm="fracas")
```

```
output Too large to include
```

3.614.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Timed out}$$

```
input integrate(cot(d*x+c)**(5/2)*(a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)
```

```
output Timed out
```

3.614.7 Maxima [F]

$$\begin{aligned} & \int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \cot(dx + c)^{\frac{5}{2}} dx \end{aligned}$$

```
input integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algo
rithm="maxima")
```

```
output integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*cot(d*x + c)^(5/2)
, x)
```

3.614.8 Giac [F(-2)]

Exception generated.

$$\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Exception raised: TypeError}$$

```
input integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algo
rithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(con
st gen &
```


3.614.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{5/2} (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx$$

input `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`output `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2), x)`

3.615 $\int \cot^{\frac{3}{2}}(c+dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.615.1 Optimal result	5883
3.615.2 Mathematica [A] (verified)	5884
3.615.3 Rubi [A] (verified)	5884
3.615.4 Maple [B] (warning: unable to verify)	5888
3.615.5 Fricas [B] (verification not implemented)	5888
3.615.6 Sympy [F]	5888
3.615.7 Maxima [F]	5889
3.615.8 Giac [F(-2)]	5889
3.615.9 Mupad [F(-1)]	5889

3.615.1 Optimal result

Integrand size = 35, antiderivative size = 194

$$\int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= -\frac{\sqrt{ia - b}(A + iB) \arctan\left(\frac{\sqrt{ia - b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{\sqrt{ia + b}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia + b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d}$$

$$- \frac{2A \sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}}{d}$$

output

```
-(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a-b)^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a+b)^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-2*A*cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/d
```

3.615.2 Mathematica [A] (verified)

Time = 0.52 (sec) , antiderivative size = 189, normalized size of antiderivative = 0.97

$$\int \cot^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx =$$

$$\frac{\sqrt{\cot(c+dx)}\left(\sqrt[4]{-1}\sqrt{-a+ib}(A-iB)\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)\sqrt{\tan(c+dx)}+\sqrt[4]{-1}\sqrt{a+ib}\right)}{d}$$

input `Integrate[Cot[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `-((Sqrt[Cot[c + d*x]]*((-1)^(1/4)*Sqrt[-a + I*b]*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]*Sqrt[Tan[c + d*x]] + (-1)^(1/4)*Sqrt[a + I*b]*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]*Sqrt[Tan[c + d*x]] + 2*A*Sqrt[a + b*Tan[c + d*x]))/d)`

3.615.3 Rubi [A] (verified)Time = 1.04 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.98, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 4729, 3042, 4091, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow 3042$$

$$\int \cot(c+dx)^{3/2}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$$

$$\downarrow 4729$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\int \frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)}dx$$

$$\downarrow 3042$$

3.615. $\int \cot^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\tan(c+dx)^{3/2}} dx$$

↓ 4091

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-2 \int -\frac{Ab+aB-(aA-bB)\tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{2A\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{Ab+aB-(aA-bB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{2A\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{Ab+aB-(aA-bB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{2A\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right)$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{1}{2}(a+ib)(-B+iA) \int \frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}(a-ib)(B+iA) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{1}{2}(a+ib)(-B+iA) \int \frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}(a-ib)(B+iA) \right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(a-ib)(B+iA) \int \frac{1}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} d\tan(c+dx)}{2d} - \frac{(a+ib)(-B+iA) \int \frac{1}{(1+i\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} d\tan(c+dx)}{2d} \right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{(a+ib)(-B+iA) \int \frac{1}{\frac{(ia-b)\tan(c+dx)}{a+b\tan(c+dx)}+1} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} + \frac{(a-ib)(B+iA) \int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(a-ib)(B+iA) \int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} - \frac{(a+ib)(-B+iA) \arctan\left(\frac{\sqrt{-b+ia}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} \right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{(a+ib)(-B+iA) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} + \frac{(a-ib)(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{b+ia}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}} \right)$$

input `Int[Cot[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-(((a + I*b)*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a - b]*d) + ((a - I*b)*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a + b]*d) - (2*A*Sqrt[a + b*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))`

3.615.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4091 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4729 `Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.615.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.29 (sec) , antiderivative size = 2178676, normalized size of antiderivative = 11230.29

output too large to display

input `int(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x)`

output `result too large to display`

3.615.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7949 vs. $2(154) = 308$.

Time = 1.43 (sec) , antiderivative size = 7949, normalized size of antiderivative = 40.97

$$\int \cot^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `Too large to include`

3.615.6 Sympy [F]

$$\begin{aligned} & \int \cot^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx \\ &= \int (A+B \tan(c+dx)) \sqrt{a+b \tan(c+dx)} \cot^{\frac{3}{2}}(c+dx) dx \end{aligned}$$

input `integrate(cot(d*x+c)**(3/2)*(a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))*cot(c + d*x)**(3/2), x)`

3.615. $\int \cot^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$

3.615.7 Maxima [F]

$$\int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a} \cot(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*cot(d*x + c)^(3/2)
, x)`

3.615.8 Giac [F(-2)]

Exception generated.

$$\int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx = \text{Exception raised: TypeError}$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algo
rithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(con
st gen &`

3.615.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{3/2} (A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)} dx$$

input `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`

output `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2), x)`

3.616 $\int \sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$

3.616.1 Optimal result	5891
3.616.2 Mathematica [A] (verified)	5892
3.616.3 Rubi [A] (verified)	5892
3.616.4 Maple [B] (warning: unable to verify)	5896
3.616.5 Fricas [B] (verification not implemented)	5897
3.616.6 Sympy [F]	5897
3.616.7 Maxima [F]	5897
3.616.8 Giac [F(-2)]	5898
3.616.9 Mupad [F(-1)]	5898

3.616.1 Optimal result

Integrand size = 35, antiderivative size = 229

$$\int \sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)} (A + B \tan(c + dx)) dx$$

$$= -\frac{\sqrt{ia - b}(iA - B) \arctan\left(\frac{\sqrt{ia - b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{2\sqrt{b}B \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d}$$

$$- \frac{\sqrt{ia + b}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia + b}\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d}$$

output

```
-(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a-b)^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+2*B*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*b^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*
(I*a+b)^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d
```

3.616.2 Mathematica [A] (verified)

Time = 0.86 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.04

$$\int \sqrt{\cot(c+dx)} \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$= \frac{\sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)} \left(\sqrt[4]{-1} \left(\sqrt{-a+ib} (iA+B) \arctan \left(\frac{\sqrt[4]{-1} \sqrt{-a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right) \right) + \sqrt{a+ib} (-iA) \right)}{d \sqrt{a+}}$$

input `Integrate[Sqrt[Cot[c + d*x]]*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-1)^(1/4)*(Sqrt[-a + I*b]*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])] + Sqrt[a + I*b]*((-I)*A + B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]*Sqrt[a + b*Tan[c + d*x]] + 2*Sqrt[a]*Sqrt[b]*B*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a]))/(d*Sqrt[a + b*Tan[c + d*x]])]`

3.616.3 Rubi [A] (verified)

Time = 1.34 (sec) , antiderivative size = 200, normalized size of antiderivative = 0.87, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 4729, 3042, 4097, 3042, 4099, 3042, 4098, 104, 216, 219, 4117, 65, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\cot(c+dx)} \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$\downarrow 3042$$

$$\int \sqrt{\cot(c+dx)} \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$$

$$\downarrow 4729$$

$$\sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \int \frac{\sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

$$\downarrow 3042$$

3.616. $\int \sqrt{\cot(c+dx)} \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

↓ 4097

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{aA-bB+(Ab+aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + bB \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{aA-bB+(Ab+aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + bB \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \right)$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2}(a+ib)(A+iB) \int \frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}(a-ib)(A-iB) \int \frac{1+i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2}(a+ib)(A+iB) \int \frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{1}{2}(a-ib)(A-iB) \int \frac{1+i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(a-ib)(A-iB) \int \frac{1}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} d\tan(c+dx)}{2d} + \frac{(a+ib)(A+iB) \int \frac{1}{(1+i\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} d\tan(c+dx)}{2d} \right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(a+ib)(A+iB) \int \frac{1}{\frac{(ia-b)\tan(c+dx)}{a+b\tan(c+dx)}+1} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} + \frac{(a-ib)(A-iB) \int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(a-ib)(A-iB) \int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} + bB \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \right)$$

↓ 219

3.616. $\int \sqrt{\cot(c+dx)}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(bB\int\frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx+\frac{(a+ib)(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}}\right)$$

↓ 4117

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{bB\int\frac{1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}d\tan(c+dx)}{d}+\frac{(a+ib)(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}}\right)$$

↓ 65

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2bB\int\frac{1}{1-\frac{b\tan(c+dx)}{a+b\tan(c+dx)}}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d}+\frac{(a+ib)(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}}\right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(a+ib)(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}}+\frac{(a-ib)(A-iB)\operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}}\right)$$

input `Int[Sqrt[Cot[c + d*x]]*Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]),x]`

output `((a + I*b)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a - b]*d) + (2*Sqrt[b]*B*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/d + ((a - I*b)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d))*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]`

3.616.3.1 Defintions of rubi rules used

rule 65 `Int[1/(Sqrt[(b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`

- rule 104 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4097 `Int[(Sqrt[(a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])]/Sqrt[(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Int[Simp[a*A - b*B + (A*b + a*B)*Tan[e + f*x], x]/(Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]), x] + Simp[b*B Int[(1 + Tan[e + f*x]^2)/(Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

```
rule 4099 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2,
0]
```

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.616.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.41 (sec) , antiderivative size = 2178428, normalized size of antiderivative = 9512.79

output too large to display

```
input int(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x)
```

```
output result too large to display
```

3.616.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8075 vs. $2(178) = 356$.

Time = 2.44 (sec) , antiderivative size = 16183, normalized size of antiderivative = 70.67

$$\int \sqrt{\cot(c+dx)}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.616.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{\cot(c+dx)}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx \\ &= \int (A+B\tan(c+dx))\sqrt{a+b\tan(c+dx)}\sqrt{\cot(c+dx)}dx \end{aligned}$$

input `integrate(cot(d*x+c)**(1/2)*(a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))*sqrt(cot(c + d*x)), x)`

3.616.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{\cot(c+dx)}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx \\ &= \int (B\tan(dx+c)+A)\sqrt{b\tan(dx+c)+a}\sqrt{\cot(dx+c)}dx \end{aligned}$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)*sqrt(cot(d*x + c)), x)`

3.616.8 Giac [F(-2)]

Exception generated.

$$\int \sqrt{\cot(c+dx)} \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx = \text{Exception raised: TypeError}$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c)),x, algorith="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen &`

3.616.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \sqrt{\cot(c+dx)} \sqrt{a+b \tan(c+dx)} (A+B \tan(c+dx)) dx \\ &= \int \sqrt{\cot(c+dx)} (A+B \tan(c+dx)) \sqrt{a+b \tan(c+dx)} dx \end{aligned}$$

input `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2),x)`

output `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2), x)`

3.617
$$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.617.1 Optimal result 5899
 3.617.2 Mathematica [A] (verified) 5900
 3.617.3 Rubi [A] (verified) 5900
 3.617.4 Maple [B] (warning: unable to verify) 5903
 3.617.5 Fricas [B] (verification not implemented) 5904
 3.617.6 Sympy [F] 5904
 3.617.7 Maxima [F] 5904
 3.617.8 Giac [F(-2)] 5905
 3.617.9 Mupad [F(-1)] 5905

3.617.1 Optimal result

Integrand size = 35, antiderivative size = 261

$$\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

$$= \frac{\sqrt{ia-b}(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d}$$

$$+ \frac{(2Ab+aB) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{\sqrt{bd}}$$

$$- \frac{\sqrt{ia+b}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d}$$

$$+ \frac{B\sqrt{a+b \tan(c+dx)}}{d\sqrt{\cot(c+dx)}}$$

output

```
(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a-b)^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(2*A*b+B*a)*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/b^(1/2)-(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*(I*a+b)^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+B*(a+b*tan(d*x+c))^(1/2)/d/cot(d*x+c)^(1/2)
```

3.617.2 Mathematica [A] (verified)

Time = 3.77 (sec) , antiderivative size = 293, normalized size of antiderivative = 1.12

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(\sqrt[4]{-1} \sqrt{-a + ib} (A - iB) \arctan \left(\frac{\sqrt[4]{-1} \sqrt{-a + ib} \sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}} \right) \right) \sqrt{a + b \tan(c + dx)}}{\dots}$$

input `Integrate[(Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-1)^(1/4)*Sqrt[-a + I*b]*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]*Sqrt[a + b*Tan[c + d*x]] + (-1)^(1/4)*Sqrt[a + I*b]*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]*Sqrt[a + b*Tan[c + d*x]] + B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x]) + ((2*A*b + a*B)*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*(a + b*Tan[c + d*x]))/(Sqrt[a]*Sqrt[b]*Sqrt[1 + (b*Tan[c + d*x])/a]))/(d*Sqrt[a + b*Tan[c + d*x]])`

3.617.3 Rubi [A] (verified)

Time = 1.16 (sec) , antiderivative size = 220, normalized size of antiderivative = 0.84, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4729, 3042, 4093, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

↓ 3042

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

↓ 4729

3.617. $\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$

$$\begin{aligned} & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx \\ & \quad \downarrow \text{3042} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx \\ & \quad \downarrow \text{4093} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{-((2Ab+aB)\tan^2(c+dx)) - 2(aA-bB)\tan(c+dx) + aB}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{B\sqrt{\tan(c+dx)}}{d} \right) \\ & \quad \downarrow \text{27} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{1}{2} \int \frac{-((2Ab+aB)\tan^2(c+dx)) - 2(aA-bB)\tan(c+dx) + aB}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \right) \\ & \quad \downarrow \text{3042} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{1}{2} \int \frac{-((2Ab+aB)\tan(c+dx)^2) - 2(aA-bB)\tan(c+dx) + aB}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \right) \\ & \quad \downarrow \text{4138} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{\int \frac{-((2Ab+aB)\tan^2(c+dx)) - 2(aA-bB)\tan(c+dx) + aB}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}}{2d} \right) \\ & \quad \downarrow \text{2035} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{\int \frac{-((2Ab+aB)\tan^2(c+dx)) - 2(aA-bB)\tan(c+dx) + aB}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}}{d} \right) \\ & \quad \downarrow \text{2257} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{\int \left(\frac{-2Ab-aB}{\sqrt{a+b\tan(c+dx)}} + \frac{2(Ab+aB-(aA-bB)\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} \right) dx}{d} \right) \\ & \quad \downarrow \text{2009} \end{aligned}$$

3.617. $\int \frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{-\sqrt{-b+ia}(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d}\right)$$

input `Int[(Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-((-(Sqrt[I*a - b]*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])] - ((2*A*b + a*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/Sqrt[b] + Sqrt[I*a + b]*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/d) + (B*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d)`

3.617.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2035 `Int[(F_x_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[Fx, x]`

rule 2257 `Int[(P_x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4093 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[B*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^n/(f*(m + n))), x] + Simp
[1/(m + n) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n - 1)*
Simp[a*A*c*(m + n) - B*(b*c*m + a*d*n) + (A*b*c + a*B*c + a*A*d - b*B*d)*(m
+ n)*Tan[e + f*x] + (A*b*d*(m + n) + B*(a*d*m + b*c*n))*Tan[e + f*x]^2, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[
a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[0, m, 1] && LtQ[0, n, 1]
```

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.617.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.21 (sec) , antiderivative size = 2180708, normalized size of antiderivative = 8355.20

output too large to display

```
input int((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x)
```

```
output result too large to display
```

$$3.617. \quad \int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.617.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8069 vs. 2(209) = 418.

Time = 3.38 (sec) , antiderivative size = 16171, normalized size of antiderivative = 61.96

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algo
rithm="fricas")`

output Too large to include

3.617.6 Sympy [F]

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \int \frac{(A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)}}{\sqrt{\cot(c + dx)}} dx$$

input `integrate((a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))/sqrt(cot(c + d*x)),
x)`

3.617.7 Maxima [F]

$$\begin{aligned} & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\ &= \int \frac{(B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a}}{\sqrt{\cot(dx + c)}} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)/sqrt(cot(d*x + c))
, x)`

3.617. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

3.617.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorith="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen &`

3.617.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\ &= \int \frac{(A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)}}{\sqrt{\cot(c + dx)}} dx \end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2))/cot(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2))/cot(c + d*x)^(1/2),x)`

$$3.618 \quad \int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$$

3.618.1 Optimal result	5906
3.618.2 Mathematica [A] (verified)	5907
3.618.3 Rubi [A] (verified)	5907
3.618.4 Maple [B] (warning: unable to verify)	5911
3.618.5 Fricas [B] (verification not implemented)	5912
3.618.6 Sympy [F]	5912
3.618.7 Maxima [F]	5912
3.618.8 Giac [F(-2)]	5913
3.618.9 Mupad [F(-1)]	5913

3.618.1 Optimal result

Integrand size = 35, antiderivative size = 324

$$\begin{aligned} & \int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx \\ &= \frac{\sqrt{ia-b}(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d} \\ &+ \frac{(4aAb - a^2B - 8b^2B) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{4b^{3/2}d} \\ &+ \frac{\sqrt{ia+b}(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d} \\ &+ \frac{(4Ab - aB) \sqrt{a+b \tan(c+dx)}}{4bd \sqrt{\cot(c+dx)}} + \frac{B(a+b \tan(c+dx))^{3/2}}{2bd \sqrt{\cot(c+dx)}} \end{aligned}$$

output $1/4*(4*A*a*b-B*a^2-8*B*b^2)*\operatorname{arctanh}(b^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/b^{(3/2)}/d+(I*A-B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})*(I*a-b)^{(1/2)}*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/d+(I*A+B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})*(I*a+b)^{(1/2)}*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/d+1/4*(4*A*b-B*a)*(a+b*\tan(d*x+c))^{(1/2)}/b/d/\cot(d*x+c)^{(1/2)}+1/2*B*(a+b*\tan(d*x+c))^{(3/2)}/b/d/\cot(d*x+c)^{(1/2)}$

$$3.618. \quad \int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$$

3.618.2 Mathematica [A] (verified)

Time = 5.20 (sec) , antiderivative size = 356, normalized size of antiderivative = 1.10

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{\sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}\left(-\left((-4aAb + a^2B + 8b^2B) \operatorname{arcsinh}\left(\frac{\sqrt{b}\sqrt{\tan(c + dx)}}{\sqrt{a}}\right)\right)(a + b \tan(c + dx))\right) + \sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)}$$

input `Integrate[(Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(3/2),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-((-4*a*A*b + a^2*B + 8*b^2*B)*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*(a + b*Tan[c + d*x])) + Sqrt[a]*Sqrt[b]*Sqrt[1 + (b*Tan[c + d*x])/a]*(-4*(-1)^(1/4)*Sqrt[-a + I*b]*b*(I*A + B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])*Sqrt[a + b*Tan[c + d*x]] + 4*(-1)^(3/4)*Sqrt[a + I*b]*b*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])*Sqrt[a + b*Tan[c + d*x]] + Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x]))*(4*A*b + a*B + 2*b*B*Tan[c + d*x])))/(4*Sqrt[a]*b^(3/2)*d*Sqrt[a + b*Tan[c + d*x]]*Sqrt[1 + (b*Tan[c + d*x])/a])`

3.618.3 Rubi [A] (verified)Time = 1.59 (sec) , antiderivative size = 286, normalized size of antiderivative = 0.88, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 4729, 3042, 4090, 27, 3042, 4130, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\cot(c + dx)^{3/2}} dx$$

$$\downarrow \text{4729}$$

3.618. $\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$

$$\begin{aligned}
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \tan(c+dx)^{3/2}\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))dx \\
& \quad \downarrow \text{4090} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int -\frac{\sqrt{a+b\tan(c+dx)}(-((4Ab-aB)\tan^2(c+dx))+4bB\tan(c+dx)+aB)}{2\sqrt{\tan(c+dx)}}dx}{2b} + \frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))}{2bc} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2bd} - \frac{\int \frac{\sqrt{a+b\tan(c+dx)}(-((4Ab-aB)\tan^2(c+dx))+4bB\tan(c+dx)+aB)}{\sqrt{\tan(c+dx)}}}{4b} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2bd} - \frac{\int \frac{\sqrt{a+b\tan(c+dx)}(-((4Ab-aB)\tan(c+dx)^2)+4bB\tan(c+dx)+aB)}{\sqrt{\tan(c+dx)}}}{4b} \right) \\
& \quad \downarrow \text{4130} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2bd} - \frac{\int \frac{-((-Ba^2+4Aba-8b^2B)\tan^2(c+dx))+8b(Ab+aB)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}}{2b} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2bd} - \frac{\frac{1}{2} \int \frac{-((-Ba^2+4Aba-8b^2B)\tan^2(c+dx))+8b(Ab+aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}}{2b} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2bd} - \frac{\frac{1}{2} \int \frac{-((-Ba^2+4Aba-8b^2B)\tan(c+dx)^2)+8b(Ab+aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}}{2b} \right) \\
& \quad \downarrow \text{4138}
\end{aligned}$$

3.618. $\int \frac{\sqrt{a+b\tan(c+dx)}(A+B\tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2bd} - \frac{\int \frac{-((-Ba^2+4Aba-8b^2B)\tan^2(c+dx))+8b(Ab+aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} dx}{2d}\right)$$

↓ 2035

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2bd} - \frac{\int \frac{-((-Ba^2+4Aba-8b^2B)\tan^2(c+dx))+8b(Ab+aB)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} dx}{d}\right)$$

↓ 2257

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2bd} - \frac{\int \left(\frac{Ba^2-4Aba+8b^2B}{\sqrt{a+b\tan(c+dx)}} + \frac{8(b(aA-bB)+b(Ab+aB)\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}\right) dx}{d}\right)$$

↓ 2009

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2bd} - \frac{(4Ab-aB)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} + \frac{(a^2+4Ab^2)}{4b}\right)$$

input `Int[(Sqrt[a + b*Tan[c + d*x]]*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(3/2),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2))/(2*b*d) - ((-4*Sqrt[I*a - b]*b*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - ((4*a*A*b - a^2*B - 8*b^2*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[b] - 4*b*Sqrt[I*a + b]*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/d - ((4*A*b - a*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d)/(4*b))`

3.618.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2035 `Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[Fx, x]`
- rule 2257 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4130 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))
```

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.618.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.36 (sec) , antiderivative size = 2181027, normalized size of antiderivative = 6731.56

output too large to display

```
input int((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x)
```

```
output result too large to display
```

$$3.618. \quad \int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$$

3.618.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8173 vs. $2(260) = 520$.

Time = 3.67 (sec) , antiderivative size = 16383, normalized size of antiderivative = 50.56

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algo
rithm="fricas")`

output Too large to include

3.618.6 Sympy [F]

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)}}{\cot^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(a + b*tan(c + d*x))/cot(c + d*x)**(3/2)
, x)`

3.618.7 Maxima [F]

$$\begin{aligned} & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(B \tan(dx + c) + A) \sqrt{b \tan(dx + c) + a}}{\cot(dx + c)^{\frac{3}{2}}} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(b*tan(d*x + c) + a)/cot(d*x + c)^(3/2)
, x)`

3.618. $\int \frac{\sqrt{a+b \tan(c+dx)}(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

3.618.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*tan(d*x+c))^(1/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen &`

3.618.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{\sqrt{a + b \tan(c + dx)}(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(A + B \tan(c + dx)) \sqrt{a + b \tan(c + dx)}}{\cot(c + dx)^{3/2}} dx \end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2))/cot(c + d*x)^(3/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(1/2))/cot(c + d*x)^(3/2),x)`

3.619 $\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.619.1 Optimal result	5914
3.619.2 Mathematica [A] (verified)	5916
3.619.3 Rubi [A] (verified)	5917
3.619.4 Maple [B] (warning: unable to verify)	5926
3.619.5 Fricas [B] (verification not implemented)	5926
3.619.6 Sympy [F(-1)]	5927
3.619.7 Maxima [F]	5927
3.619.8 Giac [F(-1)]	5927
3.619.9 Mupad [F(-1)]	5928

3.619.1 Optimal result

Integrand size = 35, antiderivative size = 422

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \frac{(ia - b)^{3/2}(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d} - \frac{(ia + b)^{3/2}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d} - \frac{2(315a^4A - 63a^2Ab^2 + 8Ab^4 - 420a^3bB - 18ab^3B) \sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}}{315a^3d} + \frac{2(126a^2Ab + 4Ab^3 + 105a^3B - 9ab^2B) \cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{315a^2d} + \frac{2(21a^2A - Ab^2 - 24abB) \cot^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{105ad} - \frac{2(10Ab + 9aB) \cot^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{63d} - \frac{2aA \cot^{\frac{9}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{9d}$$

output $(I*a-b)^{(3/2)}*(I*A-B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)/d}-(I*a+b)^{(3/2)}*(I*A+B)*\arctan(h((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)/d}+2/315*(126*A*a^2*b+4*A*b^3+105*B*a^3-9*B*a*b^2)*\cot(d*x+c)^{(3/2)}*(a+b*\tan(d*x+c))^{(1/2)/a^2/d}+2/105*(21*A*a^2-A*b^2-24*B*a*b)*\cot(d*x+c)^{(5/2)}*(a+b*\tan(d*x+c))^{(1/2)/a/d}-2/63*(10*A*b+9*B*a)*\cot(d*x+c)^{(7/2)}*(a+b*\tan(d*x+c))^{(1/2)/d}-2/9*a*A*\cot(d*x+c)^{(9/2)}*(a+b*\tan(d*x+c))^{(1/2)/d}-2/315*(315*A*a^4-63*A*a^2*b^2+8*A*b^4-420*B*a^3*b-18*B*a*b^3)*\cot(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(1/2)/a^3/d}$

3.619.2 Mathematica [A] (verified)

Time = 6.73 (sec) , antiderivative size = 495, normalized size of antiderivative = 1.17

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A$$

$$+ B \tan(c + dx)) dx = \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} - \frac{bB \sqrt{a + b \tan(c + dx)}}{4d \tan^{\frac{9}{2}}(c + dx)}$$

$$3.619. \quad \int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$$

input `Integrate[Cot[c + d*x]^(11/2)*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-1/4*(b*B*Sqrt[a + b*Tan[c + d*x]])/(d*Tan[c + d*x]^(9/2)) + (-1/9*((8*a*A - 9*b*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Tan[c + d*x]^(9/2)) + (2*((-4*a*(10*A*b + 9*a*B)*Sqrt[a + b*Tan[c + d*x]])/(7*d*Tan[c + d*x]^(7/2)) - (2*((-6*a*(21*a^2*A - A*b^2 - 24*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) - (2*((a*(126*a^2*A*b + 4*A*b^3 + 105*a^3*B - 9*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Tan[c + d*x]^(3/2)) - (2*((-945*a^4*((-1)^(1/4)*(-a + I*b)^(3/2)*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]) - (-1)^(1/4)*(a + I*b)^(3/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]))/(4*d) + (3*a*(315*a^4*A - 63*a^2*A*b^2 + 8*A*b^4 - 420*a^3*b*B - 18*a*b^3*B)*Sqrt[a + b*Tan[c + d*x]])/(2*d*Sqrt[Tan[c + d*x]])))/(3*a)))/(5*a)))/(7*a)))/(9*a))/4`

3.619.3 Rubi [A] (verified)

Time = 3.20 (sec) , antiderivative size = 451, normalized size of antiderivative = 1.07, number of steps used = 25, number of rules used = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.686$, Rules used = {3042, 4729, 3042, 4088, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\int \cot(c + dx)^{11/2}(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$$

$$\downarrow 4729$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx$$

$$\downarrow 3042$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\tan(c + dx)^{11/2}} dx$$

$$\downarrow 4088$$

3.619. $\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2}{9}\int\frac{-b(8aA-9bB)\tan^2(c+dx)-9(Aa^2-2bBa-Ab^2)\tan(c+dx)+a(10Ab+2a^2)}{2\tan^{\frac{9}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}dx\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{9}\int\frac{-b(8aA-9bB)\tan^2(c+dx)-9(Aa^2-2bBa-Ab^2)\tan(c+dx)+a(10Ab+2a^2)}{\tan^{\frac{9}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}dx\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{9}\int\frac{-b(8aA-9bB)\tan(c+dx)^2-9(Aa^2-2bBa-Ab^2)\tan(c+dx)+a(10Ab+2a^2)}{\tan(c+dx)^{9/2}\sqrt{a+b\tan(c+dx)}}dx\right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{9}\left(-\frac{2\int\frac{3(2ab(10Ab+9aB)\tan^2(c+dx)+21a(Ba^2+2Aba-b^2B)\tan(c+dx)+a(21Aa^2-24bBa-Ab^2))}{2\tan^{\frac{7}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}dx}{7a}\right)\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{9}\left(-\frac{3\int\frac{2ab(10Ab+9aB)\tan^2(c+dx)+21a(Ba^2+2Aba-b^2B)\tan(c+dx)+a(21Aa^2-24bBa-Ab^2)}{\tan^{\frac{7}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}dx}{7a}\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{9}\left(-\frac{3\int\frac{2ab(10Ab+9aB)\tan(c+dx)^2+21a(Ba^2+2Aba-b^2B)\tan(c+dx)+a(21Aa^2-24bBa-Ab^2)}{\tan(c+dx)^{7/2}\sqrt{a+b\tan(c+dx)}}dx}{7a}\right)\right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{9}\left(\left(3\left(-\frac{2\int\frac{-105(Aa^2-2bBa-Ab^2)\tan(c+dx)a^2-4b(21Aa^2-24bBa-Ab^2)\tan^2(c+dx)a+(105Ba^3+126Ab^2a)}{2\tan^{\frac{5}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}dx\right)}{5a}\right)\right)\right)$$

↓ 27

3.619. $\int \cot^{\frac{11}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{9} \left(3 \left(\frac{\int \frac{-105(Aa^2-2bBa-Ab^2)\tan(c+dx)a^2-4b(21Aa^2-24bBa-Ab^2)\tan^2(c+dx)a+(105Ba^3+126Aba^2-9b^2a)}{\tan^{\frac{5}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}}{5a} \right) \right) \right) - 7a$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{9} \left(3 \left(\frac{\int \frac{-105(Aa^2-2bBa-Ab^2)\tan(c+dx)a^2-4b(21Aa^2-24bBa-Ab^2)\tan(c+dx)^2a+(105Ba^3+126Aba^2-9b^2a)}{\tan(c+dx)^{5/2}\sqrt{a+b\tan(c+dx)}}}{5a} \right) \right) \right) - 7a$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{9} \left(3 \left(\frac{2 \int \frac{315(Ba^2+2Aba-b^2B)\tan(c+dx)a^3+2b(105Ba^3+126Aba^2-9b^2Ba+4Ab^3)\tan^2(c+dx)a+(315Aa^4-4b^2a^2)}{2 \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}}{3a} \right) \right) \right) - 5a$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{9} \left(3 \left(\frac{\int \frac{315(Ba^2+2Aba-b^2B)\tan(c+dx)a^3+2b(105Ba^3+126Aba^2-9b^2Ba+4Ab^3)\tan^2(c+dx)a+(315Aa^4-4b^2a^2)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}}{3a} \right) \right) \right) - 5a$$

↓ 3042

3.619. $\int \cot^{\frac{11}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{9} \left(3 \left(\int \frac{315(Ba^2+2Aba-b^2B)\tan(c+dx)a^3+2b(105Ba^3+126Aba^2-9b^2Ba+4Ab^3)\tan(c+dx)^2a+(315Aa^4-4Ab^4)}{\tan(c+dx)^{3/2}\sqrt{a+b\tan(c+dx)}} dx - \frac{5a}{5a} \right) \right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{9} \left(3 \left(\int \frac{2 \int \frac{315(a^4(Ba^2+2Aba-b^2B)-a^4(Aa^2-2bBa-Ab^2))\tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{2(315a^4A-420a^3bB-63a^2Ab^2-18a^2b^3)}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{5a}{5a} \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{9} \left(3 \left(\int \frac{315 \int \frac{a^4(Ba^2+2Aba-b^2B)-a^4(Aa^2-2bBa-Ab^2))\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{2(315a^4A-420a^3bB-63a^2Ab^2-18a^2b^3)}{d\sqrt{\tan(c+dx)}}}{3a} - \frac{5a}{5a} \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{9} \left(3 \left(\frac{315 \int \frac{a^4(Ba^2+2Aba-b^2B)-a^4(Aa^2-2bBa-Ab^2)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{2(315a^4A-420a^3bB-63a^2Ab^2-18a^2B^2)}{3a} - \frac{d\sqrt{\tan(c+dx)}}{5a} \right) \right) \right)$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} \left(-\frac{2(9aB+10Ab)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} - \frac{2(21a^2)}{3} \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} \left(-\frac{2(9aB+10Ab)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} - \frac{2(21a^2)}{3} \right) \right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} - \frac{2(9aB+10Ab)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} - \frac{2(21a^2)}{3} \right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} - \frac{2(9aB+10Ab)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} - \frac{2(21a^2)}{3} \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} - \frac{2(9aB+10Ab)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} - \frac{2(21a^2)}{3} \right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} - \frac{2(9aB+10Ab)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} - \frac{2(21a^2)}{3} \right)$$

input `Int[Cot[c + d*x]^(11/2)*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*a*A*Sqrt[a + b*Tan[c + d*x]])/(
9*d*Tan[c + d*x]^(9/2)) + ((-2*(10*A*b + 9*a*B)*Sqrt[a + b*Tan[c + d*x]])/
(7*d*Tan[c + d*x]^(7/2)) - (3*((-2*(21*a^2*A - A*b^2 - 24*a*b*B)*Sqrt[a +
b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) + ((-2*(126*a^2*A*b + 4*A*b^3 +
105*a^3*B - 9*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2))
- ((315*(-((a^4*(a + I*b)^2*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d
*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a - b]*d)) + (a^4*(a - I*b)^2*(I*
A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]
]/(Sqrt[I*a + b]*d)))/a - (2*(315*a^4*A - 63*a^2*A*b^2 + 8*A*b^4 - 420*a^
3*b*B - 18*a*b^3*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]])))/(3*a
))/(5*a))/(7*a))/9)
```

3.619.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 104 Int[(((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)))/((e_.) + (f_.)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.619.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.38 (sec) , antiderivative size = 2403002, normalized size of antiderivative = 5694.32

output too large to display

```
input int(cot(d*x+c)^(11/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x)
```

```
output result too large to display
```

3.619.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12342 vs. $2(356) = 712$.

Time = 2.36 (sec) , antiderivative size = 12342, normalized size of antiderivative = 29.25

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^(11/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, alg
orithm="fricas")
```

```
output Too large to include
```

3.619.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(11/2)*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.619.7 Maxima [F]

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{3}{2}} \cot(dx + c)^{\frac{11}{2}} dx$$

input `integrate(cot(d*x+c)^(11/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^(11/2), x)`

3.619.8 Giac [F(-1)]

Timed out.

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(11/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.619.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{11/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2} dx$$

input `int(cot(c + d*x)^(11/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2),x)`

output `int(cot(c + d*x)^(11/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2), x)`

3.620 $\int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.620.1 Optimal result	5929
3.620.2 Mathematica [A] (verified)	5930
3.620.3 Rubi [A] (verified)	5931
3.620.4 Maple [B] (warning: unable to verify)	5937
3.620.5 Fricas [B] (verification not implemented)	5937
3.620.6 Sympy [F(-1)]	5938
3.620.7 Maxima [F]	5938
3.620.8 Giac [F(-1)]	5938
3.620.9 Mupad [F(-1)]	5939

3.620.1 Optimal result

Integrand size = 35, antiderivative size = 351

$$\begin{aligned}
 & \int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \\
 & \frac{(ia-b)^{3/2}(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{d} \\
 & - \frac{(ia+b)^{3/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{d} \\
 & + \frac{2(140a^2Ab+6Ab^3+105a^3B-21ab^2B) \sqrt{\cot(c+dx)}\sqrt{a+b \tan(c+dx)}}{105a^2d} \\
 & + \frac{2(35a^2A-3Ab^2-42abB) \cot^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}}{105ad} \\
 & - \frac{2(8Ab+7aB) \cot^{\frac{5}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}}{35d} \\
 & - \frac{2aA \cot^{\frac{7}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}}{7d}
 \end{aligned}$$

output $-(I*a-b)^{(3/2)}*(A+I*B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)/d}-(I*a+b)^{(3/2)}*(A-I*B)*\arctan((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)/d}+2/105*(35*A*a^2-3*A*b^2-42*B*a*b)*\cot(d*x+c)^{(3/2)}*(a+b*\tan(d*x+c))^{(1/2)/a}+d-2/35*(8*A*b+7*B*a)*\cot(d*x+c)^{(5/2)}*(a+b*\tan(d*x+c))^{(1/2)/d}-2/7*a*A*\cot(d*x+c)^{(7/2)}*(a+b*\tan(d*x+c))^{(1/2)/d}+2/105*(140*A*a^2*b+6*A*b^3+105*B*a^3-21*B*a*b^2)*\cot(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(1/2)/a^2/d}$

3.620.2 Mathematica [A] (verified)

Time = 5.53 (sec) , antiderivative size = 346, normalized size of antiderivative = 0.99

$$\int \cot^{\frac{9}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx =$$

$$\cot^{\frac{7}{2}}(c+dx) \left(35a^3bB\sqrt{a+b\tan(c+dx)} + 5a^3(6aA-7bB)\sqrt{a+b\tan(c+dx)} + a\tan(c+dx) \left(105(-\right.$$

input `Integrate[Cot[c + d*x]^(9/2)*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output $-1/105*(\text{Cot}[c + d*x]^{(7/2)}*(35*a^3*b*B*\text{Sqrt}[a + b*\text{Tan}[c + d*x]] + 5*a^3*(6*a*A - 7*b*B)*\text{Sqrt}[a + b*\text{Tan}[c + d*x]] + a*\text{Tan}[c + d*x]*(105*(-1)^{(3/4)}*a^2*((-a + I*b)^{(3/2)}*(A - I*B)*\text{ArcTan}[((-1)^{(1/4)}*\text{Sqrt}[-a + I*b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a + b*\text{Tan}[c + d*x]]] + (a + I*b)^{(3/2)}*(A + I*B)*\text{ArcTan}[((-1)^{(1/4)}*\text{Sqrt}[a + I*b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a + b*\text{Tan}[c + d*x]])]*\text{Tan}[c + d*x]^{(5/2)} + 6*a^2*(8*A*b + 7*a*B)*\text{Sqrt}[a + b*\text{Tan}[c + d*x]] - 2*a*(35*a^2*A - 3*A*b^2 - 42*a*b*B)*\text{Tan}[c + d*x]*\text{Sqrt}[a + b*\text{Tan}[c + d*x]] - 2*(140*a^2*A*b + 6*A*b^3 + 105*a^3*B - 21*a*b^2*B)*\text{Tan}[c + d*x]^2*\text{Sqrt}[a + b*\text{Tan}[c + d*x]])))/(a^3*d)$

3.620.3 Rubi [A] (verified)

Time = 2.55 (sec) , antiderivative size = 373, normalized size of antiderivative = 1.06, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 4729, 3042, 4088, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{9/2}(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan(c+dx)^{9/2}} dx$$

$$\downarrow \text{4088}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2}{7} \int \frac{-b(6aA-7bB) \tan^2(c+dx) - 7(Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(8Ab+7A^2)}{2 \tan^{\frac{7}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx \right)$$

$$\downarrow \text{27}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \int \frac{-b(6aA-7bB) \tan^2(c+dx) - 7(Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(8Ab+7A^2)}{\tan^{\frac{7}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx \right)$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \int \frac{-b(6aA-7bB) \tan(c+dx)^2 - 7(Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(8Ab+7A^2)}{\tan(c+dx)^{7/2} \sqrt{a+b \tan(c+dx)}} dx \right)$$

$$\downarrow \text{4132}$$

3.620. $\int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(-\frac{2\int\frac{4ab(8Ab+7aB)\tan^2(c+dx)+35a(Ba^2+2Aba-b^2B)\tan(c+dx)+a(35Aa^2-42bBa-3Ab^2)}{2\tan^{\frac{5}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}dx}{5a}\right)\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(-\frac{\int\frac{4ab(8Ab+7aB)\tan^2(c+dx)+35a(Ba^2+2Aba-b^2B)\tan(c+dx)+a(35Aa^2-42bBa-3Ab^2)}{\tan^{\frac{5}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}dx}{5a}\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(-\frac{\int\frac{4ab(8Ab+7aB)\tan(c+dx)^2+35a(Ba^2+2Aba-b^2B)\tan(c+dx)+a(35Aa^2-42bBa-3Ab^2)}{\tan(c+dx)^{5/2}\sqrt{a+b\tan(c+dx)}}dx}{5a}\right)\right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(-\frac{2\int\frac{-105(Aa^2-2bBa-Ab^2)\tan(c+dx)a^2-2b(35Aa^2-42bBa-3Ab^2)\tan^2(c+dx)a+(105Ba^3+140Aba^2-21b^2B)}{2\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}dx}{3a}\right)\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(-\frac{\int\frac{-105(Aa^2-2bBa-Ab^2)\tan(c+dx)a^2-2b(35Aa^2-42bBa-3Ab^2)\tan^2(c+dx)a+(105Ba^3+140Aba^2-21b^2B)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}dx}{3a}\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(-\frac{\int\frac{-105(Aa^2-2bBa-Ab^2)\tan(c+dx)a^2-2b(35Aa^2-42bBa-3Ab^2)\tan(c+dx)^2a+(105Ba^3+140Aba^2-21b^2B)}{\tan(c+dx)^{3/2}\sqrt{a+b\tan(c+dx)}}dx}{3a}\right)\right)$$

↓ 4132

3.620. $\int \cot^{\frac{9}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(-\frac{2 \int \frac{105 \left((Aa^2 - 2bBa - Ab^2)a^3 + (Ba^2 + 2Aba - b^2B) \tan(c+dx)a^3 \right) dx}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}}}{a} - \frac{2(105a^3B + 140a^2Ab - 21ab^2B + 6Ab^3)}{d\sqrt{\tan(c+dx)}} \right) \right) \frac{1}{3a} \frac{1}{5a}$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(-\frac{105 \int \frac{(Aa^2 - 2bBa - Ab^2)a^3 + (Ba^2 + 2Aba - b^2B) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(105a^3B + 140a^2Ab - 21ab^2B + 6Ab^3)}{d\sqrt{\tan(c+dx)}} \right) \right) \frac{1}{3a} \frac{1}{5a}$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \left(-\frac{105 \int \frac{(Aa^2 - 2bBa - Ab^2)a^3 + (Ba^2 + 2Aba - b^2B) \tan(c+dx)a^3}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(105a^3B + 140a^2Ab - 21ab^2B + 6Ab^3)}{d\sqrt{\tan(c+dx)}} \right) \right) \frac{1}{3a} \frac{1}{5a}$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA\sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(-\frac{2(7aB + 8Ab)\sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A - 42abB)}{d \tan^{\frac{3}{2}}(c+dx)} \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA\sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(-\frac{2(7aB + 8Ab)\sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A - 42abB)}{d \tan^{\frac{3}{2}}(c+dx)} \right) \right)$$

↓ 4098

3.620. $\int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(-\frac{2(7aB+8Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-42a^2B)}{5d^2\tan^{\frac{3}{2}}(c+dx)} \right) \right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(-\frac{2(7aB+8Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-42a^2B)}{5d^2\tan^{\frac{3}{2}}(c+dx)} \right) \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(-\frac{2(7aB+8Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-42a^2B)}{5d^2\tan^{\frac{3}{2}}(c+dx)} \right) \right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(-\frac{2(7aB+8Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)} - \frac{2(35a^2A-42a^2B)}{5d^2\tan^{\frac{3}{2}}(c+dx)} \right) \right)$$

input `Int[Cot[c + d*x]^(9/2)*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*a*A*Sqrt[a + b*Tan[c + d*x]])/(7*d*Tan[c + d*x]^(7/2)) + ((-2*(8*A*b + 7*a*B)*Sqrt[a + b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) - ((-2*(35*a^2*A - 3*A*b^2 - 42*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) + ((-105*((a^3*(a + I*b)^2*(A + I*B)*ArcTan[(Sqrt[I*a - b])*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a - b]*d) + (a^3*(a - I*b)^2*(A - I*B)*ArcTanh[(Sqrt[I*a + b])*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d)))/a - (2*(140*a^2*A*b + 6*A*b^3 + 105*a^3*B - 21*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]])/(3*a)/(5*a)/7)`

3.620.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.620.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.45 (sec) , antiderivative size = 2403427, normalized size of antiderivative = 6847.37

output too large to display

input `int(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x)`

output `result too large to display`

3.620.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12268 vs. $2(291) = 582$.

Time = 2.42 (sec) , antiderivative size = 12268, normalized size of antiderivative = 34.95

$$\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorith="fricas")`

output `Too large to include`

3.620.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(9/2)*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.620.7 Maxima [F]

$$\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{3}{2}} \cot(dx + c)^{\frac{9}{2}} dx$$

input `integrate(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^(9/2), x)`

3.620.8 Giac [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.620. $\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.620.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{9/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2} dx$$

input `int(cot(c + d*x)^(9/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2),x)`output `int(cot(c + d*x)^(9/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2), x)`

3.621 $\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.621.1 Optimal result	5940
3.621.2 Mathematica [A] (verified)	5941
3.621.3 Rubi [A] (verified)	5941
3.621.4 Maple [B] (warning: unable to verify)	5947
3.621.5 Fricas [B] (verification not implemented)	5947
3.621.6 Sympy [F(-1)]	5947
3.621.7 Maxima [F]	5948
3.621.8 Giac [F(-1)]	5948
3.621.9 Mupad [F(-1)]	5948

3.621.1 Optimal result

Integrand size = 35, antiderivative size = 299

$$\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{(a+ib)^2(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{\sqrt{ia-bd}} + \frac{(ia+b)^{3/2}(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d} + \frac{2(15a^2A-3Ab^2-20abB) \sqrt{\cot(c+dx)} \sqrt{a+b \tan(c+dx)}}{15ad} - \frac{2(6Ab+5aB) \cot^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}{15d} - \frac{2aA \cot^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}{5d}$$

output $(I*a+b)^{(3/2)}*(I*A+B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/d+(a+I*b)^2*(I*A-B)*\operatorname{arctan}((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/d/(I*a-b)^{(1/2)}-2/15*(6*A*b+5*B*a)*\cot(d*x+c)^{(3/2)}*(a+b*\tan(d*x+c))^{(1/2)}/d-2/5*a*A*\cot(d*x+c)^{(5/2)}*(a+b*\tan(d*x+c))^{(1/2)}/d+2/15*(15*A*a^2-3*A*b^2-20*B*a*b)*\cot(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(1/2)}/a/d$

3.621.2 Mathematica [A] (verified)

Time = 3.21 (sec) , antiderivative size = 286, normalized size of antiderivative = 0.96

$$\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx =$$

$$\cot^{\frac{5}{2}}(c+dx) \left(30\sqrt[4]{-1}a \left((-a+ib)^{3/2}(A-iB) \arctan \left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right) - (a+ib)^{3/2}(A+iB) \arctan \left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right) \right) \right)$$

input `Integrate[Cot[c + d*x]^(7/2)*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `-1/30*(Cot[c + d*x]^(5/2)*(30*(-1)^(1/4)*a*((-a + I*b)^(3/2)*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]) - (a + I*b)^(3/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(5/2) + 15*a*b*B*Sqrt[a + b*Tan[c + d*x]] + 3*a*(4*a*A - 5*b*B)*Sqrt[a + b*Tan[c + d*x]] + 4*a*(6*A*b + 5*a*B)*Tan[c + d*x]*Sqrt[a + b*Tan[c + d*x]] - 4*(15*a^2*A - 3*A*b^2 - 20*a*b*B)*Tan[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]))/(a*d)`

3.621.3 Rubi [A] (verified)

Time = 2.01 (sec) , antiderivative size = 310, normalized size of antiderivative = 1.04, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.514$, Rules used = {3042, 4729, 3042, 4088, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{7/2}(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx$$

3.621. $\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan(c + dx)^{7/2}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan(c + dx)^{7/2}} dx \\
& \quad \downarrow \text{4088} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{2}{5} \int \frac{-b(4aA - 5bB) \tan^2(c + dx) - 5(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(6Ab + 5a^2)}{2 \tan^{5/2}(c + dx) \sqrt{a + b \tan(c + dx)}} dx \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{5} \int \frac{-b(4aA - 5bB) \tan^2(c + dx) - 5(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(6Ab + 5a^2)}{\tan^{5/2}(c + dx) \sqrt{a + b \tan(c + dx)}} dx \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{5} \int \frac{-b(4aA - 5bB) \tan(c + dx)^2 - 5(Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(6Ab + 5a^2)}{\tan(c + dx)^{5/2} \sqrt{a + b \tan(c + dx)}} dx \right) \\
& \quad \downarrow \text{4132} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{5} \left(- \frac{2 \int \frac{2ab(6Ab + 5aB) \tan^2(c + dx) + 15a(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(15Aa^2 - 20bBa - 3Ab^2)}{2 \tan^{3/2}(c + dx) \sqrt{a + b \tan(c + dx)}} dx}{3a} \right) \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{5} \left(- \frac{\int \frac{2ab(6Ab + 5aB) \tan^2(c + dx) + 15a(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(15Aa^2 - 20bBa - 3Ab^2)}{\tan^{3/2}(c + dx) \sqrt{a + b \tan(c + dx)}} dx}{3a} \right) \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{5} \left(- \frac{\int \frac{2ab(6Ab + 5aB) \tan(c + dx)^2 + 15a(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(15Aa^2 - 20bBa - 3Ab^2)}{\tan(c + dx)^{3/2} \sqrt{a + b \tan(c + dx)}} dx}{3a} \right) \right) \\
& \quad \downarrow \text{4132}
\end{aligned}$$

3.621. $\int \cot^{7/2}(c + dx) (a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(-\frac{2\int\frac{15(a^2(Ba^2+2Aba-b^2B)-a^2(Aa^2-2bBa-Ab^2)\tan(c+dx))}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx}{a}-\frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right)\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(-\frac{15\int\frac{a^2(Ba^2+2Aba-b^2B)-a^2(Aa^2-2bBa-Ab^2)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx}{a}-\frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(-\frac{15\int\frac{a^2(Ba^2+2Aba-b^2B)-a^2(Aa^2-2bBa-Ab^2)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx}{a}-\frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right)\right)$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}+\frac{1}{5}\left(-\frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}+\frac{1}{5}\left(-\frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right)\right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}+\frac{1}{5}\left(-\frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{2(15a^2A-20abB-3Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}\right)\right)$$

3.621. $\int \cot^{\frac{7}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}+\frac{1}{5}\left(-\frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{2(15a^2A-20abB)}{3d^2}\right)\right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}+\frac{1}{5}\left(-\frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{2(15a^2A-20abB)}{3d^2}\right)\right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}+\frac{1}{5}\left(-\frac{2(5aB+6Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{2(15a^2A-20abB)}{3d^2}\right)\right)$$

```
input Int[Cot[c + d*x]^(7/2)*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*a*A*Sqrt[a + b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) + ((-2*(6*A*b + 5*a*B)*Sqrt[a + b*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - ((15*((a^2*(a + I*b)^2*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d) + (a^2*(a - I*b)^2*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d)))/a - (2*(15*a^2*A - 3*A*b^2 - 20*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a)/5)
```

3.621. $\int \cot^{\frac{7}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx$

3.621.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && !ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0]))`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.621.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.49 (sec) , antiderivative size = 2401287, normalized size of antiderivative = 8031.06

output too large to display

input `int(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x)`

output `result too large to display`

3.621.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12233 vs. $2(243) = 486$.

Time = 2.37 (sec) , antiderivative size = 12233, normalized size of antiderivative = 40.91

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `Too large to include`

3.621.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(7/2)*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.621.7 Maxima [F]

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{3}{2}} \cot(dx + c)^{\frac{7}{2}} dx$$

input `integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^(7/
2), x)`

3.621.8 Giac [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algo
rithm="giac")`

output `Timed out`

3.621.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{7/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2} dx$$

input `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2),x)`

output `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2), x)`

3.621. $\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.622 $\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.622.1 Optimal result	5949
3.622.2 Mathematica [A] (verified)	5950
3.622.3 Rubi [A] (verified)	5950
3.622.4 Maple [B] (warning: unable to verify)	5955
3.622.5 Fricas [B] (verification not implemented)	5955
3.622.6 Sympy [F(-1)]	5956
3.622.7 Maxima [F]	5956
3.622.8 Giac [F(-1)]	5956
3.622.9 Mupad [F(-1)]	5957

3.622.1 Optimal result

Integrand size = 35, antiderivative size = 236

$$\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx = \frac{(ia-b)^{3/2}(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d} + \frac{(ia+b)^{3/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d} - \frac{2(4Ab+3aB) \sqrt{\cot(c+dx)} \sqrt{a+b \tan(c+dx)}}{3d} - \frac{2aA \cot^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}{3d}$$

output

```
(I*a-b)^(3/2)*(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(I*a+b)^(3/2)*(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-2/3*a*A*cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(1/2)/d-2/3*(4*A*b+3*B*a)*cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/d
```

3.622.2 Mathematica [A] (verified)

Time = 1.11 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.03

$$\int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx = \frac{\sqrt{\cot(c+dx)} \left(3\sqrt[4]{-1} \left((-a+ib)^{3/2}(iA+B) \arctan \left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}} \right) \right) + i(a+B\tan(c+dx)) \right)}{3d}$$

input `Integrate[Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*(3*(-1)^(1/4)*((-a + I*b)^(3/2)*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + I*(a + I*b)^(3/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])*Sqrt[Tan[c + d*x]] - 2*(4*A*b + 3*a*B)*Sqrt[a + b*Tan[c + d*x]] - 3*b*B*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]] + (-2*a*A + 3*b*B)*Cot[c + d*x]*Sqrt[a + b*Tan[c + d*x]])/(3*d)`

3.622.3 Rubi [A] (verified)Time = 1.52 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.04, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 4729, 3042, 4088, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \cot(c+dx)^{5/2}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx \\ & \quad \downarrow \text{4729} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\tan^{\frac{5}{2}}(c+dx)} dx \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.622. $\int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\tan(c+dx)^{5/2}} dx$$

↓ 4088

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2}{3} \int \frac{-b(2aA-3bB) \tan^2(c+dx) - 3(Aa^2-2bBa-Ab^2) \tan(c+dx) + a(4Ab+3a^2)}{2 \tan^{3/2}(c+dx) \sqrt{a+b \tan(c+dx)}} dx \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \int \frac{-b(2aA-3bB) \tan^2(c+dx) - 3(Aa^2-2bBa-Ab^2) \tan(c+dx) + a(4Ab+3a^2)}{\tan^{3/2}(c+dx) \sqrt{a+b \tan(c+dx)}} dx \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \int \frac{-b(2aA-3bB) \tan(c+dx)^2 - 3(Aa^2-2bBa-Ab^2) \tan(c+dx) + a(4Ab+3a^2)}{\tan(c+dx)^{3/2} \sqrt{a+b \tan(c+dx)}} dx \right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \left(-\frac{2 \int \frac{3(a(Aa^2-2bBa-Ab^2)+a(Ba^2+2Aba-b^2B) \tan(c+dx))}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(3aB+4Ab)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \left(-\frac{3 \int \frac{a(Aa^2-2bBa-Ab^2)+a(Ba^2+2Aba-b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(3aB+4Ab)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \left(-\frac{3 \int \frac{a(Aa^2-2bBa-Ab^2)+a(Ba^2+2Aba-b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(3aB+4Ab)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} \right) \right)$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA\sqrt{a+b \tan(c+dx)}}{3d \tan^{3/2}(c+dx)} + \frac{1}{3} \left(-\frac{2(3aB+4Ab)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\tan(c+dx)}} - \frac{3\left(\frac{1}{2}a(a-b)\right)}{\dots} \right) \right)$$

↓ 3042

3.622. $\int \cot^{5/2}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}+\frac{1}{3}\left(-\frac{2(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-\frac{3\left(\frac{1}{2}a(ib-a)\right)}{\dots}\right)\right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}+\frac{1}{3}\left(-\frac{2(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-\frac{3\left(\frac{a(a-ib)^2(A+B\tan(c+dx))}{\dots}\right)}{\dots}\right)\right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}+\frac{1}{3}\left(-\frac{2(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-\frac{3\left(\frac{a(a-ib)^2(A+B\tan(c+dx))}{\dots}\right)}{\dots}\right)\right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}+\frac{1}{3}\left(-\frac{2(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-\frac{3\left(\frac{a(a-ib)^2(A+B\tan(c+dx))}{\dots}\right)}{\dots}\right)\right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}+\frac{1}{3}\left(-\frac{2(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-\frac{3\left(\frac{a(a+ib)^2(A+B\tan(c+dx))}{\dots}\right)}{\dots}\right)\right)$$

input `Int[Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*a*A*Sqrt[a + b*Tan[c + d*x]])/(
3*d*Tan[c + d*x]^(3/2)) + ((-3*((a*(a + I*b)^2*(A + I*B)*ArcTan[(Sqrt[I*a
- b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d) + (a
*(a - I*b)^2*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a +
b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d)))/a - (2*(4*A*b + 3*a*B)*Sqrt[a + b*T
an[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/3
```

3.622.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 104 Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))/((e_) + (f_)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

```
rule 219 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```


rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.622.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.49 (sec) , antiderivative size = 2399199, normalized size of antiderivative = 10166.10

output too large to display

```
input int(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x)
```

```
output result too large to display
```

3.622.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12148 vs. $2(188) = 376$.

Time = 2.43 (sec) , antiderivative size = 12148, normalized size of antiderivative = 51.47

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algo
rithm="fricas")
```

```
output Too large to include
```

3.622.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(5/2)*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.622.7 Maxima [F]

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{3}{2}} \cot(dx + c)^{\frac{5}{2}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^(5/2), x)`

3.622.8 Giac [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.622.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{5/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2} dx$$

input `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2),x)`output `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2), x)`

3.623 $\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx$

3.623.1 Optimal result	5958
3.623.2 Mathematica [A] (verified)	5959
3.623.3 Rubi [A] (verified)	5959
3.623.4 Maple [B] (warning: unable to verify)	5962
3.623.5 Fricas [B] (verification not implemented)	5963
3.623.6 Sympy [F(-1)]	5963
3.623.7 Maxima [F(-1)]	5963
3.623.8 Giac [F(-1)]	5964
3.623.9 Mupad [F(-1)]	5964

3.623.1 Optimal result

Integrand size = 35, antiderivative size = 269

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx)) dx =$$

$$\frac{(a+ib)^2(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{\sqrt{ia-bd}}$$

$$+ \frac{2b^{3/2} \operatorname{Barctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d}$$

$$- \frac{(ia+b)^{3/2}(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d}$$

$$- \frac{2aA \sqrt{\cot(c+dx)} \sqrt{a+b \tan(c+dx)}}{d}$$

```
output 2*b^(3/2)*B*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d
*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-(I*a+b)^(3/2)*(I*A+B)*arctanh((I*a+b)^(1/2)
*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2
)/d-(a+I*b)^2*(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c
))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*a-b)^(1/2)-2*a*A*cot(d*x+
c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/d
```

3.623.2 Mathematica [A] (verified)

Time = 2.63 (sec) , antiderivative size = 266, normalized size of antiderivative = 0.99

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx = \frac{\sqrt{\cot(c+dx)} \left(\sqrt[4]{-1}(-a+ib)^{3/2}(A-iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) \right) \sqrt{\tan(c+dx)}}{\sqrt{\cot(c+dx)}}$$

input `Integrate[Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*((-1)^(1/4)*(-a + I*b)^(3/2)*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]*Sqrt[Tan[c + d*x]] - (-1)^(1/4)*(a + I*b)^(3/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]*Sqrt[Tan[c + d*x]] - 2*a*A*Sqrt[a + b*Tan[c + d*x]] + (2*Sqrt[a]*b^(3/2)*B*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[Tan[c + d*x]]*Sqrt[1 + (b*Tan[c + d*x])/a])/Sqrt[a + b*Tan[c + d*x]])/d`

3.623.3 Rubi [A] (verified)Time = 1.22 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.82, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4729, 3042, 4088, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{3/2}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

3.623. $\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx$

$$\begin{aligned}
& \int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\tan(c + dx)^{3/2}} dx \\
& \quad \downarrow \text{4088} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(2 \int \frac{b^2 B \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)}{2\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx - \frac{2aA}{d} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\int \frac{b^2 B \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx - \frac{2aA}{d} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\int \frac{b^2 B \tan(c + dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx - \frac{2aA}{d} \right) \\
& \quad \downarrow \text{4138} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{\int \frac{b^2 B \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} d \tan(c + dx)}{d} - \frac{2aA \sqrt{a + b \tan(c + dx)}}{d \sqrt{\tan(c + dx)}} \right) \\
& \quad \downarrow \text{2035} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{2 \int \frac{b^2 B \tan^2(c + dx) - (Aa^2 - 2bBa - Ab^2) \tan(c + dx) + a(2Ab + aB)}{\sqrt{a + b \tan(c + dx)} (\tan^2(c + dx) + 1)} d \sqrt{\tan(c + dx)}}{d} - \frac{2aA \sqrt{a + b \tan(c + dx)}}{d \sqrt{\tan(c + dx)}} \right) \\
& \quad \downarrow \text{2257} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{2 \int \left(\frac{Bb^2}{\sqrt{a + b \tan(c + dx)}} + \frac{Ba^2 + 2Aba - b^2 B - (Aa^2 - 2bBa - Ab^2) \tan(c + dx)}{\sqrt{a + b \tan(c + dx)} (\tan^2(c + dx) + 1)} \right) d \sqrt{\tan(c + dx)}}{d} - \frac{2aA \sqrt{a + b \tan(c + dx)}}{d \sqrt{\tan(c + dx)}} \right) \\
& \quad \downarrow \text{2009} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(-\frac{2aA \sqrt{a + b \tan(c + dx)}}{d \sqrt{\tan(c + dx)}} + \frac{2 \left(\frac{1}{2} (-b + ia)^{3/2} (-B + iA) \arctan \left(\frac{\sqrt{-b + ia} \sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}} \right) \right)}{d} \right)
\end{aligned}$$

3.623. $\int \cot^{\frac{3}{2}}(c + dx) (a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx$

input `Int[Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((2*(((I*a - b)^(3/2)*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/2 + b^(3/2)*B*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) - ((I*a + b)^(3/2)*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/2))/d - (2*a*A*Sqrt[a + b*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))`

3.623.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2035 `Int[(F_x_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[F_x, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[F_x, x]`

rule 2257 `Int[(P_x)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[P_x*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[P_x, x] && IntegerQ[p]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`


```
rule 4088 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x
])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2))
Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*
(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n
+ 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[
e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n
+ 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] &
& LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])
```

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)])^2, x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.623.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.41 (sec) , antiderivative size = 2397600, normalized size of antiderivative = 8913.01

output too large to display

```
input int(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x)
```

```
output result too large to display
```

$$3.623. \quad \int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$$

3.623.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12204 vs. $2(215) = 430$.

Time = 4.70 (sec) , antiderivative size = 24440, normalized size of antiderivative = 90.86

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algo
rithm="fricas")`

output Too large to include

3.623.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(3/2)*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output Timed out

3.623.7 Maxima [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algo
rithm="maxima")`

output Timed out

3.623.8 Giac [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.623.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{3/2}(A + B \tan(c + dx))(a + b \tan(c + dx))^{3/2} dx$$

input `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2),x)`

output `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2), x)`

3.624 $\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$

3.624.1 Optimal result	5965
3.624.2 Mathematica [A] (verified)	5966
3.624.3 Rubi [A] (verified)	5966
3.624.4 Maple [B] (warning: unable to verify)	5969
3.624.5 Fricas [B] (verification not implemented)	5970
3.624.6 Sympy [F]	5970
3.624.7 Maxima [F]	5970
3.624.8 Giac [F(-1)]	5971
3.624.9 Mupad [F(-1)]	5971

3.624.1 Optimal result

Integrand size = 35, antiderivative size = 264

$$\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx =$$

$$\frac{(ia - b)^{3/2}(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{\sqrt{b}(2Ab + 3aB) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d}$$

$$- \frac{(ia + b)^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{bB \sqrt{a + b \tan(c + dx)}}{d \sqrt{\cot(c + dx)}}$$

```
output -(I*a-b)^(3/2)*(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-(I*a+b)^(3/2)*(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(2*A*b+3*B*a)*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*b^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+b*B*(a+b*tan(d*x+c))^(1/2)/d/cot(d*x+c)^(1/2)
```

3.624.2 Mathematica [A] (verified)

Time = 1.41 (sec) , antiderivative size = 263, normalized size of antiderivative = 1.00

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx = \frac{\sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}\left(-\sqrt[4]{-1}(-a+ib)^{3/2}(iA+B)\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)+B\tan(c+dx)\right)}{dx}$$

input `Integrate[Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-((-1)^(1/4)*(-a + I*b)^(3/2)*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) - (-1)^(3/4)*(a + I*b)^(3/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) + b*B*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]] + (Sqrt[a]*Sqrt[b]*(2*A*b + 3*a*B)*ArcSin[h[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/Sqrt[a + b*Tan[c + d*x]]))/d`

3.624.3 Rubi [A] (verified)Time = 1.29 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.84, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4729, 3042, 4090, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx \\ & \quad \downarrow \text{4729} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\sqrt{\tan(c+dx)}}dx \end{aligned}$$

3.624. $\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx$

$$\begin{aligned}
& \int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\
& \quad \downarrow \text{4090} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\int \frac{b(2Ab + 3aB) \tan^2(c + dx) + 2(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(2aA - bB)}{2\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{2} \int \frac{b(2Ab + 3aB) \tan^2(c + dx) + 2(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(2aA - bB)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{2} \int \frac{b(2Ab + 3aB) \tan(c + dx)^2 + 2(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(2aA - bB)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \right) \\
& \quad \downarrow \text{4138} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{\int \frac{b(2Ab + 3aB) \tan^2(c + dx) + 2(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(2aA - bB)}{\sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)} (\tan^2(c + dx) + 1)} d \tan(c + dx)}{2d} + \frac{bB \sqrt{\tan(c + dx)}}{d} \right) \\
& \quad \downarrow \text{2035} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{\int \frac{b(2Ab + 3aB) \tan^2(c + dx) + 2(Ba^2 + 2Aba - b^2B) \tan(c + dx) + a(2aA - bB)}{\sqrt{a + b \tan(c + dx)} (\tan^2(c + dx) + 1)} d \sqrt{\tan(c + dx)}}{d} + \frac{bB \sqrt{\tan(c + dx)}}{d} \right) \\
& \quad \downarrow \text{2257} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{\int \left(\frac{b(2Ab + 3aB)}{\sqrt{a + b \tan(c + dx)}} + \frac{2(Aa^2 - 2bBa - Ab^2 + (Ba^2 + 2Aba - b^2B) \tan(c + dx))}{\sqrt{a + b \tan(c + dx)} (\tan^2(c + dx) + 1)} \right) d \sqrt{\tan(c + dx)}}{d} + \frac{bB \sqrt{\tan(c + dx)}}{d} \right) \\
& \quad \downarrow \text{2009} \\
& \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{bB \sqrt{\tan(c + dx)} \sqrt{a + b \tan(c + dx)}}{d} + \frac{-((-b + ia)^{3/2} (A + iB) \arctan \left(\frac{\sqrt{-b + ia} \sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}} \right)}}{d} \right)
\end{aligned}$$

3.624. $\int \sqrt{\cot(c + dx)} (a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx$

input `Int[Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((I*a - b)^(3/2)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + Sqrt[b]*(2*A*b + 3*a*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])] - (I*a + b)^(3/2)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/d + (b*B*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d)`

3.624.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2035 `Int[(F_x_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[F_x, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[F_x, x]`

rule 2257 `Int[(P_x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[P_x*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[P_x, x] && IntegerQ[p]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4090 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.624.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.50 (sec) , antiderivative size = 2398750, normalized size of antiderivative = 9086.17

output too large to display

```
input int(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x)
```

```
output result too large to display
```

$$3.624. \quad \int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx)) dx$$

3.624.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12219 vs. $2(212) = 424$.

Time = 4.89 (sec) , antiderivative size = 24471, normalized size of antiderivative = 92.69

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.624.6 Sympy [F]

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx = \int (A + B\tan(c+dx))(a+b\tan(c+dx))^{\frac{3}{2}}\sqrt{\cot(c+dx)}dx$$

input `integrate(cot(d*x+c)**(1/2)*(a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)*sqrt(cot(c + d*x)), x)`

3.624.7 Maxima [F]

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{3/2}(A + B\tan(c+dx))dx = \int (B\tan(dx+c) + A)(b\tan(dx+c) + a)^{\frac{3}{2}}\sqrt{\cot(dx+c)}dx$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)*sqrt(cot(d*x + c)), x)`

3.624.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.624.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx)) dx = \int \sqrt{\cot(c+dx)}(A+B\tan(c+dx))(a+b\tan(c+dx))^{3/2} dx$$

input `int(cot(c+d*x)^(1/2)*(A+B*tan(c+d*x))*(a+b*tan(c+d*x))^(3/2),x)`

output `int(cot(c+d*x)^(1/2)*(A+B*tan(c+d*x))*(a+b*tan(c+d*x))^(3/2),x)`

3.625
$$\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.625.1 Optimal result	5972
3.625.2 Mathematica [A] (verified)	5973
3.625.3 Rubi [A] (verified)	5973
3.625.4 Maple [B] (warning: unable to verify)	5977
3.625.5 Fricas [B] (verification not implemented)	5977
3.625.6 Sympy [F]	5978
3.625.7 Maxima [F]	5978
3.625.8 Giac [F(-1)]	5978
3.625.9 Mupad [F(-1)]	5979

3.625.1 Optimal result

Integrand size = 35, antiderivative size = 328

$$\int \frac{(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \frac{(a + ib)^2(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}}{\sqrt{ia - bd}}$$

$$+ \frac{(12aAb + 3a^2B - 8b^2B) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{4\sqrt{bd}}$$

$$+ \frac{(ia + b)^{3/2}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{bB\sqrt{a + b \tan(c + dx)}}{2d \cot^{\frac{3}{2}}(c + dx)} + \frac{(4Ab + 5aB)\sqrt{a + b \tan(c + dx)}}{4d\sqrt{\cot(c + dx)}}$$

output

```
(I*a+b)^(3/2)*(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(a+I*b)^2*(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*a-b)^(1/2)+1/4*(12*A*a*b+3*B*a^2-8*B*b^2)*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/b^(1/2)+1/2*b*B*(a+b*tan(d*x+c))^(1/2)/d/cot(d*x+c)^(3/2)+1/4*(4*A*b+5*B*a)*(a+b*tan(d*x+c))^(1/2)/d/cot(d*x+c)^(1/2)
```

3.625.2 Mathematica [A] (verified)

Time = 3.54 (sec) , antiderivative size = 310, normalized size of antiderivative = 0.95

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(-4\sqrt[4]{-1}(-a + ib)^{3/2} (A -$$

input `Integrate[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]], x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-4*(-1)^(1/4)*(-a + I*b)^(3/2)*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]) + 4*(-1)^(1/4)*(a + I*b)^(3/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]) + (4*A*b + 5*a*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]] + 2*b*B*Tan[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]] + (Sqrt[a]*(12*a*A*b + 3*a^2*B - 8*b^2*B)*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/Sqrt[b]*Sqrt[a + b*Tan[c + d*x]])))/(4*d)`

3.625.3 Rubi [A] (verified)Time = 1.74 (sec) , antiderivative size = 284, normalized size of antiderivative = 0.87, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 4729, 3042, 4090, 27, 3042, 4130, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \sqrt{\tan(c + dx)} (a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx)) dx$$

$$\begin{aligned} & \downarrow 3042 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))dx \\ & \downarrow 4090 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \int \frac{\sqrt{\tan(c+dx)}(b(4Ab+5aB)\tan^2(c+dx)+4(Ba^2+2Aba-b^2B)\tan(c+dx))}{2\sqrt{a+b\tan(c+dx)}} dx \right) \\ & \downarrow 27 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \int \frac{\sqrt{\tan(c+dx)}(b(4Ab+5aB)\tan^2(c+dx)+4(Ba^2+2Aba-b^2B)\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx \right) \\ & \downarrow 3042 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \int \frac{\sqrt{\tan(c+dx)}(b(4Ab+5aB)\tan(c+dx)^2+4(Ba^2+2Aba-b^2B)\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx \right) \\ & \downarrow 4130 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \left(\frac{\int -\frac{b(3Ba^2+12Aba-8b^2B)\tan^2(c+dx)-8b(Aa^2-2bBa-Ab^2)\tan(c+dx)+ab(4Ab+5aB)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{b} + \dots \right) \right) \\ & \downarrow 27 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \left(\frac{(5aB+4Ab)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{\int \frac{-b(3Ba^2+12Aba-8b^2B)\tan^2(c+dx)+ab(4Ab+5aB)}{\sqrt{\tan(c+dx)}} dx}{b} \right) \right) \\ & \downarrow 3042 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \left(\frac{(5aB+4Ab)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{\int \frac{-b(3Ba^2+12Aba-8b^2B)\tan(c+dx)+ab(4Ab+5aB)}{\sqrt{\tan(c+dx)}} dx}{b} \right) \right) \\ & \downarrow 4138 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \left(\frac{(5aB+4Ab)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d} - \frac{\int \frac{-b(3Ba^2+12Aba-8b^2B)\tan^2(c+dx)+ab(4Ab+5aB)}{\sqrt{\tan(c+dx)}} dx}{b} \right) \right) \end{aligned}$$

3.625. $\int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

↓ 2035

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{(5aB+4Ab)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d}-\int\frac{-b(3Ba^2+12Aba-8b^2B)\tan^2(c+dx)}{\sqrt{a+b\tan(c+dx)}}dx\right)\right)$$

↓ 2257

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{(5aB+4Ab)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d}-\int\left(\frac{8(b(Ba^2+2Aba-b^2B)-b(Aa^2-2Ab^2))}{\sqrt{a+b\tan(c+dx)}(\tan(c+dx)+1)}\right)dx\right)\right)$$

↓ 2009

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{bB\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}{2d}+\frac{1}{4}\left(\frac{(5aB+4Ab)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d}\right)\right)$$

input `Int[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((b*B*Tan[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]])/(2*d) + (-(4*(I*a - b)^(3/2)*b*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - Sqrt[b]*(12*a*A*b + 3*a^2*B - 8*b^2*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - 4*b*(I*a + b)^(3/2)*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(b*d) + ((4*A*b + 5*a*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d)/4)`

3.625.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2035 `Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]`

rule 2257 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol
] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a
, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.
) + (f_.)*(x_)])^2, x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))`

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.625.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.39 (sec) , antiderivative size = 2398591, normalized size of antiderivative = 7312.78

output too large to display

```
input int((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x)
```

```
output result too large to display
```

3.625.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12263 vs. 2(264) = 528.

Time = 5.02 (sec) , antiderivative size = 24563, normalized size of antiderivative = 74.89

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algo
rithm="fracas")
```

```
output Too large to include
```

3.625. $\int \frac{(a+b \tan(c+dx))^{3/2} (A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

3.625.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2}}{\sqrt{\cot(c + dx)}} dx$$

input `integrate((a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)/sqrt(cot(c + d*x)), x)`

3.625.7 Maxima [F]

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^{3/2}}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)/sqrt(cot(d*x + c)), x)`

3.625.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algo
rithm="giac")`

output `Timed out`

3.625.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2}}{\sqrt{\cot(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/cot(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/cot(c + d*x)^(1/2),x)`

3.626 $\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

3.626.1 Optimal result	5980
3.626.2 Mathematica [A] (verified)	5981
3.626.3 Rubi [A] (verified)	5981
3.626.4 Maple [B] (warning: unable to verify)	5986
3.626.5 Fricas [B] (verification not implemented)	5986
3.626.6 Sympy [F]	5987
3.626.7 Maxima [F]	5987
3.626.8 Giac [F(-1)]	5987
3.626.9 Mupad [F(-1)]	5988

3.626.1 Optimal result

Integrand size = 35, antiderivative size = 383

$$\int \frac{(a + b \tan(c + dx))^{3/2}(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx = \frac{(ia - b)^{3/2}(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}}{d}$$

$$+ \frac{(6a^2Ab - 16Ab^3 - a^3B - 24ab^2B) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{8b^{3/2}d}$$

$$+ \frac{(ia + b)^{3/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{(6aAb - a^2B - 8b^2B) \sqrt{a + b \tan(c + dx)}}{8bd\sqrt{\cot(c + dx)}}$$

$$+ \frac{(6Ab - aB)(a + b \tan(c + dx))^{3/2}}{12bd\sqrt{\cot(c + dx)}} + \frac{B(a + b \tan(c + dx))^{5/2}}{3bd\sqrt{\cot(c + dx)}}$$

```
output (I*a-b)^(3/2)*(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)
)^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+1/8*(6*A*a^2*b-16*A*b^3-B*a^
3-24*B*a*b^2)*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)^(1/2))*cot
(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/b^(3/2)/d+(I*a+b)^(3/2)*(A-I*B)*arctanh((I*
a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)^(1/2))*cot(d*x+c)^(1/2)*tan(d
*x+c)^(1/2)/d+1/8*(6*A*a*b-B*a^2-8*B*b^2)*(a+b*tan(d*x+c))^(1/2)/b/d/cot(d
*x+c)^(1/2)+1/12*(6*A*b-B*a)*(a+b*tan(d*x+c))^(3/2)/b/d/cot(d*x+c)^(1/2)+1
/3*B*(a+b*tan(d*x+c))^(5/2)/b/d/cot(d*x+c)^(1/2)
```

3.626. $\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

3.626.2 Mathematica [A] (verified)

Time = 6.17 (sec) , antiderivative size = 367, normalized size of antiderivative = 0.96

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\cot^{3/2}(c + dx)} dx = \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(24 \sqrt[4]{-1} (-a + ib)^{3/2} b (iA + \dots \right)}{\dots}$$

input `Integrate[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(3/2),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(24*(-1)^(1/4)*(-a + I*b)^(3/2)*b*(I*A + B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]] + 24*(-1)^(3/4)*(a + I*b)^(3/2)*b*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]] - 3*(-6*a*A*b + a^2*B + 8*b^2*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]] + 2*(6*A*b - a*B)*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2) + 8*B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(5/2) - (3*Sqrt[a]*(-6*a^2*A*b + 16*A*b^3 + a^3*B + 24*a*b^2*B)*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/Sqrt[b]*Sqrt[a + b*Tan[c + d*x]]))/(24*b*d)`

3.626.3 Rubi [A] (verified)

Time = 2.15 (sec) , antiderivative size = 347, normalized size of antiderivative = 0.91, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 4729, 3042, 4090, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\cot^{3/2}(c + dx)} dx \xrightarrow{3042} \int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\cot(c + dx)^{3/2}} dx \xrightarrow{4729}$$

$$\begin{aligned} & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))dx \\ & \qquad \qquad \qquad \downarrow \text{3042} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \tan(c+dx)^{3/2}(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))dx \\ & \qquad \qquad \qquad \downarrow \text{4090} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int -\frac{(a+b \tan(c+dx))^{3/2}(-((6Ab-aB) \tan^2(c+dx)+6bB \tan(c+dx)+aB))dx}{2\sqrt{\tan(c+dx)}}}{3b} + \frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}{3b} \right) \\ & \qquad \qquad \qquad \downarrow \text{27} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}}{3bd} - \frac{\int \frac{(a+b \tan(c+dx))^{3/2}(-((6Ab-aB) \tan^2(c+dx))+6bB \tan(c+dx)+aB)}{\sqrt{\tan(c+dx)}}}{6b} \right) \\ & \qquad \qquad \qquad \downarrow \text{3042} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}}{3bd} - \frac{\int \frac{(a+b \tan(c+dx))^{3/2}(-((6Ab-aB) \tan(c+dx)^2)+6bB \tan(c+dx)+aB)}{\sqrt{\tan(c+dx)}}}{6b} \right) \\ & \qquad \qquad \qquad \downarrow \text{4130} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}}{3bd} - \frac{\frac{1}{2} \int \frac{3\sqrt{a+b \tan(c+dx)}(-((-Ba^2+6Aba-8b^2B) \tan^2(c+dx)+6aB \tan(c+dx)+a^2))}{2\sqrt{\tan(c+dx)}}}{6b} \right) \\ & \qquad \qquad \qquad \downarrow \text{27} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}}{3bd} - \frac{\frac{3}{4} \int \frac{\sqrt{a+b \tan(c+dx)}(-((-Ba^2+6Aba-8b^2B) \tan^2(c+dx)+6aB \tan(c+dx)+a^2))}{\sqrt{\tan(c+dx)}}}{6b} \right) \\ & \qquad \qquad \qquad \downarrow \text{3042} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}}{3bd} - \frac{\frac{3}{4} \int \frac{\sqrt{a+b \tan(c+dx)}(-((-Ba^2+6Aba-8b^2B) \tan^2(c+dx)+6aB \tan(c+dx)+a^2))}{\sqrt{\tan(c+dx)}}}{6b} \right) \\ & \qquad \qquad \qquad \downarrow \text{4130} \end{aligned}$$

3.626. $\int \frac{(a+b \tan(c+dx))^{3/2}(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd}-\frac{\frac{3}{4}\left(\int\frac{-((-Ba^3+6Aba^2-24b^2Ba-16Ab^3)\tan^2(c+dx)+\dots)}{2\sqrt{\tan(c+dx)}}\right)}{\dots}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd}-\frac{\frac{3}{4}\left(\frac{1}{2}\int\frac{-((-Ba^3+6Aba^2-24b^2Ba-16Ab^3)\tan^2(c+dx)+\dots)}{\sqrt{\tan(c+dx)}}\right)}{\dots}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd}-\frac{\frac{3}{4}\left(\frac{1}{2}\int\frac{-((-Ba^3+6Aba^2-24b^2Ba-16Ab^3)\tan(c+dx)+\dots)}{\sqrt{\tan(c+dx)}}\right)}{\dots}\right)$$

↓ 4138

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd}-\frac{\frac{3}{4}\left(\int\frac{-((-Ba^3+6Aba^2-24b^2Ba-16Ab^3)\tan^2(c+dx)+\dots)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}\right)}{\dots}\right)$$

↓ 2035

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd}-\frac{\frac{3}{4}\left(\int\frac{-((-Ba^3+6Aba^2-24b^2Ba-16Ab^3)\tan^2(c+dx)+\dots)}{\sqrt{a+b\tan(c+dx)}}\right)}{\dots}\right)$$

↓ 2257

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd} - \frac{3}{4} \left(\int \left(\frac{Ba^3-6Aba^2+24b^2Ba+16Ab^3}{\sqrt{a+b\tan(c+dx)}} + \frac{16(b(Aa^2-2bBa-b^3))}{\sqrt{a+b\tan(c+dx)}} \right) dx \right) \right)$$

↓ 2009

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3bd} - \frac{(6Ab-aB)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d} + \frac{3}{4} \left(\int \left(\frac{Ba^3-6Aba^2+24b^2Ba+16Ab^3}{\sqrt{a+b\tan(c+dx)}} + \frac{16(b(Aa^2-2bBa-b^3))}{\sqrt{a+b\tan(c+dx)}} \right) dx \right) \right)$$

```
input Int[((a + b*Tan[c + d*x])^(3/2)*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(3/2),x]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(5/2))/(3*b*d) - (-1/2*((6*A*b - a*B)*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2))/d + (3*((-8*(I*a - b)^(3/2)*b*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - ((6*a^2*A*b - 16*A*b^3 - a^3*B - 24*a*b^2*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/Sqrt[b] - 8*b*(I*a + b)^(3/2)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/d - ((6*a*A*b - a^2*B - 8*b^2*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d))/4)/(6*b))
```

3.626.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

3.626. $\int \frac{(a+b\tan(c+dx))^{3/2}(A+B\tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

rule 2035 `Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]`

rule 2257 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol
] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a
, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.
) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))`


```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.626.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.64 (sec) , antiderivative size = 2403527, normalized size of antiderivative = 6275.53

output too large to display

```
input int((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x)
```

```
output result too large to display
```

3.626.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12332 vs. $2(312) = 624$.

Time = 5.51 (sec) , antiderivative size = 24697, normalized size of antiderivative = 64.48

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\cot^{3/2}(c + dx)} dx = \text{Too large to display}$$

```
input integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algo
rithm="fracas")
```

```
output Too large to include
```

3.626. $\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\cot^{3/2}(c + dx)} dx$

3.626.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\cot^{3/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2}}{\cot^{3/2}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**(3/2)/cot(c + d*x)**(3/2), x)`

3.626.7 Maxima [F]

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\cot^{3/2}(c + dx)} dx = \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^{3/2}}{\cot(dx + c)^{3/2}} dx$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algo rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(3/2)/cot(d*x + c)^(3/2), x)`

3.626.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\cot^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(3/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algo rithm="giac")`

output `Timed out`

3.626.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{3/2} (A + B \tan(c + dx))}{\cot^{3/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{3/2}}{\cot(c + dx)^{3/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/cot(c + d*x)^(3/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(3/2))/cot(c + d*x)^(3/2),x)`

3.627 $\int \cot^{\frac{13}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.627.1 Optimal result	5989
3.627.2 Mathematica [A] (verified)	5991
3.627.3 Rubi [A] (verified)	5992
3.627.4 Maple [B] (warning: unable to verify)	6002
3.627.5 Fricas [B] (verification not implemented)	6002
3.627.6 Sympy [F(-1)]	6003
3.627.7 Maxima [F]	6003
3.627.8 Giac [F(-1)]	6003
3.627.9 Mupad [F(-1)]	6004

3.627.1 Optimal result

Integrand size = 35, antiderivative size = 500

$$\int \cot^{\frac{13}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx =$$

$$\frac{(ia - b)^{5/2}(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d}$$

$$- \frac{(ia + b)^{5/2}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d}$$

$$- \frac{2(8085a^4 Ab - 495a^2 Ab^3 + 40Ab^5 + 3465a^5 B - 5313a^3 b^2 B - 110ab^4 B) \sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}}{3465a^3 d}$$

$$- \frac{2(1155a^4 A - 1485a^2 Ab^2 - 20Ab^4 - 2541a^3 b B + 55ab^3 B) \cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{3465a^2 d}$$

$$+ \frac{2(495a^2 Ab - 5Ab^3 + 231a^3 B - 275ab^2 B) \cot^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{1155ad}$$

$$+ \frac{2(99a^2 A - 113Ab^2 - 209abB) \cot^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{693d}$$

$$- \frac{2a(14Ab + 11aB) \cot^{\frac{9}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{99d}$$

$$- \frac{2aA \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}}{11d}$$

3.627. $\int \cot^{\frac{13}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

output $-(I*a-b)^{(5/2)}*(I*A-B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)/d}-(I*a+b)^{(5/2)}*(I*A+B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)/d}-2/3465*(1155*A*a^4-1485*A*a^2*b^2-20*A*b^4-2541*B*a^3*b+55*B*a*b^3)*\cot(d*x+c)^{(3/2)}*(a+b*\tan(d*x+c))^{(1/2)}/a^2/d+2/1155*(495*A*a^2*b-5*A*b^3+231*B*a^3-275*B*a*b^2)*\cot(d*x+c)^{(5/2)}*(a+b*\tan(d*x+c))^{(1/2)}/a/d+2/693*(99*A*a^2-113*A*b^2-209*B*a*b)*\cot(d*x+c)^{(7/2)}*(a+b*\tan(d*x+c))^{(1/2)}/d-2/99*a*(14*A*b+11*B*a)*\cot(d*x+c)^{(9/2)}*(a+b*\tan(d*x+c))^{(1/2)}/d-2/3465*(8085*A*a^4*b-495*A*a^2*b^3+40*A*b^5+3465*B*a^5-5313*B*a^3*b^2-110*B*a*b^4)*\cot(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(1/2)}/a^3/d-2/11*a*A*\cot(d*x+c)^{(11/2)}*(a+b*\tan(d*x+c))^{(3/2)}/d$

3.627.2 Mathematica [A] (verified)

Time = 7.22 (sec) , antiderivative size = 653, normalized size of antiderivative = 1.31

$$\int \cot^{\frac{13}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A$$

$$3.627. \quad \int \cot^{\frac{13}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

input `Integrate[Cot[c + d*x]^(13/2)*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-1/4*(b*B*(a + b*Tan[c + d*x])^(3/2))/(d*Tan[c + d*x]^(11/2)) + (-1/10*(b*(8*A*b + 5*a*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Tan[c + d*x]^(11/2)) + (-1/22*((80*a^2*A - 88*A*b^2 - 165*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Tan[c + d*x]^(11/2)) - (2*((5*a*(184*a*A*b + 88*a^2*B - 99*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(18*d*Tan[c + d*x]^(9/2)) - (2*((10*a^2*(99*a^2*A - 113*A*b^2 - 209*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(7*d*Tan[c + d*x]^(7/2)) - (2*((-3*a^2*(495*a^2*A*b - 5*A*b^3 + 231*a^3*B - 275*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Tan[c + d*x]^(5/2)) - (2*((-5*a^2*(1155*a^4*A - 1485*a^2*A*b^2 - 20*A*b^4 - 2541*a^3*b*B + 55*a*b^3*B)*Sqrt[a + b*Tan[c + d*x]])/(2*d*Tan[c + d*x]^(3/2)) - (2*((51975*a^5*((-1)^(3/4)*(-a + I*b)^(5/2)*(A - I*B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) - (-1)^(3/4)*(a + I*b)^(5/2)*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])))/(8*d) + (15*a^2*(8085*a^4*A*b - 495*a^2*A*b^3 + 40*A*b^5 + 3465*a^5*B - 5313*a^3*b^2*B - 110*a*b^4*B)*Sqrt[a + b*Tan[c + d*x]])/(4*d*Sqrt[Tan[c + d*x]])))/(3*a)))/(5*a)))/(7*a)))/(9*a)))/(11*a))/5)/4`

3.627.3 Rubi [A] (verified)

Time = 3.99 (sec) , antiderivative size = 530, normalized size of antiderivative = 1.06, number of steps used = 28, number of rules used = 27, $\frac{\text{number of rules}}{\text{integrand size}} = 0.771$, Rules used = {3042, 4729, 3042, 4088, 27, 3042, 4128, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{13}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

↓ 3042

$$\int \cot(c + dx)^{13/2}(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

↓ 4729

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{13}{2}}(c + dx)} dx$$

↓ 3042

3.627. $\int \cot^{\frac{13}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\tan(c+dx)^{13/2}} dx$$

↓ 4088

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2}{11} \int \frac{\sqrt{a+b\tan(c+dx)}(-b(8aA-11bB)\tan^2(c+dx)-11(Aa^2-2bBa-Ab^2))}{2\tan^{\frac{11}{2}}(c+dx)} dx \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{11} \int \frac{\sqrt{a+b\tan(c+dx)}(-b(8aA-11bB)\tan^2(c+dx)-11(Aa^2-2bBa-Ab^2))}{\tan^{\frac{11}{2}}(c+dx)} dx \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{11} \int \frac{\sqrt{a+b\tan(c+dx)}(-b(8aA-11bB)\tan(c+dx)^2-11(Aa^2-2bBa-Ab^2))}{\tan(c+dx)^{11/2}} dx \right)$$

↓ 4128

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{11} \left(\frac{2}{9} \int -\frac{b(88Ba^2+184Aba-99b^2B)\tan^2(c+dx)+99(Ba^3+3Aba^2-3b^2Ba-Ab^2)}{2\tan^{\frac{9}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}} dx \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{11} \left(-\frac{1}{9} \int \frac{b(88Ba^2+184Aba-99b^2B)\tan^2(c+dx)+99(Ba^3+3Aba^2-3b^2Ba-Ab^2)}{\tan^{\frac{9}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}} dx \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{11} \left(-\frac{1}{9} \int \frac{b(88Ba^2+184Aba-99b^2B)\tan(c+dx)^2+99(Ba^3+3Aba^2-3b^2Ba-Ab^2)}{\tan(c+dx)^{9/2}\sqrt{a+b\tan(c+dx)}} dx \right) \right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{11} \left(\frac{1}{9} \left(2 \int -\frac{3(-2ab(99Aa^2-209bBa-113Ab^2)\tan^2(c+dx)-231a(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx)+99(Aa^3-3bBa^2-3Ab^2a+b^3B))}{2\tan^{\frac{7}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}} dx \right) \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{2(99a^2A-209abB-113Ab^2)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)}-\frac{3\int\frac{-2ab(99Aa^2-209bBa)}{\dots}}{\dots}\right)\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{2(99a^2A-209abB-113Ab^2)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)}-\frac{3\int\frac{-2ab(99Aa^2-209bBa)}{\dots}}{\dots}\right)\right)\right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{2(99a^2A-209abB-113Ab^2)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)}-\frac{3\left(\frac{2\int\frac{1155(Ba^3+3Aba^2)}{\dots}}{\dots}\right)}{\dots}\right)\right)\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{2(99a^2A-209abB-113Ab^2)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)}-\frac{3\left(\frac{\int\frac{1155(Ba^3+3Aba^2)}{\dots}}{\dots}\right)}{\dots}\right)\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{11}\left(\frac{1}{9}\left(\frac{2(99a^2A-209abB-113Ab^2)\sqrt{a+b\tan(c+dx)}}{7d\tan^{\frac{7}{2}}(c+dx)}-\frac{3\left(\frac{\int\frac{1155(Ba^3+3Aba^2)}{\dots}}{\dots}\right)}{\dots}\right)\right)\right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{11} \left(\frac{1}{9} \frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} - \frac{3 \left(\int \frac{-3465(Aa^3 - 3Ba^2)}{\dots}}{\dots} \right)}{\dots} \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{11} \left(\frac{1}{9} \frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} - \frac{3 \left(\int \frac{-3465(Aa^3 - 3Ba^2)}{\dots}}{\dots} \right)}{\dots} \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{11} \left(\frac{1}{9} \frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} - \frac{3 \left(\int \frac{-3465(Aa^3 - 3Ba^2)}{\dots}}{\dots} \right)}{\dots} \right) \right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{11} \right) \left(\frac{1}{9} \right) \frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} - \frac{3 \left(\frac{2 \int \frac{3465((Aa^3-3bBa^2)}{\dots}}{\dots}}{\dots} \right)}{\dots}$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{11} \right) \left(\frac{1}{9} \right) \frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} - \frac{3 \left(\frac{3465 \int \frac{(Aa^3-3bBa^2)}{\dots}}{\dots} \right)}{\dots}$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{11} \right) \left(\frac{1}{9} \right) \frac{2(99a^2A - 209abB - 113Ab^2) \sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} - \frac{3 \left(\frac{3465 \int \frac{(Aa^3-3bBa^2)}{\dots}}{\dots} \right)}{\dots}$$

↓ 4099

3.627. $\int \cot^{\frac{13}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{11d\tan^{\frac{11}{2}}(c+dx)} + \frac{1}{11} \right) - \frac{2a(11aB+14Ab)\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} \left(\frac{2}{9} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{11d\tan^{\frac{11}{2}}(c+dx)} + \frac{1}{11} \right) - \frac{2a(11aB+14Ab)\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} \left(\frac{2}{9} \right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{11d\tan^{\frac{11}{2}}(c+dx)} + \frac{1}{11} \right) - \frac{2a(11aB+14Ab)\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} \left(\frac{2}{9} \right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{11d\tan^{\frac{11}{2}}(c+dx)} + \frac{1}{11} - \frac{2a(11aB+14Ab)\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{11d\tan^{\frac{11}{2}}(c+dx)} + \frac{1}{11} - \frac{2a(11aB+14Ab)\sqrt{a+b\tan(c+dx)}}{9d\tan^{\frac{9}{2}}(c+dx)} + \frac{1}{9} \right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{11d\tan^{11/2}(c+dx)} + \frac{1}{11} - \frac{2a(11aB+14Ab)\sqrt{a+b\tan(c+dx)}}{9d\tan^{9/2}(c+dx)} + \frac{1}{9} \right)$$

```
input Int[Cot[c + d*x]^(13/2)*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*a*A*(a + b*Tan[c + d*x])^(3/2))
/(11*d*Tan[c + d*x]^(11/2)) + ((-2*a*(14*A*b + 11*a*B)*Sqrt[a + b*Tan[c +
d*x]])/(9*d*Tan[c + d*x]^(9/2)) + ((2*(99*a^2*A - 113*A*b^2 - 209*a*b*B)*S
qrt[a + b*Tan[c + d*x]])/(7*d*Tan[c + d*x]^(7/2)) - (3*((-2*(495*a^2*A*b -
5*A*b^3 + 231*a^3*B - 275*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(5*d*Tan[c +
d*x]^(5/2)) - ((-2*(1155*a^4*A - 1485*a^2*A*b^2 - 20*A*b^4 - 2541*a^3*b*B
+ 55*a*b^3*B)*Sqrt[a + b*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) + ((-346
5*((a^4*(a + I*b)^3*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sq
rt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d) + (a^4*(a - I*b)^3*(A - I*B)*Ar
cTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[
I*a + b]*d)))/a - (2*(8085*a^4*A*b - 495*a^2*A*b^3 + 40*A*b^5 + 3465*a^5*B
- 5313*a^3*b^2*B - 110*a*b^4*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Sqrt[Tan[c +
d*x]])))/(3*a)/(5*a))/(7*a))/9)/11)
```

3.627.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && ! (ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.627.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.45 (sec) , antiderivative size = 2660853, normalized size of antiderivative = 5321.71

output too large to display

input `int(cot(d*x+c)^(13/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x)`

output `result too large to display`

3.627.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19121 vs. $2(428) = 856$.

Time = 4.96 (sec) , antiderivative size = 19121, normalized size of antiderivative = 38.24

$$\int \cot^{\frac{13}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(13/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fracas")`

output `Too large to include`

3.627.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{13}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(13/2)*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.627.7 Maxima [F]

$$\int \cot^{\frac{13}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{5}{2}} \cot(dx + c)^{\frac{13}{2}} dx$$

input `integrate(cot(d*x+c)^(13/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^(13/2), x)`

3.627.8 Giac [F(-1)]

Timed out.

$$\int \cot^{\frac{13}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(13/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.627. $\int \cot^{\frac{13}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.627.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{13}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{13/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2} dx$$

input `int(cot(c + d*x)^(13/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)`

output `int(cot(c + d*x)^(13/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2), x)`

3.628 $\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.628.1 Optimal result	6005
3.628.2 Mathematica [A] (verified)	6007
3.628.3 Rubi [A] (verified)	6008
3.628.4 Maple [B] (warning: unable to verify)	6015
3.628.5 Fricas [B] (verification not implemented)	6015
3.628.6 Sympy [F(-1)]	6016
3.628.7 Maxima [F]	6016
3.628.8 Giac [F(-1)]	6016
3.628.9 Mupad [F(-1)]	6017

3.628.1 Optimal result

Integrand size = 35, antiderivative size = 418

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{(ia - b)^{5/2}(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d} - \frac{(ia + b)^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d} - \frac{2(315a^4A - 483a^2Ab^2 - 10Ab^4 - 735a^3bB + 45ab^3B) \sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}}{315a^2d} + \frac{2(231a^2Ab - 5Ab^3 + 105a^3B - 135ab^2B) \cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{315ad} + \frac{2(21a^2A - 25Ab^2 - 45abB) \cot^{\frac{5}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{105d} - \frac{2a(4Ab + 3aB) \cot^{\frac{7}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}}{21d} - \frac{2aA \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}}{9d}$$

output $(I*a-b)^{(5/2)}*(A+I*B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)/d}-(I*a+b)^{(5/2)}*(A-I*B)*\arctan(h((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)/d}+2/315*(231*A*a^2*b-5*A*b^3+105*B*a^3-135*B*a*b^2)*\cot(d*x+c)^{(3/2)}*(a+b*\tan(d*x+c))^{(1/2)}/a/d+2/105*(21*A*a^2-25*A*b^2-45*B*a*b)*\cot(d*x+c)^{(5/2)}*(a+b*\tan(d*x+c))^{(1/2)/d}-2/21*a*(4*A*b+3*B*a)*\cot(d*x+c)^{(7/2)}*(a+b*\tan(d*x+c))^{(1/2)/d}-2/315*(315*A*a^4-483*A*a^2*b^2-10*A*b^4-735*B*a^3*b+45*B*a*b^3)*\cot(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(1/2)}/a^2/d-2/9*a*A*\cot(d*x+c)^{(9/2)}*(a+b*\tan(d*x+c))^{(3/2)}/d$

3.628.2 Mathematica [A] (verified)

Time = 7.01 (sec) , antiderivative size = 564, normalized size of antiderivative = 1.35

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A$$

$$+ B \tan(c + dx)) dx = \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} - \frac{bB(a + b \tan(c + dx))^{3/2}}{3d \tan^{\frac{9}{2}}(c + dx)}$$

$$3.628. \quad \int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

input `Integrate[Cot[c + d*x]^(11/2)*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-1/3*(b*B*(a + b*Tan[c + d*x])^(3/2))/(d*Tan[c + d*x]^(9/2)) + ((-3*b*(2*A*b + a*B)*Sqrt[a + b*Tan[c + d*x]])/(8*d*Tan[c + d*x]^(9/2)) + (-1/6*((16*a^2*A - 18*A*b^2 - 33*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Tan[c + d*x]^(9/2)) - (2*((6*a*(38*a*A*b + 18*a^2*B - 21*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(7*d*Tan[c + d*x]^(7/2)) - (2*((18*a^2*(21*a^2*A - 25*A*b^2 - 45*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) - (2*((-3*a^2*(231*a^2*A*b - 5*A*b^3 + 105*a^3*B - 135*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Tan[c + d*x]^(3/2)) - (2*((-2835*a^4*((-1)^(1/4)*(-a + I*b)^(5/2)*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]) + (-1)^(1/4)*(a + I*b)^(5/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])))/(4*d) - (9*a^2*(315*a^4*A - 483*a^2*A*b^2 - 10*A*b^4 - 735*a^3*b*B + 45*a*b^3*B)*Sqrt[a + b*Tan[c + d*x]])/(2*d*Sqrt[Tan[c + d*x]])))/(3*a))/(5*a))/(7*a))/(9*a))/4)/3`

3.628.3 Rubi [A] (verified)

Time = 3.22 (sec) , antiderivative size = 449, normalized size of antiderivative = 1.07, number of steps used = 25, number of rules used = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.686$, Rules used = {3042, 4729, 3042, 4088, 27, 3042, 4128, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

↓ 3042

$$\int \cot(c + dx)^{11/2}(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

↓ 4729

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\tan^{\frac{11}{2}}(c + dx)} dx$$

↓ 3042

3.628. $\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\tan(c+dx)^{11/2}} dx$$

↓ 4088

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2}{9} \int \frac{3\sqrt{a+b\tan(c+dx)}(-b(2aA-3bB)\tan^2(c+dx)-3(Aa^2-2bBa-Ab^2)\tan(c+dx))}{2\tan^{9/2}(c+dx)} dx \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \int \frac{\sqrt{a+b\tan(c+dx)}(-b(2aA-3bB)\tan^2(c+dx)-3(Aa^2-2bBa-Ab^2)\tan(c+dx))}{\tan^{9/2}(c+dx)} dx \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \int \frac{\sqrt{a+b\tan(c+dx)}(-b(2aA-3bB)\tan(c+dx)^2-3(Aa^2-2bBa-Ab^2)\tan(c+dx))}{\tan(c+dx)^{9/2}} dx \right)$$

↓ 4128

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \left(\frac{2}{7} \int -\frac{b(18Ba^2+38Aba-21b^2B)\tan^2(c+dx)+21(Ba^3+3Aba^2-3b^2Ba-Ab^2)\tan(c+dx)}{2\tan^{7/2}(c+dx)\sqrt{a+b\tan(c+dx)}} dx \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \left(-\frac{1}{7} \int \frac{b(18Ba^2+38Aba-21b^2B)\tan^2(c+dx)+21(Ba^3+3Aba^2-3b^2Ba-Ab^2)\tan(c+dx)}{\tan^{7/2}(c+dx)\sqrt{a+b\tan(c+dx)}} dx \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \left(-\frac{1}{7} \int \frac{b(18Ba^2+38Aba-21b^2B)\tan(c+dx)^2+21(Ba^3+3Aba^2-3b^2Ba-Ab^2)\tan(c+dx)}{\tan(c+dx)^{7/2}\sqrt{a+b\tan(c+dx)}} dx \right) \right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \left(\frac{1}{7} \left(2 \int -\frac{-4ab(21Aa^2-45bBa-25Ab^2)\tan^2(c+dx)-105a(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx)+21(Ba^3+3Aba^2-3b^2Ba-Ab^2)\tan(c+dx)}{2\tan^{5/2}(c+dx)\sqrt{a+b\tan(c+dx)}} dx \right) \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{1}{7}\left(\frac{2(21a^2A-45abB-25Ab^2)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}-\int\frac{-4ab(21Aa^2-45bBa-25Ab^2)}{\dots}\right)\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{1}{7}\left(\frac{2(21a^2A-45abB-25Ab^2)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}-\int\frac{-4ab(21Aa^2-45bBa-25Ab^2)}{\dots}\right)\right)\right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{1}{7}\left(\frac{2(21a^2A-45abB-25Ab^2)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}-\int\frac{315(Ba^3+3Aba^2-3b^2Ba-A)}{\dots}\right)\right)\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{1}{7}\left(\frac{2(21a^2A-45abB-25Ab^2)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}-\int\frac{315(Ba^3+3Aba^2-3b^2Ba-A)}{\dots}\right)\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{1}{7}\left(\frac{2(21a^2A-45abB-25Ab^2)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}-\int\frac{315(Ba^3+3Aba^2-3b^2Ba-A)}{\dots}\right)\right)\right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{3}\left(\frac{1}{7}\left(\frac{2(21a^2A-45abB-25Ab^2)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}-\int\frac{315(a^3(Ba^3+3Aba^2-3b^2Ba-A))}{\dots}\right)\right)\right)$$

$$\begin{aligned} & \downarrow \color{blue}{27} \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} & \left(\frac{1}{3} \left(\frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2) \sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{315 \int \frac{a^3 (Ba^3 + 3Aba^2 - 3b^2Ba - \dots}{\sqrt{a+b \tan(c+dx)}} dx}{\dots} \right) \right) \right) \\ & \downarrow \color{blue}{3042} \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} & \left(\frac{1}{3} \left(\frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2) \sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{315 \int \frac{a^3 (Ba^3 + 3Aba^2 - 3b^2Ba - \dots}{\sqrt{a+b \tan(c+dx)}} dx}{\dots} \right) \right) \right) \\ & \downarrow \color{blue}{4099} \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} & \left(-\frac{2aA(a+b \tan(c+dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c+dx)} + \frac{1}{3} \left(-\frac{2a(3aB + 4Ab)\sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2) \sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{315 \int \frac{a^3 (Ba^3 + 3Aba^2 - 3b^2Ba - \dots}{\sqrt{a+b \tan(c+dx)}} dx}{\dots} \right) \right) \right) \\ & \downarrow \color{blue}{3042} \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} & \left(-\frac{2aA(a+b \tan(c+dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c+dx)} + \frac{1}{3} \left(-\frac{2a(3aB + 4Ab)\sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2) \sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{315 \int \frac{a^3 (Ba^3 + 3Aba^2 - 3b^2Ba - \dots}{\sqrt{a+b \tan(c+dx)}} dx}{\dots} \right) \right) \right) \\ & \downarrow \color{blue}{4098} \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} & \left(-\frac{2aA(a+b \tan(c+dx))^{3/2}}{9d \tan^{\frac{9}{2}}(c+dx)} + \frac{1}{3} \left(-\frac{2a(3aB + 4Ab)\sqrt{a+b \tan(c+dx)}}{7d \tan^{\frac{7}{2}}(c+dx)} + \frac{1}{7} \left(\frac{2(21a^2A - 45abB - 25Ab^2) \sqrt{a+b \tan(c+dx)}}{5d \tan^{\frac{5}{2}}(c+dx)} - \frac{315 \int \frac{a^3 (Ba^3 + 3Aba^2 - 3b^2Ba - \dots}{\sqrt{a+b \tan(c+dx)}} dx}{\dots} \right) \right) \right) \end{aligned}$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{9d\tan^{9/2}(c+dx)} + \frac{1}{3} - \frac{2a(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{7d\tan^{7/2}(c+dx)} + \frac{1}{7} \right) \frac{2(21}{$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{9d\tan^{9/2}(c+dx)} + \frac{1}{3} - \frac{2a(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{7d\tan^{7/2}(c+dx)} + \frac{1}{7} \right) \frac{2(21}{$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{9d\tan^{9/2}(c+dx)} + \frac{1}{3} - \frac{2a(3aB+4Ab)\sqrt{a+b\tan(c+dx)}}{7d\tan^{7/2}(c+dx)} + \frac{1}{7} \right) \frac{2(21}{$$

input `Int[Cot[c + d*x]^(11/2)*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

```

output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*a*A*(a + b*Tan[c + d*x])^(3/2))
/(9*d*Tan[c + d*x]^(9/2)) + ((-2*a*(4*A*b + 3*a*B)*Sqrt[a + b*Tan[c + d*x]
])/ (7*d*Tan[c + d*x]^(7/2)) + ((2*(21*a^2*A - 25*A*b^2 - 45*a*b*B)*Sqrt[a
+ b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) - ((-2*(231*a^2*A*b - 5*A*b^3
+ 105*a^3*B - 135*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/
2)) - ((315*(-((a^3*(a + I*b)^3*(I*A - B)*ArcTan[(Sqrt[I*a - b])*Sqrt[Tan[c
+ d*x]])/Sqrt[a + b*Tan[c + d*x]]])/ (Sqrt[I*a - b]*d)) + (a^3*(a - I*b)^3
*(I*A + B)*ArcTanh[(Sqrt[I*a + b])*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d
*x]]])/ (Sqrt[I*a + b]*d))/a - (2*(315*a^4*A - 483*a^2*A*b^2 - 10*A*b^4 -
735*a^3*b*B + 45*a*b^3*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]
))/(3*a)/(5*a))/7)/3)

```

3.628.3.1 Defintions of rubi rules used

```

rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]

```

```

rule 104 Int[(((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)))/((e_.) + (f_.)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

```

```

rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])

```

```

rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])

```

```

rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]

```

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.628.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.64 (sec) , antiderivative size = 2658291, normalized size of antiderivative = 6359.55

output too large to display

```
input int(cot(d*x+c)^(11/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x)
```

```
output result too large to display
```

3.628.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19062 vs. $2(352) = 704$.

Time = 4.98 (sec) , antiderivative size = 19062, normalized size of antiderivative = 45.60

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

```
input integrate(cot(d*x+c)^(11/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, alg
orithm="fricas")
```

$$3.628. \quad \int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$$

output Too large to include

3.628.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(11/2)*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output Timed out

3.628.7 Maxima [F]

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{5}{2}} \cot(dx + c)^{\frac{11}{2}} dx$$

input `integrate(cot(d*x+c)^(11/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^(11/2), x)`

3.628.8 Giac [F(-1)]

Timed out.

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(11/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output Timed out

3.628. $\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.628.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{11}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{11/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2} dx$$

input `int(cot(c + d*x)^(11/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)`

output `int(cot(c + d*x)^(11/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2), x)`

3.629 $\int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.629.1 Optimal result	6018
3.629.2 Mathematica [A] (verified)	6019
3.629.3 Rubi [A] (verified)	6020
3.629.4 Maple [B] (verified)	6026
3.629.5 Fricas [B] (verification not implemented)	6027
3.629.6 Sympy [F(-1)]	6027
3.629.7 Maxima [F]	6027
3.629.8 Giac [F(-1)]	6028
3.629.9 Mupad [F(-1)]	6028

3.629.1 Optimal result

Integrand size = 35, antiderivative size = 349

$$\int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \frac{(ia-b)^{5/2}(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d} + \frac{(ia+b)^{5/2}(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d} + \frac{2(245a^2Ab - 15Ab^3 + 105a^3B - 161ab^2B) \sqrt{\cot(c+dx)} \sqrt{a+b \tan(c+dx)}}{105ad} + \frac{2(35a^2A - 45Ab^2 - 77abB) \cot^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}{105d} - \frac{2a(10Ab + 7aB) \cot^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}{35d} - \frac{2aA \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}}{7d}$$

output $(I*a-b)^{(5/2)}*(I*A-B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)/d+(I*a+b)^{(5/2)}*(I*A+B)*\arctan(h((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)/d+2/105*(35*A*a^2-45*A*b^2-77*B*a*b)*\cot(d*x+c)^{(3/2)}*(a+b*\tan(d*x+c))^{(1/2)/d-2/35*a*(10*A*b+7*B*a)*\cot(d*x+c)^{(5/2)}*(a+b*\tan(d*x+c))^{(1/2)/d+2/105*(245*A*a^2*b-15*A*b^3+105*B*a^3-161*B*a*b^2)*\cot(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(1/2)/a/d-2/7*a*A*\cot(d*x+c)^{(7/2)}*(a+b*\tan(d*x+c))^{(3/2)/d}$

3.629.2 Mathematica [A] (verified)

Time = 5.72 (sec) , antiderivative size = 382, normalized size of antiderivative = 1.09

$$\int \cot^{\frac{9}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx =$$

$$\cot^{\frac{7}{2}}(c+dx) \left(35ab(4Ab+aB)\sqrt{a+b\tan(c+dx)} + 5a(24a^2A-28Ab^2-49abB)\sqrt{a+b\tan(c+dx)} + \dots \right)$$

input `Integrate[Cot[c + d*x]^(9/2)*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output $-1/420*(\text{Cot}[c + d*x]^{(7/2)}*(35*a*b*(4*A*b + a*B)*\text{Sqrt}[a + b*\text{Tan}[c + d*x]] + 5*a*(24*a^2*A - 28*A*b^2 - 49*a*b*B)*\text{Sqrt}[a + b*\text{Tan}[c + d*x]] + 6*a*(60*a*A*b + 28*a^2*B - 35*b^2*B)*\text{Tan}[c + d*x]*\text{Sqrt}[a + b*\text{Tan}[c + d*x]] + 210*a*b*B*(a + b*\text{Tan}[c + d*x])^{(3/2)} - 4*\text{Tan}[c + d*x]^2*(105*(-1)^{(1/4)}*a*((-a + I*b)^{(5/2)}*(I*A + B)*\text{ArcTan}[((-1)^{(1/4)}*\text{Sqrt}[-a + I*b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a + b*\text{Tan}[c + d*x]]] + (a + I*b)^{(5/2)}*((-I)*A + B)*\text{ArcTan}[((-1)^{(1/4)}*\text{Sqrt}[a + I*b]*\text{Sqrt}[\text{Tan}[c + d*x]])/\text{Sqrt}[a + b*\text{Tan}[c + d*x]])]*\text{Tan}[c + d*x]^{(3/2)} + 2*a*(35*a^2*A - 45*A*b^2 - 77*a*b*B)*\text{Sqrt}[a + b*\text{Tan}[c + d*x]] + 2*(245*a^2*A*b - 15*A*b^3 + 105*a^3*B - 161*a*b^2*B)*\text{Tan}[c + d*x]*\text{Sqrt}[a + b*\text{Tan}[c + d*x]])))/(a*d)$

3.629.3 Rubi [A] (verified)

Time = 2.61 (sec) , antiderivative size = 371, normalized size of antiderivative = 1.06, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 4729, 3042, 4088, 27, 3042, 4128, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{9/2}(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{\frac{9}{2}}(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan(c+dx)^{9/2}} dx$$

$$\downarrow \text{4088}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2}{7} \int \frac{\sqrt{a+b \tan(c+dx)}(-b(4aA-7bB) \tan^2(c+dx) - 7(Aa^2 - 2bBa - Ab^2) \tan(c+dx))}{2 \tan^{\frac{7}{2}}(c+dx)} dx \right.$$

$$\downarrow \text{27}$$

$$\left. \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \int \frac{\sqrt{a+b \tan(c+dx)}(-b(4aA-7bB) \tan^2(c+dx) - 7(Aa^2 - 2bBa - Ab^2) \tan(c+dx))}{\tan^{\frac{7}{2}}(c+dx)} dx \right) \right.$$

$$\downarrow \text{3042}$$

$$\left. \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{7} \int \frac{\sqrt{a+b \tan(c+dx)}(-b(4aA-7bB) \tan(c+dx)^2 - 7(Aa^2 - 2bBa - Ab^2) \tan(c+dx))}{\tan(c+dx)^{7/2}} dx \right) \right.$$

$$\downarrow \text{4128}$$

3.629. $\int \cot^{\frac{9}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{2}{5}\int-\frac{b(28Ba^2+60Aba-35b^2B)\tan^2(c+dx)+35(Ba^3+3Aba^2-3b^2Ba-3b^3)}{2\tan^{\frac{5}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}dx\right)\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(-\frac{1}{5}\int\frac{b(28Ba^2+60Aba-35b^2B)\tan^2(c+dx)+35(Ba^3+3Aba^2-3b^2Ba-3b^3)}{\tan^{\frac{5}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}dx\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(-\frac{1}{5}\int\frac{b(28Ba^2+60Aba-35b^2B)\tan(c+dx)^2+35(Ba^3+3Aba^2-3b^2Ba-3b^3)}{\tan(c+dx)^{5/2}\sqrt{a+b\tan(c+dx)}}dx\right)\right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{2\int-\frac{-2ab(35Aa^2-77bBa-45Ab^2)\tan^2(c+dx)-105a(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx)+105a^2(Aa^2-3bBa-3Ab^2)}{2\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}}{3a}\right)\right)\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{2(35a^2A-77abB-45Ab^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\int\frac{-2ab(35Aa^2-77bBa-45Ab^2)}{3d\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}dx\right)\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{2(35a^2A-77abB-45Ab^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\int\frac{-2ab(35Aa^2-77bBa-45Ab^2)}{3d\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}dx\right)\right)\right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{2(35a^2A-77abB-45Ab^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{2\int\frac{105((Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx)+105a^2(Aa^2-3bBa-3Ab^2))}{2\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}}dx}{3a}\right)\right)\right)$$

↓ 27

3.629. $\int \cot^{\frac{9}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{2(35a^2A-77abB-45Ab^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{105\int\frac{(Aa^3-3bBa^2-3Ab^2a+b^3)}{\sqrt{\dots}}}{\dots}}\right)\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{7}\left(\frac{1}{5}\left(\frac{2(35a^2A-77abB-45Ab^2)\sqrt{a+b\tan(c+dx)}}{3d\tan^{\frac{3}{2}}(c+dx)}-\frac{105\int\frac{(Aa^3-3bBa^2-3Ab^2a+b^3)}{\sqrt{\dots}}}{\dots}}\right)\right)\right)$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{7d\tan^{\frac{7}{2}}(c+dx)}+\frac{1}{7}\left(-\frac{2a(7aB+10Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}+\frac{1}{5}\left(2(3\right)\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{7d\tan^{\frac{7}{2}}(c+dx)}+\frac{1}{7}\left(-\frac{2a(7aB+10Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}+\frac{1}{5}\left(2(3\right)\right)\right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{7d\tan^{\frac{7}{2}}(c+dx)}+\frac{1}{7}\left(-\frac{2a(7aB+10Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{\frac{5}{2}}(c+dx)}+\frac{1}{5}\left(2(3\right)\right)\right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{7d\tan^{7/2}(c+dx)} + \frac{1}{7} - \frac{2a(7aB+10Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{5/2}(c+dx)} + \frac{1}{5} \right) \frac{2(3}{$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{7d\tan^{7/2}(c+dx)} + \frac{1}{7} - \frac{2a(7aB+10Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{5/2}(c+dx)} + \frac{1}{5} \right) \frac{2(3}{$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{7d\tan^{7/2}(c+dx)} + \frac{1}{7} - \frac{2a(7aB+10Ab)\sqrt{a+b\tan(c+dx)}}{5d\tan^{5/2}(c+dx)} + \frac{1}{5} \right) \frac{2(3}{$$

input `Int[Cot[c + d*x]^(9/2)*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*a*A*(a + b*Tan[c + d*x])^(3/2))/(7*d*Tan[c + d*x]^(7/2)) + ((-2*a*(10*A*b + 7*a*B)*Sqrt[a + b*Tan[c + d*x]])/(5*d*Tan[c + d*x]^(5/2)) + ((2*(35*a^2*A - 45*A*b^2 - 77*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) - ((-105*((a^2*(a + I*b)^3*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a - b]*d) + (a^2*(a - I*b)^3*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d)))/a - (2*(245*a^2*A*b - 15*A*b^3 + 105*a^3*B - 161*a*b^2*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/(3*a))/5)/7`

3.629.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4088 `Int[(((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && ! (ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`


```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.629.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2681 vs. $2(295) = 590$.

Time = 9.66 (sec) , antiderivative size = 2682, normalized size of antiderivative = 7.68

method	result	size
derivativedivides	Expression too large to display	2682
default	Expression too large to display	2682

```
input int(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RET
URNVERBOSE)
```

```
output -1/420/d*((b+a*cot(d*x+c))/cot(d*x+c))^(1/2)*cot(d*x+c)^(1/2)*(120*A*b^3*(
b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)-1260*A*arctan((2*(b+a*
cot(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*b)^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(
1/2))*a^2*b^2+1260*B*arctan(((2*(a^2+b^2)^(1/2)+2*b)^(1/2)-2*(b+a*cot(d*x+
c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*a^3*b-420*B*arctan(((2*(a^2+b^2)
^(1/2)+2*b)^(1/2)-2*(b+a*cot(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))
*a*b^3-1260*B*arctan((2*(b+a*cot(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*b)^(1/
2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*a^3*b+420*B*arctan((2*(b+a*cot(d*x+c))^(
1/2)+(2*(a^2+b^2)^(1/2)+2*b)^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*a*b^3-
420*B*arctan(((2*(a^2+b^2)^(1/2)+2*b)^(1/2)-2*(b+a*cot(d*x+c))^(1/2))/(2*(
a^2+b^2)^(1/2)-2*b)^(1/2))*(a^2+b^2)^(1/2)*a^3+420*B*arctan((2*(b+a*cot(d*
x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*b)^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*
(a^2+b^2)^(1/2)*a^3-840*B*a^3*(b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)-2*
b)^(1/2)+1260*A*arctan(((2*(a^2+b^2)^(1/2)+2*b)^(1/2)-2*(b+a*cot(d*x+c))^(
1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*a^2*b^2+120*A*a^3*(b+a*cot(d*x+c))^(1
/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*cot(d*x+c)^3+168*B*a^3*(b+a*cot(d*x+c))^(
1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*cot(d*x+c)^2-280*A*a^3*(b+a*cot(d*x+c)
)^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*cot(d*x+c)+105*A*ln((b+a*cot(d*x+c))
^(1/2)*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)-a*cot(d*x+c)-b-(a^2+b^2)^(1/2))*(2*(a
^2+b^2)^(1/2)+2*b)^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*b^3-105*A*ln(a*c...
```

3.629.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18996 vs. $2(289) = 578$.

Time = 4.97 (sec) , antiderivative size = 18996, normalized size of antiderivative = 54.43

$$\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.629.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(9/2)*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output Timed out

3.629.7 Maxima [F]

$$\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{5}{2}} \cot(dx + c)^{\frac{9}{2}} dx$$

input `integrate(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^(9/2), x)`

3.629. $\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.629.8 Giac [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(9/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.629.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{9}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{9/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2} dx$$

input `int(cot(c + d*x)^(9/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)`

output `int(cot(c + d*x)^(9/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2), x)`

3.630 $\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.630.1 Optimal result	6029
3.630.2 Mathematica [A] (verified)	6030
3.630.3 Rubi [A] (verified)	6030
3.630.4 Maple [B] (verified)	6035
3.630.5 Fricas [B] (verification not implemented)	6036
3.630.6 Sympy [F(-1)]	6037
3.630.7 Maxima [F]	6037
3.630.8 Giac [F(-1)]	6037
3.630.9 Mupad [F(-1)]	6038

3.630.1 Optimal result

Integrand size = 35, antiderivative size = 287

$$\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx =$$

$$\frac{(ia-b)^{5/2}(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{d}$$

$$+ \frac{(ia+b)^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{d}$$

$$+ \frac{2(15a^2A-23Ab^2-35abB) \sqrt{\cot(c+dx)}\sqrt{a+b \tan(c+dx)}}{15d}$$

$$- \frac{2a(8Ab+5aB) \cot^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}}{15d}$$

$$- \frac{2aA \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}}{5d}$$

output

```
-(I*a-b)^(5/2)*(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(I*a+b)^(5/2)*(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-2/15*a*(8*A*b+5*B*a)*cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(1/2)/d+2/15*(15*A*a^2-23*A*b^2-35*B*a*b)*cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/d-2/5*a*A*cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(3/2)/d
```

3.630. $\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.630.2 Mathematica [A] (verified)

Time = 3.10 (sec) , antiderivative size = 321, normalized size of antiderivative = 1.12

$$\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \frac{\cot^{\frac{5}{2}}(c+dx) \left(60\sqrt[4]{-1} \left((-a+ib)^{5/2}(A-iB) \arctan \left(\frac{\sqrt[4]{-1} \sqrt{-a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right) \right) + (a+B \tan(c+dx)) \right)}{60d}$$

input `Integrate[Cot[c + d*x]^(7/2)*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(Cot[c + d*x]^(5/2)*(60*(-1)^(1/4)*((-a + I*b)^(5/2)*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]) + (a + I*b)^(5/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(5/2) + 15*b*(-2*A*b + a*B)*Sqrt[a + b*Tan[c + d*x]] - 3*(8*a^2*A - 10*A*b^2 - 15*a*b*B)*Sqrt[a + b*Tan[c + d*x]] - 4*(22*a*A*b + 10*a^2*B - 15*b^2*B)*Tan[c + d*x]*Sqrt[a + b*Tan[c + d*x]] + 8*(15*a^2*A - 23*A*b^2 - 35*a*b*B)*Tan[c + d*x]^2*Sqrt[a + b*Tan[c + d*x]] - 60*b*B*(a + b*Tan[c + d*x])^(3/2)))/(60*d)`

3.630.3 Rubi [A] (verified)

Time = 2.05 (sec) , antiderivative size = 304, normalized size of antiderivative = 1.06, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.514$, Rules used = {3042, 4729, 3042, 4088, 27, 3042, 4128, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{7/2}(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$$

$$\downarrow \text{4729}$$

3.630. $\int \cot^{\frac{7}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\tan^{7/2}(c+dx)} dx$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\tan(c+dx)^{7/2}} dx$$

↓ 4088

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2}{5} \int \frac{\sqrt{a+b\tan(c+dx)}(-b(2aA-5bB)\tan^2(c+dx)-5(Aa^2-2bBa-Ab^2)\tan(c+dx))}{2\tan^{5/2}(c+dx)} dx \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \int \frac{\sqrt{a+b\tan(c+dx)}(-b(2aA-5bB)\tan^2(c+dx)-5(Aa^2-2bBa-Ab^2)\tan(c+dx))}{\tan^{5/2}(c+dx)} dx \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \int \frac{\sqrt{a+b\tan(c+dx)}(-b(2aA-5bB)\tan(c+dx)^2-5(Aa^2-2bBa-Ab^2)\tan(c+dx))}{\tan(c+dx)^{5/2}} dx \right)$$

↓ 4128

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \left(\frac{2}{3} \int -\frac{b(10Ba^2+22Aba-15b^2B)\tan^2(c+dx)+15(Ba^3+3Aba^2-3b^2Ba-Ab^3)-a(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx)}{2\tan^{3/2}(c+dx)\sqrt{a+b\tan(c+dx)}} dx \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \left(-\frac{1}{3} \int \frac{b(10Ba^2+22Aba-15b^2B)\tan^2(c+dx)+15(Ba^3+3Aba^2-3b^2Ba-Ab^3)-a(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx)}{\tan^{3/2}(c+dx)\sqrt{a+b\tan(c+dx)}} dx \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \left(-\frac{1}{3} \int \frac{b(10Ba^2+22Aba-15b^2B)\tan(c+dx)^2+15(Ba^3+3Aba^2-3b^2Ba-Ab^3)-a(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx)}{\tan(c+dx)^{3/2}\sqrt{a+b\tan(c+dx)}} dx \right) \right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{5} \left(\frac{1}{3} \left(\frac{2 \int -\frac{15(a(Ba^3+3Aba^2-3b^2Ba-Ab^3))-a(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a} + \frac{2(15(b(10Ba^2+22Aba-15b^2B)\tan(c+dx)^2+15(Ba^3+3Aba^2-3b^2Ba-Ab^3)-a(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx))}{3a} \right) \right) \right)$$

3.630. $\int \cot^{7/2}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\left(\frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-\frac{15\int\frac{a(Ba^3+3Aba^2-3b^2Ba-}{\sqrt{\tan(c+dx)}}dx}{\sqrt{\tan(c+dx)}}}\right)\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{5}\left(\frac{1}{3}\left(\frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-\frac{15\int\frac{a(Ba^3+3Aba^2-3b^2Ba-}{\sqrt{\tan(c+dx)}}dx}{\sqrt{\tan(c+dx)}}}\right)\right)\right)$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{5d\tan^{5/2}(c+dx)}+\frac{1}{5}\left(-\frac{2a(5aB+8Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)}+\frac{1}{3}\left(\frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-\frac{15\int\frac{a(Ba^3+3Aba^2-3b^2Ba-}{\sqrt{\tan(c+dx)}}dx}{\sqrt{\tan(c+dx)}}}\right)\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{5d\tan^{5/2}(c+dx)}+\frac{1}{5}\left(-\frac{2a(5aB+8Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)}+\frac{1}{3}\left(\frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-\frac{15\int\frac{a(Ba^3+3Aba^2-3b^2Ba-}{\sqrt{\tan(c+dx)}}dx}{\sqrt{\tan(c+dx)}}}\right)\right)\right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{5d\tan^{5/2}(c+dx)}+\frac{1}{5}\left(-\frac{2a(5aB+8Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)}+\frac{1}{3}\left(\frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-\frac{15\int\frac{a(Ba^3+3Aba^2-3b^2Ba-}{\sqrt{\tan(c+dx)}}dx}{\sqrt{\tan(c+dx)}}}\right)\right)\right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{5d\tan^{5/2}(c+dx)}+\frac{1}{5}\left(-\frac{2a(5aB+8Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)}+\frac{1}{3}\left(\frac{2(15a^2A-35abB-23Ab^2)\sqrt{a+b\tan(c+dx)}}{d\sqrt{\tan(c+dx)}}-\frac{15\int\frac{a(Ba^3+3Aba^2-3b^2Ba-}{\sqrt{\tan(c+dx)}}dx}{\sqrt{\tan(c+dx)}}}\right)\right)\right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{5d\tan^{5/2}(c+dx)}+\frac{1}{5}\left(-\frac{2a(5aB+8Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)}+\frac{1}{3}\left(\frac{2(15}{\right.\right.\right.$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2aA(a+b\tan(c+dx))^{3/2}}{5d\tan^{5/2}(c+dx)}+\frac{1}{5}\left(-\frac{2a(5aB+8Ab)\sqrt{a+b\tan(c+dx)}}{3d\tan^{3/2}(c+dx)}+\frac{1}{3}\left(\frac{2(15}{\right.\right.\right.$$

input `Int[Cot[c + d*x]^(7/2)*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*a*A*(a + b*Tan[c + d*x])^(3/2))/(5*d*Tan[c + d*x]^(5/2)) + ((-2*a*(8*A*b + 5*a*B)*Sqrt[a + b*Tan[c + d*x]])/(3*d*Tan[c + d*x]^(3/2)) + ((-15*(-((a*(a + I*b)^3*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a - b]*d)) + (a*(a - I*b)^3*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a + b]*d)))/a + (2*(15*a^2*A - 23*A*b^2 - 35*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]))/3)/5`

3.630.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 104 `Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

```
rule 4128 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.630.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2517 vs. $2(239) = 478$.

Time = 1.62 (sec) , antiderivative size = 2518, normalized size of antiderivative = 8.77

method	result	size
derivativedivides	Expression too large to display	2518
default	Expression too large to display	2518

input `int(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RETURNVERBOSE)`

output `1/60/d*((b+a*cot(d*x+c))/cot(d*x+c))^(1/2)*cot(d*x+c)^(1/2)*(-24*A*a^3*(b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*cot(d*x+c)^2-40*B*a^3*(b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*cot(d*x+c)+15*A*ln((b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)-a*cot(d*x+c)-b-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*a^3-15*A*ln(a*cot(d*x+c)+b+(b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*a^3+15*B*ln((b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)-a*cot(d*x+c)-b-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*b^3-15*B*ln(a*cot(d*x+c)+b+(b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*b^3-60*A*arctan(((2*(a^2+b^2)^(1/2)+2*b)^(1/2)-2*(b+a*cot(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*(a^2+b^2)^(1/2)*a*b^2+60*A*arctan((2*(b+a*cot(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*b)^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*(a^2+b^2)^(1/2)*a*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)-120*B*arctan(((2*(a^2+b^2)^(1/2)+2*b)^(1/2)-2*(b+a*cot(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*(a^2+b^2)^(1/2)*a^2*b+120*B*arctan((2*(b+a*cot(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*b)^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*(a^2+b^2)^(1/2)*a^2*b-280*B*a^2*b*(b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)-180*A*arctan(((2*(a...`

3.630.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18942 vs. $2(233) = 466$.

Time = 4.95 (sec) , antiderivative size = 18942, normalized size of antiderivative = 66.00

$$\int \cot^{\frac{7}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,algorithm="fricas")`

output Too large to include

3.630. $\int \cot^{\frac{7}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$

3.630.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(7/2)*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.630.7 Maxima [F]

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{5}{2}} \cot(dx + c)^{\frac{7}{2}} dx$$

input `integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^(7/2), x)`

3.630.8 Giac [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(7/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.630.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{7}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{7/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2} dx$$

input `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)`output `int(cot(c + d*x)^(7/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2), x)`

3.631 $\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{\frac{5}{2}}(A+B \tan(c+dx)) dx$

3.631.1 Optimal result	6039
3.631.2 Mathematica [C] (verified)	6040
3.631.3 Rubi [A] (verified)	6040
3.631.4 Maple [B] (verified)	6044
3.631.5 Fricas [B] (verification not implemented)	6045
3.631.6 Sympy [F(-1)]	6046
3.631.7 Maxima [F]	6046
3.631.8 Giac [F(-1)]	6046
3.631.9 Mupad [F(-1)]	6047

3.631.1 Optimal result

Integrand size = 35, antiderivative size = 300

$$\int \cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{\frac{5}{2}}(A+B \tan(c+dx)) dx =$$

$$\frac{(ia-b)^{\frac{5}{2}}(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{d}$$

$$+ \frac{2b^{\frac{5}{2}}B \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{d}$$

$$- \frac{(ia+b)^{\frac{5}{2}}(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{d}$$

$$- \frac{2a(2Ab+aB)\sqrt{\cot(c+dx)}\sqrt{a+b \tan(c+dx)}}{d}$$

$$- \frac{2aA \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{\frac{3}{2}}}{3d}$$

```
output -(I*a-b)^(5/2)*(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+2*b^(5/2)*B*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-(I*a+b)^(5/2)*(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-2*a*(2*A*b+B*a)*cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/d-2/3*a*A*cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(3/2)/d
```

3.631.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 5.72 (sec) , antiderivative size = 417, normalized size of antiderivative = 1.39

$$\int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx = \frac{\cot^{\frac{3}{2}}(c+dx) \left(iab(A+iB) \operatorname{Hypergeometric2F1} \left(-\frac{3}{2}, -\frac{3}{2}, -\frac{1}{2}, -\frac{b\tan(c+dx)}{a} \right) \sqrt{a+b\tan(c+dx)} \right)}{\dots}$$

input `Integrate[Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(Cot[c + d*x]^(3/2)*(I*a*b*(A + I*B)*Hypergeometric2F1[-3/2, -3/2, -1/2, -((b*Tan[c + d*x])/a)]*Sqrt[a + b*Tan[c + d*x]] - a*b*(I*A + B)*Hypergeometric2F1[-3/2, -3/2, -1/2, -((b*Tan[c + d*x])/a)]*Sqrt[a + b*Tan[c + d*x]] + (I*a + b)*(A - I*B)*Sqrt[1 + (b*Tan[c + d*x])/a]*(3*(-1)^(1/4)*(-a + I*b)^(3/2)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(3/2) + (I*a + (-3*a + (4*I)*b)*Tan[c + d*x])*Sqrt[a + b*Tan[c + d*x]]) + (I*a - b)*(A + I*B)*Sqrt[1 + (b*Tan[c + d*x])/a]*(3*(-1)^(1/4)*(a + I*b)^(3/2)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]])*Tan[c + d*x]^(3/2) + (I*a + (3*a + (4*I)*b)*Tan[c + d*x])*Sqrt[a + b*Tan[c + d*x]])))/(3*d*Sqrt[1 + (b*Tan[c + d*x])/a])`

3.631.3 Rubi [A] (verified)

Time = 1.69 (sec) , antiderivative size = 262, normalized size of antiderivative = 0.87, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 4729, 3042, 4088, 27, 3042, 4128, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$$

↓ 3042

3.631. $\int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$

$$\begin{aligned}
& \int \cot(c+dx)^{5/2}(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx \\
& \quad \downarrow 4729 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan^{5/2}(c+dx)} dx \\
& \quad \downarrow 3042 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\tan(c+dx)^{5/2}} dx \\
& \quad \downarrow 4088 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \left(\frac{2}{3} \int \frac{3\sqrt{a+b \tan(c+dx)}(b^2 B \tan^2(c+dx) - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a)}{2 \tan^{3/2}(c+dx)} dx \right. \\
& \quad \downarrow 27 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \left(\int \frac{\sqrt{a+b \tan(c+dx)}(b^2 B \tan^2(c+dx) - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2A - B))}{\tan^{3/2}(c+dx)} dx \right. \\
& \quad \downarrow 3042 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \left(\int \frac{\sqrt{a+b \tan(c+dx)}(b^2 B \tan(c+dx)^2 - (Aa^2 - 2bBa - Ab^2) \tan(c+dx) + a(2A - B))}{\tan(c+dx)^{3/2}} dx \right. \\
& \quad \downarrow 4128 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \left(2 \int \frac{-B \tan^2(c+dx)b^3 + a(Aa^2 - 3bBa - 3Ab^2) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx \right. \\
& \quad \downarrow 27 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \left(- \int \frac{-B \tan^2(c+dx)b^3 + a(Aa^2 - 3bBa - 3Ab^2) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx \right. \\
& \quad \downarrow 3042 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \left(- \int \frac{-B \tan(c+dx)^2 b^3 + a(Aa^2 - 3bBa - 3Ab^2) + (Ba^3 + 3Aba^2 - 3b^2Ba - Ab^3)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx \right. \\
& \quad \downarrow 4138
\end{aligned}$$

3.631. $\int \cot^{5/2}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{\int\frac{-B\tan^2(c+dx)b^3+a(Aa^2-3bBa-3Ab^2)+(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}d\tan(c+dx)}{d}\right)$$

↓ 2035

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2\int\frac{-B\tan^2(c+dx)b^3+a(Aa^2-3bBa-3Ab^2)+(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}d\sqrt{\tan(c+dx)}}{d}\right)$$

↓ 2257

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2\int\left(\frac{Aa^3-3bBa^2-3Ab^2a+b^3B+(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}-\frac{b^3B}{\sqrt{a+b\tan(c+dx)}}\right)d\sqrt{\tan(c+dx)}}{d}\right)$$

↓ 2009

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2\left(\frac{1}{2}(-b+ia)^{5/2}(-B+iA)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)+\frac{1}{2}(b+ia)^{5/2}(B+iA)\arctan\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)\right)}{d}\right)$$

input `Int[Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*((I*a - b)^(5/2)*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/2 - b^(5/2)*B*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) + ((I*a + b)^(5/2)*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/2))/d - (2*a*(2*A*b + a*B)*Sqrt[a + b*Tan[c + d*x]])/(d*Sqrt[Tan[c + d*x]]) - (2*a*A*(a + b*Tan[c + d*x])^(3/2))/(3*d*Tan[c + d*x]^(3/2))`

3.631.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2035 `Int[(F_x_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[Fx, x]`
- rule 2257 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

```
rule 4128 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e +
f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Sim
p[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e
+ f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*
(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b
*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x],
x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ
[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]
```

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.631.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2437 vs. $2(248) = 496$.

Time = 1.41 (sec) , antiderivative size = 2438, normalized size of antiderivative = 8.13

method	result	size
derivativedivides	Expression too large to display	2438
default	Expression too large to display	2438

```
input int(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RET
URNVERBOSE)
```

$$3.631. \quad \int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{\frac{5}{2}}(A + B \tan(c + dx)) dx$$

output $1/12/d*((b+a*\cot(d*x+c))/\cot(d*x+c))^{(1/2)}*\cot(d*x+c)^{(1/2)}*(-36*A*\arctan((2*(b+a*\cot(d*x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*b)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*b)^{(1/2)})*a^2*b^2+36*B*\arctan(((2*(a^2+b^2)^{(1/2)}+2*b)^{(1/2)}-2*(b+a*\cot(d*x+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*b)^{(1/2)})*a^3*b-12*B*\arctan(((2*(a^2+b^2)^{(1/2)}+2*b)^{(1/2)}-2*(b+a*\cot(d*x+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*b)^{(1/2)})*a*b^3-36*B*\arctan((2*(b+a*\cot(d*x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*b)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*b)^{(1/2)})*a^3*b+12*B*\arctan((2*(b+a*\cot(d*x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*b)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*b)^{(1/2)})*a*b^3-12*B*\arctan(((2*(a^2+b^2)^{(1/2)}+2*b)^{(1/2)}-2*(b+a*\cot(d*x+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*b)^{(1/2)})*(a^2+b^2)^{(1/2)}*a^3+12*B*\arctan((2*(b+a*\cot(d*x+c))^{(1/2)}+(2*(a^2+b^2)^{(1/2)}+2*b)^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*b)^{(1/2)})*(a^2+b^2)^{(1/2)}*a^3-24*B*a^3*(b+a*\cot(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}-2*b)^{(1/2)}+36*A*\arctan(((2*(a^2+b^2)^{(1/2)}+2*b)^{(1/2)}-2*(b+a*\cot(d*x+c))^{(1/2)})/(2*(a^2+b^2)^{(1/2)}-2*b)^{(1/2)})*a^2*b^2-8*A*a^3*(b+a*\cot(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}-2*b)^{(1/2)}*\cot(d*x+c)+3*A*\ln((b+a*\cot(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*b)^{(1/2)}-a*\cot(d*x+c)-b-(a^2+b^2)^{(1/2)})*(2*(a^2+b^2)^{(1/2)}+2*b)^{(1/2)}*(2*(a^2+b^2)^{(1/2)}-2*b)^{(1/2)}*b^3-3*A*\ln(a*\cot(d*x+c)+b+(b+a*\cot(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*b)^{(1/2)}+(a^2+b^2)^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*b)^{(1/2)}*(2*(a^2+b^2)^{(1/2)}-2*b)^{(1/2)}*b^3-3*B*\ln((b+a*\cot(d*x+c))^{(1/2)}*(2*(a^2+b^2)^{(1/2)}+2*b)^{(1/2)}-a*\cot(d*x+c)-b-(a^2+b^2)^{(1/2)}...$

3.631.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17767 vs. $2(242) = 484$.

Time = 7.26 (sec) , antiderivative size = 35564, normalized size of antiderivative = 118.55

$$\int \cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algo rithm="fracas")`

output Too large to include

3.631.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(5/2)*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.631.7 Maxima [F]

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^{\frac{5}{2}} \cot(dx + c)^{\frac{5}{2}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^(5/2), x)`

3.631.8 Giac [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.631. $\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.631.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{5/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2} dx$$

input `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)`output `int(cot(c + d*x)^(5/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2), x)`

3.632 $\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

3.632.1 Optimal result	6048
3.632.2 Mathematica [A] (verified)	6049
3.632.3 Rubi [A] (verified)	6049
3.632.4 Maple [B] (verified)	6053
3.632.5 Fricas [B] (verification not implemented)	6054
3.632.6 Sympy [F(-1)]	6055
3.632.7 Maxima [F(-1)]	6055
3.632.8 Giac [F(-1)]	6055
3.632.9 Mupad [F(-1)]	6056

3.632.1 Optimal result

Integrand size = 35, antiderivative size = 301

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx = \frac{(ia-b)^{5/2}(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d} + \frac{b^{3/2}(2Ab+5aB) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d} - \frac{(ia+b)^{5/2}(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d} + \frac{b(2aA+bB)\sqrt{a+b \tan(c+dx)}}{d\sqrt{\cot(c+dx)}} - \frac{2aA\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^{3/2}}{d}$$

```
output (I*a-b)^(5/2)*(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+b^(3/2)*(2*A*b+5*B*a)*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d-(I*a+b)^(5/2)*(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+b*(2*A*a+B*b)*(a+b*tan(d*x+c))^(1/2)/d/cot(d*x+c)^(1/2)-2*a*A*cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(3/2)/d
```

3.632.2 Mathematica [A] (verified)

Time = 3.40 (sec) , antiderivative size = 363, normalized size of antiderivative = 1.21

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx = \frac{\sqrt{\cot(c+dx)} \left((-1)^{3/4}(-a+ib)^{5/2}(iA+B) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) \sqrt{\tan(c+dx)} \right)}{\sqrt{\cot(c+dx)}} + B \tan(c+dx)$$

input `Integrate[Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*((-1)^(3/4)*(-a + I*b)^(5/2)*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]*Sqrt[Tan[c + d*x]] - (-1)^(1/4)*(a + I*b)^(5/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]*Sqrt[Tan[c + d*x]] - (a - I*b)^2*(A - I*B)*Sqrt[a + b*Tan[c + d*x]] - (a + I*b)^2*(A + I*B)*Sqrt[a + b*Tan[c + d*x]] + b*B*(a + b*Tan[c + d*x])^(3/2) + (b*(2*A*b + 5*a*B)*Sqrt[a + b*Tan[c + d*x]]*(Sqrt[b]*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[Tan[c + d*x]] - Sqrt[a]*Sqrt[1 + (b*Tan[c + d*x])/a]))/(Sqrt[a]*Sqrt[1 + (b*Tan[c + d*x])/a]))/d`

3.632.3 Rubi [A] (verified)Time = 1.73 (sec) , antiderivative size = 258, normalized size of antiderivative = 0.86, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 4729, 3042, 4088, 27, 3042, 4130, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{3/2}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$$

$$\downarrow \text{4729}$$

3.632. $\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\tan^{3/2}(c+dx)} dx$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\tan(c+dx)^{3/2}} dx$$

↓ 4088

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(2 \int \frac{\sqrt{a+b\tan(c+dx)}(b(2aA+bB)\tan^2(c+dx) - (Aa^2 - 2bBa - Ab^2)\tan(c+dx))}{2\sqrt{\tan(c+dx)}} dx \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{\sqrt{a+b\tan(c+dx)}(b(2aA+bB)\tan^2(c+dx) - (Aa^2 - 2bBa - Ab^2)\tan(c+dx))}{\sqrt{\tan(c+dx)}} dx \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{\sqrt{a+b\tan(c+dx)}(b(2aA+bB)\tan(c+dx)^2 - (Aa^2 - 2bBa - Ab^2)\tan(c+dx))}{\sqrt{\tan(c+dx)}} dx \right)$$

↓ 4130

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{b^2(2Ab+5aB)\tan^2(c+dx) - 2(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)\tan(c+dx) + a(2Ba^2 + 6Aba - b^2B)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \int \frac{b^2(2Ab+5aB)\tan^2(c+dx) - 2(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)\tan(c+dx) + a(2Ba^2 + 6Aba - b^2B)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \int \frac{b^2(2Ab+5aB)\tan(c+dx)^2 - 2(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)\tan(c+dx) + a(2Ba^2 + 6Aba - b^2B)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \right)$$

↓ 4138

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{b^2(2Ab+5aB)\tan^2(c+dx) - 2(Aa^3 - 3bBa^2 - 3Ab^2a + b^3B)\tan(c+dx) + a(2Ba^2 + 6Aba - b^2B)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} d\tan(c+dx)}{2d} \right)$$

3.632. $\int \cot^{3/2}(c+dx)(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$

$$\begin{aligned} & \downarrow \text{2035} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{b^2(2Ab+5aB)\tan^2(c+dx)-2(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx)+a(2Ba^2+6Aba-b^2B)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} d\sqrt{\tan(c+dx)}}{d} \right) \\ & \downarrow \text{2257} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \left(\frac{(2Ab+5aB)b^2}{\sqrt{a+b\tan(c+dx)}} + \frac{2(Ba^3+3Aba^2-3b^2Ba-Ab^3-(Aa^3-3bBa^2-3Ab^2a+b^3B)\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} \right) d\sqrt{\tan(c+dx)}}{d} \right) \\ & \downarrow \text{2009} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-b+ia)^{5/2}(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) + b^{3/2}(5aB+2Ab)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d} \right) \end{aligned}$$

input `Int[Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((I*a - b)^(5/2)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + b^(3/2)*(2*A*b + 5*a*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] - (I*a + b)^(5/2)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/d + (b*(2*a*A + b*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d - (2*a*A*(a + b*Tan[c + d*x])^(3/2))/(d*Sqrt[Tan[c + d*x]]))`

3.632.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2035 `Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]`

rule 2257 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol
] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a
, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x
)^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2))
Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*
(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n
+ 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[
e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n
+ 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] &&
NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] &
& LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4130 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_)
+ (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_)
) + (f_)*(x_)^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[
e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a
+ b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C
*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*
m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] &&
NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[
c, 0] && NeQ[a, 0])))`

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.632.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2667 vs. $2(251) = 502$.

Time = 0.44 (sec) , antiderivative size = 2668, normalized size of antiderivative = 8.86

method	result	size
derivativedivides	Expression too large to display	2668
default	Expression too large to display	2668

```
input int(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RET
URNVERBOSE)
```

output

```

-1/4/d*((b+a*cot(d*x+c))/cot(d*x+c))^(1/2)/cot(d*x+c)^(1/2)*(4*A*arctan(((
2*(a^2+b^2)^(1/2)+2*b)^(1/2)-2*(b+a*cot(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-
2*b)^(1/2))*b^(7/2)*a*cot(d*x+c)-4*A*arctan((2*(b+a*cot(d*x+c))^(1/2)+(2*(
a^2+b^2)^(1/2)+2*b)^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*b^(7/2)*a*cot(d*
x+c)+12*B*arctan(((2*(a^2+b^2)^(1/2)+2*b)^(1/2)-2*(b+a*cot(d*x+c))^(1/2))/
(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*b^(5/2)*a^2*cot(d*x+c)-12*B*arctan((2*(b+a*
cot(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*b)^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(
1/2))*b^(5/2)*a^2*cot(d*x+c)-12*A*arctan(((2*(a^2+b^2)^(1/2)+2*b)^(1/2)-2*
(b+a*cot(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*b^(3/2)*a^3*cot(d*x
+c)+12*A*arctan((2*(b+a*cot(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*b)^(1/2))/(
2*(a^2+b^2)^(1/2)-2*b)^(1/2))*b^(3/2)*a^3*cot(d*x+c)-4*B*(2*(a^2+b^2)^(1/2
)-2*b)^(1/2)*(b+a*cot(d*x+c))^(1/2)*b^(5/2)*a-4*B*arctan(((2*(a^2+b^2)^(1/
2)+2*b)^(1/2)-2*(b+a*cot(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*b^(
1/2)*a^4*cot(d*x+c)+4*B*arctan((2*(b+a*cot(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2
)+2*b)^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*b^(1/2)*a^4*cot(d*x+c)-3*A*ln
((b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)-a*cot(d*x+c)-b-(a^2+
b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*b^(
5/2)*a*cot(d*x+c)+3*A*ln(a*cot(d*x+c)+b+(b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^
2)^(1/2)+2*b)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)*(2*(a^2
+b^2)^(1/2)-2*b)^(1/2))*b^(5/2)*a*cot(d*x+c)-B*ln((b+a*cot(d*x+c))^(1/2)...

```

3.632.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17670 vs. $2(245) = 490$.

Time = 7.58 (sec) , antiderivative size = 35373, normalized size of antiderivative = 117.52

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algo
rithm="fricas")`

output Too large to include

3.632.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(3/2)*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.632.7 Maxima [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorith="maxima")`

output `Timed out`

3.632.8 Giac [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorith="giac")`

output `Timed out`

3.632.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \int \cot(c + dx)^{3/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2} dx$$

input `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2),x)`output `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2), x)`

3.633 $\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx$

3.633.1 Optimal result	6057
3.633.2 Mathematica [A] (verified)	6058
3.633.3 Rubi [A] (verified)	6058
3.633.4 Maple [B] (verified)	6062
3.633.5 Fricas [B] (verification not implemented)	6063
3.633.6 Sympy [F(-1)]	6063
3.633.7 Maxima [F]	6063
3.633.8 Giac [F(-1)]	6064
3.633.9 Mupad [F(-1)]	6064

3.633.1 Optimal result

Integrand size = 35, antiderivative size = 320

$$\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx)) dx = \frac{(ia - b)^{5/2}(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d} + \frac{\sqrt{b}(20aAb + 15a^2B - 8b^2B) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{4d} + \frac{(ia + b)^{5/2}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d} + \frac{b(4Ab + 7aB)\sqrt{a + b \tan(c + dx)}}{4d\sqrt{\cot(c + dx)}} + \frac{bB(a + b \tan(c + dx))^{3/2}}{2d\sqrt{\cot(c + dx)}}$$

```
output (I*a-b)^(5/2)*(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(I*a+b)^(5/2)*(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+1/4*(20*A*a*b+15*B*a^2-8*B*b^2)*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*b^(1/2)*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+1/4*b*(4*A*b+7*B*a)*(a+b*tan(d*x+c))^(1/2)/d/cot(d*x+c)^(1/2)+1/2*b*B*(a+b*tan(d*x+c))^(3/2)/d/cot(d*x+c)^(1/2)
```


3.633.2 Mathematica [A] (verified)

Time = 3.77 (sec) , antiderivative size = 311, normalized size of antiderivative = 0.97

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))dx = \frac{\sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}\left(4\sqrt[4]{-1}(-a+ib)^{5/2}(iA+B)\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)+B\tan(c+dx)\right)}{4d}$$

input `Integrate[Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(4*(-1)^(1/4)*(-a + I*b)^(5/2)*(I*A + B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]] - 4*(-1)^(3/4)*(a + I*b)^(5/2)*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]] + b*(4*A*b + 7*a*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]] + 2*b*B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2) + (Sqrt[a]*Sqrt[b]*(20*a*A*b + 15*a^2*B - 8*b^2*B)*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/Sqrt[a + b*Tan[c + d*x]]))/(4*d)`

3.633.3 Rubi [A] (verified)Time = 1.76 (sec) , antiderivative size = 278, normalized size of antiderivative = 0.87, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 4729, 3042, 4090, 27, 3042, 4130, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))dx \\ & \quad \downarrow \text{4729} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\sqrt{\tan(c+dx)}}dx \end{aligned}$$

$$3.633. \quad \int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))dx$$

$$\begin{aligned} & \downarrow 3042 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx \\ & \downarrow 4090 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \int \frac{\sqrt{a+b \tan(c+dx)}(b(4Ab+7aB) \tan^2(c+dx) + 4(Ba^2+2Aba-b^2B) \tan(c+dx) + a)}{2\sqrt{\tan(c+dx)}} dx \right) \\ & \downarrow 27 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \int \frac{\sqrt{a+b \tan(c+dx)}(b(4Ab+7aB) \tan^2(c+dx) + 4(Ba^2+2Aba-b^2B) \tan(c+dx) + a)}{\sqrt{\tan(c+dx)}} dx \right) \\ & \downarrow 3042 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \int \frac{\sqrt{a+b \tan(c+dx)}(b(4Ab+7aB) \tan(c+dx)^2 + 4(Ba^2+2Aba-b^2B) \tan(c+dx) + a)}{\sqrt{\tan(c+dx)}} dx \right) \\ & \downarrow 4130 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \left(\int \frac{b(15Ba^2+20Aba-8b^2B) \tan^2(c+dx) + 8(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx) + a}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx \right) \right) \\ & \downarrow 27 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \left(\frac{1}{2} \int \frac{b(15Ba^2+20Aba-8b^2B) \tan^2(c+dx) + 8(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx) + a}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx \right) \right) \\ & \downarrow 3042 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \left(\frac{1}{2} \int \frac{b(15Ba^2+20Aba-8b^2B) \tan(c+dx)^2 + 8(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx) + a}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx \right) \right) \\ & \downarrow 4138 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{4} \left(\int \frac{b(15Ba^2+20Aba-8b^2B) \tan^2(c+dx) + 8(Ba^3+3Aba^2-3b^2Ba-Ab^3) \tan(c+dx) + a(8Aa^2-9bBa-4b^2)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}(\tan^2(c+dx)+1)} dx \right) \right) \\ & \downarrow 2035 \end{aligned}$$

3.633. $\int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx)) dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{\int\frac{b(15Ba^2+20Aba-8b^2B)\tan^2(c+dx)+8(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx)+a(8Aa^2-9bBa-4a^2)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}dx}{d}\right)\right)$$

↓ 2257

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{4}\left(\frac{\int\left(\frac{b(15Ba^2+20Aba-8b^2B)}{\sqrt{a+b\tan(c+dx)}}+\frac{8(Aa^3-3bBa^2-3Ab^2a+b^3B+(Ba^3+3Aba^2-3b^2Ba-Ab^3)\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}\right)dx}{d}\right)\right)$$

↓ 2009

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{bB\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d}+\frac{1}{4}\left(\frac{b(7aB+4Ab)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{d}\right)\right)$$

input `Int[Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((b*B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2))/(2*d) + ((4*(I*a - b)^(5/2)*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + Sqrt[b]*(20*a*A*b + 15*a^2*B - 8*b^2*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]] + 4*(I*a + b)^(5/2)*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/d + (b*(4*A*b + 7*a*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d)/4`

3.633.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2035 `Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[Fx, x]`

3.633. $\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))dx$

rule 2257 `Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4090 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4130 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4138 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)] + (f_)*(x_)^2)^(n_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)] + (C_)*tan[(e_) + (f_)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.633.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2818 vs. $2(264) = 528$.

Time = 0.51 (sec) , antiderivative size = 2819, normalized size of antiderivative = 8.81

method	result	size
derivativedivides	Expression too large to display	2819
default	Expression too large to display	2819

```
input int(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x,method=_RET
URNVERBOSE)
```

```
output 1/4/d*((b+a*cot(d*x+c))/cot(d*x+c))^(1/2)/cot(d*x+c)^(3/2)*(2*B*(2*(a^2+b^
2)^(1/2)-2*b)^(1/2)*(b+a*cot(d*x+c))^(1/2)*b^(5/2)*a-8*A*arctan((2*(b+a*co
t(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*b)^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/
2))*(a^2+b^2)^(1/2)*b^(3/2)*a^2*cot(d*x+c)^2-4*B*arctan(((2*(a^2+b^2)^(1/2
)+2*b)^(1/2)-2*(b+a*cot(d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*(a^2
+b^2)^(1/2)*b^(5/2)*a*cot(d*x+c)^2+4*B*arctan((2*(b+a*cot(d*x+c))^(1/2)+(2
*(a^2+b^2)^(1/2)+2*b)^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*(a^2+b^2)^(1/2
)*b^(5/2)*a*cot(d*x+c)^2-A*ln((b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*
b)^(1/2)-a*cot(d*x+c)-b-(a^2+b^2)^(1/2))*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)*(2*
(a^2+b^2)^(1/2)-2*b)^(1/2)*b^(7/2)*cot(d*x+c)^2+A*ln(a*cot(d*x+c)+b+(b+a*c
ot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)+(a^2+b^2)^(1/2))*(2*(a^2+b^
2)^(1/2)+2*b)^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*b^(7/2)*cot(d*x+c)^2+8*A
*arctan(((2*(a^2+b^2)^(1/2)+2*b)^(1/2)-2*(b+a*cot(d*x+c))^(1/2))/(2*(a^2+b
^2)^(1/2)-2*b)^(1/2))*(a^2+b^2)^(1/2)*b^(3/2)*a^2*cot(d*x+c)^2+4*A*a*b^(5/
2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*(b+a*cot(d*x+c))^(1/2)*cot(d*x+c)+9*B*a^2
*b^(3/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*(b+a*cot(d*x+c))^(1/2)*cot(d*x+c)+2
0*A*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*arctanh((b+a*cot(d*x+c))^(1/2)/b^(1/2))*
a^2*b^2*cot(d*x+c)^2+4*B*arctan(((2*(a^2+b^2)^(1/2)+2*b)^(1/2)-2*(b+a*cot(
d*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*(a^2+b^2)^(1/2)*b^(1/2)*a^3*
cot(d*x+c)^2-4*B*arctan((2*(b+a*cot(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*...
```

3.633.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17723 vs. $2(258) = 516$.

Time = 7.67 (sec) , antiderivative size = 35483, normalized size of antiderivative = 110.88

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output Too large to include

3.633.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(1/2)*(a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c)),x)`

output Timed out

3.633.7 Maxima [F]

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx = \int (B\tan(dx+c)+A)(b\tan(dx+c)+a)^{5/2}\sqrt{\cot(dx+c)} dx$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x+c)+A)*(b*tan(d*x+c)+a)^(5/2)*sqrt(cot(d*x+c)),x)`

3.633. $\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx$

3.633.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `Timed out`

3.633.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx)) dx = \int \sqrt{\cot(c+dx)}(A+B\tan(c+dx))(a+b\tan(c+dx))^{5/2} dx$$

input `int(cot(c+d*x)^(1/2)*(A+B*tan(c+d*x))*(a+b*tan(c+d*x))^(5/2),x)`

output `int(cot(c+d*x)^(1/2)*(A+B*tan(c+d*x))*(a+b*tan(c+d*x))^(5/2),x)`

3.634
$$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.634.1 Optimal result 6065
 3.634.2 Mathematica [A] (verified) 6066
 3.634.3 Rubi [A] (verified) 6066
 3.634.4 Maple [B] (warning: unable to verify) 6071
 3.634.5 Fricas [B] (verification not implemented) 6071
 3.634.6 Sympy [F(-1)] 6071
 3.634.7 Maxima [F] 6072
 3.634.8 Giac [F(-1)] 6072
 3.634.9 Mupad [F(-1)] 6072

3.634.1 Optimal result

Integrand size = 35, antiderivative size = 376

$$\int \frac{(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx =$$

$$\frac{(ia - b)^{5/2}(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{(30a^2Ab - 16Ab^3 + 5a^3B - 40ab^2B) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{8\sqrt{bd}}$$

$$+ \frac{(ia + b)^{5/2}(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{(14aAb + 5a^2B - 8b^2B) \sqrt{a + b \tan(c + dx)}}{8d\sqrt{\cot(c + dx)}}$$

$$+ \frac{bB(a + b \tan(c + dx))^{3/2}}{3d \cot^{3/2}(c + dx)} + \frac{(2Ab + 3aB)(a + b \tan(c + dx))^{3/2}}{4d\sqrt{\cot(c + dx)}}$$

output

```
-(I*a-b)^(5/2)*(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+(I*a+b)^(5/2)*(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d+1/8*(30*A*a^2*b-16*A*b^3+5*B*a^3-40*B*a*b^2)*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/b^(1/2)+1/8*(14*A*a*b+5*B*a^2-8*B*b^2)*(a+b*tan(d*x+c))^(1/2)/d/cot(d*x+c)^(1/2)+1/3*b*B*(a+b*tan(d*x+c))^(3/2)/d/cot(d*x+c)^(3/2)+1/4*(2*A*b+3*B*a)*(a+b*tan(d*x+c))^(3/2)/d/cot(d*x+c)^(1/2)
```

3.634.
$$\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.634.2 Mathematica [A] (verified)

Time = 6.84 (sec) , antiderivative size = 365, normalized size of antiderivative = 0.97

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(24 \sqrt[4]{-1} (-a + ib)^{5/2} (A - iB) \right)}{\dots}$$

input `Integrate[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]], x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(24*(-1)^(1/4)*(-a + I*b)^(5/2)*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]] + 24*(-1)^(1/4)*(a + I*b)^(5/2)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]] + 3*(14*a*A*b + 5*a^2*B - 8*b^2*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]] + 6*(2*A*b + 3*a*B)*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2) + 8*b*B*Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2) + (3*Sqrt[a]*(30*a^2*A*b - 16*A*b^3 + 5*a^3*B - 40*a*b^2*B)*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/(Sqrt[b]*Sqrt[a + b*Tan[c + d*x]]))/(24*d)`

3.634.3 Rubi [A] (verified)

Time = 2.26 (sec) , antiderivative size = 346, normalized size of antiderivative = 0.92, number of steps used = 17, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.457$, Rules used = {3042, 4729, 3042, 4090, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

↓ 4729

3.634. $\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))dx$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))dx$$

↓ 4090

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{3} \int \frac{3}{2} \sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(b(2Ab+3aB)\tan^2(c+dx)+2(Ba^2+2Ab))dx \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \int \sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(b(2Ab+3aB)\tan^2(c+dx)+2(Ba^2+2Ab))dx \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \int \sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(b(2Ab+3aB)\tan(c+dx)^2+2(Ba^2+2Ab))dx \right)$$

↓ 4130

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \left(\frac{\int -\frac{\sqrt{a+b\tan(c+dx)}(-b(5Ba^2+14Aba-8b^2B)\tan^2(c+dx)-8b(Aa^2-2bBa-Ab^2)\tan(c+dx)+ab(2Aa^2+2Ab))}{2\sqrt{\tan(c+dx)}}}{2b} \right) \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \left(\frac{(3aB+2Ab)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d} - \frac{\int \frac{\sqrt{a+b\tan(c+dx)}(-b(5Ba^2+14Aba-8b^2B)\tan^2(c+dx)-8b(Aa^2-2bBa-Ab^2)\tan(c+dx)+ab(2Aa^2+2Ab))}{2\sqrt{\tan(c+dx)}}}{2b} \right) \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2} \left(\frac{(3aB+2Ab)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d} - \frac{\int \frac{\sqrt{a+b\tan(c+dx)}(-b(5Ba^2+14Aba-8b^2B)\tan^2(c+dx)-8b(Aa^2-2bBa-Ab^2)\tan(c+dx)+ab(2Aa^2+2Ab))}{2\sqrt{\tan(c+dx)}}}{2b} \right) \right)$$

↓ 4130

3.634. $\int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{(3aB+2Ab)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d}-\int\frac{-b(5Ba^3+30Aba^2-40b^2Ba-16Ab^3)}{\sqrt{a+b\tan(c+dx)}}dx\right)\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{(3aB+2Ab)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d}-\frac{1}{2}\int\frac{-b(5Ba^3+30Aba^2-40b^2Ba-16Ab^3)}{\sqrt{a+b\tan(c+dx)}}dx\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{(3aB+2Ab)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d}-\frac{1}{2}\int\frac{-b(5Ba^3+30Aba^2-40b^2Ba-16Ab^3)}{\sqrt{a+b\tan(c+dx)}}dx\right)\right)$$

↓ 4138

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{(3aB+2Ab)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d}-\int\frac{-b(5Ba^3+30Aba^2-40b^2Ba-16Ab^3)}{\sqrt{a+b\tan(c+dx)}}dx\right)\right)$$

↓ 2035

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{(3aB+2Ab)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d}-\int\frac{-b(5Ba^3+30Aba^2-40b^2Ba-16Ab^3)}{\sqrt{a+b\tan(c+dx)}}dx\right)\right)$$

↓ 2257

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\left(\frac{(3aB+2Ab)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d}-\int\frac{16(b(Ba^3+3Aba^2-3b^2Ba-Ab^3))}{\sqrt{a+b\tan(c+dx)}}dx\right)\right)$$

↓ 2009

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{bB\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}}{3d}+\frac{1}{2}\left(\frac{(3aB+2Ab)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}{2d}\right)\right)$$

input `Int[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((b*B*Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2))/(3*d) + (((2*A*b + 3*a*B)*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^(3/2))/(2*d) - ((8*(I*a - b)^(5/2)*b*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])]/Sqrt[a + b*Tan[c + d*x]]) - Sqrt[b]*(30*a^2*A*b - 16*A*b^3 + 5*a^3*B - 40*a*b^2*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]) - 8*b*(I*a + b)^(5/2)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/d - (b*(14*a*A*b + 5*a^2*B - 8*b^2*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/d)/(4*b))/2`

3.634.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2035 `Int[(F_x_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[F_x, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[F_x, x]`

rule 2257 `Int[(P_x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[P_x*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[P_x, x] && IntegerQ[p]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4138 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.634.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.42 (sec) , antiderivative size = 2657129, normalized size of antiderivative = 7066.83

output too large to display

input `int((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x)`

output `result too large to display`

3.634.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17750 vs. 2(308) = 616.

Time = 7.92 (sec) , antiderivative size = 35533, normalized size of antiderivative = 94.50

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="fricas")`

output `Too large to include`

3.634.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)**(1/2),x)`

output `Timed out`

3.634.7 Maxima [F]

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^{5/2}}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorith="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)/sqrt(cot(d*x + c)), x)`

3.634.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorith="giac")`

output `Timed out`

3.634.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2}}{\sqrt{\cot(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/cot(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/cot(c + d*x)^(1/2),x)`

3.634. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$

$$3.635 \quad \int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$$

3.635.1 Optimal result	6073
3.635.2 Mathematica [A] (verified)	6074
3.635.3 Rubi [A] (verified)	6075
3.635.4 Maple [B] (verified)	6080
3.635.5 Fricas [B] (verification not implemented)	6081
3.635.6 Sympy [F(-1)]	6081
3.635.7 Maxima [F]	6081
3.635.8 Giac [F(-1)]	6082
3.635.9 Mupad [F(-1)]	6082

3.635.1 Optimal result

Integrand size = 35, antiderivative size = 457

$$\int \frac{(a + b \tan(c + dx))^{5/2}(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{(ia - b)^{5/2}(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{(40a^3 Ab - 320aAb^3 - 5a^4 B - 240a^2 b^2 B + 128b^4 B) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{64b^{3/2}d}$$

$$- \frac{(ia + b)^{5/2}(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{d}$$

$$+ \frac{(40a^2 Ab - 64Ab^3 - 5a^3 B - 112ab^2 B) \sqrt{a + b \tan(c + dx)}}{64bd\sqrt{\cot(c + dx)}}$$

$$+ \frac{(40aAb - 5a^2 B - 48b^2 B) (a + b \tan(c + dx))^{3/2}}{96bd\sqrt{\cot(c + dx)}}$$

$$+ \frac{(8Ab - aB)(a + b \tan(c + dx))^{5/2}}{24bd\sqrt{\cot(c + dx)}} + \frac{B(a + b \tan(c + dx))^{7/2}}{4bd\sqrt{\cot(c + dx)}}$$

3.635. $\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

output $-(I*a-b)^{(5/2)}*(I*A-B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/d+1/64*(40*A*a^3*b-320*A*a*b^3-5*B*a^4-240*B*a^2*b^2+128*B*b^4)*\operatorname{arctanh}(b^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/b^{(3/2)}/d-(I*a+b)^{(5/2)}*(I*A+B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/d+1/64*(40*A*a^2*b-64*A*b^3-5*B*a^3-112*B*a*b^2)*(a+b*\tan(d*x+c))^{(1/2)}/b/d/\cot(d*x+c)^{(1/2)}+1/96*(40*A*a*b-5*B*a^2-48*B*b^2)*(a+b*\tan(d*x+c))^{(3/2)}/b/d/\cot(d*x+c)^{(1/2)}+1/24*(8*A*b-B*a)*(a+b*\tan(d*x+c))^{(5/2)}/b/d/\cot(d*x+c)^{(1/2)}+1/4*B*(a+b*\tan(d*x+c))^{(7/2)}/b/d/\cot(d*x+c)^{(1/2)}$

3.635.2 Mathematica [A] (verified)

Time = 6.25 (sec) , antiderivative size = 431, normalized size of antiderivative = 0.94

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\cot^{3/2}(c + dx)} dx = \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(-192 \sqrt[4]{-1} (-a + ib)^{5/2} b (i \dots) \right)}{\dots}$$

input `Integrate[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(3/2),x]`

output $(\operatorname{Sqrt}[\operatorname{Cot}[c + d*x]]*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]*(-192*(-1)^{(1/4)}*(-a + I*b)^{(5/2)}*b*(I*A + B)*\operatorname{ArcTan}[((-1)^{(1/4)}*\operatorname{Sqrt}[-a + I*b]*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]])/\operatorname{Sqrt}[a + b*\operatorname{Tan}[c + d*x]]] + 192*(-1)^{(3/4)}*(a + I*b)^{(5/2)}*b*(A + I*B)*\operatorname{ArcTan}[((-1)^{(1/4)}*\operatorname{Sqrt}[a + I*b]*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]])/\operatorname{Sqrt}[a + b*\operatorname{Tan}[c + d*x]]] - 3*(-40*a^2*A*b + 64*A*b^3 + 5*a^3*B + 112*a*b^2*B)*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]*\operatorname{Sqrt}[a + b*\operatorname{Tan}[c + d*x]] - 2*(-40*a*A*b + 5*a^2*B + 48*b^2*B)*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]*(a + b*\operatorname{Tan}[c + d*x])^{(3/2)} + 8*(8*A*b - a*B)*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]*(a + b*\operatorname{Tan}[c + d*x])^{(5/2)} + 48*B*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]]*(a + b*\operatorname{Tan}[c + d*x])^{(7/2)} - (3*\operatorname{Sqrt}[a]*(-40*a^3*A*b + 320*a*A*b^3 + 5*a^4*B + 240*a^2*b^2*B - 128*b^4*B)*\operatorname{ArcSinh}[(\operatorname{Sqrt}[b]*\operatorname{Sqrt}[\operatorname{Tan}[c + d*x]])/\operatorname{Sqrt}[a]]*\operatorname{Sqrt}[1 + (b*\operatorname{Tan}[c + d*x])/a])]/(\operatorname{Sqrt}[b]*\operatorname{Sqrt}[a + b*\operatorname{Tan}[c + d*x]])))/(192*b*d)$

3.635.3 Rubi [A] (verified)

Time = 2.76 (sec) , antiderivative size = 421, normalized size of antiderivative = 0.92, number of steps used = 20, number of rules used = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.543$, Rules used = {3042, 4729, 3042, 4090, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4130, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

↓ 3042

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\cot(c + dx)^{3/2}} dx$$

↓ 4729

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \tan^{\frac{3}{2}}(c + dx) (a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx$$

↓ 3042

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \tan(c + dx)^{3/2} (a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx)) dx$$

↓ 4090

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{\int -\frac{(a + b \tan(c + dx))^{5/2} (-(8Ab - aB) \tan^2(c + dx) + 8bB \tan(c + dx) + aB)}{2\sqrt{\tan(c + dx)}} dx}{4b} + \frac{B\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{5/2}}{4b} \right)$$

↓ 27

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{B\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{7/2}}{4bd} - \frac{\int \frac{(a + b \tan(c + dx))^{5/2} (-(8Ab - aB) \tan^2(c + dx) + 8bB \tan(c + dx) + aB)}{\sqrt{\tan(c + dx)}} dx}{8b} \right)$$

↓ 3042

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{B\sqrt{\tan(c + dx)}(a + b \tan(c + dx))^{7/2}}{4bd} - \frac{\int \frac{(a + b \tan(c + dx))^{5/2} (-(8Ab - aB) \tan(c + dx)^2 + 8bB \tan(c + dx) + aB)}{\sqrt{\tan(c + dx)}} dx}{8b} \right)$$

↓ 4130

3.635. $\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd}-\frac{\frac{1}{3}\int\frac{(a+b\tan(c+dx))^{3/2}(-((-5Ba^2+40Aba-48b^2B))}{2\sqrt{\tan(c+dx)}}dx}{\sqrt{\tan(c+dx)}}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd}-\frac{\frac{1}{6}\int\frac{(a+b\tan(c+dx))^{3/2}(-((-5Ba^2+40Aba-48b^2B))}{\sqrt{\tan(c+dx)}}dx}{\sqrt{\tan(c+dx)}}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd}-\frac{\frac{1}{6}\int\frac{(a+b\tan(c+dx))^{3/2}(-((-5Ba^2+40Aba-48b^2B))}{\sqrt{\tan(c+dx)}}dx}{\sqrt{\tan(c+dx)}}\right)$$

↓ 4130

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd}-\frac{\frac{1}{6}\left(\frac{1}{2}\int\frac{3\sqrt{a+b\tan(c+dx)}(-((-5Ba^3+40Aba^2-112b^2B))}{\sqrt{\tan(c+dx)}}dx\right)}{\sqrt{\tan(c+dx)}}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd}-\frac{\frac{1}{6}\left(\frac{3}{4}\int\frac{\sqrt{a+b\tan(c+dx)}(-((-5Ba^3+40Aba^2-112b^2B))}{\sqrt{\tan(c+dx)}}dx\right)}{\sqrt{\tan(c+dx)}}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd}-\frac{\frac{1}{6}\left(\frac{3}{4}\int\frac{\sqrt{a+b\tan(c+dx)}(-((-5Ba^3+40Aba^2-112b^2B))}{\sqrt{\tan(c+dx)}}dx\right)}{\sqrt{\tan(c+dx)}}\right)$$

↓ 4130

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd}-\frac{\frac{1}{6}\left(\frac{3}{4}\left(\int\frac{-((-5Ba^4+40Aba^3-240b^2Ba^2-320Ab^3a))}{\sqrt{\tan(c+dx)}}dx\right)\right)}{\sqrt{\tan(c+dx)}}\right)$$

3.635. $\int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd}-\frac{1}{6}\left(\frac{3}{4}\left(\frac{1}{2}\int\frac{-((-5Ba^4+40Aba^3-240b^2Ba^2-320Ab^3a+128b^4))}{\sqrt{a+b\tan(c+dx)}}dx\right)\right)\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd}-\frac{1}{6}\left(\frac{3}{4}\left(\frac{1}{2}\int\frac{-((-5Ba^4+40Aba^3-240b^2Ba^2-320Ab^3a+128b^4))}{\sqrt{a+b\tan(c+dx)}}dx\right)\right)\right)$$

↓ 4138

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd}-\frac{1}{6}\left(\frac{3}{4}\left(\int\frac{-((-5Ba^4+40Aba^3-240b^2Ba^2-320Ab^3a+128b^4))}{\sqrt{a+b\tan(c+dx)}}dx\right)\right)\right)$$

↓ 2035

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd}-\frac{1}{6}\left(\frac{3}{4}\left(\int\frac{-((-5Ba^4+40Aba^3-240b^2Ba^2-320Ab^3a+128b^4))}{\sqrt{a+b\tan(c+dx)}}dx\right)\right)\right)$$

↓ 2257

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd}-\frac{1}{6}\left(\frac{3}{4}\left(\int\frac{(5Ba^4-40Aba^3+240b^2Ba^2+320Ab^3a-128b^4)}{\sqrt{a+b\tan(c+dx)}}dx\right)\right)\right)$$

↓ 2009

3.635. $\int \frac{(a+b\tan(c+dx))^{5/2}(A+B\tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{7/2}}{4bd} - \frac{(8Ab-aB)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}}{3d} + \frac{1}{6} \right)$$

```
input Int[((a + b*Tan[c + d*x])^(5/2)*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(3/2),x
]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c
+ d*x])^(7/2))/(4*b*d) - (-1/3*((8*A*b - a*B)*Sqrt[Tan[c + d*x]]*(a + b*Ta
n[c + d*x])^(5/2))/d + (-1/2*((40*a*A*b - 5*a^2*B - 48*b^2*B)*Sqrt[Tan[c +
d*x]]*(a + b*Tan[c + d*x])^(3/2))/d + (3*((64*(I*a - b)^(5/2)*b*(I*A - B)
*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])] - ((4
0*a^3*A*b - 320*a*A*b^3 - 5*a^4*B - 240*a^2*b^2*B + 128*b^4*B)*ArcTanh[(Sq
rt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/Sqrt[b] + 64*b*(I*a +
b)^(5/2)*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*
Tan[c + d*x]])]/d - ((40*a^2*A*b - 64*A*b^3 - 5*a^3*B - 112*a*b^2*B)*Sqrt[
Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x])/d))/4)/6)/(8*b))
```

3.635.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2035 Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]
```

```
rule 2257 Int[(Px_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_.), x_Symbol
] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a
, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]
```

3.635. $\int \frac{(a+b \tan(c+dx))^{5/2}(A+B \tan(c+dx))}{\cot^{3/2}(c+dx)} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4090 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m + n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4130 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[C*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m + n + 1))), x] + Simp[1/(d*(m + n + 1)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^n*Simp[a*A*d*(m + n + 1) - C*(b*c*m + a*d*(n + 1)) + d*(A*b + a*B - b*C)*(m + n + 1)*Tan[e + f*x] - (C*m*(b*c - a*d) - b*B*d*(m + n + 1))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && !(IGtQ[n, 0] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4138 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.635.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3239 vs. 2(389) = 778.

Time = 1.05 (sec) , antiderivative size = 3240, normalized size of antiderivative = 7.09

method	result	size
derivativivedivides	Expression too large to display	3240
default	Expression too large to display	3240

```
input  int((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x,method=_RET
URNVERBOSE)
```

```
output  1/192/d*((b+a*cot(d*x+c))/cot(d*x+c))^(1/2)*(384*B*arctanh((b+a*cot(d*x+c)
)^(1/2)/b^(1/2))*a*b^4*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*cot(d*x+c)^4-48*A*ln(
a*cot(d*x+c)+b+(b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)+(a^2+b
^2)^(1/2))*b^(9/2)*(2*(a^2+b^2)^(1/2)+2*b)^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(
1/2)*cot(d*x+c)^4-384*A*arctan(((2*(a^2+b^2)^(1/2)+2*b)^(1/2)-2*(b+a*cot(d
*x+c))^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*a^2*b^(5/2)*(a^2+b^2)^(1/2)*c
ot(d*x+c)^4+384*A*arctan(((2*(b+a*cot(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*b)
^(1/2))/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*a^2*b^(5/2)*(a^2+b^2)^(1/2)*cot(d*x
+c)^4-192*B*arctan(((2*(a^2+b^2)^(1/2)+2*b)^(1/2)-2*(b+a*cot(d*x+c))^(1/2)
)/(2*(a^2+b^2)^(1/2)-2*b)^(1/2))*a^3*b^(3/2)*(a^2+b^2)^(1/2)*cot(d*x+c)^4+
192*B*arctan(((2*(a^2+b^2)^(1/2)+2*b)^(1/2)-2*(b+a*cot(d*x+c))^(1/2))/(2*(
a^2+b^2)^(1/2)-2*b)^(1/2))*a*b^(7/2)*(a^2+b^2)^(1/2)*cot(d*x+c)^4+192*B*ar
ctan((2*(b+a*cot(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*b)^(1/2))/(2*(a^2+b^2)
^(1/2)-2*b)^(1/2))*a^3*b^(3/2)*(a^2+b^2)^(1/2)*cot(d*x+c)^4-192*B*arctan(((
2*(b+a*cot(d*x+c))^(1/2)+(2*(a^2+b^2)^(1/2)+2*b)^(1/2))/(2*(a^2+b^2)^(1/2)
-2*b)^(1/2))*a*b^(7/2)*(a^2+b^2)^(1/2)*cot(d*x+c)^4+264*A*a^3*b^(3/2)*(b+a
*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*cot(d*x+c)^3-192*A*a*b^(7
/2)*(b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*cot(d*x+c)^3-432*
B*a^2*b^(5/2)*(b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(1/2)*cot(d*x
+c)^3+208*A*a^2*b^(5/2)*(b+a*cot(d*x+c))^(1/2)*(2*(a^2+b^2)^(1/2)-2*b)^(...
```

3.635.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17849 vs. $2(382) = 764$.

Time = 9.42 (sec) , antiderivative size = 35731, normalized size of antiderivative = 78.19

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\cot^{3/2}(c + dx)} dx = \text{Too large to display}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algo
rithm="fricas")`

output Too large to include

3.635.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\cot^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))**(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)**(3/2),x)`

output Timed out

3.635.7 Maxima [F]

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\cot^{3/2}(c + dx)} dx = \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^{5/2}}{\cot(dx + c)^{3/2}} dx$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^(5/2)/cot(d*x + c)^(3/
2), x)`

3.635. $\int \frac{(a+b \tan(c+dx))^{5/2} (A+B \tan(c+dx))}{\cot^{3/2}(c+dx)} dx$

3.635.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\cot^{3/2}(c + dx)} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^(5/2)*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algo
rithm="giac")`

output `Timed out`

3.635.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^{5/2} (A + B \tan(c + dx))}{\cot^{3/2}(c + dx)} dx = \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^{5/2}}{\cot(c + dx)^{3/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/cot(c + d*x)^(3/2),x
)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^(5/2))/cot(c + d*x)^(3/2),
x)`

3.636 $\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

3.636.1 Optimal result 6083
 3.636.2 Mathematica [A] (verified) 6084
 3.636.3 Rubi [A] (verified) 6085
 3.636.4 Maple [B] (warning: unable to verify) 6091
 3.636.5 Fricas [B] (verification not implemented) 6091
 3.636.6 Sympy [F(-1)] 6091
 3.636.7 Maxima [F] 6092
 3.636.8 Giac [F(-2)] 6092
 3.636.9 Mupad [F(-1)] 6092

3.636.1 Optimal result

Integrand size = 35, antiderivative size = 296

$$\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

$$= \frac{(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{\sqrt{ia - bd}}$$

$$- \frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{\sqrt{ia + bd}}$$

$$+ \frac{2(15a^2A - 8Ab^2 + 10abB) \sqrt{\cot(c+dx)}\sqrt{a+b \tan(c+dx)}}{15a^3d}$$

$$+ \frac{2(4Ab - 5aB) \cot^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}{15a^2d} - \frac{2A \cot^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}{5ad}$$

```
output (I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*a-b)^(1/2)-(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*a+b)^(1/2)+2/15*(4*A*b-5*B*a)*cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^(1/2)/a^2/d-2/5*A*cot(d*x+c)^(5/2)*(a+b*tan(d*x+c))^(1/2)/a/d+2/15*(15*A*a^2-8*A*b^2+10*B*a*b)*cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/a^3/d
```

3.636.2 Mathematica [A] (verified)

Time = 6.31 (sec) , antiderivative size = 444, normalized size of antiderivative = 1.50

$$\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

$$= \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)} \left(-\frac{(A-iB)\sqrt{a+b \tan(c+dx)}}{5a \tan^{\frac{5}{2}}(c+dx)} - \frac{(A+iB)\sqrt{a+b \tan(c+dx)}}{5a \tan^{\frac{5}{2}}(c+dx)} + \frac{(A+iB) \frac{(5ia+4b)\sqrt{a+b \tan(c+dx)}}{a \tan^{\frac{3}{2}}(c+dx)}}{\dots} \right)$$

```
input Integrate[(Cot[c + d*x]^(7/2)*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]
```

```
output (Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-1/5*((A - I*B)*Sqrt[a + b*Tan[c + d*x]]))/(a*Tan[c + d*x]^(5/2)) - ((A + I*B)*Sqrt[a + b*Tan[c + d*x]])/(5*a*Tan[c + d*x]^(5/2)) + ((A + I*B)*(((5*I)*a + 4*b)*Sqrt[a + b*Tan[c + d*x]]))/(a*Tan[c + d*x]^(3/2)) - (I*((15*(-1)^(3/4)*a^2*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[a + I*b] + (((15*I)*a^2 + 10*a*b - (8*I)*b^2)*Sqrt[a + b*Tan[c + d*x]])/(a*Sqrt[Tan[c + d*x]]))/a)/(15*a) - ((A - I*B)*(((5*I - (4*b)/a)*Sqrt[a + b*Tan[c + d*x]])/Tan[c + d*x]^(3/2) - (I*((15*(-1)^(3/4)*a^2*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[-a + I*b] + ((10*a*b - I*(15*a^2 - 8*b^2))*Sqrt[a + b*Tan[c + d*x]])/(a*Sqrt[Tan[c + d*x]]))/a)/(15*a))/d
```

3.636.3 Rubi [A] (verified)

Time = 1.85 (sec) , antiderivative size = 303, normalized size of antiderivative = 1.02, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.514$, Rules used = {3042, 4729, 3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

↓ 3042

$$\int \frac{\cot(c+dx)^{7/2}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

↓ 4729

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan^{\frac{7}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^{7/2}\sqrt{a+b \tan(c+dx)}} dx$$

↓ 4092

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2 \int \frac{4Ab \tan^2(c+dx)+5aA \tan(c+dx)+4Ab-5aB}{2 \tan^{\frac{5}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2A\sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{4Ab \tan^2(c+dx)+5aA \tan(c+dx)+4Ab-5aB}{\tan^{\frac{5}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2A\sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{4Ab \tan(c+dx)^2+5aA \tan(c+dx)+4Ab-5aB}{\tan(c+dx)^{5/2}\sqrt{a+b \tan(c+dx)}} dx}{5a} - \frac{2A\sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} \right)$$

3.636. $\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\begin{array}{c} \downarrow 4132 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int -\frac{15Aa^2+15B \tan(c+dx)a^2+10bBa-8Ab^2-2b(4Ab-5aB) \tan^2(c+dx)}{2 \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(4Ab-5aB)\sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) \\ \hline 5a \end{array}$$

$$\begin{array}{c} \downarrow 27 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{15Aa^2+15B \tan(c+dx)a^2+10bBa-8Ab^2-2b(4Ab-5aB) \tan^2(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(4Ab-5aB)\sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) \\ \hline 5a \end{array}$$

$$\begin{array}{c} \downarrow 3042 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{15Aa^2+15B \tan(c+dx)a^2+10bBa-8Ab^2-2b(4Ab-5aB) \tan(c+dx)^2}{\tan(c+dx)^{3/2}\sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2(4Ab-5aB)\sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) \\ \hline 5a \end{array}$$

$$\begin{array}{c} \downarrow 4132 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int -\frac{15(a^3B-a^3A \tan(c+dx))}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(15a^2A+10abB-8Ab^2)\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}{3a} - \frac{2(4Ab-5aB)\sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) \\ \hline 5a \end{array}$$

$$\begin{array}{c} \downarrow 27 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{15 \int \frac{a^3B-a^3A \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(15a^2A+10abB-8Ab^2)\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}{3a} - \frac{2(4Ab-5aB)\sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) \\ \hline 5a \end{array}$$

$$\begin{array}{c} \downarrow 3042 \end{array}$$

3.636. $\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{15 \int \frac{a^3 B - a^3 A \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(15a^2 A + 10abB - 8Ab^2)\sqrt{a+b \tan(c+dx)}}{3a ad \sqrt{\tan(c+dx)}} - \frac{2(4Ab - 5aB)\sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) \frac{1}{5a}$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2A\sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab - 5aB)\sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} + \frac{2(15a^2 A + 10abB - 8Ab^2)\sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2A\sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab - 5aB)\sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} + \frac{2(15a^2 A + 10abB - 8Ab^2)\sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} \right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2A\sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab - 5aB)\sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} + \frac{2(15a^2 A + 10abB - 8Ab^2)\sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} \right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2A\sqrt{a+b \tan(c+dx)}}{5ad \tan^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab - 5aB)\sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} + \frac{2(15a^2 A + 10abB - 8Ab^2)\sqrt{a+b \tan(c+dx)}}{ad \sqrt{\tan(c+dx)}} \right)$$

3.636. $\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\begin{array}{c}
 \downarrow 216 \\
 \left(\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \right) \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} + \frac{2(15a^2A+10abB-8Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right) \\
 \downarrow 219 \\
 \left(\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \right) \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{5ad\tan^{\frac{5}{2}}(c+dx)} - \frac{2(4Ab-5aB)\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} + \frac{2(15a^2A+10abB-8Ab^2)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right)
 \end{array}$$

input `Int[(Cot[c + d*x]^(7/2)*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A*Sqrt[a + b*Tan[c + d*x]])/(5*a*d*Tan[c + d*x]^(5/2)) - ((-2*(4*A*b - 5*a*B)*Sqrt[a + b*Tan[c + d*x]])/(3*a*d*Tan[c + d*x]^(3/2)) + ((15*(-((a^3*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(Sqrt[I*a - b]*d)) + (a^3*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d)))/a - (2*(15*a^2*A - 8*A*b^2 + 10*a*b*B)*Sqrt[a + b*Tan[c + d*x]])/(a*d*Sqrt[Tan[c + d*x]]))/(3*a))/(5*a))`

3.636.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4092 `Int[(((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

3.636.
$$\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && ! (ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.636.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 2.07 (sec) , antiderivative size = 1890883, normalized size of antiderivative = 6388.12

output too large to display

input `int(cot(d*x+c)^(7/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x)`

output `result too large to display`

3.636.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10265 vs. 2(242) = 484.

Time = 2.93 (sec) , antiderivative size = 10265, normalized size of antiderivative = 34.68

$$\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(7/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="fricas")`

output `Too large to include`

3.636.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(7/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)`

output `Timed out`

3.636.7 Maxima [F]

$$\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{7}{2}}}{\sqrt{b\tan(dx+c)+a}} dx$$

input `integrate(cot(d*x+c)^(7/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(7/2)/sqrt(b*tan(d*x + c) + a)
, x)`

3.636.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Exception raised: TypeError}$$

input `integrate(cot(d*x+c)^(7/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(con
st gen &`

3.636.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{\cot(c+dx)^{7/2}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

input `int((cot(c + d*x)^(7/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x
)`

output `int((cot(c + d*x)^(7/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),
x)`

3.636. $\int \frac{\cot^{\frac{7}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

$$3.637 \quad \int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

3.637.1 Optimal result	6093
3.637.2 Mathematica [A] (verified)	6094
3.637.3 Rubi [A] (verified)	6094
3.637.4 Maple [B] (warning: unable to verify)	6099
3.637.5 Fricas [B] (verification not implemented)	6099
3.637.6 Sympy [F(-1)]	6100
3.637.7 Maxima [F]	6100
3.637.8 Giac [F(-2)]	6101
3.637.9 Mupad [F(-1)]	6101

3.637.1 Optimal result

Integrand size = 35, antiderivative size = 243

$$\begin{aligned} & \int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx \\ &= -\frac{(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{\sqrt{ia-bd}} \\ & \quad - \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{\sqrt{ia+bd}} \\ & \quad + \frac{2(2Ab-3aB) \sqrt{\cot(c+dx)} \sqrt{a+b \tan(c+dx)}}{3a^2d} \\ & \quad - \frac{2A \cot^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}}{3ad} \end{aligned}$$

output $-(A+I*B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/d/(I*a-b)^{(1/2)}-(A-I*B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)}/(a+b*\tan(d*x+c))^{(1/2)})*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)}/d/(I*a+b)^{(1/2)}-2/3*A*\cot(d*x+c)^{(3/2)}*(a+b*\tan(d*x+c))^{(1/2)}/a/d+2/3*(2*A*b-3*B*a)*\cot(d*x+c)^{(1/2)}*(a+b*\tan(d*x+c))^{(1/2)}/a^2/d$

$$3.637. \quad \int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

3.637.2 Mathematica [A] (verified)

Time = 2.92 (sec) , antiderivative size = 213, normalized size of antiderivative = 0.88

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

$$= \frac{\sqrt{\cot(c+dx)} \left(\frac{3 \sqrt[4]{-1} (iA+B) \arctan\left(\frac{\sqrt[4]{-1} \sqrt{-a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\tan(c+dx)}}{\sqrt{-a+ib}} + \frac{3(-1)^{3/4} (A+iB) \arctan\left(\frac{\sqrt[4]{-1} \sqrt{a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{a+ib}} \right)}{3d}$$

input `Integrate[(Cot[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `(Sqrt[Cot[c + d*x]]*((3*(-1)^(1/4)*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]*Sqrt[Tan[c + d*x]])/Sqrt[-a + I*b] + (3*(-1)^(3/4)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]*Sqrt[Tan[c + d*x]])/Sqrt[a + I*b] - (2*(-2*A*b + 3*a*B + a*A*Cot[c + d*x])*Sqrt[a + b*Tan[c + d*x]])/a^2)/(3*d)`

3.637.3 Rubi [A] (verified)Time = 1.38 (sec) , antiderivative size = 240, normalized size of antiderivative = 0.99, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 4729, 3042, 4092, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{\cot(c+dx)^{5/2}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx$$

3.637. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\begin{aligned}
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^{5/2} \sqrt{a+b \tan(c+dx)}} dx \\
& \downarrow 4092 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2 \int \frac{2Ab \tan^2(c+dx)+3aA \tan(c+dx)+2Ab-3aB}{2 \tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2A \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) \\
& \downarrow 27 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{2Ab \tan^2(c+dx)+3aA \tan(c+dx)+2Ab-3aB}{\tan^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2A \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) \\
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{2Ab \tan(c+dx)^2+3aA \tan(c+dx)+2Ab-3aB}{\tan(c+dx)^{3/2} \sqrt{a+b \tan(c+dx)}} dx}{3a} - \frac{2A \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) \\
& \downarrow 4132 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2 \int -\frac{3(Aa^2+B \tan(c+dx)a^2)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(2Ab-3aB)\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}{3a} - \frac{2A \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) \\
& \downarrow 27 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{3 \int \frac{Aa^2+B \tan(c+dx)a^2}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(2Ab-3aB)\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}{3a} - \frac{2A \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right) \\
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{3 \int \frac{Aa^2+B \tan(c+dx)a^2}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2(2Ab-3aB)\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}}}{3a} - \frac{2A \sqrt{a+b \tan(c+dx)}}{3ad \tan^{\frac{3}{2}}(c+dx)} \right)
\end{aligned}$$

3.637. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{2(2Ab-3aB)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{3\left(\frac{1}{2}a^2(A+iB)\int\frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}\right)}{3a} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{2(2Ab-3aB)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{3\left(\frac{1}{2}a^2(A+iB)\int\frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}\right)}{3a} \right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{2(2Ab-3aB)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{3\left(\frac{a^2(A-iB)\int\frac{1-i\tan(c+dx)}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}}\right)}{3a} \right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{2(2Ab-3aB)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{3\left(\frac{a^2(A+iB)\int\frac{1}{(ia-b)\tan(c+dx)+1}}{\frac{a+b\tan(c+dx)}{d}}\right)}{3a} \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{2(2Ab-3aB)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{3\left(\frac{a^2(A-iB)\int\frac{1}{1-(ia+b)\tan(c+dx)}}{\frac{a+b\tan(c+dx)}{d}}\right)}{3a} \right)$$

3.637. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{3ad\tan^{\frac{3}{2}}(c+dx)} - \frac{-2(2Ab-3aB)\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{3\left(\frac{a^2(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}}\right)}{3a} \right)$$

input `Int[(Cot[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A*Sqrt[a + b*Tan[c + d*x]])/(3*a*d*Tan[c + d*x]^(3/2)) - ((3*((a^2*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a - b]*d) + (a^2*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d)))/a - (2*(2*A*b - 3*a*B)*Sqrt[a + b*Tan[c + d*x]])/(a*d*Sqrt[Tan[c + d*x]])))/(3*a))`

3.637.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[(((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[(((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

3.637. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.637.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.31 (sec) , antiderivative size = 1890882, normalized size of antiderivative = 7781.41

output too large to display

```
input int(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x)
```

```
output result too large to display
```

3.637.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10191 vs. 2(195) = 390.

Time = 2.94 (sec) , antiderivative size = 10191, normalized size of antiderivative = 41.94

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = \text{Too large to display}$$

3.637. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="fricas")`

output Too large to include

3.637.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)`

output Timed out

3.637.7 Maxima [F]

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{5}{2}}}{\sqrt{b\tan(dx+c)+a}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(5/2)/sqrt(b*tan(d*x + c) + a)
, x)`

3.637.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Exception raised: TypeError}$$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen &`

3.637.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{\cot(c+dx)^{5/2}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

input `int((cot(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x)`

output `int((cot(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x)`

3.638
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

3.638.1 Optimal result 6102
 3.638.2 Mathematica [A] (verified) 6103
 3.638.3 Rubi [A] (verified) 6103
 3.638.4 Maple [B] (warning: unable to verify) 6107
 3.638.5 Fricas [B] (verification not implemented) 6107
 3.638.6 Sympy [F] 6107
 3.638.7 Maxima [F] 6108
 3.638.8 Giac [F] 6108
 3.638.9 Mupad [F(-1)] 6108

3.638.1 Optimal result

Integrand size = 35, antiderivative size = 199

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

$$= -\frac{(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{\sqrt{ia-bd}}$$

$$+ \frac{(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{\sqrt{ia+bd}}$$

$$- \frac{2A \sqrt{\cot(c+dx)} \sqrt{a+b \tan(c+dx)}}{ad}$$

output

```
-(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot
(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*a-b)^(1/2)+(I*A+B)*arctanh((I*a+b)^(1/
2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1
/2)/d/(I*a+b)^(1/2)-2*A*cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^(1/2)/a/d
```

3.638.2 Mathematica [A] (verified)

Time = 1.77 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.97

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx = \frac{\sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)} \left(-\frac{\sqrt[4]{-1}(A-iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} + \frac{\sqrt[4]{-1}(A+iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{a+ib}} \right)}{d}$$

```
input Integrate[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]
```

```
output -((Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-(((1/4)*(-1)^1/4*(A - I*B)*ArcTan[(-1)^1/4*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[-a + I*b] + ((1/4)*(-1)^1/4*(A + I*B)*ArcTan[(-1)^1/4*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[a + I*b] + (2*A*Sqrt[a + b*Tan[c + d*x]])/(a*Sqrt[Tan[c + d*x]]))/d)
```

3.638.3 Rubi [A] (verified)

Time = 1.04 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.94, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 4729, 3042, 4092, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

↓ 3042

$$\int \frac{\cot(c+dx)^{3/2}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

↓ 4729

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx$$

$$\begin{aligned}
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^{3/2} \sqrt{a+b \tan(c+dx)}} dx \\
& \downarrow 4092 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2 \int -\frac{aB-aA \tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2A\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right) \\
& \downarrow 27 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{aB-aA \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2A\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right) \\
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{aB-aA \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a} - \frac{2A\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} \right) \\
& \downarrow 4099 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{\frac{1}{2}a(B+iA) \int \frac{i \tan(c+dx)+1}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2}a(-B+iA)}{a} \right) \\
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{\frac{1}{2}a(B+iA) \int \frac{i \tan(c+dx)+1}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx - \frac{1}{2}a(-B+iA)}{a} \right) \\
& \downarrow 4098 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{a(B+iA) \int \frac{1}{(1-i \tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} d \tan(c+dx) - \frac{a}{a+b \tan(c+dx)}}{2d} \right) \\
& \downarrow 104 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b \tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{a(B+iA) \int \frac{1}{1-\frac{(ia+b) \tan(c+dx)}{a+b \tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} - \frac{a(-B+iA) \int \frac{1}{(ia-b) \tan(c+dx)}}{a+b \tan(c+dx)}}{a} \right)
\end{aligned}$$

3.638. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\begin{aligned} & \downarrow 216 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{a(B+iA)\int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} - \frac{a(-B+iA)\arctan\left(\frac{\sqrt{-b+ia}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} \right) \\ & \downarrow 219 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A\sqrt{a+b\tan(c+dx)}}{ad\sqrt{\tan(c+dx)}} + \frac{a(B+iA)\operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}} - \frac{a(-B+iA)\arctan\left(\frac{\sqrt{-b+ia}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} \right) \end{aligned}$$

input `Int[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-((a*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d)) + (a*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d))/a - (2*A*Sqrt[a + b*Tan[c + d*x]])/(a*d*Sqrt[Tan[c + d*x]]))`

3.638.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 104 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

3.638. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`
- rule 4098 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`
- rule 4099 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`
- rule 4729 `Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.638.
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

3.638.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.14 (sec) , antiderivative size = 1886187, normalized size of antiderivative = 9478.33

output too large to display

input `int(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x)`

output `result too large to display`

3.638.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10120 vs. $2(156) = 312$.

Time = 3.02 (sec) , antiderivative size = 10120, normalized size of antiderivative = 50.85

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="fricas")`

output `Too large to include`

3.638.6 Sympy [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(A+B\tan(c+dx))\cot^{\frac{3}{2}}(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx$$

input `integrate(cot(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**(3/2)/sqrt(a + b*tan(c + d*x))
, x)`

3.638. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

3.638.7 Maxima [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{\sqrt{b\tan(dx+c)+a}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(3/2)/sqrt(b*tan(d*x + c) + a)
, x)`

3.638.8 Giac [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{\sqrt{b\tan(dx+c)+a}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(3/2)/sqrt(b*tan(d*x + c) + a)
, x)`

3.638.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{\cot(c+dx)^{3/2}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

input `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x
)`

output `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),
x)`

3.638. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

$$3.639 \quad \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

3.639.1 Optimal result	6109
3.639.2 Mathematica [A] (verified)	6109
3.639.3 Rubi [A] (verified)	6110
3.639.4 Maple [B] (warning: unable to verify)	6113
3.639.5 Fricas [B] (verification not implemented)	6113
3.639.6 Sympy [F]	6113
3.639.7 Maxima [F]	6114
3.639.8 Giac [F(-2)]	6114
3.639.9 Mupad [F(-1)]	6114

3.639.1 Optimal result

Integrand size = 35, antiderivative size = 163

$$\begin{aligned} & \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx \\ &= \frac{(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{\sqrt{ia-bd}} \\ & \quad + \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{\sqrt{ia+bd}} \end{aligned}$$

output $(A+I*B)*\arctan((I*a-b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)}*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)/d/(I*a-b)^{(1/2)}+(A-I*B)*\operatorname{arctanh}((I*a+b)^{(1/2)}*\tan(d*x+c)^{(1/2)/(a+b*\tan(d*x+c))}^{(1/2)}*\cot(d*x+c)^{(1/2)}*\tan(d*x+c)^{(1/2)/d/(I*a+b)^{(1/2)}$

3.639.2 Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.96

$$\begin{aligned} & \int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx \\ &= \frac{\sqrt[4]{-1} \left(-\frac{(iA+B) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} + \frac{(-iA+B) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{a+ib}} \right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{d} \end{aligned}$$

3.639. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

input `Integrate[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `((-1)^(1/4)*(-(((I*A + B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[-a + I*b]) + (((-I)*A + B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[a + I*b])*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]/d`

3.639.3 Rubi [A] (verified)

Time = 0.79 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.88, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.257$, Rules used = {3042, 4729, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

↓ 3042

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$$

↓ 4729

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2}(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}(A-iB) \int \frac{i \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2}(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx + \frac{1}{2}(A-iB) \int \frac{i \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx \right)$$

3.639. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{\sqrt{a+b \tan(c+dx)}} dx$

$$\begin{aligned}
& \downarrow 4098 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(A-iB) \int \frac{1}{(1-i \tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{2d} + \frac{(A+iB) \int \frac{1}{(1+i \tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} d \tan(c+dx)}{2d} \right) \\
& \downarrow 104 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(A+iB) \int \frac{1}{\frac{(ia-b) \tan(c+dx)}{a+b \tan(c+dx)} + 1} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} + \frac{(A-iB) \int \frac{1}{1 - \frac{(ia+b) \tan(c+dx)}{a+b \tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} \right) \\
& \downarrow 216 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(A-iB) \int \frac{1}{1 - \frac{(ia+b) \tan(c+dx)}{a+b \tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d} + \frac{(A+iB) \arctan \left(\frac{\sqrt{-b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right)}{d\sqrt{-b+ia}} \right) \\
& \downarrow 219 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(A+iB) \arctan \left(\frac{\sqrt{-b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right)}{d\sqrt{-b+ia}} + \frac{(A-iB) \operatorname{arctanh} \left(\frac{\sqrt{b+ia} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right)}{d\sqrt{b+ia}} \right)
\end{aligned}$$

input `Int[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/Sqrt[a + b*Tan[c + d*x]],x]`

output `((A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a - b]*d) + ((A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d)*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]`

3.639.3.1 Defintions of rubi rules used

- rule 104 `Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`
- rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`
- rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.639.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.23 (sec) , antiderivative size = 1880801, normalized size of antiderivative = 11538.66

output too large to display

input `int(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x)`

output `result too large to display`

3.639.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10053 vs. $2(127) = 254$.

Time = 2.89 (sec) , antiderivative size = 10053, normalized size of antiderivative = 61.67

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="fricas")`

output `Too large to include`

3.639.6 Sympy [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(A+B\tan(c+dx))\sqrt{\cot(c+dx)}}{\sqrt{a+b\tan(c+dx)}} dx$$

input `integrate(cot(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(cot(c + d*x))/sqrt(a + b*tan(c + d*x)),
x)`

3.639.7 Maxima [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{(B\tan(dx+c)+A)\sqrt{\cot(dx+c)}}{\sqrt{b\tan(dx+c)+a}} dx$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorith="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(cot(d*x + c))/sqrt(b*tan(d*x + c) + a), x)`

3.639.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \text{Exception raised: TypeError}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(1/2),x, algorith="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen &`

3.639.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx = \int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$$

input `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x)`

output `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(1/2),x)`

3.639. $\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx$

3.640 $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}\sqrt{a+b \tan(c+dx)}} dx$

3.640.1 Optimal result 6115
 3.640.2 Mathematica [A] (verified) 6116
 3.640.3 Rubi [A] (verified) 6116
 3.640.4 Maple [B] (warning: unable to verify) 6120
 3.640.5 Fricas [B] (verification not implemented) 6121
 3.640.6 Sympy [F] 6121
 3.640.7 Maxima [F] 6121
 3.640.8 Giac [F(-2)] 6122
 3.640.9 Mupad [F(-1)] 6122

3.640.1 Optimal result

Integrand size = 35, antiderivative size = 228

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}\sqrt{a + b \tan(c + dx)}} dx$$

$$= \frac{(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{\sqrt{ia - bd}}$$

$$+ \frac{2B \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{\sqrt{bd}}$$

$$- \frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{\sqrt{ia + bd}}$$

```
output (I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(
d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*a-b)^(1/2)+2*B*arctanh(b^(1/2)*tan(d*x+
c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/b^(1/
2)-(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*
cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*a+b)^(1/2)
```

3.640.2 Mathematica [A] (verified)

Time = 1.60 (sec) , antiderivative size = 225, normalized size of antiderivative = 0.99

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}} dx$$

$$= \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(\sqrt[4]{-1} \left(-\frac{(A - iB) \arctan\left(\frac{\sqrt[4]{-1} \sqrt{-a + ib} \sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{\sqrt{-a + ib}} \right) + \frac{(A + iB) \arctan\left(\frac{\sqrt[4]{-1} \sqrt{-a + ib} \sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}}\right)}{\sqrt{a + ib}} \right)}{d}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*Sqrt[a + b*Tan[c + d*x]]),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-1)^(1/4)*(-((A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[-a + I*b]) + ((A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[a + I*b]) + (2*Sqrt[a]*B*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/(Sqrt[b]*Sqrt[a + b*Tan[c + d*x]]))/d`

3.640.3 Rubi [A] (verified)

Time = 1.34 (sec) , antiderivative size = 189, normalized size of antiderivative = 0.83, number of steps used = 15, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3042, 4729, 3042, 4097, 3042, 4099, 3042, 4098, 104, 216, 219, 4117, 65, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}} dx$$

↓ 4729

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\sqrt{\tan(c + dx)} (A + B \tan(c + dx))}{\sqrt{a + b \tan(c + dx)}} dx$$

$$\begin{aligned}
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\sqrt{\tan(c+dx)}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx \\
& \downarrow 4097 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{A\tan(c+dx)-B}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + B \int \frac{\tan^2(c+dx)+1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \right) \\
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\int \frac{A\tan(c+dx)-B}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + B \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \right) \\
& \downarrow 4099 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2}(-B+iA) \int \frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{1}{2}(B+iA) \int \frac{i\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx \right) \\
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{1}{2}(-B+iA) \int \frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{1}{2}(B+iA) \int \frac{i\tan(c+dx)}{\sqrt{\tan(c+dx)}} dx \right) \\
& \downarrow 4098 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{(B+iA) \int \frac{1}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} d\tan(c+dx)}{2d} + \frac{(-B+iA) \int \frac{1}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}} d\tan(c+dx)}{2d} \right) \\
& \downarrow 104 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA) \int \frac{1}{\frac{(ia-b)\tan(c+dx)}{a+b\tan(c+dx)}+1} d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}} {d} - \frac{(B+iA) \int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}} {d} \right) \\
& \downarrow 216 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{(B+iA) \int \frac{1}{1-\frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}} {d} + B \int \frac{\tan(c+dx)^2+1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx \right)
\end{aligned}$$

3.640. $\int \frac{A+B\tan(c+dx)}{\sqrt{\cot(c+dx)}\sqrt{a+b\tan(c+dx)}} dx$

$$\begin{aligned} & \downarrow 219 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(B \int \frac{\tan(c+dx)^2 + 1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx + \frac{(-B+iA) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} \right) \\ & \downarrow 4117 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B \int \frac{1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} d\tan(c+dx)}{d} + \frac{(-B+iA) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} \right) \\ & \downarrow 65 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2B \int \frac{1}{1-\frac{b\tan(c+dx)}{a+b\tan(c+dx)}} d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} + \frac{(-B+iA) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} - \frac{(B-iA) \operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}} \right) \\ & \downarrow 219 \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{(-B+iA) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} - \frac{(B+iA) \operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{b+ia}} + \frac{2B}{d} \right) \end{aligned}$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*Sqrt[a + b*Tan[c + d*x]]),x]`

output `((I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a - b]*d) + (2*B*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[b]*d) - ((I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d)*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]`

3.640.3.1 Defintions of rubi rules used

- rule 65 `Int[1/(Sqrt[(b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[b*x]/Sqrt[c + d*x]], x] /; FreeQ[{b, c, d}, x] && !GtQ[c, 0]`
- rule 104 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4097 `Int[(Sqrt[(a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]]*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]))/Sqrt[(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Int[Simp[a*A - b*B + (A*b + a*B)*Tan[e + f*x], x]/(Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]), x] + Simp[b*B Int[(1 + Tan[e + f*x]^2)/(Sqrt[a + b*Tan[e + f*x]]*Sqrt[c + d*Tan[e + f*x]]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

```
rule 4099 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
  (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2,
0]
```

```
rule 4117 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_.) + (d_.)*tan[(e_.)
+ (f_.)*(x_)])^(n_.)*((A_) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] :=
Simp[A/f Subst[Int[(a + b*x)^m*(c + d*x)^n, x], x, Tan[e + f*x]], x] /;
FreeQ[{a, b, c, d, e, f, A, C, m, n}, x] && EqQ[A, C]
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.640.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 0.98 (sec) , antiderivative size = 1887832, normalized size of antiderivative = 8279.96

output too large to display

```
input int((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2),x)
```

```
output result too large to display
```

3.640.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10230 vs. $2(180) = 360$.

Time = 4.35 (sec) , antiderivative size = 20493, normalized size of antiderivative = 89.88

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="fricas")`

output Too large to include

3.640.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)} \sqrt{\cot(c + dx)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(1/2)/(a+b*tan(d*x+c))**(1/2),x)`

output `Integral((A + B*tan(c + d*x))/(sqrt(a + b*tan(c + d*x))*sqrt(cot(c + d*x))
, x)`

3.640.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}} dx = \int \frac{B \tan(dx + c) + A}{\sqrt{b \tan(dx + c) + a} \sqrt{\cot(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)/(sqrt(b*tan(d*x + c) + a)*sqrt(cot(d*x + c)
)), x)`

3.640.8 Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen &`

3.640.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(1/2)),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(1/2)),x)`

3.641
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} dx$$

3.641.1 Optimal result 6123
 3.641.2 Mathematica [A] (verified) 6124
 3.641.3 Rubi [A] (verified) 6124
 3.641.4 Maple [B] (warning: unable to verify) 6127
 3.641.5 Fricas [B] (verification not implemented) 6128
 3.641.6 Sympy [F] 6128
 3.641.7 Maxima [F] 6128
 3.641.8 Giac [F(-2)] 6129
 3.641.9 Mupad [F(-1)] 6129

3.641.1 Optimal result

Integrand size = 35, antiderivative size = 266

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)\sqrt{a + b \tan(c + dx)}} dx$$

$$= -\frac{(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{\sqrt{ia - bd}}$$

$$+ \frac{(2Ab - aB) \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{b^{3/2}d}$$

$$- \frac{(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)}}{\sqrt{ia + bd}}$$

$$+ \frac{B\sqrt{a + b \tan(c + dx)}}{bd\sqrt{\cot(c + dx)}}$$

```
output (2*A*b-B*a)*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d
*x+c)^(1/2)*tan(d*x+c)^(1/2)/b^(3/2)/d-(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*
x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*
a-b)^(1/2)-(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c)
)^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*a+b)^(1/2)+B*(a+b*tan(d*x+c
))^(1/2)/b/d/cot(d*x+c)^(1/2)
```

3.641.2 Mathematica [A] (verified)

Time = 2.69 (sec) , antiderivative size = 354, normalized size of antiderivative = 1.33

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx$$

$$= \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(-\sqrt{a} \sqrt{-a + ib} \sqrt{a + ib} (-2Ab + aB) \operatorname{arcsinh} \left(\frac{\sqrt{b} \sqrt{\tan(c + dx)}}{\sqrt{a}} \right) \sqrt{1 + \frac{b \tan(c + dx)}{a}} \right)}{\dots}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]]),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-(Sqrt[a]*Sqrt[-a + I*b]*Sqrt[a + I*b]*(-2*A*b + a*B)*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a]) + Sqrt[b]*((-1)^(1/4)*Sqrt[a + I*b]*b*(I*A + B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]*Sqrt[a + b*Tan[c + d*x]] + Sqrt[-a + I*b]*((-1)^(3/4)*b*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]]*Sqrt[a + b*Tan[c + d*x]] + Sqrt[a + I*b]*B*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])))/(Sqrt[-a + I*b]*Sqrt[a + I*b]*b^(3/2)*d*Sqrt[a + b*Tan[c + d*x]])`

3.641.3 Rubi [A] (verified)Time = 1.16 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.86, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4729, 3042, 4090, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2} \sqrt{a + b \tan(c + dx)}} dx$$

$$\downarrow \text{4729}$$

3.641. $\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx$

$$\begin{aligned}
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\tan(c+dx)^{3/2}(A+B\tan(c+dx))}{\sqrt{a+b\tan(c+dx)}} dx \\
& \quad \downarrow \text{4090} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int -\frac{((2Ab-aB)\tan^2(c+dx)+2bB\tan(c+dx)+aB)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{b} + \frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd} - \frac{\int -\frac{((2Ab-aB)\tan^2(c+dx)+2bB\tan(c+dx)+aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{2b} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd} - \frac{\int -\frac{((2Ab-aB)\tan(c+dx)^2+2bB\tan(c+dx)+aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{2b} \right) \\
& \quad \downarrow \text{4138} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd} - \frac{\int -\frac{((2Ab-aB)\tan^2(c+dx)+2bB\tan(c+dx)+aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} d\tan}{2bd} \right) \\
& \quad \downarrow \text{2035} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd} - \frac{\int -\frac{((2Ab-aB)\tan^2(c+dx)+2bB\tan(c+dx)+aB)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} d\sqrt{t}}{bd} \right) \\
& \quad \downarrow \text{2257} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd} - \frac{\int \left(\frac{aB-2Ab}{\sqrt{a+b\tan(c+dx)}} + \frac{2(Ab+B\tan(c+dx)b)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} \right) dx}{bd} \right)
\end{aligned}$$

3.641. $\int \frac{A+B\tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}} dx$

↓ 2009

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{B\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}{bd} - \frac{b(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{-b+ia}} - \frac{(2Ab-aB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{-b+ia}} \right)$$

input `Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-(((b*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/Sqrt[I*a - b] - ((2*A*b - a*B)*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[b] + (b*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[I*a + b])/(b*d) + (B*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/(b*d))`

3.641.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2035 `Int[(F_x_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[F_x, x, k], x], x, x^(1/k)], x] /; FractionQ[m] && AlgebraicFunctionQ[F_x, x]`

rule 2257 `Int[(P_x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[P_x*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a, c, d, e, q}, x] && PolyQ[P_x, x] && IntegerQ[p]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.641. $\int \frac{A+B\tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)\sqrt{a+b\tan(c+dx)}} dx$

```
rule 4090 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[b*B*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(m +
n))), x] + Simp[1/(d*(m + n)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Ta
n[e + f*x])^n*Simp[a^2*A*d*(m + n) - b*B*(b*c*(m - 1) + a*d*(n + 1)) + d*(m
+ n)*(2*a*A*b + B*(a^2 - b^2))*Tan[e + f*x] - (b*B*(b*c - a*d)*(m - 1) - b
*(A*b + a*B)*d*(m + n))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e,
f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2
, 0] && GtQ[m, 1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(IGtQ[n, 1]
&& (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4138 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^
2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f
, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 +
d^2, 0]
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.641.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.25 (sec) , antiderivative size = 1890515, normalized size of antiderivative = 7107.20

output too large to display

```
input int((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(1/2),x)
```

```
output result too large to display
```

3.641.
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx) \sqrt{a+b \tan(c+dx)}} dx$$

3.641.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10269 vs. $2(214) = 428$.

Time = 5.04 (sec) , antiderivative size = 20571, normalized size of antiderivative = 77.33

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="fricas")`

output Too large to include

3.641.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{a + b \tan(c + dx)} \cot^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(3/2)/(a+b*tan(d*x+c))^(1/2),x)`

output `Integral((A + B*tan(c + d*x))/(sqrt(a + b*tan(c + d*x))*cot(c + d*x)**(3/2)), x)`

3.641.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \int \frac{B \tan(dx + c) + A}{\sqrt{b \tan(dx + c) + a} \cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)/(sqrt(b*tan(d*x + c) + a)*cot(d*x + c)^(3/2)), x)`

3.641.8 Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(1/2),x, algo
rithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(con
st gen &`

3.641.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2} \sqrt{a + b \tan(c + dx)}} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(1/2)),x
)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(1/2)),
x)`

3.642
$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.642.1 Optimal result 6130
 3.642.2 Mathematica [A] (verified) 6131
 3.642.3 Rubi [A] (verified) 6131
 3.642.4 Maple [B] (warning: unable to verify) 6137
 3.642.5 Fricas [B] (verification not implemented) 6137
 3.642.6 Sympy [F(-1)] 6138
 3.642.7 Maxima [F] 6138
 3.642.8 Giac [F(-2)] 6139
 3.642.9 Mupad [F(-1)] 6139

3.642.1 Optimal result

Integrand size = 35, antiderivative size = 316

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx =$$

$$\frac{(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{(ia - b)^{3/2}d}$$

$$- \frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{(ia + b)^{3/2}d}$$

$$+ \frac{2b(5a^2Ab + 8Ab^3 - 3a^3B - 6ab^2B)}{3a^3(a^2 + b^2)d\sqrt{\cot(c+dx)}\sqrt{a+b \tan(c+dx)}}$$

$$+ \frac{2(4Ab - 3aB)\sqrt{\cot(c+dx)}}{3a^2d\sqrt{a+b \tan(c+dx)}} - \frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad\sqrt{a+b \tan(c+dx)}}$$

output

```
-(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot
(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a-b)^(3/2)/d-(I*A+B)*arctanh((I*a+b)^(1/
2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1
/2)/(I*a+b)^(3/2)/d-2/3*A*cot(d*x+c)^(3/2)/a/d/(a+b*tan(d*x+c))^(1/2)+2/3*
b*(5*A*a^2*b+8*A*b^3-3*B*a^3-6*B*a*b^2)/a^3/(a^2+b^2)/d/cot(d*x+c)^(1/2)/((
a+b*tan(d*x+c))^(1/2)+2/3*(4*A*b-3*B*a)*cot(d*x+c)^(1/2)/a^2/d/(a+b*tan(d*
x+c))^(1/2)
```

3.642.
$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.642.2 Mathematica [A] (verified)

Time = 4.47 (sec) , antiderivative size = 301, normalized size of antiderivative = 0.95

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{\sqrt{\cot(c+dx)} \left(3 \sqrt[4]{-1} a \left(\frac{(a+ib)^{iA+B} \arctan \left(\frac{\sqrt[4]{-1} \sqrt{-a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}} \right)}{\sqrt{-a+ib}} \right) + \dots \right)}{a^2+b^2}$$

```
input Integrate[(Cot[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2), x]
```

```
output (Sqrt[Cot[c + d*x]]*((3*(-1)^(1/4)*a*(((a + I*b)*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/Sqrt[-a + I*b] + ((I*a + b)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/Sqrt[a + I*b])*Sqrt[Tan[c + d*x]])/(a^2 + b^2) + (8*A*b - 6*a*B)/(a*Sqrt[a + b*Tan[c + d*x]]) - (2*A*Cot[c + d*x])/Sqrt[a + b*Tan[c + d*x]] + (2*b*(5*a^2*A*b + 8*A*b^3 - 3*a^3*B - 6*a*b^2*B)*Tan[c + d*x])/(a^2*(a^2 + b^2)*Sqrt[a + b*Tan[c + d*x]])))/(3*a*d)
```

3.642.3 Rubi [A] (verified)

Time = 2.08 (sec) , antiderivative size = 337, normalized size of antiderivative = 1.07, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.514$, Rules used = {3042, 4729, 3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx \xrightarrow{3042} \int \frac{\cot(c+dx)^{5/2}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

$$\begin{aligned}
& \downarrow 4729 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx \\
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^{5/2}(a+b \tan(c+dx))^{3/2}} dx \\
& \downarrow 4092 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2 \int \frac{4Ab \tan^2(c+dx)+3aA \tan(c+dx)+4Ab-3aB}{2 \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx}{3a} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} \right) \\
& \downarrow 27 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{4Ab \tan^2(c+dx)+3aA \tan(c+dx)+4Ab-3aB}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx}{3a} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} \right) \\
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{4Ab \tan(c+dx)^2+3aA \tan(c+dx)+4Ab-3aB}{\tan(c+dx)^{3/2}(a+b \tan(c+dx))^{3/2}} dx}{3a} - \frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} \right) \\
& \downarrow 4132 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2 \int -\frac{3Aa^2+3B \tan(c+dx)a^2+6bBa-8Ab^2-2b(4Ab-3aB) \tan^2(c+dx)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{a} - \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \right) \\
& \downarrow 27 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{3Aa^2+3B \tan(c+dx)a^2+6bBa-8Ab^2-2b(4Ab-3aB) \tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{a} - \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \right) \\
& \downarrow 3042
\end{aligned}$$

3.642. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{3Aa^2+3B \tan(c+dx)a^2+6bBa-8Ab^2-2b(4Ab-3aB) \tan(c+dx)^2}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{a} - \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \right) \frac{1}{3a}$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2 \int \frac{3(a^3(aA+bB)-a^3(Ab-aB) \tan(c+dx))}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-3a^3B+5a^2Ab-6ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{a} - \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \right) \frac{1}{3a}$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{3 \int \frac{a^3(aA+bB)-a^3(Ab-aB) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-3a^3B+5a^2Ab-6ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{a} - \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \right) \frac{1}{3a}$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{3 \int \frac{a^3(aA+bB)-a^3(Ab-aB) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-3a^3B+5a^2Ab-6ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{a} - \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \right) \frac{1}{3a}$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} - \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} + \frac{2b(-3a^3B+5a^2Ab-6ab^2B+8Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \right)$$

↓ 3042

3.642. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} - \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} + \frac{2b(-3a^3B+ad^2)}{ad^2} \right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} - \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} + \frac{2b(-3a^3B+ad^2)}{ad^2} \right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} - \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} + \frac{2b(-3a^3B+ad^2)}{ad^2} \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} - \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} + \frac{2b(-3a^3B+ad^2)}{ad^2} \right)$$

↓ 219

3.642. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{3}{2}}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)\sqrt{a+b \tan(c+dx)}} - \frac{2(4Ab-3aB)}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} + \frac{2b(-3a^3B+ad^3)}{ad^3} \right)$$

input `Int[(Cot[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A)/(3*a*d*Tan[c + d*x]^(3/2)*Sqrt[a + b*Tan[c + d*x]]) - ((-2*(4*A*b - 3*a*B))/(a*d*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])) + ((3*((a^3*(a - I*b)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d) + (a^3*(a + I*b)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d)))/(a*(a^2 + b^2)) - (2*b*(5*a^2*A*b + 8*A*b^3 - 3*a^3*B - 6*a*b^2*B)*Sqrt[Tan[c + d*x]])/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]))/a)/(3*a)`

3.642.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 104 `Int[(((a_.) + (b_.)*(x_))^(m_))*((c_.) + (d_.)*(x_))^(n_)]/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4098 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

3.642.
$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.642.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 2.00 (sec) , antiderivative size = 1564143, normalized size of antiderivative = 4949.82

output too large to display

```
input int(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x)
```

```
output result too large to display
```

3.642.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18876 vs. $2(259) = 518$.

Time = 7.08 (sec) , antiderivative size = 18876, normalized size of antiderivative = 59.73

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

3.642. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algo
rithm="fricas")`

output Too large to include

3.642.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)`

output Timed out

3.642.7 Maxima [F]

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{5}{2}}}{(b\tan(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(5/2)/(b*tan(d*x + c) + a)^(3/
2), x)`

3.642.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algo
rithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(con
st gen &`

3.642.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{\cot(c+dx)^{5/2}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$$

input `int((cot(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x
)`

output `int((cot(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),
x)`

3.643 $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

3.643.1 Optimal result 6140
 3.643.2 Mathematica [A] (verified) 6141
 3.643.3 Rubi [A] (verified) 6141
 3.643.4 Maple [B] (warning: unable to verify) 6146
 3.643.5 Fricas [B] (verification not implemented) 6146
 3.643.6 Sympy [F] 6147
 3.643.7 Maxima [F] 6147
 3.643.8 Giac [F] 6148
 3.643.9 Mupad [F(-1)] 6148

3.643.1 Optimal result

Integrand size = 35, antiderivative size = 256

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{(ia-b)^{3/2}d} - \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{(ia+b)^{3/2}d} - \frac{2b(a^2A+2Ab^2-abB)}{a^2(a^2+b^2)d\sqrt{\cot(c+dx)}\sqrt{a+b \tan(c+dx)}} - \frac{2A\sqrt{\cot(c+dx)}}{ad\sqrt{a+b \tan(c+dx)}}$$

output `(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a-b)^(3/2)/d-(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a+b)^(3/2)/d-2*b*(A*a^2+2*A*b^2-B*a*b)/a^2/(a^2+b^2)/d/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2)-2*A*cot(d*x+c)^(1/2)/a/d/(a+b*tan(d*x+c))^(1/2)`

3.643.2 Mathematica [A] (verified)

Time = 2.53 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.00

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{\sqrt{\cot(c+dx)} \left(\frac{\sqrt[4]{-1} a \left(\frac{(a+ib)(A-ib) \arctan\left(\frac{\sqrt[4]{-1} \sqrt{-a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} \right) - (a-ib) \arctan\left(\frac{\sqrt[4]{-1} \sqrt{-a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{a^2+b^2} \right)}{\dots}$$

```
input Integrate[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2), x]
```

```
output (Sqrt[Cot[c + d*x]]*((( -1)^(1/4)*a*(((a + I*b)*(A - I*B)*ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[-a + I*b] - ((a - I*b)*(A + I*B)*ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]/(a^2 + b^2) - (2*A)/Sqrt[a + b*Tan[c + d*x]] - (2*b*(a^2*A + 2*A*b^2 - a*b*B)*Tan[c + d*x])/(a*(a^2 + b^2)*Sqrt[a + b*Tan[c + d*x]])))/(a*d)
```

3.643.3 Rubi [A] (verified)

Time = 1.57 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.09, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 4729, 3042, 4092, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx \xrightarrow{3042} \int \frac{\cot(c+dx)^{3/2}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx \xrightarrow{4729} \dots$$

$$\begin{aligned}
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^{3/2}(a+b \tan(c+dx))^{3/2}} dx \\
& \quad \downarrow \text{4092} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2 \int \frac{2Ab \tan^2(c+dx)+aA \tan(c+dx)+2Ab-aB}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{a} - \frac{2A}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{2Ab \tan^2(c+dx)+aA \tan(c+dx)+2Ab-aB}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{a} - \frac{2A}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{2Ab \tan(c+dx)^2+aA \tan(c+dx)+2Ab-aB}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{a} - \frac{2A}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \right) \\
& \quad \downarrow \text{4132} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\frac{2 \int \frac{(Ab-aB)a^2+(aA+bB) \tan(c+dx)a^2}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{a} - \frac{2A}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\frac{\int \frac{(Ab-aB)a^2+(aA+bB) \tan(c+dx)a^2}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{a} - \frac{2A}{ad\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} \right) \\
& \quad \downarrow \text{3042}
\end{aligned}$$

3.643. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{(Ab-aB)a^2+(aA+bB)\tan(c+dx)a^2}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \frac{2A}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} \right)$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} - \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{\frac{1}{2}a^2(b+ia)(A+iB)}{a^2+b^2} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} - \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{\frac{1}{2}a^2(b+ia)(A+iB)}{a^2+b^2} \right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} - \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{a^2(b+ia)(A+iB)}{a^2+b^2} \right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} - \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{a^2(b+ia)(A+iB)}{a^2+b^2} \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} - \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{a^2(b+ia)(A+iB)}{a^2+b^2} \right)$$

↓ 219

3.643. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{ad\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} - \frac{2b(a^2A-abB+2Ab^2)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{a^2(b+ia)(A+iB)}{\dots} \right)$$

input `Int[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A)/(a*d*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]]) - (((a^2*(I*a + b)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d) - (a^2*(I*a - b)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d))/(a*(a^2 + b^2)) + (2*b*(a^2*A + 2*A*b^2 - a*b*B)*Sqrt[Tan[c + d*x]])/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]))/a`

3.643.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

3.643. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`


```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.643.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.97 (sec) , antiderivative size = 1560397, normalized size of antiderivative = 6095.30

output too large to display

```
input int(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x)
```

```
output result too large to display
```

3.643.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18757 vs. 2(212) = 424.

Time = 7.14 (sec) , antiderivative size = 18757, normalized size of antiderivative = 73.27

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

3.643. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="fricas")`

output Too large to include

3.643.6 Sympy [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(A+B\tan(c+dx))\cot^{\frac{3}{2}}(c+dx)}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx$$

input `integrate(cot(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*cot(c + d*x)**(3/2)/(a + b*tan(c + d*x))**(3/2), x)`

3.643.7 Maxima [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{(b\tan(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(3/2)/(b*tan(d*x + c) + a)^(3/2), x)`

3.643.8 Giac [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{(b\tan(dx+c)+a)^{\frac{3}{2}}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(3/2)/(b*tan(d*x + c) + a)^(3/2), x)`

3.643.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{\cot(c+dx)^{3/2}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$$

input `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x)`

output `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x)`

3.644
$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.644.1 Optimal result 6149
 3.644.2 Mathematica [A] (verified) 6150
 3.644.3 Rubi [A] (verified) 6150
 3.644.4 Maple [B] (warning: unable to verify) 6154
 3.644.5 Fricas [B] (verification not implemented) 6154
 3.644.6 Sympy [F] 6154
 3.644.7 Maxima [F] 6155
 3.644.8 Giac [F(-2)] 6155
 3.644.9 Mupad [F(-1)] 6155

3.644.1 Optimal result

Integrand size = 35, antiderivative size = 215

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{(ia - b)^{3/2}d}$$

$$+ \frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)}}{(ia + b)^{3/2}d}$$

$$+ \frac{2b(Ab - aB)}{a(a^2 + b^2)d\sqrt{\cot(c+dx)}\sqrt{a+b \tan(c+dx)}}$$

output

```
(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a-b)^(3/2)/d+(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a+b)^(3/2)/d+2*b*(A*b-B*a)/a/(a^2+b^2)/d/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2)
```

3.644.2 Mathematica [A] (verified)

Time = 1.30 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.03

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \frac{\sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)} \left(\frac{\sqrt[4]{-1}(-ia+b)(A-iB) \arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+b}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{-a+ib}} \right)}{(a+b\tan(c+dx))^{3/2}}$$

input `Integrate[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2), x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((−1)^(1/4))*((−I)*a + b)*(A − I*B)*ArcTan[(((−1)^(1/4))*Sqrt[−a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/Sqrt[−a + I*b] + ((−1)^(1/4)*(a − I*b)*((−I)*A + B)*ArcTan[(((−1)^(1/4))*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/Sqrt[a + I*b] + (2*b*(A*b − a*B)*Sqrt[Tan[c + d*x]])/(a*Sqrt[a + b*Tan[c + d*x]])/((a^2 + b^2)*d)`

3.644.3 Rubi [A] (verified)

Time = 1.18 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.04, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 4729, 3042, 4092, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{4729} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B\tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \end{aligned}$$

3.644. $\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$

$$\begin{aligned} & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx \\ & \quad \downarrow \text{4092} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int \frac{a(aA+bB)-a(Ab-aB) \tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \right) \\ & \quad \downarrow \text{27} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(aA+bB)-a(Ab-aB) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \right) \\ & \quad \downarrow \text{3042} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{a(aA+bB)-a(Ab-aB) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} \right) \\ & \quad \downarrow \text{4099} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{\frac{1}{2}a(a-ib)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} \right) \\ & \quad \downarrow \text{3042} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{\frac{1}{2}a(a-ib)(A+iB) \int \frac{1-i \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} \right) \\ & \quad \downarrow \text{4098} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{\frac{a(a+ib)(A-iB) \int \frac{1}{(1-i \tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{2d}}{a(a^2+b^2)} \right) \\ & \quad \downarrow \text{104} \\ & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}} + \frac{\frac{a(a-ib)(A+iB) \int \frac{1}{\frac{(ia-b) \tan(c+dx)}{a+b \tan(c+dx)}+1} d - \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}}{d}}{a(a^2+b^2)} + \frac{a(a+ib)(A-iB) \int \frac{1}{(1-i \tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{2d}}{a(a^2+b^2)} \right) \end{aligned}$$

3.644. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2b(Ab - aB)\sqrt{\tan(c+dx)}}{ad(a^2 + b^2)\sqrt{a + b\tan(c+dx)}} + \frac{a(a+ib)(A-iB) \int \frac{1}{1 - \frac{(ia+b)\tan(c+dx)}{a+b\tan(c+dx)}} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}}{d} + \frac{a(a-iB)(A+iB)}{a(a^2 + b^2)} \right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2b(Ab - aB)\sqrt{\tan(c+dx)}}{ad(a^2 + b^2)\sqrt{a + b\tan(c+dx)}} + \frac{a(a-ib)(A+iB) \arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}} + \frac{a(a+ib)(A-iB)}{a(a^2 + b^2)} \right)$$

```
input Int[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2), x]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((a*(a - I*b)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d) + (a*(a + I*b)*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d))/(a*(a^2 + b^2)) + (2*b*(A*b - a*B)*Sqrt[Tan[c + d*x]])/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]))
```

3.644.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

```
rule 104 Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

3.644. $\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$

- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`
- rule 4098 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`
- rule 4099 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`
- rule 4729 `Int[(cot[(a_) + (b_)*(x_)])*(c_)^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.644.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.96 (sec) , antiderivative size = 1561166, normalized size of antiderivative = 7261.24

output too large to display

input `int(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x)`

output `result too large to display`

3.644.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18703 vs. 2(175) = 350.

Time = 7.11 (sec) , antiderivative size = 18703, normalized size of antiderivative = 86.99

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algorithm="fracas")`

output `Too large to include`

3.644.6 Sympy [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(A+B\tan(c+dx))\sqrt{\cot(c+dx)}}{(a+b\tan(c+dx))^{\frac{3}{2}}} dx$$

input `integrate(cot(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(cot(c + d*x))/(a + b*tan(c + d*x))**(3/2), x)`

3.644.7 Maxima [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{(B\tan(dx+c)+A)\sqrt{\cot(dx+c)}}{(b\tan(dx+c)+a)^{3/2}} dx$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(cot(d*x + c))/(b*tan(d*x + c) + a)^(3/
2), x)`

3.644.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x, algo
rithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(con
st gen &`

3.644.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$$

input `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),x
)`

output `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/2),
x)`

3.644. $\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx$

3.645
$$\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))^{3/2}} dx$$

3.645.1 Optimal result 6156
 3.645.2 Mathematica [A] (verified) 6157
 3.645.3 Rubi [A] (verified) 6157
 3.645.4 Maple [B] (warning: unable to verify) 6161
 3.645.5 Fracas [B] (verification not implemented) 6161
 3.645.6 Sympy [F] 6161
 3.645.7 Maxima [F] 6162
 3.645.8 Giac [F(-1)] 6162
 3.645.9 Mupad [F(-1)] 6162

3.645.1 Optimal result

Integrand size = 35, antiderivative size = 210

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))^{3/2}} dx =$$

$$\frac{(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{(ia - b)^{3/2}d}$$

$$+ \frac{(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{(ia + b)^{3/2}d}$$

$$- \frac{2(AB - aB)}{(a^2 + b^2) d \sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}}$$

```
output -(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot
(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a-b)^(3/2)/d+(A-I*B)*arctanh((I*a+b)^(1/
2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1
/2)/(I*a+b)^(3/2)/d-2*(A*b-B*a)/(a^2+b^2)/d/cot(d*x+c)^(1/2)/(a+b*tan(d*x+
c))^(1/2)
```

3.645.2 Mathematica [A] (verified)

Time = 2.35 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.23

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \frac{\sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(-\frac{\sqrt[4]{-1} a(a+ib)(A-iB) \arctan\left(\frac{\sqrt[4]{-1} \sqrt{\tan(c+dx)}}{\sqrt{a+ib}}\right)}{\sqrt{-a+ib}} \right)}{\dots}$$

```
input Integrate[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)),x]
```

```
output (Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((( (-1)^(1/4)*a*(a + I*b)*(A - I*B)*ArcTan[ ((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]] )/Sqrt[-a + I*b]) + ((-1)^(1/4)*a*(a - I*b)*(A + I*B)*ArcTan[ ((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]] )/Sqrt[a + I*b] + (2*b*(A*b - a*B)*Tan[c + d*x]^(3/2))/Sqrt[a + b*Tan[c + d*x]] + 2*(-(A*b) + a*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]] )/(a*(a^2 + b^2)*d)
```

3.645.3 Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.06, number of steps used = 13, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$, Rules used = {3042, 4729, 3042, 4091, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx$$

↓ 4729

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\sqrt{\tan(c + dx)}(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{3/2}} dx$$

$$\begin{aligned}
 & \int \frac{\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}}{b(a^2+b^2)} \left(-\frac{2\int \frac{-b(Ab-aB)+b(aA+bB)\tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{b(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \right) dx \\
 & \quad \downarrow \text{4091} \\
 & \int \frac{\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}}{b(a^2+b^2)} \left(\frac{\int \frac{b(Ab-aB)+b(aA+bB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{b(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \right) dx \\
 & \quad \downarrow \text{27} \\
 & \int \frac{\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}}{b(a^2+b^2)} \left(\frac{\int \frac{b(Ab-aB)+b(aA+bB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{b(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \right) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}}{b(a^2+b^2)} \left(\frac{\int \frac{b(Ab-aB)+b(aA+bB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{b(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \right) dx \\
 & \quad \downarrow \text{4099} \\
 & \int \frac{\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}}{b(a^2+b^2)} \left(-\frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{\frac{1}{2}b(b+ia)(A+iB)\int \frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{b(a^2+b^2)} \right) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}}{b(a^2+b^2)} \left(-\frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{\frac{1}{2}b(b+ia)(A+iB)\int \frac{1-i\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{b(a^2+b^2)} \right) dx \\
 & \quad \downarrow \text{4098} \\
 & \int \frac{\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}}{b(a^2+b^2)} \left(-\frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{\frac{b(b+ia)(A+iB)\int \frac{1}{(i\tan(c+dx)+1)\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{2d}}{b(a^2+b^2)} \right) dx \\
 & \quad \downarrow \text{104} \\
 & \int \frac{\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}}{b(a^2+b^2)} \left(-\frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{\frac{b(b+ia)(A+iB)\int \frac{1}{\frac{(ia-b)\tan(c+dx)}{a+b\tan(c+dx)}+1} d \frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}} dx}{d}}{b(a^2+b^2)} \right) dx
 \end{aligned}$$

3.645. $\int \frac{A+B\tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{3/2}} dx$

$$\downarrow 216$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}}+\frac{b(b+ia)(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}}-\frac{b(-b+ia)}{b(a^2+b^2)}\right)$$

$$\downarrow 219$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}}+\frac{b(b+ia)(A+iB)\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d\sqrt{-b+ia}}-\frac{b(-b+ia)}{b(a^2+b^2)}\right)$$

input `Int[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((b*(I*a + b)*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d) - ((I*a - b)*b*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(Sqrt[I*a + b]*d))/(b*(a^2 + b^2) - (2*(A*b - a*B)*Sqrt[Tan[c + d*x]])/((a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]))`

3.645.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

3.645. $\int \frac{A+B\tan(c+dx)}{\sqrt{\cot(c+dx)(a+b\tan(c+dx))^{3/2}}} dx$

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4091 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegerQ[2*m, 2*n])`

rule 4098 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4729 `Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.645.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.32 (sec) , antiderivative size = 1559507, normalized size of antiderivative = 7426.22

output too large to display

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x)`

output `result too large to display`

3.645.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18672 vs. 2(170) = 340.

Time = 7.01 (sec) , antiderivative size = 18672, normalized size of antiderivative = 88.91

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x, algorithm="fracas")`

output `Too large to include`

3.645.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{\frac{3}{2}} \sqrt{\cot(c + dx)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(1/2)/(a+b*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))**(3/2)*sqrt(cot(c + d*x))), x)`

3.645.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a)^{\frac{3}{2}} \sqrt{\cot(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x, algorith="maxima")`

output `integrate((B*tan(d*x + c) + A)/((b*tan(d*x + c) + a)^(3/2)*sqrt(cot(d*x + c))), x)`

3.645.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x, algorith="giac")`

output `Timed out`

3.645.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(3/2)),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(3/2)),x)`

3.646
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$$

3.646.1 Optimal result 6163
 3.646.2 Mathematica [A] (verified) 6164
 3.646.3 Rubi [A] (verified) 6164
 3.646.4 Maple [B] (warning: unable to verify) 6167
 3.646.5 Fricas [B] (verification not implemented) 6168
 3.646.6 Sympy [F] 6168
 3.646.7 Maxima [F] 6168
 3.646.8 Giac [F(-2)] 6169
 3.646.9 Mupad [F(-1)] 6169

3.646.1 Optimal result

Integrand size = 35, antiderivative size = 279

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx =$$

$$\frac{(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{(ia - b)^{3/2}d}$$

$$+ \frac{2B \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{b^{3/2}d}$$

$$- \frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{(ia + b)^{3/2}d}$$

$$+ \frac{2a(Ab - aB)}{b(a^2 + b^2) d \sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}}$$

output

```
-(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot
(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a-b)^(3/2)/d+2*B*arctanh(b^(1/2)*tan(d*x
+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/b^(3/2
)/d-(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))
*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a+b)^(3/2)/d+2*a*(A*b-B*a)/b/(a^2+b^
2)/d/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2)
```

3.646.2 Mathematica [A] (verified)

Time = 2.45 (sec) , antiderivative size = 361, normalized size of antiderivative = 1.29

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left(2\sqrt{a} \sqrt{-a + ib} \sqrt{a + ib} (a^2 + b^2) \operatorname{Barcsinh} \left(\frac{\sqrt{b} \sqrt{\tan(c + dx)}}{\sqrt{a}} \right) \sqrt{1 + \frac{b \tan(c + dx)}{a}} + \dots \right)$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2)),x]`

output `-((Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(2*Sqrt[a]*Sqrt[-a + I*b]*Sqrt[a + I*b]*(a^2 + b^2)*B*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a] + Sqrt[b]*((-1)^(1/4)*(a + I*b)^(3/2)*b*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]*Sqrt[a + b*Tan[c + d*x]] + Sqrt[-a + I*b]*(2*a*Sqrt[a + I*b]*(A*b - a*B)*Sqrt[Tan[c + d*x]] + (-1)^(1/4)*b*(I*a + b)*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]*Sqrt[a + b*Tan[c + d*x]]))/((-a + I*b)^(3/2)*(a + I*b)^(3/2)*b^(3/2)*d*Sqrt[a + b*Tan[c + d*x]])`

3.646.3 Rubi [A] (verified)

Time = 1.35 (sec) , antiderivative size = 274, normalized size of antiderivative = 0.98, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3042, 4729, 3042, 4088, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx \xrightarrow{3042} \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2}(a + b \tan(c + dx))^{3/2}} dx \xrightarrow{4729}$$

3.646. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$

$$\begin{aligned}
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\tan(c+dx)^{3/2}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx \\
& \quad \downarrow \text{4088} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int \frac{-((a^2+b^2)B\tan^2(c+dx)-b(Ab-aB)\tan(c+dx)+a(Ab-aB))}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{b(a^2+b^2)} + \frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \right) \\
& \quad \downarrow \text{27} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \frac{\int \frac{-((a^2+b^2)B\tan^2(c+dx)-b(Ab-aB)\tan(c+dx)+a(Ab-aB))}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}}{b(a^2+b^2)} \right) \\
& \quad \downarrow \text{3042} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \frac{\int \frac{-((a^2+b^2)B\tan(c+dx)^2)-b(Ab-aB)\tan(c+dx)+a(Ab-aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}}{b(a^2+b^2)} \right) \\
& \quad \downarrow \text{4138} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \frac{\int \frac{-((a^2+b^2)B\tan^2(c+dx))-b(Ab-aB)\tan(c+dx)+a(Ab-aB)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}}{bd(a^2+b^2)} \right) \\
& \quad \downarrow \text{2035} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \frac{2 \int \frac{-((a^2+b^2)B\tan^2(c+dx))-b(Ab-aB)\tan(c+dx)+a(Ab-aB)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)}}{bd(a^2+b^2)} \right) \\
& \quad \downarrow \text{2257} \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \frac{2 \int \left(\frac{b(aA+bB)-b(Ab-aB)\tan(c+dx)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} - \frac{(a^2+b^2)B}{\sqrt{a+b\tan(c+dx)}} \right)}{bd(a^2+b^2)} \right)
\end{aligned}$$

3.646. $\int \frac{A+B\tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}} dx$

↓ 2009

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} - \frac{2 \left(-\frac{B(a^2+b^2)\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{b}} + \frac{b(a-ib)(A+ib)}{bd(a^2+b^2)} \right)}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \right)$$

```
input Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2)),x
]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*((a - I*b)*b*(A + I*B)*ArcTan[
(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(2*Sqrt[I*a
- b]) - ((a^2 + b^2)*B*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan
[c + d*x]])/Sqrt[b] + ((a + I*b)*b*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*Sqrt[
Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(2*Sqrt[I*a + b])))/(b*(a^2 + b^
2)*d) + (2*a*(A*b - a*B)*Sqrt[Tan[c + d*x]])/(b*(a^2 + b^2)*d*Sqrt[a + b*T
an[c + d*x])))
```

3.646.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2035 Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]
```

```
rule 2257 Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol
] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a
, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]
```

3.646. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4138 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.646.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 2.03 (sec) , antiderivative size = 1560634, normalized size of antiderivative = 5593.67

output too large to display

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2), x)`

$$3.646. \quad \int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{\frac{3}{2}}} dx$$

output result too large to display

3.646.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18891 vs. 2(224) = 448.
Time = 11.54 (sec) , antiderivative size = 37815, normalized size of antiderivative = 135.54

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x, algorith="fricas")`

output Too large to include

3.646.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{\frac{3}{2}} \cot^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(3/2)/(a+b*tan(d*x+c))**(3/2),x)`

output `Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))**(3/2)*cot(c + d*x)**(3/2)), x)`

3.646.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a)^{\frac{3}{2}} \cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x, algorith="maxima")`

output `integrate((B*tan(d*x + c) + A)/((b*tan(d*x + c) + a)^(3/2)*cot(d*x + c)^(3/2)), x)`

3.646. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$

3.646.8 Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x, algo
rithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(con
st gen &`

3.646.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2} (a + b \tan(c + dx))^{3/2}} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(3/2)),x
)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(3/2)),
x)`

3.647 $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

3.647.1 Optimal result 6170
 3.647.2 Mathematica [A] (verified) 6171
 3.647.3 Rubi [A] (verified) 6171
 3.647.4 Maple [B] (warning: unable to verify) 6178
 3.647.5 Fricas [B] (verification not implemented) 6178
 3.647.6 Sympy [F(-1)] 6179
 3.647.7 Maxima [F] 6179
 3.647.8 Giac [F(-2)] 6180
 3.647.9 Mupad [F(-1)] 6180

3.647.1 Optimal result

Integrand size = 35, antiderivative size = 399

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{(ia-b)^{5/2}d}$$

$$+ \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{(ia+b)^{5/2}d}$$

$$+ \frac{2b(7a^2Ab+8Ab^3-3a^3B-4ab^2B)}{3a^3(a^2+b^2)d\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^{3/2}} + \frac{2(2Ab-aB)\sqrt{\cot(c+dx)}}{a^2d(a+b \tan(c+dx))^{3/2}}$$

$$- \frac{2A \cot^{\frac{3}{2}}(c+dx)}{3ad(a+b \tan(c+dx))^{3/2}} + \frac{2b(8a^4Ab+30a^2Ab^3+16Ab^5-3a^5B-17a^3b^2B-8ab^4B)}{3a^4(a^2+b^2)^2d\sqrt{\cot(c+dx)}\sqrt{a+b \tan(c+dx)}}$$

output

```
(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a-b)^(5/2)/d+(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a+b)^(5/2)/d+2/3*b*(8*A*a^4*b+30*A*a^2*b^3+16*A*b^5-3*B*a^5-17*B*a^3*b^2-8*B*a*b^4)/a^4/(a^2+b^2)^2/d/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2)-2/3*A*cot(d*x+c)^(3/2)/a/d/(a+b*tan(d*x+c))^(3/2)+2/3*b*(7*A*a^2*b+8*A*b^3-3*B*a^3-4*B*a*b^2)/a^3/(a^2+b^2)/d/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2)+2*(2*A*b-B*a)*cot(d*x+c)^(1/2)/a^2/d/(a+b*tan(d*x+c))^(3/2)
```

3.647. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

3.647.2 Mathematica [A] (verified)

Time = 4.44 (sec) , antiderivative size = 385, normalized size of antiderivative = 0.96

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{5}{2}}} dx = \sqrt{\cot(c+dx)} \left(\frac{6(6Ab-3aB)}{a(a+b \tan(c+dx))^{\frac{3}{2}}} - \frac{6A \cot(c+dx)}{(a+b \tan(c+dx))^{\frac{3}{2}}} + \frac{6b(7a^2Ab+8Ab^3)}{a^2(a^2+b^2)} \right)$$

input `Integrate[(Cot[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2), x]`

output `(Sqrt[Cot[c + d*x]]*((6*(6*A*b - 3*a*B))/(a*(a + b*Tan[c + d*x])^(3/2)) - (6*A*Cot[c + d*x])/(a + b*Tan[c + d*x])^(3/2) + (6*b*(7*a^2*A*b + 8*A*b^3 - 3*a^3*B - 4*a*b^2*B)*Tan[c + d*x])/(a^2*(a^2 + b^2)*(a + b*Tan[c + d*x])^(3/2)) + (Sqrt[Tan[c + d*x]]*(9*(-1)^(3/4)*a^4*((a + I*b)^2*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[-a + I*b] + ((a - I*b)^2*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[a + I*b]) + (6*b*(8*a^4*A*b + 30*a^2*A*b^3 + 16*A*b^5 - 3*a^5*B - 17*a^3*b^2*B - 8*a*b^4*B)*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]))/(a^3*(a^2 + b^2)^2))/(9*a*d)`

3.647.3 Rubi [A] (verified)

Time = 2.86 (sec) , antiderivative size = 445, normalized size of antiderivative = 1.12, number of steps used = 22, number of rules used = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 4729, 3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{5}{2}}} dx$$

↓ 3042

3.647. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{\frac{5}{2}}} dx$

$$\begin{aligned}
& \int \frac{\cot(c+dx)^{5/2}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx \\
& \quad \downarrow 4729 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan^{5/2}(c+dx)(a+b \tan(c+dx))^{5/2}} dx \\
& \quad \downarrow 3042 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^{5/2}(a+b \tan(c+dx))^{5/2}} dx \\
& \quad \downarrow 4092 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \left(-\frac{2 \int \frac{3(2Ab \tan^2(c+dx)+aA \tan(c+dx)+2Ab-aB)}{2 \tan^{3/2}(c+dx)(a+b \tan(c+dx))^{5/2}} dx}{3a} - \frac{2A}{3ad \tan^{3/2}(c+dx)(a+b \tan(c+dx))} \right) \\
& \quad \downarrow 27 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \left(-\frac{\int \frac{2Ab \tan^2(c+dx)+aA \tan(c+dx)+2Ab-aB}{\tan^{3/2}(c+dx)(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2A}{3ad \tan^{3/2}(c+dx)(a+b \tan(c+dx))^{3/2}} \right) \\
& \quad \downarrow 3042 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \left(-\frac{\int \frac{2Ab \tan(c+dx)^2+aA \tan(c+dx)+2Ab-aB}{\tan(c+dx)^{3/2}(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2A}{3ad \tan^{3/2}(c+dx)(a+b \tan(c+dx))^{3/2}} \right) \\
& \quad \downarrow 4132 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \left(-\frac{2 \int -\frac{Aa^2+B \tan(c+dx)a^2+4bBa-8Ab^2-4b(2Ab-aB) \tan^2(c+dx)}{2 \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2(2Ab-aB)}{ad \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \right) \\
& \quad \downarrow 27 \\
& \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \left(-\frac{\int \frac{Aa^2+B \tan(c+dx)a^2+4bBa-8Ab^2-4b(2Ab-aB) \tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2(2Ab-aB)}{ad \sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \right) \\
& \quad \downarrow 3042
\end{aligned}$$

3.647. $\int \frac{\cot^{5/2}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{Aa^2+B \tan(c+dx)a^2+4bBa-8Ab^2-4b(2Ab-aB) \tan(c+dx)^2}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2(2Ab-aB)}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int \frac{3Aa^4+9bBa^3-3(Ab-aB) \tan(c+dx)a^3-14Ab^2a^2+8b^3Ba-16Ab^4-2b(-3Ba^3+7Aba^2-4b^2Ba+8Ab^3) \tan^2(c+dx)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}}{3a(a^2+b^2)} - \frac{a}{a} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{3Aa^4+9bBa^3-3(Ab-aB) \tan(c+dx)a^3-14Ab^2a^2+8b^3Ba-16Ab^4-2b(-3Ba^3+7Aba^2-4b^2Ba+8Ab^3) \tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}}{3a(a^2+b^2)} - \frac{a}{a} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{3Aa^4+9bBa^3-3(Ab-aB) \tan(c+dx)a^3-14Ab^2a^2+8b^3Ba-16Ab^4-2b(-3Ba^3+7Aba^2-4b^2Ba+8Ab^3) \tan(c+dx)^2}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}}}{3a(a^2+b^2)} - \frac{a}{a} \right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int \frac{3(a^4(Aa^2+2bBa-Ab^2)-a^4(-Ba^2+2Aba+b^2B) \tan(c+dx))}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-3a^5B+8a^4Ab-17a^3b^2B+30a^2Ab^3-8ab^4)}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{3a(a^2+b^2)} - \frac{a}{a} \right)$$

3.647. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\begin{array}{c} \downarrow 27 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{3 \int \frac{a^4(Aa^2+2bBa-Ab^2)-a^4(-Ba^2+2Aba+b^2B)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{2b(-3a^5B+8a^4Ab-17a^3b^2B+30a^2Ab^3-8ab^4B)}{a(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{3a(a^2+b^2)} \right) \end{array}$$

$$\begin{array}{c} \downarrow 3042 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{3 \int \frac{a^4(Aa^2+2bBa-Ab^2)-a^4(-Ba^2+2Aba+b^2B)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{2b(-3a^5B+8a^4Ab-17a^3b^2B+30a^2Ab^3-8ab^4B)}{a(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{3a(a^2+b^2)} \right) \end{array}$$

$$\begin{array}{c} \downarrow 4099 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}} - \frac{2(2Ab-aB)}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} + \frac{2b(-3a^5B+8a^4Ab-17a^3b^2B+30a^2Ab^3-8ab^4B)}{a(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \right) \end{array}$$

$$\begin{array}{c} \downarrow 3042 \\ \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}} - \frac{2(2Ab-aB)}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} + \frac{2b(-3a^5B+8a^4Ab-17a^3b^2B+30a^2Ab^3-8ab^4B)}{a(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \right) \end{array}$$

$$\downarrow 4098$$

3.647. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} - \frac{2(2Ab-aB)}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} + \frac{2b(-3)}{\dots} \right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} - \frac{2(2Ab-aB)}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} + \frac{2b(-3)}{\dots} \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} - \frac{2(2Ab-aB)}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} + \frac{2b(-3)}{\dots} \right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{3ad \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} - \frac{2(2Ab-aB)}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} + \frac{-2b(-3}{\dots} \right)$$

```
input Int[(Cot[c + d*x]^(5/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x
]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A)/(3*a*d*Tan[c + d*x]^(3/2)*(a
+ b*Tan[c + d*x])^(3/2)) - ((-2*(2*A*b - a*B))/(a*d*Sqrt[Tan[c + d*x]]*(a
+ b*Tan[c + d*x])^(3/2)) + ((-2*b*(7*a^2*A*b + 8*A*b^3 - 3*a^3*B - 4*a*b^
2*B)*Sqrt[Tan[c + d*x]])/(3*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) +
((3*((a^4*(a - I*b)^2*(A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/
Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a - b]*d) + (a^4*(a + I*b)^2*(A - I*B)*
ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqr
t[I*a + b]*d)))/(a*(a^2 + b^2)) - (2*b*(8*a^4*A*b + 30*a^2*A*b^3 + 16*A*b^
5 - 3*a^5*B - 17*a^3*b^2*B - 8*a*b^4*B)*Sqrt[Tan[c + d*x]])/(a*(a^2 + b^2)
*d*Sqrt[a + b*Tan[c + d*x]]))/(3*a*(a^2 + b^2)))/a/a
```

3.647.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 104 Int[(((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_))/((e_) + (f_)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

3.647. $\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4092 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`
- rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`
- rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

3.647.
$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$


```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.647.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 6.02 (sec) , antiderivative size = 2981214, normalized size of antiderivative = 7471.71

output too large to display

```
input int(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x)
```

```
output result too large to display
```

3.647.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27987 vs. 2(340) = 680.

Time = 14.37 (sec) , antiderivative size = 27987, normalized size of antiderivative = 70.14

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

$$3.647. \quad \int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="fricas")`

output Too large to include

3.647.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output Timed out

3.647.7 Maxima [F]

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{5}{2}}}{(b\tan(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(5/2)/(b*tan(d*x + c) + a)^(5/
2), x)`

3.647.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx = \text{Exception raised: TypeError}$$

input `integrate(cot(d*x+c)^(5/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(con
st gen &`

3.647.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{5}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx = \int \frac{\cot(c+dx)^{\frac{5}{2}}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{\frac{5}{2}}} dx$$

input `int((cot(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x
)`

output `int((cot(c + d*x)^(5/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),
x)`

3.648
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

3.648.1 Optimal result 6181
 3.648.2 Mathematica [A] (verified) 6182
 3.648.3 Rubi [A] (verified) 6182
 3.648.4 Maple [B] (warning: unable to verify) 6188
 3.648.5 Fricas [B] (verification not implemented) 6188
 3.648.6 Sympy [F(-1)] 6189
 3.648.7 Maxima [F] 6189
 3.648.8 Giac [F] 6190
 3.648.9 Mupad [F(-1)] 6190

3.648.1 Optimal result

Integrand size = 35, antiderivative size = 341

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \frac{(iA-B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{(ia-b)^{5/2}d} - \frac{(iA+B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{(ia+b)^{5/2}d} - \frac{2b(3a^2A+4Ab^2-abB)}{3a^2(a^2+b^2)d\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^{3/2}} - \frac{2A\sqrt{\cot(c+dx)}}{ad(a+b \tan(c+dx))^{3/2}} - \frac{2b(3a^4A+17a^2Ab^2+8Ab^4-8a^3bB-2ab^3B)}{3a^3(a^2+b^2)^2d\sqrt{\cot(c+dx)}\sqrt{a+b \tan(c+dx)}}$$

output

```
(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a-b)^(5/2)/d-(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a+b)^(5/2)/d-2/3*b*(3*A*a^4+17*A*a^2*b^2+8*A*b^4-8*B*a^3*b-2*B*a*b^3)/a^3/(a^2+b^2)^2/d/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2)-2/3*b*(3*A*a^2+4*A*b^2-B*a*b)/a^2/(a^2+b^2)/d/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2)-2*A*cot(d*x+c)^(1/2)/a/d/(a+b*tan(d*x+c))^(3/2)
```

3.648.2 Mathematica [A] (verified)

Time = 4.43 (sec) , antiderivative size = 334, normalized size of antiderivative = 0.98

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \sqrt{\cot(c+dx)} \left(-\frac{6A}{(a+b \tan(c+dx))^{3/2}} - \frac{2b(3a^2A+4Ab^2-abB) \tan(c+dx)}{a(a^2+b^2)(a+b \tan(c+dx))^{3/2}} + \dots \right)$$

```
input Integrate[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2), x]
```

```
output (Sqrt[Cot[c + d*x]]*((-6*A)/(a + b*Tan[c + d*x])^(3/2) - (2*b*(3*a^2*A + 4*A*b^2 - a*b*B)*Tan[c + d*x])/(a*(a^2 + b^2)*(a + b*Tan[c + d*x])^(3/2)) + (Sqrt[Tan[c + d*x]]*(3*(-1)^(1/4)*a^3*(((a + I*b)^2*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/Sqrt[-a + I*b] - ((a - I*b)^2*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/Sqrt[a + I*b]) - (2*b*(3*a^4*A + 17*a^2*A*b^2 + 8*A*b^4 - 8*a^3*b*B - 2*a*b^3*B)*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]))/(a^2*(a^2 + b^2)^2))/(3*a*d)
```

3.648.3 Rubi [A] (verified)

Time = 2.22 (sec) , antiderivative size = 382, normalized size of antiderivative = 1.12, number of steps used = 19, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.514$, Rules used = {3042, 4729, 3042, 4092, 27, 3042, 4132, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx \xrightarrow{3042} \int \frac{\cot(c+dx)^{3/2}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

3.648. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \downarrow 4729 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx \\
 & \downarrow 3042 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\tan(c+dx)^{3/2}(a+b \tan(c+dx))^{5/2}} dx \\
 & \downarrow 4092 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2 \int \frac{4Ab \tan^2(c+dx)+aA \tan(c+dx)+4Ab-aB}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2A}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \right) \\
 & \downarrow 27 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{4Ab \tan^2(c+dx)+aA \tan(c+dx)+4Ab-aB}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2A}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \right) \\
 & \downarrow 3042 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{4Ab \tan(c+dx)^2+aA \tan(c+dx)+4Ab-aB}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx}{a} - \frac{2A}{ad\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} \right) \\
 & \downarrow 4132 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2 \int \frac{-3Ba^3+9Aba^2+3(aA+bB) \tan(c+dx)a^2-2b^2Ba+8Ab^3+2b(3Aa^2-bBa+4Ab^2) \tan^2(c+dx)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3a(a^2+b^2)} + \frac{2b(3a^2A-a)}{3ad(a^2+b^2)} \right) \\
 & \downarrow 27 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\int \frac{-3Ba^3+9Aba^2+3(aA+bB) \tan(c+dx)a^2-2b^2Ba+8Ab^3+2b(3Aa^2-bBa+4Ab^2) \tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3a(a^2+b^2)} + \frac{2b(3a^2A-a)}{3ad(a^2+b^2)} \right) \\
 & \downarrow 3042
 \end{aligned}$$

3.648. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(- \frac{\int \frac{-3Ba^3+9Aba^2+3(aA+bB)\tan(c+dx)a^2-2b^2Ba+8Ab^3+2b(3Aa^2-bBa+4Ab^2)\tan(c+dx)^2}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} dx}{3a(a^2+b^2)} + \frac{2b(3a^2A-abB)}{3ad(a^2+b^2)} \right)$$

↓ 4132

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(- \frac{2 \int \frac{3((-Ba^2+2Aba+b^2B)a^3+(Aa^2+2bBa-Ab^2)\tan(c+dx)a^3)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(3a^4A-8a^3bB+17a^2Ab^2-2ab^3B+8Ab^4)}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{3a(a^2+b^2)} \right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(- \frac{3 \int \frac{(-Ba^2+2Aba+b^2B)a^3+(Aa^2+2bBa-Ab^2)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(3a^4A-8a^3bB+17a^2Ab^2-2ab^3B+8Ab^4)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{3a(a^2+b^2)} \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(- \frac{3 \int \frac{(-Ba^2+2Aba+b^2B)a^3+(Aa^2+2bBa-Ab^2)\tan(c+dx)a^3}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(3a^4A-8a^3bB+17a^2Ab^2-2ab^3B+8Ab^4)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{3a(a^2+b^2)} \right)$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(- \frac{2A}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} - \frac{2b(3a^2A-abB+4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{2b(3a^4A-8a^3bB+17a^2Ab^2-2ab^3B+8Ab^4)}{3ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}} \right)$$

↓ 3042

3.648. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} - \frac{2b(3a^2A-abB+4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{2b(3a^4A-8a^2b^2B+4b^4)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} \right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} - \frac{2b(3a^2A-abB+4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{2b(3a^4A-8a^2b^2B+4b^4)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} \right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} - \frac{2b(3a^2A-abB+4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{2b(3a^4A-8a^2b^2B+4b^4)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} \right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} - \frac{2b(3a^2A-abB+4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{2b(3a^4A-8a^2b^2B+4b^4)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} \right)$$

↓ 219

3.648. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2A}{ad\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} - \frac{2b(3a^2A-abB+4Ab^2)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{2b(3a^4A-8a^3B+4a^2B^2)}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}} \right)$$

```
input Int[(Cot[c + d*x]^(3/2)*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x
]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*A)/(a*d*Sqrt[Tan[c + d*x]]*(a +
b*Tan[c + d*x])^(3/2)) - ((2*b*(3*a^2*A + 4*A*b^2 - a*b*B)*Sqrt[Tan[c + d
*x]])/(3*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) + ((3*((a^3*(a - I*b)
^2*(I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c +
d*x]]])/(Sqrt[I*a - b]*d) - (a^3*(a + I*b)^2*(I*A + B)*ArcTanh[(Sqrt[I*a +
b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d)))/(a*
(a^2 + b^2)) + (2*b*(3*a^4*A + 17*a^2*A*b^2 + 8*A*b^4 - 8*a^3*b*B - 2*a*b^
3*B)*Sqrt[Tan[c + d*x]])/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]))/(3*a*
(a^2 + b^2)))/a
```

3.648.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 104 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

3.648. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4098 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

3.648.
$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.648.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 2.84 (sec) , antiderivative size = 2976685, normalized size of antiderivative = 8729.28

output too large to display

```
input int(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x)
```

```
output result too large to display
```

3.648.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27817 vs. $2(289) = 578$.

Time = 14.41 (sec) , antiderivative size = 27817, normalized size of antiderivative = 81.57

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

3.648. $\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="fricas")`

output Too large to include

3.648.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output Timed out

3.648.7 Maxima [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{(b\tan(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(3/2)/(b*tan(d*x + c) + a)^(5/
2), x)`

3.648.8 Giac [F]

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(B\tan(dx+c)+A)\cot(dx+c)^{\frac{3}{2}}}{(b\tan(dx+c)+a)^{\frac{5}{2}}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*cot(d*x + c)^(3/2)/(b*tan(d*x + c) + a)^(5/2), x)`

3.648.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{\cot(c+dx)^{3/2}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$$

input `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)`

output `int((cot(c + d*x)^(3/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x)`

3.649
$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$$

3.649.1 Optimal result 6191
 3.649.2 Mathematica [A] (verified) 6192
 3.649.3 Rubi [A] (verified) 6192
 3.649.4 Maple [B] (warning: unable to verify) 6197
 3.649.5 Fracas [B] (verification not implemented) 6197
 3.649.6 Sympy [F] 6198
 3.649.7 Maxima [F] 6198
 3.649.8 Giac [F(-2)] 6199
 3.649.9 Mupad [F(-1)] 6199

3.649.1 Optimal result

Integrand size = 35, antiderivative size = 287

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx =$$

$$\frac{(A+iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{(ia-b)^{5/2}d}$$

$$- \frac{(A-iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{(ia+b)^{5/2}d}$$

$$+ \frac{2b(Ab-aB)}{3a(a^2+b^2)d\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^{3/2}}$$

$$+ \frac{2b(8a^2Ab+2Ab^3-5a^3B+ab^2B)}{3a^2(a^2+b^2)^2d\sqrt{\cot(c+dx)}\sqrt{a+b \tan(c+dx)}}$$

```
output -(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot
(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a-b)^(5/2)/d-(A-I*B)*arctanh((I*a+b)^(1/
2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1
/2)/(I*a+b)^(5/2)/d+2/3*b*(8*A*a^2*b+2*A*b^3-5*B*a^3+B*a*b^2)/a^2/(a^2+b^2
)^2/d/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2)+2/3*b*(A*b-B*a)/a/(a^2+b^2)/
d/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2)
```

3.649.2 Mathematica [A] (verified)

Time = 3.90 (sec) , antiderivative size = 293, normalized size of antiderivative = 1.02

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \frac{\sqrt{\cot(c+dx)}\sqrt{\tan(c+dx)} \left(-3\sqrt[4]{-1} \left(\frac{(a+ib)^2(iA+B)\arctan\left(\frac{\sqrt[4]{-1}}{\sqrt{-a+ib}}\right)}{\sqrt{-a+ib}} \right) \right)}{\dots}$$

input `Integrate[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2), x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-3*(-1)^(1/4)*((a + I*b)^2*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[-a + I*b] + (I*(a - I*b)^2*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[a + I*b] + (2*b*(a^2 + b^2)*(A*b - a*B)*Sqrt[Tan[c + d*x]])/(a*(a + b*Tan[c + d*x])^(3/2)) + (2*b*(8*a^2*A*b + 2*A*b^3 - 5*a^3*B + a*b^2*B)*Sqrt[Tan[c + d*x]])/(a^2*Sqrt[a + b*Tan[c + d*x]])))/(3*(a^2 + b^2)^2*d)`

3.649.3 Rubi [A] (verified)Time = 1.77 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.11, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 4729, 3042, 4092, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$$

↓ 4729

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B\tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{5/2}} dx$$

$$\begin{aligned}
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{A+B \tan(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{5/2}} dx \\
& \downarrow 4092 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int \frac{3Aa^2+bBa-3(Ab-aB) \tan(c+dx)a+2Ab^2+2b(Ab-aB) \tan^2(c+dx)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3a(a^2+b^2)} + \frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))} \right) \\
& \downarrow 27 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{3Aa^2+bBa-3(Ab-aB) \tan(c+dx)a+2Ab^2+2b(Ab-aB) \tan^2(c+dx)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3a(a^2+b^2)} + \frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))} \right) \\
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int \frac{3Aa^2+bBa-3(Ab-aB) \tan(c+dx)a+2Ab^2+2b(Ab-aB) \tan(c+dx)^2}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3a(a^2+b^2)} + \frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b \tan(c+dx))} \right) \\
& \downarrow 4132 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int \frac{3(a^2(Aa^2+2bBa-Ab^2)-a^2(-Ba^2+2Aba+b^2B) \tan(c+dx))}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(-5a^3B+8a^2Ab+ab^2B+2Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{3a(a^2+b^2)} \right) \\
& \downarrow 27 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{3 \int \frac{a^2(Aa^2+2bBa-Ab^2)-a^2(-Ba^2+2Aba+b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(-5a^3B+8a^2Ab+ab^2B+2Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{3a(a^2+b^2)} \right) \\
& \downarrow 3042 \\
& \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{3 \int \frac{a^2(Aa^2+2bBa-Ab^2)-a^2(-Ba^2+2Aba+b^2B) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} + \frac{2b(-5a^3B+8a^2Ab+ab^2B+2Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{3a(a^2+b^2)} \right)
\end{aligned}$$

3.649. $\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}}+\frac{2b(-5a^3B+8a^2Ab+ab^2B+2Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}+\frac{3\left(\frac{1}{2}a^2\right)}{3}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}}+\frac{2b(-5a^3B+8a^2Ab+ab^2B+2Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}+\frac{3\left(\frac{1}{2}a^2\right)}{3}\right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}}+\frac{2b(-5a^3B+8a^2Ab+ab^2B+2Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}+\frac{3\left(\frac{a^2}{2}\right)}{3}\right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}}+\frac{2b(-5a^3B+8a^2Ab+ab^2B+2Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}+\frac{3\left(\frac{a^2}{2}\right)}{3}\right)$$

↓ 216

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2b(Ab-aB)\sqrt{\tan(c+dx)}}{3ad(a^2+b^2)(a+b\tan(c+dx))^{3/2}}+\frac{2b(-5a^3B+8a^2Ab+ab^2B+2Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}+\frac{3\left(\frac{a^2}{2}\right)}{3}\right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2b(Ab - aB)\sqrt{\tan(c+dx)}}{3ad(a^2 + b^2)(a + b\tan(c+dx))^{3/2}} + \frac{2b(-5a^3B + 8a^2Ab + ab^2B + 2Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2 + b^2)\sqrt{a + b\tan(c+dx)}} + \dots \right)$$

```
input Int[(Sqrt[Cot[c + d*x]]*(A + B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(5/2),x
]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((2*b*(A*b - a*B)*Sqrt[Tan[c + d*x]]
)/(3*a*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) + ((3*((a^2*(a - I*b)^2*(
A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]
]])/(Sqrt[I*a - b]*d) + (a^2*(a + I*b)^2*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*
Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d)))/(a*(a^2
+ b^2)) + (2*b*(8*a^2*A*b + 2*A*b^3 - 5*a^3*B + a*b^2*B)*Sqrt[Tan[c + d*x
]])/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]))/(3*a*(a^2 + b^2))
```

3.649.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 104 Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))/((e_) + (f_)*(x
_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)
/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x]
/; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && L
tQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

```
rule 216 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4092 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[b*B*(b*c*(m + 1) + a*d*(n + 1)) + A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) - (A*b - a*B)*(b*c - a*d)*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n]) && !(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4098 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

```
rule 4132 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.)
+ (f_.)*(x_)]^2), x_Symbol] :> Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e +
f*x])^(m + 1)*((c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 +
b^2))), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e +
f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*
(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d
)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Ta
n[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ
[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] &&
!(ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.649.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 2.74 (sec) , antiderivative size = 2978354, normalized size of antiderivative = 10377.54

output too large to display

```
input int(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2), x)
```

```
output result too large to display
```

3.649.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27790 vs. 2(240) = 480.

Time = 14.62 (sec) , antiderivative size = 27790, normalized size of antiderivative = 96.83

$$\int \frac{\sqrt{\cot(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="fricas")`

output Too large to include

3.649.6 Sympy [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(A+B\tan(c+dx))\sqrt{\cot(c+dx)}}{(a+b\tan(c+dx))^{5/2}} dx$$

input `integrate(cot(d*x+c)**(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))*sqrt(cot(c + d*x))/(a + b*tan(c + d*x))**(5/
2), x)`

3.649.7 Maxima [F]

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{(B\tan(dx+c)+A)\sqrt{\cot(dx+c)}}{(b\tan(dx+c)+a)^{5/2}} dx$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*sqrt(cot(d*x + c))/(b*tan(d*x + c) + a)^(5/
2), x)`

3.649.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(cot(d*x+c)^(1/2)*(A+B*tan(d*x+c))/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(con
st gen &`

3.649.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx = \int \frac{\sqrt{\cot(c+dx)}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx$$

input `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),x
)`

output `int((cot(c + d*x)^(1/2)*(A + B*tan(c + d*x)))/(a + b*tan(c + d*x))^(5/2),
x)`

3.650 $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))}^{5/2}} dx$

3.650.1 Optimal result 6200
 3.650.2 Mathematica [A] (verified) 6201
 3.650.3 Rubi [A] (verified) 6201
 3.650.4 Maple [B] (warning: unable to verify) 6207
 3.650.5 Fricas [B] (verification not implemented) 6207
 3.650.6 Sympy [F] 6207
 3.650.7 Maxima [F] 6208
 3.650.8 Giac [F(-1)] 6208
 3.650.9 Mupad [F(-1)] 6208

3.650.1 Optimal result

Integrand size = 35, antiderivative size = 284

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))}^{5/2}} dx =$$

$$\frac{(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{(ia - b)^{5/2}d}$$

$$+ \frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{(ia + b)^{5/2}d}$$

$$- \frac{2(Ab - aB)}{3(a^2 + b^2)d\sqrt{\cot(c + dx)(a + b \tan(c + dx))}^{3/2}}$$

$$- \frac{2(5a^2Ab - Ab^3 - 2a^3B + 4ab^2B)}{3a(a^2 + b^2)^2 d\sqrt{\cot(c + dx)}\sqrt{a + b \tan(c + dx)}}$$

output

```
-(I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot
(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a-b)^(5/2)/d+(I*A+B)*arctanh((I*a+b)^(1/
2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1
/2)/(I*a+b)^(5/2)/d-2/3*(5*A*a^2*b-A*b^3-2*B*a^3+4*B*a*b^2)/a/(a^2+b^2)^2/
d/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2)-2/3*(A*b-B*a)/(a^2+b^2)/d/cot(d*
x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2)
```

3.650.2 Mathematica [A] (verified)

Time = 4.91 (sec) , antiderivative size = 340, normalized size of antiderivative = 1.20

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{5/2}} dx = \frac{\sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)} \left(\frac{2b(Ab - aB) \tan^{3/2}(c + dx)}{(a + b \tan(c + dx))^{3/2}} + \frac{6b(2aAb - a^2B)}{(a^2 + b^2)\sqrt{\cot(c + dx)}} \right)}{(a^2 + b^2)^{3/2}}$$

input `Integrate[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^(5/2)),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((2*b*(A*b - a*B)*Tan[c + d*x]^(3/2))/(a + b*Tan[c + d*x])^(3/2) + (6*b*(2*a*A*b - a^2*B + b^2*B)*Tan[c + d*x]^(3/2))/((a^2 + b^2)*Sqrt[a + b*Tan[c + d*x]]) + (3*(-((-1)^(1/4)*a*(a + I*b)^2*(A - I*B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[-a + I*b]) + ((-1)^(1/4)*a*(a - I*b)^2*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[a + I*b] + 2*(-2*a*A*b + a^2*B - b^2*B)*Sqrt[Tan[c + d*x]]*Sqrt[a + b*Tan[c + d*x]])/(a^2 + b^2))/(3*a*(a^2 + b^2)*d)`

3.650.3 Rubi [A] (verified)

Time = 1.72 (sec) , antiderivative size = 318, normalized size of antiderivative = 1.12, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 4729, 3042, 4091, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{5/2}} dx$$

3.650. $\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{5/2}} dx$

$$\begin{aligned}
 & \downarrow 4729 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx \\
 & \downarrow 3042 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\sqrt{\tan(c+dx)}(A+B \tan(c+dx))}{(a+b \tan(c+dx))^{5/2}} dx \\
 & \downarrow 4091 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2 \int -\frac{2b(Ab-aB) \tan^2(c+dx)+3b(aA+bB) \tan(c+dx)+b(Ab-aB)}{2\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3b(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{3d(a^2+b^2)(a+b \tan(c+dx))} \right) \\
 & \downarrow 27 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int -\frac{2b(Ab-aB) \tan^2(c+dx)+3b(aA+bB) \tan(c+dx)+b(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3b(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{3d(a^2+b^2)(a+b \tan(c+dx))} \right) \\
 & \downarrow 3042 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\int -\frac{2b(Ab-aB) \tan(c+dx)^2+3b(aA+bB) \tan(c+dx)+b(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^{3/2}} dx}{3b(a^2+b^2)} - \frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{3d(a^2+b^2)(a+b \tan(c+dx))} \right) \\
 & \downarrow 4132 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int \frac{3(ab(-Ba^2+2Aba+b^2B)+ab(Aa^2+2bBa-Ab^2) \tan(c+dx))}{2\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-2a^3B+5a^2Ab+4ab^2B-Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{3b(a^2+b^2)} \right) \\
 & \downarrow 27 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{3 \int \frac{ab(-Ba^2+2Aba+b^2B)+ab(Aa^2+2bBa-Ab^2) \tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b \tan(c+dx)}} dx}{a(a^2+b^2)} - \frac{2b(-2a^3B+5a^2Ab+4ab^2B-Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b \tan(c+dx)}}}{3b(a^2+b^2)} \right) \\
 & \downarrow 3042
 \end{aligned}$$

3.650. $\int \frac{A+B \tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b \tan(c+dx))^{5/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{3 \int \frac{ab(-Ba^2+2Aba+b^2B)+ab(Aa^2+2bBa-Ab^2)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx - \frac{2b(-2a^3B+5a^2Ab+4ab^2B-Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{3b(a^2+b^2)} \right)$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{3d(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{-\frac{2b(-2a^3B+5a^2Ab+4ab^2B-Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{3} + \dots \right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{3d(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{-\frac{2b(-2a^3B+5a^2Ab+4ab^2B-Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{3} + \dots \right)$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{3d(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{-\frac{2b(-2a^3B+5a^2Ab+4ab^2B-Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{3} + \dots \right)$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2(Ab-aB)\sqrt{\tan(c+dx)}}{3d(a^2+b^2)(a+b\tan(c+dx))^{3/2}} + \frac{-\frac{2b(-2a^3B+5a^2Ab+4ab^2B-Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{3} + \dots \right)$$

↓ 216

3.650. $\int \frac{A+B\tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{5/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2(Ab - aB)\sqrt{\tan(c+dx)}}{3d(a^2 + b^2)(a + b\tan(c+dx))^{3/2}} + \frac{-2b(-2a^3B + 5a^2Ab + 4ab^2B - Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2 + b^2)\sqrt{a + b\tan(c+dx)}} + \dots \right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{2(Ab - aB)\sqrt{\tan(c+dx)}}{3d(a^2 + b^2)(a + b\tan(c+dx))^{3/2}} + \frac{-2b(-2a^3B + 5a^2Ab + 4ab^2B - Ab^3)\sqrt{\tan(c+dx)}}{ad(a^2 + b^2)\sqrt{a + b\tan(c+dx)}} + \dots \right)$$

```
input Int[(A + B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^(5/2)),x
]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-2*(A*b - a*B)*Sqrt[Tan[c + d*x]])
/(3*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) + ((3*((a*(a - I*b)^2*b*(I*A
- B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])
/(Sqrt[I*a - b]*d) - (a*(a + I*b)^2*b*(I*A + B)*ArcTanh[(Sqrt[I*a + b]*Sqr
t[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d))/(a*(a^2 +
b^2)) - (2*b*(5*a^2*A*b - A*b^3 - 2*a^3*B + 4*a*b^2*B)*Sqrt[Tan[c + d*x]])
/(a*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]))/(3*b*(a^2 + b^2)))
```

3.650.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4091 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(A*b - a*B)*(a + b*Tan[e + f*x])^(m + 1)*((c + d*Tan[e + f*x])^n/(f*(m + 1)*(a^2 + b^2))), x] + Simp[1/(b*(m + 1)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n - 1)*Simp[b*B*(b*c*(m + 1) + a*d*n) + A*b*(a*c*(m + 1) - b*d*n) - b*(A*(b*c - a*d) - B*(a*c + b*d))*(m + 1)*Tan[e + f*x] - b*d*(A*b - a*B)*(m + n + 1)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && LtQ[0, n, 1] && (IntegerQ[m] || IntegerQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)]^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && ! (ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.650.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 2.01 (sec) , antiderivative size = 2978186, normalized size of antiderivative = 10486.57

output too large to display

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(5/2),x)`

output `result too large to display`

3.650.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27743 vs. 2(232) = 464.

Time = 14.81 (sec) , antiderivative size = 27743, normalized size of antiderivative = 97.69

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="fricas")`

output `Too large to include`

3.650.6 Sympy [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{(a + b \tan(c + dx))^{5/2} \sqrt{\cot(c + dx)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(1/2)/(a+b*tan(d*x+c))**(5/2),x)`

output `Integral((A + B*tan(c + d*x))/((a + b*tan(c + d*x))**(5/2)*sqrt(cot(c + d*
x))), x)`

3.650.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{5/2}} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a)^{5/2} \sqrt{\cot(dx + c)}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(5/2),x, algorith="maxima")`

output `integrate((B*tan(d*x + c) + A)/((b*tan(d*x + c) + a)^(5/2)*sqrt(cot(d*x + c))), x)`

3.650.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(5/2),x, algorith="giac")`

output `Timed out`

3.650.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{5/2}} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(5/2)),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(5/2)),x)`

3.651
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$$

3.651.1 Optimal result 6209
 3.651.2 Mathematica [A] (verified) 6210
 3.651.3 Rubi [A] (verified) 6210
 3.651.4 Maple [B] (warning: unable to verify) 6216
 3.651.5 Fricas [B] (verification not implemented) 6216
 3.651.6 Sympy [F(-1)] 6216
 3.651.7 Maxima [F] 6217
 3.651.8 Giac [F(-2)] 6217
 3.651.9 Mupad [F(-1)] 6217

3.651.1 Optimal result

Integrand size = 35, antiderivative size = 284

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \frac{(A + iB) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{(ia - b)^{5/2}d}$$

$$+ \frac{(A - iB) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{(ia + b)^{5/2}d}$$

$$+ \frac{2a(Ab - aB)}{3b(a^2 + b^2) d \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}}$$

$$+ \frac{2(2a^2 Ab - 4Ab^3 + a^3 B + 7ab^2 B)}{3b(a^2 + b^2)^2 d \sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}}$$

output

```
(A+I*B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a-b)^(5/2)/d+(A-I*B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a+b)^(5/2)/d+2/3*(2*A*a^2*b-4*A*b^3+B*a^3+7*B*a*b^2)/b/(a^2+b^2)^2/d/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2)+2/3*a*(A*b-B*a)/b/(a^2+b^2)/d/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2)
```


3.651.2 Mathematica [A] (verified)

Time = 4.01 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.15

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \frac{\sqrt{\cot(c + dx)}\sqrt{\tan(c + dx)} \left(-\frac{3B\sqrt{\tan(c+dx)}}{(a+b \tan(c+dx))^{3/2}} + \frac{(2aAb+a^2B+3b^2B)}{(a^2+b^2)(a+b \tan(c+dx))^{3/2}} \right)}{(a^2 + b^2)(a + b \tan(c + dx))^{3/2}}$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(5/2)),x]`

output `(Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-3*B*Sqrt[Tan[c + d*x]])/(a + b*Tan[c + d*x])^(3/2) + ((2*a*A*b + a^2*B + 3*b^2*B)*Sqrt[Tan[c + d*x]])/((a^2 + b^2)*(a + b*Tan[c + d*x])^(3/2)) + (3*(-1)^(1/4)*b*((a + I*b)^2*(I*A + B)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[-a + I*b] + (I*(a - I*b)^2*(A + I*B)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[a + I*b]) + (2*(2*a^2*A*b - 4*A*b^3 + a^3*B + 7*a*b^2*B)*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/(a^2 + b^2)^2)/(3*b*d)`

3.651.3 Rubi [A] (verified)

Time = 1.75 (sec) , antiderivative size = 314, normalized size of antiderivative = 1.11, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3042, 4729, 3042, 4088, 27, 3042, 4132, 27, 3042, 4099, 3042, 4098, 104, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx \xrightarrow{3042} \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2}(a + b \tan(c + dx))^{5/2}} dx$$

3.651. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$

$$\begin{aligned}
 & \downarrow 4729 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\tan^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx \\
 & \downarrow 3042 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\tan(c+dx)^{3/2}(A+B\tan(c+dx))}{(a+b\tan(c+dx))^{5/2}} dx \\
 & \downarrow 4088 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2 \int \frac{-((Ba^2+2Aba+3b^2B)\tan^2(c+dx))-3b(Ab-aB)\tan(c+dx)+a(Ab-aB)}{2\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}} dx}{3b(a^2+b^2)} + \frac{2a(Ab-a)}{3bd(a^2+b^2)} \right) \\
 & \downarrow 27 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \frac{\int \frac{-((Ba^2+2Aba+3b^2B)\tan^2(c+dx))-3b(Ab-aB)\tan(c+dx)+a(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}}{3b(a^2+b^2)} \right) \\
 & \downarrow 3042 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \frac{\int \frac{-((Ba^2+2Aba+3b^2B)\tan(c+dx)^2)-3b(Ab-aB)\tan(c+dx)+a(Ab-aB)}{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^{3/2}}}{3b(a^2+b^2)} \right) \\
 & \downarrow 4132 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \frac{2 \int \frac{3(ab(Aa^2+2bBa-Ab^2))-ab(-Ba^2+2Aba+b^2B)\tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}}{a(a^2+b^2)} \right) \\
 & \downarrow 27 \\
 & \sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \frac{3 \int \frac{ab(Aa^2+2bBa-Ab^2))-ab(-Ba^2+2Aba+b^2B)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}}{a(a^2+b^2)} \right) \\
 & \downarrow 3042
 \end{aligned}$$

3.651. $\int \frac{A+B\tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \frac{3 \int \frac{ab(Aa^2+2bBa-Ab^2)-ab(-Ba^2+2Aba+b^2B)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} dx}{a(a^2+b^2)} \right) - \frac{3b(a^2+b^2)}{3b(a^2+b^2)}$$

↓ 4099

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \frac{2(a^3B+2a^2Ab+7ab^2B-4Ab^3)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{3\left(\frac{1}{2}ab\right)}{3\left(\frac{1}{2}ab\right)} \right) - \frac{3b(a^2+b^2)}{3b(a^2+b^2)}$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \frac{2(a^3B+2a^2Ab+7ab^2B-4Ab^3)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{3\left(\frac{1}{2}ab\right)}{3\left(\frac{1}{2}ab\right)} \right) - \frac{3b(a^2+b^2)}{3b(a^2+b^2)}$$

↓ 4098

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \frac{2(a^3B+2a^2Ab+7ab^2B-4Ab^3)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{3\left(\frac{ab(c)}{2}\right)}{3\left(\frac{ab(c)}{2}\right)} \right) - \frac{3b(a^2+b^2)}{3b(a^2+b^2)}$$

↓ 104

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\sqrt{\tan(c+dx)}}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \frac{2(a^3B+2a^2Ab+7ab^2B-4Ab^3)\sqrt{\tan(c+dx)}}{d(a^2+b^2)\sqrt{a+b\tan(c+dx)}} + \frac{3\left(\frac{ab(c)}{2}\right)}{3\left(\frac{ab(c)}{2}\right)} \right) - \frac{3b(a^2+b^2)}{3b(a^2+b^2)}$$

↓ 216

3.651. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab - aB)\sqrt{\tan(c+dx)}}{3bd(a^2 + b^2)(a + b\tan(c+dx))^{3/2}} - \frac{2(a^3B + 2a^2Ab + 7ab^2B - 4Ab^3)\sqrt{\tan(c+dx)}}{d(a^2 + b^2)\sqrt{a + b\tan(c+dx)}} + \frac{ab(c+dx)}{3} \right)$$

↓ 219

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab - aB)\sqrt{\tan(c+dx)}}{3bd(a^2 + b^2)(a + b\tan(c+dx))^{3/2}} - \frac{2(a^3B + 2a^2Ab + 7ab^2B - 4Ab^3)\sqrt{\tan(c+dx)}}{d(a^2 + b^2)\sqrt{a + b\tan(c+dx)}} + \frac{ab(c+dx)}{3} \right)$$

```
input Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(5/2)),x
]
```

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((2*a*(A*b - a*B)*Sqrt[Tan[c + d*x]]
)/(3*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) - ((3*((a*(a - I*b)^2*b*(
A + I*B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]
]])/(Sqrt[I*a - b]*d) + (a*(a + I*b)^2*b*(A - I*B)*ArcTanh[(Sqrt[I*a + b]*
Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(Sqrt[I*a + b]*d)))/(a*(a^2
+ b^2)) - (2*(2*a^2*A*b - 4*A*b^3 + a^3*B + 7*a*b^2*B)*Sqrt[Tan[c + d*x]]
)/((a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]))/(3*b*(a^2 + b^2)))
```

3.651.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 104 `Int[(((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)))/((e_) + (f_)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`
- rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4088 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4098 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && EqQ[A^2 + B^2, 0]`

rule 4099 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[A^2 + B^2, 0]`

rule 4132 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)]) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol] := Simp[(A*b^2 - a*(b*B - a*C))*(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^(n + 1)/(f*(m + 1)*(b*c - a*d)*(a^2 + b^2)), x] + Simp[1/((m + 1)*(b*c - a*d)*(a^2 + b^2)) Int[(a + b*Tan[e + f*x])^(m + 1)*(c + d*Tan[e + f*x])^n*Simp[A*(a*(b*c - a*d)*(m + 1) - b^2*d*(m + n + 2)) + (b*B - a*C)*(b*c*(m + 1) + a*d*(n + 1)) - (m + 1)*(b*c - a*d)*(A*b - a*B - b*C)*Tan[e + f*x] - d*(A*b^2 - a*(b*B - a*C))*(m + n + 2)*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && LtQ[m, -1] && (!ILtQ[n, -1] && (!IntegerQ[m] || (EqQ[c, 0] && NeQ[a, 0])))`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.651.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 2.70 (sec) , antiderivative size = 2975109, normalized size of antiderivative = 10475.74

output too large to display

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(5/2),x)`

output `result too large to display`

3.651.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27726 vs. $2(236) = 472$.

Time = 14.56 (sec) , antiderivative size = 27726, normalized size of antiderivative = 97.63

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(5/2),x, algorith="fracas")`

output `Too large to include`

3.651.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(3/2)/(a+b*tan(d*x+c))**(5/2),x)`

output `Timed out`

3.651.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a)^{\frac{5}{2}} \cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(5/2),x, algorith="maxima")`

output `integrate((B*tan(d*x + c) + A)/((b*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^(3/2)), x)`

3.651.8 Giac [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(5/2),x, algorith="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen &`

3.651.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{3/2} (a + b \tan(c + dx))^{5/2}} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(5/2)),x)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(5/2)),x)`

3.652
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{\frac{5}{2}}} dx$$

3.652.1 Optimal result 6218
 3.652.2 Mathematica [A] (verified) 6219
 3.652.3 Rubi [A] (verified) 6220
 3.652.4 Maple [B] (warning: unable to verify) 6225
 3.652.5 Fricas [B] (verification not implemented) 6225
 3.652.6 Sympy [F(-1)] 6225
 3.652.7 Maxima [F] 6226
 3.652.8 Giac [F(-1)] 6226
 3.652.9 Mupad [F(-1)] 6226

3.652.1 Optimal result

Integrand size = 35, antiderivative size = 342

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{\frac{5}{2}}} dx = \frac{(iA - B) \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b}\tan(c+dx)}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{(ia - b)^{\frac{5}{2}}d}$$

$$+ \frac{2B \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b}\tan(c+dx)}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{b^{\frac{5}{2}}d}$$

$$- \frac{(iA + B) \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b}\tan(c+dx)}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{(ia + b)^{\frac{5}{2}}d}$$

$$+ \frac{2a(Ab - aB)}{3b(a^2 + b^2) d \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{\frac{3}{2}}}$$

$$+ \frac{2a(2Ab^3 - a(a^2 + 3b^2) B)}{b^2(a^2 + b^2)^2 d \sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}}$$

```
output (I*A-B)*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(
d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a-b)^(5/2)/d+2*B*arctanh(b^(1/2)*tan(d*x+
c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/b^(5/2)
/d-(I*A+B)*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*
cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/(I*a+b)^(5/2)/d+2*a*(2*A*b^3-a*(a^2+3*b^
2)*B)/b^2/(a^2+b^2)^2/d/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2)+2/3*a*(A*b
-B*a)/b/(a^2+b^2)/d/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2)
```

3.652.
$$\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{\frac{5}{2}}} dx$$

3.652.2 Mathematica [A] (verified)

Time = 6.44 (sec) , antiderivative size = 617, normalized size of antiderivative = 1.80

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left((A - iB) \left(\frac{3 \sqrt[4]{-1} \arctan\left(\frac{\sqrt[4]{-1} \sqrt{a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(-a+ib)^{5/2}} \right. \right.$$

$$\left. \left. (A + iB) \left(\frac{3 \sqrt[4]{-1} \arctan\left(\frac{\sqrt[4]{-1} \sqrt{a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{(a+ib)^{5/2}} - \frac{\tan^{\frac{3}{2}}(c+dx)}{(a+ib)(a+b \tan(c+dx))^{3/2}} - \frac{3i \sqrt{\tan(c+dx)}}{(a+ib)^2 \sqrt{a+b \tan(c+dx)}} \right) \right.$$

$$\left. + \frac{3d}{3b^3 d \sqrt{\tan(c + dx)}} \left((iA - B) \sqrt{a + b \tan(c + dx)} \left(\frac{b^2 \tan^2(c+dx)}{(a+b \tan(c+dx))^2} + \frac{3b \tan(c+dx)}{a+b \tan(c+dx)} - \frac{3\sqrt{b} \operatorname{arcsinh}\left(\frac{\sqrt{b} \sqrt{\tan(c+dx)}}{\sqrt{a}}\right) \sqrt{\tan(c+dx)}}{\sqrt{a} \sqrt{1 + \frac{b \tan(c+dx)}{a}}}\right) \right.$$

$$\left. \left. (iA + B) \sqrt{a + b \tan(c + dx)} \left(\frac{b^2 \tan^2(c+dx)}{(a+b \tan(c+dx))^2} + \frac{3b \tan(c+dx)}{a+b \tan(c+dx)} - \frac{3\sqrt{b} \operatorname{arcsinh}\left(\frac{\sqrt{b} \sqrt{\tan(c+dx)}}{\sqrt{a}}\right) \sqrt{\tan(c+dx)}}{\sqrt{a} \sqrt{1 + \frac{b \tan(c+dx)}{a}}}\right) \right) \right)$$

input `Integrate[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^(5/2)),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((A - I*B)*((3*(-1)^(1/4)*ArcTan[((-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(-a + I*b)^(5/2) + Tan[c + d*x]^(3/2)/((a - I*b)*(a + b*Tan[c + d*x])^(3/2)) - ((3*I)*Sqrt[Tan[c + d*x]])/((a - I*b)^2*Sqrt[a + b*Tan[c + d*x]])))/(3*d) - ((A + I*B)*((3*(-1)^(1/4)*ArcTan[((-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])]/(a + I*b)^(5/2) - Tan[c + d*x]^(3/2)/((a + I*b)*(a + b*Tan[c + d*x])^(3/2)) - ((3*I)*Sqrt[Tan[c + d*x]])/((a + I*b)^2*Sqrt[a + b*Tan[c + d*x]])))/(3*d) + ((I*A - B)*Sqrt[a + b*Tan[c + d*x]]*((b^2*Tan[c + d*x]^2)/(a + b*Tan[c + d*x])^2 + (3*b*Tan[c + d*x])/(a + b*Tan[c + d*x]) - (3*Sqrt[b]*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])*Sqrt[Tan[c + d*x]])/(Sqrt[a]*Sqrt[1 + (b*Tan[c + d*x])/a]))/(3*b^3*d*Sqrt[Tan[c + d*x]]) - ((I*A + B)*Sqrt[a + b*Tan[c + d*x]]*((b^2*Tan[c + d*x]^2)/(a + b*Tan[c + d*x])^2 + (3*b*Tan[c + d*x])/(a + b*Tan[c + d*x]) - (3*Sqrt[b]*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]])*Sqrt[Tan[c + d*x]])/(Sqrt[a]*Sqrt[1 + (b*Tan[c + d*x])/a]))/(3*b^3*d*Sqrt[Tan[c + d*x]]))`

3.652.3 Rubi [A] (verified)

Time = 2.00 (sec) , antiderivative size = 365, normalized size of antiderivative = 1.07, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.371$, Rules used = {3042, 4729, 3042, 4088, 27, 3042, 4128, 27, 3042, 4138, 2035, 2257, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{\frac{5}{2}}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{\frac{5}{2}}(a + b \tan(c + dx))^{\frac{5}{2}}} dx \\
 & \quad \downarrow \text{4729} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\tan^{\frac{5}{2}}(c + dx)(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{\frac{5}{2}}} dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\tan(c + dx)^{\frac{5}{2}}(A + B \tan(c + dx))}{(a + b \tan(c + dx))^{\frac{5}{2}}} dx \\
 & \quad \downarrow \text{4088}
 \end{aligned}$$

3.652. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{\frac{5}{2}}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2\int\frac{-3\sqrt{\tan(c+dx)}(-((a^2+b^2)B\tan^2(c+dx))-b(Ab-aB)\tan(c+dx)+a(Ab-aB))}{2(a+b\tan(c+dx))^{3/2}}dx}{3b(a^2+b^2)}+\frac{2a(Ab-aB)}{3bd(a^2+b^2)}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}}-\frac{\int\frac{\sqrt{\tan(c+dx)}(-((a^2+b^2)B\tan^2(c+dx))-b(Ab-aB)\tan(c+dx)+a(Ab-aB))}{(a+b\tan(c+dx))^{3/2}}dx}{b(a^2+b^2)}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}}-\frac{\int\frac{\sqrt{\tan(c+dx)}(-((a^2+b^2)B\tan^2(c+dx))-b(Ab-aB)\tan(c+dx)+a(Ab-aB))}{(a+b\tan(c+dx))^{3/2}}dx}{b(a^2+b^2)}\right)$$

↓ 4128

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}}-\frac{2\int\frac{(Aa^2+2bBa-Ab^2)\tan(c+dx)b^2-(a^2+b^2)^2B\tan^2(c+dx)+a(Ab-aB)\tan(c+dx)}{2\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx}{b(a^2+b^2)}\right)$$

↓ 27

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}}-\frac{\int\frac{(Aa^2+2bBa-Ab^2)\tan(c+dx)b^2-(a^2+b^2)^2B\tan^2(c+dx)+a(Ab-aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx}{b(a^2+b^2)}\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{2a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}}-\frac{\int\frac{(Aa^2+2bBa-Ab^2)\tan(c+dx)b^2-(a^2+b^2)^2B\tan^2(c+dx)+a(Ab-aB)\tan(c+dx)}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}}dx}{b(a^2+b^2)}\right)$$

↓ 4138

3.652. $\int \frac{A+B\tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^{5/2}} dx$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \frac{\int \frac{(Aa^2+2bBa-Ab^2)\tan(c+dx)b^2-(a^2+b^2)^2B\tan^2(c+dx)+a}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} dx}{bd(a^2+b^2)} \right)$$

↓ 2035

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \frac{2\int \frac{(Aa^2+2bBa-Ab^2)\tan(c+dx)b^2-(a^2+b^2)^2B\tan^2(c+dx)+a}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} dx}{bd(a^2+b^2)} \right)$$

↓ 2257

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \frac{2\int \left(\frac{(-Ba^2+2Aba+b^2B)b^2+(Aa^2+2bBa-Ab^2)\tan(c+dx)b^2}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} \right) dx}{bd(a^2+b^2)} \right)$$

↓ 2009

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{2a(Ab-aB)\tan^{\frac{3}{2}}(c+dx)}{3bd(a^2+b^2)(a+b\tan(c+dx))^{3/2}} - \frac{-\frac{2a(2Ab^3-aB(a^2+3b^2))\sqrt{\tan(c+dx)}}{bd(a^2+b^2)\sqrt{a+b\tan(c+dx)}}}{bd(a^2+b^2)} + \frac{2\left(-\frac{B(a^2+b^2)}{bd(a^2+b^2)}\right)}{bd(a^2+b^2)} \right)$$

input `Int[(A + B*Tan[c + d*x])/(Cot[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^(5/2)),x]`

```
output Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((2*a*(A*b - a*B)*Tan[c + d*x]^(3/2)
)/(3*b*(a^2 + b^2)*d*(a + b*Tan[c + d*x])^(3/2)) - ((2*(((a - I*b)^2*b^2*(
I*A - B)*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]
]])/(2*Sqrt[I*a - b]) - ((a^2 + b^2)^2*B*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]
]]/Sqrt[a + b*Tan[c + d*x]]])/Sqrt[b] - ((a + I*b)^2*b^2*(I*A + B)*ArcTan
h[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]]])/(2*Sqrt[I*
a + b]))/(b*(a^2 + b^2)*d) - (2*a*(2*A*b^3 - a*(a^2 + 3*b^2)*B)*Sqrt[Tan[
c + d*x]]/(b*(a^2 + b^2)*d*Sqrt[a + b*Tan[c + d*x]]))/(b*(a^2 + b^2)))
```

3.652.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2035 Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]
```

```
rule 2257 Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol
] := Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x] /; FreeQ[{a
, c, d, e, q}, x] && PolyQ[Px, x] && IntegerQ[p]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4088 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(b*c - a*d)*(B*c - A*d)*(a + b*Tan[e + f*x])^(m - 1)*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 2)*(c + d*Tan[e + f*x])^(n + 1)*Simp[a*A*d*(b*d*(m - 1) - a*c*(n + 1)) + (b*B*c - (A*b + a*B)*d)*(b*c*(m - 1) + a*d*(n + 1)) - d*((a*A - b*B)*(b*c - a*d) + (A*b + a*B)*(a*c + b*d))*(n + 1)*Tan[e + f*x] - b*(d*(A*b*c + a*B*c - a*A*d)*(m + n) - b*B*(c^2*(m - 1) - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[m] || IntegersQ[2*m, 2*n])`

rule 4128 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := Simp[(A*d^2 + c*(c*C - B*d))*(a + b*Tan[e + f*x])^m*((c + d*Tan[e + f*x])^(n + 1)/(d*f*(n + 1)*(c^2 + d^2))), x] - Simp[1/(d*(n + 1)*(c^2 + d^2)) Int[(a + b*Tan[e + f*x])^(m - 1)*(c + d*Tan[e + f*x])^(n + 1)*Simp[A*d*(b*d*m - a*c*(n + 1)) + (c*C - B*d)*(b*c*m + a*d*(n + 1)) - d*(n + 1)*((A - C)*(b*c - a*d) + B*(a*c + b*d))*Tan[e + f*x] - b*(d*(B*c - A*d)*(m + n + 1) - C*(c^2*m - d^2*(n + 1)))*Tan[e + f*x]^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0] && GtQ[m, 0] && LtQ[n, -1]`

rule 4138 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)] + (C_.)*tan[(e_.) + (f_.)*(x_)]^2), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*(c + d*ff*x)^n*((A + B*ff*x + C*ff^2*x^2)/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, A, B, C, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.652.4 Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 2.86 (sec) , antiderivative size = 2981291, normalized size of antiderivative = 8717.23

output too large to display

input `int((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c))^(5/2),x)`

output `result too large to display`

3.652.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 28011 vs. $2(284) = 568$.

Time = 24.50 (sec) , antiderivative size = 56055, normalized size of antiderivative = 163.90

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{\frac{5}{2}}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c))^(5/2),x, algorithm="fracas")`

output `Too large to include`

3.652.6 Sympy [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)**(5/2)/(a+b*tan(d*x+c))**(5/2),x)`

output `Timed out`

3.652.7 Maxima [F]

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \int \frac{B \tan(dx + c) + A}{(b \tan(dx + c) + a)^{\frac{5}{2}} \cot(dx + c)^{\frac{5}{2}}} dx$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="maxima")`

output `integrate((B*tan(d*x + c) + A)/((b*tan(d*x + c) + a)^(5/2)*cot(d*x + c)^(5
/2)), x)`

3.652.8 Giac [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(d*x+c))/cot(d*x+c)^(5/2)/(a+b*tan(d*x+c))^(5/2),x, algo
rithm="giac")`

output `Timed out`

3.652.9 Mupad [F(-1)]

Timed out.

$$\int \frac{A + B \tan(c + dx)}{\cot^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^{5/2}} dx = \int \frac{A + B \tan(c + dx)}{\cot(c + dx)^{5/2} (a + b \tan(c + dx))^{5/2}} dx$$

input `int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + b*tan(c + d*x))^(5/2)),x
)`

output `int((A + B*tan(c + d*x))/(cot(c + d*x)^(5/2)*(a + b*tan(c + d*x))^(5/2)),
x)`

3.652. $\int \frac{A+B \tan(c+dx)}{\cot^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^{5/2}} dx$

3.653
$$\int \frac{\sqrt{\cot(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$$

3.653.1 Optimal result	6227
3.653.2 Mathematica [A] (verified)	6227
3.653.3 Rubi [A] (verified)	6228
3.653.4 Maple [B] (warning: unable to verify)	6230
3.653.5 Fricas [B] (verification not implemented)	6231
3.653.6 Sympy [F]	6232
3.653.7 Maxima [F]	6233
3.653.8 Giac [F(-2)]	6233
3.653.9 Mupad [F(-1)]	6233

3.653.1 Optimal result

Integrand size = 38, antiderivative size = 151

$$\int \frac{\sqrt{\cot(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{B \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{\sqrt{ia-bd}} + \frac{B \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c+dx)} \sqrt{\tan(c+dx)}}{\sqrt{ia+bd}}$$

output

```
B*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*a-b)^(1/2)+B*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*a+b)^(1/2)
```

3.653.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.96

$$\int \frac{\sqrt{\cot(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx = \frac{(-1)^{3/4} B \left(-\frac{\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} - \frac{\arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{a+ib}} \right)}{d}$$

input `Integrate[(Sqrt[Cot[c + d*x]]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]`

output `((-1)^(3/4)*B*(-(ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]]/Sqrt[-a + I*b]) - ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]]/Sqrt[a + I*b])*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]/d`

3.653.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.85, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.184$, Rules used = {2011, 3042, 4729, 3042, 4058, 615, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\cot(c+dx)}(aB + bB \tan(c+dx))}{(a + b \tan(c+dx))^{3/2}} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{\sqrt{\cot(c+dx)}}{\sqrt{a + b \tan(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{\sqrt{\cot(c+dx)}}{\sqrt{a + b \tan(c+dx)}} dx \\
 & \quad \downarrow \text{4729} \\
 & B \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \int \frac{1}{\sqrt{\tan(c+dx)} \sqrt{a + b \tan(c+dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \int \frac{1}{\sqrt{\tan(c+dx)} \sqrt{a + b \tan(c+dx)}} dx \\
 & \quad \downarrow \text{4058} \\
 & \frac{B \sqrt{\tan(c+dx)} \sqrt{\cot(c+dx)} \int \frac{1}{\sqrt{\tan(c+dx)} \sqrt{a + b \tan(c+dx)} (\tan^2(c+dx) + 1)} d \tan(c+dx)}{d} \\
 & \quad \downarrow \text{615}
 \end{aligned}$$

3.653. $\int \frac{\sqrt{\cot(c+dx)}(aB + bB \tan(c+dx))}{(a + b \tan(c+dx))^{3/2}} dx$

$$\frac{B\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \left(\frac{i}{2(i-\tan(c+dx))\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} + \frac{i}{2\sqrt{\tan(c+dx)}(\tan(c+dx)+i)\sqrt{a+b\tan(c+dx)}} \right) dx}{d}$$

↓ 2009

$$\frac{B\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(\frac{\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{-b+ia}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{b+ia}} \right)}{d}$$

```
input Int[(Sqrt[Cot[c + d*x]]*(a*B + b*B*Tan[c + d*x]))/(a + b*Tan[c + d*x])^(3/2),x]
```

```
output (B*(ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[I*a - b] + ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[I*a + b])*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]])/d
```

3.653.3.1 Defintions of rubi rules used

```
rule 615 Int[((e_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(e*x)^m*(c + d*x)^n*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && ILtQ[p, 0]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2011 Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4058 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^m*((c + d*ff*x)^n/(1 + ff^2*x^2)), x], x,
Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]
```

```
rule 4729 Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] :> Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]
```

3.653.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 761 vs. 2(123) = 246.

Time = 16.24 (sec) , antiderivative size = 762, normalized size of antiderivative = 5.05

method	result
default	$B \sin(dx+c) \left(\sqrt{b+\sqrt{a^2+b^2}} \ln \left(\frac{a \cot(dx+c) \cos(dx+c) - 2a \cot(dx+c) - 2\sqrt{(\cot(dx+c)^2 a - 2a \cot(dx+c) \csc(dx+c) + a \csc(dx+c)^2 - 2b \csc(dx+c))}}{\dots} \right) \right)$

```
input int(cot(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,method=
_RETURNVERBOSE)
```

3.653. $\int \frac{\sqrt{\cot(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

output $\frac{1}{2}B/d/(a^2+b^2)^{1/2}\sin(dx+c)*((b+(a^2+b^2)^{1/2})^{1/2})*\ln((a*\cot(dx+c)*\cos(dx+c)-2*a*\cot(dx+c)-2*((\cot(dx+c)^2*a-2*a*\cot(dx+c)*\csc(dx+c)+a*\csc(dx+c)^2-2*b*\csc(dx+c)+2*\cot(dx+c)*b-a)*(\cos(dx+c)-1)*\csc(dx+c))^{1/2}*(b+(a^2+b^2)^{1/2})^{1/2}\sin(dx+c)+a*\csc(dx+c)+2*(a^2+b^2)^{1/2}\cos(dx+c)+2*b*\cos(dx+c)-a*\sin(dx+c)-2*(a^2+b^2)^{1/2}-2*b)/(\cos(dx+c)-1))-2*(-b+(a^2+b^2)^{1/2})^{1/2}*\arctan(1/(-b+(a^2+b^2)^{1/2})^{1/2})*((b+(a^2+b^2)^{1/2})^{1/2}\cos(dx+c)-(-2*(\cos(dx+c)^2*b-\cos(dx+c)*\sin(dx+c)*a-b)/(\cos(dx+c)+1)^{1/2}\sin(dx+c)-(b+(a^2+b^2)^{1/2})^{1/2}))/(\cos(dx+c)-1)-(b+(a^2+b^2)^{1/2})^{1/2}*\ln((a*\cot(dx+c)*\cos(dx+c)-2*a*\cot(dx+c)+2*((\cot(dx+c)^2*a-2*a*\cot(dx+c)*\csc(dx+c)+a*\csc(dx+c)^2-2*b*\csc(dx+c)+2*\cot(dx+c)*b-a)*(\cos(dx+c)-1)*\csc(dx+c))^{1/2}*(b+(a^2+b^2)^{1/2})^{1/2}\sin(dx+c)+a*\csc(dx+c)+2*(a^2+b^2)^{1/2}\cos(dx+c)+2*b*\cos(dx+c)-a*\sin(dx+c)-2*(a^2+b^2)^{1/2}-2*b)/(\cos(dx+c)-1))+2*(-b+(a^2+b^2)^{1/2})^{1/2}*\arctan(1/(-b+(a^2+b^2)^{1/2})^{1/2})*((b+(a^2+b^2)^{1/2})^{1/2}\cos(dx+c)+(-2*(\cos(dx+c)^2*b-\cos(dx+c)*\sin(dx+c)*a-b)/(\cos(dx+c)+1)^{1/2}\sin(dx+c)-(b+(a^2+b^2)^{1/2})^{1/2}))/(\cos(dx+c)-1))*\cot(dx+c)^{1/2}*(a+b*\tan(dx+c))^{1/2}/(\cos(dx+c)+1)/(-2*(\cos(dx+c)^2*b-\cos(dx+c)*\sin(dx+c)*a-b)/(\cos(dx+c)+1)^{1/2})$

3.653.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4609 vs. $2(119) = 238$.

Time = 0.79 (sec) , antiderivative size = 4609, normalized size of antiderivative = 30.52

$$\int \frac{\sqrt{\cot(c+dx)}(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = \text{Too large to display}$$

input `integrate(cot(dx+c)^(1/2)*(B*a+b*B*tan(dx+c))/(a+b*tan(dx+c))^(3/2),x,
algorithm="fricas")`

output $1/8*\sqrt{((a^2 + b^2)*\sqrt{-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)}*d^2 + B^2*b)/((a^2 + b^2)*d^2)}*\log(1/2*((2*(B^2*a^3*b^3 + 4*B^2*a*b^5)*d*\tan(dx + c))^2 + 2*(B^2*a^6 + 5*B^2*a^4*b^2 + 8*B^2*a^2*b^4)*d*\tan(dx + c) + 2*(B^2*a^5*b + 2*B^2*a^3*b^3)*d - ((a^7 + 8*a^5*b^2 + 19*a^3*b^4 + 12*a*b^6)*d^3*\tan(dx + c)^2 + 2*(a^6*b + 2*a^4*b^3 - 3*a^2*b^5 - 4*b^7)*d^3*\tan(dx + c) - (a^7 + 4*a^5*b^2 + 7*a^3*b^4 + 4*a*b^6)*d^3)*\sqrt{-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)}))*\sqrt{((a^2 + b^2)*\sqrt{-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)}*d^2 + B^2*b)/((a^2 + b^2)*d^2)} + 2*((B^3*a^5 + 3*B^3*a^3*b^2 + 4*B^3*a*b^4)*\tan(dx + c)^2 + 2*(B^3*a^4*b + 2*B^3*a^2*b^3)*\tan(dx + c) - (2*(B*a^5*b + 3*B*a^3*b^3 + 2*B*a*b^5)*d^2*\tan(dx + c)^2 - (B*a^6 + 4*B*a^4*b^2 + 7*B*a^2*b^4 + 4*B*b^6)*d^2*\tan(dx + c))*\sqrt{-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)}))*\sqrt{b*\tan(dx + c) + a}/\sqrt{\tan(dx + c)})/(tan(dx + c)^2 + 1) + 1/8*\sqrt{((a^2 + b^2)*\sqrt{-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)}*d^2 + B^2*b)/((a^2 + b^2)*d^2)}*\log(-1/2*((2*(B^2*a^3*b^3 + 4*B^2*a*b^5)*d*\tan(dx + c))^2 + 2*(B^2*a^6 + 5*B^2*a^4*b^2 + 8*B^2*a^2*b^4)*d*\tan(dx + c) + 2*(B^2*a^5*b + 2*B^2*a^3*b^3)*d - ((a^7 + 8*a^5*b^2 + 19*a^3*b^4 + 12*a*b^6)*d^3*\tan(dx + c)^2 + 2*(a^6*b + 2*a^4*b^3 - 3*a^2*b^5 - 4*b^7)*d^3*\tan(dx + c) - (a^7 + 4*a^5*b^2 + 7*a^3*b^4 + 4*a*b^6)*d^3)*\sqrt{-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)}))*\sqrt{((a^2 + b^2)*\sqrt{-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4)}*d^2 + B^2*b)/((a^2 + b^2)*d^2)} + ...$

3.653.6 Sympy [F]

$$\int \frac{\sqrt{\cot(c+dx)}(aB+bB\tan(c+dx))}{(a+b\tan(c+dx))^{3/2}} dx = B \int \frac{\sqrt{\cot(c+dx)}}{\sqrt{a+b\tan(c+dx)}} dx$$

input `integrate(cot(d*x+c)**(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))**(3/2),x)`

output `B*Integral(sqrt(cot(c + d*x))/sqrt(a + b*tan(c + d*x)), x)`

3.653.7 Maxima [F]

$$\int \frac{\sqrt{\cot(c+dx)}(aB + bB \tan(c+dx))}{(a + b \tan(c+dx))^{3/2}} dx = \int \frac{(Bb \tan(dx+c) + Ba) \sqrt{\cot(dx+c)}}{(b \tan(dx+c) + a)^{\frac{3}{2}}} dx$$

input `integrate(cot(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,
algorithm="maxima")`

output `integrate((B*b*tan(d*x + c) + B*a)*sqrt(cot(d*x + c))/(b*tan(d*x + c) + a)
^(3/2), x)`

3.653.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\cot(c+dx)}(aB + bB \tan(c+dx))}{(a + b \tan(c+dx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(cot(d*x+c)^(1/2)*(B*a+b*B*tan(d*x+c))/(a+b*tan(d*x+c))^(3/2),x,
algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(con
st gen &`

3.653.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cot(c+dx)}(aB + bB \tan(c+dx))}{(a + b \tan(c+dx))^{3/2}} dx = \int \frac{\sqrt{\cot(c+dx)}(Ba + Bb \tan(c+dx))}{(a + b \tan(c+dx))^{3/2}} dx$$

input `int((cot(c + d*x)^(1/2)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/
2),x)`

output `int((cot(c + d*x)^(1/2)*(B*a + B*b*tan(c + d*x)))/(a + b*tan(c + d*x))^(3/
2), x)`

3.653. $\int \frac{\sqrt{\cot(c+dx)}(aB+bB \tan(c+dx))}{(a+b \tan(c+dx))^{3/2}} dx$

3.654
$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))}^{3/2}} dx$$

3.654.1 Optimal result	6234
3.654.2 Mathematica [A] (verified)	6234
3.654.3 Rubi [A] (verified)	6235
3.654.4 Maple [B] (warning: unable to verify)	6237
3.654.5 Fricas [B] (verification not implemented)	6238
3.654.6 Sympy [F]	6239
3.654.7 Maxima [F]	6240
3.654.8 Giac [F(-1)]	6240
3.654.9 Mupad [F(-1)]	6240

3.654.1 Optimal result

Integrand size = 38, antiderivative size = 157

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))}^{3/2}} dx = \frac{iB \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{\sqrt{ia - bd}} - \frac{iB \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{\sqrt{ia + bd}}$$

output `I*B*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*a-b)^(1/2)-I*B*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*a+b)^(1/2)`

3.654.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.92

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))}^{3/2}} dx = \frac{\sqrt[4]{-1}B \left(-\frac{\arctan\left(\frac{\sqrt[4]{-1}\sqrt{-a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} + \frac{\arctan\left(\frac{\sqrt[4]{-1}\sqrt{a+ib}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{a+ib}} \right)}{d}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)),x]`

output `((-1)^(1/4)*B*(-(ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[-a + I*b]) + ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]/Sqrt[a + I*b])*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]])/d`

3.654.3 Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.85, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.237$, Rules used = {2011, 3042, 4729, 3042, 4058, 613, 104, 218, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx \\
 & \quad \downarrow \text{2011} \\
 & B \int \frac{1}{\sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \int \frac{1}{\sqrt{\cot(c + dx)} \sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4729} \\
 & B \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & B \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)}} dx \\
 & \quad \downarrow \text{4058} \\
 & \frac{B \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \frac{\sqrt{\tan(c + dx)}}{\sqrt{a + b \tan(c + dx)} (\tan^2(c + dx) + 1)} d \tan(c + dx)}{d} \\
 & \quad \downarrow \text{613}
 \end{aligned}$$

3.654. $\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx$

$$\frac{B\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}\int\frac{1}{\sqrt{\tan(c+dx)}(\tan(c+dx)+i)\sqrt{a+b\tan(c+dx)}}d\tan(c+dx)-\frac{1}{2}\int\frac{1}{(i-\tan(c+dx))\sqrt{\tan(c+dx)}}d\tan(c+dx)\right)}{d}$$

↓ 104

$$\frac{B\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\int\frac{1}{\frac{(a-ib)\tan(c+dx)}{a+b\tan(c+dx)}+i}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}-\int\frac{1}{i-\frac{(a+ib)\tan(c+dx)}{a+b\tan(c+dx)}}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{d}$$

↓ 218

$$\frac{B\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\int\frac{1}{\frac{(a-ib)\tan(c+dx)}{a+b\tan(c+dx)}+i}d\frac{\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}+\frac{i\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{-b+ia}}\right)}{d}$$

↓ 221

$$\frac{B\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{i\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{-b+ia}}-\frac{i\operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{b+ia}}\right)}{d}$$

input `Int[(a*B + b*B*Tan[c + d*x])/(Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^(3/2)),x]`

output `(B*((I*ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[I*a - b] - (I*ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[I*a + b])*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]])/d`

3.654.3.1 Defintions of rubi rules used

rule 104 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))/((e_.) + (f_.)*(x_)), x_] := With[{q = Denominator[m]}, Simp[q Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q], x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

3.654. $\int \frac{aB+bB\tan(c+dx)}{\sqrt{\cot(c+dx)}(a+b\tan(c+dx))^{3/2}} dx$

- rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 613 `Int[Sqrt[(e_)*(x_)]/(Sqrt[(c_) + (d_)*(x_)]*((a_) + (b_)*(x_)^2)), x_Symbol] := Simp[e/(2*b) Int[1/(Sqrt[e*x]*Sqrt[c + d*x]*(Rt[-a/b, 2] + x)), x] - Simp[e/(2*b) Int[1/(Sqrt[e*x]*Sqrt[c + d*x]*(Rt[-a/b, 2] - x)), x] /; FreeQ[{a, b, c, d, e}, x]`
- rule 2011 `Int[(u_)*((a_) + (b_)*(v_))^(m_)*((c_) + (d_)*(v_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4058 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Simp[ff/f Subst[Int[(a + b*ff*x)^m*((c + d*ff*x)^n/(1 + ff^2*x^2)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`
- rule 4729 `Int[(cot[(a_) + (b_)*(x_)]*(c_))^(m_)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.654.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 1301 vs. $2(127) = 254$.

Time = 13.34 (sec) , antiderivative size = 1302, normalized size of antiderivative = 8.29

method	result	size
default	Expression too large to display	1302

3.654.
$$\int \frac{aB + bB \tan(c+dx)}{\sqrt{\cot(c+dx)(a+b \tan(c+dx))^{3/2}}} dx$$

```
input int((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x,method=
_RETURNVERBOSE)
```

```
output -1/4*B/d*2^(1/2)/a/(a^2+b^2)^(1/2)/(-b+(a^2+b^2)^(1/2))^(1/2)*((-cos(d*x+c)
)+1)^2*csc(d*x+c)^2-1)*((a*(-cos(d*x+c)+1)^2*csc(d*x+c)^2-2*b*(csc(d*x+c)-
cot(d*x+c))-a)/((-cos(d*x+c)+1)^2*csc(d*x+c)^2-1))^(1/2)*((a^2+b^2)^(1/2)*
(b+(a^2+b^2)^(1/2))^(1/2)*(-b+(a^2+b^2)^(1/2))^(1/2)*ln(-1/(-cos(d*x+c)+1)
*(-a*(-cos(d*x+c)+1)^2*csc(d*x+c)+2*(a^2+b^2)^(1/2)*(-cos(d*x+c)+1)-2*(-(-
cos(d*x+c)+1)*(a*(-cos(d*x+c)+1)^2*csc(d*x+c)^2-2*b*(csc(d*x+c)-cot(d*x+c)
)-a)*csc(d*x+c))^(1/2)*(b+(a^2+b^2)^(1/2))^(1/2)*sin(d*x+c)+2*b*(-cos(d*x+
c)+1)+a*sin(d*x+c))-a^2+b^2)^(1/2)*(b+(a^2+b^2)^(1/2))^(1/2)*(-b+(a^2+b^
2)^(1/2))^(1/2)*ln(1/(-cos(d*x+c)+1)*(-a*(-cos(d*x+c)+1)^2*csc(d*x+c)+2*(a
^2+b^2)^(1/2)*(-cos(d*x+c)+1)+2*(-(-cos(d*x+c)+1)*(a*(-cos(d*x+c)+1)^2*csc
(d*x+c)^2-2*b*(csc(d*x+c)-cot(d*x+c))-a)*csc(d*x+c))^(1/2)*(b+(a^2+b^2)^(1
/2))^(1/2)*sin(d*x+c)+2*b*(-cos(d*x+c)+1)+a*sin(d*x+c))-ln(-1/(-cos(d*x+c)
)+1)*(-a*(-cos(d*x+c)+1)^2*csc(d*x+c)+2*(a^2+b^2)^(1/2)*(-cos(d*x+c)+1)-2*
(-(-cos(d*x+c)+1)*(a*(-cos(d*x+c)+1)^2*csc(d*x+c)^2-2*b*(csc(d*x+c)-cot(d*
x+c))-a)*csc(d*x+c))^(1/2)*(b+(a^2+b^2)^(1/2))^(1/2)*sin(d*x+c)+2*b*(-cos(
d*x+c)+1)+a*sin(d*x+c))*b*(b+(a^2+b^2)^(1/2))^(1/2)*(-b+(a^2+b^2)^(1/2))^(
1/2)+ln(1/(-cos(d*x+c)+1)*(-a*(-cos(d*x+c)+1)^2*csc(d*x+c)+2*(a^2+b^2)^(1
/2)*(-cos(d*x+c)+1)+2*(-(-cos(d*x+c)+1)*(a*(-cos(d*x+c)+1)^2*csc(d*x+c)^2-
2*b*(csc(d*x+c)-cot(d*x+c))-a)*csc(d*x+c))^(1/2)*(b+(a^2+b^2)^(1/2))^(1/2)
*sin(d*x+c)+2*b*(-cos(d*x+c)+1)+a*sin(d*x+c))*b*(b+(a^2+b^2)^(1/2))^(1...
```

3.654.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4649 vs. 2(121) = 242.

Time = 0.76 (sec) , antiderivative size = 4649, normalized size of antiderivative = 29.61

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

```
input integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x,
algorithm="fricas")
```

output

```
-1/8*sqrt(-((a^2 + b^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^2 +
B^2*b)/((a^2 + b^2)*d^2))*log(((B^2*a^6 + 7*B^2*a^4*b^2 + 12*B^2*a^2*b^4
)*d*tan(d*x + c)^2 + 2*(B^2*a^5*b + B^2*a^3*b^3 - 4*B^2*a*b^5)*d*tan(d*x +
c) - (B^2*a^6 + 3*B^2*a^4*b^2 + 4*B^2*a^2*b^4)*d + 2*((a^4*b^3 + 5*a^2*b^
5 + 4*b^7)*d^3*tan(d*x + c)^2 + (a^7 + 6*a^5*b^2 + 13*a^3*b^4 + 8*a*b^6)*d
^3*tan(d*x + c) + (a^6*b + 3*a^4*b^3 + 2*a^2*b^5)*d^3)*sqrt(-B^4*a^2/((a^4
+ 2*a^2*b^2 + b^4)*d^4))*sqrt(-((a^2 + b^2)*sqrt(-B^4*a^2/((a^4 + 2*a^2*
b^2 + b^4)*d^4))*d^2 + B^2*b)/((a^2 + b^2)*d^2)) + 2*((B^3*a^5 + 3*B^3*a^3
*b^2 + 4*B^3*a*b^4)*tan(d*x + c)^2 + 2*(B^3*a^4*b + 2*B^3*a^2*b^3)*tan(d*x
+ c) - (2*(B*a^5*b + 3*B*a^3*b^3 + 2*B*a*b^5)*d^2*tan(d*x + c)^2 - (B*a^6
+ 4*B*a^4*b^2 + 7*B*a^2*b^4 + 4*B*b^6)*d^2*tan(d*x + c))*sqrt(-B^4*a^2/((
a^4 + 2*a^2*b^2 + b^4)*d^4))*sqrt(b*tan(d*x + c) + a)/sqrt(tan(d*x + c)))
/(tan(d*x + c)^2 + 1)) - 1/8*sqrt(-((a^2 + b^2)*sqrt(-B^4*a^2/((a^4 + 2*a^
2*b^2 + b^4)*d^4))*d^2 + B^2*b)/((a^2 + b^2)*d^2))*log(-(((B^2*a^6 + 7*B^2
*a^4*b^2 + 12*B^2*a^2*b^4)*d*tan(d*x + c)^2 + 2*(B^2*a^5*b + B^2*a^3*b^3 -
4*B^2*a*b^5)*d*tan(d*x + c) - (B^2*a^6 + 3*B^2*a^4*b^2 + 4*B^2*a^2*b^4)*d
+ 2*((a^4*b^3 + 5*a^2*b^5 + 4*b^7)*d^3*tan(d*x + c)^2 + (a^7 + 6*a^5*b^2
+ 13*a^3*b^4 + 8*a*b^6)*d^3*tan(d*x + c) + (a^6*b + 3*a^4*b^3 + 2*a^2*b^5)
*d^3)*sqrt(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*sqrt(-((a^2 + b^2)*sqr
t(-B^4*a^2/((a^4 + 2*a^2*b^2 + b^4)*d^4))*d^2 + B^2*b)/((a^2 + b^2)*d^2...
```

3.654.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)(a + b \tan(c + dx))}^{3/2}} dx = B \int \frac{1}{\sqrt{a + b \tan(c + dx)} \sqrt{\cot(c + dx)}} dx$$

input

```
integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)**(1/2)/(a+b*tan(d*x+c))**(3/2),x
)
```

output

```
B*Integral(1/(sqrt(a + b*tan(c + d*x))*sqrt(cot(c + d*x))), x)
```

3.654.7 Maxima [F]

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \int \frac{Bb \tan(dx + c) + Ba}{(b \tan(dx + c) + a)^{\frac{3}{2}} \sqrt{\cot(dx + c)}} dx$$

input `integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x,
algorithm="maxima")`

output `integrate((B*b*tan(d*x + c) + B*a)/((b*tan(d*x + c) + a)^(3/2)*sqrt(cot(d*
x + c))), x)`

3.654.8 Giac [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(3/2),x,
algorithm="giac")`

output `Timed out`

3.654.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx = \int \frac{Ba + Bb \tan(c + dx)}{\sqrt{\cot(c + dx)}(a + b \tan(c + dx))^{3/2}} dx$$

input `int((B*a + B*b*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(3/2
)),x)`

output `int((B*a + B*b*tan(c + d*x))/(cot(c + d*x)^(1/2)*(a + b*tan(c + d*x))^(3/2
)), x)`

3.655
$$\int \frac{aB+bB \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{\frac{3}{2}}} dx$$

3.655.1 Optimal result 6241
 3.655.2 Mathematica [A] (verified) 6242
 3.655.3 Rubi [A] (verified) 6242
 3.655.4 Maple [B] (warning: unable to verify) 6244
 3.655.5 Fricas [B] (verification not implemented) 6245
 3.655.6 Sympy [F] 6246
 3.655.7 Maxima [F(-1)] 6246
 3.655.8 Giac [F(-2)] 6246
 3.655.9 Mupad [F(-1)] 6247

3.655.1 Optimal result

Integrand size = 38, antiderivative size = 215

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{\frac{3}{2}}} dx =$$

$$\frac{B \arctan\left(\frac{\sqrt{ia-b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{\sqrt{ia - bd}}$$

$$+ \frac{2B \operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{\sqrt{bd}}$$

$$- \frac{B \operatorname{arctanh}\left(\frac{\sqrt{ia+b}\sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right) \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)}}{\sqrt{ia + bd}}$$

output

```
-B*arctan((I*a-b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*a-b)^(1/2)+2*B*arctanh(b^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/b^(1/2)-B*arctanh((I*a+b)^(1/2)*tan(d*x+c)^(1/2)/(a+b*tan(d*x+c))^(1/2))*cot(d*x+c)^(1/2)*tan(d*x+c)^(1/2)/d/(I*a+b)^(1/2)
```


3.655.2 Mathematica [A] (verified)

Time = 1.06 (sec) , antiderivative size = 210, normalized size of antiderivative = 0.98

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \frac{B \sqrt{\cot(c + dx)} \sqrt{\tan(c + dx)} \left((-1)^{3/4} \left(\frac{\arctan\left(\frac{\sqrt[4]{-1} \sqrt{-a+ib} \sqrt{\tan(c+dx)}}{\sqrt{a+b \tan(c+dx)}}\right)}{\sqrt{-a+ib}} \right) \right)}{\dots}$$

input `Integrate[(a*B + b*B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2)),x]`

output `(B*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*((-1)^(3/4)*(ArcTan[(-1)^(1/4)*Sqrt[-a + I*b]*Sqrt[Tan[c + d*x]]]/Sqrt[a + b*Tan[c + d*x]]/Sqrt[-a + I*b] + ArcTan[(-1)^(1/4)*Sqrt[a + I*b]*Sqrt[Tan[c + d*x]]/Sqrt[a + b*Tan[c + d*x]]]/Sqrt[a + I*b]) + (2*Sqrt[a]*ArcSinh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a]]*Sqrt[1 + (b*Tan[c + d*x])/a])/(Sqrt[b]*Sqrt[a + b*Tan[c + d*x]]))/d`

3.655.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.78, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.184$, Rules used = {2011, 3042, 4729, 3042, 4058, 614, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx \\ & \quad \downarrow \text{2011} \\ & B \int \frac{1}{\cot^{\frac{3}{2}}(c + dx) \sqrt{a + b \tan(c + dx)}} dx \\ & \quad \downarrow \text{3042} \\ & B \int \frac{1}{\cot(c + dx)^{3/2} \sqrt{a + b \tan(c + dx)}} dx \\ & \quad \downarrow \text{4729} \end{aligned}$$

3.655. $\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx$

$$\begin{aligned}
& B\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\tan^{\frac{3}{2}}(c+dx)}{\sqrt{a+b\tan(c+dx)}} dx \\
& \quad \downarrow \text{3042} \\
& B\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\tan(c+dx)^{3/2}}{\sqrt{a+b\tan(c+dx)}} dx \\
& \quad \downarrow \text{4058} \\
& \frac{B\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{\tan^{\frac{3}{2}}(c+dx)}{\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} d\tan(c+dx)}{d} \\
& \quad \downarrow \text{614} \\
& \frac{B\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \left(\frac{1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}} - \frac{1}{\sqrt{\tan(c+dx)}\sqrt{a+b\tan(c+dx)}(\tan^2(c+dx)+1)} \right) d\tan(c+dx)}{d} \\
& \quad \downarrow \text{2009} \\
& \frac{B\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \left(-\frac{\arctan\left(\frac{\sqrt{-b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{-b+ia}} + \frac{2\operatorname{arctanh}\left(\frac{\sqrt{b}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{b}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b+ia}\sqrt{\tan(c+dx)}}{\sqrt{a+b\tan(c+dx)}}\right)}{\sqrt{b+ia}} \right)}{d}
\end{aligned}$$

input `Int[(a*B + b*B*Tan[c + d*x])/(Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^(3/2)),x]`

output `(B*(-(ArcTan[(Sqrt[I*a - b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[I*a - b]) + (2*ArcTanh[(Sqrt[b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[b] - ArcTanh[(Sqrt[I*a + b]*Sqrt[Tan[c + d*x]])/Sqrt[a + b*Tan[c + d*x]])/Sqrt[I*a + b])*Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]])/d`

3.655.3.1 Defintions of rubi rules used

rule 614 `Int[((e_.)*(x_))^(m_)/(Sqrt[(c_) + (d_.)*(x_)]*((a_) + (b_.)*(x_)^2)), x_Symbol] := Simp[e^(m + 1/2) Int[ExpandIntegrand[1/(Sqrt[e*x]*Sqrt[c + d*x]), x^(m + 1/2)/(a + b*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[m - 1/2, 0]`

$$3.655. \quad \int \frac{aB+bB\tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^{3/2}} dx$$

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2011 `Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :=
Simp[(b/d)^m Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x
&& EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x
, a + b*x])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4058 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((c_) + (d_.)*tan[(e_.) +
(f_.)*(x_)])^(n_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, S
imp[ff/f Subst[Int[(a + b*ff*x)^(m)*((c + d*ff*x)^n/(1 + ff^2*x^2)), x], x,
Tan[e + f*x]/ff], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && NeQ[b*c - a
*d, 0] && NeQ[a^2 + b^2, 0] && NeQ[c^2 + d^2, 0]`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a
+ b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x],
x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u,
x]`

3.655.4 Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 1147 vs. 2(175) = 350.

Time = 13.46 (sec) , antiderivative size = 1148, normalized size of antiderivative = 5.34

method	result	size
default	Expression too large to display	1148

input `int((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x,method=
_RETURNVERBOSE)`

3.655.
$$\int \frac{aB+bB \tan(c+dx)}{\cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^{3/2}} dx$$

output

```
-1/4*B/d*x^2^(1/2)/b^(1/2)/(a^2+b^2)^(1/2)/(-b+(a^2+b^2)^(1/2))^(1/2)*(-ln((
a*cot(d*x+c)*cos(d*x+c)-2*a*cot(d*x+c)+2*((cot(d*x+c)^2*a-2*a*cot(d*x+c)*c
sc(d*x+c)+a*csc(d*x+c)^2-2*b*csc(d*x+c)+2*cot(d*x+c)*b-a)*(cos(d*x+c)-1)*c
sc(d*x+c))^(1/2)*(b+(a^2+b^2)^(1/2))^(1/2)*sin(d*x+c)+a*csc(d*x+c)+2*(a^2+
b^2)^(1/2)*cos(d*x+c)+2*b*cos(d*x+c)-a*sin(d*x+c)-2*(a^2+b^2)^(1/2)-2*b)/(
cos(d*x+c)-1))*2^(1/2)*b^(1/2)*(-b+(a^2+b^2)^(1/2))^(1/2)*(b+(a^2+b^2)^(1/
2))^(1/2)+ln((a*cot(d*x+c)*cos(d*x+c)-2*a*cot(d*x+c)-2*((cot(d*x+c)^2*a-2*
a*cot(d*x+c)*csc(d*x+c)+a*csc(d*x+c)^2-2*b*csc(d*x+c)+2*cot(d*x+c)*b-a)*(c
os(d*x+c)-1)*csc(d*x+c))^(1/2)*(b+(a^2+b^2)^(1/2))^(1/2)*sin(d*x+c)+a*csc(
d*x+c)+2*(a^2+b^2)^(1/2)*cos(d*x+c)+2*b*cos(d*x+c)-a*sin(d*x+c)-2*(a^2+b^2
)^(1/2)-2*b)/(cos(d*x+c)-1))*2^(1/2)*b^(1/2)*(-b+(a^2+b^2)^(1/2))^(1/2)*(b
+(a^2+b^2)^(1/2))^(1/2)-2*arctan(1/(-b+(a^2+b^2)^(1/2))^(1/2))*((b+(a^2+b^2
)^(1/2))^(1/2)*cos(d*x+c)+(-2*(cos(d*x+c)^2*b-cos(d*x+c)*sin(d*x+c)*a-b)/(
cos(d*x+c)+1)^2)^(1/2)*sin(d*x+c)-(b+(a^2+b^2)^(1/2))^(1/2))/(cos(d*x+c)-1
))*2^(1/2)*b^(3/2)+2*arctan(1/(-b+(a^2+b^2)^(1/2))^(1/2))*((b+(a^2+b^2)^(1/
2))^(1/2)*cos(d*x+c)-(-2*(cos(d*x+c)^2*b-cos(d*x+c)*sin(d*x+c)*a-b)/(cos(d
*x+c)+1)^2)^(1/2)*sin(d*x+c)-(b+(a^2+b^2)^(1/2))^(1/2))/(cos(d*x+c)-1))*2^
(1/2)*b^(3/2)+2*arctan(1/(-b+(a^2+b^2)^(1/2))^(1/2))*((b+(a^2+b^2)^(1/2))^(
1/2)*cos(d*x+c)+(-2*(cos(d*x+c)^2*b-cos(d*x+c)*sin(d*x+c)*a-b)/(cos(d*x+c)
+1)^2)^(1/2)*sin(d*x+c)-(b+(a^2+b^2)^(1/2))^(1/2))/(cos(d*x+c)-1))*2^(1...
```

3.655.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 4702 vs. $2(171) = 342$.

Time = 1.12 (sec) , antiderivative size = 9437, normalized size of antiderivative = 43.89

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Too large to display}$$

input

```
integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x,
algorithm="fricas")
```

output

```
Too large to include
```

3.655. $\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx$

3.655.6 Sympy [F]

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = B \int \frac{1}{\sqrt{a + b \tan(c + dx)} \cot^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)**(3/2)/(a+b*tan(d*x+c))**(3/2),x)`

output `B*Integral(1/(sqrt(a + b*tan(c + d*x))*cot(c + d*x)**(3/2)), x)`

3.655.7 Maxima [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Timed out}$$

input `integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x,algorithm="maxima")`

output `Timed out`

3.655.8 Giac [F(-2)]

Exception generated.

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((B*a+b*B*tan(d*x+c))/cot(d*x+c)^(3/2)/(a+b*tan(d*x+c))^(3/2),x,algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen &`

3.655. $\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx$

3.655.9 Mupad [F(-1)]

Timed out.

$$\int \frac{aB + bB \tan(c + dx)}{\cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^{3/2}} dx = \int \frac{B a + B b \tan(c + dx)}{\cot(c + dx)^{3/2} (a + b \tan(c + dx))^{3/2}} dx$$

input `int((B*a + B*b*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(3/2)),x)`

output `int((B*a + B*b*tan(c + d*x))/(cot(c + d*x)^(3/2)*(a + b*tan(c + d*x))^(3/2)), x)`

3.656 $\int \cot^m(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.656.1 Optimal result	6248
3.656.2 Mathematica [F]	6249
3.656.3 Rubi [A] (verified)	6249
3.656.4 Maple [F]	6251
3.656.5 Fricas [F]	6252
3.656.6 Sympy [F(-1)]	6252
3.656.7 Maxima [F]	6252
3.656.8 Giac [F]	6253
3.656.9 Mupad [F(-1)]	6253

3.656.1 Optimal result

Integrand size = 31, antiderivative size = 195

$$\int \cot^m(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \frac{(A+iB) \operatorname{AppellF1}\left(1-m, -n, 1, 2-m, -\frac{b \tan(c+dx)}{a}, -i \tan(c+dx)\right) \cot^{-1+m}(c+dx)(a+b \tan(c+dx))}{2d(1-m)} + \frac{(A-iB) \operatorname{AppellF1}\left(1-m, -n, 1, 2-m, -\frac{b \tan(c+dx)}{a}, i \tan(c+dx)\right) \cot^{-1+m}(c+dx)(a+b \tan(c+dx))}{2d(1-m)}$$

output

```
1/2*(A+I*B)*AppellF1(1-m,1,-n,2-m,-I*tan(d*x+c),-b*tan(d*x+c)/a)*cot(d*x+c)^(-1+m)*(a+b*tan(d*x+c))^n/d/(1-m)/((1+b*tan(d*x+c)/a)^n)+1/2*(A-I*B)*AppellF1(1-m,1,-n,2-m,I*tan(d*x+c),-b*tan(d*x+c)/a)*cot(d*x+c)^(-1+m)*(a+b*tan(d*x+c))^n/d/(1-m)/((1+b*tan(d*x+c)/a)^n)
```

3.656.2 Mathematica [F]

$$\int \cot^m(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \cot^m(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

input `Integrate[Cot[c + d*x]^m*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

output `Integrate[Cot[c + d*x]^m*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]), x]`

3.656.3 Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.11, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.258$, Rules used = {3042, 4729, 3042, 4086, 3042, 4085, 152, 150}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^m(c + dx)(A + B \tan(c + dx))(a + b \tan(c + dx))^n dx$$

$$\downarrow 3042$$

$$\int \cot(c + dx)^m(A + B \tan(c + dx))(a + b \tan(c + dx))^n dx$$

$$\downarrow 4729$$

$$\tan^m(c + dx) \cot^m(c + dx) \int \tan^{-m}(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$\downarrow 3042$$

$$\tan^m(c + dx) \cot^m(c + dx) \int \tan(c + dx)^{-m}(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$\downarrow 4086$$

$$\tan^m(c + dx) \cot^m(c + dx) \left(\frac{1}{2}(A + iB) \int (1 - i \tan(c + dx)) \tan^{-m}(c + dx)(a + b \tan(c + dx))^n dx + \frac{1}{2}(A - iB) \int (i \tan(c + dx) + 1) \tan^{-m}(c + dx)(a + b \tan(c + dx))^n dx \right)$$

$$\downarrow 3042$$

$$\begin{aligned}
& \tan^m(c+dx) \cot^m(c+dx) \\
dx) & \left(\frac{1}{2}(A+iB) \int (1-i \tan(c+dx)) \tan(c+dx)^{-m} (a+b \tan(c+dx))^n dx + \frac{1}{2}(A-iB) \int (i \tan(c+dx)+1) \tan(c+dx)^{-m} (a+b \tan(c+dx))^n dx \right) \\
& \quad \downarrow \text{4085} \\
dx) & \left(\frac{(A-iB) \int \frac{\tan^{-m}(c+dx)(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d \tan(c+dx)}{2d} + \frac{(A+iB) \int \frac{\tan^{-m}(c+dx)(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d \tan(c+dx)}{2d} \right) \\
& \quad \downarrow \text{152} \\
dx) & \left(\frac{(A-iB)(a+b \tan(c+dx))^n \left(\frac{b \tan(c+dx)}{a} + 1 \right)^{-n} \int \frac{\tan^{-m}(c+dx) \left(\frac{b \tan(c+dx)}{a} + 1 \right)^n}{1-i \tan(c+dx)} d \tan(c+dx)}{2d} + \frac{(A+iB)(a+b \tan(c+dx))^n \int \frac{\tan^{-m}(c+dx) \left(\frac{b \tan(c+dx)}{a} + 1 \right)^n}{i \tan(c+dx)+1} d \tan(c+dx)}{2d} \right) \\
& \quad \downarrow \text{150} \\
dx) & \left(\frac{(A+iB) \tan^{1-m}(c+dx) (a+b \tan(c+dx))^n \left(\frac{b \tan(c+dx)}{a} + 1 \right)^{-n} \text{AppellF1} \left(1-m, -n, 1, 2-m, -\frac{b \tan(c+dx)}{a} \right)}{2d(1-m)} \right)
\end{aligned}$$

input `Int[Cot[c + d*x]^m*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `Cot[c + d*x]^m*Tan[c + d*x]^m*((A + I*B)*AppellF1[1 - m, -n, 1, 2 - m, -(b*Tan[c + d*x])/a], (-I)*Tan[c + d*x]]*Tan[c + d*x]^(1 - m)*(a + b*Tan[c + d*x])^n)/(2*d*(1 - m)*(1 + (b*Tan[c + d*x])/a)^n) + ((A - I*B)*AppellF1[1 - m, -n, 1, 2 - m, -(b*Tan[c + d*x])/a], I*Tan[c + d*x]]*Tan[c + d*x]^(1 - m)*(a + b*Tan[c + d*x])^n)/(2*d*(1 - m)*(1 + (b*Tan[c + d*x])/a)^n)`

3.656.3.1 Defintions of rubi rules used

rule 150 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_] :> Simp[c^n*e^p*((b*x)^(m + 1)/(b*(m + 1)))*AppellF1[m + 1, -n, -p, m + 2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])`

- rule 152 `Int[((b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_))*((e_) + (f_.)*(x_)^(p_)), x_] := Simp[c^IntPart[n]*((c + d*x)^FracPart[n]/(1 + d*(x/c))^FracPart[n]) Int[(b*x)^m*(1 + d*(x/c))^n*(e + f*x)^p, x], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && !GtQ[c, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4085 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && EqQ[A^2 + B^2, 0]`
- rule 4086 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`
- rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)])*(c_.)^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.656.4 Maple [F]

$$\int \cot(dx + c)^m (a + b \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(cot(d*x+c)^m*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(cot(d*x+c)^m*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.656. $\int \cot^m(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$

3.656.5 Fricas [F]

$$\int \cot^m(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c)^m dx$$

input `integrate(cot(d*x+c)^m*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*cot(d*x + c)^m, x)`

3.656.6 Sympy [F(-1)]

Timed out.

$$\int \cot^m(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**m*(a+b*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.656.7 Maxima [F]

$$\int \cot^m(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c)^m dx$$

input `integrate(cot(d*x+c)^m*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*cot(d*x + c)^m, x)`

3.656.8 Giac [F]

$$\int \cot^m(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c)^m dx$$

input `integrate(cot(d*x+c)^m*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*cot(d*x + c)^m, x)`

3.656.9 Mupad [F(-1)]

Timed out.

$$\int \cot^m(c + dx)(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^m (A + B \tan(c + dx)) (a + b \tan(c + dx))^n dx$$

input `int(cot(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n,x)`

output `int(cot(c + d*x)^m*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n, x)`

3.657 $\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.657.1 Optimal result	6254
3.657.2 Mathematica [F]	6255
3.657.3 Rubi [A] (warning: unable to verify)	6255
3.657.4 Maple [F]	6258
3.657.5 Fricas [F]	6258
3.657.6 Sympy [F(-1)]	6258
3.657.7 Maxima [F]	6259
3.657.8 Giac [F]	6259
3.657.9 Mupad [F(-1)]	6259

3.657.1 Optimal result

Integrand size = 33, antiderivative size = 169

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx =$$

$$\frac{(A+iB) \operatorname{AppellF1}\left(-\frac{1}{2}, 1, -n, \frac{1}{2}, -i \tan(c+dx), -\frac{b \tan(c+dx)}{a}\right) \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^n \left(1 + \frac{b \tan(c+dx)}{a}\right)^{\frac{1}{2}}}{d} - \frac{(A-iB) \operatorname{AppellF1}\left(-\frac{1}{2}, 1, -n, \frac{1}{2}, i \tan(c+dx), -\frac{b \tan(c+dx)}{a}\right) \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^n \left(1 + \frac{b \tan(c+dx)}{a}\right)^{\frac{1}{2}}}{d}$$

```
output -(A+I*B)*AppellF1(-1/2,1,-n,1/2,-I*tan(d*x+c),-b*tan(d*x+c)/a)*cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^n/d/((1+b*tan(d*x+c)/a)^n)-(A-I*B)*AppellF1(-1/2,1,-n,1/2,I*tan(d*x+c),-b*tan(d*x+c)/a)*cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^n/d/((1+b*tan(d*x+c)/a)^n)
```

3.657.2 Mathematica [F]

$$\int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^n(A+B\tan(c+dx))dx$$

$$= \int \cot^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^n(A+B\tan(c+dx))dx$$

input `Integrate[Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `Integrate[Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

3.657.3 Rubi [A] (warning: unable to verify)

Time = 0.76 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.08, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 4729, 3042, 4086, 3042, 4085, 148, 395, 394}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \cot^{\frac{3}{2}}(c+dx)(A+B\tan(c+dx))(a+b\tan(c+dx))^n dx$$

$$\downarrow \text{3042}$$

$$\int \cot(c+dx)^{3/2}(A+B\tan(c+dx))(a+b\tan(c+dx))^n dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b\tan(c+dx))^n(A+B\tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b\tan(c+dx))^n(A+B\tan(c+dx))}{\tan(c+dx)^{3/2}} dx$$

$$\downarrow \text{4086}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}(A+iB)\int\frac{(1-i\tan(c+dx))(a+b\tan(c+dx))^n}{\tan^{\frac{3}{2}}(c+dx)}dx+\frac{1}{2}(A-iB)\int\frac{(i\tan(c+dx)+1)(a+b\tan(c+dx))^n}{\tan^{\frac{3}{2}}(c+dx)}dx\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}(A+iB)\int\frac{(1-i\tan(c+dx))(a+b\tan(c+dx))^n}{\tan(c+dx)^{3/2}}dx+\frac{1}{2}(A-iB)\int\frac{(i\tan(c+dx)+1)(a+b\tan(c+dx))^n}{\tan(c+dx)^{3/2}}dx\right)$$

↓ 4085

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A-iB)\int\frac{(a+b\tan(c+dx))^n}{(1-i\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)}d\tan(c+dx)}{2d}+\frac{(A+iB)\int\frac{(a+b\tan(c+dx))^n}{(i\tan(c+dx)+1)\tan^{\frac{3}{2}}(c+dx)}d\tan(c+dx)}{2d}\right)$$

↓ 148

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A-iB)\int\frac{\cot^2(c+dx)(a+b\tan(c+dx))^n}{1-i\tan(c+dx)}d\sqrt{\tan(c+dx)}}{d}+\frac{(A+iB)\int\frac{\cot^2(c+dx)(a+b\tan(c+dx))^n}{i\tan(c+dx)+1}d\sqrt{\tan(c+dx)}}{d}\right)$$

↓ 395

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A-iB)(a+b\tan(c+dx))^n\left(\frac{b\tan(c+dx)}{a}+1\right)^{-n}\int\frac{\cot^2(c+dx)\left(\frac{b\tan(c+dx)}{a}+1\right)^n}{1-i\tan(c+dx)}d\sqrt{\tan(c+dx)}}{d}\right)$$

↓ 394

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(-\frac{(A+iB)\cot(c+dx)(a+b\tan(c+dx))^n\left(\frac{b\tan(c+dx)}{a}+1\right)^{-n}\text{AppellF1}\left(-\frac{1}{2},1,-n,-\frac{1}{2},(-I)\tan(c+dx),-\frac{(b\tan(c+dx))}{a}\right)*\cot(c+dx)*(a+b\tan(c+dx))^n}{d}\right)$$

input `Int[Cot[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(-(((A + I*B)*AppellF1[-1/2, 1, -n, 1/2, (-I)*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Cot[c + d*x]*(a + b*Tan[c + d*x])^n)/(d*(1 + (b*Tan[c + d*x])/a)^n)) - ((A - I*B)*AppellF1[-1/2, 1, -n, 1/2, I*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Cot[c + d*x]*(a + b*Tan[c + d*x])^n)/(d*(1 + (b*Tan[c + d*x])/a)^n))`

3.657.3.1 Defintions of rubi rules used

rule 148 `Int[((b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_) + (f_.)*(x_))^(p_.), x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c + d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x]] /; FreeQ[{b, c, d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`

rule 394 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*((e*x)^(m + 1)/(e*(m + 1)))*AppellF1[(m + 1)/2, -p, -q, 1 + (m + 1)/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, e, m, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, 1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 395 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(e*x)^m*(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, e, m, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, 1] && !(IntegerQ[p] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4085 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && EqQ[A^2 + B^2, 0]`

rule 4086 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] :> Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.657.4 Maple [F]

$$\int \cot(dx + c)^{\frac{3}{2}} (a + b \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.657.5 Fricas [F]

$$\begin{aligned} & \int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm m="fricas")`

output `integral((B*cot(d*x + c)*tan(d*x + c) + A*cot(d*x + c))*(b*tan(d*x + c) + a)^n*sqrt(cot(d*x + c)), x)`

3.657.6 Sympy [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(cot(d*x+c)**(3/2)*(a+b*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.657. $\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$

3.657.7 Maxima [F]

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm m="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*cot(d*x + c)^(3/2), x)`

3.657.8 Giac [F]

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \cot(dx + c)^{\frac{3}{2}} dx$$

input `integrate(cot(d*x+c)^(3/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*cot(d*x + c)^(3/2), x)`

3.657.9 Mupad [F(-1)]

Timed out.

$$\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int \cot(c + dx)^{3/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^n dx$$

input `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n,x)`

output `int(cot(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n, x)`

3.657. $\int \cot^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$

3.658 $\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$

3.658.1 Optimal result	6260
3.658.2 Mathematica [F]	6261
3.658.3 Rubi [A] (verified)	6261
3.658.4 Maple [F]	6264
3.658.5 Fricas [F]	6264
3.658.6 Sympy [F]	6264
3.658.7 Maxima [F]	6265
3.658.8 Giac [F]	6265
3.658.9 Mupad [F(-1)]	6266

3.658.1 Optimal result

Integrand size = 33, antiderivative size = 167

$$\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \frac{(A + iB) \operatorname{AppellF1}\left(\frac{1}{2}, 1, -n, \frac{3}{2}, -i \tan(c + dx), -\frac{b \tan(c + dx)}{a}\right) (a + b \tan(c + dx))^n \left(1 + \frac{b \tan(c + dx)}{a}\right)^{-n}}{d \sqrt{\cot(c + dx)}} + \frac{(A - iB) \operatorname{AppellF1}\left(\frac{1}{2}, 1, -n, \frac{3}{2}, i \tan(c + dx), -\frac{b \tan(c + dx)}{a}\right) (a + b \tan(c + dx))^n \left(1 + \frac{b \tan(c + dx)}{a}\right)^{-n}}{d \sqrt{\cot(c + dx)}}$$

output $(A+iB)*\operatorname{AppellF1}(1/2,1,-n,3/2,-I*\tan(d*x+c),-b*\tan(d*x+c)/a)*(a+b*\tan(d*x+c))^n/d/\cot(d*x+c)^{(1/2)/((1+b*\tan(d*x+c)/a)^n)+(A-I*B)*\operatorname{AppellF1}(1/2,1,-n,3/2,I*\tan(d*x+c),-b*\tan(d*x+c)/a)*(a+b*\tan(d*x+c))^n/d/\cot(d*x+c)^{(1/2)/((1+b*\tan(d*x+c)/a)^n)}$

3.658.2 Mathematica [F]

$$\int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \int \sqrt{\cot(c+dx)}(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

input `Integrate[Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `Integrate[Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

3.658.3 Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.13, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 4729, 3042, 4086, 3042, 4085, 148, 334, 333}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\cot(c+dx)}(A+B \tan(c+dx))(a+b \tan(c+dx))^n dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\cot(c+dx)}(A+B \tan(c+dx))(a+b \tan(c+dx))^n dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b \tan(c+dx))^n(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)} \int \frac{(a+b \tan(c+dx))^n(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

$$\downarrow \text{4086}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}(A+iB)\int\frac{(1-i\tan(c+dx))(a+b\tan(c+dx))^n}{\sqrt{\tan(c+dx)}}dx+\frac{1}{2}(A-iB)\int\frac{(i\tan(c+dx)+1)(a+b\tan(c+dx))^n}{\sqrt{\tan(c+dx)}}dx\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}(A+iB)\int\frac{(1-i\tan(c+dx))(a+b\tan(c+dx))^n}{\sqrt{\tan(c+dx)}}dx+\frac{1}{2}(A-iB)\int\frac{(i\tan(c+dx)+1)(a+b\tan(c+dx))^n}{\sqrt{\tan(c+dx)}}dx\right)$$

↓ 4085

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A-iB)\int\frac{(a+b\tan(c+dx))^n}{(1-i\tan(c+dx))\sqrt{\tan(c+dx)}}d\tan(c+dx)}{2d}+\frac{(A+iB)\int\frac{(a+b\tan(c+dx))^n}{(i\tan(c+dx)+1)\sqrt{\tan(c+dx)}}d\tan(c+dx)}{2d}\right)$$

↓ 148

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A-iB)\int\frac{(a+b\tan(c+dx))^n}{1-i\tan(c+dx)}d\sqrt{\tan(c+dx)}}{d}+\frac{(A+iB)\int\frac{(a+b\tan(c+dx))^n}{i\tan(c+dx)+1}d\sqrt{\tan(c+dx)}}{d}\right)$$

↓ 334

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A-iB)(a+b\tan(c+dx))^n\left(\frac{b\tan(c+dx)}{a}+1\right)^{-n}\int\frac{\left(\frac{b\tan(c+dx)}{a}+1\right)^n}{1-i\tan(c+dx)}d\sqrt{\tan(c+dx)}}{d}\right)$$

↓ 333

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A+iB)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^n\left(\frac{b\tan(c+dx)}{a}+1\right)^{-n}\text{AppellF1}\left(\frac{1}{2},1,-n,\frac{3}{2},-I\tan(c+dx),-\frac{(b\tan(c+dx))}{a}\right)}{d}\right)$$

input `Int[Sqrt[Cot[c + d*x]]*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((A + I*B)*AppellF1[1/2, 1, -n, 3/2, (-I)*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^n)/(d*(1 + (b*Tan[c + d*x])/a)^n) + ((A - I*B)*AppellF1[1/2, 1, -n, 3/2, I*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^n)/(d*(1 + (b*Tan[c + d*x])/a)^n))`

3.658.3.1 Defintions of rubi rules used

rule 148 `Int[((b_.)*(x_)^(m_)*((c_) + (d_.)*(x_)^(n_.))*((e_) + (f_.)*(x_)^(p_.),
x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c
+ d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x]] /; FreeQ[{b, c,
d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`

rule 333 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Sim
p[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; F
reeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a,
0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 334 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Sim
p[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[
(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] &&
NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4085 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x
]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n
] && EqQ[A^2 + B^2, 0]`

rule 4086 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] &
& !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.658.4 Maple [F]

$$\int \sqrt{\cot(dx+c)} (a+b\tan(dx+c))^n (A+B\tan(dx+c)) dx$$

input `int(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)), x)`

output `int(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)), x)`

3.658.5 Fricas [F]

$$\begin{aligned} & \int \sqrt{\cot(c+dx)} (a+b\tan(c+dx))^n (A+B\tan(c+dx)) dx \\ &= \int (B\tan(dx+c)+A)(b\tan(dx+c)+a)^n \sqrt{\cot(dx+c)} dx \end{aligned}$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)), x, algorithm="fricas")`

output `integral((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*sqrt(cot(d*x + c)), x)`

3.658.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{\cot(c+dx)} (a+b\tan(c+dx))^n (A+B\tan(c+dx)) dx \\ &= \int (A+B\tan(c+dx)) (a+b\tan(c+dx))^n \sqrt{\cot(c+dx)} dx \end{aligned}$$

input `integrate(cot(d*x+c)**(1/2)*(a+b*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**n*sqrt(cot(c + d*x)),
x)`

3.658.7 Maxima [F]

$$\begin{aligned} & \int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \sqrt{\cot(dx + c)} dx \end{aligned}$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm
m="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*sqrt(cot(d*x + c)),
x)`

3.658.8 Giac [F]

$$\begin{aligned} & \int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^n(A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \sqrt{\cot(dx + c)} dx \end{aligned}$$

input `integrate(cot(d*x+c)^(1/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm
m="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*sqrt(cot(d*x + c)),
x)`

3.658.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cot(c + dx)}(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int \sqrt{\cot(c + dx)} (A + B \tan(c + dx)) (a + b \tan(c + dx))^n dx$$

input `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n,x)`output `int(cot(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n, x)`

$$3.659 \quad \int \frac{(a+b \tan(c+dx))^n(A+B \tan(c+dx))}{\sqrt{\cot(c+dx)}} dx$$

3.659.1 Optimal result	6267
3.659.2 Mathematica [F]	6267
3.659.3 Rubi [A] (verified)	6268
3.659.4 Maple [F]	6271
3.659.5 Fracas [F]	6271
3.659.6 Sympy [F]	6271
3.659.7 Maxima [F]	6272
3.659.8 Giac [F]	6272
3.659.9 Mupad [F(-1)]	6272

3.659.1 Optimal result

Integrand size = 33, antiderivative size = 173

$$\int \frac{(a + b \tan(c + dx))^n(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \frac{(A + iB) \operatorname{AppellF1}\left(\frac{3}{2}, 1, -n, \frac{5}{2}, -i \tan(c + dx), -\frac{b \tan(c + dx)}{a}\right) (a + b \tan(c + dx))^n \left(1 + \frac{b \tan(c + dx)}{a}\right)^{-n}}{3d \cot^{\frac{3}{2}}(c + dx)}$$

$$+ \frac{(A - iB) \operatorname{AppellF1}\left(\frac{3}{2}, 1, -n, \frac{5}{2}, i \tan(c + dx), -\frac{b \tan(c + dx)}{a}\right) (a + b \tan(c + dx))^n \left(1 + \frac{b \tan(c + dx)}{a}\right)^{-n}}{3d \cot^{\frac{3}{2}}(c + dx)}$$

```
output 1/3*(A+I*B)*AppellF1(3/2,1,-n,5/2,-I*tan(d*x+c),-b*tan(d*x+c)/a)*(a+b*tan(d*x+c))^n/d/cot(d*x+c)^(3/2)/((1+b*tan(d*x+c)/a)^n)+1/3*(A-I*B)*AppellF1(3/2,1,-n,5/2,I*tan(d*x+c),-b*tan(d*x+c)/a)*(a+b*tan(d*x+c))^n/d/cot(d*x+c)^(3/2)/((1+b*tan(d*x+c)/a)^n)
```

3.659.2 Mathematica [F]

$$\int \frac{(a + b \tan(c + dx))^n(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(a + b \tan(c + dx))^n(A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

input `Integrate[((a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `Integrate[((a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]], x]`

3.659.3 Rubi [A] (verified)

Time = 0.79 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.12, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 4729, 3042, 4086, 3042, 4085, 148, 395, 394}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))^n}{\sqrt{\cot(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))^n}{\sqrt{\cot(c + dx)}} dx \\
 & \quad \downarrow \text{4729} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \sqrt{\tan(c + dx)} (a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \sqrt{\tan(c + dx)} (a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx \\
 & \quad \downarrow \text{4086} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{2} (A + iB) \int (1 - i \tan(c + dx)) \sqrt{\tan(c + dx)} (a + b \tan(c + dx))^n dx + \frac{1}{2} (A - iB) \int (1 + i \tan(c + dx)) \sqrt{\tan(c + dx)} (a + b \tan(c + dx))^n dx \right) \\
 & \quad \downarrow \text{3042} \\
 & \sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \left(\frac{1}{2} (A + iB) \int (1 - i \tan(c + dx)) \sqrt{\tan(c + dx)} (a + b \tan(c + dx))^n dx + \frac{1}{2} (A - iB) \int (1 + i \tan(c + dx)) \sqrt{\tan(c + dx)} (a + b \tan(c + dx))^n dx \right) \\
 & \quad \downarrow \text{4085}
 \end{aligned}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A-iB)\int\frac{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^n}{1-i\tan(c+dx)}d\tan(c+dx)}{2d}+\frac{(A+iB)\int\frac{\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^n}{i\tan(c+dx)+1}}{2d}\right)$$

↓ 148

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A-iB)\int\frac{\tan(c+dx)(a+b\tan(c+dx))^n}{1-i\tan(c+dx)}d\sqrt{\tan(c+dx)}}{d}+\frac{(A+iB)\int\frac{\tan(c+dx)(a+b\tan(c+dx))^n}{i\tan(c+dx)+1}}{d}\right)$$

↓ 395

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A-iB)(a+b\tan(c+dx))^n\left(\frac{b\tan(c+dx)}{a}+1\right)^{-n}\int\frac{\tan(c+dx)\left(\frac{b\tan(c+dx)}{a}+1\right)^n}{1-i\tan(c+dx)}d\sqrt{\tan(c+dx)}}{d}\right)$$

↓ 394

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A+iB)\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^n\left(\frac{b\tan(c+dx)}{a}+1\right)^{-n}\text{AppellF1}\left(\frac{3}{2},1,-n,\right)}{3d}\right)$$

input `Int[((a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Sqrt[Cot[c + d*x]],x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((A + I*B)*AppellF1[3/2, 1, -n, 5/2, (-I)*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^n)/(3*d*(1 + (b*Tan[c + d*x])/a)^n) + ((A - I*B)*AppellF1[3/2, 1, -n, 5/2, I*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^n)/(3*d*(1 + (b*Tan[c + d*x])/a)^n))`

3.659.3.1 Defintions of rubi rules used

rule 148 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_.)*((e_) + (f_.)*(x_))^(p_.), x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c + d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x]] /; FreeQ[{b, c, d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`

- rule 394 `Int[((e._)*(x._))^(m._)*((a._) + (b._)*(x._)^2)^(p._)*((c._) + (d._)*(x._)^2)^(q._), x_Symbol] := Simp[a^p*c^q*((e*x)^(m + 1)/(e*(m + 1)))*AppellF1[(m + 1)/2, -p, -q, 1 + (m + 1)/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, e, m, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, 1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`
- rule 395 `Int[((e._)*(x._))^(m._)*((a._) + (b._)*(x._)^2)^(p._)*((c._) + (d._)*(x._)^2)^(q._), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(e*x)^m*(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, e, m, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, 1] && !(IntegerQ[p] || GtQ[a, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4085 `Int[((a._) + (b._)*tan[(e._) + (f._)*(x._)])^(m._)*((A._) + (B._)*tan[(e._) + (f._)*(x._)])*((c._) + (d._)*tan[(e._) + (f._)*(x._)])^(n._), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && EqQ[A^2 + B^2, 0]`
- rule 4086 `Int[((a._) + (b._)*tan[(e._) + (f._)*(x._)])^(m._)*((A._) + (B._)*tan[(e._) + (f._)*(x._)])*((c._) + (d._)*tan[(e._) + (f._)*(x._)])^(n._), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`
- rule 4729 `Int[(cot[(a._) + (b._)*(x._)]*(c._))^(m._)*(u._), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.659.4 Maple [F]

$$\int \frac{(a + b \tan(dx + c))^n (A + B \tan(dx + c))}{\sqrt{\cot(dx + c)}} dx$$

input `int((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x)`

output `int((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x)`

3.659.5 Fricas [F]

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\ &= \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^n}{\sqrt{\cot(dx + c)}} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="fricas")`

output `integral((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n/sqrt(cot(d*x + c)), x)`

3.659.6 Sympy [F]

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx \\ &= \int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))^n}{\sqrt{\cot(c + dx)}} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))**n*(A+B*tan(d*x+c))/cot(d*x+c)**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**n/sqrt(cot(c + d*x)), x)`

3.659.7 Maxima [F]

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^n}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n/sqrt(cot(d*x + c)), x)`

3.659.8 Giac [F]

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^n}{\sqrt{\cot(dx + c)}} dx$$

input `integrate((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(1/2),x, algorithm="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n/sqrt(cot(d*x + c)), x)`

3.659.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\cot(c + dx)}} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^n}{\sqrt{\cot(c + dx)}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n)/cot(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n)/cot(c + d*x)^(1/2), x)`

3.660
$$\int \frac{(a+b \tan(c+dx))^n(A+B \tan(c+dx))}{\cot^{\frac{3}{2}}(c+dx)} dx$$

3.660.1 Optimal result 6274
 3.660.2 Mathematica [F] 6275
 3.660.3 Rubi [A] (verified) 6275
 3.660.4 Maple [F] 6278
 3.660.5 Fricas [F] 6278
 3.660.6 Sympy [F] 6279
 3.660.7 Maxima [F] 6279
 3.660.8 Giac [F] 6279
 3.660.9 Mupad [F(-1)] 6280

3.660.1 Optimal result

Integrand size = 33, antiderivative size = 173

$$\int \frac{(a + b \tan(c + dx))^n(A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{(A + iB) \operatorname{AppellF1}\left(\frac{5}{2}, 1, -n, \frac{7}{2}, -i \tan(c + dx), -\frac{b \tan(c + dx)}{a}\right) (a + b \tan(c + dx))^n \left(1 + \frac{b \tan(c + dx)}{a}\right)^{-n}}{5d \cot^{\frac{5}{2}}(c + dx)}$$

$$+ \frac{(A - iB) \operatorname{AppellF1}\left(\frac{5}{2}, 1, -n, \frac{7}{2}, i \tan(c + dx), -\frac{b \tan(c + dx)}{a}\right) (a + b \tan(c + dx))^n \left(1 + \frac{b \tan(c + dx)}{a}\right)^{-n}}{5d \cot^{\frac{5}{2}}(c + dx)}$$

output

```
1/5*(A+I*B)*AppellF1(5/2,1,-n,7/2,-I*tan(d*x+c),-b*tan(d*x+c)/a)*(a+b*tan(d*x+c))^n/d/cot(d*x+c)^(5/2)/((1+b*tan(d*x+c)/a)^n)+1/5*(A-I*B)*AppellF1(5/2,1,-n,7/2,I*tan(d*x+c),-b*tan(d*x+c)/a)*(a+b*tan(d*x+c))^n/d/cot(d*x+c)^(5/2)/((1+b*tan(d*x+c)/a)^n)
```

3.660.2 Mathematica [F]

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

input `Integrate[((a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(3/2), x]`

output `Integrate[((a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(3/2), x]`

3.660.3 Rubi [A] (verified)

Time = 0.79 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.12, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3042, 4729, 3042, 4086, 3042, 4085, 148, 395, 394}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))^n}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))^n}{\cot(c + dx)^{3/2}} dx$$

$$\downarrow \text{4729}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \tan^{\frac{3}{2}}(c + dx) (a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$\downarrow \text{3042}$$

$$\sqrt{\tan(c + dx)} \sqrt{\cot(c + dx)} \int \tan(c + dx)^{3/2} (a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$\downarrow \text{4086}$$

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}(A+iB)\int(1-i\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^n dx + \frac{1}{2}(A-iB)\int(1+i\tan(c+dx))\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^n dx\right)$$

↓ 3042

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{1}{2}(A+iB)\int(1-i\tan(c+dx))\tan(c+dx)^{3/2}(a+b\tan(c+dx))^n dx + \frac{1}{2}(A-iB)\int(1+i\tan(c+dx))\tan(c+dx)^{3/2}(a+b\tan(c+dx))^n dx\right)$$

↓ 4085

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A-iB)\int\frac{\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^n}{1-i\tan(c+dx)}d\tan(c+dx)}{2d} + \frac{(A+iB)\int\frac{\tan^{\frac{3}{2}}(c+dx)(a+b\tan(c+dx))^n}{i\tan(c+dx)+1}}{2d}\right)$$

↓ 148

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A-iB)\int\frac{\tan^2(c+dx)(a+b\tan(c+dx))^n}{1-i\tan(c+dx)}d\sqrt{\tan(c+dx)}}{d} + \frac{(A+iB)\int\frac{\tan^2(c+dx)(a+b\tan(c+dx))^n}{i\tan(c+dx)+1}}{d}\right)$$

↓ 395

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A-iB)(a+b\tan(c+dx))^n\left(\frac{b\tan(c+dx)}{a}+1\right)^{-n}\int\frac{\tan^2(c+dx)\left(\frac{b\tan(c+dx)}{a}+1\right)^n}{1-i\tan(c+dx)}d\sqrt{\tan(c+dx)}}{d} + \frac{(A+iB)(a+b\tan(c+dx))^n\left(\frac{b\tan(c+dx)}{a}+1\right)^{-n}\int\frac{\tan^2(c+dx)\left(\frac{b\tan(c+dx)}{a}+1\right)^n}{i\tan(c+dx)+1}}{d}\right)$$

↓ 394

$$\sqrt{\tan(c+dx)}\sqrt{\cot(c+dx)}\left(\frac{(A+iB)\tan^{\frac{5}{2}}(c+dx)(a+b\tan(c+dx))^n\left(\frac{b\tan(c+dx)}{a}+1\right)^{-n}\text{AppellF1}\left(\frac{5}{2}, 1, -n, \frac{b\tan(c+dx)}{a}+1\right)}{5d}\right)$$

input `Int[((a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Cot[c + d*x]^(3/2),x]`

output `Sqrt[Cot[c + d*x]]*Sqrt[Tan[c + d*x]]*(((A + I*B)*AppellF1[5/2, 1, -n, 7/2, (-I)*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^n)/(5*d*(1 + (b*Tan[c + d*x])/a)^n) + ((A - I*B)*AppellF1[5/2, 1, -n, 7/2, I*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^n)/(5*d*(1 + (b*Tan[c + d*x])/a)^n))`

3.660.3.1 Defintions of rubi rules used

rule 148 `Int[((b_.)*(x_)^(m_)*((c_) + (d_.)*(x_)^(n_.))*((e_) + (f_.)*(x_)^(p_.),
x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c
+ d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x]] /; FreeQ[{b, c,
d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`

rule 394 `Int[((e_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_
, x_Symbol] := Simp[a^p*c^q*((e*x)^(m + 1)/(e*(m + 1)))*AppellF1[(m + 1)/2
, -p, -q, 1 + (m + 1)/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c,
d, e, m, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, 1] && (Int
egerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 395 `Int[((e_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_
, x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^
FracPart[p]) Int[(e*x)^m*(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ
[{a, b, c, d, e, m, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m,
1] && !(IntegerQ[p] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4085 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x
]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n
] && EqQ[A^2 + B^2, 0]`

rule 4086 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] &
& !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`

rule 4729 `Int[(cot[(a_.) + (b_.)*(x_.)]*(c_.))^(m_.)*(u_), x_Symbol] := Simp[(c*Cot[a + b*x])^m*(c*Tan[a + b*x])^m Int[ActivateTrig[u]/(c*Tan[a + b*x])^m, x], x] /; FreeQ[{a, b, c, m}, x] && !IntegerQ[m] && KnownTangentIntegrandQ[u, x]`

3.660.4 Maple [F]

$$\int \frac{(a + b \tan(dx + c))^n (A + B \tan(dx + c))}{\cot(dx + c)^{\frac{3}{2}}} dx$$

input `int((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x)`

output `int((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x)`

3.660.5 Fracas [F]

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^n}{\cot(dx + c)^{\frac{3}{2}}} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algorithm m="fracas")`

output `integral((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n/cot(d*x + c)^(3/2), x)`

3.660.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^n}{\cot^{\frac{3}{2}}(c + dx)} dx$$

input `integrate((a+b*tan(d*x+c))**n*(A+B*tan(d*x+c))/cot(d*x+c)**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**n/cot(c + d*x)**(3/2), x)`

3.660.7 Maxima [F]

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^n}{\cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+b*tan(d*x+c))~n*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algorithm m="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n/cot(d*x + c)^(3/2), x)`

3.660.8 Giac [F]

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^n}{\cot(dx + c)^{\frac{3}{2}}} dx$$

input `integrate((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/cot(d*x+c)^(3/2),x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n/cot(d*x + c)^(3/2), x)`

3.660.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\cot^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^n}{\cot(c + dx)^{3/2}} dx$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n)/cot(c + d*x)^(3/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n)/cot(c + d*x)^(3/2), x)`

3.661 $\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

3.661.1 Optimal result	6281
3.661.2 Mathematica [F]	6282
3.661.3 Rubi [A] (verified)	6282
3.661.4 Maple [F]	6285
3.661.5 Fricas [F]	6285
3.661.6 Sympy [F(-1)]	6285
3.661.7 Maxima [F]	6286
3.661.8 Giac [F]	6286
3.661.9 Mupad [F(-1)]	6286

3.661.1 Optimal result

Integrand size = 33, antiderivative size = 173

$$\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \frac{(A+iB) \operatorname{AppellF1}\left(\frac{5}{2}, 1, -n, \frac{7}{2}, -i \tan(c+dx), -\frac{b \tan(c+dx)}{a}\right) \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^n \left(1 + \frac{b \tan(c+dx)}{a}\right)}{5d} + \frac{(A-iB) \operatorname{AppellF1}\left(\frac{5}{2}, 1, -n, \frac{7}{2}, i \tan(c+dx), -\frac{b \tan(c+dx)}{a}\right) \tan^{\frac{5}{2}}(c+dx)(a+b \tan(c+dx))^n \left(1 + \frac{b \tan(c+dx)}{a}\right)}{5d}$$

```
output 1/5*(A+I*B)*AppellF1(5/2,1,-n,7/2,-I*tan(d*x+c),-b*tan(d*x+c)/a)*tan(d*x+c)
^(5/2)*(a+b*tan(d*x+c))^n/d/((1+b*tan(d*x+c)/a)^n)+1/5*(A-I*B)*AppellF1(5
/2,1,-n,7/2,I*tan(d*x+c),-b*tan(d*x+c)/a)*tan(d*x+c)^(5/2)*(a+b*tan(d*x+c)
)^n/d/((1+b*tan(d*x+c)/a)^n)
```


3.661.2 Mathematica [F]

$$\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

$$= \int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$$

input `Integrate[Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `Integrate[Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

3.661.3 Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$, Rules used = {3042, 4086, 3042, 4085, 148, 395, 394}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \tan^{\frac{3}{2}}(c+dx)(A+B \tan(c+dx))(a+b \tan(c+dx))^n dx$$

$$\downarrow \text{3042}$$

$$\int \tan(c+dx)^{3/2}(A+B \tan(c+dx))(a+b \tan(c+dx))^n dx$$

$$\downarrow \text{4086}$$

$$\frac{1}{2}(A+iB) \int (1-i \tan(c+dx)) \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^n dx + \frac{1}{2}(A-iB) \int (i \tan(c+dx) + 1) \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^n dx$$

$$\downarrow \text{3042}$$

$$\frac{1}{2}(A+iB) \int (1-i \tan(c+dx)) \tan(c+dx)^{3/2}(a+b \tan(c+dx))^n dx + \frac{1}{2}(A-iB) \int (i \tan(c+dx) + 1) \tan(c+dx)^{3/2}(a+b \tan(c+dx))^n dx$$

$$\downarrow \text{4085}$$

3.661. $\int \tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^n(A+B \tan(c+dx)) dx$

$$\begin{aligned}
& \frac{(A - iB) \int \frac{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d \tan(c+dx)}{2d} + \\
& \frac{(A + iB) \int \frac{\tan^{\frac{3}{2}}(c+dx)(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d \tan(c+dx)}{2d} \\
& \quad \downarrow 148 \\
& \frac{(A - iB) \int \frac{\tan^2(c+dx)(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d \sqrt{\tan(c+dx)}}{d} + \\
& \frac{(A + iB) \int \frac{\tan^2(c+dx)(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d \sqrt{\tan(c+dx)}}{d} \\
& \quad \downarrow 395 \\
& \frac{(A - iB)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1\right)^{-n} \int \frac{\tan^2(c+dx) \left(\frac{b \tan(c+dx)}{a} + 1\right)^n}{1-i \tan(c+dx)} d \sqrt{\tan(c+dx)}}{d} + \\
& \frac{(A + iB)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1\right)^{-n} \int \frac{\tan^2(c+dx) \left(\frac{b \tan(c+dx)}{a} + 1\right)^n}{i \tan(c+dx)+1} d \sqrt{\tan(c+dx)}}{d} \\
& \quad \downarrow 394 \\
& \frac{(A + iB) \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1\right)^{-n} \text{AppellF1}\left(\frac{5}{2}, 1, -n, \frac{7}{2}, -i \tan(c + dx), -\frac{b \tan(c+dx)}{a}\right)}{5d} \\
& \frac{(A - iB) \tan^{\frac{5}{2}}(c + dx)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1\right)^{-n} \text{AppellF1}\left(\frac{5}{2}, 1, -n, \frac{7}{2}, i \tan(c + dx), -\frac{b \tan(c+dx)}{a}\right)}{5d}
\end{aligned}$$

input `Int[Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `((A + I*B)*AppellF1[5/2, 1, -n, 7/2, (-I)*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^n)/(5*d*(1 + (b*Tan[c + d*x])/a)^n) + ((A - I*B)*AppellF1[5/2, 1, -n, 7/2, I*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Tan[c + d*x]^(5/2)*(a + b*Tan[c + d*x])^n)/(5*d*(1 + (b*Tan[c + d*x])/a)^n)`

3.661.3.1 Defintions of rubi rules used

rule 148 `Int[((b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_) + (f_.)*(x_))^(p_.), x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c + d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x]] /; FreeQ[{b, c, d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`

rule 394 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*((e*x)^(m + 1)/(e*(m + 1)))*AppellF1[(m + 1)/2, -p, -q, 1 + (m + 1)/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, e, m, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, 1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 395 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(e*x)^m*(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, e, m, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, 1] && !(IntegerQ[p] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4085 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && EqQ[A^2 + B^2, 0]`

rule 4086 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`

3.661.4 Maple [F]

$$\int \tan(dx + c)^{\frac{3}{2}} (a + b \tan(dx + c))^n (A + B \tan(dx + c)) dx$$

input `int(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.661.5 Fricas [F]

$$\begin{aligned} & \int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx \\ &= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^{\frac{3}{2}} dx \end{aligned}$$

input `integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*tan(d*x + c)^2 + A*tan(d*x + c))*(b*tan(d*x + c) + a)^n*sqrt(tan(d*x + c)), x)`

3.661.6 Sympy [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx = \text{Timed out}$$

input `integrate(tan(d*x+c)**(3/2)*(a+b*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Timed out`

3.661.7 Maxima [F]

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^{\frac{3}{2}} dx$$

input `integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm m="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*tan(d*x + c)^(3/2), x)`

3.661.8 Giac [F]

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int (B \tan(dx + c) + A)(b \tan(dx + c) + a)^n \tan(dx + c)^{\frac{3}{2}} dx$$

input `integrate(tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*tan(d*x + c)^(3/2), x)`

3.661.9 Mupad [F(-1)]

Timed out.

$$\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \int \tan(c + dx)^{3/2} (A + B \tan(c + dx)) (a + b \tan(c + dx))^n dx$$

input `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n,x)`

output `int(tan(c + d*x)^(3/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n, x)`

3.661. $\int \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$

3.662 $\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$

3.662.1 Optimal result	6287
3.662.2 Mathematica [F]	6288
3.662.3 Rubi [A] (verified)	6288
3.662.4 Maple [F]	6291
3.662.5 Fricas [F]	6291
3.662.6 Sympy [F]	6291
3.662.7 Maxima [F]	6292
3.662.8 Giac [F]	6292
3.662.9 Mupad [F(-1)]	6292

3.662.1 Optimal result

Integrand size = 33, antiderivative size = 173

$$\int \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^n (A + B \tan(c + dx)) dx$$

$$= \frac{(A + iB) \operatorname{AppellF1}\left(\frac{3}{2}, 1, -n, \frac{5}{2}, -i \tan(c + dx), -\frac{b \tan(c + dx)}{a}\right) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n \left(1 + \frac{b \tan(c + dx)}{a}\right)}{3d} + \frac{(A - iB) \operatorname{AppellF1}\left(\frac{3}{2}, 1, -n, \frac{5}{2}, i \tan(c + dx), -\frac{b \tan(c + dx)}{a}\right) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n \left(1 + \frac{b \tan(c + dx)}{a}\right)}{3d}$$

```
output 1/3*(A+I*B)*AppellF1(3/2,1,-n,5/2,-I*tan(d*x+c),-b*tan(d*x+c)/a)*tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^n/d/((1+b*tan(d*x+c)/a)^n)+1/3*(A-I*B)*AppellF1(3/2,1,-n,5/2,I*tan(d*x+c),-b*tan(d*x+c)/a)*tan(d*x+c)^(3/2)*(a+b*tan(d*x+c))^n/d/((1+b*tan(d*x+c)/a)^n)
```

3.662.2 Mathematica [F]

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^n(A+B\tan(c+dx))dx$$

$$= \int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^n(A+B\tan(c+dx))dx$$

input `Integrate[Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `Integrate[Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

3.662.3 Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$, Rules used = {3042, 4086, 3042, 4085, 148, 395, 394}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\tan(c+dx)}(A+B\tan(c+dx))(a+b\tan(c+dx))^n dx$$

$$\downarrow \text{3042}$$

$$\int \sqrt{\tan(c+dx)}(A+B\tan(c+dx))(a+b\tan(c+dx))^n dx$$

$$\downarrow \text{4086}$$

$$\frac{1}{2}(A+iB) \int (1-i\tan(c+dx))\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^n dx + \frac{1}{2}(A-iB) \int (i\tan(c+dx)+1)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^n dx$$

$$\downarrow \text{3042}$$

$$\frac{1}{2}(A+iB) \int (1-i\tan(c+dx))\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^n dx + \frac{1}{2}(A-iB) \int (i\tan(c+dx)+1)\sqrt{\tan(c+dx)}(a+b\tan(c+dx))^n dx$$

$$\downarrow \text{4085}$$

$$\begin{aligned}
 & \frac{(A - iB) \int \frac{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d \tan(c + dx)}{2d} + \\
 & \frac{(A + iB) \int \frac{\sqrt{\tan(c+dx)}(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d \tan(c + dx)}{2d} \\
 & \quad \downarrow \text{148} \\
 & \frac{(A - iB) \int \frac{\tan(c+dx)(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d \sqrt{\tan(c + dx)}}{d} + \\
 & \frac{(A + iB) \int \frac{\tan(c+dx)(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d \sqrt{\tan(c + dx)}}{d} \\
 & \quad \downarrow \text{395} \\
 & \frac{(A - iB)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1\right)^{-n} \int \frac{\tan(c+dx) \left(\frac{b \tan(c+dx)}{a} + 1\right)^n}{1-i \tan(c+dx)} d \sqrt{\tan(c + dx)}}{d} + \\
 & \frac{(A + iB)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1\right)^{-n} \int \frac{\tan(c+dx) \left(\frac{b \tan(c+dx)}{a} + 1\right)^n}{i \tan(c+dx)+1} d \sqrt{\tan(c + dx)}}{d} \\
 & \quad \downarrow \text{394} \\
 & \frac{(A + iB) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1\right)^{-n} \text{AppellF1}\left(\frac{3}{2}, 1, -n, \frac{5}{2}, -i \tan(c + dx), -\frac{b \tan(c+dx)}{a}\right)}{3d} \\
 & \frac{(A - iB) \tan^{\frac{3}{2}}(c + dx)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1\right)^{-n} \text{AppellF1}\left(\frac{3}{2}, 1, -n, \frac{5}{2}, i \tan(c + dx), -\frac{b \tan(c+dx)}{a}\right)}{3d}
 \end{aligned}$$

input `Int[Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]),x]`

output `((A + I*B)*AppellF1[3/2, 1, -n, 5/2, (-I)*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^n)/(3*d*(1 + (b*Tan[c + d*x])/a)^n) + ((A - I*B)*AppellF1[3/2, 1, -n, 5/2, I*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Tan[c + d*x]^(3/2)*(a + b*Tan[c + d*x])^n)/(3*d*(1 + (b*Tan[c + d*x])/a)^n)`

3.662.3.1 Defintions of rubi rules used

rule 148 `Int[((b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_) + (f_.)*(x_))^(p_.), x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c + d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x]] /; FreeQ[{b, c, d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`

rule 394 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*((e*x)^(m + 1)/(e*(m + 1)))*AppellF1[(m + 1)/2, -p, -q, 1 + (m + 1)/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, e, m, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, 1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 395 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(e*x)^m*(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, e, m, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, 1] && !(IntegerQ[p] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4085 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && EqQ[A^2 + B^2, 0]`

rule 4086 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*Tan[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`

3.662.4 Maple [F]

$$\int \sqrt{\tan(dx+c)} (a+b\tan(dx+c))^n (A+B\tan(dx+c)) dx$$

input `int(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

output `int(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x)`

3.662.5 Fricas [F]

$$\begin{aligned} & \int \sqrt{\tan(c+dx)} (a+b\tan(c+dx))^n (A+B\tan(c+dx)) dx \\ &= \int (B\tan(dx+c)+A)(b\tan(dx+c)+a)^n \sqrt{\tan(dx+c)} dx \end{aligned}$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm="fricas")`

output `integral((B*tan(d*x+c)+A)*(b*tan(d*x+c)+a)^n*sqrt(tan(d*x+c)),x)`

3.662.6 Sympy [F]

$$\begin{aligned} & \int \sqrt{\tan(c+dx)} (a+b\tan(c+dx))^n (A+B\tan(c+dx)) dx \\ &= \int (A+B\tan(c+dx)) (a+b\tan(c+dx))^n \sqrt{\tan(c+dx)} dx \end{aligned}$$

input `integrate(tan(d*x+c)**(1/2)*(a+b*tan(d*x+c))**n*(A+B*tan(d*x+c)),x)`

output `Integral((A+B*tan(c+d*x))*(a+b*tan(c+d*x))**n*sqrt(tan(c+d*x)),x)`

3.662.7 Maxima [F]

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^n(A+B\tan(c+dx))dx$$

$$= \int (B\tan(dx+c)+A)(b\tan(dx+c)+a)^n\sqrt{\tan(dx+c)}dx$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm m="maxima")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*sqrt(tan(d*x + c)), x)`

3.662.8 Giac [F]

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^n(A+B\tan(c+dx))dx$$

$$= \int (B\tan(dx+c)+A)(b\tan(dx+c)+a)^n\sqrt{\tan(dx+c)}dx$$

input `integrate(tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^n*(A+B*tan(d*x+c)),x, algorithm m="giac")`

output `integrate((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n*sqrt(tan(d*x + c)), x)`

3.662.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^n(A+B\tan(c+dx))dx$$

$$= \int \sqrt{\tan(c+dx)}(A+B\tan(c+dx))(a+b\tan(c+dx))^n dx$$

input `int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n,x)`

output `int(tan(c + d*x)^(1/2)*(A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n, x)`

3.662. $\int \sqrt{\tan(c+dx)}(a+b\tan(c+dx))^n(A+B\tan(c+dx))dx$

3.663
$$\int \frac{(a+b \tan(c+dx))^n(A+B \tan(c+dx))}{\sqrt{\tan(c+dx)}} dx$$

3.663.1 Optimal result 6293
 3.663.2 Mathematica [F] 6293
 3.663.3 Rubi [A] (verified) 6294
 3.663.4 Maple [F] 6296
 3.663.5 Fricas [F] 6297
 3.663.6 Sympy [F] 6297
 3.663.7 Maxima [F(-1)] 6297
 3.663.8 Giac [F(-1)] 6298
 3.663.9 Mupad [F(-1)] 6298

3.663.1 Optimal result

Integrand size = 33, antiderivative size = 167

$$\int \frac{(a + b \tan(c + dx))^n(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \frac{(A + iB) \operatorname{AppellF1}\left(\frac{1}{2}, 1, -n, \frac{3}{2}, -i \tan(c + dx), -\frac{b \tan(c + dx)}{a}\right) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^n \left(1 + \frac{b \tan(c + dx)}{a}\right)^n}{d} + \frac{(A - iB) \operatorname{AppellF1}\left(\frac{1}{2}, 1, -n, \frac{3}{2}, i \tan(c + dx), -\frac{b \tan(c + dx)}{a}\right) \sqrt{\tan(c + dx)}(a + b \tan(c + dx))^n \left(1 + \frac{b \tan(c + dx)}{a}\right)^n}{d}$$

output `(A+I*B)*AppellF1(1/2,1,-n,3/2,-I*tan(d*x+c),-b*tan(d*x+c)/a)*tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^n/d/((1+b*tan(d*x+c)/a)^n)+(A-I*B)*AppellF1(1/2,1,-n,3/2,I*tan(d*x+c),-b*tan(d*x+c)/a)*tan(d*x+c)^(1/2)*(a+b*tan(d*x+c))^n/d/((1+b*tan(d*x+c)/a)^n)`

3.663.2 Mathematica [F]

$$\int \frac{(a + b \tan(c + dx))^n(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \int \frac{(a + b \tan(c + dx))^n(A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

input `Integrate[((a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `Integrate[((a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]], x]`

3.663.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$, Rules used = {3042, 4086, 3042, 4085, 148, 334, 333}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))^n}{\sqrt{\tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))^n}{\sqrt{\tan(c + dx)}} dx \\
 & \quad \downarrow \text{4086} \\
 & \frac{1}{2}(A + iB) \int \frac{(1 - i \tan(c + dx))(a + b \tan(c + dx))^n}{\sqrt{\tan(c + dx)}} dx + \frac{1}{2}(A - \\
 & \quad iB) \int \frac{(i \tan(c + dx) + 1)(a + b \tan(c + dx))^n}{\sqrt{\tan(c + dx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \frac{1}{2}(A + iB) \int \frac{(1 - i \tan(c + dx))(a + b \tan(c + dx))^n}{\sqrt{\tan(c + dx)}} dx + \frac{1}{2}(A - \\
 & \quad iB) \int \frac{(i \tan(c + dx) + 1)(a + b \tan(c + dx))^n}{\sqrt{\tan(c + dx)}} dx \\
 & \quad \downarrow \text{4085} \\
 & \frac{(A - iB) \int \frac{(a + b \tan(c + dx))^n}{(1 - i \tan(c + dx)) \sqrt{\tan(c + dx)}} d \tan(c + dx)}{2d} + \\
 & \quad \frac{(A + iB) \int \frac{(a + b \tan(c + dx))^n}{(i \tan(c + dx) + 1) \sqrt{\tan(c + dx)}} d \tan(c + dx)}{2d} \\
 & \quad \downarrow \text{148}
 \end{aligned}$$

3.663. $\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$

$$\begin{aligned}
& \frac{(A - iB) \int \frac{(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d\sqrt{\tan(c+dx)}}{d} + \frac{(A + iB) \int \frac{(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} \\
& \quad \downarrow \text{334} \\
& \frac{(A - iB)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1 \right)^{-n} \int \frac{\left(\frac{b \tan(c+dx)}{a} + 1 \right)^n}{1-i \tan(c+dx)} d\sqrt{\tan(c+dx)}}{d} + \\
& \frac{(A + iB)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1 \right)^{-n} \int \frac{\left(\frac{b \tan(c+dx)}{a} + 1 \right)^n}{i \tan(c+dx)+1} d\sqrt{\tan(c+dx)}}{d} \\
& \quad \downarrow \text{333} \\
& \frac{(A + iB) \sqrt{\tan(c + dx)} (a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1 \right)^{-n} \text{AppellF1} \left(\frac{1}{2}, 1, -n, \frac{3}{2}, -i \tan(c + dx), -\frac{b \tan(c+dx)}{a} \right)}{d} \\
& \frac{(A - iB) \sqrt{\tan(c + dx)} (a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1 \right)^{-n} \text{AppellF1} \left(\frac{1}{2}, 1, -n, \frac{3}{2}, i \tan(c + dx), -\frac{b \tan(c+dx)}{a} \right)}{d}
\end{aligned}$$

input `Int[((a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Sqrt[Tan[c + d*x]],x]`

output `((A + I*B)*AppellF1[1/2, 1, -n, 3/2, (-I)*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^n/(d*(1 + (b*Tan[c + d*x])/a)^n) + ((A - I*B)*AppellF1[1/2, 1, -n, 3/2, I*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Sqrt[Tan[c + d*x]]*(a + b*Tan[c + d*x])^n/(d*(1 + (b*Tan[c + d*x])/a)^n)`

3.663.3.1 Defintions of rubi rules used

rule 148 `Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_.)*((e_) + (f_.)*(x_))^(p_.), x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c + d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x]] /; FreeQ[{b, c, d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`

rule 333 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[a^p*c^q*x*AppellF1[1/2, -p, -q, 3/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

```
rule 334 Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Sim
p[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a)^FracPart[p]) Int[
(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p, q}, x] &&
NeQ[b*c - a*d, 0] && !(IntegerQ[p] || GtQ[a, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4085 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Sim
p[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x
]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n
] && EqQ[A^2 + B^2, 0]
```

```
rule 4086 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] &
& !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]
```

3.663.4 Maple [F]

$$\int \frac{(a + b \tan(dx + c))^n (A + B \tan(dx + c))}{\sqrt{\tan(dx + c)}} dx$$

```
input int((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2), x)
```

```
output int((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2), x)
```

3.663.5 Fracas [F]

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^n}{\sqrt{\tan(dx + c)}} dx$$

input `integrate((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm m="fricas")`

output `integral((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n/sqrt(tan(d*x + c)), x)`

3.663.6 Sympy [F]

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx$$

$$= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^n}{\sqrt{\tan(c + dx)}} dx$$

input `integrate((a+b*tan(d*x+c))**n*(A+B*tan(d*x+c))/tan(d*x+c)**(1/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**n/sqrt(tan(c + d*x)), x)`

3.663.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm m="maxima")`

output `Timed out`

3.663.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx = \text{Timed out}$$

input `integrate((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(1/2),x, algorithm m="giac")`

output `Timed out`

3.663.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\sqrt{\tan(c + dx)}} dx \\ &= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^n}{\sqrt{\tan(c + dx)}} dx \end{aligned}$$

input `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n)/tan(c + d*x)^(1/2),x)`

output `int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n)/tan(c + d*x)^(1/2), x)`

3.664 $\int \frac{(a+b \tan(c+dx))^n(A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

3.664.1 Optimal result 6299
 3.664.2 Mathematica [F] 6300
 3.664.3 Rubi [A] (warning: unable to verify) 6300
 3.664.4 Maple [F] 6303
 3.664.5 Fricas [F] 6303
 3.664.6 Sympy [F] 6303
 3.664.7 Maxima [F(-1)] 6304
 3.664.8 Giac [F(-1)] 6304
 3.664.9 Mupad [F(-1)] 6304

3.664.1 Optimal result

Integrand size = 33, antiderivative size = 169

$$\int \frac{(a + b \tan(c + dx))^n(A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx =$$

$$\frac{(A + iB) \operatorname{AppellF1}\left(-\frac{1}{2}, 1, -n, \frac{1}{2}, -i \tan(c + dx), -\frac{b \tan(c + dx)}{a}\right) (a + b \tan(c + dx))^n \left(1 + \frac{b \tan(c + dx)}{a}\right)^{-n}}{d \sqrt{\tan(c + dx)}} -$$

$$\frac{(A - iB) \operatorname{AppellF1}\left(-\frac{1}{2}, 1, -n, \frac{1}{2}, i \tan(c + dx), -\frac{b \tan(c + dx)}{a}\right) (a + b \tan(c + dx))^n \left(1 + \frac{b \tan(c + dx)}{a}\right)^{-n}}{d \sqrt{\tan(c + dx)}}$$

```
output -(A+I*B)*AppellF1(-1/2,1,-n,1/2,-I*tan(d*x+c),-b*tan(d*x+c)/a)*(a+b*tan(d*
x+c))^n/d/tan(d*x+c)^(1/2)/((1+b*tan(d*x+c)/a)^n)-(A-I*B)*AppellF1(-1/2,1,
-n,1/2,I*tan(d*x+c),-b*tan(d*x+c)/a)*(a+b*tan(d*x+c))^n/d/tan(d*x+c)^(1/2)
/((1+b*tan(d*x+c)/a)^n)
```

3.664.2 Mathematica [F]

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$= \int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx$$

input `Integrate[((a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2), x]`

output `Integrate[((a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2), x]`

3.664.3 Rubi [A] (warning: unable to verify)

Time = 0.61 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.95, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$, Rules used = {3042, 4086, 3042, 4085, 148, 395, 394}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))^n}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))^n}{\tan(c + dx)^{3/2}} dx$$

$$\downarrow \text{4086}$$

$$\frac{1}{2}(A + iB) \int \frac{(1 - i \tan(c + dx))(a + b \tan(c + dx))^n}{\tan^{\frac{3}{2}}(c + dx)} dx + \frac{1}{2}(A - iB) \int \frac{(i \tan(c + dx) + 1)(a + b \tan(c + dx))^n}{\tan^{\frac{3}{2}}(c + dx)} dx$$

$$\downarrow \text{3042}$$

$$\frac{1}{2}(A + iB) \int \frac{(1 - i \tan(c + dx))(a + b \tan(c + dx))^n}{\tan(c + dx)^{3/2}} dx + \frac{1}{2}(A - iB) \int \frac{(i \tan(c + dx) + 1)(a + b \tan(c + dx))^n}{\tan(c + dx)^{3/2}} dx$$

3.664. $\int \frac{(a+b \tan(c+dx))^n (A+B \tan(c+dx))}{\tan^{\frac{3}{2}}(c+dx)} dx$

$$\begin{aligned}
& \downarrow \text{4085} \\
& \frac{(A - iB) \int \frac{(a+b \tan(c+dx))^n}{(1-i \tan(c+dx)) \tan^{\frac{3}{2}}(c+dx)} d \tan(c+dx)}{2d} + \frac{(A + iB) \int \frac{(a+b \tan(c+dx))^n}{(i \tan(c+dx)+1) \tan^{\frac{3}{2}}(c+dx)} d \tan(c+dx)}{2d} \\
& \downarrow \text{148} \\
& \frac{(A - iB) \int \frac{\cot^2(c+dx)(a+b \tan(c+dx))^n}{1-i \tan(c+dx)} d \sqrt{\tan(c+dx)}}{d} + \\
& \frac{(A + iB) \int \frac{\cot^2(c+dx)(a+b \tan(c+dx))^n}{i \tan(c+dx)+1} d \sqrt{\tan(c+dx)}}{d} \\
& \downarrow \text{395} \\
& \frac{(A - iB)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1 \right)^{-n} \int \frac{\cot^2(c+dx) \left(\frac{b \tan(c+dx)}{a} + 1 \right)^n}{1-i \tan(c+dx)} d \sqrt{\tan(c+dx)}}{d} + \\
& \frac{(A + iB)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1 \right)^{-n} \int \frac{\cot^2(c+dx) \left(\frac{b \tan(c+dx)}{a} + 1 \right)^n}{i \tan(c+dx)+1} d \sqrt{\tan(c+dx)}}{d} \\
& \downarrow \text{394} \\
& \frac{(A + iB) \cot(c + dx)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1 \right)^{-n} \text{AppellF1} \left(-\frac{1}{2}, 1, -n, \frac{1}{2}, -i \tan(c + dx), -\frac{b \tan(c+dx)}{a} \right)}{d} \\
& \frac{(A - iB) \cot(c + dx)(a + b \tan(c + dx))^n \left(\frac{b \tan(c+dx)}{a} + 1 \right)^{-n} \text{AppellF1} \left(-\frac{1}{2}, 1, -n, \frac{1}{2}, i \tan(c + dx), -\frac{b \tan(c+dx)}{a} \right)}{d}
\end{aligned}$$

input `Int[((a + b*Tan[c + d*x])^n*(A + B*Tan[c + d*x]))/Tan[c + d*x]^(3/2),x]`

output `-(((A + I*B)*AppellF1[-1/2, 1, -n, 1/2, (-I)*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Cot[c + d*x]*(a + b*Tan[c + d*x])^n)/(d*(1 + (b*Tan[c + d*x])/a)^n)) - ((A - I*B)*AppellF1[-1/2, 1, -n, 1/2, I*Tan[c + d*x], -((b*Tan[c + d*x])/a)]*Cot[c + d*x]*(a + b*Tan[c + d*x])^n)/(d*(1 + (b*Tan[c + d*x])/a)^n)`

3.664.3.1 Defintions of rubi rules used

rule 148 `Int[((b_.)*(x_)^(m_)*((c_) + (d_.)*(x_)^(n_.))*((e_) + (f_.)*(x_)^(p_.),
x_] := With[{k = Denominator[m]}, Simp[k/b Subst[Int[x^(k*(m + 1) - 1)*(c
+ d*(x^k/b))^n*(e + f*(x^k/b))^p, x], x, (b*x)^(1/k)], x]] /; FreeQ[{b, c,
d, e, f, n, p}, x] && FractionQ[m] && IntegerQ[p]`

rule 394 `Int[((e_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_
, x_Symbol] := Simp[a^p*c^q*((e*x)^(m + 1)/(e*(m + 1)))*AppellF1[(m + 1)/2
, -p, -q, 1 + (m + 1)/2, (-b)*(x^2/a), (-d)*(x^2/c)], x] /; FreeQ[{a, b, c,
d, e, m, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, 1] && (Int
egerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])`

rule 395 `Int[((e_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_
, x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^
FracPart[p]) Int[(e*x)^m*(1 + b*(x^2/a))^p*(c + d*x^2)^q, x], x] /; FreeQ
[{a, b, c, d, e, m, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m,
1] && !(IntegerQ[p] || GtQ[a, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4085 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Sim
p[A^2/f Subst[Int[(a + b*x)^m*((c + d*x)^n/(A - B*x)), x], x, Tan[e + f*x
]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && NeQ[b*c - a*d, 0] &&
NeQ[a^2 + b^2, 0] && !IntegerQ[m] && !IntegerQ[n] && !IntegersQ[2*m, 2*n
] && EqQ[A^2 + B^2, 0]`

rule 4086 `Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Si
mp[(A + I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d*Tan[e + f*x])^n*(1 - I*T
an[e + f*x]), x], x] + Simp[(A - I*B)/2 Int[(a + b*Tan[e + f*x])^m*(c + d
*Tan[e + f*x])^n*(1 + I*Tan[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f, A
, B, m, n}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 + b^2, 0] && !IntegerQ[m] &
& !IntegerQ[n] && !IntegersQ[2*m, 2*n] && NeQ[A^2 + B^2, 0]`

3.664.4 Maple [F]

$$\int \frac{(a + b \tan(dx + c))^n (A + B \tan(dx + c))}{\tan(dx + c)^{\frac{3}{2}}} dx$$

input `int((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x)`

output `int((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x)`

3.664.5 Fricas [F]

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(B \tan(dx + c) + A)(b \tan(dx + c) + a)^n}{\tan(dx + c)^{\frac{3}{2}}} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm="fricas")`

output `integral((B*tan(d*x + c) + A)*(b*tan(d*x + c) + a)^n/tan(d*x + c)^(3/2), x)`

3.664.6 Sympy [F]

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(A + B \tan(c + dx))(a + b \tan(c + dx))^n}{\tan^{\frac{3}{2}}(c + dx)} dx \end{aligned}$$

input `integrate((a+b*tan(d*x+c))**n*(A+B*tan(d*x+c))/tan(d*x+c)**(3/2),x)`

output `Integral((A + B*tan(c + d*x))*(a + b*tan(c + d*x))**n/tan(c + d*x)**(3/2), x)`

3.664.7 Maxima [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm
m="maxima")
```

```
output Timed out
```

3.664.8 Giac [F(-1)]

Timed out.

$$\int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx = \text{Timed out}$$

```
input integrate((a+b*tan(d*x+c))^n*(A+B*tan(d*x+c))/tan(d*x+c)^(3/2),x, algorithm
m="giac")
```

```
output Timed out
```

3.664.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{(a + b \tan(c + dx))^n (A + B \tan(c + dx))}{\tan^{\frac{3}{2}}(c + dx)} dx \\ &= \int \frac{(A + B \tan(c + dx)) (a + b \tan(c + dx))^n}{\tan(c + dx)^{3/2}} dx \end{aligned}$$

```
input int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n)/tan(c + d*x)^(3/2),x)
```

```
output int(((A + B*tan(c + d*x))*(a + b*tan(c + d*x))^n)/tan(c + d*x)^(3/2), x)
```

3.665 $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx$

3.665.1 Optimal result	6305
3.665.2 Mathematica [A] (verified)	6305
3.665.3 Rubi [A] (verified)	6306
3.665.4 Maple [A] (verified)	6307
3.665.5 Fricas [A] (verification not implemented)	6308
3.665.6 Sympy [B] (verification not implemented)	6308
3.665.7 Maxima [B] (verification not implemented)	6309
3.665.8 Giac [F]	6310
3.665.9 Mupad [B] (verification not implemented)	6310

3.665.1 Optimal result

Integrand size = 39, antiderivative size = 63

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx$$

$$= \frac{a(iA + B)(c - ic \tan(e + fx))^n}{fn} - \frac{aB(c - ic \tan(e + fx))^{1+n}}{cf(1 + n)}$$

output `a*(I*A+B)*(c-I*c*tan(f*x+e))^n/f/n-a*B*(c-I*c*tan(f*x+e))^(1+n)/c/f/(1+n)`

3.665.2 Mathematica [A] (verified)

Time = 1.00 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.79

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx$$

$$= \frac{ia(c - ic \tan(e + fx))^n(A - iB + An + Bn \tan(e + fx))}{fn(1 + n)}$$

input `Integrate[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^n,x]`

output `(I*a*(c - I*c*Tan[e + f*x])^n*(A - I*B + A*n + B*n*Tan[e + f*x]))/(f*n*(1 + n))`

3.665.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.103$, Rules used = {3042, 4071, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int (A + B \tan(e + fx))(c - ic \tan(e + fx))^{n-1} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{53} \\
 & \frac{ac \int \left((A - iB)(c - ic \tan(e + fx))^{n-1} + \frac{iB(c - ic \tan(e + fx))^n}{c} \right) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{ac \left(\frac{(B + iA)(c - ic \tan(e + fx))^n}{cn} - \frac{B(c - ic \tan(e + fx))^{n+1}}{c^2(n+1)} \right)}{f}
 \end{aligned}$$

input `Int[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^n,x]`

output `(a*c*(((I*A + B)*(c - I*c*Tan[e + f*x])^n)/(c*n) - (B*(c - I*c*Tan[e + f*x])^(1 + n))/(c^2*(1 + n))))/f`

3.665.3.1 Defintions of rubi rules used

```
rule 53 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && IGtQ[m, 0] && ( !IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0])
|| LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.665.4 Maple [A] (verified)

Time = 1.22 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.27

method	result	size
derivativedivides	$\frac{(iAa+iAa+Ba)e^{n \ln(c-ic \tan(fx+e))}}{fn(1+n)} + \frac{iBa \tan(fx+e)e^{n \ln(c-ic \tan(fx+e))}}{f(1+n)}$	80
default	$\frac{(iAa+iAa+Ba)e^{n \ln(c-ic \tan(fx+e))}}{fn(1+n)} + \frac{iBa \tan(fx+e)e^{n \ln(c-ic \tan(fx+e))}}{f(1+n)}$	80
norman	$\frac{(iAa+iAa+Ba)e^{n \ln(c-ic \tan(fx+e))}}{fn(1+n)} + \frac{iBa \tan(fx+e)e^{n \ln(c-ic \tan(fx+e))}}{f(1+n)}$	80
risch	Expression too large to display	1058

```
input int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x,method=_RET
URNVERBOSE)
```

```
output 1/f/n/(1+n)*(I*A*a*n+I*A*a+B*a)*exp(n*ln(c-I*c*tan(f*x+e)))+I*B*a/f/(1+n)*
tan(f*x+e)*exp(n*ln(c-I*c*tan(f*x+e)))
```

3.665. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx$

3.665.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.48

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^n dx$$

$$= \frac{((iA - B)an + (iA + B)a + ((iA + B)an + (iA + B)a)e^{(2i fx + 2i e)}) \left(\frac{2c}{e^{(2i fx + 2i e)} + 1}\right)^n}{fn^2 + fn + (fn^2 + fn)e^{(2i fx + 2i e)}}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x, algo rithm="fricas")`

output `((I*A - B)*a*n + (I*A + B)*a + ((I*A + B)*a*n + (I*A + B)*a)*e^(2*I*f*x + 2*I*e))*(2*c/(e^(2*I*f*x + 2*I*e) + 1))^n/(f*n^2 + f*n + (f*n^2 + f*n)*e^(2*I*f*x + 2*I*e))`

3.665.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 394 vs. 2(48) = 96.

Time = 0.57 (sec) , antiderivative size = 394, normalized size of antiderivative = 6.25

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^n dx$$

$$= \begin{cases} x(A + B \tan(e)) (ia \tan(e) + a) (-ictan(e) + c)^n \\ \frac{2Aa}{2cf \tan(e+fx)+2icf} + \frac{2iBafx \tan(e+fx)}{2cf \tan(e+fx)+2icf} - \frac{2Bafx}{2cf \tan(e+fx)+2icf} - \frac{Ba \log(\tan^2(e+fx)+1) \tan(e+fx)}{2cf \tan(e+fx)+2icf} - \frac{iBa \log(\tan^2(e+fx)+1)}{2cf \tan(e+fx)+2icf} \\ Aax + \frac{iAa \log(\tan^2(e+fx)+1)}{2f} - iBax + \frac{Ba \log(\tan^2(e+fx)+1)}{2f} + \frac{iBa \tan(e+fx)}{f} \\ \frac{iAan(-ictan(e+fx)+c)^n}{fn^2+fn} + \frac{iAa(-ictan(e+fx)+c)^n}{fn^2+fn} + \frac{iBan(-ictan(e+fx)+c)^n \tan(e+fx)}{fn^2+fn} + \frac{Ba(-ictan(e+fx)+c)^n}{fn^2+fn} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x)`

```
output Piecewise((x*(A + B*tan(e))*(I*a*tan(e) + a)*(-I*c*tan(e) + c)**n, Eq(f, 0
)), (2*A*a/(2*c*f*tan(e + f*x) + 2*I*c*f) + 2*I*B*a*f*x*tan(e + f*x)/(2*c*
f*tan(e + f*x) + 2*I*c*f) - 2*B*a*f*x/(2*c*f*tan(e + f*x) + 2*I*c*f) - B*a
*log(tan(e + f*x)**2 + 1)*tan(e + f*x)/(2*c*f*tan(e + f*x) + 2*I*c*f) - I*
B*a*log(tan(e + f*x)**2 + 1)/(2*c*f*tan(e + f*x) + 2*I*c*f) - 2*I*B*a/(2*c
*f*tan(e + f*x) + 2*I*c*f), Eq(n, -1)), (A*a*x + I*A*a*log(tan(e + f*x)**2
+ 1)/(2*f) - I*B*a*x + B*a*log(tan(e + f*x)**2 + 1)/(2*f) + I*B*a*tan(e +
f*x)/f, Eq(n, 0)), (I*A*a*n*(-I*c*tan(e + f*x) + c)**n/(f*n**2 + f*n) + I
*A*a*(-I*c*tan(e + f*x) + c)**n/(f*n**2 + f*n) + I*B*a*n*(-I*c*tan(e + f*x
) + c)**n*tan(e + f*x)/(f*n**2 + f*n) + B*a*(-I*c*tan(e + f*x) + c)**n/(f*
n**2 + f*n), True))
```

3.665.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 311 vs. $2(57) = 114$.

Time = 0.45 (sec) , antiderivative size = 311, normalized size of antiderivative = 4.94

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^n dx$$

$$= \frac{((A - iB)ac^n n + (A - iB)ac^n)2^n \cos(-2fx + n \arctan(\sin(2fx + 2e), \cos(2fx + 2e) + 1) - 2e) + ((A + iB)ac^n n + (A - iB)ac^n)2^n \cos(n \arctan(\sin(2fx + 2e), \cos(2fx + 2e) + 1)) - ((I*A + B)ac^n n + (I*A + B)ac^n)2^n \sin(-2fx + n \arctan(\sin(2fx + 2e), \cos(2fx + 2e) + 1)) - ((I*A - B)ac^n n + (I*A + B)ac^n)2^n \sin(n \arctan(\sin(2fx + 2e), \cos(2fx + 2e) + 1)))}{((-I*n^2 + (-I*n^2 - I*n)*\cos(2*f*x + 2*e) + (n^2 + n)*\sin(2*f*x + 2*e) - I*n)*(\cos(2*f*x + 2*e)^2 + \sin(2*f*x + 2*e)^2 + 2*\cos(2*f*x + 2*e) + 1)^(1/2*n)*f)}$$

```
input integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x, algo
rithm="maxima")
```

```
output (((A - I*B)*a*c^n*n + (A - I*B)*a*c^n)*2^n*cos(-2*f*x + n*arctan2(sin(2*f*
x + 2*e), cos(2*f*x + 2*e) + 1) - 2*e) + ((A + I*B)*a*c^n*n + (A - I*B)*a*
c^n)*2^n*cos(n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)) - ((I*A +
B)*a*c^n*n + (I*A + B)*a*c^n)*2^n*sin(-2*f*x + n*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e) + 1) - 2*e) - ((I*A - B)*a*c^n*n + (I*A + B)*a*c^n)*2^n*
sin(n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)))/((-I*n^2 + (-I*n^2
- I*n)*cos(2*f*x + 2*e) + (n^2 + n)*sin(2*f*x + 2*e) - I*n)*(cos(2*f*x +
2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^(1/2*n)*f)
```

3.665. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^n dx$

3.665.8 Giac [F]

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ict \tan(e + fx))^n dx$$

$$= \int (B \tan(fx + e) + A)(ia \tan(fx + e) + a)(-ict \tan(fx + e) + c)^n dx$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x, algo rithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)*(-I*c*tan(f*x + e) + c)^n, x)`

3.665.9 Mupad [B] (verification not implemented)

Time = 1.20 (sec) , antiderivative size = 128, normalized size of antiderivative = 2.03

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ict \tan(e + fx))^n dx =$$

$$\frac{(\cos(e + fx) - \sin(e + fx) i) \left(c - \frac{c \sin(e + fx) i}{\cos(e + fx)} \right)^n \left(\frac{a(A - B i + A n + B n i)}{f n (n i + i)} + \frac{a(A - B i) (\cos(2e + 2fx) + \sin(2e + 2fx) i)}{f n (n i + i)} \right)}{2 \cos(e + fx)}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)*(c - c*tan(e + f*x)*1i)^n ,x)`

output `-((cos(e + f*x) - sin(e + f*x)*1i)*(c - (c*sin(e + f*x)*1i)/cos(e + f*x))^n*((a*(A - B*1i + A*n + B*n*1i))/(f*n*(n*1i + 1i)) + (a*(A - B*1i)*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i)*(n + 1))/(f*n*(n*1i + 1i)))/(2*cos(e + f*x))`

3.666 $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^4 dx$

3.666.1 Optimal result	6311
3.666.2 Mathematica [A] (verified)	6311
3.666.3 Rubi [A] (verified)	6312
3.666.4 Maple [A] (verified)	6313
3.666.5 Fricas [B] (verification not implemented)	6314
3.666.6 Sympy [B] (verification not implemented)	6314
3.666.7 Maxima [A] (verification not implemented)	6315
3.666.8 Giac [B] (verification not implemented)	6315
3.666.9 Mupad [B] (verification not implemented)	6316

3.666.1 Optimal result

Integrand size = 39, antiderivative size = 59

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^4 dx$$

$$= \frac{a(iA + B)c^4(1 - i \tan(e + fx))^4}{4f} - \frac{aBc^4(1 - i \tan(e + fx))^5}{5f}$$

output `1/4*a*(I*A+B)*c^4*(1-I*tan(f*x+e))^4/f-1/5*a*B*c^4*(1-I*tan(f*x+e))^5/f`

3.666.2 Mathematica [A] (verified)

Time = 1.45 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.31

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^4 dx$$

$$= \frac{iac^4(B(i + \tan(e + fx))^5 + \frac{5}{4}(A - iB) \tan(e + fx) (-4i - 6 \tan(e + fx) + 4i \tan^2(e + fx) + \tan^3(e + fx)))}{5f}$$

input `Integrate[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^4,x]`

output `((I/5)*a*c^4*(B*(I + Tan[e + f*x])^5 + (5*(A - I*B)*Tan[e + f*x]*(-4*I - 6*Tan[e + f*x] + (4*I)*Tan[e + f*x]^2 + Tan[e + f*x]^3))/4))/f`

3.666.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.90, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.128$, Rules used = {3042, 4071, 27, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(e + fx))(c - ic \tan(e + fx))^4 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(e + fx))(c - ic \tan(e + fx))^4 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int c^3 (1 - i \tan(e + fx))^3 (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{ac^4 \int (1 - i \tan(e + fx))^3 (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{49} \\
 & \frac{ac^4 \int (iB(1 - i \tan(e + fx))^4 + (A - iB)(1 - i \tan(e + fx))^3) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{ac^4 (\frac{1}{4}(B + iA)(1 - i \tan(e + fx))^4 - \frac{1}{5}B(1 - i \tan(e + fx))^5)}{f}
 \end{aligned}$$

input `Int[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^4,x]`

output `(a*c^4*(((I*A + B)*(1 - I*Tan[e + f*x])^4)/4 - (B*(1 - I*Tan[e + f*x])^5)/5))/f`

3.666.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.666.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.95

method	result
risch	$\frac{4a^4(5iAe^{2i(fx+e)}+5Be^{2i(fx+e)}+5iA-3B)}{5f(e^{2i(fx+e)}+1)^5}$
derivativedivides	$\frac{ia^4\left(\frac{B \tan(fx+e)^5}{5} + \frac{(3iB+A) \tan(fx+e)^4}{4} + \frac{(3iA-3B) \tan(fx+e)^3}{3} + \frac{(-iB-3A) \tan(fx+e)^2}{2} - i \tan(fx+e)A\right)}{f}$
default	$\frac{ia^4\left(\frac{B \tan(fx+e)^5}{5} + \frac{(3iB+A) \tan(fx+e)^4}{4} + \frac{(3iA-3B) \tan(fx+e)^3}{3} + \frac{(-iB-3A) \tan(fx+e)^2}{2} - i \tan(fx+e)A\right)}{f}$
norman	$\frac{Aa^4 \tan(fx+e)}{f} - \frac{(-iAa^4+3Ba^4) \tan(fx+e)^4}{4f} + \frac{(-3iAa^4+Ba^4) \tan(fx+e)^2}{2f} - \frac{(iBa^4+Aa^4) \tan(fx+e)}{f}$
parallelrisch	$\frac{4iBa^4 \tan(fx+e)^5+5iA \tan(fx+e)^4 a^4-20iB \tan(fx+e)^3 a^4-15B \tan(fx+e)^4 a^4-30iA \tan(fx+e)^2 a^4-20A \tan(fx+e) a^4}{20f}$
parts	$\frac{(-3iAa^4+Ba^4) \ln(1+\tan(fx+e)^2)}{2f} + \frac{(-3iBa^4-2Aa^4)(\tan(fx+e)-\arctan(\tan(fx+e)))}{f} + \frac{(-2iAa^4-2Ba^4) \tan(fx+e)}{f}$

3.666. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^4 dx$


```
input int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4,x,method=_RET
URNVERBOSE)
```

```
output 4/5*a*c^4*(5*I*A*exp(2*I*(f*x+e))+5*B*exp(2*I*(f*x+e))+5*I*A-3*B)/f/(exp(2
*I*(f*x+e))+1)^5
```

3.666.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 100 vs. $2(49) = 98$.

Time = 0.24 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.69

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^4 dx =$$

$$\frac{4(5(-iA - B)ac^4e^{(2ifx+2ie)} + (-5iA + 3B)ac^4)}{5(fe^{10ifx+10ie} + 5fe^{8ifx+8ie} + 10fe^{6ifx+6ie} + 10fe^{4ifx+4ie} + 5fe^{2ifx+2ie} + f)}$$

```
input integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4,x, algo
rithm="fricas")
```

```
output -4/5*(5*(-I*A - B)*a*c^4*e^(2*I*f*x + 2*I*e) + (-5*I*A + 3*B)*a*c^4)/(f*e^
(10*I*f*x + 10*I*e) + 5*f*e^(8*I*f*x + 8*I*e) + 10*f*e^(6*I*f*x + 6*I*e) +
10*f*e^(4*I*f*x + 4*I*e) + 5*f*e^(2*I*f*x + 2*I*e) + f)
```

3.666.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 155 vs. $2(46) = 92$.

Time = 0.31 (sec) , antiderivative size = 155, normalized size of antiderivative = 2.63

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^4 dx$$

$$= \frac{20iAac^4 - 12Bac^4 + (20iAac^4e^{2ie} + 20Bac^4e^{2ie})e^{2ifx}}{5fe^{10ie}e^{10ifx} + 25fe^{8ie}e^{8ifx} + 50fe^{6ie}e^{6ifx} + 50fe^{4ie}e^{4ifx} + 25fe^{2ie}e^{2ifx} + 5f}$$

```
input integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**4,x)
```

3.666. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^4 dx$

output $(20IAac^{**4} - 12Bac^{**4} + (20IAac^{**4} \exp(2Ie) + 20Bac^{**4} \exp(2Ie)) \exp(2Ifx)) / (5f \exp(10Ie) \exp(10Ifx) + 25f \exp(8Ie) \exp(8Ifx) + 50f \exp(6Ie) \exp(6Ifx) + 50f \exp(4Ie) \exp(4Ifx) + 25f \exp(2Ie) \exp(2Ifx) + 5f)$

3.666.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.61

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^4 dx = \frac{-4i Bac^4 \tan(fx + e)^5 + 5(-iA + 3B)ac^4 \tan(fx + e)^4 + 20(A + iB)ac^4 \tan(fx + e)^3 + 10(3iA - 20Aac^4 \tan(fx + e))}{20f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4,x, algorithm="maxima")`

output $-1/20*(-4IBac^4 \tan(fx + e)^5 + 5(-IA + 3B)ac^4 \tan(fx + e)^4 + 20(A + IB)ac^4 \tan(fx + e)^3 + 10(3IA - B)ac^4 \tan(fx + e)^2 - 20Aac^4 \tan(fx + e)) / f$

3.666.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 112 vs. $2(49) = 98$.

Time = 0.68 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.90

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^4 dx = \frac{4(-5iAac^4 e^{(2ifx+2ie)} - 5Bac^4 e^{(2ifx+2ie)} - 5iAac^4 + 3Bac^4)}{5(fe^{(10ifx+10ie)} + 5fe^{(8ifx+8ie)} + 10fe^{(6ifx+6ie)} + 10fe^{(4ifx+4ie)} + 5fe^{(2ifx+2ie)} + f)}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4,x, algorithm="giac")`

output $-4/5*(-5IAac^4 e^{(2Ifx + 2Ie)} - 5Bac^4 e^{(2Ifx + 2Ie)} - 5IAac^4 + 3Bac^4) / (f e^{(10Ifx + 10Ie)} + 5f e^{(8Ifx + 8Ie)} + 10f e^{(6Ifx + 6Ie)} + 10f e^{(4Ifx + 4Ie)} + 5f e^{(2Ifx + 2Ie)} + f)$

3.666. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^4 dx$

3.666.9 Mupad [B] (verification not implemented)

Time = 9.16 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.69

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ict \tan(e + fx))^4 dx$$

$$= \frac{\frac{1iBac^4 \tan(e+fx)^5}{5} + \frac{1ia(A+B3i)c^4 \tan(e+fx)^4}{4} + 1ia(-B + A1i) c^4 \tan(e + fx)^3 - \frac{1ia(3A+B1i)c^4 \tan(e+fx)^2}{2} + \dots}{f}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)*(c - c*tan(e + f*x)*1i)^4, x)`

output `(A*a*c^4*tan(e + f*x) + (a*c^4*tan(e + f*x)^4*(A + B*3i)*1i)/4 + (B*a*c^4*tan(e + f*x)^5*1i)/5 + a*c^4*tan(e + f*x)^3*(A*1i - B)*1i - (a*c^4*tan(e + f*x)^2*(3*A + B*1i)*1i)/2)/f`

3.667 $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^3 dx$

3.667.1 Optimal result	6317
3.667.2 Mathematica [A] (verified)	6317
3.667.3 Rubi [A] (verified)	6318
3.667.4 Maple [A] (verified)	6319
3.667.5 Fricas [A] (verification not implemented)	6320
3.667.6 Sympy [B] (verification not implemented)	6320
3.667.7 Maxima [A] (verification not implemented)	6321
3.667.8 Giac [B] (verification not implemented)	6321
3.667.9 Mupad [B] (verification not implemented)	6322

3.667.1 Optimal result

Integrand size = 39, antiderivative size = 59

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^3 dx$$

$$= \frac{a(iA + B)c^3(1 - i \tan(e + fx))^3}{3f} - \frac{aBc^3(1 - i \tan(e + fx))^4}{4f}$$

output `1/3*a*(I*A+B)*c^3*(1-I*tan(f*x+e))^3/f-1/4*a*B*c^3*(1-I*tan(f*x+e))^4/f`

3.667.2 Mathematica [A] (verified)

Time = 3.13 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.19

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^3 dx =$$

$$\frac{ac^3(3B - 12A \tan(e + fx) + (12iA - 6B) \tan^2(e + fx) + 4(A + 2iB) \tan^3(e + fx) + 3B \tan^4(e + fx))}{12f}$$

input `Integrate[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^3,x]`

output `-1/12*(a*c^3*(3*B - 12*A*Tan[e + f*x] + ((12*I)*A - 6*B)*Tan[e + f*x]^2 + 4*(A + (2*I)*B)*Tan[e + f*x]^3 + 3*B*Tan[e + f*x]^4))/f`

3.667.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.90, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.128$, Rules used = {3042, 4071, 27, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(e + fx))(c - ic \tan(e + fx))^3 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(e + fx))(c - ic \tan(e + fx))^3 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int c^2 (1 - i \tan(e + fx))^2 (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{ac^3 \int (1 - i \tan(e + fx))^2 (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{49} \\
 & \frac{ac^3 \int (iB(1 - i \tan(e + fx))^3 + (A - iB)(1 - i \tan(e + fx))^2) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{ac^3 (\frac{1}{3}(B + iA)(1 - i \tan(e + fx))^3 - \frac{1}{4}B(1 - i \tan(e + fx))^4)}{f}
 \end{aligned}$$

input `Int[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^3,x]`

output `(a*c^3*(((I*A + B)*(1 - I*Tan[e + f*x])^3)/3 - (B*(1 - I*Tan[e + f*x])^4)/4))/f`

3.667.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.667.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.95

method	result
risch	$\frac{4a^3(2iAe^{2i(fx+e)}+2Be^{2i(fx+e)}+2iA-B)}{3f(e^{2i(fx+e)}+1)^4}$
derivativedivides	$\frac{ac^3\left(-\frac{B\tan(fx+e)^4}{4}-\frac{(2iB+A)\tan(fx+e)^3}{3}-\frac{(2iA-B)\tan(fx+e)^2}{2}+A\tan(fx+e)\right)}{f}$
default	$\frac{ac^3\left(-\frac{B\tan(fx+e)^4}{4}-\frac{(2iB+A)\tan(fx+e)^3}{3}-\frac{(2iA-B)\tan(fx+e)^2}{2}+A\tan(fx+e)\right)}{f}$
norman	$\frac{Aac^3\tan(fx+e)}{f}-\frac{(2iBa^3+Aa^3)\tan(fx+e)^3}{3f}+\frac{(-2iAa^3+Ba^3)\tan(fx+e)^2}{2f}-\frac{Ba^3\tan(fx+e)^4}{4f}$
parallelrisch	$-\frac{8iB\tan(fx+e)^3a^3+3B\tan(fx+e)^4a^3+12iA\tan(fx+e)^2a^3+4A\tan(fx+e)^3a^3-6B\tan(fx+e)^2a^3-12A\tan(fx+e)a^3}{12f}$
parts	$\frac{(-2iAa^3+Ba^3)\ln(1+\tan(fx+e)^2)}{2f}+\frac{(-2iBa^3-Aa^3)\left(\frac{\tan(fx+e)^3}{3}-\tan(fx+e)+\arctan(\tan(fx+e))\right)}{f}+A$

3.667. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^3 dx$

input `int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output $\frac{4}{3}ac^3(2IA\exp(2I(fx+e))+2B\exp(2I(fx+e))+2IA-B)/f/(\exp(2I(fx+e))+1)^4$

3.667.5 Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.46

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^3 dx$$

$$= -\frac{4(2(-iA - B)ac^3e^{2ifx+2ie} + (-2iA + B)ac^3)}{3(fe^{8ifx+8ie} + 4fe^{6ifx+6ie} + 6fe^{4ifx+4ie} + 4fe^{2ifx+2ie} + f)}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3,x, algorithm="fricas")`

output $-\frac{4}{3}(2(-IA - B)ac^3e^{2Ifx+2Ie} + (-2IA + B)ac^3)/(fe^{8Ifx+8Ie} + 4ffe^{6Ifx+6Ie} + 6ffe^{4Ifx+4Ie} + 4ffe^{2Ifx+2Ie} + f)$

3.667.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 136 vs. $2(46) = 92$.

Time = 0.28 (sec) , antiderivative size = 136, normalized size of antiderivative = 2.31

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^3 dx$$

$$= \frac{8iAac^3 - 4Bac^3 + (8iAac^3e^{2ie} + 8Bac^3e^{2ie})e^{2ifx}}{3fe^{8ie}e^{8ifx} + 12fe^{6ie}e^{6ifx} + 18fe^{4ie}e^{4ifx} + 12fe^{2ie}e^{2ifx} + 3f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**3,x)`

output $(8IAac^3 - 4Bac^3 + (8iAac^3\exp(2Ie) + 8Bac^3\exp(2Ie))\exp(2Ifx))/((3f\exp(8Ie)\exp(8Ifx) + 12f\exp(6Ie)\exp(6Ifx) + 18f\exp(4Ie)\exp(4Ifx) + 12f\exp(2Ie)\exp(2Ifx) + 3f)$

3.667. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^3 dx$

3.667.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.22

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ict \tan(e + fx))^3 dx = \frac{3 Bac^3 \tan(fx + e)^4 + 4(A + 2iB)ac^3 \tan(fx + e)^3 - 6(-2iA + B)ac^3 \tan(fx + e)^2 - 12Aac^3 \tan(fx + e)}{12f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3,x, algorithm="maxima")`

output `-1/12*(3*B*a*c^3*tan(f*x + e)^4 + 4*(A + 2*I*B)*a*c^3*tan(f*x + e)^3 - 6*(-2*I*A + B)*a*c^3*tan(f*x + e)^2 - 12*A*a*c^3*tan(f*x + e))/f`

3.667.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 99 vs. $2(49) = 98$.

Time = 0.52 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.68

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ict \tan(e + fx))^3 dx = -\frac{4(-2iAac^3e^{(2i fx+2i e)} - 2Bac^3e^{(2i fx+2i e)} - 2iAac^3 + Bac^3)}{3(fe^{(8i fx+8i e)} + 4fe^{(6i fx+6i e)} + 6fe^{(4i fx+4i e)} + 4fe^{(2i fx+2i e)} + f)}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3,x, algorithm="giac")`

output `-4/3*(-2*I*A*a*c^3*e^(2*I*f*x + 2*I*e) - 2*B*a*c^3*e^(2*I*f*x + 2*I*e) - 2*I*A*a*c^3 + B*a*c^3)/(f*e^(8*I*f*x + 8*I*e) + 4*f*e^(6*I*f*x + 6*I*e) + 6*f*e^(4*I*f*x + 4*I*e) + 4*f*e^(2*I*f*x + 2*I*e) + f)`

3.667.9 Mupad [B] (verification not implemented)

Time = 8.93 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.29

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^3 dx$$

$$= -\frac{\frac{B a c^3 \tan(e+fx)^4}{4} + \frac{a(A+B2i)c^3 \tan(e+fx)^3}{3} + \frac{a(-B+A2i)c^3 \tan(e+fx)^2}{2} - A a c^3 \tan(e + fx)}{f}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)*(c - c*tan(e + f*x)*1i)^3, x)`

output `-((a*c^3*tan(e + f*x)^3*(A + B*2i))/3 - A*a*c^3*tan(e + f*x) + (B*a*c^3*tan(e + f*x)^4)/4 + (a*c^3*tan(e + f*x)^2*(A*2i - B))/2)/f`

3.668 $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx$

3.668.1 Optimal result	6323
3.668.2 Mathematica [A] (verified)	6323
3.668.3 Rubi [A] (verified)	6324
3.668.4 Maple [A] (verified)	6325
3.668.5 Fricas [A] (verification not implemented)	6326
3.668.6 Sympy [B] (verification not implemented)	6326
3.668.7 Maxima [A] (verification not implemented)	6327
3.668.8 Giac [A] (verification not implemented)	6327
3.668.9 Mupad [B] (verification not implemented)	6328

3.668.1 Optimal result

Integrand size = 39, antiderivative size = 66

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx$$

$$= \frac{aAc^2 \tan(e + fx)}{f} - \frac{a(iA - B)c^2 \tan^2(e + fx)}{2f} - \frac{iaBc^2 \tan^3(e + fx)}{3f}$$

output `a*A*c^2*tan(f*x+e)/f-1/2*a*(I*A-B)*c^2*tan(f*x+e)^2/f-1/3*I*a*B*c^2*tan(f*x+e)^3/f`

3.668.2 Mathematica [A] (verified)

Time = 1.94 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.82

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx$$

$$= \frac{ac^2(-2B + 6A \tan(e + fx) + 3(-iA + B) \tan^2(e + fx) - 2iB \tan^3(e + fx))}{6f}$$

input `Integrate[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^2,x]`

output $(a*c^2*(-2*B + 6*A*\text{Tan}[e + f*x] + 3*((-I)*A + B)*\text{Tan}[e + f*x]^2 - (2*I)*B*\text{Tan}[e + f*x]^3))/(6*f)$

3.668.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.80, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.128$, Rules used = {3042, 4071, 27, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))(c - ict \tan(e + fx))^2 (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))(c - ict \tan(e + fx))^2 (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int c(1 - i \tan(e + fx))(A + B \tan(e + fx)) d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{ac^2 \int (1 - i \tan(e + fx))(A + B \tan(e + fx)) d \tan(e + fx)}{f} \\ & \quad \downarrow \text{49} \\ & \frac{ac^2 \int (-iB \tan^2(e + fx) + (B - iA) \tan(e + fx) + A) d \tan(e + fx)}{f} \\ & \quad \downarrow \text{2009} \\ & \frac{ac^2 \left(-\frac{1}{2}(-B + iA) \tan^2(e + fx) + A \tan(e + fx) - \frac{1}{3}iB \tan^3(e + fx) \right)}{f} \end{aligned}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])*(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^2, x]$

output $(a*c^2*(A*\text{Tan}[e + f*x] - ((I*A - B)*\text{Tan}[e + f*x]^2)/2 - (I/3)*B*\text{Tan}[e + f*x]^3))/f$

3.668. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ict \tan(e + fx))^2 dx$

3.668.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.668.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.74

method	result
derivativedivides	$-\frac{ia c^2 \left(\frac{B \tan^3(fx+e)}{3} + \frac{(iB+A) \tan^2(fx+e)}{2} + i \tan(fx+e)A \right)}{f}$
default	$-\frac{ia c^2 \left(\frac{B \tan^3(fx+e)}{3} + \frac{(iB+A) \tan^2(fx+e)}{2} + i \tan(fx+e)A \right)}{f}$
risch	$\frac{2a c^2 (3iA e^{2i(fx+e)} + 3B e^{2i(fx+e)} + 3iA - B)}{3f (e^{2i(fx+e)} + 1)^3}$
norman	$\frac{aA c^2 \tan(fx+e)}{f} + \frac{(-iAa c^2 + Ba c^2) \tan^2(fx+e)}{2f} - \frac{iaB c^2 \tan^3(fx+e)}{3f}$
parallelrisch	$-\frac{2iaB c^2 \tan^3(fx+e) + 3iA \tan^2(fx+e)^2 a c^2 - 3B \tan^2(fx+e)^2 a c^2 - 6A \tan(fx+e) a c^2}{6f}$
parts	$\frac{(-iAa c^2 + Ba c^2) \ln(1 + \tan(fx+e)^2)}{2f} + \frac{(-iAa c^2 + Ba c^2) \left(\frac{\tan^2(fx+e)}{2} - \frac{\ln(1 + \tan^2(fx+e))}{2} \right)}{f} + \frac{(-iBa c^2 + Aa c^2)}{f}$

3.668. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^2 dx$

input `int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2,x,method=_RETURNERVERBOSE)`

output `-I/f*a*c^2*(1/3*B*tan(f*x+e)^3+1/2*(A+I*B)*tan(f*x+e)^2+I*A*tan(f*x+e))`

3.668.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.12

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx$$

$$= -\frac{2(3(-iA - B)ac^2e^{2ifx+2ie} + (-3iA + B)ac^2)}{3(fe^{6ifx+6ie} + 3fe^{4ifx+4ie} + 3fe^{2ifx+2ie} + f)}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2,x, algorithm="fricas")`

output `-2/3*(3*(-I*A - B)*a*c^2*e^(2*I*f*x + 2*I*e) + (-3*I*A + B)*a*c^2)/(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)`

3.668.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 117 vs. 2(56) = 112.

Time = 0.21 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.77

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx$$

$$= \frac{6iAac^2 - 2Bac^2 + (6iAac^2e^{2ie} + 6Bac^2e^{2ie})e^{2ifx}}{3fe^{6ie}e^{6ifx} + 9fe^{4ie}e^{4ifx} + 9fe^{2ie}e^{2ifx} + 3f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**2,x)`

output `(6*I*A*a*c**2 - 2*B*a*c**2 + (6*I*A*a*c**2*exp(2*I*e) + 6*B*a*c**2*exp(2*I*e))*exp(2*I*f*x))/(3*f*exp(6*I*e)*exp(6*I*f*x) + 9*f*exp(4*I*e)*exp(4*I*f*x) + 9*f*exp(2*I*e)*exp(2*I*f*x) + 3*f)`

3.668. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx$

3.668.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.83

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx$$

$$= \frac{-2i Bac^2 \tan(fx + e)^3 - 3(iA - B)ac^2 \tan(fx + e)^2 + 6Aac^2 \tan(fx + e)}{6f}$$

```
input integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2,x, algo
rithm="maxima")
```

```
output 1/6*(-2*I*B*a*c^2*tan(f*x + e)^3 - 3*(I*A - B)*a*c^2*tan(f*x + e)^2 + 6*A*
a*c^2*tan(f*x + e))/f
```

3.668.8 Giac [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.32

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx$$

$$= \frac{2(-3iAac^2e^{(2ifx+2ie)} - 3Bac^2e^{(2ifx+2ie)} - 3iAac^2 + Bac^2)}{3(fe^{(6ifx+6ie)} + 3fe^{(4ifx+4ie)} + 3fe^{(2ifx+2ie)} + f)}$$

```
input integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2,x, algo
rithm="giac")
```

```
output -2/3*(-3*I*A*a*c^2*e^(2*I*f*x + 2*I*e) - 3*B*a*c^2*e^(2*I*f*x + 2*I*e) - 3
*I*A*a*c^2 + B*a*c^2)/(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3
*f*e^(2*I*f*x + 2*I*e) + f)
```

3.668.9 Mupad [B] (verification not implemented)

Time = 8.93 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.76

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^2 dx$$

$$= \frac{a c^2 \tan(e + fx) (6 A - A \tan(e + fx) 3i + 3 B \tan(e + fx) - B \tan(e + fx)^2 2i)}{6 f}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)*(c - c*tan(e + f*x)*1i)^2, x)`

output `(a*c^2*tan(e + f*x)*(6*A - A*tan(e + f*x)*3i + 3*B*tan(e + f*x) - B*tan(e + f*x)^2*2i))/(6*f)`

3.669 $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx)) dx$

3.669.1 Optimal result	6329
3.669.2 Mathematica [A] (verified)	6329
3.669.3 Rubi [A] (verified)	6330
3.669.4 Maple [A] (verified)	6331
3.669.5 Fricas [C] (verification not implemented)	6331
3.669.6 Sympy [C] (verification not implemented)	6332
3.669.7 Maxima [A] (verification not implemented)	6332
3.669.8 Giac [B] (verification not implemented)	6333
3.669.9 Mupad [B] (verification not implemented)	6333

3.669.1 Optimal result

Integrand size = 37, antiderivative size = 32

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx)) dx$$

$$= \frac{aActan(e + fx)}{f} + \frac{aBc \tan^2(e + fx)}{2f}$$

output `a*A*c*tan(f*x+e)/f+1/2*a*B*c*tan(f*x+e)^2/f`

3.669.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx)) dx$$

$$= \frac{aBc \sec^2(e + fx)}{2f} + \frac{aActan(e + fx)}{f}$$

input `Integrate[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x]),x]`

output `(a*B*c*Sec[e + f*x]^2)/(2*f) + (a*A*c*Tan[e + f*x])/f`

3.669. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx)) dx$

3.669.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.081$, Rules used = {3042, 4071, 17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))(c - ictan(e + fx))(A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))(c - ictan(e + fx))(A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\ & \quad \downarrow \text{17} \\ & \frac{ac(A + B \tan(e + fx))^2}{2Bf} \end{aligned}$$

input `Int[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x]),x]`

output `(a*c*(A + B*Tan[e + f*x])^2)/(2*B*f)`

3.669.3.1 Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_.))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.669.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.84

method	result
derivativedivides	$\frac{ac \left(\frac{B \tan(fx+e)^2}{2} + A \tan(fx+e) \right)}{f}$
default	$\frac{ac \left(\frac{B \tan(fx+e)^2}{2} + A \tan(fx+e) \right)}{f}$
parallelrisch	$\frac{Bac \tan(fx+e)^2 + 2acA \tan(fx+e)}{2f}$
norman	$\frac{aAc \tan(fx+e)}{f} + \frac{aBc \tan(fx+e)^2}{2f}$
risch	$\frac{2ac(iAe^{2i(fx+e)} + Be^{2i(fx+e)} + iA)}{f(e^{2i(fx+e)} + 1)^2}$
parts	$acAx + \frac{Bac \ln(1 + \tan(fx+e)^2)}{2f} + \frac{Bac \left(\frac{\tan(fx+e)^2}{2} - \frac{\ln(1 + \tan(fx+e)^2)}{2} \right)}{f} + \frac{acA(\tan(fx+e) - \arctan(\tan(fx+e)))}{f}$

```
input int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x,method=_RETUR
NVERBOSE)
```

```
output 1/f*a*c*(1/2*B*tan(f*x+e)^2+A*tan(f*x+e))
```

3.669.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.24 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.69

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx)) dx$$

$$= -\frac{2((-iA - B)ace^{(2i fx + 2ie)} - iAac)}{fe^{(4i fx + 4ie)} + 2fe^{(2i fx + 2ie)} + f}$$

3.669. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx)) dx$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x, algorithm="fricas")`

output `-2*((-I*A - B)*a*c*e^(2*I*f*x + 2*I*e) - I*A*a*c)/(f*e^(4*I*f*x + 4*I*e) + 2*f*e^(2*I*f*x + 2*I*e) + f)`

3.669.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 82, normalized size of antiderivative = 2.56

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx)) dx$$

$$= \frac{2iAac + (2iAace^{2ie} + 2Bace^{2ie})e^{2ifx}}{fe^{4ie}e^{4ifx} + 2fe^{2ie}e^{2ifx} + f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x)`

output `(2*I*A*a*c + (2*I*A*a*c*exp(2*I*e) + 2*B*a*c*exp(2*I*e))*exp(2*I*f*x))/(f*exp(4*I*e)*exp(4*I*f*x) + 2*f*exp(2*I*e)*exp(2*I*f*x) + f)`

3.669.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx)) dx$$

$$= \frac{Bactan(fx + e)^2 + 2Aactan(fx + e)}{2f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x, algorithm="maxima")`

output `1/2*(B*a*c*tan(f*x + e)^2 + 2*A*a*c*tan(f*x + e))/f`

3.669.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 106 vs. $2(30) = 60$.

Time = 0.35 (sec) , antiderivative size = 106, normalized size of antiderivative = 3.31

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx)) dx$$

$$= \frac{B a c \tan(fx)^2 \tan(e)^2 - 2 A a c \tan(fx)^2 \tan(e) - 2 A a c \tan(fx) \tan(e)^2 + B a c \tan(fx)^2 + B a c \tan(e)^2}{2 (f \tan(fx)^2 \tan(e)^2 - 2 f \tan(fx) \tan(e) + f)}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x, algorithm="giac")`

output `1/2*(B*a*c*tan(f*x)^2*tan(e)^2 - 2*A*a*c*tan(f*x)^2*tan(e) - 2*A*a*c*tan(f*x)*tan(e)^2 + B*a*c*tan(f*x)^2 + B*a*c*tan(e)^2 + 2*A*a*c*tan(f*x) + 2*A*a*c*tan(e) + B*a*c)/(f*tan(f*x)^2*tan(e)^2 - 2*f*tan(f*x)*tan(e) + f)`

3.669.9 Mupad [B] (verification not implemented)

Time = 8.98 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.78

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx)) dx$$

$$= \frac{a c \tan(e + fx) (2 A + B \tan(e + fx))}{2 f}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)*(c - c*tan(e + f*x)*1i),x)`

output `(a*c*tan(e + f*x)*(2*A + B*tan(e + f*x)))/(2*f)`

3.670 $\int (a + ia \tan(e + fx))(A + B \tan(e + fx)) dx$

3.670.1 Optimal result	6334
3.670.2 Mathematica [A] (verified)	6334
3.670.3 Rubi [A] (verified)	6335
3.670.4 Maple [A] (verified)	6336
3.670.5 Fricas [A] (verification not implemented)	6336
3.670.6 Sympy [A] (verification not implemented)	6337
3.670.7 Maxima [A] (verification not implemented)	6337
3.670.8 Giac [B] (verification not implemented)	6338
3.670.9 Mupad [B] (verification not implemented)	6338

3.670.1 Optimal result

Integrand size = 24, antiderivative size = 46

$$\begin{aligned} & \int (a + ia \tan(e + fx))(A + B \tan(e + fx)) dx \\ &= a(A - iB)x - \frac{a(iA + B) \log(\cos(e + fx))}{f} + \frac{iaB \tan(e + fx)}{f} \end{aligned}$$

output `a*(A-I*B)*x-a*(I*A+B)*ln(cos(f*x+e))/f+I*a*B*tan(f*x+e)/f`

3.670.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.43

$$\begin{aligned} & \int (a + ia \tan(e + fx))(A + B \tan(e + fx)) dx \\ &= aAx - \frac{iaB \arctan(\tan(e + fx))}{f} - \frac{iaA \log(\cos(e + fx))}{f} \\ & \quad - \frac{aB \log(\cos(e + fx))}{f} + \frac{iaB \tan(e + fx)}{f} \end{aligned}$$

input `Integrate[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]),x]`

output `a*A*x - (I*a*B*ArcTan[Tan[e + f*x]])/f - (I*a*A*Log[Cos[e + f*x]])/f - (a*B*Log[Cos[e + f*x]])/f + (I*a*B*Tan[e + f*x])/f`

3.670.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3042, 4008, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))(A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))(A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4008} \\ & a(B + iA) \int \tan(e + fx) dx + ax(A - iB) + \frac{iaB \tan(e + fx)}{f} \\ & \quad \downarrow \text{3042} \\ & a(B + iA) \int \tan(e + fx) dx + ax(A - iB) + \frac{iaB \tan(e + fx)}{f} \\ & \quad \downarrow \text{3956} \\ & -\frac{a(B + iA) \log(\cos(e + fx))}{f} + ax(A - iB) + \frac{iaB \tan(e + fx)}{f} \end{aligned}$$

input `Int[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]),x]`

output `a*(A - I*B)*x - (a*(I*A + B)*Log[Cos[e + f*x]])/f + (I*a*B*Tan[e + f*x])/f`

3.670.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

```
rule 4008 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*
(x_)]), x_Symbol] :> Simp[(a*c - b*d)*x, x] + (Simp[b*d*(Tan[e + f*x]/f),
x] + Simp[(b*c + a*d) Int[Tan[e + f*x], x], x]) /; FreeQ[{a, b, c, d, e,
f}, x] && NeQ[b*c - a*d, 0] && NeQ[b*c + a*d, 0]
```

3.670.4 Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.09

method	result	size
derivativedivides	$\frac{a \left(iB \tan(fx+e) + \frac{(iA+B) \ln(1+\tan(fx+e)^2)}{2} + (-iB+A) \arctan(\tan(fx+e)) \right)}{f}$	50
default	$\frac{a \left(iB \tan(fx+e) + \frac{(iA+B) \ln(1+\tan(fx+e)^2)}{2} + (-iB+A) \arctan(\tan(fx+e)) \right)}{f}$	50
norman	$(-iaB + Aa)x + \frac{iaB \tan(fx+e)}{f} + \frac{(iAa+Ba) \ln(1+\tan(fx+e)^2)}{2f}$	52
parts	$Aax + \frac{(iAa+Ba) \ln(1+\tan(fx+e)^2)}{2f} + \frac{iBa(\tan(fx+e) - \arctan(\tan(fx+e)))}{f}$	55
parallelrisch	$\frac{-2iBxaf + ia \ln(1+\tan(fx+e)^2)a + 2Aaxf + 2iaB \tan(fx+e) + B \ln(1+\tan(fx+e)^2)a}{2f}$	61
risch	$\frac{2iaBe}{f} - \frac{2aAe}{f} - \frac{2aB}{f(e^{2i(fx+e)}+1)} - \frac{a \ln(e^{2i(fx+e)}+1)B}{f} - \frac{ia \ln(e^{2i(fx+e)}+1)A}{f}$	78

```
input int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 1/f*a*(I*B*tan(f*x+e)+1/2*(I*A+B)*ln(1+tan(f*x+e)^2)+(A-I*B)*arctan(tan(f*
x+e)))
```

3.670.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.39

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx)) dx$$

$$= -\frac{2Ba - ((-iA - B)ae^{(2i fx+2ie)} + (-iA - B)a) \log(e^{(2i fx+2ie)} + 1)}{fe^{(2i fx+2ie)} + f}$$

```
input integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e)),x, algorithm="fricas")
```

3.670. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx)) dx$

output $-(2*B*a - ((-I*A - B)*a*e^{(2*I*f*x + 2*I*e)} + (-I*A - B)*a)*\log(e^{(2*I*f*x + 2*I*e)} + 1))/(f*e^{(2*I*f*x + 2*I*e)} + f)$

3.670.6 Sympy [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.15

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx)) dx$$

$$= -\frac{2Ba}{fe^{2ie}e^{2ifx} + f} - \frac{ia(A - iB) \log(e^{2ifx} + e^{-2ie})}{f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e)),x)`

output $-2*B*a/(f*\exp(2*I*e)*\exp(2*I*f*x) + f) - I*a*(A - I*B)*\log(\exp(2*I*f*x) + \exp(-2*I*e))/f$

3.670.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.09

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx)) dx$$

$$= \frac{2(fx + e)(A - iB)a - (-iA - B)a \log(\tan(fx + e)^2 + 1) + 2iBa \tan(fx + e)}{2f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e)),x, algorithm="maxima")`

output $1/2*(2*(f*x + e)*(A - I*B)*a - (-I*A - B)*a*\log(\tan(f*x + e)^2 + 1) + 2*I*B*a*\tan(f*x + e))/f$

3.670.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 103 vs. $2(40) = 80$.

Time = 0.34 (sec) , antiderivative size = 103, normalized size of antiderivative = 2.24

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx)) dx$$

$$= \frac{-i A a e^{(2i f x + 2i e)} \log(e^{(2i f x + 2i e)} + 1) - B a e^{(2i f x + 2i e)} \log(e^{(2i f x + 2i e)} + 1) - i A a \log(e^{(2i f x + 2i e)} + 1) - B a \log(e^{(2i f x + 2i e)} + 1)}{f e^{(2i f x + 2i e)} + f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e)),x, algorithm="giac")`

output `(-I*A*a*e^(2*I*f*x + 2*I*e)*log(e^(2*I*f*x + 2*I*e) + 1) - B*a*e^(2*I*f*x + 2*I*e)*log(e^(2*I*f*x + 2*I*e) + 1) - I*A*a*log(e^(2*I*f*x + 2*I*e) + 1) - B*a*log(e^(2*I*f*x + 2*I*e) + 1) - 2*B*a)/(f*e^(2*I*f*x + 2*I*e) + f)`

3.670.9 Mupad [B] (verification not implemented)

Time = 8.88 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.83

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx)) dx$$

$$= \frac{\ln(\tan(e + fx) + 1i) (B a + A a 1i)}{f} + \frac{B a \tan(e + fx) 1i}{f}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i),x)`

output `(log(tan(e + f*x) + 1i)*(A*a*1i + B*a))/f + (B*a*tan(e + f*x)*1i)/f`

3.671 $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{c-ic \tan(e+fx)} dx$

3.671.1 Optimal result 6339
 3.671.2 Mathematica [A] (verified) 6339
 3.671.3 Rubi [A] (verified) 6340
 3.671.4 Maple [A] (verified) 6341
 3.671.5 Fricas [A] (verification not implemented) 6342
 3.671.6 Sympy [A] (verification not implemented) 6342
 3.671.7 Maxima [F(-2)] 6343
 3.671.8 Giac [B] (verification not implemented) 6343
 3.671.9 Mupad [B] (verification not implemented) 6344

3.671.1 Optimal result

Integrand size = 39, antiderivative size = 54

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx$$

$$= \frac{iaBx}{c} + \frac{aB \log(\cos(e + fx))}{cf} + \frac{a(A - iB)}{cf(i + \tan(e + fx))}$$

output `I*a*B*x/c+a*B*ln(cos(f*x+e))/c/f+a*(A-I*B)/c/f/(I+tan(f*x+e))`

3.671.2 Mathematica [A] (verified)

Time = 2.24 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.81

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx = -\frac{a \left(B \log(i + \tan(e + fx)) - \frac{A - iB}{i + \tan(e + fx)} \right)}{cf}$$

input `Integrate[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x]),x]`

output `-((a*(B*Log[I + Tan[e + f*x]] - (A - I*B)/(I + Tan[e + f*x])))/(c*f))`

3.671.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.80, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.128$, Rules used = {3042, 4071, 27, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A+B \tan(e+fx)}{c^2(1-i \tan(e+fx))^2} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a \int \frac{A+B \tan(e+fx)}{(1-i \tan(e+fx))^2} d \tan(e + fx)}{cf} \\
 & \quad \downarrow \text{49} \\
 & \frac{a \int \left(\frac{iB-A}{(\tan(e+fx)+i)^2} - \frac{B}{\tan(e+fx)+i} \right) d \tan(e + fx)}{cf} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a \left(\frac{A-iB}{\tan(e+fx)+i} - B \log(\tan(e + fx) + i) \right)}{cf}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x]),x]`

output `(a*(-(B*Log[I + Tan[e + f*x]]) + (A - I*B)/(I + Tan[e + f*x])))/(c*f)`

3.671.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.671.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.37

method	result	size
risch	$-\frac{e^{2i(fx+e)}aB}{2cf} - \frac{ie^{2i(fx+e)}Aa}{2cf} - \frac{2iBae}{cf} + \frac{Ba \ln(e^{2i(fx+e)}+1)}{cf}$	74
derivativedivides	$-\frac{iaB}{fc(i+\tan(fx+e))} + \frac{aA}{fc(i+\tan(fx+e))} - \frac{aB \ln(1+\tan(fx+e)^2)}{2fc} + \frac{iaB \arctan(\tan(fx+e))}{fc}$	83
default	$-\frac{iaB}{fc(i+\tan(fx+e))} + \frac{aA}{fc(i+\tan(fx+e))} - \frac{aB \ln(1+\tan(fx+e)^2)}{2fc} + \frac{iaB \arctan(\tan(fx+e))}{fc}$	83
norman	$\frac{(-iaB+Aa)\tan(fx+e) + \frac{iaBx}{c} + \frac{iaBx \tan(fx+e)^2}{c} - \frac{iAa+Ba}{cf} - \frac{aB \ln(1+\tan(fx+e)^2)}{2fc}}{1+\tan(fx+e)^2}$	102

input `int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x,method=_RETURNVERBOSE)`

3.671.
$$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{c-ic \tan(e+fx)} dx$$

output $-1/2/c/f*\exp(2*I*(f*x+e))*a*B-1/2*I/c/f*\exp(2*I*(f*x+e))*A*a-2*I/c/f*B*a*e+1/c/f*B*a*\ln(\exp(2*I*(f*x+e))+1)$

3.671.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.80

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{c - ict \tan(e + fx)} dx$$

$$= \frac{(-iA - B)ae^{(2i fx + 2ie)} + 2Ba \log(e^{(2i fx + 2ie)} + 1)}{2cf}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x, algorithm="fricas")`

output $1/2*((-I*A - B)*a*e^{(2*I*f*x + 2*I*e)} + 2*B*a*\log(e^{(2*I*f*x + 2*I*e)} + 1))/(c*f)$

3.671.6 Sympy [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.63

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{c - ict \tan(e + fx)} dx$$

$$= \frac{Ba \log(e^{2ifx} + e^{-2ie})}{cf} + \begin{cases} \frac{(-iAae^{2ie} - Bae^{2ie})e^{2ifx}}{2cf} & \text{for } cf \neq 0 \\ \frac{x(Aae^{2ie} - iBae^{2ie})}{c} & \text{otherwise} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x)`

output $B*a*\log(\exp(2*I*f*x) + \exp(-2*I*e))/(c*f) + \text{Piecewise}(((-I*A*a*\exp(2*I*e) - B*a*\exp(2*I*e))*\exp(2*I*f*x)/(2*c*f), \text{Ne}(c*f, 0)), (x*(A*a*\exp(2*I*e) - I*B*a*\exp(2*I*e))/c, \text{True}))$

3.671.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{c - ict \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.671.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 123 vs. $2(48) = 96$.

Time = 0.39 (sec) , antiderivative size = 123, normalized size of antiderivative = 2.28

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{c - ict \tan(e + fx)} dx$$

$$= \frac{Ba \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)}{c} - \frac{2Ba \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + i)}{c} + \frac{Ba \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)}{c} + \frac{3Ba \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 2Aa \tan(\frac{1}{2}fx + \frac{1}{2}e) + 8i}{c(\tan(\frac{1}{2}fx + \frac{1}{2}e) + i)^2}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x, algorithm="giac")`

output `(B*a*log(tan(1/2*f*x + 1/2*e) + 1)/c - 2*B*a*log(tan(1/2*f*x + 1/2*e) + I)/c + B*a*log(tan(1/2*f*x + 1/2*e) - 1)/c + (3*B*a*tan(1/2*f*x + 1/2*e)^2 - 2*A*a*tan(1/2*f*x + 1/2*e) + 8*I*B*a*tan(1/2*f*x + 1/2*e) - 3*B*a)/(c*(tan(1/2*f*x + 1/2*e) + I)^2))/f`

3.671.9 Mupad [B] (verification not implemented)

Time = 8.86 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.94

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx$$

$$= \frac{\frac{Aa}{c} - \frac{Ba1i}{c}}{f (\tan(e + fx) + 1i)} - \frac{Ba \ln(\tan(e + fx) + 1i)}{cf}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i))/(c - c*tan(e + f*x)*1i),x)`

output `((A*a)/c - (B*a*1i)/c)/(f*(tan(e + f*x) + 1i)) - (B*a*log(tan(e + f*x) + 1i))/(c*f)`

$$3.672 \quad \int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^2} dx$$

3.672.1 Optimal result	6345
3.672.2 Mathematica [A] (verified)	6345
3.672.3 Rubi [A] (verified)	6346
3.672.4 Maple [A] (verified)	6347
3.672.5 Fricas [A] (verification not implemented)	6348
3.672.6 Sympy [B] (verification not implemented)	6348
3.672.7 Maxima [F(-2)]	6349
3.672.8 Giac [A] (verification not implemented)	6349
3.672.9 Mupad [B] (verification not implemented)	6349

3.672.1 Optimal result

Integrand size = 39, antiderivative size = 46

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx = \frac{a(A + B \tan(e + fx))^2}{2(iA + B)c^2 f(1 - i \tan(e + fx))^2}$$

output `1/2*a*(A+B*tan(f*x+e))^2/(I*A+B)/c^2/f/(1-I*tan(f*x+e))^2`

3.672.2 Mathematica [A] (verified)

Time = 2.56 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.96

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx = -\frac{a(A + B \tan(e + fx))^2}{2(iA + B)c^2 f(i + \tan(e + fx))^2}$$

input `Integrate[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^2,x]`

output `-1/2*(a*(A + B*Tan[e + f*x])^2)/((I*A + B)*c^2*f*(I + Tan[e + f*x])^2)`

3.672.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.103$, Rules used = {3042, 4071, 27, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A+B \tan(e+fx)}{c^3(1-i \tan(e+fx))^3} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a \int \frac{A+B \tan(e+fx)}{(1-i \tan(e+fx))^3} d \tan(e + fx)}{c^2 f} \\
 & \quad \downarrow \text{48} \\
 & \frac{a(A + B \tan(e + fx))^2}{2c^2 f(B + iA)(1 - i \tan(e + fx))^2}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^2, x]`

output `(a*(A + B*Tan[e + f*x])^2)/(2*(I*A + B)*c^2*f*(1 - I*Tan[e + f*x])^2)`

3.672.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.672.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

method	result	size
derivativedivides	$a \left(\frac{-iA-B}{2(i+\tan(fx+e))^2} + \frac{iB}{i+\tan(fx+e)} \right) / f c^2$	46
default	$a \left(\frac{-iA-B}{2(i+\tan(fx+e))^2} + \frac{iB}{i+\tan(fx+e)} \right) / f c^2$	46
risch	$-\frac{a e^{4i(fx+e)} B}{8c^2 f} - \frac{ia e^{4i(fx+e)} A}{8c^2 f} + \frac{a e^{2i(fx+e)} B}{4c^2 f} - \frac{ia e^{2i(fx+e)} A}{4c^2 f}$	80
norman	$\frac{\frac{Aa \tan(fx+e)}{cf} + \frac{iaB \tan(fx+e)^3}{cf} - \frac{iAa+Ba}{2cf} + \frac{(iAa+3Ba) \tan(fx+e)^2}{2cf}}{c(1+\tan(fx+e))^2}$	95

input `int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `1/f*a/c^2*(-1/2*(-I*A-B)/(I+tan(f*x+e))^2+I*B/(I+tan(f*x+e)))`

3.672. $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^2} dx$

3.672.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx$$

$$= \frac{(-iA - B)ae^{(4ifx+4ie)} - 2(iA - B)ae^{(2ifx+2ie)}}{8c^2f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x, algo
rithm="fricas")`

output `1/8*((-I*A - B)*a*e^(4*I*f*x + 4*I*e) - 2*(I*A - B)*a*e^(2*I*f*x + 2*I*e))
/(c^2*f)`

3.672.6 Sympy [B] (verification not implemented)Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 153 vs. $2(36) = 72$.

Time = 0.20 (sec) , antiderivative size = 153, normalized size of antiderivative = 3.33

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx$$

$$= \begin{cases} \frac{(-8iAac^2fe^{2ie}+8Bac^2fe^{2ie})e^{2ifx}+(-4iAac^2fe^{4ie}-4Bac^2fe^{4ie})e^{4ifx}}{32c^4f^2} & \text{for } c^4f^2 \neq 0 \\ \frac{x(Aae^{4ie}+Aae^{2ie}-iBae^{4ie}+iBae^{2ie})}{2c^2} & \text{otherwise} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**2,x)`

output `Piecewise(((((-8*I*A*a*c**2*f*exp(2*I*e) + 8*B*a*c**2*f*exp(2*I*e))*exp(2*I
*f*x) + (-4*I*A*a*c**2*f*exp(4*I*e) - 4*B*a*c**2*f*exp(4*I*e))*exp(4*I*f*x
)))/(32*c**4*f**2), Ne(c**4*f**2, 0)), (x*(A*a*exp(4*I*e) + A*a*exp(2*I*e)
- I*B*a*exp(4*I*e) + I*B*a*exp(2*I*e))/(2*c**2), True))`

3.672.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x, algo
rithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.672.8 Giac [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.72

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx =$$

$$\frac{2 \left(Aa \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^3 + i Aa \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^2 - Ba \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right)^2 - Aa \tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) \right)}{c^2 f \left(\tan \left(\frac{1}{2} fx + \frac{1}{2} e \right) + i \right)^4}$$

```
input integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x, algo
rithm="giac")
```

```
output -2*(A*a*tan(1/2*f*x + 1/2*e)^3 + I*A*a*tan(1/2*f*x + 1/2*e)^2 - B*a*tan(1/
2*f*x + 1/2*e)^2 - A*a*tan(1/2*f*x + 1/2*e))/(c^2*f*(tan(1/2*f*x + 1/2*e)
+ I)^4)
```

3.672.9 Mupad [B] (verification not implemented)

Time = 9.03 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.11

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx = \frac{\frac{a(-B+Ai)}{2} + Ba \tan(e + fx) \operatorname{li}}{c^2 f (\tan(e + fx)^2 + \tan(e + fx) 2i - 1)}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i))/(c - c*tan(e + f*x)*1i)
^2,x)`

output `((a*(A*1i - B))/2 + B*a*tan(e + f*x)*1i)/(c^2*f*(tan(e + f*x)*2i + tan(e +
f*x)^2 - 1))`

3.672. $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^2} dx$

3.673 $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^3} dx$

3.673.1 Optimal result 6351
 3.673.2 Mathematica [A] (verified) 6351
 3.673.3 Rubi [A] (verified) 6352
 3.673.4 Maple [A] (verified) 6353
 3.673.5 Fricas [A] (verification not implemented) 6354
 3.673.6 Sympy [B] (verification not implemented) 6354
 3.673.7 Maxima [F(-2)] 6355
 3.673.8 Giac [B] (verification not implemented) 6355
 3.673.9 Mupad [B] (verification not implemented) 6356

3.673.1 Optimal result

Integrand size = 39, antiderivative size = 55

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx$$

$$= -\frac{a(A - iB)}{3c^3 f(i + \tan(e + fx))^3} - \frac{aB}{2c^3 f(i + \tan(e + fx))^2}$$

output `-1/3*a*(A-I*B)/c^3/f/(I+tan(f*x+e))^3-1/2*a*B/c^3/f/(I+tan(f*x+e))^2`

3.673.2 Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.75

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx = -\frac{a(2A + iB + 3B \tan(e + fx))}{6c^3 f(i + \tan(e + fx))^3}$$

input `Integrate[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^3,x]`

output `-1/6*(a*(2*A + I*B + 3*B*Tan[e + f*x]))/(c^3*f*(I + Tan[e + f*x])^3)`

3.673.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.89, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.128$, Rules used = {3042, 4071, 27, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A+B \tan(e+fx)}{c^4(1-i \tan(e+fx))^4} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a \int \frac{A+B \tan(e+fx)}{(1-i \tan(e+fx))^4} d \tan(e + fx)}{c^3 f} \\
 & \quad \downarrow \text{53} \\
 & \frac{a \int \left(\frac{A-iB}{(\tan(e+fx)+i)^4} + \frac{B}{(\tan(e+fx)+i)^3} \right) d \tan(e + fx)}{c^3 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a \left(-\frac{A-iB}{3(\tan(e+fx)+i)^3} - \frac{B}{2(\tan(e+fx)+i)^2} \right)}{c^3 f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^3, x]`

output `(a*(-1/3*(A - I*B)/(I + Tan[e + f*x])^3 - B/(2*(I + Tan[e + f*x])^2)))/(c^3*f)`

3.673.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.673.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.78

method	result	size
derivativedivides	$\frac{a\left(-\frac{B}{2(i+\tan(fx+e))^2} - \frac{-iB+A}{3(i+\tan(fx+e))^3}\right)}{fc^3}$	43
default	$\frac{a\left(-\frac{B}{2(i+\tan(fx+e))^2} - \frac{-iB+A}{3(i+\tan(fx+e))^3}\right)}{fc^3}$	43
risch	$-\frac{ae^{6i(fx+e)}B}{24c^3f} - \frac{iae^{6i(fx+e)}A}{24c^3f} - \frac{iAae^{4i(fx+e)}}{8c^3f} + \frac{ae^{2i(fx+e)}B}{8c^3f} - \frac{iae^{2i(fx+e)}A}{8c^3f}$	100

input `int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

3.673.
$$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^3} dx$$

output $1/f*a/c^3*(-1/2*B/(I+\tan(f*x+e))^2-1/3*(A-I*B)/(I+\tan(f*x+e))^3)$

3.673.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.07

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^3} dx$$

$$= \frac{(-iA - B)ae^{(6i fx + 6ie)} - 3iAae^{(4i fx + 4ie)} - 3(iA - B)ae^{(2i fx + 2ie)}}{24c^3f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x, algo rithm="fricas")`

output $1/24*((-I*A - B)*a*e^{(6*I*f*x + 6*I*e)} - 3*I*A*a*e^{(4*I*f*x + 4*I*e)} - 3*(I*A - B)*a*e^{(2*I*f*x + 2*I*e)})/(c^3*f)$

3.673.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 201 vs. $2(44) = 88$.

Time = 0.25 (sec) , antiderivative size = 201, normalized size of antiderivative = 3.65

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^3} dx$$

$$= \begin{cases} \frac{-192iAac^6 f^2 e^{4ie} e^{4ifx} + (-192iAac^6 f^2 e^{2ie} + 192Bac^6 f^2 e^{2ie}) e^{2ifx} + (-64iAac^6 f^2 e^{6ie} - 64Bac^6 f^2 e^{6ie}) e^{6ifx}}{1536c^9 f^3} & \text{for } c^9 f^3 \neq 0 \\ \frac{x(Aae^{6ie} + 2Aae^{4ie} + Aae^{2ie} - iBae^{6ie} + iBae^{2ie})}{4c^3} & \text{otherwise} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**3,x)`

output `Piecewise(((-192*I*A*a*c**6*f**2*exp(4*I*e)*exp(4*I*f*x) + (-192*I*A*a*c**6*f**2*exp(2*I*e) + 192*B*a*c**6*f**2*exp(2*I*e))*exp(2*I*f*x) + (-64*I*A*a*c**6*f**2*exp(6*I*e) - 64*B*a*c**6*f**2*exp(6*I*e))*exp(6*I*f*x))/(1536*c**9*f**3), Ne(c**9*f**3, 0)), (x*(A*a*exp(6*I*e) + 2*A*a*exp(4*I*e) + A*a*exp(2*I*e) - I*B*a*exp(6*I*e) + I*B*a*exp(2*I*e))/(4*c**3), True))`

3.673. $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^3} dx$

3.673.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x, algo
rithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.`

3.673.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 140 vs. $2(45) = 90$.

Time = 0.64 (sec) , antiderivative size = 140, normalized size of antiderivative = 2.55

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx =$$

$$\frac{2 \left(3 A a \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^5 + 6 i A a \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^4 - 3 B a \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^4 - 10 A a \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^3 \right)}{3 c^3 f \left(\tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) + i \right)^6}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x, algo
rithm="giac")`

output `-2/3*(3*A*a*tan(1/2*f*x + 1/2*e)^5 + 6*I*A*a*tan(1/2*f*x + 1/2*e)^4 - 3*B*
a*tan(1/2*f*x + 1/2*e)^4 - 10*A*a*tan(1/2*f*x + 1/2*e)^3 - 2*I*B*a*tan(1/2
*f*x + 1/2*e)^3 - 6*I*A*a*tan(1/2*f*x + 1/2*e)^2 + 3*B*a*tan(1/2*f*x + 1/2
*e)^2 + 3*A*a*tan(1/2*f*x + 1/2*e))/(c^3*f*(tan(1/2*f*x + 1/2*e) + I)^6)`

3.673.9 Mupad [B] (verification not implemented)

Time = 8.92 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.15

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx$$

$$= \frac{\frac{a(2A+B1i)}{6} + \frac{Ba \tan(e+fx)}{2}}{c^3 f (-\tan(e + fx)^3 - \tan(e + fx)^2 3i + 3 \tan(e + fx) + 1i)}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i))/(c - c*tan(e + f*x)*1i)
^3,x)`

output `((a*(2*A + B*1i))/6 + (B*a*tan(e + f*x))/2)/(c^3*f*(3*tan(e + f*x) - tan(e
+ f*x)^2*3i - tan(e + f*x)^3 + 1i))`

3.674 $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^4} dx$

3.674.1 Optimal result 6357
 3.674.2 Mathematica [A] (verified) 6357
 3.674.3 Rubi [A] (verified) 6358
 3.674.4 Maple [A] (verified) 6359
 3.674.5 Fricas [A] (verification not implemented) 6360
 3.674.6 Sympy [B] (verification not implemented) 6360
 3.674.7 Maxima [F(-2)] 6361
 3.674.8 Giac [B] (verification not implemented) 6361
 3.674.9 Mupad [B] (verification not implemented) 6362

3.674.1 Optimal result

Integrand size = 39, antiderivative size = 57

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx$$

$$= -\frac{a(iA + B)}{4c^4 f(i + \tan(e + fx))^4} - \frac{iaB}{3c^4 f(i + \tan(e + fx))^3}$$

output `-1/4*a*(I*A+B)/c^4/f/(I+tan(f*x+e))^4-1/3*I*a*B/c^4/f/(I+tan(f*x+e))^3`

3.674.2 Mathematica [A] (verified)

Time = 0.53 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.72

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx = \frac{a(-3iA + B - 4iB \tan(e + fx))}{12c^4 f(i + \tan(e + fx))^4}$$

input `Integrate[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^4,x]`

output `(a*((-3*I)*A + B - (4*I)*B*Tan[e + f*x]))/(12*c^4*f*(I + Tan[e + f*x])^4)`

3.674.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.89, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.128$, Rules used = {3042, 4071, 27, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A+B \tan(e+fx)}{c^5(1-i \tan(e+fx))^5} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a \int \frac{A+B \tan(e+fx)}{(1-i \tan(e+fx))^5} d \tan(e + fx)}{c^4 f} \\
 & \quad \downarrow \text{53} \\
 & \frac{a \int \left(\frac{iB}{(\tan(e+fx)+i)^4} + \frac{iA+B}{(\tan(e+fx)+i)^5} \right) d \tan(e + fx)}{c^4 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a \left(-\frac{B+iA}{4(\tan(e+fx)+i)^4} - \frac{iB}{3(\tan(e+fx)+i)^3} \right)}{c^4 f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^4, x]`

output `(a*(-1/4*(I*A + B)/(I + Tan[e + f*x])^4 - ((I/3)*B)/(I + Tan[e + f*x])^3)/(c^4*f)`

3.674.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx) /; FreeQ[b, x]]`

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.674.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.77

method	result
derivativedivides	$\frac{a \left(-\frac{iB}{3(i+\tan(fx+e))^3} - \frac{iA+B}{4(i+\tan(fx+e))^4} \right)}{f c^4}$
default	$\frac{a \left(-\frac{iB}{3(i+\tan(fx+e))^3} - \frac{iA+B}{4(i+\tan(fx+e))^4} \right)}{f c^4}$
risch	$-\frac{a e^{8i(fx+e)} B}{64c^4 f} - \frac{ia e^{8i(fx+e)} A}{64c^4 f} - \frac{e^{6i(fx+e)} aB}{48c^4 f} - \frac{ie^{6i(fx+e)} Aa}{16c^4 f} + \frac{e^{4i(fx+e)} aB}{32c^4 f} - \frac{3ie^{4i(fx+e)} Aa}{32c^4 f} + \frac{a e^{2i(fx+e)}}{16c^4 f}$

input `int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x,method=_RET URNVERBOSE)`

$$3.674. \quad \int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^4} dx$$

output $1/f*a/c^4*(-1/3*I*B/(I+\tan(f*x+e))^3-1/4*(I*A+B)/(I+\tan(f*x+e))^4)$

3.674.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.42

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ictan(e + fx))^4} dx = \frac{3(iA + B)ae^{(8ifx+8ie)} + 4(3iA + B)ae^{(6ifx+6ie)} + 6(3iA - B)ae^{(4ifx+4ie)} + 12(iA - B)ae^{(2ifx+2ie)}}{192c^4f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x, algo rithm="fricas")`

output $-1/192*(3*(I*A + B)*a*e^{(8*I*f*x + 8*I*e)} + 4*(3*I*A + B)*a*e^{(6*I*f*x + 6*I*e)} + 6*(3*I*A - B)*a*e^{(4*I*f*x + 4*I*e)} + 12*(I*A - B)*a*e^{(2*I*f*x + 2*I*e)})/(c^4*f)$

3.674.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 304 vs. $2(46) = 92$.

Time = 0.31 (sec) , antiderivative size = 304, normalized size of antiderivative = 5.33

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ictan(e + fx))^4} dx = \begin{cases} \frac{(-98304iAac^{12}f^3e^{2ie}+98304Bac^{12}f^3e^{2ie})e^{2ifx}+(-147456iAac^{12}f^3e^{4ie}+49152Bac^{12}f^3e^{4ie})e^{4ifx}+(-98304iAac^{12}f^3e^{6ie}-32768Bac^{12}f^3e^{6ie})e^{6ifx}+(-98304iAac^{12}f^3e^{8ie}-32768Bac^{12}f^3e^{8ie})e^{8ifx}}{1572864c^{16}f^4} \\ \frac{x(Aae^{8ie}+3Aae^{6ie}+3Aae^{4ie}+Aae^{2ie}-iBae^{8ie}-iBae^{6ie}+iBae^{4ie}+iBae^{2ie})}{8c^4} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**4,x)`

output `Piecewise(((((-98304*I*A*a*c**12*f**3*exp(2*I*e) + 98304*B*a*c**12*f**3*exp(2*I*e))*exp(2*I*f*x) + (-147456*I*A*a*c**12*f**3*exp(4*I*e) + 49152*B*a*c**12*f**3*exp(4*I*e))*exp(4*I*f*x) + (-98304*I*A*a*c**12*f**3*exp(6*I*e) - 32768*B*a*c**12*f**3*exp(6*I*e))*exp(6*I*f*x) + (-24576*I*A*a*c**12*f**3*exp(8*I*e) - 24576*B*a*c**12*f**3*exp(8*I*e))*exp(8*I*f*x))/(1572864*c**16*f**4), Ne(c**16*f**4, 0)), (x*(A*a*exp(8*I*e) + 3*A*a*exp(6*I*e) + 3*A*a*exp(4*I*e) + A*a*exp(2*I*e) - I*B*a*exp(8*I*e) - I*B*a*exp(6*I*e) + I*B*a*exp(4*I*e) + I*B*a*exp(2*I*e))/(8*c**4), True))`

3.674.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x, algo rithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.674.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 200 vs. $2(45) = 90$.

Time = 0.69 (sec) , antiderivative size = 200, normalized size of antiderivative = 3.51

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx = \frac{2 \left(3Aa \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^7 + 9iAa \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^6 - 3Ba \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^6 - 21Aa \tan\left(\frac{1}{2}fx + \frac{1}{2}e\right)^5 \right)}{\dots}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x, algo rithm="giac")`

3.674. $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^4} dx$

output
$$\frac{-2/3*(3*A*a*\tan(1/2*f*x + 1/2*e)^7 + 9*I*A*a*\tan(1/2*f*x + 1/2*e)^6 - 3*B*a*\tan(1/2*f*x + 1/2*e)^6 - 21*A*a*\tan(1/2*f*x + 1/2*e)^5 - 4*I*B*a*\tan(1/2*f*x + 1/2*e)^5 - 24*I*A*a*\tan(1/2*f*x + 1/2*e)^4 + 8*B*a*\tan(1/2*f*x + 1/2*e)^4 + 21*A*a*\tan(1/2*f*x + 1/2*e)^3 + 4*I*B*a*\tan(1/2*f*x + 1/2*e)^3 + 9*I*A*a*\tan(1/2*f*x + 1/2*e)^2 - 3*B*a*\tan(1/2*f*x + 1/2*e)^2 - 3*A*a*\tan(1/2*f*x + 1/2*e))/(c^4*f*(\tan(1/2*f*x + 1/2*e) + I)^8)}$$

3.674.9 Mupad [B] (verification not implemented)

Time = 8.59 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.28

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^4} dx$$

$$= \frac{\frac{a(-B+Ai)}{12} + \frac{B a \tan(e+fx) i}{3}}{c^4 f (\tan(e + fx)^4 + \tan(e + fx)^3 4i - 6 \tan(e + fx)^2 - \tan(e + fx) 4i + 1)}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i))/(c - c*tan(e + f*x)*1i)^4,x)`

output
$$-\left(\frac{a(A*3i - B)}{12} + \frac{B*a*\tan(e + f*x)*1i}{3}\right)/(c^4*f*(\tan(e + f*x)^3*4i - 6*\tan(e + f*x)^2 - \tan(e + f*x)*4i + \tan(e + f*x)^4 + 1))$$

3.675 $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^5} dx$

3.675.1 Optimal result 6363
 3.675.2 Mathematica [A] (verified) 6363
 3.675.3 Rubi [A] (verified) 6364
 3.675.4 Maple [A] (verified) 6365
 3.675.5 Fricas [B] (verification not implemented) 6366
 3.675.6 Sympy [B] (verification not implemented) 6366
 3.675.7 Maxima [F(-2)] 6367
 3.675.8 Giac [B] (verification not implemented) 6367
 3.675.9 Mupad [B] (verification not implemented) 6368

3.675.1 Optimal result

Integrand size = 39, antiderivative size = 55

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx$$

$$= \frac{a(A - iB)}{5c^5 f(i + \tan(e + fx))^5} + \frac{aB}{4c^5 f(i + \tan(e + fx))^4}$$

output `1/5*a*(A-I*B)/c^5/f/(I+tan(f*x+e))^5+1/4*a*B/c^5/f/(I+tan(f*x+e))^4`

3.675.2 Mathematica [A] (verified)

Time = 1.68 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.75

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx = \frac{a(4A + iB + 5B \tan(e + fx))}{20c^5 f(i + \tan(e + fx))^5}$$

input `Integrate[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^5,x]`

output `(a*(4*A + I*B + 5*B*Tan[e + f*x]))/(20*c^5*f*(I + Tan[e + f*x])^5)`

3.675.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.89, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.128$, Rules used = {3042, 4071, 27, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A+B \tan(e+fx)}{c^6(1-i \tan(e+fx))^6} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a \int \frac{A+B \tan(e+fx)}{(1-i \tan(e+fx))^6} d \tan(e + fx)}{c^5 f} \\
 & \quad \downarrow \text{53} \\
 & \frac{a \int \left(\frac{iB-A}{(\tan(e+fx)+i)^6} - \frac{B}{(\tan(e+fx)+i)^5} \right) d \tan(e + fx)}{c^5 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a \left(\frac{A-iB}{5(\tan(e+fx)+i)^5} + \frac{B}{4(\tan(e+fx)+i)^4} \right)}{c^5 f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^5, x]`

output `(a*((A - I*B)/(5*(I + Tan[e + f*x])^5) + B/(4*(I + Tan[e + f*x])^4)))/(c^5*f)`

3.675.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.675.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.82

method	result
derivativedivides	$\frac{a\left(-\frac{iB-A}{5(i+\tan(fx+e))^5} + \frac{B}{4(i+\tan(fx+e))^4}\right)}{f c^5}$
default	$\frac{a\left(-\frac{iB-A}{5(i+\tan(fx+e))^5} + \frac{B}{4(i+\tan(fx+e))^4}\right)}{f c^5}$
risch	$-\frac{ae^{10i(fx+e)}B}{160c^5f} - \frac{iae^{10i(fx+e)}A}{160c^5f} - \frac{e^{8i(fx+e)}aB}{64c^5f} - \frac{ie^{8i(fx+e)}Aa}{32c^5f} - \frac{iAae^{6i(fx+e)}}{16c^5f} + \frac{e^{4i(fx+e)}aB}{32c^5f} - \frac{ie^{4i(fx+e)}Aa}{16c^5f}$

input `int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^5,x,method=_RET URNVERBOSE)`

3.675.
$$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^5} dx$$

output $1/f*a/c^5*(-1/5*(-A+I*B)/(I+\tan(f*x+e))^5+1/4*B/(I+\tan(f*x+e))^4)$

3.675.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 94 vs. $2(45) = 90$.

Time = 0.24 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.71

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ictan(e + fx))^5} dx = \frac{2(iA + B)ae^{(10i fx + 10ie)} + 5(2iA + B)ae^{(8i fx + 8ie)} + 20iAae^{(6i fx + 6ie)} + 10(2iA - B)ae^{(4i fx + 4ie)} + 10(I*A - B)*a*e^{(2*I*f*x + 2*I*e)}}{320 c^5 f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^5,x, algorithm="fricas")`

output $-1/320*(2*(I*A + B)*a*e^{(10*I*f*x + 10*I*e)} + 5*(2*I*A + B)*a*e^{(8*I*f*x + 8*I*e)} + 20*I*A*a*e^{(6*I*f*x + 6*I*e)} + 10*(2*I*A - B)*a*e^{(4*I*f*x + 4*I*e)} + 10*(I*A - B)*a*e^{(2*I*f*x + 2*I*e)})/(c^5*f)$

3.675.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 348 vs. $2(42) = 84$.

Time = 0.39 (sec) , antiderivative size = 348, normalized size of antiderivative = 6.33

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ictan(e + fx))^5} dx = \left\{ \frac{-10485760iAac^{20}f^4e^{6ie}e^{6ifx} + (-5242880iAac^{20}f^4e^{2ie} + 5242880Bac^{20}f^4e^{2ie})e^{2ifx} + (-10485760iAac^{20}f^4e^{4ie} + 5242880Bac^{20}f^4e^{4ie})e^{4ifx}}{167772160c^{25}f^5}, \frac{x(Aae^{10ie} + 4Aae^{8ie} + 6Aae^{6ie} + 4Aae^{4ie} + Aae^{2ie} - iBae^{10ie} - 2iBae^{8ie} + 2iBae^{4ie} + iBae^{2ie})}{16c^5} \right\}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**5,x)`

output `Piecewise(((−10485760*I*A*a*c**20*f**4*exp(6*I*e)*exp(6*I*f*x) + (−5242880*I*A*a*c**20*f**4*exp(2*I*e) + 5242880*B*a*c**20*f**4*exp(2*I*e))*exp(2*I*f*x) + (−10485760*I*A*a*c**20*f**4*exp(4*I*e) + 5242880*B*a*c**20*f**4*exp(4*I*e))*exp(4*I*f*x) + (−5242880*I*A*a*c**20*f**4*exp(8*I*e) − 2621440*B*a*c**20*f**4*exp(8*I*e))*exp(8*I*f*x) + (−1048576*I*A*a*c**20*f**4*exp(10*I*e) − 1048576*B*a*c**20*f**4*exp(10*I*e))*exp(10*I*f*x))/(167772160*c**25*f**5), Ne(c**25*f**5, 0)), (x*(A*a*exp(10*I*e) + 4*A*a*exp(8*I*e) + 6*A*a*exp(6*I*e) + 4*A*a*exp(4*I*e) + A*a*exp(2*I*e) − I*B*a*exp(10*I*e) − 2*I*B*a*exp(8*I*e) + 2*I*B*a*exp(4*I*e) + I*B*a*exp(2*I*e))/(16*c**5), True))`

3.675.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^5,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.675.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 260 vs. $2(45) = 90$.

Time = 0.97 (sec) , antiderivative size = 260, normalized size of antiderivative = 4.73

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx =$$

$$\frac{2 \left(5 A a \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^9 + 20 i A a \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^8 - 5 B a \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^8 - 60 A a \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) \right)}{16 c^5}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^5,x, algorithm="giac")`

3.675. $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^5} dx$

output
$$\frac{-2/5*(5*A*a*\tan(1/2*f*x + 1/2*e)^9 + 20*I*A*a*\tan(1/2*f*x + 1/2*e)^8 - 5*B*a*\tan(1/2*f*x + 1/2*e)^8 - 60*A*a*\tan(1/2*f*x + 1/2*e)^7 - 10*I*B*a*\tan(1/2*f*x + 1/2*e)^7 - 100*I*A*a*\tan(1/2*f*x + 1/2*e)^6 + 25*B*a*\tan(1/2*f*x + 1/2*e)^6 + 126*A*a*\tan(1/2*f*x + 1/2*e)^5 + 24*I*B*a*\tan(1/2*f*x + 1/2*e)^5 + 100*I*A*a*\tan(1/2*f*x + 1/2*e)^4 - 25*B*a*\tan(1/2*f*x + 1/2*e)^4 - 60*A*a*\tan(1/2*f*x + 1/2*e)^3 - 10*I*B*a*\tan(1/2*f*x + 1/2*e)^3 - 20*I*A*a*\tan(1/2*f*x + 1/2*e)^2 + 5*B*a*\tan(1/2*f*x + 1/2*e)^2 + 5*A*a*\tan(1/2*f*x + 1/2*e))/(c^5*f*(\tan(1/2*f*x + 1/2*e) + I)^{10}}$$

3.675.9 Mupad [B] (verification not implemented)

Time = 8.66 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.49

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ictan(e + fx))^5} dx$$

$$= \frac{\frac{a(4A+B1i)}{20} + \frac{Batan(e+fx)}{4}}{c^5 f (\tan(e + fx)^5 + \tan(e + fx)^4 5i - 10 \tan(e + fx)^3 - \tan(e + fx)^2 10i + 5 \tan(e + fx) + 1i)}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i))/(c - c*tan(e + f*x)*1i)^5,x)`

output
$$((a*(4*A + B*1i))/20 + (B*a*\tan(e + f*x))/4)/(c^5*f*(5*\tan(e + f*x) - \tan(e + f*x)^2*10i - 10*\tan(e + f*x)^3 + \tan(e + f*x)^4*5i + \tan(e + f*x)^5 + 1i))$$

3.676 $\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ictan(e+fx))^n dx$

3.676.1 Optimal result	6369
3.676.2 Mathematica [A] (verified)	6369
3.676.3 Rubi [A] (verified)	6370
3.676.4 Maple [A] (verified)	6372
3.676.5 Fricas [B] (verification not implemented)	6372
3.676.6 Sympy [B] (verification not implemented)	6373
3.676.7 Maxima [B] (verification not implemented)	6373
3.676.8 Giac [F]	6374
3.676.9 Mupad [B] (verification not implemented)	6375

3.676.1 Optimal result

Integrand size = 41, antiderivative size = 109

$$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ictan(e+fx))^n dx = \frac{2a^2(iA+B)(c-ictan(e+fx))^n}{fn} - \frac{a^2(iA+3B)(c-ictan(e+fx))^{1+n}}{cf(1+n)} + \frac{a^2B(c-ictan(e+fx))^{2+n}}{c^2f(2+n)}$$

```
output 2*a^2*(I*A+B)*(c-I*c*tan(f*x+e))^n/f/n-a^2*(I*A+3*B)*(c-I*c*tan(f*x+e))^(1+n)/c/f/(1+n)+a^2*B*(c-I*c*tan(f*x+e))^(2+n)/c^2/f/(2+n)
```

3.676.2 Mathematica [A] (verified)

Time = 2.18 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.82

$$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ictan(e+fx))^n dx = \frac{a^2(c-ictan(e+fx))^n(-iA(2+n)^2-B(4+n)+n(A(2+n)-iB(4+n))\tan(e+fx)+Bn(1+n))}{fn(1+n)(2+n)}$$

```
input Integrate[(a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^n,x]
```


output $-\left(a^2(c - I*c*\text{Tan}[e + f*x])^n*((-I)*A*(2 + n)^2 - B*(4 + n) + n*(A*(2 + n) - I*B*(4 + n))*\text{Tan}[e + f*x] + B*n*(1 + n)*\text{Tan}[e + f*x]^2)/(f*n*(1 + n)*(2 + n))\right)$

3.676.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.94, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^n dx$$

↓ 3042

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^n dx$$

↓ 4071

$$\frac{ac \int a(i \tan(e + fx) + 1)(A + B \tan(e + fx))(c - ictan(e + fx))^{n-1} d \tan(e + fx)}{f}$$

↓ 27

$$\frac{a^2c \int (i \tan(e + fx) + 1)(A + B \tan(e + fx))(c - ictan(e + fx))^{n-1} d \tan(e + fx)}{f}$$

↓ 86

$$\frac{a^2c \int \left(2(A - iB)(c - ictan(e + fx))^{n-1} + \frac{(3iB - A)(c - ictan(e + fx))^n}{c} - \frac{iB(c - ictan(e + fx))^{n+1}}{c^2}\right) d \tan(e + fx)}{f}$$

↓ 2009

$$\frac{a^2c \left(-\frac{(3B + iA)(c - ictan(e + fx))^{n+1}}{c^2(n+1)} + \frac{2(B + iA)(c - ictan(e + fx))^n}{cn} + \frac{B(c - ictan(e + fx))^{n+2}}{c^3(n+2)}\right)}{f}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^2*(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^n, x]$

3.676. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^n dx$

```
output (a^2*c*((2*(I*A + B)*(c - I*c*Tan[e + f*x])^n)/(c*n) - ((I*A + 3*B)*(c - I
*c*Tan[e + f*x])^(1 + n))/(c^2*(1 + n)) + (B*(c - I*c*Tan[e + f*x])^(2 + n
))/(c^3*(2 + n))))/f
```

3.676.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.676.4 Maple [A] (verified)

Time = 1.04 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.53

method	result
derivativedivides	$\frac{(iA a^2 n^2 + 4iA a^2 n + 4iA a^2 + B a^2 n + 4B a^2) e^{n \ln(c - ic \tan(fx + e))}}{(1+n)fn(2+n)} - \frac{B a^2 \tan(fx + e)^2 e^{n \ln(c - ic \tan(fx + e))}}{f(2+n)} - \frac{a^2(-iB)}{f(2+n)}$
default	$\frac{(iA a^2 n^2 + 4iA a^2 n + 4iA a^2 + B a^2 n + 4B a^2) e^{n \ln(c - ic \tan(fx + e))}}{(1+n)fn(2+n)} - \frac{B a^2 \tan(fx + e)^2 e^{n \ln(c - ic \tan(fx + e))}}{f(2+n)} - \frac{a^2(-iB)}{f(2+n)}$
norman	$\frac{(iA a^2 n^2 + 4iA a^2 n + 4iA a^2 + B a^2 n + 4B a^2) e^{n \ln(c - ic \tan(fx + e))}}{(1+n)fn(2+n)} - \frac{B a^2 \tan(fx + e)^2 e^{n \ln(c - ic \tan(fx + e))}}{f(2+n)} - \frac{a^2(-iB)}{f(2+n)}$
risch	Expression too large to display

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x,method=_RETURVERBOSE)`

output `1/(1+n)/f/n*(I*A*a^2*n^2+4*I*A*a^2*n+4*I*A*a^2+B*a^2*n+4*B*a^2)/(2+n)*exp(n*ln(c-I*c*tan(f*x+e)))-B*a^2/f/(2+n)*tan(f*x+e)^2*exp(n*ln(c-I*c*tan(f*x+e)))-a^2*(-I*B*n+A*n-4*I*B+2*A)/f/(1+n)/(2+n)*tan(f*x+e)*exp(n*ln(c-I*c*tan(f*x+e)))`

3.676.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 206 vs. 2(99) = 198.

Time = 0.26 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.89

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx = \frac{2((-iA + B)a^2n + 2(-iA - B)a^2 + ((-iA - B)a^2n^2 + 3(-iA - B)a^2n + 2(-iA - B)a^2)e^{4i fx + 4ie}}{fn^3 + 3fn^2 + 2fn + (fn^3 + 3fn^2 + 2fn)e^{4i fx + 4ie}} + \dots$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x, algorithm="fricas")`

output `-2*((-I*A + B)*a^2*n + 2*(-I*A - B)*a^2 + ((-I*A - B)*a^2*n^2 + 3*(-I*A - B)*a^2*n + 2*(-I*A - B)*a^2)*e^(4*I*f*x + 4*I*e) + ((-I*A + B)*a^2*n^2 - 4*I*A*a^2*n + 4*(-I*A - B)*a^2)*e^(2*I*f*x + 2*I*e)*(2*c/(e^(2*I*f*x + 2*I*e) + 1))^n/(f*n^3 + 3*f*n^2 + 2*f*n + (f*n^3 + 3*f*n^2 + 2*f*n)*e^(4*I*f*x + 4*I*e) + 2*(f*n^3 + 3*f*n^2 + 2*f*n)*e^(2*I*f*x + 2*I*e))`

3.676. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx$

3.676.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1482 vs. $2(87) = 174$.

Time = 1.25 (sec) , antiderivative size = 1482, normalized size of antiderivative = 13.60

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^n dx = \text{Too large to display}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**n,x)`

output `Piecewise((x*(A + B*tan(e))*(I*a*tan(e) + a)**2*(-I*c*tan(e) + c)**n, Eq(f, 0)), (-2*A*a**2*tan(e + f*x)/(2*c**2*f*tan(e + f*x)**2 + 4*I*c**2*f*tan(e + f*x) - 2*c**2*f) - 2*I*B*a**2*f*x*tan(e + f*x)**2/(2*c**2*f*tan(e + f*x)**2 + 4*I*c**2*f*tan(e + f*x) - 2*c**2*f) + 4*B*a**2*f*x*tan(e + f*x)/(2*c**2*f*tan(e + f*x)**2 + 4*I*c**2*f*tan(e + f*x) - 2*c**2*f) + 2*I*B*a**2*f*x/(2*c**2*f*tan(e + f*x)**2 + 4*I*c**2*f*tan(e + f*x) - 2*c**2*f) + B*a**2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)**2/(2*c**2*f*tan(e + f*x)**2 + 4*I*c**2*f*tan(e + f*x) - 2*c**2*f) + 2*I*B*a**2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)/(2*c**2*f*tan(e + f*x)**2 + 4*I*c**2*f*tan(e + f*x) - 2*c**2*f) - B*a**2*log(tan(e + f*x)**2 + 1)/(2*c**2*f*tan(e + f*x)**2 + 4*I*c**2*f*tan(e + f*x) - 2*c**2*f) + 6*I*B*a**2*tan(e + f*x)/(2*c**2*f*tan(e + f*x)**2 + 4*I*c**2*f*tan(e + f*x) - 2*c**2*f) - 4*B*a**2/(2*c**2*f*tan(e + f*x)**2 + 4*I*c**2*f*tan(e + f*x) - 2*c**2*f), Eq(n, -2)), (-2*A*a**2*f*x*tan(e + f*x)/(2*c*f*tan(e + f*x) + 2*I*c*f) - 2*I*A*a**2*f*x/(2*c*f*tan(e + f*x) + 2*I*c*f) - I*A*a**2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)/(2*c*f*tan(e + f*x) + 2*I*c*f) + A*a**2*log(tan(e + f*x)**2 + 1)/(2*c*f*tan(e + f*x) + 2*I*c*f) + 4*A*a**2/(2*c*f*tan(e + f*x) + 2*I*c*f) + 6*I*B*a**2*f*x*tan(e + f*x)/(2*c*f*tan(e + f*x) + 2*I*c*f) - 6*B*a**2*f*x/(2*c*f*tan(e + f*x) + 2*I*c*f) - 3*B*a**2*log(tan(e + f*x)**2 + 1)*tan(e + f*x)/(2*c*f*tan(e + f*x) + 2*I*c*f) - 3*I*B*a**2*log(tan(e + f*x)**2 + 1)/(2*c*f*tan(e + ...`

3.676.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 660 vs. $2(99) = 198$.

$$3.676. \quad \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^n dx$$

Time = 0.45 (sec) , antiderivative size = 660, normalized size of antiderivative = 6.06

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^n dx$$

$$= \frac{2(((A + iB)a^2c^n n^2 + 4Aa^2c^n n + 4(A - iB)a^2c^n)2^n \cos(-2fx + n \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + ((A - iB)a^2c^n n^2 + 3(A - iB)a^2c^n n + 2(A - iB)a^2c^n)2^n \cos(-4fx + n \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + ((A + iB)a^2c^n n + 2(A - iB)a^2c^n)2^n \cos(n \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 1) - ((I*A - B)a^2c^n n^2 + 4*I*A*a^2c^n n + 4*(I*A + B)a^2c^n)2^n \sin(-2fx + n \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 1) - 2e) - ((I*A + B)a^2c^n n^2 + 3*(I*A + B)a^2c^n n + 2*(I*A + B)a^2c^n)2^n \sin(-4fx + n \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 1) - 4e) - ((I*A - B)a^2c^n n + 2*(I*A + B)a^2c^n)2^n \sin(n \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 1)))/((-I*n^3 - 3*I*n^2 - 2*I*n)*(cos(2fx + 2e)^2 + sin(2fx + 2e)^2 + 2*cos(2fx + 2e) + 1)^(1/2*n)*cos(4fx + 4e) + (n^3 + 3*n^2 + 2*n)*(cos(2fx + 2e)^2 + sin(2fx + 2e)^2 + 2*cos(2fx + 2e) + 1)^(1/2*n)*sin(4fx + 4e) + (-I*n^3 - 3*I*n^2 - 2*(I*n^3 + 3*I*n^2 + 2*I*n)*cos(2fx + 2e) + 2*(n^3 + 3*n^2 + 2*n)*sin(2fx + 2e) - 2*I*n)*(cos(2fx + 2e)^2 + sin(2fx + 2e)^2 + 2*cos(2fx + 2e) + 1)^(1/2*n))*f)$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x, algorithm="maxima")`

output `2*(((A + I*B)*a^2*c^n*n^2 + 4*A*a^2*c^n*n + 4*(A - I*B)*a^2*c^n)*2^n*cos(-2*f*x + n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - 2*e) + ((A - I*B)*a^2*c^n*n^2 + 3*(A - I*B)*a^2*c^n*n + 2*(A - I*B)*a^2*c^n)*2^n*cos(-4*f*x + n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - 4*e) + ((A + I*B)*a^2*c^n*n + 2*(A - I*B)*a^2*c^n)*2^n*cos(n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)) - ((I*A - B)*a^2*c^n*n^2 + 4*I*A*a^2*c^n*n + 4*(I*A + B)*a^2*c^n)*2^n*sin(-2*f*x + n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - 2*e) - ((I*A + B)*a^2*c^n*n^2 + 3*(I*A + B)*a^2*c^n*n + 2*(I*A + B)*a^2*c^n)*2^n*sin(-4*f*x + n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - 4*e) - ((I*A - B)*a^2*c^n*n + 2*(I*A + B)*a^2*c^n)*2^n*sin(n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)))/((-I*n^3 - 3*I*n^2 - 2*I*n)*(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^(1/2*n)*cos(4*f*x + 4*e) + (n^3 + 3*n^2 + 2*n)*(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^(1/2*n)*sin(4*f*x + 4*e) + (-I*n^3 - 3*I*n^2 - 2*(I*n^3 + 3*I*n^2 + 2*I*n)*cos(2*f*x + 2*e) + 2*(n^3 + 3*n^2 + 2*n)*sin(2*f*x + 2*e) - 2*I*n)*(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^(1/2*n))*f)`

3.676.8 Giac [F]

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^n dx$$

$$= \int (B \tan(fx + e) + A)(ia \tan(fx + e) + a)^2 (-ictan(fx + e) + c)^n dx$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^2*(-I*c*tan(f*x + e) + c)^n, x)`

3.676.9 Mupad [B] (verification not implemented)

Time = 11.54 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.77

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^n dx = \frac{e^{-e2i - fx2i} \left(c - \frac{c \sin(e+fx) 1i}{\cos(e+fx)} \right)^n \left(\frac{2a^2 (2A - B2i + An + Bn 1i)}{fn (n^2 1i + n 3i + 2i)} + \frac{2a^2 e^{e4i + fx4i} (A - B 1i) (n^2 + 3n + 2)}{fn (n^2 1i + n 3i + 2i)} + \frac{2a^2 e^{e2i + fx2i} (n+2)}{fn (n^2 1i + n 3i + 2i)} \right)}{4 \cos(e + fx)^2}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)^n,x)`

output `-(exp(- e*2i - f*x*2i)*(c - (c*sin(e + f*x)*1i)/cos(e + f*x))^n*((2*a^2*(2*A - B*2i + A*n + B*n*1i))/(f*n*(n*3i + n^2*1i + 2i)) + (2*a^2*exp(e*4i + f*x*4i)*(A - B*1i)*(3*n + n^2 + 2))/(f*n*(n*3i + n^2*1i + 2i)) + (2*a^2*exp(e*2i + f*x*2i)*(n + 2)*(2*A - B*2i + A*n + B*n*1i))/(f*n*(n*3i + n^2*1i + 2i))))/(4*cos(e + f*x)^2)`

3.677
$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^5 dx$$

3.677.1 Optimal result 6376
 3.677.2 Mathematica [B] (verified) 6376
 3.677.3 Rubi [A] (verified) 6377
 3.677.4 Maple [A] (verified) 6379
 3.677.5 Fricas [A] (verification not implemented) 6379
 3.677.6 Sympy [B] (verification not implemented) 6380
 3.677.7 Maxima [A] (verification not implemented) 6380
 3.677.8 Giac [B] (verification not implemented) 6381
 3.677.9 Mupad [B] (verification not implemented) 6381

3.677.1 Optimal result

Integrand size = 41, antiderivative size = 99

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^5 dx$$

$$= \frac{2a^2(iA + B)c^5(1 - i \tan(e + fx))^5}{5f} - \frac{a^2(iA + 3B)c^5(1 - i \tan(e + fx))^6}{6f} + \frac{a^2Bc^5(1 - i \tan(e + fx))^7}{7f}$$

output `2/5*a^2*(I*A+B)*c^5*(1-I*tan(f*x+e))^5/f-1/6*a^2*(I*A+3*B)*c^5*(1-I*tan(f*x+e))^6/f+1/7*a^2*B*c^5*(1-I*tan(f*x+e))^7/f`

3.677.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 270 vs. 2(99) = 198.

Time = 1.99 (sec) , antiderivative size = 270, normalized size of antiderivative = 2.73

$$\begin{aligned} & \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ic \tan(e + fx))^5 dx \\ &= \frac{a^2 Ac^5 \tan(e + fx)}{f} - \frac{3ia^2 Ac^5 \tan^2(e + fx)}{2f} + \frac{a^2 Bc^5 \tan^2(e + fx)}{2f} \\ & \quad - \frac{2a^2 Ac^5 \tan^3(e + fx)}{3f} - \frac{ia^2 Bc^5 \tan^3(e + fx)}{f} - \frac{ia^2 Ac^5 \tan^4(e + fx)}{2f} \\ & \quad - \frac{a^2 Bc^5 \tan^4(e + fx)}{2f} - \frac{3a^2 Ac^5 \tan^5(e + fx)}{5f} - \frac{2ia^2 Bc^5 \tan^5(e + fx)}{5f} \\ & \quad + \frac{ia^2 Ac^5 \tan^6(e + fx)}{6f} - \frac{a^2 Bc^5 \tan^6(e + fx)}{2f} + \frac{ia^2 Bc^5 \tan^7(e + fx)}{7f} \end{aligned}$$

input `Integrate[(a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^5,x]`

output `(a^2*A*c^5*Tan[e + f*x])/f - (((3*I)/2)*a^2*A*c^5*Tan[e + f*x]^2)/f + (a^2*B*c^5*Tan[e + f*x]^2)/(2*f) - (2*a^2*A*c^5*Tan[e + f*x]^3)/(3*f) - (I*a^2*B*c^5*Tan[e + f*x]^3)/f - ((I/2)*a^2*A*c^5*Tan[e + f*x]^4)/f - (a^2*B*c^5*Tan[e + f*x]^4)/(2*f) - (3*a^2*A*c^5*Tan[e + f*x]^5)/(5*f) - (((2*I)/5)*a^2*B*c^5*Tan[e + f*x]^5)/f + ((I/6)*a^2*A*c^5*Tan[e + f*x]^6)/f - (a^2*B*c^5*Tan[e + f*x]^6)/(2*f) + ((I/7)*a^2*B*c^5*Tan[e + f*x]^7)/f`

3.677.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.83, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^5 (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^5 (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \end{aligned}$$

$$\frac{ac \int ac^4(1 - i \tan(e + fx))^4(i \tan(e + fx) + 1)(A + B \tan(e + fx))d \tan(e + fx)}{f}$$

↓ 27

$$\frac{a^2c^5 \int (1 - i \tan(e + fx))^4(i \tan(e + fx) + 1)(A + B \tan(e + fx))d \tan(e + fx)}{f}$$

↓ 86

$$\frac{a^2c^5 \int (-iB(1 - i \tan(e + fx))^6 + (3iB - A)(1 - i \tan(e + fx))^5 + 2(A - iB)(1 - i \tan(e + fx))^4) d \tan(e + fx)}{f}$$

↓ 2009

$$\frac{a^2c^5(-\frac{1}{6}(3B + iA)(1 - i \tan(e + fx))^6 + \frac{2}{5}(B + iA)(1 - i \tan(e + fx))^5 + \frac{1}{7}B(1 - i \tan(e + fx))^7)}{f}$$

input `Int[(a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^5, x]`

output `(a^2*c^5*((2*(I*A + B)*(1 - I*Tan[e + f*x])^5)/5 - ((I*A + 3*B)*(1 - I*Tan[e + f*x])^6)/6 + (B*(1 - I*Tan[e + f*x])^7)/7))/f`

3.677.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.677. $\int (a + ia \tan(e + fx))^2(A + B \tan(e + fx))(c - ictan(e + fx))^5 dx$

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x],
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.677.4 Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.84

method	result
risch	$\frac{32c^5a^2(42iAe^{4i(fx+e)}+42Be^{4i(fx+e)}+49iAe^{2i(fx+e)}-21Be^{2i(fx+e)}+7iA-3B)}{105f(e^{2i(fx+e)}+1)^7}$
derivativedivides	$-\frac{ic^5a^2\left(-\frac{B\tan(fx+e)^7}{7}+\frac{(-3iB-A)\tan(fx+e)^6}{6}+\frac{(iA+4i(iB-A)+6B)\tan(fx+e)^5}{5}+\frac{(-2iB+2A)\tan(fx+e)^4}{4}+\frac{(-6iA-4i(iB-A)+3A^2)\tan(fx+e)^3}{3}+\frac{(-2iA+2A^2)\tan(fx+e)^2}{2}+(-iA+A^2)\tan(fx+e)+A^2\right)}{f}$
default	$-\frac{ic^5a^2\left(-\frac{B\tan(fx+e)^7}{7}+\frac{(-3iB-A)\tan(fx+e)^6}{6}+\frac{(iA+4i(iB-A)+6B)\tan(fx+e)^5}{5}+\frac{(-2iB+2A)\tan(fx+e)^4}{4}+\frac{(-6iA-4i(iB-A)+3A^2)\tan(fx+e)^3}{3}+\frac{(-2iA+2A^2)\tan(fx+e)^2}{2}+(-iA+A^2)\tan(fx+e)+A^2\right)}{f}$
norman	$\frac{Aa^2c^5\tan(fx+e)}{f}-\frac{(-iAa^2c^5+3Ba^2c^5)\tan(fx+e)^6}{6f}-\frac{(2iBa^2c^5+3Aa^2c^5)\tan(fx+e)^5}{5f}-\frac{(3iBa^2c^5+2Aa^2c^5)\tan(fx+e)^4}{3f}$
parallelrisch	$30iBa^2c^5\tan(fx+e)^7+35iA\tan(fx+e)^6a^2c^5-84iB\tan(fx+e)^5a^2c^5-105B\tan(fx+e)^6a^2c^5-105iA\tan(fx+e)^4a^2c^5$
parts	$\frac{(-5iAa^2c^5-Ba^2c^5)\left(\frac{\tan(fx+e)^2}{2}-\frac{\ln(1+\tan(fx+e)^2)}{2}\right)}{f}+\frac{(-5iBa^2c^5-5Aa^2c^5)\left(\frac{\tan(fx+e)^3}{3}-\tan(fx+e)+\arctan(\tan(fx+e))\right)}{f}$

```
input int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5,x,method=_R
ETURNVERBOSE)
```

```
output 32/105*c^5*a^2*(42*I*A*exp(4*I*(f*x+e))+42*B*exp(4*I*(f*x+e))+49*I*A*exp(2
*I*(f*x+e))-21*B*exp(2*I*(f*x+e))+7*I*A-3*B)/f/(exp(2*I*(f*x+e))+1)^7
```

3.677.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.54

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^5 dx =$$

$$-\frac{32(42(-iA - B)a^2c^5e^{4i fx+4ie} + 7(-7iA + 3B)a^2c^5e^{2i fx+2ie} + (-7iA + 3B)a^2c^5e^{2i fx+2ie} + (-7iA + 3B)a^2c^5e^{2i fx+2ie} + (-7iA + 3B)a^2c^5e^{2i fx+2ie})}{105(fe^{14i fx+14ie} + 7fe^{12i fx+12ie} + 21fe^{10i fx+10ie} + 35fe^{8i fx+8ie} + 35fe^{6i fx+6ie} + 21fe^{4i fx+4ie})}$$

3.677. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^5 dx$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5,x, algorithm="fricas")`

output `-32/105*(42*(-I*A - B)*a^2*c^5*e^(4*I*f*x + 4*I*e) + 7*(-7*I*A + 3*B)*a^2*c^5*e^(2*I*f*x + 2*I*e) + (-7*I*A + 3*B)*a^2*c^5)/(f*e^(14*I*f*x + 14*I*e) + 7*f*e^(12*I*f*x + 12*I*e) + 21*f*e^(10*I*f*x + 10*I*e) + 35*f*e^(8*I*f*x + 8*I*e) + 35*f*e^(6*I*f*x + 6*I*e) + 21*f*e^(4*I*f*x + 4*I*e) + 7*f*e^(2*I*f*x + 2*I*e) + f)`

3.677.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 243 vs. $2(80) = 160$.

Time = 0.61 (sec) , antiderivative size = 243, normalized size of antiderivative = 2.45

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^5 dx$$

$$= \frac{224iAa^2c^5 - 96Ba^2c^5 + (1568iAa^2c^5e^{2ie} - 672Ba^2c^5e^{2ie})e^{2ifx} + (1344iAa^2c^5e^{4ie} + 1344Ba^2c^5e^{4ie})e^{4ifx} + (105fe^{14ie}e^{14ifx} + 735fe^{12ie}e^{12ifx} + 2205fe^{10ie}e^{10ifx} + 3675fe^{8ie}e^{8ifx} + 3675fe^{6ie}e^{6ifx} + 2205fe^{4ie}e^{4ifx} + 735fe^{2ie}e^{2ifx} + 105f)}{105fe^{14ie}e^{14ifx} + 735fe^{12ie}e^{12ifx} + 2205fe^{10ie}e^{10ifx} + 3675fe^{8ie}e^{8ifx} + 3675fe^{6ie}e^{6ifx} + 2205fe^{4ie}e^{4ifx} + 735fe^{2ie}e^{2ifx} + 105f}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**5,x)`

output `(224*I*A*a**2*c**5 - 96*B*a**2*c**5 + (1568*I*A*a**2*c**5*exp(2*I*e) - 672*B*a**2*c**5*exp(2*I*e))*exp(2*I*f*x) + (1344*I*A*a**2*c**5*exp(4*I*e) + 1344*B*a**2*c**5*exp(4*I*e))*exp(4*I*f*x))/(105*f*exp(14*I*e)*exp(14*I*f*x) + 735*f*exp(12*I*e)*exp(12*I*f*x) + 2205*f*exp(10*I*e)*exp(10*I*f*x) + 3675*f*exp(8*I*e)*exp(8*I*f*x) + 3675*f*exp(6*I*e)*exp(6*I*f*x) + 2205*f*exp(4*I*e)*exp(4*I*f*x) + 735*f*exp(2*I*e)*exp(2*I*f*x) + 105*f)`

3.677.7 Maxima [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.53

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^5 dx$$

$$= \frac{30iBa^2c^5 \tan(fx + e)^7 + 35(iA - 3B)a^2c^5 \tan(fx + e)^6 - 42(3A + 2iB)a^2c^5 \tan(fx + e)^5 + 105(-iA + 3B)a^2c^5 \tan(fx + e)^4 + 105f \tan(fx + e)^3}{105f \tan(fx + e)^3 + 3675f \tan(fx + e)^2 + 2205f \tan(fx + e) + 105f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5,x, algorithm="maxima")`

output $\frac{1}{210}*(30*I*B*a^2*c^5*\tan(f*x + e)^7 + 35*(I*A - 3*B)*a^2*c^5*\tan(f*x + e)^6 - 42*(3*A + 2*I*B)*a^2*c^5*\tan(f*x + e)^5 + 105*(-I*A - B)*a^2*c^5*\tan(f*x + e)^4 - 70*(2*A + 3*I*B)*a^2*c^5*\tan(f*x + e)^3 + 105*(-3*I*A + B)*a^2*c^5*\tan(f*x + e)^2 + 210*A*a^2*c^5*\tan(f*x + e))/f$

3.677.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 180 vs. $2(83) = 166$.

Time = 1.02 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.82

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^5 dx = \frac{32(-42i Aa^2 c^5 e^{(4i fx + 4ie)} - 42 Ba^2 c^5 e^{(4i fx + 4ie)} - 49i Aa^2 c^5 e^{(2i fx + 2ie)} + 21 Ba^2 c^5 e^{(2i fx + 2ie)} - 7i Aa^2 c^5 e^{(4i fx + 4ie)} - 7i Ba^2 c^5 e^{(4i fx + 4ie)} + 7i Aa^2 c^5 e^{(2i fx + 2ie)} + 7i Ba^2 c^5 e^{(2i fx + 2ie)})}{105(fe^{(14i fx + 14ie)} + 7fe^{(12i fx + 12ie)} + 21fe^{(10i fx + 10ie)} + 35fe^{(8i fx + 8ie)} + 35fe^{(6i fx + 6ie)} + 21fe^{(4i fx + 4ie)})}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5,x, algorithm="giac")`

output $\frac{-32/105*(-42*I*A*a^2*c^5*e^{(4*I*f*x + 4*I*e)} - 42*B*a^2*c^5*e^{(4*I*f*x + 4*I*e)} - 49*I*A*a^2*c^5*e^{(2*I*f*x + 2*I*e)} + 21*B*a^2*c^5*e^{(2*I*f*x + 2*I*e)} - 7*I*A*a^2*c^5 + 3*B*a^2*c^5)/(f*e^{(14*I*f*x + 14*I*e)} + 7*f*e^{(12*I*f*x + 12*I*e)} + 21*f*e^{(10*I*f*x + 10*I*e)} + 35*f*e^{(8*I*f*x + 8*I*e)} + 35*f*e^{(6*I*f*x + 6*I*e)} + 21*f*e^{(4*I*f*x + 4*I*e)} + 7*f*e^{(2*I*f*x + 2*I*e)} + f)$

3.677.9 Mupad [B] (verification not implemented)

Time = 9.22 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.60

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^5 dx = \frac{Aa^2 c^5 \tan(e + fx) + \frac{a^2 c^5 \tan(e + fx)^3 (-3B + A2i) li}{3} + \frac{a^2 c^5 \tan(e + fx)^5 (-2B + A3i) li}{5} - \frac{a^2 c^5 \tan(e + fx)^2 (3A + B li) li}{2} - \frac{a^2 c^5 \tan(e + fx)^4 (3A + B li) li}{2}}{f}$$

3.677. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^5 dx$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)^5,x)`

output `((a^2*c^5*tan(e + f*x)^3*(A*2i - 3*B)*1i)/3 - (a^2*c^5*tan(e + f*x)^2*(3*A + B*1i)*1i)/2 + (a^2*c^5*tan(e + f*x)^5*(A*3i - 2*B)*1i)/5 + A*a^2*c^5*tan(e + f*x) - (a^2*c^5*tan(e + f*x)^4*(A - B*1i)*1i)/2 + (a^2*c^5*tan(e + f*x)^6*(A + B*3i)*1i)/6 + (B*a^2*c^5*tan(e + f*x)^7*1i)/7)/f`

3.678 $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^4 dx$

3.678.1 Optimal result	6383
3.678.2 Mathematica [A] (verified)	6383
3.678.3 Rubi [A] (verified)	6384
3.678.4 Maple [A] (verified)	6386
3.678.5 Fricas [A] (verification not implemented)	6386
3.678.6 Sympy [B] (verification not implemented)	6387
3.678.7 Maxima [A] (verification not implemented)	6387
3.678.8 Giac [B] (verification not implemented)	6388
3.678.9 Mupad [B] (verification not implemented)	6388

3.678.1 Optimal result

Integrand size = 41, antiderivative size = 99

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^4 dx$$

$$= \frac{a^2(iA + B)c^4(1 - i \tan(e + fx))^4}{2f} - \frac{a^2(iA + 3B)c^4(1 - i \tan(e + fx))^5}{5f} + \frac{a^2Bc^4(1 - i \tan(e + fx))^6}{6f}$$

```
output 1/2*a^2*(I*A+B)*c^4*(1-I*tan(f*x+e))^4/f-1/5*a^2*(I*A+3*B)*c^4*(1-I*tan(f*x+e))^5/f+1/6*a^2*B*c^4*(1-I*tan(f*x+e))^6/f
```

3.678.2 Mathematica [A] (verified)

Time = 2.31 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.04

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^4 dx =$$

$$\frac{a^2c^4(6iA - 7B - 30A \tan(e + fx) + (30iA - 15B) \tan^2(e + fx) + 20iB \tan^3(e + fx) + 15iA \tan^4(e + fx))}{30f}$$

```
input Integrate[(a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^4,x]
```

output
$$\frac{-1/30*(a^2*c^4*((6*I)*A - 7*B - 30*A*\text{Tan}[e + f*x] + ((30*I)*A - 15*B)*\text{Tan}[e + f*x]^2 + (20*I)*B*\text{Tan}[e + f*x]^3 + (15*I)*A*\text{Tan}[e + f*x]^4 + 6*(A + (2*I)*B)*\text{Tan}[e + f*x]^5 + 5*B*\text{Tan}[e + f*x]^6))/f}$$

3.678.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.83, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))^2 (c - ictan(e + fx))^4 (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))^2 (c - ictan(e + fx))^4 (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int ac^3 (1 - i \tan(e + fx))^3 (i \tan(e + fx) + 1) (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{a^2 c^4 \int (1 - i \tan(e + fx))^3 (i \tan(e + fx) + 1) (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\ & \quad \downarrow \text{86} \\ & \frac{a^2 c^4 \int (-iB(1 - i \tan(e + fx))^5 + (3iB - A)(1 - i \tan(e + fx))^4 + 2(A - iB)(1 - i \tan(e + fx))^3) d \tan(e + fx)}{f} \\ & \quad \downarrow \text{2009} \\ & \frac{a^2 c^4 (-\frac{1}{5}(3B + iA)(1 - i \tan(e + fx))^5 + \frac{1}{2}(B + iA)(1 - i \tan(e + fx))^4 + \frac{1}{6}B(1 - i \tan(e + fx))^6)}{f} \end{aligned}$$

input
$$\text{Int}[(a + I*a*\text{Tan}[e + f*x])^2*(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^4, x]$$

3.678.
$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^4 dx$$

output $(a^2c^4(((I*A + B)*(1 - I*\text{Tan}[e + f*x])^4)/2 - ((I*A + 3*B)*(1 - I*\text{Tan}[e + f*x])^5)/5 + (B*(1 - I*\text{Tan}[e + f*x])^6)/6))/f$

3.678.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.678.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.84

method	result
risch	$\frac{8c^4 a^2 (15iA e^{4i(fx+e)} + 15B e^{4i(fx+e)} + 18iA e^{2i(fx+e)} - 6B e^{2i(fx+e)} + 3iA - B)}{15f (e^{2i(fx+e)} + 1)^6}$
derivativedivides	$c^4 a^2 \left(-\frac{B \tan(fx+e)^6}{6} + \frac{(-2iB - A) \tan(fx+e)^5}{5} + \frac{(iA + 3i(iB - A) + 3B) \tan(fx+e)^4}{4} - \frac{2iB \tan(fx+e)^3}{3} + \frac{(-3iA - i(iB - A)) \tan(fx+e)^2}{2} \right) \frac{1}{f}$
default	$c^4 a^2 \left(-\frac{B \tan(fx+e)^6}{6} + \frac{(-2iB - A) \tan(fx+e)^5}{5} + \frac{(iA + 3i(iB - A) + 3B) \tan(fx+e)^4}{4} - \frac{2iB \tan(fx+e)^3}{3} + \frac{(-3iA - i(iB - A)) \tan(fx+e)^2}{2} \right) \frac{1}{f}$
norman	$\frac{A a^2 c^4 \tan(fx+e)}{f} - \frac{(2iB a^2 c^4 + A a^2 c^4) \tan(fx+e)^5}{5f} + \frac{(-2iA a^2 c^4 + B a^2 c^4) \tan(fx+e)^2}{2f} - \frac{B a^2 c^4 \tan(fx+e)^6}{6f}$
parallelrisch	$-\frac{12iB \tan(fx+e)^5 a^2 c^4 + 5B \tan(fx+e)^6 a^2 c^4 + 15iA a^2 c^4 \tan(fx+e)^4 + 6A \tan(fx+e)^5 a^2 c^4 + 20iB a^2 c^4 \tan(fx+e)^3 + 3A a^2 c^4}{30f}$
parts	$\frac{(-4iA a^2 c^4 + B a^2 c^4) \left(\frac{\tan(fx+e)^2}{2} - \frac{\ln(1 + \tan(fx+e)^2)}{2} \right)}{f} + \frac{(-4iB a^2 c^4 - A a^2 c^4) \left(\frac{\tan(fx+e)^3}{3} - \tan(fx+e) + \arctan(\tan(fx+e)) \right)}{f}$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4,x,method=_R ETURNVERBOSE)`

output `8/15*c^4*a^2*(15*I*A*exp(4*I*(f*x+e))+15*B*exp(4*I*(f*x+e))+18*I*A*exp(2*I*(f*x+e))-6*B*exp(2*I*(f*x+e))+3*I*A-B)/f/(exp(2*I*(f*x+e))+1)^6`

3.678.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.37

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^4 dx =$$

$$-\frac{8(15(-iA - B)a^2 c^4 e^{(4i fx + 4i e)} + 6(-3iA + B)a^2 c^4 e^{(2i fx + 2i e)} + (-3iA + B)a^2 c^4)}{15(fe^{(12i fx + 12i e)} + 6fe^{(10i fx + 10i e)} + 15fe^{(8i fx + 8i e)} + 20fe^{(6i fx + 6i e)} + 15fe^{(4i fx + 4i e)} + 6fe^{(2i fx + 2i e)})}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4,x, algorithm="fricas")`

output `-8/15*(15*(-I*A - B)*a^2*c^4*e^(4*I*f*x + 4*I*e) + 6*(-3*I*A + B)*a^2*c^4*e^(2*I*f*x + 2*I*e) + (-3*I*A + B)*a^2*c^4)/(f*e^(12*I*f*x + 12*I*e) + 6*f*e^(10*I*f*x + 10*I*e) + 15*f*e^(8*I*f*x + 8*I*e) + 20*f*e^(6*I*f*x + 6*I*e) + 15*f*e^(4*I*f*x + 4*I*e) + 6*f*e^(2*I*f*x + 2*I*e) + f)`

3.678. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^4 dx$

3.678.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 224 vs. $2(78) = 156$.

Time = 0.49 (sec) , antiderivative size = 224, normalized size of antiderivative = 2.26

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ict \tan(e + fx))^4 dx$$

$$= \frac{24iAa^2c^4 - 8Ba^2c^4 + (144iAa^2c^4e^{2ie} - 48Ba^2c^4e^{2ie})e^{2ifx} + (120iAa^2c^4e^{4ie} + 120Ba^2c^4e^{4ie})e^{4ifx}}{15fe^{12ie}e^{12ifx} + 90fe^{10ie}e^{10ifx} + 225fe^{8ie}e^{8ifx} + 300fe^{6ie}e^{6ifx} + 225fe^{4ie}e^{4ifx} + 90fe^{2ie}e^{2ifx} + 15f}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**4,x)`

output `(24*I*A*a**2*c**4 - 8*B*a**2*c**4 + (144*I*A*a**2*c**4*exp(2*I*e) - 48*B*a**2*c**4*exp(2*I*e))*exp(2*I*f*x) + (120*I*A*a**2*c**4*exp(4*I*e) + 120*B*a**2*c**4*exp(4*I*e))*exp(4*I*f*x))/(15*f*exp(12*I*e)*exp(12*I*f*x) + 90*f*exp(10*I*e)*exp(10*I*f*x) + 225*f*exp(8*I*e)*exp(8*I*f*x) + 300*f*exp(6*I*e)*exp(6*I*f*x) + 225*f*exp(4*I*e)*exp(4*I*f*x) + 90*f*exp(2*I*e)*exp(2*I*f*x) + 15*f)`

3.678.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.17

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ict \tan(e + fx))^4 dx =$$

$$\frac{5Ba^2c^4 \tan(fx + e)^6 + 6(A + 2iB)a^2c^4 \tan(fx + e)^5 + 15iAa^2c^4 \tan(fx + e)^4 + 20iBa^2c^4 \tan(fx + e)^3 + 15(2iA - B)a^2c^4 \tan(fx + e)^2 - 30Aa^2c^4 \tan(fx + e)}{30f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4,x, algorithm="maxima")`

output `-1/30*(5*B*a^2*c^4*tan(f*x + e)^6 + 6*(A + 2*I*B)*a^2*c^4*tan(f*x + e)^5 + 15*I*A*a^2*c^4*tan(f*x + e)^4 + 20*I*B*a^2*c^4*tan(f*x + e)^3 + 15*(2*I*A - B)*a^2*c^4*tan(f*x + e)^2 - 30*A*a^2*c^4*tan(f*x + e))/f`

3.678.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 167 vs. $2(83) = 166$.

Time = 0.89 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.69

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^4 dx =$$

$$\frac{8(-15i Aa^2 c^4 e^{(4i fx + 4i e)} - 15 B a^2 c^4 e^{(4i fx + 4i e)} - 18i A a^2 c^4 e^{(2i fx + 2i e)} + 6 B a^2 c^4 e^{(2i fx + 2i e)} - 3i A a^2 c^4 + 15 f e^{(12i fx + 12i e)} + 6 f e^{(10i fx + 10i e)} + 15 f e^{(8i fx + 8i e)} + 20 f e^{(6i fx + 6i e)} + 15 f e^{(4i fx + 4i e)} + 6 f e^{(2i fx + 2i e)})}{15 f e^{(12i fx + 12i e)} + 6 f e^{(10i fx + 10i e)} + 15 f e^{(8i fx + 8i e)} + 20 f e^{(6i fx + 6i e)} + 15 f e^{(4i fx + 4i e)} + 6 f e^{(2i fx + 2i e)}}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4,x, algorithm="giac")`

output `-8/15*(-15*I*A*a^2*c^4*e^(4*I*f*x + 4*I*e) - 15*B*a^2*c^4*e^(4*I*f*x + 4*I*e) - 18*I*A*a^2*c^4*e^(2*I*f*x + 2*I*e) + 6*B*a^2*c^4*e^(2*I*f*x + 2*I*e) - 3*I*A*a^2*c^4 + B*a^2*c^4)/(f*e^(12*I*f*x + 12*I*e) + 6*f*e^(10*I*f*x + 10*I*e) + 15*f*e^(8*I*f*x + 8*I*e) + 20*f*e^(6*I*f*x + 6*I*e) + 15*f*e^(4*I*f*x + 4*I*e) + 6*f*e^(2*I*f*x + 2*I*e) + f)`

3.678.9 Mupad [B] (verification not implemented)

Time = 9.28 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.21

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^4 dx =$$

$$-\frac{a^2 c^4 \tan(e+fx)^2 (-B+A 2i)}{2} - A a^2 c^4 \tan(e + fx) + \frac{a^2 c^4 \tan(e+fx)^5 (A+B 2i)}{5} + \frac{B a^2 c^4 \tan(e+fx)^6}{6} + \frac{A a^2 c^4 \tan(e+fx)}{2} f$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)^4,x)`

output `-((a^2*c^4*tan(e + f*x)^2*(A*2i - B))/2 - A*a^2*c^4*tan(e + f*x) + (A*a^2*c^4*tan(e + f*x)^4*1i)/2 + (a^2*c^4*tan(e + f*x)^5*(A + B*2i))/5 + (B*a^2*c^4*tan(e + f*x)^3*2i)/3 + (B*a^2*c^4*tan(e + f*x)^6)/6)/f`

3.679 $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^3 dx$

3.679.1 Optimal result	6389
3.679.2 Mathematica [A] (verified)	6389
3.679.3 Rubi [A] (verified)	6390
3.679.4 Maple [A] (verified)	6392
3.679.5 Fricas [A] (verification not implemented)	6392
3.679.6 Sympy [B] (verification not implemented)	6393
3.679.7 Maxima [A] (verification not implemented)	6393
3.679.8 Giac [A] (verification not implemented)	6394
3.679.9 Mupad [B] (verification not implemented)	6394

3.679.1 Optimal result

Integrand size = 41, antiderivative size = 99

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^3 dx$$

$$= \frac{2a^2(iA + B)c^3(1 - i \tan(e + fx))^3}{3f} - \frac{a^2(iA + 3B)c^3(1 - i \tan(e + fx))^4}{4f} + \frac{a^2Bc^3(1 - i \tan(e + fx))^5}{5f}$$

```
output 2/3*a^2*(I*A+B)*c^3*(1-I*tan(f*x+e))^3/f-1/4*a^2*(I*A+3*B)*c^3*(1-I*tan(f*x+e))^4/f+1/5*a^2*B*c^3*(1-I*tan(f*x+e))^5/f
```

3.679.2 Mathematica [A] (verified)

Time = 5.62 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.96

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^3 dx$$

$$= \frac{a^2c^3(5iA + 11B + 60A \tan(e + fx) + 30(-iA + B) \tan^2(e + fx) + 20(A - iB) \tan^3(e + fx) + 15(-iA - B) \tan^4(e + fx))}{60f}$$

```
input Integrate[(a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^3,x]
```

output $(a^2c^3((5I)A + 11B + 60A \operatorname{Tan}[e + f*x] + 30((-I)A + B) \operatorname{Tan}[e + f*x]^2 + 20(A - I*B) \operatorname{Tan}[e + f*x]^3 + 15((-I)A + B) \operatorname{Tan}[e + f*x]^4 - (12*I) * B \operatorname{Tan}[e + f*x]^5))/(60*f)$

3.679.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.83, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))^2 (c - ictan(e + fx))^3 (A + B \tan(e + fx)) dx$$

$$\downarrow 3042$$

$$\int (a + ia \tan(e + fx))^2 (c - ictan(e + fx))^3 (A + B \tan(e + fx)) dx$$

$$\downarrow 4071$$

$$\frac{ac \int ac^2 (1 - i \tan(e + fx))^2 (i \tan(e + fx) + 1) (A + B \tan(e + fx)) d \tan(e + fx)}{f}$$

$$\downarrow 27$$

$$\frac{a^2c^3 \int (1 - i \tan(e + fx))^2 (i \tan(e + fx) + 1) (A + B \tan(e + fx)) d \tan(e + fx)}{f}$$

$$\downarrow 86$$

$$\frac{a^2c^3 \int (-iB(1 - i \tan(e + fx))^4 + (3iB - A)(1 - i \tan(e + fx))^3 + 2(A - iB)(1 - i \tan(e + fx))^2) d \tan(e + fx)}{f}$$

$$\downarrow 2009$$

$$\frac{a^2c^3 (-\frac{1}{4}(3B + iA)(1 - i \tan(e + fx))^4 + \frac{2}{3}(B + iA)(1 - i \tan(e + fx))^3 + \frac{1}{5}B(1 - i \tan(e + fx))^5)}{f}$$

input $\operatorname{Int}[(a + I*a*\operatorname{Tan}[e + f*x])^2*(A + B*\operatorname{Tan}[e + f*x])*(c - I*c*\operatorname{Tan}[e + f*x])^3, x]$

$$3.679. \quad \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^3 dx$$

output $(a^2c^3((2*(I*A + B)*(1 - I*\tan[e + f*x])^3)/3 - ((I*A + 3*B)*(1 - I*\tan[e + f*x])^4)/4 + (B*(1 - I*\tan[e + f*x])^5)/5))/f$

3.679.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.679.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.84

method	result
risch	$\frac{4c^3 a^2 (20iA e^{4i(fx+e)} + 20B e^{4i(fx+e)} + 25iA e^{2i(fx+e)} - 5B e^{2i(fx+e)} + 5iA - B)}{15f (e^{2i(fx+e)} + 1)^5}$
derivativedivides	$\frac{ic^3 a^2 \left(-\frac{B \tan(fx+e)^5}{5} + \frac{(-iB-A) \tan(fx+e)^4}{4} + \frac{(iA+2i(iB-A)+B) \tan(fx+e)^3}{3} + \frac{(-iB-A) \tan(fx+e)^2}{2} - i \tan(fx+e)A \right)}{f}$
default	$\frac{ic^3 a^2 \left(-\frac{B \tan(fx+e)^5}{5} + \frac{(-iB-A) \tan(fx+e)^4}{4} + \frac{(iA+2i(iB-A)+B) \tan(fx+e)^3}{3} + \frac{(-iB-A) \tan(fx+e)^2}{2} - i \tan(fx+e)A \right)}{f}$
norman	$\frac{A a^2 c^3 \tan(fx+e)}{f} + \frac{(-iA a^2 c^3 + B a^2 c^3) \tan(fx+e)^2}{2f} + \frac{(-iA a^2 c^3 + B a^2 c^3) \tan(fx+e)^4}{4f} + \frac{(-iB a^2 c^3 + A a^2 c^3) \tan(fx+e)^5}{3f}$
parallelrisch	$-\frac{12iB a^2 c^3 \tan(fx+e)^5 + 15iA \tan(fx+e)^4 a^2 c^3 + 20iB \tan(fx+e)^3 a^2 c^3 - 15B \tan(fx+e)^4 a^2 c^3 + 30iA \tan(fx+e)^2 a^2 c^3}{60f}$
parts	$\frac{(-2iA a^2 c^3 + 2B a^2 c^3) \left(\frac{\tan(fx+e)^2}{2} - \frac{\ln(1+\tan(fx+e)^2)}{2} \right)}{f} + \frac{(-2iB a^2 c^3 + A a^2 c^3) \left(\frac{\tan(fx+e)^3}{3} - \tan(fx+e) + \arctan(\tan(fx+e)) \right)}{f}$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3,x,method=_R
ETURNVERBOSE)`

output `4/15*c^3*a^2*(20*I*A*exp(4*I*(f*x+e))+20*B*exp(4*I*(f*x+e))+25*I*A*exp(2*I
*(f*x+e))-5*B*exp(2*I*(f*x+e))+5*I*A-B)/f/(exp(2*I*(f*x+e))+1)^5`

3.679.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.25

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^3 dx$$

$$= -\frac{4(20(-iA - B)a^2 c^3 e^{(4i fx + 4i e)} + 5(-5iA + B)a^2 c^3 e^{(2i fx + 2i e)} + (-5iA + B)a^2 c^3)}{15(fe^{(10i fx + 10i e)} + 5fe^{(8i fx + 8i e)} + 10fe^{(6i fx + 6i e)} + 10fe^{(4i fx + 4i e)} + 5fe^{(2i fx + 2i e)} + f)}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3,x, al
gorithm="fracas")`

output `-4/15*(20*(-I*A - B)*a^2*c^3*e^(4*I*f*x + 4*I*e) + 5*(-5*I*A + B)*a^2*c^3*
e^(2*I*f*x + 2*I*e) + (-5*I*A + B)*a^2*c^3)/(f*e^(10*I*f*x + 10*I*e) + 5*f
*e^(8*I*f*x + 8*I*e) + 10*f*e^(6*I*f*x + 6*I*e) + 10*f*e^(4*I*f*x + 4*I*e)
+ 5*f*e^(2*I*f*x + 2*I*e) + f)`

3.679. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^3 dx$

3.679.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 206 vs. $2(80) = 160$.

Time = 0.39 (sec) , antiderivative size = 206, normalized size of antiderivative = 2.08

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^3 dx$$

$$= \frac{20iAa^2c^3 - 4Ba^2c^3 + (100iAa^2c^3e^{2ie} - 20Ba^2c^3e^{2ie})e^{2ifx} + (80iAa^2c^3e^{4ie} + 80Ba^2c^3e^{4ie})e^{4ifx}}{15fe^{10ie}e^{10ifx} + 75fe^{8ie}e^{8ifx} + 150fe^{6ie}e^{6ifx} + 150fe^{4ie}e^{4ifx} + 75fe^{2ie}e^{2ifx} + 15f}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**3,x)`

output `(20*I*A*a**2*c**3 - 4*B*a**2*c**3 + (100*I*A*a**2*c**3*exp(2*I*e) - 20*B*a**2*c**3*exp(2*I*e))*exp(2*I*f*x) + (80*I*A*a**2*c**3*exp(4*I*e) + 80*B*a**2*c**3*exp(4*I*e))*exp(4*I*f*x))/(15*f*exp(10*I*e)*exp(10*I*f*x) + 75*f*exp(8*I*e)*exp(8*I*f*x) + 150*f*exp(6*I*e)*exp(6*I*f*x) + 150*f*exp(4*I*e)*exp(4*I*f*x) + 75*f*exp(2*I*e)*exp(2*I*f*x) + 15*f)`

3.679.7 Maxima [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.02

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^3 dx =$$

$$\frac{12iBa^2c^3 \tan(fx + e)^5 - 15(-iA + B)a^2c^3 \tan(fx + e)^4 - 20(A - iB)a^2c^3 \tan(fx + e)^3 - 30(-iA + B)a^2c^3 \tan(fx + e)^2 - 60Aa^2c^3 \tan(fx + e)}{60f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3,x, algorithm="maxima")`

output `-1/60*(12*I*B*a^2*c^3*tan(f*x + e)^5 - 15*(-I*A + B)*a^2*c^3*tan(f*x + e)^4 - 20*(A - I*B)*a^2*c^3*tan(f*x + e)^3 - 30*(-I*A + B)*a^2*c^3*tan(f*x + e)^2 - 60*A*a^2*c^3*tan(f*x + e))/f`

3.679.8 Giac [A] (verification not implemented)

Time = 0.66 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.57

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ict \tan(e + fx))^3 dx =$$

$$\frac{4(-20i Aa^2c^3e^{4ifx+4ie} - 20Ba^2c^3e^{4ifx+4ie} - 25i Aa^2c^3e^{2ifx+2ie} + 5Ba^2c^3e^{2ifx+2ie} - 5i Aa^2c^3 + 15fe^{10ifx+10ie} + 5fe^{8ifx+8ie} + 10fe^{6ifx+6ie} + 10fe^{4ifx+4ie} + 5fe^{2ifx+2ie} + f)}{15fe^{10ifx+10ie} + 5fe^{8ifx+8ie} + 10fe^{6ifx+6ie} + 10fe^{4ifx+4ie} + 5fe^{2ifx+2ie} + f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3,x, algorithm="giac")`

output `-4/15*(-20*I*A*a^2*c^3*e^(4*I*f*x + 4*I*e) - 20*B*a^2*c^3*e^(4*I*f*x + 4*I*e) - 25*I*A*a^2*c^3*e^(2*I*f*x + 2*I*e) + 5*B*a^2*c^3*e^(2*I*f*x + 2*I*e) - 5*I*A*a^2*c^3 + B*a^2*c^3)/(f*e^(10*I*f*x + 10*I*e) + 5*f*e^(8*I*f*x + 8*I*e) + 10*f*e^(6*I*f*x + 6*I*e) + 10*f*e^(4*I*f*x + 4*I*e) + 5*f*e^(2*I*f*x + 2*I*e) + f)`

3.679.9 Mupad [B] (verification not implemented)

Time = 9.43 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.09

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ict \tan(e + fx))^3 dx =$$

$$\frac{-Aa^2c^3 \tan(e + fx) + \frac{a^2c^3 \tan(e+fx)^2 (A+B \operatorname{li}) \operatorname{li}}{2} + \frac{a^2c^3 \tan(e+fx)^3 (B+A \operatorname{li}) \operatorname{li}}{3} + \frac{a^2c^3 \tan(e+fx)^4 (A+B \operatorname{li}) \operatorname{li}}{4} + \frac{Ba^2c^3 \tan(e+fx)^5 \operatorname{li}}{5}}{f}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)^3,x)`

output `-((a^2*c^3*tan(e + f*x)^2*(A + B*1i)*1i)/2 - A*a^2*c^3*tan(e + f*x) + (a^2*c^3*tan(e + f*x)^3*(A*1i + B)*1i)/3 + (a^2*c^3*tan(e + f*x)^4*(A + B*1i)*1i)/4 + (B*a^2*c^3*tan(e + f*x)^5*1i)/5)/f`

$$3.680 \quad \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx$$

3.680.1 Optimal result	6395
3.680.2 Mathematica [A] (verified)	6395
3.680.3 Rubi [A] (verified)	6396
3.680.4 Maple [A] (verified)	6398
3.680.5 Fricas [C] (verification not implemented)	6398
3.680.6 Sympy [C] (verification not implemented)	6399
3.680.7 Maxima [A] (verification not implemented)	6399
3.680.8 Giac [B] (verification not implemented)	6400
3.680.9 Mupad [B] (verification not implemented)	6400

3.680.1 Optimal result

Integrand size = 41, antiderivative size = 62

$$\begin{aligned} & \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx \\ &= \frac{a^2 B c^2 \sec^4(e + fx)}{4f} + \frac{a^2 A c^2 \tan(e + fx)}{f} + \frac{a^2 A c^2 \tan^3(e + fx)}{3f} \end{aligned}$$

output `1/4*a^2*B*c^2*sec(f*x+e)^4/f+a^2*A*c^2*tan(f*x+e)/f+1/3*a^2*A*c^2*tan(f*x+e)^3/f`

3.680.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.85

$$\begin{aligned} & \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx \\ &= \frac{a^2 B c^2 \sec^4(e + fx)}{4f} + \frac{a^2 A c^2 (\tan(e + fx) + \frac{1}{3} \tan^3(e + fx))}{f} \end{aligned}$$

input `Integrate[(a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^2,x]`

$$3.680. \quad \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx$$

output $(a^2 B c^2 \text{Sec}[e + f x]^4)/(4 f) + (a^2 A c^2 (\text{Tan}[e + f x] + \text{Tan}[e + f x]^3/3))/f$

3.680.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.79, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.146$, Rules used = {3042, 4071, 27, 82, 455, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(e + fx))^2 (c - ictan(e + fx))^2 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(e + fx))^2 (c - ictan(e + fx))^2 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int ac(1 - i \tan(e + fx))(i \tan(e + fx) + 1)(A + B \tan(e + fx)) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^2 c^2 \int (1 - i \tan(e + fx))(i \tan(e + fx) + 1)(A + B \tan(e + fx)) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{82} \\
 & \frac{a^2 c^2 \int (A + B \tan(e + fx)) (\tan^2(e + fx) + 1) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{455} \\
 & \frac{a^2 c^2 \left(A \int (\tan^2(e + fx) + 1) d \tan(e + fx) + \frac{1}{4} B (\tan^2(e + fx) + 1)^2 \right)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^2 c^2 \left(A \left(\frac{1}{3} \tan^3(e + fx) + \tan(e + fx) \right) + \frac{1}{4} B (\tan^2(e + fx) + 1)^2 \right)}{f}
 \end{aligned}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^2*(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^2, x]$

3.680. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^2 dx$

output $(a^2c^2((B(1 + \tan[e + fx])^2)^2)/4 + A(\tan[e + fx] + \tan[e + fx]^3/3))/f$

3.680.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 82 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 455 `Int[((c_) + (d_)*(x_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[d*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] + Simp[c Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, p}, x] && !LeQ[p, -1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.680.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.85

method	result
derivativedivides	$\frac{c^2 a^2 \left(\frac{B \tan(fx+e)^4}{4} + \frac{A \tan(fx+e)^3}{3} + \frac{B \tan(fx+e)^2}{2} + A \tan(fx+e) \right)}{f}$
default	$\frac{c^2 a^2 \left(\frac{B \tan(fx+e)^4}{4} + \frac{A \tan(fx+e)^3}{3} + \frac{B \tan(fx+e)^2}{2} + A \tan(fx+e) \right)}{f}$
risch	$\frac{4c^2 a^2 (3iA e^{4i(fx+e)} + 3B e^{4i(fx+e)} + 4iA e^{2i(fx+e)} + iA)}{3f(e^{2i(fx+e)} + 1)^4}$
parallelrisch	$\frac{3B a^2 c^2 \tan(fx+e)^4 + 4A a^2 c^2 \tan(fx+e)^3 + 6B a^2 c^2 \tan(fx+e)^2 + 12A a^2 c^2 \tan(fx+e)}{12f}$
norman	$\frac{a^2 A c^2 \tan(fx+e)}{f} + \frac{B a^2 c^2 \tan(fx+e)^2}{2f} + \frac{B a^2 c^2 \tan(fx+e)^4}{4f} + \frac{a^2 A c^2 \tan(fx+e)^3}{3f}$
parts	$A a^2 c^2 x + \frac{A a^2 c^2 \left(\frac{\tan(fx+e)^3}{3} - \tan(fx+e) + \arctan(\tan(fx+e)) \right)}{f} + \frac{B a^2 c^2 \ln(1 + \tan(fx+e)^2)}{2f} + \frac{B a^2 c^2 \left(\frac{\tan(fx+e)}{1 + \tan(fx+e)^2} \right)}{f}$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2,x,method=_R
ETURNVERBOSE)`

output `1/f*c^2*a^2*(1/4*B*tan(f*x+e)^4+1/3*A*tan(f*x+e)^3+1/2*B*tan(f*x+e)^2+A*ta
n(f*x+e))`

3.680.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.22 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.69

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx$$

$$= -\frac{4(3(-iA - B)a^2 c^2 e^{(4i fx + 4i e)} - 4i A a^2 c^2 e^{(2i fx + 2i e)} - i A a^2 c^2)}{3(fe^{(8i fx + 8i e)} + 4fe^{(6i fx + 6i e)} + 6fe^{(4i fx + 4i e)} + 4fe^{(2i fx + 2i e)} + f)}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2,x, al
gorithm="fracas")`

output
$$-4/3*(3*(-I*A - B)*a^2*c^2*e^{(4*I*f*x + 4*I*e)} - 4*I*A*a^2*c^2*e^{(2*I*f*x + 2*I*e)} - I*A*a^2*c^2)/(f*e^{(8*I*f*x + 8*I*e)} + 4*f*e^{(6*I*f*x + 6*I*e)} + 6*f*e^{(4*I*f*x + 4*I*e)} + 4*f*e^{(2*I*f*x + 2*I*e)} + f)$$

3.680.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 160, normalized size of antiderivative = 2.58

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx$$

$$= \frac{16iAa^2c^2e^{2ie}e^{2ifx} + 4iAa^2c^2 + (12iAa^2c^2e^{4ie} + 12Ba^2c^2e^{4ie})e^{4ifx}}{3fe^{8ie}e^{8ifx} + 12fe^{6ie}e^{6ifx} + 18fe^{4ie}e^{4ifx} + 12fe^{2ie}e^{2ifx} + 3f}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**2,x)`

output
$$(16*I*A*a**2*c**2*\exp(2*I*e)*\exp(2*I*f*x) + 4*I*A*a**2*c**2 + (12*I*A*a**2*c**2*\exp(4*I*e) + 12*B*a**2*c**2*\exp(4*I*e))*\exp(4*I*f*x))/(3*f*\exp(8*I*e)*\exp(8*I*f*x) + 12*f*\exp(6*I*e)*\exp(6*I*f*x) + 18*f*\exp(4*I*e)*\exp(4*I*f*x) + 12*f*\exp(2*I*e)*\exp(2*I*f*x) + 3*f)$$

3.680.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.16

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^2 dx$$

$$= \frac{3Ba^2c^2 \tan(fx + e)^4 + 4Aa^2c^2 \tan(fx + e)^3 + 6Ba^2c^2 \tan(fx + e)^2 + 12Aa^2c^2 \tan(fx + e)}{12f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2,x, algorithm="maxima")`

output
$$1/12*(3*B*a^2*c^2*\tan(f*x + e)^4 + 4*A*a^2*c^2*\tan(f*x + e)^3 + 6*B*a^2*c^2*\tan(f*x + e)^2 + 12*A*a^2*c^2*\tan(f*x + e))/f$$

3.680.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 391 vs. 2(58) = 116.

Time = 0.54 (sec) , antiderivative size = 391, normalized size of antiderivative = 6.31

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ic \tan(e + fx))^2 dx$$

$$= \frac{3Ba^2c^2 \tan(fx)^4 \tan(e)^4 - 12Aa^2c^2 \tan(fx)^4 \tan(e)^3 - 12Aa^2c^2 \tan(fx)^3 \tan(e)^4 + 6Ba^2c^2 \tan(fx)^4}{12}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2,x, algorithm="giac")`

output `1/12*(3*B*a^2*c^2*tan(f*x)^4*tan(e)^4 - 12*A*a^2*c^2*tan(f*x)^4*tan(e)^3 - 12*A*a^2*c^2*tan(f*x)^3*tan(e)^4 + 6*B*a^2*c^2*tan(f*x)^4*tan(e)^2 + 6*B*a^2*c^2*tan(f*x)^2*tan(e)^4 - 4*A*a^2*c^2*tan(f*x)^4*tan(e) + 24*A*a^2*c^2*tan(f*x)^3*tan(e)^2 + 24*A*a^2*c^2*tan(f*x)^2*tan(e)^3 - 4*A*a^2*c^2*tan(f*x)*tan(e)^4 + 3*B*a^2*c^2*tan(f*x)^4 + 12*B*a^2*c^2*tan(f*x)^2*tan(e)^2 + 3*B*a^2*c^2*tan(e)^4 + 4*A*a^2*c^2*tan(f*x)^3 - 24*A*a^2*c^2*tan(f*x)^2*tan(e) - 24*A*a^2*c^2*tan(f*x)*tan(e)^2 + 4*A*a^2*c^2*tan(e)^3 + 6*B*a^2*c^2*tan(f*x)^2 + 6*B*a^2*c^2*tan(e)^2 + 12*A*a^2*c^2*tan(f*x) + 12*A*a^2*c^2*tan(e) + 3*B*a^2*c^2)/(f*tan(f*x)^4*tan(e)^4 - 4*f*tan(f*x)^3*tan(e)^3 + 6*f*tan(f*x)^2*tan(e)^2 - 4*f*tan(f*x)*tan(e) + f)`

3.680.9 Mupad [B] (verification not implemented)

Time = 8.74 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.32

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ic \tan(e + fx))^2 dx$$

$$= \frac{a^2 c^2 \sin(e + fx) (12 A \cos(e + fx)^3 + 6 B \cos(e + fx)^2 \sin(e + fx) + 4 A \cos(e + fx) \sin(e + fx)^2)}{12 f \cos(e + fx)^4}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)^2,x)`

output `(a^2*c^2*sin(e + f*x)*(12*A*cos(e + f*x)^3 + 3*B*sin(e + f*x)^3 + 4*A*cos(e + f*x)*sin(e + f*x)^2 + 6*B*cos(e + f*x)^2*sin(e + f*x)))/(12*f*cos(e + f*x)^4)`

3.680. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ic \tan(e + fx))^2 dx$

3.681 $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx)) dx$

3.681.1 Optimal result	6401
3.681.2 Mathematica [A] (verified)	6401
3.681.3 Rubi [A] (verified)	6402
3.681.4 Maple [A] (verified)	6403
3.681.5 Fricas [A] (verification not implemented)	6404
3.681.6 Sympy [B] (verification not implemented)	6404
3.681.7 Maxima [A] (verification not implemented)	6405
3.681.8 Giac [B] (verification not implemented)	6405
3.681.9 Mupad [B] (verification not implemented)	6406

3.681.1 Optimal result

Integrand size = 39, antiderivative size = 64

$$\begin{aligned} & \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx)) dx \\ &= \frac{a^2 A c \tan(e + fx)}{f} + \frac{a^2 (iA + B) c \tan^2(e + fx)}{2f} + \frac{ia^2 B c \tan^3(e + fx)}{3f} \end{aligned}$$

output $a^2 A c \tan(fx+e)/f + 1/2 a^2 (iA+B) c \tan(fx+e)^2/f + 1/3 i a^2 B c \tan(fx+e)^3/f$

3.681.2 Mathematica [A] (verified)

Time = 1.08 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.84

$$\begin{aligned} & \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx)) dx \\ &= \frac{a^2 c (-2B + 6A \tan(e + fx) + 3(iA + B) \tan^2(e + fx) + 2iB \tan^3(e + fx))}{6f} \end{aligned}$$

input `Integrate[(a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x]),x]`

3.681. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx)) dx$

output $(a^2*c*(-2*B + 6*A*\text{Tan}[e + f*x] + 3*(I*A + B)*\text{Tan}[e + f*x]^2 + (2*I)*B*\text{Tan}[e + f*x]^3))/(6*f)$

3.681.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.80, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.128$, Rules used = {3042, 4071, 27, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))^2 (c - ic \tan(e + fx)) (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))^2 (c - ic \tan(e + fx)) (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int a(i \tan(e + fx) + 1)(A + B \tan(e + fx)) d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{a^2 c \int (i \tan(e + fx) + 1)(A + B \tan(e + fx)) d \tan(e + fx)}{f} \\ & \quad \downarrow \text{49} \\ & \frac{a^2 c \int (iB \tan^2(e + fx) + (iA + B) \tan(e + fx) + A) d \tan(e + fx)}{f} \\ & \quad \downarrow \text{2009} \\ & \frac{a^2 c (\frac{1}{2}(B + iA) \tan^2(e + fx) + A \tan(e + fx) + \frac{1}{3}iB \tan^3(e + fx))}{f} \end{aligned}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^2*(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x]),x]$

output $(a^2*c*(A*\text{Tan}[e + f*x] + ((I*A + B)*\text{Tan}[e + f*x]^2)/2 + (I/3)*B*\text{Tan}[e + f*x]^3))/f$

3.681. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ic \tan(e + fx)) dx$

3.681.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.681.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.80

method	result
derivativedivides	$-\frac{ia^2c\left(-\frac{B \tan(fx+e)^3}{3} + \frac{(iB-A) \tan(fx+e)^2}{2} + i \tan(fx+e)A\right)}{f}$
default	$-\frac{ia^2c\left(-\frac{B \tan(fx+e)^3}{3} + \frac{(iB-A) \tan(fx+e)^2}{2} + i \tan(fx+e)A\right)}{f}$
norman	$\frac{a^2Ac \tan(fx+e)}{f} + \frac{(ia^2cA+a^2cB) \tan(fx+e)^2}{2f} + \frac{ia^2Bc \tan(fx+e)^3}{3f}$
parallelrisch	$\frac{2ia^2Bc \tan(fx+e)^3 + 3iA \tan(fx+e)^2 a^2c + 3B \tan(fx+e)^2 a^2c + 6A \tan(fx+e) a^2c}{6f}$
risch	$\frac{2a^2c(6iA e^{4i(fx+e)} + 6B e^{4i(fx+e)} + 9iA e^{2i(fx+e)} + 3B e^{2i(fx+e)} + 3iA + B)}{3f(e^{2i(fx+e)} + 1)^3}$
parts	$\frac{(iB a^2c + a^2cA) (\tan(fx+e) - \arctan(\tan(fx+e)))}{f} + \frac{(ia^2cA + a^2cB) \ln(1 + \tan(fx+e)^2)}{2f} + \frac{(ia^2cA + a^2cB) \left(\frac{\tan(fx+e)}{2}\right)}{f}$

3.681. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ict \tan(e + fx)) dx$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output `-I/f*a^2*c*(-1/3*B*tan(f*x+e)^3+1/2*(-A+I*B)*tan(f*x+e)^2+I*A*tan(f*x+e))`

3.681.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.53

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx)) dx$$

$$= -\frac{2(6(-iA - B)a^2ce^{4ifx+4ie} + 3(-3iA - B)a^2ce^{2ifx+2ie} + (-3iA - B)a^2c)}{3(fe^{6ifx+6ie} + 3fe^{4ifx+4ie} + 3fe^{2ifx+2ie} + f)}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x, algorithm="fricas")`

output `-2/3*(6*(-I*A - B)*a^2*c*e^(4*I*f*x + 4*I*e) + 3*(-3*I*A - B)*a^2*c*e^(2*I*f*x + 2*I*e) + (-3*I*A - B)*a^2*c)/(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)`

3.681.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 158 vs. 2(56) = 112.

Time = 0.21 (sec) , antiderivative size = 158, normalized size of antiderivative = 2.47

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx)) dx$$

$$= \frac{6iAa^2c + 2Ba^2c + (18iAa^2ce^{2ie} + 6Ba^2ce^{2ie})e^{2ifx} + (12iAa^2ce^{4ie} + 12Ba^2ce^{4ie})e^{4ifx}}{3fe^{6ie}e^{6ifx} + 9fe^{4ie}e^{4ifx} + 9fe^{2ie}e^{2ifx} + 3f}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x)`

output `(6*I*A*a**2*c + 2*B*a**2*c + (18*I*A*a**2*c*exp(2*I*e) + 6*B*a**2*c*exp(2*I*e))*exp(2*I*f*x) + (12*I*A*a**2*c*exp(4*I*e) + 12*B*a**2*c*exp(4*I*e))*exp(4*I*f*x))/(3*f*exp(6*I*e)*exp(6*I*f*x) + 9*f*exp(4*I*e)*exp(4*I*f*x) + 9*f*exp(2*I*e)*exp(2*I*f*x) + 3*f)`

3.681.7 Maxima [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.83

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx)) dx$$

$$= -\frac{-2i Ba^2 c \tan(fx + e)^3 - 3(iA + B)a^2 c \tan(fx + e)^2 - 6Aa^2 c \tan(fx + e)}{6f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x, algorithm="maxima")`

output `-1/6*(-2*I*B*a^2*c*tan(f*x + e)^3 - 3*(I*A + B)*a^2*c*tan(f*x + e)^2 - 6*A*a^2*c*tan(f*x + e))/f`

3.681.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 120 vs. 2(56) = 112.

Time = 0.44 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.88

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx)) dx =$$

$$-\frac{2(-6iAa^2ce^{(4i fx+4ie)} - 6Ba^2ce^{(4i fx+4ie)} - 9iAa^2ce^{(2i fx+2ie)} - 3Ba^2ce^{(2i fx+2ie)} - 3iAa^2c - Ba^2c)}{3(fe^{(6i fx+6ie)} + 3fe^{(4i fx+4ie)} + 3fe^{(2i fx+2ie)} + f)}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x, algorithm="giac")`

output `-2/3*(-6*I*A*a^2*c*e^(4*I*f*x + 4*I*e) - 6*B*a^2*c*e^(4*I*f*x + 4*I*e) - 9*I*A*a^2*c*e^(2*I*f*x + 2*I*e) - 3*B*a^2*c*e^(2*I*f*x + 2*I*e) - 3*I*A*a^2*c - B*a^2*c)/(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)`

3.681.9 Mupad [B] (verification not implemented)

Time = 8.33 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.78

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ic \tan(e + fx)) dx$$

$$= \frac{a^2 c \tan(e + fx) (6A + A \tan(e + fx) 3i + 3B \tan(e + fx) + B \tan(e + fx)^2 2i)}{6f}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i),x)`

output `(a^2*c*tan(e + f*x)*(6*A + A*tan(e + f*x)*3i + 3*B*tan(e + f*x) + B*tan(e + f*x)^2*2i))/(6*f)`

3.682 $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) dx$

3.682.1 Optimal result	6407
3.682.2 Mathematica [A] (verified)	6407
3.682.3 Rubi [A] (verified)	6408
3.682.4 Maple [A] (verified)	6409
3.682.5 Fricas [A] (verification not implemented)	6410
3.682.6 Sympy [A] (verification not implemented)	6410
3.682.7 Maxima [A] (verification not implemented)	6411
3.682.8 Giac [B] (verification not implemented)	6411
3.682.9 Mupad [B] (verification not implemented)	6412

3.682.1 Optimal result

Integrand size = 26, antiderivative size = 80

$$\begin{aligned} & \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) dx \\ &= 2a^2(A - iB)x - \frac{2a^2(iA + B) \log(\cos(e + fx))}{f} \\ & \quad - \frac{a^2(A - iB) \tan(e + fx)}{f} + \frac{B(a + ia \tan(e + fx))^2}{2f} \end{aligned}$$

output `2*a^2*(A-I*B)*x-2*a^2*(I*A+B)*ln(cos(f*x+e))/f-a^2*(A-I*B)*tan(f*x+e)/f+1/2*B*(a+I*a*tan(f*x+e))^2/f`

3.682.2 Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.72

$$\begin{aligned} & \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) dx \\ &= \frac{a^2(B + 4(iA + B) \log(i + \tan(e + fx)) - 2(A - 2iB) \tan(e + fx) - B \tan^2(e + fx))}{2f} \end{aligned}$$

input `Integrate[(a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]),x]`

output `(a^2*(B + 4*(I*A + B)*Log[I + Tan[e + f*x]] - 2*(A - (2*I)*B)*Tan[e + f*x] - B*Tan[e + f*x]^2))/(2*f)`

3.682.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.88, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3042, 4010, 3042, 3958, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{4010} \\
 & (A - iB) \int (i \tan(e + fx)a + a)^2 dx + \frac{B(a + ia \tan(e + fx))^2}{2f} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \int (i \tan(e + fx)a + a)^2 dx + \frac{B(a + ia \tan(e + fx))^2}{2f} \\
 & \quad \downarrow \text{3958} \\
 & (A - iB) \left(2ia^2 \int \tan(e + fx) dx - \frac{a^2 \tan(e + fx)}{f} + 2a^2 x \right) + \frac{B(a + ia \tan(e + fx))^2}{2f} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \left(2ia^2 \int \tan(e + fx) dx - \frac{a^2 \tan(e + fx)}{f} + 2a^2 x \right) + \frac{B(a + ia \tan(e + fx))^2}{2f} \\
 & \quad \downarrow \text{3956} \\
 & (A - iB) \left(-\frac{a^2 \tan(e + fx)}{f} - \frac{2ia^2 \log(\cos(e + fx))}{f} + 2a^2 x \right) + \frac{B(a + ia \tan(e + fx))^2}{2f}
 \end{aligned}$$

input `Int[(a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]),x]`

output `(B*(a + I*a*Tan[e + f*x])^2)/(2*f) + (A - I*B)*(2*a^2*x - ((2*I)*a^2*Log[Cos[e + f*x]])/f - (a^2*Tan[e + f*x])/f)`

3.682.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 3958 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^2, x_Symbol] := Simp[(a^2 - b^2)*x, x] + (Simp[b^2*(Tan[c + d*x]/d), x] + Simp[2*a*b Int[Tan[c + d*x], x], x]) /; FreeQ[{a, b, c, d}, x]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^(m)/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

3.682.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.95

method	result
derivativedivides	$\frac{a^2 \left(-\frac{B \tan^2(fx+e)}{2} - A \tan(fx+e) + 2i \tan(fx+e) B + \frac{(2iA+2B) \ln\left(\frac{1+\tan(fx+e)}{2}\right)}{2} + (-2iB+2A) \arctan(\tan(fx+e)) \right)}{f}$
default	$\frac{a^2 \left(-\frac{B \tan^2(fx+e)}{2} - A \tan(fx+e) + 2i \tan(fx+e) B + \frac{(2iA+2B) \ln\left(\frac{1+\tan(fx+e)}{2}\right)}{2} + (-2iB+2A) \arctan(\tan(fx+e)) \right)}{f}$
norman	$(-2iB a^2 + 2A a^2) x - \frac{(-2iB a^2 + A a^2) \tan(fx+e)}{f} - \frac{B a^2 \tan^2(fx+e)}{2f} + \frac{(iA a^2 + B a^2) \ln(1+\tan(fx+e))}{f}$
parallelrisch	$\frac{-4iBx a^2 f + 2iA \ln(1+\tan(fx+e))^2 a^2 + 4Ax a^2 f + 4iB \tan(fx+e) a^2 - B \tan^2(fx+e) a^2 - 2A \tan(fx+e) a^2 + 2B \ln(1+\tan(fx+e))}{2f}$
parts	$A a^2 x + \frac{(2iA a^2 + B a^2) \ln(1+\tan(fx+e))}{2f} + \frac{(2iB a^2 - A a^2) (\tan(fx+e) - \arctan(\tan(fx+e)))}{f} - \frac{B a^2 \left(\frac{\tan(fx+e)}{2} \right)}{f}$
risch	$\frac{4ia^2 B e}{f} - \frac{4a^2 A e}{f} - \frac{2a^2 (iA e^{2i(fx+e)} + 3B e^{2i(fx+e)} + iA + 2B)}{f(e^{2i(fx+e)} + 1)^2} - \frac{2a^2 \ln(e^{2i(fx+e)} + 1) B}{f} - \frac{2ia^2 \ln(e^{2i(fx+e)} + 1)}{f}$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e)),x,method=_RETURNVERBOSE)`

3.682. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) dx$

output $1/f*a^2*(-1/2*B*\tan(f*x+e)^2-A*\tan(f*x+e)+2*I*\tan(f*x+e)*B+1/2*(2*B+2*I*A)*\ln(1+\tan(f*x+e)^2)+(2*A-2*I*B)*\arctan(\tan(f*x+e)))$

3.682.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.51

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) dx = \frac{2((iA + 3B)a^2 e^{(2i fx + 2ie)} + (iA + 2B)a^2 + ((iA + B)a^2 e^{(4i fx + 4ie)} + 2(iA + B)a^2 e^{(2i fx + 2ie)} + (iA + B)a^2 e^{(4i fx + 4ie)} + 2fe^{(2i fx + 2ie)} + f)}{fe^{(4i fx + 4ie)} + 2fe^{(2i fx + 2ie)} + f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e)),x, algorithm="fricas")`

output $-2*((I*A + 3*B)*a^2*e^{(2*I*f*x + 2*I*e)} + (I*A + 2*B)*a^2 + ((I*A + B)*a^2*e^{(4*I*f*x + 4*I*e)} + 2*(I*A + B)*a^2*e^{(2*I*f*x + 2*I*e)} + (I*A + B)*a^2)*\log(e^{(2*I*f*x + 2*I*e)} + 1))/(f*e^{(4*I*f*x + 4*I*e)} + 2*f*e^{(2*I*f*x + 2*I*e)} + f)$

3.682.6 Sympy [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.52

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) dx = -\frac{2ia^2(A - iB) \log(e^{2ifx} + e^{-2ie})}{f} + \frac{-2iAa^2 - 4Ba^2 + (-2iAa^2 e^{2ie} - 6Ba^2 e^{2ie}) e^{2ifx}}{fe^{4ie} e^{4ifx} + 2fe^{2ie} e^{2ifx} + f}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e)),x)`

output $-2*I*a**2*(A - I*B)*\log(\exp(2*I*f*x) + \exp(-2*I*e))/f + (-2*I*A*a**2 - 4*B*a**2 + (-2*I*A*a**2*\exp(2*I*e) - 6*B*a**2*\exp(2*I*e))*\exp(2*I*f*x))/(f*\exp(4*I*e)*\exp(4*I*f*x) + 2*f*\exp(2*I*e)*\exp(2*I*f*x) + f)$

3.682.7 Maxima [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.89

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) dx = \frac{Ba^2 \tan(fx + e)^2 - 4(fx + e)(A - iB)a^2 - 2(iA + B)a^2 \log(\tan(fx + e)^2 + 1) + 2(A - 2iB)a^2 \tan(fx + e)}{2f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e)),x, algorithm="maxima")`

output `-1/2*(B*a^2*tan(f*x + e)^2 - 4*(f*x + e)*(A - I*B)*a^2 - 2*(I*A + B)*a^2*log(tan(f*x + e)^2 + 1) + 2*(A - 2*I*B)*a^2*tan(f*x + e))/f`

3.682.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 214 vs. 2(70) = 140.

Time = 0.41 (sec) , antiderivative size = 214, normalized size of antiderivative = 2.68

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) dx = \frac{2(iAa^2 e^{(4i fx + 4i e)} \log(e^{(2i fx + 2i e)} + 1) + Ba^2 e^{(4i fx + 4i e)} \log(e^{(2i fx + 2i e)} + 1) + 2iAa^2 e^{(2i fx + 2i e)} \log(e^{(2i fx + 2i e)} + 1) + 2iAa^2 e^{(4i fx + 4i e)} \log(e^{(2i fx + 2i e)} + 1) + 2iBa^2 e^{(4i fx + 4i e)} \log(e^{(2i fx + 2i e)} + 1) + 2iBa^2 e^{(2i fx + 2i e)} \log(e^{(2i fx + 2i e)} + 1) + I*A*a^2 * e^{(4*I*f*x + 4*I*e)} * \log(e^{(2*I*f*x + 2*I*e)} + 1) + B*a^2 * e^{(4*I*f*x + 4*I*e)} * \log(e^{(2*I*f*x + 2*I*e)} + 1) + 2*I*A*a^2 * e^{(2*I*f*x + 2*I*e)} * \log(e^{(2*I*f*x + 2*I*e)} + 1) + 2*B*a^2 * e^{(2*I*f*x + 2*I*e)} * \log(e^{(2*I*f*x + 2*I*e)} + 1) + I*A*a^2 * e^{(2*I*f*x + 2*I*e)} + 3*B*a^2 * e^{(2*I*f*x + 2*I*e)} + I*A*a^2 * \log(e^{(2*I*f*x + 2*I*e)} + 1) + B*a^2 * \log(e^{(2*I*f*x + 2*I*e)} + 1) + I*A*a^2 + 2*B*a^2) / (f * e^{(4*I*f*x + 4*I*e)} + 2*f * e^{(2*I*f*x + 2*I*e)} + f)$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e)),x, algorithm="giac")`

output `-2*(I*A*a^2*e^(4*I*f*x + 4*I*e)*log(e^(2*I*f*x + 2*I*e) + 1) + B*a^2*e^(4*I*f*x + 4*I*e)*log(e^(2*I*f*x + 2*I*e) + 1) + 2*I*A*a^2*e^(2*I*f*x + 2*I*e)*log(e^(2*I*f*x + 2*I*e) + 1) + 2*B*a^2*e^(2*I*f*x + 2*I*e)*log(e^(2*I*f*x + 2*I*e) + 1) + I*A*a^2*e^(2*I*f*x + 2*I*e) + 3*B*a^2*e^(2*I*f*x + 2*I*e) + I*A*a^2*log(e^(2*I*f*x + 2*I*e) + 1) + B*a^2*log(e^(2*I*f*x + 2*I*e) + 1) + I*A*a^2 + 2*B*a^2)/(f*e^(4*I*f*x + 4*I*e) + 2*f*e^(2*I*f*x + 2*I*e) + f)`

3.682.9 Mupad [B] (verification not implemented)

Time = 8.43 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.95

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) dx$$

$$= \frac{\ln(\tan(e + fx) + 1i) (2B a^2 + A a^2 2i)}{f}$$

$$+ \frac{\tan(e + fx) (a^2 (B + A 1i) 1i + B a^2 1i)}{f} - \frac{B a^2 \tan(e + fx)^2}{2f}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2,x)`output `(log(tan(e + f*x) + 1i)*(A*a^2*2i + 2*B*a^2))/f + (tan(e + f*x)*(a^2*(A*1i + B)*1i + B*a^2*1i))/f - (B*a^2*tan(e + f*x)^2)/(2*f)`

3.683
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{c-ictan(e+fx)} dx$$

3.683.1 Optimal result 6413
 3.683.2 Mathematica [A] (verified) 6413
 3.683.3 Rubi [A] (verified) 6414
 3.683.4 Maple [A] (verified) 6416
 3.683.5 Fricas [A] (verification not implemented) 6416
 3.683.6 Sympy [A] (verification not implemented) 6417
 3.683.7 Maxima [F(-2)] 6417
 3.683.8 Giac [B] (verification not implemented) 6417
 3.683.9 Mupad [B] (verification not implemented) 6418

3.683.1 Optimal result

Integrand size = 41, antiderivative size = 93

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{c - ictan(e + fx)} dx$$

$$= -\frac{a^2(A - 3iB)x}{c} + \frac{a^2(iA + 3B) \log(\cos(e + fx))}{cf}$$

$$- \frac{ia^2B \tan(e + fx)}{cf} + \frac{2a^2(A - iB)}{cf(i + \tan(e + fx))}$$

output `-a^2*(A-3*I*B)*x/c+a^2*(I*A+3*B)*ln(cos(f*x+e))/c/f-I*a^2*B*tan(f*x+e)/c/f+2*a^2*(A-I*B)/c/f/(I+tan(f*x+e))`

3.683.2 Mathematica [A] (verified)

Time = 5.53 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.90

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{c - ictan(e + fx)} dx$$

$$= \frac{B(a + ia \tan(e + fx))^2}{f(c - ictan(e + fx))} - \frac{a^2(iA + 3B) \left(\log(i + \tan(e + fx)) + \frac{2i}{i + \tan(e + fx)} \right)}{cf}$$

input `Integrate[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x]),x]`

output $(B*(a + I*a*\text{Tan}[e + f*x])^2)/(f*(c - I*c*\text{Tan}[e + f*x])) - (a^2*(I*A + 3*B) * (\text{Log}[I + \text{Tan}[e + f*x]] + (2*I)/(I + \text{Tan}[e + f*x])))/(c*f)$

3.683.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.70, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{a(i \tan(e+fx)+1)(A+B \tan(e+fx))}{c^2(1-i \tan(e+fx))^2} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{a^2 \int \frac{(i \tan(e+fx)+1)(A+B \tan(e+fx))}{(1-i \tan(e+fx))^2} d \tan(e + fx)}{cf} \\ & \quad \downarrow \text{86} \\ & \frac{a^2 \int \left(-\frac{2(A-iB)}{(\tan(e+fx)+i)^2} - iB - \frac{i(A-3iB)}{\tan(e+fx)+i} \right) d \tan(e + fx)}{cf} \\ & \quad \downarrow \text{2009} \\ & \frac{a^2 \left(\frac{2(A-iB)}{\tan(e+fx)+i} - (3B + iA) \log(\tan(e + fx) + i) - iB \tan(e + fx) \right)}{cf} \end{aligned}$$

input $\text{Int}[\frac{(a + I*a*\text{Tan}[e + f*x])^2*(A + B*\text{Tan}[e + f*x])}{(c - I*c*\text{Tan}[e + f*x])}, x]$

output $(a^2 * (-((I * A + 3 * B) * \text{Log}[I + \text{Tan}[e + f * x]]) - I * B * \text{Tan}[e + f * x] + (2 * (A - I * B)) / (I + \text{Tan}[e + f * x]))) / (c * f)$

3.683.3.1 Defintions of rubi rules used

rule 27 $\text{Int}[(a_)*(F_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[F, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F, (b_)*(G_)] /; \text{FreeQ}[b, x]$

rule 86 $\text{Int}[(a_ + (b_)*(x_))*((c_ + (d_)*(x_))^{(n_)}*(e_ + (f_)*(x_))^{(p_)}), x_] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \ \&\& \ ((\text{ILtQ}[n, 0] \ \&\& \ \text{ILtQ}[p, 0]) \ || \ \text{EqQ}[p, 1] \ || \ (\text{IGtQ}[p, 0] \ \&\& \ (!\text{IntegerQ}[n] \ || \ \text{LeQ}[9*p + 5*(n + 2), 0] \ || \ \text{GeQ}[n + p + 1, 0] \ || \ (\text{GeQ}[n + p + 2, 0] \ \&\& \ \text{RationalQ}[a, b, c, d, e, f])))$

rule 2009 $\text{Int}[u_, x_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4071 $\text{Int}[(a_ + (b_)*\text{tan}[(e_ + (f_)*(x_)]))^{(m_)}*((A_ + (B_)*\text{tan}[(e_ + (f_)*(x_)]))^{(n_)}*(c_ + (d_)*\text{tan}[(e_ + (f_)*(x_)]))^{(n_)}), x_Symbol] \rightarrow \text{Simp}[a*(c/f) \ \text{Subst}[\text{Int}[(a + b*x)^{(m-1)}*(c + d*x)^{(n-1)}*(A + B*x), x], x, \text{Tan}[e + f*x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f, A, B, m, n\}, x] \ \&\& \ \text{EqQ}[b*c + a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0]$

3.683.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.57

method	result
risch	$-\frac{e^{2i(fx+e)}a^2B}{cf} - \frac{ie^{2i(fx+e)}Aa^2}{cf} - \frac{6ia^2Be}{cf} + \frac{2a^2Ae}{cf} + \frac{2a^2B}{fc(e^{2i(fx+e)}+1)} + \frac{3a^2 \ln(e^{2i(fx+e)}+1)B}{cf} + \frac{ia^2 \ln(e^{2i(fx+e)}+1)A}{cf}$
derivativedivides	$-\frac{ia^2B \tan(fx+e)}{cf} - \frac{2ia^2B}{fc(i+\tan(fx+e))} + \frac{2a^2A}{fc(i+\tan(fx+e))} - \frac{ia^2A \ln(1+\tan(fx+e)^2)}{2fc} - \frac{3a^2B \ln(1+\tan(fx+e)^2)}{2fc}$
default	$-\frac{ia^2B \tan(fx+e)}{cf} - \frac{2ia^2B}{fc(i+\tan(fx+e))} + \frac{2a^2A}{fc(i+\tan(fx+e))} - \frac{ia^2A \ln(1+\tan(fx+e)^2)}{2fc} - \frac{3a^2B \ln(1+\tan(fx+e)^2)}{2fc}$
norman	$\frac{(-3iB a^2+2A a^2) \tan(fx+e)}{cf} - \frac{(-3iB a^2+A a^2)x}{c} - \frac{2iA a^2+2B a^2}{cf} - \frac{(-3iB a^2+A a^2)x \tan(fx+e)^2}{c} - \frac{iB a^2 \tan(fx+e)^3}{cf} - (iA a^2) \ln(1+\tan(fx+e)^2)$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output $-1/c/f*\exp(2*I*(f*x+e))*a^2*B-I/c/f*\exp(2*I*(f*x+e))*A*a^2-6*I*a^2/c/f*B*e+2*a^2/c/f*A*e+2/f/c*a^2*B/(\exp(2*I*(f*x+e))+1)+3*a^2/c/f*\ln(\exp(2*I*(f*x+e))+1)*B+I*a^2/c/f*\ln(\exp(2*I*(f*x+e))+1)*A$

3.683.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.19

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx$$

$$= \frac{(-iA - B)a^2e^{(4i fx+4i e)} + (-iA - B)a^2e^{(2i fx+2i e)} + 2Ba^2 + ((iA + 3B)a^2e^{(2i fx+2i e)} + (iA + 3B)a^2) \ln(e^{(2i fx+2i e)} + 1)}{cfe^{(2i fx+2i e)} + cf}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x, algorithm="fracas")`

output $((-I*A - B)*a^2*e^{(4*I*f*x + 4*I*e)} + (-I*A - B)*a^2*e^{(2*I*f*x + 2*I*e)} + 2*B*a^2 + ((I*A + 3*B)*a^2*e^{(2*I*f*x + 2*I*e)} + (I*A + 3*B)*a^2)*\log(e^{(2*I*f*x + 2*I*e)} + 1))/(c*f*e^{(2*I*f*x + 2*I*e)} + c*f)$

3.683.6 Sympy [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.44

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx$$

$$= \frac{2Ba^2}{cfe^{2ie}e^{2ifx} + cf} + \frac{ia^2(A - 3iB) \log(e^{2ifx} + e^{-2ie})}{cf} + \begin{cases} \frac{(-iAa^2e^{2ie} - Ba^2e^{2ie})e^{2ifx}}{cf} & \text{for } cf \neq 0 \\ \frac{x(2Aa^2e^{2ie} - 2iBa^2e^{2ie})}{c} & \text{otherwise} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x)`

output `2*B*a**2/(c*f*exp(2*I*e)*exp(2*I*f*x) + c*f) + I*a**2*(A - 3*I*B)*log(exp(2*I*f*x) + exp(-2*I*e))/(c*f) + Piecewise(((-I*A*a**2*exp(2*I*e) - B*a**2*exp(2*I*e))*exp(2*I*f*x)/(c*f), Ne(c*f, 0)), (x*(2*A*a**2*exp(2*I*e) - 2*I*B*a**2*exp(2*I*e))/c, True))`

3.683.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x, algorith="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.683.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 271 vs. $2(83) = 166$.

Time = 0.47 (sec) , antiderivative size = 271, normalized size of antiderivative = 2.91

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx$$

$$= \frac{(iAa^2 + 3Ba^2) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)}{c} + \frac{2(-iAa^2 - 3Ba^2) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + i)}{c} - \frac{(-iAa^2 - 3Ba^2) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)}{c} - \frac{iAa^2 \tan(e + fx)}{c}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x, algorithm="giac")`

output `((I*A*a^2 + 3*B*a^2)*log(tan(1/2*f*x + 1/2*e) + 1)/c + 2*(-I*A*a^2 - 3*B*a^2)*log(tan(1/2*f*x + 1/2*e) + I)/c - (-I*A*a^2 - 3*B*a^2)*log(tan(1/2*f*x + 1/2*e) - 1)/c - (I*A*a^2*tan(1/2*f*x + 1/2*e)^2 + 3*B*a^2*tan(1/2*f*x + 1/2*e)^2 - 2*I*B*a^2*tan(1/2*f*x + 1/2*e) - I*A*a^2 - 3*B*a^2)/((tan(1/2*f*x + 1/2*e)^2 - 1)*c) - (-3*I*A*a^2*tan(1/2*f*x + 1/2*e)^2 - 9*B*a^2*tan(1/2*f*x + 1/2*e)^2 + 10*A*a^2*tan(1/2*f*x + 1/2*e) - 22*I*B*a^2*tan(1/2*f*x + 1/2*e) + 3*I*A*a^2 + 9*B*a^2)/(c*(tan(1/2*f*x + 1/2*e) + I)^2))/f`

3.683.9 Mupad [B] (verification not implemented)

Time = 8.40 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.13

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx$$

$$= -\frac{\ln(\tan(e + fx) + 1i) \left(\frac{3Ba^2}{c} + \frac{Aa^2 1i}{c} \right)}{f} + \frac{\frac{Aa^2 + Ba^2 1i}{c} + \frac{Aa^2 - Ba^2 3i}{c}}{f(\tan(e + fx) + 1i)} - \frac{Ba^2 \tan(e + fx) 1i}{cf}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2)/(c - c*tan(e + f*x)*1i),x)`

output `((A*a^2 + B*a^2*1i)/c + (A*a^2 - B*a^2*3i)/c)/(f*(tan(e + f*x) + 1i)) - (log(tan(e + f*x) + 1i)*((A*a^2*1i)/c + (3*B*a^2)/c))/f - (B*a^2*tan(e + f*x)*1i)/(c*f)`

3.684
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^2} dx$$

3.684.1 Optimal result	6419
3.684.2 Mathematica [A] (verified)	6419
3.684.3 Rubi [A] (verified)	6420
3.684.4 Maple [A] (verified)	6421
3.684.5 Fricas [A] (verification not implemented)	6422
3.684.6 Sympy [A] (verification not implemented)	6422
3.684.7 Maxima [F(-2)]	6423
3.684.8 Giac [B] (verification not implemented)	6423
3.684.9 Mupad [B] (verification not implemented)	6424

3.684.1 Optimal result

Integrand size = 41, antiderivative size = 91

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx$$

$$= -\frac{ia^2 Bx}{c^2} - \frac{a^2 B \log(\cos(e + fx))}{c^2 f} + \frac{a^2(iA + B)}{c^2 f(i + \tan(e + fx))^2} - \frac{a^2(A - 3iB)}{c^2 f(i + \tan(e + fx))}$$

output `-I*a^2*B*x/c^2-a^2*B*ln(cos(f*x+e))/c^2/f+a^2*(I*A+B)/c^2/f/(I+tan(f*x+e))
^2-a^2*(A-3*I*B)/c^2/f/(I+tan(f*x+e))`

3.684.2 Mathematica [A] (verified)

Time = 4.53 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.62

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx$$

$$= \frac{a^2 \left(B \log(i + \tan(e + fx)) - \frac{2B+(A-3iB) \tan(e+fx)}{(i+\tan(e+fx))^2} \right)}{c^2 f}$$

input `Integrate[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^2,x]`

output `(a^2*(B*Log[I + Tan[e + f*x]] - (2*B + (A - (3*I)*B)*Tan[e + f*x])/(I + Tan[e + f*x]^2))/(c^2*f)`

3.684.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.71, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{a(i \tan(e+fx)+1)(A+B \tan(e+fx))}{c^3(1-i \tan(e+fx))^3} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^2 \int \frac{(i \tan(e+fx)+1)(A+B \tan(e+fx))}{(1-i \tan(e+fx))^3} d \tan(e + fx)}{c^2 f} \\
 & \quad \downarrow \text{86} \\
 & \frac{a^2 \int \left(-\frac{2i(A-iB)}{(\tan(e+fx)+i)^3} + \frac{B}{\tan(e+fx)+i} + \frac{A-3iB}{(\tan(e+fx)+i)^2} \right) d \tan(e + fx)}{c^2 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^2 \left(-\frac{A-3iB}{\tan(e+fx)+i} + \frac{B+iA}{(\tan(e+fx)+i)^2} + B \log(\tan(e + fx) + i) \right)}{c^2 f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^2,x]`

output `(a^2*(B*Log[I + Tan[e + f*x]] + (I*A + B)/(I + Tan[e + f*x])^2 - (A - (3*I)*B)/(I + Tan[e + f*x]))/(c^2*f)`

3.684.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.684.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.13

method	result
risch	$-\frac{e^{4i(fx+e)}a^2B}{4c^2f} - \frac{ie^{4i(fx+e)}Aa^2}{4c^2f} + \frac{Ba^2e^{2i(fx+e)}}{c^2f} + \frac{2iBa^2e}{c^2f} - \frac{Ba^2 \ln(e^{2i(fx+e)}+1)}{c^2f}$
derivativedivides	$\frac{ia^2A}{fc^2(i+\tan(fx+e))^2} + \frac{a^2B}{fc^2(i+\tan(fx+e))^2} + \frac{3ia^2B}{fc^2(i+\tan(fx+e))} - \frac{a^2A}{fc^2(i+\tan(fx+e))} + \frac{a^2B \ln(1+\tan(fx+e))}{2fc^2}$
default	$\frac{ia^2A}{fc^2(i+\tan(fx+e))^2} + \frac{a^2B}{fc^2(i+\tan(fx+e))^2} + \frac{3ia^2B}{fc^2(i+\tan(fx+e))} - \frac{a^2A}{fc^2(i+\tan(fx+e))} + \frac{a^2B \ln(1+\tan(fx+e))}{2fc^2}$

```
input int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x,method=_R
ETURNVERBOSE)
```

3.684.
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^2} dx$$

output
$$-1/4/c^2/f*\exp(4*I*(f*x+e))*a^2*B-1/4*I/c^2/f*\exp(4*I*(f*x+e))*A*a^2+B*a^2/c^2/f*\exp(2*I*(f*x+e))+2*I*B*a^2/c^2/f*e-B*a^2/c^2/f*\ln(\exp(2*I*(f*x+e))+1)$$

3.684.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.68

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx$$

$$= \frac{(-iA - B)a^2 e^{(4i fx + 4ie)} + 4Ba^2 e^{(2i fx + 2ie)} - 4Ba^2 \log(e^{(2i fx + 2ie)} + 1)}{4c^2 f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x, algorithm="fricas")`

output
$$1/4*((-I*A - B)*a^2*e^{(4*I*f*x + 4*I*e)} + 4*B*a^2*e^{(2*I*f*x + 2*I*e)} - 4*B*a^2*\log(e^{(2*I*f*x + 2*I*e)} + 1))/(c^2*f)$$

3.684.6 Sympy [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.76

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx$$

$$= -\frac{Ba^2 \log(e^{2ifx} + e^{-2ie})}{c^2 f} + \begin{cases} \frac{4Ba^2 c^2 f e^{2ie} e^{2ifx} + (-iAa^2 c^2 f e^{4ie} - Ba^2 c^2 f e^{4ie}) e^{4ifx}}{4c^4 f^2} & \text{for } c^4 f^2 \neq 0 \\ \frac{x(Aa^2 e^{4ie} - iBa^2 e^{4ie} + 2iBa^2 e^{2ie})}{c^2} & \text{otherwise} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**2,x)`

output
$$-B*a**2*\log(\exp(2*I*f*x) + \exp(-2*I*e))/(c**2*f) + \text{Piecewise}(((4*B*a**2*c**2*f*\exp(2*I*e)*\exp(2*I*f*x) + (-I*A*a**2*c**2*f*\exp(4*I*e) - B*a**2*c**2*f*\exp(4*I*e))*\exp(4*I*f*x))/(4*c**4*f**2), \text{Ne}(c**4*f**2, 0)), (x*(A*a**2*\exp(4*I*e) - I*B*a**2*\exp(4*I*e) + 2*I*B*a**2*\exp(2*I*e))/c**2, \text{True}))$$

3.684.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.684.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 191 vs. $2(81) = 162$.

Time = 0.54 (sec) , antiderivative size = 191, normalized size of antiderivative = 2.10

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx =$$

$$\frac{6Ba^2 \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)}{c^2} - \frac{12Ba^2 \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + i)}{c^2} + \frac{6Ba^2 \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)}{c^2} + \frac{25Ba^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^4 + 12Aa^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 - 198Ba^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 12Aa^2 \tan(\frac{1}{2}fx + \frac{1}{2}e) - 112I*Ba^2 \tan(\frac{1}{2}fx + \frac{1}{2}e) + 25Ba^2}{(c^2 * (\tan(\frac{1}{2}fx + \frac{1}{2}e) + I)^4)} / f$$

6 f

```
input integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x, algorithm="giac")
```

```
output -1/6*(6*B*a^2*log(tan(1/2*f*x + 1/2*e) + 1)/c^2 - 12*B*a^2*log(tan(1/2*f*x + 1/2*e) + I)/c^2 + 6*B*a^2*log(tan(1/2*f*x + 1/2*e) - 1)/c^2 + (25*B*a^2 *tan(1/2*f*x + 1/2*e)^4 + 12*A*a^2*tan(1/2*f*x + 1/2*e)^3 + 112*I*B*a^2*tan(1/2*f*x + 1/2*e)^3 - 198*B*a^2*tan(1/2*f*x + 1/2*e)^2 - 12*A*a^2*tan(1/2*f*x + 1/2*e) - 112*I*B*a^2*tan(1/2*f*x + 1/2*e) + 25*B*a^2)/(c^2*(tan(1/2*f*x + 1/2*e) + I)^4))/f
```

3.684.9 Mupad [B] (verification not implemented)

Time = 8.44 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.14

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx =$$

$$\frac{a^2 (B 2i + A \tan(e + fx) 1i + 3 B \tan(e + fx) + B \ln(\tan(e + fx) + 1i) 1i + 2 B \ln(\tan(e + fx) + 1i))}{c^2 f (-1 + \tan(e + fx) 1i)^2}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2)/(c - c*tan(e + f*x)*1i)^2,x)`

output `-(a^2*(B*2i + A*tan(e + f*x)*1i + 3*B*tan(e + f*x) + B*log(tan(e + f*x) + 1i)*1i + 2*B*log(tan(e + f*x) + 1i)*tan(e + f*x) - B*log(tan(e + f*x) + 1i)*tan(e + f*x)^2*1i*1i)/(c^2*f*(tan(e + f*x)*1i - 1)^2)`

3.685
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^3} dx$$

3.685.1 Optimal result	6425
3.685.2 Mathematica [A] (verified)	6425
3.685.3 Rubi [A] (verified)	6426
3.685.4 Maple [A] (verified)	6427
3.685.5 Fricas [A] (verification not implemented)	6428
3.685.6 Sympy [B] (verification not implemented)	6428
3.685.7 Maxima [F(-2)]	6429
3.685.8 Giac [B] (verification not implemented)	6429
3.685.9 Mupad [B] (verification not implemented)	6430

3.685.1 Optimal result

Integrand size = 41, antiderivative size = 93

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx$$

$$= -\frac{2a^2(A - iB)}{3c^3 f(i + \tan(e + fx))^3} - \frac{a^2(iA + 3B)}{2c^3 f(i + \tan(e + fx))^2} - \frac{ia^2 B}{c^3 f(i + \tan(e + fx))}$$

output
$$-2/3*a^2*(A-I*B)/c^3/f/(I+\tan(f*x+e))^3-1/2*a^2*(I*A+3*B)/c^3/f/(I+\tan(f*x+e))^2-I*a^2*B/c^3/f/(I+\tan(f*x+e))$$

3.685.2 Mathematica [A] (verified)

Time = 3.94 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.67

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx$$

$$= \frac{a^2(-A + iB + 3(-iA + B) \tan(e + fx) - 6iB \tan^2(e + fx))}{6c^3 f(i + \tan(e + fx))^3}$$

input `Integrate[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^3,x]`

output
$$(a^2*(-A + I*B + 3*((-I)*A + B)*Tan[e + f*x] - (6*I)*B*Tan[e + f*x]^2))/(6*c^3*f*(I + Tan[e + f*x])^3)$$

3.685.
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^3} dx$$

3.685.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.82, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{a(i \tan(e+fx)+1)(A+B \tan(e+fx))}{c^4(1-i \tan(e+fx))^4} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^2 \int \frac{(i \tan(e+fx)+1)(A+B \tan(e+fx))}{(1-i \tan(e+fx))^4} d \tan(e + fx)}{c^3 f} \\
 & \quad \downarrow \text{86} \\
 & \frac{a^2 \int \left(\frac{2(A-iB)}{(\tan(e+fx)+i)^4} + \frac{iB}{(\tan(e+fx)+i)^2} + \frac{iA+3B}{(\tan(e+fx)+i)^3} \right) d \tan(e + fx)}{c^3 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^2 \left(-\frac{2(A-iB)}{3(\tan(e+fx)+i)^3} - \frac{3B+iA}{2(\tan(e+fx)+i)^2} - \frac{iB}{\tan(e+fx)+i} \right)}{c^3 f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^3,x]`

output `(a^2*((-2*(A - I*B))/(3*(I + Tan[e + f*x])^3) - (I*A + 3*B)/(2*(I + Tan[e + f*x])^2) - (I*B)/(I + Tan[e + f*x]))/(c^3*f)`

3.685.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.685.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.74

method	result
derivativedivides	$\frac{a^2 \left(-\frac{iA+3B}{2(i+\tan(fx+e))^2} - \frac{iB}{i+\tan(fx+e)} - \frac{-2iB+2A}{3(i+\tan(fx+e))^3} \right)}{f c^3}$
default	$\frac{a^2 \left(-\frac{iA+3B}{2(i+\tan(fx+e))^2} - \frac{iB}{i+\tan(fx+e)} - \frac{-2iB+2A}{3(i+\tan(fx+e))^3} \right)}{f c^3}$
risch	$-\frac{a^2 e^{6i(fx+e)} B}{12c^3 f} - \frac{ia^2 e^{6i(fx+e)} A}{12c^3 f} + \frac{a^2 e^{4i(fx+e)} B}{8c^3 f} - \frac{ia^2 e^{4i(fx+e)} A}{8c^3 f}$
norman	$\frac{\frac{2iA a^2 \tan(fx+e)^2}{cf} + \frac{A a^2 \tan(fx+e)}{cf} - \frac{iA a^2 + B a^2}{6cf} - \frac{5(-iB a^2 + A a^2) \tan(fx+e)^3}{3cf} - \frac{(iA a^2 + 5B a^2) \tan(fx+e)^4}{2cf} - \frac{iB a^2 \tan(fx+e)}{cf}}{c^2 (1+\tan(fx+e))^2}$

3.685.
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^3} dx$$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x,method=_RETURVERBOSE)`

output `1/f*a^2/c^3*(-1/2*(I*A+3*B)/(I+tan(f*x+e))^2-I*B/(I+tan(f*x+e))-1/3*(2*A-2*I*B)/(I+tan(f*x+e))^3)`

3.685.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.53

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^3} dx$$

$$= -\frac{2(iA + B)a^2 e^{(6i fx + 6ie)} + 3(iA - B)a^2 e^{(4i fx + 4ie)}}{24c^3 f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x, algorithm="fricas")`

output `-1/24*(2*(I*A + B)*a^2*e^(6*I*f*x + 6*I*e) + 3*(I*A - B)*a^2*e^(4*I*f*x + 4*I*e))/(c^3*f)`

3.685.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 167 vs. $2(75) = 150$.

Time = 0.28 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.80

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^3} dx$$

$$= \begin{cases} \frac{(-12iAa^2c^3fe^{4ie} + 12Ba^2c^3fe^{4ie})e^{4ifx} + (-8iAa^2c^3fe^{6ie} - 8Ba^2c^3fe^{6ie})e^{6ifx}}{96c^6f^2} & \text{for } c^6f^2 \neq 0 \\ \frac{x(Aa^2e^{6ie} + Aa^2e^{4ie} - iBa^2e^{6ie} + iBa^2e^{4ie})}{2c^3} & \text{otherwise} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**3,x)`

```
output Piecewise(((((-12*I*A*a**2*c**3*f*exp(4*I*e) + 12*B*a**2*c**3*f*exp(4*I*e))
*exp(4*I*f*x) + (-8*I*A*a**2*c**3*f*exp(6*I*e) - 8*B*a**2*c**3*f*exp(6*I*e)
))*exp(6*I*f*x))/(96*c**6*f**2), Ne(c**6*f**2, 0)), (x*(A*a**2*exp(6*I*e)
+ A*a**2*exp(4*I*e) - I*B*a**2*exp(6*I*e) + I*B*a**2*exp(4*I*e))/(2*c**3),
True))
```

3.685.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x, al
gorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.685.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 156 vs. $2(77) = 154$.

Time = 0.71 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.68

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx =$$

$$\frac{2 \left(3 A a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^5 + 3 i A a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^4 - 3 B a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^4 - 8 A a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^3 + 2 i B a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^3 - 3 i A a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^2 + 3 B a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^2 + 3 A a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) \right)}{3 c^3 f (\tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) + i)^6}$$

```
input integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x, al
gorithm="giac")
```

```
output -2/3*(3*A*a^2*tan(1/2*f*x + 1/2*e)^5 + 3*I*A*a^2*tan(1/2*f*x + 1/2*e)^4 -
3*B*a^2*tan(1/2*f*x + 1/2*e)^4 - 8*A*a^2*tan(1/2*f*x + 1/2*e)^3 + 2*I*B*a^
2*tan(1/2*f*x + 1/2*e)^3 - 3*I*A*a^2*tan(1/2*f*x + 1/2*e)^2 + 3*B*a^2*tan(
1/2*f*x + 1/2*e)^2 + 3*A*a^2*tan(1/2*f*x + 1/2*e))/(c^3*f*(tan(1/2*f*x + 1
/2*e) + I)^6)
```

3.685. $\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^3} dx$

3.685.9 Mupad [B] (verification not implemented)

Time = 8.32 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.94

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx$$

$$= \frac{\frac{a^2(A - B1i)}{6} + \frac{a^2 \tan(e + fx)(-3B + A3i)}{6} + B a^2 \tan(e + fx)^2 1i}{c^3 f (-\tan(e + fx)^3 - \tan(e + fx)^2 3i + 3 \tan(e + fx) + 1i)}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2)/(c - c*tan(e + f*x)*1i)^3,x)`

output `((a^2*(A - B*1i))/6 + (a^2*tan(e + f*x)*(A*3i - 3*B))/6 + B*a^2*tan(e + f*x)^2*1i)/(c^3*f*(3*tan(e + f*x) - tan(e + f*x)^2*3i - tan(e + f*x)^3 + 1i))`

3.686
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^4} dx$$

3.686.1 Optimal result 6431
 3.686.2 Mathematica [A] (verified) 6431
 3.686.3 Rubi [A] (verified) 6432
 3.686.4 Maple [A] (verified) 6433
 3.686.5 Fricas [A] (verification not implemented) 6434
 3.686.6 Sympy [B] (verification not implemented) 6434
 3.686.7 Maxima [F(-2)] 6435
 3.686.8 Giac [B] (verification not implemented) 6435
 3.686.9 Mupad [B] (verification not implemented) 6436

3.686.1 Optimal result

Integrand size = 41, antiderivative size = 91

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx$$

$$= -\frac{a^2(iA + B)}{2c^4 f(i + \tan(e + fx))^4} + \frac{a^2(A - 3iB)}{3c^4 f(i + \tan(e + fx))^3} + \frac{a^2B}{2c^4 f(i + \tan(e + fx))^2}$$

output
$$-1/2*a^2*(I*A+B)/c^4/f/(I+\tan(f*x+e))^4+1/3*a^2*(A-3*I*B)/c^4/f/(I+\tan(f*x+e))^3+1/2*a^2*B/c^4/f/(I+\tan(f*x+e))^2$$

3.686.2 Mathematica [A] (verified)

Time = 5.30 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.56

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx$$

$$= \frac{a^2(-iA + 2A \tan(e + fx) + 3B \tan^2(e + fx))}{6c^4 f(i + \tan(e + fx))^4}$$

input `Integrate[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^4,x]`

output
$$(a^2*((-I)*A + 2*A*\tan[e + f*x] + 3*B*\tan[e + f*x]^2))/(6*c^4*f*(I + \tan[e + f*x])^4)$$

3.686.
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^4} dx$$

3.686.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.81, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx$$

↓ 4071

$$\frac{ac \int \frac{a(i \tan(e+fx)+1)(A+B \tan(e+fx))}{c^5(1-i \tan(e+fx))^5} d \tan(e + fx)}{f}$$

↓ 27

$$\frac{a^2 \int \frac{(i \tan(e+fx)+1)(A+B \tan(e+fx))}{(1-i \tan(e+fx))^5} d \tan(e + fx)}{c^4 f}$$

↓ 86

$$\frac{a^2 \int \left(\frac{3iB-A}{(\tan(e+fx)+i)^4} - \frac{B}{(\tan(e+fx)+i)^3} + \frac{2(iA+B)}{(\tan(e+fx)+i)^5} \right) d \tan(e + fx)}{c^4 f}$$

↓ 2009

$$\frac{a^2 \left(\frac{A-3iB}{3(\tan(e+fx)+i)^3} - \frac{B+iA}{2(\tan(e+fx)+i)^4} + \frac{B}{2(\tan(e+fx)+i)^2} \right)}{c^4 f}$$

input `Int[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^4,x]`

output `(a^2*(-1/2*(I*A + B)/(I + Tan[e + f*x])^4 + (A - (3*I)*B)/(3*(I + Tan[e + f*x])^3) + B/(2*(I + Tan[e + f*x])^2))/(c^4*f)`

3.686.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.686.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.75

method	result	size
derivativedivides	$\frac{a^2 \left(-\frac{2iA+2B}{4(i+\tan(fx+e))^4} - \frac{3iB-A}{3(i+\tan(fx+e))^3} + \frac{B}{2(i+\tan(fx+e))^2} \right)}{f c^4}$	68
default	$\frac{a^2 \left(-\frac{2iA+2B}{4(i+\tan(fx+e))^4} - \frac{3iB-A}{3(i+\tan(fx+e))^3} + \frac{B}{2(i+\tan(fx+e))^2} \right)}{f c^4}$	68
risch	$-\frac{a^2 e^{8i(fx+e)} B}{32c^4 f} - \frac{ia^2 e^{8i(fx+e)} A}{32c^4 f} - \frac{iA a^2 e^{6i(fx+e)}}{12c^4 f} + \frac{a^2 e^{4i(fx+e)} B}{16c^4 f} - \frac{ia^2 e^{4i(fx+e)} A}{16c^4 f}$	110

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x,method=_R ETURNVERBOSE)`

$$3.686. \quad \int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^4} dx$$

output $\frac{1}{f a^2 c^4} \left(-\frac{1}{4} \frac{(2B+2IA)}{(I+\tan(fx+e))^4} - \frac{1}{3} \frac{(-A+3IB)}{(I+\tan(fx+e))^3} + \frac{1}{2} \frac{B}{(I+\tan(fx+e))^2} \right)$

3.686.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.70

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx$$

$$= -\frac{3(iA + B)a^2 e^{(8i fx + 8ie)} + 8i A a^2 e^{(6i fx + 6ie)} + 6(iA - B)a^2 e^{(4i fx + 4ie)}}{96 c^4 f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x, algorithm="fracas")`

output $-\frac{1}{96} \frac{(3(IA + B)a^2 e^{(8I*fx + 8I*e)} + 8I*A*a^2 e^{(6I*fx + 6I*e)} + 6(IA - B)a^2 e^{(4I*fx + 4I*e)})}{c^4 f}$

3.686.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 218 vs. $2(73) = 146$.

Time = 0.33 (sec) , antiderivative size = 218, normalized size of antiderivative = 2.40

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx$$

$$= \begin{cases} \frac{-512iAa^2 c^8 f^2 e^{6ie} e^{6ifx} + (-384iAa^2 c^8 f^2 e^{4ie} + 384Ba^2 c^8 f^2 e^{4ie}) e^{4ifx} + (-192iAa^2 c^8 f^2 e^{8ie} - 192Ba^2 c^8 f^2 e^{8ie}) e^{8ifx}}{6144c^{12} f^3} & \text{for } c^{12} f^3 \neq 0 \\ \frac{x(Aa^2 e^{8ie} + 2Aa^2 e^{6ie} + Aa^2 e^{4ie} - iBa^2 e^{8ie} + iBa^2 e^{4ie})}{4c^4} & \text{otherwise} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**4,x)`

output `Piecewise(((((-512*I*A*a**2*c**8*f**2*exp(6*I*e)*exp(6*I*fx) + (-384*I*A*a**2*c**8*f**2*exp(4*I*e) + 384*B*a**2*c**8*f**2*exp(4*I*e))*exp(4*I*fx) + (-192*I*A*a**2*c**8*f**2*exp(8*I*e) - 192*B*a**2*c**8*f**2*exp(8*I*e))*exp(8*I*fx))/(6144*c**12*f**3), Ne(c**12*f**3, 0)), (x*(A*a**2*exp(8*I*e) + 2*A*a**2*exp(6*I*e) + A*a**2*exp(4*I*e) - I*B*a**2*exp(8*I*e) + I*B*a**2*exp(4*I*e))/(4*c**4), True))`

3.686. $\int \frac{(a+ia \tan(e+fx))^2 (A+B \tan(e+fx))}{(c-ic \tan(e+fx))^4} dx$

3.686.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.686.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 190 vs. $2(75) = 150$.

Time = 0.89 (sec) , antiderivative size = 190, normalized size of antiderivative = 2.09

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx =$$

$$\frac{2 \left(3 A a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^7 + 6 i A a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^6 - 3 B a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^6 - 17 A a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^5 \right)}{(c - ic \tan(e + fx))^4}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x, algorithm="giac")`

output `-2/3*(3*A*a^2*tan(1/2*f*x + 1/2*e)^7 + 6*I*A*a^2*tan(1/2*f*x + 1/2*e)^6 - 3*B*a^2*tan(1/2*f*x + 1/2*e)^6 - 17*A*a^2*tan(1/2*f*x + 1/2*e)^5 - 16*I*A*a^2*tan(1/2*f*x + 1/2*e)^4 + 6*B*a^2*tan(1/2*f*x + 1/2*e)^4 + 17*A*a^2*tan(1/2*f*x + 1/2*e)^3 + 6*I*A*a^2*tan(1/2*f*x + 1/2*e)^2 - 3*B*a^2*tan(1/2*f*x + 1/2*e)^2 - 3*A*a^2*tan(1/2*f*x + 1/2*e))/(c^4*f*(tan(1/2*f*x + 1/2*e) + I)^8)`

3.686.9 Mupad [B] (verification not implemented)

Time = 8.45 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.86

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx$$

$$= \frac{a^2 (3B \tan(e + fx)^2 + 2A \tan(e + fx) - A i)}{6c^4 f (\tan(e + fx)^4 + \tan(e + fx)^3 4i - 6 \tan(e + fx)^2 - \tan(e + fx) 4i + 1)}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2)/(c - c*tan(e + f*x)*1i)^4,x)`

output `(a^2*(2*A*tan(e + f*x) - A*1i + 3*B*tan(e + f*x)^2))/(6*c^4*f*(tan(e + f*x)^3*4i - 6*tan(e + f*x)^2 - tan(e + f*x)*4i + tan(e + f*x)^4 + 1))`

3.687
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^5} dx$$

3.687.1 Optimal result	6437
3.687.2 Mathematica [A] (verified)	6437
3.687.3 Rubi [A] (verified)	6438
3.687.4 Maple [A] (verified)	6439
3.687.5 Fricas [A] (verification not implemented)	6440
3.687.6 Sympy [B] (verification not implemented)	6440
3.687.7 Maxima [F(-2)]	6441
3.687.8 Giac [B] (verification not implemented)	6441
3.687.9 Mupad [B] (verification not implemented)	6442

3.687.1 Optimal result

Integrand size = 41, antiderivative size = 95

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx$$

$$= \frac{2a^2(A - iB)}{5c^5 f(i + \tan(e + fx))^5} + \frac{a^2(iA + 3B)}{4c^5 f(i + \tan(e + fx))^4} + \frac{ia^2B}{3c^5 f(i + \tan(e + fx))^3}$$

output $2/5*a^2*(A-I*B)/c^5/f/(I+\tan(f*x+e))^5+1/4*a^2*(I*A+3*B)/c^5/f/(I+\tan(f*x+e))^4+1/3*I*a^2*B/c^5/f/(I+\tan(f*x+e))^3$

3.687.2 Mathematica [A] (verified)

Time = 5.36 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.65

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx$$

$$= \frac{a^2(9A + iB + 5(3iA + B) \tan(e + fx) + 20iB \tan^2(e + fx))}{60c^5 f(i + \tan(e + fx))^5}$$

input `Integrate[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^5,x]`

output $(a^2*(9*A + I*B + 5*((3*I)*A + B)*Tan[e + f*x] + (20*I)*B*Tan[e + f*x]^2))/(60*c^5*f*(I + Tan[e + f*x])^5)$

3.687.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.82, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{a(i \tan(e+fx)+1)(A+B \tan(e+fx))}{c^6(1-i \tan(e+fx))^6} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^2 \int \frac{(i \tan(e+fx)+1)(A+B \tan(e+fx))}{(1-i \tan(e+fx))^6} d \tan(e + fx)}{c^5 f} \\
 & \quad \downarrow \text{86} \\
 & \frac{a^2 \int \left(-\frac{2(A-iB)}{(\tan(e+fx)+i)^6} - \frac{iB}{(\tan(e+fx)+i)^4} - \frac{i(A-3iB)}{(\tan(e+fx)+i)^5} \right) d \tan(e + fx)}{c^5 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^2 \left(\frac{2(A-iB)}{5(\tan(e+fx)+i)^5} + \frac{3B+iA}{4(\tan(e+fx)+i)^4} + \frac{iB}{3(\tan(e+fx)+i)^3} \right)}{c^5 f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^5,x]`

output `(a^2*((2*(A - I*B))/(5*(I + Tan[e + f*x])^5) + (I*A + 3*B)/(4*(I + Tan[e + f*x])^4) + ((I/3)*B)/(I + Tan[e + f*x]^3))/(c^5*f)`

3.687.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.687.4 Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.73

method	result
derivativedivides	$\frac{a^2 \left(-\frac{2iB-2A}{5(i+\tan(fx+e))^5} + \frac{iB}{3(i+\tan(fx+e))^3} - \frac{-iA-3B}{4(i+\tan(fx+e))^4} \right)}{f c^5}$
default	$\frac{a^2 \left(-\frac{2iB-2A}{5(i+\tan(fx+e))^5} + \frac{iB}{3(i+\tan(fx+e))^3} - \frac{-iA-3B}{4(i+\tan(fx+e))^4} \right)}{f c^5}$
risch	$-\frac{a^2 e^{10i(fx+e)} B}{80c^5 f} - \frac{ia^2 e^{10i(fx+e)} A}{80c^5 f} - \frac{e^{8i(fx+e)} B a^2}{64c^5 f} - \frac{3ie^{8i(fx+e)} A a^2}{64c^5 f} + \frac{e^{6i(fx+e)} B a^2}{48c^5 f} - \frac{ie^{6i(fx+e)} A a^2}{16c^5 f} +$
norman	$\frac{A a^2 \tan(fx+e) + \frac{-9iA a^2 + B a^2}{60cf} - \frac{(-7iB a^2 + 12A a^2) \tan(fx+e)^3}{3cf} + \frac{(iA a^2 + 7B a^2) \tan(fx+e)^6}{4cf} + \frac{7(-8iB a^2 + 3A a^2) \tan(fx+e)}{15cf}}{(1+\tan(fx+e))^5 c^4}$

3.687. $\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^5} dx$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^5,x,method=_R
ETURNVERBOSE)`

output `1/f*a^2/c^5*(-1/5*(-2*A+2*I*B)/(I+tan(f*x+e))^5+1/3*I*B/(I+tan(f*x+e))^3-1
/4*(-I*A-3*B)/(I+tan(f*x+e))^4)`

3.687.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.94

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx =$$

$$\frac{12(iA + B)a^2 e^{(10i fx + 10ie)} + 15(3iA + B)a^2 e^{(8i fx + 8ie)} + 20(3iA - B)a^2 e^{(6i fx + 6ie)} + 30(iA - B)a^2 e^{(4i fx + 4ie)}}{960 c^5 f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^5,x, al
gorithm="fricas")`

output `-1/960*(12*(I*A + B)*a^2*e^(10*I*f*x + 10*I*e) + 15*(3*I*A + B)*a^2*e^(8*I
*f*x + 8*I*e) + 20*(3*I*A - B)*a^2*e^(6*I*f*x + 6*I*e) + 30*(I*A - B)*a^2*
e^(4*I*f*x + 4*I*e))/(c^5*f)`

3.687.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 332 vs. $2(76) = 152$.

Time = 0.41 (sec) , antiderivative size = 332, normalized size of antiderivative = 3.49

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx$$

$$= \begin{cases} \frac{(-245760iAa^2c^{15}f^3e^{4ie} + 245760Ba^2c^{15}f^3e^{4ie})e^{4ifx} + (-491520iAa^2c^{15}f^3e^{6ie} + 163840Ba^2c^{15}f^3e^{6ie})e^{6ifx} + (-368640iAa^2c^{15}f^3e^{8ie} - 122880Ba^2c^{15}f^3e^{8ie})e^{8ifx} + (-245760iAa^2c^{15}f^3e^{10ie} + 245760Ba^2c^{15}f^3e^{10ie})e^{10ifx}}{7864320c^{20}f^4} \\ \frac{x(Aa^2e^{10ie} + 3Aa^2e^{8ie} + 3Aa^2e^{6ie} + Aa^2e^{4ie} - iBa^2e^{10ie} - iBa^2e^{8ie} + iBa^2e^{6ie} + iBa^2e^{4ie})}{8c^5} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**5,x)`

3.687. $\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^5} dx$

output `Piecewise(((((-245760*I*A*a**2*c**15*f**3*exp(4*I*e) + 245760*B*a**2*c**15*f**3*exp(4*I*e))*exp(4*I*f*x) + (-491520*I*A*a**2*c**15*f**3*exp(6*I*e) + 163840*B*a**2*c**15*f**3*exp(6*I*e))*exp(6*I*f*x) + (-368640*I*A*a**2*c**15*f**3*exp(8*I*e) - 122880*B*a**2*c**15*f**3*exp(8*I*e))*exp(8*I*f*x) + (-98304*I*A*a**2*c**15*f**3*exp(10*I*e) - 98304*B*a**2*c**15*f**3*exp(10*I*e))*exp(10*I*f*x))/(7864320*c**20*f**4), Ne(c**20*f**4, 0)), (x*(A*a**2*exp(10*I*e) + 3*A*a**2*exp(8*I*e) + 3*A*a**2*exp(6*I*e) + A*a**2*exp(4*I*e) - I*B*a**2*exp(10*I*e) - I*B*a**2*exp(8*I*e) + I*B*a**2*exp(6*I*e) + I*B*a**2*exp(4*I*e))/(8*c**5), True))`

3.687.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^5,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.687.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 292 vs. $2(77) = 154$.

Time = 1.13 (sec) , antiderivative size = 292, normalized size of antiderivative = 3.07

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx =$$

$$\frac{2 \left(15 A a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^9 + 45 i A a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^8 - 15 B a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^8 - 150 A a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^7 \right)}{(c - ic \tan(e + fx))^5}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^5,x, algorithm="giac")`

3.687. $\int \frac{(a+ia \tan(e+fx))^2 (A+B \tan(e+fx))}{(c-ic \tan(e+fx))^5} dx$

output
$$\begin{aligned} & -2/15*(15*A*a^2*\tan(1/2*f*x + 1/2*e)^9 + 45*I*A*a^2*\tan(1/2*f*x + 1/2*e)^8 \\ & - 15*B*a^2*\tan(1/2*f*x + 1/2*e)^8 - 150*A*a^2*\tan(1/2*f*x + 1/2*e)^7 - 10 \\ & *I*B*a^2*\tan(1/2*f*x + 1/2*e)^7 - 225*I*A*a^2*\tan(1/2*f*x + 1/2*e)^6 + 55* \\ & B*a^2*\tan(1/2*f*x + 1/2*e)^6 + 306*A*a^2*\tan(1/2*f*x + 1/2*e)^5 + 24*I*B*a \\ & ^2*\tan(1/2*f*x + 1/2*e)^5 + 225*I*A*a^2*\tan(1/2*f*x + 1/2*e)^4 - 55*B*a^2* \\ & \tan(1/2*f*x + 1/2*e)^4 - 150*A*a^2*\tan(1/2*f*x + 1/2*e)^3 - 10*I*B*a^2*\tan \\ & (1/2*f*x + 1/2*e)^3 - 45*I*A*a^2*\tan(1/2*f*x + 1/2*e)^2 + 15*B*a^2*\tan(1/2 \\ & *f*x + 1/2*e)^2 + 15*A*a^2*\tan(1/2*f*x + 1/2*e))/(c^5*f*(\tan(1/2*f*x + 1/2 \\ & *e) + I)^{10}) \end{aligned}$$

3.687.9 Mupad [B] (verification not implemented)

Time = 8.65 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.14

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx$$

$$= \frac{\frac{a^2 (9A + B i i)}{60} + \frac{a^2 \tan(e + fx) (5B + A 15i)}{60} + \frac{B a^2 \tan(e + fx)^2 i i}{3}}{c^5 f (\tan(e + fx)^5 + \tan(e + fx)^4 5i - 10 \tan(e + fx)^3 - \tan(e + fx)^2 10i + 5 \tan(e + fx) + i i)}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2)/(c - c*tan(e + f*x)*1i)^5,x)`

output
$$\frac{(a^2*(9*A + B*1i))/60 + (a^2*\tan(e + f*x)*(A*15i + 5*B))/60 + (B*a^2*\tan(e + f*x)^2*1i)/3}{(c^5*f*(5*\tan(e + f*x) - \tan(e + f*x)^2*10i - 10*\tan(e + f*x)^3 + \tan(e + f*x)^4*5i + \tan(e + f*x)^5 + 1i))}$$

3.688 $\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^6} dx$

3.688.1 Optimal result	6443
3.688.2 Mathematica [A] (verified)	6443
3.688.3 Rubi [A] (verified)	6444
3.688.4 Maple [A] (verified)	6445
3.688.5 Fricas [A] (verification not implemented)	6446
3.688.6 Sympy [B] (verification not implemented)	6446
3.688.7 Maxima [F(-2)]	6447
3.688.8 Giac [B] (verification not implemented)	6447
3.688.9 Mupad [B] (verification not implemented)	6448

3.688.1 Optimal result

Integrand size = 41, antiderivative size = 91

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx$$

$$= \frac{a^2(iA + B)}{3c^6 f(i + \tan(e + fx))^6} - \frac{a^2(A - 3iB)}{5c^6 f(i + \tan(e + fx))^5} - \frac{a^2 B}{4c^6 f(i + \tan(e + fx))^4}$$

output $1/3*a^2*(I*A+B)/c^6/f/(I+\tan(f*x+e))^6-1/5*a^2*(A-3*I*B)/c^6/f/(I+\tan(f*x+e))^5-1/4*a^2*B/c^6/f/(I+\tan(f*x+e))^4$

3.688.2 Mathematica [A] (verified)

Time = 5.41 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.66

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx$$

$$= -\frac{a^2(-8iA + B + 6(2A - iB) \tan(e + fx) + 15B \tan^2(e + fx))}{60c^6 f(i + \tan(e + fx))^6}$$

input `Integrate[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^6,x]`

output $-1/60*(a^2*((-8*I)*A + B + 6*(2*A - I*B)*Tan[e + f*x] + 15*B*Tan[e + f*x]^2))/(c^6*f*(I + Tan[e + f*x])^6)$

3.688. $\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^6} dx$

3.688.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.81, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{a(i \tan(e+fx)+1)(A+B \tan(e+fx))}{c^7(1-i \tan(e+fx))^7} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^2 \int \frac{(i \tan(e+fx)+1)(A+B \tan(e+fx))}{(1-i \tan(e+fx))^7} d \tan(e + fx)}{c^6 f} \\
 & \quad \downarrow \text{86} \\
 & \frac{a^2 \int \left(-\frac{2i(A-iB)}{(\tan(e+fx)+i)^7} + \frac{B}{(\tan(e+fx)+i)^5} + \frac{A-3iB}{(\tan(e+fx)+i)^6} \right) d \tan(e + fx)}{c^6 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^2 \left(-\frac{A-3iB}{5(\tan(e+fx)+i)^5} + \frac{B+iA}{3(\tan(e+fx)+i)^6} - \frac{B}{4(\tan(e+fx)+i)^4} \right)}{c^6 f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^6,x]`

output `(a^2*((I*A + B)/(3*(I + Tan[e + f*x])^6) - (A - (3*I)*B)/(5*(I + Tan[e + f*x])^5) - B/(4*(I + Tan[e + f*x])^4)))/(c^6*f)`

3.688.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.688.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.73

method	result
derivativedivides	$\frac{a^2 \left(-\frac{-3iB+A}{5(i+\tan(fx+e))^5} - \frac{B}{4(i+\tan(fx+e))^4} - \frac{-2iA-2B}{6(i+\tan(fx+e))^6} \right)}{f c^6}$
default	$\frac{a^2 \left(-\frac{-3iB+A}{5(i+\tan(fx+e))^5} - \frac{B}{4(i+\tan(fx+e))^4} - \frac{-2iA-2B}{6(i+\tan(fx+e))^6} \right)}{f c^6}$
risch	$-\frac{a^2 e^{12i(fx+e)} B}{192c^6 f} - \frac{ia^2 e^{12i(fx+e)} A}{192c^6 f} - \frac{e^{10i(fx+e)} B a^2}{80c^6 f} - \frac{ie^{10i(fx+e)} A a^2}{40c^6 f} - \frac{3iA a^2 e^{8i(fx+e)}}{64c^6 f} + \frac{e^{6i(fx+e)} B a^2}{48c^6 f}$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^6,x,method=_R ETURNVERBOSE)`

$$3.688. \quad \int \frac{(a+ia \tan(e+fx))^2 (A+B \tan(e+fx))}{(c-ic \tan(e+fx))^6} dx$$

output $1/f*a^2/c^6*(-1/5*(A-3*I*B)/(I+\tan(f*x+e))^5-1/4*B/(I+\tan(f*x+e))^4-1/6*(-2*I*A-2*B)/(I+\tan(f*x+e))^6)$

3.688.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.14

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx = \frac{5(iA + B)a^2e^{(12ifx+12ie)} + 12(2iA + B)a^2e^{(10ifx+10ie)} + 45iAa^2e^{(8ifx+8ie)} + 20(2iA - B)a^2e^{(6ifx+6ie)}}{960c^6f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^6,x, algorithm="fricas")`

output $-1/960*(5*(I*A + B)*a^2*e^{(12*I*f*x + 12*I*e)} + 12*(2*I*A + B)*a^2*e^{(10*I*f*x + 10*I*e)} + 45*I*A*a^2*e^{(8*I*f*x + 8*I*e)} + 20*(2*I*A - B)*a^2*e^{(6*I*f*x + 6*I*e)} + 15*(I*A - B)*a^2*e^{(4*I*f*x + 4*I*e)})/(c^6*f)$

3.688.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 379 vs. 2(73) = 146.

Time = 0.52 (sec) , antiderivative size = 379, normalized size of antiderivative = 4.16

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx = \left\{ \begin{array}{l} \frac{-141557760iAa^2c^{24}f^4e^{8ie}e^{8ifx} + (-47185920iAa^2c^{24}f^4e^{4ie} + 47185920Ba^2c^{24}f^4e^{4ie})e^{4ifx} + (-125829120iAa^2c^{24}f^4e^{6ie} + 62914560Ba^2c^{24}f^4e^{6ie})e^{2ifx} + 3019898880iAa^2c^{24}f^4e^{2ie}}{16c^6} \\ x(Aa^2e^{12ie} + 4Aa^2e^{10ie} + 6Aa^2e^{8ie} + 4Aa^2e^{6ie} + Aa^2e^{4ie} - iBa^2e^{12ie} - 2iBa^2e^{10ie} + 2iBa^2e^{6ie} + iBa^2e^{4ie}) \end{array} \right.$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**6,x)`

output `Piecewise(((−141557760*I*A**2*c**24*f**4*exp(8*I*e)*exp(8*I*f*x) + (−47185920*I*A**2*c**24*f**4*exp(4*I*e) + 47185920*B**2*c**24*f**4*exp(4*I*e))*exp(4*I*f*x) + (−125829120*I*A**2*c**24*f**4*exp(6*I*e) + 62914560*B**2*c**24*f**4*exp(6*I*e))*exp(6*I*f*x) + (−75497472*I*A**2*c**24*f**4*exp(10*I*e) − 37748736*B**2*c**24*f**4*exp(10*I*e))*exp(10*I*f*x) + (−15728640*I*A**2*c**24*f**4*exp(12*I*e) − 15728640*B**2*c**24*f**4*exp(12*I*e))*exp(12*I*f*x))/(3019898880*c**30*f**5), Ne(c**30*f**5, 0)), (x*(A**2*exp(12*I*e) + 4*A**2*exp(10*I*e) + 6*A**2*exp(8*I*e) + 4*A**2*exp(6*I*e) + A**2*exp(4*I*e) − I*B**2*exp(12*I*e) − 2*I*B**2*exp(10*I*e) + 2*I*B**2*exp(6*I*e) + I*B**2*exp(4*I*e))/(16*c**6), True))`

3.688.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^6,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.688.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 360 vs. $2(75) = 150$.

Time = 1.25 (sec) , antiderivative size = 360, normalized size of antiderivative = 3.96

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx = \frac{2 \left(15 A a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^{11} + 60 i A a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^{10} - 15 B a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^{10} - 235 A a^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^9 \right)}{(c - ic \tan(e + fx))^6}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^6,x, algorithm="giac")`

3.688. $\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx$

output
$$\begin{aligned} & -2/15*(15*A*a^2*\tan(1/2*f*x + 1/2*e)^{11} + 60*I*A*a^2*\tan(1/2*f*x + 1/2*e)^{10} - 15*B*a^2*\tan(1/2*f*x + 1/2*e)^{10} - 235*A*a^2*\tan(1/2*f*x + 1/2*e)^9 - \\ & 20*I*B*a^2*\tan(1/2*f*x + 1/2*e)^9 - 480*I*A*a^2*\tan(1/2*f*x + 1/2*e)^8 + 90*B*a^2*\tan(1/2*f*x + 1/2*e)^8 + 822*A*a^2*\tan(1/2*f*x + 1/2*e)^7 + 84*I* \\ & B*a^2*\tan(1/2*f*x + 1/2*e)^7 + 904*I*A*a^2*\tan(1/2*f*x + 1/2*e)^6 - 158*B* \\ & a^2*\tan(1/2*f*x + 1/2*e)^6 - 822*A*a^2*\tan(1/2*f*x + 1/2*e)^5 - 84*I*B*a^2 \\ & * \tan(1/2*f*x + 1/2*e)^5 - 480*I*A*a^2*\tan(1/2*f*x + 1/2*e)^4 + 90*B*a^2* \tan \\ & n(1/2*f*x + 1/2*e)^4 + 235*A*a^2*\tan(1/2*f*x + 1/2*e)^3 + 20*I*B*a^2*\tan(1 \\ & /2*f*x + 1/2*e)^3 + 60*I*A*a^2*\tan(1/2*f*x + 1/2*e)^2 - 15*B*a^2*\tan(1/2*f \\ & *x + 1/2*e)^2 - 15*A*a^2*\tan(1/2*f*x + 1/2*e))/ (c^6*f*(\tan(1/2*f*x + 1/2*e) \\ &) + I)^{12} \end{aligned}$$

3.688.9 Mupad [B] (verification not implemented)

Time = 8.48 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.30

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx =$$

$$-\frac{\frac{B a^2 \tan(e+fx)^2}{4} + \frac{a^2 \tan(e+fx) (12A - B 6i)}{60} - \frac{a^2 (-B + A 8i)}{60}}{c^6 f (\tan(e + fx)^6 + \tan(e + fx)^5 6i - 15 \tan(e + fx)^4 - \tan(e + fx)^3 20i + 15 \tan(e + fx)^2 + \tan(e + fx) - 1)}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2)/(c - c*tan(e + f*x)*1i)^6,x)`

output
$$-\frac{(a^2*\tan(e + f*x)*(12*A - B*6i))/60 - (a^2*(A*8i - B))/60 + (B*a^2*\tan(e + f*x)^2)/4}{(c^6*f*(\tan(e + f*x)*6i + 15*\tan(e + f*x)^2 - \tan(e + f*x)^3 *20i - 15*\tan(e + f*x)^4 + \tan(e + f*x)^5*6i + \tan(e + f*x)^6 - 1))}$$

3.689 $\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx))^n dx$

3.689.1 Optimal result	6449
3.689.2 Mathematica [A] (verified)	6449
3.689.3 Rubi [A] (verified)	6450
3.689.4 Maple [A] (verified)	6452
3.689.5 Fricas [B] (verification not implemented)	6452
3.689.6 Sympy [B] (verification not implemented)	6453
3.689.7 Maxima [B] (verification not implemented)	6454
3.689.8 Giac [F]	6455
3.689.9 Mupad [B] (verification not implemented)	6455

3.689.1 Optimal result

Integrand size = 41, antiderivative size = 151

$$\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))(c - ictan(e + fx))^n dx$$

$$= \frac{4a^3(iA + B)(c - ictan(e + fx))^n}{fn} - \frac{4a^3(iA + 2B)(c - ictan(e + fx))^{1+n}}{cf(1 + n)}$$

$$+ \frac{a^3(iA + 5B)(c - ictan(e + fx))^{2+n}}{c^2f(2 + n)} - \frac{a^3B(c - ictan(e + fx))^{3+n}}{c^3f(3 + n)}$$

```
output 4*a^3*(I*A+B)*(c-I*c*tan(f*x+e))^n/f/n-4*a^3*(I*A+2*B)*(c-I*c*tan(f*x+e))^(1+n)/c/f/(1+n)+a^3*(I*A+5*B)*(c-I*c*tan(f*x+e))^(2+n)/c^2/f/(2+n)-a^3*B*(c-I*c*tan(f*x+e))^(3+n)/c^3/f/(3+n)
```

3.689.2 Mathematica [A] (verified)

Time = 5.19 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.02

$$\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))(c - ictan(e + fx))^n dx =$$

$$- \frac{ia^3(c - ictan(e + fx))^n (iB(24 + 9n + n^2) - A(24 + 23n + 8n^2 + n^3) - in(2A(3 + n)^2 - iB(24 + 9n + n^2)))}{fn(1 + n)(2 + n)}$$

input `Integrate[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^n,x]`

output `((-I)*a^3*(c - I*c*Tan[e + f*x])^n*(I*B*(24 + 9*n + n^2) - A*(24 + 23*n + 8*n^2 + n^3) - I*n*(2*A*(3 + n)^2 - I*B*(24 + 9*n + n^2))*Tan[e + f*x] + n*(1 + n)*(A*(3 + n) - I*B*(9 + 2*n))*Tan[e + f*x]^2 + B*n*(2 + 3*n + n^2)*Tan[e + f*x]^3)/(f*n*(1 + n)*(2 + n)*(3 + n))`

3.689.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.91, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ict \tan(e + fx))^n dx$$

$$\downarrow 3042$$

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ict \tan(e + fx))^n dx$$

$$\downarrow 4071$$

$$\frac{ac \int a^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))(c - ict \tan(e + fx))^{n-1} d \tan(e + fx)}{f}$$

$$\downarrow 27$$

$$\frac{a^3 c \int (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))(c - ict \tan(e + fx))^{n-1} d \tan(e + fx)}{f}$$

$$\downarrow 86$$

$$\frac{a^3 c \int \left(4(A - iB)(c - ict \tan(e + fx))^{n-1} - \frac{4(A - 2iB)(c - ict \tan(e + fx))^n}{c} + \frac{(A - 5iB)(c - ict \tan(e + fx))^{n+1}}{c^2} + \frac{iB(c - ict \tan(e + fx))^{n+3}}{c^3} \right)}{f}$$

$$\downarrow 2009$$

$$\frac{a^3 c \left(\frac{(5B + iA)(c - ict \tan(e + fx))^{n+2}}{c^3(n+2)} - \frac{4(2B + iA)(c - ict \tan(e + fx))^{n+1}}{c^2(n+1)} + \frac{4(B + iA)(c - ict \tan(e + fx))^n}{cn} - \frac{B(c - ict \tan(e + fx))^{n+3}}{c^4(n+3)} \right)}{f}$$

3.689. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ict \tan(e + fx))^n dx$

input `Int[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^n, x]`

output `(a^3*c*((4*(I*A + B)*(c - I*c*Tan[e + f*x])^n)/(c*n) - (4*(I*A + 2*B)*(c - I*c*Tan[e + f*x])^(1 + n))/(c^2*(1 + n)) + ((I*A + 5*B)*(c - I*c*Tan[e + f*x])^(2 + n))/(c^3*(2 + n)) - (B*(c - I*c*Tan[e + f*x])^(3 + n))/(c^4*(3 + n))))/f`

3.689.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.689.4 Maple [A] (verified)

Time = 0.96 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.79

method	result
derivativedivides	$\frac{i(Aa^3n^3 - iBa^3n^2 + 8Aa^3n^2 - 9iBa^3n + 23a^3An - 24iBa^3 + 24a^3A)e^{n \ln(c - ic \tan(fx + e))}}{(n^2 + 3n + 2)fn(3+n)} - \frac{a^3(-iBn^2 + 2An^2 - 9iBn}$
default	$\frac{i(Aa^3n^3 - iBa^3n^2 + 8Aa^3n^2 - 9iBa^3n + 23a^3An - 24iBa^3 + 24a^3A)e^{n \ln(c - ic \tan(fx + e))}}{(n^2 + 3n + 2)fn(3+n)} - \frac{a^3(-iBn^2 + 2An^2 - 9iBn}$
risch	Expression too large to display

input `int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x,method=_RETURVERBOSE)`

output `I/(n^2+3*n+2)/f/n*(-9*I*B*a^3*n+23*a^3*A*n-24*I*B*a^3+A*a^3*n^3+8*A*a^3*n^2-I*B*a^3*n^2+24*a^3*A)/(3+n)*exp(n*ln(c-I*c*tan(f*x+e)))-a^3*(-I*B*n^2+2*A*n^2-9*I*B*n+12*A*n-24*I*B+18*A)/f/(n^2+3*n+2)/(3+n)*tan(f*x+e)*exp(n*ln(c-I*c*tan(f*x+e)))-(I*A*n+3*I*A+2*B*n+9*B)*a^3/f/(3+n)/(2+n)*tan(f*x+e)^2*exp(n*ln(c-I*c*tan(f*x+e)))-I/f/(3+n)*B*a^3*tan(f*x+e)^3*exp(n*ln(c-I*c*tan(f*x+e)))`

3.689.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 335 vs. 2(137) = 274.

Time = 0.26 (sec) , antiderivative size = 335, normalized size of antiderivative = 2.22

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx = \frac{4(2(-iA + B)a^3n + 6(-iA - B)a^3 + ((-iA - B)a^3n^3 + 6(-iA - B)a^3n^2 + 11(-iA - B)a^3n + 6(-iA - B)a^3))}{fn^4 + 6fn^3 + 11fn^2 + 6fn + f}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x, algorithm="fracas")`

```
output -4*(2*(-I*A + B)*a^3*n + 6*(-I*A - B)*a^3 + ((-I*A - B)*a^3*n^3 + 6*(-I*A
- B)*a^3*n^2 + 11*(-I*A - B)*a^3*n + 6*(-I*A - B)*a^3)*e^(6*I*f*x + 6*I*e)
+ ((-I*A + B)*a^3*n^3 + 2*(-4*I*A + B)*a^3*n^2 + 3*(-7*I*A - 3*B)*a^3*n +
18*(-I*A - B)*a^3)*e^(4*I*f*x + 4*I*e) + 2*((-I*A + B)*a^3*n^2 - 6*I*A*a^
3*n + 9*(-I*A - B)*a^3)*e^(2*I*f*x + 2*I*e))*(2*c/(e^(2*I*f*x + 2*I*e) + 1
))^n/(f*n^4 + 6*f*n^3 + 11*f*n^2 + 6*f*n + (f*n^4 + 6*f*n^3 + 11*f*n^2 + 6
*f*n)*e^(6*I*f*x + 6*I*e) + 3*(f*n^4 + 6*f*n^3 + 11*f*n^2 + 6*f*n)*e^(4*I*
f*x + 4*I*e) + 3*(f*n^4 + 6*f*n^3 + 11*f*n^2 + 6*f*n)*e^(2*I*f*x + 2*I*e))
```

3.689.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 3665 vs. $2(122) = 244$.

Time = 3.36 (sec) , antiderivative size = 3665, normalized size of antiderivative = 24.27

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^n dx = \text{Too large to display}$$

```
input integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**n,x)
```

```
output Piecewise((x*(A + B*tan(e))*(I*a*tan(e) + a)**3*(-I*c*tan(e) + c)**n, Eq(f
, 0)), (6*A*a**3*tan(e + f*x)**2/(6*c**3*f*tan(e + f*x)**3 + 18*I*c**3*f*t
an(e + f*x)**2 - 18*c**3*f*tan(e + f*x) - 6*I*c**3*f) - 2*A*a**3/(6*c**3*f
*tan(e + f*x)**3 + 18*I*c**3*f*tan(e + f*x)**2 - 18*c**3*f*tan(e + f*x) -
6*I*c**3*f) + 6*I*B*a**3*f*x*tan(e + f*x)**3/(6*c**3*f*tan(e + f*x)**3 + 1
8*I*c**3*f*tan(e + f*x)**2 - 18*c**3*f*tan(e + f*x) - 6*I*c**3*f) - 18*B*a
**3*f*x*tan(e + f*x)**2/(6*c**3*f*tan(e + f*x)**3 + 18*I*c**3*f*tan(e + f*
x)**2 - 18*c**3*f*tan(e + f*x) - 6*I*c**3*f) - 18*I*B*a**3*f*x*tan(e + f*x
)/(6*c**3*f*tan(e + f*x)**3 + 18*I*c**3*f*tan(e + f*x)**2 - 18*c**3*f*tan(
e + f*x) - 6*I*c**3*f) + 6*B*a**3*f*x/(6*c**3*f*tan(e + f*x)**3 + 18*I*c**
3*f*tan(e + f*x)**2 - 18*c**3*f*tan(e + f*x) - 6*I*c**3*f) - 3*B*a**3*log(
tan(e + f*x)**2 + 1)*tan(e + f*x)**3/(6*c**3*f*tan(e + f*x)**3 + 18*I*c**3
*f*tan(e + f*x)**2 - 18*c**3*f*tan(e + f*x) - 6*I*c**3*f) - 9*I*B*a**3*log
(tan(e + f*x)**2 + 1)*tan(e + f*x)**2/(6*c**3*f*tan(e + f*x)**3 + 18*I*c**
3*f*tan(e + f*x)**2 - 18*c**3*f*tan(e + f*x) - 6*I*c**3*f) + 9*B*a**3*log(
tan(e + f*x)**2 + 1)*tan(e + f*x)/(6*c**3*f*tan(e + f*x)**3 + 18*I*c**3*f*
tan(e + f*x)**2 - 18*c**3*f*tan(e + f*x) - 6*I*c**3*f) + 3*I*B*a**3*log(ta
n(e + f*x)**2 + 1)/(6*c**3*f*tan(e + f*x)**3 + 18*I*c**3*f*tan(e + f*x)**2
- 18*c**3*f*tan(e + f*x) - 6*I*c**3*f) - 30*I*B*a**3*tan(e + f*x)**2/(6*c
**3*f*tan(e + f*x)**3 + 18*I*c**3*f*tan(e + f*x)**2 - 18*c**3*f*tan(e + ...
```

3.689. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^n dx$

3.689.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1056 vs. $2(137) = 274$.

Time = 0.58 (sec) , antiderivative size = 1056, normalized size of antiderivative = 6.99

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^n dx = \text{Too large to display}$$

```
input integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x, algorithm="maxima")
```

```
output 4*(2*((A + I*B)*a^3*c^n*n^2 + 6*A*a^3*c^n*n + 9*(A - I*B)*a^3*c^n)*2^n*cos(-2*f*x + n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - 2*e) + ((A + I*B)*a^3*c^n*n^3 + 2*(4*A + I*B)*a^3*c^n*n^2 + 3*(7*A - 3*I*B)*a^3*c^n*n + 18*(A - I*B)*a^3*c^n)*2^n*cos(-4*f*x + n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - 4*e) + ((A - I*B)*a^3*c^n*n^3 + 6*(A - I*B)*a^3*c^n*n^2 + 11*(A - I*B)*a^3*c^n*n + 6*(A - I*B)*a^3*c^n)*2^n*cos(-6*f*x + n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - 6*e) + 2*((A + I*B)*a^3*c^n*n + 3*(A - I*B)*a^3*c^n)*2^n*cos(n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)) - 2*((I*A - B)*a^3*c^n*n^2 + 6*I*A*a^3*c^n*n + 9*(I*A + B)*a^3*c^n)*2^n*sin(-2*f*x + n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - 2*e) - ((I*A - B)*a^3*c^n*n^3 + 2*(4*I*A - B)*a^3*c^n*n^2 + 3*(7*I*A + 3*B)*a^3*c^n*n + 18*(I*A + B)*a^3*c^n)*2^n*sin(-4*f*x + n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - 4*e) - ((I*A + B)*a^3*c^n*n^3 + 6*(I*A + B)*a^3*c^n*n^2 + 11*(I*A + B)*a^3*c^n*n + 6*(I*A + B)*a^3*c^n)*2^n*sin(-6*f*x + n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - 6*e) - 2*((I*A - B)*a^3*c^n*n + 3*(I*A + B)*a^3*c^n)*2^n*sin(n*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1)))/((-I*n^4 - 6*I*n^3 - 11*I*n^2 - 6*I*n)*(cos(2*f*x + 2*e))^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^(1/2*n)*cos(6*f*x + 6*e) - 3*(I*n^4 + 6*I*n^3 + 11*I*n^2 + 6*I*n)*(cos(2*f*x + 2*e))^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1)^(1/2*n)*cos(4*f*x + 4*e) + (n^4 + 6...
```

3.689.8 Giac [F]

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx$$

$$= \int (B \tan(fx + e) + A)(ia \tan(fx + e) + a)^3 (-ic \tan(fx + e) + c)^n dx$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^3*(-I*c*tan(f*x + e) + c)^n, x)`

3.689.9 Mupad [B] (verification not implemented)

Time = 13.21 (sec) , antiderivative size = 323, normalized size of antiderivative = 2.14

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx =$$

$$\frac{\left(c + \frac{c(e^{2i+fx} - 1) - i}{e^{2i+fx} + 1}\right)^n \left(\frac{8a^3(3A - B3i + An + Bn1i)}{fn(n^31i + n^26i + n11i + 6i)} + \frac{4a^3e^{e4i+fx4i}(n^2+5n+6)(3A - B3i + An + Bn1i)}{fn(n^31i + n^26i + n11i + 6i)} + \frac{4a^3e^{e6i+fx6i}}{fn(n^31i + n^26i + n11i + 6i)}\right)}{3e^{e2i+fx2i} + 3e^{e4i+fx4i} + e^{e6i+fx6i} + 1}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i)^n,x)`

output `-((c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^n*((8*a^3*(3*A - B*3i + A*n + B*n*1i))/(f*n*(n*11i + n^2*6i + n^3*1i + 6i)) + (4*a^3*exp(e*4i + f*x*4i)*(5*n + n^2 + 6)*(3*A - B*3i + A*n + B*n*1i))/(f*n*(n*11i + n^2*6i + n^3*1i + 6i)) + (4*a^3*exp(e*6i + f*x*6i)*(A - B*1i)*(11*n + 6*n^2 + n^3 + 6))/(f*n*(n*11i + n^2*6i + n^3*1i + 6i)) + (8*a^3*exp(e*2i + f*x*2i)*(n + 3)*(3*A - B*3i + A*n + B*n*1i))/(f*n*(n*11i + n^2*6i + n^3*1i + 6i))))/(3*exp(e*2i + f*x*2i) + 3*exp(e*4i + f*x*4i) + exp(e*6i + f*x*6i) + 1)`

3.690 $\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx))^6 dx$

3.690.1 Optimal result	6456
3.690.2 Mathematica [A] (verified)	6456
3.690.3 Rubi [A] (verified)	6457
3.690.4 Maple [A] (verified)	6459
3.690.5 Fricas [A] (verification not implemented)	6459
3.690.6 Sympy [B] (verification not implemented)	6460
3.690.7 Maxima [A] (verification not implemented)	6460
3.690.8 Giac [B] (verification not implemented)	6461
3.690.9 Mupad [B] (verification not implemented)	6462

3.690.1 Optimal result

Integrand size = 41, antiderivative size = 135

$$\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))(c - ictan(e + fx))^6 dx$$

$$= \frac{2a^3(iA + B)c^6(1 - i \tan(e + fx))^6}{3f} - \frac{4a^3(iA + 2B)c^6(1 - i \tan(e + fx))^7}{7f}$$

$$+ \frac{a^3(iA + 5B)c^6(1 - i \tan(e + fx))^8}{8f} - \frac{a^3Bc^6(1 - i \tan(e + fx))^9}{9f}$$

```
output 2/3*a^3*(I*A+B)*c^6*(1-I*tan(f*x+e))^6/f-4/7*a^3*(I*A+2*B)*c^6*(1-I*tan(f*x+e))^7/f+1/8*a^3*(I*A+5*B)*c^6*(1-I*tan(f*x+e))^8/f-1/9*a^3*B*c^6*(1-I*tan(f*x+e))^9/f
```

3.690.2 Mathematica [A] (verified)

Time = 5.89 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.11

$$\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))(c - ictan(e + fx))^6 dx$$

$$= \frac{a^3c^6 \sec^9(e + fx)(126(-3iA + B) \cos(e + fx) + 168(-iA + B) \cos(3(e + fx)) + 84A \sin(3(e + fx)) - 8$$

input `Integrate[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^6,x]`

output `(a^3*c^6*Sec[e + f*x]^9*(126*((-3*I)*A + B)*Cos[e + f*x] + 168*((-I)*A + B)*Cos[3*(e + f*x)] + 84*A*Sin[3*(e + f*x)] - (84*I)*B*Sin[3*(e + f*x)] + 108*A*Sin[5*(e + f*x)] + (36*I)*B*Sin[5*(e + f*x)] + 27*A*Sin[7*(e + f*x)] + (9*I)*B*Sin[7*(e + f*x)] + 3*A*Sin[9*(e + f*x)] + I*B*Sin[9*(e + f*x)])/(1008*f)`

3.690.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.81, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(e + fx))^3 (c - ictan(e + fx))^6 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(e + fx))^3 (c - ictan(e + fx))^6 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{a^3 c^6 \int (1 - i \tan(e + fx))^5 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^3 c^6 \int (1 - i \tan(e + fx))^5 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{86} \\
 & \frac{a^3 c^6 \int (iB(1 - i \tan(e + fx))^8 + (A - 5iB)(1 - i \tan(e + fx))^7 - 4(A - 2iB)(1 - i \tan(e + fx))^6 + 4(A - iB)(1 - i \tan(e + fx))^5)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^3 c^6 (\frac{1}{8}(5B + iA)(1 - i \tan(e + fx))^8 - \frac{4}{7}(2B + iA)(1 - i \tan(e + fx))^7 + \frac{2}{3}(B + iA)(1 - i \tan(e + fx))^6 - \frac{1}{9}B(1 - i \tan(e + fx))^5)}{f}
 \end{aligned}$$

3.690. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ictan(e + fx))^6 dx$

input `Int[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^6, x]`

output `(a^3*c^6*((2*(I*A + B)*(1 - I*Tan[e + f*x])^6)/3 - (4*(I*A + 2*B)*(1 - I*Tan[e + f*x])^7)/7 + ((I*A + 5*B)*(1 - I*Tan[e + f*x])^8)/8 - (B*(1 - I*Tan[e + f*x])^9)/9))/f`

3.690.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.690.4 Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.80

method	result
risch	$\frac{32c^6 a^3 (84iA e^{6i(fx+e)} + 84B e^{6i(fx+e)} + 108iA e^{4i(fx+e)} - 36B e^{4i(fx+e)} + 27iA e^{2i(fx+e)} - 9B e^{2i(fx+e)} + 3iA - B)}{63f (e^{2i(fx+e)} + 1)^9}$
derivativedivides	$ic^6 a^3 \left(\frac{B \tan(fx+e)^9}{9} + \frac{(3iB+A) \tan(fx+e)^8}{8} + \frac{(-11B-2iA+5i(-2iB+A)) \tan(fx+e)^7}{7} + \frac{(-11A+5i(-2iA-B)+10iB) \tan(fx+e)^6}{6} \right)$
default	$ic^6 a^3 \left(\frac{B \tan(fx+e)^9}{9} + \frac{(3iB+A) \tan(fx+e)^8}{8} + \frac{(-11B-2iA+5i(-2iB+A)) \tan(fx+e)^7}{7} + \frac{(-11A+5i(-2iA-B)+10iB) \tan(fx+e)^6}{6} \right)$
norman	$\frac{A a^3 c^6 \tan(fx+e)}{f} - \frac{(3iB a^3 c^6 + A a^3 c^6) \tan(fx+e)^3}{3f} - \frac{(5iA a^3 c^6 + B a^3 c^6) \tan(fx+e)^4}{4f} - \frac{(-iA a^3 c^6 + 3B a^3 c^6) \tan(fx+e)^5}{8f}$
parallelrisch	$-504iB \tan(fx+e)^3 a^3 c^6 - 72iB \tan(fx+e)^7 a^3 c^6 - 84iA \tan(fx+e)^6 a^3 c^6 - 189B \tan(fx+e)^8 a^3 c^6 - 504iB \tan(fx+e)^5 a^3 c^6$
parts	$\frac{(-8iB a^3 c^6 - 6A a^3 c^6) \left(\frac{\tan(fx+e)^3}{3} - \tan(fx+e) + \arctan(\tan(fx+e)) \right)}{f} + \frac{(-6iA a^3 c^6 - 6B a^3 c^6) \left(\frac{\tan(fx+e)^4}{4} - \frac{\tan(fx+e)^5}{5} \right)}{f}$

input `int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^6,x,method=_R
ETURNVERBOSE)`

output `32/63*c^6*a^3*(84*I*A*exp(6*I*(f*x+e))+84*B*exp(6*I*(f*x+e))+108*I*A*exp(4
I(f*x+e))-36*B*exp(4*I*(f*x+e))+27*I*A*exp(2*I*(f*x+e))-9*B*exp(2*I*(f*x
+e))+3*I*A-B)/f/(exp(2*I*(f*x+e))+1)^9`

3.690.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.44

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ict \tan(e + fx))^6 dx =$$

$$\frac{32 (84 (-iA - B) a^3 c^6 e^{(6i fx + 6ie)} + 36 (-3iA + B) a^3 c^6 e^{(4i fx + 4ie)} + 9 (-3iA + B) a^3 c^6 e^{(2i fx + 2ie)}}{63 (f e^{(18i fx + 18ie)} + 9 f e^{(16i fx + 16ie)} + 36 f e^{(14i fx + 14ie)} + 84 f e^{(12i fx + 12ie)} + 126 f e^{(10i fx + 10ie)} + 126 f e^{(8i fx + 8ie)} + 9 f e^{(6i fx + 6ie)} + 3 f e^{(4i fx + 4ie)} + f e^{(2i fx + 2ie)})}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^6,x, al
gorithm="fracas")`

output
$$\frac{-32/63*(84*(-I*A - B)*a^3*c^6*e^{(6*I*f*x + 6*I*e)} + 36*(-3*I*A + B)*a^3*c^6*e^{(4*I*f*x + 4*I*e)} + 9*(-3*I*A + B)*a^3*c^6*e^{(2*I*f*x + 2*I*e)} + (-3*I*A + B)*a^3*c^6)/(f*e^{(18*I*f*x + 18*I*e)} + 9*f*e^{(16*I*f*x + 16*I*e)} + 36*f*e^{(14*I*f*x + 14*I*e)} + 84*f*e^{(12*I*f*x + 12*I*e)} + 126*f*e^{(10*I*f*x + 10*I*e)} + 126*f*e^{(8*I*f*x + 8*I*e)} + 84*f*e^{(6*I*f*x + 6*I*e)} + 36*f*e^{(4*I*f*x + 4*I*e)} + 9*f*e^{(2*I*f*x + 2*I*e)} + f)$$

3.690.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 325 vs. $2(110) = 220$.

Time = 0.93 (sec) , antiderivative size = 325, normalized size of antiderivative = 2.41

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^6 dx$$

$$= \frac{96iAa^3c^6 - 32Ba^3c^6 + (864iAa^3c^6e^{2ie} - 288Ba^3c^6e^{2ie})e^{2ifx} + (3456iAa^3c^6e^{4ie} - 1152Ba^3c^6e^{4ie})e^{4ifx} + (2688iAa^3c^6e^{6ie} - 2688Ba^3c^6e^{6ie})e^{6ifx} + (1296iAa^3c^6e^{8ie} - 1296Ba^3c^6e^{8ie})e^{8ifx} + (5184iAa^3c^6e^{10ie} - 5184Ba^3c^6e^{10ie})e^{10ifx} + (15552iAa^3c^6e^{12ie} - 15552Ba^3c^6e^{12ie})e^{12ifx} + (373248iAa^3c^6e^{14ie} - 373248Ba^3c^6e^{14ie})e^{14ifx} + (848832iAa^3c^6e^{16ie} - 848832Ba^3c^6e^{16ie})e^{16ifx} + 63f^2e^{18ie}}{63fe^{18ie}e^{18ifx} + 567fe^{16ie}e^{16ifx} + 2268fe^{14ie}e^{14ifx} + 5292fe^{12ie}e^{12ifx} + 7938fe^{10ie}e^{10ifx} + 7938fe^{8ie}e^{8ifx} + 5184fe^{6ie}e^{6ifx} + 1296fe^{4ie}e^{4ifx} + 1296fe^{2ie}e^{2ifx} + 63f^2}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**6,x)`

output
$$(96*I*A*a**3*c**6 - 32*B*a**3*c**6 + (864*I*A*a**3*c**6*exp(2*I*e) - 288*B*a**3*c**6*exp(2*I*e))*exp(2*I*f*x) + (3456*I*A*a**3*c**6*exp(4*I*e) - 1152*B*a**3*c**6*exp(4*I*e))*exp(4*I*f*x) + (2688*I*A*a**3*c**6*exp(6*I*e) + 2688*B*a**3*c**6*exp(6*I*e))*exp(6*I*f*x)/(63*f*exp(18*I*e)*exp(18*I*f*x) + 567*f*exp(16*I*e)*exp(16*I*f*x) + 2268*f*exp(14*I*e)*exp(14*I*f*x) + 5292*f*exp(12*I*e)*exp(12*I*f*x) + 7938*f*exp(10*I*e)*exp(10*I*f*x) + 7938*f*exp(8*I*e)*exp(8*I*f*x) + 5292*f*exp(6*I*e)*exp(6*I*f*x) + 2268*f*exp(4*I*e)*exp(4*I*f*x) + 567*f*exp(2*I*e)*exp(2*I*f*x) + 63*f)$$

3.690.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.43

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^6 dx =$$

$$\frac{-56iBa^3c^6 \tan(fx + e)^9 + 63(-iA + 3B)a^3c^6 \tan(fx + e)^8 + 72(3A + iB)a^3c^6 \tan(fx + e)^7 + 84(3A + iB)a^3c^6 \tan(fx + e)^6 + 56iBa^3c^6 \tan(fx + e)^5 + 63(-iA + 3B)a^3c^6 \tan(fx + e)^4 + 72(3A + iB)a^3c^6 \tan(fx + e)^3 + 84(3A + iB)a^3c^6 \tan(fx + e)^2 + 56iBa^3c^6 \tan(fx + e) + 63(-iA + 3B)a^3c^6 \tan(fx + e) + 72(3A + iB)a^3c^6 \tan(fx + e) + 84(3A + iB)a^3c^6}{63f^2e^{18ie}}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^6,x, algorithm="maxima")`

output
$$\frac{-1/504*(-56*I*B*a^3*c^6*\tan(f*x + e)^9 + 63*(-I*A + 3*B)*a^3*c^6*\tan(f*x + e)^8 + 72*(3*A + I*B)*a^3*c^6*\tan(f*x + e)^7 + 84*(I*A + 5*B)*a^3*c^6*\tan(f*x + e)^6 + 504*(A + I*B)*a^3*c^6*\tan(f*x + e)^5 + 126*(5*I*A + B)*a^3*c^6*\tan(f*x + e)^4 + 168*(A + 3*I*B)*a^3*c^6*\tan(f*x + e)^3 + 252*(3*I*A - B)*a^3*c^6*\tan(f*x + e)^2 - 504*A*a^3*c^6*\tan(f*x + e))/f$$

3.690.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 239 vs. $2(113) = 226$.

Time = 1.29 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.77

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^6 dx = \frac{32(-84i Aa^3c^6e^{(6i fx+6i e)} - 84 Ba^3c^6e^{(6i fx+6i e)} - 108i Aa^3c^6e^{(4i fx+4i e)} + 36 Ba^3c^6e^{(4i fx+4i e)} - 27i Aa^3c^6e^{(2i fx+2i e)} + 9 Ba^3c^6e^{(2i fx+2i e)} - 3i Aa^3c^6 + Ba^3c^6)/(fe^{(18i fx+18i e)} + 9 fe^{(16i fx+16i e)} + 36 fe^{(14i fx+14i e)} + 84 fe^{(12i fx+12i e)} + 126 fe^{(10i fx+10i e)} + 126 fe^{(8i fx+8i e)} + 84 fe^{(6i fx+6i e)} + 9 fe^{(4i fx+4i e)} + 36 fe^{(2i fx+2i e)} + f)$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^6,x, algorithm="giac")`

output
$$-32/63*(-84*I*A*a^3*c^6*e^{(6*I*f*x + 6*I*e)} - 84*B*a^3*c^6*e^{(6*I*f*x + 6*I*e)} - 108*I*A*a^3*c^6*e^{(4*I*f*x + 4*I*e)} + 36*B*a^3*c^6*e^{(4*I*f*x + 4*I*e)} - 27*I*A*a^3*c^6*e^{(2*I*f*x + 2*I*e)} + 9*B*a^3*c^6*e^{(2*I*f*x + 2*I*e)} - 3*I*A*a^3*c^6 + B*a^3*c^6)/(f*e^{(18*I*f*x + 18*I*e)} + 9*f*e^{(16*I*f*x + 16*I*e)} + 36*f*e^{(14*I*f*x + 14*I*e)} + 84*f*e^{(12*I*f*x + 12*I*e)} + 126*f*e^{(10*I*f*x + 10*I*e)} + 126*f*e^{(8*I*f*x + 8*I*e)} + 84*f*e^{(6*I*f*x + 6*I*e)} + 36*f*e^{(4*I*f*x + 4*I*e)} + 9*f*e^{(2*I*f*x + 2*I*e)} + f)$$

3.690.9 Mupad [B] (verification not implemented)

Time = 8.72 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.54

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^6 dx$$

$$= \frac{A a^3 c^6 \tan(e + fx) + \frac{a^3 c^6 \tan(e+fx)^3 (-3B+A 1i) 1i}{3} + a^3 c^6 \tan(e + fx)^5 (-B + A 1i) 1i - \frac{a^3 c^6 \tan(e+fx)^4 (5A-}{4}}$$

```
input int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i)^6,x)
```

```
output ((a^3*c^6*tan(e + f*x)^3*(A*1i - 3*B)*1i)/3 - (a^3*c^6*tan(e + f*x)^2*(3*A + B*1i)*1i)/2 + a^3*c^6*tan(e + f*x)^5*(A*1i - B)*1i - (a^3*c^6*tan(e + f*x)^4*(5*A - B*1i)*1i)/4 + (a^3*c^6*tan(e + f*x)^7*(A*3i - B)*1i)/7 + A*a^3*c^6*tan(e + f*x) - (a^3*c^6*tan(e + f*x)^6*(A - B*5i)*1i)/6 + (a^3*c^6*tan(e + f*x)^8*(A + B*3i)*1i)/8 + (B*a^3*c^6*tan(e + f*x)^9*1i)/9)/f
```

3.691 $\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx))^5 dx$

3.691.1 Optimal result	6463
3.691.2 Mathematica [A] (verified)	6463
3.691.3 Rubi [A] (verified)	6464
3.691.4 Maple [A] (verified)	6466
3.691.5 Fricas [A] (verification not implemented)	6466
3.691.6 Sympy [B] (verification not implemented)	6467
3.691.7 Maxima [A] (verification not implemented)	6467
3.691.8 Giac [B] (verification not implemented)	6468
3.691.9 Mupad [B] (verification not implemented)	6469

3.691.1 Optimal result

Integrand size = 41, antiderivative size = 135

$$\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))(c - ictan(e + fx))^5 dx$$

$$= \frac{4a^3(iA + B)c^5(1 - i \tan(e + fx))^5}{5f} - \frac{2a^3(iA + 2B)c^5(1 - i \tan(e + fx))^6}{3f}$$

$$+ \frac{a^3(iA + 5B)c^5(1 - i \tan(e + fx))^7}{7f} - \frac{a^3Bc^5(1 - i \tan(e + fx))^8}{8f}$$

```
output 4/5*a^3*(I*A+B)*c^5*(1-I*tan(f*x+e))^5/f-2/3*a^3*(I*A+2*B)*c^5*(1-I*tan(f*x+e))^6/f+1/7*a^3*(I*A+5*B)*c^5*(1-I*tan(f*x+e))^7/f-1/8*a^3*B*c^5*(1-I*tan(f*x+e))^8/f
```

3.691.2 Mathematica [A] (verified)

Time = 5.69 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.73

$$\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))(c - ictan(e + fx))^5 dx$$

$$= \frac{a^3c^5 \sec^8(e + fx)(140(-iA + B) \cos(2(e + fx)) + 84(A - iB) \sin(2(e + fx)) + (4A + iB)(-35i + 28 \sin(2(e + fx))))}{840f}$$

input `Integrate[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^5,x]`

output `(a^3*c^5*Sec[e + f*x]^8*(140*((-I)*A + B)*Cos[2*(e + f*x)] + 84*(A - I*B)*Sin[2*(e + f*x)] + (4*A + I*B)*(-35*I + 28*Sin[4*(e + f*x)] + 8*Sin[6*(e + f*x)] + Sin[8*(e + f*x)])))/(840*f)`

3.691.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.81, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))^3 (c - ictan(e + fx))^5 (A + B \tan(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (a + ia \tan(e + fx))^3 (c - ictan(e + fx))^5 (A + B \tan(e + fx)) dx$$

$$\downarrow \text{4071}$$

$$\frac{ac \int a^2 c^4 (1 - i \tan(e + fx))^4 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) d \tan(e + fx)}{f}$$

$$\downarrow \text{27}$$

$$\frac{a^3 c^5 \int (1 - i \tan(e + fx))^4 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) d \tan(e + fx)}{f}$$

$$\downarrow \text{86}$$

$$\frac{a^3 c^5 \int (iB(1 - i \tan(e + fx))^7 + (A - 5iB)(1 - i \tan(e + fx))^6 - 4(A - 2iB)(1 - i \tan(e + fx))^5 + 4(A - iB)(1 - i \tan(e + fx))^4 - 4iA(1 - i \tan(e + fx))^3 + 4iA(1 - i \tan(e + fx))^2 - 4iA(1 - i \tan(e + fx)) + 4iA) d \tan(e + fx)}{f}$$

$$\downarrow \text{2009}$$

$$\frac{a^3 c^5 (\frac{1}{7}(5B + iA)(1 - i \tan(e + fx))^7 - \frac{2}{3}(2B + iA)(1 - i \tan(e + fx))^6 + \frac{4}{5}(B + iA)(1 - i \tan(e + fx))^5 - \frac{1}{8}B(1 - i \tan(e + fx))^4 + \frac{1}{4}iA(1 - i \tan(e + fx))^3 - \frac{1}{4}iA(1 - i \tan(e + fx))^2 + \frac{1}{4}iA(1 - i \tan(e + fx)) - \frac{1}{4}iA)}{f}$$

3.691. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ictan(e + fx))^5 dx$

input `Int[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^5, x]`

output `(a^3*c^5*((4*(I*A + B)*(1 - I*Tan[e + f*x])^5)/5 - (2*(I*A + 2*B)*(1 - I*Tan[e + f*x])^6)/3 + ((I*A + 5*B)*(1 - I*Tan[e + f*x])^7)/7 - (B*(1 - I*Tan[e + f*x])^8)/8))/f`

3.691.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.691.4 Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.80

method	result
risch	$\frac{32c^5 a^3 (84iA e^{6i(fx+e)} + 84B e^{6i(fx+e)} + 112iA e^{4i(fx+e)} - 28B e^{4i(fx+e)} + 32iA e^{2i(fx+e)} - 8B e^{2i(fx+e)} + 4iA - B)}{105f(e^{2i(fx+e)} + 1)^8}$
derivativedivides	$\frac{c^5 a^3 \left(-\frac{B \tan(fx+e)^8}{8} - \frac{(2iB+A) \tan(fx+e)^7}{7} - \frac{(-7B-2iA+4i(-2iB+A)) \tan(fx+e)^6}{6} - \frac{(-7A+4i(-2iA-B)+8iB) \tan(fx+e)^5}{5} \right)}{f}$
default	$\frac{c^5 a^3 \left(-\frac{B \tan(fx+e)^8}{8} - \frac{(2iB+A) \tan(fx+e)^7}{7} - \frac{(-7B-2iA+4i(-2iB+A)) \tan(fx+e)^6}{6} - \frac{(-7A+4i(-2iA-B)+8iB) \tan(fx+e)^5}{5} \right)}{f}$
norman	$\frac{A a^3 c^5 \tan(fx+e)}{f} - \frac{(2iA a^3 c^5 + B a^3 c^5) \tan(fx+e)^6}{6f} - \frac{(2iB a^3 c^5 + A a^3 c^5) \tan(fx+e)^7}{7f} - \frac{(4iB a^3 c^5 + A a^3 c^5) \tan(fx+e)^8}{5f}$
parallelrisch	$-\frac{240iB \tan(fx+e)^7 a^3 c^5 + 105B \tan(fx+e)^8 a^3 c^5 + 280iA \tan(fx+e)^6 a^3 c^5 + 120A \tan(fx+e)^7 a^3 c^5 + 672iB \tan(fx+e)^8 a^3 c^5}{f}$
parts	$\frac{(-6iA a^3 c^5 + 2B a^3 c^5) \left(\frac{\tan(fx+e)^2}{2} - \frac{\ln(1+\tan(fx+e)^2)}{2} \right)}{f} + \frac{(-6iB a^3 c^5 - 2A a^3 c^5) \left(\frac{\tan(fx+e)^5}{5} - \frac{\tan(fx+e)^3}{3} + \tan(fx+e) \right)}{f}$

input `int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5,x,method=_R
ETURNVERBOSE)`

output `32/105*c^5*a^3*(84*I*A*exp(6*I*(f*x+e))+84*B*exp(6*I*(f*x+e))+112*I*A*exp(4*I*(f*x+e))-28*B*exp(4*I*(f*x+e))+32*I*A*exp(2*I*(f*x+e))-8*B*exp(2*I*(f*x+e))+4*I*A-B)/f/(exp(2*I*(f*x+e))+1)^8`

3.691.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.35

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ict \tan(e + fx))^5 dx =$$

$$\frac{32 (84 (-iA - B) a^3 c^5 e^{(6i fx + 6ie)} + 28 (-4iA + B) a^3 c^5 e^{(4i fx + 4ie)} + 8 (-4iA + B) a^3 c^5 e^{(2i fx + 2ie)}}{105 (fe^{(16i fx + 16ie)} + 8 fe^{(14i fx + 14ie)} + 28 fe^{(12i fx + 12ie)} + 56 fe^{(10i fx + 10ie)} + 70 fe^{(8i fx + 8ie)} + 56 fe^{(6i fx + 6ie)})}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5,x, algorithm="fricas")`

output
$$\frac{-32/105(84(-I*A - B)*a^3*c^5*e^{(6*I*f*x + 6*I*e)} + 28*(-4*I*A + B)*a^3*c^5*e^{(4*I*f*x + 4*I*e)} + 8*(-4*I*A + B)*a^3*c^5*e^{(2*I*f*x + 2*I*e)} + (-4*I*A + B)*a^3*c^5)/(f*e^{(16*I*f*x + 16*I*e)} + 8*f*e^{(14*I*f*x + 14*I*e)} + 2*8*f*e^{(12*I*f*x + 12*I*e)} + 56*f*e^{(10*I*f*x + 10*I*e)} + 70*f*e^{(8*I*f*x + 8*I*e)} + 56*f*e^{(6*I*f*x + 6*I*e)} + 28*f*e^{(4*I*f*x + 4*I*e)} + 8*f*e^{(2*I*f*x + 2*I*e)} + f)$$

3.691.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 306 vs. $2(110) = 220$.

Time = 0.83 (sec) , antiderivative size = 306, normalized size of antiderivative = 2.27

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^5 dx$$

$$= \frac{128iAa^3c^5 - 32Ba^3c^5 + (1024iAa^3c^5e^{2ie} - 256Ba^3c^5e^{2ie})e^{2ifx} + (3584iAa^3c^5e^{4ie} - 896Ba^3c^5e^{4ie})e^{4ifx} + \dots}{105fe^{16ie}e^{16ifx} + 840fe^{14ie}e^{14ifx} + 2940fe^{12ie}e^{12ifx} + 5880fe^{10ie}e^{10ifx} + 7350fe^{8ie}e^{8ifx} + 5880fe^{6ie}e^{6ifx} + \dots}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**5,x)`

output
$$(128*I*A*a**3*c**5 - 32*B*a**3*c**5 + (1024*I*A*a**3*c**5*\exp(2*I*e) - 256*B*a**3*c**5*\exp(2*I*e))*\exp(2*I*f*x) + (3584*I*A*a**3*c**5*\exp(4*I*e) - 896*B*a**3*c**5*\exp(4*I*e))*\exp(4*I*f*x) + (2688*I*A*a**3*c**5*\exp(6*I*e) + 2688*B*a**3*c**5*\exp(6*I*e))*\exp(6*I*f*x))/(105*f*\exp(16*I*e)*\exp(16*I*f*x) + 840*f*\exp(14*I*e)*\exp(14*I*f*x) + 2940*f*\exp(12*I*e)*\exp(12*I*f*x) + 5880*f*\exp(10*I*e)*\exp(10*I*f*x) + 7350*f*\exp(8*I*e)*\exp(8*I*f*x) + 5880*f*\exp(6*I*e)*\exp(6*I*f*x) + 2940*f*\exp(4*I*e)*\exp(4*I*f*x) + 840*f*\exp(2*I*e)*\exp(2*I*f*x) + 105*f)$$

3.691.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.23

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^5 dx =$$

$$\frac{105 Ba^3c^5 \tan^8(fx + e) + 120 (A + 2iB)a^3c^5 \tan^7(fx + e) - 140(-2iA - B)a^3c^5 \tan^6(fx + e) + 160(-2iA - B)a^3c^5 \tan^5(fx + e) - 120(A + 2iB)a^3c^5 \tan^4(fx + e) + 105Ba^3c^5 \tan^3(fx + e)}{105f^6 \exp(16ie) \exp(16ifx) + 840f^5 \exp(14ie) \exp(14ifx) + 2940f^4 \exp(12ie) \exp(12ifx) + 5880f^3 \exp(10ie) \exp(10ifx) + 7350f^2 \exp(8ie) \exp(8ifx) + 5880f \exp(6ie) \exp(6ifx) + 105 \exp(4ie) \exp(4ifx) + 105 \exp(2ie) \exp(2ifx) + 105}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5,x, algorithm="maxima")`

output
$$-1/840*(105*B*a^3*c^5*\tan(f*x + e)^8 + 120*(A + 2*I*B)*a^3*c^5*\tan(f*x + e)^7 - 140*(-2*I*A - B)*a^3*c^5*\tan(f*x + e)^6 + 168*(A + 4*I*B)*a^3*c^5*\tan(f*x + e)^5 - 210*(-4*I*A + B)*a^3*c^5*\tan(f*x + e)^4 - 280*(A - 2*I*B)*a^3*c^5*\tan(f*x + e)^3 - 420*(-2*I*A + B)*a^3*c^5*\tan(f*x + e)^2 - 840*A*a^3*c^5*\tan(f*x + e))/f$$

3.691.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 227 vs. $2(113) = 226$.

Time = 1.20 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.68

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^5 dx = \frac{32(-84i Aa^3c^5e^{(6ifx+6ie)} - 84Ba^3c^5e^{(6ifx+6ie)} - 112i Aa^3c^5e^{(4ifx+4ie)} + 28Ba^3c^5e^{(4ifx+4ie)} - 32i Aa^3c^5e^{(2ifx+2ie)} + 8Ba^3c^5e^{(2ifx+2ie)} - 4iAa^3c^5 + Ba^3c^5)/(fe^{(16ifx+16ie)} + 8fe^{(14ifx+14ie)} + 28fe^{(12ifx+12ie)} + 56fe^{(10ifx+10ie)} + 70fe^{(8ifx+8ie)} + 56fe^{(6ifx+6ie)} + 28fe^{(4ifx+4ie)} + 8fe^{(2ifx+2ie)} + f)$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5,x, algorithm="giac")`

output
$$-32/105*(-84*I*A*a^3*c^5*e^{(6*I*f*x + 6*I*e)} - 84*B*a^3*c^5*e^{(6*I*f*x + 6*I*e)} - 112*I*A*a^3*c^5*e^{(4*I*f*x + 4*I*e)} + 28*B*a^3*c^5*e^{(4*I*f*x + 4*I*e)} - 32*I*A*a^3*c^5*e^{(2*I*f*x + 2*I*e)} + 8*B*a^3*c^5*e^{(2*I*f*x + 2*I*e)} - 4*I*A*a^3*c^5 + B*a^3*c^5)/(f*e^{(16*I*f*x + 16*I*e)} + 8*f*e^{(14*I*f*x + 14*I*e)} + 28*f*e^{(12*I*f*x + 12*I*e)} + 56*f*e^{(10*I*f*x + 10*I*e)} + 70*f*e^{(8*I*f*x + 8*I*e)} + 56*f*e^{(6*I*f*x + 6*I*e)} + 28*f*e^{(4*I*f*x + 4*I*e)} + 8*f*e^{(2*I*f*x + 2*I*e)} + f)$$

3.691. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^5 dx$

3.691.9 Mupad [B] (verification not implemented)

Time = 8.60 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.29

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^5 dx =$$

$$-\frac{\frac{a^3 c^5 \tan(e+fx)^2 (-B+A2i)}{2} + \frac{a^3 c^5 \tan(e+fx)^4 (-B+A4i)}{4} - A a^3 c^5 \tan(e + fx) - \frac{a^3 c^5 \tan(e+fx)^3 (A-B2i)}{3} + \frac{a^3 c^5 \tan(e+fx)^5 (A+B2i)}{5}}{f}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i)^5,x)`

output `-((a^3*c^5*tan(e + f*x)^2*(A*2i - B))/2 + (a^3*c^5*tan(e + f*x)^4*(A*4i - B))/4 - A*a^3*c^5*tan(e + f*x) - (a^3*c^5*tan(e + f*x)^3*(A - B*2i))/3 + (a^3*c^5*tan(e + f*x)^6*(A*2i + B))/6 + (a^3*c^5*tan(e + f*x)^5*(A + B*4i))/5 + (a^3*c^5*tan(e + f*x)^7*(A + B*2i))/7 + (B*a^3*c^5*tan(e + f*x)^8)/8)/f`

3.692 $\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx))^4 dx$

3.692.1 Optimal result	6470
3.692.2 Mathematica [A] (verified)	6470
3.692.3 Rubi [A] (verified)	6471
3.692.4 Maple [A] (verified)	6473
3.692.5 Fricas [A] (verification not implemented)	6473
3.692.6 Sympy [B] (verification not implemented)	6474
3.692.7 Maxima [A] (verification not implemented)	6474
3.692.8 Giac [A] (verification not implemented)	6475
3.692.9 Mupad [B] (verification not implemented)	6475

3.692.1 Optimal result

Integrand size = 41, antiderivative size = 132

$$\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))(c - ictan(e + fx))^4 dx$$

$$= \frac{a^3(iA + B)c^4(1 - i \tan(e + fx))^4}{f} - \frac{4a^3(iA + 2B)c^4(1 - i \tan(e + fx))^5}{5f}$$

$$+ \frac{a^3(iA + 5B)c^4(1 - i \tan(e + fx))^6}{6f} - \frac{a^3Bc^4(1 - i \tan(e + fx))^7}{7f}$$

```
output a^3*(I*A+B)*c^4*(1-I*tan(f*x+e))^4/f-4/5*a^3*(I*A+2*B)*c^4*(1-I*tan(f*x+e))^5/f+1/6*a^3*(I*A+5*B)*c^4*(1-I*tan(f*x+e))^6/f-1/7*a^3*B*c^4*(1-I*tan(f*x+e))^7/f
```

3.692.2 Mathematica [A] (verified)

Time = 2.09 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.99

$$\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))(c - ictan(e + fx))^4 dx$$

$$= \frac{a^3c^4(7iA + 29B + 210A \tan(e + fx) + 105(-iA + B) \tan^2(e + fx) + 70(2A - iB) \tan^3(e + fx) + 105(-$$

input `Integrate[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^4,x]`

output `(a^3*c^4*((7*I)*A + 29*B + 210*A*Tan[e + f*x] + 105*((-I)*A + B)*Tan[e + f*x]^2 + 70*(2*A - I*B)*Tan[e + f*x]^3 + 105*((-I)*A + B)*Tan[e + f*x]^4 + 42*(A - (2*I)*B)*Tan[e + f*x]^5 + 35*((-I)*A + B)*Tan[e + f*x]^6 - (30*I)*B*Tan[e + f*x]^7))/(210*f)`

3.692.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.80, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(e + fx))^3 (c - ictan(e + fx))^4 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(e + fx))^3 (c - ictan(e + fx))^4 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int a^2 c^3 (1 - i \tan(e + fx))^3 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^3 c^4 \int (1 - i \tan(e + fx))^3 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{86} \\
 & \frac{a^3 c^4 \int (iB(1 - i \tan(e + fx))^6 + (A - 5iB)(1 - i \tan(e + fx))^5 - 4(A - 2iB)(1 - i \tan(e + fx))^4 + 4(A - iB)(1 - i \tan(e + fx))^3)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^3 c^4 (\frac{1}{6}(5B + iA)(1 - i \tan(e + fx))^6 - \frac{4}{5}(2B + iA)(1 - i \tan(e + fx))^5 + (B + iA)(1 - i \tan(e + fx))^4 - \frac{1}{7}B(1 - i \tan(e + fx))^3)}{f}
 \end{aligned}$$

3.692. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ictan(e + fx))^4 dx$

input `Int[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^4, x]`

output `(a^3*c^4*((I*A + B)*(1 - I*Tan[e + f*x])^4 - (4*(I*A + 2*B)*(1 - I*Tan[e + f*x])^5)/5 + ((I*A + 5*B)*(1 - I*Tan[e + f*x])^6)/6 - (B*(1 - I*Tan[e + f*x])^7)/7))/f`

3.692.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.692.4 Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.82

method	result
risch	$\frac{16c^4 a^3 (105iA e^{6i(fx+e)} + 105B e^{6i(fx+e)} + 147iA e^{4i(fx+e)} - 21B e^{4i(fx+e)} + 49iA e^{2i(fx+e)} - 7B e^{2i(fx+e)} + 7iA - B)}{105f(e^{2i(fx+e)} + 1)^7}$
derivativedivides	$- \frac{ic^4 a^3 \left(\frac{B \tan(fx+e)^7}{7} + \frac{(iB+A) \tan(fx+e)^6}{6} + \frac{(-4B-2iA+3i(-2iB+A)) \tan(fx+e)^5}{5} + \frac{(-4A+3i(-2iA-B)+5iB) \tan(fx+e)^4}{4} \right)}{f}$
default	$- \frac{ic^4 a^3 \left(\frac{B \tan(fx+e)^7}{7} + \frac{(iB+A) \tan(fx+e)^6}{6} + \frac{(-4B-2iA+3i(-2iB+A)) \tan(fx+e)^5}{5} + \frac{(-4A+3i(-2iA-B)+5iB) \tan(fx+e)^4}{4} \right)}{f}$
norman	$\frac{A a^3 c^4 \tan(fx+e)}{f} + \frac{(-iA a^3 c^4 + B a^3 c^4) \tan(fx+e)^2}{2f} + \frac{(-iA a^3 c^4 + B a^3 c^4) \tan(fx+e)^4}{2f} + \frac{(-iA a^3 c^4 + B a^3 c^4) \tan(fx+e)^6}{6f}$
parallelrisch	$- \frac{30iB a^3 c^4 \tan(fx+e)^7 + 35iA \tan(fx+e)^6 a^3 c^4 + 84iB \tan(fx+e)^5 a^3 c^4 - 35B \tan(fx+e)^6 a^3 c^4 + 105iA \tan(fx+e)^4 a^3 c^4}{f}$
parts	$\frac{(-3iA a^3 c^4 + 3B a^3 c^4) \left(\frac{\tan(fx+e)^2}{2} - \frac{\ln(1+\tan(fx+e)^2)}{2} \right)}{f} + \frac{(-3iA a^3 c^4 + 3B a^3 c^4) \left(\frac{\tan(fx+e)^4}{4} - \frac{\tan(fx+e)^2}{2} + \frac{\ln(1+\tan(fx+e)^2)}{2} \right)}{f}$

input `int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4,x,method=_R
ETURNVERBOSE)`

output `16/105*c^4*a^3*(105*I*A*exp(6*I*(f*x+e))+105*B*exp(6*I*(f*x+e))+147*I*A*exp
p(4*I*(f*x+e))-21*B*exp(4*I*(f*x+e))+49*I*A*exp(2*I*(f*x+e))-7*B*exp(2*I*(
f*x+e))+7*I*A-B)/f/(exp(2*I*(f*x+e))+1)^7`

3.692.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.29

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ict \tan(e + fx))^4 dx =$$

$$- \frac{16 (105 (-iA - B) a^3 c^4 e^{(6i fx + 6i e)} + 21 (-7iA + B) a^3 c^4 e^{(4i fx + 4i e)} + 7 (-7iA + B) a^3 c^4 e^{(2i fx + 2i e)}}{105 (f e^{(14i fx + 14i e)} + 7 f e^{(12i fx + 12i e)} + 21 f e^{(10i fx + 10i e)} + 35 f e^{(8i fx + 8i e)} + 35 f e^{(6i fx + 6i e)} + 21 f e^{(4i fx + 4i e)})}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4,x, al
gorithm="fricas")`

output
$$\frac{-16/105*(105*(-I*A - B)*a^3*c^4*e^{(6*I*f*x + 6*I*e)} + 21*(-7*I*A + B)*a^3*c^4*e^{(4*I*f*x + 4*I*e)} + 7*(-7*I*A + B)*a^3*c^4*e^{(2*I*f*x + 2*I*e)} + (-7*I*A + B)*a^3*c^4)/(f*e^{(14*I*f*x + 14*I*e)} + 7*f*e^{(12*I*f*x + 12*I*e)} + 21*f*e^{(10*I*f*x + 10*I*e)} + 35*f*e^{(8*I*f*x + 8*I*e)} + 35*f*e^{(6*I*f*x + 6*I*e)} + 21*f*e^{(4*I*f*x + 4*I*e)} + 7*f*e^{(2*I*f*x + 2*I*e)} + f)$$

3.692.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 287 vs. $2(107) = 214$.

Time = 0.64 (sec) , antiderivative size = 287, normalized size of antiderivative = 2.17

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^4 dx$$

$$= \frac{112iAa^3c^4 - 16Ba^3c^4 + (784iAa^3c^4e^{2ie} - 112Ba^3c^4e^{2ie})e^{2ifx} + (2352iAa^3c^4e^{4ie} - 336Ba^3c^4e^{4ie})e^{4ifx} + (1680iAa^3c^4e^{6ie} - 168Ba^3c^4e^{6ie})e^{6ifx} + (735f^2e^{8ie}e^{8ifx} + 2205fe^{10ie}e^{10ifx} + 3675f^2e^{12ie}e^{12ifx} + 2205fe^{14ie}e^{14ifx})}{105fe^{14ie}e^{14ifx} + 735f^2e^{12ie}e^{12ifx} + 2205fe^{10ie}e^{10ifx} + 3675f^2e^{8ie}e^{8ifx} + 3675fe^{6ie}e^{6ifx} + 2205fe^{4ie}e^{4ifx} + 735f^2e^{2ie}e^{2ifx} + 105f}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**4,x)`

output
$$\frac{(112*I*A*a**3*c**4 - 16*B*a**3*c**4 + (784*I*A*a**3*c**4*exp(2*I*e) - 112*B*a**3*c**4*exp(2*I*e))*exp(2*I*f*x) + (2352*I*A*a**3*c**4*exp(4*I*e) - 336*B*a**3*c**4*exp(4*I*e))*exp(4*I*f*x) + (1680*I*A*a**3*c**4*exp(6*I*e) + 1680*B*a**3*c**4*exp(6*I*e))*exp(6*I*f*x))/(105*f*exp(14*I*e)*exp(14*I*f*x) + 735*f*exp(12*I*e)*exp(12*I*f*x) + 2205*f*exp(10*I*e)*exp(10*I*f*x) + 3675*f*exp(8*I*e)*exp(8*I*f*x) + 3675*f*exp(6*I*e)*exp(6*I*f*x) + 2205*f*exp(4*I*e)*exp(4*I*f*x) + 735*f*exp(2*I*e)*exp(2*I*f*x) + 105*f)$$

3.692.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.14

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^4 dx =$$

$$\frac{30iBa^3c^4 \tan^7(fx + e) + 35(iA - B)a^3c^4 \tan^6(fx + e) - 42(A - 2iB)a^3c^4 \tan^5(fx + e) + 105(iA - B)a^3c^4 \tan^4(fx + e) - 35(iA + B)a^3c^4 \tan^3(fx + e) + 30iBa^3c^4 \tan^2(fx + e) - 105(iA - B)a^3c^4 \tan(fx + e) + 105(iA + B)a^3c^4}{105f^2e^{14ie}e^{14ifx} + 735f^2e^{12ie}e^{12ifx} + 2205fe^{10ie}e^{10ifx} + 3675f^2e^{8ie}e^{8ifx} + 3675fe^{6ie}e^{6ifx} + 2205fe^{4ie}e^{4ifx} + 735f^2e^{2ie}e^{2ifx} + 105f}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4,x, algorithm="maxima")`

3.692. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^4 dx$

output
$$\frac{-1/210*(30*I*B*a^3*c^4*\tan(f*x + e)^7 + 35*(I*A - B)*a^3*c^4*\tan(f*x + e)^6 - 42*(A - 2*I*B)*a^3*c^4*\tan(f*x + e)^5 + 105*(I*A - B)*a^3*c^4*\tan(f*x + e)^4 - 70*(2*A - I*B)*a^3*c^4*\tan(f*x + e)^3 + 105*(I*A - B)*a^3*c^4*\tan(f*x + e)^2 - 210*A*a^3*c^4*\tan(f*x + e))/f$$

3.692.8 Giac [A] (verification not implemented)

Time = 0.96 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.63

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^4 dx =$$

$$-\frac{16(-105i Aa^3c^4e^{(6ifx+6ie)} - 105Ba^3c^4e^{(6ifx+6ie)} - 147i Aa^3c^4e^{(4ifx+4ie)} + 21Ba^3c^4e^{(4ifx+4ie)} - 49i Aa^3c^4e^{(2ifx+2ie)} + 7Ba^3c^4e^{(2ifx+2ie)} - 7I*A*a^3*c^4 + B*a^3*c^4)/(f*e^{(14ifx+14ie)} + 7*f*e^{(12ifx+12ie)} + 21*f*e^{(10ifx+10ie)} + 35*f*e^{(8ifx+8ie)} + 35*f*e^{(6ifx+6ie)} + f)$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4,x, algorithm="giac")`

output
$$\frac{-16/105*(-105*I*A*a^3*c^4*e^{(6*I*f*x + 6*I*e)} - 105*B*a^3*c^4*e^{(6*I*f*x + 6*I*e)} - 147*I*A*a^3*c^4*e^{(4*I*f*x + 4*I*e)} + 21*B*a^3*c^4*e^{(4*I*f*x + 4*I*e)} - 49*I*A*a^3*c^4*e^{(2*I*f*x + 2*I*e)} + 7*B*a^3*c^4*e^{(2*I*f*x + 2*I*e)} - 7*I*A*a^3*c^4 + B*a^3*c^4)/(f*e^{(14*I*f*x + 14*I*e)} + 7*f*e^{(12*I*f*x + 12*I*e)} + 21*f*e^{(10*I*f*x + 10*I*e)} + 35*f*e^{(8*I*f*x + 8*I*e)} + 35*f*e^{(6*I*f*x + 6*I*e)} + 21*f*e^{(4*I*f*x + 4*I*e)} + 7*f*e^{(2*I*f*x + 2*I*e)} + f)$$

3.692.9 Mupad [B] (verification not implemented)

Time = 8.48 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.18

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^4 dx =$$

$$-\frac{a^3c^4 \tan(e+fx)^5 (2B+A \operatorname{li}) \operatorname{li}}{5} - Aa^3c^4 \tan(e+fx) + \frac{a^3c^4 \tan(e+fx)^2 (A+B \operatorname{li}) \operatorname{li}}{2} + \frac{a^3c^4 \tan(e+fx)^3 (B+A2i) \operatorname{li}}{3} + \frac{a^3c^4}{f}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i)^4,x)`

output $-\left(\frac{a^3 c^4 \tan(e + f x)^5 (A i + 2 B) i}{5} - A a^3 c^4 \tan(e + f x) + \frac{a^3 c^4 \tan(e + f x)^2 (A + B i) i}{2} + \frac{a^3 c^4 \tan(e + f x)^3 (A^2 i + B) i}{3} + \frac{a^3 c^4 \tan(e + f x)^4 (A + B i) i}{2} + \frac{a^3 c^4 \tan(e + f x)^6 (A + B i) i}{6} + \frac{B a^3 c^4 \tan(e + f x)^7 i}{7}\right) / f$

3.693 $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^3 dx$

3.693.1 Optimal result	6477
3.693.2 Mathematica [A] (verified)	6477
3.693.3 Rubi [A] (verified)	6478
3.693.4 Maple [A] (verified)	6480
3.693.5 Fricas [C] (verification not implemented)	6480
3.693.6 Sympy [C] (verification not implemented)	6481
3.693.7 Maxima [A] (verification not implemented)	6481
3.693.8 Giac [B] (verification not implemented)	6482
3.693.9 Mupad [B] (verification not implemented)	6483

3.693.1 Optimal result

Integrand size = 41, antiderivative size = 84

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^3 dx$$

$$= \frac{a^3 B c^3 \sec^6(e + fx)}{6f} + \frac{a^3 A c^3 \tan(e + fx)}{f} + \frac{2a^3 A c^3 \tan^3(e + fx)}{3f} + \frac{a^3 A c^3 \tan^5(e + fx)}{5f}$$

```
output 1/6*a^3*B*c^3*sec(f*x+e)^6/f+a^3*A*c^3*tan(f*x+e)/f+2/3*a^3*A*c^3*tan(f*x+e)^3/f+1/5*a^3*A*c^3*tan(f*x+e)^5/f
```

3.693.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.77

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^3 dx$$

$$= \frac{a^3 B c^3 \sec^6(e + fx)}{6f} + \frac{a^3 A c^3 (\tan(e + fx) + \frac{2}{3} \tan^3(e + fx) + \frac{1}{5} \tan^5(e + fx))}{f}$$

```
input Integrate[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^3,x]
```

output $(a^3 B c^3 \text{Sec}[e + f x]^6)/(6 f) + (a^3 A c^3 (\text{Tan}[e + f x] + (2 \text{Tan}[e + f x]^3)/3 + \text{Tan}[e + f x]^5/5))/f$

3.693.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.73, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.171$, Rules used = {3042, 4071, 27, 82, 455, 210, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(e + fx))^3 (c - ictan(e + fx))^3 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(e + fx))^3 (c - ictan(e + fx))^3 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int a^2 c^2 (1 - i \tan(e + fx))^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^3 c^3 \int (1 - i \tan(e + fx))^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{82} \\
 & \frac{a^3 c^3 \int (A + B \tan(e + fx)) (\tan^2(e + fx) + 1)^2 d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{455} \\
 & \frac{a^3 c^3 \left(A \int (\tan^2(e + fx) + 1)^2 d \tan(e + fx) + \frac{1}{6} B (\tan^2(e + fx) + 1)^3 \right)}{f} \\
 & \quad \downarrow \text{210} \\
 & \frac{a^3 c^3 \left(A \int (\tan^4(e + fx) + 2 \tan^2(e + fx) + 1) d \tan(e + fx) + \frac{1}{6} B (\tan^2(e + fx) + 1)^3 \right)}{f} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

3.693. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ictan(e + fx))^3 dx$

$$\frac{a^3 c^3 \left(A \left(\frac{1}{5} \tan^5(e + fx) + \frac{2}{3} \tan^3(e + fx) + \tan(e + fx) \right) + \frac{1}{6} B (\tan^2(e + fx) + 1)^3 \right)}{f}$$

input `Int[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^3, x]`

output `(a^3*c^3*((B*(1 + Tan[e + f*x]^2)^3)/6 + A*(Tan[e + f*x] + (2*Tan[e + f*x]^3)/3 + Tan[e + f*x]^5/5)))/f`

3.693.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 82 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 210 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^2)^p, x], x] /; FreeQ[{a, b}, x] && IGtQ[p, 0]`

rule 455 `Int[((c_) + (d_.)*(x_))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Simp[d*((a + b*x^2)^(p + 1)/(2*b*(p + 1))), x] + Simp[c Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, p}, x] && !LeQ[p, -1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x],
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.693.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.89

method	result
derivativedivides	$\frac{c^3 a^3 \left(\frac{B \tan(fx+e)^6}{6} + \frac{A \tan(fx+e)^5}{5} + \frac{B \tan(fx+e)^4}{2} + \frac{2A \tan(fx+e)^3}{3} + \frac{B \tan(fx+e)^2}{2} + A \tan(fx+e) \right)}{f}$
default	$\frac{c^3 a^3 \left(\frac{B \tan(fx+e)^6}{6} + \frac{A \tan(fx+e)^5}{5} + \frac{B \tan(fx+e)^4}{2} + \frac{2A \tan(fx+e)^3}{3} + \frac{B \tan(fx+e)^2}{2} + A \tan(fx+e) \right)}{f}$
risch	$\frac{16c^3 a^3 (10iA e^{6i(fx+e)} + 10B e^{6i(fx+e)} + 15iA e^{4i(fx+e)} + 6iA e^{2i(fx+e)} + iA)}{15f (e^{2i(fx+e)} + 1)^6}$
parallelrisc	$\frac{5B a^3 c^3 \tan(fx+e)^6 + 6A a^3 c^3 \tan(fx+e)^5 + 15B a^3 c^3 \tan(fx+e)^4 + 20A a^3 c^3 \tan(fx+e)^3 + 15B a^3 c^3 \tan(fx+e)^2 + 30A a^3 c^3 \tan(fx+e)}{30f}$
norman	$\frac{a^3 A c^3 \tan(fx+e)}{f} + \frac{B a^3 c^3 \tan(fx+e)^2}{2f} + \frac{B a^3 c^3 \tan(fx+e)^4}{2f} + \frac{B a^3 c^3 \tan(fx+e)^6}{6f} + \frac{2a^3 A c^3 \tan(fx+e)^3}{3f} +$
parts	$A a^3 c^3 x + \frac{A a^3 c^3 \left(\frac{\tan(fx+e)^5}{5} - \frac{\tan(fx+e)^3}{3} + \tan(fx+e) - \arctan(\tan(fx+e)) \right)}{f} + \frac{B a^3 c^3 \ln(1 + \tan(fx+e)^2)}{2f} +$

```
input int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3,x,method=_R
ETURNVERBOSE)
```

```
output 1/f*c^3*a^3*(1/6*B*tan(f*x+e)^6+1/5*A*tan(f*x+e)^5+1/2*B*tan(f*x+e)^4+2/3*
A*tan(f*x+e)^3+1/2*B*tan(f*x+e)^2+A*tan(f*x+e))
```

3.693.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.75

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ict \tan(e + fx))^3 dx =$$

$$\frac{16 (10 (-iA - B) a^3 c^3 e^{6i fx + 6ie} - 15i A a^3 c^3 e^{4i fx + 4ie} - 6i A a^3 c^3 e^{2i fx + 2ie} - i A a^3 c^3)}{15 (f e^{12i fx + 12ie} + 6 f e^{10i fx + 10ie} + 15 f e^{8i fx + 8ie} + 20 f e^{6i fx + 6ie} + 15 f e^{4i fx + 4ie} + 6 f e^{2i fx + 2ie})}$$

3.693. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ict \tan(e + fx))^3 dx$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3,x, algorithm="fricas")`

output `-16/15*(10*(-I*A - B)*a^3*c^3*e^(6*I*f*x + 6*I*e) - 15*I*A*a^3*c^3*e^(4*I*f*x + 4*I*e) - 6*I*A*a^3*c^3*e^(2*I*f*x + 2*I*e) - I*A*a^3*c^3)/(f*e^(12*I*f*x + 12*I*e) + 6*f*e^(10*I*f*x + 10*I*e) + 15*f*e^(8*I*f*x + 8*I*e) + 20*f*e^(6*I*f*x + 6*I*e) + 15*f*e^(4*I*f*x + 4*I*e) + 6*f*e^(2*I*f*x + 2*I*e) + f)`

3.693.6 Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.49 (sec) , antiderivative size = 224, normalized size of antiderivative = 2.67

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ict \tan(e + fx))^3 dx$$

$$= \frac{240iAa^3c^3e^{4ie}e^{4ifx} + 96iAa^3c^3e^{2ie}e^{2ifx} + 16iAa^3c^3 + (160iAa^3c^3e^{6ie} + 160Ba^3c^3e^{6ie})e^{6ifx}}{15fe^{12ie}e^{12ifx} + 90fe^{10ie}e^{10ifx} + 225fe^{8ie}e^{8ifx} + 300fe^{6ie}e^{6ifx} + 225fe^{4ie}e^{4ifx} + 90fe^{2ie}e^{2ifx} + 15f}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**3,x)`

output `(240*I*A*a**3*c**3*exp(4*I*e)*exp(4*I*f*x) + 96*I*A*a**3*c**3*exp(2*I*e)*exp(2*I*f*x) + 16*I*A*a**3*c**3 + (160*I*A*a**3*c**3*exp(6*I*e) + 160*B*a**3*c**3*exp(6*I*e))*exp(6*I*f*x))/(15*f*exp(12*I*e)*exp(12*I*f*x) + 90*f*exp(10*I*e)*exp(10*I*f*x) + 225*f*exp(8*I*e)*exp(8*I*f*x) + 300*f*exp(6*I*e)*exp(6*I*f*x) + 225*f*exp(4*I*e)*exp(4*I*f*x) + 90*f*exp(2*I*e)*exp(2*I*f*x) + 15*f)`

3.693.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.26

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ict \tan(e + fx))^3 dx$$

$$= \frac{5Ba^3c^3 \tan(fx + e)^6 + 6Aa^3c^3 \tan(fx + e)^5 + 15Ba^3c^3 \tan(fx + e)^4 + 20Aa^3c^3 \tan(fx + e)^3 + 15Ba^3c^3 \tan(fx + e)^2 + 6Aa^3c^3 \tan(fx + e) + 5Ba^3c^3}{30f}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3,x, algorithm="maxima")`

output $\frac{1}{30}*(5*B*a^3*c^3*\tan(f*x + e)^6 + 6*A*a^3*c^3*\tan(f*x + e)^5 + 15*B*a^3*c^3*\tan(f*x + e)^4 + 20*A*a^3*c^3*\tan(f*x + e)^3 + 15*B*a^3*c^3*\tan(f*x + e)^2 + 30*A*a^3*c^3*\tan(f*x + e))/f$

3.693.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 754 vs. $2(78) = 156$.

Time = 0.90 (sec) , antiderivative size = 754, normalized size of antiderivative = 8.98

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^3 dx$$

$$= \frac{5Ba^3c^3 \tan(fx)^6 \tan(e)^6 - 30Aa^3c^3 \tan(fx)^6 \tan(e)^5 - 30Aa^3c^3 \tan(fx)^5 \tan(e)^6 + 15Ba^3c^3 \tan(fx)^4 \tan(e)^6 - 60Aa^3c^3 \tan(fx)^4 \tan(e)^5 - 15Ba^3c^3 \tan(fx)^4 \tan(e)^4 + 20Aa^3c^3 \tan(fx)^3 \tan(e)^6 - 30Aa^3c^3 \tan(fx)^3 \tan(e)^5 - 15Ba^3c^3 \tan(fx)^3 \tan(e)^4 + 30Aa^3c^3 \tan(fx)^2 \tan(e)^6 - 60Aa^3c^3 \tan(fx)^2 \tan(e)^5 - 15Ba^3c^3 \tan(fx)^2 \tan(e)^4 + 20Aa^3c^3 \tan(fx) \tan(e)^6 - 30Aa^3c^3 \tan(fx) \tan(e)^5 - 15Ba^3c^3 \tan(fx) \tan(e)^4 + 30Aa^3c^3 \tan(e)^6 - 60Aa^3c^3 \tan(e)^5 - 15Ba^3c^3 \tan(e)^4 + 30Aa^3c^3}{(f \tan(fx) + e)^6}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3,x, algorithm="giac")`

output $\frac{1}{30}*(5*B*a^3*c^3*\tan(f*x)^6*\tan(e)^6 - 30*A*a^3*c^3*\tan(f*x)^6*\tan(e)^5 - 30*A*a^3*c^3*\tan(f*x)^5*\tan(e)^6 + 15*B*a^3*c^3*\tan(f*x)^6*\tan(e)^4 + 15*B*a^3*c^3*\tan(f*x)^4*\tan(e)^6 - 20*A*a^3*c^3*\tan(f*x)^6*\tan(e)^3 + 90*A*a^3*c^3*\tan(f*x)^5*\tan(e)^4 + 90*A*a^3*c^3*\tan(f*x)^4*\tan(e)^5 - 20*A*a^3*c^3*\tan(f*x)^3*\tan(e)^6 + 15*B*a^3*c^3*\tan(f*x)^6*\tan(e)^2 + 45*B*a^3*c^3*\tan(f*x)^4*\tan(e)^4 + 15*B*a^3*c^3*\tan(f*x)^2*\tan(e)^6 - 6*A*a^3*c^3*\tan(f*x)^6*\tan(e) + 30*A*a^3*c^3*\tan(f*x)^5*\tan(e)^2 - 180*A*a^3*c^3*\tan(f*x)^4*\tan(e)^3 - 180*A*a^3*c^3*\tan(f*x)^3*\tan(e)^4 + 30*A*a^3*c^3*\tan(f*x)^2*\tan(e)^5 - 6*A*a^3*c^3*\tan(f*x)*\tan(e)^6 + 5*B*a^3*c^3*\tan(f*x)^6 + 45*B*a^3*c^3*\tan(f*x)^4*\tan(e)^2 + 45*B*a^3*c^3*\tan(f*x)^2*\tan(e)^4 + 5*B*a^3*c^3*\tan(e)^6 + 6*A*a^3*c^3*\tan(f*x)^5 - 30*A*a^3*c^3*\tan(f*x)^4*\tan(e) + 180*A*a^3*c^3*\tan(f*x)^3*\tan(e)^2 + 180*A*a^3*c^3*\tan(f*x)^2*\tan(e)^3 - 30*A*a^3*c^3*\tan(f*x)*\tan(e)^4 + 6*A*a^3*c^3*\tan(e)^5 + 15*B*a^3*c^3*\tan(f*x)^4 + 45*B*a^3*c^3*\tan(f*x)^2*\tan(e)^2 + 15*B*a^3*c^3*\tan(e)^4 + 20*A*a^3*c^3*\tan(f*x)^3 - 90*A*a^3*c^3*\tan(f*x)^2*\tan(e) - 90*A*a^3*c^3*\tan(f*x)*\tan(e)^2 + 20*A*a^3*c^3*\tan(e)^3 + 15*B*a^3*c^3*\tan(f*x)^2 + 15*B*a^3*c^3*\tan(e)^2 + 30*A*a^3*c^3*\tan(f*x) + 30*A*a^3*c^3*\tan(e) + 5*B*a^3*c^3)/(f*\tan(f*x)^6*\tan(e)^6 - 6*f*\tan(f*x)^5*\tan(e)^5 + 15*f*\tan(f*x)^4*\tan(e)^4 - 20*f*\tan(f*x)^3*\tan(e)^3 + 15*f*\tan(f*x)^2*\tan(e)^2 - 6*f*\tan(f*x)*\tan(e) + f)$

3.693. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^3 dx$

3.693.9 Mupad [B] (verification not implemented)

Time = 8.33 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.43

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^3 dx$$

$$= \frac{a^3 c^3 \sin(e + fx) (30 A \cos(e + fx)^5 + 15 B \cos(e + fx)^4 \sin(e + fx) + 20 A \cos(e + fx)^3 \sin(e + fx) + 6 A \cos(e + fx)^2 \sin(e + fx)^2 + 15 B \cos(e + fx)^2 \sin(e + fx)^3 + 6 A \cos(e + fx) \sin(e + fx)^4 + 15 B \cos(e + fx)^4 \sin(e + fx))}{30 f \cos(e + fx)^6}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i)^3,x)`

output `(a^3*c^3*sin(e + f*x)*(30*A*cos(e + f*x)^5 + 5*B*sin(e + f*x)^5 + 20*A*cos(e + f*x)^3*sin(e + f*x)^2 + 15*B*cos(e + f*x)^2*sin(e + f*x)^3 + 6*A*cos(e + f*x)*sin(e + f*x)^4 + 15*B*cos(e + f*x)^4*sin(e + f*x)))/(30*f*cos(e + f*x)^6)`

3.694 $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^2 dx$

3.694.1 Optimal result	6484
3.694.2 Mathematica [A] (verified)	6484
3.694.3 Rubi [A] (verified)	6485
3.694.4 Maple [A] (verified)	6487
3.694.5 Fricas [A] (verification not implemented)	6487
3.694.6 Sympy [B] (verification not implemented)	6488
3.694.7 Maxima [A] (verification not implemented)	6488
3.694.8 Giac [B] (verification not implemented)	6489
3.694.9 Mupad [B] (verification not implemented)	6489

3.694.1 Optimal result

Integrand size = 41, antiderivative size = 101

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^2 dx$$

$$= -\frac{2a^3(iA - B)c^2(1 + i \tan(e + fx))^3}{3f} + \frac{a^3(iA - 3B)c^2(1 + i \tan(e + fx))^4}{4f} + \frac{a^3Bc^2(1 + i \tan(e + fx))^5}{5f}$$

```
output -2/3*a^3*(I*A-B)*c^2*(1+I*tan(f*x+e))^3/f+1/4*a^3*(I*A-3*B)*c^2*(1+I*tan(f
*x+e))^4/f+1/5*a^3*B*c^2*(1+I*tan(f*x+e))^5/f
```

3.694.2 Mathematica [A] (verified)

Time = 5.54 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.90

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^2 dx$$

$$= \frac{a^3 c^2 \sec^5(e + fx)(30(iA + B) \cos(e + fx) + 2(25A + 7iB) \cos(2(e + fx)) + (5A - iB) \cos(3(e + fx)))}{120f}$$

```
input Integrate[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f
*x])^2,x]
```

output $(a^3 c^2 \text{Sec}[e + f x]^5 (30 (I A + B) \text{Cos}[e + f x] + 2 (25 A + (7 I) B + 6 (5 A - I B) \text{Cos}[2 (e + f x)] + (5 A - I B) \text{Cos}[4 (e + f x)]) \text{Sin}[e + f x]) / (120 f)$

3.694.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.83, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + i a \tan(e + f x))^3 (c - i c \tan(e + f x))^2 (A + B \tan(e + f x)) dx$$

$$\downarrow 3042$$

$$\int (a + i a \tan(e + f x))^3 (c - i c \tan(e + f x))^2 (A + B \tan(e + f x)) dx$$

$$\downarrow 4071$$

$$\frac{a c \int a^2 c (1 - i \tan(e + f x)) (i \tan(e + f x) + 1)^2 (A + B \tan(e + f x)) d \tan(e + f x)}{f}$$

$$\downarrow 27$$

$$\frac{a^3 c^2 \int (1 - i \tan(e + f x)) (i \tan(e + f x) + 1)^2 (A + B \tan(e + f x)) d \tan(e + f x)}{f}$$

$$\downarrow 86$$

$$\frac{a^3 c^2 \int (i B (i \tan(e + f x) + 1)^4 + (-A - 3 i B) (i \tan(e + f x) + 1)^3 + 2 (A + i B) (i \tan(e + f x) + 1)^2) d \tan(e + f x)}{f}$$

$$\downarrow 2009$$

$$\frac{a^3 c^2 (\frac{1}{4} (-3 B + i A) (1 + i \tan(e + f x))^4 - \frac{2}{3} (-B + i A) (1 + i \tan(e + f x))^3 + \frac{1}{5} B (1 + i \tan(e + f x))^5)}{f}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^3*(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^2, x]$

3.694. $\int (a + i a \tan(e + f x))^3 (A + B \tan(e + f x)) (c - i c \tan(e + f x))^2 dx$

output $(a^3 c^2 ((-2(A - B)(1 + i \tan(e + fx))^3)/3 + ((A - 3B)(1 + i \tan(e + fx))^4)/4 + (B(1 + i \tan(e + fx))^5)/5))/f$

3.694.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 86 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.694.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.93

method	result
derivativedivides	$\frac{ic^2 a^3 \left(\frac{B \tan(fx+e)^5}{5} + \frac{(-iB+A) \tan(fx+e)^4}{4} + \frac{(iA-2i(iB+A)-B) \tan(fx+e)^3}{3} + \frac{(-iB+A) \tan(fx+e)^2}{2} - i \tan(fx+e)A \right)}{f}$
default	$\frac{ic^2 a^3 \left(\frac{B \tan(fx+e)^5}{5} + \frac{(-iB+A) \tan(fx+e)^4}{4} + \frac{(iA-2i(iB+A)-B) \tan(fx+e)^3}{3} + \frac{(-iB+A) \tan(fx+e)^2}{2} - i \tan(fx+e)A \right)}{f}$
risch	$\frac{4c^2 a^3 (30iA e^{6i(fx+e)} + 30B e^{6i(fx+e)} + 50iA e^{4i(fx+e)} + 10B e^{4i(fx+e)} + 25iA e^{2i(fx+e)} + 5B e^{2i(fx+e)} + 5iA + B)}{15f (e^{2i(fx+e)} + 1)^5}$
norman	$\frac{A a^3 c^2 \tan(fx+e)}{f} + \frac{(iA a^3 c^2 + B a^3 c^2) \tan(fx+e)^2}{2f} + \frac{(iA a^3 c^2 + B a^3 c^2) \tan(fx+e)^4}{4f} + \frac{(iB a^3 c^2 + A a^3 c^2) \tan(fx+e)^5}{3f}$
parallelrisch	$\frac{12iB a^3 c^2 \tan(fx+e)^5 + 15iA \tan(fx+e)^4 a^3 c^2 + 20iB \tan(fx+e)^3 a^3 c^2 + 15B \tan(fx+e)^4 a^3 c^2 + 30iA \tan(fx+e)^2 a^3 c^2 + 5iA a^3 c^2}{60f}$
parts	$\frac{(iA a^3 c^2 + B a^3 c^2) \ln(1 + \tan(fx+e)^2)}{2f} + \frac{(iA a^3 c^2 + B a^3 c^2) \left(\frac{\tan(fx+e)^4}{4} - \frac{\tan(fx+e)^2}{2} + \frac{\ln(1 + \tan(fx+e)^2)}{2} \right)}{f} + \frac{(iB a^3 c^2 + A a^3 c^2) \tan(fx+e)^5}{3f}$

input `int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2,x,method=_RETURVERBOSE)`

output `I/f*c^2*a^3*(1/5*B*tan(f*x+e)^5+1/4*(A-I*B)*tan(f*x+e)^4+1/3*(I*A-2*I*(A+I*B)-B)*tan(f*x+e)^3+1/2*(A-I*B)*tan(f*x+e)^2-I*tan(f*x+e)*A)`

3.694.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.50

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^2 dx = \frac{4(30(-iA - B)a^3 c^2 e^{(6i fx + 6i e)} + 10(-5iA - B)a^3 c^2 e^{(4i fx + 4i e)} + 5(-5iA - B)a^3 c^2 e^{(2i fx + 2i e)} + (-5iA - B)a^3 c^2)}{15(fe^{(10i fx + 10i e)} + 5fe^{(8i fx + 8i e)} + 10fe^{(6i fx + 6i e)} + 10fe^{(4i fx + 4i e)} + 5fe^{(2i fx + 2i e)} + f)}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2,x, algorithm="fracas")`

output
$$\begin{aligned} & -4/15*(30*(-I*A - B)*a^3*c^2*e^{(6*I*f*x + 6*I*e)} + 10*(-5*I*A - B)*a^3*c^2 \\ & *e^{(4*I*f*x + 4*I*e)} + 5*(-5*I*A - B)*a^3*c^2*e^{(2*I*f*x + 2*I*e)} + (-5*I* \\ & A - B)*a^3*c^2)/(f*e^{(10*I*f*x + 10*I*e)} + 5*f*e^{(8*I*f*x + 8*I*e)} + 10*f* \\ & e^{(6*I*f*x + 6*I*e)} + 10*f*e^{(4*I*f*x + 4*I*e)} + 5*f*e^{(2*I*f*x + 2*I*e)} + \\ & f) \end{aligned}$$

3.694.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 250 vs. $2(80) = 160$.

Time = 0.41 (sec) , antiderivative size = 250, normalized size of antiderivative = 2.48

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^2 dx$$

$$= \frac{20iAa^3c^2 + 4Ba^3c^2 + (100iAa^3c^2e^{2ie} + 20Ba^3c^2e^{2ie})e^{2ifx} + (200iAa^3c^2e^{4ie} + 40Ba^3c^2e^{4ie})e^{4ifx} + (120iAa^3c^2e^{6ie} + 20Ba^3c^2e^{6ie})e^{6ifx} + (120iAa^3c^2e^{8ie} + 40Ba^3c^2e^{8ie})e^{8ifx} + 15f^2e^{10ie}e^{10ifx} + 75f^2e^{8ie}e^{8ifx} + 150f^2e^{6ie}e^{6ifx} + 150f^2e^{4ie}e^{4ifx} + 75f^2e^{2ie}e^{2ifx} + 15f^2}{15fe^{10ie}e^{10ifx} + 75fe^{8ie}e^{8ifx} + 150fe^{6ie}e^{6ifx} + 150fe^{4ie}e^{4ifx} + 75fe^{2ie}e^{2ifx} + 15f}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**2,x)`

output
$$\begin{aligned} & (20*I*A*a**3*c**2 + 4*B*a**3*c**2 + (100*I*A*a**3*c**2*\exp(2*I*e) + 20*B*a \\ & **3*c**2*\exp(2*I*e))*\exp(2*I*f*x) + (200*I*A*a**3*c**2*\exp(4*I*e) + 40*B*a \\ & **3*c**2*\exp(4*I*e))*\exp(4*I*f*x) + (120*I*A*a**3*c**2*\exp(6*I*e) + 120*B* \\ & a**3*c**2*\exp(6*I*e))*\exp(6*I*f*x))/(15*f*\exp(10*I*e)*\exp(10*I*f*x) + 75*f \\ & *\exp(8*I*e)*\exp(8*I*f*x) + 150*f*\exp(6*I*e)*\exp(6*I*f*x) + 150*f*\exp(4*I*e) \\ &)*\exp(4*I*f*x) + 75*f*\exp(2*I*e)*\exp(2*I*f*x) + 15*f) \end{aligned}$$

3.694.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.04

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^2 dx$$

$$= \frac{12iBa^3c^2 \tan(fx + e)^5 - 15(-iA - B)a^3c^2 \tan(fx + e)^4 + 20(A + iB)a^3c^2 \tan(fx + e)^3 - 30(-iA - B)a^3c^2 \tan(fx + e)^2 + 15f^2 \tan(fx + e)}{60f}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2,x, algorithm="maxima")`

output $\frac{1}{60}*(12*I*B*a^3*c^2*\tan(f*x + e)^5 - 15*(-I*A - B)*a^3*c^2*\tan(f*x + e)^4 + 20*(A + I*B)*a^3*c^2*\tan(f*x + e)^3 - 30*(-I*A - B)*a^3*c^2*\tan(f*x + e)^2 + 60*A*a^3*c^2*\tan(f*x + e))/f$

3.694.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 192 vs. $2(85) = 170$.

Time = 0.64 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.90

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^2 dx = \frac{4(-30i Aa^3c^2e^{(6i fx+6ie)} - 30Ba^3c^2e^{(6i fx+6ie)} - 50i Aa^3c^2e^{(4i fx+4ie)} - 10Ba^3c^2e^{(4i fx+4ie)} - 25i Aa^3c^2e^{(2i fx+2ie)} - 5Ba^3c^2e^{(2i fx+2ie)} - 5Aa^3c^2 - Ba^3c^2)/(fe^{(10i fx+10ie)} + 5fe^{(8i fx+8ie)} + 10fe^{(6i fx+6ie)} + 10fe^{(4i fx+4ie)} + 5fe^{(2i fx+2ie)} + f)}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2,x, algorithm="giac")`

output $\frac{-4/15*(-30*I*A*a^3*c^2*e^{(6*I*f*x + 6*I*e)} - 30*B*a^3*c^2*e^{(6*I*f*x + 6*I*e)} - 50*I*A*a^3*c^2*e^{(4*I*f*x + 4*I*e)} - 10*B*a^3*c^2*e^{(4*I*f*x + 4*I*e)} - 25*I*A*a^3*c^2*e^{(2*I*f*x + 2*I*e)} - 5*B*a^3*c^2*e^{(2*I*f*x + 2*I*e)} - 5*I*A*a^3*c^2 - B*a^3*c^2)/(f*e^{(10*I*f*x + 10*I*e)} + 5*f*e^{(8*I*f*x + 8*I*e)} + 10*f*e^{(6*I*f*x + 6*I*e)} + 10*f*e^{(4*I*f*x + 4*I*e)} + 5*f*e^{(2*I*f*x + 2*I*e)} + f)}$

3.694.9 Mupad [B] (verification not implemented)

Time = 8.31 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.07

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^2 dx = \frac{Aa^3c^2 \tan(e + fx) - \frac{a^3c^2 \tan(e+fx)^3 (-B+A \operatorname{li}) \operatorname{li}}{3} + \frac{a^3c^2 \tan(e+fx)^2 (A-B \operatorname{li}) \operatorname{li}}{2} + \frac{a^3c^2 \tan(e+fx)^4 (A-B \operatorname{li}) \operatorname{li}}{4} + \frac{Ba^3c^2}{4}}{f}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i)^2,x)`

output $(Aa^3c^2\tan(e + fx) - (a^3c^2\tan(e + fx)^3(A + B)\tan(e + fx))/3 + (a^3c^2\tan(e + fx)^2(A - B)\tan(e + fx))/2 + (a^3c^2\tan(e + fx)^4(A - B)\tan(e + fx))/4 + (Ba^3c^2\tan(e + fx)^5)/5)/f$

3.695 $\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx)) dx$

3.695.1 Optimal result	6491
3.695.2 Mathematica [A] (verified)	6491
3.695.3 Rubi [A] (verified)	6492
3.695.4 Maple [A] (verified)	6493
3.695.5 Fricas [B] (verification not implemented)	6494
3.695.6 Sympy [B] (verification not implemented)	6494
3.695.7 Maxima [A] (verification not implemented)	6495
3.695.8 Giac [B] (verification not implemented)	6495
3.695.9 Mupad [B] (verification not implemented)	6496

3.695.1 Optimal result

Integrand size = 39, antiderivative size = 61

$$\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))(c - ictan(e + fx)) dx$$

$$= -\frac{a^3(iA - B)c(1 + i \tan(e + fx))^3}{3f} - \frac{a^3Bc(1 + i \tan(e + fx))^4}{4f}$$

output `-1/3*a^3*(I*A-B)*c*(1+I*tan(f*x+e))^3/f-1/4*a^3*B*c*(1+I*tan(f*x+e))^4/f`

3.695.2 Mathematica [A] (verified)

Time = 3.28 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.15

$$\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))(c - ictan(e + fx)) dx =$$

$$-\frac{a^3c(3B - 12A \tan(e + fx) + (-12iA - 6B) \tan^2(e + fx) + 4(A - 2iB) \tan^3(e + fx) + 3B \tan^4(e + fx))}{12f}$$

input `Integrate[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x]),x]`

output `-1/12*(a^3*c*(3*B - 12*A*Tan[e + f*x] + ((-12*I)*A - 6*B)*Tan[e + f*x]^2 + 4*(A - (2*I)*B)*Tan[e + f*x]^3 + 3*B*Tan[e + f*x]^4))/f`

3.695. $\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))(c - ictan(e + fx)) dx$

3.695.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.90, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.128$, Rules used = {3042, 4071, 27, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(e + fx))^3 (c - ictan(e + fx))(A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(e + fx))^3 (c - ictan(e + fx))(A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{a^3 c \int a^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^3 c \int (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{49} \\
 & \frac{a^3 c \int ((A + iB)(i \tan(e + fx) + 1)^2 - iB(i \tan(e + fx) + 1)^3) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^3 c (-\frac{1}{3}(-B + iA)(1 + i \tan(e + fx))^3 - \frac{1}{4}B(1 + i \tan(e + fx))^4)}{f}
 \end{aligned}$$

input `Int[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x]),x]`

output `(a^3*c*(-1/3*((I*A - B)*(1 + I*Tan[e + f*x])^3) - (B*(1 + I*Tan[e + f*x])^4)/4))/f`

3.695.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.695.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.03

method	result
derivativedivides	$\frac{a^3 c \left(-\frac{B \tan(fx+e)^4}{4} - \frac{(-2iB+A) \tan(fx+e)^3}{3} - \frac{(-2iA-B) \tan(fx+e)^2}{2} + A \tan(fx+e) \right)}{f}$
default	$\frac{a^3 c \left(-\frac{B \tan(fx+e)^4}{4} - \frac{(-2iB+A) \tan(fx+e)^3}{3} - \frac{(-2iA-B) \tan(fx+e)^2}{2} + A \tan(fx+e) \right)}{f}$
norman	$\frac{a^3 c A \tan(fx+e)}{f} - \frac{(-2iB a^3 c + a^3 c A) \tan(fx+e)^3}{3f} + \frac{(2iA a^3 c + B a^3 c) \tan(fx+e)^2}{2f} - \frac{B a^3 c \tan(fx+e)^4}{4f}$
parallelrisch	$\frac{8iB \tan(fx+e)^3 a^3 c - 3B \tan(fx+e)^4 a^3 c + 12iA \tan(fx+e)^2 a^3 c - 4A \tan(fx+e)^3 a^3 c + 6B \tan(fx+e)^2 a^3 c + 12A \tan(fx+e) a^3 c}{12f}$
risch	$\frac{4a^3 c (6iA e^{6i(fx+e)} + 6B e^{6i(fx+e)} + 12iA e^{4i(fx+e)} + 6B e^{4i(fx+e)} + 8iA e^{2i(fx+e)} + 4B e^{2i(fx+e)} + 2iA + B)}{3f (e^{2i(fx+e)} + 1)^4}$
parts	$\frac{(2iA a^3 c + B a^3 c) \ln(1 + \tan(fx+e)^2)}{2f} + \frac{(2iB a^3 c - a^3 c A) \left(\frac{\tan(fx+e)^3}{3} - \tan(fx+e) + \arctan(\tan(fx+e)) \right)}{f} + a^3 c A$

3.695. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ict \tan(e + fx)) dx$

```
input int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x,method=_RET
URNVERBOSE)
```

```
output 1/f*a^3*c*(-1/4*B*tan(f*x+e)^4-1/3*(A-2*I*B)*tan(f*x+e)^3-1/2*(-2*I*A-B)*t
an(f*x+e)^2+A*tan(f*x+e))
```

3.695.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 132 vs. $2(51) = 102$.

Time = 0.23 (sec) , antiderivative size = 132, normalized size of antiderivative = 2.16

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ic \tan(e + fx)) dx =$$

$$-\frac{4(6(-iA - B)a^3ce^{(6ifx+6ie)} + 6(-2iA - B)a^3ce^{(4ifx+4ie)} + 4(-2iA - B)a^3ce^{(2ifx+2ie)} + (-2iA - B)a^3c)}{3(fe^{(8ifx+8ie)} + 4fe^{(6ifx+6ie)} + 6fe^{(4ifx+4ie)} + 4fe^{(2ifx+2ie)} + f)}$$

```
input integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x, algo
rithm="fricas")
```

```
output -4/3*(6*(-I*A - B)*a^3*c*e^(6*I*f*x + 6*I*e) + 6*(-2*I*A - B)*a^3*c*e^(4*I
*f*x + 4*I*e) + 4*(-2*I*A - B)*a^3*c*e^(2*I*f*x + 2*I*e) + (-2*I*A - B)*a^
3*c)/(f*e^(8*I*f*x + 8*I*e) + 4*f*e^(6*I*f*x + 6*I*e) + 6*f*e^(4*I*f*x + 4
*I*e) + 4*f*e^(2*I*f*x + 2*I*e) + f)
```

3.695.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 218 vs. $2(48) = 96$.

Time = 0.28 (sec) , antiderivative size = 218, normalized size of antiderivative = 3.57

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ic \tan(e + fx)) dx$$

$$= \frac{8iAa^3c + 4Ba^3c + (32iAa^3ce^{2ie} + 16Ba^3ce^{2ie})e^{2ifx} + (48iAa^3ce^{4ie} + 24Ba^3ce^{4ie})e^{4ifx} + (24iAa^3ce^{6ie} + 12Ba^3ce^{6ie})e^{6ifx} + (8iAa^3ce^{8ie} + 4Ba^3ce^{8ie})e^{8ifx}}{3fe^{8ie}e^{8ifx} + 12fe^{6ie}e^{6ifx} + 18fe^{4ie}e^{4ifx} + 12fe^{2ie}e^{2ifx} + 3f}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x)`

output $(8*I*A*a**3*c + 4*B*a**3*c + (32*I*A*a**3*c*\exp(2*I*e) + 16*B*a**3*c*\exp(2*I*e))*\exp(2*I*f*x) + (48*I*A*a**3*c*\exp(4*I*e) + 24*B*a**3*c*\exp(4*I*e))*\exp(4*I*f*x) + (24*I*A*a**3*c*\exp(6*I*e) + 24*B*a**3*c*\exp(6*I*e))*\exp(6*I*f*x))/(3*f*\exp(8*I*e)*\exp(8*I*f*x) + 12*f*\exp(6*I*e)*\exp(6*I*f*x) + 18*f*\exp(4*I*e)*\exp(4*I*f*x) + 12*f*\exp(2*I*e)*\exp(2*I*f*x) + 3*f)$

3.695.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.18

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx)) dx = \frac{3Ba^3c \tan(fx + e)^4 + 4(A - 2iB)a^3c \tan(fx + e)^3 - 6(2iA + B)a^3c \tan(fx + e)^2 - 12Aa^3c \tan(fx + e) - 3a^3c}{12f}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x, algorithm="maxima")`

output $-1/12*(3*B*a^3*c*\tan(f*x + e)^4 + 4*(A - 2*I*B)*a^3*c*\tan(f*x + e)^3 - 6*(2*I*A + B)*a^3*c*\tan(f*x + e)^2 - 12*A*a^3*c*\tan(f*x + e))/f$

3.695.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 164 vs. $2(51) = 102$.

Time = 0.54 (sec) , antiderivative size = 164, normalized size of antiderivative = 2.69

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx)) dx = \frac{4(-6iAa^3ce^{(6ifx+6ie)} - 6Ba^3ce^{(6ifx+6ie)} - 12iAa^3ce^{(4ifx+4ie)} - 6Ba^3ce^{(4ifx+4ie)} - 8iAa^3ce^{(2ifx+2ie)})}{3(fe^{(8ifx+8ie)} + 4fe^{(6ifx+6ie)} + 6fe^{(4ifx+4ie)} + 4fe^{(2ifx+2ie)}) + 3}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e)),x, algorithm="giac")`

3.695. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx)) dx$

output
$$\frac{-4/3*(-6*I*A*a^3*c*e^{(6*I*f*x + 6*I*e)} - 6*B*a^3*c*e^{(6*I*f*x + 6*I*e)} - 12*I*A*a^3*c*e^{(4*I*f*x + 4*I*e)} - 6*B*a^3*c*e^{(4*I*f*x + 4*I*e)} - 8*I*A*a^3*c*e^{(2*I*f*x + 2*I*e)} - 4*B*a^3*c*e^{(2*I*f*x + 2*I*e)} - 2*I*A*a^3*c - B*a^3*c)/(f*e^{(8*I*f*x + 8*I*e)} + 4*f*e^{(6*I*f*x + 6*I*e)} + 6*f*e^{(4*I*f*x + 4*I*e)} + 4*f*e^{(2*I*f*x + 2*I*e)} + f)$$

3.695.9 Mupad [B] (verification not implemented)

Time = 8.34 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.18

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ic \tan(e + fx)) dx$$

$$= \frac{-\frac{Bca^3 \tan(e+fx)^4}{4} - \frac{c(A-B2i)a^3 \tan(e+fx)^3}{3} + \frac{c(B+A2i)a^3 \tan(e+fx)^2}{2} + Aca^3 \tan(e+fx)}{f}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i),x)`

output
$$(A*a^3*c*\tan(e + f*x) + (a^3*c*\tan(e + f*x)^2*(A*2i + B))/2 - (a^3*c*\tan(e + f*x)^3*(A - B*2i))/3 - (B*a^3*c*\tan(e + f*x)^4)/4)/f$$

3.696 $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) dx$

3.696.1 Optimal result	6497
3.696.2 Mathematica [A] (verified)	6497
3.696.3 Rubi [A] (verified)	6498
3.696.4 Maple [A] (verified)	6500
3.696.5 Fricas [A] (verification not implemented)	6500
3.696.6 Sympy [A] (verification not implemented)	6501
3.696.7 Maxima [A] (verification not implemented)	6501
3.696.8 Giac [B] (verification not implemented)	6502
3.696.9 Mupad [B] (verification not implemented)	6502

3.696.1 Optimal result

Integrand size = 26, antiderivative size = 110

$$\begin{aligned} & \int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) dx \\ &= 4a^3(A - iB)x - \frac{4a^3(iA + B) \log(\cos(e + fx))}{f} - \frac{2a^3(A - iB) \tan(e + fx)}{f} \\ & \quad + \frac{a(iA + B)(a + ia \tan(e + fx))^2}{2f} + \frac{B(a + ia \tan(e + fx))^3}{3f} \end{aligned}$$

output `4*a^3*(A-I*B)*x-4*a^3*(I*A+B)*ln(cos(f*x+e))/f-2*a^3*(A-I*B)*tan(f*x+e)/f+1/2*a*(I*A+B)*(a+I*a*tan(f*x+e))^2/f+1/3*B*(a+I*a*tan(f*x+e))^3/f`

3.696.2 Mathematica [A] (verified)

Time = 0.92 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.66

$$\begin{aligned} & \int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) dx \\ &= \frac{B(a + ia \tan(e + fx))^3 + \frac{3}{2}a^3(iA + B) (8 \log(i + \tan(e + fx)) + 6i \tan(e + fx) - \tan^2(e + fx))}{3f} \end{aligned}$$

input `Integrate[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]),x]`

output `(B*(a + I*a*Tan[e + f*x])^3 + (3*a^3*(I*A + B)*(8*Log[I + Tan[e + f*x]] + (6*I)*Tan[e + f*x] - Tan[e + f*x]^2))/2)/(3*f)`

3.696. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) dx$

3.696.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.90, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3042, 4010, 3042, 3959, 3042, 3958, 3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) dx \\
 & \quad \downarrow \text{4010} \\
 & (A - iB) \int (i \tan(e + fx)a + a)^3 dx + \frac{B(a + ia \tan(e + fx))^3}{3f} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \int (i \tan(e + fx)a + a)^3 dx + \frac{B(a + ia \tan(e + fx))^3}{3f} \\
 & \quad \downarrow \text{3959} \\
 & (A - iB) \left(2a \int (i \tan(e + fx)a + a)^2 dx + \frac{ia(a + ia \tan(e + fx))^2}{2f} \right) + \frac{B(a + ia \tan(e + fx))^3}{3f} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \left(2a \int (i \tan(e + fx)a + a)^2 dx + \frac{ia(a + ia \tan(e + fx))^2}{2f} \right) + \frac{B(a + ia \tan(e + fx))^3}{3f} \\
 & \quad \downarrow \text{3958} \\
 & (A - iB) \left(2a \left(2ia^2 \int \tan(e + fx) dx - \frac{a^2 \tan(e + fx)}{f} + 2a^2 x \right) + \frac{ia(a + ia \tan(e + fx))^2}{2f} \right) + \\
 & \quad \frac{B(a + ia \tan(e + fx))^3}{3f} \\
 & \quad \downarrow \text{3042} \\
 & (A - iB) \left(2a \left(2ia^2 \int \tan(e + fx) dx - \frac{a^2 \tan(e + fx)}{f} + 2a^2 x \right) + \frac{ia(a + ia \tan(e + fx))^2}{2f} \right) + \\
 & \quad \frac{B(a + ia \tan(e + fx))^3}{3f} \\
 & \quad \downarrow \text{3956}
 \end{aligned}$$

3.696. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) dx$

$$(A - iB) \left(2a \left(-\frac{a^2 \tan(e + fx)}{f} - \frac{2ia^2 \log(\cos(e + fx))}{f} + 2a^2 x \right) + \frac{ia(a + ia \tan(e + fx))^2}{2f} \right) + \frac{B(a + ia \tan(e + fx))^3}{3f}$$

input `Int[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]),x]`

output `(B*(a + I*a*Tan[e + f*x])^3)/(3*f) + (A - I*B)*(((I/2)*a*(a + I*a*Tan[e + f*x])^2)/f + 2*a*(2*a^2*x - ((2*I)*a^2*Log[Cos[e + f*x]])/f - (a^2*Tan[e + f*x])/f))`

3.696.3.1 Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

rule 3958 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_.)])^2, x_Symbol] := Simp[(a^2 - b^2)*x, x] + (Simp[b^2*(Tan[c + d*x]/d), x] + Simp[2*a*b Int[Tan[c + d*x], x], x]) /; FreeQ[{a, b, c, d}, x]`

rule 3959 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Simp[b*((a + b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] + Simp[2*a Int[(a + b*Tan[c + d*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && GtQ[n, 1]`

rule 4010 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] := Simp[d*((a + b*Tan[e + f*x])^m/(f*m)), x] + Simp[(b*c + a*d)/b Int[(a + b*Tan[e + f*x])^m, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && !LtQ[m, 0]`

3.696.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.91

method	result
derivativedivides	$\frac{a^3 \left(-\frac{iB \tan(fx+e)^3}{3} - \frac{iA \tan(fx+e)^2}{2} + 4i \tan(fx+e)B - \frac{3B \tan(fx+e)^2}{2} - 3A \tan(fx+e) + \frac{(4iA+4B) \ln(1+\tan(fx+e)^2)}{2} \right) + (-4iB a^3 + 4a^3 A) x - \frac{(iA a^3 + 3B a^3) \tan(fx+e)^2}{2f} - \frac{(-4iB a^3 + 3a^3 A) \tan(fx+e)}{f} - \frac{iB a^3 \tan(fx+e)^3}{3f} + \dots}{f}$
default	$\frac{a^3 \left(-\frac{iB \tan(fx+e)^3}{3} - \frac{iA \tan(fx+e)^2}{2} + 4i \tan(fx+e)B - \frac{3B \tan(fx+e)^2}{2} - 3A \tan(fx+e) + \frac{(4iA+4B) \ln(1+\tan(fx+e)^2)}{2} \right) + (-4iB a^3 + 4a^3 A) x - \frac{(iA a^3 + 3B a^3) \tan(fx+e)^2}{2f} - \frac{(-4iB a^3 + 3a^3 A) \tan(fx+e)}{f} - \frac{iB a^3 \tan(fx+e)^3}{3f} + \dots}{f}$
norman	$(-4iB a^3 + 4a^3 A) x - \frac{(iA a^3 + 3B a^3) \tan(fx+e)^2}{2f} - \frac{(-4iB a^3 + 3a^3 A) \tan(fx+e)}{f} - \frac{iB a^3 \tan(fx+e)^3}{3f} + \dots$
parallelrisch	$\frac{-2iB a^3 \tan(fx+e)^3 - 3iA \tan(fx+e)^2 a^3 - 24iB x a^3 f + 12iA \ln(1+\tan(fx+e)^2) a^3 + 24A x a^3 f + 24iB \tan(fx+e) a^3 - 9B a^3}{6f}$
risch	$\frac{8ia^3 B e}{f} - \frac{8a^3 A e}{f} - \frac{2a^3 (12iA e^{4i(fx+e)} + 24B e^{4i(fx+e)} + 21iA e^{2i(fx+e)} + 33B e^{2i(fx+e)} + 9iA + 13B)}{3f(e^{2i(fx+e)} + 1)^3} - \frac{4a^3 \ln(e^{2i(fx+e)} + 1)}{3f}$
parts	$A a^3 x + \frac{(-iA a^3 - 3B a^3) \left(\frac{\tan(fx+e)^2}{2} - \frac{\ln(1+\tan(fx+e)^2)}{2} \right)}{f} + \frac{(3iA a^3 + B a^3) \ln(1+\tan(fx+e)^2)}{2f} + \frac{(3iB a^3 - 3A a^3)}{2f}$

input `int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output `1/f*a^3*(-1/3*I*B*tan(f*x+e)^3-1/2*I*A*tan(f*x+e)^2+4*I*B*tan(f*x+e)-3/2*B*tan(f*x+e)^2-3*A*tan(f*x+e)+1/2*(4*I*A+4*B)*ln(1+tan(f*x+e)^2)+(-4*I*B+4*A)*arctan(tan(f*x+e)))`

3.696.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.59

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) dx = \frac{2(12(iA + 2B)a^3 e^{4i fx + 4ie} + 3(7iA + 11B)a^3 e^{2i fx + 2ie} + (9iA + 13B)a^3 + 6((iA + B)a^3 e^{6i fx + 6ie} + 3iA a^3 e^{4i fx + 4ie}))}{3(fe^{6i fx + 6ie} + 3fe^{4i fx + 4ie})}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e)),x, algorithm="fracas")`

output
$$-2/3*(12*(I*A + 2*B)*a^3*e^(4*I*f*x + 4*I*e) + 3*(7*I*A + 11*B)*a^3*e^(2*I*f*x + 2*I*e) + (9*I*A + 13*B)*a^3 + 6*((I*A + B)*a^3*e^(6*I*f*x + 6*I*e) + 3*(I*A + B)*a^3*e^(4*I*f*x + 4*I*e) + 3*(I*A + B)*a^3*e^(2*I*f*x + 2*I*e) + (I*A + B)*a^3)*log(e^(2*I*f*x + 2*I*e) + 1)/(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)$$

3.696.6 Sympy [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.67

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) dx = -\frac{4ia^3(A - iB) \log(e^{2ifx} + e^{-2ie})}{f} + \frac{-18iAa^3 - 26Ba^3 + (-42iAa^3e^{2ie} - 66Ba^3e^{2ie})e^{2ifx} + (-24iAa^3e^{4ie} - 48Ba^3e^{4ie})e^{4ifx}}{3fe^{6ie}e^{6ifx} + 9fe^{4ie}e^{4ifx} + 9fe^{2ie}e^{2ifx} + 3f}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e)),x)`

output
$$-4*I*a**3*(A - I*B)*log(exp(2*I*f*x) + exp(-2*I*e))/f + (-18*I*A*a**3 - 26*B*a**3 + (-42*I*A*a**3*exp(2*I*e) - 66*B*a**3*exp(2*I*e))*exp(2*I*f*x) + (-24*I*A*a**3*exp(4*I*e) - 48*B*a**3*exp(4*I*e))*exp(4*I*f*x))/(3*f*exp(6*I*e)*exp(6*I*f*x) + 9*f*exp(4*I*e)*exp(4*I*f*x) + 9*f*exp(2*I*e)*exp(2*I*f*x) + 3*f)$$

3.696.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.87

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) dx = \frac{2iBa^3 \tan(fx + e)^3 + 3(iA + 3B)a^3 \tan(fx + e)^2 - 24(fx + e)(A - iB)a^3 + 12(-iA - B)a^3 \log(\tan(fx + e)^2 + 1)}{6f}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e)),x, algorithm="maxima")`

output
$$-1/6*(2*I*B*a^3*tan(f*x + e)^3 + 3*(I*A + 3*B)*a^3*tan(f*x + e)^2 - 24*(fx + e)*(A - I*B)*a^3 + 12*(-I*A - B)*a^3*log(tan(f*x + e)^2 + 1) + 6*(3*A - 4*I*B)*a^3*tan(f*x + e))/f$$

3.696.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 312 vs. $2(94) = 188$.

Time = 0.49 (sec) , antiderivative size = 312, normalized size of antiderivative = 2.84

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) dx =$$

$$\frac{2(6i Aa^3 e^{(6i fx + 6i e)} \log(e^{(2i fx + 2i e)} + 1) + 6Ba^3 e^{(6i fx + 6i e)} \log(e^{(2i fx + 2i e)} + 1) + 18i Aa^3 e^{(4i fx + 4i e)} \log$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e)),x, algorithm="giac")`

output

$$\begin{aligned} & -2/3*(6*I*A*a^3*e^{(6*I*f*x + 6*I*e)}*\log(e^{(2*I*f*x + 2*I*e)} + 1) + 6*B*a^3 \\ & *e^{(6*I*f*x + 6*I*e)}*\log(e^{(2*I*f*x + 2*I*e)} + 1) + 18*I*A*a^3*e^{(4*I*f*x \\ & + 4*I*e)}*\log(e^{(2*I*f*x + 2*I*e)} + 1) + 18*B*a^3*e^{(4*I*f*x + 4*I*e)}*\log(e \\ & ^{(2*I*f*x + 2*I*e)} + 1) + 18*I*A*a^3*e^{(2*I*f*x + 2*I*e)}*\log(e^{(2*I*f*x + \\ & 2*I*e)} + 1) + 18*B*a^3*e^{(2*I*f*x + 2*I*e)}*\log(e^{(2*I*f*x + 2*I*e)} + 1) + \\ & 12*I*A*a^3*e^{(4*I*f*x + 4*I*e)} + 24*B*a^3*e^{(4*I*f*x + 4*I*e)} + 21*I*A*a^3 \\ & *e^{(2*I*f*x + 2*I*e)} + 33*B*a^3*e^{(2*I*f*x + 2*I*e)} + 6*I*A*a^3*\log(e^{(2*I \\ & *f*x + 2*I*e)} + 1) + 6*B*a^3*\log(e^{(2*I*f*x + 2*I*e)} + 1) + 9*I*A*a^3 + 13 \\ & *B*a^3)/(f*e^{(6*I*f*x + 6*I*e)} + 3*f*e^{(4*I*f*x + 4*I*e)} + 3*f*e^{(2*I*f*x \\ & + 2*I*e)} + f) \end{aligned}$$
3.696.9 Mupad [B] (verification not implemented)

Time = 8.48 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.14

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) dx$$

$$= -\frac{\tan(e + fx)^2 \left(\frac{Ba^3}{2} + \frac{a^3(2B + A1i)}{2} \right)}{f} + \frac{\ln(\tan(e + fx) + 1i) (4Ba^3 + Aa^3 4i)}{f}$$

$$+ \frac{\tan(e + fx) (Ba^3 1i - a^3(2A - B1i) + a^3(2B + A1i) 1i)}{f} - \frac{Ba^3 \tan(e + fx)^3 1i}{3f}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3,x)`

output

$$\begin{aligned} & (\log(\tan(e + f*x) + 1i)*(A*a^3*4i + 4*B*a^3))/f - (\tan(e + f*x)^2*((B*a^3) \\ & /2 + (a^3*(A*1i + 2*B))/2))/f + (\tan(e + f*x)*(B*a^3*1i - a^3*(2*A - B*1i) \\ & + a^3*(A*1i + 2*B)*1i))/f - (B*a^3*\tan(e + f*x)^3*1i)/(3*f) \end{aligned}$$

3.697
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{c-ic \tan(e+fx)} dx$$

3.697.1 Optimal result 6503
 3.697.2 Mathematica [A] (verified) 6503
 3.697.3 Rubi [A] (verified) 6504
 3.697.4 Maple [A] (verified) 6506
 3.697.5 Fricas [A] (verification not implemented) 6506
 3.697.6 Sympy [A] (verification not implemented) 6507
 3.697.7 Maxima [F(-2)] 6507
 3.697.8 Giac [B] (verification not implemented) 6508
 3.697.9 Mupad [B] (verification not implemented) 6508

3.697.1 Optimal result

Integrand size = 41, antiderivative size = 119

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx \\ &= -\frac{4a^3(A - 2iB)x}{c} + \frac{4a^3(iA + 2B) \log(\cos(e + fx))}{cf} \\ & \quad + \frac{a^3(A - 4iB) \tan(e + fx)}{cf} + \frac{a^3B \tan^2(e + fx)}{2cf} + \frac{4a^3(A - iB)}{cf(i + \tan(e + fx))} \end{aligned}$$

output `-4*a^3*(A-2*I*B)*x/c+4*a^3*(I*A+2*B)*ln(cos(f*x+e))/c/f+a^3*(A-4*I*B)*tan(f*x+e)/c/f+1/2*a^3*B*tan(f*x+e)^2/c/f+4*a^3*(A-I*B)/c/f/(I+tan(f*x+e))`

3.697.2 Mathematica [A] (verified)

Time = 5.68 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.01

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx \\ &= \frac{a^3(14A - 27iB + 8(A - 2iB) \log(i + \tan(e + fx)) + (-4iA - 11B - 8i(A - 2iB) \log(i + \tan(e + fx))))}{2cf(i + \tan(e + fx))} \end{aligned}$$

input `Integrate[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x]),x]`

output $(a^3(14A - (27I)B + 8(A - (2I)B)\text{Log}[I + \text{Tan}[e + f*x]] + ((-4I)*A - 11B - (8I)*(A - (2I)B)\text{Log}[I + \text{Tan}[e + f*x]])*\text{Tan}[e + f*x] + (2A - (7I)B)*\text{Tan}[e + f*x]^2 + B*\text{Tan}[e + f*x]^3)/(2*c*f*(I + \text{Tan}[e + f*x]))$

3.697.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.68, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{a^2 (i \tan(e+fx)+1)^2 (A+B \tan(e+fx))}{c^2 (1-i \tan(e+fx))^2} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^3 \int \frac{(i \tan(e+fx)+1)^2 (A+B \tan(e+fx))}{(1-i \tan(e+fx))^2} d \tan(e + fx)}{cf} \\
 & \quad \downarrow \text{86} \\
 & \frac{a^3 \int \left(-\frac{4(A-iB)}{(\tan(e+fx)+i)^2} + A \left(1 - \frac{4iB}{A} \right) + B \tan(e + fx) - \frac{4i(A-2iB)}{\tan(e+fx)+i} \right) d \tan(e + fx)}{cf} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^3 \left((A - 4iB) \tan(e + fx) + \frac{4(A-iB)}{\tan(e+fx)+i} - 4(2B + iA) \log(\tan(e + fx) + i) + \frac{1}{2} B \tan^2(e + fx) \right)}{cf}
 \end{aligned}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^3*(A + B*\text{Tan}[e + f*x])]/(c - I*c*\text{Tan}[e + f*x]),x]$

output $(a^3(-4(IA + 2B)\text{Log}[I + \text{Tan}[e + fx]] + (A - (4I)B)\text{Tan}[e + fx] + (B\text{Tan}[e + fx]^2)/2 + (4(A - IB))/(I + \text{Tan}[e + fx]))/(c*f)$

3.697.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.697.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.50

method	result
risch	$-\frac{2e^{2i(fx+e)}a^3B}{cf} - \frac{2ie^{2i(fx+e)}a^3A}{cf} - \frac{16ia^3Be}{fc} + \frac{8a^3Ae}{fc} + \frac{2a^3(iAe^{2i(fx+e)}+5Be^{2i(fx+e)}+iA+4B)}{cf(e^{2i(fx+e)}+1)^2} + \frac{8a^3B}{cf}$
derivativedivides	$\frac{a^3A \tan(fx+e)}{fc} - \frac{4ia^3 \tan(fx+e)B}{fc} + \frac{a^3B \tan(fx+e)^2}{2cf} - \frac{4a^3A \arctan(\tan(fx+e))}{fc} - \frac{2ia^3A \ln(1+\tan(fx+e)^2)}{fc}$
default	$\frac{a^3A \tan(fx+e)}{fc} - \frac{4ia^3 \tan(fx+e)B}{fc} + \frac{a^3B \tan(fx+e)^2}{2cf} - \frac{4a^3A \arctan(\tan(fx+e))}{fc} - \frac{2ia^3A \ln(1+\tan(fx+e)^2)}{fc}$
norman	$\frac{(-4iBa^3+a^3A) \tan(fx+e)^3}{cf} + \frac{(-8iBa^3+5a^3A) \tan(fx+e)}{cf} - \frac{4(-2iBa^3+a^3A)x}{c} - \frac{8iAa^3+9Ba^3}{2cf} - \frac{4(-2iBa^3+a^3A)x \tan(fx+e)}{c} \frac{1}{1+\tan(fx+e)^2}$

input `int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output `-2/c/f*exp(2*I*(f*x+e))*a^3*B-2*I/c/f*exp(2*I*(f*x+e))*a^3*A-16*I*a^3/f/c*B*e+8*a^3/f/c*A*e+2*a^3*(I*A*exp(2*I*(f*x+e))+5*B*exp(2*I*(f*x+e))+I*A+4*B)/c/f/(exp(2*I*(f*x+e))+1)^2+8*a^3/f/c*ln(exp(2*I*(f*x+e))+1)*B+4*I*a^3/f/c*ln(exp(2*I*(f*x+e))+1)*A`

3.697.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.38

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx = \frac{2((iA + B)a^3e^{(6i fx + 6ie)} + 2(iA + B)a^3e^{(4i fx + 4ie)} - 4Ba^3e^{(2i fx + 2ie)} + (-iA - 4B)a^3 + 2((-iA - 4B)a^3 \log(e^{(2i fx + 2ie)} + 1))}{cfe^{(4i fx + 4ie)} + 2cfe^{(2i fx + 2ie)} + c}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x, algorithm="fricas")`

output `-2*((I*A + B)*a^3*e^(6*I*f*x + 6*I*e) + 2*(I*A + B)*a^3*e^(4*I*f*x + 4*I*e) - 4*B*a^3*e^(2*I*f*x + 2*I*e) + (-I*A - 4*B)*a^3 + 2*((-I*A - 2*B)*a^3*e^(4*I*f*x + 4*I*e) + 2*(-I*A - 2*B)*a^3*e^(2*I*f*x + 2*I*e) + (-I*A - 2*B)*a^3)*log(e^(2*I*f*x + 2*I*e) + 1)/(c*f*e^(4*I*f*x + 4*I*e) + 2*c*f*e^(2*I*f*x + 2*I*e) + c*f)`

3.697.
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{c-ic \tan(e+fx)} dx$$

3.697.6 Sympy [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.73

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx$$

$$= \frac{4ia^3(A - 2iB) \log(e^{2ifx} + e^{-2ie})}{cf} + \frac{2iAa^3 + 8Ba^3 + (2iAa^3 e^{2ie} + 10Ba^3 e^{2ie}) e^{2ifx}}{cfe^{4ie} e^{4ifx} + 2cfe^{2ie} e^{2ifx} + cf}$$

$$+ \begin{cases} \frac{(-2iAa^3 e^{2ie} - 2Ba^3 e^{2ie}) e^{2ifx}}{cf} & \text{for } cf \neq 0 \\ \frac{x(4Aa^3 e^{2ie} - 4iBa^3 e^{2ie})}{c} & \text{otherwise} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x)`output `4*I*a**3*(A - 2*I*B)*log(exp(2*I*f*x) + exp(-2*I*e))/(c*f) + (2*I*A*a**3 + 8*B*a**3 + (2*I*A*a**3*exp(2*I*e) + 10*B*a**3*exp(2*I*e))*exp(2*I*f*x))/(c*f*exp(4*I*e)*exp(4*I*f*x) + 2*c*f*exp(2*I*e)*exp(2*I*f*x) + c*f) + Piecewise(((-2*I*A*a**3*exp(2*I*e) - 2*B*a**3*exp(2*I*e))*exp(2*I*f*x)/(c*f), Ne(c*f, 0)), (x*(4*A*a**3*exp(2*I*e) - 4*I*B*a**3*exp(2*I*e))/c, True))`**3.697.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x, algorithm="maxima")`output `Exception raised: RuntimeError >> ECL says: exp: undefined: 0 to a negative exponent.`

3.697.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 306 vs. $2(107) = 214$.

Time = 0.58 (sec) , antiderivative size = 306, normalized size of antiderivative = 2.57

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx =$$

$$2 \left(\frac{2(-iAa^3 - 2Ba^3) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)}{c} + \frac{4(iAa^3 + 2Ba^3) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + i)}{c} + \frac{2(-iAa^3 - 2Ba^3) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)}{c} \right)$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e)),x, algo rithm="giac")`

output `-2*(2*(-I*A*a^3 - 2*B*a^3)*log(tan(1/2*f*x + 1/2*e) + 1)/c + 4*(I*A*a^3 + 2*B*a^3)*log(tan(1/2*f*x + 1/2*e) + I)/c + 2*(-I*A*a^3 - 2*B*a^3)*log(tan(1/2*f*x + 1/2*e) - 1)/c + (5*A*a^3*tan(1/2*f*x + 1/2*e)^5 - 8*I*B*a^3*tan(1/2*f*x + 1/2*e)^5 + 2*I*A*a^3*tan(1/2*f*x + 1/2*e)^4 + 7*B*a^3*tan(1/2*f*x + 1/2*e)^4 - 10*A*a^3*tan(1/2*f*x + 1/2*e)^3 + 14*I*B*a^3*tan(1/2*f*x + 1/2*e)^3 - 2*I*A*a^3*tan(1/2*f*x + 1/2*e)^2 - 7*B*a^3*tan(1/2*f*x + 1/2*e)^2 + 5*A*a^3*tan(1/2*f*x + 1/2*e) - 8*I*B*a^3*tan(1/2*f*x + 1/2*e))/(tan(1/2*f*x + 1/2*e)^3 + I*tan(1/2*f*x + 1/2*e)^2 - tan(1/2*f*x + 1/2*e) - I)^2*c)/f`

3.697.9 Mupad [B] (verification not implemented)

Time = 8.63 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.17

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{c - ic \tan(e + fx)} dx$$

$$= \frac{B a^3 \tan(e + fx)^2}{2 c f} + \frac{4 A a^3 - B a^3 8i + B a^3 4i}{f (\tan(e + fx) + 1i)}$$

$$- \frac{\tan(e + fx) \left(\frac{B a^3 2i}{c} + \frac{a^3 (2B + A 1i) 1i}{c} \right)}{f} - \frac{\ln(\tan(e + fx) + 1i) \left(\frac{8 B a^3}{c} + \frac{A a^3 4i}{c} \right)}{f}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3)/(c - c*tan(e + f*x)*1 i),x)`

3.697. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{c-ic \tan(e+fx)} dx$

output $((4Aa^3 - B a^3 8i)/c + (B a^3 4i)/c)/(f(\tan(e + fx) + 1i)) - (\log(\tan(e + fx) + 1i)((A a^3 4i)/c + (8B a^3)/c))/f - (\tan(e + fx)((B a^3 2i)/c + (a^3(A 1i + 2B) 1i)/c))/f + (B a^3 \tan(e + fx)^2)/(2c f)$

3.698
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^2} dx$$

3.698.1 Optimal result 6510
 3.698.2 Mathematica [A] (verified) 6510
 3.698.3 Rubi [A] (verified) 6511
 3.698.4 Maple [A] (verified) 6513
 3.698.5 Fricas [A] (verification not implemented) 6513
 3.698.6 Sympy [A] (verification not implemented) 6514
 3.698.7 Maxima [F(-2)] 6514
 3.698.8 Giac [B] (verification not implemented) 6515
 3.698.9 Mupad [B] (verification not implemented) 6515

3.698.1 Optimal result

Integrand size = 41, antiderivative size = 123

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx$$

$$= \frac{a^3(A - 5iB)x}{c^2} - \frac{a^3(iA + 5B) \log(\cos(e + fx))}{c^2 f} + \frac{ia^3 B \tan(e + fx)}{c^2 f}$$

$$+ \frac{2a^3(iA + B)}{c^2 f(i + \tan(e + fx))^2} - \frac{4a^3(A - 2iB)}{c^2 f(i + \tan(e + fx))}$$

output

```
a^3*(A-5*I*B)*x/c^2-a^3*(I*A+5*B)*ln(cos(f*x+e))/c^2/f+I*a^3*B*tan(f*x+e)/
c^2/f+2*a^3*(I*A+B)/c^2/f/(I+tan(f*x+e))^2-4*a^3*(A-2*I*B)/c^2/f/(I+tan(f*
x+e))
```

3.698.2 Mathematica [A] (verified)

Time = 5.60 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.73

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx$$

$$= \frac{B(a+ia \tan(e+fx))^3}{(c-ic \tan(e+fx))^2} + \frac{a^3(iA+5B) \left(\log(i+\tan(e+fx)) + \frac{-2+4i \tan(e+fx)}{(i+\tan(e+fx))^2} \right)}{c^2}$$

f

input `Integrate[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^2,x]`

output `((B*(a + I*a*Tan[e + f*x])^3)/(c - I*c*Tan[e + f*x])^2 + (a^3*(I*A + 5*B)*(Log[I + Tan[e + f*x]] + (-2 + (4*I)*Tan[e + f*x])/(I + Tan[e + f*x])^2))/c^2)/f`

3.698.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.69, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{a^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{c^3 (1 - i \tan(e + fx))^3} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^3 \int \frac{(i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{(1 - i \tan(e + fx))^3} d \tan(e + fx)}{c^2 f} \\
 & \quad \downarrow \text{86} \\
 & \frac{a^3 \int \left(-\frac{4i(A - iB)}{(\tan(e + fx) + i)^3} + iB + \frac{i(A - 5iB)}{\tan(e + fx) + i} + \frac{4(A - 2iB)}{(\tan(e + fx) + i)^2} \right) d \tan(e + fx)}{c^2 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^3 \left(-\frac{4(A - 2iB)}{\tan(e + fx) + i} + \frac{2(B + iA)}{(\tan(e + fx) + i)^2} + (5B + iA) \log(\tan(e + fx) + i) + iB \tan(e + fx) \right)}{c^2 f}
 \end{aligned}$$

3.698. $\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^2} dx$

input `Int[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^2,x]`

output `(a^3*((I*A + 5*B)*Log[I + Tan[e + f*x]] + I*B*Tan[e + f*x] + (2*(I*A + B))/(I + Tan[e + f*x])^2 - (4*(A - (2*I)*B))/(I + Tan[e + f*x]))/(c^2*f)`

3.698.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.698.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.54

method	result
risch	$-\frac{e^{4i(fx+e)}a^3B}{2c^2f} - \frac{ie^{4i(fx+e)}a^3A}{2c^2f} + \frac{3e^{2i(fx+e)}a^3B}{c^2f} + \frac{ie^{2i(fx+e)}a^3A}{c^2f} + \frac{10ia^3Be}{c^2f} - \frac{2a^3Ae}{c^2f} - \frac{2Ba^3}{fc^2(e^{2i(fx+e)})}$
derivativedivides	$\frac{ia^3B \tan(fx+e)}{c^2f} + \frac{8ia^3B}{fc^2(i+\tan(fx+e))} - \frac{4a^3A}{fc^2(i+\tan(fx+e))} + \frac{ia^3A \ln(1+\tan(fx+e)^2)}{2fc^2} + \frac{5a^3B \ln(1+\tan(fx+e))}{2fc^2}$
default	$\frac{ia^3B \tan(fx+e)}{c^2f} + \frac{8ia^3B}{fc^2(i+\tan(fx+e))} - \frac{4a^3A}{fc^2(i+\tan(fx+e))} + \frac{ia^3A \ln(1+\tan(fx+e)^2)}{2fc^2} + \frac{5a^3B \ln(1+\tan(fx+e))}{2fc^2}$
norman	$\frac{(-5iBa^3+a^3A)x}{c} + \frac{2iAa^3+6Ba^3}{cf} + \frac{(-5iBa^3+a^3A)x \tan(fx+e)^4}{c} + \frac{iBa^3 \tan(fx+e)^5}{cf} + \frac{2(-5iBa^3+a^3A)x \tan(fx+e)^2}{c} - \frac{2(-5iBa^3+a^3A)x}{c} \bigg/ c(1+\tan(fx+e)^2)^2$

input `int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x,method=_R
ETURNVERBOSE)`

output `-1/2/c^2/f*exp(4*I*(f*x+e))*a^3*B-1/2*I/c^2/f*exp(4*I*(f*x+e))*a^3*A+3/c^2
/f*exp(2*I*(f*x+e))*a^3*B+I/c^2/f*exp(2*I*(f*x+e))*a^3*A+10*I*a^3/c^2/f*B*
e-2*a^3/c^2/f*A*e-2/f/c^2*B*a^3/(exp(2*I*(f*x+e))+1)-5*a^3/c^2/f*ln(exp(2*
I*(f*x+e))+1)*B-I*a^3/c^2/f*ln(exp(2*I*(f*x+e))+1)*A`

3.698.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.12

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx$$

$$= \frac{(-iA - B)a^3e^{(6ifx+6ie)} + (iA + 5B)a^3e^{(4ifx+4ie)} - 2(-iA - 3B)a^3e^{(2ifx+2ie)} - 4Ba^3 - 2((iA + 5B)a^3)}{2(c^2fe^{(2ifx+2ie)} + c^2f)}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x, al
gorithm="fracas")`

output `1/2*((-I*A - B)*a^3*e^(6*I*f*x + 6*I*e) + (I*A + 5*B)*a^3*e^(4*I*f*x + 4*I
e) - 2(-I*A - 3*B)*a^3*e^(2*I*f*x + 2*I*e) - 4*B*a^3 - 2*((I*A + 5*B)*a^
3*e^(2*I*f*x + 2*I*e) + (I*A + 5*B)*a^3)*log(e^(2*I*f*x + 2*I*e) + 1))/(c^
2*f*e^(2*I*f*x + 2*I*e) + c^2*f)`

3.698. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^2} dx$

3.698.6 Sympy [A] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.92

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx$$

$$= -\frac{2Ba^3}{c^2 f e^{2ie} e^{2ifx} + c^2 f} - \frac{ia^3 (A - 5iB) \log(e^{2ifx} + e^{-2ie})}{c^2 f}$$

$$+ \begin{cases} \frac{(2iAa^3 c^2 f e^{2ie} + 6Ba^3 c^2 f e^{2ie}) e^{2ifx} + (-iAa^3 c^2 f e^{4ie} - Ba^3 c^2 f e^{4ie}) e^{4ifx}}{2c^4 f^2} & \text{for } c^4 f^2 \neq 0 \\ \frac{x(2Aa^3 e^{4ie} - 2Aa^3 e^{2ie} - 2iBa^3 e^{4ie} + 6iBa^3 e^{2ie})}{c^2} & \text{otherwise} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**2,x)`output `-2*B*a**3/(c**2*f*exp(2*I*e)*exp(2*I*f*x) + c**2*f) - I*a**3*(A - 5*I*B)*log(exp(2*I*f*x) + exp(-2*I*e))/(c**2*f) + Piecewise((((2*I*A*a**3*c**2*f*exp(2*I*e) + 6*B*a**3*c**2*f*exp(2*I*e))*exp(2*I*f*x) + (-I*A*a**3*c**2*f*exp(4*I*e) - B*a**3*c**2*f*exp(4*I*e))*exp(4*I*f*x))/(2*c**4*f**2), Ne(c**4*f**2, 0)), (x*(2*A*a**3*exp(4*I*e) - 2*A*a**3*exp(2*I*e) - 2*I*B*a**3*exp(4*I*e) + 6*I*B*a**3*exp(2*I*e))/c**2, True))`**3.698.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x, algorithm="maxima")`output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.698.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 341 vs. $2(109) = 218$.

Time = 0.64 (sec) , antiderivative size = 341, normalized size of antiderivative = 2.77

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx$$

$$= \frac{6(-iAa^3 - 5Ba^3) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)}{c^2} + \frac{12(iAa^3 + 5Ba^3) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + i)}{c^2} - \frac{6(iAa^3 + 5Ba^3) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)}{c^2} - \frac{6(-iAa^3 - 5Ba^3) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - i)}{c^2}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x, algorithm="giac")`

output `1/6*(6*(-I*A*a^3 - 5*B*a^3)*log(tan(1/2*f*x + 1/2*e) + 1)/c^2 + 12*(I*A*a^3 + 5*B*a^3)*log(tan(1/2*f*x + 1/2*e) + I)/c^2 - 6*(I*A*a^3 + 5*B*a^3)*log(tan(1/2*f*x + 1/2*e) - 1)/c^2 - 6*(-I*A*a^3*tan(1/2*f*x + 1/2*e)^2 - 5*B*a^3*tan(1/2*f*x + 1/2*e)^2 + 2*I*B*a^3*tan(1/2*f*x + 1/2*e) + I*A*a^3 + 5*B*a^3)/((tan(1/2*f*x + 1/2*e)^2 - 1)*c^2) - (25*I*A*a^3*tan(1/2*f*x + 1/2*e)^4 + 125*B*a^3*tan(1/2*f*x + 1/2*e)^4 - 100*A*a^3*tan(1/2*f*x + 1/2*e)^3 + 548*I*B*a^3*tan(1/2*f*x + 1/2*e)^3 - 198*I*A*a^3*tan(1/2*f*x + 1/2*e)^2 - 894*B*a^3*tan(1/2*f*x + 1/2*e)^2 + 100*A*a^3*tan(1/2*f*x + 1/2*e) - 548*I*B*a^3*tan(1/2*f*x + 1/2*e) + 25*I*A*a^3 + 125*B*a^3)/(c^2*(tan(1/2*f*x + 1/2*e) + I)^4))/f`

3.698.9 Mupad [B] (verification not implemented)

Time = 9.10 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.48

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^2} dx =$$

$$\frac{a^3 (7B \tan(e + fx) + B6i + A \tan(e + fx) 4i - 2A + B \tan(e + fx)^2 2i + B \tan(e + fx)^3 - A \ln(t$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3)/(c - c*tan(e + f*x)*1i)^2,x)`

output $-(a^3(B6i - 2A + A\tan(e + f*x)*4i + 7*B*\tan(e + f*x) + B*\tan(e + f*x)^2*2i + B*\tan(e + f*x)^3 - A*\log(\tan(e + f*x) + 1i) + B*\log(\tan(e + f*x) + 1i)*5i + A*\log(\tan(e + f*x) + 1i)*\tan(e + f*x)*2i + 10*B*\log(\tan(e + f*x) + 1i)*\tan(e + f*x) + A*\log(\tan(e + f*x) + 1i)*\tan(e + f*x)^2 - B*\log(\tan(e + f*x) + 1i)*\tan(e + f*x)^2*5i*1i)/(c^2*f*(\tan(e + f*x)*1i - 1)^2)$

3.698. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^2} dx$

3.699
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^3} dx$$

3.699.1 Optimal result 6517
 3.699.2 Mathematica [A] (verified) 6517
 3.699.3 Rubi [A] (verified) 6518
 3.699.4 Maple [A] (verified) 6520
 3.699.5 Fricas [A] (verification not implemented) 6520
 3.699.6 Sympy [A] (verification not implemented) 6521
 3.699.7 Maxima [F(-2)] 6521
 3.699.8 Giac [B] (verification not implemented) 6522
 3.699.9 Mupad [B] (verification not implemented) 6522

3.699.1 Optimal result

Integrand size = 41, antiderivative size = 129

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx$$

$$= \frac{ia^3 Bx}{c^3} + \frac{a^3 B \log(\cos(e + fx))}{c^3 f} - \frac{a^3(iA + B)(1 + i \tan(e + fx))^3}{6c^3 f(1 - i \tan(e + fx))^3}$$

$$- \frac{2a^3 B}{c^3 f(i + \tan(e + fx))^2} - \frac{4ia^3 B}{c^3 f(i + \tan(e + fx))}$$

output `I*a^3*B*x/c^3+a^3*B*ln(cos(f*x+e))/c^3/f-1/6*a^3*(I*A+B)*(1+I*tan(f*x+e))^3/c^3/f/(1-I*tan(f*x+e))^3-2*a^3*B/c^3/f/(I+tan(f*x+e))^2-4*I*a^3*B/c^3/f/(I+tan(f*x+e))`

3.699.2 Mathematica [A] (verified)

Time = 4.96 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.57

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx$$

$$= - \frac{a^3 \left(3B \log(i + \tan(e + fx)) + \frac{A - 7iB - 18B \tan(e + fx) - 3(A - 5iB) \tan^2(e + fx)}{(i + \tan(e + fx))^3} \right)}{3c^3 f}$$

input `Integrate[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^3,x]`

output `-1/3*(a^3*(3*B*Log[I + Tan[e + f*x]] + (A - (7*I)*B - 18*B*Tan[e + f*x] - 3*(A - (5*I)*B)*Tan[e + f*x]^2)/(I + Tan[e + f*x]^3))/(c^3*f)`

3.699.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.78, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.146$, Rules used = {3042, 4071, 27, 87, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{a^2(i \tan(e+fx)+1)^2(A+B \tan(e+fx))}{c^4(1-i \tan(e+fx))^4} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^3 \int \frac{(i \tan(e+fx)+1)^2(A+B \tan(e+fx))}{(1-i \tan(e+fx))^4} d \tan(e + fx)}{c^3 f} \\
 & \quad \downarrow \text{87} \\
 & \frac{a^3 \left(iB \int \frac{(i \tan(e+fx)+1)^2}{(1-i \tan(e+fx))^3} d \tan(e + fx) - \frac{(B+iA)(1+i \tan(e+fx))^3}{6(1-i \tan(e+fx))^3} \right)}{c^3 f} \\
 & \quad \downarrow \text{49} \\
 & \frac{a^3 \left(iB \int \left(\frac{i}{\tan(e+fx)+i} + \frac{4}{(\tan(e+fx)+i)^2} - \frac{4i}{(\tan(e+fx)+i)^3} \right) d \tan(e + fx) - \frac{(B+iA)(1+i \tan(e+fx))^3}{6(1-i \tan(e+fx))^3} \right)}{c^3 f} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

3.699. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^3} dx$

$$\frac{a^3 \left(iB \left(-\frac{4}{\tan(e+fx)+i} + \frac{2i}{(\tan(e+fx)+i)^2} + i \log(\tan(e+fx)+i) \right) - \frac{(B+iA)(1+i \tan(e+fx))^3}{6(1-i \tan(e+fx))^3} \right)}{c^3 f}$$

input `Int[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^3,x]`

output `(a^3*(-1/6*((I*A + B)*(1 + I*Tan[e + f*x])^3)/(1 - I*Tan[e + f*x])^3 + I*B*(I*Log[I + Tan[e + f*x]] + (2*I)/(I + Tan[e + f*x])^2 - 4/(I + Tan[e + f*x]))) / (c^3*f)`

3.699.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 49 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x],
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.699.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.96

method	result
risch	$-\frac{e^{6i(fx+e)}a^3B}{6c^3f} - \frac{ie^{6i(fx+e)}a^3A}{6c^3f} + \frac{Ba^3e^{4i(fx+e)}}{2c^3f} - \frac{Ba^3e^{2i(fx+e)}}{c^3f} - \frac{2iBa^3e}{c^3f} + \frac{Ba^3 \ln(e^{2i(fx+e)}+1)}{c^3f}$
derivativedivides	$\frac{4ia^3B}{3fc^3(i+\tan(fx+e))^3} - \frac{4a^3A}{3fc^3(i+\tan(fx+e))^3} - \frac{2ia^3A}{fc^3(i+\tan(fx+e))^2} - \frac{4a^3B}{c^3f(i+\tan(fx+e))^2} - \frac{5ia^3B}{fc^3(i+\tan(fx+e))}$
default	$\frac{4ia^3B}{3fc^3(i+\tan(fx+e))^3} - \frac{4a^3A}{3fc^3(i+\tan(fx+e))^3} - \frac{2ia^3A}{fc^3(i+\tan(fx+e))^2} - \frac{4a^3B}{c^3f(i+\tan(fx+e))^2} - \frac{5ia^3B}{fc^3(i+\tan(fx+e))}$
norman	$\frac{(-iBa^3+a^3A)\tan(fx+e)}{cf} + \frac{(-5iBa^3+a^3A)\tan(fx+e)^5}{cf} + \frac{iBa^3x}{c} + \frac{iBa^3x\tan(fx+e)^6}{c} - \frac{iAa^3+7Ba^3}{3cf} - \frac{2(iBa^3+5a^3A)\tan(fx+e)}{3cf}$ $c^2(1+\tan(fx+e))$

```
input int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x,method=_R
ETURNVERBOSE)
```

```
output -1/6/c^3/f*exp(6*I*(f*x+e))*a^3*B-1/6*I/c^3/f*exp(6*I*(f*x+e))*a^3*A+1/2/c
^3/f*B*a^3*exp(4*I*(f*x+e))-1/c^3/f*B*a^3*exp(2*I*(f*x+e))-2*I/c^3/f*B*a^3
*e+1/c^3/f*B*a^3*ln(exp(2*I*(f*x+e))+1)
```

3.699.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.60

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^3} dx$$

$$= \frac{(-iA - B)a^3e^{(6ifx+6ie)} + 3Ba^3e^{(4ifx+4ie)} - 6Ba^3e^{(2ifx+2ie)} + 6Ba^3 \log(e^{(2ifx+2ie)} + 1)}{6c^3f}$$

```
input integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x, al
gorithm="fracas")
```

3.699.
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^3} dx$$

output $1/6*((-I*A - B)*a^3*e^{(6*I*f*x + 6*I*e)} + 3*B*a^3*e^{(4*I*f*x + 4*I*e)} - 6*B*a^3*e^{(2*I*f*x + 2*I*e)} + 6*B*a^3*\log(e^{(2*I*f*x + 2*I*e)} + 1))/(c^3*f)$

3.699.6 Sympy [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.64

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx$$

$$= \frac{Ba^3 \log(e^{2ifx} + e^{-2ie})}{c^3 f}$$

$$+ \begin{cases} \frac{6Ba^3 c^6 f^2 e^{4ie} e^{4ifx} - 12Ba^3 c^6 f^2 e^{2ie} e^{2ifx} + (-2iAa^3 c^6 f^2 e^{6ie} - 2Ba^3 c^6 f^2 e^{6ie}) e^{6ifx}}{12c^9 f^3} & \text{for } c^9 f^3 \neq 0 \\ \frac{x(Aa^3 e^{6ie} - iBa^3 e^{6ie} + 2iBa^3 e^{4ie} - 2iBa^3 e^{2ie})}{c^3} & \text{otherwise} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**3,x)`

output `B*a**3*log(exp(2*I*f*x) + exp(-2*I*e))/(c**3*f) + Piecewise(((6*B*a**3*c**6*f**2*exp(4*I*e)*exp(4*I*f*x) - 12*B*a**3*c**6*f**2*exp(2*I*e)*exp(2*I*f*x) + (-2*I*A*a**3*c**6*f**2*exp(6*I*e) - 2*B*a**3*c**6*f**2*exp(6*I*e))*exp(6*I*f*x))/(12*c**9*f**3), Ne(c**9*f**3, 0)), (x*(A*a**3*exp(6*I*e) - I*B*a**3*exp(6*I*e) + 2*I*B*a**3*exp(4*I*e) - 2*I*B*a**3*exp(2*I*e))/c**3, True))`

3.699.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.699.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 242 vs. $2(113) = 226$.

Time = 0.87 (sec) , antiderivative size = 242, normalized size of antiderivative = 1.88

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx$$

$$= \frac{30 Ba^3 \log(\tan(\frac{1}{2} fx + \frac{1}{2} e) + 1)}{c^3} - \frac{60 Ba^3 \log(\tan(\frac{1}{2} fx + \frac{1}{2} e) + i)}{c^3} + \frac{30 Ba^3 \log(\tan(\frac{1}{2} fx + \frac{1}{2} e) - 1)}{c^3} + \frac{147 Ba^3 \tan(\frac{1}{2} fx + \frac{1}{2} e)^6 - 60 Aa^3 \tan(\frac{1}{2} fx + \frac{1}{2} e)^5 + 942 I B a^3 \tan(\frac{1}{2} fx + \frac{1}{2} e)^4 - 2445 B a^3 \tan(\frac{1}{2} fx + \frac{1}{2} e)^3 + 200 A a^3 \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 - 3620 I B a^3 \tan(\frac{1}{2} fx + \frac{1}{2} e) + 2445 B a^3 \tan(\frac{1}{2} fx + \frac{1}{2} e) - 60 A a^3}{c^3 (\tan(\frac{1}{2} fx + \frac{1}{2} e) + I)^6} / f$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x, algorithm="giac")`

output `1/30*(30*B*a^3*log(tan(1/2*f*x + 1/2*e) + 1)/c^3 - 60*B*a^3*log(tan(1/2*f*x + 1/2*e) + I)/c^3 + 30*B*a^3*log(tan(1/2*f*x + 1/2*e) - 1)/c^3 + (147*B*a^3*tan(1/2*f*x + 1/2*e)^6 - 60*A*a^3*tan(1/2*f*x + 1/2*e)^5 + 942*I*B*a^3*tan(1/2*f*x + 1/2*e)^4 - 2445*B*a^3*tan(1/2*f*x + 1/2*e)^3 + 200*A*a^3*tan(1/2*f*x + 1/2*e)^2 - 3620*I*B*a^3*tan(1/2*f*x + 1/2*e) + 2445*B*a^3*tan(1/2*f*x + 1/2*e) - 60*A*a^3)/(c^3*(tan(1/2*f*x + 1/2*e) + I)^6))/f`

3.699.9 Mupad [B] (verification not implemented)

Time = 8.69 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.09

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^3} dx =$$

$$\frac{a^3 (15 B \tan(e + fx)^2 - 7 B + B \tan(e + fx) 18i + A \tan(e + fx)^2 3i - A 1i - 3 B \ln(\tan(e + fx) + i))}{(c - ic \tan(e + fx))^3}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3)/(c - c*tan(e + f*x)*1i)^3,x)`

output `-(a^3*(B*tan(e + f*x)*18i - 7*B - A*1i + A*tan(e + f*x)^2*3i + 15*B*tan(e + f*x)^2 - 3*B*log(tan(e + f*x) + 1i) + B*log(tan(e + f*x) + 1i)*tan(e + f*x)*9i + 9*B*log(tan(e + f*x) + 1i)*tan(e + f*x)^2 - B*log(tan(e + f*x) + 1i)*tan(e + f*x)^3*3i))/(3*c^3*f*(tan(e + f*x)*1i - 1)^3)`

3.699. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^3} dx$

3.700
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^4} dx$$

3.700.1 Optimal result	6523
3.700.2 Mathematica [A] (verified)	6523
3.700.3 Rubi [A] (verified)	6524
3.700.4 Maple [A] (verified)	6525
3.700.5 Fricas [A] (verification not implemented)	6526
3.700.6 Sympy [B] (verification not implemented)	6527
3.700.7 Maxima [F(-2)]	6527
3.700.8 Giac [B] (verification not implemented)	6528
3.700.9 Mupad [B] (verification not implemented)	6528

3.700.1 Optimal result

Integrand size = 41, antiderivative size = 99

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx$$

$$= -\frac{a^3(iA + B)(1 + i \tan(e + fx))^3}{8c^4 f(1 - i \tan(e + fx))^4} - \frac{a^3(iA - 7B)(1 + i \tan(e + fx))^3}{48c^4 f(1 - i \tan(e + fx))^3}$$

output
$$-1/8*a^3*(I*A+B)*(1+I*\tan(f*x+e))^3/c^4/f/(1-I*\tan(f*x+e))^4-1/48*a^3*(I*A-7*B)*(1+I*\tan(f*x+e))^3/c^4/f/(1-I*\tan(f*x+e))^3$$

3.700.2 Mathematica [A] (verified)

Time = 3.53 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.80

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx$$

$$= \frac{a^3(-iA + B + 2(A - 2iB) \tan(e + fx) + 3i(A + iB) \tan^2(e + fx) + 6iB \tan^3(e + fx))}{6c^4 f(i + \tan(e + fx))^4}$$

input
$$\text{Integrate}[(a + I*a*\text{Tan}[e + f*x])^3*(A + B*\text{Tan}[e + f*x])/(c - I*c*\text{Tan}[e + f*x])^4,x]$$

output
$$(a^3*((-I)*A + B + 2*(A - (2*I)*B)*\text{Tan}[e + f*x] + (3*I)*(A + I*B)*\text{Tan}[e + f*x]^2 + (6*I)*B*\text{Tan}[e + f*x]^3))/(6*c^4*f*(I + \text{Tan}[e + f*x])^4)$$

3.700.
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^4} dx$$

3.700.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.92, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{a^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{c^5 (1 - i \tan(e + fx))^5} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^3 \int \frac{(i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{(1 - i \tan(e + fx))^5} d \tan(e + fx)}{c^4 f} \\
 & \quad \downarrow \text{87} \\
 & \frac{a^3 \left(\frac{1}{8} (A + 7iB) \int \frac{(i \tan(e + fx) + 1)^2}{(1 - i \tan(e + fx))^4} d \tan(e + fx) - \frac{(B + iA)(1 + i \tan(e + fx))^3}{8(1 - i \tan(e + fx))^4} \right)}{c^4 f} \\
 & \quad \downarrow \text{48} \\
 & \frac{a^3 \left(-\frac{i(A + 7iB)(1 + i \tan(e + fx))^3}{48(1 - i \tan(e + fx))^3} - \frac{(B + iA)(1 + i \tan(e + fx))^3}{8(1 - i \tan(e + fx))^4} \right)}{c^4 f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^4,x]`

output `(a^3*(-1/8*((I*A + B)*(1 + I*Tan[e + f*x])^3)/(1 - I*Tan[e + f*x])^4 - ((I/48)*(A + (7*I)*B)*(1 + I*Tan[e + f*x])^3)/(1 - I*Tan[e + f*x])^3))/(c^4*f)`

3.700.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.700.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.89

method	result
risch	$-\frac{a^3 e^{8i(fx+e)} B}{16c^4 f} - \frac{ia^3 e^{8i(fx+e)} A}{16c^4 f} + \frac{a^3 e^{6i(fx+e)} B}{12c^4 f} - \frac{ia^3 e^{6i(fx+e)} A}{12c^4 f}$
derivativedivides	$a^3 \left(\frac{iB}{i+\tan(fx+e)} - \frac{4iA+4B}{4(i+\tan(fx+e))^4} - \frac{-iA-5B}{2(i+\tan(fx+e))^2} - \frac{8iB-4A}{3(i+\tan(fx+e))^3} \right) \frac{1}{f c^4}$
default	$a^3 \left(\frac{iB}{i+\tan(fx+e)} - \frac{4iA+4B}{4(i+\tan(fx+e))^4} - \frac{-iA-5B}{2(i+\tan(fx+e))^2} - \frac{8iB-4A}{3(i+\tan(fx+e))^3} \right) \frac{1}{f c^4}$
norman	$\frac{a^3 A \tan(fx+e)}{fc} + \frac{iB a^3 \tan(fx+e)^7}{cf} + \frac{-iA a^3 + B a^3}{6cf} + \frac{(iA a^3 + 7B a^3) \tan(fx+e)^6}{2cf} + \frac{(17iA a^3 + 7B a^3) \tan(fx+e)^2}{6cf} + \frac{7(-2iB a^3 + A^2)}{(1+\tan(fx+e))^2} \frac{1}{c^3}$

input `int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x,method=_RETURNNVERBOSE)`

output
$$-1/16*a^3/c^4/f*\exp(8*I*(f*x+e))*B-1/16*I*a^3/c^4/f*\exp(8*I*(f*x+e))*A+1/12*a^3/c^4/f*\exp(6*I*(f*x+e))*B-1/12*I*a^3/c^4/f*\exp(6*I*(f*x+e))*A$$

3.700.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.49

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx$$

$$= -\frac{3(iA + B)a^3 e^{(8i fx + 8ie)} + 4(iA - B)a^3 e^{(6i fx + 6ie)}}{48 c^4 f}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x, algorithm="fracas")`

output
$$-1/48*(3*(I*A + B)*a^3*e^{(8*I*f*x + 8*I*e)} + 4*(I*A - B)*a^3*e^{(6*I*f*x + 6*I*e)})/(c^4*f)$$

3.700.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 167 vs. $2(80) = 160$.

Time = 0.37 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.69

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^4} dx$$

$$= \begin{cases} \frac{(-16iAa^3c^4fe^{6ie} + 16Ba^3c^4fe^{6ie})e^{6ifx} + (-12iAa^3c^4fe^{8ie} - 12Ba^3c^4fe^{8ie})e^{8ifx}}{192c^8f^2} & \text{for } c^8f^2 \neq 0 \\ \frac{x(Aa^3e^{8ie} + Aa^3e^{6ie} - iBa^3e^{8ie} + iBa^3e^{6ie})}{2c^4} & \text{otherwise} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**4,x)`

output `Piecewise(((((-16*I*A*a**3*c**4*f*exp(6*I*e) + 16*B*a**3*c**4*f*exp(6*I*e))*exp(6*I*f*x) + (-12*I*A*a**3*c**4*f*exp(8*I*e) - 12*B*a**3*c**4*f*exp(8*I*e))*exp(8*I*f*x))/(192*c**8*f**2), Ne(c**8*f**2, 0)), (x*(A*a**3*exp(8*I*e) + A*a**3*exp(6*I*e) - I*B*a**3*exp(8*I*e) + I*B*a**3*exp(6*I*e))/(2*c**4), True))`

3.700.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^4} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.700.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 224 vs. $2(83) = 166$.

Time = 1.03 (sec) , antiderivative size = 224, normalized size of antiderivative = 2.26

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx =$$

$$\frac{2 \left(3 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^7 + 3i A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^6 - 3 B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^6 - 17 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^5 \right)}{(c^4 f (\tan(e + fx)^4 + \tan(e + fx)^3 4i - 6 \tan(e + fx)^2 - \tan(e + fx) 4i + 1))}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x, algorithm="giac")`

output `-2/3*(3*A*a^3*tan(1/2*f*x + 1/2*e)^7 + 3*I*A*a^3*tan(1/2*f*x + 1/2*e)^6 - 3*B*a^3*tan(1/2*f*x + 1/2*e)^6 - 17*A*a^3*tan(1/2*f*x + 1/2*e)^5 + 4*I*B*a^3*tan(1/2*f*x + 1/2*e)^5 - 10*I*A*a^3*tan(1/2*f*x + 1/2*e)^4 + 10*B*a^3*tan(1/2*f*x + 1/2*e)^4 + 17*A*a^3*tan(1/2*f*x + 1/2*e)^3 - 4*I*B*a^3*tan(1/2*f*x + 1/2*e)^3 + 3*I*A*a^3*tan(1/2*f*x + 1/2*e)^2 - 3*B*a^3*tan(1/2*f*x + 1/2*e)^2 - 3*A*a^3*tan(1/2*f*x + 1/2*e))/(c^4*f*(tan(1/2*f*x + 1/2*e) + I)^8)`

3.700.9 Mupad [B] (verification not implemented)

Time = 8.70 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.19

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^4} dx$$

$$= \frac{-\frac{a^3(-B+Ai)}{6} + \frac{a^3 \tan(e+fx)(2A-B4i)}{6} + B a^3 \tan(e+fx)^3 i + \frac{a^3 \tan(e+fx)^2(-3B+A3i)}{6}}{c^4 f (\tan(e+fx)^4 + \tan(e+fx)^3 4i - 6 \tan(e+fx)^2 - \tan(e+fx) 4i + 1)}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3)/(c - c*tan(e + f*x)*1i)^4,x)`

output `((a^3*tan(e + f*x)*(2*A - B*4i))/6 - (a^3*(A*1i - B))/6 + B*a^3*tan(e + f*x)^3*1i + (a^3*tan(e + f*x)^2*(A*3i - 3*B))/6)/(c^4*f*(tan(e + f*x)^3*4i - 6*tan(e + f*x)^2 - tan(e + f*x)*4i + tan(e + f*x)^4 + 1))`

3.700. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^4} dx$

3.701
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^5} dx$$

3.701.1 Optimal result 6529
 3.701.2 Mathematica [A] (verified) 6529
 3.701.3 Rubi [A] (verified) 6530
 3.701.4 Maple [A] (verified) 6532
 3.701.5 Fricas [A] (verification not implemented) 6532
 3.701.6 Sympy [B] (verification not implemented) 6533
 3.701.7 Maxima [F(-2)] 6533
 3.701.8 Giac [B] (verification not implemented) 6534
 3.701.9 Mupad [B] (verification not implemented) 6534

3.701.1 Optimal result

Integrand size = 41, antiderivative size = 122

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx$$

$$= \frac{4a^3(A - iB)}{5c^5 f(i + \tan(e + fx))^5} + \frac{a^3(iA + 2B)}{c^5 f(i + \tan(e + fx))^4}$$

$$- \frac{a^3(A - 5iB)}{3c^5 f(i + \tan(e + fx))^3} - \frac{a^3 B}{2c^5 f(i + \tan(e + fx))^2}$$

output $4/5*a^3*(A-I*B)/c^5/f/(I+\tan(f*x+e))^5+a^3*(I*A+2*B)/c^5/f/(I+\tan(f*x+e))^4-1/3*a^3*(A-5*I*B)/c^5/f/(I+\tan(f*x+e))^3-1/2*a^3*B/c^5/f/(I+\tan(f*x+e))^2$

3.701.2 Mathematica [A] (verified)

Time = 5.64 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.64

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx$$

$$= \frac{a^3(4A + iB + 5(2iA + B) \tan(e + fx) + (-10A + 5iB) \tan^2(e + fx) - 15B \tan^3(e + fx))}{30c^5 f(i + \tan(e + fx))^5}$$

input `Integrate[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^5,x]`

output $(a^3(4A + I*B + 5*((2*I)*A + B)*\text{Tan}[e + f*x] + (-10*A + (5*I)*B)*\text{Tan}[e + f*x]^2 - 15*B*\text{Tan}[e + f*x]^3))/(30*c^5*f*(I + \text{Tan}[e + f*x])^5)$

3.701.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.79, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{a^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{c^6 (1 - i \tan(e + fx))^6} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{a^3 \int \frac{(i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{(1 - i \tan(e + fx))^6} d \tan(e + fx)}{c^5 f} \\ & \quad \downarrow \text{86} \\ & \frac{a^3 \int \left(-\frac{4(A - iB)}{(\tan(e + fx) + i)^6} + \frac{B}{(\tan(e + fx) + i)^3} + \frac{A - 5iB}{(\tan(e + fx) + i)^4} - \frac{4i(A - 2iB)}{(\tan(e + fx) + i)^5} \right) d \tan(e + fx)}{c^5 f} \\ & \quad \downarrow \text{2009} \\ & \frac{a^3 \left(\frac{4(A - iB)}{5(\tan(e + fx) + i)^5} - \frac{A - 5iB}{3(\tan(e + fx) + i)^3} + \frac{2B + iA}{(\tan(e + fx) + i)^4} - \frac{B}{2(\tan(e + fx) + i)^2} \right)}{c^5 f} \end{aligned}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^3*(A + B*\text{Tan}[e + f*x])]/(c - I*c*\text{Tan}[e + f*x])^5, x]$

3.701. $\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx$

```
output (a^3*((4*(A - I*B))/(5*(I + Tan[e + f*x])^5) + (I*A + 2*B)/(I + Tan[e + f*
x])^4 - (A - (5*I)*B)/(3*(I + Tan[e + f*x])^3) - B/(2*(I + Tan[e + f*x])^2
)))/(c^5*f)
```

3.701.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.701.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.71

method	result
derivativedivides	$\frac{a^3 \left(-\frac{4iB-4A}{5(i+\tan(fx+e))^5} - \frac{-4iA-8B}{4(i+\tan(fx+e))^4} - \frac{-5iB+A}{3(i+\tan(fx+e))^3} - \frac{B}{2(i+\tan(fx+e))^2} \right)}{f c^5}$
default	$\frac{a^3 \left(-\frac{4iB-4A}{5(i+\tan(fx+e))^5} - \frac{-4iA-8B}{4(i+\tan(fx+e))^4} - \frac{-5iB+A}{3(i+\tan(fx+e))^3} - \frac{B}{2(i+\tan(fx+e))^2} \right)}{f c^5}$
risch	$-\frac{a^3 e^{10i(fx+e)} B}{40c^5 f} - \frac{ia^3 e^{10i(fx+e)} A}{40c^5 f} - \frac{ia^3 A e^{8i(fx+e)}}{16c^5 f} + \frac{a^3 e^{6i(fx+e)} B}{24c^5 f} - \frac{ia^3 e^{6i(fx+e)} A}{24c^5 f}$
norman	$\frac{a^3 A \tan(fx+e)}{fc} + \frac{-4iA a^3 + B a^3}{30cf} - \frac{(-8iB a^3 + 19a^3 A) \tan(fx+e)^3}{3cf} + \frac{7(-16iB a^3 + 11a^3 A) \tan(fx+e)^5}{15cf} - \frac{(-8iB a^3 + a^3 A) \tan(fx+e)}{3cf} (1+\tan(fx+e))$

input `int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^5,x,method=_RETURVERBOSE)`

output `1/f*a^3/c^5*(-1/5*(-4*A+4*I*B)/(I+tan(f*x+e))^5-1/4*(-4*I*A-8*B)/(I+tan(f*x+e))^4-1/3*(A-5*I*B)/(I+tan(f*x+e))^3-1/2*B/(I+tan(f*x+e))^2)`

3.701.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.52

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ictan(e + fx))^5} dx$$

$$= -\frac{6(iA + B)a^3 e^{(10i fx + 10ie)} + 15iAa^3 e^{(8i fx + 8ie)} + 10(iA - B)a^3 e^{(6i fx + 6ie)}}{240 c^5 f}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^5,x, algorithm="fracas")`

output `-1/240*(6*(I*A + B)*a^3*e^(10*I*f*x + 10*I*e) + 15*I*A*a^3*e^(8*I*f*x + 8*I*e) + 10*(I*A - B)*a^3*e^(6*I*f*x + 6*I*e))/(c^5*f)`

3.701.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 218 vs. $2(100) = 200$.

Time = 0.46 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.79

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx$$

$$= \begin{cases} \frac{-960iAa^3c^{10}f^2e^{8ie}e^{8ifx} + (-640iAa^3c^{10}f^2e^{6ie} + 640Ba^3c^{10}f^2e^{6ie})e^{6ifx} + (-384iAa^3c^{10}f^2e^{10ie} - 384Ba^3c^{10}f^2e^{10ie})e^{10ifx}}{15360c^{15}f^3} & \text{for } c^{15}f^3 \\ \frac{x(Aa^3e^{10ie} + 2Aa^3e^{8ie} + Aa^3e^{6ie} - iBa^3e^{10ie} + iBa^3e^{6ie})}{4c^5} & \text{otherwise} \end{cases}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**5,x)`

output `Piecewise(((−960*I*A*a**3*c**10*f**2*exp(8*I*e)*exp(8*I*f*x) + (−640*I*A*a**3*c**10*f**2*exp(6*I*e) + 640*B*a**3*c**10*f**2*exp(6*I*e))*exp(6*I*f*x) + (−384*I*A*a**3*c**10*f**2*exp(10*I*e) − 384*B*a**3*c**10*f**2*exp(10*I*e))*exp(10*I*f*x))/(15360*c**15*f**3), Ne(c**15*f**3, 0)), (x*(A*a**3*exp(10*I*e) + 2*A*a**3*exp(8*I*e) + A*a**3*exp(6*I*e) − I*B*a**3*exp(10*I*e) + I*B*a**3*exp(6*I*e))/(4*c**5), True))`

3.701.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^5,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.701.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 292 vs. $2(102) = 204$.

Time = 1.30 (sec) , antiderivative size = 292, normalized size of antiderivative = 2.39

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx =$$

$$\frac{2 \left(15 A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^9 + 30 i A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^8 - 15 B a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^8 - 140 A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^7 + 10 i B a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^7 - 170 i A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^6 + 65 B a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^6 + 282 A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^5 - 12 i B a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^5 + 170 i A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^4 - 65 B a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^4 - 140 A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^3 + 10 i B a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^3 - 30 i A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^2 + 15 B a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^2 + 15 A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) \right) / (c^5 f (\tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) + i)^{10}}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^5,x, algorithm="giac")`

output `-2/15*(15*A*a^3*tan(1/2*f*x + 1/2*e)^9 + 30*I*A*a^3*tan(1/2*f*x + 1/2*e)^8 - 15*B*a^3*tan(1/2*f*x + 1/2*e)^8 - 140*A*a^3*tan(1/2*f*x + 1/2*e)^7 + 10*I*B*a^3*tan(1/2*f*x + 1/2*e)^7 - 170*I*A*a^3*tan(1/2*f*x + 1/2*e)^6 + 65*B*a^3*tan(1/2*f*x + 1/2*e)^6 + 282*A*a^3*tan(1/2*f*x + 1/2*e)^5 - 12*I*B*a^3*tan(1/2*f*x + 1/2*e)^5 + 170*I*A*a^3*tan(1/2*f*x + 1/2*e)^4 - 65*B*a^3*tan(1/2*f*x + 1/2*e)^4 - 140*A*a^3*tan(1/2*f*x + 1/2*e)^3 + 10*I*B*a^3*tan(1/2*f*x + 1/2*e)^3 - 30*I*A*a^3*tan(1/2*f*x + 1/2*e)^2 + 15*B*a^3*tan(1/2*f*x + 1/2*e)^2 + 15*A*a^3*tan(1/2*f*x + 1/2*e))/(c^5*f*(tan(1/2*f*x + 1/2*e) + I)^10)`

3.701.9 Mupad [B] (verification not implemented)

Time = 8.70 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.05

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^5} dx$$

$$= \frac{\frac{a^3 (4A + B i)}{30} + \frac{a^3 \tan(e + f x) (5B + A i)}{30} - \frac{B a^3 \tan(e + f x)^3}{2} - \frac{a^3 \tan(e + f x)^2 (10A - B i)}{30}}{c^5 f (\tan(e + f x)^5 + \tan(e + f x)^4 i - 10 \tan(e + f x)^3 - \tan(e + f x)^2 i + 5 \tan(e + f x) + i)}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*i)^3)/(c - c*tan(e + f*x)*i)^5,x)`

output $((a^3(4A + B1i))/30 + (a^3\tan(e + f*x)*(A*10i + 5*B))/30 - (B*a^3\tan(e + f*x)^3)/2 - (a^3\tan(e + f*x)^2*(10*A - B*5i))/30)/(c^5*f*(5*\tan(e + f*x) - \tan(e + f*x)^2*10i - 10*\tan(e + f*x)^3 + \tan(e + f*x)^4*5i + \tan(e + f*x)^5 + 1i))$

3.701. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^5} dx$

3.702
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^6} dx$$

3.702.1 Optimal result 6536
 3.702.2 Mathematica [A] (verified) 6536
 3.702.3 Rubi [A] (verified) 6537
 3.702.4 Maple [A] (verified) 6539
 3.702.5 Fricas [A] (verification not implemented) 6539
 3.702.6 Sympy [B] (verification not implemented) 6540
 3.702.7 Maxima [F(-2)] 6540
 3.702.8 Giac [B] (verification not implemented) 6541
 3.702.9 Mupad [B] (verification not implemented) 6541

3.702.1 Optimal result

Integrand size = 41, antiderivative size = 127

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx$$

$$= \frac{2a^3(iA + B)}{3c^6 f(i + \tan(e + fx))^6} - \frac{4a^3(A - 2iB)}{5c^6 f(i + \tan(e + fx))^5}$$

$$- \frac{a^3(iA + 5B)}{4c^6 f(i + \tan(e + fx))^4} - \frac{ia^3 B}{3c^6 f(i + \tan(e + fx))^3}$$

output $2/3*a^3*(I*A+B)/c^6/f/(I+\tan(f*x+e))^6-4/5*a^3*(A-2*I*B)/c^6/f/(I+\tan(f*x+e))^5-1/4*a^3*(I*A+5*B)/c^6/f/(I+\tan(f*x+e))^4-1/3*I*a^3*B/c^6/f/(I+\tan(f*x+e))^3$

3.702.2 Mathematica [A] (verified)

Time = 5.67 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.63

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx =$$

$$\frac{ia^3(-7A - iB + (-18iA - 6B) \tan(e + fx) + 15(A - iB) \tan^2(e + fx) + 20B \tan^3(e + fx))}{60c^6 f(i + \tan(e + fx))^6}$$

input `Integrate[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^6,x]`

output $((-1/60*I)*a^3*(-7*A - I*B + ((-18*I)*A - 6*B)*\text{Tan}[e + f*x] + 15*(A - I*B)*\text{Tan}[e + f*x]^2 + 20*B*\text{Tan}[e + f*x]^3))/(c^6*f*(I + \text{Tan}[e + f*x])^6)$

3.702.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.80, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{a^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{c^7 (1 - i \tan(e + fx))^7} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{a^3 \int \frac{(i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{(1 - i \tan(e + fx))^7} d \tan(e + fx)}{c^6 f} \\ & \quad \downarrow \text{86} \\ & \frac{a^3 \int \left(-\frac{4i(A - iB)}{(\tan(e + fx) + i)^7} + \frac{iB}{(\tan(e + fx) + i)^4} + \frac{iA + 5B}{(\tan(e + fx) + i)^5} + \frac{4(A - 2iB)}{(\tan(e + fx) + i)^6} \right) d \tan(e + fx)}{c^6 f} \\ & \quad \downarrow \text{2009} \\ & \frac{a^3 \left(-\frac{4(A - 2iB)}{5(\tan(e + fx) + i)^5} - \frac{5B + iA}{4(\tan(e + fx) + i)^4} + \frac{2(B + iA)}{3(\tan(e + fx) + i)^6} - \frac{iB}{3(\tan(e + fx) + i)^3} \right)}{c^6 f} \end{aligned}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^3*(A + B*\text{Tan}[e + f*x])]/(c - I*c*\text{Tan}[e + f*x])^6, x]$

3.702. $\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx$


```
output (a^3*((2*(I*A + B))/(3*(I + Tan[e + f*x])^6) - (4*(A - (2*I)*B))/(5*(I + Tan[e + f*x])^5) - (I*A + 5*B)/(4*(I + Tan[e + f*x])^4) - ((I/3)*B)/(I + Tan[e + f*x]^3))/(c^6*f)
```

3.702.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.702.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.71

method	result
derivativedivides	$\frac{a^3 \left(-\frac{iB}{3(i+\tan(fx+e))^3} - \frac{-4iA-4B}{6(i+\tan(fx+e))^6} - \frac{iA+5B}{4(i+\tan(fx+e))^4} - \frac{-8iB+4A}{5(i+\tan(fx+e))^5} \right)}{f c^6}$
default	$\frac{a^3 \left(-\frac{iB}{3(i+\tan(fx+e))^3} - \frac{-4iA-4B}{6(i+\tan(fx+e))^6} - \frac{iA+5B}{4(i+\tan(fx+e))^4} - \frac{-8iB+4A}{5(i+\tan(fx+e))^5} \right)}{f c^6}$
risch	$-\frac{a^3 e^{12i(fx+e)} B}{96c^6 f} - \frac{ia^3 e^{12i(fx+e)} A}{96c^6 f} - \frac{e^{10i(fx+e)} B a^3}{80c^6 f} - \frac{3ie^{10i(fx+e)} a^3 A}{80c^6 f} + \frac{e^{8i(fx+e)} B a^3}{64c^6 f} - \frac{3ie^{8i(fx+e)} a^3 A}{64c^6 f}$

```
input int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^6,x,method=_R
ETURNVERBOSE)
```

```
output 1/f*a^3/c^6*(-1/3*I*B/(I+tan(f*x+e))^3-1/6*(-4*B-4*I*A)/(I+tan(f*x+e))^6-1
/4*(I*A+5*B)/(I+tan(f*x+e))^4-1/5*(-8*I*B+4*A)/(I+tan(f*x+e))^5)
```

3.702.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.70

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx =$$

$$-\frac{10(iA + B)a^3 e^{(12i fx + 12i e)} + 12(3iA + B)a^3 e^{(10i fx + 10i e)} + 15(3iA - B)a^3 e^{(8i fx + 8i e)} + 20(iA - B)a^3 e^{(6i fx + 6i e)}}{960 c^6 f}$$

```
input integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^6,x, al
gorithm="fracas")
```

```
output -1/960*(10*(I*A + B)*a^3*e^(12*I*f*x + 12*I*e) + 12*(3*I*A + B)*a^3*e^(10*
I*f*x + 10*I*e) + 15*(3*I*A - B)*a^3*e^(8*I*f*x + 8*I*e) + 20*(I*A - B)*a^
3*e^(6*I*f*x + 6*I*e))/(c^6*f)
```

3.702.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 332 vs. $2(105) = 210$.

Time = 0.53 (sec) , antiderivative size = 332, normalized size of antiderivative = 2.61

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx$$

$$= \left\{ \frac{(-491520iAa^3c^{18}f^3e^{6ie} + 491520Ba^3c^{18}f^3e^{6ie})e^{6ifx} + (-1105920iAa^3c^{18}f^3e^{8ie} + 368640Ba^3c^{18}f^3e^{8ie})e^{8ifx} + (-884736iAa^3c^{18}f^3e^{10ie} - 294912Ba^3c^{18}f^3e^{10ie})e^{10ifx} + (-245760iAa^3c^{18}f^3e^{12ie} - 245760Ba^3c^{18}f^3e^{12ie})e^{12ifx}}{23592960c^{24}f^4} \right.$$

$$\left. \frac{x(Aa^3e^{12ie} + 3Aa^3e^{10ie} + 3Aa^3e^{8ie} + Aa^3e^{6ie} - iBa^3e^{12ie} - iBa^3e^{10ie} + iBa^3e^{8ie} + iBa^3e^{6ie})}{8c^6} \right\}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**6,x)`

output `Piecewise(((((-491520*I*A*a**3*c**18*f**3*exp(6*I*e) + 491520*B*a**3*c**18*f**3*exp(6*I*e))*exp(6*I*f*x) + (-1105920*I*A*a**3*c**18*f**3*exp(8*I*e) + 368640*B*a**3*c**18*f**3*exp(8*I*e))*exp(8*I*f*x) + (-884736*I*A*a**3*c**18*f**3*exp(10*I*e) - 294912*B*a**3*c**18*f**3*exp(10*I*e))*exp(10*I*f*x) + (-245760*I*A*a**3*c**18*f**3*exp(12*I*e) - 245760*B*a**3*c**18*f**3*exp(12*I*e))*exp(12*I*f*x))/(23592960*c**24*f**4), Ne(c**24*f**4, 0)), (x*(A*a**3*exp(12*I*e) + 3*A*a**3*exp(10*I*e) + 3*A*a**3*exp(8*I*e) + A*a**3*exp(6*I*e) - I*B*a**3*exp(12*I*e) - I*B*a**3*exp(10*I*e) + I*B*a**3*exp(8*I*e) + I*B*a**3*exp(6*I*e))/(8*c**6), True))`

3.702.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^6,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.702.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 326 vs. $2(103) = 206$.

Time = 1.58 (sec) , antiderivative size = 326, normalized size of antiderivative = 2.57

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^6} dx =$$

$$\frac{2 \left(15 A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^{11} + 45 i A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^{10} - 15 B a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^{10} - 215 A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^9 - 390 i A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^8 + 90 B a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^8 + 738 A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^7 + 24 i B a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^7 + 746 i A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^6 - 158 B a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^6 - 738 A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^5 - 24 i B a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^5 - 390 i A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^4 + 90 B a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^4 + 215 A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^3 + 45 i A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^2 - 15 B a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^2 - 15 A a^3 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) \right) / (c^6 f (\tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) + i)^{12}}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^6,x, algorithm="giac")`

output `-2/15*(15*A*a^3*tan(1/2*f*x + 1/2*e)^11 + 45*I*A*a^3*tan(1/2*f*x + 1/2*e)^10 - 15*B*a^3*tan(1/2*f*x + 1/2*e)^10 - 215*A*a^3*tan(1/2*f*x + 1/2*e)^9 - 390*I*A*a^3*tan(1/2*f*x + 1/2*e)^8 + 90*B*a^3*tan(1/2*f*x + 1/2*e)^8 + 738*A*a^3*tan(1/2*f*x + 1/2*e)^7 + 24*I*B*a^3*tan(1/2*f*x + 1/2*e)^7 + 746*I*A*a^3*tan(1/2*f*x + 1/2*e)^6 - 158*B*a^3*tan(1/2*f*x + 1/2*e)^6 - 738*A*a^3*tan(1/2*f*x + 1/2*e)^5 - 24*I*B*a^3*tan(1/2*f*x + 1/2*e)^5 - 390*I*A*a^3*tan(1/2*f*x + 1/2*e)^4 + 90*B*a^3*tan(1/2*f*x + 1/2*e)^4 + 215*A*a^3*tan(1/2*f*x + 1/2*e)^3 + 45*I*A*a^3*tan(1/2*f*x + 1/2*e)^2 - 15*B*a^3*tan(1/2*f*x + 1/2*e)^2 - 15*A*a^3*tan(1/2*f*x + 1/2*e))/(c^6*f*(tan(1/2*f*x + 1/2*e) + I)^12)`

3.702.9 Mupad [B] (verification not implemented)

Time = 8.71 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.10

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^6} dx =$$

$$-\frac{a^3 (-B + A 7i)}{60} + \frac{a^3 \tan(e + fx) (18 A - B 6i)}{60} + \frac{B a^3 \tan(e + fx)^3 i}{3} + \frac{a^3 \tan(e + fx)^2 (15 B + A 15i)}{60}$$

$$c^6 f (\tan(e + fx)^6 + \tan(e + fx)^5 6i - 15 \tan(e + fx)^4 - \tan(e + fx)^3 20i + 15 \tan(e + fx)^2 + \tan(e + fx) + i)$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3)/(c - c*tan(e + f*x)*1i)^6,x)`

output $-\frac{(a^3 \tan(e + fx)(18A - B6i))}{60} - \frac{(a^3(A7i - B))}{60} + \frac{(B a^3 \tan(e + fx)^3 1i)}{3} + \frac{(a^3 \tan(e + fx)^2 (A15i + 15B))}{60} / (c^6 f (\tan(e + fx)^6 i + 15 \tan(e + fx)^2 - \tan(e + fx)^3 20i - 15 \tan(e + fx)^4 + \tan(e + fx)^5 6i + \tan(e + fx)^6 - 1))$

3.702. $\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^6} dx$

3.703
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^7} dx$$

3.703.1 Optimal result 6543
 3.703.2 Mathematica [A] (verified) 6543
 3.703.3 Rubi [A] (verified) 6544
 3.703.4 Maple [A] (verified) 6546
 3.703.5 Fricas [A] (verification not implemented) 6546
 3.703.6 Sympy [B] (verification not implemented) 6547
 3.703.7 Maxima [F(-2)] 6547
 3.703.8 Giac [B] (verification not implemented) 6548
 3.703.9 Mupad [B] (verification not implemented) 6548

3.703.1 Optimal result

Integrand size = 41, antiderivative size = 125

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^7} dx \\ &= -\frac{4a^3(A - iB)}{7c^7 f(i + \tan(e + fx))^7} - \frac{2a^3(iA + 2B)}{3c^7 f(i + \tan(e + fx))^6} \\ & \quad + \frac{a^3(A - 5iB)}{5c^7 f(i + \tan(e + fx))^5} + \frac{a^3 B}{4c^7 f(i + \tan(e + fx))^4} \end{aligned}$$

output
$$-4/7*a^3*(A-I*B)/c^7/f/(I+\tan(f*x+e))^7-2/3*a^3*(I*A+2*B)/c^7/f/(I+\tan(f*x+e))^6+1/5*a^3*(A-5*I*B)/c^7/f/(I+\tan(f*x+e))^5+1/4*a^3*B/c^7/f/(I+\tan(f*x+e))^4$$

3.703.2 Mathematica [A] (verified)

Time = 5.53 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.64

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^7} dx \\ &= \frac{a^3(-44A - 5iB + (-112iA - 35B) \tan(e + fx) + 21(4A - 5iB) \tan^2(e + fx) + 105B \tan^3(e + fx))}{420c^7 f(i + \tan(e + fx))^7} \end{aligned}$$

input
$$\text{Integrate}[\frac{(a + I*a*\text{Tan}[e + f*x])^3*(A + B*\text{Tan}[e + f*x])}{(c - I*c*\text{Tan}[e + f*x])^7}, x]$$

output $(a^3*(-44*A - (5*I)*B + ((-112*I)*A - 35*B)*\text{Tan}[e + f*x] + 21*(4*A - (5*I)*B)*\text{Tan}[e + f*x]^2 + 105*B*\text{Tan}[e + f*x]^3))/(420*c^7*f*(I + \text{Tan}[e + f*x])^7)$

3.703.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.79, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^7} dx \\ & \quad \downarrow 3042 \\ & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^7} dx \\ & \quad \downarrow 4071 \\ & \frac{ac \int \frac{a^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{c^8 (1 - i \tan(e + fx))^8} d \tan(e + fx)}{f} \\ & \quad \downarrow 27 \\ & \frac{a^3 \int \frac{(i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{(1 - i \tan(e + fx))^8} d \tan(e + fx)}{c^7 f} \\ & \quad \downarrow 86 \\ & \frac{a^3 \int \left(\frac{4(A - iB)}{(\tan(e + fx) + i)^8} - \frac{B}{(\tan(e + fx) + i)^5} + \frac{5iB - A}{(\tan(e + fx) + i)^6} + \frac{4(iA + 2B)}{(\tan(e + fx) + i)^7} \right) d \tan(e + fx)}{c^7 f} \\ & \quad \downarrow 2009 \\ & \frac{a^3 \left(-\frac{4(A - iB)}{7(\tan(e + fx) + i)^7} + \frac{A - 5iB}{5(\tan(e + fx) + i)^5} - \frac{2(2B + iA)}{3(\tan(e + fx) + i)^6} + \frac{B}{4(\tan(e + fx) + i)^4} \right)}{c^7 f} \end{aligned}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^3*(A + B*\text{Tan}[e + f*x])]/(c - I*c*\text{Tan}[e + f*x])^7, x]$

3.703. $\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^7} dx$

```
output (a^3*((-4*(A - I*B))/(7*(I + Tan[e + f*x])^7) - (2*(I*A + 2*B))/(3*(I + Tan[e + f*x])^6) + (A - (5*I)*B)/(5*(I + Tan[e + f*x])^5) + B/(4*(I + Tan[e + f*x])^4))/(c^7*f)
```

3.703.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]
```


3.703.4 Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.71

method	result
derivativedivides	$\frac{a^3 \left(\frac{B}{4(i+\tan(fx+e))^4} - \frac{5iB-A}{5(i+\tan(fx+e))^5} - \frac{4iA+8B}{6(i+\tan(fx+e))^6} - \frac{-4iB+4A}{7(i+\tan(fx+e))^7} \right)}{f c^7}$
default	$\frac{a^3 \left(\frac{B}{4(i+\tan(fx+e))^4} - \frac{5iB-A}{5(i+\tan(fx+e))^5} - \frac{4iA+8B}{6(i+\tan(fx+e))^6} - \frac{-4iB+4A}{7(i+\tan(fx+e))^7} \right)}{f c^7}$
risch	$-\frac{a^3 e^{14i(fx+e)} B}{224c^7 f} - \frac{ia^3 e^{14i(fx+e)} A}{224c^7 f} - \frac{e^{12i(fx+e)} B a^3}{96c^7 f} - \frac{ie^{12i(fx+e)} a^3 A}{48c^7 f} - \frac{3ia^3 A e^{10i(fx+e)}}{80c^7 f} + \frac{e^{8i(fx+e)} B a^3}{64c^7 f}$
norman	$\frac{a^3 A \tan(fx+e)}{fc} + \frac{-44iA a^3 + 5B a^3}{420cf} + \frac{B a^3 \tan(fx+e)^{10}}{4cf} + \frac{2(-125iB a^3 + 139a^3 A) \tan(fx+e)^5}{15cf} - \frac{6(-85iB a^3 + 36a^3 A) \tan(fx+e)^7}{35cf}$

input `int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^7,x,method=_R
ETURNVERBOSE)`

output `1/f*a^3/c^7*(1/4*B/(I+tan(f*x+e))^4-1/5*(-A+5*I*B)/(I+tan(f*x+e))^5-1/6*(4
*I*A+8*B)/(I+tan(f*x+e))^6-1/7*(-4*I*B+4*A)/(I+tan(f*x+e))^7)`

3.703.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.83

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^7} dx =$$

$$-\frac{30(iA + B)a^3 e^{(14i fx + 14ie)} + 70(2iA + B)a^3 e^{(12i fx + 12ie)} + 252iAa^3 e^{(10i fx + 10ie)} + 105(2iA - B)a^3 e^{(8i fx + 8ie)}}{6720 c^7 f}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^7,x, al
gorithm="fracas")`

output `-1/6720*(30*(I*A + B)*a^3*e^(14*I*f*x + 14*I*e) + 70*(2*I*A + B)*a^3*e^(12
*I*f*x + 12*I*e) + 252*I*A*a^3*e^(10*I*f*x + 10*I*e) + 105*(2*I*A - B)*a^3
*e^(8*I*f*x + 8*I*e) + 70*(I*A - B)*a^3*e^(6*I*f*x + 6*I*e))/(c^7*f)`

3.703.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 379 vs. $2(104) = 208$.

Time = 0.66 (sec) , antiderivative size = 379, normalized size of antiderivative = 3.03

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^7} dx$$

$$= \left\{ \begin{array}{l} \frac{-396361728iAa^3c^{28}f^4e^{10ie}e^{10ifx} + (-110100480iAa^3c^{28}f^4e^{6ie} + 110100480Ba^3c^{28}f^4e^{6ie})e^{6ifx} + (-330301440iAa^3c^{28}f^4e^{8ie} + 165150720B \\ x(Aa^3e^{14ie} + 4Aa^3e^{12ie} + 6Aa^3e^{10ie} + 4Aa^3e^{8ie} + Aa^3e^{6ie} - iBa^3e^{14ie} - 2iBa^3e^{12ie} + 2iBa^3e^{8ie} + iBa^3e^{6ie})}{16c^7} \end{array} \right.$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**7,x)`

output `Piecewise(((-396361728*I*A*a**3*c**28*f**4*exp(10*I*e)*exp(10*I*f*x) + (-110100480*I*A*a**3*c**28*f**4*exp(6*I*e) + 110100480*B*a**3*c**28*f**4*exp(6*I*e))*exp(6*I*f*x) + (-330301440*I*A*a**3*c**28*f**4*exp(8*I*e) + 165150720*B*a**3*c**28*f**4*exp(8*I*e))*exp(8*I*f*x) + (-220200960*I*A*a**3*c**28*f**4*exp(12*I*e) - 110100480*B*a**3*c**28*f**4*exp(12*I*e))*exp(12*I*f*x) + (-47185920*I*A*a**3*c**28*f**4*exp(14*I*e) - 47185920*B*a**3*c**28*f**4*exp(14*I*e))*exp(14*I*f*x))/(10569646080*c**35*f**5), Ne(c**35*f**5, 0)), (x*(A*a**3*exp(14*I*e) + 4*A*a**3*exp(12*I*e) + 6*A*a**3*exp(10*I*e) + 4*A*a**3*exp(8*I*e) + A*a**3*exp(6*I*e) - I*B*a**3*exp(14*I*e) - 2*I*B*a**3*exp(12*I*e) + 2*I*B*a**3*exp(8*I*e) + I*B*a**3*exp(6*I*e))/(16*c**7), True))`

3.703.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^7} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^7,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.703. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^7} dx$

3.703.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 428 vs. $2(103) = 206$.

Time = 1.07 (sec) , antiderivative size = 428, normalized size of antiderivative = 3.42

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^7} dx =$$

$$\frac{2 \left(105 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{13} + 420i A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{12} - 105 B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{12} - 2170 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{11} - 70i B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{11} - 5180i A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{10} + 875 B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{10} + 11431 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^9 + 700i B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^9 + 15904 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^8 - 2380 B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^8 - 19436 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^7 - 1340i B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^7 - 15904 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^6 + 2380 B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^6 + 11431 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^5 + 700i B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^5 + 5180 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^4 - 875 B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^4 - 2170 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^3 - 70i B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^3 - 420 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 + 105 B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 + 105 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right) \right)}{(c^7 f (\tan\left(\frac{1}{2} fx + \frac{1}{2} e\right) + i)^{14}}$$

```
input integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^7,x, algorithm="giac")
```

```
output -2/105*(105*A*a^3*tan(1/2*f*x + 1/2*e)^13 + 420*I*A*a^3*tan(1/2*f*x + 1/2*e)^12 - 105*B*a^3*tan(1/2*f*x + 1/2*e)^12 - 2170*A*a^3*tan(1/2*f*x + 1/2*e)^11 - 70*I*B*a^3*tan(1/2*f*x + 1/2*e)^11 - 5180*I*A*a^3*tan(1/2*f*x + 1/2*e)^10 + 875*B*a^3*tan(1/2*f*x + 1/2*e)^10 + 11431*A*a^3*tan(1/2*f*x + 1/2*e)^9 + 700*I*B*a^3*tan(1/2*f*x + 1/2*e)^9 + 15904*A*a^3*tan(1/2*f*x + 1/2*e)^8 - 2380*B*a^3*tan(1/2*f*x + 1/2*e)^8 - 19436*A*a^3*tan(1/2*f*x + 1/2*e)^7 - 1340*I*B*a^3*tan(1/2*f*x + 1/2*e)^7 - 15904*A*a^3*tan(1/2*f*x + 1/2*e)^6 + 2380*B*a^3*tan(1/2*f*x + 1/2*e)^6 + 11431*A*a^3*tan(1/2*f*x + 1/2*e)^5 + 700*I*B*a^3*tan(1/2*f*x + 1/2*e)^5 + 5180*A*a^3*tan(1/2*f*x + 1/2*e)^4 - 875*B*a^3*tan(1/2*f*x + 1/2*e)^4 - 2170*A*a^3*tan(1/2*f*x + 1/2*e)^3 - 70*I*B*a^3*tan(1/2*f*x + 1/2*e)^3 - 420*A*a^3*tan(1/2*f*x + 1/2*e)^2 + 105*B*a^3*tan(1/2*f*x + 1/2*e)^2 + 105*A*a^3*tan(1/2*f*x + 1/2*e))/(c^7*f*(tan(1/2*f*x + 1/2*e) + I)^14)
```

3.703.9 Mupad [B] (verification not implemented)

Time = 8.94 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.21

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^7} dx$$

$$= \frac{a^3 (44 A + B 5i)}{420} + \frac{a^3 \tan(e + fx) (35 B + A 112i)}{420} - \frac{B a^3 \tan(e + fx)^3}{4} - \frac{a^3 \tan(e + fx)^2 (84 A - 112 B i)}{420} - \frac{a^3 \tan(e + fx)}{420} - \frac{a^3}{420} - \frac{a^3 \tan(e + fx)^3}{4} - \frac{a^3 \tan(e + fx)^2 (84 A - 112 B i)}{420} - \frac{a^3 \tan(e + fx)}{420} - \frac{a^3}{420}$$

```
input int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3)/(c - c*tan(e + f*x)*1i)^7,x)
```

output $((a^3(44A + B5i))/420 + (a^3 \tan(e + f*x)(A*112i + 35*B))/420 - (B*a^3 \tan(e + f*x)^3)/4 - (a^3 \tan(e + f*x)^2(84*A - B*105i))/420)/(c^7*f*(7*\tan(e + f*x) - \tan(e + f*x)^2*21i - 35*\tan(e + f*x)^3 + \tan(e + f*x)^4*35i + 21*\tan(e + f*x)^5 - \tan(e + f*x)^6*7i - \tan(e + f*x)^7 + 1i))$

3.703. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^7} dx$

3.704
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^8} dx$$

3.704.1 Optimal result 6550
 3.704.2 Mathematica [A] (verified) 6550
 3.704.3 Rubi [A] (verified) 6551
 3.704.4 Maple [A] (verified) 6553
 3.704.5 Fricas [A] (verification not implemented) 6553
 3.704.6 Sympy [B] (verification not implemented) 6554
 3.704.7 Maxima [F(-2)] 6554
 3.704.8 Giac [B] (verification not implemented) 6555
 3.704.9 Mupad [B] (verification not implemented) 6556

3.704.1 Optimal result

Integrand size = 41, antiderivative size = 127

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^8} dx \\ &= -\frac{a^3(iA + B)}{2c^8 f(i + \tan(e + fx))^8} + \frac{4a^3(A - 2iB)}{7c^8 f(i + \tan(e + fx))^7} \\ &+ \frac{a^3(iA + 5B)}{6c^8 f(i + \tan(e + fx))^6} + \frac{ia^3 B}{5c^8 f(i + \tan(e + fx))^5} \end{aligned}$$

output `-1/2*a^3*(I*A+B)/c^8/f/(I+tan(f*x+e))^8+4/7*a^3*(A-2*I*B)/c^8/f/(I+tan(f*x+e))^7+1/6*a^3*(I*A+5*B)/c^8/f/(I+tan(f*x+e))^6+1/5*I*a^3*B/c^8/f/(I+tan(f*x+e))^5`

3.704.2 Mathematica [A] (verified)

Time = 5.55 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.66

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^8} dx \\ &= \frac{a^3(2(-10iA + B) + 2(25A - 8iB) \tan(e + fx) + 7(5iA + 7B) \tan^2(e + fx) + 42iB \tan^3(e + fx))}{210c^8 f(i + \tan(e + fx))^8} \end{aligned}$$

input `Integrate[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^8,x]`

3.704.
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^8} dx$$

output $(a^3(2*((-10*I)*A + B) + 2*(25*A - (8*I)*B)*\text{Tan}[e + f*x] + 7*((5*I)*A + 7*B)*\text{Tan}[e + f*x]^2 + (42*I)*B*\text{Tan}[e + f*x]^3)/(210*c^8*f*(I + \text{Tan}[e + f*x])^8)$

3.704.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.80, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^8} dx \\ & \quad \downarrow 3042 \\ & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^8} dx \\ & \quad \downarrow 4071 \\ & \frac{ac \int \frac{a^2(i \tan(e+fx)+1)^2(A+B \tan(e+fx))}{c^9(1-i \tan(e+fx))^9} d \tan(e + fx)}{f} \\ & \quad \downarrow 27 \\ & \frac{a^3 \int \frac{(i \tan(e+fx)+1)^2(A+B \tan(e+fx))}{(1-i \tan(e+fx))^9} d \tan(e + fx)}{c^8 f} \\ & \quad \downarrow 86 \\ & \frac{a^3 \int \left(-\frac{4(A-2iB)}{(\tan(e+fx)+i)^8} - \frac{iB}{(\tan(e+fx)+i)^6} - \frac{i(A-5iB)}{(\tan(e+fx)+i)^7} + \frac{4(iA+B)}{(\tan(e+fx)+i)^9} \right) d \tan(e + fx)}{c^8 f} \\ & \quad \downarrow 2009 \\ & \frac{a^3 \left(\frac{4(A-2iB)}{7(\tan(e+fx)+i)^7} + \frac{5B+iA}{6(\tan(e+fx)+i)^6} - \frac{B+iA}{2(\tan(e+fx)+i)^8} + \frac{iB}{5(\tan(e+fx)+i)^5} \right)}{c^8 f} \end{aligned}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^3*(A + B*\text{Tan}[e + f*x])]/(c - I*c*\text{Tan}[e + f*x])^8,x]$

```
output (a^3*(-1/2*(I*A + B)/(I + Tan[e + f*x])^8 + (4*(A - (2*I)*B))/(7*(I + Tan[
e + f*x])^7) + (I*A + 5*B)/(6*(I + Tan[e + f*x])^6) + ((I/5)*B)/(I + Tan[e
+ f*x])^5))/(c^8*f)
```

3.704.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.704.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.71

method	result
derivativedivides	$\frac{a^3 \left(-\frac{8iB-4A}{7(i+\tan(fx+e))^7} - \frac{-iA-5B}{6(i+\tan(fx+e))^6} + \frac{iB}{5(i+\tan(fx+e))^5} - \frac{4iA+4B}{8(i+\tan(fx+e))^8} \right)}{f c^8}$
default	$\frac{a^3 \left(-\frac{8iB-4A}{7(i+\tan(fx+e))^7} - \frac{-iA-5B}{6(i+\tan(fx+e))^6} + \frac{iB}{5(i+\tan(fx+e))^5} - \frac{4iA+4B}{8(i+\tan(fx+e))^8} \right)}{f c^8}$
risch	$-\frac{a^3 e^{16i(fx+e)} B}{512c^8 f} - \frac{ia^3 e^{16i(fx+e)} A}{512c^8 f} - \frac{3e^{14i(fx+e)} B a^3}{448c^8 f} - \frac{5ie^{14i(fx+e)} a^3 A}{448c^8 f} - \frac{e^{12i(fx+e)} B a^3}{192c^8 f} - \frac{5ie^{12i(fx+e)} a^3}{192c^8 f}$

```
input int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^8,x,method=_R
ETURNVERBOSE)
```

```
output 1/f*a^3/c^8*(-1/7*(8*I*B-4*A)/(I+tan(f*x+e))^7-1/6*(-I*A-5*B)/(I+tan(f*x+e)
))^6+1/5*I*B/(I+tan(f*x+e))^5-1/8*(4*I*A+4*B)/(I+tan(f*x+e))^8)
```

3.704.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.03

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^8} dx =$$

$$-\frac{105(iA + B)a^3 e^{(16i fx + 16ie)} + 120(5iA + 3B)a^3 e^{(14i fx + 14ie)} + 280(5iA + B)a^3 e^{(12i fx + 12ie)} + 336(5iA - B)a^3 e^{(10i fx + 10ie)} + 210(5iA - 3B)a^3 e^{(8i fx + 8ie)} + 280(IA - B)a^3 e^{(6i fx + 6ie)}}{53760 c^8 f}$$

```
input integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^8,x, al
gorithm="fracas")
```

```
output -1/53760*(105*(I*A + B)*a^3*e^(16*I*f*x + 16*I*e) + 120*(5*I*A + 3*B)*a^3*
e^(14*I*f*x + 14*I*e) + 280*(5*I*A + B)*a^3*e^(12*I*f*x + 12*I*e) + 336*(5
*I*A - B)*a^3*e^(10*I*f*x + 10*I*e) + 210*(5*I*A - 3*B)*a^3*e^(8*I*f*x + 8
*I*e) + 280*(I*A - B)*a^3*e^(6*I*f*x + 6*I*e))/(c^8*f)
```

3.704. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^8} dx$

3.704.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 496 vs. $2(104) = 208$.

Time = 0.72 (sec) , antiderivative size = 496, normalized size of antiderivative = 3.91

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^8} dx$$

$$= \left\{ \frac{(-1803886264320iAa^3c^{40}f^5e^{6ie} + 1803886264320Ba^3c^{40}f^5e^{6ie})e^{6ifx} + (-6764573491200iAa^3c^{40}f^5e^{8ie} + 4058744094720Ba^3c^{40}f^5e^{8ie})e^{8ifx} + (-10823317585920iAa^3c^{40}f^5e^{10ie} + 2164663517184Ba^3c^{40}f^5e^{10ie})e^{10ifx} + (-9019431321600iAa^3c^{40}f^5e^{12ie} - 1803886264320Ba^3c^{40}f^5e^{12ie})e^{12ifx} + (-3865470566400iAa^3c^{40}f^5e^{14ie} - 2319282339840Ba^3c^{40}f^5e^{14ie})e^{14ifx} + (-6764573491200iAa^3c^{40}f^5e^{16ie} - 6764573491200Ba^3c^{40}f^5e^{16ie})e^{16ifx}}{32c^8} \right.$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**8,x)`

output `Piecewise(((((-1803886264320*I*A*a**3*c**40*f**5*exp(6*I*e) + 1803886264320*B*a**3*c**40*f**5*exp(6*I*e))*exp(6*I*f*x) + (-6764573491200*I*A*a**3*c**40*f**5*exp(8*I*e) + 4058744094720*B*a**3*c**40*f**5*exp(8*I*e))*exp(8*I*f*x) + (-10823317585920*I*A*a**3*c**40*f**5*exp(10*I*e) + 2164663517184*B*a**3*c**40*f**5*exp(10*I*e))*exp(10*I*f*x) + (-9019431321600*I*A*a**3*c**40*f**5*exp(12*I*e) - 1803886264320*B*a**3*c**40*f**5*exp(12*I*e))*exp(12*I*f*x) + (-3865470566400*I*A*a**3*c**40*f**5*exp(14*I*e) - 2319282339840*B*a**3*c**40*f**5*exp(14*I*e))*exp(14*I*f*x) + (-6764573491200*I*A*a**3*c**40*f**5*exp(16*I*e) - 6764573491200*B*a**3*c**40*f**5*exp(16*I*e))*exp(16*I*f*x))/(346346162749440*c**48*f**6), Ne(c**48*f**6, 0)), (x*(A*a**3*exp(16*I*e) + 5*A*a**3*exp(14*I*e) + 10*A*a**3*exp(12*I*e) + 10*A*a**3*exp(10*I*e) + 5*A*a**3*exp(8*I*e) + A*a**3*exp(6*I*e) - I*B*a**3*exp(16*I*e) - 3*I*B*a**3*exp(14*I*e) - 2*I*B*a**3*exp(12*I*e) + 2*I*B*a**3*exp(10*I*e) + 3*I*B*a**3*exp(8*I*e) + I*B*a**3*exp(6*I*e))/(32*c**8), True))`

3.704.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^8} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^8,x, algorithm="maxima")`

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

3.704.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 496 vs. $2(103) = 206$.

Time = 1.06 (sec) , antiderivative size = 496, normalized size of antiderivative = 3.91

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^8} dx =$$

$$\frac{2 \left(105 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{15} + 525i A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{14} - 105 B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{14} - 2975 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{13} - 140i B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{13} - 8750i A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{12} + 1190 B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{12} + 22365 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{11} + 1596i B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{11} + 39235i A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{10} - 4711 B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^{10} - 58075 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^9 - 4600i B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^9 - 63300i A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^8 + 7380 B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^8 + 58075 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^7 + 4600i B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^7 + 39235i A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^6 - 4711 B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^6 - 22365 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^5 - 1596i B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^5 - 8750i A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^4 + 1190 B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^4 + 2975 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^3 + 140i B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^3 + 525i A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - 105 B a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right)^2 - 105 A a^3 \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right) \right)}{(c^8 f \tan\left(\frac{1}{2} fx + \frac{1}{2} e\right) + I)^{16}}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^8,x, algorithm="giac")`

output

$$\frac{-2/105*(105*A*a^3*\tan(1/2*f*x + 1/2*e)^{15} + 525*I*A*a^3*\tan(1/2*f*x + 1/2*e)^{14} - 105*B*a^3*\tan(1/2*f*x + 1/2*e)^{14} - 2975*A*a^3*\tan(1/2*f*x + 1/2*e)^{13} - 140*I*B*a^3*\tan(1/2*f*x + 1/2*e)^{13} - 8750*I*A*a^3*\tan(1/2*f*x + 1/2*e)^{12} + 1190*B*a^3*\tan(1/2*f*x + 1/2*e)^{12} + 22365*A*a^3*\tan(1/2*f*x + 1/2*e)^{11} + 1596*I*B*a^3*\tan(1/2*f*x + 1/2*e)^{11} + 39235*I*A*a^3*\tan(1/2*f*x + 1/2*e)^{10} - 4711*B*a^3*\tan(1/2*f*x + 1/2*e)^{10} - 58075*A*a^3*\tan(1/2*f*x + 1/2*e)^9 - 4600*I*B*a^3*\tan(1/2*f*x + 1/2*e)^9 - 63300*I*A*a^3*\tan(1/2*f*x + 1/2*e)^8 + 7380*B*a^3*\tan(1/2*f*x + 1/2*e)^8 + 58075*A*a^3*\tan(1/2*f*x + 1/2*e)^7 + 4600*I*B*a^3*\tan(1/2*f*x + 1/2*e)^7 + 39235*I*A*a^3*\tan(1/2*f*x + 1/2*e)^6 - 4711*B*a^3*\tan(1/2*f*x + 1/2*e)^6 - 22365*A*a^3*\tan(1/2*f*x + 1/2*e)^5 - 1596*I*B*a^3*\tan(1/2*f*x + 1/2*e)^5 - 8750*I*A*a^3*\tan(1/2*f*x + 1/2*e)^4 + 1190*B*a^3*\tan(1/2*f*x + 1/2*e)^4 + 2975*A*a^3*\tan(1/2*f*x + 1/2*e)^3 + 140*I*B*a^3*\tan(1/2*f*x + 1/2*e)^3 + 525*I*A*a^3*\tan(1/2*f*x + 1/2*e)^2 - 105*B*a^3*\tan(1/2*f*x + 1/2*e)^2 - 105*A*a^3*\tan(1/2*f*x + 1/2*e))/(c^8*f*(\tan(1/2*f*x + 1/2*e) + I)^{16}}$$

3.704.9 Mupad [B] (verification not implemented)

Time = 9.22 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.26

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^8} dx$$

$$= \frac{-\frac{a^3(-2B + A20i)}{210} + \frac{a^3 \tan(e + fx)(50A - B16i)}{210} + \frac{B a^3 \tan(e + fx)^3 1i}{5} + \frac{a^3 \tan(e + fx)^5 56i}{5} + \frac{a^3 \tan(e + fx)^7 8i}{5} + \frac{a^3 \tan(e + fx)^9 1i}{5}}{c^8 f (\tan(e + fx)^8 + \tan(e + fx)^7 8i - 28 \tan(e + fx)^6 - \tan(e + fx)^5 56i + 70 \tan(e + fx)^4 + \tan(e + fx)^3 56i - 28 \tan(e + fx)^2 - \tan(e + fx) 8i + 70 \tan(e + fx) + 1)}$$

```
input int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3)/(c - c*tan(e + f*x)*1i)^8,x)
```

```
output ((a^3*tan(e + f*x)*(50*A - B*16i))/210 - (a^3*(A*20i - 2*B))/210 + (B*a^3*tan(e + f*x)^3*1i)/5 + (a^3*tan(e + f*x)^2*(A*35i + 49*B))/210)/(c^8*f*(tan(e + f*x)^3*56i - 28*tan(e + f*x)^2 - tan(e + f*x)*8i + 70*tan(e + f*x)^4 - tan(e + f*x)^5*56i - 28*tan(e + f*x)^6 + tan(e + f*x)^7*8i + tan(e + f*x)^8 + 1))
```

$$3.705 \quad \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^n}{a+ia \tan(e+fx)} dx$$

3.705.1 Optimal result	6557
3.705.2 Mathematica [A] (verified)	6557
3.705.3 Rubi [A] (verified)	6558
3.705.4 Maple [F]	6560
3.705.5 Fricas [F]	6560
3.705.6 Sympy [F]	6560
3.705.7 Maxima [F(-2)]	6561
3.705.8 Giac [F]	6561
3.705.9 Mupad [F(-1)]	6561

3.705.1 Optimal result

Integrand size = 41, antiderivative size = 115

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{a + ia \tan(e + fx)} dx$$

$$= \frac{(iA(1 - n) + B(1 + n)) \operatorname{Hypergeometric2F1}\left(1, n, 1 + n, \frac{1}{2}(1 - i \tan(e + fx))\right) (c - ic \tan(e + fx))^n}{4afn}$$

$$+ \frac{(iA - B)(c - ic \tan(e + fx))^n}{2af(1 + i \tan(e + fx))}$$

output `1/4*(I*A*(1-n)+B*(1+n))*hypergeom([1, n],[1+n],1/2-1/2*I*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/a/f/n+1/2*(I*A-B)*(c-I*c*tan(f*x+e))^n/a/f/(1+I*tan(f*x+e))`

3.705.2 Mathematica [A] (verified)

Time = 1.42 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.85

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{a + ia \tan(e + fx)} dx$$

$$= \frac{(c - ic \tan(e + fx))^n (2(A + iB)n + (-iA(-1 + n) + B(1 + n))) \operatorname{Hypergeometric2F1}\left(1, n, 1 + n, -\frac{1}{2}i(i \tan(e + fx) + 1)\right)}{4afn(-i + \tan(e + fx))}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^n)/(a + I*a*Tan[e + f*x]),x]`

output $((c - I*c*\text{Tan}[e + f*x])^n*(2*(A + I*B)*n + ((-I)*A*(-1 + n) + B*(1 + n))*\text{Hypergeometric2F1}[1, n, 1 + n, (-1/2*I)*(I + \text{Tan}[e + f*x])]*(-I + \text{Tan}[e + f*x])))/(4*a*f*n*(-I + \text{Tan}[e + f*x]))$

3.705.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 87, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{a + ia \tan(e + fx)} dx$$

↓ 3042

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{a + ia \tan(e + fx)} dx$$

↓ 4071

$$\frac{ac \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{n-1}}{a^2(i \tan(e + fx) + 1)^2} d \tan(e + fx)}{f}$$

↓ 27

$$\frac{c \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{n-1}}{(i \tan(e + fx) + 1)^2} d \tan(e + fx)}{af}$$

↓ 87

$$\frac{c \left(\frac{1}{2} (A(1 - n) - iB(n + 1)) \int \frac{(c - ic \tan(e + fx))^{n-1}}{i \tan(e + fx) + 1} d \tan(e + fx) + \frac{(-B + iA)(c - ic \tan(e + fx))^n}{2c(1 + i \tan(e + fx))} \right)}{af}$$

↓ 78

$$\frac{c \left(\frac{(i(A(1 - n) - iB(n + 1))(c - ic \tan(e + fx))^n \text{Hypergeometric2F1}(1, n, n + 1, \frac{1}{2}(1 - i \tan(e + fx)))}{4cn} + \frac{(-B + iA)(c - ic \tan(e + fx))^n}{2c(1 + i \tan(e + fx))} \right)}{af}$$

input $\text{Int}[(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^n/(a + I*a*\text{Tan}[e + f*x]), x]$

3.705. $\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{a + ia \tan(e + fx)} dx$

```
output (c*((I/4)*(A*(1 - n) - I*B*(1 + n))*Hypergeometric2F1[1, n, 1 + n, (1 - I
*Tan[e + f*x])/2]*(c - I*c*Tan[e + f*x])^n)/(c*n) + ((I*A - B)*(c - I*c*Ta
n[e + f*x])^n)/(2*c*(1 + I*Tan[e + f*x])))/(a*f)
```

3.705.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 78 Int[((a_) + (b_.)*(x_)^(m_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(b
*c - a*d)^n*((a + b*x)^(m + 1)/(b^(n + 1)*(m + 1)))*Hypergeometric2F1[-n, m
+ 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x]
&& !IntegerQ[m] && IntegerQ[n]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p
_.)), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.))*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.705.4 Maple [F]

$$\int \frac{(A + B \tan(fx + e))(c - ic \tan(fx + e))^n}{a + ia \tan(fx + e)} dx$$

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e)),x)`

output `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e)),x)`

3.705.5 Fricas [F]

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{a + ia \tan(e + fx)} dx \\ &= \int \frac{(B \tan(fx + e) + A)(-ic \tan(fx + e) + c)^n}{ia \tan(fx + e) + a} dx \end{aligned}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`

output `integral(1/2*((A - I*B)*e^(2*I*f*x + 2*I*e) + A + I*B)*(2*c/(e^(2*I*f*x + 2*I*e) + 1))^n*e^(-2*I*f*x - 2*I*e)/a, x)`

3.705.6 Sympy [F]

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{a + ia \tan(e + fx)} dx \\ &= - \frac{i \left(\int \frac{A(-ic \tan(e+fx)+c)^n}{\tan(e+fx)-i} dx + \int \frac{B(-ic \tan(e+fx)+c)^n \tan(e+fx)}{\tan(e+fx)-i} dx \right)}{a} \end{aligned}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e)),x)`

output `-I*(Integral(A*(-I*c*tan(e + f*x) + c)**n/(tan(e + f*x) - I), x) + Integral(B*(-I*c*tan(e + f*x) + c)**n*tan(e + f*x)/(tan(e + f*x) - I), x))/a`

3.705.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^n}{a + i a \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.705.8 Giac [F]

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^n}{a + i a \tan(e + fx)} dx \\ &= \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^n}{i a \tan(fx + e) + a} dx \end{aligned}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^n/(I*a*tan(f*x + e) + a), x)`

3.705.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^n}{a + i a \tan(e + fx)} dx \\ &= \int \frac{(A + B \tan(e + fx)) (c - c \tan(e + fx) li)^n}{a + a \tan(e + fx) li} dx \end{aligned}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*li)^n)/(a + a*tan(e + f*x)*li),x)`

output `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^n)/(a + a*tan(e + f*x)*1i), x)`

3.705. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^n}{a+ia \tan(e+fx)} dx$

3.706
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^4}{a+ia \tan(e+fx)} dx$$

3.706.1 Optimal result 6563
 3.706.2 Mathematica [A] (verified) 6563
 3.706.3 Rubi [A] (verified) 6564
 3.706.4 Maple [A] (verified) 6566
 3.706.5 Fricas [B] (verification not implemented) 6566
 3.706.6 Sympy [A] (verification not implemented) 6567
 3.706.7 Maxima [F(-2)] 6568
 3.706.8 Giac [B] (verification not implemented) 6568
 3.706.9 Mupad [B] (verification not implemented) 6569

3.706.1 Optimal result

Integrand size = 41, antiderivative size = 157

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{a + ia \tan(e + fx)} dx$$

$$= -\frac{4(3A + 5iB)c^4x}{a} - \frac{4(3iA - 5B)c^4 \log(\cos(e + fx))}{af} - \frac{8(A + iB)c^4}{af(i - \tan(e + fx))}$$

$$+ \frac{(5A + 12iB)c^4 \tan(e + fx)}{af} - \frac{(iA - 5B)c^4 \tan^2(e + fx)}{2af} - \frac{iBc^4 \tan^3(e + fx)}{3af}$$

output `-4*(3*A+5*I*B)*c^4*x/a-4*(3*I*A-5*B)*c^4*ln(cos(f*x+e))/a/f-8*(A+I*B)*c^4/a/f/(I-tan(f*x+e))+(5*A+12*I*B)*c^4*tan(f*x+e)/a/f-1/2*(I*A-5*B)*c^4*tan(f*x+e)^2/a/f-1/3*I*B*c^4*tan(f*x+e)^3/a/f`

3.706.2 Mathematica [A] (verified)

Time = 1.51 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.96

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{a + ia \tan(e + fx)} dx$$

$$= \frac{c^4(123A + 203iB + 24(3A + 5iB) \log(i - \tan(e + fx)) + (45iA - 83B + 24i(3A + 5iB) \log(i - \tan(e + \tan$$

$6af(-i + \tan$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^4)/(a + I*a*Tan[e + f*x]),x]`

output $(c^4*(123*A + (203*I)*B + 24*(3*A + (5*I)*B)*\text{Log}[I - \text{Tan}[e + f*x]] + ((45*I)*A - 83*B + (24*I)*(3*A + (5*I)*B)*\text{Log}[I - \text{Tan}[e + f*x]])*\text{Tan}[e + f*x] + 3*(9*A + (19*I)*B)*\text{Tan}[e + f*x]^2 + ((-3*I)*A + 13*B)*\text{Tan}[e + f*x]^3 - (2*I)*B*\text{Tan}[e + f*x]^4)/(6*a*f*(-I + \text{Tan}[e + f*x]))$

3.706.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.70, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - i c \tan(e + f x))^4 (A + B \tan(e + f x))}{a + i a \tan(e + f x)} dx$$

↓ 3042

$$\int \frac{(c - i c \tan(e + f x))^4 (A + B \tan(e + f x))}{a + i a \tan(e + f x)} dx$$

↓ 4071

$$\frac{a c \int \frac{c^3 (1 - i \tan(e + f x))^3 (A + B \tan(e + f x))}{a^2 (i \tan(e + f x) + 1)^2} d \tan(e + f x)}{f}$$

↓ 27

$$\frac{c^4 \int \frac{(1 - i \tan(e + f x))^3 (A + B \tan(e + f x))}{(i \tan(e + f x) + 1)^2} d \tan(e + f x)}{a f}$$

↓ 86

$$\frac{c^4 \int \left(-i B \tan^2(e + f x) + (5 B - i A) \tan(e + f x) + 5 A \left(\frac{12 i B}{5 A} + 1 \right) + \frac{4 i (3 A + 5 i B)}{\tan(e + f x) - i} - \frac{8 (A + i B)}{(\tan(e + f x) - i)^2} \right) d \tan(e + f x)}{a f}$$

↓ 2009

$$\frac{c^4 \left(-\frac{1}{2} (-5 B + i A) \tan^2(e + f x) + (5 A + 12 i B) \tan(e + f x) - \frac{8 (A + i B)}{-\tan(e + f x) + i} + 4 (-5 B + 3 i A) \log(-\tan(e + f x)) \right)}{a f}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^4)/(a + I*a*Tan[e + f*x]),x]`

output `(c^4*(4*((3*I)*A - 5*B)*Log[I - Tan[e + f*x]] - (8*(A + I*B))/(I - Tan[e + f*x]) + (5*A + (12*I)*B)*Tan[e + f*x] - ((I*A - 5*B)*Tan[e + f*x]^2)/2 - (I/3)*B*Tan[e + f*x]^3))/(a*f)`

3.706.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.706.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.43

method	result
risch	$-\frac{4c^4 e^{-2i(fx+e)} B}{af} + \frac{4ic^4 e^{-2i(fx+e)} A}{af} - \frac{40ic^4 Bx}{a} - \frac{24c^4 Ax}{a} - \frac{40ic^4 Be}{fa} - \frac{24c^4 Ae}{fa} - \frac{2c^4 (-12iA e^{4i(fx+e)} + \dots)}{a}$
norman	$\frac{(20ic^4 B + 13c^4 A) \tan(fx+e) - 4(5ic^4 B + 3c^4 A)x - 17ic^4 A + 21c^4 B + 5(7ic^4 B + 3c^4 A) \tan(fx+e)^3 - 4(5ic^4 B + 3c^4 A)x \tan(fx+e)}{1 + \tan(fx+e)^2}$
derivativedivides	$\frac{5c^4 B \tan(fx+e)^2}{2fa} - \frac{iB c^4 \tan(fx+e)^3}{3af} + \frac{5c^4 A \tan(fx+e)}{fa} - \frac{ic^4 A \tan(fx+e)^2}{2fa} + \frac{12ic^4 \tan(fx+e) B}{fa} - \frac{12c^4 A a}{fa}$
default	$\frac{5c^4 B \tan(fx+e)^2}{2fa} - \frac{iB c^4 \tan(fx+e)^3}{3af} + \frac{5c^4 A \tan(fx+e)}{fa} - \frac{ic^4 A \tan(fx+e)^2}{2fa} + \frac{12ic^4 \tan(fx+e) B}{fa} - \frac{12c^4 A a}{fa}$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output -4*c^4/a/f*exp(-2*I*(f*x+e))*B+4*I*c^4/a/f*exp(-2*I*(f*x+e))*A-40*I*c^4/a*B*x-24*c^4/a*A*x-40*I*c^4/f/a*B*e-24*c^4/f/a*A*e-2/3*c^4*(-12*I*A*exp(4*I*(f*x+e))+24*B*exp(4*I*(f*x+e))-27*I*A*exp(2*I*(f*x+e))+57*B*exp(2*I*(f*x+e))-15*I*A+37*B)/f/a/(exp(2*I*(f*x+e))+1)^3+20*c^4/f/a*ln(exp(2*I*(f*x+e))+1)*B-12*I*c^4/f/a*ln(exp(2*I*(f*x+e))+1)*A
```

3.706.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 298 vs. 2(137) = 274.

Time = 0.25 (sec) , antiderivative size = 298, normalized size of antiderivative = 1.90

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{a + ia \tan(e + fx)} dx = \frac{2(12(3A + 5iB)c^4 fxe^{(8i fx + 8ie)} + 6(-iA + B)c^4 + 6(6(3A + 5iB)c^4 fx + (-3iA + 5B)c^4)e^{(6i fx + 6ie)}}{a + ia \tan(e + fx)}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4/(a+I*a*tan(f*x+e)),x, algorith="fricas")
```

output
$$-2/3*(12*(3*A + 5*I*B)*c^4*f*x*e^(8*I*f*x + 8*I*e) + 6*(-I*A + B)*c^4 + 6*(6*(3*A + 5*I*B)*c^4*f*x + (-3*I*A + 5*B)*c^4)*e^(6*I*f*x + 6*I*e) + 3*(12*(3*A + 5*I*B)*c^4*f*x + 5*(-3*I*A + 5*B)*c^4)*e^(4*I*f*x + 4*I*e) + (12*(3*A + 5*I*B)*c^4*f*x + 11*(-3*I*A + 5*B)*c^4)*e^(2*I*f*x + 2*I*e) + 6*((3*I*A - 5*B)*c^4*e^(8*I*f*x + 8*I*e) + 3*(3*I*A - 5*B)*c^4*e^(6*I*f*x + 6*I*e) + 3*(3*I*A - 5*B)*c^4*e^(4*I*f*x + 4*I*e) + (3*I*A - 5*B)*c^4*e^(2*I*f*x + 2*I*e))*log(e^(2*I*f*x + 2*I*e) + 1)/(a*f*e^(8*I*f*x + 8*I*e) + 3*a*f*e^(6*I*f*x + 6*I*e) + 3*a*f*e^(4*I*f*x + 4*I*e) + a*f*e^(2*I*f*x + 2*I*e))$$

3.706.6 Sympy [A] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 328, normalized size of antiderivative = 2.09

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{a + ia \tan(e + fx)} dx$$

$$= \frac{30iAc^4 - 74Bc^4 + (54iAc^4e^{2ie} - 114Bc^4e^{2ie})e^{2ifx} + (24iAc^4e^{4ie} - 48Bc^4e^{4ie})e^{4ifx}}{3afe^{6ie}e^{6ifx} + 9afe^{4ie}e^{4ifx} + 9afe^{2ie}e^{2ifx} + 3af}$$

$$+ \begin{cases} \frac{(4iAc^4 - 4Bc^4)e^{-2ie}e^{-2ifx}}{af} & \text{for } afe^{2ie} \neq 0 \\ x \left(-\frac{24Ac^4 - 40iBc^4}{a} + \frac{(-24Ac^4e^{2ie} + 8Ac^4 - 40iBc^4e^{2ie} + 8iBc^4)e^{-2ie}}{a} \right) & \text{otherwise} \end{cases}$$

$$- \frac{4ic^4 \cdot (3A + 5iB) \log(e^{2ifx} + e^{-2ie})}{af} + \frac{x(-24Ac^4 - 40iBc^4)}{a}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**4/(a+I*a*tan(f*x+e)),x)`

output
$$(30*I*A*c**4 - 74*B*c**4 + (54*I*A*c**4*\exp(2*I*e) - 114*B*c**4*\exp(2*I*e))*\exp(2*I*f*x) + (24*I*A*c**4*\exp(4*I*e) - 48*B*c**4*\exp(4*I*e))*\exp(4*I*f*x))/(3*a*f*\exp(6*I*e)*\exp(6*I*f*x) + 9*a*f*\exp(4*I*e)*\exp(4*I*f*x) + 9*a*f*\exp(2*I*e)*\exp(2*I*f*x) + 3*a*f) + \text{Piecewise}(((4*I*A*c**4 - 4*B*c**4)*\exp(-2*I*e)*\exp(-2*I*f*x)/(a*f), \text{Ne}(a*f*\exp(2*I*e), 0)), (x*(-(-24*A*c**4 - 40*I*B*c**4)/a + (-24*A*c**4*\exp(2*I*e) + 8*A*c**4 - 40*I*B*c**4*\exp(2*I*e) + 8*I*B*c**4)*\exp(-2*I*e)/a), \text{True})) - 4*I*c**4*(3*A + 5*I*B)*\log(\exp(2*I*f*x) + \exp(-2*I*e))/(a*f) + x*(-24*A*c**4 - 40*I*B*c**4)/a$$

3.706.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{a + ia \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4/(a+I*a*tan(f*x+e)),x, algo
rithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.706.8 Giac [B] (verification not implemented)Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 424 vs. $2(137) = 274$.

Time = 0.66 (sec) , antiderivative size = 424, normalized size of antiderivative = 2.70

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{a + ia \tan(e + fx)} dx$$

$$= \frac{2 \left(\frac{6(-3iAc^4 + 5Bc^4) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)}{a} - \frac{12(-3iAc^4 + 5Bc^4) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - i)}{a} - \frac{6(3iAc^4 - 5Bc^4) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)}{a} \right)}{a}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4/(a+I*a*tan(f*x+e)),x, algo
rithm="giac")
```

```
output 2/3*(6*(-3*I*A*c^4 + 5*B*c^4)*log(tan(1/2*f*x + 1/2*e) + 1)/a - 12*(-3*I*A
*c^4 + 5*B*c^4)*log(tan(1/2*f*x + 1/2*e) - I)/a - 6*(3*I*A*c^4 - 5*B*c^4)*
log(tan(1/2*f*x + 1/2*e) - 1)/a - 6*(9*I*A*c^4*tan(1/2*f*x + 1/2*e)^2 - 15
*B*c^4*tan(1/2*f*x + 1/2*e)^2 + 22*A*c^4*tan(1/2*f*x + 1/2*e) + 34*I*B*c^4
*tan(1/2*f*x + 1/2*e) - 9*I*A*c^4 + 15*B*c^4)/(a*(tan(1/2*f*x + 1/2*e) - I
)^2) - (-33*I*A*c^4*tan(1/2*f*x + 1/2*e)^6 + 55*B*c^4*tan(1/2*f*x + 1/2*e)
^6 + 15*A*c^4*tan(1/2*f*x + 1/2*e)^5 + 36*I*B*c^4*tan(1/2*f*x + 1/2*e)^5 +
102*I*A*c^4*tan(1/2*f*x + 1/2*e)^4 - 180*B*c^4*tan(1/2*f*x + 1/2*e)^4 - 3
0*A*c^4*tan(1/2*f*x + 1/2*e)^3 - 76*I*B*c^4*tan(1/2*f*x + 1/2*e)^3 - 102*I
*A*c^4*tan(1/2*f*x + 1/2*e)^2 + 180*B*c^4*tan(1/2*f*x + 1/2*e)^2 + 15*A*c^
4*tan(1/2*f*x + 1/2*e) + 36*I*B*c^4*tan(1/2*f*x + 1/2*e) + 33*I*A*c^4 - 55
*B*c^4)/((tan(1/2*f*x + 1/2*e)^2 - 1)^3*a))/f
```

3.706. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^4}{a+ia \tan(e+fx)} dx$

3.706.9 Mupad [B] (verification not implemented)

Time = 8.43 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.31

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{a + ia \tan(e + fx)} dx$$

$$= \frac{\ln(\tan(e + fx) - i) \left(-\frac{20Bc^4}{a} + \frac{Ac^4 12i}{a} \right)}{f} - \frac{\tan(e + fx)^2 \left(-\frac{Bc^4}{a} + \frac{c^4(A+B3i)1i}{2a} \right)}{f} - \frac{\left(\frac{4Ac^4+Bc^4 12i}{a} 1i - \frac{(12Ac^4+Bc^4 20i) 1i}{a} \right)}{f(1 + \tan(e + fx) 1i)}$$

$$+ \frac{\tan(e + fx) \left(\frac{2c^4(A+B3i)}{a} + \frac{Bc^4 3i}{a} - \frac{c^4(-B+A1i)3i}{a} \right)}{f} - \frac{Bc^4 \tan(e + fx)^3 1i}{3af}$$

```
input int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^4)/(a + a*tan(e + f*x)*1i),x)
```

```
output (log(tan(e + f*x) - 1i)*((A*c^4*12i)/a - (20*B*c^4)/a))/f - (tan(e + f*x)^2*((c^4*(A + B*3i)*1i)/(2*a) - (B*c^4)/a))/f - (((4*A*c^4 + B*c^4*12i)*1i)/a - ((12*A*c^4 + B*c^4*20i)*1i)/a)/(f*(tan(e + f*x)*1i + 1)) + (tan(e + f*x)*((2*c^4*(A + B*3i))/a + (B*c^4*3i)/a - (c^4*(A*1i - B)*3i)/a))/f - (B*c^4*tan(e + f*x)^3*1i)/(3*a*f)
```


3.707
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^3}{a+ia \tan(e+fx)} dx$$

3.707.1 Optimal result 6570
 3.707.2 Mathematica [A] (verified) 6570
 3.707.3 Rubi [A] (verified) 6571
 3.707.4 Maple [A] (verified) 6573
 3.707.5 Fricas [B] (verification not implemented) 6573
 3.707.6 Sympy [A] (verification not implemented) 6574
 3.707.7 Maxima [F(-2)] 6575
 3.707.8 Giac [B] (verification not implemented) 6575
 3.707.9 Mupad [B] (verification not implemented) 6576

3.707.1 Optimal result

Integrand size = 41, antiderivative size = 121

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{a + ia \tan(e + fx)} dx$$

$$= -\frac{4(A + 2iB)c^3x}{a} - \frac{4(iA - 2B)c^3 \log(\cos(e + fx))}{af} - \frac{4(A + iB)c^3}{af(i - \tan(e + fx))} + \frac{(A + 4iB)c^3 \tan(e + fx)}{af} + \frac{Bc^3 \tan^2(e + fx)}{2af}$$

output `-4*(A+2*I*B)*c^3*x/a-4*(I*A-2*B)*c^3*ln(cos(f*x+e))/a/f-4*(A+I*B)*c^3/a/f/(I-tan(f*x+e))+(A+4*I*B)*c^3*tan(f*x+e)/a/f+1/2*B*c^3*tan(f*x+e)^2/a/f`

3.707.2 Mathematica [A] (verified)

Time = 5.11 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.02

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{a + ia \tan(e + fx)} dx$$

$$= \frac{c^3(14A + 27iB + 8(A + 2iB) \log(i - \tan(e + fx)) + (4iA - 11B + 8i(A + 2iB) \log(i - \tan(e + fx))) \tan(e + fx))}{2af(-i + \tan(e + fx))}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^3)/(a + I*a*Tan[e + f*x]),x]`

output $(c^3*(14*A + (27*I)*B + 8*(A + (2*I)*B)*\text{Log}[I - \text{Tan}[e + f*x]] + ((4*I)*A - 11*B + (8*I)*(A + (2*I)*B)*\text{Log}[I - \text{Tan}[e + f*x]])*\text{Tan}[e + f*x] + (2*A + (7*I)*B)*\text{Tan}[e + f*x]^2 + B*\text{Tan}[e + f*x]^3)/(2*a*f*(-I + \text{Tan}[e + f*x]))$

3.707.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.70, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - ic \tan(e + fx))^3 (A + B \tan(e + fx))}{a + ia \tan(e + fx)} dx$$

↓ 3042

$$\int \frac{(c - ic \tan(e + fx))^3 (A + B \tan(e + fx))}{a + ia \tan(e + fx)} dx$$

↓ 4071

$$\frac{ac \int \frac{c^2(1-i \tan(e+fx))^2(A+B \tan(e+fx))}{a^2(i \tan(e+fx)+1)^2} d \tan(e + fx)}{f}$$

↓ 27

$$\frac{c^3 \int \frac{(1-i \tan(e+fx))^2(A+B \tan(e+fx))}{(i \tan(e+fx)+1)^2} d \tan(e + fx)}{af}$$

↓ 86

$$\frac{c^3 \int \left(-\frac{4(A+iB)}{(\tan(e+fx)-i)^2} + A\left(\frac{4iB}{A} + 1\right) + B \tan(e + fx) + \frac{4i(A+2iB)}{\tan(e+fx)-i} \right) d \tan(e + fx)}{af}$$

↓ 2009

$$\frac{c^3 \left((A + 4iB) \tan(e + fx) - \frac{4(A+iB)}{-\tan(e+fx)+i} + 4(-2B + iA) \log(-\tan(e + fx) + i) + \frac{1}{2}B \tan^2(e + fx) \right)}{af}$$

input $\text{Int}[(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^3/(a + I*a*\text{Tan}[e + f*x]),x]$

3.707. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^3}{a+ia \tan(e+fx)} dx$

```
output (c^3*(4*(I*A - 2*B)*Log[I - Tan[e + f*x]] - (4*(A + I*B))/(I - Tan[e + f*x
])) + (A + (4*I)*B)*Tan[e + f*x] + (B*Tan[e + f*x]^2)/2)/(a*f)
```

3.707.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```


output $-2*(4*(A + 2*I*B)*c^3*f*x*e^{(6*I*f*x + 6*I*e)} + (-I*A + B)*c^3 + 2*(4*(A + 2*I*B)*c^3*f*x + (-I*A + 2*B)*c^3)*e^{(4*I*f*x + 4*I*e)} + (4*(A + 2*I*B)*c^3*f*x + 3*(-I*A + 2*B)*c^3)*e^{(2*I*f*x + 2*I*e)} + 2*((I*A - 2*B)*c^3*e^{(6*I*f*x + 6*I*e)} + 2*(I*A - 2*B)*c^3*e^{(4*I*f*x + 4*I*e)} + (I*A - 2*B)*c^3*e^{(2*I*f*x + 2*I*e)})*\log(e^{(2*I*f*x + 2*I*e)} + 1)/(a*f*e^{(6*I*f*x + 6*I*e)} + 2*a*f*e^{(4*I*f*x + 4*I*e)} + a*f*e^{(2*I*f*x + 2*I*e)})$

3.707.6 Sympy [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 265, normalized size of antiderivative = 2.19

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{a + ia \tan(e + fx)} dx$$

$$= \frac{2iAc^3 - 8Bc^3 + (2iAc^3e^{2ie} - 6Bc^3e^{2ie})e^{2ifx}}{afe^{4ie}e^{4ifx} + 2afe^{2ie}e^{2ifx} + af}$$

$$+ \begin{cases} \frac{(2iAc^3 - 2Bc^3)e^{-2ie}e^{-2ifx}}{af} & \text{for } afe^{2ie} \neq 0 \\ x \left(-\frac{8Ac^3 - 16iBc^3}{a} + \frac{(-8Ac^3e^{2ie} + 4Ac^3 - 16iBc^3e^{2ie} + 4iBc^3)e^{-2ie}}{a} \right) & \text{otherwise} \end{cases}$$

$$- \frac{4ic^3(A + 2iB) \log(e^{2ifx} + e^{-2ie})}{af} + \frac{x(-8Ac^3 - 16iBc^3)}{a}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**3/(a+I*a*tan(f*x+e)),x)`

output $(2*I*A*c**3 - 8*B*c**3 + (2*I*A*c**3*\exp(2*I*e) - 6*B*c**3*\exp(2*I*e))*\exp(2*I*f*x))/(a*f*\exp(4*I*e)*\exp(4*I*f*x) + 2*a*f*\exp(2*I*e)*\exp(2*I*f*x) + a*f) + \text{Piecewise}(((2*I*A*c**3 - 2*B*c**3)*\exp(-2*I*e)*\exp(-2*I*f*x)/(a*f), \text{Ne}(a*f*\exp(2*I*e), 0)), (x*(-(-8*A*c**3 - 16*I*B*c**3)/a + (-8*A*c**3*\exp(2*I*e) + 4*A*c**3 - 16*I*B*c**3*\exp(2*I*e) + 4*I*B*c**3)*\exp(-2*I*e)/a), \text{True})) - 4*I*c**3*(A + 2*I*B)*\log(\exp(2*I*f*x) + \exp(-2*I*e))/(a*f) + x*(-8*A*c**3 - 16*I*B*c**3)/a$

3.707.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{a + ia \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3/(a+I*a*tan(f*x+e)),x, algorith="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.707.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 307 vs. $2(107) = 214$.

Time = 0.60 (sec) , antiderivative size = 307, normalized size of antiderivative = 2.54

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{a + ia \tan(e + fx)} dx$$

$$= \frac{2 \left(\frac{2(-iAc^3 + 2Bc^3) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)}{a} - \frac{4(-iAc^3 + 2Bc^3) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - i)}{a} - \frac{2(iAc^3 - 2Bc^3) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)}{a} \right)}{f}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3/(a+I*a*tan(f*x+e)),x, algorith="giac")`

output `2*(2*(-I*A*c^3 + 2*B*c^3)*log(tan(1/2*f*x + 1/2*e) + 1)/a - 4*(-I*A*c^3 + 2*B*c^3)*log(tan(1/2*f*x + 1/2*e) - I)/a - 2*(I*A*c^3 - 2*B*c^3)*log(tan(1/2*f*x + 1/2*e) - 1)/a - (5*A*c^3*tan(1/2*f*x + 1/2*e)^5 + 8*I*B*c^3*tan(1/2*f*x + 1/2*e)^5 - 2*I*A*c^3*tan(1/2*f*x + 1/2*e)^4 + 7*B*c^3*tan(1/2*f*x + 1/2*e)^4 - 10*A*c^3*tan(1/2*f*x + 1/2*e)^3 - 14*I*B*c^3*tan(1/2*f*x + 1/2*e)^3 + 2*I*A*c^3*tan(1/2*f*x + 1/2*e)^2 - 7*B*c^3*tan(1/2*f*x + 1/2*e)^2 + 5*A*c^3*tan(1/2*f*x + 1/2*e) + 8*I*B*c^3*tan(1/2*f*x + 1/2*e))/((tan(1/2*f*x + 1/2*e)^3 - I*tan(1/2*f*x + 1/2*e)^2 - tan(1/2*f*x + 1/2*e) + I)^2*a)/f`

3.707. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^3}{a+ia \tan(e+fx)} dx$

3.707.9 Mupad [B] (verification not implemented)

Time = 8.48 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.12

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{a + ia \tan(e + fx)} dx$$

$$= \frac{\ln(\tan(e + fx) - i) \left(-\frac{8Bc^3}{a} + \frac{Ac^3 4i}{a} \right)}{f} + \frac{\tan(e + fx) \left(\frac{c^3(A+B2i)}{a} + \frac{Bc^3 2i}{a} \right)}{f}$$

$$+ \frac{\frac{4Bc^3}{a} + \frac{(4Ac^3 + Bc^3 8i) li}{a}}{f(1 + \tan(e + fx) li)} + \frac{Bc^3 \tan(e + fx)^2}{2af}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^3)/(a + a*tan(e + f*x)*1i),x)`

output `(log(tan(e + f*x) - 1i)*((A*c^3*4i)/a - (8*B*c^3)/a))/f + (tan(e + f*x)*((c^3*(A + B*2i))/a + (B*c^3*2i)/a))/f + (((4*A*c^3 + B*c^3*8i)*1i)/a + (4*B*c^3)/a)/(f*(tan(e + f*x)*1i + 1)) + (B*c^3*tan(e + f*x)^2)/(2*a*f)`

3.708 $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^2}{a+ia \tan(e+fx)} dx$

3.708.1 Optimal result	6577
3.708.2 Mathematica [A] (verified)	6577
3.708.3 Rubi [A] (verified)	6578
3.708.4 Maple [A] (verified)	6580
3.708.5 Fricas [A] (verification not implemented)	6580
3.708.6 Sympy [A] (verification not implemented)	6581
3.708.7 Maxima [F(-2)]	6581
3.708.8 Giac [B] (verification not implemented)	6582
3.708.9 Mupad [B] (verification not implemented)	6582

3.708.1 Optimal result

Integrand size = 41, antiderivative size = 96

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{a + ia \tan(e + fx)} dx$$

$$= -\frac{(A + 3iB)c^2x}{a} - \frac{(iA - 3B)c^2 \log(\cos(e + fx))}{af}$$

$$- \frac{2(A + iB)c^2}{af(i - \tan(e + fx))} + \frac{iBc^2 \tan(e + fx)}{af}$$

output `-(A+3*I*B)*c^2*x/a-(I*A-3*B)*c^2*ln(cos(f*x+e))/a/f-2*(A+I*B)*c^2/a/f/(I-tan(f*x+e))+I*B*c^2*tan(f*x+e)/a/f`

3.708.2 Mathematica [A] (verified)

Time = 4.42 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.86

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{a + ia \tan(e + fx)} dx$$

$$= \frac{\frac{B(c-ic \tan(e+fx))^2}{a+ia \tan(e+fx)} + \frac{(iA-3B)c^2(\log(i-\tan(e+fx))-\frac{2i}{-i+\tan(e+fx)})}{a}}{f}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^2)/(a + I*a*Tan[e + f*x]),x]`

output $((B*(c - I*c*\text{Tan}[e + f*x])^2)/(a + I*a*\text{Tan}[e + f*x]) + ((I*A - 3*B)*c^2*(\text{Log}[I - \text{Tan}[e + f*x]] - (2*I)/(-I + \text{Tan}[e + f*x])))/a)/f$

3.708.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.71, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c - ictan(e + fx))^2(A + B \tan(e + fx))}{a + ia \tan(e + fx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c - ictan(e + fx))^2(A + B \tan(e + fx))}{a + ia \tan(e + fx)} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{c(1-i \tan(e+fx))(A+B \tan(e+fx))}{a^2(i \tan(e+fx)+1)^2} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{c^2 \int \frac{(1-i \tan(e+fx))(A+B \tan(e+fx))}{(i \tan(e+fx)+1)^2} d \tan(e + fx)}{af} \\ & \quad \downarrow \text{86} \\ & \frac{c^2 \int \left(-\frac{2(A+iB)}{(\tan(e+fx)-i)^2} + iB + \frac{i(A+3iB)}{\tan(e+fx)-i} \right) d \tan(e + fx)}{af} \\ & \quad \downarrow \text{2009} \\ & \frac{c^2 \left(-\frac{2(A+iB)}{-\tan(e+fx)+i} + (-3B + iA) \log(-\tan(e + fx) + i) + iB \tan(e + fx) \right)}{af} \end{aligned}$$

input $\text{Int}[(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^2/(a + I*a*\text{Tan}[e + f*x]),x]$

```
output (c^2*((I*A - 3*B)*Log[I - Tan[e + f*x]] - (2*(A + I*B))/(I - Tan[e + f*x])
+ I*B*Tan[e + f*x]))/(a*f)
```

3.708.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.708.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.60

method	result
derivativedivides	$\frac{iB c^2 \tan(fx+e)}{af} + \frac{2ic^2B}{fa(-i+\tan(fx+e))} + \frac{2c^2A}{fa(-i+\tan(fx+e))} + \frac{ic^2A \ln(1+\tan(fx+e)^2)}{2fa} - \frac{3c^2B \ln(1+\tan(fx+e)^2)}{2fa}$
default	$\frac{iB c^2 \tan(fx+e)}{af} + \frac{2ic^2B}{fa(-i+\tan(fx+e))} + \frac{2c^2A}{fa(-i+\tan(fx+e))} + \frac{ic^2A \ln(1+\tan(fx+e)^2)}{2fa} - \frac{3c^2B \ln(1+\tan(fx+e)^2)}{2fa}$
norman	$\frac{(3ic^2B+2c^2A) \tan(fx+e) + ic^2B \tan(fx+e)^3 - (3ic^2B+c^2A)x}{af} - \frac{-2iAc^2+2c^2B}{af} - \frac{(3ic^2B+c^2A)x \tan(fx+e)^2}{a} - \frac{(-iAc^2+3c^2B)x}{a}$ $\frac{1+\tan(fx+e)^2}{1+\tan(fx+e)^2}$
risch	$-\frac{c^2e^{-2i(fx+e)}B}{af} + \frac{ic^2e^{-2i(fx+e)}A}{af} - \frac{6ic^2Bx}{a} - \frac{2c^2Ax}{a} - \frac{6ic^2Be}{af} - \frac{2c^2Ae}{af} - \frac{2c^2B}{fa(e^{2i(fx+e)}+1)} + \frac{3c^2 \ln(e^{2i(fx+e)}+1)}{fa(e^{2i(fx+e)}+1)}$

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output `I*B*c^2*tan(f*x+e)/a/f+2*I/f*c^2/a/(-I+tan(f*x+e))*B+2/f*c^2/a/(-I+tan(f*x+e))*A+1/2*I/f*c^2/a*A*ln(1+tan(f*x+e)^2)-3/2/f*c^2/a*B*ln(1+tan(f*x+e)^2)-1/f*c^2/a*A*arctan(tan(f*x+e))-3*I/f*c^2/a*B*arctan(tan(f*x+e))`

3.708.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.59

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{a + ia \tan(e + fx)} dx = \frac{2(A + 3iB)c^2 fxe^{(4i fx + 4i e)} - (iA - B)c^2 + (2(A + 3iB)c^2 fx - (iA - 3B)c^2)e^{(2i fx + 2i e)} - ((-iA + 3B)c^2 fxe^{(4i fx + 4i e)} + afe^{(2i fx + 2i e)})}{afe^{(4i fx + 4i e)} + afe^{(2i fx + 2i e)}}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`

output `-(2*(A + 3*I*B)*c^2*f*x*e^(4*I*f*x + 4*I*e) - (I*A - B)*c^2 + (2*(A + 3*I*B)*c^2*f*x - (I*A - 3*B)*c^2)*e^(2*I*f*x + 2*I*e) - ((-I*A + 3*B)*c^2*e^(4*I*f*x + 4*I*e) + (-I*A + 3*B)*c^2*e^(2*I*f*x + 2*I*e))*log(e^(2*I*f*x + 2*I*e) + 1)/(a*f*e^(4*I*f*x + 4*I*e) + a*f*e^(2*I*f*x + 2*I*e))`

3.708.6 Sympy [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 194, normalized size of antiderivative = 2.02

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{a + ia \tan(e + fx)} dx$$

$$= -\frac{2Bc^2}{af e^{2ie} e^{2ifx} + af} + \begin{cases} \frac{(iAc^2 - Bc^2)e^{-2ie} e^{-2ifx}}{af} & \text{for } af e^{2ie} \neq 0 \\ x \left(-\frac{-2Ac^2 - 6iBc^2}{a} + \frac{(-2Ac^2 e^{2ie} + 2Ac^2 - 6iBc^2 e^{2ie} + 2iBc^2) e^{-2ie}}{a} \right) & \text{otherwise} \end{cases}$$

$$- \frac{ic^2(A + 3iB) \log(e^{2ifx} + e^{-2ie})}{af} + \frac{x(-2Ac^2 - 6iBc^2)}{a}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**2/(a+I*a*tan(f*x+e)),x)
```

```
output -2*B*c**2/(a*f*exp(2*I*e)*exp(2*I*f*x) + a*f) + Piecewise(((I*A*c**2 - B*c**2)*exp(-2*I*e)*exp(-2*I*f*x)/(a*f), Ne(a*f*exp(2*I*e), 0)), (x*(-(-2*A*c**2 - 6*I*B*c**2)/a + (-2*A*c**2*exp(2*I*e) + 2*A*c**2 - 6*I*B*c**2*exp(2*I*e) + 2*I*B*c**2)*exp(-2*I*e)/a), True)) - I*c**2*(A + 3*I*B)*log(exp(2*I*f*x) + exp(-2*I*e))/(a*f) + x*(-2*A*c**2 - 6*I*B*c**2)/a
```

3.708.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{a + ia \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2/(a+I*a*tan(f*x+e)),x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.708.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 271 vs. $2(84) = 168$.

Time = 0.51 (sec) , antiderivative size = 271, normalized size of antiderivative = 2.82

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{a + ia \tan(e + fx)} dx$$

$$= \frac{(-iAc^2 + 3Bc^2) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)}{a} + \frac{2(iAc^2 - 3Bc^2) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - i)}{a} - \frac{(iAc^2 - 3Bc^2) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)}{a} - \frac{-iAc^2 \tan(\frac{1}{2}fx + \frac{1}{2}e)}{a}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

output `((-I*A*c^2 + 3*B*c^2)*log(tan(1/2*f*x + 1/2*e) + 1)/a + 2*(I*A*c^2 - 3*B*c^2)*log(tan(1/2*f*x + 1/2*e) - I)/a - (I*A*c^2 - 3*B*c^2)*log(tan(1/2*f*x + 1/2*e) - 1)/a - (-I*A*c^2*tan(1/2*f*x + 1/2*e)^2 + 3*B*c^2*tan(1/2*f*x + 1/2*e)^2 + 2*I*B*c^2*tan(1/2*f*x + 1/2*e) + I*A*c^2 - 3*B*c^2)/((tan(1/2*f*x + 1/2*e)^2 - 1)*a) - (3*I*A*c^2*tan(1/2*f*x + 1/2*e)^2 - 9*B*c^2*tan(1/2*f*x + 1/2*e)^2 + 10*A*c^2*tan(1/2*f*x + 1/2*e) + 22*I*B*c^2*tan(1/2*f*x + 1/2*e) - 3*I*A*c^2 + 9*B*c^2)/(a*(tan(1/2*f*x + 1/2*e) - I)^2))/f`

3.708.9 Mupad [B] (verification not implemented)

Time = 8.81 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.15

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{a + ia \tan(e + fx)} dx$$

$$= \frac{\ln(\tan(e + fx) - i) \left(-\frac{3Bc^2}{a} + \frac{Ac^2 li}{a} \right)}{f} + \frac{(Ac^2 - Bc^2 li) li}{a} + \frac{(Ac^2 + Bc^2 3i) li}{a} + \frac{Bc^2 \tan(e + fx) li}{af}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^2)/(a + a*tan(e + f*x)*1i),x)`

output $(\log(\tan(e + f*x) - 1i)*((A*c^2*1i)/a - (3*B*c^2)/a))/f + (((A*c^2 - B*c^2*1i)*1i)/a + ((A*c^2 + B*c^2*3i)*1i)/a)/(f*(\tan(e + f*x)*1i + 1)) + (B*c^2*\tan(e + f*x)*1i)/(a*f)$

3.708. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^2}{a+ia \tan(e+fx)} dx$

3.709
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))}{a+ia \tan(e+fx)} dx$$

3.709.1 Optimal result 6584
 3.709.2 Mathematica [A] (verified) 6584
 3.709.3 Rubi [A] (verified) 6585
 3.709.4 Maple [A] (verified) 6586
 3.709.5 Fricas [A] (verification not implemented) 6587
 3.709.6 Sympy [A] (verification not implemented) 6587
 3.709.7 Maxima [F(-2)] 6588
 3.709.8 Giac [B] (verification not implemented) 6588
 3.709.9 Mupad [B] (verification not implemented) 6589

3.709.1 Optimal result

Integrand size = 39, antiderivative size = 57

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))}{a + ia \tan(e + fx)} dx$$

$$= -\frac{iBcx}{a} + \frac{Bc \log(\cos(e + fx))}{af} - \frac{(A + iB)c}{af(i - \tan(e + fx))}$$

output `-I*B*c*x/a+B*c*ln(cos(f*x+e))/a/f-(A+I*B)*c/a/f/(I-tan(f*x+e))`

3.709.2 Mathematica [A] (verified)

Time = 0.74 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.82

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))}{a + ia \tan(e + fx)} dx = -\frac{c \left(B \log(i - \tan(e + fx)) + \frac{A+iB}{i-\tan(e+fx)} \right)}{af}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x]))/(a + I*a*Tan[e + f*x]),x]`

output `-((c*(B*Log[I - Tan[e + f*x]] + (A + I*B)/(I - Tan[e + f*x])))/(a*f))`

3.709.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.84, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.128$, Rules used = {3042, 4071, 27, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - ic \tan(e + fx))(A + B \tan(e + fx))}{a + ia \tan(e + fx)} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c - ic \tan(e + fx))(A + B \tan(e + fx))}{a + ia \tan(e + fx)} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A+B \tan(e+fx)}{a^2(i \tan(e+fx)+1)^2} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{c \int \frac{A+B \tan(e+fx)}{(i \tan(e+fx)+1)^2} d \tan(e + fx)}{af} \\
 & \quad \downarrow \text{49} \\
 & \frac{c \int \left(\frac{-A-iB}{(\tan(e+fx)-i)^2} - \frac{B}{\tan(e+fx)-i} \right) d \tan(e + fx)}{af} \\
 & \quad \downarrow \text{2009} \\
 & \frac{c \left(-\frac{A+iB}{-\tan(e+fx)+i} - B \log(-\tan(e + fx) + i) \right)}{af}
 \end{aligned}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x]))/(a + I*a*Tan[e + f*x]),x]`

output `(c*(-(B*Log[I - Tan[e + f*x]]) - (A + I*B)/(I - Tan[e + f*x])))/(a*f)`

3.709.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.709.4 Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.46

method	result	size
derivativedivides	$\frac{icB}{fa(-i+\tan(fx+e))} + \frac{cA}{fa(-i+\tan(fx+e))} - \frac{cB \ln(1+\tan(fx+e)^2)}{2fa} - \frac{icB \arctan(\tan(fx+e))}{fa}$	83
default	$\frac{icB}{fa(-i+\tan(fx+e))} + \frac{cA}{fa(-i+\tan(fx+e))} - \frac{cB \ln(1+\tan(fx+e)^2)}{2fa} - \frac{icB \arctan(\tan(fx+e))}{fa}$	83
risch	$-\frac{ce^{-2i(fx+e)}B}{2af} + \frac{ice^{-2i(fx+e)}A}{2af} - \frac{2icBx}{a} - \frac{2icBe}{af} + \frac{cB \ln(e^{2i(fx+e)}+1)}{af}$	83
norman	$\frac{(icB+cA) \tan(fx+e) - icA+cB}{af} - \frac{icBx}{a} - \frac{icBx \tan(fx+e)^2}{a} - \frac{cB \ln(1+\tan(fx+e)^2)}{2fa}$	102

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)`

3.709. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))}{a+ia \tan(e+fx)} dx$

output $I/f*c/a/(-I+\tan(f*x+e))*B+1/f*c/a/(-I+\tan(f*x+e))*A-1/2/f*c/a*B*\ln(1+\tan(f*x+e)^2)-I/f*c/a*B*\arctan(\tan(f*x+e))$

3.709.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.18

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))}{a + ia \tan(e + fx)} dx$$

$$= \frac{(-4i B c f x e^{(2i f x + 2i e)} + 2 B c e^{(2i f x + 2i e)} \log(e^{(2i f x + 2i e)} + 1) + (i A - B) c) e^{(-2i f x - 2i e)}}{2 a f}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`

output $1/2*(-4*I*B*c*f*x*e^{(2*I*f*x + 2*I*e)} + 2*B*c*e^{(2*I*f*x + 2*I*e)}*\log(e^{(2*I*f*x + 2*I*e)} + 1) + (I*A - B)*c)*e^{(-2*I*f*x - 2*I*e)}/(a*f)$

3.709.6 Sympy [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.96

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))}{a + ia \tan(e + fx)} dx$$

$$= -\frac{2i B c x}{a} + \frac{B c \log(e^{2i f x} + e^{-2i e})}{a f} + \begin{cases} \frac{(i A c - B c) e^{-2i e} e^{-2i f x}}{2 a f} & \text{for } a f e^{2i e} \neq 0 \\ x \left(\frac{2i B c}{a} + \frac{(A c - 2i B c e^{2i e} + i B c) e^{-2i e}}{a} \right) & \text{otherwise} \end{cases}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e)),x)`

output $-2*I*B*c*x/a + B*c*\log(\exp(2*I*f*x) + \exp(-2*I*e))/(a*f) + \text{Piecewise}(((I*A*c - B*c)*\exp(-2*I*e)*\exp(-2*I*f*x)/(2*a*f), \text{Ne}(a*f*\exp(2*I*e), 0)), (x*(2*I*B*c/a + (A*c - 2*I*B*c*\exp(2*I*e) + I*B*c)*\exp(-2*I*e)/a), \text{True}))$

3.709.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))}{a + i a \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.709.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 123 vs. $2(48) = 96$.

Time = 0.39 (sec) , antiderivative size = 123, normalized size of antiderivative = 2.16

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))}{a + i a \tan(e + fx)} dx$$

$$= \frac{B c \log(\tan(\frac{1}{2} f x + \frac{1}{2} e) + 1)}{a} - \frac{2 B c \log(\tan(\frac{1}{2} f x + \frac{1}{2} e) - i)}{a} + \frac{B c \log(\tan(\frac{1}{2} f x + \frac{1}{2} e) - 1)}{a} + \frac{3 B c \tan(\frac{1}{2} f x + \frac{1}{2} e)^2 - 2 A c \tan(\frac{1}{2} f x + \frac{1}{2} e) - 8 i}{a (\tan(\frac{1}{2} f x + \frac{1}{2} e) - i)^2} f$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

output `(B*c*log(tan(1/2*f*x + 1/2*e) + 1)/a - 2*B*c*log(tan(1/2*f*x + 1/2*e) - I)/a + B*c*log(tan(1/2*f*x + 1/2*e) - 1)/a + (3*B*c*tan(1/2*f*x + 1/2*e)^2 - 2*A*c*tan(1/2*f*x + 1/2*e) - 8*I*B*c*tan(1/2*f*x + 1/2*e) - 3*B*c)/(a*(tan(1/2*f*x + 1/2*e) - I)^2))/f`

3.709.9 Mupad [B] (verification not implemented)

Time = 8.23 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.95

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))}{a + i a \tan(e + fx)} dx$$

$$= \frac{-\frac{Bc}{a} + \frac{Ac \operatorname{li}}{a}}{f(1 + \tan(e + fx) i)} - \frac{Bc \ln(\tan(e + fx) - i)}{a f}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i))/(a + a*tan(e + f*x)*1i),x)`

output `((A*c*1i)/a - (B*c)/a)/(f*(tan(e + f*x)*1i + 1)) - (B*c*log(tan(e + f*x) - 1i))/(a*f)`

3.710 $\int \frac{A+B \tan(e+fx)}{a+ia \tan(e+fx)} dx$

3.710.1 Optimal result	6590
3.710.2 Mathematica [A] (verified)	6590
3.710.3 Rubi [A] (verified)	6591
3.710.4 Maple [A] (verified)	6592
3.710.5 Fricas [A] (verification not implemented)	6592
3.710.6 Sympy [A] (verification not implemented)	6593
3.710.7 Maxima [F(-2)]	6593
3.710.8 Giac [B] (verification not implemented)	6593
3.710.9 Mupad [B] (verification not implemented)	6594

3.710.1 Optimal result

Integrand size = 26, antiderivative size = 47

$$\int \frac{A + B \tan(e + fx)}{a + ia \tan(e + fx)} dx = \frac{(A - iB)x}{2a} + \frac{iA - B}{2f(a + ia \tan(e + fx))}$$

output `1/2*(A-I*B)*x/a+1/2*(I*A-B)/f/(a+I*a*tan(f*x+e))`

3.710.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.19

$$\int \frac{A + B \tan(e + fx)}{a + ia \tan(e + fx)} dx = \frac{(A - iB) \arctan(\tan(e + fx))}{2af} - \frac{A + iB}{2af(i - \tan(e + fx))}$$

input `Integrate[(A + B*Tan[e + f*x])/(a + I*a*Tan[e + f*x]),x]`

output `((A - I*B)*ArcTan[Tan[e + f*x]]/(2*a*f) - (A + I*B)/(2*a*f*(I - Tan[e + f*x])))`

3.710.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {3042, 4009, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(e + fx)}{a + ia \tan(e + fx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(e + fx)}{a + ia \tan(e + fx)} dx \\ & \quad \downarrow \text{4009} \\ & \frac{(A - iB) \int 1 dx}{2a} + \frac{-B + iA}{2f(a + ia \tan(e + fx))} \\ & \quad \downarrow \text{24} \\ & \frac{-B + iA}{2f(a + ia \tan(e + fx))} + \frac{x(A - iB)}{2a} \end{aligned}$$

input `Int[(A + B*Tan[e + f*x])/(a + I*a*Tan[e + f*x]),x]`

output `((A - I*B)*x)/(2*a) + (I*A - B)/(2*f*(a + I*a*Tan[e + f*x]))`

3.710.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4009 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a
*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x
], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2
, 0] && LtQ[m, 0]
```

3.710.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.15

method	result	size
risch	$-\frac{ixB}{2a} + \frac{xA}{2a} - \frac{e^{-2i(fx+e)}B}{4af} + \frac{ie^{-2i(fx+e)}A}{4af}$	54
derivativedivides	$\frac{A}{2fa(-i+\tan(fx+e))} + \frac{iB}{2fa(-i+\tan(fx+e))} + \frac{A \arctan(\tan(fx+e))}{2fa} - \frac{iB \arctan(\tan(fx+e))}{2fa}$	76
default	$\frac{A}{2fa(-i+\tan(fx+e))} + \frac{iB}{2fa(-i+\tan(fx+e))} + \frac{A \arctan(\tan(fx+e))}{2fa} - \frac{iB \arctan(\tan(fx+e))}{2fa}$	76
norman	$\frac{\frac{(-iB+A)x}{2a} - \frac{-iA+B}{2af} + \frac{(iB+A)\tan(fx+e)}{2af} + \frac{(-iB+A)x \tan(fx+e)^2}{2a}}{1+\tan(fx+e)^2}$	81

```
input int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output -1/2*I*x/a*B+1/2*x/a*A-1/4/a/f*exp(-2*I*(f*x+e))*B+1/4*I/a/f*exp(-2*I*(f*x
+e))*A
```

3.710.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.89

$$\int \frac{A + B \tan(e + fx)}{a + ia \tan(e + fx)} dx = \frac{(2(A - iB)fx e^{(2i fx + 2ie)} + iA - B)e^{(-2i fx - 2ie)}}{4af}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e)),x, algorithm="fricas")
```

```
output 1/4*(2*(A - I*B)*f*x*e^(2*I*f*x + 2*I*e) + I*A - B)*e^(-2*I*f*x - 2*I*e)/(
a*f)
```

3.710.6 Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.85

$$\int \frac{A + B \tan(e + fx)}{a + ia \tan(e + fx)} dx$$

$$= \begin{cases} \frac{(iA-B)e^{-2ie}e^{-2ifx}}{4af} & \text{for } afe^{2ie} \neq 0 \\ x\left(-\frac{A-iB}{2a} + \frac{(Ae^{2ie}+A-iBe^{2ie}+iB)e^{-2ie}}{2a}\right) & \text{otherwise} \end{cases} + \frac{x(A-iB)}{2a}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e)),x)`output `Piecewise(((I*A - B)*exp(-2*I*e)*exp(-2*I*f*x)/(4*a*f), Ne(a*f*exp(2*I*e), 0)), (x*(-(A - I*B)/(2*a) + (A*exp(2*I*e) + A - I*B*exp(2*I*e) + I*B)*exp(-2*I*e)/(2*a)), True)) + x*(A - I*B)/(2*a)`**3.710.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{a + ia \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`**3.710.8 Giac [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 83 vs. 2(35) = 70.

Time = 0.38 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.77

$$\int \frac{A + B \tan(e + fx)}{a + ia \tan(e + fx)} dx$$

$$= -\frac{\frac{(-iA-B)\log(\tan(fx+e)+i)}{a} + \frac{(iA+B)\log(\tan(fx+e)-i)}{a} + \frac{-iA\tan(fx+e)-B\tan(fx+e)-3A-iB}{a(\tan(fx+e)-i)}}{4f}$$

3.710. $\int \frac{A+B \tan(e+fx)}{a+ia \tan(e+fx)} dx$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

output `-1/4*((-I*A - B)*log(tan(f*x + e) + I)/a + (I*A + B)*log(tan(f*x + e) - I)/a + (-I*A*tan(f*x + e) - B*tan(f*x + e) - 3*A - I*B)/(a*(tan(f*x + e) - I)))/f`

3.710.9 Mupad [B] (verification not implemented)

Time = 8.37 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.96

$$\int \frac{A + B \tan(e + fx)}{a + ia \tan(e + fx)} dx = -\frac{x(B + A \operatorname{li}) \operatorname{li}}{2a} + \frac{-\frac{B}{2a} + \frac{A \operatorname{li}}{2a}}{f(1 + \tan(e + fx) \operatorname{li})}$$

input `int((A + B*tan(e + f*x))/(a + a*tan(e + f*x)*1i),x)`

output `((A*1i)/(2*a) - B/(2*a))/(f*(tan(e + f*x)*1i + 1)) - (x*(A*1i + B)*1i)/(2*a)`

3.711 $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))} dx$

3.711.1 Optimal result 6595
 3.711.2 Mathematica [A] (verified) 6595
 3.711.3 Rubi [A] (verified) 6596
 3.711.4 Maple [A] (verified) 6598
 3.711.5 Fricas [C] (verification not implemented) 6598
 3.711.6 Sympy [A] (verification not implemented) 6599
 3.711.7 Maxima [F(-2)] 6599
 3.711.8 Giac [A] (verification not implemented) 6600
 3.711.9 Mupad [B] (verification not implemented) 6600

3.711.1 Optimal result

Integrand size = 41, antiderivative size = 45

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))} dx = \frac{Ax}{2ac} - \frac{\cos^2(e + fx)(B - A \tan(e + fx))}{2acf}$$

output `1/2*A*x/a/c-1/2*cos(f*x+e)^2*(B-A*tan(f*x+e))/a/c/f`

3.711.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.96

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))} dx = \frac{-2B \cos^2(e + fx) + A(2(e + fx) + \sin(2(e + fx)))}{4acf}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])*(c - I*c*Tan[e + f*x])),x]`

output `(-2*B*Cos[e + f*x]^2 + A*(2*(e + f*x) + Sin[2*(e + f*x)]))/(4*a*c*f)`

3.711.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.11, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.146$, Rules used = {3042, 4071, 27, 82, 454, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A + B \tan(e + fx)}{a^2 c^2 (1 - i \tan(e + fx))^2 (i \tan(e + fx) + 1)^2} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{A + B \tan(e + fx)}{(1 - i \tan(e + fx))^2 (i \tan(e + fx) + 1)^2} d \tan(e + fx)}{acf} \\
 & \quad \downarrow \text{82} \\
 & \frac{\int \frac{A + B \tan(e + fx)}{(\tan^2(e + fx) + 1)^2} d \tan(e + fx)}{acf} \\
 & \quad \downarrow \text{454} \\
 & \frac{\frac{1}{2} A \int \frac{1}{\tan^2(e + fx) + 1} d \tan(e + fx) - \frac{B - A \tan(e + fx)}{2(\tan^2(e + fx) + 1)}}{acf} \\
 & \quad \downarrow \text{216} \\
 & \frac{\frac{1}{2} A \arctan(\tan(e + fx)) - \frac{B - A \tan(e + fx)}{2(\tan^2(e + fx) + 1)}}{acf}
 \end{aligned}$$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])*(c - I*c*Tan[e + f*x])),x]`

output `((A*ArcTan[Tan[e + f*x]])/2 - (B - A*Tan[e + f*x])/(2*(1 + Tan[e + f*x]^2)))/(a*c*f)`

3.711. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))} dx$

3.711.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 82 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 454 `Int[((c_) + (d_)*(x_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[((a*d - b*c*x)/(2*a*b*(p + 1)))*(a + b*x^2)^(p + 1), x] + Simp[c*((2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d}, x] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.711.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.20

method	result
risch	$\frac{Ax}{2ac} - \frac{B \cos(2fx+2e)}{4acf} + \frac{A \sin(2fx+2e)}{4acf}$
norman	$\frac{\frac{Ax}{2ac} - \frac{B}{2acf} + \frac{A \tan(fx+e)}{2acf} + \frac{Ax \tan(fx+e)^2}{2ac}}{1+\tan(fx+e)^2}$
derivativedivides	$\frac{A \arctan(\tan(fx+e))}{2fac} + \frac{A}{4fac(-i+\tan(fx+e))} + \frac{iB}{4fac(-i+\tan(fx+e))} + \frac{A}{4fac(i+\tan(fx+e))} - \frac{iB}{4fac(i+\tan(fx+e))}$
default	$\frac{A \arctan(\tan(fx+e))}{2fac} + \frac{A}{4fac(-i+\tan(fx+e))} + \frac{iB}{4fac(-i+\tan(fx+e))} + \frac{A}{4fac(i+\tan(fx+e))} - \frac{iB}{4fac(i+\tan(fx+e))}$

```
input int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 1/2*A*x/a/c-1/4*B/a/c/f*cos(2*f*x+2*e)+1/4*A/a/c/f*sin(2*f*x+2*e)
```

3.711.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.29

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))} dx$$

$$= \frac{(4Afxe^{2i fx+2ie}) + (-iA - B)e^{(4i fx+4ie)} + iA - B)e^{(-2i fx-2ie)}}{8acf}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e)),x, algorithm="fracas")
```

```
output 1/8*(4*A*f*x*e^(2*I*f*x + 2*I*e) + (-I*A - B)*e^(4*I*f*x + 4*I*e) + I*A - B)*e^(-2*I*f*x - 2*I*e)/(a*c*f)
```

3.711.6 Sympy [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 165, normalized size of antiderivative = 3.67

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))} dx$$

$$= \frac{Ax}{2ac} + \begin{cases} \frac{((8iAacf - 8Bacf)e^{-2ifx} + (-8iAacfe^{4ie} - 8Bacfe^{4ie})e^{2ifx})e^{-2ie}}{64a^2c^2f^2} & \text{for } a^2c^2f^2e^{2ie} \neq 0 \\ x\left(-\frac{A}{2ac} + \frac{(Ae^{4ie} + 2Ae^{2ie} + A - iBe^{4ie} + iB)e^{-2ie}}{4ac}\right) & \text{otherwise} \end{cases}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e)),x)`output `A*x/(2*a*c) + Piecewise((((8*I*A*a*c*f - 8*B*a*c*f)*exp(-2*I*f*x) + (-8*I*A*a*c*f*exp(4*I*e) - 8*B*a*c*f*exp(4*I*e))*exp(2*I*f*x))*exp(-2*I*e)/(64*a**2*c**2*f**2), Ne(a**2*c**2*f**2*exp(2*I*e), 0)), (x*(-A/(2*a*c) + (A*exp(4*I*e) + 2*A*exp(2*I*e) + A - I*B*exp(4*I*e) + I*B)*exp(-2*I*e)/(4*a*c)), True))`**3.711.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e)),x, algorithm="maxima")`output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.711.8 Giac [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.11

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))} dx = \frac{\frac{(fx+e)A}{ac} + \frac{A \tan(fx+e) - B}{(\tan(fx+e)^2 + 1)ac}}{2f}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e)),x, algorithm="giac")`

output `1/2*((f*x + e)*A/(a*c) + (A*tan(f*x + e) - B)/((tan(f*x + e)^2 + 1)*a*c))/f`

3.711.9 Mupad [B] (verification not implemented)

Time = 8.88 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.89

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))} dx = \frac{\frac{A \sin(2e+2fx)}{2} - \frac{B \cos(2e+2fx)}{2} + Afx}{2ac}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)*(c - c*tan(e + f*x)*1i)),x)`

output `((A*sin(2*e + 2*f*x))/2 - (B*cos(2*e + 2*f*x))/2 + A*f*x)/(2*a*c*f)`

3.712
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^2} dx$$

3.712.1 Optimal result 6601
 3.712.2 Mathematica [A] (verified) 6601
 3.712.3 Rubi [A] (verified) 6602
 3.712.4 Maple [A] (verified) 6604
 3.712.5 Fricas [A] (verification not implemented) 6604
 3.712.6 Sympy [A] (verification not implemented) 6605
 3.712.7 Maxima [F(-2)] 6605
 3.712.8 Giac [A] (verification not implemented) 6606
 3.712.9 Mupad [B] (verification not implemented) 6606

3.712.1 Optimal result

Integrand size = 41, antiderivative size = 113

$$\begin{aligned} & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^2} dx \\ &= \frac{(3A + iB)x}{8ac^2} - \frac{A + iB}{8ac^2 f(i - \tan(e + fx))} \\ & \quad + \frac{iA + B}{8ac^2 f(i + \tan(e + fx))^2} + \frac{A}{4ac^2 f(i + \tan(e + fx))} \end{aligned}$$

output `1/8*(3*A+I*B)*x/a/c^2+1/8*(-A-I*B)/a/c^2/f/(I-tan(f*x+e))+1/8*(I*A+B)/a/c^2/f/(I+tan(f*x+e))^2+1/4*A/a/c^2/f/(I+tan(f*x+e))`

3.712.2 Mathematica [A] (verified)

Time = 5.63 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.09

$$\begin{aligned} & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^2} dx \\ &= \frac{\sec^2(e + fx)(5A - iB - (A + 3iB) \cos(2(e + fx)) + 3iA \sin(2(e + fx)) - B \sin(2(e + fx)) + 2(3A + iB) \tan(e + fx))}{16ac^2 f(-i + \tan(e + fx))(i + \tan(e + fx))^2} \end{aligned}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^2),x]`

output $(\text{Sec}[e + f*x]^2*(5*A - I*B - (A + (3*I)*B)*\text{Cos}[2*(e + f*x)] + (3*I)*A*\text{Sin}[2*(e + f*x)] - B*\text{Sin}[2*(e + f*x)] + 2*(3*A + I*B)*\text{ArcTan}[\text{Tan}[e + f*x]]*(I + \text{Tan}[e + f*x]))/(16*a*c^2*f*(-I + \text{Tan}[e + f*x])*(I + \text{Tan}[e + f*x])^2)$

3.712.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.85, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^2} dx \\ & \quad \downarrow 3042 \\ & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^2} dx \\ & \quad \downarrow 4071 \\ & \frac{ac \int \frac{A + B \tan(e + fx)}{a^2 c^3 (1 - i \tan(e + fx))^3 (i \tan(e + fx) + 1)^2} d \tan(e + fx)}{f} \\ & \quad \downarrow 27 \\ & \frac{\int \frac{A + B \tan(e + fx)}{(1 - i \tan(e + fx))^3 (i \tan(e + fx) + 1)^2} d \tan(e + fx)}{ac^2 f} \\ & \quad \downarrow 86 \\ & \frac{\int \left(-\frac{A}{4(\tan(e + fx) + i)^2} + \frac{3A + iB}{8(\tan^2(e + fx) + 1)} + \frac{-A - iB}{8(\tan(e + fx) - i)^2} - \frac{i(A - iB)}{4(\tan(e + fx) + i)^3} \right) d \tan(e + fx)}{ac^2 f} \\ & \quad \downarrow 2009 \\ & \frac{\frac{1}{8}(3A + iB) \arctan(\tan(e + fx)) - \frac{A + iB}{8(-\tan(e + fx) + i)} + \frac{B + iA}{8(\tan(e + fx) + i)^2} + \frac{A}{4(\tan(e + fx) + i)}}{ac^2 f} \end{aligned}$$

input $\text{Int}[(A + B*\text{Tan}[e + f*x])/((a + I*a*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^2), x]$

output $((3A + I*B)*\text{ArcTan}[\text{Tan}[e + f*x]])/8 - (A + I*B)/(8*(I - \text{Tan}[e + f*x])) + (I*A + B)/(8*(I + \text{Tan}[e + f*x])^2) + A/(4*(I + \text{Tan}[e + f*x]))/(a*c^2*f)$

3.712.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.712.4 Maple [A] (verified)

Time = 0.19 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.15

method	result
risch	$\frac{ixB}{8ac^2} + \frac{3xA}{8ac^2} - \frac{e^{4i(fx+e)}B}{32ac^2f} - \frac{ie^{4i(fx+e)}A}{32ac^2f} - \frac{\cos(2fx+2e)B}{8ac^2f} - \frac{i\cos(2fx+2e)A}{8ac^2f} + \frac{A\sin(2fx+2e)}{4ac^2f}$
norman	$\frac{\frac{(iB+3A)x}{8ac} - \frac{iA+B}{4acf} + \frac{(iB+3A)\tan(fx+e)^3}{8acf} + \frac{(iB+3A)x\tan(fx+e)^2}{4ac} + \frac{(iB+3A)x\tan(fx+e)^4}{8ac} + \frac{(-iB+5A)\tan(fx+e)}{8acf}}{c(1+\tan(fx+e))^2}$
derivativedivides	$\frac{3A\arctan(\tan(fx+e))}{8fa^2c^2} + \frac{iB\arctan(\tan(fx+e))}{8fa^2c^2} + \frac{A}{4ac^2f(i+\tan(fx+e))} + \frac{iA}{8fa^2c^2(i+\tan(fx+e))^2} + \frac{1}{8fa^2c^2(i+\tan(fx+e))}$
default	$\frac{3A\arctan(\tan(fx+e))}{8fa^2c^2} + \frac{iB\arctan(\tan(fx+e))}{8fa^2c^2} + \frac{A}{4ac^2f(i+\tan(fx+e))} + \frac{iA}{8fa^2c^2(i+\tan(fx+e))^2} + \frac{1}{8fa^2c^2(i+\tan(fx+e))}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `1/8*I*x/a/c^2*B+3/8*x/a/c^2*A-1/32/a/c^2/f*exp(4*I*(f*x+e))*B-1/32*I/a/c^2/f*exp(4*I*(f*x+e))*A-1/8/a/c^2/f*cos(2*f*x+2*e)*B-1/8*I/a/c^2/f*cos(2*f*x+2*e)*A+1/4*A/a/c^2/f*sin(2*f*x+2*e)`

3.712.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.71

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^2} dx$$

$$= \frac{(4(3A + iB)fx e^{(2i fx + 2ie)} + (-iA - B)e^{(6i fx + 6ie)} - 2(3iA + B)e^{(4i fx + 4ie)} + 2iA - 2B)e^{(-2i fx - 2ie)}}{32ac^2f}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x, algorithm="fracas")`

output `1/32*(4*(3*A + I*B)*f*x*e^(2*I*f*x + 2*I*e) + (-I*A - B)*e^(6*I*f*x + 6*I*e) - 2*(3*I*A + B)*e^(4*I*f*x + 4*I*e) + 2*I*A - 2*B)*e^(-2*I*f*x - 2*I*e)/(a*c^2*f)`

3.712.6 Sympy [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 284, normalized size of antiderivative = 2.51

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^2} dx$$

$$= \begin{cases} \frac{((512iAa^2c^4f^2 - 512Ba^2c^4f^2)e^{-2ifx} + (-1536iAa^2c^4f^2e^{4ie} - 512Ba^2c^4f^2e^{4ie})e^{2ifx} + (-256iAa^2c^4f^2e^{6ie} - 256Ba^2c^4f^2e^{6ie})e^{4ifx})e^{-2ie}}{8192a^3c^6f^3} \\ x \left(-\frac{3A+iB}{8ac^2} + \frac{(Ae^{6ie} + 3Ae^{4ie} + 3Ae^{2ie} + A - iBe^{6ie} - iBe^{4ie} + iBe^{2ie} + iB)e^{-2ie}}{8ac^2} \right) \\ + \frac{x(3A + iB)}{8ac^2} \end{cases}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))**2,x)`

output `Piecewise((((512*I*A*a**2*c**4*f**2 - 512*B*a**2*c**4*f**2)*exp(-2*I*f*x) + (-1536*I*A*a**2*c**4*f**2*exp(4*I*e) - 512*B*a**2*c**4*f**2*exp(4*I*e))*exp(2*I*f*x) + (-256*I*A*a**2*c**4*f**2*exp(6*I*e) - 256*B*a**2*c**4*f**2*exp(6*I*e))*exp(4*I*f*x))*exp(-2*I*e)/(8192*a**3*c**6*f**3), Ne(a**3*c**6*f**3*exp(2*I*e), 0)), (x*(-(3*A + I*B)/(8*a*c**2) + (A*exp(6*I*e) + 3*A*exp(4*I*e) + 3*A*exp(2*I*e) + A - I*B*exp(6*I*e) - I*B*exp(4*I*e) + I*B*exp(2*I*e) + I*B)*exp(-2*I*e)/(8*a*c**2)), True)) + x*(3*A + I*B)/(8*a*c**2)`

3.712.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.712.8 Giac [A] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.41

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^2} dx$$

$$= \frac{\frac{2(3iA-B) \log(\tan(fx+e)+i)}{ac^2} + \frac{2(-3iA+B) \log(\tan(fx+e)-i)}{ac^2} - \frac{2(3A \tan(fx+e)+iB \tan(fx+e)-5iA+3B)}{ac^2(i \tan(fx+e)+1)} + \frac{-9iA \tan(fx+e)^2+3B}{32f}}{32f}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^2,x, algo
rithm="giac")
```

```
output 1/32*(2*(3*I*A - B)*log(tan(f*x + e) + I)/(a*c^2) + 2*(-3*I*A + B)*log(tan
(f*x + e) - I)/(a*c^2) - 2*(3*A*tan(f*x + e) + I*B*tan(f*x + e) - 5*I*A +
3*B)/(a*c^2*(I*tan(f*x + e) + 1)) + (-9*I*A*tan(f*x + e)^2 + 3*B*tan(f*x +
e)^2 + 26*A*tan(f*x + e) + 6*I*B*tan(f*x + e) + 21*I*A + B)/(a*c^2*(tan(f
*x + e) + I)^2))/f
```

3.712.9 Mupad [B] (verification not implemented)

Time = 9.06 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.14

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^2} dx$$

$$= \frac{\tan(e + fx) \left(-\frac{B}{8ac^2} + \frac{A3i}{8ac^2}\right) + \tan(e + fx)^2 \left(\frac{3A}{8ac^2} + \frac{B1i}{8ac^2}\right) + \frac{A}{4ac^2} - \frac{B1i}{4ac^2}}{f \left(\tan(e + fx)^3 + \tan(e + fx)^2 1i + \tan(e + fx) + 1i\right)}$$

$$- \frac{x(-B + A3i) 1i}{8ac^2}$$

```
input int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)*(c - c*tan(e + f*x)*1i)^
2),x)
```

```
output (tan(e + f*x)*((A*3i)/(8*a*c^2) - B/(8*a*c^2)) + tan(e + f*x)^2*((3*A)/(8*
a*c^2) + (B*1i)/(8*a*c^2)) + A/(4*a*c^2) - (B*1i)/(4*a*c^2))/(f*(tan(e + f
*x) + tan(e + f*x)^2*1i + tan(e + f*x)^3 + 1i)) - (x*(A*3i - B)*1i)/(8*a*c
^2)
```

3.713
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^3} dx$$

3.713.1 Optimal result 6607
 3.713.2 Mathematica [A] (verified) 6607
 3.713.3 Rubi [A] (verified) 6608
 3.713.4 Maple [A] (verified) 6610
 3.713.5 Fricas [A] (verification not implemented) 6610
 3.713.6 Sympy [A] (verification not implemented) 6611
 3.713.7 Maxima [F(-2)] 6611
 3.713.8 Giac [A] (verification not implemented) 6612
 3.713.9 Mupad [B] (verification not implemented) 6612

3.713.1 Optimal result

Integrand size = 41, antiderivative size = 149

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^3} dx$$

$$= \frac{(2A + iB)x}{8ac^3} - \frac{A + iB}{16ac^3 f(i - \tan(e + fx))} - \frac{A - iB}{12ac^3 f(i + \tan(e + fx))^3}$$

$$+ \frac{iA}{8ac^3 f(i + \tan(e + fx))^2} + \frac{3A + iB}{16ac^3 f(i + \tan(e + fx))}$$

output `1/8*(2*A+I*B)*x/a/c^3+1/16*(-A-I*B)/a/c^3/f/(I-tan(f*x+e))+1/12*(-A+I*B)/a/c^3/f/(I+tan(f*x+e))^3+1/8*I*A/a/c^3/f/(I+tan(f*x+e))^2+1/16*(3*A+I*B)/a/c^3/f/(I+tan(f*x+e))`

3.713.2 Mathematica [A] (verified)

Time = 5.81 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.00

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^3} dx$$

$$= \frac{\sec^3(e + fx)(9iA \cos(e + fx) + (-1 + 2 \cos(2(e + fx)))((-iA + 2B) \cos(e + fx) - (2A + iB) \sin(e + fx)))}{24ac^3 f(-i + \tan(e + fx))(i + \tan(e + fx))}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^3),x]`

3.713.
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^3} dx$$

output $(\text{Sec}[e + f*x]^3*((9*I)*A*\text{Cos}[e + f*x] + (-1 + 2*\text{Cos}[2*(e + f*x)])*((-I)*A + 2*B)*\text{Cos}[e + f*x] - (2*A + I*B)*\text{Sin}[e + f*x]) - 3*(2*A + I*B)*\text{ArcTan}[\text{Tan}[e + f*x]]*\text{Sec}[e + f*x]*(\text{Cos}[2*(e + f*x)] - I*\text{Sin}[2*(e + f*x)])))/(24*a*c^3*f*(-I + \text{Tan}[e + f*x])*(I + \text{Tan}[e + f*x])^3)$

3.713.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.83, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^3} dx$$

↓ 4071

$$\frac{ac \int \frac{A + B \tan(e + fx)}{a^2 c^4 (1 - i \tan(e + fx))^4 (i \tan(e + fx) + 1)^2} d \tan(e + fx)}{f}$$

↓ 27

$$\int \frac{A + B \tan(e + fx)}{(1 - i \tan(e + fx))^4 (i \tan(e + fx) + 1)^2} d \tan(e + fx)$$

$ac^3 f$

↓ 86

$$\int \left(-\frac{iA}{4(\tan(e + fx) + i)^3} + \frac{2A + iB}{8(\tan^2(e + fx) + 1)} + \frac{-A - iB}{16(\tan(e + fx) - i)^2} + \frac{-3A - iB}{16(\tan(e + fx) + i)^2} + \frac{A - iB}{4(\tan(e + fx) + i)^4} \right) d \tan(e + fx)$$

$ac^3 f$

↓ 2009

$$\frac{\frac{1}{8}(2A + iB) \arctan(\tan(e + fx)) - \frac{A + iB}{16(-\tan(e + fx) + i)} + \frac{3A + iB}{16(\tan(e + fx) + i)} - \frac{A - iB}{12(\tan(e + fx) + i)^3} + \frac{iA}{8(\tan(e + fx) + i)^2}}{ac^3 f}$$

3.713. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^3} dx$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^3), x]`

output `((2*A + I*B)*ArcTan[Tan[e + f*x]]/8 - (A + I*B)/(16*(I - Tan[e + f*x])) - (A - I*B)/(12*(I + Tan[e + f*x])^3) + ((I/8)*A)/(I + Tan[e + f*x])^2 + (3*A + I*B)/(16*(I + Tan[e + f*x])))/(a*c^3*f)`

3.713.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.713.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.31

method	result
risch	$\frac{ixB}{8ac^3} + \frac{xA}{4ac^3} - \frac{e^{6i(fx+e)}B}{96ac^3f} - \frac{ie^{6i(fx+e)}A}{96ac^3f} - \frac{e^{4i(fx+e)}B}{32ac^3f} - \frac{ie^{4i(fx+e)}A}{16ac^3f} - \frac{\cos(2fx+2e)B}{32ac^3f} - \frac{5i\cos(2fx+2e)A}{32ac^3f}$
derivativedivides	$\frac{iA}{8ac^3f(i+\tan(fx+e))^2} - \frac{A}{12fac^3(i+\tan(fx+e))^3} + \frac{iB}{12fac^3(i+\tan(fx+e))^3} + \frac{3A}{16fac^3(i+\tan(fx+e))} + \frac{3iB}{16fac^3(i+\tan(fx+e))}$
default	$\frac{iA}{8ac^3f(i+\tan(fx+e))^2} - \frac{A}{12fac^3(i+\tan(fx+e))^3} + \frac{iB}{12fac^3(i+\tan(fx+e))^3} + \frac{3A}{16fac^3(i+\tan(fx+e))} + \frac{3iB}{16fac^3(i+\tan(fx+e))}$
norman	$\frac{(iB+2A)x}{8ac} - \frac{4iA+B}{12acf} + \frac{B \tan(fx+e)^2}{4acf} + \frac{(-iB+6A) \tan(fx+e)}{8acf} + \frac{(iB+2A) \tan(fx+e)^3}{3acf} + \frac{(iB+2A) \tan(fx+e)^5}{8acf} + \frac{3(iB+2A)x \tan(fx+e)}{8ac}$ $\frac{\hspace{10em}}{(1+\tan(fx+e))^2} c^2$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output `1/8*I*x/a/c^3*B+1/4*x/a/c^3*A-1/96/a/c^3/f*exp(6*I*(f*x+e))*B-1/96*I/a/c^3/f*exp(6*I*(f*x+e))*A-1/32/a/c^3/f*exp(4*I*(f*x+e))*B-1/16*I/a/c^3/f*exp(4*I*(f*x+e))*A-1/32/a/c^3/f*cos(2*f*x+2*e)*B-5/32*I/a/c^3/f*cos(2*f*x+2*e)*A+1/32*I/a/c^3/f*sin(2*f*x+2*e)*B+7/32/a/c^3/f*sin(2*f*x+2*e)*A`

3.713.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.62

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^3} dx$$

$$= \frac{(12(2A + iB)fxe^{(2i fx+2ie)} + (-iA - B)e^{(8i fx+8ie)} - 3(2iA + B)e^{(6i fx+6ie)} - 18iAe^{(4i fx+4ie)} + 3iA - 3B)e^{(-2i fx - 2ie)}}{96ac^3f}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x, algorith="fracas")`

output `1/96*(12*(2*A + I*B)*f*x*e^(2*I*f*x + 2*I*e) + (-I*A - B)*e^(8*I*f*x + 8*I*e) - 3*(2*I*A + B)*e^(6*I*f*x + 6*I*e) - 18*I*A*e^(4*I*f*x + 4*I*e) + 3*I*A - 3*B)*e^(-2*I*f*x - 2*I*e)/(a*c^3*f)`

3.713.6 Sympy [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 328, normalized size of antiderivative = 2.20

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^3} dx$$

$$= \left\{ \frac{(-294912iAa^3c^9f^3e^{4ie}e^{2ifx} + (49152iAa^3c^9f^3 - 49152Ba^3c^9f^3)e^{-2ifx} + (-98304iAa^3c^9f^3e^{6ie} - 49152Ba^3c^9f^3e^{6ie})e^{4ifx} + (-16384iAa^3c^9f^3e^{8ie} - 16384Ba^3c^9f^3e^{8ie})e^{2ifx} + (-16384iAa^3c^9f^3e^{10ie} - 16384Ba^3c^9f^3e^{10ie})e^{0ifx})}{1572864a^4c^{12}f^4} \right.$$

$$\left. x \left(-\frac{2A+iB}{8ac^3} + \frac{(Ae^{8ie} + 4Ae^{6ie} + 6Ae^{4ie} + 4Ae^{2ie} + A - iBe^{8ie} - 2iBe^{6ie} + 2iBe^{2ie} + iB)e^{-2ie}}{16ac^3} \right) \right.$$

$$\left. + \frac{x(2A + iB)}{8ac^3} \right.$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))**3,x)
```

```
output Piecewise((( -294912*I*A*a**3*c**9*f**3*exp(4*I*e)*exp(2*I*f*x) + (49152*I*
A*a**3*c**9*f**3 - 49152*B*a**3*c**9*f**3)*exp(-2*I*f*x) + (-98304*I*A*a**
3*c**9*f**3*exp(6*I*e) - 49152*B*a**3*c**9*f**3*exp(6*I*e))*exp(4*I*f*x) +
(-16384*I*A*a**3*c**9*f**3*exp(8*I*e) - 16384*B*a**3*c**9*f**3*exp(8*I*e)
)*exp(6*I*f*x))*exp(-2*I*e)/(1572864*a**4*c**12*f**4), Ne(a**4*c**12*f**4*
exp(2*I*e), 0)), (x*(-(2*A + I*B)/(8*a*c**3) + (A*exp(8*I*e) + 4*A*exp(6*I
*e) + 6*A*exp(4*I*e) + 4*A*exp(2*I*e) + A - I*B*exp(8*I*e) - 2*I*B*exp(6*I
*e) + 2*I*B*exp(2*I*e) + I*B)*exp(-2*I*e)/(16*a*c**3)), True)) + x*(2*A +
I*B)/(8*a*c**3)
```

3.713.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x, algo
rithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.713.8 Giac [A] (verification not implemented)

Time = 0.61 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.21

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^3} dx =$$

$$\frac{6(-2iA+B)\log(\tan(fx+e)+i)}{ac^3} + \frac{6(2iA-B)\log(\tan(fx+e)-i)}{ac^3} + \frac{6(-2iA\tan(fx+e)+B\tan(fx+e)-3A-2iB)}{ac^3(\tan(fx+e)-i)} + \frac{22iA\tan(fx+e)^3}{ac^3(\tan(fx+e)+i)}$$

96 f

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^3,x, algo
rithm="giac")
```

```
output -1/96*(6*(-2*I*A + B)*log(tan(f*x + e) + I)/(a*c^3) + 6*(2*I*A - B)*log(ta
n(f*x + e) - I)/(a*c^3) + 6*(-2*I*A*tan(f*x + e) + B*tan(f*x + e) - 3*A -
2*I*B)/(a*c^3*(tan(f*x + e) - I)) + (22*I*A*tan(f*x + e)^3 - 11*B*tan(f*x
+ e)^3 - 84*A*tan(f*x + e)^2 - 39*I*B*tan(f*x + e)^2 - 114*I*A*tan(f*x + e
) + 45*B*tan(f*x + e) + 60*A + 9*I*B)/(a*c^3*(tan(f*x + e) + I)^3))/f
```

3.713.9 Mupad [B] (verification not implemented)

Time = 9.24 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.08

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^3} dx$$

$$= \frac{B}{12ac^3} + \tan(e + fx)^2 \left(-\frac{B}{4ac^3} + \frac{A1i}{2ac^3} \right) + \tan(e + fx)^3 \left(\frac{A}{4ac^3} + \frac{B1i}{8ac^3} \right) - \tan(e + fx) \left(\frac{A}{12ac^3} + \frac{B1i}{24ac^3} \right) + \frac{A}{3ac^3}$$

$$\frac{x(-B + A2i)1i}{8ac^3}$$

```
input int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)*(c - c*tan(e + f*x)*1i)^
3),x)
```

```
output (tan(e + f*x)^2*((A*1i)/(2*a*c^3) - B/(4*a*c^3)) - tan(e + f*x)*(A/(12*a*c
^3) + (B*1i)/(24*a*c^3)) + tan(e + f*x)^3*(A/(4*a*c^3) + (B*1i)/(8*a*c^3))
+ (A*1i)/(3*a*c^3) + B/(12*a*c^3))/(f*(tan(e + f*x)*2i + tan(e + f*x)^3*2
i + tan(e + f*x)^4 - 1)) - (x*(A*2i - B)*1i)/(8*a*c^3)
```

$$3.714 \quad \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^4} dx$$

3.714.1 Optimal result	6613
3.714.2 Mathematica [A] (verified)	6613
3.714.3 Rubi [A] (verified)	6614
3.714.4 Maple [A] (verified)	6616
3.714.5 Fricas [A] (verification not implemented)	6616
3.714.6 Sympy [A] (verification not implemented)	6617
3.714.7 Maxima [F(-2)]	6617
3.714.8 Giac [A] (verification not implemented)	6618
3.714.9 Mupad [B] (verification not implemented)	6618

3.714.1 Optimal result

Integrand size = 41, antiderivative size = 181

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^4} dx$$

$$= \frac{(5A + 3iB)x}{32ac^4} - \frac{A + iB}{32ac^4 f(i - \tan(e + fx))} - \frac{iA + B}{16ac^4 f(i + \tan(e + fx))^4}$$

$$- \frac{A}{12ac^4 f(i + \tan(e + fx))^3} + \frac{3iA - B}{32ac^4 f(i + \tan(e + fx))^2} + \frac{2A + iB}{16ac^4 f(i + \tan(e + fx))}$$

```
output 1/32*(5*A+3*I*B)*x/a/c^4+1/32*(-A-I*B)/a/c^4/f/(I-tan(f*x+e))+1/16*(-I*A-B)
)/a/c^4/f/(I+tan(f*x+e))^4-1/12*A/a/c^4/f/(I+tan(f*x+e))^3+1/32*(3*I*A-B)/
a/c^4/f/(I+tan(f*x+e))^2+1/16*(2*A+I*B)/a/c^4/f/(I+tan(f*x+e))
```

3.714.2 Mathematica [A] (verified)

Time = 3.78 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.78

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^4} dx$$

$$= \frac{i \left(3(-5iA + 3B) \arctan(\tan(e + fx)) + \frac{32iA+3(5A+3iB) \tan(e+fx)+7i(5A+3iB) \tan^2(e+fx)+9(5A+3iB) \tan^3(e+fx)+3(-i+\tan(e+fx))(i+\tan(e+fx))^4}{(-i+\tan(e+fx))(i+\tan(e+fx))^4} \right)}{96ac^4 f}$$

```
input Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])*(c - I*c*Tan[e + f*
x]))^4, x]
```

3.714. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^4} dx$

output $((I/96)*(3*((-5*I)*A + 3*B)*ArcTan[Tan[e + f*x]] + ((32*I)*A + 3*(5*A + (3*I)*B)*Tan[e + f*x] + (7*I)*(5*A + (3*I)*B)*Tan[e + f*x]^2 + 9*(5*A + (3*I)*B)*Tan[e + f*x]^3 + 3*((-5*I)*A + 3*B)*Tan[e + f*x]^4)/((-I + Tan[e + f*x])*(I + Tan[e + f*x])^4))/(a*c^4*f)$

3.714.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.81, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^4} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^4} dx$$

↓ 4071

$$\frac{ac \int \frac{A + B \tan(e + fx)}{a^2 c^5 (1 - i \tan(e + fx))^5 (i \tan(e + fx) + 1)^2} d \tan(e + fx)}{f}$$

↓ 27

$$\int \frac{A + B \tan(e + fx)}{(1 - i \tan(e + fx))^5 (i \tan(e + fx) + 1)^2} d \tan(e + fx)$$

$ac^4 f$

↓ 86

$$\int \left(\frac{A}{4(\tan(e + fx) + i)^4} + \frac{5A + 3iB}{32(\tan^2(e + fx) + 1)} + \frac{-A - iB}{32(\tan(e + fx) - i)^2} + \frac{-2A - iB}{16(\tan(e + fx) + i)^2} + \frac{B - 3iA}{16(\tan(e + fx) + i)^3} + \frac{iA + B}{4(\tan(e + fx) + i)^5} \right) d \tan(e + fx)$$

$ac^4 f$

↓ 2009

$$\frac{\frac{1}{32}(5A + 3iB) \arctan(\tan(e + fx)) - \frac{A + iB}{32(-\tan(e + fx) + i)} + \frac{2A + iB}{16(\tan(e + fx) + i)} + \frac{-B + 3iA}{32(\tan(e + fx) + i)^2} - \frac{B + iA}{16(\tan(e + fx) + i)^4} - \frac{1}{12(\tan(e + fx) + i)^6}}{ac^4 f}$$

3.714. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^4} dx$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^4), x]`

output `((5*A + (3*I)*B)*ArcTan[Tan[e + f*x]]/32 - (A + I*B)/(32*(I - Tan[e + f*x])) - (I*A + B)/(16*(I + Tan[e + f*x])^4) - A/(12*(I + Tan[e + f*x])^3) + ((3*I)*A - B)/(32*(I + Tan[e + f*x])^2) + (2*A + I*B)/(16*(I + Tan[e + f*x]))) / (a*c^4*f)`

3.714.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.714.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.31

method	result
risch	$\frac{3ixB}{32ac^4} + \frac{5xA}{32ac^4} - \frac{e^{8i(fx+e)}B}{256ac^4f} - \frac{ie^{8i(fx+e)}A}{256ac^4f} - \frac{e^{6i(fx+e)}B}{64ac^4f} - \frac{5ie^{6i(fx+e)}A}{192ac^4f} - \frac{e^{4i(fx+e)}B}{64ac^4f} - \frac{5ie^{4i(fx+e)}A}{64ac^4f}$
derivativedivides	$-\frac{A}{12ac^4f(i+\tan(fx+e))^3} + \frac{3iB \arctan(\tan(fx+e))}{32fa c^4} + \frac{A}{8fa c^4(i+\tan(fx+e))} + \frac{iB}{32fa c^4(-i+\tan(fx+e))} + \frac{5A}{12ac^4f(i+\tan(fx+e))^3}$
default	$-\frac{A}{12ac^4f(i+\tan(fx+e))^3} + \frac{3iB \arctan(\tan(fx+e))}{32fa c^4} + \frac{A}{8fa c^4(i+\tan(fx+e))} + \frac{iB}{32fa c^4(-i+\tan(fx+e))} + \frac{5A}{12ac^4f(i+\tan(fx+e))^3}$
norman	$\frac{(3iB+5A)x}{32ac} - \frac{iA}{3acf} + \frac{(iA+3B)\tan(fx+e)^2}{6acf} + \frac{11(3iB+5A)\tan(fx+e)^5}{96acf} + \frac{(3iB+5A)\tan(fx+e)^7}{32acf} + \frac{(3iB+5A)x \tan(fx+e)^2}{8ac} + \frac{3(3iB+5A)}{12ac^4f(i+\tan(fx+e))^3}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x,method=_RETURNVERBOSE)`

output `3/32*I*x/a/c^4*B+5/32*x/a/c^4*A-1/256/a/c^4/f*exp(8*I*(f*x+e))*B-1/256*I/a/c^4/f*exp(8*I*(f*x+e))*A-1/64/a/c^4/f*exp(6*I*(f*x+e))*B-5/192*I/a/c^4/f*exp(6*I*(f*x+e))*A-1/64/a/c^4/f*exp(4*I*(f*x+e))*B-5/64*I/a/c^4/f*exp(4*I*(f*x+e))*A+1/64/a/c^4/f*cos(2*f*x+2*e)*B-9/64*I/a/c^4/f*cos(2*f*x+2*e))*A+3/64*I/a/c^4/f*sin(2*f*x+2*e)*B+11/64/a/c^4/f*sin(2*f*x+2*e))*A`

3.714.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.64

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^4} dx = \frac{(24(5A + 3iB)fx e^{(2i fx + 2i e)} - 3(iA + B)e^{(10i fx + 10i e)} - 4(5iA + 3B)e^{(8i fx + 8i e)} - 12(5iA + B)e^{(6i fx + 6i e)} + 12iA - 12iB)e^{(-2i fx - 2i e)}}{768 ac^4 f}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x, algorith="fracas")`

output `1/768*(24*(5*A + 3*I*B)*f*x*e^(2*I*f*x + 2*I*e) - 3*(I*A + B)*e^(10*I*f*x + 10*I*e) - 4*(5*I*A + 3*B)*e^(8*I*f*x + 8*I*e) - 12*(5*I*A + B)*e^(6*I*f*x + 6*I*e) - 24*(5*I*A - B)*e^(4*I*f*x + 4*I*e) + 12*I*A - 12*I*B)*e^(-2*I*f*x - 2*I*e)/(a*c^4*f)`

3.714. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^4} dx$

3.714.6 Sympy [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 439, normalized size of antiderivative = 2.43

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^4} dx$$

$$= \left\{ \begin{array}{l} \left((100663296iAa^4c^{16}f^4 - 100663296Ba^4c^{16}f^4)e^{-2ifx} + (-1006632960iAa^4c^{16}f^4e^{4ie} + 201326592Ba^4c^{16}f^4e^{4ie})e^{2ifx} + (-503316480iAa^4c^{16}f^4e^{6ie} + 1006632960iAa^4c^{16}f^4e^{6ie})e^{4ifx} + (-167772160iAa^4c^{16}f^4e^{8ie} + 1006632960iAa^4c^{16}f^4e^{8ie})e^{2ifx} + (-25165824iAa^4c^{16}f^4e^{10ie} - 251658240iAa^4c^{16}f^4e^{10ie})e^{ifx} \right) \exp(-2ie) \\ x \left(-\frac{5A+3iB}{32ac^4} + \frac{(Ae^{10ie}+5Ae^{8ie}+10Ae^{6ie}+10Ae^{4ie}+5Ae^{2ie}+A-iBe^{10ie}-3iBe^{8ie}-2iBe^{6ie}+2iBe^{4ie}+3iBe^{2ie}+iB)e^{-2ie}}{32ac^4} \right) \\ + \frac{x(5A+3iB)}{32ac^4} \end{array} \right.$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))**4,x)`

output `Piecewise((((100663296*I*A*a**4*c**16*f**4 - 100663296*B*a**4*c**16*f**4)*exp(-2*I*f*x) + (-1006632960*I*A*a**4*c**16*f**4*exp(4*I*e) + 201326592*B*a**4*c**16*f**4*exp(4*I*e))*exp(2*I*f*x) + (-503316480*I*A*a**4*c**16*f**4*exp(6*I*e) - 100663296*B*a**4*c**16*f**4*exp(6*I*e))*exp(4*I*f*x) + (-167772160*I*A*a**4*c**16*f**4*exp(8*I*e) - 100663296*B*a**4*c**16*f**4*exp(8*I*e))*exp(6*I*f*x) + (-25165824*I*A*a**4*c**16*f**4*exp(10*I*e) - 25165824*B*a**4*c**16*f**4*exp(10*I*e))*exp(8*I*f*x))*exp(-2*I*e)/(6442450944*a**5*c**20*f**5), Ne(a**5*c**20*f**5*exp(2*I*e), 0)), (x*(-(5*A + 3*I*B)/(32*a*c**4) + (A*exp(10*I*e) + 5*A*exp(8*I*e) + 10*A*exp(6*I*e) + 10*A*exp(4*I*e) + 5*A*exp(2*I*e) + A - I*B*exp(10*I*e) - 3*I*B*exp(8*I*e) - 2*I*B*exp(6*I*e) + 2*I*B*exp(4*I*e) + 3*I*B*exp(2*I*e) + I*B)*exp(-2*I*e)/(32*a*c**4)), True)) + x*(5*A + 3*I*B)/(32*a*c**4)`

3.714.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^4} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x, algo rithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.714. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^4} dx$

3.714.8 Giac [A] (verification not implemented)

Time = 0.82 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.14

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^4} dx$$

$$= \frac{12(5iA - 3B) \log(\tan(fx + e) + i)}{ac^4} + \frac{12(-5iA + 3B) \log(\tan(fx + e) - i)}{ac^4} + \frac{12(5A \tan(fx + e) + 3iB \tan(fx + e) - 7iA + 5B)}{ac^4(-i \tan(fx + e) - 1)} + \frac{-125iA \tan(fx + e)^4 + 75B \tan(fx + e)^4 + 596A \tan(fx + e)^3 + 348iB \tan(fx + e)^3 + 1110iA \tan(fx + e)^2 - 618B \tan(fx + e)^2 - 996A \tan(fx + e) - 492iB \tan(fx + e) - 405iA + 99B}{ac^4(\tan(fx + e) + i)^4} / f$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^4,x, algorithm="giac")
```

```
output 1/768*(12*(5*I*A - 3*B)*log(tan(f*x + e) + I)/(a*c^4) + 12*(-5*I*A + 3*B)*log(tan(f*x + e) - I)/(a*c^4) + 12*(5*A*tan(f*x + e) + 3*I*B*tan(f*x + e) - 7*I*A + 5*B)/(a*c^4*(-I*tan(f*x + e) - 1)) + (-125*I*A*tan(f*x + e)^4 + 75*B*tan(f*x + e)^4 + 596*A*tan(f*x + e)^3 + 348*I*B*tan(f*x + e)^3 + 1110*I*A*tan(f*x + e)^2 - 618*B*tan(f*x + e)^2 - 996*A*tan(f*x + e) - 492*I*B*tan(f*x + e) - 405*I*A + 99*B)/(a*c^4*(tan(f*x + e) + I)^4))/f
```

3.714.9 Mupad [B] (verification not implemented)

Time = 9.83 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.13

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^4} dx =$$

$$-\frac{\tan(e + fx) \left(-\frac{3B}{32ac^4} + \frac{A5i}{32ac^4}\right) + \tan(e + fx)^4 \left(\frac{5A}{32ac^4} + \frac{B3i}{32ac^4}\right) + \tan(e + fx)^3 \left(-\frac{9B}{32ac^4} + \frac{A15i}{32ac^4}\right) - \tan(e + fx)^2 \left(\frac{7A}{32ac^4} + \frac{B7i}{32ac^4}\right) + \tan(e + fx) \left(-\frac{3B}{32ac^4} + \frac{A5i}{32ac^4}\right) + \frac{1}{32ac^4}}{f \left(-\tan(e + fx)^5 - \tan(e + fx)^4 3i + 2 \tan(e + fx)^3 - \tan(e + fx)^2 2i + 3 \tan(e + fx) + 1\right)}$$

```
input int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)*(c - c*tan(e + f*x)*1i)^4),x)
```

```
output - (tan(e + f*x)*((A*5i)/(32*a*c^4) - (3*B)/(32*a*c^4)) + tan(e + f*x)^4*((5*A)/(32*a*c^4) + (B*3i)/(32*a*c^4)) + tan(e + f*x)^3*((A*15i)/(32*a*c^4) - (9*B)/(32*a*c^4)) - tan(e + f*x)^2*((35*A)/(96*a*c^4) + (B*7i)/(32*a*c^4)) - A/(3*a*c^4))/(f*(3*tan(e + f*x) - tan(e + f*x)^2*2i + 2*tan(e + f*x)^3 - tan(e + f*x)^4*3i - tan(e + f*x)^5 + 1i)) - (x*(A*5i - 3*B)*1i)/(32*a*c^4)
```

3.714. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^4} dx$

$$3.715 \quad \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^n}{(a+ia \tan(e+fx))^2} dx$$

3.715.1 Optimal result	6619
3.715.2 Mathematica [A] (verified)	6619
3.715.3 Rubi [A] (verified)	6620
3.715.4 Maple [F]	6622
3.715.5 Fricas [F]	6622
3.715.6 Sympy [F]	6622
3.715.7 Maxima [F(-2)]	6623
3.715.8 Giac [F]	6623
3.715.9 Mupad [F(-1)]	6623

3.715.1 Optimal result

Integrand size = 41, antiderivative size = 115

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{(iA(2 - n) + B(2 + n)) \operatorname{Hypergeometric2F1}\left(2, n, 1 + n, \frac{1}{2}(1 - i \tan(e + fx))\right) (c - ic \tan(e + fx))^n}{16a^2fn} + \frac{(iA - B)(c - ic \tan(e + fx))^n}{4a^2f(1 + i \tan(e + fx))^2}$$

output

```
1/16*(I*A*(2-n)+B*(2+n))*hypergeom([2, n],[1+n],1/2-1/2*I*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/a^2/f/n+1/4*(I*A-B)*(c-I*c*tan(f*x+e))^n/a^2/f/(1+I*tan(f*x+e))^2
```

3.715.2 Mathematica [A] (verified)

Time = 2.48 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.32

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{(c - ic \tan(e + fx))^n \left(-iA(-3 + n) + B(1 + n) + \frac{i(-1+n)(A(-2+n)+iB(2+n)) \operatorname{Hypergeometric2F1}\left(1, n, 1+n, -\frac{1}{2}i(i+\tan(e+fx))\right)}{n} \right)}{16a^2f}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^n)/(a + I*a*Tan[e + f*x])^2,x]`

output `((c - I*c*Tan[e + f*x])^n*((-I)*A*(-3 + n) + B*(1 + n) + (I*(-1 + n)*(A*(-2 + n) + I*B*(2 + n))*Hypergeometric2F1[1, n, 1 + n, (-1/2*I)*(I + Tan[e + f*x])])/n + ((-I)*A*(-3 + n) + B*(1 + n))*(Cos[2*(e + f*x)] - I*Sin[2*(e + f*x)]) - (2*(A + I*B)*(I + Tan[e + f*x]))/(-I + Tan[e + f*x])^2))/(16*a^2*f)`

3.715.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 87, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{n-1}}{a^3(i \tan(e+fx)+1)^3} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{c \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{n-1}}{(i \tan(e+fx)+1)^3} d \tan(e + fx)}{a^2 f} \\
 & \quad \downarrow \text{87} \\
 & \frac{c \left(\frac{1}{4} (A(2-n) - iB(n+2)) \int \frac{(c-ic \tan(e+fx))^{n-1}}{(i \tan(e+fx)+1)^2} d \tan(e + fx) + \frac{(-B+iA)(c-ic \tan(e+fx))^n}{4c(1+i \tan(e+fx))^2} \right)}{a^2 f} \\
 & \quad \downarrow \text{78}
 \end{aligned}$$

3.715. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^n}{(a+ia \tan(e+fx))^2} dx$

$$\frac{c \left(\frac{i(A(2-n) - iB(n+2))(c - i \tan(e+fx))^n \operatorname{Hypergeometric2F1}(2, n, n+1, \frac{1}{2}(1 - i \tan(e+fx)))}{16cn} + \frac{(-B+iA)(c - i \tan(e+fx))^n}{4c(1+i \tan(e+fx))^2} \right)}{a^2 f}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^n)/(a + I*a*Tan[e + f*x])^2,x]`

output `(c*(((I/16)*(A*(2 - n) - I*B*(2 + n))*Hypergeometric2F1[2, n, 1 + n, (1 - I*Tan[e + f*x])/2]*(c - I*c*Tan[e + f*x])^n)/(c*n) + ((I*A - B)*(c - I*c*Tan[e + f*x])^n)/(4*c*(1 + I*Tan[e + f*x])^2)))/(a^2*f)`

3.715.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 78 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[(b*c - a*d)^n*(a + b*x)^(m + 1)/(b^(n + 1)*(m + 1))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.715.4 Maple [F]

$$\int \frac{(A + B \tan(fx + e))(c - ic \tan(fx + e))^n}{(a + ia \tan(fx + e))^2} dx$$

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e))^2,x)`

output `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e))^2,x)`

3.715.5 Fracas [F]

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^2} dx \\ &= \int \frac{(B \tan(fx + e) + A)(-ic \tan(fx + e) + c)^n}{(ia \tan(fx + e) + a)^2} dx \end{aligned}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e))^2,x, algorithm="fracas")`

output `integral(1/4*((A - I*B)*e^(4*I*f*x + 4*I*e) + 2*A*e^(2*I*f*x + 2*I*e) + A + I*B)*(2*c/(e^(2*I*f*x + 2*I*e) + 1))^n*e^(-4*I*f*x - 4*I*e)/a^2, x)`

3.715.6 Sympy [F]

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^2} dx \\ &= -\frac{\int \frac{A(-ic \tan(e+fx)+c)^n}{\tan^2(e+fx)-2i \tan(e+fx)-1} dx + \int \frac{B(-ic \tan(e+fx)+c)^n \tan(e+fx)}{\tan^2(e+fx)-2i \tan(e+fx)-1} dx}{a^2} \end{aligned}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e))^2,x)`

output `-(Integral(A*(-I*c*tan(e + f*x) + c)**n/(tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x) + Integral(B*(-I*c*tan(e + f*x) + c)**n*tan(e + f*x)/(tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x))/a**2`

3.715.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^n}{(a + i a \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.715.8 Giac [F]

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^n}{(a + i a \tan(e + fx))^2} dx \\ &= \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^n}{(i a \tan(fx + e) + a)^2} dx \end{aligned}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^n/(I*a*tan(f*x + e) + a)^2, x)`

3.715.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^n}{(a + i a \tan(e + fx))^2} dx \\ &= \int \frac{(A + B \tan(e + fx)) (c - c \tan(e + fx) li)^n}{(a + a \tan(e + fx) li)^2} dx \end{aligned}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*li)^n)/(a + a*tan(e + f*x)*li)^2,x)`

output `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^n)/(a + a*tan(e + f*x)*1i)^2, x)`

3.715. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^n}{(a+ia \tan(e+fx))^2} dx$

3.716
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^5}{(a+ia \tan(e+fx))^2} dx$$

3.716.1 Optimal result 6625
 3.716.2 Mathematica [A] (verified) 6626
 3.716.3 Rubi [A] (verified) 6626
 3.716.4 Maple [A] (verified) 6628
 3.716.5 Fricas [A] (verification not implemented) 6629
 3.716.6 Sympy [A] (verification not implemented) 6629
 3.716.7 Maxima [F(-2)] 6630
 3.716.8 Giac [B] (verification not implemented) 6630
 3.716.9 Mupad [B] (verification not implemented) 6631

3.716.1 Optimal result

Integrand size = 41, antiderivative size = 194

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^5}{(a + ia \tan(e + fx))^2} dx \\ &= \frac{8(3A + 7iB)c^5 x}{a^2} + \frac{8(3iA - 7B)c^5 \log(\cos(e + fx))}{a^2 f} - \frac{8(iA - B)c^5}{a^2 f (i - \tan(e + fx))^2} \\ &+ \frac{16(2A + 3iB)c^5}{a^2 f (i - \tan(e + fx))} - \frac{(7A + 24iB)c^5 \tan(e + fx)}{a^2 f} \\ &+ \frac{(iA - 7B)c^5 \tan^2(e + fx)}{2a^2 f} + \frac{iBc^5 \tan^3(e + fx)}{3a^2 f} \end{aligned}$$

output `8*(3*A+7*I*B)*c^5*x/a^2+8*(3*I*A-7*B)*c^5*ln(cos(f*x+e))/a^2/f-8*(I*A-B)*c^5/a^2/f/(I-tan(f*x+e))^2+16*(2*A+3*I*B)*c^5/a^2/f/(I-tan(f*x+e))-(7*A+24*I*B)*c^5*tan(f*x+e)/a^2/f+1/2*(I*A-7*B)*c^5*tan(f*x+e)^2/a^2/f+1/3*I*B*c^5*tan(f*x+e)^3/a^2/f`

3.716.2 Mathematica [A] (verified)

Time = 6.12 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.02

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^5}{(a + i a \tan(e + fx))^2} dx$$

$$= \frac{c^5(315iA - 737B + 48i(3A + 7iB) \log(i - \tan(e + fx)) - 2(246A + 569iB + 48(3A + 7iB) \log(i - \tan(e + fx)))}{(a + i a \tan(e + fx))^2}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^5)/(a + I*a*Tan[e + f*x])^2,x]`

output `(c^5*((315*I)*A - 737*B + (48*I)*(3*A + (7*I)*B)*Log[I - Tan[e + f*x]] - 2*(246*A + (569*I)*B + 48*(3*A + (7*I)*B)*Log[I - Tan[e + f*x]])*Tan[e + f*x] + 2*(5*((-9*I)*A + 23*B) + 24*((-3*I)*A + 7*B)*Log[I - Tan[e + f*x]])*Tan[e + f*x]^2 - 4*(9*A + (26*I)*B)*Tan[e + f*x]^3 + ((3*I)*A - 17*B)*Tan[e + f*x]^4 + (2*I)*B*Tan[e + f*x]^5)/(6*a^2*f*(-I + Tan[e + f*x])^2)`

3.716.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.71, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - i c \tan(e + fx))^5 (A + B \tan(e + fx))}{(a + i a \tan(e + fx))^2} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(c - i c \tan(e + fx))^5 (A + B \tan(e + fx))}{(a + i a \tan(e + fx))^2} dx$$

$$\downarrow \text{4071}$$

$$\frac{ac \int \frac{c^4(1-i \tan(e+fx))^4(A+B \tan(e+fx))}{a^3(i \tan(e+fx)+1)^3} d \tan(e + fx)}{f}$$

$$\downarrow \text{27}$$

3.716. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^5}{(a+ia \tan(e+fx))^2} dx$

$$\frac{c^5 \int \frac{(1-i \tan(e+fx))^4 (A+B \tan(e+fx))}{(i \tan(e+fx)+1)^3} d \tan(e+fx)}{a^2 f}$$

↓ 86

$$\frac{c^5 \int \left(iB \tan^2(e+fx) + i(A+7iB) \tan(e+fx) - 7A \left(\frac{24iB}{7A} + 1 \right) - \frac{8i(3A+7iB)}{\tan(e+fx)-i} + \frac{16(2A+3iB)}{(\tan(e+fx)-i)^2} + \frac{16i(A+iB)}{(\tan(e+fx)-i)^3} \right) dx}{a^2 f}$$

↓ 2009

$$\frac{c^5 \left(\frac{1}{2}(-7B+iA) \tan^2(e+fx) - (7A+24iB) \tan(e+fx) + \frac{16(2A+3iB)}{-\tan(e+fx)+i} - \frac{8(-B+iA)}{(-\tan(e+fx)+i)^2} - 8(-7B+3iA) \log \right)}{a^2 f}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^5)/(a + I*a*Tan[e + f*x])^2,x]`

output `(c^5*(-8*((3*I)*A - 7*B)*Log[I - Tan[e + f*x]] - (8*(I*A - B))/(I - Tan[e + f*x])^2 + (16*(2*A + (3*I)*B))/(I - Tan[e + f*x]) - (7*A + (24*I)*B)*Tan[e + f*x] + ((I*A - 7*B)*Tan[e + f*x]^2)/2 + (I/3)*B*Tan[e + f*x]^3))/(a^2*f)`

3.716.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.716. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^5}{(a+ia \tan(e+fx))^2} dx$

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.716.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.38

method	result
risch	$\frac{20c^5 e^{-2i(fx+e)} B}{a^2 f} - \frac{12ic^5 e^{-2i(fx+e)} A}{a^2 f} - \frac{2c^5 e^{-4i(fx+e)} B}{a^2 f} + \frac{2ic^5 e^{-4i(fx+e)} A}{a^2 f} + \frac{112ic^5 Bx}{a^2} + \frac{48c^5 Ax}{a^2} + \frac{112ic^5 B}{f a^2}$
derivativedivides	$\frac{iB c^5 \tan(fx+e)^3}{3a^2 f} + \frac{ic^5 A \tan(fx+e)^2}{2f a^2} - \frac{24ic^5 B \tan(fx+e)}{f a^2} - \frac{7c^5 B \tan(fx+e)^2}{2f a^2} - \frac{7c^5 A \tan(fx+e)}{f a^2} + \frac{24c^5 A}{f a^2}$
default	$\frac{iB c^5 \tan(fx+e)^3}{3a^2 f} + \frac{ic^5 A \tan(fx+e)^2}{2f a^2} - \frac{24ic^5 B \tan(fx+e)}{f a^2} - \frac{7c^5 B \tan(fx+e)^2}{2f a^2} - \frac{7c^5 A \tan(fx+e)}{f a^2} + \frac{24c^5 A}{f a^2}$
norman	$\frac{-25ic^5 A + 47c^5 B}{af} + \frac{8(7ic^5 B + 3c^5 A)x}{a} - \frac{(-ic^5 A + 7c^5 B) \tan(fx+e)^6}{2af} - \frac{7(10ic^5 B + 3c^5 A) \tan(fx+e)^5}{3af} + \frac{(-83ic^5 A + 133c^5 B) \tan(fx+e)^4}{2af}$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5/(a+I*a*tan(f*x+e))^2,x,method=_R
ETURNVERBOSE)
```

```
output 20*c^5/a^2/f*exp(-2*I*(f*x+e))*B-12*I*c^5/a^2/f*exp(-2*I*(f*x+e))*A-2*c^5/
a^2/f*exp(-4*I*(f*x+e))*B+2*I*c^5/a^2/f*exp(-4*I*(f*x+e))*A+112*I*c^5/a^2*
B*x+48*c^5/a^2*A*x+112*I*c^5/f/a^2*B*e+48*c^5/f/a^2*A*e+2/3*c^5*(-18*I*A*e
xp(4*I*(f*x+e))+54*B*exp(4*I*(f*x+e))-39*I*A*exp(2*I*(f*x+e))+123*B*exp(2*
I*(f*x+e))-21*I*A+73*B)/f/a^2/(exp(2*I*(f*x+e))+1)^3-56*c^5/f/a^2*ln(exp(2
*I*(f*x+e))+1)*B+24*I*c^5/f/a^2*ln(exp(2*I*(f*x+e))+1)*A
```

$$3.716. \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^5}{(a+ia \tan(e+fx))^2} dx$$

3.716.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 329, normalized size of antiderivative = 1.70

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^5}{(a + i a \tan(e + fx))^2} dx$$

$$= \frac{2(24(3A + 7iB)c^5 f x e^{(10i f x + 10i e)} - 3(3iA - 7B)c^5 e^{(2i f x + 2i e)} - 3(-iA + B)c^5 + 12(6(3A + 7iB)c^5 f$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")
```

```
output 2/3*(24*(3*A + 7*I*B)*c^5*f*x*e^(10*I*f*x + 10*I*e) - 3*(3*I*A - 7*B)*c^5*
e^(2*I*f*x + 2*I*e) - 3*(-I*A + B)*c^5 + 12*(6*(3*A + 7*I*B)*c^5*f*x - (3*
I*A - 7*B)*c^5)*e^(8*I*f*x + 8*I*e) + 6*(12*(3*A + 7*I*B)*c^5*f*x - 5*(3*I
*A - 7*B)*c^5)*e^(6*I*f*x + 6*I*e) + 2*(12*(3*A + 7*I*B)*c^5*f*x - 11*(3*I
*A - 7*B)*c^5)*e^(4*I*f*x + 4*I*e) - 12*((-3*I*A + 7*B)*c^5*e^(10*I*f*x +
10*I*e) + 3*(-3*I*A + 7*B)*c^5*e^(8*I*f*x + 8*I*e) + 3*(-3*I*A + 7*B)*c^5*
e^(6*I*f*x + 6*I*e) + (-3*I*A + 7*B)*c^5*e^(4*I*f*x + 4*I*e))*log(e^(2*I*f
*x + 2*I*e) + 1)/(a^2*f*e^(10*I*f*x + 10*I*e) + 3*a^2*f*e^(8*I*f*x + 8*I*
e) + 3*a^2*f*e^(6*I*f*x + 6*I*e) + a^2*f*e^(4*I*f*x + 4*I*e))
```

3.716.6 Sympy [A] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 445, normalized size of antiderivative = 2.29

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^5}{(a + i a \tan(e + fx))^2} dx$$

$$= \frac{-42iAc^5 + 146Bc^5 + (-78iAc^5 e^{2ie} + 246Bc^5 e^{2ie}) e^{2ifx} + (-36iAc^5 e^{4ie} + 108Bc^5 e^{4ie}) e^{4ifx}}{3a^2 f e^{6ie} e^{6ifx} + 9a^2 f e^{4ie} e^{4ifx} + 9a^2 f e^{2ie} e^{2ifx} + 3a^2 f}$$

$$+ \begin{cases} \frac{((2iAa^2 c^5 f e^{2ie} - 2Ba^2 c^5 f e^{2ie}) e^{-4ifx} + (-12iAa^2 c^5 f e^{4ie} + 20Ba^2 c^5 f e^{4ie}) e^{-2ifx}) e^{-6ie}}{a^4 f^2} & \text{for } a^4 f^2 e^{6ie} \neq 0 \\ x \left(-\frac{48Ac^5 + 112iBc^5}{a^2} + \frac{(48Ac^5 e^{4ie} - 24Ac^5 e^{2ie} + 8Ac^5 + 112iBc^5 e^{4ie} - 40iBc^5 e^{2ie} + 8iBc^5) e^{-4ie}}{a^2} \right) & \text{otherwise} \end{cases}$$

$$+ \frac{8ic^5 \cdot (3A + 7iB) \log(e^{2ifx} + e^{-2ie})}{a^2 f} + \frac{x(48Ac^5 + 112iBc^5)}{a^2}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**5/(a+I*a*tan(f*x+e))**2,x)
```

```
output (-42*I*A*c**5 + 146*B*c**5 + (-78*I*A*c**5*exp(2*I*e) + 246*B*c**5*exp(2*I
*e))*exp(2*I*f*x) + (-36*I*A*c**5*exp(4*I*e) + 108*B*c**5*exp(4*I*e))*exp(
4*I*f*x))/(3*a**2*f*exp(6*I*e)*exp(6*I*f*x) + 9*a**2*f*exp(4*I*e)*exp(4*I*
f*x) + 9*a**2*f*exp(2*I*e)*exp(2*I*f*x) + 3*a**2*f) + Piecewise((((2*I*A*a
**2*c**5*f*exp(2*I*e) - 2*B*a**2*c**5*f*exp(2*I*e))*exp(-4*I*f*x) + (-12*I
*A*a**2*c**5*f*exp(4*I*e) + 20*B*a**2*c**5*f*exp(4*I*e))*exp(-2*I*f*x))*ex
p(-6*I*e)/(a**4*f**2), Ne(a**4*f**2*exp(6*I*e), 0)), (x*(-(48*A*c**5 + 112
*I*B*c**5)/a**2 + (48*A*c**5*exp(4*I*e) - 24*A*c**5*exp(2*I*e) + 8*A*c**5
+ 112*I*B*c**5*exp(4*I*e) - 40*I*B*c**5*exp(2*I*e) + 8*I*B*c**5)*exp(-4*I*
e)/a**2), True)) + 8*I*c**5*(3*A + 7*I*B)*log(exp(2*I*f*x) + exp(-2*I*e))/
(a**2*f) + x*(48*A*c**5 + 112*I*B*c**5)/a**2
```

3.716.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^5}{(a + ia \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5/(a+I*a*tan(f*x+e))^2,x, al
gorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.716.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 492 vs. $2(166) = 332$.

Time = 1.03 (sec) , antiderivative size = 492, normalized size of antiderivative = 2.54

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^5}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{2 \left(\frac{12(3iAc^5 - 7Bc^5) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)}{a^2} - \frac{24(3iAc^5 - 7Bc^5) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - i)}{a^2} - \frac{12(-3iAc^5 + 7Bc^5) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)}{a^2} \right)}{a^2}$$

3.716. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^5}{(a+ia \tan(e+fx))^2} dx$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output
$$\frac{2/3*(12*(3*I*A*c^5 - 7*B*c^5)*\log(\tan(1/2*f*x + 1/2*e) + 1)/a^2 - 24*(3*I*A*c^5 - 7*B*c^5)*\log(\tan(1/2*f*x + 1/2*e) - I)/a^2 - 12*(-3*I*A*c^5 + 7*B*c^5)*\log(\tan(1/2*f*x + 1/2*e) - 1)/a^2 - (66*I*A*c^5*\tan(1/2*f*x + 1/2*e)^6 - 154*B*c^5*\tan(1/2*f*x + 1/2*e)^6 - 21*A*c^5*\tan(1/2*f*x + 1/2*e)^5 - 72*I*B*c^5*\tan(1/2*f*x + 1/2*e)^5 - 201*I*A*c^5*\tan(1/2*f*x + 1/2*e)^4 + 483*B*c^5*\tan(1/2*f*x + 1/2*e)^4 + 42*A*c^5*\tan(1/2*f*x + 1/2*e)^3 + 148*I*B*c^5*\tan(1/2*f*x + 1/2*e)^3 + 201*I*A*c^5*\tan(1/2*f*x + 1/2*e)^2 - 483*B*c^5*\tan(1/2*f*x + 1/2*e)^2 - 21*A*c^5*\tan(1/2*f*x + 1/2*e) - 72*I*B*c^5*\tan(1/2*f*x + 1/2*e) - 66*I*A*c^5 + 154*B*c^5)/((\tan(1/2*f*x + 1/2*e)^2 - 1)^3*a^2) - 2*(-75*I*A*c^5*\tan(1/2*f*x + 1/2*e)^4 + 175*B*c^5*\tan(1/2*f*x + 1/2*e)^4 - 324*A*c^5*\tan(1/2*f*x + 1/2*e)^3 - 748*I*B*c^5*\tan(1/2*f*x + 1/2*e)^3 + 522*I*A*c^5*\tan(1/2*f*x + 1/2*e)^2 - 1170*B*c^5*\tan(1/2*f*x + 1/2*e)^2 + 324*A*c^5*\tan(1/2*f*x + 1/2*e) + 748*I*B*c^5*\tan(1/2*f*x + 1/2*e) - 75*I*A*c^5 + 175*B*c^5)/(a^2*(\tan(1/2*f*x + 1/2*e) - I)^4))/f$$

3.716.9 Mupad [B] (verification not implemented)

Time = 8.53 (sec) , antiderivative size = 282, normalized size of antiderivative = 1.45

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^5}{(a + ia \tan(e + fx))^2} dx$$

$$= -\frac{\ln(\tan(e + fx) - i) \left(-\frac{56 B c^5}{a^2} + \frac{A c^5 24i}{a^2} \right)}{f} + \frac{\tan(e + fx)^2 \left(-\frac{3 B c^5}{2 a^2} + \frac{c^5 (A + B 4i) li}{2 a^2} \right)}{f}$$

$$- \frac{\tan(e + fx) \left(\frac{3 c^5 (A + B 4i)}{a^2} + \frac{B c^5 6i}{a^2} - \frac{c^5 (-3 B + A 2i) 2i}{a^2} \right)}{f}$$

$$+ \frac{-\frac{(-24 B c^5 + A c^5 8i) li}{2 a^2} + \frac{16 A c^5 + B c^5 64i}{2 a^2} + \frac{(-56 B c^5 + A c^5 24i) 3i}{2 a^2} + \tan(e + fx) \left(\frac{(16 A c^5 + B c^5 64i) li}{a^2} - \frac{2(-56 B c^5 + A c^5 24i) li}{a^2} \right)}{f (\tan(e + fx)^2 li + 2 \tan(e + fx) - i)}$$

$$+ \frac{B c^5 \tan(e + fx)^3 li}{3 a^2 f}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^5)/(a + a*tan(e + f*x)*1i)^2,x)`

output $(\tan(e + fx))^2 \left(\frac{c^5(A + B4i)1i}{2a^2} - \frac{3Bc^5}{2a^2} \right) / f - \log(\tan(e + fx) - 1i) \left(\frac{A^5c^5 24i}{a^2} - \frac{56Bc^5}{a^2} \right) / f - (\tan(e + fx) \left(\frac{3c^5(A + B4i)}{a^2} + \frac{Bc^5 6i}{a^2} - \frac{c^5(A^2i - 3B)2i}{a^2} \right) / f + \left(\frac{16A^5c^5 + Bc^5 64i}{2a^2} - \frac{(A^5c^5 8i - 24Bc^5)1i}{2a^2} + \frac{(A^5c^5 24i - 56Bc^5)3i}{2a^2} + \tan(e + fx) \left(\frac{16A^5c^5 + Bc^5 64i}{a^2} - \frac{2(A^5c^5 24i - 56Bc^5)}{a^2} \right) / (f(2\tan(e + fx) + \tan(e + fx)^2 1i - 1i)) + \frac{Bc^5 \tan(e + fx)^3 1i}{3a^2 f} \right)$

3.716. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^5}{(a+ia \tan(e+fx))^2} dx$

$$3.717 \quad \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^4}{(a+ia \tan(e+fx))^2} dx$$

3.717.1 Optimal result	6633
3.717.2 Mathematica [A] (verified)	6633
3.717.3 Rubi [A] (verified)	6634
3.717.4 Maple [A] (verified)	6636
3.717.5 Fricas [A] (verification not implemented)	6636
3.717.6 Sympy [A] (verification not implemented)	6637
3.717.7 Maxima [F(-2)]	6638
3.717.8 Giac [B] (verification not implemented)	6638
3.717.9 Mupad [B] (verification not implemented)	6639

3.717.1 Optimal result

Integrand size = 41, antiderivative size = 158

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^2} dx \\ &= \frac{6(A + 3iB)c^4x}{a^2} + \frac{6(iA - 3B)c^4 \log(\cos(e + fx))}{a^2 f} - \frac{4(iA - B)c^4}{a^2 f(i - \tan(e + fx))^2} \\ &+ \frac{4(3A + 5iB)c^4}{a^2 f(i - \tan(e + fx))} - \frac{(A + 6iB)c^4 \tan(e + fx)}{a^2 f} - \frac{Bc^4 \tan^2(e + fx)}{2a^2 f} \end{aligned}$$

```
output 6*(A+3*I*B)*c^4*x/a^2+6*(I*A-3*B)*c^4*ln(cos(f*x+e))/a^2/f-4*(I*A-B)*c^4/a^2/f/(I-tan(f*x+e))^2+4*(3*A+5*I*B)*c^4/a^2/f/(I-tan(f*x+e))-(A+6*I*B)*c^4*tan(f*x+e)/a^2/f-1/2*B*c^4*tan(f*x+e)^2/a^2/f
```

3.717.2 Mathematica [A] (verified)

Time = 6.03 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.82

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^2} dx \\ &= \frac{c^4 \left(\frac{2(A+3iB)(i+\tan(e+fx))^3}{(a+ia \tan(e+fx))^2} + \frac{B(i+\tan(e+fx))^4}{(a+ia \tan(e+fx))^2} + \frac{12(-iA+3B) \left(\log(i-\tan(e+fx)) + \frac{-2-4i \tan(e+fx)}{(-i+\tan(e+fx))^2} \right)}{a^2} \right)}{2f} \end{aligned}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^4)/(a + I*a*Tan[e + f*x])^2,x]`

output `(c^4*((2*(A + (3*I)*B)*(I + Tan[e + f*x])^3)/(a + I*a*Tan[e + f*x])^2 + (B*(I + Tan[e + f*x])^4)/(a + I*a*Tan[e + f*x])^2 + (12*((-I)*A + 3*B)*(Log[I - Tan[e + f*x]] + (-2 - (4*I)*Tan[e + f*x])/(-I + Tan[e + f*x])^2))/a^2)/(2*f)`

3.717.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.72, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - ict \tan(e + fx))^4 (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c - ict \tan(e + fx))^4 (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{c^3 (1 - i \tan(e + fx))^3 (A + B \tan(e + fx))}{a^3 (i \tan(e + fx) + 1)^3} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{c^4 \int \frac{(1 - i \tan(e + fx))^3 (A + B \tan(e + fx))}{(i \tan(e + fx) + 1)^3} d \tan(e + fx)}{a^2 f} \\
 & \quad \downarrow \text{86} \\
 & \frac{c^4 \int \left(\frac{8i(A + iB)}{(\tan(e + fx) - i)^3} - A \left(\frac{6iB}{A} + 1 \right) - B \tan(e + fx) - \frac{6i(A + 3iB)}{\tan(e + fx) - i} + \frac{4(3A + 5iB)}{(\tan(e + fx) - i)^2} \right) d \tan(e + fx)}{a^2 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{c^4 \left(-(A + 6iB) \tan(e + fx) + \frac{4(3A + 5iB)}{-\tan(e + fx) + i} - \frac{4(-B + iA)}{(-\tan(e + fx) + i)^2} - 6(-3B + iA) \log(-\tan(e + fx) + i) - \frac{1}{2} B \tan^2(e + fx) \right)}{a^2 f}
 \end{aligned}$$

3.717. $\int \frac{(A + B \tan(e + fx))(c - ict \tan(e + fx))^4}{(a + ia \tan(e + fx))^2} dx$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^4)/(a + I*a*Tan[e + f*x])^2,x]`

output `(c^4*(-6*(I*A - 3*B)*Log[I - Tan[e + f*x]] - (4*(I*A - B))/(I - Tan[e + f*x])^2 + (4*(3*A + (5*I)*B))/(I - Tan[e + f*x]) - (A + (6*I)*B)*Tan[e + f*x] - (B*Tan[e + f*x]^2)/2))/(a^2*f)`

3.717.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.717.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.51

method	result
derivativedivides	$-\frac{Bc^4 \tan(fx+e)^2}{2a^2f} - \frac{c^4 A \tan(fx+e)}{fa^2} - \frac{6ic^4 B \tan(fx+e)}{fa^2} - \frac{20ic^4 B}{fa^2(-i+\tan(fx+e))} - \frac{12c^4 A}{fa^2(-i+\tan(fx+e))} + \dots$
default	$-\frac{Bc^4 \tan(fx+e)^2}{2a^2f} - \frac{c^4 A \tan(fx+e)}{fa^2} - \frac{6ic^4 B \tan(fx+e)}{fa^2} - \frac{20ic^4 B}{fa^2(-i+\tan(fx+e))} - \frac{12c^4 A}{fa^2(-i+\tan(fx+e))} + \dots$
risch	$\frac{8c^4 e^{-2i(fx+e)} B}{a^2 f} - \frac{4ic^4 e^{-2i(fx+e)} A}{a^2 f} - \frac{c^4 e^{-4i(fx+e)} B}{a^2 f} + \frac{ic^4 e^{-4i(fx+e)} A}{a^2 f} + \frac{36ic^4 Bx}{a^2} + \frac{12c^4 Ax}{a^2} + \frac{36ic^4 B}{fa^2} + \dots$
norman	$\frac{-8ic^4 A + 17c^4 B}{af} + \frac{6(3ic^4 B + c^4 A)x}{a} - \frac{(6ic^4 B + c^4 A) \tan(fx+e)^5}{af} + \frac{(-32ic^4 A + 51c^4 B) \tan(fx+e)^2}{2af} + \frac{12(3ic^4 B + c^4 A)x \tan(fx+e)^2}{a} + \dots$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4/(a+I*a*tan(f*x+e))^2,x,method=_R
ETURNVERBOSE)
```

```
output -1/2*B*c^4*tan(f*x+e)^2/a^2/f-1/f*c^4/a^2*A*tan(f*x+e)-6*I/f*c^4/a^2*B*tan
(f*x+e)-20*I/f*c^4/a^2/(-I+tan(f*x+e))*B-12/f*c^4/a^2/(-I+tan(f*x+e))*A+6/
f*c^4/a^2*A*arctan(tan(f*x+e))-3*I/f*c^4/a^2*A*ln(1+tan(f*x+e)^2)+18*I/f*c
^4/a^2*B*arctan(tan(f*x+e))+9/f*c^4/a^2*B*ln(1+tan(f*x+e)^2)-4*I/f*c^4/a^2
/(-I+tan(f*x+e))^2*A+4/f*c^4/a^2/(-I+tan(f*x+e))^2*B
```

3.717.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.58

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{12(A + 3iB)c^4 f x e^{(8i f x + 8i e)} - 2(iA - 3B)c^4 e^{(2i f x + 2i e)} + (iA - B)c^4 + 6(4(A + 3iB)c^4 f x - (iA - 3B)c^4)}{(a + ia \tan(e + fx))^2}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4/(a+I*a*tan(f*x+e))^2,x, al
gorithm="fricas")
```

```
output (12*(A + 3*I*B)*c^4*f*x*e^(8*I*f*x + 8*I*e) - 2*(I*A - 3*B)*c^4*e^(2*I*f*x
+ 2*I*e) + (I*A - B)*c^4 + 6*(4*(A + 3*I*B)*c^4*f*x - (I*A - 3*B)*c^4)*e^
(6*I*f*x + 6*I*e) + 3*(4*(A + 3*I*B)*c^4*f*x - 3*(I*A - 3*B)*c^4)*e^(4*I*f
*x + 4*I*e) - 6*((-I*A + 3*B)*c^4*e^(8*I*f*x + 8*I*e) + 2*(-I*A + 3*B)*c^4
*e^(6*I*f*x + 6*I*e) + (-I*A + 3*B)*c^4*e^(4*I*f*x + 4*I*e))*log(e^(2*I*f*
x + 2*I*e) + 1))/(a^2*f*e^(8*I*f*x + 8*I*e) + 2*a^2*f*e^(6*I*f*x + 6*I*e)
+ a^2*f*e^(4*I*f*x + 4*I*e))
```

3.717.6 Sympy [A] (verification not implemented)

Time = 0.58 (sec) , antiderivative size = 377, normalized size of antiderivative = 2.39

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^4}{(a + i a \tan(e + fx))^2} dx$$

$$= \frac{-2iAc^4 + 12Bc^4 + (-2iAc^4e^{2ie} + 10Bc^4e^{2ie})e^{2ifx}}{a^2fe^{4ie}e^{4ifx} + 2a^2fe^{2ie}e^{2ifx} + a^2f}$$

$$+ \begin{cases} \frac{((iAa^2c^4fe^{2ie} - Ba^2c^4fe^{2ie})e^{-4ifx} + (-4iAa^2c^4fe^{4ie} + 8Ba^2c^4fe^{4ie})e^{-2ifx})e^{-6ie}}{a^4f^2} & \text{for } a^4f^2e^{6ie} \neq 0 \\ x \left(-\frac{12Ac^4 + 36iBc^4}{a^2} + \frac{(12Ac^4e^{4ie} - 8Ac^4e^{2ie} + 4Ac^4 + 36iBc^4e^{4ie} - 16iBc^4e^{2ie} + 4iBc^4)e^{-4ie}}{a^2} \right) & \text{otherwise} \end{cases}$$

$$+ \frac{6ic^4(A + 3iB) \log(e^{2ifx} + e^{-2ie})}{a^2f} + \frac{x(12Ac^4 + 36iBc^4)}{a^2}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**4/(a+I*a*tan(f*x+e))**2,x)
```

```
output (-2*I*A*c**4 + 12*B*c**4 + (-2*I*A*c**4*exp(2*I*e) + 10*B*c**4*exp(2*I*e))
*exp(2*I*f*x))/(a**2*f*exp(4*I*e)*exp(4*I*f*x) + 2*a**2*f*exp(2*I*e)*exp(2
*I*f*x) + a**2*f) + Piecewise((((I*A*a**2*c**4*f*exp(2*I*e) - B*a**2*c**4*
f*exp(2*I*e))*exp(-4*I*f*x) + (-4*I*A*a**2*c**4*f*exp(4*I*e) + 8*B*a**2*c
**4*f*exp(4*I*e))*exp(-2*I*f*x))*exp(-6*I*e)/(a**4*f**2), Ne(a**4*f**2*exp(
6*I*e), 0)), (x*(-(12*A*c**4 + 36*I*B*c**4)/a**2 + (12*A*c**4*exp(4*I*e) -
8*A*c**4*exp(2*I*e) + 4*A*c**4 + 36*I*B*c**4*exp(4*I*e) - 16*I*B*c**4*exp
(2*I*e) + 4*I*B*c**4)*exp(-4*I*e)/a**2), True)) + 6*I*c**4*(A + 3*I*B)*log
(exp(2*I*f*x) + exp(-2*I*e))/(a**2*f) + x*(12*A*c**4 + 36*I*B*c**4)/a**2
```

3.717.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.717.8 Giac [B] (verification not implemented)Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 423 vs. $2(136) = 272$.

Time = 0.78 (sec) , antiderivative size = 423, normalized size of antiderivative = 2.68

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^2} dx$$

$$\frac{6(iAc^4 - 3Bc^4) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)}{a^2} - \frac{12(iAc^4 - 3Bc^4) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - i)}{a^2} - \frac{6(-iAc^4 + 3Bc^4) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)}{a^2} - \frac{9iAc^4}{a^2}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")
```

```
output (6*(I*A*c^4 - 3*B*c^4)*log(tan(1/2*f*x + 1/2*e) + 1)/a^2 - 12*(I*A*c^4 - 3*B*c^4)*log(tan(1/2*f*x + 1/2*e) - I)/a^2 - 6*(-I*A*c^4 + 3*B*c^4)*log(tan(1/2*f*x + 1/2*e) - 1)/a^2 - (9*I*A*c^4*tan(1/2*f*x + 1/2*e)^4 - 27*B*c^4*tan(1/2*f*x + 1/2*e)^4 - 2*A*c^4*tan(1/2*f*x + 1/2*e)^3 - 12*I*B*c^4*tan(1/2*f*x + 1/2*e)^3 - 18*I*A*c^4*tan(1/2*f*x + 1/2*e)^2 + 56*B*c^4*tan(1/2*f*x + 1/2*e)^2 + 2*A*c^4*tan(1/2*f*x + 1/2*e) + 12*I*B*c^4*tan(1/2*f*x + 1/2*e) + 9*I*A*c^4 - 27*B*c^4)/((tan(1/2*f*x + 1/2*e)^2 - 1)^2*a^2) - (-25*I*A*c^4*tan(1/2*f*x + 1/2*e)^4 + 75*B*c^4*tan(1/2*f*x + 1/2*e)^4 - 108*A*c^4*tan(1/2*f*x + 1/2*e)^3 - 324*I*B*c^4*tan(1/2*f*x + 1/2*e)^3 + 182*I*A*c^4*tan(1/2*f*x + 1/2*e)^2 - 514*B*c^4*tan(1/2*f*x + 1/2*e)^2 + 108*A*c^4*tan(1/2*f*x + 1/2*e) + 324*I*B*c^4*tan(1/2*f*x + 1/2*e) - 25*I*A*c^4 + 75*B*c^4)/(a^2*(tan(1/2*f*x + 1/2*e) - I)^4))/f
```

3.717. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^4}{(a+ia \tan(e+fx))^2} dx$

3.717.9 Mupad [B] (verification not implemented)

Time = 8.48 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.31

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^2} dx$$

$$= -\frac{\ln(\tan(e + fx) - i) \left(-\frac{18 B c^4}{a^2} + \frac{A c^4 6i}{a^2}\right)}{f} - \frac{\tan(e + fx) \left(\frac{c^4 (A + B 3i)}{a^2} + \frac{B c^4 3i}{a^2}\right)}{f}$$

$$- \frac{\frac{(-6 B c^4 + A c^4 2i) 1i}{2 a^2} - \frac{(-18 B c^4 + A c^4 6i) 3i}{2 a^2} + \tan(e + fx) \left(\frac{2(-18 B c^4 + A c^4 6i)}{a^2} + \frac{16 B c^4}{a^2}\right) - \frac{B c^4 8i}{a^2}}{f (\tan(e + fx)^2 1i + 2 \tan(e + fx) - i)}$$

$$- \frac{B c^4 \tan(e + fx)^2}{2 a^2 f}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^4)/(a + a*tan(e + f*x)*1i)^2,x)`

output `- (log(tan(e + f*x) - 1i)*((A*c^4*6i)/a^2 - (18*B*c^4)/a^2))/f - (tan(e + f*x)*((c^4*(A + B*3i))/a^2 + (B*c^4*3i)/a^2))/f - (((A*c^4*2i - 6*B*c^4)*1i)/(2*a^2) - ((A*c^4*6i - 18*B*c^4)*3i)/(2*a^2) + tan(e + f*x)*((2*(A*c^4*6i - 18*B*c^4))/a^2 + (16*B*c^4)/a^2) - (B*c^4*8i)/a^2)/(f*(2*tan(e + f*x) + tan(e + f*x)^2*1i - 1i)) - (B*c^4*tan(e + f*x)^2)/(2*a^2*f)`

3.718
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^3}{(a+ia \tan(e+fx))^2} dx$$

3.718.1 Optimal result	6640
3.718.2 Mathematica [A] (verified)	6640
3.718.3 Rubi [A] (verified)	6641
3.718.4 Maple [A] (verified)	6643
3.718.5 Fricas [A] (verification not implemented)	6643
3.718.6 Sympy [A] (verification not implemented)	6644
3.718.7 Maxima [F(-2)]	6644
3.718.8 Giac [B] (verification not implemented)	6645
3.718.9 Mupad [B] (verification not implemented)	6645

3.718.1 Optimal result

Integrand size = 41, antiderivative size = 128

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{(a + ia \tan(e + fx))^2} dx \\ &= \frac{(A + 5iB)c^3x}{a^2} + \frac{(iA - 5B)c^3 \log(\cos(e + fx))}{a^2 f} \\ & \quad - \frac{2(iA - B)c^3}{a^2 f(i - \tan(e + fx))^2} + \frac{4(A + 2iB)c^3}{a^2 f(i - \tan(e + fx))} - \frac{iBc^3 \tan(e + fx)}{a^2 f} \end{aligned}$$

output

```
(A+5*I*B)*c^3*x/a^2+(I*A-5*B)*c^3*ln(cos(f*x+e))/a^2/f-2*(I*A-B)*c^3/a^2/f
/(I-tan(f*x+e))^2+4*(A+2*I*B)*c^3/a^2/f/(I-tan(f*x+e))-I*B*c^3*tan(f*x+e)/
a^2/f
```

3.718.2 Mathematica [A] (verified)

Time = 5.64 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.72

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{(a + ia \tan(e + fx))^2} dx \\ &= \frac{\frac{B(c-ic \tan(e+fx))^3}{(a+ia \tan(e+fx))^2} + \frac{(-iA+5B)c^3 \left(\log(i - \tan(e+fx)) + \frac{-2-4i \tan(e+fx)}{(-i + \tan(e+fx))^2} \right)}{a^2}}{f} \end{aligned}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^3)/(a + I*a*Tan[e + f*x])^2,x]`

output `((B*(c - I*c*Tan[e + f*x])^3)/(a + I*a*Tan[e + f*x])^2 + (((-I)*A + 5*B)*c^3*(Log[I - Tan[e + f*x]] + (-2 - (4*I)*Tan[e + f*x])/(-I + Tan[e + f*x])^2))/a^2)/f`

3.718.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.73, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - ic \tan(e + fx))^3 (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c - ic \tan(e + fx))^3 (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{c^2(1-i \tan(e+fx))^2(A+B \tan(e+fx))}{a^3(i \tan(e+fx)+1)^3} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{c^3 \int \frac{(1-i \tan(e+fx))^2(A+B \tan(e+fx))}{(i \tan(e+fx)+1)^3} d \tan(e + fx)}{a^2 f} \\
 & \quad \downarrow \text{86} \\
 & \frac{c^3 \int \left(\frac{4i(A+iB)}{(\tan(e+fx)-i)^3} - iB - \frac{i(A+5iB)}{\tan(e+fx)-i} + \frac{4(A+2iB)}{(\tan(e+fx)-i)^2} \right) d \tan(e + fx)}{a^2 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{c^3 \left(-\frac{2(-B+ia)}{(-\tan(e+fx)+i)^2} + \frac{4(A+2iB)}{-\tan(e+fx)+i} - (-5B + iA) \log(-\tan(e + fx) + i) - iB \tan(e + fx) \right)}{a^2 f}
 \end{aligned}$$

3.718. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^3}{(a+ia \tan(e+fx))^2} dx$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^3)/(a + I*a*Tan[e + f*x])^2,x]`

output `(c^3*(-((I*A - 5*B)*Log[I - Tan[e + f*x]]) - (2*(I*A - B))/(I - Tan[e + f*x]))^2 + (4*(A + (2*I)*B))/(I - Tan[e + f*x]) - I*B*Tan[e + f*x]))/(a^2*f)`

3.718.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x_, x], x] /; FreeQ[a, x] && !MatchQ[F_x_, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.718.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.56

method	result
derivativedivides	$-\frac{iBc^3 \tan(fx+e)}{a^2 f} - \frac{8ic^3 B}{fa^2(-i+\tan(fx+e))} - \frac{4c^3 A}{fa^2(-i+\tan(fx+e))} - \frac{2ic^3 A}{fa^2(-i+\tan(fx+e))^2} + \frac{2c^3 B}{fa^2(-i+\tan(fx+e))}$
default	$-\frac{iBc^3 \tan(fx+e)}{a^2 f} - \frac{8ic^3 B}{fa^2(-i+\tan(fx+e))} - \frac{4c^3 A}{fa^2(-i+\tan(fx+e))} - \frac{2ic^3 A}{fa^2(-i+\tan(fx+e))^2} + \frac{2c^3 B}{fa^2(-i+\tan(fx+e))}$
risch	$\frac{3c^3 e^{-2i(fx+e)} B}{a^2 f} - \frac{ic^3 e^{-2i(fx+e)} A}{a^2 f} - \frac{c^3 e^{-4i(fx+e)} B}{2a^2 f} + \frac{ic^3 e^{-4i(fx+e)} A}{2a^2 f} + \frac{10ic^3 Bx}{a^2} + \frac{2c^3 Ax}{a^2} + \frac{10ic^3 Bx}{a^2 f} + \frac{(5ic^3 B+c^3 A)x}{a} + \frac{-2ic^3 A+6c^3 B}{af} + \frac{(5ic^3 B+c^3 A)x \tan(fx+e)^4}{a} + \frac{2(5ic^3 B+c^3 A)x \tan(fx+e)^2}{a} - \frac{2(5ic^3 B+2c^3 A) \tan(fx+e)^3}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^3}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^5}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^7}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^9}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{11}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{13}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{15}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{17}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{19}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{21}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{23}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{25}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{27}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{29}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{31}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{33}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{35}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{37}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{39}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{41}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{43}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{45}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{47}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{49}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{51}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{53}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{55}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{57}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{59}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{61}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{63}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{65}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{67}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{69}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{71}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{73}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{75}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{77}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{79}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{81}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{83}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{85}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{87}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{89}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{91}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{93}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{95}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{97}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{99}}{af}$
norman	$\frac{(5ic^3 B+c^3 A)x}{a} + \frac{-2ic^3 A+6c^3 B}{af} + \frac{(5ic^3 B+c^3 A)x \tan(fx+e)^4}{a} + \frac{2(5ic^3 B+c^3 A)x \tan(fx+e)^2}{a} - \frac{2(5ic^3 B+2c^3 A) \tan(fx+e)^3}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^3}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^5}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^7}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^9}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{11}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{13}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{15}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{17}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{19}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{21}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{23}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{25}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{27}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{29}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{31}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{33}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{35}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{37}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{39}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{41}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{43}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{45}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{47}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{49}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{51}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{53}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{55}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{57}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{59}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{61}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{63}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{65}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{67}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{69}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{71}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{73}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{75}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{77}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{79}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{81}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{83}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{85}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{87}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{89}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{91}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{93}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{95}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{97}}{af} + \frac{2(5ic^3 B+c^3 A) \tan(fx+e)^{99}}{af}$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3/(a+I*a*tan(f*x+e))^2,x,method=_R
ETURNVERBOSE)
```

```
output -I*B*c^3*tan(f*x+e)/a^2/f-8*I/f*c^3/a^2/(-I+tan(f*x+e))*B-4/f*c^3/a^2/(-I+
tan(f*x+e))*A-2*I/f*c^3/a^2/(-I+tan(f*x+e))^2*A+2/f*c^3/a^2/(-I+tan(f*x+e)
)^2*B-1/2*I/f*c^3/a^2*A*ln(1+tan(f*x+e)^2)+5/2/f*c^3/a^2*B*ln(1+tan(f*x+e)
^2)+1/f*c^3/a^2*A*arctan(tan(f*x+e))+5*I/f*c^3/a^2*B*arctan(tan(f*x+e))
```

3.718.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.38

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{4(A + 5iB)c^3 f x e^{(6i f x + 6i e)} + (-iA + 5B)c^3 e^{(2i f x + 2i e)} + (iA - B)c^3 + 2(2(A + 5iB)c^3 f x - (iA - 5B)c^3)}{2(a^2 f e^{(6i f x + 6i e)} + a^2 f)}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3/(a+I*a*tan(f*x+e))^2,x, al
gorithm="fricas")
```

```
output 1/2*(4*(A + 5*I*B)*c^3*f*x*e^(6*I*f*x + 6*I*e) + (-I*A + 5*B)*c^3*e^(2*I*f
*x + 2*I*e) + (I*A - B)*c^3 + 2*(2*(A + 5*I*B)*c^3*f*x - (I*A - 5*B)*c^3)*
e^(4*I*f*x + 4*I*e) - 2*((-I*A + 5*B)*c^3*e^(6*I*f*x + 6*I*e) + (-I*A + 5*
B)*c^3*e^(4*I*f*x + 4*I*e))*log(e^(2*I*f*x + 2*I*e) + 1)/(a^2*f*e^(6*I*f*
x + 6*I*e) + a^2*f*e^(4*I*f*x + 4*I*e))
```

3.718. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^3}{(a+ia \tan(e+fx))^2} dx$

3.718.6 Sympy [A] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 309, normalized size of antiderivative = 2.41

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{(a + ia \tan(e + fx))^2} dx = \frac{2Bc^3}{a^2 f e^{2ie} e^{2ifx} + a^2 f}$$

$$+ \begin{cases} \frac{((iAa^2c^3fe^{2ie} - Ba^2c^3fe^{2ie})e^{-4ifx} + (-2iAa^2c^3fe^{4ie} + 6Ba^2c^3fe^{4ie})e^{-2ifx})e^{-6ie}}{2a^4f^2} & \text{for } a^4 f^2 e^{6ie} \neq 0 \\ x \left(-\frac{2Ac^3 + 10iBc^3}{a^2} + \frac{(2Ac^3e^{4ie} - 2Ac^3e^{2ie} + 2Ac^3 + 10iBc^3e^{4ie} - 6iBc^3e^{2ie} + 2iBc^3)e^{-4ie}}{a^2} \right) & \text{otherwise} \end{cases}$$

$$+ \frac{ic^3(A + 5iB) \log(e^{2ifx} + e^{-2ie})}{a^2 f} + \frac{x(2Ac^3 + 10iBc^3)}{a^2}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**3/(a+I*a*tan(f*x+e))**2,x)`

output `2*B*c**3/(a**2*f*exp(2*I*e)*exp(2*I*f*x) + a**2*f) + Piecewise((((I*A*a**2*c**3*f*exp(2*I*e) - B*a**2*c**3*f*exp(2*I*e))*exp(-4*I*f*x) + (-2*I*A*a**2*c**3*f*exp(4*I*e) + 6*B*a**2*c**3*f*exp(4*I*e))*exp(-2*I*f*x))*exp(-6*I*e)/(2*a**4*f**2), Ne(a**4*f**2*exp(6*I*e), 0)), (x*(-(2*A*c**3 + 10*I*B*c**3)/a**2 + (2*A*c**3*exp(4*I*e) - 2*A*c**3*exp(2*I*e) + 2*A*c**3 + 10*I*B*c**3*exp(4*I*e) - 6*I*B*c**3*exp(2*I*e) + 2*I*B*c**3)*exp(-4*I*e)/a**2), True)) + I*c**3*(A + 5*I*B)*log(exp(2*I*f*x) + exp(-2*I*e))/(a**2*f) + x*(2*A*c**3 + 10*I*B*c**3)/a**2`

3.718.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{(a + ia \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.718.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 341 vs. $2(108) = 216$.

Time = 0.66 (sec) , antiderivative size = 341, normalized size of antiderivative = 2.66

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^3}{(a + i a \tan(e + fx))^2} dx$$

$$= \frac{6(i A c^3 - 5 B c^3) \log(\tan(\frac{1}{2} f x + \frac{1}{2} e) + 1)}{a^2} + \frac{12(-i A c^3 + 5 B c^3) \log(\tan(\frac{1}{2} f x + \frac{1}{2} e) - i)}{a^2} - \frac{6(-i A c^3 + 5 B c^3) \log(\tan(\frac{1}{2} f x + \frac{1}{2} e) - 1)}{a^2} - \frac{6(i A c^3 - 5 B c^3) \log(\tan(\frac{1}{2} f x + \frac{1}{2} e) + i)}{a^2}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output `1/6*(6*(I*A*c^3 - 5*B*c^3)*log(tan(1/2*f*x + 1/2*e) + 1)/a^2 + 12*(-I*A*c^3 + 5*B*c^3)*log(tan(1/2*f*x + 1/2*e) - I)/a^2 - 6*(-I*A*c^3 + 5*B*c^3)*log(tan(1/2*f*x + 1/2*e) - 1)/a^2 - 6*(I*A*c^3*tan(1/2*f*x + 1/2*e)^2 - 5*B*c^3*tan(1/2*f*x + 1/2*e)^2 - 2*I*B*c^3*tan(1/2*f*x + 1/2*e) - I*A*c^3 + 5*B*c^3)/((tan(1/2*f*x + 1/2*e)^2 - 1)*a^2) - (-25*I*A*c^3*tan(1/2*f*x + 1/2*e)^4 + 125*B*c^3*tan(1/2*f*x + 1/2*e)^4 - 100*A*c^3*tan(1/2*f*x + 1/2*e)^3 - 548*I*B*c^3*tan(1/2*f*x + 1/2*e)^3 + 198*I*A*c^3*tan(1/2*f*x + 1/2*e)^2 - 894*B*c^3*tan(1/2*f*x + 1/2*e)^2 + 100*A*c^3*tan(1/2*f*x + 1/2*e) + 548*I*B*c^3*tan(1/2*f*x + 1/2*e) - 25*I*A*c^3 + 125*B*c^3)/(a^2*(tan(1/2*f*x + 1/2*e) - I)^4))/f`

3.718.9 Mupad [B] (verification not implemented)

Time = 9.07 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.52

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^3}{(a + i a \tan(e + fx))^2} dx$$

$$= \frac{c^3 (6 B - A 2i + 4 A \tan(e + fx) + B \tan(e + fx) 7i - A \ln(-1 - \tan(e + fx) 1i) 1i + 5 B \ln(-1 - \tan(e + fx) 1i))}{(a + i a \tan(e + fx))^2}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^3)/(a + a*tan(e + f*x)*1i)^2,x)`

3.718. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^3}{(a+ia \tan(e+fx))^2} dx$

output $(c^3(6B - A^2i + 4A\tan(e + fx) + B\tan(e + fx)*7i - A\log(-\tan(e + fx)*1i - 1)*1i + 5B\log(-\tan(e + fx)*1i - 1) + 2B\tan(e + fx)^2 + B\tan(e + fx)^3*1i + A\tan(e + fx)^2\log(-\tan(e + fx)*1i - 1)*1i - 5B\tan(e + fx)^2\log(-\tan(e + fx)*1i - 1) + 2A\tan(e + fx)\log(-\tan(e + fx)*1i - 1) + B\tan(e + fx)\log(-\tan(e + fx)*1i - 1)*10i))/(a^2f*(\tan(e + fx)*1i + 1)^2)$

3.718. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^3}{(a+ia \tan(e+fx))^2} dx$

$$3.719 \quad \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^2}{(a+ia \tan(e+fx))^2} dx$$

3.719.1 Optimal result	6647
3.719.2 Mathematica [A] (verified)	6647
3.719.3 Rubi [A] (verified)	6648
3.719.4 Maple [A] (verified)	6649
3.719.5 Fracas [A] (verification not implemented)	6650
3.719.6 Sympy [A] (verification not implemented)	6650
3.719.7 Maxima [F(-2)]	6651
3.719.8 Giac [B] (verification not implemented)	6651
3.719.9 Mupad [B] (verification not implemented)	6652

3.719.1 Optimal result

Integrand size = 41, antiderivative size = 97

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{iBc^2x}{a^2} - \frac{Bc^2 \log(\cos(e + fx))}{a^2f} - \frac{(iA - B)c^2}{a^2f(i - \tan(e + fx))^2} + \frac{(A + 3iB)c^2}{a^2f(i - \tan(e + fx))}$$

```
output I*B*c^2*x/a^2-B*c^2*ln(cos(f*x+e))/a^2/f-(I*A-B)*c^2/a^2/f/(I-tan(f*x+e))^2+(A+3*I*B)*c^2/a^2/f/(I-tan(f*x+e))
```

3.719.2 Mathematica [A] (verified)

Time = 5.64 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.60

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{c^2 \left(B \log(i - \tan(e + fx)) - \frac{2B+(A+3iB)\tan(e+fx)}{(-i+\tan(e+fx))^2} \right)}{a^2f}$$

```
input Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^2)/(a + I*a*Tan[e + f*x])^2,x]
```

```
output (c^2*(B*Log[I - Tan[e + f*x]] - (2*B + (A + (3*I)*B)*Tan[e + f*x])/(-I + Tan[e + f*x])^2))/(a^2*f)
```

3.719.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.75, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - ic \tan(e + fx))^2 (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c - ic \tan(e + fx))^2 (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{c(1-i \tan(e+fx))(A+B \tan(e+fx))}{a^3(i \tan(e+fx)+1)^3} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{c^2 \int \frac{(1-i \tan(e+fx))(A+B \tan(e+fx))}{(i \tan(e+fx)+1)^3} d \tan(e + fx)}{a^2 f} \\
 & \quad \downarrow \text{86} \\
 & \frac{c^2 \int \left(\frac{2i(A+iB)}{(\tan(e+fx)-i)^3} + \frac{B}{\tan(e+fx)-i} + \frac{A+3iB}{(\tan(e+fx)-i)^2} \right) d \tan(e + fx)}{a^2 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{c^2 \left(-\frac{-B+iA}{(-\tan(e+fx)+i)^2} + \frac{A+3iB}{-\tan(e+fx)+i} + B \log(-\tan(e + fx) + i) \right)}{a^2 f}
 \end{aligned}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^2)/(a + I*a*Tan[e + f*x])^2,x]`

output `(c^2*(B*Log[I - Tan[e + f*x]] - (I*A - B)/(I - Tan[e + f*x])^2 + (A + (3*I)*B)/(I - Tan[e + f*x]))/(a^2*f)`

3.719.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.719.4 Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.18

method	result
risch	$\frac{c^2 B e^{-2i(fx+e)}}{a^2 f} - \frac{c^2 e^{-4i(fx+e)} B}{4a^2 f} + \frac{ic^2 e^{-4i(fx+e)} A}{4a^2 f} + \frac{2iB c^2 x}{a^2} + \frac{2ic^2 B e}{a^2 f} - \frac{c^2 B \ln(e^{2i(fx+e)} + 1)}{a^2 f}$
derivativedivides	$-\frac{3ic^2 B}{f a^2 (-i + \tan(fx+e))} - \frac{c^2 A}{f a^2 (-i + \tan(fx+e))} + \frac{c^2 B \ln(1 + \tan(fx+e)^2)}{2f a^2} + \frac{ic^2 B \arctan(\tan(fx+e))}{f a^2} - \frac{f a^2(-i + \tan(fx+e))}{f a^2(-i + \tan(fx+e))}$
default	$-\frac{3ic^2 B}{f a^2 (-i + \tan(fx+e))} - \frac{c^2 A}{f a^2 (-i + \tan(fx+e))} + \frac{c^2 B \ln(1 + \tan(fx+e)^2)}{2f a^2} + \frac{ic^2 B \arctan(\tan(fx+e))}{f a^2} - \frac{f a^2(-i + \tan(fx+e))}{f a^2(-i + \tan(fx+e))}$

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2/(a+I*a*tan(f*x+e))^2,x,method=_R
ETURNVERBOSE)`

$$3.719. \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^2}{(a+ia \tan(e+fx))^2} dx$$

output $c^2 B/a^2/f \exp(-2I*(f*x+e)) - 1/4*c^2/a^2/f \exp(-4I*(f*x+e))*B + 1/4*I*c^2/a^2/f \exp(-4I*(f*x+e))*A + 2*I*B*c^2*x/a^2 + 2*I*c^2*B/a^2/f * e^{-c^2*B/a^2/f * \ln(\exp(2I*(f*x+e))+1)}$

3.719.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.91

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^2}{(a + i a \tan(e + fx))^2} dx$$

$$= \frac{(8i B c^2 f x e^{(4i f x + 4i e)} - 4 B c^2 e^{(4i f x + 4i e)} \log(e^{(2i f x + 2i e)} + 1) + 4 B c^2 e^{(2i f x + 2i e)} + (i A - B) c^2) e^{(-4i f x - 4i e)}}{4 a^2 f}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`

output $1/4*(8*I*B*c^2*f*x*e^{(4*I*f*x + 4*I*e)} - 4*B*c^2*e^{(4*I*f*x + 4*I*e)}*\log(e^{(2*I*f*x + 2*I*e)} + 1) + 4*B*c^2*e^{(2*I*f*x + 2*I*e)} + (I*A - B)*c^2)*e^{(-4*I*f*x - 4*I*e)}/(a^2*f)$

3.719.6 Sympy [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 206, normalized size of antiderivative = 2.12

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^2}{(a + i a \tan(e + fx))^2} dx$$

$$= \frac{2i B c^2 x}{a^2} - \frac{B c^2 \log(e^{2i f x} + e^{-2i e})}{a^2 f}$$

$$+ \begin{cases} \frac{(4B a^2 c^2 f e^{4i e} e^{-2i f x} + (i A a^2 c^2 f e^{2i e} - B a^2 c^2 f e^{2i e}) e^{-4i f x}) e^{-6i e}}{4 a^4 f^2} & \text{for } a^4 f^2 e^{6i e} \neq 0 \\ x \left(-\frac{2i B c^2}{a^2} + \frac{(A c^2 + 2i B c^2 e^{4i e} - 2i B c^2 e^{2i e} + i B c^2) e^{-4i e}}{a^2} \right) & \text{otherwise} \end{cases}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**2/(a+I*a*tan(f*x+e))**2,x)`

output `2*I*B*c**2*x/a**2 - B*c**2*log(exp(2*I*f*x) + exp(-2*I*e))/(a**2*f) + Piecewise(((4*B*a**2*c**2*f*exp(4*I*e)*exp(-2*I*f*x) + (I*A*a**2*c**2*f*exp(2*I*e) - B*a**2*c**2*f*exp(2*I*e))*exp(-4*I*f*x))*exp(-6*I*e)/(4*a**4*f**2), Ne(a**4*f**2*exp(6*I*e), 0)), (x*(-2*I*B*c**2/a**2 + (A*c**2 + 2*I*B*c**2*exp(4*I*e) - 2*I*B*c**2*exp(2*I*e) + I*B*c**2)*exp(-4*I*e)/a**2), True))`

3.719.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{(a + ia \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.719.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 191 vs. 2(84) = 168.

Time = 0.54 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.97

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{(a + ia \tan(e + fx))^2} dx = \frac{6 Bc^2 \log(\tan(\frac{1}{2} fx + \frac{1}{2} e) + 1)}{a^2} - \frac{12 Bc^2 \log(\tan(\frac{1}{2} fx + \frac{1}{2} e) - i)}{a^2} + \frac{6 Bc^2 \log(\tan(\frac{1}{2} fx + \frac{1}{2} e) - 1)}{a^2} + \frac{25 Bc^2 \tan(\frac{1}{2} fx + \frac{1}{2} e)^4 + 12 Ac^2 \tan(\frac{1}{2} fx + \frac{1}{2} e)^3 + 6 A^2 c^2 \tan(\frac{1}{2} fx + \frac{1}{2} e)^2 + 6 A^2 c^2 \tan(\frac{1}{2} fx + \frac{1}{2} e) + 6 A^2 c^2}{a^2}$$

6f

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output
$$\frac{-1/6*(6*B*c^2*\log(\tan(1/2*f*x + 1/2*e) + 1)/a^2 - 12*B*c^2*\log(\tan(1/2*f*x + 1/2*e) - 1)/a^2 + 6*B*c^2*\log(\tan(1/2*f*x + 1/2*e) - 1)/a^2 + (25*B*c^2*\tan(1/2*f*x + 1/2*e)^4 + 12*A*c^2*\tan(1/2*f*x + 1/2*e)^3 - 112*I*B*c^2*\tan(1/2*f*x + 1/2*e)^3 - 198*B*c^2*\tan(1/2*f*x + 1/2*e)^2 - 12*A*c^2*\tan(1/2*f*x + 1/2*e) + 112*I*B*c^2*\tan(1/2*f*x + 1/2*e) + 25*B*c^2)/(a^2*(\tan(1/2*f*x + 1/2*e) - I)^4))/f}$$

3.719.9 Mupad [B] (verification not implemented)

Time = 8.84 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.07

$$\int \frac{(A + B \tan(e + f x))(c - i c \tan(e + f x))^2}{(a + i a \tan(e + f x))^2} dx$$

$$= \frac{c^2 (2B + A \tan(e + f x) + B \tan(e + f x) 3i + B \ln(-1 - \tan(e + f x) 1i) - B \tan(e + f x)^2 \ln(-1 - \tan(e + f x) 1i))}{a^2 f (1 + \tan(e + f x) 1i)^2}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^2)/(a + a*tan(e + f*x)*1i)^2,x)`

output
$$(c^2*(2*B + A*\tan(e + f*x) + B*\tan(e + f*x)*3i + B*\log(-\tan(e + f*x)*1i - 1) - B*\tan(e + f*x)^2*\log(-\tan(e + f*x)*1i - 1) + B*\tan(e + f*x)*\log(-\tan(e + f*x)*1i - 1)*2i))/(a^2*f*(\tan(e + f*x)*1i + 1)^2)$$

3.720 $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))}{(a+ia \tan(e+fx))^2} dx$

3.720.1 Optimal result 6653
 3.720.2 Mathematica [A] (verified) 6653
 3.720.3 Rubi [A] (verified) 6654
 3.720.4 Maple [A] (verified) 6655
 3.720.5 Fricas [A] (verification not implemented) 6656
 3.720.6 Sympy [B] (verification not implemented) 6656
 3.720.7 Maxima [F(-2)] 6657
 3.720.8 Giac [A] (verification not implemented) 6657
 3.720.9 Mupad [B] (verification not implemented) 6657

3.720.1 Optimal result

Integrand size = 39, antiderivative size = 48

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx = -\frac{c(A + B \tan(e + fx))^2}{2a^2(iA - B)f(1 + i \tan(e + fx))^2}$$

output `-1/2*c*(A+B*tan(f*x+e))^2/a^2/(I*A-B)/f/(1+I*tan(f*x+e))^2`

3.720.2 Mathematica [A] (verified)

Time = 1.21 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.25

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx = \frac{c(A \cos(e + fx) + B \sin(e + fx))^2(i \cos(2(e + fx)) + \sin(2(e + fx)))}{2a^2(A + iB)f}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x]))/(a + I*a*Tan[e + f*x])^2,x]`

output `(c*(A*Cos[e + f*x] + B*Sin[e + f*x])^2*(I*Cos[2*(e + f*x)] + Sin[2*(e + f*x)]))/(2*a^2*(A + I*B)*f)`

3.720.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.103$, Rules used = {3042, 4071, 27, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c - ic \tan(e + fx))(A + B \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c - ic \tan(e + fx))(A + B \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{A+B \tan(e+fx)}{a^3(i \tan(e+fx)+1)^3} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{c \int \frac{A+B \tan(e+fx)}{(i \tan(e+fx)+1)^3} d \tan(e + fx)}{a^2 f} \\ & \quad \downarrow \text{48} \\ & -\frac{c(A + B \tan(e + fx))^2}{2a^2 f(-B + iA)(1 + i \tan(e + fx))^2} \end{aligned}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x]))/(a + I*a*Tan[e + f*x])^2, x]`

output `-1/2*(c*(A + B*Tan[e + f*x])^2)/(a^2*(I*A - B)*f*(1 + I*Tan[e + f*x])^2`

3.720.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.720.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.96

method	result	size
derivativedivides	$c \left(\frac{-\frac{iA-B}{2(-i+\tan(fx+e))^2} - \frac{iB}{-i+\tan(fx+e)}}{f a^2} \right)$	46
default	$c \left(\frac{-\frac{iA-B}{2(-i+\tan(fx+e))^2} - \frac{iB}{-i+\tan(fx+e)}}{f a^2} \right)$	46
risch	$\frac{c e^{-2i(fx+e)} B}{4a^2 f} + \frac{i c e^{-2i(fx+e)} A}{4a^2 f} - \frac{c e^{-4i(fx+e)} B}{8a^2 f} + \frac{i c e^{-4i(fx+e)} A}{8a^2 f}$	80
norman	$\frac{\frac{cA \tan(fx+e)}{af} + \frac{i c A + c B}{2af} + \frac{(-i c A + 3c B) \tan(fx+e)^2}{2af} - \frac{i c B \tan(fx+e)^3}{af}}{a(1+\tan(fx+e)^2)^2}$	95

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `1/f*c/a^2*(-1/2*(I*A-B)/(-I+tan(f*x+e))^2-I*B/(-I+tan(f*x+e)))`

3.720.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))}{(a+ia \tan(e+fx))^2} dx$$

3.720.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.98

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx$$

$$= -\frac{(2(-iA - B)ce^{2ifx+2ie}) - (iA - B)c)e^{(-4ifx-4ie)}}{8a^2f}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x, algo
rithm="fricas")`

output `-1/8*(2*(-I*A - B)*c*e^(2*I*f*x + 2*I*e) - (I*A - B)*c)*e^(-4*I*f*x - 4*I*
e)/(a^2*f)`

3.720.6 Sympy [B] (verification not implemented)Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 158 vs. $2(37) = 74$.

Time = 0.20 (sec) , antiderivative size = 158, normalized size of antiderivative = 3.29

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx$$

$$= \begin{cases} \frac{((4iAa^2cfe^{2ie} - 4Ba^2cfe^{2ie})e^{-4ifx} + (8iAa^2cfe^{4ie} + 8Ba^2cfe^{4ie})e^{-2ifx})e^{-6ie}}{32a^4f^2} & \text{for } a^4f^2e^{6ie} \neq 0 \\ \frac{x(Ace^{2ie} + Ac - iBce^{2ie} + iBc)e^{-4ie}}{2a^2} & \text{otherwise} \end{cases}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e))**2,x)`

output `Piecewise((((4*I*A*a**2*c*f*exp(2*I*e) - 4*B*a**2*c*f*exp(2*I*e))*exp(-4*I*
*f*x) + (8*I*A*a**2*c*f*exp(4*I*e) + 8*B*a**2*c*f*exp(4*I*e))*exp(-2*I*f*x
))*exp(-6*I*e)/(32*a**4*f**2), Ne(a**4*f**2*exp(6*I*e), 0)), (x*(A*c*exp(2
*I*e) + A*c - I*B*c*exp(2*I*e) + I*B*c)*exp(-4*I*e)/(2*a**2), True))`

3.720.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))}{(a + i a \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x, algo
rithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.720.8 Giac [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.65

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))}{(a + i a \tan(e + fx))^2} dx =$$

$$\frac{2 \left(A c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^3 - i A c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^2 - B c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^2 - A c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) \right)}{a^2 f \left(\tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) - i \right)^4}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x, algo
rithm="giac")
```

```
output -2*(A*c*tan(1/2*f*x + 1/2*e)^3 - I*A*c*tan(1/2*f*x + 1/2*e)^2 - B*c*tan(1/
2*f*x + 1/2*e)^2 - A*c*tan(1/2*f*x + 1/2*e))/(a^2*f*(tan(1/2*f*x + 1/2*e)
- I)^4)
```

3.720.9 Mupad [B] (verification not implemented)

Time = 8.31 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.04

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))}{(a + i a \tan(e + fx))^2} dx$$

$$= \frac{\frac{c(A-Bli)}{2} + B c \tan(e + fx)}{a^2 f \left(\tan(e + fx)^2 li + 2 \tan(e + fx) - i \right)}$$

3.720. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))}{(a+ia \tan(e+fx))^2} dx$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i))/(a + a*tan(e + f*x)*1i)
^2,x)`

output `((c*(A - B*1i))/2 + B*c*tan(e + f*x))/(a^2*f*(2*tan(e + f*x) + tan(e + f*x)
)^2*1i - 1i))`

3.721 $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2} dx$

3.721.1 Optimal result 6659
 3.721.2 Mathematica [A] (verified) 6659
 3.721.3 Rubi [A] (verified) 6660
 3.721.4 Maple [A] (verified) 6661
 3.721.5 Fricas [A] (verification not implemented) 6662
 3.721.6 Sympy [A] (verification not implemented) 6662
 3.721.7 Maxima [F(-2)] 6663
 3.721.8 Giac [A] (verification not implemented) 6663
 3.721.9 Mupad [B] (verification not implemented) 6663

3.721.1 Optimal result

Integrand size = 26, antiderivative size = 80

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2} dx = \frac{(A - iB)x}{4a^2} + \frac{iA - B}{4f(a + ia \tan(e + fx))^2} + \frac{iA + B}{4f(a^2 + ia^2 \tan(e + fx))}$$

output `1/4*(A-I*B)*x/a^2+1/4*(I*A-B)/f/(a+I*a*tan(f*x+e))^2+1/4*(I*A+B)/f/(a^2+I*a^2*tan(f*x+e))`

3.721.2 Mathematica [A] (verified)

Time = 0.80 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.11

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2} dx = \frac{(A - iB) \arctan(\tan(e + fx))}{4a^2 f} - \frac{iA - B}{4a^2 f(i - \tan(e + fx))^2} - \frac{A - iB}{4a^2 f(i - \tan(e + fx))}$$

input `Integrate[(A + B*Tan[e + f*x])/(a + I*a*Tan[e + f*x])^2,x]`

output `((A - I*B)*ArcTan[Tan[e + f*x]])/(4*a^2*f) - (I*A - B)/(4*a^2*f*(I - Tan[e + f*x])^2) - (A - I*B)/(4*a^2*f*(I - Tan[e + f*x]))`

3.721. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2} dx$

3.721.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.99, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$, Rules used = {3042, 4009, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4009} \\
 & \frac{(A - iB) \int \frac{1}{i \tan(e+fx)a+a} dx}{2a} + \frac{-B + iA}{4f(a + ia \tan(e + fx))^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A - iB) \int \frac{1}{i \tan(e+fx)a+a} dx}{2a} + \frac{-B + iA}{4f(a + ia \tan(e + fx))^2} \\
 & \quad \downarrow \text{3960} \\
 & \frac{(A - iB) \left(\frac{\int 1 dx}{2a} + \frac{i}{2f(a+ia \tan(e+fx))} \right)}{2a} + \frac{-B + iA}{4f(a + ia \tan(e + fx))^2} \\
 & \quad \downarrow \text{24} \\
 & \frac{-B + iA}{4f(a + ia \tan(e + fx))^2} + \frac{(A - iB) \left(\frac{x}{2a} + \frac{i}{2f(a+ia \tan(e+fx))} \right)}{2a}
 \end{aligned}$$

input `Int[(A + B*Tan[e + f*x])/(a + I*a*Tan[e + f*x])^2,x]`

output `(I*A - B)/(4*f*(a + I*a*Tan[e + f*x])^2) + ((A - I*B)*(x/(2*a) + (I/2)/(f*(a + I*a*Tan[e + f*x]))))/(2*a)`

3.721.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

3.721.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.91

method	result
risch	$-\frac{ixB}{4a^2} + \frac{xA}{4a^2} + \frac{iAe^{-2i(fx+e)}}{4a^2f} - \frac{e^{-4i(fx+e)}B}{16a^2f} + \frac{ie^{-4i(fx+e)}A}{16a^2f}$
derivativedivides	$\frac{B}{4fa^2(-i+\tan(fx+e))^2} - \frac{iA}{4fa^2(-i+\tan(fx+e))^2} + \frac{A}{4fa^2(-i+\tan(fx+e))} - \frac{iB}{4fa^2(-i+\tan(fx+e))} + \frac{A \arctan(\dots)}{4fa^2(-i+\tan(fx+e))}$
default	$\frac{B}{4fa^2(-i+\tan(fx+e))^2} - \frac{iA}{4fa^2(-i+\tan(fx+e))^2} + \frac{A}{4fa^2(-i+\tan(fx+e))} - \frac{iB}{4fa^2(-i+\tan(fx+e))} + \frac{A \arctan(\dots)}{4fa^2(-i+\tan(fx+e))}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `-1/4*I*x/a^2*B+1/4*x/a^2*A+1/4*I*A/a^2/f*exp(-2*I*(f*x+e))-1/16/a^2/f*exp(-4*I*(f*x+e))*B+1/16*I/a^2/f*exp(-4*I*(f*x+e))*A`

3.721.
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2} dx$$

3.721.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.68

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{(4(A - iB)fx e^{(4i fx + 4i e)} + 4i A e^{(2i fx + 2i e)} + i A - B) e^{(-4i fx - 4i e)}}{16 a^2 f}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`output `1/16*(4*(A - I*B)*f*x*e^(4*I*f*x + 4*I*e) + 4*I*A*e^(2*I*f*x + 2*I*e) + I*A - B)*e^(-4*I*f*x - 4*I*e)/(a^2*f)`**3.721.6 Sympy [A] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 162, normalized size of antiderivative = 2.02

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2} dx$$

$$= \begin{cases} \frac{(16iAa^2 f e^{4ie} e^{-2ifx} + (4iAa^2 f e^{2ie} - 4Ba^2 f e^{2ie}) e^{-4ifx}) e^{-6ie}}{64a^4 f^2} & \text{for } a^4 f^2 e^{6ie} \neq 0 \\ x \left(-\frac{A-iB}{4a^2} + \frac{(Ae^{4ie} + 2Ae^{2ie} + A - iBe^{4ie} + iB) e^{-4ie}}{4a^2} \right) & \text{otherwise} \end{cases} + \frac{x(A - iB)}{4a^2}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**2,x)`output `Piecewise((((16*I*A*a**2*f*exp(4*I*e)*exp(-2*I*f*x) + (4*I*A*a**2*f*exp(2*I*e) - 4*B*a**2*f*exp(2*I*e))*exp(-4*I*f*x))*exp(-6*I*e)/(64*a**4*f**2), Ne(a**4*f**2*exp(6*I*e), 0)), (x*(-(A - I*B)/(4*a**2) + (A*exp(4*I*e) + 2*A*exp(2*I*e) + A - I*B*exp(4*I*e) + I*B)*exp(-4*I*e)/(4*a**2)), True)) + x*(A - I*B)/(4*a**2)`

3.721.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.721.8 Giac [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.38

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2} dx = \frac{\frac{2(-iA-B)\log(\tan(fx+e)+i)}{a^2} - \frac{2(-iA-B)\log(\tan(fx+e)-i)}{a^2} - \frac{3iA \tan(fx+e)^2 + 3B \tan(fx+e)^2 + 10A \tan(fx+e) - 10iB \tan(fx+e)}{a^2(\tan(fx+e)-i)^2}}{16f}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")
```

```
output -1/16*(2*(-I*A - B)*log(tan(f*x + e) + I)/a^2 - 2*(-I*A - B)*log(tan(f*x + e) - I)/a^2 - (3*I*A*tan(f*x + e)^2 + 3*B*tan(f*x + e)^2 + 10*A*tan(f*x + e) - 10*I*B*tan(f*x + e) - 11*I*A - 3*B)/(a^2*(tan(f*x + e) - I)^2))/f
```

3.721.9 Mupad [B] (verification not implemented)

Time = 8.29 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.88

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2} dx = \frac{\frac{A}{2a^2} + \tan(e + fx) \left(\frac{B}{4a^2} + \frac{A1i}{4a^2} \right)}{f (\tan(e + fx)^2 1i + 2 \tan(e + fx) - i)} - \frac{x (B + A 1i) 1i}{4a^2}$$

```
input int((A + B*tan(e + f*x))/(a + a*tan(e + f*x)*1i)^2,x)
```

```
output (A/(2*a^2) + tan(e + f*x)*((A*1i)/(4*a^2) + B/(4*a^2)))/(f*(2*tan(e + f*x) + tan(e + f*x)^2*1i - 1i)) - (x*(A*1i + B)*1i)/(4*a^2)
```

3.721. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2} dx$

3.722
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))} dx$$

3.722.1 Optimal result 6664
 3.722.2 Mathematica [A] (verified) 6664
 3.722.3 Rubi [A] (verified) 6665
 3.722.4 Maple [A] (verified) 6667
 3.722.5 Fricas [A] (verification not implemented) 6667
 3.722.6 Sympy [A] (verification not implemented) 6668
 3.722.7 Maxima [F(-2)] 6668
 3.722.8 Giac [A] (verification not implemented) 6669
 3.722.9 Mupad [B] (verification not implemented) 6669

3.722.1 Optimal result

Integrand size = 41, antiderivative size = 117

$$\begin{aligned} & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2(c - ic \tan(e + fx))} dx \\ &= \frac{(3A - iB)x}{8a^2c} - \frac{iA - B}{8a^2cf(i - \tan(e + fx))^2} \\ & \quad - \frac{A}{4a^2cf(i - \tan(e + fx))} + \frac{A - iB}{8a^2cf(i + \tan(e + fx))} \end{aligned}$$

output `1/8*(3*A-I*B)*x/a^2/c+1/8*(-I*A+B)/a^2/c/f/(I-tan(f*x+e))^2-1/4*A/a^2/c/f/(I-tan(f*x+e))+1/8*(A-I*B)/a^2/c/f/(I+tan(f*x+e))`

3.722.2 Mathematica [A] (verified)

Time = 5.69 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.05

$$\begin{aligned} & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2(c - ic \tan(e + fx))} dx \\ &= \frac{\sec^2(e + fx)(5A + iB - (A - 3iB) \cos(2(e + fx)) - 3iA \sin(2(e + fx)) - B \sin(2(e + fx)) + 2(3A - iB) \tan(e + fx))}{16a^2cf(-i + \tan(e + fx))^2(i + \tan(e + fx))} \end{aligned}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])),x]`

output $(\text{Sec}[e + f*x]^2*(5*A + I*B - (A - (3*I)*B)*\text{Cos}[2*(e + f*x)] - (3*I)*A*\text{Sin}[2*(e + f*x)] - B*\text{Sin}[2*(e + f*x)] + 2*(3*A - I*B)*\text{ArcTan}[\text{Tan}[e + f*x]]*(-I + \text{Tan}[e + f*x]))/(16*a^2*c*f*(-I + \text{Tan}[e + f*x])^2*(I + \text{Tan}[e + f*x]))$

3.722.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.85, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))} dx$$

↓ 4071

$$\frac{ac \int \frac{A + B \tan(e + fx)}{a^3 c^2 (1 - i \tan(e + fx))^2 (i \tan(e + fx) + 1)^3} d \tan(e + fx)}{f}$$

↓ 27

$$\frac{\int \frac{A + B \tan(e + fx)}{(1 - i \tan(e + fx))^2 (i \tan(e + fx) + 1)^3} d \tan(e + fx)}{a^2 c f}$$

↓ 86

$$\frac{\int \left(-\frac{A}{4(\tan(e + fx) - i)^2} + \frac{3A - iB}{8(\tan^2(e + fx) + 1)} + \frac{iB - A}{8(\tan(e + fx) + i)^2} + \frac{i(A + iB)}{4(\tan(e + fx) - i)^3} \right) d \tan(e + fx)}{a^2 c f}$$

↓ 2009

$$\frac{\frac{1}{8}(3A - iB) \arctan(\tan(e + fx)) + \frac{A - iB}{8(\tan(e + fx) + i)} - \frac{-B + iA}{8(-\tan(e + fx) + i)^2} - \frac{A}{4(-\tan(e + fx) + i)}}{a^2 c f}$$

input $\text{Int}[(A + B*\text{Tan}[e + f*x])/((a + I*a*\text{Tan}[e + f*x])^2*(c - I*c*\text{Tan}[e + f*x])),x]$


```
output ((3*A - I*B)*ArcTan[Tan[e + f*x]]/8 - (I*A - B)/(8*(I - Tan[e + f*x])^2)
- A/(4*(I - Tan[e + f*x])) + (A - I*B)/(8*(I + Tan[e + f*x])))/(a^2*c*f)
```

3.722.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.722.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.11

method	result
risch	$-\frac{ixB}{8a^2c} + \frac{3xA}{8a^2c} - \frac{e^{-4i(fx+e)}B}{32a^2cf} + \frac{ie^{-4i(fx+e)}A}{32a^2cf} - \frac{\cos(2fx+2e)B}{8a^2cf} + \frac{i\cos(2fx+2e)A}{8a^2cf} + \frac{A\sin(2fx+2e)}{4a^2cf}$
norman	$\frac{\frac{(-iB+3A)x}{8ac} - \frac{-iA+B}{4acf} + \frac{(-iB+3A)\tan(fx+e)^3}{8acf} + \frac{(-iB+3A)x\tan(fx+e)^2}{4ac} + \frac{(-iB+3A)x\tan(fx+e)^4}{8ac} + \frac{(iB+5A)\tan(fx+e)}{8acf}}{a(1+\tan(fx+e))^2}$
derivativedivides	$\frac{3A\arctan(\tan(fx+e))}{8fa^2c} - \frac{iB\arctan(\tan(fx+e))}{8fa^2c} + \frac{A}{8fa^2c(i+\tan(fx+e))} - \frac{iB}{8fa^2c(i+\tan(fx+e))} + \frac{A}{4fa^2c(-i+\tan(fx+e))} - \frac{iB}{4fa^2c(-i+\tan(fx+e))}$
default	$\frac{3A\arctan(\tan(fx+e))}{8fa^2c} - \frac{iB\arctan(\tan(fx+e))}{8fa^2c} + \frac{A}{8fa^2c(i+\tan(fx+e))} - \frac{iB}{8fa^2c(i+\tan(fx+e))} + \frac{A}{4fa^2c(-i+\tan(fx+e))} - \frac{iB}{4fa^2c(-i+\tan(fx+e))}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output `-1/8*I*x/a^2/c*B+3/8*x/a^2/c*A-1/32/a^2/c/f*exp(-4*I*(f*x+e))*B+1/32*I/a^2/c/f*exp(-4*I*(f*x+e))*A-1/8/a^2/c/f*cos(2*f*x+2*e)*B+1/8*I/a^2/c/f*cos(2*f*x+2*e)*A+1/4*A/a^2/c/f*sin(2*f*x+2*e)`

3.722.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.68

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ictan(e + fx))} dx$$

$$= \frac{(4(3A - iB)fx e^{(4i fx + 4i e)} - 2(iA + B)e^{(6i fx + 6i e)} - 2(-3iA + B)e^{(2i fx + 2i e)} + iA - B)e^{(-4i fx - 4i e)}}{32 a^2 c f}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e)),x,algorithm="fracas")`

output `1/32*(4*(3*A - I*B)*f*x*e^(4*I*f*x + 4*I*e) - 2*(I*A + B)*e^(6*I*f*x + 6*I*e) - 2*(-3*I*A + B)*e^(2*I*f*x + 2*I*e) + I*A - B)*e^(-4*I*f*x - 4*I*e)/(a^2*c*f)`

3.722.6 Sympy [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 296, normalized size of antiderivative = 2.53

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ictan(e + fx))} dx$$

$$= \begin{cases} \frac{((256iAa^4c^2f^2e^{2ie} - 256Ba^4c^2f^2e^{2ie})e^{-4ifx} + (1536iAa^4c^2f^2e^{4ie} - 512Ba^4c^2f^2e^{4ie})e^{-2ifx} + (-512iAa^4c^2f^2e^{8ie} - 512Ba^4c^2f^2e^{8ie})e^{2ifx})}{8192a^6c^3f^3} \\ x \left(-\frac{3A-iB}{8a^2c} + \frac{(Ae^{6ie} + 3Ae^{4ie} + 3Ae^{2ie} + A - iBe^{6ie} - iBe^{4ie} + iBe^{2ie} + iB)e^{-4ie}}{8a^2c} \right) \\ + \frac{x(3A - iB)}{8a^2c} \end{cases}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**2/(c-I*c*tan(f*x+e)),x)`

output `Piecewise((((256*I*A*a**4*c**2*f**2*exp(2*I*e) - 256*B*a**4*c**2*f**2*exp(2*I*e))*exp(-4*I*f*x) + (1536*I*A*a**4*c**2*f**2*exp(4*I*e) - 512*B*a**4*c**2*f**2*exp(4*I*e))*exp(-2*I*f*x) + (-512*I*A*a**4*c**2*f**2*exp(8*I*e) - 512*B*a**4*c**2*f**2*exp(8*I*e))*exp(2*I*f*x))*exp(-6*I*e)/(8192*a**6*c**3*f**3), Ne(a**6*c**3*f**3*exp(6*I*e), 0)), (x*(-(3*A - I*B)/(8*a**2*c) + (A*exp(6*I*e) + 3*A*exp(4*I*e) + 3*A*exp(2*I*e) + A - I*B*exp(6*I*e) - I*B*exp(4*I*e) + I*B*exp(2*I*e) + I*B)*exp(-4*I*e)/(8*a**2*c)), True)) + x*(3*A - I*B)/(8*a**2*c)`

3.722.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ictan(e + fx))} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e)),x, algo rithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.722.8 Giac [A] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.36

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ictan(e + fx))} dx$$

$$= \frac{\frac{2(3iA+B)\log(\tan(fx+e)+i)}{a^2c} + \frac{2(-3iA-B)\log(\tan(fx+e)-i)}{a^2c} - \frac{2(3A\tan(fx+e)-iB\tan(fx+e)+5iA+3B)}{a^2c(-i\tan(fx+e)+1)} + \frac{9iA\tan(fx+e)^2+3B\tan(fx+e)}{32f}}{32f}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e)),x, algo rithm="giac")`

output `1/32*(2*(3*I*A + B)*log(tan(f*x + e) + I)/(a^2*c) + 2*(-3*I*A - B)*log(tan(f*x + e) - I)/(a^2*c) - 2*(3*A*tan(f*x + e) - I*B*tan(f*x + e) + 5*I*A + 3*B)/(a^2*c*(-I*tan(f*x + e) + 1)) + (9*I*A*tan(f*x + e)^2 + 3*B*tan(f*x + e)^2 + 26*A*tan(f*x + e) - 6*I*B*tan(f*x + e) - 21*I*A + B)/(a^2*c*(tan(f*x + e) - I)^2))/f`

3.722.9 Mupad [B] (verification not implemented)

Time = 8.59 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.10

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ictan(e + fx))} dx$$

$$= \frac{\tan(e + fx) \left(\frac{3A}{8a^2c} - \frac{B1i}{8a^2c} \right) + \tan(e + fx)^2 \left(\frac{B}{8a^2c} + \frac{A3i}{8a^2c} \right) - \frac{B}{4a^2c} + \frac{A1i}{4a^2c}}{f \left(\tan(e + fx)^3 1i + \tan(e + fx)^2 + \tan(e + fx) 1i + 1 \right)} - \frac{x(B + A3i) 1i}{8a^2c}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)),x)`

output `(tan(e + f*x)*((3*A)/(8*a^2*c) - (B*1i)/(8*a^2*c)) + tan(e + f*x)^2*((A*3i)/(8*a^2*c) + B/(8*a^2*c)) + (A*1i)/(4*a^2*c) - B/(4*a^2*c))/(f*(tan(e + f*x)*1i + tan(e + f*x)^2 + tan(e + f*x)^3*1i + 1)) - (x*(A*3i + B)*1i)/(8*a^2*c)`

3.723
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^2} dx$$

3.723.1 Optimal result 6670
 3.723.2 Mathematica [A] (verified) 6670
 3.723.3 Rubi [A] (verified) 6671
 3.723.4 Maple [A] (verified) 6673
 3.723.5 Fricas [C] (verification not implemented) 6673
 3.723.6 Sympy [A] (verification not implemented) 6674
 3.723.7 Maxima [F(-2)] 6674
 3.723.8 Giac [A] (verification not implemented) 6675
 3.723.9 Mupad [B] (verification not implemented) 6675

3.723.1 Optimal result

Integrand size = 41, antiderivative size = 71

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2(c - ic \tan(e + fx))^2} dx$$

$$= \frac{3Ax}{8a^2c^2} + \frac{3A \cos(e + fx) \sin(e + fx)}{8a^2c^2f} - \frac{\cos^4(e + fx)(B - A \tan(e + fx))}{4a^2c^2f}$$

output `3/8*A*x/a^2/c^2+3/8*A*cos(f*x+e)*sin(f*x+e)/a^2/c^2/f-1/4*cos(f*x+e)^4*(B-A*tan(f*x+e))/a^2/c^2/f`

3.723.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.75

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2(c - ic \tan(e + fx))^2} dx$$

$$= \frac{-8B \cos^4(e + fx) + A(12(e + fx) + 8 \sin(2(e + fx)) + \sin(4(e + fx)))}{32a^2c^2f}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^2),x]`

output `(-8*B*Cos[e + f*x]^4 + A*(12*(e + f*x) + 8*Sin[2*(e + f*x)] + Sin[4*(e + f*x)]))/(32*a^2*c^2*f)`

3.723.
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^2} dx$$

3.723.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.08, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.171$, Rules used = {3042, 4071, 27, 82, 454, 215, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A+B \tan(e+fx)}{a^3 c^3 (1-i \tan(e+fx))^3 (i \tan(e+fx)+1)^3} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{A+B \tan(e+fx)}{(1-i \tan(e+fx))^3 (i \tan(e+fx)+1)^3} d \tan(e + fx)}{a^2 c^2 f} \\
 & \quad \downarrow \text{82} \\
 & \frac{\int \frac{A+B \tan(e+fx)}{(\tan^2(e+fx)+1)^3} d \tan(e + fx)}{a^2 c^2 f} \\
 & \quad \downarrow \text{454} \\
 & \frac{\frac{3}{4} A \int \frac{1}{(\tan^2(e+fx)+1)^2} d \tan(e + fx) - \frac{B-A \tan(e+fx)}{4(\tan^2(e+fx)+1)^2}}{a^2 c^2 f} \\
 & \quad \downarrow \text{215} \\
 & \frac{\frac{3}{4} A \left(\frac{1}{2} \int \frac{1}{\tan^2(e+fx)+1} d \tan(e + fx) + \frac{\tan(e+fx)}{2(\tan^2(e+fx)+1)} \right) - \frac{B-A \tan(e+fx)}{4(\tan^2(e+fx)+1)^2}}{a^2 c^2 f} \\
 & \quad \downarrow \text{216} \\
 & \frac{\frac{3}{4} A \left(\frac{1}{2} \arctan(\tan(e + fx)) + \frac{\tan(e+fx)}{2(\tan^2(e+fx)+1)} \right) - \frac{B-A \tan(e+fx)}{4(\tan^2(e+fx)+1)^2}}{a^2 c^2 f}
 \end{aligned}$$

3.723. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 (c-ic \tan(e+fx))^2} dx$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^2),x]`

output `(-1/4*(B - A*Tan[e + f*x])/(1 + Tan[e + f*x]^2)^2 + (3*A*(ArcTan[Tan[e + f*x]])/2 + Tan[e + f*x]/(2*(1 + Tan[e + f*x]^2))))/4/(a^2*c^2*f)`

3.723.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 82 `Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 215 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 454 `Int[((c_) + (d_.)*(x_))*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[((a*d - b*c*x)/(2*a*b*(p + 1))*(a + b*x^2)^(p + 1), x] + Simp[c*((2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d}, x] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.723.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.35

method	result
risch	$\frac{3Ax}{8a^2c^2} - \frac{B \cos(4fx+4e)}{32a^2c^2f} + \frac{A \sin(4fx+4e)}{32a^2c^2f} - \frac{B \cos(2fx+2e)}{8a^2c^2f} + \frac{A \sin(2fx+2e)}{4a^2c^2f}$
norman	$\frac{\frac{3Ax}{8ac} - \frac{B}{4acf} + \frac{5A \tan(fx+e)}{8acf} + \frac{3A \tan(fx+e)^3}{8acf} + \frac{3Ax \tan(fx+e)^2}{4ac} + \frac{3Ax \tan(fx+e)^4}{8ac}}{(1+\tan(fx+e))^2} ac$
derivativedivides	$\frac{3A}{16f a^2c^2(-i+\tan(fx+e))} + \frac{iB}{16f a^2c^2(-i+\tan(fx+e))} - \frac{iA}{16f a^2c^2(-i+\tan(fx+e))^2} + \frac{B}{16f a^2c^2(-i+\tan(fx+e))}$
default	$\frac{3A}{16f a^2c^2(-i+\tan(fx+e))} + \frac{iB}{16f a^2c^2(-i+\tan(fx+e))} - \frac{iA}{16f a^2c^2(-i+\tan(fx+e))^2} + \frac{B}{16f a^2c^2(-i+\tan(fx+e))}$

```
input int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^2,x,method=_R
ETURNVERBOSE)
```

```
output 3/8*A*x/a^2/c^2-1/32*B/a^2/c^2/f*cos(4*f*x+4*e)+1/32*A/a^2/c^2/f*sin(4*f*x
+4*e)-1/8*B/a^2/c^2/f*cos(2*f*x+2*e)+1/4*A/a^2/c^2/f*sin(2*f*x+2*e)
```

3.723.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.27

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2(c - ic \tan(e + fx))^2} dx$$

$$= \frac{(24 A f x e^{(4i f x + 4i e)} + (-i A - B) e^{(8i f x + 8i e)} - 4(2i A + B) e^{(6i f x + 6i e)} - 4(-2i A + B) e^{(2i f x + 2i e)} + i A - i B)}{64 a^2 c^2 f}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^2,x, al
gorithm="fracas")
```

3.723. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^2} dx$

output $\frac{1}{64} \cdot (24 \cdot A \cdot f \cdot x \cdot e^{(4 \cdot I \cdot f \cdot x + 4 \cdot I \cdot e)} + (-I \cdot A - B) \cdot e^{(8 \cdot I \cdot f \cdot x + 8 \cdot I \cdot e)} - 4 \cdot (2 \cdot I \cdot A + B) \cdot e^{(6 \cdot I \cdot f \cdot x + 6 \cdot I \cdot e)} - 4 \cdot (-2 \cdot I \cdot A + B) \cdot e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} + I \cdot A - B) \cdot e^{(-4 \cdot I \cdot f \cdot x - 4 \cdot I \cdot e)} / (a^2 \cdot c^2 \cdot f)$

3.723.6 Sympy [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 360, normalized size of antiderivative = 5.07

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^2} dx = \frac{3Ax}{8a^2c^2} + \left\{ \frac{((16384iAa^6c^6f^3e^{2ie} - 16384Ba^6c^6f^3e^{2ie})e^{-4ifx} + (131072iAa^6c^6f^3e^{4ie} - 65536Ba^6c^6f^3e^{4ie})e^{-2ifx} + (-131072iAa^6c^6f^3e^{8ie} - 65536Ba^6c^6f^3e^{8ie})e^{-4ifx})}{1048576a^8c^8f^4} x \left(-\frac{3A}{8a^2c^2} + \frac{(Ae^{8ie} + 4Ae^{6ie} + 6Ae^{4ie} + 4Ae^{2ie} + A - iBe^{8ie} - 2iBe^{6ie} + 2iBe^{2ie} + iB)e^{-4ie}}{16a^2c^2} \right) \right\}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**2/(c-I*c*tan(f*x+e))**2,x)`

output `3*A*x/(8*a**2*c**2) + Piecewise((((16384*I*A*a**6*c**6*f**3*exp(2*I*e) - 16384*B*a**6*c**6*f**3*exp(2*I*e))*exp(-4*I*f*x) + (131072*I*A*a**6*c**6*f**3*exp(4*I*e) - 65536*B*a**6*c**6*f**3*exp(4*I*e))*exp(-2*I*f*x) + (-131072*I*A*a**6*c**6*f**3*exp(8*I*e) - 65536*B*a**6*c**6*f**3*exp(8*I*e))*exp(2*I*f*x) + (-16384*I*A*a**6*c**6*f**3*exp(10*I*e) - 16384*B*a**6*c**6*f**3*exp(10*I*e))*exp(4*I*f*x))*exp(-6*I*e)/(1048576*a**8*c**8*f**4), Ne(a**8*c**8*f**4*exp(6*I*e), 0)), (x*(-3*A/(8*a**2*c**2) + (A*exp(8*I*e) + 4*A*exp(6*I*e) + 6*A*exp(4*I*e) + 4*A*exp(2*I*e) + A - I*B*exp(8*I*e) - 2*I*B*exp(6*I*e) + 2*I*B*exp(2*I*e) + I*B)*exp(-4*I*e)/(16*a**2*c**2)), True))`

3.723.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.723. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 (c-ic \tan(e+fx))^2} dx$

3.723.8 Giac [A] (verification not implemented)

Time = 0.56 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.89

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^2} dx = \frac{\frac{3(fx+e)A}{a^2 c^2} + \frac{3A \tan(fx+e)^3 + 5A \tan(fx+e) - 2B}{(\tan(fx+e)^2 + 1)^2 a^2 c^2}}{8f}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^2,x, algorithm="giac")`

output `1/8*(3*(f*x + e)*A/(a^2*c^2) + (3*A*tan(f*x + e)^3 + 5*A*tan(f*x + e) - 2*B)/((tan(f*x + e)^2 + 1)^2*a^2*c^2))/f`

3.723.9 Mupad [B] (verification not implemented)

Time = 8.62 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.75

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^2} dx$$

$$= \frac{3Ax}{8a^2 c^2} + \frac{\cos(e + fx)^4 \left(\frac{3A \tan(e+fx)^3}{8} + \frac{5A \tan(e+fx)}{8} - \frac{B}{4} \right)}{a^2 c^2 f}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)^2),x)`

output `(3*A*x)/(8*a^2*c^2) + (cos(e + f*x)^4*((5*A*tan(e + f*x))/8 - B/4 + (3*A*tan(e + f*x)^3)/8))/(a^2*c^2*f)`

$$3.724 \quad \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^3} dx$$

3.724.1 Optimal result	6676
3.724.2 Mathematica [A] (verified)	6676
3.724.3 Rubi [A] (verified)	6677
3.724.4 Maple [A] (verified)	6679
3.724.5 Fricas [A] (verification not implemented)	6679
3.724.6 Sympy [A] (verification not implemented)	6680
3.724.7 Maxima [F(-2)]	6680
3.724.8 Giac [A] (verification not implemented)	6681
3.724.9 Mupad [B] (verification not implemented)	6681

3.724.1 Optimal result

Integrand size = 41, antiderivative size = 183

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2(c - ic \tan(e + fx))^3} dx$$

$$= \frac{(5A + iB)x}{16a^2c^3} - \frac{iA - B}{32a^2c^3 f(i - \tan(e + fx))^2} - \frac{2A + iB}{16a^2c^3 f(i - \tan(e + fx))}$$

$$- \frac{A - iB}{24a^2c^3 f(i + \tan(e + fx))^3} + \frac{3iA + B}{32a^2c^3 f(i + \tan(e + fx))^2} + \frac{3A}{16a^2c^3 f(i + \tan(e + fx))}$$

output `1/16*(5*A+I*B)*x/a^2/c^3+1/32*(-I*A+B)/a^2/c^3/f/(I-tan(f*x+e))^2+1/16*(-2*A-I*B)/a^2/c^3/f/(I-tan(f*x+e))+1/24*(-A+I*B)/a^2/c^3/f/(I+tan(f*x+e))^3+1/32*(3*I*A+B)/a^2/c^3/f/(I+tan(f*x+e))^2+3/16*A/a^2/c^3/f/(I+tan(f*x+e))`

3.724.2 Mathematica [A] (verified)

Time = 6.00 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.95

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2(c - ic \tan(e + fx))^3} dx$$

$$= \frac{\sec^4(e + fx)(47A - 5iB - 2(7A + 11iB) \cos(2(e + fx)) - A \cos(4(e + fx)) - 5iB \cos(4(e + fx)) + 40i)}{192a^2c^3 f($$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^3),x]`

3.724. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^3} dx$

output $(\text{Sec}[e + f*x]^4*(47*A - (5*I)*B - 2*(7*A + (11*I)*B)*\text{Cos}[2*(e + f*x)] - A*\text{Cos}[4*(e + f*x)] - (5*I)*B*\text{Cos}[4*(e + f*x)] + (40*I)*A*\text{Sin}[2*(e + f*x)] - 8*B*\text{Sin}[2*(e + f*x)] + (5*I)*A*\text{Sin}[4*(e + f*x)] - B*\text{Sin}[4*(e + f*x)] + 12*(5*A + I*B)*\text{ArcTan}[\text{Tan}[e + f*x]]*(I + \text{Tan}[e + f*x]))/(192*a^2*c^3*f*(-I + \text{Tan}[e + f*x])^2*(I + \text{Tan}[e + f*x])^3)$

3.724.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.81, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^3} dx$$

↓ 4071

$$\frac{ac \int \frac{A + B \tan(e + fx)}{a^3 c^4 (1 - i \tan(e + fx))^4 (i \tan(e + fx) + 1)^3} d \tan(e + fx)}{f}$$

↓ 27

$$\int \frac{A + B \tan(e + fx)}{(1 - i \tan(e + fx))^4 (i \tan(e + fx) + 1)^3} d \tan(e + fx)}{a^2 c^3 f}$$

↓ 86

$$\int \left(-\frac{3A}{16(\tan(e + fx) + i)^2} + \frac{5A + iB}{16(\tan^2(e + fx) + 1)} + \frac{-2A - iB}{16(\tan(e + fx) - i)^2} + \frac{i(A + iB)}{16(\tan(e + fx) - i)^3} - \frac{i(3A - iB)}{16(\tan(e + fx) + i)^3} + \frac{A - iB}{8(\tan(e + fx) + i)^4} \right) d \tan(e + fx)}{a^2 c^3 f}$$

↓ 2009

$$\frac{\frac{1}{16}(5A + iB) \arctan(\tan(e + fx)) - \frac{2A + iB}{16(-\tan(e + fx) + i)} - \frac{-B + iA}{32(-\tan(e + fx) + i)^2} + \frac{B + 3iA}{32(\tan(e + fx) + i)^2} - \frac{A - iB}{24(\tan(e + fx) + i)^3} + \frac{1}{16}}{a^2 c^3 f}$$

3.724. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^3} dx$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^3),x]`

output `((5*A + I*B)*ArcTan[Tan[e + f*x]]/16 - (I*A - B)/(32*(I - Tan[e + f*x])^2) - (2*A + I*B)/(16*(I - Tan[e + f*x])) - (A - I*B)/(24*(I + Tan[e + f*x])^3) + ((3*I)*A + B)/(32*(I + Tan[e + f*x])^2) + (3*A)/(16*(I + Tan[e + f*x]))) / (a^2*c^3*f)`

3.724.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.724.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.14

method	result
norman	$\frac{\frac{(iB+5A)x}{16ac} - \frac{iA+B}{6acf} + \frac{(-iB+11A)\tan(fx+e)}{16acf} + \frac{(iB+5A)\tan(fx+e)^3}{6acf} + \frac{(iB+5A)\tan(fx+e)^5}{16acf} + \frac{3(iB+5A)x\tan(fx+e)^2}{16ac} + \frac{3(iB+5A)}{16ac}}{(1+\tan(fx+e))^3 a c^2}$
risch	$\frac{i x B}{16 a^2 c^3} + \frac{5 x A}{16 a^2 c^3} - \frac{e^{6 i(f x+e)} B}{192 a^2 c^3 f} - \frac{i e^{6 i(f x+e)} A}{192 a^2 c^3 f} - \frac{\cos(4 f x+4 e) B}{32 a^2 c^3 f} - \frac{i \cos(4 f x+4 e) A}{32 a^2 c^3 f} - \frac{i \sin(4 f x+4 e) B}{64 a^2 c^3 f} + \frac{3 B}{24 f a^2 c^3}$
derivativedivides	$\frac{i B}{16 f a^2 c^3(-i+\tan(f x+e))} + \frac{A}{8 f a^2 c^3(-i+\tan(f x+e))} + \frac{i B \arctan(\tan(f x+e))}{16 f a^2 c^3} + \frac{5 A \arctan(\tan(f x+e))}{16 f a^2 c^3} + \frac{3 B}{24 f a^2 c^3}$
default	$\frac{i B}{16 f a^2 c^3(-i+\tan(f x+e))} + \frac{A}{8 f a^2 c^3(-i+\tan(f x+e))} + \frac{i B \arctan(\tan(f x+e))}{16 f a^2 c^3} + \frac{5 A \arctan(\tan(f x+e))}{16 f a^2 c^3} + \frac{3 B}{24 f a^2 c^3}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^3,x,method=_R
ETURNVERBOSE)`

output `(1/16*(5*A+I*B)/a/c*x-1/6*(I*A+B)/a/c/f+1/16*(-I*B+11*A)/a/c/f*tan(f*x+e)+
1/6*(5*A+I*B)/a/c/f*tan(f*x+e)^3+1/16*(5*A+I*B)/a/c/f*tan(f*x+e)^5+3/16*(5
*A+I*B)/a/c*x*tan(f*x+e)^2+3/16*(5*A+I*B)/a/c*x*tan(f*x+e)^4+1/16*(5*A+I*B
)/a/c*x*tan(f*x+e)^6)/(1+tan(f*x+e)^2)^3/a/c^2`

3.724.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.63

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^3} dx$$

$$= \frac{(24(5A + iB)fx e^{(4i fx + 4ie)} - 2(iA + B)e^{(10i fx + 10ie)} - 3(5iA + 3B)e^{(8i fx + 8ie)} - 12(5iA + B)e^{(6i fx + 6ie)})}{384 a^2 c^3 f}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^3,x, al
gorithm="fricas")`

output `1/384*(24*(5*A + I*B)*f*x*e^(4*I*f*x + 4*I*e) - 2*(I*A + B)*e^(10*I*f*x +
10*I*e) - 3*(5*I*A + 3*B)*e^(8*I*f*x + 8*I*e) - 12*(5*I*A + B)*e^(6*I*f*x
+ 6*I*e) - 6*(-5*I*A + 3*B)*e^(2*I*f*x + 2*I*e) + 3*I*A - 3*B)*e^(-4*I*f*x
- 4*I*e)/(a^2*c^3*f)`

3.724.
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^3} dx$$

3.724.6 Sympy [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 454, normalized size of antiderivative = 2.48

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^3} dx$$

$$= \left\{ \begin{array}{l} \left((50331648iAa^8c^{12}f^4e^{2ie} - 50331648Ba^8c^{12}f^4e^{2ie})e^{-4ifx} + (503316480iAa^8c^{12}f^4e^{4ie} - 301989888Ba^8c^{12}f^4e^{4ie})e^{-2ifx} + (-1006632960iAa^8c^{12}f^4e^{6ie} + 503316480iAa^8c^{12}f^4e^{6ie})e^{-ifx} + (-1006632960iAa^8c^{12}f^4e^{8ie} + 503316480iAa^8c^{12}f^4e^{8ie})e^{-ifx} + (-1006632960iAa^8c^{12}f^4e^{10ie} + 503316480iAa^8c^{12}f^4e^{10ie})e^{-ifx} + (-1006632960iAa^8c^{12}f^4e^{12ie} + 503316480iAa^8c^{12}f^4e^{12ie})e^{-ifx} \right) \\ x \left(-\frac{5A+iB}{16a^2c^3} + \frac{(Ae^{10ie} + 5Ae^{8ie} + 10Ae^{6ie} + 10Ae^{4ie} + 5Ae^{2ie} + A - iBe^{10ie} - 3iBe^{8ie} - 2iBe^{6ie} + 2iBe^{4ie} + 3iBe^{2ie} + iB)e^{-4ie}}{32a^2c^3} \right) \\ + \frac{x(5A + iB)}{16a^2c^3} \end{array} \right.$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**2/(c-I*c*tan(f*x+e))**3,x)`output `Piecewise((((50331648*I*A*a**8*c**12*f**4*exp(2*I*e) - 50331648*B*a**8*c**12*f**4*exp(2*I*e))*exp(-4*I*f*x) + (503316480*I*A*a**8*c**12*f**4*exp(4*I*e) - 301989888*B*a**8*c**12*f**4*exp(4*I*e))*exp(-2*I*f*x) + (-1006632960*I*A*a**8*c**12*f**4*exp(8*I*e) - 201326592*B*a**8*c**12*f**4*exp(8*I*e))*exp(2*I*f*x) + (-251658240*I*A*a**8*c**12*f**4*exp(10*I*e) - 150994944*B*a**8*c**12*f**4*exp(10*I*e))*exp(4*I*f*x) + (-33554432*I*A*a**8*c**12*f**4*exp(12*I*e) - 33554432*B*a**8*c**12*f**4*exp(12*I*e))*exp(6*I*f*x))*exp(-6*I*e)/(6442450944*a**10*c**15*f**5), Ne(a**10*c**15*f**5*exp(6*I*e), 0)), (x*(-(5*A + I*B)/(16*a**2*c**3) + (A*exp(10*I*e) + 5*A*exp(8*I*e) + 10*A*exp(6*I*e) + 10*A*exp(4*I*e) + 5*A*exp(2*I*e) + A - I*B*exp(10*I*e) - 3*I*B*exp(8*I*e) - 2*I*B*exp(6*I*e) + 2*I*B*exp(4*I*e) + 3*I*B*exp(2*I*e) + I*B)*exp(-4*I*e)/(32*a**2*c**3)), True)) + x*(5*A + I*B)/(16*a**2*c**3)`**3.724.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^3,x, algorithm="maxima")`output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

$$3.724. \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^3} dx$$

3.724.8 Giac [A] (verification not implemented)

Time = 0.79 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.12

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^3} dx =$$

$$\frac{6(-5iA+B)\log(\tan(fx+e)+i)}{a^2c^3} + \frac{6(5iA-B)\log(\tan(fx+e)-i)}{a^2c^3} + \frac{3(15iA\tan(fx+e)^2 - 3B\tan(fx+e)^2 + 38A\tan(fx+e) + 10iB\tan(fx+e) - 25iA + 9B)}{a^2c^3(i\tan(fx+e)+1)^2}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^3,x, algorithm="giac")
```

```
output -1/192*(6*(-5*I*A + B)*log(tan(f*x + e) + I)/(a^2*c^3) + 6*(5*I*A - B)*log(tan(f*x + e) - I)/(a^2*c^3) + 3*(15*I*A*tan(f*x + e)^2 - 3*B*tan(f*x + e)^2 + 38*A*tan(f*x + e) + 10*I*B*tan(f*x + e) - 25*I*A + 9*B)/(a^2*c^3*(I*tan(f*x + e) + 1)^2) + (55*I*A*tan(f*x + e)^3 - 11*B*tan(f*x + e)^3 - 201*A*tan(f*x + e)^2 - 33*I*B*tan(f*x + e)^2 - 255*I*A*tan(f*x + e) + 27*B*tan(f*x + e) + 117*A - 3*I*B)/(a^2*c^3*(tan(f*x + e) + I)^3)/f
```

3.724.9 Mupad [B] (verification not implemented)

Time = 9.09 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.14

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^3} dx$$

$$= \frac{\tan(e + fx) \left(-\frac{5B}{48a^2c^3} + \frac{A25i}{48a^2c^3} \right) + \tan(e + fx)^3 \left(-\frac{B}{16a^2c^3} + \frac{A5i}{16a^2c^3} \right) + \tan(e + fx)^4 \left(\frac{5A}{16a^2c^3} + \frac{B1i}{16a^2c^3} \right) + f \left(\tan(e + fx)^5 + \tan(e + fx)^4 1i + 2 \tan(e + fx)^3 + \tan(e + fx)^2 2i + \tan(e + fx) + 1 \right)}{16a^2c^3}$$

```
input int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)^3),x)
```

```
output (tan(e + f*x)*((A*25i)/(48*a^2*c^3) - (5*B)/(48*a^2*c^3)) + tan(e + f*x)^3*((A*5i)/(16*a^2*c^3) - B/(16*a^2*c^3)) + tan(e + f*x)^4*((5*A)/(16*a^2*c^3) + (B*1i)/(16*a^2*c^3)) + tan(e + f*x)^2*((25*A)/(48*a^2*c^3) + (B*5i)/(48*a^2*c^3)) + A/(6*a^2*c^3) - (B*1i)/(6*a^2*c^3))/(f*(tan(e + f*x) + tan(e + f*x)^2*2i + 2*tan(e + f*x)^3 + tan(e + f*x)^4*1i + tan(e + f*x)^5 + 1i)) - (x*(A*5i - B)*1i)/(16*a^2*c^3)
```

3.724. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^3} dx$

3.725
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^4} dx$$

3.725.1 Optimal result 6682
 3.725.2 Mathematica [A] (verified) 6683
 3.725.3 Rubi [A] (verified) 6683
 3.725.4 Maple [A] (verified) 6685
 3.725.5 Fricas [A] (verification not implemented) 6685
 3.725.6 Sympy [A] (verification not implemented) 6686
 3.725.7 Maxima [F(-2)] 6687
 3.725.8 Giac [A] (verification not implemented) 6687
 3.725.9 Mupad [B] (verification not implemented) 6688

3.725.1 Optimal result

Integrand size = 41, antiderivative size = 221

$$\begin{aligned} & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2(c - ic \tan(e + fx))^4} dx \\ &= \frac{5(3A + iB)x}{64a^2c^4} - \frac{iA - B}{64a^2c^4 f(i - \tan(e + fx))^2} - \frac{5A + 3iB}{64a^2c^4 f(i - \tan(e + fx))} \\ & \quad - \frac{iA + B}{32a^2c^4 f(i + \tan(e + fx))^4} - \frac{3A - iB}{48a^2c^4 f(i + \tan(e + fx))^3} \\ & \quad + \frac{3iA}{32a^2c^4 f(i + \tan(e + fx))^2} + \frac{5A + iB}{32a^2c^4 f(i + \tan(e + fx))} \end{aligned}$$

```
output 5/64*(3*A+I*B)*x/a^2/c^4+1/64*(-I*A+B)/a^2/c^4/f/(I-tan(f*x+e))^2+1/64*(-5
*A-3*I*B)/a^2/c^4/f/(I-tan(f*x+e))+1/32*(-I*A-B)/a^2/c^4/f/(I+tan(f*x+e))^
4+1/48*(-3*A+I*B)/a^2/c^4/f/(I+tan(f*x+e))^3+3/32*I*A/a^2/c^4/f/(I+tan(f*x
+e))^2+1/32*(5*A+I*B)/a^2/c^4/f/(I+tan(f*x+e))
```

3.725.2 Mathematica [A] (verified)

Time = 6.08 (sec) , antiderivative size = 213, normalized size of antiderivative = 0.96

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ict \tan(e + fx))^4} dx$$

$$= \frac{\sec^5(e + fx)(240iA \cos(e + fx) + 5(-9iA + 11B) \cos(3(e + fx)) - 3iA \cos(5(e + fx)) + 9B \cos(5(e + fx)))}{768a^2c^4f(-I + \tan(e + fx))^2(I + \tan(e + fx))^4}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^4),x]`

output `(Sec[e + f*x]^5*((240*I)*A*Cos[e + f*x] + 5*((-9*I)*A + 11*B)*Cos[3*(e + f*x)] - (3*I)*A*Cos[5*(e + f*x)] + 9*B*Cos[5*(e + f*x)] + 102*A*Sin[e + f*x] + (34*I)*B*Sin[e + f*x] - 60*(3*A + I*B)*ArcTan[Tan[e + f*x]]*Sec[e + f*x]*(Cos[2*(e + f*x)] - I*Sin[2*(e + f*x)]) - 87*A*Sin[3*(e + f*x)] - (29*I)*B*Sin[3*(e + f*x)] - 9*A*Sin[5*(e + f*x)] - (3*I)*B*Sin[5*(e + f*x)]))/((768*a^2*c^4*f*(-I + Tan[e + f*x])^2*(I + Tan[e + f*x])^4)`

3.725.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.80, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ict \tan(e + fx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ict \tan(e + fx))^4} dx$$

$$\downarrow \text{4071}$$

$$\frac{ac \int \frac{A + B \tan(e + fx)}{a^3 c^5 (1 - i \tan(e + fx))^5 (i \tan(e + fx) + 1)^3} d \tan(e + fx)}{f}$$

$$\downarrow \text{27}$$

3.725. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ict \tan(e + fx))^4} dx$

$$\int \frac{A+B \tan(e+fx)}{(1-i \tan(e+fx))^5 (i \tan(e+fx)+1)^3} d \tan(e+fx)$$

↓ 86

$$\int \left(-\frac{3iA}{16(\tan(e+fx)+i)^3} + \frac{5(3A+iB)}{64(\tan^2(e+fx)+1)} + \frac{-5A-3iB}{64(\tan(e+fx)-i)^2} + \frac{-5A-iB}{32(\tan(e+fx)+i)^2} + \frac{i(A+iB)}{32(\tan(e+fx)-i)^3} + \frac{3A-iB}{16(\tan(e+fx)+i)^4} + \dots \right) d \tan(e+fx)$$

↓ 2009

$$\frac{5}{64}(3A+iB) \arctan(\tan(e+fx)) - \frac{5A+3iB}{64(-\tan(e+fx)+i)} + \frac{5A+iB}{32(\tan(e+fx)+i)} - \frac{-B+iA}{64(-\tan(e+fx)+i)^2} - \frac{3A-iB}{48(\tan(e+fx)+i)^3} - \frac{3A-iB}{32(\tan(e+fx)+i)^4} + \dots$$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^4), x]`

output `((5*(3*A + I*B)*ArcTan[Tan[e + f*x]])/64 - (I*A - B)/(64*(I - Tan[e + f*x])^2) - (5*A + (3*I)*B)/(64*(I - Tan[e + f*x])) - (I*A + B)/(32*(I + Tan[e + f*x])^4) - (3*A - I*B)/(48*(I + Tan[e + f*x])^3) + (((3*I)/32)*A)/(I + Tan[e + f*x])^2 + (5*A + I*B)/(32*(I + Tan[e + f*x]))) / (a^2*c^4*f)`

3.725.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 86 `Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.725. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^4} dx$

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x],
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.725.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.27

method	result
norman	$\frac{5(iB+3A)x}{64ac} - \frac{3iA+B}{12acf} + \frac{B \tan(fx+e)^2}{6acf} + \frac{73(iB+3A) \tan(fx+e)^3}{192acf} + \frac{55(iB+3A) \tan(fx+e)^5}{192acf} + \frac{5(iB+3A) \tan(fx+e)^7}{64acf} + \frac{5(iB+3A)x \tan(fx+e)^8}{16a^2c^2f} + \frac{a^3 \left(1 + \tan(fx+e)\right)^8}{16a^2c^2f}$
risch	$\frac{5ixB}{64a^2c^4} + \frac{15xA}{64a^2c^4} - \frac{e^{8i(fx+e)}B}{512a^2c^4f} - \frac{ie^{8i(fx+e)}A}{512a^2c^4f} - \frac{e^{6i(fx+e)}B}{96a^2c^4f} - \frac{ie^{6i(fx+e)}A}{64a^2c^4f} - \frac{3 \cos(4fx+4e)B}{128a^2c^4f} - \frac{7i \cos(4fx+4e)A}{128a^2c^4f}$
derivativedivides	$\frac{iB}{32f a^2c^4(i+\tan(fx+e))} + \frac{3iA}{32a^2c^4f(i+\tan(fx+e))^2} + \frac{15A \arctan(\tan(fx+e))}{64f a^2c^4} + \frac{5A}{64f a^2c^4(-i+\tan(fx+e))} - \frac{3A}{32f a^2c^4}$
default	$\frac{iB}{32f a^2c^4(i+\tan(fx+e))} + \frac{3iA}{32a^2c^4f(i+\tan(fx+e))^2} + \frac{15A \arctan(\tan(fx+e))}{64f a^2c^4} + \frac{5A}{64f a^2c^4(-i+\tan(fx+e))} - \frac{3A}{32f a^2c^4}$

```
input int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^4,x,method=_R
ETURNVERBOSE)
```

```
output (5/64*(3*A+I*B)/a/c*x-1/12*(3*I*A+B)/a/c/f+1/6/a/c/f*B*tan(f*x+e)^2+73/192
*(3*A+I*B)/a/c/f*tan(f*x+e)^3+55/192*(3*A+I*B)/a/c/f*tan(f*x+e)^5+5/64*(3*
A+I*B)/a/c/f*tan(f*x+e)^7+5/16*(3*A+I*B)/a/c*x*tan(f*x+e)^2+15/32*(3*A+I*B
)/a/c*x*tan(f*x+e)^4+5/16*(3*A+I*B)/a/c*x*tan(f*x+e)^6+5/64*(3*A+I*B)/a/c*
x*tan(f*x+e)^8+1/64*(49*A-5*I*B)/a/c/f*tan(f*x+e)/a/c^3/(1+tan(f*x+e)^2)^
4
```

3.725.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.57

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2(c - ictan(e + fx))^4} dx$$

$$= \frac{(120(3A + iB)fx e^{(4i fx + 4ie)} - 3(iA + B)e^{(12i fx + 12ie)} - 8(3iA + 2B)e^{(10i fx + 10ie)} - 30(3iA + B)e^{(8i fx + 8ie)})}{1536 a^2 c^4 f}$$

3.725. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^4} dx$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^4,x, algorithm="fricas")`

output `1/1536*(120*(3*A + I*B)*f*x*e^(4*I*f*x + 4*I*e) - 3*(I*A + B)*e^(12*I*f*x + 12*I*e) - 8*(3*I*A + 2*B)*e^(10*I*f*x + 10*I*e) - 30*(3*I*A + B)*e^(8*I*f*x + 8*I*e) - 240*I*A*e^(6*I*f*x + 6*I*e) - 24*(-3*I*A + 2*B)*e^(2*I*f*x + 2*I*e) + 6*I*A - 6*B)*e^(-4*I*f*x - 4*I*e)/(a^2*c^4*f)`

3.725.6 Sympy [A] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 498, normalized size of antiderivative = 2.25

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^4} dx$$

$$= \begin{cases} (-2061584302080iAa^{10}c^{20}f^5e^{8ie}e^{2ifx} + (51539607552iAa^{10}c^{20}f^5e^{2ie} - 51539607552Ba^{10}c^{20}f^5e^{2ie})e^{-4ifx} + (618475290624iAa^{10}c^{20}f^5e^{4ie} - 412316860416B*a^{10}c^{20}f^5*exp(4*I*e))*exp(-2*I*f*x) + (-773094113280*I*A*a^{10}c^{20}f^5*exp(10*I*e) - 257698037760*B*a^{10}c^{20}f^5*exp(10*I*e))*exp(4*I*f*x) + (-206158430208*I*A*a^{10}c^{20}f^5*exp(12*I*e) - 137438953472*B*a^{10}c^{20}f^5*exp(12*I*e))*exp(6*I*f*x) + (-25769803776*I*A*a^{10}c^{20}f^5*exp(14*I*e) - 25769803776*B*a^{10}c^{20}f^5*exp(14*I*e))*exp(8*I*f*x))*exp(-6*I*e)/(13194139533312*a^{12}c^{24}f^6), \\ x\left(-\frac{15A+5iB}{64a^2c^4} + \frac{(Ae^{12ie}+6Ae^{10ie}+15Ae^{8ie}+20Ae^{6ie}+15Ae^{4ie}+6Ae^{2ie}+A-iBe^{12ie}-4iBe^{10ie}-5iBe^{8ie}+5iBe^{4ie}+4iBe^{2ie}+iB)e^{-4ie}}{64a^2c^4}\right) \\ + \frac{x(15A + 5iB)}{64a^2c^4} \end{cases}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**2/(c-I*c*tan(f*x+e))**4,x)`

output `Piecewise(((-2061584302080*I*A*a**10*c**20*f**5*exp(8*I*e)*exp(2*I*f*x) + (51539607552*I*A*a**10*c**20*f**5*exp(2*I*e) - 51539607552*B*a**10*c**20*f**5*exp(2*I*e))*exp(-4*I*f*x) + (618475290624*I*A*a**10*c**20*f**5*exp(4*I*e) - 412316860416*B*a**10*c**20*f**5*exp(4*I*e))*exp(-2*I*f*x) + (-773094113280*I*A*a**10*c**20*f**5*exp(10*I*e) - 257698037760*B*a**10*c**20*f**5*exp(10*I*e))*exp(4*I*f*x) + (-206158430208*I*A*a**10*c**20*f**5*exp(12*I*e) - 137438953472*B*a**10*c**20*f**5*exp(12*I*e))*exp(6*I*f*x) + (-25769803776*I*A*a**10*c**20*f**5*exp(14*I*e) - 25769803776*B*a**10*c**20*f**5*exp(14*I*e))*exp(8*I*f*x))*exp(-6*I*e)/(13194139533312*a**12*c**24*f**6), Ne(a**12*c**24*f**6*exp(6*I*e), 0)), (x*(-(15*A + 5*I*B)/(64*a**2*c**4) + (A*exp(12*I*e) + 6*A*exp(10*I*e) + 15*A*exp(8*I*e) + 20*A*exp(6*I*e) + 15*A*exp(4*I*e) + 6*A*exp(2*I*e) + A - I*B*exp(12*I*e) - 4*I*B*exp(10*I*e) - 5*I*B*exp(8*I*e) + 5*I*B*exp(4*I*e) + 4*I*B*exp(2*I*e) + I*B)*exp(-4*I*e)/(64*a**2*c**4)), True)) + x*(15*A + 5*I*B)/(64*a**2*c**4)`

3.725.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^4} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^4,x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.725.8 Giac [A] (verification not implemented)

Time = 0.94 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.02

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^4} dx$$

$$= \frac{60(3iA - B) \log(\tan(fx + e) + i)}{a^2 c^4} + \frac{60(-3iA + B) \log(\tan(fx + e) - i)}{a^2 c^4} - \frac{6(-45iA \tan(fx + e)^2 + 15B \tan(fx + e)^2 - 110A \tan(fx + e) - 42iB \tan(fx + e) + 69iA - 31B)}{a^2 c^4 (\tan(fx + e) - i)^2}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^4,x, algorithm="giac")
```

```
output 1/1536*(60*(3*I*A - B)*log(tan(f*x + e) + I)/(a^2*c^4) + 60*(-3*I*A + B)*log(tan(f*x + e) - I)/(a^2*c^4) - 6*(-45*I*A*tan(f*x + e)^2 + 15*B*tan(f*x + e)^2 - 110*A*tan(f*x + e) - 42*I*B*tan(f*x + e) + 69*I*A - 31*B)/(a^2*c^4*(tan(f*x + e) - I)^2) + (-375*I*A*tan(f*x + e)^4 + 125*B*tan(f*x + e)^4 + 1740*A*tan(f*x + e)^3 + 548*I*B*tan(f*x + e)^3 + 3114*I*A*tan(f*x + e)^2 - 894*B*tan(f*x + e)^2 - 2604*A*tan(f*x + e) - 612*I*B*tan(f*x + e) - 903*I*A + 93*B)/(a^2*c^4*(tan(f*x + e) + I)^4))/f
```

3.725.9 Mupad [B] (verification not implemented)

Time = 10.03 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.12

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^4} dx$$

$$= \frac{\frac{B}{12a^2c^4} + \tan(e + fx)^4 \left(-\frac{5B}{32a^2c^4} + \frac{A15i}{32a^2c^4}\right) + \tan(e + fx)^3 \left(\frac{5A}{32a^2c^4} + \frac{B5i}{96a^2c^4}\right) + \tan(e + fx)^5 \left(\frac{15A}{64a^2c^4} + \frac{B}{64a^2c^4}\right)}{f (\tan(e + fx)^6 + \tan(e + fx)^5 2i + \tan(e + fx)^4 + \tan(e + fx)^3 2i + \tan(e + fx)^2 + \tan(e + fx) + 1)} + \frac{5x(3A + B1i)}{64a^2c^4}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)^4),x)`

output `(tan(e + f*x)^4*((A*15i)/(32*a^2*c^4) - (5*B)/(32*a^2*c^4)) - tan(e + f*x)*((17*A)/(64*a^2*c^4) + (B*17i)/(192*a^2*c^4)) + tan(e + f*x)^3*((5*A)/(32*a^2*c^4) + (B*5i)/(96*a^2*c^4)) + tan(e + f*x)^5*((15*A)/(64*a^2*c^4) + (B*5i)/(64*a^2*c^4)) + tan(e + f*x)^2*((A*25i)/(32*a^2*c^4) - (25*B)/(96*a^2*c^4)) + (A*1i)/(4*a^2*c^4) + B/(12*a^2*c^4))/(f*(tan(e + f*x)*2i - tan(e + f*x)^2 + tan(e + f*x)^3*4i + tan(e + f*x)^4 + tan(e + f*x)^5*2i + tan(e + f*x)^6 - 1)) + (5*x*(3*A + B*1i))/(64*a^2*c^4)`

3.726 $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^5} dx$

3.726.1 Optimal result 6689
 3.726.2 Mathematica [A] (verified) 6690
 3.726.3 Rubi [A] (verified) 6690
 3.726.4 Maple [A] (verified) 6692
 3.726.5 Fricas [A] (verification not implemented) 6693
 3.726.6 Sympy [A] (verification not implemented) 6693
 3.726.7 Maxima [F(-2)] 6694
 3.726.8 Giac [A] (verification not implemented) 6695
 3.726.9 Mupad [B] (verification not implemented) 6695

3.726.1 Optimal result

Integrand size = 41, antiderivative size = 251

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2(c - ic \tan(e + fx))^5} dx$$

$$= \frac{3(7A + 3iB)x}{128a^2c^5} - \frac{iA - B}{128a^2c^5 f(i - \tan(e + fx))^2} - \frac{3A + 2iB}{64a^2c^5 f(i - \tan(e + fx))}$$

$$+ \frac{A - iB}{40a^2c^5 f(i + \tan(e + fx))^5} - \frac{3iA + B}{64a^2c^5 f(i + \tan(e + fx))^4} - \frac{A}{16a^2c^5 f(i + \tan(e + fx))^3}$$

$$+ \frac{5iA - B}{64a^2c^5 f(i + \tan(e + fx))^2} + \frac{5(3A + iB)}{128a^2c^5 f(i + \tan(e + fx))}$$

```
output 3/128*(7*A+3*I*B)*x/a^2/c^5+1/128*(-I*A+B)/a^2/c^5/f/(I-tan(f*x+e))^2+1/64
*(-3*A-2*I*B)/a^2/c^5/f/(I-tan(f*x+e))+1/40*(A-I*B)/a^2/c^5/f/(I+tan(f*x+e
))^5+1/64*(-3*I*A-B)/a^2/c^5/f/(I+tan(f*x+e))^4-1/16*A/a^2/c^5/f/(I+tan(f*
x+e))^3+1/64*(5*I*A-B)/a^2/c^5/f/(I+tan(f*x+e))^2+5/128*(3*A+I*B)/a^2/c^5/
f/(I+tan(f*x+e))
```


3.726.2 Mathematica [A] (verified)

Time = 6.53 (sec) , antiderivative size = 233, normalized size of antiderivative = 0.93

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ictan(e + fx))^5} dx$$

$$= \frac{\sec^6(e + fx)(-241A + 11iB - 8(71A + 9iB) \cos(2(e + fx)) + 3(33A + 37iB) \cos(4(e + fx)) + 6A \cos(6(e + fx)))}{(2560a^2c^5f(-I + \tan(e + fx))^2(I + \tan(e + fx))^5)}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^5),x]`

output `(Sec[e + f*x]^6*(-241*A + (11*I)*B - 8*(71*A + (9*I)*B)*Cos[2*(e + f*x)] + 3*(33*A + (37*I)*B)*Cos[4*(e + f*x)] + 6*A*Cos[6*(e + f*x)] + (14*I)*B*Cos[6*(e + f*x)] + (350*I)*A*Sin[2*(e + f*x)] - 150*B*Sin[2*(e + f*x)] + 60*((-7*I)*A + 3*B)*ArcTan[Tan[e + f*x]]*Sec[e + f*x]*(Cos[3*(e + f*x)] - I*Sin[3*(e + f*x)]) - (161*I)*A*Sin[4*(e + f*x)] + 69*B*Sin[4*(e + f*x)] - (14*I)*A*Sin[6*(e + f*x)] + 6*B*Sin[6*(e + f*x)])/(2560*a^2*c^5*f*(-I + Tan[e + f*x])^2*(I + Tan[e + f*x])^5)`

3.726.3 Rubi [A] (verified)Time = 0.46 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.79, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ictan(e + fx))^5} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ictan(e + fx))^5} dx$$

$$\downarrow \text{4071}$$

$$ac \int \frac{A+B \tan(e+fx)}{a^3 c^6 (1-i \tan(e+fx))^6 (i \tan(e+fx)+1)^3} d \tan(e + fx)$$

$$\downarrow \text{27}$$

3.726. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 (c-ic \tan(e+fx))^5} dx$

$$\int \frac{A+B \tan(e+fx)}{(1-i \tan(e+fx))^6(i \tan(e+fx)+1)^3} d \tan(e+fx)$$

$a^2 c^5 f$

↓ 86

$$\int \left(\frac{3A}{16(\tan(e+fx)+i)^4} + \frac{3(7A+3iB)}{128(\tan^2(e+fx)+1)} + \frac{-3A-2iB}{64(\tan(e+fx)-i)^2} - \frac{5(3A+iB)}{128(\tan(e+fx)+i)^2} + \frac{i(A+iB)}{64(\tan(e+fx)-i)^3} + \frac{B-5iA}{32(\tan(e+fx)+i)^3} + \dots \right) \frac{1}{a^2 c^5 f}$$

↓ 2009

$$\frac{3}{128}(7A+3iB) \arctan(\tan(e+fx)) - \frac{3A+2iB}{64(-\tan(e+fx)+i)} + \frac{5(3A+iB)}{128(\tan(e+fx)+i)} - \frac{-B+iA}{128(-\tan(e+fx)+i)^2} + \frac{-B+5iA}{64(\tan(e+fx)+i)^2} - \dots \frac{1}{a^2 c^5 f}$$

```
input Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^5),x]
```

```
output ((3*(7*A + (3*I)*B)*ArcTan[Tan[e + f*x]])/128 - (I*A - B)/(128*(I - Tan[e + f*x])^2) - (3*A + (2*I)*B)/(64*(I - Tan[e + f*x])) + (A - I*B)/(40*(I + Tan[e + f*x])^5) - ((3*I)*A + B)/(64*(I + Tan[e + f*x])^4) - A/(16*(I + Tan[e + f*x])^3) + ((5*I)*A - B)/(64*(I + Tan[e + f*x])^2) + (5*(3*A + I*B))/(128*(I + Tan[e + f*x])))/(a^2*c^5*f)
```

3.726.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

3.726. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^5} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.726.4 Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 303, normalized size of antiderivative = 1.21

method	result
risch	$\frac{9ixB}{128a^2c^5} + \frac{21xA}{128a^2c^5} - \frac{e^{10i(fx+e)}B}{1280a^2c^5f} - \frac{ie^{10i(fx+e)}A}{1280a^2c^5f} - \frac{5e^{8i(fx+e)}B}{1024a^2c^5f} - \frac{7ie^{8i(fx+e)}A}{1024a^2c^5f} - \frac{3e^{6i(fx+e)}B}{256a^2c^5f} - \frac{7ie^{6i(fx+e)}A}{256a^2c^5f}$
norman	$\frac{3(3iB+7A)x}{128ac} - \frac{11iA+B}{40acf} + \frac{(-9iB+107A)\tan(fx+e)}{128acf} + \frac{(3iB+7A)\tan(fx+e)^5}{5acf} + \frac{7(3iB+7A)\tan(fx+e)^7}{64acf} + \frac{3(3iB+7A)\tan(fx+e)^9}{128acf}$
derivativedivides	$\frac{9iB \arctan(\tan(fx+e))}{128f a^2 c^5} + \frac{21A \arctan(\tan(fx+e))}{128f a^2 c^5} - \frac{iB}{40f a^2 c^5 (i+\tan(fx+e))^5} + \frac{iB}{32f a^2 c^5 (-i+\tan(fx+e))} + \dots$
default	$\frac{9iB \arctan(\tan(fx+e))}{128f a^2 c^5} + \frac{21A \arctan(\tan(fx+e))}{128f a^2 c^5} - \frac{iB}{40f a^2 c^5 (i+\tan(fx+e))^5} + \frac{iB}{32f a^2 c^5 (-i+\tan(fx+e))} + \dots$

input `int((A+B*tan(f*x+e))/(a+i*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^5,x,method=_RETURVERBOSE)`

output `9/128*I*x/a^2/c^5*B+21/128*x/a^2/c^5*A-1/1280/a^2/c^5/f*exp(10*I*(f*x+e))*B-1/1280*I/a^2/c^5/f*exp(10*I*(f*x+e))*A-5/1024/a^2/c^5/f*exp(8*I*(f*x+e))*B-7/1024*I/a^2/c^5/f*exp(8*I*(f*x+e))*A-3/256/a^2/c^5/f*exp(6*I*(f*x+e))*B-7/256*I/a^2/c^5/f*exp(6*I*(f*x+e))*A-3/256/a^2/c^5/f*cos(4*f*x+4*e)*B-17/256*I/a^2/c^5/f*cos(4*f*x+4*e)*A-1/128*I/a^2/c^5/f*sin(4*f*x+4*e)*B+9/128/a^2/c^5/f*sin(4*f*x+4*e)*A-7/64*I*A/a^2/c^5/f*cos(2*f*x+2*e)+5/128*I/a^2/c^5/f*sin(2*f*x+2*e)*B+21/128/a^2/c^5/f*sin(2*f*x+2*e)*A`

3.726. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^5} dx$

3.726.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.60

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^5} dx$$

$$= \frac{(120(7A + 3iB)fx e^{(4i fx + 4ie)} - 4(iA + B)e^{(14i fx + 14ie)} - 5(7iA + 5B)e^{(12i fx + 12ie)} - 20(7iA + 3B)e^{(10i fx + 10ie)} - 50(7iA + B)e^{(8i fx + 8ie)} - 100(7iA - B)e^{(6i fx + 6ie)} - 20(-7iA + 5B)e^{(2i fx + 2ie)} + 10iA - 10B)e^{(-4i fx - 4ie)}}{(a^2 c^5 f)}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^5,x, algorithm="fricas")
```

```
output 1/5120*(120*(7*A + 3*I*B)*f*x*e^(4*I*f*x + 4*I*e) - 4*(I*A + B)*e^(14*I*f*x + 14*I*e) - 5*(7*I*A + 5*B)*e^(12*I*f*x + 12*I*e) - 20*(7*I*A + 3*B)*e^(10*I*f*x + 10*I*e) - 50*(7*I*A + B)*e^(8*I*f*x + 8*I*e) - 100*(7*I*A - B)*e^(6*I*f*x + 6*I*e) - 20*(-7*I*A + 5*B)*e^(2*I*f*x + 2*I*e) + 10*I*A - 10*B)*e^(-4*I*f*x - 4*I*e)/(a^2*c^5*f)
```

3.726.6 Sympy [A] (verification not implemented)

Time = 0.58 (sec) , antiderivative size = 605, normalized size of antiderivative = 2.41

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^5} dx$$

$$= \left\{ \frac{((11258999068426240iAa^{12}c^{30}f^6e^{2ie} - 11258999068426240Ba^{12}c^{30}f^6e^{2ie})e^{-4ifx} + (157625986957967360iAa^{12}c^{30}f^6e^{4ie} - 11258999068426240Ba^{12}c^{30}f^6e^{4ie})e^{-12ifx} + (157625986957967360iAa^{12}c^{30}f^6e^{6ie} - 11258999068426240Ba^{12}c^{30}f^6e^{6ie})e^{-20ifx} + (157625986957967360iAa^{12}c^{30}f^6e^{8ie} - 11258999068426240Ba^{12}c^{30}f^6e^{8ie})e^{-28ifx} + (157625986957967360iAa^{12}c^{30}f^6e^{10ie} - 11258999068426240Ba^{12}c^{30}f^6e^{10ie})e^{-36ifx} + (157625986957967360iAa^{12}c^{30}f^6e^{12ie} - 11258999068426240Ba^{12}c^{30}f^6e^{12ie})e^{-44ifx} + (157625986957967360iAa^{12}c^{30}f^6e^{14ie} - 11258999068426240Ba^{12}c^{30}f^6e^{14ie})e^{-52ifx} + (157625986957967360iAa^{12}c^{30}f^6e^{16ie} - 11258999068426240Ba^{12}c^{30}f^6e^{16ie})e^{-60ifx} + (157625986957967360iAa^{12}c^{30}f^6e^{18ie} - 11258999068426240Ba^{12}c^{30}f^6e^{18ie})e^{-68ifx} + (157625986957967360iAa^{12}c^{30}f^6e^{20ie} - 11258999068426240Ba^{12}c^{30}f^6e^{20ie})e^{-76ifx}}{128a^2c^5} \right\} + \frac{x(21A + 9iB)}{128a^2c^5}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**2/(c-I*c*tan(f*x+e))**5,x)
```

```
output Piecewise((((11258999068426240*I*A**12*c**30*f**6*exp(2*I*e) - 112589990
68426240*B*a**12*c**30*f**6*exp(2*I*e))*exp(-4*I*f*x) + (15762598695796736
0*I*A**12*c**30*f**6*exp(4*I*e) - 112589990684262400*B*a**12*c**30*f**6*
exp(4*I*e))*exp(-2*I*f*x) + (-788129934789836800*I*A**12*c**30*f**6*exp(
8*I*e) + 112589990684262400*B*a**12*c**30*f**6*exp(8*I*e))*exp(2*I*f*x) +
(-394064967394918400*I*A**12*c**30*f**6*exp(10*I*e) - 56294995342131200*
B*a**12*c**30*f**6*exp(10*I*e))*exp(4*I*f*x) + (-157625986957967360*I*A**
12*c**30*f**6*exp(12*I*e) - 67553994410557440*B*a**12*c**30*f**6*exp(12*I
*e))*exp(6*I*f*x) + (-39406496739491840*I*A**12*c**30*f**6*exp(14*I*e) -
28147497671065600*B*a**12*c**30*f**6*exp(14*I*e))*exp(8*I*f*x) + (-450359
9627370496*I*A**12*c**30*f**6*exp(16*I*e) - 4503599627370496*B*a**12*c**
30*f**6*exp(16*I*e))*exp(10*I*f*x))*exp(-6*I*e)/(5764607523034234880*a**14
*c**35*f**7), Ne(a**14*c**35*f**7*exp(6*I*e), 0)), (x*(-(21*A + 9*I*B)/(12
8*a**2*c**5) + (A*exp(14*I*e) + 7*A*exp(12*I*e) + 21*A*exp(10*I*e) + 35*A*
exp(8*I*e) + 35*A*exp(6*I*e) + 21*A*exp(4*I*e) + 7*A*exp(2*I*e) + A - I*B*
exp(14*I*e) - 5*I*B*exp(12*I*e) - 9*I*B*exp(10*I*e) - 5*I*B*exp(8*I*e) + 5
*I*B*exp(6*I*e) + 9*I*B*exp(4*I*e) + 5*I*B*exp(2*I*e) + I*B)*exp(-4*I*e)/(
128*a**2*c**5)), True)) + x*(21*A + 9*I*B)/(128*a**2*c**5)
```

3.726.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^5} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^5,x, al
gorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.726.8 Giac [A] (verification not implemented)

Time = 0.79 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.00

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ictan(e + fx))^5} dx =$$

$$\frac{60(-7iA+3B)\log(\tan(fx+e)+i)}{a^2c^5} + \frac{60(7iA-3B)\log(\tan(fx+e)-i)}{a^2c^5} + \frac{10(63iA\tan(fx+e)^2-27B\tan(fx+e)^2+150A\tan(fx+e)+70B)}{a^2c^5(-i\tan(fx+e)-1)^2}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^5,x, algorithm="giac")`

output `-1/5120*(60*(-7*I*A + 3*B)*log(tan(f*x + e) + I)/(a^2*c^5) + 60*(7*I*A - 3*B)*log(tan(f*x + e) - I)/(a^2*c^5) + 10*(63*I*A*tan(f*x + e)^2 - 27*B*tan(f*x + e)^2 + 150*A*tan(f*x + e) + 70*I*B*tan(f*x + e) - 91*I*A + 47*B)/(a^2*c^5*(-I*tan(f*x + e) - 1)^2) + (959*I*A*tan(f*x + e)^5 - 411*B*tan(f*x + e)^5 - 5395*A*tan(f*x + e)^4 - 2255*I*B*tan(f*x + e)^4 - 12390*I*A*tan(f*x + e)^3 + 4990*B*tan(f*x + e)^3 + 14710*A*tan(f*x + e)^2 + 5550*I*B*tan(f*x + e)^2 + 9275*I*A*tan(f*x + e) - 3015*B*tan(f*x + e) - 2647*A - 483*I*B)/(a^2*c^5*(tan(f*x + e) + I)^5))/f`

3.726.9 Mupad [B] (verification not implemented)

Time = 10.57 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.16

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ictan(e + fx))^5} dx$$

$$= \frac{\tan(e + fx) \left(-\frac{3B}{640a^2c^5} + \frac{A7i}{640a^2c^5} \right) + \tan(e + fx)^4 \left(\frac{7A}{32a^2c^5} + \frac{B3i}{32a^2c^5} \right) - \tan(e + fx)^3 \left(-\frac{9B}{32a^2c^5} + \frac{A21i}{32a^2c^5} \right)}{f \left(-\tan(e + fx)^7 - \tan(e + fx)^6 3i + \tan(e + fx)^5 \right)} + \frac{3x(7A + B3i)}{128a^2c^5}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*i)^2*(c - c*tan(e + f*x)*i)^5),x)`

output $(\tan(e + f*x)*((A*7i)/(640*a^2*c^5) - (3*B)/(640*a^2*c^5)) + \tan(e + f*x)^4*((7*A)/(32*a^2*c^5) + (B*3i)/(32*a^2*c^5)) - \tan(e + f*x)^3*((A*21i)/(32*a^2*c^5) - (9*B)/(32*a^2*c^5)) - \tan(e + f*x)^6*((21*A)/(128*a^2*c^5) + (B*9i)/(128*a^2*c^5)) - \tan(e + f*x)^5*((A*63i)/(128*a^2*c^5) - (27*B)/(128*a^2*c^5)) + \tan(e + f*x)^2*((469*A)/(640*a^2*c^5) + (B*201i)/(640*a^2*c^5)) + (11*A)/(40*a^2*c^5) - (B*1i)/(40*a^2*c^5))/(f*(3*\tan(e + f*x) - \tan(e + f*x)^2*1i + 5*\tan(e + f*x)^3 - \tan(e + f*x)^4*5i + \tan(e + f*x)^5 - \tan(e + f*x)^6*3i - \tan(e + f*x)^7 + 1i)) + (3*x*(7*A + B*3i))/(128*a^2*c^5)$

3.726. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^5} dx$

$$3.727 \quad \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^n}{(a+ia \tan(e+fx))^3} dx$$

3.727.1 Optimal result	6697
3.727.2 Mathematica [A] (verified)	6697
3.727.3 Rubi [A] (verified)	6698
3.727.4 Maple [F]	6700
3.727.5 Fricas [F]	6700
3.727.6 Sympy [F]	6700
3.727.7 Maxima [F(-2)]	6701
3.727.8 Giac [F]	6701
3.727.9 Mupad [F(-1)]	6701

3.727.1 Optimal result

Integrand size = 41, antiderivative size = 115

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{(iA(3 - n) + B(3 + n)) \operatorname{Hypergeometric2F1}\left(3, n, 1 + n, \frac{1}{2}(1 - i \tan(e + fx))\right) (c - ic \tan(e + fx))^n}{48a^3fn} + \frac{(iA - B)(c - ic \tan(e + fx))^n}{6a^3f(1 + i \tan(e + fx))^3}$$

output `1/48*(I*A*(3-n)+B*(3+n))*hypergeom([3, n],[1+n],1/2-1/2*I*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/a^3/f/n+1/6*(I*A-B)*(c-I*c*tan(f*x+e))^n/a^3/f/(1+I*tan(f*x+e))^3`

3.727.2 Mathematica [A] (verified)

Time = 2.58 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.83

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{(c - ic \tan(e + fx))^n \left(-B(-7 + n^2) + iA(11 - 6n + n^2) + \frac{(2-3n+n^2)(-iA(-3+n)+B(3+n)) \operatorname{Hypergeometric2F1}(1, n, 2, \frac{1}{2}(1 - i \tan(e + fx)))}{n} \right)}{6a^3f(1 + i \tan(e + fx))^3}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^n)/(a + I*a*Tan[e + f*x])^3,x]`

output `((c - I*c*Tan[e + f*x])^n*(-(B*(-7 + n^2)) + I*A*(11 - 6*n + n^2) + ((2 - 3*n + n^2)*((-I)*A*(-3 + n) + B*(3 + n))*Hypergeometric2F1[1, n, 1 + n, (-1/2*I)*(I + Tan[e + f*x])])/n + ((8*I)*(A + I*B)*(I + Tan[e + f*x]))/(-I + Tan[e + f*x])^3 + (2*(A*(-5 + n) + I*B*(1 + n))*(I + Tan[e + f*x]))/(-I + Tan[e + f*x])^2 + ((B*(-7 + n^2) - I*A*(11 - 6*n + n^2))*(I + Tan[e + f*x]))/(-I + Tan[e + f*x]))/(96*a^3*f)`

3.727.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 87, 78}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{n-1}}{a^4(i \tan(e + fx) + 1)^4} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{c \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{n-1}}{(i \tan(e + fx) + 1)^4} d \tan(e + fx)}{a^3 f} \\
 & \quad \downarrow \text{87} \\
 & \frac{c \left(\frac{1}{6} (A(3 - n) - iB(n + 3)) \int \frac{(c - ic \tan(e + fx))^{n-1}}{(i \tan(e + fx) + 1)^3} d \tan(e + fx) + \frac{(-B + iA)(c - ic \tan(e + fx))^n}{6c(1 + i \tan(e + fx))^3} \right)}{a^3 f} \\
 & \quad \downarrow \text{78}
 \end{aligned}$$

3.727. $\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^3} dx$

$$\frac{c \left(\frac{i(A(3-n) - iB(n+3))(c - i \tan(e+fx))^n \operatorname{Hypergeometric2F1}\left(3, n, n+1, \frac{1}{2}(1 - i \tan(e+fx))\right)}{48cn} + \frac{(-B+iA)(c - i \tan(e+fx))^n}{6c(1+i \tan(e+fx))^3} \right)}{a^3 f}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^n)/(a + I*a*Tan[e + f*x])^3,x]`

output `(c*(((I/48)*(A*(3 - n) - I*B*(3 + n))*Hypergeometric2F1[3, n, 1 + n, (1 - I*Tan[e + f*x])/2]*(c - I*c*Tan[e + f*x])^n)/(c*n) + ((I*A - B)*(c - I*c*Tan[e + f*x])^n)/(6*c*(1 + I*Tan[e + f*x])^3)))/(a^3*f)`

3.727.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 78 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[(b*c - a*d)^n*(a + b*x)^(m + 1)/(b^(n + 1)*(m + 1))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m}, x] && !IntegerQ[m] && IntegerQ[n]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.727.4 Maple [F]

$$\int \frac{(A + B \tan(fx + e))(c - ic \tan(fx + e))^n}{(a + ia \tan(fx + e))^3} dx$$

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e))^3,x)`

output `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e))^3,x)`

3.727.5 Fricas [F]

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^3} dx \\ &= \int \frac{(B \tan(fx + e) + A)(-ic \tan(fx + e) + c)^n}{(ia \tan(fx + e) + a)^3} dx \end{aligned}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output `integral(1/8*((A - I*B)*e^(6*I*f*x + 6*I*e) + (3*A - I*B)*e^(4*I*f*x + 4*I*e) + (3*A + I*B)*e^(2*I*f*x + 2*I*e) + A + I*B)*(2*c/(e^(2*I*f*x + 2*I*e) + 1))^n*e^(-6*I*f*x - 6*I*e)/a^3, x)`

3.727.6 Sympy [F]

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^n}{(a + ia \tan(e + fx))^3} dx \\ &= \frac{i \left(\int \frac{A(-ic \tan(e+fx)+c)^n}{\tan^3(e+fx)-3i \tan^2(e+fx)-3 \tan(e+fx)+i} dx + \int \frac{B(-ic \tan(e+fx)+c)^n \tan(e+fx)}{\tan^3(e+fx)-3i \tan^2(e+fx)-3 \tan(e+fx)+i} dx \right)}{a^3} \end{aligned}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e))^3,x)`

output `I*(Integral(A*(-I*c*tan(e + f*x) + c))^n/(tan(e + f*x)**3 - 3*I*tan(e + f*x)**2 - 3*tan(e + f*x) + I), x) + Integral(B*(-I*c*tan(e + f*x) + c)**n*tan(e + f*x)/(tan(e + f*x)**3 - 3*I*tan(e + f*x)**2 - 3*tan(e + f*x) + I), x))/a**3`

3.727.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^n}{(a + i a \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.727.8 Giac [F]

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^n}{(a + i a \tan(e + fx))^3} dx \\ &= \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^n}{(i a \tan(fx + e) + a)^3} dx \end{aligned}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^n/(I*a*tan(f*x + e) + a)^3, x)`

3.727.9 Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^n}{(a + i a \tan(e + fx))^3} dx \\ &= \int \frac{(A + B \tan(e + fx)) (c - c \tan(e + fx) li)^n}{(a + a \tan(e + fx) li)^3} dx \end{aligned}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*li)^n)/(a + a*tan(e + f*x)*li)^3,x)`

output `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^n)/(a + a*tan(e + f*x)*1i)^3, x)`

3.727.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^n}{(a+ia \tan(e+fx))^3} dx$$

3.728
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^5}{(a+ia \tan(e+fx))^3} dx$$

3.728.1 Optimal result 6703
 3.728.2 Mathematica [A] (verified) 6704
 3.728.3 Rubi [A] (verified) 6704
 3.728.4 Maple [A] (verified) 6706
 3.728.5 Fricas [A] (verification not implemented) 6707
 3.728.6 Sympy [A] (verification not implemented) 6707
 3.728.7 Maxima [F(-2)] 6708
 3.728.8 Giac [B] (verification not implemented) 6708
 3.728.9 Mupad [B] (verification not implemented) 6709

3.728.1 Optimal result

Integrand size = 41, antiderivative size = 191

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^5}{(a + ia \tan(e + fx))^3} dx \\ &= -\frac{8(A + 4iB)c^5 x}{a^3} - \frac{8(iA - 4B)c^5 \log(\cos(e + fx))}{a^3 f} \\ & \quad + \frac{16(A + iB)c^5}{3a^3 f(i - \tan(e + fx))^3} + \frac{8(2iA - 3B)c^5}{a^3 f(i - \tan(e + fx))^2} \\ & \quad - \frac{8(3A + 7iB)c^5}{a^3 f(i - \tan(e + fx))} + \frac{(A + 8iB)c^5 \tan(e + fx)}{a^3 f} + \frac{Bc^5 \tan^2(e + fx)}{2a^3 f} \end{aligned}$$

output

```
-8*(A+4*I*B)*c^5*x/a^3-8*(I*A-4*B)*c^5*ln(cos(f*x+e))/a^3/f+16/3*(A+I*B)*c^5/a^3/f/(I-tan(f*x+e))^3+8*(2*I*A-3*B)*c^5/a^3/f/(I-tan(f*x+e))^2-8*(3*A+7*I*B)*c^5/a^3/f/(I-tan(f*x+e))+(A+8*I*B)*c^5*tan(f*x+e)/a^3/f+1/2*B*c^5*tan(f*x+e)^2/a^3/f
```

3.728.2 Mathematica [A] (verified)

Time = 5.01 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.80

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^5}{(a + i a \tan(e + fx))^3} dx$$

$$= \frac{\frac{2(A+4iB)c^5(i+\tan(e+fx))^4}{a^3(-i+\tan(e+fx))^3} + \frac{B(c-ic \tan(e+fx))^5}{(a+ia \tan(e+fx))^3} - \frac{16i(A+4iB)c^5 \left(-\log(i-\tan(e+fx)) + \frac{2(-4i+9 \tan(e+fx)+9i \tan^2(e+fx))}{3(-i+\tan(e+fx))^3} \right)}{a^3}}{2f}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^5)/(a + I*a*Tan[e + f*x])^3,x]`

output `((2*(A + (4*I)*B)*c^5*(I + Tan[e + f*x])^4)/(a^3*(-I + Tan[e + f*x])^3) + (B*(c - I*c*Tan[e + f*x])^5)/(a + I*a*Tan[e + f*x])^3 - ((16*I)*(A + (4*I)*B)*c^5*(-Log[I - Tan[e + f*x]] + (2*(-4*I + 9*Tan[e + f*x] + (9*I)*Tan[e + f*x]^2))/(3*(-I + Tan[e + f*x])^3)))/a^3)/(2*f)`

3.728.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.72, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - i c \tan(e + fx))^5 (A + B \tan(e + fx))}{(a + i a \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{(c - i c \tan(e + fx))^5 (A + B \tan(e + fx))}{(a + i a \tan(e + fx))^3} dx$$

↓ 4071

$$ac \int \frac{c^4(1-i \tan(e+fx))^4(A+B \tan(e+fx))}{a^4(i \tan(e+fx)+1)^4} d \tan(e + fx)$$

↓ 27

$$\frac{c^5 \int \frac{(1-i \tan(e+fx))^4 (A+B \tan(e+fx))}{(i \tan(e+fx)+1)^4} d \tan(e+fx)}{a^3 f}$$

↓ 86

$$\frac{c^5 \int \left(\frac{16(A+iB)}{(\tan(e+fx)-i)^4} + A \left(\frac{8iB}{A} + 1 \right) + B \tan(e+fx) + \frac{8i(A+4iB)}{\tan(e+fx)-i} - \frac{8(3A+7iB)}{(\tan(e+fx)-i)^2} + \frac{16(3B-2iA)}{(\tan(e+fx)-i)^3} \right) d \tan(e+fx)}{a^3 f}$$

↓ 2009

$$\frac{c^5 \left((A + 8iB) \tan(e+fx) - \frac{8(3A+7iB)}{-\tan(e+fx)+i} + \frac{8(-3B+2iA)}{(-\tan(e+fx)+i)^2} + \frac{16(A+iB)}{3(-\tan(e+fx)+i)^3} + 8(-4B + iA) \log(-\tan(e+fx)) \right)}{a^3 f}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^5)/(a + I*a*Tan[e + f*x])^3,x]`

output `(c^5*(8*(I*A - 4*B)*Log[I - Tan[e + f*x]] + (16*(A + I*B))/(3*(I - Tan[e + f*x])^3) + (8*((2*I)*A - 3*B))/(I - Tan[e + f*x])^2 - (8*(3*A + (7*I)*B))/(I - Tan[e + f*x]) + (A + (8*I)*B)*Tan[e + f*x] + (B*Tan[e + f*x]^2)/2))/(a^3*f)`

3.728.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.728. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^5}{(a+ia \tan(e+fx))^3} dx$


```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.728.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.49

method	result
derivativedivides	$\frac{c^5 A \tan(fx+e)}{f a^3} + \frac{8ic^5 \tan(fx+e)B}{f a^3} + \frac{B c^5 \tan(fx+e)^2}{2a^3 f} - \frac{8c^5 A \arctan(\tan(fx+e))}{f a^3} + \frac{4ic^5 A \ln(1+\tan(fx+e)^2)}{f a^3}$
default	$\frac{c^5 A \tan(fx+e)}{f a^3} + \frac{8ic^5 \tan(fx+e)B}{f a^3} + \frac{B c^5 \tan(fx+e)^2}{2a^3 f} - \frac{8c^5 A \arctan(\tan(fx+e))}{f a^3} + \frac{4ic^5 A \ln(1+\tan(fx+e)^2)}{f a^3}$
risch	$-\frac{18c^5 e^{-2i(fx+e)} B}{a^3 f} + \frac{6ic^5 e^{-2i(fx+e)} A}{a^3 f} + \frac{4c^5 e^{-4i(fx+e)} B}{a^3 f} - \frac{2ic^5 e^{-4i(fx+e)} A}{a^3 f} - \frac{2c^5 e^{-6i(fx+e)} B}{3a^3 f} + \frac{2ic^5 e^{-6i(fx+e)} A}{3a^3 f}$
norman	$\frac{(8ic^5 B+c^5 A) \tan(fx+e)^7}{af} + \frac{(32ic^5 B+9c^5 A) \tan(fx+e)}{af} + \frac{(80ic^5 B+27c^5 A) \tan(fx+e)^5}{af} - \frac{8(4ic^5 B+c^5 A)x}{a} - \frac{80ic^5 A+233c^5 B}{6af}$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5/(a+I*a*tan(f*x+e))^3,x,method=_R
ETURNVERBOSE)
```

```
output 1/f*c^5/a^3*A*tan(f*x+e)+8*I/f*c^5/a^3*tan(f*x+e)*B+1/2*B*c^5*tan(f*x+e)^2
/a^3/f-8/f*c^5/a^3*A*arctan(tan(f*x+e))+4*I/f*c^5/a^3*A*ln(1+tan(f*x+e)^2)
-32*I/f*c^5/a^3*B*arctan(tan(f*x+e))-16/f*c^5/a^3*B*ln(1+tan(f*x+e)^2)+56*
I/f*c^5/a^3/(-I+tan(f*x+e))*B+24/f*c^5/a^3/(-I+tan(f*x+e))*A+16*I/f*c^5/a^
3/(-I+tan(f*x+e))^2*A-24/f*c^5/a^3/(-I+tan(f*x+e))^2*B-16/3/f*c^5/a^3/(-I+
tan(f*x+e))^3*A-16/3*I/f*c^5/a^3/(-I+tan(f*x+e))^3*B
```

3.728. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^5}{(a+ia \tan(e+fx))^3} dx$

3.728.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.40

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^5}{(a + i a \tan(e + fx))^3} dx =$$

$$\frac{2(24(A + 4iB)c^5 f x e^{(10i f x + 10i e)} + 4(-iA + 4B)c^5 e^{(4i f x + 4i e)} + (iA - 4B)c^5 e^{(2i f x + 2i e)} + (-iA + B)c^5 e^{(0i f x + 0i e)})}{(a + i a \tan(e + fx))^3}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output `-2/3*(24*(A + 4*I*B)*c^5*f*x*e^(10*I*f*x + 10*I*e) + 4*(-I*A + 4*B)*c^5*e^(4*I*f*x + 4*I*e) + (I*A - 4*B)*c^5*e^(2*I*f*x + 2*I*e) + (-I*A + B)*c^5 + 12*(4*(A + 4*I*B)*c^5*f*x + (-I*A + 4*B)*c^5)*e^(8*I*f*x + 8*I*e) + 6*(4*(A + 4*I*B)*c^5*f*x + 3*(-I*A + 4*B)*c^5)*e^(6*I*f*x + 6*I*e) + 12*((I*A - 4*B)*c^5*e^(10*I*f*x + 10*I*e) + 2*(I*A - 4*B)*c^5*e^(8*I*f*x + 8*I*e) + (I*A - 4*B)*c^5*e^(6*I*f*x + 6*I*e))*log(e^(2*I*f*x + 2*I*e) + 1)/(a^3*f*e^(10*I*f*x + 10*I*e) + 2*a^3*f*e^(8*I*f*x + 8*I*e) + a^3*f*e^(6*I*f*x + 6*I*e))`

3.728.6 Sympy [A] (verification not implemented)

Time = 0.77 (sec) , antiderivative size = 473, normalized size of antiderivative = 2.48

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^5}{(a + i a \tan(e + fx))^3} dx$$

$$= \frac{2iAc^5 - 16Bc^5 + (2iAc^5 e^{2ie} - 14Bc^5 e^{2ie}) e^{2ifx}}{a^3 f e^{4ie} e^{4ifx} + 2a^3 f e^{2ie} e^{2ifx} + a^3 f}$$

$$+ \left\{ \frac{((2iAa^6 c^5 f^2 e^{6ie} - 2Ba^6 c^5 f^2 e^{6ie}) e^{-6ifx} + (-6iAa^6 c^5 f^2 e^{8ie} + 12Ba^6 c^5 f^2 e^{8ie}) e^{-4ifx} + (18iAa^6 c^5 f^2 e^{10ie} - 54Ba^6 c^5 f^2 e^{10ie}) e^{-2ifx}) e^{-12ie}}{3a^9 f^3} \right.$$

$$\left. + x \left(-\frac{16Ac^5 - 64iBc^5}{a^3} + \frac{(-16Ac^5 e^{6ie} + 12Ac^5 e^{4ie} - 8Ac^5 e^{2ie} + 4Ac^5 - 64iBc^5 e^{6ie} + 36iBc^5 e^{4ie} - 16iBc^5 e^{2ie} + 4iBc^5) e^{-6ie}}{a^3} \right) \right.$$

$$- \frac{8ic^5(A + 4iB) \log(e^{2ifx} + e^{-2ie})}{a^3 f} + \frac{x(-16Ac^5 - 64iBc^5)}{a^3}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**5/(a+I*a*tan(f*x+e))**3,x)`

output `(2*I*A*c**5 - 16*B*c**5 + (2*I*A*c**5*exp(2*I*e) - 14*B*c**5*exp(2*I*e))*exp(2*I*f*x))/(a**3*f*exp(4*I*e)*exp(4*I*f*x) + 2*a**3*f*exp(2*I*e)*exp(2*I*f*x) + a**3*f) + Piecewise((((2*I*A*a**6*c**5*f**2*exp(6*I*e) - 2*B*a**6*c**5*f**2*exp(6*I*e))*exp(-6*I*f*x) + (-6*I*A*a**6*c**5*f**2*exp(8*I*e) + 12*B*a**6*c**5*f**2*exp(8*I*e))*exp(-4*I*f*x) + (18*I*A*a**6*c**5*f**2*exp(10*I*e) - 54*B*a**6*c**5*f**2*exp(10*I*e))*exp(-2*I*f*x))*exp(-12*I*e)/(3*a**9*f**3), Ne(a**9*f**3*exp(12*I*e), 0)), (x*(-(-16*A*c**5 - 64*I*B*c**5)/a**3 + (-16*A*c**5*exp(6*I*e) + 12*A*c**5*exp(4*I*e) - 8*A*c**5*exp(2*I*e) + 4*A*c**5 - 64*I*B*c**5*exp(6*I*e) + 36*I*B*c**5*exp(4*I*e) - 16*I*B*c**5*exp(2*I*e) + 4*I*B*c**5)*exp(-6*I*e)/a**3), True)) - 8*I*c**5*(A + 4*I*B)*log(exp(2*I*f*x) + exp(-2*I*e))/(a**3*f) + x*(-16*A*c**5 - 64*I*B*c**5)/a**3`

3.728.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^5}{(a + ia \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.728.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 491 vs. $2(163) = 326$.

Time = 1.23 (sec) , antiderivative size = 491, normalized size of antiderivative = 2.57

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^5}{(a + ia \tan(e + fx))^3} dx$$

$$= 2 \left(\frac{60(-iAc^5 + 4Bc^5) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)}{a^3} - \frac{120(-iAc^5 + 4Bc^5) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - i)}{a^3} - \frac{60(iAc^5 - 4Bc^5) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)}{a^3} \right)$$

3.728. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^5}{(a+ia \tan(e+fx))^3} dx$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^5/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")`

output `2/15*(60*(-I*A*c^5 + 4*B*c^5)*log(tan(1/2*f*x + 1/2*e) + 1)/a^3 - 120*(-I*A*c^5 + 4*B*c^5)*log(tan(1/2*f*x + 1/2*e) - I)/a^3 - 60*(I*A*c^5 - 4*B*c^5)*log(tan(1/2*f*x + 1/2*e) - 1)/a^3 + 15*(6*I*A*c^5*tan(1/2*f*x + 1/2*e)^4 - 24*B*c^5*tan(1/2*f*x + 1/2*e)^4 - A*c^5*tan(1/2*f*x + 1/2*e)^3 - 8*I*B*c^5*tan(1/2*f*x + 1/2*e)^3 - 12*I*A*c^5*tan(1/2*f*x + 1/2*e)^2 + 49*B*c^5*tan(1/2*f*x + 1/2*e)^2 + A*c^5*tan(1/2*f*x + 1/2*e) + 8*I*B*c^5*tan(1/2*f*x + 1/2*e) + 6*I*A*c^5 - 24*B*c^5)/((tan(1/2*f*x + 1/2*e)^2 - 1)^2*a^3) - 2*(147*I*A*c^5*tan(1/2*f*x + 1/2*e)^6 - 588*B*c^5*tan(1/2*f*x + 1/2*e)^6 + 942*A*c^5*tan(1/2*f*x + 1/2*e)^5 + 3708*I*B*c^5*tan(1/2*f*x + 1/2*e)^5 - 2445*I*A*c^5*tan(1/2*f*x + 1/2*e)^4 + 9660*B*c^5*tan(1/2*f*x + 1/2*e)^4 - 3460*A*c^5*tan(1/2*f*x + 1/2*e)^3 - 13240*I*B*c^5*tan(1/2*f*x + 1/2*e)^3 + 2445*I*A*c^5*tan(1/2*f*x + 1/2*e)^2 - 9660*B*c^5*tan(1/2*f*x + 1/2*e)^2 + 942*A*c^5*tan(1/2*f*x + 1/2*e) + 3708*I*B*c^5*tan(1/2*f*x + 1/2*e) - 147*I*A*c^5 + 588*B*c^5)/(a^3*(tan(1/2*f*x + 1/2*e) - I)^6))/f`

3.728.9 Mupad [B] (verification not implemented)

Time = 8.79 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.22

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^5}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{\ln(\tan(e + fx) - i) \left(-\frac{32Bc^5}{a^3} + \frac{Ac^5 8i}{a^3} \right)}{f} + \frac{\tan(e + fx) \left(\frac{c^5(A+B4i)}{a^3} + \frac{Bc^5 4i}{a^3} \right)}{f}$$

$$+ \frac{\frac{5(-32Bc^5 + Ac^5 8i)}{3a^3} + \tan(e + fx) \left(\frac{(-32Bc^5 + Ac^5 8i) 4i}{a^3} + \frac{Bc^5 40i}{a^3} \right) - \tan(e + fx)^2 \left(\frac{3(-32Bc^5 + Ac^5 8i)}{a^3} + \frac{40Bc^5}{a^3} \right)}{f(-\tan(e + fx)^3 li - 3 \tan(e + fx)^2 + \tan(e + fx) 3i + 1)}$$

$$+ \frac{Bc^5 \tan(e + fx)^2}{2a^3 f}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^5)/(a + a*tan(e + f*x)*1i)^3,x)`

```

output (log(tan(e + f*x) - 1i)*((A*c^5*8i)/a^3 - (32*B*c^5)/a^3))/f + (tan(e + f*
x)*((c^5*(A + B*4i))/a^3 + (B*c^5*4i)/a^3))/f + ((5*(A*c^5*8i - 32*B*c^5))
/(3*a^3) + tan(e + f*x)*((A*c^5*8i - 32*B*c^5)*4i)/a^3 + (B*c^5*40i)/a^3)
- tan(e + f*x)^2*((3*(A*c^5*8i - 32*B*c^5))/a^3 + (40*B*c^5)/a^3) + (16*B
*c^5)/a^3)/(f*(tan(e + f*x)*3i - 3*tan(e + f*x)^2 - tan(e + f*x)^3*1i + 1)
) + (B*c^5*tan(e + f*x)^2)/(2*a^3*f)

```

3.728.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^5}{(a+ia \tan(e+fx))^3} dx$$

3.729
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^4}{(a+ia \tan(e+fx))^3} dx$$

3.729.1 Optimal result 6711
 3.729.2 Mathematica [A] (verified) 6711
 3.729.3 Rubi [A] (verified) 6712
 3.729.4 Maple [A] (verified) 6714
 3.729.5 Fricas [A] (verification not implemented) 6714
 3.729.6 Sympy [A] (verification not implemented) 6715
 3.729.7 Maxima [F(-2)] 6716
 3.729.8 Giac [B] (verification not implemented) 6716
 3.729.9 Mupad [B] (verification not implemented) 6717

3.729.1 Optimal result

Integrand size = 41, antiderivative size = 164

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^3} dx$$

$$= -\frac{(A + 7iB)c^4x}{a^3} - \frac{(iA - 7B)c^4 \log(\cos(e + fx))}{a^3 f} + \frac{8(A + iB)c^4}{3a^3 f(i - \tan(e + fx))^3}$$

$$+ \frac{2(3iA - 5B)c^4}{a^3 f(i - \tan(e + fx))^2} - \frac{6(A + 3iB)c^4}{a^3 f(i - \tan(e + fx))} + \frac{iBc^4 \tan(e + fx)}{a^3 f}$$

```
output -(A+7*I*B)*c^4*x/a^3-(I*A-7*B)*c^4*ln(cos(f*x+e))/a^3/f+8/3*(A+I*B)*c^4/a^3/f/(I-tan(f*x+e))^3+2*(3*I*A-5*B)*c^4/a^3/f/(I-tan(f*x+e))^2-6*(A+3*I*B)*c^4/a^3/f/(I-tan(f*x+e))+I*B*c^4*tan(f*x+e)/a^3/f
```

3.729.2 Mathematica [A] (verified)

Time = 5.71 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.65

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{c^4 \left(\frac{B(i + \tan(e + fx))^4}{(a + ia \tan(e + fx))^3} - \frac{i(A + 7iB) \left(-\log(i - \tan(e + fx)) + \frac{2(-4i + 9 \tan(e + fx) + 9i \tan^2(e + fx))}{3(-i + \tan(e + fx))^3} \right)}{a^3} \right)}{f}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^4)/(a + I*a*Tan[e + f*x])^3,x]`

output `(c^4*((B*(I + Tan[e + f*x])^4)/(a + I*a*Tan[e + f*x])^3 - (I*(A + (7*I)*B)*(-Log[I - Tan[e + f*x]] + (2*(-4*I + 9*Tan[e + f*x] + (9*I)*Tan[e + f*x]^2))/(3*(-I + Tan[e + f*x])^3)))/a^3)/f`

3.729.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.72, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - ic \tan(e + fx))^4 (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c - ic \tan(e + fx))^4 (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{c^3 (1 - i \tan(e + fx))^3 (A + B \tan(e + fx))}{a^4 (i \tan(e + fx) + 1)^4} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{c^4 \int \frac{(1 - i \tan(e + fx))^3 (A + B \tan(e + fx))}{(i \tan(e + fx) + 1)^4} d \tan(e + fx)}{a^3 f} \\
 & \quad \downarrow \text{86} \\
 & \frac{c^4 \int \left(\frac{8(A + iB)}{(\tan(e + fx) - i)^4} + iB + \frac{i(A + 7iB)}{\tan(e + fx) - i} - \frac{6(A + 3iB)}{(\tan(e + fx) - i)^2} + \frac{4(5B - 3iA)}{(\tan(e + fx) - i)^3} \right) d \tan(e + fx)}{a^3 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{c^4 \left(\frac{2(-5B + 3iA)}{(-\tan(e + fx) + i)^2} - \frac{6(A + 3iB)}{-\tan(e + fx) + i} + \frac{8(A + iB)}{3(-\tan(e + fx) + i)^3} + (-7B + iA) \log(-\tan(e + fx) + i) + iB \tan(e + fx) \right)}{a^3 f}
 \end{aligned}$$

3.729. $\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^3} dx$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^4)/(a + I*a*Tan[e + f*x])^3,x]`

output `(c^4*((I*A - 7*B)*Log[I - Tan[e + f*x]] + (8*(A + I*B))/(3*(I - Tan[e + f*x]))^3) + (2*((3*I)*A - 5*B))/(I - Tan[e + f*x])^2 - (6*(A + (3*I)*B))/(I - Tan[e + f*x]) + I*B*Tan[e + f*x]))/(a^3*f)`

3.729.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x_, x], x] /; FreeQ[a, x] && !MatchQ[F_x_, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.729.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.51

method	result
derivativedivides	$\frac{iBc^4 \tan(fx+e)}{a^3 f} - \frac{8c^4 A}{3f a^3 (-i+\tan(fx+e))^3} - \frac{8ic^4 B}{3f a^3 (-i+\tan(fx+e))^3} + \frac{18ic^4 B}{f a^3 (-i+\tan(fx+e))} + \frac{6c^4 A}{f a^3 (-i+\tan(fx+e))}$
default	$\frac{iBc^4 \tan(fx+e)}{a^3 f} - \frac{8c^4 A}{3f a^3 (-i+\tan(fx+e))^3} - \frac{8ic^4 B}{3f a^3 (-i+\tan(fx+e))^3} + \frac{18ic^4 B}{f a^3 (-i+\tan(fx+e))} + \frac{6c^4 A}{f a^3 (-i+\tan(fx+e))}$
risch	$-\frac{5c^4 e^{-2i(fx+e)} B}{a^3 f} + \frac{ic^4 e^{-2i(fx+e)} A}{a^3 f} + \frac{3c^4 e^{-4i(fx+e)} B}{2a^3 f} - \frac{ic^4 e^{-4i(fx+e)} A}{2a^3 f} - \frac{c^4 e^{-6i(fx+e)} B}{3a^3 f} + \frac{ic^4 e^{-6i(fx+e)} A}{3a^3 f}$
norman	$\frac{(7ic^4 B+2c^4 A) \tan(fx+e)}{af} + \frac{ic^4 B \tan(fx+e)^7}{af} - \frac{(7ic^4 B+c^4 A)x}{a} - \frac{-8ic^4 A+32c^4 B}{3af} - \frac{3(7ic^4 B+c^4 A)x \tan(fx+e)^2}{a} - \frac{3(7ic^4 B+c^4 A)}{6(a^2)}$

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4/(a+I*a*tan(f*x+e))^3,x,method=_R
ETURNVERBOSE)`

output `I*B*c^4*tan(f*x+e)/a^3/f-8/3/f*c^4/a^3/(-I+tan(f*x+e))^3*A-8/3*I/f*c^4/a^3
/(-I+tan(f*x+e))^3*B+18*I/f*c^4/a^3/(-I+tan(f*x+e))*B+6/f*c^4/a^3/(-I+tan(
f*x+e))*A+1/2*I/f*c^4/a^3*A*ln(1+tan(f*x+e)^2)-7/2/f*c^4/a^3*B*ln(1+tan(f*
x+e)^2)-1/f*c^4/a^3*A*arctan(tan(f*x+e))-7*I/f*c^4/a^3*B*arctan(tan(f*x+e)
) +6*I/f*c^4/a^3/(-I+tan(f*x+e))^2*A-10/f*c^4/a^3/(-I+tan(f*x+e))^2*B`

3.729.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.20

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^3} dx =$$

$$\frac{12(A + 7iB)c^4 f x e^{(8i f x + 8i e)} + 3(-iA + 7B)c^4 e^{(4i f x + 4i e)} - (-iA + 7B)c^4 e^{(2i f x + 2i e)} + 2(-iA + B)c^4 e^{(i f x + i e)}}{6(a^2)}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4/(a+I*a*tan(f*x+e))^3,x, al
gorithm="fricas")`

output
$$-1/6*(12*(A + 7*I*B)*c^4*f*x*e^(8*I*f*x + 8*I*e) + 3*(-I*A + 7*B)*c^4*e^(4*I*f*x + 4*I*e) - (-I*A + 7*B)*c^4*e^(2*I*f*x + 2*I*e) + 2*(-I*A + B)*c^4 + 6*(2*(A + 7*I*B)*c^4*f*x + (-I*A + 7*B)*c^4)*e^(6*I*f*x + 6*I*e) + 6*((I*A - 7*B)*c^4*e^(8*I*f*x + 8*I*e) + (I*A - 7*B)*c^4*e^(6*I*f*x + 6*I*e))*\log(e^(2*I*f*x + 2*I*e) + 1)/(a^3*f*e^(8*I*f*x + 8*I*e) + a^3*f*e^(6*I*f*x + 6*I*e))$$

3.729.6 Sympy [A] (verification not implemented)

Time = 0.64 (sec) , antiderivative size = 403, normalized size of antiderivative = 2.46

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^3} dx = -\frac{2Bc^4}{a^3 f e^{2ie} e^{2ifx} + a^3 f} + \left\{ \frac{((2iAa^6c^4f^2e^{6ie} - 2Ba^6c^4f^2e^{6ie})e^{-6ifx} + (-3iAa^6c^4f^2e^{8ie} + 9Ba^6c^4f^2e^{8ie})e^{-4ifx} + (6iAa^6c^4f^2e^{10ie} - 30Ba^6c^4f^2e^{10ie})e^{-2ifx})e^{-12ie}}{6a^9f^3} \right. \\ \left. + x \left(-\frac{2Ac^4 - 14iBc^4}{a^3} + \frac{(-2Ac^4e^{6ie} + 2Ac^4e^{4ie} - 2Ac^4e^{2ie} + 2Ac^4 - 14iBc^4e^{6ie} + 10iBc^4e^{4ie} - 6iBc^4e^{2ie} + 2iBc^4)e^{-6ie}}{a^3} \right) \right. \\ \left. - \frac{ic^4(A + 7iB) \log(e^{2ifx} + e^{-2ie})}{a^3 f} + \frac{x(-2Ac^4 - 14iBc^4)}{a^3} \right.$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**4/(a+I*a*tan(f*x+e))**3,x)`

output
$$-2*B*c**4/(a**3*f*\exp(2*I*e)*\exp(2*I*f*x) + a**3*f) + \text{Piecewise}(\left((2*I*A*a**6*c**4*f**2*\exp(6*I*e) - 2*B*a**6*c**4*f**2*\exp(6*I*e))*\exp(-6*I*f*x) + (-3*I*A*a**6*c**4*f**2*\exp(8*I*e) + 9*B*a**6*c**4*f**2*\exp(8*I*e))*\exp(-4*I*f*x) + (6*I*A*a**6*c**4*f**2*\exp(10*I*e) - 30*B*a**6*c**4*f**2*\exp(10*I*e))*\exp(-2*I*f*x) \right)*\exp(-12*I*e)/(6*a**9*f**3), \text{Ne}(a**9*f**3*\exp(12*I*e), 0)), (x*(-(-2*A*c**4 - 14*I*B*c**4)/a**3 + (-2*A*c**4*\exp(6*I*e) + 2*A*c**4*\exp(4*I*e) - 2*A*c**4*\exp(2*I*e) + 2*A*c**4 - 14*I*B*c**4*\exp(6*I*e) + 10*I*B*c**4*\exp(4*I*e) - 6*I*B*c**4*\exp(2*I*e) + 2*I*B*c**4)*\exp(-6*I*e)/a**3), \text{True})) - I*c**4*(A + 7*I*B)*\log(\exp(2*I*f*x) + \exp(-2*I*e))/(a**3*f) + x*(-2*A*c**4 - 14*I*B*c**4)/a**3$$

3.729.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.729.8 Giac [B] (verification not implemented)Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 409 vs. $2(138) = 276$.

Time = 1.04 (sec) , antiderivative size = 409, normalized size of antiderivative = 2.49

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{30(-iAc^4 + 7Bc^4) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)}{a^3} + \frac{60(iAc^4 - 7Bc^4) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - i)}{a^3} - \frac{30(iAc^4 - 7Bc^4) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)}{a^3} - \frac{30(iAc^4 - 7Bc^4) \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + i)}{a^3}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^4/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")
```

```
output 1/30*(30*(-I*A*c^4 + 7*B*c^4)*log(tan(1/2*f*x + 1/2*e) + 1)/a^3 + 60*(I*A*c^4 - 7*B*c^4)*log(tan(1/2*f*x + 1/2*e) - I)/a^3 - 30*(I*A*c^4 - 7*B*c^4)*log(tan(1/2*f*x + 1/2*e) - 1)/a^3 - 30*(-I*A*c^4*tan(1/2*f*x + 1/2*e)^2 + 7*B*c^4*tan(1/2*f*x + 1/2*e)^2 + 2*I*B*c^4*tan(1/2*f*x + 1/2*e) + I*A*c^4 - 7*B*c^4)/((tan(1/2*f*x + 1/2*e)^2 - 1)*a^3) - (147*I*A*c^4*tan(1/2*f*x + 1/2*e)^6 - 1029*B*c^4*tan(1/2*f*x + 1/2*e)^6 + 1002*A*c^4*tan(1/2*f*x + 1/2*e)^5 + 6534*I*B*c^4*tan(1/2*f*x + 1/2*e)^5 - 2445*I*A*c^4*tan(1/2*f*x + 1/2*e)^4 + 17115*B*c^4*tan(1/2*f*x + 1/2*e)^4 - 3820*A*c^4*tan(1/2*f*x + 1/2*e)^3 - 23860*I*B*c^4*tan(1/2*f*x + 1/2*e)^3 + 2445*I*A*c^4*tan(1/2*f*x + 1/2*e)^2 - 17115*B*c^4*tan(1/2*f*x + 1/2*e)^2 + 1002*A*c^4*tan(1/2*f*x + 1/2*e) + 6534*I*B*c^4*tan(1/2*f*x + 1/2*e) - 147*I*A*c^4 + 1029*B*c^4)/(a^3*(tan(1/2*f*x + 1/2*e) - I)^6))/f
```

3.729. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^4}{(a+ia \tan(e+fx))^3} dx$

3.729.9 Mupad [B] (verification not implemented)

Time = 10.61 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.62

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^4}{(a + ia \tan(e + fx))^3} dx =$$

$$\frac{c^4 (25 B \tan(e + fx) - \frac{B 32i}{3} - A \tan(e + fx) 6i - \frac{8A}{3} - A \ln(-1 - \tan(e + fx) 1i) - B \ln(-1 - \tan(e + fx) 1i))}{(a + ia \tan(e + fx))^3}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^4)/(a + a*tan(e + f*x)*1i)^3,x)`

output `-(c^4*(25*B*tan(e + f*x) - (B*32i)/3 - A*tan(e + f*x)*6i - (8*A)/3 - A*log(-tan(e + f*x)*1i - 1) - B*log(-tan(e + f*x)*1i - 1)*7i + 6*A*tan(e + f*x)^2 + B*tan(e + f*x)^2*15i + 3*B*tan(e + f*x)^3 + B*tan(e + f*x)^4*1i + 3*A*tan(e + f*x)^2*log(-tan(e + f*x)*1i - 1) + A*tan(e + f*x)^3*log(-tan(e + f*x)*1i - 1)*1i + B*tan(e + f*x)^2*log(-tan(e + f*x)*1i - 1)*21i - 7*B*tan(e + f*x)^3*log(-tan(e + f*x)*1i - 1) - A*tan(e + f*x)*log(-tan(e + f*x)*1i - 1)*3i + 21*B*tan(e + f*x)*log(-tan(e + f*x)*1i - 1)*1i)/(a^3*f*(tan(e + f*x)*1i + 1)^3)`

3.730
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^3}{(a+ia \tan(e+fx))^3} dx$$

3.730.1 Optimal result 6718
 3.730.2 Mathematica [A] (verified) 6718
 3.730.3 Rubi [A] (verified) 6719
 3.730.4 Maple [A] (verified) 6721
 3.730.5 Fricas [A] (verification not implemented) 6721
 3.730.6 Sympy [A] (verification not implemented) 6722
 3.730.7 Maxima [F(-2)] 6722
 3.730.8 Giac [B] (verification not implemented) 6723
 3.730.9 Mupad [B] (verification not implemented) 6723

3.730.1 Optimal result

Integrand size = 41, antiderivative size = 135

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{(a + ia \tan(e + fx))^3} dx$$

$$= -\frac{iBc^3x}{a^3} + \frac{Bc^3 \log(\cos(e + fx))}{a^3 f} - \frac{2Bc^3}{a^3 f (i - \tan(e + fx))^2}$$

$$- \frac{4iBc^3}{a^3 f (i - \tan(e + fx))} + \frac{(iA - B)c^3 (1 - i \tan(e + fx))^3}{6a^3 f (1 + i \tan(e + fx))^3}$$

output `-I*B*c^3*x/a^3+B*c^3*ln(cos(f*x+e))/a^3/f-2*B*c^3/a^3/f/(I-tan(f*x+e))^2-4
 *I*B*c^3/a^3/f/(I-tan(f*x+e))+1/6*(I*A-B)*c^3*(1-I*tan(f*x+e))^3/a^3/f/(1+
 I*tan(f*x+e))^3`

3.730.2 Mathematica [A] (verified)

Time = 5.91 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.56

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{(a + ia \tan(e + fx))^3} dx$$

$$= -\frac{c^3 \left(3B \log(i - \tan(e + fx)) + \frac{A+7iB-18B \tan(e+fx)-3(A+5iB) \tan^2(e+fx)}{(-i+\tan(e+fx))^3} \right)}{3a^3 f}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^3)/(a + I*a*Tan[e + f*x])^3,x]`

output `-1/3*(c^3*(3*B*Log[I - Tan[e + f*x]] + (A + (7*I)*B - 18*B*Tan[e + f*x] - 3*(A + (5*I)*B)*Tan[e + f*x]^2)/(-I + Tan[e + f*x])^3))/(a^3*f)`

3.730.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.81, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.146$, Rules used = {3042, 4071, 27, 87, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - ic \tan(e + fx))^3 (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c - ic \tan(e + fx))^3 (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{c^2(1-i \tan(e+fx))^2(A+B \tan(e+fx))}{a^4(i \tan(e+fx)+1)^4} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{c^3 \int \frac{(1-i \tan(e+fx))^2(A+B \tan(e+fx))}{(i \tan(e+fx)+1)^4} d \tan(e + fx)}{a^3 f} \\
 & \quad \downarrow \text{87} \\
 & \frac{c^3 \left(\frac{(-B+iA)(1-i \tan(e+fx))^3}{6(1+i \tan(e+fx))^3} - iB \int \frac{(1-i \tan(e+fx))^2}{(i \tan(e+fx)+1)^3} d \tan(e + fx) \right)}{a^3 f} \\
 & \quad \downarrow \text{49} \\
 & \frac{c^3 \left(\frac{(-B+iA)(1-i \tan(e+fx))^3}{6(1+i \tan(e+fx))^3} - iB \int \left(-\frac{i}{\tan(e+fx)-i} + \frac{4}{(\tan(e+fx)-i)^2} + \frac{4i}{(\tan(e+fx)-i)^3} \right) d \tan(e + fx) \right)}{a^3 f} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

3.730. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^3}{(a+ia \tan(e+fx))^3} dx$

$$\frac{c^3 \left(\frac{(-B+iA)(1-i \tan(e+fx))^3}{6(1+i \tan(e+fx))^3} - iB \left(\frac{4}{-\tan(e+fx)+i} - \frac{2i}{(-\tan(e+fx)+i)^2} - i \log(-\tan(e+fx)+i) \right) \right)}{a^3 f}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^3)/(a + I*a*Tan[e + f*x])^3,x]`

output `(c^3*((-I)*B*((-I)*Log[I - Tan[e + f*x]] - (2*I)/(I - Tan[e + f*x])^2 + 4/(I - Tan[e + f*x])) + ((I*A - B)*(1 - I*Tan[e + f*x])^3)/(6*(1 + I*Tan[e + f*x])^3)))/(a^3*f)`

3.730.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x],
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.730.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.00

method	result
risch	$-\frac{c^3 B e^{-2i(fx+e)}}{a^3 f} + \frac{c^3 B e^{-4i(fx+e)}}{2a^3 f} - \frac{c^3 e^{-6i(fx+e)} B}{6a^3 f} + \frac{ic^3 e^{-6i(fx+e)} A}{6a^3 f} - \frac{2ic^3 B x}{a^3} - \frac{2ic^3 B e}{a^3 f} + \frac{c^3 B \ln(e^{2i(fx+e)})}{a^3 f}$
derivativedivides	$\frac{2ic^3 A}{f a^3 (-i + \tan(fx+e))^2} - \frac{4c^3 B}{f a^3 (-i + \tan(fx+e))^2} - \frac{c^3 B \ln(1 + \tan(fx+e)^2)}{2f a^3} - \frac{ic^3 B \arctan(\tan(fx+e))}{f a^3} - \frac{c^3 B \ln(e^{2i(fx+e)})}{3f a^3}$
default	$\frac{2ic^3 A}{f a^3 (-i + \tan(fx+e))^2} - \frac{4c^3 B}{f a^3 (-i + \tan(fx+e))^2} - \frac{c^3 B \ln(1 + \tan(fx+e)^2)}{2f a^3} - \frac{ic^3 B \arctan(\tan(fx+e))}{f a^3} - \frac{c^3 B \ln(e^{2i(fx+e)})}{3f a^3}$
norman	$\frac{(ic^3 B + c^3 A) \tan(fx+e)}{af} + \frac{(5ic^3 B + c^3 A) \tan(fx+e)^5}{af} - \frac{-ic^3 A + 7c^3 B}{3af} - \frac{2(-ic^3 B + 5c^3 A) \tan(fx+e)^3}{3af} - \frac{3(-ic^3 A + 3c^3 B) \tan(fx+e)}{af} - \frac{c^3 B \ln(e^{2i(fx+e)})}{a^2 (1 + \tan(fx+e)^2)^3}$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3/(a+I*a*tan(f*x+e))^3,x,method=_R
ETURNVERBOSE)
```

```
output -1/a^3/f*c^3*B*exp(-2*I*(f*x+e))+1/2/a^3/f*c^3*B*exp(-4*I*(f*x+e))-1/6*c^3
/a^3/f*exp(-6*I*(f*x+e))*B+1/6*I*c^3/a^3/f*exp(-6*I*(f*x+e))*A-2*I/a^3*c^3
*B*x-2*I/a^3/f*c^3*B*e+1/a^3/f*c^3*B*ln(exp(2*I*(f*x+e))+1)
```

3.730.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.76

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{(-12i B c^3 f x e^{(6i f x + 6i e)} + 6 B c^3 e^{(6i f x + 6i e)} \log(e^{(2i f x + 2i e)} + 1) - 6 B c^3 e^{(4i f x + 4i e)} + 3 B c^3 e^{(2i f x + 2i e)} + (i A - c^3)) e^{(6i f x + 6i e)}}{6 a^3 f}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3/(a+I*a*tan(f*x+e))^3,x, al
gorithm="fricas")
```

3.730. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^3}{(a+ia \tan(e+fx))^3} dx$

output $1/6*(-12*I*B*c^3*f*x*e^{(6*I*f*x + 6*I*e)} + 6*B*c^3*e^{(6*I*f*x + 6*I*e)}*\log(e^{(2*I*f*x + 2*I*e)} + 1) - 6*B*c^3*e^{(4*I*f*x + 4*I*e)} + 3*B*c^3*e^{(2*I*f*x + 2*I*e)} + (I*A - B)*c^3)*e^{(-6*I*f*x - 6*I*e)}/(a^3*f)$

3.730.6 Sympy [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.91

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{(a + ia \tan(e + fx))^3} dx = -\frac{2iBc^3x}{a^3} + \frac{Bc^3 \log(e^{2ifx} + e^{-2ie})}{a^3 f}$$

$$+ \begin{cases} \frac{(-12Ba^6c^3f^2e^{10ie}e^{-2ifx} + 6Ba^6c^3f^2e^{8ie}e^{-4ifx} + (2iAa^6c^3f^2e^{6ie} - 2Ba^6c^3f^2e^{6ie})e^{-6ifx})e^{-12ie}}{12a^9f^3} & \text{for } a^9f^3e^{12ie} \neq 0 \\ x \left(\frac{2iBc^3}{a^3} + \frac{(Ac^3 - 2iBc^3e^{6ie} + 2iBc^3e^{4ie} - 2iBc^3e^{2ie} + iBc^3)e^{-6ie}}{a^3} \right) & \text{otherwise} \end{cases}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**3/(a+I*a*tan(f*x+e))**3,x)`

output $-2*I*B*c**3*x/a**3 + B*c**3*\log(\exp(2*I*f*x) + \exp(-2*I*e))/(a**3*f) + \text{Piecewise}(((-12*B*a**6*c**3*f**2*\exp(10*I*e)*\exp(-2*I*f*x) + 6*B*a**6*c**3*f**2*\exp(8*I*e)*\exp(-4*I*f*x) + (2*I*A*a**6*c**3*f**2*\exp(6*I*e) - 2*B*a**6*c**3*f**2*\exp(6*I*e))*\exp(-6*I*f*x))*\exp(-12*I*e)/(12*a**9*f**3), \text{Ne}(a**9*f**3*\exp(12*I*e), 0)), (x*(2*I*B*c**3/a**3 + (A*c**3 - 2*I*B*c**3*\exp(6*I*e) + 2*I*B*c**3*\exp(4*I*e) - 2*I*B*c**3*\exp(2*I*e) + I*B*c**3)*\exp(-6*I*e)/a**3), \text{True}))$

3.730.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{(a + ia \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.730.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 242 vs. $2(113) = 226$.

Time = 0.88 (sec) , antiderivative size = 242, normalized size of antiderivative = 1.79

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{30 Bc^3 \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) + 1)}{a^3} - \frac{60 Bc^3 \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - i)}{a^3} + \frac{30 Bc^3 \log(\tan(\frac{1}{2}fx + \frac{1}{2}e) - 1)}{a^3} + \frac{147 Bc^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)^6 - 60 Ac^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)^5 - 942 I Bc^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)^4 + 200 A^2 c^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)^3 + 3620 I Bc^3 \tan(\frac{1}{2}fx + \frac{1}{2}e)^2 - 60 A^2 c^3 \tan(\frac{1}{2}fx + \frac{1}{2}e) - 942 I Bc^3 \tan(\frac{1}{2}fx + \frac{1}{2}e) - 147 Bc^3}{a^3 (\tan(\frac{1}{2}fx + \frac{1}{2}e) - I)^6} / f$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^3/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")`

output `1/30*(30*B*c^3*log(tan(1/2*f*x + 1/2*e) + 1)/a^3 - 60*B*c^3*log(tan(1/2*f*x + 1/2*e) - I)/a^3 + 30*B*c^3*log(tan(1/2*f*x + 1/2*e) - 1)/a^3 + (147*B*c^3*tan(1/2*f*x + 1/2*e)^6 - 60*A*c^3*tan(1/2*f*x + 1/2*e)^5 - 942*I*B*c^3*tan(1/2*f*x + 1/2*e)^4 + 200*A^2*c^3*tan(1/2*f*x + 1/2*e)^3 + 3620*I*B*c^3*tan(1/2*f*x + 1/2*e)^2 + 2445*B*c^3*tan(1/2*f*x + 1/2*e) - 60*A*c^3*tan(1/2*f*x + 1/2*e) - 942*I*B*c^3*tan(1/2*f*x + 1/2*e) - 147*B*c^3)/(a^3*(tan(1/2*f*x + 1/2*e) - I)^6))/f`

3.730.9 Mupad [B] (verification not implemented)

Time = 8.68 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.10

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^3}{(a + ia \tan(e + fx))^3} dx =$$

$$\frac{c^3 (18 B \tan(e + fx) - B 7i - A - B \ln(-1 - \tan(e + fx) 1i) 3i + 3 A \tan(e + fx)^2 + B \tan(e + fx) \ln(-1 - \tan(e + fx) 1i))}{(3 a^3 f (\tan(e + fx) 1i + 1)^3)}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^3)/(a + a*tan(e + f*x)*1i)^3,x)`

output `-(c^3*(18*B*tan(e + f*x) - B*7i - A - B*log(-tan(e + f*x)*1i - 1)*3i + 3*A*tan(e + f*x)^2 + B*tan(e + f*x)^2*15i + B*tan(e + f*x)^2*log(-tan(e + f*x)*1i - 1)*9i - 3*B*tan(e + f*x)^3*log(-tan(e + f*x)*1i - 1) + 9*B*tan(e + f*x)*log(-tan(e + f*x)*1i - 1))*1i)/(3*a^3*f*(tan(e + f*x)*1i + 1)^3)`

3.730. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^3}{(a+ia \tan(e+fx))^3} dx$

3.731
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^2}{(a+ia \tan(e+fx))^3} dx$$

3.731.1 Optimal result	6724
3.731.2 Mathematica [A] (verified)	6724
3.731.3 Rubi [A] (verified)	6725
3.731.4 Maple [A] (verified)	6726
3.731.5 Fricas [A] (verification not implemented)	6727
3.731.6 Sympy [B] (verification not implemented)	6727
3.731.7 Maxima [F(-2)]	6728
3.731.8 Giac [B] (verification not implemented)	6728
3.731.9 Mupad [B] (verification not implemented)	6729

3.731.1 Optimal result

Integrand size = 41, antiderivative size = 99

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{2(A + iB)c^2}{3a^3 f(i - \tan(e + fx))^3} + \frac{(iA - 3B)c^2}{2a^3 f(i - \tan(e + fx))^2} - \frac{iBc^2}{a^3 f(i - \tan(e + fx))}$$

output $2/3*(A+i*B)*c^2/a^3/f/(I-\tan(f*x+e))^3+1/2*(I*A-3*B)*c^2/a^3/f/(I-\tan(f*x+e))^2-I*B*c^2/a^3/f/(I-\tan(f*x+e))$

3.731.2 Mathematica [A] (verified)

Time = 5.33 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.63

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{c^2(-A - iB + 3(iA + B) \tan(e + fx) + 6iB \tan^2(e + fx))}{6a^3 f(-i + \tan(e + fx))^3}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^2)/(a + I*a*Tan[e + f*x])^3,x]`

output $(c^2*(-A - I*B + 3*(I*A + B)*Tan[e + f*x] + (6*I)*B*Tan[e + f*x]^2))/(6*a^3*f*(-I + Tan[e + f*x])^3)$

3.731.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.83, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - ic \tan(e + fx))^2 (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c - ic \tan(e + fx))^2 (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{c(1-i \tan(e+fx))(A+B \tan(e+fx))}{a^4(i \tan(e+fx)+1)^4} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{c^2 \int \frac{(1-i \tan(e+fx))(A+B \tan(e+fx))}{(i \tan(e+fx)+1)^4} d \tan(e + fx)}{a^3 f} \\
 & \quad \downarrow \text{86} \\
 & \frac{c^2 \int \left(\frac{2(A+iB)}{(\tan(e+fx)-i)^4} - \frac{iB}{(\tan(e+fx)-i)^2} + \frac{3B-iA}{(\tan(e+fx)-i)^3} \right) d \tan(e + fx)}{a^3 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{c^2 \left(\frac{-3B+iA}{2(-\tan(e+fx)+i)^2} + \frac{2(A+iB)}{3(-\tan(e+fx)+i)^3} - \frac{iB}{-\tan(e+fx)+i} \right)}{a^3 f}
 \end{aligned}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^2)/(a + I*a*Tan[e + f*x])^3,x]`

output `(c^2*((2*(A + I*B))/(3*(I - Tan[e + f*x])^3) + (I*A - 3*B)/(2*(I - Tan[e + f*x])^2) - (I*B)/(I - Tan[e + f*x]))/(a^3*f)`

3.731.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.731.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.70

method	result
derivativedivides	$\frac{c^2 \left(\frac{iB}{-i+\tan(fx+e)} - \frac{-iA+3B}{2(-i+\tan(fx+e))^2} - \frac{2iB+2A}{3(-i+\tan(fx+e))^3} \right)}{f a^3}$
default	$\frac{c^2 \left(\frac{iB}{-i+\tan(fx+e)} - \frac{-iA+3B}{2(-i+\tan(fx+e))^2} - \frac{2iB+2A}{3(-i+\tan(fx+e))^3} \right)}{f a^3}$
risch	$\frac{c^2 e^{-4i(fx+e)} B}{8a^3 f} + \frac{ic^2 e^{-4i(fx+e)} A}{8a^3 f} - \frac{c^2 e^{-6i(fx+e)} B}{12a^3 f} + \frac{ic^2 e^{-6i(fx+e)} A}{12a^3 f}$
norman	$\frac{-\frac{2iA c^2 \tan(fx+e)^2}{af} + \frac{c^2 A \tan(fx+e)}{af} + \frac{ic^2 B \tan(fx+e)^5}{af} - \frac{-iA c^2 + c^2 B}{6af} - \frac{5(ic^2 B + c^2 A) \tan(fx+e)^3}{3af} - \frac{(-iA c^2 + 5c^2 B) \tan(fx+e)}{2af}}{a^2 (1+\tan(fx+e))^3}$

3.731.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^2}{(a+ia \tan(e+fx))^3} dx$$

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2/(a+I*a*tan(f*x+e))^3,x,method=_RETURVERBOSE)`

output $\frac{1}{f*c^2/a^3*(I*B/(-I+\tan(f*x+e))-1/2*(-I*A+3*B)/(-I+\tan(f*x+e))^2-1/3*(2*A+2*I*B)/(-I+\tan(f*x+e))^3)}$

3.731.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.49

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^2}{(a + i a \tan(e + fx))^3} dx$$

$$= -\frac{(3(-iA - B)c^2 e^{(2i fx + 2ie)} + 2(-iA + B)c^2) e^{(-6i fx - 6ie)}}{24 a^3 f}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output $-\frac{1}{24} \frac{(3(-iA - B)c^2 e^{(2i fx + 2ie)} + 2(-iA + B)c^2) e^{(-6i fx - 6ie)}}{a^3 f}$

3.731.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 172 vs. $2(78) = 156$.

Time = 0.28 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.74

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^2}{(a + i a \tan(e + fx))^3} dx$$

$$= \begin{cases} \frac{((8iAa^3c^2fe^{4ie} - 8Ba^3c^2fe^{4ie})e^{-6ifx} + (12iAa^3c^2fe^{6ie} + 12Ba^3c^2fe^{6ie})e^{-4ifx})e^{-10ie}}{96a^6f^2} & \text{for } a^6 f^2 e^{10ie} \neq 0 \\ \frac{x(Ac^2e^{2ie} + Ac^2 - iBc^2e^{2ie} + iBc^2)e^{-6ie}}{2a^3} & \text{otherwise} \end{cases}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**2/(a+I*a*tan(f*x+e))**3,x)`

output `Piecewise((((8*I*A*a**3*c**2*f*exp(4*I*e) - 8*B*a**3*c**2*f*exp(4*I*e))*exp(-6*I*f*x) + (12*I*A*a**3*c**2*f*exp(6*I*e) + 12*B*a**3*c**2*f*exp(6*I*e))*exp(-4*I*f*x))*exp(-10*I*e)/(96*a**6*f**2), Ne(a**6*f**2*exp(10*I*e), 0)), (x*(A*c**2*exp(2*I*e) + A*c**2 - I*B*c**2*exp(2*I*e) + I*B*c**2)*exp(-6*I*e)/(2*a**3), True))`

3.731.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{(a + ia \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.731.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 156 vs. $2(77) = 154$.

Time = 0.73 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.58

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{(a + ia \tan(e + fx))^3} dx = \frac{2 \left(3 A c^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^5 - 3 i A c^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^4 - 3 B c^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^4 - 8 A c^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^3 + 3 i A c^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^2 + 3 B c^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^2 + 3 A c^2 \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) \right)}{3 a^3 f (\tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) - i)^6}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^2/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")`

output `-2/3*(3*A*c^2*tan(1/2*f*x + 1/2*e)^5 - 3*I*A*c^2*tan(1/2*f*x + 1/2*e)^4 - 3*B*c^2*tan(1/2*f*x + 1/2*e)^4 - 8*A*c^2*tan(1/2*f*x + 1/2*e)^3 - 2*I*B*c^2*tan(1/2*f*x + 1/2*e)^3 + 3*I*A*c^2*tan(1/2*f*x + 1/2*e)^2 + 3*B*c^2*tan(1/2*f*x + 1/2*e)^2 + 3*A*c^2*tan(1/2*f*x + 1/2*e))/(a^3*f*(tan(1/2*f*x + 1/2*e) - I)^6)`

3.731. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^2}{(a+ia \tan(e+fx))^3} dx$

3.731.9 Mupad [B] (verification not implemented)

Time = 8.52 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.88

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^2}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{\frac{c^2(-B+Ai)}{6} + \frac{c^2 \tan(e+fx)(3A-B3i)}{6} + Bc^2 \tan(e+fx)^2}{a^3 f (-\tan(e+fx)^3 1i - 3 \tan(e+fx)^2 + \tan(e+fx) 3i + 1)}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^2)/(a + a*tan(e + f*x)*1i)^3,x)`

output `((c^2*(A*1i - B))/6 + (c^2*tan(e + f*x)*(3*A - B*3i))/6 + B*c^2*tan(e + f*x)^2)/(a^3*f*(tan(e + f*x)*3i - 3*tan(e + f*x)^2 - tan(e + f*x)^3*1i + 1))`

3.732 $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))}{(a+ia \tan(e+fx))^3} dx$

3.732.1 Optimal result 6730
 3.732.2 Mathematica [A] (verified) 6730
 3.732.3 Rubi [A] (verified) 6731
 3.732.4 Maple [A] (verified) 6732
 3.732.5 Fricas [A] (verification not implemented) 6733
 3.732.6 Sympy [B] (verification not implemented) 6733
 3.732.7 Maxima [F(-2)] 6734
 3.732.8 Giac [B] (verification not implemented) 6734
 3.732.9 Mupad [B] (verification not implemented) 6735

3.732.1 Optimal result

Integrand size = 39, antiderivative size = 59

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{(A + iB)c}{3a^3 f(i - \tan(e + fx))^3} - \frac{Bc}{2a^3 f(i - \tan(e + fx))^2}$$

output `1/3*(A+i*B)*c/a^3/f/(I-tan(f*x+e))^3-1/2*B*c/a^3/f/(I-tan(f*x+e))^2`

3.732.2 Mathematica [A] (verified)

Time = 5.14 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.69

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx = \frac{c(-2A + iB - 3B \tan(e + fx))}{6a^3 f(-i + \tan(e + fx))^3}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x]))/(a + I*a*Tan[e + f*x])^3,x]`

output `(c*(-2*A + I*B - 3*B*Tan[e + f*x]))/(6*a^3*f*(-I + Tan[e + f*x])^3)`

3.732.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.90, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.128$, Rules used = {3042, 4071, 27, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - ic \tan(e + fx))(A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c - ic \tan(e + fx))(A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A+B \tan(e+fx)}{a^4(i \tan(e+fx)+1)^4} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{c \int \frac{A+B \tan(e+fx)}{(i \tan(e+fx)+1)^4} d \tan(e + fx)}{a^3 f} \\
 & \quad \downarrow \text{53} \\
 & \frac{c \int \left(\frac{A+iB}{(\tan(e+fx)-i)^4} + \frac{B}{(\tan(e+fx)-i)^3} \right) d \tan(e + fx)}{a^3 f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{c \left(\frac{A+iB}{3(-\tan(e+fx)+i)^3} - \frac{B}{2(-\tan(e+fx)+i)^2} \right)}{a^3 f}
 \end{aligned}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x]))/(a + I*a*Tan[e + f*x])^3, x]`

output `(c*((A + I*B)/(3*(I - Tan[e + f*x])^3) - B/(2*(I - Tan[e + f*x])^2)))/(a^3*f)`

3.732.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.732.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.73

method	result	size
derivativedivides	$\frac{c\left(-\frac{iB+A}{3(-i+\tan(fx+e))^3}-\frac{B}{2(-i+\tan(fx+e))^2}\right)}{fa^3}$	43
default	$\frac{c\left(-\frac{iB+A}{3(-i+\tan(fx+e))^3}-\frac{B}{2(-i+\tan(fx+e))^2}\right)}{fa^3}$	43
risch	$\frac{ce^{-2i(fx+e)}B}{8a^3f} + \frac{ice^{-2i(fx+e)}A}{8a^3f} + \frac{icAe^{-4i(fx+e)}}{8a^3f} - \frac{ce^{-6i(fx+e)}B}{24a^3f} + \frac{ice^{-6i(fx+e)}A}{24a^3f}$	100

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

$$3.732. \quad \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))}{(a+ia \tan(e+fx))^3} dx$$

output $1/f*c/a^3*(-1/3*(A+I*B)/(-I+\tan(f*x+e))^3-1/2*B/(-I+\tan(f*x+e))^2)$

3.732.5 Fricas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.02

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))}{(a + i a \tan(e + fx))^3} dx$$

$$= -\frac{(3(-iA - B)ce^{4ifx+4ie} - 3iAce^{2ifx+2ie} - (iA - B)c)e^{-6ifx-6ie}}{24a^3f}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output $-1/24*(3*(-I*A - B)*c*e^{(4*I*f*x + 4*I*e)} - 3*I*A*c*e^{(2*I*f*x + 2*I*e)} - (I*A - B)*c)*e^{(-6*I*f*x - 6*I*e)}/(a^3*f)$

3.732.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 206 vs. $2(44) = 88$.

Time = 0.27 (sec) , antiderivative size = 206, normalized size of antiderivative = 3.49

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))}{(a + i a \tan(e + fx))^3} dx$$

$$= \begin{cases} \frac{(192iAa^6cf^2e^{8ie}e^{-4ifx} + (64iAa^6cf^2e^{6ie} - 64Ba^6cf^2e^{6ie})e^{-6ifx} + (192iAa^6cf^2e^{10ie} + 192Ba^6cf^2e^{10ie})e^{-2ifx})e^{-12ie}}{1536a^9f^3} & \text{for } a^9f^3e^{12ie} \neq 0 \\ \frac{x(Ace^{4ie} + 2Ace^{2ie} + Ac - iBce^{4ie} + iBc)e^{-6ie}}{4a^3} & \text{otherwise} \end{cases}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e))**3,x)`

output `Piecewise((((192*I*A*a**6*c*f**2*exp(8*I*e)*exp(-4*I*f*x) + (64*I*A*a**6*c*f**2*exp(6*I*e) - 64*B*a**6*c*f**2*exp(6*I*e))*exp(-6*I*f*x) + (192*I*A*a**6*c*f**2*exp(10*I*e) + 192*B*a**6*c*f**2*exp(10*I*e))*exp(-2*I*f*x))*exp(-12*I*e)/(1536*a**9*f**3), Ne(a**9*f**3*exp(12*I*e), 0)), (x*(A*c*exp(4*I*e) + 2*A*c*exp(2*I*e) + A*c - I*B*c*exp(4*I*e) + I*B*c)*exp(-6*I*e)/(4*a**3), True))`

3.732. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))}{(a+ia \tan(e+fx))^3} dx$

3.732.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))}{(a + i a \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e))^3,x, algorith="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.732.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 140 vs. $2(45) = 90$.

Time = 0.64 (sec) , antiderivative size = 140, normalized size of antiderivative = 2.37

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))}{(a + i a \tan(e + fx))^3} dx = \frac{2 \left(3 A c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^5 - 6 i A c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^4 - 3 B c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^4 - 10 A c \tan \left(\frac{1}{2} f x + \frac{1}{2} e \right)^3 \right)}{3 a^3 f \left(\tan \left(\frac{1}{2} f x + \frac{1}{2} e \right) - i \right)^6}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))/(a+I*a*tan(f*x+e))^3,x, algorith="giac")`

output `-2/3*(3*A*c*tan(1/2*f*x + 1/2*e)^5 - 6*I*A*c*tan(1/2*f*x + 1/2*e)^4 - 3*B*c*tan(1/2*f*x + 1/2*e)^4 - 10*A*c*tan(1/2*f*x + 1/2*e)^3 + 2*I*B*c*tan(1/2*f*x + 1/2*e)^3 + 6*I*A*c*tan(1/2*f*x + 1/2*e)^2 + 3*B*c*tan(1/2*f*x + 1/2*e)^2 + 3*A*c*tan(1/2*f*x + 1/2*e))/(a^3*f*(tan(1/2*f*x + 1/2*e) - I)^6)`

3.732.9 Mupad [B] (verification not implemented)

Time = 8.47 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.05

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{\frac{c(B+A2i)}{6} + \frac{Bc \tan(e+fx) 1i}{2}}{a^3 f (-\tan(e + fx)^3 1i - 3 \tan(e + fx)^2 + \tan(e + fx) 3i + 1)}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i))/(a + a*tan(e + f*x)*1i)^3,x)`

output `((c*(A*2i + B))/6 + (B*c*tan(e + f*x)*1i)/2)/(a^3*f*(tan(e + f*x)*3i - 3*tan(e + f*x)^2 - tan(e + f*x)^3*1i + 1))`

3.733 $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3} dx$

3.733.1 Optimal result	6736
3.733.2 Mathematica [A] (verified)	6736
3.733.3 Rubi [A] (verified)	6737
3.733.4 Maple [A] (verified)	6738
3.733.5 Fricas [A] (verification not implemented)	6739
3.733.6 Sympy [A] (verification not implemented)	6739
3.733.7 Maxima [F(-2)]	6740
3.733.8 Giac [A] (verification not implemented)	6740
3.733.9 Mupad [B] (verification not implemented)	6741

3.733.1 Optimal result

Integrand size = 26, antiderivative size = 112

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3} dx = \frac{(A - iB)x}{8a^3} + \frac{iA - B}{6f(a + ia \tan(e + fx))^3} + \frac{iA + B}{8af(a + ia \tan(e + fx))^2} + \frac{iA + B}{8f(a^3 + ia^3 \tan(e + fx))}$$

```
output 1/8*(A-I*B)*x/a^3+1/6*(I*A-B)/f/(a+I*a*tan(f*x+e))^3+1/8*(I*A+B)/a/f/(a+I*a*tan(f*x+e))^2+1/8*(I*A+B)/f/(a^3+I*a^3*tan(f*x+e))
```

3.733.2 Mathematica [A] (verified)

Time = 1.01 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.97

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3} dx = \frac{-10A + 2iB + 3(iA + B) \arctan(\tan(e + fx)) \sec^3(e + fx) (\cos(3(e + fx)) + i \sin(3(e + fx))) + (-9iA - 9iB) \tan(e + fx)}{24a^3 f (-i + \tan(e + fx))^3}$$

```
input Integrate[(A + B*Tan[e + f*x])/(a + I*a*Tan[e + f*x])^3,x]
```

```
output (-10*A + (2*I)*B + 3*(I*A + B)*ArcTan[Tan[e + f*x]]*Sec[e + f*x]^3*(Cos[3*(e + f*x)] + I*Sin[3*(e + f*x)]) + ((-9*I)*A - 9*B)*Tan[e + f*x] + 3*(A - I*B)*Tan[e + f*x]^2)/(24*a^3*f*(-I + Tan[e + f*x])^3)
```

3.733.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.99, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.269$, Rules used = {3042, 4009, 3042, 3960, 3042, 3960, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{4009} \\
 & \frac{(A - iB) \int \frac{1}{(i \tan(e + fx)a + a)^2} dx}{2a} + \frac{-B + iA}{6f(a + ia \tan(e + fx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A - iB) \int \frac{1}{(i \tan(e + fx)a + a)^2} dx}{2a} + \frac{-B + iA}{6f(a + ia \tan(e + fx))^3} \\
 & \quad \downarrow \text{3960} \\
 & \frac{(A - iB) \left(\frac{\int \frac{1}{i \tan(e + fx)a + a} dx}{2a} + \frac{i}{4f(a + ia \tan(e + fx))^2} \right)}{2a} + \frac{-B + iA}{6f(a + ia \tan(e + fx))^3} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(A - iB) \left(\frac{\int \frac{1}{i \tan(e + fx)a + a} dx}{2a} + \frac{i}{4f(a + ia \tan(e + fx))^2} \right)}{2a} + \frac{-B + iA}{6f(a + ia \tan(e + fx))^3} \\
 & \quad \downarrow \text{3960} \\
 & \frac{(A - iB) \left(\frac{\int \frac{1 dx}{2a} + \frac{i}{2f(a + ia \tan(e + fx))}}{2a} + \frac{i}{4f(a + ia \tan(e + fx))^2} \right)}{2a} + \frac{-B + iA}{6f(a + ia \tan(e + fx))^3} \\
 & \quad \downarrow \text{24} \\
 & \frac{-B + iA}{6f(a + ia \tan(e + fx))^3} + \frac{(A - iB) \left(\frac{\frac{x}{2a} + \frac{i}{2f(a + ia \tan(e + fx))}}{2a} + \frac{i}{4f(a + ia \tan(e + fx))^2} \right)}{2a}
 \end{aligned}$$

input `Int[(A + B*Tan[e + f*x])/(a + I*a*Tan[e + f*x])^3,x]`

output `(I*A - B)/(6*f*(a + I*a*Tan[e + f*x])^3) + ((A - I*B)*((I/4)/(f*(a + I*a*Tan[e + f*x])^2) + (x/(2*a) + (I/2)/(f*(a + I*a*Tan[e + f*x]))/(2*a)))/(2*a)`

3.733.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3960 `Int[((a_) + (b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Simp[1/(2*a) Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

rule 4009 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] := Simp[(-b*c - a*d)*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

3.733.4 Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.14

method	result
risch	$-\frac{ixB}{8a^3} + \frac{xA}{8a^3} + \frac{e^{-2i(fx+e)}B}{16a^3f} + \frac{3ie^{-2i(fx+e)}A}{16a^3f} - \frac{e^{-4i(fx+e)}B}{32a^3f} + \frac{3ie^{-4i(fx+e)}A}{32a^3f} - \frac{e^{-6i(fx+e)}B}{48a^3f} + \frac{ie^{-6i(fx+e)}A}{48a^3f}$
derivativedivides	$-\frac{iB \arctan(\tan(fx+e))}{8fa^3} + \frac{A \arctan(\tan(fx+e))}{8fa^3} - \frac{iA}{8fa^3(-i+\tan(fx+e))^2} - \frac{B}{8fa^3(-i+\tan(fx+e))^2} + \frac{ie^{-6i(fx+e)}A}{8fa^3}$
default	$-\frac{iB \arctan(\tan(fx+e))}{8fa^3} + \frac{A \arctan(\tan(fx+e))}{8fa^3} - \frac{iA}{8fa^3(-i+\tan(fx+e))^2} - \frac{B}{8fa^3(-i+\tan(fx+e))^2} + \frac{ie^{-6i(fx+e)}A}{8fa^3}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

3.733. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3} dx$

output
$$-1/8*I*x/a^3*B+1/8*x/a^3*A+1/16/a^3/f*\exp(-2*I*(f*x+e))*B+3/16*I/a^3/f*\exp(-2*I*(f*x+e))*A-1/32/a^3/f*\exp(-4*I*(f*x+e))*B+3/32*I/a^3/f*\exp(-4*I*(f*x+e))*A-1/48/a^3/f*\exp(-6*I*(f*x+e))*B+1/48*I/a^3/f*\exp(-6*I*(f*x+e))*A$$

3.733.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.68

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{(12(A - iB)fx e^{(6i fx + 6ie)} - 6(-3iA - B)e^{(4i fx + 4ie)} - 3(-3iA + B)e^{(2i fx + 2ie)} + 2iA - 2B)e^{(-6i fx - 6ie)}}{96a^3 f}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3,x, algorithm="fracas")`

output
$$1/96*(12*(A - I*B)*f*x*e^{(6*I*f*x + 6*I*e)} - 6*(-3*I*A - B)*e^{(4*I*f*x + 4*I*e)} - 3*(-3*I*A + B)*e^{(2*I*f*x + 2*I*e)} + 2*I*A - 2*B)*e^{(-6*I*f*x - 6*I*e)}/(a^3*f)$$

3.733.6 Sympy [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 258, normalized size of antiderivative = 2.30

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3} dx$$

$$= \left\{ \frac{((512iAa^6 f^2 e^{6ie} - 512Ba^6 f^2 e^{6ie})e^{-6ifx} + (2304iAa^6 f^2 e^{8ie} - 768Ba^6 f^2 e^{8ie})e^{-4ifx} + (4608iAa^6 f^2 e^{10ie} + 1536Ba^6 f^2 e^{10ie})e^{-2ifx})e^{-12ie}}{24576a^9 f^3} \right.$$

$$\left. x \left(-\frac{A-iB}{8a^3} + \frac{(Ae^{6ie} + 3Ae^{4ie} + 3Ae^{2ie} + A - iBe^{6ie} - iBe^{4ie} + iBe^{2ie} + iB)e^{-6ie}}{8a^3} \right) \right.$$

$$\left. + \frac{x(A - iB)}{8a^3} \right.$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**3,x)`

```
output Piecewise((((512*I*A*a**6*f**2*exp(6*I*e) - 512*B*a**6*f**2*exp(6*I*e))*exp(-6*I*f*x) + (2304*I*A*a**6*f**2*exp(8*I*e) - 768*B*a**6*f**2*exp(8*I*e))*exp(-4*I*f*x) + (4608*I*A*a**6*f**2*exp(10*I*e) + 1536*B*a**6*f**2*exp(10*I*e))*exp(-2*I*f*x))*exp(-12*I*e)/(24576*a**9*f**3), Ne(a**9*f**3*exp(12*I*e), 0)), (x*(-(A - I*B)/(8*a**3) + (A*exp(6*I*e) + 3*A*exp(4*I*e) + 3*A*exp(2*I*e) + A - I*B*exp(6*I*e) - I*B*exp(4*I*e) + I*B*exp(2*I*e) + I*B)*exp(-6*I*e)/(8*a**3)), True)) + x*(A - I*B)/(8*a**3)
```

3.733.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.733.8 Giac [A] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.15

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3} dx = \frac{\frac{6(-iA-B)\log(\tan(fx+e)+i)}{a^3} + \frac{6(iA+B)\log(\tan(fx+e)-i)}{a^3} + \frac{-11iA\tan(fx+e)^3 - 11B\tan(fx+e)^3 - 45A\tan(fx+e)^2 + 45iB\tan(fx+e)}{a^3(\tan(fx+e)-i)^5}}{96f}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")
```

```
output -1/96*(6*(-I*A - B)*log(tan(f*x + e) + I)/a^3 + 6*(I*A + B)*log(tan(f*x + e) - I)/a^3 + (-11*I*A*tan(f*x + e)^3 - 11*B*tan(f*x + e)^3 - 45*A*tan(f*x + e)^2 + 45*I*B*tan(f*x + e)^2 + 69*I*A*tan(f*x + e) + 69*B*tan(f*x + e) + 51*A - 19*I*B)/(a^3*(tan(f*x + e) - I)^3))/f
```

3.733.9 Mupad [B] (verification not implemented)

Time = 8.56 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.99

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3} dx$$

$$= -\frac{\tan(e + fx)^2 \left(\frac{B}{8a^3} + \frac{A1i}{8a^3}\right) - \frac{A5i}{12a^3} - \frac{B}{12a^3} + \tan(e + fx) \left(\frac{3A}{8a^3} - \frac{B3i}{8a^3}\right)}{f \left(-\tan(e + fx)^3 1i - 3 \tan(e + fx)^2 + \tan(e + fx) 3i + 1\right)} - \frac{x(B + A 1i) 1i}{8a^3}$$

input `int((A + B*tan(e + f*x))/(a + a*tan(e + f*x)*1i)^3,x)`output `- (tan(e + f*x)^2*((A*1i)/(8*a^3) + B/(8*a^3)) - (A*5i)/(12*a^3) - B/(12*a^3) + tan(e + f*x)*((3*A)/(8*a^3) - (B*3i)/(8*a^3)))/(f*(tan(e + f*x)*3i - 3*tan(e + f*x)^2 - tan(e + f*x)^3*1i + 1)) - (x*(A*1i + B)*1i)/(8*a^3)`

3.734 $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))} dx$

3.734.1 Optimal result 6742
 3.734.2 Mathematica [A] (verified) 6742
 3.734.3 Rubi [A] (verified) 6743
 3.734.4 Maple [A] (verified) 6745
 3.734.5 Fricas [A] (verification not implemented) 6745
 3.734.6 Sympy [A] (verification not implemented) 6746
 3.734.7 Maxima [F(-2)] 6746
 3.734.8 Giac [A] (verification not implemented) 6747
 3.734.9 Mupad [B] (verification not implemented) 6747

3.734.1 Optimal result

Integrand size = 41, antiderivative size = 153

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3(c - ic \tan(e + fx))} dx$$

$$= \frac{(2A - iB)x}{8a^3c} + \frac{A + iB}{12a^3cf(i - \tan(e + fx))^3} - \frac{iA}{8a^3cf(i - \tan(e + fx))^2}$$

$$- \frac{3A - iB}{16a^3cf(i - \tan(e + fx))} + \frac{A - iB}{16a^3cf(i + \tan(e + fx))}$$

output `1/8*(2*A-I*B)*x/a^3/c+1/12*(A+I*B)/a^3/c/f/(I-tan(f*x+e))^3-1/8*I*A/a^3/c/f/(I-tan(f*x+e))^2+1/16*(-3*A+I*B)/a^3/c/f/(I-tan(f*x+e))+1/16*(A-I*B)/a^3/c/f/(I+tan(f*x+e))`

3.734.2 Mathematica [A] (verified)

Time = 5.83 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.97

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3(c - ic \tan(e + fx))} dx$$

$$= \frac{\sec^3(e + fx)(-9iA \cos(e + fx) + (-1 + 2 \cos(2(e + fx))))((iA + 2B) \cos(e + fx) + (-2A + iB) \sin(e + fx))}{24a^3cf(-i + \tan(e + fx))^3(i + \tan(e + fx))}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])),x]`

3.734. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))} dx$

output $(\text{Sec}[e + f*x]^3*((-9*I)*A*\text{Cos}[e + f*x] + (-1 + 2*\text{Cos}[2*(e + f*x)])*((I*A + 2*B)*\text{Cos}[e + f*x] + (-2*A + I*B)*\text{Sin}[e + f*x]) - 3*(2*A - I*B)*\text{ArcTan}[\text{Tan}[e + f*x]]*\text{Sec}[e + f*x]*(\text{Cos}[2*(e + f*x)] + I*\text{Sin}[2*(e + f*x)])))/(24*a^3*c*f*(-I + \text{Tan}[e + f*x])^3*(I + \text{Tan}[e + f*x]))$

3.734.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.83, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))} dx$$

↓ 4071

$$\frac{ac \int \frac{A + B \tan(e + fx)}{a^4 c^2 (1 - i \tan(e + fx))^2 (i \tan(e + fx) + 1)^4} d \tan(e + fx)}{f}$$

↓ 27

$$\int \frac{A + B \tan(e + fx)}{(1 - i \tan(e + fx))^2 (i \tan(e + fx) + 1)^4} d \tan(e + fx)$$

$a^3 c f$

↓ 86

$$\int \left(\frac{iA}{4(\tan(e + fx) - i)^3} + \frac{2A - iB}{8(\tan^2(e + fx) + 1)} + \frac{iB - 3A}{16(\tan(e + fx) - i)^2} + \frac{iB - A}{16(\tan(e + fx) + i)^2} + \frac{A + iB}{4(\tan(e + fx) - i)^4} \right) d \tan(e + fx)$$

$a^3 c f$

↓ 2009

$$\frac{\frac{1}{8}(2A - iB) \arctan(\tan(e + fx)) - \frac{3A - iB}{16(-\tan(e + fx) + i)} + \frac{A - iB}{16(\tan(e + fx) + i)} + \frac{A + iB}{12(-\tan(e + fx) + i)^3} - \frac{iA}{8(-\tan(e + fx) + i)^2}}{a^3 c f}$$

3.734. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))} dx$

```
input Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x]))
,x]
```

```
output (((2*A - I*B)*ArcTan[Tan[e + f*x]])/8 + (A + I*B)/(12*(I - Tan[e + f*x])^3
) - ((I/8)*A)/(I - Tan[e + f*x])^2 - (3*A - I*B)/(16*(I - Tan[e + f*x])) +
(A - I*B)/(16*(I + Tan[e + f*x])))/(a^3*c*f)
```

3.734.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.734.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.27

method	result
risch	$-\frac{ixB}{8a^3c} + \frac{xA}{4a^3c} - \frac{e^{-4i(fx+e)}B}{32a^3cf} + \frac{ie^{-4i(fx+e)}A}{16a^3cf} - \frac{e^{-6i(fx+e)}B}{96a^3cf} + \frac{ie^{-6i(fx+e)}A}{96a^3cf} - \frac{\cos(2fx+2e)B}{32a^3cf} + \frac{5i \cos(2fx+2e)A}{32a^3cf}$
derivativedivides	$-\frac{iA}{8fa^3c(-i+\tan(fx+e))^2} + \frac{3A}{16fa^3c(-i+\tan(fx+e))} - \frac{iB}{16fa^3c(-i+\tan(fx+e))} - \frac{A}{12fa^3c(-i+\tan(fx+e))^3}$
default	$-\frac{iA}{8fa^3c(-i+\tan(fx+e))^2} + \frac{3A}{16fa^3c(-i+\tan(fx+e))} - \frac{iB}{16fa^3c(-i+\tan(fx+e))} - \frac{A}{12fa^3c(-i+\tan(fx+e))^3}$
norman	$\frac{(-iB+2A)x - \frac{-4iA+B}{12acf} + \frac{B \tan(fx+e)^2}{4acf} + \frac{(-iB+2A) \tan(fx+e)^3}{3acf} + \frac{(-iB+2A) \tan(fx+e)^5}{8acf} + \frac{3(-iB+2A)x \tan(fx+e)^2}{8ac} + \frac{3(-iB+2A) \tan(fx+e)^4}{8ac}}{(1+\tan(fx+e)^2)^3 a^2}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output
$$-1/8*I*x/a^3/c*B+1/4*x/a^3/c*A-1/32/a^3/c/f*\exp(-4*I*(f*x+e))*B+1/16*I/a^3/c/f*\exp(-4*I*(f*x+e))*A-1/96/a^3/c/f*\exp(-6*I*(f*x+e))*B+1/96*I/a^3/c/f*\exp(-6*I*(f*x+e))*A-1/32/a^3/c/f*\cos(2*f*x+2*e)*B+5/32*I/a^3/c/f*\cos(2*f*x+2*e)*A-1/32*I/a^3/c/f*\sin(2*f*x+2*e)*B+7/32/a^3/c/f*\sin(2*f*x+2*e)*A$$

3.734.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.59

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))} dx$$

$$= \frac{(12(2A - iB)fx e^{6i fx + 6ie} - 3(iA + B)e^{8i fx + 8ie} + 18iAe^{4i fx + 4ie} - 3(-2iA + B)e^{2i fx + 2ie} + iA - B)}{96a^3cf}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e)),x, algorithm="fracas")`

output
$$1/96*(12*(2*A - I*B)*f*x*e^{(6*I*f*x + 6*I*e)} - 3*(I*A + B)*e^{(8*I*f*x + 8*I*e)} + 18*I*A*e^{(4*I*f*x + 4*I*e)} - 3*(-2*I*A + B)*e^{(2*I*f*x + 2*I*e)} + I*A - B)*e^{(-6*I*f*x - 6*I*e)}/(a^3*c*f)$$

3.734.
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))} dx$$

3.734.6 Sympy [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 340, normalized size of antiderivative = 2.22

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ict \tan(e + fx))} dx$$

$$= \begin{cases} \frac{(294912iAa^9c^3f^3e^{10ie}e^{-2ifx} + (16384iAa^9c^3f^3e^{6ie} - 16384Ba^9c^3f^3e^{6ie})e^{-6ifx} + (98304iAa^9c^3f^3e^{8ie} - 49152Ba^9c^3f^3e^{8ie})e^{-4ifx} + (-49152Ba^9c^3f^3e^{8ie} + 16384Ba^9c^3f^3e^{6ie})e^{-2ifx} + (-49152Ba^9c^3f^3e^{8ie} - 16384Ba^9c^3f^3e^{6ie}))e^{-6ifx}}{1572864a^{12}c^4f^4} \\ x \left(-\frac{2A - iB}{8a^3c} + \frac{(Ae^{8ie} + 4Ae^{6ie} + 6Ae^{4ie} + 4Ae^{2ie} + A - iBe^{8ie} - 2iBe^{6ie} + 2iBe^{2ie} + iB)e^{-6ie}}{16a^3c} \right) \\ + \frac{x(2A - iB)}{8a^3c} \end{cases}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**3/(c-I*c*tan(f*x+e)),x)`

output `Piecewise(((294912*I*A*a**9*c**3*f**3*exp(10*I*e)*exp(-2*I*f*x) + (16384*I*A*a**9*c**3*f**3*exp(6*I*e) - 16384*B*a**9*c**3*f**3*exp(6*I*e))*exp(-6*I*f*x) + (98304*I*A*a**9*c**3*f**3*exp(8*I*e) - 49152*B*a**9*c**3*f**3*exp(8*I*e))*exp(-4*I*f*x) + (-49152*I*A*a**9*c**3*f**3*exp(14*I*e) - 49152*B*a**9*c**3*f**3*exp(14*I*e))*exp(2*I*f*x))*exp(-12*I*e)/(1572864*a**12*c**4*f**4), Ne(a**12*c**4*f**4*exp(12*I*e), 0)), (x*(-(2*A - I*B)/(8*a**3*c) + (A*exp(8*I*e) + 4*A*exp(6*I*e) + 6*A*exp(4*I*e) + 4*A*exp(2*I*e) + A - I*B*exp(8*I*e) - 2*I*B*exp(6*I*e) + 2*I*B*exp(2*I*e) + I*B)*exp(-6*I*e)/(16*a**3*c)), True)) + x*(2*A - I*B)/(8*a**3*c)`

3.734.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ict \tan(e + fx))} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e)),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.734.8 Giac [A] (verification not implemented)

Time = 0.63 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.18

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))} dx =$$

$$\frac{\frac{6(-2iA-B)\log(\tan(fx+e)+i)}{a^3c} + \frac{6(2iA+B)\log(\tan(fx+e)-i)}{a^3c} + \frac{6(2iA\tan(fx+e)+B\tan(fx+e)-3A+2iB)}{a^3c(\tan(fx+e)+i)} + \frac{-22iA\tan(fx+e)^3}{a^3c}}{96f}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e)),x, algo
rithm="giac")
```

```
output -1/96*(6*(-2*I*A - B)*log(tan(f*x + e) + I)/(a^3*c) + 6*(2*I*A + B)*log(ta
n(f*x + e) - I)/(a^3*c) + 6*(2*I*A*tan(f*x + e) + B*tan(f*x + e) - 3*A + 2
*I*B)/(a^3*c*(tan(f*x + e) + I)) + (-22*I*A*tan(f*x + e)^3 - 11*B*tan(f*x
+ e)^3 - 84*A*tan(f*x + e)^2 + 39*I*B*tan(f*x + e)^2 + 114*I*A*tan(f*x + e
) + 45*B*tan(f*x + e) + 60*A - 9*I*B)/(a^3*c*(tan(f*x + e) - I)^3))/f
```

3.734.9 Mupad [B] (verification not implemented)

Time = 8.73 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.05

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))} dx$$

$$= \frac{\frac{A}{3a^3c} + \tan(e + fx)^2 \left(\frac{A}{2a^3c} - \frac{B i i}{4a^3c} \right) + \tan(e + fx)^3 \left(\frac{B}{8a^3c} + \frac{A i i}{4a^3c} \right) - \tan(e + fx) \left(\frac{B}{24a^3c} + \frac{A i i}{12a^3c} \right) + \frac{B i i}{12a^3c}}{f (\tan(e + fx)^4 i i + 2 \tan(e + fx)^3 + 2 \tan(e + fx) - i)}$$

$$- \frac{x (B + A 2i) i i}{8 a^3 c}$$

```
input int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*ii)^3*(c - c*tan(e + f*x)*ii
)),x)
```

```
output (tan(e + f*x)^2*(A/(2*a^3*c) - (B*ii)/(4*a^3*c)) - tan(e + f*x)*((A*ii)/(1
2*a^3*c) + B/(24*a^3*c)) + tan(e + f*x)^3*((A*ii)/(4*a^3*c) + B/(8*a^3*c))
+ A/(3*a^3*c) + (B*ii)/(12*a^3*c))/(f*(2*tan(e + f*x) + 2*tan(e + f*x)^3
+ tan(e + f*x)^4*ii - ii)) - (x*(A*2i + B)*ii)/(8*a^3*c)
```

3.735
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^2} dx$$

3.735.1 Optimal result	6748
3.735.2 Mathematica [A] (verified)	6748
3.735.3 Rubi [A] (verified)	6749
3.735.4 Maple [A] (verified)	6751
3.735.5 Fricas [A] (verification not implemented)	6751
3.735.6 Sympy [A] (verification not implemented)	6752
3.735.7 Maxima [F(-2)]	6752
3.735.8 Giac [A] (verification not implemented)	6753
3.735.9 Mupad [B] (verification not implemented)	6753

3.735.1 Optimal result

Integrand size = 41, antiderivative size = 185

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3(c - ic \tan(e + fx))^2} dx$$

$$= \frac{(5A - iB)x}{16a^3c^2} + \frac{A + iB}{24a^3c^2 f(i - \tan(e + fx))^3} - \frac{3iA - B}{32a^3c^2 f(i - \tan(e + fx))^2}$$

$$- \frac{3A}{16a^3c^2 f(i - \tan(e + fx))} + \frac{iA + B}{32a^3c^2 f(i + \tan(e + fx))^2} + \frac{2A - iB}{16a^3c^2 f(i + \tan(e + fx))}$$

output `1/16*(5*A-I*B)*x/a^3/c^2+1/24*(A+I*B)/a^3/c^2/f/(I-tan(f*x+e))^3+1/32*(-3*I*A+B)/a^3/c^2/f/(I-tan(f*x+e))^2-3/16*A/a^3/c^2/f/(I-tan(f*x+e))+1/32*(I*A+B)/a^3/c^2/f/(I+tan(f*x+e))^2+1/16*(2*A-I*B)/a^3/c^2/f/(I+tan(f*x+e))`

3.735.2 Mathematica [A] (verified)

Time = 6.06 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.93

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3(c - ic \tan(e + fx))^2} dx$$

$$= \frac{\sec^4(e + fx)(47A + 5iB + (-14A + 22iB) \cos(2(e + fx)) - A \cos(4(e + fx)) + 5iB \cos(4(e + fx)) - 40iB \sin(4(e + fx)))}{192a^3c^2}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^2), x]`

3.735.
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^2} dx$$

output $(\text{Sec}[e + f*x]^4*(47*A + (5*I)*B + (-14*A + (22*I)*B)*\text{Cos}[2*(e + f*x)] - A*\text{Cos}[4*(e + f*x)] + (5*I)*B*\text{Cos}[4*(e + f*x)] - (40*I)*A*\text{Sin}[2*(e + f*x)] - 8*B*\text{Sin}[2*(e + f*x)] - (5*I)*A*\text{Sin}[4*(e + f*x)] - B*\text{Sin}[4*(e + f*x)] + 12*(5*A - I*B)*\text{ArcTan}[\text{Tan}[e + f*x]]*(-I + \text{Tan}[e + f*x]))/(192*a^3*c^2*f*(-I + \text{Tan}[e + f*x])^3*(I + \text{Tan}[e + f*x])^2)$

3.735.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.81, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^2} dx$$

↓ 4071

$$\frac{ac \int \frac{A + B \tan(e + fx)}{a^4 c^3 (1 - i \tan(e + fx))^3 (i \tan(e + fx) + 1)^4} d \tan(e + fx)}{f}$$

↓ 27

$$\int \frac{A + B \tan(e + fx)}{(1 - i \tan(e + fx))^3 (i \tan(e + fx) + 1)^4} d \tan(e + fx)}{a^3 c^2 f}$$

↓ 86

$$\int \left(-\frac{3A}{16(\tan(e + fx) - i)^2} + \frac{5A - iB}{16(\tan^2(e + fx) + 1)} + \frac{iB - 2A}{16(\tan(e + fx) + i)^2} + \frac{i(3A + iB)}{16(\tan(e + fx) - i)^3} - \frac{i(A - iB)}{16(\tan(e + fx) + i)^3} + \frac{A + iB}{8(\tan(e + fx) - i)^4} \right) d \tan(e + fx)}{a^3 c^2 f}$$

↓ 2009

$$\frac{\frac{1}{16}(5A - iB) \arctan(\tan(e + fx)) + \frac{2A - iB}{16(\tan(e + fx) + i)} - \frac{-B + 3iA}{32(-\tan(e + fx) + i)^2} + \frac{B + iA}{32(\tan(e + fx) + i)^2} + \frac{A + iB}{24(-\tan(e + fx) + i)^3} - \frac{1}{16}}{a^3 c^2 f}$$

3.735. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^2} dx$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^2),x]`

output `((((5*A - I*B)*ArcTan[Tan[e + f*x]])/16 + (A + I*B)/(24*(I - Tan[e + f*x])^3) - ((3*I)*A - B)/(32*(I - Tan[e + f*x])^2) - (3*A)/(16*(I - Tan[e + f*x])) + (I*A + B)/(32*(I + Tan[e + f*x])^2) + (2*A - I*B)/(16*(I + Tan[e + f*x]))) / (a^3*c^2*f)`

3.735.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.735.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.13

method	result
norman	$\frac{(-iB+5A)x - \frac{-iA+B}{6acf} + \frac{(-iB+5A)\tan(fx+e)^3}{6acf} + \frac{(-iB+5A)\tan(fx+e)^5}{16acf} + \frac{3(-iB+5A)x\tan(fx+e)^2}{16ac} + \frac{3(-iB+5A)x\tan(fx+e)^4}{16ac}}{(1+\tan(fx+e))^2} a^2 c$
risch	$-\frac{iB}{16a^3c^2} + \frac{5xA}{16a^3c^2} - \frac{e^{-6i(fx+e)}B}{192a^3c^2f} + \frac{ie^{-6i(fx+e)}A}{192a^3c^2f} - \frac{\cos(4fx+4e)B}{32a^3c^2f} + \frac{i\cos(4fx+4e)A}{32a^3c^2f} + \frac{i\sin(4fx+4e)B}{64a^3c^2f}$
derivativedivides	$-\frac{3iA}{32fa^3c^2(-i+\tan(fx+e))^2} + \frac{A}{8fa^3c^2(i+\tan(fx+e))} - \frac{iB}{16fa^3c^2(i+\tan(fx+e))} + \frac{5A\arctan(\tan(fx+e))}{16fa^3c^2} - i$
default	$-\frac{3iA}{32fa^3c^2(-i+\tan(fx+e))^2} + \frac{A}{8fa^3c^2(i+\tan(fx+e))} - \frac{iB}{16fa^3c^2(i+\tan(fx+e))} + \frac{5A\arctan(\tan(fx+e))}{16fa^3c^2} - i$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^2,x,method=_R
ETURNVERBOSE)`

output `(1/16*(5*A-I*B)/a/c*x-1/6*(-I*A+B)/a/c/f+1/6*(5*A-I*B)/a/c/f*tan(f*x+e)^3+
1/16*(5*A-I*B)/a/c/f*tan(f*x+e)^5+3/16*(5*A-I*B)/a/c*x*tan(f*x+e)^2+3/16*(
5*A-I*B)/a/c*x*tan(f*x+e)^4+1/16*(5*A-I*B)/a/c*x*tan(f*x+e)^6+1/16*(11*A+I
*B)/a/c/f*tan(f*x+e))/(1+tan(f*x+e))^2)^3/a^2/c`

3.735.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.62

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^2} dx$$

$$= \frac{(24(5A - iB)fx e^{(6i fx + 6ie)} - 3(iA + B)e^{(10i fx + 10ie)} - 6(5iA + 3B)e^{(8i fx + 8ie)} - 12(-5iA + B)e^{(4i fx + 4ie)})}{384a^3c^2f}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^2,x, al
gorithm="fracas")`

output `1/384*(24*(5*A - I*B)*f*x*e^(6*I*f*x + 6*I*e) - 3*(I*A + B)*e^(10*I*f*x +
10*I*e) - 6*(5*I*A + 3*B)*e^(8*I*f*x + 8*I*e) - 12*(-5*I*A + B)*e^(4*I*f*x
+ 4*I*e) - 3*(-5*I*A + 3*B)*e^(2*I*f*x + 2*I*e) + 2*I*A - 2*B)*e^(-6*I*f*x
x - 6*I*e)/(a^3*c^2*f)`

3.735.
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^2} dx$$

3.735.6 Sympy [A] (verification not implemented)

Time = 0.46 (sec) , antiderivative size = 452, normalized size of antiderivative = 2.44

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^2} dx$$

$$= \left\{ \begin{aligned} & \frac{((33554432iAa^{12}c^8f^4e^{6ie} - 33554432Ba^{12}c^8f^4e^{6ie})e^{-6ifx} + (251658240iAa^{12}c^8f^4e^{8ie} - 150994944Ba^{12}c^8f^4e^{8ie})e^{-4ifx} + (1006632960iAa^{12}c^8f^4e^{10ie} - 1006632960Ba^{12}c^8f^4e^{10ie})e^{-2ifx} + (-503316480iAa^{12}c^8f^4e^{12ie} + 301989888B a^{12}c^8f^4e^{12ie})e^{-ifx} + (-50331648iAa^{12}c^8f^4e^{14ie} + 301989888B a^{12}c^8f^4e^{14ie})e^{ifx} + (-50331648iAa^{12}c^8f^4e^{16ie} + 301989888B a^{12}c^8f^4e^{16ie})e^{3ifx} + (-50331648B a^{12}c^8f^4e^{16ie})e^{5ifx}}{32a^3c^2} \\ & + \frac{x(5A - iB)}{16a^3c^2} \end{aligned} \right.$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**3/(c-I*c*tan(f*x+e))**2,x)
```

```
output Piecewise((((33554432*I*A*a**12*c**8*f**4*exp(6*I*e) - 33554432*B*a**12*c**8*f**4*exp(6*I*e))*exp(-6*I*f*x) + (251658240*I*A*a**12*c**8*f**4*exp(8*I*e) - 150994944*B*a**12*c**8*f**4*exp(8*I*e))*exp(-4*I*f*x) + (1006632960*I*A*a**12*c**8*f**4*exp(10*I*e) - 201326592*B*a**12*c**8*f**4*exp(10*I*e))*exp(-2*I*f*x) + (-503316480*I*A*a**12*c**8*f**4*exp(14*I*e) - 301989888*B*a**12*c**8*f**4*exp(14*I*e))*exp(2*I*f*x) + (-50331648*I*A*a**12*c**8*f**4*exp(16*I*e) - 50331648*B*a**12*c**8*f**4*exp(16*I*e))*exp(4*I*f*x))*exp(-12*I*e)/(6442450944*a**15*c**10*f**5), Ne(a**15*c**10*f**5*exp(12*I*e), 0)), (x*(-(5*A - I*B)/(16*a**3*c**2) + (A*exp(10*I*e) + 5*A*exp(8*I*e) + 10*A*exp(6*I*e) + 10*A*exp(4*I*e) + 5*A*exp(2*I*e) + A - I*B*exp(10*I*e) - 3*I*B*exp(8*I*e) - 2*I*B*exp(6*I*e) + 2*I*B*exp(4*I*e) + 3*I*B*exp(2*I*e) + I*B)*exp(-6*I*e)/(32*a**3*c**2)), True)) + x*(5*A - I*B)/(16*a**3*c**2)
```

3.735.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^2,x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

3.735. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^2} dx$

3.735.8 Giac [A] (verification not implemented)

Time = 0.80 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.11

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^2} dx =$$

$$\frac{6(-5iA - B) \log(\tan(fx + e) + i)}{a^3 c^2} + \frac{6(5iA + B) \log(\tan(fx + e) - i)}{a^3 c^2} + \frac{3(-15iA \tan(fx + e)^2 - 3B \tan(fx + e)^2 + 38A \tan(fx + e) - 10iB \tan(fx + e) + 11A + 3iB)}{a^3 c^2 (-i \tan(fx + e) + 1)^2}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^2,x, algorithm="giac")
```

```
output -1/192*(6*(-5*I*A - B)*log(tan(f*x + e) + I)/(a^3*c^2) + 6*(5*I*A + B)*log(tan(f*x + e) - I)/(a^3*c^2) + 3*(-15*I*A*tan(f*x + e)^2 - 3*B*tan(f*x + e)^2 + 38*A*tan(f*x + e) - 10*I*B*tan(f*x + e) + 25*I*A + 9*B)/(a^3*c^2*(-I*tan(f*x + e) + 1)^2) + (-55*I*A*tan(f*x + e)^3 - 11*B*tan(f*x + e)^3 - 201*A*tan(f*x + e)^2 + 33*I*B*tan(f*x + e)^2 + 255*I*A*tan(f*x + e) + 27*B*tan(f*x + e) + 117*A + 3*I*B)/(a^3*c^2*(tan(f*x + e) - I)^3))/f
```

3.735.9 Mupad [B] (verification not implemented)

Time = 9.36 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.12

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^2} dx$$

$$= \frac{\tan(e + fx) \left(\frac{25A}{48a^3c^2} - \frac{B5i}{48a^3c^2} \right) + \tan(e + fx)^3 \left(\frac{5A}{16a^3c^2} - \frac{B1i}{16a^3c^2} \right) + \tan(e + fx)^4 \left(\frac{B}{16a^3c^2} + \frac{A5i}{16a^3c^2} \right) + \tan(e + fx)^5 \operatorname{li} + \tan(e + fx)^4 + \tan(e + fx)^3 2i + 2 \tan(e + fx)^2 + \tan(e + fx) + 1}{f (\tan(e + fx)^5 \operatorname{li} + \tan(e + fx)^4 + \tan(e + fx)^3 2i + 2 \tan(e + fx)^2 + \tan(e + fx) + 1)} - \frac{x(B + A5i) \operatorname{li}}{16a^3c^2}$$

```
input int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i)^2),x)
```

```
output (tan(e + f*x)*((25*A)/(48*a^3*c^2) - (B*5i)/(48*a^3*c^2)) + tan(e + f*x)^3*((5*A)/(16*a^3*c^2) - (B*1i)/(16*a^3*c^2)) + tan(e + f*x)^4*((A*5i)/(16*a^3*c^2) + B/(16*a^3*c^2)) + tan(e + f*x)^2*((A*25i)/(48*a^3*c^2) + (5*B)/(48*a^3*c^2)) + (A*1i)/(6*a^3*c^2) - B/(6*a^3*c^2))/(f*(tan(e + f*x)*1i + 2*tan(e + f*x)^2 + tan(e + f*x)^3*2i + tan(e + f*x)^4 + tan(e + f*x)^5*1i + 1)) - (x*(A*5i + B)*1i)/(16*a^3*c^2)
```

3.735. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^2} dx$

3.736 $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^3} dx$

3.736.1 Optimal result	6754
3.736.2 Mathematica [A] (verified)	6754
3.736.3 Rubi [A] (verified)	6755
3.736.4 Maple [A] (verified)	6757
3.736.5 Fricas [C] (verification not implemented)	6758
3.736.6 Sympy [A] (verification not implemented)	6758
3.736.7 Maxima [F(-2)]	6759
3.736.8 Giac [A] (verification not implemented)	6759
3.736.9 Mupad [B] (verification not implemented)	6760

3.736.1 Optimal result

Integrand size = 41, antiderivative size = 99

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3(c - ic \tan(e + fx))^3} dx$$

$$= \frac{5Ax}{16a^3c^3} + \frac{5A \cos(e + fx) \sin(e + fx)}{16a^3c^3f}$$

$$+ \frac{5A \cos^3(e + fx) \sin(e + fx)}{24a^3c^3f} - \frac{\cos^6(e + fx)(B - A \tan(e + fx))}{6a^3c^3f}$$

output `5/16*A*x/a^3/c^3+5/16*A*cos(f*x+e)*sin(f*x+e)/a^3/c^3/f+5/24*A*cos(f*x+e)^3*sin(f*x+e)/a^3/c^3/f-1/6*cos(f*x+e)^6*(B-A*tan(f*x+e))/a^3/c^3/f`

3.736.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.64

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3(c - ic \tan(e + fx))^3} dx$$

$$= \frac{-32B \cos^6(e + fx) + A(60e + 60fx + 45 \sin(2(e + fx)) + 9 \sin(4(e + fx)) + \sin(6(e + fx)))}{192a^3c^3f}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^3), x]`

3.736. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^3} dx$

output $(-32*B*\text{Cos}[e + f*x]^6 + A*(60*e + 60*f*x + 45*\text{Sin}[2*(e + f*x)] + 9*\text{Sin}[4*(e + f*x)] + \text{Sin}[6*(e + f*x)]))/(192*a^3*c^3*f)$

3.736.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.05, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.195$, Rules used = {3042, 4071, 27, 82, 454, 215, 215, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^3} dx \\ & \quad \downarrow 3042 \\ & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^3} dx \\ & \quad \downarrow 4071 \\ & \frac{ac \int \frac{A + B \tan(e + fx)}{a^4 c^4 (1 - i \tan(e + fx))^4 (i \tan(e + fx) + 1)^4} d \tan(e + fx)}{f} \\ & \quad \downarrow 27 \\ & \frac{\int \frac{A + B \tan(e + fx)}{(1 - i \tan(e + fx))^4 (i \tan(e + fx) + 1)^4} d \tan(e + fx)}{a^3 c^3 f} \\ & \quad \downarrow 82 \\ & \frac{\int \frac{A + B \tan(e + fx)}{(\tan^2(e + fx) + 1)^4} d \tan(e + fx)}{a^3 c^3 f} \\ & \quad \downarrow 454 \\ & \frac{\frac{5}{6} A \int \frac{1}{(\tan^2(e + fx) + 1)^3} d \tan(e + fx) - \frac{B - A \tan(e + fx)}{6(\tan^2(e + fx) + 1)^3}}{a^3 c^3 f} \\ & \quad \downarrow 215 \\ & \frac{\frac{5}{6} A \left(\frac{3}{4} \int \frac{1}{(\tan^2(e + fx) + 1)^2} d \tan(e + fx) + \frac{\tan(e + fx)}{4(\tan^2(e + fx) + 1)^2} \right) - \frac{B - A \tan(e + fx)}{6(\tan^2(e + fx) + 1)^3}}{a^3 c^3 f} \\ & \quad \downarrow 215 \end{aligned}$$

3.736. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^3} dx$

$$\frac{\frac{5}{6}A\left(\frac{3}{4}\left(\frac{1}{2}\int\frac{1}{\tan^2(e+fx)+1}d\tan(e+fx)+\frac{\tan(e+fx)}{2(\tan^2(e+fx)+1)}\right)+\frac{\tan(e+fx)}{4(\tan^2(e+fx)+1)^2}\right)-\frac{B-A\tan(e+fx)}{6(\tan^2(e+fx)+1)^3}}{a^3c^3f}$$

↓ 216

$$\frac{\frac{5}{6}A\left(\frac{3}{4}\left(\frac{1}{2}\arctan(\tan(e+fx))+\frac{\tan(e+fx)}{2(\tan^2(e+fx)+1)}\right)+\frac{\tan(e+fx)}{4(\tan^2(e+fx)+1)^2}\right)-\frac{B-A\tan(e+fx)}{6(\tan^2(e+fx)+1)^3}}{a^3c^3f}$$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^3), x]`

output `(-1/6*(B - A*Tan[e + f*x])/(1 + Tan[e + f*x]^2)^3 + (5*A*(Tan[e + f*x]/(4*(1 + Tan[e + f*x]^2)^2) + (3*(ArcTan[Tan[e + f*x]]/2 + Tan[e + f*x]/(2*(1 + Tan[e + f*x]^2))))/4))/6)/(a^3*c^3*f)`

3.736.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 82 `Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 215 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 454 `Int[((c_) + (d_)*(x_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[((a*d - b*c*x)/(2*a*b*(p + 1))*(a + b*x^2)^(p + 1), x] + Simp[c*((2*p + 3)/(2*a*(p + 1))) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d}, x] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.736.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.39

method	result
risch	$\frac{5Ax}{16a^3c^3} - \frac{B \cos(6fx+6e)}{192f c^3a^3} + \frac{A \sin(6fx+6e)}{192f c^3a^3} - \frac{B \cos(4fx+4e)}{32f c^3a^3} + \frac{3A \sin(4fx+4e)}{64f c^3a^3} - \frac{5B \cos(2fx+2e)}{64f c^3a^3} + \frac{15A \sin(2fx+2e)}{64f c^3a^3}$
norman	$\frac{5Ax}{16ac} - \frac{B}{6acf} + \frac{11A \tan(fx+e)}{16acf} + \frac{5A \tan(fx+e)^3}{6acf} + \frac{5A \tan(fx+e)^5}{16acf} + \frac{15Ax \tan(fx+e)^2}{16ac} + \frac{15Ax \tan(fx+e)^4}{16ac} + \frac{5Ax \tan(fx+e)^6}{16ac}$ $(1+\tan(fx+e)^2)^3 a^2 c^2$
derivativedivides	$-\frac{iA}{16f a^3 c^3 (-i+\tan(fx+e))^2} + \frac{5A \arctan(\tan(fx+e))}{16f a^3 c^3} + \frac{5A}{32f a^3 c^3 (-i+\tan(fx+e))} + \frac{iB}{32f a^3 c^3 (-i+\tan(fx+e))}$
default	$-\frac{iA}{16f a^3 c^3 (-i+\tan(fx+e))^2} + \frac{5A \arctan(\tan(fx+e))}{16f a^3 c^3} + \frac{5A}{32f a^3 c^3 (-i+\tan(fx+e))} + \frac{iB}{32f a^3 c^3 (-i+\tan(fx+e))}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^3,x,method=_R ETURNVERBOSE)`

output `5/16*A*x/a^3/c^3-1/192*B/f/c^3/a^3*cos(6*f*x+6*e)+1/192*A/f/c^3/a^3*sin(6*f*x+6*e)-1/32*B/f/c^3/a^3*cos(4*f*x+4*e)+3/64*A/f/c^3/a^3*sin(4*f*x+4*e)-5/64*B/f/c^3/a^3*cos(2*f*x+2*e)+15/64*A/f/c^3/a^3*sin(2*f*x+2*e)`

3.736. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^3} dx$

3.736.5 Fracas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.27

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^3} dx$$

$$= \frac{(120 A f x e^{(6i f x + 6i e)} + (-i A - B) e^{(12i f x + 12i e)} - 3 (3i A + 2 B) e^{(10i f x + 10i e)} - 15 (3i A + B) e^{(8i f x + 8i e)} - 15 (-3i A + B) e^{(4i f x + 4i e)} - 3 (-3i A + 2 B) e^{(2i f x + 2i e)} + i A - B) e^{(-6i f x - 6i e)}}{384 a^3 c^3 f}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^3,x, algorithm="fracas")
```

```
output 1/384*(120*A*f*x*e^(6*I*f*x + 6*I*e) + (-I*A - B)*e^(12*I*f*x + 12*I*e) - 3*(3*I*A + 2*B)*e^(10*I*f*x + 10*I*e) - 15*(3*I*A + B)*e^(8*I*f*x + 8*I*e) - 15*(-3*I*A + B)*e^(4*I*f*x + 4*I*e) - 3*(-3*I*A + 2*B)*e^(2*I*f*x + 2*I*e) + I*A - B)*e^(-6*I*f*x - 6*I*e)/(a^3*c^3*f)
```

3.736.6 Sympy [A] (verification not implemented)

Time = 0.55 (sec) , antiderivative size = 508, normalized size of antiderivative = 5.13

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^3} dx = \frac{5Ax}{16a^3c^3}$$

$$+ \left\{ \frac{((103079215104iAa^{15}c^{15}f^5e^{6ie} - 103079215104Ba^{15}c^{15}f^5e^{6ie})e^{-6ifx} + (927712935936iAa^{15}c^{15}f^5e^{8ie} - 618475290624Ba^{15}c^{15}f^5e^{8ie})e^{-8ifx} + \dots)}{64a^3c^3} \right.$$

$$\left. x \left(-\frac{5A}{16a^3c^3} + \frac{(Ae^{12ie} + 6Ae^{10ie} + 15Ae^{8ie} + 20Ae^{6ie} + 15Ae^{4ie} + 6Ae^{2ie} + A - iBe^{12ie} - 4iBe^{10ie} - 5iBe^{8ie} + 5iBe^{4ie} + 4iBe^{2ie} + iB)e^{-6ie}}{64a^3c^3} \right) \right.$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**3/(c-I*c*tan(f*x+e))**3,x)
```

output `5*A*x/(16*a**3*c**3) + Piecewise((((103079215104*I*A*a**15*c**15*f**5*exp(6*I*e) - 103079215104*B*a**15*c**15*f**5*exp(6*I*e))*exp(-6*I*f*x) + (927712935936*I*A*a**15*c**15*f**5*exp(8*I*e) - 618475290624*B*a**15*c**15*f**5*exp(8*I*e))*exp(-4*I*f*x) + (4638564679680*I*A*a**15*c**15*f**5*exp(10*I*e) - 1546188226560*B*a**15*c**15*f**5*exp(10*I*e))*exp(-2*I*f*x) + (-4638564679680*I*A*a**15*c**15*f**5*exp(14*I*e) - 1546188226560*B*a**15*c**15*f**5*exp(14*I*e))*exp(2*I*f*x) + (-927712935936*I*A*a**15*c**15*f**5*exp(16*I*e) - 618475290624*B*a**15*c**15*f**5*exp(16*I*e))*exp(4*I*f*x) + (-103079215104*I*A*a**15*c**15*f**5*exp(18*I*e) - 103079215104*B*a**15*c**15*f**5*exp(18*I*e))*exp(6*I*f*x))*exp(-12*I*e)/(39582418599936*a**18*c**18*f**6), Ne(a**18*c**18*f**6*exp(12*I*e), 0)), (x*(-5*A/(16*a**3*c**3) + (A*exp(12*I*e) + 6*A*exp(10*I*e) + 15*A*exp(8*I*e) + 20*A*exp(6*I*e) + 15*A*exp(4*I*e) + 6*A*exp(2*I*e) + A - I*B*exp(12*I*e) - 4*I*B*exp(10*I*e) - 5*I*B*exp(8*I*e) + 5*I*B*exp(4*I*e) + 4*I*B*exp(2*I*e) + I*B)*exp(-6*I*e)/(64*a**3*c**3)), True))`

3.736.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.736.8 Giac [A] (verification not implemented)

Time = 0.77 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.75

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^3} dx$$

$$= \frac{\frac{15(fx+e)A}{a^3c^3} + \frac{15A \tan(fx+e)^5 + 40A \tan(fx+e)^3 + 33A \tan(fx+e) - 8B}{(\tan(fx+e)^2 + 1)^3 a^3c^3}}{48f}$$

3.736. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^3} dx$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^3,x, algorithm="giac")`

output `1/48*(15*(f*x + e)*A/(a^3*c^3) + (15*A*tan(f*x + e)^5 + 40*A*tan(f*x + e)^3 + 33*A*tan(f*x + e) - 8*B)/((tan(f*x + e)^2 + 1)^3*a^3*c^3))/f`

3.736.9 Mupad [B] (verification not implemented)

Time = 8.73 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.65

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^3} dx$$

$$= \frac{5Ax}{16a^3c^3} + \frac{\cos(e + fx)^6 \left(\frac{5A \tan(e+fx)^5}{16} + \frac{5A \tan(e+fx)^3}{6} + \frac{11A \tan(e+fx)}{16} - \frac{B}{6} \right)}{a^3c^3f}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*i)^3*(c - c*tan(e + f*x)*i)^3),x)`

output `(5*A*x)/(16*a^3*c^3) + (cos(e + f*x)^6*((11*A*tan(e + f*x))/16 - B/6 + (5*A*tan(e + f*x)^3)/6 + (5*A*tan(e + f*x)^5)/16))/(a^3*c^3*f)`

3.737 $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^4} dx$

3.737.1 Optimal result 6761
 3.737.2 Mathematica [A] (verified) 6762
 3.737.3 Rubi [A] (verified) 6762
 3.737.4 Maple [A] (verified) 6764
 3.737.5 Fricas [A] (verification not implemented) 6765
 3.737.6 Sympy [A] (verification not implemented) 6765
 3.737.7 Maxima [F(-2)] 6766
 3.737.8 Giac [A] (verification not implemented) 6767
 3.737.9 Mupad [B] (verification not implemented) 6767

3.737.1 Optimal result

Integrand size = 41, antiderivative size = 251

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3(c - ic \tan(e + fx))^4} dx$$

$$= \frac{5(7A + iB)x}{128a^3c^4} + \frac{A + iB}{96a^3c^4f(i - \tan(e + fx))^3} - \frac{5iA - 3B}{128a^3c^4f(i - \tan(e + fx))^2}$$

$$- \frac{5(3A + iB)}{128a^3c^4f(i - \tan(e + fx))} - \frac{iA + B}{64a^3c^4f(i + \tan(e + fx))^4}$$

$$- \frac{2A - iB}{48a^3c^4f(i + \tan(e + fx))^3} + \frac{5iA + B}{64a^3c^4f(i + \tan(e + fx))^2} + \frac{5A}{32a^3c^4f(i + \tan(e + fx))}$$

```
output 5/128*(7*A+I*B)*x/a^3/c^4+1/96*(A+I*B)/a^3/c^4/f/(I-tan(f*x+e))^3+1/128*(-
5*I*A+3*B)/a^3/c^4/f/(I-tan(f*x+e))^2-5/128*(3*A+I*B)/a^3/c^4/f/(I-tan(f*x
+e))+1/64*(-I*A-B)/a^3/c^4/f/(I+tan(f*x+e))^4+1/48*(-2*A+I*B)/a^3/c^4/f/(I
+tan(f*x+e))^3+1/64*(5*I*A+B)/a^3/c^4/f/(I+tan(f*x+e))^2+5/32*A/a^3/c^4/f/
(I+tan(f*x+e))
```


3.737.2 Mathematica [A] (verified)

Time = 6.40 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.88

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ict \tan(e + fx))^4} dx$$

$$= \frac{\sec^6(e + fx)(319A - 23iB - (113A + 119iB) \cos(2(e + fx)) - 13A \cos(4(e + fx)) - 43iB \cos(4(e + fx)) - 7A \cos(6(e + fx)) - (7iB) \cos(6(e + fx)) + (315i)A \sin(2(e + fx)) - 45B \sin(2(e + fx)) + (63i)A \sin(4(e + fx)) - 9B \sin(4(e + fx)) + (7i)A \sin(6(e + fx)) - B \sin(6(e + fx)) + 60(7A + iB) \operatorname{ArcTan}[\tan(e + fx)](I + \tan(e + fx)))}{(1536a^3c^4f(-I + \tan(e + fx))^3(I + \tan(e + fx))^4)}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^4),x]`

output `(Sec[e + f*x]^6*(319*A - (23*I)*B - (113*A + (119*I)*B)*Cos[2*(e + f*x)] - 13*A*Cos[4*(e + f*x)] - (43*I)*B*Cos[4*(e + f*x)] - A*Cos[6*(e + f*x)] - (7*I)*B*Cos[6*(e + f*x)] + (315*I)*A*Sin[2*(e + f*x)] - 45*B*Sin[2*(e + f*x)] + (63*I)*A*Sin[4*(e + f*x)] - 9*B*Sin[4*(e + f*x)] + (7*I)*A*Sin[6*(e + f*x)] - B*Sin[6*(e + f*x)] + 60*(7*A + I*B)*ArcTan[Tan[e + f*x]]*(I + Tan[e + f*x]))/(1536*a^3*c^4*f*(-I + Tan[e + f*x])^3*(I + Tan[e + f*x])^4)`

3.737.3 Rubi [A] (verified)Time = 0.44 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.79, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ict \tan(e + fx))^4} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ict \tan(e + fx))^4} dx$$

$$\downarrow \text{4071}$$

$$\frac{ac \int \frac{A + B \tan(e + fx)}{a^4 c^5 (1 - i \tan(e + fx))^5 (i \tan(e + fx) + 1)^4} d \tan(e + fx)}{f}$$

$$\downarrow \text{27}$$

3.737. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ict \tan(e + fx))^4} dx$

$$\int \frac{A+B \tan(e+fx)}{(1-i \tan(e+fx))^5(i \tan(e+fx)+1)^4} d \tan(e+fx)$$

↓ 86

$$\int \left(-\frac{5A}{32(\tan(e+fx)+i)^2} + \frac{5(7A+iB)}{128(\tan^2(e+fx)+1)} - \frac{5(3A+iB)}{128(\tan(e+fx)-i)^2} + \frac{i(5A+3iB)}{64(\tan(e+fx)-i)^3} - \frac{i(5A-iB)}{32(\tan(e+fx)+i)^3} + \frac{A+iB}{32(\tan(e+fx)-i)^4} \right) \frac{d \tan(e+fx)}{a^3 c^4 f}$$

↓ 2009

$$\frac{5}{128}(7A+iB) \arctan(\tan(e+fx)) - \frac{5(3A+iB)}{128(-\tan(e+fx)+i)} - \frac{-3B+5iA}{128(-\tan(e+fx)+i)^2} + \frac{B+5iA}{64(\tan(e+fx)+i)^2} + \frac{A+iB}{96(-\tan(e+fx)+i)^3} \frac{d \tan(e+fx)}{a^3 c^4 f}$$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^4), x]`

output `((5*(7*A + I*B)*ArcTan[Tan[e + f*x]])/128 + (A + I*B)/(96*(I - Tan[e + f*x])^3) - ((5*I)*A - 3*B)/(128*(I - Tan[e + f*x])^2) - (5*(3*A + I*B))/(128*(I - Tan[e + f*x])) - (I*A + B)/(64*(I + Tan[e + f*x])^4) - (2*A - I*B)/(48*(I + Tan[e + f*x])^3) + ((5*I)*A + B)/(64*(I + Tan[e + f*x])^2) + (5*A)/(32*(I + Tan[e + f*x]))) / (a^3*c^4*f)`

3.737.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x_, x], x] /; FreeQ[a, x] && !MatchQ[F_x_, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.737. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^4} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.737.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.04

method	result
norman	$\frac{5(iB+7A)x}{128ac} - \frac{iA+B}{8acf} + \frac{(-5iB+93A)\tan(fx+e)}{128acf} + \frac{73(iB+7A)\tan(fx+e)^3}{384acf} + \frac{55(iB+7A)\tan(fx+e)^5}{384acf} + \frac{5(iB+7A)\tan(fx+e)^7}{128acf} + \frac{5(iB+7A)\tan(fx+e)^9}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{11}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{13}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{15}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{17}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{19}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{21}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{23}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{25}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{27}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{29}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{31}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{33}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{35}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{37}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{39}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{41}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{43}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{45}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{47}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{49}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{51}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{53}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{55}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{57}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{59}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{61}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{63}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{65}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{67}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{69}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{71}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{73}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{75}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{77}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{79}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{81}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{83}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{85}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{87}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{89}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{91}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{93}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{95}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{97}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{99}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{101}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{103}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{105}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{107}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{109}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{111}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{113}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{115}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{117}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{119}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{121}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{123}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{125}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{127}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{129}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{131}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{133}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{135}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{137}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{139}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{141}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{143}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{145}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{147}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{149}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{151}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{153}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{155}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{157}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{159}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{161}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{163}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{165}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{167}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{169}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{171}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{173}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{175}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{177}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{179}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{181}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{183}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{185}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{187}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{189}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{191}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{193}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{195}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{197}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{199}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{201}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{203}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{205}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{207}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{209}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{211}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{213}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{215}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{217}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{219}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{221}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{223}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{225}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{227}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{229}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{231}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{233}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{235}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{237}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{239}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{241}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{243}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{245}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{247}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{249}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{251}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{253}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{255}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{257}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{259}}{128acf} + \frac{5(iB+7A)\tan(fx+e)^{261}}{128acf}$
risch	$\frac{i \sin(2fx+2e)B}{64a^3c^4f} + \frac{35xA}{128a^3c^4} - \frac{e^{8i(fx+e)}B}{1024a^3c^4f} - \frac{7i \cos(2fx+2e)A}{128a^3c^4f} - \frac{\cos(6fx+6e)B}{128a^3c^4f} - \frac{i \cos(6fx+6e)A}{128a^3c^4f} - \frac{7i \cos(4fx+4e)B}{256a^3c^4f} + \frac{7i \cos(4fx+4e)A}{256a^3c^4f}$
derivativedivides	$-\frac{A}{96fa^3c^4(-i+\tan(fx+e))^3} + \frac{iB}{48fa^3c^4(i+\tan(fx+e))^3} + \frac{5iA}{64fa^3c^4(i+\tan(fx+e))^2} + \frac{15A}{128fa^3c^4(-i+\tan(fx+e))^2}$
default	$-\frac{A}{96fa^3c^4(-i+\tan(fx+e))^3} + \frac{iB}{48fa^3c^4(i+\tan(fx+e))^3} + \frac{5iA}{64fa^3c^4(i+\tan(fx+e))^2} + \frac{15A}{128fa^3c^4(-i+\tan(fx+e))^2}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^4,x,method=_RETURVERBOSE)`

output
$$\frac{5}{128} \frac{(7A+I*B)}{a/c*x-1/8*(I*A+B)/a/c/f+1/128*(-5*I*B+93*A)/a/c/f*\tan(f*x+e)+73/384*(7A+I*B)/a/c/f*\tan(f*x+e)^3+55/384*(7A+I*B)/a/c/f*\tan(f*x+e)^5+5/128*(7A+I*B)/a/c/f*\tan(f*x+e)^7+5/32*(7A+I*B)/a/c*x*\tan(f*x+e)^2+15/64*(7A+I*B)/a/c*x*\tan(f*x+e)^4+5/32*(7A+I*B)/a/c*x*\tan(f*x+e)^6+5/128*(7A+I*B)/a/c*x*\tan(f*x+e)^8/a^2/c^3/(1+\tan(f*x+e)^2)^4$$

3.737.
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^4} dx$$

3.737.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.60

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^4} dx$$

$$= \frac{(120(7A + iB)fx e^{(6i fx + 6ie)} - 3(iA + B)e^{(14i fx + 14ie)} - 4(7iA + 5B)e^{(12i fx + 12ie)} - 18(7iA + 3B)e^{(10i fx + 10ie)} - 6(-7iA + 5B)e^{(8i fx + 8ie)} - 36(-7iA + 3B)e^{(4i fx + 4ie)} - 6(-7iA + 5B)e^{(2i fx + 2ie)} + 4iA - 4B)e^{(-6i fx - 6ie)}}{(a^3 c^4 f)}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^4,x, algorithm="fricas")
```

```
output 1/3072*(120*(7*A + I*B)*f*x*e^(6*I*f*x + 6*I*e) - 3*(I*A + B)*e^(14*I*f*x + 14*I*e) - 4*(7*I*A + 5*B)*e^(12*I*f*x + 12*I*e) - 18*(7*I*A + 3*B)*e^(10*I*f*x + 10*I*e) - 60*(7*I*A + B)*e^(8*I*f*x + 8*I*e) - 36*(-7*I*A + 3*B)*e^(4*I*f*x + 4*I*e) - 6*(-7*I*A + 5*B)*e^(2*I*f*x + 2*I*e) + 4*I*A - 4*B)*e^(-6*I*f*x - 6*I*e)/(a^3*c^4*f)
```

3.737.6 Sympy [A] (verification not implemented)

Time = 0.63 (sec) , antiderivative size = 605, normalized size of antiderivative = 2.41

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^4} dx$$

$$= \left\{ \frac{((13510798882111488iAa^{18}c^{24}f^6e^{6ie} - 13510798882111488Ba^{18}c^{24}f^6e^{6ie})e^{-6ifx} + (141863388262170624iAa^{18}c^{24}f^6e^{8ie} - 1013309916158368iBa^{18}c^{24}f^6e^{8ie})e^{-12ifx} + (141863388262170624iAa^{18}c^{24}f^6e^{10ie} - 1013309916158368iBa^{18}c^{24}f^6e^{10ie})e^{-18ifx} + (141863388262170624iAa^{18}c^{24}f^6e^{12ie} - 1013309916158368iBa^{18}c^{24}f^6e^{12ie})e^{-24ifx} + (141863388262170624iAa^{18}c^{24}f^6e^{14ie} - 1013309916158368iBa^{18}c^{24}f^6e^{14ie})e^{-30ifx} + (141863388262170624iAa^{18}c^{24}f^6e^{16ie} - 1013309916158368iBa^{18}c^{24}f^6e^{16ie})e^{-36ifx} + (141863388262170624iAa^{18}c^{24}f^6e^{18ie} - 1013309916158368iBa^{18}c^{24}f^6e^{18ie})e^{-42ifx} + (141863388262170624iAa^{18}c^{24}f^6e^{20ie} - 1013309916158368iBa^{18}c^{24}f^6e^{20ie})e^{-48ifx} + (141863388262170624iAa^{18}c^{24}f^6e^{22ie} - 1013309916158368iBa^{18}c^{24}f^6e^{22ie})e^{-54ifx} + (141863388262170624iAa^{18}c^{24}f^6e^{24ie} - 1013309916158368iBa^{18}c^{24}f^6e^{24ie})e^{-60ifx}}{128a^3c^4} \right.$$

$$\left. + \frac{x(35A + 5iB)}{128a^3c^4} \right.$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**3/(c-I*c*tan(f*x+e))**4,x)
```

```
output Piecewise((((13510798882111488*I*A*a**18*c**24*f**6*exp(6*I*e) - 135107988
82111488*B*a**18*c**24*f**6*exp(6*I*e))*exp(-6*I*f*x) + (14186338826217062
4*I*A*a**18*c**24*f**6*exp(8*I*e) - 101330991615836160*B*a**18*c**24*f**6*
exp(8*I*e))*exp(-4*I*f*x) + (851180329573023744*I*A*a**18*c**24*f**6*exp(1
0*I*e) - 364791569817010176*B*a**18*c**24*f**6*exp(10*I*e))*exp(-2*I*f*x)
+ (-1418633882621706240*I*A*a**18*c**24*f**6*exp(14*I*e) - 202661983231672
320*B*a**18*c**24*f**6*exp(14*I*e))*exp(2*I*f*x) + (-425590164786511872*I*
A*a**18*c**24*f**6*exp(16*I*e) - 182395784908505088*B*a**18*c**24*f**6*exp
(16*I*e))*exp(4*I*f*x) + (-94575592174780416*I*A*a**18*c**24*f**6*exp(18*I
*e) - 67553994410557440*B*a**18*c**24*f**6*exp(18*I*e))*exp(6*I*f*x) + (-1
0133099161583616*I*A*a**18*c**24*f**6*exp(20*I*e) - 10133099161583616*B*a
**18*c**24*f**6*exp(20*I*e))*exp(8*I*f*x))*exp(-12*I*e)/(103762935414616227
84*a**21*c**28*f**7), Ne(a**21*c**28*f**7*exp(12*I*e), 0)), (x*(-(35*A + 5
*I*B)/(128*a**3*c**4) + (A*exp(14*I*e) + 7*A*exp(12*I*e) + 21*A*exp(10*I*e
) + 35*A*exp(8*I*e) + 35*A*exp(6*I*e) + 21*A*exp(4*I*e) + 7*A*exp(2*I*e) +
A - I*B*exp(14*I*e) - 5*I*B*exp(12*I*e) - 9*I*B*exp(10*I*e) - 5*I*B*exp(8
*I*e) + 5*I*B*exp(6*I*e) + 9*I*B*exp(4*I*e) + 5*I*B*exp(2*I*e) + I*B)*exp(
-6*I*e)/(128*a**3*c**4)), True)) + x*(35*A + 5*I*B)/(128*a**3*c**4)
```

3.737.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^4} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^4,x, al
gorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.737.8 Giac [A] (verification not implemented)

Time = 0.85 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.00

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ict \tan(e + fx))^4} dx$$

$$= \frac{60(7iA - B) \log(\tan(fx + e) + i)}{a^3 c^4} - \frac{60(7iA - B) \log(\tan(fx + e) - i)}{a^3 c^4} + \frac{2(385A \tan(fx + e)^3 + 55iB \tan(fx + e)^3 - 1335iA \tan(fx + e)^2 + 225B \tan(fx + e)^2 - 1575iA \tan(fx + e) - 321iB \tan(fx + e) + 641iA - 167iB)}{a^3 c^4 (i \tan(fx + e) + 1)^3} + \frac{(-875iA \tan(fx + e)^4 + 125B \tan(fx + e)^4 + 3980A \tan(fx + e)^3 + 500iB \tan(fx + e)^3 + 6930iA \tan(fx + e)^2 - 702B \tan(fx + e)^2 - 5548A \tan(fx + e) - 340iB \tan(fx + e) - 1771iA - 35iB)}{a^3 c^4 (\tan(fx + e) + i)^4} / f$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^4,x, algorithm="giac")
```

```
output 1/3072*(60*(7*I*A - B)*log(tan(f*x + e) + I)/(a^3*c^4) - 60*(7*I*A - B)*log(tan(f*x + e) - I)/(a^3*c^4) + 2*(385*A*tan(f*x + e)^3 + 55*I*B*tan(f*x + e)^3 - 1335*I*A*tan(f*x + e)^2 + 225*B*tan(f*x + e)^2 - 1575*A*tan(f*x + e) - 321*I*B*tan(f*x + e) + 641*I*A - 167*B)/(a^3*c^4*(I*tan(f*x + e) + 1)^3) + (-875*I*A*tan(f*x + e)^4 + 125*B*tan(f*x + e)^4 + 3980*A*tan(f*x + e)^3 + 500*I*B*tan(f*x + e)^3 + 6930*I*A*tan(f*x + e)^2 - 702*B*tan(f*x + e)^2 - 5548*A*tan(f*x + e) - 340*I*B*tan(f*x + e) - 1771*I*A - 35*B)/(a^3*c^4*(tan(f*x + e) + I)^4))/f
```

3.737.9 Mupad [B] (verification not implemented)

Time = 10.53 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.14

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ict \tan(e + fx))^4} dx$$

$$= \frac{\tan(e + fx) \left(-\frac{11B}{128a^3c^4} + \frac{A77i}{128a^3c^4} \right) + \tan(e + fx)^3 \left(-\frac{5B}{48a^3c^4} + \frac{A35i}{48a^3c^4} \right) + \tan(e + fx)^4 \left(\frac{35A}{48a^3c^4} + \frac{B5i}{48a^3c^4} \right) - f \left(\tan(e + fx)^7 + \tan(e + fx)^6 \operatorname{li} + 3 \tan(e + fx)^5 + \frac{5x(7A + B \operatorname{li})}{128a^3c^4} \right)}{f \left(\tan(e + fx)^7 + \tan(e + fx)^6 \operatorname{li} + 3 \tan(e + fx)^5 + \frac{5x(7A + B \operatorname{li})}{128a^3c^4} \right)}$$

```
input int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*i)^3*(c - c*tan(e + f*x)*i)^4),x)
```

output

$$\begin{aligned}
& (\tan(e + f*x)*((A*77i)/(128*a^3*c^4) - (11*B)/(128*a^3*c^4)) + \tan(e + f*x) \\
&)^3*((A*35i)/(48*a^3*c^4) - (5*B)/(48*a^3*c^4)) + \tan(e + f*x)^4*((35*A)/(\\
& 48*a^3*c^4) + (B*5i)/(48*a^3*c^4)) + \tan(e + f*x)^5*((A*35i)/(128*a^3*c^4) \\
& - (5*B)/(128*a^3*c^4)) + \tan(e + f*x)^6*((35*A)/(128*a^3*c^4) + (B*5i)/(1 \\
& 28*a^3*c^4)) + \tan(e + f*x)^2*((77*A)/(128*a^3*c^4) + (B*11i)/(128*a^3*c^4 \\
&)) + A/(8*a^3*c^4) - (B*1i)/(8*a^3*c^4))/(f*(\tan(e + f*x) + \tan(e + f*x)^2 \\
& *3i + 3*\tan(e + f*x)^3 + \tan(e + f*x)^4*3i + 3*\tan(e + f*x)^5 + \tan(e + f* \\
& x)^6*1i + \tan(e + f*x)^7 + 1i)) + (5*x*(7*A + B*1i))/(128*a^3*c^4)
\end{aligned}$$

3.737.
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^4} dx$$

3.738 $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^5} dx$

3.738.1 Optimal result	6769
3.738.2 Mathematica [A] (verified)	6770
3.738.3 Rubi [A] (verified)	6770
3.738.4 Maple [A] (verified)	6772
3.738.5 Fricas [A] (verification not implemented)	6773
3.738.6 Sympy [A] (verification not implemented)	6773
3.738.7 Maxima [F(-2)]	6774
3.738.8 Giac [A] (verification not implemented)	6775
3.738.9 Mupad [B] (verification not implemented)	6775

3.738.1 Optimal result

Integrand size = 41, antiderivative size = 287

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3(c - ic \tan(e + fx))^5} dx$$

$$= \frac{7(4A + iB)x}{128a^3c^5} + \frac{A + iB}{192a^3c^5 f(i - \tan(e + fx))^3} - \frac{3iA - 2B}{128a^3c^5 f(i - \tan(e + fx))^2}$$

$$- \frac{3(7A + 3iB)}{256a^3c^5 f(i - \tan(e + fx))} + \frac{A - iB}{80a^3c^5 f(i + \tan(e + fx))^5} - \frac{2iA + B}{64a^3c^5 f(i + \tan(e + fx))^4}$$

$$- \frac{5A - iB}{96a^3c^5 f(i + \tan(e + fx))^3} + \frac{5iA}{64a^3c^5 f(i + \tan(e + fx))^2} + \frac{5(7A + iB)}{256a^3c^5 f(i + \tan(e + fx))}$$

```
output 7/128*(4*A+I*B)*x/a^3/c^5+1/192*(A+I*B)/a^3/c^5/f/(I-tan(f*x+e))^3+1/128*(
-3*I*A+2*B)/a^3/c^5/f/(I-tan(f*x+e))^2-3/256*(7*A+3*I*B)/a^3/c^5/f/(I-tan(
f*x+e))+1/80*(A-I*B)/a^3/c^5/f/(I+tan(f*x+e))^5+1/64*(-2*I*A-B)/a^3/c^5/f/
(I+tan(f*x+e))^4+1/96*(-5*A+I*B)/a^3/c^5/f/(I+tan(f*x+e))^3+5/64*I*A/a^3/c
^5/f/(I+tan(f*x+e))^2+5/256*(7*A+I*B)/a^3/c^5/f/(I+tan(f*x+e))
```


3.738.2 Mathematica [A] (verified)

Time = 6.79 (sec) , antiderivative size = 261, normalized size of antiderivative = 0.91

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ictan(e + fx))^5} dx$$

$$= \frac{\sec^7(e + fx)(2100iA \cos(e + fx) + 63(-8iA + 7B) \cos(3(e + fx)) - 56iA \cos(5(e + fx)) + 119B \cos(5(e + fx)))}{(a + ia \tan(e + fx))^3 (c - ictan(e + fx))^5}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^5),x]`

output `(Sec[e + f*x]^7*((2100*I)*A*Cos[e + f*x] + 63*((-8*I)*A + 7*B)*Cos[3*(e + f*x)] - (56*I)*A*Cos[5*(e + f*x)] + 119*B*Cos[5*(e + f*x)] - (4*I)*A*Cos[7*(e + f*x)] + 16*B*Cos[7*(e + f*x)] + 872*A*Sin[e + f*x] + (218*I)*B*Sin[e + f*x] - 420*(4*A + I*B)*ArcTan[Tan[e + f*x]]*Sec[e + f*x]*(Cos[2*(e + f*x)] - I*Sin[2*(e + f*x)]) - 956*A*Sin[3*(e + f*x)] - (239*I)*B*Sin[3*(e + f*x)] - 164*A*Sin[5*(e + f*x)] - (41*I)*B*Sin[5*(e + f*x)] - 16*A*Sin[7*(e + f*x)] - (4*I)*B*Sin[7*(e + f*x)]))/(7680*a^3*c^5*f*(-I + Tan[e + f*x])^3*(I + Tan[e + f*x])^5)`

3.738.3 Rubi [A] (verified)Time = 0.48 (sec) , antiderivative size = 225, normalized size of antiderivative = 0.78, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ictan(e + fx))^5} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ictan(e + fx))^5} dx$$

$$\downarrow \text{4071}$$

$$\frac{ac \int \frac{A+B \tan(e+fx)}{a^4 c^6 (1-i \tan(e+fx))^6 (i \tan(e+fx)+1)^4} d \tan(e + fx)}{f}$$

3.738. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3 (c-ic \tan(e+fx))^5} dx$

$$\int \frac{A+B \tan(e+fx)}{(1-i \tan(e+fx))^6 (i \tan(e+fx)+1)^4} d \tan(e+fx)$$

$$\frac{\int \left(-\frac{5iA}{32(\tan(e+fx)+i)^3} + \frac{7(4A+iB)}{128(\tan^2(e+fx)+1)} - \frac{3(7A+3iB)}{256(\tan(e+fx)-i)^2} - \frac{5(7A+iB)}{256(\tan(e+fx)+i)^2} + \frac{i(3A+2iB)}{64(\tan(e+fx)-i)^3} + \frac{A+iB}{64(\tan(e+fx)-i)^4} \right)}{a^3 c^5 f}$$

$$\frac{\frac{7}{128}(4A+iB) \arctan(\tan(e+fx)) - \frac{3(7A+3iB)}{256(-\tan(e+fx)+i)} + \frac{5(7A+iB)}{256(\tan(e+fx)+i)} - \frac{-2B+3iA}{128(-\tan(e+fx)+i)^2} + \frac{A+iB}{192(-\tan(e+fx)+i)^3}}{a^3 c^5 f}$$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^5), x]`

output `((7*(4*A + I*B)*ArcTan[Tan[e + f*x]]/128 + (A + I*B)/(192*(I - Tan[e + f*x])^3) - ((3*I)*A - 2*B)/(128*(I - Tan[e + f*x])^2) - (3*(7*A + (3*I)*B))/(256*(I - Tan[e + f*x])) + (A - I*B)/(80*(I + Tan[e + f*x])^5) - ((2*I)*A + B)/(64*(I + Tan[e + f*x])^4) - (5*A - I*B)/(96*(I + Tan[e + f*x])^3) + (((5*I)/64)*A)/(I + Tan[e + f*x])^2 + (5*(7*A + I*B))/(256*(I + Tan[e + f*x]))) / (a^3*c^5*f)`

3.738.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

$$3.738. \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3 (c-ic \tan(e+fx))^5} dx$$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.738.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 335, normalized size of antiderivative = 1.17

method	result
norman	$\frac{7(iB+4A)x}{128ac} - \frac{8iA+3B}{40acf} + \frac{B \tan(fx+e)^2}{8acf} + \frac{79(iB+4A) \tan(fx+e)^3}{192acf} + \frac{7(iB+4A) \tan(fx+e)^5}{15acf} + \frac{49(iB+4A) \tan(fx+e)^7}{192acf} + \frac{7(iB+4A) \tan(fx+e)^9}{128acf}$
risch	$-\frac{i \sin(4fx+4e)B}{128a^3c^5f} + \frac{7xA}{32a^3c^5} - \frac{e^{10i(fx+e)}B}{2560a^3c^5f} + \frac{7ixB}{128a^3c^5} - \frac{3e^{8i(fx+e)}B}{1024a^3c^5f} - \frac{13i \sin(6fx+6e)B}{1536a^3c^5f} - \frac{5 \cos(6fx+6e)B}{512a^3c^5f}$
derivativedivides	$-\frac{iA}{32fa^3c^5(i+\tan(fx+e))^4} + \frac{A}{80fa^3c^5(i+\tan(fx+e))^5} + \frac{9iB}{256fa^3c^5(-i+\tan(fx+e))} - \frac{5A}{96fa^3c^5(i+\tan(fx+e))^3}$
default	$-\frac{iA}{32fa^3c^5(i+\tan(fx+e))^4} + \frac{A}{80fa^3c^5(i+\tan(fx+e))^5} + \frac{9iB}{256fa^3c^5(-i+\tan(fx+e))} - \frac{5A}{96fa^3c^5(i+\tan(fx+e))^3}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^5,x,method=_RETURNVERBOSE)`

output $(7/128*(4*A+I*B)/a/c*x-1/40*(3*B+8*I*A)/a/c/f+1/8/a/c/f*B*\tan(f*x+e)^2+79/192*(4*A+I*B)/a/c/f*\tan(f*x+e)^3+7/15*(4*A+I*B)/a/c/f*\tan(f*x+e)^5+49/192*(4*A+I*B)/a/c/f*\tan(f*x+e)^7+7/128*(4*A+I*B)/a/c/f*\tan(f*x+e)^9+35/128*(4*A+I*B)/a/c*x*\tan(f*x+e)^2+35/64*(4*A+I*B)/a/c*x*\tan(f*x+e)^4+35/64*(4*A+I*B)/a/c*x*\tan(f*x+e)^6+35/128*(4*A+I*B)/a/c*x*\tan(f*x+e)^8+7/128*(4*A+I*B)/a/c*x*\tan(f*x+e)^{10}+1/128*(100*A-7*I*B)/a/c/f*\tan(f*x+e))/(1+\tan(f*x+e))^2)^5/a^2/c^4$

3.738. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^5} dx$

3.738.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.55

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^5} dx$$

$$= \frac{(840(4A + iB)fx e^{6i fx + 6ie} - 6(iA + B)e^{16i fx + 16ie} - 15(4iA + 3B)e^{14i fx + 14ie} - 140(2iA + B)e^{12i fx + 12ie} - 210(4iA + B)e^{10i fx + 10ie} - 2100iA e^{8i fx + 8ie} - 420(-2iA + B)e^{4i fx + 4ie} - 30(-4iA + 3B)e^{2i fx + 2ie} + 10iA - 10B)e^{-6i fx - 6ie}}{a^3 c^5 f}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^5,x, algorithm="fricas")
```

```
output 1/15360*(840*(4*A + I*B)*f*x*e^(6*I*f*x + 6*I*e) - 6*(I*A + B)*e^(16*I*f*x + 16*I*e) - 15*(4*I*A + 3*B)*e^(14*I*f*x + 14*I*e) - 140*(2*I*A + B)*e^(12*I*f*x + 12*I*e) - 210*(4*I*A + B)*e^(10*I*f*x + 10*I*e) - 2100*I*A*e^(8*I*f*x + 8*I*e) - 420*(-2*I*A + B)*e^(4*I*f*x + 4*I*e) - 30*(-4*I*A + 3*B)*e^(2*I*f*x + 2*I*e) + 10*I*A - 10*B)*e^(-6*I*f*x - 6*I*e)/(a^3*c^5*f)
```

3.738.6 Sympy [A] (verification not implemented)

Time = 0.80 (sec) , antiderivative size = 646, normalized size of antiderivative = 2.25

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^5} dx$$

$$= \left\{ \frac{(-7263405479023135948800iAa^{21}c^{35}f^7e^{14ie}e^{2ifx} + (34587645138205409280iAa^{21}c^{35}f^7e^{6ie} - 34587645138205409280Ba^{21}c^{35}f^7e^{6ie})e^{-6ifx}}{256a^3c^5} \right.$$

$$+ \frac{x(28A + 7iB)}{128a^3c^5}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**3/(c-I*c*tan(f*x+e))**5,x)
```

```
output Piecewise((( -7263405479023135948800*I*A*a**21*c**35*f**7*exp(14*I*e)*exp(2
*I*f*x) + (34587645138205409280*I*A*a**21*c**35*f**7*exp(6*I*e) - 34587645
138205409280*B*a**21*c**35*f**7*exp(6*I*e))*exp(-6*I*f*x) + (4150517416584
64911360*I*A*a**21*c**35*f**7*exp(8*I*e) - 311288806243848683520*B*a**21*c
**35*f**7*exp(8*I*e))*exp(-4*I*f*x) + (2905362191609254379520*I*A*a**21*c
**35*f**7*exp(10*I*e) - 1452681095804627189760*B*a**21*c**35*f**7*exp(10*I
e))*exp(-2*I*f*x) + (-2905362191609254379520*I*A*a**21*c**35*f**7*exp(16*I
*e) - 726340547902313594880*B*a**21*c**35*f**7*exp(16*I*e))*exp(4*I*f*x) +
(-968454063869751459840*I*A*a**21*c**35*f**7*exp(18*I*e) - 48422703193487
5729920*B*a**21*c**35*f**7*exp(18*I*e))*exp(6*I*f*x) + (-20752587082923245
5680*I*A*a**21*c**35*f**7*exp(20*I*e) - 155644403121924341760*B*a**21*c**3
5*f**7*exp(20*I*e))*exp(8*I*f*x) + (-20752587082923245568*I*A*a**21*c**35*
f**7*exp(22*I*e) - 20752587082923245568*B*a**21*c**35*f**7*exp(22*I*e))*ex
p(10*I*f*x))*exp(-12*I*e)/(53126622932283508654080*a**24*c**40*f**8), Ne(a
**24*c**40*f**8*exp(12*I*e), 0)), (x*(-(28*A + 7*I*B)/(128*a**3*c**5) + (A
*exp(16*I*e) + 8*A*exp(14*I*e) + 28*A*exp(12*I*e) + 56*A*exp(10*I*e) + 70*
A*exp(8*I*e) + 56*A*exp(6*I*e) + 28*A*exp(4*I*e) + 8*A*exp(2*I*e) + A - I*
B*exp(16*I*e) - 6*I*B*exp(14*I*e) - 14*I*B*exp(12*I*e) - 14*I*B*exp(10*I*
e) + 14*I*B*exp(6*I*e) + 14*I*B*exp(4*I*e) + 6*I*B*exp(2*I*e) + I*B)*exp(-6
*I*e)/(256*a**3*c**5)), True)) + x*(28*A + 7*I*B)/(128*a**3*c**5)
```

3.738.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^5} dx = \text{Exception raised: RuntimeError}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^5,x, al
gorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negati
ve exponent.
```

3.738. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3 (c-ic \tan(e+fx))^5} dx$

3.738.8 Giac [A] (verification not implemented)

Time = 0.89 (sec) , antiderivative size = 269, normalized size of antiderivative = 0.94

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^5} dx =$$

$$\frac{420(-4iA+B)\log(\tan(fx+e)+i)}{a^3c^5} + \frac{420(4iA-B)\log(\tan(fx+e)-i)}{a^3c^5} + \frac{10(-308iA\tan(fx+e)^3+77B\tan(fx+e)^3-1050A\tan(fx+e)^2}{a^3c^5}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^5,x, algorithm="giac")`

output `-1/15360*(420*(-4*I*A + B)*log(tan(f*x + e) + I)/(a^3*c^5) + 420*(4*I*A - B)*log(tan(f*x + e) - I)/(a^3*c^5) + 10*(-308*I*A*tan(f*x + e)^3 + 77*B*tan(f*x + e)^3 - 1050*A*tan(f*x + e)^2 - 285*I*B*tan(f*x + e)^2 + 1212*I*A*tan(f*x + e) - 363*B*tan(f*x + e) + 478*A + 163*I*B)/(a^3*c^5*(tan(f*x + e) - I)^3) + (3836*I*A*tan(f*x + e)^5 - 959*B*tan(f*x + e)^5 - 21280*A*tan(f*x + e)^4 - 5095*I*B*tan(f*x + e)^4 - 47960*I*A*tan(f*x + e)^3 + 10790*B*tan(f*x + e)^3 + 55360*A*tan(f*x + e)^2 + 11230*I*B*tan(f*x + e)^2 + 33260*I*A*tan(f*x + e) - 5435*B*tan(f*x + e) - 8608*A - 667*I*B)/(a^3*c^5*(tan(f*x + e) + I)^5))/f`

3.738.9 Mupad [B] (verification not implemented)

Time = 10.41 (sec) , antiderivative size = 319, normalized size of antiderivative = 1.11

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^5} dx$$

$$= \frac{\frac{3B}{40a^3c^5} + \tan(e + fx)^4 \left(-\frac{7B}{24a^3c^5} + \frac{A7i}{6a^3c^5}\right) + \tan(e + fx)^6 \left(-\frac{7B}{64a^3c^5} + \frac{A7i}{16a^3c^5}\right) + \tan(e + fx)^7 \left(\frac{7A}{32a^3c^5} + \frac{1}{16a^3c^5}\right)}{f \left(\tan(e + fx)^8 + \tan(e + fx)^7 2i + 2\right)}$$

$$+ \frac{7x(4A + B1i)}{128a^3c^5}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i)^5),x)`

output $(\tan(e + f*x)^4*((A*7i)/(6*a^3*c^5) - (7*B)/(24*a^3*c^5)) - \tan(e + f*x)*((61*A)/(160*a^3*c^5) + (B*61i)/(640*a^3*c^5)) + \tan(e + f*x)^6*((A*7i)/(16*a^3*c^5) - (7*B)/(64*a^3*c^5)) + \tan(e + f*x)^7*((7*A)/(32*a^3*c^5) + (B*7i)/(128*a^3*c^5)) + \tan(e + f*x)^5*((35*A)/(96*a^3*c^5) + (B*35i)/(384*a^3*c^5)) + \tan(e + f*x)^2*((A*77i)/(80*a^3*c^5) - (77*B)/(320*a^3*c^5)) - \tan(e + f*x)^3*((49*A)/(480*a^3*c^5) + (B*49i)/(1920*a^3*c^5)) + (A*1i)/(5*a^3*c^5) + (3*B)/(40*a^3*c^5))/(f*(\tan(e + f*x)*2i - 2*\tan(e + f*x)^2 + \tan(e + f*x)^3*6i + \tan(e + f*x)^5*6i + 2*\tan(e + f*x)^6 + \tan(e + f*x)^7*2i + \tan(e + f*x)^8 - 1)) + (7*x*(4*A + B*1i))/(128*a^3*c^5)$

3.738. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^5} dx$

$$3.739 \quad \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^6} dx$$

3.739.1 Optimal result	6777
3.739.2 Mathematica [A] (verified)	6778
3.739.3 Rubi [A] (verified)	6778
3.739.4 Maple [A] (verified)	6780
3.739.5 Fricas [A] (verification not implemented)	6781
3.739.6 Sympy [A] (verification not implemented)	6781
3.739.7 Maxima [F(-2)]	6782
3.739.8 Giac [A] (verification not implemented)	6783
3.739.9 Mupad [B] (verification not implemented)	6783

3.739.1 Optimal result

Integrand size = 41, antiderivative size = 319

$$\begin{aligned} & \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^6} dx \\ &= \frac{7(3A+iB)x}{128a^3c^6} + \frac{A+iB}{384a^3c^6 f(i-\tan(e+fx))^3} - \frac{7iA-5B}{512a^3c^6 f(i-\tan(e+fx))^2} \\ & - \frac{7(2A+iB)}{256a^3c^6 f(i-\tan(e+fx))} + \frac{iA+B}{96a^3c^6 f(i+\tan(e+fx))^6} + \frac{2A-iB}{80a^3c^6 f(i+\tan(e+fx))^5} \\ & - \frac{128a^3c^6 f(i+\tan(e+fx))^4}{5(7iA-B)} - \frac{96a^3c^6 f(i+\tan(e+fx))^3}{7(4A+iB)} \\ & + \frac{512a^3c^6 f(i+\tan(e+fx))^2}{256a^3c^6 f(i+\tan(e+fx))} \end{aligned}$$

```
output 7/128*(3*A+I*B)*x/a^3/c^6+1/384*(A+I*B)/a^3/c^6/f/(I-tan(f*x+e))^3+1/512*(
-7*I*A+5*B)/a^3/c^6/f/(I-tan(f*x+e))^2-7/256*(2*A+I*B)/a^3/c^6/f/(I-tan(f*
x+e))+1/96*(I*A+B)/a^3/c^6/f/(I+tan(f*x+e))^6+1/80*(2*A-I*B)/a^3/c^6/f/(I+
tan(f*x+e))^5+1/128*(-5*I*A-B)/a^3/c^6/f/(I+tan(f*x+e))^4-5/96*A/a^3/c^6/f
/(I+tan(f*x+e))^3+5/512*(7*I*A-B)/a^3/c^6/f/(I+tan(f*x+e))^2+7/256*(4*A+I*
B)/a^3/c^6/f/(I+tan(f*x+e))
```


3.739.2 Mathematica [A] (verified)

Time = 7.20 (sec) , antiderivative size = 278, normalized size of antiderivative = 0.87

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^6} dx$$

$$= \frac{\sec^8(e + fx)(-1271A + 43iB - 8(391A + 37iB) \cos(2(e + fx)) + (734A + 618iB) \cos(4(e + fx)) + 76A \cos(6(e + fx)))}{(15360a^3c^6f(-I + \tan[e + fx])^3(I + \tan[e + fx])^6)}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^6),x]`

output `(Sec[e + f*x]^8*(-1271*A + (43*I)*B - 8*(391*A + (37*I)*B)*Cos[2*(e + f*x)] + (734*A + (618*I)*B)*Cos[4*(e + f*x)] + 76*A*Cos[6*(e + f*x)] + (132*I)*B*Cos[6*(e + f*x)] + 5*A*Cos[8*(e + f*x)] + (15*I)*B*Cos[8*(e + f*x)] + (1890*I)*A*Sin[2*(e + f*x)] - 630*B*Sin[2*(e + f*x)] + 840*((-3*I)*A + B)*ArcTan[Tan[e + f*x]]*Sec[e + f*x]*(Cos[3*(e + f*x)] - I*Sin[3*(e + f*x)]) - (1176*I)*A*Sin[4*(e + f*x)] + 392*B*Sin[4*(e + f*x)] - (174*I)*A*Sin[6*(e + f*x)] + 58*B*Sin[6*(e + f*x)] - (15*I)*A*Sin[8*(e + f*x)] + 5*B*Sin[8*(e + f*x)]))/(15360*a^3*c^6*f*(-I + Tan[e + f*x])^3*(I + Tan[e + f*x])^6)`

3.739.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 248, normalized size of antiderivative = 0.78, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^6} dx$$

$$\downarrow 3042$$

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^6} dx$$

$$\downarrow 4071$$

$$\frac{ac \int \frac{A+B \tan(e+fx)}{a^4 c^7 (1-i \tan(e+fx))^7 (i \tan(e+fx)+1)^4} d \tan(e+fx)}{f}$$

3.739. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3 (c-ic \tan(e+fx))^6} dx$

$$\begin{array}{c}
 \downarrow 27 \\
 \int \frac{A+B \tan(e+fx)}{(1-i \tan(e+fx))^7(i \tan(e+fx)+1)^4} d \tan(e+fx) \\
 a^3 c^6 f \\
 \downarrow 86 \\
 \int \left(\frac{5A}{32(\tan(e+fx)+i)^4} + \frac{7(3A+iB)}{128(\tan^2(e+fx)+1)} - \frac{7(2A+iB)}{256(\tan(e+fx)-i)^2} - \frac{7(4A+iB)}{256(\tan(e+fx)+i)^2} + \frac{i(7A+5iB)}{256(\tan(e+fx)-i)^3} + \frac{5(B-7iA)}{256(\tan(e+fx)+i)^3} \right) \\
 a^3 c^6 f \\
 \downarrow 2009 \\
 \frac{7}{128}(3A+iB) \arctan(\tan(e+fx)) - \frac{7(2A+iB)}{256(-\tan(e+fx)+i)} + \frac{7(4A+iB)}{256(\tan(e+fx)+i)} - \frac{-5B+7iA}{512(-\tan(e+fx)+i)^2} + \frac{5(-B+7iA)}{512(\tan(e+fx)+i)^2} + \\
 a^3 c^6 f
 \end{array}$$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^6), x]`

output `((7*(3*A + I*B)*ArcTan[Tan[e + f*x]])/128 + (A + I*B)/(384*(I - Tan[e + f*x])^3) - ((7*I)*A - 5*B)/(512*(I - Tan[e + f*x])^2) - (7*(2*A + I*B))/(256*(I - Tan[e + f*x])) + (I*A + B)/(96*(I + Tan[e + f*x])^6) + (2*A - I*B)/(80*(I + Tan[e + f*x])^5) - ((5*I)*A + B)/(128*(I + Tan[e + f*x])^4) - (5*A)/(96*(I + Tan[e + f*x])^3) + (5*((7*I)*A - B))/(512*(I + Tan[e + f*x])^2) + (7*(4*A + I*B))/(256*(I + Tan[e + f*x]))) / (a^3*c^6*f)`

3.739.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

3.739. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^6} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.739.4 Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 392, normalized size of antiderivative = 1.23

method	result
norman	$\frac{7(iB+3A)x}{128ac} - \frac{7iA+B}{30acf} + \frac{281(iB+3A)\tan(fx+e)^5}{320acf} + \frac{231(iB+3A)\tan(fx+e)^7}{320acf} + \frac{119(iB+3A)\tan(fx+e)^9}{384acf} + \frac{7(iB+3A)\tan(fx+e)^{11}}{128acf} + \dots$
risch	$-\frac{9i \sin(6fx+6e)B}{1024a^3c^6f} + \frac{21xA}{128a^3c^6} - \frac{e^{12i(fx+e)}B}{6144a^3c^6f} - \frac{9ie^{10i(fx+e)}A}{5120a^3c^6f} - \frac{7e^{10i(fx+e)}B}{5120a^3c^6f} - \frac{117i \cos(4fx+4e)A}{2048a^3c^6f} - \frac{5e^{10i(fx+e)}A}{1024a^3c^6f}$
derivativedivides	$-\frac{A}{384fa^3c^6(-i+\tan(fx+e))^3} + \frac{7A}{128fa^3c^6(-i+\tan(fx+e))} + \frac{5B}{512fa^3c^6(-i+\tan(fx+e))^2} + \frac{7A}{64fa^3c^6(i+\tan(fx+e))}$
default	$-\frac{A}{384fa^3c^6(-i+\tan(fx+e))^3} + \frac{7A}{128fa^3c^6(-i+\tan(fx+e))} + \frac{5B}{512fa^3c^6(-i+\tan(fx+e))^2} + \frac{7A}{64fa^3c^6(i+\tan(fx+e))}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^6,x,method=_RETURNVERBOSE)`

output $(7/128*(3*A+I*B)/a/c*x-1/30*(7*I*A+B)/a/c/f+281/320*(3*A+I*B)/a/c/f*\tan(f*x+e)^5+231/320*(3*A+I*B)/a/c/f*\tan(f*x+e)^7+119/384*(3*A+I*B)/a/c/f*\tan(f*x+e)^9+7/128*(3*A+I*B)/a/c/f*\tan(f*x+e)^{11}+21/64*(3*A+I*B)/a/c*x*\tan(f*x+e)^2+105/128*(3*A+I*B)/a/c*x*\tan(f*x+e)^4+35/32*(3*A+I*B)/a/c*x*\tan(f*x+e)^6+105/128*(3*A+I*B)/a/c*x*\tan(f*x+e)^8+21/64*(3*A+I*B)/a/c*x*\tan(f*x+e)^{10}+7/128*(3*A+I*B)/a/c*x*\tan(f*x+e)^{12}+1/128*(107*A-7*I*B)/a/c/f*\tan(f*x+e)+1/384*(667*A+265*I*B)/a/c/f*\tan(f*x+e)^3+1/10/a/c/f*(I*A+3*B)*\tan(f*x+e)^2)/(1+\tan(f*x+e)^2)^6/a^2/c^5$

3.739. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^6} dx$

3.739.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.58

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^6} dx$$

$$= \frac{(1680(3A + iB)fx e^{(6i fx + 6ie)} - 5(iA + B)e^{(18i fx + 18ie)} - 6(9iA + 7B)e^{(16i fx + 16ie)} - 30(9iA + 5B)e^{(14i fx + 14ie)} - 280(3iA + B)e^{(12i fx + 12ie)} - 210(9iA + B)e^{(10i fx + 10ie)} - 420(9iA - B)e^{(8i fx + 8ie)} - 120(-9iA + 5B)e^{(4i fx + 4ie)} - 15(-9iA + 7B)e^{(2i fx + 2ie)} + 10iA - 10B)e^{(-6i fx - 6ie)}}{(a^3 c^6 f)}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^6,x, algorithm="fricas")`

output `1/30720*(1680*(3*A + I*B)*f*x*e^(6*I*f*x + 6*I*e) - 5*(I*A + B)*e^(18*I*f*x + 18*I*e) - 6*(9*I*A + 7*B)*e^(16*I*f*x + 16*I*e) - 30*(9*I*A + 5*B)*e^(14*I*f*x + 14*I*e) - 280*(3*I*A + B)*e^(12*I*f*x + 12*I*e) - 210*(9*I*A + B)*e^(10*I*f*x + 10*I*e) - 420*(9*I*A - B)*e^(8*I*f*x + 8*I*e) - 120*(-9*I*A + 5*B)*e^(4*I*f*x + 4*I*e) - 15*(-9*I*A + 7*B)*e^(2*I*f*x + 2*I*e) + 10*I*A - 10*B)*e^(-6*I*f*x - 6*I*e)/(a^3*c^6*f)`

3.739.6 Sympy [A] (verification not implemented)

Time = 0.84 (sec) , antiderivative size = 753, normalized size of antiderivative = 2.36

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^6} dx$$

$$= \left\{ \begin{array}{l} ((6800207735332289107722240iAa^{24}c^{48}f^8e^{6ie} - 6800207735332289107722240Ba^{24}c^{48}f^8e^{6ie})e^{-6ifx} + (91802804426985902954250240iAa^{24}c^{48}f^8e^{6ie})e^{-6ifx} \\ x \left(-\frac{21A+7iB}{128a^3c^6} + \frac{(Ae^{18ie}+9Ae^{16ie}+36Ae^{14ie}+84Ae^{12ie}+126Ae^{10ie}+126Ae^{8ie}+84Ae^{6ie}+36Ae^{4ie}+9Ae^{2ie}+A-iBe^{18ie}-7iBe^{16ie}-20iBe^{14ie}-14iBe^{12ie}-9iBe^{10ie}-7iBe^{8ie}-5iBe^{6ie}-3iBe^{4ie}-iBe^{2ie})}{512a^3c^6} \right) \\ + \frac{x(21A + 7iB)}{128a^3c^6} \end{array} \right.$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**3/(c-I*c*tan(f*x+e))**6,x)`

output `Piecewise((((6800207735332289107722240*I*A*a**24*c**48*f**8*exp(6*I*e) - 6800207735332289107722240*B*a**24*c**48*f**8*exp(6*I*e))*exp(-6*I*f*x) + (91802804426985902954250240*I*A*a**24*c**48*f**8*exp(8*I*e) - 71402181220989035631083520*B*a**24*c**48*f**8*exp(8*I*e))*exp(-4*I*f*x) + (734422435415887223634001920*I*A*a**24*c**48*f**8*exp(10*I*e) - 408012464119937346463334400*B*a**24*c**48*f**8*exp(10*I*e))*exp(-2*I*f*x) + (-2570478523955605282719006720*I*A*a**24*c**48*f**8*exp(14*I*e) + 285608724883956142524334080*B*a**24*c**48*f**8*exp(14*I*e))*exp(2*I*f*x) + (-1285239261977802641359503360*I*A*a**24*c**48*f**8*exp(16*I*e) - 142804362441978071262167040*B*a**24*c**48*f**8*exp(16*I*e))*exp(4*I*f*x) + (-571217449767912285048668160*I*A*a**24*c**48*f**8*exp(18*I*e) - 190405816589304095016222720*B*a**24*c**48*f**8*exp(18*I*e))*exp(6*I*f*x) + (-183605608853971805908500480*I*A*a**24*c**48*f**8*exp(20*I*e) - 102003116029984336615833600*B*a**24*c**48*f**8*exp(20*I*e))*exp(8*I*f*x) + (-36721121770794361181700096*I*A*a**24*c**48*f**8*exp(22*I*e) - 28560872488395614252433408*B*a**24*c**48*f**8*exp(22*I*e))*exp(10*I*f*x) + (-3400103867666144553861120*I*A*a**24*c**48*f**8*exp(24*I*e) - 3400103867666144553861120*B*a**24*c**48*f**8*exp(24*I*e))*exp(12*I*f*x))*exp(-12*I*e)/(20890238162940792138922721280*a**27*c**54*f**9), Ne(a**27*c**54*f**9*exp(12*I*e), 0)), (x*(-(21*A + 7*I*B)/(128*a**3*c**6) + (A*exp(18*I*e) + 9*A*exp(16*I*e) + 36*A*exp(14*I*e) + 84*A*exp(12*I*e) + 126*A...`

3.739.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^6} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^6,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: exp: undefined: 0 to a negative exponent.`

3.739.8 Giac [A] (verification not implemented)

Time = 0.95 (sec) , antiderivative size = 295, normalized size of antiderivative = 0.92

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^6} dx$$

$$= \frac{420(3iA - B) \log(\tan(fx + e) + i)}{a^3 c^6} - \frac{420(3iA - B) \log(\tan(fx + e) - i)}{a^3 c^6} - \frac{10(231A \tan(fx + e)^3 + 77iB \tan(fx + e)^3 - 777iA \tan(fx + e)^2 + 273B \tan(fx + e)^2 - 330iB \tan(fx + e) + 340iA - 138B)}{a^3 c^6 (-i \tan(fx + e) - 1)^3} + \frac{-3087iA \tan(fx + e)^6 + 1029B \tan(fx + e)^6 + 20202A \tan(fx + e)^5 + 6594iB \tan(fx + e)^5 + 55755iA \tan(fx + e)^4 - 17685B \tan(fx + e)^4 - 83540A \tan(fx + e)^3 - 25380iB \tan(fx + e)^3 - 72405iA \tan(fx + e)^2 + 20415B \tan(fx + e)^2 + 35106A \tan(fx + e) + 8442iB \tan(fx + e) + 7761iA - 1127B}{a^3 c^6 (\tan(fx + e) + i)^6} / f$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^6,x, algorithm="giac")`

output `1/15360*(420*(3*I*A - B)*log(tan(f*x + e) + I)/(a^3*c^6) - 420*(3*I*A - B)*log(tan(f*x + e) - I)/(a^3*c^6) - 10*(231*A*tan(f*x + e)^3 + 77*I*B*tan(f*x + e)^3 - 777*I*A*tan(f*x + e)^2 + 273*B*tan(f*x + e)^2 - 882*A*tan(f*x + e) - 330*I*B*tan(f*x + e) + 340*I*A - 138*B)/(a^3*c^6*(-I*tan(f*x + e) - 1)^3) + (-3087*I*A*tan(f*x + e)^6 + 1029*B*tan(f*x + e)^6 + 20202*A*tan(f*x + e)^5 + 6594*I*B*tan(f*x + e)^5 + 55755*I*A*tan(f*x + e)^4 - 17685*B*tan(f*x + e)^4 - 83540*A*tan(f*x + e)^3 - 25380*I*B*tan(f*x + e)^3 - 72405*I*A*tan(f*x + e)^2 + 20415*B*tan(f*x + e)^2 + 35106*A*tan(f*x + e) + 8442*I*B*tan(f*x + e) + 7761*I*A - 1127*B)/(a^3*c^6*(tan(f*x + e) + I)^6))/f`

3.739.9 Mupad [B] (verification not implemented)

Time = 10.58 (sec) , antiderivative size = 352, normalized size of antiderivative = 1.10

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^6} dx$$

$$= \frac{\tan(e + fx) \left(-\frac{29B}{640a^3c^6} + \frac{A87i}{640a^3c^6} \right) - \tan(e + fx)^8 \left(\frac{21A}{128a^3c^6} + \frac{B7i}{128a^3c^6} \right) - \tan(e + fx)^7 \left(-\frac{21B}{128a^3c^6} + \frac{A63i}{128a^3c^6} \right) - \tan(e + fx)^6 \left(\frac{7A}{128a^3c^6} + \frac{B7i}{128a^3c^6} \right) - \tan(e + fx)^5 \left(-\frac{7B}{128a^3c^6} + \frac{A7i}{128a^3c^6} \right) - \tan(e + fx)^4 \left(\frac{7A}{128a^3c^6} + \frac{B7i}{128a^3c^6} \right) - \tan(e + fx)^3 \left(-\frac{7B}{128a^3c^6} + \frac{A7i}{128a^3c^6} \right) - \tan(e + fx)^2 \left(\frac{7A}{128a^3c^6} + \frac{B7i}{128a^3c^6} \right) - \tan(e + fx) \left(-\frac{7B}{128a^3c^6} + \frac{A7i}{128a^3c^6} \right) - \frac{7x(3A + B1i)}{128a^3c^6}}{f(-\tan(e + fx))^9}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i)^6),x)`

output $(\tan(e + f*x)*((A*87i)/(640*a^3*c^6) - (29*B)/(640*a^3*c^6)) - \tan(e + f*x)^8*((21*A)/(128*a^3*c^6) + (B*7i)/(128*a^3*c^6)) - \tan(e + f*x)^7*((A*63i)/(128*a^3*c^6) - (21*B)/(128*a^3*c^6)) + \tan(e + f*x)^2*((129*A)/(128*a^3*c^6) + (B*43i)/(128*a^3*c^6)) - \tan(e + f*x)^5*((A*147i)/(128*a^3*c^6) - (49*B)/(128*a^3*c^6)) + \tan(e + f*x)^6*((7*A)/(128*a^3*c^6) + (B*7i)/(384*a^3*c^6)) + \tan(e + f*x)^4*((609*A)/(640*a^3*c^6) + (B*203i)/(640*a^3*c^6)) - \tan(e + f*x)^3*((A*413i)/(640*a^3*c^6) - (413*B)/(1920*a^3*c^6)) + (7*A)/(30*a^3*c^6) - (B*1i)/(30*a^3*c^6))/(f*(3*\tan(e + f*x) + 8*\tan(e + f*x)^3 - \tan(e + f*x)^4*6i + 6*\tan(e + f*x)^5 - \tan(e + f*x)^6*8i - \tan(e + f*x)^8*3i - \tan(e + f*x)^9 + 1i)) + (7*x*(3*A + B*1i))/(128*a^3*c^6)$

3.739. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^6} dx$

3.740 $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx$

3.740.1 Optimal result	6785
3.740.2 Mathematica [A] (verified)	6785
3.740.3 Rubi [A] (verified)	6786
3.740.4 Maple [A] (verified)	6787
3.740.5 Fricas [B] (verification not implemented)	6788
3.740.6 Sympy [F]	6788
3.740.7 Maxima [A] (verification not implemented)	6789
3.740.8 Giac [F]	6789
3.740.9 Mupad [B] (verification not implemented)	6790

3.740.1 Optimal result

Integrand size = 41, antiderivative size = 62

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx = \frac{2a(iA + B)(c - ic \tan(e + fx))^{7/2}}{7f} - \frac{2aB(c - ic \tan(e + fx))^{9/2}}{9cf}$$

```
output 2/7*a*(I*A+B)*(c-I*c*tan(f*x+e))^(7/2)/f-2/9*a*B*(c-I*c*tan(f*x+e))^(9/2)/c/f
```

3.740.2 Mathematica [A] (verified)

Time = 3.57 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.94

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx = \frac{2ac^3(i + \tan(e + fx))^3(9A - 2iB + 7B \tan(e + fx))\sqrt{c - ic \tan(e + fx)}}{63f}$$

```
input Integrate[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2),x]
```


output $(-2*a*c^3*(I + \text{Tan}[e + f*x])^3*(9*A - (2*I)*B + 7*B*\text{Tan}[e + f*x])*Sqrt[c - I*c*\text{Tan}[e + f*x]]/(63*f)$

3.740.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.098$, Rules used = {3042, 4071, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))(c - ictan(e + fx))^{7/2}(A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))(c - ictan(e + fx))^{7/2}(A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int (A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{53} \\ & \frac{ac \int \left(\frac{iB(c - ictan(e + fx))^{7/2}}{c} + (A - iB)(c - ictan(e + fx))^{5/2} \right) d \tan(e + fx)}{f} \\ & \quad \downarrow \text{2009} \\ & \frac{ac \left(\frac{2(B + iA)(c - ictan(e + fx))^{7/2}}{7c} - \frac{2B(c - ictan(e + fx))^{9/2}}{9c^2} \right)}{f} \end{aligned}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])*(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^{(7/2)}, x]$

output $(a*c*((2*(I*A + B)*(c - I*c*\text{Tan}[e + f*x])^{(7/2)})/(7*c) - (2*B*(c - I*c*\text{Tan}[e + f*x])^{(9/2)})/(9*c^2)))/f$

3.740.3.1 Defintions of rubi rules used

- rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int [ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x , Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.740.4 Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.89

method	result
derivativedivides	$\frac{2ia \left(\frac{iB(c-ic \tan(fx+e))^{\frac{9}{2}}}{9} + \frac{(-iBc+cA)(c-ic \tan(fx+e))^{\frac{7}{2}}}{7} \right)}{fc}$
default	$\frac{2ia \left(\frac{iB(c-ic \tan(fx+e))^{\frac{9}{2}}}{9} + \frac{(-iBc+cA)(c-ic \tan(fx+e))^{\frac{7}{2}}}{7} \right)}{fc}$
parts	$\frac{2iAac \left(-\frac{(c-ic \tan(fx+e))^{\frac{5}{2}}}{5} - \frac{2c(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} - 4\sqrt{c-ic \tan(fx+e)} c^2 + 4c^{\frac{5}{2}} \sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}} \right) \right)}{f} +$

```
input int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x,method=_RETURNVERBOSE)
```

```
output 2*I/f*a/c*(1/9*I*B*(c-I*c*tan(f*x+e))^(9/2)+1/7*(-I*B*c+c*A)*(c-I*c*tan(f*x+e))^(7/2))
```

3.740. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx$

3.740.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 108 vs. $2(48) = 96$.

Time = 0.35 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.74

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx =$$

$$\frac{16\sqrt{2}(9(-iA - B)ac^3e^{(2i fx + 2ie)} + (-9iA + 5B)ac^3)\sqrt{\frac{c}{e^{(2i fx + 2ie)} + 1}}}{63(fe^{(8i fx + 8ie)} + 4fe^{(6i fx + 6ie)} + 6fe^{(4i fx + 4ie)} + 4fe^{(2i fx + 2ie)} + f)}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x,
algorithm="fricas")`

output `-16/63*sqrt(2)*(9*(-I*A - B)*a*c^3*e^(2*I*f*x + 2*I*e) + (-9*I*A + 5*B)*a*
c^3)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(f*e^(8*I*f*x + 8*I*e) + 4*f*e^(6*I
*f*x + 6*I*e) + 6*f*e^(4*I*f*x + 4*I*e) + 4*f*e^(2*I*f*x + 2*I*e) + f)`

3.740.6 Sympy [F]

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx = ia \left(\int (-iAc^3 \sqrt{-ic \tan(e + fx) + c}) dx \right.$$

$$+ \int (-2Ac^3 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx)) dx$$

$$+ \int (-2Ac^3 \sqrt{-ic \tan(e + fx) + c} \tan^3(e + fx)) dx$$

$$+ \int (-2Bc^3 \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx)) dx$$

$$+ \int (-2Bc^3 \sqrt{-ic \tan(e + fx) + c} \tan^4(e + fx)) dx$$

$$+ \int iAc^3 \sqrt{-ic \tan(e + fx) + c} \tan^4(e + fx) dx$$

$$+ \int (-iBc^3 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx)) dx$$

$$\left. + \int iBc^3 \sqrt{-ic \tan(e + fx) + c} \tan^5(e + fx) dx \right)$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(7/2),x)`

output `I*a*(Integral(-I*A*c**3*sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-2*A*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(-2*A*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x) + Integral(-2*B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x) + Integral(-2*B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4, x) + Integral(I*A*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4, x) + Integral(-I*B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(I*B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**5, x))`

3.740.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.77

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx = \frac{2i \left(7i (-ictan(fx + e) + c)^{\frac{9}{2}} Ba + 9 (-ictan(fx + e) + c)^{\frac{7}{2}} (A - iB)ac \right)}{63cf}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="maxima")`

output `2/63*I*(7*I*(-I*c*tan(f*x + e) + c)^(9/2)*B*a + 9*(-I*c*tan(f*x + e) + c)^(7/2)*(A - I*B)*a*c)/(c*f)`

3.740.8 Giac [F]

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx = \int (B \tan(fx + e) + A)(ia \tan(fx + e) + a)(-ictan(fx + e) + c)^{\frac{7}{2}} dx$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)*(-I*c*tan(f*x + e) + c)^(7/2), x)`

3.740.9 Mupad [B] (verification not implemented)

Time = 12.35 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.63

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx = \frac{16 a c^3 \sqrt{c + \frac{c(e^{e 2i + f x 2i} - 1) - 1}{e^{e 2i + f x 2i} + 1}} (A 9i - 5 B + A e^{e 2i + f x 2i} 9i + 9 B e^{e 2i + f x 2i})}{63 f (e^{e 2i + f x 2i} + 1)^4}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)*(c - c*tan(e + f*x)*1i)^(7/2),x)`

output `(16*a*c^3*(c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(1/2)*(A*9i - 5*B + A*exp(e*2i + f*x*2i)*9i + 9*B*exp(e*2i + f*x*2i)))/(63*f*(exp(e*2i + f*x*2i) + 1)^4)`

3.741 $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx$

3.741.1 Optimal result	6791
3.741.2 Mathematica [A] (verified)	6791
3.741.3 Rubi [A] (verified)	6792
3.741.4 Maple [A] (verified)	6793
3.741.5 Fricas [A] (verification not implemented)	6794
3.741.6 Sympy [F]	6794
3.741.7 Maxima [A] (verification not implemented)	6795
3.741.8 Giac [F]	6795
3.741.9 Mupad [B] (verification not implemented)	6796

3.741.1 Optimal result

Integrand size = 41, antiderivative size = 62

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx = \frac{2a(iA + B)(c - ictan(e + fx))^{5/2}}{5f} - \frac{2aB(c - ictan(e + fx))^{7/2}}{7cf}$$

```
output 2/5*a*(I*A+B)*(c-I*c*tan(f*x+e))^(5/2)/f-2/7*a*B*(c-I*c*tan(f*x+e))^(7/2)/c/f
```

3.741.2 Mathematica [A] (verified)

Time = 2.23 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.97

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx = \frac{2iac^2(i + \tan(e + fx))^2(7A - 2iB + 5B \tan(e + fx))\sqrt{c - ictan(e + fx)}}{35f}$$

```
input Integrate[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2), x]
```

output $(((-2*I)/35)*a*c^2*(I + \text{Tan}[e + f*x])^2*(7*A - (2*I)*B + 5*B*\text{Tan}[e + f*x])$
 $*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])/f$

3.741.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.098$, Rules used = {3042, 4071, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))(c - ictan(e + fx))^{5/2}(A + B \tan(e + fx)) dx$$

↓ 3042

$$\int (a + ia \tan(e + fx))(c - ictan(e + fx))^{5/2}(A + B \tan(e + fx)) dx$$

↓ 4071

$$\frac{ac \int (A + B \tan(e + fx))(c - ictan(e + fx))^{3/2} d \tan(e + fx)}{f}$$

↓ 53

$$\frac{ac \int \left(\frac{iB(c - ictan(e + fx))^{5/2}}{c} + (A - iB)(c - ictan(e + fx))^{3/2} \right) d \tan(e + fx)}{f}$$

↓ 2009

$$\frac{ac \left(\frac{2(B + iA)(c - ictan(e + fx))^{5/2}}{5c} - \frac{2B(c - ictan(e + fx))^{7/2}}{7c^2} \right)}{f}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])*(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^(5/2), x]$

output $(a*c*((2*(I*A + B)*(c - I*c*\text{Tan}[e + f*x])^(5/2))/(5*c) - (2*B*(c - I*c*\text{Tan}[e + f*x])^(7/2))/(7*c^2)))/f$

3.741. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx$

3.741.3.1 Defintions of rubi rules used

- rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int [ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x , Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.741.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.89

method	result
derivativedivides	$\frac{2ia \left(\frac{iB(c-ic \tan(fx+e))^{\frac{7}{2}}}{7} + \frac{(-iBc+cA)(c-ic \tan(fx+e))^{\frac{5}{2}}}{5} \right)}{fc}$
default	$\frac{2ia \left(\frac{iB(c-ic \tan(fx+e))^{\frac{7}{2}}}{7} + \frac{(-iBc+cA)(c-ic \tan(fx+e))^{\frac{5}{2}}}{5} \right)}{fc}$
parts	$\frac{2iAac \left(-\frac{(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} - 2c\sqrt{c-ic \tan(fx+e)} + 2c^{\frac{3}{2}}\sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(fx+e)}\sqrt{2}}{2\sqrt{c}} \right) \right)}{f} + \frac{a(iA+B) \left(\frac{2(c-ic \tan(fx+e))^{\frac{5}{2}}}{5} \right)}{f}$

input `int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

output `2*I/f*a/c*(1/7*I*B*(c-I*c*tan(f*x+e))^(7/2)+1/5*(-I*B*c+c*A)*(c-I*c*tan(f*x+e))^(5/2))`

3.741. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx$

3.741.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.55

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx =$$

$$\frac{8\sqrt{2}(7(-iA - B)ac^2e^{(2ifx+2ie)} + (-7iA + 3B)ac^2)\sqrt{\frac{c}{e^{(2ifx+2ie)}+1}}}{35(fe^{(6ifx+6ie)} + 3fe^{(4ifx+4ie)} + 3fe^{(2ifx+2ie)} + f)}$$

```
input integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x,
algorithm="fricas")
```

```
output -8/35*sqrt(2)*(7*(-I*A - B)*a*c^2*e^(2*I*f*x + 2*I*e) + (-7*I*A + 3*B)*a*c
^2)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*
f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)
```

3.741.6 Sympy [F]

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx = ia \left(\int (-iAc^2 \sqrt{-ictan(e + fx) + c}) dx \right.$$

$$+ \int (-Ac^2 \sqrt{-ictan(e + fx) + c} \tan(e + fx)) dx$$

$$+ \int (-Ac^2 \sqrt{-ictan(e + fx) + c} \tan^3(e + fx)) dx$$

$$+ \int (-Bc^2 \sqrt{-ictan(e + fx) + c} \tan^2(e + fx)) dx$$

$$+ \int (-Bc^2 \sqrt{-ictan(e + fx) + c} \tan^4(e + fx)) dx$$

$$+ \int (-iAc^2 \sqrt{-ictan(e + fx) + c} \tan^2(e + fx)) dx$$

$$+ \int (-iBc^2 \sqrt{-ictan(e + fx) + c} \tan(e + fx)) dx$$

$$\left. + \int (-iBc^2 \sqrt{-ictan(e + fx) + c} \tan^3(e + fx)) dx \right)$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(5/2),x)`

output `I*a*(Integral(-I*A*c**2*sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-A*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(-A*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x) + Integral(-B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x) + Integral(-B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4, x) + Integral(-I*A*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x) + Integral(-I*B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(-I*B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x))`

3.741.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.77

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx = \frac{2i \left(5i (-ictan(fx + e) + c)^{7/2} Ba + 7(-ictan(fx + e) + c)^{5/2} (A - iB)ac \right)}{35cf}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `2/35*I*(5*I*(-I*c*tan(f*x + e) + c)^(7/2)*B*a + 7*(-I*c*tan(f*x + e) + c)^(5/2)*(A - I*B)*a*c)/(c*f)`

3.741.8 Giac [F]

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx) + fx)^{5/2} dx = \int (B \tan(fx + e) + A)(ia \tan(fx + e) + a)(-ictan(fx + e) + c)^{5/2} dx$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)*(-I*c*tan(f*x + e) + c)^(5/2), x)`

3.741.9 Mupad [B] (verification not implemented)

Time = 13.27 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.63

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2} dx = \frac{8ac^2 \sqrt{c + \frac{c(e^{2i+fx2i} - 1) - 1}{e^{2i+fx2i} + 1}} (A7i - 3B + Ae^{2i+fx2i}7i + 7Be^{2i+fx2i})}{35f(e^{2i+fx2i} + 1)^3}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)*(c - c*tan(e + f*x)*1i)^(5/2),x)`

output `(8*a*c^2*(c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(1/2)*(A*7i - 3*B + A*exp(e*2i + f*x*2i)*7i + 7*B*exp(e*2i + f*x*2i)))/(35*f*(exp(e*2i + f*x*2i) + 1)^3)`

3.742 $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx$

3.742.1 Optimal result	6797
3.742.2 Mathematica [A] (verified)	6797
3.742.3 Rubi [A] (verified)	6798
3.742.4 Maple [A] (verified)	6799
3.742.5 Fricas [A] (verification not implemented)	6800
3.742.6 Sympy [F]	6800
3.742.7 Maxima [A] (verification not implemented)	6801
3.742.8 Giac [F]	6801
3.742.9 Mupad [B] (verification not implemented)	6801

3.742.1 Optimal result

Integrand size = 41, antiderivative size = 62

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx = \frac{2a(iA + B)(c - ic \tan(e + fx))^{3/2}}{3f} - \frac{2aB(c - ic \tan(e + fx))^{5/2}}{5cf}$$

```
output 2/3*a*(I*A+B)*(c-I*c*tan(f*x+e))^(3/2)/f-2/5*a*B*(c-I*c*tan(f*x+e))^(5/2)/c/f
```

3.742.2 Mathematica [A] (verified)

Time = 1.55 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.87

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx = \frac{2ac(i + \tan(e + fx))(5A - 2iB + 3B \tan(e + fx))\sqrt{c - ic \tan(e + fx)}}{15f}$$

```
input Integrate[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2), x]
```

output $(2*a*c*(I + \tan[e + f*x])*(5*A - (2*I)*B + 3*B*\tan[e + f*x])*Sqrt[c - I*c*\tan[e + f*x]]/(15*f)$

3.742.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.098$, Rules used = {3042, 4071, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))(c - ictan(e + fx))^{3/2}(A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))(c - ictan(e + fx))^{3/2}(A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{53} \\ & \frac{ac \int \left(\frac{iB(c - ictan(e + fx))^{3/2}}{c} + (A - iB) \sqrt{c - ictan(e + fx)} \right) d \tan(e + fx)}{f} \\ & \quad \downarrow \text{2009} \\ & \frac{ac \left(\frac{2(B + iA)(c - ictan(e + fx))^{3/2}}{3c} - \frac{2B(c - ictan(e + fx))^{5/2}}{5c^2} \right)}{f} \end{aligned}$$

input $\text{Int}[(a + I*a*\tan[e + f*x])*(A + B*\tan[e + f*x])*(c - I*c*\tan[e + f*x])^(3/2), x]$

output $(a*c*((2*(I*A + B)*(c - I*c*\tan[e + f*x])^(3/2))/(3*c) - (2*B*(c - I*c*\tan[e + f*x])^(5/2))/(5*c^2)))/f$

3.742. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^{3/2} dx$

3.742.3.1 Defintions of rubi rules used

- rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int [ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x , Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.742.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.89

method	result
derivativedivides	$\frac{2ia \left(\frac{iB(c-ic \tan(fx+e))^{\frac{5}{2}}}{5} + \frac{(-iBc+cA)(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} \right)}{fc}$
default	$\frac{2ia \left(\frac{iB(c-ic \tan(fx+e))^{\frac{5}{2}}}{5} + \frac{(-iBc+cA)(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} \right)}{fc}$
parts	$\frac{2iAac \left(-\sqrt{c-ic \tan(fx+e)} + \sqrt{c} \sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}} \right) \right)}{f} + \frac{a(iA+B) \left(\frac{2(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} + 2c\sqrt{c-ic \tan(fx+e)} \right)}{f}$

```
input int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
```

```
output 2*I/f*a/c*(1/5*I*B*(c-I*c*tan(f*x+e))^(5/2)+1/3*(-I*B*c+c*A)*(c-I*c*tan(f*x+e))^(3/2))
```

3.742. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx$

3.742.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.26

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^{3/2} dx = \frac{4\sqrt{2}(5(-iA - B)ace^{(2i fx + 2i e)} + (-5iA + B)ac)\sqrt{\frac{c}{e^{(2i fx + 2i e)} + 1}}}{15(fe^{(4i fx + 4i e)} + 2fe^{(2i fx + 2i e)} + f)}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x,
algorithm="fricas")`

output `-4/15*sqrt(2)*(5*(-I*A - B)*a*c*e^(2*I*f*x + 2*I*e) + (-5*I*A + B)*a*c)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(f*e^(4*I*f*x + 4*I*e) + 2*f*e^(2*I*f*x + 2*I*e) + f)`

3.742.6 Sympy [F]

$$\begin{aligned} & \int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ictan(e + fx))^{3/2} dx = ia \left(\int (-iAc\sqrt{-ictan(e + fx) + c}) dx \right. \\ & + \int (-iAc\sqrt{-ictan(e + fx) + c}\tan^2(e + fx)) dx \\ & + \int (-iBc\sqrt{-ictan(e + fx) + c}\tan(e + fx)) dx \\ & \left. + \int (-iBc\sqrt{-ictan(e + fx) + c}\tan^3(e + fx)) dx \right) \end{aligned}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(3/2),x)`

output `I*a*(Integral(-I*A*c*sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-I*A*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x) + Integral(-I*B*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(-I*B*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x))`

3.742.7 Maxima [A] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.77

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx = \frac{2i \left(3i (-ic \tan(fx + e) + c)^{5/2} Ba + 5 (-ic \tan(fx + e) + c)^{3/2} (A - iB)ac \right)}{15cf}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x,
algorithm="maxima")`

output `2/15*I*(3*I*(-I*c*tan(f*x + e) + c)^(5/2)*B*a + 5*(-I*c*tan(f*x + e) + c)^(3/2)*(A - I*B)*a*c)/(c*f)`

3.742.8 Giac [F]

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx = \int (B \tan(fx + e) + A)(ia \tan(fx + e) + a)(-ic \tan(fx + e) + c)^{3/2} dx$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x,
algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)*(-I*c*tan(f*x + e) + c)^(3/2), x)`

3.742.9 Mupad [B] (verification not implemented)

Time = 10.87 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.60

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx = \frac{4ac \sqrt{c + \frac{c(e^{e^{2i+fx^{2i}} - 1} - i)}{e^{e^{2i+fx^{2i}} + 1}}}}{15f(e^{e^{2i+fx^{2i}} + 1})^2} (A5i - B + Ae^{e^{2i+fx^{2i}}}5i + 5Be^{e^{2i+fx^{2i}}})$$

3.742. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)*(c - c*tan(e + f*x)*1i)^(3/2),x)`

output `(4*a*c*(c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(1/2)*(A*5i - B + A*exp(e*2i + f*x*2i)*5i + 5*B*exp(e*2i + f*x*2i)))/(15*f*(exp(e*2i + f*x*2i) + 1)^2)`

3.743 $\int (a+ia \tan(e+fx))(A+B \tan(e+fx))\sqrt{c-ictan(e+fx)}$

3.743.1 Optimal result	6803
3.743.2 Mathematica [A] (verified)	6803
3.743.3 Rubi [A] (verified)	6804
3.743.4 Maple [A] (verified)	6805
3.743.5 Fricas [A] (verification not implemented)	6806
3.743.6 Sympy [F]	6806
3.743.7 Maxima [A] (verification not implemented)	6807
3.743.8 Giac [F]	6807
3.743.9 Mupad [B] (verification not implemented)	6807

3.743.1 Optimal result

Integrand size = 41, antiderivative size = 60

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)} dx$$

$$= \frac{2a(iA + B)\sqrt{c - ictan(e + fx)}}{f} - \frac{2aB(c - ictan(e + fx))^{3/2}}{3cf}$$

output `2*a*(I*A+B)*(c-I*c*tan(f*x+e))^(1/2)/f-2/3*a*B*(c-I*c*tan(f*x+e))^(3/2)/c/f`

3.743.2 Mathematica [A] (verified)

Time = 1.18 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.75

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)} dx$$

$$= \frac{2a(3iA + 2B + iB \tan(e + fx))\sqrt{c - ictan(e + fx)}}{3f}$$

input `Integrate[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]],x]`

output `(2*a*((3*I)*A + 2*B + I*B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])/(3*f)`

3.743.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.098$, Rules used = {3042, 4071, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} (A + B \tan(e + fx)) dx$$

$$\downarrow \text{3042}$$

$$\int (a + ia \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} (A + B \tan(e + fx)) dx$$

$$\downarrow \text{4071}$$

$$\frac{ac \int \frac{A + B \tan(e + fx)}{\sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{f}$$

$$\downarrow \text{53}$$

$$\frac{ac \int \left(\frac{A - iB}{\sqrt{c - ic \tan(e + fx)}} + \frac{iB \sqrt{c - ic \tan(e + fx)}}{c} \right) d \tan(e + fx)}{f}$$

$$\downarrow \text{2009}$$

$$\frac{ac \left(\frac{2(B + iA) \sqrt{c - ic \tan(e + fx)}}{c} - \frac{2B(c - ic \tan(e + fx))^{3/2}}{3c^2} \right)}{f}$$

input `Int[(a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]],x]`

output `(a*c*((2*(I*A + B)*Sqrt[c - I*c*Tan[e + f*x]])/c - (2*B*(c - I*c*Tan[e + f*x])^(3/2))/(3*c^2)))/f`

3.743.3.1 Defintions of rubi rules used

```
rule 53 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && IGtQ[m, 0] && ( !IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0])
|| LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.743.4 Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.10

method	result
derivativedivides	$\frac{2ia \left(\frac{iB(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} - i\sqrt{c-ic \tan(fx+e)} Bc + \sqrt{c-ic \tan(fx+e)} cA \right)}{fc}$
default	$\frac{2ia \left(\frac{iB(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} - i\sqrt{c-ic \tan(fx+e)} Bc + \sqrt{c-ic \tan(fx+e)} cA \right)}{fc}$
parts	$\frac{a(iA+B) \left(2\sqrt{c-ic \tan(fx+e)} - \sqrt{c}\sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(fx+e)}\sqrt{2}}{2\sqrt{c}} \right) \right)}{f} + \frac{iAa\sqrt{c}\sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(fx+e)}\sqrt{2}}{2\sqrt{c}} \right)}{f}$

```
input int((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))*(A+B*tan(f*x+e)),x,method=
_RETURNVERBOSE)
```

```
output 2*I/f*a/c*(1/3*I*B*(c-I*c*tan(f*x+e))^(3/2)-I*(c-I*c*tan(f*x+e))^(1/2)*B*c
+(c-I*c*tan(f*x+e))^(1/2)*c*A)
```

3.743. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx))\sqrt{c - ic \tan(e + fx)} dx$

3.743.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.10

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))\sqrt{c - ic \tan(e + fx)} dx$$

$$= -\frac{2\sqrt{2}(3(-iA - B)ae^{(2i fx + 2i e)} + (-3iA - B)a)\sqrt{\frac{c}{e^{(2i fx + 2i e)} + 1}}}{3(fe^{(2i fx + 2i e)} + f)}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))*(A+B*tan(f*x+e)),x,
algorithm="fricas")`

output `-2/3*sqrt(2)*(3*(-I*A - B)*a*e^(2*I*f*x + 2*I*e) + (-3*I*A - B)*a)*sqrt(c/
(e^(2*I*f*x + 2*I*e) + 1))/(f*e^(2*I*f*x + 2*I*e) + f)`

3.743.6 Sympy [F]

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx))\sqrt{c - ic \tan(e + fx)} dx$$

$$= ia \left(\int (-iA\sqrt{-ic \tan(e + fx) + c}) dx + \int A\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) dx \right.$$

$$\quad \left. + \int B\sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx) dx \right.$$

$$\quad \left. + \int (-iB\sqrt{-ic \tan(e + fx) + c} \tan(e + fx)) dx \right)$$

input `integrate((c-I*c*tan(f*x+e))**(1/2)*(a+I*a*tan(f*x+e))*(A+B*tan(f*x+e)),x)`

output `I*a*(Integral(-I*A*sqrt(-I*c*tan(e + f*x) + c), x) + Integral(A*sqrt(-I*c*
tan(e + f*x) + c)*tan(e + f*x), x) + Integral(B*sqrt(-I*c*tan(e + f*x) + c
) *tan(e + f*x)**2, x) + Integral(-I*B*sqrt(-I*c*tan(e + f*x) + c)*tan(e +
f*x), x))`

3.743.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.80

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx$$

$$= \frac{2i \left(i(-ic \tan(fx + e) + c)^{\frac{3}{2}} Ba + 3 \sqrt{-ic \tan(fx + e) + c} (A - iB) ac \right)}{3cf}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))*(A+B*tan(f*x+e)),x,
algorithm="maxima")`

output `2/3*I*(I*(-I*c*tan(f*x + e) + c)^(3/2)*B*a + 3*sqrt(-I*c*tan(f*x + e) + c)
*(A - I*B)*a*c)/(c*f)`

3.743.8 Giac [F]

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx$$

$$= \int (B \tan(fx + e) + A)(ia \tan(fx + e) + a) \sqrt{-ic \tan(fx + e) + c} dx$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))*(A+B*tan(f*x+e)),x,
algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)*sqrt(-I*c*tan(f*x +
e) + c), x)`

3.743.9 Mupad [B] (verification not implemented)

Time = 0.57 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.70

$$\int (a + ia \tan(e + fx))(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx$$

$$= \frac{a \sqrt{-\frac{c(-2 \cos(e+fx)^2 + \sin(2e+2fx) 1i)}{2 \cos(e+fx)^2}} (A 3i + 2 B + A (2 \cos(e + fx)^2 - 1) 3i + 2 B (2 \cos(e + fx)^2 - 1) + \dots}{3 f \cos(e + fx)^2}$$

3.743. $\int (a + ia \tan(e + fx))(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)*(c - c*tan(e + f*x)*1i)^(1/2),x)`

output `(a*(-(c*(sin(2*e + 2*f*x)*1i - 2*cos(e + f*x)^2))/(2*cos(e + f*x)^2))^(1/2)*(A*3i + 2*B + A*(2*cos(e + f*x)^2 - 1)*3i + 2*B*(2*cos(e + f*x)^2 - 1) + B*sin(2*e + 2*f*x)*1i))/(3*f*cos(e + f*x)^2)`

3.744 $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$

3.744.1 Optimal result 6809
 3.744.2 Mathematica [A] (verified) 6809
 3.744.3 Rubi [A] (verified) 6810
 3.744.4 Maple [A] (verified) 6811
 3.744.5 Fricas [A] (verification not implemented) 6812
 3.744.6 Sympy [F] 6812
 3.744.7 Maxima [A] (verification not implemented) 6813
 3.744.8 Giac [F] 6813
 3.744.9 Mupad [B] (verification not implemented) 6813

3.744.1 Optimal result

Integrand size = 41, antiderivative size = 58

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx$$

$$= -\frac{2a(iA + B)}{f\sqrt{c - ic \tan(e + fx)}} - \frac{2aB\sqrt{c - ic \tan(e + fx)}}{cf}$$

output `-2*a*(I*A+B)/f/(c-I*c*tan(f*x+e))^(1/2)-2*a*B*(c-I*c*tan(f*x+e))^(1/2)/c/f`

3.744.2 Mathematica [A] (verified)

Time = 1.59 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.72

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx = \frac{2ia(-A + 2iB + B \tan(e + fx))}{f\sqrt{c - ic \tan(e + fx)}}$$

input `Integrate[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/Sqrt[c - I*c*Tan[e + f*x]],x]`

output `((2*I)*a*(-A + (2*I)*B + B*Tan[e + f*x]))/(f*Sqrt[c - I*c*Tan[e + f*x]])`

3.744.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.098$, Rules used = {3042, 4071, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A+B \tan(e+fx)}{(c-ic \tan(e+fx))^{3/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{53} \\
 & \frac{ac \int \left(\frac{A-iB}{(c-ic \tan(e+fx))^{3/2}} + \frac{iB}{c\sqrt{c-ic \tan(e+fx)}} \right) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{ac \left(-\frac{2(B+iA)}{c\sqrt{c-ic \tan(e+fx)}} - \frac{2B\sqrt{c-ic \tan(e+fx)}}{c^2} \right)}{f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/Sqrt[c - I*c*Tan[e + f*x]],x]`

output `(a*c*((-2*(I*A + B))/(c*Sqrt[c - I*c*Tan[e + f*x]]) - (2*B*Sqrt[c - I*c*Tan[e + f*x]])/c^2))/f`

3.744.3.1 Defintions of rubi rules used

- rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int [ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x , Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.744.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.91

method	result
derivativedivides	$\frac{2ia \left(i\sqrt{c-ic \tan(fx+e)} B - \frac{c(-iB+A)}{\sqrt{c-ic \tan(fx+e)}} \right)}{fc}$
default	$\frac{2ia \left(i\sqrt{c-ic \tan(fx+e)} B - \frac{c(-iB+A)}{\sqrt{c-ic \tan(fx+e)}} \right)}{fc}$
parts	$\frac{2iAac \left(-\frac{1}{2c\sqrt{c-ic \tan(fx+e)}} + \frac{\sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}} \right)}{4c^{\frac{3}{2}}} \right)}{f} + \frac{a(iA+B) \left(-\frac{1}{\sqrt{c-ic \tan(fx+e)}} - \frac{\sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}} \right)}{2\sqrt{c}} \right)}{f}$

```
input int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x,method=
_RETURNVERBOSE)
```

```
output 2*I/f*a/c*(I*(c-I*c*tan(f*x+e))^(1/2)*B-c*(A-I*B)/(c-I*c*tan(f*x+e))^(1/2)
)
```

3.744.
$$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$$

3.744.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.95

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{\sqrt{c - ict \tan(e + fx)}} dx$$

$$= \frac{\sqrt{2}((-iA - B)ae^{(2i fx + 2ie)} + (-iA - 3B)a)\sqrt{\frac{c}{e^{(2i fx + 2ie)} + 1}}}{cf}$$

```
input integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x,
algorithm="fricas")
```

```
output sqrt(2)*((-I*A - B)*a*e^(2*I*f*x + 2*I*e) + (-I*A - 3*B)*a)*sqrt(c/(e^(2*I
*f*x + 2*I*e) + 1))/(c*f)
```

3.744.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{\sqrt{c - ict \tan(e + fx)}} dx$$

$$= ia \left(\int \left(-\frac{iA}{\sqrt{-ict \tan(e + fx) + c}} \right) dx + \int \frac{A \tan(e + fx)}{\sqrt{-ict \tan(e + fx) + c}} dx \right.$$

$$\left. + \int \frac{B \tan^2(e + fx)}{\sqrt{-ict \tan(e + fx) + c}} dx + \int \left(-\frac{iB \tan(e + fx)}{\sqrt{-ict \tan(e + fx) + c}} \right) dx \right)$$

```
input integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(1/2),x)
```

```
output I*a*(Integral(-I*A/sqrt(-I*c*tan(e + f*x) + c), x) + Integral(A*tan(e + f*
x)/sqrt(-I*c*tan(e + f*x) + c), x) + Integral(B*tan(e + f*x)**2/sqrt(-I*c*
tan(e + f*x) + c), x) + Integral(-I*B*tan(e + f*x)/sqrt(-I*c*tan(e + f*x)
+ c), x))
```

3.744.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.83

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx$$

$$= \frac{2i \left(i \sqrt{-ic \tan(fx + e) + c} Ba - \frac{(A - iB)ac}{\sqrt{-ic \tan(fx + e) + c}} \right)}{cf}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x,
algorithm="maxima")`

output `2*I*(I*sqrt(-I*c*tan(f*x + e) + c)*B*a - (A - I*B)*a*c/sqrt(-I*c*tan(f*x +
e) + c))/(c*f)`

3.744.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx$$

$$= \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)}{\sqrt{-ic \tan(fx + e) + c}} dx$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x,
algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)/sqrt(-I*c*tan(f*x +
e) + c), x)`

3.744.9 Mupad [B] (verification not implemented)

Time = 8.43 (sec) , antiderivative size = 164, normalized size of antiderivative = 2.83

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx =$$

$$\frac{a \sqrt{\frac{2c}{e^{e^{2i+fx} 2i+1}}} \left(A \operatorname{li} + 3B + A \left(\frac{e^{-e^{2i-fx} 2i}}{2} + \frac{e^{e^{2i+fx} 2i}}{2} \right) \operatorname{li} - A \left(\frac{e^{-e^{2i-fx} 2i}}{2} \operatorname{li} - \frac{e^{e^{2i+fx} 2i}}{2} \operatorname{li} \right) + B \left(\frac{e^{-e^{2i-fx} 2i}}{2} \right)}{cf}$$

3.744. $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i))/(c - c*tan(e + f*x)*1i)^(1/2),x)`

output `-(a*((2*c)/(exp(e*2i + f*x*2i) + 1))^(1/2)*(A*1i + 3*B + A*(exp(- e*2i - f*x*2i)/2 + exp(e*2i + f*x*2i)/2)*1i - A*((exp(- e*2i - f*x*2i)*1i)/2 - (exp(e*2i + f*x*2i)*1i)/2) + B*(exp(- e*2i - f*x*2i)/2 + exp(e*2i + f*x*2i)/2) + B*((exp(- e*2i - f*x*2i)*1i)/2 - (exp(e*2i + f*x*2i)*1i)/2)*1i))/(c*f)`

3.745
$$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$$

3.745.1 Optimal result 6815
 3.745.2 Mathematica [A] (verified) 6815
 3.745.3 Rubi [A] (verified) 6816
 3.745.4 Maple [A] (verified) 6817
 3.745.5 Fricas [A] (verification not implemented) 6818
 3.745.6 Sympy [F] 6818
 3.745.7 Maxima [A] (verification not implemented) 6819
 3.745.8 Giac [F] 6819
 3.745.9 Mupad [B] (verification not implemented) 6819

3.745.1 Optimal result

Integrand size = 41, antiderivative size = 60

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = -\frac{2a(iA + B)}{3f(c - ic \tan(e + fx))^{3/2}} + \frac{2aB}{cf\sqrt{c - ic \tan(e + fx)}}$$

output `2*a*B/c/f/(c-I*c*tan(f*x+e))^(1/2)-2/3*a*(I*A+B)/f/(c-I*c*tan(f*x+e))^(3/2)`

3.745.2 Mathematica [A] (verified)

Time = 2.55 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.95

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = -\frac{2a\left(\frac{iA+B}{(c-ic \tan(e+fx))^{3/2}} - \frac{3B}{c\sqrt{c-ic \tan(e+fx)}}\right)}{3f}$$

input `Integrate[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(3/2), x]`

output `(-2*a*((I*A + B)/(c - I*c*Tan[e + f*x])^(3/2) - (3*B)/(c*Sqrt[c - I*c*Tan[e + f*x]])))/(3*f)`

3.745.
$$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$$

3.745.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.098$, Rules used = {3042, 4071, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A+B \tan(e+fx)}{(c-ic \tan(e+fx))^{5/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{53} \\
 & \frac{ac \int \left(\frac{A-iB}{(c-ic \tan(e+fx))^{5/2}} + \frac{iB}{c(c-ic \tan(e+fx))^{3/2}} \right) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{ac \left(\frac{2B}{c^2 \sqrt{c-ic \tan(e+fx)}} - \frac{2(B+iA)}{3c(c-ic \tan(e+fx))^{3/2}} \right)}{f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(a*c*((-2*(I*A + B))/(3*c*(c - I*c*Tan[e + f*x])^(3/2)) + (2*B)/(c^2*sqrt[c - I*c*Tan[e + f*x]])))/f`

3.745.3.1 Defintions of rubi rules used

```
rule 53 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && IGtQ[m, 0] && ( !IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0])
|| LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.745.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.88

method	result
derivativedivides	$\frac{2ia \left(-\frac{c(-iB+A)}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{iB}{\sqrt{c-ic \tan(fx+e)}} \right)}{fc}$
default	$\frac{2ia \left(-\frac{c(-iB+A)}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{iB}{\sqrt{c-ic \tan(fx+e)}} \right)}{fc}$
risch	$-\frac{a(iAe^{2i(fx+e)}+Be^{2i(fx+e)}+iA-5B)\sqrt{2}}{6c\sqrt{\frac{c}{e^{2i(fx+e)}+1}}}f$
parts	$\frac{2iAac \left(-\frac{1}{4c^2\sqrt{c-ic \tan(fx+e)}} - \frac{1}{6c(c-ic \tan(fx+e))^{\frac{3}{2}}} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)}\sqrt{2}}{2\sqrt{c}}\right)}{8c^{\frac{5}{2}}} \right)}{f} + \frac{a(iA+B) \left(-\frac{1}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} \right)}{f}$

```
input int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x,method=
_RETURNVERBOSE)
```

3.745. $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$

output $2*I/f*a/c*(-1/3*c*(A-I*B)/(c-I*c*\tan(f*x+e))^(3/2)-I*B/(c-I*c*\tan(f*x+e))^(1/2))$

3.745.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.25

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \frac{\sqrt{2}((-iA - B)ae^{(4i fx + 4i e)} - 2(iA - 2B)ae^{(2i fx + 2i e)} + 2iAe) + (-iA + 5B)a\sqrt{c/(e^{(2i fx + 2i e)} + 1)}}{6c^2 f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output $1/6*\sqrt{2}*((-I*A - B)*a*e^{(4*I*f*x + 4*I*e)} - 2*(I*A - 2*B)*a*e^{(2*I*f*x + 2*I*e)} + (-I*A + 5*B)*a)*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)}/(c^2*f)$

3.745.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = ia \left(\int \left(-\frac{iA}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} \right. \right. \\ \left. \left. + \int \frac{A \tan(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} dx \right. \right. \\ \left. \left. + \int \frac{B \tan^2(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} dx \right. \right. \\ \left. \left. + \int \left(-\frac{iB \tan(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} \right) dx \right)$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(3/2),x)`

output `I*a*(Integral(-I*A/(-I*c*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(A*tan(e + f*x)/(-I*c*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(B*tan(e + f*x)**2/(-I*c*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(-I*B*tan(e + f*x)/(-I*c*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x)`

3.745. $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$

3.745.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.75

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx =$$

$$\frac{2i(3i(-ic \tan(fx + e) + c)Ba + (A - iB)ac)}{3(-ic \tan(fx + e) + c)^{3/2}cf}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x,
algorithm="maxima")`

output `-2/3*I*(3*I*(-I*c*tan(f*x + e) + c)*B*a + (A - I*B)*a*c)/((-I*c*tan(f*x +
e) + c)^(3/2)*c*f)`

3.745.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)}{(-ic \tan(fx + e) + c)^{3/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x,
algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)/(-I*c*tan(f*x + e) +
c)^(3/2), x)`

3.745.9 Mupad [B] (verification not implemented)

Time = 9.06 (sec) , antiderivative size = 170, normalized size of antiderivative = 2.83

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx =$$

$$a \sqrt{-\frac{c(-2 \cos(e+fx)^2 + \sin(2e+2fx) i)}{2 \cos(e+fx)^2}} (B(2 \cos(2e + 2fx)^2 - 1) - 4B(2 \cos(e + fx)^2 - 1) - 2A \sin(2e + 2fx))$$

3.745. $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i))/(c - c*tan(e + f*x)*1i)
^(3/2),x)`

output `-(a*(-(c*(sin(2*e + 2*f*x)*1i - 2*cos(e + f*x)^2))/(2*cos(e + f*x)^2))^(1/
2)*(A*1i - 5*B + A*(2*cos(e + f*x)^2 - 1)*2i - 4*B*(2*cos(e + f*x)^2 - 1)
- 2*A*sin(2*e + 2*f*x) - A*sin(4*e + 4*f*x) - B*sin(2*e + 2*f*x)*4i + B*si
n(4*e + 4*f*x)*1i + A*(2*cos(2*e + 2*f*x)^2 - 1)*1i + B*(2*cos(2*e + 2*f*x
)^2 - 1)))/(6*c^2*f)`

3.746
$$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{5/2}} dx$$

3.746.1 Optimal result 6821
 3.746.2 Mathematica [A] (verified) 6821
 3.746.3 Rubi [A] (verified) 6822
 3.746.4 Maple [A] (verified) 6823
 3.746.5 Fricas [A] (verification not implemented) 6824
 3.746.6 Sympy [F] 6824
 3.746.7 Maxima [A] (verification not implemented) 6825
 3.746.8 Giac [F] 6825
 3.746.9 Mupad [B] (verification not implemented) 6826

3.746.1 Optimal result

Integrand size = 41, antiderivative size = 62

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{5/2}} dx = -\frac{2a(iA + B)}{5f(c - ict \tan(e + fx))^{5/2}} + \frac{2aB}{3cf(c - ict \tan(e + fx))^{3/2}}$$

output `-2/5*a*(I*A+B)/f/(c-I*c*tan(f*x+e))^(5/2)+2/3*a*B/c/f/(c-I*c*tan(f*x+e))^(3/2)`

3.746.2 Mathematica [A] (verified)

Time = 3.72 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.94

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{5/2}} dx = -\frac{2a\left(\frac{3(iA+B)}{(c-ict \tan(e+fx))^{5/2}} - \frac{5B}{c(c-ict \tan(e+fx))^{3/2}}\right)}{15f}$$

input `Integrate[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(5/2), x]`

output `(-2*a*((3*(I*A + B))/(c - I*c*Tan[e + f*x])^(5/2) - (5*B)/(c*(c - I*c*Tan[e + f*x])^(3/2))))/(15*f)`

3.746.
$$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{5/2}} dx$$

3.746.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.098$, Rules used = {3042, 4071, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A+B \tan(e+fx)}{(c-ic \tan(e+fx))^{7/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{53} \\
 & \frac{ac \int \left(\frac{A-iB}{(c-ic \tan(e+fx))^{7/2}} + \frac{iB}{c(c-ic \tan(e+fx))^{5/2}} \right) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{ac \left(\frac{2B}{3c^2(c-ic \tan(e+fx))^{3/2}} - \frac{2(B+iA)}{5c(c-ic \tan(e+fx))^{5/2}} \right)}{f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(5/2),x]`

output `(a*c*((-2*(I*A + B))/(5*c*(c - I*c*Tan[e + f*x])^(5/2)) + (2*B)/(3*c^2*(c - I*c*Tan[e + f*x])^(3/2))))/f`

3.746.3.1 Defintions of rubi rules used

- rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int [ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x , Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.746.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.85

method	result
derivativedivides	$\frac{2ia \left(-\frac{iB}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{c(-iB+A)}{5(c-ic \tan(fx+e))^{\frac{5}{2}}} \right)}{fc}$
default	$\frac{2ia \left(-\frac{iB}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{c(-iB+A)}{5(c-ic \tan(fx+e))^{\frac{5}{2}}} \right)}{fc}$
risch	$-\frac{a(3iA e^{4i(fx+e)} + 3B e^{4i(fx+e)} + 6iA e^{2i(fx+e)} - 4B e^{2i(fx+e)} + 3iA - 7B)\sqrt{2}}{60c^2 \sqrt{\frac{c}{e^{2i(fx+e)} + 1}}} f$
parts	$\frac{2iAac \left(-\frac{1}{8c^3 \sqrt{c-ic \tan(fx+e)}} - \frac{1}{12c^2 (c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{1}{10c(c-ic \tan(fx+e))^{\frac{5}{2}}} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{16c^{\frac{7}{2}}} \right)}{f} + \dots$

```
input int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)
```

3.746. $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$

output $2*I/f*a/c*(-1/3*I*B/(c-I*c*\tan(f*x+e))^(3/2)-1/5*c*(A-I*B)/(c-I*c*\tan(f*x+e))^(5/2))$

3.746.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.48

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \frac{\sqrt{2}(3(iA + B)ae^{(6ifx+6ie)} - (-9iA + B)ae^{(4ifx+4ie)} - (-9iA + 11B)ae^{(2ifx+2ie)} - (-3iA + 7B)a)\sqrt{c/(e^{(2ifx+2ie)} + 1)}}{60c^3f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="fricas")`

output $-1/60*\text{sqrt}(2)*(3*(I*A + B)*a*e^{(6*I*f*x + 6*I*e)} - (-9*I*A + B)*a*e^{(4*I*f*x + 4*I*e)} - (-9*I*A + 11*B)*a*e^{(2*I*f*x + 2*I*e)} - (-3*I*A + 7*B)*a)*\text{sqrt}(c/(e^{(2*I*f*x + 2*I*e)} + 1))/(c^3*f)$

3.746.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = ia \left(\int \left(-\frac{A \tan(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 2ic} \right. \right. \\ + \int \frac{A \tan(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 2ic^2 \sqrt{-ic \tan(e + fx) + c \tan(e + fx) + c^2 \sqrt{-ic \tan(e + fx)}}} \\ + \int \frac{B \tan^2(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 2ic^2 \sqrt{-ic \tan(e + fx) + c \tan(e + fx) + c^2 \sqrt{-ic \tan(e + fx)}}} \\ \left. + \int \left(-\frac{iB \tan(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 2ic^2 \sqrt{-ic \tan(e + fx) + c \tan(e + fx) + c^2 \sqrt{-ic \tan(e + fx)}}} \right) \right)$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(5/2),x)`

```
output I*a*(Integral(-I*A/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 2*
I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**2*sqrt(-I*c*tan(e + f
*x) + c)), x) + Integral(A*tan(e + f*x)/(-c**2*sqrt(-I*c*tan(e + f*x) + c)
*tan(e + f*x)**2 - 2*I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**
2*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(B*tan(e + f*x)**2/(-c**2*sqr
t(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 2*I*c**2*sqrt(-I*c*tan(e + f*x)
+ c)*tan(e + f*x) + c**2*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(-I*B
*tan(e + f*x)/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 2*I*c**
2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**2*sqrt(-I*c*tan(e + f*x) +
c)), x))
```

3.746.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.74

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ictan(e + fx))^{5/2}} dx = \frac{2i(5i(-ictan(fx + e) + c)Ba + 3(A - iB)ac)}{15(-ictan(fx + e) + c)^{5/2}cf}$$

```
input integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,
algorithm="maxima")
```

```
output -2/15*I*(5*I*(-I*c*tan(f*x + e) + c)*B*a + 3*(A - I*B)*a*c)/((-I*c*tan(f*x
+ e) + c)^(5/2)*c*f)
```

3.746.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ictan(e + fx))^{5/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)}{(-ictan(fx + e) + c)^{5/2}} dx$$

```
input integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,
algorithm="giac")
```

```
output integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)/(-I*c*tan(f*x + e) +
c)^(5/2), x)
```

3.746. $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ictan(e+fx))^{5/2}} dx$

3.746.9 Mupad [B] (verification not implemented)

Time = 9.96 (sec) , antiderivative size = 232, normalized size of antiderivative = 3.74

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ictan(e + fx))^{5/2}} dx = \frac{a \sqrt{-\frac{c(-2 \cos(e + fx)^2 + \sin(2e + 2fx) 1i)}{2 \cos(e + fx)^2}}}{(7B + 11B(2 \cos(e + fx) - 1))} + \dots$$

```
input int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i))/(c - c*tan(e + f*x)*1i)
^(5/2),x)
```

```
output (a*(-(c*(sin(2*e + 2*f*x)*1i - 2*cos(e + f*x)^2))/(2*cos(e + f*x)^2))^(1/2)
)*(7*B - A*3i - A*(2*cos(e + f*x)^2 - 1)*9i + 11*B*(2*cos(e + f*x)^2 - 1)
+ 9*A*sin(2*e + 2*f*x) + 9*A*sin(4*e + 4*f*x) + 3*A*sin(6*e + 6*f*x) + B*si
in(2*e + 2*f*x)*11i + B*sin(4*e + 4*f*x)*1i - B*sin(6*e + 6*f*x)*3i - A*(2
*cos(2*e + 2*f*x)^2 - 1)*9i - A*(2*cos(3*e + 3*f*x)^2 - 1)*3i + B*(2*cos(2
*e + 2*f*x)^2 - 1) - 3*B*(2*cos(3*e + 3*f*x)^2 - 1))/(60*c^3*f)
```

3.747
$$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{7/2}} dx$$

3.747.1 Optimal result 6827
 3.747.2 Mathematica [A] (verified) 6827
 3.747.3 Rubi [A] (verified) 6828
 3.747.4 Maple [A] (verified) 6829
 3.747.5 Fricas [B] (verification not implemented) 6830
 3.747.6 Sympy [F] 6830
 3.747.7 Maxima [A] (verification not implemented) 6831
 3.747.8 Giac [F] 6832
 3.747.9 Mupad [B] (verification not implemented) 6832

3.747.1 Optimal result

Integrand size = 41, antiderivative size = 62

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{7/2}} dx = -\frac{2a(iA + B)}{7f(c - ict \tan(e + fx))^{7/2}} + \frac{2aB}{5cf(c - ict \tan(e + fx))^{5/2}}$$

output `-2/7*a*(I*A+B)/f/(c-I*c*tan(f*x+e))^(7/2)+2/5*a*B/c/f/(c-I*c*tan(f*x+e))^(5/2)`

3.747.2 Mathematica [A] (verified)

Time = 5.64 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.94

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{7/2}} dx = -\frac{2a\left(\frac{5(iA+B)}{(c-ict \tan(e+fx))^{7/2}} - \frac{7B}{c(c-ict \tan(e+fx))^{5/2}}\right)}{35f}$$

input `Integrate[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(7/2), x]`

output `(-2*a*((5*(I*A + B))/(c - I*c*Tan[e + f*x])^(7/2) - (7*B)/(c*(c - I*c*Tan[e + f*x])^(5/2))))/(35*f)`

3.747.
$$\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{7/2}} dx$$

3.747.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.02, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.098$, Rules used = {3042, 4071, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx$$

↓ 4071

$$ac \int \frac{A + B \tan(e + fx)}{(c - ic \tan(e + fx))^{9/2}} d \tan(e + fx)$$

f
↓ 53

$$ac \int \left(\frac{A - iB}{(c - ic \tan(e + fx))^{9/2}} + \frac{iB}{c(c - ic \tan(e + fx))^{7/2}} \right) d \tan(e + fx)$$

f
↓ 2009

$$ac \left(\frac{2B}{5c^2(c - ic \tan(e + fx))^{5/2}} - \frac{2(B + iA)}{7c(c - ic \tan(e + fx))^{7/2}} \right)$$

f

input `Int[((a + I*a*Tan[e + f*x])*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(7/2),x]`

output `(a*c*((-2*(I*A + B))/(7*c*(c - I*c*Tan[e + f*x])^(7/2)) + (2*B)/(5*c^2*(c - I*c*Tan[e + f*x])^(5/2))))/f`

3.747.3.1 Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int [ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x , Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.747.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.85

method	result
derivativedivides	$\frac{2ia \left(-\frac{c(-iB+A)}{7(c-ic \tan(fx+e))^{\frac{7}{2}}} - \frac{iB}{5(c-ic \tan(fx+e))^{\frac{5}{2}}} \right)}{fc}$
default	$\frac{2ia \left(-\frac{c(-iB+A)}{7(c-ic \tan(fx+e))^{\frac{7}{2}}} - \frac{iB}{5(c-ic \tan(fx+e))^{\frac{5}{2}}} \right)}{fc}$
risch	$-\frac{a(5iA e^{6i(fx+e)} + 5B e^{6i(fx+e)} + 15iA e^{4i(fx+e)} + B e^{4i(fx+e)} + 15iA e^{2i(fx+e)} - 13B e^{2i(fx+e)} + 5iA - 9B)\sqrt{2}}{280c^3 \sqrt{\frac{c}{e^{2i(fx+e)} + 1}}} f$
parts	$\frac{2iAac \left(-\frac{1}{16c^4 \sqrt{c-ic \tan(fx+e)}} - \frac{1}{24c^3 (c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{1}{20c^2 (c-ic \tan(fx+e))^{\frac{5}{2}}} - \frac{1}{14c (c-ic \tan(fx+e))^{\frac{7}{2}}} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)}}{\sqrt{c-ic \tan(fx+e)} + 1}\right)}{f} \right)}{f}$

input `int((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x,method=_RETURNVERBOSE)`

$$3.747. \int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$$

output `2*I/f*a/c*(-1/7*c*(A-I*B)/(c-I*c*tan(f*x+e))^(7/2)-1/5*I*B/(c-I*c*tan(f*x+e))^(5/2))`

3.747.5 Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 113 vs. $2(48) = 96$.

Time = 0.28 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.82

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \frac{\sqrt{2}(5(iA + B)ae^{(8i fx + 8ie)} + 2(10iA + 3B)ae^{(6i fx + 6ie)} + 6(5iA - 2B)ae^{(4i fx + 4ie)} + 2(10iA - 11B)ae^{(2i fx + 2ie)} - (-5iA + 9B)a)\sqrt{c/(e^{(2i fx + 2ie)} + 1)}}{280 c^4 f}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x, algorithm="fricas")`

output `-1/280*sqrt(2)*(5*(I*A + B)*a*e^(8*I*f*x + 8*I*e) + 2*(10*I*A + 3*B)*a*e^(6*I*f*x + 6*I*e) + 6*(5*I*A - 2*B)*a*e^(4*I*f*x + 4*I*e) + 2*(10*I*A - 11*B)*a*e^(2*I*f*x + 2*I*e) - (-5*I*A + 9*B)*a)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^4*f)`

3.747.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = ia \left(\int \left(-\frac{A \tan(e + fx)}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)}}{B \tan^2(e + fx)} \right. \right. \\ \left. \left. + \int \left(-\frac{iB \tan(e + fx)}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)}} \right) \right)$$

3.747. $\int \frac{(a+ia \tan(e+fx))(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(7/2),x)`

output `I*a*(Integral(-I*A/(I*c**3*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x)**3 - 3*c**3*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x)**2 - 3*I*c**3*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x) + c**3*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(A*tan(e + f*x)/(I*c**3*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x)**3 - 3*c**3*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x)**2 - 3*I*c**3*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x) + c**3*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(B*tan(e + f*x)**2/(I*c**3*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x)**3 - 3*c**3*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x)**2 - 3*I*c**3*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x) + c**3*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(-I*B*tan(e + f*x)/(I*c**3*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x)**3 - 3*c**3*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x)**2 - 3*I*c**3*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x) + c**3*sqrt(-I*c*tan(e + f*x) + c)), x))`

3.747.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.74

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \frac{2i(7i(-ic \tan(fx + e) + c)Ba + 5(A - iB)ac)}{35(-ic \tan(fx + e) + c)^{7/2}cf}$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x, algorithm="maxima")`

output `-2/35*I*(7*I*(-I*c*tan(f*x + e) + c)*B*a + 5*(A - I*B)*a*c)/((-I*c*tan(f*x + e) + c)^(7/2)*c*f)`

3.747.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)}{(-ic \tan(fx + e) + c)^{7/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x,
algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)/(-I*c*tan(f*x + e) +
c)^(7/2), x)`

3.747.9 Mupad [B] (verification not implemented)

Time = 10.38 (sec) , antiderivative size = 157, normalized size of antiderivative = 2.53

$$\int \frac{(a + ia \tan(e + fx))(A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx =$$

$$-\sqrt{c - \frac{c \sin(e + fx)}{\cos(e + fx)}} \left(\frac{a(5A + B9i)}{280c^4 f} \right) \text{li}$$

$$+ \frac{ae^{e8i+fx8i}(A - B1i)}{56c^4 f} \text{li} + \frac{ae^{e4i+fx4i}(5A + B2i)}{140c^4 f} \text{li}$$

$$+ \frac{ae^{e2i+fx2i}(10A + B11i)}{140c^4 f} \text{li} + \frac{ae^{e6i+fx6i}(10A - B3i)}{140c^4 f} \text{li}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i))/(c - c*tan(e + f*x)*1i)
^(7/2),x)`

output `-(c - (c*sin(e + f*x)*1i)/cos(e + f*x))^(1/2)*((a*(5*A + B*9i)*1i)/(280*c^4*f) + (a*exp(e*8i + f*x*8i)*(A - B*1i)*1i)/(56*c^4*f) + (a*exp(e*4i + f*x*4i)*(5*A + B*2i)*3i)/(140*c^4*f) + (a*exp(e*2i + f*x*2i)*(10*A + B*11i)*1i)/(140*c^4*f) + (a*exp(e*6i + f*x*6i)*(10*A - B*3i)*1i)/(140*c^4*f))`

3.748 $\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ictan(e+fx))^{7/2} dx$

3.748.1 Optimal result	6833
3.748.2 Mathematica [A] (verified)	6833
3.748.3 Rubi [A] (verified)	6834
3.748.4 Maple [A] (verified)	6836
3.748.5 Fricas [A] (verification not implemented)	6836
3.748.6 Sympy [F]	6837
3.748.7 Maxima [A] (verification not implemented)	6838
3.748.8 Giac [F]	6838
3.748.9 Mupad [B] (verification not implemented)	6839

3.748.1 Optimal result

Integrand size = 43, antiderivative size = 105

$$\int (a + ia \tan(e + fx))^2(A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx = \frac{4a^2(iA + B)(c - ictan(e + fx))^{7/2}}{7f} - \frac{2a^2(iA + 3B)(c - ictan(e + fx))^{9/2}}{9cf} + \frac{2a^2B(c - ictan(e + fx))^{11/2}}{11c^2f}$$

output `4/7*a^2*(I*A+B)*(c-I*c*tan(f*x+e))^(7/2)/f-2/9*a^2*(I*A+3*B)*(c-I*c*tan(f*x+e))^(9/2)/c/f+2/11*a^2*B*(c-I*c*tan(f*x+e))^(11/2)/c^2/f`

3.748.2 Mathematica [A] (verified)

Time = 5.76 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.99

$$\int (a + ia \tan(e + fx))^2(A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx = \frac{a^2c^4 \sec^6(e + fx)(121iA - 33B + (121iA + 93B) \cos(2(e + fx)) + (-77A + 105iB) \sin(2(e + fx)))}{693f \sqrt{c - ictan(e + fx)}}$$

input `Integrate[(a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2), x]`

output $(a^2c^4\text{Sec}[e + f*x]^6*((121*I)*A - 33*B + ((121*I)*A + 93*B)*\text{Cos}[2*(e + f*x)] + (-77*A + (105*I)*B)*\text{Sin}[2*(e + f*x)]*(\text{Cos}[4*(e + f*x)] - I*\text{Sin}[4*(e + f*x)]))/(693*f*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])$

3.748.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.93, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))^2 (c - ictan(e + fx))^{7/2} (A + B \tan(e + fx)) dx$$

↓ 3042

$$\int (a + ia \tan(e + fx))^2 (c - ictan(e + fx))^{7/2} (A + B \tan(e + fx)) dx$$

↓ 4071

$$\frac{a \int a(i \tan(e + fx) + 1)(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} d \tan(e + fx)}{f}$$

↓ 27

$$\frac{a^2 c \int (i \tan(e + fx) + 1)(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} d \tan(e + fx)}{f}$$

↓ 86

$$\frac{a^2 c \int \left(-\frac{iB(c - ictan(e + fx))^{9/2}}{c^2} + \frac{(3iB - A)(c - ictan(e + fx))^{7/2}}{c} + 2(A - iB)(c - ictan(e + fx))^{5/2} \right) d \tan(e + fx)}{f}$$

↓ 2009

$$\frac{a^2 c \left(-\frac{2(3B + iA)(c - ictan(e + fx))^{9/2}}{9c^2} + \frac{4(B + iA)(c - ictan(e + fx))^{7/2}}{7c} + \frac{2B(c - ictan(e + fx))^{11/2}}{11c^3} \right)}{f}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^2*(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^(7/2),x]$

3.748. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx$

```
output (a^2*c*((4*(I*A + B)*(c - I*c*Tan[e + f*x])^(7/2))/(7*c) - (2*(I*A + 3*B)*
(c - I*c*Tan[e + f*x])^(9/2))/(9*c^2) + (2*B*(c - I*c*Tan[e + f*x])^(11/2)
)/(11*c^3))/f
```

3.748.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.748.4 Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.80

method	result
derivativedivides	$\frac{2ia^2 \left(-\frac{iB(c-ic \tan(fx+e))^{11}}{11} + \frac{(3iBc-cA)(c-ic \tan(fx+e))^9}{9} + \frac{2(-iBc+cA)c(c-ic \tan(fx+e))^{7/2}}{7} \right)}{f c^2}$
default	$\frac{2ia^2 \left(-\frac{iB(c-ic \tan(fx+e))^{11}}{11} + \frac{(3iBc-cA)(c-ic \tan(fx+e))^9}{9} + \frac{2(-iBc+cA)c(c-ic \tan(fx+e))^{7/2}}{7} \right)}{f c^2}$
parts	$\frac{2iA a^2 c \left(-\frac{(c-ic \tan(fx+e))^{5/2}}{5} - \frac{2c(c-ic \tan(fx+e))^{3/2}}{3} - 4\sqrt{c-ic \tan(fx+e)} c^2 + 4c^{5/2} \sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}} \right) \right)}{f}$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x,method=_RETURNVERBOSE)`

output `2*I/f*a^2/c^2*(-1/11*I*B*(c-I*c*tan(f*x+e))^(11/2)+1/9*(3*I*B*c-c*A)*(c-I*c*tan(f*x+e))^(9/2)+2/7*(-I*B*c+c*A)*c*(c-I*c*tan(f*x+e))^(7/2))`

3.748.5 Fracas [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.42

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx = \frac{32 \sqrt{2} (99(-iA - B)a^2 c^3 e^{4i fx + 4ie} + 11(-11iA + 3B)a^2 c^3 e^{2i fx + 2ie} + 2(-11iA + 3B)a^2 c^3) \sqrt{\frac{c}{e^{2i fx + 2ie}}}}{693 (f e^{10i fx + 10ie} + 5 f e^{8i fx + 8ie} + 10 f e^{6i fx + 6ie} + 10 f e^{4i fx + 4ie} + 5 f e^{2i fx + 2ie} + f)}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="fracas")`

output `-32/693*sqrt(2)*(99*(-I*A - B)*a^2*c^3*e^(4*I*f*x + 4*I*e) + 11*(-11*I*A + 3*B)*a^2*c^3*e^(2*I*f*x + 2*I*e) + 2*(-11*I*A + 3*B)*a^2*c^3)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(f*e^(10*I*f*x + 10*I*e) + 5*f*e^(8*I*f*x + 8*I*e) + 10*f*e^(6*I*f*x + 6*I*e) + 10*f*e^(4*I*f*x + 4*I*e) + 5*f*e^(2*I*f*x + 2*I*e) + f)`

3.748. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx$

3.748.6 Sympy [F]

$$\begin{aligned}
& \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \\
& -a^2 \left(\int (-Ac^3 \sqrt{-ic \tan(e + fx) + c}) dx \right. \\
& + \int (-2Ac^3 \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx)) dx \\
& + \int (-Ac^3 \sqrt{-ic \tan(e + fx) + c} \tan^4(e + fx)) dx \\
& + \int (-Bc^3 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx)) dx \\
& + \int (-2Bc^3 \sqrt{-ic \tan(e + fx) + c} \tan^3(e + fx)) dx \\
& + \int (-Bc^3 \sqrt{-ic \tan(e + fx) + c} \tan^5(e + fx)) dx \\
& + \int iAc^3 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx) dx \\
& + \int 2iAc^3 \sqrt{-ic \tan(e + fx) + c} \tan^3(e + fx) dx \\
& + \int iAc^3 \sqrt{-ic \tan(e + fx) + c} \tan^5(e + fx) dx \\
& + \int iBc^3 \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx) dx \\
& + \int 2iBc^3 \sqrt{-ic \tan(e + fx) + c} \tan^4(e + fx) dx \\
& \left. + \int iBc^3 \sqrt{-ic \tan(e + fx) + c} \tan^6(e + fx) dx \right)
\end{aligned}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(7/2),x)`

output `-a**2*(Integral(-A*c**3*sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-2*A*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x) + Integral(-A*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4, x) + Integral(-B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(-2*B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x) + Integral(-B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**5, x) + Integral(I*A*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(2*I*A*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x) + Integral(I*A*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**5, x) + Integral(I*B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x) + Integral(2*I*B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4, x) + Integral(I*B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**6, x))`

3.748.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.74

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx = \frac{2i \left(63i(-ictan(fx + e) + c)^{\frac{11}{2}} Ba^2 + 77(-ictan(fx + e) + c)^{\frac{9}{2}} (A - 3iB)a^2c - 198(-ictan(fx + e) + c)^{\frac{7}{2}} (A - 3iB)a^2c^2 \right)}{693c^2f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2), x, algorithm="maxima")`

output `-2/693*I*(63*I*(-I*c*tan(f*x + e) + c)^(11/2)*B*a^2 + 77*(-I*c*tan(f*x + e) + c)^(9/2)*(A - 3*I*B)*a^2*c - 198*(-I*c*tan(f*x + e) + c)^(7/2)*(A - I*B)*a^2*c^2)/(c^2*f)`

3.748.8 Giac [F]

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx = \int (B \tan(fx + e) + A)(ia \tan(fx + e) + a)^2 (-ictan(fx + e) + c)^{\frac{7}{2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2), x, algorithm="giac")`

3.748. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx$

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^2*(-I*c*tan(f*x + e) + c)^(7/2), x)`

3.748.9 Mupad [B] (verification not implemented)

Time = 11.86 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.26

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx = \frac{32 a^2 c^3 \sqrt{c + \frac{c(e^{e^{2i+fx} 2i} 1i - i) 1i}{e^{e^{2i+fx} 2i} + 1}} (A 22i - 6 B + A e^{e^{2i+fx} 2i} 121i + A e^{e^{4i+fx} 4i} 99i - 33 B e^{e^{2i+fx} 2i})}{693 f (e^{e^{2i+fx} 2i} + 1)^5}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)^(7/2),x)`

output `(32*a^2*c^3*(c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(1/2)*(A*22i - 6*B + A*exp(e*2i + f*x*2i)*121i + A*exp(e*4i + f*x*4i)*99i - 33*B*exp(e*2i + f*x*2i) + 99*B*exp(e*4i + f*x*4i))/(693*f*(exp(e*2i + f*x*2i) + 1)^5)`

3.749 $\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ictan(e+fx))^{5/2} dx$

3.749.1 Optimal result	6840
3.749.2 Mathematica [A] (verified)	6840
3.749.3 Rubi [A] (verified)	6841
3.749.4 Maple [A] (verified)	6843
3.749.5 Fricas [A] (verification not implemented)	6843
3.749.6 Sympy [F]	6844
3.749.7 Maxima [A] (verification not implemented)	6844
3.749.8 Giac [F]	6845
3.749.9 Mupad [B] (verification not implemented)	6845

3.749.1 Optimal result

Integrand size = 43, antiderivative size = 105

$$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ictan(e+fx))^{5/2} dx = \frac{4a^2(iA+B)(c-ictan(e+fx))^{5/2}}{5f} - \frac{2a^2(iA+3B)(c-ictan(e+fx))^{7/2}}{7cf} + \frac{2a^2B(c-ictan(e+fx))^{9/2}}{9c^2f}$$

output `4/5*a^2*(I*A+B)*(c-I*c*tan(f*x+e))^(5/2)/f-2/7*a^2*(I*A+3*B)*(c-I*c*tan(f*x+e))^(7/2)/c/f+2/9*a^2*B*(c-I*c*tan(f*x+e))^(9/2)/c^2/f`

3.749.2 Mathematica [A] (verified)

Time = 5.30 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.99

$$\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ictan(e+fx))^{5/2} dx = \frac{a^2c^3 \sec^5(e+fx)(81iA-9B+(81iA+61B) \cos(2(e+fx)) + (-45A+65iB) \sin(2(e+fx)))}{315f \sqrt{c-ictan(e+fx)}}$$

input `Integrate[(a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2), x]`

output $(a^2c^3\text{Sec}[e + f*x]^5*((81*I)*A - 9*B + ((81*I)*A + 61*B)*\text{Cos}[2*(e + f*x)] + (-45*A + (65*I)*B)*\text{Sin}[2*(e + f*x)]*(\text{Cos}[3*(e + f*x)] - I*\text{Sin}[3*(e + f*x)]))/((315*f*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]]))$

3.749.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.93, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))^2 (c - ictan(e + fx))^{5/2} (A + B \tan(e + fx)) dx$$

↓ 3042

$$\int (a + ia \tan(e + fx))^2 (c - ictan(e + fx))^{5/2} (A + B \tan(e + fx)) dx$$

↓ 4071

$$\frac{ac \int a(i \tan(e + fx) + 1)(A + B \tan(e + fx))(c - ictan(e + fx))^{3/2} d \tan(e + fx)}{f}$$

↓ 27

$$\frac{a^2c \int (i \tan(e + fx) + 1)(A + B \tan(e + fx))(c - ictan(e + fx))^{3/2} d \tan(e + fx)}{f}$$

↓ 86

$$\frac{a^2c \int \left(-\frac{iB(c-ictan(e+fx))^{7/2}}{c^2} + \frac{(3iB-A)(c-ictan(e+fx))^{5/2}}{c} + 2(A-iB)(c-ictan(e+fx))^{3/2} \right) d \tan(e + fx)}{f}$$

↓ 2009

$$\frac{a^2c \left(-\frac{2(3B+iA)(c-ictan(e+fx))^{7/2}}{7c^2} + \frac{4(B+iA)(c-ictan(e+fx))^{5/2}}{5c} + \frac{2B(c-ictan(e+fx))^{9/2}}{9c^3} \right)}{f}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^2*(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^(5/2),x]$

3.749. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx$


```
output (a^2*c*((4*(I*A + B)*(c - I*c*Tan[e + f*x])^(5/2))/(5*c) - (2*(I*A + 3*B)*
(c - I*c*Tan[e + f*x])^(7/2))/(7*c^2) + (2*B*(c - I*c*Tan[e + f*x])^(9/2))
/(9*c^3)))/f
```

3.749.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.749.4 Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.80

method	result
derivativedivides	$\frac{2ia^2 \left(-\frac{iB(c-ic \tan(fx+e))^{9/2}}{9} + \frac{(3iBc-cA)(c-ic \tan(fx+e))^{7/2}}{7} + \frac{2(-iBc+cA)c(c-ic \tan(fx+e))^{5/2}}{5} \right)}{fc^2}$
default	$\frac{2ia^2 \left(-\frac{iB(c-ic \tan(fx+e))^{9/2}}{9} + \frac{(3iBc-cA)(c-ic \tan(fx+e))^{7/2}}{7} + \frac{2(-iBc+cA)c(c-ic \tan(fx+e))^{5/2}}{5} \right)}{fc^2}$
parts	$\frac{2iAa^2c \left(-\frac{(c-ic \tan(fx+e))^{3/2}}{3} - 2c\sqrt{c-ic \tan(fx+e)} + 2c^{3/2}\sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(fx+e)}\sqrt{2}}{2\sqrt{c}} \right) \right)}{f} + \frac{a^2(2iA+B) \left(\frac{2(c-ic \tan(fx+e))^{5/2}}{5} \right)}{f}$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

output `2*I/f*a^2/c^2*(-1/9*I*B*(c-I*c*tan(f*x+e))^(9/2)+1/7*(3*I*B*c-c*A)*(c-I*c*tan(f*x+e))^(7/2)+2/5*(-I*B*c+c*A)*c*(c-I*c*tan(f*x+e))^(5/2))`

3.749.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.27

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2} dx = \frac{16\sqrt{2}(63(-iA - B)a^2c^2e^{4ifx+4ie} + 9(-9iA + B)a^2c^2e^{2ifx+2ie} + 2(-9iA + B)a^2c^2)\sqrt{\frac{c}{e^{(2ifx+2ie)}+1}}}{315(fe^{(8ifx+8ie)} + 4fe^{(6ifx+6ie)} + 6fe^{(4ifx+4ie)} + 4fe^{(2ifx+2ie)} + f)}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="fracas")`

output `-16/315*sqrt(2)*(63*(-I*A - B)*a^2*c^2*e^(4*I*f*x + 4*I*e) + 9*(-9*I*A + B)*a^2*c^2*e^(2*I*f*x + 2*I*e) + 2*(-9*I*A + B)*a^2*c^2)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(f*e^(8*I*f*x + 8*I*e) + 4*f*e^(6*I*f*x + 6*I*e) + 6*f*e^(4*I*f*x + 4*I*e) + 4*f*e^(2*I*f*x + 2*I*e) + f)`

3.749. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2} dx$

3.749.6 Sympy [F]

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx =$$

$$-a^2 \left(\int (-Ac^2 \sqrt{-ic \tan(e + fx) + c}) dx \right.$$

$$+ \int (-2Ac^2 \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx)) dx$$

$$+ \int (-Ac^2 \sqrt{-ic \tan(e + fx) + c} \tan^4(e + fx)) dx$$

$$+ \int (-Bc^2 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx)) dx$$

$$+ \int (-2Bc^2 \sqrt{-ic \tan(e + fx) + c} \tan^3(e + fx)) dx$$

$$\left. + \int (-Bc^2 \sqrt{-ic \tan(e + fx) + c} \tan^5(e + fx)) dx \right)$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(5/2),x)`

output `-a**2*(Integral(-A*c**2*sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-2*A*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x) + Integral(-A*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4, x) + Integral(-B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(-2*B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x) + Integral(-B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**5, x))`

3.749.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.74

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx =$$

$$\frac{2i \left(35i (-ic \tan(fx + e) + c)^{\frac{9}{2}} Ba^2 + 45 (-ic \tan(fx + e) + c)^{\frac{7}{2}} (A - 3iB)a^2c - 126 (-ic \tan(fx + e) - \dots \right)}{315c^2f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="maxima")`

3.749. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx$

output
$$\frac{-2/315 \cdot I \cdot (35 \cdot I \cdot (-I \cdot c \cdot \tan(f \cdot x + e) + c)^{9/2} \cdot B \cdot a^2 + 45 \cdot (-I \cdot c \cdot \tan(f \cdot x + e) + c)^{7/2} \cdot (A - 3 \cdot I \cdot B) \cdot a^2 \cdot c - 126 \cdot (-I \cdot c \cdot \tan(f \cdot x + e) + c)^{5/2} \cdot (A - I \cdot B) \cdot a^2 \cdot c^2)}{(c^2 \cdot f)}$$

3.749.8 Giac [F]

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^{5/2} dx = \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^2 (-ictan(fx + e) + c)^{5/2} dx$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^2*(-I*c*tan(f*x + e) + c)^(5/2), x)`

3.749.9 Mupad [B] (verification not implemented)

Time = 12.74 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.26

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^{5/2} dx = \frac{16 a^2 c^2 \sqrt{c + \frac{c(e^{e 2i + f x 2i} 1i - i) 1i}{e^{e 2i + f x 2i} + 1}} (A 18i - 2B + A e^{e 2i + f x 2i} 81i + A e^{e 4i + f x 4i} 63i - 9B e^{e 2i + f x 2i})}{315 f (e^{e 2i + f x 2i} + 1)^4}$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)^(5/2),x)`

output
$$\frac{(16 \cdot a^2 \cdot c^2 \cdot (c + (c \cdot (\exp(e \cdot 2i + f \cdot x \cdot 2i) \cdot 1i - 1i) \cdot 1i) / (\exp(e \cdot 2i + f \cdot x \cdot 2i) + 1)))^{1/2} \cdot (A \cdot 18i - 2 \cdot B + A \cdot \exp(e \cdot 2i + f \cdot x \cdot 2i) \cdot 81i + A \cdot \exp(e \cdot 4i + f \cdot x \cdot 4i) \cdot 63i - 9 \cdot B \cdot \exp(e \cdot 2i + f \cdot x \cdot 2i) + 63 \cdot B \cdot \exp(e \cdot 4i + f \cdot x \cdot 4i))}{315 \cdot f \cdot (\exp(e \cdot 2i + f \cdot x \cdot 2i) + 1)^4}$$

3.750 $\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))(c-ictan(e+fx))^{3/2} dx$

3.750.1 Optimal result	6846
3.750.2 Mathematica [A] (verified)	6846
3.750.3 Rubi [A] (verified)	6847
3.750.4 Maple [A] (verified)	6849
3.750.5 Fricas [A] (verification not implemented)	6849
3.750.6 Sympy [F]	6850
3.750.7 Maxima [A] (verification not implemented)	6851
3.750.8 Giac [F]	6851
3.750.9 Mupad [B] (verification not implemented)	6852

3.750.1 Optimal result

Integrand size = 43, antiderivative size = 105

$$\int (a + ia \tan(e + fx))^2(A + B \tan(e + fx))(c - ictan(e + fx))^{3/2} dx = \frac{4a^2(iA + B)(c - ictan(e + fx))^{3/2}}{3f} - \frac{2a^2(iA + 3B)(c - ictan(e + fx))^{5/2}}{5cf} + \frac{2a^2B(c - ictan(e + fx))^{7/2}}{7c^2f}$$

output `4/3*a^2*(I*A+B)*(c-I*c*tan(f*x+e))^(3/2)/f-2/5*a^2*(I*A+3*B)*(c-I*c*tan(f*x+e))^(5/2)/c/f+2/7*a^2*B*(c-I*c*tan(f*x+e))^(7/2)/c^2/f`

3.750.2 Mathematica [A] (verified)

Time = 4.24 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.00

$$\int (a + ia \tan(e + fx))^2(A + B \tan(e + fx))(c - ictan(e + fx))^{3/2} dx = \frac{a^2c^2 \sec^4(e + fx)(\cos(2(e + fx)) - i \sin(2(e + fx)))(7(7iA + B) + (49iA + 37B) \cos(2(e + fx)))}{105f \sqrt{c - ictan(e + fx)}}$$

input `Integrate[(a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2), x]`

output $(a^2c^2\text{Sec}[e + f*x]^4(\text{Cos}[2*(e + f*x)] - I*\text{Sin}[2*(e + f*x)])*(7*((7*I)*A + B) + ((49*I)*A + 37*B)*\text{Cos}[2*(e + f*x)] + (-21*A + (33*I)*B)*\text{Sin}[2*(e + f*x)]))/(105*f*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])$

3.750.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.93, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))^2 (c - ictan(e + fx))^{3/2} (A + B \tan(e + fx)) dx$$

↓ 3042

$$\int (a + ia \tan(e + fx))^2 (c - ictan(e + fx))^{3/2} (A + B \tan(e + fx)) dx$$

↓ 4071

$$\frac{ac \int a(i \tan(e + fx) + 1)(A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} d \tan(e + fx)}{f}$$

↓ 27

$$\frac{a^2c \int (i \tan(e + fx) + 1)(A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} d \tan(e + fx)}{f}$$

↓ 86

$$\frac{a^2c \int \left(-\frac{iB(c - ictan(e + fx))^{5/2}}{c^2} + \frac{(3iB - A)(c - ictan(e + fx))^{3/2}}{c} + 2(A - iB) \sqrt{c - ictan(e + fx)} \right) d \tan(e + fx)}{f}$$

↓ 2009

$$\frac{a^2c \left(-\frac{2(3B + iA)(c - ictan(e + fx))^{5/2}}{5c^2} + \frac{4(B + iA)(c - ictan(e + fx))^{3/2}}{3c} + \frac{2B(c - ictan(e + fx))^{7/2}}{7c^3} \right)}{f}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^2*(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^(3/2), x]$

3.750. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx$

```
output (a^2*c*((4*(I*A + B)*(c - I*c*Tan[e + f*x])^(3/2))/(3*c) - (2*(I*A + 3*B)*
(c - I*c*Tan[e + f*x])^(5/2))/(5*c^2) + (2*B*(c - I*c*Tan[e + f*x])^(7/2))
/(7*c^3)))/f
```

3.750.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.750.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.80

method	result
derivativedivides	$\frac{2ia^2 \left(-\frac{iB(c-ic \tan(fx+e))^{\frac{7}{2}}}{7} + \frac{(3iBc-cA)(c-ic \tan(fx+e))^{\frac{5}{2}}}{5} + \frac{2(-iBc+cA)c(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} \right)}{f c^2}$
default	$\frac{2ia^2 \left(-\frac{iB(c-ic \tan(fx+e))^{\frac{7}{2}}}{7} + \frac{(3iBc-cA)(c-ic \tan(fx+e))^{\frac{5}{2}}}{5} + \frac{2(-iBc+cA)c(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} \right)}{f c^2}$
parts	$\frac{2iA a^2 c \left(-\sqrt{c-ic \tan(fx+e)} + \sqrt{c} \sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}} \right) \right)}{f} + \frac{a^2(2iA+B) \left(\frac{2(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} + 2c\sqrt{c-ic \tan(fx+e)} \right)}{f}$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

output `2*I/f*a^2/c^2*(-1/7*I*B*(c-I*c*tan(f*x+e))^(7/2)+1/5*(3*I*B*c-c*A)*(c-I*c*tan(f*x+e))^(5/2)+2/3*(-I*B*c+c*A)*c*(c-I*c*tan(f*x+e))^(3/2))`

3.750.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.13

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx = \frac{8\sqrt{2}(35(-iA - B)a^2ce^{(4i fx + 4i e)} + 7(-7iA - B)a^2ce^{(2i fx + 2i e)} + 2(-7iA - B)a^2c)\sqrt{\frac{c}{e^{(2i fx + 2i e)} + 1}}}{105(fe^{(6i fx + 6i e)} + 3fe^{(4i fx + 4i e)} + 3fe^{(2i fx + 2i e)} + f)}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x, algorithm="fracas")`

output `-8/105*sqrt(2)*(35*(-I*A - B)*a^2*c*e^(4*I*f*x + 4*I*e) + 7*(-7*I*A - B)*a^2*c*e^(2*I*f*x + 2*I*e) + 2*(-7*I*A - B)*a^2*c)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)`

3.750. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx$

3.750.6 Sympy [F]

$$\begin{aligned}
& \int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} dx = \\
& -a^2 \left(\int (-Ac \sqrt{-ic \tan(e + fx) + c}) dx \right. \\
& + \int (-Ac \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx)) dx \\
& + \int (-Bc \sqrt{-ic \tan(e + fx) + c} \tan(e + fx)) dx \\
& + \int (-Bc \sqrt{-ic \tan(e + fx) + c} \tan^3(e + fx)) dx \\
& + \int (-iAc \sqrt{-ic \tan(e + fx) + c} \tan(e + fx)) dx \\
& + \int (-iAc \sqrt{-ic \tan(e + fx) + c} \tan^3(e + fx)) dx \\
& + \int (-iBc \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx)) dx \\
& \left. + \int (-iBc \sqrt{-ic \tan(e + fx) + c} \tan^4(e + fx)) dx \right)
\end{aligned}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(3/2),x)`

output `-a**2*(Integral(-A*c*sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-A*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x) + Integral(-B*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(-B*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x) + Integral(-I*A*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(-I*A*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x) + Integral(-I*B*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x) + Integral(-I*B*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4, x))`

3.750.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.74

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} dx =$$

$$\frac{2i \left(15i (-ic \tan(fx + e) + c)^{7/2} B a^2 + 21 (-ic \tan(fx + e) + c)^{5/2} (A - 3i B) a^2 c - 70 (-ic \tan(fx + e) + c)^{3/2} (A - I*B) a^2 c^2 \right)}{105 c^2 f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x
, algorithm="maxima")`

output `-2/105*I*(15*I*(-I*c*tan(f*x + e) + c)^(7/2)*B*a^2 + 21*(-I*c*tan(f*x + e) + c)^(5/2)*(A - 3*I*B)*a^2*c - 70*(-I*c*tan(f*x + e) + c)^(3/2)*(A - I*B)*a^2*c^2)/(c^2*f)`

3.750.8 Giac [F]

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} dx = \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^2 (-ic \tan(fx + e) + c)^{3/2} dx$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^2*(-I*c*tan(f*x + e) + c)^(3/2), x)`

3.750.9 Mupad [B] (verification not implemented)

Time = 11.70 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.24

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx))(c - ictan(e + fx))^{3/2} dx = \frac{8a^2 c \sqrt{c + \frac{c(e^{e^{2i} + fx^{2i}} - 1) - 1}{e^{e^{2i} + fx^{2i}} + 1}}} (A 14i + 2B + A e^{e^{2i} + fx^{2i}} 49i + A e^{e^{4i} + fx^{4i}} 35i + 7B e^{e^{2i} + fx^{2i}} + 105 f (e^{e^{2i} + fx^{2i}} + 1)^3$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)^(3/2),x)`

output `(8*a^2*c*(c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(1/2)*(A*14i + 2*B + A*exp(e*2i + f*x*2i)*49i + A*exp(e*4i + f*x*4i)*35i + 7*B*exp(e*2i + f*x*2i) + 35*B*exp(e*4i + f*x*4i))/(105*f*(exp(e*2i + f*x*2i) + 1)^3)`

3.751 $\int (a+ia \tan(e+fx))^2(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)} dx$

3.751.1 Optimal result	6853
3.751.2 Mathematica [A] (verified)	6853
3.751.3 Rubi [A] (verified)	6854
3.751.4 Maple [A] (verified)	6856
3.751.5 Fricas [A] (verification not implemented)	6856
3.751.6 Sympy [F]	6857
3.751.7 Maxima [A] (verification not implemented)	6857
3.751.8 Giac [F]	6858
3.751.9 Mupad [B] (verification not implemented)	6858

3.751.1 Optimal result

Integrand size = 43, antiderivative size = 103

$$\int (a + ia \tan(e + fx))^2(A + B \tan(e + fx))\sqrt{c - ic \tan(e + fx)} dx$$

$$= \frac{4a^2(iA + B)\sqrt{c - ic \tan(e + fx)}}{f} - \frac{2a^2(iA + 3B)(c - ic \tan(e + fx))^{3/2}}{3cf} + \frac{2a^2B(c - ic \tan(e + fx))^{5/2}}{5c^2f}$$

output `4*a^2*(I*A+B)*(c-I*c*tan(f*x+e))^(1/2)/f-2/3*a^2*(I*A+3*B)*(c-I*c*tan(f*x+e))^(3/2)/c/f+2/5*a^2*B*(c-I*c*tan(f*x+e))^(5/2)/c^2/f`

3.751.2 Mathematica [A] (verified)

Time = 3.19 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.74

$$\int (a + ia \tan(e + fx))^2(A + B \tan(e + fx))\sqrt{c - ic \tan(e + fx)} dx$$

$$= \frac{2a^2c(i + \tan(e + fx))(25A - 18iB + (5iA + 9B) \tan(e + fx) + 3iB \tan^2(e + fx))}{15f\sqrt{c - ic \tan(e + fx)}}$$

input `Integrate[(a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]],x]`

output $(2*a^2*c*(I + \text{Tan}[e + f*x])*(25*A - (18*I)*B + ((5*I)*A + 9*B)*\text{Tan}[e + f*x] + (3*I)*B*\text{Tan}[e + f*x]^2))/(15*f*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])$

3.751.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.93, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))^2 \sqrt{c - ictan(e + fx)} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))^2 \sqrt{c - ictan(e + fx)} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{a(i \tan(e+fx)+1)(A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{a^2c \int \frac{(i \tan(e+fx)+1)(A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{86} \\ & \frac{a^2c \int \left(-\frac{iB(c-ictan(e+fx))^{3/2}}{c^2} + \frac{(3iB-A)\sqrt{c-ictan(e+fx)}}{c} + \frac{2(A-iB)}{\sqrt{c-ictan(e+fx)}} \right) d \tan(e + fx)}{f} \\ & \quad \downarrow \text{2009} \\ & \frac{a^2c \left(-\frac{2(3B+iA)(c-ictan(e+fx))^{3/2}}{3c^2} + \frac{4(B+iA)\sqrt{c-ictan(e+fx)}}{c} + \frac{2B(c-ictan(e+fx))^{5/2}}{5c^3} \right)}{f} \end{aligned}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^2*(A + B*\text{Tan}[e + f*x])* \text{Sqrt}[c - I*c*\text{Tan}[e + f*x]],x]$

```
output (a^2*c*((4*(I*A + B)*Sqrt[c - I*c*Tan[e + f*x]])/c - (2*(I*A + 3*B)*(c - I
*c*Tan[e + f*x])^(3/2))/(3*c^2) + (2*B*(c - I*c*Tan[e + f*x])^(5/2))/(5*c^
3)))/f
```

3.751.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.751.4 Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.82

method	result
derivativedivides	$\frac{2ia^2 \left(-\frac{iB(c-ic \tan(fx+e))^{5/2}}{5} + \frac{(3iBc-cA)(c-ic \tan(fx+e))^{3/2}}{3} + 2\sqrt{c-ic \tan(fx+e)}(-iBc+cA)c \right)}{f c^2}$
default	$\frac{2ia^2 \left(-\frac{iB(c-ic \tan(fx+e))^{5/2}}{5} + \frac{(3iBc-cA)(c-ic \tan(fx+e))^{3/2}}{3} + 2\sqrt{c-ic \tan(fx+e)}(-iBc+cA)c \right)}{f c^2}$
parts	$-\frac{2ia^2(-2iB+A) \left(\frac{(c-ic \tan(fx+e))^{3/2}}{3} - \frac{c^{3/2} \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{2} \right)}{f c} + \frac{a^2(2iA+B) \left(2\sqrt{c-ic \tan(fx+e)} - \sqrt{\frac{c}{e^{2i fx+2ie}+1}} \right)}{f c}$

input `int((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output `2*I/f*a^2/c^2*(-1/5*I*B*(c-I*c*tan(f*x+e))^(5/2)+1/3*(3*I*B*c-c*A)*(c-I*c*tan(f*x+e))^(3/2)+2*(c-I*c*tan(f*x+e))^(1/2)*(-I*B*c+c*A)*c)`

3.751.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.01

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx = \frac{4\sqrt{2}(15(-iA - B)a^2 e^{(4i fx+4ie)} + 5(-5iA - 3B)a^2 e^{(2i fx+2ie)} + 2(-5iA - 3B)a^2) \sqrt{\frac{c}{e^{(2i fx+2ie)}+1}}}{15(fe^{(4i fx+4ie)} + 2fe^{(2i fx+2ie)} + f)}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e)),x,algorithm="fricas")`

output `-4/15*sqrt(2)*(15*(-I*A - B)*a^2*e^(4*I*f*x + 4*I*e) + 5*(-5*I*A - 3*B)*a^2*e^(2*I*f*x + 2*I*e) + 2*(-5*I*A - 3*B)*a^2)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(f*e^(4*I*f*x + 4*I*e) + 2*f*e^(2*I*f*x + 2*I*e) + f)`

3.751.6 Sympy [F]

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx$$

$$= -a^2 \left(\int \left(-A \sqrt{-ictan(e + fx) + c} \right) dx + \int A \sqrt{-ictan(e + fx) + c} \tan^2(e + fx) dx \right.$$

$$+ \int \left(-B \sqrt{-ictan(e + fx) + c} \tan(e + fx) \right) dx$$

$$+ \int B \sqrt{-ictan(e + fx) + c} \tan^3(e + fx) dx$$

$$+ \int \left(-2iA \sqrt{-ictan(e + fx) + c} \tan(e + fx) \right) dx$$

$$\left. + \int \left(-2iB \sqrt{-ictan(e + fx) + c} \tan^2(e + fx) \right) dx \right)$$

input `integrate((c-I*c*tan(f*x+e))**(1/2)*(a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e)),x)`

output `-a**2*(Integral(-A*sqrt(-I*c*tan(e + f*x) + c), x) + Integral(A*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x) + Integral(-B*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(B*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x) + Integral(-2*I*A*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(-2*I*B*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x))`

3.751.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.76

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx =$$

$$\frac{2i \left(3i (-ictan(fx + e) + c)^{\frac{5}{2}} Ba^2 + 5 (-ictan(fx + e) + c)^{\frac{3}{2}} (A - 3iB) a^2 c - 30 \sqrt{-ictan(fx + e)} \right)}{15c^2 f}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e)),x, algorithm="maxima")`

output `-2/15*I*(3*I*(-I*c*tan(f*x + e) + c)^(5/2)*B*a^2 + 5*(-I*c*tan(f*x + e) + c)^(3/2)*(A - 3*I*B)*a^2*c - 30*sqrt(-I*c*tan(f*x + e) + c)*(A - I*B)*a^2*c^2)/(c^2*f)`

3.751. $\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx$

3.751.8 Giac [F]

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx$$

$$= \int (B \tan(fx + e) + A)(ia \tan(fx + e) + a)^2 \sqrt{-ic \tan(fx + e) + c} dx$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e)),x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^2*sqrt(-I*c*tan(f*x
+ e) + c), x)`

3.751.9 Mupad [B] (verification not implemented)

Time = 10.35 (sec) , antiderivative size = 241, normalized size of antiderivative = 2.34

$$\int (a + ia \tan(e + fx))^2 (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx$$

$$= \frac{2a^2 \sqrt{\frac{c(\cos(2e+2fx)+1) - \sin(2e+2fx)i}{\cos(2e+2fx)+1}} (A 250i + 174B + A \cos(2e + 2fx) 375i + A \cos(4e + 4fx) 150i -$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)
^(1/2),x)`

output `(2*a^2*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x)
+ 1))^(1/2)*(A*250i + 174*B + A*cos(2*e + 2*f*x)*375i + A*cos(4*e + 4*f*x)
)*150i + A*cos(6*e + 6*f*x)*25i + 267*B*cos(2*e + 2*f*x) + 114*B*cos(4*e +
4*f*x) + 21*B*cos(6*e + 6*f*x) - 25*A*sin(2*e + 2*f*x) - 20*A*sin(4*e + 4
*f*x) - 5*A*sin(6*e + 6*f*x) + B*sin(2*e + 2*f*x)*45i + B*sin(4*e + 4*f*x)
*36i + B*sin(6*e + 6*f*x)*9i))/(15*f*(15*cos(2*e + 2*f*x) + 6*cos(4*e + 4*
f*x) + cos(6*e + 6*f*x) + 10))`

3.752
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$$

3.752.1 Optimal result	6859
3.752.2 Mathematica [A] (verified)	6859
3.752.3 Rubi [A] (verified)	6860
3.752.4 Maple [A] (verified)	6862
3.752.5 Fricas [A] (verification not implemented)	6862
3.752.6 Sympy [F]	6863
3.752.7 Maxima [A] (verification not implemented)	6863
3.752.8 Giac [F]	6864
3.752.9 Mupad [B] (verification not implemented)	6864

3.752.1 Optimal result

Integrand size = 43, antiderivative size = 101

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx \\ &= -\frac{4a^2(iA + B)}{f\sqrt{c - ic \tan(e + fx)}} - \frac{2a^2(iA + 3B)\sqrt{c - ic \tan(e + fx)}}{cf} \\ & \quad + \frac{2a^2B(c - ic \tan(e + fx))^{3/2}}{3c^2f} \end{aligned}$$

output `-4*a^2*(I*A+B)/f/(c-I*c*tan(f*x+e))^(1/2)-2*a^2*(I*A+3*B)*(c-I*c*tan(f*x+e))^(1/2)/c/f+2/3*a^2*B*(c-I*c*tan(f*x+e))^(3/2)/c^2/f`

3.752.2 Mathematica [A] (verified)

Time = 4.09 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.61

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx \\ &= -\frac{2a^2(9iA + 14B + (3A - 7iB) \tan(e + fx) + B \tan^2(e + fx))}{3f\sqrt{c - ic \tan(e + fx)}} \end{aligned}$$

input `Integrate[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/Sqrt[c - I*c*Tan[e + f*x]],x]`

output $(-2*a^2*((9*I)*A + 14*B + (3*A - (7*I)*B)*\text{Tan}[e + f*x] + B*\text{Tan}[e + f*x]^2)/((3*f*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]]))$

3.752.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.93, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx \\ & \quad \downarrow 3042 \\ & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx \\ & \quad \downarrow 4071 \\ & \frac{ac \int \frac{a(i \tan(e + fx) + 1)(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{f} \\ & \quad \downarrow 27 \\ & \frac{a^2 c \int \frac{(i \tan(e + fx) + 1)(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{f} \\ & \quad \downarrow 86 \\ & \frac{a^2 c \int \left(-\frac{2(A - iB)}{(c - ic \tan(e + fx))^{3/2}} - \frac{iB \sqrt{c - ic \tan(e + fx)}}{c^2} + \frac{3iB - A}{c \sqrt{c - ic \tan(e + fx)}} \right) d \tan(e + fx)}{f} \\ & \quad \downarrow 2009 \\ & \frac{a^2 c \left(-\frac{2(3B + iA) \sqrt{c - ic \tan(e + fx)}}{c^2} - \frac{4(B + iA)}{c \sqrt{c - ic \tan(e + fx)}} + \frac{2B(c - ic \tan(e + fx))^{3/2}}{3c^3} \right)}{f} \end{aligned}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^2*(A + B*\text{Tan}[e + f*x])/(\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]]), x]$

3.752. $\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx$

```
output (a^2*c*((-4*(I*A + B))/(c*Sqrt[c - I*c*Tan[e + f*x]]) - (2*(I*A + 3*B)*Sqr
t[c - I*c*Tan[e + f*x]])/c^2 + (2*B*(c - I*c*Tan[e + f*x])^(3/2))/(3*c^3)
)/f
```

3.752.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.752.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.93

method	result
derivativedivides	$\frac{2ia^2 \left(-\frac{iB(c-ic \tan(fx+e))^{3/2}}{3} + 3i\sqrt{c-ic \tan(fx+e)} Bc - \sqrt{c-ic \tan(fx+e)} cA - \frac{2c^2(-iB+A)}{\sqrt{c-ic \tan(fx+e)}} \right)}{f c^2}$
default	$\frac{2ia^2 \left(-\frac{iB(c-ic \tan(fx+e))^{3/2}}{3} + 3i\sqrt{c-ic \tan(fx+e)} Bc - \sqrt{c-ic \tan(fx+e)} cA - \frac{2c^2(-iB+A)}{\sqrt{c-ic \tan(fx+e)}} \right)}{f c^2}$
parts	$\frac{2iA a^2 c \left(-\frac{1}{2c\sqrt{c-ic \tan(fx+e)}} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)}\sqrt{2}}{2\sqrt{c}}\right)}{4c^{3/2}} \right)}{f} + \frac{a^2(2iA+B) \left(-\frac{1}{\sqrt{c-ic \tan(fx+e)}} - \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)}\sqrt{2}}{2\sqrt{c}}\right)}{4c^{3/2}} \right)}{f}$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `2*I/f*a^2/c^2*(-1/3*I*B*(c-I*c*tan(f*x+e))^(3/2)+3*I*(c-I*c*tan(f*x+e))^(1/2)*B*c-(c-I*c*tan(f*x+e))^(1/2)*c*A-2*c^2*(A-I*B)/(c-I*c*tan(f*x+e))^(1/2))`

3.752.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.92

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx = \frac{2\sqrt{2}(3(iA + B)a^2 e^{(4i fx + 4i e)} + 3(3iA + 5B)a^2 e^{(2i fx + 2i e)} + 2(3iA + 5B)a^2) \sqrt{\frac{c}{e^{(2i fx + 2i e)} + 1}}}{3(cfe^{(2i fx + 2i e)} + cf)}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x,algorithm="fracas")`

output `-2/3*sqrt(2)*(3*(I*A + B)*a^2*e^(4*I*f*x + 4*I*e) + 3*(3*I*A + 5*B)*a^2*e^(2*I*f*x + 2*I*e) + 2*(3*I*A + 5*B)*a^2)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c*f*e^(2*I*f*x + 2*I*e) + c*f)`

3.752. $\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx$

3.752.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx$$

$$= -a^2 \left(\int \left(-\frac{A}{\sqrt{-ictan(e + fx) + c}} \right) dx + \int \frac{A \tan^2(e + fx)}{\sqrt{-ictan(e + fx) + c}} dx \right.$$

$$+ \int \left(-\frac{B \tan(e + fx)}{\sqrt{-ictan(e + fx) + c}} \right) dx + \int \frac{B \tan^3(e + fx)}{\sqrt{-ictan(e + fx) + c}} dx$$

$$\left. + \int \left(-\frac{2iA \tan(e + fx)}{\sqrt{-ictan(e + fx) + c}} \right) dx + \int \left(-\frac{2iB \tan^2(e + fx)}{\sqrt{-ictan(e + fx) + c}} \right) dx \right)$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(1/2),x)`

output `-a**2*(Integral(-A/sqrt(-I*c*tan(e + f*x) + c), x) + Integral(A*tan(e + f*x)**2/sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-B*tan(e + f*x)/sqrt(-I*c*tan(e + f*x) + c), x) + Integral(B*tan(e + f*x)**3/sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-2*I*A*tan(e + f*x)/sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-2*I*B*tan(e + f*x)**2/sqrt(-I*c*tan(e + f*x) + c), x))`

3.752.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.80

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx$$

$$= -\frac{2i \left(\frac{6(A-iB)a^2c}{\sqrt{-ictan(fx+e)+c}} + \frac{i(-ictan(fx+e)+c)^{\frac{3}{2}}Ba^2+3\sqrt{-ictan(fx+e)+c}(A-3iB)a^2c}{c} \right)}{3cf}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x, algorithm="maxima")`

output `-2/3*I*(6*(A - I*B)*a^2*c/sqrt(-I*c*tan(f*x + e) + c) + (I*(-I*c*tan(f*x + e) + c)^(3/2)*B*a^2 + 3*sqrt(-I*c*tan(f*x + e) + c)*(A - 3*I*B)*a^2*c)/(c*f)`

3.752. $\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} dx$

3.752.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx$$

$$= \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^2}{\sqrt{-ic \tan(fx + e) + c}} dx$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^2/sqrt(-I*c*tan(f*x
+ e) + c), x)`

3.752.9 Mupad [B] (verification not implemented)

Time = 9.19 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.74

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx =$$

$$\frac{2\sqrt{2}a^2 \sqrt{\frac{c}{\cos(2e+2fx)+1+\sin(2e+2fx)i}} (A6i + 10B + A \cos(2e + 2fx) 9i + A \cos(4e + 4fx) 3i + 15$$

3cf (c

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2)/(c - c*tan(e + f*x)*1
i)^(1/2),x)`

output `-(2*2^(1/2)*a^2*(c/(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))^(1/2)*(A*
6i + 10*B + A*cos(2*e + 2*f*x)*9i + A*cos(4*e + 4*f*x)*3i + 15*B*cos(2*e +
2*f*x) + 3*B*cos(4*e + 4*f*x) - 9*A*sin(2*e + 2*f*x) - 3*A*sin(4*e + 4*f*
x) + B*sin(2*e + 2*f*x)*15i + B*sin(4*e + 4*f*x)*3i))/(3*c*f*(cos(2*e + 2*
f*x) + sin(2*e + 2*f*x)*1i + 1))`

3.753
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ictan(e+fx))^{3/2}} dx$$

3.753.1 Optimal result 6865
 3.753.2 Mathematica [A] (verified) 6865
 3.753.3 Rubi [A] (verified) 6866
 3.753.4 Maple [A] (verified) 6867
 3.753.5 Fricas [A] (verification not implemented) 6868
 3.753.6 Sympy [F] 6868
 3.753.7 Maxima [A] (verification not implemented) 6869
 3.753.8 Giac [F] 6870
 3.753.9 Mupad [B] (verification not implemented) 6870

3.753.1 Optimal result

Integrand size = 43, antiderivative size = 101

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ictan(e + fx))^{3/2}} dx = -\frac{4a^2(iA + B)}{3f(c - ictan(e + fx))^{3/2}} + \frac{2a^2(iA + 3B)}{cf\sqrt{c - ictan(e + fx)}} + \frac{2a^2B\sqrt{c - ictan(e + fx)}}{c^2f}$$

output `2*a^2*(I*A+3*B)/c/f/(c-I*c*tan(f*x+e))^(1/2)+2*a^2*B*(c-I*c*tan(f*x+e))^(1/2)/c^2/f-4/3*a^2*(I*A+B)/f/(c-I*c*tan(f*x+e))^(3/2)`

3.753.2 Mathematica [A] (verified)

Time = 5.54 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.80

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ictan(e + fx))^{3/2}} dx = \frac{2a^2(-A + 10iB + 3(iA + 5B) \tan(e + fx) - 3iB \tan^2(e + fx))}{3cf(i + \tan(e + fx))\sqrt{c - ictan(e + fx)}}$$

input `Integrate[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(2*a^2*(-A + (10*I)*B + 3*(I*A + 5*B)*Tan[e + f*x] - (3*I)*B*Tan[e + f*x]^2))/(3*c*f*(I + Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])`

3.753.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.93, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{a(i \tan(e+fx)+1)(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^2 c \int \frac{(i \tan(e+fx)+1)(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{86} \\
 & \frac{a^2 c \int \left(\frac{2(A-iB)}{(c-ic \tan(e+fx))^{5/2}} - \frac{iB}{c^2 \sqrt{c-ic \tan(e+fx)}} + \frac{3iB-A}{c(c-ic \tan(e+fx))^{3/2}} \right) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^2 c \left(\frac{2(3B+iA)}{c^2 \sqrt{c-ic \tan(e+fx)}} - \frac{4(B+iA)}{3c(c-ic \tan(e+fx))^{3/2}} + \frac{2B \sqrt{c-ic \tan(e+fx)}}{c^3} \right)}{f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(a^2*c*((-4*(I*A + B))/(3*c*(c - I*c*Tan[e + f*x])^(3/2)) + (2*(I*A + 3*B))/(c^2*sqrt[c - I*c*Tan[e + f*x]]) + (2*B*sqrt[c - I*c*Tan[e + f*x]])/c^3)/f`

3.753.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.753.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.78

method	result
derivativedivides	$\frac{2ia^2 \left(-i\sqrt{c-ic \tan(fx+e)} B + \frac{c(-3iB+A)}{\sqrt{c-ic \tan(fx+e)}} - \frac{2c^2(-iB+A)}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} \right)}{f c^2}$
default	$\frac{2ia^2 \left(-i\sqrt{c-ic \tan(fx+e)} B + \frac{c(-3iB+A)}{\sqrt{c-ic \tan(fx+e)}} - \frac{2c^2(-iB+A)}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} \right)}{f c^2}$
parts	$\frac{2iA a^2 c \left(-\frac{1}{4c^2 \sqrt{c-ic \tan(fx+e)}} - \frac{1}{6c(c-ic \tan(fx+e))^{\frac{3}{2}}} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{8c^{\frac{5}{2}}} \right)}{f} + \frac{a^2(2iA+B)}{3(c-ic \tan(fx+e))}$

3.753.
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

output `2*I/f*a^2/c^2*(-I*(c-I*c*tan(f*x+e))^(1/2)*B+c*(A-3*I*B)/(c-I*c*tan(f*x+e))^(1/2)-2/3*c^2*(A-I*B)/(c-I*c*tan(f*x+e))^(3/2))`

3.753.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.80

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \frac{\sqrt{2}((-iA - B)a^2 e^{(4i fx + 4i e)} + (iA + 7B)a^2 e^{(2i fx + 2i e)})}{3c^2 f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x,algorithm="fracas")`

output `1/3*sqrt(2)*((-I*A - B)*a^2*e^(4*I*f*x + 4*I*e) + (I*A + 7*B)*a^2*e^(2*I*f*x + 2*I*e) - 2*(-I*A - 7*B)*a^2)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^2*f)`

3.753.6 Sympy [F]

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \\ & -a^2 \left(\int \left(-\frac{A}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} \right) dx \right. \\ & + \int \frac{A \tan^2(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} dx \\ & + \int \left(-\frac{B \tan(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} \right) dx \\ & + \int \frac{B \tan^3(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} dx \\ & + \int \left(-\frac{2iA \tan(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} \right) dx \\ & \left. + \int \left(-\frac{2iB \tan^2(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} \right) dx \right) \end{aligned}$$

3.753. $\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(3/2),x)`

output `-a**2*(Integral(-A/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(A*tan(e + f*x)**2/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(-B*tan(e + f*x)/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(B*tan(e + f*x)**3/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(-2*I*A*tan(e + f*x)/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(-2*I*B*tan(e + f*x)**2/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x))`

3.753.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.78

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \frac{2i \left(\frac{3i \sqrt{-ic \tan(fx+e)+c} Ba^2}{c} - \frac{3(-ic \tan(fx+e)+c)(A-3iB)a^2 - 2(A-iB)a^2 c}{(-ic \tan(fx+e)+c)^{3/2}} \right)}{3cf}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `-2/3*I*(3*I*sqrt(-I*c*tan(f*x + e) + c)*B*a^2/c - (3*(-I*c*tan(f*x + e) + c)*(A - 3*I*B)*a^2 - 2*(A - I*B)*a^2*c)/(-I*c*tan(f*x + e) + c)^(3/2))/(c*f)`

3.753.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^2}{(-ic \tan(fx + e) + c)^{3/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^2/(-I*c*tan(f*x + e)
+ c)^(3/2), x)`

3.753.9 Mupad [B] (verification not implemented)

Time = 9.01 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.56

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \frac{a^2 \sqrt{\frac{c(\cos(2e+2fx)+1) - \sin(2e+2fx)1i}{\cos(2e+2fx)+1}}}{(A2i + 14B + A \cos(2e + 2fx))^{3/2}}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2)/(c - c*tan(e + f*x)*1
i)^(3/2),x)`

output `(a^2*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) +
1))^(1/2)*(A*2i + 14*B + A*cos(2*e + 2*f*x)*1i - A*cos(4*e + 4*f*x)*1i +
7*B*cos(2*e + 2*f*x) - B*cos(4*e + 4*f*x) - A*sin(2*e + 2*f*x) + A*sin(4*e
+ 4*f*x) + B*sin(2*e + 2*f*x)*7i - B*sin(4*e + 4*f*x)*1i))/(3*c^2*f)`

3.754
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ictan(e+fx))^{5/2}} dx$$

3.754.1 Optimal result 6871
 3.754.2 Mathematica [A] (verified) 6871
 3.754.3 Rubi [A] (verified) 6872
 3.754.4 Maple [A] (verified) 6873
 3.754.5 Fricas [A] (verification not implemented) 6874
 3.754.6 Sympy [F] 6875
 3.754.7 Maxima [A] (verification not implemented) 6876
 3.754.8 Giac [F] 6876
 3.754.9 Mupad [B] (verification not implemented) 6876

3.754.1 Optimal result

Integrand size = 43, antiderivative size = 103

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ictan(e + fx))^{5/2}} dx = -\frac{4a^2(iA + B)}{5f(c - ictan(e + fx))^{5/2}} + \frac{2a^2(iA + 3B)}{3cf(c - ictan(e + fx))^{3/2}} - \frac{2a^2B}{c^2f\sqrt{c - ictan(e + fx)}}$$

output `-2*a^2*B/c^2/f/(c-I*c*tan(f*x+e))^(1/2)-4/5*a^2*(I*A+B)/f/(c-I*c*tan(f*x+e))^(5/2)+2/3*a^2*(I*A+3*B)/c/f/(c-I*c*tan(f*x+e))^(3/2)`

3.754.2 Mathematica [A] (verified)

Time = 5.74 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.75

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ictan(e + fx))^{5/2}} dx = \frac{2a^2(-iA - 6B + 5(A + 3iB) \tan(e + fx) + 15B \tan^2(e + fx))}{15c^2f(i + \tan(e + fx))^2\sqrt{c - ictan(e + fx)}}$$

input `Integrate[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(5/2),x]`

output `(-2*a^2*((-I)*A - 6*B + 5*(A + (3*I)*B)*Tan[e + f*x] + 15*B*Tan[e + f*x]^2))/(15*c^2*f*(I + Tan[e + f*x])^2*Sqrt[c - I*c*Tan[e + f*x]])`

3.754.
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ictan(e+fx))^{5/2}} dx$$

3.754.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.93, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{a(i \tan(e+fx)+1)(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^2 c \int \frac{(i \tan(e+fx)+1)(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{86} \\
 & \frac{a^2 c \int \left(\frac{2(A-iB)}{(c-ic \tan(e+fx))^{7/2}} - \frac{iB}{c^2(c-ic \tan(e+fx))^{3/2}} + \frac{3iB-A}{c(c-ic \tan(e+fx))^{5/2}} \right) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^2 c \left(\frac{2(3B+iA)}{3c^2(c-ic \tan(e+fx))^{3/2}} - \frac{4(B+iA)}{5c(c-ic \tan(e+fx))^{5/2}} - \frac{2B}{c^3 \sqrt{c-ic \tan(e+fx)}} \right)}{f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(5/2),x]`

output `(a^2*c*((-4*(I*A + B))/(5*c*(c - I*c*Tan[e + f*x])^(5/2)) + (2*(I*A + 3*B))/(3*c^2*(c - I*c*Tan[e + f*x])^(3/2)) - (2*B)/(c^3*sqrt[c - I*c*Tan[e + f*x]])))/f`

3.754. $\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$

3.754.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.754.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.78

method	result
derivativedivides	$\frac{2ia^2 \left(-\frac{2c^2(-iB+A)}{5(c-ic \tan(fx+e))^{\frac{5}{2}}} + \frac{iB}{\sqrt{c-ic \tan(fx+e)}} + \frac{c(-3iB+A)}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} \right)}{f c^2}$
default	$\frac{2ia^2 \left(-\frac{2c^2(-iB+A)}{5(c-ic \tan(fx+e))^{\frac{5}{2}}} + \frac{iB}{\sqrt{c-ic \tan(fx+e)}} + \frac{c(-3iB+A)}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} \right)}{f c^2}$
risch	$-\frac{a^2(3iA e^{4i(fx+e)} + 3B e^{4i(fx+e)} + iA e^{2i(fx+e)} - 9B e^{2i(fx+e)} - 2iA + 18B)\sqrt{2}}{30c^2 \sqrt{e^{2i(fx+e)} + 1}} f$
parts	$\frac{2iA a^2 c \left(-\frac{1}{8c^3 \sqrt{c-ic \tan(fx+e)}} - \frac{1}{12c^2 (c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{1}{10c (c-ic \tan(fx+e))^{\frac{5}{2}}} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{16c^{\frac{7}{2}}} \right)}{f} +$

3.754. $\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

output `2*I/f*a^2/c^2*(-2/5*c^2*(A-I*B)/(c-I*c*tan(f*x+e))^(5/2)+I*B/(c-I*c*tan(f*x+e))^(1/2)+1/3*c*(A-3*I*B)/(c-I*c*tan(f*x+e))^(3/2))`

3.754.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.99

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx =$$

$$\frac{\sqrt{2}(3(iA + B)a^2 e^{(6ifx+6ie)} + 2(2iA - 3B)a^2 e^{(4ifx+4ie)} - (iA - 9B)a^2 e^{(2ifx+2ie)} + 2(-iA + 9B)a^2)}{30c^3 f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,algorithm="fracas")`

output `-1/30*sqrt(2)*(3*(I*A + B)*a^2*e^(6*I*f*x + 6*I*e) + 2*(2*I*A - 3*B)*a^2*e^(4*I*f*x + 4*I*e) - (I*A - 9*B)*a^2*e^(2*I*f*x + 2*I*e) + 2*(-I*A + 9*B)*a^2)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^3*f)`

3.754.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx =$$

$$-a^2 \left(\int \left(-\frac{A}{-c^2 \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx) - 2ic^2 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c^2 \sqrt{-ic \tan(e + fx) + c}} \right. \right.$$

$$+ \int \frac{A \tan^2(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx) - 2ic^2 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c^2 \sqrt{-ic \tan(e + fx) + c}} dx$$

$$+ \int \left(-\frac{B \tan(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx) - 2ic^2 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c^2 \sqrt{-ic \tan(e + fx) + c}} \right.$$

$$+ \int \frac{B \tan^3(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx) - 2ic^2 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c^2 \sqrt{-ic \tan(e + fx) + c}} dx$$

$$+ \int \left(-\frac{2iA \tan(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx) - 2ic^2 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c^2 \sqrt{-ic \tan(e + fx) + c}} \right.$$

$$+ \int \left(-\frac{2iB \tan^2(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx) - 2ic^2 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c^2 \sqrt{-ic \tan(e + fx) + c}} \right.$$

```
input integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(5/2),x)
```

```
output -a**2*(Integral(-A/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 2*I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**2*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(A*tan(e + f*x)**2/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 2*I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**2*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(-B*tan(e + f*x)/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 2*I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**2*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(B*tan(e + f*x)**3/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 2*I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**2*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(-2*I*A*tan(e + f*x)/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 2*I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**2*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(-2*I*B*tan(e + f*x)**2/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 2*I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**2*sqrt(-I*c*tan(e + f*x) + c)), x))
```

3.754.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.74

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \frac{2i (15i (-ic \tan(fx + e) + c)^2 Ba^2 + 5 (-ic \tan(fx + e) + c) (A - 3iB) a^2 c - 6(A - iB) a^2 c^2)}{15 (-ic \tan(fx + e) + c)^{5/2} c^2 f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x
, algorithm="maxima")`

output `2/15*I*(15*I*(-I*c*tan(f*x + e) + c)^2*B*a^2 + 5*(-I*c*tan(f*x + e) + c)*(
A - 3*I*B)*a^2*c - 6*(A - I*B)*a^2*c^2)/((-I*c*tan(f*x + e) + c)^(5/2)*c^2
*f)`

3.754.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^2}{(-ic \tan(fx + e) + c)^{5/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^2/(-I*c*tan(f*x + e)
+ c)^(5/2), x)`

3.754.9 Mupad [B] (verification not implemented)

Time = 9.77 (sec) , antiderivative size = 208, normalized size of antiderivative = 2.02

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \frac{a^2 \sqrt{\frac{c(\cos(2e+2fx)+1)-\sin(2e+2fx)1i}{\cos(2e+2fx)+1}} (18B - A2i - A \cos(2e + 2fx) 1i + A \cos(4e + 4fx) 4i + A \cos(6e + 6fx) 6i)}{15 (-ic \tan(fx + e) + c)^{5/2} c^2 f}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2)/(c - c*tan(e + f*x)*1i)^(5/2),x)`

output `-(a^2*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(18*B - A*2i - A*cos(2*e + 2*f*x)*1i + A*cos(4*e + 4*f*x)*4i + A*cos(6*e + 6*f*x)*3i + 9*B*cos(2*e + 2*f*x) - 6*B*cos(4*e + 4*f*x) + 3*B*cos(6*e + 6*f*x) + A*sin(2*e + 2*f*x) - 4*A*sin(4*e + 4*f*x) - 3*A*sin(6*e + 6*f*x) + B*sin(2*e + 2*f*x)*9i - B*sin(4*e + 4*f*x)*6i + B*sin(6*e + 6*f*x)*3i))/(30*c^3*f)`

3.755
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ictan(e+fx))^{7/2}} dx$$

3.755.1 Optimal result	6878
3.755.2 Mathematica [A] (verified)	6878
3.755.3 Rubi [A] (verified)	6879
3.755.4 Maple [A] (verified)	6880
3.755.5 Fricas [A] (verification not implemented)	6881
3.755.6 Sympy [F]	6882
3.755.7 Maxima [A] (verification not implemented)	6883
3.755.8 Giac [F]	6884
3.755.9 Mupad [B] (verification not implemented)	6884

3.755.1 Optimal result

Integrand size = 43, antiderivative size = 105

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx = -\frac{4a^2(iA + B)}{7f(c - ictan(e + fx))^{7/2}} + \frac{2a^2(iA + 3B)}{5cf(c - ictan(e + fx))^{5/2}} - \frac{2a^2B}{3c^2f(c - ictan(e + fx))^{3/2}}$$

output
$$-4/7*a^2*(I*A+B)/f/(c-I*c*tan(f*x+e))^(7/2)+2/5*a^2*(I*A+3*B)/c/f/(c-I*c*tan(f*x+e))^(5/2)-2/3*a^2*B/c^2/f/(c-I*c*tan(f*x+e))^(3/2)$$

3.755.2 Mathematica [A] (verified)

Time = 6.02 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.75

$$\int \frac{(a + ia \tan(e + fx))^2(A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx = \frac{2a^2(-9A + 2iB + 7(-3iA + B) \tan(e + fx) - 35iB \tan(e + fx)^2)}{105c^3f(i + \tan(e + fx))^3\sqrt{c - ictan(e + fx)}}$$

input
$$\text{Integrate}[\frac{(a + I*a*\text{Tan}[e + f*x])^2*(A + B*\text{Tan}[e + f*x])}{(c - I*c*\text{Tan}[e + f*x])^{7/2}}, x]$$

output
$$\frac{(2*a^2*(-9*A + (2*I)*B + 7*((-3*I)*A + B)*\text{Tan}[e + f*x] - (35*I)*B*\text{Tan}[e + f*x]^2))/(105*c^3*f*(I + \text{Tan}[e + f*x])^3*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])}{1}$$

3.755.
$$\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ictan(e+fx))^{7/2}} dx$$

3.755.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.93, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{a(i \tan(e+fx)+1)(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{9/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^2 c \int \frac{(i \tan(e+fx)+1)(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{9/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{86} \\
 & \frac{a^2 c \int \left(\frac{2(A-iB)}{(c-ic \tan(e+fx))^{9/2}} - \frac{iB}{c^2(c-ic \tan(e+fx))^{5/2}} + \frac{3iB-A}{c(c-ic \tan(e+fx))^{7/2}} \right) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^2 c \left(\frac{2(3B+iA)}{5c^2(c-ic \tan(e+fx))^{5/2}} - \frac{4(B+iA)}{7c(c-ic \tan(e+fx))^{7/2}} - \frac{2B}{3c^3(c-ic \tan(e+fx))^{3/2}} \right)}{f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])^2*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(7/2),x]`

output `(a^2*c*((-4*(I*A + B))/(7*c*(c - I*c*Tan[e + f*x])^(7/2)) + (2*(I*A + 3*B))/(5*c^2*(c - I*c*Tan[e + f*x])^(5/2)) - (2*B)/(3*c^3*(c - I*c*Tan[e + f*x])^(3/2)))/f`

3.755.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.755.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.76

method	result
derivativedivides	$\frac{2ia^2 \left(\frac{iB}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{2c^2(-iB+A)}{7(c-ic \tan(fx+e))^{\frac{7}{2}}} + \frac{c(-3iB+A)}{5(c-ic \tan(fx+e))^{\frac{5}{2}}} \right)}{f c^2}$
default	$\frac{2ia^2 \left(\frac{iB}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{2c^2(-iB+A)}{7(c-ic \tan(fx+e))^{\frac{7}{2}}} + \frac{c(-3iB+A)}{5(c-ic \tan(fx+e))^{\frac{5}{2}}} \right)}{f c^2}$
risch	$-\frac{a^2(15iA e^{6i(fx+e)} + 15B e^{6i(fx+e)} + 24iA e^{4i(fx+e)} - 18B e^{4i(fx+e)} + 3iA e^{2i(fx+e)} - 11B e^{2i(fx+e)} - 6iA + 22B)\sqrt{2}}{420c^3 \sqrt{\frac{c}{e^{2i(fx+e)} + 1}}} f$
parts	$2iA a^2 c \left(-\frac{1}{16c^4 \sqrt{c-ic \tan(fx+e)}} - \frac{1}{24c^3 (c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{1}{20c^2 (c-ic \tan(fx+e))^{\frac{5}{2}}} - \frac{1}{14c (c-ic \tan(fx+e))^{\frac{7}{2}}} + \frac{\sqrt{2} \operatorname{arctanh}\left(\sqrt{\frac{c-ic \tan(fx+e)}{e^{2i(fx+e)} + 1}}\right)}{f} \right)$

3.755. $\int \frac{(a+ia \tan(e+fx))^2(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$

input `int((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x,method=_RETURNVERBOSE)`

output `2*I/f*a^2/c^2*(1/3*I*B/(c-I*c*tan(f*x+e))^(3/2)-2/7*c^2*(A-I*B)/(c-I*c*tan(f*x+e))^(7/2)+1/5*c*(A-3*I*B)/(c-I*c*tan(f*x+e))^(5/2))`

3.755.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.17

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx =$$

$$\frac{\sqrt{2}(15(iA + B)a^2 e^{(8i fx + 8ie)} + 3(13iA - B)a^2 e^{(6i fx + 6ie)} - (-27iA + 29B)a^2 e^{(4i fx + 4ie)} - (3iA - 11B)a^2 e^{(2i fx + 2ie)})}{420 c^4 f}$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x,algorithm="fracas")`

output `-1/420*sqrt(2)*(15*(I*A + B)*a^2*e^(8*I*f*x + 8*I*e) + 3*(13*I*A - B)*a^2*e^(6*I*f*x + 6*I*e) - (-27*I*A + 29*B)*a^2*e^(4*I*f*x + 4*I*e) - (3*I*A - 11*B)*a^2*e^(2*I*f*x + 2*I*e) + 2*(-3*I*A + 11*B)*a^2)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^4*f)`

3.755.6 Sympy [F]

$$\begin{aligned}
& \int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \\
& -a^2 \left(\int \left(-\frac{A}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx)}} \right. \right. \\
& + \int \frac{A \tan^2(e + fx)}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx)}} \\
& + \int \left(-\frac{B \tan(e + fx)}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx)}} \right. \\
& + \int \frac{B \tan^3(e + fx)}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx)}} \\
& + \int \left(-\frac{2iA \tan(e + fx)}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx)}} \right. \\
& \left. \left. + \int \left(-\frac{2iB \tan^2(e + fx)}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx)}} \right) \right) \right)
\end{aligned}$$

input `integrate((a+I*a*tan(f*x+e))**2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(7/2),x)`

```

output -a**2*(Integral(-A/(I*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3 - 3
*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 3*I*c**3*sqrt(-I*c*tan
(e + f*x) + c)*tan(e + f*x) + c**3*sqrt(-I*c*tan(e + f*x) + c)), x) + Inte
gral(A*tan(e + f*x)**2/(I*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3
- 3*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 3*I*c**3*sqrt(-I*c
*tan(e + f*x) + c)*tan(e + f*x) + c**3*sqrt(-I*c*tan(e + f*x) + c)), x) +
Integral(-B*tan(e + f*x)/(I*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)*
**3 - 3*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 3*I*c**3*sqrt(-I
*c*tan(e + f*x) + c)*tan(e + f*x) + c**3*sqrt(-I*c*tan(e + f*x) + c)), x)
+ Integral(B*tan(e + f*x)**3/(I*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f
*x)**3 - 3*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 3*I*c**3*sqr
t(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**3*sqrt(-I*c*tan(e + f*x) + c)),
x) + Integral(-2*I*A*tan(e + f*x)/(I*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan
(e + f*x)**3 - 3*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 3*I*c*
**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**3*sqrt(-I*c*tan(e + f*x)
+ c)), x) + Integral(-2*I*B*tan(e + f*x)**2/(I*c**3*sqrt(-I*c*tan(e + f*x)
+ c)*tan(e + f*x)**3 - 3*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2
- 3*I*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**3*sqrt(-I*c*tan(
e + f*x) + c)), x))

```

3.755.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.72

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \frac{2i (35i (-ic \tan(fx + e) + c)^2 Ba^2 + 21 (-ic \tan(fx + e) - 105 (-ic \tan(fx + e) -$$

```

input integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x
, algorithm="maxima")

```

```

output 2/105*I*(35*I*(-I*c*tan(f*x + e) + c)^2*B*a^2 + 21*(-I*c*tan(f*x + e) + c)
*(A - 3*I*B)*a^2*c - 30*(A - I*B)*a^2*c^2)/((-I*c*tan(f*x + e) + c)^(7/2)*
c^2*f)

```

3.755.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^2}{(-ic \tan(fx + e) + c)^{7/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^2*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^2/(-I*c*tan(f*x + e)
+ c)^(7/2), x)`

3.755.9 Mupad [B] (verification not implemented)

Time = 10.12 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.59

$$\int \frac{(a + ia \tan(e + fx))^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx =$$

$$-\sqrt{c - \frac{c \sin(e + fx)}{\cos(e + fx)}} \left(-\frac{a^2 (3A + B 11i) 1i}{210 c^4 f} \right.$$

$$-\frac{a^2 e^{2i+fx 2i} (3A + B 11i) 1i}{420 c^4 f} + \frac{a^2 e^{6i+fx 6i} (13A + B 1i) 1i}{140 c^4 f}$$

$$\left. + \frac{a^2 e^{4i+fx 4i} (27A + B 29i) 1i}{420 c^4 f} + \frac{a^2 e^{8i+fx 8i} (A - B 1i) 1i}{28 c^4 f} \right)$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^2)/(c - c*tan(e + f*x)*1
i)^(7/2),x)`

output `-(c - (c*sin(e + f*x)*1i)/cos(e + f*x))^(1/2)*((a^2*exp(e*6i + f*x*6i)*(13
*A + B*1i)*1i)/(140*c^4*f) - (a^2*exp(e*2i + f*x*2i)*(3*A + B*11i)*1i)/(42
0*c^4*f) - (a^2*(3*A + B*11i)*1i)/(210*c^4*f) + (a^2*exp(e*4i + f*x*4i)*(2
7*A + B*29i)*1i)/(420*c^4*f) + (a^2*exp(e*8i + f*x*8i)*(A - B*1i)*1i)/(28*
c^4*f))`

3.756 $\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx))^{7/2} dx$

3.756.1 Optimal result	6885
3.756.2 Mathematica [A] (verified)	6886
3.756.3 Rubi [A] (verified)	6886
3.756.4 Maple [A] (verified)	6888
3.756.5 Fricas [A] (verification not implemented)	6888
3.756.6 Sympy [F]	6889
3.756.7 Maxima [A] (verification not implemented)	6890
3.756.8 Giac [F]	6890
3.756.9 Mupad [B] (verification not implemented)	6891

3.756.1 Optimal result

Integrand size = 43, antiderivative size = 144

$$\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx))^{7/2} dx = \frac{8a^3(iA+B)(c-ictan(e+fx))^{7/2}}{7f} - \frac{8a^3(iA+2B)(c-ictan(e+fx))^{9/2}}{9cf} + \frac{2a^3(iA+5B)(c-ictan(e+fx))^{11/2}}{11c^2f} - \frac{2a^3B(c-ictan(e+fx))^{13/2}}{13c^3f}$$

```
output 8/7*a^3*(I*A+B)*(c-I*c*tan(f*x+e))^(7/2)/f-8/9*a^3*(I*A+2*B)*(c-I*c*tan(f*x+e))^(9/2)/c/f+2/11*a^3*(I*A+5*B)*(c-I*c*tan(f*x+e))^(11/2)/c^2/f-2/13*a^3*B*(c-I*c*tan(f*x+e))^(13/2)/c^3/f
```

3.756.2 Mathematica [A] (verified)

Time = 6.80 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.85

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ictan(e + fx))^{7/2} dx = \frac{a^3 c^4 \sec^6(e + fx) (i \cos(4(e + fx)) + \sin(4(e + fx))) (2(572A + 737iB + 7(169iA + 86B) \tan(e + fx)) + \cos[2(e + fx)])}{9009 f \sqrt{c - ictan(e + fx)}}$$

input `Integrate[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2),x]`

output `(a^3*c^4*Sec[e + f*x]^6*(I*Cos[4*(e + f*x)] + Sin[4*(e + f*x)])*(2*(572*A + (737*I)*B + 7*((169*I)*A + 86*B)*Tan[e + f*x]) + Cos[2*(e + f*x)]*(2782*A - (2558*I)*B + 14*((169*I)*A + 185*B)*Tan[e + f*x]))/(9009*f*Sqrt[c - I*c*Tan[e + f*x]])`

3.756.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.91, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))^3 (c - ictan(e + fx))^{7/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))^3 (c - ictan(e + fx))^{7/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int a^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^{5/2} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{a^3 c \int (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) (c - ictan(e + fx))^{5/2} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{86} \end{aligned}$$

3.756. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ictan(e + fx))^{7/2} dx$

$$a^3 c \int \left(\frac{iB(c-ictan(e+fx))^{11/2}}{c^3} + \frac{(A-5iB)(c-ictan(e+fx))^{9/2}}{c^2} - \frac{4(A-2iB)(c-ictan(e+fx))^{7/2}}{c} + 4(A-iB)(c-ictan(e+fx)) \right) dx$$

↓ 2009

$$a^3 c \left(\frac{2(5B+iA)(c-ictan(e+fx))^{11/2}}{11c^3} - \frac{8(2B+iA)(c-ictan(e+fx))^{9/2}}{9c^2} + \frac{8(B+iA)(c-ictan(e+fx))^{7/2}}{7c} - \frac{2B(c-ictan(e+fx))^{13/2}}{13c^4} \right) dx$$

input `Int[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2), x]`

output `(a^3*c*((8*(I*A + B)*(c - I*c*Tan[e + f*x])^(7/2))/(7*c) - (8*(I*A + 2*B)*(c - I*c*Tan[e + f*x])^(9/2))/(9*c^2) + (2*(I*A + 5*B)*(c - I*c*Tan[e + f*x])^(11/2))/(11*c^3) - (2*B*(c - I*c*Tan[e + f*x])^(13/2))/(13*c^4))/f`

3.756.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x],
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.756.4 Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.84

method	result
derivativedivides	$\frac{2ia^3 \left(\frac{iB(c-ic \tan(fx+e))^{13}}{13} + \frac{(-5iBc+cA)(c-ic \tan(fx+e))^{11}}{11} + \frac{(-4(-iBc+cA)c+4iBc^2)(c-ic \tan(fx+e))^9}{9} + \frac{4(-iBc+cA)c^7}{7} \right)}{fc^3}$
default	$\frac{2ia^3 \left(\frac{iB(c-ic \tan(fx+e))^{13}}{13} + \frac{(-5iBc+cA)(c-ic \tan(fx+e))^{11}}{11} + \frac{(-4(-iBc+cA)c+4iBc^2)(c-ic \tan(fx+e))^9}{9} + \frac{4(-iBc+cA)c^7}{7} \right)}{fc^3}$
parts	$\frac{2ia^3 Ac \left(-\frac{(c-ic \tan(fx+e))^{\frac{5}{2}}}{5} - \frac{2c(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} - 4\sqrt{c-ic \tan(fx+e)}c^2 + 4c^{\frac{5}{2}}\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)}\sqrt{2}}{2\sqrt{c}}\right) \right)}{f}$

```
input int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, metho
d=_RETURNVERBOSE)
```

```
output 2*I/f*a^3/c^3*(1/13*I*B*(c-I*c*tan(f*x+e))^(13/2)+1/11*(-5*I*B*c+c*A)*(c-I
*c*tan(f*x+e))^(11/2)+1/9*(-4*(-I*B*c+c*A)*c+4*I*B*c^2)*(c-I*c*tan(f*x+e))
^(9/2)+4/7*(-I*B*c+c*A)*c^2*(c-I*c*tan(f*x+e))^(7/2))
```

3.756.5 Fracas [A] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.24

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx =$$

$$\frac{64\sqrt{2}(1287(-iA - B)a^3c^3e^{(6ifx+6ie)} + 143(-13iA + B)a^3c^3e^{(4ifx+4ie)} + 52(-13iA + B)a^3c^3e^{(2ifx+2ie)} + 15fe^{(4ifx+4ie)})}{9009(fe^{(12ifx+12ie)} + 6fe^{(10ifx+10ie)} + 15fe^{(8ifx+8ie)} + 20fe^{(6ifx+6ie)} + 15fe^{(4ifx+4ie)})}$$

```
input integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x
, algorithm="fricas")
```

3.756. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx$

output `-64/9009*sqrt(2)*(1287*(-I*A - B)*a^3*c^3*e^(6*I*f*x + 6*I*e) + 143*(-13*I*A + B)*a^3*c^3*e^(4*I*f*x + 4*I*e) + 52*(-13*I*A + B)*a^3*c^3*e^(2*I*f*x + 2*I*e) + 8*(-13*I*A + B)*a^3*c^3)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(f*e^(12*I*f*x + 12*I*e) + 6*f*e^(10*I*f*x + 10*I*e) + 15*f*e^(8*I*f*x + 8*I*e) + 20*f*e^(6*I*f*x + 6*I*e) + 15*f*e^(4*I*f*x + 4*I*e) + 6*f*e^(2*I*f*x + 2*I*e) + f)`

3.756.6 Sympy [F]

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx =$$

$$-ia^3 \left(\int iAc^3 \sqrt{-ic \tan(e + fx) + c} dx \right.$$

$$+ \int 3iAc^3 \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx) dx$$

$$+ \int 3iAc^3 \sqrt{-ic \tan(e + fx) + c} \tan^4(e + fx) dx$$

$$+ \int iAc^3 \sqrt{-ic \tan(e + fx) + c} \tan^6(e + fx) dx$$

$$+ \int iBc^3 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx) dx$$

$$+ \int 3iBc^3 \sqrt{-ic \tan(e + fx) + c} \tan^3(e + fx) dx$$

$$+ \int 3iBc^3 \sqrt{-ic \tan(e + fx) + c} \tan^5(e + fx) dx$$

$$\left. + \int iBc^3 \sqrt{-ic \tan(e + fx) + c} \tan^7(e + fx) dx \right)$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(7/2),x)`

output `-I*a**3*(Integral(I*A*c**3*sqrt(-I*c*tan(e + f*x) + c), x) + Integral(3*I*A*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x) + Integral(3*I*A*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4, x) + Integral(I*A*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**6, x) + Integral(I*B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(3*I*B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x) + Integral(3*I*B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**5, x) + Integral(I*B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**7, x))`

3.756. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx$

3.756.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.72

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ictan(e + fx))^{7/2} dx = \frac{2i \left(693i (-ictan(fx + e) + c)^{\frac{13}{2}} B a^3 + 819 (-ictan(fx + e) + c)^{\frac{11}{2}} (A - 5iB) a^3 c - 4004 (-ictan(fx + e) + c)^{\frac{9}{2}} (A - 2iB) a^3 c^2 + 5148 (-ictan(fx + e) + c)^{\frac{7}{2}} (A - iB) a^3 c^3 \right)}{9009 c^3 f}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="maxima")`

output `2/9009*I*(693*I*(-I*c*tan(f*x + e) + c)^(13/2)*B*a^3 + 819*(-I*c*tan(f*x + e) + c)^(11/2)*(A - 5*I*B)*a^3*c - 4004*(-I*c*tan(f*x + e) + c)^(9/2)*(A - 2*I*B)*a^3*c^2 + 5148*(-I*c*tan(f*x + e) + c)^(7/2)*(A - I*B)*a^3*c^3)/(c^3*f)`

3.756.8 Giac [F]

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ictan(e + fx))^{7/2} dx = \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^3 (-ictan(fx + e) + c)^{\frac{7}{2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^3*(-I*c*tan(f*x + e) + c)^(7/2), x)`

3.756.9 Mupad [B] (verification not implemented)

Time = 12.90 (sec) , antiderivative size = 349, normalized size of antiderivative = 2.42

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx =$$

$$-\frac{\left(\frac{a^3 c^3 (A-B 1i) 64i}{9f} + \frac{a^3 c^3 (A-B 3i) 64i}{9f}\right) \sqrt{c + \frac{c(e^{e 2i+fx 2i} 1i-i)}{e^{e 2i+fx 2i}+1}}}{(e^{e 2i+fx 2i} + 1)^4}$$

$$-\frac{\left(\frac{a^3 c^3 (A-B 1i) 64i}{13f} - \frac{a^3 c^3 (A+B 1i) 64i}{13f}\right) \sqrt{c + \frac{c(e^{e 2i+fx 2i} 1i-i)}{e^{e 2i+fx 2i}+1}}}{(e^{e 2i+fx 2i} + 1)^6}$$

$$+\frac{\left(\frac{256 B a^3 c^3}{11f} + \frac{a^3 c^3 (A-B 1i) 64i}{11f}\right) \sqrt{c + \frac{c(e^{e 2i+fx 2i} 1i-i)}{e^{e 2i+fx 2i}+1}}}{(e^{e 2i+fx 2i} + 1)^5}$$

$$+\frac{a^3 c^3 (A - B 1i) \sqrt{c + \frac{c(e^{e 2i+fx 2i} 1i-i)}{e^{e 2i+fx 2i}+1}} 64i}{7 f (e^{e 2i+fx 2i} + 1)^3}$$

```
input int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i)
^(7/2),x)
```

```
output (((a^3*c^3*(A - B*1i)*64i)/(11*f) + (256*B*a^3*c^3)/(11*f))*(c + (c*(exp(e
*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(1/2))/(exp(e*2i + f*
x*2i) + 1)^5 - (((a^3*c^3*(A - B*1i)*64i)/(13*f) - (a^3*c^3*(A + B*1i)*64i
)/(13*f))*(c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1
))^(1/2))/(exp(e*2i + f*x*2i) + 1)^6 - (((a^3*c^3*(A - B*1i)*64i)/(9*f) +
(a^3*c^3*(A - B*3i)*64i)/(9*f))*(c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(
exp(e*2i + f*x*2i) + 1))^(1/2))/(exp(e*2i + f*x*2i) + 1)^4 + (a^3*c^3*(A -
B*1i)*(c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(
1/2)*64i)/(7*f*(exp(e*2i + f*x*2i) + 1)^3)
```

3.757 $\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx))^{5/2} dx$

3.757.1 Optimal result	6892
3.757.2 Mathematica [A] (verified)	6893
3.757.3 Rubi [A] (verified)	6893
3.757.4 Maple [A] (verified)	6895
3.757.5 Fricas [A] (verification not implemented)	6895
3.757.6 Sympy [F]	6896
3.757.7 Maxima [A] (verification not implemented)	6897
3.757.8 Giac [F]	6897
3.757.9 Mupad [B] (verification not implemented)	6898

3.757.1 Optimal result

Integrand size = 43, antiderivative size = 144

$$\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx = \frac{8a^3(iA + B)(c - ictan(e + fx))^{5/2}}{5f} - \frac{8a^3(iA + 2B)(c - ictan(e + fx))^{7/2}}{7cf} + \frac{2a^3(iA + 5B)(c - ictan(e + fx))^{9/2}}{9c^2f} - \frac{2a^3B(c - ictan(e + fx))^{11/2}}{11c^3f}$$

```
output 8/5*a^3*(I*A+B)*(c-I*c*tan(f*x+e))^(5/2)/f-8/7*a^3*(I*A+2*B)*(c-I*c*tan(f*x+e))^(7/2)/c/f+2/9*a^3*(I*A+5*B)*(c-I*c*tan(f*x+e))^(9/2)/c^2/f-2/11*a^3*B*(c-I*c*tan(f*x+e))^(11/2)/c^3/f
```

3.757.2 Mathematica [A] (verified)

Time = 5.97 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.89

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ict \tan(e + fx))^{5/2} dx = \frac{a^3 c^3 \sec^6(e + fx) (143(11iA + B) \cos(e + fx) + (781iA + 701B) \cos(3(e + fx)) - 10(121A - 11B) \cos(5(e + fx)))}{3465f \sqrt{c - ict \tan(e + fx)}}$$

input `Integrate[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2), x]`

output `(a^3*c^3*Sec[e + f*x]^6*(143*((11*I)*A + B)*Cos[e + f*x] + ((781*I)*A + 701*B)*Cos[3*(e + f*x)] - 10*(121*A - (74*I)*B + (121*A - (137*I)*B)*Cos[2*(e + f*x)])*Sin[e + f*x]*(Cos[3*(e + f*x)] - I*Sin[3*(e + f*x)])/(3465*f*Sqrt[c - I*c*Tan[e + f*x]])`

3.757.3 Rubi [A] (verified)Time = 0.38 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.91, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))^3 (c - ict \tan(e + fx))^{5/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))^3 (c - ict \tan(e + fx))^{5/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int a^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) (c - ict \tan(e + fx))^{3/2} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{a^3 c \int (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) (c - ict \tan(e + fx))^{3/2} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{86} \end{aligned}$$

3.757. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ict \tan(e + fx))^{5/2} dx$

$$a^3 c \int \left(\frac{iB(c-ictan(e+fx))^{9/2}}{c^3} + \frac{(A-5iB)(c-ictan(e+fx))^{7/2}}{c^2} - \frac{4(A-2iB)(c-ictan(e+fx))^{5/2}}{c} + 4(A-iB)(c-ictan(e+fx)) \right) f$$

↓ 2009

$$a^3 c \left(\frac{2(5B+iA)(c-ictan(e+fx))^{9/2}}{9c^3} - \frac{8(2B+iA)(c-ictan(e+fx))^{7/2}}{7c^2} + \frac{8(B+iA)(c-ictan(e+fx))^{5/2}}{5c} - \frac{2B(c-ictan(e+fx))^{11/2}}{11c^4} \right) f$$

input `Int[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2), x]`

output `(a^3*c*((8*(I*A + B)*(c - I*c*Tan[e + f*x])^(5/2))/(5*c) - (8*(I*A + 2*B)*(c - I*c*Tan[e + f*x])^(7/2))/(7*c^2) + (2*(I*A + 5*B)*(c - I*c*Tan[e + f*x])^(9/2))/(9*c^3) - (2*B*(c - I*c*Tan[e + f*x])^(11/2))/(11*c^4))/f`

3.757.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.757.4 Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.84

method	result
derivativeldivides	$\frac{2ia^3 \left(\frac{iB(c-ic \tan(fx+e))^{11}}{11} + \frac{(-5iBc+cA)(c-ic \tan(fx+e))^9}{9} + \frac{(-4(-iBc+cA)c+4iBc^2)(c-ic \tan(fx+e))^7}{7} + \frac{4(-iBc+cA)c^2}{4} \right)}{f c^3}$
default	$\frac{2ia^3 \left(\frac{iB(c-ic \tan(fx+e))^{11}}{11} + \frac{(-5iBc+cA)(c-ic \tan(fx+e))^9}{9} + \frac{(-4(-iBc+cA)c+4iBc^2)(c-ic \tan(fx+e))^7}{7} + \frac{4(-iBc+cA)c^2}{4} \right)}{f c^3}$
parts	$\frac{2ia^3 Ac \left(-\frac{(c-ic \tan(fx+e))^3}{3} - 2c\sqrt{c-ic \tan(fx+e)} + 2c^{\frac{3}{2}}\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)}\sqrt{2}}{2\sqrt{c}}\right) \right)}{f} + \frac{a^3(3iA+B) \left(\frac{2(c-ic \tan(fx+e))^{11/2}}{11} + \frac{(-5iBc+cA)(c-ic \tan(fx+e))^9}{9} + \frac{(-4(-iBc+cA)c+4iBc^2)(c-ic \tan(fx+e))^7}{7} + \frac{4(-iBc+cA)c^2}{4} \right)}{f c^3}$

```
input int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, metho
d=_RETURNVERBOSE)
```

```
output 2*I/f*a^3/c^3*(1/11*I*B*(c-I*c*tan(f*x+e))^(11/2)+1/9*(-5*I*B*c+c*A)*(c-I*
c*tan(f*x+e))^(9/2)+1/7*(-4*(-I*B*c+c*A)*c+4*I*B*c^2)*(c-I*c*tan(f*x+e))^(
7/2)+4/5*(-I*B*c+c*A)*c^2*(c-I*c*tan(f*x+e))^(5/2))
```

3.757.5 Fracas [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 173, normalized size of antiderivative = 1.20

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx =$$

$$\frac{32\sqrt{2}(693(-iA - B)a^3c^2e^{(6ifx+6ie)} + 99(-11iA - B)a^3c^2e^{(4ifx+4ie)} + 44(-11iA - B)a^3c^2e^{(2ifx+2ie)} + 4(-11iA - B)a^3c^2e^{(ifx+ie)} + 4(-11iA - B)a^3c^2e^{(ifx+ie)})}{3465(fe^{(10ifx+10ie)} + 5fe^{(8ifx+8ie)} + 10fe^{(6ifx+6ie)} + 10fe^{(4ifx+4ie)} + 5fe^{(2ifx+2ie)})}$$

```
input integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x
, algorithm="fracas")
```

3.757. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx$

output $-32/3465\sqrt{2}*(693*(-I*A - B)*a^3*c^2*e^{(6*I*f*x + 6*I*e)} + 99*(-11*I*A - B)*a^3*c^2*e^{(4*I*f*x + 4*I*e)} + 44*(-11*I*A - B)*a^3*c^2*e^{(2*I*f*x + 2*I*e)} + 8*(-11*I*A - B)*a^3*c^2)*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)}/(f*e^{(10*I*f*x + 10*I*e)} + 5*f*e^{(8*I*f*x + 8*I*e)} + 10*f*e^{(6*I*f*x + 6*I*e)} + 10*f*e^{(4*I*f*x + 4*I*e)} + 5*f*e^{(2*I*f*x + 2*I*e)} + f)$

3.757.6 Sympy [F]

$$\begin{aligned} & \int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx = \\ & -ia^3 \left(\int iAc^2 \sqrt{-ic \tan(e + fx) + c} dx \right. \\ & + \int \left(-Ac^2 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx) \right) dx \\ & + \int \left(-2Ac^2 \sqrt{-ic \tan(e + fx) + c} \tan^3(e + fx) \right) dx \\ & + \int \left(-Ac^2 \sqrt{-ic \tan(e + fx) + c} \tan^5(e + fx) \right) dx \\ & + \int \left(-Bc^2 \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx) \right) dx \\ & + \int \left(-2Bc^2 \sqrt{-ic \tan(e + fx) + c} \tan^4(e + fx) \right) dx \\ & + \int \left(-Bc^2 \sqrt{-ic \tan(e + fx) + c} \tan^6(e + fx) \right) dx \\ & + \int 2iAc^2 \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx) dx \\ & + \int iAc^2 \sqrt{-ic \tan(e + fx) + c} \tan^4(e + fx) dx \\ & + \int iBc^2 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx) dx \\ & + \int 2iBc^2 \sqrt{-ic \tan(e + fx) + c} \tan^3(e + fx) dx \\ & \left. + \int iBc^2 \sqrt{-ic \tan(e + fx) + c} \tan^5(e + fx) dx \right) \end{aligned}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(5/2),x)`

```
output -I*a**3*(Integral(I*A*c**2*sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-A*c
**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(-2*A*c**2*sqrt
(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x) + Integral(-A*c**2*sqrt(-I*c*t
an(e + f*x) + c)*tan(e + f*x)**5, x) + Integral(-B*c**2*sqrt(-I*c*tan(e +
f*x) + c)*tan(e + f*x)**2, x) + Integral(-2*B*c**2*sqrt(-I*c*tan(e + f*x)
+ c)*tan(e + f*x)**4, x) + Integral(-B*c**2*sqrt(-I*c*tan(e + f*x) + c)*ta
n(e + f*x)**6, x) + Integral(2*I*A*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e
+ f*x)**2, x) + Integral(I*A*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)
**4, x) + Integral(I*B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) +
Integral(2*I*B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x) + Int
egral(I*B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**5, x))
```

3.757.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.72

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx = \frac{2i \left(315i (-ictan(fx + e) + c)^{\frac{11}{2}} Ba^3 + 385 (-ictan(fx + e) + c)^{\frac{9}{2}} (A - 5iB)a^3c - 1980 (-ictan(fx + e) + c)^{\frac{7}{2}} (A - 2iB)a^3c^2 + 2772 (-ictan(fx + e) + c)^{\frac{5}{2}} (A - iB)a^3c^3 \right)}{3465 c^3 f}$$

```
input integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x
, algorithm="maxima")
```

```
output 2/3465*I*(315*I*(-I*c*tan(f*x + e) + c)^(11/2)*B*a^3 + 385*(-I*c*tan(f*x +
e) + c)^(9/2)*(A - 5*I*B)*a^3*c - 1980*(-I*c*tan(f*x + e) + c)^(7/2)*(A -
2*I*B)*a^3*c^2 + 2772*(-I*c*tan(f*x + e) + c)^(5/2)*(A - I*B)*a^3*c^3)/(c
^3*f)
```

3.757.8 Giac [F]

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx = \int (B \tan(fx + e) + A)(ia \tan(fx + e) + a)^3 (-ictan(fx + e) + c)^{\frac{5}{2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^3*(-I*c*tan(f*x + e)
+ c)^(5/2), x)`

3.757.9 Mupad [B] (verification not implemented)

Time = 12.11 (sec) , antiderivative size = 349, normalized size of antiderivative = 2.42

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2} dx =$$

$$-\frac{\left(\frac{a^3 c^2 (A-B 1i) 32i}{7f} + \frac{a^3 c^2 (A-B 3i) 32i}{7f}\right) \sqrt{c + \frac{c(e^{e 2i+fx 2i} 1i-i) 1i}{e^{e 2i+fx 2i+1}}}}{(e^{e 2i+fx 2i} + 1)^3}$$

$$-\frac{\left(\frac{a^3 c^2 (A-B 1i) 32i}{11f} - \frac{a^3 c^2 (A+B 1i) 32i}{11f}\right) \sqrt{c + \frac{c(e^{e 2i+fx 2i} 1i-i) 1i}{e^{e 2i+fx 2i+1}}}}{(e^{e 2i+fx 2i} + 1)^5}$$

$$+\frac{\left(\frac{128 B a^3 c^2}{9f} + \frac{a^3 c^2 (A-B 1i) 32i}{9f}\right) \sqrt{c + \frac{c(e^{e 2i+fx 2i} 1i-i) 1i}{e^{e 2i+fx 2i+1}}}}{(e^{e 2i+fx 2i} + 1)^4}$$

$$+\frac{a^3 c^2 (A - B 1i) \sqrt{c + \frac{c(e^{e 2i+fx 2i} 1i-i) 1i}{e^{e 2i+fx 2i+1}}}}{5f (e^{e 2i+fx 2i} + 1)^2} 32i$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i)
^(5/2),x)`

output `((a^3*c^2*(A - B*1i)*32i)/(9*f) + (128*B*a^3*c^2)/(9*f))*(c + (c*(exp(e*2
i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(1/2))/(exp(e*2i + f*x*
2i) + 1)^4 - (((a^3*c^2*(A - B*1i)*32i)/(11*f) - (a^3*c^2*(A + B*1i)*32i)/
(11*f))*(c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))
^(1/2))/(exp(e*2i + f*x*2i) + 1)^5 - (((a^3*c^2*(A - B*1i)*32i)/(7*f) + (a
^3*c^2*(A - B*3i)*32i)/(7*f))*(c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(ex
p(e*2i + f*x*2i) + 1))^(1/2))/(exp(e*2i + f*x*2i) + 1)^3 + (a^3*c^2*(A - B
1i)(c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(1
/2)*32i)/(5*f*(exp(e*2i + f*x*2i) + 1)^2)`

3.758 $\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))(c-ictan(e+fx))^{3/2} dx$

3.758.1 Optimal result	6899
3.758.2 Mathematica [A] (verified)	6900
3.758.3 Rubi [A] (verified)	6900
3.758.4 Maple [A] (verified)	6902
3.758.5 Fricas [A] (verification not implemented)	6902
3.758.6 Sympy [F]	6903
3.758.7 Maxima [A] (verification not implemented)	6904
3.758.8 Giac [F]	6904
3.758.9 Mupad [B] (verification not implemented)	6905

3.758.1 Optimal result

Integrand size = 43, antiderivative size = 144

$$\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))(c - ictan(e + fx))^{3/2} dx = \frac{8a^3(iA + B)(c - ictan(e + fx))^{3/2}}{3f} - \frac{8a^3(iA + 2B)(c - ictan(e + fx))^{5/2}}{5cf} + \frac{2a^3(iA + 5B)(c - ictan(e + fx))^{7/2}}{7c^2f} - \frac{2a^3B(c - ictan(e + fx))^{9/2}}{9c^3f}$$

```
output 8/3*a^3*(I*A+B)*(c-I*c*tan(f*x+e))^(3/2)/f-8/5*a^3*(I*A+2*B)*(c-I*c*tan(f*x+e))^(5/2)/c/f+2/7*a^3*(I*A+5*B)*(c-I*c*tan(f*x+e))^(7/2)/c^2/f-2/9*a^3*B*(c-I*c*tan(f*x+e))^(9/2)/c^3/f
```

3.758.2 Mathematica [A] (verified)

Time = 4.88 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.89

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ict \tan(e + fx))^{3/2} dx = \frac{a^3 c^2 \sec^5(e + fx) (99(3iA + B) \cos(e + fx) + (129iA + 113B) \cos(3(e + fx)) - 2(81A - 62iB) \cos(5(e + fx))) \sin(e + fx) (\cos(2(e + fx)) - i \sin(2(e + fx)))}{315 f \sqrt{c - ict \tan(e + fx)}}$$

input `Integrate[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(a^3*c^2*Sec[e + f*x]^5*(99*((3*I)*A + B)*Cos[e + f*x] + ((129*I)*A + 113*B)*Cos[3*(e + f*x)] - 2*(81*A - (62*I)*B + (81*A - (97*I)*B)*Cos[2*(e + f*x)])*Sin[e + f*x]*(Cos[2*(e + f*x)] - I*Sin[2*(e + f*x)])/(315*f*Sqrt[c - I*c*Tan[e + f*x]])`

3.758.3 Rubi [A] (verified)Time = 0.39 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.91, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))^3 (c - ict \tan(e + fx))^{3/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))^3 (c - ict \tan(e + fx))^{3/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int a^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) \sqrt{c - ict \tan(e + fx)} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{a^3 c \int (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx)) \sqrt{c - ict \tan(e + fx)} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{86} \end{aligned}$$

3.758. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ict \tan(e + fx))^{3/2} dx$

$$\frac{a^3 c \int \left(\frac{iB(c-ictan(e+fx))^{7/2}}{c^3} + \frac{(A-5iB)(c-ictan(e+fx))^{5/2}}{c^2} - \frac{4(A-2iB)(c-ictan(e+fx))^{3/2}}{c} + 4(A-iB)\sqrt{c-ictan(e+fx)} \right)}{f}$$

↓ 2009

$$\frac{a^3 c \left(\frac{2(5B+iA)(c-ictan(e+fx))^{7/2}}{7c^3} - \frac{8(2B+iA)(c-ictan(e+fx))^{5/2}}{5c^2} + \frac{8(B+iA)(c-ictan(e+fx))^{3/2}}{3c} - \frac{2B(c-ictan(e+fx))^{9/2}}{9c^4} \right)}{f}$$

input `Int[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2), x]`

output `(a^3*c*((8*(I*A + B)*(c - I*c*Tan[e + f*x])^(3/2))/(3*c) - (8*(I*A + 2*B)*(c - I*c*Tan[e + f*x])^(5/2))/(5*c^2) + (2*(I*A + 5*B)*(c - I*c*Tan[e + f*x])^(7/2))/(7*c^3) - (2*B*(c - I*c*Tan[e + f*x])^(9/2))/(9*c^4))/f`

3.758.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.758.4 Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.84

method	result
derivativedivides	$\frac{2ia^3 \left(\frac{iB(c-ic \tan(fx+e))^{\frac{9}{2}}}{9} + \frac{(-5iBc+cA)(c-ic \tan(fx+e))^{\frac{7}{2}}}{7} + \frac{(-4(-iBc+cA)c+4iBc^2)(c-ic \tan(fx+e))^{\frac{5}{2}}}{5} + \frac{4(-iBc+cA)c^2}{4} \right)}{fc^3}$
default	$\frac{2ia^3 \left(\frac{iB(c-ic \tan(fx+e))^{\frac{9}{2}}}{9} + \frac{(-5iBc+cA)(c-ic \tan(fx+e))^{\frac{7}{2}}}{7} + \frac{(-4(-iBc+cA)c+4iBc^2)(c-ic \tan(fx+e))^{\frac{5}{2}}}{5} + \frac{4(-iBc+cA)c^2}{4} \right)}{fc^3}$
parts	$\frac{2ia^3 Ac \left(-\sqrt{c-ic \tan(fx+e)} + \sqrt{c} \sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}} \right) \right)}{f} + \frac{a^3(3iA+B) \left(\frac{2(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} + 2c\sqrt{c-ic \tan(fx+e)} \right)}{f}$

```
input int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x, metho
d=_RETURNVERBOSE)
```

```
output 2*I/f*a^3/c^3*(1/9*I*B*(c-I*c*tan(f*x+e))^(9/2)+1/7*(-5*I*B*c+c*A)*(c-I*c*
tan(f*x+e))^(7/2)+1/5*(-4*(-I*B*c+c*A)*c+4*I*B*c^2)*(c-I*c*tan(f*x+e))^(5/
2)+4/3*(-I*B*c+c*A)*c^2*(c-I*c*tan(f*x+e))^(3/2))
```

3.758.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.06

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx =$$

$$\frac{16\sqrt{2}(105(-iA - B)a^3ce^{(6i fx+6ie)} + 63(-3iA - B)a^3ce^{(4i fx+4ie)} + 36(-3iA - B)a^3ce^{(2i fx+2ie)} + 8(-3iA - B)a^3ce^{(0i fx+0ie)})}{315(fe^{(8i fx+8ie)} + 4fe^{(6i fx+6ie)} + 6fe^{(4i fx+4ie)} + 4fe^{(2i fx+2ie)} + f)}$$

```
input integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x
, algorithm="fricas")
```

3.758. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx$

output $-16/315\sqrt{2}*(105*(-I*A - B)*a^3*c*e^{(6*I*f*x + 6*I*e)} + 63*(-3*I*A - B)*a^3*c*e^{(4*I*f*x + 4*I*e)} + 36*(-3*I*A - B)*a^3*c*e^{(2*I*f*x + 2*I*e)} + 8*(-3*I*A - B)*a^3*c)*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)})/(f*e^{(8*I*f*x + 8*I*e)} + 4*f*e^{(6*I*f*x + 6*I*e)} + 6*f*e^{(4*I*f*x + 4*I*e)} + 4*f*e^{(2*I*f*x + 2*I*e)} + f)$

3.758.6 Sympy [F]

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx =$$

$$-ia^3 \left(\int iAc \sqrt{-ictan(e + fx) + c} dx \right.$$

$$+ \int \left(-2Ac \sqrt{-ictan(e + fx) + c} \tan(e + fx) \right) dx$$

$$+ \int \left(-2Ac \sqrt{-ictan(e + fx) + c} \tan^3(e + fx) \right) dx$$

$$+ \int \left(-2Bc \sqrt{-ictan(e + fx) + c} \tan^2(e + fx) \right) dx$$

$$+ \int \left(-2Bc \sqrt{-ictan(e + fx) + c} \tan^4(e + fx) \right) dx$$

$$+ \int \left(-iAc \sqrt{-ictan(e + fx) + c} \tan^4(e + fx) \right) dx$$

$$+ \int iBc \sqrt{-ictan(e + fx) + c} \tan(e + fx) dx$$

$$+ \left. \int \left(-iBc \sqrt{-ictan(e + fx) + c} \tan^5(e + fx) \right) dx \right)$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(3/2),x)`

output `-I*a**3*(Integral(I*A*c*sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-2*A*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(-2*A*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x) + Integral(-2*B*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x) + Integral(-2*B*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4, x) + Integral(-I*A*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4, x) + Integral(I*B*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(-I*B*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**5, x))`

3.758.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.72

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx = \frac{2i \left(35i (-ictan(fx + e) + c)^{\frac{9}{2}} Ba^3 + 45 (-ictan(fx + e) + c)^{\frac{7}{2}} (A - 5iB) a^3 c - 252 (-ictan(fx + e) + c)^{\frac{5}{2}} (A - 2iB) a^3 c^2 + 420 (-ictan(fx + e) + c)^{\frac{3}{2}} (A - iB) a^3 c^3 \right)}{315 c^3 f}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `2/315*I*(35*I*(-I*c*tan(f*x + e) + c)^(9/2)*B*a^3 + 45*(-I*c*tan(f*x + e) + c)^(7/2)*(A - 5*I*B)*a^3*c - 252*(-I*c*tan(f*x + e) + c)^(5/2)*(A - 2*I*B)*a^3*c^2 + 420*(-I*c*tan(f*x + e) + c)^(3/2)*(A - I*B)*a^3*c^3)/(c^3*f)`

3.758.8 Giac [F]

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx = \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^3 (-ictan(fx + e) + c)^{\frac{3}{2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^3*(-I*c*tan(f*x + e) + c)^(3/2), x)`

3.758.9 Mupad [B] (verification not implemented)

Time = 11.70 (sec) , antiderivative size = 335, normalized size of antiderivative = 2.33

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx =$$

$$-\frac{\left(\frac{a^3 c(A-B)16i}{5f} + \frac{a^3 c(A-B)3i16i}{5f}\right) \sqrt{c + \frac{c(e^{e^{2i+fx}2i}1i-i)}{e^{e^{2i+fx}2i+1}}}}{(e^{e^{2i+fx}2i} + 1)^2}$$

$$-\frac{\left(\frac{a^3 c(A-B)16i}{9f} - \frac{a^3 c(A+B)16i}{9f}\right) \sqrt{c + \frac{c(e^{e^{2i+fx}2i}1i-i)}{e^{e^{2i+fx}2i+1}}}}{(e^{e^{2i+fx}2i} + 1)^4}$$

$$+\frac{\left(\frac{64Ba^3c}{7f} + \frac{a^3c(A-B)16i}{7f}\right) \sqrt{c + \frac{c(e^{e^{2i+fx}2i}1i-i)}{e^{e^{2i+fx}2i+1}}}}{(e^{e^{2i+fx}2i} + 1)^3}$$

$$+\frac{a^3c(A-B)16i \sqrt{c + \frac{c(e^{e^{2i+fx}2i}1i-i)}{e^{e^{2i+fx}2i+1}}}}{3f(e^{e^{2i+fx}2i} + 1)}$$

```
input int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i)
^(3/2),x)
```

```
output (((a^3*c*(A - B*1i)*16i)/(7*f) + (64*B*a^3*c)/(7*f))*(c + (c*(exp(e*2i + f
*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(1/2))/(exp(e*2i + f*x*2i) +
1)^3 - (((a^3*c*(A - B*1i)*16i)/(9*f) - (a^3*c*(A + B*1i)*16i)/(9*f))*(c
+ (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(1/2))/(ex
p(e*2i + f*x*2i) + 1)^4 - (((a^3*c*(A - B*1i)*16i)/(5*f) + (a^3*c*(A - B*3
i)*16i)/(5*f))*(c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i
) + 1))^(1/2))/(exp(e*2i + f*x*2i) + 1)^2 + (a^3*c*(A - B*1i)*(c + (c*(exp
(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(1/2)*16i)/(3*f*(ex
p(e*2i + f*x*2i) + 1))
```


3.759 $\int (a+ia \tan(e+fx))^3(A+B \tan(e+fx))\sqrt{c-ictan(e+fx)}$

3.759.1 Optimal result	6906
3.759.2 Mathematica [A] (verified)	6906
3.759.3 Rubi [A] (verified)	6907
3.759.4 Maple [A] (verified)	6909
3.759.5 Fricas [A] (verification not implemented)	6909
3.759.6 Sympy [F]	6910
3.759.7 Maxima [A] (verification not implemented)	6911
3.759.8 Giac [F]	6911
3.759.9 Mupad [B] (verification not implemented)	6912

3.759.1 Optimal result

Integrand size = 43, antiderivative size = 142

$$\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)} dx$$

$$= \frac{8a^3(iA + B)\sqrt{c - ictan(e + fx)}}{f} - \frac{8a^3(iA + 2B)(c - ictan(e + fx))^{3/2}}{3cf}$$

$$+ \frac{2a^3(iA + 5B)(c - ictan(e + fx))^{5/2}}{5c^2f} - \frac{2a^3B(c - ictan(e + fx))^{7/2}}{7c^3f}$$

```
output 8*a^3*(I*A+B)*(c-I*c*tan(f*x+e))^(1/2)/f-8/3*a^3*(I*A+2*B)*(c-I*c*tan(f*x+
e))^(3/2)/c/f+2/5*a^3*(I*A+5*B)*(c-I*c*tan(f*x+e))^(5/2)/c^2/f-2/7*a^3*B*(
c-I*c*tan(f*x+e))^(7/2)/c^3/f
```

3.759.2 Mathematica [A] (verified)

Time = 3.96 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.65

$$\int (a + ia \tan(e + fx))^3(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)} dx =$$

$$\frac{2a^3c(i + \tan(e + fx))(-301A + 230iB + (-98iA - 115B) \tan(e + fx) + 3(7A - 20iB) \tan^2(e + fx))}{105f\sqrt{c - ictan(e + fx)}}$$

```
input Integrate[(a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e
+ f*x]],x]
```

output $(-2*a^3*c*(I + \text{Tan}[e + f*x])*(-301*A + (230*I)*B + ((-98*I)*A - 115*B)*\text{Tan}[e + f*x] + 3*(7*A - (20*I)*B)*\text{Tan}[e + f*x]^2 + 15*B*\text{Tan}[e + f*x]^3))/(105*f*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])$

3.759.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.91, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))^3 \sqrt{c - ictan(e + fx)} (A + B \tan(e + fx)) dx$$

↓ 3042

$$\int (a + ia \tan(e + fx))^3 \sqrt{c - ictan(e + fx)} (A + B \tan(e + fx)) dx$$

↓ 4071

$$\frac{ac \int \frac{a^2(i \tan(e+fx)+1)^2(A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} d \tan(e + fx)}{f}$$

↓ 27

$$\frac{a^3c \int \frac{(i \tan(e+fx)+1)^2(A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} d \tan(e + fx)}{f}$$

↓ 86

$$\frac{a^3c \int \left(\frac{iB(c-ictan(e+fx))^{5/2}}{c^3} + \frac{(A-5iB)(c-ictan(e+fx))^{3/2}}{c^2} - \frac{4(A-2iB)\sqrt{c-ictan(e+fx)}}{c} + \frac{4(A-iB)}{\sqrt{c-ictan(e+fx)}} \right) d \tan(e + fx)}{f}$$

↓ 2009

$$\frac{a^3c \left(\frac{2(5B+iA)(c-ictan(e+fx))^{5/2}}{5c^3} - \frac{8(2B+iA)(c-ictan(e+fx))^{3/2}}{3c^2} + \frac{8(B+iA)\sqrt{c-ictan(e+fx)}}{c} - \frac{2B(c-ictan(e+fx))^{7/2}}{7c^4} \right)}{f}$$

input $\text{Int}[(a + I*a*\text{Tan}[e + f*x])^3*(A + B*\text{Tan}[e + f*x])*Sqrt[c - I*c*\text{Tan}[e + f*x]],x]$

3.759. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx$

```
output (a^3*c*((8*(I*A + B)*Sqrt[c - I*c*Tan[e + f*x]])/c - (8*(I*A + 2*B)*(c - I
*c*Tan[e + f*x])^(3/2))/(3*c^2) + (2*(I*A + 5*B)*(c - I*c*Tan[e + f*x])^(5
/2))/(5*c^3) - (2*B*(c - I*c*Tan[e + f*x])^(7/2))/(7*c^4))/f
```

3.759.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.759.4 Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.85

method	result
derivativedivides	$2ia^3 \frac{\left(\frac{iB(c-ic \tan(fx+e))^{\frac{7}{2}}}{7} + \frac{(-5iBc+cA)(c-ic \tan(fx+e))^{\frac{5}{2}}}{5} + \frac{(-4(-iBc+cA)c+4iBc^2)(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} + 4\sqrt{c-ic \tan(fx+e)} \right)}{fc^3}$
default	$2ia^3 \left(\frac{iB(c-ic \tan(fx+e))^{\frac{7}{2}}}{7} + \frac{(-5iBc+cA)(c-ic \tan(fx+e))^{\frac{5}{2}}}{5} + \frac{(-4(-iBc+cA)c+4iBc^2)(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} + 4\sqrt{c-ic \tan(fx+e)} \right) \frac{1}{fc^3}$
parts	$\frac{6ia^3(-iB+A) \left(\frac{(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} - \frac{c^{\frac{3}{2}}\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)}\sqrt{2}}{2\sqrt{c}}\right)}{2} \right)}{fc} - \frac{2a^3(iA+3B) \left(-\frac{(c-ic \tan(fx+e))^{\frac{5}{2}}}{5} + \dots \right)}{fc}$

input `int((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output `2*I/f*a^3/c^3*(1/7*I*B*(c-I*c*tan(f*x+e))^(7/2)+1/5*(-5*I*B*c+c*A)*(c-I*c*tan(f*x+e))^(5/2)+1/3*(-4*(-I*B*c+c*A)*c+4*I*B*c^2)*(c-I*c*tan(f*x+e))^(3/2)+4*(c-I*c*tan(f*x+e))^(1/2)*(-I*B*c+c*A)*c^2)`

3.759.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.96

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx = \frac{8\sqrt{2}(105(-iA - B)a^3e^{(6ifx+6ie)} + 35(-7iA - 5B)a^3e^{(4ifx+4ie)} + 28(-7iA - 5B)a^3e^{(2ifx+2ie)} + 8(-7iA - 5B)a^3) \sqrt{c/(e^{(2ifx+2ie)} + 1)} + 3fe^{(6ifx+6ie)} + 3fe^{(4ifx+4ie)} + 3fe^{(2ifx+2ie)} + f}{105(fe^{(6ifx+6ie)} + 3fe^{(4ifx+4ie)} + 3fe^{(2ifx+2ie)} + f)}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e)),x,algorithm="fricas")`

output `-8/105*sqrt(2)*(105*(-I*A - B)*a^3*e^(6*I*f*x + 6*I*e) + 35*(-7*I*A - 5*B)*a^3*e^(4*I*f*x + 4*I*e) + 28*(-7*I*A - 5*B)*a^3*e^(2*I*f*x + 2*I*e) + 8*(-7*I*A - 5*B)*a^3)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)`

3.759. $\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx$

3.759.6 Sympy [F]

$$\begin{aligned}
& \int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx \\
&= -ia^3 \left(\int iA \sqrt{-ic \tan(e + fx) + c} dx \right. \\
&\quad + \int \left(-3A \sqrt{-ic \tan(e + fx) + c} \tan(e + fx) \right) dx \\
&\quad + \int A \sqrt{-ic \tan(e + fx) + c} \tan^3(e + fx) dx \\
&\quad + \int \left(-3B \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx) \right) dx \\
&\quad + \int B \sqrt{-ic \tan(e + fx) + c} \tan^4(e + fx) dx \\
&\quad + \int \left(-3iA \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx) \right) dx \\
&\quad + \int iB \sqrt{-ic \tan(e + fx) + c} \tan(e + fx) dx \\
&\quad \left. + \int \left(-3iB \sqrt{-ic \tan(e + fx) + c} \tan^3(e + fx) \right) dx \right)
\end{aligned}$$

input `integrate((c-I*c*tan(f*x+e))**(1/2)*(a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e)),x)`

output `-I*a**3*(Integral(I*A*sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-3*A*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(A*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x) + Integral(-3*B*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x) + Integral(B*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4, x) + Integral(-3*I*A*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2, x) + Integral(I*B*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x), x) + Integral(-3*I*B*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3, x))`

3.759.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.73

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx$$

$$= \frac{2i \left(15i (-ic \tan(fx + e) + c)^{\frac{7}{2}} B a^3 + 21 (-ic \tan(fx + e) + c)^{\frac{5}{2}} (A - 5i B) a^3 c - 140 (-ic \tan(fx + e) + c)^{\frac{3}{2}} (A - 2i B) a^3 c^2 + 420 \sqrt{-ic \tan(fx + e) + c} (A - i B) a^3 c^3 \right)}{105 c^3 f}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e)),x
, algorithm="maxima")`

output `2/105*I*(15*I*(-I*c*tan(f*x + e) + c)^(7/2)*B*a^3 + 21*(-I*c*tan(f*x + e)
+ c)^(5/2)*(A - 5*I*B)*a^3*c - 140*(-I*c*tan(f*x + e) + c)^(3/2)*(A - 2*I*
B)*a^3*c^2 + 420*sqrt(-I*c*tan(f*x + e) + c)*(A - I*B)*a^3*c^3)/(c^3*f)`

3.759.8 Giac [F]

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx$$

$$= \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^3 \sqrt{-ic \tan(fx + e) + c} dx$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e)),x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^3*sqrt(-I*c*tan(f*x
+ e) + c), x)`

3.759.9 Mupad [B] (verification not implemented)

Time = 12.55 (sec) , antiderivative size = 313, normalized size of antiderivative = 2.20

$$\int (a + ia \tan(e + fx))^3 (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx$$

$$= - \frac{\sqrt{c + \frac{c(e^{e^{2i} + f x^{2i}} - 1i - i)}{e^{e^{2i} + f x^{2i}} + 1}} \frac{1i}{e^{e^{2i} + f x^{2i}} + 1} \left(\frac{a^3 (A - B 1i) 8i}{3f} + \frac{a^3 (A - B 3i) 8i}{3f} \right)}{e^{e^{2i} + f x^{2i}} + 1}$$

$$- \frac{\sqrt{c + \frac{c(e^{e^{2i} + f x^{2i}} - 1i - i)}{e^{e^{2i} + f x^{2i}} + 1}} \frac{1i}{e^{e^{2i} + f x^{2i}} + 1} \left(\frac{a^3 (A - B 1i) 8i}{7f} - \frac{a^3 (A + B 1i) 8i}{7f} \right)}{(e^{e^{2i} + f x^{2i}} + 1)^3}$$

$$+ \frac{\sqrt{c + \frac{c(e^{e^{2i} + f x^{2i}} - 1i - i)}{e^{e^{2i} + f x^{2i}} + 1}} \frac{1i}{e^{e^{2i} + f x^{2i}} + 1} \left(\frac{32 B a^3}{5f} + \frac{a^3 (A - B 1i) 8i}{5f} \right)}{(e^{e^{2i} + f x^{2i}} + 1)^2} + \frac{a^3 (A - B 1i) \sqrt{c + \frac{c(e^{e^{2i} + f x^{2i}} - 1i - i)}{e^{e^{2i} + f x^{2i}} + 1}} \frac{1i}{e^{e^{2i} + f x^{2i}} + 1} 8i}{f}$$

```
input int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i)
^(1/2),x)
```

```
output ((c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(1/2)*
((a^3*(A - B*1i)*8i)/(5*f) + (32*B*a^3)/(5*f)))/(exp(e*2i + f*x*2i) + 1)^2
- ((c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(1/2)*
((a^3*(A - B*1i)*8i)/(7*f) - (a^3*(A + B*1i)*8i)/(7*f)))/(exp(e*2i + f*
x*2i) + 1)^3 - ((c + (c*(exp(e*2i + f*x*2i)*1i - 1i)*1i)/(exp(e*2i + f*x*2
i) + 1))^(1/2)*((a^3*(A - B*1i)*8i)/(3*f) + (a^3*(A - B*3i)*8i)/(3*f)))/(e
xp(e*2i + f*x*2i) + 1) + (a^3*(A - B*1i)*(c + (c*(exp(e*2i + f*x*2i)*1i -
1i)*1i)/(exp(e*2i + f*x*2i) + 1))^(1/2)*8i)/f
```

3.760
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$$

3.760.1 Optimal result 6913
 3.760.2 Mathematica [A] (verified) 6913
 3.760.3 Rubi [A] (verified) 6914
 3.760.4 Maple [A] (verified) 6916
 3.760.5 Fricas [A] (verification not implemented) 6916
 3.760.6 Sympy [F] 6917
 3.760.7 Maxima [A] (verification not implemented) 6917
 3.760.8 Giac [F] 6918
 3.760.9 Mupad [B] (verification not implemented) 6918

3.760.1 Optimal result

Integrand size = 43, antiderivative size = 140

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx$$

$$= -\frac{8a^3(iA + B)}{f\sqrt{c - ic \tan(e + fx)}} - \frac{8a^3(iA + 2B)\sqrt{c - ic \tan(e + fx)}}{cf}$$

$$+ \frac{2a^3(iA + 5B)(c - ic \tan(e + fx))^{3/2}}{3c^2f} - \frac{2a^3B(c - ic \tan(e + fx))^{5/2}}{5c^3f}$$

output `-8*a^3*(I*A+B)/f/(c-I*c*tan(f*x+e))^(1/2)-8*a^3*(I*A+2*B)*(c-I*c*tan(f*x+e))^(1/2)/c/f+2/3*a^3*(I*A+5*B)*(c-I*c*tan(f*x+e))^(3/2)/c^2/f-2/5*a^3*B*(c-I*c*tan(f*x+e))^(5/2)/c^3/f`

3.760.2 Mathematica [A] (verified)

Time = 5.19 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.59

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx =$$

$$-\frac{2ia^3(115A - 158iB + (-50iA - 79B) \tan(e + fx) + (5A - 16iB) \tan^2(e + fx) + 3B \tan^3(e + fx))}{15f\sqrt{c - ic \tan(e + fx)}}$$

input `Integrate[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/Sqrt[c - I*c*Tan[e + f*x]],x]`

output `(((-2*I)/15)*a^3*(115*A - (158*I)*B + ((-50*I)*A - 79*B)*Tan[e + f*x] + (5*A - (16*I)*B)*Tan[e + f*x]^2 + 3*B*Tan[e + f*x]^3))/(f*Sqrt[c - I*c*Tan[e + f*x]])`

3.760.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.91, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{a^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^3 c \int \frac{(i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{86} \\
 & \frac{a^3 c \int \left(\frac{iB(c - ic \tan(e + fx))^{3/2}}{c^3} + \frac{(A - 5iB)\sqrt{c - ic \tan(e + fx)}}{c^2} - \frac{4(A - 2iB)}{c\sqrt{c - ic \tan(e + fx)}} + \frac{4(A - iB)}{(c - ic \tan(e + fx))^{3/2}} \right) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^3 c \left(\frac{2(5B + iA)(c - ic \tan(e + fx))^{3/2}}{3c^3} - \frac{8(2B + iA)\sqrt{c - ic \tan(e + fx)}}{c^2} - \frac{8(B + iA)}{c\sqrt{c - ic \tan(e + fx)}} - \frac{2B(c - ic \tan(e + fx))^{5/2}}{5c^4} \right)}{f}
 \end{aligned}$$

3.760. $\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx$

input `Int[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/Sqrt[c - I*c*Tan[e + f*x]],x]`

output `(a^3*c*((-8*(I*A + B))/(c*Sqrt[c - I*c*Tan[e + f*x]]) - (8*(I*A + 2*B)*Sqrt[c - I*c*Tan[e + f*x]])/c^2 + (2*(I*A + 5*B)*(c - I*c*Tan[e + f*x])^(3/2))/(3*c^3) - (2*B*(c - I*c*Tan[e + f*x])^(5/2))/(5*c^4))/f`

3.760.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.760.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.96

method	result
derivativedivides	$\frac{2ia^3 \left(\frac{iB(c-ic \tan(fx+e))^{5/2}}{5} - \frac{5iBc(c-ic \tan(fx+e))^{3/2}}{3} + \frac{Ac(c-ic \tan(fx+e))^{3/2}}{3} + 8i\sqrt{c-ic \tan(fx+e)} Bc^2 - 4\sqrt{c-ic \tan(fx+e)} \right)}{fc^3}$
default	$\frac{2ia^3 \left(\frac{iB(c-ic \tan(fx+e))^{5/2}}{5} - \frac{5iBc(c-ic \tan(fx+e))^{3/2}}{3} + \frac{Ac(c-ic \tan(fx+e))^{3/2}}{3} + 8i\sqrt{c-ic \tan(fx+e)} Bc^2 - 4\sqrt{c-ic \tan(fx+e)} \right)}{fc^3}$
parts	$2ia^3 Ac \left(-\frac{1}{2c\sqrt{c-ic \tan(fx+e)}} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{4c^{3/2}} \right) + a^3(3iA+B) \left(-\frac{1}{\sqrt{c-ic \tan(fx+e)}} - \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{f} \right)$

input `int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output `2*I/f*a^3/c^3*(1/5*I*B*(c-I*c*tan(f*x+e))^(5/2)-5/3*I*B*c*(c-I*c*tan(f*x+e))^(3/2)+1/3*A*c*(c-I*c*tan(f*x+e))^(3/2)+8*I*(c-I*c*tan(f*x+e))^(1/2)*B*c^2-4*(c-I*c*tan(f*x+e))^(1/2)*A*c^2-4*c^3*(A-I*B)/(c-I*c*tan(f*x+e))^(1/2))`

3.760.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.91

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx = \frac{4\sqrt{2}(15(iA + B)a^3 e^{(6i fx + 6i e)} + 15(5iA + 7B)a^3 e^{(4i fx + 4i e)} + 20(5iA + 7B)a^3 e^{(2i fx + 2i e)} + 8(5iA + 7B)a^3) \sqrt{c/(e^{(2i fx + 2i e)} + 1)} + 15(cfe^{(4i fx + 4i e)} + 2cfe^{(2i fx + 2i e)} + cf)}{15(cfe^{(4i fx + 4i e)} + 2cfe^{(2i fx + 2i e)} + cf)}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x,algorithm="fracas")`

output `-4/15*sqrt(2)*(15*(I*A + B)*a^3*e^(6*I*f*x + 6*I*e) + 15*(5*I*A + 7*B)*a^3*e^(4*I*f*x + 4*I*e) + 20*(5*I*A + 7*B)*a^3*e^(2*I*f*x + 2*I*e) + 8*(5*I*A + 7*B)*a^3)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c*f*e^(4*I*f*x + 4*I*e) + 2*c*f*e^(2*I*f*x + 2*I*e) + c*f)`

3.760. $\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx$

3.760.6 Sympy [F]

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx \\ &= -ia^3 \left(\int \frac{iA}{\sqrt{-ictan(e + fx) + c}} dx + \int \left(-\frac{3A \tan(e + fx)}{\sqrt{-ictan(e + fx) + c}} \right) dx \right. \\ & \quad + \int \frac{A \tan^3(e + fx)}{\sqrt{-ictan(e + fx) + c}} dx + \int \left(-\frac{3B \tan^2(e + fx)}{\sqrt{-ictan(e + fx) + c}} \right) dx \\ & \quad + \int \frac{B \tan^4(e + fx)}{\sqrt{-ictan(e + fx) + c}} dx + \int \left(-\frac{3iA \tan^2(e + fx)}{\sqrt{-ictan(e + fx) + c}} \right) dx \\ & \quad \left. + \int \frac{iB \tan(e + fx)}{\sqrt{-ictan(e + fx) + c}} dx + \int \left(-\frac{3iB \tan^3(e + fx)}{\sqrt{-ictan(e + fx) + c}} \right) dx \right) \end{aligned}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(1/2),x)`

output `-I*a**3*(Integral(I*A/sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-3*A*tan(e + f*x)/sqrt(-I*c*tan(e + f*x) + c), x) + Integral(A*tan(e + f*x)**3/sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-3*B*tan(e + f*x)**2/sqrt(-I*c*tan(e + f*x) + c), x) + Integral(B*tan(e + f*x)**4/sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-3*I*A*tan(e + f*x)**2/sqrt(-I*c*tan(e + f*x) + c), x) + Integral(I*B*tan(e + f*x)/sqrt(-I*c*tan(e + f*x) + c), x) + Integral(-3*I*B*tan(e + f*x)**3/sqrt(-I*c*tan(e + f*x) + c), x))`

3.760.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.77

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx = \\ & \frac{2i \left(\frac{60(A-iB)a^3c}{\sqrt{-ictan(fx+e)+c}} - \frac{3i(-ictan(fx+e)+c)^{\frac{5}{2}}Ba^3+5(-ictan(fx+e)+c)^{\frac{3}{2}}(A-5iB)a^3c-60\sqrt{-ictan(fx+e)+c}(A-2iB)a^3c^2}{c^2} \right)}{15cf} \end{aligned}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x, algorithm="maxima")`

3.760. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} dx$

output
$$-2/15*I*(60*(A - I*B)*a^3*c/\sqrt{-I*c*\tan(f*x + e) + c} - (3*I*(-I*c*\tan(f*x + e) + c)^{(5/2)}*B*a^3 + 5*(-I*c*\tan(f*x + e) + c)^{(3/2)}*(A - 5*I*B)*a^3*c - 60*\sqrt{-I*c*\tan(f*x + e) + c}*(A - 2*I*B)*a^3*c^2)/c^2)/(c*f)$$

3.760.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx$$

$$= \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^3}{\sqrt{-ictan(fx + e) + c}} dx$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^3/sqrt(-I*c*tan(f*x + e) + c), x)`

3.760.9 Mupad [B] (verification not implemented)

Time = 11.89 (sec) , antiderivative size = 351, normalized size of antiderivative = 2.51

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx$$

$$= -\sqrt{c + \frac{c(e^{e2i+fx2i} li - i) li}{e^{e2i+fx2i} + 1}} \left(\frac{a^3 (A - B li) 4i}{cf} + \frac{a^3 e^{e2i+fx2i} (A - B li) 4i}{cf} \right)$$

$$- \left(\frac{a^3 (A - B li) 4i}{cf} + \frac{a^3 (A - B 3i) 4i}{cf} \right) \sqrt{c + \frac{c(e^{e2i+fx2i} li - i) li}{e^{e2i+fx2i} + 1}}$$

$$- \frac{\left(\frac{a^3 (A - B li) 4i}{5cf} - \frac{a^3 (A + B li) 4i}{5cf} \right) \sqrt{c + \frac{c(e^{e2i+fx2i} li - i) li}{e^{e2i+fx2i} + 1}}}{(e^{e2i+fx2i} + 1)^2}$$

$$+ \frac{\left(\frac{16Ba^3}{3cf} + \frac{a^3 (A - B li) 4i}{3cf} \right) \sqrt{c + \frac{c(e^{e2i+fx2i} li - i) li}{e^{e2i+fx2i} + 1}}}{e^{e2i+fx2i} + 1}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*li)^3)/(c - c*tan(e + f*x)*li)^(1/2),x)`

3.760.
$$\int \frac{(a+ia \tan(e+fx))^3 (A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} dx$$

output $((a^3(A - B \tan(e+fx))^4)/(3cf) + (16Ba^3)/(3cf))(c + (c(\exp(e^2 + f^2x^2) \tan(e+fx) - 1) \tan(e+fx))/(\exp(e^2 + f^2x^2) + 1))^{1/2})/(\exp(e^2 + f^2x^2) + 1) - ((a^3(A - B \tan(e+fx))^4)/(cf) + (a^3(A - B \tan(e+fx))^4)/(cf))(c + (c(\exp(e^2 + f^2x^2) \tan(e+fx) - 1) \tan(e+fx))/(\exp(e^2 + f^2x^2) + 1))^{1/2} - (((a^3(A - B \tan(e+fx))^4)/(5cf) - (a^3(A + B \tan(e+fx))^4)/(5cf))(c + (c(\exp(e^2 + f^2x^2) \tan(e+fx) - 1) \tan(e+fx))/(\exp(e^2 + f^2x^2) + 1))^{1/2})/(\exp(e^2 + f^2x^2) + 1)^2 - (c + (c(\exp(e^2 + f^2x^2) \tan(e+fx) - 1) \tan(e+fx))/(\exp(e^2 + f^2x^2) + 1))^{1/2}) * ((a^3(A - B \tan(e+fx))^4)/(cf) + (a^3 \exp(e^2 + f^2x^2) (A - B \tan(e+fx))^4)/(cf))$

3.760. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$

3.761
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ictan(e+fx))^{3/2}} dx$$

3.761.1 Optimal result 6920
 3.761.2 Mathematica [A] (verified) 6920
 3.761.3 Rubi [A] (verified) 6921
 3.761.4 Maple [A] (verified) 6923
 3.761.5 Fricas [A] (verification not implemented) 6923
 3.761.6 Sympy [F] 6924
 3.761.7 Maxima [A] (verification not implemented) 6925
 3.761.8 Giac [F] 6925
 3.761.9 Mupad [B] (verification not implemented) 6925

3.761.1 Optimal result

Integrand size = 43, antiderivative size = 140

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ictan(e + fx))^{3/2}} dx =$$

$$-\frac{8a^3(iA + B)}{3f(c - ictan(e + fx))^{3/2}} + \frac{8a^3(iA + 2B)}{cf\sqrt{c - ictan(e + fx)}}$$

$$+ \frac{2a^3(iA + 5B)\sqrt{c - ictan(e + fx)}}{c^2f} - \frac{2a^3B(c - ictan(e + fx))^{3/2}}{3c^3f}$$

output `8*a^3*(I*A+2*B)/c/f/((c-I*c*tan(f*x+e))^(1/2)+2*a^3*(I*A+5*B)*(c-I*c*tan(f*x+e))^(1/2)/c^2/f-8/3*a^3*(I*A+B)/f/(c-I*c*tan(f*x+e))^(3/2)-2/3*a^3*B*(c-I*c*tan(f*x+e))^(3/2)/c^3/f`

3.761.2 Mathematica [A] (verified)

Time = 5.83 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.68

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ictan(e + fx))^{3/2}} dx = \frac{2a^3(-11A + 34iB + 3(6iA + 17B) \tan(e + fx) + 3(A - 3cf(i + \tan(e + fx))\sqrt{c - ic$$

input `Integrate[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(3/2), x]`

```
output (2*a^3*(-11*A + (34*I)*B + 3*((6*I)*A + 17*B)*Tan[e + f*x] + 3*(A - (4*I)*
B)*Tan[e + f*x]^2 + B*Tan[e + f*x]^3))/(3*c*f*(I + Tan[e + f*x])*Sqrt[c -
I*c*Tan[e + f*x]])
```

3.761.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.91, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx$$

↓ 4071

$$\frac{ac \int \frac{a^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} d \tan(e + fx)}{f}$$

↓ 27

$$\frac{a^3 c \int \frac{(i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} d \tan(e + fx)}{f}$$

↓ 86

$$\frac{a^3 c \int \left(\frac{4(A - iB)}{(c - ic \tan(e + fx))^{5/2}} + \frac{iB \sqrt{c - ic \tan(e + fx)}}{c^3} + \frac{A - 5iB}{c^2 \sqrt{c - ic \tan(e + fx)}} - \frac{4(A - 2iB)}{c(c - ic \tan(e + fx))^{3/2}} \right) d \tan(e + fx)}{f}$$

↓ 2009

$$\frac{a^3 c \left(\frac{2(5B + iA) \sqrt{c - ic \tan(e + fx)}}{c^3} + \frac{8(2B + iA)}{c^2 \sqrt{c - ic \tan(e + fx)}} - \frac{8(B + iA)}{3c(c - ic \tan(e + fx))^{3/2}} - \frac{2B(c - ic \tan(e + fx))^{3/2}}{3c^4} \right)}{f}$$

```
input Int[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])
^(3/2),x]
```

3.761. $\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx$


```
output (a^3*c*((-8*(I*A + B))/(3*c*(c - I*c*Tan[e + f*x])^(3/2)) + (8*(I*A + 2*B)
)/(c^2*Sqrt[c - I*c*Tan[e + f*x]]) + (2*(I*A + 5*B)*Sqrt[c - I*c*Tan[e + f
*x]])/c^3 - (2*B*(c - I*c*Tan[e + f*x])^(3/2))/(3*c^4))/f
```

3.761.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 86 Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_
.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;
FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1
] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p
+ 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.761.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.84

method	result
derivativedivides	$\frac{2ia^3 \left(\frac{iB(c-ic \tan(fx+e))^{3/2}}{3} - 5i\sqrt{c-ic \tan(fx+e)} Bc + \sqrt{c-ic \tan(fx+e)} cA + \frac{4c^2(-2iB+A)}{\sqrt{c-ic \tan(fx+e)}} - \frac{4c^3(-iB+A)}{3(c-ic \tan(fx+e))^{3/2}} \right)}{f c^3}$
default	$\frac{2ia^3 \left(\frac{iB(c-ic \tan(fx+e))^{3/2}}{3} - 5i\sqrt{c-ic \tan(fx+e)} Bc + \sqrt{c-ic \tan(fx+e)} cA + \frac{4c^2(-2iB+A)}{\sqrt{c-ic \tan(fx+e)}} - \frac{4c^3(-iB+A)}{3(c-ic \tan(fx+e))^{3/2}} \right)}{f c^3}$
parts	$2ia^3 Ac \left(-\frac{1}{4c^2 \sqrt{c-ic \tan(fx+e)}} - \frac{1}{6c(c-ic \tan(fx+e))^{3/2}} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{8c^{5/2}} \right) + a^3(3iA+B) \left(-\frac{1}{3(c-ic \tan(fx+e))^{3/2}} \right)$

input `int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

output `2*I/f*a^3/c^3*(1/3*I*B*(c-I*c*tan(f*x+e))^(3/2)-5*I*(c-I*c*tan(f*x+e))^(1/2)*B*c+(c-I*c*tan(f*x+e))^(1/2)*c*A+4*c^2*(A-2*I*B)/(c-I*c*tan(f*x+e))^(1/2)-4/3*c^3*(A-I*B)/(c-I*c*tan(f*x+e))^(3/2))`

3.761.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.84

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \frac{2\sqrt{2}((iA + B)a^3 e^{(6i fx + 6ie)} + 3(-iA - 3B)a^3 e^{(4i fx + 4ie)} + 12(-iA - 3B)a^3 e^{(2i fx + 2ie)} + 8(-iA - 3B)a^3) \sqrt{c/(e^{(2i fx + 2ie)} + 1)}}{3(c^2 f e^{(2i fx + 2ie)} + c^2 f)}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x,algorithm="fracas")`

output `-2/3*sqrt(2)*((I*A + B)*a^3*e^(6*I*f*x + 6*I*e) + 3*(-I*A - 3*B)*a^3*e^(4*I*f*x + 4*I*e) + 12*(-I*A - 3*B)*a^3*e^(2*I*f*x + 2*I*e) + 8*(-I*A - 3*B)*a^3)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^2*f*e^(2*I*f*x + 2*I*e) + c^2*f)`

3.761. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$

3.761.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx =$$

$$-ia^3 \left(\int \frac{iA}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} dx \right.$$

$$+ \int \left(-\frac{3A \tan(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} \right) dx$$

$$+ \int \frac{A \tan^3(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} dx$$

$$+ \int \left(-\frac{3B \tan^2(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} \right) dx$$

$$+ \int \frac{B \tan^4(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} dx$$

$$+ \int \left(-\frac{3iA \tan^2(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} \right) dx$$

$$+ \int \frac{iB \tan(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} dx$$

$$+ \int \left(-\frac{3iB \tan^3(e + fx)}{-ic\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) + c\sqrt{-ic \tan(e + fx) + c}} \right) dx \Big)$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(3/2),x)`

output `-I*a**3*(Integral(I*A/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(-3*A*tan(e + f*x)/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(A*tan(e + f*x)**3/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(-3*B*tan(e + f*x)**2/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(B*tan(e + f*x)**4/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(-3*I*A*tan(e + f*x)**2/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(I*B*tan(e + f*x)/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(-3*I*B*tan(e + f*x)**3/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c*sqrt(-I*c*tan(e + f*x) + c)), x))`

3.761. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$

3.761.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.75

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \frac{2i \left(\frac{4(3(-ic \tan(fx+e)+c)(A-2iB)a^3 - (A-iB)a^3c)}{(-ic \tan(fx+e)+c)^{3/2}} + \frac{i(-ic \tan(fx+e)+c)}{3cf} \right)}{3cf}$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2), x, algorithm="maxima")`

output `2/3*I*(4*(3*(-I*c*tan(f*x + e) + c)*(A - 2*I*B)*a^3 - (A - I*B)*a^3*c)/(-I*c*tan(f*x + e) + c)^(3/2) + (I*(-I*c*tan(f*x + e) + c)^(3/2)*B*a^3 + 3*sqrt(-I*c*tan(f*x + e) + c)*(A - 5*I*B)*a^3*c)/c^2)/(c*f)`

3.761.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^3}{(-ic \tan(fx + e) + c)^{3/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2), x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^3/(-I*c*tan(f*x + e) + c)^(3/2), x)`

3.761.9 Mupad [B] (verification not implemented)

Time = 10.46 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.58

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \frac{a^3 \sqrt{\frac{c(\cos(2e+2fx)+1-\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}}}{(A 20i + 60 B + A \cos(2e+2fx))^{3/2}}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3)/(c - c*tan(e + f*x)*1i)^(3/2), x)`

3.761. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$

output $(a^3((c(\cos(2e + 2fx) - \sin(2e + 2fx)*1i + 1))/(\cos(2e + 2fx) + 1))^{1/2}*(A*20i + 60*B + A*\cos(2e + 2fx)*23i + A*\cos(4e + 4fx)*2i - A*\cos(6e + 6fx)*1i + 69*B*\cos(2e + 2fx) + 8*B*\cos(4e + 4fx) - B*\cos(6e + 6fx) - 7*A*\sin(2e + 2fx) - 2*A*\sin(4e + 4fx) + A*\sin(6e + 6fx) + B*\sin(2e + 2fx)*21i + B*\sin(4e + 4fx)*8i - B*\sin(6e + 6fx)*1i))/(3*c^2*f*(\cos(2e + 2fx) + 1))$

3.761. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{3/2}} dx$

3.762
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ictan(e+fx))^{5/2}} dx$$

3.762.1 Optimal result 6927
 3.762.2 Mathematica [A] (verified) 6927
 3.762.3 Rubi [A] (verified) 6928
 3.762.4 Maple [A] (verified) 6929
 3.762.5 Fricas [A] (verification not implemented) 6930
 3.762.6 Sympy [F] 6931
 3.762.7 Maxima [A] (verification not implemented) 6932
 3.762.8 Giac [F] 6933
 3.762.9 Mupad [B] (verification not implemented) 6933

3.762.1 Optimal result

Integrand size = 43, antiderivative size = 140

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ictan(e + fx))^{5/2}} dx = -\frac{8a^3(iA + B)}{5f(c - ictan(e + fx))^{5/2}} + \frac{8a^3(iA + 2B)}{3cf(c - ictan(e + fx))^{3/2}} - \frac{2a^3(iA + 5B)}{c^2f\sqrt{c - ictan(e + fx)}} - \frac{2a^3B\sqrt{c - ictan(e + fx)}}{c^3f}$$

```
output -2*a^3*(I*A+5*B)/c^2/f/(c-I*c*tan(f*x+e))^(1/2)-2*a^3*B*(c-I*c*tan(f*x+e))
^(1/2)/c^3/f-8/5*a^3*(I*A+B)/f/(c-I*c*tan(f*x+e))^(5/2)+8/3*a^3*(I*A+2*B)/
c/f/(c-I*c*tan(f*x+e))^(3/2)
```

3.762.2 Mathematica [A] (verified)

Time = 6.25 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.71

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ictan(e + fx))^{5/2}} dx = \frac{2a^3(7iA + 62B + 5(2A - 31iB) \tan(e + fx) - 15i(A - (8I)*B)*\tan[e + f*x]^2 + (15*I)*B*\tan[e + f*x]^3))/(15*c^2*f*(I + \tan[e + f*x]))^2*\sqrt{c - I*c*\tan[e + f*x]}}$$

```
input Integrate[(((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e +
f*x])^(5/2), x]
```

```
output (2*a^3*((7*I)*A + 62*B + 5*(2*A - (31*I)*B)*Tan[e + f*x] - (15*I)*(A - (8*
I)*B)*Tan[e + f*x]^2 + (15*I)*B*Tan[e + f*x]^3))/(15*c^2*f*(I + Tan[e + f*
x]))^2*Sqrt[c - I*c*Tan[e + f*x]]
```

3.762.
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ictan(e+fx))^{5/2}} dx$$

3.762.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.91, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{a^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^3 c \int \frac{(i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{86} \\
 & \frac{a^3 c \int \left(\frac{4(A - iB)}{(c - ic \tan(e + fx))^{7/2}} + \frac{iB}{c^3 \sqrt{c - ic \tan(e + fx)}} + \frac{A - 5iB}{c^2 (c - ic \tan(e + fx))^{3/2}} - \frac{4(A - 2iB)}{c (c - ic \tan(e + fx))^{5/2}} \right) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^3 c \left(-\frac{2(5B + iA)}{c^3 \sqrt{c - ic \tan(e + fx)}} + \frac{8(2B + iA)}{3c^2 (c - ic \tan(e + fx))^{3/2}} - \frac{8(B + iA)}{5c (c - ic \tan(e + fx))^{5/2}} - \frac{2B \sqrt{c - ic \tan(e + fx)}}{c^4} \right)}{f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(5/2),x]`

output `(a^3*c*((-8*(I*A + B))/(5*c*(c - I*c*Tan[e + f*x])^(5/2)) + (8*(I*A + 2*B))/(3*c^2*(c - I*c*Tan[e + f*x])^(3/2)) - (2*(I*A + 5*B))/(c^3*Sqrt[c - I*c*Tan[e + f*x]]) - (2*B*Sqrt[c - I*c*Tan[e + f*x]])/c^4)/f`

3.762. $\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx$

3.762.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.762.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.75

method	result
derivativedivides	$\frac{2ia^3 \left(i\sqrt{c-ic \tan(fx+e)} B + \frac{4c^2(-2iB+A)}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{c(-5iB+A)}{\sqrt{c-ic \tan(fx+e)}} - \frac{4c^3(-iB+A)}{5(c-ic \tan(fx+e))^{\frac{5}{2}}} \right)}{f c^3}$
default	$\frac{2ia^3 \left(i\sqrt{c-ic \tan(fx+e)} B + \frac{4c^2(-2iB+A)}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{c(-5iB+A)}{\sqrt{c-ic \tan(fx+e)}} - \frac{4c^3(-iB+A)}{5(c-ic \tan(fx+e))^{\frac{5}{2}}} \right)}{f c^3}$
parts	$\frac{2ia^3 Ac \left(-\frac{1}{8c^3 \sqrt{c-ic \tan(fx+e)}} - \frac{1}{12c^2(c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{1}{10c(c-ic \tan(fx+e))^{\frac{5}{2}}} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{16c^{\frac{7}{2}}} \right)}{f} +$

3.762. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$


```
input int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)
```

```
output 2*I/f*a^3/c^3*(I*(c-I*c*tan(f*x+e))^(1/2)*B+4/3*c^2*(A-2*I*B)/(c-I*c*tan(f*x+e))^(3/2)-c*(A-5*I*B)/(c-I*c*tan(f*x+e))^(1/2)-4/5*c^3*(A-I*B)/(c-I*c*tan(f*x+e))^(5/2))
```

3.762.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.73

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx =$$

$$\frac{\sqrt{2}(3(iA + B)a^3 e^{(6i fx + 6i e)} - (iA + 11B)a^3 e^{(4i fx + 4i e)} + 4(iA + 11B)a^3 e^{(2i fx + 2i e)} + 8(iA + 11B)a^3)}{15c^3 f}$$

```
input integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,algorithm="fricas")
```

```
output -1/15*sqrt(2)*(3*(I*A + B)*a^3*e^(6*I*f*x + 6*I*e) - (I*A + 11*B)*a^3*e^(4*I*f*x + 4*I*e) + 4*(I*A + 11*B)*a^3*e^(2*I*f*x + 2*I*e) + 8*(I*A + 11*B)*a^3)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^3*f)
```

3.762.6 Sympy [F]

$$\begin{aligned}
& \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \\
& -ia^3 \left(\int \frac{iA}{-c^2 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 2ic^2 \sqrt{-ic \tan(e + fx) + c \tan(e + fx)} + c^2 \sqrt{-ic \tan(e + fx)}} \right. \\
& + \int \left(-\frac{3A \tan(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 2ic^2 \sqrt{-ic \tan(e + fx) + c \tan(e + fx)} + c^2 \sqrt{-ic \tan(e + fx)}} \right. \\
& + \int \frac{A \tan^3(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 2ic^2 \sqrt{-ic \tan(e + fx) + c \tan(e + fx)} + c^2 \sqrt{-ic \tan(e + fx)}} \\
& + \int \left(-\frac{3B \tan^2(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 2ic^2 \sqrt{-ic \tan(e + fx) + c \tan(e + fx)} + c^2 \sqrt{-ic \tan(e + fx)}} \right. \\
& + \int \frac{B \tan^4(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 2ic^2 \sqrt{-ic \tan(e + fx) + c \tan(e + fx)} + c^2 \sqrt{-ic \tan(e + fx)}} \\
& + \int \left(-\frac{3iA \tan^2(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 2ic^2 \sqrt{-ic \tan(e + fx) + c \tan(e + fx)} + c^2 \sqrt{-ic \tan(e + fx)}} \right. \\
& + \int \frac{iB \tan(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 2ic^2 \sqrt{-ic \tan(e + fx) + c \tan(e + fx)} + c^2 \sqrt{-ic \tan(e + fx)}} \\
& \left. + \int \left(-\frac{3iB \tan^3(e + fx)}{-c^2 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 2ic^2 \sqrt{-ic \tan(e + fx) + c \tan(e + fx)} + c^2 \sqrt{-ic \tan(e + fx)}} \right) \right)
\end{aligned}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(5/2),x)`

output

```
-I*a**3*(Integral(I*A/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 -
  2*I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**2*sqrt(-I*c*tan(e
+ f*x) + c)), x) + Integral(-3*A*tan(e + f*x)/(-c**2*sqrt(-I*c*tan(e + f*x
) + c)*tan(e + f*x)**2 - 2*I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)
+ c**2*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(A*tan(e + f*x)**3/(-c*
**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 2*I*c**2*sqrt(-I*c*tan(e
+ f*x) + c)*tan(e + f*x) + c**2*sqrt(-I*c*tan(e + f*x) + c)), x) + Integra
l(-3*B*tan(e + f*x)**2/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2
- 2*I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**2*sqrt(-I*c*tan(e
+ f*x) + c)), x) + Integral(B*tan(e + f*x)**4/(-c**2*sqrt(-I*c*tan(e + f*
x) + c)*tan(e + f*x)**2 - 2*I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x
) + c**2*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(-3*I*A*tan(e + f*x)**
2/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 2*I*c**2*sqrt(-I*c*
tan(e + f*x) + c)*tan(e + f*x) + c**2*sqrt(-I*c*tan(e + f*x) + c)), x) + I
ntegral(I*B*tan(e + f*x)/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**
2 - 2*I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**2*sqrt(-I*c*tan
(e + f*x) + c)), x) + Integral(-3*I*B*tan(e + f*x)**3/(-c**2*sqrt(-I*c*tan
(e + f*x) + c)*tan(e + f*x)**2 - 2*I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(
e + f*x) + c**2*sqrt(-I*c*tan(e + f*x) + c)), x))
```

3.762.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.76

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx =$$

$$\frac{2i \left(-\frac{15i \sqrt{-ic \tan(fx+e)+c} B a^3}{c^2} + \frac{15(-ic \tan(fx+e)+c)^2 (A-5iB)a^3 - 20(-ic \tan(fx+e)+c)(A-2iB)a^3 c + 12(A-iB)a^3 c^2}{(-ic \tan(fx+e)+c)^{5/2} c} \right)}{15cf}$$

input

```
integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x
, algorithm="maxima")
```

output

```
-2/15*I*(-15*I*sqrt(-I*c*tan(f*x + e) + c)*B*a^3/c^2 + (15*(-I*c*tan(f*x +
e) + c)^2*(A - 5*I*B)*a^3 - 20*(-I*c*tan(f*x + e) + c)*(A - 2*I*B)*a^3*c
+ 12*(A - I*B)*a^3*c^2)/((-I*c*tan(f*x + e) + c)^(5/2)*c))/(c*f)
```

3.762.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^3}{(-ic \tan(fx + e) + c)^{5/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^3/(-I*c*tan(f*x + e)
+ c)^(5/2), x)`

3.762.9 Mupad [B] (verification not implemented)

Time = 9.89 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.49

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx =$$

$$\frac{a^3 \sqrt{\frac{c(\cos(2e+2fx)+1)-\sin(2e+2fx) \operatorname{li}}{\cos(2e+2fx)+1}} (A 8i + 88 B + A \cos(2e + 2fx) 4i - A \cos(4e + 4fx) \operatorname{li} + A \cos(6e + 6fx) 3i + 44 B \cos(2e + 2fx) - 11 B \cos(4e + 4fx) + 3 B \cos(6e + 6fx) - 4 A \sin(2e + 2fx) + A \sin(4e + 4fx) - 3 A \sin(6e + 6fx) + B \sin(2e + 2fx) 44i - B \sin(4e + 4fx) 11i + B \sin(6e + 6fx) 3i)}{(15c^3 f)}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3)/(c - c*tan(e + f*x)*1
i)^(5/2),x)`

output `-(a^3*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x)
+ 1))^(1/2)*(A*8i + 88*B + A*cos(2*e + 2*f*x)*4i - A*cos(4*e + 4*f*x)*1i +
A*cos(6*e + 6*f*x)*3i + 44*B*cos(2*e + 2*f*x) - 11*B*cos(4*e + 4*f*x) + 3
*B*cos(6*e + 6*f*x) - 4*A*sin(2*e + 2*f*x) + A*sin(4*e + 4*f*x) - 3*A*sin(
6*e + 6*f*x) + B*sin(2*e + 2*f*x)*44i - B*sin(4*e + 4*f*x)*11i + B*sin(6*e
+ 6*f*x)*3i))/(15*c^3*f)`

3.763
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ictan(e+fx))^{7/2}} dx$$

3.763.1 Optimal result 6934
 3.763.2 Mathematica [A] (verified) 6934
 3.763.3 Rubi [A] (verified) 6935
 3.763.4 Maple [A] (verified) 6936
 3.763.5 Fricas [A] (verification not implemented) 6937
 3.763.6 Sympy [F] 6938
 3.763.7 Maxima [A] (verification not implemented) 6939
 3.763.8 Giac [F] 6940
 3.763.9 Mupad [B] (verification not implemented) 6940

3.763.1 Optimal result

Integrand size = 43, antiderivative size = 142

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx = -\frac{8a^3(iA + B)}{7f(c - ictan(e + fx))^{7/2}} + \frac{8a^3(iA + 2B)}{5cf(c - ictan(e + fx))^{5/2}} - \frac{2a^3(iA + 5B)}{3c^2f(c - ictan(e + fx))^{3/2}} + \frac{2a^3B}{c^3f\sqrt{c - ictan(e + fx)}}$$

output `2*a^3*B/c^3/f/(c-I*c*tan(f*x+e))^(1/2)-8/7*a^3*(I*A+B)/f/(c-I*c*tan(f*x+e))^(7/2)+8/5*a^3*(I*A+2*B)/c/f/(c-I*c*tan(f*x+e))^(5/2)-2/3*a^3*(I*A+5*B)/c^2/f/(c-I*c*tan(f*x+e))^(3/2)`

3.763.2 Mathematica [A] (verified)

Time = 6.37 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.67

$$\int \frac{(a + ia \tan(e + fx))^3(A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx = \frac{2a^3(-11A - 38iB + (-14iA - 133B) \tan(e + fx) + 35)}{105c^3f(i + \tan(e + fx))^3\sqrt{}}$$

input `Integrate[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(7/2), x]`

output `(2*a^3*(-11*A - (38*I)*B + ((-14*I)*A - 133*B)*Tan[e + f*x] + 35*(A + (4*I)*B)*Tan[e + f*x]^2 + 105*B*Tan[e + f*x]^3)/(105*c^3*f*(I + Tan[e + f*x])^3*Sqrt[c - I*c*Tan[e + f*x]])`

3.763.
$$\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ictan(e+fx))^{7/2}} dx$$

3.763.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.91, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.116$, Rules used = {3042, 4071, 27, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{a^2 (i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{a^3 c \int \frac{(i \tan(e + fx) + 1)^2 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{86} \\
 & \frac{a^3 c \int \left(\frac{4(A - iB)}{(c - ic \tan(e + fx))^{9/2}} + \frac{iB}{c^3 (c - ic \tan(e + fx))^{3/2}} + \frac{A - 5iB}{c^2 (c - ic \tan(e + fx))^{5/2}} - \frac{4(A - 2iB)}{c (c - ic \tan(e + fx))^{7/2}} \right) d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{2009} \\
 & \frac{a^3 c \left(-\frac{2(5B + iA)}{3c^3 (c - ic \tan(e + fx))^{3/2}} + \frac{8(2B + iA)}{5c^2 (c - ic \tan(e + fx))^{5/2}} - \frac{8(B + iA)}{7c (c - ic \tan(e + fx))^{7/2}} + \frac{2B}{c^4 \sqrt{c - ic \tan(e + fx)}} \right)}{f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])^3*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(7/2),x]`

output `(a^3*c*((-8*(I*A + B))/(7*c*(c - I*c*Tan[e + f*x])^(7/2)) + (8*(I*A + 2*B))/(5*c^2*(c - I*c*Tan[e + f*x])^(5/2)) - (2*(I*A + 5*B))/(3*c^3*(c - I*c*Tan[e + f*x])^(3/2)) + (2*B)/(c^4*Sqrt[c - I*c*Tan[e + f*x]])))/f`

3.763. $\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx$

3.763.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.763.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.74

method	result
derivativedivides	$\frac{2ia^3 \left(\frac{4c^2(-2iB+A)}{5(c-ic \tan(fx+e))^{\frac{5}{2}}} - \frac{iB}{\sqrt{c-ic \tan(fx+e)}} - \frac{4c^3(-iB+A)}{7(c-ic \tan(fx+e))^{\frac{7}{2}}} - \frac{c(-5iB+A)}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} \right)}{f c^3}$
default	$\frac{2ia^3 \left(\frac{4c^2(-2iB+A)}{5(c-ic \tan(fx+e))^{\frac{5}{2}}} - \frac{iB}{\sqrt{c-ic \tan(fx+e)}} - \frac{4c^3(-iB+A)}{7(c-ic \tan(fx+e))^{\frac{7}{2}}} - \frac{c(-5iB+A)}{3(c-ic \tan(fx+e))^{\frac{3}{2}}} \right)}{f c^3}$
risch	$-\frac{a^3(15iA e^{6i(fx+e)} + 15B e^{6i(fx+e)} + 3iA e^{4i(fx+e)} - 39B e^{4i(fx+e)} - 4iA e^{2i(fx+e)} + 52B e^{2i(fx+e)} + 8iA - 104B)\sqrt{2}}{210c^3 \sqrt{\frac{c}{e^{2i(fx+e)}+1}} f}$
parts	$\frac{2ia^3 Ac \left(-\frac{1}{16c^4 \sqrt{c-ic \tan(fx+e)}} - \frac{1}{24c^3 (c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{1}{20c^2 (c-ic \tan(fx+e))^{\frac{5}{2}}} - \frac{1}{14c (c-ic \tan(fx+e))^{\frac{7}{2}}} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)}}{e^{i(fx+e)}}\right)}{f} \right)}{f}$

3.763. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$

```
input int((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x,method=_RETURNVERBOSE)
```

```
output 2*I/f*a^3/c^3*(4/5*c^2*(A-2*I*B)/(c-I*c*tan(f*x+e))^(5/2)-I*B/(c-I*c*tan(f*x+e))^(1/2)-4/7*c^3*(A-I*B)/(c-I*c*tan(f*x+e))^(7/2)-1/3*c*(A-5*I*B)/(c-I*c*tan(f*x+e))^(3/2))
```

3.763.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.87

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx =$$

$$\frac{\sqrt{2}(15(iA + B)a^3 e^{(8i fx + 8ie)} + 6(3iA - 4B)a^3 e^{(6i fx + 6ie)} - (iA - 13B)a^3 e^{(4i fx + 4ie)} + 4(iA - 13B)a^3 e^{(2i fx + 2ie)})}{210 c^4 f}$$

```
input integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x,algorithm="fricas")
```

```
output -1/210*sqrt(2)*(15*(I*A + B)*a^3*e^(8*I*f*x + 8*I*e) + 6*(3*I*A - 4*B)*a^3*e^(6*I*f*x + 6*I*e) - (I*A - 13*B)*a^3*e^(4*I*f*x + 4*I*e) + 4*(I*A - 13*B)*a^3*e^(2*I*f*x + 2*I*e) + 8*(I*A - 13*B)*a^3)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^4*f)
```


3.763.6 Sympy [F]

$$\begin{aligned}
& \int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \\
& -ia^3 \left(\int \frac{iA}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)}} dx \right. \\
& + \int \left(-\frac{3A \tan(e + fx)}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)}} \right. \\
& + \int \frac{A \tan^3(e + fx)}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)}} dx \\
& + \int \left(-\frac{3B \tan^2(e + fx)}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)}} \right. \\
& + \int \frac{B \tan^4(e + fx)}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)}} dx \\
& + \int \left(-\frac{3iA \tan^2(e + fx)}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)}} \right. \\
& + \int \frac{iB \tan(e + fx)}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)}} dx \\
& \left. + \int \left(-\frac{3iB \tan^3(e + fx)}{ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)} - 3c^3 \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx)} - 3ic^3 \sqrt{-ic \tan(e + fx) + c \tan^3(e + fx)}} \right) dx \right)
\end{aligned}$$

input `integrate((a+I*a*tan(f*x+e))**3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(7/2),x)`

```

output -I*a**3*(Integral(I*A/(I*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3
- 3*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 3*I*c**3*sqrt(-I*c*
tan(e + f*x) + c)*tan(e + f*x) + c**3*sqrt(-I*c*tan(e + f*x) + c)), x) + I
ntegral(-3*A*tan(e + f*x)/(I*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)
**3 - 3*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 3*I*c**3*sqrt(-
I*c*tan(e + f*x) + c)*tan(e + f*x) + c**3*sqrt(-I*c*tan(e + f*x) + c)), x)
+ Integral(A*tan(e + f*x)**3/(I*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e +
f*x)**3 - 3*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 3*I*c**3*sq
rt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**3*sqrt(-I*c*tan(e + f*x) + c))
, x) + Integral(-3*B*tan(e + f*x)**2/(I*c**3*sqrt(-I*c*tan(e + f*x) + c)*t
an(e + f*x)**3 - 3*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - 3*I*
c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**3*sqrt(-I*c*tan(e + f*x
) + c)), x) + Integral(B*tan(e + f*x)**4/(I*c**3*sqrt(-I*c*tan(e + f*x) +
c)*tan(e + f*x)**3 - 3*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 -
3*I*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**3*sqrt(-I*c*tan(e +
f*x) + c)), x) + Integral(-3*I*A*tan(e + f*x)**2/(I*c**3*sqrt(-I*c*tan(e
+ f*x) + c)*tan(e + f*x)**3 - 3*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f
*x)**2 - 3*I*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + c**3*sqrt(-I*
c*tan(e + f*x) + c)), x) + Integral(I*B*tan(e + f*x)/(I*c**3*sqrt(-I*c*tan
(e + f*x) + c)*tan(e + f*x)**3 - 3*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan...

```

3.763.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.72

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \frac{2i (105i (-ic \tan(fx + e) + c)^3 Ba^3 + 35 (-ic \tan(fx + e) + c)^2 (A - 5iB) a^3 c - 84 (-ic \tan(fx + e) + c) + 60 (A - iB) a^3 c^3)}{105 (-ic \tan(fx + e) + c)^{7/2} c^3 f}$$

```

input integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x
, algorithm="maxima")

```

```

output -2/105*I*(105*I*(-I*c*tan(f*x + e) + c)^3*B*a^3 + 35*(-I*c*tan(f*x + e) +
c)^2*(A - 5*I*B)*a^3*c - 84*(-I*c*tan(f*x + e) + c)*(A - 2*I*B)*a^3*c^2 +
60*(A - I*B)*a^3*c^3)/((-I*c*tan(f*x + e) + c)^(7/2)*c^3*f)

```

3.763. $\int \frac{(a+ia \tan(e+fx))^3(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$

3.763.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^3}{(-ic \tan(fx + e) + c)^{7/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^3*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^3/(-I*c*tan(f*x + e)
+ c)^(7/2), x)`

3.763.9 Mupad [B] (verification not implemented)

Time = 10.11 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.13

$$\int \frac{(a + ia \tan(e + fx))^3 (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx =$$

$$-\sqrt{c - \frac{c \sin(e + fx)}{\cos(e + fx)}} \left(\frac{a^3 (A + B 13i) 4i}{105 c^4 f} + \frac{a^3 e^{e 6i + f x 6i} (3A + B 4i) 1i}{35 c^4 f} \right)$$

$$+ \frac{a^3 e^{e 2i + f x 2i} (A + B 13i) 2i}{105 c^4 f} + \frac{a^3 e^{e 8i + f x 8i} (A - B 1i) 1i}{14 c^4 f} - \frac{a^3 e^{e 4i + f x 4i} (A + B 13i) 1i}{210 c^4 f}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^3)/(c - c*tan(e + f*x)*1
i)^(7/2),x)`

output `-(c - (c*sin(e + f*x)*1i)/cos(e + f*x))^(1/2)*((a^3*(A + B*13i)*4i)/(105*c
^4*f) + (a^3*exp(e*6i + f*x*6i)*(3*A + B*4i)*1i)/(35*c^4*f) + (a^3*exp(e*2
i + f*x*2i)*(A + B*13i)*2i)/(105*c^4*f) + (a^3*exp(e*8i + f*x*8i)*(A - B*1
i)*1i)/(14*c^4*f) - (a^3*exp(e*4i + f*x*4i)*(A + B*13i)*1i)/(210*c^4*f))`

3.764
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{a+ia \tan(e+fx)} dx$$

3.764.1 Optimal result 6941
 3.764.2 Mathematica [A] (verified) 6942
 3.764.3 Rubi [A] (verified) 6942
 3.764.4 Maple [A] (verified) 6945
 3.764.5 Fricas [B] (verification not implemented) 6945
 3.764.6 Sympy [F] 6946
 3.764.7 Maxima [A] (verification not implemented) 6947
 3.764.8 Giac [F] 6947
 3.764.9 Mupad [B] (verification not implemented) 6948

3.764.1 Optimal result

Integrand size = 43, antiderivative size = 220

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2}}{a + ia \tan(e + fx)} dx =$$

$$\frac{2\sqrt{2}(5iA - 9B)c^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{af} + \frac{2(5iA - 9B)c^3 \sqrt{c - ic \tan(e + fx)}}{af} + \frac{(5iA - 9B)c^2 (c - ic \tan(e + fx))^{3/2}}{3af}$$

$$+ \frac{(5iA - 9B)c (c - ic \tan(e + fx))^{5/2}}{10af} + \frac{(iA - B)(c - ic \tan(e + fx))^{7/2}}{2af(1 + i \tan(e + fx))}$$

output

```
-2*(5*I*A-9*B)*c^(7/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))
)*2^(1/2)/a/f+2*(5*I*A-9*B)*c^3*(c-I*c*tan(f*x+e))^(1/2)/a/f+1/3*(5*I*A-9
*B)*c^2*(c-I*c*tan(f*x+e))^(3/2)/a/f+1/10*(5*I*A-9*B)*c*(c-I*c*tan(f*x+e))
^(5/2)/a/f+1/2*(I*A-B)*(c-I*c*tan(f*x+e))^(7/2)/a/f/(1+I*tan(f*x+e))
```

3.764.2 Mathematica [A] (verified)

Time = 7.75 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.67

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2}}{a + ia \tan(e + fx)} dx = \frac{2c^3 \left(15\sqrt{2}(-5iA + 9B)\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right) + \dots \right)}{a + ia \tan(e + fx)}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2))/(a + I*a*Tan[e + f*x]),x]`

output `(2*c^3*(15*Sqrt[2]*((-5*I)*A + 9*B)*Sqrt[c]*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])]) + (Sqrt[c - I*c*Tan[e + f*x]]*(95*A + (168*I)*B + ((60*I)*A - 117*B)*Tan[e + f*x] + (5*A + (18*I)*B)*Tan[e + f*x]^2 + 3*B*Tan[e + f*x]^3))/(-I + Tan[e + f*x]))/(15*a*f)`

3.764.3 Rubi [A] (verified)Time = 0.41 (sec) , antiderivative size = 188, normalized size of antiderivative = 0.85, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.209$, Rules used = {3042, 4071, 27, 87, 60, 60, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c - ic \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{a + ia \tan(e + fx)} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c - ic \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{a + ia \tan(e + fx)} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{a^2(i \tan(e+fx)+1)^2} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{c \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(i \tan(e+fx)+1)^2} d \tan(e + fx)}{af} \\ & \quad \downarrow \text{87} \end{aligned}$$

3.764. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{a+ia \tan(e+fx)} dx$

$$\begin{aligned}
& \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{7/2}}{2c(1+i\tan(e+fx))} - \frac{1}{4}(5A+9iB) \int \frac{(c-ictan(e+fx))^{5/2}}{i\tan(e+fx)+1} d\tan(e+fx)\right)}{af} \\
& \quad \downarrow 60 \\
& \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{7/2}}{2c(1+i\tan(e+fx))} - \frac{1}{4}(5A+9iB) \left(2c \int \frac{(c-ictan(e+fx))^{3/2}}{i\tan(e+fx)+1} d\tan(e+fx) - \frac{2}{5}i(c-ictan(e+fx))^{5/2}\right)\right)}{af} \\
& \quad \downarrow 60 \\
& \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{7/2}}{2c(1+i\tan(e+fx))} - \frac{1}{4}(5A+9iB) \left(2c\left(2c \int \frac{\sqrt{c-ictan(e+fx)}}{i\tan(e+fx)+1} d\tan(e+fx) - \frac{2}{3}i(c-ictan(e+fx))^{3/2}\right) - \frac{2}{5}i(c-ictan(e+fx))^{5/2}\right)\right)}{af} \\
& \quad \downarrow 60 \\
& \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{7/2}}{2c(1+i\tan(e+fx))} - \frac{1}{4}(5A+9iB) \left(2c\left(2c\left(2c \int \frac{1}{(i\tan(e+fx)+1)\sqrt{c-ictan(e+fx)}} d\tan(e+fx) - 2i\sqrt{c-ictan(e+fx)}\right) - \frac{2}{5}i(c-ictan(e+fx))^{5/2}\right) - \frac{2}{3}i(c-ictan(e+fx))^{3/2}\right)\right)}{af} \\
& \quad \downarrow 73 \\
& \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{7/2}}{2c(1+i\tan(e+fx))} - \frac{1}{4}(5A+9iB) \left(2c\left(2c\left(4i \int \frac{1}{2-\frac{c-ictan(e+fx)}{c}} d\sqrt{c-ictan(e+fx)} - 2i\sqrt{c-ictan(e+fx)}\right) - \frac{2}{5}i(c-ictan(e+fx))^{5/2}\right) - \frac{2}{3}i(c-ictan(e+fx))^{3/2}\right)\right)}{af} \\
& \quad \downarrow 219 \\
& \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{7/2}}{2c(1+i\tan(e+fx))} - \frac{1}{4}(5A+9iB) \left(2c\left(2c\left(2i\sqrt{2}\sqrt{c}\operatorname{arctanh}\left(\frac{\sqrt{c-ictan(e+fx)}}{\sqrt{2}\sqrt{c}}\right) - 2i\sqrt{c-ictan(e+fx)}\right) - \frac{2}{5}i(c-ictan(e+fx))^{5/2}\right) - \frac{2}{3}i(c-ictan(e+fx))^{3/2}\right)\right)}{af}
\end{aligned}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2))/(a + I*a*Tan[e + f*x]), x]`

output `(c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(7/2))/(2*c*(1 + I*Tan[e + f*x])) - ((5*A + (9*I)*B)*(((-2*I)/5)*(c - I*c*Tan[e + f*x])^(5/2) + 2*c*(((-2*I)/3)*(c - I*c*Tan[e + f*x])^(3/2) + 2*c*((2*I)*Sqrt[2]*Sqrt[c]*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c]]) - (2*I)*Sqrt[c - I*c*Tan[e + f*x]]))))/4)/(a*f)`

3.764. $\int \frac{(A+B \tan(e+fx))(c-ictan(e+fx))^{7/2}}{a+ia \tan(e+fx)} dx$

3.764.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 60 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 73 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_))*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.764.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.87

method	result
derivativedivides	$\frac{2ic \left(\frac{iB(c-ic \tan(fx+e))^{5/2}}{5} + iBc(c-ic \tan(fx+e))^{3/2} + \frac{Ac(c-ic \tan(fx+e))^{3/2}}{3} + 8i\sqrt{c-ic \tan(fx+e)} Bc^2 + 4\sqrt{c-ic \tan(fx+e)} \right)}{fa}$
default	$\frac{2ic \left(\frac{iB(c-ic \tan(fx+e))^{5/2}}{5} + iBc(c-ic \tan(fx+e))^{3/2} + \frac{Ac(c-ic \tan(fx+e))^{3/2}}{3} + 8i\sqrt{c-ic \tan(fx+e)} Bc^2 + 4\sqrt{c-ic \tan(fx+e)} \right)}{fa}$

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)`

output
$$2*I/f/a*c*(1/5*I*B*(c-I*c*tan(f*x+e))^(5/2)+I*B*c*(c-I*c*tan(f*x+e))^(3/2)+1/3*A*c*(c-I*c*tan(f*x+e))^(3/2)+8*I*(c-I*c*tan(f*x+e))^(1/2)*B*c^2+4*(c-I*c*tan(f*x+e))^(1/2)*A*c^2-4*c^3*((-1/4*I*B-1/4*A)*(c-I*c*tan(f*x+e))^(1/2)/(1/2*c+1/2*I*c*tan(f*x+e))+1/2*(9/2*I*B+5/2*A)*2^(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))))$$

3.764.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 466 vs. 2(173) = 346.

Time = 0.30 (sec) , antiderivative size = 466, normalized size of antiderivative = 2.12

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2}}{a + ia \tan(e + fx)} dx =$$

$$15\sqrt{2}\sqrt{-\frac{(25A^2+90iAB-81B^2)c^7}{a^2f^2}}(afe^{(6ifx+6ie)} + 2afe^{(4ifx+4ie)} + afe^{(2ifx+2ie)}) \log \left(\frac{8 \left((5iA-9B)c^4 + \sqrt{-\frac{(25A^2+90iAB-81B^2)c^7}{a^2f^2}} \right)}{\dots} \right)$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e)),x,algorithm="fracas")`

output
$$\begin{aligned} & -1/15*(15*\sqrt{2})*\sqrt{-(25*A^2 + 90*I*A*B - 81*B^2)*c^7/(a^2*f^2)}*(a*f*e^{(6*I*f*x + 6*I*e)} + 2*a*f*e^{(4*I*f*x + 4*I*e)} + a*f*e^{(2*I*f*x + 2*I*e)})* \\ & \log(-8*((5*I*A - 9*B)*c^4 + \sqrt{-(25*A^2 + 90*I*A*B - 81*B^2)*c^7/(a^2*f^2)})*(a*f*e^{(2*I*f*x + 2*I*e)} + a*f)*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)}))e^{(-I*f*x - I*e)/(a*f)} - 15*\sqrt{2}*\sqrt{-(25*A^2 + 90*I*A*B - 81*B^2)*c^7/(a^2*f^2)}*(a*f*e^{(6*I*f*x + 6*I*e)} + 2*a*f*e^{(4*I*f*x + 4*I*e)} + a*f*e^{(2*I*f*x + 2*I*e)})* \\ & \log(-8*((5*I*A - 9*B)*c^4 - \sqrt{-(25*A^2 + 90*I*A*B - 81*B^2)*c^7/(a^2*f^2)})*(a*f*e^{(2*I*f*x + 2*I*e)} + a*f)*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)}))e^{(-I*f*x - I*e)/(a*f)} + 2*\sqrt{2}*(15*(-5*I*A + 9*B)*c^3*e^{(6*I*f*x + 6*I*e)} + 35*(-5*I*A + 9*B)*c^3*e^{(4*I*f*x + 4*I*e)} + 23*(-5*I*A + 9*B)*c^3*e^{(2*I*f*x + 2*I*e)} + 15*(-I*A + B)*c^3)*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)})/(a*f*e^{(6*I*f*x + 6*I*e)} + 2*a*f*e^{(4*I*f*x + 4*I*e)} + a*f*e^{(2*I*f*x + 2*I*e)}) \end{aligned}$$

3.764.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{a + i a \tan(e + fx)} dx =$$

$$i \left(\int \frac{A c^3 \sqrt{-i c \tan(e + fx) + c}}{\tan(e + fx) - i} dx + \int \left(-\frac{3 A c^3 \sqrt{-i c \tan(e + fx) + c \tan^2(e + fx)}}{\tan(e + fx) - i} \right) dx + \int \frac{B c^3 \sqrt{-i c \tan(e + fx) + c \tan(e + fx)}}{\tan(e + fx) - i} dx + \int \right)$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(7/2)/(a+I*a*tan(f*x+e)),x)`

output
$$\begin{aligned} & -I*(Integral(A*c**3*\sqrt{-I*c*\tan(e + f*x) + c}/(\tan(e + f*x) - I), x) + I \\ & ntegral(-3*A*c**3*\sqrt{-I*c*\tan(e + f*x) + c)*\tan(e + f*x)**2/(\tan(e + f*x) \\ &) - I), x) + Integral(B*c**3*\sqrt{-I*c*\tan(e + f*x) + c)*\tan(e + f*x)/(\tan \\ & (e + f*x) - I), x) + Integral(-3*B*c**3*\sqrt{-I*c*\tan(e + f*x) + c)*\tan(e \\ & + f*x)**3/(\tan(e + f*x) - I), x) + Integral(-3*I*A*c**3*\sqrt{-I*c*\tan(e + \\ & f*x) + c)*\tan(e + f*x)/(\tan(e + f*x) - I), x) + Integral(I*A*c**3*\sqrt{-I* \\ & c*\tan(e + f*x) + c)*\tan(e + f*x)**3/(\tan(e + f*x) - I), x) + Integral(-3*I \\ & *B*c**3*\sqrt{-I*c*\tan(e + f*x) + c)*\tan(e + f*x)**2/(\tan(e + f*x) - I), x) \\ & + Integral(I*B*c**3*\sqrt{-I*c*\tan(e + f*x) + c)*\tan(e + f*x)**4/(\tan(e + \\ & f*x) - I), x))/a \end{aligned}$$

3.764.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.86

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{a + i a \tan(e + fx)} dx = \frac{i \left(\frac{15 \sqrt{2} (5A + 9iB) c^{9/2} \log\left(\frac{-\sqrt{2}\sqrt{c} - \sqrt{-i c \tan(fx+e)+c}}{\sqrt{2}\sqrt{c} + \sqrt{-i c \tan(fx+e)+c}}\right)}{a} - \frac{60 \sqrt{-i c \tan(fx+e)+c}}{(-i c \tan(fx+e)+c)} \right)}{a}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e)),x,
algorithm="maxima")`

output `1/15*I*(15*sqrt(2)*(5*A + 9*I*B)*c^(9/2)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c
*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a - 6
0*sqrt(-I*c*tan(f*x + e) + c)*(A + I*B)*c^5/((-I*c*tan(f*x + e) + c)*a - 2
*a*c) + 2*(3*I*(-I*c*tan(f*x + e) + c)^(5/2)*B*c^2 + 5*(-I*c*tan(f*x + e)
+ c)^(3/2)*(A + 3*I*B)*c^3 + 60*sqrt(-I*c*tan(f*x + e) + c)*(A + 2*I*B)*c^
4)/a)/(c*f)`

3.764.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{a + i a \tan(e + fx)} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{7/2}}{i a \tan(fx + e) + a} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e)),x,
algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(7/2)/(I*a*tan(f*x
+ e) + a), x)`

3.764.9 Mupad [B] (verification not implemented)

Time = 1.37 (sec) , antiderivative size = 298, normalized size of antiderivative = 1.35

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{a + i a \tan(e + fx)} dx = \frac{4 B c^4 \sqrt{c - c \tan(e + fx)} \operatorname{li}}{a f (c - c \tan(e + fx) \operatorname{li}) - 2 a c f}$$

$$+ \frac{A c^3 \sqrt{c - c \tan(e + fx)} \operatorname{li} 8i}{a f} + \frac{A c^2 (c - c \tan(e + fx) \operatorname{li})^{3/2} 2i}{3 a f}$$

$$- \frac{16 B c^3 \sqrt{c - c \tan(e + fx)} \operatorname{li}}{a f} - \frac{2 B c^2 (c - c \tan(e + fx) \operatorname{li})^{3/2}}{a f}$$

$$- \frac{2 B c (c - c \tan(e + fx) \operatorname{li})^{5/2}}{5 a f} + \frac{\sqrt{2} A (-c)^{7/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx)} \operatorname{li}}{2 \sqrt{-c}}\right) 10i}{a f}$$

$$- \frac{\sqrt{2} B c^{7/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx)} \operatorname{li}}{2 \sqrt{c}}\right) 18i}{a f} + \frac{A c^4 \sqrt{c - c \tan(e + fx)} \operatorname{li} 4i}{a f (c + c \tan(e + fx) \operatorname{li})}$$

```
input int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(7/2))/(a + a*tan(e + f*x)*1i),x)
```

```
output (4*B*c^4*(c - c*tan(e + f*x)*1i)^(1/2))/(a*f*(c - c*tan(e + f*x)*1i) - 2*a*c*f) + (A*c^3*(c - c*tan(e + f*x)*1i)^(1/2)*8i)/(a*f) + (A*c^2*(c - c*tan(e + f*x)*1i)^(3/2)*2i)/(3*a*f) - (16*B*c^3*(c - c*tan(e + f*x)*1i)^(1/2))/(a*f) - (2*B*c^2*(c - c*tan(e + f*x)*1i)^(3/2))/(a*f) - (2*B*c*(c - c*tan(e + f*x)*1i)^(5/2))/(5*a*f) + (2^(1/2)*A*(-c)^(7/2)*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*(-c)^(1/2)))*10i)/(a*f) - (2^(1/2)*B*c^(7/2)*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2)*1i)/(2*c^(1/2)))*18i)/(a*f) + (A*c^4*(c - c*tan(e + f*x)*1i)^(1/2)*4i)/(a*f*(c + c*tan(e + f*x)*1i))
```

3.765
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{a+ia \tan(e+fx)} dx$$

3.765.1 Optimal result 6949
 3.765.2 Mathematica [A] (verified) 6949
 3.765.3 Rubi [A] (verified) 6950
 3.765.4 Maple [A] (verified) 6952
 3.765.5 Fricas [B] (verification not implemented) 6953
 3.765.6 Sympy [F] 6954
 3.765.7 Maxima [A] (verification not implemented) 6954
 3.765.8 Giac [F] 6955
 3.765.9 Mupad [B] (verification not implemented) 6955

3.765.1 Optimal result

Integrand size = 43, antiderivative size = 180

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{a + ia \tan(e + fx)} dx =$$

$$-\frac{\sqrt{2}(3iA - 7B)c^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{af} + \frac{(3iA - 7B)c^2 \sqrt{c - ic \tan(e + fx)}}{af}$$

$$+ \frac{(3iA - 7B)c(c - ic \tan(e + fx))^{3/2}}{6af} + \frac{(iA - B)(c - ic \tan(e + fx))^{5/2}}{2af(1 + i \tan(e + fx))}$$

output

```
-(3*I*A-7*B)*c^(5/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))
*2^(1/2)/a/f+(3*I*A-7*B)*c^2*(c-I*c*tan(f*x+e))^(1/2)/a/f+1/6*(3*I*A-7*B)*
c*(c-I*c*tan(f*x+e))^(3/2)/a/f+1/2*(I*A-B)*(c-I*c*tan(f*x+e))^(5/2)/a/f/(1
+I*tan(f*x+e))
```

3.765.2 Mathematica [A] (verified)

Time = 6.18 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.74

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{a + ia \tan(e + fx)} dx = \frac{c^2 \left(3\sqrt{2}(-3iA + 7B)\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right) + \frac{2\sqrt{c-ic \tan(e+fx)}}{3af} \right)}{3af}$$

input

```
Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2))/(a + I*a*Tan
[e + f*x]),x]
```

3.765.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{a+ia \tan(e+fx)} dx$$

output $(c^2(3\sqrt{2}((-3i)A + 7B)\sqrt{c}\operatorname{ArcTanh}[\sqrt{c - Ic\tan[e + fx]}/(\sqrt{2}\sqrt{c})]) + (2\sqrt{c - Ic\tan[e + fx]})(6A + (13i)B + (3i)(A + (3i)B)\tan[e + fx] + I B \tan[e + fx]^2))/(-I + \tan[e + fx]))/(3af)$

3.765.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.89, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.186$, Rules used = {3042, 4071, 27, 87, 60, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - ic \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{a + ia \tan(e + fx)} dx$$

↓ 3042

$$\int \frac{(c - ic \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{a + ia \tan(e + fx)} dx$$

↓ 4071

$$\frac{ac \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{a^2(i \tan(e + fx) + 1)^2} d \tan(e + fx)}{f}$$

↓ 27

$$\frac{c \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(i \tan(e + fx) + 1)^2} d \tan(e + fx)}{af}$$

↓ 87

$$\frac{c \left(\frac{(-B + iA)(c - ic \tan(e + fx))^{5/2}}{2c(1 + i \tan(e + fx))} - \frac{1}{4}(3A + 7iB) \int \frac{(c - ic \tan(e + fx))^{3/2}}{i \tan(e + fx) + 1} d \tan(e + fx) \right)}{af}$$

↓ 60

$$\frac{c \left(\frac{(-B + iA)(c - ic \tan(e + fx))^{5/2}}{2c(1 + i \tan(e + fx))} - \frac{1}{4}(3A + 7iB) \left(2c \int \frac{\sqrt{c - ic \tan(e + fx)}}{i \tan(e + fx) + 1} d \tan(e + fx) - \frac{2}{3}i(c - ic \tan(e + fx))^{3/2} \right) \right)}{af}$$

↓ 60

3.765. $\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{a + ia \tan(e + fx)} dx$

$$\frac{c\left(\frac{(-B+iA)(c-ic\tan(e+fx))^{5/2}}{2c(1+i\tan(e+fx))} - \frac{1}{4}(3A+7iB)\right)\left(2c\left(2c\int\frac{1}{(i\tan(e+fx)+1)\sqrt{c-ic\tan(e+fx)}}d\tan(e+fx) - 2i\sqrt{c-ic\tan(e+fx)}\right)\right)}{af}$$

↓ 73

$$\frac{c\left(\frac{(-B+iA)(c-ic\tan(e+fx))^{5/2}}{2c(1+i\tan(e+fx))} - \frac{1}{4}(3A+7iB)\right)\left(2c\left(4i\int\frac{1}{2-\frac{c-ic\tan(e+fx)}{c}}d\sqrt{c-ic\tan(e+fx)} - 2i\sqrt{c-ic\tan(e+fx)}\right)\right)}{af}$$

↓ 219

$$\frac{c\left(\frac{(-B+iA)(c-ic\tan(e+fx))^{5/2}}{2c(1+i\tan(e+fx))} - \frac{1}{4}(3A+7iB)\right)\left(2c\left(2i\sqrt{2}\sqrt{c}\operatorname{arctanh}\left(\frac{\sqrt{c-ic\tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right) - 2i\sqrt{c-ic\tan(e+fx)}\right)\right) - \frac{2}{3}i}{af}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2))/(a + I*a*Tan[e + f*x]),x]`

output `(c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(5/2))/(2*c*(1 + I*Tan[e + f*x])) - ((3*A + (7*I)*B)*(((-2*I)/3)*(c - I*c*Tan[e + f*x])^(3/2) + 2*c*((2*I)*Sqrt[2]*Sqrt[c]*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c]]) - (2*I)*Sqrt[c - I*c*Tan[e + f*x]])))/4))/(a*f)`

3.765.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
 .), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
 + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
 + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
 /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
 rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
 ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
 Q[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
 Q[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
 (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
 mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
 , Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
 + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.765.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.83

method	result
derivativedivides	$\frac{2ic \left(\frac{iB(c-ic \tan(fx+e))}{3} \right)^{\frac{3}{2}} + 3i\sqrt{c-ic \tan(fx+e)} Bc + \sqrt{c-ic \tan(fx+e)} cA - 4c^2 \left(\frac{(-\frac{iB}{8} - \frac{A}{8})\sqrt{c-ic \tan(fx+e)}}{\frac{c}{2} + \frac{ic \tan(fx+e)}{2}} + \frac{(\frac{7iB}{2} + 3)}{\dots} \right)}{fa}$
default	$\frac{2ic \left(\frac{iB(c-ic \tan(fx+e))}{3} \right)^{\frac{3}{2}} + 3i\sqrt{c-ic \tan(fx+e)} Bc + \sqrt{c-ic \tan(fx+e)} cA - 4c^2 \left(\frac{(-\frac{iB}{8} - \frac{A}{8})\sqrt{c-ic \tan(fx+e)}}{\frac{c}{2} + \frac{ic \tan(fx+e)}{2}} + \frac{(\frac{7iB}{2} + 3)}{\dots} \right)}{fa}$

3.765. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{a+ia \tan(e+fx)} dx$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e)),x,method=
_RETURNVERBOSE)
```

```
output 2*I/f/a*c*(1/3*I*B*(c-I*c*tan(f*x+e))^(3/2)+3*I*(c-I*c*tan(f*x+e))^(1/2)*B
*c+(c-I*c*tan(f*x+e))^(1/2)*c*A-4*c^2*((-1/8*I*B-1/8*A)*(c-I*c*tan(f*x+e))
^(1/2)/(1/2*c+1/2*I*c*tan(f*x+e))+1/4*(7/2*I*B+3/2*A)*2^(1/2)/c^(1/2)*arct
anh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2)))
```

3.765.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 406 vs. $2(140) = 280$.

Time = 0.27 (sec) , antiderivative size = 406, normalized size of antiderivative = 2.26

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{a + ia \tan(e + fx)} dx =$$

$$3\sqrt{2}(afe^{4ifx+4ie} + afe^{2ifx+2ie})\sqrt{-\frac{(9A^2+42iAB-49B^2)c^5}{a^2f^2}} \log \left(-\frac{4 \left((3iA-7B)c^3 + (afe^{2ifx+2ie} + af)\sqrt{-\frac{(9A^2+42iAB-49B^2)c^5}{a^2f^2}} \right)}{af} \right)$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e)),x,
algorithm="fricas")
```

```
output -1/6*(3*sqrt(2)*(a*f*e^(4*I*f*x + 4*I*e) + a*f*e^(2*I*f*x + 2*I*e))*sqrt(-
(9*A^2 + 42*I*A*B - 49*B^2)*c^5/(a^2*f^2))*log(-4*((3*I*A - 7*B)*c^3 + (a*
f*e^(2*I*f*x + 2*I*e) + a*f))*sqrt(-(9*A^2 + 42*I*A*B - 49*B^2)*c^5/(a^2*f^
2))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))*e^(-I*f*x - I*e)/(a*f)) - 3*sqrt(2)
*(a*f*e^(4*I*f*x + 4*I*e) + a*f*e^(2*I*f*x + 2*I*e))*sqrt(-(9*A^2 + 42*I*A
*B - 49*B^2)*c^5/(a^2*f^2))*log(-4*((3*I*A - 7*B)*c^3 - (a*f*e^(2*I*f*x +
2*I*e) + a*f))*sqrt(-(9*A^2 + 42*I*A*B - 49*B^2)*c^5/(a^2*f^2))*sqrt(c/(e^(
2*I*f*x + 2*I*e) + 1)))*e^(-I*f*x - I*e)/(a*f)) + 2*sqrt(2)*(3*(-3*I*A + 7
*B)*c^2*e^(4*I*f*x + 4*I*e) + 4*(-3*I*A + 7*B)*c^2*e^(2*I*f*x + 2*I*e) + 3
*(-I*A + B)*c^2)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(a*f*e^(4*I*f*x + 4*I*
e) + a*f*e^(2*I*f*x + 2*I*e))
```

3.765. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{a+ia \tan(e+fx)} dx$

3.765.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{a + i a \tan(e + fx)} dx =$$

$$i \left(\int \frac{A c^2 \sqrt{-i c \tan(e + fx) + c}}{\tan(e + fx) - i} dx + \int \left(-\frac{A c^2 \sqrt{-i c \tan(e + fx) + c} \tan^2(e + fx)}{\tan(e + fx) - i} \right) dx + \int \frac{B c^2 \sqrt{-i c \tan(e + fx) + c} \tan(e + fx)}{\tan(e + fx) - i} dx + \int \right.$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(5/2)/(a+I*a*tan(f*x+e)),x)`

output `-I*(Integral(A*c**2*sqrt(-I*c*tan(e + f*x) + c)/(tan(e + f*x) - I), x) + Integral(-A*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2/(tan(e + f*x) - I), x) + Integral(B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)/(tan(e + f*x) - I), x) + Integral(-B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3/(tan(e + f*x) - I), x) + Integral(-2*I*A*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)/(tan(e + f*x) - I), x) + Integral(-2*I*B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2/(tan(e + f*x) - I), x))/a`

3.765.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.93

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{a + i a \tan(e + fx)} dx = i \left(\frac{3 \sqrt{2} (3A + 7iB) c^{7/2} \log \left(-\frac{\sqrt{2} \sqrt{c} - \sqrt{-i c \tan(fx+e) + c}}{\sqrt{2} \sqrt{c} + \sqrt{-i c \tan(fx+e) + c}} \right)}{a} - \frac{12 \sqrt{-i c \tan(fx+e) + c}}{(-i c \tan(fx+e) + c)} \right)$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output `1/6*I*(3*sqrt(2)*(3*A + 7*I*B)*c^(7/2)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a - 12*sqrt(-I*c*tan(f*x + e) + c)*(A + I*B)*c^4/((-I*c*tan(f*x + e) + c)*a - 2*a*c) + 4*(I*(-I*c*tan(f*x + e) + c)^(3/2)*B*c^2 + 3*sqrt(-I*c*tan(f*x + e) + c)*(A + 3*I*B)*c^3)/a)/(c*f)`

3.765.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{a + i a \tan(e + fx)} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{5/2}}{i a \tan(fx + e) + a} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e)),x,
algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(5/2)/(I*a*tan(f*x
+ e) + a), x)`

3.765.9 Mupad [B] (verification not implemented)

Time = 9.39 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.36

$$\begin{aligned} \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{a + i a \tan(e + fx)} dx &= \frac{2 B c^3 \sqrt{c - c \tan(e + fx)} \operatorname{li}}{a f (c - c \tan(e + fx) \operatorname{li}) - 2 a c f} \\ &+ \frac{A c^2 \sqrt{c - c \tan(e + fx)} \operatorname{li} 2i}{a f} - \frac{6 B c^2 \sqrt{c - c \tan(e + fx)} \operatorname{li}}{a f} \\ &- \frac{2 B c (c - c \tan(e + fx) \operatorname{li})^{3/2}}{3 a f} - \frac{\sqrt{2} A (-c)^{5/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx)} \operatorname{li}}{2 \sqrt{-c}}\right) 3i}{a f} \\ &- \frac{\sqrt{2} B c^{5/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx)} \operatorname{li} \operatorname{li}}{2 \sqrt{c}}\right) 7i}{a f} + \frac{A c^3 \sqrt{c - c \tan(e + fx)} \operatorname{li} 2i}{a f (c + c \tan(e + fx) \operatorname{li})} \end{aligned}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(5/2))/(a + a*tan(e + f*
x)*1i),x)`

output `(2*B*c^3*(c - c*tan(e + f*x)*1i)^(1/2))/(a*f*(c - c*tan(e + f*x)*1i) - 2*a
*c*f) + (A*c^2*(c - c*tan(e + f*x)*1i)^(1/2)*2i)/(a*f) - (6*B*c^2*(c - c*t
an(e + f*x)*1i)^(1/2))/(a*f) - (2*B*c*(c - c*tan(e + f*x)*1i)^(3/2))/(3*a*
f) - (2^(1/2)*A*(-c)^(5/2)*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2
*(-c)^(1/2)))*3i)/(a*f) - (2^(1/2)*B*c^(5/2)*atan((2^(1/2)*(c - c*tan(e +
f*x)*1i)^(1/2)*1i)/(2*c^(1/2)))*7i)/(a*f) + (A*c^3*(c - c*tan(e + f*x)*1i)
^(1/2)*2i)/(a*f*(c + c*tan(e + f*x)*1i))`

3.766
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{a+ia \tan(e+fx)} dx$$

3.766.1 Optimal result	6956
3.766.2 Mathematica [A] (verified)	6956
3.766.3 Rubi [A] (verified)	6957
3.766.4 Maple [A] (verified)	6959
3.766.5 Fricas [B] (verification not implemented)	6960
3.766.6 Sympy [F]	6961
3.766.7 Maxima [A] (verification not implemented)	6961
3.766.8 Giac [F]	6962
3.766.9 Mupad [B] (verification not implemented)	6962

3.766.1 Optimal result

Integrand size = 43, antiderivative size = 144

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{a + ia \tan(e + fx)} dx =$$

$$\frac{(iA - 5B)c^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{\sqrt{2}af} + \frac{(iA - 5B)c\sqrt{c - ic \tan(e + fx)}}{2af} + \frac{(iA - B)(c - ic \tan(e + fx))^{3/2}}{2af(1 + i \tan(e + fx))}$$

output

```
-1/2*(I*A-5*B)*c^(3/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2)))/a/f*2^(1/2)+1/2*(I*A-5*B)*c*(c-I*c*tan(f*x+e))^(1/2)/a/f+1/2*(I*A-B)*(c-I*c*tan(f*x+e))^(3/2)/a/f/(1+I*tan(f*x+e))
```

3.766.2 Mathematica [A] (verified)

Time = 3.90 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.75

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{a + ia \tan(e + fx)} dx = \frac{c\left(\sqrt{2}(-iA + 5B)\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right) + \frac{2(A+3iB)}{2af}\right)}{2af}$$

input

```
Integrate[(((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2))/(a + I*a*Tan[e + f*x])),x]
```

3.766.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{a+ia \tan(e+fx)} dx$$

output $(c*(\text{Sqrt}[2]*((-I)*A + 5*B)*\text{Sqrt}[c]*\text{ArcTanh}[\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]]]/(\text{Sqrt}[2]*\text{Sqrt}[c])) + (2*(A + (3*I)*B - 2*B*\text{Tan}[e + f*x])*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])/(-I + \text{Tan}[e + f*x]))/(2*a*f)$

3.766.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.92, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.163$, Rules used = {3042, 4071, 27, 87, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{a + ia \tan(e + fx)} dx$$

↓ 3042

$$\int \frac{(c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{a + ia \tan(e + fx)} dx$$

↓ 4071

$$\frac{ac \int \frac{(A+B \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}}{a^2(i \tan(e+fx)+1)^2} d \tan(e + fx)}{f}$$

↓ 27

$$\frac{c \int \frac{(A+B \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}}{(i \tan(e+fx)+1)^2} d \tan(e + fx)}{af}$$

↓ 87

$$\frac{c \left(\frac{(-B+IA)(c-ic \tan(e+fx))^{3/2}}{2c(1+i \tan(e+fx))} - \frac{1}{4}(A + 5iB) \int \frac{\sqrt{c-ic \tan(e+fx)}}{i \tan(e+fx)+1} d \tan(e + fx) \right)}{af}$$

↓ 60

$$\frac{c \left(\frac{(-B+IA)(c-ic \tan(e+fx))^{3/2}}{2c(1+i \tan(e+fx))} - \frac{1}{4}(A + 5iB) \left(2c \int \frac{1}{(i \tan(e+fx)+1) \sqrt{c-ic \tan(e+fx)}} d \tan(e + fx) - 2i \sqrt{c - ic \tan(e + fx)} \right) \right)}{af}$$

↓ 73

3.766. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{a+ia \tan(e+fx)} dx$

$$\frac{c \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{3/2}}{2c(1+i \tan(e+fx))} - \frac{1}{4}(A+5iB) \left(4i \int \frac{1}{2-\frac{c-ic \tan(e+fx)}{c}} d\sqrt{c-ic \tan(e+fx)} - 2i\sqrt{c-ic \tan(e+fx)} \right) \right)}{af}$$

↓ 219

$$\frac{c \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{3/2}}{2c(1+i \tan(e+fx))} - \frac{1}{4}(A+5iB) \left(2i\sqrt{2}\sqrt{c} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}} \right) - 2i\sqrt{c-ic \tan(e+fx)} \right) \right)}{af}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2))/(a + I*a*Tan[e + f*x]),x]`

output `(c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(3/2))/(2*c*(1 + I*Tan[e + f*x])) - ((A + (5*I)*B)*((2*I)*Sqrt[2]*Sqrt[c]*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])] - (2*I)*Sqrt[c - I*c*Tan[e + f*x]]))/4))/(a*f)`

3.766.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(- (b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.766.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.77

method	result	size
derivativedivides	$\frac{2ic \left(i\sqrt{c-ic \tan(fx+e)} B + c \left(\frac{\left(\frac{iB}{4} + \frac{A}{4}\right) \sqrt{c-ic \tan(fx+e)}}{\frac{c}{2} + \frac{ic \tan(fx+e)}{2}} - \frac{\left(\frac{A}{2} + \frac{5iB}{2}\right) \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{2\sqrt{c}} \right)}{fa} \right)}{fa}$	111
default	$\frac{2ic \left(i\sqrt{c-ic \tan(fx+e)} B + c \left(\frac{\left(\frac{iB}{4} + \frac{A}{4}\right) \sqrt{c-ic \tan(fx+e)}}{\frac{c}{2} + \frac{ic \tan(fx+e)}{2}} - \frac{\left(\frac{A}{2} + \frac{5iB}{2}\right) \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{2\sqrt{c}} \right)}{fa} \right)}{fa}$	111

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e)),x,method=
_RETURNVERBOSE)
```

$$3.766. \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{a+ia \tan(e+fx)} dx$$

```
output 2*I/f/a*c*(I*(c-I*c*tan(f*x+e))^(1/2)*B+c*((1/4*I*B+1/4*A)*(c-I*c*tan(f*x+
e))^(1/2)/(1/2*c+1/2*I*c*tan(f*x+e))-1/2*(1/2*A+5/2*I*B)*2^(1/2)/c^(1/2)*a
rctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))))
```

3.766.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 330 vs. $2(111) = 222$.

Time = 0.25 (sec) , antiderivative size = 330, normalized size of antiderivative = 2.29

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{a + ia \tan(e + fx)} dx =$$

$$\left(\sqrt{2}af \sqrt{-\frac{(A^2 + 10iAB - 25B^2)c^3}{a^2f^2}} e^{(2ifx + 2ie)} \log \left(-\frac{2 \left((iA - 5B)c^2 + (afe^{(2ifx + 2ie)} + af) \sqrt{-\frac{(A^2 + 10iAB - 25B^2)c^3}{a^2f^2}} \sqrt{\frac{c}{e^{(2ifx + 2ie)}}}} \right)}{af} \right) \right)$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e)),x,
algorithm="fricas")
```

```
output -1/4*(sqrt(2)*a*f*sqrt(-(A^2 + 10*I*A*B - 25*B^2)*c^3/(a^2*f^2))*e^(2*I*f*
x + 2*I*e)*log(-2*((I*A - 5*B)*c^2 + (a*f*e^(2*I*f*x + 2*I*e) + a*f)*sqrt(
-(A^2 + 10*I*A*B - 25*B^2)*c^3/(a^2*f^2))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1
)))*e^(-I*f*x - I*e)/(a*f)) - sqrt(2)*a*f*sqrt(-(A^2 + 10*I*A*B - 25*B^2)*c
^3/(a^2*f^2))*e^(2*I*f*x + 2*I*e)*log(-2*((I*A - 5*B)*c^2 - (a*f*e^(2*I*f*
x + 2*I*e) + a*f)*sqrt(-(A^2 + 10*I*A*B - 25*B^2)*c^3/(a^2*f^2))*sqrt(c/(e
^(2*I*f*x + 2*I*e) + 1)))*e^(-I*f*x - I*e)/(a*f)) + 2*sqrt(2)*((-I*A + 5*B
)*c*e^(2*I*f*x + 2*I*e) + (-I*A + B)*c)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))
*e^(-2*I*f*x - 2*I*e)/(a*f)
```

3.766.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{a + i a \tan(e + fx)} dx =$$

$$\frac{i \left(\int \frac{A c \sqrt{-i c \tan(e + fx) + c}}{\tan(e + fx) - i} dx + \int \frac{B c \sqrt{-i c \tan(e + fx) + c} \tan(e + fx)}{\tan(e + fx) - i} dx + \int \left(-\frac{i A c \sqrt{-i c \tan(e + fx) + c} \tan(e + fx)}{\tan(e + fx) - i} \right) dx + \int \left(-\frac{4 \sqrt{-i c \tan(e + fx) + c}}{(-i c \tan(e + fx) + c)} \right) dx \right)}{a}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(3/2)/(a+I*a*tan(f*x+e)),x)`

output `-I*(Integral(A*c*sqrt(-I*c*tan(e + f*x) + c)/(tan(e + f*x) - I), x) + Integral(B*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)/(tan(e + f*x) - I), x) + Integral(-I*A*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)/(tan(e + f*x) - I), x) + Integral(-I*B*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2/(tan(e + f*x) - I), x))/a`

3.766.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.96

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{a + i a \tan(e + fx)} dx = \frac{i \left(\frac{\sqrt{2}(A+5iB)c^{5/2} \log\left(-\frac{\sqrt{2}\sqrt{c}-\sqrt{-i c \tan(fx+e)+c}}{\sqrt{2}\sqrt{c}+\sqrt{-i c \tan(fx+e)+c}}\right)}{a} - \frac{4 \sqrt{-i c \tan(fx+e)+c}}{(-i c \tan(fx+e)+c)} \right)}{4 c f}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output `1/4*I*(sqrt(2)*(A + 5*I*B)*c^(5/2)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a - 4*sqrt(-I*c*tan(f*x + e) + c)*(A + I*B)*c^3/((-I*c*tan(f*x + e) + c)*a - 2*a*c) + 8*I*sqrt(-I*c*tan(f*x + e) + c)*B*c^2/a)/(c*f)`

3.766.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{a + i a \tan(e + fx)} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{3/2}}{i a \tan(fx + e) + a} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e)),x,
algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(3/2)/(I*a*tan(f*x
+ e) + a), x)`

3.766.9 Mupad [B] (verification not implemented)

Time = 9.20 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.31

$$\begin{aligned} \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{a + i a \tan(e + fx)} dx &= \frac{B c^2 \sqrt{c - c \tan(e + fx)} \operatorname{li}}{a f (c - c \tan(e + fx) \operatorname{li}) - 2 a c f} \\ &- \frac{2 B c \sqrt{c - c \tan(e + fx)} \operatorname{li}}{a f} + \frac{\sqrt{2} A (-c)^{3/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx)} \operatorname{li}}{2 \sqrt{-c}}\right) \operatorname{li}}{2 a f} \\ &+ \frac{5 \sqrt{2} B c^{3/2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx)} \operatorname{li}}{2 \sqrt{-c}}\right)}{2 a f} + \frac{A c^2 \sqrt{c - c \tan(e + fx)} \operatorname{li} \operatorname{li}}{a f (c + c \tan(e + fx) \operatorname{li})} \end{aligned}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(3/2))/(a + a*tan(e + f*
x)*1i),x)`

output `(B*c^2*(c - c*tan(e + f*x)*1i)^(1/2))/(a*f*(c - c*tan(e + f*x)*1i) - 2*a*c
*f) - (2*B*c*(c - c*tan(e + f*x)*1i)^(1/2))/(a*f) + (2^(1/2)*A*(-c)^(3/2)*
atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*(-c)^(1/2)))*1i)/(2*a*f) +
(5*2^(1/2)*B*c^(3/2)*atanh((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*c^(
1/2))))/(2*a*f) + (A*c^2*(c - c*tan(e + f*x)*1i)^(1/2)*1i)/(a*f*(c + c*tan
(e + f*x)*1i))`

$$3.767 \quad \int \frac{(A+B \tan(e+fx))\sqrt{c-ictan(e+fx)}}{a+ia \tan(e+fx)} dx$$

3.767.1 Optimal result	6963
3.767.2 Mathematica [A] (verified)	6963
3.767.3 Rubi [A] (verified)	6964
3.767.4 Maple [A] (verified)	6966
3.767.5 Fricas [B] (verification not implemented)	6966
3.767.6 Sympy [F]	6967
3.767.7 Maxima [A] (verification not implemented)	6967
3.767.8 Giac [F]	6968
3.767.9 Mupad [B] (verification not implemented)	6968

3.767.1 Optimal result

Integrand size = 43, antiderivative size = 109

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)}}{a + ia \tan(e + fx)} dx$$

$$= \frac{(iA + 3B)\sqrt{c}\operatorname{arctanh}\left(\frac{\sqrt{c-ictan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{2\sqrt{2}af} + \frac{(iA - B)\sqrt{c - ictan(e + fx)}}{2af(1 + i \tan(e + fx))}$$

output `1/4*(I*A+3*B)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))*c^(1/2)/a/f*2^(1/2)+1/2*(I*A-B)*(c-I*c*tan(f*x+e))^(1/2)/a/f/(1+I*tan(f*x+e))`

3.767.2 Mathematica [A] (verified)

Time = 2.89 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.90

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)}}{a + ia \tan(e + fx)} dx$$

$$= \frac{\sqrt{2}(iA + 3B)\sqrt{c}\operatorname{arctanh}\left(\frac{\sqrt{c-ictan(e+fx)}}{\sqrt{2}\sqrt{c}}\right) + \frac{2(A+iB)\sqrt{c-ictan(e+fx)}}{-i+\tan(e+fx)}}{4af}$$

input `Integrate[((A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])/(a + I*a*Tan[e + f*x]),x]`

output $(\text{Sqrt}[2]*(I*A + 3*B)*\text{Sqrt}[c]*\text{ArcTanh}[\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]]]/(\text{Sqrt}[2]*\text{Sqrt}[c])) + (2*(A + I*B)*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])/(-I + \text{Tan}[e + f*x]))/(4*a*f)$

3.767.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.99, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.140$, Rules used = {3042, 4071, 27, 87, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{c - ic \tan(e + fx)}(A + B \tan(e + fx))}{a + ia \tan(e + fx)} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{\sqrt{c - ic \tan(e + fx)}(A + B \tan(e + fx))}{a + ia \tan(e + fx)} dx \\
 & \quad \downarrow 4071 \\
 & \frac{ac \int \frac{A + B \tan(e + fx)}{a^2(i \tan(e + fx) + 1)^2 \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{f} \\
 & \quad \downarrow 27 \\
 & \frac{c \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx) + 1)^2 \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{af} \\
 & \quad \downarrow 87 \\
 & \frac{c \left(\frac{1}{4}(A - 3iB) \int \frac{1}{(i \tan(e + fx) + 1) \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) + \frac{(-B + iA) \sqrt{c - ic \tan(e + fx)}}{2c(1 + i \tan(e + fx))} \right)}{af} \\
 & \quad \downarrow 73 \\
 & \frac{c \left(\frac{i(A - 3iB) \int \frac{1}{2 - \frac{c - ic \tan(e + fx)}{c}} d \sqrt{c - ic \tan(e + fx)}}{2c} + \frac{(-B + iA) \sqrt{c - ic \tan(e + fx)}}{2c(1 + i \tan(e + fx))} \right)}{af} \\
 & \quad \downarrow 219
 \end{aligned}$$

3.767. $\int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{a + ia \tan(e + fx)} dx$

$$\frac{c \left(\frac{i(A-3iB)\operatorname{arctanh}\left(\frac{\sqrt{c-ic\tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{2\sqrt{2}\sqrt{c}} + \frac{(-B+iA)\sqrt{c-ic\tan(e+fx)}}{2c(1+i\tan(e+fx))} \right)}{af}$$

input `Int[((A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]/(a + I*a*Tan[e + f*x])),x]`

output `(c*((I/2)*(A - (3*I)*B)*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])])/(Sqrt[2]*Sqrt[c]) + ((I*A - B)*Sqrt[c - I*c*Tan[e + f*x]]/(2*c*(1 + I*Tan[e + f*x]))) / (a*f)`

3.767.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.767. $\int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{a+ia \tan(e+fx)} dx$

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.767.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$\frac{2ic \left(\frac{\left(\frac{iB}{8} + \frac{A}{8}\right) \sqrt{c - ic \tan(fx+e)}}{\frac{c}{2} + \frac{ic \tan(fx+e)}{2}} + \frac{\left(-\frac{3iB}{2} + \frac{A}{2}\right) \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{4\sqrt{c}} \right)}{fa}$	90
default	$\frac{2ic \left(\frac{\left(\frac{iB}{8} + \frac{A}{8}\right) \sqrt{c - ic \tan(fx+e)}}{\frac{c}{2} + \frac{ic \tan(fx+e)}{2}} + \frac{\left(-\frac{3iB}{2} + \frac{A}{2}\right) \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{4\sqrt{c}} \right)}{fa}$	90

```
input int((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e)),x,method=
_RETURNVERBOSE)
```

```
output 2*I/f/a*c*((1/8*I*B+1/8*A)*(c-I*c*tan(f*x+e))^(1/2)/(1/2*c+1/2*I*c*tan(f*x
+e))+1/4*(-3/2*I*B+1/2*A)*2^(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(
1/2)*2^(1/2)/c^(1/2)))
```

3.767.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 326 vs. 2(82) = 164.

Time = 0.26 (sec) , antiderivative size = 326, normalized size of antiderivative = 2.99

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{a + ia \tan(e + fx)} dx$$

$$= \left(\sqrt{\frac{1}{2}} af \sqrt{-\frac{(A^2 - 6iAB - 9B^2)c}{a^2 f^2}} e^{(2i fx + 2ie)} \log \left(\frac{\left(\sqrt{2} \sqrt{\frac{1}{2}} (af e^{(2i fx + 2ie)} + af) \sqrt{\frac{c}{e^{(2i fx + 2ie)} + 1}} \sqrt{-\frac{(A^2 - 6iAB - 9B^2)c}{a^2 f^2}} + (iA + 3B)c \right)}{af} \right) \right)$$

3.767. $\int \frac{(A+B \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}}{a+ia \tan(e+fx)} dx$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e)),x,
algorithm="fricas")`

output `1/4*(sqrt(1/2)*a*f*sqrt(-(A^2 - 6*I*A*B - 9*B^2)*c/(a^2*f^2))*e^(2*I*f*x +
2*I*e)*log((sqrt(2)*sqrt(1/2)*(a*f*e^(2*I*f*x + 2*I*e) + a*f)*sqrt(c/(e^(
2*I*f*x + 2*I*e) + 1))*sqrt(-(A^2 - 6*I*A*B - 9*B^2)*c/(a^2*f^2)) + (I*A +
3*B)*c)*e^(-I*f*x - I*e)/(a*f)) - sqrt(1/2)*a*f*sqrt(-(A^2 - 6*I*A*B - 9*
B^2)*c/(a^2*f^2))*e^(2*I*f*x + 2*I*e)*log(-(sqrt(2)*sqrt(1/2)*(a*f*e^(2*I*
f*x + 2*I*e) + a*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(A^2 - 6*I*A*B
- 9*B^2)*c/(a^2*f^2)) - (I*A + 3*B)*c)*e^(-I*f*x - I*e)/(a*f)) + sqrt(2)*
((I*A - B)*e^(2*I*f*x + 2*I*e) + I*A - B)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)`

3.767.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{a + i a \tan(e + fx)} dx$$

$$= - \frac{i \left(\int \frac{A \sqrt{-i c \tan(e + fx) + c}}{\tan(e + fx) - i} dx + \int \frac{B \sqrt{-i c \tan(e + fx) + c} \tan(e + fx)}{\tan(e + fx) - i} dx \right)}{a}$$

input `integrate((c-I*c*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e)),x)`

output `-I*(Integral(A*sqrt(-I*c*tan(e + f*x) + c)/(tan(e + f*x) - I), x) + Integr
al(B*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)/(tan(e + f*x) - I), x))/a`

3.767.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.06

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{a + i a \tan(e + fx)} dx$$

$$= - \frac{i \left(\frac{\sqrt{2}(A - 3iB)c^{\frac{3}{2}} \log\left(-\frac{\sqrt{2}\sqrt{c} - \sqrt{-i c \tan(fx+e)+c}}{\sqrt{2}\sqrt{c} + \sqrt{-i c \tan(fx+e)+c}}\right)}{a} + \frac{4 \sqrt{-i c \tan(fx+e)+c}(A + iB)c^2}{(-i c \tan(fx+e)+c)a - 2ac} \right)}{8cf}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e)),x,
algorithm="maxima")`

output `-1/8*I*(sqrt(2)*(A - 3*I*B)*c^(3/2)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a + 4*sqrt(-I*c*tan(f*x + e) + c)*(A + I*B)*c^2/((-I*c*tan(f*x + e) + c)*a - 2*a*c))/(c*f)`

3.767.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{a + i a \tan(e + fx)} dx$$

$$= \int \frac{(B \tan(fx + e) + A) \sqrt{-i c \tan(fx + e) + c}}{i a \tan(fx + e) + a} dx$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e)),x,
algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*sqrt(-I*c*tan(f*x + e) + c)/(I*a*tan(f*x + e) + a), x)`

3.767.9 Mupad [B] (verification not implemented)

Time = 8.94 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.46

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{a + i a \tan(e + fx)} dx$$

$$= -\frac{B c \sqrt{c - c \tan(e + f x)} \operatorname{li}}{2 (a c f + a c f \tan(e + f x) \operatorname{li})} + \frac{A c \sqrt{c - c \tan(e + f x)} \operatorname{li}}{2 a f (c + c \tan(e + f x) \operatorname{li})}$$

$$+ \frac{\sqrt{2} A \sqrt{-c} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + f x)} \operatorname{li}}{2 \sqrt{-c}}\right) \operatorname{li}}{4 a f} + \frac{3 \sqrt{2} B \sqrt{c} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + f x)} \operatorname{li}}{2 \sqrt{c}}\right)}{4 a f}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(1/2))/(a + a*tan(e + f*x)*1i),x)`

output $(A*c*(c - c*\tan(e + f*x)*1i)^{(1/2)*1i}/(2*a*f*(c + c*\tan(e + f*x)*1i)) - (B*c*(c - c*\tan(e + f*x)*1i)^{(1/2)})/(2*(a*c*f + a*c*f*\tan(e + f*x)*1i)) + (2^{(1/2)*A*(-c)^{(1/2)*\operatorname{atan}((2^{(1/2)*(c - c*\tan(e + f*x)*1i)^{(1/2)})/(2*(-c)^{(1/2)})})*1i)/(4*a*f)} + (3*2^{(1/2)*B*c^{(1/2)*\operatorname{atanh}((2^{(1/2)*(c - c*\tan(e + f*x)*1i)^{(1/2)})/(2*c^{(1/2)})})})/(4*a*f))$

3.768
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))\sqrt{c-ic \tan(e+fx)}} dx$$

3.768.1 Optimal result	6970
3.768.2 Mathematica [A] (verified)	6970
3.768.3 Rubi [A] (verified)	6971
3.768.4 Maple [A] (verified)	6973
3.768.5 Fracas [B] (verification not implemented)	6974
3.768.6 Sympy [F]	6975
3.768.7 Maxima [A] (verification not implemented)	6975
3.768.8 Giac [F]	6976
3.768.9 Mupad [B] (verification not implemented)	6976

3.768.1 Optimal result

Integrand size = 43, antiderivative size = 141

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))\sqrt{c - ic \tan(e + fx)}} dx$$

$$= \frac{(3iA + B)\operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{4\sqrt{2}a\sqrt{cf}} - \frac{3iA + B}{4af\sqrt{c - ic \tan(e + fx)}} + \frac{iA - B}{2af(1 + i \tan(e + fx))\sqrt{c - ic \tan(e + fx)}}$$

```
output 1/8*(3*I*A+B)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a/f*2^(1/2)/c^(1/2)+1/4*(-3*I*A-B)/a/f/(c-I*c*tan(f*x+e))^(1/2)+1/2*(I*A-B)/a/f/(c-I*c*tan(f*x+e))^(1/2)/(1+I*tan(f*x+e))
```

3.768.2 Mathematica [A] (verified)

Time = 3.91 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.80

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))\sqrt{c - ic \tan(e + fx)}} dx$$

$$= \frac{\sqrt{2}(3iA+B)\operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{\sqrt{c}} + \frac{-2A+6iB+(-6iA-2B)\tan(e+fx)}{8af(-i+\tan(e+fx))\sqrt{c-ic \tan(e+fx)}}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]),x]`

output `((Sqrt[2]*((3*I)*A + B)*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])))/Sqrt[c] + (-2*A + (6*I)*B + ((-6*I)*A - 2*B)*Tan[e + f*x])/((-I + Tan [e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])/(8*a*f)`

3.768.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.97, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.163$, Rules used = {3042, 4071, 27, 87, 61, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A + B \tan(e + fx)}{a^2 (i \tan(e + fx) + 1)^2 (c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{c \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx) + 1)^2 (c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{af} \\
 & \quad \downarrow \text{87} \\
 & \frac{c \left(\frac{1}{4} (3A - iB) \int \frac{1}{(i \tan(e + fx) + 1) (c - ic \tan(e + fx))^{3/2}} d \tan(e + fx) + \frac{-B + iA}{2c(1 + i \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}} \right)}{af} \\
 & \quad \downarrow \text{61} \\
 & \frac{c \left(\frac{1}{4} (3A - iB) \left(\frac{\int \frac{1}{(i \tan(e + fx) + 1) \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{2c} - \frac{i}{c \sqrt{c - ic \tan(e + fx)}} \right) + \frac{-B + iA}{2c(1 + i \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}} \right)}{af}
 \end{aligned}$$

3.768. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}} dx$

$$\begin{aligned}
 & \downarrow 73 \\
 & \frac{c \left(\frac{1}{4}(3A - iB) \left(\frac{i \int \frac{1}{2 - \frac{c - ic \tan(e+fx)}{c}} d\sqrt{c - ic \tan(e+fx)}}{c^2} - \frac{i}{c\sqrt{c - ic \tan(e+fx)}} \right) + \frac{-B + iA}{2c(1 + i \tan(e+fx))\sqrt{c - ic \tan(e+fx)}} \right)}{af} \\
 & \downarrow 219 \\
 & \frac{c \left(\frac{1}{4}(3A - iB) \left(\frac{i \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{\sqrt{2}c^{3/2}} - \frac{i}{c\sqrt{c - ic \tan(e+fx)}} \right) + \frac{-B + iA}{2c(1 + i \tan(e+fx))\sqrt{c - ic \tan(e+fx)}} \right)}{af}
 \end{aligned}$$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]),x]`

output `(c*((I*A - B)/(2*c*(1 + I*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]) + ((3*A - I*B)*((I*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])])/(Sqrt[2]*c^(3/2)) - I/(c*Sqrt[c - I*c*Tan[e + f*x]])))/4)/(a*f)`

3.768.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(- (b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.768.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.86

method	result	size
derivativedivides	$2ic \left(-\frac{-iB+A}{4c\sqrt{c-ic\tan(fx+e)}} + \frac{\frac{iB}{4} + \frac{A}{4}}{\frac{c}{2} + \frac{ic\tan(fx+e)}{2}} + \frac{\left(\frac{3A}{2} - \frac{iB}{2}\right)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic\tan(fx+e)}\sqrt{2}}{2\sqrt{c}}\right)}{4c} \right)$	121
default	$2ic \left(-\frac{-iB+A}{4c\sqrt{c-ic\tan(fx+e)}} + \frac{\frac{iB}{4} + \frac{A}{4}}{\frac{c}{2} + \frac{ic\tan(fx+e)}{2}} + \frac{\left(\frac{3A}{2} - \frac{iB}{2}\right)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic\tan(fx+e)}\sqrt{2}}{2\sqrt{c}}\right)}{4c} \right)$	121

```
input int((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e)),x,method=
_RETURNVERBOSE)
```

$$3.768. \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))\sqrt{c-ic \tan(e+fx)}} dx$$

```
output 2*I/f/a*c*(-1/4/c*(A-I*B)/(c-I*c*tan(f*x+e))^(1/2)+1/4/c*((1/4*I*B+1/4*A)*
(c-I*c*tan(f*x+e))^(1/2)/(1/2*c+1/2*I*c*tan(f*x+e))+1/2*(3/2*A-1/2*I*B)*2^
(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))))
```

3.768.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 355 vs. 2(108) = 216.

Time = 0.26 (sec) , antiderivative size = 355, normalized size of antiderivative = 2.52

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}} dx$$

$$\left(\sqrt{\frac{1}{2}} acf \sqrt{-\frac{9A^2 - 6iAB - B^2}{a^2cf^2}} e^{(2ifx + 2ie)} \log \left(\frac{(\sqrt{2} \sqrt{\frac{1}{2}} (afe^{(2ifx + 2ie)} + af) \sqrt{\frac{c}{e^{(2ifx + 2ie)} + 1}} \sqrt{-\frac{9A^2 - 6iAB - B^2}{a^2cf^2}} + 3iA + B)) e^{(-ifx)}}{2af} \right) \right)$$

```
input integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e)),x,
algorithm="fricas")
```

```
output 1/8*(sqrt(1/2)*a*c*f*sqrt(-(9*A^2 - 6*I*A*B - B^2)/(a^2*c*f^2))*e^(2*I*f*x
+ 2*I*e)*log(1/2*(sqrt(2)*sqrt(1/2)*(a*f*e^(2*I*f*x + 2*I*e) + a*f)*sqrt(
c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(9*A^2 - 6*I*A*B - B^2)/(a^2*c*f^2)) +
3*I*A + B)*e^(-I*f*x - I*e)/(a*f)) - sqrt(1/2)*a*c*f*sqrt(-(9*A^2 - 6*I*A*
B - B^2)/(a^2*c*f^2))*e^(2*I*f*x + 2*I*e)*log(-1/2*(sqrt(2)*sqrt(1/2)*(a*f
*e^(2*I*f*x + 2*I*e) + a*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(9*A^2
- 6*I*A*B - B^2)/(a^2*c*f^2)) - 3*I*A - B)*e^(-I*f*x - I*e)/(a*f)) - sqrt
(2)*(2*(I*A + B)*e^(4*I*f*x + 4*I*e) - (-I*A - 3*B)*e^(2*I*f*x + 2*I*e) -
I*A + B)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))e^(-2*I*f*x - 2*I*e)/(a*c*f)
```

3.768.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}} dx =$$

$$\frac{i \left(\int \frac{A}{\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) - i \sqrt{-ic \tan(e + fx) + c}} dx + \int \frac{B \tan(e + fx)}{\sqrt{-ic \tan(e + fx) + c} \tan(e + fx) - i \sqrt{-ic \tan(e + fx) + c}} dx \right)}{a}$$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(1/2)/(a+I*a*tan(f*x+e)),x)`

output `-I*(Integral(A/(sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x) - I*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(B*tan(e + f*x)/(sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) - I*sqrt(-I*c*tan(e + f*x) + c)), x))/a`

3.768.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.01

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}} dx$$

$$= \frac{i \left(\frac{\sqrt{2}(3A - iB)\sqrt{c} \log\left(-\frac{\sqrt{2}\sqrt{c} - \sqrt{-ic \tan(fx+e)+c}}{\sqrt{2}\sqrt{c} + \sqrt{-ic \tan(fx+e)+c}}\right)}{a} + \frac{4((-ic \tan(fx+e)+c)(3A - iB)c - 4(A - iB)c^2)}{(-ic \tan(fx+e)+c)^{\frac{3}{2}} a - 2\sqrt{-ic \tan(fx+e)+c}ac} \right)}{16cf}$$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

output `-1/16*I*(sqrt(2)*(3*A - I*B)*sqrt(c)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a + 4*((-I*c*tan(f*x + e) + c)*(3*A - I*B)*c - 4*(A - I*B)*c^2)/((-I*c*tan(f*x + e) + c)^(3/2)*a - 2*sqrt(-I*c*tan(f*x + e) + c)*a*c)/(c*f)`

3.768.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}} dx$$

$$= \int \frac{B \tan(fx + e) + A}{(ia \tan(fx + e) + a) \sqrt{-ic \tan(fx + e) + c}} dx$$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e)),x,
algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)*sqrt(-I*c*tan(f*x +
e) + c)), x)`

3.768.9 Mupad [B] (verification not implemented)

Time = 9.29 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.50

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}} dx$$

$$= \frac{Bc - \frac{B(c - c \tan(e + fx) li)}{4}}{af(c - c \tan(e + fx) li)^{3/2} - 2acf \sqrt{c - c \tan(e + fx) li}}$$

$$+ \frac{\frac{A(c - c \tan(e + fx) li) 3i}{4af} - \frac{Ac li}{af}}{2c \sqrt{c - c \tan(e + fx) li} - (c - c \tan(e + fx) li)^{3/2}}$$

$$- \frac{\sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) li}}{2\sqrt{-c}}\right) 3i}{8a \sqrt{-c} f} + \frac{\sqrt{2} B \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) li}}{2\sqrt{c}}\right)}{8a \sqrt{c} f}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)*(c - c*tan(e + f*x)*1i)^(1/2)),x)`

output `(B*c - (B*(c - c*tan(e + f*x)*1i))/4)/(a*f*(c - c*tan(e + f*x)*1i)^(3/2) -
2*a*c*f*(c - c*tan(e + f*x)*1i)^(1/2)) + ((A*(c - c*tan(e + f*x)*1i)*3i)/
(4*a*f) - (A*c*1i)/(a*f))/(2*c*(c - c*tan(e + f*x)*1i)^(1/2) - (c - c*tan(
e + f*x)*1i)^(3/2)) - (2^(1/2)*A*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/
2)))/(2*(-c)^(1/2)))*3i)/(8*a*(-c)^(1/2)*f) + (2^(1/2)*B*atanh((2^(1/2)*(c
- c*tan(e + f*x)*1i)^(1/2))/(2*c^(1/2)))/(8*a*c^(1/2)*f)`

3.768. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}} dx$

3.769
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} dx$$

3.769.1 Optimal result	6977
3.769.2 Mathematica [C] (verified)	6977
3.769.3 Rubi [A] (verified)	6978
3.769.4 Maple [A] (verified)	6981
3.769.5 Fricas [B] (verification not implemented)	6981
3.769.6 Sympy [F]	6982
3.769.7 Maxima [A] (verification not implemented)	6982
3.769.8 Giac [F]	6983
3.769.9 Mupad [B] (verification not implemented)	6983

3.769.1 Optimal result

Integrand size = 43, antiderivative size = 184

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^{3/2}} dx = \frac{(5iA - B) \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{8\sqrt{2}ac^{3/2}f} - \frac{5iA - B}{12af(c - ic \tan(e + fx))^{3/2}} + \frac{iA - B}{2af(1 + i \tan(e + fx))(c - ic \tan(e + fx))^{3/2}} - \frac{5iA - B}{8acf\sqrt{c - ic \tan(e + fx)}}$$

```
output 1/16*(5*I*A-B)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a/c^(3/2)/f*2^(1/2)+1/8*(-5*I*A+B)/a/c/f/(c-I*c*tan(f*x+e))^(1/2)+1/12*(-5*I*A+B)/a/f/(c-I*c*tan(f*x+e))^(3/2)+1/2*(I*A-B)/a/f/(1+I*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2)
```

3.769.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 4.71 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.56

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^{3/2}} dx = \frac{3(-5iA + B) \operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -\frac{1}{2}i(i + \tan(e + fx))\right)}{\dots}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2)),x]`

output `(3*((-5*I)*A + B)*Hypergeometric2F1[-1/2, 1, 1/2, (-1/2*I)*(I + Tan[e + f*x])] + (2*I)*Cos[e + f*x]*((A + (5*I)*B)*Cos[e + f*x] + ((-5*I)*A + B)*Sin[e + f*x]))/(24*a*c*f*Sqrt[c - I*c*Tan[e + f*x]])`

3.769.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.93, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.186$, Rules used = {3042, 4071, 27, 87, 61, 61, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ictan(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ictan(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A+B \tan(e+fx)}{a^2(i \tan(e+fx)+1)^2(c-ic \tan(e+fx))^{5/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{c \int \frac{A+B \tan(e+fx)}{(i \tan(e+fx)+1)^2(c-ic \tan(e+fx))^{5/2}} d \tan(e + fx)}{af} \\
 & \quad \downarrow \text{87} \\
 & \frac{c \left(\frac{1}{4} (5A + iB) \int \frac{1}{(i \tan(e+fx)+1)(c-ic \tan(e+fx))^{5/2}} d \tan(e + fx) + \frac{-B+iA}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} \right)}{af} \\
 & \quad \downarrow \text{61} \\
 & \frac{c \left(\frac{1}{4} (5A + iB) \left(\frac{\int \frac{1}{(i \tan(e+fx)+1)(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{2c} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} \right) + \frac{-B+iA}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} \right)}{af}
 \end{aligned}$$

3.769. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} dx$

$$\begin{array}{c} \downarrow 61 \\ c \left(\frac{1}{4}(5A + iB) \left(\frac{\int \frac{1}{(i \tan(e+fx)+1)\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{2c} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} \right) + \frac{-B+iA}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} \right) \end{array}$$

af

$$\begin{array}{c} \downarrow 73 \\ c \left(\frac{1}{4}(5A + iB) \left(\frac{\int \frac{1}{2 - \frac{c-ic \tan(e+fx)}{c}} d\sqrt{c-ic \tan(e+fx)}}{c^2} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} \right) + \frac{-B+iA}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} \right) \end{array}$$

af

$$\begin{array}{c} \downarrow 219 \\ c \left(\frac{1}{4}(5A + iB) \left(\frac{i \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{\sqrt{2}c^{3/2}} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} \right) + \frac{-B+iA}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} \right) \end{array}$$

af

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2)),x]`

output `(c*((I*A - B)/(2*c*(1 + I*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2)) + ((5*A + I*B)*((-1/3*I)/(c*(c - I*c*Tan[e + f*x])^(3/2)) + ((I*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])])/(Sqrt[2]*c^(3/2)) - I/(c*Sqrt[c - I*c*Tan[e + f*x]])))/(2*c)))/4)/(a*f)`

3.769.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 61 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p_.)), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.769.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.77

method	result
derivativedivides	$2ic \left(\frac{\left(\frac{iB}{8} + \frac{A}{8}\right) \sqrt{c-ic \tan(fx+e)} + \left(\frac{iB}{2} + \frac{5A}{2}\right) \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{\frac{c}{2} + \frac{ic \tan(fx+e)}{2}} + \frac{}{4c^2} - \frac{A}{4c^2 \sqrt{c-ic \tan(fx+e)}} - \frac{-iB+A}{12c(c-ic \tan(fx+e))} \right) \frac{1}{fa}$
default	$2ic \left(\frac{\left(\frac{iB}{8} + \frac{A}{8}\right) \sqrt{c-ic \tan(fx+e)} + \left(\frac{iB}{2} + \frac{5A}{2}\right) \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{\frac{c}{2} + \frac{ic \tan(fx+e)}{2}} + \frac{}{4c^2} - \frac{A}{4c^2 \sqrt{c-ic \tan(fx+e)}} - \frac{-iB+A}{12c(c-ic \tan(fx+e))} \right) \frac{1}{fa}$

```
input int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
```

```
output 2*I/f/a*c*(1/4/c^2*((1/8*I*B+1/8*A)*(c-I*c*tan(f*x+e))^(1/2)/(1/2*c+1/2*I*c*tan(f*x+e))+1/4*(1/2*I*B+5/2*A)*2^(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2)))-1/4/c^2*A/(c-I*c*tan(f*x+e))^(1/2)-1/12/c*(A-I*B)/(c-I*c*tan(f*x+e))^(3/2))
```

3.769.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 388 vs. 2(139) = 278.

Time = 0.26 (sec) , antiderivative size = 388, normalized size of antiderivative = 2.11

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^{3/2}} dx = \left(3 \sqrt{\frac{1}{2}} ac^2 f \sqrt{\frac{-25 A^2 + 10i AB - B^2}{a^2 c^3 f^2}} e^{(2i fx + 2i e)} \log \left(\frac{\sqrt{2} \sqrt{\frac{1}{2}} (a + ia \tan(e + fx))}{\dots} \right) \right)$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x,algorithm="fricas")
```

```
output 1/48*(3*sqrt(1/2)*a*c^2*f*sqrt(-(25*A^2 + 10*I*A*B - B^2)/(a^2*c^3*f^2))*e
^(2*I*f*x + 2*I*e)*log(1/4*(sqrt(2)*sqrt(1/2)*(a*c*f*e^(2*I*f*x + 2*I*e) +
a*c*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(25*A^2 + 10*I*A*B - B^2)/
(a^2*c^3*f^2)) + 5*I*A - B)*e^(-I*f*x - I*e)/(a*c*f)) - 3*sqrt(1/2)*a*c^2*
f*sqrt(-(25*A^2 + 10*I*A*B - B^2)/(a^2*c^3*f^2))*e^(2*I*f*x + 2*I*e)*log(-
1/4*(sqrt(2)*sqrt(1/2)*(a*c*f*e^(2*I*f*x + 2*I*e) + a*c*f)*sqrt(c/(e^(2*I*
f*x + 2*I*e) + 1))*sqrt(-(25*A^2 + 10*I*A*B - B^2)/(a^2*c^3*f^2)) - 5*I*A
+ B)*e^(-I*f*x - I*e)/(a*c*f)) - sqrt(2)*(2*(I*A + B)*e^(6*I*f*x + 6*I*e)
+ 4*(4*I*A + B)*e^(4*I*f*x + 4*I*e) - (-11*I*A - 5*B)*e^(2*I*f*x + 2*I*e)
- 3*I*A + 3*B)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*e^(-2*I*f*x - 2*I*e)/(a*
c^2*f)
```

3.769.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^{3/2}} dx =$$

$$\frac{i \left(\int \frac{A}{-ic\sqrt{-ic \tan(e+fx)+c \tan^2(e+fx)-ic\sqrt{-ic \tan(e+fx)+c}} dx + \int \frac{B \tan(e+fx)}{-ic\sqrt{-ic \tan(e+fx)+c \tan^2(e+fx)-ic\sqrt{-ic \tan(e+fx)+c}} dx \right)}{a}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))**(3/2),x)
```

```
output -I*(Integral(A/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - I*c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(B*tan(e + f*x)/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - I*c*sqrt(-I*c*tan(e + f*x) + c)), x))/a
```

3.769.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.91

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^{3/2}} dx =$$

$$i \left(\frac{3\sqrt{2}(5A+iB) \log\left(\frac{-\sqrt{2}\sqrt{c}-\sqrt{-ic \tan(fx+e)+c}}{\sqrt{2}\sqrt{c}+\sqrt{-ic \tan(fx+e)+c}}\right)}{a\sqrt{c}} + \frac{4\left(3(-ic \tan(fx+e)+c)^2(5A+iB)-4(-ic \tan(fx+e)+c)(5A+iB)c-8(A-iB)c^2\right)}{(-ic \tan(fx+e)+c)^{\frac{5}{2}}a-2(-ic \tan(fx+e)+c)^{\frac{3}{2}}ac} \right)$$

96 cf

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x,
algorithm="maxima")
```

$$3.769. \quad \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} dx$$

output
$$\frac{-1/96 I (3 \sqrt{2} (5A + I B) \log(-\sqrt{2} \sqrt{c} - \sqrt{-I c \tan(fx + e) + c}) / (\sqrt{2} \sqrt{c} + \sqrt{-I c \tan(fx + e) + c})) / (a \sqrt{c}) + 4 * (3 * (-I c \tan(fx + e) + c)^2 (5A + I B) - 4 * (-I c \tan(fx + e) + c) (5A + I B) c - 8 * (A - I B) c^2) / ((-I c \tan(fx + e) + c)^{5/2} a - 2 * (-I c \tan(fx + e) + c)^{3/2} a c)}{c f}$$

3.769.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + i a \tan(e + fx))(c - i c \tan(e + fx))^{3/2}} dx = \int \frac{B \tan(fx + e) + A}{(i a \tan(fx + e) + a)(-i c \tan(fx + e) + c)^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)*(-I*c*tan(f*x + e) + c)^(3/2)), x)`

3.769.9 Mupad [B] (verification not implemented)

Time = 9.70 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.42

$$\int \frac{A + B \tan(e + fx)}{(a + i a \tan(e + fx))(c - i c \tan(e + fx))^{3/2}} dx = \frac{\frac{Bc}{3} - \frac{B(c - c \tan(e + fx) \operatorname{li})}{6} + \frac{B(c - c \tan(e + fx) \operatorname{li})^2}{8c}}{a f (c - c \tan(e + fx) \operatorname{li})^{5/2} - 2 a c f (c - c \tan(e + fx) \operatorname{li})} - \frac{\frac{A(c - c \tan(e + fx) \operatorname{li}) \operatorname{li}}{6 a f} + \frac{A c \operatorname{li}}{3 a f} - \frac{A(c - c \tan(e + fx) \operatorname{li})^2 \operatorname{li}}{8 a c f}}{2 c (c - c \tan(e + fx) \operatorname{li})^{3/2} - (c - c \tan(e + fx) \operatorname{li})^{5/2}} + \frac{\sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) \operatorname{li}}}{2 \sqrt{-c}}\right) \operatorname{li}}{16 a (-c)^{3/2} f} - \frac{\sqrt{2} B \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) \operatorname{li}}}{2 \sqrt{c}}\right)}{16 a c^{3/2} f}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*li)*(c - c*tan(e + f*x)*li)^(3/2)),x)`

output $((B*c)/3 - (B*(c - c*\tan(e + f*x)*1i))/6 + (B*(c - c*\tan(e + f*x)*1i)^2)/(8*c))/(a*f*(c - c*\tan(e + f*x)*1i)^{(5/2)} - 2*a*c*f*(c - c*\tan(e + f*x)*1i)^{(3/2)}) - ((A*(c - c*\tan(e + f*x)*1i)*5i)/(6*a*f) + (A*c*1i)/(3*a*f) - (A*(c - c*\tan(e + f*x)*1i)^2*5i)/(8*a*c*f))/(2*c*(c - c*\tan(e + f*x)*1i)^{(3/2)} - (c - c*\tan(e + f*x)*1i)^{(5/2)}) + (2^{(1/2)}*A*atan(2^{(1/2)}*(c - c*\tan(e + f*x)*1i)^{(1/2)})/(2*(-c)^{(1/2)}))*5i/(16*a*(-c)^{(3/2)}*f) - (2^{(1/2)}*B*atanh(2^{(1/2)}*(c - c*\tan(e + f*x)*1i)^{(1/2)})/(2*c^{(1/2)}))/(16*a*c^{(3/2)}*f)$

3.769. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} dx$

$$3.770 \quad \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^{5/2}} dx$$

3.770.1 Optimal result 6985
 3.770.2 Mathematica [C] (verified) 6985
 3.770.3 Rubi [A] (verified) 6986
 3.770.4 Maple [A] (verified) 6989
 3.770.5 Fricas [B] (verification not implemented) 6989
 3.770.6 Sympy [F] 6990
 3.770.7 Maxima [A] (verification not implemented) 6990
 3.770.8 Giac [F] 6991
 3.770.9 Mupad [B] (verification not implemented) 6991

3.770.1 Optimal result

Integrand size = 43, antiderivative size = 223

$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^{5/2}} dx = \frac{(7iA-3B)\operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{16\sqrt{2}ac^{5/2}f} - \frac{7iA-3B}{20af(c-ic \tan(e+fx))^{5/2}} + \frac{iA-B}{2af(1+i \tan(e+fx))(c-ic \tan(e+fx))^{5/2}} - \frac{7iA-3B}{24acf(c-ic \tan(e+fx))^{3/2}} - \frac{7iA-3B}{16ac^2f\sqrt{c-ic \tan(e+fx)}}$$

```
output 1/32*(7*I*A-3*B)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a/c
^(5/2)/f*2^(1/2)+1/16*(-7*I*A+3*B)/a/c^2/f/(c-I*c*tan(f*x+e))^(1/2)+1/20*(
-7*I*A+3*B)/a/f/(c-I*c*tan(f*x+e))^(5/2)+1/2*(I*A-B)/a/f/(1+I*tan(f*x+e))/
(c-I*c*tan(f*x+e))^(5/2)+1/24*(-7*I*A+3*B)/a/c/f/(c-I*c*tan(f*x+e))^(3/2)
```

3.770.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 5.85 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.57

$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^{5/2}} dx = \frac{5(7A+3iB) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, -\frac{1}{2}i(i+\tan(e+fx))\right)}{120ac^2f(-i+\tan(e+fx))}$$

3.770. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^{5/2}} dx$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2)),x]`

output `(5*(7*A + (3*I)*B)*Hypergeometric2F1[-3/2, 1, -1/2, (-1/2*I)*(I + Tan[e + f*x])] * Sec[e + f*x]^2 + (6*I)*((3*I)*A - 7*B + (7*A + (3*I)*B)*Tan[e + f*x]))/(120*a*c^2*f*(-I + Tan[e + f*x])*(I + Tan[e + f*x])^2*sqrt[c - I*c*Tan[e + f*x]])`

3.770.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.92, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.209$, Rules used = {3042, 4071, 27, 87, 61, 61, 61, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ict \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^{5/2}} dx$$

↓ 4071

$$ac \int \frac{A + B \tan(e + fx)}{a^2(i \tan(e + fx) + 1)^2(c - ic \tan(e + fx))^{7/2}} d \tan(e + fx)$$

f
↓ 27

$$c \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx) + 1)^2(c - ic \tan(e + fx))^{7/2}} d \tan(e + fx)$$

af
↓ 87

$$c \left(\frac{1}{4}(7A + 3iB) \int \frac{1}{(i \tan(e + fx) + 1)(c - ic \tan(e + fx))^{7/2}} d \tan(e + fx) + \frac{-B + iA}{2c(1 + i \tan(e + fx))(c - ic \tan(e + fx))^{5/2}} \right)$$

af
↓ 61

$$c \left(\frac{1}{4}(7A + 3iB) \left(\frac{\int \frac{1}{(i \tan(e + fx) + 1)(c - ic \tan(e + fx))^{5/2}} d \tan(e + fx)}{2c} - \frac{i}{5c(c - ic \tan(e + fx))^{5/2}} \right) + \frac{-B + iA}{2c(1 + i \tan(e + fx))(c - ic \tan(e + fx))^{5/2}} \right)$$

af

3.770. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^{5/2}} dx$

↓ 61

$$c \left(\frac{1}{4}(7A + 3iB) \left(\frac{\int \frac{1}{(i \tan(e+fx)+1)(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{2c} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} - \frac{i}{5c(c-ic \tan(e+fx))^{5/2}} \right) + \frac{1}{2c(1+i \tan(e+fx))} \right)$$

af

↓ 61

$$c \left(\frac{1}{4}(7A + 3iB) \left(\frac{\int \frac{1}{(i \tan(e+fx)+1)\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{2c} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} - \frac{i}{5c(c-ic \tan(e+fx))^{5/2}} \right) + \frac{1}{2c(1+i \tan(e+fx))} \right)$$

af

↓ 73

$$c \left(\frac{1}{4}(7A + 3iB) \left(\frac{i \int \frac{1}{2 - \frac{c-ic \tan(e+fx)}{c}} d\sqrt{c-ic \tan(e+fx)}}{c^2} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} - \frac{i}{5c(c-ic \tan(e+fx))^{5/2}} \right) + \frac{1}{2c(1+i \tan(e+fx))} \right)$$

af

↓ 219

$$c \left(\frac{1}{4}(7A + 3iB) \left(\frac{i \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{\sqrt{2}c^{3/2}} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} - \frac{i}{5c(c-ic \tan(e+fx))^{5/2}} \right) + \frac{1}{2c(1+i \tan(e+fx))} \right)$$

af

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2)),x]`

output `(c*((I*A - B)/(2*c*(1 + I*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2)) + ((7*A + (3*I)*B)*((-1/5*I)/(c*(c - I*c*Tan[e + f*x])^(5/2)) + ((-1/3*I)/(c*(c - I*c*Tan[e + f*x])^(3/2)) + ((I*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])])/(Sqrt[2]*c^(3/2)) - I/(c*Sqrt[c - I*c*Tan[e + f*x]])))/(2*c)))/(2*c))/4)/(a*f)`

3.770. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^{5/2}} dx$

3.770.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 61 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p_.)), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.770.4 Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.75

method	result
derivativedivides	$2ic \left(\frac{\left(\frac{iB}{4} + \frac{A}{4}\right) \sqrt{c-ic \tan(fx+e)} + \frac{\left(\frac{7A}{2} + \frac{3iB}{2}\right) \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{\frac{c}{2} + \frac{ic \tan(fx+e)}{2}} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{2\sqrt{c}} \right) - \frac{iB+3A}{16c^3 \sqrt{c-ic \tan(fx+e)}} - \frac{A}{12c^2(c-ic \tan(fx+e))}$
default	$2ic \left(\frac{\left(\frac{iB}{4} + \frac{A}{4}\right) \sqrt{c-ic \tan(fx+e)} + \frac{\left(\frac{7A}{2} + \frac{3iB}{2}\right) \sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{\frac{c}{2} + \frac{ic \tan(fx+e)}{2}} + \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}}\right)}{2\sqrt{c}} \right) - \frac{iB+3A}{16c^3 \sqrt{c-ic \tan(fx+e)}} - \frac{A}{12c^2(c-ic \tan(fx+e))}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

output `2*I/f/a*c*(1/16/c^3*((1/4*I*B+1/4*A)*(c-I*c*tan(f*x+e))^(1/2)/(1/2*c+1/2*I*c*tan(f*x+e))+1/2*(7/2*A+3/2*I*B)*2^(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2)))-1/16/c^3*(3*A+I*B)/(c-I*c*tan(f*x+e))^(1/2)-1/12/c^2*A/(c-I*c*tan(f*x+e))^(3/2)-1/20/c*(A-I*B)/(c-I*c*tan(f*x+e))^(5/2))`

3.770.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 418 vs. 2(172) = 344.

Time = 0.29 (sec) , antiderivative size = 418, normalized size of antiderivative = 1.87

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^{5/2}} dx = \left(15 \sqrt{\frac{1}{2}} ac^3 f \sqrt{-\frac{49A^2 + 42iAB - 9B^2}{a^2 c^5 f^2}} e^{(2i fx + 2i e)} \log \left(\frac{\sqrt{2} \sqrt{c - ic \tan(e + fx)}}{2\sqrt{c}} \right) \right)$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,algorithm="fricas")`

3.770. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^{5/2}} dx$

output `1/480*(15*sqrt(1/2)*a*c^3*f*sqrt(-(49*A^2 + 42*I*A*B - 9*B^2)/(a^2*c^5*f^2)))*e^(2*I*f*x + 2*I*e)*log(1/8*(sqrt(2)*sqrt(1/2)*(a*c^2*f*e^(2*I*f*x + 2*I*e) + a*c^2*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(49*A^2 + 42*I*A*B - 9*B^2)/(a^2*c^5*f^2)) + 7*I*A - 3*B)*e^(-I*f*x - I*e)/(a*c^2*f)) - 15*sqrt(1/2)*a*c^3*f*sqrt(-(49*A^2 + 42*I*A*B - 9*B^2)/(a^2*c^5*f^2))*e^(2*I*f*x + 2*I*e)*log(-1/8*(sqrt(2)*sqrt(1/2)*(a*c^2*f*e^(2*I*f*x + 2*I*e) + a*c^2*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(49*A^2 + 42*I*A*B - 9*B^2)/(a^2*c^5*f^2)) - 7*I*A + 3*B)*e^(-I*f*x - I*e)/(a*c^2*f)) - sqrt(2)*(6*(I*A + B)*e^(8*I*f*x + 8*I*e) + 2*(19*I*A + 9*B)*e^(6*I*f*x + 6*I*e) + 4*(37*I*A - 3*B)*e^(4*I*f*x + 4*I*e) - (-101*I*A + 9*B)*e^(2*I*f*x + 2*I*e) - 15*I*A + 15*B)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*e^(-2*I*f*x - 2*I*e)/(a*c^3*f)`

3.770.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^{5/2}} dx =$$

$$i \left(\int \frac{A}{-c^2 \sqrt{-ic \tan(e+fx)+c} \tan^3(e+fx) - ic^2 \sqrt{-ic \tan(e+fx)+c} \tan^2(e+fx) - c^2 \sqrt{-ic \tan(e+fx)+c} \tan(e+fx) - ic^2 \sqrt{-ic \tan(e+fx)+c}} dx \right) + \dots$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))**(5/2), x)`

output `-I*(Integral(A/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3 - I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) - I*c**2*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(B*tan(e + f*x)/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3 - I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2 - c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) - I*c**2*sqrt(-I*c*tan(e + f*x) + c)), x))/a`

3.770.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 195, normalized size of antiderivative = 0.87

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ic \tan(e + fx))^{5/2}} dx =$$

$$i \left(\frac{4 \left(15(-ic \tan(fx+e)+c)^3(7A+3iB) - 20(-ic \tan(fx+e)+c)^2(7A+3iB)c - 8(-ic \tan(fx+e)+c)(7A+3iB)c^2 - 48(A-iB)c^3 \right)}{(-ic \tan(fx+e)+c)^{7/2} ac - 2(-ic \tan(fx+e)+c)^{5/2} ac^2} + \frac{15 \sqrt{2}}{960 cf} \right)$$

$$3.770. \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^{5/2}} dx$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,
algorithm="maxima")`

output `-1/960*I*(4*(15*(-I*c*tan(f*x + e) + c)^3*(7*A + 3*I*B) - 20*(-I*c*tan(f*x
+ e) + c)^2*(7*A + 3*I*B)*c - 8*(-I*c*tan(f*x + e) + c)*(7*A + 3*I*B)*c^2
- 48*(A - I*B)*c^3)/((-I*c*tan(f*x + e) + c)^(7/2)*a*c - 2*(-I*c*tan(f*x
+ e) + c)^(5/2)*a*c^2) + 15*sqrt(2)*(7*A + 3*I*B)*log(-(sqrt(2)*sqrt(c) -
sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c
)))/(a*c^(3/2)))/(c*f)`

3.770.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ictan(e + fx))^{5/2}} dx = \int \frac{B \tan(fx + e) + A}{(ia \tan(fx + e) + a)(-ictan(fx + e) + c)^{5/2}} dx$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,
algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)*(-I*c*tan(f*x + e)
+ c)^(5/2)), x)`

3.770.9 Mupad [B] (verification not implemented)

Time = 10.10 (sec) , antiderivative size = 308, normalized size of antiderivative = 1.38

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))(c - ictan(e + fx))^{5/2}} dx = \frac{\frac{Bc}{5} - \frac{B(c - c \tan(e + fx) li)}{10} - \frac{B(c - c \tan(e + fx) li)^2}{4c}}{af(c - c \tan(e + fx) li)^{7/2} - 2acf(c - c \tan(e + fx) li)} + \frac{3B(c - c \tan(e + fx) li)}{16c}$$

$$- \frac{\frac{A(c - c \tan(e + fx) li) \gamma i}{30af} + \frac{Ac li}{5af} + \frac{A(c - c \tan(e + fx) li)^2 \gamma i}{12acf} - \frac{A(c - c \tan(e + fx) li)^3 \gamma i}{16ac^2 f}}{2c(c - c \tan(e + fx) li)^{5/2} - (c - c \tan(e + fx) li)^{7/2}}$$

$$- \frac{\sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) li}}{2\sqrt{-c}}\right) \gamma i}{32a(-c)^{5/2} f} - \frac{3\sqrt{2} B \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) li}}{2\sqrt{c}}\right)}{32ac^{5/2} f}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*li)*(c - c*tan(e + f*x)*li)^(5/2)),x)`

3.770. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ictan(e+fx))^{5/2}} dx$

output $((B*c)/5 - (B*(c - c*\tan(e + f*x)*1i))/10 - (B*(c - c*\tan(e + f*x)*1i)^2)/(4*c) + (3*B*(c - c*\tan(e + f*x)*1i)^3)/(16*c^2))/(a*f*(c - c*\tan(e + f*x)*1i)^{7/2} - 2*a*c*f*(c - c*\tan(e + f*x)*1i)^{5/2}) - ((A*(c - c*\tan(e + f*x)*1i)*7i)/(30*a*f) + (A*c*1i)/(5*a*f) + (A*(c - c*\tan(e + f*x)*1i)^2*7i)/(12*a*c*f) - (A*(c - c*\tan(e + f*x)*1i)^3*7i)/(16*a*c^2*f))/(2*c*(c - c*\tan(e + f*x)*1i)^{5/2} - (c - c*\tan(e + f*x)*1i)^{7/2}) - (2^{1/2}*A*atan((2^{1/2}*(c - c*\tan(e + f*x)*1i)^{1/2})/(2*(-c)^{1/2}))*7i)/(32*a*(-c)^{5/2}*f) - (3*2^{1/2}*B*atanh((2^{1/2}*(c - c*\tan(e + f*x)*1i)^{1/2})/(2*c^{1/2}))))/(32*a*c^{5/2}*f)$

3.770. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))(c-ic \tan(e+fx))^{5/2}} dx$

3.771
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^2} dx$$

3.771.1 Optimal result	6993
3.771.2 Mathematica [A] (verified)	6994
3.771.3 Rubi [A] (verified)	6994
3.771.4 Maple [A] (verified)	6997
3.771.5 Fracas [B] (verification not implemented)	6998
3.771.6 Sympy [F(-1)]	6999
3.771.7 Maxima [A] (verification not implemented)	6999
3.771.8 Giac [F]	7000
3.771.9 Mupad [B] (verification not implemented)	7000

3.771.1 Optimal result

Integrand size = 43, antiderivative size = 275

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{9/2}}{(a + ia \tan(e + fx))^2} dx = \frac{7(5iA - 13B)c^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{\sqrt{2}a^2 f} - \frac{7(5iA - 13B)c^4 \sqrt{c - ic \tan(e + fx)}}{2a^2 f} - \frac{7(5iA - 13B)c^3 (c - ic \tan(e + fx))^{3/2}}{12a^2 f} - \frac{7(5iA - 13B)c^2 (c - ic \tan(e + fx))^{5/2}}{40a^2 f} - \frac{(5iA - 13B)c(c - ic \tan(e + fx))^{7/2}}{8a^2 f(1 + i \tan(e + fx))} + \frac{(iA - B)(c - ic \tan(e + fx))^{9/2}}{4a^2 f(1 + i \tan(e + fx))^2}$$

```
output 7/2*(5*I*A-13*B)*c^(9/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a^2/f*2^(1/2)-7/2*(5*I*A-13*B)*c^4*(c-I*c*tan(f*x+e))^(1/2)/a^2/f-7/12*(5*I*A-13*B)*c^3*(c-I*c*tan(f*x+e))^(3/2)/a^2/f-7/40*(5*I*A-13*B)*c^2*(c-I*c*tan(f*x+e))^(5/2)/a^2/f-1/8*(5*I*A-13*B)*c*(c-I*c*tan(f*x+e))^(7/2)/a^2/f/(1+I*tan(f*x+e))+1/4*(I*A-B)*(c-I*c*tan(f*x+e))^(9/2)/a^2/f/(1+I*tan(f*x+e))^2
```


3.771.2 Mathematica [A] (verified)

Time = 7.92 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.71

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^2} dx = \frac{105\sqrt{2}(-5iA + 13B)c^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{c - i c \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right) \sec^2(e + fx)}{(a + i a \tan(e + fx))^2}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(9/2))/(a + I*a*Tan[e + f*x])^2,x]`

output `(105*Sqrt[2]*((-5*I)*A + 13*B)*c^(9/2)*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])]*Sec[e + f*x]^2*(Cos[2*(e + f*x)] + I*Sin[2*(e + f*x)]) - 2*c^4*Sqrt[c - I*c*Tan[e + f*x]]*((-325*I)*A + 851*B + 5*(113*A + (289*I)*B)*Tan[e + f*x] + ((170*I)*A - 478*B)*Tan[e + f*x]^2 + 10*(A + (5*I)*B)*Tan[e + f*x]^3 + 6*B*Tan[e + f*x]^4))/(30*a^2*f*(-I + Tan[e + f*x])^2)`

3.771.3 Rubi [A] (verified)Time = 0.42 (sec) , antiderivative size = 229, normalized size of antiderivative = 0.83, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.233$, Rules used = {3042, 4071, 27, 87, 51, 60, 60, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c - i c \tan(e + fx))^{9/2}(A + B \tan(e + fx))}{(a + i a \tan(e + fx))^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c - i c \tan(e + fx))^{9/2}(A + B \tan(e + fx))}{(a + i a \tan(e + fx))^2} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{a^3(i \tan(e + fx) + 1)^3} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{c \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(i \tan(e + fx) + 1)^3} d \tan(e + fx)}{a^2 f} \end{aligned}$$

3.771. $\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^2} dx$

$$\begin{aligned}
 & \downarrow 87 \\
 & \frac{c \left(\frac{(-B+iA)(c-ictan(e+fx))^{9/2}}{4c(1+i \tan(e+fx))^2} - \frac{1}{8}(5A + 13iB) \int \frac{(c-ictan(e+fx))^{7/2}}{(i \tan(e+fx)+1)^2} d \tan(e + fx) \right)}{a^2 f} \\
 & \downarrow 51 \\
 & \frac{c \left(\frac{(-B+iA)(c-ictan(e+fx))^{9/2}}{4c(1+i \tan(e+fx))^2} - \frac{1}{8}(5A + 13iB) \left(\frac{i(c-ictan(e+fx))^{7/2}}{1+i \tan(e+fx)} - \frac{7}{2}c \int \frac{(c-ictan(e+fx))^{5/2}}{i \tan(e+fx)+1} d \tan(e + fx) \right) \right)}{a^2 f} \\
 & \downarrow 60 \\
 & \frac{c \left(\frac{(-B+iA)(c-ictan(e+fx))^{9/2}}{4c(1+i \tan(e+fx))^2} - \frac{1}{8}(5A + 13iB) \left(\frac{i(c-ictan(e+fx))^{7/2}}{1+i \tan(e+fx)} - \frac{7}{2}c \left(2c \int \frac{(c-ictan(e+fx))^{3/2}}{i \tan(e+fx)+1} d \tan(e + fx) - \frac{2}{5}i(c-ictan(e+fx))^{5/2} \right) \right) \right)}{a^2 f} \\
 & \downarrow 60 \\
 & \frac{c \left(\frac{(-B+iA)(c-ictan(e+fx))^{9/2}}{4c(1+i \tan(e+fx))^2} - \frac{1}{8}(5A + 13iB) \left(\frac{i(c-ictan(e+fx))^{7/2}}{1+i \tan(e+fx)} - \frac{7}{2}c \left(2c \left(2c \int \frac{\sqrt{c-ictan(e+fx)}}{i \tan(e+fx)+1} d \tan(e + fx) - \frac{2}{3}i(c-ictan(e+fx))^{3/2} \right) \right) \right) \right)}{a^2 f} \\
 & \downarrow 60 \\
 & \frac{c \left(\frac{(-B+iA)(c-ictan(e+fx))^{9/2}}{4c(1+i \tan(e+fx))^2} - \frac{1}{8}(5A + 13iB) \left(\frac{i(c-ictan(e+fx))^{7/2}}{1+i \tan(e+fx)} - \frac{7}{2}c \left(2c \left(2c \int \frac{1}{(i \tan(e+fx)+1)\sqrt{c-ictan(e+fx)}} d \tan(e + fx) - \frac{2}{5}i(c-ictan(e+fx))^{5/2} \right) \right) \right) \right)}{a^2 f} \\
 & \downarrow 73 \\
 & \frac{c \left(\frac{(-B+iA)(c-ictan(e+fx))^{9/2}}{4c(1+i \tan(e+fx))^2} - \frac{1}{8}(5A + 13iB) \left(\frac{i(c-ictan(e+fx))^{7/2}}{1+i \tan(e+fx)} - \frac{7}{2}c \left(2c \left(2c \left(4i \int \frac{1}{2-\frac{c-ictan(e+fx)}{c}} d \sqrt{c-ictan(e+fx)} - \frac{2}{5}i(c-ictan(e+fx))^{5/2} \right) \right) \right) \right) \right)}{a^2 f} \\
 & \downarrow 219 \\
 & \frac{c \left(\frac{(-B+iA)(c-ictan(e+fx))^{9/2}}{4c(1+i \tan(e+fx))^2} - \frac{1}{8}(5A + 13iB) \left(\frac{i(c-ictan(e+fx))^{7/2}}{1+i \tan(e+fx)} - \frac{7}{2}c \left(2c \left(2c \left(2i\sqrt{2}\sqrt{c} \operatorname{arctanh} \left(\frac{\sqrt{c-ictan(e+fx)}}{\sqrt{2}\sqrt{c}} \right) - \frac{2}{5}i(c-ictan(e+fx))^{5/2} \right) \right) \right) \right) \right)}{a^2 f}
 \end{aligned}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(9/2))/(a + I*a*Tan[e + f*x])^2,x]`

3.771. $\int \frac{(A+B \tan(e+fx))(c-ictan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^2} dx$

```
output (c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(9/2))/(4*c*(1 + I*Tan[e + f*x])^2)
- ((5*A + (13*I)*B)*((I*(c - I*c*Tan[e + f*x])^(7/2))/(1 + I*Tan[e + f*x])
- (7*c*((( -2*I)/5)*(c - I*c*Tan[e + f*x])^(5/2) + 2*c*((( -2*I)/3)*(c - I*
c*Tan[e + f*x])^(3/2) + 2*c*((2*I)*Sqrt[2]*Sqrt[c]*ArcTanh[Sqrt[c - I*c*Ta
n[e + f*x]]/(Sqrt[2]*Sqrt[c])) - (2*I)*Sqrt[c - I*c*Tan[e + f*x]]))))/2))/
8))/(a^2*f)
```

3.771.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 51 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x
] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]
```

```
rule 60 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(
b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.771.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 220, normalized size of antiderivative = 0.80

method	result
derivativedivides	$2ic^2 \left(-\frac{iB(c-ic \tan(fx+e))^{5/2}}{5} - \frac{5iBc(c-ic \tan(fx+e))^{3/2}}{3} - \frac{Ac(c-ic \tan(fx+e))^{3/2}}{3} - 18i\sqrt{c-ic \tan(fx+e)} Bc^2 - 6\sqrt{c-ic \tan(fx+e)} Bc \right)$
default	$2ic^2 \left(-\frac{iB(c-ic \tan(fx+e))^{5/2}}{5} - \frac{5iBc(c-ic \tan(fx+e))^{3/2}}{3} - \frac{Ac(c-ic \tan(fx+e))^{3/2}}{3} - 18i\sqrt{c-ic \tan(fx+e)} Bc^2 - 6\sqrt{c-ic \tan(fx+e)} Bc \right)$

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2)/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `2*I/f/a^2*c^2*(-1/5*I*B*(c-I*c*tan(f*x+e))^(5/2)-5/3*I*B*c*(c-I*c*tan(f*x+e))^(3/2)-1/3*A*c*(c-I*c*tan(f*x+e))^(3/2)-18*I*(c-I*c*tan(f*x+e))^(1/2)*B*c^2-6*(c-I*c*tan(f*x+e))^(1/2)*A*c^2+8*c^3*(4*((21/64*I*B+13/64*A)*(c-I*c*tan(f*x+e))^(3/2)+(-19/32*I*B*c-11/32*c*A)*(c-I*c*tan(f*x+e))^(1/2)))/(c+I*c*tan(f*x+e))^2+7/8*(13/4*I*B+5/4*A)*2^(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2)))`

3.771.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^2} dx$$

3.771.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 513 vs. $2(216) = 432$.

Time = 0.28 (sec) , antiderivative size = 513, normalized size of antiderivative = 1.87

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{9/2}}{(a + ia \tan(e + fx))^2} dx =$$

$$105 \sqrt{2} \sqrt{-\frac{(25A^2 + 130iAB - 169B^2)c^9}{a^4 f^2}} (a^2 f e^{(8i fx + 8i e)} + 2a^2 f e^{(6i fx + 6i e)} + a^2 f e^{(4i fx + 4i e)}) \log \left(-\frac{14 \left((-5iA + 13B) \right)}{\dots} \right)$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2)/(a+I*a*tan(f*x+e))^2,x
, algorithm="fricas")`

output `-1/60*(105*sqrt(2)*sqrt(-(25*A^2 + 130*I*A*B - 169*B^2)*c^9/(a^4*f^2))*(a^2*f*e^(8*I*f*x + 8*I*e) + 2*a^2*f*e^(6*I*f*x + 6*I*e) + a^2*f*e^(4*I*f*x + 4*I*e))*log(-14*((-5*I*A + 13*B)*c^5 + sqrt(-(25*A^2 + 130*I*A*B - 169*B^2)*c^9/(a^4*f^2))*(a^2*f*e^(2*I*f*x + 2*I*e) + a^2*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))*e^(-I*f*x - I*e)/(a^2*f)) - 105*sqrt(2)*sqrt(-(25*A^2 + 130*I*A*B - 169*B^2)*c^9/(a^4*f^2))*(a^2*f*e^(8*I*f*x + 8*I*e) + 2*a^2*f*e^(6*I*f*x + 6*I*e) + a^2*f*e^(4*I*f*x + 4*I*e))*log(-14*((-5*I*A + 13*B)*c^5 - sqrt(-(25*A^2 + 130*I*A*B - 169*B^2)*c^9/(a^4*f^2))*(a^2*f*e^(2*I*f*x + 2*I*e) + a^2*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))*e^(-I*f*x - I*e)/(a^2*f)) + 2*sqrt(2)*(105*(5*I*A - 13*B)*c^4*e^(8*I*f*x + 8*I*e) + 245*(5*I*A - 13*B)*c^4*e^(6*I*f*x + 6*I*e) + 161*(5*I*A - 13*B)*c^4*e^(4*I*f*x + 4*I*e) + 15*(5*I*A - 13*B)*c^4*e^(2*I*f*x + 2*I*e) + 30*(-I*A + B)*c^4)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(a^2*f*e^(8*I*f*x + 8*I*e) + 2*a^2*f*e^(6*I*f*x + 6*I*e) + a^2*f*e^(4*I*f*x + 4*I*e))`

3.771.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^2} dx = \text{Timed out}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(9/2)/(a+I*a*tan(f*x+e))**2
,x)
```

```
output Timed out
```

3.771.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 244, normalized size of antiderivative = 0.89

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^2} dx =$$

$$i \left(\frac{105 \sqrt{2} (5A + 13iB) c^{\frac{11}{2}} \log\left(-\frac{\sqrt{2}\sqrt{c} - \sqrt{-i c \tan(fx+e)+c}}{\sqrt{2}\sqrt{c} + \sqrt{-i c \tan(fx+e)+c}}\right)}{a^2} - \frac{60 \left((-i c \tan(fx+e)+c\right)^{\frac{3}{2}} (13A + 21iB) c^6 - 2 \sqrt{-i c \tan(fx+e)+c} (11A + 19iB) c^5}{(-i c \tan(fx+e)+c)^2 a^2 - 4(-i c \tan(fx+e)+c) a^2 c + 4 a^2 c^2} \right)$$

60 cf

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2)/(a+I*a*tan(f*x+e))^2,x
, algorithm="maxima")
```

```
output -1/60*I*(105*sqrt(2)*(5*A + 13*I*B)*c^(11/2)*log(-(sqrt(2)*sqrt(c) - sqrt(
-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a
^2 - 60*((-I*c*tan(f*x + e) + c)^(3/2)*(13*A + 21*I*B)*c^6 - 2*sqrt(-I*c*t
an(f*x + e) + c)*(11*A + 19*I*B)*c^7)/((-I*c*tan(f*x + e) + c)^2*a^2 - 4*(
-I*c*tan(f*x + e) + c)*a^2*c + 4*a^2*c^2) + 8*(3*I*(-I*c*tan(f*x + e) + c)
^(5/2)*B*c^3 + 5*(-I*c*tan(f*x + e) + c)^(3/2)*(A + 5*I*B)*c^4 + 90*sqrt(-
I*c*tan(f*x + e) + c)*(A + 3*I*B)*c^5)/a^2)/(c*f)
```

3.771.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^2} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{9/2}}{(i a \tan(fx + e) + a)^2} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2)/(a+I*a*tan(f*x+e))^2,x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(9/2)/(I*a*tan(f*x
+ e) + a)^2, x)`

3.771.9 Mupad [B] (verification not implemented)

Time = 8.95 (sec) , antiderivative size = 402, normalized size of antiderivative = 1.46

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^2} dx = \frac{38 B c^6 \sqrt{c - c \tan(e + f x) i} - 21 B c^5 (c - c \tan(e + f x) i)^{3/2}}{4 a^2 c^2 f + a^2 f (c - c \tan(e + f x) i)^2 - 4 a^2 c f (c - c \tan(e + f x) i)} + \frac{A c^6 \sqrt{c - c \tan(e + f x) i} 22 i - A c^5 (c - c \tan(e + f x) i)^{3/2} 13 i}{a^2 f} - \frac{(c - c \tan(e + f x) i)^2 - 4 c (c - c \tan(e + f x) i) + 4 c^2}{a^2 f} + \frac{A c^4 \sqrt{c - c \tan(e + f x) i} 12 i - A c^3 (c - c \tan(e + f x) i)^{3/2} 2 i}{3 a^2 f} + \frac{36 B c^4 \sqrt{c - c \tan(e + f x) i}}{a^2 f} + \frac{10 B c^3 (c - c \tan(e + f x) i)^{3/2}}{3 a^2 f} + \frac{2 B c^2 (c - c \tan(e + f x) i)^{5/2}}{5 a^2 f} + \frac{\sqrt{2} A (-c)^{9/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + f x) i}}{2 \sqrt{-c}}\right) 35 i}{2 a^2 f} + \frac{\sqrt{2} B c^{9/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + f x) i} i}{2 \sqrt{c}}\right) 91 i}{2 a^2 f}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*i)^(9/2))/(a + a*tan(e + f*
x)*i)^2,x)`

output

$$\begin{aligned}
& (38*B*c^6*(c - c*\tan(e + f*x)*1i)^{(1/2)} - 21*B*c^5*(c - c*\tan(e + f*x)*1i)^{(3/2)}) / (4*a^2*c^2*f + a^2*f*(c - c*\tan(e + f*x)*1i)^2 - 4*a^2*c*f*(c - c*\tan(e + f*x)*1i)) - ((A*c^6*(c - c*\tan(e + f*x)*1i)^{(1/2)*22i} / (a^2*f) - (A*c^5*(c - c*\tan(e + f*x)*1i)^{(3/2)*13i} / (a^2*f))) / ((c - c*\tan(e + f*x)*1i)^2 - 4*c*(c - c*\tan(e + f*x)*1i) + 4*c^2) - (A*c^4*(c - c*\tan(e + f*x)*1i)^{(1/2)*12i} / (a^2*f) - (A*c^3*(c - c*\tan(e + f*x)*1i)^{(3/2)*2i} / (3*a^2*f) + (36*B*c^4*(c - c*\tan(e + f*x)*1i)^{(1/2}) / (a^2*f) + (10*B*c^3*(c - c*\tan(e + f*x)*1i)^{(3/2}) / (3*a^2*f) + (2*B*c^2*(c - c*\tan(e + f*x)*1i)^{(5/2}) / (5*a^2*f) + (2^{(1/2)}*A*(-c)^{(9/2)}*atan((2^{(1/2)}*(c - c*\tan(e + f*x)*1i)^{(1/2)}) / (2*(-c)^{(1/2)})))*35i) / (2*a^2*f) + (2^{(1/2)}*B*c^{(9/2)}*atan((2^{(1/2)}*(c - c*\tan(e + f*x)*1i)^{(1/2)*1i} / (2*c^{(1/2)})))*91i) / (2*a^2*f)
\end{aligned}$$

3.772
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^2} dx$$

3.772.1 Optimal result	7002
3.772.2 Mathematica [A] (verified)	7003
3.772.3 Rubi [A] (verified)	7003
3.772.4 Maple [A] (verified)	7006
3.772.5 Fracas [B] (verification not implemented)	7006
3.772.6 Sympy [F]	7007
3.772.7 Maxima [A] (verification not implemented)	7008
3.772.8 Giac [F]	7008
3.772.9 Mupad [B] (verification not implemented)	7009

3.772.1 Optimal result

Integrand size = 43, antiderivative size = 238

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2}}{(a + ia \tan(e + fx))^2} dx = \frac{5(3iA - 11B)c^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{2\sqrt{2}a^2 f} - \frac{5(3iA - 11B)c^3 \sqrt{c - ic \tan(e + fx)}}{4a^2 f} - \frac{5(3iA - 11B)c^2 (c - ic \tan(e + fx))^{3/2}}{24a^2 f} - \frac{(3iA - 11B)c(c - ic \tan(e + fx))^{5/2}}{8a^2 f(1 + i \tan(e + fx))} + \frac{(iA - B)(c - ic \tan(e + fx))^{7/2}}{4a^2 f(1 + i \tan(e + fx))^2}$$

```
output 5/4*(3*I*A-11*B)*c^(7/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a^2/f*2^(1/2)-5/4*(3*I*A-11*B)*c^3*(c-I*c*tan(f*x+e))^(1/2)/a^2/f-5/24*(3*I*A-11*B)*c^2*(c-I*c*tan(f*x+e))^(3/2)/a^2/f-1/8*(3*I*A-11*B)*c*(c-I*c*tan(f*x+e))^(5/2)/a^2/f/(1+I*tan(f*x+e))+1/4*(I*A-B)*(c-I*c*tan(f*x+e))^(7/2)/a^2/f/(1+I*tan(f*x+e))^2
```

3.772.2 Mathematica [A] (verified)

Time = 6.89 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.76

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^2} dx = \frac{15\sqrt{2}(-3iA + 11B)c^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{c - i c \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right) \sec^2(e + fx)}{(a + i a \tan(e + fx))^2}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2))/(a + I*a*Tan[e + f*x])^2,x]`

output `(15*Sqrt[2]*((-3*I)*A + 11*B)*c^(7/2)*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])]*Sec[e + f*x]^2*(Cos[2*(e + f*x)] + I*Sin[2*(e + f*x)]) + 2*c^3*Sqrt[c - I*c*Tan[e + f*x]]*((27*I)*A - 103*B - (51*A + (175*I)*B)*Tan[e + f*x] + 4*((-3*I)*A + 14*B)*Tan[e + f*x]^2 - (4*I)*B*Tan[e + f*x]^3)/(12*a^2*f*(-I + Tan[e + f*x])^2)`

3.772.3 Rubi [A] (verified)Time = 0.41 (sec) , antiderivative size = 202, normalized size of antiderivative = 0.85, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.209$, Rules used = {3042, 4071, 27, 87, 51, 60, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c - i c \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(a + i a \tan(e + fx))^2} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c - i c \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(a + i a \tan(e + fx))^2} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{a^3 (i \tan(e + fx) + 1)^3} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{c \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(i \tan(e + fx) + 1)^3} d \tan(e + fx)}{a^2 f} \end{aligned}$$

3.772. $\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^2} dx$

$$\begin{aligned}
 & \downarrow 87 \\
 & \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{7/2}}{4c(1+i\tan(e+fx))^2} - \frac{1}{8}(3A+11iB) \int \frac{(c-ictan(e+fx))^{5/2}}{(i\tan(e+fx)+1)^2} d\tan(e+fx)\right)}{a^2f} \\
 & \downarrow 51 \\
 & \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{7/2}}{4c(1+i\tan(e+fx))^2} - \frac{1}{8}(3A+11iB) \left(\frac{i(c-ictan(e+fx))^{5/2}}{1+i\tan(e+fx)} - \frac{5}{2}c \int \frac{(c-ictan(e+fx))^{3/2}}{i\tan(e+fx)+1} d\tan(e+fx)\right)\right)}{a^2f} \\
 & \downarrow 60 \\
 & \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{7/2}}{4c(1+i\tan(e+fx))^2} - \frac{1}{8}(3A+11iB) \left(\frac{i(c-ictan(e+fx))^{5/2}}{1+i\tan(e+fx)} - \frac{5}{2}c \left(2c \int \frac{\sqrt{c-ictan(e+fx)}}{i\tan(e+fx)+1} d\tan(e+fx) - \frac{2}{3}i(c-i\tan(e+fx))\right)\right)\right)}{a^2f} \\
 & \downarrow 60 \\
 & \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{7/2}}{4c(1+i\tan(e+fx))^2} - \frac{1}{8}(3A+11iB) \left(\frac{i(c-ictan(e+fx))^{5/2}}{1+i\tan(e+fx)} - \frac{5}{2}c \left(2c \left(2c \int \frac{1}{(i\tan(e+fx)+1)\sqrt{c-ictan(e+fx)}} d\tan(e+fx) - \frac{2}{3}i(c-i\tan(e+fx))\right)\right)\right)\right)}{a^2f} \\
 & \downarrow 73 \\
 & \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{7/2}}{4c(1+i\tan(e+fx))^2} - \frac{1}{8}(3A+11iB) \left(\frac{i(c-ictan(e+fx))^{5/2}}{1+i\tan(e+fx)} - \frac{5}{2}c \left(2c \left(4i \int \frac{1}{2-\frac{c-ictan(e+fx)}{c}} d\sqrt{c-ictan(e+fx)} - \frac{2}{3}i(c-i\tan(e+fx))\right)\right)\right)\right)}{a^2f} \\
 & \downarrow 219 \\
 & \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{7/2}}{4c(1+i\tan(e+fx))^2} - \frac{1}{8}(3A+11iB) \left(\frac{i(c-ictan(e+fx))^{5/2}}{1+i\tan(e+fx)} - \frac{5}{2}c \left(2c \left(2i\sqrt{2}\sqrt{c}\operatorname{arctanh}\left(\frac{\sqrt{c-ictan(e+fx)}}{\sqrt{2}\sqrt{c}}\right) - \frac{2}{3}i(c-i\tan(e+fx))\right)\right)\right)\right)}{a^2f}
 \end{aligned}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2))/(a + I*a*Tan[e + f*x])^2, x]`

output `(c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(7/2))/(4*c*(1 + I*Tan[e + f*x])^2) - ((3*A + (11*I)*B)*((I*(c - I*c*Tan[e + f*x])^(5/2))/(1 + I*Tan[e + f*x]) - (5*c*(((-2*I)/3)*(c - I*c*Tan[e + f*x])^(3/2) + 2*c*((2*I)*Sqrt[2]*Sqrt[c]*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])]) - (2*I)*Sqrt[c - I*c*Tan[e + f*x]])))/2))/8)/(a^2*f)`

3.772. $\int \frac{(A+B \tan(e+fx))(c-ictan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^2} dx$

3.772.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 51 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`
- rule 60 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p_.)), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x],
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.772.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.75

method	result
derivativedivides	$2ic^2 \left(-\frac{iB(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} - 5i\sqrt{c-ic \tan(fx+e)} Bc - \sqrt{c-ic \tan(fx+e)} cA + 2c^2 \left(\frac{4\left(\frac{17iB}{32} + \frac{9A}{32}\right)(c-ic \tan(fx+e))^{\frac{3}{2}} + 4}{(c+ic \tan(fx+e))^{\frac{3}{2}} + 4} \right) \right) \frac{1}{fa^2}$
default	$2ic^2 \left(-\frac{iB(c-ic \tan(fx+e))^{\frac{3}{2}}}{3} - 5i\sqrt{c-ic \tan(fx+e)} Bc - \sqrt{c-ic \tan(fx+e)} cA + 2c^2 \left(\frac{4\left(\frac{17iB}{32} + \frac{9A}{32}\right)(c-ic \tan(fx+e))^{\frac{3}{2}} + 4}{(c+ic \tan(fx+e))^{\frac{3}{2}} + 4} \right) \right) \frac{1}{fa^2}$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^2,x,method=
_RETURNVERBOSE)
```

```
output 2*I/f/a^2*c^2*(-1/3*I*B*(c-I*c*tan(f*x+e))^(3/2)-5*I*(c-I*c*tan(f*x+e))^(1
/2)*B*c-(c-I*c*tan(f*x+e))^(1/2)*c*A+2*c^2*(4*((17/32*I*B+9/32*A)*(c-I*c*t
an(f*x+e))^(3/2)+(-15/16*I*B*c-7/16*c*A)*(c-I*c*tan(f*x+e))^(1/2)))/(c+I*c*
tan(f*x+e))^2+5/4*(11/4*I*B+3/4*A)*2^(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(
f*x+e))^(1/2)*2^(1/2)/c^(1/2)))
```

3.772.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 458 vs. 2(185) = 370.

Time = 0.27 (sec) , antiderivative size = 458, normalized size of antiderivative = 1.92

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2}}{(a + ia \tan(e + fx))^2} dx =$$

$$15 \sqrt{\frac{1}{2}} (a^2 f e^{(6i fx + 6i e)} + a^2 f e^{(4i fx + 4i e)}) \sqrt{-\frac{(9A^2 + 66i AB - 121B^2)c^7}{a^4 f^2}} \log \left(-\frac{5 \left((-3iA + 11B)c^4 + \sqrt{2} \sqrt{\frac{1}{2}} (a^2 f e^{(2i fx + 2i e)} + \dots \right)} \right)$$

3.772. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^2} dx$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^2,x
, algorithm="fricas")
```

```
output -1/12*(15*sqrt(1/2)*(a^2*f*e^(6*I*f*x + 6*I*e) + a^2*f*e^(4*I*f*x + 4*I*e)
)*sqrt(-(9*A^2 + 66*I*A*B - 121*B^2)*c^7/(a^4*f^2))*log(-5*((-3*I*A + 11*B
)*c^4 + sqrt(2)*sqrt(1/2)*(a^2*f*e^(2*I*f*x + 2*I*e) + a^2*f)*sqrt(-(9*A^2
+ 66*I*A*B - 121*B^2)*c^7/(a^4*f^2))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))e
^(-I*f*x - I*e)/(a^2*f)) - 15*sqrt(1/2)*(a^2*f*e^(6*I*f*x + 6*I*e) + a^2*f
*e^(4*I*f*x + 4*I*e))*sqrt(-(9*A^2 + 66*I*A*B - 121*B^2)*c^7/(a^4*f^2))*lo
g(-5*((-3*I*A + 11*B)*c^4 - sqrt(2)*sqrt(1/2)*(a^2*f*e^(2*I*f*x + 2*I*e) +
a^2*f)*sqrt(-(9*A^2 + 66*I*A*B - 121*B^2)*c^7/(a^4*f^2))*sqrt(c/(e^(2*I*f
*x + 2*I*e) + 1)))e^(-I*f*x - I*e)/(a^2*f)) + sqrt(2)*(15*(3*I*A - 11*B)*
c^3*e^(6*I*f*x + 6*I*e) + 20*(3*I*A - 11*B)*c^3*e^(4*I*f*x + 4*I*e) + 3*(3
*I*A - 11*B)*c^3*e^(2*I*f*x + 2*I*e) + 6*(-I*A + B)*c^3)*sqrt(c/(e^(2*I*f*
x + 2*I*e) + 1)))/(a^2*f*e^(6*I*f*x + 6*I*e) + a^2*f*e^(4*I*f*x + 4*I*e))
```

3.772.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2}}{(a + ia \tan(e + fx))^2} dx =$$

$$\int \frac{Ac^3 \sqrt{-ic \tan(e + fx) + c}}{\tan^2(e + fx) - 2i \tan(e + fx) - 1} dx + \int \left(-\frac{3Ac^3 \sqrt{-ic \tan(e + fx) + c} \tan^2(e + fx)}{\tan^2(e + fx) - 2i \tan(e + fx) - 1} \right) dx + \int \frac{Bc^3 \sqrt{-ic \tan(e + fx) + c} \tan(e + fx)}{\tan^2(e + fx) - 2i \tan(e + fx) - 1} dx +$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(7/2)/(a+I*a*tan(f*x+e))**2
,x)
```

```
output -(Integral(A*c**3*sqrt(-I*c*tan(e + f*x) + c)/(tan(e + f*x)**2 - 2*I*tan(e
+ f*x) - 1), x) + Integral(-3*A*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e +
f*x)**2/(tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x) + Integral(B*c**3*sqr
t(-I*c*tan(e + f*x) + c)*tan(e + f*x)/(tan(e + f*x)**2 - 2*I*tan(e + f*x)
- 1), x) + Integral(-3*B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3/
(tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x) + Integral(-3*I*A*c**3*sqrt(-
I*c*tan(e + f*x) + c)*tan(e + f*x)/(tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1
), x) + Integral(I*A*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3/(tan
(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x) + Integral(-3*I*B*c**3*sqrt(-I*c*
tan(e + f*x) + c)*tan(e + f*x)**2/(tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1)
, x) + Integral(I*B*c**3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4/(tan(
e + f*x)**2 - 2*I*tan(e + f*x) - 1), x))/a**2
```

3.772. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^2} dx$

3.772.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.93

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^2} dx =$$

$$i \left(\frac{15 \sqrt{2} (3A + 11iB) c^{9/2} \log\left(\frac{\sqrt{2}\sqrt{c} - \sqrt{-i c \tan(fx+e)+c}}{\sqrt{2}\sqrt{c} + \sqrt{-i c \tan(fx+e)+c}}\right)}{a^2} - \frac{12 \left((-i c \tan(fx+e)+c)^{3/2} (9A + 17iB) c^5 - 2 \sqrt{-i c \tan(fx+e)+c} (7A + 15iB) c^6 \right)}{(-i c \tan(fx+e)+c)^2 a^2 - 4(-i c \tan(fx+e)+c) a^2 c + 4 a^2 c^2} \right)$$

24cf

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^2,x
, algorithm="maxima")`

output `-1/24*I*(15*sqrt(2)*(3*A + 11*I*B)*c^(9/2)*log(-(sqrt(2)*sqrt(c) - sqrt(-I
*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a^2
- 12*((-I*c*tan(f*x + e) + c)^(3/2)*(9*A + 17*I*B)*c^5 - 2*sqrt(-I*c*tan(
f*x + e) + c)*(7*A + 15*I*B)*c^6)/((-I*c*tan(f*x + e) + c)^2*a^2 - 4*(-I*c
*tan(f*x + e) + c)*a^2*c + 4*a^2*c^2) + 16*(I*(-I*c*tan(f*x + e) + c)^(3/2
) * B*c^3 + 3*sqrt(-I*c*tan(f*x + e) + c)*(A + 5*I*B)*c^4)/a^2)/(c*f)`

3.772.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^2} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{7/2}}{(i a \tan(fx + e) + a)^2} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^2,x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(7/2)/(I*a*tan(f*x
+ e) + a)^2, x)`

3.772.9 Mupad [B] (verification not implemented)

Time = 8.91 (sec) , antiderivative size = 349, normalized size of antiderivative = 1.47

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^2} dx = \frac{15 B c^5 \sqrt{c - c \tan(e + fx) \operatorname{li}} - \frac{17 B c^4 (c - c \tan(e + fx) \operatorname{li})}{2}}{4 a^2 c^2 f + a^2 f (c - c \tan(e + fx) \operatorname{li})^2 - 4 a^2 c f (c - c \tan(e + fx) \operatorname{li})} - \frac{\frac{A c^5 \sqrt{c - c \tan(e + fx) \operatorname{li}} 7i}{a^2 f} - \frac{A c^4 (c - c \tan(e + fx) \operatorname{li})^{3/2} 9i}{2 a^2 f}}{(c - c \tan(e + fx) \operatorname{li})^2 - 4 c (c - c \tan(e + fx) \operatorname{li}) + 4 c^2} - \frac{A c^3 \sqrt{c - c \tan(e + fx) \operatorname{li}} 2i}{a^2 f} + \frac{10 B c^3 \sqrt{c - c \tan(e + fx) \operatorname{li}}}{a^2 f} + \frac{2 B c^2 (c - c \tan(e + fx) \operatorname{li})^{3/2}}{3 a^2 f} - \frac{\sqrt{2} A (-c)^{7/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) \operatorname{li}}}{2 \sqrt{-c}}\right) 15i}{4 a^2 f} + \frac{\sqrt{2} B c^{7/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) \operatorname{li}}}{2 \sqrt{c}}\right) 55i}{4 a^2 f}$$

```
input int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(7/2))/(a + a*tan(e + f*x)*1i)^2,x)
```

```
output (15*B*c^5*(c - c*tan(e + f*x)*1i)^(1/2) - (17*B*c^4*(c - c*tan(e + f*x)*1i)^(3/2))/2)/(4*a^2*c^2*f + a^2*f*(c - c*tan(e + f*x)*1i)^2 - 4*a^2*c*f*(c - c*tan(e + f*x)*1i)) - ((A*c^5*(c - c*tan(e + f*x)*1i)^(1/2)*7i)/(a^2*f) - (A*c^4*(c - c*tan(e + f*x)*1i)^(3/2)*9i)/(2*a^2*f))/((c - c*tan(e + f*x)*1i)^2 - 4*c*(c - c*tan(e + f*x)*1i) + 4*c^2) - (A*c^3*(c - c*tan(e + f*x)*1i)^(1/2)*2i)/(a^2*f) + (10*B*c^3*(c - c*tan(e + f*x)*1i)^(1/2))/(a^2*f) + (2*B*c^2*(c - c*tan(e + f*x)*1i)^(3/2))/(3*a^2*f) - (2^(1/2)*A*(-c)^(7/2)*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*(-c)^(1/2)))*15i)/(4*a^2*f) + (2^(1/2)*B*c^(7/2)*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2)*1i)/(2*c^(1/2)))*55i)/(4*a^2*f)
```


3.773
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^2} dx$$

3.773.1 Optimal result	7010
3.773.2 Mathematica [A] (verified)	7010
3.773.3 Rubi [A] (verified)	7011
3.773.4 Maple [A] (verified)	7014
3.773.5 Fricas [B] (verification not implemented)	7014
3.773.6 Sympy [F]	7015
3.773.7 Maxima [A] (verification not implemented)	7016
3.773.8 Giac [F]	7016
3.773.9 Mupad [B] (verification not implemented)	7017

3.773.1 Optimal result

Integrand size = 43, antiderivative size = 199

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^2} dx = \frac{3(iA - 9B)c^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{4\sqrt{2}a^2 f} - \frac{3(iA - 9B)c^2 \sqrt{c - ic \tan(e + fx)}}{8a^2 f} - \frac{(iA - 9B)c(c - ic \tan(e + fx))^{3/2}}{8a^2 f(1 + i \tan(e + fx))} + \frac{(iA - B)(c - ic \tan(e + fx))^{5/2}}{4a^2 f(1 + i \tan(e + fx))^2}$$

output `3/8*(I*A-9*B)*c^(5/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a^2/f*2^(1/2)-3/8*(I*A-9*B)*c^2*(c-I*c*tan(f*x+e))^(1/2)/a^2/f-1/8*(I*A-9*B)*c*(c-I*c*tan(f*x+e))^(3/2)/a^2/f/(1+I*tan(f*x+e))+1/4*(I*A-B)*(c-I*c*tan(f*x+e))^(5/2)/a^2/f/(1+I*tan(f*x+e))^2`

3.773.2 Mathematica [A] (verified)

Time = 6.20 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.84

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^2} dx = \frac{c^2 \sec^2(e + fx) \left(3\sqrt{2}(-iA + 9B)\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)\right)}{(a + ia \tan(e + fx))^2}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2))/(a + I*a*Tan[e + f*x])^2,x]`

3.773.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^2} dx$$

output $(c^2 \sec[e + f*x]^2 (3 \sqrt{2} ((-1)*A + 9*B) \sqrt{c} \operatorname{ArcTanh}[\sqrt{c - I*c \tan[e + f*x]}] / (\sqrt{2} \sqrt{c})) * (\cos[2*(e + f*x)] + I \sin[2*(e + f*x)]) + I*(A + (9*I)*B + (A + (25*I)*B) \cos[2*(e + f*x)] + ((5*I)*A - 29*B) \sin[2*(e + f*x)]) \sqrt{c - I*c \tan[e + f*x]}) / (8*a^2*f*(-1 + \tan[e + f*x])^2)$

3.773.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.87, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.186$, Rules used = {3042, 4071, 27, 87, 51, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - ic \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c - ic \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{a^3 (i \tan(e + fx) + 1)^3} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{c \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(i \tan(e + fx) + 1)^3} d \tan(e + fx)}{a^2 f} \\
 & \quad \downarrow \text{87} \\
 & \frac{c \left(\frac{(-B + iA)(c - ic \tan(e + fx))^{5/2}}{4c(1 + i \tan(e + fx))^2} - \frac{1}{8} (A + 9iB) \int \frac{(c - ic \tan(e + fx))^{3/2}}{(i \tan(e + fx) + 1)^2} d \tan(e + fx) \right)}{a^2 f} \\
 & \quad \downarrow \text{51} \\
 & \frac{c \left(\frac{(-B + iA)(c - ic \tan(e + fx))^{5/2}}{4c(1 + i \tan(e + fx))^2} - \frac{1}{8} (A + 9iB) \left(\frac{i(c - ic \tan(e + fx))^{3/2}}{1 + i \tan(e + fx)} - \frac{3}{2} c \int \frac{\sqrt{c - ic \tan(e + fx)}}{i \tan(e + fx) + 1} d \tan(e + fx) \right) \right)}{a^2 f} \\
 & \quad \downarrow \text{60}
 \end{aligned}$$

3.773. $\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^2} dx$

$$\frac{c \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{4c(1+i \tan(e+fx))^2} - \frac{1}{8}(A+9iB) \left(\frac{i(c-ic \tan(e+fx))^{3/2}}{1+i \tan(e+fx)} - \frac{3}{2}c \int \frac{1}{(i \tan(e+fx)+1)\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) \right) \right)}{a^2 f}$$

↓ 73

$$\frac{c \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{4c(1+i \tan(e+fx))^2} - \frac{1}{8}(A+9iB) \left(\frac{i(c-ic \tan(e+fx))^{3/2}}{1+i \tan(e+fx)} - \frac{3}{2}c \left(4i \int \frac{1}{2-\frac{c-ic \tan(e+fx)}{e}} d\sqrt{c-ic \tan(e+fx)} - 2i \sqrt{c-ic \tan(e+fx)} \right) \right) \right)}{a^2 f}$$

↓ 219

$$\frac{c \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{4c(1+i \tan(e+fx))^2} - \frac{1}{8}(A+9iB) \left(\frac{i(c-ic \tan(e+fx))^{3/2}}{1+i \tan(e+fx)} - \frac{3}{2}c \left(2i\sqrt{2}\sqrt{c} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}} \right) - 2i\sqrt{c-ic \tan(e+fx)} \right) \right) \right)}{a^2 f}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2))/(a + I*a*Tan[e + f*x])^2,x]`

output `(c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(5/2))/(4*c*(1 + I*Tan[e + f*x])^2) - ((A + (9*I)*B)*((I*(c - I*c*Tan[e + f*x])^(3/2))/(1 + I*Tan[e + f*x]) - (3*c*((2*I)*Sqrt[2]*Sqrt[c]*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])) - (2*I)*Sqrt[c - I*c*Tan[e + f*x]]))/2))/8)/(a^2*f)`

3.773.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

- rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.773.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.70

method	result
derivativedivides	$\frac{2ic^2 \left(-i\sqrt{c-ic \tan(fx+e)} B + c \left(\frac{4 \left(\frac{13iB}{32} + \frac{5A}{32} \right) (c-ic \tan(fx+e))^{\frac{3}{2}} + 4 \left(-\frac{11}{16} iBc - \frac{3}{16} cA \right) \sqrt{c-ic \tan(fx+e)} + \frac{3 \left(\frac{9iB}{4} + \frac{A}{4} \right) \sqrt{2}}{4} \right)}{(c+ic \tan(fx+e))^2} \right)}{fa^2}$
default	$\frac{2ic^2 \left(-i\sqrt{c-ic \tan(fx+e)} B + c \left(\frac{4 \left(\frac{13iB}{32} + \frac{5A}{32} \right) (c-ic \tan(fx+e))^{\frac{3}{2}} + 4 \left(-\frac{11}{16} iBc - \frac{3}{16} cA \right) \sqrt{c-ic \tan(fx+e)} + \frac{3 \left(\frac{9iB}{4} + \frac{A}{4} \right) \sqrt{2}}{4} \right)}{(c+ic \tan(fx+e))^2} \right)}{fa^2}$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output 2*I/f/a^2*c^2*(-I*(c-I*c*tan(f*x+e))^(1/2)*B+c*(4*((13/32*I*B+5/32*A)*(c-I*c*tan(f*x+e))^(3/2)+(-11/16*I*B*c-3/16*c*A)*(c-I*c*tan(f*x+e))^(1/2)))/(c+I*c*tan(f*x+e))^2+3/4*(9/4*I*B+1/4*A)*2^(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2)))
```

3.773.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 381 vs. 2(154) = 308.

Time = 0.27 (sec) , antiderivative size = 381, normalized size of antiderivative = 1.91

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^2} dx =$$

$$\left(3 \sqrt{\frac{1}{2}} a^2 f \sqrt{-\frac{(A^2 + 18iAB - 81B^2)c^5}{a^4 f^2}} e^{(4i fx + 4ie)} \log \left(-\frac{3 \left((-iA + 9B)c^3 + \sqrt{2} \sqrt{\frac{1}{2}} (a^2 f e^{(2i fx + 2ie)} + a^2 f) \sqrt{-\frac{(A^2 + 18iAB - 81B^2)c^5}{a^4 f^2}} \right)}{2 a^2 f} \right) \right)$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^2,x,algorithm="fracas")
```

3.773. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^2} dx$

output
$$\begin{aligned} & -1/8*(3*\sqrt{1/2}*a^{2*f}*\sqrt{-(A^2 + 18*I*A*B - 81*B^2)*c^5/(a^4*f^2)})*e^{(4*I*f*x + 4*I*e)}*\log(-3/2*((-I*A + 9*B)*c^3 + \sqrt{2}*\sqrt{1/2}*(a^{2*f}*e^{(2*I*f*x + 2*I*e)} + a^{2*f})*\sqrt{-(A^2 + 18*I*A*B - 81*B^2)*c^5/(a^4*f^2)})*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)})) * e^{-I*f*x - I*e}/(a^{2*f}) - 3*\sqrt{1/2}*a^{2*f}*\sqrt{-(A^2 + 18*I*A*B - 81*B^2)*c^5/(a^4*f^2)}*e^{(4*I*f*x + 4*I*e)}*\log(-3/2*((-I*A + 9*B)*c^3 - \sqrt{2}*\sqrt{1/2}*(a^{2*f}*e^{(2*I*f*x + 2*I*e)} + a^{2*f})*\sqrt{-(A^2 + 18*I*A*B - 81*B^2)*c^5/(a^4*f^2)})*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)})) * e^{-I*f*x - I*e}/(a^{2*f}) + \sqrt{2}*(3*(I*A - 9*B)*c^2*e^{(4*I*f*x + 4*I*e)} - (-I*A + 9*B)*c^2*e^{(2*I*f*x + 2*I*e)} + 2*(-I*A + B)*c^2)*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)})) * e^{(-4*I*f*x - 4*I*e)/(a^{2*f})} \end{aligned}$$

3.773.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(a + i a \tan(e + fx))^2} dx = \int \frac{A c^2 \sqrt{-i c \tan(e + fx) + c}}{\tan^2(e + fx) - 2i \tan(e + fx) - 1} dx + \int \left(-\frac{A c^2 \sqrt{-i c \tan(e + fx) + c} \tan^2(e + fx)}{\tan^2(e + fx) - 2i \tan(e + fx) - 1} \right) dx + \int \frac{B c^2 \sqrt{-i c \tan(e + fx) + c} \tan(e + fx)}{\tan^2(e + fx) - 2i \tan(e + fx) - 1} dx +$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(5/2)/(a+I*a*tan(f*x+e))**2,x)`

output
$$\begin{aligned} & -(Integral(A*c**2*\sqrt{-I*c*tan(e + f*x) + c}/(\tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x) + Integral(-A*c**2*\sqrt{-I*c*tan(e + f*x) + c}*\tan(e + f*x)**2/(\tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x) + Integral(B*c**2*\sqrt{-I*c*tan(e + f*x) + c}*\tan(e + f*x)/(\tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x) + Integral(-B*c**2*\sqrt{-I*c*tan(e + f*x) + c}*\tan(e + f*x)**3/(\tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x) + Integral(-2*I*A*c**2*\sqrt{-I*c*tan(e + f*x) + c}*\tan(e + f*x)/(\tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x) + Integral(-2*I*B*c**2*\sqrt{-I*c*tan(e + f*x) + c}*\tan(e + f*x)**2/(\tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x))/a**2 \end{aligned}$$

3.773.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.97

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(a + i a \tan(e + fx))^2} dx =$$

$$i \left(\frac{3\sqrt{2}(A+9iB)c^{7/2} \log\left(-\frac{\sqrt{2}\sqrt{c}-\sqrt{-ic\tan(fx+e)+c}}{\sqrt{2}\sqrt{c}+\sqrt{-ic\tan(fx+e)+c}}\right)}{a^2} + \frac{32i\sqrt{-ic\tan(fx+e)+c}Bc^3}{a^2} - \frac{4\left((-ic\tan(fx+e)+c\right)^{3/2}(5A+13iB)c^4-2\sqrt{-ic\tan(fx+e)+c}\right)}{(-ic\tan(fx+e)+c)^2a^2-4(-ic\tan(fx+e)+c)} \right)$$

$$16cf$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^2,x
, algorithm="maxima")`

output `-1/16*I*(3*sqrt(2)*(A + 9*I*B)*c^(7/2)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a^2 + 32*I*sqrt(-I*c*tan(f*x + e) + c)*B*c^3/a^2 - 4*((-I*c*tan(f*x + e) + c)^(3/2)*(5*A + 13*I*B)*c^4 - 2*sqrt(-I*c*tan(f*x + e) + c)*(3*A + 11*I*B)*c^5)/((-I*c*tan(f*x + e) + c)^2*a^2 - 4*(-I*c*tan(f*x + e) + c)*a^2*c + 4*a^2*c^2))/(c*f)`

3.773.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(a + i a \tan(e + fx))^2} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{5/2}}{(i a \tan(fx + e) + a)^2} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^2,x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(5/2)/(I*a*tan(f*x + e) + a)^2, x)`

3.773.9 Mupad [B] (verification not implemented)

Time = 8.72 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.48

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(a + i a \tan(e + fx))^2} dx = \frac{\frac{11 B c^4 \sqrt{c - c \tan(e + fx) i i}}{2} - \frac{13 B c^3 (c - c \tan(e + fx))}{4}}{4 a^2 c^2 f + a^2 f (c - c \tan(e + fx) i i)^2 - 4 a^2 c f (c - c \tan(e + fx) i i)^2 - 4 c (c - c \tan(e + fx) i i) + 4 c^2} + \frac{\frac{A c^4 \sqrt{c - c \tan(e + fx) i i} 3i}{2 a^2 f} - \frac{A c^3 (c - c \tan(e + fx) i i)^{3/2} 5i}{4 a^2 f}}{a^2 f} + \frac{2 B c^2 \sqrt{c - c \tan(e + fx) i i}}{a^2 f} + \frac{\sqrt{2} A (-c)^{5/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) i i}}{2 \sqrt{-c}}\right) 3i}{8 a^2 f} - \frac{27 \sqrt{2} B c^{5/2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) i i}}{2 \sqrt{c}}\right)}{8 a^2 f}$$

```
input int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(5/2))/(a + a*tan(e + f*x)*1i)^2,x)
```

```
output ((11*B*c^4*(c - c*tan(e + f*x)*1i)^(1/2))/2 - (13*B*c^3*(c - c*tan(e + f*x)*1i)^(3/2))/4)/(4*a^2*c^2*f + a^2*f*(c - c*tan(e + f*x)*1i)^2 - 4*a^2*c*f*(c - c*tan(e + f*x)*1i)) - ((A*c^4*(c - c*tan(e + f*x)*1i)^(1/2)*3i)/(2*a^2*f) - (A*c^3*(c - c*tan(e + f*x)*1i)^(3/2)*5i)/(4*a^2*f))/((c - c*tan(e + f*x)*1i)^2 - 4*c*(c - c*tan(e + f*x)*1i) + 4*c^2) + (2*B*c^2*(c - c*tan(e + f*x)*1i)^(1/2))/(a^2*f) + (2^(1/2)*A*(-c)^(5/2)*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*(-c)^(1/2)))*3i)/(8*a^2*f) - (27*2^(1/2)*B*c^(5/2)*atanh((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*c^(1/2))))/(8*a^2*f)
```


3.774
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^2} dx$$

3.774.1 Optimal result	7018
3.774.2 Mathematica [A] (verified)	7018
3.774.3 Rubi [A] (verified)	7019
3.774.4 Maple [A] (verified)	7021
3.774.5 Fricas [B] (verification not implemented)	7022
3.774.6 Sympy [F]	7022
3.774.7 Maxima [A] (verification not implemented)	7023
3.774.8 Giac [F]	7023
3.774.9 Mupad [B] (verification not implemented)	7024

3.774.1 Optimal result

Integrand size = 43, antiderivative size = 160

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^2} dx =$$

$$\frac{(iA + 7B)c^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{8\sqrt{2}a^2 f} + \frac{(iA + 7B)c\sqrt{c - ic \tan(e + fx)}}{8a^2 f(1 + i \tan(e + fx))} + \frac{(iA - B)(c - ic \tan(e + fx))^{3/2}}{4a^2 f(1 + i \tan(e + fx))^2}$$

output

```
-1/16*(I*A+7*B)*c^(3/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a^2/f*2^(1/2)+1/8*(I*A+7*B)*c*(c-I*c*tan(f*x+e))^(1/2)/a^2/f/(1+I*tan(f*x+e))+1/4*(I*A-B)*(c-I*c*tan(f*x+e))^(3/2)/a^2/f/(1+I*tan(f*x+e))^2
```

3.774.2 Mathematica [A] (verified)

Time = 6.22 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.90

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^2} dx = \frac{\sqrt{2}(iA + 7B)c^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right) \sec^2(e + fx)}{(a + ia \tan(e + fx))^2}$$

input

```
Integrate[(((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2))/(a + I*a*Tan[e + f*x])^2,x]
```

output $(\text{Sqrt}[2]*(I*A + 7*B)*c^{(3/2)}*\text{ArcTanh}[\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]]]/(\text{Sqrt}[2]*\text{Sqrt}[c]))*\text{Sec}[e + f*x]^2*(\text{Cos}[2*(e + f*x)] + I*\text{Sin}[2*(e + f*x)]) - 2*c*((3*I)*A + 5*B + (A + (9*I)*B)*\text{Tan}[e + f*x])* \text{Sqrt}[c - I*c*\text{Tan}[e + f*x]]/(16*a^2*f*(-I + \text{Tan}[e + f*x])^2)$

3.774.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.91, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.163$, Rules used = {3042, 4071, 27, 87, 51, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx$$

↓ 3042

$$\int \frac{(c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx$$

↓ 4071

$$\frac{ac \int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{a^3 (i \tan(e + fx) + 1)^3} d \tan(e + fx)}{f}$$

↓ 27

$$\frac{c \int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{(i \tan(e + fx) + 1)^3} d \tan(e + fx)}{a^2 f}$$

↓ 87

$$\frac{c \left(\frac{1}{8} (A - 7iB) \int \frac{\sqrt{c - ic \tan(e + fx)}}{(i \tan(e + fx) + 1)^2} d \tan(e + fx) + \frac{(-B + iA)(c - ic \tan(e + fx))^{3/2}}{4c(1 + i \tan(e + fx))^2} \right)}{a^2 f}$$

↓ 51

$$\frac{c \left(\frac{1}{8} (A - 7iB) \left(\frac{i \sqrt{c - ic \tan(e + fx)}}{1 + i \tan(e + fx)} - \frac{1}{2} c \int \frac{1}{(i \tan(e + fx) + 1) \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) \right) + \frac{(-B + iA)(c - ic \tan(e + fx))^{3/2}}{4c(1 + i \tan(e + fx))^2} \right)}{a^2 f}$$

↓ 73

3.774. $\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^2} dx$

$$\frac{c \left(\frac{1}{8}(A - 7iB) \left(\frac{i\sqrt{c-ic \tan(e+fx)}}{1+i \tan(e+fx)} - i \int \frac{1}{2-c-ic \tan(e+fx)} d\sqrt{c-ic \tan(e+fx)} \right) + \frac{(-B+iA)(c-ic \tan(e+fx))^{3/2}}{4c(1+i \tan(e+fx))^2} \right)}{a^2 f}$$

↓ 219

$$\frac{c \left(\frac{1}{8}(A - 7iB) \left(\frac{i\sqrt{c-ic \tan(e+fx)}}{1+i \tan(e+fx)} - \frac{i\sqrt{c} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}} \right)}{\sqrt{2}} \right) + \frac{(-B+iA)(c-ic \tan(e+fx))^{3/2}}{4c(1+i \tan(e+fx))^2} \right)}{a^2 f}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2))/(a + I*a*Tan[e + f*x])^2,x]`

output `(c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(3/2))/(4*c*(1 + I*Tan[e + f*x])^2) + ((A - (7*I)*B)*(((- I)*Sqrt[c]*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])])]/Sqrt[2] + (I*Sqrt[c - I*c*Tan[e + f*x]])/(1 + I*Tan[e + f*x]))/8))/(a^2*f)`

3.774.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.774.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.74

method	result
derivativedivides	$2ic^2 \frac{\left(\frac{4\left(\frac{9iB}{64} + \frac{A}{64}\right)(c-ic \tan(fx+e))^{\frac{3}{2}} + 4\left(-\frac{7}{32}iBc + \frac{1}{32}cA\right)\sqrt{c-ic \tan(fx+e)}}{(c+ic \tan(fx+e))^2} - \frac{\left(-\frac{7iB}{4} + \frac{A}{4}\right)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)}\sqrt{2}}{2\sqrt{c}}\right)}{8\sqrt{c}} \right)}{fa^2}$
default	$2ic^2 \frac{\left(\frac{4\left(\frac{9iB}{64} + \frac{A}{64}\right)(c-ic \tan(fx+e))^{\frac{3}{2}} + 4\left(-\frac{7}{32}iBc + \frac{1}{32}cA\right)\sqrt{c-ic \tan(fx+e)}}{(c+ic \tan(fx+e))^2} - \frac{\left(-\frac{7iB}{4} + \frac{A}{4}\right)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)}\sqrt{2}}{2\sqrt{c}}\right)}{8\sqrt{c}} \right)}{fa^2}$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^2,x, metho
d=_RETURNVERBOSE)
```

$$3.774. \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^2} dx$$

output $2*I/f/a^2*c^2*(4*((9/64*I*B+1/64*A)*(c-I*c*\tan(f*x+e))^{3/2}+(-7/32*I*B*c+1/32*c*A)*(c-I*c*\tan(f*x+e))^{1/2}))/((c+I*c*\tan(f*x+e))^2-1/8*(-7/4*I*B+1/4*A)*2^{1/2}/c^{1/2})*\operatorname{arctanh}(1/2*(c-I*c*\tan(f*x+e))^{1/2}*2^{1/2}/c^{1/2}))$

3.774.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 372 vs. $2(123) = 246$.

Time = 0.27 (sec) , antiderivative size = 372, normalized size of antiderivative = 2.32

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^2} dx = \left(\sqrt{\frac{1}{2}} a^2 f \sqrt{-\frac{(A^2 - 14i AB - 49 B^2)c^3}{a^4 f^2}} e^{(4i fx + 4i e)} \log \left(\frac{(-i A - \dots)}{\dots} \right) \right)$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^2,x, algorithm="fracas")`

output $1/16*(\operatorname{sqrt}(1/2)*a^2*f*\operatorname{sqrt}(-(A^2 - 14*I*A*B - 49*B^2)*c^3/(a^4*f^2)))*e^{(4*I*f*x + 4*I*e)}*\log(1/4*((-I*A - 7*B)*c^2 + \operatorname{sqrt}(2)*\operatorname{sqrt}(1/2)*(a^2*f*e^{(2*I*f*x + 2*I*e)} + a^2*f)*\operatorname{sqrt}(c/(e^{(2*I*f*x + 2*I*e)} + 1))*\operatorname{sqrt}(-(A^2 - 14*I*A*B - 49*B^2)*c^3/(a^4*f^2))))*e^{(-I*f*x - I*e)/(a^2*f)} - \operatorname{sqrt}(1/2)*a^2*f*\operatorname{sqrt}(-(A^2 - 14*I*A*B - 49*B^2)*c^3/(a^4*f^2)))*e^{(4*I*f*x + 4*I*e)}*\log(1/4*((-I*A - 7*B)*c^2 - \operatorname{sqrt}(2)*\operatorname{sqrt}(1/2)*(a^2*f*e^{(2*I*f*x + 2*I*e)} + a^2*f)*\operatorname{sqrt}(c/(e^{(2*I*f*x + 2*I*e)} + 1))*\operatorname{sqrt}(-(A^2 - 14*I*A*B - 49*B^2)*c^3/(a^4*f^2))))*e^{(-I*f*x - I*e)/(a^2*f)} + \operatorname{sqrt}(2)*((I*A + 7*B)*c*e^{(4*I*f*x + 4*I*e)} + (3*I*A + 5*B)*c*e^{(2*I*f*x + 2*I*e)} - 2*(-I*A + B)*c)*\operatorname{sqrt}(c/(e^{(2*I*f*x + 2*I*e)} + 1)))*e^{(-4*I*f*x - 4*I*e)/(a^2*f)}$

3.774.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^2} dx = \frac{\int \frac{Ac\sqrt{-i c \tan(e+fx)+c}}{\tan^2(e+fx)-2i \tan(e+fx)-1} dx + \int \frac{Bc\sqrt{-i c \tan(e+fx)+c \tan(e+fx)}}{\tan^2(e+fx)-2i \tan(e+fx)-1} dx + \int \left(-\frac{iAc\sqrt{-i c \tan(e+fx)+c \tan(e+fx)}}{\tan^2(e+fx)-2i \tan(e+fx)-1} \right) dx}{a^2}$$

3.774. $\int \frac{(A+B \tan(e+fx))(c-i c \tan(e+fx))^{3/2}}{(a+i a \tan(e+fx))^2} dx$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(3/2)/(a+I*a*tan(f*x+e))**2,x)`

output `-(Integral(A*c*sqrt(-I*c*tan(e + f*x) + c)/(tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x) + Integral(B*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)/(tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x) + Integral(-I*A*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)/(tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x) + Integral(-I*B*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2/(tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x))/a**2`

3.774.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.04

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^2} dx = \frac{i \left(\frac{\sqrt{2}(A - 7iB)c^{5/2} \log\left(-\frac{\sqrt{2}\sqrt{c} - \sqrt{-i c \tan(fx+e)+c}}{\sqrt{2}\sqrt{c} + \sqrt{-i c \tan(fx+e)+c}}\right)}{a^2} + \frac{4((-i c \tan(fx+e) + c))^{3/2}}{(-i c \tan(fx+e) + c)} \right)}{32 c f}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output `1/32*I*(sqrt(2)*(A - 7*I*B)*c^(5/2)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a^2 + 4*((-I*c*tan(f*x + e) + c)^(3/2)*(A + 9*I*B)*c^3 + 2*sqrt(-I*c*tan(f*x + e) + c)*(A - 7*I*B)*c^4)/((-I*c*tan(f*x + e) + c)^2*a^2 - 4*(-I*c*tan(f*x + e) + c)*a^2*c + 4*a^2*c^2)/(c*f)`

3.774.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^2} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{3/2}}{(i a \tan(fx + e) + a)^2} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(3/2)/(I*a*tan(f*x + e) + a)^2, x)`

3.774.9 Mupad [B] (verification not implemented)

Time = 8.72 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.67

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^2} dx = \frac{\frac{7 B c^3 \sqrt{c - c \tan(e + fx) i} \operatorname{li}}{4} - \frac{9 B c^2 (c - c \tan(e + fx) i)}{8}}{4 a^2 c^2 f + a^2 f (c - c \tan(e + fx) i)^2 - 4 a^2 c f (c - c \tan(e + fx) i)} + \frac{\frac{A c^3 \sqrt{c - c \tan(e + fx) i} \operatorname{li}}{4 a^2 f} + \frac{A c^2 (c - c \tan(e + fx) i)^{3/2} \operatorname{li}}{8 a^2 f}}{(c - c \tan(e + fx) i)^2 - 4 c (c - c \tan(e + fx) i) + 4 c^2} + \frac{\sqrt{2} A (-c)^{3/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) i}}{2 \sqrt{-c}}\right) \operatorname{li}}{16 a^2 f} - \frac{7 \sqrt{2} B c^{3/2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) i}}{2 \sqrt{c}}\right)}{16 a^2 f}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(3/2))/(a + a*tan(e + f*x)*1i)^2,x)`

output `((7*B*c^3*(c - c*tan(e + f*x)*1i)^(1/2))/4 - (9*B*c^2*(c - c*tan(e + f*x)*1i)^(3/2))/8)/(4*a^2*c^2*f + a^2*f*(c - c*tan(e + f*x)*1i)^2 - 4*a^2*c*f*(c - c*tan(e + f*x)*1i)) + ((A*c^3*(c - c*tan(e + f*x)*1i)^(1/2)*1i)/(4*a^2*f) + (A*c^2*(c - c*tan(e + f*x)*1i)^(3/2)*1i)/(8*a^2*f))/((c - c*tan(e + f*x)*1i)^2 - 4*c*(c - c*tan(e + f*x)*1i) + 4*c^2) + (2^(1/2)*A*(-c)^(3/2)*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*(-c)^(1/2)))*1i)/(16*a^2*f) - (7*2^(1/2)*B*c^(3/2)*atanh((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*c^(1/2)))/16*a^2*f)`

$$3.775 \quad \int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{(a+ia \tan(e+fx))^2} dx$$

3.775.1 Optimal result	7025
3.775.2 Mathematica [A] (verified)	7025
3.775.3 Rubi [A] (verified)	7026
3.775.4 Maple [A] (verified)	7028
3.775.5 Fracas [B] (verification not implemented)	7029
3.775.6 Sympy [F]	7030
3.775.7 Maxima [A] (verification not implemented)	7030
3.775.8 Giac [F]	7031
3.775.9 Mupad [B] (verification not implemented)	7031

3.775.1 Optimal result

Integrand size = 43, antiderivative size = 159

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))\sqrt{c - ic \tan(e + fx)}}{(a + ia \tan(e + fx))^2} dx \\ &= \frac{(3iA + 5B)\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{16\sqrt{2}a^2 f} \\ & \quad + \frac{(iA - B)\sqrt{c - ic \tan(e + fx)}}{4a^2 f(1 + i \tan(e + fx))^2} + \frac{(3iA + 5B)\sqrt{c - ic \tan(e + fx)}}{16a^2 f(1 + i \tan(e + fx))} \end{aligned}$$

output `1/32*(3*I*A+5*B)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))*c^(1/2)/a^2/f*2^(1/2)+1/4*(I*A-B)*(c-I*c*tan(f*x+e))^(1/2)/a^2/f/(1+I*tan(f*x+e))^2+1/16*(3*I*A+5*B)*(c-I*c*tan(f*x+e))^(1/2)/a^2/f/(1+I*tan(f*x+e))`

3.775.2 Mathematica [A] (verified)

Time = 5.53 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.91

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))\sqrt{c - ic \tan(e + fx)}}{(a + ia \tan(e + fx))^2} dx \\ &= \frac{\sqrt{2}(3A - 5iB)\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right) \sec^2(e + fx)(-i \cos(2(e + fx)) + \sin(2(e + fx))) + 2(-7iA - 32a^2 f(-i + \tan(e + fx))^2)}{32a^2 f(-i + \tan(e + fx))^2} \end{aligned}$$

3.775. $\int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{(a+ia \tan(e+fx))^2} dx$

input `Integrate[((A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]/(a + I*a*Tan[e + f*x])^2,x]`

output `(Sqrt[2]*(3*A - (5*I)*B)*Sqrt[c]*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])]*Sec[e + f*x]^2*((-I)*Cos[2*(e + f*x)] + Sin[2*(e + f*x)]) + 2*((-7*I)*A - B + (3*A - (5*I)*B)*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]/(32*a^2*f*(-I + Tan[e + f*x])^2)`

3.775.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.97, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.163$, Rules used = {3042, 4071, 27, 87, 52, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{c - ic \tan(e + fx)}(A + B \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{\sqrt{c - ic \tan(e + fx)}(A + B \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx \\
 & \quad \downarrow 4071 \\
 & \frac{ac \int \frac{A + B \tan(e + fx)}{a^3(i \tan(e + fx) + 1)^3 \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{f} \\
 & \quad \downarrow 27 \\
 & \frac{c \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx) + 1)^3 \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{a^2 f} \\
 & \quad \downarrow 87 \\
 & \frac{c \left(\frac{1}{8}(3A - 5iB) \int \frac{1}{(i \tan(e + fx) + 1)^2 \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) + \frac{(-B + iA) \sqrt{c - ic \tan(e + fx)}}{4c(1 + i \tan(e + fx))^2} \right)}{a^2 f} \\
 & \quad \downarrow 52 \\
 & \frac{c \left(\frac{1}{8}(3A - 5iB) \left(\frac{1}{4} \int \frac{1}{(i \tan(e + fx) + 1) \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) + \frac{i \sqrt{c - ic \tan(e + fx)}}{2c(1 + i \tan(e + fx))} \right) + \frac{(-B + iA) \sqrt{c - ic \tan(e + fx)}}{4c(1 + i \tan(e + fx))^2} \right)}{a^2 f}
 \end{aligned}$$

3.775. $\int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{(a + ia \tan(e + fx))^2} dx$

$$\begin{aligned}
 & \downarrow 73 \\
 & \frac{c \left(\frac{1}{8}(3A - 5iB) \left(\frac{i \int \frac{1}{2 - c - i c \tan(e+fx)} d\sqrt{c - i c \tan(e+fx)}}{2c} + \frac{i\sqrt{c - i c \tan(e+fx)}}{2c(1+i \tan(e+fx))} \right) + \frac{(-B+iA)\sqrt{c - i c \tan(e+fx)}}{4c(1+i \tan(e+fx))^2} \right)}{a^2 f} \\
 & \downarrow 219 \\
 & \frac{c \left(\frac{1}{8}(3A - 5iB) \left(\frac{i \operatorname{arctanh}\left(\frac{\sqrt{c - i c \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{2\sqrt{2}\sqrt{c}} + \frac{i\sqrt{c - i c \tan(e+fx)}}{2c(1+i \tan(e+fx))} \right) + \frac{(-B+iA)\sqrt{c - i c \tan(e+fx)}}{4c(1+i \tan(e+fx))^2} \right)}{a^2 f}
 \end{aligned}$$

input `Int[((A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]/(a + I*a*Tan[e + f*x]))^2,x]`

output `(c*(((I*A - B)*Sqrt[c - I*c*Tan[e + f*x]]/(4*c*(1 + I*Tan[e + f*x])^2) + ((3*A - (5*I)*B)*(((I/2)*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])]))/(Sqrt[2]*Sqrt[c]) + ((I/2)*Sqrt[c - I*c*Tan[e + f*x]]/(c*(1 + I*Tan[e + f*x]))))/8))/(a^2*f)`

3.775.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 52 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 73 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.775.4 Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.75

method	result
derivativedivides	$\frac{2ic^2 \left(\frac{(-5iB+3A)(c-ic \tan(fx+e))^{\frac{3}{2}} + 4\left(\frac{5A}{64} - \frac{3iB}{64}\right)\sqrt{c-ic \tan(fx+e)}}{(c+ic \tan(fx+e))^2} + \frac{(-5iB+3A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)}\sqrt{2}}{2\sqrt{c}}\right)}{64c^{\frac{3}{2}}} \right)}{fa^2}$
default	$\frac{2ic^2 \left(\frac{(-5iB+3A)(c-ic \tan(fx+e))^{\frac{3}{2}} + 4\left(\frac{5A}{64} - \frac{3iB}{64}\right)\sqrt{c-ic \tan(fx+e)}}{(c+ic \tan(fx+e))^2} + \frac{(-5iB+3A)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(fx+e)}\sqrt{2}}{2\sqrt{c}}\right)}{64c^{\frac{3}{2}}} \right)}{fa^2}$

```
input int((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x, metho
d=_RETURNVERBOSE)
```

$$3.775. \int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{(a+ia \tan(e+fx))^2} dx$$

output $2*I/f/a^2*c^2*(4*(-1/128/c*(3*A-5*I*B)*(c-I*c*\tan(f*x+e))^{(3/2)}+(5/64*A-3/64*I*B)*(c-I*c*\tan(f*x+e))^{(1/2)})/(c+I*c*\tan(f*x+e))^{2+1/64}/c^{(3/2)}*(3*A-5*I*B)*2^{(1/2)}*\operatorname{arctanh}(1/2*(c-I*c*\tan(f*x+e))^{(1/2)}*2^{(1/2)}/c^{(1/2)})$

3.775.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 362 vs. $2(122) = 244$.

Time = 0.28 (sec) , antiderivative size = 362, normalized size of antiderivative = 2.28

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{(a + i a \tan(e + fx))^2} dx$$

$$= \left(\sqrt{\frac{1}{2}} a^2 f \sqrt{-\frac{(9A^2 - 30iAB - 25B^2)c}{a^4 f^2}} e^{(4i f x + 4i e)} \log \left(\frac{\left(\sqrt{2} \sqrt{\frac{1}{2}} (a^2 f e^{(2i f x + 2i e)} + a^2 f) \sqrt{\frac{c}{e^{(2i f x + 2i e)} + 1}} \sqrt{-\frac{(9A^2 - 30iAB - 25B^2)c}{a^4 f^2}} \right)}{8 a^2 f} \right) \right)$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x, algorithm="fracas")`

output $1/32*(\operatorname{sqrt}(1/2)*a^2*f*\operatorname{sqrt}(-(9*A^2 - 30*I*A*B - 25*B^2)*c/(a^4*f^2)))*e^{(4*I*f*x + 4*I*e)}*\log(1/8*(\operatorname{sqrt}(2)*\operatorname{sqrt}(1/2)*(a^2*f*e^{(2*I*f*x + 2*I*e)} + a^2*f)*\operatorname{sqrt}(c/(e^{(2*I*f*x + 2*I*e)} + 1)))*\operatorname{sqrt}(-(9*A^2 - 30*I*A*B - 25*B^2)*c/(a^4*f^2)) + (3*I*A + 5*B)*c)*e^{(-I*f*x - I*e)/(a^2*f)} - \operatorname{sqrt}(1/2)*a^2*f*\operatorname{sqrt}(-(9*A^2 - 30*I*A*B - 25*B^2)*c/(a^4*f^2))*e^{(4*I*f*x + 4*I*e)}*\log(-1/8*(\operatorname{sqrt}(2)*\operatorname{sqrt}(1/2)*(a^2*f*e^{(2*I*f*x + 2*I*e)} + a^2*f)*\operatorname{sqrt}(c/(e^{(2*I*f*x + 2*I*e)} + 1)))*\operatorname{sqrt}(-(9*A^2 - 30*I*A*B - 25*B^2)*c/(a^4*f^2)) - (3*I*A + 5*B)*c)*e^{(-I*f*x - I*e)/(a^2*f)} + \operatorname{sqrt}(2)*((5*I*A + 3*B)*e^{(4*I*f*x + 4*I*e)} + (7*I*A + B)*e^{(2*I*f*x + 2*I*e)} + 2*I*A - 2*B)*\operatorname{sqrt}(c/(e^{(2*I*f*x + 2*I*e)} + 1)))*e^{(-4*I*f*x - 4*I*e)/(a^2*f)}$

3.775.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{(a + ia \tan(e + fx))^2} dx$$

$$= - \frac{\int \frac{A \sqrt{-ic \tan(e+fx)+c}}{\tan^2(e+fx)-2i \tan(e+fx)-1} dx + \int \frac{B \sqrt{-ic \tan(e+fx)+c} \tan(e+fx)}{\tan^2(e+fx)-2i \tan(e+fx)-1} dx}{a^2}$$

input `integrate((c-I*c*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**2, x)`

output `-(Integral(A*sqrt(-I*c*tan(e + f*x) + c)/(tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x) + Integral(B*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)/(tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x))/a**2`

3.775.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.08

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{(a + ia \tan(e + fx))^2} dx =$$

$$i \left(\frac{\sqrt{2}(3A - 5iB)c^{\frac{3}{2}} \log\left(-\frac{\sqrt{2}\sqrt{c} - \sqrt{-ic \tan(fx+e)+c}}{\sqrt{2}\sqrt{c} + \sqrt{-ic \tan(fx+e)+c}}\right)}{a^2} + \frac{4 \left((-ic \tan(fx+e)+c)^{\frac{3}{2}} (3A - 5iB)c^2 - 2\sqrt{-ic \tan(fx+e)+c} (5A - 3iB)c^3 \right)}{(-ic \tan(fx+e)+c)^2 a^2 - 4(-ic \tan(fx+e)+c)a^2 c + 4a^2 c^2} \right)$$

$$= \frac{\hspace{15em}}{64cf}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2, x, algorithm="maxima")`

output `-1/64*I*(sqrt(2)*(3*A - 5*I*B)*c^(3/2)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a^2 + 4*((-I*c*tan(f*x + e) + c)^(3/2)*(3*A - 5*I*B)*c^2 - 2*sqrt(-I*c*tan(f*x + e) + c)*(5*A - 3*I*B)*c^3)/((-I*c*tan(f*x + e) + c)^2*a^2 - 4*(-I*c*tan(f*x + e) + c)*a^2*c + 4*a^2*c^2)/(c*f)`

3.775.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{(a + i a \tan(e + fx))^2} dx$$

$$= \int \frac{(B \tan(fx + e) + A) \sqrt{-i c \tan(fx + e) + c}}{(i a \tan(fx + e) + a)^2} dx$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*sqrt(-I*c*tan(f*x + e) + c)/(I*a*tan(f*x + e) + a)^2, x)`

3.775.9 Mupad [B] (verification not implemented)

Time = 8.83 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.66

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{(a + i a \tan(e + fx))^2} dx$$

$$= -\frac{\frac{5 B c (c - c \tan(e + f x) i i)^{3/2}}{16} - \frac{3 B c^2 \sqrt{c - c \tan(e + f x) i i}}{8}}{4 a^2 c^2 f + a^2 f (c - c \tan(e + f x) i i)^2 - 4 a^2 c f (c - c \tan(e + f x) i i)}$$

$$+ \frac{\frac{A c^2 \sqrt{c - c \tan(e + f x) i i} 5 i}{8 a^2 f} - \frac{A c (c - c \tan(e + f x) i i)^{3/2} 3 i}{16 a^2 f}}{(c - c \tan(e + f x) i i)^2 - 4 c (c - c \tan(e + f x) i i) + 4 c^2}$$

$$+ \frac{\sqrt{2} A \sqrt{-c} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + f x) i i}}{2 \sqrt{-c}}\right) 3 i}{32 a^2 f} + \frac{5 \sqrt{2} B \sqrt{c} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + f x) i i}}{2 \sqrt{c}}\right)}{32 a^2 f}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(1/2))/(a + a*tan(e + f*x)*1i)^2,x)`

output `((A*c^2*(c - c*tan(e + f*x)*1i)^(1/2)*5i)/(8*a^2*f) - (A*c*(c - c*tan(e + f*x)*1i)^(3/2)*3i)/(16*a^2*f))/((c - c*tan(e + f*x)*1i)^2 - 4*c*(c - c*tan(e + f*x)*1i) + 4*c^2) - ((5*B*c*(c - c*tan(e + f*x)*1i)^(3/2))/16 - (3*B*c^2*(c - c*tan(e + f*x)*1i)^(1/2))/8)/(4*a^2*c^2*f + a^2*f*(c - c*tan(e + f*x)*1i)^2 - 4*a^2*c*f*(c - c*tan(e + f*x)*1i)) + (2^(1/2)*A*(-c)^(1/2)*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*(-c)^(1/2)))*3i)/(32*a^2*f) + (5*2^(1/2)*B*c^(1/2)*atanh((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*c^(1/2)))/(32*a^2*f)`

3.775. $\int \frac{(A+B \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}}{(a+ia \tan(e+fx))^2} dx$

3.776
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 \sqrt{c-ic \tan(e+fx)}} dx$$

3.776.1 Optimal result 7032
 3.776.2 Mathematica [A] (verified) 7033
 3.776.3 Rubi [A] (verified) 7033
 3.776.4 Maple [A] (verified) 7036
 3.776.5 Fricas [B] (verification not implemented) 7037
 3.776.6 Sympy [F] 7037
 3.776.7 Maxima [A] (verification not implemented) 7038
 3.776.8 Giac [F] 7038
 3.776.9 Mupad [B] (verification not implemented) 7039

3.776.1 Optimal result

Integrand size = 43, antiderivative size = 195

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 \sqrt{c - ic \tan(e + fx)}} dx$$

$$= \frac{3(5iA + 3B) \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{32\sqrt{2}a^2\sqrt{c}f} - \frac{3(5iA + 3B)}{32a^2 f \sqrt{c - ic \tan(e + fx)}}$$

$$+ \frac{iA - B}{4a^2 f (1 + i \tan(e + fx))^2 \sqrt{c - ic \tan(e + fx)}}$$

$$+ \frac{5iA + 3B}{16a^2 f (1 + i \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}$$

```
output 3/64*(5*I*A+3*B)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a^2
/f*2^(1/2)/c^(1/2)-3/32*(5*I*A+3*B)/a^2/f/(c-I*c*tan(f*x+e))^(1/2)+1/4*(I*
A-B)/a^2/f/(c-I*c*tan(f*x+e))^(1/2)/(1+I*tan(f*x+e))^2+1/16*(5*I*A+3*B)/a^
2/f/(c-I*c*tan(f*x+e))^(1/2)/(1+I*tan(f*x+e))
```

3.776.2 Mathematica [A] (verified)

Time = 6.24 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.78

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 \sqrt{c - ic \tan(e + fx)}} dx$$

$$= \frac{3\sqrt{2}a(5iA+3B)\operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{\sqrt{c}} + \frac{2a\sqrt{c-ic \tan(e+fx)}(3A+11iB+(-20iA-12B)\tan(e+fx)+3(5A-3iB)\tan^2(e+fx))}{c(-i+\tan(e+fx))^2(i+\tan(e+fx))}$$

$$64a^3f$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*Sqrt[c - I*c*Tan[e + f*x]]),x]`

output `((3*Sqrt[2]*a*((5*I)*A + 3*B)*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])])/Sqrt[c] + (2*a*Sqrt[c - I*c*Tan[e + f*x]]*(3*A + (11*I)*B + ((-20*I)*A - 12*B)*Tan[e + f*x] + 3*(5*A - (3*I)*B)*Tan[e + f*x]^2))/(c*(-I + Tan[e + f*x])^2*(I + Tan[e + f*x]))/(64*a^3*f)`

3.776.3 Rubi [A] (verified)Time = 0.41 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.93, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.186$, Rules used = {3042, 4071, 27, 87, 52, 61, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 \sqrt{c - ic \tan(e + fx)}} dx$$

$$\downarrow 3042$$

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 \sqrt{c - ic \tan(e + fx)}} dx$$

$$\downarrow 4071$$

$$ac \int \frac{A+B \tan(e+fx)}{a^3(i \tan(e+fx)+1)^3(c-ic \tan(e+fx))^{3/2}} d \tan(e + fx)$$

$$\downarrow 27$$

3.776. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 \sqrt{c-ic \tan(e+fx)}} dx$

$$\begin{aligned}
& \frac{c \int \frac{A+B \tan(e+fx)}{(i \tan(e+fx)+1)^3 (c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{a^2 f} \\
& \quad \downarrow 87 \\
& \frac{c \left(\frac{1}{8} (5A - 3iB) \int \frac{1}{(i \tan(e+fx)+1)^2 (c-ic \tan(e+fx))^{3/2}} d \tan(e+fx) + \frac{-B+iA}{4c(1+i \tan(e+fx))^2 \sqrt{c-ic \tan(e+fx)}} \right)}{a^2 f} \\
& \quad \downarrow 52 \\
& \frac{c \left(\frac{1}{8} (5A - 3iB) \left(\frac{3}{4} \int \frac{1}{(i \tan(e+fx)+1)(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx) + \frac{i}{2c(1+i \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}} \right) + \frac{1}{4c(1+i \tan(e+fx))} \right)}{a^2 f} \\
& \quad \downarrow 61 \\
& \frac{c \left(\frac{1}{8} (5A - 3iB) \left(\frac{3}{4} \left(\frac{\int \frac{1}{(i \tan(e+fx)+1) \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{2c} - \frac{i}{c \sqrt{c-ic \tan(e+fx)}} \right) + \frac{i}{2c(1+i \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}} \right) + \frac{1}{4c(1+i \tan(e+fx))} \right)}{a^2 f} \\
& \quad \downarrow 73 \\
& \frac{c \left(\frac{1}{8} (5A - 3iB) \left(\frac{3}{4} \left(\frac{i \int \frac{1}{2 - \frac{c-ic \tan(e+fx)}{e}} d \sqrt{c-ic \tan(e+fx)}}{c^2} - \frac{i}{c \sqrt{c-ic \tan(e+fx)}} \right) + \frac{i}{2c(1+i \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}} \right) + \frac{1}{4c(1+i \tan(e+fx))} \right)}{a^2 f} \\
& \quad \downarrow 219 \\
& \frac{c \left(\frac{1}{8} (5A - 3iB) \left(\frac{3}{4} \left(\frac{i \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2} \sqrt{c}} \right)}{\sqrt{2} c^{3/2}} - \frac{i}{c \sqrt{c-ic \tan(e+fx)}} \right) + \frac{i}{2c(1+i \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}} \right) + \frac{1}{4c(1+i \tan(e+fx))} \right)}{a^2 f}
\end{aligned}$$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*Sqrt[c - I*c*Tan[e + f*x]]),x]`

output `(c*((I*A - B)/(4*c*(1 + I*Tan[e + f*x])^2*Sqrt[c - I*c*Tan[e + f*x]]) + ((5*A - (3*I)*B)*((I/2)/(c*(1 + I*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]) + (3*((I*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])])/(Sqrt[2]*c^(3/2)) - I/(c*Sqrt[c - I*c*Tan[e + f*x]])))/4)/8)/(a^2*f)`

3.776. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 \sqrt{c-ic \tan(e+fx)}} dx$

3.776.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 52 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`
- rule 61 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x],
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.776.4 Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.76

method	result
derivativedivides	$2ic^2 \left(-\frac{-iB+A}{8c^2\sqrt{c-ic\tan(fx+e)}} + \frac{4\left(\frac{iB}{32}-\frac{7A}{32}\right)(c-ic\tan(fx+e))^{\frac{3}{2}}+4\left(\frac{9}{16}cA+\frac{1}{16}iBc\right)\sqrt{c-ic\tan(fx+e)}}{(c+ic\tan(fx+e))^2} + \frac{3\left(-\frac{3iB}{4}+\frac{5A}{4}\right)\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{c-ic\tan(fx+e)}}{c+ic\tan(fx+e)}\right)}{4\sqrt{c^2+a^2}} \right) \frac{1}{fa^2}$
default	$2ic^2 \left(-\frac{-iB+A}{8c^2\sqrt{c-ic\tan(fx+e)}} + \frac{4\left(\frac{iB}{32}-\frac{7A}{32}\right)(c-ic\tan(fx+e))^{\frac{3}{2}}+4\left(\frac{9}{16}cA+\frac{1}{16}iBc\right)\sqrt{c-ic\tan(fx+e)}}{(c+ic\tan(fx+e))^2} + \frac{3\left(-\frac{3iB}{4}+\frac{5A}{4}\right)\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{c-ic\tan(fx+e)}}{c+ic\tan(fx+e)}\right)}{4\sqrt{c^2+a^2}} \right) \frac{1}{fa^2}$

```
input int((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
output 2*I/f/a^2*c^2*(-1/8/c^2*(A-I*B)/(c-I*c*tan(f*x+e))^(1/2)+1/8/c^2*(4*((1/32
*I*B-7/32*A)*(c-I*c*tan(f*x+e))^(3/2)+(9/16*c*A+1/16*I*B*c)*(c-I*c*tan(f*x
+e))^(1/2)))/(c+I*c*tan(f*x+e))^2+3/4*(-3/4*I*B+5/4*A)*2^(1/2)/c^(1/2)*arct
anh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2)))
```

$$3.776. \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 \sqrt{c-ic \tan(e+fx)}} dx$$

3.776.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 390 vs. $2(150) = 300$.

Time = 0.26 (sec) , antiderivative size = 390, normalized size of antiderivative = 2.00

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 \sqrt{c - ic \tan(e + fx)}} dx$$

$$\left(3 \sqrt{\frac{1}{2}} a^2 c f \sqrt{-\frac{25A^2 - 30iAB - 9B^2}{a^4 c f^2}} e^{(4i fx + 4i e)} \log \left(\frac{3 \left(\sqrt{2} \sqrt{\frac{1}{2}} (a^2 f e^{(2i fx + 2i e)} + a^2 f) \sqrt{\frac{c}{e^{(2i fx + 2i e)} + 1}} \sqrt{-\frac{25A^2 - 30iAB - 9B^2}{a^4 c f^2}} + 5 \right)}{16 a^2 f} \right) \right)$$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^2,x
, algorithm="fricas")`

output `1/64*(3*sqrt(1/2)*a^2*c*f*sqrt(-(25*A^2 - 30*I*A*B - 9*B^2)/(a^4*c*f^2))*e
^(4*I*f*x + 4*I*e)*log(3/16*(sqrt(2)*sqrt(1/2)*(a^2*f*e^(2*I*f*x + 2*I*e)
+ a^2*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(25*A^2 - 30*I*A*B - 9*B^2)/(a^4*c*f^2)) + 5*I*A + 3*B)*e^(-I*f*x - I*e)/(a^2*f)) - 3*sqrt(1/2)*a^2
*c*f*sqrt(-(25*A^2 - 30*I*A*B - 9*B^2)/(a^4*c*f^2))*e^(4*I*f*x + 4*I*e)*lo
g(-3/16*(sqrt(2)*sqrt(1/2)*(a^2*f*e^(2*I*f*x + 2*I*e) + a^2*f)*sqrt(c/(e^(
2*I*f*x + 2*I*e) + 1))*sqrt(-(25*A^2 - 30*I*A*B - 9*B^2)/(a^4*c*f^2)) - 5*
I*A - 3*B)*e^(-I*f*x - I*e)/(a^2*f)) - sqrt(2)*(8*(I*A + B)*e^(6*I*f*x + 6
*I*e) - (I*A - 9*B)*e^(4*I*f*x + 4*I*e) - (11*I*A - 3*B)*e^(2*I*f*x + 2*I*
e) - 2*I*A + 2*B)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*e^(-4*I*f*x - 4*I*e)/
(a^2*c*f)`

3.776.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 \sqrt{c - ic \tan(e + fx)}} dx =$$

$$\frac{\int \frac{A}{\sqrt{-ic \tan(e+fx)+c \tan^2(e+fx)-2i\sqrt{-ic \tan(e+fx)+c \tan(e+fx)-\sqrt{-ic \tan(e+fx)+c}}}} dx + \int \frac{\sqrt{-ic \tan(e+fx)+c \tan^2(e+fx)-}}{a^2}}$$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(1/2)/(a+I*a*tan(f*x+e))**2
,x)`

3.776. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 \sqrt{c-ic \tan(e+fx)}} dx$

output $-(\text{Integral}(A/(\sqrt{-I*c*\tan(e + f*x) + c})*\tan(e + f*x)**2 - 2*I*\sqrt{-I*c*\tan(e + f*x) + c})*\tan(e + f*x) - \sqrt{-I*c*\tan(e + f*x) + c}), x) + \text{Integral}(B*\tan(e + f*x)/(\sqrt{-I*c*\tan(e + f*x) + c})*\tan(e + f*x)**2 - 2*I*\sqrt{-I*c*\tan(e + f*x) + c})*\tan(e + f*x) - \sqrt{-I*c*\tan(e + f*x) + c}), x)/a**2$

3.776.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.00

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 \sqrt{c - ic \tan(e + fx)}} dx =$$

$$i \left(\frac{3\sqrt{2}(5A - 3iB)\sqrt{c} \log\left(\frac{\sqrt{2}\sqrt{c} - \sqrt{-ic \tan(fx+e)+c}}{\sqrt{2}\sqrt{c} + \sqrt{-ic \tan(fx+e)+c}}\right)}{a^2} + \frac{4(3(-ic \tan(fx+e)+c)^2(5A - 3iB)c - 10(-ic \tan(fx+e)+c)(5A - 3iB)c^2 + 3(-ic \tan(fx+e)+c)^3)}{(-ic \tan(fx+e)+c)^{\frac{5}{2}}a^2 - 4(-ic \tan(fx+e)+c)^{\frac{3}{2}}a^2c + 4\sqrt{-ic \tan(fx+e)+c}} \right)$$

$128cf$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output $-1/128*I*(3*\sqrt{2}*(5*A - 3*I*B)*\sqrt{c}*\log(-(\sqrt{2}*\sqrt{c} - \sqrt{-I*c*\tan(f*x + e) + c}))/(\sqrt{2}*\sqrt{c} + \sqrt{-I*c*\tan(f*x + e) + c}))/a^2 + 4*(3*(-I*c*\tan(f*x + e) + c)^2*(5*A - 3*I*B)*c - 10*(-I*c*\tan(f*x + e) + c)*(5*A - 3*I*B)*c^2 + 32*(A - I*B)*c^3)/((-I*c*\tan(f*x + e) + c)^(5/2)*a^2 - 4*(-I*c*\tan(f*x + e) + c)^(3/2)*a^2*c + 4*\sqrt{-I*c*\tan(f*x + e) + c}*a^2*c^2))/(c*f)$

3.776.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 \sqrt{c - ic \tan(e + fx)}} dx$$

$$= \int \frac{B \tan(fx + e) + A}{(ia \tan(fx + e) + a)^2 \sqrt{-ic \tan(fx + e) + c}} dx$$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

3.776. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 \sqrt{c-ic \tan(e+fx)}} dx$

output `integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)^2*sqrt(-I*c*tan(f*x + e) + c)), x)`

3.776.9 Mupad [B] (verification not implemented)

Time = 9.05 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.56

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 \sqrt{c - ic \tan(e + fx)}} dx =$$

$$\frac{\frac{A(c - c \tan(e + fx) li)^2 15i}{32 a^2 f} + \frac{A c^2 li}{a^2 f} - \frac{A c(c - c \tan(e + fx) li) 25i}{16 a^2 f}}{-4 c(c - c \tan(e + fx) li)^{3/2} + (c - c \tan(e + fx) li)^{5/2} + 4 c^2 \sqrt{c - c \tan(e + fx) li}}$$

$$\frac{B c^2 + \frac{9 B(c - c \tan(e + fx) li)^2}{32} - \frac{15 B c(c - c \tan(e + fx) li)}{16}}{a^2 f(c - c \tan(e + fx) li)^{5/2} - 4 a^2 c f(c - c \tan(e + fx) li)^{3/2} + 4 a^2 c^2 f \sqrt{c - c \tan(e + fx) li}}$$

$$- \frac{\sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) li}}{2 \sqrt{-c}} li\right) 15i}{64 a^2 \sqrt{-c} f} + \frac{9 \sqrt{2} B \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) li}}{2 \sqrt{c}} li\right)}{64 a^2 \sqrt{c} f}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)^(1/2)),x)`

output `(9*2^(1/2)*B*atanh((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*c^(1/2))))/(64*a^2*c^(1/2)*f) - (B*c^2 + (9*B*(c - c*tan(e + f*x)*1i)^2)/32 - (15*B*c*(c - c*tan(e + f*x)*1i))/16)/(a^2*f*(c - c*tan(e + f*x)*1i)^(5/2) - 4*a^2*c*f*(c - c*tan(e + f*x)*1i)^(3/2) + 4*a^2*c^2*f*(c - c*tan(e + f*x)*1i)^(1/2)) - (2^(1/2)*A*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*(-c)^(1/2)))*15i)/(64*a^2*(-c)^(1/2)*f) - ((A*(c - c*tan(e + f*x)*1i)^2*15i)/(32*a^2*f) + (A*c^2*1i)/(a^2*f) - (A*c*(c - c*tan(e + f*x)*1i)*25i)/(16*a^2*f))/((c - c*tan(e + f*x)*1i)^(5/2) - 4*c*(c - c*tan(e + f*x)*1i)^(3/2) + 4*c^2*(c - c*tan(e + f*x)*1i)^(1/2))`

$$3.777 \quad \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^{3/2}} dx$$

3.777.1 Optimal result	7040
3.777.2 Mathematica [C] (verified)	7040
3.777.3 Rubi [A] (verified)	7041
3.777.4 Maple [A] (verified)	7044
3.777.5 Fracas [B] (verification not implemented)	7044
3.777.6 Sympy [F]	7045
3.777.7 Maxima [A] (verification not implemented)	7046
3.777.8 Giac [F]	7046
3.777.9 Mupad [B] (verification not implemented)	7047

3.777.1 Optimal result

Integrand size = 43, antiderivative size = 226

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2(c - ic \tan(e + fx))^{3/2}} dx = \frac{5(7iA + B) \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{64\sqrt{2}a^2c^{3/2}f} - \frac{5(7iA + B)}{96a^2f(c - ic \tan(e + fx))^{3/2}} + \frac{iA - B}{4a^2f(1 + i \tan(e + fx))^2(c - ic \tan(e + fx))^{3/2}} + \frac{7iA + B}{16a^2f(1 + i \tan(e + fx))(c - ic \tan(e + fx))^{3/2}} - \frac{5(7iA + B)}{64a^2cf\sqrt{c - ic \tan(e + fx)}}$$

output `5/128*(7*I*A+B)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a^2/c^(3/2)/f*2^(1/2)-5/64*(7*I*A+B)/a^2/c/f/(c-I*c*tan(f*x+e))^(1/2)-5/96*(7*I*A+B)/a^2/f/(c-I*c*tan(f*x+e))^(3/2)+1/4*(I*A-B)/a^2/f/(1+I*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^(3/2)+1/16*(7*I*A+B)/a^2/f/(1+I*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2)`

3.777.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 5.95 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.69

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2(c - ic \tan(e + fx))^{3/2}} dx = \frac{\sec^2(e + fx) (-66A - 18iB + 4(A - 7iB) \cos(2(e + fx)) + 28iA \sin(2(e + fx)) + 4B \sin(2(e + fx)) + 192a^2cf(-i + \tan(e + fx))^2(i + \tan(e + fx)))}{192a^2cf(-i + \tan(e + fx))^2(i + \tan(e + fx))}$$

3.777. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^{3/2}} dx$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^(3/2)),x]`

output `-1/192*(Sec[e + f*x]^2*(-66*A - (18*I)*B + 4*(A - (7*I)*B)*Cos[2*(e + f*x)] + (28*I)*A*Sin[2*(e + f*x)] + 4*B*Sin[2*(e + f*x)] + 15*((7*I)*A + B)*Hypergeometric2F1[-1/2, 1, 1/2, (-1/2*I)*(I + Tan[e + f*x])]*(-I + Tan[e + f*x])))/(a^2*c*f*(-I + Tan[e + f*x])^2*(I + Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])]`

3.777.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.96, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.209$, Rules used = {3042, 4071, 27, 87, 52, 61, 61, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & ac \int \frac{A + B \tan(e + fx)}{a^3 (i \tan(e + fx) + 1)^3 (c - ic \tan(e + fx))^{5/2}} d \tan(e + fx) \\
 & \quad \downarrow \text{27} \\
 & \frac{c \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx) + 1)^3 (c - ic \tan(e + fx))^{5/2}} d \tan(e + fx)}{a^2 f} \\
 & \quad \downarrow \text{87} \\
 & \frac{c \left(\frac{1}{8} (7A - iB) \int \frac{1}{(i \tan(e + fx) + 1)^2 (c - ic \tan(e + fx))^{5/2}} d \tan(e + fx) + \frac{-B + iA}{4c(1 + i \tan(e + fx))^2 (c - ic \tan(e + fx))^{3/2}} \right)}{a^2 f} \\
 & \quad \downarrow \text{52}
 \end{aligned}$$

3.777. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^{3/2}} dx$

$$c \left(\frac{1}{8}(7A - iB) \left(\frac{5}{4} \int \frac{1}{(i \tan(e+fx)+1)(c-ic \tan(e+fx))^{5/2}} d \tan(e+fx) + \frac{i}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} \right) + \frac{i}{4c(1+i \tan(e+fx))} \right) \frac{1}{a^2 f}$$

↓ 61

$$c \left(\frac{1}{8}(7A - iB) \left(\frac{5}{4} \left(\int \frac{1}{(i \tan(e+fx)+1)(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx) - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} \right) + \frac{i}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} \right) \right) \frac{1}{a^2 f}$$

↓ 61

$$c \left(\frac{1}{8}(7A - iB) \left(\frac{5}{4} \left(\frac{\int \frac{1}{(i \tan(e+fx)+1)\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{2c} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} \right) + \frac{i}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} \right) \right) \frac{1}{a^2 f}$$

↓ 73

$$c \left(\frac{1}{8}(7A - iB) \left(\frac{5}{4} \left(\frac{\int \frac{1}{2 - \frac{c-ic \tan(e+fx)}{c}} d \sqrt{c-ic \tan(e+fx)}}{c^2} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} \right) + \frac{i}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} \right) \right) \frac{1}{a^2 f}$$

↓ 219

$$c \left(\frac{1}{8}(7A - iB) \left(\frac{5}{4} \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{\sqrt{2}c^{3/2}} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} \right) + \frac{i}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))^{3/2}} \right) \right) \frac{1}{a^2 f}$$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^(3/2)),x]`

output `(c*((I*A - B)/(4*c*(1 + I*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^(3/2)) + ((7*A - I*B)*((I/2)/(c*(1 + I*Tan[e + f*x]))*(c - I*c*Tan[e + f*x])^(3/2)) + (5*((-1/3*I)/(c*(c - I*c*Tan[e + f*x])^(3/2)) + ((I*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])])/(Sqrt[2]*c^(3/2)) - I/(c*Sqrt[c - I*c*Tan[e + f*x]]))/(2*c)))/4)/8))/(a^2*f)`

3.777. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^{3/2}} dx$

3.777.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 52 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`
- rule 61 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p_.)), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x],
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.777.4 Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.78

method	result
derivativedivides	$2ic^2 \left(-\frac{-iB+3A}{16c^3\sqrt{c-ic\tan(fx+e)}} - \frac{-iB+A}{24c^2(c-ic\tan(fx+e))^{\frac{3}{2}}} + \frac{4\left(-\frac{3iB}{32} - \frac{11A}{32}\right)(c-ic\tan(fx+e))^{\frac{3}{2}} + 4\left(\frac{13}{16}cA + \frac{5}{16}iBc\right)\sqrt{c-ic\tan(fx+e)}}{(c+ic\tan(fx+e))^2} \right) \frac{fa^2}{16c^3}$
default	$2ic^2 \left(-\frac{-iB+3A}{16c^3\sqrt{c-ic\tan(fx+e)}} - \frac{-iB+A}{24c^2(c-ic\tan(fx+e))^{\frac{3}{2}}} + \frac{4\left(-\frac{3iB}{32} - \frac{11A}{32}\right)(c-ic\tan(fx+e))^{\frac{3}{2}} + 4\left(\frac{13}{16}cA + \frac{5}{16}iBc\right)\sqrt{c-ic\tan(fx+e)}}{(c+ic\tan(fx+e))^2} \right) \frac{fa^2}{16c^3}$

```
input int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
```

```
output 2*I/f/a^2*c^2*(-1/16/c^3*(3*A-I*B)/(c-I*c*tan(f*x+e))^(1/2)-1/24/c^2*(A-I*B)/(c-I*c*tan(f*x+e))^(3/2)+1/16/c^3*(4*((-3/32*I*B-11/32*A)*(c-I*c*tan(f*x+e))^(3/2)+(13/16*c*A+5/16*I*B*c)*(c-I*c*tan(f*x+e))^(1/2))/(c+I*c*tan(f*x+e))^2+5/4*(7/4*A-1/4*I*B)*2^(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))))
```

3.777.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 420 vs. 2(181) = 362.

Time = 0.27 (sec) , antiderivative size = 420, normalized size of antiderivative = 1.86

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^{3/2}} dx = \frac{\left(15 \sqrt{\frac{1}{2}} a^2 c^2 f \sqrt{-\frac{49 A^2 - 14 i A B - B^2}{a^4 c^3 f^2}} e^{(4i f x + 4i e)} \log \left(\frac{5 \left(\sqrt{2} \right)}{\dots} \right) \right)}{\dots}$$

3.777. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 (c-ic \tan(e+fx))^{3/2}} dx$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^(3/2),x
, algorithm="fricas")`

output `1/384*(15*sqrt(1/2)*a^2*c^2*f*sqrt(-(49*A^2 - 14*I*A*B - B^2)/(a^4*c^3*f^2
) * e^(4*I*f*x + 4*I*e) * log(5/32*(sqrt(2)*sqrt(1/2)*(a^2*c*f*e^(2*I*f*x + 2
*I*e) + a^2*c*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(49*A^2 - 14*I*A*B - B^2)/(a^4*c^3*f^2)) + 7*I*A + B)*e^(-I*f*x - I*e)/(a^2*c*f)) - 15*sqrt
(1/2)*a^2*c^2*f*sqrt(-(49*A^2 - 14*I*A*B - B^2)/(a^4*c^3*f^2))*e^(4*I*f*x
+ 4*I*e) * log(-5/32*(sqrt(2)*sqrt(1/2)*(a^2*c*f*e^(2*I*f*x + 2*I*e) + a^2*c
*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(49*A^2 - 14*I*A*B - B^2)/(a^4
*c^3*f^2)) - 7*I*A - B)*e^(-I*f*x - I*e)/(a^2*c*f)) - sqrt(2)*(8*(I*A + B)
*e^(8*I*f*x + 8*I*e) + 8*(11*I*A + 5*B)*e^(6*I*f*x + 6*I*e) - (-41*I*A - 4
7*B)*e^(4*I*f*x + 4*I*e) + 3*(-15*I*A + 7*B)*e^(2*I*f*x + 2*I*e) - 6*I*A +
6*B)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))) * e^(-4*I*f*x - 4*I*e)/(a^2*c^2*f)`

3.777.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^{3/2}} dx =$$

$$\frac{\int \frac{A}{-ic\sqrt{-ic \tan(e+fx)+c} \tan^3(e+fx) - c\sqrt{-ic \tan(e+fx)+c} \tan^2(e+fx) - ic\sqrt{-ic \tan(e+fx)+c} \tan(e+fx) - c\sqrt{-ic \tan(e+fx)+c}} dx + \int \frac{B}{a^2} dx}{a^2}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**2/(c-I*c*tan(f*x+e))**3/2
,x)`

output `-(Integral(A/(-I*c*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x)**3 - c*sqrt(-I
*c*tan(e + f*x) + c))*tan(e + f*x)**2 - I*c*sqrt(-I*c*tan(e + f*x) + c))*tan
(e + f*x) - c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(B*tan(e + f*x)/(
-I*c*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x)**3 - c*sqrt(-I*c*tan(e + f*x
) + c))*tan(e + f*x)**2 - I*c*sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x) - c
*sqrt(-I*c*tan(e + f*x) + c)), x))/a**2`

3.777. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 (c-ic \tan(e+fx))^{3/2}} dx$

3.777.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 217, normalized size of antiderivative = 0.96

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^{3/2}} dx =$$

$$i \left(\frac{4 \left(15(-ic \tan(fx+e)+c)^3(7A-iB) - 50(-ic \tan(fx+e)+c)^2(7A-iB)c + 32(-ic \tan(fx+e)+c)(7A-iB)c^2 + 64(A-iB)c^3 \right)}{(-ic \tan(fx+e)+c)^{7/2} a^2 - 4(-ic \tan(fx+e)+c)^{5/2} a^2 c + 4(-ic \tan(fx+e)+c)^{3/2} a^2 c^2} \right) + \frac{15\sqrt{2}(7A-iB)}{768cf}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^(3/2),x
, algorithm="maxima")
```

```
output -1/768*I*(4*(15*(-I*c*tan(f*x + e) + c)^3*(7*A - I*B) - 50*(-I*c*tan(f*x +
e) + c)^2*(7*A - I*B)*c + 32*(-I*c*tan(f*x + e) + c)*(7*A - I*B)*c^2 + 64
*(A - I*B)*c^3)/((-I*c*tan(f*x + e) + c)^(7/2)*a^2 - 4*(-I*c*tan(f*x + e)
+ c)^(5/2)*a^2*c + 4*(-I*c*tan(f*x + e) + c)^(3/2)*a^2*c^2) + 15*sqrt(2)*(
7*A - I*B)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*s
qrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/(a^2*sqrt(c))/(c*f)
```

3.777.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^{3/2}} dx = \int \frac{B \tan(fx + e) + A}{(ia \tan(fx + e) + a)^2 (-ic \tan(fx + e) + c)^{3/2}} dx$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^(3/2),x
, algorithm="giac")
```

```
output integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)^2*(-I*c*tan(f*x + e)
) + c)^(3/2)), x)
```

3.777.9 Mupad [B] (verification not implemented)

Time = 9.30 (sec) , antiderivative size = 353, normalized size of antiderivative = 1.56

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^{3/2}} dx =$$

$$\frac{\frac{Bc^2}{3} - \frac{25B(c - c \tan(e + fx) li)^2}{96} + \frac{5B(c - c \tan(e + fx) li)^3}{64c} + \frac{Bc(c - c \tan(e + fx) li)}{6}}{a^2 f (c - c \tan(e + fx) li)^{7/2} - 4a^2 c f (c - c \tan(e + fx) li)^{5/2} + 4a^2 c^2 f (c - c \tan(e + fx) li)^{3/2}}$$

$$- \frac{-\frac{A(c - c \tan(e + fx) li)^2 175i}{96a^2 f} + \frac{Ac^2 li}{3a^2 f} + \frac{A(c - c \tan(e + fx) li)^3 35i}{64a^2 c f} + \frac{Ac(c - c \tan(e + fx) li) 7i}{6a^2 f}}{-4c(c - c \tan(e + fx) li)^{5/2} + (c - c \tan(e + fx) li)^{7/2} + 4c^2 (c - c \tan(e + fx) li)^{3/2}}$$

$$+ \frac{\sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) li}}{2\sqrt{-c}}\right) 35i}{128a^2 (-c)^{3/2} f} + \frac{5\sqrt{2} B \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) li}}{2\sqrt{c}}\right)}{128a^2 c^{3/2} f}$$

```
input int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i)^(3/2)),x)
```

```
output (2^(1/2)*A*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*(-c)^(1/2)))*35i)/(128*a^2*(-c)^(3/2)*f) - ((A*c^2*1i)/(3*a^2*f) - (A*(c - c*tan(e + f*x)*1i)^2*175i)/(96*a^2*f) + (A*(c - c*tan(e + f*x)*1i)^3*35i)/(64*a^2*c*f) + (A*c*(c - c*tan(e + f*x)*1i)*7i)/(6*a^2*f))/((c - c*tan(e + f*x)*1i)^(7/2) - 4*c*(c - c*tan(e + f*x)*1i)^(5/2) + 4*c^2*(c - c*tan(e + f*x)*1i)^(3/2)) - ((B*c^2)/3 - (25*B*(c - c*tan(e + f*x)*1i)^2)/96 + (5*B*(c - c*tan(e + f*x)*1i)^3)/(64*c) + (B*c*(c - c*tan(e + f*x)*1i))/6)/(a^2*f*(c - c*tan(e + f*x)*1i)^(7/2) - 4*a^2*c*f*(c - c*tan(e + f*x)*1i)^(5/2) + 4*a^2*c^2*f*(c - c*tan(e + f*x)*1i)^(3/2)) + (5*2^(1/2)*B*atanh((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*c^(1/2))))/(128*a^2*c^(3/2)*f)
```

3.778
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^{5/2}} dx$$

3.778.1 Optimal result 7048
 3.778.2 Mathematica [C] (verified) 7049
 3.778.3 Rubi [A] (verified) 7049
 3.778.4 Maple [A] (verified) 7053
 3.778.5 Fricas [B] (verification not implemented) 7053
 3.778.6 Sympy [F] 7054
 3.778.7 Maxima [A] (verification not implemented) 7054
 3.778.8 Giac [F] 7055
 3.778.9 Mupad [B] (verification not implemented) 7055

3.778.1 Optimal result

Integrand size = 43, antiderivative size = 273

$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^{5/2}} dx = \frac{7(9iA-B) \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{128\sqrt{2}a^2c^{5/2}f} - \frac{7(9iA-B)}{160a^2f(c-ic \tan(e+fx))^{5/2}} + \frac{iA-B}{4a^2f(1+i \tan(e+fx))^2(c-ic \tan(e+fx))^{5/2}} + \frac{16a^2f(1+i \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{9iA-B} - \frac{7(9iA-B)}{192a^2cf(c-ic \tan(e+fx))^{3/2}} - \frac{7(9iA-B)}{128a^2c^2f\sqrt{c-ic \tan(e+fx)}}$$

```
output 7/256*(9*I*A-B)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a^2/c^(5/2)/f*2^(1/2)-7/128*(9*I*A-B)/a^2/c^2/f/(c-I*c*tan(f*x+e))^(1/2)-7/160*(9*I*A-B)/a^2/f/(c-I*c*tan(f*x+e))^(5/2)+1/4*(I*A-B)/a^2/f/(1+I*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^(5/2)+1/16*(9*I*A-B)/a^2/f/(1+I*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2)-7/192*(9*I*A-B)/a^2/c/f/(c-I*c*tan(f*x+e))^(3/2)
```

3.778.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 5.84 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.52

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ict \tan(e + fx))^{5/2}} dx = \frac{\cos(e + fx) (35(-9iA + B) \text{Hypergeometric2F1}(-\frac{3}{2}, 1, -1/2, (-1/2 * I) * (I + \tan(e + fx))) * (\cos(e + fx) + I \sin(e + fx)) + 6 * \cos(e + fx) * ((65 * I) * A - 25 * B + (2 * I) * (A + (9 * I) * B) * \cos[2 * (e + fx)] + 2 * (9 * A + I * B) * \sin[2 * (e + fx)]))}}{(960 * a^2 * c^2 * f * \text{Sqrt}[c - I * c * \tan(e + fx)])}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^(5/2)),x]`

output `(Cos[e + f*x]*(35*((-9*I)*A + B)*Hypergeometric2F1[-3/2, 1, -1/2, (-1/2*I)*(I + Tan[e + f*x]])*(Cos[e + f*x] + I*Sin[e + f*x]) + 6*Cos[e + f*x]*((65*I)*A - 25*B + (2*I)*(A + (9*I)*B)*Cos[2*(e + f*x)] + 2*(9*A + I*B)*Sin[2*(e + f*x)])))/(960*a^2*c^2*f*Sqrt[c - I*c*Tan[e + f*x]])`

3.778.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 250, normalized size of antiderivative = 0.92, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.233$, Rules used = {3042, 4071, 27, 87, 52, 61, 61, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ict \tan(e + fx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ict \tan(e + fx))^{5/2}} dx \\ & \quad \downarrow \text{4071} \\ & ac \int \frac{A + B \tan(e + fx)}{a^3 (i \tan(e + fx) + 1)^3 (c - ict \tan(e + fx))^{7/2}} d \tan(e + fx) \\ & \quad \downarrow \text{27} \\ & c \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx) + 1)^3 (c - ict \tan(e + fx))^{7/2}} d \tan(e + fx) \\ & \quad \quad \quad a^2 f \end{aligned}$$

3.778. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ict \tan(e + fx))^{5/2}} dx$

$$\frac{c \left(\frac{1}{8}(9A + iB) \int \frac{1}{(i \tan(e+fx)+1)^2 (c-ic \tan(e+fx))^{7/2}} d \tan(e+fx) + \frac{-B+iA}{4c(1+i \tan(e+fx))^2 (c-ic \tan(e+fx))^{5/2}} \right)}{a^2 f}$$

↓ 87

$$\frac{c \left(\frac{1}{8}(9A + iB) \left(\frac{7}{4} \int \frac{1}{(i \tan(e+fx)+1)(c-ic \tan(e+fx))^{7/2}} d \tan(e+fx) + \frac{i}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))^{5/2}} \right) + \frac{i}{4c(1+i \tan(e+fx))^{5/2}} \right)}{a^2 f}$$

↓ 52

$$\frac{c \left(\frac{1}{8}(9A + iB) \left(\frac{7}{4} \left(\int \frac{1}{(i \tan(e+fx)+1)(c-ic \tan(e+fx))^{5/2}} d \tan(e+fx) - \frac{i}{5c(c-ic \tan(e+fx))^{5/2}} \right) + \frac{i}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))^{5/2}} \right) \right)}{a^2 f}$$

↓ 61

$$\frac{c \left(\frac{1}{8}(9A + iB) \left(\frac{7}{4} \left(\frac{\int \frac{1}{(i \tan(e+fx)+1)(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{2c} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} - \frac{i}{5c(c-ic \tan(e+fx))^{5/2}} \right) + \frac{i}{2c(1+i \tan(e+fx))^{5/2}} \right) \right)}{a^2 f}$$

↓ 61

$$\frac{c \left(\frac{1}{8}(9A + iB) \left(\frac{7}{4} \left(\frac{\int \frac{1}{(i \tan(e+fx)+1)\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{2c} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} - \frac{i}{5c(c-ic \tan(e+fx))^{5/2}} \right) \right) \right)}{a^2 f}$$

↓ 61

$$\frac{c \left(\frac{1}{8}(9A + iB) \left(\frac{7}{4} \left(\frac{i \int \frac{1}{2 - \frac{c-ic \tan(e+fx)}{c}} d \sqrt{c-ic \tan(e+fx)}}{c^2} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} - \frac{i}{5c(c-ic \tan(e+fx))^{5/2}} \right) \right) \right)}{a^2 f}$$

↓ 73

↓ 219

3.778. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 (c-ic \tan(e+fx))^{5/2}} dx$

$$c \left(\frac{1}{8}(9A + iB) \left(\frac{7}{4} \left(\frac{\operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{\sqrt{2}c^{3/2}} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}} - \frac{i}{5c(c-ic \tan(e+fx))^{5/2}} \right) + \frac{i}{2c(1+i \tan(e+fx))} \right) \right) \frac{1}{a^2 f}$$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^(5/2)),x]`

output `(c*((I*A - B)/(4*c*(1 + I*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^(5/2)) + ((9*A + I*B)*((I/2)/(c*(1 + I*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2)) + (7*((-1/5*I)/(c*(c - I*c*Tan[e + f*x])^(5/2)) + ((-1/3*I)/(c*(c - I*c*Tan[e + f*x])^(3/2)) + ((I*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])])/(Sqrt[2]*c^(3/2)) - I/(c*Sqrt[c - I*c*Tan[e + f*x]])/(2*c))/(2*c)))/4))/8)/(a^2*f)`

3.778.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 52 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 61 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

- rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
 .), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
 + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
 + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
 /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
 rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
 ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
 Q[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
 Q[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
 (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
 mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
 , Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
 + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.778.4 Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.72

method	result
derivativedivides	$2ic^2 \left(\frac{4 \left(-\frac{7iB}{64} - \frac{15A}{64} \right) (c - ic \tan(fx+e))^{\frac{3}{2}} + 4 \left(\frac{9}{32} iBc + \frac{17}{32} cA \right) \sqrt{c - ic \tan(fx+e)}}{(c + ic \tan(fx+e))^2} + \frac{7 \left(\frac{iB}{4} + \frac{9A}{4} \right) \sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{c - ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}} \right)}{8\sqrt{c}} \right) \frac{f a^2}{16c^4}$
default	$2ic^2 \left(\frac{4 \left(-\frac{7iB}{64} - \frac{15A}{64} \right) (c - ic \tan(fx+e))^{\frac{3}{2}} + 4 \left(\frac{9}{32} iBc + \frac{17}{32} cA \right) \sqrt{c - ic \tan(fx+e)}}{(c + ic \tan(fx+e))^2} + \frac{7 \left(\frac{iB}{4} + \frac{9A}{4} \right) \sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{c - ic \tan(fx+e)} \sqrt{2}}{2\sqrt{c}} \right)}{8\sqrt{c}} \right) \frac{f a^2}{16c^4}$

```
input int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)
```

```
output 2*I/f/a^2*c^2*(1/16/c^4*(4*((-7/64*I*B-15/64*A)*(c-I*c*tan(f*x+e))^(3/2)+(9/32*I*B*c+17/32*c*A)*(c-I*c*tan(f*x+e))^(1/2)))/(c+I*c*tan(f*x+e))^2+7/8*(1/4*I*B+9/4*A)*2^(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))-3/16/c^4*A/(c-I*c*tan(f*x+e))^(1/2)-1/48/c^3*(3*A-I*B)/(c-I*c*tan(f*x+e))^(3/2)-1/40/c^2*(A-I*B)/(c-I*c*tan(f*x+e))^(5/2))
```

3.778.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 446 vs. 2(202) = 404.

Time = 0.28 (sec) , antiderivative size = 446, normalized size of antiderivative = 1.63

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^{5/2}} dx = \left(105 \sqrt{\frac{1}{2}} a^2 c^3 f \sqrt{-\frac{81 A^2 + 18i AB - B^2}{a^4 c^5 f^2}} e^{(4i fx + 4i e)} \log \left(\frac{7 \left(\sqrt{c - ic \tan(e + fx)} \sqrt{2} \right)}{2\sqrt{c}} \right) \right)$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^(5/2),x,algorithm="fracas")
```

output $1/3840*(105*\sqrt{1/2}*a^2*c^3*f*\sqrt{-(81*A^2 + 18*I*A*B - B^2)/(a^4*c^5*f^2)})*e^{(4*I*f*x + 4*I*e)}*\log(7/64*(\sqrt{2})*\sqrt{1/2}*(a^2*c^2*f*e^{(2*I*f*x + 2*I*e)} + a^2*c^2*f)*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{-(81*A^2 + 18*I*A*B - B^2)/(a^4*c^5*f^2)}) + 9*I*A - B)*e^{(-I*f*x - I*e)/(a^2*c^2*f)} - 105*\sqrt{1/2}*a^2*c^3*f*\sqrt{-(81*A^2 + 18*I*A*B - B^2)/(a^4*c^5*f^2)}*e^{(4*I*f*x + 4*I*e)}*\log(-7/64*(\sqrt{2})*\sqrt{1/2}*(a^2*c^2*f*e^{(2*I*f*x + 2*I*e)} + a^2*c^2*f)*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{-(81*A^2 + 18*I*A*B - B^2)/(a^4*c^5*f^2)}) - 9*I*A + B)*e^{(-I*f*x - I*e)/(a^2*c^2*f)} - \sqrt{2}*(24*(I*A + B)*e^{(10*I*f*x + 10*I*e)} + 16*(12*I*A + 7*B)*e^{(8*I*f*x + 8*I*e)} + 8*(129*I*A + 19*B)*e^{(6*I*f*x + 6*I*e)} - (-609*I*A - 199*B)*e^{(4*I*f*x + 4*I*e)} + 15*(-19*I*A + 11*B)*e^{(2*I*f*x + 2*I*e)} - 30*I*A + 30*B)*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)})*e^{(-4*I*f*x - 4*I*e)/(a^2*c^3*f)}$

3.778.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^{5/2}} dx = \frac{\int \frac{A}{-c^2 \sqrt{-ic \tan(e+fx)+c} \tan^4(e+fx) - 2c^2 \sqrt{-ic \tan(e+fx)+c} \tan^2(e+fx) - c^2 \sqrt{-ic \tan(e+fx)+c}} dx + \int \frac{B}{-c^2 \sqrt{-ic \tan(e+fx)+c} \tan^4(e+fx) - 2c^2 \sqrt{-ic \tan(e+fx)+c} \tan^2(e+fx) - c^2 \sqrt{-ic \tan(e+fx)+c}} dx}{a^2}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**2/(c-I*c*tan(f*x+e))**(5/2),x)`

output $-(\text{Integral}(A/(-c**2*\sqrt{-I*c*\tan(e + f*x) + c})*\tan(e + f*x)**4 - 2*c**2*\sqrt{-I*c*\tan(e + f*x) + c})*\tan(e + f*x)**2 - c**2*\sqrt{-I*c*\tan(e + f*x) + c})), x) + \text{Integral}(B*\tan(e + f*x)/(-c**2*\sqrt{-I*c*\tan(e + f*x) + c})*\tan(e + f*x)**4 - 2*c**2*\sqrt{-I*c*\tan(e + f*x) + c})*\tan(e + f*x)**2 - c**2*\sqrt{-I*c*\tan(e + f*x) + c})), x))/a**2$

3.778.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 245, normalized size of antiderivative = 0.90

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^{5/2}} dx = i \left(\frac{4 \left(105 (-ic \tan(fx+e)+c)^4 (9A+iB) - 350 (-ic \tan(fx+e)+c)^3 (9A+iB)c + 224 (-ic \tan(fx+e)+c)^2 (9A+iB)c^2 + 64 (-ic \tan(fx+e)+c) \right)}{(-ic \tan(fx+e)+c)^{\frac{9}{2}} a^2 c - 4 (-ic \tan(fx+e)+c)^{\frac{7}{2}} a^2 c^2 + 4 (-ic \tan(fx+e)+c)^{\frac{5}{2}} a^2 c^3} \right)$$

7680 cf

3.778. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 (c-ic \tan(e+fx))^{5/2}} dx$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^(5/2),x
, algorithm="maxima")`

output `-1/7680*I*(4*(105*(-I*c*tan(f*x + e) + c)^4*(9*A + I*B) - 350*(-I*c*tan(f*
x + e) + c)^3*(9*A + I*B)*c + 224*(-I*c*tan(f*x + e) + c)^2*(9*A + I*B)*c^
2 + 64*(-I*c*tan(f*x + e) + c)*(9*A + I*B)*c^3 + 384*(A - I*B)*c^4)/((-I*c
*tan(f*x + e) + c)^(9/2)*a^2*c - 4*(-I*c*tan(f*x + e) + c)^(7/2)*a^2*c^2 +
4*(-I*c*tan(f*x + e) + c)^(5/2)*a^2*c^3) + 105*sqrt(2)*(9*A + I*B)*log(-(
sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*
c*tan(f*x + e) + c)))/(a^2*c^(3/2)))/(c*f)`

3.778.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^{5/2}} dx = \int \frac{B \tan(fx + e) + A}{(ia \tan(fx + e) + a)^2 (-ic \tan(fx + e) + c)^{5/2}} dx$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^(5/2),x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)^2*(-I*c*tan(f*x + e)
) + c)^(5/2)), x)`

3.778.9 Mupad [B] (verification not implemented)

Time = 9.88 (sec) , antiderivative size = 399, normalized size of antiderivative = 1.46

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^2 (c - ic \tan(e + fx))^{5/2}} dx = \frac{-\frac{Bc^2}{5} + \frac{7B(c - c \tan(e + fx) li)^2}{60} - \frac{35B(c - c \tan(e + fx) li)^3}{192c}}{a^2 f (c - c \tan(e + fx) li)^{9/2} - 4a^2 c f (c - c \tan(e + fx) li)^{7/2} + (c - c \tan(e + fx) li)^{9/2} + 4c^2 (c - c \tan(e + fx) li)^{5/2}}$$

$$- \frac{\frac{A(c - c \tan(e + fx) li)^2 2li}{20a^2 f} + \frac{Ac^2 li}{5a^2 f} - \frac{A(c - c \tan(e + fx) li)^3 105i}{64a^2 c f} + \frac{A(c - c \tan(e + fx) li)^4 63i}{128a^2 c^2 f} + \frac{Ac(c - c \tan(e + fx) li) 3i}{10a^2 f}}{256a^2 (-c)^{5/2} f} - \frac{7\sqrt{2} B \operatorname{atanh}\left(\frac{\sqrt{2}\sqrt{c - c \tan(e + fx) li}}{2\sqrt{c}}\right)}{256a^2 c^{5/2} f}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^2*(c - c*tan(e + f*x)*1i
)^(5/2)),x)`

3.778. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2 (c-ic \tan(e+fx))^{5/2}} dx$

output

$$\begin{aligned} & ((7*B*(c - c*\tan(e + f*x)*1i)^2)/60 - (B*c^2)/5 - (35*B*(c - c*\tan(e + f*x) \\ &)*1i)^3)/(192*c) + (7*B*(c - c*\tan(e + f*x)*1i)^4)/(128*c^2) + (B*c*(c - c \\ & * \tan(e + f*x)*1i))/30)/(a^2*f*(c - c*\tan(e + f*x)*1i)^{(9/2)} - 4*a^2*c*f*(c \\ & - c*\tan(e + f*x)*1i)^{(7/2)} + 4*a^2*c^2*f*(c - c*\tan(e + f*x)*1i)^{(5/2)}) - \\ & ((A*(c - c*\tan(e + f*x)*1i)^2*21i)/(20*a^2*f) + (A*c^2*1i)/(5*a^2*f) - (A \\ & *(c - c*\tan(e + f*x)*1i)^3*105i)/(64*a^2*c*f) + (A*(c - c*\tan(e + f*x)*1i) \\ & ^4*63i)/(128*a^2*c^2*f) + (A*c*(c - c*\tan(e + f*x)*1i)*3i)/(10*a^2*f))/((c \\ & - c*\tan(e + f*x)*1i)^{(9/2)} - 4*c*(c - c*\tan(e + f*x)*1i)^{(7/2)} + 4*c^2*(c \\ & - c*\tan(e + f*x)*1i)^{(5/2)}) - (2^{(1/2)}*A*atan((2^{(1/2)}*(c - c*\tan(e + f*x) \\ &)*1i)^{(1/2)})/(2*(-c)^{(1/2)}))*63i)/(256*a^2*(-c)^{(5/2)}*f) - (7*2^{(1/2)}*B*at \\ & anh((2^{(1/2)}*(c - c*\tan(e + f*x)*1i)^{(1/2)})/(2*c^{(1/2)})))/(256*a^2*c^{(5/2)} \\ & *f) \end{aligned}$$

3.778. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^2(c-ic \tan(e+fx))^{5/2}} dx$

3.779
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^3} dx$$

3.779.1 Optimal result 7057
 3.779.2 Mathematica [A] (verified) 7058
 3.779.3 Rubi [A] (verified) 7058
 3.779.4 Maple [A] (verified) 7061
 3.779.5 Fricas [B] (verification not implemented) 7062
 3.779.6 Sympy [F(-1)] 7062
 3.779.7 Maxima [A] (verification not implemented) 7063
 3.779.8 Giac [F] 7063
 3.779.9 Mupad [B] (verification not implemented) 7064

3.779.1 Optimal result

Integrand size = 43, antiderivative size = 291

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{9/2}}{(a + ia \tan(e + fx))^3} dx =$$

$$-\frac{35(iA - 5B)c^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{4\sqrt{2}a^3 f} + \frac{35(iA - 5B)c^4 \sqrt{c - ic \tan(e + fx)}}{8a^3 f}$$

$$+ \frac{35(iA - 5B)c^3 (c - ic \tan(e + fx))^{3/2}}{48a^3 f} + \frac{7(iA - 5B)c^2 (c - ic \tan(e + fx))^{5/2}}{16a^3 f (1 + i \tan(e + fx))}$$

$$- \frac{(iA - 5B)c (c - ic \tan(e + fx))^{7/2}}{8a^3 f (1 + i \tan(e + fx))^2} + \frac{(iA - B)(c - ic \tan(e + fx))^{9/2}}{6a^3 f (1 + i \tan(e + fx))^3}$$

```
output -35/8*(I*A-5*B)*c^(9/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a^3/f*2^(1/2)+35/8*(I*A-5*B)*c^4*(c-I*c*tan(f*x+e))^(1/2)/a^3/f+35/48*(I*A-5*B)*c^3*(c-I*c*tan(f*x+e))^(3/2)/a^3/f+7/16*(I*A-5*B)*c^2*(c-I*c*tan(f*x+e))^(5/2)/a^3/f/(1+I*tan(f*x+e))-1/8*(I*A-5*B)*c*(c-I*c*tan(f*x+e))^(7/2)/a^3/f/(1+I*tan(f*x+e))^2+1/6*(I*A-B)*(c-I*c*tan(f*x+e))^(9/2)/a^3/f/(1+I*tan(f*x+e))^3
```


3.779.2 Mathematica [A] (verified)

Time = 7.27 (sec) , antiderivative size = 201, normalized size of antiderivative = 0.69

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{9/2}}{(a + ia \tan(e + fx))^3} dx = \frac{105\sqrt{2}(A + 5iB)c^{9/2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right) \sec^3(e + fx)}{a^3}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(9/2))/(a + I*a*Tan[e + f*x])^3,x]`

output `(105*sqrt[2]*(A + (5*I)*B)*c^(9/2)*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(sqrt[2]*sqrt[c])]*Sec[e + f*x]^3*(Cos[3*(e + f*x)] + I*Sin[3*(e + f*x)]) + 2*c^4*sqrt[c - I*c*Tan[e + f*x]]*(-67*A - (327*I)*B + 2*((-85*I)*A + 441*B)*Tan[e + f*x] + 3*(53*A + (249*I)*B)*Tan[e + f*x]^2 + (8*I)*(3*A + (19*I)*B)*Tan[e + f*x]^3 + (8*I)*B*Tan[e + f*x]^4)/(24*a^3*f*(-I + Tan[e + f*x])^3)`

3.779.3 Rubi [A] (verified)Time = 0.42 (sec) , antiderivative size = 243, normalized size of antiderivative = 0.84, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.233$, Rules used = {3042, 4071, 27, 87, 51, 51, 60, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c - ic \tan(e + fx))^{9/2}(A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c - ic \tan(e + fx))^{9/2}(A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{a^4(i \tan(e+fx)+1)^4} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{c \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(i \tan(e+fx)+1)^4} d \tan(e + fx)}{a^3 f} \end{aligned}$$

3.779. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^3} dx$

$$\begin{aligned}
& \downarrow 87 \\
& \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{9/2}}{6c(1+i\tan(e+fx))^3} - \frac{1}{4}(A+5iB) \int \frac{(c-ictan(e+fx))^{7/2}}{(i\tan(e+fx)+1)^3} d\tan(e+fx)\right)}{a^3 f} \\
& \downarrow 51 \\
& \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{9/2}}{6c(1+i\tan(e+fx))^3} - \frac{1}{4}(A+5iB) \left(\frac{i(c-ictan(e+fx))^{7/2}}{2(1+i\tan(e+fx))^2} - \frac{7}{4}c \int \frac{(c-ictan(e+fx))^{5/2}}{(i\tan(e+fx)+1)^2} d\tan(e+fx)\right)\right)}{a^3 f} \\
& \downarrow 51 \\
& \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{9/2}}{6c(1+i\tan(e+fx))^3} - \frac{1}{4}(A+5iB) \left(\frac{i(c-ictan(e+fx))^{7/2}}{2(1+i\tan(e+fx))^2} - \frac{7}{4}c \left(\frac{i(c-ictan(e+fx))^{5/2}}{1+i\tan(e+fx)} - \frac{5}{2}c \int \frac{(c-ictan(e+fx))^{3/2}}{i\tan(e+fx)+1} d\tan(e+fx)\right)\right)\right)}{a^3 f} \\
& \downarrow 60 \\
& \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{9/2}}{6c(1+i\tan(e+fx))^3} - \frac{1}{4}(A+5iB) \left(\frac{i(c-ictan(e+fx))^{7/2}}{2(1+i\tan(e+fx))^2} - \frac{7}{4}c \left(\frac{i(c-ictan(e+fx))^{5/2}}{1+i\tan(e+fx)} - \frac{5}{2}c \left(2c \int \frac{\sqrt{c-ictan(e+fx)}}{i\tan(e+fx)+1} d\tan(e+fx)\right)\right)\right)\right)}{a^3 f} \\
& \downarrow 60 \\
& \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{9/2}}{6c(1+i\tan(e+fx))^3} - \frac{1}{4}(A+5iB) \left(\frac{i(c-ictan(e+fx))^{7/2}}{2(1+i\tan(e+fx))^2} - \frac{7}{4}c \left(\frac{i(c-ictan(e+fx))^{5/2}}{1+i\tan(e+fx)} - \frac{5}{2}c \left(2c \left(2c \int \frac{1}{(i\tan(e+fx)+1)} d\tan(e+fx)\right)\right)\right)\right)\right)}{a^3 f} \\
& \downarrow 73 \\
& \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{9/2}}{6c(1+i\tan(e+fx))^3} - \frac{1}{4}(A+5iB) \left(\frac{i(c-ictan(e+fx))^{7/2}}{2(1+i\tan(e+fx))^2} - \frac{7}{4}c \left(\frac{i(c-ictan(e+fx))^{5/2}}{1+i\tan(e+fx)} - \frac{5}{2}c \left(2c \left(4i \int \frac{1}{2-\frac{c-ictan(e+fx)}{c}} d\tan(e+fx)\right)\right)\right)\right)\right)}{a^3 f} \\
& \downarrow 219 \\
& \frac{c\left(\frac{(-B+iA)(c-ictan(e+fx))^{9/2}}{6c(1+i\tan(e+fx))^3} - \frac{1}{4}(A+5iB) \left(\frac{i(c-ictan(e+fx))^{7/2}}{2(1+i\tan(e+fx))^2} - \frac{7}{4}c \left(\frac{i(c-ictan(e+fx))^{5/2}}{1+i\tan(e+fx)} - \frac{5}{2}c \left(2c \left(2i\sqrt{2}\sqrt{c}\operatorname{arctanh}\left(\frac{1}{2-\frac{c-ictan(e+fx)}{c}}\right)\right)\right)\right)\right)\right)}{a^3 f}
\end{aligned}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(9/2))/(a + I*a*Tan[e + f*x])^3,x]`

3.779. $\int \frac{(A+B \tan(e+fx))(c-ictan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^3} dx$

```
output (c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(9/2))/(6*c*(1 + I*Tan[e + f*x])^3)
- ((A + (5*I)*B)*(((I/2)*(c - I*c*Tan[e + f*x])^(7/2))/(1 + I*Tan[e + f*x]
)^2 - (7*c*(((I*(c - I*c*Tan[e + f*x])^(5/2))/(1 + I*Tan[e + f*x]) - (5*c*(
((-2*I)/3)*(c - I*c*Tan[e + f*x])^(3/2) + 2*c*((2*I)*Sqrt[2]*Sqrt[c]*ArcTa
nh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])]) - (2*I)*Sqrt[c - I*c*Tan[
e + f*x]])))/2))/4))/4))/(a^3*f)
```

3.779.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 51 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x
] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]
```

```
rule 60 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(
b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.779.4 Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 207, normalized size of antiderivative = 0.71

method	result
derivativedivides	$2ic^3 \left(\frac{iB(c-ic \tan(fx+e))^{3/2}}{3} + 7i\sqrt{c-ic \tan(fx+e)} Bc + \sqrt{c-ic \tan(fx+e)} cA - 8c^2 \left(\frac{8(-\frac{81iB}{512} - \frac{29A}{512})(c-ic \tan(fx+e))^{5/2} + 8}{3} \right) \right)$
default	$2ic^3 \left(\frac{iB(c-ic \tan(fx+e))^{3/2}}{3} + 7i\sqrt{c-ic \tan(fx+e)} Bc + \sqrt{c-ic \tan(fx+e)} cA - 8c^2 \left(\frac{8(-\frac{81iB}{512} - \frac{29A}{512})(c-ic \tan(fx+e))^{5/2} + 8}{3} \right) \right)$

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2)/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output
$$2*I/f/a^3*c^3*(1/3*I*B*(c-I*c*tan(f*x+e))^(3/2)+7*I*(c-I*c*tan(f*x+e))^(1/2)*B*c+(c-I*c*tan(f*x+e))^(1/2)*c*A-8*c^2*(8*((-81/512*I*B-29/512*A)*(c-I*c*tan(f*x+e))^(5/2)+(53/96*I*B*c+17/96*c*A)*(c-I*c*tan(f*x+e))^(3/2)+(-63/128*I*B*c^2-19/128*c^2*A)*(c-I*c*tan(f*x+e))^(1/2)))/(c+I*c*tan(f*x+e))^3+35/16*(1/8*A+5/8*I*B)*2^(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2)))$$

3.779.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^3} dx$$

3.779.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 471 vs. $2(228) = 456$.

Time = 0.28 (sec) , antiderivative size = 471, normalized size of antiderivative = 1.62

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^3} dx =$$

$$105 \sqrt{\frac{1}{2}} (a^3 f e^{(8i f x + 8i e)} + a^3 f e^{(6i f x + 6i e)}) \sqrt{-\frac{(A^2 + 10i AB - 25 B^2)c^9}{a^6 f^2}} \log \left(\frac{35 \left((i A - 5 B)c^5 + \sqrt{2} \sqrt{\frac{1}{2}} (a^3 f e^{(2i f x + 2i e)} + a^3 f e^{(4i f x + 4i e)}) \right)}{\dots} \right)$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2)/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output `-1/24*(105*sqrt(1/2)*(a^3*f*e^(8*I*f*x + 8*I*e) + a^3*f*e^(6*I*f*x + 6*I*e))*sqrt(-(A^2 + 10*I*A*B - 25*B^2)*c^9/(a^6*f^2))*log(-35/2*((I*A - 5*B)*c^5 + sqrt(2)*sqrt(1/2)*(a^3*f*e^(2*I*f*x + 2*I*e) + a^3*f)*sqrt(-(A^2 + 10*I*A*B - 25*B^2)*c^9/(a^6*f^2))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))*e^(-I*f*x - I*e)/(a^3*f)) - 105*sqrt(1/2)*(a^3*f*e^(8*I*f*x + 8*I*e) + a^3*f*e^(6*I*f*x + 6*I*e))*sqrt(-(A^2 + 10*I*A*B - 25*B^2)*c^9/(a^6*f^2))*log(-35/2*((I*A - 5*B)*c^5 - sqrt(2)*sqrt(1/2)*(a^3*f*e^(2*I*f*x + 2*I*e) + a^3*f)*sqrt(-(A^2 + 10*I*A*B - 25*B^2)*c^9/(a^6*f^2))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))*e^(-I*f*x - I*e)/(a^3*f)) + sqrt(2)*(105*(-I*A + 5*B)*c^4*e^(8*I*f*x + 8*I*e) + 140*(-I*A + 5*B)*c^4*e^(6*I*f*x + 6*I*e) + 21*(-I*A + 5*B)*c^4*e^(4*I*f*x + 4*I*e) + 6*(I*A - 5*B)*c^4*e^(2*I*f*x + 2*I*e) + 8*(-I*A + B)*c^4)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(a^3*f*e^(8*I*f*x + 8*I*e) + a^3*f*e^(6*I*f*x + 6*I*e))`

3.779.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^3} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(9/2)/(a+I*a*tan(f*x+e))**3,x)`

3.779. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^3} dx$

output Timed out

3.779.7 Maxima [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 266, normalized size of antiderivative = 0.91

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^3} dx = \frac{i \left(\frac{105 \sqrt{2} (A + 5i B) c^{11/2} \log\left(-\frac{\sqrt{2}\sqrt{c} - \sqrt{-i c \tan(fx+e)+c}}{\sqrt{2}\sqrt{c} + \sqrt{-i c \tan(fx+e)+c}}\right)}{a^3} - \frac{4(3(-i c \tan(fx+e) + c)^{5/2} (29A + 81i B) c^6 - 16(-i c \tan(fx+e) + c)^{3/2} (17A + 53i B) c^7 + 12 \sqrt{-i c \tan(fx+e) + c} (19A + 63i B) c^8)}{((-i c \tan(fx+e) + c)^3 a^3 - 6(-i c \tan(fx+e) + c)^2 a^3 c + 12(-i c \tan(fx+e) + c) a^3 c^2 - 8a^3 c^3) + 32(i(-i c \tan(fx+e) + c)^{3/2} B c^4 + 3 \sqrt{-i c \tan(fx+e) + c} (A + 7i B) c^5)}{a^3} \right)}{c f}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2)/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`

output `1/48*I*(105*sqrt(2)*(A + 5*I*B)*c^(11/2)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a^3 - 4*(3*(-I*c*tan(f*x + e) + c)^(5/2)*(29*A + 81*I*B)*c^6 - 16*(-I*c*tan(f*x + e) + c)^(3/2)*(17*A + 53*I*B)*c^7 + 12*sqrt(-I*c*tan(f*x + e) + c)*(19*A + 63*I*B)*c^8)/((-I*c*tan(f*x + e) + c)^3*a^3 - 6*(-I*c*tan(f*x + e) + c)^2*a^3*c + 12*(-I*c*tan(f*x + e) + c)*a^3*c^2 - 8*a^3*c^3) + 32*(I*(-I*c*tan(f*x + e) + c)^(3/2)*B*c^4 + 3*sqrt(-I*c*tan(f*x + e) + c)*(A + 7*I*B)*c^5)/a^3)/(c*f)`

3.779.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^3} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{9/2}}{(i a \tan(fx + e) + a)^3} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2)/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(9/2)/(I*a*tan(f*x + e) + a)^3, x)`

3.779.9 Mupad [B] (verification not implemented)

Time = 9.17 (sec) , antiderivative size = 441, normalized size of antiderivative = 1.52

$$\int \frac{(A + B \tan(e + fx))(c - i \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^3} dx = \frac{\frac{A c^7 \sqrt{c - c \tan(e + fx) i} 19i}{a^3 f} - \frac{A c^6 (c - c \tan(e + fx) i)^{3/2} 68i}{3 a^3 f}}{6 c (c - c \tan(e + fx) i)^2 - 12 c^2 (c - c \tan(e + fx) i) - (c - c \tan(e + fx) i)^3 + 8 c^3} - \frac{63 B c^7 \sqrt{c - c \tan(e + fx) i} - \frac{212 B c^6 (c - c \tan(e + fx) i)^{3/2}}{3} + \frac{81 B c^5 (c - c \tan(e + fx) i)^{5/2}}{4}}{8 a^3 c^3 f - a^3 f (c - c \tan(e + fx) i)^3 + 6 a^3 c f (c - c \tan(e + fx) i)^2 - 12 a^3 c^2 f (c - c \tan(e + fx) i)} + \frac{A c^4 \sqrt{c - c \tan(e + fx) i} 2i}{a^3 f} - \frac{14 B c^4 \sqrt{c - c \tan(e + fx) i}}{a^3 f} - \frac{2 B c^3 (c - c \tan(e + fx) i)^{3/2}}{3 a^3 f} - \frac{\sqrt{2} A (-c)^{9/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) i}}{2 \sqrt{-c}}\right)}{8 a^3 f} 35i - \frac{\sqrt{2} B c^{9/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) i} i}{2 \sqrt{c}}\right)}{8 a^3 f} 175i$$

```
input int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*i)^(9/2))/(a + a*tan(e + f*x)*i)^3,x)
```

```
output ((A*c^7*(c - c*tan(e + f*x)*i)^(1/2)*19i)/(a^3*f) - (A*c^6*(c - c*tan(e + f*x)*i)^(3/2)*68i)/(3*a^3*f) + (A*c^5*(c - c*tan(e + f*x)*i)^(5/2)*29i)/(4*a^3*f))/(6*c*(c - c*tan(e + f*x)*i)^2 - 12*c^2*(c - c*tan(e + f*x)*i) - (c - c*tan(e + f*x)*i)^3 + 8*c^3) - (63*B*c^7*(c - c*tan(e + f*x)*i)^(1/2) - (212*B*c^6*(c - c*tan(e + f*x)*i)^(3/2))/3 + (81*B*c^5*(c - c*tan(e + f*x)*i)^(5/2))/4)/(8*a^3*c^3*f - a^3*f*(c - c*tan(e + f*x)*i)^3 + 6*a^3*c*f*(c - c*tan(e + f*x)*i)^2 - 12*a^3*c^2*f*(c - c*tan(e + f*x)*i)) + (A*c^4*(c - c*tan(e + f*x)*i)^(1/2)*2i)/(a^3*f) - (14*B*c^4*(c - c*tan(e + f*x)*i)^(1/2))/(a^3*f) - (2*B*c^3*(c - c*tan(e + f*x)*i)^(3/2))/(3*a^3*f) - (2^(1/2)*A*(-c)^(9/2)*atan((2^(1/2)*(c - c*tan(e + f*x)*i)^(1/2))/(2*(-c)^(1/2)))*35i)/(8*a^3*f) - (2^(1/2)*B*c^(9/2)*atan((2^(1/2)*(c - c*tan(e + f*x)*i)^(1/2)*i)/(2*c^(1/2)))*175i)/(8*a^3*f)
```

3.780
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^3} dx$$

3.780.1 Optimal result 7065
 3.780.2 Mathematica [A] (verified) 7066
 3.780.3 Rubi [A] (verified) 7066
 3.780.4 Maple [A] (verified) 7069
 3.780.5 Fricas [B] (verification not implemented) 7069
 3.780.6 Sympy [F] 7070
 3.780.7 Maxima [A] (verification not implemented) 7071
 3.780.8 Giac [F] 7071
 3.780.9 Mupad [B] (verification not implemented) 7072

3.780.1 Optimal result

Integrand size = 43, antiderivative size = 252

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2}}{(a + ia \tan(e + fx))^3} dx =$$

$$\frac{5(iA - 13B)c^{7/2} \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{8\sqrt{2}a^3 f}$$

$$+ \frac{5(iA - 13B)c^3 \sqrt{c - ic \tan(e + fx)}}{16a^3 f} + \frac{5(iA - 13B)c^2 (c - ic \tan(e + fx))^{3/2}}{48a^3 f (1 + i \tan(e + fx))}$$

$$- \frac{(iA - 13B)c (c - ic \tan(e + fx))^{5/2}}{24a^3 f (1 + i \tan(e + fx))^2} + \frac{(iA - B)(c - ic \tan(e + fx))^{7/2}}{6a^3 f (1 + i \tan(e + fx))^3}$$

output

```
-5/16*(I*A-13*B)*c^(7/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a^3/f*2^(1/2)+5/16*(I*A-13*B)*c^3*(c-I*c*tan(f*x+e))^(1/2)/a^3/f+5/48*(I*A-13*B)*c^2*(c-I*c*tan(f*x+e))^(3/2)/a^3/f/(1+I*tan(f*x+e))-1/24*(I*A-13*B)*c*(c-I*c*tan(f*x+e))^(5/2)/a^3/f/(1+I*tan(f*x+e))^2+1/6*(I*A-B)*(c-I*c*tan(f*x+e))^(7/2)/a^3/f/(1+I*tan(f*x+e))^3
```


3.780.2 Mathematica [A] (verified)

Time = 6.71 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.77

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^3} dx = \frac{c^3 \sec^3(e + fx) \left(15\sqrt{2}(A + 13iB)\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{c - i c \tan(e + fx)}}{\sqrt{2}\sqrt{a + i a \tan(e + fx)}}\right)\right)}{(a + i a \tan(e + fx))^3}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2))/(a + I*a*Tan[e + f*x])^3,x]`

output `(c^3*Sec[e + f*x]^3*(15*Sqrt[2]*(A + (13*I)*B)*Sqrt[c]*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])]*(Cos[3*(e + f*x)] + I*Sin[3*(e + f*x)]) - (3*(A + (13*I)*B)*Cos[e + f*x] + (23*A + (203*I)*B)*Cos[3*(e + f*x)] + (2*I)*(7*A + (139*I)*B + (7*A + (187*I)*B)*Cos[2*(e + f*x)])*Sin[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])/(48*a^3*f*(-I + Tan[e + f*x])^3)`

3.780.3 Rubi [A] (verified)Time = 0.39 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.86, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.209$, Rules used = {3042, 4071, 27, 87, 51, 51, 60, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c - i c \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{(a + i a \tan(e + fx))^3} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c - i c \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{(a + i a \tan(e + fx))^3} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{a^4(i \tan(e + fx) + 1)^4} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{27} \\ & \frac{c \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(i \tan(e + fx) + 1)^4} d \tan(e + fx)}{a^3 f} \end{aligned}$$

3.780. $\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^3} dx$

$$\begin{aligned}
& \downarrow 87 \\
& \frac{c \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{6c(1+i \tan(e+fx))^3} - \frac{1}{12}(A+13iB) \int \frac{(c-ic \tan(e+fx))^{5/2}}{(i \tan(e+fx)+1)^3} d \tan(e+fx) \right)}{a^3 f} \\
& \downarrow 51 \\
& \frac{c \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{6c(1+i \tan(e+fx))^3} - \frac{1}{12}(A+13iB) \left(\frac{i(c-ic \tan(e+fx))^{5/2}}{2(1+i \tan(e+fx))^2} - \frac{5}{4}c \int \frac{(c-ic \tan(e+fx))^{3/2}}{(i \tan(e+fx)+1)^2} d \tan(e+fx) \right) \right)}{a^3 f} \\
& \downarrow 51 \\
& \frac{c \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{6c(1+i \tan(e+fx))^3} - \frac{1}{12}(A+13iB) \left(\frac{i(c-ic \tan(e+fx))^{5/2}}{2(1+i \tan(e+fx))^2} - \frac{5}{4}c \left(\frac{i(c-ic \tan(e+fx))^{3/2}}{1+i \tan(e+fx)} - \frac{3}{2}c \int \frac{\sqrt{c-ic \tan(e+fx)}}{i \tan(e+fx)+1} d \tan(e+fx) \right) \right) \right)}{a^3 f} \\
& \downarrow 60 \\
& \frac{c \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{6c(1+i \tan(e+fx))^3} - \frac{1}{12}(A+13iB) \left(\frac{i(c-ic \tan(e+fx))^{5/2}}{2(1+i \tan(e+fx))^2} - \frac{5}{4}c \left(\frac{i(c-ic \tan(e+fx))^{3/2}}{1+i \tan(e+fx)} - \frac{3}{2}c \left(2c \int \frac{1}{(i \tan(e+fx)+1)\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) \right) \right) \right) \right)}{a^3 f} \\
& \downarrow 73 \\
& \frac{c \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{6c(1+i \tan(e+fx))^3} - \frac{1}{12}(A+13iB) \left(\frac{i(c-ic \tan(e+fx))^{5/2}}{2(1+i \tan(e+fx))^2} - \frac{5}{4}c \left(\frac{i(c-ic \tan(e+fx))^{3/2}}{1+i \tan(e+fx)} - \frac{3}{2}c \left(4i \int \frac{1}{2 - \frac{c-ic \tan(e+fx)}{c}} d \tan(e+fx) \right) \right) \right) \right)}{a^3 f} \\
& \downarrow 219 \\
& \frac{c \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{6c(1+i \tan(e+fx))^3} - \frac{1}{12}(A+13iB) \left(\frac{i(c-ic \tan(e+fx))^{5/2}}{2(1+i \tan(e+fx))^2} - \frac{5}{4}c \left(\frac{i(c-ic \tan(e+fx))^{3/2}}{1+i \tan(e+fx)} - \frac{3}{2}c \left(2i\sqrt{2}\sqrt{c} \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(e+fx)}}{c} \right) \right) \right) \right) \right)}{a^3 f}
\end{aligned}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2))/(a + I*a*Tan[e + f*x])^3, x]`

output `(c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(7/2))/(6*c*(1 + I*Tan[e + f*x])^3) - ((A + (13*I)*B)*(((I/2)*(c - I*c*Tan[e + f*x])^(5/2))/(1 + I*Tan[e + f*x])^2 - (5*c*((I*(c - I*c*Tan[e + f*x])^(3/2))/(1 + I*Tan[e + f*x]) - (3*c*((2*I)*Sqrt[2]*Sqrt[c]*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])]) - (2*I)*Sqrt[c - I*c*Tan[e + f*x]]))/2))/4))/12))/(a^3*f)`

3.780. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^3} dx$

3.780.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 51 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`
- rule 60 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x],
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.780.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.67

method	result
derivativedivides	$2ic^3 \left(i\sqrt{c-ic \tan(fx+e)} B + c \left(\frac{8 \left(\frac{47iB}{128} + \frac{11A}{128} \right) (c-ic \tan(fx+e))^{\frac{5}{2}} + 8 \left(-\frac{29}{24} iBc - \frac{5}{24} cA \right) (c-ic \tan(fx+e))^{\frac{3}{2}} + 8 \left(\frac{33}{32} iB c^2 + \frac{5}{32} c^2 \right)}{(c+ic \tan(fx+e))^3} \right) \right) f a^3$
default	$2ic^3 \left(i\sqrt{c-ic \tan(fx+e)} B + c \left(\frac{8 \left(\frac{47iB}{128} + \frac{11A}{128} \right) (c-ic \tan(fx+e))^{\frac{5}{2}} + 8 \left(-\frac{29}{24} iBc - \frac{5}{24} cA \right) (c-ic \tan(fx+e))^{\frac{3}{2}} + 8 \left(\frac{33}{32} iB c^2 + \frac{5}{32} c^2 \right)}{(c+ic \tan(fx+e))^3} \right) \right) f a^3$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^3,x,metho
d=_RETURNVERBOSE)
```

```
output 2*I/f/a^3*c^3*(I*(c-I*c*tan(f*x+e))^(1/2)*B+c*(8*((47/128*I*B+11/128*A)*(c
-I*c*tan(f*x+e))^(5/2)+(-29/24*I*B*c-5/24*c*A)*(c-I*c*tan(f*x+e))^(3/2)+(3
3/32*I*B*c^2+5/32*c^2*A)*(c-I*c*tan(f*x+e))^(1/2))/(c+I*c*tan(f*x+e))^3-5/
4*(13/8*I*B+1/8*A)*2^(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^
(1/2)/c^(1/2))))
```

3.780.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 402 vs. 2(197) = 394.

Time = 0.27 (sec) , antiderivative size = 402, normalized size of antiderivative = 1.60

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2}}{(a + ia \tan(e + fx))^3} dx =$$

$$\left(15 \sqrt{\frac{1}{2}} a^3 f \sqrt{-\frac{(A^2 + 26iAB - 169B^2)c^7}{a^6 f^2}} e^{(6i fx + 6ie)} \log \left(-\frac{5 \left((iA - 13B)c^4 + \sqrt{2} \sqrt{\frac{1}{2}} (a^3 f e^{(2i fx + 2ie)} + a^3 f) \sqrt{-\frac{(A^2 + 26iAB - 169B^2)c^7}{a^6 f^2}} \right)}{4 a^3 f} \right) \right)$$

3.780. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^3} dx$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output
$$\begin{aligned} & -1/48*(15*\sqrt{1/2}*a^3*f*\sqrt{-(A^2 + 26*I*A*B - 169*B^2)*c^7/(a^6*f^2)}* \\ & e^{(6*I*f*x + 6*I*e)}*\log(-5/4*((I*A - 13*B)*c^4 + \sqrt{2}*\sqrt{1/2}*(a^3*f* \\ & e^{(2*I*f*x + 2*I*e)} + a^3*f)*\sqrt{-(A^2 + 26*I*A*B - 169*B^2)*c^7/(a^6*f^2)} \\ &))*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)})*e^{-(I*f*x - I*e)/(a^3*f)} - 15*\sqrt{1/2}* \\ & a^3*f*\sqrt{-(A^2 + 26*I*A*B - 169*B^2)*c^7/(a^6*f^2)}*e^{(6*I*f*x + 6*I*e)}* \\ & \log(-5/4*((I*A - 13*B)*c^4 - \sqrt{2}*\sqrt{1/2}*(a^3*f*e^{(2*I*f*x + 2*I*e)} + \\ & a^3*f)*\sqrt{-(A^2 + 26*I*A*B - 169*B^2)*c^7/(a^6*f^2)})*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)})* \\ & e^{-(I*f*x - I*e)/(a^3*f)} + \sqrt{2}*(15*(-I*A + 13*B)*c^3*e^{(6*I*f*x + 6*I*e)} + \\ & 5*(-I*A + 13*B)*c^3*e^{(4*I*f*x + 4*I*e)} + 2*(I*A - 13*B)*c^3*e^{(2*I*f*x + 2*I*e)} + \\ & 8*(-I*A + B)*c^3)*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)})*e^{-(6*I*f*x - 6*I*e)/(a^3*f)} \end{aligned}$$

3.780.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^3} dx = i \left(\int \frac{A c^3 \sqrt{-i c \tan(e + fx) + c}}{\tan^3(e + fx) - 3i \tan^2(e + fx) - 3 \tan(e + fx) + i} dx + \int \left(-\frac{3A}{\tan^3(e + fx)} \right) dx \right)$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(7/2)/(a+I*a*tan(f*x+e))**3,x)`

output
$$\begin{aligned} & I*(\text{Integral}(A*c**3*\sqrt{-I*c*\tan(e + f*x) + c}/(\tan(e + f*x)**3 - 3*I*\tan(e + f*x)**2 - 3*\tan(e + f*x) + I), x) + \text{Integral}(-3*A*c**3*\sqrt{-I*c*\tan(e + f*x) + c}*\tan(e + f*x)**2/(\tan(e + f*x)**3 - 3*I*\tan(e + f*x)**2 - 3*\tan(e + f*x) + I), x) + \text{Integral}(B*c**3*\sqrt{-I*c*\tan(e + f*x) + c}*\tan(e + f*x)/(\tan(e + f*x)**3 - 3*I*\tan(e + f*x)**2 - 3*\tan(e + f*x) + I), x) + \text{Integral}(-3*B*c**3*\sqrt{-I*c*\tan(e + f*x) + c}*\tan(e + f*x)**3/(\tan(e + f*x)**3 - 3*I*\tan(e + f*x)**2 - 3*\tan(e + f*x) + I), x) + \text{Integral}(-3*I*A*c**3*\sqrt{-I*c*\tan(e + f*x) + c}*\tan(e + f*x)/(\tan(e + f*x)**3 - 3*I*\tan(e + f*x)**2 - 3*\tan(e + f*x) + I), x) + \text{Integral}(I*A*c**3*\sqrt{-I*c*\tan(e + f*x) + c}*\tan(e + f*x)**3/(\tan(e + f*x)**3 - 3*I*\tan(e + f*x)**2 - 3*\tan(e + f*x) + I), x) + \text{Integral}(-3*I*B*c**3*\sqrt{-I*c*\tan(e + f*x) + c}*\tan(e + f*x)**2/(\tan(e + f*x)**3 - 3*I*\tan(e + f*x)**2 - 3*\tan(e + f*x) + I), x) + \text{Integral}(I*B*c**3*\sqrt{-I*c*\tan(e + f*x) + c}*\tan(e + f*x)**4/(\tan(e + f*x)**3 - 3*I*\tan(e + f*x)**2 - 3*\tan(e + f*x) + I), x))/a**3 \end{aligned}$$

3.780.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^3} dx$$

3.780.7 Maxima [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 240, normalized size of antiderivative = 0.95

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^3} dx = \frac{i \left(\frac{15 \sqrt{2} (A + 13i B) c^{9/2} \log \left(\frac{-\sqrt{2} \sqrt{c} - \sqrt{-i c \tan(fx + e) + c}}{\sqrt{2} \sqrt{c} + \sqrt{-i c \tan(fx + e) + c}} \right)}{a^3} + \frac{192i \sqrt{-i c \tan(fx + e) + c}}{a^3} \right)}{a^3}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^3,x
, algorithm="maxima")`

output `1/96*I*(15*sqrt(2)*(A + 13*I*B)*c^(9/2)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a^3 + 192*I*sqrt(-I*c*tan(f*x + e) + c)*B*c^4/a^3 - 4*(3*(-I*c*tan(f*x + e) + c)^(5/2)*(11*A + 47*I*B)*c^5 - 16*(-I*c*tan(f*x + e) + c)^(3/2)*(5*A + 29*I*B)*c^6 + 12*sqrt(-I*c*tan(f*x + e) + c)*(5*A + 33*I*B)*c^7)/((-I*c*tan(f*x + e) + c)^3*a^3 - 6*(-I*c*tan(f*x + e) + c)^2*a^3*c + 12*(-I*c*tan(f*x + e) + c)*a^3*c^2 - 8*a^3*c^3)/(c*f)`

3.780.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^3} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{7/2}}{(i a \tan(fx + e) + a)^3} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^3,x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(7/2)/(I*a*tan(f*x + e) + a)^3, x)`

3.780.9 Mupad [B] (verification not implemented)

Time = 9.28 (sec) , antiderivative size = 386, normalized size of antiderivative = 1.53

$$\int \frac{(A + B \tan(e + fx))(c - i \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^3} dx = \frac{\frac{A c^6 \sqrt{c - c \tan(e + fx) i} 5i}{2 a^3 f} - \frac{A c^5 (c - c \tan(e + fx) i)^{3/2} 10i}{3 a^3 f}}{6 c (c - c \tan(e + fx) i)^2 - 12 c^2 (c - c \tan(e + fx) i)} - \frac{\frac{33 B c^6 \sqrt{c - c \tan(e + fx) i}}{2} - \frac{58 B c^5 (c - c \tan(e + fx) i)^{3/2}}{3} + \frac{47 B c^4 (c - c \tan(e + fx) i)^{5/2}}{8}}{8 a^3 c^3 f - a^3 f (c - c \tan(e + fx) i)^3 + 6 a^3 c f (c - c \tan(e + fx) i)^2 - 12 a^3 c^2 f (c - c \tan(e + fx) i)} - \frac{2 B c^3 \sqrt{c - c \tan(e + fx) i}}{a^3 f} + \frac{\sqrt{2} A (-c)^{7/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) i}}{2 \sqrt{-c}}\right) 5i}{16 a^3 f} + \frac{65 \sqrt{2} B c^{7/2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) i}}{2 \sqrt{c}}\right)}{16 a^3 f}$$

```
input int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(7/2))/(a + a*tan(e + f*x)*1i)^3,x)
```

```
output ((A*c^6*(c - c*tan(e + f*x)*1i)^(1/2)*5i)/(2*a^3*f) - (A*c^5*(c - c*tan(e + f*x)*1i)^(3/2)*10i)/(3*a^3*f) + (A*c^4*(c - c*tan(e + f*x)*1i)^(5/2)*11i)/(8*a^3*f))/(6*c*(c - c*tan(e + f*x)*1i)^2 - 12*c^2*(c - c*tan(e + f*x)*1i) - (c - c*tan(e + f*x)*1i)^3 + 8*c^3) - ((33*B*c^6*(c - c*tan(e + f*x)*1i)^(1/2))/2 - (58*B*c^5*(c - c*tan(e + f*x)*1i)^(3/2))/3 + (47*B*c^4*(c - c*tan(e + f*x)*1i)^(5/2))/8)/(8*a^3*c^3*f - a^3*f*(c - c*tan(e + f*x)*1i)^3 + 6*a^3*c*f*(c - c*tan(e + f*x)*1i)^2 - 12*a^3*c^2*f*(c - c*tan(e + f*x)*1i)) - (2*B*c^3*(c - c*tan(e + f*x)*1i)^(1/2))/(a^3*f) + (2^(1/2)*A*(-c)^(7/2)*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*(-c)^(1/2)))*5i)/(16*a^3*f) + (65*2^(1/2)*B*c^(7/2)*atanh((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*c^(1/2)))/(16*a^3*f))
```

3.781
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^3} dx$$

3.781.1 Optimal result 7073
 3.781.2 Mathematica [A] (verified) 7073
 3.781.3 Rubi [A] (verified) 7074
 3.781.4 Maple [A] (verified) 7077
 3.781.5 Fricas [B] (verification not implemented) 7077
 3.781.6 Sympy [F] 7078
 3.781.7 Maxima [A] (verification not implemented) 7079
 3.781.8 Giac [F] 7079
 3.781.9 Mupad [B] (verification not implemented) 7080

3.781.1 Optimal result

Integrand size = 43, antiderivative size = 213

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^3} dx = \frac{(iA + 11B)c^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{16\sqrt{2}a^3 f} - \frac{(iA + 11B)c^2 \sqrt{c - ic \tan(e + fx)}}{16a^3 f(1 + i \tan(e + fx))} + \frac{(iA + 11B)c(c - ic \tan(e + fx))^{3/2}}{24a^3 f(1 + i \tan(e + fx))^2} + \frac{(iA - B)(c - ic \tan(e + fx))^{5/2}}{6a^3 f(1 + i \tan(e + fx))^3}$$

```
output 1/32*(I*A+11*B)*c^(5/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a^3/f*2^(1/2)-1/16*(I*A+11*B)*c^2*(c-I*c*tan(f*x+e))^(1/2)/a^3/f/(1+I*tan(f*x+e))+1/24*(I*A+11*B)*c*(c-I*c*tan(f*x+e))^(3/2)/a^3/f/(1+I*tan(f*x+e))^2+1/6*(I*A-B)*(c-I*c*tan(f*x+e))^(5/2)/a^3/f/(1+I*tan(f*x+e))^3
```

3.781.2 Mathematica [A] (verified)

Time = 6.50 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.82

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^3} dx = \frac{c^2 \sec^3(e + fx) \left(3\sqrt{2}(A - 11iB)\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right) (\cos(3(e + fx)) + i \sin(3(e + fx))) + 2 \cos(e + fx) \right)}{96a^3 f(-i + \dots)}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2))/(a + I*a*Tan[e + f*x])^3,x]`

output `-1/96*(c^2*Sec[e + f*x]^3*(3*Sqrt[2]*(A - (11*I)*B)*Sqrt[c]*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])]*(Cos[3*(e + f*x)] + I*Sin[3*(e + f*x)]) + 2*Cos[e + f*x]*(2*(A - (11*I)*B) + (5*A + (41*I)*B)*Cos[2*(e + f*x)]) + ((-11*I)*A - 25*B)*Sin[2*(e + f*x)])*Sqrt[c - I*c*Tan[e + f*x]])/(a^3*f*(-I + Tan[e + f*x])^3)`

3.781.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 189, normalized size of antiderivative = 0.89, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.186$, Rules used = {3042, 4071, 27, 87, 51, 51, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - ict \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow 3042 \\
 & \int \frac{(c - ict \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow 4071 \\
 & \frac{ac \int \frac{(A + B \tan(e + fx))(c - ict \tan(e + fx))^{3/2}}{a^4 (i \tan(e + fx) + 1)^4} d \tan(e + fx)}{f} \\
 & \quad \downarrow 27 \\
 & \frac{c \int \frac{(A + B \tan(e + fx))(c - ict \tan(e + fx))^{3/2}}{(i \tan(e + fx) + 1)^4} d \tan(e + fx)}{a^3 f} \\
 & \quad \downarrow 87 \\
 & \frac{c \left(\frac{1}{12} (A - 11iB) \int \frac{(c - ict \tan(e + fx))^{3/2}}{(i \tan(e + fx) + 1)^3} d \tan(e + fx) + \frac{(-B + iA)(c - ict \tan(e + fx))^{5/2}}{6c(1 + i \tan(e + fx))^3} \right)}{a^3 f} \\
 & \quad \downarrow 51
 \end{aligned}$$

3.781. $\int \frac{(A + B \tan(e + fx))(c - ict \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^3} dx$

$$\frac{c \left(\frac{1}{12} (A - 11iB) \left(\frac{i(c - ic \tan(e+fx))^{3/2}}{2(1+i \tan(e+fx))^2} - \frac{3}{4} c \int \frac{\sqrt{c - ic \tan(e+fx)}}{(i \tan(e+fx) + 1)^2} d \tan(e+fx) \right) + \frac{(-B + iA)(c - ic \tan(e+fx))^{5/2}}{6c(1+i \tan(e+fx))^3} \right)}{a^3 f}$$

↓ 51

$$\frac{c \left(\frac{1}{12} (A - 11iB) \left(\frac{i(c - ic \tan(e+fx))^{3/2}}{2(1+i \tan(e+fx))^2} - \frac{3}{4} c \left(\frac{i \sqrt{c - ic \tan(e+fx)}}{1+i \tan(e+fx)} - \frac{1}{2} c \int \frac{1}{(i \tan(e+fx) + 1) \sqrt{c - ic \tan(e+fx)}} d \tan(e+fx) \right) \right) + \frac{(-B + iA)(c - ic \tan(e+fx))^{5/2}}{6c(1+i \tan(e+fx))^3} \right)}{a^3 f}$$

↓ 73

$$\frac{c \left(\frac{1}{12} (A - 11iB) \left(\frac{i(c - ic \tan(e+fx))^{3/2}}{2(1+i \tan(e+fx))^2} - \frac{3}{4} c \left(\frac{i \sqrt{c - ic \tan(e+fx)}}{1+i \tan(e+fx)} - i \int \frac{1}{2 - \frac{c - ic \tan(e+fx)}{c}} d \sqrt{c - ic \tan(e+fx)} \right) \right) + \frac{(-B + iA)(c - ic \tan(e+fx))^{5/2}}{6c(1+i \tan(e+fx))^3} \right)}{a^3 f}$$

↓ 219

$$\frac{c \left(\frac{1}{12} (A - 11iB) \left(\frac{i(c - ic \tan(e+fx))^{3/2}}{2(1+i \tan(e+fx))^2} - \frac{3}{4} c \left(\frac{i \sqrt{c - ic \tan(e+fx)}}{1+i \tan(e+fx)} - \frac{i \sqrt{c} \operatorname{arctanh} \left(\frac{\sqrt{c - ic \tan(e+fx)}}{\sqrt{2} \sqrt{c}} \right)}{\sqrt{2}} \right) \right) + \frac{(-B + iA)(c - ic \tan(e+fx))^{5/2}}{6c(1+i \tan(e+fx))^3} \right)}{a^3 f}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2))/(a + I*a*Tan[e + f*x])^3,x]`

output `(c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(5/2))/(6*c*(1 + I*Tan[e + f*x])^3) + ((A - (11*I)*B)*(((I/2)*(c - I*c*Tan[e + f*x])^(3/2))/(1 + I*Tan[e + f*x])^2 - (3*c*(((-I)*Sqrt[c]*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])))/Sqrt[2] + (I*Sqrt[c - I*c*Tan[e + f*x]])/(1 + I*Tan[e + f*x])))/4)/12))/(a^3*f)`

3.781.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 51 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`
- rule 73 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.781.4 Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.69

method	result
derivativedivides	$\frac{2ic^3 \left(\frac{8 \left(\frac{21iB}{256} + \frac{A}{256} \right) (c-ic \tan(fx+e))^{\frac{5}{2}} + 8 \left(-\frac{11}{48} iBc + \frac{1}{48} cA \right) (c-ic \tan(fx+e))^{\frac{3}{2}} + 8 \left(\frac{11}{64} iBc^2 - \frac{1}{64} c^2A \right) \sqrt{c-ic \tan(fx+e)} + \left(-\frac{11i}{8} \right)}{(c+ic \tan(fx+e))^3} \right)}{fa^3}$
default	$\frac{2ic^3 \left(\frac{8 \left(\frac{21iB}{256} + \frac{A}{256} \right) (c-ic \tan(fx+e))^{\frac{5}{2}} + 8 \left(-\frac{11}{48} iBc + \frac{1}{48} cA \right) (c-ic \tan(fx+e))^{\frac{3}{2}} + 8 \left(\frac{11}{64} iBc^2 - \frac{1}{64} c^2A \right) \sqrt{c-ic \tan(fx+e)} + \left(-\frac{11i}{8} \right)}{(c+ic \tan(fx+e))^3} \right)}{fa^3}$

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output `2*I/f/a^3*c^3*(8*((21/256*I*B+1/256*A)*(c-I*c*tan(f*x+e))^(5/2)+(-11/48*I*B*c+1/48*c*A)*(c-I*c*tan(f*x+e))^(3/2)+(11/64*I*B*c^2-1/64*c^2*A)*(c-I*c*tan(f*x+e))^(1/2))/(c+I*c*tan(f*x+e))^3+1/8*(-11/8*I*B+1/8*A)*2^(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2)))`

3.781.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 403 vs. 2(166) = 332.

Time = 0.27 (sec) , antiderivative size = 403, normalized size of antiderivative = 1.89

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^3} dx = \frac{\left(3 \sqrt{\frac{1}{2}} a^3 f \sqrt{-\frac{(A^2 - 22iAB - 121B^2)c^5}{a^6 f^2}} e^{(6i fx + 6i e)} \log \left(\frac{(iA - \dots)}{\dots} \right) \right)}{\dots}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^3,x,algorithm="fracas")`

```
output 1/96*(3*sqrt(1/2)*a^3*f*sqrt(-(A^2 - 22*I*A*B - 121*B^2)*c^5/(a^6*f^2))*e^
(6*I*f*x + 6*I*e)*log(1/8*((I*A + 11*B)*c^3 + sqrt(2)*sqrt(1/2)*(a^3*f*e^(
2*I*f*x + 2*I*e) + a^3*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(A^2 - 2
2*I*A*B - 121*B^2)*c^5/(a^6*f^2))))*e^(-I*f*x - I*e)/(a^3*f) - 3*sqrt(1/2)
*a^3*f*sqrt(-(A^2 - 22*I*A*B - 121*B^2)*c^5/(a^6*f^2))*e^(6*I*f*x + 6*I*e)
*log(1/8*((I*A + 11*B)*c^3 - sqrt(2)*sqrt(1/2)*(a^3*f*e^(2*I*f*x + 2*I*e)
+ a^3*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(A^2 - 22*I*A*B - 121*B^2
)*c^5/(a^6*f^2))))*e^(-I*f*x - I*e)/(a^3*f) - sqrt(2)*(3*(I*A + 11*B)*c^2*
e^(6*I*f*x + 6*I*e) - (-I*A - 11*B)*c^2*e^(4*I*f*x + 4*I*e) + 2*(-5*I*A -
7*B)*c^2*e^(2*I*f*x + 2*I*e) + 8*(-I*A + B)*c^2)*sqrt(c/(e^(2*I*f*x + 2*I*
e) + 1)))e^(-6*I*f*x - 6*I*e)/(a^3*f)
```

3.781.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(a + i a \tan(e + fx))^3} dx = i \left(\int \frac{Ac^2 \sqrt{-i c \tan(e + fx) + c}}{\tan^3(e + fx) - 3i \tan^2(e + fx) - 3 \tan(e + fx) + i} dx + \int \left(-\frac{Ac}{\tan^3(e + fx)} \right) dx \right)$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(5/2)/(a+I*a*tan(f*x+e))**3
,x)
```

```
output I*(Integral(A*c**2*sqrt(-I*c*tan(e + f*x) + c)/(tan(e + f*x)**3 - 3*I*tan(
e + f*x)**2 - 3*tan(e + f*x) + I), x) + Integral(-A*c**2*sqrt(-I*c*tan(e +
f*x) + c)*tan(e + f*x)**2/(tan(e + f*x)**3 - 3*I*tan(e + f*x)**2 - 3*tan(
e + f*x) + I), x) + Integral(B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*
x)/(tan(e + f*x)**3 - 3*I*tan(e + f*x)**2 - 3*tan(e + f*x) + I), x) + Inte
gral(-B*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3/(tan(e + f*x)**3
- 3*I*tan(e + f*x)**2 - 3*tan(e + f*x) + I), x) + Integral(-2*I*A*c**2*sqr
t(-I*c*tan(e + f*x) + c)*tan(e + f*x)/(tan(e + f*x)**3 - 3*I*tan(e + f*x)*
*2 - 3*tan(e + f*x) + I), x) + Integral(-2*I*B*c**2*sqrt(-I*c*tan(e + f*x)
+ c)*tan(e + f*x)**2/(tan(e + f*x)**3 - 3*I*tan(e + f*x)**2 - 3*tan(e + f
*x) + I), x))/a**3
```

3.781. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^3} dx$

3.781.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.00

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(a + i a \tan(e + fx))^3} dx =$$

$$i \left(\frac{3\sqrt{2}(A-11iB)c^{\frac{7}{2}} \log\left(-\frac{\sqrt{2}\sqrt{c}-\sqrt{-ic \tan(fx+e)+c}}{\sqrt{2}\sqrt{c}+\sqrt{-ic \tan(fx+e)+c}}\right)}{a^3} + \frac{4\left(3(-ic \tan(fx+e)+c)^{\frac{5}{2}}(A+21iB)c^4+16(-ic \tan(fx+e)+c)^{\frac{3}{2}}(A-11iB)c^5-12(-ic \tan(fx+e)+c)^{\frac{1}{2}}(A-11iB)c^6\right)}{(-ic \tan(fx+e)+c)^3 a^3 - 6(-ic \tan(fx+e)+c)^2 a^3 c + 12(-ic \tan(fx+e)+c) a^3 c^2 - 8a^3 c^3} \right) / (192 c f)$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^3,x
, algorithm="maxima")
```

```
output -1/192*I*(3*sqrt(2)*(A - 11*I*B)*c^(7/2)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c
*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a^3 +
4*(3*(-I*c*tan(f*x + e) + c)^(5/2)*(A + 21*I*B)*c^4 + 16*(-I*c*tan(f*x +
e) + c)^(3/2)*(A - 11*I*B)*c^5 - 12*sqrt(-I*c*tan(f*x + e) + c)*(A - 11*I*
B)*c^6)/((-I*c*tan(f*x + e) + c)^3*a^3 - 6*(-I*c*tan(f*x + e) + c)^2*a^3*c
+ 12*(-I*c*tan(f*x + e) + c)*a^3*c^2 - 8*a^3*c^3)/(c*f)
```

3.781.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(a + i a \tan(e + fx))^3} dx = \int \frac{(B \tan(fx + e) + A)(-ic \tan(fx + e) + c)^{\frac{5}{2}}}{(ia \tan(fx + e) + a)^3} dx$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^3,x
, algorithm="giac")
```

```
output integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(5/2)/(I*a*tan(f*x
+ e) + a)^3, x)
```

3.781.9 Mupad [B] (verification not implemented)

Time = 9.35 (sec) , antiderivative size = 360, normalized size of antiderivative = 1.69

$$\int \frac{(A + B \tan(e + fx))(c - i \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^3} dx = \frac{-\frac{A c^5 \sqrt{c - c \tan(e + fx) i} i}{4 a^3 f} + \frac{A c^4 (c - c \tan(e + fx) i)^{3/2}}{3 a^3 f}}{6 c (c - c \tan(e + fx) i)^2 - 12 c^2 (c - c \tan(e + fx) i)} - \frac{\frac{11 B c^5 \sqrt{c - c \tan(e + fx) i}}{4} - \frac{11 B c^4 (c - c \tan(e + fx) i)^{3/2}}{3} + \frac{21 B c^3 (c - c \tan(e + fx) i)^{5/2}}{16}}{8 a^3 c^3 f - a^3 f (c - c \tan(e + fx) i)^3 + 6 a^3 c f (c - c \tan(e + fx) i)^2 - 12 a^3 c^2 f (c - c \tan(e + fx) i)} + \frac{\sqrt{2} A (-c)^{5/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) i}}{2 \sqrt{-c}}\right) i}{32 a^3 f} + \frac{11 \sqrt{2} B c^{5/2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) i}}{2 \sqrt{c}}\right)}{32 a^3 f}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(5/2))/(a + a*tan(e + f*x)*1i)^3,x)`

output `((A*c^4*(c - c*tan(e + f*x)*1i)^(3/2)*1i)/(3*a^3*f) - (A*c^5*(c - c*tan(e + f*x)*1i)^(1/2)*1i)/(4*a^3*f) + (A*c^3*(c - c*tan(e + f*x)*1i)^(5/2)*1i)/(16*a^3*f))/(6*c*(c - c*tan(e + f*x)*1i)^2 - 12*c^2*(c - c*tan(e + f*x)*1i) - (c - c*tan(e + f*x)*1i)^3 + 8*c^3) - ((11*B*c^5*(c - c*tan(e + f*x)*1i)^(1/2))/4 - (11*B*c^4*(c - c*tan(e + f*x)*1i)^(3/2))/3 + (21*B*c^3*(c - c*tan(e + f*x)*1i)^(5/2))/16)/(8*a^3*c^3*f - a^3*f*(c - c*tan(e + f*x)*1i)^3 + 6*a^3*c*f*(c - c*tan(e + f*x)*1i)^2 - 12*a^3*c^2*f*(c - c*tan(e + f*x)*1i)) + (2^(1/2)*A*(-c)^(5/2)*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*(-c)^(1/2)))*1i)/(32*a^3*f) + (11*2^(1/2)*B*c^(5/2)*atanh((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*c^(1/2)))/(32*a^3*f)`

3.782
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^3} dx$$

3.782.1 Optimal result 7081
 3.782.2 Mathematica [A] (verified) 7081
 3.782.3 Rubi [A] (verified) 7082
 3.782.4 Maple [A] (verified) 7085
 3.782.5 Fricas [B] (verification not implemented) 7085
 3.782.6 Sympy [F] 7086
 3.782.7 Maxima [A] (verification not implemented) 7086
 3.782.8 Giac [F] 7087
 3.782.9 Mupad [B] (verification not implemented) 7087

3.782.1 Optimal result

Integrand size = 43, antiderivative size = 211

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^3} dx =$$

$$-\frac{(iA + 3B)c^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{32\sqrt{2}a^3 f} + \frac{(iA + 3B)c\sqrt{c - ic \tan(e + fx)}}{8a^3 f(1 + i \tan(e + fx))^2}$$

$$-\frac{(iA + 3B)c\sqrt{c - ic \tan(e + fx)}}{32a^3 f(1 + i \tan(e + fx))} + \frac{(iA - B)(c - ic \tan(e + fx))^{3/2}}{6a^3 f(1 + i \tan(e + fx))^3}$$

output

```
-1/64*(I*A+3*B)*c^(3/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a^3/f*2^(1/2)+1/8*(I*A+3*B)*c*(c-I*c*tan(f*x+e))^(1/2)/a^3/f/(1+I*tan(f*x+e))^2-1/32*(I*A+3*B)*c*(c-I*c*tan(f*x+e))^(1/2)/a^3/f/(1+I*tan(f*x+e))+1/6*(I*A-B)*(c-I*c*tan(f*x+e))^(3/2)/a^3/f/(1+I*tan(f*x+e))^3
```

3.782.2 Mathematica [A] (verified)

Time = 6.44 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.82

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^3} dx = \frac{c \sec^3(e + fx) \left(3\sqrt{2}(A - 3iB)\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)\right)}{(a + ia \tan(e + fx))^3}$$

input

```
Integrate[(((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2))/(a + I*a*Tan[e + f*x])^3,x]
```

3.782.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^3} dx$$

output $(c*\text{Sec}[e + f*x]^3*(3*\text{Sqrt}[2]*(A - (3*I)*B)*\text{Sqrt}[c]*\text{ArcTanh}[\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]]]/(\text{Sqrt}[2]*\text{Sqrt}[c]))*(\text{Cos}[3*(e + f*x)] + I*\text{Sin}[3*(e + f*x)]) - 2*\text{Cos}[e + f*x]*(2*(7*A - (5*I)*B) + (11*A - I*B)*\text{Cos}[2*(e + f*x)] + ((-5*I)*A + 17*B)*\text{Sin}[2*(e + f*x)])*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])/((192*a^3*f*(-I + \text{Tan}[e + f*x])^3)$

3.782.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.93, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.186$, Rules used = {3042, 4071, 27, 87, 51, 52, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx$$

↓ 3042

$$\int \frac{(c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx$$

↓ 4071

$$\frac{ac \int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{a^4 (i \tan(e + fx) + 1)^4} d \tan(e + fx)}{f}$$

↓ 27

$$\frac{c \int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{(i \tan(e + fx) + 1)^4} d \tan(e + fx)}{a^3 f}$$

↓ 87

$$\frac{c \left(\frac{1}{4} (A - 3iB) \int \frac{\sqrt{c - ic \tan(e + fx)}}{(i \tan(e + fx) + 1)^3} d \tan(e + fx) + \frac{(-B + iA)(c - ic \tan(e + fx))^{3/2}}{6c(1 + i \tan(e + fx))^3} \right)}{a^3 f}$$

↓ 51

$$\frac{c \left(\frac{1}{4} (A - 3iB) \left(\frac{i \sqrt{c - ic \tan(e + fx)}}{2(1 + i \tan(e + fx))^2} - \frac{1}{4} c \int \frac{1}{(i \tan(e + fx) + 1)^2 \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) \right) + \frac{(-B + iA)(c - ic \tan(e + fx))^{3/2}}{6c(1 + i \tan(e + fx))^3} \right)}{a^3 f}$$

↓ 52

3.782. $\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^3} dx$

$$\frac{c\left(\frac{1}{4}(A-3iB)\left(\frac{i\sqrt{c-ictan(e+fx)}}{2(1+i\tan(e+fx))^2} - \frac{1}{4}c\left(\frac{1}{4}\int\frac{1}{(i\tan(e+fx)+1)\sqrt{c-ictan(e+fx)}}d\tan(e+fx) + \frac{i\sqrt{c-ictan(e+fx)}}{2c(1+i\tan(e+fx))}\right)\right)\right) + \frac{(-B+iA)(c-ictan(e+fx))^{3/2}}{6c(1+i\tan(e+fx))^3}}{a^3f}$$

↓ 73

$$\frac{c\left(\frac{1}{4}(A-3iB)\left(\frac{i\sqrt{c-ictan(e+fx)}}{2(1+i\tan(e+fx))^2} - \frac{1}{4}c\left(\frac{i\int\frac{1}{2-\frac{c-ictan(e+fx)}{c}}d\sqrt{c-ictan(e+fx)}}{2c} + \frac{i\sqrt{c-ictan(e+fx)}}{2c(1+i\tan(e+fx))}\right)\right)\right) + \frac{(-B+iA)(c-ictan(e+fx))^{3/2}}{6c(1+i\tan(e+fx))^3}}{a^3f}$$

↓ 219

$$\frac{c\left(\frac{1}{4}(A-3iB)\left(\frac{i\sqrt{c-ictan(e+fx)}}{2(1+i\tan(e+fx))^2} - \frac{1}{4}c\left(\frac{i\operatorname{arctanh}\left(\frac{\sqrt{c-ictan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{2\sqrt{2}\sqrt{c}} + \frac{i\sqrt{c-ictan(e+fx)}}{2c(1+i\tan(e+fx))}\right)\right)\right) + \frac{(-B+iA)(c-ictan(e+fx))^{3/2}}{6c(1+i\tan(e+fx))^3}}{a^3f}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2))/(a + I*a*Tan[e + f*x])^3,x]`

output `(c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(3/2))/(6*c*(1 + I*Tan[e + f*x])^3) + ((A - (3*I)*B)*(((I/2)*Sqrt[c - I*c*Tan[e + f*x]])/(1 + I*Tan[e + f*x])^2 - (c*(((I/2)*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])])/(Sqrt[2]*Sqrt[c]) + ((I/2)*Sqrt[c - I*c*Tan[e + f*x]])/(c*(1 + I*Tan[e + f*x]))))/4))/4)/(a^3*f)`

3.782.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !EqQ[e, 0] || !EqQ[c, 0] || LtQ[p, n]))`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.782.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.66

method	result
derivativedivides	$2ic^3 \left(\frac{-\frac{(-3iB+A)(c-ic \tan(fx+e))^{\frac{5}{2}}}{64c} + 8\left(\frac{iB}{96} + \frac{A}{96}\right)(c-ic \tan(fx+e))^{\frac{3}{2}} + \frac{c(-3iB+A)\sqrt{c-ic \tan(fx+e)}}{16}}{(c+ic \tan(fx+e))^3} - \frac{(-3iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{c-ic \tan(fx+e)}{c+ic \tan(fx+e)}\right)}{128c} \right) \frac{1}{fa^3}$
default	$2ic^3 \left(\frac{-\frac{(-3iB+A)(c-ic \tan(fx+e))^{\frac{5}{2}}}{64c} + 8\left(\frac{iB}{96} + \frac{A}{96}\right)(c-ic \tan(fx+e))^{\frac{3}{2}} + \frac{c(-3iB+A)\sqrt{c-ic \tan(fx+e)}}{16}}{(c+ic \tan(fx+e))^3} - \frac{(-3iB+A)\sqrt{2} \operatorname{arctanh}\left(\frac{c-ic \tan(fx+e)}{c+ic \tan(fx+e)}\right)}{128c} \right) \frac{1}{fa^3}$

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output `2*I/f/a^3*c^3*(8*(-1/512/c*(A-3*I*B)*(c-I*c*tan(f*x+e))^(5/2)+(1/96*I*B+1/96*A)*(c-I*c*tan(f*x+e))^(3/2)+1/128*c*(A-3*I*B)*(c-I*c*tan(f*x+e))^(1/2))/(c+I*c*tan(f*x+e))^3-1/128/c^(3/2)*(A-3*I*B)*2^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))`

3.782.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 395 vs. 2(164) = 328.

Time = 0.26 (sec) , antiderivative size = 395, normalized size of antiderivative = 1.87

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^3} dx = \left(3 \sqrt{\frac{1}{2}} a^3 f \sqrt{-\frac{(A^2 - 6iAB - 9B^2)c^3}{a^6 f^2}} e^{(6i fx + 6i e)} \log \left(\frac{(-iA - 3B \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^3} \right) \right)$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^3,x,algorithm="fracas")`

```
output 1/192*(3*sqrt(1/2)*a^3*f*sqrt(-(A^2 - 6*I*A*B - 9*B^2)*c^3/(a^6*f^2))*e^(6
*I*f*x + 6*I*e)*log(1/16*((-I*A - 3*B)*c^2 + sqrt(2)*sqrt(1/2)*(a^3*f*e^(2
*I*f*x + 2*I*e) + a^3*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(A^2 - 6*
I*A*B - 9*B^2)*c^3/(a^6*f^2)))*e^(-I*f*x - I*e)/(a^3*f)) - 3*sqrt(1/2)*a^3
*f*sqrt(-(A^2 - 6*I*A*B - 9*B^2)*c^3/(a^6*f^2))*e^(6*I*f*x + 6*I*e)*log(1/
16*((-I*A - 3*B)*c^2 - sqrt(2)*sqrt(1/2)*(a^3*f*e^(2*I*f*x + 2*I*e) + a^3*
f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(A^2 - 6*I*A*B - 9*B^2)*c^3/(a^
6*f^2)))*e^(-I*f*x - I*e)/(a^3*f)) - sqrt(2)*(3*(-I*A - 3*B)*c*e^(6*I*f*x
+ 6*I*e) - (17*I*A + 19*B)*c*e^(4*I*f*x + 4*I*e) + 2*(-11*I*A - B)*c*e^(2*
I*f*x + 2*I*e) + 8*(-I*A + B)*c)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))*e^(-6*
I*f*x - 6*I*e)/(a^3*f)
```

3.782.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - i \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^3} dx = i \left(\int \frac{Ac\sqrt{-i \tan(e + fx) + c}}{\tan^3(e + fx) - 3i \tan^2(e + fx) - 3 \tan(e + fx) + i} dx + \int \frac{Bc\sqrt{-i \tan(e + fx) + c}}{\tan^3(e + fx) - 3i \tan^2(e + fx) - 3 \tan(e + fx) + i} dx \right)$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(3/2)/(a+I*a*tan(f*x+e))**3
,x)
```

```
output I*(Integral(A*c*sqrt(-I*c*tan(e + f*x) + c)/(tan(e + f*x)**3 - 3*I*tan(e +
f*x)**2 - 3*tan(e + f*x) + I), x) + Integral(B*c*sqrt(-I*c*tan(e + f*x) +
c)*tan(e + f*x)/(tan(e + f*x)**3 - 3*I*tan(e + f*x)**2 - 3*tan(e + f*x) +
I), x) + Integral(-I*A*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)/(tan(e
+ f*x)**3 - 3*I*tan(e + f*x)**2 - 3*tan(e + f*x) + I), x) + Integral(-I*B*
c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**2/(tan(e + f*x)**3 - 3*I*tan(e
+ f*x)**2 - 3*tan(e + f*x) + I), x))/a**3
```

3.782.7 Maxima [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.00

$$\int \frac{(A + B \tan(e + fx))(c - i \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^3} dx = \frac{i \left(\frac{3\sqrt{2}(A-3iB)c^{5/2} \log\left(\frac{-\sqrt{2}\sqrt{c}-\sqrt{-i \tan(fx+e)+c}}{\sqrt{2}\sqrt{c}+\sqrt{-i \tan(fx+e)+c}}\right)}{a^3} + \frac{4(3(-i \tan(fx+e)+c))^{3/2}}{a^3} \right)}{a^3}$$

3.782. $\int \frac{(A+B \tan(e+fx))(c-i \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^3} dx$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`

output `1/384*I*(3*sqrt(2)*(A - 3*I*B)*c^(5/2)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a^3 + 4*(3*(-I*c*tan(f*x + e) + c)^(5/2)*(A - 3*I*B)*c^3 - 16*(-I*c*tan(f*x + e) + c)^(3/2)*(A + I*B)*c^4 - 12*sqrt(-I*c*tan(f*x + e) + c)*(A - 3*I*B)*c^5)/((-I*c*tan(f*x + e) + c)^3*a^3 - 6*(-I*c*tan(f*x + e) + c)^2*a^3*c + 12*(-I*c*tan(f*x + e) + c)*a^3*c^2 - 8*a^3*c^3)/(c*f)`

3.782.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^3} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{3/2}}{(i a \tan(fx + e) + a)^3} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(3/2)/(I*a*tan(f*x + e) + a)^3, x)`

3.782.9 Mupad [B] (verification not implemented)

Time = 9.10 (sec) , antiderivative size = 360, normalized size of antiderivative = 1.71

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^3} dx = \frac{A c^4 \sqrt{c - c \tan(e + f x) i} i + A c^3 (c - c \tan(e + f x) i)^{3/2}}{8 a^3 f} + \frac{A c^3 (c - c \tan(e + f x) i)^{3/2}}{6 a^3 f} - \frac{3 B c^4 \sqrt{c - c \tan(e + f x) i}}{8} + \frac{B c^3 (c - c \tan(e + f x) i)^{3/2}}{6} + \frac{3 B c^2 (c - c \tan(e + f x) i)^{5/2}}{32} - \frac{8 a^3 c^3 f - a^3 f (c - c \tan(e + f x) i)^3 + 6 a^3 c f (c - c \tan(e + f x) i)^2 - 12 a^3 c^2 f (c - c \tan(e + f x) i)}{64 a^3 f} + \frac{\sqrt{2} A (-c)^{3/2} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + f x) i}}{2 \sqrt{-c}}\right) i}{64 a^3 f} - \frac{3 \sqrt{2} B c^{3/2} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + f x) i}}{2 \sqrt{c}}\right)}{64 a^3 f}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*i)^(3/2))/(a + a*tan(e + f*x)*i)^3,x)`

3.782. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^3} dx$

output

$$\begin{aligned} & ((A*c^4*(c - c*\tan(e + f*x)*1i)^{(1/2)*1i)/(8*a^3*f) + (A*c^3*(c - c*\tan(e \\ & + f*x)*1i)^{(3/2)*1i)/(6*a^3*f) - (A*c^2*(c - c*\tan(e + f*x)*1i)^{(5/2)*1i)/ \\ & (32*a^3*f))/(6*c*(c - c*\tan(e + f*x)*1i)^2 - 12*c^2*(c - c*\tan(e + f*x)*1i \\ &) - (c - c*\tan(e + f*x)*1i)^3 + 8*c^3) - ((B*c^3*(c - c*\tan(e + f*x)*1i)^{(\\ & 3/2)})/6 - (3*B*c^4*(c - c*\tan(e + f*x)*1i)^{(1/2)})/8 + (3*B*c^2*(c - c*\tan(\\ & e + f*x)*1i)^{(5/2)})/32)/(8*a^3*c^3*f - a^3*f*(c - c*\tan(e + f*x)*1i)^3 + 6 \\ & *a^3*c*f*(c - c*\tan(e + f*x)*1i)^2 - 12*a^3*c^2*f*(c - c*\tan(e + f*x)*1i)) \\ & + (2^{(1/2)}*A*(-c)^{(3/2)}*atan((2^{(1/2)}*(c - c*\tan(e + f*x)*1i)^{(1/2)})/(2*(\\ & -c)^{(1/2)}))*1i)/(64*a^3*f) - (3*2^{(1/2)}*B*c^{(3/2)}*atanh((2^{(1/2)}*(c - c*ta \\ & n(e + f*x)*1i)^{(1/2)})/(2*c^{(1/2)})))/(64*a^3*f) \end{aligned}$$

3.783
$$\int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{(a+ia \tan(e+fx))^3} dx$$

3.783.1 Optimal result 7089
 3.783.2 Mathematica [A] (verified) 7089
 3.783.3 Rubi [A] (verified) 7090
 3.783.4 Maple [A] (verified) 7093
 3.783.5 Fricas [B] (verification not implemented) 7093
 3.783.6 Sympy [F] 7094
 3.783.7 Maxima [A] (verification not implemented) 7094
 3.783.8 Giac [F] 7095
 3.783.9 Mupad [B] (verification not implemented) 7095

3.783.1 Optimal result

Integrand size = 43, antiderivative size = 209

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - ic \tan(e + fx)}}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{(5iA + 7B)\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{64\sqrt{2}a^3 f} + \frac{(iA - B)\sqrt{c - ic \tan(e + fx)}}{6a^3 f(1 + i \tan(e + fx))^3}$$

$$+ \frac{(5iA + 7B)\sqrt{c - ic \tan(e + fx)}}{48a^3 f(1 + i \tan(e + fx))^2} + \frac{(5iA + 7B)\sqrt{c - ic \tan(e + fx)}}{64a^3 f(1 + i \tan(e + fx))}$$

```
output 1/128*(5*I*A+7*B)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))*c^(1/2)/a^3/f*2^(1/2)+1/6*(I*A-B)*(c-I*c*tan(f*x+e))^(1/2)/a^3/f/(1+I*tan(f*x+e))^3+1/48*(5*I*A+7*B)*(c-I*c*tan(f*x+e))^(1/2)/a^3/f/(1+I*tan(f*x+e))^2+1/64*(5*I*A+7*B)*(c-I*c*tan(f*x+e))^(1/2)/a^3/f/(1+I*tan(f*x+e))
```

3.783.2 Mathematica [A] (verified)

Time = 5.76 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.83

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - ic \tan(e + fx)}}{(a + ia \tan(e + fx))^3} dx =$$

$$\frac{\sec^3(e + fx) \left(3\sqrt{2}(5A - 7iB)\sqrt{c} \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right) (\cos(3(e + fx)) + i \sin(3(e + fx))) + 2 \cos(3(e + fx)) \right)}{384a^3 f(-i + \dots)}$$

input `Integrate[((A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])/(a + I*a*Tan[e + f*x])^3,x]`

output `-1/384*(Sec[e + f*x]^3*(3*Sqrt[2]*(5*A - (7*I)*B)*Sqrt[c]*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])]*(Cos[3*(e + f*x)] + I*Sin[3*(e + f*x)]) + 2*Cos[e + f*x]*(26*A + (2*I)*B + (41*A - (19*I)*B)*Cos[2*(e + f*x)] + 5*((5*I)*A + 7*B)*Sin[2*(e + f*x)]*Sqrt[c - I*c*Tan[e + f*x]]))/(a^3*f*(-I + Tan[e + f*x])^3)`

3.783.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 200, normalized size of antiderivative = 0.96, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.186$, Rules used = {3042, 4071, 27, 87, 52, 52, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{c - ic \tan(e + fx)}(A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{c - ic \tan(e + fx)}(A + B \tan(e + fx))}{(a + ia \tan(e + fx))^3} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A + B \tan(e + fx)}{a^4 (i \tan(e + fx) + 1)^4 \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{27} \\
 & \frac{c \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx) + 1)^4 \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{a^3 f} \\
 & \quad \downarrow \text{87} \\
 & \frac{c \left(\frac{1}{12} (5A - 7iB) \int \frac{1}{(i \tan(e + fx) + 1)^3 \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) + \frac{(-B + iA) \sqrt{c - ic \tan(e + fx)}}{6c(1 + i \tan(e + fx))^3} \right)}{a^3 f} \\
 & \quad \downarrow \text{52}
 \end{aligned}$$

3.783. $\int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{(a + ia \tan(e + fx))^3} dx$

$$\frac{c \left(\frac{1}{12} (5A - 7iB) \left(\frac{3}{8} \int \frac{1}{(i \tan(e+fx)+1)^2 \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i \sqrt{c-ic \tan(e+fx)}}{4c(1+i \tan(e+fx))^2} \right) + \frac{(-B+iA) \sqrt{c-ic \tan(e+fx)}}{6c(1+i \tan(e+fx))^3} \right)}{a^3 f}$$

↓ 52

$$\frac{c \left(\frac{1}{12} (5A - 7iB) \left(\frac{3}{8} \left(\frac{1}{4} \int \frac{1}{(i \tan(e+fx)+1) \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i \sqrt{c-ic \tan(e+fx)}}{2c(1+i \tan(e+fx))} \right) + \frac{i \sqrt{c-ic \tan(e+fx)}}{4c(1+i \tan(e+fx))^2} \right) + \frac{(-B+iA) \sqrt{c-ic \tan(e+fx)}}{6c(1+i \tan(e+fx))^3} \right)}{a^3 f}$$

↓ 73

$$\frac{c \left(\frac{1}{12} (5A - 7iB) \left(\frac{3}{8} \left(\frac{i \int \frac{1}{2 - \frac{c-ic \tan(e+fx)}{c}} d \sqrt{c-ic \tan(e+fx)}}{2c} + \frac{i \sqrt{c-ic \tan(e+fx)}}{2c(1+i \tan(e+fx))} \right) + \frac{i \sqrt{c-ic \tan(e+fx)}}{4c(1+i \tan(e+fx))^2} \right) + \frac{(-B+iA) \sqrt{c-ic \tan(e+fx)}}{6c(1+i \tan(e+fx))^3} \right)}{a^3 f}$$

↓ 219

$$\frac{c \left(\frac{1}{12} (5A - 7iB) \left(\frac{3}{8} \left(\frac{i \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2} \sqrt{c}} \right)}{2\sqrt{2} \sqrt{c}} + \frac{i \sqrt{c-ic \tan(e+fx)}}{2c(1+i \tan(e+fx))} \right) + \frac{i \sqrt{c-ic \tan(e+fx)}}{4c(1+i \tan(e+fx))^2} \right) + \frac{(-B+iA) \sqrt{c-ic \tan(e+fx)}}{6c(1+i \tan(e+fx))^3} \right)}{a^3 f}$$

input `Int[((A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])/(a + I*a*Tan[e + f*x])^3,x]`

output `(c*(((I*A - B)*Sqrt[c - I*c*Tan[e + f*x]])/(6*c*(1 + I*Tan[e + f*x])^3) + ((5*A - (7*I)*B)*(((I/4)*Sqrt[c - I*c*Tan[e + f*x]])/(c*(1 + I*Tan[e + f*x])^2) + (3*(((I/2)*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])])/(Sqrt[2]*Sqrt[c]) + ((I/2)*Sqrt[c - I*c*Tan[e + f*x]])/(c*(1 + I*Tan[e + f*x]))))/8))/12))/(a^3*f)`

3.783.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 52 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`
- rule 73 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !EqQ[e, 0] || !EqQ[c, 0] || LtQ[p, n]))`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.783.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.70

method	result
derivativedivides	$2ic^3 \left(\frac{(-7iB+5A)(c-ic \tan(fx+e))^{\frac{5}{2}}}{128c^2} - \frac{(-7iB+5A)(c-ic \tan(fx+e))^{\frac{3}{2}}}{24c} + 8 \left(\frac{11A}{256} - \frac{9iB}{256} \right) \sqrt{c-ic \tan(fx+e)} + \frac{(-7iB+5A)\sqrt{2} \arctan\left(\frac{c-ic \tan(fx+e)}{c+ic \tan(fx+e)}\right)}{(c+ic \tan(fx+e))^3} \right) \frac{1}{fa^3}$
default	$2ic^3 \left(\frac{(-7iB+5A)(c-ic \tan(fx+e))^{\frac{5}{2}}}{128c^2} - \frac{(-7iB+5A)(c-ic \tan(fx+e))^{\frac{3}{2}}}{24c} + 8 \left(\frac{11A}{256} - \frac{9iB}{256} \right) \sqrt{c-ic \tan(fx+e)} + \frac{(-7iB+5A)\sqrt{2} \arctan\left(\frac{c-ic \tan(fx+e)}{c+ic \tan(fx+e)}\right)}{(c+ic \tan(fx+e))^3} \right) \frac{1}{fa^3}$

```
input int((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
output 2*I/f/a^3*c^3*(8*(1/1024/c^2*(5*A-7*I*B)*(c-I*c*tan(f*x+e))^(5/2)-1/192/c*(5*A-7*I*B)*(c-I*c*tan(f*x+e))^(3/2)+(11/256*A-9/256*I*B)*(c-I*c*tan(f*x+e))^(1/2))/(c+I*c*tan(f*x+e))^3+1/256/c^(5/2)*(5*A-7*I*B)*2^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2)))
```

3.783.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 386 vs. 2(162) = 324.

Time = 0.26 (sec) , antiderivative size = 386, normalized size of antiderivative = 1.85

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{(a + ia \tan(e + fx))^3} dx$$

$$= \left(3 \sqrt{\frac{1}{2}} a^3 f \sqrt{-\frac{(25A^2 - 70iAB - 49B^2)c}{a^6 f^2}} e^{(6i fx + 6i e)} \log \left(\frac{\left(\sqrt{2} \sqrt{\frac{1}{2}} (a^3 f e^{(2i fx + 2i e)} + a^3 f) \sqrt{\frac{c}{e^{(2i fx + 2i e)} + 1}} \sqrt{-\frac{(25A^2 - 70iAB - 49B^2)c}{a^6 f^2}} \right)}{32 a^3 f} \right) \right)$$

```
input integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3,x,algorithm="fracas")
```

3.783. $\int \frac{(A+B \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}}{(a+ia \tan(e+fx))^3} dx$

```
output 1/384*(3*sqrt(1/2)*a^3*f*sqrt(-(25*A^2 - 70*I*A*B - 49*B^2)*c/(a^6*f^2))*e
^(6*I*f*x + 6*I*e)*log(1/32*(sqrt(2)*sqrt(1/2)*(a^3*f*e^(2*I*f*x + 2*I*e)
+ a^3*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(25*A^2 - 70*I*A*B - 49*B
^2)*c/(a^6*f^2)) + (5*I*A + 7*B)*c)*e^(-I*f*x - I*e)/(a^3*f)) - 3*sqrt(1/2
)*a^3*f*sqrt(-(25*A^2 - 70*I*A*B - 49*B^2)*c/(a^6*f^2))*e^(6*I*f*x + 6*I*e
)*log(-1/32*(sqrt(2)*sqrt(1/2)*(a^3*f*e^(2*I*f*x + 2*I*e) + a^3*f)*sqrt(c/
(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(25*A^2 - 70*I*A*B - 49*B^2)*c/(a^6*f^2))
- (5*I*A + 7*B)*c)*e^(-I*f*x - I*e)/(a^3*f)) - sqrt(2)*(3*(-11*I*A - 9*B)
*e^(6*I*f*x + 6*I*e) - (59*I*A + 25*B)*e^(4*I*f*x + 4*I*e) + 2*(-17*I*A +
5*B)*e^(2*I*f*x + 2*I*e) - 8*I*A + 8*B)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))
*e^(-6*I*f*x - 6*I*e)/(a^3*f)
```

3.783.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - i c \tan(e + fx)}}{(a + i a \tan(e + fx))^3} dx$$

$$= \frac{i \left(\int \frac{A\sqrt{-i c \tan(e + fx) + c}}{\tan^3(e + fx) - 3i \tan^2(e + fx) - 3 \tan(e + fx) + i} dx + \int \frac{B\sqrt{-i c \tan(e + fx) + c} \tan(e + fx)}{\tan^3(e + fx) - 3i \tan^2(e + fx) - 3 \tan(e + fx) + i} dx \right)}{a^3}$$

```
input integrate((c-I*c*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**3
,x)
```

```
output I*(Integral(A*sqrt(-I*c*tan(e + f*x) + c)/(tan(e + f*x)**3 - 3*I*tan(e + f
*x)**2 - 3*tan(e + f*x) + I), x) + Integral(B*sqrt(-I*c*tan(e + f*x) + c)*
tan(e + f*x)/(tan(e + f*x)**3 - 3*I*tan(e + f*x)**2 - 3*tan(e + f*x) + I),
x))/a**3
```

3.783.7 Maxima [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.05

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - i c \tan(e + fx)}}{(a + i a \tan(e + fx))^3} dx =$$

$$i \left(\frac{3\sqrt{2}(5A - 7iB)c^{\frac{3}{2}} \log\left(\frac{-\sqrt{2}\sqrt{c} - \sqrt{-i c \tan(fx+e) + c}}{\sqrt{2}\sqrt{c} + \sqrt{-i c \tan(fx+e) + c}}\right)}{a^3} + \frac{4 \left(3(-i c \tan(fx+e) + c)^{\frac{5}{2}}(5A - 7iB)c^2 - 16(-i c \tan(fx+e) + c)^{\frac{3}{2}}(5A - 7iB)c^3 \right)}{(-i c \tan(fx+e) + c)^3 a^3 - 6(-i c \tan(fx+e) + c)^2 a^3 c + 12(-i c \tan(fx+e) + c) a^3} \right)$$

768 cf

3.783. $\int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{(a+ia \tan(e+fx))^3} dx$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3,x
, algorithm="maxima")`

output `-1/768*I*(3*sqrt(2)*(5*A - 7*I*B)*c^(3/2)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/a^3
+ 4*(3*(-I*c*tan(f*x + e) + c)^(5/2)*(5*A - 7*I*B)*c^2 - 16*(-I*c*tan(f*x + e) + c)^(3/2)*(5*A - 7*I*B)*c^3 + 12*sqrt(-I*c*tan(f*x + e) + c)*(11*A - 9*I*B)*c^4)/((-I*c*tan(f*x + e) + c)^3*a^3 - 6*(-I*c*tan(f*x + e) + c)^2*a^3*c + 12*(-I*c*tan(f*x + e) + c)*a^3*c^2 - 8*a^3*c^3)/(c*f)`

3.783.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{(a + i a \tan(e + fx))^3} dx$$

$$= \int \frac{(B \tan(fx + e) + A) \sqrt{-i c \tan(fx + e) + c}}{(i a \tan(fx + e) + a)^3} dx$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3,x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*sqrt(-I*c*tan(f*x + e) + c)/(I*a*tan(f*x + e) + a)^3, x)`

3.783.9 Mupad [B] (verification not implemented)

Time = 9.05 (sec) , antiderivative size = 355, normalized size of antiderivative = 1.70

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{(a + i a \tan(e + fx))^3} dx$$

$$= \frac{\frac{7 B c (c - c \tan(e + f x) \operatorname{li})^{5/2}}{64} + \frac{9 B c^3 \sqrt{c - c \tan(e + f x) \operatorname{li}}}{16} - \frac{7 B c^2 (c - c \tan(e + f x) \operatorname{li})^{3/2}}{12}}{8 a^3 c^3 f - a^3 f (c - c \tan(e + f x) \operatorname{li})^3 + 6 a^3 c f (c - c \tan(e + f x) \operatorname{li})^2 - 12 a^3 c^2 f (c - c \tan(e + f x) \operatorname{li})}$$

$$+ \frac{\frac{A c^3 \sqrt{c - c \tan(e + f x) \operatorname{li}} \operatorname{li}}{16 a^3 f} - \frac{A c^2 (c - c \tan(e + f x) \operatorname{li})^{3/2} 5i}{12 a^3 f} + \frac{A c (c - c \tan(e + f x) \operatorname{li})^{5/2} 5i}{64 a^3 f}}{6 c (c - c \tan(e + f x) \operatorname{li})^2 - 12 c^2 (c - c \tan(e + f x) \operatorname{li}) - (c - c \tan(e + f x) \operatorname{li})^3 + 8 c^3}$$

$$+ \frac{\sqrt{2} A \sqrt{-c} \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + f x) \operatorname{li}}}{2 \sqrt{-c}}\right) 5i}{128 a^3 f} + \frac{7 \sqrt{2} B \sqrt{c} \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + f x) \operatorname{li}}}{2 \sqrt{c}}\right)}{128 a^3 f}$$

3.783. $\int \frac{(A+B \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}}{(a+ia \tan(e+fx))^3} dx$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(1/2))/(a + a*tan(e + f*x)*1i)^3,x)`

output `((7*B*c*(c - c*tan(e + f*x)*1i)^(5/2))/64 + (9*B*c^3*(c - c*tan(e + f*x)*1i)^(1/2))/16 - (7*B*c^2*(c - c*tan(e + f*x)*1i)^(3/2))/12)/(8*a^3*c^3*f - a^3*f*(c - c*tan(e + f*x)*1i)^3 + 6*a^3*c*f*(c - c*tan(e + f*x)*1i)^2 - 12*a^3*c^2*f*(c - c*tan(e + f*x)*1i)) + ((A*c^3*(c - c*tan(e + f*x)*1i)^(1/2)*11i)/(16*a^3*f) - (A*c^2*(c - c*tan(e + f*x)*1i)^(3/2)*5i)/(12*a^3*f) + (A*c*(c - c*tan(e + f*x)*1i)^(5/2)*5i)/(64*a^3*f))/(6*c*(c - c*tan(e + f*x)*1i)^2 - 12*c^2*(c - c*tan(e + f*x)*1i) - (c - c*tan(e + f*x)*1i)^3 + 8*c^3) + (2^(1/2)*A*(-c)^(1/2)*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*(-c)^(1/2)))*5i)/(128*a^3*f) + (7*2^(1/2)*B*c^(1/2)*atanh((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*c^(1/2)))/(128*a^3*f)`

3.784
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3 \sqrt{c-ic \tan(e+fx)}} dx$$

3.784.1 Optimal result 7097
 3.784.2 Mathematica [A] (verified) 7098
 3.784.3 Rubi [A] (verified) 7098
 3.784.4 Maple [A] (verified) 7101
 3.784.5 Fricas [B] (verification not implemented) 7102
 3.784.6 Sympy [F] 7102
 3.784.7 Maxima [A] (verification not implemented) 7103
 3.784.8 Giac [F] 7103
 3.784.9 Mupad [B] (verification not implemented) 7104

3.784.1 Optimal result

Integrand size = 43, antiderivative size = 245

$$\begin{aligned} & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 \sqrt{c - ic \tan(e + fx)}} dx \\ &= \frac{5(7iA + 5B) \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{128\sqrt{2}a^3\sqrt{cf}} - \frac{5(7iA + 5B)}{128a^3 f \sqrt{c - ic \tan(e + fx)}} \\ &+ \frac{iA - B}{6a^3 f (1 + i \tan(e + fx))^3 \sqrt{c - ic \tan(e + fx)}} \\ &+ \frac{7iA + 5B}{48a^3 f (1 + i \tan(e + fx))^2 \sqrt{c - ic \tan(e + fx)}} \\ &+ \frac{5(7iA + 5B)}{192a^3 f (1 + i \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}} \end{aligned}$$

```
output 5/256*(7*I*A+5*B)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a^3/f*2^(1/2)/c^(1/2)-5/128*(7*I*A+5*B)/a^3/f/(c-I*c*tan(f*x+e))^(1/2)+1/6*(I*A-B)/a^3/f/(c-I*c*tan(f*x+e))^(1/2)/(1+I*tan(f*x+e))^3+1/48*(7*I*A+5*B)/a^3/f/(c-I*c*tan(f*x+e))^(1/2)/(1+I*tan(f*x+e))^2+5/192*(7*I*A+5*B)/a^3/f/(c-I*c*tan(f*x+e))^(1/2)/(1+I*tan(f*x+e))
```


3.784.2 Mathematica [A] (verified)

Time = 5.07 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.86

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 \sqrt{c - ic \tan(e + fx)}} dx$$

$$= \frac{\sec^3(e + fx) \left(2\sqrt{c}((-125A + 7iB) \cos(e + fx) + 8(5A - 7iB) \cos(3(e + fx)) - i(7A - 5iB)(7 \sin(e + fx) + \dots)) \right)}{768a^3 \sqrt{c} f (-i - \dots)}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*Sqrt[c - I*c*Tan[e + f*x]]),x]`

output `(Sec[e + f*x]^3*(2*Sqrt[c]*((-125*A + (7*I)*B)*Cos[e + f*x] + 8*(5*A - (7*I)*B)*Cos[3*(e + f*x)] - I*(7*A - (5*I)*B)*(7*Sin[e + f*x] - 8*Sin[3*(e + f*x)])) - 15*Sqrt[2]*(7*A - (5*I)*B)*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])]*(Cos[3*(e + f*x)] + I*Sin[3*(e + f*x)])*Sqrt[c - I*c*Tan[e + f*x]])/(768*a^3*Sqrt[c]*f*(-I + Tan[e + f*x])^3*Sqrt[c - I*c*Tan[e + f*x]])`

3.784.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 227, normalized size of antiderivative = 0.93, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.209$, Rules used = {3042, 4071, 27, 87, 52, 52, 61, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 \sqrt{c - ic \tan(e + fx)}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 \sqrt{c - ic \tan(e + fx)}} dx$$

↓ 4071

$$ac \int \frac{A+B \tan(e+fx)}{a^4(i \tan(e+fx)+1)^4(c-ic \tan(e+fx))^{3/2}} d \tan(e + fx)$$

f

↓ 27

3.784. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3 \sqrt{c-ic \tan(e+fx)}} dx$

$$\begin{aligned}
& \frac{c \int \frac{A+B \tan(e+fx)}{(i \tan(e+fx)+1)^4 (c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{a^3 f} \\
& \quad \downarrow 87 \\
& \frac{c \left(\frac{1}{12} (7A - 5iB) \int \frac{1}{(i \tan(e+fx)+1)^3 (c-ic \tan(e+fx))^{3/2}} d \tan(e+fx) + \frac{-B+iA}{6c(1+i \tan(e+fx))^3 \sqrt{c-ic \tan(e+fx)}} \right)}{a^3 f} \\
& \quad \downarrow 52 \\
& \frac{c \left(\frac{1}{12} (7A - 5iB) \left(\frac{5}{8} \int \frac{1}{(i \tan(e+fx)+1)^2 (c-ic \tan(e+fx))^{3/2}} d \tan(e+fx) + \frac{i}{4c(1+i \tan(e+fx))^2 \sqrt{c-ic \tan(e+fx)}} \right) + \frac{1}{6c(1+i \tan(e+fx))} \right)}{a^3 f} \\
& \quad \downarrow 52 \\
& \frac{c \left(\frac{1}{12} (7A - 5iB) \left(\frac{5}{8} \left(\frac{3}{4} \int \frac{1}{(i \tan(e+fx)+1) (c-ic \tan(e+fx))^{3/2}} d \tan(e+fx) + \frac{i}{2c(1+i \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}} \right) + \frac{1}{4c(1+i \tan(e+fx))} \right) \right)}{a^3 f} \\
& \quad \downarrow 61 \\
& \frac{c \left(\frac{1}{12} (7A - 5iB) \left(\frac{5}{8} \left(\frac{3}{4} \left(\frac{\int \frac{1}{(i \tan(e+fx)+1) \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{2c} - \frac{i}{c \sqrt{c-ic \tan(e+fx)}} \right) + \frac{i}{2c(1+i \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}} \right) \right) \right)}{a^3 f} \\
& \quad \downarrow 73 \\
& \frac{c \left(\frac{1}{12} (7A - 5iB) \left(\frac{5}{8} \left(\frac{3}{4} \left(\frac{i \int \frac{1}{2 - \frac{c-ic \tan(e+fx)}{c}} d \sqrt{c-ic \tan(e+fx)}}{c^2} - \frac{i}{c \sqrt{c-ic \tan(e+fx)}} \right) + \frac{i}{2c(1+i \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}} \right) \right) \right)}{a^3 f} \\
& \quad \downarrow 219 \\
& \frac{c \left(\frac{1}{12} (7A - 5iB) \left(\frac{5}{8} \left(\frac{3}{4} \left(\frac{i \operatorname{arctanh} \left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2c}} \right)}{\sqrt{2c^{3/2}}} - \frac{i}{c \sqrt{c-ic \tan(e+fx)}} \right) + \frac{i}{2c(1+i \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}} \right) + \frac{1}{4c(1+i \tan(e+fx))} \right) \right)}{a^3 f}
\end{aligned}$$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*Sqrt[c - I*c*Tan[e + f*x]]),x]`

3.784. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3 \sqrt{c-ic \tan(e+fx)}} dx$

output $(c*((I*A - B)/(6*c*(1 + I*Tan[e + f*x])^3*sqrt[c - I*c*Tan[e + f*x]]) + ((7*A - (5*I)*B)*((I/4)/(c*(1 + I*Tan[e + f*x])^2*sqrt[c - I*c*Tan[e + f*x]]) + (5*((I/2)/(c*(1 + I*Tan[e + f*x])*sqrt[c - I*c*Tan[e + f*x]]) + (3*((I*ArcTanh[sqrt[c - I*c*Tan[e + f*x]]/(sqrt[2]*sqrt[c])))/(sqrt[2]*c^(3/2)) - I/(c*sqrt[c - I*c*Tan[e + f*x])))/4)/8)/12))/(a^3*f)$

3.784.3.1 Defintions of rubi rules used

rule 27 $Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] \&\& !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]$

rule 52 $Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] \&\& ILtQ[m, -1] \&\& FractionQ[n] \&\& LtQ[n, 0]$

rule 61 $Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] \&\& LtQ[m, -1] \&\& !(LtQ[n, -1] \&\& (EqQ[a, 0] || (NeQ[c, 0] \&\& LtQ[m - n, 0] \&\& IntegerQ[n]))) \&\& IntLinearQ[a, b, c, d, m, n, x]$

rule 73 $Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] \&\& LtQ[-1, m, 0] \&\& LeQ[-1, n, 0] \&\& LeQ[Denominator[n], Denominator[m]] \&\& IntLinearQ[a, b, c, d, m, n, x]$

rule 87 $Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] \&\& LtQ[p, -1] \&\& (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))$

rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.784.4 Maple [A] (verified)

Time = 0.29 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.73

method	result
derivativedivides	$2ic^3 \left(-\frac{-iB+A}{16c^3 \sqrt{c-ic \tan(fx+e)}} + \frac{8 \left(-\frac{9iB}{128} + \frac{19A}{128} \right) (c-ic \tan(fx+e))^{\frac{5}{2}} + 8 \left(\frac{7}{24} iBc - \frac{17}{24} cA \right) (c-ic \tan(fx+e))^{\frac{3}{2}} + 8 \left(-\frac{7}{32} iBc^2 + \frac{29}{32} c^2 A \right) (c+ic \tan(fx+e))^3}{16c^3} \right) \frac{1}{fa^3}$
default	$2ic^3 \left(-\frac{-iB+A}{16c^3 \sqrt{c-ic \tan(fx+e)}} + \frac{8 \left(-\frac{9iB}{128} + \frac{19A}{128} \right) (c-ic \tan(fx+e))^{\frac{5}{2}} + 8 \left(\frac{7}{24} iBc - \frac{17}{24} cA \right) (c-ic \tan(fx+e))^{\frac{3}{2}} + 8 \left(-\frac{7}{32} iBc^2 + \frac{29}{32} c^2 A \right) (c+ic \tan(fx+e))^3}{16c^3} \right) \frac{1}{fa^3}$

input `int((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

output `2*I/f/a^3*c^3*(-1/16/c^3*(A-I*B)/(c-I*c*tan(f*x+e))^(1/2)+1/16/c^3*(8*((-9/128*I*B+19/128*A)*(c-I*c*tan(f*x+e))^(5/2)+(7/24*I*B*c-17/24*c*A)*(c-I*c*tan(f*x+e))^(3/2)+(-7/32*I*B*c^2+29/32*c^2*A)*(c-I*c*tan(f*x+e))^(1/2))/(c+I*c*tan(f*x+e))^3+5/4*(-5/8*I*B+7/8*A)*2^(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))))`

$$3.784. \quad \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3 \sqrt{c-ic \tan(e+fx)}} dx$$

3.784.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 408 vs. $2(190) = 380$.

Time = 0.27 (sec) , antiderivative size = 408, normalized size of antiderivative = 1.67

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 \sqrt{c - ic \tan(e + fx)}} dx$$

$$= \left(15 \sqrt{\frac{1}{2}} a^3 c f \sqrt{-\frac{49A^2 - 70iAB - 25B^2}{a^6 c f^2}} e^{(6i fx + 6i e)} \log \left(\frac{5 \left(\sqrt{2} \sqrt{\frac{1}{2}} (a^3 f e^{(2i fx + 2i e)} + a^3 f) \sqrt{\frac{c}{e^{(2i fx + 2i e)} + 1}} \sqrt{-\frac{49A^2 - 70iAB - 25B^2}{a^6 c f^2}} \right)}{64 a^3 f} \right) \right)$$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

output `1/768*(15*sqrt(1/2)*a^3*c*f*sqrt(-(49*A^2 - 70*I*A*B - 25*B^2)/(a^6*c*f^2)))*e^(6*I*f*x + 6*I*e)*log(5/64*(sqrt(2)*sqrt(1/2)*(a^3*f*e^(2*I*f*x + 2*I*e) + a^3*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(49*A^2 - 70*I*A*B - 25*B^2)/(a^6*c*f^2)) + 7*I*A + 5*B)*e^(-I*f*x - I*e)/(a^3*f)) - 15*sqrt(1/2)*a^3*c*f*sqrt(-(49*A^2 - 70*I*A*B - 25*B^2)/(a^6*c*f^2))*e^(6*I*f*x + 6*I*e)*log(-5/64*(sqrt(2)*sqrt(1/2)*(a^3*f*e^(2*I*f*x + 2*I*e) + a^3*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(49*A^2 - 70*I*A*B - 25*B^2)/(a^6*c*f^2)) - 7*I*A - 5*B)*e^(-I*f*x - I*e)/(a^3*f)) - sqrt(2)*(48*(I*A + B)*e^(8*I*f*x + 8*I*e) + 3*(-13*I*A + 9*B)*e^(6*I*f*x + 6*I*e) - (125*I*A + 7*B)*e^(4*I*f*x + 4*I*e) + 2*(-23*I*A + 11*B)*e^(2*I*f*x + 2*I*e) - 8*I*A + 8*B)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*e^(-6*I*f*x - 6*I*e)/(a^3*c*f)`

3.784.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 \sqrt{c - ic \tan(e + fx)}} dx$$

$$= \frac{i \left(\int \frac{A}{\sqrt{-ic \tan(e + fx) + c \tan^3(e + fx) - 3i \sqrt{-ic \tan(e + fx) + c \tan^2(e + fx) - 3 \sqrt{-ic \tan(e + fx) + c \tan(e + fx) + i \sqrt{-ic \tan(e + fx) + c}}}} dx + \int \frac{B}{a^3} dx \right)}{a^3}$$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(1/2)/(a+I*a*tan(f*x+e))**3, x)`

3.784. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 \sqrt{c - ic \tan(e + fx)}} dx$

```
output I*(Integral(A/(sqrt(-I*c*tan(e + f*x) + c))*tan(e + f*x)**3 - 3*I*sqrt(-I*c
*tan(e + f*x) + c)*tan(e + f*x)**2 - 3*sqrt(-I*c*tan(e + f*x) + c)*tan(e +
f*x) + I*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(B*tan(e + f*x)/(sqrt
(-I*c*tan(e + f*x) + c))*tan(e + f*x)**3 - 3*I*sqrt(-I*c*tan(e + f*x) + c)*
tan(e + f*x)**2 - 3*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + I*sqrt(-I*c
*tan(e + f*x) + c)), x))/a**3
```

3.784.7 Maxima [A] (verification not implemented)

Time = 0.49 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.98

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 \sqrt{c - ic \tan(e + fx)}} dx =$$

$$i \left(\frac{4 \left(15 (-ic \tan(fx+e)+c)^3 (7A-5iB)c - 80 (-ic \tan(fx+e)+c)^2 (7A-5iB)c^2 + 132 (-ic \tan(fx+e)+c) (7A-5iB)c^3 - 384 (A-iB)c^4 \right)}{(-ic \tan(fx+e)+c)^{\frac{7}{2}} a^3 - 6 (-ic \tan(fx+e)+c)^{\frac{5}{2}} a^3 c + 12 (-ic \tan(fx+e)+c)^{\frac{3}{2}} a^3 c^2 - 8 \sqrt{-ic \tan(fx+e)+c} a^3 c^3} \right)$$

1536 cf

```
input integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^3,x
, algorithm="maxima")
```

```
output -1/1536*I*(4*(15*(-I*c*tan(f*x + e) + c)^3*(7*A - 5*I*B)*c - 80*(-I*c*tan(
f*x + e) + c)^2*(7*A - 5*I*B)*c^2 + 132*(-I*c*tan(f*x + e) + c)*(7*A - 5*I
*B)*c^3 - 384*(A - I*B)*c^4)/((-I*c*tan(f*x + e) + c)^(7/2)*a^3 - 6*(-I*c*
tan(f*x + e) + c)^(5/2)*a^3*c + 12*(-I*c*tan(f*x + e) + c)^(3/2)*a^3*c^2 -
8*sqrt(-I*c*tan(f*x + e) + c)*a^3*c^3) + 15*sqrt(2)*(7*A - 5*I*B)*sqrt(c)
*log(-sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + s
qrt(-I*c*tan(f*x + e) + c))/a^3)/(c*f)
```

3.784.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 \sqrt{c - ic \tan(e + fx)}} dx$$

$$= \int \frac{B \tan(fx + e) + A}{(ia \tan(fx + e) + a)^3 \sqrt{-ic \tan(fx + e) + c}} dx$$

3.784. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3 \sqrt{c-ic \tan(e+fx)}} dx$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^3,x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)^3*sqrt(-I*c*tan(f*x
+ e) + c)), x)`

3.784.9 Mupad [B] (verification not implemented)

Time = 9.20 (sec) , antiderivative size = 394, normalized size of antiderivative = 1.61

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 \sqrt{c - ic \tan(e + fx)}} dx$$

$$= \frac{B c^3 - \frac{25 B (c - c \tan(e + fx) li)^3}{128} + \frac{25 B c (c - c \tan(e + fx) li)^2}{24} - \frac{55 B c^2 (c - c \tan(e + fx) li)}{32}}{a^3 f (c - c \tan(e + fx) li)^{7/2} - 6 a^3 c f (c - c \tan(e + fx) li)^{5/2} - 8 a^3 c^3 f \sqrt{c - c \tan(e + fx) li} + 12 a^3 c^2 f \sqrt{c - c \tan(e + fx) li}}$$

$$+ \frac{\frac{A (c - c \tan(e + fx) li)^3 35i}{128 a^3 f} - \frac{A c^3 li}{a^3 f} - \frac{A c (c - c \tan(e + fx) li)^2 35i}{24 a^3 f} + \frac{A c^2 (c - c \tan(e + fx) li) 77i}{32 a^3 f}}{6 c (c - c \tan(e + fx) li)^{5/2} - (c - c \tan(e + fx) li)^{7/2} + 8 c^3 \sqrt{c - c \tan(e + fx) li} - 12 c^2 (c - c \tan(e + fx) li) \sqrt{c - c \tan(e + fx) li}}$$

$$- \frac{\sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) li}}{2 \sqrt{-c}}\right) 35i}{256 a^3 \sqrt{-c} f} + \frac{25 \sqrt{2} B \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) li}}{2 \sqrt{c}}\right)}{256 a^3 \sqrt{c} f}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^3*(c - c*tan(e + f*x)*1i
)^^(1/2)),x)`

output `(B*c^3 - (25*B*(c - c*tan(e + f*x)*1i)^3)/128 + (25*B*c*(c - c*tan(e + f*x)
)*1i)^2)/24 - (55*B*c^2*(c - c*tan(e + f*x)*1i))/32)/(a^3*f*(c - c*tan(e +
f*x)*1i)^(7/2) - 6*a^3*c*f*(c - c*tan(e + f*x)*1i)^(5/2) - 8*a^3*c^3*f*(c
- c*tan(e + f*x)*1i)^(1/2) + 12*a^3*c^2*f*(c - c*tan(e + f*x)*1i)^(3/2))
+ ((A*(c - c*tan(e + f*x)*1i)^3*35i)/(128*a^3*f) - (A*c^3*1i)/(a^3*f) - (A
c(c - c*tan(e + f*x)*1i)^2*35i)/(24*a^3*f) + (A*c^2*(c - c*tan(e + f*x)*
1i)*77i)/(32*a^3*f))/(6*c*(c - c*tan(e + f*x)*1i)^(5/2) - (c - c*tan(e + f
*x)*1i)^(7/2) + 8*c^3*(c - c*tan(e + f*x)*1i)^(1/2) - 12*c^2*(c - c*tan(e
+ f*x)*1i)^(3/2)) - (2^(1/2)*A*atan((2^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2)
)/(2*(-c)^(1/2)))*35i)/(256*a^3*(-c)^(1/2)*f) + (25*2^(1/2)*B*atanh((2^(1/
2)*(c - c*tan(e + f*x)*1i)^(1/2))/(2*c^(1/2))))/(256*a^3*c^(1/2)*f)`

3.785
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^{3/2}} dx$$

3.785.1 Optimal result 7105
 3.785.2 Mathematica [C] (verified) 7106
 3.785.3 Rubi [A] (verified) 7106
 3.785.4 Maple [A] (verified) 7109
 3.785.5 Fracas [A] (verification not implemented) 7110
 3.785.6 Sympy [F] 7111
 3.785.7 Maxima [A] (verification not implemented) 7111
 3.785.8 Giac [F] 7112
 3.785.9 Mupad [B] (verification not implemented) 7112

3.785.1 Optimal result

Integrand size = 43, antiderivative size = 274

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3(c - ic \tan(e + fx))^{3/2}} dx = \frac{35(3iA + B) \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{256\sqrt{2}a^3c^{3/2}f} - \frac{35(3iA + B)}{384a^3f(c - ic \tan(e + fx))^{3/2}} + \frac{iA - B}{6a^3f(1 + i \tan(e + fx))^3(c - ic \tan(e + fx))^{3/2}} + \frac{3iA + B}{16a^3f(1 + i \tan(e + fx))^2(c - ic \tan(e + fx))^{3/2}} + \frac{7(3iA + B)}{64a^3f(1 + i \tan(e + fx))(c - ic \tan(e + fx))^{3/2}} - \frac{35(3iA + B)}{256a^3cf\sqrt{c - ic \tan(e + fx)}}$$

```
output 35/512*(3*I*A+B)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a^3/c^(3/2)/f*2^(1/2)-35/256*(3*I*A+B)/a^3/c/f/(c-I*c*tan(f*x+e))^(1/2)-35/384*(3*I*A+B)/a^3/f/(c-I*c*tan(f*x+e))^(3/2)+1/6*(I*A-B)/a^3/f/(1+I*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^(3/2)+1/16*(3*I*A+B)/a^3/f/(1+I*tan(f*x+e))^2/(c-I*c*tan(f*x+e))^(3/2)+7/64*(3*I*A+B)/a^3/f/(1+I*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2)
```


3.785.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.14 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.70

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{3/2}} dx = \frac{\sec^4(e + fx) (105(3iA + B) \text{Hypergeometric2F1}(-\frac{1}{2}, 1, 1/2, (-1/2 * I + \tan(e + fx)) * (\cos[2*(e + fx)] + I \sin[2*(e + fx)]) + 2 * \cos[e + fx] * (3 * ((-55 * I) * A + 3 * B) * \cos[e + fx] + 8 * (I * A + 3 * B) * \cos[3*(e + fx)] + (3 * A - I * B) * (27 * \sin[e + fx] - 8 * \sin[3*(e + fx)])))))) / (768 * a^3 * c * f * (-I + \tan[e + fx])^3 * (I + \tan[e + fx]) * \text{Sqrt}[c - I * c * \tan[e + fx]])}{}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^(3/2)),x]`

output `(Sec[e + f*x]^4*(105*((3*I)*A + B)*Hypergeometric2F1[-1/2, 1, 1/2, (-1/2*I + Tan[e + f*x])*(Cos[2*(e + f*x)] + I*Sin[2*(e + f*x)]) + 2*Cos[e + f*x]*(3*((-55*I)*A + 3*B)*Cos[e + f*x] + 8*(I*A + 3*B)*Cos[3*(e + f*x)] + (3*A - I*B)*(27*Sin[e + f*x] - 8*Sin[3*(e + f*x)])))/((768*a^3*c*f*(-I + Tan[e + f*x])^3*(I + Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])`

3.785.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 261, normalized size of antiderivative = 0.95, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.233$, Rules used = {3042, 4071, 27, 87, 52, 52, 61, 61, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{4071} \\ & ac \int \frac{A + B \tan(e + fx)}{a^4 (i \tan(e + fx) + 1)^4 (c - ic \tan(e + fx))^{5/2}} d \tan(e + fx) \\ & \quad \downarrow \text{27} \\ & c \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx) + 1)^4 (c - ic \tan(e + fx))^{5/2}} d \tan(e + fx) \\ & \quad \quad \quad a^3 f \end{aligned}$$

3.785. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{3/2}} dx$

$$\begin{aligned}
 & \downarrow 87 \\
 & \frac{c\left(\frac{1}{4}(3A - iB) \int \frac{1}{(i \tan(e+fx)+1)^3(c-ic \tan(e+fx))^{5/2}} d \tan(e+fx) + \frac{-B+iA}{6c(1+i \tan(e+fx))^3(c-ic \tan(e+fx))^{3/2}}\right)}{a^3 f} \\
 & \downarrow 52 \\
 & \frac{c\left(\frac{1}{4}(3A - iB) \left(\frac{7}{8} \int \frac{1}{(i \tan(e+fx)+1)^2(c-ic \tan(e+fx))^{5/2}} d \tan(e+fx) + \frac{i}{4c(1+i \tan(e+fx))^2(c-ic \tan(e+fx))^{3/2}}\right) + \frac{i}{6c(1+i \tan(e+fx))}\right)}{a^3 f} \\
 & \downarrow 52 \\
 & \frac{c\left(\frac{1}{4}(3A - iB) \left(\frac{7}{8} \left(\frac{5}{4} \int \frac{1}{(i \tan(e+fx)+1)(c-ic \tan(e+fx))^{5/2}} d \tan(e+fx) + \frac{i}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}\right) + \frac{i}{4c(1+i \tan(e+fx))}\right)\right)}{a^3 f} \\
 & \downarrow 61 \\
 & \frac{c\left(\frac{1}{4}(3A - iB) \left(\frac{7}{8} \left(\frac{5}{4} \left(\frac{\int \frac{1}{(i \tan(e+fx)+1)(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{2c} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}}\right) + \frac{i}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))}\right)\right)\right)}{a^3 f} \\
 & \downarrow 61 \\
 & \frac{c\left(\frac{1}{4}(3A - iB) \left(\frac{7}{8} \left(\frac{5}{4} \left(\frac{\int \frac{1}{(i \tan(e+fx)+1)\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{2c} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}}\right) + \frac{i}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))}\right)\right)\right)}{a^3 f} \\
 & \downarrow 73 \\
 & \frac{c\left(\frac{1}{4}(3A - iB) \left(\frac{7}{8} \left(\frac{5}{4} \left(\frac{i \int \frac{1}{2-c-ic \tan(e+fx)} d\sqrt{c-ic \tan(e+fx)}}{c^2} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}}\right) + \frac{i}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))}\right)\right)\right)}{a^3 f} \\
 & \downarrow 219 \\
 & \frac{c\left(\frac{1}{4}(3A - iB) \left(\frac{7}{8} \left(\frac{5}{4} \left(\frac{i \operatorname{arctanh}\left(\frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{\sqrt{2}c^{3/2}} - \frac{i}{c\sqrt{c-ic \tan(e+fx)}} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}}\right) + \frac{i}{2c(1+i \tan(e+fx))(c-ic \tan(e+fx))}\right)\right)\right)}{a^3 f}
 \end{aligned}$$

3.785. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^{3/2}} dx$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^(3/2)),x]`

output `(c*((I*A - B)/(6*c*(1 + I*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^(3/2)) + ((3*A - I*B)*((I/4)/(c*(1 + I*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^(3/2)) + (7*((I/2)/(c*(1 + I*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2)) + (5*(-1/3*I)/(c*(c - I*c*Tan[e + f*x])^(3/2)) + ((I*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c]))]/(Sqrt[2]*c^(3/2)) - I/(c*Sqrt[c - I*c*Tan[e + f*x]]))/(2*c)))/4))/8))/4)/(a^3*f)`

3.785.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*(c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x]
, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.785.4 Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.75

method	result
derivativedivides	$2ic^3 \left(\frac{8 \left(-\frac{3iB}{256} + \frac{41A}{256} \right) (c - ic \tan(fx+e))^{\frac{5}{2}} + 8 \left(\frac{1}{48} iBc - \frac{35}{48} cA \right) (c - ic \tan(fx+e))^{\frac{3}{2}} + 8 \left(\frac{3}{64} iBc^2 + \frac{55}{64} c^2A \right) \sqrt{c - ic \tan(fx+e)} + \frac{35 \left(\frac{3A}{8} \right)}{16c^4} \right)}{f a^3}$
default	$2ic^3 \left(\frac{8 \left(-\frac{3iB}{256} + \frac{41A}{256} \right) (c - ic \tan(fx+e))^{\frac{5}{2}} + 8 \left(\frac{1}{48} iBc - \frac{35}{48} cA \right) (c - ic \tan(fx+e))^{\frac{3}{2}} + 8 \left(\frac{3}{64} iBc^2 + \frac{55}{64} c^2A \right) \sqrt{c - ic \tan(fx+e)} + \frac{35 \left(\frac{3A}{8} \right)}{16c^4} \right)}{f a^3}$

```
input int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^(3/2),x,method=
_RETURNVERBOSE)
```

$$3.785. \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^{3/2}} dx$$

output $2*I/f/a^3*c^3*(1/16/c^4*(8*((-3/256*I*B+41/256*A)*(c-I*c*\tan(f*x+e))^(5/2) + (1/48*I*B*c-35/48*c*A)*(c-I*c*\tan(f*x+e))^(3/2)+(3/64*I*B*c^2+55/64*c^2*A) * (c-I*c*\tan(f*x+e))^(1/2))/(c+I*c*\tan(f*x+e))^3+35/8*(3/8*A-1/8*I*B)*2^(1/2)/c^(1/2)*\operatorname{arctanh}(1/2*(c-I*c*\tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2)))-1/16/c^4*(2*A-I*B)/(c-I*c*\tan(f*x+e))^(1/2)-1/48/c^3*(A-I*B)/(c-I*c*\tan(f*x+e))^(3/2))$

3.785.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 438, normalized size of antiderivative = 1.60

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{3/2}} dx = \frac{\left(105 \sqrt{\frac{1}{2}} a^3 c^2 f \sqrt{-\frac{9A^2 - 6iAB - B^2}{a^6 c^3 f^2}} e^{(6i fx + 6ie)} \log \left(\frac{35(\sqrt{2}}{2} \right) \right)}{\dots}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^(3/2), x, algorithm="fricas")`

output $1/1536*(105*\sqrt{1/2}*a^3*c^2*f*\sqrt{-(9*A^2 - 6*I*A*B - B^2)/(a^6*c^3*f^2)})*e^{(6*I*f*x + 6*I*e)}*\log(35/128*(\sqrt{2}*\sqrt{1/2}*(a^3*c*f*e^{(2*I*f*x + 2*I*e)} + a^3*c*f)*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{-(9*A^2 - 6*I*A*B - B^2)/(a^6*c^3*f^2)}) + 3*I*A + B)*e^{(-I*f*x - I*e)/(a^3*c*f)} - 105*\sqrt{1/2}*a^3*c^2*f*\sqrt{-(9*A^2 - 6*I*A*B - B^2)/(a^6*c^3*f^2)})*e^{(6*I*f*x + 6*I*e)}*\log(-35/128*(\sqrt{2}*\sqrt{1/2}*(a^3*c*f*e^{(2*I*f*x + 2*I*e)} + a^3*c*f)*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{-(9*A^2 - 6*I*A*B - B^2)/(a^6*c^3*f^2)}) - 3*I*A - B)*e^{(-I*f*x - I*e)/(a^3*c*f)} - \sqrt{2}*(16*(I*A + B)*e^{(10*I*f*x + 10*I*e)} + 32*(7*I*A + 4*B)*e^{(8*I*f*x + 8*I*e)} - (-43*I*A - 121*B)*e^{(6*I*f*x + 6*I*e)} + 5*(-43*I*A + 7*B)*e^{(4*I*f*x + 4*I*e)} + 2*(-29*I*A + 17*B)*e^{(2*I*f*x + 2*I*e)} - 8*I*A + 8*B)*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)})*e^{(-6*I*f*x - 6*I*e)/(a^3*c^2*f)}$

3.785.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{3/2}} dx = i \left(\int \frac{A}{-ic \sqrt{-ic \tan(e + fx) + c} \tan^4(e + fx) - 2c \sqrt{-ic \tan(e + fx) + c} \tan^3(e + fx) + c^2 \tan^2(e + fx) - c^2} dx \right)$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**3/(c-I*c*tan(f*x+e))**(3/2),x)`

output `I*(Integral(A/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4 - 2*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3 - 2*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + I*c*sqrt(-I*c*tan(e + f*x) + c)), x) + Integral(B*tan(e + f*x)/(-I*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4 - 2*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3 - 2*c*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + I*c*sqrt(-I*c*tan(e + f*x) + c)), x))/a**3`

3.785.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.96

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{3/2}} dx = \frac{i \left(\frac{4 \left(105 (-ic \tan(fx+e)+c)^4 (3A-iB) - 560 (-ic \tan(fx+e)+c)^3 (3A-iB)c + 924 (-ic \tan(fx+e)+c)^2 (3A-iB)c^2 - 384 (-ic \tan(fx+e)+c) \right)}{(-ic \tan(fx+e)+c)^{\frac{9}{2}} a^3 - 6 (-ic \tan(fx+e)+c)^{\frac{7}{2}} a^3 c + 12 (-ic \tan(fx+e)+c)^{\frac{5}{2}} a^3 c^2 - 8 (-ic \tan(fx+e)+c)^{\frac{3}{2}} a^3 c^3} \right)}{3072 cf}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `-1/3072*I*(4*(105*(-I*c*tan(f*x + e) + c)^4*(3*A - I*B) - 560*(-I*c*tan(f*x + e) + c)^3*(3*A - I*B)*c + 924*(-I*c*tan(f*x + e) + c)^2*(3*A - I*B)*c^2 - 384*(-I*c*tan(f*x + e) + c)*(3*A - I*B)*c^3 - 256*(A - I*B)*c^4)/((-I*c*tan(f*x + e) + c)^(9/2)*a^3 - 6*(-I*c*tan(f*x + e) + c)^(7/2)*a^3*c + 12*(-I*c*tan(f*x + e) + c)^(5/2)*a^3*c^2 - 8*(-I*c*tan(f*x + e) + c)^(3/2)*a^3*c^3) + 105*sqrt(2)*(3*A - I*B)*log(-sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c))/(a^3*sqrt(c))/(c*f)`

3.785. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3 (c-ic \tan(e+fx))^{3/2}} dx$

3.785.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{3/2}} dx = \int \frac{B \tan(fx + e) + A}{(ia \tan(fx + e) + a)^3 (-ic \tan(fx + e) + c)^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^(3/2),x
, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)^3*(-I*c*tan(f*x + e)
) + c)^(3/2)), x)`

3.785.9 Mupad [B] (verification not implemented)

Time = 9.49 (sec) , antiderivative size = 443, normalized size of antiderivative = 1.62

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{3/2}} dx =$$

$$-\frac{\frac{A(c - c \tan(e + fx) \operatorname{li})^3 35i}{16 a^3 f} + \frac{A c^3 \operatorname{li}}{3 a^3 f} - \frac{A(c - c \tan(e + fx) \operatorname{li})^4 105i}{256 a^3 c f} - \frac{A c(c - c \tan(e + fx) \operatorname{li})^2 231i}{64 a^3 f} + \frac{A c^2 (c - c \tan(e + fx) \operatorname{li})}{2 a^3 f}}{6 c(c - c \tan(e + fx) \operatorname{li})^{7/2} - (c - c \tan(e + fx) \operatorname{li})^{9/2} + 8 c^3 (c - c \tan(e + fx) \operatorname{li})^{3/2} - 12 c^2 (c - c \tan(e + fx) \operatorname{li})^{1/2}}$$

$$+ \frac{\frac{B c^3}{3} + \frac{35 B (c - c \tan(e + fx) \operatorname{li})^3}{48} - \frac{77 B c (c - c \tan(e + fx) \operatorname{li})^2}{64} + \frac{B c^2 (c - c \tan(e + fx) \operatorname{li})}{2} - \frac{35 B (c - c \tan(e + fx) \operatorname{li})}{2}}{a^3 f (c - c \tan(e + fx) \operatorname{li})^{9/2} - 6 a^3 c f (c - c \tan(e + fx) \operatorname{li})^{7/2} - 8 a^3 c^3 f (c - c \tan(e + fx) \operatorname{li})^{3/2} + 12 a^3 c^2 f (c - c \tan(e + fx) \operatorname{li})^{1/2}}$$

$$+ \frac{\sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) \operatorname{li}}}{2 \sqrt{-c}}\right) 105i}{512 a^3 (-c)^{3/2} f} + \frac{35 \sqrt{2} B \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) \operatorname{li}}}{2 \sqrt{c}}\right)}{512 a^3 c^{3/2} f}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*I)^3*(c - c*tan(e + f*x)*I)
)^(3/2)),x)`

output $((B*c^3)/3 + (35*B*(c - c*\tan(e + f*x)*1i)^3)/48 - (77*B*c*(c - c*\tan(e + f*x)*1i)^2)/64 + (B*c^2*(c - c*\tan(e + f*x)*1i))/2 - (35*B*(c - c*\tan(e + f*x)*1i)^4)/(256*c))/(a^3*f*(c - c*\tan(e + f*x)*1i)^{(9/2)} - 6*a^3*c*f*(c - c*\tan(e + f*x)*1i)^{(7/2)} - 8*a^3*c^3*f*(c - c*\tan(e + f*x)*1i)^{(3/2)} + 12*a^3*c^2*f*(c - c*\tan(e + f*x)*1i)^{(5/2)}) - ((A*(c - c*\tan(e + f*x)*1i)^3*35i)/(16*a^3*f) + (A*c^3*1i)/(3*a^3*f) - (A*(c - c*\tan(e + f*x)*1i)^4*105i)/(256*a^3*c*f) - (A*c*(c - c*\tan(e + f*x)*1i)^2*231i)/(64*a^3*f) + (A*c^2*(c - c*\tan(e + f*x)*1i)*3i)/(2*a^3*f))/(6*c*(c - c*\tan(e + f*x)*1i)^{(7/2)} - (c - c*\tan(e + f*x)*1i)^{(9/2)} + 8*c^3*(c - c*\tan(e + f*x)*1i)^{(3/2)} - 12*c^2*(c - c*\tan(e + f*x)*1i)^{(5/2)}) + (2^{(1/2)}*A*atan((2^{(1/2)}*(c - c*\tan(e + f*x)*1i)^{(1/2)})/(2*(-c)^{(1/2)})))*105i)/(512*a^3*(-c)^{(3/2)}*f) + (35*2^{(1/2)}*B*atanh((2^{(1/2)}*(c - c*\tan(e + f*x)*1i)^{(1/2)})/(2*c^{(1/2)})))/(512*a^3*c^{(3/2)}*f)$

3.785. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^{3/2}} dx$

3.786
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ictan(e+fx))^{5/2}} dx$$

3.786.1 Optimal result 7114
 3.786.2 Mathematica [C] (verified) 7115
 3.786.3 Rubi [A] (verified) 7115
 3.786.4 Maple [A] (verified) 7119
 3.786.5 Fricas [A] (verification not implemented) 7119
 3.786.6 Sympy [F] 7120
 3.786.7 Maxima [A] (verification not implemented) 7121
 3.786.8 Giac [F] 7121
 3.786.9 Mupad [B] (verification not implemented) 7122

3.786.1 Optimal result

Integrand size = 43, antiderivative size = 311

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3(c - ictan(e + fx))^{5/2}} dx = \frac{21(11iA + B) \operatorname{arctanh}\left(\frac{\sqrt{c - ictan(e + fx)}}{\sqrt{2}\sqrt{c}}\right)}{512\sqrt{2}a^3c^{5/2}f} - \frac{21(11iA + B)}{640a^3f(c - ictan(e + fx))^{5/2}} + \frac{iA - B}{6a^3f(1 + i \tan(e + fx))^3(c - ictan(e + fx))^{5/2}} + \frac{11iA + B}{48a^3f(1 + i \tan(e + fx))^2(c - ictan(e + fx))^{5/2}} + \frac{3(11iA + B)}{64a^3f(1 + i \tan(e + fx))(c - ictan(e + fx))^{5/2}} - \frac{7(11iA + B)}{256a^3cf(c - ictan(e + fx))^{3/2}} - \frac{21(11iA + B)}{512a^3c^2f\sqrt{c - ictan(e + fx)}}$$

```
output 21/1024*(11*I*A+B)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2))/a
^3/c^(5/2)/f*2^(1/2)-21/512*(11*I*A+B)/a^3/c^2/f/(c-I*c*tan(f*x+e))^(1/2)-
21/640*(11*I*A+B)/a^3/f/(c-I*c*tan(f*x+e))^(5/2)+1/6*(I*A-B)/a^3/f/(1+I*ta
n(f*x+e))^3/(c-I*c*tan(f*x+e))^(5/2)+1/48*(11*I*A+B)/a^3/f/(1+I*tan(f*x+e)
)^2/(c-I*c*tan(f*x+e))^(5/2)+3/64*(11*I*A+B)/a^3/f/(1+I*tan(f*x+e))/(c-I*c
*tan(f*x+e))^(5/2)-7/256*(11*I*A+B)/a^3/c/f/(c-I*c*tan(f*x+e))^(3/2)
```

3.786.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.41 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.60

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{5/2}} dx = \frac{\sec^3(e + fx) (30(71A + 11iB) \cos(e + fx) - 16(A - 1$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^(5/2)),x]`

output `(Sec[e + f*x]^3*(30*(71*A + (11*I)*B)*Cos[e + f*x] - 16*(A - (11*I)*B)*Cos[3*(e + f*x)] - 105*(11*A - I*B)*Hypergeometric2F1[-3/2, 1, -1/2, (-1/2*I)*(I + Tan[e + f*x])] * Sec[e + f*x] * (Cos[2*(e + f*x)] + I*Sin[2*(e + f*x)]) + 2*((11*I)*A + B)*(55*Sin[e + f*x] - 8*Sin[3*(e + f*x)])))/(3840*a^3*c^2*f*(-I + Tan[e + f*x])^3*(I + Tan[e + f*x])^2*Sqrt[c - I*c*Tan[e + f*x]])`

3.786.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 295, normalized size of antiderivative = 0.95, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.256$, Rules used = {3042, 4071, 27, 87, 52, 52, 61, 61, 61, 73, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{5/2}} dx \\ & \quad \downarrow \text{4071} \\ & ac \int \frac{A + B \tan(e + fx)}{a^4 (i \tan(e + fx) + 1)^4 (c - ic \tan(e + fx))^{7/2}} d \tan(e + fx) \\ & \quad \downarrow \text{27} \\ & c \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx) + 1)^4 (c - ic \tan(e + fx))^{7/2}} d \tan(e + fx) \\ & \quad \downarrow \\ & \frac{c \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx) + 1)^4 (c - ic \tan(e + fx))^{7/2}} d \tan(e + fx)}{a^3 f} \end{aligned}$$

3.786. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{5/2}} dx$

↓ 87

$$\frac{c \left(\frac{1}{12} (11A - iB) \int \frac{1}{(i \tan(e+fx)+1)^3 (c-ic \tan(e+fx))^{7/2}} d \tan(e+fx) + \frac{-B+iA}{6c(1+i \tan(e+fx))^3 (c-ic \tan(e+fx))^{5/2}} \right)}{a^3 f}$$

↓ 52

$$\frac{c \left(\frac{1}{12} (11A - iB) \left(\frac{9}{8} \int \frac{1}{(i \tan(e+fx)+1)^2 (c-ic \tan(e+fx))^{7/2}} d \tan(e+fx) + \frac{i}{4c(1+i \tan(e+fx))^2 (c-ic \tan(e+fx))^{5/2}} \right) \right) + \frac{1}{6c(1+i \tan(e+fx))}}{a^3 f}$$

↓ 52

$$\frac{c \left(\frac{1}{12} (11A - iB) \left(\frac{9}{8} \left(\frac{7}{4} \int \frac{1}{(i \tan(e+fx)+1) (c-ic \tan(e+fx))^{7/2}} d \tan(e+fx) + \frac{i}{2c(1+i \tan(e+fx)) (c-ic \tan(e+fx))^{5/2}} \right) \right) \right) + \frac{1}{4c(1+i \tan(e+fx))}}{a^3 f}$$

↓ 61

$$\frac{c \left(\frac{1}{12} (11A - iB) \left(\frac{9}{8} \left(\frac{7}{4} \left(\frac{\int \frac{1}{(i \tan(e+fx)+1) (c-ic \tan(e+fx))^{5/2}} d \tan(e+fx)}{2c} - \frac{i}{5c(c-ic \tan(e+fx))^{5/2}} \right) \right) \right) \right) + \frac{i}{2c(1+i \tan(e+fx)) (c-ic \tan(e+fx))}}{a^3 f}$$

↓ 61

$$\frac{c \left(\frac{1}{12} (11A - iB) \left(\frac{9}{8} \left(\frac{7}{4} \left(\frac{\frac{\int \frac{1}{(i \tan(e+fx)+1) (c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{2c} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}}}{2c} - \frac{i}{5c(c-ic \tan(e+fx))^{5/2}} \right) \right) \right) \right) + \frac{1}{2c(1+i \tan(e+fx))}}{a^3 f}$$

↓ 61

$$\frac{c \left(\frac{1}{12} (11A - iB) \left(\frac{9}{8} \left(\frac{7}{4} \left(\frac{\frac{\frac{\int \frac{1}{(i \tan(e+fx)+1) \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{2c} - \frac{i}{c \sqrt{c-ic \tan(e+fx)}}}{2c} - \frac{i}{3c(c-ic \tan(e+fx))^{3/2}}}{2c} - \frac{i}{5c(c-ic \tan(e+fx))^{5/2}} \right) \right) \right) \right) + \frac{1}{2c(1+i \tan(e+fx))}}{a^3 f}$$

↓ 73

3.786. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3 (c-ic \tan(e+fx))^{5/2}} dx$

$$c \left(\frac{1}{12}(11A - iB) \left(\frac{9}{8} \left(\frac{7}{4} \left(\frac{i \int \frac{1}{2 - c - ic \tan(e+fx)} d\sqrt{c - ic \tan(e+fx)}}{c^2} - \frac{i}{c\sqrt{c - ic \tan(e+fx)}} - \frac{i}{3c(c - ic \tan(e+fx))^{3/2}} - \frac{i}{5c(c - ic \tan(e+fx))^{5/2}} \right) \right) \right) \right) a^3$$

↓ 219

$$c \left(\frac{1}{12}(11A - iB) \left(\frac{9}{8} \left(\frac{7}{4} \left(\frac{i \operatorname{arctanh}\left(\frac{\sqrt{c - ic \tan(e+fx)}}{\sqrt{2}\sqrt{c}}\right)}{\sqrt{2}c^{3/2}} - \frac{i}{c\sqrt{c - ic \tan(e+fx)}} - \frac{i}{3c(c - ic \tan(e+fx))^{3/2}} - \frac{i}{5c(c - ic \tan(e+fx))^{5/2}} \right) \right) \right) \right) + \frac{1}{2c} a^3 f$$

```
input Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^(5/2)),x]
```

```
output (c*((I*A - B)/(6*c*(1 + I*Tan[e + f*x])^3*(c - I*c*Tan[e + f*x])^(5/2)) + ((11*A - I*B)*((I/4)/(c*(1 + I*Tan[e + f*x])^2*(c - I*c*Tan[e + f*x])^(5/2))) + (9*((I/2)/(c*(1 + I*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2)) + (7*((-1/5*I)/(c*(c - I*c*Tan[e + f*x])^(5/2)) + ((-1/3*I)/(c*(c - I*c*Tan[e + f*x])^(3/2)) + ((I*ArcTanh[Sqrt[c - I*c*Tan[e + f*x]]/(Sqrt[2]*Sqrt[c])])/(Sqrt[2]*c^(3/2)) - I/(c*Sqrt[c - I*c*Tan[e + f*x]]))/(2*c))/(2*c)))/4))/8))/12))/(a^3*f)
```

3.786.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 52 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]
```

3.786. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^3(c-ic \tan(e+fx))^{5/2}} dx$

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.786.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 232, normalized size of antiderivative = 0.75

method	result
derivativedivides	$2ic^3 \left(-\frac{-iB+5A}{32c^5 \sqrt{c-ic \tan(fx+e)}} - \frac{-iB+2A}{48c^4 (c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{-iB+A}{80c^3 (c-ic \tan(fx+e))^{\frac{5}{2}}} + \frac{8 \left(\frac{11iB}{256} + \frac{71A}{256} \right) (c-ic \tan(fx+e))^{\frac{5}{2}} + 8 \left(-\frac{59}{48} \right)}{f a} \right)$
default	$2ic^3 \left(-\frac{-iB+5A}{32c^5 \sqrt{c-ic \tan(fx+e)}} - \frac{-iB+2A}{48c^4 (c-ic \tan(fx+e))^{\frac{3}{2}}} - \frac{-iB+A}{80c^3 (c-ic \tan(fx+e))^{\frac{5}{2}}} + \frac{8 \left(\frac{11iB}{256} + \frac{71A}{256} \right) (c-ic \tan(fx+e))^{\frac{5}{2}} + 8 \left(-\frac{59}{48} \right)}{f a} \right)$

```
input int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)
```

```
output 2*I/f/a^3*c^3*(-1/32/c^5*(5*A-I*B)/(c-I*c*tan(f*x+e))^(1/2)-1/48/c^4*(2*A-I*B)/(c-I*c*tan(f*x+e))^(3/2)-1/80/c^3*(A-I*B)/(c-I*c*tan(f*x+e))^(5/2)+1/32/c^5*(8*((11/256*I*B+71/256*A)*(c-I*c*tan(f*x+e))^(5/2)+(-59/48*c*A-11/48*I*B*c)*(c-I*c*tan(f*x+e))^(3/2)+(21/64*I*B*c^2+89/64*c^2*A)*(c-I*c*tan(f*x+e))^(1/2)))/(c+I*c*tan(f*x+e))^3+21/8*(-1/8*I*B+11/8*A)*2^(1/2)/c^(1/2)*arctanh(1/2*(c-I*c*tan(f*x+e))^(1/2)*2^(1/2)/c^(1/2)))
```

3.786.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 464, normalized size of antiderivative = 1.49

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{5/2}} dx = \left(315 \sqrt{\frac{1}{2}} a^3 c^3 f \sqrt{-\frac{121 A^2 - 22i AB - B^2}{a^6 c^5 f^2}} e^{(6i fx + 6i e)} \log \left(\frac{21}{\dots} \right) \right)$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^(5/2),x,algorithm="fricas")
```

```
output 1/15360*(315*sqrt(1/2)*a^3*c^3*f*sqrt(-(121*A^2 - 22*I*A*B - B^2)/(a^6*c^5
*f^2))*e^(6*I*f*x + 6*I*e)*log(21/256*(sqrt(2)*sqrt(1/2)*(a^3*c^2*f*e^(2*I
*f*x + 2*I*e) + a^3*c^2*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(121*A^
2 - 22*I*A*B - B^2)/(a^6*c^5*f^2)) + 11*I*A + B)*e^(-I*f*x - I*e)/(a^3*c^2
*f)) - 315*sqrt(1/2)*a^3*c^3*f*sqrt(-(121*A^2 - 22*I*A*B - B^2)/(a^6*c^5*f
^2))*e^(6*I*f*x + 6*I*e)*log(-21/256*(sqrt(2)*sqrt(1/2)*(a^3*c^2*f*e^(2*I
*f*x + 2*I*e) + a^3*c^2*f)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-(121*A^2
- 22*I*A*B - B^2)/(a^6*c^5*f^2)) - 11*I*A - B)*e^(-I*f*x - I*e)/(a^3*c^2*
f)) - sqrt(2)*(48*(I*A + B)*e^(12*I*f*x + 12*I*e) + 16*(29*I*A + 19*B)*e^(
10*I*f*x + 10*I*e) + 16*(199*I*A + 59*B)*e^(8*I*f*x + 8*I*e) - (-1433*I*A
- 1003*B)*e^(6*I*f*x + 6*I*e) + 5*(-329*I*A + 101*B)*e^(4*I*f*x + 4*I*e) +
10*(-35*I*A + 23*B)*e^(2*I*f*x + 2*I*e) - 40*I*A + 40*B)*sqrt(c/(e^(2*I*f
*x + 2*I*e) + 1)))e^(-6*I*f*x - 6*I*e)/(a^3*c^3*f)
```

3.786.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{5/2}} dx = i \left(\int \frac{-c^2 \sqrt{-ic \tan(e + fx) + c \tan^5(e + fx) + ic^2 \sqrt{-ic \tan(e + fx) + c \tan^4(e + fx)}}}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{5/2}} dx \right)$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**3/(c-I*c*tan(f*x+e))**
(5/2),x)
```

```
output I*(Integral(A/(-c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**5 + I*c**2*
sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**4 - 2*c**2*sqrt(-I*c*tan(e + f*x)
+ c)*tan(e + f*x)**3 + 2*I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)
**2 - c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x) + I*c**2*sqrt(-I*c*tan
(e + f*x) + c)), x) + Integral(B*tan(e + f*x)/(-c**2*sqrt(-I*c*tan(e + f*x)
+ c)*tan(e + f*x)**5 + I*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**
4 - 2*c**2*sqrt(-I*c*tan(e + f*x) + c)*tan(e + f*x)**3 + 2*I*c**2*sqrt(-I*
c*tan(e + f*x) + c)*tan(e + f*x)**2 - c**2*sqrt(-I*c*tan(e + f*x) + c)*tan
(e + f*x) + I*c**2*sqrt(-I*c*tan(e + f*x) + c)), x))/a**3
```

3.786.7 Maxima [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 291, normalized size of antiderivative = 0.94

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{5/2}} dx =$$

$$i \left(\frac{4 \left(315 (-ic \tan(fx+e)+c)^5 (11A-iB) - 1680 (-ic \tan(fx+e)+c)^4 (11A-iB)c + 2772 (-ic \tan(fx+e)+c)^3 (11A-iB)c^2 - 1152 (-ic \tan(fx+e)+c)^2 a^3 c^3 - 256 (-ic \tan(fx+e)+c) a^3 c^2 + 12 (-ic \tan(fx+e)+c) a^3 c - 6 (-ic \tan(fx+e)+c)^{11/2} a^3 c - 6 (-ic \tan(fx+e)+c)^{9/2} a^3 c^2 + 12 (-ic \tan(fx+e)+c)^{7/2} a^3 c^3 - 6 (-ic \tan(fx+e)+c)^{5/2} a^3 c^2 + 12 (-ic \tan(fx+e)+c)^{3/2} a^3 c - 6 (-ic \tan(fx+e)+c)^{1/2} a^3 c \right)}{(-ic \tan(fx+e)+c)^{11/2} a^3 c - 6 (-ic \tan(fx+e)+c)^{9/2} a^3 c^2 + 12 (-ic \tan(fx+e)+c)^{7/2} a^3 c^3 - 6 (-ic \tan(fx+e)+c)^{5/2} a^3 c^2 + 12 (-ic \tan(fx+e)+c)^{3/2} a^3 c - 6 (-ic \tan(fx+e)+c)^{1/2} a^3 c}$$

30720

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^(5/2),x
, algorithm="maxima")
```

```
output -1/30720*I*(4*(315*(-I*c*tan(f*x + e) + c)^5*(11*A - I*B) - 1680*(-I*c*tan
(f*x + e) + c)^4*(11*A - I*B)*c + 2772*(-I*c*tan(f*x + e) + c)^3*(11*A - I
*B)*c^2 - 1152*(-I*c*tan(f*x + e) + c)^2*(11*A - I*B)*c^3 - 256*(-I*c*tan(
f*x + e) + c)*(11*A - I*B)*c^4 - 1536*(A - I*B)*c^5)/((-I*c*tan(f*x + e) +
c)^(11/2)*a^3*c - 6*(-I*c*tan(f*x + e) + c)^(9/2)*a^3*c^2 + 12*(-I*c*tan(
f*x + e) + c)^(7/2)*a^3*c^3 - 8*(-I*c*tan(f*x + e) + c)^(5/2)*a^3*c^4) + 3
15*sqrt(2)*(11*A - I*B)*log(-(sqrt(2)*sqrt(c) - sqrt(-I*c*tan(f*x + e) + c
)))/(sqrt(2)*sqrt(c) + sqrt(-I*c*tan(f*x + e) + c)))/(a^3*c^(3/2)))/(c*f)
```

3.786.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - ic \tan(e + fx))^{5/2}} dx = \int \frac{B \tan(fx + e) + A}{(ia \tan(fx + e) + a)^3 (-ic \tan(fx + e) + c)^{5/2}} dx$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^3/(c-I*c*tan(f*x+e))^(5/2),x
, algorithm="giac")
```

```
output integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)^3*(-I*c*tan(f*x + e)
) + c)^(5/2)), x)
```


3.786.9 Mupad [B] (verification not implemented)

Time = 9.99 (sec) , antiderivative size = 490, normalized size of antiderivative = 1.58

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^3 (c - i c \tan(e + fx))^{5/2}} dx =$$

$$\frac{-\frac{A(c - c \tan(e + fx) li)^3 254 li}{640 a^3 f} + \frac{A c^3 li}{5 a^3 f} + \frac{A(c - c \tan(e + fx) li)^4 77 i}{32 a^3 c f} - \frac{A(c - c \tan(e + fx) li)^5 231 i}{512 a^3 c^2 f} + \frac{A c(c - c \tan(e + fx) li)^2 33 i}{20 a^3 f} + \frac{A c^2(c - c \tan(e + fx) li)^4}{32 c} + \frac{B c^3}{5} - \frac{231 B(c - c \tan(e + fx) li)^3}{640} + \frac{3 B c(c - c \tan(e + fx) li)^2}{20} + \frac{B c^2(c - c \tan(e + fx) li)}{30} + \frac{7 B(c - c \tan(e + fx) li)^4}{32 c}}{6 c(c - c \tan(e + fx) li)^{9/2} - (c - c \tan(e + fx) li)^{11/2} + 8 c^3(c - c \tan(e + fx) li)^{5/2} - 12 c^2(c - c \tan(e + fx) li)^{7/2} - 8 a^3 c^3 f(c - c \tan(e + fx) li)^{5/2} + a^3 f(c - c \tan(e + fx) li)^{11/2} - 6 a^3 c f(c - c \tan(e + fx) li)^{9/2} - 8 a^3 c^3 f(c - c \tan(e + fx) li)^{5/2} + \sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) li}}{2 \sqrt{-c}}\right) 231 li - \frac{21 \sqrt{2} B \operatorname{atanh}\left(\frac{\sqrt{2} \sqrt{c - c \tan(e + fx) li}}{2 \sqrt{c}}\right)}{1024 a^3 c^{5/2} f}}$$

```
input int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*li)^3*(c - c*tan(e + f*x)*li)^(5/2)),x)
```

```
output ((B*c^3)/5 - (231*B*(c - c*tan(e + f*x)*li)^3)/640 + (3*B*c*(c - c*tan(e + f*x)*li)^2)/20 + (B*c^2*(c - c*tan(e + f*x)*li))/30 + (7*B*(c - c*tan(e + f*x)*li)^4)/(32*c) - (21*B*(c - c*tan(e + f*x)*li)^5)/(512*c^2))/(a^3*f*(c - c*tan(e + f*x)*li)^(11/2) - 6*a^3*c*f*(c - c*tan(e + f*x)*li)^(9/2) - 8*a^3*c^3*f*(c - c*tan(e + f*x)*li)^(5/2) + 12*a^3*c^2*f*(c - c*tan(e + f*x)*li)^(7/2)) - ((A*c^3*li)/(5*a^3*f) - (A*(c - c*tan(e + f*x)*li)^3*254 li)/(640*a^3*f) + (A*(c - c*tan(e + f*x)*li)^4*77 i)/(32*a^3*c*f) - (A*(c - c*tan(e + f*x)*li)^5*231 i)/(512*a^3*c^2*f) + (A*c*(c - c*tan(e + f*x)*li)^2*33 i)/(20*a^3*f) + (A*c^2*(c - c*tan(e + f*x)*li)*li)/(30*a^3*f))/(6*c*(c - c*tan(e + f*x)*li)^(9/2) - (c - c*tan(e + f*x)*li)^(11/2) + 8*c^3*(c - c*tan(e + f*x)*li)^(5/2) - 12*c^2*(c - c*tan(e + f*x)*li)^(7/2)) - (2^(1/2)*A*atan((2^(1/2)*(c - c*tan(e + f*x)*li)^(1/2))/(2*(-c)^(1/2)))*231 i)/(1024*a^3*(-c)^(5/2)*f) + (21*2^(1/2)*B*atanh((2^(1/2)*(c - c*tan(e + f*x)*li)^(1/2))/(2*c^(1/2))))/(1024*a^3*c^(5/2)*f)
```

3.787 $\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx$

3.787.1 Optimal result	7123
3.787.2 Mathematica [A] (verified)	7124
3.787.3 Rubi [A] (verified)	7124
3.787.4 Maple [A] (verified)	7127
3.787.5 Fricas [B] (verification not implemented)	7128
3.787.6 Sympy [F(-1)]	7129
3.787.7 Maxima [B] (verification not implemented)	7129
3.787.8 Giac [F(-1)]	7130
3.787.9 Mupad [F(-1)]	7131

3.787.1 Optimal result

Integrand size = 45, antiderivative size = 272

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx =$$

$$\frac{5\sqrt{a}(4iA - 3B)c^{7/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{4f}$$

$$- \frac{5(4iA - 3B)c^3 \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}}{8f}$$

$$- \frac{5(4iA - 3B)c^2 \sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}}{24f}$$

$$- \frac{(4iA - 3B)c \sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{5/2}}{12f}$$

$$+ \frac{B \sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{7/2}}{4f}$$

output

```
-5/4*(4*I*A-3*B)*c^(7/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))*a^(1/2)/f-5/8*(4*I*A-3*B)*c^3*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)/f-5/24*(4*I*A-3*B)*c^2*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(3/2)/f-1/12*(4*I*A-3*B)*c*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(5/2)/f+1/4*B*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(7/2)/f
```

3.787.2 Mathematica [A] (verified)

Time = 6.62 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.63

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx = \frac{30\sqrt{a}(-4iA + 3B)c^{7/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right) + c^3 \sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}}{2}$$

input `Integrate[Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2),x]`

output `(30*Sqrt[a]*((-4*I)*A + 3*B)*c^(7/2)*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])] + c^3*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]]*((-88*I)*A + 72*B - 9*(4*A + (5*I)*B)*Tan[e + f*x] + (8*I)*(A + (3*I)*B)*Tan[e + f*x]^2 + (6*I)*B*Tan[e + f*x]^3)/(24*f)`

3.787.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.97, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.178$, Rules used = {3042, 4071, 90, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{a + ia \tan(e + fx)}(c - ictan(e + fx))^{7/2}(A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{a + ia \tan(e + fx)}(c - ictan(e + fx))^{7/2}(A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(A+B \tan(e+fx))(c-ictan(e+fx))^{5/2}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{90} \end{aligned}$$

3.787. $\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx$

$$\begin{aligned}
 & \frac{ac\left(\frac{1}{4}(4A + 3iB) \int \frac{(c-ic \tan(e+fx))^{5/2}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e + fx) + \frac{B\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{7/2}}{4ac}\right)}{f} \\
 & \quad \downarrow 60 \\
 & \frac{ac\left(\frac{1}{4}(4A + 3iB) \left(\frac{5}{3}c \int \frac{(c-ic \tan(e+fx))^{3/2}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e + fx) - \frac{i\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{5/2}}{3a}\right) + \frac{B\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{7/2}}{4ac}\right)}{f} \\
 & \quad \downarrow 60 \\
 & \frac{ac\left(\frac{1}{4}(4A + 3iB) \left(\frac{5}{3}c \left(\frac{3}{2}c \int \frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e + fx) - \frac{i\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{3/2}}{2a}\right) - \frac{i\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{5/2}}{3a}\right)\right)}{f} \\
 & \quad \downarrow 60 \\
 & \frac{ac\left(\frac{1}{4}(4A + 3iB) \left(\frac{5}{3}c \left(\frac{3}{2}c \left(c \int \frac{1}{\sqrt{i \tan(e+fx)a+a}\sqrt{c-ic \tan(e+fx)}} d \tan(e + fx) - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a}\right) - \frac{i\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{3/2}}{2a}\right)\right)\right)}{f} \\
 & \quad \downarrow 45 \\
 & \frac{ac\left(\frac{1}{4}(4A + 3iB) \left(\frac{5}{3}c \left(\frac{3}{2}c \left(2c \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a}\right) - \frac{i\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{3/2}}{2a}\right)\right)\right)}{f} \\
 & \quad \downarrow 218 \\
 & \frac{ac\left(\frac{1}{4}(4A + 3iB) \left(\frac{5}{3}c \left(\frac{3}{2}c \left(-\frac{2i\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right) - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a}\right) - \frac{i\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{3/2}}{2a}\right)\right)\right)}{f}
 \end{aligned}$$

input `Int[Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2),x]`

output `(a*c*((B*Sqrt[a + I*a*Tan[e + f*x]]*(c - I*c*Tan[e + f*x])^(7/2))/(4*a*c) + ((4*A + (3*I)*B)*(((-1/3*I)*Sqrt[a + I*a*Tan[e + f*x]]*(c - I*c*Tan[e + f*x])^(5/2))/a + (5*c*(((-1/2*I)*Sqrt[a + I*a*Tan[e + f*x]]*(c - I*c*Tan[e + f*x])^(3/2))/a + (3*c*(((-2*I)*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[a] - (I*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/a))/2))/3))/4))/f`

3.787.3.1 Defintions of rubi rules used

- rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 60 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 90 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.787.4 Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 349, normalized size of antiderivative = 1.28

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} c^3\left(6 i B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)} \tan (f x+e)^3+8 i A \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}\right)}{6 f \sqrt{a c\left(1+\tan (f x+e)^2\right)}}$
default	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} c^3\left(6 i B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)} \tan (f x+e)^3+8 i A \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}\right)}{6 f \sqrt{a c\left(1+\tan (f x+e)^2\right)}}$
parts	$\frac{A \sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} c^3\left(2 i \tan (f x+e)^2 \sqrt{a c\left(1+\tan (f x+e)^2\right)} \sqrt{a c}-22 i \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}\right)}{6 f \sqrt{a c\left(1+\tan (f x+e)^2\right)}}$

input `int((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x,method=_RETURNVERBOSE)`

output `1/24/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*c^3*(6*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^3+8*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+45*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c-45*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-24*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2-88*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)+60*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c-36*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+72*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2)/(a*c*(1+tan(f*x+e)^2))^(1/2)`

$$3.787. \quad \int \sqrt{a+ia \tan (e+f x)}(A+B \tan (e+f x))(c-i c \tan (e+f x))^{7 / 2} d x$$

3.787.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 604 vs. $2(206) = 412$.

Time = 0.27 (sec) , antiderivative size = 604, normalized size of antiderivative = 2.22

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx = \frac{15 \sqrt{\frac{(16A^2 + 24iAB - 9B^2)ac^7}{f^2}} (fe^{(6ifx+6ie)} + 3fe^{(4ifx+4ie)} + 3fe^{(2ifx+2ie)} + f) \log \left(\frac{4 \left(2 \left((4iA - 3B) \right) \right)}{\dots} \right)}{\dots}$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="fricas")`

output `1/48*(15*sqrt((16*A^2 + 24*I*A*B - 9*B^2)*a*c^7/f^2)*(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)*log(4*(2*((4*I*A - 3*B)*c^3*e^(3*I*f*x + 3*I*e) + (4*I*A - 3*B)*c^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + sqrt((16*A^2 + 24*I*A*B - 9*B^2)*a*c^7/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((4*I*A - 3*B)*c^3*e^(2*I*f*x + 2*I*e) + (4*I*A - 3*B)*c^3) - 15*sqrt((16*A^2 + 24*I*A*B - 9*B^2)*a*c^7/f^2)*(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)*log(4*(2*((4*I*A - 3*B)*c^3*e^(3*I*f*x + 3*I*e) + (4*I*A - 3*B)*c^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - sqrt((16*A^2 + 24*I*A*B - 9*B^2)*a*c^7/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((4*I*A - 3*B)*c^3*e^(2*I*f*x + 2*I*e) + (4*I*A - 3*B)*c^3) - 4*(15*(4*I*A - 3*B)*c^3*e^(7*I*f*x + 7*I*e) + 55*(4*I*A - 3*B)*c^3*e^(5*I*f*x + 5*I*e) + 73*(4*I*A - 3*B)*c^3*e^(3*I*f*x + 3*I*e) + 3*(44*I*A - 49*B)*c^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)`

3.787.6 Sympy [F(-1)]

Timed out.

$$\int \sqrt{a + ia \tan(e + fx)} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(7/2),x)`

output `Timed out`

3.787.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1343 vs. $2(206) = 412$.

Time = 1.29 (sec) , antiderivative size = 1343, normalized size of antiderivative = 4.94

$$\int \sqrt{a + ia \tan(e + fx)} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \text{Too large to display}$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="maxima")`

output

```
-96*(60*(4*A + 3*I*B)*c^3*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 220*(4*A + 3*I*B)*c^3*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 292*(4*A + 3*I*B)*c^3*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 12*(44*A + 49*I*B)*c^3*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 60*(4*I*A - 3*B)*c^3*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 220*(4*I*A - 3*B)*c^3*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 292*(4*I*A - 3*B)*c^3*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 12*(44*I*A - 49*B)*c^3*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 30*((4*A + 3*I*B)*c^3*cos(8*f*x + 8*e) + 4*(4*A + 3*I*B)*c^3*cos(6*f*x + 6*e) + 6*(4*A + 3*I*B)*c^3*cos(4*f*x + 4*e) + 4*(4*A + 3*I*B)*c^3*cos(2*f*x + 2*e) + (4*I*A - 3*B)*c^3*sin(8*f*x + 8*e) + 4*(4*I*A - 3*B)*c^3*sin(6*f*x + 6*e) + 6*(4*I*A - 3*B)*c^3*sin(4*f*x + 4*e) + 4*(4*I*A - 3*B)*c^3*sin(2*f*x + 2*e) + (4*A + 3*I*B)*c^3*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) + 30*((4*A + 3*I*B)*c^3*cos(8*f*x + 8*e) + 4*(4*A + 3*I*B)*c^3*cos(6*f*x + 6*e) + 6*(4*A + 3*I*B)*c^3*cos(4*f*x + 4*e) + 4*(4*A + 3*I*B)*c^3*cos(2*f*x + 2*e) + (4*I*A - 3*B)*c^3*sin(8*f*x + 8*e) + 4*(4*I*A - 3*B)*c^3*sin(6*f*x + 6*e) + 6*(4*I*A - 3*B)*c^3*sin(4*f*x + 4*e) + 4*(4*I*A - 3*B)*c^3*sin(2*f*x + 2*e) + (4*A + 3*I*B)*c^3)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), -sin(1/2*a...
```

3.787.8 Giac [F(-1)]

Timed out.

$$\int \sqrt{a + ia \tan(e + fx)} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \text{Timed out}$$

input

```
integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="giac")
```

output

```
Timed out
```

3.787.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + ia \tan(e + fx)} (A + B \tan(e + fx)) (c - ictan(e + fx))^{7/2} dx = \int (A + B \tan(e + fx)) \sqrt{a + a \tan(e + fx)} \operatorname{li}(c - c \tan(e + fx) \operatorname{li})^{7/2} dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(1/2)*(c - c*tan(e + f*x)*1i)^(7/2),x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(1/2)*(c - c*tan(e + f*x)*1i)^(7/2), x)`

3.788 $\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2} dx$

3.788.1 Optimal result	7132
3.788.2 Mathematica [A] (verified)	7133
3.788.3 Rubi [A] (verified)	7133
3.788.4 Maple [A] (verified)	7136
3.788.5 Fricas [B] (verification not implemented)	7136
3.788.6 Sympy [F]	7137
3.788.7 Maxima [B] (verification not implemented)	7138
3.788.8 Giac [F]	7139
3.788.9 Mupad [F(-1)]	7139

3.788.1 Optimal result

Integrand size = 45, antiderivative size = 217

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2} dx =$$

$$\frac{\sqrt{a}(3iA - 2B)c^{5/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{f}$$

$$- \frac{(3iA - 2B)c^2 \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}}{2f}$$

$$- \frac{(3iA - 2B)c \sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}}{6f}$$

$$+ \frac{B \sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{5/2}}{3f}$$

output

```
-(3*I*A-2*B)*c^(5/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))*a^(1/2)/f-1/2*(3*I*A-2*B)*c^2*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)/f-1/6*(3*I*A-2*B)*c*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(3/2)/f+1/3*B*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(5/2)/f
```

3.788.2 Mathematica [A] (verified)

Time = 5.31 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.69

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ict \tan(e + fx))^{5/2} dx = \frac{6\sqrt{a}(-3iA + 2B)c^{5/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ict \tan(e+fx)}}\right) - c^2 \sqrt{a + ia \tan(e + fx)} \sqrt{c - ict \tan(e + fx)}}{6f}$$

input `Integrate[Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2), x]`

output `(6*Sqrt[a]*((-3*I)*A + 2*B)*c^(5/2)*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])] - c^2*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]]*((12*I)*A - 10*B + 3*(A + (2*I)*B)*Tan[e + f*x] + 2*B*Tan[e + f*x]^2))/(6*f)`

3.788.3 Rubi [A] (verified)Time = 0.42 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 90, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{a + ia \tan(e + fx)}(c - ict \tan(e + fx))^{5/2}(A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{a + ia \tan(e + fx)}(c - ict \tan(e + fx))^{5/2}(A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(A+B \tan(e+fx))(c-ict \tan(e+fx))^{3/2}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{90} \\ & \frac{ac\left(\frac{1}{3}(3A + 2iB) \int \frac{(c-ict \tan(e+fx))^{3/2}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e + fx) + \frac{B\sqrt{a+ia \tan(e+fx)}(c-ict \tan(e+fx))^{5/2}}{3ac}\right)}{f} \end{aligned}$$

3.788. $\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ict \tan(e + fx))^{5/2} dx$

↓ 60

$$ac \left(\frac{1}{3}(3A + 2iB) \left(\frac{3}{2}c \int \frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e + fx) - \frac{i\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{3/2}}{2a} \right) + \frac{B\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{3/2}}{3ac} \right) / f$$

↓ 60

$$ac \left(\frac{1}{3}(3A + 2iB) \left(\frac{3}{2}c \left(c \int \frac{1}{\sqrt{i \tan(e+fx)a+a} \sqrt{c-ic \tan(e+fx)}} d \tan(e + fx) - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a} \right) \right) - \frac{i\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{3/2}}{3ac} \right) / f$$

↓ 45

$$ac \left(\frac{1}{3}(3A + 2iB) \left(\frac{3}{2}c \left(2c \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a} \right) \right) - \frac{i\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{3/2}}{3ac} \right) / f$$

↓ 218

$$ac \left(\frac{1}{3}(3A + 2iB) \left(\frac{3}{2}c \left(-\frac{2i\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{\sqrt{a}} - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a} \right) \right) - \frac{i\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{3/2}}{3ac} \right) / f$$

input `Int[Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2),x]`

output `(a*c*((B*Sqrt[a + I*a*Tan[e + f*x]]*(c - I*c*Tan[e + f*x])^(5/2))/(3*a*c) + ((3*A + (2*I)*B)*(((-1/2*I)*Sqrt[a + I*a*Tan[e + f*x]]*(c - I*c*Tan[e + f*x])^(3/2))/a + (3*c*((-2*I)*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[a] - (I*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/a))/2))/3)/f`

3.788.3.1 Defintions of rubi rules used

- rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 60 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(
b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]`
- rule 90 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
), x] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p
+ 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n,
p}, x] && NeQ[n + p + 2, 0]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`
- rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]`

3.788.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.31

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} c^2 \left(-6iB \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right) ac+6iB\sqrt{ac} \sqrt{ac}}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} c^2 \left(-6iB \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right) ac+6iB\sqrt{ac} \sqrt{ac}}{\dots}$
parts	$\frac{A\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} c^2 \left(4i\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} + \tan(fx+e)\sqrt{ac(1+\tan(fx+e)^2)} \right) \sqrt{ac}}{2f\sqrt{ac(1+\tan(fx+e)^2)} \sqrt{ac}}$

input `int((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

output `-1/6/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*c^2*(-6*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)))^(1/2))/(a*c)^(1/2))*a*c+6*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+2*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+12*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)-9*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)))^(1/2))/(a*c)^(1/2))*a*c+3*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-10*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2)/(a*c*(1+tan(f*x+e)^2))^(1/2)`

3.788.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 547 vs. 2(164) = 328.

3.788. $\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2} dx$

Time = 0.27 (sec) , antiderivative size = 547, normalized size of antiderivative = 2.52

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx = \frac{3 \sqrt{\frac{(9A^2 + 12iAB - 4B^2)ac^5}{f^2}} (fe^{(4i fx + 4i e)} + 2 fe^{(2i fx + 2i e)} + f) \log \left(\frac{4 \left(2((3iA - 2B)c^2 e^{(3i fx + 3i e)} + (3iA - 2B)c^2 e^{(i fx + i e)} + (-3iA + 2B)c^2) \right)}{\dots} \right)}{\dots}$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="fricas")`

output `1/12*(3*sqrt((9*A^2 + 12*I*A*B - 4*B^2)*a*c^5/f^2)*(f*e^(4*I*f*x + 4*I*e) + 2*f*e^(2*I*f*x + 2*I*e) + f)*log(-4*(2*((3*I*A - 2*B)*c^2*e^(3*I*f*x + 3*I*e) + (3*I*A - 2*B)*c^2*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1)))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + sqrt((9*A^2 + 12*I*A*B - 4*B^2)*a*c^5/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((-3*I*A + 2*B)*c^2*e^(2*I*f*x + 2*I*e) + (-3*I*A + 2*B)*c^2) - 3*sqrt((9*A^2 + 12*I*A*B - 4*B^2)*a*c^5/f^2)*(f*e^(4*I*f*x + 4*I*e) + 2*f*e^(2*I*f*x + 2*I*e) + f)*log(-4*(2*((3*I*A - 2*B)*c^2*e^(3*I*f*x + 3*I*e) + (3*I*A - 2*B)*c^2*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1)))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - sqrt((9*A^2 + 12*I*A*B - 4*B^2)*a*c^5/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((-3*I*A + 2*B)*c^2*e^(2*I*f*x + 2*I*e) + (-3*I*A + 2*B)*c^2) - 4*(3*(3*I*A - 2*B)*c^2*e^(5*I*f*x + 5*I*e) + 8*(3*I*A - 2*B)*c^2*e^(3*I*f*x + 3*I*e) + 3*(5*I*A - 6*B)*c^2*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(f*e^(4*I*f*x + 4*I*e) + 2*f*e^(2*I*f*x + 2*I*e) + f)`

3.788.6 Sympy [F]

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx = \int \sqrt{ia(\tan(e + fx) - i)}(-ic(\tan(e + fx) + i))^{5/2} (A + B \tan(e + fx)) dx$$

input `integrate((a+I*a*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(5/2),x)`

3.788. $\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx$

output `Integral(sqrt(I*a*(tan(e + f*x) - I))*(-I*c*(tan(e + f*x) + I))**(5/2)*(A + B*tan(e + f*x)), x)`

3.788.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1085 vs. $2(164) = 328$.

Time = 0.85 (sec) , antiderivative size = 1085, normalized size of antiderivative = 5.00

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2} dx = \text{Too large to display}$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `-6*(12*(3*A + 2*I*B)*c^2*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 32*(3*A + 2*I*B)*c^2*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 12*(5*A + 6*I*B)*c^2*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 12*(3*I*A - 2*B)*c^2*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 32*(3*I*A - 2*B)*c^2*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 12*(5*I*A - 6*B)*c^2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 6*((3*A + 2*I*B)*c^2*cos(6*f*x + 6*e) + 3*(3*A + 2*I*B)*c^2*cos(4*f*x + 4*e) + 3*(3*A + 2*I*B)*c^2*cos(2*f*x + 2*e) + (3*I*A - 2*B)*c^2*sin(6*f*x + 6*e) + 3*(3*I*A - 2*B)*c^2*sin(4*f*x + 4*e) + 3*(3*I*A - 2*B)*c^2*sin(2*f*x + 2*e) + (3*A + 2*I*B)*c^2)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) + 6*((3*A + 2*I*B)*c^2*cos(6*f*x + 6*e) + 3*(3*A + 2*I*B)*c^2*cos(4*f*x + 4*e) + 3*(3*A + 2*I*B)*c^2*cos(2*f*x + 2*e) + (3*I*A - 2*B)*c^2*sin(6*f*x + 6*e) + 3*(3*I*A - 2*B)*c^2*sin(4*f*x + 4*e) + 3*(3*I*A - 2*B)*c^2*sin(2*f*x + 2*e) + (3*A + 2*I*B)*c^2)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), -sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) + 3*((3*I*A - 2*B)*c^2*cos(6*f*x + 6*e) + 3*(3*I*A - 2*B)*c^2*cos(4*f*x + 4*e) + 3*(3*I*A - 2*B)*c^2*cos(2*f*x + 2*e) - (3*A + 2*I*B)*c^2*sin(6*f*x + 6*e) - 3*(3*A + 2*I*B)*c^2*sin(4*f*x + 4*e) - 3*(3*A + 2*I*B)*c^2*sin(2*f*x + 2*e) + (3*I*A - 2*B)*c^2)*log(cos(1/2*arct...`

3.788.8 Giac [F]

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx = \int (B \tan(fx + e) + A) \sqrt{ia \tan(fx + e) + a} (-ictan(fx + e) + c)^{5/2} dx$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `sage0*x`

3.788.9 Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx = \int (A + B \tan(e + fx)) \sqrt{a + a \tan(e + fx)} \operatorname{li}(c - c \tan(e + fx) \operatorname{li})^{5/2} dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(1/2)*(c - c*tan(e + f*x)*1i)^(5/2),x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(1/2)*(c - c*tan(e + f*x)*1i)^(5/2), x)`

3.789 $\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx$

3.789.1 Optimal result 7140
 3.789.2 Mathematica [A] (verified) 7141
 3.789.3 Rubi [A] (verified) 7141
 3.789.4 Maple [A] (verified) 7143
 3.789.5 Fricas [B] (verification not implemented) 7144
 3.789.6 Sympy [F] 7145
 3.789.7 Maxima [B] (verification not implemented) 7145
 3.789.8 Giac [B] (verification not implemented) 7146
 3.789.9 Mupad [F(-1)] 7147

3.789.1 Optimal result

Integrand size = 45, antiderivative size = 164

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx =$$

$$\frac{\sqrt{a}(2iA - B)c^{3/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{f}$$

$$- \frac{(2iA - B)c\sqrt{a + ia \tan(e + fx)}\sqrt{c - ic \tan(e + fx)}}{2f}$$

$$+ \frac{B\sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}}{2f}$$

output

```
-(2*I*A-B)*c^(3/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*
tan(f*x+e))^(1/2))*a^(1/2)/f-1/2*(2*I*A-B)*c*(a+I*a*tan(f*x+e))^(1/2)*(c-I
*c*tan(f*x+e))^(1/2)/f+1/2*B*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(
3/2)/f
```

3.789.2 Mathematica [A] (verified)

Time = 3.71 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.79

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ict \tan(e + fx))^{3/2} dx = \frac{2\sqrt{a}(-2iA + B)c^{3/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ict \tan(e+fx)}}\right) + c\sqrt{a + ia \tan(e + fx)}(-2iA + 2B - iB \tan(e + fx))}{2f}$$

input `Integrate[Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2), x]`

output `(2*Sqrt[a]*((-2*I)*A + B)*c^(3/2)*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])] + c*Sqrt[a + I*a*Tan[e + f*x]]*((-2*I)*A + 2*B - I*B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]/(2*f)`

3.789.3 Rubi [A] (verified)Time = 0.40 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.02, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 90, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{a + ia \tan(e + fx)}(c - ict \tan(e + fx))^{3/2}(A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int \sqrt{a + ia \tan(e + fx)}(c - ict \tan(e + fx))^{3/2}(A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(A+B \tan(e+fx))\sqrt{c-ict \tan(e+fx)}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{90} \\ & \frac{ac \left(\frac{1}{2}(2A + iB) \int \frac{\sqrt{c-ict \tan(e+fx)}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e + fx) + \frac{B\sqrt{a+ia \tan(e+fx)}(c-ict \tan(e+fx))^{3/2}}{2ac} \right)}{f} \end{aligned}$$

3.789. $\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ict \tan(e + fx))^{3/2} dx$

$$\begin{aligned}
 & \downarrow 60 \\
 & \frac{ac \left(\frac{1}{2}(2A + iB) \left(c \int \frac{1}{\sqrt{i \tan(e+fx)a+a} \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) - \frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{a} \right) + \frac{B \sqrt{a+ia \tan(e+fx)}}{2ac} \right)}{f} \\
 & \downarrow 45 \\
 & \frac{ac \left(\frac{1}{2}(2A + iB) \left(2c \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} - \frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{a} \right) + \frac{B \sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))}{2ac} \right)}{f} \\
 & \downarrow 218 \\
 & \frac{ac \left(\frac{1}{2}(2A + iB) \left(-\frac{2i\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{\sqrt{a}} - \frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{a} \right) + \frac{B \sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))}{2ac} \right)}{f}
 \end{aligned}$$

input `Int[Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(a*c*((B*Sqrt[a + I*a*Tan[e + f*x]]*(c - I*c*Tan[e + f*x])^(3/2))/(2*a*c) + ((2*A + I*B)*(((-2*I)*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[a] - (I*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/a))/2)/f`

3.789.3.1 Defintions of rubi rules used

rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

3.789. $\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx$

```
rule 90 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p
+ 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n,
p}, x] && NeQ[n + p + 2, 0]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.789.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.36

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} c\left(i B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)} \tan (f x+e)-i B \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c}}{\sqrt{a}}\right)}{2 f}$
default	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} c\left(i B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)} \tan (f x+e)-i B \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c}}{\sqrt{a}}\right)}{2 f}$
parts	$\frac{A\left(-i \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}+a c \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}}\right)\right) \sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)}}{f \sqrt{a c\left(1+\tan (f x+e)^2\right)} \sqrt{a c}}$

```
input int((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x,m
ethod=_RETURNVERBOSE)
```

$$3.789. \int \sqrt{a+i a \tan (e+f x)}(A+B \tan (e+f x))(c-i c \tan (e+f x))^{3 / 2} d x$$

output
$$-1/2/f*(a*(1+I*\tan(f*x+e)))^{(1/2)}*(-c*(I*\tan(f*x+e)-1))^{(1/2)}*c*(I*B*(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)}*\tan(f*x+e)-I*B*\ln((a*c*\tan(f*x+e)+(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)))/(a*c)^{(1/2)})*a*c+2*I*A*(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)}-2*A*\ln((a*c*\tan(f*x+e)+(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)))/(a*c)^{(1/2)})*a*c-2*B*(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)))/(a*c*(1+\tan(f*x+e)^2))^{(1/2)}/(a*c)^{(1/2)}$$

3.789.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 459 vs. $2(119) = 238$.

Time = 0.27 (sec) , antiderivative size = 459, normalized size of antiderivative = 2.80

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ictan(e + fx))^{3/2} dx = \frac{\sqrt{\frac{(4A^2 + 4iAB - B^2)ac^3}{f^2}}(fe^{(2i fx + 2ie)} + f) \log \left(\frac{4 \left(2((2iA - B)ce^{(3i fx + 3ie)} + (2iA - B)ce^{(i fx + ie)}) \sqrt{\frac{a}{e^{(2i fx + 2ie)}}} \right)}{(-2iA + B)ce^{(2i fx + 2ie)}} \right)}{(-2iA + B)ce^{(2i fx + 2ie)}}$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x, algorithm="fracas")`

output
$$1/4*(\text{sqrt}((4*A^2 + 4*I*A*B - B^2)*a*c^3/f^2)*(f*e^{(2*I*f*x + 2*I*e)} + f)*\log(-4*(2*((2*I*A - B)*c*e^{(3*I*f*x + 3*I*e)} + (2*I*A - B)*c*e^{(I*f*x + I*e)}))\text{sqrt}(a/(e^{(2*I*f*x + 2*I*e)} + 1))\text{sqrt}(c/(e^{(2*I*f*x + 2*I*e)} + 1)) + \text{sqrt}((4*A^2 + 4*I*A*B - B^2)*a*c^3/f^2)*(f*e^{(2*I*f*x + 2*I*e)} - f))/((-2*I*A + B)*c*e^{(2*I*f*x + 2*I*e)} + (-2*I*A + B)*c) - \text{sqrt}((4*A^2 + 4*I*A*B - B^2)*a*c^3/f^2)*(f*e^{(2*I*f*x + 2*I*e)} + f)*\log(-4*(2*((2*I*A - B)*c*e^{(3*I*f*x + 3*I*e)} + (2*I*A - B)*c*e^{(I*f*x + I*e)}))\text{sqrt}(a/(e^{(2*I*f*x + 2*I*e)} + 1))\text{sqrt}(c/(e^{(2*I*f*x + 2*I*e)} + 1)) - \text{sqrt}((4*A^2 + 4*I*A*B - B^2)*a*c^3/f^2)*(f*e^{(2*I*f*x + 2*I*e)} - f))/((-2*I*A + B)*c*e^{(2*I*f*x + 2*I*e)} + (-2*I*A + B)*c) - 4*((2*I*A - B)*c*e^{(3*I*f*x + 3*I*e)} + (2*I*A - 3*B)*c*e^{(I*f*x + I*e)})\text{sqrt}(a/(e^{(2*I*f*x + 2*I*e)} + 1))\text{sqrt}(c/(e^{(2*I*f*x + 2*I*e)} + 1)))/(f*e^{(2*I*f*x + 2*I*e)} + f)$$

3.789.6 Sympy [F]

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx = \int \sqrt{ia(\tan(e + fx) - i)}(-ic(\tan(e + fx) + i))^{3/2} (A + B \tan(e + fx)) dx$$

input `integrate((a+I*a*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(3/2),x)`

output `Integral(sqrt(I*a*(tan(e + f*x) - I))*(-I*c*(tan(e + f*x) + I))**(3/2)*(A + B*tan(e + f*x)), x)`

3.789.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 771 vs. $2(119) = 238$.

Time = 0.63 (sec) , antiderivative size = 771, normalized size of antiderivative = 4.70

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx = \text{Too large to display}$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output

```
-4*(4*(2*A + I*B)*c*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) +
  4*(2*A + 3*I*B)*c*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) +
  4*(2*I*A - B)*c*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 4*(
  2*I*A - 3*B)*c*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 2*((
  2*A + I*B)*c*cos(4*f*x + 4*e) + 2*(2*A + I*B)*c*cos(2*f*x + 2*e) + (2*I*A
  - B)*c*sin(4*f*x + 4*e) + 2*(2*I*A - B)*c*sin(2*f*x + 2*e) + (2*A + I*B)*c
  )*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*ar
  ctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) + 2*((2*A + I*B)*c*cos(4*f
  *x + 4*e) + 2*(2*A + I*B)*c*cos(2*f*x + 2*e) + (2*I*A - B)*c*sin(4*f*x + 4
  *e) + 2*(2*I*A - B)*c*sin(2*f*x + 2*e) + (2*A + I*B)*c)*arctan2(cos(1/2*ar
  ctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), -sin(1/2*arctan2(sin(2*f*x + 2
  *e), cos(2*f*x + 2*e))) + 1) + ((2*I*A - B)*c*cos(4*f*x + 4*e) + 2*(2*I*A
  - B)*c*cos(2*f*x + 2*e) - (2*A + I*B)*c*sin(4*f*x + 4*e) - 2*(2*A + I*B)*c
  *sin(2*f*x + 2*e) + (2*I*A - B)*c)*log(cos(1/2*arctan2(sin(2*f*x + 2*e), c
  os(2*f*x + 2*e)))^2 + sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))
  ^2 + 2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) + ((-2*I*
  A + B)*c*cos(4*f*x + 4*e) + 2*(-2*I*A + B)*c*cos(2*f*x + 2*e) + (2*A + I*B
  )*c*sin(4*f*x + 4*e) + 2*(2*A + I*B)*c*sin(2*f*x + 2*e) + (-2*I*A + B)*c)*
  log(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + sin(1/2*arcta
  n2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 - 2*sin(1/2*arctan2(sin(2*f*x...
```

3.789.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1747 vs. $2(119) = 238$.

Time = 14.17 (sec) , antiderivative size = 1747, normalized size of antiderivative = 10.65

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2} dx = \text{Too large to display}$$

input

```
integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/
2),x, algorithm="giac")
```

```
output 1/4*(4*(a^4*c^4 + 3*a^4*c^3 + 3*a^3*c^4 + 3*a^4*c^2 + 3*a^3*c^3 + 3*a^2*c^4 + a^4*c + a^3*c^2 + a^2*c^3 + a*c^4 + a^3*c - 2*a^2*c^2 + a*c^3)*sqrt(-a*c)*A*e^(9*I*f*x + 9*I*e) - 7*(a^4*c^4 + 3*a^4*c^3 + 3*a^3*c^4 + 3*a^4*c^2 + 3*a^3*c^3 + 3*a^2*c^4 + a^4*c + a^3*c^2 + a^2*c^3 + a*c^4 + a^3*c - 2*a^2*c^2 + a*c^3)*sqrt(a*c)*B*e^(9*I*f*x + 9*I*e) - 8*(a^4*c^4 + 3*a^4*c^3 + 3*a^3*c^4 + 3*a^4*c^2 + 3*a^3*c^3 + 3*a^2*c^4 + a^4*c + a^3*c^2 + a^2*c^3 + a*c^4 + a^3*c - 2*a^2*c^2 + a*c^3)*sqrt(-a*c)*A*e^(7*I*f*x + 7*I*e) - 2*6*(a^4*c^4 + 3*a^4*c^3 + 3*a^3*c^4 + 3*a^4*c^2 + 3*a^3*c^3 + 3*a^2*c^4 + a^4*c + a^3*c^2 + a^2*c^3 + a*c^4 + a^3*c - 2*a^2*c^2 + a*c^3)*sqrt(a*c)*B*e^(7*I*f*x + 7*I*e) - 16*(a^4*c^4 + 3*a^4*c^3 + 3*a^3*c^4 + 3*a^4*c^2 + 3*a^3*c^3 + 3*a^2*c^4 + a^4*c + a^3*c^2 + a^2*c^3 + a*c^4 + a^3*c - 2*a^2*c^2 + a*c^3)*sqrt(-a*c)*A*e^(5*I*f*x + 5*I*e) + 12*(a^4*c^4 + 3*a^4*c^3 + 3*a^3*c^4 + 3*a^4*c^2 + 3*a^3*c^3 + 3*a^2*c^4 + a^4*c + a^3*c^2 + a^2*c^3 + a*c^4 + a^3*c - 2*a^2*c^2 + a*c^3)*sqrt(a*c)*B*e^(5*I*f*x + 5*I*e) - 8*(a^4*c^4 + 3*a^4*c^3 + 3*a^3*c^4 + 3*a^4*c^2 + 3*a^3*c^3 + 3*a^2*c^4 + a^4*c + a^3*c^2 + a^2*c^3 + a*c^4 + a^3*c - 2*a^2*c^2 + a*c^3)*sqrt(-a*c)*A*e^(3*I*f*x + 3*I*e) - 12*(a^4*c^4 + 3*a^4*c^3 + 3*a^3*c^4 + 3*a^4*c^2 + 3*a^3*c^3 + 3*a^2*c^4 + a^4*c + a^3*c^2 + a^2*c^3 + a*c^4 + a^3*c - 2*a^2*c^2 + a*c^3)*sqrt(a*c)*B*e^(3*I*f*x + 3*I*e) - 4*(a^4*c^4 + 3*a^4*c^3 + 3*a^3*c^4 + 3*a^4*c^2 + 3*a^3*c^3 + 3*a^2*c^4 + a^4*c + a^3*c^2 + a^2*c^3 + a*c...
```

3.789.9 Mupad [**F(-1)**]

Timed out.

$$\int \sqrt{a + ia \tan(e + fx)} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} dx = \int (A + B \tan(e + fx)) \sqrt{a + a \tan(e + fx)} \operatorname{li}(c - c \tan(e + fx) \operatorname{li})^{3/2} dx$$

```
input int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(1/2)*(c - c*tan(e + f*x)*1i)^(3/2), x)
```

```
output int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(1/2)*(c - c*tan(e + f*x)*1i)^(3/2), x)
```

3.790 $\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))\sqrt{c - ict \tan(e + fx)}$

3.790.1 Optimal result	7148
3.790.2 Mathematica [A] (verified)	7148
3.790.3 Rubi [A] (verified)	7149
3.790.4 Maple [A] (verified)	7151
3.790.5 Fricas [B] (verification not implemented)	7151
3.790.6 Sympy [F]	7152
3.790.7 Maxima [B] (verification not implemented)	7152
3.790.8 Giac [F]	7153
3.790.9 Mupad [B] (verification not implemented)	7153

3.790.1 Optimal result

Integrand size = 45, antiderivative size = 104

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))\sqrt{c - ict \tan(e + fx)} dx$$

$$= -\frac{2i\sqrt{a}A\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ict \tan(e+fx)}}\right)}{f} + \frac{B\sqrt{a + ia \tan(e + fx)}\sqrt{c - ict \tan(e + fx)}}{f}$$

output `-2*I*A*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))*a^(1/2)*c^(1/2)/f+B*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)/f`

3.790.2 Mathematica [A] (verified)

Time = 2.42 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.00

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))\sqrt{c - ict \tan(e + fx)} dx$$

$$= -\frac{2i\sqrt{a}A\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ict \tan(e+fx)}}\right)}{f} + \frac{B\sqrt{a + ia \tan(e + fx)}\sqrt{c - ict \tan(e + fx)}}{f}$$

input `Integrate[Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]],x]`

output $((-2*I)*\text{Sqrt}[a]*A*\text{Sqrt}[c]*\text{ArcTan}[(\text{Sqrt}[c]*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]])/(\text{Sqrt}[a]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])])/f + (B*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])/f$

3.790.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.06, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4071, 90, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)} (A + B \tan(e + fx)) dx$$

↓ 3042

$$\int \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)} (A + B \tan(e + fx)) dx$$

↓ 4071

$$\frac{ac \int \frac{A + B \tan(e + fx)}{\sqrt{i \tan(e + fx) a + a} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{f}$$

↓ 90

$$\frac{ac \left(A \int \frac{1}{\sqrt{i \tan(e + fx) a + a} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) + \frac{B \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}}{ac} \right)}{f}$$

↓ 45

$$\frac{ac \left(2A \int \frac{1}{ia + \frac{ic(i \tan(e + fx) a + a)}{c - ic \tan(e + fx)}} d \frac{\sqrt{i \tan(e + fx) a + a}}{\sqrt{c - ic \tan(e + fx)}} + \frac{B \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}}{ac} \right)}{f}$$

↓ 218

$$\frac{ac \left(\frac{B \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}}{ac} - \frac{2iA \arctan \left(\frac{\sqrt{c} \sqrt{a + ia \tan(e + fx)}}{\sqrt{a} \sqrt{c - ic \tan(e + fx)}} \right)}{\sqrt{a} \sqrt{c}} \right)}{f}$$

input $\text{Int}[\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]*(A + B*\text{Tan}[e + f*x])* \text{Sqrt}[c - I*c*\text{Tan}[e + f*x]], x]$

3.790. $\int \sqrt{a + ia \tan(e + fx)} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx$

```
output (a*c*(((2*I)*A*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[
c - I*c*Tan[e + f*x]])]/(Sqrt[a]*Sqrt[c]) + (B*Sqrt[a + I*a*Tan[e + f*x]]
*Sqrt[c - I*c*Tan[e + f*x]])/(a*c)))/f
```

3.790.3.1 Defintions of rubi rules used

```
rule 45 Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; Fre
eQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]
```

```
rule 90 Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
_), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p
+ 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n,
p}, x] && NeQ[n + p + 2, 0]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.790.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.16

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} \left(A \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}} \right) a c+B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{f \sqrt{a c\left(1+\tan (f x+e)^2\right)} \sqrt{a c}}$
default	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} \left(A \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}} \right) a c+B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{f \sqrt{a c\left(1+\tan (f x+e)^2\right)} \sqrt{a c}}$
parts	$\frac{A \sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a c \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}} \right)}{f \sqrt{a c\left(1+\tan (f x+e)^2\right)} \sqrt{a c}} + \frac{B \sqrt{a(1+i \tan (f x+e))}}{f}$

input `int((a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)),x,m
method=_RETURNVERBOSE)`

output `1/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*(A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c+B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c*(1+tan(f*x+e)^2))^(1/2)/(a*c)^(1/2)`

3.790.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 293 vs. 2(78) = 156.

Time = 0.26 (sec) , antiderivative size = 293, normalized size of antiderivative = 2.82

$$\int \sqrt{a+ia \tan (e+f x)}(A+B \tan (e+f x)) \sqrt{c-ic \tan (e+f x)} d x$$

$$4 B \sqrt{\frac{a}{e^{(2 i f x+2 i e)}+1}} \sqrt{\frac{c}{e^{(2 i f x+2 i e)}+1}} e^{(i f x+i e)} + \sqrt{\frac{A^2 a c}{f^2}} f \log \left(\frac{4 \left(2 \left(A e^{(3 i f x+3 i e)}+A e^{(i f x+i e)} \right) \sqrt{\frac{a}{e^{(2 i f x+2 i e)}+1}} \sqrt{\frac{c}{e^{(2 i f x+2 i e)}+1}} \right)}{A e^{(2 i f x+2 i e)}+A} \right)$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)),x, algorithm="fracas")`

3.790. $\int \sqrt{a+ia \tan (e+f x)}(A+B \tan (e+f x)) \sqrt{c-ic \tan (e+f x)} d x$

```
output 1/2*(4*B*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))
)*e^(I*f*x + I*e) + sqrt(A^2*a*c/f^2)*f*log(4*(2*(A*e^(3*I*f*x + 3*I*e) +
A*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x +
2*I*e) + 1)) - sqrt(A^2*a*c/f^2)*(I*f*e^(2*I*f*x + 2*I*e) - I*f))/(A*e^(2
*I*f*x + 2*I*e) + A)) - sqrt(A^2*a*c/f^2)*f*log(4*(2*(A*e^(3*I*f*x + 3*I*e)
) + A*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*
x + 2*I*e) + 1)) - sqrt(A^2*a*c/f^2)*(-I*f*e^(2*I*f*x + 2*I*e) + I*f))/(A*
e^(2*I*f*x + 2*I*e) + A))/f
```

3.790.6 Sympy [F]

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)} dx$$

$$= \int \sqrt{ia(\tan(e + fx) - i)}\sqrt{-ic(\tan(e + fx) + i)}(A + B \tan(e + fx)) dx$$

```
input integrate((a+I*a*tan(f*x+e))**(1/2)*(c-I*c*tan(f*x+e))**(1/2)*(A+B*tan(f*x
+e)),x)
```

```
output Integral(sqrt(I*a*(tan(e + f*x) - I))*sqrt(-I*c*(tan(e + f*x) + I))*(A + B
*tan(e + f*x)), x)
```

3.790.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 447 vs. $2(78) = 156$.

Time = 0.55 (sec) , antiderivative size = 447, normalized size of antiderivative = 4.30

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)} dx =$$

$$\frac{\left(2(A \cos(2fx + 2e) + iA \sin(2fx + 2e) + A) \arctan\left(\cos\left(\frac{1}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)\right)\right)}{\dots}$$

```
input integrate((a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e
)),x, algorithm="maxima")
```

3.790. $\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)} dx$

output $-(2*(A*\cos(2*f*x + 2*e) + I*A*\sin(2*f*x + 2*e) + A)*\arctan2(\cos(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))), \sin(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 1) + 2*(A*\cos(2*f*x + 2*e) + I*A*\sin(2*f*x + 2*e) + A)*\arctan2(\cos(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))), -\sin(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 1) + 4*I*B*\cos(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) - (-I*A*\cos(2*f*x + 2*e) + A*\sin(2*f*x + 2*e) - I*A)*\log(\cos(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e)))^2 + \sin(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e)))^2 + 2*\sin(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 1) - (I*A*\cos(2*f*x + 2*e) - A*\sin(2*f*x + 2*e) + I*A)*\log(\cos(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e)))^2 + \sin(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e)))^2 - 2*\sin(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 1) - 4*B*\sin(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))))*\sqrt{a}*\sqrt{c}/(f*(-2*I*\cos(2*f*x + 2*e) + 2*\sin(2*f*x + 2*e) - 2*I))$

3.790.8 Giac [F]

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)} dx$$

$$= \int (B \tan(fx + e) + A)\sqrt{ia \tan(fx + e) + a}\sqrt{-ictan(fx + e) + c} dx$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e)),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*sqrt(I*a*tan(f*x + e) + a)*sqrt(-I*c*tan(f*x + e) + c), x)`

3.790.9 Mupad [B] (verification not implemented)

Time = 10.46 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.28

$$\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)} dx$$

$$= -\frac{A \sqrt{a} \sqrt{c} \operatorname{atan}\left(\frac{\sqrt{c}(\sqrt{a+a \tan(e+fx)} \operatorname{li}-\sqrt{a})}{\sqrt{a}(\sqrt{c-c \tan(e+fx)} \operatorname{li}-\sqrt{c})}\right)}{f} + \frac{\sqrt{2} B \sqrt{\frac{c}{2 \cos(e+fx)^2 + \sin(2e+2fx) \operatorname{li}}} \sqrt{\frac{a(2 \cos(e+fx)^2 + \sin(2e+2fx) \operatorname{li})}{2 \cos(e+fx)^2}}}{f}$$

3.790. $\int \sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)} dx$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2),x)`

output `(2^(1/2)*B*(c/(sin(2*e + 2*f*x)*1i + 2*cos(e + f*x)^2))^(1/2)*((a*(sin(2*e + 2*f*x)*1i + 2*cos(e + f*x)^2))/(2*cos(e + f*x)^2))^(1/2))/f - (A*a^(1/2)*c^(1/2)*atan((c^(1/2)*((a + a*tan(e + f*x)*1i)^(1/2) - a^(1/2)))/(a^(1/2)*((c - c*tan(e + f*x)*1i)^(1/2) - c^(1/2))))*4i)/f`

3.791
$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} dx$$

3.791.1 Optimal result	7155
3.791.2 Mathematica [A] (verified)	7155
3.791.3 Rubi [A] (verified)	7156
3.791.4 Maple [B] (verified)	7158
3.791.5 Fricas [B] (verification not implemented)	7159
3.791.6 Sympy [F]	7159
3.791.7 Maxima [F(-2)]	7160
3.791.8 Giac [F]	7160
3.791.9 Mupad [B] (verification not implemented)	7161

3.791.1 Optimal result

Integrand size = 45, antiderivative size = 109

$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} dx$$

$$= \frac{2\sqrt{a}B \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{\sqrt{cf}} - \frac{(iA+B)\sqrt{a+ia \tan(e+fx)}}{f\sqrt{c-ictan(e+fx)}}$$

output `2*B*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))*a^(1/2)/f/c^(1/2)-(I*A+B)*(a+I*a*tan(f*x+e))^(1/2)/f/(c-I*c*tan(f*x+e))^(1/2)`

3.791.2 Mathematica [A] (verified)

Time = 3.72 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.01

$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} dx$$

$$= \frac{2\sqrt{a}B \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{\sqrt{cf}} + \frac{(-iA-B)\sqrt{a+ia \tan(e+fx)}}{f\sqrt{c-ictan(e+fx)}}$$

input `Integrate[(Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x]))/Sqrt[c - I*c*Tan[e + f*x]],x]`

3.791.
$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} dx$$

output $(2*\text{Sqrt}[a]*B*\text{ArcTan}[(\text{Sqrt}[c]*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]])/(\text{Sqrt}[a]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])]/(\text{Sqrt}[c]*f) + (((-I)*A - B)*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]])/(f*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])$

3.791.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.06, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4071, 87, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx \\
 & \quad \downarrow \text{4071} \\
 & ac \int \frac{A + B \tan(e + fx)}{\sqrt{i \tan(e + fx) a + a(c - ic \tan(e + fx))^{3/2}}} d \tan(e + fx) \\
 & \quad \downarrow \text{87} \\
 & ac \left(\frac{iB \int \frac{1}{\sqrt{i \tan(e + fx) a + a(c - ic \tan(e + fx))}} d \tan(e + fx)}{c} - \frac{(B + iA) \sqrt{a + ia \tan(e + fx)}}{ac \sqrt{c - ic \tan(e + fx)}} \right) \\
 & \quad \downarrow \text{45} \\
 & ac \left(\frac{2iB \int \frac{1}{ia + \frac{ic(i \tan(e + fx) a + a)}{c - ic \tan(e + fx)}} d \frac{\sqrt{i \tan(e + fx) a + a}}{\sqrt{c - ic \tan(e + fx)}}}{c} - \frac{(B + iA) \sqrt{a + ia \tan(e + fx)}}{ac \sqrt{c - ic \tan(e + fx)}} \right) \\
 & \quad \downarrow \text{218} \\
 & ac \left(\frac{2B \arctan\left(\frac{\sqrt{c} \sqrt{a + ia \tan(e + fx)}}{\sqrt{a} \sqrt{c - ic \tan(e + fx)}}\right)}{\sqrt{ac}^{3/2}} - \frac{(B + iA) \sqrt{a + ia \tan(e + fx)}}{ac \sqrt{c - ic \tan(e + fx)}} \right)
 \end{aligned}$$

3.791. $\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx$

input `Int[(Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x]))/Sqrt[c - I*c*Tan[e + f*x]],x]`

output `(a*c*((2*B*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/(Sqrt[a]*c^(3/2)) - ((I*A + B)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c*Sqrt[c - I*c*Tan[e + f*x]]))/f`

3.791.3.1 Defintions of rubi rules used

rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.791.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 320 vs. 2(88) = 176.

Time = 0.39 (sec) , antiderivative size = 321, normalized size of antiderivative = 2.94

method	result
derivativedivides	$\frac{i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}\left(-2iB\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}}\right)ac\tan(fx+e)-B\ln\right)}{\dots}$
default	$\frac{i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}\left(-2iB\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}}\right)ac\tan(fx+e)-B\ln\right)}{\dots}$
parts	$-\frac{iA\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(i\tan(fx+e)-1)}{fc(i+\tan(fx+e))^2} + \frac{iB\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}}{\dots}$

```
input int((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -I/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)/c*(-2*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)-B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)^2+I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)+B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c+B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c*(1+tan(f*x+e)^2))^(1/2)/(I+tan(f*x+e))^2/(a*c)^(1/2)
```

3.791.
$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$$

3.791.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 332 vs. $2(83) = 166$.

Time = 0.27 (sec) , antiderivative size = 332, normalized size of antiderivative = 3.05

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx =$$

$$cf \sqrt{-\frac{B^2 a}{cf^2}} \log \left(\frac{4 \left(2 (Be^{(3i fx + 3i e)} + Be^{(i fx + i e)}) \sqrt{\frac{a}{e^{(2i fx + 2i e)} + 1}} \sqrt{\frac{c}{e^{(2i fx + 2i e)} + 1}} + (cfe^{(2i fx + 2i e)} - cf) \sqrt{-\frac{B^2 a}{cf^2}} \right)}{Be^{(2i fx + 2i e)} + B} \right) - cf \sqrt{-\frac{B^2 a}{cf^2}}$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x, algorithm="fracas")`

output `-1/2*(c*f*sqrt(-B^2*a/(c*f^2))*log(4*(2*(B*e^(3*I*f*x + 3*I*e) + B*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + (c*f*e^(2*I*f*x + 2*I*e) - c*f)*sqrt(-B^2*a/(c*f^2)))/(B*e^(2*I*f*x + 2*I*e) + B)) - c*f*sqrt(-B^2*a/(c*f^2))*log(4*(2*(B*e^(3*I*f*x + 3*I*e) + B*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - (c*f*e^(2*I*f*x + 2*I*e) - c*f)*sqrt(-B^2*a/(c*f^2)))/(B*e^(2*I*f*x + 2*I*e) + B)) + 2*((I*A + B)*e^(3*I*f*x + 3*I*e) + (I*A + B)*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(c*f)`

3.791.6 Sympy [F]

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx$$

$$= \int \frac{\sqrt{ia (\tan(e + fx) - i)}(A + B \tan(e + fx))}{\sqrt{-ic (\tan(e + fx) + i)}} dx$$

input `integrate((a+I*a*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(1/2),x)`

output `Integral(sqrt(I*a*(tan(e + f*x) - I))*(A + B*tan(e + f*x))/sqrt(-I*c*(tan(e + f*x) + I)), x)`

3.791. $\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$

3.791.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.791.8 Giac [F]

$$\begin{aligned} & \int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx \\ &= \int \frac{(B \tan(fx + e) + A) \sqrt{ia \tan(fx + e) + a}}{\sqrt{-ic \tan(fx + e) + c}} dx \end{aligned}$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*sqrt(I*a*tan(f*x + e) + a)/sqrt(-I*c*tan(f*x + e) + c), x)`

3.791.9 Mupad [B] (verification not implemented)

Time = 11.89 (sec) , antiderivative size = 266, normalized size of antiderivative = 2.44

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx = \frac{4B\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{c}(\sqrt{a+a \tan(e+fx) \operatorname{li}-\sqrt{a}})}{\sqrt{a}(\sqrt{c-c \tan(e+fx) \operatorname{li}-\sqrt{c}})}\right)}{\sqrt{c} f} + \frac{A\sqrt{a+a \tan(e+fx) \operatorname{li}}\sqrt{c-c \tan(e+fx) \operatorname{li}}}{cf(\tan(e+fx) + \operatorname{li})} - \frac{4Ba(\sqrt{a+a \tan(e+fx) \operatorname{li}} - \sqrt{a})}{cf(\sqrt{c-c \tan(e+fx) \operatorname{li}} - \sqrt{c})\left(\frac{a}{c} - \frac{(\sqrt{a+a \tan(e+fx) \operatorname{li}-\sqrt{a}})^2}{(\sqrt{c-c \tan(e+fx) \operatorname{li}-\sqrt{c}})^2} + \frac{2\sqrt{a}(\sqrt{a+a \tan(e+fx) \operatorname{li}-\sqrt{a}})}{\sqrt{c}(\sqrt{c-c \tan(e+fx) \operatorname{li}-\sqrt{c}})}\right)}$$

```
input int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(1/2))/(c - c*tan(e + f*x)*1i)^(1/2),x)
```

```
output (4*B*a^(1/2)*atan((c^(1/2)*((a + a*tan(e + f*x)*1i)^(1/2) - a^(1/2)))/(a^(1/2)*((c - c*tan(e + f*x)*1i)^(1/2) - c^(1/2))))/(c^(1/2)*f) + (A*(a + a*tan(e + f*x)*1i)^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2))/(c*f*(tan(e + f*x) + 1i)) - (4*B*a*((a + a*tan(e + f*x)*1i)^(1/2) - a^(1/2)))/(c*f*((c - c*tan(e + f*x)*1i)^(1/2) - c^(1/2))*(a/c - ((a + a*tan(e + f*x)*1i)^(1/2) - a^(1/2))^2/((c - c*tan(e + f*x)*1i)^(1/2) - c^(1/2))^2 + (2*a^(1/2)*((a + a*tan(e + f*x)*1i)^(1/2) - a^(1/2)))/(c^(1/2)*((c - c*tan(e + f*x)*1i)^(1/2) - c^(1/2))))))
```


3.792
$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$$

3.792.1 Optimal result	7162
3.792.2 Mathematica [A] (verified)	7162
3.792.3 Rubi [A] (verified)	7163
3.792.4 Maple [A] (verified)	7164
3.792.5 Fricas [A] (verification not implemented)	7165
3.792.6 Sympy [F]	7165
3.792.7 Maxima [F(-2)]	7166
3.792.8 Giac [F]	7166
3.792.9 Mupad [B] (verification not implemented)	7166

3.792.1 Optimal result

Integrand size = 45, antiderivative size = 102

$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx = -\frac{(iA+B)\sqrt{a+ia \tan(e+fx)}}{3f(c-ic \tan(e+fx))^{3/2}} - \frac{(iA-2B)\sqrt{a+ia \tan(e+fx)}}{3cf\sqrt{c-ic \tan(e+fx)}}$$

output `-1/3*(I*A-2*B)*(a+I*a*tan(f*x+e))^(1/2)/c/f/(c-I*c*tan(f*x+e))^(1/2)-1/3*(I*A+B)*(a+I*a*tan(f*x+e))^(1/2)/f/(c-I*c*tan(f*x+e))^(3/2)`

3.792.2 Mathematica [A] (verified)

Time = 4.94 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.79

$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx = \frac{\sqrt{a+ia \tan(e+fx)}(2A+iB+(-iA+2B)\tan(e+fx))}{3cf(i+\tan(e+fx))\sqrt{c-ic \tan(e+fx)}}$$

input `Integrate[(Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(Sqrt[a + I*a*Tan[e + f*x]]*(2*A + I*B + ((-I)*A + 2*B)*Tan[e + f*x]))/(3*c*f*(I + Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])`

3.792.
$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$$

3.792.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.089$, Rules used = {3042, 4071, 87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx \\
 \downarrow 3042 \\
 \int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx \\
 \downarrow 4071 \\
 ac \int \frac{A + B \tan(e + fx)}{\sqrt{i \tan(e + fx) a + a(c - ic \tan(e + fx))^{5/2}} d \tan(e + fx)} \\
 \downarrow 87 \\
 ac \left(\frac{(A + 2iB) \int \frac{1}{\sqrt{i \tan(e + fx) a + a(c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{3c} - \frac{(B + iA) \sqrt{a + ia \tan(e + fx)}}{3ac(c - ic \tan(e + fx))^{3/2}} \right) \\
 \downarrow 48 \\
 ac \left(-\frac{i(A + 2iB) \sqrt{a + ia \tan(e + fx)}}{3ac^2 \sqrt{c - ic \tan(e + fx)}} - \frac{(B + iA) \sqrt{a + ia \tan(e + fx)}}{3ac(c - ic \tan(e + fx))^{3/2}} \right)
 \end{array}$$

input `Int[(Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(a*c*(-1/3*((I*A + B)*Sqrt[a + I*a*Tan[e + f*x]]))/(a*c*(c - I*c*Tan[e + f*x])^(3/2)) - ((I/3)*(A + (2*I)*B)*Sqrt[a + I*a*Tan[e + f*x]]/(a*c^2*Sqrt[c - I*c*Tan[e + f*x]]))/f`

3.792.3.1 Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.792.4 Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.82

method	result
risch	$-\frac{\sqrt{\frac{ae^{2i(fx+e)}}{e^{2i(fx+e)}+1}}(iAe^{2i(fx+e)}+Be^{2i(fx+e)}+3iA-3B)}{6c\sqrt{\frac{c}{e^{2i(fx+e)}+1}}}f$
derivativedivides	$\frac{\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(2iB\tan(fx+e)^2+3iA\tan(fx+e)+A\tan(fx+e)^2-iB-3B\tan(fx+e)-2A)}{3fc^2(i+\tan(fx+e))^3}$
default	$\frac{\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(2iB\tan(fx+e)^2+3iA\tan(fx+e)+A\tan(fx+e)^2-iB-3B\tan(fx+e)-2A)}{3fc^2(i+\tan(fx+e))^3}$
parts	$\frac{A\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(3i\tan(fx+e)+\tan(fx+e)^2-2)}{3fc^2(i+\tan(fx+e))^3} + \frac{iB\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}}{3fc^2(i+\tan(fx+e))^3}$

3.792.
$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{3/2}} dx$$

```
input int((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
```

```
output -1/6/c*(a*exp(2*I*(f*x+e))/(exp(2*I*(f*x+e))+1))^(1/2)/(c/(exp(2*I*(f*x+e))+1))^(1/2)*(I*A*exp(2*I*(f*x+e))+B*exp(2*I*(f*x+e))+3*I*A-3*B)/f
```

3.792.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.94

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{3/2}} dx = \frac{((-iA - B)e^{(5ifx+5ie)} - 2(2iA - B)e^{(3ifx+3ie)} - 3(iA - B)e^{(ifx+ie)})\sqrt{a/(e^{(2ifx+2ie)} + 1)}\sqrt{c/(e^{(2ifx+2ie)} + 1)}}{6c^2f}$$

```
input integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="fracas")
```

```
output 1/6*((-I*A - B)*e^(5*I*f*x + 5*I*e) - 2*(2*I*A - B)*e^(3*I*f*x + 3*I*e) - 3*(I*A - B)*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^2*f)
```

3.792.6 Sympy [F]

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{3/2}} dx = \int \frac{\sqrt{ia(\tan(e + fx) - i)}(A + B \tan(e + fx))}{(-ic(\tan(e + fx) + i))^{3/2}} dx$$

```
input integrate((a+I*a*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(3/2),x)
```

```
output Integral(sqrt(I*a*(tan(e + f*x) - I))*(A + B*tan(e + f*x))/(-I*c*(tan(e + f*x) + I))**(3/2), x)
```

3.792.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.792.8 Giac [F]

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{3/2}} dx = \int \frac{(B \tan(fx + e) + A)\sqrt{ia \tan(fx + e) + a}}{(-ictan(fx + e) + c)^{\frac{3}{2}}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*sqrt(I*a*tan(f*x + e) + a)/(-I*c*tan(f*x + e) + c)^(3/2), x)`

3.792.9 Mupad [B] (verification not implemented)

Time = 1.23 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.42

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{3/2}} dx =$$

$$\frac{\sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}}(A3i - 3B + A \cos(2e + 2fx) 1i + B \cos(2e + 2fx) - A \sin(2e + 2fx))}{6cf \sqrt{\frac{c(\cos(2e+2fx)+1-\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}}}$$

3.792. $\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{3/2}} dx$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(1/2))/(c - c*tan(e + f*x)*1i)^(3/2),x)`

output `-(((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*3i - 3*B + A*cos(2*e + 2*f*x)*1i + B*cos(2*e + 2*f*x) - A*sin(2*e + 2*f*x) + B*sin(2*e + 2*f*x)*1i))/(6*c*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

3.792.
$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{3/2}} dx$$

3.793
$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{5/2}} dx$$

3.793.1 Optimal result 7168
 3.793.2 Mathematica [A] (verified) 7168
 3.793.3 Rubi [A] (verified) 7169
 3.793.4 Maple [A] (verified) 7171
 3.793.5 Fricas [A] (verification not implemented) 7171
 3.793.6 Sympy [F] 7172
 3.793.7 Maxima [F(-2)] 7172
 3.793.8 Giac [F] 7172
 3.793.9 Mupad [B] (verification not implemented) 7173

3.793.1 Optimal result

Integrand size = 45, antiderivative size = 155

$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{5/2}} dx = -\frac{(iA+B)\sqrt{a+ia \tan(e+fx)}}{5f(c-ictan(e+fx))^{5/2}} - \frac{(2iA-3B)\sqrt{a+ia \tan(e+fx)}}{15cf(c-ictan(e+fx))^{3/2}} - \frac{(2iA-3B)\sqrt{a+ia \tan(e+fx)}}{15c^2f\sqrt{c-ictan(e+fx)}}$$

output `-1/15*(2*I*A-3*B)*(a+I*a*tan(f*x+e))^(1/2)/c^2/f/(c-I*c*tan(f*x+e))^(1/2)-1/5*(I*A+B)*(a+I*a*tan(f*x+e))^(1/2)/f/(c-I*c*tan(f*x+e))^(5/2)-1/15*(2*I*A-3*B)*(a+I*a*tan(f*x+e))^(1/2)/c/f/(c-I*c*tan(f*x+e))^(3/2)`

3.793.2 Mathematica [A] (verified)

Time = 6.20 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.71

$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{5/2}} dx = \frac{a(-i+\tan(e+fx))(-7A-3iB+(6iA-9B)\tan(e+fx))}{15c^2f(i+\tan(e+fx))^2\sqrt{a+ia \tan(e+fx)}}$$

input `Integrate[(Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(5/2),x]`

output `(a*(-I + Tan[e + f*x])*(-7*A - (3*I)*B + ((6*I)*A - 9*B)*Tan[e + f*x] + (2*A + (3*I)*B)*Tan[e + f*x]^2))/(15*c^2*f*(I + Tan[e + f*x])^2*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])`

3.793.
$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{5/2}} dx$$

3.793.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.08, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4071, 87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx \\
 \downarrow 3042 \\
 \int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx \\
 \downarrow 4071 \\
 ac \int \frac{A+B \tan(e+fx)}{\sqrt{i \tan(e+fx)a+a(c-ic \tan(e+fx))^{7/2}} d \tan(e+fx)} \\
 \downarrow 87 \\
 ac \left(\frac{(2A+3iB) \int \frac{1}{\sqrt{i \tan(e+fx)a+a(c-ic \tan(e+fx))^{5/2}} d \tan(e+fx)}{5c} - \frac{(B+iA)\sqrt{a+ia \tan(e+fx)}}{5ac(c-ic \tan(e+fx))^{5/2}} \right) \\
 \downarrow 55 \\
 ac \left(\frac{(2A+3iB) \left(\frac{\int \frac{1}{\sqrt{i \tan(e+fx)a+a(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{3c} - \frac{i\sqrt{a+ia \tan(e+fx)}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{5c} - \frac{(B+iA)\sqrt{a+ia \tan(e+fx)}}{5ac(c-ic \tan(e+fx))^{5/2}} \right) \\
 \downarrow 48 \\
 ac \left(\frac{(2A+3iB) \left(-\frac{i\sqrt{a+ia \tan(e+fx)}}{3ac^2\sqrt{c-ic \tan(e+fx)}} - \frac{i\sqrt{a+ia \tan(e+fx)}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{5c} - \frac{(B+iA)\sqrt{a+ia \tan(e+fx)}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)
 \end{array}$$

input `Int[(Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(5/2),x]`

$$3.793. \quad \int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$$


```
output (a*c*(-1/5*((I*A + B)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c*(c - I*c*Tan[e + f*x])^(5/2)) + ((2*A + (3*I)*B)*((-1/3*I)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c*(c - I*c*Tan[e + f*x])^(3/2)) - ((I/3)*Sqrt[a + I*a*Tan[e + f*x]]/(a*c^2*Sqrt[c - I*c*Tan[e + f*x]])))/(5*c))/f
```

3.793.3.1 Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 55 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.793.4 Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.63

method	result
risch	$-\frac{\sqrt{\frac{ae^{2i(fx+e)}}{e^{2i(fx+e)}+1}} (3iAe^{4i(fx+e)}+3Be^{4i(fx+e)}+10iAe^{2i(fx+e)}+15iA-15B)}{60c^2\sqrt{\frac{c}{e^{2i(fx+e)}+1}}f}$
derivativedivides	$-\frac{i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(2iA\tan(fx+e)^3-12iB\tan(fx+e)^2-3B\tan(fx+e)^3-13iA\tan(fx+e))}{15fc^3(i+\tan(fx+e))^4}$
default	$-\frac{i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(2iA\tan(fx+e)^3-12iB\tan(fx+e)^2-3B\tan(fx+e)^3-13iA\tan(fx+e))}{15fc^3(i+\tan(fx+e))^4}$
parts	$\frac{A\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(8i\tan(fx+e)^2+2\tan(fx+e)^3-7i-13\tan(fx+e))}{15fc^3(i+\tan(fx+e))^4} + \frac{iB\sqrt{a(1+i\tan(fx+e))}}{15fc^3(i+\tan(fx+e))^4}$

input `int((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

output `-1/60/c^2*(a*exp(2*I*(f*x+e))/(exp(2*I*(f*x+e))+1))^(1/2)/(c/(exp(2*I*(f*x+e))+1))^(1/2)*(3*I*A*exp(4*I*(f*x+e))+3*B*exp(4*I*(f*x+e))+10*I*A*exp(2*I*(f*x+e))+15*I*A-15*B)/f`

3.793.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.73

$$\int \frac{\sqrt{a+ia\tan(e+fx)}(A+B\tan(e+fx))}{(c-ictan(e+fx))^{5/2}} dx = \frac{(3(iA+B)e^{(7ifx+7ie)} - (-13iA-3B)e^{(5ifx+5ie)} + 5(5iA-3B)e^{(3ifx+3ie)} + 15(iA-B)e^{(ifx+ie)})\sqrt{a}}{60c^3f}$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,algorithm="fracas")`

output `-1/60*(3*(I*A+B)*e^(7*I*f*x+7*I*e) - (-13*I*A-3*B)*e^(5*I*f*x+5*I*e) + 5*(5*I*A-3*B)*e^(3*I*f*x+3*I*e) + 15*(I*A-B)*e^(I*f*x+I*e))*sqrt(a/(e^(2*I*f*x+2*I*e)+1))*sqrt(c/(e^(2*I*f*x+2*I*e)+1))/(c^3*f)`

3.793.
$$\int \frac{\sqrt{a+ia\tan(e+fx)}(A+B\tan(e+fx))}{(c-ictan(e+fx))^{5/2}} dx$$

3.793.6 Sympy [F]

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{5/2}} dx = \int \frac{\sqrt{ia (\tan(e + fx) - i)}(A + B \tan(e + fx))}{(-ic (\tan(e + fx) + i))^{5/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(5/2),x)`

output `Integral(sqrt(I*a*(tan(e + f*x) - I))*(A + B*tan(e + f*x))/(-I*c*(tan(e + f*x) + I))**(5/2), x)`

3.793.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.793.8 Giac [F]

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{5/2}} dx = \int \frac{(B \tan(fx + e) + A)\sqrt{ia \tan(fx + e) + a}}{(-ic \tan(fx + e) + c)^{5/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*sqrt(I*a*tan(f*x + e) + a)/(-I*c*tan(f*x + e) + c)^(5/2), x)`

3.793. $\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{5/2}} dx$

3.793.9 Mupad [B] (verification not implemented)

Time = 9.36 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.10

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{5/2}} dx =$$

$$\frac{\sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}} (A 15i - 15 B + A \cos(2e + 2fx) 10i + A \cos(4e + 4fx) 3i + 3 B \cos(4e + 4fx) 3i)}{60 c^2 f \sqrt{\frac{c(\cos(2e+2fx)+1-\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}}}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(1/2))/(c - c*tan(e + f*x)*1i)^(5/2),x)`

output `-(((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*15i - 15*B + A*cos(2*e + 2*f*x)*10i + A*cos(4*e + 4*f*x)*3i + 3*B*cos(4*e + 4*f*x) - 10*A*sin(2*e + 2*f*x) - 3*A*sin(4*e + 4*f*x) + B*sin(4*e + 4*f*x)*3i))/(60*c^2*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

3.794
$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{7/2}} dx$$

3.794.1 Optimal result	7174
3.794.2 Mathematica [A] (verified)	7174
3.794.3 Rubi [A] (verified)	7175
3.794.4 Maple [A] (verified)	7177
3.794.5 Fricas [A] (verification not implemented)	7178
3.794.6 Sympy [F]	7178
3.794.7 Maxima [F(-2)]	7179
3.794.8 Giac [F]	7179
3.794.9 Mupad [B] (verification not implemented)	7179

3.794.1 Optimal result

Integrand size = 45, antiderivative size = 208

$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{7/2}} dx =$$

$$\frac{(iA+B)\sqrt{a+ia \tan(e+fx)}}{7f(c-ictan(e+fx))^{7/2}} - \frac{(3iA-4B)\sqrt{a+ia \tan(e+fx)}}{35cf(c-ictan(e+fx))^{5/2}}$$

$$- \frac{2(3iA-4B)\sqrt{a+ia \tan(e+fx)}}{105c^2f(c-ictan(e+fx))^{3/2}} - \frac{2(3iA-4B)\sqrt{a+ia \tan(e+fx)}}{105c^3f\sqrt{c-ictan(e+fx)}}$$

output

```
-2/105*(3*I*A-4*B)*(a+I*a*tan(f*x+e))^(1/2)/c^3/f/(c-I*c*tan(f*x+e))^(1/2)
-1/7*(I*A+B)*(a+I*a*tan(f*x+e))^(1/2)/f/(c-I*c*tan(f*x+e))^(7/2)-1/35*(3*I
*A-4*B)*(a+I*a*tan(f*x+e))^(1/2)/c/f/(c-I*c*tan(f*x+e))^(5/2)-2/105*(3*I*A
-4*B)*(a+I*a*tan(f*x+e))^(1/2)/c^2/f/(c-I*c*tan(f*x+e))^(3/2)
```

3.794.2 Mathematica [A] (verified)

Time = 6.61 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.63

$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{7/2}} dx = \frac{a(-i+\tan(e+fx))(-36iA+13B-13(3A+4iB)\tan(e+fx))}{105c^3f(i+\tan(e+fx))^3\sqrt{c-ictan(e+fx)}}$$

input

```
Integrate[(Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e
+ f*x])^(7/2), x]
```

3.794.
$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{7/2}} dx$$

output $(a*(-I + \text{Tan}[e + f*x])*((-36*I)*A + 13*B - 13*(3*A + (4*I)*B)*\text{Tan}[e + f*x] + (8*I)*(3*A + (4*I)*B)*\text{Tan}[e + f*x]^2 + (6*A + (8*I)*B)*\text{Tan}[e + f*x]^3) / (105*c^3*f*(I + \text{Tan}[e + f*x])^3*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])$

3.794.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 87, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx$$

↓ 3042

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx$$

↓ 4071

$$ac \int \frac{A + B \tan(e + fx)}{\sqrt{i \tan(e + fx)a + a(c - ic \tan(e + fx))^{9/2}}} d \tan(e + fx)$$

f
↓ 87

$$ac \left(\frac{(3A + 4iB) \int \frac{1}{\sqrt{i \tan(e + fx)a + a(c - ic \tan(e + fx))^{7/2}}} d \tan(e + fx)}{7c} - \frac{(B + iA) \sqrt{a + ia \tan(e + fx)}}{7ac(c - ic \tan(e + fx))^{7/2}} \right)$$

f
↓ 55

$$ac \left(\frac{(3A + 4iB) \left(\frac{2 \int \frac{1}{\sqrt{i \tan(e + fx)a + a(c - ic \tan(e + fx))^{5/2}}} d \tan(e + fx)}{5c} - \frac{i \sqrt{a + ia \tan(e + fx)}}{5ac(c - ic \tan(e + fx))^{5/2}} \right)}{7c} - \frac{(B + iA) \sqrt{a + ia \tan(e + fx)}}{7ac(c - ic \tan(e + fx))^{7/2}} \right)$$

f
↓ 55

3.794. $\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx$

$$\begin{array}{c}
 \left(\frac{(3A+4iB) \left(\frac{2 \left(\frac{\int \frac{1}{\sqrt{i \tan(e+fx)a+a(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{3c} - \frac{i \sqrt{a+ia \tan(e+fx)}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{5c} - \frac{i \sqrt{a+ia \tan(e+fx)}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)}{7c} - \frac{(B+iA) \sqrt{a+ia \tan(e+fx)}}{7ac(c-ic \tan(e+fx))^{7/2}} \right)}{f} \\
 \downarrow 48 \\
 \left(\frac{(3A+4iB) \left(\frac{2 \left(-\frac{i \sqrt{a+ia \tan(e+fx)}}{3ac^2 \sqrt{c-ic \tan(e+fx)}} - \frac{i \sqrt{a+ia \tan(e+fx)}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{5c} - \frac{i \sqrt{a+ia \tan(e+fx)}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)}{7c} - \frac{(B+iA) \sqrt{a+ia \tan(e+fx)}}{7ac(c-ic \tan(e+fx))^{7/2}} \right)}{f}
 \end{array}$$

input `Int[(Sqrt[a + I*a*Tan[e + f*x]]*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(7/2), x]`

output `(a*c*(-1/7*((I*A + B)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c*(c - I*c*Tan[e + f*x])^(7/2)) + ((3*A + (4*I)*B)*((-1/5*I)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c*(c - I*c*Tan[e + f*x])^(5/2)) + (2*(((1/3)*I)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c*(c - I*c*Tan[e + f*x])^(3/2)) - ((I/3)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c^2*Sqrt[c - I*c*Tan[e + f*x]])))/(5*c))/(7*c))/f`

3.794.3.1 Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

3.794. $\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$

```
rule 55 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x]
, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.794.4 Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.65

method	result
risch	$-\frac{\sqrt{\frac{ae^{2i(fx+e)}}{e^{2i(fx+e)}+1}} (15iAe^{6i(fx+e)}+15Be^{6i(fx+e)}+63iAe^{4i(fx+e)}+21Be^{4i(fx+e)}+105iAe^{2i(fx+e)}-35Be^{2i(fx+e)}+105)}{840c^3\sqrt{\frac{c}{e^{2i(fx+e)}+1}}}f$
derivativedivides	$\frac{\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(8iB\tan(fx+e)^4+30iA\tan(fx+e)^3+6A\tan(fx+e)^4-84iB\tan(fx+e)^2-105fc^4(i+\tan(fx+e))^5)}{105fc^4(i+\tan(fx+e))^5}$
default	$\frac{\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(8iB\tan(fx+e)^4+30iA\tan(fx+e)^3+6A\tan(fx+e)^4-84iB\tan(fx+e)^2-105fc^4(i+\tan(fx+e))^5)}{105fc^4(i+\tan(fx+e))^5}$
parts	$\frac{A\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(10i\tan(fx+e)^3+2\tan(fx+e)^4-25i\tan(fx+e)-21\tan(fx+e)^2+12)}{35fc^4(i+\tan(fx+e))^5} +$

3.794.
$$\int \frac{\sqrt{a+ia\tan(e+fx)}(A+B\tan(e+fx))}{(c-ictan(e+fx))^{7/2}} dx$$


```
input int((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x,method=_RETURNVERBOSE)
```

```
output -1/840/c^3*(a*exp(2*I*(f*x+e))/(exp(2*I*(f*x+e))+1))^(1/2)/(c/(exp(2*I*(f*x+e))+1))^(1/2)*(15*I*A*exp(6*I*(f*x+e))+15*B*exp(6*I*(f*x+e))+63*I*A*exp(4*I*(f*x+e))+21*B*exp(4*I*(f*x+e))+105*I*A*exp(2*I*(f*x+e))-35*B*exp(2*I*(f*x+e))+105*I*A-105*B)/f
```

3.794.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.63

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx =$$

$$\frac{(15(iA + B)e^{(9ifx+9ie)} + 6(13iA + 6B)e^{(7ifx+7ie)} + 14(12iA - B)e^{(5ifx+5ie)} + 70(3iA - 2B)e^{(3ifx+3ie)} + 105(IA - B)e^{(Ifx + Ie)})\sqrt{a/(e^{(2Ifx + 2Ie)} + 1)}\sqrt{c/(e^{(2Ifx + 2Ie)} + 1)}}{840c^4f}$$

```
input integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x, algorithm="fracas")
```

```
output -1/840*(15*(I*A + B)*e^(9*I*f*x + 9*I*e) + 6*(13*I*A + 6*B)*e^(7*I*f*x + 7*I*e) + 14*(12*I*A - B)*e^(5*I*f*x + 5*I*e) + 70*(3*I*A - 2*B)*e^(3*I*f*x + 3*I*e) + 105*(I*A - B)*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^4*f)
```

3.794.6 Sympy [F]

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx = \int \frac{\sqrt{ia(\tan(e + fx) - i)}(A + B \tan(e + fx))}{(-ic(\tan(e + fx) + i))^{7/2}} dx$$

```
input integrate((a+I*a*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(7/2),x)
```

```
output Integral(sqrt(I*a*(tan(e + f*x) - I))*(A + B*tan(e + f*x))/(-I*c*(tan(e + f*x) + I))**(7/2), x)
```

3.794. $\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{7/2}} dx$

3.794.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.794.8 Giac [F]

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx = \int \frac{(B \tan(fx + e) + A)\sqrt{ia \tan(fx + e) + a}}{(-ictan(fx + e) + c)^{7/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*sqrt(I*a*tan(f*x + e) + a)/(-I*c*tan(f*x + e) + c)^(7/2), x)`

3.794.9 Mupad [B] (verification not implemented)

Time = 10.13 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.18

$$\int \frac{\sqrt{a + ia \tan(e + fx)}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx =$$

$$\sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)li)}{\cos(2e+2fx)+1}} (A 105i - 105 B + A \cos(2e + 2fx) 105i + A \cos(4e + 4fx) 63i + A c$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(1/2))/(c - c*tan(e + f*x)*1i)^(7/2),x)`

output `-(((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*105i - 105*B + A*cos(2*e + 2*f*x)*105i + A*cos(4*e + 4*f*x)*63i + A*cos(6*e + 6*f*x)*15i - 35*B*cos(2*e + 2*f*x) + 21*B*cos(4*e + 4*f*x) + 15*B*cos(6*e + 6*f*x) - 105*A*sin(2*e + 2*f*x) - 63*A*sin(4*e + 4*f*x) - 15*A*sin(6*e + 6*f*x) - B*sin(2*e + 2*f*x)*35i + B*sin(4*e + 4*f*x)*21i + B*sin(6*e + 6*f*x)*15i))/(840*c^3*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

3.794.
$$\int \frac{\sqrt{a+ia \tan(e+fx)}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{7/2}} dx$$

3.795 $\int (a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{7/2} dx$

3.795.1 Optimal result	7181
3.795.2 Mathematica [A] (verified)	7182
3.795.3 Rubi [A] (verified)	7182
3.795.4 Maple [A] (verified)	7185
3.795.5 Fricas [B] (verification not implemented)	7186
3.795.6 Sympy [F(-1)]	7187
3.795.7 Maxima [B] (verification not implemented)	7187
3.795.8 Giac [F(-1)]	7188
3.795.9 Mupad [F(-1)]	7189

3.795.1 Optimal result

Integrand size = 45, antiderivative size = 279

$$\int (a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{7/2} dx =$$

$$\frac{a^{3/2}(5iA-2B)c^{7/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{4f}$$

$$+ \frac{a(5A+2iB)c^3 \tan(e+fx) \sqrt{a+ia \tan(e+fx)} \sqrt{c-ictan(e+fx)}}{8f}$$

$$- \frac{(5iA-2B)c^2(a+ia \tan(e+fx))^{3/2}(c-ictan(e+fx))^{3/2}}{12f}$$

$$- \frac{(5iA-2B)c(a+ia \tan(e+fx))^{3/2}(c-ictan(e+fx))^{5/2}}{20f}$$

$$+ \frac{B(a+ia \tan(e+fx))^{3/2}(c-ictan(e+fx))^{7/2}}{5f}$$

output

```
-1/4*a^(3/2)*(5*I*A-2*B)*c^(7/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a
^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/f+1/8*a*(5*A+2*I*B)*c^3*(a+I*a*tan(f*x+e)
)^(1/2)*(c-I*c*tan(f*x+e))^(1/2)*tan(f*x+e)/f-1/12*(5*I*A-2*B)*c^2*(a+I*a*
tan(f*x+e))^(3/2)*(c-I*c*tan(f*x+e))^(3/2)/f-1/20*(5*I*A-2*B)*c*(a+I*a*tan
(f*x+e))^(3/2)*(c-I*c*tan(f*x+e))^(5/2)/f+1/5*B*(a+I*a*tan(f*x+e))^(3/2)*(
c-I*c*tan(f*x+e))^(7/2)/f
```

3.795.2 Mathematica [A] (verified)

Time = 11.54 (sec) , antiderivative size = 456, normalized size of antiderivative = 1.63

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{7/2} dx = \frac{B(a + ia \tan(e + fx))^{3/2} (c - ictan(e + fx))^{7/2}}{5f}$$

$$+ \frac{-\frac{a(5iA-2B)c(a+ia \tan(e+fx))^{3/2}(c-ictan(e+fx))^{5/2}}{4f} + \frac{-5a^2(5iA-2B)c^2(a+ia \tan(e+fx))^{3/2}(c-ictan(e+fx))^{3/2}}{3f} - \frac{15a^3(5iA-2B)c^3(a+ia \tan(e+fx))^{3/2}(c-ictan(e+fx))^{1/2}}{3f}}{1}$$

input `Integrate[(a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2),x]`

output `(B*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(7/2))/(5*f) + (-1/4*(a*((5*I)*A - 2*B)*c*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(5/2))/f + ((-5*a^2*((5*I)*A - 2*B)*c^2*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(3/2))/(3*f) + ((-15*a^3*((5*I)*A - 2*B)*c^3*(a + I*a*Tan[e + f*x])^(3/2)*Sqrt[c - I*c*Tan[e + f*x]])/(2*f) + ((-15*a^4*((5*I)*A - 2*B)*c^4*(a + I*a*Tan[e + f*x])^(3/2))/(f*Sqrt[c - I*c*Tan[e + f*x]]) + (30*a^6*((5*I)*A - 2*B)*c^4*(1 - I*Tan[e + f*x])*((1 + I*Tan[e + f*x])/(1 - I*Tan[e + f*x]) - (ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a]])*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[1 - I*Tan[e + f*x]])))/(f*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/(2*a))/(3*a)/(4*a))/(5*a)`

3.795.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.95, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.178$, Rules used = {3042, 4071, 90, 59, 59, 40, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))^{3/2} (c - ictan(e + fx))^{7/2} (A + B \tan(e + fx)) dx$$

3.795. $\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{7/2} dx$

↓ 3042

$$\int (a + ia \tan(e + fx))^{3/2} (c - ictan(e + fx))^{7/2} (A + B \tan(e + fx)) dx$$

↓ 4071

$$\frac{ac \int \sqrt{i \tan(e + fx)a + a} (A + B \tan(e + fx)) (c - ictan(e + fx))^{5/2} d \tan(e + fx)}{f}$$

↓ 90

$$\frac{ac \left(\frac{1}{5} (5A + 2iB) \int \sqrt{i \tan(e + fx)a + a} (c - ictan(e + fx))^{5/2} d \tan(e + fx) + \frac{B(a + ia \tan(e + fx))^{3/2} (c - ictan(e + fx))^{7/2}}{5ac} \right)}{f}$$

↓ 59

$$\frac{ac \left(\frac{1}{5} (5A + 2iB) \left(\frac{5}{4} c \int \sqrt{i \tan(e + fx)a + a} (c - ictan(e + fx))^{3/2} d \tan(e + fx) - \frac{i(a + ia \tan(e + fx))^{3/2} (c - ictan(e + fx))^{7/2}}{4a} \right) \right)}{f}$$

↓ 59

$$\frac{ac \left(\frac{1}{5} (5A + 2iB) \left(\frac{5}{4} c \left(c \int \sqrt{i \tan(e + fx)a + a} \sqrt{c - ictan(e + fx)} d \tan(e + fx) - \frac{i(a + ia \tan(e + fx))^{3/2} (c - ictan(e + fx))^{7/2}}{3a} \right) \right) \right)}{f}$$

↓ 40

$$\frac{ac \left(\frac{1}{5} (5A + 2iB) \left(\frac{5}{4} c \left(c \left(\frac{1}{2} ac \int \frac{1}{\sqrt{i \tan(e + fx)a + a} \sqrt{c - ictan(e + fx)}} d \tan(e + fx) + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ictan(e + fx)} \right) \right) \right) \right)}$$

↓ 45

$$\frac{ac \left(\frac{1}{5} (5A + 2iB) \left(\frac{5}{4} c \left(c \left(ac \int \frac{1}{ia + \frac{ic(i \tan(e + fx)a + a)}{c - ictan(e + fx)}} d \frac{\sqrt{i \tan(e + fx)a + a}}{\sqrt{c - ictan(e + fx)}} + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ictan(e + fx)} \right) \right) \right) \right)}$$

↓ 218

$$\frac{ac \left(\frac{1}{5} (5A + 2iB) \left(\frac{5}{4} c \left(c \left(\frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ictan(e + fx)} - i\sqrt{a}\sqrt{c} \arctan \left(\frac{\sqrt{c}\sqrt{a + ia \tan(e + fx)}}{\sqrt{a}\sqrt{c - ictan(e + fx)}} \right) \right) \right) \right) \right)}$$

input `Int[(a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2),x]`

output `(a*c*((B*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(7/2))/(5*a*c) + ((5*A + (2*I)*B)*((-1/4*I)*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(5/2))/a + (5*c*((-1/3*I)*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(3/2))/a + c*((-I)*Sqrt[a]*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])]) + (Tan[e + f*x]*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/2))/4)/5)/f`

3.795.3.1 Defintions of rubi rules used

rule 40 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^n/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`

rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 59 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[2*c*(n/(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0] && IGtQ[n + 1/2, 0] && LtQ[m, n]`

rule 90 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.795.4 Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 412, normalized size of antiderivative = 1.48

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} c^3 a \left(60 i B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)} \tan (f x+e)^3+24 B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}\right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} c^3 a \left(60 i B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)} \tan (f x+e)^3+24 B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}\right)}{\dots}$
parts	$\frac{A \sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} c^3 a \left(16 i \tan (f x+e)^2 \sqrt{a c\left(1+\tan (f x+e)^2\right)} \sqrt{a c}+6 \tan (f x+e)^3 \sqrt{a c\left(1+\tan (f x+e)^2\right)}\right)}{\dots}$

input `int((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x,method=_RETURNVERBOSE)`


```
output -1/120/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*c^3*a*(60*
I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^3+24*B*(a*c)^(1/2)
*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^4+80*I*A*(a*c)^(1/2)*(a*c*(1+tan(
f*x+e)^2))^(1/2)*tan(f*x+e)^2+30*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)
)*tan(f*x+e)^3-30*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)
)^(1/2))/(a*c)^(1/2))*a*c+30*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*
tan(f*x+e)-32*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+80*I
*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)-75*A*ln((a*c*tan(f*x+e)+(a*c)^(
1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c-45*A*(a*c)^(1/2)*(a*c
*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-56*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)
)^(1/2))/(a*c)^(1/2)/(a*c*(1+tan(f*x+e)^2))^(1/2)
```

3.795.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 682 vs. 2(213) = 426.

Time = 0.28 (sec) , antiderivative size = 682, normalized size of antiderivative = 2.44

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx = \frac{15 \sqrt{\frac{(25 A^2 + 20 i AB - 4 B^2) a^3 c^7}{f^2}} (f e^{(8i fx + 8i e)} + 4 f e^{(6i fx + 6i e)} + 6 f e^{(4i fx + 4i e)} + 4 f e^{(2i fx + 2i e)})}{\dots}$$

```
input integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/
2),x, algorithm="fricas")
```

output $1/240*(15*\sqrt{(25*A^2 + 20*I*A*B - 4*B^2)*a^3*c^7/f^2}*(f*e^{(8*I*f*x + 8*I*e)} + 4*f*e^{(6*I*f*x + 6*I*e)} + 6*f*e^{(4*I*f*x + 4*I*e)} + 4*f*e^{(2*I*f*x + 2*I*e)} + f)*\log(-4*(2*((5*I*A - 2*B)*a*c^3*e^{(3*I*f*x + 3*I*e)} + (5*I*A - 2*B)*a*c^3*e^{(I*f*x + I*e)}))*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} + \sqrt{(25*A^2 + 20*I*A*B - 4*B^2)*a^3*c^7/f^2}*(f*e^{(2*I*f*x + 2*I*e)} - f))/((-5*I*A + 2*B)*a*c^3*e^{(2*I*f*x + 2*I*e)} + (-5*I*A + 2*B)*a*c^3) - 15*\sqrt{(25*A^2 + 20*I*A*B - 4*B^2)*a^3*c^7/f^2}*(f*e^{(8*I*f*x + 8*I*e)} + 4*f*e^{(6*I*f*x + 6*I*e)} + 6*f*e^{(4*I*f*x + 4*I*e)} + 4*f*e^{(2*I*f*x + 2*I*e)} + f)*\log(-4*(2*((5*I*A - 2*B)*a*c^3*e^{(3*I*f*x + 3*I*e)} + (5*I*A - 2*B)*a*c^3*e^{(I*f*x + I*e)}))*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} - \sqrt{(25*A^2 + 20*I*A*B - 4*B^2)*a^3*c^7/f^2}*(f*e^{(2*I*f*x + 2*I*e)} - f))/((-5*I*A + 2*B)*a*c^3*e^{(2*I*f*x + 2*I*e)} + (-5*I*A + 2*B)*a*c^3) - 4*(15*(5*I*A - 2*B)*a*c^3*e^{(9*I*f*x + 9*I*e)} + 70*(5*I*A - 2*B)*a*c^3*e^{(7*I*f*x + 7*I*e)} + 128*(5*I*A - 2*B)*a*c^3*e^{(5*I*f*x + 5*I*e)} + 10*(29*I*A - 50*B)*a*c^3*e^{(3*I*f*x + 3*I*e)} + 15*(-5*I*A + 2*B)*a*c^3*e^{(I*f*x + I*e)})*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)))/(f*e^{(8*I*f*x + 8*I*e)} + 4*f*e^{(6*I*f*x + 6*I*e)} + 6*f*e^{(4*I*f*x + 4*I*e)} + 4*f*e^{(2*I*f*x + 2*I*e)} + f)$

3.795.6 Sympy [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(7/2),x)`

output Timed out

3.795.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1655 vs. $2(213) = 426$.

Time = 2.24 (sec) , antiderivative size = 1655, normalized size of antiderivative = 5.93

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \text{Too large to display}$$

3.795. $\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="maxima")`

output `-480*(60*(5*A + 2*I*B)*a*c^3*cos(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 280*(5*A + 2*I*B)*a*c^3*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 512*(5*A + 2*I*B)*a*c^3*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 40*(29*A + 50*I*B)*a*c^3*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 60*(5*A + 2*I*B)*a*c^3*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 60*(5*I*A - 2*B)*a*c^3*sin(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 280*(5*I*A - 2*B)*a*c^3*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 512*(5*I*A - 2*B)*a*c^3*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 40*(29*I*A - 50*B)*a*c^3*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 60*(-5*I*A + 2*B)*a*c^3*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 30*((5*A + 2*I*B)*a*c^3*cos(10*f*x + 10*e) + 5*(5*A + 2*I*B)*a*c^3*cos(8*f*x + 8*e) + 10*(5*A + 2*I*B)*a*c^3*cos(6*f*x + 6*e) + 10*(5*A + 2*I*B)*a*c^3*cos(4*f*x + 4*e) + 5*(5*A + 2*I*B)*a*c^3*cos(2*f*x + 2*e) + (5*I*A - 2*B)*a*c^3*sin(10*f*x + 10*e) + 5*(5*I*A - 2*B)*a*c^3*sin(8*f*x + 8*e) + 10*(5*I*A - 2*B)*a*c^3*sin(6*f*x + 6*e) + 10*(5*I*A - 2*B)*a*c^3*sin(4*f*x + 4*e) + 5*(5*I*A - 2*B)*a*c^3*sin(2*f*x + 2*e) + (5*A + 2*I*B)*a*c^3*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) + 30*((5*A + 2*I*B)*a*c^3*cos(10*f*x + 10*e) + 5*(5*A + 2*I*B)*a*c^3*cos(8*f*x + 8*e) + 10*(5*A + 2*I*B)*a*c^3*cos(6*f*x + ...`

3.795.8 Giac [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="giac")`

output `Timed out`

3.795.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{7/2} dx = \int (A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{3/2} (c - c \tan(e + fx) li)^{7/2} dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(3/2)*(c - c*tan(e + f*x)*1i)^(7/2), x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(3/2)*(c - c*tan(e + f*x)*1i)^(7/2), x)`

3.796 $\int (a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{5/2} dx$

3.796.1 Optimal result	7190
3.796.2 Mathematica [A] (verified)	7191
3.796.3 Rubi [A] (verified)	7191
3.796.4 Maple [A] (verified)	7194
3.796.5 Fricas [B] (verification not implemented)	7195
3.796.6 Sympy [F(-1)]	7196
3.796.7 Maxima [B] (verification not implemented)	7196
3.796.8 Giac [F]	7197
3.796.9 Mupad [F(-1)]	7198

3.796.1 Optimal result

Integrand size = 45, antiderivative size = 226

$$\int (a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{5/2} dx =$$

$$\frac{a^{3/2}(4iA - B)c^{5/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{4f}$$

$$+ \frac{a(4A + iB)c^2 \tan(e+fx) \sqrt{a+ia \tan(e+fx)} \sqrt{c-ictan(e+fx)}}{8f}$$

$$- \frac{(4iA - B)c(a+ia \tan(e+fx))^{3/2}(c-ictan(e+fx))^{3/2}}{12f}$$

$$+ \frac{B(a+ia \tan(e+fx))^{3/2}(c-ictan(e+fx))^{5/2}}{4f}$$

output

```
-1/4*a^(3/2)*(4*I*A-B)*c^(5/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/f+1/8*a*(4*A+I*B)*c^2*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)*tan(f*x+e)/f-1/12*(4*I*A-B)*c*(a+I*a*tan(f*x+e))^(3/2)*(c-I*c*tan(f*x+e))^(3/2)/f+1/4*B*(a+I*a*tan(f*x+e))^(3/2)*(c-I*c*tan(f*x+e))^(5/2)/f
```

3.796.2 Mathematica [A] (verified)

Time = 7.77 (sec) , antiderivative size = 206, normalized size of antiderivative = 0.91

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx = \frac{a^{3/2} c^3 \sqrt{1 - i \tan(e + fx)} \left(6(-4iA + B) \arcsin \left(\frac{\sqrt{a + ia \tan(e + fx)}}{\sqrt{2}\sqrt{a}} \right) \sqrt{a + ia \tan(e + fx)} \right)}{}$$

input `Integrate[(a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2),x]`

output `(a^(3/2)*c^3*Sqrt[1 - I*Tan[e + f*x]]*(6*((-4*I)*A + B)*ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a])]*Sqrt[a + I*a*Tan[e + f*x]] + Sqrt[a]*Sqrt[1 - I*Tan[e + f*x]]*(-I + Tan[e + f*x])*(8*(A + I*B) + 3*((4*I)*A + B)*Tan[e + f*x] + 8*(A + I*B)*Tan[e + f*x]^2 + 6*B*Tan[e + f*x]^3))/(24*f*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])`

3.796.3 Rubi [A] (verified)Time = 0.41 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.96, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 90, 59, 40, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{5/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{5/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \sqrt{i \tan(e + fx)a + a} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{90} \end{aligned}$$

$$\frac{ac\left(\frac{1}{4}(4A + iB) \int \sqrt{i \tan(e + fx)a + a} (c - ic \tan(e + fx))^{3/2} d \tan(e + fx) + \frac{B(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{5/2}}{4ac}\right)}{f}$$

↓ 59

$$\frac{ac\left(\frac{1}{4}(4A + iB) \left(c \int \sqrt{i \tan(e + fx)a + a} \sqrt{c - ic \tan(e + fx)} d \tan(e + fx) - \frac{i(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2}}{3a}\right)\right)}{f}$$

↓ 40

$$\frac{ac\left(\frac{1}{4}(4A + iB) \left(c \left(\frac{1}{2} ac \int \frac{1}{\sqrt{i \tan(e + fx)a + a} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}\right)\right)\right)}{f}$$

↓ 45

$$\frac{ac\left(\frac{1}{4}(4A + iB) \left(c \left(ac \int \frac{1}{ia + \frac{ic(i \tan(e + fx)a + a)}{c - ic \tan(e + fx)}} d \frac{\sqrt{i \tan(e + fx)a + a}}{\sqrt{c - ic \tan(e + fx)}} + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}\right)\right)\right)}{f}$$

↓ 218

$$\frac{ac\left(\frac{1}{4}(4A + iB) \left(c \left(\frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)} - i \sqrt{a} \sqrt{c} \arctan\left(\frac{\sqrt{c} \sqrt{a + ia \tan(e + fx)}}{\sqrt{a} \sqrt{c - ic \tan(e + fx)}}\right)\right)\right)\right)}{f}$$

input `Int[(a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2),x]`

output `(a*c*((B*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(5/2))/(4*a*c) + ((4*A + I*B)*(((-1/3*I)*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(3/2))/a + c*((-I)*Sqrt[a]*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])]) + (Tan[e + f*x]*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/2))/4)/f`

3.796.3.1 Defintions of rubi rules used

- rule 40 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^m/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`
- rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 59 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[2*c*(n/(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0] && IGtQ[n + 1/2, 0] && LtQ[m, n]`
- rule 90 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.796.4 Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 350, normalized size of antiderivative = 1.55

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} c^2 a \left(6iB\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \tan(fx+e)^3 + 8iA\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} c^2 a \left(6iB\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \tan(fx+e)^3 + 8iA\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \right)}{\dots}$
parts	$\frac{A\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} c^2 a \left(2i \tan(fx+e)^2 \sqrt{ac(1+\tan(fx+e)^2)} \sqrt{ac} + 2i\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \right)}{6f\sqrt{ac(1+\tan(fx+e)^2)}}$

input `int((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x,m
method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/24/f*(a*(1+I*\tan(f*x+e)))^{(1/2)}*(-c*(I*\tan(f*x+e)-1))^{(1/2)}*c^2*a*(6*I* \\ & B*(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)}*\tan(f*x+e)^3+8*I*A*(a*c)^{(1/2)}* \\ & (a*c*(1+\tan(f*x+e)^2))^{(1/2)}*\tan(f*x+e)^2-3*I*B*\ln((a*c*\tan(f*x+e)+(a*c)^{(1/2)}* \\ & (a*c*(1+\tan(f*x+e)^2))^{(1/2)})/(a*c)^{(1/2)})*a*c+3*I*B*(a*c)^{(1/2)}*(a*c \\ & *(1+\tan(f*x+e)^2))^{(1/2)}*\tan(f*x+e)-8*B*(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2)) \\ & ^{(1/2)}*\tan(f*x+e)^2+8*I*A*(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)}-12*A*\ln \\ & ((a*c*\tan(f*x+e)+(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)})/(a*c)^{(1/2)})*a* \\ & c-12*A*(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)}*\tan(f*x+e)-8*B*(a*c)^{(1/2)} \\ & *(a*c*(1+\tan(f*x+e)^2))^{(1/2)})/(a*c)^{(1/2)}/(a*c*(1+\tan(f*x+e)^2))^{(1/2)} \end{aligned}$$

3.796.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 614 vs. $2(170) = 340$.

Time = 0.28 (sec) , antiderivative size = 614, normalized size of antiderivative = 2.72

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx = \frac{3 \sqrt{\frac{(16A^2 + 8iAB - B^2)a^3c^5}{f^2}} (fe^{(6ifx+6ie)} + 3fe^{(4ifx+4ie)} + 3fe^{(2ifx+2ie)} + f) \log \left(-\frac{4}{\dots} \right)}{\dots}$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="fricas")`

output `1/48*(3*sqrt((16*A^2 + 8*I*A*B - B^2)*a^3*c^5/f^2)*(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)*log(-4*(2*((4*I*A - B)*a*c^2*e^(3*I*f*x + 3*I*e) + (4*I*A - B)*a*c^2*e^(I*f*x + I*e)))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + sqrt((16*A^2 + 8*I*A*B - B^2)*a^3*c^5/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((-4*I*A + B)*a*c^2*e^(2*I*f*x + 2*I*e) + (-4*I*A + B)*a*c^2) - 3*sqrt((16*A^2 + 8*I*A*B - B^2)*a^3*c^5/f^2)*(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)*log(-4*(2*((4*I*A - B)*a*c^2*e^(3*I*f*x + 3*I*e) + (4*I*A - B)*a*c^2*e^(I*f*x + I*e)))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - sqrt((16*A^2 + 8*I*A*B - B^2)*a^3*c^5/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((-4*I*A + B)*a*c^2*e^(2*I*f*x + 2*I*e) + (-4*I*A + B)*a*c^2) - 4*(3*(4*I*A - B)*a*c^2*e^(7*I*f*x + 7*I*e) + 11*(4*I*A - B)*a*c^2*e^(5*I*f*x + 5*I*e) - (-20*I*A + 53*B)*a*c^2*e^(3*I*f*x + 3*I*e) + 3*(-4*I*A + B)*a*c^2*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)`

3.796.6 Sympy [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(5/2),x)`

output `Timed out`

3.796.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1375 vs. $2(170) = 340$.

Time = 1.13 (sec) , antiderivative size = 1375, normalized size of antiderivative = 6.08

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx = \text{Too large to display}$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output

```
-96*(12*(4*A + I*B)*a*c^2*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 44*(4*A + I*B)*a*c^2*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 4*(20*A + 53*I*B)*a*c^2*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 12*(4*A + I*B)*a*c^2*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 12*(4*I*A - B)*a*c^2*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 44*(4*I*A - B)*a*c^2*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 4*(20*I*A - 53*B)*a*c^2*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 12*(-4*I*A + B)*a*c^2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 6*((4*A + I*B)*a*c^2*cos(8*f*x + 8*e) + 4*(4*A + I*B)*a*c^2*cos(6*f*x + 6*e) + 6*(4*A + I*B)*a*c^2*cos(4*f*x + 4*e) + 4*(4*A + I*B)*a*c^2*cos(2*f*x + 2*e) + (4*I*A - B)*a*c^2*sin(8*f*x + 8*e) + 4*(4*I*A - B)*a*c^2*sin(6*f*x + 6*e) + 6*(4*I*A - B)*a*c^2*sin(4*f*x + 4*e) + 4*(4*I*A - B)*a*c^2*sin(2*f*x + 2*e) + (4*A + I*B)*a*c^2)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) + 6*((4*A + I*B)*a*c^2*cos(8*f*x + 8*e) + 4*(4*A + I*B)*a*c^2*cos(6*f*x + 6*e) + 6*(4*A + I*B)*a*c^2*cos(4*f*x + 4*e) + 4*(4*A + I*B)*a*c^2*cos(2*f*x + 2*e) + (4*I*A - B)*a*c^2*sin(8*f*x + 8*e) + 4*(4*I*A - B)*a*c^2*sin(6*f*x + 6*e) + 6*(4*I*A - B)*a*c^2*sin(4*f*x + 4*e) + 4*(4*I*A - B)*a*c^2*sin(2*f*x + 2*e) + (4*A + I*B)*a*c^2)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), -sin(1/2*arct...
```

3.796.8 Giac [F]

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{5/2} dx = \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^{3/2} (-ictan(fx + e) + c)^{5/2} dx$$

input

```
integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="giac")
```

output

```
sage0*x
```

3.796.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{5/2} dx = \int (A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{3/2} (c - c \tan(e + fx) li)^{5/2} dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(3/2)*(c - c*tan(e + f*x)*1i)^(5/2), x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(3/2)*(c - c*tan(e + f*x)*1i)^(5/2), x)`

3.797 $\int (a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{3/2} dx$

3.797.1 Optimal result	7199
3.797.2 Mathematica [A] (verified)	7200
3.797.3 Rubi [A] (verified)	7200
3.797.4 Maple [A] (verified)	7202
3.797.5 Fricas [B] (verification not implemented)	7203
3.797.6 Sympy [F]	7204
3.797.7 Maxima [B] (verification not implemented)	7204
3.797.8 Giac [F]	7205
3.797.9 Mupad [F(-1)]	7206

3.797.1 Optimal result

Integrand size = 45, antiderivative size = 157

$$\int (a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{3/2} dx =$$

$$-\frac{ia^{3/2}Ac^{3/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{f}$$

$$+\frac{aAc \tan(e+fx)\sqrt{a+ia \tan(e+fx)}\sqrt{c-ictan(e+fx)}}{2f}$$

$$+\frac{B(a+ia \tan(e+fx))^{3/2}(c-ictan(e+fx))^{3/2}}{3f}$$

output

```
-I*a^(3/2)*A*c^(3/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/f+1/2*a*A*c*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)*tan(f*x+e)/f+1/3*B*(a+I*a*tan(f*x+e))^(3/2)*(c-I*c*tan(f*x+e))^(3/2)/f
```

3.797.2 Mathematica [A] (verified)

Time = 5.46 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.15

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} dx = \frac{a^{3/2} c^2 (i + \tan(e + fx)) \left(-6A \arcsin \left(\frac{\sqrt{a + ia \tan(e + fx)}}{\sqrt{2}\sqrt{a}} \right) \sqrt{a + ia \tan(e + fx)} + \sqrt{a} \right)}{6f \sqrt{1 - i \tan(e + fx)} \sqrt{a + ia \tan(e + fx)}}$$

input `Integrate[(a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(a^(3/2)*c^2*(I + Tan[e + f*x])*(-6*A*ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a])]*Sqrt[a + I*a*Tan[e + f*x]] + Sqrt[a]*Sqrt[1 - I*Tan[e + f*x]])*(-I + Tan[e + f*x])*(2*B + 3*A*Tan[e + f*x] + 2*B*Tan[e + f*x]^2))/(6*f*Sqrt[1 - I*Tan[e + f*x]]*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])`

3.797.3 Rubi [A] (verified)Time = 0.40 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.01, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 90, 40, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \sqrt{i \tan(e + fx)a + a} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{90} \end{aligned}$$

$$\frac{ac \left(A \int \sqrt{i \tan(e+fx)a+a} \sqrt{c-ictan(e+fx)} d \tan(e+fx) + \frac{B(a+ia \tan(e+fx))^{3/2} (c-ictan(e+fx))^{3/2}}{3ac} \right)}{f}$$

↓ 40

$$\frac{ac \left(A \left(\frac{1}{2} ac \int \frac{1}{\sqrt{i \tan(e+fx)a+a} \sqrt{c-ictan(e+fx)}} d \tan(e+fx) + \frac{1}{2} \tan(e+fx) \sqrt{a+ia \tan(e+fx)} \sqrt{c-ictan(e+fx)} \right) \right)}{f}$$

↓ 45

$$\frac{ac \left(A \left(ac \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ictan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ictan(e+fx)}} + \frac{1}{2} \tan(e+fx) \sqrt{a+ia \tan(e+fx)} \sqrt{c-ictan(e+fx)} \right) \right) + \frac{B(a+ia \tan(e+fx))^{3/2} (c-ictan(e+fx))^{3/2}}{3ac}}{f}$$

↓ 218

$$\frac{ac \left(A \left(\frac{1}{2} \tan(e+fx) \sqrt{a+ia \tan(e+fx)} \sqrt{c-ictan(e+fx)} - i \sqrt{a} \sqrt{c} \arctan \left(\frac{\sqrt{c} \sqrt{a+ia \tan(e+fx)}}{\sqrt{a} \sqrt{c-ictan(e+fx)}} \right) \right) \right) + \frac{B(a+ia \tan(e+fx))^{3/2} (c-ictan(e+fx))^{3/2}}{3ac}}{f}$$

input `Int[(a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(a*c*((B*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(3/2))/(3*a*c) + A*((-I)*Sqrt[a]*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])] + (Tan[e + f*x]*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/2)))/f`

3.797.3.1 Defintions of rubi rules used

rule 40 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^m/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`

rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`


```
rule 90 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]

rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.797.4 Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.14

method	result
parts	$\frac{A\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}ac\left(\tan(fx+e)\sqrt{ac(1+\tan(fx+e)^2)}\sqrt{ac}+ac\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{a}}{\sqrt{ac}}\right)\right)}{2f\sqrt{ac(1+\tan(fx+e)^2)}\sqrt{ac}}$
derivativedivides	$\frac{\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}ac\left(2B\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}\tan(fx+e)^2+3A\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{a}}{\sqrt{ac}}\right)\right)}{6f\sqrt{ac(1+\tan(fx+e)^2)}}$
default	$\frac{\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}ac\left(2B\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}\tan(fx+e)^2+3A\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{a}}{\sqrt{ac}}\right)\right)}{6f\sqrt{ac(1+\tan(fx+e)^2)}}$

```
input int((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x,m
method=_RETURNVERBOSE)
```

3.797. $\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx$

output
$$\begin{aligned} & -1/2*A/f*(a*(1+I*\tan(f*x+e)))^{(1/2)}*(-c*(I*\tan(f*x+e)-1))^{(1/2)}*a*c*(\tan(f \\ & *x+e)*(a*c*(1+\tan(f*x+e)^2))^{(1/2)}*(a*c)^{(1/2)}+a*c*\ln((a*c*\tan(f*x+e)+(a*c \\ &)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)))/(a*c)^{(1/2)))/(a*c*(1+\tan(f*x+e)^2)) \\ & ^{(1/2)}/(a*c)^{(1/2)}-1/3*B/f*(a*(1+I*\tan(f*x+e)))^{(1/2)}*(-c*(I*\tan(f*x+e)-1) \\ &)^{(1/2)}*a*c*(1+\tan(f*x+e)^2) \end{aligned}$$

3.797.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 433 vs. $2(119) = 238$.

Time = 0.28 (sec) , antiderivative size = 433, normalized size of antiderivative = 2.76

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx = \frac{3 \sqrt{\frac{A^2 a^3 c^3}{f^2}} (f e^{4i fx + 4i e} + 2 f e^{2i fx + 2i e} + f) \log \left(\frac{4 \left(2 (A a c e^{3i fx + 3i e} + A a c e^{i fx + i e}) \sqrt{e^{2i e}} \right)}{A} \right) - ictan(e + fx))^{3/2} dx =$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output
$$\begin{aligned} & 1/12*(3*\sqrt{A^2*a^3*c^3/f^2}*(f*e^{(4*I*f*x + 4*I*e)} + 2*f*e^{(2*I*f*x + 2*I*e)} + f)*\log(4*(2*(A*a*c*e^{(3*I*f*x + 3*I*e)} + A*a*c*e^{(I*f*x + I*e)}))*\sqrt{ \\ & a/(e^{(2*I*f*x + 2*I*e)} + 1))*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} - \sqrt{A^2*a^3*c^3/f^2}*(I*f*e^{(2*I*f*x + 2*I*e)} - I*f))/(A*a*c*e^{(2*I*f*x + 2*I*e)} \\ & + A*a*c) - 3*\sqrt{A^2*a^3*c^3/f^2}*(f*e^{(4*I*f*x + 4*I*e)} + 2*f*e^{(2*I*f*x + 2*I*e)} + f)*\log(4*(2*(A*a*c*e^{(3*I*f*x + 3*I*e)} + A*a*c*e^{(I*f*x + I*e)}))*\sqrt{ \\ & a/(e^{(2*I*f*x + 2*I*e)} + 1))*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} - \sqrt{A^2*a^3*c^3/f^2}*(-I*f*e^{(2*I*f*x + 2*I*e)} + I*f))/(A*a*c*e^{(2*I*f*x + 2*I*e)} + A*a*c) - 4*(3*I*A*a*c*e^{(5*I*f*x + 5*I*e)} - 8*B*a*c*e^{(3*I*f*x + 3*I*e)} - 3*I*A*a*c*e^{(I*f*x + I*e)})*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1))*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1))}/(f*e^{(4*I*f*x + 4*I*e)} + 2*f*e^{(2*I*f*x + 2*I*e)} + f) \end{aligned}$$

3.797.6 Sympy [F]

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} dx = \int (ia(\tan(e + fx) - i))^{3/2} (-ic(\tan(e + fx) + i))^{3/2} (A + B \tan(e + fx)) dx$$

input `integrate((a+I*a*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(3/2),x)`

output `Integral((I*a*(tan(e + f*x) - I))**(3/2)*(-I*c*(tan(e + f*x) + I))**(3/2)*(A + B*tan(e + f*x)), x)`

3.797.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 853 vs. 2(119) = 238.

Time = 0.45 (sec) , antiderivative size = 853, normalized size of antiderivative = 5.43

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} dx = \text{Too large to display}$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output

```

-(12*A*a*c*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 32*I*B*a
*c*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 12*A*a*c*cos(1/2
*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 12*I*A*a*c*sin(5/2*arctan2
(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 32*B*a*c*sin(3/2*arctan2(sin(2*f*x
+ 2*e), cos(2*f*x + 2*e))) - 12*I*A*a*c*sin(1/2*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e))) + 6*(A*a*c*cos(6*f*x + 6*e) + 3*A*a*c*cos(4*f*x + 4*e)
+ 3*A*a*c*cos(2*f*x + 2*e) + I*A*a*c*sin(6*f*x + 6*e) + 3*I*A*a*c*sin(4*f
*x + 4*e) + 3*I*A*a*c*sin(2*f*x + 2*e) + A*a*c)*arctan2(cos(1/2*arctan2(si
n(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(
2*f*x + 2*e))) + 1) + 6*(A*a*c*cos(6*f*x + 6*e) + 3*A*a*c*cos(4*f*x + 4*e)
+ 3*A*a*c*cos(2*f*x + 2*e) + I*A*a*c*sin(6*f*x + 6*e) + 3*I*A*a*c*sin(4*f
*x + 4*e) + 3*I*A*a*c*sin(2*f*x + 2*e) + A*a*c)*arctan2(cos(1/2*arctan2(si
n(2*f*x + 2*e), cos(2*f*x + 2*e))), -sin(1/2*arctan2(sin(2*f*x + 2*e), cos
(2*f*x + 2*e))) + 1) + 3*(I*A*a*c*cos(6*f*x + 6*e) + 3*I*A*a*c*cos(4*f*x +
4*e) + 3*I*A*a*c*cos(2*f*x + 2*e) - A*a*c*sin(6*f*x + 6*e) - 3*A*a*c*sin(
4*f*x + 4*e) - 3*A*a*c*sin(2*f*x + 2*e) + I*A*a*c)*log(cos(1/2*arctan2(sin
(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + sin(1/2*arctan2(sin(2*f*x + 2*e), co
s(2*f*x + 2*e)))^2 + 2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))
) + 1) + 3*(-I*A*a*c*cos(6*f*x + 6*e) - 3*I*A*a*c*cos(4*f*x + 4*e) - 3*I*A
*a*c*cos(2*f*x + 2*e) + A*a*c*sin(6*f*x + 6*e) + 3*A*a*c*sin(4*f*x + 4*...

```

3.797.8 Giac [F]

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx = \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^{3/2} (-ictan(fx + e) + c)^{3/2} dx$$

input

```

integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/
2),x, algorithm="giac")

```

output

```

sage0*x

```

3.797.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx = \int (A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{3/2} (c - c \tan(e + fx) li)^{3/2} dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(3/2)*(c - c*tan(e + f*x)*1i)^(3/2),x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(3/2)*(c - c*tan(e + f*x)*1i)^(3/2), x)`

3.798 $\int (a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))\sqrt{c-ictan(e+fx)}$

3.798.1 Optimal result	7207
3.798.2 Mathematica [A] (verified)	7208
3.798.3 Rubi [A] (verified)	7208
3.798.4 Maple [A] (verified)	7210
3.798.5 Fricas [B] (verification not implemented)	7211
3.798.6 Sympy [F]	7212
3.798.7 Maxima [B] (verification not implemented)	7212
3.798.8 Giac [B] (verification not implemented)	7213
3.798.9 Mupad [F(-1)]	7214

3.798.1 Optimal result

Integrand size = 45, antiderivative size = 160

$$\int (a + ia \tan(e + fx))^{3/2}(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)} dx =$$

$$-\frac{a^{3/2}(2iA + B)\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{f}$$

$$+ \frac{a(2iA + B)\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}}{2f}$$

$$+ \frac{B(a + ia \tan(e + fx))^{3/2}\sqrt{c - ictan(e + fx)}}{2f}$$

```
output -a^(3/2)*(2*I*A+B)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*
tan(f*x+e))^(1/2))*c^(1/2)/f+1/2*a*(2*I*A+B)*(a+I*a*tan(f*x+e))^(1/2)*(c-I
*c*tan(f*x+e))^(1/2)/f+1/2*B*(c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(
3/2)/f
```

3.798.2 Mathematica [A] (verified)

Time = 4.37 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.14

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx = \frac{a^{3/2} c (i + \tan(e + fx)) \left(-2(2A - iB) \arcsin \left(\frac{\sqrt{a + ia \tan(e + fx)}}{\sqrt{2}\sqrt{a}} \right) \sqrt{c - ic \tan(e + fx)} \right)}{2f \sqrt{1 - i \tan(e + fx)}}$$

```
input Integrate[(a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]],x]
```

```
output (a^(3/2)*c*(I + Tan[e + f*x])*(-2*(2*A - I*B)*ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a])]*Sqrt[a + I*a*Tan[e + f*x]] + Sqrt[a]*Sqrt[1 - I*Tan[e + f*x]]*(1 + I*Tan[e + f*x])*(2*A - (2*I)*B + B*Tan[e + f*x]))/(2*f*Sqrt[1 - I*Tan[e + f*x]]*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])
```

3.798.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.04, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 90, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)} (A + B \tan(e + fx)) dx$$

↓ 3042

$$\int (a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)} (A + B \tan(e + fx)) dx$$

↓ 4071

$$\frac{ac \int \frac{\sqrt{i \tan(e + fx) a + a(A + B \tan(e + fx))}}{\sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{f}$$

↓ 90

$$\begin{aligned}
 & \frac{ac \left(\frac{1}{2} (2A - iB) \int \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{B(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2ac} \right)}{f} \\
 & \quad \downarrow 60 \\
 & \frac{ac \left(\frac{1}{2} (2A - iB) \left(a \int \frac{1}{\sqrt{i \tan(e+fx)a+a} \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{c} \right) + \frac{B(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2ac} \right)}{f} \\
 & \quad \downarrow 45 \\
 & \frac{ac \left(\frac{1}{2} (2A - iB) \left(2a \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} + \frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{c} \right) + \frac{B(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2ac} \right)}{f} \\
 & \quad \downarrow 218 \\
 & \frac{ac \left(\frac{1}{2} (2A - iB) \left(\frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{c} - \frac{2i \sqrt{a} \arctan \left(\frac{\sqrt{c} \sqrt{a+ia \tan(e+fx)}}{\sqrt{a} \sqrt{c-ic \tan(e+fx)}} \right)}{\sqrt{c}} \right) + \frac{B(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2ac} \right)}{f}
 \end{aligned}$$

input `Int[(a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]],x]`

output `(a*c*((B*(a + I*a*Tan[e + f*x])^(3/2)*Sqrt[c - I*c*Tan[e + f*x]])/(2*a*c) + ((2*A - I*B)*(((-2*I)*Sqrt[a]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])]/Sqrt[c] + (I*Sqrt[a + I*a*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])/c))/2))/f`

3.798.3.1 Defintions of rubi rules used

rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.798.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.39

method	result
derivativedivides	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} a \left(iB\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \tan(fx+e) - iB \ln \left(\frac{ac \tan(fx+e) + \sqrt{ac} \sqrt{ac}}{\sqrt{ac}} \right) \right)}{2f\sqrt{ac}}$
default	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} a \left(iB\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \tan(fx+e) - iB \ln \left(\frac{ac \tan(fx+e) + \sqrt{ac} \sqrt{ac}}{\sqrt{ac}} \right) \right)}{2f\sqrt{ac}}$
parts	$\frac{A\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} a \left(i\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} + ac \ln \left(\frac{ac \tan(fx+e) + \sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right)}{f\sqrt{ac(1+\tan(fx+e)^2)} \sqrt{ac}}$

```
input int((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)),x,m
method=_RETURNVERBOSE)
```

```
output 1/2/f*(-c*(I*tan(f*x+e)-1))^(1/2)*(a*(1+I*tan(f*x+e)))^(1/2)*a*(I*B*(a*c)^(
(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-I*B*ln((a*c*tan(f*x+e)+(a*c)
^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c+2*I*A*(a*c)^(1/2)*(a
*c*(1+tan(f*x+e)^2))^(1/2)+2*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(
f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c+2*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(
(1/2))/(a*c*(1+tan(f*x+e)^2))^(1/2)/(a*c)^(1/2)
```

3.798.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 459 vs. 2(120) = 240.

Time = 0.28 (sec) , antiderivative size = 459, normalized size of antiderivative = 2.87

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx =$$

$$\sqrt{\frac{4A^2 - 4iAB - B^2}{f^2} a^3 c} (f e^{(2i fx + 2i e)} + f) \log \left(- \frac{4 \left(2((-2iA - B)ae^{(3i fx + 3i e)} + (-2iA - B)ae^{(i fx + i e)}) \sqrt{\frac{a}{e^{(2i fx + 2i e)} + 1}} \sqrt{e^{(2i fx + 2i e)}} \right)}{(2iA + B)ae^{(2i fx + 2i e)} + (2iA + B)} \right)$$

```
input integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e
)),x, algorithm="fracas")
```

3.798. $\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx$

output `-1/4*(sqrt((4*A^2 - 4*I*A*B - B^2)*a^3*c/f^2)*(f*e^(2*I*f*x + 2*I*e) + f)*
log(-4*(2*(-2*I*A - B)*a*e^(3*I*f*x + 3*I*e) + (-2*I*A - B)*a*e^(I*f*x +
I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))
+ sqrt((4*A^2 - 4*I*A*B - B^2)*a^3*c/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((2
*I*A + B)*a*e^(2*I*f*x + 2*I*e) + (2*I*A + B)*a) - sqrt((4*A^2 - 4*I*A*B
- B^2)*a^3*c/f^2)*(f*e^(2*I*f*x + 2*I*e) + f)*log(-4*(2*(-2*I*A - B)*a*e^(
3*I*f*x + 3*I*e) + (-2*I*A - B)*a*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2
*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - sqrt((4*A^2 - 4*I*A*B - B^
2)*a^3*c/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((2*I*A + B)*a*e^(2*I*f*x + 2*I
*e) + (2*I*A + B)*a) + 4*((-2*I*A - 3*B)*a*e^(3*I*f*x + 3*I*e) + (-2*I*A
- B)*a*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f
*x + 2*I*e) + 1)))/(f*e^(2*I*f*x + 2*I*e) + f)`

3.798.6 Sympy [F]

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx = \int (ia(\tan(e + fx) - i))^{3/2} \sqrt{-ic(\tan(e + fx) + i)} (A + B \tan(e + fx)) dx$$

input `integrate((c-I*c*tan(f*x+e))**(1/2)*(a+I*a*tan(f*x+e))**(3/2)*(A+B*tan(f*x
+e)),x)`

output `Integral((I*a*(tan(e + f*x) - I))**(3/2)*sqrt(-I*c*(tan(e + f*x) + I))*(A
+ B*tan(e + f*x)), x)`

3.798.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 771 vs. $2(120) = 240$.

Time = 0.47 (sec) , antiderivative size = 771, normalized size of antiderivative = 4.82

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx = \text{Too large to display}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)),x, algorithm="maxima")`

output `4*(4*(2*A - 3*I*B)*a*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 4*(2*A - I*B)*a*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 4*(2*I*A + 3*B)*a*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 4*(2*I*A + B)*a*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) - 2*((2*A - I*B)*a*cos(4*f*x + 4*e) + 2*(2*A - I*B)*a*cos(2*f*x + 2*e) - (-2*I*A - B)*a*sin(4*f*x + 4*e) - 2*(-2*I*A - B)*a*sin(2*f*x + 2*e) + (2*A - I*B)*a*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) - 2*((2*A - I*B)*a*cos(4*f*x + 4*e) + 2*(2*A - I*B)*a*cos(2*f*x + 2*e) - (-2*I*A - B)*a*sin(4*f*x + 4*e) - 2*(-2*I*A - B)*a*sin(2*f*x + 2*e) + (2*A - I*B)*a*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), -sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) + ((-2*I*A - B)*a*cos(4*f*x + 4*e) + 2*(-2*I*A - B)*a*cos(2*f*x + 2*e) + (2*A - I*B)*a*sin(4*f*x + 4*e) + 2*(2*A - I*B)*a*sin(2*f*x + 2*e) + (-2*I*A - B)*a*log(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))^2 + sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))^2 + 2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) + ((2*I*A + B)*a*cos(4*f*x + 4*e) + 2*(2*I*A + B)*a*cos(2*f*x + 2*e) - (2*A - I*B)*a*sin(4*f*x + 4*e) - 2*(2*A - I*B)*a*sin(2*f*x + 2*e) + (2*I*A + B)*a*log(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))^2 + sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))^2 - 2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))`

3.798.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 533 vs. $2(120) = 240$.

Time = 1.53 (sec) , antiderivative size = 533, normalized size of antiderivative = 3.33

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx = \frac{-3i Ba^{\frac{3}{2}} \sqrt{c} e^{(8i fx + 8i e)} \log(e^{(i fx + i e)} + i) - 12i Ba^{\frac{3}{2}} \sqrt{c} e^{(6i fx + 6i e)}}{4f} - \frac{i \left(\left(8Aa^{\frac{3}{2}} \sqrt{c} - i Ba^{\frac{3}{2}} \sqrt{c} \right) \arctan(e^{(i fx + i e)}) - \frac{8Aa^{\frac{3}{2}} \sqrt{c} e^{(3i fx + 3i e)} - 7i Ba^{\frac{3}{2}} \sqrt{c} e^{(3i fx + 3i e)} + 8Aa^{\frac{3}{2}} \sqrt{c} e^{(i fx + i e)} - i Ba^{\frac{3}{2}} \sqrt{c} e^{(i fx + i e)}}{(e^{(2i fx + 2i e)} + 1)^2}}{4f}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e)),x, algorithm="giac")`

3.798. $\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx$

output

```

1/8*(-3*I*B*a^(3/2)*sqrt(c)*e^(8*I*f*x + 8*I*e)*log(e^(I*f*x + I*e) + I) -
12*I*B*a^(3/2)*sqrt(c)*e^(6*I*f*x + 6*I*e)*log(e^(I*f*x + I*e) + I) - 18*
I*B*a^(3/2)*sqrt(c)*e^(4*I*f*x + 4*I*e)*log(e^(I*f*x + I*e) + I) - 12*I*B*
a^(3/2)*sqrt(c)*e^(2*I*f*x + 2*I*e)*log(e^(I*f*x + I*e) + I) + 3*I*B*a^(3/
2)*sqrt(c)*e^(8*I*f*x + 8*I*e)*log(e^(I*f*x + I*e) - I) + 12*I*B*a^(3/2)*s
qrt(c)*e^(6*I*f*x + 6*I*e)*log(e^(I*f*x + I*e) - I) + 18*I*B*a^(3/2)*sqrt(
c)*e^(4*I*f*x + 4*I*e)*log(e^(I*f*x + I*e) - I) + 12*I*B*a^(3/2)*sqrt(c)*e
^(2*I*f*x + 2*I*e)*log(e^(I*f*x + I*e) - I) + 10*B*a^(3/2)*sqrt(c)*e^(7*I*
f*x + 7*I*e) + 26*B*a^(3/2)*sqrt(c)*e^(5*I*f*x + 5*I*e) + 22*B*a^(3/2)*sqr
t(c)*e^(3*I*f*x + 3*I*e) + 6*B*a^(3/2)*sqrt(c)*e^(I*f*x + I*e) - 3*I*B*a^(
3/2)*sqrt(c)*log(e^(I*f*x + I*e) + I) + 3*I*B*a^(3/2)*sqrt(c)*log(e^(I*f*x
+ I*e) - I))/(f*e^(8*I*f*x + 8*I*e) + 4*f*e^(6*I*f*x + 6*I*e) + 6*f*e^(4*
I*f*x + 4*I*e) + 4*f*e^(2*I*f*x + 2*I*e) + f) - 1/4*I*((8*A*a^(3/2)*sqrt(c
) - I*B*a^(3/2)*sqrt(c))*arctan(e^(I*f*x + I*e)) - (8*A*a^(3/2)*sqrt(c)*e^
(3*I*f*x + 3*I*e) - 7*I*B*a^(3/2)*sqrt(c)*e^(3*I*f*x + 3*I*e) + 8*A*a^(3/2
)*sqrt(c)*e^(I*f*x + I*e) - I*B*a^(3/2)*sqrt(c)*e^(I*f*x + I*e))/(e^(2*I*f
*x + 2*I*e) + 1)^2)/f

```

3.798.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx = \int (A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{3/2} \sqrt{c - ctan(e + fx) li} dx$$

input

```

int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*li)^(3/2)*(c - c*tan(e + f*x)
*li)^(1/2),x)

```

output

```

int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*li)^(3/2)*(c - c*tan(e + f*x)
*li)^(1/2), x)

```

$$3.799 \quad \int \frac{(a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} dx$$

3.799.1 Optimal result	7215
3.799.2 Mathematica [A] (verified)	7215
3.799.3 Rubi [A] (verified)	7216
3.799.4 Maple [B] (verified)	7218
3.799.5 Fricas [B] (verification not implemented)	7219
3.799.6 Sympy [F]	7220
3.799.7 Maxima [B] (verification not implemented)	7220
3.799.8 Giac [F]	7221
3.799.9 Mupad [F(-1)]	7222

3.799.1 Optimal result

Integrand size = 45, antiderivative size = 169

$$\int \frac{(a + ia \tan(e + fx))^{3/2}(A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx = \frac{2a^{3/2}(iA + 2B) \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{\sqrt{c}f} - \frac{(iA + B)(a + ia \tan(e + fx))^{3/2}}{f\sqrt{c - ictan(e + fx)}} - \frac{a(iA + 2B)\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}}{cf}$$

output `2*a^(3/2)*(I*A+2*B)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/f/c^(1/2)-a*(I*A+2*B)*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)/c/f-(I*A+B)*(a+I*a*tan(f*x+e))^(3/2)/f/(c-I*c*tan(f*x+e))^(1/2)`

3.799.2 Mathematica [A] (verified)

Time = 5.59 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.88

$$\int \frac{(a + ia \tan(e + fx))^{3/2}(A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx = \frac{2a^{3/2}(iA + 2B) \arcsin\left(\frac{\sqrt{a+ia \tan(e+fx)}}{\sqrt{2}\sqrt{a}}\right) \sqrt{1 - i \tan(e + fx)}}{f\sqrt{a +}}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x]))/Sqrt[c - I*c*Tan[e + f*x]],x]`

output $(2*a^{(3/2)}*(I*A + 2*B)*ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a])]*Sqrt[1 - I*Tan[e + f*x]]*Sqrt[a + I*a*Tan[e + f*x]] - a^2*(-I + Tan[e + f*x])*(-2*A + (3*I)*B + B*Tan[e + f*x])]/(f*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])$

3.799.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.01, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 87, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{\sqrt{i \tan(e + fx) a + a} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{87} \\
 & \frac{ac \left(-\frac{(A - 2iB) \int \frac{\sqrt{i \tan(e + fx) a + a}}{\sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{c} - \frac{(B + iA)(a + ia \tan(e + fx))^{3/2}}{ac \sqrt{c - ic \tan(e + fx)}} \right)}{f} \\
 & \quad \downarrow \text{60} \\
 & \frac{ac \left(-\frac{(A - 2iB) \left(a \int \frac{1}{\sqrt{i \tan(e + fx) a + a} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) + \frac{i \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}}{c} \right)}{c} - \frac{(B + iA)(a + ia \tan(e + fx))^{3/2}}{ac \sqrt{c - ic \tan(e + fx)}} \right)}{f} \\
 & \quad \downarrow \text{45}
 \end{aligned}$$

3.799. $\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx$

$$\begin{array}{c}
 \left(\frac{(A-2iB) \left(2a \int \frac{1}{ia + \frac{ic \tan(e+fx)a+a}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} + \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{c} \right)}{c} - \frac{(B+ia)(a+ia \tan(e+fx))^{3/2}}{ac\sqrt{c-ic \tan(e+fx)}} \right) \\
 \hline
 f \\
 \downarrow \text{218} \\
 \left(\frac{(A-2iB) \left(\frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{c} - \frac{2i\sqrt{a} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{\sqrt{c}} \right)}{c} - \frac{(B+ia)(a+ia \tan(e+fx))^{3/2}}{ac\sqrt{c-ic \tan(e+fx)}} \right) \\
 \hline
 f
 \end{array}$$

input `Int[((a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x]))/Sqrt[c - I*c*Tan[e + f*x]],x]`

output `(a*c*(-(((I*A + B)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c*Sqrt[c - I*c*Tan[e + f*x]]))) - ((A - (2*I)*B)*(((-2*I)*Sqrt[a]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[c] + (I*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/c))/c)/f`

3.799.3.1 Defintions of rubi rules used

rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`


```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p
_.), x_] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e)), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1))]/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.799.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 496 vs. 2(140) = 280.

Time = 0.35 (sec) , antiderivative size = 497, normalized size of antiderivative = 2.94

method	result
derivativedivides	$\left(2iB \ln \left(\frac{ac \tan(fx+e) + \sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right)\right) ac \tan(fx+e)^2 - 2iA \ln \left(\frac{ac \tan(fx+e) + \sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) ac \tan(fx+e)$
default	$\left(2iB \ln \left(\frac{ac \tan(fx+e) + \sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right)\right) ac \tan(fx+e)^2 - 2iA \ln \left(\frac{ac \tan(fx+e) + \sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) ac \tan(fx+e)$
parts	$iA \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a \left(i \ln \left(\frac{ac \tan(fx+e) + \sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right) \tan(fx+e)^2 ac - i \ln \left(\frac{ac \tan(fx+e) + \sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) ac \tan(fx+e)$

3.799.
$$\int \frac{(a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} dx$$

```
input int((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x,m
method=_RETURNVERBOSE)
```

```
output 1/f*(2*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a
*c)^(1/2))*a*c*tan(f*x+e)^2-2*I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+t
an(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)-A*ln((a*c*tan(f*x+e)+(a*c
)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)^2-2*I*B*
ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*
a*c-4*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-4*B*ln((a*c*
tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(
f*x+e)-B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+2*I*A*(a*c
)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)+A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c
(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c+2*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+
e)^2))^(1/2)*tan(f*x+e)+3*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*(
1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a/c/(a*c*(1+tan(f*x+e)
^2))^(1/2)/(I+tan(f*x+e))^2/(a*c)^(1/2)
```

3.799.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 438 vs. 2(131) = 262.

Time = 0.28 (sec) , antiderivative size = 438, normalized size of antiderivative = 2.59

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx = \frac{c \sqrt{\frac{(A^2 - 4iAB - 4B^2)a^3}{cf^2}} f \log \left(- \frac{4 \left(2((-iA - 2B)ae^{(3i fx + 3i e)} + (- \right)}{2((-iA - 2B)ae^{(3i fx + 3i e)} + (- \right)} \right)}{2((-iA - 2B)ae^{(3i fx + 3i e)} + (- \right)}$$

```
input integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/
2),x, algorithm="fricas")
```

output $\frac{1}{2} * (c * \sqrt{(A^2 - 4 * I * A * B - 4 * B^2) * a^3 / (c * f^2)}) * f * \log(-4 * (2 * ((-I * A - 2 * B) * a * e^{(3 * I * f * x + 3 * I * e)} + (-I * A - 2 * B) * a * e^{(I * f * x + I * e)})) * \sqrt{a / (e^{(2 * I * f * x + 2 * I * e)} + 1)}) * \sqrt{c / (e^{(2 * I * f * x + 2 * I * e)} + 1)}) + (c * f * e^{(2 * I * f * x + 2 * I * e)} - c * f) * \sqrt{(A^2 - 4 * I * A * B - 4 * B^2) * a^3 / (c * f^2)}) / ((I * A + 2 * B) * a * e^{(2 * I * f * x + 2 * I * e)} + (I * A + 2 * B) * a)) - c * \sqrt{(A^2 - 4 * I * A * B - 4 * B^2) * a^3 / (c * f^2)}) * f * \log(-4 * (2 * ((-I * A - 2 * B) * a * e^{(3 * I * f * x + 3 * I * e)} + (-I * A - 2 * B) * a * e^{(I * f * x + I * e)})) * \sqrt{a / (e^{(2 * I * f * x + 2 * I * e)} + 1)}) * \sqrt{c / (e^{(2 * I * f * x + 2 * I * e)} + 1)}) - (c * f * e^{(2 * I * f * x + 2 * I * e)} - c * f) * \sqrt{(A^2 - 4 * I * A * B - 4 * B^2) * a^3 / (c * f^2)}) / ((I * A + 2 * B) * a * e^{(2 * I * f * x + 2 * I * e)} + (I * A + 2 * B) * a)) - 4 * ((I * A + B) * a * e^{(3 * I * f * x + 3 * I * e)} + (I * A + 2 * B) * a * e^{(I * f * x + I * e)}) * \sqrt{a / (e^{(2 * I * f * x + 2 * I * e)} + 1)}) * \sqrt{c / (e^{(2 * I * f * x + 2 * I * e)} + 1)}) / (c * f)$

3.799.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx = \int \frac{(ia(\tan(e + fx) - i))^{3/2} (A + B \tan(e + fx))}{\sqrt{-ic(\tan(e + fx) + i)}} dx$$

input `integrate((a+I*a*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(1/2),x)`

output `Integral((I*a*(tan(e + f*x) - I))**(3/2)*(A + B*tan(e + f*x))/sqrt(-I*c*(tan(e + f*x) + I)), x)`

3.799.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 599 vs. $2(131) = 262$.

Time = 0.47 (sec) , antiderivative size = 599, normalized size of antiderivative = 3.54

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx = \frac{(2((A - 2iB)a \cos(2fx + 2e) - (-iA - 2B)a \sin(2fx + 2e)) * \sqrt{a^3 / (c - ic \tan(e + fx))})}{2 * \sqrt{c - ic \tan(e + fx)}}$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x, algorithm="maxima")`

3.799. $\int \frac{(a+ia \tan(e+fx))^{3/2} (A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$

output `(2*((A - 2*I*B)*a*cos(2*f*x + 2*e) - (-I*A - 2*B)*a*sin(2*f*x + 2*e) + (A - 2*I*B)*a)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) + 2*((A - 2*I*B)*a*cos(2*f*x + 2*e) - (-I*A - 2*B)*a*sin(2*f*x + 2*e) + (A - 2*I*B)*a)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), -sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) - 4*((A - I*B)*a*cos(2*f*x + 2*e) + (I*A + B)*a*sin(2*f*x + 2*e) + (A - 2*I*B)*a)*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + ((I*A + 2*B)*a*cos(2*f*x + 2*e) - (A - 2*I*B)*a*sin(2*f*x + 2*e) + (I*A + 2*B)*a)*log(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))^2 + sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))^2 + 2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) + ((-I*A - 2*B)*a*cos(2*f*x + 2*e) + (A - 2*I*B)*a*sin(2*f*x + 2*e) + (-I*A - 2*B)*a)*log(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))^2 + sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))^2 - 2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) - 4*((I*A + B)*a*cos(2*f*x + 2*e) - (A - I*B)*a*sin(2*f*x + 2*e) + (I*A + 2*B)*a)*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sqrt(a)*sqrt(c)/((-2*I*c*cos(2*f*x + 2*e) + 2*c*sin(2*f*x + 2*e) - 2*I*c)*f)`

3.799.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{3/2}}{\sqrt{-ic \tan(fx + e) + c}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(3/2)/sqrt(-I*c*tan(f*x + e) + c), x)`

3.799.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx = \int \frac{(A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{3/2}}{\sqrt{c - c \tan(e + fx) li}} dx$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(3/2))/(c - c*tan(e + f*x)*1i)^(1/2), x)`

output `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(3/2))/(c - c*tan(e + f*x)*1i)^(1/2), x)`

3.800
$$\int \frac{(a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{3/2}} dx$$

3.800.1 Optimal result	7223
3.800.2 Mathematica [A] (verified)	7223
3.800.3 Rubi [A] (verified)	7224
3.800.4 Maple [B] (verified)	7226
3.800.5 Fricas [B] (verification not implemented)	7227
3.800.6 Sympy [F]	7228
3.800.7 Maxima [A] (verification not implemented)	7228
3.800.8 Giac [F]	7229
3.800.9 Mupad [F(-1)]	7229

3.800.1 Optimal result

Integrand size = 45, antiderivative size = 155

$$\int \frac{(a + ia \tan(e + fx))^{3/2}(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{3/2}} dx =$$

$$-\frac{2a^{3/2}B \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ict \tan(e+fx)}}\right)}{c^{3/2}f}$$

$$-\frac{(iA + B)(a + ia \tan(e + fx))^{3/2}}{3f(c - ict \tan(e + fx))^{3/2}} + \frac{2aB\sqrt{a + ia \tan(e + fx)}}{cf\sqrt{c - ict \tan(e + fx)}}$$

output `-2*a^(3/2)*B*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/c^(3/2)/f+2*a*B*(a+I*a*tan(f*x+e))^(1/2)/c/f/(c-I*c*tan(f*x+e))^(1/2)-1/3*(I*A+B)*(a+I*a*tan(f*x+e))^(3/2)/f/(c-I*c*tan(f*x+e))^(3/2)`

3.800.2 Mathematica [A] (verified)

Time = 7.25 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.41

$$\int \frac{(a + ia \tan(e + fx))^{3/2}(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{3/2}} dx = \frac{a^{3/2} \sec^2(e + fx) \left(\sqrt{a}(6iB + (A - iB) \cos(2(e + fx))) \right)}{c^2}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(3/2),x]`

output $(a^{3/2} \operatorname{Sec}[e + f*x]^2 (\operatorname{Sqrt}[a] ((6I)B + (A - I*B) \operatorname{Cos}[2*(e + f*x)] + (I*A + B) \operatorname{Sin}[2*(e + f*x)]) \operatorname{Sqrt}[1 - I \operatorname{Tan}[e + f*x]] - (6I)B \operatorname{ArcSin}[\operatorname{Sqrt}[a + I*a \operatorname{Tan}[e + f*x]] / (\operatorname{Sqrt}[2] \operatorname{Sqrt}[a])] (\operatorname{Cos}[2*(e + f*x)] - I \operatorname{Sin}[2*(e + f*x)]) \operatorname{Sqrt}[a + I*a \operatorname{Tan}[e + f*x]])) / (3*c*f \operatorname{Sqrt}[1 - I \operatorname{Tan}[e + f*x]] (I + \operatorname{Tan}[e + f*x]) \operatorname{Sqrt}[a + I*a \operatorname{Tan}[e + f*x]] \operatorname{Sqrt}[c - I*c \operatorname{Tan}[e + f*x]])$

3.800.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.08, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 87, 57, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{\sqrt{i \tan(e+fx)a+a} (A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{87} \\
 & \frac{ac \left(\frac{iB \int \frac{\sqrt{i \tan(e+fx)a+a}}{(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{c} - \frac{(B+iA)(a+ia \tan(e+fx))^{3/2}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{f} \\
 & \quad \downarrow \text{57} \\
 & \frac{ac \left(\frac{iB \left(-\frac{a \int \frac{1}{\sqrt{i \tan(e+fx)a+a} \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{c} - \frac{2i \sqrt{a+ia \tan(e+fx)}}{c \sqrt{c-ic \tan(e+fx)}} \right)}{c} - \frac{(B+iA)(a+ia \tan(e+fx))^{3/2}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{f} \\
 & \quad \downarrow \text{45}
 \end{aligned}$$

3.800. $\int \frac{(a+ia \tan(e+fx))^{3/2} (A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$

$$\begin{array}{c}
 \left(\frac{ac \left(\frac{2a \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} dx \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} - \frac{2i \sqrt{a+ia \tan(e+fx)}}{c \sqrt{c-ic \tan(e+fx)}} \right)}{c} - \frac{(B+ia)(a+ia \tan(e+fx))^{3/2}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{f} \\
 \downarrow \text{218} \\
 \left(\frac{ac \left(\frac{2i \sqrt{a} \arctan\left(\frac{\sqrt{c} \sqrt{a+ia \tan(e+fx)}}{\sqrt{a} \sqrt{c-ic \tan(e+fx)}}\right)}{c^{3/2}} - \frac{2i \sqrt{a+ia \tan(e+fx)}}{c \sqrt{c-ic \tan(e+fx)}} \right)}{c} - \frac{(B+ia)(a+ia \tan(e+fx))^{3/2}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{f}
 \end{array}$$

```
input Int[((a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(3/2),x]
```

```
output (a*c*(-1/3*((I*A + B)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c*(c - I*c*Tan[e + f*x])^(3/2)) + (I*B*((2*I)*Sqrt[a]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/c^(3/2) - ((2*I)*Sqrt[a + I*a*Tan[e + f*x]])/(c*Sqrt[c - I*c*Tan[e + f*x]]))/c)/f
```

3.800.3.1 Defintions of rubi rules used

```
rule 45 Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]
```

```
rule 57 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))*Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]
```

3.800. $\int \frac{(a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Simp[(- (b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.800.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 405 vs. 2(126) = 252.

Time = 0.39 (sec) , antiderivative size = 406, normalized size of antiderivative = 2.62

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a \left(3 i B \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}}\right)\right)}{3 f c^2(i+\tan (f x+e))^3} a c \tan (f x+e)^3-9 i B \ln$
default	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a \left(3 i B \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}}\right)\right)}{3 f c^2(i+\tan (f x+e))^3} a c \tan (f x+e)^3-9 i B \ln$
parts	$\frac{A \sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a\left(1+\tan (f x+e)^2\right)}{3 f c^2(i+\tan (f x+e))^3} - \frac{i B \sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a}{3 f c^2(i+\tan (f x+e))^3}$

3.800.
$$\int \frac{(a+ia \tan (e+fx))^{3 / 2}(A+B \tan (e+fx))}{(c-ic \tan (e+fx))^{3 / 2}} dx$$

```
input int((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x,m
method=_RETURNVERBOSE)
```

```
output -1/3/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a/c^2*(3*I*B
*I*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))
*a*c*tan(f*x+e)^3-9*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^
2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)-9*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*
(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)^2-7*I*B*(a*c)^(1
/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+A*(a*c*(1+tan(f*x+e)^2))^(1/
2)*(a*c)^(1/2)*tan(f*x+e)^2+3*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan
(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c+5*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2
))^(1/2)+12*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+A*(a*c)^(
1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c*(1+tan(f*x+e)^2))^(1/2)/(I+tan(f*
x+e))^3/(a*c)^(1/2)
```

3.800.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 380 vs. $2(121) = 242$.

Time = 0.29 (sec) , antiderivative size = 380, normalized size of antiderivative = 2.45

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \frac{3c^2 f \sqrt{-\frac{B^2 a^3}{c^3 f^2}} \log \left(\frac{4 \left(2 (Bae^{(3i fx + 3i e)} + Bae^{(i fx + i e)}) \sqrt{\frac{a}{e^{(2i fx + 2i e)}}} \right)}{Ba} \right)}{}$$

```
input integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/
2),x, algorithm="fracas")
```

```
output 1/6*(3*c^2*f*sqrt(-B^2*a^3/(c^3*f^2))*log(4*(2*(B*a*e^(3*I*f*x + 3*I*e) +
B*a*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x
+ 2*I*e) + 1)) + (c^2*f*e^(2*I*f*x + 2*I*e) - c^2*f)*sqrt(-B^2*a^3/(c^3*f^
2)))/(B*a*e^(2*I*f*x + 2*I*e) + B*a)) - 3*c^2*f*sqrt(-B^2*a^3/(c^3*f^2))*l
og(4*(2*(B*a*e^(3*I*f*x + 3*I*e) + B*a*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x
+ 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - (c^2*f*e^(2*I*f*x + 2*
I*e) - c^2*f)*sqrt(-B^2*a^3/(c^3*f^2)))/(B*a*e^(2*I*f*x + 2*I*e) + B*a)) -
2*((I*A + B)*a*e^(5*I*f*x + 5*I*e) + (I*A - 5*B)*a*e^(3*I*f*x + 3*I*e) -
6*B*a*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*
x + 2*I*e) + 1)))/(c^2*f)
```

3.800. $\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx$

3.800.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \int \frac{(ia(\tan(e + fx) - i))^{3/2} (A + B \tan(e + fx))}{(-ic(\tan(e + fx) + i))^{3/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(3/2),x)`

output `Integral((I*a*(tan(e + f*x) - I))**(3/2)*(A + B*tan(e + f*x))/(-I*c*(tan(e + f*x) + I))**(3/2), x)`

3.800.7 Maxima [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.10

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx =$$

$$\frac{6Ba \arctan(\cos(fx + e), \sin(fx + e) + 1) + 6Ba \arctan(\cos(fx + e), -\sin(fx + e) + 1) - 2(-iA - B)a \cos(3fx + 3e) - 12B^2 a \cos(fx + e) + 3I^2 B^2 a \log(\cos(fx + e)^2 + \sin(fx + e)^2 + 2\sin(fx + e) + 1) - 3I^2 B^2 a \log(\cos(fx + e)^2 + \sin(fx + e)^2 - 2\sin(fx + e) + 1) - 2(A - I^2 B)a \sin(3fx + 3e) - 12I^2 B^2 a \sin(fx + e) \sqrt{a}}{(c^2)^{3/2} f}$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `-1/6*(6*B*a*arctan2(cos(f*x + e), sin(f*x + e) + 1) + 6*B*a*arctan2(cos(f*x + e), -sin(f*x + e) + 1) - 2*(-I*A - B)*a*cos(3*f*x + 3*e) - 12*B*a*cos(f*x + e) + 3*I*B*a*log(cos(f*x + e)^2 + sin(f*x + e)^2 + 2*sin(f*x + e) + 1) - 3*I*B*a*log(cos(f*x + e)^2 + sin(f*x + e)^2 - 2*sin(f*x + e) + 1) - 2*(A - I*B)*a*sin(3*f*x + 3*e) - 12*I*B*a*sin(f*x + e))*sqrt(a)/(c^(3/2)*f)`

3.800.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{3/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{3/2}}{(-ict \tan(fx + e) + c)^{3/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(3/2)/(-I*c*tan(f*x + e) + c)^(3/2), x)`

3.800.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{3/2}} dx = \int \frac{(A + B \tan(e + fx)) (a + a \tan(e + fx) i)^{3/2}}{(c - c \tan(e + fx) i)^{3/2}} dx$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(3/2))/(c - c*tan(e + f*x)*1i)^(3/2),x)`

output `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(3/2))/(c - c*tan(e + f*x)*1i)^(3/2), x)`

3.801
$$\int \frac{(a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$$

3.801.1 Optimal result	7230
3.801.2 Mathematica [A] (verified)	7230
3.801.3 Rubi [A] (verified)	7231
3.801.4 Maple [A] (verified)	7232
3.801.5 Fricas [A] (verification not implemented)	7233
3.801.6 Sympy [F]	7233
3.801.7 Maxima [A] (verification not implemented)	7234
3.801.8 Giac [F]	7234
3.801.9 Mupad [B] (verification not implemented)	7235

3.801.1 Optimal result

Integrand size = 45, antiderivative size = 102

$$\int \frac{(a + ia \tan(e + fx))^{3/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx =$$

$$-\frac{(iA + B)(a + ia \tan(e + fx))^{3/2}}{5f(c - ic \tan(e + fx))^{5/2}} - \frac{(iA - 4B)(a + ia \tan(e + fx))^{3/2}}{15cf(c - ic \tan(e + fx))^{3/2}}$$

output `-1/5*(I*A+B)*(a+I*a*tan(f*x+e))^(3/2)/f/(c-I*c*tan(f*x+e))^(5/2)-1/15*(I*A-4*B)*(a+I*a*tan(f*x+e))^(3/2)/c/f/(c-I*c*tan(f*x+e))^(3/2)`

3.801.2 Mathematica [A] (verified)

Time = 7.02 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.09

$$\int \frac{(a + ia \tan(e + fx))^{3/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \frac{a^2 \sec^2(e + fx)(\cos(2(e + fx)) + i \sin(2(e + fx)))(4iA + B + (4i)B \tan(e + fx))}{15c^2 f (i + \tan(e + fx))^2 \sqrt{a + ia \tan(e + fx)}}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(5/2),x]`

output `(a^2*Sec[e + f*x]^2*(Cos[2*(e + f*x)] + I*Sin[2*(e + f*x)])*((4*I)*A - B + (A + (4*I)*B)*Tan[e + f*x])/(15*c^2*f*(I + Tan[e + f*x])^2*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])`

3.801.
$$\int \frac{(a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$$

3.801.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.089$, Rules used = {3042, 4071, 87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{\sqrt{i \tan(e+fx)a+a}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{87} \\
 & \frac{ac \left(\frac{(A+4iB) \int \frac{\sqrt{i \tan(e+fx)a+a}}{(c-ic \tan(e+fx))^{5/2}} d \tan(e+fx)}{5c} - \frac{(B+iA)(a+ia \tan(e+fx))^{3/2}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)}{f} \\
 & \quad \downarrow \text{48} \\
 & \frac{ac \left(-\frac{i(A+4iB)(a+ia \tan(e+fx))^{3/2}}{15ac^2(c-ic \tan(e+fx))^{3/2}} - \frac{(B+iA)(a+ia \tan(e+fx))^{3/2}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)}{f}
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(5/2), x]`

output `(a*c*(-1/5*((I*A + B)*(a + I*a*Tan[e + f*x])^(3/2)))/(a*c*(c - I*c*Tan[e + f*x])^(5/2)) - ((I/15)*(A + (4*I)*B)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c^2*(c - I*c*Tan[e + f*x])^(3/2)))/f`

3.801.3.1 Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*(c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.801.4 Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.88

method	result
derivativedivides	$\frac{i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a(1+\tan(fx+e)^2)(-4A+iA\tan(fx+e)-iB-4B\tan(fx+e))}{15fc^3(i+\tan(fx+e))^4}$
default	$\frac{i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a(1+\tan(fx+e)^2)(-4A+iA\tan(fx+e)-iB-4B\tan(fx+e))}{15fc^3(i+\tan(fx+e))^4}$
risch	$-\frac{a\sqrt{\frac{ae^{2i(fx+e)}}{e^{2i(fx+e)}+1}}(3iAe^{4i(fx+e)}+3Be^{4i(fx+e)}+5iAe^{2i(fx+e)}-5Be^{2i(fx+e)})}{30c^2\sqrt{\frac{c}{e^{2i(fx+e)}+1}}f}$
parts	$-\frac{A\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a(1+\tan(fx+e)^2)(4i+\tan(fx+e))}{15fc^3(i+\tan(fx+e))^4} - \frac{iB\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i$

3.801. $\int \frac{(a+ia\tan(e+fx))^{3/2}(A+B\tan(e+fx))}{(c-ictan(e+fx))^{5/2}} dx$

```
input int((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)
```

```
output 1/15*I/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a/c^3*(1+tan(f*x+e)^2)*(-4*A+I*A*tan(f*x+e)-I*B-4*B*tan(f*x+e))/(I+tan(f*x+e))^4
```

3.801.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.96

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \frac{(3(iA + B)ae^{(7i fx + 7ie)} + 2(4iA - B)ae^{(5i fx + 5ie)} + 5(iA - B)ae^{(3i fx + 3ie)}) \sqrt{\frac{a}{e^{(2i fx + 2ie)} + 1}} \sqrt{\frac{c}{e^{(2i fx + 2ie)} + 1}}}{30c^3 f}$$

```
input integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="fricas")
```

```
output -1/30*(3*(I*A + B)*a*e^(7*I*f*x + 7*I*e) + 2*(4*I*A - B)*a*e^(5*I*f*x + 5*I*e) + 5*(I*A - B)*a*e^(3*I*f*x + 3*I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^3*f)
```

3.801.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \int \frac{(ia(\tan(e + fx) - i))^{\frac{3}{2}} (A + B \tan(e + fx))}{(-ic(\tan(e + fx) + i))^{\frac{5}{2}}} dx$$

```
input integrate((a+I*a*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(5/2),x)
```

```
output Integral((I*a*(tan(e + f*x) - I))**(3/2)*(A + B*tan(e + f*x))/(-I*c*(tan(e + f*x) + I))**(5/2), x)
```

3.801. $\int \frac{(a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$

3.801.7 Maxima [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.51

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx =$$

$$\frac{30(3(A - iB)a \cos(7fx + 7e) + 2(4A + iB)a \cos(5fx + 5e) + 5(A + iB)a \cos(3fx + 3e) - 3(-iA + B)a \sin(7fx + 7e) - 2(-4iA + B)a \sin(5fx + 5e) - 5(-iA + B)a \sin(3fx + 3e)) \sqrt{a} \sqrt{c}}{-900(i c^3 \cos(2fx + 2e) - c^3 \sin(2fx + 2e))}$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `-30*(3*(A - I*B)*a*cos(7*f*x + 7*e) + 2*(4*A + I*B)*a*cos(5*f*x + 5*e) + 5*(A + I*B)*a*cos(3*f*x + 3*e) - 3*(-I*A - B)*a*sin(7*f*x + 7*e) - 2*(-4*I*A + B)*a*sin(5*f*x + 5*e) - 5*(-I*A + B)*a*sin(3*f*x + 3*e))*sqrt(a)*sqrt(c)/((-900*I*c^3*cos(2*f*x + 2*e) + 900*c^3*sin(2*f*x + 2*e) - 900*I*c^3)*f)`

3.801.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{3/2}}{(-ic \tan(fx + e) + c)^{5/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(3/2)/(-I*c*tan(f*x + e) + c)^(5/2), x)`

3.801.9 Mupad [B] (verification not implemented)

Time = 9.82 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.86

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx =$$

$$\frac{a \sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}} (A \cos(2e + 2fx) 5i + A \cos(4e + 4fx) 3i - 5B \cos(2e + 2fx) + 3B \cos(4e + 4fx) - 5A \sin(2e + 2fx) - 3A \sin(4e + 4fx) - B \sin(2e + 2fx) 5i + B \sin(4e + 4fx) 3i)}{30 c^2 f \sqrt{\frac{c(\cos(2e + 2fx) - \sin(2e + 2fx) 1i + 1)}{\cos(2e + 2fx) + 1}}}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(3/2))/(c - c*tan(e + f*x)*1i)^(5/2),x)`

output `-(a*((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*cos(2*e + 2*f*x)*5i + A*cos(4*e + 4*f*x)*3i - 5*B*cos(2*e + 2*f*x) + 3*B*cos(4*e + 4*f*x) - 5*A*sin(2*e + 2*f*x) - 3*A*sin(4*e + 4*f*x) - B*sin(2*e + 2*f*x)*5i + B*sin(4*e + 4*f*x)*3i))/(30*c^2*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

3.802
$$\int \frac{(a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{7/2}} dx$$

3.802.1 Optimal result	7236
3.802.2 Mathematica [A] (verified)	7236
3.802.3 Rubi [A] (verified)	7237
3.802.4 Maple [A] (verified)	7239
3.802.5 Fricas [A] (verification not implemented)	7239
3.802.6 Sympy [F]	7240
3.802.7 Maxima [A] (verification not implemented)	7240
3.802.8 Giac [F]	7241
3.802.9 Mupad [B] (verification not implemented)	7241

3.802.1 Optimal result

Integrand size = 45, antiderivative size = 155

$$\int \frac{(a + ia \tan(e + fx))^{3/2}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx = -\frac{(iA + B)(a + ia \tan(e + fx))^{3/2}}{7f(c - ictan(e + fx))^{7/2}} - \frac{(2iA - 5B)(a + ia \tan(e + fx))^{3/2}}{35cf(c - ictan(e + fx))^{5/2}} - \frac{(2iA - 5B)(a + ia \tan(e + fx))^{3/2}}{105c^2f(c - ictan(e + fx))^{3/2}}$$

output

```
-1/7*(I*A+B)*(a+I*a*tan(f*x+e))^(3/2)/f/(c-I*c*tan(f*x+e))^(7/2)-1/35*(2*I
*A-5*B)*(a+I*a*tan(f*x+e))^(3/2)/c/f/(c-I*c*tan(f*x+e))^(5/2)-1/105*(2*I*A
-5*B)*(a+I*a*tan(f*x+e))^(3/2)/c^2/f/(c-I*c*tan(f*x+e))^(3/2)
```

3.802.2 Mathematica [A] (verified)

Time = 6.89 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.84

$$\int \frac{(a + ia \tan(e + fx))^{3/2}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx = \frac{a^2 \sec^4(e + fx)(\cos(2(e + fx)) + i \sin(2(e + fx)))(21A + 5(5A + 2iB) \cos(2(e + fx)) + 5(-2iA + 5B) \sin(2(e + fx)))}{210c^3 f (i + \tan(e + fx))^3 \sqrt{a + ia \tan(e + fx)} \sqrt{c - ictan(e + fx)}}$$

input

```
Integrate[(((a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan
[e + f*x])^(7/2),x]
```

output
$$\frac{-1/210*(a^2*\text{Sec}[e + f*x]^4*(\text{Cos}[2*(e + f*x)] + I*\text{Sin}[2*(e + f*x)])*(21*A + 5*(5*A + (2*I)*B)*\text{Cos}[2*(e + f*x)] + 5*((-2*I)*A + 5*B)*\text{Sin}[2*(e + f*x)])}{(c^3*f*(I + \text{Tan}[e + f*x])^3*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])}$$

3.802.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.08, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4071, 87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx$$

↓ 4071

$$\frac{ac \int \frac{\sqrt{i \tan(e + fx) a + a} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} d \tan(e + fx)}{f}$$

↓ 87

$$\frac{ac \left(\frac{(2A + 5iB) \int \frac{\sqrt{i \tan(e + fx) a + a}}{(c - ic \tan(e + fx))^{7/2}} d \tan(e + fx)}{7c} - \frac{(B + iA)(a + ia \tan(e + fx))^{3/2}}{7ac(c - ic \tan(e + fx))^{7/2}} \right)}{f}$$

↓ 55

$$\frac{ac \left(\frac{(2A + 5iB) \left(\frac{\int \frac{\sqrt{i \tan(e + fx) a + a}}{(c - ic \tan(e + fx))^{5/2}} d \tan(e + fx)}{5c} - \frac{i(a + ia \tan(e + fx))^{3/2}}{5ac(c - ic \tan(e + fx))^{5/2}} \right)}{7c} - \frac{(B + iA)(a + ia \tan(e + fx))^{3/2}}{7ac(c - ic \tan(e + fx))^{7/2}} \right)}{f}$$

↓ 48

3.802.
$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx$$

$$\frac{ac \left(\frac{(2A+5iB) \left(-\frac{i(a+ia \tan(e+fx))^{3/2}}{15ac^2(c-ic \tan(e+fx))^{3/2}} - \frac{i(a+ia \tan(e+fx))^{3/2}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)}{7c} - \frac{(B+iA)(a+ia \tan(e+fx))^{3/2}}{7ac(c-ic \tan(e+fx))^{7/2}} \right)}{f}$$

input `Int[((a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(7/2), x]`

output `(a*c*(-1/7*((I*A + B)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c*(c - I*c*Tan[e + f*x])^(7/2)) + ((2*A + (5*I)*B)*((-1/5*I)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c*(c - I*c*Tan[e + f*x])^(5/2)) - ((I/15)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c^2*(c - I*c*Tan[e + f*x])^(3/2)))/(7*c))/f`

3.802.3.1 Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x],
, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.802.4 Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.73

method	result
derivativedivides	$\frac{i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a(1+\tan(fx+e)^2)(5B-25i\tan(fx+e)B-5B\tan(fx+e)^2-23iA-10A\tan(fx+e))}{105f c^4(i+\tan(fx+e))^5}$
default	$\frac{i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a(1+\tan(fx+e)^2)(5B-25i\tan(fx+e)B-5B\tan(fx+e)^2-23iA-10A\tan(fx+e))}{105f c^4(i+\tan(fx+e))^5}$
risch	$-\frac{a\sqrt{\frac{a e^{2i(fx+e)}}{e^{2i(fx+e)}+1}}(15iA e^{6i(fx+e)}+15B e^{6i(fx+e)}+42iA e^{4i(fx+e)}+35iA e^{2i(fx+e)}-35B e^{2i(fx+e)})}{420c^3\sqrt{\frac{c}{e^{2i(fx+e)}+1}}f}$
parts	$-\frac{A\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a(1+\tan(fx+e)^2)(-23+10i\tan(fx+e)+2\tan(fx+e)^2)}{105f c^4(i+\tan(fx+e))^5} - \frac{iB\sqrt{a(1+i\tan(fx+e))}}{105f c^4(i+\tan(fx+e))^5}$

```
input int((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x,m
ethod=_RETURNVERBOSE)
```

```
output 1/105*I/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a/c^4*(1+
tan(f*x+e)^2)*(5*B-25*I*B*tan(f*x+e)-5*B*tan(f*x+e)^2-23*I*A-10*A*tan(f*x+
e)+2*I*A*tan(f*x+e)^2)/(I+tan(f*x+e))^5
```

3.802.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.75

$$\int \frac{(a + ia \tan(e + fx))^{3/2}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx =$$

$$\frac{(15(iA + B)ae^{(9i fx+9ie)} + 3(19iA + 5B)ae^{(7i fx+7ie)} + 7(11iA - 5B)ae^{(5i fx+5ie)} + 35(iA - B)ae^{(3i fx+3ie)})}{420c^4f}$$

3.802. $\int \frac{(a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{7/2}} dx$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x, algorithm="fricas")`

output `-1/420*(15*(I*A + B)*a*e^(9*I*f*x + 9*I*e) + 3*(19*I*A + 5*B)*a*e^(7*I*f*x + 7*I*e) + 7*(11*I*A - 5*B)*a*e^(5*I*f*x + 5*I*e) + 35*(I*A - B)*a*e^(3*I*f*x + 3*I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^4*f)`

3.802.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \int \frac{(ia(\tan(e + fx) - i))^{3/2} (A + B \tan(e + fx))}{(-ic(\tan(e + fx) + i))^{7/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(7/2),x)`

output `Integral((I*a*(tan(e + f*x) - I))**(3/2)*(A + B*tan(e + f*x))/(-I*c*(tan(e + f*x) + I))**(7/2), x)`

3.802.7 Maxima [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.20

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \frac{(15(-iA - B)a \cos(\frac{7}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)), \cos(2fx + 2e))}{(c - ic \tan(e + fx))^{7/2}}$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x, algorithm="maxima")`

output `1/420*(15*(-I*A - B)*a*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 42*I*A*a*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 35*(-I*A + B)*a*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 15*(A - I*B)*a*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 42*A*a*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 35*(A + I*B)*a*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sqrt(a)/(c^(7/2)*f)`

3.802. $\int \frac{(a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$

3.802.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{3/2}}{(-ic \tan(fx + e) + c)^{7/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(3/2)/(-I*c*tan(f*x + e) + c)^(7/2), x)`

3.802.9 Mupad [B] (verification not implemented)

Time = 10.30 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.39

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx =$$

$$a \sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}} (A \cos(2e + 2fx) 35i + A \cos(4e + 4fx) 42i + A \cos(6e + 6fx) 15i$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(3/2))/(c - c*tan(e + f*x)*1i)^(7/2),x)`

output `-(a*((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*cos(2*e + 2*f*x)*35i + A*cos(4*e + 4*f*x)*42i + A*cos(6*e + 6*f*x)*15i - 35*B*cos(2*e + 2*f*x) + 15*B*cos(6*e + 6*f*x) - 35*A*sin(2*e + 2*f*x) - 42*A*sin(4*e + 4*f*x) - 15*A*sin(6*e + 6*f*x) - B*sin(2*e + 2*f*x)*35i + B*sin(6*e + 6*f*x)*15i))/(420*c^3*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

3.803
$$\int \frac{(a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{9/2}} dx$$

3.803.1 Optimal result	7242
3.803.2 Mathematica [A] (verified)	7242
3.803.3 Rubi [A] (verified)	7243
3.803.4 Maple [A] (verified)	7245
3.803.5 Fricas [A] (verification not implemented)	7246
3.803.6 Sympy [F(-1)]	7246
3.803.7 Maxima [A] (verification not implemented)	7247
3.803.8 Giac [F]	7247
3.803.9 Mupad [B] (verification not implemented)	7248

3.803.1 Optimal result

Integrand size = 45, antiderivative size = 208

$$\int \frac{(a + ia \tan(e + fx))^{3/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx =$$

$$\frac{(iA + B)(a + ia \tan(e + fx))^{3/2}}{9f(c - ic \tan(e + fx))^{9/2}} - \frac{(iA - 2B)(a + ia \tan(e + fx))^{3/2}}{21cf(c - ic \tan(e + fx))^{7/2}}$$

$$- \frac{2(iA - 2B)(a + ia \tan(e + fx))^{3/2}}{105c^2f(c - ic \tan(e + fx))^{5/2}} - \frac{2(iA - 2B)(a + ia \tan(e + fx))^{3/2}}{315c^3f(c - ic \tan(e + fx))^{3/2}}$$

```
output -1/9*(I*A+B)*(a+I*a*tan(f*x+e))^(3/2)/f/(c-I*c*tan(f*x+e))^(9/2)-1/21*(I*A
-2*B)*(a+I*a*tan(f*x+e))^(3/2)/c/f/(c-I*c*tan(f*x+e))^(7/2)-2/105*(I*A-2*B
)*(a+I*a*tan(f*x+e))^(3/2)/c^2/f/(c-I*c*tan(f*x+e))^(5/2)-2/315*(I*A-2*B)*
(a+I*a*tan(f*x+e))^(3/2)/c^3/f/(c-I*c*tan(f*x+e))^(3/2)
```

3.803.2 Mathematica [A] (verified)

Time = 7.06 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.63

$$\int \frac{(a + ia \tan(e + fx))^{3/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx =$$

$$\frac{a^2(-i + \tan(e + fx))^2(-58iA + 11B - 33(A + 2iB) \tan(e + fx) + 12i(A + 2iB) \tan^2(e + fx) + 2(A +$$

$$315c^4f(i + \tan(e + fx))^4\sqrt{a + ia \tan(e + fx)}\sqrt{c - ic \tan(e + fx)})}{315c^4f(i + \tan(e + fx))^4\sqrt{a + ia \tan(e + fx)}\sqrt{c - ic \tan(e + fx)}}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(9/2),x]`

output `-1/315*(a^2*(-I + Tan[e + f*x])^2*((-58*I)*A + 11*B - 33*(A + (2*I)*B)*Tan[e + f*x] + (12*I)*(A + (2*I)*B)*Tan[e + f*x]^2 + 2*(A + (2*I)*B)*Tan[e + f*x]^3)/(c^4*f*(I + Tan[e + f*x])^4*sqrt[a + I*a*Tan[e + f*x]]*sqrt[c - I*c*Tan[e + f*x]])`

3.803.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.05, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 87, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{\sqrt{i \tan(e+fx)a+a}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{11/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{87} \\
 & \frac{ac \left(\frac{(A+2iB) \int \frac{\sqrt{i \tan(e+fx)a+a}}{(c-ic \tan(e+fx))^{9/2}} d \tan(e+fx)}{3c} - \frac{(B+iA)(a+ia \tan(e+fx))^{3/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{f} \\
 & \quad \downarrow \text{55} \\
 & \frac{ac \left(\frac{(A+2iB) \left(\frac{2 \int \frac{\sqrt{i \tan(e+fx)a+a}}{(c-ic \tan(e+fx))^{7/2}} d \tan(e+fx)}{7c} - \frac{i(a+ia \tan(e+fx))^{3/2}}{7ac(c-ic \tan(e+fx))^{7/2}} \right)}{3c} - \frac{(B+iA)(a+ia \tan(e+fx))^{3/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{f}
 \end{aligned}$$

3.803. $\int \frac{(a+ia \tan(e+fx))^{3/2} (A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{9/2}} dx$

$$\begin{array}{c}
 \downarrow 55 \\
 \left(\frac{(A+2iB) \left(\frac{2 \left(\int \frac{\sqrt{i \tan(e+fx)a+a}}{(c-ic \tan(e+fx))^{5/2}} d \tan(e+fx) - \frac{i(a+ia \tan(e+fx))^{3/2}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)}{7c} - \frac{i(a+ia \tan(e+fx))^{3/2}}{7ac(c-ic \tan(e+fx))^{7/2}} \right)}{3c} - \frac{(B+iA)(a+ia \tan(e+fx))^{3/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{ac} \right) \\
 \hline
 f \\
 \downarrow 48 \\
 \left(\frac{(A+2iB) \left(\frac{2 \left(-\frac{i(a+ia \tan(e+fx))^{3/2}}{15ac^2(c-ic \tan(e+fx))^{3/2}} - \frac{i(a+ia \tan(e+fx))^{3/2}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)}{7c} - \frac{i(a+ia \tan(e+fx))^{3/2}}{7ac(c-ic \tan(e+fx))^{7/2}} \right)}{3c} - \frac{(B+iA)(a+ia \tan(e+fx))^{3/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{ac} \right) \\
 \hline
 f
 \end{array}$$

input `Int[((a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(9/2), x]`

output `(a*c*(-1/9*((I*A + B)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c*(c - I*c*Tan[e + f*x])^(9/2)) + ((A + (2*I)*B)*((-1/7*I)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c*(c - I*c*Tan[e + f*x])^(7/2)) + (2*((-1/5*I)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c*(c - I*c*Tan[e + f*x])^(5/2)) - ((I/15)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c^2*(c - I*c*Tan[e + f*x])^(3/2)))/(7*c))/(3*c))/f`

3.803.3.1 Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

```
rule 55 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*(c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.803.4 Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.65

method	result
derivativedivides	$\frac{i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a(1+\tan(fx+e)^2)\left(2iA\tan(fx+e)^3-24iB\tan(fx+e)^2-4B\tan(fx+e)^3-315fc^5(i+\tan(fx+e))^6\right)}{315fc^5(i+\tan(fx+e))^6}$
default	$\frac{i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a(1+\tan(fx+e)^2)\left(2iA\tan(fx+e)^3-24iB\tan(fx+e)^2-4B\tan(fx+e)^3-315fc^5(i+\tan(fx+e))^6\right)}{315fc^5(i+\tan(fx+e))^6}$
risch	$-\frac{a\sqrt{\frac{ae^{2i(fx+e)}}{e^{2i(fx+e)}+1}}(35iAe^{8i(fx+e)}+35Be^{8i(fx+e)}+135iAe^{6i(fx+e)}+45Be^{6i(fx+e)}+189iAe^{4i(fx+e)}-63Be^{4i(fx+e)})}{2520c^4\sqrt{\frac{c}{e^{2i(fx+e)}+1}}f}$
parts	$\frac{iA\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a(1+\tan(fx+e)^2)\left(58-33i\tan(fx+e)-12\tan(fx+e)^2+2i\tan(fx+e)^3\right)}{315fc^5(i+\tan(fx+e))^6}$

3.803.
$$\int \frac{(a+ia\tan(e+fx))^{3/2}(A+B\tan(e+fx))}{(c-ictan(e+fx))^{9/2}} dx$$

```
input int((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(9/2),x,method=_RETURNVERBOSE)
```

```
output 1/315*I/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a/c^5*(1+tan(f*x+e)^2)*(2*I*A*tan(f*x+e)^3-24*I*B*tan(f*x+e)^2-4*B*tan(f*x+e)^3-33*I*A*tan(f*x+e)-12*A*tan(f*x+e)^2+11*I*B+66*B*tan(f*x+e)+58*A)/(I+tan(f*x+e))^6
```

3.803.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.65

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx =$$

$$\frac{(35(iA + B)ae^{(11ifx+11ie)} + 10(17iA + 8B)ae^{(9ifx+9ie)} + 18(18iA - B)ae^{(7ifx+7ie)} + 42(7iA - 4B)ae^{(5ifx+5ie)} + 105(IA - B)ae^{(3ifx+3ie)})\sqrt{a/(e^{(2ifx+2ie)} + 1)}\sqrt{c/(e^{(2ifx+2ie)} + 1)}}{2520c^5f}$$

```
input integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(9/2),x, algorithm="fricas")
```

```
output -1/2520*(35*(I*A + B)*a*e^(11*I*f*x + 11*I*e) + 10*(17*I*A + 8*B)*a*e^(9*I*f*x + 9*I*e) + 18*(18*I*A - B)*a*e^(7*I*f*x + 7*I*e) + 42*(7*I*A - 4*B)*a*e^(5*I*f*x + 5*I*e) + 105*(I*A - B)*a*e^(3*I*f*x + 3*I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^5*f)
```

3.803.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx = \text{Timed out}$$

```
input integrate((a+I*a*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(9/2),x)
```

```
output Timed out
```

3.803. $\int \frac{(a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{9/2}} dx$

3.803.7 Maxima [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 260, normalized size of antiderivative = 1.25

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx = \frac{(35(-iA - B)a \cos(\frac{9}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 45(-3iA - B)a \cos(\frac{7}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 63(-3iA + B)a \cos(\frac{5}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 105(-iA + B)a \cos(\frac{3}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e))) + 35(A - iB)a \sin(\frac{9}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 45(3A - iB)a \sin(\frac{7}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 63(3A + iB)a \sin(\frac{5}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 105(A + iB)a \sin(\frac{3}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e))}{(c - ic \tan(e + fx))^{9/2}}$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(9/2),x, algorithm="maxima")`

output `1/2520*(35*(-I*A - B)*a*cos(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 45*(-3*I*A - B)*a*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 63*(-3*I*A + B)*a*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 105*(-I*A + B)*a*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 35*(A - I*B)*a*sin(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 45*(3*A - I*B)*a*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 63*(3*A + I*B)*a*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 105*(A + I*B)*a*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sqrt(a)/(c^(9/2)*f)`

3.803.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{3/2}}{(-ic \tan(fx + e) + c)^{9/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(9/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(3/2)/(-I*c*tan(f*x + e) + c)^(9/2), x)`

3.803.9 Mupad [B] (verification not implemented)

Time = 11.05 (sec) , antiderivative size = 290, normalized size of antiderivative = 1.39

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx =$$

$$a \sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)i)}{\cos(2e+2fx)+1}} (A \cos(2e + 2fx) 105i + A \cos(4e + 4fx) 189i + A \cos(6e + 6fx))$$

```
input int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(3/2))/(c - c*tan(e + f*
x)*1i)^(9/2),x)
```

```
output -(a*((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) +
1))^(1/2)*(A*cos(2*e + 2*f*x)*105i + A*cos(4*e + 4*f*x)*189i + A*cos(6*e +
6*f*x)*135i + A*cos(8*e + 8*f*x)*35i - 105*B*cos(2*e + 2*f*x) - 63*B*cos(
4*e + 4*f*x) + 45*B*cos(6*e + 6*f*x) + 35*B*cos(8*e + 8*f*x) - 105*A*sin(2
*e + 2*f*x) - 189*A*sin(4*e + 4*f*x) - 135*A*sin(6*e + 6*f*x) - 35*A*sin(8
*e + 8*f*x) - B*sin(2*e + 2*f*x)*105i - B*sin(4*e + 4*f*x)*63i + B*sin(6*e
+ 6*f*x)*45i + B*sin(8*e + 8*f*x)*35i))/(2520*c^4*f*((c*(cos(2*e + 2*f*x)
- sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))
```

3.804
$$\int \frac{(a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{11/2}} dx$$

3.804.1 Optimal result	7249
3.804.2 Mathematica [A] (verified)	7250
3.804.3 Rubi [A] (verified)	7250
3.804.4 Maple [A] (verified)	7253
3.804.5 Fricas [A] (verification not implemented)	7254
3.804.6 Sympy [F(-1)]	7254
3.804.7 Maxima [A] (verification not implemented)	7255
3.804.8 Giac [F]	7255
3.804.9 Mupad [B] (verification not implemented)	7256

3.804.1 Optimal result

Integrand size = 45, antiderivative size = 261

$$\int \frac{(a + ia \tan(e + fx))^{3/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{11/2}} dx =$$

$$\frac{(iA + B)(a + ia \tan(e + fx))^{3/2}}{11f(c - ic \tan(e + fx))^{11/2}} - \frac{(4iA - 7B)(a + ia \tan(e + fx))^{3/2}}{99cf(c - ic \tan(e + fx))^{9/2}}$$

$$- \frac{(4iA - 7B)(a + ia \tan(e + fx))^{3/2}}{231c^2f(c - ic \tan(e + fx))^{7/2}} - \frac{2(4iA - 7B)(a + ia \tan(e + fx))^{3/2}}{1155c^3f(c - ic \tan(e + fx))^{5/2}}$$

$$- \frac{2(4iA - 7B)(a + ia \tan(e + fx))^{3/2}}{3465c^4f(c - ic \tan(e + fx))^{3/2}}$$

output

```
-1/11*(I*A+B)*(a+I*a*tan(f*x+e))^(3/2)/f/(c-I*c*tan(f*x+e))^(11/2)-1/99*(4
*I*A-7*B)*(a+I*a*tan(f*x+e))^(3/2)/c/f/(c-I*c*tan(f*x+e))^(9/2)-1/231*(4*I
*A-7*B)*(a+I*a*tan(f*x+e))^(3/2)/c^2/f/(c-I*c*tan(f*x+e))^(7/2)-2/1155*(4*
I*A-7*B)*(a+I*a*tan(f*x+e))^(3/2)/c^3/f/(c-I*c*tan(f*x+e))^(5/2)-2/3465*(4
*I*A-7*B)*(a+I*a*tan(f*x+e))^(3/2)/c^4/f/(c-I*c*tan(f*x+e))^(3/2)
```


3.804.2 Mathematica [A] (verified)

Time = 7.70 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.59

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx =$$

$$\frac{a^2(-i + \tan(e + fx))^2 (547A + 91iB + 91(-4iA + 7B) \tan(e + fx) - 45(4A + 7iB) \tan^2(e + fx) + (56A + 91iB) \tan^3(e + fx))}{3465c^5 f(i + \tan(e + fx))^5 \sqrt{a + ia \tan(e + fx)} \sqrt{c - ictan(e + fx)}}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(11/2),x]`

output `-1/3465*(a^2*(-I + Tan[e + f*x])^2*(547*A + (91*I)*B + 91*((-4*I)*A + 7*B)*Tan[e + f*x] - 45*(4*A + (7*I)*B)*Tan[e + f*x]^2 + ((56*I)*A - 98*B)*Tan[e + f*x]^3 + 2*(4*A + (7*I)*B)*Tan[e + f*x]^4))/(c^5*f*(I + Tan[e + f*x])^5*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])`

3.804.3 Rubi [A] (verified)Time = 0.46 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 87, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx$$

$$\downarrow \text{3042}$$

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx$$

$$\downarrow \text{4071}$$

$$\frac{ac \int \frac{\sqrt{i \tan(e+fx)a+a(A+B \tan(e+fx))}}{(c-ictan(e+fx))^{13/2}} d \tan(e + fx)}{f}$$

$$\downarrow \text{87}$$

3.804. $\int \frac{(a+ia \tan(e+fx))^{3/2} (A+B \tan(e+fx))}{(c-ictan(e+fx))^{11/2}} dx$

$$\begin{array}{c}
 \frac{ac \left(\frac{(4A+7iB) \int \frac{\sqrt{i \tan(e+fx)a+a}}{(c-ic \tan(e+fx))^{11/2}} d \tan(e+fx)}{11c} - \frac{(B+iA)(a+ia \tan(e+fx))^{3/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{f} \\
 \downarrow 55 \\
 \frac{ac \left(\frac{(4A+7iB) \left(\frac{\int \frac{\sqrt{i \tan(e+fx)a+a}}{(c-ic \tan(e+fx))^{9/2}} d \tan(e+fx)}{3c} - \frac{i(a+ia \tan(e+fx))^{3/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{(B+iA)(a+ia \tan(e+fx))^{3/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{f} \\
 \downarrow 55 \\
 \frac{ac \left(\frac{(4A+7iB) \left(\frac{2 \int \frac{\sqrt{i \tan(e+fx)a+a}}{(c-ic \tan(e+fx))^{7/2}} d \tan(e+fx)}{7c} - \frac{i(a+ia \tan(e+fx))^{3/2}}{7ac(c-ic \tan(e+fx))^{7/2}} - \frac{i(a+ia \tan(e+fx))^{3/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{(B+iA)(a+ia \tan(e+fx))^{3/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{f} \\
 \downarrow 55 \\
 \frac{ac \left(\frac{(4A+7iB) \left(\frac{2 \left(\frac{\int \frac{\sqrt{i \tan(e+fx)a+a}}{(c-ic \tan(e+fx))^{5/2}} d \tan(e+fx)}{5c} - \frac{i(a+ia \tan(e+fx))^{3/2}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)}{7c} - \frac{i(a+ia \tan(e+fx))^{3/2}}{7ac(c-ic \tan(e+fx))^{7/2}} - \frac{i(a+ia \tan(e+fx))^{3/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{(B+iA)(a+ia \tan(e+fx))^{3/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{f} \\
 \downarrow 48
 \end{array}$$

3.804. $\int \frac{(a+ia \tan(e+fx))^{3/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{11/2}} dx$

$$ac \left(\frac{(4A+7iB) \left(\frac{2 \left(-\frac{i(a+ia \tan(e+fx))^{3/2}}{15ac^2(c-ic \tan(e+fx))^{3/2}} - \frac{i(a+ia \tan(e+fx))^{3/2}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)}{7c} - \frac{i(a+ia \tan(e+fx))^{3/2}}{3c} - \frac{i(a+ia \tan(e+fx))^{3/2}}{7ac(c-ic \tan(e+fx))^{7/2}} - \frac{i(a+ia \tan(e+fx))^{3/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{(B+iA)(a+ia \tan(e+fx))^{3/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right) dx$$

```
input Int[((a + I*a*Tan[e + f*x])^(3/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(11/2), x]
```

```
output (a*c*(-1/11*((I*A + B)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c*(c - I*c*Tan[e + f*x])^(11/2)) + (((4*A + (7*I)*B)*((-1/9*I)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c*(c - I*c*Tan[e + f*x])^(9/2)) + (((-1/7*I)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c*(c - I*c*Tan[e + f*x])^(7/2)) + (2*((-1/5*I)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c*(c - I*c*Tan[e + f*x])^(5/2)) - ((I/15)*(a + I*a*Tan[e + f*x])^(3/2))/(a*c^2*(c - I*c*Tan[e + f*x])^(3/2))))/(7*c))/(3*c))/(11*c))/f
```

3.804.3.1 Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 55 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e)), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1))]/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.804.4 Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.61

method	result
derivativedivides	$-\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a\left(1+\tan (f x+e)^2\right)\left(14 i B \tan (f x+e)^4+56 i A \tan (f x+e)^3+8 A \tan (f x+e)^2\right)}{3465 f c^6(i+\tan (f x+e))}$
default	$-\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a\left(1+\tan (f x+e)^2\right)\left(14 i B \tan (f x+e)^4+56 i A \tan (f x+e)^3+8 A \tan (f x+e)^2\right)}{3465 f c^6(i+\tan (f x+e))}$
risch	$-\frac{a \sqrt{\frac{a e^{2 i(f x+e)}}{e^{2 i(f x+e)}+1}}\left(315 i A e^{10 i(f x+e)}+315 B e^{10 i(f x+e)}+1540 i A e^{8 i(f x+e)}+770 B e^{8 i(f x+e)}+2970 i A e^{6 i(f x+e)}+2772 i A e^{4 i(f x+e)}+180 B e^{4 i(f x+e)}\right)}{55440 c^5 \sqrt{\frac{c}{e^{2 i(f x+e)}+1}} f}$
parts	$-\frac{A \sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a\left(1+\tan (f x+e)^2\right)\left(56 i \tan (f x+e)^3+8 \tan (f x+e)^4-364 i \tan (f x+e)^2+91\right)}{3465 f c^6(i+\tan (f x+e))^7}$

```
input int((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(11/2), x,
method=_RETURNVERBOSE)
```

```
output -1/3465/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a/c^6*(1+
tan(f*x+e)^2)*(14*I*B*tan(f*x+e)^4+56*I*A*tan(f*x+e)^3+8*A*tan(f*x+e)^2-31
5*I*B*tan(f*x+e)-98*B*tan(f*x+e)^3-364*I*A*tan(f*x+e)-180*A*tan(f*x+e)^2
+91*I*B+637*B*tan(f*x+e)+547*A)/(I+tan(f*x+e))^7
```

$$3.804. \int \frac{(a+ia \tan (e+fx))^{3/2}(A+B \tan (e+fx))}{(c-ic \tan (e+fx))^{11/2}} dx$$

3.804.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 155, normalized size of antiderivative = 0.59

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx =$$

$$(315 (i A + B) a e^{(13i fx + 13i e)} + 35 (53i A + 31 B) a e^{(11i fx + 11i e)} + 110 (41i A + 7 B) a e^{(9i fx + 9i e)} + 198 (29i A - 7 B) a e^{(7i fx + 7i e)} + 231 (17i A - 11 B) a e^{(5i fx + 5i e)} + 1155 (i A - B) a e^{(3i fx + 3i e)}) \sqrt{a/(e^{(2i fx + 2i e)} + 1)} \sqrt{c/(e^{(2i fx + 2i e)} + 1)} / (c^6 f)$$

55

input `integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(11/2),x, algorithm="fracas")`

output `-1/55440*(315*(I*A + B)*a*e^(13*I*f*x + 13*I*e) + 35*(53*I*A + 31*B)*a*e^(11*I*f*x + 11*I*e) + 110*(41*I*A + 7*B)*a*e^(9*I*f*x + 9*I*e) + 198*(29*I*A - 7*B)*a*e^(7*I*f*x + 7*I*e) + 231*(17*I*A - 11*B)*a*e^(5*I*f*x + 5*I*e) + 1155*(I*A - B)*a*e^(3*I*f*x + 3*I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^6*f)`

3.804.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(11/2),x)`

output `Timed out`

3.804.7 Maxima [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 312, normalized size of antiderivative = 1.20

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx = \frac{(315(-iA - B)a \cos(\frac{11}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 770(-2iA - B)a \cos(\frac{9}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) - 2970iAa \cos(\frac{7}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 1386(-2iA + B)a \cos(\frac{5}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 1155(-iA + B)a \cos(\frac{3}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 315(A - iB)a \sin(\frac{11}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 770(2A - iB)a \sin(\frac{9}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 2970Aa \sin(\frac{7}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 1386(2A + iB)a \sin(\frac{5}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 1155(A + iB)a \sin(\frac{3}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e))) \sqrt{a}}{(c^{11/2}) * f}$$

```
input integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(11/2),x, algorithm="maxima")
```

```
output 1/55440*(315*(-I*A - B)*a*cos(11/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 770*(-2*I*A - B)*a*cos(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 2970*I*A*a*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1386*(-2*I*A + B)*a*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1155*(-I*A + B)*a*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 315*(A - I*B)*a*sin(11/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 770*(2*A - I*B)*a*sin(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 2970*A*a*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1386*(2*A + I*B)*a*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1155*(A + I*B)*a*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sqrt(a)/(c^(11/2)*f)
```

3.804.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{3/2}}{(-ictan(fx + e) + c)^{11/2}} dx$$

```
input integrate((a+I*a*tan(f*x+e))^(3/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(11/2),x, algorithm="giac")
```

```
output integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(3/2)/(-I*c*tan(f*x + e) + c)^(11/2), x)
```

3.804.9 Mupad [B] (verification not implemented)

Time = 12.51 (sec) , antiderivative size = 315, normalized size of antiderivative = 1.21

$$\int \frac{(a + ia \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx =$$

$$a \sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}} (A \cos(2e + 2fx) 1155i + A \cos(4e + 4fx) 2772i + A \cos(6e + 6fx) 2970i + A \cos(8e + 8fx) 1540i + A \cos(10e + 10fx) 315i - 1155B \cos(2e + 2fx) - 1386B \cos(4e + 4fx) + 770B \cos(8e + 8fx) + 315B \cos(10e + 10fx) - 1155A \sin(2e + 2fx) - 2772A \sin(4e + 4fx) - 2970A \sin(6e + 6fx) - 1540A \sin(8e + 8fx) - 315A \sin(10e + 10fx) - B \sin(2e + 2fx) 1155i - B \sin(4e + 4fx) 1386i + B \sin(8e + 8fx) 770i + B \sin(10e + 10fx) 315i) / (55440c^5f((c(\cos(2e + 2fx) - \sin(2e + 2fx)1i + 1)) / (\cos(2e + 2fx) + 1))^{1/2})$$

```
input int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(3/2))/(c - c*tan(e + f*x)*1i)^(11/2),x)
```

```
output -(a*((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*cos(2*e + 2*f*x)*1155i + A*cos(4*e + 4*f*x)*2772i + A*cos(6*e + 6*f*x)*2970i + A*cos(8*e + 8*f*x)*1540i + A*cos(10*e + 10*f*x)*315i - 1155*B*cos(2*e + 2*f*x) - 1386*B*cos(4*e + 4*f*x) + 770*B*cos(8*e + 8*f*x) + 315*B*cos(10*e + 10*f*x) - 1155*A*sin(2*e + 2*f*x) - 2772*A*sin(4*e + 4*f*x) - 2970*A*sin(6*e + 6*f*x) - 1540*A*sin(8*e + 8*f*x) - 315*A*sin(10*e + 10*f*x) - B*sin(2*e + 2*f*x)*1155i - B*sin(4*e + 4*f*x)*1386i + B*sin(8*e + 8*f*x)*770i + B*sin(10*e + 10*f*x)*315i))/(55440*c^5*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))
```

3.805 $\int (a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{7/2} dx$

3.805.1 Optimal result	7257
3.805.2 Mathematica [A] (verified)	7258
3.805.3 Rubi [A] (verified)	7258
3.805.4 Maple [B] (verified)	7261
3.805.5 Fricas [B] (verification not implemented)	7262
3.805.6 Sympy [F(-1)]	7263
3.805.7 Maxima [B] (verification not implemented)	7263
3.805.8 Giac [F(-1)]	7264
3.805.9 Mupad [F(-1)]	7265

3.805.1 Optimal result

Integrand size = 45, antiderivative size = 288

$$\int (a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))(c - ictan(e + fx))^{7/2} dx =$$

$$\frac{a^{5/2}(6iA - B)c^{7/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{8f}$$

$$+ \frac{a^2(6A + iB)c^3 \tan(e + fx)\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}}{16f}$$

$$+ \frac{a(6A + iB)c^2 \tan(e + fx)(a + ia \tan(e + fx))^{3/2}(c - ictan(e + fx))^{3/2}}{24f}$$

$$- \frac{(6iA - B)c(a + ia \tan(e + fx))^{5/2}(c - ictan(e + fx))^{5/2}}{30f}$$

$$+ \frac{B(a + ia \tan(e + fx))^{5/2}(c - ictan(e + fx))^{7/2}}{6f}$$

output

```
-1/8*a^(5/2)*(6*I*A-B)*c^(7/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/f+1/16*a^2*(6*A+I*B)*c^3*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)*tan(f*x+e)/f+1/24*a*(6*A+I*B)*c^2*tan(f*x+e)*(a+I*a*tan(f*x+e))^(3/2)*(c-I*c*tan(f*x+e))^(3/2)/f-1/30*(6*I*A-B)*c*(a+I*a*tan(f*x+e))^(5/2)*(c-I*c*tan(f*x+e))^(5/2)/f+1/6*B*(a+I*a*tan(f*x+e))^(5/2)*(c-I*c*tan(f*x+e))^(7/2)/f
```


3.805.2 Mathematica [A] (verified)

Time = 11.11 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.80

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \frac{a^{5/2} c^4 \sqrt{1 - i \tan(e + fx)} \left(30(-6iA + B) \arcsin \left(\frac{\sqrt{a + ia \tan(e + fx)}}{\sqrt{2}\sqrt{a}} \right) \sqrt{a + ia \tan(e + fx)} \right)}{\dots}$$

input `Integrate[(a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2),x]`

output `(a^(5/2)*c^4*Sqrt[1 - I*Tan[e + f*x]]*(30*((-6*I)*A + B)*ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a])]*Sqrt[a + I*a*Tan[e + f*x]] + (Sqrt[a]*Sec[e + f*x]^4*Sqrt[1 - I*Tan[e + f*x]]*(1 + I*Tan[e + f*x])*(384*((-I)*A + B) + 5*(102*A - (47*I)*B + 20*(6*A + I*B)*Cos[2*(e + f*x)] + 3*(6*A + I*B)*Cos[4*(e + f*x)])*Tan[e + f*x]))/8)/(240*f*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])`

3.805.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 267, normalized size of antiderivative = 0.93, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.178$, Rules used = {3042, 4071, 90, 59, 40, 40, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{7/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{7/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int (i \tan(e + fx)a + a)^{3/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{90} \end{aligned}$$

$$\frac{ac\left(\frac{1}{6}(6A + iB) \int (i \tan(e + fx)a + a)^{3/2}(c - ictan(e + fx))^{5/2}d \tan(e + fx) + \frac{B(a+ia \tan(e+fx))^{5/2}(c-ictan(e+fx))^{7/2}}{6ac}\right)}{f}$$

↓ 59

$$\frac{ac\left(\frac{1}{6}(6A + iB) \left(c \int (i \tan(e + fx)a + a)^{3/2}(c - ictan(e + fx))^{3/2}d \tan(e + fx) - \frac{i(a+ia \tan(e+fx))^{5/2}(c-ictan(e+fx))}{5a}\right)\right)}{f}$$

↓ 40

$$\frac{ac\left(\frac{1}{6}(6A + iB) \left(c\left(\frac{3}{4}ac \int \sqrt{i \tan(e + fx)a + a}\sqrt{c - ictan(e + fx)}d \tan(e + fx) + \frac{1}{4} \tan(e + fx)(a + ia \tan(e + fx))\right)\right)\right)}{f}$$

↓ 40

$$\frac{ac\left(\frac{1}{6}(6A + iB) \left(c\left(\frac{3}{4}ac\left(\frac{1}{2}ac \int \frac{1}{\sqrt{i \tan(e+fx)a+a}\sqrt{c-ictan(e+fx)}}d \tan(e + fx) + \frac{1}{2} \tan(e + fx)\sqrt{a + ia \tan(e + fx)}\right)\right)\right)\right)}{f}$$

↓ 45

$$\frac{ac\left(\frac{1}{6}(6A + iB) \left(c\left(\frac{3}{4}ac\left(ac \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ictan(e+fx)}}d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ictan(e+fx)}} + \frac{1}{2} \tan(e + fx)\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}\right)\right)\right)\right)}{f}$$

↓ 218

$$\frac{ac\left(\frac{1}{6}(6A + iB) \left(c\left(\frac{3}{4}ac\left(\frac{1}{2} \tan(e + fx)\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)} - i\sqrt{a}\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)\right)\right)\right)\right)}{f}$$

input `Int[(a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2),x]`

output `(a*c*((B*(a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e + f*x])^(7/2))/(6*a*c) + ((6*A + I*B)*(((-1/5*I)*(a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e + f*x])^(5/2))/a + c*((Tan[e + f*x]*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(3/2))/4 + (3*a*c*((-I)*Sqrt[a]*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])] + (Tan[e + f*x]*Sqrt[a + I*a*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])/2))/4))/6))/f`

3.805.3.1 Defintions of rubi rules used

- rule 40 $\text{Int}[(a + b \cdot x)^m \cdot (c + d \cdot x)^m, x_Symbol] \rightarrow \text{Simp}[x \cdot (a + b \cdot x)^m \cdot (c + d \cdot x)^m / (2 \cdot m + 1), x] + \text{Simp}[2 \cdot a \cdot c \cdot m / (2 \cdot m + 1) \text{Int}[(a + b \cdot x)^{m-1} \cdot (c + d \cdot x)^{m-1}, x], x] /;$ FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]
- rule 45 $\text{Int}[1/(\text{Sqrt}[a + b \cdot x] \cdot \text{Sqrt}[c + d \cdot x]), x_Symbol] \rightarrow \text{Simp}[2 \text{Subst}[\text{Int}[1/(b - d \cdot x^2), x], x, \text{Sqrt}[a + b \cdot x] / \text{Sqrt}[c + d \cdot x]], x] /;$ FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]
- rule 59 $\text{Int}[(a + b \cdot x)^m \cdot (c + d \cdot x)^n, x_Symbol] \rightarrow \text{Simp}[(a + b \cdot x)^{m+1} \cdot (c + d \cdot x)^n / (b \cdot (m + n + 1)), x] + \text{Simp}[2 \cdot c \cdot n / (m + n + 1) \text{Int}[(a + b \cdot x)^m \cdot (c + d \cdot x)^{n-1}, x], x] /;$ FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0] && IGtQ[n + 1/2, 0] && LtQ[m, n]
- rule 90 $\text{Int}[(a + b \cdot x)^n \cdot (c + d \cdot x)^p \cdot (e + f \cdot x)^q, x] \rightarrow \text{Simp}[b \cdot (c + d \cdot x)^{n+1} \cdot (e + f \cdot x)^q / (d \cdot f \cdot (n + p + 2)), x] + \text{Simp}[(a \cdot d \cdot f \cdot (n + p + 2) - b \cdot (d \cdot e \cdot (n + 1) + c \cdot f \cdot (p + 1))) / (d \cdot f \cdot (n + p + 2)) \text{Int}[(c + d \cdot x)^n \cdot (e + f \cdot x)^q, x], x] /;$ FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]
- rule 218 $\text{Int}[(a + b \cdot x)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a) \cdot \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /;$ FreeQ[{a, b}, x] && PosQ[a/b]
- rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$ FunctionOfTrigOfLinearQ[u, x]
- rule 4071 $\text{Int}[(a + b \cdot \tan[e + f \cdot x])^m \cdot (c + d \cdot \tan[e + f \cdot x])^n, x_Symbol] \rightarrow \text{Simp}[a \cdot (c/f) \text{Subst}[\text{Int}[(a + b \cdot x)^{m-1} \cdot (c + d \cdot x)^{n-1} \cdot (A + B \cdot x), x], x, \text{Tan}[e + f \cdot x]], x] /;$ FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]

3.805.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 477 vs. 2(236) = 472.

Time = 0.48 (sec) , antiderivative size = 478, normalized size of antiderivative = 1.66

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^2 c^3 \left(40iB \sqrt{ac(1+\tan(fx+e)^2)} \sqrt{ac} \tan(fx+e)^5 + 48iA \sqrt{ac(1+\tan(fx+e)^2)} \right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^2 c^3 \left(40iB \sqrt{ac(1+\tan(fx+e)^2)} \sqrt{ac} \tan(fx+e)^5 + 48iA \sqrt{ac(1+\tan(fx+e)^2)} \right)}{\dots}$
parts	$\frac{A \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^2 c^3 \left(8i \tan(fx+e)^4 \sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} + 16i \tan(fx+e)^2 \sqrt{ac} \right)}{\dots}$

```
input int((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x,method=_RETURNVERBOSE)
```

```
output -1/240/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^2*c^3*(40*I*B*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)*tan(f*x+e)^5+48*I*A*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)*tan(f*x+e)^4+70*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^3-48*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2-60*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^3-15*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)))^(1/2))/(a*c)^(1/2))*a*c+15*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-96*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+48*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)-90*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)))^(1/2))/(a*c)^(1/2))*a*c-150*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-48*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2)/(a*c*(1+tan(f*x+e)^2))^(1/2)
```

3.805. $\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{7/2} dx$

3.805.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 756 vs. $2(220) = 440$.

Time = 0.28 (sec) , antiderivative size = 756, normalized size of antiderivative = 2.62

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \frac{15 \sqrt{\frac{(36A^2 + 12iAB - B^2)a^5c^7}{f^2}} (fe^{(10ifx+10ie)} + 5fe^{(8ifx+8ie)} + 10fe^{(6ifx+6ie)} + 10fe^{(4ifx+4ie)} - ic \tan(e + fx))^{7/2}}{\dots}$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="fracas")`

output `1/480*(15*sqrt((36*A^2 + 12*I*A*B - B^2)*a^5*c^7/f^2)*(f*e^(10*I*f*x + 10*I*e) + 5*f*e^(8*I*f*x + 8*I*e) + 10*f*e^(6*I*f*x + 6*I*e) + 10*f*e^(4*I*f*x + 4*I*e) + 5*f*e^(2*I*f*x + 2*I*e) + f)*log(-4*(2*((6*I*A - B)*a^2*c^3*e^(3*I*f*x + 3*I*e) + (6*I*A - B)*a^2*c^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + sqrt((36*A^2 + 12*I*A*B - B^2)*a^5*c^7/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((-6*I*A + B)*a^2*c^3*e^(2*I*f*x + 2*I*e) + (-6*I*A + B)*a^2*c^3) - 15*sqrt((36*A^2 + 12*I*A*B - B^2)*a^5*c^7/f^2)*(f*e^(10*I*f*x + 10*I*e) + 5*f*e^(8*I*f*x + 8*I*e) + 10*f*e^(6*I*f*x + 6*I*e) + 10*f*e^(4*I*f*x + 4*I*e) + 5*f*e^(2*I*f*x + 2*I*e) + f)*log(-4*(2*((6*I*A - B)*a^2*c^3*e^(3*I*f*x + 3*I*e) + (6*I*A - B)*a^2*c^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))) - sqrt((36*A^2 + 12*I*A*B - B^2)*a^5*c^7/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((-6*I*A + B)*a^2*c^3*e^(2*I*f*x + 2*I*e) + (-6*I*A + B)*a^2*c^3) - 4*(15*(6*I*A - B)*a^2*c^3*e^(11*I*f*x + 11*I*e) + 85*(6*I*A - B)*a^2*c^3*e^(9*I*f*x + 9*I*e) + 198*(6*I*A - B)*a^2*c^3*e^(7*I*f*x + 7*I*e) + 6*(58*I*A - 223*B)*a^2*c^3*e^(5*I*f*x + 5*I*e) + 85*(-6*I*A + B)*a^2*c^3*e^(3*I*f*x + 3*I*e) + 15*(-6*I*A + B)*a^2*c^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(f*e^(10*I*f*x + 10*I*e) + 5*f*e^(8*I*f*x + 8*I*e) + 10*f*e^(6*I*f*x + 6*I*e) + 10*f*e^(4*I*f*x + 4*I*e) + 5*f*e^(2*I*f*x + 2*I*e) + f)`

3.805.6 Sympy [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(7/2),x)`

output `Timed out`

3.805.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2033 vs. $2(220) = 440$.

Time = 4.58 (sec) , antiderivative size = 2033, normalized size of antiderivative = 7.06

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \text{Too large to display}$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="maxima")`

output

```
-3840*(60*(6*A + I*B)*a^2*c^3*cos(11/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 340*(6*A + I*B)*a^2*c^3*cos(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 792*(6*A + I*B)*a^2*c^3*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 24*(58*A + 223*I*B)*a^2*c^3*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 340*(6*A + I*B)*a^2*c^3*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 60*(6*A + I*B)*a^2*c^3*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 60*(6*I*A - B)*a^2*c^3*sin(11/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 340*(6*I*A - B)*a^2*c^3*sin(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 792*(6*I*A - B)*a^2*c^3*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 24*(58*I*A - 223*B)*a^2*c^3*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 340*(-6*I*A + B)*a^2*c^3*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 60*(-6*I*A + B)*a^2*c^3*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 30*((6*A + I*B)*a^2*c^3*cos(12*f*x + 12*e) + 6*(6*A + I*B)*a^2*c^3*cos(10*f*x + 10*e) + 15*(6*A + I*B)*a^2*c^3*cos(8*f*x + 8*e) + 20*(6*A + I*B)*a^2*c^3*cos(6*f*x + 6*e) + 15*(6*A + I*B)*a^2*c^3*cos(4*f*x + 4*e) + 6*(6*A + I*B)*a^2*c^3*cos(2*f*x + 2*e) + (6*I*A - B)*a^2*c^3*sin(12*f*x + 12*e) + 6*(6*I*A - B)*a^2*c^3*sin(10*f*x + 10*e) + 15*(6*I*A - B)*a^2*c^3*sin(8*f*x + 8*e) + 20*(6*I*A - B)*a^2*c^3*sin(6*f*x + 6*e) + 15*(6*I*A - B)*a^2*c^3*sin(4*f*x + 4*e) + 6*(6*I*A - B)*a^2*c^3*sin(2*f*x + 2*e) + (6*A + I...
```

3.805.8 Giac [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{7/2} dx = \text{Timed out}$$

input

```
integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="giac")
```

output

```
Timed out
```

3.805.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{7/2} dx = \int (A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{5/2} (c - c \tan(e + fx) li)^{7/2} dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(5/2)*(c - c*tan(e + f*x)*1i)^(7/2),x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(5/2)*(c - c*tan(e + f*x)*1i)^(7/2), x)`

3.806 $\int (a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{5/2} dx$

3.806.1 Optimal result	7266
3.806.2 Mathematica [A] (verified)	7267
3.806.3 Rubi [A] (verified)	7267
3.806.4 Maple [A] (verified)	7270
3.806.5 Fricas [B] (verification not implemented)	7270
3.806.6 Sympy [F(-1)]	7271
3.806.7 Maxima [B] (verification not implemented)	7272
3.806.8 Giac [F(-1)]	7273
3.806.9 Mupad [F(-1)]	7273

3.806.1 Optimal result

Integrand size = 45, antiderivative size = 213

$$\int (a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))(c - ictan(e + fx))^{5/2} dx =$$

$$\frac{3ia^{5/2}Ac^{5/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{4f}$$

$$+ \frac{3a^2Ac^2 \tan(e + fx)\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}}{8f}$$

$$+ \frac{aActan(e + fx)(a + ia \tan(e + fx))^{3/2}(c - ictan(e + fx))^{3/2}}{4f}$$

$$+ \frac{B(a + ia \tan(e + fx))^{5/2}(c - ictan(e + fx))^{5/2}}{5f}$$

output

```
-3/4*I*a^(5/2)*A*c^(5/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/f+3/8*a^2*A*c^2*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)*tan(f*x+e)/f+1/4*a*A*c*tan(f*x+e)*(a+I*a*tan(f*x+e))^(3/2)*(c-I*c*tan(f*x+e))^(3/2)/f+1/5*B*(a+I*a*tan(f*x+e))^(5/2)*(c-I*c*tan(f*x+e))^(5/2)/f
```

3.806.2 Mathematica [A] (verified)

Time = 8.53 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.69

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx = \frac{a^{5/2} c^3 \left(-\frac{240A \arcsin\left(\frac{\sqrt{a+ia \tan(e+fx)}}{\sqrt{2}\sqrt{a}}\right) (i+\tan(e+fx))}{\sqrt{1-i \tan(e+fx)}} + \frac{\sqrt{a} \sec^6(e+fx) (64B+70A \sin(2(e+fx))+15A \sin(4(e+fx)))}{\sqrt{a+ia \tan(e+fx)}} \right)}{320f \sqrt{c - ic \tan(e + fx)}}$$

input `Integrate[(a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2),x]`

output `(a^(5/2)*c^3*((-240*A*ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a])]*(I + Tan[e + f*x]))/Sqrt[1 - I*Tan[e + f*x]] + (Sqrt[a]*Sec[e + f*x]^6*(64*B + 70*A*Sin[2*(e + f*x)] + 15*A*Sin[4*(e + f*x)]))/Sqrt[a + I*a*Tan[e + f*x]]))/(320*f*Sqrt[c - I*c*Tan[e + f*x]])`

3.806.3 Rubi [A] (verified)Time = 0.41 (sec) , antiderivative size = 210, normalized size of antiderivative = 0.99, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 90, 40, 40, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{5/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{5/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int (i \tan(e + fx) a + a)^{3/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{90} \end{aligned}$$

$$\frac{ac \left(A \int (i \tan(e + fx)a + a)^{3/2} (c - ic \tan(e + fx))^{3/2} d \tan(e + fx) + \frac{B(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{5/2}}{5ac} \right)}{f}$$

↓ 40

$$\frac{ac \left(A \left(\frac{3}{4} ac \int \sqrt{i \tan(e + fx)a + a} \sqrt{c - ic \tan(e + fx)} d \tan(e + fx) + \frac{1}{4} \tan(e + fx) (a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2} \right) \right)}{f}$$

↓ 40

$$\frac{ac \left(A \left(\frac{3}{4} ac \left(\frac{1}{2} ac \int \frac{1}{\sqrt{i \tan(e + fx)a + a} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)} \right) \right) \right)}{f}$$

↓ 45

$$\frac{ac \left(A \left(\frac{3}{4} ac \left(ac \int \frac{1}{ia + \frac{ic(i \tan(e + fx)a + a)}{c - ic \tan(e + fx)}} d \frac{\sqrt{i \tan(e + fx)a + a}}{\sqrt{c - ic \tan(e + fx)}} + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)} \right) \right) \right)}{f}$$

↓ 218

$$\frac{ac \left(A \left(\frac{3}{4} ac \left(\frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)} - i \sqrt{a} \sqrt{c} \arctan \left(\frac{\sqrt{c} \sqrt{a + ia \tan(e + fx)}}{\sqrt{a} \sqrt{c - ic \tan(e + fx)}} \right) \right) \right) \right) + \frac{1}{4} \tan(e + fx)}{f}$$

input `Int[(a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2),x]`

output `(a*c*((B*(a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e + f*x])^(5/2))/(5*a*c) + A*((Tan[e + f*x]*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(3/2))/4 + (3*a*c*((-I)*Sqrt[a]*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])] + (Tan[e + f*x]*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/2))/4))/f`

3.806.3.1 Defintions of rubi rules used

- rule 40 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^m/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`
- rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.806.4 Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.04

method	result
parts	$\frac{A\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^2c^2\left(2\tan(fx+e)^3\sqrt{ac(1+\tan(fx+e)^2)}\sqrt{ac+3ac}\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac(1+\tan(fx+e)^2)}}{8f\sqrt{ac(1+\tan(fx+e)^2)}\sqrt{ac}}\right)\right)}{8f\sqrt{ac(1+\tan(fx+e)^2)}\sqrt{ac}}$
derivativelimit	$\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^2c^2\left(8B\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}\tan(fx+e)^4+10A\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}\right)$
default	$\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^2c^2\left(8B\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}\tan(fx+e)^4+10A\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}\right)$

input `int((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

output `1/8*A/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^2*c^2*(2*tan(f*x+e)^3*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)+3*a*c*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))+5*tan(f*x+e)*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2))/(a*c*(1+tan(f*x+e)^2))^(1/2)/(a*c)^(1/2)+1/5*B/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^2*c^2*(1+tan(f*x+e)^2)^2`

3.806.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 585 vs. 2(163) = 326.

Time = 0.28 (sec) , antiderivative size = 585, normalized size of antiderivative = 2.75

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2} dx = \frac{15 \sqrt{\frac{A^2 a^5 c^5}{f^2}} (f e^{(8i fx + 8i e)} + 4 f e^{(6i fx + 6i e)} + 6 f e^{(4i fx + 4i e)} + 4 f e^{(2i fx + 2i e)} + f) \log \left(\frac{a + ia \tan(e + fx)}{c - ic \tan(e + fx)} \right)}{15 \sqrt{\frac{A^2 a^5 c^5}{f^2}}}$$

```
input integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="fricas")
```

```
output 1/80*(15*sqrt(A^2*a^5*c^5/f^2)*(f*e^(8*I*f*x + 8*I*e) + 4*f*e^(6*I*f*x + 6*I*e) + 6*f*e^(4*I*f*x + 4*I*e) + 4*f*e^(2*I*f*x + 2*I*e) + f)*log(4*(2*(A*a^2*c^2*e^(3*I*f*x + 3*I*e) + A*a^2*c^2*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - sqrt(A^2*a^5*c^5/f^2)*(I*f*e^(2*I*f*x + 2*I*e) - I*f))/(A*a^2*c^2*e^(2*I*f*x + 2*I*e) + A*a^2*c^2) - 15*sqrt(A^2*a^5*c^5/f^2)*(f*e^(8*I*f*x + 8*I*e) + 4*f*e^(6*I*f*x + 6*I*e) + 6*f*e^(4*I*f*x + 4*I*e) + 4*f*e^(2*I*f*x + 2*I*e) + f)*log(4*(2*(A*a^2*c^2*e^(3*I*f*x + 3*I*e) + A*a^2*c^2*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - sqrt(A^2*a^5*c^5/f^2)*(-I*f*e^(2*I*f*x + 2*I*e) + I*f))/(A*a^2*c^2*e^(2*I*f*x + 2*I*e) + A*a^2*c^2) + 4*(-15*I*A*a^2*c^2*e^(9*I*f*x + 9*I*e) - 70*I*A*a^2*c^2*e^(7*I*f*x + 7*I*e) + 128*B*a^2*c^2*e^(5*I*f*x + 5*I*e) + 70*I*A*a^2*c^2*e^(3*I*f*x + 3*I*e) + 15*I*A*a^2*c^2*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(f*e^(8*I*f*x + 8*I*e) + 4*f*e^(6*I*f*x + 6*I*e) + 6*f*e^(4*I*f*x + 4*I*e) + 4*f*e^(2*I*f*x + 2*I*e) + f)
```

3.806.6 Sympy [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx = \text{Timed out}$$

```
input integrate((a+I*a*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(5/2),x)
```

```
output Timed out
```

3.806.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1445 vs. $2(163) = 326$.

Time = 0.94 (sec) , antiderivative size = 1445, normalized size of antiderivative = 6.78

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx = \text{Too large to display}$$

```
input integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="maxima")
```

```
output -(60*A*a^2*c^2*cos(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 280*A*a^2*c^2*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 512*I*B*a^2*c^2*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 280*A*a^2*c^2*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 60*A*a^2*c^2*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 60*I*A*a^2*c^2*sin(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 280*I*A*a^2*c^2*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 512*B*a^2*c^2*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 280*I*A*a^2*c^2*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 60*I*A*a^2*c^2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 30*(A*a^2*c^2*cos(10*f*x + 10*e) + 5*A*a^2*c^2*cos(8*f*x + 8*e) + 10*A*a^2*c^2*cos(6*f*x + 6*e) + 10*A*a^2*c^2*cos(4*f*x + 4*e) + 5*A*a^2*c^2*cos(2*f*x + 2*e) + I*A*a^2*c^2*sin(10*f*x + 10*e) + 5*I*A*a^2*c^2*sin(8*f*x + 8*e) + 10*I*A*a^2*c^2*sin(6*f*x + 6*e) + 10*I*A*a^2*c^2*sin(4*f*x + 4*e) + 5*I*A*a^2*c^2*sin(2*f*x + 2*e) + A*a^2*c^2)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) + 30*(A*a^2*c^2*cos(10*f*x + 10*e) + 5*A*a^2*c^2*cos(8*f*x + 8*e) + 10*A*a^2*c^2*cos(6*f*x + 6*e) + 10*A*a^2*c^2*cos(4*f*x + 4*e) + 5*A*a^2*c^2*cos(2*f*x + 2*e) + I*A*a^2*c^2*sin(10*f*x + 10*e) + 5*I*A*a^2*c^2*sin(8*f*x + 8*e) + 10*I*A*a^2*c^2*sin(6*f*x + 6*e) + 10*I*A*a^2*c^2*sin(4*f*x + 4*e) + 5*I*A*a^2*c^2*sin(2*f*x...
```

3.806.8 Giac [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `Timed out`

3.806.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx = \int (A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{5/2} (c - c \tan(e + fx) li)^{5/2} dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*li)^(5/2)*(c - c*tan(e + f*x)*li)^(5/2),x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*li)^(5/2)*(c - c*tan(e + f*x)*li)^(5/2), x)`

3.807 $\int (a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{3/2} dx$

3.807.1 Optimal result	7274
3.807.2 Mathematica [A] (verified)	7275
3.807.3 Rubi [A] (verified)	7275
3.807.4 Maple [A] (verified)	7278
3.807.5 Fricas [B] (verification not implemented)	7279
3.807.6 Sympy [F(-1)]	7280
3.807.7 Maxima [B] (verification not implemented)	7280
3.807.8 Giac [F]	7281
3.807.9 Mupad [F(-1)]	7282

3.807.1 Optimal result

Integrand size = 45, antiderivative size = 222

$$\int (a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{3/2} dx =$$

$$\frac{a^{5/2}(4iA+B)c^{3/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{4f}$$

$$+ \frac{a^2(4A-iB)c \tan(e+fx) \sqrt{a+ia \tan(e+fx)} \sqrt{c-ictan(e+fx)}}{8f}$$

$$+ \frac{a(4iA+B)(a+ia \tan(e+fx))^{3/2}(c-ictan(e+fx))^{3/2}}{12f}$$

$$+ \frac{B(a+ia \tan(e+fx))^{5/2}(c-ictan(e+fx))^{3/2}}{4f}$$

output

```
-1/4*a^(5/2)*(4*I*A+B)*c^(3/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/f+1/8*a^2*(4*A-I*B)*c*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)*tan(f*x+e)/f+1/12*a*(4*I*A+B)*(a+I*a*tan(f*x+e))^(3/2)*(c-I*c*tan(f*x+e))^(3/2)/f+1/4*B*(a+I*a*tan(f*x+e))^(5/2)*(c-I*c*tan(f*x+e))^(3/2)/f
```

3.807.2 Mathematica [A] (verified)

Time = 7.93 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.00

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} dx = \frac{a^{5/2} c^2 (i + \tan(e + fx)) \left(-6(4A - iB) \arcsin \left(\frac{\sqrt{a + ia \tan(e + fx)}}{\sqrt{2}\sqrt{a}} \right) \sqrt{a + ia \tan(e + fx)} \right)}{24f\sqrt{a}}$$

input `Integrate[(a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(a^(5/2)*c^2*(I + Tan[e + f*x])*(-6*(4*A - I*B)*ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a]])*Sqrt[a + I*a*Tan[e + f*x]] + Sqrt[a]*Sqrt[1 - I*Tan[e + f*x]]*(-I + Tan[e + f*x])*(8*(I*A + B) + 3*(4*A + I*B)*Tan[e + f*x] + 8*(I*A + B)*Tan[e + f*x]^2 + (6*I)*B*Tan[e + f*x]^3))/(24*f*Sqrt[1 - I*Tan[e + f*x]]*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])`

3.807.3 Rubi [A] (verified)Time = 0.41 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.97, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 90, 59, 40, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int (i \tan(e + fx) a + a)^{3/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{90} \end{aligned}$$

$$\frac{ac\left(\frac{1}{4}(4A - iB) \int (i \tan(e + fx)a + a)^{3/2} \sqrt{c - ic \tan(e + fx)} d \tan(e + fx) + \frac{B(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2}}{4ac}\right)}{f}$$

↓ 59

$$\frac{ac\left(\frac{1}{4}(4A - iB) \left(a \int \sqrt{i \tan(e + fx)a + a} \sqrt{c - ic \tan(e + fx)} d \tan(e + fx) + \frac{i(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2}}{3c}\right)\right)}{f}$$

↓ 40

$$\frac{ac\left(\frac{1}{4}(4A - iB) \left(a \left(\frac{1}{2} ac \int \frac{1}{\sqrt{i \tan(e + fx)a + a} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}\right)\right)\right)}{f}$$

↓ 45

$$\frac{ac\left(\frac{1}{4}(4A - iB) \left(a \left(ac \int \frac{1}{ia + \frac{ic(i \tan(e + fx)a + a)}{c - ic \tan(e + fx)}} d \frac{\sqrt{i \tan(e + fx)a + a}}{\sqrt{c - ic \tan(e + fx)}} + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}\right)\right)\right)}{f}$$

↓ 218

$$\frac{ac\left(\frac{1}{4}(4A - iB) \left(a \left(\frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)} - i\sqrt{a}\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a + ia \tan(e + fx)}}{\sqrt{a}\sqrt{c - ic \tan(e + fx)}}\right)\right)\right)\right)}{f}$$

input `Int[(a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(a*c*((B*(a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e + f*x])^(3/2))/(4*a*c) + ((4*A - I*B)*(((I/3)*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(3/2))/c + a*((-I)*Sqrt[a]*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])]) + (Tan[e + f*x]*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/2))))/4)/f`

3.807.3.1 Defintions of rubi rules used

- rule 40 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^m/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`
- rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 59 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[2*c*(n/(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0] && IGtQ[n + 1/2, 0] && LtQ[m, n]`
- rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.807.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 350, normalized size of antiderivative = 1.58

method	result
derivativedivides	$\frac{\sqrt{a(1+i\tan(fx+e))} \sqrt{-c(i\tan(fx+e)-1)} a^2 c \left(6iB\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \tan(fx+e)^3 + 8iA\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \right)}{\dots}$
default	$\frac{\sqrt{a(1+i\tan(fx+e))} \sqrt{-c(i\tan(fx+e)-1)} a^2 c \left(6iB\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \tan(fx+e)^3 + 8iA\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \right)}{\dots}$
parts	$\frac{A\sqrt{a(1+i\tan(fx+e))} \sqrt{-c(i\tan(fx+e)-1)} a^2 c \left(2i \tan(fx+e)^2 \sqrt{ac(1+\tan(fx+e)^2)} \sqrt{ac+2i\sqrt{ac}} \sqrt{ac(1+\tan(fx+e)^2)} \right)}{6f\sqrt{ac(1+\tan(fx+e)^2)}}$

```
input int((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
```

```
output 1/24/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^2*c*(6*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^3+8*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2-3*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c+3*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+8*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+8*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)+12*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c+12*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+8*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2)/(a*c*(1+tan(f*x+e)^2))^(1/2)
```

3.807.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 614 vs. $2(168) = 336$.

Time = 0.29 (sec) , antiderivative size = 614, normalized size of antiderivative = 2.77

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} dx =$$

$$3 \sqrt{\frac{(16A^2 - 8iAB - B^2)a^5 c^3}{f^2}} (f e^{(6i fx + 6i e)} + 3 f e^{(4i fx + 4i e)} + 3 f e^{(2i fx + 2i e)} + f) \log \left(- \frac{4 \left(2 \left((-4iA - B) a^2 c e^{(3i fx + 3i e)} + \dots \right) \right)}{\dots} \right)$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output `-1/48*(3*sqrt((16*A^2 - 8*I*A*B - B^2)*a^5*c^3/f^2)*(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)*log(-4*(2*((-4*I*A - B)*a^2*c*e^(3*I*f*x + 3*I*e) + (-4*I*A - B)*a^2*c*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + sqrt((16*A^2 - 8*I*A*B - B^2)*a^5*c^3/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((4*I*A + B)*a^2*c*e^(2*I*f*x + 2*I*e) + (4*I*A + B)*a^2*c) - 3*sqrt((16*A^2 - 8*I*A*B - B^2)*a^5*c^3/f^2)*(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)*log(-4*(2*((-4*I*A - B)*a^2*c*e^(3*I*f*x + 3*I*e) + (-4*I*A - B)*a^2*c*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - sqrt((16*A^2 - 8*I*A*B - B^2)*a^5*c^3/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((4*I*A + B)*a^2*c*e^(2*I*f*x + 2*I*e) + (4*I*A + B)*a^2*c) + 4*(3*(4*I*A + B)*a^2*c*e^(7*I*f*x + 7*I*e) - (20*I*A + 53*B)*a^2*c*e^(5*I*f*x + 5*I*e) + 11*(-4*I*A - B)*a^2*c*e^(3*I*f*x + 3*I*e) + 3*(-4*I*A - B)*a^2*c*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)`

3.807.6 Sympy [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(3/2),x)`

output `Timed out`

3.807.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1375 vs. $2(168) = 336$.

Time = 1.10 (sec) , antiderivative size = 1375, normalized size of antiderivative = 6.19

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} dx = \text{Too large to display}$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output

```

-96*(12*(4*A - I*B)*a^2*c*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*
e))) - 4*(20*A - 53*I*B)*a^2*c*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x
+ 2*e))) - 44*(4*A - I*B)*a^2*c*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f
*x + 2*e))) - 12*(4*A - I*B)*a^2*c*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2
*f*x + 2*e))) - 12*(-4*I*A - B)*a^2*c*sin(7/2*arctan2(sin(2*f*x + 2*e), co
s(2*f*x + 2*e))) - 4*(20*I*A + 53*B)*a^2*c*sin(5/2*arctan2(sin(2*f*x + 2*e
), cos(2*f*x + 2*e))) - 44*(4*I*A + B)*a^2*c*sin(3/2*arctan2(sin(2*f*x + 2
*e), cos(2*f*x + 2*e))) - 12*(4*I*A + B)*a^2*c*sin(1/2*arctan2(sin(2*f*x +
2*e), cos(2*f*x + 2*e))) + 6*((4*A - I*B)*a^2*c*cos(8*f*x + 8*e) + 4*(4*A
- I*B)*a^2*c*cos(6*f*x + 6*e) + 6*(4*A - I*B)*a^2*c*cos(4*f*x + 4*e) + 4*
(4*A - I*B)*a^2*c*cos(2*f*x + 2*e) - (-4*I*A - B)*a^2*c*sin(8*f*x + 8*e) -
4*(-4*I*A - B)*a^2*c*sin(6*f*x + 6*e) - 6*(-4*I*A - B)*a^2*c*sin(4*f*x +
4*e) - 4*(-4*I*A - B)*a^2*c*sin(2*f*x + 2*e) + (4*A - I*B)*a^2*c)*arctan2(
cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(
2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) + 6*((4*A - I*B)*a^2*c*cos(8*f*x + 8
*e) + 4*(4*A - I*B)*a^2*c*cos(6*f*x + 6*e) + 6*(4*A - I*B)*a^2*c*cos(4*f*x
+ 4*e) + 4*(4*A - I*B)*a^2*c*cos(2*f*x + 2*e) - (-4*I*A - B)*a^2*c*sin(8*
f*x + 8*e) - 4*(-4*I*A - B)*a^2*c*sin(6*f*x + 6*e) - 6*(-4*I*A - B)*a^2*c*
sin(4*f*x + 4*e) - 4*(-4*I*A - B)*a^2*c*sin(2*f*x + 2*e) + (4*A - I*B)*a^2
*c)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), -sin(...

```

3.807.8 Giac [F]

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx = \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^{5/2} (-ictan(fx + e) + c)^{3/2} dx$$

input

```

integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/
2),x, algorithm="giac")

```

output

```

sage0*x

```


3.807.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx = \int (A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{5/2} (c - c \tan(e + fx) li)^{3/2} dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(5/2)*(c - c*tan(e + f*x)*1i)^(3/2), x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(5/2)*(c - c*tan(e + f*x)*1i)^(3/2), x)`

3.808 $\int (a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}$

3.808.1 Optimal result	7283
3.808.2 Mathematica [A] (verified)	7284
3.808.3 Rubi [A] (verified)	7284
3.808.4 Maple [A] (verified)	7287
3.808.5 Fracas [B] (verification not implemented)	7287
3.808.6 Sympy [F]	7288
3.808.7 Maxima [B] (verification not implemented)	7289
3.808.8 Giac [F]	7290
3.808.9 Mupad [F(-1)]	7290

3.808.1 Optimal result

Integrand size = 45, antiderivative size = 217

$$\int (a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))\sqrt{c - ic \tan(e + fx)} dx =$$

$$-\frac{a^{5/2}(3iA + 2B)\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{f}$$

$$+ \frac{a^2(3iA + 2B)\sqrt{a + ia \tan(e + fx)}\sqrt{c - ic \tan(e + fx)}}{2f}$$

$$+ \frac{a(3iA + 2B)(a + ia \tan(e + fx))^{3/2}\sqrt{c - ic \tan(e + fx)}}{6f}$$

$$+ \frac{B(a + ia \tan(e + fx))^{5/2}\sqrt{c - ic \tan(e + fx)}}{3f}$$

```
output -a^(5/2)*(3*I*A+2*B)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*
c*tan(f*x+e))^(1/2))*c^(1/2)/f+1/2*a^2*(3*I*A+2*B)*(a+I*a*tan(f*x+e))^(1/2
)*(c-I*c*tan(f*x+e))^(1/2)/f+1/6*a*(3*I*A+2*B)*(c-I*c*tan(f*x+e))^(1/2)*(a
+I*a*tan(f*x+e))^(3/2)/f+1/3*B*(c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e)
^(5/2)/f
```

3.808.2 Mathematica [A] (verified)

Time = 5.74 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.91

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx =$$

$$\frac{a^{5/2} c (i + \tan(e + fx)) \left(6(3A - 2iB) \arcsin \left(\frac{\sqrt{a + ia \tan(e + fx)}}{\sqrt{2}\sqrt{a}} \right) \sqrt{a + ia \tan(e + fx)} + \sqrt{a} \sqrt{1 - i \tan(e + fx)} \right)}{6f \sqrt{1 - i \tan(e + fx)} \sqrt{a + ia \tan(e + fx)}}$$

input `Integrate[(a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]],x]`

output `-1/6*(a^(5/2)*c*(I + Tan[e + f*x])*(6*(3*A - (2*I)*B)*ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a]])*Sqrt[a + I*a*Tan[e + f*x]] + Sqrt[a]*Sqrt[1 - I*Tan[e + f*x]]*(-I + Tan[e + f*x])*((-12*I)*A - 10*B + 3*(A - (2*I)*B)*Tan[e + f*x] + 2*B*Tan[e + f*x]^2))/(f*Sqrt[1 - I*Tan[e + f*x]]*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])`

3.808.3 Rubi [A] (verified)Time = 0.41 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 90, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))^{5/2} \sqrt{c - ic \tan(e + fx)} (A + B \tan(e + fx)) dx$$

$$\downarrow 3042$$

$$\int (a + ia \tan(e + fx))^{5/2} \sqrt{c - ic \tan(e + fx)} (A + B \tan(e + fx)) dx$$

$$\downarrow 4071$$

$$\frac{ac \int \frac{(i \tan(e + fx)a + a)^{3/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{f}$$

$$\downarrow 90$$

$$\begin{array}{c}
 \frac{ac\left(\frac{1}{3}(3A - 2iB) \int \frac{(i \tan(e+fx)a+a)^{3/2}}{\sqrt{c-ictan(e+fx)}} d \tan(e+fx) + \frac{B(a+ia \tan(e+fx))^{5/2} \sqrt{c-ictan(e+fx)}}{3ac}\right)}{f} \\
 \downarrow 60 \\
 \frac{ac\left(\frac{1}{3}(3A - 2iB) \left(\frac{3}{2}a \int \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ictan(e+fx)}} d \tan(e+fx) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ictan(e+fx)}}{2c}\right) + \frac{B(a+ia \tan(e+fx))^{5/2} \sqrt{c-ictan(e+fx)}}{3ac}\right)}{f} \\
 \downarrow 60 \\
 \frac{ac\left(\frac{1}{3}(3A - 2iB) \left(\frac{3}{2}a \left(a \int \frac{1}{\sqrt{i \tan(e+fx)a+a} \sqrt{c-ictan(e+fx)}} d \tan(e+fx) + \frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ictan(e+fx)}}{c}\right) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ictan(e+fx)}}{2c}\right) + \frac{B(a+ia \tan(e+fx))^{5/2} \sqrt{c-ictan(e+fx)}}{3ac}\right)}{f} \\
 \downarrow 45 \\
 \frac{ac\left(\frac{1}{3}(3A - 2iB) \left(\frac{3}{2}a \left(2a \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ictan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ictan(e+fx)}} + \frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ictan(e+fx)}}{c}\right) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ictan(e+fx)}}{2c}\right) + \frac{B(a+ia \tan(e+fx))^{5/2} \sqrt{c-ictan(e+fx)}}{3ac}\right)}{f} \\
 \downarrow 218 \\
 \frac{ac\left(\frac{1}{3}(3A - 2iB) \left(\frac{3}{2}a \left(\frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ictan(e+fx)}}{c} - \frac{2i \sqrt{a} \arctan\left(\frac{\sqrt{c} \sqrt{a+ia \tan(e+fx)}}{\sqrt{a} \sqrt{c-ictan(e+fx)}}\right)}{\sqrt{c}}\right) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ictan(e+fx)}}{2c}\right) + \frac{B(a+ia \tan(e+fx))^{5/2} \sqrt{c-ictan(e+fx)}}{3ac}\right)}{f}
 \end{array}$$

input `Int[(a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]],x]`

output `(a*c*((B*(a + I*a*Tan[e + f*x])^(5/2)*Sqrt[c - I*c*Tan[e + f*x]])/(3*a*c) + (((3*A - (2*I)*B)*(((I/2)*(a + I*a*Tan[e + f*x])^(3/2)*Sqrt[c - I*c*Tan[e + f*x]])/c + (3*a*(((2*I)*Sqrt[a]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[c] + (I*Sqrt[a + I*a*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]]/c))/2))/3)/f`

3.808.3.1 Defintions of rubi rules used

- rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 60 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(
b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]`
- rule 90 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
), x] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p
+ 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n,
p}, x] && NeQ[n + p + 2, 0]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`
- rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]`

3.808.4 Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.31

method	result
derivativedivides	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} a^2 \left(-6iB \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right) ac+6iB\sqrt{ac} \sqrt{ac(1+i \tan(fx+e))}}{\dots}$
default	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} a^2 \left(-6iB \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right) ac+6iB\sqrt{ac} \sqrt{ac(1+i \tan(fx+e))}}{\dots}$
parts	$\frac{A\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} a^2 \left(4i\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} - \tan(fx+e) \sqrt{ac(1+\tan(fx+e)^2)} \right) \sqrt{ac}}{2f\sqrt{ac(1+\tan(fx+e)^2)} \sqrt{ac}}$

```
input int((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 1/6/f*(-c*(I*tan(f*x+e)-1))^(1/2)*(a*(1+I*tan(f*x+e)))^(1/2)*a^2*(-6*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c+6*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-2*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+12*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)+9*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c-3*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+10*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2)/(a*c*(1+tan(f*x+e)^2))^(1/2)
```

3.808.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 547 vs. 2(165) = 330.

3.808. $\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx$

Time = 0.28 (sec) , antiderivative size = 547, normalized size of antiderivative = 2.52

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx =$$

$$3 \sqrt{\frac{(9A^2 - 12iAB - 4B^2)a^5c}{f^2}} (fe^{(4ifx + 4ie)} + 2fe^{(2ifx + 2ie)} + f) \log \left(- \frac{4 \left(2((-3iA - 2B)a^2e^{(3ifx + 3ie)} + (-3iA - 2B)a^2e^{(ifx + ie)} \right)}{\dots} \right)$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)),x, algorithm="fricas")`

output `-1/12*(3*sqrt((9*A^2 - 12*I*A*B - 4*B^2)*a^5*c/f^2))*(f*e^(4*I*f*x + 4*I*e) + 2*f*e^(2*I*f*x + 2*I*e) + f)*log(-4*(2*((-3*I*A - 2*B)*a^2*e^(3*I*f*x + 3*I*e) + (-3*I*A - 2*B)*a^2*e^(I*f*x + I*e)))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + sqrt((9*A^2 - 12*I*A*B - 4*B^2)*a^5*c/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((3*I*A + 2*B)*a^2*e^(2*I*f*x + 2*I*e) + (3*I*A + 2*B)*a^2)) - 3*sqrt((9*A^2 - 12*I*A*B - 4*B^2)*a^5*c/f^2)*(f*e^(4*I*f*x + 4*I*e) + 2*f*e^(2*I*f*x + 2*I*e) + f)*log(-4*(2*((-3*I*A - 2*B)*a^2*e^(3*I*f*x + 3*I*e) + (-3*I*A - 2*B)*a^2*e^(I*f*x + I*e)))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - sqrt((9*A^2 - 12*I*A*B - 4*B^2)*a^5*c/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((3*I*A + 2*B)*a^2*e^(2*I*f*x + 2*I*e) + (3*I*A + 2*B)*a^2)) + 4*(3*(-5*I*A - 6*B)*a^2*e^(5*I*f*x + 5*I*e) + 8*(-3*I*A - 2*B)*a^2*e^(3*I*f*x + 3*I*e) + 3*(-3*I*A - 2*B)*a^2*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(f*e^(4*I*f*x + 4*I*e) + 2*f*e^(2*I*f*x + 2*I*e) + f)`

3.808.6 Sympy [F]

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx = \int (ia(\tan(e + fx) - i))^{5/2} \sqrt{-ic(\tan(e + fx) + i)} (A + B \tan(e + fx)) dx$$

input `integrate((c-I*c*tan(f*x+e))**(1/2)*(a+I*a*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e)),x)`

3.808. $\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx$

output `Integral((I*a*(tan(e + f*x) - I))**(5/2)*sqrt(-I*c*(tan(e + f*x) + I))*(A + B*tan(e + f*x)), x)`

3.808.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1087 vs. $2(165) = 330$.

Time = 0.58 (sec) , antiderivative size = 1087, normalized size of antiderivative = 5.01

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx = \text{Too large to display}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)),x, algorithm="maxima")`

output `6*(12*(5*A - 6*I*B)*a^2*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 32*(3*A - 2*I*B)*a^2*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 12*(3*A - 2*I*B)*a^2*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 12*(5*I*A + 6*B)*a^2*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 32*(3*I*A + 2*B)*a^2*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 12*(3*I*A + 2*B)*a^2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 6*((3*A - 2*I*B)*a^2*cos(6*f*x + 6*e) + 3*(3*A - 2*I*B)*a^2*cos(4*f*x + 4*e) + 3*(3*A - 2*I*B)*a^2*cos(2*f*x + 2*e) - (-3*I*A - 2*B)*a^2*sin(6*f*x + 6*e) - 3*(-3*I*A - 2*B)*a^2*sin(4*f*x + 4*e) - 3*(-3*I*A - 2*B)*a^2*sin(2*f*x + 2*e) + (3*A - 2*I*B)*a^2)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) - 6*((3*A - 2*I*B)*a^2*cos(6*f*x + 6*e) + 3*(3*A - 2*I*B)*a^2*cos(4*f*x + 4*e) + 3*(3*A - 2*I*B)*a^2*cos(2*f*x + 2*e) - (-3*I*A - 2*B)*a^2*sin(6*f*x + 6*e) - 3*(-3*I*A - 2*B)*a^2*sin(4*f*x + 4*e) - 3*(-3*I*A - 2*B)*a^2*sin(2*f*x + 2*e) + (3*A - 2*I*B)*a^2)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), -sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) + 3*((-3*I*A - 2*B)*a^2*cos(6*f*x + 6*e) + 3*(-3*I*A - 2*B)*a^2*cos(4*f*x + 4*e) + 3*(-3*I*A - 2*B)*a^2*cos(2*f*x + 2*e) + (3*A - 2*I*B)*a^2*sin(6*f*x + 6*e) + 3*(3*A - 2*I*B)*a^2*sin(4*f*x + 4*e) + 3*(3*A - 2*I*B)*a^2*sin(2*f*x + 2*e) + (-3*I*A - 2*B)*a^2)*log(cos...`

3.808.8 Giac [F]

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx = \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^{5/2} \sqrt{-ic \tan(fx + e)}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e)),x, algorithm="giac")`

output `sage0*x`

3.808.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx = \int (A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{5/2} \sqrt{c - c \tan(e + fx)} li dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(5/2)*(c - c*tan(e + f*x)*1i)^(1/2),x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(5/2)*(c - c*tan(e + f*x)*1i)^(1/2), x)`

$$3.809 \quad \int \frac{(a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} dx$$

3.809.1 Optimal result	7291
3.809.2 Mathematica [A] (verified)	7292
3.809.3 Rubi [A] (verified)	7292
3.809.4 Maple [B] (verified)	7295
3.809.5 Fricas [B] (verification not implemented)	7296
3.809.6 Sympy [F]	7297
3.809.7 Maxima [B] (verification not implemented)	7297
3.809.8 Giac [F]	7298
3.809.9 Mupad [F(-1)]	7299

3.809.1 Optimal result

Integrand size = 45, antiderivative size = 227

$$\int \frac{(a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx = \frac{3a^{5/2}(2iA + 3B) \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{\sqrt{cf}}$$

$$- \frac{(iA + B)(a + ia \tan(e + fx))^{5/2}}{f\sqrt{c - ictan(e + fx)}}$$

$$- \frac{3a^2(2iA + 3B)\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}}{2cf}$$

$$- \frac{a(2iA + 3B)(a + ia \tan(e + fx))^{3/2}\sqrt{c - ictan(e + fx)}}{2cf}$$

```
output 3*a^(5/2)*(2*I*A+3*B)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I
*c*tan(f*x+e))^(1/2))/f/c^(1/2)-3/2*a^2*(2*I*A+3*B)*(a+I*a*tan(f*x+e))^(1/
2)*(c-I*c*tan(f*x+e))^(1/2)/c/f-1/2*a*(2*I*A+3*B)*(c-I*c*tan(f*x+e))^(1/2)
*(a+I*a*tan(f*x+e))^(3/2)/c/f-(I*A+B)*(a+I*a*tan(f*x+e))^(5/2)/f/(c-I*c*ta
n(f*x+e))^(1/2)
```

3.809.2 Mathematica [A] (verified)

Time = 7.33 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.75

$$\int \frac{(a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx = \frac{a^{5/2} \left(\frac{6(2A - 3iB) \arcsin\left(\frac{\sqrt{a + ia \tan(e + fx)}}{\sqrt{2}\sqrt{a}}\right)(i + \tan(e + fx))}{\sqrt{1 - i \tan(e + fx)}} - \frac{i\sqrt{a}(-i + \tan(e + fx))}{2f\sqrt{c - ictan(e + fx)}} \right)}{2f\sqrt{c - ictan(e + fx)}}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/Sqrt[c - I*c*Tan[e + f*x]],x]`

output `(a^(5/2)*((6*(2*A - (3*I)*B)*ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a]])*(I + Tan[e + f*x]))/Sqrt[1 - I*Tan[e + f*x]] - (I*Sqrt[a]*(-I + Tan[e + f*x])*(2*((5*I)*A + 7*B) + (2*A - (5*I)*B)*Tan[e + f*x] + B*Tan[e + f*x]^2))/Sqrt[a + I*a*Tan[e + f*x]]))/(2*f*Sqrt[c - I*c*Tan[e + f*x]])`

3.809.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.97, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 87, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(i \tan(e + fx)a + a)^{3/2}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{3/2}} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{87} \end{aligned}$$

$$ac \left(- \frac{(2A-3iB) \int \frac{(i \tan(e+fx)a+a)^{3/2}}{\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{c} - \frac{(B+iA)(a+ia \tan(e+fx))^{5/2}}{ac \sqrt{c-ic \tan(e+fx)}} \right)$$

f
↓ 60

$$ac \left(- \frac{(2A-3iB) \left(\frac{3}{2} a \int \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2c} \right)}{c} - \frac{(B+iA)(a+ia \tan(e+fx))^{5/2}}{ac \sqrt{c-ic \tan(e+fx)}} \right)$$

f
↓ 60

$$ac \left(- \frac{(2A-3iB) \left(\frac{3}{2} a \left(a \int \frac{1}{\sqrt{i \tan(e+fx)a+a} \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{c} \right) \right) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2c}}{c}$$

f

↓ 45

$$ac \left(- \frac{(2A-3iB) \left(\frac{3}{2} a \left(2a \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} + \frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{c} \right) \right) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2c}}{c}$$

f

↓ 218

$$ac \left(- \frac{(2A-3iB) \left(\frac{3}{2} a \left(\frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{c} - \frac{2i \sqrt{a} \arctan \left(\frac{\sqrt{c} \sqrt{a+ia \tan(e+fx)}}{\sqrt{a} \sqrt{c-ic \tan(e+fx)}} \right)}{\sqrt{c}} \right) \right) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2c}}{c} - \frac{(B+iA)(a+ia \tan(e+fx))^{5/2}}{ac \sqrt{c-ic \tan(e+fx)}} \right)$$

f

input `Int[((a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/Sqrt[c - I*c*Tan[e + f*x]],x]`

```
output (a*c*(-(((I*A + B)*(a + I*a*Tan[e + f*x])^(5/2))/(a*c*Sqrt[c - I*c*Tan[e +
f*x]])) - ((2*A - (3*I)*B)*(((I/2)*(a + I*a*Tan[e + f*x])^(3/2)*Sqrt[c -
I*c*Tan[e + f*x]])/c + (3*a*((( -2*I)*Sqrt[a]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*
Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[c] + (I*Sqrt[a
+ I*a*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]/c))/2))/c)/f
```

3.809.3.1 Defintions of rubi rules used

```
rule 45 Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]
```

```
rule 60 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(
b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && ( !Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```

```
rule 87 Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
_), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && ( !LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.809.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 564 vs. $2(187) = 374$.

Time = 0.42 (sec) , antiderivative size = 565, normalized size of antiderivative = 2.49

method	result
derivativedivides	$i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^2\left(6iA\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}}\right)ac\tan(fx+e)^2+18iB\right)$
default	$i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^2\left(6iA\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}}\right)ac\tan(fx+e)^2+18iB\right)$
parts	$iA\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^2\left(3i\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}}\right)\tan(fx+e)^2ac-3i\ln\left(\right)\right)$

```
input int((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x,m
ethod=_RETURNVERBOSE)
```

```
output 1/2*I/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^2/c*(6*I*
A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2)
)*a*c*tan(f*x+e)^2+18*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)
)^2))^(1/2))/(a*c)^(1/2)*a*c*tan(f*x+e)+4*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x
+e)^2))^(1/2)*tan(f*x+e)^2+9*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(
f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)^2-B*(a*c)^(1/2)*(a*c*(1+tan(
f*x+e)^2))^(1/2)*tan(f*x+e)^3-6*I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1
+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c-12*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x
+e)^2))^(1/2)*tan(f*x+e)-12*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f
*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c*tan(f*x+e)-2*A*(a*c*(1+tan(f*x+e)^2))^(1
/2)*(a*c)^(1/2)*tan(f*x+e)^2-14*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/
2)-9*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(
1/2))*a*c-19*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+10*A*(
a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c*(1+tan(f*x+e)^2))^(1/2)/(I+t
an(f*x+e))^2/(a*c)^(1/2)
```

3.809.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 530 vs. $2(175) = 350$.

Time = 0.28 (sec) , antiderivative size = 530, normalized size of antiderivative = 2.33

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx = \frac{3 \sqrt{\frac{(4A^2 - 12iAB - 9B^2)a^5}{cf^2}} (cfe^{(2i fx + 2ie)} + cf) \log \left(\frac{4 \left(2(- \right)}{\right)}{\right)}{\right)}$$

```
input integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/
2),x, algorithm="fracas")
```

output $\frac{1}{4} \cdot (3 \sqrt{(4A^2 - 12IAB - 9B^2)a^5/(cf^2)}) \cdot (cfe^{(2Ifx + 2Ie)} + cf) \cdot \log(4 \cdot (2 \cdot ((-2IA - 3B)a^2e^{(3Ifx + 3Ie)} + (-2IA - 3B)a^2e^{(Ifx + Ie)})) \cdot \sqrt{a/(e^{(2Ifx + 2Ie)} + 1)}) \cdot \sqrt{c/(e^{(2Ifx + 2Ie)} + 1)}) + \sqrt{(4A^2 - 12IAB - 9B^2)a^5/(cf^2)} \cdot (cfe^{(2Ifx + 2Ie)} - cf) / ((-2IA - 3B)a^2e^{(2Ifx + 2Ie)} + (-2IA - 3B)a^2) - 3 \sqrt{(4A^2 - 12IAB - 9B^2)a^5/(cf^2)} \cdot (cfe^{(2Ifx + 2Ie)} + cf) \cdot \log(4 \cdot (2 \cdot ((-2IA - 3B)a^2e^{(3Ifx + 3Ie)} + (-2IA - 3B)a^2e^{(Ifx + Ie)})) \cdot \sqrt{a/(e^{(2Ifx + 2Ie)} + 1)}) \cdot \sqrt{c/(e^{(2Ifx + 2Ie)} + 1)}) - \sqrt{(4A^2 - 12IAB - 9B^2)a^5/(cf^2)} \cdot (cfe^{(2Ifx + 2Ie)} - cf) / ((-2IA - 3B)a^2e^{(2Ifx + 2Ie)} + (-2IA - 3B)a^2) - 4 \cdot (4 \cdot (IA + B)a^2e^{(5Ifx + 5Ie)} + 5 \cdot (2IA + 3B)a^2e^{(3Ifx + 3Ie)} + 3 \cdot (2IA + 3B)a^2e^{(Ifx + Ie)}) \cdot \sqrt{a/(e^{(2Ifx + 2Ie)} + 1)}) \cdot \sqrt{c/(e^{(2Ifx + 2Ie)} + 1)}) / (cfe^{(2Ifx + 2Ie)} + cf)$

3.809.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx = \int \frac{(ia(\tan(e + fx) - i))^{5/2} (A + B \tan(e + fx))}{\sqrt{-ic(\tan(e + fx) + i)}} dx$$

input `integrate((a+I*a*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(1/2),x)`

output `Integral((I*a*(tan(e + f*x) - I))**(5/2)*(A + B*tan(e + f*x))/sqrt(-I*c*(tan(e + f*x) + I)), x)`

3.809.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 993 vs. $2(175) = 350$.

Time = 0.51 (sec) , antiderivative size = 993, normalized size of antiderivative = 4.37

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx = \text{Too large to display}$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x, algorithm="maxima")`

3.809. $\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx$

output

```

-4*(4*(2*A - 7*I*B)*a^2*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))
)) + 4*(2*I*A + 7*B)*a^2*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e
))) - 6*((2*A - 3*I*B)*a^2*cos(4*f*x + 4*e) + 2*(2*A - 3*I*B)*a^2*cos(2*f*
x + 2*e) - (-2*I*A - 3*B)*a^2*sin(4*f*x + 4*e) - 2*(-2*I*A - 3*B)*a^2*sin(
2*f*x + 2*e) + (2*A - 3*I*B)*a^2)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e)
, cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))
+ 1) - 6*((2*A - 3*I*B)*a^2*cos(4*f*x + 4*e) + 2*(2*A - 3*I*B)*a^2*cos(2*
f*x + 2*e) - (-2*I*A - 3*B)*a^2*sin(4*f*x + 4*e) - 2*(-2*I*A - 3*B)*a^2*si
n(2*f*x + 2*e) + (2*A - 3*I*B)*a^2)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*
e), cos(2*f*x + 2*e))), -sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e
))) + 1) + 4*(4*(A - I*B)*a^2*cos(4*f*x + 4*e) + 8*(A - I*B)*a^2*cos(2*f*x
+ 2*e) + 4*(I*A + B)*a^2*sin(4*f*x + 4*e) + 8*(I*A + B)*a^2*sin(2*f*x + 2
*e) + 3*(2*A - 3*I*B)*a^2)*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2
*e))) + 3*((-2*I*A - 3*B)*a^2*cos(4*f*x + 4*e) + 2*(-2*I*A - 3*B)*a^2*cos(
2*f*x + 2*e) + (2*A - 3*I*B)*a^2*sin(4*f*x + 4*e) + 2*(2*A - 3*I*B)*a^2*si
n(2*f*x + 2*e) + (-2*I*A - 3*B)*a^2)*log(cos(1/2*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e)))^2 + sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)
))^2 + 2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) + 3*((2
*I*A + 3*B)*a^2*cos(4*f*x + 4*e) + 2*(2*I*A + 3*B)*a^2*cos(2*f*x + 2*e) -
(2*A - 3*I*B)*a^2*sin(4*f*x + 4*e) - 2*(2*A - 3*I*B)*a^2*sin(2*f*x + 2*...

```

3.809.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{5/2}}{\sqrt{-ic \tan(fx + e) + c}} dx$$

input

```

integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/
2),x, algorithm="giac")

```

output

```

integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(5/2)/sqrt(-I*c*tan(
f*x + e) + c), x)

```

3.809.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx = \int \frac{(A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{5/2}}{\sqrt{c - c \tan(e + fx) li}} dx$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(5/2))/(c - c*tan(e + f*x)*1i)^(1/2),x)`

output `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(5/2))/(c - c*tan(e + f*x)*1i)^(1/2), x)`

3.810 $\int \frac{(a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$

3.810.1 Optimal result 7300
 3.810.2 Mathematica [A] (verified) 7301
 3.810.3 Rubi [A] (verified) 7301
 3.810.4 Maple [B] (verified) 7304
 3.810.5 Fricas [B] (verification not implemented) 7305
 3.810.6 Sympy [F] 7306
 3.810.7 Maxima [B] (verification not implemented) 7306
 3.810.8 Giac [F] 7307
 3.810.9 Mupad [F(-1)] 7308

3.810.1 Optimal result

Integrand size = 45, antiderivative size = 226

$$\int \frac{(a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx =$$

$$-\frac{2a^{5/2}(iA + 4B) \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{c^{3/2}f}$$

$$-\frac{(iA + B)(a + ia \tan(e + fx))^{5/2}}{3f(c - ic \tan(e + fx))^{3/2}} + \frac{2a(iA + 4B)(a + ia \tan(e + fx))^{3/2}}{3cf\sqrt{c - ic \tan(e + fx)}}$$

$$+ \frac{a^2(iA + 4B)\sqrt{a + ia \tan(e + fx)}\sqrt{c - ic \tan(e + fx)}}{c^2f}$$

```
output -2*a^(5/2)*(I*A+4*B)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/c^(3/2)/f+a^2*(I*A+4*B)*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)/c^2/f+2/3*a*(I*A+4*B)*(a+I*a*tan(f*x+e))^(3/2)/c/f/(c-I*c*tan(f*x+e))^(1/2)-1/3*(I*A+B)*(a+I*a*tan(f*x+e))^(5/2)/f/(c-I*c*tan(f*x+e))^(3/2)
```

3.810.2 Mathematica [A] (verified)

Time = 7.63 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.79

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \frac{a^2 \cos^2(e + fx) \sqrt{a + ia \tan(e + fx)} ((-3 - 3i)(A - 4$$

input `Integrate[((a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(a^2*Cos[e + f*x]^2*Sqrt[a + I*a*Tan[e + f*x]]*((-3 - 3*I)*(A - (4*I)*B)*A rcSin[(1/2 + I/2)*Sqrt[-I + Tan[e + f*x]]]*Sqrt[2 - (2*I)*Tan[e + f*x]]*Sqrt[-I + Tan[e + f*x]]*(I + Tan[e + f*x]) + (-I + Tan[e + f*x])*(-4*A + (19*I)*B + ((8*I)*A + 26*B)*Tan[e + f*x] - (3*I)*B*Tan[e + f*x]^2))/(3*c*f*Sqrt[c - I*c*Tan[e + f*x]])`

3.810.3 Rubi [A] (verified)Time = 0.43 (sec) , antiderivative size = 222, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 87, 57, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(i \tan(e+fx)a+a)^{3/2} (A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{87} \end{aligned}$$

3.810. $\int \frac{(a+ia \tan(e+fx))^{5/2} (A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$

$$ac \left(- \frac{(A-4iB) \int \frac{(i \tan(e+fx)a+a)^{3/2}}{(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{3c} - \frac{(B+iA)(a+ia \tan(e+fx))^{5/2}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)$$

f
↓ 57

$$ac \left(- \frac{(A-4iB) \left(- \frac{3a \int \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{3c} - \frac{2i(a+ia \tan(e+fx))^{3/2}}{c\sqrt{c-ic \tan(e+fx)}} \right)}{3c} - \frac{(B+iA)(a+ia \tan(e+fx))^{5/2}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)$$

f
↓ 60

$$ac \left(- \frac{(A-4iB) \left(- \frac{3a \left(a \int \frac{1}{\sqrt{i \tan(e+fx)a+a} \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{c} \right)}{3c} - \frac{2i(a+ia \tan(e+fx))^{3/2}}{c\sqrt{c-ic \tan(e+fx)}} \right)}{3c} - \frac{(B+iA)(a+ia \tan(e+fx))^{5/2}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)$$

f

↓ 45

$$ac \left(- \frac{(A-4iB) \left(- \frac{3a \left(2a \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} + \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{c} \right)}{3c} - \frac{2i(a+ia \tan(e+fx))^{3/2}}{c\sqrt{c-ic \tan(e+fx)}} \right)}{3c} - \frac{(B+iA)(a+ia \tan(e+fx))^{5/2}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)$$

f

↓ 218

$$ac \left(- \frac{(A-4iB) \left(- \frac{3a \left(\frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{c} - \frac{2i\sqrt{a} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{\sqrt{c}} \right)}{3c} - \frac{2i(a+ia \tan(e+fx))^{3/2}}{c\sqrt{c-ic \tan(e+fx)}} \right)}{3c} - \frac{(B+iA)(a+ia \tan(e+fx))^{5/2}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)$$

f

3.810. $\int \frac{(a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$

input `Int[((a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(a*c*(-1/3*((I*A + B)*(a + I*a*Tan[e + f*x])^(5/2)))/(a*c*(c - I*c*Tan[e + f*x])^(3/2)) - ((A - (4*I)*B)*((-2*I)*(a + I*a*Tan[e + f*x])^(3/2))/(c*Sqrt[c - I*c*Tan[e + f*x]]) - (3*a*((-2*I)*Sqrt[a]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[c] + (I*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/c)/(3*c))/f`

3.810.3.1 Defintions of rubi rules used

rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 57 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

rule 60 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.810.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 666 vs. $2(186) = 372$.

Time = 0.35 (sec) , antiderivative size = 667, normalized size of antiderivative = 2.95

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^2 \left(-12 i B \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}}\right)\right) a c \tan (f x+e)^3+9 i A}{}$
default	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^2 \left(-12 i B \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}}\right)\right) a c \tan (f x+e)^3+9 i A}{}$
parts	$\frac{A \sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^2 \left(9 i \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}}\right)\right) \tan (f x+e)^2 a c+3 \ln \left(\frac{a}{c-i c \tan (f x+e)}\right)}{}$

input `int((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x,m
method=_RETURNVERBOSE)`

$$3.810. \quad \int \frac{(a+ia \tan (e+fx))^{5/2}(A+B \tan (e+fx))}{(c-ic \tan (e+fx))^{3/2}} dx$$

```

output 1/3/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^2/c^2*(-12*
I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/
2))*a*c*tan(f*x+e)^3+9*I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+
e)^2))^(1/2)))/(a*c)^(1/2))*a*c*tan(f*x+e)^2+3*A*ln((a*c*tan(f*x+e)+(a*c)^(
1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c*tan(f*x+e)^3+36*I*B*ln
((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*
c*tan(f*x+e)+29*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+
36*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1
/2))*a*c*tan(f*x+e)^2+3*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x
+e)^3-3*I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(
a*c)^(1/2))*a*c-12*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)
-9*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1
/2))*a*c*tan(f*x+e)-8*A*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)*tan(f*x+e)
)^2-19*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)-12*B*ln((a*c*tan(f*x+e)
)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c-45*B*(a*c)^(1
/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+4*A*(a*c)^(1/2)*(a*c*(1+tan(f*
x+e)^2))^(1/2))/(a*c*(1+tan(f*x+e)^2))^(1/2)/(a*c)^(1/2)/(I+tan(f*x+e))^3

```

3.810.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 493 vs. $2(176) = 352$.

Time = 0.26 (sec) , antiderivative size = 493, normalized size of antiderivative = 2.18

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx =$$

$$3c^2 \sqrt{\frac{(A^2 - 8iAB - 16B^2)a^5}{c^3 f^2}} f \log \left(\frac{4 \left(2((-iA - 4B)a^2 e^{(3i fx + 3ie)} + (-iA - 4B)a^2 e^{(i fx + ie)}) \sqrt{\frac{a}{e^{(2i fx + 2ie)} + 1}} \sqrt{\frac{c}{e^{(2i fx + 2ie)} + 1}} + (c^2 f \right)}{(iA + 4B)a^2 e^{(2i fx + 2ie)} + (iA + 4B)a^2} \right)$$

```

input integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/
2),x, algorithm="fracas")

```

3.810. $\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx$

output
$$\begin{aligned} & -1/6*(3*c^2*\sqrt{(A^2 - 8*I*A*B - 16*B^2)*a^5/(c^3*f^2)})*f*\log(-4*(2*((-I*A - 4*B)*a^2*e^{(3*I*f*x + 3*I*e)} + (-I*A - 4*B)*a^2*e^{(I*f*x + I*e)}))*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1))*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} + (c^2*f*e^{(2*I*f*x + 2*I*e)} - c^2*f)*\sqrt{(A^2 - 8*I*A*B - 16*B^2)*a^5/(c^3*f^2))})/(\\ & ((I*A + 4*B)*a^2*e^{(2*I*f*x + 2*I*e)} + (I*A + 4*B)*a^2)) - 3*c^2*\sqrt{(A^2 - 8*I*A*B - 16*B^2)*a^5/(c^3*f^2)})*f*\log(-4*(2*((-I*A - 4*B)*a^2*e^{(3*I*f*x + 3*I*e)} + (-I*A - 4*B)*a^2*e^{(I*f*x + I*e)}))*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1))*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} - (c^2*f*e^{(2*I*f*x + 2*I*e)} - c^2*f)*\sqrt{(A^2 - 8*I*A*B - 16*B^2)*a^5/(c^3*f^2))})/(\\ & ((I*A + 4*B)*a^2*e^{(2*I*f*x + 2*I*e)} + (I*A + 4*B)*a^2)) + 4*((I*A + B)*a^2*e^{(5*I*f*x + 5*I*e)} + 2*(-I*A - 4*B)*a^2*e^{(3*I*f*x + 3*I*e)} + 3*(-I*A - 4*B)*a^2*e^{(I*f*x + I*e)}))*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1))*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1))})/(\\ & c^2*f) \end{aligned}$$

3.810.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \int \frac{(ia(\tan(e + fx) - i))^{5/2} (A + B \tan(e + fx))}{(-ic(\tan(e + fx) + i))^{3/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(3/2),x)`

output `Integral((I*a*(tan(e + f*x) - I))**(5/2)*(A + B*tan(e + f*x))/(-I*c*(tan(e + f*x) + I))**(3/2), x)`

3.810.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 795 vs. $2(176) = 352$.

Time = 0.49 (sec) , antiderivative size = 795, normalized size of antiderivative = 3.52

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="maxima")`

3.810.
$$\int \frac{(a+ia \tan(e+fx))^{5/2} (A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$$

output

```

-3*(6*((A - 4*I*B)*a^2*cos(2*f*x + 2*e) - (-I*A - 4*B)*a^2*sin(2*f*x + 2*
e) + (A - 4*I*B)*a^2)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x +
2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) + 6*((A
- 4*I*B)*a^2*cos(2*f*x + 2*e) - (-I*A - 4*B)*a^2*sin(2*f*x + 2*e) + (A -
4*I*B)*a^2)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))),
-sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) + 4*((A - I*B)*
a^2*cos(2*f*x + 2*e) - (-I*A - B)*a^2*sin(2*f*x + 2*e) + (A - I*B)*a^2)*co
s(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 12*((A - 3*I*B)*a^2*c
os(2*f*x + 2*e) + (I*A + 3*B)*a^2*sin(2*f*x + 2*e) + (A - 4*I*B)*a^2)*cos(
1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 3*((-I*A - 4*B)*a^2*cos
(2*f*x + 2*e) + (A - 4*I*B)*a^2*sin(2*f*x + 2*e) + (-I*A - 4*B)*a^2)*log(c
os(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + sin(1/2*arctan2(si
n(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 2*sin(1/2*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e))) + 1) - 3*((I*A + 4*B)*a^2*cos(2*f*x + 2*e) - (A - 4*I*
B)*a^2*sin(2*f*x + 2*e) + (I*A + 4*B)*a^2)*log(cos(1/2*arctan2(sin(2*f*x +
2*e), cos(2*f*x + 2*e)))^2 + sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x
+ 2*e)))^2 - 2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) -
4*((-I*A - B)*a^2*cos(2*f*x + 2*e) + (A - I*B)*a^2*sin(2*f*x + 2*e) + (-I
*A - B)*a^2)*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 12*((I
*A + 3*B)*a^2*cos(2*f*x + 2*e) - (A - 3*I*B)*a^2*sin(2*f*x + 2*e) + (I*...

```

3.810.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{5/2}}{(-ic \tan(fx + e) + c)^{3/2}} dx$$

input

```

integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/
2),x, algorithm="giac")

```

output

```

integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(5/2)/(-I*c*tan(f*x
+ e) + c)^(3/2), x)

```

3.810.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{3/2}} dx = \int \frac{(A + B \tan(e + fx)) (a + a \tan(e + fx) \text{li})^{5/2}}{(c - c \tan(e + fx) \text{li})^{3/2}} dx$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(5/2))/(c - c*tan(e + f*x)*1i)^(3/2), x)`

output `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(5/2))/(c - c*tan(e + f*x)*1i)^(3/2), x)`

3.811
$$\int \frac{(a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{5/2}} dx$$

3.811.1 Optimal result 7309
 3.811.2 Mathematica [A] (verified) 7309
 3.811.3 Rubi [A] (verified) 7310
 3.811.4 Maple [B] (verified) 7313
 3.811.5 Fricas [B] (verification not implemented) 7314
 3.811.6 Sympy [F] 7315
 3.811.7 Maxima [A] (verification not implemented) 7315
 3.811.8 Giac [F] 7316
 3.811.9 Mupad [F(-1)] 7316

3.811.1 Optimal result

Integrand size = 45, antiderivative size = 203

$$\int \frac{(a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{5/2}} dx = \frac{2a^{5/2}B \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ict \tan(e+fx)}}\right)}{c^{5/2}f} - \frac{(iA + B)(a + ia \tan(e + fx))^{5/2}}{5f(c - ict \tan(e + fx))^{5/2}} + \frac{2aB(a + ia \tan(e + fx))^{3/2}}{3cf(c - ict \tan(e + fx))^{3/2}} - \frac{2a^2B\sqrt{a + ia \tan(e + fx)}}{c^2f\sqrt{c - ict \tan(e + fx)}}$$

output `2*a^(5/2)*B*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/c^(5/2)/f-2*a^2*B*(a+I*a*tan(f*x+e))^(1/2)/c^2/f/(c-I*c*tan(f*x+e))^(1/2)-1/5*(I*A+B)*(a+I*a*tan(f*x+e))^(5/2)/f/(c-I*c*tan(f*x+e))^(5/2)+2/3*a*B*(a+I*a*tan(f*x+e))^(3/2)/c/f/(c-I*c*tan(f*x+e))^(3/2)`

3.811.2 Mathematica [A] (verified)

Time = 17.95 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.00

$$\int \frac{(a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{5/2}} dx = \frac{a^2 \cos^2(e + fx) (\cos(\frac{1}{2}(e - 2fx)) - i \sin(\frac{1}{2}(e - 2fx)))}{(c - ict \tan(e + fx))^{5/2}}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(5/2), x]`

3.811.
$$\int \frac{(a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{5/2}} dx$$

output $(a^2 \cos[e + f x]^2 (\cos[(e - 2 f x)/2] - I \sin[(e - 2 f x)/2]) (\cos[(e - 2 f x)/2] + I \sin[(e - 2 f x)/2]) (-10 B + ((3 I) A + 33 B) \cos[2(e + f x)]) - 3 A \sin[2(e + f x)] - (27 I) B \sin[2(e + f x)] - 30 B \operatorname{ArcTan}[E^{(I(e + f x))}] (\cos[3(e + f x)] - I \sin[3(e + f x)]) (-I + \tan[e + f x])^2 \operatorname{Sqrt}[a + I a \tan[e + f x]] / (15 c^2 f \operatorname{Sqrt}[c - I c \tan[e + f x]])$

3.811.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 87, 57, 57, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx$$

↓ 4071

$$ac \int \frac{(i \tan(e + fx) a + a)^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} d \tan(e + fx)$$

f
↓ 87

$$ac \left(\frac{iB \int \frac{(i \tan(e + fx) a + a)^{3/2}}{(c - ic \tan(e + fx))^{5/2}} d \tan(e + fx)}{c} - \frac{(B + iA)(a + ia \tan(e + fx))^{5/2}}{5ac(c - ic \tan(e + fx))^{5/2}} \right)$$

f
↓ 57

$$ac \left(\frac{iB \left(-\frac{a \int \frac{\sqrt{i \tan(e + fx) a + a}}{(c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{c} - \frac{2i(a + ia \tan(e + fx))^{3/2}}{3c(c - ic \tan(e + fx))^{3/2}} \right)}{c} - \frac{(B + iA)(a + ia \tan(e + fx))^{5/2}}{5ac(c - ic \tan(e + fx))^{5/2}} \right)$$

f
↓ 57

3.811. $\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx$

$$ac \left(\frac{iB \left(a \left(\frac{\int \frac{1}{\sqrt{i \tan(e+fx)a+a\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{c} - \frac{2i\sqrt{a+ia \tan(e+fx)}}{c\sqrt{c-ic \tan(e+fx)}} \right) - \frac{2i(a+ia \tan(e+fx))^{3/2}}{3c(c-ic \tan(e+fx))^{3/2}} \right)}{c} - \frac{(B+ia)(a+ia \tan(e+fx))^{5/2}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)$$

f

↓ 45

$$ac \left(\frac{iB \left(a \left(\frac{\int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}}}{c} - \frac{2i\sqrt{a+ia \tan(e+fx)}}{c\sqrt{c-ic \tan(e+fx)}} \right) - \frac{2i(a+ia \tan(e+fx))^{3/2}}{3c(c-ic \tan(e+fx))^{3/2}} \right)}{c} - \frac{(B+ia)(a+ia \tan(e+fx))^{5/2}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)$$

f

↓ 218

$$ac \left(\frac{iB \left(a \left(\frac{2i\sqrt{a} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{c^{3/2}} - \frac{2i\sqrt{a+ia \tan(e+fx)}}{c\sqrt{c-ic \tan(e+fx)}} \right) - \frac{2i(a+ia \tan(e+fx))^{3/2}}{3c(c-ic \tan(e+fx))^{3/2}} \right)}{c} - \frac{(B+ia)(a+ia \tan(e+fx))^{5/2}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)$$

f

```
input Int[((a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(5/2), x]
```

3.811. $\int \frac{(a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$

output $(a*c*(-1/5*((I*A + B)*(a + I*a*\text{Tan}[e + f*x])^{5/2}))/ (a*c*(c - I*c*\text{Tan}[e + f*x])^{5/2}) + (I*B*((((-2*I)/3)*(a + I*a*\text{Tan}[e + f*x])^{3/2}))/ (c*(c - I*c*\text{Tan}[e + f*x])^{3/2}) - (a*(((2*I)*\text{Sqrt}[a]*\text{ArcTan}[(\text{Sqrt}[c]*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]])/(\text{Sqrt}[a]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])]))/c^{3/2} - ((2*I)*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]])/(c*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])))/c)/c)/f$

3.811.3.1 Defintions of rubi rules used

rule 45 $\text{Int}[1/(\text{Sqrt}[(a_) + (b_)*(x_)]*\text{Sqrt}[(c_) + (d_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[2 \text{ Subst}[\text{Int}[1/(b - d*x^2), x], x, \text{Sqrt}[a + b*x]/\text{Sqrt}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{EqQ}[b*c + a*d, 0] \&\& !\text{GtQ}[c, 0]$

rule 57 $\text{Int}[(a_) + (b_)*(x_)^{(m_)}*((c_) + (d_)*(x_)^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + 1))), x] - \text{Simp}[d*(n/(b*(m + 1))) \text{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^{(n - 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{GtQ}[n, 0] \&\& \text{LtQ}[m, -1] \&\& !(IntegerQ[n] \&\& !IntegerQ[m]) \&\& !(ILeQ[m + n + 2, 0] \&\& (\text{FractionQ}[m] || \text{GeQ}[2*n + m + 1, 0])) \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 87 $\text{Int}[(a_) + (b_)*(x_)*((c_) + (d_)*(x_))^{(n_)}*((e_) + (f_)*(x_))^{(p_)}, x_] \rightarrow \text{Simp}[(-b*e - a*f)*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)})/(f*(p + 1)*(c*f - d*e)), x] - \text{Simp}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) \text{Int}[(c + d*x)^n*(e + f*x)^{(p + 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \&\& \text{LtQ}[p, -1] \&\& (!\text{LtQ}[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || \text{LtQ}[p, n]))))$

rule 218 $\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.811.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 506 vs. 2(166) = 332.

Time = 0.35 (sec) , antiderivative size = 507, normalized size of antiderivative = 2.50

method	result
parts	$-\frac{A\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^2(1+\tan(fx+e)^2)(i-\tan(fx+e))}{5fc^3(i+\tan(fx+e))^4} + \frac{iB\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}}{5fc^3(i+\tan(fx+e))^4}$
derivativedivides	$-\frac{\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^2\left(-15iB\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}}\right)\right)actan(fx+e)^4+9}{5fc^3(i+\tan(fx+e))^4}$
default	$-\frac{\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^2\left(-15iB\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}}\right)\right)actan(fx+e)^4+9}{5fc^3(i+\tan(fx+e))^4}$

```
input int((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,m
ethod=_RETURNVERBOSE)
```

$$3.811. \int \frac{(a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$$

output

```

-1/5*A/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^2/c^3*(1
+tan(f*x+e)^2)*(I-tan(f*x+e))/(I+tan(f*x+e))^4+1/15*I*B/f*(a*(1+I*tan(f*x+
e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^2/c^3*(60*I*ln((a*c*tan(f*x+e)+(a
*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)))^(1/2))/(a*c)^(1/2))*tan(f*x+e)^3*a*c+15*I
n((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)))^(1/2))/(a*c)^(1/2))*t
an(f*x+e)^4*a*c-60*I*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))
^(1/2))/(a*c)^(1/2))*tan(f*x+e)*a*c-97*I*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c
)^(1/2)*tan(f*x+e)^2-90*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^
2)))^(1/2))/(a*c)^(1/2))*tan(f*x+e)^2*a*c-43*tan(f*x+e)^3*(a*c*(1+tan(f*x+e
)^2))^(1/2)*(a*c)^(1/2)+23*I*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)+15*a
*c*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)))^(1/2))/(a*c)^(1/2
))+77*tan(f*x+e)*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2))/(a*c*(1+tan(f*x
+e)^2))^(1/2)/(I+tan(f*x+e))^4/(a*c)^(1/2)

```

3.811.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 418 vs. $2(159) = 318$.

Time = 0.27 (sec) , antiderivative size = 418, normalized size of antiderivative = 2.06

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx =$$

$$15 c^3 f \sqrt{-\frac{B^2 a^5}{c^5 f^2}} \log \left(\frac{4 \left(2 (Ba^2 e^{(3i fx + 3i e)} + Ba^2 e^{(i fx + i e)}) \sqrt{\frac{a}{e^{(2i fx + 2i e)} + 1}} \sqrt{\frac{c}{e^{(2i fx + 2i e)} + 1}} + (c^3 f e^{(2i fx + 2i e)} - c^3 f) \sqrt{-\frac{B^2 a^5}{c^5 f^2}} \right)}{Ba^2 e^{(2i fx + 2i e)} + Ba^2} \right)$$

input

```

integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/
2),x, algorithm="fracas")

```

output
$$\begin{aligned} & -1/30*(15*c^3*f*\sqrt{-B^2*a^5/(c^5*f^2)}*\log(4*(2*(B*a^2*e^{(3*I*f*x + 3*I*e)} + B*a^2*e^{(I*f*x + I*e)})*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} + (c^3*f*e^{(2*I*f*x + 2*I*e)} - c^3*f)*\sqrt{-B^2*a^5/(c^5*f^2)}))/(B*a^2*e^{(2*I*f*x + 2*I*e)} + B*a^2)) - 15*c^3*f*\sqrt{-B^2*a^5/(c^5*f^2)}*\log(4*(2*(B*a^2*e^{(3*I*f*x + 3*I*e)} + B*a^2*e^{(I*f*x + I*e)})*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} - (c^3*f*e^{(2*I*f*x + 2*I*e)} - c^3*f)*\sqrt{-B^2*a^5/(c^5*f^2)}))/(B*a^2*e^{(2*I*f*x + 2*I*e)} + B*a^2)) + 2*(3*(I*A + B)*a^2*e^{(7*I*f*x + 7*I*e)} + (3*I*A - 7*B)*a^2*e^{(5*I*f*x + 5*I*e)} + 20*B*a^2*e^{(3*I*f*x + 3*I*e)} + 30*B*a^2*e^{(I*f*x + I*e)})*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)))/(c^3*f) \end{aligned}$$

3.811.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \int \frac{(ia(\tan(e + fx) - i))^{5/2} (A + B \tan(e + fx))}{(-ic(\tan(e + fx) + i))^{5/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(5/2),x)`

output `Integral((I*a*(tan(e + f*x) - I))**(5/2)*(A + B*tan(e + f*x))/(-I*c*(tan(e + f*x) + I))**(5/2), x)`

3.811.7 Maxima [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.05

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \frac{(30 B a^2 \arctan(\cos(fx + e), \sin(fx + e) + 1) + 30 B$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `1/30*(30*B*a^2*arctan2(cos(f*x + e), sin(f*x + e) + 1) + 30*B*a^2*arctan2(cos(f*x + e), -sin(f*x + e) + 1) - 6*(I*A + B)*a^2*cos(5*f*x + 5*e) + 20*B*a^2*cos(3*f*x + 3*e) - 60*B*a^2*cos(f*x + e) + 15*I*B*a^2*log(cos(f*x + e)^2 + sin(f*x + e)^2 + 2*sin(f*x + e) + 1) - 15*I*B*a^2*log(cos(f*x + e)^2 + sin(f*x + e)^2 - 2*sin(f*x + e) + 1) + 6*(A - I*B)*a^2*sin(5*f*x + 5*e) + 20*I*B*a^2*sin(3*f*x + 3*e) - 60*I*B*a^2*sin(f*x + e))*sqrt(a)/(c^(5/2)*f)`

3.811.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{5/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{5/2}}{(-ict \tan(fx + e) + c)^{5/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(5/2)/(-I*c*tan(f*x + e) + c)^(5/2), x)`

3.811.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{5/2}} dx = \int \frac{(A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{5/2}}{(c - c \tan(e + fx) li)^{5/2}} dx$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*li)^(5/2))/(c - c*tan(e + f*x)*li)^(5/2),x)`

output `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*li)^(5/2))/(c - c*tan(e + f*x)*li)^(5/2), x)`

3.812
$$\int \frac{(a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{7/2}} dx$$

3.812.1 Optimal result	7317
3.812.2 Mathematica [A] (verified)	7317
3.812.3 Rubi [A] (verified)	7318
3.812.4 Maple [A] (verified)	7319
3.812.5 Fricas [A] (verification not implemented)	7320
3.812.6 Sympy [F]	7320
3.812.7 Maxima [B] (verification not implemented)	7321
3.812.8 Giac [F]	7321
3.812.9 Mupad [B] (verification not implemented)	7322

3.812.1 Optimal result

Integrand size = 45, antiderivative size = 102

$$\int \frac{(a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{7/2}} dx =$$

$$-\frac{(iA + B)(a + ia \tan(e + fx))^{5/2}}{7f(c - ict \tan(e + fx))^{7/2}} - \frac{(iA - 6B)(a + ia \tan(e + fx))^{5/2}}{35cf(c - ict \tan(e + fx))^{5/2}}$$

output
$$-1/7*(I*A+B)*(a+I*a*\tan(f*x+e))^(5/2)/f/(c-I*c*\tan(f*x+e))^(7/2)-1/35*(I*A-6*B)*(a+I*a*\tan(f*x+e))^(5/2)/c/f/(c-I*c*\tan(f*x+e))^(5/2)$$

3.812.2 Mathematica [A] (verified)

Time = 6.90 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.11

$$\int \frac{(a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{7/2}} dx = \frac{ia^3 \sec^3(e + fx)(\cos(3(e + fx)) + i \sin(3(e + fx)))(6i(A - B + (A + (6I)B)\tan(e + fx)))}{35c^3 f (i + \tan(e + fx))^3 \sqrt{a + ia \tan(e + fx)}}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(7/2),x]`

output
$$((I/35)*a^3*Sec[e + f*x]^3*(Cos[3*(e + f*x)] + I*Sin[3*(e + f*x)])*((6*I)*A - B + (A + (6*I)*B)*Tan[e + f*x]))/(c^3*f*(I + Tan[e + f*x])^3*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])$$

3.812.
$$\int \frac{(a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{7/2}} dx$$

3.812.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.089$, Rules used = {3042, 4071, 87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx \\
 \downarrow \text{3042} \\
 \int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx \\
 \downarrow \text{4071} \\
 \frac{ac \int \frac{(i \tan(e + fx)a + a)^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} d \tan(e + fx)}{f} \\
 \downarrow \text{87} \\
 \frac{ac \left(\frac{(A + 6iB) \int \frac{(i \tan(e + fx)a + a)^{3/2}}{(c - ic \tan(e + fx))^{7/2}} d \tan(e + fx)}{7c} - \frac{(B + iA)(a + ia \tan(e + fx))^{5/2}}{7ac(c - ic \tan(e + fx))^{7/2}} \right)}{f} \\
 \downarrow \text{48} \\
 \frac{ac \left(-\frac{i(A + 6iB)(a + ia \tan(e + fx))^{5/2}}{35ac^2(c - ic \tan(e + fx))^{5/2}} - \frac{(B + iA)(a + ia \tan(e + fx))^{5/2}}{7ac(c - ic \tan(e + fx))^{7/2}} \right)}{f}
 \end{array}$$

input `Int[((a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(7/2), x]`

output `(a*c*(-1/7*((I*A + B)*(a + I*a*Tan[e + f*x])^(5/2)))/(a*c*(c - I*c*Tan[e + f*x])^(7/2)) - ((I/35)*(A + (6*I)*B)*(a + I*a*Tan[e + f*x])^(5/2))/(a*c^2*(c - I*c*Tan[e + f*x])^(5/2)))/f`

3.812. $\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx$

3.812.3.1 Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.812.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.04

method	result
risch	$-\frac{a^2 \sqrt{\frac{a e^{2i(fx+e)}}{e^{2i(fx+e)+1}}} (5iA e^{6i(fx+e)} + 5B e^{6i(fx+e)} + 7iA e^{4i(fx+e)} - 7B e^{4i(fx+e)})}{70c^3 \sqrt{\frac{c}{e^{2i(fx+e)+1}}} f}$
derivativedivides	$-\frac{i \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^2 (1+\tan(fx+e)^2) (iA \tan(fx+e)^2 + 5i \tan(fx+e)B - 6B \tan(fx+e)^2 + 6A)}{35f c^4 (i+\tan(fx+e))^5}$
default	$-\frac{i \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^2 (1+\tan(fx+e)^2) (iA \tan(fx+e)^2 + 5i \tan(fx+e)B - 6B \tan(fx+e)^2 + 6A)}{35f c^4 (i+\tan(fx+e))^5}$
parts	$\frac{A \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^2 (1+\tan(fx+e)^2) (5i \tan(fx+e) + \tan(fx+e)^2 + 6)}{35f c^4 (i+\tan(fx+e))^5} - \frac{iB \sqrt{a(1+i \tan(fx+e))}}{35f c^4 (i+\tan(fx+e))^5}$

3.812.
$$\int \frac{(a+ia \tan(e+fx))^{5/2} (A+B \tan(e+fx))}{(c-ictan(e+fx))^{7/2}} dx$$

```
input int((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x,method=_RETURNVERBOSE)
```

```
output -1/70*a^2/c^3*(a*exp(2*I*(f*x+e))/(exp(2*I*(f*x+e))+1))^(1/2)/(c/(exp(2*I*(f*x+e))+1))^(1/2)/f*(5*I*A*exp(6*I*(f*x+e))+5*B*exp(6*I*(f*x+e))+7*I*A*exp(4*I*(f*x+e))-7*B*exp(4*I*(f*x+e)))
```

3.812.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.02

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \frac{(5(iA + B)a^2 e^{(9i fx + 9i e)} + 2(6iA - B)a^2 e^{(7i fx + 7i e)} + 7(iA - B)a^2 e^{(5i fx + 5i e)}) \sqrt{\frac{a}{e^{(2i fx + 2i e)} + 1}} \sqrt{\frac{c}{e^{(2i fx + 2i e)} + 1}}}{70 c^4 f}$$

```
input integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x, algorithm="fricas")
```

```
output -1/70*(5*(I*A + B)*a^2*e^(9*I*f*x + 9*I*e) + 2*(6*I*A - B)*a^2*e^(7*I*f*x + 7*I*e) + 7*(I*A - B)*a^2*e^(5*I*f*x + 5*I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^4*f)
```

3.812.6 Sympy [F]

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \int \frac{(ia(\tan(e + fx) - i))^{5/2} (A + B \tan(e + fx))}{(-ic(\tan(e + fx) + i))^{7/2}} dx$$

```
input integrate((a+I*a*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(7/2),x)
```

```
output Integral((I*a*(tan(e + f*x) - I))**(5/2)*(A + B*tan(e + f*x))/(-I*c*(tan(e + f*x) + I))**(7/2), x)
```

3.812. $\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx$

3.812.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 166 vs. $2(78) = 156$.

Time = 0.43 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.63

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx =$$

$$\frac{70(5(A - iB)a^2 \cos(9fx + 9e) + 2(6A + iB)a^2 \cos(7fx + 7e) + 7(A + iB)a^2 \cos(5fx + 5e) - 5(-I^2A - B)a^2 \sin(9fx + 9e) - 2(-6I^2A + B)a^2 \sin(7fx + 7e) - 7(-I^2A + B)a^2 \sin(5fx + 5e)) \sqrt{a} \sqrt{c}}{-4900(i c^4 \cos(2fx + 2e) - 4900 I^2 c^4 \sin(2fx + 2e))} + \dots$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x, algorithm="maxima")`

output `-70*(5*(A - I*B)*a^2*cos(9*f*x + 9*e) + 2*(6*A + I*B)*a^2*cos(7*f*x + 7*e) + 7*(A + I*B)*a^2*cos(5*f*x + 5*e) - 5*(-I*A - B)*a^2*sin(9*f*x + 9*e) - 2*(-6*I*A + B)*a^2*sin(7*f*x + 7*e) - 7*(-I*A + B)*a^2*sin(5*f*x + 5*e))*sqrt(a)*sqrt(c)/((-4900*I*c^4*cos(2*f*x + 2*e) + 4900*c^4*sin(2*f*x + 2*e) - 4900*I*c^4)*f)`

3.812.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{5/2}}{(-ic \tan(fx + e) + c)^{7/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(5/2)/(-I*c*tan(f*x + e) + c)^(7/2), x)`

3.812.9 Mupad [B] (verification not implemented)

Time = 9.96 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.88

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx =$$

$$\frac{a^2 \sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)i)}{\cos(2e+2fx)+1}} (A \cos(4e+4fx) 7i + A \cos(6e+6fx) 5i - 7B \cos(4e+4fx) + 5B \cos(6e+6fx) - 7A \sin(4e+4fx) - 5A \sin(6e+6fx) - B \sin(4e+4fx) 7i + B \sin(6e+6fx) 5i)}{70 c^3 f \sqrt{\frac{c(\cos(2e+2fx)+1+\sin(2e+2fx)i)}{\cos(2e+2fx)+1}}}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(5/2))/(c - c*tan(e + f*x)*1i)^(7/2),x)`

output `-(a^2*((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*cos(4*e + 4*f*x)*7i + A*cos(6*e + 6*f*x)*5i - 7*B*cos(4*e + 4*f*x) + 5*B*cos(6*e + 6*f*x) - 7*A*sin(4*e + 4*f*x) - 5*A*sin(6*e + 6*f*x) - B*sin(4*e + 4*f*x)*7i + B*sin(6*e + 6*f*x)*5i))/(70*c^3*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

3.813
$$\int \frac{(a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{9/2}} dx$$

3.813.1 Optimal result 7323
 3.813.2 Mathematica [A] (verified) 7323
 3.813.3 Rubi [A] (verified) 7324
 3.813.4 Maple [A] (verified) 7326
 3.813.5 Fricas [A] (verification not implemented) 7326
 3.813.6 Sympy [F(-1)] 7327
 3.813.7 Maxima [A] (verification not implemented) 7327
 3.813.8 Giac [F] 7328
 3.813.9 Mupad [B] (verification not implemented) 7328

3.813.1 Optimal result

Integrand size = 45, antiderivative size = 155

$$\int \frac{(a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{9/2}} dx = -\frac{(iA + B)(a + ia \tan(e + fx))^{5/2}}{9f(c - ict \tan(e + fx))^{9/2}} - \frac{(2iA - 7B)(a + ia \tan(e + fx))^{5/2}}{63cf(c - ict \tan(e + fx))^{7/2}} - \frac{(2iA - 7B)(a + ia \tan(e + fx))^{5/2}}{315c^2f(c - ict \tan(e + fx))^{5/2}}$$

output

```
-1/9*(I*A+B)*(a+I*a*tan(f*x+e))^(5/2)/f/(c-I*c*tan(f*x+e))^(9/2)-1/63*(2*I
*A-7*B)*(a+I*a*tan(f*x+e))^(5/2)/c/f/(c-I*c*tan(f*x+e))^(7/2)-1/315*(2*I*A
-7*B)*(a+I*a*tan(f*x+e))^(5/2)/c^2/f/(c-I*c*tan(f*x+e))^(5/2)
```

3.813.2 Mathematica [A] (verified)

Time = 7.02 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.84

$$\int \frac{(a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{9/2}} dx = \frac{a^3 \sec^5(e + fx)(45A + 7(7A + 2iB) \cos(2(e + fx)) + \dots)}{630c^4 f(i + \tan(e + fx))^4 \sqrt{\dots}}$$

input

```
Integrate[((a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan
[e + f*x])^(9/2),x]
```

output $(a^3 \sec[e + fx]^5 (45A + 7(7A + (2I)B) \cos[2(e + fx)] + 7((-2I)A + 7B) \sin[2(e + fx)]) \cdot ((-I) \cos[3(e + fx)] + \sin[3(e + fx)]) / (630c^4 f (I + \tan[e + fx])^4 \sqrt{a + I a \tan[e + fx]} \sqrt{c - I c \tan[e + fx]})$

3.813.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.08, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4071, 87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{9/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{9/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{(i \tan(e + fx) a + a)^{3/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{11/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{87} \\
 & \frac{ac \left(\frac{(2A + 7iB) \int \frac{(i \tan(e + fx) a + a)^{3/2}}{(c - ict \tan(e + fx))^{9/2}} d \tan(e + fx)}{9c} - \frac{(B + iA)(a + ia \tan(e + fx))^{5/2}}{9ac(c - ict \tan(e + fx))^{9/2}} \right)}{f} \\
 & \quad \downarrow \text{55} \\
 & \frac{ac \left(\frac{(2A + 7iB) \left(\frac{\int \frac{(i \tan(e + fx) a + a)^{3/2}}{(c - ict \tan(e + fx))^{7/2}} d \tan(e + fx)}{7c} - \frac{i(a + ia \tan(e + fx))^{5/2}}{7ac(c - ict \tan(e + fx))^{7/2}} \right)}{9c} - \frac{(B + iA)(a + ia \tan(e + fx))^{5/2}}{9ac(c - ict \tan(e + fx))^{9/2}} \right)}{f} \\
 & \quad \downarrow \text{48}
 \end{aligned}$$

3.813. $\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{9/2}} dx$

$$\frac{ac \left(\frac{(2A+7iB) \left(-\frac{i(a+ia \tan(e+fx))^{5/2}}{35ac^2(c-ic \tan(e+fx))^{5/2}} - \frac{i(a+ia \tan(e+fx))^{5/2}}{7ac(c-ic \tan(e+fx))^{7/2}} \right)}{9c} - \frac{(B+iA)(a+ia \tan(e+fx))^{5/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{f}$$

input `Int[((a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(9/2), x]`

output `(a*c*(-1/9*((I*A + B)*(a + I*a*Tan[e + f*x])^(5/2))/(a*c*(c - I*c*Tan[e + f*x])^(9/2)) + ((2*A + (7*I)*B)*((-1/7*I)*(a + I*a*Tan[e + f*x])^(5/2))/(a*c*(c - I*c*Tan[e + f*x])^(7/2)) - ((I/35)*(a + I*a*Tan[e + f*x])^(5/2))/(a*c^2*(c - I*c*Tan[e + f*x])^(5/2)))/(9*c))/f`

3.813.3.1 Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x],
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.813.4 Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.77

method	result
risch	$-\frac{a^2 \sqrt{\frac{a e^{2i(fx+e)}}{e^{2i(fx+e)}+1}} (35iA e^{8i(fx+e)}+35B e^{8i(fx+e)}+90iA e^{6i(fx+e)}+63iA e^{4i(fx+e)}-63B e^{4i(fx+e)})}{1260c^4 \sqrt{\frac{c}{e^{2i(fx+e)}+1}} f}$
derivativedivides	$-\frac{i\sqrt{a(1+i\tan(fx+e))} \sqrt{-c(i\tan(fx+e)-1)} a^2 (1+\tan(fx+e)^2) (-47A+2iA\tan(fx+e)^3-12A\tan(fx+e)^2-33iA\tan(fx+e))}{315f c^5 (i+\tan(fx+e))^6}$
default	$-\frac{i\sqrt{a(1+i\tan(fx+e))} \sqrt{-c(i\tan(fx+e)-1)} a^2 (1+\tan(fx+e)^2) (-47A+2iA\tan(fx+e)^3-12A\tan(fx+e)^2-33iA\tan(fx+e))}{315f c^5 (i+\tan(fx+e))^6}$
parts	$-\frac{iA\sqrt{a(1+i\tan(fx+e))} \sqrt{-c(i\tan(fx+e)-1)} a^2 (1+\tan(fx+e)^2) (-47-33i\tan(fx+e)-12\tan(fx+e)^2+2i\tan(fx+e)^3)}{315f c^5 (i+\tan(fx+e))^6}$

```
input int((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(9/2),x,m
ethod=_RETURNVERBOSE)
```

```
output -1/1260*a^2/c^4*(a*exp(2*I*(f*x+e))/(exp(2*I*(f*x+e))+1))^(1/2)/(c/(exp(2*
I*(f*x+e))+1))^(1/2)/f*(35*I*A*exp(8*I*(f*x+e))+35*B*exp(8*I*(f*x+e))+90*I
*A*exp(6*I*(f*x+e))+63*I*A*exp(4*I*(f*x+e))-63*B*exp(4*I*(f*x+e)))
```

3.813.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.81

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx =$$

$$\frac{(35(iA + B)a^2 e^{(11i fx + 11ie)} + 5(25iA + 7B)a^2 e^{(9i fx + 9ie)} + 9(17iA - 7B)a^2 e^{(7i fx + 7ie)} + 63(iA - B)a^2 e^{(5i fx + 5ie)})}{1260 c^5 f}$$

3.813. $\int \frac{(a+ia \tan(e+fx))^{5/2} (A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{9/2}} dx$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(9/2),x, algorithm="fricas")`

output `-1/1260*(35*(I*A + B)*a^2*e^(11*I*f*x + 11*I*e) + 5*(25*I*A + 7*B)*a^2*e^(9*I*f*x + 9*I*e) + 9*(17*I*A - 7*B)*a^2*e^(7*I*f*x + 7*I*e) + 63*(I*A - B)*a^2*e^(5*I*f*x + 5*I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^5*f)`

3.813.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{9/2}} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(9/2),x)`

output `Timed out`

3.813.7 Maxima [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.28

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{9/2}} dx = \frac{(35(-iA - B)a^2 \cos(\frac{9}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e))) - 90IAa^2 \cos(\frac{7}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e))) + 63(-IA + B)a^2 \cos(\frac{5}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e))) + 35(A - IB)a^2 \sin(\frac{9}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e))) + 90Aa^2 \sin(\frac{7}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e))) + 63(A + IB)a^2 \sin(\frac{5}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)))}{c^{9/2}}$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(9/2),x, algorithm="maxima")`

output `1/1260*(35*(-I*A - B)*a^2*cos(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 90*I*A*a^2*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 63*(-I*A + B)*a^2*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 35*(A - I*B)*a^2*sin(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 90*A*a^2*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 63*(A + I*B)*a^2*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sqrt(a)/(c^(9/2)*f)`

3.813. $\int \frac{(a+ia \tan(e+fx))^{5/2} (A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{9/2}} dx$

3.813.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{5/2}}{(-ic \tan(fx + e) + c)^{9/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(9/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(5/2)/(-I*c*tan(f*x + e) + c)^(9/2), x)`

3.813.9 Mupad [B] (verification not implemented)

Time = 10.56 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.40

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx =$$

$$a^2 \sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)i)}{\cos(2e+2fx)+1}} (A \cos(4e + 4fx) 63i + A \cos(6e + 6fx) 90i + A \cos(8e + 8fx) 35i - 63B \cos(4e + 4fx) + 35B \cos(8e + 8fx) - 63A \sin(4e + 4fx) - 90A \sin(6e + 6fx) - 35A \sin(8e + 8fx) - B \sin(4e + 4fx) + B \sin(8e + 8fx) 35i) / (1260c^4 f ((c(\cos(2e + 2fx) - \sin(2e + 2fx)i + 1)) / (\cos(2e + 2fx) + 1))^{1/2})$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(5/2))/(c - c*tan(e + f*x)*1i)^(9/2),x)`

output `-(a^2*((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*cos(4*e + 4*f*x)*63i + A*cos(6*e + 6*f*x)*90i + A*cos(8*e + 8*f*x)*35i - 63*B*cos(4*e + 4*f*x) + 35*B*cos(8*e + 8*f*x) - 63*A*sin(4*e + 4*f*x) - 90*A*sin(6*e + 6*f*x) - 35*A*sin(8*e + 8*f*x) - B*sin(4*e + 4*f*x)*63i + B*sin(8*e + 8*f*x)*35i))/(1260*c^4*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

3.814
$$\int \frac{(a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{11/2}} dx$$

3.814.1 Optimal result	7329
3.814.2 Mathematica [A] (verified)	7329
3.814.3 Rubi [A] (verified)	7330
3.814.4 Maple [A] (verified)	7333
3.814.5 Fricas [A] (verification not implemented)	7333
3.814.6 Sympy [F(-1)]	7334
3.814.7 Maxima [A] (verification not implemented)	7334
3.814.8 Giac [F]	7335
3.814.9 Mupad [B] (verification not implemented)	7335

3.814.1 Optimal result

Integrand size = 45, antiderivative size = 208

$$\int \frac{(a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{11/2}} dx =$$

$$\frac{(iA + B)(a + ia \tan(e + fx))^{5/2}}{11f(c - ic \tan(e + fx))^{11/2}} - \frac{(3iA - 8B)(a + ia \tan(e + fx))^{5/2}}{99cf(c - ic \tan(e + fx))^{9/2}}$$

$$- \frac{2(3iA - 8B)(a + ia \tan(e + fx))^{5/2}}{693c^2f(c - ic \tan(e + fx))^{7/2}} - \frac{2(3iA - 8B)(a + ia \tan(e + fx))^{5/2}}{3465c^3f(c - ic \tan(e + fx))^{5/2}}$$

output

```
-1/11*(I*A+B)*(a+I*a*tan(f*x+e))^(5/2)/f/(c-I*c*tan(f*x+e))^(11/2)-1/99*(3
*I*A-8*B)*(a+I*a*tan(f*x+e))^(5/2)/c/f/(c-I*c*tan(f*x+e))^(9/2)-2/693*(3*I
*A-8*B)*(a+I*a*tan(f*x+e))^(5/2)/c^2/f/(c-I*c*tan(f*x+e))^(7/2)-2/3465*(3*
I*A-8*B)*(a+I*a*tan(f*x+e))^(5/2)/c^3/f/(c-I*c*tan(f*x+e))^(5/2)
```

3.814.2 Mathematica [A] (verified)

Time = 16.38 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.75

$$\int \frac{(a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{11/2}} dx = \frac{a^2 \cos(e + fx)(55(-24iA + B) \cos(e + fx) + 63(-8iA + B) \sin(e + fx))}{(c - ic \tan(e + fx))^{11/2}}$$

input

```
Integrate[((a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan
[e + f*x])^(11/2), x]
```


output $(a^2 \cos[e + f x] * (55 * ((-24 * I) * A + B) * \cos[e + f x] + 63 * ((-8 * I) * A + 3 * B) * \cos[3 * (e + f x)] - (3 * A + (8 * I) * B) * (55 * \sin[e + f x] + 63 * \sin[3 * (e + f x)])) * (\cos[8 * e + 10 * f x] + I * \sin[8 * e + 10 * f x]) * \sqrt{a + I * a * \tan[e + f x]} * \sqrt{c - I * c * \tan[e + f x]} / (13860 * c^6 * f * (\cos[f x] + I * \sin[f x])^2)$

3.814.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 87, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{11/2}} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{11/2}} dx$$

↓ 4071

$$\frac{ac \int \frac{(i \tan(e + fx) a + a)^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} d \tan(e + fx)}{f}$$

↓ 87

$$ac \left(\frac{(3A + 8iB) \int \frac{(i \tan(e + fx) a + a)^{3/2}}{(c - ic \tan(e + fx))^{11/2}} d \tan(e + fx)}{11c} - \frac{(B + iA)(a + ia \tan(e + fx))^{5/2}}{11ac(c - ic \tan(e + fx))^{11/2}} \right)$$

↓ 55

$$ac \left(\frac{(3A + 8iB) \left(\frac{2 \int \frac{(i \tan(e + fx) a + a)^{3/2}}{(c - ic \tan(e + fx))^{9/2}} d \tan(e + fx)}{9c} - \frac{i(a + ia \tan(e + fx))^{5/2}}{9ac(c - ic \tan(e + fx))^{9/2}} \right)}{11c} - \frac{(B + iA)(a + ia \tan(e + fx))^{5/2}}{11ac(c - ic \tan(e + fx))^{11/2}} \right)$$

↓ 55

3.814. $\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{11/2}} dx$

$$\begin{array}{c}
 \left(\frac{(3A+8iB) \left(\frac{2 \left(\int \frac{(i \tan(e+fx)a+a)^{3/2}}{(c-ic \tan(e+fx))^{7/2}} d \tan(e+fx) - \frac{i(a+ia \tan(e+fx))^{5/2}}{7ac(c-ic \tan(e+fx))^{7/2}} \right)}{9c} - \frac{i(a+ia \tan(e+fx))^{5/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{(B+iA)(a+ia \tan(e+fx))^{5/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{ac} \\
 \hline
 f \\
 \downarrow 48 \\
 \left(\frac{(3A+8iB) \left(\frac{2 \left(-\frac{i(a+ia \tan(e+fx))^{5/2}}{35ac^2(c-ic \tan(e+fx))^{5/2}} - \frac{i(a+ia \tan(e+fx))^{5/2}}{7ac(c-ic \tan(e+fx))^{7/2}} \right)}{9c} - \frac{i(a+ia \tan(e+fx))^{5/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{(B+iA)(a+ia \tan(e+fx))^{5/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{ac} \\
 \hline
 f
 \end{array}$$

input `Int[((a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(11/2), x]`

output `(a*c*(-1/11*((I*A + B)*(a + I*a*Tan[e + f*x])^(5/2))/(a*c*(c - I*c*Tan[e + f*x])^(11/2)) + ((3*A + (8*I)*B)*(((1/9*I)*(a + I*a*Tan[e + f*x])^(5/2))/(a*c*(c - I*c*Tan[e + f*x])^(9/2)) + (2*(((1/7*I)*(a + I*a*Tan[e + f*x])^(5/2))/(a*c*(c - I*c*Tan[e + f*x])^(7/2)) - ((I/35)*(a + I*a*Tan[e + f*x])^(5/2))/(a*c^2*(c - I*c*Tan[e + f*x])^(5/2))))/(9*c)))/(11*c))/f`

3.814.3.1 Defintions of rubi rules used

- rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`
- rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
.), x] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]`

3.814.4 Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.75

method	result
risch	$-\frac{a^2 \sqrt{\frac{a e^{2i(fx+e)}}{e^{2i(fx+e)}+1}} (315iA e^{10i(fx+e)}+315B e^{10i(fx+e)}+1155iA e^{8i(fx+e)}+385B e^{8i(fx+e)}+1485iA e^{6i(fx+e)}-495B e^{4i(fx+e)})}{27720c^5 \sqrt{\frac{c}{e^{2i(fx+e)}+1}} f}$
derivativedivides	$-\frac{i \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^2 (1+\tan(fx+e)^2) (6iA \tan(fx+e)^4-112iB \tan(fx+e)^3-16B \tan(fx+e)^2)}{3465 f c^6 (i+\tan(fx+e))^7}$
default	$-\frac{i \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^2 (1+\tan(fx+e)^2) (6iA \tan(fx+e)^4-112iB \tan(fx+e)^3-16B \tan(fx+e)^2)}{3465 f c^6 (i+\tan(fx+e))^7}$
parts	$-\frac{iA \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^2 (1+\tan(fx+e)^2) (2i \tan(fx+e)^4-45i \tan(fx+e)^2-14 \tan(fx+e)^3)}{1155 f c^6 (i+\tan(fx+e))^7}$

input `int((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(11/2),x,
method=_RETURNVERBOSE)`

output `-1/27720*a^2/c^5*(a*exp(2*I*(f*x+e))/(exp(2*I*(f*x+e))+1))^(1/2)/(c/(exp(2
I(f*x+e))+1))^(1/2)/f*(315*I*A*exp(10*I*(f*x+e))+315*B*exp(10*I*(f*x+e))
+1155*I*A*exp(8*I*(f*x+e))+385*B*exp(8*I*(f*x+e))+1485*I*A*exp(6*I*(f*x+e))
)-495*B*exp(6*I*(f*x+e))+693*I*A*exp(4*I*(f*x+e))-693*B*exp(4*I*(f*x+e))`

3.814.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.70

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx =$$

$$\frac{(315(iA + B)a^2 e^{13i fx + 13ie} + 70(21iA + 10B)a^2 e^{(11i fx + 11ie)} + 110(24iA - B)a^2 e^{(9i fx + 9ie)} + 198(11iA + B)a^2 e^{(7i fx + 7ie)})}{27720 c^6 f}$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(11
/2),x, algorithm="fracas")`

3.814. $\int \frac{(a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{11/2}} dx$

output $-1/27720*(315*(I*A + B)*a^2*e^{(13*I*f*x + 13*I*e)} + 70*(21*I*A + 10*B)*a^2*e^{(11*I*f*x + 11*I*e)} + 110*(24*I*A - B)*a^2*e^{(9*I*f*x + 9*I*e)} + 198*(11*I*A - 6*B)*a^2*e^{(7*I*f*x + 7*I*e)} + 693*(I*A - B)*a^2*e^{(5*I*f*x + 5*I*e)})*sqrt(a/(e^{(2*I*f*x + 2*I*e)} + 1))*sqrt(c/(e^{(2*I*f*x + 2*I*e)} + 1))/(c^{6*f})$

3.814.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{11/2}} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(11/2),x)`

output Timed out

3.814.7 Maxima [A] (verification not implemented)

Time = 0.61 (sec) , antiderivative size = 276, normalized size of antiderivative = 1.33

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{11/2}} dx = \frac{(315(-iA - B)a^2 \cos(\frac{11}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 385(-3IA - B)a^2 \cos(9/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 495(-3IA + B)a^2 \cos(7/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 693(-IA + B)a^2 \cos(5/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 315(A - IB)a^2 \sin(11/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 385(3A - IB)a^2 \sin(9/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 495(3A + IB)a^2 \sin(7/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 693(A + IB)a^2 \sin(5/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e))) * sqrt(a) / (c^{(11/2)*f})$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(11/2),x, algorithm="maxima")`

output $1/27720*(315*(-I*A - B)*a^2*\cos(11/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 385*(-3*I*A - B)*a^2*\cos(9/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 495*(-3*I*A + B)*a^2*\cos(7/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 693*(-I*A + B)*a^2*\cos(5/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 315*(A - I*B)*a^2*\sin(11/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 385*(3*A - I*B)*a^2*\sin(9/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 495*(3*A + I*B)*a^2*\sin(7/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 693*(A + I*B)*a^2*\sin(5/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))))*sqrt(a)/(c^{(11/2)*f})$

3.814.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{5/2}}{(-ictan(fx + e) + c)^{11/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(11/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(5/2)/(-I*c*tan(f*x + e) + c)^(11/2), x)`

3.814.9 Mupad [B] (verification not implemented)

Time = 12.41 (sec) , antiderivative size = 292, normalized size of antiderivative = 1.40

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx =$$

$$a^2 \sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}} (A \cos(4e + 4fx) 693i + A \cos(6e + 6fx) 1485i + A \cos(8e + 8fx) 1155i + A \cos(10e + 10fx) 315i - 693B \cos(4e + 4fx) - 495B \cos(6e + 6fx) + 385B \cos(8e + 8fx) + 315B \cos(10e + 10fx) - 693A \sin(4e + 4fx) - 1485A \sin(6e + 6fx) - 1155A \sin(8e + 8fx) - 315A \sin(10e + 10fx) - B \sin(4e + 4fx) 693i - B \sin(6e + 6fx) 495i + B \sin(8e + 8fx) 385i + B \sin(10e + 10fx) 315i) / (27720c^5 f ((c*(\cos(2e + 2fx) - \sin(2e + 2fx)*1i + 1)) / (\cos(2e + 2fx) + 1))^{1/2})$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(5/2))/(c - c*tan(e + f*x)*1i)^(11/2),x)`

output `-(a^2*((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*cos(4*e + 4*f*x)*693i + A*cos(6*e + 6*f*x)*1485i + A*cos(8*e + 8*f*x)*1155i + A*cos(10*e + 10*f*x)*315i - 693*B*cos(4*e + 4*f*x) - 495*B*cos(6*e + 6*f*x) + 385*B*cos(8*e + 8*f*x) + 315*B*cos(10*e + 10*f*x) - 693*A*sin(4*e + 4*f*x) - 1485*A*sin(6*e + 6*f*x) - 1155*A*sin(8*e + 8*f*x) - 315*A*sin(10*e + 10*f*x) - B*sin(4*e + 4*f*x)*693i - B*sin(6*e + 6*f*x)*495i + B*sin(8*e + 8*f*x)*385i + B*sin(10*e + 10*f*x)*315i))/(27720*c^5*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

3.815
$$\int \frac{(a+ia \tan(e+fx))^{5/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{13/2}} dx$$

3.815.1 Optimal result 7336
 3.815.2 Mathematica [B] (verified) 7337
 3.815.3 Rubi [A] (verified) 7337
 3.815.4 Maple [A] (verified) 7341
 3.815.5 Fricas [A] (verification not implemented) 7341
 3.815.6 Sympy [F(-1)] 7342
 3.815.7 Maxima [A] (verification not implemented) 7342
 3.815.8 Giac [F] 7343
 3.815.9 Mupad [B] (verification not implemented) 7343

3.815.1 Optimal result

Integrand size = 45, antiderivative size = 261

$$\int \frac{(a + ia \tan(e + fx))^{5/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} dx =$$

$$\frac{(iA + B)(a + ia \tan(e + fx))^{5/2}}{13f(c - ic \tan(e + fx))^{13/2}} - \frac{(4iA - 9B)(a + ia \tan(e + fx))^{5/2}}{143cf(c - ic \tan(e + fx))^{11/2}}$$

$$- \frac{(4iA - 9B)(a + ia \tan(e + fx))^{5/2}}{429c^2f(c - ic \tan(e + fx))^{9/2}} - \frac{2(4iA - 9B)(a + ia \tan(e + fx))^{5/2}}{3003c^3f(c - ic \tan(e + fx))^{7/2}}$$

$$- \frac{2(4iA - 9B)(a + ia \tan(e + fx))^{5/2}}{15015c^4f(c - ic \tan(e + fx))^{5/2}}$$

output

```
-1/13*(I*A+B)*(a+I*a*tan(f*x+e))^(5/2)/f/(c-I*c*tan(f*x+e))^(13/2)-1/143*(
4*I*A-9*B)*(a+I*a*tan(f*x+e))^(5/2)/c/f/(c-I*c*tan(f*x+e))^(11/2)-1/429*(4
*I*A-9*B)*(a+I*a*tan(f*x+e))^(5/2)/c^2/f/(c-I*c*tan(f*x+e))^(9/2)-2/3003*(
4*I*A-9*B)*(a+I*a*tan(f*x+e))^(5/2)/c^3/f/(c-I*c*tan(f*x+e))^(7/2)-2/15015
*(4*I*A-9*B)*(a+I*a*tan(f*x+e))^(5/2)/c^4/f/(c-I*c*tan(f*x+e))^(5/2)
```

3.815.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 577 vs. $2(261) = 522$.

Time = 19.17 (sec) , antiderivative size = 577, normalized size of antiderivative = 2.21

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} dx = \frac{\cos^3(e + fx) \left((-iA + B) \cos(4fx) \left(\frac{\cos(2e)}{160c^7} + \frac{i \sin(2e)}{160c^7} \right) \right)}{}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(13/2),x]`

output `(Cos[e + f*x]^3*(((-I)*A + B)*Cos[4*f*x]*(Cos[2*e]/(160*c^7) + ((I/160)*Sin[2*e])/c^7) + ((-27*I)*A + 17*B)*Cos[6*f*x]*(Cos[4*e]/(1120*c^7) + ((I/1120)*Sin[4*e])/c^7) + ((-13*I)*A + 3*B)*Cos[8*f*x]*(Cos[6*e]/(336*c^7) + ((I/336)*Sin[6*e])/c^7) + (17*A - (3*I)*B)*Cos[10*f*x]*(((-1/528*I)*Cos[8*e])/c^7 + Sin[8*e]/(528*c^7)) + (63*A - (37*I)*B)*Cos[12*f*x]*(((-1/4576*I)*Cos[10*e])/c^7 + Sin[10*e]/(4576*c^7)) + (A - I*B)*Cos[14*f*x]*(((-1/416*I)*Cos[12*e])/c^7 + Sin[12*e]/(416*c^7)) + (A + I*B)*(Cos[2*e]/(160*c^7) + ((I/160)*Sin[2*e])/c^7)*Sin[4*f*x] + (27*A + (17*I)*B)*(Cos[4*e]/(1120*c^7) + ((I/1120)*Sin[4*e])/c^7)*Sin[6*f*x] + (13*A + (3*I)*B)*(Cos[6*e]/(336*c^7) + ((I/336)*Sin[6*e])/c^7)*Sin[8*f*x] + (17*A - (3*I)*B)*(Cos[8*e]/(528*c^7) + ((I/528)*Sin[8*e])/c^7)*Sin[10*f*x] + (63*A - (37*I)*B)*(Cos[10*e]/(4576*c^7) + ((I/4576)*Sin[10*e])/c^7)*Sin[12*f*x] + (A - I*B)*(Cos[12*e]/(416*c^7) + ((I/416)*Sin[12*e])/c^7)*Sin[14*f*x])*Sqrt[Sec[e + f*x]*(c*Cos[e + f*x] - I*c*Sin[e + f*x])]*(a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/(f*(Cos[f*x] + I*Sin[f*x])^2*(A*Cos[e + f*x] + B*Sin[e + f*x]))]`

3.815.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 87, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} dx$$

3.815. $\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} dx$

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \quad ac \int \frac{(i \tan(e + fx)a + a)^{3/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{15/2}} d \tan(e + fx) \\
 & \quad \quad \quad \downarrow \text{4071} \\
 & \quad \quad \quad ac \left(\frac{(4A + 9iB) \int \frac{(i \tan(e + fx)a + a)^{3/2}}{(c - ic \tan(e + fx))^{13/2}} d \tan(e + fx)}{13c} - \frac{(B + iA)(a + ia \tan(e + fx))^{5/2}}{13ac(c - ic \tan(e + fx))^{13/2}} \right) \\
 & \quad \quad \quad \quad \quad \quad \downarrow \text{87} \\
 & \quad \quad \quad \quad \quad \quad ac \left(\frac{(4A + 9iB) \left(\frac{3 \int \frac{(i \tan(e + fx)a + a)^{3/2}}{(c - ic \tan(e + fx))^{11/2}} d \tan(e + fx)}{11c} - \frac{i(a + ia \tan(e + fx))^{5/2}}{11ac(c - ic \tan(e + fx))^{11/2}} \right)}{13c} - \frac{(B + iA)(a + ia \tan(e + fx))^{5/2}}{13ac(c - ic \tan(e + fx))^{13/2}} \right) \\
 & \quad \quad \quad \quad \quad \quad \quad \quad \quad \downarrow \text{55} \\
 & \quad \quad \quad \quad \quad \quad \quad \quad \quad ac \left(\frac{(4A + 9iB) \left(\frac{3 \left(\frac{2 \int \frac{(i \tan(e + fx)a + a)^{3/2}}{(c - ic \tan(e + fx))^{9/2}} d \tan(e + fx)}{9c} - \frac{i(a + ia \tan(e + fx))^{5/2}}{9ac(c - ic \tan(e + fx))^{9/2}} \right)}{11c} - \frac{i(a + ia \tan(e + fx))^{5/2}}{11ac(c - ic \tan(e + fx))^{11/2}} \right)}{13c} - \frac{(B + iA)(a + ia \tan(e + fx))^{5/2}}{13ac(c - ic \tan(e + fx))^{13/2}} \right) \\
 & \quad \quad \quad \quad \quad \quad \quad \quad \quad \quad \quad \quad \downarrow \text{55}
 \end{aligned}$$

3.815. $\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} dx$

$$ac \left(\frac{(4A+9iB) \left(\frac{2 \left(\int \frac{(i \tan(e+fx)a+a)^{3/2}}{(c-ic \tan(e+fx))^{7/2}} d \tan(e+fx) - \frac{i(a+ia \tan(e+fx))^{5/2}}{7ac(c-ic \tan(e+fx))^{7/2}} \right)}{9c} - \frac{i(a+ia \tan(e+fx))^{5/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{i(a+ia \tan(e+fx))^{5/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{13c} \right) - \frac{(B+iA)(a+ia \tan(e+fx))^{5/2}}{13ac(c-ic \tan(e+fx))^{13/2}}$$

f

↓ 48

$$ac \left(\frac{(4A+9iB) \left(\frac{2 \left(-\frac{i(a+ia \tan(e+fx))^{5/2}}{35ac^2(c-ic \tan(e+fx))^{5/2}} - \frac{i(a+ia \tan(e+fx))^{5/2}}{7ac(c-ic \tan(e+fx))^{7/2}} \right)}{9c} - \frac{i(a+ia \tan(e+fx))^{5/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{i(a+ia \tan(e+fx))^{5/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{13c} \right) - \frac{(B+iA)(a+ia \tan(e+fx))^{5/2}}{13ac(c-ic \tan(e+fx))^{13/2}}$$

f

```
input Int[((a + I*a*Tan[e + f*x])^(5/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(13/2), x]
```

```
output (a*c*(-1/13*((I*A + B)*(a + I*a*Tan[e + f*x])^(5/2))/(a*c*(c - I*c*Tan[e +
f*x])^(13/2)) + ((4*A + (9*I)*B)*((-1/11*I)*(a + I*a*Tan[e + f*x])^(5/2)
)/(a*c*(c - I*c*Tan[e + f*x])^(11/2)) + (3*((-1/9*I)*(a + I*a*Tan[e + f*x
])^(5/2))/(a*c*(c - I*c*Tan[e + f*x])^(9/2)) + (2*((-1/7*I)*(a + I*a*Tan[
e + f*x])^(5/2))/(a*c*(c - I*c*Tan[e + f*x])^(7/2)) - ((I/35)*(a + I*a*Tan
[e + f*x])^(5/2))/(a*c^2*(c - I*c*Tan[e + f*x])^(5/2))))/(9*c)))/(11*c)))/(
(13*c)))/f
```

3.815.3.1 Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 55 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.815.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.65

method	result
risch	$-\frac{a^2 \sqrt{\frac{a e^{2i(fx+e)}}{e^{2i(fx+e)}+1}} (1155iA e^{12i(fx+e)}+1155B e^{12i(fx+e)}+5460iA e^{10i(fx+e)}+2730B e^{10i(fx+e)}+10010iA e^{8i(fx+e)}+8580iA e^{6i(fx+e)}-4290B e^{6i(fx+e)}+3003iA e^{4i(fx+e)}-3003B e^{4i(fx+e)})}{240240c^6 \sqrt{\frac{c}{e^{2i(fx+e)}+1}} f}$
derivativedivides	$\frac{\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^2 (1+\tan(fx+e)^2) (18iB \tan(fx+e)^5+64iA \tan(fx+e)^4+8A \tan(fx+e)^5)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^2 (1+\tan(fx+e)^2) (18iB \tan(fx+e)^5+64iA \tan(fx+e)^4+8A \tan(fx+e)^5)}{\dots}$
parts	$-\frac{iA \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^2 (1+\tan(fx+e)^2) (8i \tan(fx+e)^5-236i \tan(fx+e)^3-64 \tan(fx+e))}{15015f c^7 (i+\tan(fx+e))^8}$

```
input int((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(13/2),x,
method=_RETURNVERBOSE)
```

```
output -1/240240*a^2/c^6*(a*exp(2*I*(f*x+e))/(exp(2*I*(f*x+e))+1))^(1/2)/(c/(exp(
2*I*(f*x+e))+1))^(1/2)/f*(1155*I*A*exp(12*I*(f*x+e))+1155*B*exp(12*I*(f*x+
e))+5460*I*A*exp(10*I*(f*x+e))+2730*B*exp(10*I*(f*x+e))+10010*I*A*exp(8*I*
(f*x+e))+8580*I*A*exp(6*I*(f*x+e))-4290*B*exp(6*I*(f*x+e))+3003*I*A*exp(4*
I*(f*x+e))-3003*B*exp(4*I*(f*x+e)))
```

3.815.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.64

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} dx =$$

$$\frac{(1155 (i A + B) a^2 e^{(15i fx + 15i e)} + 105 (63i A + 37 B) a^2 e^{(13i fx + 13i e)} + 910 (17i A + 3 B) a^2 e^{(11i fx + 11i e)} + 14 \dots)}{\dots}$$

```
input integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(13
/2),x, algorithm="fracas")
```

output $-1/240240*(1155*(I*A + B)*a^2*e^{(15*I*f*x + 15*I*e)} + 105*(63*I*A + 37*B)*a^2*e^{(13*I*f*x + 13*I*e)} + 910*(17*I*A + 3*B)*a^2*e^{(11*I*f*x + 11*I*e)} + 1430*(13*I*A - 3*B)*a^2*e^{(9*I*f*x + 9*I*e)} + 429*(27*I*A - 17*B)*a^2*e^{(7*I*f*x + 7*I*e)} + 3003*(I*A - B)*a^2*e^{(5*I*f*x + 5*I*e)})*sqrt(a/(e^{(2*I*f*x + 2*I*e)} + 1))*sqrt(c/(e^{(2*I*f*x + 2*I*e)} + 1))/(c^7*f)$

3.815.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(13/2),x)`

output Timed out

3.815.7 Maxima [A] (verification not implemented)

Time = 0.60 (sec) , antiderivative size = 332, normalized size of antiderivative = 1.27

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} dx = \frac{(1155(-iA - B)a^2 \cos(\frac{13}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 2730(-2iA - B)a^2 \cos(11/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) - 10010iAa^2 \cos(9/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 4290(-2iA + B)a^2 \cos(7/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 3003(-iA + B)a^2 \cos(5/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 1155(A - iB)a^2 \sin(13/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 2730(2A - iB)a^2 \sin(11/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 10010Aa^2 \sin(9/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 4290(2A + iB)a^2 \sin(7/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 3003(A + iB)a^2 \sin(5/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e))) * sqrt(a) / (c^{(13/2)} * f)}$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(13/2),x, algorithm="maxima")`

output $1/240240*(1155*(-I*A - B)*a^2*\cos(13/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 2730*(-2*I*A - B)*a^2*\cos(11/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) - 10010*I*A*a^2*\cos(9/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 4290*(-2*I*A + B)*a^2*\cos(7/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 3003*(-I*A + B)*a^2*\cos(5/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 1155*(A - I*B)*a^2*\sin(13/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 2730*(2*A - I*B)*a^2*\sin(11/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 10010*A*a^2*\sin(9/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 4290*(2*A + I*B)*a^2*\sin(7/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 3003*(A + I*B)*a^2*\sin(5/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))))*sqrt(a)/(c^{(13/2)}*f)$

3.815. $\int \frac{(a+ia \tan(e+fx))^{5/2} (A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{13/2}} dx$

3.815.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{13/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{5/2}}{(-ictan(fx + e) + c)^{13/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(5/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(13/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(5/2)/(-I*c*tan(f*x + e) + c)^(13/2), x)`

3.815.9 Mupad [B] (verification not implemented)

Time = 13.40 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.73

$$\int \frac{(a + ia \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{13/2}} dx = \frac{\sqrt{a + \frac{a \sin(e+fx) \operatorname{li}}{\cos(e+fx)}} \left(\frac{a^2 e^{6i+fx 6i} (2A+B \operatorname{li}) \operatorname{li}}{56 c^6 f} + \frac{a^2 e^{10i+fx 10i} (2A-B \operatorname{li}) \operatorname{li}}{88 c^6 f} + \frac{A a^2 e^{8i+fx 8i} \operatorname{li}}{24 c^6 f} + \frac{a^2 e^{4i+fx 4i} (A+B \operatorname{li}) \operatorname{li}}{80 c^6 f} + \dots \right)}{\sqrt{c - \frac{c \sin(e+fx) \operatorname{li}}{\cos(e+fx)}}}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(5/2))/(c - c*tan(e + f*x)*1i)^(13/2),x)`

output `-((a + (a*sin(e + f*x)*1i)/cos(e + f*x))^(1/2))*((a^2*exp(e*6i + f*x*6i))*(2*A + B*1i)*1i)/(56*c^6*f) + (a^2*exp(e*10i + f*x*10i)*(2*A - B*1i)*1i)/(88*c^6*f) + (A*a^2*exp(e*8i + f*x*8i)*1i)/(24*c^6*f) + (a^2*exp(e*4i + f*x*4i)*(A + B*1i)*1i)/(80*c^6*f) + (a^2*exp(e*12i + f*x*12i)*(A - B*1i)*1i)/(208*c^6*f))/((c - (c*sin(e + f*x)*1i)/cos(e + f*x))^(1/2))`

3.816 $\int (a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{9/2} dx$

3.816.1 Optimal result	7344
3.816.2 Mathematica [A] (verified)	7345
3.816.3 Rubi [A] (verified)	7345
3.816.4 Maple [B] (verified)	7348
3.816.5 Fricas [B] (verification not implemented)	7349
3.816.6 Sympy [F(-1)]	7350
3.816.7 Maxima [B] (verification not implemented)	7351
3.816.8 Giac [F]	7352
3.816.9 Mupad [F(-1)]	7352

3.816.1 Optimal result

Integrand size = 45, antiderivative size = 350

$$\int (a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{9/2} dx =$$

$$-\frac{5a^{7/2}(8iA-B)c^{9/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{64f}$$

$$+\frac{5a^3(8A+iB)c^4 \tan(e+fx)\sqrt{a+ia \tan(e+fx)}\sqrt{c-ictan(e+fx)}}{128f}$$

$$+\frac{5a^2(8A+iB)c^3 \tan(e+fx)(a+ia \tan(e+fx))^{3/2}(c-ictan(e+fx))^{3/2}}{192f}$$

$$+\frac{a(8A+iB)c^2 \tan(e+fx)(a+ia \tan(e+fx))^{5/2}(c-ictan(e+fx))^{5/2}}{48f}$$

$$-\frac{(8iA-B)c(a+ia \tan(e+fx))^{7/2}(c-ictan(e+fx))^{7/2}}{56f}$$

$$+\frac{B(a+ia \tan(e+fx))^{7/2}(c-ictan(e+fx))^{9/2}}{8f}$$

output
$$\begin{aligned} & -5/64*a^{(7/2)}*(8*I*A-B)*c^{(9/2)}*\arctan(c^{(1/2)}*(a+I*a*\tan(f*x+e))^{(1/2)}/a^{(1/2)})/(c-I*c*\tan(f*x+e))^{(1/2)}/f+5/128*a^3*(8*A+I*B)*c^4*(a+I*a*\tan(f*x+e))^{(1/2)}*(c-I*c*\tan(f*x+e))^{(1/2)}*\tan(f*x+e)/f+5/192*a^2*(8*A+I*B)*c^3*\tan(f*x+e)*(a+I*a*\tan(f*x+e))^{(3/2)}*(c-I*c*\tan(f*x+e))^{(3/2)}/f+1/48*a*(8*A+I*B)*c^2*\tan(f*x+e)*(a+I*a*\tan(f*x+e))^{(5/2)}*(c-I*c*\tan(f*x+e))^{(5/2)}/f-1/56*(8*I*A-B)*c*(a+I*a*\tan(f*x+e))^{(7/2)}*(c-I*c*\tan(f*x+e))^{(7/2)}/f+1/8*B*(a+I*a*\tan(f*x+e))^{(7/2)}*(c-I*c*\tan(f*x+e))^{(9/2)}/f \end{aligned}$$

3.816.2 Mathematica [A] (verified)

Time = 9.43 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.69

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{9/2} dx = \frac{a^{7/2} c^5 \sqrt{1 - i \tan(e + fx)} \left(210(-8iA + B) \arcsin\left(\frac{\sqrt{a + ia \tan(e + fx)}}{\sqrt{2}\sqrt{a}}\right) \sqrt{a + ia \tan(e + fx)} \right)}{}$$

input `Integrate[(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(9/2),x]`

output
$$\begin{aligned} & (a^{(7/2)}*c^5*\text{Sqrt}[1 - I*\text{Tan}[e + f*x]]*(210*((-8*I)*A + B)*\text{ArcSin}[\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]/(\text{Sqrt}[2]*\text{Sqrt}[a])]*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]] + (\text{Sqrt}[a]*\text{Sec}[e + f*x]^6*\text{Sqrt}[1 - I*\text{Tan}[e + f*x]]*(1 + I*\text{Tan}[e + f*x])*(24576*((-I)*A + B) + 7*(8*A + I*B)*\text{Sec}[e + f*x]*(383*\text{Sin}[3*(e + f*x)] + 115*\text{Sin}[5*(e + f*x)] + 15*\text{Sin}[7*(e + f*x)]) + 7*(2264*A - (2789*I)*B)*\text{Tan}[e + f*x])]/64))/ (2688*f*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]]) \end{aligned}$$

3.816.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 318, normalized size of antiderivative = 0.91, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 4071, 90, 59, 40, 40, 40, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))^{7/2} (c - ic \tan(e + fx))^{9/2} (A + B \tan(e + fx)) dx$$

3.816. $\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{9/2} dx$

↓ 3042

$$\int (a + ia \tan(e + fx))^{7/2} (c - ic \tan(e + fx))^{9/2} (A + B \tan(e + fx)) dx$$

↓ 4071

$$\frac{ac \int (i \tan(e + fx)a + a)^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} d \tan(e + fx)}{f}$$

↓ 90

$$\frac{ac \left(\frac{1}{8} (8A + iB) \int (i \tan(e + fx)a + a)^{5/2} (c - ic \tan(e + fx))^{7/2} d \tan(e + fx) + \frac{B(a + ia \tan(e + fx))^{7/2} (c - ic \tan(e + fx))^{9/2}}{8ac} \right)}{f}$$

↓ 59

$$\frac{ac \left(\frac{1}{8} (8A + iB) \left(c \int (i \tan(e + fx)a + a)^{5/2} (c - ic \tan(e + fx))^{5/2} d \tan(e + fx) - \frac{i(a + ia \tan(e + fx))^{7/2} (c - ic \tan(e + fx))^{9/2}}{7a} \right) \right)}{f}$$

↓ 40

$$\frac{ac \left(\frac{1}{8} (8A + iB) \left(c \left(\frac{5}{6} ac \int (i \tan(e + fx)a + a)^{3/2} (c - ic \tan(e + fx))^{3/2} d \tan(e + fx) + \frac{1}{6} \tan(e + fx) (a + ia \tan(e + fx)) \right) \right) \right)}{f}$$

↓ 40

$$\frac{ac \left(\frac{1}{8} (8A + iB) \left(c \left(\frac{5}{6} ac \left(\frac{3}{4} ac \int \sqrt{i \tan(e + fx)a + a} \sqrt{c - ic \tan(e + fx)} d \tan(e + fx) + \frac{1}{4} \tan(e + fx) (a + ia \tan(e + fx)) \right) \right) \right) \right)}{f}$$

↓ 40

$$\frac{ac \left(\frac{1}{8} (8A + iB) \left(c \left(\frac{5}{6} ac \left(\frac{3}{4} ac \left(\frac{1}{2} ac \int \frac{1}{\sqrt{i \tan(e + fx)a + a} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \right) \right) \right) \right) \right)}{f}$$

↓ 45

$$\frac{ac \left(\frac{1}{8} (8A + iB) \left(c \left(\frac{5}{6} ac \left(\frac{3}{4} ac \left(ac \int \frac{1}{ia + \frac{ic(i \tan(e + fx)a + a)}{c - ic \tan(e + fx)}} d \frac{\sqrt{i \tan(e + fx)a + a}}{\sqrt{c - ic \tan(e + fx)}} + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)} \right) \right) \right) \right) \right)}{f}$$

↓ 218

$$ac\left(\frac{1}{8}(8A + iB)\left(c\left(\frac{5}{6}ac\left(\frac{3}{4}ac\left(\frac{1}{2}\tan(e + fx)\sqrt{a + ia\tan(e + fx)}\sqrt{c - ictan(e + fx)} - i\sqrt{a}\sqrt{c}\arctan\left(\frac{\sqrt{c}\sqrt{a+ia\tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)\right)\right)\right)\right)$$

input `Int[(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(9/2),x]`

output `(a*c*((B*(a + I*a*Tan[e + f*x])^(7/2)*(c - I*c*Tan[e + f*x])^(9/2))/(8*a*c) + ((8*A + I*B)*((-1/7*I)*(a + I*a*Tan[e + f*x])^(7/2)*(c - I*c*Tan[e + f*x])^(7/2))/a + c*((Tan[e + f*x]*(a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e + f*x])^(5/2))/6 + (5*a*c*((Tan[e + f*x]*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(3/2))/4 + (3*a*c*((-I)*Sqrt[a]*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])] + (Tan[e + f*x]*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/2))/4)/6))/8)/f`

3.816.3.1 Defintions of rubi rules used

rule 40 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^m/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`

rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 59 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[2*c*(n/(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0] && IGtQ[n + 1/2, 0] && LtQ[m, n]`

rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

3.816. $\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{9/2} dx$

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.816.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 603 vs. 2(289) = 578.

Time = 0.50 (sec) , antiderivative size = 604, normalized size of antiderivative = 1.73

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^3 c^4 \left(826 i B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)} \tan (f x+e)^3+105 i B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}\right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^3 c^4 \left(826 i B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)} \tan (f x+e)^3+105 i B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}\right)}{\dots}$
parts	$\frac{A \sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^3 c^4 \left(48 i \tan (f x+e)^6 \sqrt{a c\left(1+\tan (f x+e)^2\right)} \sqrt{a c+144 i \tan (f x+e)^4} \sqrt{a c}\right)}{\dots}$

input `int((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2),x,m
method=_RETURNVERBOSE)`

output

```

-1/2688/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^3*c^4*(
826*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^3+105*I*B*(a*c
)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+1152*I*A*(a*c)^(1/2)*(a*c*
(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^4-384*B*tan(f*x+e)^6*(a*c*(1+tan(f*x+e)
^2))^(1/2)*(a*c)^(1/2)+1152*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*t
an(f*x+e)^2-448*A*tan(f*x+e)^5*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)+38
4*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^6-1152*B*(a*c)^(
1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^4+336*I*B*(a*c)^(1/2)*(a*c*(1
+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^7-1456*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2
))^(1/2)*tan(f*x+e)^3+384*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)-105
*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1
/2))*a*c-1152*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+952*
I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^5-840*A*ln((a*c*ta
n(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c-1848*A
*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-384*B*(a*c)^(1/2)*(a*
c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2)/(a*c*(1+tan(f*x+e)^2))^(1/2)

```

3.816.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 882 vs. $2(270) = 540$.

Time = 0.29 (sec) , antiderivative size = 882, normalized size of antiderivative = 2.52

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{9/2} dx = \text{Too large to display}$$

input

```

integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/
2),x, algorithm="fracas")

```

output `1/5376*(105*sqrt((64*A^2 + 16*I*A*B - B^2)*a^7*c^9/f^2)*(f*e^(14*I*f*x + 14*I*e) + 7*f*e^(12*I*f*x + 12*I*e) + 21*f*e^(10*I*f*x + 10*I*e) + 35*f*e^(8*I*f*x + 8*I*e) + 35*f*e^(6*I*f*x + 6*I*e) + 21*f*e^(4*I*f*x + 4*I*e) + 7*f*e^(2*I*f*x + 2*I*e) + f)*log(4*(2*((8*I*A - B)*a^3*c^4*e^(3*I*f*x + 3*I*e) + (8*I*A - B)*a^3*c^4*e^(I*f*x + I*e)))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1)))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + sqrt((64*A^2 + 16*I*A*B - B^2)*a^7*c^9/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((8*I*A - B)*a^3*c^4*e^(2*I*f*x + 2*I*e) + (8*I*A - B)*a^3*c^4) - 105*sqrt((64*A^2 + 16*I*A*B - B^2)*a^7*c^9/f^2)*(f*e^(14*I*f*x + 14*I*e) + 7*f*e^(12*I*f*x + 12*I*e) + 21*f*e^(10*I*f*x + 10*I*e) + 35*f*e^(8*I*f*x + 8*I*e) + 35*f*e^(6*I*f*x + 6*I*e) + 21*f*e^(4*I*f*x + 4*I*e) + 7*f*e^(2*I*f*x + 2*I*e) + f)*log(4*(2*((8*I*A - B)*a^3*c^4*e^(3*I*f*x + 3*I*e) + (8*I*A - B)*a^3*c^4*e^(I*f*x + I*e)))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1)))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - sqrt((64*A^2 + 16*I*A*B - B^2)*a^7*c^9/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((8*I*A - B)*a^3*c^4*e^(2*I*f*x + 2*I*e) + (8*I*A - B)*a^3*c^4) - 4*(105*(8*I*A - B)*a^3*c^4*e^(15*I*f*x + 15*I*e) + 805*(8*I*A - B)*a^3*c^4*e^(13*I*f*x + 13*I*e) + 2681*(8*I*A - B)*a^3*c^4*e^(11*I*f*x + 11*I*e) + 5053*(8*I*A - B)*a^3*c^4*e^(9*I*f*x + 9*I*e) - (-8728*I*A + 44099*B)*a^3*c^4*e^(7*I*f*x + 7*I*e) + 2681*(-8*I*A + B)*a^3*c^4*e^(5*I*f*x + 5*I*e) + 805*(-8*I*A + B)*a^3*c^4*e^(3*I*f*x + 3*I*e) + 105*(-8*I*A + B)*a^3*c^4*e^(I*f*x + I*e))*sqr...`

3.816.6 Sympy [**F(-1)**]

Timed out.

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{9/2} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(9/2),x)`

output `Timed out`

3.816.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2603 vs. $2(270) = 540$.

Time = 19.78 (sec) , antiderivative size = 2603, normalized size of antiderivative = 7.44

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{9/2} dx = \text{Too large to display}$$

```
input integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2),x, algorithm="maxima")
```

```
output -86016*(420*(8*A + I*B)*a^3*c^4*cos(15/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 3220*(8*A + I*B)*a^3*c^4*cos(13/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 10724*(8*A + I*B)*a^3*c^4*cos(11/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 20212*(8*A + I*B)*a^3*c^4*cos(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 4*(8728*A + 44099*I*B)*a^3*c^4*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 10724*(8*A + I*B)*a^3*c^4*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 3220*(8*A + I*B)*a^3*c^4*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 420*(8*A + I*B)*a^3*c^4*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 420*(8*I*A - B)*a^3*c^4*sin(15/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 3220*(8*I*A - B)*a^3*c^4*sin(13/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 10724*(8*I*A - B)*a^3*c^4*sin(11/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 20212*(8*I*A - B)*a^3*c^4*sin(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 4*(8728*I*A - 44099*B)*a^3*c^4*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 10724*(-8*I*A + B)*a^3*c^4*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 3220*(-8*I*A + B)*a^3*c^4*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 420*(-8*I*A + B)*a^3*c^4*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 210*((8*A + I*B)*a^3*c^4*cos(16*f*x + 16*e) + 8*(8*A + I*B)*a^3*c^4*cos(14*f*x + 14*e) + 28*(8*A + I*B)*a^3*c^4*cos(12*f*x + 12*e) + 56*(8*A + I*B)*a^3*c^4*cos(10*f*x...
```

3.816.8 Giac [F]

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{9/2} dx = \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^{7/2} (-ic \tan(fx + e) + c)^{9/2} dx$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2),x, algorithm="giac")`

output `sage0*x`

3.816.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{9/2} dx = \int (A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{7/2} (c - c \tan(e + fx) li)^{9/2} dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2)*(c - c*tan(e + f*x)*1i)^(9/2),x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2)*(c - c*tan(e + f*x)*1i)^(9/2), x)`

3.817 $\int (a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{7/2} dx$

3.817.1 Optimal result	7353
3.817.2 Mathematica [A] (verified)	7354
3.817.3 Rubi [A] (verified)	7354
3.817.4 Maple [A] (verified)	7357
3.817.5 Fricas [B] (verification not implemented)	7357
3.817.6 Sympy [F(-1)]	7358
3.817.7 Maxima [B] (verification not implemented)	7359
3.817.8 Giac [F]	7360
3.817.9 Mupad [F(-1)]	7360

3.817.1 Optimal result

Integrand size = 45, antiderivative size = 267

$$\int (a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{7/2} dx =$$

$$\frac{5ia^{7/2}Ac^{7/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{8f}$$

$$+ \frac{5a^3Ac^3 \tan(e+fx)\sqrt{a+ia \tan(e+fx)}\sqrt{c-ictan(e+fx)}}{16f}$$

$$+ \frac{5a^2Ac^2 \tan(e+fx)(a+ia \tan(e+fx))^{3/2}(c-ictan(e+fx))^{3/2}}{24f}$$

$$+ \frac{aActan(e+fx)(a+ia \tan(e+fx))^{5/2}(c-ictan(e+fx))^{5/2}}{6f}$$

$$+ \frac{B(a+ia \tan(e+fx))^{7/2}(c-ictan(e+fx))^{7/2}}{7f}$$

output

```
-5/8*I*a^(7/2)*A*c^(7/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/f+5/16*a^3*A*c^3*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)*tan(f*x+e)/f+5/24*a^2*A*c^2*tan(f*x+e)*(a+I*a*tan(f*x+e))^(3/2)*(c-I*c*tan(f*x+e))^(3/2)/f+1/6*a*A*c*tan(f*x+e)*(a+I*a*tan(f*x+e))^(5/2)*(c-I*c*tan(f*x+e))^(5/2)/f+1/7*B*(a+I*a*tan(f*x+e))^(7/2)*(c-I*c*tan(f*x+e))^(7/2)/f
```


3.817.2 Mathematica [A] (verified)

Time = 8.56 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.60

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \frac{a^{7/2} c^4 \left(-\frac{6720A \arcsin\left(\frac{\sqrt{a+ia \tan(e+fx)}}{\sqrt{2}\sqrt{a}}\right) (i+\tan(e+fx))}{\sqrt{1-i \tan(e+fx)}} + \frac{\sqrt{a} \sec^8(e+fx) (1536B+1981A \sin(2(e+fx))+700A \sin(4(e+fx))+105A \sin(6(e+fx)))}{\sqrt{a+ia \tan(e+fx)}} \right)}{10752f \sqrt{c - ic \tan(e + fx)}}$$

```
input Integrate[(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2),x]
```

```
output (a^(7/2)*c^4*((-6720*A*ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a])]*(I + Tan[e + f*x]))/Sqrt[1 - I*Tan[e + f*x]] + (Sqrt[a]*Sec[e + f*x]^8*(1536*B + 1981*A*Sin[2*(e + f*x)] + 700*A*Sin[4*(e + f*x)] + 105*A*Sin[6*(e + f*x)]))/Sqrt[a + I*a*Tan[e + f*x]]))/(10752*f*Sqrt[c - I*c*Tan[e + f*x]])
```

3.817.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 261, normalized size of antiderivative = 0.98, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.178$, Rules used = {3042, 4071, 90, 40, 40, 40, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (a + ia \tan(e + fx))^{7/2} (c - ic \tan(e + fx))^{7/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{3042} \\ & \int (a + ia \tan(e + fx))^{7/2} (c - ic \tan(e + fx))^{7/2} (A + B \tan(e + fx)) dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int (i \tan(e + fx) a + a)^{5/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{90} \end{aligned}$$

$$\frac{ac \left(A \int (i \tan(e + fx)a + a)^{5/2} (c - ic \tan(e + fx))^{5/2} d \tan(e + fx) + \frac{B(a + ia \tan(e + fx))^{7/2} (c - ic \tan(e + fx))^{7/2}}{7ac} \right)}{f}$$

↓ 40

$$\frac{ac \left(A \left(\frac{5}{6} ac \int (i \tan(e + fx)a + a)^{3/2} (c - ic \tan(e + fx))^{3/2} d \tan(e + fx) + \frac{1}{6} \tan(e + fx) (a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{5/2} \right) \right)}{f}$$

↓ 40

$$\frac{ac \left(A \left(\frac{5}{6} ac \left(\frac{3}{4} ac \int \sqrt{i \tan(e + fx)a + a} \sqrt{c - ic \tan(e + fx)} d \tan(e + fx) + \frac{1}{4} \tan(e + fx) (a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2} \right) \right) \right)}$$

↓ 40

$$\frac{ac \left(A \left(\frac{5}{6} ac \left(\frac{3}{4} ac \left(\frac{1}{2} ac \int \frac{1}{\sqrt{i \tan(e + fx)a + a} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)} \right) \right) \right) \right)}$$

↓ 45

$$\frac{ac \left(A \left(\frac{5}{6} ac \left(\frac{3}{4} ac \left(ac \int \frac{1}{ia + \frac{ic(i \tan(e + fx)a + a)}{c - ic \tan(e + fx)}} d \frac{\sqrt{i \tan(e + fx)a + a}}{\sqrt{c - ic \tan(e + fx)}} + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)} \right) \right) \right) \right)}$$

↓ 218

$$\frac{ac \left(A \left(\frac{5}{6} ac \left(\frac{3}{4} ac \left(\frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)} - i \sqrt{a} \sqrt{c} \arctan \left(\frac{\sqrt{c} \sqrt{a + ia \tan(e + fx)}}{\sqrt{a} \sqrt{c - ic \tan(e + fx)}} \right) \right) \right) \right) \right) +$$

input `Int[(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2),x]`

output `(a*c*((B*(a + I*a*Tan[e + f*x])^(7/2)*(c - I*c*Tan[e + f*x])^(7/2))/(7*a*c) + A*((Tan[e + f*x]*(a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e + f*x])^(5/2))/6 + (5*a*c*((Tan[e + f*x]*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(3/2))/4 + (3*a*c*((-I)*Sqrt[a]*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])] + (Tan[e + f*x]*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/2))/4))/6))/f`

3.817.3.1 Defintions of rubi rules used

- rule 40 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^m/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`
- rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 90 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.817.4 Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.01

method	result
parts	$A\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^3c^3\left(8\tan(fx+e)^5\sqrt{ac(1+\tan(fx+e)^2)}\sqrt{ac+26\tan(fx+e)^3}\sqrt{ac(1+\tan(fx+e)^2)}\right)$
derivatividedivides	$\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^3c^3\left(48B\tan(fx+e)^6\sqrt{ac(1+\tan(fx+e)^2)}\sqrt{ac+56A\tan(fx+e)^5}\sqrt{ac(1+\tan(fx+e)^2)}\right)$
default	$\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^3c^3\left(48B\tan(fx+e)^6\sqrt{ac(1+\tan(fx+e)^2)}\sqrt{ac+56A\tan(fx+e)^5}\sqrt{ac(1+\tan(fx+e)^2)}\right)$

input `int((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x,method=_RETURNVERBOSE)`

output `1/48*A/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^3*c^3*(8*tan(f*x+e)^5*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)+26*tan(f*x+e)^3*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)+15*a*c*ln((a*c*tan(f*x+e)+(a*c)^(1/2))*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))+33*tan(f*x+e)*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2))/(a*c*(1+tan(f*x+e)^2))^(1/2)/(a*c)^(1/2)+1/7*B/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^3*c^3*(1+tan(f*x+e)^2)*(tan(f*x+e)^4+2*tan(f*x+e)^2+1)`

3.817.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 693 vs. 2(207) = 414.

Time = 0.27 (sec) , antiderivative size = 693, normalized size of antiderivative = 2.60

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx = \frac{105 \sqrt{\frac{A^2 a^7 c^7}{f^2}} (f e^{(12i fx + 12i e)} + 6 f e^{(10i fx + 10i e)} + 15 f e^{(8i fx + 8i e)} + 20 f e^{(6i fx + 6i e)} + 15 f e^{(4i fx + 4i e)} + 6 f e^{(2i fx + 2i e)} + f e^{(0i fx + 0i e)})}{f^2}$$

3.817. $\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2} dx$

```
input integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="fricas")
```

```
output 1/672*(105*sqrt(A^2*a^7*c^7/f^2)*(f*e^(12*I*f*x + 12*I*e) + 6*f*e^(10*I*f*x + 10*I*e) + 15*f*e^(8*I*f*x + 8*I*e) + 20*f*e^(6*I*f*x + 6*I*e) + 15*f*e^(4*I*f*x + 4*I*e) + 6*f*e^(2*I*f*x + 2*I*e) + f)*log(4*(2*(A*a^3*c^3*e^(3*I*f*x + 3*I*e) + A*a^3*c^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - sqrt(A^2*a^7*c^7/f^2)*(I*f*e^(2*I*f*x + 2*I*e) - I*f))/(A*a^3*c^3*e^(2*I*f*x + 2*I*e) + A*a^3*c^3)) - 105*sqrt(A^2*a^7*c^7/f^2)*(f*e^(12*I*f*x + 12*I*e) + 6*f*e^(10*I*f*x + 10*I*e) + 15*f*e^(8*I*f*x + 8*I*e) + 20*f*e^(6*I*f*x + 6*I*e) + 15*f*e^(4*I*f*x + 4*I*e) + 6*f*e^(2*I*f*x + 2*I*e) + f)*log(4*(2*(A*a^3*c^3*e^(3*I*f*x + 3*I*e) + A*a^3*c^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - sqrt(A^2*a^7*c^7/f^2)*(-I*f*e^(2*I*f*x + 2*I*e) + I*f))/(A*a^3*c^3*e^(2*I*f*x + 2*I*e) + A*a^3*c^3)) + 4*(-105*I*A*a^3*c^3*e^(13*I*f*x + 13*I*e) - 700*I*A*a^3*c^3*e^(11*I*f*x + 11*I*e) - 1981*I*A*a^3*c^3*e^(9*I*f*x + 9*I*e) + 3072*B*a^3*c^3*e^(7*I*f*x + 7*I*e) + 1981*I*A*a^3*c^3*e^(5*I*f*x + 5*I*e) + 700*I*A*a^3*c^3*e^(3*I*f*x + 3*I*e) + 105*I*A*a^3*c^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(f*e^(12*I*f*x + 12*I*e) + 6*f*e^(10*I*f*x + 10*I*e) + 15*f*e^(8*I*f*x + 8*I*e) + 20*f*e^(6*I*f*x + 6*I*e) + 15*f*e^(4*I*f*x + 4*I*e) + 6*f*e^(2*I*f*x + 2*I*e) + f)
```

3.817.6 Sympy [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{7/2} dx = \text{Timed out}$$

```
input integrate((a+I*a*tan(f*x+e))**(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(7/2),x)
```

```
output Timed out
```

3.817.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1901 vs. $2(207) = 414$.

Time = 2.74 (sec) , antiderivative size = 1901, normalized size of antiderivative = 7.12

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \text{Too large to display}$$

```
input integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="maxima")
```

```
output -(420*A*a^3*c^3*cos(13/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 2800*A*a^3*c^3*cos(11/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 7924*A*a^3*c^3*cos(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 12288*I*B*a^3*c^3*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 7924*A*a^3*c^3*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 2800*A*a^3*c^3*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 420*A*a^3*c^3*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 420*I*A*a^3*c^3*sin(13/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 2800*I*A*a^3*c^3*sin(11/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 7924*I*A*a^3*c^3*sin(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 12288*B*a^3*c^3*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 7924*I*A*a^3*c^3*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 2800*I*A*a^3*c^3*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 420*I*A*a^3*c^3*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 210*(A*a^3*c^3*cos(14*f*x + 14*e) + 7*A*a^3*c^3*cos(12*f*x + 12*e) + 21*A*a^3*c^3*cos(10*f*x + 10*e) + 35*A*a^3*c^3*cos(8*f*x + 8*e) + 35*A*a^3*c^3*cos(6*f*x + 6*e) + 21*A*a^3*c^3*cos(4*f*x + 4*e) + 7*A*a^3*c^3*cos(2*f*x + 2*e) + I*A*a^3*c^3*sin(14*f*x + 14*e) + 7*I*A*a^3*c^3*sin(12*f*x + 12*e) + 21*I*A*a^3*c^3*sin(10*f*x + 10*e) + 35*I*A*a^3*c^3*sin(8*f*x + 8*e) + 35*I*A*a^3*c^3*sin(6*f*x + 6*e) + 21*I*A*a^3*c^3*sin(4*f*x + 4*e) + 7*I*A*a^3*c^3*sin(2*f*x + 2*e) + A*a^3*c^3)...
```

3.817.8 Giac [F]

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^{7/2} (-ic \tan(fx + e) + c)^{7/2} dx$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2),x, algorithm="giac")`

output `sage0*x`

3.817.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{7/2} dx = \int (A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{7/2} (c - c \tan(e + fx) li)^{7/2} dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2)*(c - c*tan(e + f*x)*1i)^(7/2),x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2)*(c - c*tan(e + f*x)*1i)^(7/2), x)`

3.818 $\int (a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{5/2} dx$

3.818.1 Optimal result	7361
3.818.2 Mathematica [A] (verified)	7362
3.818.3 Rubi [A] (verified)	7362
3.818.4 Maple [B] (verified)	7365
3.818.5 Fricas [B] (verification not implemented)	7366
3.818.6 Sympy [F(-1)]	7367
3.818.7 Maxima [B] (verification not implemented)	7368
3.818.8 Giac [F]	7369
3.818.9 Mupad [F(-1)]	7369

3.818.1 Optimal result

Integrand size = 45, antiderivative size = 284

$$\int (a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{5/2} dx =$$

$$\frac{a^{7/2}(6iA+B)c^{5/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{8f}$$

$$+ \frac{a^3(6A-iB)c^2 \tan(e+fx) \sqrt{a+ia \tan(e+fx)} \sqrt{c-ictan(e+fx)}}{16f}$$

$$+ \frac{a^2(6A-iB)c \tan(e+fx)(a+ia \tan(e+fx))^{3/2}(c-ictan(e+fx))^{3/2}}{24f}$$

$$+ \frac{a(6iA+B)(a+ia \tan(e+fx))^{5/2}(c-ictan(e+fx))^{5/2}}{30f}$$

$$+ \frac{B(a+ia \tan(e+fx))^{7/2}(c-ictan(e+fx))^{5/2}}{6f}$$

output

```
-1/8*a^(7/2)*(6*I*A+B)*c^(5/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/f+1/16*a^3*(6*A-I*B)*c^2*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)*tan(f*x+e)/f+1/24*a^2*(6*A-I*B)*c*tan(f*x+e)*(a+I*a*tan(f*x+e))^(3/2)*(c-I*c*tan(f*x+e))^(3/2)/f+1/30*a*(6*I*A+B)*(a+I*a*tan(f*x+e))^(5/2)*(c-I*c*tan(f*x+e))^(5/2)/f+1/6*B*(a+I*a*tan(f*x+e))^(7/2)*(c-I*c*tan(f*x+e))^(5/2)/f
```


3.818.2 Mathematica [A] (verified)

Time = 11.71 (sec) , antiderivative size = 510, normalized size of antiderivative = 1.80

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{5/2} dx = \frac{B(a + ia \tan(e + fx))^{7/2} (c - ictan(e + fx))^{5/2}}{6f}$$

$$+ \frac{a(6iA+B)c(a+ia \tan(e+fx))^{7/2}(c-ictan(e+fx))^{3/2}}{5f} + \frac{-3a^2(6iA+B)c^2(a+ia \tan(e+fx))^{7/2}\sqrt{c-ictan(e+fx)}}{4f} + \frac{-a^3(6iA+B)c^3(a+ia \tan(e+fx))^{7/2}}{f\sqrt{c-ictan(e+fx)}}$$

input `Integrate[(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2),x]`

output `(B*(a + I*a*Tan[e + f*x])^(7/2)*(c - I*c*Tan[e + f*x])^(5/2))/(6*f) + (-1/5*(a*((6*I)*A + B)*c*(a + I*a*Tan[e + f*x])^(7/2)*(c - I*c*Tan[e + f*x])^(3/2))/f + ((-3*a^2*((6*I)*A + B)*c^2*(a + I*a*Tan[e + f*x])^(7/2)*Sqrt[c - I*c*Tan[e + f*x]])/(4*f) + (-((a^3*((6*I)*A + B)*c^3*(a + I*a*Tan[e + f*x])^(7/2))/(f*Sqrt[c - I*c*Tan[e + f*x]])) + ((-3*a^4*((6*I)*A + B)*c^4*(a + I*a*Tan[e + f*x])^(5/2))/(2*f*Sqrt[c - I*c*Tan[e + f*x]]) + ((-15*a^5*((6*I)*A + B)*c^5*(a + I*a*Tan[e + f*x])^(3/2))/(f*Sqrt[c - I*c*Tan[e + f*x]]) + ((90*I)*a^7*(6*A - I*B)*c^5*(1 - I*Tan[e + f*x])*((1 + I*Tan[e + f*x])/(1 - I*Tan[e + f*x]) - (ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a]])*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[1 - I*Tan[e + f*x]])))/(f*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]]))/(2*c)/(3*c)/(4*a)/(5*a))/(6*a)`

3.818.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 267, normalized size of antiderivative = 0.94, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.178$, Rules used = {3042, 4071, 90, 59, 40, 40, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.818. $\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{5/2} dx$

$$\int (a + ia \tan(e + fx))^{7/2} (c - ictan(e + fx))^{5/2} (A + B \tan(e + fx)) dx$$

↓ 3042

$$\int (a + ia \tan(e + fx))^{7/2} (c - ictan(e + fx))^{5/2} (A + B \tan(e + fx)) dx$$

↓ 4071

$$\frac{ac \int (i \tan(e + fx)a + a)^{5/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} d \tan(e + fx)}{f}$$

↓ 90

$$\frac{ac \left(\frac{1}{6} (6A - iB) \int (i \tan(e + fx)a + a)^{5/2} (c - ictan(e + fx))^{3/2} d \tan(e + fx) + \frac{B(a + ia \tan(e + fx))^{7/2} (c - ictan(e + fx))^{5/2}}{6ac} \right)}{f}$$

↓ 59

$$\frac{ac \left(\frac{1}{6} (6A - iB) \left(a \int (i \tan(e + fx)a + a)^{3/2} (c - ictan(e + fx))^{3/2} d \tan(e + fx) + \frac{i(a + ia \tan(e + fx))^{5/2} (c - ictan(e + fx))}{5c} \right) \right)}{f}$$

↓ 40

$$\frac{ac \left(\frac{1}{6} (6A - iB) \left(a \left(\frac{3}{4} ac \int \sqrt{i \tan(e + fx)a + a} \sqrt{c - ictan(e + fx)} d \tan(e + fx) + \frac{1}{4} \tan(e + fx) (a + ia \tan(e + fx)) \right) \right) \right)}{f}$$

↓ 40

$$\frac{ac \left(\frac{1}{6} (6A - iB) \left(a \left(\frac{3}{4} ac \left(\frac{1}{2} ac \int \frac{1}{\sqrt{i \tan(e + fx)a + a} \sqrt{c - ictan(e + fx)}} d \tan(e + fx) + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ictan(e + fx)} \right) \right) \right) \right)}{f}$$

↓ 45

$$\frac{ac \left(\frac{1}{6} (6A - iB) \left(a \left(\frac{3}{4} ac \left(ac \int \frac{1}{ia + \frac{ic(i \tan(e + fx)a + a)}{c - ictan(e + fx)}} d \frac{\sqrt{i \tan(e + fx)a + a}}{\sqrt{c - ictan(e + fx)}} + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ictan(e + fx)} \right) \right) \right) \right)}{f}$$

↓ 218

$$\frac{ac \left(\frac{1}{6} (6A - iB) \left(a \left(\frac{3}{4} ac \left(\frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ictan(e + fx)} - i\sqrt{a}\sqrt{c} \arctan \left(\frac{\sqrt{c}\sqrt{a + ia \tan(e + fx)}}{\sqrt{a}\sqrt{c - ictan(e + fx)}} \right) \right) \right) \right) \right)}{f}$$

3.818. $\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{5/2} dx$

input `Int[(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2),x]`

output `(a*c*((B*(a + I*a*Tan[e + f*x])^(7/2)*(c - I*c*Tan[e + f*x])^(5/2))/(6*a*c) + (((6*A - I*B)*(((I/5)*(a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e + f*x])^(5/2))/c + a*((Tan[e + f*x]*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(3/2))/4 + (3*a*c*((-I)*Sqrt[a]*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])] + (Tan[e + f*x]*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/2))/4))/6))/f`

3.818.3.1 Defintions of rubi rules used

rule 40 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^n/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`

rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 59 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[2*c*(n/(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0] && IGtQ[n + 1/2, 0] && LtQ[m, n]`

rule 90 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.818.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 477 vs. $2(232) = 464$.

Time = 0.49 (sec) , antiderivative size = 478, normalized size of antiderivative = 1.68

method	result
derivativedivides	$\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^3 c^2 \left(40iB \sqrt{ac(1+\tan(fx+e)^2)} \sqrt{ac} \tan(fx+e)^5 + 48iA \sqrt{ac(1+\tan(fx+e)^2)} \right)$
default	$\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^3 c^2 \left(40iB \sqrt{ac(1+\tan(fx+e)^2)} \sqrt{ac} \tan(fx+e)^5 + 48iA \sqrt{ac(1+\tan(fx+e)^2)} \right)$
parts	$A \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^3 c^2 \left(8i \tan(fx+e)^4 \sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} + 16i \tan(fx+e)^2 \sqrt{ac(1+\tan(fx+e)^2)} \right)$

input `int((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x,m
ethod=_RETURNVERBOSE)`

output

```

1/240/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^3*c^2*(40
*I*B*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)*tan(f*x+e)^5+48*I*A*(a*c*(1+
tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)*tan(f*x+e)^4+70*I*B*(a*c)^(1/2)*(a*c*(1+t
an(f*x+e)^2))^(1/2)*tan(f*x+e)^3+48*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(
1/2)*tan(f*x+e)^4+96*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+
e)^2+60*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^3-15*I*B*ln(
(a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c
+15*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+96*B*(a*c)^(1/
2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+48*I*A*(a*c)^(1/2)*(a*c*(1+ta
n(f*x+e)^2))^(1/2)+90*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^
2))^(1/2))/(a*c)^(1/2))*a*c+150*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)
*tan(f*x+e)+48*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2)/(a*
c*(1+tan(f*x+e)^2))^(1/2)

```

3.818.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 756 vs. $2(218) = 436$.

Time = 0.29 (sec) , antiderivative size = 756, normalized size of antiderivative = 2.66

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx =$$

$$15 \sqrt{\frac{(36A^2 - 12iAB - B^2)a^7c^5}{f^2}} (fe^{(10i fx + 10ie)} + 5fe^{(8i fx + 8ie)} + 10fe^{(6i fx + 6ie)} + 10fe^{(4i fx + 4ie)} + 5fe^{(2i fx + 2ie)})$$

input

```

integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/
2),x, algorithm="fricas")

```

3.818. $\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx$

output

```
-1/480*(15*sqrt((36*A^2 - 12*I*A*B - B^2)*a^7*c^5/f^2)*(f*e^(10*I*f*x + 10
*I*e) + 5*f*e^(8*I*f*x + 8*I*e) + 10*f*e^(6*I*f*x + 6*I*e) + 10*f*e^(4*I*f
*x + 4*I*e) + 5*f*e^(2*I*f*x + 2*I*e) + f)*log(-4*(2*((-6*I*A - B)*a^3*c^2
*e^(3*I*f*x + 3*I*e) + (-6*I*A - B)*a^3*c^2*e^(I*f*x + I*e))*sqrt(a/(e^(2*
I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + sqrt((36*A^2 - 12
*I*A*B - B^2)*a^7*c^5/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((6*I*A + B)*a^3*c
^2*e^(2*I*f*x + 2*I*e) + (6*I*A + B)*a^3*c^2)) - 15*sqrt((36*A^2 - 12*I*A*
B - B^2)*a^7*c^5/f^2)*(f*e^(10*I*f*x + 10*I*e) + 5*f*e^(8*I*f*x + 8*I*e) +
10*f*e^(6*I*f*x + 6*I*e) + 10*f*e^(4*I*f*x + 4*I*e) + 5*f*e^(2*I*f*x + 2*
I*e) + f)*log(-4*(2*((-6*I*A - B)*a^3*c^2*e^(3*I*f*x + 3*I*e) + (-6*I*A -
B)*a^3*c^2*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2
*I*f*x + 2*I*e) + 1)) - sqrt((36*A^2 - 12*I*A*B - B^2)*a^7*c^5/f^2)*(f*e^(
2*I*f*x + 2*I*e) - f))/((6*I*A + B)*a^3*c^2*e^(2*I*f*x + 2*I*e) + (6*I*A +
B)*a^3*c^2)) + 4*(15*(6*I*A + B)*a^3*c^2*e^(11*I*f*x + 11*I*e) + 85*(6*I*
A + B)*a^3*c^2*e^(9*I*f*x + 9*I*e) + 6*(-58*I*A - 223*B)*a^3*c^2*e^(7*I*f*
x + 7*I*e) + 198*(-6*I*A - B)*a^3*c^2*e^(5*I*f*x + 5*I*e) + 85*(-6*I*A - B
)*a^3*c^2*e^(3*I*f*x + 3*I*e) + 15*(-6*I*A - B)*a^3*c^2*e^(I*f*x + I*e))*s
qrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(f*e^(
10*I*f*x + 10*I*e) + 5*f*e^(8*I*f*x + 8*I*e) + 10*f*e^(6*I*f*x + 6*I*e) +
10*f*e^(4*I*f*x + 4*I*e) + 5*f*e^(2*I*f*x + 2*I*e) + f)
```

3.818.6 Sympy [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx = \text{Timed out}$$

input

```
integrate((a+I*a*tan(f*x+e))**(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(
5/2),x)
```

output

```
Timed out
```

3.818.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2033 vs. $2(218) = 436$.

Time = 5.96 (sec) , antiderivative size = 2033, normalized size of antiderivative = 7.16

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx = \text{Too large to display}$$

```
input integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="maxima")
```

```
output -3840*(60*(6*A - I*B)*a^3*c^2*cos(11/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 340*(6*A - I*B)*a^3*c^2*cos(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 24*(58*A - 223*I*B)*a^3*c^2*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 792*(6*A - I*B)*a^3*c^2*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 340*(6*A - I*B)*a^3*c^2*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 60*(6*A - I*B)*a^3*c^2*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 60*(-6*I*A - B)*a^3*c^2*sin(11/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 340*(-6*I*A - B)*a^3*c^2*sin(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 24*(58*I*A + 223*B)*a^3*c^2*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 792*(6*I*A + B)*a^3*c^2*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 340*(6*I*A + B)*a^3*c^2*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 60*(6*I*A + B)*a^3*c^2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 30*((6*A - I*B)*a^3*c^2*cos(12*f*x + 12*e) + 6*(6*A - I*B)*a^3*c^2*cos(10*f*x + 10*e) + 15*(6*A - I*B)*a^3*c^2*cos(8*f*x + 8*e) + 20*(6*A - I*B)*a^3*c^2*cos(6*f*x + 6*e) + 15*(6*A - I*B)*a^3*c^2*cos(4*f*x + 4*e) + 6*(6*A - I*B)*a^3*c^2*cos(2*f*x + 2*e) - (-6*I*A - B)*a^3*c^2*sin(12*f*x + 12*e) - 6*(-6*I*A - B)*a^3*c^2*sin(10*f*x + 10*e) - 15*(-6*I*A - B)*a^3*c^2*sin(8*f*x + 8*e) - 20*(-6*I*A - B)*a^3*c^2*sin(6*f*x + 6*e) - 15*(-6*I*A - B)*a^3*c^2*sin(4*f*x + 4*e) - 6*(-6*I*A - B)*a^3*c^2*sin(2*f*x + 2*e) + (6...
```

3.818.8 Giac [F]

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx = \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^{7/2} (-ic \tan(fx + e) + c)^{5/2} dx$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `sage0*x`

3.818.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{5/2} dx = \int (A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{7/2} (c - c \tan(e + fx) li)^{5/2} dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2)*(c - c*tan(e + f*x)*1i)^(5/2),x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2)*(c - c*tan(e + f*x)*1i)^(5/2), x)`

3.819 $\int (a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))(c-ictan(e+fx))^{3/2} dx$

3.819.1 Optimal result	7370
3.819.2 Mathematica [A] (verified)	7371
3.819.3 Rubi [A] (verified)	7371
3.819.4 Maple [A] (verified)	7374
3.819.5 Fricas [B] (verification not implemented)	7375
3.819.6 Sympy [F(-1)]	7376
3.819.7 Maxima [B] (verification not implemented)	7376
3.819.8 Giac [F]	7377
3.819.9 Mupad [F(-1)]	7378

3.819.1 Optimal result

Integrand size = 45, antiderivative size = 279

$$\int (a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))(c - ictan(e + fx))^{3/2} dx =$$

$$\frac{a^{7/2}(5iA + 2B)c^{3/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{4f}$$

$$+ \frac{a^3(5A - 2iB)c \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ictan(e + fx)}}{8f}$$

$$+ \frac{a^2(5iA + 2B)(a + ia \tan(e + fx))^{3/2}(c - ictan(e + fx))^{3/2}}{12f}$$

$$+ \frac{a(5iA + 2B)(a + ia \tan(e + fx))^{5/2}(c - ictan(e + fx))^{3/2}}{20f}$$

$$+ \frac{B(a + ia \tan(e + fx))^{7/2}(c - ictan(e + fx))^{3/2}}{5f}$$

output

```
-1/4*a^(7/2)*(5*I*A+2*B)*c^(3/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a
^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/f+1/8*a^3*(5*A-2*I*B)*c*(a+I*a*tan(f*x+e)
)^(1/2)*(c-I*c*tan(f*x+e))^(1/2)*tan(f*x+e)/f+1/12*a^2*(5*I*A+2*B)*(a+I*a*
tan(f*x+e))^(3/2)*(c-I*c*tan(f*x+e))^(3/2)/f+1/20*a*(5*I*A+2*B)*(a+I*a*
tan(f*x+e))^(5/2)*(c-I*c*tan(f*x+e))^(3/2)/f+1/5*B*(a+I*a*tan(f*x+e))^(7/2)*(
c-I*c*tan(f*x+e))^(3/2)/f
```

3.819.2 Mathematica [A] (verified)

Time = 11.53 (sec) , antiderivative size = 458, normalized size of antiderivative = 1.64

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx = \frac{B(a + ia \tan(e + fx))^{7/2} (c - ictan(e + fx))^{3/2}}{5f}$$

$$+ \frac{a(5iA+2B)c(a+ia \tan(e+fx))^{7/2} \sqrt{c-ictan(e+fx)}}{4f} + \frac{a^2(5iA+2B)c^2(a+ia \tan(e+fx))^{7/2}}{3f\sqrt{c-ictan(e+fx)}} + \frac{-a^3(5iA+2B)c^3(a+ia \tan(e+fx))^{5/2}}{2f\sqrt{c-ictan(e+fx)}} + \frac{-5a^4(5iA+2B)c^4(a+ia \tan(e+fx))^{3/2}}{f}$$

5a

input `Integrate[(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(B*(a + I*a*Tan[e + f*x])^(7/2)*(c - I*c*Tan[e + f*x])^(3/2))/(5*f) + (-1/4*(a*((5*I)*A + 2*B)*c*(a + I*a*Tan[e + f*x])^(7/2)*Sqrt[c - I*c*Tan[e + f*x]])/f + (-1/3*(a^2*((5*I)*A + 2*B)*c^2*(a + I*a*Tan[e + f*x])^(7/2))/(f*Sqrt[c - I*c*Tan[e + f*x]]) + (-1/2*(a^3*((5*I)*A + 2*B)*c^3*(a + I*a*Tan[e + f*x])^(5/2))/(f*Sqrt[c - I*c*Tan[e + f*x]]) + ((-5*a^4*((5*I)*A + 2*B)*c^4*(a + I*a*Tan[e + f*x])^(3/2))/(f*Sqrt[c - I*c*Tan[e + f*x]]) + ((30*I)*a^6*(5*A - (2*I)*B)*c^4*(1 - I*Tan[e + f*x])*((1 + I*Tan[e + f*x])/(1 - I*Tan[e + f*x]) - (ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a])]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[1 - I*Tan[e + f*x]])))/(f*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/(2*c))/(3*c))/(4*a))/(5*a)`

3.819.3 Rubi [A] (verified)Time = 0.44 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.95, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.178$, Rules used = {3042, 4071, 90, 59, 59, 40, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))^{7/2} (c - ictan(e + fx))^{3/2} (A + B \tan(e + fx)) dx$$

↓ 3042

3.819. $\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx$

$$\int (a + ia \tan(e + fx))^{7/2} (c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx)) dx$$

↓ 4071

$$\frac{ac \int (i \tan(e + fx)a + a)^{5/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} d \tan(e + fx)}{f}$$

↓ 90

$$\frac{ac \left(\frac{1}{5} (5A - 2iB) \int (i \tan(e + fx)a + a)^{5/2} \sqrt{c - ic \tan(e + fx)} d \tan(e + fx) + \frac{B(a + ia \tan(e + fx))^{7/2} (c - ic \tan(e + fx))^{3/2}}{5ac} \right)}{f}$$

↓ 59

$$\frac{ac \left(\frac{1}{5} (5A - 2iB) \left(\frac{5}{4} a \int (i \tan(e + fx)a + a)^{3/2} \sqrt{c - ic \tan(e + fx)} d \tan(e + fx) + \frac{i(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2}}{4c} \right) \right)}{f}$$

↓ 59

$$\frac{ac \left(\frac{1}{5} (5A - 2iB) \left(\frac{5}{4} a \left(a \int \sqrt{i \tan(e + fx)a + a} \sqrt{c - ic \tan(e + fx)} d \tan(e + fx) + \frac{i(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2}}{3c} \right) \right) \right)}{f}$$

↓ 40

$$\frac{ac \left(\frac{1}{5} (5A - 2iB) \left(\frac{5}{4} a \left(a \left(\frac{1}{2} ac \int \frac{1}{\sqrt{i \tan(e + fx)a + a} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)} \right) \right) \right) \right)}{f}$$

↓ 45

$$\frac{ac \left(\frac{1}{5} (5A - 2iB) \left(\frac{5}{4} a \left(a \left(ac \int \frac{1}{ia + \frac{ic(i \tan(e + fx)a + a)}{c - ic \tan(e + fx)}} d \frac{\sqrt{i \tan(e + fx)a + a}}{\sqrt{c - ic \tan(e + fx)}} + \frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)} \right) \right) \right) \right)}{f}$$

↓ 218

$$\frac{ac \left(\frac{1}{5} (5A - 2iB) \left(\frac{5}{4} a \left(a \left(\frac{1}{2} \tan(e + fx) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)} - i\sqrt{a}\sqrt{c} \arctan \left(\frac{\sqrt{c}\sqrt{a + ia \tan(e + fx)}}{\sqrt{a}\sqrt{c - ic \tan(e + fx)}} \right) \right) \right) \right) \right)}{f}$$

input `Int[(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2),x]`

```
output (a*c*((B*(a + I*a*Tan[e + f*x])^(7/2)*(c - I*c*Tan[e + f*x])^(3/2))/(5*a*c
) + ((5*A - (2*I)*B)*(((I/4)*(a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e +
f*x])^(3/2))/c + (5*a*(((I/3)*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e
+ f*x])^(3/2))/c + a*((-I)*Sqrt[a]*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*T
an[e + f*x])]/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])] + (Tan[e + f*x]*Sqrt[a
+ I*a*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]))/2))))/4))/5))/f
```

3.819.3.1 Defintions of rubi rules used

```
rule 40 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[x*
(a + b*x)^m*((c + d*x)^(n)/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a
+ b*x)^(m - 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[
b*c + a*d, 0] && IGtQ[m + 1/2, 0]
```

```
rule 45 Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; Fre
eQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]
```

```
rule 59 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a
+ b*x)^(m + 1)*((c + d*x)^(n)/(b*(m + n + 1))), x] + Simp[2*c*(n/(m + n + 1)
) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] &&
EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0] && IGtQ[n + 1/2, 0] && LtQ[m, n]
```

```
rule 90 Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
_), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p
+ 2)) Int[(c + d*x)^(n)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n,
p}, x] && NeQ[n + p + 2, 0]
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.819.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 412, normalized size of antiderivative = 1.48

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^3 c\left(60 i B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)} \tan (f x+e)^3-24 B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}\right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^3 c\left(60 i B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)} \tan (f x+e)^3-24 B \sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}\right)}{\dots}$
parts	$\frac{A \sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^3 c\left(16 i \tan (f x+e)^2 \sqrt{a c\left(1+\tan (f x+e)^2\right)} \sqrt{a c}-6 \tan (f x+e)^3 \sqrt{a c\left(1+\tan (f x+e)^2\right)}\right)}{\dots}$

```
input int((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x,m
method=_RETURNVERBOSE)
```

```
output 1/120/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^3*c*(60*I
*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^3-24*B*(a*c)^(1/2)*
(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^4+80*I*A*(a*c)^(1/2)*(a*c*(1+tan(f
*x+e)^2))^(1/2)*tan(f*x+e)^2-30*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)
*tan(f*x+e)^3-30*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))
^(1/2)))/(a*c)^(1/2))*a*c+30*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*t
an(f*x+e)+32*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+80*I*
A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)+75*A*ln((a*c*tan(f*x+e)+(a*c)^(
1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c+45*A*(a*c)^(1/2)*(a*c
*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+56*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)
^(1/2))/(a*c)^(1/2)/(a*c*(1+tan(f*x+e)^2))^(1/2)
```

3.819. $\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))(c - ictan(e + fx))^{3/2} dx$

3.819.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 682 vs. $2(213) = 426$.

Time = 0.29 (sec) , antiderivative size = 682, normalized size of antiderivative = 2.44

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} dx =$$

$$15 \sqrt{\frac{(25A^2 - 20iAB - 4B^2)a^7c^3}{f^2}} (fe^{(8ifx+8ie)} + 4fe^{(6ifx+6ie)} + 6fe^{(4ifx+4ie)} + 4fe^{(2ifx+2ie)} + f) \log \left(-\frac{4 \left(2 \left((-5iA - 2B) a^3 c e^{(3ifx+3ie)} + (-5iA - 2B) a^3 c e^{(ifx+ie)} \right) \sqrt{a/(e^{(2ifx+2ie)} + 1)} \sqrt{c/(e^{(2ifx+2ie)} + 1)} + \sqrt{(25A^2 - 20iAB - 4B^2)a^7c^3/f^2} \right) (fe^{(2ifx+2ie)} - f) / ((5iA + 2B)a^3c e^{(2ifx+2ie)} + (5iA + 2B)a^3c) - 15 \sqrt{(25A^2 - 20iAB - 4B^2)a^7c^3/f^2} (fe^{(8ifx+8ie)} + 4fe^{(6ifx+6ie)} + 6fe^{(4ifx+4ie)} + 4fe^{(2ifx+2ie)} + f) \log(-4 \left(2 \left((-5iA - 2B) a^3 c e^{(3ifx+3ie)} + (-5iA - 2B) a^3 c e^{(ifx+ie)} \right) \sqrt{a/(e^{(2ifx+2ie)} + 1)} \sqrt{c/(e^{(2ifx+2ie)} + 1)} - \sqrt{(25A^2 - 20iAB - 4B^2)a^7c^3/f^2} \right) (fe^{(2ifx+2ie)} - f) / ((5iA + 2B)a^3c e^{(2ifx+2ie)} + (5iA + 2B)a^3c) + 4 \left(15(5iA + 2B)a^3c e^{(9ifx+9ie)} + 10(-29iA - 50B)a^3c e^{(7ifx+7ie)} + 128(-5iA - 2B)a^3c e^{(5ifx+5ie)} + 70(-5iA - 2B)a^3c e^{(3ifx+3ie)} + 15(-5iA - 2B)a^3c e^{(ifx+ie)} \right) \sqrt{a/(e^{(2ifx+2ie)} + 1)} \sqrt{c/(e^{(2ifx+2ie)} + 1)} \right) / (fe^{(8ifx+8ie)} + 4fe^{(6ifx+6ie)} + 6fe^{(4ifx+4ie)} + 4fe^{(2ifx+2ie)} + f) \right)$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output `-1/240*(15*sqrt((25*A^2 - 20*I*A*B - 4*B^2)*a^7*c^3/f^2)*(f*e^(8*I*f*x + 8*I*e) + 4*f*e^(6*I*f*x + 6*I*e) + 6*f*e^(4*I*f*x + 4*I*e) + 4*f*e^(2*I*f*x + 2*I*e) + f)*log(-4*(2*((-5*I*A - 2*B)*a^3*c*e^(3*I*f*x + 3*I*e) + (-5*I*A - 2*B)*a^3*c*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + sqrt((25*A^2 - 20*I*A*B - 4*B^2)*a^7*c^3/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((5*I*A + 2*B)*a^3*c*e^(2*I*f*x + 2*I*e) + (5*I*A + 2*B)*a^3*c)) - 15*sqrt((25*A^2 - 20*I*A*B - 4*B^2)*a^7*c^3/f^2)*(f*e^(8*I*f*x + 8*I*e) + 4*f*e^(6*I*f*x + 6*I*e) + 6*f*e^(4*I*f*x + 4*I*e) + 4*f*e^(2*I*f*x + 2*I*e) + f)*log(-4*(2*((-5*I*A - 2*B)*a^3*c*e^(3*I*f*x + 3*I*e) + (-5*I*A - 2*B)*a^3*c*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - sqrt((25*A^2 - 20*I*A*B - 4*B^2)*a^7*c^3/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((5*I*A + 2*B)*a^3*c*e^(2*I*f*x + 2*I*e) + (5*I*A + 2*B)*a^3*c) + 4*(15*(5*I*A + 2*B)*a^3*c*e^(9*I*f*x + 9*I*e) + 10*(-29*I*A - 50*B)*a^3*c*e^(7*I*f*x + 7*I*e) + 128*(-5*I*A - 2*B)*a^3*c*e^(5*I*f*x + 5*I*e) + 70*(-5*I*A - 2*B)*a^3*c*e^(3*I*f*x + 3*I*e) + 15*(-5*I*A - 2*B)*a^3*c*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(f*e^(8*I*f*x + 8*I*e) + 4*f*e^(6*I*f*x + 6*I*e) + 6*f*e^(4*I*f*x + 4*I*e) + 4*f*e^(2*I*f*x + 2*I*e) + f)`

3.819.6 Sympy [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(3/2),x)`

output `Timed out`

3.819.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1657 vs. $2(213) = 426$.

Time = 2.78 (sec) , antiderivative size = 1657, normalized size of antiderivative = 5.94

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{3/2} dx = \text{Too large to display}$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output

```
-480*(60*(5*A - 2*I*B)*a^3*c*cos(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 40*(29*A - 50*I*B)*a^3*c*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 512*(5*A - 2*I*B)*a^3*c*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 280*(5*A - 2*I*B)*a^3*c*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 60*(5*A - 2*I*B)*a^3*c*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 60*(-5*I*A - 2*B)*a^3*c*sin(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 40*(29*I*A + 50*B)*a^3*c*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 512*(5*I*A + 2*B)*a^3*c*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 280*(5*I*A + 2*B)*a^3*c*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 60*(5*I*A + 2*B)*a^3*c*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 30*((5*A - 2*I*B)*a^3*c*cos(10*f*x + 10*e) + 5*(5*A - 2*I*B)*a^3*c*cos(8*f*x + 8*e) + 10*(5*A - 2*I*B)*a^3*c*cos(6*f*x + 6*e) + 10*(5*A - 2*I*B)*a^3*c*cos(4*f*x + 4*e) + 5*(5*A - 2*I*B)*a^3*c*cos(2*f*x + 2*e) - (-5*I*A - 2*B)*a^3*c*sin(10*f*x + 10*e) - 5*(-5*I*A - 2*B)*a^3*c*sin(8*f*x + 8*e) - 10*(-5*I*A - 2*B)*a^3*c*sin(6*f*x + 6*e) - 10*(-5*I*A - 2*B)*a^3*c*sin(4*f*x + 4*e) - 5*(-5*I*A - 2*B)*a^3*c*sin(2*f*x + 2*e) + (5*A - 2*I*B)*a^3*c*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) + 30*((5*A - 2*I*B)*a^3*c*cos(10*f*x + 10*e) + 5*(5*A - 2*I*B)*a^3*c*cos(8*f*x + 8*e) + 10*(5*A - 2*I*B)*a^3*c*cos(6*f...
```

3.819.8 Giac [F]

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx = \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^{7/2} (-ictan(fx + e) + c)^{3/2} dx$$

input

```
integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2),x, algorithm="giac")
```

output

```
sage0*x
```


3.819.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) (c - ictan(e + fx))^{3/2} dx = \int (A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{7/2} (c - c \tan(e + fx) li)^{3/2} dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2)*(c - c*tan(e + f*x)*1i)^(3/2),x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2)*(c - c*tan(e + f*x)*1i)^(3/2), x)`

3.820 $\int (a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))\sqrt{c-ictan(e+fx)}$

3.820.1 Optimal result	7379
3.820.2 Mathematica [A] (verified)	7380
3.820.3 Rubi [A] (verified)	7380
3.820.4 Maple [A] (verified)	7383
3.820.5 Fricas [B] (verification not implemented)	7384
3.820.6 Sympy [F(-1)]	7385
3.820.7 Maxima [B] (verification not implemented)	7385
3.820.8 Giac [F(-1)]	7386
3.820.9 Mupad [F(-1)]	7387

3.820.1 Optimal result

Integrand size = 45, antiderivative size = 272

$$\int (a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)} dx =$$

$$\frac{5a^{7/2}(4iA + 3B)\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{4f}$$

$$+ \frac{5a^3(4iA + 3B)\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}}{8f}$$

$$+ \frac{5a^2(4iA + 3B)(a + ia \tan(e + fx))^{3/2}\sqrt{c - ictan(e + fx)}}{24f}$$

$$+ \frac{a(4iA + 3B)(a + ia \tan(e + fx))^{5/2}\sqrt{c - ictan(e + fx)}}{12f}$$

$$+ \frac{B(a + ia \tan(e + fx))^{7/2}\sqrt{c - ictan(e + fx)}}{4f}$$

output

```
-5/4*a^(7/2)*(4*I*A+3*B)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))*c^(1/2)/f+5/8*a^3*(4*I*A+3*B)*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)/f+5/24*a^2*(4*I*A+3*B)*(c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(3/2)/f+1/12*a*(4*I*A+3*B)*(c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(5/2)/f+1/4*B*(c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(7/2)/f
```

3.820.2 Mathematica [A] (verified)

Time = 7.96 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.81

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx = \frac{a^{7/2} c (i + \tan(e + fx)) \left(-30(4A - 3iB) \arcsin \left(\frac{\sqrt{a + ia \tan(e + fx)}}{\sqrt{2}\sqrt{a}} \right) \right)}{\dots}$$

```
input Integrate[(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]],x]
```

```
output (a^(7/2)*c*(I + Tan[e + f*x])*(-30*(4*A - (3*I)*B)*ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a]])*Sqrt[a + I*a*Tan[e + f*x]] + Sqrt[a]*Sqrt[1 - I*Tan[e + f*x]]*(-I + Tan[e + f*x])*((88*I)*A + 72*B - 9*(4*A - (5*I)*B)*Tan[e + f*x] - (8*I)*(A - (3*I)*B)*Tan[e + f*x]^2 - (6*I)*B*Tan[e + f*x]^3))/((24*f*Sqrt[1 - I*Tan[e + f*x]]*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])
```

3.820.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.97, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.178$, Rules used = {3042, 4071, 90, 60, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + ia \tan(e + fx))^{7/2} \sqrt{c - ic \tan(e + fx)} (A + B \tan(e + fx)) dx$$

↓ 3042

$$\int (a + ia \tan(e + fx))^{7/2} \sqrt{c - ic \tan(e + fx)} (A + B \tan(e + fx)) dx$$

↓ 4071

$$\frac{ac \int \frac{(i \tan(e + fx)a + a)^{5/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{f}$$

↓ 90

3.820. $\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx$

$$\begin{aligned}
& \frac{ac \left(\frac{1}{4}(4A - 3iB) \int \frac{(i \tan(e+fx)a+a)^{5/2}}{\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{B(a+ia \tan(e+fx))^{7/2} \sqrt{c-ic \tan(e+fx)}}{4ac} \right)}{f} \\
& \quad \downarrow 60 \\
& \frac{ac \left(\frac{1}{4}(4A - 3iB) \left(\frac{5}{3}a \int \frac{(i \tan(e+fx)a+a)^{3/2}}{\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}}{3c} \right) + \frac{B(a+ia \tan(e+fx))^{7/2} \sqrt{c-ic \tan(e+fx)}}{4ac} \right)}{f} \\
& \quad \downarrow 60 \\
& \frac{ac \left(\frac{1}{4}(4A - 3iB) \left(\frac{5}{3}a \left(\frac{3}{2}a \int \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2c} \right) + \frac{i(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}}{3c} \right) \right)}{f} \\
& \quad \downarrow 60 \\
& \frac{ac \left(\frac{1}{4}(4A - 3iB) \left(\frac{5}{3}a \left(\frac{3}{2}a \left(a \int \frac{1}{\sqrt{i \tan(e+fx)a+a} \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{c} \right) \right) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{3c} \right) \right)}{f} \\
& \quad \downarrow 45 \\
& \frac{ac \left(\frac{1}{4}(4A - 3iB) \left(\frac{5}{3}a \left(\frac{3}{2}a \left(2a \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} + \frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{c} \right) \right) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{3c} \right) \right)}{f} \\
& \quad \downarrow 218 \\
& \frac{ac \left(\frac{1}{4}(4A - 3iB) \left(\frac{5}{3}a \left(\frac{3}{2}a \left(\frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{c} - \frac{2i\sqrt{a} \arctan\left(\frac{\sqrt{c} \sqrt{a+ia \tan(e+fx)}}{\sqrt{a} \sqrt{c-ic \tan(e+fx)}}\right)}{\sqrt{c}} \right) \right) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2c} \right) \right)}{f}
\end{aligned}$$

input `Int[(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]],x]`

output `(a*c*((B*(a + I*a*Tan[e + f*x])^(7/2)*Sqrt[c - I*c*Tan[e + f*x]])/(4*a*c) + ((4*A - (3*I)*B)*(((I/3)*(a + I*a*Tan[e + f*x])^(5/2)*Sqrt[c - I*c*Tan[e + f*x]])/c + (5*a*(((I/2)*(a + I*a*Tan[e + f*x])^(3/2)*Sqrt[c - I*c*Tan[e + f*x]])/c + (3*a*(((-2*I)*Sqrt[a]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[c] + (I*Sqrt[a + I*a*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]/c))/2))/3))/4)/f`

3.820. $\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx$

3.820.3.1 Defintions of rubi rules used

- rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 60 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 90 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)) Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.820.4 Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 349, normalized size of antiderivative = 1.28

method	result
derivativedivides	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} a^3 \left(-6iB\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \tan(fx+e)^3 - 8iA\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \right)}{}$
default	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} a^3 \left(-6iB\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \tan(fx+e)^3 - 8iA\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \right)}{}$
parts	$\frac{A\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} a^3 \left(2i \tan(fx+e)^2 \sqrt{ac(1+\tan(fx+e)^2)} \sqrt{ac} - 22i\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \right)}{6f\sqrt{ac(1+\tan(fx+e)^2)}}$

```
input int((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e)),x,method=_RETURNVERBOSE)
```

```
output 1/24/f*(-c*(I*tan(f*x+e)-1))^(1/2)*(a*(1+I*tan(f*x+e)))^(1/2)*a^3*(-6*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^3-8*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2-45*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c+45*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-24*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+60*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c+88*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)-36*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+72*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2)/(a*c*(1+tan(f*x+e)^2))^(1/2)
```

3.820.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 604 vs. $2(206) = 412$.

Time = 0.27 (sec) , antiderivative size = 604, normalized size of antiderivative = 2.22

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx =$$

$$15 \sqrt{\frac{(16A^2 - 24iAB - 9B^2)a^7c}{f^2}} (fe^{(6ifx+6ie)} + 3fe^{(4ifx+4ie)} + 3fe^{(2ifx+2ie)} + f) \log \left(\frac{4 \left(2((-4iA-3B)a^3e^{(3ifx+3ie)} \right)}{\dots} \right)$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e)),x, algorithm="fricas")`

output `-1/48*(15*sqrt((16*A^2 - 24*I*A*B - 9*B^2)*a^7*c/f^2)*(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)*log(4*(2*((-4*I*A - 3*B)*a^3*e^(3*I*f*x + 3*I*e) + (-4*I*A - 3*B)*a^3*e^(I*f*x + I*e)))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + sqrt((16*A^2 - 24*I*A*B - 9*B^2)*a^7*c/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((-4*I*A - 3*B)*a^3*e^(2*I*f*x + 2*I*e) + (-4*I*A - 3*B)*a^3) - 15*sqrt((16*A^2 - 24*I*A*B - 9*B^2)*a^7*c/f^2)*(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)*log(4*(2*((-4*I*A - 3*B)*a^3*e^(3*I*f*x + 3*I*e) + (-4*I*A - 3*B)*a^3*e^(I*f*x + I*e)))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - sqrt((16*A^2 - 24*I*A*B - 9*B^2)*a^7*c/f^2)*(f*e^(2*I*f*x + 2*I*e) - f))/((-4*I*A - 3*B)*a^3*e^(2*I*f*x + 2*I*e) + (-4*I*A - 3*B)*a^3) + 4*(3*(-44*I*A - 49*B)*a^3*e^(7*I*f*x + 7*I*e) + 73*(-4*I*A - 3*B)*a^3*e^(5*I*f*x + 5*I*e) + 55*(-4*I*A - 3*B)*a^3*e^(3*I*f*x + 3*I*e) + 15*(-4*I*A - 3*B)*a^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(f*e^(6*I*f*x + 6*I*e) + 3*f*e^(4*I*f*x + 4*I*e) + 3*f*e^(2*I*f*x + 2*I*e) + f)`

3.820.6 Sympy [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx = \text{Timed out}$$

input `integrate((c-I*c*tan(f*x+e))**(1/2)*(a+I*a*tan(f*x+e))**(7/2)*(A+B*tan(f*x+e)),x)`

output `Timed out`

3.820.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1345 vs. $2(206) = 412$.

Time = 1.06 (sec) , antiderivative size = 1345, normalized size of antiderivative = 4.94

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)} dx = \text{Too large to display}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e)),x, algorithm="maxima")`

output

```

96*(12*(44*A - 49*I*B)*a^3*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2
*e))) + 292*(4*A - 3*I*B)*a^3*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x
+ 2*e))) + 220*(4*A - 3*I*B)*a^3*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f
*x + 2*e))) + 60*(4*A - 3*I*B)*a^3*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2
*f*x + 2*e))) + 12*(44*I*A + 49*B)*a^3*sin(7/2*arctan2(sin(2*f*x + 2*e), c
os(2*f*x + 2*e))) + 292*(4*I*A + 3*B)*a^3*sin(5/2*arctan2(sin(2*f*x + 2*e)
, cos(2*f*x + 2*e))) + 220*(4*I*A + 3*B)*a^3*sin(3/2*arctan2(sin(2*f*x + 2
*e), cos(2*f*x + 2*e))) + 60*(4*I*A + 3*B)*a^3*sin(1/2*arctan2(sin(2*f*x +
2*e), cos(2*f*x + 2*e))) - 30*((4*A - 3*I*B)*a^3*cos(8*f*x + 8*e) + 4*(4*
A - 3*I*B)*a^3*cos(6*f*x + 6*e) + 6*(4*A - 3*I*B)*a^3*cos(4*f*x + 4*e) + 4
*(4*A - 3*I*B)*a^3*cos(2*f*x + 2*e) - (-4*I*A - 3*B)*a^3*sin(8*f*x + 8*e)
- 4*(-4*I*A - 3*B)*a^3*sin(6*f*x + 6*e) - 6*(-4*I*A - 3*B)*a^3*sin(4*f*x +
4*e) - 4*(-4*I*A - 3*B)*a^3*sin(2*f*x + 2*e) + (4*A - 3*I*B)*a^3)*arctan2
(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(sin
(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) - 30*((4*A - 3*I*B)*a^3*cos(8*f*x +
8*e) + 4*(4*A - 3*I*B)*a^3*cos(6*f*x + 6*e) + 6*(4*A - 3*I*B)*a^3*cos(4*f
*x + 4*e) + 4*(4*A - 3*I*B)*a^3*cos(2*f*x + 2*e) - (-4*I*A - 3*B)*a^3*sin(
8*f*x + 8*e) - 4*(-4*I*A - 3*B)*a^3*sin(6*f*x + 6*e) - 6*(-4*I*A - 3*B)*a^
3*sin(4*f*x + 4*e) - 4*(-4*I*A - 3*B)*a^3*sin(2*f*x + 2*e) + (4*A - 3*I*B)
*a^3)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), -si...

```

3.820.8 Giac [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx = \text{Timed out}$$

input

```

integrate((c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e
)),x, algorithm="giac")

```

output

```

Timed out

```

3.820.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx)) \sqrt{c - ictan(e + fx)} dx = \int (A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{7/2} \sqrt{c - c \tan(e + fx) li} dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2)*(c - c*tan(e + f*x)*1i)^(1/2),x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2)*(c - c*tan(e + f*x)*1i)^(1/2), x)`

3.821
$$\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{\sqrt{c-ictan(e+fx)}} dx$$

3.821.1 Optimal result 7388
 3.821.2 Mathematica [A] (verified) 7389
 3.821.3 Rubi [A] (verified) 7389
 3.821.4 Maple [B] (verified) 7392
 3.821.5 Fricas [B] (verification not implemented) 7393
 3.821.6 Sympy [F(-1)] 7394
 3.821.7 Maxima [B] (verification not implemented) 7394
 3.821.8 Giac [F] 7395
 3.821.9 Mupad [F(-1)] 7396

3.821.1 Optimal result

Integrand size = 45, antiderivative size = 283

$$\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx = \frac{5a^{7/2}(3iA + 4B) \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{\sqrt{cf}}$$

$$- \frac{(iA + B)(a + ia \tan(e + fx))^{7/2}}{f\sqrt{c - ictan(e + fx)}}$$

$$- \frac{5a^3(3iA + 4B)\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}}{2cf}$$

$$- \frac{5a^2(3iA + 4B)(a + ia \tan(e + fx))^{3/2}\sqrt{c - ictan(e + fx)}}{6cf}$$

$$- \frac{a(3iA + 4B)(a + ia \tan(e + fx))^{5/2}\sqrt{c - ictan(e + fx)}}{3cf}$$

output

```
5*a^(7/2)*(3*I*A+4*B)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/f/c^(1/2)-5/2*a^3*(3*I*A+4*B)*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)/c/f-5/6*a^2*(3*I*A+4*B)*(c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(3/2)/c/f-1/3*a*(3*I*A+4*B)*(c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(5/2)/c/f-(I*A+B)*(a+I*a*tan(f*x+e))^(7/2)/f/(c-I*c*tan(f*x+e))^(1/2)
```

3.821.2 Mathematica [A] (verified)

Time = 8.51 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.65

$$\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx = \frac{a^{7/2} \left(\frac{30(3A - 4iB) \arcsin\left(\frac{\sqrt{a + ia \tan(e + fx)}}{\sqrt{2}\sqrt{a}}\right)(i + \tan(e + fx))}{\sqrt{1 - i \tan(e + fx)}} + \frac{\sqrt{a}(-i + \tan(e + fx))}{\sqrt{1 - i \tan(e + fx)}} \right)}{6f}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/Sqrt[c - I*c*Tan[e + f*x]],x]`

output `(a^(7/2)*((30*(3*A - (4*I)*B)*ArcSin[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a])]*(I + Tan[e + f*x]))/Sqrt[1 - I*Tan[e + f*x]] + (Sqrt[a]*(-I + Tan[e + f*x])*(72*A - (94*I)*B + ((-21*I)*A - 34*B)*Tan[e + f*x] + (3*A - (10*I)*B)*Tan[e + f*x]^2 + 2*B*Tan[e + f*x]^3))/Sqrt[a + I*a*Tan[e + f*x]]))/(6*f*Sqrt[c - I*c*Tan[e + f*x]])`

3.821.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 270, normalized size of antiderivative = 0.95, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.178$, Rules used = {3042, 4071, 87, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(i \tan(e + fx)a + a)^{5/2}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{3/2}} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{87} \end{aligned}$$

3.821. $\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx$

$$ac \left(-\frac{(3A-4iB) \int \frac{(i \tan(e+fx)a+a)^{5/2}}{\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{ac \sqrt{c-ic \tan(e+fx)}} \right)$$

f
↓ 60

$$ac \left(-\frac{(3A-4iB) \left(\frac{5}{3} a \int \frac{(i \tan(e+fx)a+a)^{3/2}}{\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}}{3c} \right)}{c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{ac \sqrt{c-ic \tan(e+fx)}} \right)$$

f
↓ 60

$$ac \left(-\frac{(3A-4iB) \left(\frac{5}{3} a \left(\frac{3}{2} a \int \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2c} \right) + \frac{i(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}}{3c} \right)}{c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{ac \sqrt{c-ic \tan(e+fx)}} \right)$$

f
↓ 60

$$ac \left(-\frac{(3A-4iB) \left(\frac{5}{3} a \left(\frac{3}{2} a \left(a \int \frac{1}{\sqrt{i \tan(e+fx)a+a} \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{c} \right) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2c} \right) \right)}{c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{ac \sqrt{c-ic \tan(e+fx)}} \right)$$

f
↓ 45

$$ac \left(-\frac{(3A-4iB) \left(\frac{5}{3} a \left(\frac{3}{2} a \left(2a \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} + \frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{c} \right) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2c} \right) \right)}{c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{ac \sqrt{c-ic \tan(e+fx)}} \right)$$

f
↓ 218

$$ac \left(-\frac{(3A-4iB) \left(\frac{5}{3} a \left(\frac{3}{2} a \left(\frac{i \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{c} - \frac{2i \sqrt{a} \arctan \left(\frac{\sqrt{c} \sqrt{a+ia \tan(e+fx)}}{\sqrt{a} \sqrt{c-ic \tan(e+fx)}} \right)}{\sqrt{c}} \right) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2c} \right) \right)}{c} + \frac{i(a+ia \tan(e+fx))^{7/2}}{ac \sqrt{c-ic \tan(e+fx)}} \right)$$

f

3.821. $\int \frac{(a+ia \tan(e+fx))^{7/2} (A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$

input `Int[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/Sqrt[c - I*c*Tan[e + f*x]],x]`

output `(a*c*(-(((I*A + B)*(a + I*a*Tan[e + f*x])^(7/2))/(a*c*Sqrt[c - I*c*Tan[e + f*x]])) - ((3*A - (4*I)*B)*((I/3)*(a + I*a*Tan[e + f*x])^(5/2)*Sqrt[c - I*c*Tan[e + f*x]])/c + (5*a*((I/2)*(a + I*a*Tan[e + f*x])^(3/2)*Sqrt[c - I*c*Tan[e + f*x]])/c + (3*a*((-2*I)*Sqrt[a]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[c] + (I*Sqrt[a + I*a*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]/c))/2))/3)/c)/f`

3.821.3.1 Defintions of rubi rules used

rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 60 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.821.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 626 vs. 2(234) = 468.

Time = 0.40 (sec) , antiderivative size = 627, normalized size of antiderivative = 2.22

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^3 \left(-60 i B \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}}\right) a c \tan (f x+e)^2+8\right)}{}$
default	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^3 \left(-60 i B \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}}\right) a c \tan (f x+e)^2+8\right)}{}$
parts	$\frac{A \sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^3 \left(30 i \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}}\right) \tan (f x+e) a c+6 i \tan (f x+e)\right)}{}$

```
input int((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x,m
ethod=_RETURNVERBOSE)
```

3.821.
$$\int \frac{(a+ia \tan (e+f x))^{7 / 2}(A+B \tan (e+f x))}{\sqrt{c-i c \tan (e+f x)}} d x$$

output

```

-1/6/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^3/c*(-60*I
*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2
))*a*c*tan(f*x+e)^2+8*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x
+e)^3-2*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^4+90*I*A*ln(
(a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c
*tan(f*x+e)+18*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+4
5*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/
2))*a*c*tan(f*x+e)^2-3*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+
e)^3+60*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(
a*c)^(1/2))*a*c+128*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e
)+120*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)
^(1/2))*a*c*tan(f*x+e)+24*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f
*x+e)^2-72*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)-45*A*ln((a*c*tan(f
*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c-93*A*(a*c
)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-94*B*(a*c)^(1/2)*(a*c*(1+t
an(f*x+e)^2))^(1/2))/(a*c*(1+tan(f*x+e)^2))^(1/2)/(I+tan(f*x+e))^2/(a*c)^(
1/2)

```

3.821.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 590 vs. $2(219) = 438$.

Time = 0.27 (sec) , antiderivative size = 590, normalized size of antiderivative = 2.08

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx = \frac{15 \sqrt{\frac{(9A^2 - 24iAB - 16B^2)a^7}{cf^2}} (cfe^{(4i)fx + 4ie} + 2cfe^{(2i)fx + 2ie})}{15 \sqrt{\frac{(9A^2 - 24iAB - 16B^2)a^7}{cf^2}}}$$

input

```

integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/
2),x, algorithm="fracas")

```


output $1/12*(15*\sqrt{(9*A^2 - 24*I*A*B - 16*B^2)*a^7/(c*f^2)})*(c*f*e^{(4*I*f*x + 4*I*e)} + 2*c*f*e^{(2*I*f*x + 2*I*e)} + c*f)*\log(4*(2*((-3*I*A - 4*B)*a^3*e^{(3*I*f*x + 3*I*e)} + (-3*I*A - 4*B)*a^3*e^{(I*f*x + I*e)}))*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} + \sqrt{(9*A^2 - 24*I*A*B - 16*B^2)*a^7/(c*f^2)}*(c*f*e^{(2*I*f*x + 2*I*e)} - c*f)/((-3*I*A - 4*B)*a^3*e^{(2*I*f*x + 2*I*e)} + (-3*I*A - 4*B)*a^3)) - 15*\sqrt{(9*A^2 - 24*I*A*B - 16*B^2)*a^7/(c*f^2)}*(c*f*e^{(4*I*f*x + 4*I*e)} + 2*c*f*e^{(2*I*f*x + 2*I*e)} + c*f)*\log(4*(2*((-3*I*A - 4*B)*a^3*e^{(3*I*f*x + 3*I*e)} + (-3*I*A - 4*B)*a^3*e^{(I*f*x + I*e)}))*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} - \sqrt{(9*A^2 - 24*I*A*B - 16*B^2)*a^7/(c*f^2)}*(c*f*e^{(2*I*f*x + 2*I*e)} - c*f)/((-3*I*A - 4*B)*a^3*e^{(2*I*f*x + 2*I*e)} + (-3*I*A - 4*B)*a^3)) - 4*(24*(I*A + B)*a^3*e^{(7*I*f*x + 7*I*e)} + 33*(3*I*A + 4*B)*a^3*e^{(5*I*f*x + 5*I*e)} + 40*(3*I*A + 4*B)*a^3*e^{(3*I*f*x + 3*I*e)} + 15*(3*I*A + 4*B)*a^3*e^{(I*f*x + I*e)}))*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)})/(c*f*e^{(4*I*f*x + 4*I*e)} + 2*c*f*e^{(2*I*f*x + 2*I*e)} + c*f)$

3.821.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{\sqrt{c - ict \tan(e + fx)}} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(1/2),x)`

output Timed out

3.821.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1329 vs. $2(219) = 438$.

Time = 0.71 (sec) , antiderivative size = 1329, normalized size of antiderivative = 4.70

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{\sqrt{c - ict \tan(e + fx)}} dx = \text{Too large to display}$$

3.821. $\int \frac{(a+ia \tan(e+fx))^{7/2} (A+B \tan(e+fx))}{\sqrt{c-ict \tan(e+fx)}} dx$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x, algorithm="maxima")`

output `-6*(12*(9*A - 20*I*B)*a^3*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 32*(6*A - 11*I*B)*a^3*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 12*(9*I*A + 20*B)*a^3*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 32*(6*I*A + 11*B)*a^3*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 30*((3*A - 4*I*B)*a^3*cos(6*f*x + 6*e) + 3*(3*A - 4*I*B)*a^3*cos(4*f*x + 4*e) + 3*(3*A - 4*I*B)*a^3*cos(2*f*x + 2*e) - (-3*I*A - 4*B)*a^3*sin(6*f*x + 6*e) - 3*(-3*I*A - 4*B)*a^3*sin(4*f*x + 4*e) - 3*(-3*I*A - 4*B)*a^3*sin(2*f*x + 2*e) + (3*A - 4*I*B)*a^3)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) - 30*((3*A - 4*I*B)*a^3*cos(6*f*x + 6*e) + 3*(3*A - 4*I*B)*a^3*cos(4*f*x + 4*e) + 3*(3*A - 4*I*B)*a^3*cos(2*f*x + 2*e) - (-3*I*A - 4*B)*a^3*sin(6*f*x + 6*e) - 3*(-3*I*A - 4*B)*a^3*sin(4*f*x + 4*e) - 3*(-3*I*A - 4*B)*a^3*sin(2*f*x + 2*e) + (3*A - 4*I*B)*a^3)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), -sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) + 12*(8*(A - I*B)*a^3*cos(6*f*x + 6*e) + 24*(A - I*B)*a^3*cos(4*f*x + 4*e) + 24*(A - I*B)*a^3*cos(2*f*x + 2*e) + 8*(I*A + B)*a^3*sin(6*f*x + 6*e) + 24*(I*A + B)*a^3*sin(4*f*x + 4*e) + 24*(I*A + B)*a^3*sin(2*f*x + 2*e) + 5*(3*A - 4*I*B)*a^3*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 15*((-3*I*A - 4*B)*a^3*cos(6*f*x + 6*e) + 3*(-3*I*A - 4*B)*a^3*cos(4*f*x + 4*e) + 3*(-3*I*A - 4*B)*a^3*cos(2*f*x + ...`

3.821.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{\sqrt{c - ic \tan(e + fx)}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{7/2}}{\sqrt{-ic \tan(fx + e) + c}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(7/2)/sqrt(-I*c*tan(f*x + e) + c), x)`

3.821. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{\sqrt{c-ic \tan(e+fx)}} dx$

3.821.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{\sqrt{c - ictan(e + fx)}} dx = \int \frac{(A + B \tan(e + fx)) (a + a \tan(e + fx) li)^{7/2}}{\sqrt{c - c \tan(e + fx) li}} dx$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2))/(c - c*tan(e + f*x)*1i)^(1/2),x)`

output `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2))/(c - c*tan(e + f*x)*1i)^(1/2), x)`

3.822
$$\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$$

3.822.1 Optimal result	7397
3.822.2 Mathematica [A] (verified)	7398
3.822.3 Rubi [A] (verified)	7398
3.822.4 Maple [B] (verified)	7402
3.822.5 Fricas [B] (verification not implemented)	7403
3.822.6 Sympy [F(-1)]	7404
3.822.7 Maxima [B] (verification not implemented)	7404
3.822.8 Giac [F]	7405
3.822.9 Mupad [F(-1)]	7406

3.822.1 Optimal result

Integrand size = 45, antiderivative size = 285

$$\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx =$$

$$\frac{5a^{7/2}(2iA + 5B) \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{c^{3/2}f}$$

$$- \frac{(iA + B)(a + ia \tan(e + fx))^{7/2}}{3f(c - ic \tan(e + fx))^{3/2}} + \frac{2a(2iA + 5B)(a + ia \tan(e + fx))^{5/2}}{3cf\sqrt{c - ic \tan(e + fx)}}$$

$$+ \frac{5a^3(2iA + 5B)\sqrt{a + ia \tan(e + fx)}\sqrt{c - ic \tan(e + fx)}}{2c^2f}$$

$$+ \frac{5a^2(2iA + 5B)(a + ia \tan(e + fx))^{3/2}\sqrt{c - ic \tan(e + fx)}}{6c^2f}$$

```
output -5*a^(7/2)*(2*I*A+5*B)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/c^(3/2)/f+5/2*a^3*(2*I*A+5*B)*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)/c^2/f+5/6*a^2*(2*I*A+5*B)*(c-I*c*tan(f*x+e))^(1/2)*(a+I*a*tan(f*x+e))^(3/2)/c^2/f+2/3*a*(2*I*A+5*B)*(a+I*a*tan(f*x+e))^(5/2)/c/f/(c-I*c*tan(f*x+e))^(1/2)-1/3*(I*A+B)*(a+I*a*tan(f*x+e))^(7/2)/f/(c-I*c*tan(f*x+e))^(3/2)
```

3.822.2 Mathematica [A] (verified)

Time = 9.27 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.69

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \frac{a^3 \cos^2(e + fx) \sqrt{a + ia \tan(e + fx)} \left((-15 - 15i)(2A + B) \operatorname{ArcSin}\left[\frac{1}{2} + \frac{I}{2} \sqrt{-I + \tan(e + fx)}\right] \sqrt{2 - (2I) \tan(e + fx)} \right) + (-I + \tan(e + fx)) \left(-46A + (118I)B + ((68I)A + 161B) \tan(e + fx) + 6(A - (4I)B) \tan(e + fx)^2 + 3B \tan(e + fx)^3 \right)}{(6c f \sqrt{c - I c \tan(e + fx)})}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(3/2),x]`

output `(a^3*Cos[e + f*x]^2*Sqrt[a + I*a*Tan[e + f*x]]*((-15 - 15*I)*(2*A - (5*I)*B)*ArcSin[(1/2 + I/2)*Sqrt[-I + Tan[e + f*x]]]*Sqrt[2 - (2*I)*Tan[e + f*x]]*Sqrt[-I + Tan[e + f*x]]*(I + Tan[e + f*x]) + (-I + Tan[e + f*x])*(-46*A + (118*I)*B + ((68*I)*A + 161*B)*Tan[e + f*x] + 6*(A - (4*I)*B)*Tan[e + f*x]^2 + 3*B*Tan[e + f*x]^3)))/(6*c*f*Sqrt[c - I*c*Tan[e + f*x]])`

3.822.3 Rubi [A] (verified)Time = 0.46 (sec) , antiderivative size = 273, normalized size of antiderivative = 0.96, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.178$, Rules used = {3042, 4071, 87, 57, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx \\ \downarrow \text{3042} \\ \int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx \\ \downarrow \text{4071} \\ \frac{ac \int \frac{(i \tan(e+fx)a+a)^{5/2} (A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} d \tan(e + fx)}{f} \\ \downarrow \text{87} \end{array}$$

3.822. $\int \frac{(a+ia \tan(e+fx))^{7/2} (A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{ac \left(-\frac{(2A-5iB) \int \frac{(i \tan(e+fx)a+a)^{5/2}}{(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{3c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{f} \\
 & \quad \downarrow 57 \\
 & \frac{ac \left(-\frac{(2A-5iB) \left(-\frac{5a \int \frac{(i \tan(e+fx)a+a)^{3/2}}{\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{c} - \frac{2i(a+ia \tan(e+fx))^{5/2}}{c\sqrt{c-ic \tan(e+fx)}} \right)}{3c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{f} \\
 & \quad \downarrow 60 \\
 & \frac{ac \left(-\frac{(2A-5iB) \left(-\frac{5a \left(\frac{3}{2} a \int \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2c} \right)}{c} - \frac{2i(a+ia \tan(e+fx))^{5/2}}{c\sqrt{c-ic \tan(e+fx)}} \right)}{3c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{f} \\
 & \quad \downarrow 60 \\
 & \frac{ac \left(-\frac{(2A-5iB) \left(-\frac{5a \left(\frac{3}{2} a \left(a \int \frac{1}{\sqrt{i \tan(e+fx)a+a} \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i\sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{c} \right) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2c} \right)}{c} \right)}{3c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{f} \\
 & \quad \downarrow 45 \\
 & \frac{ac \left(-\frac{(2A-5iB) \left(-\frac{5a \left(\frac{3}{2} a \left(2a \int \frac{1}{\frac{ia+ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} + \frac{i\sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}}{c} \right) + \frac{i(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}}{2c} \right)}{c} \right)}{3c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{f} \\
 & \quad \downarrow 218
 \end{aligned}$$

3.822. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{3/2}} dx$

$$ac \left(\frac{(2A-5iB) \left(\frac{5a \left(\frac{3}{2} a \left(\frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{c} - \frac{2i\sqrt{a} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{\sqrt{c}} \right) + \frac{i(a+ia \tan(e+fx))^{3/2}\sqrt{c-ic \tan(e+fx)}}{2c} \right)}{c} \right)}{3c} \right) - \frac{2i}{c} \right) dx$$

input `Int[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(3/2), x]`

output `(a*c*(-1/3*((I*A + B)*(a + I*a*Tan[e + f*x])^(7/2)))/(a*c*(c - I*c*Tan[e + f*x])^(3/2)) - ((2*A - (5*I)*B)*((-2*I)*(a + I*a*Tan[e + f*x])^(5/2))/(c*Sqrt[c - I*c*Tan[e + f*x]]) - (5*a*(((I/2)*(a + I*a*Tan[e + f*x])^(3/2)*Sqrt[c - I*c*Tan[e + f*x]])/c + (3*a*(((2*I)*Sqrt[a]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[c] + (I*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/c))/2)/c)/(3*c)))/f`

3.822.3.1 Defintions of rubi rules used

rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 57 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))*Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.822.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 730 vs. 2(234) = 468.

Time = 0.38 (sec) , antiderivative size = 731, normalized size of antiderivative = 2.56

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^3 \left(-30 i A \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}}\right) a c-114 i A \sqrt{a c} \sqrt{a c}\right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^3 \left(-30 i A \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}}\right) a c-114 i A \sqrt{a c} \sqrt{a c}\right)}{\dots}$
parts	$\frac{A \sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)} a^3 \left(45 i \ln \left(\frac{a c \tan (f x+e)+\sqrt{a c} \sqrt{a c\left(1+\tan (f x+e)^2\right)}}{\sqrt{a c}}\right) \tan (f x+e)^2 a c+3 i \sqrt{a c}\right)}{\dots}$

```
input int((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x,m
method=_RETURNVERBOSE)
```

```
output 1/6/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^3/c^2*(-30*
I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/
2))*a*c-114*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+225*I*
B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2)
)*a*c*tan(f*x+e)+185*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+
e)^2+30*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*
c)^(1/2)*a*c*tan(f*x+e)^3+6*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*
tan(f*x+e)^3+3*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^4+2
25*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1
/2))*a*c*tan(f*x+e)^2+21*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*
x+e)^3-75*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)
)/(a*c)^(1/2))*a*c*tan(f*x+e)^3+90*I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*
(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c*tan(f*x+e)^2-90*A*ln((a*c*tan(f*
x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c*tan(f*x+e)
-74*A*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)*tan(f*x+e)^2-118*I*B*(a*c)^(
1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)-75*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*
c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c-279*B*(a*c)^(1/2)*(a*c*(1+tan(
f*x+e)^2))^(1/2)*tan(f*x+e)+46*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)
)/(a*c*(1+tan(f*x+e)^2))^(1/2)/(a*c)^(1/2)/(I+tan(f*x+e))^3
```

$$3.822. \int \frac{(a+ia \tan (e+fx))^{7/2}(A+B \tan (e+fx))}{(c-ic \tan (e+fx))^{3/2}} dx$$

3.822.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 571 vs. $2(221) = 442$.

Time = 0.28 (sec) , antiderivative size = 571, normalized size of antiderivative = 2.00

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx =$$

$$15 (c^2 f e^{(2i f x + 2i e)} + c^2 f) \sqrt{\frac{(4A^2 - 20iAB - 25B^2)a^7}{c^3 f^2}} \log \left(\frac{4 \left(2((-2iA - 5B)a^3 e^{(3i f x + 3i e)} + (-2iA - 5B)a^3 e^{(i f x + i e)}) \sqrt{\frac{a}{e^{(2i f x + 2i e)}}}}{(-2iA - 5B)a^3 e^{(2i f x + 2i e)}} \right)}{(-2iA - 5B)a^3 e^{(2i f x + 2i e)}} \right)$$

```
input integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="fricas")
```

```
output -1/12*(15*(c^2*f*e^(2*I*f*x + 2*I*e) + c^2*f)*sqrt((4*A^2 - 20*I*A*B - 25*B^2)*a^7/(c^3*f^2))*log(4*(2*((-2*I*A - 5*B)*a^3*e^(3*I*f*x + 3*I*e) + (-2*I*A - 5*B)*a^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + (c^2*f*e^(2*I*f*x + 2*I*e) - c^2*f)*sqrt((4*A^2 - 20*I*A*B - 25*B^2)*a^7/(c^3*f^2)))/((-2*I*A - 5*B)*a^3*e^(2*I*f*x + 2*I*e) + (-2*I*A - 5*B)*a^3)) - 15*(c^2*f*e^(2*I*f*x + 2*I*e) + c^2*f)*sqrt((4*A^2 - 20*I*A*B - 25*B^2)*a^7/(c^3*f^2))*log(4*(2*((-2*I*A - 5*B)*a^3*e^(3*I*f*x + 3*I*e) + (-2*I*A - 5*B)*a^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - (c^2*f*e^(2*I*f*x + 2*I*e) - c^2*f)*sqrt((4*A^2 - 20*I*A*B - 25*B^2)*a^7/(c^3*f^2)))/((-2*I*A - 5*B)*a^3*e^(2*I*f*x + 2*I*e) + (-2*I*A - 5*B)*a^3)) + 4*(4*(I*A + B)*a^3*e^(7*I*f*x + 7*I*e) + 8*(-2*I*A - 5*B)*a^3*e^(5*I*f*x + 5*I*e) + 25*(-2*I*A - 5*B)*a^3*e^(3*I*f*x + 3*I*e) + 15*(-2*I*A - 5*B)*a^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(c^2*f*e^(2*I*f*x + 2*I*e) + c^2*f)
```

3.822.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(3/2),x)`

output `Timed out`

3.822.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1176 vs. $2(221) = 442$.

Time = 0.56 (sec) , antiderivative size = 1176, normalized size of antiderivative = 4.13

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output

```

-2*(30*((2*A - 5*I*B)*a^3*cos(4*f*x + 4*e) + 2*(2*A - 5*I*B)*a^3*cos(2*f*x
+ 2*e) - (-2*I*A - 5*B)*a^3*sin(4*f*x + 4*e) - 2*(-2*I*A - 5*B)*a^3*sin(2
*f*x + 2*e) + (2*A - 5*I*B)*a^3)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))
+ 1) + 30*((2*A - 5*I*B)*a^3*cos(4*f*x + 4*e) + 2*(2*A - 5*I*B)*a^3*cos(2*
f*x + 2*e) - (-2*I*A - 5*B)*a^3*sin(4*f*x + 4*e) - 2*(-2*I*A - 5*B)*a^3*si
n(2*f*x + 2*e) + (2*A - 5*I*B)*a^3)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*
e), cos(2*f*x + 2*e))), -sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e
))) + 1) + 4*(4*(A - I*B)*a^3*cos(4*f*x + 4*e) + 8*(A - I*B)*a^3*cos(2*f*x
+ 2*e) - 4*(-I*A - B)*a^3*sin(4*f*x + 4*e) - 8*(-I*A - B)*a^3*sin(2*f*x +
2*e) - (2*A - 29*I*B)*a^3*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x +
2*e))) - 12*(8*(A - 2*I*B)*a^3*cos(4*f*x + 4*e) + 16*(A - 2*I*B)*a^3*cos(2
*f*x + 2*e) + 8*(I*A + 2*B)*a^3*sin(4*f*x + 4*e) + 16*(I*A + 2*B)*a^3*sin(
2*f*x + 2*e) + 5*(2*A - 5*I*B)*a^3*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(
2*f*x + 2*e))) - 15*((-2*I*A - 5*B)*a^3*cos(4*f*x + 4*e) + 2*(-2*I*A - 5*B
)*a^3*cos(2*f*x + 2*e) + (2*A - 5*I*B)*a^3*sin(4*f*x + 4*e) + 2*(2*A - 5*I
*B)*a^3*sin(2*f*x + 2*e) + (-2*I*A - 5*B)*a^3)*log(cos(1/2*arctan2(sin(2*f
*x + 2*e), cos(2*f*x + 2*e)))^2 + sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*
f*x + 2*e)))^2 + 2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) +
1) - 15*((2*I*A + 5*B)*a^3*cos(4*f*x + 4*e) + 2*(2*I*A + 5*B)*a^3*cos(2...

```

3.822.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{3/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{7/2}}{(-ic \tan(fx + e) + c)^{3/2}} dx$$

input

```

integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(3/
2),x, algorithm="giac")

```

output

```

integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(7/2)/(-I*c*tan(f*x
+ e) + c)^(3/2), x)

```

3.822.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{3/2}} dx = \int \frac{(A + B \tan(e + fx)) (a + a \tan(e + fx) \text{li})^{7/2}}{(c - c \tan(e + fx) \text{li})^{3/2}} dx$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2))/(c - c*tan(e + f*x)*1i)^(3/2), x)`

output `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2))/(c - c*tan(e + f*x)*1i)^(3/2), x)`

3.823
$$\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{5/2}} dx$$

3.823.1 Optimal result 7407
 3.823.2 Mathematica [A] (warning: unable to verify) 7408
 3.823.3 Rubi [A] (verified) 7408
 3.823.4 Maple [B] (verified) 7413
 3.823.5 Fricas [B] (verification not implemented) 7414
 3.823.6 Sympy [F(-1)] 7415
 3.823.7 Maxima [B] (verification not implemented) 7415
 3.823.8 Giac [F] 7416
 3.823.9 Mupad [F(-1)] 7417

3.823.1 Optimal result

Integrand size = 45, antiderivative size = 283

$$\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{5/2}} dx = \frac{2a^{7/2}(iA+6B) \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{c^{5/2}f} - \frac{(iA+B)(a+ia \tan(e+fx))^{7/2}}{5f(c-ictan(e+fx))^{5/2}} + \frac{2a(iA+6B)(a+ia \tan(e+fx))^{5/2}}{15cf(c-ictan(e+fx))^{3/2}} - \frac{2a^2(iA+6B)(a+ia \tan(e+fx))^{3/2}}{3c^2f\sqrt{c-ictan(e+fx)}} - \frac{a^3(iA+6B)\sqrt{a+ia \tan(e+fx)}\sqrt{c-ictan(e+fx)}}{c^3f}$$

output

```
2*a^(7/2)*(I*A+6*B)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c
*tan(f*x+e))^(1/2))/c^(5/2)/f-a^3*(I*A+6*B)*(a+I*a*tan(f*x+e))^(1/2)*(c-I*
c*tan(f*x+e))^(1/2)/c^3/f-2/3*a^2*(I*A+6*B)*(a+I*a*tan(f*x+e))^(3/2)/c^2/f
/(c-I*c*tan(f*x+e))^(1/2)-1/5*(I*A+B)*(a+I*a*tan(f*x+e))^(7/2)/f/(c-I*c*ta
n(f*x+e))^(5/2)+2/15*a*(I*A+6*B)*(a+I*a*tan(f*x+e))^(5/2)/c/f/(c-I*c*tan(f
*x+e))^(3/2)
```

3.823.2 Mathematica [A] (warning: unable to verify)

Time = 19.93 (sec) , antiderivative size = 528, normalized size of antiderivative = 1.87

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \frac{2(iA + 6B)e^{-i(4e+fx)} \sqrt{e^{ifx}} \sqrt{\frac{e^{i(e+fx)}}{1+e^{2i(e+fx)}}} \arctan(e^{i(e+fx)}}}{c^2 \sqrt{\frac{c}{1+e^{2i(e+fx)}}} f \sec^{\frac{9}{2}}(e + fx) (\cos(fx) + i \sin(fx))} + \frac{\cos^4(e + fx) \left((A - 6iB) \cos(2fx) \left(-\frac{2i \cos(e)}{3c^3} - \frac{2 \sin(e)}{3c^3} \right) + (iA + 6B) \cos(4fx) \left(\frac{2 \cos(e)}{15c^3} + \frac{2i \sin(e)}{15c^3} \right) + (A - 6iB) \cos(6fx) \left(-\frac{2i \cos(e)}{15c^3} - \frac{2 \sin(e)}{15c^3} \right) \right)}{c^2 \sqrt{\frac{c}{1+e^{2i(e+fx)}}} f \sec^{\frac{9}{2}}(e + fx) (\cos(fx) + i \sin(fx))}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(5/2),x]`

output `(2*(I*A + 6*B)*Sqrt[E^(I*f*x)]*Sqrt[E^(I*(e + f*x))/(1 + E^((2*I)*(e + f*x)))]*ArcTan[E^(I*(e + f*x))]*(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c^2*E^(I*(4*e + f*x))*Sqrt[c/(1 + E^((2*I)*(e + f*x)))]*f*Sec[e + f*x]^(9/2)*(Cos[f*x] + I*Sin[f*x])^(7/2)*(A*Cos[e + f*x] + B*Sin[e + f*x])) + (Cos[e + f*x]^4*((A - (6*I)*B)*Cos[2*f*x]*(((2*I)/3)*Cos[e])/c^3 - (2*Sin[e])/(3*c^3)) + (I*A + 6*B)*Cos[4*f*x]*(((2*I)/3)*Cos[e]/(15*c^3) + ((2*I)/15)*Sin[e])/c^3 + (A - (6*I)*B)*(((2*I)/3)*Cos[e]/c^3 - Sin[3*e]/c^3) + (A - I*B)*Cos[6*f*x]*(((2*I)/3)*Cos[e]/c^3 + Sin[3*e]/(5*c^3)) + (A - (6*I)*B)*(((2*I)/3)*Cos[e]/(3*c^3) - ((2*I)/3)*Sin[e]/c^3)*Sin[2*f*x] + (A - (6*I)*B)*((-2*cos[e])/(15*c^3) - ((2*I)/15)*Sin[e]/c^3)*Sin[4*f*x] + (A - I*B)*(Cos[3*e]/(5*c^3) + ((I/5)*Sin[3*e])/c^3)*Sin[6*f*x])*Sqrt[Sec[e + f*x]*(c*cos[e + f*x] - I*c*sin[e + f*x])]*(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(f*(Cos[f*x] + I*Sin[f*x])^3*(A*Cos[e + f*x] + B*Sin[e + f*x]))`

3.823.3 Rubi [A] (verified)Time = 0.44 (sec) , antiderivative size = 274, normalized size of antiderivative = 0.97, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.178$, Rules used = {3042, 4071, 87, 57, 57, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx$$

3.823. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$

$$\begin{array}{c}
 \downarrow 3042 \\
 \int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx \\
 \downarrow 4071 \\
 ac \int \frac{(i \tan(e + fx)a + a)^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} d \tan(e + fx) \\
 \downarrow 87 \\
 ac \left(\frac{(A - 6iB) \int \frac{(i \tan(e + fx)a + a)^{5/2}}{(c - ic \tan(e + fx))^{5/2}} d \tan(e + fx)}{5c} - \frac{(B + iA)(a + ia \tan(e + fx))^{7/2}}{5ac(c - ic \tan(e + fx))^{5/2}} \right) \\
 \downarrow 57 \\
 ac \left(\frac{(A - 6iB) \left(\frac{5a \int \frac{(i \tan(e + fx)a + a)^{3/2}}{(c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{3c} - \frac{2i(a + ia \tan(e + fx))^{5/2}}{3c(c - ic \tan(e + fx))^{3/2}} \right)}{5c} - \frac{(B + iA)(a + ia \tan(e + fx))^{7/2}}{5ac(c - ic \tan(e + fx))^{5/2}} \right) \\
 \downarrow 57 \\
 ac \left(\frac{(A - 6iB) \left(\frac{5a \left(\frac{3a \int \frac{\sqrt{i \tan(e + fx)a + a}}{\sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{c} - \frac{2i(a + ia \tan(e + fx))^{3/2}}{c \sqrt{c - ic \tan(e + fx)}} \right)}{3c} - \frac{2i(a + ia \tan(e + fx))^{5/2}}{3c(c - ic \tan(e + fx))^{3/2}} \right)}{5c} - \frac{(B + iA)(a + ia \tan(e + fx))^{7/2}}{5ac(c - ic \tan(e + fx))^{5/2}} \right) \\
 \downarrow 60
 \end{array}$$

3.823. $\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx$

$$ac \left((A-6iB) \left[\frac{5a \left(\frac{3a \left(a \int \frac{1}{\sqrt{i \tan(e+fx)a+a\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) + \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{c} \right)}{c} - \frac{2i(a+ia \tan(e+fx))^{3/2}}{c\sqrt{c-ic \tan(e+fx)}} \right)}{3c} \right] - \frac{2i(a+ia \tan(e+fx))^{3/2}}{3c(c-i)} \right)$$

f

↓ 45

$$ac \left((A-6iB) \left[\frac{5a \left(\frac{3a \left(2a \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} + \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{c} \right)}{c} - \frac{2i(a+ia \tan(e+fx))^{3/2}}{c\sqrt{c-ic \tan(e+fx)}} \right)}{3c} \right] - \frac{2i(a+ia \tan(e+fx))^{3/2}}{3c(c-i)} \right)$$

f

↓ 218

3.823. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$

3.823.3.1 Defintions of rubi rules used

- rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 57 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`
- rule 60 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !EqQ[e, 0] || !EqQ[c, 0] || LtQ[p, n]))`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.823.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 832 vs. 2(234) = 468.

Time = 0.37 (sec) , antiderivative size = 833, normalized size of antiderivative = 2.94

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^3 \left(246iB\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \tan(fx+e)^3 - 474iB\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \right)}{\dots}$
default	$\frac{\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^3 \left(246iB\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \tan(fx+e)^3 - 474iB\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)} \right)}{\dots}$
parts	Expression too large to display

```
input int((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x,m
ethod=_RETURNVERBOSE)
```

3.823. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{5/2}} dx$

output

```

-1/15/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^3/c^3*(24
6*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^3-474*I*B*(a*c)^(
1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+15*A*ln((a*c*tan(f*x+e)+(a*c
)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)^4+60*I*A
*a*c*tan(f*x+e)^3-94*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+
e)^2+360*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a
*c)^(1/2))*a*c*tan(f*x+e)^3+15*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*
tan(f*x+e)^4+540*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))
^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)^2-60*I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2
))*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)-90*A*ln((a*c*t
an(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f
*x+e)^2-46*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^3+26*I*A*
(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)-90*I*B*ln((a*c*tan(f*x+e)+(a*c)^(
1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)^4-360*B*ln(
(a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c
*tan(f*x+e)-564*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2-90
*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1
/2))*a*c+15*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))
/(a*c)^(1/2))*a*c+74*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x...
    
```

3.823.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 515 vs. 2(221) = 442.

Time = 0.27 (sec) , antiderivative size = 515, normalized size of antiderivative = 1.82

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \frac{15 c^3 \sqrt{\frac{(A^2 - 12i AB - 36 B^2) a^7}{c^5 f^2}} f \log \left(- \frac{4 \left(2 ((-i A - 6 B) a^3 e^{(3i f x - \dots)} \right)}{\dots} \right)}{\dots}$$

input

```

integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/
2),x, algorithm="fracas")
    
```

output $\frac{1}{30} \cdot (15 \cdot c^3 \cdot \sqrt{(A^2 - 12 \cdot I \cdot A \cdot B - 36 \cdot B^2) \cdot a^7 / (c^5 \cdot f^2)}) \cdot f \cdot \log(-4 \cdot (2 \cdot ((-I \cdot A - 6 \cdot B) \cdot a^3 \cdot e^{(3 \cdot I \cdot f \cdot x + 3 \cdot I \cdot e)} + (-I \cdot A - 6 \cdot B) \cdot a^3 \cdot e^{(I \cdot f \cdot x + I \cdot e)})) \cdot \sqrt{a / (e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} + 1)}) \cdot \sqrt{c / (e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} + 1)}) + (c^3 \cdot f \cdot e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} - c^3 \cdot f) \cdot \sqrt{(A^2 - 12 \cdot I \cdot A \cdot B - 36 \cdot B^2) \cdot a^7 / (c^5 \cdot f^2)}) / ((I \cdot A + 6 \cdot B) \cdot a^3 \cdot e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} + (I \cdot A + 6 \cdot B) \cdot a^3)) - 15 \cdot c^3 \cdot \sqrt{(A^2 - 12 \cdot I \cdot A \cdot B - 36 \cdot B^2) \cdot a^7 / (c^5 \cdot f^2)}) \cdot f \cdot \log(-4 \cdot (2 \cdot ((-I \cdot A - 6 \cdot B) \cdot a^3 \cdot e^{(3 \cdot I \cdot f \cdot x + 3 \cdot I \cdot e)} + (-I \cdot A - 6 \cdot B) \cdot a^3 \cdot e^{(I \cdot f \cdot x + I \cdot e)})) \cdot \sqrt{a / (e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} + 1)}) \cdot \sqrt{c / (e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} + 1)}) - (c^3 \cdot f \cdot e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} - c^3 \cdot f) \cdot \sqrt{(A^2 - 12 \cdot I \cdot A \cdot B - 36 \cdot B^2) \cdot a^7 / (c^5 \cdot f^2)}) / ((I \cdot A + 6 \cdot B) \cdot a^3 \cdot e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} + (I \cdot A + 6 \cdot B) \cdot a^3)) - 4 \cdot (3 \cdot (I \cdot A + B) \cdot a^3 \cdot e^{(7 \cdot I \cdot f \cdot x + 7 \cdot I \cdot e)} + 2 \cdot (-I \cdot A - 6 \cdot B) \cdot a^3 \cdot e^{(5 \cdot I \cdot f \cdot x + 5 \cdot I \cdot e)} + 10 \cdot (I \cdot A + 6 \cdot B) \cdot a^3 \cdot e^{(3 \cdot I \cdot f \cdot x + 3 \cdot I \cdot e)} + 15 \cdot (I \cdot A + 6 \cdot B) \cdot a^3 \cdot e^{(I \cdot f \cdot x + I \cdot e)}) \cdot \sqrt{a / (e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} + 1)}) \cdot \sqrt{c / (e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} + 1)}) / (c^3 \cdot f)$

3.823.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(5/2),x)`

output Timed out

3.823.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 936 vs. $2(221) = 442$.

Time = 0.50 (sec) , antiderivative size = 936, normalized size of antiderivative = 3.31

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="maxima")`

3.823. $\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx$

output

```

15*(30*((A - 6*I*B)*a^3*cos(2*f*x + 2*e) - (-I*A - 6*B)*a^3*sin(2*f*x + 2*
e) + (A - 6*I*B)*a^3)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x
+ 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) + 30*(
(A - 6*I*B)*a^3*cos(2*f*x + 2*e) - (-I*A - 6*B)*a^3*sin(2*f*x + 2*e) + (A
- 6*I*B)*a^3)*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))
, -sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) - 12*((A - I*
B)*a^3*cos(2*f*x + 2*e) + (I*A + B)*a^3*sin(2*f*x + 2*e) + (A - I*B)*a^3)*
cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 20*((A - 3*I*B)*a^3
*cos(2*f*x + 2*e) - (-I*A - 3*B)*a^3*sin(2*f*x + 2*e) + (A - 3*I*B)*a^3)*c
os(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 60*((A - 5*I*B)*a^3*
cos(2*f*x + 2*e) + (I*A + 5*B)*a^3*sin(2*f*x + 2*e) + (A - 6*I*B)*a^3)*cos
(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 15*((-I*A - 6*B)*a^3*c
os(2*f*x + 2*e) + (A - 6*I*B)*a^3*sin(2*f*x + 2*e) + (-I*A - 6*B)*a^3)*log
(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))^2 + sin(1/2*arctan2(
sin(2*f*x + 2*e), cos(2*f*x + 2*e)))^2 + 2*sin(1/2*arctan2(sin(2*f*x + 2*e
), cos(2*f*x + 2*e))) + 1) - 15*((I*A + 6*B)*a^3*cos(2*f*x + 2*e) - (A - 6
*I*B)*a^3*sin(2*f*x + 2*e) + (I*A + 6*B)*a^3)*log(cos(1/2*arctan2(sin(2*f*
x + 2*e), cos(2*f*x + 2*e))))^2 + sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f
*x + 2*e)))^2 - 2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1
) - 12*((I*A + B)*a^3*cos(2*f*x + 2*e) - (A - I*B)*a^3*sin(2*f*x + 2*e)...

```

3.823.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{5/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{7/2}}{(-ic \tan(fx + e) + c)^{5/2}} dx$$

input

```

integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(5/
2),x, algorithm="giac")

```

output

```

integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(7/2)/(-I*c*tan(f*x
+ e) + c)^(5/2), x)

```

3.823.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{5/2}} dx = \int \frac{(A + B \tan(e + fx)) (a + a \tan(e + fx) \text{li})^{7/2}}{(c - c \tan(e + fx) \text{li})^{5/2}} dx$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2))/(c - c*tan(e + f*x)*1i)^(5/2), x)`

output `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2))/(c - c*tan(e + f*x)*1i)^(5/2), x)`

3.824
$$\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$$

3.824.1 Optimal result	7418
3.824.2 Mathematica [B] (warning: unable to verify)	7419
3.824.3 Rubi [A] (verified)	7420
3.824.4 Maple [B] (verified)	7425
3.824.5 Fricas [B] (verification not implemented)	7426
3.824.6 Sympy [F(-1)]	7426
3.824.7 Maxima [A] (verification not implemented)	7427
3.824.8 Giac [F]	7427
3.824.9 Mupad [F(-1)]	7428

3.824.1 Optimal result

Integrand size = 45, antiderivative size = 251

$$\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx =$$

$$\frac{2a^{7/2} B \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{c^{7/2} f}$$

$$- \frac{(iA + B)(a + ia \tan(e + fx))^{7/2}}{7f(c - ic \tan(e + fx))^{7/2}} + \frac{2aB(a + ia \tan(e + fx))^{5/2}}{5cf(c - ic \tan(e + fx))^{5/2}}$$

$$- \frac{2a^2 B(a + ia \tan(e + fx))^{3/2}}{3c^2 f(c - ic \tan(e + fx))^{3/2}} + \frac{2a^3 B \sqrt{a + ia \tan(e + fx)}}{c^3 f \sqrt{c - ic \tan(e + fx)}}$$

output

```
-2*a^(7/2)*B*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/c^(7/2)/f+2*a^3*B*(a+I*a*tan(f*x+e))^(1/2)/c^3/f/(c-I*c*tan(f*x+e))^(1/2)-1/7*(I*A+B)*(a+I*a*tan(f*x+e))^(7/2)/f/(c-I*c*tan(f*x+e))^(7/2)+2/5*a*B*(a+I*a*tan(f*x+e))^(5/2)/c/f/(c-I*c*tan(f*x+e))^(5/2)-2/3*a^2*B*(a+I*a*tan(f*x+e))^(3/2)/c^2/f/(c-I*c*tan(f*x+e))^(3/2)
```

3.824.2 Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 570 vs. $2(251) = 502$.

Time = 19.88 (sec) , antiderivative size = 570, normalized size of antiderivative = 2.27

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx =$$

$$\frac{2Be^{-i(4e+fx)} \sqrt{e^{ifx}} \sqrt{\frac{e^{i(e+fx)}}{1+e^{2i(e+fx)}}} \arctan(e^{i(e+fx)}) (a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{c^3 \sqrt{\frac{c}{1+e^{2i(e+fx)}}} f \sec^{\frac{9}{2}}(e + fx) (\cos(fx) + i \sin(fx))^{7/2} (A \cos(e + fx) + B \sin(e + fx))}$$

$$+ \frac{\cos^4(e + fx) \left(\frac{B \cos(3e)}{c^4} + \cos(4fx) \left(-\frac{2B \cos(e)}{15c^4} - \frac{2iB \sin(e)}{15c^4} \right) + \cos(2fx) \left(\frac{2B \cos(e)}{3c^4} - \frac{2iB \sin(e)}{3c^4} \right) - \frac{iB \sin(3e)}{c^4} + \dots \right)}{}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(7/2),x]`

output `(-2*B*Sqrt[E^(I*f*x)]*Sqrt[E^(I*(e + f*x))/(1 + E^((2*I)*(e + f*x)))]*ArcTan[E^(I*(e + f*x))]*(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c^3*E^(I*(4*e + f*x))*Sqrt[c/(1 + E^((2*I)*(e + f*x)))]*f*Sec[e + f*x]^(9/2)*(Cos[f*x] + I*Sin[f*x])^(7/2)*(A*Cos[e + f*x] + B*Sin[e + f*x])) + (Cos[e + f*x]^4*((B*Cos[3*e])/c^4 + Cos[4*f*x]*((-2*B*Cos[e])/(15*c^4) - ((2*I)/15)*B*Sin[e])/c^4) + Cos[2*f*x]*((2*B*Cos[e])/(3*c^4) - ((2*I)/3)*B*Sin[e])/c^4 - (I*B*Sin[3*e])/c^4 + ((-5*I)*A + 9*B)*Cos[6*f*x]*(Cos[3*e]/(70*c^4) + ((I/70)*Sin[3*e])/c^4) + (A - I*B)*Cos[8*f*x]*(((1/14*I)*Cos[5*e])/c^4 + Sin[5*e]/(14*c^4)) + (((2*I)/3)*B*Cos[e])/c^4 + (2*B*Sin[e])/(3*c^4))*Sin[2*f*x] + ((((-2*I)/15)*B*Cos[e])/c^4 + (2*B*Sin[e])/(15*c^4))*Sin[4*f*x] + (5*A + (9*I)*B)*(Cos[3*e]/(70*c^4) + ((I/70)*Sin[3*e])/c^4)*Sin[6*f*x] + (A - I*B)*(Cos[5*e]/(14*c^4) + ((I/14)*Sin[5*e])/c^4)*Sin[8*f*x])*Sqrt[Sec[e + f*x]*(c*Cos[e + f*x] - I*c*Sin[e + f*x])*(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x])]/(f*(Cos[f*x] + I*Sin[f*x])^3*(A*Cos[e + f*x] + B*Sin[e + f*x]))`

3.824.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.07, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.178$, Rules used = {3042, 4071, 87, 57, 57, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx \\
 \downarrow \text{3042} \\
 \int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx \\
 \downarrow \text{4071} \\
 \frac{ac \int \frac{(i \tan(e + fx)a + a)^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} d \tan(e + fx)}{f} \\
 \downarrow \text{87} \\
 \frac{ac \left(\frac{iB \int \frac{(i \tan(e + fx)a + a)^{5/2}}{(c - ic \tan(e + fx))^{7/2}} d \tan(e + fx)}{c} - \frac{(B + iA)(a + ia \tan(e + fx))^{7/2}}{7ac(c - ic \tan(e + fx))^{7/2}} \right)}{f} \\
 \downarrow \text{57} \\
 \frac{ac \left(\frac{iB \left(-\frac{a \int \frac{(i \tan(e + fx)a + a)^{3/2}}{(c - ic \tan(e + fx))^{5/2}} d \tan(e + fx)}{c} - \frac{2i(a + ia \tan(e + fx))^{5/2}}{5c(c - ic \tan(e + fx))^{5/2}} \right)}{c} - \frac{(B + iA)(a + ia \tan(e + fx))^{7/2}}{7ac(c - ic \tan(e + fx))^{7/2}} \right)}{f} \\
 \downarrow \text{57}
 \end{array}$$

3.824. $\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx$

$$\left(\begin{array}{l} iB \\ ac \end{array} \right) \left(\begin{array}{l} a \left(-\frac{a \int \frac{\sqrt{i \tan(e+fx)a+a}}{(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{c} - \frac{2i(a+ia \tan(e+fx))^{3/2}}{3c(c-ic \tan(e+fx))^{3/2}} \right) \\ - \frac{2i(a+ia \tan(e+fx))^{5/2}}{5c(c-ic \tan(e+fx))^{5/2}} \end{array} \right) - \frac{(B+IA)(a+ia \tan(e+fx))^{7/2}}{7ac(c-ic \tan(e+fx))^{7/2}}$$

f
↓ 57

$$\left(\begin{array}{l} iB \\ ac \end{array} \right) \left(\begin{array}{l} a \left(-\frac{a \int \frac{1}{\sqrt{i \tan(e+fx)a+a\sqrt{c-ic \tan(e+fx)}}} d \tan(e+fx)}{c} - \frac{2i\sqrt{a+ia \tan(e+fx)}}{c\sqrt{c-ic \tan(e+fx)}} \right) \\ - \frac{2i(a+ia \tan(e+fx))^{3/2}}{3c(c-ic \tan(e+fx))^{3/2}} \end{array} \right) - \frac{2i(a+ia \tan(e+fx))^{5/2}}{5c(c-ic \tan(e+fx))^{5/2}}$$

f
↓ 45

3.824. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$

$$\left(\begin{array}{l}
 \left(\begin{array}{l}
 \left(\begin{array}{l}
 2a \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} \\
 - \frac{2i \sqrt{a+ia \tan(e+fx)}}{c \sqrt{c-ic \tan(e+fx)}}
 \end{array} \right) \\
 - \frac{2i(a+ia \tan(e+fx))^{3/2}}{3c(c-ic \tan(e+fx))^{3/2}}
 \end{array} \right) \\
 - \frac{2i(a+ia \tan(e+fx))^{5/2}}{5c(c-ic \tan(e+fx))^{5/2}}
 \end{array} \right) \\
 - \frac{2i(a+ia \tan(e+fx))^{7/2}}{7c(c-ic \tan(e+fx))^{7/2}}
 \end{array} \right)$$

218

3.824. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$

$$\begin{aligned}
 & \left(\frac{a \left(\frac{2i\sqrt{a} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right) - \frac{2i\sqrt{a+ia \tan(e+fx)}}{c\sqrt{c-ic \tan(e+fx)}}}{c^{3/2}} \right) - \frac{2i(a+ia \tan(e+fx))^{3/2}}{3c(c-ic \tan(e+fx))^{3/2}}}{c} \right) \\
 & \frac{iB}{c} - \frac{2i(a+ia \tan(e+fx))^{5/2}}{5c(c-ic \tan(e+fx))^{5/2}} \\
 & \frac{aC}{c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{7ac(c-ic \tan(e+fx))^{7/2}}
 \end{aligned}$$

f

input `Int[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(7/2),x]`

output `(a*c*(-1/7*((I*A + B)*(a + I*a*Tan[e + f*x])^(7/2)))/(a*c*(c - I*c*Tan[e + f*x])^(7/2)) + (I*B*((((-2*I)/5)*(a + I*a*Tan[e + f*x])^(5/2))/(c*(c - I*c*Tan[e + f*x])^(5/2)) - (a*((((-2*I)/3)*(a + I*a*Tan[e + f*x])^(3/2))/(c*(c - I*c*Tan[e + f*x])^(3/2)) - (a*(((2*I)*Sqrt[a]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/c^(3/2) - ((2*I)*Sqrt[a + I*a*Tan[e + f*x]])/(c*Sqrt[c - I*c*Tan[e + f*x]])))/c))/c)/c)/f`

3.824.3.1 Defintions of rubi rules used

- rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`
- rule 57 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] &
& GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m
+ n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c
, d, m, n, x]`
- rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`
- rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]`

3.824.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 597 vs. 2(206) = 412.

Time = 0.34 (sec) , antiderivative size = 598, normalized size of antiderivative = 2.38

method	result
parts	$\frac{A\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^3(1+\tan(fx+e)^2)(-\tan(fx+e)^2+2i\tan(fx+e)+1)}{7fc^4(i+\tan(fx+e))^5} - \frac{iB\sqrt{a(1+i\tan(fx+e))}}{c}$
derivativedivides	$\frac{\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^3\left(-105iB\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}}\right)\right)ac\tan(fx+e)^5+10}{c}$
default	$\frac{\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}a^3\left(-105iB\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}}\right)\right)ac\tan(fx+e)^5+10}{c}$

```
input int((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x,method=_RETURNVERBOSE)
```

```
output 1/7*A/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^3/c^4*(1+tan(f*x+e)^2)*(-tan(f*x+e)^2+2*I*tan(f*x+e)+1)/(I+tan(f*x+e))^5-1/105*I*B/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)*a^3/c^4*(525*I*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)))^(1/2))/(a*c)^(1/2))*tan(f*x+e)^4*a*c+105*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)))^(1/2))/(a*c)^(1/2))*tan(f*x+e)^5*a*c-1050*I*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)))^(1/2))/(a*c)^(1/2))*tan(f*x+e)^2*a*c-1050*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)))^(1/2))/(a*c)^(1/2))*tan(f*x+e)^3*a*c-950*I*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)*tan(f*x+e)^3-337*tan(f*x+e)^4*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)+105*I*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)))^(1/2))/(a*c)^(1/2))*a*c+525*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2)))^(1/2))/(a*c)^(1/2))*tan(f*x+e)*a*c+730*I*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)*tan(f*x+e)+1176*tan(f*x+e)^2*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)-167*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c*(1+tan(f*x+e)^2))^(1/2)/(I+tan(f*x+e))^5/(a*c)^(1/2)
```

3.824.
$$\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{7/2}} dx$$

3.824.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 434 vs. $2(197) = 394$.

Time = 0.28 (sec) , antiderivative size = 434, normalized size of antiderivative = 1.73

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \frac{105 c^4 f \sqrt{-\frac{B^2 a^7}{c^7 f^2}} \log \left(\frac{4 \left(2 (Ba^3 e^{(3i fx + 3i e)} + Ba^3 e^{(i fx + i e)}) \sqrt{\frac{a}{e^{(2i fx + 2i e)} + 1}} \right)}{\dots} \right)}{\dots}$$

```
input integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x, algorithm="fricas")
```

```
output 1/210*(105*c^4*f*sqrt(-B^2*a^7/(c^7*f^2))*log(4*(2*(B*a^3*e^(3*I*f*x + 3*I*e) + B*a^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + (c^4*f*e^(2*I*f*x + 2*I*e) - c^4*f)*sqrt(-B^2*a^7/(c^7*f^2)))/(B*a^3*e^(2*I*f*x + 2*I*e) + B*a^3)) - 105*c^4*f*sqrt(-B^2*a^7/(c^7*f^2))*log(4*(2*(B*a^3*e^(3*I*f*x + 3*I*e) + B*a^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - (c^4*f*e^(2*I*f*x + 2*I*e) - c^4*f)*sqrt(-B^2*a^7/(c^7*f^2)))/(B*a^3*e^(2*I*f*x + 2*I*e) + B*a^3)) - 2*(15*(I*A + B)*a^3*e^(9*I*f*x + 9*I*e) + 3*(5*I*A - 9*B)*a^3*e^(7*I*f*x + 7*I*e) + 28*B*a^3*e^(5*I*f*x + 5*I*e) - 140*B*a^3*e^(3*I*f*x + 3*I*e) - 210*B*a^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(c^4*f)
```

3.824.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \text{Timed out}$$

```
input integrate((a+I*a*tan(f*x+e))**(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(7/2),x)
```

```
output Timed out
```

3.824.7 Maxima [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 246, normalized size of antiderivative = 0.98

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx =$$

$$(210 Ba^3 \arctan(\cos(fx + e), \sin(fx + e) + 1) + 210 Ba^3 \arctan(\cos(fx + e), -\sin(fx + e) + 1) - 30$$

```
input integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x, algorithm="maxima")
```

```
output -1/210*(210*B*a^3*arctan2(cos(f*x + e), sin(f*x + e) + 1) + 210*B*a^3*arctan2(cos(f*x + e), -sin(f*x + e) + 1) - 30*(-I*A - B)*a^3*cos(7*f*x + 7*e) - 84*B*a^3*cos(5*f*x + 5*e) + 140*B*a^3*cos(3*f*x + 3*e) - 420*B*a^3*cos(f*x + e) + 105*I*B*a^3*log(cos(f*x + e)^2 + sin(f*x + e)^2 + 2*sin(f*x + e) + 1) - 105*I*B*a^3*log(cos(f*x + e)^2 + sin(f*x + e)^2 - 2*sin(f*x + e) + 1) - 30*(A - I*B)*a^3*sin(7*f*x + 7*e) - 84*I*B*a^3*sin(5*f*x + 5*e) + 140*I*B*a^3*sin(3*f*x + 3*e) - 420*I*B*a^3*sin(f*x + e))*sqrt(a)/(c^(7/2)*f)
```

3.824.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{7/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{7/2}}{(-ic \tan(fx + e) + c)^{7/2}} dx$$

```
input integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(7/2),x, algorithm="giac")
```

```
output integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(7/2)/(-I*c*tan(f*x + e) + c)^(7/2), x)
```

3.824.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{7/2}} dx = \int \frac{(A + B \tan(e + fx)) (a + a \tan(e + fx) \text{li})^{7/2}}{(c - c \tan(e + fx) \text{li})^{7/2}} dx$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2))/(c - c*tan(e + f*x)*1i)^(7/2),x)`

output `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2))/(c - c*tan(e + f*x)*1i)^(7/2), x)`

3.825
$$\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{9/2}} dx$$

3.825.1 Optimal result	7429
3.825.2 Mathematica [A] (verified)	7429
3.825.3 Rubi [A] (verified)	7430
3.825.4 Maple [A] (verified)	7431
3.825.5 Fricas [A] (verification not implemented)	7432
3.825.6 Sympy [F(-1)]	7432
3.825.7 Maxima [B] (verification not implemented)	7433
3.825.8 Giac [F]	7433
3.825.9 Mupad [B] (verification not implemented)	7434

3.825.1 Optimal result

Integrand size = 45, antiderivative size = 102

$$\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx =$$

$$-\frac{(iA + B)(a + ia \tan(e + fx))^{7/2}}{9f(c - ic \tan(e + fx))^{9/2}} - \frac{(iA - 8B)(a + ia \tan(e + fx))^{7/2}}{63cf(c - ic \tan(e + fx))^{7/2}}$$

output

```
-1/9*(I*A+B)*(a+I*a*tan(f*x+e))^(7/2)/f/(c-I*c*tan(f*x+e))^(9/2)-1/63*(I*A
-8*B)*(a+I*a*tan(f*x+e))^(7/2)/c/f/(c-I*c*tan(f*x+e))^(7/2)
```

3.825.2 Mathematica [A] (verified)

Time = 6.96 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.09

$$\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx =$$

$$\frac{a^4 \sec^4(e + fx)(\cos(4(e + fx)) + i \sin(4(e + fx)))(8iA - B + (A + 8iB) \tan(e + fx))}{63c^4 f (i + \tan(e + fx))^4 \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}}$$

input

```
Integrate[(((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan
[e + f*x])^(9/2), x]
```

output
$$-1/63*(a^4*\text{Sec}[e + f*x]^4*(\text{Cos}[4*(e + f*x)] + I*\text{Sin}[4*(e + f*x)])*((8*I)*A - B + (A + (8*I)*B)*\text{Tan}[e + f*x]))/(c^4*f*(I + \text{Tan}[e + f*x])^4*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])$$

3.825.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.089$, Rules used = {3042, 4071, 87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{9/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{9/2}} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(i \tan(e+fx)a+a)^{5/2} (A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{11/2}} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{87} \\ & \frac{ac \left(\frac{(A+8iB) \int \frac{(i \tan(e+fx)a+a)^{5/2} d \tan(e+fx)}{(c-ict \tan(e+fx))^{9/2}}}{9c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{9ac(c-ict \tan(e+fx))^{9/2}} \right)}{f} \\ & \quad \downarrow \text{48} \\ & \frac{ac \left(-\frac{i(A+8iB)(a+ia \tan(e+fx))^{7/2}}{63ac^2(c-ict \tan(e+fx))^{7/2}} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{9ac(c-ict \tan(e+fx))^{9/2}} \right)}{f} \end{aligned}$$

input
$$\text{Int}[(a + I*a*\text{Tan}[e + f*x])^{(7/2)}*(A + B*\text{Tan}[e + f*x])]/(c - I*c*\text{Tan}[e + f*x])^{(9/2)}, x]$$

output
$$(a*c*(-1/9*((I*A + B)*(a + I*a*\text{Tan}[e + f*x])^{(7/2)}))/(a*c*(c - I*c*\text{Tan}[e + f*x])^{(9/2)}) - ((I/63)*(A + (8*I)*B)*(a + I*a*\text{Tan}[e + f*x])^{(7/2)}))/(a*c^2*(c - I*c*\text{Tan}[e + f*x])^{(7/2)}))/f$$

3.825.
$$\int \frac{(a+ia \tan(e+fx))^{7/2} (A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{9/2}} dx$$

3.825.3.1 Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.825.4 Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.04

method	result
risch	$-\frac{a^3 \sqrt{\frac{a e^{2i(fx+e)}}{e^{2i(fx+e)}+1}} (7iA e^{8i(fx+e)}+7B e^{8i(fx+e)}+9iA e^{6i(fx+e)}-9B e^{6i(fx+e)})}{126c^4 \sqrt{\frac{c}{e^{2i(fx+e)}+1}} f}$
derivativedivides	$-\frac{\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^3 (1+\tan(fx+e)^2) (8iB \tan(fx+e)^3+6iA \tan(fx+e)^2+A \tan(fx+e)^3-63f c^5(i+\tan(fx+e))^6)}{63f c^5(i+\tan(fx+e))^6}$
default	$-\frac{\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^3 (1+\tan(fx+e)^2) (8iB \tan(fx+e)^3+6iA \tan(fx+e)^2+A \tan(fx+e)^3-63f c^5(i+\tan(fx+e))^6)}{63f c^5(i+\tan(fx+e))^6}$
parts	$-\frac{A \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^3 (1+\tan(fx+e)^2) (6i \tan(fx+e)^2+\tan(fx+e)^3-8i+15 \tan(fx+e))}{63f c^5(i+\tan(fx+e))^6}$

3.825.
$$\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{9/2}} dx$$

```
input int((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(9/2),x,m
method=_RETURNVERBOSE)
```

```
output -1/126*a^3/c^4*(a*exp(2*I*(f*x+e))/(exp(2*I*(f*x+e))+1))^(1/2)/(c/(exp(2*I
*(f*x+e))+1))^(1/2)/f*(7*I*A*exp(8*I*(f*x+e))+7*B*exp(8*I*(f*x+e))+9*I*A*exp
(6*I*(f*x+e))-9*B*exp(6*I*(f*x+e)))
```

3.825.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.02

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx =$$

$$\frac{(7(iA + B)a^3 e^{(11i fx + 11ie)} + 2(8iA - B)a^3 e^{(9i fx + 9ie)} + 9(iA - B)a^3 e^{(7i fx + 7ie)}) \sqrt{\frac{a}{e^{(2i fx + 2ie)} + 1}} \sqrt{\frac{c}{e^{(2i fx + 2ie)} + 1}}}{126 c^5 f}$$

```
input integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(9/
2),x, algorithm="fricas")
```

```
output -1/126*(7*(I*A + B)*a^3*e^(11*I*f*x + 11*I*e) + 2*(8*I*A - B)*a^3*e^(9*I*f
*x + 9*I*e) + 9*(I*A - B)*a^3*e^(7*I*f*x + 7*I*e))*sqrt(a/(e^(2*I*f*x + 2*
I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^5*f)
```

3.825.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx = \text{Timed out}$$

```
input integrate((a+I*a*tan(f*x+e))**(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(
9/2),x)
```

```
output Timed out
```

3.825. $\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx$

3.825.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 166 vs. $2(78) = 156$.

Time = 0.44 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.63

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx =$$

$$\frac{126(7(A - iB)a^3 \cos(11fx + 11e) + 2(8A + iB)a^3 \cos(9fx + 9e) + 9(A + iB)a^3 \cos(7fx + 7e) - 7(-IA - B)a^3 \sin(11fx + 11e) - 2(-8IA + B)a^3 \sin(9fx + 9e) - 9(-IA + B)a^3 \sin(7fx + 7e)) \sqrt{a} \sqrt{c}}{-15876(i c^5 \cos(2fx + 2e) + 15876 I c^5 \sin(2fx + 2e)) \sqrt{c} + 15876 c^5 \sin(2fx + 2e)}$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(9/2),x, algorithm="maxima")`

output `-126*(7*(A - I*B)*a^3*cos(11*f*x + 11*e) + 2*(8*A + I*B)*a^3*cos(9*f*x + 9*e) + 9*(A + I*B)*a^3*cos(7*f*x + 7*e) - 7*(-I*A - B)*a^3*sin(11*f*x + 11*e) - 2*(-8*I*A + B)*a^3*sin(9*f*x + 9*e) - 9*(-I*A + B)*a^3*sin(7*f*x + 7*e))*sqrt(a)*sqrt(c)/((-15876*I*c^5*cos(2*f*x + 2*e) + 15876*c^5*sin(2*f*x + 2*e) - 15876*I*c^5)*f)`

3.825.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{7/2}}{(-ic \tan(fx + e) + c)^{9/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(9/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(7/2)/(-I*c*tan(f*x + e) + c)^(9/2), x)`

3.825.9 Mupad [B] (verification not implemented)

Time = 10.60 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.88

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{9/2}} dx =$$

$$\frac{a^3 \sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)i)}{\cos(2e+2fx)+1}} (A \cos(6e+6fx) 9i + A \cos(8e+8fx) 7i - 9B \cos(6e+6fx) + 7B \cos(8e+8fx) - 9A \sin(6e+6fx) - 7A \sin(8e+8fx) - B \sin(6e+6fx) 9i + B \sin(8e+8fx) 7i)}{126 c^4 f \sqrt{\frac{c(\cos(2e+2fx)-1+i \sin(2e+2fx))}{c(\cos(2e+2fx)+1)}}}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2))/(c - c*tan(e + f*x)*1i)^(9/2),x)`

output `-(a^3*((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*cos(6*e + 6*f*x)*9i + A*cos(8*e + 8*f*x)*7i - 9*B*cos(6*e + 6*f*x) + 7*B*cos(8*e + 8*f*x) - 9*A*sin(6*e + 6*f*x) - 7*A*sin(8*e + 8*f*x) - B*sin(6*e + 6*f*x)*9i + B*sin(8*e + 8*f*x)*7i))/(126*c^4*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

3.826
$$\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{11/2}} dx$$

3.826.1 Optimal result	7435
3.826.2 Mathematica [B] (verified)	7435
3.826.3 Rubi [A] (verified)	7436
3.826.4 Maple [A] (verified)	7438
3.826.5 Fricas [A] (verification not implemented)	7439
3.826.6 Sympy [F(-1)]	7439
3.826.7 Maxima [A] (verification not implemented)	7439
3.826.8 Giac [F]	7440
3.826.9 Mupad [B] (verification not implemented)	7440

3.826.1 Optimal result

Integrand size = 45, antiderivative size = 155

$$\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{11/2}} dx = -\frac{(iA + B)(a + ia \tan(e + fx))^{7/2}}{11f(c - ic \tan(e + fx))^{11/2}} - \frac{(2iA - 9B)(a + ia \tan(e + fx))^{7/2}}{99cf(c - ic \tan(e + fx))^{9/2}} - \frac{(2iA - 9B)(a + ia \tan(e + fx))^{7/2}}{693c^2f(c - ic \tan(e + fx))^{7/2}}$$

output `-1/11*(I*A+B)*(a+I*a*tan(f*x+e))^(7/2)/f/(c-I*c*tan(f*x+e))^(11/2)-1/99*(2*I*A-9*B)*(a+I*a*tan(f*x+e))^(7/2)/c/f/(c-I*c*tan(f*x+e))^(9/2)-1/693*(2*I*A-9*B)*(a+I*a*tan(f*x+e))^(7/2)/c^2/f/(c-I*c*tan(f*x+e))^(7/2)`

3.826.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 417 vs. 2(155) = 310.

Time = 18.82 (sec) , antiderivative size = 417, normalized size of antiderivative = 2.69

$$\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{11/2}} dx = \frac{\cos^4(e + fx) \left((-iA + B) \cos(6fx) \left(\frac{\cos(3e)}{56c^6} + \frac{i \sin(3e)}{56c^6} \right) \right)}{1}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(11/2), x]`

output $(\text{Cos}[e + f*x]^4 * ((-I)*A + B) * \text{Cos}[6*f*x] * (\text{Cos}[3*e]/(56*c^6) + ((I/56)*\text{Sin}[3*e])/c^6) + ((-23*I)*A + 9*B) * \text{Cos}[8*f*x] * (\text{Cos}[5*e]/(504*c^6) + ((I/504)*\text{Sin}[5*e])/c^6) + (31*A - (9*I)*B) * \text{Cos}[10*f*x] * (((-1/792*I)*\text{Cos}[7*e])/c^6 + \text{Sin}[7*e]/(792*c^6)) + (A - I*B) * \text{Cos}[12*f*x] * (((-1/88*I)*\text{Cos}[9*e])/c^6 + \text{Sin}[9*e]/(88*c^6)) + (A + I*B) * (\text{Cos}[3*e]/(56*c^6) + ((I/56)*\text{Sin}[3*e])/c^6) * \text{Sin}[6*f*x] + (23*A + (9*I)*B) * (\text{Cos}[5*e]/(504*c^6) + ((I/504)*\text{Sin}[5*e])/c^6) * \text{Sin}[8*f*x] + (31*A - (9*I)*B) * (\text{Cos}[7*e]/(792*c^6) + ((I/792)*\text{Sin}[7*e])/c^6) * \text{Sin}[10*f*x] + (A - I*B) * (\text{Cos}[9*e]/(88*c^6) + ((I/88)*\text{Sin}[9*e])/c^6) * \text{Sin}[12*f*x] * \text{Sqrt}[\text{Sec}[e + f*x] * (c * \text{Cos}[e + f*x] - I * c * \text{Sin}[e + f*x])] * (a + I * a * \text{Tan}[e + f*x])^{7/2} * (A + B * \text{Tan}[e + f*x])]) / (f * (\text{Cos}[f*x] + I * \text{Sin}[f*x])^3 * (A * \text{Cos}[e + f*x] + B * \text{Sin}[e + f*x]))$

3.826.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.08, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4071, 87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{11/2}} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{11/2}} dx$$

↓ 4071

$$ac \int \frac{(i \tan(e + fx)a + a)^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} d \tan(e + fx)$$

f

↓ 87

$$ac \left(\frac{(2A + 9iB) \int \frac{(i \tan(e + fx)a + a)^{5/2}}{(c - ic \tan(e + fx))^{11/2}} d \tan(e + fx)}{11c} - \frac{(B + iA)(a + ia \tan(e + fx))^{7/2}}{11ac(c - ic \tan(e + fx))^{11/2}} \right)$$

f

↓ 55

3.826. $\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{11/2}} dx$

$$ac \left(\frac{(2A+9iB) \left(\frac{\int \frac{(i \tan(e+fx)a+a)^{5/2}}{(c-ic \tan(e+fx))^{9/2}} d \tan(e+fx)}{9c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)$$

f
↓
48

$$ac \left(\frac{(2A+9iB) \left(-\frac{i(a+ia \tan(e+fx))^{7/2}}{63ac^2(c-ic \tan(e+fx))^{7/2}} - \frac{i(a+ia \tan(e+fx))^{7/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)$$

f

input `Int[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(11/2), x]`

output `(a*c*(-1/11*((I*A + B)*(a + I*a*Tan[e + f*x])^(7/2))/(a*c*(c - I*c*Tan[e + f*x])^(11/2)) + ((2*A + (9*I)*B)*(((-1/9*I)*(a + I*a*Tan[e + f*x])^(7/2))/(a*c*(c - I*c*Tan[e + f*x])^(9/2)) - ((I/63)*(a + I*a*Tan[e + f*x])^(7/2))/(a*c^2*(c - I*c*Tan[e + f*x])^(7/2))))/(11*c))/f`

3.826.3.1 Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.826.4 Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.77

method	result
risch	$-\frac{a^3 \sqrt{\frac{a e^{2i(fx+e)}}{e^{2i(fx+e)}+1}} (63iA e^{10i(fx+e)}+63B e^{10i(fx+e)}+154iA e^{8i(fx+e)}+99iA e^{6i(fx+e)}-99B e^{6i(fx+e)})}{2772c^5 \sqrt{\frac{c}{e^{2i(fx+e)}+1}} f}$
derivativedivides	$\frac{i \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^3 (1+\tan(fx+e)^2) (2iA \tan(fx+e)^4-63iB \tan(fx+e)^3-9B \tan(fx+e)^4)}{693f c^6 (i+\tan(fx+e))}$
default	$\frac{i \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^3 (1+\tan(fx+e)^2) (2iA \tan(fx+e)^4-63iB \tan(fx+e)^3-9B \tan(fx+e)^4)}{693f c^6 (i+\tan(fx+e))}$
parts	$\frac{iA \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^3 (1+\tan(fx+e)^2) (2i \tan(fx+e)^4-45i \tan(fx+e)^2-14 \tan(fx+e)^3+7)}{693f c^6 (i+\tan(fx+e))^7}$

```
input int((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(11/2), x,
method=_RETURNVERBOSE)
```

```
output -1/2772*a^3/c^5*(a*exp(2*I*(f*x+e))/(exp(2*I*(f*x+e))+1))^(1/2)/(c/(exp(2*
I*(f*x+e))+1))^(1/2)/f*(63*I*A*exp(10*I*(f*x+e))+63*B*exp(10*I*(f*x+e))+15
4*I*A*exp(8*I*(f*x+e))+99*I*A*exp(6*I*(f*x+e))-99*B*exp(6*I*(f*x+e)))
```

3.826. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{11/2}} dx$

3.826.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.81

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx =$$

$$\frac{(63(iA + B)a^3 e^{(13i fx + 13ie)} + 7(31iA + 9B)a^3 e^{(11i fx + 11ie)} + 11(23iA - 9B)a^3 e^{(9i fx + 9ie)} + 99(iA - B)a^3 e^{(7i fx + 7ie)}) \sqrt{a/(e^{(2i fx + 2ie)} + 1)} \sqrt{c/(e^{(2i fx + 2ie)} + 1)}}{2772 c^6 f}$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(11/2),x, algorithm="fricas")`

output `-1/2772*(63*(I*A + B)*a^3*e^(13*I*f*x + 13*I*e) + 7*(31*I*A + 9*B)*a^3*e^(11*I*f*x + 11*I*e) + 11*(23*I*A - 9*B)*a^3*e^(9*I*f*x + 9*I*e) + 99*(I*A - B)*a^3*e^(7*I*f*x + 7*I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))/(c^6*f)`

3.826.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(11/2),x)`

output `Timed out`

3.826.7 Maxima [A] (verification not implemented)

Time = 0.63 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.28

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx = \frac{(63(-iA - B)a^3 \cos(\frac{11}{2} \arctan(\sin(2fx + 2e)), \cos(\frac{11}{2} \arctan(\sin(2fx + 2e))))}{2772 c^6 f}$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(11/2),x, algorithm="maxima")`

output `1/2772*(63*(-I*A - B)*a^3*cos(11/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) - 154*I*A*a^3*cos(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 99*(-I*A + B)*a^3*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 63*(A - I*B)*a^3*sin(11/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 154*A*a^3*sin(9/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 99*(A + I*B)*a^3*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*sqrt(a)/(c^(11/2)*f)`

3.826.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{7/2}}{(-ictan(fx + e) + c)^{11/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(11/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(7/2)/(-I*c*tan(f*x + e) + c)^(11/2), x)`

3.826.9 Mupad [B] (verification not implemented)

Time = 10.82 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.40

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{11/2}} dx =$$

$$a^3 \sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}} (A \cos(6e + 6fx) 99i + A \cos(8e + 8fx) 154i + A \cos(10e + 10fx) 99i)$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2))/(c - c*tan(e + f*x)*1i)^(11/2),x)`

3.826. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{11/2}} dx$

output $-(a^3((a(\cos(2e + 2fx) + \sin(2e + 2fx)*1i + 1))/(\cos(2e + 2fx) + 1))^{1/2}(A\cos(6e + 6fx)*99i + A\cos(8e + 8fx)*154i + A\cos(10e + 10fx)*63i - 99B\cos(6e + 6fx) + 63B\cos(10e + 10fx) - 99A\sin(6e + 6fx) - 154A\sin(8e + 8fx) - 63A\sin(10e + 10fx) - B\sin(6e + 6fx)*99i + B\sin(10e + 10fx)*63i))/(2772*c^5*f*((c(\cos(2e + 2fx) - \sin(2e + 2fx)*1i + 1))/(\cos(2e + 2fx) + 1))^{1/2})$

3.826. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{11/2}} dx$

$$3.827 \quad \int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{13/2}} dx$$

3.827.1 Optimal result	7442
3.827.2 Mathematica [B] (verified)	7442
3.827.3 Rubi [A] (verified)	7443
3.827.4 Maple [A] (verified)	7446
3.827.5 Fricas [A] (verification not implemented)	7446
3.827.6 Sympy [F(-1)]	7447
3.827.7 Maxima [A] (verification not implemented)	7447
3.827.8 Giac [F]	7448
3.827.9 Mupad [B] (verification not implemented)	7448

3.827.1 Optimal result

Integrand size = 45, antiderivative size = 208

$$\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} dx =$$

$$\frac{(iA + B)(a + ia \tan(e + fx))^{7/2}}{13f(c - ic \tan(e + fx))^{13/2}} - \frac{(3iA - 10B)(a + ia \tan(e + fx))^{7/2}}{143cf(c - ic \tan(e + fx))^{11/2}}$$

$$- \frac{2(3iA - 10B)(a + ia \tan(e + fx))^{7/2}}{1287c^2f(c - ic \tan(e + fx))^{9/2}} - \frac{2(3iA - 10B)(a + ia \tan(e + fx))^{7/2}}{9009c^3f(c - ic \tan(e + fx))^{7/2}}$$

output
$$-1/13*(I*A+B)*(a+I*a*\tan(f*x+e))^(7/2)/f/(c-I*c*\tan(f*x+e))^(13/2)-1/143*(3*I*A-10*B)*(a+I*a*\tan(f*x+e))^(7/2)/c/f/(c-I*c*\tan(f*x+e))^(11/2)-2/1287*(3*I*A-10*B)*(a+I*a*\tan(f*x+e))^(7/2)/c^2/f/(c-I*c*\tan(f*x+e))^(9/2)-2/9009*(3*I*A-10*B)*(a+I*a*\tan(f*x+e))^(7/2)/c^3/f/(c-I*c*\tan(f*x+e))^(7/2)$$

3.827.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 495 vs. 2(208) = 416.

Time = 19.12 (sec) , antiderivative size = 495, normalized size of antiderivative = 2.38

$$\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} dx = \frac{\cos^4(e + fx) \left((-iA + B) \cos(6fx) \left(\frac{\cos(3e)}{112c^7} + \frac{i \sin(3e)}{112c^7} \right) \right)}{1}$$

3.827.
$$\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{13/2}} dx$$

input `Integrate[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(13/2),x]`

output `(Cos[e + f*x]^4*((-I)*A + B)*Cos[6*f*x]*(Cos[3*e]/(112*c^7) + ((I/112)*Sin[3*e])/c^7) + ((-15*I)*A + 8*B)*Cos[8*f*x]*(Cos[5*e]/(504*c^7) + ((I/504)*Sin[5*e])/c^7) + ((-30*I)*A + B)*Cos[10*f*x]*(Cos[7*e]/(792*c^7) + ((I/792)*Sin[7*e])/c^7) + (25*A - (12*I)*B)*Cos[12*f*x]*(((-1/1144*I)*Cos[9*e])/c^7 + Sin[9*e]/(1144*c^7)) + (A - I*B)*Cos[14*f*x]*(((-1/208*I)*Cos[11*e])/c^7 + Sin[11*e]/(208*c^7)) + (A + I*B)*(Cos[3*e]/(112*c^7) + ((I/112)*Sin[3*e])/c^7)*Sin[6*f*x] + (15*A + (8*I)*B)*(Cos[5*e]/(504*c^7) + ((I/504)*Sin[5*e])/c^7)*Sin[8*f*x] + (30*A + I*B)*(Cos[7*e]/(792*c^7) + ((I/792)*Sin[7*e])/c^7)*Sin[10*f*x] + (25*A - (12*I)*B)*(Cos[9*e]/(1144*c^7) + ((I/1144)*Sin[9*e])/c^7)*Sin[12*f*x] + (A - I*B)*(Cos[11*e]/(208*c^7) + ((I/208)*Sin[11*e])/c^7)*Sin[14*f*x])*Sqrt[Sec[e + f*x]*(c*Cos[e + f*x] - I*c*Sin[e + f*x])]*(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(f*(Cos[f*x] + I*Sin[f*x])^3*(A*Cos[e + f*x] + B*Sin[e + f*x]))`

3.827.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 87, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{13/2}} dx$$

↓ 3042

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{13/2}} dx$$

↓ 4071

$$\frac{ac \int \frac{(i \tan(e+fx)a+a)^{5/2} (A+B \tan(e+fx))}{(c-ictan(e+fx))^{15/2}} d \tan(e + fx)}{f}$$

↓ 87

3.827. $\int \frac{(a+ia \tan(e+fx))^{7/2} (A+B \tan(e+fx))}{(c-ictan(e+fx))^{13/2}} dx$

$$\begin{aligned}
 & ac \left(\frac{(3A+10iB) \int \frac{(i \tan(e+fx)a+a)^{5/2}}{(c-ic \tan(e+fx))^{13/2}} d \tan(e+fx)}{13c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{13ac(c-ic \tan(e+fx))^{13/2}} \right) \\
 & \quad \downarrow f \quad 55 \\
 & ac \left(\frac{(3A+10iB) \left(\frac{2 \int \frac{(i \tan(e+fx)a+a)^{5/2}}{(c-ic \tan(e+fx))^{11/2}} d \tan(e+fx)}{11c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{13c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{13ac(c-ic \tan(e+fx))^{13/2}} \right) \\
 & \quad \downarrow f \quad 55 \\
 & ac \left(\frac{(3A+10iB) \left(\frac{2 \left(\frac{\int \frac{(i \tan(e+fx)a+a)^{5/2}}{(c-ic \tan(e+fx))^{9/2}} d \tan(e+fx)}{9c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{13c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{13ac(c-ic \tan(e+fx))^{13/2}} \right) \\
 & \quad \downarrow f \quad 48 \\
 & ac \left(\frac{(3A+10iB) \left(\frac{2 \left(-\frac{i(a+ia \tan(e+fx))^{7/2}}{63ac^2(c-ic \tan(e+fx))^{7/2}} - \frac{i(a+ia \tan(e+fx))^{7/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{13c} - \frac{(B+iA)(a+ia \tan(e+fx))^{7/2}}{13ac(c-ic \tan(e+fx))^{13/2}} \right) \\
 & \quad \downarrow f
 \end{aligned}$$

input `Int[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(13/2), x]`

3.827. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{13/2}} dx$

output $(a*c*(-1/13*((I*A + B)*(a + I*a*Tan[e + f*x])^(7/2))/(a*c*(c - I*c*Tan[e + f*x])^(13/2)) + ((3*A + (10*I)*B)*((-1/11*I)*(a + I*a*Tan[e + f*x])^(7/2)))/(a*c*(c - I*c*Tan[e + f*x])^(11/2)) + (2*((-1/9*I)*(a + I*a*Tan[e + f*x])^(7/2))/(a*c*(c - I*c*Tan[e + f*x])^(9/2)) - ((I/63)*(a + I*a*Tan[e + f*x])^(7/2))/(a*c^2*(c - I*c*Tan[e + f*x])^(7/2)))/(11*c))/(13*c))/f$

3.827.3.1 Defintions of rubi rules used

rule 48 $\text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{m+1} * (c + d*x)^{n+1} / ((b*c - a*d) * (m+1)), x] /;$ $\text{FreeQ}\{a, b, c, d, m, n\}, x\} \ \&\& \ \text{EqQ}[m + n + 2, 0] \ \&\& \ \text{NeQ}[m, -1]$

rule 55 $\text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{m+1} * (c + d*x)^{n+1} / ((b*c - a*d) * (m+1)), x] - \text{Simp}[d * (\text{Simplify}[m + n + 2] / ((b*c - a*d) * (m+1))) \text{Int}[(a + b*x)^{\text{Simplify}[m+1]} * (c + d*x)^n, x], x] /;$ $\text{FreeQ}\{a, b, c, d, m, n\}, x\} \ \&\& \ \text{ILtQ}[\text{Simplify}[m + n + 2], 0] \ \&\& \ \text{NeQ}[m, -1] \ \&\& \ !(\text{LtQ}[m, -1] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ (\text{EqQ}[a, 0] \ || \ (\text{NeQ}[c, 0] \ \&\& \ \text{LtQ}[m - n, 0] \ \&\& \ \text{IntegerQ}[n]))) \ \&\& \ (\text{SumSimplerQ}[m, 1] \ || \ !\text{SumSimplerQ}[n, 1])$

rule 87 $\text{Int}[(a + b*x)^m * (c + d*x)^n * (e + f*x)^p, x] \rightarrow \text{Simp}[(-b*e - a*f) * (c + d*x)^{n+1} * (e + f*x)^{p+1} / (f * (p+1) * (c*f - d*e)), x] - \text{Simp}[(a*d*f * (n + p + 2) - b * (d*e * (n + 1) + c*f * (p + 1))) / (f * (p + 1) * (c*f - d*e)) \text{Int}[(c + d*x)^n * (e + f*x)^{p+1}, x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, n\}, x\} \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ (!\text{LtQ}[n, -1] \ || \ \text{IntegerQ}[p] \ || \ !(\text{IntegerQ}[n] \ || \ !(\text{EqQ}[e, 0] \ || \ !(\text{EqQ}[c, 0] \ || \ \text{LtQ}[p, n])))$

rule 3042 $\text{Int}[u, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /;$ $\text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4071 $\text{Int}[(a + b*x)^m * (c + d*x)^n * (A + B*x)^p * \tan[e + f*x], x] \rightarrow \text{Simp}[a * (c/f) \text{Subst}[\text{Int}[(a + b*x)^{m-1} * (c + d*x)^{n-1} * (A + B*x)^p, x], x, \text{Tan}[e + f*x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, A, B, m, n\}, x\} \ \&\& \ \text{EqQ}[b*c + a*d, 0] \ \&\& \ \text{EqQ}[a^2 + b^2, 0]$

3.827.4 Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.75

method	result
risch	$\frac{a^3 \sqrt{\frac{a e^{2i(fx+e)}}{e^{2i(fx+e)}+1}} (693iA e^{12i(fx+e)}+693B e^{12i(fx+e)}+2457iA e^{10i(fx+e)}+819B e^{10i(fx+e)}+3003iA e^{8i(fx+e)}-1001B e^{8i(fx+e)}+1287iA e^{6i(fx+e)}-1287B e^{6i(fx+e)})}{72072c^6 \sqrt{\frac{c}{e^{2i(fx+e)}+1}} f}$
derivativedivides	$\frac{i\sqrt{a(1+i\tan(fx+e))} \sqrt{-c(i\tan(fx+e)-1)} a^3 (1+\tan(fx+e))^2 (6iA \tan(fx+e)^5 - 160iB \tan(fx+e)^4 - 20B \tan(fx+e)^3 - 160iA \tan(fx+e)^2 + 160iB \tan(fx+e) - 160A)}{72072c^6 \sqrt{\frac{c}{e^{2i(fx+e)}+1}} f}$
default	$\frac{i\sqrt{a(1+i\tan(fx+e))} \sqrt{-c(i\tan(fx+e)-1)} a^3 (1+\tan(fx+e))^2 (6iA \tan(fx+e)^5 - 160iB \tan(fx+e)^4 - 20B \tan(fx+e)^3 - 160iA \tan(fx+e)^2 + 160iB \tan(fx+e) - 160A)}{72072c^6 \sqrt{\frac{c}{e^{2i(fx+e)}+1}} f}$
parts	$\frac{iA \sqrt{a(1+i\tan(fx+e))} \sqrt{-c(i\tan(fx+e)-1)} a^3 (1+\tan(fx+e))^2 (2i \tan(fx+e)^5 - 59i \tan(fx+e)^3 - 16 \tan(fx+e)^4 - 5 \tan(fx+e)^2 + 5 \tan(fx+e) - 5)}{3003f c^7 (i+\tan(fx+e))^8}$

input `int((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(13/2),x,method=_RETURNVERBOSE)`

output `-1/72072*a^3/c^6*(a*exp(2*I*(f*x+e))/(exp(2*I*(f*x+e))+1))^(1/2)/(c/(exp(2*I*(f*x+e))+1))^(1/2)/f*(693*I*A*exp(12*I*(f*x+e))+693*B*exp(12*I*(f*x+e))+2457*I*A*exp(10*I*(f*x+e))+819*B*exp(10*I*(f*x+e))+3003*I*A*exp(8*I*(f*x+e))-1001*B*exp(8*I*(f*x+e))+1287*I*A*exp(6*I*(f*x+e))-1287*B*exp(6*I*(f*x+e)))`

3.827.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.70

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} dx = \frac{(693(iA + B)a^3 e^{(15i fx + 15ie)} + 126(25iA + 12B)a^3 e^{(13i fx + 13ie)} + 182(30iA - B)a^3 e^{(11i fx + 11ie)} + 286(10iA - 6B)a^3 e^{(9i fx + 9ie)} + 1287iA e^{(7i fx + 7ie)} - 1287B e^{(7i fx + 7ie)})}{72072 c^7 f}$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(13/2),x, algorithm="fricas")`

output $-1/72072*(693*(I*A + B)*a^3*e^{(15*I*f*x + 15*I*e)} + 126*(25*I*A + 12*B)*a^3*e^{(13*I*f*x + 13*I*e)} + 182*(30*I*A - B)*a^3*e^{(11*I*f*x + 11*I*e)} + 286*(15*I*A - 8*B)*a^3*e^{(9*I*f*x + 9*I*e)} + 1287*(I*A - B)*a^3*e^{(7*I*f*x + 7*I*e)})*sqrt(a/(e^{(2*I*f*x + 2*I*e)} + 1))*sqrt(c/(e^{(2*I*f*x + 2*I*e)} + 1))/(c^7*f)$

3.827.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(13/2),x)`

output Timed out

3.827.7 Maxima [A] (verification not implemented)

Time = 0.67 (sec) , antiderivative size = 276, normalized size of antiderivative = 1.33

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{13/2}} dx = \frac{(693(-iA - B)a^3 \cos(\frac{13}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 819(-3IA - B)a^3 \cos(11/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 1001(-3IA + B)a^3 \cos(9/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 1287(-IA + B)a^3 \cos(7/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 693(A - IB)a^3 \sin(13/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 819(3A - IB)a^3 \sin(11/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 1001(3A + IB)a^3 \sin(9/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 1287(A + IB)a^3 \sin(7/2 \arctan(\sin(2fx + 2e)), \cos(2fx + 2e))) * sqrt(a) / (c^{13/2} * f)}$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(13/2),x, algorithm="maxima")`

output $1/72072*(693*(-I*A - B)*a^3*\cos(13/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 819*(-3*I*A - B)*a^3*\cos(11/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 1001*(-3*I*A + B)*a^3*\cos(9/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 1287*(-I*A + B)*a^3*\cos(7/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 693*(A - I*B)*a^3*\sin(13/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 819*(3*A - I*B)*a^3*\sin(11/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 1001*(3*A + I*B)*a^3*\sin(9/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 1287*(A + I*B)*a^3*\sin(7/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))))*sqrt(a)/(c^(13/2)*f)$

3.827. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{13/2}} dx$

3.827.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{13/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{7/2}}{(-ictan(fx + e) + c)^{13/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(13/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(7/2)/(-I*c*tan(f*x + e) + c)^(13/2), x)`

3.827.9 Mupad [B] (verification not implemented)

Time = 13.01 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.80

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{13/2}} dx = \frac{\sqrt{a + \frac{a \sin(e+fx) \operatorname{li}}{\cos(e+fx)}} \left(\frac{a^3 e^{e 8i + f x 8i} (3A+B \operatorname{li}) \operatorname{li}}{72 c^6 f} + \frac{a^3 e^{e 10i + f x 10i} (3A-B \operatorname{li}) \operatorname{li}}{88 c^6 f} + \frac{a^3 e^{e 6i + f x 6i} (A+B \operatorname{li}) \operatorname{li}}{56 c^6 f} + \frac{a^3 e^{e 12i + f x 12i} (A-B \operatorname{li}) \operatorname{li}}{104 c^6 f} \right)}{\sqrt{c - \frac{c \sin(e+fx) \operatorname{li}}{\cos(e+fx)}}}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2))/(c - c*tan(e + f*x)*1i)^(13/2),x)`

output `-((a + (a*sin(e + f*x)*1i)/cos(e + f*x))^(1/2))*((a^3*exp(e*8i + f*x*8i))*(3*A + B*1i)*1i)/(72*c^6*f) + (a^3*exp(e*10i + f*x*10i)*(3*A - B*1i)*1i)/(88*c^6*f) + (a^3*exp(e*6i + f*x*6i)*(A + B*1i)*1i)/(56*c^6*f) + (a^3*exp(e*12i + f*x*12i)*(A - B*1i)*1i)/(104*c^6*f))/((c - (c*sin(e + f*x)*1i)/cos(e + f*x))^(1/2))`

3.828
$$\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{15/2}} dx$$

3.828.1 Optimal result	7449
3.828.2 Mathematica [B] (verified)	7450
3.828.3 Rubi [A] (verified)	7450
3.828.4 Maple [A] (verified)	7454
3.828.5 Fricas [A] (verification not implemented)	7454
3.828.6 Sympy [F(-1)]	7455
3.828.7 Maxima [A] (verification not implemented)	7455
3.828.8 Giac [F]	7456
3.828.9 Mupad [B] (verification not implemented)	7456

3.828.1 Optimal result

Integrand size = 45, antiderivative size = 261

$$\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{15/2}} dx =$$

$$\frac{(iA + B)(a + ia \tan(e + fx))^{7/2}}{15f(c - ic \tan(e + fx))^{15/2}} - \frac{(4iA - 11B)(a + ia \tan(e + fx))^{7/2}}{195cf(c - ic \tan(e + fx))^{13/2}}$$

$$- \frac{(4iA - 11B)(a + ia \tan(e + fx))^{7/2}}{715c^2f(c - ic \tan(e + fx))^{11/2}} - \frac{2(4iA - 11B)(a + ia \tan(e + fx))^{7/2}}{6435c^3f(c - ic \tan(e + fx))^{9/2}}$$

$$- \frac{2(4iA - 11B)(a + ia \tan(e + fx))^{7/2}}{45045c^4f(c - ic \tan(e + fx))^{7/2}}$$

output

```
-1/15*(I*A+B)*(a+I*a*tan(f*x+e))^(7/2)/f/(c-I*c*tan(f*x+e))^(15/2)-1/195*(
4*I*A-11*B)*(a+I*a*tan(f*x+e))^(7/2)/c/f/(c-I*c*tan(f*x+e))^(13/2)-1/715*(
4*I*A-11*B)*(a+I*a*tan(f*x+e))^(7/2)/c^2/f/(c-I*c*tan(f*x+e))^(11/2)-2/643
5*(4*I*A-11*B)*(a+I*a*tan(f*x+e))^(7/2)/c^3/f/(c-I*c*tan(f*x+e))^(9/2)-2/4
5045*(4*I*A-11*B)*(a+I*a*tan(f*x+e))^(7/2)/c^4/f/(c-I*c*tan(f*x+e))^(7/2)
```


3.828.2 Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 577 vs. $2(261) = 522$.

Time = 19.38 (sec) , antiderivative size = 577, normalized size of antiderivative = 2.21

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{15/2}} dx = \frac{\cos^4(e + fx) \left((-iA + B) \cos(6fx) \left(\frac{\cos(3e)}{224c^8} + \frac{i \sin(3e)}{224c^8} \right) \right)}{}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(15/2),x]`

output `(Cos[e + f*x]^4*(((I)*A + B)*Cos[6*f*x]*(Cos[3*e]/(224*c^8) + ((I/224)*Sin[3*e])/c^8) + ((-37*I)*A + 23*B)*Cos[8*f*x]*(Cos[5*e]/(2016*c^8) + ((I/2016)*Sin[5*e])/c^8) + ((-49*I)*A + 11*B)*Cos[10*f*x]*(Cos[7*e]/(1584*c^8) + ((I/1584)*Sin[7*e])/c^8) + (61*A - (11*I)*B)*Cos[12*f*x]*(((I/2288)*Cos[9*e])/c^8 + Sin[9*e]/(2288*c^8)) + (73*A - (43*I)*B)*Cos[14*f*x]*(((I/6240)*Cos[11*e])/c^8 + Sin[11*e]/(6240*c^8)) + (A - I*B)*Cos[16*f*x]*(((I/480)*Cos[13*e])/c^8 + Sin[13*e]/(480*c^8)) + (A + I*B)*(Cos[3*e]/(224*c^8) + ((I/224)*Sin[3*e])/c^8)*Sin[6*f*x] + (37*A + (23*I)*B)*(Cos[5*e]/(2016*c^8) + ((I/2016)*Sin[5*e])/c^8)*Sin[8*f*x] + (49*A + (11*I)*B)*(Cos[7*e]/(1584*c^8) + ((I/1584)*Sin[7*e])/c^8)*Sin[10*f*x] + (61*A - (11*I)*B)*(Cos[9*e]/(2288*c^8) + ((I/2288)*Sin[9*e])/c^8)*Sin[12*f*x] + (73*A - (43*I)*B)*(Cos[11*e]/(6240*c^8) + ((I/6240)*Sin[11*e])/c^8)*Sin[14*f*x] + (A - I*B)*(Cos[13*e]/(480*c^8) + ((I/480)*Sin[13*e])/c^8)*Sin[16*f*x])*Sqrt[Sec[e + f*x]*(c*cos[e + f*x] - I*c*sin[e + f*x])]*(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(f*(Cos[f*x] + I*SIN[f*x])^3*(A*cos[e + f*x] + B*Sin[e + f*x]))`

3.828.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 87, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.828. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{15/2}} dx$

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{15/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{15/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{(i \tan(e + fx)a + a)^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{17/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{87} \\
 & \frac{ac \left(\frac{(4A + 11iB) \int \frac{(i \tan(e + fx)a + a)^{5/2}}{(c - ic \tan(e + fx))^{15/2}} d \tan(e + fx)}{15c} - \frac{(B + iA)(a + ia \tan(e + fx))^{7/2}}{15ac(c - ic \tan(e + fx))^{15/2}} \right)}{f} \\
 & \quad \downarrow \text{55} \\
 & \frac{ac \left(\frac{(4A + 11iB) \left(\frac{3 \int \frac{(i \tan(e + fx)a + a)^{5/2}}{(c - ic \tan(e + fx))^{13/2}} d \tan(e + fx)}{13c} - \frac{i(a + ia \tan(e + fx))^{7/2}}{13ac(c - ic \tan(e + fx))^{13/2}} \right)}{15c} - \frac{(B + iA)(a + ia \tan(e + fx))^{7/2}}{15ac(c - ic \tan(e + fx))^{15/2}} \right)}{f} \\
 & \quad \downarrow \text{55} \\
 & \frac{ac \left(\frac{(4A + 11iB) \left(\frac{3 \left(\frac{2 \int \frac{(i \tan(e + fx)a + a)^{5/2}}{(c - ic \tan(e + fx))^{11/2}} d \tan(e + fx)}{11c} - \frac{i(a + ia \tan(e + fx))^{7/2}}{11ac(c - ic \tan(e + fx))^{11/2}} \right)}{13c} - \frac{i(a + ia \tan(e + fx))^{7/2}}{13ac(c - ic \tan(e + fx))^{13/2}} \right)}{15c} - \frac{(B + iA)(a + ia \tan(e + fx))^{7/2}}{15ac(c - ic \tan(e + fx))^{15/2}} \right)}{f} \\
 & \quad \downarrow \text{55}
 \end{aligned}$$

3.828. $\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{15/2}} dx$

$$ac \left(\frac{(4A+11iB) \left(\frac{2 \left(\frac{\int \frac{(i \tan(e+fx)a+a)^{5/2}}{(c-ic \tan(e+fx))^{9/2}} d \tan(e+fx)}{9c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{13c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{13ac(c-ic \tan(e+fx))^{13/2}} \right)}{15c} \right) - \frac{(B+iA)}{15ac}$$

f

↓ 48

$$ac \left(\frac{(4A+11iB) \left(\frac{2 \left(-\frac{i(a+ia \tan(e+fx))^{7/2}}{63ac^2(c-ic \tan(e+fx))^{7/2}} - \frac{i(a+ia \tan(e+fx))^{7/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{13c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{13ac(c-ic \tan(e+fx))^{13/2}} \right)}{15c} \right) - \frac{(B+iA)}{15ac}$$

f

```
input Int[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(15/2), x]
```

```
output (a*c*(-1/15*((I*A + B)*(a + I*a*Tan[e + f*x])^(7/2))/(a*c*(c - I*c*Tan[e +
f*x])^(15/2)) + ((4*A + (11*I)*B)*((-1/13*I)*(a + I*a*Tan[e + f*x])^(7/2
)))/(a*c*(c - I*c*Tan[e + f*x])^(13/2)) + (3*((-1/11*I)*(a + I*a*Tan[e + f
*x])^(7/2))/(a*c*(c - I*c*Tan[e + f*x])^(11/2)) + (2*((-1/9*I)*(a + I*a*T
an[e + f*x])^(7/2))/(a*c*(c - I*c*Tan[e + f*x])^(9/2)) - ((I/63)*(a + I*a*
Tan[e + f*x])^(7/2))/(a*c^2*(c - I*c*Tan[e + f*x])^(7/2))))/(11*c)))/(13*c
)))/(15*c))/f
```

3.828.3.1 Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 55 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.828.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.65

method	result
risch	$a^3 \frac{\sqrt{\frac{a e^{2i(fx+e)}}{e^{2i(fx+e)}+1}} (3003iA e^{14i(fx+e)}+3003B e^{14i(fx+e)}+13860iA e^{12i(fx+e)}+6930B e^{12i(fx+e)}+24570iA e^{10i(fx+e)}+20020iA e^{8i(fx+e)}-10010B e^{8i(fx+e)}+6435iA e^{6i(fx+e)}-6435B e^{6i(fx+e)})}{720720c^7 \sqrt{\frac{c}{e^{2i(fx+e)}+1}} f}$
derivativedivides	$-\frac{\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^3 (1+\tan(fx+e)^2) (22iB \tan(fx+e)^6+72iA \tan(fx+e)^5+8A \tan(fx+e)^4)}{\dots}$
default	$-\frac{\sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^3 (1+\tan(fx+e)^2) (22iB \tan(fx+e)^6+72iA \tan(fx+e)^5+8A \tan(fx+e)^4)}{\dots}$
parts	$\frac{iA \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} a^3 (1+\tan(fx+e)^2) (-4243i+6858 \tan(fx+e)+1455i \tan(fx+e)^2+780 \tan(fx+e)^3)}{45045 f c^8 (i+\tan(fx+e))^9}$

```
input int((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(15/2),x,
method=_RETURNVERBOSE)
```

```
output -1/720720*a^3/c^7*(a*exp(2*I*(f*x+e))/(exp(2*I*(f*x+e))+1))^(1/2)/(c/(exp(
2*I*(f*x+e))+1))^(1/2)/f*(3003*I*A*exp(14*I*(f*x+e))+3003*B*exp(14*I*(f*x+
e))+13860*I*A*exp(12*I*(f*x+e))+6930*B*exp(12*I*(f*x+e))+24570*I*A*exp(10*
I*(f*x+e))+20020*I*A*exp(8*I*(f*x+e))-10010*B*exp(8*I*(f*x+e))+6435*I*A*ex
p(6*I*(f*x+e))-6435*B*exp(6*I*(f*x+e)))
```

3.828.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.64

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{15/2}} dx =$$

$$\frac{(3003 (i A + B) a^3 e^{(17i fx+17i e)} + 231 (73i A + 43 B) a^3 e^{(15i fx+15i e)} + 630 (61i A + 11 B) a^3 e^{(13i fx+13i e)} + 9 \dots)}{\dots}$$

```
input integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(15
/2),x, algorithm="fricas")
```

output $-1/720720*(3003*(I*A + B)*a^3*e^{(17*I*f*x + 17*I*e)} + 231*(73*I*A + 43*B)*a^3*e^{(15*I*f*x + 15*I*e)} + 630*(61*I*A + 11*B)*a^3*e^{(13*I*f*x + 13*I*e)} + 910*(49*I*A - 11*B)*a^3*e^{(11*I*f*x + 11*I*e)} + 715*(37*I*A - 23*B)*a^3*e^{(9*I*f*x + 9*I*e)} + 6435*(I*A - B)*a^3*e^{(7*I*f*x + 7*I*e)})*sqrt(a/(e^{(2*I*f*x + 2*I*e)} + 1))*sqrt(c/(e^{(2*I*f*x + 2*I*e)} + 1))/(c^8*f)$

3.828.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{15/2}} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(15/2),x)`

output Timed out

3.828.7 Maxima [A] (verification not implemented)

Time = 1.09 (sec) , antiderivative size = 332, normalized size of antiderivative = 1.27

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{15/2}} dx = \frac{(3003(-iA - B)a^3 \cos(\frac{15}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 6930(-2iA - B)a^3 \cos(\frac{13}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) - 24570iAa^3 \cos(\frac{11}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 10010(-2iA + B)a^3 \cos(\frac{9}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 6435(-iA + B)a^3 \cos(\frac{7}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 3003(A - iB)a^3 \sin(\frac{15}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 6930(2A - iB)a^3 \sin(\frac{13}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 24570Aa^3 \sin(\frac{11}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 10010(2A + iB)a^3 \sin(\frac{9}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)) + 6435(A + iB)a^3 \sin(\frac{7}{2} \arctan(\sin(2fx + 2e)), \cos(2fx + 2e)))*sqrt(a)/(c^{(15/2)*f})$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(15/2),x, algorithm="maxima")`

output $1/720720*(3003*(-I*A - B)*a^3*\cos(15/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 6930*(-2*I*A - B)*a^3*\cos(13/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) - 24570*I*A*a^3*\cos(11/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 10010*(-2*I*A + B)*a^3*\cos(9/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 6435*(-I*A + B)*a^3*\cos(7/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 3003*(A - I*B)*a^3*\sin(15/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 6930*(2*A - I*B)*a^3*\sin(13/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 24570*A*a^3*\sin(11/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 10010*(2*A + I*B)*a^3*\sin(9/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 6435*(A + I*B)*a^3*\sin(7/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))))*sqrt(a)/(c^{(15/2)*f})$

3.828. $\int \frac{(a+ia \tan(e+fx))^{7/2} (A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{15/2}} dx$

3.828.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{15/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{7/2}}{(-ictan(fx + e) + c)^{15/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(15/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(7/2)/(-I*c*tan(f*x + e) + c)^(15/2), x)`

3.828.9 Mupad [B] (verification not implemented)

Time = 12.60 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.73

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{15/2}} dx = \frac{\sqrt{a + \frac{a \sin(e+fx) \operatorname{li}}{\cos(e+fx)}} \left(\frac{a^3 e^{8i+fx} 8i (2A+B \operatorname{li}) \operatorname{li}}{72 c^7 f} + \frac{a^3 e^{12i+fx} 12i (2A-B \operatorname{li}) \operatorname{li}}{104 c^7 f} + \frac{A a^3 e^{10i+fx} 10i 3i}{88 c^7 f} + \frac{a^3 e^{6i+fx} 6i (A+B \operatorname{li}) \operatorname{li}}{112 c^7 f} \right)}{\sqrt{c - \frac{c \sin(e+fx) \operatorname{li}}{\cos(e+fx)}}}$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2))/(c - c*tan(e + f*x)*1i)^(15/2),x)`

output `-((a + (a*sin(e + f*x)*1i)/cos(e + f*x))^(1/2))*((a^3*exp(e*8i + f*x*8i))*(2*A + B*1i)*1i)/(72*c^7*f) + (a^3*exp(e*12i + f*x*12i)*(2*A - B*1i)*1i)/(104*c^7*f) + (A*a^3*exp(e*10i + f*x*10i)*3i)/(88*c^7*f) + (a^3*exp(e*6i + f*x*6i)*(A + B*1i)*1i)/(112*c^7*f) + (a^3*exp(e*14i + f*x*14i)*(A - B*1i)*1i)/(240*c^7*f))/((c - (c*sin(e + f*x)*1i)/cos(e + f*x))^(1/2))`

3.829
$$\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{17/2}} dx$$

3.829.1 Optimal result 7457
 3.829.2 Mathematica [B] (warning: unable to verify) 7458
 3.829.3 Rubi [A] (verified) 7458
 3.829.4 Maple [A] (verified) 7464
 3.829.5 Fricas [A] (verification not implemented) 7464
 3.829.6 Sympy [F(-1)] 7465
 3.829.7 Maxima [A] (verification not implemented) 7465
 3.829.8 Giac [F] 7466
 3.829.9 Mupad [B] (verification not implemented) 7466

3.829.1 Optimal result

Integrand size = 45, antiderivative size = 314

$$\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{17/2}} dx =$$

$$\frac{(iA + B)(a + ia \tan(e + fx))^{7/2}}{17f(c - ic \tan(e + fx))^{17/2}} - \frac{(5iA - 12B)(a + ia \tan(e + fx))^{7/2}}{255cf(c - ic \tan(e + fx))^{15/2}}$$

$$- \frac{4(5iA - 12B)(a + ia \tan(e + fx))^{7/2}}{3315c^2f(c - ic \tan(e + fx))^{13/2}} - \frac{4(5iA - 12B)(a + ia \tan(e + fx))^{7/2}}{12155c^3f(c - ic \tan(e + fx))^{11/2}}$$

$$- \frac{8(5iA - 12B)(a + ia \tan(e + fx))^{7/2}}{109395c^4f(c - ic \tan(e + fx))^{9/2}} - \frac{8(5iA - 12B)(a + ia \tan(e + fx))^{7/2}}{765765c^5f(c - ic \tan(e + fx))^{7/2}}$$

output

```
-1/17*(I*A+B)*(a+I*a*tan(f*x+e))^(7/2)/f/(c-I*c*tan(f*x+e))^(17/2)-1/255*(
5*I*A-12*B)*(a+I*a*tan(f*x+e))^(7/2)/c/f/(c-I*c*tan(f*x+e))^(15/2)-4/3315*
(5*I*A-12*B)*(a+I*a*tan(f*x+e))^(7/2)/c^2/f/(c-I*c*tan(f*x+e))^(13/2)-4/12
155*(5*I*A-12*B)*(a+I*a*tan(f*x+e))^(7/2)/c^3/f/(c-I*c*tan(f*x+e))^(11/2)-
8/109395*(5*I*A-12*B)*(a+I*a*tan(f*x+e))^(7/2)/c^4/f/(c-I*c*tan(f*x+e))^(9
/2)-8/765765*(5*I*A-12*B)*(a+I*a*tan(f*x+e))^(7/2)/c^5/f/(c-I*c*tan(f*x+e)
)^(7/2)
```


3.829.2 Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 655 vs. $2(314) = 628$.

Time = 19.83 (sec) , antiderivative size = 655, normalized size of antiderivative = 2.09

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ict \tan(e + fx))^{17/2}} dx = \frac{\cos^4(e + fx) \left((-iA + B) \cos(6fx) \left(\frac{\cos(3e)}{448c^9} + \frac{i \sin(3e)}{448c^9} \right) \right)}{}$$

input `Integrate[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(17/2),x]`

output `(Cos[e + f*x]^4*(((-I)*A + B)*Cos[6*f*x]*(Cos[3*e]/(448*c^9) + ((I/448)*Sin[3*e])/c^9) + ((-22*I)*A + 15*B)*Cos[8*f*x]*(Cos[5*e]/(2016*c^9) + ((I/2016)*Sin[5*e])/c^9) + ((-145*I)*A + 51*B)*Cos[10*f*x]*(Cos[7*e]/(6336*c^9) + ((I/6336)*Sin[7*e])/c^9) + ((-60*I)*A + B)*Cos[12*f*x]*(Cos[9*e]/(2288*c^9) + ((I/2288)*Sin[9*e])/c^9) + (215*A - (69*I)*B)*Cos[14*f*x]*(((-1/12480*I)*Cos[11*e])/c^9 + Sin[11*e]/(12480*c^9)) + (50*A - (33*I)*B)*Cos[16*f*x]*(((-1/8160*I)*Cos[13*e])/c^9 + Sin[13*e]/(8160*c^9)) + (A - I*B)*Cos[18*f*x]*(((-1/1088*I)*Cos[15*e])/c^9 + Sin[15*e]/(1088*c^9)) + (A + I*B)*(Cos[3*e]/(448*c^9) + ((I/448)*Sin[3*e])/c^9)*Sin[6*f*x] + (22*A + (15*I)*B)*(Cos[5*e]/(2016*c^9) + ((I/2016)*Sin[5*e])/c^9)*Sin[8*f*x] + (145*A + (51*I)*B)*(Cos[7*e]/(6336*c^9) + ((I/6336)*Sin[7*e])/c^9)*Sin[10*f*x] + (60*A + I*B)*(Cos[9*e]/(2288*c^9) + ((I/2288)*Sin[9*e])/c^9)*Sin[12*f*x] + (215*A - (69*I)*B)*(Cos[11*e]/(12480*c^9) + ((I/12480)*Sin[11*e])/c^9)*Sin[14*f*x] + (50*A - (33*I)*B)*(Cos[13*e]/(8160*c^9) + ((I/8160)*Sin[13*e])/c^9)*Sin[16*f*x] + (A - I*B)*(Cos[15*e]/(1088*c^9) + ((I/1088)*Sin[15*e])/c^9)*Sin[18*f*x]*Sqrt[Sec[e + f*x]*(c*cos[e + f*x] - I*c*sin[e + f*x])]*(a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(f*(Cos[f*x] + I*SIN[f*x])^3*(A*cos[e + f*x] + B*sin[e + f*x]))`

3.829.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 329, normalized size of antiderivative = 1.05, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.178$, Rules used = {3042, 4071, 87, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

3.829. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ict \tan(e+fx))^{17/2}} dx$

$$\begin{aligned}
 & \int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{17/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{17/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{(i \tan(e + fx)a + a)^{5/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{19/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{87} \\
 & \frac{ac \left(\frac{(5A + 12iB) \int \frac{(i \tan(e + fx)a + a)^{5/2}}{(c - ic \tan(e + fx))^{17/2}} d \tan(e + fx)}{17c} - \frac{(B + iA)(a + ia \tan(e + fx))^{7/2}}{17ac(c - ic \tan(e + fx))^{17/2}} \right)}{f} \\
 & \quad \downarrow \text{55} \\
 & \frac{ac \left(\frac{(5A + 12iB) \left(\frac{4 \int \frac{(i \tan(e + fx)a + a)^{5/2}}{(c - ic \tan(e + fx))^{15/2}} d \tan(e + fx)}{15c} - \frac{i(a + ia \tan(e + fx))^{7/2}}{15ac(c - ic \tan(e + fx))^{15/2}} \right)}{17c} - \frac{(B + iA)(a + ia \tan(e + fx))^{7/2}}{17ac(c - ic \tan(e + fx))^{17/2}} \right)}{f} \\
 & \quad \downarrow \text{55} \\
 & \frac{ac \left(\frac{(5A + 12iB) \left(\frac{4 \left(\frac{3 \int \frac{(i \tan(e + fx)a + a)^{5/2}}{(c - ic \tan(e + fx))^{13/2}} d \tan(e + fx)}{13c} - \frac{i(a + ia \tan(e + fx))^{7/2}}{13ac(c - ic \tan(e + fx))^{13/2}} \right)}{15c} - \frac{i(a + ia \tan(e + fx))^{7/2}}{15ac(c - ic \tan(e + fx))^{15/2}} \right)}{17c} - \frac{(B + iA)(a + ia \tan(e + fx))^{7/2}}{17ac(c - ic \tan(e + fx))^{17/2}} \right)}{f} \\
 & \quad \downarrow \text{55}
 \end{aligned}$$

3.829. $\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ic \tan(e + fx))^{17/2}} dx$

$$\left(\frac{(5A+12iB) \left(\frac{2 \int \frac{(i \tan(e+fx)a+a)^{5/2}}{(c-ic \tan(e+fx))^{11/2}} d \tan(e+fx)}{11c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{13c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{13ac(c-ic \tan(e+fx))^{13/2}} \right)}{15c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{15ac(c-ic \tan(e+fx))^{15/2}} \right)$$

f

↓ 55

3.829. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{17/2}} dx$

$$\left(\frac{(5A+12iB) \left(\frac{2 \left(\int \frac{(i \tan(e+fx)a+a)^{5/2}}{(c-ic \tan(e+fx))^{9/2}} d \tan(e+fx) - \frac{i(a+ia \tan(e+fx))^{7/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{4} - \frac{i(a+ia \tan(e+fx))^{7/2}}{13ac(c-ic \tan(e+fx))^{13/2}} \right)}{15c} \right)$$

$$\frac{ac}{17c}$$

f

↓ 48

3.829. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{17/2}} dx$

$$\begin{array}{c}
 \left(\frac{(5A+12iB) \left(\frac{2 \left(-\frac{i(a+ia \tan(e+fx))^{7/2}}{63ac^2(c-ic \tan(e+fx))^{7/2}} - \frac{i(a+ia \tan(e+fx))^{7/2}}{9ac(c-ic \tan(e+fx))^{9/2}} \right)}{11c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{11ac(c-ic \tan(e+fx))^{11/2}} \right)}{13c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{13ac(c-ic \tan(e+fx))^{13/2}} \right)}{15c} - \frac{i(a+ia \tan(e+fx))^{7/2}}{15ac(c-ic \tan(e+fx))^{15/2}} \right) \\
 \frac{ac}{17c}
 \end{array}$$

f

input `Int[((a + I*a*Tan[e + f*x])^(7/2)*(A + B*Tan[e + f*x]))/(c - I*c*Tan[e + f*x])^(17/2),x]`

output `(a*c*(-1/17*((I*A + B)*(a + I*a*Tan[e + f*x])^(7/2))/(a*c*(c - I*c*Tan[e + f*x])^(17/2)) + ((5*A + (12*I)*B)*((-1/15*I)*(a + I*a*Tan[e + f*x])^(7/2)))/(a*c*(c - I*c*Tan[e + f*x])^(15/2)) + (4*(((1/13*I)*(a + I*a*Tan[e + f*x])^(7/2)))/(a*c*(c - I*c*Tan[e + f*x])^(13/2)) + (3*(((1/11*I)*(a + I*a*Tan[e + f*x])^(7/2)))/(a*c*(c - I*c*Tan[e + f*x])^(11/2)) + (2*(((1/9*I)*(a + I*a*Tan[e + f*x])^(7/2)))/(a*c*(c - I*c*Tan[e + f*x])^(9/2)) - ((I/63)*(a + I*a*Tan[e + f*x])^(7/2))/(a*c^2*(c - I*c*Tan[e + f*x])^(7/2))))/(11*c)))/(13*c))/(15*c))/(17*c))/f`

3.829.3.1 Defintions of rubi rules used

- rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`
- rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
.), x] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]`

3.829.4 Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 206, normalized size of antiderivative = 0.66

method	result
risch	$a^3 \frac{\sqrt{\frac{a e^{2i(fx+e)}}{e^{2i(fx+e)}+1}}}{e^{2i(fx+e)}+1} (45045iA e^{16i(fx+e)}+45045B e^{16i(fx+e)}+255255iA e^{14i(fx+e)}+153153B e^{14i(fx+e)}+589050iA e^{12i(fx+e)}+117810B e^{12i(fx+e)}+696150iA e^{10i(fx+e)}+139230B e^{10i(fx+e)}+425425iA e^{8i(fx+e)}+255255B e^{8i(fx+e)}+109395iA e^{6i(fx+e)}-109395B e^{6i(fx+e)})$
derivativedivides	$i\sqrt{a(1+i\tan(fx+e))} \sqrt{-c(i\tan(fx+e)-1)} a^3 (1+\tan(fx+e))^2 (5871iB+12960iB \tan(fx+e)^4-96B \tan(fx+e)^7+1153153f c^9(i+\tan(fx+e))^{10})$
default	$i\sqrt{a(1+i\tan(fx+e))} \sqrt{-c(i\tan(fx+e)-1)} a^3 (1+\tan(fx+e))^2 (5871iB+12960iB \tan(fx+e)^4-96B \tan(fx+e)^7+1153153f c^9(i+\tan(fx+e))^{10})$
parts	$iA\sqrt{a(1+i\tan(fx+e))} \sqrt{-c(i\tan(fx+e)-1)} a^3 (1+\tan(fx+e))^2 (8i \tan(fx+e)^7-372i \tan(fx+e)^5-80 \tan(fx+e)^6+153153f c^9(i+\tan(fx+e))^{10})$

input `int((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(17/2),x,method=_RETURNVERBOSE)`

output `-1/24504480*a^3/c^8*(a*exp(2*I*(f*x+e))/(exp(2*I*(f*x+e))+1))^(1/2)/(c/(exp(2*I*(f*x+e))+1))^(1/2)/f*(45045*I*A*exp(16*I*(f*x+e))+45045*B*exp(16*I*(f*x+e))+255255*I*A*exp(14*I*(f*x+e))+153153*B*exp(14*I*(f*x+e))+589050*I*A*exp(12*I*(f*x+e))+117810*B*exp(12*I*(f*x+e))+696150*I*exp(10*I*(f*x+e))*A-139230*B*exp(10*I*(f*x+e))+425425*I*exp(8*I*(f*x+e))*A-255255*B*exp(8*I*(f*x+e))+109395*I*A*exp(6*I*(f*x+e))-109395*B*exp(6*I*(f*x+e)))`

3.829.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 188, normalized size of antiderivative = 0.60

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{17/2}} dx =$$

$$(45045 (iA + B)a^3 e^{(19i fx+19ie)} + 6006 (50iA + 33B)a^3 e^{(17i fx+17ie)} + 3927 (215iA + 69B)a^3 e^{(15i fx+15ie)})$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(17/2),x, algorithm="fricas")`

output $-1/24504480*(45045*(I*A + B)*a^3*e^{(19*I*f*x + 19*I*e)} + 6006*(50*I*A + 33*B)*a^3*e^{(17*I*f*x + 17*I*e)} + 3927*(215*I*A + 69*B)*a^3*e^{(15*I*f*x + 15*I*e)} + 21420*(60*I*A - B)*a^3*e^{(13*I*f*x + 13*I*e)} + 7735*(145*I*A - 51*B)*a^3*e^{(11*I*f*x + 11*I*e)} + 24310*(22*I*A - 15*B)*a^3*e^{(9*I*f*x + 9*I*e)} + 109395*(I*A - B)*a^3*e^{(7*I*f*x + 7*I*e)})*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)}/(c^9*f)$

3.829.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{17/2}} dx = \text{Timed out}$$

input `integrate((a+I*a*tan(f*x+e))**(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(17/2),x)`

output Timed out

3.829.7 Maxima [A] (verification not implemented)

Time = 0.56 (sec) , antiderivative size = 412, normalized size of antiderivative = 1.31

$$\int \frac{(a + ia \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(c - ictan(e + fx))^{17/2}} dx = \frac{(45045(-iA - B)a^3 \cos(\frac{17}{2} \arctan(\sin(2fx + 2e)), c$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(17/2),x, algorithm="maxima")`

output $1/24504480*(45045*(-I*A - B)*a^3*\cos(17/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 51051*(-5*I*A - 3*B)*a^3*\cos(15/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 117810*(-5*I*A - B)*a^3*\cos(13/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 139230*(-5*I*A + B)*a^3*\cos(11/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 85085*(-5*I*A + 3*B)*a^3*\cos(9/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 109395*(-I*A + B)*a^3*\cos(7/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 45045*(A - I*B)*a^3*\sin(17/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 51051*(5*A - 3*I*B)*a^3*\sin(15/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 117810*(5*A - I*B)*a^3*\sin(13/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 139230*(5*A + I*B)*a^3*\sin(11/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 85085*(5*A + 3*I*B)*a^3*\sin(9/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 109395*(A + I*B)*a^3*\sin(7/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))))*sqrt(a)/(c^(17/2)*f)$

3.829.8 Giac [F]

$$\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{17/2}} dx = \int \frac{(B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{7/2}}{(-ictan(fx + e) + c)^{17/2}} dx$$

input `integrate((a+I*a*tan(f*x+e))^(7/2)*(A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(17/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(7/2)/(-I*c*tan(f*x + e) + c)^(17/2), x)`

3.829.9 Mupad [B] (verification not implemented)

Time = 13.23 (sec) , antiderivative size = 229, normalized size of antiderivative = 0.73

$$\int \frac{(a + ia \tan(e + fx))^{7/2}(A + B \tan(e + fx))}{(c - ictan(e + fx))^{17/2}} dx = \frac{\sqrt{a + \frac{a \sin(e+fx)}{\cos(e+fx)} \operatorname{li} \left(\frac{a^3 e^{e 8i+fx 8i} (5A+B 3i) \operatorname{li}}{288 c^8 f} + \frac{a^3 e^{e 10i+fx 10i} (5A+B \operatorname{li}) \operatorname{li}}{176 c^8 f} + \frac{a^3 e^{e 12i+fx 12i} (5A-B \operatorname{li}) \operatorname{li}}{208 c^8 f} + \frac{a^3 e^{e 14i+fx 14i} (5A+B \operatorname{li}) \operatorname{li}}{480 c^8 f} \right)}}{\sqrt{c - \frac{c \sin(e+fx)}{\cos(e+fx)} \operatorname{li}}}$$

3.829. $\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ictan(e+fx))^{17/2}} dx$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(7/2))/(c - c*tan(e + f*x)*1i)^(17/2),x)`

output `-((a + (a*sin(e + f*x)*1i)/cos(e + f*x))^(1/2))*((a^3*exp(e*8i + f*x*8i)*(5*A + B*3i)*1i)/(288*c^8*f) + (a^3*exp(e*10i + f*x*10i)*(5*A + B*1i)*1i)/(176*c^8*f) + (a^3*exp(e*12i + f*x*12i)*(5*A - B*1i)*1i)/(208*c^8*f) + (a^3*exp(e*14i + f*x*14i)*(5*A - B*3i)*1i)/(480*c^8*f) + (a^3*exp(e*6i + f*x*6i)*(A + B*1i)*1i)/(224*c^8*f) + (a^3*exp(e*16i + f*x*16i)*(A - B*1i)*1i)/(544*c^8*f)))/(c - (c*sin(e + f*x)*1i)/cos(e + f*x))^(1/2)`

3.829.
$$\int \frac{(a+ia \tan(e+fx))^{7/2}(A+B \tan(e+fx))}{(c-ic \tan(e+fx))^{17/2}} dx$$

3.830
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{\sqrt{a+ia \tan(e+fx)}} dx$$

3.830.1 Optimal result 7468
 3.830.2 Mathematica [A] (verified) 7469
 3.830.3 Rubi [A] (verified) 7469
 3.830.4 Maple [B] (verified) 7472
 3.830.5 Fricas [B] (verification not implemented) 7473
 3.830.6 Sympy [F] 7474
 3.830.7 Maxima [B] (verification not implemented) 7474
 3.830.8 Giac [F] 7475
 3.830.9 Mupad [F(-1)] 7476

3.830.1 Optimal result

Integrand size = 45, antiderivative size = 228

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{\sqrt{a + ia \tan(e + fx)}} dx = \frac{3(2iA - 3B)c^{5/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{\sqrt{a}f} + \frac{3(2iA - 3B)c^2 \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}}{2af} + \frac{(2iA - 3B)c \sqrt{a + ia \tan(e + fx)} (c - ic \tan(e + fx))^{3/2}}{2af} + \frac{(iA - B)(c - ic \tan(e + fx))^{5/2}}{f \sqrt{a + ia \tan(e + fx)}}$$

output

```

3*(2*I*A-3*B)*c^(5/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I
*c*tan(f*x+e))^(1/2))/f/a^(1/2)+3/2*(2*I*A-3*B)*c^2*(a+I*a*tan(f*x+e))^(1/
2)*(c-I*c*tan(f*x+e))^(1/2)/a/f+1/2*(2*I*A-3*B)*c*(a+I*a*tan(f*x+e))^(1/2)
*(c-I*c*tan(f*x+e))^(3/2)/a/f+(I*A-B)*(c-I*c*tan(f*x+e))^(5/2)/f/(a+I*a*ta
n(f*x+e))^(1/2)
    
```

3.830.2 Mathematica [A] (verified)

Time = 7.74 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.71

$$\int \frac{(A + B \tan(e + fx))(c - i \tan(e + fx))^{5/2}}{\sqrt{a + ia \tan(e + fx)}} dx = \frac{c^2 \sqrt{c - i \tan(e + fx)} \left(-((2A + 3iB)(i + \tan(e + fx)) \right.$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2))/Sqrt[a + I*a*Tan[e + f*x]],x]`

output `(c^2*Sqrt[c - I*c*Tan[e + f*x]]*(-((2*A + (3*I)*B)*(I + Tan[e + f*x])) - B*(I + Tan[e + f*x])^2 + 6*((2*I)*A - 3*B)*(1 + (ArcSin[Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[2]*Sqrt[a]])*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[1 - I*Tan[e + f*x]])))/(2*f*Sqrt[a + I*a*Tan[e + f*x]])`

3.830.3 Rubi [A] (verified)Time = 0.43 (sec) , antiderivative size = 222, normalized size of antiderivative = 0.97, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 87, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c - i \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{\sqrt{a + ia \tan(e + fx)}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c - i \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{\sqrt{a + ia \tan(e + fx)}} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(A + B \tan(e + fx))(c - i \tan(e + fx))^{3/2}}{(i \tan(e + fx)a + a)^{3/2}} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{87} \end{aligned}$$

3.830. $\int \frac{(A + B \tan(e + fx))(c - i \tan(e + fx))^{5/2}}{\sqrt{a + ia \tan(e + fx)}} dx$

$$ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{ac\sqrt{a+ia \tan(e+fx)}} - \frac{(2A+3iB) \int \frac{(c-ic \tan(e+fx))^{3/2}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e+fx)}{a} \right)$$

f
↓ 60

$$ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{ac\sqrt{a+ia \tan(e+fx)}} - \frac{(2A+3iB) \left(\frac{3}{2} c \int \frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e+fx) - \frac{i\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{3/2}}{2a} \right)}{a} \right)$$

f
↓ 60

$$ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{ac\sqrt{a+ia \tan(e+fx)}} - \frac{(2A+3iB) \left(\frac{3}{2} c \left(c \int \frac{1}{\sqrt{i \tan(e+fx)a+a}\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a} \right) \right)}{a} \right)$$

f

↓ 45

$$ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{ac\sqrt{a+ia \tan(e+fx)}} - \frac{(2A+3iB) \left(\frac{3}{2} c \left(2c \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a} \right) \right)}{a} \right)$$

f

↓ 218

$$ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{ac\sqrt{a+ia \tan(e+fx)}} - \frac{(2A+3iB) \left(\frac{3}{2} c \left(-\frac{2i\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{\sqrt{a}} - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a} \right) \right)}{a} \right)$$

f

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2))/Sqrt[a + I*a*Tan[e + f*x]],x]`

3.830. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{\sqrt{a+ia \tan(e+fx)}} dx$

output $(a*c*((I*A - B)*(c - I*c*\text{Tan}[e + f*x])^{5/2})/(a*c*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]) - ((2*A + (3*I)*B)*((-1/2*I)*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]*(c - I*c*\text{Tan}[e + f*x])^{3/2})/a + (3*c*((-2*I)*\text{Sqrt}[c]*\text{ArcTan}[(\text{Sqrt}[c]*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]])/(\text{Sqrt}[a]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])])/(\text{Sqrt}[a] - (I*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])/a))/2)/a)/f$

3.830.3.1 Defintions of rubi rules used

rule 45 $\text{Int}[1/(\text{Sqrt}[(a_) + (b_)*(x_)]*\text{Sqrt}[(c_) + (d_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[2 \text{ Subst}[\text{Int}[1/(b - d*x^2), x], x, \text{Sqrt}[a + b*x]/\text{Sqrt}[c + d*x]], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{EqQ}[b*c + a*d, 0] \&\& !\text{GtQ}[c, 0]$

rule 60 $\text{Int}[(a_) + (b_)*(x_)]^{(m_)}*((c_) + (d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m+1)}*((c + d*x)^n/(b*(m+n+1))), x] + \text{Simp}[n*((b*c - a*d)/(b*(m+n+1))) \text{ Int}[(a + b*x)^m*(c + d*x)^{(n-1)}, x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{GtQ}[n, 0] \&\& \text{NeQ}[m+n+1, 0] \&\& !(\text{IGtQ}[m, 0] \&\& (!\text{IntegerQ}[n] || (\text{GtQ}[m, 0] \&\& \text{LtQ}[m-n, 0]))) \&\& !\text{ILtQ}[m+n+2, 0] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 87 $\text{Int}[(a_) + (b_)*(x_)]*((c_) + (d_)*(x_)]^{(n_)}*((e_) + (f_)*(x_)]^{(p_)}, x_] \rightarrow \text{Simp}[(-b*e - a*f)*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)})/(f*(p+1)*(c*f - d*e)), x] - \text{Simp}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1)))/(f*(p+1)*(c*f - d*e)) \text{ Int}[(c + d*x)^n*(e + f*x)^{(p+1)}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x] \&\& \text{LtQ}[p, -1] \&\& (!\text{LtQ}[n, -1] || \text{IntegerQ}[p] || !(\text{IntegerQ}[n] || !(\text{EqQ}[e, 0] || !(\text{EqQ}[c, 0] || \text{LtQ}[p, n])))))$

rule 218 $\text{Int}[(a_) + (b_)*(x_)^2]^{(-1)}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{PosQ}[a/b]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.830.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 565 vs. 2(188) = 376.

Time = 0.39 (sec) , antiderivative size = 566, normalized size of antiderivative = 2.48

method	result
derivativedivides	$i\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c^2 \left(6iA \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) ac \tan(fx+e)^2+18iB \right)$
default	$i\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c^2 \left(6iA \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) ac \tan(fx+e)^2+18iB \right)$
parts	$iA\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c^2 \left(3i \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \tan(fx+e)^2 ac-3i \ln \left(\right) \right)$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^(1/2),x,m
ethod=_RETURNVERBOSE)
```

$$3.830. \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{\sqrt{a+ia \tan(e+fx)}} dx$$

output

```

1/2*I/f*(-c*(I*tan(f*x+e)-1))^(1/2)*(a*(1+I*tan(f*x+e)))^(1/2)*c^2/a*(6*I*
A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2)
)*a*c*tan(f*x+e)^2+18*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)
)^2))^(1/2))/(a*c)^(1/2)*a*c*tan(f*x+e)+4*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x
+e)^2))^(1/2)*tan(f*x+e)^2-9*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(
f*x+e)^2))^(1/2)))/(a*c)^(1/2)*a*c*tan(f*x+e)^2+B*(a*c)^(1/2)*(a*c*(1+tan(
f*x+e)^2))^(1/2)*tan(f*x+e)^3-6*I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1
+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2)*a*c-12*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x
+e)^2))^(1/2)*tan(f*x+e)+12*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f
*x+e)^2))^(1/2)))/(a*c)^(1/2)*a*c*tan(f*x+e)+2*A*(a*c*(1+tan(f*x+e)^2))^(1
/2)*(a*c)^(1/2)*tan(f*x+e)^2-14*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/
2)+9*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(
1/2)*a*c+19*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-10*A*(
a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c*(1+tan(f*x+e)^2))^(1/2)/(I-t
an(f*x+e))^2/(a*c)^(1/2)

```

3.830.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 548 vs. $2(176) = 352$.

Time = 0.28 (sec) , antiderivative size = 548, normalized size of antiderivative = 2.40

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{\sqrt{a + ia \tan(e + fx)}} dx =$$

$$3 \sqrt{\frac{(4A^2 + 12iAB - 9B^2)c^5}{af^2}} (afe^{(3i fx + 3ie)} + afe^{(i fx + ie)}) \log \left(\frac{4 \left(2((2iA - 3B)c^2 e^{(3i fx + 3ie)} + (2iA - 3B)c^2 e^{(i fx + ie)}) \sqrt{e^{(2i fx + 2ie)}} \right)}{(2iA - 3B)c^2 e^{(2i fx + 2ie)}} \right)$$

input

```

integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^(1/
2),x, algorithm="fricas")

```



```
output -1/4*(3*sqrt((4*A^2 + 12*I*A*B - 9*B^2)*c^5/(a*f^2))*(a*f*e^(3*I*f*x + 3*I
*e) + a*f*e^(I*f*x + I*e))*log(4*(2*((2*I*A - 3*B)*c^2*e^(3*I*f*x + 3*I*e)
+ (2*I*A - 3*B)*c^2*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sq
rt(c/(e^(2*I*f*x + 2*I*e) + 1)) + sqrt((4*A^2 + 12*I*A*B - 9*B^2)*c^5/(a*f
^2))*(a*f*e^(2*I*f*x + 2*I*e) - a*f))/((2*I*A - 3*B)*c^2*e^(2*I*f*x + 2*I*
e) + (2*I*A - 3*B)*c^2)) - 3*sqrt((4*A^2 + 12*I*A*B - 9*B^2)*c^5/(a*f^2))*
(a*f*e^(3*I*f*x + 3*I*e) + a*f*e^(I*f*x + I*e))*log(4*(2*((2*I*A - 3*B)*c^
2*e^(3*I*f*x + 3*I*e) + (2*I*A - 3*B)*c^2*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*
f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - sqrt((4*A^2 + 12*I*
A*B - 9*B^2)*c^5/(a*f^2))*(a*f*e^(2*I*f*x + 2*I*e) - a*f))/((2*I*A - 3*B)*
c^2*e^(2*I*f*x + 2*I*e) + (2*I*A - 3*B)*c^2)) + 4*(3*(-2*I*A + 3*B)*c^2*e^
(4*I*f*x + 4*I*e) + 5*(-2*I*A + 3*B)*c^2*e^(2*I*f*x + 2*I*e) + 4*(-I*A + B
)*c^2)*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))
)/(a*f*e^(3*I*f*x + 3*I*e) + a*f*e^(I*f*x + I*e))
```

3.830.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{\sqrt{a + ia \tan(e + fx)}} dx = \int \frac{(-ic(\tan(e + fx) + i))^{5/2} (A + B \tan(e + fx))}{\sqrt{ia(\tan(e + fx) - i)}} dx$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(5/2)/(a+I*a*tan(f*x+e))**(
1/2),x)
```

```
output Integral((-I*c*(tan(e + f*x) + I))**(5/2)*(A + B*tan(e + f*x))/sqrt(I*a*(t
an(e + f*x) - I)), x)
```

3.830.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1319 vs. $2(176) = 352$.

Time = 0.66 (sec) , antiderivative size = 1319, normalized size of antiderivative = 5.79

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{\sqrt{a + ia \tan(e + fx)}} dx = \text{Too large to display}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^(1/
2),x, algorithm="maxima")
```

3.830. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{\sqrt{a+ia \tan(e+fx)}} dx$

output

```

4*(12*(2*A + 3*I*B)*c^2*cos(4*f*x + 4*e) + 20*(2*A + 3*I*B)*c^2*cos(2*f*x
+ 2*e) + 12*(2*I*A - 3*B)*c^2*sin(4*f*x + 4*e) + 20*(2*I*A - 3*B)*c^2*sin(
2*f*x + 2*e) + 16*(A + I*B)*c^2 + 6*((2*A + 3*I*B)*c^2*cos(5/2*arctan2(sin
(2*f*x + 2*e), cos(2*f*x + 2*e))) + 2*(2*A + 3*I*B)*c^2*cos(3/2*arctan2(si
n(2*f*x + 2*e), cos(2*f*x + 2*e))) + (2*A + 3*I*B)*c^2*cos(1/2*arctan2(sin
(2*f*x + 2*e), cos(2*f*x + 2*e))) + (2*I*A - 3*B)*c^2*sin(5/2*arctan2(sin(
2*f*x + 2*e), cos(2*f*x + 2*e))) + 2*(2*I*A - 3*B)*c^2*sin(3/2*arctan2(sin
(2*f*x + 2*e), cos(2*f*x + 2*e))) + (2*I*A - 3*B)*c^2*sin(1/2*arctan2(sin(
2*f*x + 2*e), cos(2*f*x + 2*e))))*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e)
, cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))
+ 1) + 6*((2*A + 3*I*B)*c^2*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x +
2*e))) + 2*(2*A + 3*I*B)*c^2*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x
+ 2*e))) + (2*A + 3*I*B)*c^2*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x +
2*e))) + (2*I*A - 3*B)*c^2*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x +
2*e))) + 2*(2*I*A - 3*B)*c^2*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x +
2*e))) + (2*I*A - 3*B)*c^2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x +
2*e))))*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), -sin
(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) + 3*((2*I*A - 3*B)*
c^2*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 2*(2*I*A - 3*B)
*c^2*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (2*I*A - 3*...

```

3.830.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{\sqrt{a + ia \tan(e + fx)}} dx = \int \frac{(B \tan(fx + e) + A)(-ic \tan(fx + e) + c)^{5/2}}{\sqrt{ia \tan(fx + e) + a}} dx$$

input

```

integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^(1/
2),x, algorithm="giac")

```

output

```

integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(5/2)/sqrt(I*a*tan(
f*x + e) + a), x)

```

3.830.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{\sqrt{a + i a \tan(e + fx)}} dx = \int \frac{(A + B \tan(e + fx)) (c - c \tan(e + fx) i)^{5/2}}{\sqrt{a + a \tan(e + fx) i}} dx$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(5/2))/(a + a*tan(e + f*x)*1i)^(1/2),x)`

output `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(5/2))/(a + a*tan(e + f*x)*1i)^(1/2), x)`

$$3.831 \quad \int \frac{(A+B \tan(e+fx))(c-ictan(e+fx))^{3/2}}{\sqrt{a+ia \tan(e+fx)}} dx$$

3.831.1 Optimal result	7477
3.831.2 Mathematica [A] (verified)	7477
3.831.3 Rubi [A] (verified)	7478
3.831.4 Maple [B] (verified)	7480
3.831.5 Fricas [B] (verification not implemented)	7481
3.831.6 Sympy [F]	7482
3.831.7 Maxima [B] (verification not implemented)	7482
3.831.8 Giac [F]	7483
3.831.9 Mupad [F(-1)]	7484

3.831.1 Optimal result

Integrand size = 45, antiderivative size = 169

$$\int \frac{(A + B \tan(e + fx))(c - ictan(e + fx))^{3/2}}{\sqrt{a + ia \tan(e + fx)}} dx = \frac{2(iA - 2B)c^{3/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{\sqrt{a}f} + \frac{(iA - 2B)c\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}}{af} + \frac{(iA - B)(c - ictan(e + fx))^{3/2}}{f\sqrt{a + ia \tan(e + fx)}}$$

output

```
2*(I*A-2*B)*c^(3/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/f/a^(1/2)+(I*A-2*B)*c*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)/a/f+(I*A-B)*(c-I*c*tan(f*x+e))^(3/2)/f/(a+I*a*tan(f*x+e))^(1/2)
```

3.831.2 Mathematica [A] (verified)

Time = 4.60 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.80

$$\int \frac{(A + B \tan(e + fx))(c - ictan(e + fx))^{3/2}}{\sqrt{a + ia \tan(e + fx)}} dx = \frac{\sqrt{c - ictan(e + fx)} \left(Bc(1 - i \tan(e + fx)) + 2(iA - B)c \right)}{f\sqrt{a + ia \tan(e + fx)}}$$

input

```
Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2))/Sqrt[a + I*a*Tan[e + f*x]],x]
```

output $(\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]]*(B*c*(1 - I*\text{Tan}[e + f*x]) + 2*(I*A - 2*B)*c*(1 + (\text{ArcSin}[\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]/(\text{Sqrt}[2]*\text{Sqrt}[a]])*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]])/(\text{Sqrt}[a]*\text{Sqrt}[1 - I*\text{Tan}[e + f*x]]))))/(f*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]])$

3.831.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.01, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 87, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{\sqrt{a + ia \tan(e + fx)}} dx$$

↓ 3042

$$\int \frac{(c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{\sqrt{a + ia \tan(e + fx)}} dx$$

↓ 4071

$$ac \int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{(i \tan(e + fx) a + a)^{3/2}} d \tan(e + fx)$$

↓ 87

$$ac \left(\frac{(-B + iA)(c - ic \tan(e + fx))^{3/2}}{ac \sqrt{a + ia \tan(e + fx)}} - \frac{(A + 2iB) \int \frac{\sqrt{c - ic \tan(e + fx)}}{\sqrt{i \tan(e + fx) a + a}} d \tan(e + fx)}{a} \right)$$

↓ 60

$$ac \left(\frac{(-B + iA)(c - ic \tan(e + fx))^{3/2}}{ac \sqrt{a + ia \tan(e + fx)}} - \frac{(A + 2iB) \left(c \int \frac{1}{\sqrt{i \tan(e + fx) a + a} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) - \frac{i \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}}{a} \right)}{a} \right)$$

↓ 45

3.831. $\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{\sqrt{a + ia \tan(e + fx)}} dx$

$$\frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{3/2}}{ac\sqrt{a+ia \tan(e+fx)}} - \frac{(A+2iB) \left(2c \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a} \right)}{a} \right)}{f}$$

↓ 218

$$\frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{3/2}}{ac\sqrt{a+ia \tan(e+fx)}} - \frac{(A+2iB) \left(-\frac{2i\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{\sqrt{a}} - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a} \right)}{a} \right)}{f}$$

```
input Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2))/Sqrt[a + I*a*Tan[e + f*x]],x]
```

```
output (a*c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(3/2))/(a*c*Sqrt[a + I*a*Tan[e + f*x]]) - ((A + (2*I)*B)*(((2*I)*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[a] - (I*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/a))/a)/f
```

3.831.3.1 Defintions of rubi rules used

```
rule 45 Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]
```

```
rule 60 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

3.831. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{\sqrt{a+ia \tan(e+fx)}} dx$

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.831.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 498 vs. 2(140) = 280.

Time = 0.38 (sec) , antiderivative size = 499, normalized size of antiderivative = 2.95

method	result
derivativedivides	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c \left(-2iB \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right) ac \tan(fx+e)^2 + 2iA \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right)}{\dots}$
default	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c \left(-2iB \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right) ac \tan(fx+e)^2 + 2iA \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right)}{\dots}$
parts	$\frac{iA \sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c \left(i \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right) \tan(fx+e)^2 ac - i \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right)}{\dots}$

3.831.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{\sqrt{a+ia \tan(e+fx)}} dx$$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^(1/2),x,m
method=_RETURNVERBOSE)
```

```
output 1/f*(-c*(I*tan(f*x+e)-1))^(1/2)*(a*(1+I*tan(f*x+e)))^(1/2)/a*c*(-2*I*B*ln(
(a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c
*tan(f*x+e)^2+2*I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(
1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)-A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(
1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)^2+2*I*B*ln((a*c*tan(f*
x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c+4*I*B*(a*c
)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-4*B*ln((a*c*tan(f*x+e)+(a*
c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)-B*(a*c
)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2-2*I*A*(a*c)^(1/2)*(a*c*(1
+tan(f*x+e)^2))^(1/2)+A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^
2))^(1/2))/(a*c)^(1/2))*a*c+2*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*t
an(f*x+e)+3*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c*(1+tan(f*x+e
)^2))^(1/2)/(I-tan(f*x+e))^2/(a*c)^(1/2)
```

3.831.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 456 vs. $2(131) = 262$.

Time = 0.27 (sec) , antiderivative size = 456, normalized size of antiderivative = 2.70

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{\sqrt{a + ia \tan(e + fx)}} dx =$$

$$\left(a \sqrt{\frac{(A^2 + 4iAB - 4B^2)c^3}{af^2}} f e^{(ifx + ie)} \log \left(- \frac{4 \left(2((iA - 2B)ce^{(3ifx + 3ie)} + (iA - 2B)ce^{(ifx + ie)}) \sqrt{\frac{a}{e^{(2ifx + 2ie)} + 1}} \sqrt{\frac{c}{e^{(2ifx + 2ie)} + 1}} + (-iA + 2B)ce^{(2ifx + 2ie)} + (-iA + 2B)c \right)}{(-iA + 2B)ce^{(2ifx + 2ie)} + (-iA + 2B)c} \right) \right)$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^(1/
2),x, algorithm="fracas")
```

3.831. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{\sqrt{a+ia \tan(e+fx)}} dx$

output
$$-1/2*(a*\sqrt{(A^2 + 4*I*A*B - 4*B^2)*c^3/(a*f^2)})*f*e^{(I*f*x + I*e)}*\log(-4*(2*((I*A - 2*B)*c*e^{(3*I*f*x + 3*I*e)} + (I*A - 2*B)*c*e^{(I*f*x + I*e)}))*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} + (a*f*e^{(2*I*f*x + 2*I*e)} - a*f)*\sqrt{(A^2 + 4*I*A*B - 4*B^2)*c^3/(a*f^2)})/((-I*A + 2*B)*c*e^{(2*I*f*x + 2*I*e)} + (-I*A + 2*B)*c) - a*\sqrt{(A^2 + 4*I*A*B - 4*B^2)*c^3/(a*f^2)})*f*e^{(I*f*x + I*e)}*\log(-4*(2*((I*A - 2*B)*c*e^{(3*I*f*x + 3*I*e)} + (I*A - 2*B)*c*e^{(I*f*x + I*e)}))*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} - (a*f*e^{(2*I*f*x + 2*I*e)} - a*f)*\sqrt{(A^2 + 4*I*A*B - 4*B^2)*c^3/(a*f^2)})/((-I*A + 2*B)*c*e^{(2*I*f*x + 2*I*e)} + (-I*A + 2*B)*c) + 4*((-I*A + 2*B)*c*e^{(2*I*f*x + 2*I*e)} + (-I*A + B)*c)*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1))*e^{(-I*f*x - I*e)}/(a*f)$$

3.831.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{\sqrt{a + ia \tan(e + fx)}} dx = \int \frac{(-ic(\tan(e + fx) + i))^{3/2} (A + B \tan(e + fx))}{\sqrt{ia(\tan(e + fx) - i)}} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(3/2)/(a+I*a*tan(f*x+e))**(1/2),x)`

output `Integral((-I*c*(tan(e + f*x) + I))**(3/2)*(A + B*tan(e + f*x))/sqrt(I*a*(tan(e + f*x) - I)), x)`

3.831.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 898 vs. $2(131) = 262$.

Time = 0.47 (sec) , antiderivative size = 898, normalized size of antiderivative = 5.31

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{\sqrt{a + ia \tan(e + fx)}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^(1/2),x, algorithm="maxima")`

3.831.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{\sqrt{a+ia \tan(e+fx)}} dx$$

output

```
(4*(A + 2*I*B)*c*cos(2*f*x + 2*e) + 4*(I*A - 2*B)*c*sin(2*f*x + 2*e) + 4*(
A + I*B)*c + 2*((A + 2*I*B)*c*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x
+ 2*e))) + (A + 2*I*B)*c*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e
))) + (I*A - 2*B)*c*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) +
(I*A - 2*B)*c*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*arcta
n2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(s
in(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) + 2*((A + 2*I*B)*c*cos(3/2*arctan
2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (A + 2*I*B)*c*cos(1/2*arctan2(sin
(2*f*x + 2*e), cos(2*f*x + 2*e))) + (I*A - 2*B)*c*sin(3/2*arctan2(sin(2*f*
x + 2*e), cos(2*f*x + 2*e))) + (I*A - 2*B)*c*sin(1/2*arctan2(sin(2*f*x + 2
*e), cos(2*f*x + 2*e))))*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f
*x + 2*e))), -sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) -
((-I*A + 2*B)*c*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (-I
*A + 2*B)*c*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (A + 2*
I*B)*c*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (A + 2*I*B)*
c*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*log(cos(1/2*arctan
2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))^2 + sin(1/2*arctan2(sin(2*f*x + 2*e
), cos(2*f*x + 2*e))))^2 + 2*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x +
2*e))) + 1) - ((I*A - 2*B)*c*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x +
2*e))) + (I*A - 2*B)*c*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2...
```

3.831.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{\sqrt{a + i a \tan(e + fx)}} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{3/2}}{\sqrt{i a \tan(fx + e) + a}} dx$$

input

```
integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^(1/
2),x, algorithm="giac")
```

output

```
integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(3/2)/sqrt(I*a*tan(
f*x + e) + a), x)
```

3.831.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{\sqrt{a + i a \tan(e + fx)}} dx = \int \frac{(A + B \tan(e + fx)) (c - c \tan(e + fx) i)^{3/2}}{\sqrt{a + a \tan(e + fx) i}} dx$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(3/2))/(a + a*tan(e + f*x)*1i)^(1/2),x)`

output `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(3/2))/(a + a*tan(e + f*x)*1i)^(1/2), x)`

3.832
$$\int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{\sqrt{a+ia \tan(e+fx)}} dx$$

3.832.1 Optimal result 7485
 3.832.2 Mathematica [A] (verified) 7485
 3.832.3 Rubi [A] (verified) 7486
 3.832.4 Maple [B] (verified) 7488
 3.832.5 Fracas [B] (verification not implemented) 7489
 3.832.6 Sympy [F] 7489
 3.832.7 Maxima [A] (verification not implemented) 7490
 3.832.8 Giac [F] 7490
 3.832.9 Mupad [B] (verification not implemented) 7491

3.832.1 Optimal result

Integrand size = 45, antiderivative size = 110

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - ic \tan(e + fx)}}{\sqrt{a + ia \tan(e + fx)}} dx$$

$$= -\frac{2B\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{\sqrt{a}f} + \frac{(iA - B)\sqrt{c - ic \tan(e + fx)}}{f\sqrt{a + ia \tan(e + fx)}}$$

output `-2*B*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))*c^(1/2)/f/a^(1/2)+(I*A-B)*(c-I*c*tan(f*x+e))^(1/2)/f/(a+I*a*tan(f*x+e))^(1/2)`

3.832.2 Mathematica [A] (verified)

Time = 3.02 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.08

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - ic \tan(e + fx)}}{\sqrt{a + ia \tan(e + fx)}} dx$$

$$= -\frac{2B\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{\sqrt{a}f} + \frac{(A + iB)c(i + \tan(e + fx))}{f\sqrt{a + ia \tan(e + fx)}\sqrt{c - ic \tan(e + fx)}}$$

input `Integrate[((A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])/Sqrt[a + I*a*Tan[e + f*x]],x]`

3.832.
$$\int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{\sqrt{a+ia \tan(e+fx)}} dx$$

output $(-2*B*\text{Sqrt}[c]*\text{ArcTan}[(\text{Sqrt}[c]*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]])/(\text{Sqrt}[a]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])]/(\text{Sqrt}[a]*f) + ((A + I*B)*c*(I + \text{Tan}[e + f*x]))/(f*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])$

3.832.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4071, 87, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{c - ic \tan(e + fx)}(A + B \tan(e + fx))}{\sqrt{a + ia \tan(e + fx)}} dx$$

↓ 3042

$$\int \frac{\sqrt{c - ic \tan(e + fx)}(A + B \tan(e + fx))}{\sqrt{a + ia \tan(e + fx)}} dx$$

↓ 4071

$$\frac{ac \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx)a + a)^{3/2} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{f}$$

↓ 87

$$\frac{ac \left(\frac{(-B + iA) \sqrt{c - ic \tan(e + fx)}}{ac \sqrt{a + ia \tan(e + fx)}} - \frac{iB \int \frac{1}{\sqrt{i \tan(e + fx)a + a} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{a} \right)}{f}$$

↓ 45

$$ac \left(\frac{(-B + iA) \sqrt{c - ic \tan(e + fx)}}{ac \sqrt{a + ia \tan(e + fx)}} - \frac{2iB \int \frac{1}{ia + \frac{ic(i \tan(e + fx)a + a)}{c - ic \tan(e + fx)}} d \frac{\sqrt{i \tan(e + fx)a + a}}{\sqrt{c - ic \tan(e + fx)}}}{a} \right)$$

↓ 218

$$\frac{ac \left(\frac{(-B + iA) \sqrt{c - ic \tan(e + fx)}}{ac \sqrt{a + ia \tan(e + fx)}} - \frac{2B \arctan \left(\frac{\sqrt{c} \sqrt{a + ia \tan(e + fx)}}{\sqrt{a} \sqrt{c - ic \tan(e + fx)}} \right)}{a^{3/2} \sqrt{c}} \right)}{f}$$

3.832. $\int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{\sqrt{a + ia \tan(e + fx)}} dx$

input `Int[((A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])/Sqrt[a + I*a*Tan[e + f*x]],x]`

output `(a*c*((-2*B*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/(a^(3/2)*Sqrt[c]) + ((I*A - B)*Sqrt[c - I*c*Tan[e + f*x]])/(a*c*Sqrt[a + I*a*Tan[e + f*x]]))/f`

3.832.3.1 Defintions of rubi rules used

rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.832.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 322 vs. 2(89) = 178.

Time = 0.41 (sec) , antiderivative size = 323, normalized size of antiderivative = 2.94

method	result
derivativedivides	$\frac{i\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} \left(-2iB \ln \left(\frac{ac \tan(fx+e) + \sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) ac \tan(fx+e) + B \ln \right)}{\dots}$
default	$\frac{i\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} \left(-2iB \ln \left(\frac{ac \tan(fx+e) + \sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) ac \tan(fx+e) + B \ln \right)}{\dots}$
parts	$-\frac{iA\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} (1+i \tan(fx+e))}{fa(i-\tan(fx+e))^2} + \frac{iB\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))}}{\dots}$

```
input int((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
output -I/f*(-c*(I*tan(f*x+e)-1))^(1/2)*(a*(1+I*tan(f*x+e)))^(1/2)/a*(-2*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)+B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)^2+I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)-B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c-B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c*(1+tan(f*x+e)^2))^(1/2)/(I-tan(f*x+e))^2/(a*c)^(1/2)
```

$$3.832. \int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{\sqrt{a+ia \tan(e+fx)}} dx$$

3.832.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 348 vs. $2(84) = 168$.

Time = 0.27 (sec) , antiderivative size = 348, normalized size of antiderivative = 3.16

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{\sqrt{a + i a \tan(e + fx)}} dx$$

$$= \left(a f \sqrt{-\frac{B^2 c}{a f^2}} e^{(i f x + i e)} \log \left(\frac{4 \left(2 (B e^{(3 i f x + 3 i e)} + B e^{(i f x + i e)}) \sqrt{\frac{a}{e^{(2 i f x + 2 i e)} + 1}} \sqrt{\frac{c}{e^{(2 i f x + 2 i e)} + 1}} + (a f e^{(2 i f x + 2 i e)} - a f) \sqrt{-\frac{B^2 c}{a f^2}} \right)}{B e^{(2 i f x + 2 i e)} + B} \right) \right)$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2),x, algorithm="fracas")`

output `1/2*(a*f*sqrt(-B^2*c/(a*f^2))*e^(I*f*x + I*e)*log(4*(2*(B*e^(3*I*f*x + 3*I*e) + B*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + (a*f*e^(2*I*f*x + 2*I*e) - a*f)*sqrt(-B^2*c/(a*f^2)))/(B*e^(2*I*f*x + 2*I*e) + B) - a*f*sqrt(-B^2*c/(a*f^2))*e^(I*f*x + I*e)*log(4*(2*(B*e^(3*I*f*x + 3*I*e) + B*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - (a*f*e^(2*I*f*x + 2*I*e) - a*f)*sqrt(-B^2*c/(a*f^2)))/(B*e^(2*I*f*x + 2*I*e) + B)) - 2*((-I*A + B)*e^(2*I*f*x + 2*I*e) - I*A + B)*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))*e^(-I*f*x - I*e)/(a*f)`

3.832.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{\sqrt{a + i a \tan(e + fx)}} dx$$

$$= \int \frac{\sqrt{-i c (\tan(e + fx) + i)} (A + B \tan(e + fx))}{\sqrt{i a (\tan(e + fx) - i)}} dx$$

input `integrate((c-I*c*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**(1/2),x)`

output `Integral(sqrt(-I*c*(tan(e + f*x) + I))*(A + B*tan(e + f*x))/sqrt(I*a*(tan(e + f*x) - I)), x)`

3.832. $\int \frac{(A+B \tan(e+fx)) \sqrt{c-ic \tan(e+fx)}}{\sqrt{a+ia \tan(e+fx)}} dx$

3.832.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.25

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{\sqrt{a + i a \tan(e + fx)}} dx =$$

$$\frac{(2 B \arctan(\cos(fx + e), \sin(fx + e) + 1) + 2 B \arctan(\cos(fx + e), -\sin(fx + e) + 1) - 2(i A -$$

```
input integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2),x, algorithm="maxima")
```

```
output -1/2*(2*B*arctan2(cos(f*x + e), sin(f*x + e) + 1) + 2*B*arctan2(cos(f*x + e), -sin(f*x + e) + 1) - 2*(I*A - B)*cos(f*x + e) + I*B*log(cos(f*x + e)^2 + sin(f*x + e)^2 + 2*sin(f*x + e) + 1) - I*B*log(cos(f*x + e)^2 + sin(f*x + e)^2 - 2*sin(f*x + e) + 1) - 2*(A + I*B)*sin(f*x + e))*sqrt(c)/(sqrt(a)*f)
```

3.832.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{\sqrt{a + i a \tan(e + fx)}} dx$$

$$= \int \frac{(B \tan(fx + e) + A) \sqrt{-i c \tan(fx + e) + c}}{\sqrt{i a \tan(fx + e) + a}} dx$$

```
input integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2),x, algorithm="giac")
```

```
output integrate((B*tan(f*x + e) + A)*sqrt(-I*c*tan(f*x + e) + c)/sqrt(I*a*tan(f*x + e) + a), x)
```

3.832.9 Mupad [B] (verification not implemented)

Time = 11.56 (sec) , antiderivative size = 250, normalized size of antiderivative = 2.27

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i \tan(e + fx)}}{\sqrt{a + i a \tan(e + fx)}} dx$$

$$= \frac{A \sqrt{c - c \tan(e + fx) \operatorname{li} \operatorname{li}}}{f \sqrt{a + a \tan(e + fx) \operatorname{li}}} - \frac{4 B \sqrt{c} \operatorname{atan}\left(\frac{\sqrt{c} (\sqrt{a + a \tan(e + fx) \operatorname{li} - \sqrt{a}})}{\sqrt{a} (\sqrt{c - c \tan(e + fx) \operatorname{li} - \sqrt{c}})}\right)}{\sqrt{a} f}$$

$$- \frac{4 B (\sqrt{a + a \tan(e + fx) \operatorname{li}} - \sqrt{a})}{f (\sqrt{c - c \tan(e + fx) \operatorname{li}} - \sqrt{c}) \left(-\frac{a}{c} + \frac{(\sqrt{a + a \tan(e + fx) \operatorname{li} - \sqrt{a}})^2}{(\sqrt{c - c \tan(e + fx) \operatorname{li} - \sqrt{c}})^2} + \frac{2 \sqrt{a} (\sqrt{a + a \tan(e + fx) \operatorname{li} - \sqrt{a}})}{\sqrt{c} (\sqrt{c - c \tan(e + fx) \operatorname{li} - \sqrt{c}})} \right)}$$

```
input int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(1/2))/(a + a*tan(e + f*
x)*1i)^(1/2),x)
```

```
output (A*(c - c*tan(e + f*x)*1i)^(1/2)*1i)/(f*(a + a*tan(e + f*x)*1i)^(1/2)) - (
4*B*c^(1/2)*atan((c^(1/2)*((a + a*tan(e + f*x)*1i)^(1/2) - a^(1/2)))/(a^(1
/2)*((c - c*tan(e + f*x)*1i)^(1/2) - c^(1/2)))))/(a^(1/2)*f) - (4*B*((a +
a*tan(e + f*x)*1i)^(1/2) - a^(1/2)))/(f*((c - c*tan(e + f*x)*1i)^(1/2) - c
^(1/2))*((a + a*tan(e + f*x)*1i)^(1/2) - a^(1/2))^2/((c - c*tan(e + f*x)*
1i)^(1/2) - c^(1/2))^2 - a/c + (2*a^(1/2)*((a + a*tan(e + f*x)*1i)^(1/2) -
a^(1/2)))/(c^(1/2)*((c - c*tan(e + f*x)*1i)^(1/2) - c^(1/2))))))
```

3.833
$$\int \frac{A+B \tan(e+fx)}{\sqrt{a+ia \tan(e+fx)}\sqrt{c-ictan(e+fx)}} dx$$

3.833.1 Optimal result 7492
 3.833.2 Mathematica [A] (verified) 7492
 3.833.3 Rubi [A] (verified) 7493
 3.833.4 Maple [A] (verified) 7494
 3.833.5 Fricas [A] (verification not implemented) 7495
 3.833.6 Sympy [F] 7495
 3.833.7 Maxima [A] (verification not implemented) 7496
 3.833.8 Giac [F] 7496
 3.833.9 Mupad [B] (verification not implemented) 7497

3.833.1 Optimal result

Integrand size = 45, antiderivative size = 92

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}} dx$$

$$= -\frac{iA + B}{f\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}} + \frac{iA\sqrt{c - ictan(e + fx)}}{cf\sqrt{a + ia \tan(e + fx)}}$$

output `(-I*A-B)/f/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(1/2)+I*A*(c-I*c*tan(f*x+e))^(1/2)/c/f/(a+I*a*tan(f*x+e))^(1/2)`

3.833.2 Mathematica [A] (verified)

Time = 2.49 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.54

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}} dx$$

$$= \frac{-B + A \tan(e + fx)}{f\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}}$$

input `Integrate[(A + B*Tan[e + f*x])/(Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]]),x]`

output `(-B + A*Tan[e + f*x])/(f*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])`

3.833.
$$\int \frac{A+B \tan(e+fx)}{\sqrt{a+ia \tan(e+fx)}\sqrt{c-ictan(e+fx)}} dx$$

3.833.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.10, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.089$, Rules used = {3042, 4071, 87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}} dx$$

↓ 4071

$$ac \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx)a + a)^{3/2} (c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)$$

f
↓ 87

$$ac \left(\frac{A \int \frac{1}{(i \tan(e + fx)a + a)^{3/2} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{c} - \frac{B + iA}{ac \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}} \right)$$

f
↓ 48

$$\frac{ac \left(\frac{iA \sqrt{c - ic \tan(e + fx)}}{ac^2 \sqrt{a + ia \tan(e + fx)}} - \frac{B + iA}{ac \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}} \right)}{f}$$

input `Int[(A + B*Tan[e + f*x])/(Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]]),x]`

output `(a*c*(-((I*A + B)/(a*c*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])) + (I*A*Sqrt[c - I*c*Tan[e + f*x]])/(a*c^2*Sqrt[a + I*a*Tan[e + f*x]])))/f`

3.833.3.1 Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*(c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.833.4 Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00

method	result
risch	$-\frac{iAe^{2i(fx+e)}+Be^{2i(fx+e)}-iA+B}{2\sqrt{\frac{ae^{2i(fx+e)}}{e^{2i(fx+e)}+1}}(e^{2i(fx+e)}+1)\sqrt{\frac{c}{e^{2i(fx+e)}+1}}}f$
derivativedivides	$\frac{\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(A\tan(fx+e)^3-B\tan(fx+e)^2+A\tan(fx+e)-B)}{fac(i+\tan(fx+e))^2(i-\tan(fx+e))^2}$
default	$\frac{\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(A\tan(fx+e)^3-B\tan(fx+e)^2+A\tan(fx+e)-B)}{fac(i+\tan(fx+e))^2(i-\tan(fx+e))^2}$
parts	$\frac{A\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(1+\tan(fx+e)^2)\tan(fx+e)}{fac(i-\tan(fx+e))^2(i+\tan(fx+e))^2} + \frac{B(-\tan(fx+e)^2-1)\sqrt{-c(i\tan(fx+e)-1)}}{fca(i-\tan(fx+e))^2(i+\tan(fx+e))^2}$

3.833.
$$\int \frac{A+B\tan(e+fx)}{\sqrt{a+ia\tan(e+fx)}\sqrt{c-ictan(e+fx)}} dx$$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

output
$$-1/2/(a*\exp(2*I*(f*x+e))/(\exp(2*I*(f*x+e))+1))^(1/2)/(\exp(2*I*(f*x+e))+1)/(c/(\exp(2*I*(f*x+e))+1))^(1/2)*(I*A*\exp(2*I*(f*x+e))+B*\exp(2*I*(f*x+e))-I*(A+B))/f$$

3.833.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.24

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}} dx$$

$$= \frac{((-iA - B)e^{(4i fx + 4i e)} + 2Be^{(3i fx + 3i e)} - 2Be^{(2i fx + 2i e)} + 2Be^{(i fx + i e)} + iA - B) \sqrt{\frac{a}{e^{(2i fx + 2i e)} + 1}} \sqrt{\frac{c}{e^{(2i fx + 2i e)} + 1}}}{2acf}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(1/2),x, algorithm="fricas")`

output
$$1/2*((-I*A - B)*e^{(4*I*f*x + 4*I*e)} + 2*B*e^{(3*I*f*x + 3*I*e)} - 2*B*e^{(2*I*f*x + 2*I*e)} + 2*B*e^{(I*f*x + I*e)} + I*A - B)*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)}*e^{(-I*f*x - I*e)}/(a*c*f)$$

3.833.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}} dx$$

$$= \int \frac{A + B \tan(e + fx)}{\sqrt{ia (\tan(e + fx) - i)} \sqrt{-ic (\tan(e + fx) + i)}} dx$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**(1/2)/(c-I*c*tan(f*x+e))**(1/2),x)`

output `Integral((A + B*tan(e + f*x))/(sqrt(I*a*(tan(e + f*x) - I))*sqrt(-I*c*(tan(e + f*x) + I))), x)`

3.833.
$$\int \frac{A+B \tan(e+fx)}{\sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}} dx$$

3.833.7 Maxima [A] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.35

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}} dx =$$

$$\frac{2((A - iB) \cos(4fx + 4e) - 2iB \cos(2fx + 2e) - (-iA - B) \sin(4fx + 4e) + 2B \sin(2fx + 2e))}{-4(iac \cos(3fx + 3e) + iac \cos(fx + e) - ac \sin(3fx + 3e) - ac \sin(fx + e))} f$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(1/2),x, algorithm="maxima")
```

```
output -2*((A - I*B)*cos(4*f*x + 4*e) - 2*I*B*cos(2*f*x + 2*e) - (-I*A - B)*sin(4*f*x + 4*e) + 2*B*sin(2*f*x + 2*e) - A - I*B)*sqrt(a)*sqrt(c)/((-4*I*a*c*cos(3*f*x + 3*e) - 4*I*a*c*cos(f*x + e) + 4*a*c*sin(3*f*x + 3*e) + 4*a*c*sin(f*x + e))*f)
```

3.833.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}} dx$$

$$= \int \frac{B \tan(fx + e) + A}{\sqrt{ia \tan(fx + e) + a} \sqrt{-ic \tan(fx + e) + c}} dx$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(1/2),x, algorithm="giac")
```

```
output integrate((B*tan(f*x + e) + A)/(sqrt(I*a*tan(f*x + e) + a)*sqrt(-I*c*tan(f*x + e) + c)), x)
```

3.833.9 Mupad [B] (verification not implemented)

Time = 0.80 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.55

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}} dx =$$

$$\frac{\sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}} (A 1i + B - A \cos(2e + 2fx) 1i + B \cos(2e + 2fx) - A \sin(2e + 2fx))}{2af \sqrt{\frac{c(\cos(2e+2fx)+1-\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}}}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^(1/2)*(c - c*tan(e + f*x)*1i)^(1/2)),x)`

output `-(((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*1i + B - A*cos(2*e + 2*f*x)*1i + B*cos(2*e + 2*f*x) - A*sin(2*e + 2*f*x) - B*sin(2*e + 2*f*x)*1i))/(2*a*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

3.834
$$\int \frac{A+B \tan(e+fx)}{\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{3/2}} dx$$

3.834.1 Optimal result	7498
3.834.2 Mathematica [A] (verified)	7498
3.834.3 Rubi [A] (verified)	7499
3.834.4 Maple [A] (verified)	7501
3.834.5 Fricas [A] (verification not implemented)	7501
3.834.6 Sympy [F]	7502
3.834.7 Maxima [F(-2)]	7502
3.834.8 Giac [F]	7502
3.834.9 Mupad [B] (verification not implemented)	7503

3.834.1 Optimal result

Integrand size = 45, antiderivative size = 157

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}} dx = \frac{iA - B}{f \sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}} - \frac{(2iA - B)\sqrt{a + ia \tan(e + fx)}}{3af(c - ic \tan(e + fx))^{3/2}} - \frac{(2iA - B)\sqrt{a + ia \tan(e + fx)}}{3acf \sqrt{c - ic \tan(e + fx)}}$$

output `-1/3*(2*I*A-B)*(a+I*a*tan(f*x+e))^(1/2)/a/c/f/(c-I*c*tan(f*x+e))^(1/2)+(I*A-B)/f/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(3/2)-1/3*(2*I*A-B)*(a+I*a*tan(f*x+e))^(1/2)/a/f/(c-I*c*tan(f*x+e))^(3/2)`

3.834.2 Mathematica [A] (verified)

Time = 4.82 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.62

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}} dx = \frac{A - iB + (2iA - B) \tan(e + fx) + (2A + iB) \tan^2(e + fx)}{3cf(i + \tan(e + fx))\sqrt{a + ia \tan(e + fx)}\sqrt{c - ic \tan(e + fx)}}$$

input `Integrate[(A + B*Tan[e + f*x])/(Sqrt[a + I*a*Tan[e + f*x]]*(c - I*c*Tan[e + f*x])^(3/2)),x]`

output `(A - I*B + ((2*I)*A - B)*Tan[e + f*x] + (2*A + I*B)*Tan[e + f*x]^2)/(3*c*f*(I + Tan[e + f*x])*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])`

3.834.
$$\int \frac{A+B \tan(e+fx)}{\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{3/2}} dx$$

3.834.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.04, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4071, 87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx)a + a)^{3/2}(c - ic \tan(e + fx))^{5/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{87} \\
 & \frac{ac \left(\frac{(2A + iB) \int \frac{1}{\sqrt{i \tan(e + fx)a + a}(c - ic \tan(e + fx))^{5/2}} d \tan(e + fx)}{a} + \frac{-B + iA}{ac \sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}} \right)}{f} \\
 & \quad \downarrow \text{55} \\
 & \frac{ac \left(\frac{(2A + iB) \left(\frac{\int \frac{1}{\sqrt{i \tan(e + fx)a + a}(c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{3c} - \frac{i \sqrt{a + ia \tan(e + fx)}}{3ac(c - ic \tan(e + fx))^{3/2}} \right)}{a} + \frac{-B + iA}{ac \sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}} \right)}{f} \\
 & \quad \downarrow \text{48} \\
 & \frac{ac \left(\frac{(2A + iB) \left(-\frac{i \sqrt{a + ia \tan(e + fx)}}{3ac^2 \sqrt{c - ic \tan(e + fx)}} - \frac{i \sqrt{a + ia \tan(e + fx)}}{3ac(c - ic \tan(e + fx))^{3/2}} \right)}{a} + \frac{-B + iA}{ac \sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}} \right)}{f}
 \end{aligned}$$

input `Int[(A + B*Tan[e + f*x])/(Sqrt[a + I*a*Tan[e + f*x]]*(c - I*c*Tan[e + f*x])^(3/2)),x]`

$$3.834. \quad \int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}} dx$$

output $(a*c*((I*A - B)/(a*c*Sqrt[a + I*a*Tan[e + f*x]]*(c - I*c*Tan[e + f*x])^(3/2)) + ((2*A + I*B)*((-1/3*I)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c*(c - I*c*Tan[e + f*x])^(3/2)) - ((I/3)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c^2*Sqrt[c - I*c*Tan[e + f*x]])))/a)/f$

3.834.3.1 Defintions of rubi rules used

rule 48 $\text{Int}[(a_.) + (b_.)*(x_)^(m_.)*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x] \&\& \text{EqQ}[m + n + 2, 0] \&\& \text{NeQ}[m, -1]$

rule 55 $\text{Int}[(a_.) + (b_.)*(x_)^(m_.)*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - \text{Simp}[d*(\text{Simplify}[m + n + 2]/((b*c - a*d)*(m + 1))) \text{Int}[(a + b*x)^\text{Simplify}[m + 1]*(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x] \&\& \text{ILtQ}[\text{Simplify}[m + n + 2], 0] \&\& \text{NeQ}[m, -1] \&\& !(\text{LtQ}[m, -1] \&\& \text{LtQ}[n, -1] \&\& (\text{EqQ}[a, 0] || (\text{NeQ}[c, 0] \&\& \text{LtQ}[m - n, 0] \&\& \text{IntegerQ}[n]))) \&\& (\text{SumSimplerQ}[m, 1] || !\text{SumSimplerQ}[n, 1])$

rule 87 $\text{Int}[(a_.) + (b_.)*(x_) * ((c_.) + (d_.)*(x_)^(n_.)) * ((e_.) + (f_.)*(x_)^(p_.), x_] \rightarrow \text{Simp}[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - \text{Simp}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) \text{Int}[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x] \&\& \text{LtQ}[p, -1] \&\& (!\text{LtQ}[n, -1] || \text{IntegerQ}[p] || !(\text{IntegerQ}[n] || !(\text{EqQ}[e, 0] || !(\text{EqQ}[c, 0] || \text{LtQ}[p, n]))))$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4071 $\text{Int}[(a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_)]^(m_.)*((A_.) + (B_.)*\tan[(e_.) + (f_.)*(x_)] * ((c_.) + (d_.)*\tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] \rightarrow \text{Simp}[a*(c/f) \text{Subst}[\text{Int}[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, \text{Tan}[e + f*x], x] /; \text{FreeQ}\{a, b, c, d, e, f, A, B, m, n\}, x] \&\& \text{EqQ}[b*c + a*d, 0] \&\& \text{EqQ}[a^2 + b^2, 0]$

3.834.4 Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.70

method	result
risch	$\frac{iA e^{4i(fx+e)} + B e^{4i(fx+e)} + 6iA e^{2i(fx+e)} - 3iA + 3B}{12c \sqrt{\frac{a e^{2i(fx+e)}}{e^{2i(fx+e)} + 1}} (e^{2i(fx+e)} + 1) \sqrt{\frac{c}{e^{2i(fx+e)} + 1}}} f$
derivativedivides	$\frac{i \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} (2iA \tan(fx+e)^4 - iB \tan(fx+e)^3 - B \tan(fx+e)^4 + 3iA \tan(fx+e)^2 - 2A)}{3fa c^2 (i + \tan(fx+e))^3 (i - \tan(fx+e))^2}$
default	$\frac{i \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} (2iA \tan(fx+e)^4 - iB \tan(fx+e)^3 - B \tan(fx+e)^4 + 3iA \tan(fx+e)^2 - 2A)}{3fa c^2 (i + \tan(fx+e))^3 (i - \tan(fx+e))^2}$
parts	$\frac{A \sqrt{a(1+i \tan(fx+e))} \sqrt{-c(i \tan(fx+e)-1)} (2i \tan(fx+e)^3 + 2 \tan(fx+e)^4 + 2i \tan(fx+e) + 3 \tan(fx+e)^2 + 1)}{3fa c^2 (i - \tan(fx+e))^2 (i + \tan(fx+e))^3} + \frac{iB}{f}$

```
input int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
```

```
output -1/12/c/(a*exp(2*I*(f*x+e))/(exp(2*I*(f*x+e))+1))^(1/2)/(exp(2*I*(f*x+e))+1)/(c/(exp(2*I*(f*x+e))+1))^(1/2)*(I*A*exp(4*I*(f*x+e))+B*exp(4*I*(f*x+e))+6*I*A*exp(2*I*(f*x+e))-3*I*A+3*B)/f
```

3.834.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.94

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)(c - ictan(e + fx))^{3/2}}} dx = \frac{((-iA - B)e^{(6i fx + 6i e)} + (-7iA - B)e^{(4i fx + 4i e)} - 4(-iA - B)e^{(3i fx + 3i e)} - 3(IA + B)e^{(2i fx + 2i e)} - 4(-iA - B)e^{(i fx + i e)} + 3IA - 3B) \sqrt{a/(e^{(2i fx + 2i e)} + 1)} \operatorname{sqrt}(c/(e^{(2i fx + 2i e)} + 1)) e^{(-i fx - i e)}}{(a c^2 f)}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="fricas")
```

```
output 1/12*((-I*A - B)*e^(6*I*f*x + 6*I*e) + (-7*I*A - B)*e^(4*I*f*x + 4*I*e) - 4*(-I*A - B)*e^(3*I*f*x + 3*I*e) - 3*(I*A + B)*e^(2*I*f*x + 2*I*e) - 4*(-I*A - B)*e^(I*f*x + I*e) + 3*I*A - 3*B)*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*e^(-I*f*x - I*e)/(a*c^2*f)
```

3.834.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}} dx = \int \frac{A + B \tan(e + fx)}{\sqrt{ia (\tan(e + fx) - i)} (-ic (\tan(e + fx) + i))^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**(1/2)/(c-I*c*tan(f*x+e))**(3/2),x)`

output `Integral((A + B*tan(e + f*x))/(sqrt(I*a*(tan(e + f*x) - I))*(-I*c*(tan(e + f*x) + I))**(3/2)), x)`

3.834.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.834.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}} dx = \int \frac{B \tan(fx + e) + A}{\sqrt{ia \tan(fx + e) + a}(-ic \tan(fx + e) + c)^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)/(sqrt(I*a*tan(f*x + e) + a)*(-I*c*tan(f*x + e) + c)^(3/2)), x)`

3.834. $\int \frac{A+B \tan(e+fx)}{\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{3/2}} dx$

3.834.9 Mupad [B] (verification not implemented)

Time = 0.73 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.93

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ictan(e + fx))^{3/2}} dx = \frac{\sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}} (2A \sin(2e + 2fx) + \dots)}{6acf \sqrt{\dots}}$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^(1/2)*(c - c*tan(e + f*x)*1i)^(3/2)),x)`

output `((a*cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1)^(1/2)*(A*cos(2*e + 2*f*x)*1i - A*3i - 2*B*cos(2*e + 2*f*x) + 2*A*sin(2*e + 2*f*x) + B*sin(2*e + 2*f*x)*1i))/(6*a*c*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

3.835
$$\int \frac{A+B \tan(e+fx)}{\sqrt{a+ia \tan(e+fx)}(c-ictan(e+fx))^{5/2}} dx$$

3.835.1 Optimal result 7504
 3.835.2 Mathematica [A] (verified) 7504
 3.835.3 Rubi [A] (verified) 7505
 3.835.4 Maple [A] (verified) 7507
 3.835.5 Fricas [A] (verification not implemented) 7508
 3.835.6 Sympy [F] 7508
 3.835.7 Maxima [F(-2)] 7509
 3.835.8 Giac [F] 7509
 3.835.9 Mupad [B] (verification not implemented) 7509

3.835.1 Optimal result

Integrand size = 45, antiderivative size = 213

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ictan(e + fx))^{5/2}} dx = \frac{iA - B}{f\sqrt{a + ia \tan(e + fx)}(c - ictan(e + fx))^{5/2}} - \frac{(3iA - 2B)\sqrt{a + ia \tan(e + fx)}}{5af(c - ictan(e + fx))^{5/2}} - \frac{2(3iA - 2B)\sqrt{a + ia \tan(e + fx)}}{15acf(c - ictan(e + fx))^{3/2}} - \frac{2(3iA - 2B)\sqrt{a + ia \tan(e + fx)}}{15ac^2f\sqrt{c - ictan(e + fx)}}$$

output `-2/15*(3*I*A-2*B)*(a+I*a*tan(f*x+e))^(1/2)/a/c^2/f/(c-I*c*tan(f*x+e))^(1/2)+(I*A-B)/f/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(5/2)-1/5*(3*I*A-2*B)*(a+I*a*tan(f*x+e))^(1/2)/a/f/(c-I*c*tan(f*x+e))^(5/2)-2/15*(3*I*A-2*B)*(a+I*a*tan(f*x+e))^(1/2)/a/c/f/(c-I*c*tan(f*x+e))^(3/2)`

3.835.2 Mathematica [A] (verified)

Time = 6.24 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.54

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ictan(e + fx))^{5/2}} dx = \frac{6iA + B - (3A + 2iB) \tan(e + fx) + (12iA - 8B) \tan^2(e + fx)}{15c^2 f (i + \tan(e + fx))^2 \sqrt{a + ia \tan(e + fx)}}$$

input `Integrate[(A + B*Tan[e + f*x])/(Sqrt[a + I*a*Tan[e + f*x]]*(c - I*c*Tan[e + f*x])^(5/2)),x]`

3.835.
$$\int \frac{A+B \tan(e+fx)}{\sqrt{a+ia \tan(e+fx)}(c-ictan(e+fx))^{5/2}} dx$$

output $((6*I)*A + B - (3*A + (2*I)*B)*\text{Tan}[e + f*x] + ((12*I)*A - 8*B)*\text{Tan}[e + f*x]^2 + (6*A + (4*I)*B)*\text{Tan}[e + f*x]^3)/(15*c^2*f*(I + \text{Tan}[e + f*x])^2*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])$

3.835.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.02, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 87, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{5/2}} dx$$

↓ 4071

$$ac \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx)a + a)^{3/2}(c - ic \tan(e + fx))^{7/2}} d \tan(e + fx)$$

f
↓ 87

$$ac \left(\frac{(3A + 2iB) \int \frac{1}{\sqrt{i \tan(e + fx)a + a}(c - ic \tan(e + fx))^{7/2}} d \tan(e + fx)}{a} + \frac{-B + iA}{ac \sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{5/2}} \right)$$

f
↓ 55

$$ac \left(\frac{(3A + 2iB) \left(\frac{\int \frac{1}{\sqrt{i \tan(e + fx)a + a}(c - ic \tan(e + fx))^{5/2}} d \tan(e + fx)}{5c} - \frac{i \sqrt{a + ia \tan(e + fx)}}{5ac(c - ic \tan(e + fx))^{5/2}} \right)}{a} + \frac{-B + iA}{ac \sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{5/2}} \right)$$

f
↓ 55

3.835. $\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{5/2}} dx$

$$\begin{aligned}
 & \left(\frac{(3A+2iB) \left(\frac{2 \left(\frac{\int \frac{1}{\sqrt{i \tan(e+fx)a+a(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{3c} - \frac{i\sqrt{a+ia \tan(e+fx)}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{5c} - \frac{i\sqrt{a+ia \tan(e+fx)}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)}{a} \right)}{ac} + \frac{1}{ac\sqrt{a+ia \tan(e+fx)}} \right) \\
 & \hspace{15em} f \\
 & \quad \downarrow 48 \\
 & \left(\frac{(3A+2iB) \left(\frac{2 \left(-\frac{i\sqrt{a+ia \tan(e+fx)}}{3ac^2\sqrt{c-ic \tan(e+fx)}} - \frac{i\sqrt{a+ia \tan(e+fx)}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{5c} - \frac{i\sqrt{a+ia \tan(e+fx)}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)}{a} \right)}{ac} + \frac{-B+iA}{ac\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{5/2}} \right) \\
 & \hspace{15em} f
 \end{aligned}$$

```
input Int[(A + B*Tan[e + f*x])/(Sqrt[a + I*a*Tan[e + f*x]]*(c - I*c*Tan[e + f*x])^(5/2)),x]
```

```
output (a*c*((I*A - B)/(a*c*Sqrt[a + I*a*Tan[e + f*x]]*(c - I*c*Tan[e + f*x])^(5/2)) + ((3*A + (2*I)*B)*((( -1/5*I)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c*(c - I*c*Tan[e + f*x])^(5/2)) + (2*((( -1/3*I)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c*(c - I*c*Tan[e + f*x])^(3/2)) - ((I/3)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c^2*Sqrt[c - I*c*Tan[e + f*x]]))))/(5*c))/a)/f
```

3.835.3.1 Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

$$3.835. \quad \int \frac{A+B \tan(e+fx)}{\sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))^{5/2}} dx$$

```
rule 55 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.835.4 Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.69

method	result
risch	$-\frac{3iAe^{6i(fx+e)}+3Be^{6i(fx+e)}+15iAe^{4i(fx+e)}+5Be^{4i(fx+e)}+45iAe^{2i(fx+e)}-15Be^{2i(fx+e)}-15iA+15B}{120c^2\sqrt{\frac{ae^{2i(fx+e)}}{e^{2i(fx+e)}+1}}\sqrt{\frac{c}{e^{2i(fx+e)}+1}}(e^{2i(fx+e)}+1)f}$
derivativedivides	$\frac{\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}\left(4iB\tan(fx+e)^5+12iA\tan(fx+e)^4+6A\tan(fx+e)^5+2iB\tan(fx+e)^3-8iA\right)}{15fac^3(i+\tan(fx+e))^4}$
default	$\frac{\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}\left(4iB\tan(fx+e)^5+12iA\tan(fx+e)^4+6A\tan(fx+e)^5+2iB\tan(fx+e)^3-8iA\right)}{15fac^3(i+\tan(fx+e))^4}$
parts	$\frac{A\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}\left(4i\tan(fx+e)^4+2\tan(fx+e)^5+6i\tan(fx+e)^2+\tan(fx+e)^3+2i-\tan(fx+e)\right)}{5fac^3(i+\tan(fx+e))^4(i-\tan(fx+e))^2}$

3.835.
$$\int \frac{A+B\tan(e+fx)}{\sqrt{a+ia\tan(e+fx)(c-ictan(e+fx))^{5/2}}} dx$$

```
input int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)
```

```
output -1/120/c^2/(a*exp(2*I*(f*x+e))/(exp(2*I*(f*x+e))+1))^(1/2)/(c/(exp(2*I*(f*x+e))+1))^(1/2)/(exp(2*I*(f*x+e))+1)*(3*I*A*exp(6*I*(f*x+e))+3*B*exp(6*I*(f*x+e))+15*I*A*exp(4*I*(f*x+e))+5*B*exp(4*I*(f*x+e))+45*I*A*exp(2*I*(f*x+e))-15*B*exp(2*I*(f*x+e))-15*I*A+15*B)/f
```

3.835.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.76

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ictan(e + fx))^{5/2}} dx = \frac{(3(iA + B)e^{(8ifx+8ie)} + 2(9iA + 4B)e^{(6ifx+6ie)} + 10(6iA - B)e^{(4ifx+4ie)} + 8(-6iA - B)e^{(3ifx+3ie)} + 30iAe^{(2ifx+2ie)} + 8(-6iA - B)e^{(ifx+ie)} - 15iA + 15B)\sqrt{a/(e^{(2ifx+2ie)} + 1)}\sqrt{c/(e^{(2ifx+2ie)} + 1)}}{120ac^3f}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="fricas")
```

```
output -1/120*(3*(I*A + B)*e^(8*I*f*x + 8*I*e) + 2*(9*I*A + 4*B)*e^(6*I*f*x + 6*I*e) + 10*(6*I*A - B)*e^(4*I*f*x + 4*I*e) + 8*(-6*I*A - B)*e^(3*I*f*x + 3*I*e) + 30*I*A*e^(2*I*f*x + 2*I*e) + 8*(-6*I*A - B)*e^(I*f*x + I*e) - 15*I*A + 15*B)*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*e^(-I*f*x - I*e)/(a*c^3*f)
```

3.835.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ictan(e + fx))^{5/2}} dx = \int \frac{A + B \tan(e + fx)}{\sqrt{ia(\tan(e + fx) - i)}(-ic(\tan(e + fx) + i))^{5/2}} dx$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**(1/2)/(c-I*c*tan(f*x+e))**(5/2),x)
```

```
output Integral((A + B*tan(e + f*x))/(sqrt(I*a*(tan(e + f*x) - I))*(-I*c*(tan(e + f*x) + I))**(5/2)), x)
```

3.835. $\int \frac{A+B \tan(e+fx)}{\sqrt{a+ia \tan(e+fx)}(c-ictan(e+fx))^{5/2}} dx$

3.835.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ictan(e + fx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.835.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ictan(e + fx))^{5/2}} dx = \int \frac{B \tan(fx + e) + A}{\sqrt{ia \tan(fx + e) + a}(-ic \tan(fx + e) + c)^{5/2}} dx$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)/(sqrt(I*a*tan(f*x + e) + a)*(-I*c*tan(f*x + e) + c)^(5/2)), x)`

3.835.9 Mupad [B] (verification not implemented)

Time = 9.34 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.87

$$\int \frac{A + B \tan(e + fx)}{\sqrt{a + ia \tan(e + fx)}(c - ictan(e + fx))^{5/2}} dx =$$

$$\frac{\sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}} (A 45i - 15 B + A \cos(4e + 4fx) 3i + 20 B \cos(2e + 2fx) + 3 B \cos(4e + 4fx))}{120 a c^2 f \sqrt{c(\cos(2e+2fx)+1)}}$$

3.835. $\int \frac{A+B \tan(e+fx)}{\sqrt{a+ia \tan(e+fx)}(c-ictan(e+fx))^{5/2}} dx$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^(1/2)*(c - c*tan(e + f*x)*1i)^(5/2)),x)`

output `-(((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*45i - 15*B + A*cos(4*e + 4*f*x)*3i + 20*B*cos(2*e + 2*f*x) + 3*B*cos(4*e + 4*f*x) - 30*A*sin(2*e + 2*f*x) - 3*A*sin(4*e + 4*f*x) - B*sin(2*e + 2*f*x)*10i + B*sin(4*e + 4*f*x)*3i))/(120*a*c^2*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

3.836
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^{3/2}} dx$$

3.836.1 Optimal result 7511
 3.836.2 Mathematica [C] (verified) 7512
 3.836.3 Rubi [A] (verified) 7512
 3.836.4 Maple [B] (verified) 7516
 3.836.5 Fricas [B] (verification not implemented) 7517
 3.836.6 Sympy [F(-1)] 7517
 3.836.7 Maxima [B] (verification not implemented) 7518
 3.836.8 Giac [F] 7519
 3.836.9 Mupad [F(-1)] 7519

3.836.1 Optimal result

Integrand size = 45, antiderivative size = 287

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2}}{(a + ia \tan(e + fx))^{3/2}} dx =$$

$$-\frac{5(2iA - 5B)c^{7/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{a^{3/2}f}$$

$$-\frac{5(2iA - 5B)c^3 \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}}{2a^2 f}$$

$$-\frac{5(2iA - 5B)c^2 \sqrt{a + ia \tan(e + fx)}(c - ic \tan(e + fx))^{3/2}}{6a^2 f}$$

$$-\frac{2(2iA - 5B)c(c - ic \tan(e + fx))^{5/2}}{3af \sqrt{a + ia \tan(e + fx)}} + \frac{(iA - B)(c - ic \tan(e + fx))^{7/2}}{3f(a + ia \tan(e + fx))^{3/2}}$$

```
output -5*(2*I*A-5*B)*c^(7/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-
I*c*tan(f*x+e))^(1/2))/a^(3/2)/f-5/2*(2*I*A-5*B)*c^3*(a+I*a*tan(f*x+e))^(1
/2)*(c-I*c*tan(f*x+e))^(1/2)/a^2/f-5/6*(2*I*A-5*B)*c^2*(a+I*a*tan(f*x+e))^(
1/2)*(c-I*c*tan(f*x+e))^(3/2)/a^2/f-2/3*(2*I*A-5*B)*c*(c-I*c*tan(f*x+e))^(
5/2)/a/f/(a+I*a*tan(f*x+e))^(1/2)+1/3*(I*A-B)*(c-I*c*tan(f*x+e))^(7/2)/f/
(a+I*a*tan(f*x+e))^(3/2)
```

3.836.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 8.38 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.56

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2}}{(a + ia \tan(e + fx))^{3/2}} dx = \frac{c^3 \sec(e + fx) \sqrt{c - ic \tan(e + fx)} (20i(2A + 5iB) \cos^5(e + fx) - \dots)}{(a + ia \tan(e + fx))^{3/2}}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2))/(a + I*a*Tan[e + f*x])^(3/2),x]`

output `(c^3*Sec[e + f*x]*Sqrt[c - I*c*Tan[e + f*x]]*((20*I)*(2*A + (5*I)*B)*Cos[e + f*x]^3*Hypergeometric2F1[-3/2, -3/2, -1/2, (1 + I*Tan[e + f*x])/2]*Sqrt[2 - (2*I)*Tan[e + f*x]] + 3*(Cos[3*(e + f*x)] - I*Sin[3*(e + f*x)])*((-2*I)*A + 6*B - I*B*Tan[e + f*x])))/(6*a*f*Sqrt[a + I*a*Tan[e + f*x]])`

3.836.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 275, normalized size of antiderivative = 0.96, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.178$, Rules used = {3042, 4071, 87, 57, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c - ic \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c - ic \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(i \tan(e + fx)a + a)^{5/2}} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{87} \end{aligned}$$

3.836. $\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{7/2}}{(a + ia \tan(e + fx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{3ac(a+ia \tan(e+fx))^{3/2}} - \frac{(2A+5iB) \int \frac{(c-ic \tan(e+fx))^{5/2}}{(i \tan(e+fx)a+a)^{3/2}} d \tan(e+fx)}{3a} \right)}{f} \\
 & \quad \downarrow 57 \\
 & \frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{3ac(a+ia \tan(e+fx))^{3/2}} - \frac{(2A+5iB) \left(\frac{2i(c-ic \tan(e+fx))^{5/2}}{a \sqrt{a+ia \tan(e+fx)}} - \frac{5c \int \frac{(c-ic \tan(e+fx))^{3/2}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e+fx)}{a} \right)}{3a} \right)}{f} \\
 & \quad \downarrow 60 \\
 & \frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{3ac(a+ia \tan(e+fx))^{3/2}} - \frac{(2A+5iB) \left(\frac{2i(c-ic \tan(e+fx))^{5/2}}{a \sqrt{a+ia \tan(e+fx)}} - \frac{5c \left(\frac{3}{2} c \int \frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e+fx) - \frac{i \sqrt{a+ia \tan(e+fx)}(c-ic \tan(e+fx))}{2a} \right)}{a} \right)}{3a} \right)}{f} \\
 & \quad \downarrow 60 \\
 & \frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{3ac(a+ia \tan(e+fx))^{3/2}} - \frac{(2A+5iB) \left(\frac{2i(c-ic \tan(e+fx))^{5/2}}{a \sqrt{a+ia \tan(e+fx)}} - \frac{5c \left(\frac{3}{2} c \left(c \int \frac{1}{\sqrt{i \tan(e+fx)a+a} \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) - \frac{i \sqrt{a+ia \tan(e+fx)}}{a} \right) \right)}{a} \right)}{3a} \right)}{f} \\
 & \quad \downarrow 45 \\
 & \frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{3ac(a+ia \tan(e+fx))^{3/2}} - \frac{(2A+5iB) \left(\frac{2i(c-ic \tan(e+fx))^{5/2}}{a \sqrt{a+ia \tan(e+fx)}} - \frac{5c \left(\frac{3}{2} c \left(2c \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} - \frac{i \sqrt{a+ia \tan(e+fx)}}{a} \right) \right)}{a} \right)}{3a} \right)}{f} \\
 & \quad \downarrow 218
 \end{aligned}$$

3.836. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^{3/2}} dx$

$$ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{3ac(a+ia \tan(e+fx))^{3/2}} - \frac{(2A+5iB) \left(\frac{2i(c-ic \tan(e+fx))^{5/2}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{5c \left(\frac{3}{2}c \left(-\frac{2i\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right) - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a} \right)}{a} \right)}{3a} \right)}{f} \right)$$

```
input Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2))/(a + I*a*Tan[e + f*x])^(3/2), x]
```

```
output (a*c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(7/2))/(3*a*c*(a + I*a*Tan[e + f*x])^(3/2)) - ((2*A + (5*I)*B)*(((2*I)*(c - I*c*Tan[e + f*x])^(5/2))/(a*Sqrt[a + I*a*Tan[e + f*x]]) - (5*c*(((1/2*I)*Sqrt[a + I*a*Tan[e + f*x]]*(c - I*c*Tan[e + f*x])^(3/2))/a + (3*c*(((2*I)*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[a] - (I*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/a))/2)/a))/(3*a)))/f
```

3.836.3.1 Defintions of rubi rules used

```
rule 45 Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]
```

```
rule 57 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]
```

- rule 60 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.))*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.836.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 732 vs. 2(236) = 472.

Time = 0.38 (sec) , antiderivative size = 733, normalized size of antiderivative = 2.55

method	result
derivativedivides	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c^3 \left(-30iA \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) ac+225iB \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right)}{\dots}$
default	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c^3 \left(-30iA \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) ac+225iB \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right)}{\dots}$
parts	$\frac{A \sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c^3 \left(45i \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \tan(fx+e)^2 ac+3i \sqrt{ac} \right)}{\dots}$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^(3/2),x,m
method=_RETURNVERBOSE)
```

```
output 1/6/f*(-c*(I*tan(f*x+e)-1))^(1/2)*(a*(1+I*tan(f*x+e)))^(1/2)*c^3/a^2*(-30*
I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/
2))*a*c+225*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2
)))/(a*c)^(1/2))*a*c*tan(f*x+e)-114*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(
1/2)*tan(f*x+e)+185*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+
e)^2-30*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*
c)^(1/2))*a*c*tan(f*x+e)^3+6*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*
tan(f*x+e)^3+3*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^4-2
25*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1
/2))*a*c*tan(f*x+e)^2-21*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*
x+e)^3-75*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)
)/(a*c)^(1/2))*a*c*tan(f*x+e)^3+90*I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*
(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)^2+90*A*ln((a*c*tan(f*
x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e
)+74*A*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)*tan(f*x+e)^2-118*I*B*(a*c)^(
1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)+75*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*
c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c+279*B*(a*c)^(1/2)*(a*c*(1+tan(
f*x+e)^2))^(1/2)*tan(f*x+e)-46*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)
)/(a*c*(1+tan(f*x+e)^2))^(1/2)/(a*c)^(1/2)/(I-tan(f*x+e))^3
```

$$3.836. \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^{3/2}} dx$$

3.836.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 589 vs. $2(219) = 438$.

Time = 0.27 (sec) , antiderivative size = 589, normalized size of antiderivative = 2.05

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^{3/2}} dx = \frac{15 (a^2 f e^{(5i f x + 5i e)} + a^2 f e^{(3i f x + 3i e)}) \sqrt{\frac{(4 A^2 + 20i AB - 25 B^2)}{a^3 f^2}}}{1}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output `1/12*(15*(a^2*f*e^(5*I*f*x + 5*I*e) + a^2*f*e^(3*I*f*x + 3*I*e))*sqrt((4*A^2 + 20*I*A*B - 25*B^2)*c^7/(a^3*f^2))*log(4*(2*((2*I*A - 5*B)*c^3*e^(3*I*f*x + 3*I*e) + (2*I*A - 5*B)*c^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + (a^2*f*e^(2*I*f*x + 2*I*e) - a^2*f)*sqrt((4*A^2 + 20*I*A*B - 25*B^2)*c^7/(a^3*f^2)))/((2*I*A - 5*B)*c^3*e^(2*I*f*x + 2*I*e) + (2*I*A - 5*B)*c^3) - 15*(a^2*f*e^(5*I*f*x + 5*I*e) + a^2*f*e^(3*I*f*x + 3*I*e))*sqrt((4*A^2 + 20*I*A*B - 25*B^2)*c^7/(a^3*f^2))*log(4*(2*((2*I*A - 5*B)*c^3*e^(3*I*f*x + 3*I*e) + (2*I*A - 5*B)*c^3*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) - (a^2*f*e^(2*I*f*x + 2*I*e) - a^2*f)*sqrt((4*A^2 + 20*I*A*B - 25*B^2)*c^7/(a^3*f^2)))/((2*I*A - 5*B)*c^3*e^(2*I*f*x + 2*I*e) + (2*I*A - 5*B)*c^3) - 4*(15*(2*I*A - 5*B)*c^3*e^(6*I*f*x + 6*I*e) + 25*(2*I*A - 5*B)*c^3*e^(4*I*f*x + 4*I*e) + 8*(2*I*A - 5*B)*c^3*e^(2*I*f*x + 2*I*e) + 4*(-I*A + B)*c^3)*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(a^2*f*e^(5*I*f*x + 5*I*e) + a^2*f*e^(3*I*f*x + 3*I*e))`

3.836.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^{3/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(7/2)/(a+I*a*tan(f*x+e))**(3/2),x)`

3.836. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^{3/2}} dx$

output Timed out

3.836.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1373 vs. $2(219) = 438$.

Time = 0.83 (sec) , antiderivative size = 1373, normalized size of antiderivative = 4.78

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^{3/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output

```
-12*(60*(2*A + 5*I*B)*c^3*cos(6*f*x + 6*e) + 100*(2*A + 5*I*B)*c^3*cos(4*f
*x + 4*e) + 32*(2*A + 5*I*B)*c^3*cos(2*f*x + 2*e) + 60*(2*I*A - 5*B)*c^3*s
in(6*f*x + 6*e) + 100*(2*I*A - 5*B)*c^3*sin(4*f*x + 4*e) + 32*(2*I*A - 5*B
)*c^3*sin(2*f*x + 2*e) - 16*(A + I*B)*c^3 + 30*((2*A + 5*I*B)*c^3*cos(7/2*
arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 2*(2*A + 5*I*B)*c^3*cos(5/2
*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (2*A + 5*I*B)*c^3*cos(3/2*
arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (2*I*A - 5*B)*c^3*sin(7/2*a
rctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 2*(2*I*A - 5*B)*c^3*sin(5/2*
arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (2*I*A - 5*B)*c^3*sin(3/2*a
rctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*arctan2(cos(1/2*arctan2(sin(2
*f*x + 2*e), cos(2*f*x + 2*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f
*x + 2*e))) + 1) + 30*((2*A + 5*I*B)*c^3*cos(7/2*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e))) + 2*(2*A + 5*I*B)*c^3*cos(5/2*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e))) + (2*A + 5*I*B)*c^3*cos(3/2*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e))) + (2*I*A - 5*B)*c^3*sin(7/2*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e))) + 2*(2*I*A - 5*B)*c^3*sin(5/2*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e))) + (2*I*A - 5*B)*c^3*sin(3/2*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e))))*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x +
2*e))), -sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 1) + 15*((
2*I*A - 5*B)*c^3*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + ...
```

3.836.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^{3/2}} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{7/2}}{(i a \tan(fx + e) + a)^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(7/2)/(I*a*tan(f*x + e) + a)^(3/2), x)`

3.836.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^{3/2}} dx = \int \frac{(A + B \tan(e + fx)) (c - c \tan(e + fx) 1i)^{7/2}}{(a + a \tan(e + fx) 1i)^{3/2}} dx$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(7/2))/(a + a*tan(e + f*x)*1i)^(3/2), x)`

output `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(7/2))/(a + a*tan(e + f*x)*1i)^(3/2), x)`

3.837
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^{3/2}} dx$$

3.837.1 Optimal result	7520
3.837.2 Mathematica [C] (verified)	7521
3.837.3 Rubi [A] (verified)	7521
3.837.4 Maple [B] (verified)	7524
3.837.5 Fricas [B] (verification not implemented)	7525
3.837.6 Sympy [F]	7526
3.837.7 Maxima [F(-2)]	7526
3.837.8 Giac [F]	7527
3.837.9 Mupad [F(-1)]	7527

3.837.1 Optimal result

Integrand size = 45, antiderivative size = 229

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^{3/2}} dx =$$

$$\frac{2(iA - 4B)c^{5/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{a^{3/2}f}$$

$$- \frac{(iA - 4B)c^2 \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}}{a^2 f}$$

$$- \frac{2(iA - 4B)c(c - ic \tan(e + fx))^{3/2}}{3af \sqrt{a + ia \tan(e + fx)}} + \frac{(iA - B)(c - ic \tan(e + fx))^{5/2}}{3f(a + ia \tan(e + fx))^{3/2}}$$

output

```
-2*(I*A-4*B)*c^(5/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/a^(3/2)/f-(I*A-4*B)*c^2*(a+I*a*tan(f*x+e))^(1/2)*(c-I*c*tan(f*x+e))^(1/2)/a^2/f-2/3*(I*A-4*B)*c*(c-I*c*tan(f*x+e))^(3/2)/a/f/(a+I*a*tan(f*x+e))^(1/2)+1/3*(I*A-B)*(c-I*c*tan(f*x+e))^(5/2)/f/(a+I*a*tan(f*x+e))^(3/2)
```

3.837.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.78 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.62

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^{3/2}} dx = \frac{c^2 \cos^2(e + fx) \left(3B \sec^3(e + fx) (\cos(3(e + fx))) - i \sin(3(e + fx)) \right)}{(a + ia \tan(e + fx))^{3/2}}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2))/(a + I*a*Tan[e + f*x])^(3/2),x]`

output `(c^2*Cos[e + f*x]^2*(3*B*Sec[e + f*x]^3*(Cos[3*(e + f*x)] - I*Sin[3*(e + f*x)]) + (4*I)*(A + (4*I)*B)*Hypergeometric2F1[-3/2, -3/2, -1/2, (1 + I*Tan[e + f*x])/2]*Sqrt[2 - (2*I)*Tan[e + f*x]])*Sqrt[c - I*c*Tan[e + f*x]]/(3*a*f*Sqrt[a + I*a*Tan[e + f*x]])`

3.837.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 224, normalized size of antiderivative = 0.98, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 87, 57, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c - ic \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c - ic \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(i \tan(e + fx)a + a)^{5/2}} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{87} \end{aligned}$$

3.837. $\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{3ac(a+ia \tan(e+fx))^{3/2}} - \frac{(A+4iB) \int \frac{(c-ic \tan(e+fx))^{3/2}}{(i \tan(e+fx)a+a)^{3/2}} d \tan(e+fx)}{3a} \right)}{f} \\
 & \quad \downarrow 57 \\
 & \frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{3ac(a+ia \tan(e+fx))^{3/2}} - \frac{(A+4iB) \left(\frac{2i(c-ic \tan(e+fx))^{3/2}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{3c \int \frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e+fx)}{a} \right)}{3a} \right)}{f} \\
 & \quad \downarrow 60 \\
 & \frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{3ac(a+ia \tan(e+fx))^{3/2}} - \frac{(A+4iB) \left(\frac{2i(c-ic \tan(e+fx))^{3/2}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{3c \left(c \int \frac{1}{\sqrt{i \tan(e+fx)a+a} \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx) - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a} \right)}{a} \right)}{3a} \right)}{f} \\
 & \quad \downarrow 45 \\
 & \frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{3ac(a+ia \tan(e+fx))^{3/2}} - \frac{(A+4iB) \left(\frac{2i(c-ic \tan(e+fx))^{3/2}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{3c \left(2c \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a} \right)}{a} \right)}{3a} \right)}{f} \\
 & \quad \downarrow 218 \\
 & \frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{3ac(a+ia \tan(e+fx))^{3/2}} - \frac{(A+4iB) \left(\frac{2i(c-ic \tan(e+fx))^{3/2}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{3c \left(-\frac{2i\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{\sqrt{a}} - \frac{i\sqrt{a+ia \tan(e+fx)}\sqrt{c-ic \tan(e+fx)}}{a} \right)}{a} \right)}{3a} \right)}{f}
 \end{aligned}$$

3.837. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^{3/2}} dx$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2))/(a + I*a*Tan[e + f*x])^(3/2),x]`

output `(a*c*((I*A - B)*(c - I*c*Tan[e + f*x])^(5/2))/(3*a*c*(a + I*a*Tan[e + f*x])^(3/2)) - ((A + (4*I)*B)*((2*I)*(c - I*c*Tan[e + f*x])^(3/2))/(a*Sqrt[a + I*a*Tan[e + f*x]]) - (3*c*((-2*I)*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[a] - (I*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])/a)/a)/(3*a))/f`

3.837.3.1 Defintions of rubi rules used

rule 45 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 57 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

rule 60 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.837.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 668 vs. $2(189) = 378$.

Time = 0.41 (sec) , antiderivative size = 669, normalized size of antiderivative = 2.92

method	result
derivativedivides	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c^2 \left(-12iB \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) ac \tan(fx+e)^3 + 9iA}{}$
default	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c^2 \left(-12iB \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) ac \tan(fx+e)^3 + 9iA}{}$
parts	$\frac{A \sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c^2 \left(9i \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \tan(fx+e)^2 ac - 3 \ln \left(\frac{a}{} \right)}{}$

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^(3/2),x,m
method=_RETURNVERBOSE)`

$$3.837. \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^{3/2}} dx$$

```
output 1/3/f*(-c*(I*tan(f*x+e)-1))^(1/2)*(a*(1+I*tan(f*x+e)))^(1/2)*c^2/a^2*(-12*
I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/
2))*a*c*tan(f*x+e)^3+9*I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+
e)^2))^(1/2)))/(a*c)^(1/2))*a*c*tan(f*x+e)^2-3*A*ln((a*c*tan(f*x+e)+(a*c)^(
1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c*tan(f*x+e)^3+36*I*B*ln
((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a
c*tan(f*x+e)+29*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2-
36*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1
/2))*a*c*tan(f*x+e)^2-3*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x
+e)^3-3*I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(
a*c)^(1/2))*a*c-12*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e
+9*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1
/2))*a*c*tan(f*x+e)+8*A*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)*tan(f*x+e
)^2-19*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)+12*B*ln((a*c*tan(f*x+e
)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c+45*B*(a*c)^(1
/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-4*A*(a*c)^(1/2)*(a*c*(1+tan(f*
x+e)^2))^(1/2))/(a*c*(1+tan(f*x+e)^2))^(1/2)/(a*c)^(1/2)/(I-tan(f*x+e))^3
```

3.837.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 511 vs. 2(175) = 350.

Time = 0.27 (sec) , antiderivative size = 511, normalized size of antiderivative = 2.23

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(a + i a \tan(e + fx))^{3/2}} dx = \frac{\left(3 a^2 \sqrt{\frac{(A^2 + 8i AB - 16 B^2)c^5}{a^3 f^2}} f e^{(3i fx + 3i e)} \log \left(- \frac{4 \left(2((i A - 4) \right)}{\dots} \right) \right)}{\dots}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^(3/
2),x, algorithm="fricas")
```

output $\frac{1}{6} \cdot (3a^2 \sqrt{(A^2 + 8IAB - 16B^2)} c^5 / (a^3 f^2)) f e^{(3Ifx + 3Ie)} \cdot \log(-4 \cdot (2 \cdot ((IA - 4B) c^2 e^{(3Ifx + 3Ie)} + (IA - 4B) c^2 e^{(Ifx + Ie)}) \sqrt{a / (e^{(2Ifx + 2Ie)} + 1)}) \sqrt{c / (e^{(2Ifx + 2Ie)} + 1)}) + (a^2 f e^{(2Ifx + 2Ie)} - a^2 f) \sqrt{(A^2 + 8IAB - 16B^2) c^5 / (a^3 f^2)}) / ((-IA + 4B) c^2 e^{(2Ifx + 2Ie)} + (-IA + 4B) c^2) - 3a^2 \sqrt{(A^2 + 8IAB - 16B^2)} c^5 / (a^3 f^2)) f e^{(3Ifx + 3Ie)} \cdot \log(-4 \cdot (2 \cdot ((IA - 4B) c^2 e^{(3Ifx + 3Ie)} + (IA - 4B) c^2 e^{(Ifx + Ie)}) \sqrt{a / (e^{(2Ifx + 2Ie)} + 1)}) \sqrt{c / (e^{(2Ifx + 2Ie)} + 1)}) - (a^2 f e^{(2Ifx + 2Ie)} - a^2 f) \sqrt{(A^2 + 8IAB - 16B^2) c^5 / (a^3 f^2)}) / ((-IA + 4B) c^2 e^{(2Ifx + 2Ie)} + (-IA + 4B) c^2) - 4 \cdot (3 \cdot (IA - 4B) c^2 e^{(4Ifx + 4Ie)} + 2 \cdot (IA - 4B) c^2 e^{(2Ifx + 2Ie)} + (-IA + B) c^2) \sqrt{a / (e^{(2Ifx + 2Ie)} + 1)}) \sqrt{c / (e^{(2Ifx + 2Ie)} + 1)}) e^{(-3Ifx - 3Ie)} / (a^2 f)$

3.837.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^{3/2}} dx = \int \frac{(-ic(\tan(e + fx) + i))^{5/2} (A + B \tan(e + fx))}{(ia(\tan(e + fx) - i))^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(5/2)/(a+I*a*tan(f*x+e))**(3/2),x)`

output `Integral((-I*c*(tan(e + f*x) + I))**(5/2)*(A + B*tan(e + f*x))/(I*a*(tan(e + f*x) - I))**(3/2), x)`

3.837.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is not defined.`

3.837. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^{3/2}} dx$

3.837.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(a + i a \tan(e + fx))^{3/2}} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{5/2}}{(i a \tan(fx + e) + a)^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(5/2)/(I*a*tan(f*x + e) + a)^(3/2), x)`

3.837.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(a + i a \tan(e + fx))^{3/2}} dx = \int \frac{(A + B \tan(e + fx)) (c - c \tan(e + fx) li)^{5/2}}{(a + a \tan(e + fx) li)^{3/2}} dx$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(5/2))/(a + a*tan(e + f*x)*1i)^(3/2), x)`

output `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(5/2))/(a + a*tan(e + f*x)*1i)^(3/2), x)`

3.838
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^{3/2}} dx$$

3.838.1 Optimal result	7528
3.838.2 Mathematica [A] (verified)	7528
3.838.3 Rubi [A] (verified)	7529
3.838.4 Maple [B] (verified)	7531
3.838.5 Fricas [B] (verification not implemented)	7532
3.838.6 Sympy [F]	7533
3.838.7 Maxima [A] (verification not implemented)	7533
3.838.8 Giac [F]	7534
3.838.9 Mupad [F(-1)]	7534

3.838.1 Optimal result

Integrand size = 45, antiderivative size = 157

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^{3/2}} dx = \frac{2Bc^{3/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{a^{3/2} f} + \frac{2Bc\sqrt{c-ic \tan(e+fx)}}{af\sqrt{a+ia \tan(e+fx)}} + \frac{(iA - B)(c - ic \tan(e + fx))^{3/2}}{3f(a + ia \tan(e + fx))^{3/2}}$$

output `2*B*c^(3/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/a^(3/2)/f+2*B*c*(c-I*c*tan(f*x+e))^(1/2)/a/f/(a+I*a*tan(f*x+e))^(1/2)+1/3*(I*A-B)*(c-I*c*tan(f*x+e))^(3/2)/f/(a+I*a*tan(f*x+e))^(3/2)`

3.838.2 Mathematica [A] (verified)

Time = 5.23 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.87

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^{3/2}} dx = \frac{c\sqrt{c - ic \tan(e + fx)} \left(\frac{6B \arcsin\left(\frac{\sqrt{a+ia \tan(e+fx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{1-i \tan(e+fx)}} + \frac{\sqrt{a}(A-iB)}{(-i+ia \tan(e+fx))} \right)}{3a^{3/2} f}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2))/(a + I*a*Tan[e + f*x])^(3/2),x]`

output $(c*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]]*((6*B*\text{ArcSin}[\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]/(\text{Sqrt}[2]*\text{Sqrt}[a])])/\text{Sqrt}[1 - I*\text{Tan}[e + f*x]] + (\text{Sqrt}[a]*(A - (5*I)*B + ((-I)*A + 7*B)*\text{Tan}[e + f*x]))/((-I + \text{Tan}[e + f*x])*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]])))/ (3*a^{(3/2)*f})$

3.838.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.08, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 87, 57, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{(c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx$$

↓ 4071

$$ac \int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{(i \tan(e + fx) a + a)^{5/2}} d \tan(e + fx)$$

f
↓ 87

$$ac \left(\frac{(-B + iA)(c - ic \tan(e + fx))^{3/2}}{3ac(a + ia \tan(e + fx))^{3/2}} - \frac{iB \int \frac{\sqrt{c - ic \tan(e + fx)}}{(i \tan(e + fx) a + a)^{3/2}} d \tan(e + fx)}{a} \right)$$

f
↓ 57

$$ac \left(\frac{(-B + iA)(c - ic \tan(e + fx))^{3/2}}{3ac(a + ia \tan(e + fx))^{3/2}} - \frac{iB \left(\frac{2i \sqrt{c - ic \tan(e + fx)}}{a \sqrt{a + ia \tan(e + fx)}} - \frac{c \int \frac{1}{\sqrt{i \tan(e + fx) a + a} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{a} \right)}{a} \right)$$

f
↓ 45

3.838. $\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^{3/2}} dx$

$$\begin{array}{c}
 \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{3/2}}{3ac(a+ia \tan(e+fx))^{3/2}} - \frac{iB \left(\frac{2i\sqrt{c-ic \tan(e+fx)}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{2c \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}} \right)}{a} \right) \\
 \hline
 f \\
 \downarrow 218 \\
 \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{3/2}}{3ac(a+ia \tan(e+fx))^{3/2}} - \frac{iB \left(\frac{2i\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{a^{3/2}} + \frac{2i\sqrt{c-ic \tan(e+fx)}}{a\sqrt{a+ia \tan(e+fx)}} \right)}{a} \right) \\
 \hline
 f
 \end{array}$$

input `Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2))/(a + I*a*Tan[e + f*x])^(3/2), x]`

output `(a*c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(3/2))/(3*a*c*(a + I*a*Tan[e + f*x])^(3/2)) - (I*B*(((2*I)*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/a^(3/2) + ((2*I)*Sqrt[c - I*c*Tan[e + f*x]])/(a*Sqrt[a + I*a*Tan[e + f*x]])))/a)/f`

3.838.3.1 Defintions of rubi rules used

rule 45 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]`

rule 57 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))*Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.838.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 407 vs. 2(128) = 256.

Time = 0.34 (sec) , antiderivative size = 408, normalized size of antiderivative = 2.60

method	result
derivativedivides	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c \left(-3iB \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right) a c \tan(fx+e)^3 + 9iB \ln \dots}{\dots}$
default	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c \left(-3iB \ln \left(\frac{ac \tan(fx+e)+\sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right) a c \tan(fx+e)^3 + 9iB \ln \dots}{\dots}$
parts	$\frac{A \sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c (1+\tan(fx+e)^2)}{3f a^2 (i - \tan(fx+e))^3} + \frac{iB \sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c \left(9 \dots \right)}{\dots}$

3.838.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^{3/2}} dx$$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
```

```
output 1/3/f*(-c*(I*tan(f*x+e)-1))^(1/2)*(a*(1+I*tan(f*x+e)))^(1/2)/a^2*c*(-3*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)^3+9*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)+7*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2-9*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(f*x+e)^2+A*(a*c*(1+tan(f*x+e)^2))^(1/2)*(a*c)^(1/2)*tan(f*x+e)^2-5*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)+3*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c+12*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)+A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c*(1+tan(f*x+e)^2))^(1/2)/(I-tan(f*x+e))^3/(a*c)^(1/2)
```

3.838.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 400 vs. $2(119) = 238$.

Time = 0.28 (sec) , antiderivative size = 400, normalized size of antiderivative = 2.55

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^{3/2}} dx =$$

$$\left(3a^2 f \sqrt{-\frac{B^2 c^3}{a^3 f^2}} e^{(3i fx + 3ie)} \log \left(\frac{4 \left(2 (Bce^{(3i fx + 3ie)} + Bce^{(i fx + ie)}) \sqrt{\frac{a}{e^{(2i fx + 2ie)} + 1}} \sqrt{\frac{c}{e^{(2i fx + 2ie)} + 1}} + (a^2 f e^{(2i fx + 2ie)} - a^2 f) \sqrt{-\frac{Bce^{(2i fx + 2ie)} + Bc}}{Bce^{(2i fx + 2ie)} + Bc}} \right)} \right)$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="fricas")
```

```
output -1/6*(3*a^2*f*sqrt(-B^2*c^3/(a^3*f^2))*e^(3*I*f*x + 3*I*e)*log(4*(2*(B*c*e
^(3*I*f*x + 3*I*e) + B*c*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1)
)*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + (a^2*f*e^(2*I*f*x + 2*I*e) - a^2*f)*
sqrt(-B^2*c^3/(a^3*f^2)))/(B*c*e^(2*I*f*x + 2*I*e) + B*c)) - 3*a^2*f*sqrt(
-B^2*c^3/(a^3*f^2))*e^(3*I*f*x + 3*I*e)*log(4*(2*(B*c*e^(3*I*f*x + 3*I*e)
+ B*c*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*
x + 2*I*e) + 1)) - (a^2*f*e^(2*I*f*x + 2*I*e) - a^2*f)*sqrt(-B^2*c^3/(a^3*
f^2)))/(B*c*e^(2*I*f*x + 2*I*e) + B*c)) - 2*(6*B*c*e^(4*I*f*x + 4*I*e) - (
-I*A - 5*B)*c*e^(2*I*f*x + 2*I*e) - (-I*A + B)*c)*sqrt(a/(e^(2*I*f*x + 2*I
*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*e^(-3*I*f*x - 3*I*e)/(a^2*f)
```

3.838.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^{3/2}} dx = \int \frac{(-i c (\tan(e + fx) + i))^{3/2} (A + B \tan(e + fx))}{(i a (\tan(e + fx) - i))^{3/2}} dx$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(3/2)/(a+I*a*tan(f*x+e))**(
3/2),x)
```

```
output Integral((-I*c*(tan(e + f*x) + I))**(3/2)*(A + B*tan(e + f*x))/(I*a*(tan(e
+ f*x) - I))**(3/2), x)
```

3.838.7 Maxima [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.07

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^{3/2}} dx = \frac{(6 B c \arctan(\cos(fx + e), \sin(fx + e) + 1) + 6 B c \arctan(\cos(fx + e), \sin(fx + e) - 1) - 2*(-I*A + B)*c*\cos(3*f*x + 3*e) + 12*B*c*\cos(f*x + e) + 3*I*B*c*\log(\cos(f*x + e)^2 + \sin(f*x + e)^2 + 2*\sin(f*x + e) + 1) - 3*I*B*c*\log(\cos(f*x + e)^2 + \sin(f*x + e)^2 - 2*\sin(f*x + e) + 1) + 2*(A + I*B)*c*\sin(3*f*x + 3*e) - 12*I*B*c*\sin(f*x + e))*sqrt(c)/(a^(3/2)*f)}{1}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^(3/
2),x, algorithm="maxima")
```

```
output 1/6*(6*B*c*arctan2(cos(f*x + e), sin(f*x + e) + 1) + 6*B*c*arctan2(cos(f*x
+ e), -sin(f*x + e) + 1) - 2*(-I*A + B)*c*cos(3*f*x + 3*e) + 12*B*c*cos(f
*x + e) + 3*I*B*c*log(cos(f*x + e)^2 + sin(f*x + e)^2 + 2*sin(f*x + e) + 1
) - 3*I*B*c*log(cos(f*x + e)^2 + sin(f*x + e)^2 - 2*sin(f*x + e) + 1) + 2*
(A + I*B)*c*sin(3*f*x + 3*e) - 12*I*B*c*sin(f*x + e))*sqrt(c)/(a^(3/2)*f)
```

3.838. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^{3/2}} dx$

3.838.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^{3/2}} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{3/2}}{(i a \tan(fx + e) + a)^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(3/2)/(I*a*tan(f*x + e) + a)^(3/2), x)`

3.838.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^{3/2}} dx = \int \frac{(A + B \tan(e + fx)) (c - c \tan(e + fx) 1i)^{3/2}}{(a + a \tan(e + fx) 1i)^{3/2}} dx$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(3/2))/(a + a*tan(e + f*x)*1i)^(3/2), x)`

output `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(3/2))/(a + a*tan(e + f*x)*1i)^(3/2), x)`

3.839
$$\int \frac{(A+B \tan(e+fx))\sqrt{c-ictan(e+fx)}}{(a+ia \tan(e+fx))^{3/2}} dx$$

3.839.1 Optimal result	7535
3.839.2 Mathematica [A] (verified)	7535
3.839.3 Rubi [A] (verified)	7536
3.839.4 Maple [A] (verified)	7537
3.839.5 Fricas [A] (verification not implemented)	7538
3.839.6 Sympy [F]	7538
3.839.7 Maxima [F(-2)]	7539
3.839.8 Giac [F]	7539
3.839.9 Mupad [B] (verification not implemented)	7539

3.839.1 Optimal result

Integrand size = 45, antiderivative size = 104

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)}}{(a + ia \tan(e + fx))^{3/2}} dx = \frac{(iA - B)\sqrt{c - ictan(e + fx)}}{3f(a + ia \tan(e + fx))^{3/2}} + \frac{(iA + 2B)\sqrt{c - ictan(e + fx)}}{3af\sqrt{a + ia \tan(e + fx)}}$$

output `1/3*(I*A+2*B)*(c-I*c*tan(f*x+e))^(1/2)/a/f/(a+I*a*tan(f*x+e))^(1/2)+1/3*(I*A-B)*(c-I*c*tan(f*x+e))^(1/2)/f/(a+I*a*tan(f*x+e))^(3/2)`

3.839.2 Mathematica [A] (verified)

Time = 3.02 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.78

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)}}{(a + ia \tan(e + fx))^{3/2}} dx = \frac{(2A - iB + (iA + 2B) \tan(e + fx))\sqrt{c - ictan(e + fx)}}{3af(-i + \tan(e + fx))\sqrt{a + ia \tan(e + fx)}}$$

input `Integrate[((A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])/(a + I*a*Tan[e + f*x])^(3/2),x]`

output `((2*A - I*B + (I*A + 2*B)*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])/(3*a*f*(-I + Tan[e + f*x])*Sqrt[a + I*a*Tan[e + f*x]])`

3.839.
$$\int \frac{(A+B \tan(e+fx))\sqrt{c-ictan(e+fx)}}{(a+ia \tan(e+fx))^{3/2}} dx$$

3.839.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.089$, Rules used = {3042, 4071, 87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{\sqrt{c - ic \tan(e + fx)}(A + B \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx \\
 \downarrow 3042 \\
 \int \frac{\sqrt{c - ic \tan(e + fx)}(A + B \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx \\
 \downarrow 4071 \\
 ac \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx)a + a)^{5/2} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) \\
 \downarrow 87 \\
 ac \left(\frac{(A - 2iB) \int \frac{1}{(i \tan(e + fx)a + a)^{3/2} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{3a} + \frac{(-B + iA) \sqrt{c - ic \tan(e + fx)}}{3ac(a + ia \tan(e + fx))^{3/2}} \right) \\
 \downarrow 48 \\
 ac \left(\frac{i(A - 2iB) \sqrt{c - ic \tan(e + fx)}}{3a^2 c \sqrt{a + ia \tan(e + fx)}} + \frac{(-B + iA) \sqrt{c - ic \tan(e + fx)}}{3ac(a + ia \tan(e + fx))^{3/2}} \right)
 \end{array}$$

input `Int[((A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])/(a + I*a*Tan[e + f*x])^(3/2),x]`

output `(a*c*(((I*A - B)*Sqrt[c - I*c*Tan[e + f*x]])/(3*a*c*(a + I*a*Tan[e + f*x])^(3/2)) + ((I/3)*(A - (2*I)*B)*Sqrt[c - I*c*Tan[e + f*x]])/(a^2*c*Sqrt[a + I*a*Tan[e + f*x]])))/f`

3.839.3.1 Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*(c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.839.4 Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.99

method	result
derivativedivides	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} (2iB \tan(fx+e)^2+3iA \tan(fx+e)-A \tan(fx+e)^2-iB+3B \tan(fx+e)+2A)}{3f a^2(i-\tan(fx+e))^3}$
default	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} (2iB \tan(fx+e)^2+3iA \tan(fx+e)-A \tan(fx+e)^2-iB+3B \tan(fx+e)+2A)}{3f a^2(i-\tan(fx+e))^3}$
parts	$\frac{A\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} (3i \tan(fx+e)-\tan(fx+e)^2+2)}{3f a^2(i-\tan(fx+e))^3} - \frac{iB\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))}}{3f a^2(i-\tan(fx+e))^3}$

```
input int((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2),x,m
ethod=_RETURNVERBOSE)
```

$$3.839. \int \frac{(A+B \tan(e+fx))\sqrt{c-ic \tan(e+fx)}}{(a+ia \tan(e+fx))^{3/2}} dx$$

output $\frac{1}{3}f(-c(I\tan(fx+e)-1))^{1/2}(a(1+I\tan(fx+e)))^{1/2}/a^2(2IB\tan(fx+e)^2+3IA\tan(fx+e)-A\tan(fx+e)^2-I*B+3B\tan(fx+e)+2A)/(I-\tan(fx+e))^3$

3.839.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.88

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - ic \tan(e + fx)}}{(a + ia \tan(e + fx))^{3/2}} dx = \frac{(3(-iA - B)e^{(4ifx+4ie)} + 2(-2iA - B)e^{(2ifx+2ie)} - iA + B)\sqrt{\frac{a}{e^{(2ifx+2ie)}+1}}\sqrt{\frac{c}{e^{(2ifx+2ie)}+1}}e^{(-3ifx-3ie)}}{6a^2f}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="fracas")`

output $-1/6*(3*(-I*A - B)*e^{(4*I*f*x + 4*I*e)} + 2*(-2*I*A - B)*e^{(2*I*f*x + 2*I*e)} - I*A + B)*\text{sqrt}(a/(e^{(2*I*f*x + 2*I*e)} + 1))*\text{sqrt}(c/(e^{(2*I*f*x + 2*I*e)} + 1))*e^{(-3*I*f*x - 3*I*e)}/(a^2*f)$

3.839.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - ic \tan(e + fx)}}{(a + ia \tan(e + fx))^{3/2}} dx = \int \frac{\sqrt{-ic(\tan(e + fx) + i)}(A + B \tan(e + fx))}{(ia(\tan(e + fx) - i))^{3/2}} dx$$

input `integrate((c-I*c*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**(3/2),x)`

output `Integral(sqrt(-I*c*(tan(e + f*x) + I))*(A + B*tan(e + f*x))/(I*a*(tan(e + f*x) - I))**(3/2), x)`

3.839.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{(a + i a \tan(e + fx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.839.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{(a + i a \tan(e + fx))^{3/2}} dx = \int \frac{(B \tan(fx + e) + A) \sqrt{-i c \tan(fx + e) + c}}{(i a \tan(fx + e) + a)^{3/2}} dx$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*sqrt(-I*c*tan(f*x + e) + c)/(I*a*tan(f*x + e) + a)^(3/2), x)`

3.839.9 Mupad [B] (verification not implemented)

Time = 9.69 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.88

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{(a + i a \tan(e + fx))^{3/2}} dx = \sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}} \sqrt{\frac{c(\cos(2e+2fx)+1-\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}}$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(1/2))/(a + a*tan(e + f*x)*1i)^(3/2),x)`

output $((a \cdot (\cos(2e + 2fx) + \sin(2e + 2fx) \cdot i + 1)) / (\cos(2e + 2fx) + 1))^{1/2} \cdot ((c \cdot (\cos(2e + 2fx) - \sin(2e + 2fx) \cdot i + 1)) / (\cos(2e + 2fx) + 1))^{1/2} \cdot (A \cdot 3i + 3B + A \cdot \cos(2e + 2fx) \cdot 4i + A \cdot \cos(4e + 4fx) \cdot i + 2B \cdot \cos(2e + 2fx) - B \cdot \cos(4e + 4fx) + 4A \cdot \sin(2e + 2fx) + A \cdot \sin(4e + 4fx) - B \cdot \sin(2e + 2fx) \cdot 2i + B \cdot \sin(4e + 4fx) \cdot i)) / (12 \cdot a^2 \cdot f)$

3.840 $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}} dx$

3.840.1 Optimal result 7541
 3.840.2 Mathematica [A] (verified) 7541
 3.840.3 Rubi [A] (verified) 7542
 3.840.4 Maple [A] (verified) 7544
 3.840.5 Fricas [A] (verification not implemented) 7544
 3.840.6 Sympy [F] 7545
 3.840.7 Maxima [F(-2)] 7545
 3.840.8 Giac [F] 7546
 3.840.9 Mupad [B] (verification not implemented) 7546

3.840.1 Optimal result

Integrand size = 45, antiderivative size = 152

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} dx =$$

$$\frac{iA + B}{f(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} + \frac{(2iA + B) \sqrt{c - ic \tan(e + fx)}}{3cf(a + ia \tan(e + fx))^{3/2}} + \frac{(2iA + B) \sqrt{c - ic \tan(e + fx)}}{3acf \sqrt{a + ia \tan(e + fx)}}$$

output `1/3*(2*I*A+B)*(c-I*c*tan(f*x+e))^(1/2)/a/c/f/(a+I*a*tan(f*x+e))^(1/2)+(-I*A-B)/f/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^(3/2)+1/3*(2*I*A+B)*(c-I*c*tan(f*x+e))^(1/2)/c/f/(a+I*a*tan(f*x+e))^(3/2)`

3.840.2 Mathematica [A] (verified)

Time = 3.17 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.64

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} dx = \frac{A + iB + (-2iA - B) \tan(e + fx) + (2A - iB) \tan^2(e + fx)}{3af(-i + \tan(e + fx)) \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^(3/2)*Sqrt[c - I*c*Tan[e + f*x]]),x]`

output (A + I*B + ((-2*I)*A - B)*Tan[e + f*x] + (2*A - I*B)*Tan[e + f*x]^2)/(3*a*f*(-I + Tan[e + f*x])*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])

3.840.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.07, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4071, 87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} dx$$

↓ 4071

$$\frac{ac \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx)a + a)^{5/2} (c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{f}$$

↓ 87

$$ac \left(\frac{(2A - iB) \int \frac{1}{(i \tan(e + fx)a + a)^{5/2} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{c} - \frac{B + iA}{ac(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} \right)$$

↓ 55

$$ac \left(\frac{(2A - iB) \left(\frac{\int \frac{1}{(i \tan(e + fx)a + a)^{3/2} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{3a} + \frac{i \sqrt{c - ic \tan(e + fx)}}{3ac(a + ia \tan(e + fx))^{3/2}} \right)}{c} - \frac{B + iA}{ac(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} \right)$$

↓ 48

$$ac \left(\frac{(2A - iB) \left(\frac{i \sqrt{c - ic \tan(e + fx)}}{3a^2 c \sqrt{a + ia \tan(e + fx)}} + \frac{i \sqrt{c - ic \tan(e + fx)}}{3ac(a + ia \tan(e + fx))^{3/2}} \right)}{c} - \frac{B + iA}{ac(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} \right)$$

3.840. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} dx$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^(3/2)*Sqrt[c - I*c*Tan[e + f*x]]),x]`

output `(a*c*(-((I*A + B)/(a*c*(a + I*a*Tan[e + f*x])^(3/2)*Sqrt[c - I*c*Tan[e + f*x]])) + ((2*A - I*B)*((I/3)*Sqrt[c - I*c*Tan[e + f*x]])/(a*c*(a + I*a*Tan[e + f*x])^(3/2)) + ((I/3)*Sqrt[c - I*c*Tan[e + f*x]]/(a^2*c*Sqrt[a + I*a*Tan[e + f*x]]))))/c)/f`

3.840.3.1 Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.840.4 Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{i\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} (2iA \tan(fx+e)^4 - iB \tan(fx+e)^3 + B \tan(fx+e)^4 + 3iA \tan(fx+e)^2 + 2A \tan(fx+e))}{3f a^2 c (i - \tan(fx+e))^3 (i + \tan(fx+e))^2}$
default	$\frac{i\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} (2iA \tan(fx+e)^4 - iB \tan(fx+e)^3 + B \tan(fx+e)^4 + 3iA \tan(fx+e)^2 + 2A \tan(fx+e))}{3f a^2 c (i - \tan(fx+e))^3 (i + \tan(fx+e))^2}$
parts	$\frac{A\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} (2i \tan(fx+e)^3 - 2 \tan(fx+e)^4 + 2i \tan(fx+e) - 3 \tan(fx+e)^2 - 1)}{3f a^2 c (i - \tan(fx+e))^3 (i + \tan(fx+e))^2} - \frac{iB \tan(fx+e)}{3f a^2 c (i - \tan(fx+e))^3 (i + \tan(fx+e))^2}$

```
input int((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^(3/2),x,m
ethod=_RETURNVERBOSE)
```

```
output 1/3*I/f*(-c*(I*tan(f*x+e)-1))^(1/2)*(a*(1+I*tan(f*x+e)))^(1/2)/a^2/c*(2*I*
A*tan(f*x+e)^4-I*B*tan(f*x+e)^3+B*tan(f*x+e)^4+3*I*A*tan(f*x+e)^2+2*A*tan(
f*x+e)^3-I*B*tan(f*x+e)+I*A+2*A*tan(f*x+e)-B)/(I-tan(f*x+e))^3/(I+tan(f*x+
e))^2
```

3.840.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.95

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} \sqrt{c - ictan(e + fx)}} dx = \frac{(3(iA + B)e^{(6i fx + 6ie)} + 4(iA - B)e^{(5i fx + 5ie)} + 3(-iA + B)e^{(4i fx + 4ie)} + 4(iA - B)e^{(3i fx + 3ie)} - (7iA - B)e^{(2i fx + 2ie)} + 3(-iA + B)e^{(i fx + ie)} + 3A)}{12 a^2 c f}$$

```
input integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^(3/
2),x, algorithm="fracas")
```

3.840.
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{3/2} \sqrt{c-ictan(e+fx)}} dx$$

output $-1/12*(3*(I*A + B)*e^{(6*I*f*x + 6*I*e)} + 4*(I*A - B)*e^{(5*I*f*x + 5*I*e)} + 3*(-I*A + B)*e^{(4*I*f*x + 4*I*e)} + 4*(I*A - B)*e^{(3*I*f*x + 3*I*e)} - (7*I*A - B)*e^{(2*I*f*x + 2*I*e)} - I*A + B)*\text{sqrt}(a/(e^{(2*I*f*x + 2*I*e)} + 1))*\text{sqrt}(c/(e^{(2*I*f*x + 2*I*e)} + 1))*e^{(-3*I*f*x - 3*I*e)}/(a^2*c*f)$

3.840.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} dx = \int \frac{A + B \tan(e + fx)}{(ia (\tan(e + fx) - i))^{\frac{3}{2}} \sqrt{-ic (\tan(e + fx) + i)}} dx$$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(1/2)/(a+I*a*tan(f*x+e))**(3/2),x)`

output `Integral((A + B*tan(e + f*x))/((I*a*(tan(e + f*x) - I))**(3/2)*sqrt(-I*c*(tan(e + f*x) + I))), x)`

3.840.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.840.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} dx = \int \frac{B \tan(fx + e) + A}{(ia \tan(fx + e) + a)^{3/2} \sqrt{-ic \tan(fx + e) + c}} dx$$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)^(3/2)*sqrt(-I*c*tan(f*x + e) + c)), x)`

3.840.9 Mupad [B] (verification not implemented)

Time = 9.36 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.12

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} dx = \frac{\sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)i)}{\cos(2e+2fx)+1}}}{(6A \sin(2e + 2fx) -$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^(3/2)*(c - c*tan(e + f*x)*1i)^(1/2)),x)`

output `((a*cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1)/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*cos(2*e + 2*f*x)*6i - 3*B - A*3i + A*cos(4*e + 4*f*x)*1i - B*cos(4*e + 4*f*x) + 6*A*sin(2*e + 2*f*x) + A*sin(4*e + 4*f*x) + B*sin(4*e + 4*f*x)*1i)/(12*a^2*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

$$3.841 \quad \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{3/2}(c-ict \tan(e+fx))^{3/2}} dx$$

3.841.1 Optimal result	7547
3.841.2 Mathematica [A] (verified)	7547
3.841.3 Rubi [A] (verified)	7548
3.841.4 Maple [A] (verified)	7550
3.841.5 Fricas [A] (verification not implemented)	7550
3.841.6 Sympy [F]	7551
3.841.7 Maxima [A] (verification not implemented)	7551
3.841.8 Giac [F]	7552
3.841.9 Mupad [B] (verification not implemented)	7552

3.841.1 Optimal result

Integrand size = 45, antiderivative size = 152

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2}(c - ict \tan(e + fx))^{3/2}} dx = \frac{-iA - B}{3f(a + ia \tan(e + fx))^{3/2}(c - ict \tan(e + fx))^{3/2}} + \frac{iA}{3cf(a + ia \tan(e + fx))^{3/2}\sqrt{c - ict \tan(e + fx)}} + \frac{2A \tan(e + fx)}{3acf\sqrt{a + ia \tan(e + fx)}\sqrt{c - ict \tan(e + fx)}}$$

```
output 2/3*A*tan(f*x+e)/a/c/f/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(1/2)+1
/3*I*A/c/f/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^(3/2)+1/3*(-I*A-B)/
f/(a+I*a*tan(f*x+e))^(3/2)/(c-I*c*tan(f*x+e))^(3/2)
```

3.841.2 Mathematica [A] (verified)

Time = 5.92 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.52

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2}(c - ict \tan(e + fx))^{3/2}} dx = \frac{\cos^2(e + fx) (-B + 3A \tan(e + fx) + 2A \tan^3(e + fx))}{3acf\sqrt{a + ia \tan(e + fx)}\sqrt{c - ict \tan(e + fx)}}$$

```
input Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[
e + f*x])^(3/2)), x]
```

output $(\text{Cos}[e + f*x]^2*(-B + 3*A*\text{Tan}[e + f*x] + 2*A*\text{Tan}[e + f*x]^3))/(3*a*c*f*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])$

3.841.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4071, 87, 55, 41}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2}} dx$$

↓ 4071

$$ac \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx)a + a)^{5/2} (c - ic \tan(e + fx))^{5/2}} d \tan(e + fx)$$

f
↓ 87

$$ac \left(\frac{A \int \frac{1}{(i \tan(e + fx)a + a)^{5/2} (c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{c} - \frac{B + iA}{3ac(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2}} \right)$$

f
↓ 55

$$ac \left(\frac{A \left(\frac{2 \int \frac{1}{(i \tan(e + fx)a + a)^{3/2} (c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{3a} + \frac{i}{3ac(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} \right)}{c} - \frac{B + iA}{3ac(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2}} \right)$$

f
↓ 41

$$ac \left(\frac{A \left(\frac{2 \tan(e + fx)}{3a^2 c \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}} + \frac{i}{3ac(a + ia \tan(e + fx))^{3/2} \sqrt{c - ic \tan(e + fx)}} \right)}{c} - \frac{B + iA}{3ac(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2}} \right)$$

f

3.841. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2}} dx$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(3/2)),x]`

output `(a*c*(-1/3*(I*A + B)/(a*c*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(3/2)) + (A*((I/3)/(a*c*(a + I*a*Tan[e + f*x])^(3/2)*Sqrt[c - I*c*Tan[e + f*x]]) + (2*Tan[e + f*x])/(3*a^2*c*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])))/c)/f`

3.841.3.1 Defintions of rubi rules used

rule 41 `Int[1/(((a_) + (b_)*(x_))^(3/2)*((c_) + (d_)*(x_))^(3/2)), x_Symbol] := Simp[x/(a*c*Sqrt[a + b*x]*Sqrt[c + d*x]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0]`

rule 55 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(-(b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.841.4 Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.74

method	result
derivativedivides	$-\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)}\left(2 A \tan (f x+e)^5+5 A \tan (f x+e)^3-B \tan (f x+e)^2+3 A \tan (f x+e)-B\right)}{3 f a^2 c^2(i-\tan (f x+e))^3(i+\tan (f x+e))^3}$
default	$-\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)}\left(2 A \tan (f x+e)^5+5 A \tan (f x+e)^3-B \tan (f x+e)^2+3 A \tan (f x+e)-B\right)}{3 f a^2 c^2(i-\tan (f x+e))^3(i+\tan (f x+e))^3}$
parts	$-\frac{A \sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)}\left(1+\tan (f x+e)^2\right) \tan (f x+e)\left(2 \tan (f x+e)^2+3\right)}{3 f a^2 c^2(i+\tan (f x+e))^3(i-\tan (f x+e))^3}+\frac{B\left(1+\tan (f x+e)^2\right)}{3 f c^2 a^2}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2)/(c-I*c*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

output `-1/3/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)/a^2/c^2*(2*A*tan(f*x+e)^5+5*A*tan(f*x+e)^3-B*tan(f*x+e)^2+3*A*tan(f*x+e)-B)/(I-tan(f*x+e))^3/(I+tan(f*x+e))^3`

3.841.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.99

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2}(c - ic \tan(e + fx))^{3/2}} dx = \frac{((-iA - B)e^{(8i fx + 8ie)} - 2(5iA + 2B)e^{(6i fx + 6ie)} +$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2)/(c-I*c*tan(f*x+e))^(3/2),x,algorithm="fricas")`

3.841. $\int \frac{A+B \tan (e+f x)}{(a+i a \tan (e+f x))^{3 / 2}(c-i c \tan (e+f x))^{3 / 2}} d x$

output $1/24*((-I*A - B)*e^{(8*I*f*x + 8*I*e)} - 2*(5*I*A + 2*B)*e^{(6*I*f*x + 6*I*e)} + 8*B*e^{(5*I*f*x + 5*I*e)} - 6*B*e^{(4*I*f*x + 4*I*e)} + 8*B*e^{(3*I*f*x + 3*I*e)} - 2*(-5*I*A + 2*B)*e^{(2*I*f*x + 2*I*e)} + I*A - B)*\text{sqrt}(a/(e^{(2*I*f*x + 2*I*e)} + 1))*\text{sqrt}(c/(e^{(2*I*f*x + 2*I*e)} + 1))*e^{(-3*I*f*x - 3*I*e)}/(a^2*c^2*f)$

3.841.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2}} dx = \int \frac{A + B \tan(e + fx)}{(ia (\tan(e + fx) - i))^{\frac{3}{2}} (-ic (\tan(e + fx) + i))^{\frac{3}{2}}}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**(3/2)/(c-I*c*tan(f*x+e))**(3/2),x)`

output `Integral((A + B*tan(e + f*x))/((I*a*(tan(e + f*x) - I))**(3/2)*(-I*c*(tan(e + f*x) + I))**(3/2)), x)`

3.841.7 Maxima [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.32

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2}} dx = \frac{(3(3iA - B) \cos(2fx + 2e) - 3(3A + iB) \sin(2fx + 2e))}{(a^2 c^2)^{3/2}}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2)/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output $1/24*((3*(3*I*A - B)*\cos(2*f*x + 2*e) - 3*(3*A + I*B)*\sin(2*f*x + 2*e) - 2*B)*\cos(3/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 3*(-3*I*A - B)*\cos(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + (3*(3*A + I*B)*\cos(2*f*x + 2*e) + 3*(3*I*A - B)*\sin(2*f*x + 2*e) + 2*A)*\sin(3/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))) + 3*(3*A - I*B)*\sin(1/2*\arctan2(\sin(2*f*x + 2*e), \cos(2*f*x + 2*e))))/(a^{(3/2)}*c^{(3/2)}*f)$

3.841.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2}} dx = \int \frac{B \tan(fx + e) + A}{(ia \tan(fx + e) + a)^{3/2} (-ic \tan(fx + e) + c)^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2)/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)^(3/2)*(-I*c*tan(f*x + e) + c)^(3/2)), x)`

3.841.9 Mupad [B] (verification not implemented)

Time = 9.38 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.30

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{3/2}} dx = \sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)i)}{\cos(2e+2fx)+1}} (10A \sin(2e+2fx) + \dots)$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^(3/2)*(c - c*tan(e + f*x)*1i)^(3/2)),x)`

output `((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*cos(2*e + 2*f*x)*8i - 3*B - A*9i + A*cos(4*e + 4*f*x)*1i - 4*B*cos(2*e + 2*f*x) - B*cos(4*e + 4*f*x) + 10*A*sin(2*e + 2*f*x) + A*sin(4*e + 4*f*x) + B*sin(2*e + 2*f*x)*2i + B*sin(4*e + 4*f*x)*1i))/(24*a^2*c*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

$$3.842 \quad \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{3/2}(c-ict \tan(e+fx))^{5/2}} dx$$

3.842.1 Optimal result	7553
3.842.2 Mathematica [A] (verified)	7553
3.842.3 Rubi [A] (verified)	7554
3.842.4 Maple [A] (verified)	7557
3.842.5 Fricas [A] (verification not implemented)	7557
3.842.6 Sympy [F]	7558
3.842.7 Maxima [F(-2)]	7558
3.842.8 Giac [F]	7559
3.842.9 Mupad [B] (verification not implemented)	7559

3.842.1 Optimal result

Integrand size = 45, antiderivative size = 269

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2}(c - ict \tan(e + fx))^{5/2}} dx = \frac{iA - B}{3f(a + ia \tan(e + fx))^{3/2}(c - ict \tan(e + fx))^{5/2}} + \frac{4iA - B}{3af\sqrt{a + ia \tan(e + fx)}(c - ict \tan(e + fx))^{5/2}} - \frac{(4iA - B)\sqrt{a + ia \tan(e + fx)}}{5a^2f(c - ict \tan(e + fx))^{5/2}} - \frac{2(4iA - B)\sqrt{a + ia \tan(e + fx)}}{15a^2cf(c - ict \tan(e + fx))^{3/2}} - \frac{2(4iA - B)\sqrt{a + ia \tan(e + fx)}}{15a^2c^2f\sqrt{c - ict \tan(e + fx)}}$$

```
output -2/15*(4*I*A-B)*(a+I*a*tan(f*x+e))^(1/2)/a^2/c^2/f/(c-I*c*tan(f*x+e))^(1/2)
)+1/3*(4*I*A-B)/a/f/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(5/2)-1/5*
(4*I*A-B)*(a+I*a*tan(f*x+e))^(1/2)/a^2/f/(c-I*c*tan(f*x+e))^(5/2)+1/3*(I*A
-B)/f/(a+I*a*tan(f*x+e))^(3/2)/(c-I*c*tan(f*x+e))^(5/2)-2/15*(4*I*A-B)*(a+
I*a*tan(f*x+e))^(1/2)/a^2/c/f/(c-I*c*tan(f*x+e))^(3/2)
```

3.842.2 Mathematica [A] (verified)

Time = 6.75 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.57

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2}(c - ict \tan(e + fx))^{5/2}} dx = \frac{3(A - iB) + (12iA - 3B) \tan(e + fx) + 3(4A + iB)}{15ac^2f(-i + \tan(e + fx))(i + \tan(e + fx))}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(5/2)),x]`

output `(3*(A - I*B) + ((12*I)*A - 3*B)*Tan[e + f*x] + 3*(4*A + I*B)*Tan[e + f*x]^2 + ((8*I)*A - 2*B)*Tan[e + f*x]^3 + (8*A + (2*I)*B)*Tan[e + f*x]^4)/(15*a*c^2*f*(-I + Tan[e + f*x])*(I + Tan[e + f*x])^2*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])`

3.842.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.01, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 87, 55, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & ac \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx)a + a)^{5/2} (c - ic \tan(e + fx))^{7/2}} d \tan(e + fx) \\
 & \quad \downarrow \text{87} \\
 & ac \left(\frac{(4A + iB) \int \frac{1}{(i \tan(e + fx)a + a)^{3/2} (c - ic \tan(e + fx))^{7/2}} d \tan(e + fx)}{3a} + \frac{-B + iA}{3ac(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{5/2}} \right) \\
 & \quad \downarrow \text{55} \\
 & ac \left(\frac{(4A + iB) \left(\frac{3 \int \frac{1}{\sqrt{i \tan(e + fx)a + a} (c - ic \tan(e + fx))^{7/2}} d \tan(e + fx)}{a} + \frac{i}{ac \sqrt{a + ia \tan(e + fx)} (c - ic \tan(e + fx))^{5/2}} \right)}{3a} + \frac{-B + iA}{3ac(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{5/2}} \right) \\
 & \quad \downarrow \text{55}
 \end{aligned}$$

3.842. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{5/2}} dx$

$$ac \left(\frac{(4A+iB) \left(\frac{2 \int \frac{1}{\sqrt{i \tan(e+fx)a+a(c-ic \tan(e+fx))^{5/2}} d \tan(e+fx)}{5c} - \frac{i \sqrt{a+ia \tan(e+fx)}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)}{a} + \frac{i}{ac \sqrt{a+ia \tan(e+fx)(c-ic \tan(e+fx))^{5/2}} \right)}{3a} + \frac{i}{3ac} \right)$$

f

↓ 55

$$ac \left(\frac{(4A+iB) \left(\frac{2 \left(\frac{\int \frac{1}{\sqrt{i \tan(e+fx)a+a(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{3c} - \frac{i \sqrt{a+ia \tan(e+fx)}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{5c} - \frac{i \sqrt{a+ia \tan(e+fx)}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)}{a} + \frac{i}{ac \sqrt{a+ia \tan(e+fx)(c-ic \tan(e+fx))^{5/2}} \right)}{3a} + \frac{i}{3ac} \right)$$

f

↓ 48

$$ac \left(\frac{(4A+iB) \left(\frac{2 \left(\frac{-\frac{i \sqrt{a+ia \tan(e+fx)}}{3ac^2 \sqrt{c-ic \tan(e+fx)}} - \frac{i \sqrt{a+ia \tan(e+fx)}}{3ac(c-ic \tan(e+fx))^{3/2}} \right)}{5c} - \frac{i \sqrt{a+ia \tan(e+fx)}}{5ac(c-ic \tan(e+fx))^{5/2}} \right)}{a} + \frac{i}{ac \sqrt{a+ia \tan(e+fx)(c-ic \tan(e+fx))^{5/2}} \right)}{3a} + \frac{i}{3ac} \right)$$

f

3.842. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{3/2}(c-ic \tan(e+fx))^{5/2}} dx$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(5/2)),x]`

output `(a*c*((I*A - B)/(3*a*c*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(5/2)) + ((4*A + I*B)*(I/(a*c*Sqrt[a + I*a*Tan[e + f*x]]*(c - I*c*Tan[e + f*x])^(5/2)) + (3*((-1/5*I)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c*(c - I*c*Tan[e + f*x])^(5/2)) + (2*((-1/3*I)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c*(c - I*c*Tan[e + f*x])^(3/2)) - ((I/3)*Sqrt[a + I*a*Tan[e + f*x]])/(a*c^2*Sqrt[c - I*c*Tan[e + f*x]])))/(5*c)))/a)/(3*a))/f`

3.842.3.1 Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

rule 55 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x],
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.842.4 Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 199, normalized size of antiderivative = 0.74

method	result
derivativedivides	$\frac{i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(8iA\tan(fx+e)^6-2iB\tan(fx+e)^5-2B\tan(fx+e)^6+20iA\tan(fx+e)^4-8A\tan(fx+e)^5)}{15fa^2c^3(i+\tan(fx+e))^4}$
default	$\frac{i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(8iA\tan(fx+e)^6-2iB\tan(fx+e)^5-2B\tan(fx+e)^6+20iA\tan(fx+e)^4-8A\tan(fx+e)^5)}{15fa^2c^3(i+\tan(fx+e))^4}$
parts	$-\frac{A\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(8i\tan(fx+e)^5+8\tan(fx+e)^6+20i\tan(fx+e)^3+20\tan(fx+e)^4+12i\tan(fx+e)^5)}{15fa^2c^3(i-\tan(fx+e))^3(i+\tan(fx+e))^4}$

```
input int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2)/(c-I*c*tan(f*x+e))^(5/2),x,m
ethod=_RETURNVERBOSE)
```

```
output 1/15*I/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)/a^2/c^3*(8
*I*A*tan(f*x+e)^6-2*I*B*tan(f*x+e)^5-2*B*tan(f*x+e)^6+20*I*A*tan(f*x+e)^4-
8*A*tan(f*x+e)^5-5*I*B*tan(f*x+e)^3-5*B*tan(f*x+e)^4+15*I*A*tan(f*x+e)^2-2
0*A*tan(f*x+e)^3-3*I*B*tan(f*x+e)+3*I*A-12*A*tan(f*x+e)+3*B)/(I+tan(f*x+e)
)^4/(I-tan(f*x+e))^3
```

3.842.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 181, normalized size of antiderivative = 0.67

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{5/2}} dx = \frac{(3(iA + B)e^{(10ifx+10ie)} - (-23iA - 13B)e^{(8ifx+8ie)} + 10(11iA + B)e^{(6ifx+6ie)} + 48(-iA - B)e^{(5ifx+5ie)})}{15fa^2c^3(i+\tan(fx+e))^4}$$

```
input integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2)/(c-I*c*tan(f*x+e))^(5/
2),x, algorithm="fricas")
```

3.842. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{3/2} (c-ic \tan(e+fx))^{5/2}} dx$

output
$$-1/240*(3*(I*A + B)*e^{(10*I*f*x + 10*I*e)} - (-23*I*A - 13*B)*e^{(8*I*f*x + 8*I*e)} + 10*(11*I*A + B)*e^{(6*I*f*x + 6*I*e)} + 48*(-I*A - B)*e^{(5*I*f*x + 5*I*e)} + 30*(I*A + B)*e^{(4*I*f*x + 4*I*e)} + 48*(-I*A - B)*e^{(3*I*f*x + 3*I*e)} + 5*(-13*I*A + 7*B)*e^{(2*I*f*x + 2*I*e)} - 5*I*A + 5*B)*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)}*e^{(-3*I*f*x - 3*I*e)}/(a^2*c^3*f)$$

3.842.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{5/2}} dx = \int \frac{A + B \tan(e + fx)}{(ia (\tan(e + fx) - i))^{3/2} (-ic (\tan(e + fx) + i))^{5/2}}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**(3/2)/(c-I*c*tan(f*x+e))**(5/2),x)`

output `Integral((A + B*tan(e + f*x))/((I*a*(tan(e + f*x) - I))**(3/2)*(-I*c*(tan(e + f*x) + I))**(5/2)), x)`

3.842.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2)/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.842.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{5/2}} dx = \int \frac{B \tan(fx + e) + A}{(ia \tan(fx + e) + a)^{3/2} (-ic \tan(fx + e) + c)^{5/2}} dx$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2)/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)^(3/2)*(-I*c*tan(f*x + e) + c)^(5/2)), x)`

3.842.9 Mupad [B] (verification not implemented)

Time = 9.54 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.73

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{3/2} (c - ic \tan(e + fx))^{5/2}} dx = \frac{\sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)i)}{\cos(2e+2fx)+1}}}{(40 A \sin(2e + 2fx))} dx$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^(3/2)*(c - c*tan(e + f*x)*1i)^(5/2)),x)`

output `((a*cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1)/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*cos(2*e + 2*f*x)*20i - A*45i + A*cos(4*e + 4*f*x)*1i - 20*B*cos(2*e + 2*f*x) - 4*B*cos(4*e + 4*f*x) + 40*A*sin(2*e + 2*f*x) + 4*A*sin(4*e + 4*f*x) + B*sin(2*e + 2*f*x)*10i + B*sin(4*e + 4*f*x)*1i)/(120*a^2*c^2*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

3.843
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^{5/2}} dx$$

3.843.1 Optimal result	7560
3.843.2 Mathematica [C] (verified)	7561
3.843.3 Rubi [A] (verified)	7561
3.843.4 Maple [B] (verified)	7566
3.843.5 Fracas [B] (verification not implemented)	7567
3.843.6 Sympy [F(-1)]	7568
3.843.7 Maxima [F(-2)]	7568
3.843.8 Giac [F]	7568
3.843.9 Mupad [F(-1)]	7569

3.843.1 Optimal result

Integrand size = 45, antiderivative size = 343

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{9/2}}{(a + ia \tan(e + fx))^{5/2}} dx = \frac{7(2iA - 7B)c^{9/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{a^{5/2}f}$$

$$+ \frac{7(2iA - 7B)c^4 \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}}{2a^3 f}$$

$$+ \frac{7(2iA - 7B)c^3 \sqrt{a + ia \tan(e + fx)} (c - ic \tan(e + fx))^{3/2}}{6a^3 f}$$

$$+ \frac{14(2iA - 7B)c^2 (c - ic \tan(e + fx))^{5/2}}{15a^2 f \sqrt{a + ia \tan(e + fx)}}$$

$$- \frac{2(2iA - 7B)c (c - ic \tan(e + fx))^{7/2}}{15af (a + ia \tan(e + fx))^{3/2}} + \frac{(iA - B)(c - ic \tan(e + fx))^{9/2}}{5f (a + ia \tan(e + fx))^{5/2}}$$

```
output 7*(2*I*A-7*B)*c^(9/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I
*c*tan(f*x+e))^(1/2))/a^(5/2)/f+7/2*(2*I*A-7*B)*c^4*(a+I*a*tan(f*x+e))^(1/
2)*(c-I*c*tan(f*x+e))^(1/2)/a^3/f+7/6*(2*I*A-7*B)*c^3*(a+I*a*tan(f*x+e))^(
1/2)*(c-I*c*tan(f*x+e))^(3/2)/a^3/f+14/15*(2*I*A-7*B)*c^2*(c-I*c*tan(f*x+e
))^(5/2)/a^2/f/(a+I*a*tan(f*x+e))^(1/2)-2/15*(2*I*A-7*B)*c*(c-I*c*tan(f*x+
e))^(7/2)/a/f/(a+I*a*tan(f*x+e))^(3/2)+1/5*(I*A-B)*(c-I*c*tan(f*x+e))^(9/2
)/f/(a+I*a*tan(f*x+e))^(5/2)
```

3.843.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 9.77 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.66

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^{5/2}} dx =$$

$$c^4 \sqrt{c - i c \tan(e + fx)} \left(\frac{3}{2} \sec^4(e + fx) (i \cos(2(e + fx)) + \sin(2(e + fx))) (9(2A + 7iB) + (18A + 53iB) c \right)$$

30a

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(9/2))/(a + I*a*Tan[e + f*x])^(5/2),x]`

output `-1/30*(c^4*Sqrt[c - I*c*Tan[e + f*x]]*((3*Sec[e + f*x]^4*(I*Cos[2*(e + f*x)] + Sin[2*(e + f*x)])*(9*(2*A + (7*I)*B) + (18*A + (53*I)*B)*Cos[2*(e + f*x)] + (5*I)*(2*A + (9*I)*B)*Sin[2*(e + f*x)])*Sqrt[1 - I*Tan[e + f*x]])/2 + 140*Sqrt[2]*(2*A + (7*I)*B)*Hypergeometric2F1[-3/2, -3/2, -1/2, (1 + I*Tan[e + f*x])/2]*(-I + Tan[e + f*x]))/(a^2*f*Sqrt[1 - I*Tan[e + f*x]]*(-I + Tan[e + f*x])^2*Sqrt[a + I*a*Tan[e + f*x]])`

3.843.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 327, normalized size of antiderivative = 0.95, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3042, 4071, 87, 57, 57, 60, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - i c \tan(e + fx))^{9/2} (A + B \tan(e + fx))}{(a + i a \tan(e + fx))^{5/2}} dx$$

↓ 3042

$$\int \frac{(c - i c \tan(e + fx))^{9/2} (A + B \tan(e + fx))}{(a + i a \tan(e + fx))^{5/2}} dx$$

↓ 4071

$$\frac{a c \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(i \tan(e + fx) a + a)^{7/2}} d \tan(e + fx)}{f}$$

3.843. $\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^{5/2}} dx$

$$\begin{array}{c}
 \downarrow 87 \\
 ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{9/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{(2A+7iB) \int \frac{(c-ic \tan(e+fx))^{7/2}}{(i \tan(e+fx)a+a)^{5/2}} d \tan(e+fx)}{5a} \right) \\
 \hline
 f \\
 \downarrow 57 \\
 ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{9/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{(2A+7iB) \left(\frac{2i(c-ic \tan(e+fx))^{7/2}}{3a(a+ia \tan(e+fx))^{3/2}} - \frac{7c \int \frac{(c-ic \tan(e+fx))^{5/2}}{(i \tan(e+fx)a+a)^{3/2}} d \tan(e+fx)}{3a} \right)}{5a} \right) \\
 \hline
 f \\
 \downarrow 57 \\
 ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{9/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{(2A+7iB) \left(\frac{2i(c-ic \tan(e+fx))^{7/2}}{3a(a+ia \tan(e+fx))^{3/2}} - \frac{7c \left(\frac{2i(c-ic \tan(e+fx))^{5/2}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{5c \int \frac{(c-ic \tan(e+fx))^{3/2}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e+fx)}{a} \right)}{3a} \right)}{5a} \right) \\
 \hline
 f \\
 \downarrow 60 \\
 ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{9/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{(2A+7iB) \left(\frac{2i(c-ic \tan(e+fx))^{7/2}}{3a(a+ia \tan(e+fx))^{3/2}} - \frac{7c \left(\frac{2i(c-ic \tan(e+fx))^{5/2}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{5c \left(\frac{3}{2} c \int \frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{i \tan(e+fx)a+a}} d \tan(e+fx) - \frac{i\sqrt{a}}{a} \right)}{3a} \right)}{5a} \right)}{5a} \right) \\
 \hline
 f \\
 \downarrow 60
 \end{array}$$

3.843. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^{5/2}} dx$

$$ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{9/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{(2A+7iB) \left(\frac{2i(c-ic \tan(e+fx))^{7/2}}{3a(a+ia \tan(e+fx))^{3/2}} - \frac{7c \left(\frac{2i(c-ic \tan(e+fx))^{5/2}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{5c \left(\frac{3}{2}c \int \frac{1}{\sqrt{i \tan(e+fx)a+a\sqrt{c-ic \tan(e+fx)}}} \right)}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}}} d \frac{\sqrt{i \tan(e+fx)}}{\sqrt{c-ic \tan(e+fx)}}} \right)}{5a} \right) dx$$

↓ 45

$$ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{9/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{(2A+7iB) \left(\frac{2i(c-ic \tan(e+fx))^{7/2}}{3a(a+ia \tan(e+fx))^{3/2}} - \frac{7c \left(\frac{2i(c-ic \tan(e+fx))^{5/2}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{5c \left(\frac{3}{2}c \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}}} d \frac{\sqrt{i \tan(e+fx)}}{\sqrt{c-ic \tan(e+fx)}}} \right)}{5a} \right)}{5a} \right) dx$$

↓ 218

$$\begin{aligned}
 & \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{9/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{(2A+7iB) \frac{2i(c-ic \tan(e+fx))^{7/2}}{3a(a+ia \tan(e+fx))^{3/2}} - \frac{7c \frac{2i(c-ic \tan(e+fx))^{5/2}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{5c \left(\frac{3}{2}c \left(-\frac{2i\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{\sqrt{a}} \right)}{5a}}{f} \right) \right)
 \end{aligned}$$

```
input Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(9/2))/(a + I*a*Tan[e + f*x])^(5/2), x]
```

```
output (a*c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(9/2))/(5*a*c*(a + I*a*Tan[e + f*x])^(5/2)) - ((2*A + (7*I)*B)*(((2*I)/3)*(c - I*c*Tan[e + f*x])^(7/2))/(a*(a + I*a*Tan[e + f*x])^(3/2)) - (7*c*(((2*I)*(c - I*c*Tan[e + f*x])^(5/2))/(a*Sqrt[a + I*a*Tan[e + f*x]])) - (5*c*(((1/2*I)*Sqrt[a + I*a*Tan[e + f*x]])*(c - I*c*Tan[e + f*x])^(3/2))/a + (3*c*(((2*I)*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[a] - (I*Sqrt[a + I*a*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]/a))/2)/a)/(3*a)))/(5*a))/f
```

3.843.3.1 Defintions of rubi rules used

```
rule 45 Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]
```

$$3.843. \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^{5/2}} dx$$

- rule 57 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`
- rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) + (f_.)*(x_)])*((c_.) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.843.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 898 vs. $2(283) = 566$.

Time = 0.44 (sec) , antiderivative size = 899, normalized size of antiderivative = 2.62

method	result	size
derivativedivides	Expression too large to display	899
default	Expression too large to display	899
parts	Expression too large to display	957

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2)/(a+I*a*tan(f*x+e))^(5/2),x,m
method=_RETURNVERBOSE)
```

```
output 1/30/f*(-c*(I*tan(f*x+e)-1))^(1/2)*(a*(1+I*tan(f*x+e)))^(1/2)*c^4/a^3*(-73
5*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(
1/2))*a*c*tan(f*x+e)^4+15*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan
(f*x+e)^5+2014*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^3-3
881*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)-210*A*ln((a*c*
tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2))/(a*c)^(1/2))*a*c*tan(
f*x+e)^4+840*I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/
2)))/(a*c)^(1/2))*a*c*tan(f*x+e)^3-1316*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^
2))^(1/2)*tan(f*x+e)^2-2940*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f
*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c*tan(f*x+e)^3-150*B*(a*c)^(1/2)*(a*c*(1+t
an(f*x+e)^2))^(1/2)*tan(f*x+e)^4+4410*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(
a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c*tan(f*x+e)^2-840*I*A*ln((a*c
*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c*tan
(f*x+e)+1260*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)
))/(a*c)^(1/2))*a*c*tan(f*x+e)^2+584*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(
1/2)*tan(f*x+e)^3+334*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)-735*I*B
*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))
*a*c+2940*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(
a*c)^(1/2))*a*c*tan(f*x+e)+4576*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)
*tan(f*x+e)^2+30*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e...
```

3.843.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 610 vs. $2(263) = 526$.

Time = 0.28 (sec) , antiderivative size = 610, normalized size of antiderivative = 1.78

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{9/2}}{(a + ia \tan(e + fx))^{5/2}} dx =$$

$$105 (a^3 f e^{(7i fx + 7i e)} + a^3 f e^{(5i fx + 5i e)}) \sqrt{\frac{(4A^2 + 28iAB - 49B^2)c^9}{a^5 f^2}} \log \left(\frac{4 \left(2((2iA - 7B)c^4 e^{(3i fx + 3i e)} + (2iA - 7B)c^4 e^{(i fx + i e)}) \right)}{(2iA - 7B)c^4 e^{(2i fx + 2i e)} + (2iA - 7B)c^4} \right)$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2)/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="fricas")`

output

```
-1/60*(105*(a^3*f*e^(7*I*f*x + 7*I*e) + a^3*f*e^(5*I*f*x + 5*I*e))*sqrt((4
*A^2 + 28*I*A*B - 49*B^2)*c^9/(a^5*f^2))*log(4*(2*((2*I*A - 7*B)*c^4*e^(3*
I*f*x + 3*I*e) + (2*I*A - 7*B)*c^4*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2
*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)) + (a^3*f*e^(2*I*f*x + 2*I*e)
- a^3*f)*sqrt((4*A^2 + 28*I*A*B - 49*B^2)*c^9/(a^5*f^2)))/((2*I*A - 7*B)*
c^4*e^(2*I*f*x + 2*I*e) + (2*I*A - 7*B)*c^4) - 105*(a^3*f*e^(7*I*f*x + 7*
I*e) + a^3*f*e^(5*I*f*x + 5*I*e))*sqrt((4*A^2 + 28*I*A*B - 49*B^2)*c^9/(a^
5*f^2))*log(4*(2*((2*I*A - 7*B)*c^4*e^(3*I*f*x + 3*I*e) + (2*I*A - 7*B)*c^
4*e^(I*f*x + I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x +
2*I*e) + 1)) - (a^3*f*e^(2*I*f*x + 2*I*e) - a^3*f)*sqrt((4*A^2 + 28*I*A*B
- 49*B^2)*c^9/(a^5*f^2)))/((2*I*A - 7*B)*c^4*e^(2*I*f*x + 2*I*e) + (2*I*A
- 7*B)*c^4) + 4*(105*(-2*I*A + 7*B)*c^4*e^(8*I*f*x + 8*I*e) + 175*(-2*I*A
+ 7*B)*c^4*e^(6*I*f*x + 6*I*e) + 56*(-2*I*A + 7*B)*c^4*e^(4*I*f*x + 4*I*e
) + 8*(2*I*A - 7*B)*c^4*e^(2*I*f*x + 2*I*e) + 12*(-I*A + B)*c^4)*sqrt(a/(e
^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1)))/(a^3*f*e^(7*I*
f*x + 7*I*e) + a^3*f*e^(5*I*f*x + 5*I*e))
```

3.843.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(9/2)/(a+I*a*tan(f*x+e))**(5/2),x)`

output `Timed out`

3.843.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2)/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.843.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{9/2}}{(i a \tan(fx + e) + a)^{5/2}} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(9/2)/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(9/2)/(I*a*tan(f*x + e) + a)^(5/2), x)`

3.843. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{9/2}}{(a+ia \tan(e+fx))^{5/2}} dx$

3.843.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{9/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \int \frac{(A + B \tan(e + fx)) (c - c \tan(e + fx) i)^{9/2}}{(a + a \tan(e + fx) i)^{5/2}} dx$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(9/2))/(a + a*tan(e + f*x)*1i)^(5/2),x)`

output `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(9/2))/(a + a*tan(e + f*x)*1i)^(5/2), x)`

$$3.844 \quad \int \frac{(A+B \tan(e+fx))(c-ictan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^{5/2}} dx$$

3.844.1 Optimal result	7570
3.844.2 Mathematica [C] (verified)	7571
3.844.3 Rubi [A] (verified)	7571
3.844.4 Maple [B] (verified)	7575
3.844.5 Fracas [B] (verification not implemented)	7576
3.844.6 Sympy [F(-1)]	7577
3.844.7 Maxima [B] (verification not implemented)	7577
3.844.8 Giac [F]	7578
3.844.9 Mupad [F(-1)]	7579

3.844.1 Optimal result

Integrand size = 45, antiderivative size = 284

$$\int \frac{(A+B \tan(e+fx))(c-ictan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^{5/2}} dx = \frac{2(iA-6B)c^{7/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ictan(e+fx)}}\right)}{a^{5/2}f}$$

$$+ \frac{(iA-6B)c^3 \sqrt{a+ia \tan(e+fx)} \sqrt{c-ictan(e+fx)}}{a^3 f}$$

$$+ \frac{2(iA-6B)c^2(c-ictan(e+fx))^{3/2}}{3a^2 f \sqrt{a+ia \tan(e+fx)}}$$

$$- \frac{2(iA-6B)c(c-ictan(e+fx))^{5/2}}{15af(a+ia \tan(e+fx))^{3/2}} + \frac{(iA-B)(c-ictan(e+fx))^{7/2}}{5f(a+ia \tan(e+fx))^{5/2}}$$

output

```
2*(I*A-6*B)*c^(7/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c
*tan(f*x+e))^(1/2))/a^(5/2)/f+(I*A-6*B)*c^3*(a+I*a*tan(f*x+e))^(1/2)*(c-I*
c*tan(f*x+e))^(1/2)/a^3/f+2/3*(I*A-6*B)*c^2*(c-I*c*tan(f*x+e))^(3/2)/a^2/f
/(a+I*a*tan(f*x+e))^(1/2)-2/15*(I*A-6*B)*c*(c-I*c*tan(f*x+e))^(5/2)/a/f/(a
+I*a*tan(f*x+e))^(3/2)+1/5*(I*A-B)*(c-I*c*tan(f*x+e))^(7/2)/f/(a+I*a*tan(f
*x+e))^(5/2)
```

3.844.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 7.67 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.67

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \frac{c^3 \sqrt{c - i c \tan(e + fx)} \left(-20\sqrt{2}(A + 6iB) \text{Hypergeometric2F1}[-3/2, -3/2, -1/2, (1 + I \tan[e + f*x])/2] * (-I + \tan[e + f*x]) + 3 \sec[e + f*x]^2 * (I \cos[2*(e + f*x)] + \sin[2*(e + f*x)]) * \sqrt{1 - I \tan[e + f*x]} * (-2*A - (7*I)*B + 5*B \tan[e + f*x]) \right)}{(15*a^2*f \sqrt{1 - I \tan[e + f*x]} * (-I + \tan[e + f*x])^2 \sqrt{a + I*a \tan[e + f*x]})}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2))/(a + I*a*Tan[e + f*x])^(5/2),x]`

output `(c^3*Sqrt[c - I*c*Tan[e + f*x]]*(-20*Sqrt[2]*(A + (6*I)*B)*Hypergeometric2F1[-3/2, -3/2, -1/2, (1 + I*Tan[e + f*x])/2]*(-I + Tan[e + f*x]) + 3*Sec[e + f*x]^2*(I*Cos[2*(e + f*x)] + Sin[2*(e + f*x)])*Sqrt[1 - I*Tan[e + f*x]]*(-2*A - (7*I)*B + 5*B*Tan[e + f*x]))/(15*a^2*f*Sqrt[1 - I*Tan[e + f*x]]*(-I + Tan[e + f*x])^2*Sqrt[a + I*a*Tan[e + f*x]])`

3.844.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 276, normalized size of antiderivative = 0.97, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.178$, Rules used = {3042, 4071, 87, 57, 57, 60, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c - i c \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(a + i a \tan(e + fx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{(c - i c \tan(e + fx))^{7/2} (A + B \tan(e + fx))}{(a + i a \tan(e + fx))^{5/2}} dx \\ & \quad \downarrow \text{4071} \\ & \frac{ac \int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(i \tan(e + fx)a + a)^{7/2}} d \tan(e + fx)}{f} \\ & \quad \downarrow \text{87} \end{aligned}$$

3.844. $\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^{5/2}} dx$

$$\begin{aligned}
 & \frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{(A+6iB) \int \frac{(c-ic \tan(e+fx))^{5/2}}{(i \tan(e+fx)a+a)^{5/2}} d \tan(e+fx)}{5a} \right)}{f} \\
 & \quad \downarrow 57 \\
 & \frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{(A+6iB) \left(\frac{2i(c-ic \tan(e+fx))^{5/2}}{3a(a+ia \tan(e+fx))^{3/2}} - \frac{5c \int \frac{(c-ic \tan(e+fx))^{3/2}}{(i \tan(e+fx)a+a)^{3/2}} d \tan(e+fx)}{3a} \right)}{5a} \right)}{f} \\
 & \quad \downarrow 57 \\
 & \frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{(A+6iB) \left(\frac{2i(c-ic \tan(e+fx))^{5/2}}{3a(a+ia \tan(e+fx))^{3/2}} - \frac{5c \left(\frac{2i(c-ic \tan(e+fx))^{3/2}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{3c \int \frac{\sqrt{c-ic \tan(e+fx)}}{\sqrt{i \tan(e+fx)a+a} \frac{1}{a}} d \tan(e+fx)}{3a} \right)}{3a} \right)}{5a} \right)}{f} \\
 & \quad \downarrow 60 \\
 & \frac{ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{(A+6iB) \left(\frac{2i(c-ic \tan(e+fx))^{5/2}}{3a(a+ia \tan(e+fx))^{3/2}} - \frac{5c \left(\frac{2i(c-ic \tan(e+fx))^{3/2}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{3c \int \frac{1}{\sqrt{i \tan(e+fx)a+a} \sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{3a} \right)}{3a} \right)}{5a} \right)}{f} \\
 & \quad \downarrow 45
 \end{aligned}$$

3.844. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^{5/2}} dx$

$$ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{(A+6iB) \frac{2i(c-ic \tan(e+fx))^{5/2}}{3a(a+ia \tan(e+fx))^{3/2}} - \frac{5c \left(\frac{2i(c-ic \tan(e+fx))^{3/2}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{3c \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)}}{\sqrt{c-ic \tan(e+fx)}}}{a} \right)}{3a}}{5a} \right) dx$$

↓ 218

$$ac \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{7/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{(A+6iB) \frac{2i(c-ic \tan(e+fx))^{5/2}}{3a(a+ia \tan(e+fx))^{3/2}} - \frac{5c \left(\frac{2i(c-ic \tan(e+fx))^{3/2}}{a\sqrt{a+ia \tan(e+fx)}} - \frac{3c \left(-\frac{2i\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{\sqrt{a}} - \frac{i\sqrt{c}}{a} \right)}{a} \right)}{3a}}{5a} \right) dx$$

```
input Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(7/2))/(a + I*a*Tan[e + f*x])^(5/2), x]
```

3.844. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^{5/2}} dx$

```
output (a*c*(((I*A - B)*(c - I*c*Tan[e + f*x])^(7/2))/(5*a*c*(a + I*a*Tan[e + f*x
])^(5/2)) - ((A + (6*I)*B)*(((2*I)/3)*(c - I*c*Tan[e + f*x])^(5/2))/(a*(a
+ I*a*Tan[e + f*x])^(3/2)) - (5*c*(((2*I)*(c - I*c*Tan[e + f*x])^(3/2))/(
a*Sqrt[a + I*a*Tan[e + f*x]]) - (3*c*(((2*I)*Sqrt[c]*ArcTan[(Sqrt[c]*Sqrt
[a + I*a*Tan[e + f*x]])/(Sqrt[a]*Sqrt[c - I*c*Tan[e + f*x]])])/Sqrt[a] - (
I*Sqrt[a + I*a*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x])/a))/a))/(3*a)))/f
```

3.844.3.1 Defintions of rubi rules used

```
rule 45 Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && !GtQ[c, 0]
```

```
rule 57 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] &
& GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m
+ n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c
, d, m, n, x]
```

```
rule 60 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(
b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```

```
rule 87 Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
_), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

```
rule 218 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

$$3.844. \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^{5/2}} dx$$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], Tan[e + f*x], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.844.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 834 vs. 2(235) = 470.

Time = 0.38 (sec) , antiderivative size = 835, normalized size of antiderivative = 2.94

method	result
derivativedivides	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c^3 \left(-90iB \ln \left(\frac{ac \tan(fx+e) + \sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right) ac \tan(fx+e)^4 + 246}{}$
default	$\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c^3 \left(-90iB \ln \left(\frac{ac \tan(fx+e) + \sqrt{ac} \sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}} \right) \right) ac \tan(fx+e)^4 + 246}{}$
parts	Expression too large to display

input `int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

$$3.844. \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^{5/2}} dx$$

```

output 1/15/f*(-c*(I*tan(f*x+e)-1))^(1/2)*(a*(1+I*tan(f*x+e)))^(1/2)*c^3/a^3*(-90
*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1
/2))*a*c*tan(f*x+e)^4+246*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan
(f*x+e)^3-15*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)
)/(a*c)^(1/2))*a*c*tan(f*x+e)^4-474*I*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))
^(1/2)*tan(f*x+e)+60*I*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)
^2))^(1/2)))/(a*c)^(1/2))*a*c*tan(f*x+e)^3-360*B*ln((a*c*tan(f*x+e)+(a*c)^(
1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c*tan(f*x+e)^3-15*B*(a*c
)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^4-94*I*A*(a*c)^(1/2)*(a*c
(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2+540*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)
)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c*tan(f*x+e)^2+90*A*ln((a*c
*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c*tan
(f*x+e)^2+46*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^3-60*I*
A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2)
)*a*c*tan(f*x+e)+26*I*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)+360*B*ln(
(a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1/2))*a*c
*tan(f*x+e)+564*B*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x+e)^2-90
*I*B*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)))/(a*c)^(1
/2))*a*c-15*A*ln((a*c*tan(f*x+e)+(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)
)/(a*c)^(1/2))*a*c-74*A*(a*c)^(1/2)*(a*c*(1+tan(f*x+e)^2))^(1/2)*tan(f*x...

```

3.844.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 533 vs. $2(218) = 436$.

Time = 0.27 (sec) , antiderivative size = 533, normalized size of antiderivative = 1.88

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^{5/2}} dx =$$

$$\left(15 a^3 \sqrt{\frac{(A^2 + 12i AB - 36 B^2)c^7}{a^5 f^2}} f e^{(5i fx + 5i e)} \log \left(- \frac{4 \left(2 \left((i A - 6 B)c^3 e^{(3i fx + 3i e)} + (i A - 6 B)c^3 e^{(i fx + i e)} \right) \sqrt{\frac{a}{e^{(2i fx + 2i e)} + 1}} \sqrt{e^{(2i fx + 2i e)}} \right)}{(-i A + 6 B)c^3 e^{(2i fx + 2i e)} + (-} \right.$$

```

input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^(5/
2),x, algorithm="fracas")

```

3.844. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^{5/2}} dx$

output
$$-1/30*(15*a^3*\sqrt{(A^2 + 12*I*A*B - 36*B^2)*c^7/(a^5*f^2)}*f*e^{(5*I*f*x + 5*I*e)}*\log(-4*(2*((I*A - 6*B)*c^3*e^{(3*I*f*x + 3*I*e)} + (I*A - 6*B)*c^3*e^{(I*f*x + I*e)}))*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} + (a^3*f*e^{(2*I*f*x + 2*I*e)} - a^3*f)*\sqrt{(A^2 + 12*I*A*B - 36*B^2)*c^7/(a^5*f^2)})/((-I*A + 6*B)*c^3*e^{(2*I*f*x + 2*I*e)} + (-I*A + 6*B)*c^3)) - 15*a^3*\sqrt{(A^2 + 12*I*A*B - 36*B^2)*c^7/(a^5*f^2)}*f*e^{(5*I*f*x + 5*I*e)}*\log(-4*(2*((I*A - 6*B)*c^3*e^{(3*I*f*x + 3*I*e)} + (I*A - 6*B)*c^3*e^{(I*f*x + I*e)}))*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} - (a^3*f*e^{(2*I*f*x + 2*I*e)} - a^3*f)*\sqrt{(A^2 + 12*I*A*B - 36*B^2)*c^7/(a^5*f^2)})/((-I*A + 6*B)*c^3*e^{(2*I*f*x + 2*I*e)} + (-I*A + 6*B)*c^3)) + 4*(15*(-I*A + 6*B)*c^3*e^{(6*I*f*x + 6*I*e)} + 10*(-I*A + 6*B)*c^3*e^{(4*I*f*x + 4*I*e)} + 2*(I*A - 6*B)*c^3*e^{(2*I*f*x + 2*I*e)} + 3*(-I*A + B)*c^3)*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)})*e^{(-5*I*f*x - 5*I*e)}/(a^3*f)$$

3.844.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \text{Timed out}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(7/2)/(a+I*a*tan(f*x+e))**(5/2),x)`

output Timed out

3.844.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1025 vs. $2(218) = 436$.

Time = 0.57 (sec) , antiderivative size = 1025, normalized size of antiderivative = 3.61

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \text{Too large to display}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="maxima")`

3.844.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{7/2}}{(a+ia \tan(e+fx))^{5/2}} dx$$

output

```

15*(60*(A + 6*I*B)*c^3*cos(6*f*x + 6*e) + 40*(A + 6*I*B)*c^3*cos(4*f*x + 4
*e) - 8*(A + 6*I*B)*c^3*cos(2*f*x + 2*e) + 60*(I*A - 6*B)*c^3*sin(6*f*x +
6*e) + 40*(I*A - 6*B)*c^3*sin(4*f*x + 4*e) + 8*(-I*A + 6*B)*c^3*sin(2*f*x
+ 2*e) + 12*(A + I*B)*c^3 + 30*((A + 6*I*B)*c^3*cos(7/2*arctan2(sin(2*f*x
+ 2*e), cos(2*f*x + 2*e))) + (A + 6*I*B)*c^3*cos(5/2*arctan2(sin(2*f*x + 2
*e), cos(2*f*x + 2*e)))) + (I*A - 6*B)*c^3*sin(7/2*arctan2(sin(2*f*x + 2*e)
, cos(2*f*x + 2*e))) + (I*A - 6*B)*c^3*sin(5/2*arctan2(sin(2*f*x + 2*e), c
os(2*f*x + 2*e))))*arctan2(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2
*e))), sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e)))) + 1) + 30*((A
+ 6*I*B)*c^3*cos(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (A + 6
*I*B)*c^3*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (I*A - 6*
B)*c^3*sin(7/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (I*A - 6*B)*
c^3*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))*arctan2(cos(1/2*
arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))), -sin(1/2*arctan2(sin(2*f*x +
2*e), cos(2*f*x + 2*e)))) + 1) + 15*((I*A - 6*B)*c^3*cos(7/2*arctan2(sin(2
*f*x + 2*e), cos(2*f*x + 2*e))) + (I*A - 6*B)*c^3*cos(5/2*arctan2(sin(2*f*
x + 2*e), cos(2*f*x + 2*e))) - (A + 6*I*B)*c^3*sin(7/2*arctan2(sin(2*f*x +
2*e), cos(2*f*x + 2*e))) - (A + 6*I*B)*c^3*sin(5/2*arctan2(sin(2*f*x + 2*
e), cos(2*f*x + 2*e))))*log(cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x +
2*e))))^2 + sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))^2 + 2*s...

```

3.844.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{7/2}}{(i a \tan(fx + e) + a)^{5/2}} dx$$

input

```

integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(7/2)/(a+I*a*tan(f*x+e))^(5/
2),x, algorithm="giac")

```

output

```

integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(7/2)/(I*a*tan(f*x
+ e) + a)^(5/2), x)

```

3.844.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{7/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \int \frac{(A + B \tan(e + fx)) (c - c \tan(e + fx) \text{ li})^{7/2}}{(a + a \tan(e + fx) \text{ li})^{5/2}} dx$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(7/2))/(a + a*tan(e + f*x)*1i)^(5/2),x)`

output `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(7/2))/(a + a*tan(e + f*x)*1i)^(5/2), x)`

3.845
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^{5/2}} dx$$

3.845.1 Optimal result	7580
3.845.2 Mathematica [C] (verified)	7580
3.845.3 Rubi [A] (verified)7581
3.845.4 Maple [B] (verified)7584
3.845.5 Fricas [B] (verification not implemented)7585
3.845.6 Sympy [F]7586
3.845.7 Maxima [A] (verification not implemented)7586
3.845.8 Giac [F]7587
3.845.9 Mupad [F(-1)]7587

3.845.1 Optimal result

Integrand size = 45, antiderivative size = 205

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^{5/2}} dx =$$

$$-\frac{2Bc^{5/2} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{a^{5/2}f} - \frac{2Bc^2 \sqrt{c - ic \tan(e + fx)}}{a^2 f \sqrt{a + ia \tan(e + fx)}}$$

$$+ \frac{2Bc(c - ic \tan(e + fx))^{3/2}}{3af(a + ia \tan(e + fx))^{3/2}} + \frac{(iA - B)(c - ic \tan(e + fx))^{5/2}}{5f(a + ia \tan(e + fx))^{5/2}}$$

```
output -2*B*c^(5/2)*arctan(c^(1/2)*(a+I*a*tan(f*x+e))^(1/2)/a^(1/2)/(c-I*c*tan(f*x+e))^(1/2))/a^(5/2)/f-2*B*c^2*(c-I*c*tan(f*x+e))^(1/2)/a^2/f/(a+I*a*tan(f*x+e))^(1/2)+2/3*B*c*(c-I*c*tan(f*x+e))^(3/2)/a/f/(a+I*a*tan(f*x+e))^(3/2)+1/5*(I*A-B)*(c-I*c*tan(f*x+e))^(5/2)/f/(a+I*a*tan(f*x+e))^(5/2)
```

3.845.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 7.65 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.81

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^{5/2}} dx = \frac{c^3(i + \tan(e + fx)) \left(-20\sqrt{2}B \text{Hypergeometric2F1}(-\dots)\right)}{15a^2 f \sqrt{1 - i \tan(e + fx)}}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2))/(a + I*a*Tan[e + f*x])^(5/2),x]`

output `(c^3*(I + Tan[e + f*x])*(-20*Sqrt[2]*B*Hypergeometric2F1[-3/2, -3/2, -1/2, (1 + I*Tan[e + f*x])/2]*(-I + Tan[e + f*x])) + 3*(A + I*B)*Sqrt[1 - I*Tan[e + f*x]]*(I + Tan[e + f*x])^2)/(15*a^2*f*Sqrt[1 - I*Tan[e + f*x]]*(-I + Tan[e + f*x])^2*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])`

3.845.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.156$, Rules used = {3042, 4071, 87, 57, 57, 45, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - ic \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(c - ic \tan(e + fx))^{5/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(i \tan(e + fx)a + a)^{7/2}} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{87} \\
 & ac \left(\frac{(-B + iA)(c - ic \tan(e + fx))^{5/2}}{5ac(a + ia \tan(e + fx))^{5/2}} - \frac{iB \int \frac{(c - ic \tan(e + fx))^{3/2}}{(i \tan(e + fx)a + a)^{5/2}} d \tan(e + fx)}{a} \right) \\
 & \quad \downarrow \text{57} \\
 & ac \left(\frac{(-B + iA)(c - ic \tan(e + fx))^{5/2}}{5ac(a + ia \tan(e + fx))^{5/2}} - \frac{iB \left(\frac{2i(c - ic \tan(e + fx))^{3/2}}{3a(a + ia \tan(e + fx))^{3/2}} - \frac{c \int \frac{\sqrt{c - ic \tan(e + fx)}}{(i \tan(e + fx)a + a)^{3/2}} d \tan(e + fx)}{a} \right)}{a} \right)
 \end{aligned}$$

3.845. $\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^{5/2}} dx$

↓ 57

$$aC \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{iB \left(\frac{2i(c-ic \tan(e+fx))^{3/2}}{3a(a+ia \tan(e+fx))^{3/2}} - \frac{c \int \frac{1}{\sqrt{i \tan(e+fx)a+a\sqrt{c-ic \tan(e+fx)}} d \tan(e+fx)}{a} \right)}{a} \right)$$

f

↓ 45

$$aC \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{iB \left(\frac{2i(c-ic \tan(e+fx))^{3/2}}{3a(a+ia \tan(e+fx))^{3/2}} - \frac{c \int \frac{1}{ia + \frac{ic(i \tan(e+fx)a+a)}{c-ic \tan(e+fx)}} d \frac{\sqrt{i \tan(e+fx)a+a}}{\sqrt{c-ic \tan(e+fx)}}}{a} \right)}{a} \right)$$

f

↓ 218

$$aC \left(\frac{(-B+iA)(c-ic \tan(e+fx))^{5/2}}{5ac(a+ia \tan(e+fx))^{5/2}} - \frac{iB \left(\frac{2i(c-ic \tan(e+fx))^{3/2}}{3a(a+ia \tan(e+fx))^{3/2}} - \frac{c \left(\frac{2i\sqrt{c} \arctan\left(\frac{\sqrt{c}\sqrt{a+ia \tan(e+fx)}}{\sqrt{a}\sqrt{c-ic \tan(e+fx)}}\right)}{a^{3/2}} + \frac{2i\sqrt{c-ic \tan(e+fx)}}{a\sqrt{a+ia \tan(e+fx)}} \right)}{a} \right)}{a} \right)$$

f

```
input Int[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(5/2))/(a + I*a*Tan[e + f*x])^(5/2), x]
```

3.845. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^{5/2}} dx$

output $(a*c*((I*A - B)*(c - I*c*\text{Tan}[e + f*x])^{(5/2)})/(5*a*c*(a + I*a*\text{Tan}[e + f*x])^{(5/2)}) - (I*B*(((2*I)/3)*(c - I*c*\text{Tan}[e + f*x])^{(3/2)})/(a*(a + I*a*\text{Tan}[e + f*x])^{(3/2)}) - (c*(((2*I)*\text{Sqrt}[c]*\text{ArcTan}[(\text{Sqrt}[c]*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]])/(\text{Sqrt}[a]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])]))/a^{(3/2)} + ((2*I)*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])/(a*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]])))/a)/a)/f$

3.845.3.1 Defintions of rubi rules used

rule 45 $\text{Int}[1/(\text{Sqrt}[(a_) + (b_)*(x_)]*\text{Sqrt}[(c_) + (d_)*(x_)]), x_Symbol] \rightarrow \text{Simp}[2 \text{ Subst}[\text{Int}[1/(b - d*x^2), x], x, \text{Sqrt}[a + b*x]/\text{Sqrt}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{EqQ}[b*c + a*d, 0] \&\& !\text{GtQ}[c, 0]$

rule 57 $\text{Int}[(a_) + (b_)*(x_)]^{(m_)}*((c_) + (d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + 1))), x] - \text{Simp}[d*(n/(b*(m + 1))) \text{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^{(n - 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{GtQ}[n, 0] \&\& \text{LtQ}[m, -1] \&\& !(IntegerQ[n] \&\& !IntegerQ[m]) \&\& !(ILeQ[m + n + 2, 0] \&\& (\text{FractionQ}[m] || \text{GeQ}[2*n + m + 1, 0])) \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

rule 87 $\text{Int}[(a_) + (b_)*(x_)]*((c_) + (d_)*(x_)]^{(n_)}*((e_) + (f_)*(x_)]^{(p_)}, x_] \rightarrow \text{Simp}[(-b*e - a*f)*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)})/(f*(p + 1)*(c*f - d*e)), x] - \text{Simp}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) \text{Int}[(c + d*x)^n*(e + f*x)^{(p + 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \&\& \text{LtQ}[p, -1] \&\& (!\text{LtQ}[n, -1] || \text{IntegerQ}[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || \text{LtQ}[p, n])))$

rule 218 $\text{Int}[(a_) + (b_)*(x_)^2]^{(-1)}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b]$

rule 3042 $\text{Int}[u_, x_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.845.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 508 vs. 2(168) = 336.

Time = 0.41 (sec) , antiderivative size = 509, normalized size of antiderivative = 2.48

method	result
parts	$\frac{A\sqrt{-c(i\tan(fx+e)-1)}\sqrt{a(1+i\tan(fx+e))}c^2(1+\tan(fx+e)^2)(i+\tan(fx+e))}{5fa^3(i-\tan(fx+e))^4} + \frac{iB\sqrt{-c(i\tan(fx+e)-1)}\sqrt{a(1+i\tan(fx+e))}}{5fa^3(i-\tan(fx+e))^4}$
derivativedivides	$\frac{\sqrt{-c(i\tan(fx+e)-1)}\sqrt{a(1+i\tan(fx+e))}c^2\left(-15iB\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}}\right)\right)}{5fa^3(i-\tan(fx+e))^4} + 90iB\sqrt{-c(i\tan(fx+e)-1)}\sqrt{a(1+i\tan(fx+e))}c^2\left(-15iB\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}}\right)\right)$
default	$\frac{\sqrt{-c(i\tan(fx+e)-1)}\sqrt{a(1+i\tan(fx+e))}c^2\left(-15iB\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}}\right)\right)}{5fa^3(i-\tan(fx+e))^4} + 90iB\sqrt{-c(i\tan(fx+e)-1)}\sqrt{a(1+i\tan(fx+e))}c^2\left(-15iB\ln\left(\frac{ac\tan(fx+e)+\sqrt{ac}\sqrt{ac(1+\tan(fx+e)^2)}}{\sqrt{ac}}\right)\right)$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^(5/2),x,m
ethod=_RETURNVERBOSE)
```

$$3.845. \int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^{5/2}} dx$$

output
$$\frac{1}{5}A/f*(-c*(I*\tan(f*x+e)-1))^{(1/2)}*(a*(1+I*\tan(f*x+e)))^{(1/2)}*c^2/a^3*(1+\tan(f*x+e)^2)*(I+\tan(f*x+e))/(I-\tan(f*x+e))^4+1/15*I*B/f*(-c*(I*\tan(f*x+e)-1))^{(1/2)}*(a*(1+I*\tan(f*x+e)))^{(1/2)}*c^2/a^3*(60*I*\ln((a*c*\tan(f*x+e)+(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)})/(a*c)^{(1/2)})*\tan(f*x+e)^3*a*c-15*\ln((a*c*\tan(f*x+e)+(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)})/(a*c)^{(1/2)})*\tan(f*x+e)^4*a*c-60*I*\ln((a*c*\tan(f*x+e)+(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)})/(a*c)^{(1/2)})*\tan(f*x+e)*a*c-97*I*(a*c*(1+\tan(f*x+e)^2))^{(1/2)}*(a*c)^{(1/2)}*\tan(f*x+e)^2+90*\ln((a*c*\tan(f*x+e)+(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)})/(a*c)^{(1/2)})*\tan(f*x+e)^2*a*c+43*\tan(f*x+e)^3*(a*c*(1+\tan(f*x+e)^2))^{(1/2)}*(a*c)^{(1/2)}+23*I*(a*c*(1+\tan(f*x+e)^2))^{(1/2)}*(a*c)^{(1/2)}-15*a*c*\ln((a*c*\tan(f*x+e)+(a*c)^{(1/2)}*(a*c*(1+\tan(f*x+e)^2))^{(1/2)})/(a*c)^{(1/2)})-77*\tan(f*x+e)*(a*c*(1+\tan(f*x+e)^2))^{(1/2)}*(a*c)^{(1/2)})/(a*c*(1+\tan(f*x+e)^2))^{(1/2)}/(I-\tan(f*x+e))^4/(a*c)^{(1/2)}$$

3.845.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 436 vs. $2(157) = 314$.

Time = 0.29 (sec) , antiderivative size = 436, normalized size of antiderivative = 2.13

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^{5/2}} dx = \frac{\left(15 a^3 f \sqrt{-\frac{B^2 c^5}{a^5 f^2}} e^{(5i f x + 5i e)} \log \left(\frac{4 \left(2 (B c^2 e^{(3i f x + 3i e)} + B c^2 e^{(3i f x + 3i e)} \right)}{\dots} \right)}{\dots} \right)}{\dots}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="fracas")`

output
$$\frac{1}{30}*(15*a^3*f*\sqrt{-B^2*c^5/(a^5*f^2)})*e^{(5*I*f*x + 5*I*e)}*\log(4*(2*(B*c^2*e^{(3*I*f*x + 3*I*e)} + B*c^2*e^{(I*f*x + I*e)})*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} + (a^3*f*e^{(2*I*f*x + 2*I*e)} - a^3*f)*\sqrt{-B^2*c^5/(a^5*f^2)}))/(B*c^2*e^{(2*I*f*x + 2*I*e)} + B*c^2) - 15*a^3*f*\sqrt{-B^2*c^5/(a^5*f^2)})*e^{(5*I*f*x + 5*I*e)}*\log(4*(2*(B*c^2*e^{(3*I*f*x + 3*I*e)} + B*c^2*e^{(I*f*x + I*e)})*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)} - (a^3*f*e^{(2*I*f*x + 2*I*e)} - a^3*f)*\sqrt{-B^2*c^5/(a^5*f^2)}))/(B*c^2*e^{(2*I*f*x + 2*I*e)} + B*c^2) - 2*(30*B*c^2*e^{(6*I*f*x + 6*I*e)} + 20*B*c^2*e^{(4*I*f*x + 4*I*e)} + (-3*I*A - 7*B)*c^2*e^{(2*I*f*x + 2*I*e)} + 3*(-I*A + B)*c^2)*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)))*e^{(-5*I*f*x - 5*I*e)}/(a^3*f)$$

3.845.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{5/2}}{(a+ia \tan(e+fx))^{5/2}} dx$$

3.845.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^{5/2}} dx = \int \frac{(-ic(\tan(e + fx) + i))^{5/2} (A + B \tan(e + fx))}{(ia(\tan(e + fx) - i))^{5/2}} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(5/2)/(a+I*a*tan(f*x+e))**(5/2),x)`

output `Integral((-I*c*(tan(e + f*x) + I))**(5/2)*(A + B*tan(e + f*x))/(I*a*(tan(e + f*x) - I))**(5/2), x)`

3.845.7 Maxima [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.05

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{5/2}}{(a + ia \tan(e + fx))^{5/2}} dx =$$

$$\frac{(30 Bc^2 \arctan(\cos(fx + e), \sin(fx + e) + 1) + 30 Bc^2 \arctan(\cos(fx + e), -\sin(fx + e) + 1) - 6(iA - B)c^2 \cos(5fx + 5e) - 20 Bc^2 \cos(3fx + 3e) + 60 Bc^2 \cos(fx + e) + 15 I Bc^2 \log(\cos(fx + e)^2 + \sin(fx + e)^2 + 2 \sin(fx + e) + 1) - 15 I Bc^2 \log(\cos(fx + e)^2 + \sin(fx + e)^2 - 2 \sin(fx + e) + 1) - 6(A + I B)c^2 \sin(5fx + 5e) + 20 I Bc^2 \sin(3fx + 3e) - 60 I Bc^2 \sin(fx + e)) \sqrt{c}}{(a^2)^{5/2} f}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `-1/30*(30*B*c^2*arctan2(cos(f*x + e), sin(f*x + e) + 1) + 30*B*c^2*arctan2(cos(f*x + e), -sin(f*x + e) + 1) - 6*(I*A - B)*c^2*cos(5*f*x + 5*e) - 20*B*c^2*cos(3*f*x + 3*e) + 60*B*c^2*cos(f*x + e) + 15*I*B*c^2*log(cos(f*x + e)^2 + sin(f*x + e)^2 + 2*sin(f*x + e) + 1) - 15*I*B*c^2*log(cos(f*x + e)^2 + sin(f*x + e)^2 - 2*sin(f*x + e) + 1) - 6*(A + I*B)*c^2*sin(5*f*x + 5*e) + 20*I*B*c^2*sin(3*f*x + 3*e) - 60*I*B*c^2*sin(f*x + e))*sqrt(c)/(a^(5/2)*f)`

3.845.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{5/2}}{(i a \tan(fx + e) + a)^{5/2}} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(5/2)/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(5/2)/(I*a*tan(f*x + e) + a)^(5/2), x)`

3.845.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{5/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \int \frac{(A + B \tan(e + fx)) (c - c \tan(e + fx) 1i)^{5/2}}{(a + a \tan(e + fx) 1i)^{5/2}} dx$$

input `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(5/2))/(a + a*tan(e + f*x)*1i)^(5/2), x)`

output `int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(5/2))/(a + a*tan(e + f*x)*1i)^(5/2), x)`

3.846
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^{5/2}} dx$$

3.846.1 Optimal result	7588
3.846.2 Mathematica [A] (verified)	7588
3.846.3 Rubi [A] (verified)	7589
3.846.4 Maple [A] (verified)	7590
3.846.5 Fricas [A] (verification not implemented)	7591
3.846.6 Sympy [F]	7591
3.846.7 Maxima [A] (verification not implemented)	7592
3.846.8 Giac [F]	7592
3.846.9 Mupad [B] (verification not implemented)	7592

3.846.1 Optimal result

Integrand size = 45, antiderivative size = 104

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^{5/2}} dx = \frac{(iA - B)(c - ic \tan(e + fx))^{3/2}}{5f(a + ia \tan(e + fx))^{5/2}} + \frac{(iA + 4B)(c - ic \tan(e + fx))^{3/2}}{15af(a + ia \tan(e + fx))^{3/2}}$$

output `1/5*(I*A-B)*(c-I*c*tan(f*x+e))^(3/2)/f/(a+I*a*tan(f*x+e))^(5/2)+1/15*(I*A+4*B)*(c-I*c*tan(f*x+e))^(3/2)/a/f/(a+I*a*tan(f*x+e))^(3/2)`

3.846.2 Mathematica [A] (verified)

Time = 5.37 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.88

$$\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^{5/2}} dx = \frac{ic(i + \tan(e + fx))(-4iA - B + (A - 4iB) \tan(e + fx))\sqrt{c - ic \tan(e + fx)}}{15a^2 f(-i + \tan(e + fx))^2 \sqrt{a + ia \tan(e + fx)}}$$

input `Integrate[((A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(3/2))/(a + I*a*Tan[e + f*x])^(5/2),x]`

output $((-1/15*I)*c*(I + \text{Tan}[e + f*x])*((-4*I)*A - B + (A - (4*I)*B)*\text{Tan}[e + f*x])*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]]/(a^2*f*(-I + \text{Tan}[e + f*x])^2*\text{Sqrt}[a + I*a*\text{Tan}[e + f*x]])$

3.846.3 Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.09, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.089$, Rules used = {3042, 4071, 87, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^{5/2}} dx \\ & \quad \downarrow 3042 \\ & \int \frac{(c - ic \tan(e + fx))^{3/2} (A + B \tan(e + fx))}{(a + ia \tan(e + fx))^{5/2}} dx \\ & \quad \downarrow 4071 \\ & \frac{ac \int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{(i \tan(e + fx)a + a)^{7/2}} d \tan(e + fx)}{f} \\ & \quad \downarrow 87 \\ & \frac{ac \left(\frac{(A - 4iB) \int \frac{\sqrt{c - ic \tan(e + fx)}}{(i \tan(e + fx)a + a)^{5/2}} d \tan(e + fx)}{5a} + \frac{(-B + iA)(c - ic \tan(e + fx))^{3/2}}{5ac(a + ia \tan(e + fx))^{5/2}} \right)}{f} \\ & \quad \downarrow 48 \\ & \frac{ac \left(\frac{i(A - 4iB)(c - ic \tan(e + fx))^{3/2}}{15a^2c(a + ia \tan(e + fx))^{3/2}} + \frac{(-B + iA)(c - ic \tan(e + fx))^{3/2}}{5ac(a + ia \tan(e + fx))^{5/2}} \right)}{f} \end{aligned}$$

input $\text{Int}[(A + B*\text{Tan}[e + f*x])*(c - I*c*\text{Tan}[e + f*x])^(3/2)/(a + I*a*\text{Tan}[e + f*x])^(5/2), x]$

output $(a*c*(((I*A - B)*(c - I*c*\text{Tan}[e + f*x])^(3/2))/(5*a*c*(a + I*a*\text{Tan}[e + f*x])^(5/2)) + ((I/15)*(A - (4*I)*B)*(c - I*c*\text{Tan}[e + f*x])^(3/2))/(a^2*c*(a + I*a*\text{Tan}[e + f*x])^(3/2))))/f$

3.846. $\int \frac{(A + B \tan(e + fx))(c - ic \tan(e + fx))^{3/2}}{(a + ia \tan(e + fx))^{5/2}} dx$

3.846.3.1 Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*(c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.846.4 Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.88

method	result
derivativedivides	$\frac{i\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c(1+\tan(fx+e)^2) (iA \tan(fx+e)-iB+4B \tan(fx+e)+4A)}{15f a^3(i-\tan(fx+e))^4}$
default	$\frac{i\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c(1+\tan(fx+e)^2) (iA \tan(fx+e)-iB+4B \tan(fx+e)+4A)}{15f a^3(i-\tan(fx+e))^4}$
parts	$\frac{A\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} c(1+\tan(fx+e)^2) (4i-\tan(fx+e))}{15f a^3(i-\tan(fx+e))^4} - \frac{iB\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))}}{15f}$

```
input int((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^(5/2),x,m
ethod=_RETURNVERBOSE)
```

3.846.
$$\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^{5/2}} dx$$

output $1/15*I/f*(-c*(I*\tan(f*x+e)-1))^(1/2)*(a*(1+I*\tan(f*x+e)))^(1/2)/a^3*c*(1+\tan(f*x+e)^2)*(I*A*\tan(f*x+e)-I*B+4*B*\tan(f*x+e)+4*A)/(I-\tan(f*x+e))^4$

3.846.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.94

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \frac{(5(-iA - B)ce^{(4i fx + 4i e)} + 2(-4iA - B)ce^{(2i fx + 2i e)} + 3(-iA + B)c) \sqrt{\frac{a}{e^{(2i fx + 2i e)} + 1}} \sqrt{\frac{c}{e^{(2i fx + 2i e)} + 1}} e^{(-5i fx)}}{30 a^3 f}$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="fracas")`

output $-1/30*(5*(-I*A - B)*c*e^{(4*I*f*x + 4*I*e)} + 2*(-4*I*A - B)*c*e^{(2*I*f*x + 2*I*e)} + 3*(-I*A + B)*c)*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)}*e^{(-5*I*f*x - 5*I*e)}/(a^3*f)$

3.846.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \int \frac{(-i c (\tan(e + fx) + i))^{3/2} (A + B \tan(e + fx))}{(i a (\tan(e + fx) - i))^{5/2}} dx$$

input `integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(3/2)/(a+I*a*tan(f*x+e))**(5/2),x)`

output `Integral((-I*c*(tan(e + f*x) + I))**(3/2)*(A + B*tan(e + f*x))/(I*a*(tan(e + f*x) - I))**(5/2), x)`

3.846.7 Maxima [A] (verification not implemented)

Time = 0.43 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.47

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \frac{30(5(A - i B)c \cos(4fx + 4e) + 2(4A - i B)c \cos(2fx + 2e) - 5(-iA - B)c \sin(4fx + 4e) - 2(-4iA - B)c \sin(2fx + 2e) + 3(A + iB)c) \sqrt{a} \sqrt{c}}{-900(i a^3 \cos(7fx + 7e) - 900i a^3 \cos(5fx + 5e) + 900a^3 \sin(7fx + 7e) + 900a^3 \sin(5fx + 5e)) * f}$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="maxima")
```

```
output 30*(5*(A - I*B)*c*cos(4*f*x + 4*e) + 2*(4*A - I*B)*c*cos(2*f*x + 2*e) - 5*(-I*A - B)*c*sin(4*f*x + 4*e) - 2*(-4*I*A - B)*c*sin(2*f*x + 2*e) + 3*(A + I*B)*c)*sqrt(a)*sqrt(c)/((-900*I*a^3*cos(7*f*x + 7*e) - 900*I*a^3*cos(5*f*x + 5*e) + 900*a^3*sin(7*f*x + 7*e) + 900*a^3*sin(5*f*x + 5*e))*f)
```

3.846.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \int \frac{(B \tan(fx + e) + A)(-i c \tan(fx + e) + c)^{3/2}}{(i a \tan(fx + e) + a)^{5/2}} dx$$

```
input integrate((A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(3/2)/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="giac")
```

```
output integrate((B*tan(f*x + e) + A)*(-I*c*tan(f*x + e) + c)^(3/2)/(I*a*tan(f*x + e) + a)^(5/2), x)
```

3.846.9 Mupad [B] (verification not implemented)

Time = 10.10 (sec) , antiderivative size = 240, normalized size of antiderivative = 2.31

$$\int \frac{(A + B \tan(e + fx))(c - i c \tan(e + fx))^{3/2}}{(a + i a \tan(e + fx))^{5/2}} dx = \frac{c \sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}} \sqrt{\frac{c(\cos(2e+2fx)+1-\sin(2e+2fx)1i)}{\cos(2e+2fx)+1}}}{(a + i a \tan(e + fx))^{5/2}}$$

```
input int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(3/2))/(a + a*tan(e + f*x)*1i)^(5/2),x)
```

3.846. $\int \frac{(A+B \tan(e+fx))(c-ic \tan(e+fx))^{3/2}}{(a+ia \tan(e+fx))^{5/2}} dx$

output

```
(c*((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*cos(2*e + 2*f*x)*5i + A*cos(4*e + 4*f*x)*8i + A*cos(6*e + 6*f*x)*3i + 5*B*cos(2*e + 2*f*x) + 2*B*cos(4*e + 4*f*x) - 3*B*cos(6*e + 6*f*x) + 5*A*sin(2*e + 2*f*x) + 8*A*sin(4*e + 4*f*x) + 3*A*sin(6*e + 6*f*x) - B*sin(2*e + 2*f*x)*5i - B*sin(4*e + 4*f*x)*2i + B*sin(6*e + 6*f*x)*3i)/(60*a^3*f)
```


$$3.847 \quad \int \frac{(A+B \tan(e+fx))\sqrt{c-ictan(e+fx)}}{(a+ia \tan(e+fx))^{5/2}} dx$$

3.847.1 Optimal result	7594
3.847.2 Mathematica [A] (verified)	7594
3.847.3 Rubi [A] (verified)	7595
3.847.4 Maple [A] (verified)	7597
3.847.5 Fricas [A] (verification not implemented)	7597
3.847.6 Sympy [F]	7598
3.847.7 Maxima [F(-2)]	7598
3.847.8 Giac [F]	7598
3.847.9 Mupad [B] (verification not implemented)	7599

3.847.1 Optimal result

Integrand size = 45, antiderivative size = 157

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)}}{(a + ia \tan(e + fx))^{5/2}} dx = \frac{(iA - B)\sqrt{c - ictan(e + fx)}}{5f(a + ia \tan(e + fx))^{5/2}} + \frac{(2iA + 3B)\sqrt{c - ictan(e + fx)}}{15af(a + ia \tan(e + fx))^{3/2}} + \frac{(2iA + 3B)\sqrt{c - ictan(e + fx)}}{15a^2f\sqrt{a + ia \tan(e + fx)}}$$

output `1/15*(2*I*A+3*B)*(c-I*c*tan(f*x+e))^(1/2)/a^2/f/(a+I*a*tan(f*x+e))^(1/2)+1/5*(I*A-B)*(c-I*c*tan(f*x+e))^(1/2)/f/(a+I*a*tan(f*x+e))^(5/2)+1/15*(2*I*A+3*B)*(c-I*c*tan(f*x+e))^(1/2)/a/f/(a+I*a*tan(f*x+e))^(3/2)`

3.847.2 Mathematica [A] (verified)

Time = 3.53 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.63

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)}}{(a + ia \tan(e + fx))^{5/2}} dx = \frac{\sqrt{c - ictan(e + fx)}(-7iA - 3B + (6A - 9iB) \tan(e + fx))}{15a^2f(-i + \tan(e + fx))^2\sqrt{a + ia \tan(e + fx)}}$$

input `Integrate[((A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]])/(a + I*a*Tan[e + f*x])^(5/2),x]`

output `(Sqrt[c - I*c*Tan[e + f*x]]*((-7*I)*A - 3*B + (6*A - (9*I)*B)*Tan[e + f*x] + ((2*I)*A + 3*B)*Tan[e + f*x]^2))/(15*a^2*f*(-I + Tan[e + f*x])^2*Sqrt[a + I*a*Tan[e + f*x]])`

3.847. $\int \frac{(A+B \tan(e+fx))\sqrt{c-ictan(e+fx)}}{(a+ia \tan(e+fx))^{5/2}} dx$

3.847.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.08, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {3042, 4071, 87, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{c - ic \tan(e + fx)}(A + B \tan(e + fx))}{(a + ia \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{\sqrt{c - ic \tan(e + fx)}(A + B \tan(e + fx))}{(a + ia \tan(e + fx))^{5/2}} dx \\
 & \quad \downarrow \text{4071} \\
 & ac \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx)a + a)^{7/2} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx) \\
 & \quad \downarrow \text{87} \\
 & ac \left(\frac{(2A - 3iB) \int \frac{1}{(i \tan(e + fx)a + a)^{5/2} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{5a} + \frac{(-B + iA) \sqrt{c - ic \tan(e + fx)}}{5ac(a + ia \tan(e + fx))^{5/2}} \right) \\
 & \quad \downarrow \text{55} \\
 & ac \left(\frac{(2A - 3iB) \left(\frac{\int \frac{1}{(i \tan(e + fx)a + a)^{3/2} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{3a} + \frac{i \sqrt{c - ic \tan(e + fx)}}{3ac(a + ia \tan(e + fx))^{3/2}} \right)}{5a} + \frac{(-B + iA) \sqrt{c - ic \tan(e + fx)}}{5ac(a + ia \tan(e + fx))^{5/2}} \right) \\
 & \quad \downarrow \text{48} \\
 & ac \left(\frac{(2A - 3iB) \left(\frac{i \sqrt{c - ic \tan(e + fx)}}{3a^2 \sqrt{a + ia \tan(e + fx)}} + \frac{i \sqrt{c - ic \tan(e + fx)}}{3ac(a + ia \tan(e + fx))^{3/2}} \right)}{5a} + \frac{(-B + iA) \sqrt{c - ic \tan(e + fx)}}{5ac(a + ia \tan(e + fx))^{5/2}} \right)
 \end{aligned}$$

input `Int[((A + B*Tan[e + f*x])*Sqrt[c - I*c*Tan[e + f*x]]/(a + I*a*Tan[e + f*x]))^(5/2),x]`

$$3.847. \quad \int \frac{(A + B \tan(e + fx)) \sqrt{c - ic \tan(e + fx)}}{(a + ia \tan(e + fx))^{5/2}} dx$$

```
output (a*c*((I*A - B)*Sqrt[c - I*c*Tan[e + f*x]])/(5*a*c*(a + I*a*Tan[e + f*x])
^(5/2)) + ((2*A - (3*I)*B)*(((I/3)*Sqrt[c - I*c*Tan[e + f*x]])/(a*c*(a + I
*a*Tan[e + f*x])^(3/2)) + ((I/3)*Sqrt[c - I*c*Tan[e + f*x]]/(a^2*c*Sqrt[a
+ I*a*Tan[e + f*x]])))/(5*a))/f
```

3.847.3.1 Defintions of rubi rules used

```
rule 48 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]
```

```
rule 55 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.847.4 Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.81

method	result
derivativedivides	$-\frac{i\sqrt{-c(i\tan(fx+e)-1)}\sqrt{a(1+i\tan(fx+e))}\left(2iA\tan(fx+e)^3-12iB\tan(fx+e)^2+3B\tan(fx+e)^3-13iA\tan(fx+e)\right)}{15fa^3(i-\tan(fx+e))^4}$
default	$-\frac{i\sqrt{-c(i\tan(fx+e)-1)}\sqrt{a(1+i\tan(fx+e))}\left(2iA\tan(fx+e)^3-12iB\tan(fx+e)^2+3B\tan(fx+e)^3-13iA\tan(fx+e)\right)}{15fa^3(i-\tan(fx+e))^4}$
parts	$-\frac{A\sqrt{-c(i\tan(fx+e)-1)}\sqrt{a(1+i\tan(fx+e))}\left(8i\tan(fx+e)^2-2\tan(fx+e)^3-7i+13\tan(fx+e)\right)}{15fa^3(i-\tan(fx+e))^4} + \frac{iB\sqrt{-c(i\tan(fx+e)-1)}\sqrt{a(1+i\tan(fx+e))}}{15fa^3(i-\tan(fx+e))^4}$

```
input int((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(5/2),x,m
method=_RETURNVERBOSE)
```

```
output -1/15*I/f*(-c*(I*tan(f*x+e)-1))^(1/2)*(a*(1+I*tan(f*x+e)))^(1/2)/a^3*(2*I*
A*tan(f*x+e)^3-12*I*B*tan(f*x+e)^2+3*B*tan(f*x+e)^3-13*I*A*tan(f*x+e)+8*A*
tan(f*x+e)^2+3*I*B-12*B*tan(f*x+e)-7*A)/(I-tan(f*x+e))^4
```

3.847.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.71

$$\int \frac{(A + B \tan(e + fx))\sqrt{c - ictan(e + fx)}}{(a + ia \tan(e + fx))^{5/2}} dx =$$

$$\frac{(15(-iA - B)e^{(6i fx + 6ie)} + 5(-5iA - 3B)e^{(4i fx + 4ie)} - (13iA - 3B)e^{(2i fx + 2ie)} - 3iA + 3B)\sqrt{\frac{a}{e^{(2i fx + 2ie)}}}}{60a^3 f}$$

```
input integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(5/
2),x, algorithm="fracas")
```

```
output -1/60*(15*(-I*A - B)*e^(6*I*f*x + 6*I*e) + 5*(-5*I*A - 3*B)*e^(4*I*f*x + 4
*I*e) - (13*I*A - 3*B)*e^(2*I*f*x + 2*I*e) - 3*I*A + 3*B)*sqrt(a/(e^(2*I*f
*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*e^(-5*I*f*x - 5*I*e)/(
a^3*f)
```

3.847. $\int \frac{(A+B \tan(e+fx))\sqrt{c-ictan(e+fx)}}{(a+ia \tan(e+fx))^{5/2}} dx$

3.847.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{(a + i a \tan(e + fx))^{5/2}} dx = \int \frac{\sqrt{-i c (\tan(e + fx) + i)} (A + B \tan(e + fx))}{(i a (\tan(e + fx) - i))^{5/2}} dx$$

input `integrate((c-I*c*tan(f*x+e))**(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**(5/2),x)`

output `Integral(sqrt(-I*c*(tan(e + f*x) + I))*(A + B*tan(e + f*x))/(I*a*(tan(e + f*x) - I))**(5/2), x)`

3.847.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{(a + i a \tan(e + fx))^{5/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.847.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i c \tan(e + fx)}}{(a + i a \tan(e + fx))^{5/2}} dx = \int \frac{(B \tan(fx + e) + A) \sqrt{-i c \tan(fx + e) + c}}{(i a \tan(fx + e) + a)^{5/2}} dx$$

input `integrate((c-I*c*tan(f*x+e))^(1/2)*(A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*sqrt(-I*c*tan(f*x + e) + c)/(I*a*tan(f*x + e) + a)^(5/2), x)`

3.847. $\int \frac{(A+B \tan(e+fx)) \sqrt{c-i c \tan(e+fx)}}{(a+i a \tan(e+fx))^{5/2}} dx$

3.847.9 Mupad [B] (verification not implemented)

Time = 10.32 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.57

$$\int \frac{(A + B \tan(e + fx)) \sqrt{c - i \tan(e + fx)}}{(a + i a \tan(e + fx))^{5/2}} dx = \sqrt{\frac{a(\cos(2e + 2fx) + 1 + \sin(2e + 2fx) i)}{\cos(2e + 2fx) + 1}} \sqrt{\frac{c(\cos(2e + 2fx) + 1 - \sin(2e + 2fx) i)}{\cos(2e + 2fx) + 1}}$$

```
input int(((A + B*tan(e + f*x))*(c - c*tan(e + f*x)*1i)^(1/2))/(a + a*tan(e + f*x)*1i)^(5/2),x)
```

```
output (((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*15i + 15*B + A*cos(2*e + 2*f*x)*25i + A*cos(4*e + 4*f*x)*13i + A*cos(6*e + 6*f*x)*3i + 15*B*cos(2*e + 2*f*x) - 3*B*cos(4*e + 4*f*x) - 3*B*cos(6*e + 6*f*x) + 25*A*sin(2*e + 2*f*x) + 13*A*sin(4*e + 4*f*x) + 3*A*sin(6*e + 6*f*x) - B*sin(2*e + 2*f*x)*15i + B*sin(4*e + 4*f*x)*3i + B*sin(6*e + 6*f*x)*3i))/(120*a^3*f)
```

$$3.848 \quad \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}} dx$$

3.848.1 Optimal result	7600
3.848.2 Mathematica [A] (verified)	7600
3.848.3 Rubi [A] (verified)	7601
3.848.4 Maple [A] (verified)	7603
3.848.5 Fricas [A] (verification not implemented)	7604
3.848.6 Sympy [F]	7604
3.848.7 Maxima [F(-2)]	7605
3.848.8 Giac [F]	7605
3.848.9 Mupad [B] (verification not implemented)	7605

3.848.1 Optimal result

Integrand size = 45, antiderivative size = 212

$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}} dx =$$

$$-\frac{iA+B}{f(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}} + \frac{(3iA+2B)\sqrt{c-ic \tan(e+fx)}}{5cf(a+ia \tan(e+fx))^{5/2}}$$

$$+ \frac{2(3iA+2B)\sqrt{c-ic \tan(e+fx)}}{15acf(a+ia \tan(e+fx))^{3/2}} + \frac{2(3iA+2B)\sqrt{c-ic \tan(e+fx)}}{15a^2cf\sqrt{a+ia \tan(e+fx)}}$$

output $2/15*(3*I*A+2*B)*(c-I*c*\tan(f*x+e))^{(1/2)}/a^2/c/f/(a+I*a*\tan(f*x+e))^{(1/2)}$
 $+(-I*A-B)/f/(c-I*c*\tan(f*x+e))^{(1/2)}/(a+I*a*\tan(f*x+e))^{(5/2)}+1/5*(3*I*A+2$
 $*B)*(c-I*c*\tan(f*x+e))^{(1/2)}/c/f/(a+I*a*\tan(f*x+e))^{(5/2)}+2/15*(3*I*A+2*B)$
 $*c-I*c*\tan(f*x+e))^{(1/2)}/a/c/f/(a+I*a*\tan(f*x+e))^{(3/2)}$

3.848.2 Mathematica [A] (verified)

Time = 4.51 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.54

$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}} dx = \frac{-6iA+B+(-3A+2iB)\tan(e+fx)+(-12iA-8B)\tan^2(e+fx)}{15a^2f(-i+\tan(e+fx))^2\sqrt{a+ia \tan(e+fx)}}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^(5/2)*Sqrt[c - I*c*Tan[e + f*x]]),x]`

3.848. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}} dx$

output $((-6*I)*A + B + (-3*A + (2*I)*B)*\text{Tan}[e + f*x] + ((-12*I)*A - 8*B)*\text{Tan}[e + f*x]^2 + (6*A - (4*I)*B)*\text{Tan}[e + f*x]^3)/(15*a^2*f*(-I + \text{Tan}[e + f*x])^2*\text{qrt}[a + I*a*\text{Tan}[e + f*x]]*\text{Sqrt}[c - I*c*\text{Tan}[e + f*x]])$

3.848.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.02, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 87, 55, 55, 48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} \sqrt{c - ic \tan(e + fx)}} dx$$

↓ 3042

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} \sqrt{c - ic \tan(e + fx)}} dx$$

↓ 4071

$$ac \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx)a + a)^{7/2} (c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)$$

f
↓ 87

$$ac \left(\frac{(3A - 2iB) \int \frac{1}{(i \tan(e + fx)a + a)^{7/2} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{c} - \frac{B + iA}{ac(a + ia \tan(e + fx))^{5/2} \sqrt{c - ic \tan(e + fx)}} \right)$$

f
↓ 55

$$ac \left(\frac{(3A - 2iB) \left(\frac{\int \frac{1}{(i \tan(e + fx)a + a)^{5/2} \sqrt{c - ic \tan(e + fx)}} d \tan(e + fx)}{5a} + \frac{i \sqrt{c - ic \tan(e + fx)}}{5ac(a + ia \tan(e + fx))^{5/2}} \right)}{c} - \frac{B + iA}{ac(a + ia \tan(e + fx))^{5/2} \sqrt{c - ic \tan(e + fx)}} \right)$$

f
↓ 55

3.848. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} \sqrt{c - ic \tan(e + fx)}} dx$

$$\begin{array}{c}
 \left(\frac{(3A-2iB) \left(\frac{2 \left(\frac{\int \frac{1}{(i \tan(e+fx)a+a)^{3/2} \sqrt{c-ic \tan(e+fx)} d \tan(e+fx)}{3a} + \frac{i \sqrt{c-ic \tan(e+fx)}}{3ac(a+ia \tan(e+fx))^{3/2}} \right)}{5a} + \frac{i \sqrt{c-ic \tan(e+fx)}}{5ac(a+ia \tan(e+fx))^{5/2}} \right)}{c} - \frac{1}{ac(a+ia \tan(e+fx))} \right)}{f} \\
 \downarrow 48 \\
 \left(\frac{(3A-2iB) \left(\frac{2 \left(\frac{i \sqrt{c-ic \tan(e+fx)}}{3a^2 c \sqrt{a+ia \tan(e+fx)}} + \frac{i \sqrt{c-ic \tan(e+fx)}}{3ac(a+ia \tan(e+fx))^{3/2}} \right)}{5a} + \frac{i \sqrt{c-ic \tan(e+fx)}}{5ac(a+ia \tan(e+fx))^{5/2}} \right)}{c} - \frac{B+iA}{ac(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}} \right)}{f}
 \end{array}$$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^(5/2)*Sqrt[c - I*c*Tan[e + f*x]]),x]`

output `(a*c*(-((I*A + B)/(a*c*(a + I*a*Tan[e + f*x])^(5/2)*Sqrt[c - I*c*Tan[e + f*x]])) + ((3*A - (2*I)*B)*(((I/5)*Sqrt[c - I*c*Tan[e + f*x]])/(a*c*(a + I*a*Tan[e + f*x])^(5/2)) + (2*((I/3)*Sqrt[c - I*c*Tan[e + f*x]])/(a*c*(a + I*a*Tan[e + f*x])^(3/2)) + ((I/3)*Sqrt[c - I*c*Tan[e + f*x]]/(a^2*c*Sqrt[a + I*a*Tan[e + f*x]]))))/(5*a))/c)/f`

3.848.3.1 Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

$$3.848. \quad \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}} dx$$

```
rule 55 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(
c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n +
2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[
c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimp
lerQ[n, 1])
```

```
rule 87 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4071 Int[((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((A_.) + (B_.)*tan[(e_.) +
(f_.)*(x_)])*((c_) + (d_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.848.4 Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.88

method	result
derivativedivides	$-\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} (4iB \tan(fx+e)^5 + 12iA \tan(fx+e)^4 - 6A \tan(fx+e)^5 + 2iB \tan(fx+e)^3 - 15f a^3 c(i - \tan(fx+e))}{15f a^3 c(i - \tan(fx+e))}$
default	$-\frac{\sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} (4iB \tan(fx+e)^5 + 12iA \tan(fx+e)^4 - 6A \tan(fx+e)^5 + 2iB \tan(fx+e)^3 - 15f a^3 c(i - \tan(fx+e))}{15f a^3 c(i - \tan(fx+e))}$
parts	$-\frac{A \sqrt{-c(i \tan(fx+e)-1)} \sqrt{a(1+i \tan(fx+e))} (4i \tan(fx+e)^4 - 2 \tan(fx+e)^5 + 6i \tan(fx+e)^2 - \tan(fx+e)^3 + 2i + \tan(fx+e))}{5f a^3 c(i - \tan(fx+e))^4 (i + \tan(fx+e))^2}$

3.848.
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}} dx$$

```
input int((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)
```

```
output -1/15/f*(-c*(I*tan(f*x+e)-1))^(1/2)*(a*(1+I*tan(f*x+e)))^(1/2)/a^3/c*(4*I*B*tan(f*x+e)^5+12*I*A*tan(f*x+e)^4-6*A*tan(f*x+e)^5+2*I*B*tan(f*x+e)^3+8*B*tan(f*x+e)^4+18*I*A*tan(f*x+e)^2-3*A*tan(f*x+e)^3-2*I*B*tan(f*x+e)+7*B*tan(f*x+e)^2+6*I*A+3*A*tan(f*x+e)-B)/(I-tan(f*x+e))^4/(I+tan(f*x+e))^2
```

3.848.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.76

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} \sqrt{c - ictan(e + fx)}} dx =$$

$$\frac{(15(iA + B)e^{(8ifx+8ie)} + 8(6iA - B)e^{(7ifx+7ie)} - 30iAe^{(6ifx+6ie)} + 8(6iA - B)e^{(5ifx+5ie)} + 10(-6iA - B)e^{(4ifx+4ie)} + 2(-9iA + 4B)e^{(2ifx+2ie)} - 3(iA + 3B)\sqrt{a/(e^{(2ifx+2ie)} + 1)})\sqrt{c/(e^{(2ifx+2ie)} + 1)})e^{(-5ifx - 5ie)}}{120a^3c}$$

```
input integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="fracas")
```

```
output -1/120*(15*(I*A + B)*e^(8*I*f*x + 8*I*e) + 8*(6*I*A - B)*e^(7*I*f*x + 7*I*e) - 30*I*A*e^(6*I*f*x + 6*I*e) + 8*(6*I*A - B)*e^(5*I*f*x + 5*I*e) + 10*(-6*I*A - B)*e^(4*I*f*x + 4*I*e) + 2*(-9*I*A + 4*B)*e^(2*I*f*x + 2*I*e) - 3*I*A + 3*B)*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*e^(-5*I*f*x - 5*I*e)/(a^3*c*f)
```

3.848.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} \sqrt{c - ictan(e + fx)}} dx = \int \frac{A + B \tan(e + fx)}{(ia(\tan(e + fx) - i))^{5/2} \sqrt{-ic(\tan(e + fx) + i)}} dx$$

```
input integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))**(1/2)/(a+I*a*tan(f*x+e))**(5/2),x)
```

```
output Integral((A + B*tan(e + f*x))/((I*a*(tan(e + f*x) - I))**(5/2)*sqrt(-I*c*(tan(e + f*x) + I))), x)
```

3.848. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2} \sqrt{c-ictan(e+fx)}} dx$

3.848.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} \sqrt{c - ic \tan(e + fx)}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.848.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} \sqrt{c - ic \tan(e + fx)}} dx = \int \frac{B \tan(fx + e) + A}{(ia \tan(fx + e) + a)^{5/2} \sqrt{-ic \tan(fx + e) + c}} dx$$

input `integrate((A+B*tan(f*x+e))/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)^(5/2)*sqrt(-I*c*tan(f*x + e) + c)), x)`

3.848.9 Mupad [B] (verification not implemented)

Time = 9.88 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.16

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} \sqrt{c - ic \tan(e + fx)}} dx = \frac{\sqrt{\frac{a(\cos(2e+2fx)+1)+\sin(2e+2fx)li}{\cos(2e+2fx)+1}}}{(15B \cos(2e+2fx) -$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^(5/2)*(c - c*tan(e + f*x)*1i)^(1/2)),x)`

3.848. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}} dx$

output

```
((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))
^(1/2)*(A*cos(2*e + 2*f*x)*45i - 15*B - A*15i + A*cos(4*e + 4*f*x)*15i + A
*cos(6*e + 6*f*x)*3i + 15*B*cos(2*e + 2*f*x) - 5*B*cos(4*e + 4*f*x) - 3*B*
*cos(6*e + 6*f*x) + 45*A*sin(2*e + 2*f*x) + 15*A*sin(4*e + 4*f*x) + 3*A*sin
(6*e + 6*f*x) - B*sin(2*e + 2*f*x)*15i + B*sin(4*e + 4*f*x)*5i + B*sin(6*e
+ 6*f*x)*3i))/(120*a^3*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1)
)/(cos(2*e + 2*f*x) + 1))^(1/2))
```

3.848.
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}} dx$$

$$3.849 \quad \int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2}(c-ictan(e+fx))^{3/2}} dx$$

3.849.1 Optimal result	7607
3.849.2 Mathematica [A] (verified)	7608
3.849.3 Rubi [A] (verified)	7608
3.849.4 Maple [A] (verified)	7611
3.849.5 Fricas [A] (verification not implemented)	7611
3.849.6 Sympy [F]	7612
3.849.7 Maxima [F(-2)]	7612
3.849.8 Giac [F]	7612
3.849.9 Mupad [B] (verification not implemented)	7613

3.849.1 Optimal result

Integrand size = 45, antiderivative size = 218

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2}(c - ictan(e + fx))^{3/2}} dx = \frac{-iA - B}{3f(a + ia \tan(e + fx))^{5/2}(c - ictan(e + fx))^{3/2}}$$

$$+ \frac{4iA + B}{15cf(a + ia \tan(e + fx))^{5/2}\sqrt{c - ictan(e + fx)}}$$

$$+ \frac{4iA + B}{15acf(a + ia \tan(e + fx))^{3/2}\sqrt{c - ictan(e + fx)}}$$

$$+ \frac{2(4A - iB) \tan(e + fx)}{15a^2cf\sqrt{a + ia \tan(e + fx)}\sqrt{c - ictan(e + fx)}}$$

```
output 2/15*(4*A-I*B)*tan(f*x+e)/a^2/c/f/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(1/2)+1/15*(4*I*A+B)/c/f/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^(5/2)+1/15*(4*I*A+B)/a/c/f/(c-I*c*tan(f*x+e))^(1/2)/(a+I*a*tan(f*x+e))^(3/2)+1/3*(-I*A-B)/f/(a+I*a*tan(f*x+e))^(5/2)/(c-I*c*tan(f*x+e))^(3/2)
```

3.849.2 Mathematica [A] (verified)

Time = 6.66 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.70

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2}} dx = \frac{3(A + iB) + (-12iA - 3B) \tan(e + fx) + 3(4A - iB) \tan^2(e + fx) + (-8iA - 2B) \tan^3(e + fx) + (8A - (2i)B) \tan^4(e + fx)}{15a^2 c f (-i + \tan(e + fx))^2 (i + \tan(e + fx))}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e + f*x])^(3/2)),x]`

output `(3*(A + I*B) + ((-12*I)*A - 3*B)*Tan[e + f*x] + 3*(4*A - I*B)*Tan[e + f*x]^2 + ((-8*I)*A - 2*B)*Tan[e + f*x]^3 + (8*A - (2*I)*B)*Tan[e + f*x]^4)/(15*a^2*c*f*(-I + Tan[e + f*x])^2*(I + Tan[e + f*x])*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])`

3.849.3 Rubi [A] (verified)Time = 0.43 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.03, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 87, 55, 55, 41}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2}} dx \\ & \quad \downarrow \text{4071} \\ & ac \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx) a + a)^{7/2} (c - ic \tan(e + fx))^{5/2}} d \tan(e + fx) \\ & \quad \downarrow \text{87} \\ & ac \left(\frac{(4A - iB) \int \frac{1}{(i \tan(e + fx) a + a)^{7/2} (c - ic \tan(e + fx))^{3/2}} d \tan(e + fx)}{3c} - \frac{B + iA}{3ac(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2}} \right) \\ & \quad \downarrow \text{55} \end{aligned}$$

3.849. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2}} dx$

$$ac \left(\frac{(4A-iB) \left(\frac{3 \int \frac{1}{(i \tan(e+fx)a+a)^{5/2}(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{5a} + \frac{i}{5ac(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}} \right)}{3c} \right) - \frac{B+iA}{3ac(a+ia \tan(e+fx))^{5/2}(c-ic \tan(e+fx))} \int f$$

↓ 55

$$ac \left(\frac{(4A-iB) \left(\frac{3 \left(\frac{2 \int \frac{1}{(i \tan(e+fx)a+a)^{3/2}(c-ic \tan(e+fx))^{3/2}} d \tan(e+fx)}{3a} + \frac{i}{3ac(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}} \right)}{5a} \right)}{3c} \right) + \frac{i}{5ac(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}} \int f$$

↓ 41

$$ac \left(\frac{(4A-iB) \left(\frac{3 \left(\frac{2 \tan(e+fx)}{3a^2 c \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}} + \frac{i}{3ac(a+ia \tan(e+fx))^{3/2} \sqrt{c-ic \tan(e+fx)}} \right)}{5a} \right)}{3c} \right) + \frac{i}{5ac(a+ia \tan(e+fx))^{5/2} \sqrt{c-ic \tan(e+fx)}} \int f$$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e + f*x])^(3/2)),x]`

output `(a*c*(-1/3*(I*A + B)/(a*c*(a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e + f*x])^(3/2)) + ((4*A - I*B)*((I/5)/(a*c*(a + I*a*Tan[e + f*x])^(5/2)*Sqrt[c - I*c*Tan[e + f*x]]) + (3*((I/3)/(a*c*(a + I*a*Tan[e + f*x])^(3/2)*Sqrt[c - I*c*Tan[e + f*x]]) + (2*Tan[e + f*x])/(3*a^2*c*Sqrt[a + I*a*Tan[e + f*x]])*Sqrt[c - I*c*Tan[e + f*x])))/(5*a)))/(3*c))/f`

3.849.3.1 Defintions of rubi rules used

rule 41 `Int[1/(((a_) + (b_)*(x_))^(3/2)*((c_) + (d_)*(x_))^(3/2)), x_Symbol] := Simp[x/(a*c*Sqrt[a + b*x]*Sqrt[c + d*x]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0]`

rule 55 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1))) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*(c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.849.4 Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 199, normalized size of antiderivative = 0.91

method	result
derivativedivides	$-\frac{i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(8iA\tan(fx+e)^6-2iB\tan(fx+e)^5+2B\tan(fx+e)^6+20iA\tan(fx+e)^4)}{15fa^3c^2(i-\tan(fx+e))^4}$
default	$-\frac{i\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(8iA\tan(fx+e)^6-2iB\tan(fx+e)^5+2B\tan(fx+e)^6+20iA\tan(fx+e)^4)}{15fa^3c^2(i-\tan(fx+e))^4}$
parts	$-\frac{A\sqrt{a(1+i\tan(fx+e))}\sqrt{-c(i\tan(fx+e)-1)}(8i\tan(fx+e)^5-8\tan(fx+e)^6+20i\tan(fx+e)^3-20\tan(fx+e)^4+12i\tan(fx+e)^5)}{15fa^3c^2(i+\tan(fx+e))^3(i-\tan(fx+e))^4}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(5/2)/(c-I*c*tan(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

output
$$-1/15*I/f*(a*(1+I*\tan(f*x+e)))^{1/2}*(-c*(I*\tan(f*x+e)-1))^{1/2}/a^3/c^2*(8*I*A*\tan(f*x+e)^6-2*I*B*\tan(f*x+e)^5+2*B*\tan(f*x+e)^6+20*I*A*\tan(f*x+e)^4+8*A*\tan(f*x+e)^5-5*I*B*\tan(f*x+e)^3+5*B*\tan(f*x+e)^4+15*I*A*\tan(f*x+e)^2+20*A*\tan(f*x+e)^3-3*I*B*\tan(f*x+e)+3*I*A+12*A*\tan(f*x+e)-3*B)/(I-\tan(f*x+e))^4/(I+\tan(f*x+e))^3$$

3.849.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 181, normalized size of antiderivative = 0.83

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2}} dx = \frac{(5(iA + B)e^{(10ifx+10ie)} + 5(13iA + 7B)e^{(8ifx+8ie)} + 48(iA - B)e^{(7ifx+7ie)} + 30(-iA + B)e^{(6ifx+6ie)})}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2}}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(5/2)/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output
$$-1/240*(5*(I*A + B)*e^{(10*I*f*x + 10*I*e)} + 5*(13*I*A + 7*B)*e^{(8*I*f*x + 8*I*e)} + 48*(I*A - B)*e^{(7*I*f*x + 7*I*e)} + 30*(-I*A + B)*e^{(6*I*f*x + 6*I*e)} + 48*(I*A - B)*e^{(5*I*f*x + 5*I*e)} + 10*(-11*I*A + B)*e^{(4*I*f*x + 4*I*e)} - (23*I*A - 13*B)*e^{(2*I*f*x + 2*I*e)} - 3*I*A + 3*B)*\sqrt{a/(e^{(2*I*f*x + 2*I*e)} + 1)}*\sqrt{c/(e^{(2*I*f*x + 2*I*e)} + 1)}*e^{(-5*I*f*x - 5*I*e)}/(a^3*c^2*f)$$

3.849.
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2}(c-ic \tan(e+fx))^{3/2}} dx$$

3.849.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2}} dx = \int \frac{A + B \tan(e + fx)}{(ia (\tan(e + fx) - i))^{5/2} (-ic (\tan(e + fx) + i))^{3/2}}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**(5/2)/(c-I*c*tan(f*x+e))**(3/2),x)`

output `Integral((A + B*tan(e + f*x))/((I*a*(tan(e + f*x) - I))**(5/2)*(-I*c*(tan(e + f*x) + I))**(3/2)), x)`

3.849.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(5/2)/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.`

3.849.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2}} dx = \int \frac{B \tan(fx + e) + A}{(ia \tan(fx + e) + a)^{5/2} (-ic \tan(fx + e) + c)^{3/2}} dx$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(5/2)/(c-I*c*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)^(5/2)*(-I*c*tan(f*x + e) + c)^(3/2)), x)`

3.849. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2}(c-ic \tan(e+fx))^{3/2}} dx$

3.849.9 Mupad [B] (verification not implemented)

Time = 9.97 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.14

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{3/2}} dx = \sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)i)}{\cos(2e+2fx)+1}} (95 A \sin(2e + 2fx)$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^(5/2)*(c - c*tan(e + f*x)*1i)^(3/2)),x)`

output `((a*cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1)/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*cos(2*e + 2*f*x)*85i - 30*B - A*60i + A*cos(4*e + 4*f*x)*20i + A*cos(6*e + 6*f*x)*3i - 5*B*cos(2*e + 2*f*x) - 10*B*cos(4*e + 4*f*x) - 3*B*cos(6*e + 6*f*x) + 95*A*sin(2*e + 2*f*x) + 20*A*sin(4*e + 4*f*x) + 3*A*sin(6*e + 6*f*x) - B*sin(2*e + 2*f*x)*5i + B*sin(4*e + 4*f*x)*10i + B*sin(6*e + 6*f*x)*3i)/(240*a^3*c*f*(c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)`

3.850
$$\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2}(c-ictan(e+fx))^{5/2}} dx$$

3.850.1 Optimal result 7614
 3.850.2 Mathematica [A] (verified) 7615
 3.850.3 Rubi [A] (verified) 7615
 3.850.4 Maple [A] (verified) 7618
 3.850.5 Fricas [A] (verification not implemented) 7618
 3.850.6 Sympy [F] 7619
 3.850.7 Maxima [B] (verification not implemented) 7619
 3.850.8 Giac [F] 7620
 3.850.9 Mupad [B] (verification not implemented) 7620

3.850.1 Optimal result

Integrand size = 45, antiderivative size = 206

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2}(c - ictan(e + fx))^{5/2}} dx = \frac{-iA - B}{5f(a + ia \tan(e + fx))^{5/2}(c - ictan(e + fx))^{5/2}}$$

$$+ \frac{iA}{5cf(a + ia \tan(e + fx))^{5/2}(c - ictan(e + fx))^{3/2}}$$

$$+ \frac{4A \tan(e + fx)}{15acf(a + ia \tan(e + fx))^{3/2}(c - ictan(e + fx))^{3/2}}$$

$$+ \frac{8A \tan(e + fx)}{15a^2c^2f \sqrt{a + ia \tan(e + fx)} \sqrt{c - ictan(e + fx)}}$$

output `8/15*A*tan(f*x+e)/a^2/c^2/f/(a+I*a*tan(f*x+e))^(1/2)/(c-I*c*tan(f*x+e))^(1/2)+1/5*(-I*A-B)/f/(a+I*a*tan(f*x+e))^(5/2)/(c-I*c*tan(f*x+e))^(5/2)+1/5*I*A/c/f/(a+I*a*tan(f*x+e))^(5/2)/(c-I*c*tan(f*x+e))^(3/2)+4/15*A*tan(f*x+e)/a/c/f/(a+I*a*tan(f*x+e))^(3/2)/(c-I*c*tan(f*x+e))^(3/2)`

3.850.2 Mathematica [A] (verified)

Time = 7.04 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.44

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{5/2}} dx = \frac{\cos^4(e + fx) (-3B + 15A \tan(e + fx) + 20A \tan^3(e + fx))}{15a^2 c^2 f \sqrt{a + ia \tan(e + fx)} \sqrt{c - ic \tan(e + fx)}}$$

input `Integrate[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e + f*x])^(5/2)),x]`

output `(Cos[e + f*x]^4*(-3*B + 15*A*Tan[e + f*x] + 20*A*Tan[e + f*x]^3 + 8*A*Tan[e + f*x]^5))/(15*a^2*c^2*f*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])`

3.850.3 Rubi [A] (verified)Time = 0.41 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.06, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {3042, 4071, 87, 55, 42, 41}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{5/2}} dx \\ & \quad \downarrow \text{3042} \\ & \int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{5/2}} dx \\ & \quad \downarrow \text{4071} \\ & ac \int \frac{A + B \tan(e + fx)}{(i \tan(e + fx) a + a)^{7/2} (c - ic \tan(e + fx))^{7/2}} d \tan(e + fx) \\ & \quad \downarrow \text{87} \\ & ac \left(\frac{A \int \frac{1}{(i \tan(e + fx) a + a)^{7/2} (c - ic \tan(e + fx))^{5/2}} d \tan(e + fx)}{c} - \frac{B + ia}{5ac(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{5/2}} \right) \\ & \quad \downarrow \text{55} \end{aligned}$$

3.850. $\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{5/2}} dx$

$$ac \left(\frac{A \left(\frac{4 \int \frac{1}{(i \tan(e+fx)a+a)^{5/2}(c-ic \tan(e+fx))^{5/2}} dx \tan(e+fx)}{5a} + \frac{i}{5ac(a+ia \tan(e+fx))^{5/2}(c-ic \tan(e+fx))^{3/2}} \right)}{c} - \frac{B+iA}{5ac(a+ia \tan(e+fx))^{5/2}(c-ic \tan(e+fx))^{3/2}} \right) f$$

42

$$ac \left(\frac{A \left(\frac{4 \left(\frac{2 \int \frac{1}{(i \tan(e+fx)a+a)^{3/2}(c-ic \tan(e+fx))^{3/2}} dx \tan(e+fx)}{3ac} + \frac{\tan(e+fx)}{3ac(a+ia \tan(e+fx))^{3/2}(c-ic \tan(e+fx))^{3/2}} \right)}{5a} + \frac{i}{5ac(a+ia \tan(e+fx))^{5/2}(c-ic \tan(e+fx))^{3/2}} \right)}{c} \right) f$$

41

$$ac \left(\frac{A \left(\frac{4 \left(\frac{2 \tan(e+fx)}{3a^2c^2 \sqrt{a+ia \tan(e+fx)} \sqrt{c-ic \tan(e+fx)}} + \frac{\tan(e+fx)}{3ac(a+ia \tan(e+fx))^{3/2}(c-ic \tan(e+fx))^{3/2}} \right)}{5a} + \frac{i}{5ac(a+ia \tan(e+fx))^{5/2}(c-ic \tan(e+fx))^{3/2}} \right)}{c} \right) f$$

input `Int[(A + B*Tan[e + f*x])/((a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e + f*x])^(5/2)), x]`

output `(a*c*(-1/5*(I*A + B)/(a*c*(a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e + f*x])^(5/2)) + (A*((I/5)/(a*c*(a + I*a*Tan[e + f*x])^(5/2)*(c - I*c*Tan[e + f*x])^(3/2)) + (4*(Tan[e + f*x]/(3*a*c*(a + I*a*Tan[e + f*x])^(3/2)*(c - I*c*Tan[e + f*x])^(3/2)) + (2*Tan[e + f*x]/(3*a^2*c^2*Sqrt[a + I*a*Tan[e + f*x]]*Sqrt[c - I*c*Tan[e + f*x]])))/(5*a)))/c)/f`

3.850. $\int \frac{A+B \tan(e+fx)}{(a+ia \tan(e+fx))^{5/2}(c-ic \tan(e+fx))^{5/2}} dx$

3.850.3.1 Defintions of rubi rules used

- rule 41 `Int[1/(((a_) + (b_)*(x_))^(3/2)*((c_) + (d_)*(x_))^(3/2)), x_Symbol] := Simp[x/(a*c*Sqrt[a + b*x]*Sqrt[c + d*x]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0]`
- rule 42 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[(-x)*(a + b*x)^(m + 1)*((c + d*x)^(m + 1)/(2*a*c*(m + 1))), x] + Simp[(2*m + 3)/(2*a*c*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(m + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && ILtQ[m + 3/2, 0]`
- rule 55 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*(Simplify[m + n + 2]/((b*c - a*d)*(m + 1)) Int[(a + b*x)^Simplify[m + 1]*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && ILtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])`
- rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.850.4 Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.60

method	result
derivativedivides	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)}\left(8 A \tan (f x+e)^7+28 A \tan (f x+e)^5+35 A \tan (f x+e)^3-3 B \tan (f x+e)^2+15 B\right)}{15 f a^3 c^3(i-\tan (f x+e))^4(i+\tan (f x+e))^4}$
default	$\frac{\sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)}\left(8 A \tan (f x+e)^7+28 A \tan (f x+e)^5+35 A \tan (f x+e)^3-3 B \tan (f x+e)^2+15 B\right)}{15 f a^3 c^3(i-\tan (f x+e))^4(i+\tan (f x+e))^4}$
parts	$\frac{A \sqrt{a(1+i \tan (f x+e))} \sqrt{-c(i \tan (f x+e)-1)}\left(1+\tan (f x+e)^2\right) \tan (f x+e)\left(8 \tan (f x+e)^4+20 \tan (f x+e)^2+15\right)}{15 f a^3 c^3(i+\tan (f x+e))^4(i-\tan (f x+e))^4}-\frac{B}{f}$

input `int((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(5/2)/(c-I*c*tan(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

output `1/15/f*(a*(1+I*tan(f*x+e)))^(1/2)*(-c*(I*tan(f*x+e)-1))^(1/2)/a^3/c^3*(8*A*tan(f*x+e)^7+28*A*tan(f*x+e)^5+35*A*tan(f*x+e)^3-3*B*tan(f*x+e)^2+15*A*tan(f*x+e)-3*B)/(I-tan(f*x+e))^4/(I+tan(f*x+e))^4`

3.850.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.90

$$\int \frac{A+B \tan (e+f x)}{(a+i a \tan (e+f x))^{5 / 2}(c-i c \tan (e+f x))^{5 / 2}} d x = \frac{(3(i A+B) e^{(12 i f x+12 i e)}+2(14 i A+9 B) e^{(10 i f x+10 i e)}+5(35 i A+9 B) e^{(8 i f x+8 i e)}-96 B e^{(7 i f x+7 i e)}+60 B e^{(6 i f x+6 i e)}-96 B e^{(5 i f x+5 i e)}+5(-35 i A+9 B) e^{(4 i f x+4 i e)}+2(-14 i A+9 B) e^{(2 i f x+2 i e)}-3 i A+3 B) \sqrt{a}\left(e^{(2 i f x+2 i e)}+1\right) \sqrt{c}\left(e^{(2 i f x+2 i e)}+1\right) e^{(-5 i f x-5 i e)}}{\left(a^3 c^3 f\right)}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(5/2)/(c-I*c*tan(f*x+e))^(5/2),x,algorithm="fracas")`

output `-1/480*(3*(I*A + B)*e^(12*I*f*x + 12*I*e) + 2*(14*I*A + 9*B)*e^(10*I*f*x + 10*I*e) + 5*(35*I*A + 9*B)*e^(8*I*f*x + 8*I*e) - 96*B*e^(7*I*f*x + 7*I*e) + 60*B*e^(6*I*f*x + 6*I*e) - 96*B*e^(5*I*f*x + 5*I*e) + 5*(-35*I*A + 9*B)*e^(4*I*f*x + 4*I*e) + 2*(-14*I*A + 9*B)*e^(2*I*f*x + 2*I*e) - 3*I*A + 3*B)*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(c/(e^(2*I*f*x + 2*I*e) + 1))*e^(-5*I*f*x - 5*I*e)/(a^3*c^3*f)`

3.850.
$$\int \frac{A+B \tan (e+f x)}{(a+i a \tan (e+f x))^{5 / 2}(c-i c \tan (e+f x))^{5 / 2}} d x$$

3.850.6 Sympy [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{5/2}} dx = \int \frac{A + B \tan(e + fx)}{(ia (\tan(e + fx) - i))^{5/2} (-ic (\tan(e + fx) + i))^{5/2}}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))**(5/2)/(c-I*c*tan(f*x+e))**(5/2),x)`

output `Integral((A + B*tan(e + f*x))/((I*a*(tan(e + f*x) - I))**(5/2)*(-I*c*(tan(e + f*x) + I))**(5/2)), x)`

3.850.7 Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 334 vs. 2(160) = 320.

Time = 0.59 (sec) , antiderivative size = 334, normalized size of antiderivative = 1.62

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{5/2}} dx = \frac{(30(5iA - B) \cos(4fx + 4e) + 5(5iA - 3B) \cos(2fx + 2e) - 30(5A + I*B) \sin(4fx + 4e) - 5(5A + 3I*B) \sin(2fx + 2e) - 6*B) \cos(5/2 \arctan2(\sin(2fx + 2e), \cos(2fx + 2e))) + 5*(-5I*A - 3*B) \cos(3/2 \arctan2(\sin(2fx + 2e), \cos(2fx + 2e))) + 30*(-5I*A - B) \cos(1/2 \arctan2(\sin(2fx + 2e), \cos(2fx + 2e))) + (30(5A + I*B) \cos(4fx + 4e) + 5(5A + 3I*B) \cos(2fx + 2e) + 30(5I*A - B) \sin(4fx + 4e) + 5(5I*A - 3*B) \sin(2fx + 2e) + 6*A) \sin(5/2 \arctan2(\sin(2fx + 2e), \cos(2fx + 2e))) + 5(5A - 3I*B) \sin(3/2 \arctan2(\sin(2fx + 2e), \cos(2fx + 2e))) + 30(5A - I*B) \sin(1/2 \arctan2(\sin(2fx + 2e), \cos(2fx + 2e)))}{(a^{5/2} * c^{5/2} * f)}$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(5/2)/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="maxima")`

output `1/480*((30*(5*I*A - B)*cos(4*f*x + 4*e) + 5*(5*I*A - 3*B)*cos(2*f*x + 2*e) - 30*(5*A + I*B)*sin(4*f*x + 4*e) - 5*(5*A + 3*I*B)*sin(2*f*x + 2*e) - 6*B)*cos(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 5*(-5*I*A - 3*B)*cos(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 30*(-5*I*A - B)*cos(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + (30*(5*A + I*B)*cos(4*f*x + 4*e) + 5*(5*A + 3*I*B)*cos(2*f*x + 2*e) + 30*(5*I*A - B)*sin(4*f*x + 4*e) + 5*(5*I*A - 3*B)*sin(2*f*x + 2*e) + 6*A)*sin(5/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 5*(5*A - 3*I*B)*sin(3/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))) + 30*(5*A - I*B)*sin(1/2*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e))))/(a^(5/2)*c^(5/2)*f)`

3.850.8 Giac [F]

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{5/2}} dx = \int \frac{B \tan(fx + e) + A}{(ia \tan(fx + e) + a)^{5/2} (-ic \tan(fx + e) + c)^{5/2}} dx$$

input `integrate((A+B*tan(f*x+e))/(a+I*a*tan(f*x+e))^(5/2)/(c-I*c*tan(f*x+e))^(5/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)/((I*a*tan(f*x + e) + a)^(5/2)*(-I*c*tan(f*x + e) + c)^(5/2)), x)`

3.850.9 Mupad [B] (verification not implemented)

Time = 10.48 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.21

$$\int \frac{A + B \tan(e + fx)}{(a + ia \tan(e + fx))^{5/2} (c - ic \tan(e + fx))^{5/2}} dx = \sqrt{\frac{a(\cos(2e+2fx)+1+\sin(2e+2fx)i)}{\cos(2e+2fx)+1}} (175 A \sin(2e + 2fx) + \dots)$$

input `int((A + B*tan(e + f*x))/((a + a*tan(e + f*x)*1i)^(5/2)*(c - c*tan(e + f*x)*1i)^(5/2)),x)`

output `((a*(cos(2*e + 2*f*x) + sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2)*(A*cos(2*e + 2*f*x)*125i - 30*B - A*150i + A*cos(4*e + 4*f*x)*22i + A*cos(6*e + 6*f*x)*3i - 45*B*cos(2*e + 2*f*x) - 18*B*cos(4*e + 4*f*x) - 3*B*cos(6*e + 6*f*x) + 175*A*sin(2*e + 2*f*x) + 28*A*sin(4*e + 4*f*x) + 3*A*sin(6*e + 6*f*x) + B*sin(2*e + 2*f*x)*15i + B*sin(4*e + 4*f*x)*12i + B*sin(6*e + 6*f*x)*3i)/(480*a^3*c^2*f*((c*(cos(2*e + 2*f*x) - sin(2*e + 2*f*x)*1i + 1))/(cos(2*e + 2*f*x) + 1))^(1/2))`

3.851 $\int (a+ia \tan(e+fx))^m (A+B \tan(e+fx))(c-ictan(e+fx))^n dx$

3.851.1 Optimal result	7621
3.851.2 Mathematica [A] (verified)	7621
3.851.3 Rubi [A] (verified)	7622
3.851.4 Maple [F]	7624
3.851.5 Fricas [F]	7624
3.851.6 Sympy [F]	7625
3.851.7 Maxima [F]	7625
3.851.8 Giac [F]	7625
3.851.9 Mupad [F(-1)]	7626

3.851.1 Optimal result

Integrand size = 41, antiderivative size = 150

$$\int (a + ia \tan(e + fx))^m (A + B \tan(e + fx))(c - ictan(e + fx))^n dx$$

$$= \frac{(iA + B)(a + ia \tan(e + fx))^m (c - ictan(e + fx))^n}{2fn}$$

$$- \frac{2^{-1+n}(B(m - n) + iA(m + n)) \text{Hypergeometric2F1}(m, -n, 1 + m, \frac{1}{2}(1 + i \tan(e + fx))) (1 - i \tan(e + fx))}{fmn}$$

```
output 1/2*(I*A+B)*(a+I*a*tan(f*x+e))^m*(c-I*c*tan(f*x+e))^n/f/n-2^(-1+n)*(B*(-n+m)+I*A*(n+m))*hypergeom([m, -n], [1+m], 1/2+1/2*I*tan(f*x+e))*(a+I*a*tan(f*x+e))^m*(c-I*c*tan(f*x+e))^n/f/m/n/((1-I*tan(f*x+e))^n)
```

3.851.2 Mathematica [A] (verified)

Time = 1.13 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.81

$$\int (a + ia \tan(e + fx))^m (A + B \tan(e + fx))(c - ictan(e + fx))^n dx$$

$$= \frac{2^{-1+n}((-iA - B) \text{Hypergeometric2F1}(m, 1 - n, 1 + m, \frac{1}{2}(1 + i \tan(e + fx))) + 2B \text{Hypergeometric2F1}(m, 1 - n, 1 + m, \frac{1}{2}(1 + i \tan(e + fx))))}{fn}$$

input `Integrate[(a + I*a*Tan[e + f*x])^m*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^n,x]`

output `(2^(-1 + n)*((-I)*A - B)*Hypergeometric2F1[m, 1 - n, 1 + m, (1 + I*Tan[e + f*x])/2] + 2*B*Hypergeometric2F1[m, -n, 1 + m, (1 + I*Tan[e + f*x])/2])* (a + I*a*Tan[e + f*x])^m*(c - I*c*Tan[e + f*x])^n/(f*m*(1 - I*Tan[e + f*x]))^n`

3.851.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.10, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.122$, Rules used = {3042, 4071, 88, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(e + fx))^m (A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(e + fx))^m (A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int (i \tan(e + fx)a + a)^{m-1} (A + B \tan(e + fx))(c - ic \tan(e + fx))^{n-1} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{88} \\
 & \frac{ac \left(\frac{(B+ia)(a+ia \tan(e+fx))^m (c-ic \tan(e+fx))^n}{2acn} - \frac{(-A(m+n)+iB(m-n)) \int (i \tan(e+fx)a+a)^{m-1} (c-ic \tan(e+fx))^{n-1} d \tan(e+fx)}{2cn} \right)}{f} \\
 & \quad \downarrow \text{80} \\
 & \frac{ac \left(\frac{(B+ia)(a+ia \tan(e+fx))^m (c-ic \tan(e+fx))^n}{2acn} - \frac{2^{n-1}(-A(m+n)+iB(m-n))(1-i \tan(e+fx))^{-n} (c-ic \tan(e+fx))^n \int (\frac{1}{2} - \frac{1}{2}i \tan(e+fx))}{cn} \right)}{f} \\
 & \quad \downarrow \text{79}
 \end{aligned}$$

3.851. $\int (a + ia \tan(e + fx))^m (A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx$

$$ac \left(\frac{i2^{n-1}(-A(m+n)+iB(m-n))(1-i \tan(e+fx))^{-n}(a+ia \tan(e+fx))^m(c-ictan(e+fx))^n \text{Hypergeometric2F1}(m,-n,m+1,\frac{1}{2}(i \tan(e+fx)+1),x)}{acmn} \right) f$$

input `Int[(a + I*a*Tan[e + f*x])^m*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^n, x]`

output `(a*c*(((I*A + B)*(a + I*a*Tan[e + f*x])^m*(c - I*c*Tan[e + f*x])^n)/(2*a*c*n) + (I*2^(-1 + n)*(I*B*(m - n) - A*(m + n))*Hypergeometric2F1[m, -n, 1 + m, (1 + I*Tan[e + f*x])/2]*(a + I*a*Tan[e + f*x])^m*(c - I*c*Tan[e + f*x])^n)/(a*c*m*n*(1 - I*Tan[e + f*x])^n))/f`

3.851.3.1 Defintions of rubi rules used

rule 79 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))`

rule 80 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d))], x]^n, x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])`

rule 88 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^Simplify[p + 1], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && !RationalQ[p] && SumSimplerQ[p, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.851. $\int (a + ia \tan(e + fx))^m (A + B \tan(e + fx))(c - ictan(e + fx))^n dx$

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.851.4 Maple [F]

$$\int (a + ia \tan(fx + e))^m (A + B \tan(fx + e)) (c - ic \tan(fx + e))^n dx$$

input `int((a+I*a*tan(f*x+e))^m*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x)`

output `int((a+I*a*tan(f*x+e))^m*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x)`

3.851.5 Fracas [F]

$$\begin{aligned} & \int (a + ia \tan(e + fx))^m (A + B \tan(e + fx)) (c - ic \tan(e + fx))^n dx \\ &= \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^m (-ic \tan(fx + e) + c)^n dx \end{aligned}$$

input `integrate((a+I*a*tan(f*x+e))^m*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x, algorithm="fricas")`

output `integral(((A - I*B)*e^(2*I*f*x + 2*I*e) + A + I*B)*(2*c/(e^(2*I*f*x + 2*I*e) + 1))^n*e^(2*I*f*m*x + 2*I*e*m + m*log(a/c) + m*log(2*c/(e^(2*I*f*x + 2*I*e) + 1)))/(e^(2*I*f*x + 2*I*e) + 1), x)`

3.851.6 Sympy [F]

$$\int (a + ia \tan(e + fx))^m (A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx$$

$$= \int (ia(\tan(e + fx) - i))^m (-ic(\tan(e + fx) + i))^n (A + B \tan(e + fx)) dx$$

input `integrate((a+I*a*tan(f*x+e))**m*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**n,x)`

output `Integral((I*a*(tan(e + f*x) - I))**m*(-I*c*(tan(e + f*x) + I))**n*(A + B*tan(e + f*x)), x)`

3.851.7 Maxima [F]

$$\int (a + ia \tan(e + fx))^m (A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx$$

$$= \int (B \tan(fx + e) + A)(ia \tan(fx + e) + a)^m (-ic \tan(fx + e) + c)^n dx$$

input `integrate((a+I*a*tan(f*x+e))^m*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x, algorithm="maxima")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^m*(-I*c*tan(f*x + e) + c)^n, x)`

3.851.8 Giac [F]

$$\int (a + ia \tan(e + fx))^m (A + B \tan(e + fx))(c - ic \tan(e + fx))^n dx$$

$$= \int (B \tan(fx + e) + A)(ia \tan(fx + e) + a)^m (-ic \tan(fx + e) + c)^n dx$$

input `integrate((a+I*a*tan(f*x+e))^m*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^n,x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^m*(-I*c*tan(f*x + e) + c)^n, x)`

3.851.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^m (A + B \tan(e + fx))(c - ictan(e + fx))^n dx$$

$$= \int (A + B \tan(e + fx)) (a + a \tan(e + fx) 1i)^m (c - c \tan(e + fx) 1i)^n dx$$

input `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^m*(c - c*tan(e + f*x)*1i)
^n,x)`

output `int((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^m*(c - c*tan(e + f*x)*1i)
^n, x)`

3.852 $\int (a+ia \tan(e+fx))^{1+m}(A+B \tan(e+fx))(c-ictan(e+fx))^{-1-m} dx$

3.852.1 Optimal result	7627
3.852.2 Mathematica [A] (verified)	7627
3.852.3 Rubi [A] (verified)	7628
3.852.4 Maple [F]	7630
3.852.5 Fricas [F]	7630
3.852.6 Sympy [F]	7631
3.852.7 Maxima [F]	7631
3.852.8 Giac [F]	7632
3.852.9 Mupad [F(-1)]	7633

3.852.1 Optimal result

Integrand size = 47, antiderivative size = 147

$$\int (a + ia \tan(e + fx))^{1+m}(A + B \tan(e + fx))(c - ictan(e + fx))^{-1-m} dx$$

$$= -\frac{(iA + B)(a + ia \tan(e + fx))^{1+m}(c - ictan(e + fx))^{-1-m}}{2f(1 + m)}$$

$$+ \frac{2^m a B \text{Hypergeometric2F1}(-m, -m, 1 - m, \frac{1}{2}(1 - i \tan(e + fx))) (1 + i \tan(e + fx))^{-m}(a + ia \tan(e + fx))}{cfm}$$

```
output -1/2*(I*A+B)*(a+I*a*tan(f*x+e))^(1+m)*(c-I*c*tan(f*x+e))^(-1-m)/f/(1+m)+2^m*a*B*hypergeom([-m, -m], [1-m], 1/2-1/2*I*tan(f*x+e))*(a+I*a*tan(f*x+e))^m/c/f/m/((1+I*tan(f*x+e))^m)/((c-I*c*tan(f*x+e))^m)
```

3.852.2 Mathematica [A] (verified)

Time = 5.84 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.90

$$\int (a + ia \tan(e + fx))^{1+m}(A + B \tan(e + fx))(c - ictan(e + fx))^{-1-m} dx$$

$$= \frac{a(a + ia \tan(e + fx))^m(c - ictan(e + fx))^{-m} \left(\frac{2^{1+m} B \text{Hypergeometric2F1}(-m, -m, 1 - m, -\frac{1}{2}i(i + \tan(e + fx))) (1 + i \tan(e + fx))}{m} \right)}{2cf}$$

input `Integrate[(a + I*a*Tan[e + f*x])^(1 + m)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(-1 - m),x]`

output `(a*(a + I*a*Tan[e + f*x])^m*((2^(1 + m)*B*Hypergeometric2F1[-m, -m, 1 - m, (-1/2*I)*(I + Tan[e + f*x])])/(m*(1 + I*Tan[e + f*x])^m) + ((I*A + B)*(-I + Tan[e + f*x]))/((1 + m)*(I + Tan[e + f*x]))) / (2*c*f*(c - I*c*Tan[e + f*x])^m)`

3.852.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.106$, Rules used = {3042, 4071, 88, 80, 79}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (a + ia \tan(e + fx))^{m+1} (A + B \tan(e + fx)) (c - ictan(e + fx))^{-m-1} dx \\
 & \quad \downarrow \text{3042} \\
 & \int (a + ia \tan(e + fx))^{m+1} (A + B \tan(e + fx)) (c - ictan(e + fx))^{-m-1} dx \\
 & \quad \downarrow \text{4071} \\
 & \frac{ac \int (i \tan(e + fx)a + a)^m (A + B \tan(e + fx)) (c - ictan(e + fx))^{-m-2} d \tan(e + fx)}{f} \\
 & \quad \downarrow \text{88} \\
 & \frac{ac \left(\frac{iB \int (i \tan(e + fx)a + a)^m (c - ictan(e + fx))^{-m-1} d \tan(e + fx)}{c} - \frac{(B + iA)(a + ia \tan(e + fx))^{m+1} (c - ictan(e + fx))^{-m-1}}{2ac(m+1)} \right)}{f} \\
 & \quad \downarrow \text{80} \\
 & \frac{ac \left(\frac{iB 2^m (1 + i \tan(e + fx))^{-m} (a + ia \tan(e + fx))^m \int (\frac{1}{2} i \tan(e + fx) + \frac{1}{2})^m (c - ictan(e + fx))^{-m-1} d \tan(e + fx)}{c} - \frac{(B + iA)(a + ia \tan(e + fx))^{m+1}}{2ac(m+1)} \right)}{f} \\
 & \quad \downarrow \text{79}
 \end{aligned}$$

3.852. $\int (a + ia \tan(e + fx))^{1+m} (A + B \tan(e + fx)) (c - ictan(e + fx))^{-1-m} dx$

$$ac \frac{B2^m(1+i \tan(e+fx))^{-m}(a+ia \tan(e+fx))^m(c-ictan(e+fx))^{-m} \text{Hypergeometric2F1}(-m, -m, 1-m, \frac{1}{2}(1-i \tan(e+fx)))}{c^{2m}} - \frac{(B+iA)(a+ia \tan(e+fx))}{f}$$

input `Int[(a + I*a*Tan[e + f*x])^(1 + m)*(A + B*Tan[e + f*x])*(c - I*c*Tan[e + f*x])^(-1 - m), x]`

output `(a*c*(-1/2*((I*A + B)*(a + I*a*Tan[e + f*x])^(1 + m)*(c - I*c*Tan[e + f*x])^(-1 - m)))/(a*c*(1 + m)) + (2^m*B*Hypergeometric2F1[-m, -m, 1 - m, (1 - I*Tan[e + f*x])/2]*(a + I*a*Tan[e + f*x])^m)/(c^2*m*(1 + I*Tan[e + f*x])^m*(c - I*c*Tan[e + f*x])^m))/f`

3.852.3.1 Defintions of rubi rules used

rule 79 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))`

rule 80 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(c + d*x)^FracPart[n]/((b/(b*c - a*d))^IntPart[n]*(b*((c + d*x)/(b*c - a*d)))^FracPart[n]) Int[(a + b*x)^m*Simp[b*(c/(b*c - a*d)) + b*d*(x/(b*c - a*d))], x]^n, x] /; FreeQ[{a, b, c, d, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && (RationalQ[m] || !SimplerQ[n + 1, m + 1])`

rule 88 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^Simplify[p + 1], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && !RationalQ[p] && SumSimplerQ[p, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4071 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c + a*d, 0] && EqQ[a^2 + b^2, 0]`

3.852.4 Maple [F]

$$\int (a + ia \tan(fx + e))^{1+m} (A + B \tan(fx + e)) (c - ictan(fx + e))^{-1-m} dx$$

input `int((a+I*a*tan(f*x+e))^(1+m)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(1-m),x)`

output `int((a+I*a*tan(f*x+e))^(1+m)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(1-m),x)`

3.852.5 Fracas [F]

$$\begin{aligned} & \int (a + ia \tan(e + fx))^{1+m} (A + B \tan(e + fx)) (c - ictan(e + fx))^{-1-m} dx \\ &= \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^{m+1} (-ictan(fx + e) + c)^{-m-1} dx \end{aligned}$$

input `integrate((a+I*a*tan(f*x+e))^(1+m)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(1-m),x, algorithm="fricas")`

output `integral(((A - I*B)*e^(2*I*f*x + 2*I*e) + A + I*B)*(2*c/(e^(2*I*f*x + 2*I*e) + 1))^(1-m) * e^(2*I*e*m - 2*(-I*f*m - I*f)*x + (m + 1)*log(a/c) + (m + 1)*log(2*c/(e^(2*I*f*x + 2*I*e) + 1)) + 2*I*e)/(e^(2*I*f*x + 2*I*e) + 1), x)`

3.852.6 Sympy [F]

$$\int (a + ia \tan(e + fx))^{1+m} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{-1-m} dx$$

$$= \int (ia(\tan(e + fx) - i))^{m+1} (-ic(\tan(e + fx) + i))^{-m-1} (A + B \tan(e + fx)) dx$$

input `integrate((a+I*a*tan(f*x+e))**(1+m)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))**(-1-m),x)`

output `Integral((I*a*(tan(e + f*x) - I))**(m + 1)*(-I*c*(tan(e + f*x) + I))**(-m - 1)*(A + B*tan(e + f*x)), x)`

3.852.7 Maxima [F]

$$\int (a + ia \tan(e + fx))^{1+m} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{-1-m} dx$$

$$= \int (B \tan(fx + e) + A)(ia \tan(fx + e) + a)^{m+1} (-ic \tan(fx + e) + c)^{-m-1} dx$$

input `integrate((a+I*a*tan(f*x+e))^(1+m)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(1-m),x, algorithm="maxima")`

output

```

-(2*(-I*B*a^(m + 1)*m - I*B*a^(m + 1))*cos(2*f*m*x + 2*e*m) + ((A - I*B)*a
^(m + 1)*m^2 - (A - I*B)*a^(m + 1)*m)*cos(2*e*m + 2*(f*m + 2*f)*x + 4*e) +
((A + I*B)*a^(m + 1)*m^2 - (A - I*B)*a^(m + 1)*m - 2*I*B*a^(m + 1))*cos(2
*e*m + 2*(f*m + f)*x + 2*e) - 4*(B*a^(m + 1)*c^(m + 1)*f*m^3 - B*a^(m + 1)
*c^(m + 1)*f*m + (B*a^(m + 1)*c^(m + 1)*f*m^3 - B*a^(m + 1)*c^(m + 1)*f*m)
*cos(2*f*x + 2*e) - (-I*B*a^(m + 1)*c^(m + 1)*f*m^3 + I*B*a^(m + 1)*c^(m +
1)*f*m)*sin(2*f*x + 2*e))*integrate(((cos(4*f*x + 4*e) + 2*cos(2*f*x + 2*
e) + 1)*cos(2*f*m*x + 2*e*m) + (sin(4*f*x + 4*e) + 2*sin(2*f*x + 2*e))*sin
(2*f*m*x + 2*e*m))/((c^(m + 1)*m - c^(m + 1))*cos(4*f*x + 4*e)^2 + 4*(c^(m
+ 1)*m - c^(m + 1))*cos(2*f*x + 2*e)^2 + (c^(m + 1)*m - c^(m + 1))*sin(4*
f*x + 4*e)^2 + 4*(c^(m + 1)*m - c^(m + 1))*sin(4*f*x + 4*e)*sin(2*f*x + 2*
e) + 4*(c^(m + 1)*m - c^(m + 1))*sin(2*f*x + 2*e)^2 + c^(m + 1)*m + 2*(c^(
m + 1)*m + 2*(c^(m + 1)*m - c^(m + 1))*cos(2*f*x + 2*e) - c^(m + 1))*cos(4
*f*x + 4*e) + 4*(c^(m + 1)*m - c^(m + 1))*cos(2*f*x + 2*e) - c^(m + 1)), x
) + 4*(-I*B*a^(m + 1)*c^(m + 1)*f*m^3 + I*B*a^(m + 1)*c^(m + 1)*f*m + (-I*
B*a^(m + 1)*c^(m + 1)*f*m^3 + I*B*a^(m + 1)*c^(m + 1)*f*m)*cos(2*f*x + 2*e
) + (B*a^(m + 1)*c^(m + 1)*f*m^3 - B*a^(m + 1)*c^(m + 1)*f*m)*sin(2*f*x +
2*e))*integrate(-((sin(4*f*x + 4*e) + 2*sin(2*f*x + 2*e))*cos(2*f*m*x + 2*
e*m) - (cos(4*f*x + 4*e) + 2*cos(2*f*x + 2*e) + 1)*sin(2*f*m*x + 2*e*m))/
(c^(m + 1)*m - c^(m + 1))*cos(4*f*x + 4*e)^2 + 4*(c^(m + 1)*m - c^(m + ...

```

3.852.8 Giac [F]

$$\int (a + ia \tan(e + fx))^{1+m} (A + B \tan(e + fx)) (c - ictan(e + fx))^{-1-m} dx$$

$$= \int (B \tan(fx + e) + A) (ia \tan(fx + e) + a)^{m+1} (-ictan(fx + e) + c)^{-m-1} dx$$

input

```

integrate((a+I*a*tan(f*x+e))^(1+m)*(A+B*tan(f*x+e))*(c-I*c*tan(f*x+e))^(1-
-m),x, algorithm="giac")

```

output

```

integrate((B*tan(f*x + e) + A)*(I*a*tan(f*x + e) + a)^(m + 1)*(-I*c*tan(f*
x + e) + c)^(-m - 1), x)

```

3.852.9 Mupad [F(-1)]

Timed out.

$$\int (a + ia \tan(e + fx))^{1+m} (A + B \tan(e + fx)) (c - ic \tan(e + fx))^{-1-m} dx$$

$$= \int \frac{(A + B \tan(e + fx)) (a + a \tan(e + fx) \text{li})^{m+1}}{(c - c \tan(e + fx) \text{li})^{m+1}} dx$$

input `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(m + 1))/(c - c*tan(e + f*x)*1i)^(m + 1),x)`

output `int(((A + B*tan(e + f*x))*(a + a*tan(e + f*x)*1i)^(m + 1))/(c - c*tan(e + f*x)*1i)^(m + 1), x)`

$$3.853 \quad \int \frac{(c - ic \tan(e + fx))^n (-i(2+n) + (-2+n) \tan(e + fx))}{(-i + \tan(e + fx))^2} dx$$

3.853.1 Optimal result	7634
3.853.2 Mathematica [A] (verified)	7634
3.853.3 Rubi [A] (verified)	7635
3.853.4 Maple [B] (verified)	7636
3.853.5 Fricas [A] (verification not implemented)	7636
3.853.6 Sympy [B] (verification not implemented)	7637
3.853.7 Maxima [F(-2)]	7637
3.853.8 Giac [F]	7638
3.853.9 Mupad [B] (verification not implemented)	7638

3.853.1 Optimal result

Integrand size = 46, antiderivative size = 33

$$\int \frac{(c - ic \tan(e + fx))^n (-i(2+n) + (-2+n) \tan(e + fx))}{(-i + \tan(e + fx))^2} dx = \frac{(c - ic \tan(e + fx))^n}{f(i - \tan(e + fx))^2}$$

output `(c-I*c*tan(f*x+e))^n/f/(I-tan(f*x+e))^2`

3.853.2 Mathematica [A] (verified)

Time = 5.45 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.94

$$\int \frac{(c - ic \tan(e + fx))^n (-i(2+n) + (-2+n) \tan(e + fx))}{(-i + \tan(e + fx))^2} dx = \frac{(c - ic \tan(e + fx))^n}{f(-i + \tan(e + fx))^2}$$

input `Integrate[((c - I*c*Tan[e + f*x])^n*((-I)*(2 + n) + (-2 + n)*Tan[e + f*x]))/(-I + Tan[e + f*x])^2,x]`

output `(c - I*c*Tan[e + f*x])^n/(f*(-I + Tan[e + f*x])^2)`

3.853.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.065$, Rules used = {3042, 4071, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{((n-2)\tan(e+fx) - i(n+2))(c - ictan(e+fx))^n}{(\tan(e+fx) - i)^2} dx$$

↓ 3042

$$\int \frac{((n-2)\tan(e+fx) - i(n+2))(c - ictan(e+fx))^n}{(\tan(e+fx) - i)^2} dx$$

↓ 4071

$$\frac{ic \int \frac{(c-ictan(e+fx))^{n-1}(i(n+2)+(2-n)\tan(e+fx))}{(i-\tan(e+fx))^3} d \tan(e+fx)}{f}$$

↓ 83

$$\frac{(c - ictan(e+fx))^n}{f(-\tan(e+fx) + i)^2}$$

input `Int[((c - I*c*Tan[e + f*x])^n*((-I)*(2 + n) + (-2 + n)*Tan[e + f*x]))/(-I + Tan[e + f*x])^2,x]`

output `(c - I*c*Tan[e + f*x])^n/(f*(I - Tan[e + f*x])^2)`

3.853.3.1 Defintions of rubi rules used

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

3.853. $\int \frac{(c-ictan(e+fx))^n(-i(2+n)+(-2+n)\tan(e+fx))}{(-i+\tan(e+fx))^2} dx$

```
rule 4071 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Si
mp[a*(c/f) Subst[Int[(a + b*x)^(m - 1)*(c + d*x)^(n - 1)*(A + B*x), x], x
, Tan[e + f*x]], x] /; FreeQ[{a, b, c, d, e, f, A, B, m, n}, x] && EqQ[b*c
+ a*d, 0] && EqQ[a^2 + b^2, 0]
```

3.853.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 91 vs. 2(31) = 62.

Time = 0.74 (sec) , antiderivative size = 92, normalized size of antiderivative = 2.79

method	result	size
derivativedivides	$\frac{\frac{\tan(fx+e)^2 e^{n \ln(c-ic \tan(fx+e))}}{f} - \frac{e^{n \ln(c-ic \tan(fx+e))}}{f} + \frac{2i \tan(fx+e) e^{n \ln(c-ic \tan(fx+e))}}{f}}{(1+\tan(fx+e)^2)^2}$	92
default	$\frac{\frac{\tan(fx+e)^2 e^{n \ln(c-ic \tan(fx+e))}}{f} - \frac{e^{n \ln(c-ic \tan(fx+e))}}{f} + \frac{2i \tan(fx+e) e^{n \ln(c-ic \tan(fx+e))}}{f}}{(1+\tan(fx+e)^2)^2}$	92

```
input int((c-I*c*tan(f*x+e))^n*(-I*(2+n)+(-2+n)*tan(f*x+e))/(-I+tan(f*x+e))^2,x,
method=_RETURNVERBOSE)
```

```
output (1/f*tan(f*x+e)^2*exp(n*ln(c-I*c*tan(f*x+e)))-1/f*exp(n*ln(c-I*c*tan(f*x+e
))) + 2*I/f*tan(f*x+e)*exp(n*ln(c-I*c*tan(f*x+e))))/(1+tan(f*x+e)^2)^2
```

3.853.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.64

$$\int \frac{(c - ic \tan(e + fx))^n (-i(2 + n) + (-2 + n) \tan(e + fx))}{(-i + \tan(e + fx))^2} dx$$

$$= - \frac{\left(\frac{2c}{e^{(2i fx + 2i e)} + 1}\right)^n (e^{(4i fx + 4i e)} + 2e^{(2i fx + 2i e)} + 1)e^{(-4i fx - 4i e)}}{4f}$$

```
input integrate((c-I*c*tan(f*x+e))^n*(-I*(2+n)+(-2+n)*tan(f*x+e))/(-I+tan(f*x+e)
)^2,x, algorithm="fracas")
```

3.853. $\int \frac{(c - ic \tan(e + fx))^n (-i(2 + n) + (-2 + n) \tan(e + fx))}{(-i + \tan(e + fx))^2} dx$

output
$$-1/4*(2*c/(e^{(2*I*f*x + 2*I*e)} + 1))^n*(e^{(4*I*f*x + 4*I*e)} + 2*e^{(2*I*f*x + 2*I*e)} + 1)*e^{(-4*I*f*x - 4*I*e)}/f$$

3.853.6 Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 66 vs. $2(24) = 48$.

Time = 0.64 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.00

$$\int \frac{(c - ic \tan(e + fx))^n (-i(2 + n) + (-2 + n) \tan(e + fx))}{(-i + \tan(e + fx))^2} dx$$

$$= \begin{cases} \frac{(-ic \tan(e + fx) + c)^n}{f \tan^2(e + fx) - 2if \tan(e + fx) - f} & \text{for } f \neq 0 \\ \frac{x((n-2) \tan(e) - i(n+2))(-ic \tan(e) + c)^n}{(\tan(e) - i)^2} & \text{otherwise} \end{cases}$$

input `integrate((c-I*c*tan(f*x+e))**n*(-I*(2+n)+(-2+n)*tan(f*x+e))/(-I+tan(f*x+e))**2,x)`

output `Piecewise(((-I*c*tan(e + f*x) + c)**n/(f*tan(e + f*x)**2 - 2*I*f*tan(e + f*x) - f), Ne(f, 0)), (x*((n - 2)*tan(e) - I*(n + 2))*(-I*c*tan(e) + c)**n/(tan(e) - I)**2, True))`

3.853.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(c - ic \tan(e + fx))^n (-i(2 + n) + (-2 + n) \tan(e + fx))}{(-i + \tan(e + fx))^2} dx$$

= Exception raised: RuntimeError

input `integrate((c-I*c*tan(f*x+e))^n*(-I*(2+n)+(-2+n)*tan(f*x+e))/(-I+tan(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is un defined.`

3.853.8 Giac [F]

$$\int \frac{(c - ic \tan(e + fx))^n (-i(2 + n) + (-2 + n) \tan(e + fx))}{(-i + \tan(e + fx))^2} dx$$

$$= \int \frac{((n - 2) \tan(fx + e) - in - 2i)(-ic \tan(fx + e) + c)^n}{(\tan(fx + e) - i)^2} dx$$

input `integrate((c-I*c*tan(f*x+e))^n*(-I*(2+n)+(-2+n)*tan(f*x+e))/(-I+tan(f*x+e))^2,x, algorithm="giac")`

output `integrate(((n - 2)*tan(f*x + e) - I*n - 2*I)*(-I*c*tan(f*x + e) + c)^n/(tan(f*x + e) - I)^2, x)`

3.853.9 Mupad [B] (verification not implemented)

Time = 8.46 (sec) , antiderivative size = 90, normalized size of antiderivative = 2.73

$$\int \frac{(c - ic \tan(e + fx))^n (-i(2 + n) + (-2 + n) \tan(e + fx))}{(-i + \tan(e + fx))^2} dx$$

$$= \frac{\left(-\frac{c(-2 \cos(e+fx)^2 + \sin(2e+2fx)1i)}{2 \cos(e+fx)^2} \right)^n (-4 \cos(e + fx)^2 - 2 \cos(2e + 2fx)^2 + \sin(2e + 2fx) 2i + \sin(4e + 2fx) 2i)}{4f}$$

input `int(-((c - c*tan(e + f*x)*1i)^n*(n*1i - tan(e + f*x)*(n - 2) + 2i))/(tan(e + f*x) - 1i)^2,x)`

output `((-(c*(sin(2*e + 2*f*x)*1i - 2*cos(e + f*x)^2))/(2*cos(e + f*x)^2))^n*(sin(2*e + 2*f*x)*2i + sin(4*e + 4*f*x)*1i - 2*cos(2*e + 2*f*x)^2 - 4*cos(e + f*x)^2 + 2))/(4*f)`

3.854
$$\int \frac{(A+B \tan(e+fx))(c+d \tan(e+fx))}{(a+ia \tan(e+fx))^2} dx$$

3.854.1 Optimal result	7639
3.854.2 Mathematica [A] (verified)	7639
3.854.3 Rubi [A] (verified)	7640
3.854.4 Maple [A] (verified)	7641
3.854.5 Fricas [A] (verification not implemented)	7642
3.854.6 Sympy [A] (verification not implemented)	7642
3.854.7 Maxima [F(-2)]	7643
3.854.8 Giac [B] (verification not implemented)	7643
3.854.9 Mupad [B] (verification not implemented)	7644

3.854.1 Optimal result

Integrand size = 36, antiderivative size = 104

$$\int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{(A - iB)(c - id)x}{4a^2} + \frac{B(c + 3id) + A(ic + d)}{4a^2 f(1 + i \tan(e + fx))} + \frac{(iA - B)(c + id)}{4f(a + ia \tan(e + fx))^2}$$

output `1/4*(A-I*B)*(c-I*d)*x/a^2+1/4*(B*(c+3*I*d)+A*(I*c+d))/a^2/f/(1+I*tan(f*x+e))+1/4*(I*A-B)*(c+I*d)/f/(a+I*a*tan(f*x+e))^2`

3.854.2 Mathematica [A] (verified)

Time = 2.06 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.89

$$\int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{(A - iB)(c - id) \arctan(\tan(e + fx)) + \frac{(A+iB)(-ic+d)}{(-i+\tan(e+fx))^2} + \frac{Ac-iBc-iAd+3Bd}{-i+\tan(e+fx)}}{4a^2 f}$$

input `Integrate[((A + B*Tan[e + f*x])*(c + d*Tan[e + f*x]))/(a + I*a*Tan[e + f*x])^2,x]`

output $((A - I*B)*(c - I*d)*ArcTan[Tan[e + f*x]] + ((A + I*B)*((-I)*c + d))/(-I + Tan[e + f*x])^2 + (A*c - I*B*c - I*A*d + 3*B*d)/(-I + Tan[e + f*x])/(4*a^2*f)$

3.854.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.05, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.139$, Rules used = {3042, 4073, 3042, 4009, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx \\ & \quad \downarrow 3042 \\ & \int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx \\ & \quad \downarrow 4073 \\ & \frac{(-B + iA)(c + id)}{4f(a + ia \tan(e + fx))^2} - \frac{i \int \frac{a(B(c+id)+A(ic+d))+2aBd \tan(e+fx)}{i \tan(e+fx)a+a} dx}{2a^2} \\ & \quad \downarrow 3042 \\ & \frac{(-B + iA)(c + id)}{4f(a + ia \tan(e + fx))^2} - \frac{i \int \frac{a(B(c+id)+A(ic+d))+2aBd \tan(e+fx)}{i \tan(e+fx)a+a} dx}{2a^2} \\ & \quad \downarrow 4009 \\ & \frac{(-B + iA)(c + id)}{4f(a + ia \tan(e + fx))^2} - \frac{i \left(\frac{1}{2} (B + iA)(c - id) \int 1 dx - \frac{Ac - iAd - iBc + 3Bd}{2f(1 + i \tan(e + fx))} \right)}{2a^2} \\ & \quad \downarrow 24 \\ & \frac{(-B + iA)(c + id)}{4f(a + ia \tan(e + fx))^2} - \frac{i \left(\frac{1}{2} x (B + iA)(c - id) - \frac{Ac - iAd - iBc + 3Bd}{2f(1 + i \tan(e + fx))} \right)}{2a^2} \end{aligned}$$

input $Int[((A + B*Tan[e + f*x])*(c + d*Tan[e + f*x]))/(a + I*a*Tan[e + f*x])^2,x]$

output $((-1/2*I)*(((I*A + B)*(c - I*d)*x)/2 - (A*c - I*B*c - I*A*d + 3*B*d)/(2*f*(1 + I*Tan[e + f*x]))))/a^2 + ((I*A - B)*(c + I*d))/(4*f*(a + I*a*Tan[e + f*x])^2)$

3.854.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4009 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2, 0] && LtQ[m, 0]`

rule 4073 `Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) + (f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-(A*b - a*B))*(a*c + b*d)*((a + b*Tan[e + f*x])^m/(2*a^2*f*m)), x] + Simp[1/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[A*b*c + a*B*c + a*A*d + b*B*d + 2*a*B*d*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && EqQ[a^2 + b^2, 0]`

3.854.4 Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.48

method	result
risch	$-\frac{ixAd}{4a^2} - \frac{ixBc}{4a^2} + \frac{xcA}{4a^2} - \frac{x Bd}{4a^2} + \frac{ie^{-2i(fx+e)}cA}{4fa^2} + \frac{ie^{-2i(fx+e)}Bd}{4fa^2} - \frac{e^{-4i(fx+e)}Ad}{16fa^2} - \frac{e^{-4i(fx+e)}Bc}{16fa^2} + \frac{A \arctan(\frac{a + b \tan(e + fx)}{a + ia \tan(e + fx)})}{4fa^2}$
derivativedivides	$\frac{iBd}{4fa^2(-i+\tan(fx+e))^2} - \frac{iAc}{4fa^2(-i+\tan(fx+e))^2} - \frac{iAd}{4fa^2(-i+\tan(fx+e))} - \frac{iBc}{4fa^2(-i+\tan(fx+e))} + \frac{A \arctan(\frac{a + b \tan(e + fx)}{a + ia \tan(e + fx)})}{4fa^2}$
default	$\frac{iBd}{4fa^2(-i+\tan(fx+e))^2} - \frac{iAc}{4fa^2(-i+\tan(fx+e))^2} - \frac{iAd}{4fa^2(-i+\tan(fx+e))} - \frac{iBc}{4fa^2(-i+\tan(fx+e))} + \frac{A \arctan(\frac{a + b \tan(e + fx)}{a + ia \tan(e + fx)})}{4fa^2}$

3.854. $\int \frac{(A+B \tan(e+fx))(c+d \tan(e+fx))}{(a+ia \tan(e+fx))^2} dx$

input `int((A+B*tan(f*x+e))*(c+d*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

output `-1/4*I*x/a^2*A*d-1/4*I*x/a^2*B*c+1/4*x/a^2*c*A-1/4*x/a^2*B*d+1/4*I/f/a^2*exp(-2*I*(f*x+e))*c*A+1/4*I/f/a^2*exp(-2*I*(f*x+e))*B*d-1/16/f/a^2*exp(-4*I*(f*x+e))*A*d-1/16/f/a^2*exp(-4*I*(f*x+e))*B*c+1/16*I/f/a^2*exp(-4*I*(f*x+e))*c*A-1/16*I/f/a^2*exp(-4*I*(f*x+e))*B*d`

3.854.5 Fracas [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.81

$$\int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx$$

$$= \frac{(4((A - iB)c - (iA + B)d)fxe^{(4i fx + 4ie)} + (iA - B)c - (A + iB)d - 4(-iAc - iBd)e^{(2i fx + 2ie)})e^{(-4i)}}{16a^2f}$$

input `integrate((A+B*tan(f*x+e))*(c+d*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`

output `1/16*(4*((A - I*B)*c - (I*A + B)*d)*f*x*e^(4*I*f*x + 4*I*e) + (I*A - B)*c - (A + I*B)*d - 4*(-I*A*c - I*B*d)*e^(2*I*f*x + 2*I*e))*e^(-4*I*f*x - 4*I*e)/(a^2*f)`

3.854.6 Sympy [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 296, normalized size of antiderivative = 2.85

$$\int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx$$

$$= \begin{cases} \frac{((16iAa^2cfe^{4ie} + 16iBa^2dfe^{4ie})e^{-2ifx} + (4iAa^2cfe^{2ie} - 4Aa^2dfe^{2ie} - 4Ba^2cfe^{2ie} - 4iBa^2dfe^{2ie})e^{-4ifx})e^{-6ie}}{64a^4f^2} & \text{for } a^4f^2e^{6ie} \neq 0 \\ x \left(-\frac{Ac - iAd - iBc - Bd}{4a^2} + \frac{(Ace^{4ie} + 2Ace^{2ie} + Ac - iAde^{4ie} + iAd - iBce^{4ie} + iBc - Bde^{4ie} + 2Bde^{2ie} - Bd)e^{-4ie}}{4a^2} \right) & \text{otherwise} \\ + \frac{x(Ac - iAd - iBc - Bd)}{4a^2} \end{cases}$$

3.854. $\int \frac{(A+B \tan(e+fx))(c+d \tan(e+fx))}{(a+ia \tan(e+fx))^2} dx$

input `integrate((A+B*tan(f*x+e))*(c+d*tan(f*x+e))/(a+I*a*tan(f*x+e))**2,x)`

output `Piecewise((((16*I*A*a**2*c*f*exp(4*I*e) + 16*I*B*a**2*d*f*exp(4*I*e))*exp(-2*I*f*x) + (4*I*A*a**2*c*f*exp(2*I*e) - 4*A*a**2*d*f*exp(2*I*e) - 4*B*a**2*c*f*exp(2*I*e) - 4*I*B*a**2*d*f*exp(2*I*e))*exp(-4*I*f*x))*exp(-6*I*e)/(64*a**4*f**2), Ne(a**4*f**2*exp(6*I*e), 0)), (x*(-(A*c - I*A*d - I*B*c - B*d)/(4*a**2) + (A*c*exp(4*I*e) + 2*A*c*exp(2*I*e) + A*c - I*A*d*exp(4*I*e) + I*A*d - I*B*c*exp(4*I*e) + I*B*c - B*d*exp(4*I*e) + 2*B*d*exp(2*I*e) - B*d)*exp(-4*I*e)/(4*a**2)), True)) + x*(A*c - I*A*d - I*B*c - B*d)/(4*a**2)`

3.854.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((A+B*tan(f*x+e))*(c+d*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

3.854.8 Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 185 vs. $2(80) = 160$.

Time = 0.49 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.78

$$\int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx = \frac{2(-iAc - Bc - Ad + iBd) \log(\tan(fx+e)+i)}{a^2} + \frac{2(iAc + Bc + Ad - iBd) \log(\tan(fx+e)-i)}{a^2} + \frac{-3iAc \tan(fx+e)^2 - 3Bc \tan(fx+e)^2 - 3Ad \tan(fx+e)^2 + 3iBd \tan(fx+e)^2}{a^2}$$

16 f

input `integrate((A+B*tan(f*x+e))*(c+d*tan(f*x+e))/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

3.854. $\int \frac{(A+B \tan(e+fx))(c+d \tan(e+fx))}{(a+ia \tan(e+fx))^2} dx$

output
$$\frac{-1/16*(2*(-I*A*c - B*c - A*d + I*B*d)*\log(\tan(f*x + e) + I)/a^2 + 2*(I*A*c + B*c + A*d - I*B*d)*\log(\tan(f*x + e) - I)/a^2 + (-3*I*A*c*\tan(f*x + e)^2 - 3*B*c*\tan(f*x + e)^2 - 3*A*d*\tan(f*x + e)^2 + 3*I*B*d*\tan(f*x + e)^2 - 10*A*c*\tan(f*x + e) + 10*I*B*c*\tan(f*x + e) + 10*I*A*d*\tan(f*x + e) - 6*B*d*\tan(f*x + e) + 11*I*A*c + 3*B*c + 3*A*d + 5*I*B*d)/(a^2*(\tan(f*x + e) - I)^2))/f}$$

3.854.9 Mupad [B] (verification not implemented)

Time = 8.73 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.53

$$\int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^2} dx = -\frac{B d f x - A c f x + A d f x \operatorname{li} + B c f x \operatorname{li}}{4 a^2 f} + \frac{(A c + 3 B d - A d \operatorname{li} - B c \operatorname{li}) \tan(e + f x)^3 + (2 A d + 2 B c + B d 4 i) \tan(e + f x)^2 + (3 A c + B d + f (4 a^2 \tan(e + f x)^4 + 8 a^2 \tan(e + f x)^2 + 4 a^2))}{f (4 a^2 \tan(e + f x)^4 + 8 a^2 \tan(e + f x)^2 + 4 a^2)}$$

input `int(((A + B*tan(e + f*x))*(c + d*tan(e + f*x)))/(a + a*tan(e + f*x)*1i)^2, x)`

output
$$\frac{(A*c*2i + B*d*2i + \tan(e + f*x)*(3*A*c + A*d*1i + B*c*1i + B*d) + \tan(e + f*x)^2*(2*A*d + 2*B*c + B*d*4i) + \tan(e + f*x)^3*(A*c - A*d*1i - B*c*1i + 3*B*d))/(f*(4*a^2 + 8*a^2*\tan(e + f*x)^2 + 4*a^2*\tan(e + f*x)^4)) - (A*d*f*x*1i - A*c*f*x + B*c*f*x*1i + B*d*f*x)/(4*a^2*f)}$$

3.855 $\int \frac{(A+B \tan(e+fx))(c+d \tan(e+fx))}{(a+ia \tan(e+fx))^{3/2}} dx$

3.855.1 Optimal result 7645
 3.855.2 Mathematica [A] (verified) 7645
 3.855.3 Rubi [A] (verified) 7646
 3.855.4 Maple [A] (verified) 7648
 3.855.5 Fricas [B] (verification not implemented) 7649
 3.855.6 Sympy [F] 7649
 3.855.7 Maxima [A] (verification not implemented) 7650
 3.855.8 Giac [F] 7650
 3.855.9 Mupad [B] (verification not implemented) 7651

3.855.1 Optimal result

Integrand size = 38, antiderivative size = 147

$$\int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx =$$

$$-\frac{(iA + B)(c - id) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(e+fx)}}{\sqrt{2}\sqrt{a}}\right)}{2\sqrt{2}a^{3/2}f}$$

$$+ \frac{(iA - B)(c + id)}{3f(a + ia \tan(e + fx))^{3/2}} + \frac{B(c + 3id) + A(ic + d)}{2af\sqrt{a + ia \tan(e + fx)}}$$

```
output -1/4*(I*A+B)*(c-I*d)*arctanh(1/2*(a+I*a*tan(f*x+e))^(1/2)*2^(1/2)/a^(1/2))
/a^(3/2)/f*2^(1/2)+1/2*(B*(c+3*I*d)+A*(I*c+d))/a/f/(a+I*a*tan(f*x+e))^(1/2)
)+1/3*(I*A-B)*(c+I*d)/f/(a+I*a*tan(f*x+e))^(3/2)
```

3.855.2 Mathematica [A] (verified)

Time = 2.52 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.96

$$\int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx =$$

$$i \left(\frac{3\sqrt{2}(A-iB)(c-id) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(e+fx)}}{\sqrt{2}\sqrt{a}}\right)}{\sqrt{a}} - \frac{4a(A+iB)(c+id)}{(a+ia \tan(e+fx))^{3/2}} - \frac{6(Ac-iBc-iAd+3Bd)}{\sqrt{a+ia \tan(e+fx)}} \right)$$

$$12af$$

input `Integrate[((A + B*Tan[e + f*x])*(c + d*Tan[e + f*x]))/(a + I*a*Tan[e + f*x])^(3/2),x]`

output `((-1/12*I)*((3*Sqrt[2]*(A - I*B)*(c - I*d)*ArcTanh[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a])])/Sqrt[a] - (4*a*(A + I*B)*(c + I*d))/(a + I*a*Tan[e + f*x])^(3/2) - (6*(A*c - I*B*c - I*A*d + 3*B*d))/Sqrt[a + I*a*Tan[e + f*x]]))/(a*f)`

3.855.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.184$, Rules used = {3042, 4073, 3042, 4009, 3042, 3961, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{3042} \\
 & \int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx \\
 & \quad \downarrow \text{4073} \\
 & \frac{(-B + iA)(c + id)}{3f(a + ia \tan(e + fx))^{3/2}} - \frac{i \int \frac{a(B(c+id)+A(ic+d))+2aBd \tan(e+fx)}{\sqrt{i \tan(e+fx)a+a}} dx}{2a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B + iA)(c + id)}{3f(a + ia \tan(e + fx))^{3/2}} - \frac{i \int \frac{a(B(c+id)+A(ic+d))+2aBd \tan(e+fx)}{\sqrt{i \tan(e+fx)a+a}} dx}{2a^2} \\
 & \quad \downarrow \text{4009} \\
 & \frac{(-B + iA)(c + id)}{3f(a + ia \tan(e + fx))^{3/2}} - \frac{i \left(\frac{1}{2}(B + iA)(c - id) \int \sqrt{i \tan(e + fx)a + a} dx - \frac{a(Ac - iAd - iBc + 3Bd)}{f \sqrt{a + ia \tan(e + fx)}} \right)}{2a^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{(-B + iA)(c + id)}{3f(a + ia \tan(e + fx))^{3/2}} - \frac{i \left(\frac{1}{2}(B + iA)(c - id) \int \sqrt{i \tan(e + fx)a + a} dx - \frac{a(Ac - iAd - iBc + 3Bd)}{f \sqrt{a + ia \tan(e + fx)}} \right)}{2a^2}
 \end{aligned}$$

3.855. $\int \frac{(A+B \tan(e+fx))(c+d \tan(e+fx))}{(a+ia \tan(e+fx))^{3/2}} dx$

$$\frac{(-B + iA)(c + id)}{3f(a + ia \tan(e + fx))^{3/2}} - \frac{i \left(-\frac{ia(B+iA)(c-id) \int \frac{1}{a-ia \tan(e+fx)} d\sqrt{i \tan(e+fx)a+a} - \frac{a(Ac-iAd-iBc+3Bd)}{f\sqrt{a+ia \tan(e+fx)}} \right)}{2a^2}$$

$$\frac{(-B + iA)(c + id)}{3f(a + ia \tan(e + fx))^{3/2}} - \frac{i \left(-\frac{i\sqrt{a}(B+iA)(c-id) \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(e+fx)}}{\sqrt{2}\sqrt{a}}\right) - \frac{a(Ac-iAd-iBc+3Bd)}{f\sqrt{a+ia \tan(e+fx)}} \right)}{2a^2}$$

input `Int[((A + B*Tan[e + f*x])*(c + d*Tan[e + f*x]))/(a + I*a*Tan[e + f*x])^(3/2), x]`

output `((I*A - B)*(c + I*d))/(3*f*(a + I*a*Tan[e + f*x])^(3/2)) - ((I/2)*(((-I)*Sqrt[a]*(I*A + B)*(c - I*d)*ArcTanh[Sqrt[a + I*a*Tan[e + f*x]]/(Sqrt[2]*Sqrt[a]])/(Sqrt[2]*f) - (a*(A*c - I*B*c - I*A*d + 3*B*d))/(f*Sqrt[a + I*a*Tan[e + f*x]])))/a^2`

3.855.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3961 `Int[Sqrt[(a_) + (b_.)*tan[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[-2*(b/d) Subst[Int[1/(2*a - x^2), x], x, Sqrt[a + b*Tan[c + d*x]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0]`

```
rule 4009 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((c_) + (d_)*tan[(e_) +
(f_)*(x_)]), x_Symbol] := Simp[(-(b*c - a*d))*((a + b*Tan[e + f*x])^m/(2*a
*f*m)), x] + Simp[(b*c + a*d)/(2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1), x
], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 + b^2
, 0] && LtQ[m, 0]
```

```
rule 4073 Int[((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(m_)*((A_) + (B_)*tan[(e_) +
(f_)*(x_)])*((c_) + (d_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(-(
A*b - a*B))*(a*c + b*d)*((a + b*Tan[e + f*x])^m/(2*a^2*f*m)), x] + Simp[1/(
2*a*b) Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[A*b*c + a*B*c + a*A*d + b*B*
d + 2*a*B*d*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, c, d, e, f, A, B}, x]
&& NeQ[b*c - a*d, 0] && LtQ[m, -1] && EqQ[a^2 + b^2, 0]
```

3.855.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.89

method	result
derivativedivides	$2i \left(-\frac{\frac{1}{4}iAd + \frac{1}{4}iBc - \frac{1}{4}cA - \frac{3}{4}Bd}{\sqrt{a+ia \tan(fx+e)}} + \frac{a(iAd+iBc+cA-Bd)}{6(a+ia \tan(fx+e))^{\frac{3}{2}}} - \frac{\left(-\frac{1}{4}iAd - \frac{1}{4}iBc + \frac{1}{4}cA - \frac{1}{4}Bd\right)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(fx+e)}\sqrt{2}}{2\sqrt{a}}\right)}{2\sqrt{a}} \right) \frac{1}{fa}$
default	$2i \left(-\frac{\frac{1}{4}iAd + \frac{1}{4}iBc - \frac{1}{4}cA - \frac{3}{4}Bd}{\sqrt{a+ia \tan(fx+e)}} + \frac{a(iAd+iBc+cA-Bd)}{6(a+ia \tan(fx+e))^{\frac{3}{2}}} - \frac{\left(-\frac{1}{4}iAd - \frac{1}{4}iBc + \frac{1}{4}cA - \frac{1}{4}Bd\right)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(fx+e)}\sqrt{2}}{2\sqrt{a}}\right)}{2\sqrt{a}} \right) \frac{1}{fa}$
parts	$\frac{2i c A a \left(\frac{1}{4a^2 \sqrt{a+ia \tan(fx+e)}} + \frac{1}{6a(a+ia \tan(fx+e))^{\frac{3}{2}}} - \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{a+ia \tan(fx+e)}\sqrt{2}}{2\sqrt{a}}\right)}{8a^{\frac{5}{2}}} \right)}{f} + \frac{(Ad+cB) \left(-\frac{1}{3(a+ia \tan(fx+e))} \right)}{f}$

```
input int((A+B*tan(f*x+e))*(c+d*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2),x,method=_R
ETURNVERBOSE)
```

```
output 2*I/f/a*(-(1/4*I*A*d+1/4*I*B*c-1/4*c*A-3/4*B*d)/(a+I*a*tan(f*x+e))^(1/2)+1
/6*a*(-B*d+I*A*d+I*B*c+c*A)/(a+I*a*tan(f*x+e))^(3/2)-1/2*(-1/4*I*A*d-1/4*I
*B*c+1/4*c*A-1/4*B*d)*2^(1/2)/a^(1/2)*arctanh(1/2*(a+I*a*tan(f*x+e))^(1/2)
*2^(1/2)/a^(1/2)))
```

$$3.855. \int \frac{(A+B \tan(e+fx))(c+d \tan(e+fx))}{(a+ia \tan(e+fx))^{3/2}} dx$$

3.855.5 Fracas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 619 vs. $2(108) = 216$.

Time = 0.25 (sec) , antiderivative size = 619, normalized size of antiderivative = 4.21

$$\int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx = \frac{\left(3 \sqrt{\frac{1}{2}} a^2 f \sqrt{-\frac{(A^2 - 2iAB - B^2)c^2 + 2(-iA^2 - 2AB + iB^2)cd - (A^2 - 2iAB - B^2)d^2}{a^3 f^2}} \right)}{}$$

input `integrate((A+B*tan(f*x+e))*(c+d*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="fricas")`

output `1/12*(3*sqrt(1/2)*a^2*f*sqrt(-((A^2 - 2*I*A*B - B^2)*c^2 + 2*(-I*A^2 - 2*A*B + I*B^2)*c*d - (A^2 - 2*I*A*B - B^2)*d^2)/(a^3*f^2))*e^(3*I*f*x + 3*I*e)*log(4*(sqrt(2)*sqrt(1/2)*(I*a^2*f*e^(2*I*f*x + 2*I*e) + I*a^2*f)*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-((A^2 - 2*I*A*B - B^2)*c^2 + 2*(-I*A^2 - 2*A*B + I*B^2)*c*d - (A^2 - 2*I*A*B - B^2)*d^2)/(a^3*f^2)) + ((A - I*B)*a*c + (-I*A - B)*a*d)*e^(I*f*x + I*e))*e^(-I*f*x - I*e)/((A - I*B)*c - (I*A + B)*d) - 3*sqrt(1/2)*a^2*f*sqrt(-((A^2 - 2*I*A*B - B^2)*c^2 + 2*(-I*A^2 - 2*A*B + I*B^2)*c*d - (A^2 - 2*I*A*B - B^2)*d^2)/(a^3*f^2))*e^(3*I*f*x + 3*I*e)*log(4*(sqrt(2)*sqrt(1/2)*(-I*a^2*f*e^(2*I*f*x + 2*I*e) - I*a^2*f)*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*sqrt(-((A^2 - 2*I*A*B - B^2)*c^2 + 2*(-I*A^2 - 2*A*B + I*B^2)*c*d - (A^2 - 2*I*A*B - B^2)*d^2)/(a^3*f^2)) + ((A - I*B)*a*c + (-I*A - B)*a*d)*e^(I*f*x + I*e))*e^(-I*f*x - I*e)/((A - I*B)*c - (I*A + B)*d) + sqrt(2)*((I*A - B)*c - (A + I*B)*d - 2*((-2*I*A - B)*c - (A + 4*I*B)*d))*e^(4*I*f*x + 4*I*e) + ((5*I*A + B)*c + (A + 7*I*B)*d)*e^(2*I*f*x + 2*I*e))*sqrt(a/(e^(2*I*f*x + 2*I*e) + 1))*e^(-3*I*f*x - 3*I*e)/(a^2*f)`

3.855.6 Sympy [F]

$$\int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx = \int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(ia(\tan(e + fx) - i))^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(f*x+e))*(c+d*tan(f*x+e))/(a+I*a*tan(f*x+e))**(3/2),x)`

3.855. $\int \frac{(A+B \tan(e+fx))(c+d \tan(e+fx))}{(a+ia \tan(e+fx))^{3/2}} dx$

output `Integral((A + B*tan(e + f*x))*(c + d*tan(e + f*x))/(I*a*(tan(e + f*x) - I)**(3/2), x)`

3.855.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.97

$$\int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx = \frac{i \left(\frac{3\sqrt{2}(A-iB)(c-id) \log\left(-\frac{\sqrt{2}\sqrt{a}-\sqrt{ia \tan(fx+e)+a}}{\sqrt{2}\sqrt{a}+\sqrt{ia \tan(fx+e)+a}}\right)}{\sqrt{a}} \right) + \frac{4(2(A+iB)ac+24af}{24af}}$$

input `integrate((A+B*tan(f*x+e))*(c+d*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="maxima")`

output `1/24*I*(3*sqrt(2)*(A - I*B)*(c - I*d)*log(-(sqrt(2)*sqrt(a) - sqrt(I*a*tan(f*x + e) + a))/(sqrt(2)*sqrt(a) + sqrt(I*a*tan(f*x + e) + a)))/sqrt(a) + 4*(2*(A + I*B)*a*c + 2*(I*A - B)*a*d + 3*((A - I*B)*c + (-I*A + 3*B)*d)*(I*a*tan(f*x + e) + a))/(I*a*tan(f*x + e) + a)^(3/2))/(a*f)`

3.855.8 Giac [F]

$$\int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx = \int \frac{(B \tan(fx + e) + A)(d \tan(fx + e) + c)}{(ia \tan(fx + e) + a)^{\frac{3}{2}}} dx$$

input `integrate((A+B*tan(f*x+e))*(c+d*tan(f*x+e))/(a+I*a*tan(f*x+e))^(3/2),x, algorithm="giac")`

output `integrate((B*tan(f*x + e) + A)*(d*tan(f*x + e) + c)/(I*a*tan(f*x + e) + a)^(3/2), x)`

3.855.9 Mupad [B] (verification not implemented)

Time = 1.59 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.67

$$\int \frac{(A + B \tan(e + fx))(c + d \tan(e + fx))}{(a + ia \tan(e + fx))^{3/2}} dx = \frac{\frac{(Ac + Ad \operatorname{li}) \operatorname{li}}{3f} + \frac{(Ac - Ad \operatorname{li})(a + a \tan(e + fx) \operatorname{li}) \operatorname{li}}{2af}}{(a + a \tan(e + fx) \operatorname{li})^{3/2}} - \frac{\frac{Bc + Bd \operatorname{li}}{3f} - \frac{(Bc + Bd \operatorname{li})(a + a \tan(e + fx) \operatorname{li})}{2af}}{(a + a \tan(e + fx) \operatorname{li})^{3/2}} + \frac{\sqrt{2} B \operatorname{atanh}\left(\frac{\sqrt{2} B (d + c \operatorname{li}) \sqrt{a + a \tan(e + fx) \operatorname{li}}}{2\sqrt{-a}(Bc - Bd \operatorname{li})}\right) (d + c \operatorname{li})}{4(-a)^{3/2} f} + \frac{\sqrt{2} A \operatorname{atan}\left(\frac{\sqrt{2} A (d + c \operatorname{li}) \sqrt{a + a \tan(e + fx) \operatorname{li}}}{2\sqrt{a}(Ac - Ad \operatorname{li})}\right) (d + c \operatorname{li}) \operatorname{li}}{4a^{3/2} f}$$

input `int(((A + B*tan(e + f*x))*(c + d*tan(e + f*x)))/(a + a*tan(e + f*x)*1i)^(3/2),x)`

output `((((A*c + A*d*1i)*1i)/(3*f) + ((A*c - A*d*1i)*(a + a*tan(e + f*x)*1i)*1i)/(2*a*f))/(a + a*tan(e + f*x)*1i)^(3/2) - ((B*c + B*d*1i)/(3*f) - ((B*c + B*d*3i)*(a + a*tan(e + f*x)*1i))/(2*a*f))/(a + a*tan(e + f*x)*1i)^(3/2) + (2^(1/2)*B*atanh((2^(1/2)*B*(c*1i + d)*(a + a*tan(e + f*x)*1i)^(1/2))/(2*(-a)^(1/2)*(B*c - B*d*1i)))*(c*1i + d))/(4*(-a)^(3/2)*f) + (2^(1/2)*A*atan((2^(1/2)*A*(c*1i + d)*(a + a*tan(e + f*x)*1i)^(1/2))/(2*a^(1/2)*(A*c - A*d*1i)))*(c*1i + d)*1i)/(4*a^(3/2)*f)`

APPENDIX

4.1 Listing of Grading functions	7652
--	------

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.1.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
```

```

(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A"," "}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A"," "}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.$
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
      ,
      finalresult={"F","Contains unresolved integral."}
    ]
  ];

  finalresult
]

```

```

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType,expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]],2]],
            Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3,ExpnType[expn[[1]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
            If[Head[expn]===RootSum,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

4.1.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

```

```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of
                        convert(leaf_count_result,string)," vs. $2 (" ,
                        convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(",
                        convert(leaf_count_optimal,string)," )=" ,convert(2*leaf_cou
        fi;
      fi;
    fi;
  fi;

```



```

else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  end if
end if

```

```

elif type(expn, ``~`) then
  if type(op(2,expn), 'integer') then
    ExpnType(op(1,expn))
  elif type(op(2,expn), 'rational') then
    if type(op(1,expn), 'rational') then
      1
    else
      max(2, ExpnType(op(1,expn)))
    end if
  else
    max(3, ExpnType(op(1,expn)), ExpnType(op(2,expn)))
  end if
elif type(expn, ``+`) or type(expn, ``*`) then
  max(ExpnType(op(1,expn)), max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3, ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4, apply(max, map(ExpnType, [op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5, apply(max, map(ExpnType, [op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6, apply(max, map(ExpnType, [op(expn)])))
elif op(0,expn)='int' then
  max(8, apply(max, map(ExpnType, [op(expn)]))) else
  9
end if
end proc:

```

```

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

```

```

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,

```

```

        GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
        EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
    member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u),op(2..nops(u),u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

4.1.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):

```

```
if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
        return True
    else:
        return False
else:
    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
```

```

return 1
elif isinstance(expn,list):
    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
        return 1
    else:
        return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]]
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

#main function

```

def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

    #print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```

4.1.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
```

```
        return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m
```



```

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric', 'hypergeometric_M', 'hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=", expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception, AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:  #isinstance(expn,list):

```

```

    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
elif expn.operator() == operator.pow: #isinstance(expn,Pow)
    if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
        return expnType(expn.operands()[0]) #expnType(expn.args[0])
    elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
        if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

```

```

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)",type(result))
    print("type(optimal)",type(optimal))

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal." + str(leaf
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```